

Computer Algebra Independent Integration Tests

Summer 2024

0-Independent-test-suites/1-Apostol-Problems

Nasser M. Abbasi

May 17, 2024

Compiled on May 17, 2024 at 8:57pm

Contents

1	Introduction	7
1.1	Listing of CAS systems tested	8
1.2	Results	9
1.3	Time and leaf size Performance	13
1.4	Performance based on number of rules Rubi used	15
1.5	Performance based on number of steps Rubi used	16
1.6	Solved integrals histogram based on leaf size of result	17
1.7	Solved integrals histogram based on CPU time used	18
1.8	Leaf size vs. CPU time used	19
1.9	list of integrals with no known antiderivative	20
1.10	List of integrals solved by CAS but has no known antiderivative	20
1.11	list of integrals solved by CAS but failed verification	20
1.12	Timing	21
1.13	Verification	21
1.14	Important notes about some of the results	22
1.15	Current tree layout of integration tests	25
1.16	Design of the test system	26
2	detailed summary tables of results	27
2.1	List of integrals sorted by grade for each CAS	28
2.2	Detailed conclusion table per each integral for all CAS systems	33
2.3	Detailed conclusion table specific for Rubi results	77
3	Listing of integrals	83
3.1	$\int \sqrt{1 + 2x} dx$	89
3.2	$\int x\sqrt{1 + 3x} dx$	94
3.3	$\int x^2\sqrt{1 + x} dx$	99
3.4	$\int \frac{x}{\sqrt{2-3x}} dx$	104
3.5	$\int \frac{1+x}{(2+2x+x^2)^3} dx$	109
3.6	$\int \sin^3(x) dx$	114

3.7	$\int \sqrt[3]{-1 + zz} dz$	119
3.8	$\int \cot(x) \csc^2(x) dx$	124
3.9	$\int \cos(2x) \sqrt{4 - \sin(2x)} dx$	129
3.10	$\int \frac{\sin(x)}{(3 + \cos(x))^2} dx$	134
3.11	$\int \frac{\sin(x)}{\sqrt{\cos^3(x)}} dx$	139
3.12	$\int \frac{\sin(\sqrt{1+x})}{\sqrt{1+x}} dx$	145
3.13	$\int x^{-1+n} \sin(x^n) dx$	150
3.14	$\int \frac{x^5}{\sqrt{1-x^6}} dx$	155
3.15	$\int t \sqrt[4]{1+t} dt$	160
3.16	$\int \frac{1}{(1+x^2)^{3/2}} dx$	165
3.17	$\int x^2 (27 + 8x^3)^{2/3} dx$	170
3.18	$\int \frac{\cos(x) + \sin(x)}{\sqrt[3]{-\cos(x) + \sin(x)}} dx$	175
3.19	$\int \frac{x}{\sqrt{1+x^2 + (1+x^2)^{3/2}}} dx$	180
3.20	$\int \frac{x}{\sqrt{1+x^2} \sqrt{1+\sqrt{1+x^2}}} dx$	185
3.21	$\int \frac{\sqrt[5]{1-2x+x^2}}{1-x} dx$	190
3.22	$\int x \sin(x) dx$	195
3.23	$\int x^2 \sin(x) dx$	200
3.24	$\int x^3 \cos(x) dx$	205
3.25	$\int x^3 \sin(x) dx$	211
3.26	$\int \cos(x) \sin(x) dx$	217
3.27	$\int x \cos(x) \sin(x) dx$	222
3.28	$\int \sin^2(x) dx$	227
3.29	$\int \sin^3(x) dx$	232
3.30	$\int \sin^4(x) dx$	237
3.31	$\int \sin^5(x) dx$	242
3.32	$\int \sin^6(x) dx$	247
3.33	$\int x \sin^2(x) dx$	252
3.34	$\int x \sin^3(x) dx$	257
3.35	$\int x^2 \sin^2(x) dx$	262
3.36	$\int \cos^2(x) dx$	268
3.37	$\int \cos^3(x) dx$	273
3.38	$\int \cos^4(x) dx$	278
3.39	$\int (a^2 - x^2)^{5/2} dx$	283
3.40	$\int \frac{x^5}{\sqrt{5+x^2}} dx$	289
3.41	$\int \frac{t^3}{\sqrt{4+t^3}} dt$	294
3.42	$\int \tan^2(x) dx$	300

3.43	$\int \tan^4(x) dx$	305
3.44	$\int \cot^2(x) dx$	310
3.45	$\int \cot^4(x) dx$	315
3.46	$\int (2 + 3x) \sin(5x) dx$	320
3.47	$\int x\sqrt{1+x^2} dx$	325
3.48	$\int x(-1+x^2)^9 dx$	330
3.49	$\int \frac{3+2x}{(7+6x)^3} dx$	335
3.50	$\int x^4(1+x^5)^5 dx$	340
3.51	$\int (1-x)^{20} x^4 dx$	345
3.52	$\int \frac{\sin(\frac{1}{x})}{x^2} dx$	352
3.53	$\int \sin(\sqrt[4]{-1+x}) dx$	357
3.54	$\int x \cos(x^2) \sin(x^2) dx$	363
3.55	$\int \sqrt{1+3\cos^2(x)} \sin(2x) dx$	368
3.56	$\int \frac{1}{2+3x} dx$	373
3.57	$\int \log^2(x) dx$	378
3.58	$\int x \log(x) dx$	383
3.59	$\int x \log^2(x) dx$	388
3.60	$\int \frac{1}{1+t} dt$	393
3.61	$\int \cot(x) dx$	398
3.62	$\int x^n \log(ax) dx$	403
3.63	$\int x^2 \log^2(x) dx$	408
3.64	$\int \frac{1}{x \log(x)} dx$	413
3.65	$\int \frac{\log(1-t)}{1-t} dt$	418
3.66	$\int \frac{\log(x)}{x\sqrt{1+\log(x)}} dx$	423
3.67	$\int x^3 \log^3(x) dx$	428
3.68	$\int e^{x^3} x^2 dx$	433
3.69	$\int \frac{2\sqrt{x}}{\sqrt{x}} dx$	438
3.70	$\int e^{2\sin(x)} \cos(x) dx$	443
3.71	$\int e^x \sin(x) dx$	448
3.72	$\int e^x \cos(x) dx$	453
3.73	$\int \frac{1}{1+e^x} dx$	458
3.74	$\int e^x x dx$	463
3.75	$\int e^{-x} x dx$	468
3.76	$\int e^x x^2 dx$	473
3.77	$\int e^{-2x} x^2 dx$	478
3.78	$\int e^{\sqrt{x}} dx$	483
3.79	$\int e^{-x^2} x^3 dx$	488
3.80	$\int e^{ax} \cos(bx) dx$	493

3.81	$\int e^{ax} \sin(bx) dx$	498
3.82	$\int \cot^{-1}(x) dx$	503
3.83	$\int \sec^{-1}(x) dx$	508
3.84	$\int \csc^{-1}(x) dx$	514
3.85	$\int \arcsin(x)^2 dx$	520
3.86	$\int \frac{\arcsin(x)}{x^2} dx$	525
3.87	$\int \frac{1}{\sqrt{a^2-x^2}} dx$	530
3.88	$\int \frac{1}{\sqrt{1-2x-x^2}} dx$	535
3.89	$\int \frac{1}{a^2+x^2} dx$	540
3.90	$\int \frac{1}{a+bx^2} dx$	545
3.91	$\int \frac{1}{2-x+x^2} dx$	550
3.92	$\int x \arctan(x) dx$	555
3.93	$\int x^2 \arccos(x) dx$	560
3.94	$\int x \arctan(x)^2 dx$	565
3.95	$\int \arctan(\sqrt{x}) dx$	571
3.96	$\int \frac{\arctan(\sqrt{x})}{\sqrt{x(1+x)}} dx$	577
3.97	$\int \sqrt{1-x^2} dx$	582
3.98	$\int \frac{e^{\arctan(x)} x}{(1+x^2)^{3/2}} dx$	587
3.99	$\int \frac{e^{\arctan(x)}}{(1+x^2)^{3/2}} dx$	592
3.100	$\int \frac{x^2}{(1+x^2)^2} dx$	597
3.101	$\int \frac{e^x}{1+e^{2x}} dx$	602
3.102	$\int e^{-x} \cot^{-1}(e^x) dx$	607
3.103	$\int \sqrt{\frac{a+x}{a-x}} dx$	613
3.104	$\int \sqrt{(b-x)(-a+x)} dx$	618
3.105	$\int \frac{1}{\sqrt{(b-x)(-a+x)}} dx$	624
3.106	$\int \frac{3+5x}{-3+2x+x^2} dx$	630
3.107	$\int \frac{5+2x}{-3+2x+x^2} dx$	635
3.108	$\int \frac{3x+x^3}{-3-2x+x^2} dx$	640
3.109	$\int \frac{-1+5x+2x^2}{-2x+x^2+x^3} dx$	645
3.110	$\int \frac{3+2x+x^2}{(-1+x)(1+x)^2} dx$	650
3.111	$\int \frac{-2+2x+3x^2}{-1+x^3} dx$	655
3.112	$\int \frac{2-x+2x^2-x^3+x^4}{(-1+x)(2+x^2)^2} dx$	661
3.113	$\int \frac{1}{\cos(x)+\sin(x)} dx$	667
3.114	$\int \frac{x}{4-x^2+\sqrt{4-x^2}} dx$	672
3.115	$\int \frac{3+2x}{(-2+x)(5+x)} dx$	677
3.116	$\int \frac{x}{(1+x)(2+x)(3+x)} dx$	682

3.117	$\int \frac{x}{2-3x+x^3} dx$	687
3.118	$\int \frac{-6+2x+x^4}{-2x+x^2+x^3} dx$	692
3.119	$\int \frac{7+8x^3}{(1+x)(1+2x)^3} dx$	697
3.120	$\int \frac{1+x+4x^2}{-1+x^3} dx$	702
3.121	$\int \frac{x^4}{4+5x^2+x^4} dx$	707
3.122	$\int \frac{2+x}{x+x^2} dx$	712
3.123	$\int \frac{1}{x(1+x^2)^2} dx$	717
3.124	$\int \frac{1}{(1+x)(2+x)^2(3+x)^3} dx$	722
3.125	$\int \frac{x}{(1+x)^2} dx$	728
3.126	$\int \frac{1}{-x+x^3} dx$	733
3.127	$\int \frac{x^2}{-6+x+x^2} dx$	739
3.128	$\int \frac{2+x}{4-4x+x^2} dx$	744
3.129	$\int \frac{1}{(4-4x+x^2)(5-4x+x^2)} dx$	749
3.130	$\int \frac{-3+x}{2x+3x^2+x^3} dx$	754
3.131	$\int \frac{1}{(-1+x^2)^2} dx$	759
3.132	$\int \frac{1+x}{-1+x^3} dx$	764
3.133	$\int \frac{1+x^4}{x(1+x^2)^2} dx$	769
3.134	$\int \frac{1}{-2x^3+x^4} dx$	774
3.135	$\int \frac{1-x^3}{x(1+x^2)} dx$	779
3.136	$\int \frac{1}{-1+x^4} dx$	784
3.137	$\int \frac{1}{1+x^4} dx$	789
3.138	$\int \frac{x^2}{(2+2x+x^2)^2} dx$	797
3.139	$\int \frac{-1+4x^5}{(1+x+x^5)^2} dx$	802
3.140	$\int \frac{1}{5-\cos(x)+2\sin(x)} dx$	807
3.141	$\int \frac{1}{1+a\cos(x)} dx$	813
3.142	$\int \frac{1}{1+2\cos(x)} dx$	819
3.143	$\int \frac{1}{1+\cos(x)} dx$	824
3.144	$\int \frac{\sin^2(x)}{1+\sin^2(x)} dx$	829
3.145	$\int \frac{1}{b^2\cos^2(x)+a^2\sin^2(x)} dx$	835
3.146	$\int \frac{1}{(b\cos(x)+a\sin(x))^2} dx$	841
3.147	$\int \frac{\sin(x)}{1+\cos(x)+\sin(x)} dx$	847
3.148	$\int \sqrt{3-x^2} dx$	853
3.149	$\int \frac{x}{\sqrt{3-x^2}} dx$	858
3.150	$\int \frac{\sqrt{3-x^2}}{x} dx$	863
3.151	$\int \frac{\sqrt{x+x^2}}{x} dx$	869
3.152	$\int \sqrt{5+x^2} dx$	874

3.153	$\int \frac{x}{\sqrt{1+x+x^2}} dx$	879
3.154	$\int \frac{1}{\sqrt{x+x^2}} dx$	884
3.155	$\int \frac{\sqrt{2-x-x^2}}{x^2} dx$	889
3.156	$\int \frac{\log(t)}{1+t} dt$	896
3.157	$\int \log(e^{\cos(x)}) dx$	901
3.158	$\int \frac{e^t}{t} dt$	906
3.159	$\int \frac{e^{at}}{t} dt$	911
3.160	$\int \frac{e^t}{t^2} dt$	916
3.161	$\int e^{\frac{1}{t}} dt$	921
3.162	$\int \frac{e^{-t}}{-1-a+t} dt$	926
3.163	$\int \frac{e^{t^2} t}{1+t^2} dt$	930
3.164	$\int \frac{e^t}{(1+t)^2} dt$	935
3.165	$\int e^t \log(1+t) dt$	940
3.166	$\int e^{-t} t dt$	945
3.167	$\int e^{-t} t^2 dt$	950
3.168	$\int e^{-t} t^3 dt$	955
3.169	$\int \frac{b \cos(x) + a \sin(x)}{b \cos(x) + a \sin(x)} dx$	960
3.170	$\int \frac{1}{\log(t)} dt$	967
3.171	$\int \frac{1}{\log^2(t)} dt$	971
3.172	$\int \log^{-1-n}(t) dt$	976
3.173	$\int \frac{e^{2t}}{-1+t} dt$	981
3.174	$\int \frac{e^{2x}}{2-3x+x^2} dx$	986
3.175	$\int \frac{1}{\sqrt{1+t^3}} dt$	991

4 Appendix 996

4.1	Listing of Grading functions	996
4.2	Links to plain text integration problems used in this report for each CAS	1014

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	8
1.2	Results	9
1.3	Time and leaf size Performance	13
1.4	Performance based on number of rules Rubi used	15
1.5	Performance based on number of steps Rubi used	16
1.6	Solved integrals histogram based on leaf size of result	17
1.7	Solved integrals histogram based on CPU time used	18
1.8	Leaf size vs. CPU time used	19
1.9	list of integrals with no known antiderivative	20
1.10	List of integrals solved by CAS but has no known antiderivative	20
1.11	list of integrals solved by CAS but failed verification	20
1.12	Timing	21
1.13	Verification	21
1.14	Important notes about some of the results	22
1.15	Current tree layout of integration tests	25
1.16	Design of the test system	26

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [175]. This is test number [1].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (175)	0.00 (0)
Mathematica	100.00 (175)	0.00 (0)
Fricas	99.43 (174)	0.57 (1)
Maple	98.86 (173)	1.14 (2)
Giac	97.71 (171)	2.29 (4)
Mupad	96.57 (169)	3.43 (6)
Maxima	94.86 (166)	5.14 (9)
Sympy	94.29 (165)	5.71 (10)
Reduce	91.43 (160)	8.57 (15)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Maple	93.714	1.714	3.429	1.143
Mathematica	93.143	3.429	3.429	0.000
Maxima	92.000	2.857	0.000	5.143
Giac	90.857	6.857	0.000	2.286
Fricas	90.286	8.571	0.571	0.571
Sympy	78.857	8.571	6.857	5.714
Mupad	0.000	96.571	0.000	3.429
Reduce	0.000	91.429	0.000	8.571

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	1	100.00	0.00	0.00
Maple	2	100.00	0.00	0.00
Giac	4	100.00	0.00	0.00
Mupad	6	0.00	100.00	0.00
Maxima	9	66.67	0.00	33.33
Sympy	10	100.00	0.00	0.00
Reduce	15	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.05
Fricas	0.08
Mupad	0.10
Maple	0.10
Mathematica	0.14
Giac	0.15
Reduce	0.15
Rubi	0.16
Sympy	1.15

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Maple	17.79	0.90	15.00	0.82
Maxima	18.24	0.86	14.50	0.81
Giac	20.88	0.97	16.00	0.83
Reduce	21.53	1.01	17.00	0.83
Fricas	21.76	1.04	17.00	0.85
Mathematica	21.79	1.03	18.00	1.00
Rubi	23.53	1.03	19.00	1.00
Mupad	31.83	1.11	14.00	0.79
Sympy	467.13	30.41	17.00	0.87

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram shows that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {}

Maple {75, 79, 166, 167, 168}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```


For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	28
2.2	Detailed conclusion table per each integral for all CAS systems	33
2.3	Detailed conclusion table specific for Rubi results	77

2.1 List of integrals sorted by grade for each CAS

Rubi	28
Mma	29
Maple	29
Fricas	30
Maxima	30
Giac	31
Mupad	31
Sympy	32
Reduce	32

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 42, 43, 46, 47, 48, 49, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 85, 86, 87, 89, 90, 91, 92, 93, 94, 95, 96, 97, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174 }

B grade { 50, 51, 83, 84, 88, 154 }

C grade { 41, 44, 45, 98, 113, 175 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 36, 37, 38, 39, 40, 42, 43, 44, 45, 46, 47, 48, 49, 50, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 132, 133, 134, 135, 136, 138, 139, 140, 141, 142, 143, 144, 145, 146, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 171, 173, 174 }

B grade { 51, 158, 170 }

C grade { 35, 41, 131, 137, 147, 175 }

F normal fail { 19, 172 }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 46, 47, 49, 52, 53, 54, 55, 56, 57, 58, 59, 60, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 85, 86, 87, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 115, 116, 117, 118, 119, 120, 121, 122, 123, 125, 126, 127, 128, 129, 130, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 147, 148, 149, 150, 151, 152, 153, 154, 155, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 173, 174, 175 }

B grade { 16, 44, 45, 48, 50, 51, 61, 84, 88, 113, 114, 124, 131, 145, 146 }

C grade { 172 }

F normal fail { 156 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maxima

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 42, 43, 44, 45, 46, 47, 48, 49, 50, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 100, 101, 102, 103, 106, 107, 108, 109, 110, 111, 112, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 170, 171, 172, 173 }

B grade { 51, 83, 84, 113, 169 }

C grade { }

F normal fail { 19, 41, 98, 99, 174, 175 }

F(-1) timedout fail { }

F(-2) exception fail { 104, 105, 141 }

Giac

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 42, 43, 46, 47, 48, 49, 50, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 85, 86, 87, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 106, 107, 108, 109, 110, 111, 112, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 157, 158, 159, 160, 161, 162, 163, 165, 166, 167, 168, 169, 170, 171, 173, 174 }

B grade { 44, 45, 51, 83, 84, 88, 105, 113, 136, 154, 155, 164 }

C grade { }

F normal fail { 41, 156, 172, 175 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 100, 101, 102, 103, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 166, 167, 168, 169, 170, 171, 172, 173, 175 }

C grade { }

F normal fail { }

F(-1) timedout fail { 98, 99, 104, 105, 165, 174 }

F(-2) exception fail { }

Sympy

A grade { 1, 2, 3, 5, 6, 8, 10, 11, 12, 13, 14, 15, 16, 18, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 40, 41, 43, 44, 45, 46, 49, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 82, 84, 85, 86, 88, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 102, 104, 106, 107, 108, 109, 110, 111, 112, 113, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 142, 143, 147, 148, 149, 152, 153, 154, 157, 158, 159, 160, 161, 166, 167, 168, 170, 171, 172, 175 }

B grade { 9, 17, 42, 47, 48, 50, 51, 62, 90, 101, 114, 141, 144, 145, 146 }

C grade { 4, 7, 39, 80, 81, 83, 87, 89, 105, 150, 156, 169 }

F normal fail { 19, 103, 151, 155, 162, 163, 164, 165, 173, 174 }

F(-1) timedout fail { }

F(-2) exception fail { }

Reduce

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 157, 158, 159, 160, 161, 166, 167, 168, 169, 170, 171 }

C grade { }

F normal fail { 18, 41, 55, 83, 84, 145, 156, 162, 163, 164, 165, 172, 173, 174, 175 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	13	10	9	9	8	9	13	9
N.S.	1	1.00	1.00	0.77	0.69	0.69	0.62	0.69	1.00	0.69
time (sec)	N/A	0.110	0.005	0.161	0.026	0.073	0.018	0.119	0.152	0.184

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	18	15	19	19	39	19	18	14
N.S.	1	1.00	0.67	0.56	0.70	0.70	1.44	0.70	0.67	0.52
time (sec)	N/A	0.131	0.008	0.098	0.029	0.066	0.532	0.569	0.155	0.048

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	21	18	22	22	48	22	21	19
N.S.	1	1.00	0.62	0.53	0.65	0.65	1.41	0.65	0.62	0.56
time (sec)	N/A	0.132	0.038	0.072	0.029	0.064	0.709	0.177	0.153	0.041

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	18	14	19	14	60	19	13	14
N.S.	1	1.00	0.67	0.52	0.70	0.52	2.22	0.70	0.48	0.52
time (sec)	N/A	0.130	0.007	0.085	0.037	0.061	0.523	0.109	0.154	0.033

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	13	12	22	22	12	24	12
N.S.	1	1.00	1.00	0.93	0.86	1.57	1.57	0.86	1.71	0.86
time (sec)	N/A	0.119	0.006	0.187	0.030	0.071	0.047	1.183	0.146	0.038

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	15	11	11	11	8	11	14	10
N.S.	1	1.00	1.15	0.85	0.85	0.85	0.62	0.85	1.08	0.77
time (sec)	N/A	0.150	0.030	0.258	0.024	0.094	0.023	0.748	0.152	0.090

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	18	13	15	17	92	15	17	12
N.S.	1	1.00	0.78	0.57	0.65	0.74	4.00	0.65	0.74	0.52
time (sec)	N/A	0.124	0.010	0.114	0.025	0.080	0.544	0.123	0.148	0.017

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	10	8	6	6	6
N.S.	1	1.00	1.00	0.88	0.75	1.25	1.00	0.75	0.75	0.75
time (sec)	N/A	0.153	0.004	0.085	0.028	0.080	0.042	0.118	0.141	0.235

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	13	12	18	29	12	17	12
N.S.	1	1.00	1.00	0.81	0.75	1.12	1.81	0.75	1.06	0.75
time (sec)	N/A	0.168	0.010	0.195	0.030	0.083	0.099	0.118	0.153	0.098

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	6	6	6	7	6	6	5	6	12	6
N.S.	1	1.00	1.00	1.17	1.00	1.00	0.83	1.00	2.00	1.00
time (sec)	N/A	0.161	0.011	0.090	0.039	0.071	0.167	0.122	0.155	0.025

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	12	11	10	12	12	6	9	9
N.S.	1	1.00	1.00	0.92	0.83	1.00	1.00	0.50	0.75	0.75
time (sec)	N/A	0.205	0.005	0.125	0.030	0.081	0.236	0.115	0.147	0.115

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	10	9	8	8	10	8	7	8
N.S.	1	1.00	1.00	0.90	0.80	0.80	1.00	0.80	0.70	0.80
time (sec)	N/A	0.181	0.020	0.073	0.026	0.085	0.106	0.126	0.155	0.096

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	9	9	9	10	9	9	7	9	9	9
N.S.	1	1.00	1.00	1.11	1.00	1.00	0.78	1.00	1.00	1.00
time (sec)	N/A	0.166	0.012	0.197	0.026	0.090	1.206	0.119	0.154	0.115

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	12	11	11	10	11	10	11
N.S.	1	1.00	1.00	0.80	0.73	0.73	0.67	0.73	0.67	0.73
time (sec)	N/A	0.118	0.008	0.128	0.026	0.071	0.090	0.288	0.153	0.215

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	18	13	15	15	34	15	15	12
N.S.	1	1.00	0.78	0.57	0.65	0.65	1.48	0.65	0.65	0.52
time (sec)	N/A	0.126	0.009	0.085	0.035	0.074	0.538	0.117	0.152	0.018

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	11	10	9	22	8	9	21	9
N.S.	1	1.00	1.00	0.91	0.82	2.00	0.73	0.82	1.91	0.82
time (sec)	N/A	0.113	0.016	0.103	0.025	0.091	0.349	0.124	0.150	0.042

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	12	11	11	27	11	11	11
N.S.	1	1.00	1.00	0.80	0.73	0.73	1.80	0.73	0.73	0.73
time (sec)	N/A	0.121	0.009	0.082	0.023	0.079	0.092	0.120	0.151	0.104

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	12	11	11	12	11	29	15
N.S.	1	1.00	1.00	0.80	0.73	0.73	0.80	0.73	1.93	1.00
time (sec)	N/A	0.169	0.042	0.198	0.023	0.086	0.114	0.241	0.168	0.135

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	A	B	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	32	24	32	0	0	23	0	15	42	47
N.S.	1	0.75	1.00	0.00	0.00	0.72	0.00	0.47	1.31	1.47
time (sec)	N/A	0.277	1.617	0.000	0.000	0.133	0.000	0.129	0.156	0.326

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	13	13	14	13	42	13
N.S.	1	1.00	1.00	0.82	0.76	0.76	0.82	0.76	2.47	0.76
time (sec)	N/A	0.241	0.023	0.056	0.033	0.099	0.119	0.119	0.153	0.125

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	13	10	7	12	15	12	12	9
N.S.	1	1.00	0.81	0.62	0.44	0.75	0.94	0.75	0.75	0.56
time (sec)	N/A	0.122	0.003	0.110	0.024	0.088	0.499	0.117	0.155	0.086

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	9	8	8	7	8	8	8
N.S.	1	1.00	1.00	1.12	1.00	1.00	0.88	1.00	1.00	1.00
time (sec)	N/A	0.168	0.007	0.111	0.030	0.113	0.062	0.257	0.156	0.035

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	15	17	15	15	17	15	17	15
N.S.	1	1.00	0.88	1.00	0.88	0.88	1.00	0.88	1.00	0.88
time (sec)	N/A	0.225	0.010	0.105	0.029	0.096	0.086	0.348	0.159	0.017

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	26	19	20	20	20	26	20	23	24
N.S.	1	1.13	0.83	0.87	0.87	0.87	1.13	0.87	1.00	1.04
time (sec)	N/A	0.290	0.011	0.135	0.036	0.091	0.113	0.602	0.150	0.016

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	27	20	23	21	21	26	21	24	23
N.S.	1	1.12	0.83	0.96	0.88	0.88	1.08	0.88	1.00	0.96
time (sec)	N/A	0.287	0.010	0.104	0.033	0.091	0.121	0.115	0.154	0.017

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	6	5	6	6	6
N.S.	1	1.00	1.00	0.88	0.75	0.75	0.62	0.75	0.75	0.75
time (sec)	N/A	0.137	0.001	0.082	0.026	0.078	0.035	0.123	0.151	0.011

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	28	18	15	14	17	24	14	21	18
N.S.	1	1.22	0.78	0.65	0.61	0.74	1.04	0.61	0.91	0.78
time (sec)	N/A	0.174	0.002	0.125	0.038	0.091	0.090	0.116	0.152	0.057

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	11	10	10	10	10	10	10
N.S.	1	1.00	1.00	0.79	0.71	0.71	0.71	0.71	0.71	0.71
time (sec)	N/A	0.137	0.006	0.105	0.025	0.098	0.020	0.121	0.155	0.021

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	15	11	11	11	8	11	14	10
N.S.	1	1.00	1.15	0.85	0.85	0.85	0.62	0.85	1.08	0.77
time (sec)	N/A	0.151	0.001	0.000	0.034	0.103	0.024	0.122	0.155	0.001

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	29	22	17	16	19	24	16	18	16
N.S.	1	1.21	0.92	0.71	0.67	0.79	1.00	0.67	0.75	0.67
time (sec)	N/A	0.181	0.006	0.403	0.036	0.099	0.020	0.136	0.157	0.020

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	23	17	17	17	17	17	22	17
N.S.	1	1.00	1.10	0.81	0.81	0.81	0.81	0.81	1.05	0.81
time (sec)	N/A	0.157	0.006	0.506	0.025	0.105	0.025	0.123	0.156	0.024

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	44	30	23	24	25	36	22	26	22
N.S.	1	1.29	0.88	0.68	0.71	0.74	1.06	0.65	0.76	0.65
time (sec)	N/A	0.228	0.006	0.784	0.031	0.087	0.023	0.120	0.158	0.022

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	25	20	19	19	36	19	20	19
N.S.	1	1.00	1.00	0.80	0.76	0.76	1.44	0.76	0.80	0.76
time (sec)	N/A	0.157	0.009	0.136	0.032	0.092	0.084	0.113	0.152	0.055

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	32	31	23	23	23	39	23	25	25
N.S.	1	0.97	0.94	0.70	0.70	0.70	1.18	0.70	0.76	0.76
time (sec)	N/A	0.238	0.006	0.273	0.028	0.094	0.132	0.114	0.153	0.066

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	46	29	19	26	29	56	26	31	28
N.S.	1	1.12	0.71	0.46	0.63	0.71	1.37	0.63	0.76	0.68
time (sec)	N/A	0.211	0.024	0.175	0.026	0.120	0.120	0.116	0.156	0.036

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	11	10	10	10	10	10	10
N.S.	1	1.00	1.00	0.79	0.71	0.71	0.71	0.71	0.71	0.71
time (sec)	N/A	0.145	0.005	0.088	0.034	0.095	0.018	0.119	0.152	0.015

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	11	11	9	10	8	9	12	9
N.S.	1	1.00	1.00	1.00	0.82	0.91	0.73	0.82	1.09	0.82
time (sec)	N/A	0.157	0.003	0.211	0.048	0.093	0.021	0.127	0.157	0.019

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	29	22	17	16	19	24	16	18	16
N.S.	1	1.21	0.92	0.71	0.67	0.79	1.00	0.67	0.75	0.67
time (sec)	N/A	0.186	0.005	0.306	0.025	0.090	0.018	0.118	0.145	0.016

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	94	61	54	60	60	180	50	61	37
N.S.	1	1.12	0.73	0.64	0.71	0.71	2.14	0.60	0.73	0.44
time (sec)	N/A	0.160	0.124	0.200	0.106	0.108	2.256	0.133	0.153	0.119

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	42	25	21	34	21	39	28	20	20
N.S.	1	1.11	0.66	0.55	0.89	0.55	1.03	0.74	0.53	0.53
time (sec)	N/A	0.144	0.012	0.076	0.113	0.087	0.193	0.118	0.151	0.016

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	172	172	34	17	0	17	31	0	28	301
N.S.	1	1.00	0.20	0.10	0.00	0.10	0.18	0.00	0.16	1.75
time (sec)	N/A	0.191	9.728	0.142	0.000	0.106	0.385	0.000	0.151	0.042

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	6	6	8	7	6	6	7	6	6	6
N.S.	1	1.00	1.33	1.17	1.00	1.00	1.17	1.00	1.00	1.00
time (sec)	N/A	0.140	0.004	0.020	0.103	0.094	0.026	0.123	0.151	0.039

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	16	13	12	12	19	12	12	12
N.S.	1	1.00	1.14	0.93	0.86	0.86	1.36	0.86	0.86	0.86
time (sec)	N/A	0.184	0.004	0.030	0.105	0.085	0.028	0.123	0.147	0.017

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	18	9	10	20	8	18	8	8
N.S.	1	1.00	2.25	1.12	1.25	2.50	1.00	2.25	1.00	1.00
time (sec)	N/A	0.145	0.008	0.044	0.104	0.090	0.022	0.123	0.155	0.034

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	22	11	16	48	19	34	10	10
N.S.	1	1.00	1.83	0.92	1.33	4.00	1.58	2.83	0.83	0.83
time (sec)	N/A	0.186	0.005	0.044	0.102	0.104	0.024	0.115	0.152	0.012

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	26	18	20	18	26	18	20	20
N.S.	1	1.00	1.18	0.82	0.91	0.82	1.18	0.82	0.91	0.91
time (sec)	N/A	0.189	0.052	0.170	0.025	0.098	0.071	0.121	0.153	0.058

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	13	10	9	9	22	9	13	9
N.S.	1	1.00	1.00	0.77	0.69	0.69	1.69	0.69	1.00	0.69
time (sec)	N/A	0.114	0.001	0.062	0.029	0.076	0.071	0.115	0.155	0.014

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	11	10	9	51	58	9	50	9
N.S.	1	1.00	0.85	0.77	0.69	3.92	4.46	0.69	3.85	0.69
time (sec)	N/A	0.119	0.001	0.077	0.023	0.077	0.019	0.118	0.156	0.053

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	16	14	19	19	15	14	20	14
N.S.	1	1.00	0.89	0.78	1.06	1.06	0.83	0.78	1.11	0.78
time (sec)	N/A	0.120	0.003	0.066	0.024	0.078	0.042	0.124	0.150	0.022

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	43	10	9	31	31	9	30	31
N.S.	1	1.00	3.91	0.91	0.82	2.82	2.82	0.82	2.73	2.82
time (sec)	N/A	0.118	0.001	0.052	0.030	0.086	0.018	0.120	0.148	0.016

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	140	106	106	106	131	106	105	106
N.S.	1	1.00	2.50	1.89	1.89	1.89	2.34	1.89	1.88	1.89
time (sec)	N/A	0.167	0.001	0.068	0.027	0.082	0.028	0.128	0.154	0.290

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	4	4	4	5	4	4	3	4	4	4
N.S.	1	1.00	1.00	1.25	1.00	1.00	0.75	1.00	1.00	1.00
time (sec)	N/A	0.163	0.008	0.108	0.025	0.074	0.145	0.118	0.148	0.055

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	62	67	46	49	37	37	60	37	47	41
N.S.	1	1.08	0.74	0.79	0.60	0.60	0.97	0.60	0.76	0.66
time (sec)	N/A	0.333	0.023	0.033	0.049	0.098	0.276	0.124	0.156	0.129

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	10	9	8	8	7	8	8	8
N.S.	1	1.00	1.00	0.90	0.80	0.80	0.70	0.80	0.80	0.80
time (sec)	N/A	0.135	0.002	0.177	0.023	0.090	0.084	0.124	0.156	0.029

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	13	12	12	15	12	16	12
N.S.	1	1.00	1.00	0.81	0.75	0.75	0.94	0.75	1.00	0.75
time (sec)	N/A	0.178	0.009	0.139	0.030	0.097	0.671	0.127	0.157	0.110

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	10	7	8	8	7	9	8	6
N.S.	1	1.00	1.00	0.70	0.80	0.80	0.70	0.90	0.80	0.60
time (sec)	N/A	0.110	0.000	0.056	0.026	0.069	0.019	0.120	0.155	0.040

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	17	15	16	12	15	15	15	12	12
N.S.	1	1.13	1.00	1.07	0.80	1.00	1.00	1.00	0.80	0.80
time (sec)	N/A	0.143	0.002	0.036	0.022	0.085	0.043	0.117	0.154	0.047

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	13	13	12	13	11	9
N.S.	1	1.00	1.00	0.82	0.76	0.76	0.71	0.76	0.65	0.53
time (sec)	N/A	0.128	0.002	0.017	0.028	0.079	0.038	0.117	0.156	0.018

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	28	23	17	22	22	22	17	17
N.S.	1	1.00	1.00	0.82	0.61	0.79	0.79	0.79	0.61	0.61
time (sec)	N/A	0.155	0.002	0.044	0.024	0.078	0.046	0.117	0.150	0.018

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	4	4	4	5	4	4	3	5	4	4
N.S.	1	1.00	1.00	1.25	1.00	1.00	0.75	1.25	1.00	1.00
time (sec)	N/A	0.108	0.000	0.045	0.027	0.071	0.020	0.119	0.152	0.009

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	3	3	3	4	3	11	3	4	17	3
N.S.	1	1.00	1.00	1.33	1.00	3.67	1.00	1.33	5.67	1.00
time (sec)	N/A	0.137	0.002	0.038	0.028	0.093	0.045	0.123	0.157	0.013

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	21	33	28	32	61	28	27	38
N.S.	1	1.00	0.75	1.18	1.00	1.14	2.18	1.00	0.96	1.36
time (sec)	N/A	0.141	0.005	0.069	0.022	0.096	0.293	0.118	0.153	0.121

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	33	28	23	17	22	26	22	17	17
N.S.	1	1.18	1.00	0.82	0.61	0.79	0.93	0.79	0.61	0.61
time (sec)	N/A	0.162	0.002	0.050	0.025	0.094	0.044	0.117	0.157	0.022

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	3	3	3	4	3	3	3	4	3	3
N.S.	1	1.00	1.00	1.33	1.00	1.00	1.00	1.33	1.00	1.00
time (sec)	N/A	0.134	0.004	0.016	0.034	0.091	0.039	0.120	0.153	0.042

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	12	11	10	10	8	10	10	10
N.S.	1	1.00	1.00	0.92	0.83	0.83	0.67	0.83	0.83	0.83
time (sec)	N/A	0.156	0.003	0.098	0.024	0.082	0.041	0.115	0.153	0.198

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	16	18	17	12	20	17	11	13
N.S.	1	1.00	0.70	0.78	0.74	0.52	0.87	0.74	0.48	0.57
time (sec)	N/A	0.172	0.019	0.038	0.027	0.072	0.868	0.115	0.155	0.096

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	49	39	32	23	31	37	31	23	23
N.S.	1	1.26	1.00	0.82	0.59	0.79	0.95	0.79	0.59	0.59
time (sec)	N/A	0.193	0.002	0.096	0.027	0.088	0.056	0.124	0.157	0.018

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	9	9	9	7	6	6	5	6	7	6
N.S.	1	1.00	1.00	0.78	0.67	0.67	0.56	0.67	0.78	0.67
time (sec)	N/A	0.131	0.010	0.029	0.025	0.078	0.033	0.129	0.154	0.044

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	12	12	11	10	11	10	11
N.S.	1	1.00	1.00	0.86	0.86	0.79	0.71	0.79	0.71	0.79
time (sec)	N/A	0.132	0.003	0.046	0.026	0.088	0.070	0.120	0.152	0.061

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	10	8	7	7	7	7	8	7
N.S.	1	1.00	1.00	0.80	0.70	0.70	0.70	0.70	0.80	0.70
time (sec)	N/A	0.150	0.006	0.104	0.030	0.085	0.109	0.119	0.155	0.063

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	14	12	11	13	15	11	12	11
N.S.	1	1.00	0.74	0.63	0.58	0.68	0.79	0.58	0.63	0.58
time (sec)	N/A	0.141	0.010	0.083	0.028	0.089	0.091	0.116	0.151	0.011

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	12	10	9	13	15	9	10	9
N.S.	1	1.00	0.63	0.53	0.47	0.68	0.79	0.47	0.53	0.47
time (sec)	N/A	0.143	0.007	0.079	0.025	0.089	0.086	0.118	0.153	0.010

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	13	10	10	9	9	7	9	10	9
N.S.	1	1.30	1.00	1.00	0.90	0.90	0.70	0.90	1.00	0.90
time (sec)	N/A	0.132	0.008	0.018	0.024	0.088	0.034	0.115	0.153	0.026

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	7	7	6	6	5	6	7	6
N.S.	1	1.00	0.64	0.64	0.55	0.55	0.45	0.55	0.64	0.55
time (sec)	N/A	0.141	0.007	0.017	0.023	0.082	0.033	0.127	0.157	0.010

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	11	10	9	9	7	9	11	9
N.S.	1	1.00	0.69	0.62	0.56	0.56	0.44	0.56	0.69	0.56
time (sec)	N/A	0.144	0.008	0.023	0.024	0.086	0.030	0.122	0.154	0.011

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	21	12	12	11	11	10	11	12	11
N.S.	1	1.11	0.63	0.63	0.58	0.58	0.53	0.58	0.63	0.58
time (sec)	N/A	0.170	0.009	0.020	0.029	0.094	0.036	0.119	0.158	0.012

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	19	16	16	16	17	16	19	16
N.S.	1	1.00	0.59	0.50	0.50	0.50	0.53	0.50	0.59	0.50
time (sec)	N/A	0.175	0.012	0.024	0.024	0.082	0.036	0.119	0.150	0.043

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	25	16	16	11	11	20	11	10	11
N.S.	1	1.04	0.67	0.67	0.46	0.46	0.83	0.46	0.42	0.46
time (sec)	N/A	0.159	0.005	0.011	0.026	0.083	0.087	0.117	0.151	0.013

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	16	14	13	13	12	13	16	13
N.S.	1	1.00	0.62	0.54	0.50	0.50	0.46	0.50	0.62	0.50
time (sec)	N/A	0.164	0.016	0.026	0.034	0.098	0.034	0.329	0.157	0.064

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	28	28	27	31	136	36	28	27
N.S.	1	1.00	0.68	0.68	0.66	0.76	3.32	0.88	0.68	0.66
time (sec)	N/A	0.158	0.023	0.127	0.031	0.097	0.248	1.252	0.153	0.016

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	42	29	29	29	33	139	38	29	29
N.S.	1	1.00	0.69	0.69	0.69	0.79	3.31	0.90	0.69	0.69
time (sec)	N/A	0.158	0.022	0.096	0.028	0.093	0.252	0.122	0.149	0.011

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	13	12	21	13	13
N.S.	1	1.00	1.00	0.93	0.87	0.87	0.80	1.40	0.87	0.87
time (sec)	N/A	0.139	0.001	0.062	0.024	0.088	0.067	0.124	0.155	0.060

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	B	A	C	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	64	22	35	33	17	37	4	21
N.S.	1	1.00	3.37	1.16	1.84	1.74	0.89	1.95	0.21	1.11
time (sec)	N/A	0.151	0.042	0.024	0.031	0.104	0.965	0.129	0.155	0.334

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	B	B	A	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	64	20	35	35	17	37	4	20
N.S.	1	1.00	3.76	1.18	2.06	2.06	1.00	2.18	0.24	1.18
time (sec)	N/A	0.156	0.024	0.009	0.024	0.107	0.951	0.128	0.156	0.117

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	26	25	24	23	23	22	23	22	22
N.S.	1	1.04	1.00	0.96	0.92	0.92	0.88	0.92	0.88	0.88
time (sec)	N/A	0.195	0.004	0.076	0.103	0.090	0.070	0.154	0.150	0.019

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	22	21	33	39	22	38	17	20
N.S.	1	1.00	1.00	0.95	1.50	1.77	1.00	1.73	0.77	0.91
time (sec)	N/A	0.157	0.003	0.038	0.112	0.115	0.915	0.125	0.158	0.012

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	15	6	23	19	28	6	14
N.S.	1	1.00	1.00	0.94	0.38	1.44	1.19	1.75	0.38	0.88
time (sec)	N/A	0.120	0.003	0.083	0.116	0.088	0.490	0.126	0.152	0.088

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	17	23	10	11	21	10	27	9	11
N.S.	1	1.70	2.30	1.00	1.10	2.10	1.00	2.70	0.90	1.10
time (sec)	N/A	0.128	0.045	0.153	0.104	0.086	0.283	0.126	0.158	0.055

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	10	11	10	10	20	10	10	10
N.S.	1	1.00	1.00	1.10	1.00	1.00	2.00	1.00	1.00	1.00
time (sec)	N/A	0.114	0.002	0.082	0.117	0.076	0.048	0.121	0.155	0.021

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	24	16	15	67	53	15	23	16
N.S.	1	1.00	1.00	0.67	0.62	2.79	2.21	0.62	0.96	0.67
time (sec)	N/A	0.120	0.004	0.074	0.104	0.085	0.058	0.116	0.155	0.057

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	19	17	16	16	26	16	15	16
N.S.	1	1.00	1.00	0.89	0.84	0.84	1.37	0.84	0.79	0.84
time (sec)	N/A	0.127	0.006	0.248	0.110	0.082	0.046	0.116	0.148	0.044

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	20	16	16	15	13	15	15	15	14
N.S.	1	0.95	0.76	0.76	0.71	0.62	0.71	0.71	0.71	0.67
time (sec)	N/A	0.145	0.003	0.056	0.110	0.084	0.079	0.121	0.156	0.014

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	43	30	34	33	24	32	33	31	24
N.S.	1	1.08	0.75	0.85	0.82	0.60	0.80	0.82	0.78	0.60
time (sec)	N/A	0.175	0.009	0.046	0.101	0.089	0.099	0.120	0.156	0.016

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	26	30	34	25	29	29	29	29
N.S.	1	1.00	0.74	0.86	0.97	0.71	0.83	0.83	0.83	0.83
time (sec)	N/A	0.296	0.007	0.084	0.103	0.088	0.095	0.113	0.153	0.071

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	29	18	17	16	14	19	16	13	16
N.S.	1	1.32	0.82	0.77	0.73	0.64	0.86	0.73	0.59	0.73
time (sec)	N/A	0.149	0.011	0.053	0.111	0.097	0.367	0.128	0.159	0.031

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	6	7	6	5	6
N.S.	1	1.00	1.00	0.88	0.75	0.75	0.88	0.75	0.62	0.75
time (sec)	N/A	0.170	0.004	0.063	0.025	0.097	0.266	0.128	0.149	0.374

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	37	18	17	31	15	17	16	17
N.S.	1	1.00	1.61	0.78	0.74	1.35	0.65	0.74	0.70	0.74
time (sec)	N/A	0.123	0.033	0.191	0.103	0.072	0.081	0.123	0.151	0.051

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	A	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	37	16	0	15	31	24	23	0
N.S.	1	1.00	1.68	0.73	0.00	0.68	1.41	1.09	1.05	0.00
time (sec)	N/A	0.178	0.008	0.072	0.000	0.085	8.336	0.127	0.159	0.000

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	16	0	15	31	22	23	0
N.S.	1	1.00	1.00	0.80	0.00	0.75	1.55	1.10	1.15	0.00
time (sec)	N/A	0.160	0.005	0.072	0.000	0.098	8.020	0.118	0.159	0.000

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	19	16	15	21	12	15	22	17
N.S.	1	1.00	1.00	0.84	0.79	1.11	0.63	0.79	1.16	0.89
time (sec)	N/A	0.121	0.005	0.057	0.103	0.074	0.037	0.123	0.154	0.018

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	4	4	4	4	3	3	15	3	4	3
N.S.	1	1.00	1.00	1.00	0.75	0.75	3.75	0.75	1.00	0.75
time (sec)	N/A	0.136	0.013	0.021	0.102	0.077	0.045	0.122	0.148	0.059

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	33	27	25	19	28	19	21	32	22
N.S.	1	1.22	1.00	0.93	0.70	1.04	0.70	0.78	1.19	0.81
time (sec)	N/A	0.186	0.012	0.059	0.023	0.097	1.084	0.119	0.150	0.081

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	58	67	61	49	38	0	36	32	49
N.S.	1	1.38	1.60	1.45	1.17	0.90	0.00	0.86	0.76	1.17
time (sec)	N/A	0.146	0.057	0.075	0.103	0.075	0.000	0.136	0.150	0.039

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	71	71	75	68	0	80	114	61	182	0
N.S.	1	1.00	1.06	0.96	0.00	1.13	1.61	0.86	2.56	0.00
time (sec)	N/A	0.177	0.108	0.138	0.000	0.081	1.648	0.123	0.150	0.000

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	C	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	55	28	0	43	71	61	20	0
N.S.	1	1.00	1.72	0.88	0.00	1.34	2.22	1.91	0.62	0.00
time (sec)	N/A	0.150	0.029	0.119	0.000	0.078	0.835	0.128	0.153	0.000

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	13	12	15	13	13
N.S.	1	1.00	1.00	0.93	0.87	0.87	0.80	1.00	0.87	0.87
time (sec)	N/A	0.139	0.003	0.078	0.023	0.074	0.042	0.118	0.152	0.030

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	19	14	13	13	14	15	13	13
N.S.	1	1.00	1.00	0.74	0.68	0.68	0.74	0.79	0.68	0.68
time (sec)	N/A	0.138	0.003	0.083	0.024	0.087	0.044	0.114	0.151	0.021

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	23	20	19	19	19	21	19	19
N.S.	1	1.00	1.00	0.87	0.83	0.83	0.83	0.91	0.83	0.83
time (sec)	N/A	0.185	0.003	0.089	0.028	0.072	0.056	0.125	0.150	0.023

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	23	18	17	17	17	20	17	19
N.S.	1	1.00	1.00	0.78	0.74	0.74	0.74	0.87	0.74	0.83
time (sec)	N/A	0.190	0.004	0.032	0.026	0.081	0.057	0.124	0.150	0.109

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	22	19	18	26	19	24	38	18
N.S.	1	1.00	0.92	0.79	0.75	1.08	0.79	1.00	1.58	0.75
time (sec)	N/A	0.147	0.007	0.075	0.029	0.073	0.048	0.114	0.153	0.027

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	29	28	29	28	28	3	29	27	57
N.S.	1	1.04	1.00	1.04	1.00	1.00	0.11	1.04	0.96	2.04
time (sec)	N/A	0.182	0.015	0.081	0.101	0.077	0.055	0.109	0.154	0.102

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	61	37	36	51	14	37	75	53
N.S.	1	1.00	1.24	0.76	0.73	1.04	0.29	0.76	1.53	1.08
time (sec)	N/A	0.230	0.021	0.086	0.104	0.078	0.070	0.126	0.149	0.055

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	B	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	24	19	39	38	39	37	27	21
N.S.	1	1.00	1.14	0.90	1.86	1.81	1.86	1.76	1.29	1.00
time (sec)	N/A	0.155	0.017	0.091	0.104	0.082	0.218	0.141	0.162	0.186

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	21	14	55	48	14	13	87
N.S.	1	1.00	1.00	1.31	0.88	3.44	3.00	0.88	0.81	5.44
time (sec)	N/A	0.206	0.012	0.102	0.031	0.070	1.332	0.125	0.159	0.094

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	9	9	9	9	8	11	9	9
N.S.	1	1.00	0.82	0.82	0.82	0.82	0.73	1.00	0.82	0.82
time (sec)	N/A	0.129	0.003	0.069	0.024	0.062	0.037	0.121	0.153	0.027

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	23	20	19	19	20	22	19	19
N.S.	1	1.00	1.00	0.87	0.83	0.83	0.87	0.96	0.83	0.83
time (sec)	N/A	0.138	0.004	0.073	0.025	0.068	0.057	0.122	0.149	0.064

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	30	28	21	20	27	22	22	38	18
N.S.	1	1.00	0.93	0.70	0.67	0.90	0.73	0.73	1.27	0.60
time (sec)	N/A	0.158	0.006	0.026	0.023	0.062	0.041	0.116	0.140	0.025

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	27	24	23	23	20	26	23	30
N.S.	1	1.00	1.00	0.89	0.85	0.85	0.74	0.96	0.85	1.11
time (sec)	N/A	0.187	0.004	0.032	0.024	0.068	0.055	0.116	0.145	0.059

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	24	16	20	32	17	16	42	15
N.S.	1	1.00	1.04	0.70	0.87	1.39	0.74	0.70	1.83	0.65
time (sec)	N/A	0.170	0.007	0.076	0.029	0.061	0.048	0.115	0.149	0.052

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	15	14	14	14	15	14	14
N.S.	1	1.00	1.00	0.94	0.88	0.88	0.88	0.94	0.88	0.88
time (sec)	N/A	0.153	0.003	0.090	0.102	0.068	0.044	0.120	0.144	0.022

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	18	13	12	12	14	12	12	12
N.S.	1	1.00	1.00	0.72	0.67	0.67	0.78	0.67	0.67	0.67
time (sec)	N/A	0.142	0.005	0.047	0.109	0.076	0.065	0.120	0.154	0.021

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	11	12	11	11	8	13	11	11
N.S.	1	1.00	1.00	1.09	1.00	1.00	0.73	1.18	1.00	1.00
time (sec)	N/A	0.136	0.002	0.066	0.032	0.064	0.044	0.113	0.149	0.057

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	24	21	24	32	19	29	46	20
N.S.	1	1.00	1.00	0.88	1.00	1.33	0.79	1.21	1.92	0.83
time (sec)	N/A	0.138	0.007	0.065	0.023	0.079	0.044	0.125	0.151	0.019

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	44	39	46	83	46	52	121	45
N.S.	1	1.00	0.96	0.85	1.00	1.80	1.00	1.13	2.63	0.98
time (sec)	N/A	0.152	0.011	0.080	0.025	0.074	0.087	0.122	0.152	0.050

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	10	11	10	16	8	11	20	10
N.S.	1	1.00	1.00	1.10	1.00	1.60	0.80	1.10	2.00	1.00
time (sec)	N/A	0.127	0.003	0.066	0.030	0.064	0.029	0.122	0.147	0.021

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	19	17	14	17	13	10	16	17	13
N.S.	1	1.12	1.00	0.82	1.00	0.76	0.59	0.94	1.00	0.76
time (sec)	N/A	0.135	0.003	0.082	0.028	0.060	0.040	0.117	0.150	0.067

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	15	14	14	17	16	14	14
N.S.	1	1.00	1.00	0.75	0.70	0.70	0.85	0.80	0.70	0.70
time (sec)	N/A	0.140	0.003	0.085	0.031	0.064	0.046	0.117	0.153	0.027

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	12	13	12	16	8	13	22	12
N.S.	1	1.00	0.75	0.81	0.75	1.00	0.50	0.81	1.38	0.75
time (sec)	N/A	0.137	0.003	0.083	0.022	0.065	0.036	0.123	0.154	0.020

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	20	14	15	14	17	10	14	25	14
N.S.	1	1.43	1.00	1.07	1.00	1.21	0.71	1.00	1.79	1.00
time (sec)	N/A	0.157	0.006	0.246	0.103	0.065	0.061	0.121	0.153	0.048

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	21	18	17	17	20	20	17	17
N.S.	1	1.00	1.00	0.86	0.81	0.81	0.95	0.95	0.81	0.81
time (sec)	N/A	0.155	0.004	0.029	0.025	0.075	0.065	0.120	0.153	0.036

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	27	23	23	34	20	25	41	17
N.S.	1	1.00	1.29	1.10	1.10	1.62	0.95	1.19	1.95	0.81
time (sec)	N/A	0.123	0.005	0.070	0.025	0.083	0.045	0.120	0.152	0.050

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	22	17	16	16	17	17	16	16
N.S.	1	1.00	1.00	0.77	0.73	0.73	0.77	0.77	0.73	0.73
time (sec)	N/A	0.151	0.003	0.074	0.101	0.067	0.037	0.120	0.157	0.093

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	18	10	11	14	18	8	14	22	10
N.S.	1	1.80	1.00	1.10	1.40	1.80	0.80	1.40	2.20	1.00
time (sec)	N/A	0.147	0.004	0.073	0.022	0.067	0.037	0.117	0.151	0.016

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	31	21	19	25	19	21	25	16
N.S.	1	1.00	1.00	0.68	0.61	0.81	0.61	0.68	0.81	0.52
time (sec)	N/A	0.149	0.002	0.073	0.030	0.089	0.045	0.117	0.154	0.022

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	18	17	16	16	15	17	16	24
N.S.	1	1.00	1.00	0.94	0.89	0.89	0.83	0.94	0.89	1.33
time (sec)	N/A	0.167	0.005	0.093	0.104	0.074	0.059	0.120	0.156	0.023

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	25	10	17	17	17	19	17	9
N.S.	1	1.00	1.92	0.77	1.31	1.31	1.31	1.46	1.31	0.69
time (sec)	N/A	0.121	0.003	0.077	0.100	0.075	0.056	0.121	0.157	0.016

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	90	64	22	72	60	73	72	56	33
N.S.	1	1.06	0.75	0.26	0.85	0.71	0.86	0.85	0.66	0.39
time (sec)	N/A	0.231	0.013	0.081	0.102	0.074	0.071	0.118	0.145	0.064

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	15	16	15	26	14	15	34	15
N.S.	1	1.00	0.65	0.70	0.65	1.13	0.61	0.65	1.48	0.65
time (sec)	N/A	0.145	0.006	0.123	0.102	0.077	0.042	0.122	0.144	0.048

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	11	12	11	11	8	11	14	11
N.S.	1	1.00	1.00	1.09	1.00	1.00	0.73	1.00	1.27	1.00
time (sec)	N/A	0.129	0.005	0.049	0.022	0.066	0.043	0.124	0.158	0.034

Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	26	23	20	23	36	39	47	18	21
N.S.	1	0.58	0.51	0.44	0.51	0.80	0.87	1.04	0.40	0.47
time (sec)	N/A	0.179	0.025	0.132	0.110	0.086	0.209	0.123	0.146	0.060

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	31	30	0	111	110	53	42	28
N.S.	1	1.00	0.84	0.81	0.00	3.00	2.97	1.43	1.14	0.76
time (sec)	N/A	0.168	0.022	0.079	0.000	0.090	1.327	0.127	0.167	0.178

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	20	20	16	37	50	36	35	25	15
N.S.	1	0.36	0.36	0.29	0.66	0.89	0.64	0.62	0.45	0.27
time (sec)	N/A	0.157	0.014	0.059	0.106	0.085	0.145	0.146	0.153	0.138

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	20	16	19	23	32	40	14	32
N.S.	1	1.00	0.65	0.52	0.61	0.74	1.03	1.29	0.45	1.03
time (sec)	N/A	0.158	0.012	0.060	0.106	0.075	0.123	0.122	0.148	0.128

Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	18	18	17	14	33	248	48	53	26
N.S.	1	0.50	0.50	0.47	0.39	0.92	6.89	1.33	1.47	0.72
time (sec)	N/A	0.212	0.100	0.146	0.105	0.091	21.348	0.123	0.154	0.118

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	16	15	43	71839	26	21	15
N.S.	1	1.00	1.00	1.07	1.00	2.87	4789.27	1.73	1.40	1.00
time (sec)	N/A	0.179	0.040	0.673	0.106	0.087	15.299	0.122	0.145	0.252

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	14	39	602	13	18	29
N.S.	1	1.00	1.00	0.82	0.82	2.29	35.41	0.76	1.06	1.71
time (sec)	N/A	0.158	0.026	0.250	0.031	0.080	103.861	0.116	0.149	0.356

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	30	22	20	41	11	22	25	22	34
N.S.	1	1.00	0.73	0.67	1.37	0.37	0.73	0.83	0.73	1.13
time (sec)	N/A	0.215	0.050	0.179	0.104	0.082	0.131	0.127	0.145	0.175

Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	41	23	22	34	24	22	21	22
N.S.	1	1.00	1.41	0.79	0.76	1.17	0.83	0.76	0.72	0.76
time (sec)	N/A	0.123	0.045	0.247	0.106	0.077	0.075	0.122	0.150	0.020

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	13	12	11	11	8	11	10	11
N.S.	1	1.00	1.00	0.92	0.85	0.85	0.62	0.85	0.77	0.85
time (sec)	N/A	0.121	0.001	0.073	0.023	0.074	0.060	0.116	0.142	0.093

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	43	33	30	41	40	87	47	27	35
N.S.	1	1.16	0.89	0.81	1.11	1.08	2.35	1.27	0.73	0.95
time (sec)	N/A	0.140	0.031	0.127	0.101	0.069	0.742	0.117	0.167	0.116

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	42	22	25	25	0	26	16	21
N.S.	1	1.00	1.91	1.00	1.14	1.14	0.00	1.18	0.73	0.95
time (sec)	N/A	0.137	0.032	0.132	0.024	0.070	0.000	0.126	0.144	0.046

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	33	21	20	25	24	25	26	20
N.S.	1	1.00	1.22	0.78	0.74	0.93	0.89	0.93	0.96	0.74
time (sec)	N/A	0.123	0.018	0.110	0.106	0.065	0.071	0.120	0.151	0.050

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	33	21	22	27	26	27	30	23
N.S.	1	1.00	1.22	0.78	0.81	1.00	0.96	1.00	1.11	0.85
time (sec)	N/A	0.142	0.042	0.271	0.104	0.065	0.230	0.130	0.146	0.029

Problem 154	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	39	7	15	17	15	33	10	11
N.S.	1	1.00	2.79	0.50	1.07	1.21	1.07	2.36	0.71	0.79
time (sec)	N/A	0.129	0.021	0.069	0.023	0.067	0.184	0.119	0.147	0.095

Problem 155	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	72	76	60	59	92	0	168	69	73
N.S.	1	1.06	1.12	0.88	0.87	1.35	0.00	2.47	1.01	1.07
time (sec)	N/A	0.205	0.104	0.217	0.105	0.076	0.000	0.148	0.157	0.046

Problem 156	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	F	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	13	13	12	0	73	0	21	13
N.S.	1	1.00	1.00	1.00	0.92	0.00	5.62	0.00	1.62	1.00
time (sec)	N/A	0.155	0.003	0.096	0.025	0.000	0.898	0.000	0.152	0.016

Problem 157	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	15	2	2	15	2	2	2
N.S.	1	1.00	1.00	1.00	0.13	0.13	1.00	0.13	0.13	0.13
time (sec)	N/A	0.198	0.010	0.139	0.025	0.077	0.072	0.119	0.154	0.060

Problem 158	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2	2	2	8	2	2	2	2	2	2
N.S.	1	1.00	1.00	4.00	1.00	1.00	1.00	1.00	1.00	1.00
time (sec)	N/A	0.123	0.009	0.049	0.053	0.065	0.325	0.122	0.153	0.004

Problem 159	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	4	4	4	9	4	4	3	4	4	4
N.S.	1	1.00	1.00	2.25	1.00	1.00	0.75	1.00	1.00	1.00
time (sec)	N/A	0.128	0.010	0.033	0.052	0.071	0.365	0.111	0.143	0.008

Problem 160	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	11	16	5	13	7	13	14	14
N.S.	1	1.00	1.00	1.45	0.45	1.18	0.64	1.18	1.27	1.27
time (sec)	N/A	0.155	0.010	0.028	0.045	0.065	0.464	0.118	0.162	0.014

Problem 161	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	15	9	13	10	18	14	9
N.S.	1	1.00	1.00	1.07	0.64	0.93	0.71	1.29	1.00	0.64
time (sec)	N/A	0.149	0.004	0.033	0.045	0.069	0.492	0.124	0.160	0.010

Problem 162	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	17	16	14	0	14	21	14
N.S.	1	1.00	1.00	1.13	1.07	0.93	0.00	0.93	1.40	0.93
time (sec)	N/A	0.141	0.027	0.083	0.052	0.065	0.000	0.123	0.166	0.016

Problem 163	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	13	14	13	10	0	10	16	10
N.S.	1	1.00	1.00	1.08	1.00	0.77	0.00	0.77	1.23	0.77
time (sec)	N/A	0.215	0.043	0.079	0.052	0.067	0.000	0.113	0.156	0.066

Problem 164	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	19	22	16	23	0	80	16	17
N.S.	1	1.00	1.00	1.16	0.84	1.21	0.00	4.21	0.84	0.89
time (sec)	N/A	0.165	0.038	0.067	0.053	0.061	0.000	0.122	0.154	0.075

Problem 165	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	18	19	18	19	0	16	22	0
N.S.	1	1.00	1.00	1.06	1.00	1.06	0.00	0.89	1.22	0.00
time (sec)	N/A	0.165	0.007	0.040	0.062	0.070	0.000	0.121	0.141	0.000

Problem 166	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	11	10	9	9	7	9	11	9
N.S.	1	1.00	0.69	0.62	0.56	0.56	0.44	0.56	0.69	0.56
time (sec)	N/A	0.151	0.009	0.019	0.023	0.062	0.029	0.119	0.147	0.010

Problem 167	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	26	29	16	15	14	14	12	14	16	14
N.S.	1	1.12	0.62	0.58	0.54	0.54	0.46	0.54	0.62	0.54
time (sec)	N/A	0.184	0.011	0.022	0.029	0.068	0.030	0.118	0.145	0.016

Problem 168	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	36	42	21	20	19	19	17	19	21	19
N.S.	1	1.17	0.58	0.56	0.53	0.53	0.47	0.53	0.58	0.53
time (sec)	N/A	0.219	0.011	0.031	0.031	0.066	0.030	0.122	0.151	0.013

Problem 169	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	48	39	60	181	60	360	77	46	2034
N.S.	1	1.00	0.81	1.25	3.77	1.25	7.50	1.60	0.96	42.38
time (sec)	N/A	0.196	0.144	0.348	0.114	0.103	0.341	0.126	0.152	6.174

Problem 170	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2	2	2	9	3	2	2	3	3	2
N.S.	1	1.00	1.00	4.50	1.50	1.00	1.00	1.50	1.50	1.00
time (sec)	N/A	0.119	0.037	0.009	0.064	0.068	0.178	0.122	0.148	0.005

Problem 171	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	10	17	6	14	7	11	15	10
N.S.	1	1.00	1.00	1.70	0.60	1.40	0.70	1.10	1.50	1.00
time (sec)	N/A	0.142	0.004	0.015	0.075	0.071	0.162	0.122	0.146	0.021

Problem 172	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	A	C	A	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	22	0	22	18	24	0	13	22
N.S.	1	1.00	1.00	0.00	1.00	0.82	1.09	0.00	0.59	1.00
time (sec)	N/A	0.158	0.018	0.000	0.069	0.085	0.247	0.000	0.153	0.039

Problem 173	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	10	12	11	9	0	9	13	9
N.S.	1	1.00	0.83	1.00	0.92	0.75	0.00	0.75	1.08	0.75
time (sec)	N/A	0.140	0.037	0.070	0.071	0.070	0.000	0.120	0.155	0.009

Problem 174	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	22	23	0	20	0	20	18	0
N.S.	1	1.00	1.00	1.05	0.00	0.91	0.00	0.91	0.82	0.00
time (sec)	N/A	0.190	0.116	0.069	0.000	0.069	0.000	0.115	0.152	0.000

Problem 175	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	103	103	17	14	0	6	27	0	16	155
N.S.	1	1.00	0.17	0.14	0.00	0.06	0.26	0.00	0.16	1.50
time (sec)	N/A	0.156	10.020	0.145	0.000	0.075	0.329	0.000	0.155	0.173

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [83] had the largest ratio of [2]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	1	1	1.00	9	0.111
2	A	2	2	1.00	11	0.182
3	A	2	2	1.00	11	0.182
4	A	2	2	1.00	11	0.182
5	A	1	1	1.00	14	0.071
6	A	4	3	1.00	4	0.750
7	A	2	2	1.00	9	0.222
8	A	5	4	1.00	7	0.571
9	A	4	3	1.00	17	0.176
10	A	4	3	1.00	9	0.333
11	A	8	7	1.00	11	0.636
12	A	5	4	1.00	16	0.250
13	A	4	3	1.00	10	0.300
14	A	1	1	1.00	15	0.067
15	A	2	2	1.00	9	0.222
16	A	1	1	1.00	9	0.111
17	A	1	1	1.00	15	0.067
18	A	2	2	1.00	17	0.118
19	A	4	3	0.75	20	0.150
20	A	1	1	1.00	26	0.038
21	A	1	1	1.00	20	0.050

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	4	4	1.00	4	1.000
23	A	7	7	1.00	6	1.167
24	A	10	10	1.13	6	1.667
25	A	9	9	1.12	6	1.500
26	A	4	3	1.00	5	0.600
27	A	4	4	1.22	6	0.667
28	A	3	3	1.00	4	0.750
29	A	4	3	1.00	4	0.750
30	A	5	5	1.21	4	1.250
31	A	4	3	1.00	4	0.750
32	A	7	7	1.29	4	1.750
33	A	3	3	1.00	6	0.500
34	A	6	6	0.97	6	1.000
35	A	6	6	1.12	8	0.750
36	A	3	3	1.00	4	0.750
37	A	4	3	1.00	4	0.750
38	A	5	5	1.21	4	1.250
39	A	6	5	1.12	13	0.385
40	A	4	3	1.11	13	0.231
41	A	2	2	1.00	13	0.154
42	A	3	3	1.00	4	0.750
43	A	5	5	1.00	4	1.250
44	A	3	3	1.00	4	0.750
45	A	5	5	1.00	4	1.250
46	A	4	4	1.00	10	0.400
47	A	1	1	1.00	11	0.091
48	A	1	1	1.00	9	0.111
49	A	1	1	1.00	13	0.077
50	A	1	1	1.00	11	0.091
51	A	2	2	1.00	11	0.182
52	A	4	3	1.00	8	0.375
53	A	11	10	1.08	8	1.250

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
54	A	1	1	1.00	10	0.100
55	A	5	4	1.00	17	0.235
56	A	1	1	1.00	7	0.143
57	A	2	2	1.13	4	0.500
58	A	1	1	1.00	4	0.250
59	A	2	2	1.00	6	0.333
60	A	1	1	1.00	5	0.200
61	A	3	3	1.00	2	1.500
62	A	1	1	1.00	8	0.125
63	A	2	2	1.18	8	0.250
64	A	3	2	1.00	8	0.250
65	A	3	2	1.00	14	0.143
66	A	4	3	1.00	14	0.214
67	A	3	3	1.26	8	0.375
68	A	1	1	1.00	9	0.111
69	A	1	1	1.00	13	0.077
70	A	3	2	1.00	9	0.222
71	A	1	1	1.00	6	0.167
72	A	1	1	1.00	6	0.167
73	A	5	4	1.30	7	0.571
74	A	2	2	1.00	5	0.400
75	A	2	2	1.00	7	0.286
76	A	3	3	1.11	7	0.429
77	A	3	3	1.00	9	0.333
78	A	4	3	1.04	7	0.429
79	A	2	2	1.00	11	0.182
80	A	1	1	1.00	10	0.100
81	A	1	1	1.00	10	0.100
82	A	2	2	1.00	2	1.000
83	A	5	4	1.00	2	2.000
84	A	5	4	1.00	2	2.000
85	A	3	3	1.04	4	0.750

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
86	A	5	4	1.00	6	0.667
87	A	3	2	1.00	13	0.154
88	A	3	2	1.70	14	0.143
89	A	1	1	1.00	9	0.111
90	A	1	1	1.00	9	0.111
91	A	3	2	1.00	10	0.200
92	A	3	3	0.95	4	0.750
93	A	5	4	1.08	6	0.667
94	A	5	5	1.00	6	0.833
95	A	5	4	1.32	6	0.667
96	A	1	1	1.00	17	0.059
97	A	2	2	1.00	11	0.182
98	A	1	1	1.00	15	0.067
99	A	1	1	1.00	14	0.071
100	A	2	2	1.00	11	0.182
101	A	3	2	1.00	13	0.154
102	A	7	6	1.22	10	0.600
103	A	4	3	1.38	15	0.200
104	A	5	4	1.00	15	0.267
105	A	4	3	1.00	15	0.200
106	A	2	2	1.00	16	0.125
107	A	2	2	1.00	16	0.125
108	A	3	3	1.00	18	0.167
109	A	3	3	1.00	23	0.130
110	A	2	2	1.00	19	0.105
111	A	7	6	1.04	18	0.333
112	A	4	4	1.00	31	0.129
113	A	4	3	1.00	7	0.429
114	A	4	3	1.00	22	0.136
115	A	2	2	1.00	16	0.125
116	A	2	2	1.00	17	0.118
117	A	2	2	1.00	12	0.167

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
118	A	3	3	1.00	21	0.143
119	A	2	2	1.00	20	0.100
120	A	4	4	1.00	16	0.250
121	A	3	3	1.00	16	0.188
122	A	2	2	1.00	11	0.182
123	A	4	3	1.00	11	0.273
124	A	2	2	1.00	16	0.125
125	A	2	2	1.00	7	0.286
126	A	7	6	1.12	9	0.667
127	A	2	2	1.00	12	0.167
128	A	3	3	1.00	14	0.214
129	A	5	4	1.43	21	0.190
130	A	3	3	1.00	18	0.167
131	A	2	2	1.00	7	0.286
132	A	4	4	1.00	11	0.364
133	A	4	3	1.80	16	0.188
134	A	3	3	1.00	11	0.273
135	A	2	2	1.00	18	0.111
136	A	3	3	1.00	7	0.429
137	A	9	8	1.06	7	1.143
138	A	4	3	1.00	14	0.214
139	A	1	1	1.00	16	0.062
140	A	5	4	0.58	12	0.333
141	A	4	3	1.00	8	0.375
142	A	4	3	0.36	8	0.375
143	A	2	2	1.00	10	0.200
144	A	6	5	0.50	13	0.385
145	A	4	3	1.00	19	0.158
146	A	2	2	1.00	11	0.182
147	A	6	5	1.00	11	0.455
148	A	2	2	1.00	11	0.182
149	A	1	1	1.00	13	0.077

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
150	A	5	4	1.16	15	0.267
151	A	4	3	1.00	13	0.231
152	A	2	2	1.00	9	0.222
153	A	4	3	1.00	12	0.250
154	A	3	2	1.00	9	0.222
155	A	8	7	1.06	18	0.389
156	A	2	2	1.00	8	0.250
157	A	6	6	1.00	5	1.200
158	A	1	1	1.00	7	0.143
159	A	1	1	1.00	9	0.111
160	A	2	2	1.00	7	0.286
161	A	2	2	1.00	5	0.400
162	A	1	1	1.00	14	0.071
163	A	3	2	1.00	14	0.143
164	A	2	2	1.00	9	0.222
165	A	2	2	1.00	8	0.250
166	A	2	2	1.00	7	0.286
167	A	3	3	1.12	9	0.333
168	A	4	4	1.17	9	0.444
169	A	2	2	1.00	21	0.095
170	A	1	1	1.00	4	0.250
171	A	2	2	1.00	4	0.500
172	A	3	2	1.00	8	0.250
173	A	1	1	1.00	11	0.091
174	A	2	2	1.00	16	0.125
175	A	1	1	1.00	9	0.111

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \sqrt{1+2x} dx$	89
3.2	$\int x\sqrt{1+3x} dx$	94
3.3	$\int x^2\sqrt{1+x} dx$	99
3.4	$\int \frac{x}{\sqrt{2-3x}} dx$	104
3.5	$\int \frac{1+x}{(2+2x+x^2)^3} dx$	109
3.6	$\int \sin^3(x) dx$	114
3.7	$\int \sqrt[3]{-1+zz} dz$	119
3.8	$\int \cot(x) \csc^2(x) dx$	124
3.9	$\int \cos(2x)\sqrt{4-\sin(2x)} dx$	129
3.10	$\int \frac{\sin(x)}{(3+\cos(x))^2} dx$	134
3.11	$\int \frac{\sin(x)}{\sqrt{\cos^3(x)}} dx$	139
3.12	$\int \frac{\sin(\sqrt{1+x})}{\sqrt{1+x}} dx$	145
3.13	$\int x^{-1+n} \sin(x^n) dx$	150
3.14	$\int \frac{x^5}{\sqrt{1-x^6}} dx$	155
3.15	$\int t\sqrt[4]{1+t} dt$	160
3.16	$\int \frac{1}{(1+x^2)^{3/2}} dx$	165
3.17	$\int x^2(27+8x^3)^{2/3} dx$	170
3.18	$\int \frac{\cos(x)+\sin(x)}{\sqrt[3]{-\cos(x)+\sin(x)}} dx$	175
3.19	$\int \frac{x}{\sqrt{1+x^2+(1+x^2)^{3/2}}} dx$	180
3.20	$\int \frac{x}{\sqrt{1+x^2}\sqrt{1+\sqrt{1+x^2}}} dx$	185
3.21	$\int \frac{\sqrt[5]{1-2x+x^2}}{1-x} dx$	190
3.22	$\int x \sin(x) dx$	195
3.23	$\int x^2 \sin(x) dx$	200
3.24	$\int x^3 \cos(x) dx$	205

3.25	$\int x^3 \sin(x) dx$	211
3.26	$\int \cos(x) \sin(x) dx$	217
3.27	$\int x \cos(x) \sin(x) dx$	222
3.28	$\int \sin^2(x) dx$	227
3.29	$\int \sin^3(x) dx$	232
3.30	$\int \sin^4(x) dx$	237
3.31	$\int \sin^5(x) dx$	242
3.32	$\int \sin^6(x) dx$	247
3.33	$\int x \sin^2(x) dx$	252
3.34	$\int x \sin^3(x) dx$	257
3.35	$\int x^2 \sin^2(x) dx$	262
3.36	$\int \cos^2(x) dx$	268
3.37	$\int \cos^3(x) dx$	273
3.38	$\int \cos^4(x) dx$	278
3.39	$\int (a^2 - x^2)^{5/2} dx$	283
3.40	$\int \frac{x^5}{\sqrt{5+x^2}} dx$	289
3.41	$\int \frac{t^3}{\sqrt{4+t^3}} dt$	294
3.42	$\int \tan^2(x) dx$	300
3.43	$\int \tan^4(x) dx$	305
3.44	$\int \cot^2(x) dx$	310
3.45	$\int \cot^4(x) dx$	315
3.46	$\int (2 + 3x) \sin(5x) dx$	320
3.47	$\int x\sqrt{1+x^2} dx$	325
3.48	$\int x(-1+x^2)^9 dx$	330
3.49	$\int \frac{3+2x}{(7+6x)^3} dx$	335
3.50	$\int x^4(1+x^5)^5 dx$	340
3.51	$\int (1-x)^{20} x^4 dx$	345
3.52	$\int \frac{\sin(\frac{1}{x})}{x^2} dx$	352
3.53	$\int \sin(\sqrt[4]{-1+x}) dx$	357
3.54	$\int x \cos(x^2) \sin(x^2) dx$	363
3.55	$\int \sqrt{1+3\cos^2(x)} \sin(2x) dx$	368
3.56	$\int \frac{1}{2+3x} dx$	373
3.57	$\int \log^2(x) dx$	378
3.58	$\int x \log(x) dx$	383
3.59	$\int x \log^2(x) dx$	388
3.60	$\int \frac{1}{1+t} dt$	393
3.61	$\int \cot(x) dx$	398
3.62	$\int x^n \log(ax) dx$	403

3.63	$\int x^2 \log^2(x) dx$	408
3.64	$\int \frac{1}{x \log(x)} dx$	413
3.65	$\int \frac{\log(1-t)}{1-t} dt$	418
3.66	$\int \frac{\log(x)}{x\sqrt{1+\log(x)}} dx$	423
3.67	$\int x^3 \log^3(x) dx$	428
3.68	$\int e^{x^3} x^2 dx$	433
3.69	$\int \frac{2\sqrt{x}}{\sqrt{x}} dx$	438
3.70	$\int e^{2\sin(x)} \cos(x) dx$	443
3.71	$\int e^x \sin(x) dx$	448
3.72	$\int e^x \cos(x) dx$	453
3.73	$\int \frac{1}{1+e^x} dx$	458
3.74	$\int e^x x dx$	463
3.75	$\int e^{-x} x dx$	468
3.76	$\int e^x x^2 dx$	473
3.77	$\int e^{-2x} x^2 dx$	478
3.78	$\int e^{\sqrt{x}} dx$	483
3.79	$\int e^{-x^2} x^3 dx$	488
3.80	$\int e^{ax} \cos(bx) dx$	493
3.81	$\int e^{ax} \sin(bx) dx$	498
3.82	$\int \cot^{-1}(x) dx$	503
3.83	$\int \sec^{-1}(x) dx$	508
3.84	$\int \csc^{-1}(x) dx$	514
3.85	$\int \arcsin(x)^2 dx$	520
3.86	$\int \frac{\arcsin(x)}{x^2} dx$	525
3.87	$\int \frac{1}{\sqrt{a^2-x^2}} dx$	530
3.88	$\int \frac{1}{\sqrt{1-2x-x^2}} dx$	535
3.89	$\int \frac{1}{a^2+x^2} dx$	540
3.90	$\int \frac{1}{a+bx^2} dx$	545
3.91	$\int \frac{1}{2-x+x^2} dx$	550
3.92	$\int x \arctan(x) dx$	555
3.93	$\int x^2 \arccos(x) dx$	560
3.94	$\int x \arctan(x)^2 dx$	565
3.95	$\int \arctan(\sqrt{x}) dx$	571
3.96	$\int \frac{\arctan(\sqrt{x})}{\sqrt{x(1+x)}} dx$	577
3.97	$\int \sqrt{1-x^2} dx$	582
3.98	$\int \frac{e^{\arctan(x)} x}{(1+x^2)^{3/2}} dx$	587
3.99	$\int \frac{e^{\arctan(x)}}{(1+x^2)^{3/2}} dx$	592

3.100	$\int \frac{x^2}{(1+x^2)^2} dx$	597
3.101	$\int \frac{e^x}{1+e^{2x}} dx$	602
3.102	$\int e^{-x} \cot^{-1}(e^x) dx$	607
3.103	$\int \sqrt{\frac{a+x}{a-x}} dx$	613
3.104	$\int \sqrt{(b-x)(-a+x)} dx$	618
3.105	$\int \frac{1}{\sqrt{(b-x)(-a+x)}} dx$	624
3.106	$\int \frac{3+5x}{-3+2x+x^2} dx$	630
3.107	$\int \frac{5+2x}{-3+2x+x^2} dx$	635
3.108	$\int \frac{3x+x^3}{-3-2x+x^2} dx$	640
3.109	$\int \frac{-1+5x+2x^2}{-2x+x^2+x^3} dx$	645
3.110	$\int \frac{3+2x+x^2}{(-1+x)(1+x)^2} dx$	650
3.111	$\int \frac{-2+2x+3x^2}{-1+x^3} dx$	655
3.112	$\int \frac{2-x+2x^2-x^3+x^4}{(-1+x)(2+x^2)^2} dx$	661
3.113	$\int \frac{1}{\cos(x)+\sin(x)} dx$	667
3.114	$\int \frac{x}{4-x^2+\sqrt{4-x^2}} dx$	672
3.115	$\int \frac{3+2x}{(-2+x)(5+x)} dx$	677
3.116	$\int \frac{x}{(1+x)(2+x)(3+x)} dx$	682
3.117	$\int \frac{x}{2-3x+x^3} dx$	687
3.118	$\int \frac{-6+2x+x^4}{-2x+x^2+x^3} dx$	692
3.119	$\int \frac{7+8x^3}{(1+x)(1+2x)^3} dx$	697
3.120	$\int \frac{1+x+4x^2}{-1+x^3} dx$	702
3.121	$\int \frac{x^4}{4+5x^2+x^4} dx$	707
3.122	$\int \frac{2+x}{x+x^2} dx$	712
3.123	$\int \frac{1}{x(1+x^2)^2} dx$	717
3.124	$\int \frac{1}{(1+x)(2+x)^2(3+x)^3} dx$	722
3.125	$\int \frac{x}{(1+x)^2} dx$	728
3.126	$\int \frac{1}{-x+x^3} dx$	733
3.127	$\int \frac{x^2}{-6+x+x^2} dx$	739
3.128	$\int \frac{2+x}{4-4x+x^2} dx$	744
3.129	$\int \frac{1}{(4-4x+x^2)(5-4x+x^2)} dx$	749
3.130	$\int \frac{-3+x}{2x+3x^2+x^3} dx$	754
3.131	$\int \frac{1}{(-1+x^2)^2} dx$	759
3.132	$\int \frac{1+x}{-1+x^3} dx$	764
3.133	$\int \frac{1+x^4}{x(1+x^2)^2} dx$	769
3.134	$\int \frac{1}{-2x^3+x^4} dx$	774
3.135	$\int \frac{1-x^3}{x(1+x^2)} dx$	779

3.136	$\int \frac{1}{-1+x^4} dx$	784
3.137	$\int \frac{1}{1+x^4} dx$	789
3.138	$\int \frac{x^2}{(2+2x+x^2)^2} dx$	797
3.139	$\int \frac{-1+4x^5}{(1+x+x^5)^2} dx$	802
3.140	$\int \frac{1}{5-\cos(x)+2\sin(x)} dx$	807
3.141	$\int \frac{1}{1+a\cos(x)} dx$	813
3.142	$\int \frac{1}{1+2\cos(x)} dx$	819
3.143	$\int \frac{1}{1+\frac{\cos(x)}{2}} dx$	824
3.144	$\int \frac{\sin^2(x)}{1+\sin^2(x)} dx$	829
3.145	$\int \frac{1}{b^2\cos^2(x)+a^2\sin^2(x)} dx$	835
3.146	$\int \frac{1}{(b\cos(x)+a\sin(x))^2} dx$	841
3.147	$\int \frac{\sin(x)}{1+\cos(x)+\sin(x)} dx$	847
3.148	$\int \sqrt{3-x^2} dx$	853
3.149	$\int \frac{x}{\sqrt{3-x^2}} dx$	858
3.150	$\int \frac{\sqrt{3-x^2}}{x} dx$	863
3.151	$\int \frac{\sqrt{x+x^2}}{x} dx$	869
3.152	$\int \sqrt{5+x^2} dx$	874
3.153	$\int \frac{x}{\sqrt{1+x+x^2}} dx$	879
3.154	$\int \frac{1}{\sqrt{x+x^2}} dx$	884
3.155	$\int \frac{\sqrt{2-x-x^2}}{x^2} dx$	889
3.156	$\int \frac{\log(t)}{1+t} dt$	896
3.157	$\int \log(e^{\cos(x)}) dx$	901
3.158	$\int \frac{e^t}{t} dt$	906
3.159	$\int \frac{e^{at}}{t} dt$	911
3.160	$\int \frac{e^t}{t^2} dt$	916
3.161	$\int e^{\frac{1}{t}} dt$	921
3.162	$\int \frac{e^{-t}}{-1-a+t} dt$	926
3.163	$\int \frac{e^{t^2}t}{1+t^2} dt$	930
3.164	$\int \frac{e^t}{(1+t)^2} dt$	935
3.165	$\int e^t \log(1+t) dt$	940
3.166	$\int e^{-t}t dt$	945
3.167	$\int e^{-t}t^2 dt$	950
3.168	$\int e^{-t}t^3 dt$	955
3.169	$\int \frac{b1\cos(x)+a1\sin(x)}{b\cos(x)+a\sin(x)} dx$	960
3.170	$\int \frac{1}{\log(t)} dt$	967
3.171	$\int \frac{1}{\log^2(t)} dt$	971

3.172	$\int \log^{-1-n}(t) dt$	976
3.173	$\int \frac{e^{2t}}{-1+t} dt$	981
3.174	$\int \frac{e^{2x}}{2-3x+x^2} dx$	986
3.175	$\int \frac{1}{\sqrt{1+t^3}} dt$	991

3.1 $\int \sqrt{1 + 2x} dx$

Optimal result	89
Mathematica [A] (verified)	89
Rubi [A] (verified)	90
Maple [A] (verified)	91
Fricas [A] (verification not implemented)	91
Sympy [A] (verification not implemented)	92
Maxima [A] (verification not implemented)	92
Giac [A] (verification not implemented)	92
Mupad [B] (verification not implemented)	93
Reduce [B] (verification not implemented)	93

Optimal result

Integrand size = 9, antiderivative size = 13

$$\int \sqrt{1 + 2x} dx = \frac{1}{3}(1 + 2x)^{3/2}$$

output `1/3*(1+2*x)^(3/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \sqrt{1 + 2x} dx = \frac{1}{3}(1 + 2x)^{3/2}$$

input `Integrate[Sqrt[1 + 2*x],x]`

output `(1 + 2*x)^(3/2)/3`

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{2x+1} dx$$

$$\downarrow 17$$

$$\frac{1}{3}(2x+1)^{3/2}$$

input `Int[Sqrt[1 + 2*x], x]`

output `(1 + 2*x)^(3/2)/3`

Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] :> Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

method	result	size
gospers	$\frac{(1+2x)^{\frac{3}{2}}}{3}$	10
derivativedivides	$\frac{(1+2x)^{\frac{3}{2}}}{3}$	10
default	$\frac{(1+2x)^{\frac{3}{2}}}{3}$	10
risch	$\frac{(1+2x)^{\frac{3}{2}}}{3}$	10
pseudoelliptic	$\frac{(1+2x)^{\frac{3}{2}}}{3}$	10
trager	$\left(\frac{1}{3} + \frac{2x}{3}\right) \sqrt{1+2x}$	14
orering	$\left(\frac{1}{3} + \frac{2x}{3}\right) \sqrt{1+2x}$	14
meijerg	$-\frac{\frac{4\sqrt{\pi}}{3} - 2\sqrt{\pi} \frac{(2+4x)\sqrt{1+2x}}{3}}{4\sqrt{\pi}}$	29

input `int((1+2*x)^(1/2),x,method=_RETURNVERBOSE)`output `1/3*(1+2*x)^(3/2)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int \sqrt{1+2x} dx = \frac{1}{3} (2x+1)^{\frac{3}{2}}$$

input `integrate((1+2*x)^(1/2),x, algorithm="fricas")`output `1/3*(2*x + 1)^(3/2)`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.62

$$\int \sqrt{1+2x} dx = \frac{(2x+1)^{\frac{3}{2}}}{3}$$

input `integrate((1+2*x)**(1/2),x)`

output `(2*x + 1)**(3/2)/3`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int \sqrt{1+2x} dx = \frac{1}{3} (2x+1)^{\frac{3}{2}}$$

input `integrate((1+2*x)^(1/2),x, algorithm="maxima")`

output `1/3*(2*x + 1)^(3/2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int \sqrt{1+2x} dx = \frac{1}{3} (2x+1)^{\frac{3}{2}}$$

input `integrate((1+2*x)^(1/2),x, algorithm="giac")`

output `1/3*(2*x + 1)^(3/2)`

Mupad [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int \sqrt{1+2x} dx = \frac{(2x+1)^{3/2}}{3}$$

input `int((2*x + 1)^(1/2), x)`

output `(2*x + 1)^(3/2)/3`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \sqrt{1+2x} dx = \frac{\sqrt{2x+1}(2x+1)}{3}$$

input `int((1+2*x)^(1/2), x)`

output `(sqrt(2*x + 1)*(2*x + 1))/3`

3.2 $\int x\sqrt{1+3x} dx$

Optimal result	94
Mathematica [A] (verified)	94
Rubi [A] (verified)	95
Maple [A] (verified)	96
Fricas [A] (verification not implemented)	96
Sympy [A] (verification not implemented)	97
Maxima [A] (verification not implemented)	97
Giac [A] (verification not implemented)	97
Mupad [B] (verification not implemented)	98
Reduce [B] (verification not implemented)	98

Optimal result

Integrand size = 11, antiderivative size = 27

$$\int x\sqrt{1+3x} dx = -\frac{2}{27}(1+3x)^{3/2} + \frac{2}{45}(1+3x)^{5/2}$$

output `-2/27*(1+3*x)^(3/2)+2/45*(1+3*x)^(5/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.67

$$\int x\sqrt{1+3x} dx = \frac{2}{135}(1+3x)^{3/2}(-2+9x)$$

input `Integrate[x*Sqrt[1 + 3*x],x]`

output `(2*(1 + 3*x)^(3/2)*(-2 + 9*x))/135`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x\sqrt{3x+1} dx$$

$$\downarrow 53$$

$$\int \left(\frac{1}{3}(3x+1)^{3/2} - \frac{1}{3}\sqrt{3x+1} \right) dx$$

$$\downarrow 2009$$

$$\frac{2}{45}(3x+1)^{5/2} - \frac{2}{27}(3x+1)^{3/2}$$

input `Int[x*Sqrt[1 + 3*x],x]`

output `(-2*(1 + 3*x)^(3/2))/27 + (2*(1 + 3*x)^(5/2))/45`

Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.56

method	result	size
gospers	$\frac{2(1+3x)^{\frac{3}{2}}(9x-2)}{135}$	15
orering	$\frac{2(1+3x)^{\frac{3}{2}}(9x-2)}{135}$	15
trager	$\left(\frac{2}{5}x^2 + \frac{2}{45}x - \frac{4}{135}\right)\sqrt{1+3x}$	19
derivativdivides	$-\frac{2(1+3x)^{\frac{3}{2}}}{27} + \frac{2(1+3x)^{\frac{5}{2}}}{45}$	20
default	$-\frac{2(1+3x)^{\frac{3}{2}}}{27} + \frac{2(1+3x)^{\frac{5}{2}}}{45}$	20
risch	$\frac{2(27x^2+3x-2)\sqrt{1+3x}}{135}$	20
pseudoelliptic	$\frac{2(27x^2+3x-2)\sqrt{1+3x}}{135}$	20
meijerg	$-\frac{-\frac{8\sqrt{\pi}}{15} + \frac{4\sqrt{\pi}(1+3x)^{\frac{3}{2}}(-9x+2)}{15}}{18\sqrt{\pi}}$	29

input `int(x*(1+3*x)^(1/2),x,method=_RETURNVERBOSE)`output `2/135*(1+3*x)^(3/2)*(9*x-2)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int x\sqrt{1+3x} dx = \frac{2}{135} (27x^2 + 3x - 2)\sqrt{3x+1}$$

input `integrate(x*(1+3*x)^(1/2),x, algorithm="fricas")`output `2/135*(27*x^2 + 3*x - 2)*sqrt(3*x + 1)`

Sympy [A] (verification not implemented)

Time = 0.53 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.44

$$\int x\sqrt{1+3x} dx = \frac{2x^2\sqrt{3x+1}}{5} + \frac{2x\sqrt{3x+1}}{45} - \frac{4\sqrt{3x+1}}{135}$$

input `integrate(x*(1+3*x)**(1/2),x)`output `2*x**2*sqrt(3*x + 1)/5 + 2*x*sqrt(3*x + 1)/45 - 4*sqrt(3*x + 1)/135`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int x\sqrt{1+3x} dx = \frac{2}{45} (3x+1)^{\frac{5}{2}} - \frac{2}{27} (3x+1)^{\frac{3}{2}}$$

input `integrate(x*(1+3*x)^(1/2),x, algorithm="maxima")`output `2/45*(3*x + 1)^(5/2) - 2/27*(3*x + 1)^(3/2)`**Giac [A] (verification not implemented)**

Time = 0.57 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int x\sqrt{1+3x} dx = \frac{2}{45} (3x+1)^{\frac{5}{2}} - \frac{2}{27} (3x+1)^{\frac{3}{2}}$$

input `integrate(x*(1+3*x)^(1/2),x, algorithm="giac")`output `2/45*(3*x + 1)^(5/2) - 2/27*(3*x + 1)^(3/2)`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.52

$$\int x\sqrt{1+3x} dx = \frac{2(3x+1)^{3/2}(9x-2)}{135}$$

input `int(x*(3*x + 1)^(1/2),x)`

output `(2*(3*x + 1)^(3/2)*(9*x - 2))/135`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.67

$$\int x\sqrt{1+3x} dx = \frac{2\sqrt{3x+1}(27x^2+3x-2)}{135}$$

input `int(x*(1+3*x)^(1/2),x)`

output `(2*sqrt(3*x + 1)*(27*x**2 + 3*x - 2))/135`

3.3 $\int x^2 \sqrt{1+x} dx$

Optimal result	99
Mathematica [A] (verified)	99
Rubi [A] (verified)	100
Maple [A] (verified)	101
Fricas [A] (verification not implemented)	101
Sympy [A] (verification not implemented)	102
Maxima [A] (verification not implemented)	102
Giac [A] (verification not implemented)	102
Mupad [B] (verification not implemented)	103
Reduce [B] (verification not implemented)	103

Optimal result

Integrand size = 11, antiderivative size = 34

$$\int x^2 \sqrt{1+x} dx = \frac{2}{3}(1+x)^{3/2} - \frac{4}{5}(1+x)^{5/2} + \frac{2}{7}(1+x)^{7/2}$$

output $2/3*(1+x)^{(3/2)}-4/5*(1+x)^{(5/2)}+2/7*(1+x)^{(7/2)}$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.62

$$\int x^2 \sqrt{1+x} dx = \frac{2}{105}(1+x)^{3/2} (8 - 12x + 15x^2)$$

input `Integrate[x^2*Sqrt[1 + x],x]`

output $(2*(1+x)^{(3/2)}*(8-12*x+15*x^2))/105$

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \sqrt{x+1} dx$$

$$\downarrow 53$$

$$\int \left((x+1)^{5/2} - 2(x+1)^{3/2} + \sqrt{x+1} \right) dx$$

$$\downarrow 2009$$

$$\frac{2}{7}(x+1)^{7/2} - \frac{4}{5}(x+1)^{5/2} + \frac{2}{3}(x+1)^{3/2}$$

input `Int[x^2*Sqrt[1 + x], x]`

output `(2*(1 + x)^(3/2))/3 - (4*(1 + x)^(5/2))/5 + (2*(1 + x)^(7/2))/7`

Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.53

method	result	size
gospers	$\frac{2(1+x)^{\frac{3}{2}}(15x^2-12x+8)}{105}$	18
orering	$\frac{2(1+x)^{\frac{3}{2}}(15x^2-12x+8)}{105}$	18
trager	$\left(\frac{2}{7}x^3 + \frac{2}{35}x^2 - \frac{8}{105}x + \frac{16}{105}\right)\sqrt{1+x}$	22
derivativdivides	$\frac{2(1+x)^{\frac{3}{2}}}{3} - \frac{4(1+x)^{\frac{5}{2}}}{5} + \frac{2(1+x)^{\frac{7}{2}}}{7}$	23
default	$\frac{2(1+x)^{\frac{3}{2}}}{3} - \frac{4(1+x)^{\frac{5}{2}}}{5} + \frac{2(1+x)^{\frac{7}{2}}}{7}$	23
risch	$\frac{2(15x^3+3x^2-4x+8)\sqrt{1+x}}{105}$	23
meijerg	$-\frac{\frac{32\sqrt{\pi}}{105} - \frac{4\sqrt{\pi}(1+x)^{\frac{3}{2}}(15x^2-12x+8)}{105}}{2\sqrt{\pi}}$	32

input `int(x^2*(1+x)^(1/2),x,method=_RETURNVERBOSE)`output `2/105*(1+x)^(3/2)*(15*x^2-12*x+8)`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.65

$$\int x^2\sqrt{1+x} dx = \frac{2}{105} (15x^3 + 3x^2 - 4x + 8)\sqrt{x+1}$$

input `integrate(x^2*(1+x)^(1/2),x, algorithm="fricas")`output `2/105*(15*x^3 + 3*x^2 - 4*x + 8)*sqrt(x + 1)`

Sympy [A] (verification not implemented)

Time = 0.71 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.41

$$\int x^2 \sqrt{1+x} dx = \frac{2x^3 \sqrt{x+1}}{7} + \frac{2x^2 \sqrt{x+1}}{35} - \frac{8x \sqrt{x+1}}{105} + \frac{16 \sqrt{x+1}}{105}$$

input `integrate(x**2*(1+x)**(1/2),x)`output `2*x**3*sqrt(x + 1)/7 + 2*x**2*sqrt(x + 1)/35 - 8*x*sqrt(x + 1)/105 + 16*sqrt(x + 1)/105`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.65

$$\int x^2 \sqrt{1+x} dx = \frac{2}{7} (x+1)^{\frac{7}{2}} - \frac{4}{5} (x+1)^{\frac{5}{2}} + \frac{2}{3} (x+1)^{\frac{3}{2}}$$

input `integrate(x^2*(1+x)^(1/2),x, algorithm="maxima")`output `2/7*(x + 1)^(7/2) - 4/5*(x + 1)^(5/2) + 2/3*(x + 1)^(3/2)`**Giac [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.65

$$\int x^2 \sqrt{1+x} dx = \frac{2}{7} (x+1)^{\frac{7}{2}} - \frac{4}{5} (x+1)^{\frac{5}{2}} + \frac{2}{3} (x+1)^{\frac{3}{2}}$$

input `integrate(x^2*(1+x)^(1/2),x, algorithm="giac")`output `2/7*(x + 1)^(7/2) - 4/5*(x + 1)^(5/2) + 2/3*(x + 1)^(3/2)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.56

$$\int x^2 \sqrt{1+x} dx = -\frac{2(x+1)^{3/2}(42x - 15(x+1)^2 + 7)}{105}$$

input `int(x^2*(x + 1)^(1/2),x)`output `-(2*(x + 1)^(3/2)*(42*x - 15*(x + 1)^2 + 7))/105`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.62

$$\int x^2 \sqrt{1+x} dx = \frac{2\sqrt{x+1}(15x^3 + 3x^2 - 4x + 8)}{105}$$

input `int(x^2*(1+x)^(1/2),x)`output `(2*sqrt(x + 1)*(15*x**3 + 3*x**2 - 4*x + 8))/105`

3.4 $\int \frac{x}{\sqrt{2-3x}} dx$

Optimal result	104
Mathematica [A] (verified)	104
Rubi [A] (verified)	105
Maple [A] (verified)	106
Fricas [A] (verification not implemented)	106
Sympy [C] (verification not implemented)	107
Maxima [A] (verification not implemented)	107
Giac [A] (verification not implemented)	107
Mupad [B] (verification not implemented)	108
Reduce [B] (verification not implemented)	108

Optimal result

Integrand size = 11, antiderivative size = 27

$$\int \frac{x}{\sqrt{2-3x}} dx = -\frac{4}{9}\sqrt{2-3x} + \frac{2}{27}(2-3x)^{3/2}$$

output

```
2/27*(2-3*x)^(3/2)-4/9*(2-3*x)^(1/2)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.67

$$\int \frac{x}{\sqrt{2-3x}} dx = -\frac{2}{27}\sqrt{2-3x}(4+3x)$$

input

```
Integrate[x/Sqrt[2 - 3*x], x]
```

output

```
(-2*Sqrt[2 - 3*x]*(4 + 3*x))/27
```

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\sqrt{2-3x}} dx$$

↓ 53

$$\int \left(\frac{2}{3\sqrt{2-3x}} - \frac{1}{3}\sqrt{2-3x} \right) dx$$

↓ 2009

$$\frac{2}{27}(2-3x)^{3/2} - \frac{4}{9}\sqrt{2-3x}$$

input `Int[x/Sqrt[2 - 3*x], x]`

output `(-4*Sqrt[2 - 3*x])/9 + (2*(2 - 3*x)^(3/2))/27`

Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.52

method	result	size
trager	$\left(-\frac{2x}{9} - \frac{8}{27}\right) \sqrt{2-3x}$	14
gosper	$-\frac{2(3x+4)\sqrt{2-3x}}{27}$	15
pseudoelliptic	$-\frac{2(3x+4)\sqrt{2-3x}}{27}$	15
derivativdivides	$\frac{2(2-3x)^{\frac{3}{2}}}{27} - \frac{4\sqrt{2-3x}}{9}$	20
default	$\frac{2(2-3x)^{\frac{3}{2}}}{27} - \frac{4\sqrt{2-3x}}{9}$	20
risch	$\frac{2(-2+3x)(3x+4)}{27\sqrt{2-3x}}$	20
orering	$\frac{2(-2+3x)(3x+4)}{27\sqrt{2-3x}}$	20
meijerg	$\frac{2\sqrt{2} \left(\frac{4\sqrt{\pi}}{3} - \frac{\sqrt{\pi}(6x+8)\sqrt{1-\frac{3x}{2}}}{6} \right)}{9\sqrt{\pi}}$	32

input `int(x/(2-3*x)^(1/2),x,method=_RETURNVERBOSE)`output `(-2/9*x-8/27)*(2-3*x)^(1/2)`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.52

$$\int \frac{x}{\sqrt{2-3x}} dx = -\frac{2}{27} (3x+4)\sqrt{-3x+2}$$

input `integrate(x/(2-3*x)^(1/2),x, algorithm="fricas")`output `-2/27*(3*x + 4)*sqrt(-3*x + 2)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.52 (sec) , antiderivative size = 60, normalized size of antiderivative = 2.22

$$\int \frac{x}{\sqrt{2-3x}} dx = \begin{cases} -\frac{2ix\sqrt{3x-2}}{9} - \frac{8i\sqrt{3x-2}}{27} & \text{for } |x| > \frac{2}{3} \\ -\frac{2x\sqrt{2-3x}}{9} - \frac{8\sqrt{2-3x}}{27} & \text{otherwise} \end{cases}$$

input `integrate(x/(2-3*x)**(1/2),x)`

output `Piecewise((-2*I*x*sqrt(3*x - 2)/9 - 8*I*sqrt(3*x - 2)/27, Abs(x) > 2/3), (-2*x*sqrt(2 - 3*x)/9 - 8*sqrt(2 - 3*x)/27, True))`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int \frac{x}{\sqrt{2-3x}} dx = \frac{2}{27} (-3x+2)^{\frac{3}{2}} - \frac{4}{9} \sqrt{-3x+2}$$

input `integrate(x/(2-3*x)^(1/2),x, algorithm="maxima")`

output `2/27*(-3*x + 2)^(3/2) - 4/9*sqrt(-3*x + 2)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int \frac{x}{\sqrt{2-3x}} dx = \frac{2}{27} (-3x+2)^{\frac{3}{2}} - \frac{4}{9} \sqrt{-3x+2}$$

input `integrate(x/(2-3*x)^(1/2),x, algorithm="giac")`

output `2/27*(-3*x + 2)^(3/2) - 4/9*sqrt(-3*x + 2)`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.52

$$\int \frac{x}{\sqrt{2-3x}} dx = -\frac{2\sqrt{2-3x}(3x+4)}{27}$$

input `int(x/(2 - 3*x)^(1/2),x)`

output `-(2*(2 - 3*x)^(1/2)*(3*x + 4))/27`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.48

$$\int \frac{x}{\sqrt{2-3x}} dx = \frac{2\sqrt{-3x+2}(-3x-4)}{27}$$

input `int(x/(2-3*x)^(1/2),x)`

output `(2*sqrt(- 3*x + 2)*(- 3*x - 4))/27`

3.5 $\int \frac{1+x}{(2+2x+x^2)^3} dx$

Optimal result	109
Mathematica [A] (verified)	109
Rubi [A] (verified)	110
Maple [A] (verified)	111
Fricas [A] (verification not implemented)	111
Sympy [A] (verification not implemented)	112
Maxima [A] (verification not implemented)	112
Giac [A] (verification not implemented)	112
Mupad [B] (verification not implemented)	113
Reduce [B] (verification not implemented)	113

Optimal result

Integrand size = 14, antiderivative size = 14

$$\int \frac{1+x}{(2+2x+x^2)^3} dx = -\frac{1}{4(2+2x+x^2)^2}$$

output `-1/4/(x^2+2*x+2)^2`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1+x}{(2+2x+x^2)^3} dx = -\frac{1}{4(2+2x+x^2)^2}$$

input `Integrate[(1 + x)/(2 + 2*x + x^2)^3,x]`

output `-1/4*1/(2 + 2*x + x^2)^2`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$, Rules used = {1104}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x+1}{(x^2+2x+2)^3} dx$$

$$\downarrow 1104$$

$$-\frac{1}{4(x^2+2x+2)^2}$$

input `Int[(1 + x)/(2 + 2*x + x^2)^3,x]`

output `-1/4*1/(2 + 2*x + x^2)^2`

Defintions of rubi rules used

rule 1104 `Int[((d_) + (e_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[d*((a + b*x + c*x^2)^(p + 1)/(b*(p + 1))), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[2*c*d - b*e, 0]`

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

method	result	size
gospers	$-\frac{1}{4(x^2+2x+2)^2}$	13
default	$-\frac{1}{4(x^2+2x+2)^2}$	13
norman	$-\frac{1}{4(x^2+2x+2)^2}$	13
risch	$-\frac{1}{4(x^2+2x+2)^2}$	13
parallelrisch	$-\frac{1}{4(x^2+2x+2)^2}$	13
orering	$-\frac{1}{4(x^2+2x+2)^2}$	13

input `int((1+x)/(x^2+2*x+2)^3,x,method=_RETURNVERBOSE)`

output `-1/4/(x^2+2*x+2)^2`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.57

$$\int \frac{1+x}{(2+2x+x^2)^3} dx = -\frac{1}{4(x^4+4x^3+8x^2+8x+4)}$$

input `integrate((1+x)/(x^2+2*x+2)^3,x, algorithm="fricas")`

output `-1/4/(x^4 + 4*x^3 + 8*x^2 + 8*x + 4)`

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.57

$$\int \frac{1+x}{(2+2x+x^2)^3} dx = -\frac{1}{4x^4 + 16x^3 + 32x^2 + 32x + 16}$$

input `integrate((1+x)/(x**2+2*x+2)**3,x)`output `-1/(4*x**4 + 16*x**3 + 32*x**2 + 32*x + 16)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1+x}{(2+2x+x^2)^3} dx = -\frac{1}{4(x^2+2x+2)^2}$$

input `integrate((1+x)/(x^2+2*x+2)^3,x, algorithm="maxima")`output `-1/4/(x^2 + 2*x + 2)^2`**Giac [A] (verification not implemented)**

Time = 1.18 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1+x}{(2+2x+x^2)^3} dx = -\frac{1}{4(x^2+2x+2)^2}$$

input `integrate((1+x)/(x^2+2*x+2)^3,x, algorithm="giac")`output `-1/4/(x^2 + 2*x + 2)^2`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1+x}{(2+2x+x^2)^3} dx = -\frac{1}{4(x^2+2x+2)^2}$$

input `int((x + 1)/(2*x + x^2 + 2)^3,x)`output `-1/(4*(2*x + x^2 + 2)^2)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.71

$$\int \frac{1+x}{(2+2x+x^2)^3} dx = -\frac{1}{4x^4 + 16x^3 + 32x^2 + 32x + 16}$$

input `int((1+x)/(x^2+2*x+2)^3,x)`output `(- 1)/(4*(x**4 + 4*x**3 + 8*x**2 + 8*x + 4))`

3.6 $\int \sin^3(x) dx$

Optimal result	114
Mathematica [A] (verified)	114
Rubi [A] (verified)	115
Maple [A] (verified)	116
Fricas [A] (verification not implemented)	116
Sympy [A] (verification not implemented)	117
Maxima [A] (verification not implemented)	117
Giac [A] (verification not implemented)	117
Mupad [B] (verification not implemented)	118
Reduce [B] (verification not implemented)	118

Optimal result

Integrand size = 4, antiderivative size = 13

$$\int \sin^3(x) dx = -\cos(x) + \frac{\cos^3(x)}{3}$$

output

```
-cos(x)+1/3*cos(x)^3
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \sin^3(x) dx = -\frac{3 \cos(x)}{4} + \frac{1}{12} \cos(3x)$$

input

```
Integrate[Sin[x]^3,x]
```

output

```
(-3*Cos[x])/4 + Cos[3*x]/12
```

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3113, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sin^3(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin(x)^3 dx \\ & \quad \downarrow \text{3113} \\ & - \int (1 - \cos^2(x)) d \cos(x) \\ & \quad \downarrow \text{2009} \\ & \frac{\cos^3(x)}{3} - \cos(x) \end{aligned}$$

input `Int[Sin[x]^3,x]`

output `-Cos[x] + Cos[x]^3/3`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113

```
Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
and[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x]
&& IGtQ[(n - 1)/2, 0]
```

Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

method	result	size
default	$-\frac{(2+\sin(x)^2)\cos(x)}{3}$	11
risch	$-\frac{3\cos(x)}{4} + \frac{\cos(3x)}{12}$	12
parallelrisch	$-\frac{2}{3} - \frac{3\cos(x)}{4} + \frac{\cos(3x)}{12}$	13
orering	$-\sin(x)^2\cos(x) - \frac{2\cos(x)^3}{3}$	16
norman	$\frac{-4\tan(\frac{x}{2})^2 - \frac{4}{3}}{(1+\tan(\frac{x}{2})^2)^3}$	22

input

```
int(sin(x)^3,x,method=_RETURNVERBOSE)
```

output

```
-1/3*(2+sin(x)^2)*cos(x)
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \sin^3(x) dx = \frac{1}{3} \cos(x)^3 - \cos(x)$$

input

```
integrate(sin(x)^3,x, algorithm="fricas")
```

output

```
1/3*cos(x)^3 - cos(x)
```

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.62

$$\int \sin^3(x) dx = \frac{\cos^3(x)}{3} - \cos(x)$$

input `integrate(sin(x)**3,x)`

output `cos(x)**3/3 - cos(x)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \sin^3(x) dx = \frac{1}{3} \cos(x)^3 - \cos(x)$$

input `integrate(sin(x)^3,x, algorithm="maxima")`

output `1/3*cos(x)^3 - cos(x)`

Giac [A] (verification not implemented)

Time = 0.75 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \sin^3(x) dx = \frac{1}{3} \cos(x)^3 - \cos(x)$$

input `integrate(sin(x)^3,x, algorithm="giac")`

output `1/3*cos(x)^3 - cos(x)`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \sin^3(x) dx = \frac{\cos(x) (\cos(x)^2 - 3)}{3}$$

input `int(sin(x)^3,x)`

output `(cos(x)*(cos(x)^2 - 3))/3`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.08

$$\int \sin^3(x) dx = -\frac{\cos(x) \sin(x)^2}{3} - \frac{2 \cos(x)}{3} + \frac{2}{3}$$

input `int(sin(x)^3,x)`

output `(- cos(x)*sin(x)**2 - 2*cos(x) + 2)/3`

3.7 $\int \sqrt[3]{-1 + zz} dz$

Optimal result	119
Mathematica [A] (verified)	119
Rubi [A] (verified)	120
Maple [A] (verified)	121
Fricas [A] (verification not implemented)	121
Sympy [C] (verification not implemented)	122
Maxima [A] (verification not implemented)	122
Giac [A] (verification not implemented)	122
Mupad [B] (verification not implemented)	123
Reduce [B] (verification not implemented)	123

Optimal result

Integrand size = 9, antiderivative size = 23

$$\int \sqrt[3]{-1 + zz} dz = \frac{3}{4}(-1 + z)^{4/3} + \frac{3}{7}(-1 + z)^{7/3}$$

output `3/4*(-1+z)^(4/3)+3/7*(-1+z)^(7/3)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

$$\int \sqrt[3]{-1 + zz} dz = \frac{3}{28}(7 + 4(-1 + z))(-1 + z)^{4/3}$$

input `Integrate[(-1 + z)^(1/3)*z,z]`

output `(3*(7 + 4*(-1 + z))*(-1 + z)^(4/3))/28`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt[3]{z-1} z \, dz$$

$$\downarrow 53$$

$$\int \left((z-1)^{4/3} + \sqrt[3]{z-1} \right) dz$$

$$\downarrow 2009$$

$$\frac{3}{7}(z-1)^{7/3} + \frac{3}{4}(z-1)^{4/3}$$

input `Int[(-1 + z)^(1/3)*z,z]`

output `(3*(-1 + z)^(4/3))/4 + (3*(-1 + z)^(7/3))/7`

Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.57

method	result	size
gospers	$\frac{3(-1+z)^{\frac{4}{3}}(4z+3)}{28}$	13
orering	$\frac{3(-1+z)^{\frac{4}{3}}(4z+3)}{28}$	13
derivativdivides	$\frac{3(-1+z)^{\frac{4}{3}}}{4} + \frac{3(-1+z)^{\frac{7}{3}}}{7}$	16
default	$\frac{3(-1+z)^{\frac{4}{3}}}{4} + \frac{3(-1+z)^{\frac{7}{3}}}{7}$	16
trager	$\left(\frac{3}{7}z^2 - \frac{3}{28}z - \frac{9}{28}\right)(-1+z)^{\frac{1}{3}}$	17
risch	$\frac{3(-1+z)^{\frac{1}{3}}(4z^2-z-3)}{28}$	18
meijerg	$\frac{\text{signum}(-1+z)^{\frac{1}{3}} z^2 \text{hypergeom}\left(\left[-\frac{1}{3}, 2\right], [3], z\right)}{2(-\text{signum}(-1+z))^{\frac{1}{3}}}$	27

input `int((-1+z)^(1/3)*z,z,method=_RETURNVERBOSE)`output `3/28*(-1+z)^(4/3)*(4*z+3)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \sqrt[3]{-1+zz} dz = \frac{3}{28} (4z^2 - z - 3)(z - 1)^{\frac{1}{3}}$$

input `integrate((-1+z)^(1/3)*z,z, algorithm="fricas")`output `3/28*(4*z^2 - z - 3)*(z - 1)^(1/3)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.54 (sec) , antiderivative size = 92, normalized size of antiderivative = 4.00

$$\int \sqrt[3]{-1+zz} dz = \begin{cases} \frac{3z^2 \sqrt[3]{z-1}}{7} - \frac{3z \sqrt[3]{z-1}}{28} - \frac{9 \sqrt[3]{z-1}}{28} & \text{for } |z| > 1 \\ \frac{3z^2 \sqrt[3]{1-ze^{i\pi/3}}}{7} - \frac{3z \sqrt[3]{1-ze^{i\pi/3}}}{28} - \frac{9 \sqrt[3]{1-ze^{i\pi/3}}}{28} & \text{otherwise} \end{cases}$$

input `integrate((-1+z)**(1/3)*z,z)`

output `Piecewise((3*z**2*(z - 1)**(1/3)/7 - 3*z*(z - 1)**(1/3)/28 - 9*(z - 1)**(1/3)/28, Abs(z) > 1), (3*z**2*(1 - z)**(1/3)*exp(I*pi/3)/7 - 3*z*(1 - z)**(1/3)*exp(I*pi/3)/28 - 9*(1 - z)**(1/3)*exp(I*pi/3)/28, True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int \sqrt[3]{-1+zz} dz = \frac{3}{7} (z-1)^{7/3} + \frac{3}{4} (z-1)^{4/3}$$

input `integrate((-1+z)^(1/3)*z,z, algorithm="maxima")`

output `3/7*(z - 1)^(7/3) + 3/4*(z - 1)^(4/3)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int \sqrt[3]{-1+zz} dz = \frac{3}{7} (z-1)^{7/3} + \frac{3}{4} (z-1)^{4/3}$$

input `integrate((-1+z)^(1/3)*z,z, algorithm="giac")`

output $3/7*(z - 1)^{(7/3)} + 3/4*(z - 1)^{(4/3)}$

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.52

$$\int \sqrt[3]{-1 + zz} dz = \frac{3(4z + 3)(z - 1)^{4/3}}{28}$$

input `int(z*(z - 1)^(1/3),z)`

output $(3*(4*z + 3)*(z - 1)^{(4/3)})/28$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \sqrt[3]{-1 + zz} dz = \frac{3(z - 1)^{\frac{1}{3}}(4z^2 - z - 3)}{28}$$

input `int((-1+z)^(1/3)*z,z)`

output $(3*(z - 1)**(1/3)*(4*z**2 - z - 3))/28$

3.8 $\int \cot(x) \csc^2(x) dx$

Optimal result	124
Mathematica [A] (verified)	124
Rubi [A] (verified)	125
Maple [A] (verified)	126
Fricas [A] (verification not implemented)	127
Sympy [A] (verification not implemented)	127
Maxima [A] (verification not implemented)	127
Giac [A] (verification not implemented)	128
Mupad [B] (verification not implemented)	128
Reduce [B] (verification not implemented)	128

Optimal result

Integrand size = 7, antiderivative size = 8

$$\int \cot(x) \csc^2(x) dx = -\frac{1}{2} \csc^2(x)$$

output `-1/2*csc(x)^2`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \cot(x) \csc^2(x) dx = -\frac{1}{2} \csc^2(x)$$

input `Integrate[Cot[x]*Csc[x]^2,x]`

output `-1/2*Csc[x]^2`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 25, 3086, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot(x) \csc^2(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan\left(x - \frac{\pi}{2}\right) \left(-\sec\left(x - \frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{25} \\
 & - \int \sec\left(x - \frac{\pi}{2}\right)^2 \tan\left(x - \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{3086} \\
 & - \int \csc(x) d \csc(x) \\
 & \quad \downarrow \text{15} \\
 & -\frac{1}{2} \csc^2(x)
 \end{aligned}$$

input

Int [Cot [x] *Csc [x] ^2, x]

output

-1/2*Csc [x] ^2

Definitions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)]^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativeldivides	$-\frac{1}{2\sin(x)^2}$	7
default	$-\frac{1}{2\sin(x)^2}$	7
risch	$\frac{2e^{2ix}}{(e^{2ix}-1)^2}$	17
norman	$\frac{-\frac{1}{8} - \frac{\tan(\frac{x}{2})^4}{8}}{\tan(\frac{x}{2})^2}$	18
parallelrisch	$-\frac{\tan(\frac{x}{2})^2}{8} - \frac{\cot(\frac{x}{2})^2}{8}$	18

input `int(cos(x)/sin(x)^3,x,method=_RETURNVERBOSE)`

output `-1/2/sin(x)^2`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.25

$$\int \cot(x) \csc^2(x) dx = \frac{1}{2(\cos(x)^2 - 1)}$$

input `integrate(cos(x)/sin(x)^3,x, algorithm="fricas")`

output `1/2/(cos(x)^2 - 1)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \cot(x) \csc^2(x) dx = -\frac{1}{2 \sin^2(x)}$$

input `integrate(cos(x)/sin(x)**3,x)`

output `-1/(2*sin(x)**2)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cot(x) \csc^2(x) dx = -\frac{1}{2 \sin(x)^2}$$

input `integrate(cos(x)/sin(x)^3,x, algorithm="maxima")`

output `-1/2/sin(x)^2`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cot(x) \csc^2(x) dx = -\frac{1}{2 \sin(x)^2}$$

input `integrate(cos(x)/sin(x)^3,x, algorithm="giac")`

output `-1/2/sin(x)^2`

Mupad [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cot(x) \csc^2(x) dx = -\frac{\cot(x)^2}{2}$$

input `int(cos(x)/sin(x)^3,x)`

output `-cot(x)^2/2`

Reduce [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cot(x) \csc^2(x) dx = -\frac{1}{2 \sin(x)^2}$$

input `int(cos(x)/sin(x)^3,x)`

output `(- 1)/(2*sin(x)**2)`

3.9 $\int \cos(2x) \sqrt{4 - \sin(2x)} dx$

Optimal result	129
Mathematica [A] (verified)	129
Rubi [A] (verified)	130
Maple [A] (verified)	131
Fricas [A] (verification not implemented)	131
Sympy [B] (verification not implemented)	132
Maxima [A] (verification not implemented)	132
Giac [A] (verification not implemented)	132
Mupad [B] (verification not implemented)	133
Reduce [B] (verification not implemented)	133

Optimal result

Integrand size = 17, antiderivative size = 16

$$\int \cos(2x) \sqrt{4 - \sin(2x)} dx = -\frac{1}{3}(4 - \sin(2x))^{3/2}$$

output `-1/3*(4-sin(2*x))^(3/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \cos(2x) \sqrt{4 - \sin(2x)} dx = -\frac{1}{3}(4 - \sin(2x))^{3/2}$$

input `Integrate[Cos[2*x]*Sqrt[4 - Sin[2*x]],x]`

output `-1/3*(4 - Sin[2*x])^(3/2)`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {3042, 3147, 17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{4 - \sin(2x)} \cos(2x) dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{4 - \sin(2x)} \cos(2x) dx \\ & \quad \downarrow \text{3147} \\ & -\frac{1}{2} \int \sqrt{4 - \sin(2x)} d(-\sin(2x)) \\ & \quad \downarrow \text{17} \\ & -\frac{1}{3} (4 - \sin(2x))^{3/2} \end{aligned}$$

input `Int[Cos[2*x]*Sqrt[4 - Sin[2*x]],x]`

output `-1/3*(4 - Sin[2*x])^(3/2)`

Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3147

```
Int[cos[(e_.) + (f_.)*(x_)]^(p_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Simp[1/(b^p*f) Subst[Int[(a + x)^m*(b^2 - x^2)^((p - 1)/2), x], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && IntegerQ[(p - 1)/2] && NeQ[a^2 - b^2, 0]
```

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$-\frac{(4-\sin(2x))^{\frac{3}{2}}}{3}$	13
default	$-\frac{(4-\sin(2x))^{\frac{3}{2}}}{3}$	13

input

```
int(cos(2*x)*(4-sin(2*x))^(1/2),x,method=_RETURNVERBOSE)
```

output

```
-1/3*(4-sin(2*x))^(3/2)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \cos(2x) \sqrt{4 - \sin(2x)} dx = \frac{1}{3} (\sin(2x) - 4) \sqrt{-\sin(2x) + 4}$$

input

```
integrate(cos(2*x)*(4-sin(2*x))^(1/2),x, algorithm="fricas")
```

output

```
1/3*(sin(2*x) - 4)*sqrt(-sin(2*x) + 4)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. $2(12) = 24$.

Time = 0.10 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.81

$$\int \cos(2x)\sqrt{4 - \sin(2x)} dx = \frac{\sqrt{4 - \sin(2x)} \sin(2x)}{3} - \frac{4\sqrt{4 - \sin(2x)}}{3}$$

input `integrate(cos(2*x)*(4-sin(2*x))**(1/2),x)`

output `sqrt(4 - sin(2*x))*sin(2*x)/3 - 4*sqrt(4 - sin(2*x))/3`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \cos(2x)\sqrt{4 - \sin(2x)} dx = -\frac{1}{3}(-\sin(2x) + 4)^{\frac{3}{2}}$$

input `integrate(cos(2*x)*(4-sin(2*x))^(1/2),x, algorithm="maxima")`

output `-1/3*(-sin(2*x) + 4)^(3/2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \cos(2x)\sqrt{4 - \sin(2x)} dx = -\frac{1}{3}(-\sin(2x) + 4)^{\frac{3}{2}}$$

input `integrate(cos(2*x)*(4-sin(2*x))^(1/2),x, algorithm="giac")`

output `-1/3*(-sin(2*x) + 4)^(3/2)`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \cos(2x) \sqrt{4 - \sin(2x)} dx = -\frac{(4 - \sin(2x))^{3/2}}{3}$$

input `int(cos(2*x)*(4 - sin(2*x))^(1/2),x)`

output `-(4 - sin(2*x))^(3/2)/3`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

$$\int \cos(2x) \sqrt{4 - \sin(2x)} dx = \frac{\sqrt{-\sin(2x) + 4} (\sin(2x) - 4)}{3}$$

input `int(cos(2*x)*(4-sin(2*x))^(1/2),x)`

output `(sqrt(- sin(2*x) + 4)*(sin(2*x) - 4))/3`

3.10 $\int \frac{\sin(x)}{(3+\cos(x))^2} dx$

Optimal result	134
Mathematica [A] (verified)	134
Rubi [A] (verified)	135
Maple [A] (verified)	136
Fricas [A] (verification not implemented)	136
Sympy [A] (verification not implemented)	137
Maxima [A] (verification not implemented)	137
Giac [A] (verification not implemented)	137
Mupad [B] (verification not implemented)	138
Reduce [B] (verification not implemented)	138

Optimal result

Integrand size = 9, antiderivative size = 6

$$\int \frac{\sin(x)}{(3 + \cos(x))^2} dx = \frac{1}{3 + \cos(x)}$$

output `1/(3+cos(x))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{\sin(x)}{(3 + \cos(x))^2} dx = \frac{1}{3 + \cos(x)}$$

input `Integrate[Sin[x]/(3 + Cos[x])^2,x]`

output `(3 + Cos[x])^(-1)`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3147, 17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sin(x)}{(\cos(x) + 3)^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\cos\left(x - \frac{\pi}{2}\right)}{\left(3 - \sin\left(x - \frac{\pi}{2}\right)\right)^2} dx \\ & \quad \downarrow \text{3147} \\ & - \int \frac{1}{(\cos(x) + 3)^2} d\cos(x) \\ & \quad \downarrow \text{17} \\ & \frac{1}{\cos(x) + 3} \end{aligned}$$

input

```
Int[Sin[x]/(3 + Cos[x])^2,x]
```

output

```
(3 + Cos[x])^(-1)
```

Defintions of rubi rules used

rule 17

```
Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1)
)/(b*(m + 1))], x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```


rule 3147

```
Int[cos[(e_.) + (f_.)*(x_)]^(p_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Simp[1/(b^p*f) Subst[Int[(a + x)^m*(b^2 - x^2)^((p - 1)/2), x], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && IntegerQ[(p - 1)/2] && NeQ[a^2 - b^2, 0]
```

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.17

method	result	size
derivativdivides	$\frac{1}{3+\cos(x)}$	7
default	$\frac{1}{3+\cos(x)}$	7
parallelrisch	$-\frac{1}{2 \tan\left(\frac{x}{2}\right)^2 + 4}$	15
risch	$\frac{2e^{ix}}{e^{2ix} + 6e^{ix} + 1}$	24
norman	$\frac{-\frac{\tan\left(\frac{x}{2}\right)^2}{2} - \frac{1}{2}}{\left(1 + \tan\left(\frac{x}{2}\right)^2\right)\left(\tan\left(\frac{x}{2}\right)^2 + 2\right)}$	32

input

```
int(sin(x)/(3+cos(x))^2,x,method=_RETURNVERBOSE)
```

output

```
1/(3+cos(x))
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{\sin(x)}{(3 + \cos(x))^2} dx = \frac{1}{\cos(x) + 3}$$

input

```
integrate(sin(x)/(3+cos(x))^2,x, algorithm="fricas")
```

output

```
1/(cos(x) + 3)
```

Sympy [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int \frac{\sin(x)}{(3 + \cos(x))^2} dx = \frac{1}{\cos(x) + 3}$$

input `integrate(sin(x)/(3+cos(x))**2,x)`

output `1/(cos(x) + 3)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{\sin(x)}{(3 + \cos(x))^2} dx = \frac{1}{\cos(x) + 3}$$

input `integrate(sin(x)/(3+cos(x))^2,x, algorithm="maxima")`

output `1/(cos(x) + 3)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{\sin(x)}{(3 + \cos(x))^2} dx = \frac{1}{\cos(x) + 3}$$

input `integrate(sin(x)/(3+cos(x))^2,x, algorithm="giac")`

output `1/(cos(x) + 3)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{\sin(x)}{(3 + \cos(x))^2} dx = \frac{1}{\cos(x) + 3}$$

input `int(sin(x)/(cos(x) + 3)^2,x)`

output `1/(cos(x) + 3)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 12, normalized size of antiderivative = 2.00

$$\int \frac{\sin(x)}{(3 + \cos(x))^2} dx = -\frac{\cos(x)}{3 \cos(x) + 9}$$

input `int(sin(x)/(3+cos(x))^2,x)`

output `(- cos(x))/(3*(cos(x) + 3))`

3.11 $\int \frac{\sin(x)}{\sqrt{\cos^3(x)}} dx$

Optimal result	139
Mathematica [A] (verified)	139
Rubi [A] (verified)	140
Maple [A] (verified)	142
Fricas [A] (verification not implemented)	142
Sympy [A] (verification not implemented)	142
Maxima [A] (verification not implemented)	143
Giac [A] (verification not implemented)	143
Mupad [B] (verification not implemented)	143
Reduce [B] (verification not implemented)	144

Optimal result

Integrand size = 11, antiderivative size = 12

$$\int \frac{\sin(x)}{\sqrt{\cos^3(x)}} dx = \frac{2 \cos(x)}{\sqrt{\cos^3(x)}}$$

output `2*cos(x)/(cos(x)^3)^(1/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\sin(x)}{\sqrt{\cos^3(x)}} dx = \frac{2 \cos(x)}{\sqrt{\cos^3(x)}}$$

input `Integrate[Sin[x]/Sqrt[Cos[x]^3],x]`

output `(2*Cos[x])/Sqrt[Cos[x]^3]`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {3042, 25, 3686, 25, 3042, 3045, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin(x)}{\sqrt{\cos^3(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\cos\left(x + \frac{\pi}{2}\right)}{\sqrt{\sin\left(x + \frac{\pi}{2}\right)^3}} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{\cos\left(x + \frac{\pi}{2}\right)}{\sqrt{\sin\left(x + \frac{\pi}{2}\right)^3}} dx \\
 & \quad \downarrow \text{3686} \\
 & \frac{\cos^{\frac{3}{2}}(x) \int -\frac{\sin(x)}{\cos^{\frac{3}{2}}(x)} dx}{\sqrt{\cos^3(x)}} \\
 & \quad \downarrow \text{25} \\
 & \frac{\cos^{\frac{3}{2}}(x) \int \frac{\sin(x)}{\cos^{\frac{3}{2}}(x)} dx}{\sqrt{\cos^3(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\cos^{\frac{3}{2}}(x) \int \frac{\sin(x)}{\cos(x)^{3/2}} dx}{\sqrt{\cos^3(x)}} \\
 & \quad \downarrow \text{3045} \\
 & -\frac{\cos^{\frac{3}{2}}(x) \int \frac{1}{\cos^{\frac{3}{2}}(x)} d \cos(x)}{\sqrt{\cos^3(x)}} \\
 & \quad \downarrow \text{15}
 \end{aligned}$$

$$\frac{2 \cos(x)}{\sqrt{\cos^3(x)}}$$

input `Int[Sin[x]/Sqrt[Cos[x]^3], x]`

output `(2*Cos[x])/Sqrt[Cos[x]^3]`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :=> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 25 `Int[-(Fx_), x_Symbol] :=> Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3045 `Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] :=> Simp[-(a*f)^(-1) Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])`

rule 3686 `Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] :=> With[{ff = FreeFactors[Sin[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*SIn[e + f*x]^n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p])) Int[ActivateTrig[u]*(Sin[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

method	result	size
derivativedivides	$\frac{4 \cos(x)}{\sqrt{\cos(3x)+3 \cos(x)}}$	11
default	$\frac{4 \cos(x)}{\sqrt{\cos(3x)+3 \cos(x)}}$	11

input `int(sin(x)/(cos(x)^3)^(1/2),x,method=_RETURNVERBOSE)`output `2*cos(x)/(cos(x)^3)^(1/2)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\sin(x)}{\sqrt{\cos^3(x)}} dx = \frac{2 \sqrt{\cos(x)^3}}{\cos(x)^2}$$

input `integrate(sin(x)/(cos(x)^3)^(1/2),x, algorithm="fricas")`output `2*sqrt(cos(x)^3)/cos(x)^2`**Sympy [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\sin(x)}{\sqrt{\cos^3(x)}} dx = \frac{2 \cos(x)}{\sqrt{\cos^3(x)}}$$

input `integrate(sin(x)/(cos(x)**3)**(1/2),x)`output `2*cos(x)/sqrt(cos(x)**3)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\sin(x)}{\sqrt{\cos^3(x)}} dx = \frac{2 \cos(x)}{\sqrt{\cos(x)^3}}$$

input `integrate(sin(x)/(cos(x)^3)^(1/2),x, algorithm="maxima")`output `2*cos(x)/sqrt(cos(x)^3)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.50

$$\int \frac{\sin(x)}{\sqrt{\cos^3(x)}} dx = \frac{2}{\sqrt{\cos(x)}}$$

input `integrate(sin(x)/(cos(x)^3)^(1/2),x, algorithm="giac")`output `2/sqrt(cos(x))`**Mupad [B] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

$$\int \frac{\sin(x)}{\sqrt{\cos^3(x)}} dx = \frac{2 |\cos(x)|}{\cos(x)^{3/2}}$$

input `int(sin(x)/(cos(x)^3)^(1/2),x)`output `(2*abs(cos(x)))/cos(x)^(3/2)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

$$\int \frac{\sin(x)}{\sqrt{\cos^3(x)}} dx = \frac{2\sqrt{\cos(x)}}{\cos(x)}$$

input `int(sin(x)/(cos(x)^3)^(1/2),x)`

output `(2*sqrt(cos(x)))/cos(x)`

3.12 $\int \frac{\sin(\sqrt{1+x})}{\sqrt{1+x}} dx$

Optimal result	145
Mathematica [A] (verified)	145
Rubi [A] (verified)	146
Maple [A] (verified)	147
Fricas [A] (verification not implemented)	147
Sympy [A] (verification not implemented)	148
Maxima [A] (verification not implemented)	148
Giac [A] (verification not implemented)	148
Mupad [B] (verification not implemented)	149
Reduce [B] (verification not implemented)	149

Optimal result

Integrand size = 16, antiderivative size = 10

$$\int \frac{\sin(\sqrt{1+x})}{\sqrt{1+x}} dx = -2 \cos(\sqrt{1+x})$$

output `-2*cos((1+x)^(1/2))`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\sin(\sqrt{1+x})}{\sqrt{1+x}} dx = -2 \cos(\sqrt{1+x})$$

input `Integrate[Sin[Sqrt[1 + x]]/Sqrt[1 + x],x]`

output `-2*Cos[Sqrt[1 + x]]`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3912, 30, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sin(\sqrt{x+1})}{\sqrt{x+1}} dx$$

$$\downarrow \text{3912}$$

$$2 \int \sin(\sqrt{x+1}) d\sqrt{x+1}$$

$$\downarrow \text{3042}$$

$$2 \int \sin(\sqrt{x+1}) d\sqrt{x+1}$$

$$\downarrow \text{3118}$$

$$-2 \cos(\sqrt{x+1})$$

input `Int[Sin[Sqrt[1 + x]]/Sqrt[1 + x],x]`

output `-2*Cos[Sqrt[1 + x]]`

Defintions of rubi rules used

rule 30 `Int[(u_.)*((a_.)*(x_))^(m_.)*((b_.)*(x_)^(i_.))^(p_), x_Symbol] :> Simp[b^IntPart[p]*((b*x^i)^FracPart[p]/(a^(i*IntPart[p])*(a*x)^(i*FracPart[p])))`
`Int[u*(a*x)^(m + i*p), x], x] /; FreeQ[{a, b, i, m, p}, x] && IntegerQ[i] && !IntegerQ[p]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3912 `Int[((g_.) + (h_.)*(x_)^(m_.))*((a_.) + (b_.)*Sin[(c_.) + (d_.)*((e_.) + (f_.)*(x_)^(n_)])^(p_.), x_Symbol] := Simp[1/(n*f) Subst[Int[ExpandIntegrand[(a + b*SIN[c + d*x])^p, x^(1/n - 1)*(g - e*(h/f) + h*(x^(1/n)/f))^m, x], x, (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f, g, h, m}, x] && IGtQ[p, 0] && IntegerQ[1/n]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

method	result	size
derivativedivides	$-2 \cos(\sqrt{1+x})$	9
default	$-2 \cos(\sqrt{1+x})$	9

input `int(sin((1+x)^(1/2))/(1+x)^(1/2),x,method=_RETURNVERBOSE)`

output `-2*cos((1+x)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{\sin(\sqrt{1+x})}{\sqrt{1+x}} dx = -2 \cos(\sqrt{x+1})$$

input `integrate(sin((1+x)^(1/2))/(1+x)^(1/2),x, algorithm="fricas")`

output `-2*cos(sqrt(x + 1))`

Sympy [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\sin(\sqrt{1+x})}{\sqrt{1+x}} dx = -2 \cos(\sqrt{x+1})$$

input `integrate(sin((1+x)**(1/2))/(1+x)**(1/2),x)`

output `-2*cos(sqrt(x + 1))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{\sin(\sqrt{1+x})}{\sqrt{1+x}} dx = -2 \cos(\sqrt{x+1})$$

input `integrate(sin((1+x)^(1/2))/(1+x)^(1/2),x, algorithm="maxima")`

output `-2*cos(sqrt(x + 1))`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{\sin(\sqrt{1+x})}{\sqrt{1+x}} dx = -2 \cos(\sqrt{x+1})$$

input `integrate(sin((1+x)^(1/2))/(1+x)^(1/2),x, algorithm="giac")`

output `-2*cos(sqrt(x + 1))`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{\sin(\sqrt{1+x})}{\sqrt{1+x}} dx = -2 \cos(\sqrt{x+1})$$

input `int(sin((x + 1)^(1/2))/(x + 1)^(1/2),x)`

output `-2*cos((x + 1)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int \frac{\sin(\sqrt{1+x})}{\sqrt{1+x}} dx = -2 \cos(\sqrt{x+1})$$

input `int(sin((1+x)^(1/2))/(1+x)^(1/2),x)`

output `- 2*cos(sqrt(x + 1))`

3.13 $\int x^{-1+n} \sin(x^n) dx$

Optimal result	150
Mathematica [A] (verified)	150
Rubi [A] (verified)	151
Maple [A] (verified)	152
Fricas [A] (verification not implemented)	152
Sympy [A] (verification not implemented)	153
Maxima [A] (verification not implemented)	153
Giac [A] (verification not implemented)	153
Mupad [B] (verification not implemented)	154
Reduce [B] (verification not implemented)	154

Optimal result

Integrand size = 10, antiderivative size = 9

$$\int x^{-1+n} \sin(x^n) dx = -\frac{\cos(x^n)}{n}$$

output

```
-cos(x^n)/n
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int x^{-1+n} \sin(x^n) dx = -\frac{\cos(x^n)}{n}$$

input

```
Integrate[x^(-1 + n)*Sin[x^n],x]
```

output

```
-(Cos[x^n]/n)
```

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3860, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{n-1} \sin(x^n) dx$$

$$\downarrow \text{3860}$$

$$\frac{\int \sin(x^n) dx^n}{n}$$

$$\downarrow \text{3042}$$

$$\frac{\int \sin(x^n) dx^n}{n}$$

$$\downarrow \text{3118}$$

$$\frac{-\cos(x^n)}{n}$$

input `Int[x^(-1 + n)*Sin[x^n],x]`

output `-(Cos[x^n]/n)`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3860

```
Int[(x_)^(m_.)*((a_.) + (b_.)*Sin[(c_.) + (d_.)*(x_)^(n_)])^(p_.), x_Symbol]
:> Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Sin[c + d*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))
```

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.11

method	result	size
default	$-\frac{\cos(x^n)}{n}$	10
risch	$-\frac{\cos(x^n)}{n}$	10
norman	$\frac{2 \tan\left(\frac{e^n \ln(x)}{2}\right)^2}{n \left(1 + \tan\left(\frac{e^n \ln(x)}{2}\right)^2\right)}$	30
meijerg	$\frac{\sqrt{\pi} \left(2^{1-\frac{-1+n}{n}-\frac{1}{n}} (-1)^{\frac{1}{2}-\frac{-1+n}{2n}-\frac{1}{2n}} - (-1)^{\frac{1}{2}-\frac{-1+n}{2n}-\frac{1}{2n}} 2^{1-\frac{-1+n}{n}-\frac{1}{n}} \cos(x^n) \right)}{\sqrt{\pi} \Gamma\left(3-\frac{-1+n}{n}-\frac{1}{n}\right) - \sqrt{\pi} \Gamma\left(3-\frac{-1+n}{n}-\frac{1}{n}\right)}$	126

input

```
int(x^(-1+n)*sin(x^n),x,method=_RETURNVERBOSE)
```

output

```
-cos(x^n)/n
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int x^{-1+n} \sin(x^n) dx = -\frac{\cos(x^n)}{n}$$

input

```
integrate(x^(-1+n)*sin(x^n),x, algorithm="fricas")
```

output

```
-cos(x^n)/n
```

Sympy [A] (verification not implemented)

Time = 1.21 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int x^{-1+n} \sin(x^n) dx = -\frac{\cos(x^n)}{n}$$

input `integrate(x**(-1+n)*sin(x**n),x)`

output `-cos(x**n)/n`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int x^{-1+n} \sin(x^n) dx = -\frac{\cos(x^n)}{n}$$

input `integrate(x^(-1+n)*sin(x^n),x, algorithm="maxima")`

output `-cos(x^n)/n`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int x^{-1+n} \sin(x^n) dx = -\frac{\cos(x^n)}{n}$$

input `integrate(x^(-1+n)*sin(x^n),x, algorithm="giac")`

output `-cos(x^n)/n`

Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int x^{-1+n} \sin(x^n) dx = -\frac{\cos(x^n)}{n}$$

input `int(x^(n - 1)*sin(x^n),x)`

output `-cos(x^n)/n`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int x^{-1+n} \sin(x^n) dx = -\frac{\cos(x^n)}{n}$$

input `int(x^(-1+n)*sin(x^n),x)`

output `(- cos(x**n))/n`

3.14 $\int \frac{x^5}{\sqrt{1-x^6}} dx$

Optimal result	155
Mathematica [A] (verified)	155
Rubi [A] (verified)	156
Maple [A] (verified)	157
Fricas [A] (verification not implemented)	157
Sympy [A] (verification not implemented)	158
Maxima [A] (verification not implemented)	158
Giac [A] (verification not implemented)	158
Mupad [B] (verification not implemented)	159
Reduce [B] (verification not implemented)	159

Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{x^5}{\sqrt{1-x^6}} dx = -\frac{1}{3}\sqrt{1-x^6}$$

output `-1/3*(-x^6+1)^(1/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{x^5}{\sqrt{1-x^6}} dx = -\frac{1}{3}\sqrt{1-x^6}$$

input `Integrate[x^5/Sqrt[1 - x^6],x]`

output `-1/3*Sqrt[1 - x^6]`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$, Rules used = {793}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^5}{\sqrt{1-x^6}} dx$$

↓ 793

$$-\frac{1}{3}\sqrt{1-x^6}$$

input `Int[x^5/Sqrt[1 - x^6],x]`

output `-1/3*Sqrt[1 - x^6]`

Defintions of rubi rules used

rule 793 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(a + b*x^n)^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] && NeQ[p, -1]`

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

method	result	size
derivativedivides	$-\frac{\sqrt{-x^6+1}}{3}$	12
default	$-\frac{\sqrt{-x^6+1}}{3}$	12
trager	$-\frac{\sqrt{-x^6+1}}{3}$	12
pseudoelliptic	$-\frac{\sqrt{-x^6+1}}{3}$	12
risch	$\frac{x^6-1}{3\sqrt{-x^6+1}}$	17
meijerg	$-\frac{-2\sqrt{\pi}+2\sqrt{\pi}\sqrt{-x^6+1}}{6\sqrt{\pi}}$	26
gospers	$\frac{(-1+x)(1+x)(x^2+x+1)(x^2-x+1)}{3\sqrt{-x^6+1}}$	32
orering	$\frac{(-1+x)(1+x)(x^2+x+1)(x^2-x+1)}{3\sqrt{-x^6+1}}$	32

input `int(x^5/(-x^6+1)^(1/2),x,method=_RETURNVERBOSE)`output `-1/3*(-x^6+1)^(1/2)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \frac{x^5}{\sqrt{1-x^6}} dx = -\frac{1}{3} \sqrt{-x^6+1}$$

input `integrate(x^5/(-x^6+1)^(1/2),x, algorithm="fricas")`output `-1/3*sqrt(-x^6 + 1)`

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.67

$$\int \frac{x^5}{\sqrt{1-x^6}} dx = -\frac{\sqrt{1-x^6}}{3}$$

input `integrate(x**5/(-x**6+1)**(1/2),x)`output `-sqrt(1 - x**6)/3`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \frac{x^5}{\sqrt{1-x^6}} dx = -\frac{1}{3} \sqrt{-x^6 + 1}$$

input `integrate(x^5/(-x^6+1)^(1/2),x, algorithm="maxima")`output `-1/3*sqrt(-x^6 + 1)`**Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \frac{x^5}{\sqrt{1-x^6}} dx = -\frac{1}{3} \sqrt{-x^6 + 1}$$

input `integrate(x^5/(-x^6+1)^(1/2),x, algorithm="giac")`output `-1/3*sqrt(-x^6 + 1)`

Mupad [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \frac{x^5}{\sqrt{1-x^6}} dx = -\frac{\sqrt{1-x^6}}{3}$$

input `int(x^5/(1 - x^6)^(1/2),x)`

output `-(1 - x^6)^(1/2)/3`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.67

$$\int \frac{x^5}{\sqrt{1-x^6}} dx = -\frac{\sqrt{-x^6+1}}{3}$$

input `int(x^5/(-x^6+1)^(1/2),x)`

output `(- sqrt(- x**6 + 1))/3`

3.15 $\int t\sqrt[4]{1+t} dt$

Optimal result	160
Mathematica [A] (verified)	160
Rubi [A] (verified)	161
Maple [A] (verified)	162
Fricas [A] (verification not implemented)	162
Sympy [A] (verification not implemented)	163
Maxima [A] (verification not implemented)	163
Giac [A] (verification not implemented)	163
Mupad [B] (verification not implemented)	164
Reduce [B] (verification not implemented)	164

Optimal result

Integrand size = 9, antiderivative size = 23

$$\int t\sqrt[4]{1+t} dt = -\frac{4}{5}(1+t)^{5/4} + \frac{4}{9}(1+t)^{9/4}$$

output `-4/5*(1+t)^(5/4)+4/9*(1+t)^(9/4)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

$$\int t\sqrt[4]{1+t} dt = \frac{4}{45}(1+t)^{5/4}(-9+5(1+t))$$

input `Integrate[t*(1+t)^(1/4),t]`

output `(4*(1+t)^(5/4)*(-9+5*(1+t)))/45`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int t\sqrt[4]{t+1} dt$$

$$\downarrow 53$$

$$\int \left((t+1)^{5/4} - \sqrt[4]{t+1} \right) dt$$

$$\downarrow 2009$$

$$\frac{4}{9}(t+1)^{9/4} - \frac{4}{5}(t+1)^{5/4}$$

input `Int[t*(1 + t)^(1/4),t]`

output `(-4*(1 + t)^(5/4))/5 + (4*(1 + t)^(9/4))/9`

Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.57

method	result	size
gospers	$\frac{4(1+t)^{\frac{5}{4}}(5t-4)}{45}$	13
orering	$\frac{4(1+t)^{\frac{5}{4}}(5t-4)}{45}$	13
meijerg	$\frac{t^2 \operatorname{hypergeom}\left(\left[-\frac{1}{4}, 2\right], [3], -t\right)}{2}$	15
derivativedivides	$-\frac{4(1+t)^{\frac{5}{4}}}{5} + \frac{4(1+t)^{\frac{9}{4}}}{9}$	16
default	$-\frac{4(1+t)^{\frac{5}{4}}}{5} + \frac{4(1+t)^{\frac{9}{4}}}{9}$	16
risch	$\frac{4(1+t)^{\frac{1}{4}}(5t^2+t-4)}{45}$	16
trager	$\left(\frac{4}{9}t^2 + \frac{4}{45}t - \frac{16}{45}\right)(1+t)^{\frac{1}{4}}$	17

input `int(t*(1+t)^(1/4),t,method=_RETURNVERBOSE)`output `4/45*(1+t)^(5/4)*(5*t-4)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int t\sqrt[4]{1+t} dt = \frac{4}{45} (5t^2 + t - 4)(t+1)^{\frac{1}{4}}$$

input `integrate(t*(1+t)^(1/4),t, algorithm="fricas")`output `4/45*(5*t^2 + t - 4)*(t + 1)^(1/4)`

Sympy [A] (verification not implemented)

Time = 0.54 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.48

$$\int t\sqrt[4]{1+t} dt = \frac{4t^2\sqrt[4]{t+1}}{9} + \frac{4t\sqrt[4]{t+1}}{45} - \frac{16\sqrt[4]{t+1}}{45}$$

input `integrate(t*(1+t)**(1/4),t)`output `4*t**2*(t + 1)**(1/4)/9 + 4*t*(t + 1)**(1/4)/45 - 16*(t + 1)**(1/4)/45`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int t\sqrt[4]{1+t} dt = \frac{4}{9}(t+1)^{\frac{9}{4}} - \frac{4}{5}(t+1)^{\frac{5}{4}}$$

input `integrate(t*(1+t)^(1/4),t, algorithm="maxima")`output `4/9*(t + 1)^(9/4) - 4/5*(t + 1)^(5/4)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int t\sqrt[4]{1+t} dt = \frac{4}{9}(t+1)^{\frac{9}{4}} - \frac{4}{5}(t+1)^{\frac{5}{4}}$$

input `integrate(t*(1+t)^(1/4),t, algorithm="giac")`output `4/9*(t + 1)^(9/4) - 4/5*(t + 1)^(5/4)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.52

$$\int t\sqrt[4]{1+t} dt = \frac{4(5t-4)(t+1)^{5/4}}{45}$$

input `int(t*(t + 1)^(1/4),t)`

output `(4*(5*t - 4)*(t + 1)^(5/4))/45`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int t\sqrt[4]{1+t} dt = \frac{4(t+1)^{1/4}(5t^2+t-4)}{45}$$

input `int(t*(1+t)^(1/4),t)`

output `(4*(t + 1)**(1/4)*(5*t**2 + t - 4))/45`

3.16 $\int \frac{1}{(1+x^2)^{3/2}} dx$

Optimal result	165
Mathematica [A] (verified)	165
Rubi [A] (verified)	166
Maple [A] (verified)	167
Fricas [B] (verification not implemented)	167
Sympy [A] (verification not implemented)	168
Maxima [A] (verification not implemented)	168
Giac [A] (verification not implemented)	168
Mupad [B] (verification not implemented)	169
Reduce [B] (verification not implemented)	169

Optimal result

Integrand size = 9, antiderivative size = 11

$$\int \frac{1}{(1+x^2)^{3/2}} dx = \frac{x}{\sqrt{1+x^2}}$$

output `x/(x^2+1)^(1/2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1+x^2)^{3/2}} dx = \frac{x}{\sqrt{1+x^2}}$$

input `Integrate[(1 + x^2)^(-3/2), x]`

output `x/Sqrt[1 + x^2]`

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {208}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(x^2 + 1)^{3/2}} dx$$

$$\downarrow 208$$

$$\frac{x}{\sqrt{x^2 + 1}}$$

input

```
Int[(1 + x^2)^(-3/2), x]
```

output

```
x/Sqrt[1 + x^2]
```

Defintions of rubi rules used

rule 208

```
Int[((a_) + (b_.)*(x_)^2)^(-3/2), x_Symbol] :> Simp[x/(a*Sqrt[a + b*x^2]), x] /; FreeQ[{a, b}, x]
```

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

method	result	size
gospers	$\frac{x}{\sqrt{x^2+1}}$	10
default	$\frac{x}{\sqrt{x^2+1}}$	10
trager	$\frac{x}{\sqrt{x^2+1}}$	10
meijerg	$\frac{x}{\sqrt{x^2+1}}$	10
risch	$\frac{x}{\sqrt{x^2+1}}$	10
pseudoelliptic	$\frac{x}{\sqrt{x^2+1}}$	10
orering	$\frac{x}{\sqrt{x^2+1}}$	10

input `int(1/(x^2+1)^(3/2),x,method=_RETURNVERBOSE)`

output `x/(x^2+1)^(1/2)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 22 vs. 2(9) = 18.

Time = 0.09 (sec) , antiderivative size = 22, normalized size of antiderivative = 2.00

$$\int \frac{1}{(1+x^2)^{3/2}} dx = \frac{x^2 + \sqrt{x^2+1}x + 1}{x^2 + 1}$$

input `integrate(1/(x^2+1)^(3/2),x, algorithm="fricas")`

output `(x^2 + sqrt(x^2 + 1)*x + 1)/(x^2 + 1)`

Sympy [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int \frac{1}{(1+x^2)^{3/2}} dx = \frac{x}{\sqrt{x^2+1}}$$

input `integrate(1/(x**2+1)**(3/2),x)`

output `x/sqrt(x**2 + 1)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \frac{1}{(1+x^2)^{3/2}} dx = \frac{x}{\sqrt{x^2+1}}$$

input `integrate(1/(x^2+1)^(3/2),x, algorithm="maxima")`

output `x/sqrt(x^2 + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \frac{1}{(1+x^2)^{3/2}} dx = \frac{x}{\sqrt{x^2+1}}$$

input `integrate(1/(x^2+1)^(3/2),x, algorithm="giac")`

output `x/sqrt(x^2 + 1)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \frac{1}{(1+x^2)^{3/2}} dx = \frac{x}{\sqrt{x^2+1}}$$

input `int(1/(x^2 + 1)^(3/2),x)`

output `x/(x^2 + 1)^(1/2)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.91

$$\int \frac{1}{(1+x^2)^{3/2}} dx = \frac{\sqrt{x^2+1}x + x^2 + 1}{x^2 + 1}$$

input `int(1/(x^2+1)^(3/2),x)`

output `(sqrt(x**2 + 1)*x + x**2 + 1)/(x**2 + 1)`

3.17 $\int x^2(27 + 8x^3)^{2/3} dx$

Optimal result	170
Mathematica [A] (verified)	170
Rubi [A] (verified)	171
Maple [A] (verified)	172
Fricas [A] (verification not implemented)	172
Sympy [B] (verification not implemented)	173
Maxima [A] (verification not implemented)	173
Giac [A] (verification not implemented)	173
Mupad [B] (verification not implemented)	174
Reduce [B] (verification not implemented)	174

Optimal result

Integrand size = 15, antiderivative size = 15

$$\int x^2(27 + 8x^3)^{2/3} dx = \frac{1}{40}(27 + 8x^3)^{5/3}$$

output

```
1/40*(8*x^3+27)^(5/3)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int x^2(27 + 8x^3)^{2/3} dx = \frac{1}{40}(27 + 8x^3)^{5/3}$$

input

```
Integrate[x^2*(27 + 8*x^3)^(2/3),x]
```

output

```
(27 + 8*x^3)^(5/3)/40
```

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$, Rules used = {793}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 (8x^3 + 27)^{2/3} dx$$

$$\downarrow 793$$

$$\frac{1}{40} (8x^3 + 27)^{5/3}$$

input `Int[x^2*(27 + 8*x^3)^(2/3),x]`

output `(27 + 8*x^3)^(5/3)/40`

Defintions of rubi rules used

rule 793 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a + b*x^n)^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] && NeQ[p, -1]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

method	result	size
derivativedivides	$\frac{(8x^3+27)^{\frac{5}{3}}}{40}$	12
default	$\frac{(8x^3+27)^{\frac{5}{3}}}{40}$	12
risch	$\frac{(8x^3+27)^{\frac{5}{3}}}{40}$	12
pseudoelliptic	$\frac{(8x^3+27)^{\frac{5}{3}}}{40}$	12
meijerg	$3x^3 \text{ hypergeom} \left(\left[-\frac{2}{3}, 1 \right], [2], -\frac{8x^3}{27} \right)$	17
trager	$\left(\frac{x^3}{5} + \frac{27}{40} \right) (8x^3 + 27)^{\frac{2}{3}}$	18
gosper	$\frac{(3+2x)(4x^2-6x+9)(8x^3+27)^{\frac{2}{3}}}{40}$	27
orering	$\frac{(3+2x)(4x^2-6x+9)(8x^3+27)^{\frac{2}{3}}}{40}$	27

input `int(x^2*(8*x^3+27)^(2/3),x,method=_RETURNVERBOSE)`output `1/40*(8*x^3+27)^(5/3)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int x^2(27 + 8x^3)^{2/3} dx = \frac{1}{40} (8x^3 + 27)^{\frac{5}{3}}$$

input `integrate(x^2*(8*x^3+27)^(2/3),x, algorithm="fricas")`output `1/40*(8*x^3 + 27)^(5/3)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. $2(10) = 20$.

Time = 0.09 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.80

$$\int x^2(27 + 8x^3)^{2/3} dx = \frac{x^3(8x^3 + 27)^{2/3}}{5} + \frac{27(8x^3 + 27)^{2/3}}{40}$$

input `integrate(x**2*(8*x**3+27)**(2/3),x)`

output `x**3*(8*x**3 + 27)**(2/3)/5 + 27*(8*x**3 + 27)**(2/3)/40`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int x^2(27 + 8x^3)^{2/3} dx = \frac{1}{40} (8x^3 + 27)^{5/3}$$

input `integrate(x^2*(8*x^3+27)^(2/3),x, algorithm="maxima")`

output `1/40*(8*x^3 + 27)^(5/3)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int x^2(27 + 8x^3)^{2/3} dx = \frac{1}{40} (8x^3 + 27)^{5/3}$$

input `integrate(x^2*(8*x^3+27)^(2/3),x, algorithm="giac")`

output `1/40*(8*x^3 + 27)^(5/3)`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int x^2(27 + 8x^3)^{2/3} dx = \frac{(8x^3 + 27)^{5/3}}{40}$$

input `int(x^2*(8*x^3 + 27)^(2/3),x)`

output `(8*x^3 + 27)^(5/3)/40`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int x^2(27 + 8x^3)^{2/3} dx = \frac{(8x^3 + 27)^{5/3}}{40}$$

input `int(x^2*(8*x^3+27)^(2/3),x)`

output `((8*x**3 + 27)**(2/3)*(8*x**3 + 27))/40`

$$3.18 \quad \int \frac{\cos(x) + \sin(x)}{\sqrt[3]{-\cos(x) + \sin(x)}} dx$$

Optimal result	175
Mathematica [A] (verified)	175
Rubi [A] (verified)	176
Maple [A] (verified)	177
Fricas [A] (verification not implemented)	177
Sympy [A] (verification not implemented)	177
Maxima [A] (verification not implemented)	178
Giac [A] (verification not implemented)	178
Mupad [B] (verification not implemented)	178
Reduce [F]	179

Optimal result

Integrand size = 17, antiderivative size = 15

$$\int \frac{\cos(x) + \sin(x)}{\sqrt[3]{-\cos(x) + \sin(x)}} dx = \frac{3}{2}(-\cos(x) + \sin(x))^{2/3}$$

output

```
3/2*(-cos(x)+sin(x))^(2/3)
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x) + \sin(x)}{\sqrt[3]{-\cos(x) + \sin(x)}} dx = \frac{3}{2}(-\cos(x) + \sin(x))^{2/3}$$

input

```
Integrate[(Cos[x] + Sin[x])/(-Cos[x] + Sin[x])^(1/3), x]
```

output

```
(3*(-Cos[x] + Sin[x])^(2/3))/2
```


Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {3042, 3624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sin(x) + \cos(x)}{\sqrt[3]{\sin(x) - \cos(x)}} dx$$

↓ 3042

$$\int \frac{\sin(x) + \cos(x)}{\sqrt[3]{\sin(x) - \cos(x)}} dx$$

↓ 3624

$$\frac{3}{2}(\sin(x) - \cos(x))^{2/3}$$

input `Int[(Cos[x] + Sin[x])/(-Cos[x] + Sin[x])^(1/3),x]`

output `(3*(-Cos[x] + Sin[x])^(2/3))/2`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3624 `Int[(cos[(d_.) + (e_.)*(x_)]*(b_.) + (c_.)*sin[(d_.) + (e_.)*(x_)])^(n_.)*(cos[(d_.) + (e_.)*(x_)]*(B_.) + (C_.)*sin[(d_.) + (e_.)*(x_)]), x_Symbol] :> Simp[(c*B - b*C)*((b*Cos[d + e*x] + c*Sin[d + e*x])^(n + 1)/(e*(n + 1)*(b^2 + c^2))), x] /; FreeQ[{b, c, d, e, B, C}, x] && NeQ[n, -1] && NeQ[b^2 + c^2, 0] && EqQ[b*B + c*C, 0]`

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

method	result	size
derivativedivides	$\frac{3(-\cos(x)+\sin(x))^{2/3}}{2}$	12
default	$\frac{3(-\cos(x)+\sin(x))^{2/3}}{2}$	12
risch	$\frac{(-\frac{3}{2}-\frac{3i}{2})((1+i)(-e^{4ix}+ie^{2ix}))^{1/3}(e^{ix}-ie^{-ix})}{(-8\cos(x)+8\sin(x))^{1/3}((-1-i)(e^{4ix}-ie^{2ix}))^{1/3}}$	72

input `int((cos(x)+sin(x))/(-cos(x)+sin(x))^(1/3),x,method=_RETURNVERBOSE)`

output `3/2*(-cos(x)+sin(x))^(2/3)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \frac{\cos(x) + \sin(x)}{\sqrt[3]{-\cos(x) + \sin(x)}} dx = \frac{3}{2} (-\cos(x) + \sin(x))^{2/3}$$

input `integrate((cos(x)+sin(x))/(-cos(x)+sin(x))^(1/3),x, algorithm="fricas")`

output `3/2*(-cos(x) + sin(x))^(2/3)`

Sympy [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

$$\int \frac{\cos(x) + \sin(x)}{\sqrt[3]{-\cos(x) + \sin(x)}} dx = \frac{3(\sin(x) - \cos(x))^{2/3}}{2}$$

input `integrate((cos(x)+sin(x))/(-cos(x)+sin(x))**(1/3),x)`

output `3*(sin(x) - cos(x))**(2/3)/2`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \frac{\cos(x) + \sin(x)}{\sqrt[3]{-\cos(x) + \sin(x)}} dx = \frac{3}{2} (-\cos(x) + \sin(x))^{\frac{2}{3}}$$

input `integrate((cos(x)+sin(x))/(-cos(x)+sin(x))^(1/3),x, algorithm="maxima")`

output `3/2*(-cos(x) + sin(x))^(2/3)`

Giac [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \frac{\cos(x) + \sin(x)}{\sqrt[3]{-\cos(x) + \sin(x)}} dx = \frac{3}{2} (-\cos(x) + \sin(x))^{\frac{2}{3}}$$

input `integrate((cos(x)+sin(x))/(-cos(x)+sin(x))^(1/3),x, algorithm="giac")`

output `3/2*(-cos(x) + sin(x))^(2/3)`

Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x) + \sin(x)}{\sqrt[3]{-\cos(x) + \sin(x)}} dx = \frac{3 \cdot 2^{1/3} (-\cos(x + \frac{\pi}{4}))^{2/3}}{2}$$

input `int((cos(x) + sin(x))/(sin(x) - cos(x))^(1/3),x)`

output `(3*2^(1/3)*(-cos(x + pi/4))^(2/3))/2`

Reduce [F]

$$\int \frac{\cos(x) + \sin(x)}{\sqrt[3]{-\cos(x) + \sin(x)}} dx = \int \frac{\cos(x)}{(-\cos(x) + \sin(x))^{\frac{1}{3}}} dx + \int \frac{\sin(x)}{(-\cos(x) + \sin(x))^{\frac{1}{3}}} dx$$

input `int((cos(x)+sin(x))/(-cos(x)+sin(x))^(1/3),x)`

output `int(cos(x)/(-cos(x)+sin(x))**(1/3),x) + int(sin(x)/(-cos(x)+sin(x))**(1/3),x)`

$$3.19 \quad \int \frac{x}{\sqrt{1+x^2+(1+x^2)^{3/2}}} dx$$

Optimal result	180
Mathematica [A] (verified)	180
Rubi [A] (verified)	181
Maple [F]	182
Fricas [A] (verification not implemented)	182
Sympy [F]	183
Maxima [F]	183
Giac [A] (verification not implemented)	183
Mupad [B] (verification not implemented)	184
Reduce [B] (verification not implemented)	184

Optimal result

Integrand size = 20, antiderivative size = 32

$$\int \frac{x}{\sqrt{1+x^2+(1+x^2)^{3/2}}} dx = \frac{2\sqrt{(1+x^2)(1+\sqrt{1+x^2})}}{\sqrt{1+x^2}}$$

output `2*((x^2+1)*(1+(x^2+1)^(1/2)))^(1/2)/(x^2+1)^(1/2)`

Mathematica [A] (verified)

Time = 1.62 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

$$\int \frac{x}{\sqrt{1+x^2+(1+x^2)^{3/2}}} dx = \frac{2\sqrt{(1+x^2)(1+\sqrt{1+x^2})}}{\sqrt{1+x^2}}$$

input `Integrate[x/Sqrt[1 + x^2 + (1 + x^2)^(3/2)],x]`

output `(2*Sqrt[(1 + x^2)*(1 + Sqrt[1 + x^2])])/Sqrt[1 + x^2]`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.75, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {7266, 7267, 2021}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\sqrt{x^2 + (x^2 + 1)^{3/2} + 1}} dx$$

$$\downarrow 7266$$

$$\frac{1}{2} \int \frac{1}{\sqrt{x^2 + (x^2 + 1)^{3/2} + 1}} dx^2$$

$$\downarrow 7267$$

$$\int \frac{\sqrt{x^2 + 1}}{\sqrt{x^4 (\sqrt{x^2 + 1} + 1)}} d\sqrt{x^2 + 1}$$

$$\downarrow 2021$$

$$\frac{2\sqrt{x^4 (\sqrt{x^2 + 1} + 1)}}{x^2}$$

input `Int[x/Sqrt[1 + x^2 + (1 + x^2)^(3/2)],x]`

output `(2*Sqrt[x^4*(1 + Sqrt[1 + x^2])])/x^2`

Defintions of rubi rules used

rule 2021

```
Int[(Pp_)*(Qq_)^(m_.), x_Symbol] := With[{p = Expon[Pp, x], q = Expon[Qq, x]
}], Simp[Coeff[Pp, x, p]*x^(p - q + 1)*(Qq^(m + 1)/((p + m*q + 1)*Coeff[Qq,
x, q])), x] /; NeQ[p + m*q + 1, 0] && EqQ[(p + m*q + 1)*Coeff[Qq, x, q]*Pp
, Coeff[Pp, x, p]*x^(p - q)*((p - q + 1)*Qq + (m + 1)*x*D[Qq, x])] /; Free
Q[m, x] && PolyQ[Pp, x] && PolyQ[Qq, x] && NeQ[m, -1]
```

rule 7266

```
Int[(u_)*(x_)^(m_.), x_Symbol] := Simp[1/(m + 1) Subst[Int[SubstFor[x^(m
+ 1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && Function
OfQ[x^(m + 1), u, x]
```

rule 7267

```
Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Si
mp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])]], x
] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]
```

Maple [F]

$$\int \frac{x}{\sqrt{1+x^2+(x^2+1)^{\frac{3}{2}}}} dx$$

input

```
int(x/(1+x^2+(x^2+1)^(3/2))^(1/2),x)
```

output

```
int(x/(1+x^2+(x^2+1)^(3/2))^(1/2),x)
```

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.72

$$\int \frac{x}{\sqrt{1+x^2+(1+x^2)^{3/2}}} dx = \frac{2\sqrt{x^2+(x^2+1)^{\frac{3}{2}}+1}}{\sqrt{x^2+1}}$$

input

```
integrate(x/(1+x^2+(x^2+1)^(3/2))^(1/2),x, algorithm="fricas")
```

output `2*sqrt(x^2 + (x^2 + 1)^(3/2) + 1)/sqrt(x^2 + 1)`

Sympy [F]

$$\int \frac{x}{\sqrt{1+x^2+(1+x^2)^{3/2}}} dx = \int \frac{x}{\sqrt{(x^2+1)(\sqrt{x^2+1}+1)}} dx$$

input `integrate(x/(1+x**2+(x**2+1)**(3/2))**(1/2),x)`

output `Integral(x/sqrt((x**2 + 1)*(sqrt(x**2 + 1) + 1)), x)`

Maxima [F]

$$\int \frac{x}{\sqrt{1+x^2+(1+x^2)^{3/2}}} dx = \int \frac{x}{\sqrt{x^2+(x^2+1)^{\frac{3}{2}}+1}} dx$$

input `integrate(x/(1+x^2+(x^2+1)^(3/2))^(1/2),x, algorithm="maxima")`

output `integrate(x/sqrt(x^2 + (x^2 + 1)^(3/2) + 1), x)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.47

$$\int \frac{x}{\sqrt{1+x^2+(1+x^2)^{3/2}}} dx = 2\sqrt{\sqrt{x^2+1}+1} - 2$$

input `integrate(x/(1+x^2+(x^2+1)^(3/2))^(1/2),x, algorithm="giac")`

output `2*sqrt(sqrt(x^2 + 1) + 1) - 2`

Mupad [B] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.47

$$\int \frac{x}{\sqrt{1+x^2+(1+x^2)^{3/2}}} dx = \frac{2(x^2+1)\sqrt{\sqrt{x^2+1}+1}}{\left(\sqrt{\sqrt{x^2+1}+1}+1\right)\sqrt{(x^2+1)^{3/2}+x^2+1}}$$

input `int(x/((x^2 + 1)^(3/2) + x^2 + 1)^(1/2),x)`output `(2*(x^2 + 1)*((x^2 + 1)^(1/2) + 1)^(1/2))/(((x^2 + 1)^(1/2) + 1)^(1/2) + 1)*((x^2 + 1)^(3/2) + x^2 + 1)^(1/2)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.31

$$\int \frac{x}{\sqrt{1+x^2+(1+x^2)^{3/2}}} dx = 2\sqrt{\sqrt{x^2+1}x + \sqrt{x^2+1} + x^2 + x + 1}\sqrt{\sqrt{x^2+1} + x}(\sqrt{x^2+1} - x)$$

input `int(x/(1+x^2+(x^2+1)^(3/2))^(1/2),x)`output `2*sqrt(sqrt(x**2 + 1)*x + sqrt(x**2 + 1) + x**2 + x + 1)*sqrt(sqrt(x**2 + 1) + x)*(sqrt(x**2 + 1) - x)`

$$3.20 \quad \int \frac{x}{\sqrt{1+x^2}\sqrt{1+\sqrt{1+x^2}}} dx$$

Optimal result	185
Mathematica [A] (verified)	185
Rubi [A] (verified)	186
Maple [A] (verified)	186
Fricas [A] (verification not implemented)	187
Sympy [A] (verification not implemented)	187
Maxima [A] (verification not implemented)	187
Giac [A] (verification not implemented)	188
Mupad [B] (verification not implemented)	188
Reduce [B] (verification not implemented)	188

Optimal result

Integrand size = 26, antiderivative size = 17

$$\int \frac{x}{\sqrt{1+x^2}\sqrt{1+\sqrt{1+x^2}}} dx = 2\sqrt{1+\sqrt{1+x^2}}$$

output

```
2*(1+(x^2+1)^(1/2))^(1/2)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{x}{\sqrt{1+x^2}\sqrt{1+\sqrt{1+x^2}}} dx = 2\sqrt{1+\sqrt{1+x^2}}$$

input

```
Integrate[x/(Sqrt[1 + x^2]*Sqrt[1 + Sqrt[1 + x^2]]),x]
```

output

```
2*Sqrt[1 + Sqrt[1 + x^2]]
```

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.038$, Rules used = {7237}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\sqrt{x^2+1}\sqrt{\sqrt{x^2+1}+1}} dx$$

↓ 7237

$$2\sqrt{\sqrt{x^2+1}+1}$$

input `Int[x/(Sqrt[1 + x^2]*Sqrt[1 + Sqrt[1 + x^2]]),x]`

output `2*Sqrt[1 + Sqrt[1 + x^2]]`

Defintions of rubi rules used

rule 7237 `Int[(u_)*(y_)^(m_.), x_Symbol] := With[{q = DerivativeDivides[y, u, x]}, Simp[q*(y^(m + 1)/(m + 1)), x] /; !FalseQ[q] /; FreeQ[m, x] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$2\sqrt{1 + \sqrt{x^2 + 1}}$	14
default	$2\sqrt{1 + \sqrt{x^2 + 1}}$	14

input `int(x/(x^2+1)^(1/2)/(1+(x^2+1)^(1/2))^(1/2),x,method=_RETURNVERBOSE)`

output `2*(1+(x^2+1)^(1/2))^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \frac{x}{\sqrt{1+x^2}\sqrt{1+\sqrt{1+x^2}}} dx = 2\sqrt{\sqrt{x^2+1}+1}$$

input `integrate(x/(x^2+1)^(1/2)/(1+(x^2+1)^(1/2))^(1/2),x, algorithm="fricas")`

output `2*sqrt(sqrt(x^2 + 1) + 1)`

Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \frac{x}{\sqrt{1+x^2}\sqrt{1+\sqrt{1+x^2}}} dx = 2\sqrt{\sqrt{x^2+1}+1}$$

input `integrate(x/(x**2+1)**(1/2)/(1+(x**2+1)**(1/2))**(1/2),x)`

output `2*sqrt(sqrt(x**2 + 1) + 1)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \frac{x}{\sqrt{1+x^2}\sqrt{1+\sqrt{1+x^2}}} dx = 2\sqrt{\sqrt{x^2+1}+1}$$

input `integrate(x/(x^2+1)^(1/2)/(1+(x^2+1)^(1/2))^(1/2),x, algorithm="maxima")`

output `2*sqrt(sqrt(x^2 + 1) + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \frac{x}{\sqrt{1+x^2}\sqrt{1+\sqrt{1+x^2}}} dx = 2\sqrt{\sqrt{x^2+1}+1}$$

input `integrate(x/(x^2+1)^(1/2)/(1+(x^2+1)^(1/2))^(1/2),x, algorithm="giac")`

output `2*sqrt(sqrt(x^2 + 1) + 1)`

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \frac{x}{\sqrt{1+x^2}\sqrt{1+\sqrt{1+x^2}}} dx = 2\sqrt{\sqrt{x^2+1}+1}$$

input `int(x/((x^2 + 1)^(1/2)*((x^2 + 1)^(1/2) + 1)^(1/2)),x)`

output `2*((x^2 + 1)^(1/2) + 1)^(1/2)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 42, normalized size of antiderivative = 2.47

$$\begin{aligned} & \int \frac{x}{\sqrt{1+x^2}\sqrt{1+\sqrt{1+x^2}}} dx \\ &= 2\sqrt{\sqrt{x^2+1}x + \sqrt{x^2+1} + x^2 + x + 1}\sqrt{\sqrt{x^2+1} + x}(\sqrt{x^2+1} - x) \end{aligned}$$

input `int(x/(x^2+1)^(1/2)/(1+(x^2+1)^(1/2))^(1/2),x)`

output `2*sqrt(sqrt(x**2 + 1)*x + sqrt(x**2 + 1) + x**2 + x + 1)*sqrt(sqrt(x**2 + 1) + x)*(sqrt(x**2 + 1) - x)`

$$3.21 \quad \int \frac{\sqrt[5]{1-2x+x^2}}{1-x} dx$$

Optimal result	190
Mathematica [A] (verified)	190
Rubi [A] (verified)	191
Maple [A] (verified)	191
Fricas [A] (verification not implemented)	192
Sympy [A] (verification not implemented)	193
Maxima [A] (verification not implemented)	193
Giac [A] (verification not implemented)	193
Mupad [B] (verification not implemented)	194
Reduce [B] (verification not implemented)	194

Optimal result

Integrand size = 20, antiderivative size = 16

$$\int \frac{\sqrt[5]{1-2x+x^2}}{1-x} dx = -\frac{5}{2} \sqrt[5]{1-2x+x^2}$$

output `-5/2*(x^2-2*x+1)^(1/5)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.81

$$\int \frac{\sqrt[5]{1-2x+x^2}}{1-x} dx = -\frac{5}{2} \sqrt[5]{(-1+x)^2}$$

input `Integrate[(1 - 2*x + x^2)^(1/5)/(1 - x), x]`

output `(-5*((-1 + x)^2)^(1/5))/2`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$, Rules used = {1099}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt[5]{x^2 - 2x + 1}}{1 - x} dx$$

↓ 1099

$$-\frac{5}{2} \sqrt[5]{x^2 - 2x + 1}$$

input `Int[(1 - 2*x + x^2)^(1/5)/(1 - x),x]`

output `(-5*(1 - 2*x + x^2)^(1/5))/2`

Defintions of rubi rules used

rule 1099

```
Int[((d_) + (e_.)*(x_)^(m_.))*((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Sy
mbol] := Simp[e^m*((a + b*x + c*x^2)^(p + (m + 1)/2)/(c^((m + 1)/2)*(m + 2*
p + 1))), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[b^2 - 4*a*c, 0] && EqQ[
2*c*d - b*e, 0] && IntegerQ[(m - 1)/2]
```

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.62

method	result	size
risch	$-\frac{5((-1+x)^2)^{\frac{1}{5}}}{2}$	10
pseudoelliptic	$-\frac{5((-1+x)^2)^{\frac{1}{5}}}{2}$	10
gosper	$-\frac{5(x^2-2x+1)^{\frac{1}{5}}}{2}$	13
trager	$-\frac{5(x^2-2x+1)^{\frac{1}{5}}}{2}$	13
meijerg	$\frac{\text{signum}(-1+x)^{\frac{2}{5}} x \text{ hypergeom}([\frac{3}{5}, 1], [2], x)}{(-\text{signum}(-1+x))^{\frac{2}{5}}}$	24
orering	$\frac{(-\frac{5}{2} + \frac{5x}{2})(x^2-2x+1)^{\frac{1}{5}}}{1-x}$	24

input `int((x^2-2*x+1)^(1/5)/(1-x),x,method=_RETURNVERBOSE)`

output `-5/2*((-1+x)^2)^(1/5)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt[5]{1-2x+x^2}}{1-x} dx = -\frac{5}{2} (x^2 - 2x + 1)^{\frac{1}{5}}$$

input `integrate((x^2-2*x+1)^(1/5)/(1-x),x, algorithm="fricas")`

output `-5/2*(x^2 - 2*x + 1)^(1/5)`

Sympy [A] (verification not implemented)

Time = 0.50 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{\sqrt[5]{1-2x+x^2}}{1-x} dx = -\frac{5\sqrt[5]{x^2-2x+1}}{2}$$

input `integrate((x**2-2*x+1)**(1/5)/(1-x),x)`output `-5*(x**2 - 2*x + 1)**(1/5)/2`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.44

$$\int \frac{\sqrt[5]{1-2x+x^2}}{1-x} dx = -\frac{5}{2}(x-1)^{\frac{2}{5}}$$

input `integrate((x^2-2*x+1)^(1/5)/(1-x),x, algorithm="maxima")`output `-5/2*(x - 1)^(2/5)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt[5]{1-2x+x^2}}{1-x} dx = -\frac{5}{2}(x^2-2x+1)^{\frac{1}{5}}$$

input `integrate((x^2-2*x+1)^(1/5)/(1-x),x, algorithm="giac")`output `-5/2*(x^2 - 2*x + 1)^(1/5)`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.56

$$\int \frac{\sqrt[5]{1-2x+x^2}}{1-x} dx = -\frac{5((x-1)^2)^{1/5}}{2}$$

input `int(-(x^2 - 2*x + 1)^(1/5)/(x - 1),x)`

output `-(5*((x - 1)^2)^(1/5))/2`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt[5]{1-2x+x^2}}{1-x} dx = -\frac{5(x^2-2x+1)^{1/5}}{2}$$

input `int((x^2-2*x+1)^(1/5)/(1-x),x)`

output `(- 5*(x**2 - 2*x + 1)**(1/5))/2`

3.22 $\int x \sin(x) dx$

Optimal result	195
Mathematica [A] (verified)	195
Rubi [A] (verified)	196
Maple [A] (verified)	197
Fricas [A] (verification not implemented)	198
Sympy [A] (verification not implemented)	198
Maxima [A] (verification not implemented)	198
Giac [A] (verification not implemented)	199
Mupad [B] (verification not implemented)	199
Reduce [B] (verification not implemented)	199

Optimal result

Integrand size = 4, antiderivative size = 8

$$\int x \sin(x) dx = -x \cos(x) + \sin(x)$$

output `-x*cos(x)+sin(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int x \sin(x) dx = -x \cos(x) + \sin(x)$$

input `Integrate[x*Sin[x],x]`

output `-(x*Cos[x]) + Sin[x]`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int x \sin(x) dx \\
 \downarrow \text{3042} \\
 \int x \sin(x) dx \\
 \downarrow \text{3777} \\
 \int \cos(x) dx - x \cos(x) \\
 \downarrow \text{3042} \\
 \int \sin\left(x + \frac{\pi}{2}\right) dx - x \cos(x) \\
 \downarrow \text{3117} \\
 \sin(x) - x \cos(x)
 \end{array}$$

input `Int[x*Sin[x],x]`

output `-(x*Cos[x]) + Sin[x]`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] :=> Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] :=> Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.12

method	result	size
default	$-x \cos(x) + \sin(x)$	9
risch	$-x \cos(x) + \sin(x)$	9
parallelrisch	$-x \cos(x) + \sin(x)$	9
parts	$-x \cos(x) + \sin(x)$	9
orering	$-x \cos(x) + \sin(x)$	9
meijerg	$2\sqrt{\pi} \left(-\frac{x \cos(x)}{2\sqrt{\pi}} + \frac{\sin(x)}{2\sqrt{\pi}} \right)$	22
norman	$\frac{x \tan(\frac{x}{2})^2 - x + 2 \tan(\frac{x}{2})}{1 + \tan(\frac{x}{2})^2}$	30

input `int(x*sin(x),x,method=_RETURNVERBOSE)`

output `-x*cos(x)+sin(x)`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int x \sin(x) dx = -x \cos(x) + \sin(x)$$

input `integrate(x*sin(x),x, algorithm="fricas")`

output `-x*cos(x) + sin(x)`

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int x \sin(x) dx = -x \cos(x) + \sin(x)$$

input `integrate(x*sin(x),x)`

output `-x*cos(x) + sin(x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int x \sin(x) dx = -x \cos(x) + \sin(x)$$

input `integrate(x*sin(x),x, algorithm="maxima")`

output `-x*cos(x) + sin(x)`

Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int x \sin(x) dx = -x \cos(x) + \sin(x)$$

input `integrate(x*sin(x),x, algorithm="giac")`

output `-x*cos(x) + sin(x)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int x \sin(x) dx = \sin(x) - x \cos(x)$$

input `int(x*sin(x),x)`

output `sin(x) - x*cos(x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int x \sin(x) dx = -\cos(x)x + \sin(x)$$

input `int(x*sin(x),x)`

output `-cos(x)*x + sin(x)`

3.23 $\int x^2 \sin(x) dx$

Optimal result	200
Mathematica [A] (verified)	200
Rubi [A] (verified)	201
Maple [A] (verified)	202
Fricas [A] (verification not implemented)	203
Sympy [A] (verification not implemented)	203
Maxima [A] (verification not implemented)	203
Giac [A] (verification not implemented)	204
Mupad [B] (verification not implemented)	204
Reduce [B] (verification not implemented)	204

Optimal result

Integrand size = 6, antiderivative size = 17

$$\int x^2 \sin(x) dx = 2 \cos(x) - x^2 \cos(x) + 2x \sin(x)$$

output

```
2*cos(x)-x^2*cos(x)+2*x*sin(x)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int x^2 \sin(x) dx = -((-2 + x^2) \cos(x)) + 2x \sin(x)$$

input

```
Integrate[x^2*Sin[x],x]
```

output

```
-((-2 + x^2)*Cos[x]) + 2*x*Sin[x]
```

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 1.167$, Rules used = {3042, 3777, 3042, 3777, 25, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \sin(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x^2 \sin(x) dx \\
 & \quad \downarrow \text{3777} \\
 & 2 \int x \cos(x) dx - x^2 \cos(x) \\
 & \quad \downarrow \text{3042} \\
 & 2 \int x \sin\left(x + \frac{\pi}{2}\right) dx - x^2 \cos(x) \\
 & \quad \downarrow \text{3777} \\
 & 2\left(\int -\sin(x) dx + x \sin(x)\right) - x^2 \cos(x) \\
 & \quad \downarrow \text{25} \\
 & 2(x \sin(x) - \int \sin(x) dx) - x^2 \cos(x) \\
 & \quad \downarrow \text{3042} \\
 & 2(x \sin(x) - \int \sin(x) dx) - x^2 \cos(x) \\
 & \quad \downarrow \text{3118} \\
 & 2(x \sin(x) + \cos(x)) - x^2 \cos(x)
 \end{aligned}$$

input

```
Int[x^2*Sin[x],x]
```

output $-(x^2 \cos(x)) + 2(\cos(x) + x \sin(x))$

Defintions of rubi rules used

rule 25 $\text{Int}[-(F x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F x, x], x]$

rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinear} \\ \text{Q}[u, x]$

rule 3118 $\text{Int}[\sin[(c.) + (d.)*(x)], x_Symbol] \rightarrow \text{Simp}[-\text{Cos}[c + d*x]/d, x] \text{ ; FreeQ} \\ \{c, d\}, x]$

rule 3777 $\text{Int}[((c.) + (d.)*(x_))^{(m.)}*\sin[(e.) + (f.)*(x)], x_Symbol] \rightarrow \text{Simp}[(\\ -(c + d*x)^m)*(\text{Cos}[e + f*x]/f), x] + \text{Simp}[d*(m/f) \quad \text{Int}[(c + d*x)^{(m - 1)}*C \\ \text{os}[e + f*x], x], x] \text{ ; FreeQ}\{c, d, e, f\}, x \ \&\& \text{GtQ}[m, 0]$

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

method	result	size
risch	$(-x^2 + 2) \cos(x) + 2x \sin(x)$	17
default	$2 \cos(x) - x^2 \cos(x) + 2x \sin(x)$	18
parts	$2 \cos(x) - x^2 \cos(x) + 2x \sin(x)$	18
parallelrisch	$-x^2 \cos(x) + 2x \sin(x) + 2 \cos(x) + 2$	19
meijerg	$4\sqrt{\pi} \left(-\frac{1}{2\sqrt{\pi}} + \frac{\left(-\frac{x^2}{2} + 1\right) \cos(x)}{2\sqrt{\pi}} + \frac{x \sin(x)}{2\sqrt{\pi}} \right)$	34
norman	$\frac{x^2 \tan\left(\frac{x}{2}\right)^2 - x^2 + 4x \tan\left(\frac{x}{2}\right) + 4}{1 + \tan\left(\frac{x}{2}\right)^2}$	36
orering	$\frac{4(x^2 - 1) \sin(x)}{x} - \frac{(x^2 - 2)(2x \sin(x) + x^2 \cos(x))}{x^2}$	36

input $\text{int}(x^2*\sin(x), x, \text{method}=_RETURNVERBOSE)$

output `(-x^2+2)*cos(x)+2*x*sin(x)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int x^2 \sin(x) dx = -(x^2 - 2) \cos(x) + 2x \sin(x)$$

input `integrate(x^2*sin(x),x, algorithm="fricas")`

output `-(x^2 - 2)*cos(x) + 2*x*sin(x)`

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int x^2 \sin(x) dx = -x^2 \cos(x) + 2x \sin(x) + 2 \cos(x)$$

input `integrate(x**2*sin(x),x)`

output `-x**2*cos(x) + 2*x*sin(x) + 2*cos(x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int x^2 \sin(x) dx = -(x^2 - 2) \cos(x) + 2x \sin(x)$$

input `integrate(x^2*sin(x),x, algorithm="maxima")`

output `-(x^2 - 2)*cos(x) + 2*x*sin(x)`

Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int x^2 \sin(x) dx = -(x^2 - 2) \cos(x) + 2x \sin(x)$$

input `integrate(x^2*sin(x),x, algorithm="giac")`

output `-(x^2 - 2)*cos(x) + 2*x*sin(x)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int x^2 \sin(x) dx = 2x \sin(x) - \cos(x) (x^2 - 2)$$

input `int(x^2*sin(x),x)`

output `2*x*sin(x) - cos(x)*(x^2 - 2)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int x^2 \sin(x) dx = -\cos(x) x^2 + 2 \cos(x) + 2 \sin(x) x$$

input `int(x^2*sin(x),x)`

output `- cos(x)*x**2 + 2*cos(x) + 2*sin(x)*x`

3.24 $\int x^3 \cos(x) dx$

Optimal result	205
Mathematica [A] (verified)	205
Rubi [A] (verified)	206
Maple [A] (verified)	208
Fricas [A] (verification not implemented)	208
Sympy [A] (verification not implemented)	209
Maxima [A] (verification not implemented)	209
Giac [A] (verification not implemented)	209
Mupad [B] (verification not implemented)	210
Reduce [B] (verification not implemented)	210

Optimal result

Integrand size = 6, antiderivative size = 23

$$\int x^3 \cos(x) dx = -6 \cos(x) + 3x^2 \cos(x) - 6x \sin(x) + x^3 \sin(x)$$

output `-6*cos(x)+3*x^2*cos(x)-6*x*sin(x)+x^3*sin(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int x^3 \cos(x) dx = 3(-2 + x^2) \cos(x) + x(-6 + x^2) \sin(x)$$

input `Integrate[x^3*Cos[x],x]`

output `3*(-2 + x^2)*Cos[x] + x*(-6 + x^2)*Sin[x]`

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.13, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 1.667$, Rules used = {3042, 3777, 25, 3042, 3777, 3042, 3777, 25, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \cos(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x^3 \sin\left(x + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{3777} \\
 & 3 \int -x^2 \sin(x) dx + x^3 \sin(x) \\
 & \quad \downarrow \text{25} \\
 & x^3 \sin(x) - 3 \int x^2 \sin(x) dx \\
 & \quad \downarrow \text{3042} \\
 & x^3 \sin(x) - 3 \int x^2 \sin(x) dx \\
 & \quad \downarrow \text{3777} \\
 & x^3 \sin(x) - 3 \left(2 \int x \cos(x) dx - x^2 \cos(x) \right) \\
 & \quad \downarrow \text{3042} \\
 & x^3 \sin(x) - 3 \left(2 \int x \sin\left(x + \frac{\pi}{2}\right) dx - x^2 \cos(x) \right) \\
 & \quad \downarrow \text{3777} \\
 & x^3 \sin(x) - 3 \left(2 \left(\int -\sin(x) dx + x \sin(x) \right) - x^2 \cos(x) \right) \\
 & \quad \downarrow \text{25} \\
 & x^3 \sin(x) - 3 \left(2(x \sin(x) - \int \sin(x) dx) - x^2 \cos(x) \right)
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow \text{3042} \\
 x^3 \sin(x) - 3 \left(2(x \sin(x) - \int \sin(x) dx) - x^2 \cos(x) \right) \\
 \downarrow \text{3118} \\
 x^3 \sin(x) - 3(2(x \sin(x) + \cos(x)) - x^2 \cos(x))
 \end{array}$$

input `Int[x^3*Cos[x],x]`

output `x^3*Sin[x] - 3*(-(x^2*Cos[x]) + 2*(Cos[x] + x*Sin[x]))`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] :> Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

method	result	size
risch	$3(x^2 - 2) \cos(x) + x(x^2 - 6) \sin(x)$	20
parallelrisch	$(3x^2 - 6) \cos(x) - 6 + (x^3 - 6x) \sin(x)$	23
default	$-6 \cos(x) + 3x^2 \cos(x) - 6x \sin(x) + x^3 \sin(x)$	24
parts	$-6 \cos(x) + 3x^2 \cos(x) - 6x \sin(x) + x^3 \sin(x)$	24
orering	$6(x^2 - 4) \cos(x) - \frac{(x^2 - 6)(3x^2 \cos(x) - x^3 \sin(x))}{x^2}$	36
meijerg	$8\sqrt{\pi} \left(\frac{3}{4\sqrt{\pi}} - \frac{(-\frac{3x^2}{2} + 3) \cos(x)}{4\sqrt{\pi}} - \frac{x(-\frac{x^2}{2} + 3) \sin(x)}{4\sqrt{\pi}} \right)$	41
norman	$\frac{3x^2 - 12x \tan(\frac{x}{2}) - 3x^2 \tan(\frac{x}{2})^2 + 2x^3 \tan(\frac{x}{2}) - 12}{1 + \tan(\frac{x}{2})^2}$	46

input `int(x^3*cos(x),x,method=_RETURNVERBOSE)`

output `3*(x^2-2)*cos(x)+x*(x^2-6)*sin(x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int x^3 \cos(x) dx = 3(x^2 - 2) \cos(x) + (x^3 - 6x) \sin(x)$$

input `integrate(x^3*cos(x),x, algorithm="fricas")`

output `3*(x^2 - 2)*cos(x) + (x^3 - 6*x)*sin(x)`

Sympy [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.13

$$\int x^3 \cos(x) dx = x^3 \sin(x) + 3x^2 \cos(x) - 6x \sin(x) - 6 \cos(x)$$

input `integrate(x**3*cos(x),x)`

output `x**3*sin(x) + 3*x**2*cos(x) - 6*x*sin(x) - 6*cos(x)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int x^3 \cos(x) dx = 3(x^2 - 2) \cos(x) + (x^3 - 6x) \sin(x)$$

input `integrate(x^3*cos(x),x, algorithm="maxima")`

output `3*(x^2 - 2)*cos(x) + (x^3 - 6*x)*sin(x)`

Giac [A] (verification not implemented)

Time = 0.60 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int x^3 \cos(x) dx = 3(x^2 - 2) \cos(x) + (x^3 - 6x) \sin(x)$$

input `integrate(x^3*cos(x),x, algorithm="giac")`

output `3*(x^2 - 2)*cos(x) + (x^3 - 6*x)*sin(x)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.04

$$\int x^3 \cos(x) dx = \cos(x) (3x^2 - 6) - \sin(x) (6x - x^3)$$

input `int(x^3*cos(x),x)`

output `cos(x)*(3*x^2 - 6) - sin(x)*(6*x - x^3)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int x^3 \cos(x) dx = 3 \cos(x) x^2 - 6 \cos(x) + \sin(x) x^3 - 6 \sin(x) x$$

input `int(x^3*cos(x),x)`

output `3*cos(x)*x**2 - 6*cos(x) + sin(x)*x**3 - 6*sin(x)*x`

3.25 $\int x^3 \sin(x) dx$

Optimal result	211
Mathematica [A] (verified)	211
Rubi [A] (verified)	212
Maple [A] (verified)	213
Fricas [A] (verification not implemented)	214
Sympy [A] (verification not implemented)	215
Maxima [A] (verification not implemented)	215
Giac [A] (verification not implemented)	215
Mupad [B] (verification not implemented)	216
Reduce [B] (verification not implemented)	216

Optimal result

Integrand size = 6, antiderivative size = 24

$$\int x^3 \sin(x) dx = 6x \cos(x) - x^3 \cos(x) - 6 \sin(x) + 3x^2 \sin(x)$$

output `6*x*cos(x)-x^3*cos(x)-6*sin(x)+3*x^2*sin(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int x^3 \sin(x) dx = -x(-6 + x^2) \cos(x) + 3(-2 + x^2) \sin(x)$$

input `Integrate[x^3*Sin[x],x]`

output `-(x*(-6 + x^2)*Cos[x]) + 3*(-2 + x^2)*Sin[x]`

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.12, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$, Rules used = {3042, 3777, 3042, 3777, 25, 3042, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \sin(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x^3 \sin(x) dx \\
 & \quad \downarrow \text{3777} \\
 & 3 \int x^2 \cos(x) dx - x^3 \cos(x) \\
 & \quad \downarrow \text{3042} \\
 & 3 \int x^2 \sin\left(x + \frac{\pi}{2}\right) dx - x^3 \cos(x) \\
 & \quad \downarrow \text{3777} \\
 & 3 \left(2 \int -x \sin(x) dx + x^2 \sin(x) \right) - x^3 \cos(x) \\
 & \quad \downarrow \text{25} \\
 & 3 \left(x^2 \sin(x) - 2 \int x \sin(x) dx \right) - x^3 \cos(x) \\
 & \quad \downarrow \text{3042} \\
 & 3 \left(x^2 \sin(x) - 2 \int x \sin(x) dx \right) - x^3 \cos(x) \\
 & \quad \downarrow \text{3777} \\
 & 3 \left(x^2 \sin(x) - 2 \left(\int \cos(x) dx - x \cos(x) \right) \right) - x^3 \cos(x) \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$3\left(x^2 \sin(x) - 2\left(\int \sin\left(x + \frac{\pi}{2}\right) dx - x \cos(x)\right)\right) - x^3 \cos(x)$$

↓ 3117

$$3(x^2 \sin(x) - 2(\sin(x) - x \cos(x))) - x^3 \cos(x)$$

input `Int[x^3*Sin[x],x]`

output `-(x^3*Cos[x]) + 3*(x^2*Sin[x] - 2*(-(x*Cos[x]) + Sin[x]))`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.96

method	result	size
risch	$(-x^3 + 6x) \cos(x) + 3(x^2 - 2) \sin(x)$	23
default	$6x \cos(x) - x^3 \cos(x) - 6 \sin(x) + 3x^2 \sin(x)$	25
parallelrisc	$6x \cos(x) - x^3 \cos(x) - 6 \sin(x) + 3x^2 \sin(x)$	25
parts	$6x \cos(x) - x^3 \cos(x) - 6 \sin(x) + 3x^2 \sin(x)$	25
orering	$6(x^2 - 4) \sin(x) - \frac{(x^2-6)(3x^2 \sin(x)+x^3 \cos(x))}{x^2}$	35
meijerg	$8\sqrt{\pi} \left(\frac{x \left(-\frac{5x^2}{2} + 15\right) \cos(x)}{20\sqrt{\pi}} - \frac{\left(-\frac{15x^2}{2} + 15\right) \sin(x)}{20\sqrt{\pi}} \right)$	36
norman	$\frac{x^3 \tan\left(\frac{x}{2}\right)^2 + 6x - x^3 - 6x \tan\left(\frac{x}{2}\right)^2 + 6x^2 \tan\left(\frac{x}{2}\right) - 12 \tan\left(\frac{x}{2}\right)}{1 + \tan\left(\frac{x}{2}\right)^2}$	55

input `int(x^3*sin(x),x,method=_RETURNVERBOSE)`

output `(-x^3+6*x)*cos(x)+3*(x^2-2)*sin(x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.88

$$\int x^3 \sin(x) dx = -(x^3 - 6x) \cos(x) + 3(x^2 - 2) \sin(x)$$

input `integrate(x^3*sin(x),x, algorithm="fricas")`

output `-(x^3 - 6*x)*cos(x) + 3*(x^2 - 2)*sin(x)`

Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int x^3 \sin(x) dx = -x^3 \cos(x) + 3x^2 \sin(x) + 6x \cos(x) - 6 \sin(x)$$

input `integrate(x**3*sin(x),x)`

output `-x**3*cos(x) + 3*x**2*sin(x) + 6*x*cos(x) - 6*sin(x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.88

$$\int x^3 \sin(x) dx = -(x^3 - 6x) \cos(x) + 3(x^2 - 2) \sin(x)$$

input `integrate(x^3*sin(x),x, algorithm="maxima")`

output `-(x^3 - 6*x)*cos(x) + 3*(x^2 - 2)*sin(x)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.88

$$\int x^3 \sin(x) dx = -(x^3 - 6x) \cos(x) + 3(x^2 - 2) \sin(x)$$

input `integrate(x^3*sin(x),x, algorithm="giac")`

output `-(x^3 - 6*x)*cos(x) + 3*(x^2 - 2)*sin(x)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.96

$$\int x^3 \sin(x) dx = \cos(x) (6x - x^3) + \sin(x) (3x^2 - 6)$$

input `int(x^3*sin(x),x)`

output `cos(x)*(6*x - x^3) + sin(x)*(3*x^2 - 6)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int x^3 \sin(x) dx = -\cos(x) x^3 + 6 \cos(x) x + 3 \sin(x) x^2 - 6 \sin(x)$$

input `int(x^3*sin(x),x)`

output `-cos(x)*x**3 + 6*cos(x)*x + 3*sin(x)*x**2 - 6*sin(x)`

3.26 $\int \cos(x) \sin(x) dx$

Optimal result	217
Mathematica [A] (verified)	217
Rubi [A] (verified)	218
Maple [A] (verified)	219
Fricas [A] (verification not implemented)	219
Sympy [A] (verification not implemented)	220
Maxima [A] (verification not implemented)	220
Giac [A] (verification not implemented)	220
Mupad [B] (verification not implemented)	221
Reduce [B] (verification not implemented)	221

Optimal result

Integrand size = 5, antiderivative size = 8

$$\int \cos(x) \sin(x) dx = \frac{\sin^2(x)}{2}$$

output `1/2*sin(x)^2`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \cos(x) \sin(x) dx = -\frac{1}{2} \cos^2(x)$$

input `Integrate[Cos[x]*Sin[x],x]`

output `-1/2*Cos[x]^2`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3042, 3044, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \sin(x) \cos(x) dx \\ \downarrow \text{3042} \\ \int \sin(x) \cos(x) dx \\ \downarrow \text{3044} \\ \int \sin(x) d \sin(x) \\ \downarrow \text{15} \\ \frac{\sin^2(x)}{2} \end{array}$$

input `Int[Cos[x]*Sin[x],x]`

output `Sin[x]^2/2`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3044

```
Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)]^(m_.), x_
Symbol] := Simp[1/(a*f) Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a
*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(I
ntegerQ[(m - 1)/2] && LtQ[0, m, n])
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$\frac{\sin(x)^2}{2}$	7
default	$\frac{\sin(x)^2}{2}$	7
risch	$-\frac{\cos(2x)}{4}$	7
parallelrisch	$\frac{1}{4} - \frac{\cos(2x)}{4}$	9
orering	$-\frac{\cos(x)^2}{4} + \frac{\sin(x)^2}{4}$	14
norman	$\frac{2 \tan(\frac{x}{2})^2}{(1 + \tan(\frac{x}{2})^2)^2}$	19
meijerg	$\frac{\sqrt{\pi} \left(\frac{1}{\sqrt{\pi}} - \frac{\cos(2x)}{\sqrt{\pi}} \right)}{4}$	19

input

```
int(cos(x)*sin(x), x, method=_RETURNVERBOSE)
```

output

```
1/2*sin(x)^2
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos(x) \sin(x) dx = -\frac{1}{2} \cos(x)^2$$

input

```
integrate(cos(x)*sin(x), x, algorithm="fricas")
```

output `-1/2*cos(x)^2`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int \cos(x) \sin(x) dx = \frac{\sin^2(x)}{2}$$

input `integrate(cos(x)*sin(x),x)`

output `sin(x)**2/2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos(x) \sin(x) dx = -\frac{1}{2} \cos(x)^2$$

input `integrate(cos(x)*sin(x),x, algorithm="maxima")`

output `-1/2*cos(x)^2`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos(x) \sin(x) dx = -\frac{1}{2} \cos(x)^2$$

input `integrate(cos(x)*sin(x),x, algorithm="giac")`

output `-1/2*cos(x)^2`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos(x) \sin(x) dx = \frac{\sin(x)^2}{2}$$

input `int(cos(x)*sin(x),x)`

output `sin(x)^2/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos(x) \sin(x) dx = -\frac{\cos(x)^2}{2}$$

input `int(cos(x)*sin(x),x)`

output `(- cos(x)**2)/2`

3.27 $\int x \cos(x) \sin(x) dx$

Optimal result	222
Mathematica [A] (verified)	222
Rubi [A] (verified)	223
Maple [A] (verified)	224
Fricas [A] (verification not implemented)	225
Sympy [A] (verification not implemented)	225
Maxima [A] (verification not implemented)	225
Giac [A] (verification not implemented)	226
Mupad [B] (verification not implemented)	226
Reduce [B] (verification not implemented)	226

Optimal result

Integrand size = 6, antiderivative size = 23

$$\int x \cos(x) \sin(x) dx = -\frac{x}{4} + \frac{1}{4} \cos(x) \sin(x) + \frac{1}{2} x \sin^2(x)$$

output `-1/4*x+1/4*cos(x)*sin(x)+1/2*x*sin(x)^2`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

$$\int x \cos(x) \sin(x) dx = -\frac{1}{4} x \cos(2x) + \frac{1}{8} \sin(2x)$$

input `Integrate[x*Cos[x]*Sin[x],x]`

output `-1/4*(x*Cos[2*x]) + Sin[2*x]/8`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.22, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3924, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \sin(x) \cos(x) dx \\
 & \quad \downarrow \text{3924} \\
 & \frac{1}{2} x \sin^2(x) - \frac{1}{2} \int \sin^2(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} x \sin^2(x) - \frac{1}{2} \int \sin(x)^2 dx \\
 & \quad \downarrow \text{3115} \\
 & \frac{1}{2} \left(\frac{1}{2} \sin(x) \cos(x) - \frac{\int 1 dx}{2} \right) + \frac{1}{2} x \sin^2(x) \\
 & \quad \downarrow \text{24} \\
 & \frac{1}{2} x \sin^2(x) + \frac{1}{2} \left(\frac{1}{2} \sin(x) \cos(x) - \frac{x}{2} \right)
 \end{aligned}$$

input `Int [x*Cos [x] *Sin [x] , x]`

output `(x*Sin [x]^2)/2 + (-1/2*x + (Cos [x] *Sin [x])/2)/2`

Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3924 `Int[Cos[(a_.) + (b_.)*(x_)^(n_.)]*(x_)^(m_.)*Sin[(a_.) + (b_.)*(x_)^(n_.)]^(p_.), x_Symbol] := Simp[x^(m - n + 1)*(Sin[a + b*x^n]^(p + 1)/(b*n*(p + 1))), x] - Simp[(m - n + 1)/(b*n*(p + 1)) Int[x^(m - n)*Sin[a + b*x^n]^(p + 1), x], x] /; FreeQ[{a, b, p}, x] && LtQ[0, n, m + 1] && NeQ[p, -1]`

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

method	result	size
risch	$-\frac{x \cos(2x)}{4} + \frac{\sin(2x)}{8}$	15
paralelrisch	$-\frac{x \cos(2x)}{4} + \frac{\sin(2x)}{8}$	15
default	$-\frac{x \cos(x)^2}{2} + \frac{\cos(x) \sin(x)}{4} + \frac{x}{4}$	18
oring	$\frac{\cos(x) \sin(x)}{4} + \frac{x \sin(x)^2}{4} - \frac{x \cos(x)^2}{4}$	22
meijerg	$\frac{\sqrt{\pi} \left(-\frac{x \cos(2x)}{\sqrt{\pi}} + \frac{\sin(2x)}{2\sqrt{\pi}} \right)}{4}$	26
norman	$-\frac{x}{4} - \frac{\tan\left(\frac{x}{2}\right)^3}{2} + \frac{3x \tan\left(\frac{x}{2}\right)^2}{2} - \frac{x \tan\left(\frac{x}{2}\right)^4}{4} + \frac{\tan\left(\frac{x}{2}\right)}{2}$ $(1 + \tan\left(\frac{x}{2}\right)^2)^2$	48

input `int(x*cos(x)*sin(x), x, method=_RETURNVERBOSE)`

output `-1/4*x*cos(2*x)+1/8*sin(2*x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int x \cos(x) \sin(x) dx = -\frac{1}{2} x \cos(x)^2 + \frac{1}{4} \cos(x) \sin(x) + \frac{1}{4} x$$

input `integrate(x*cos(x)*sin(x),x, algorithm="fricas")`

output `-1/2*x*cos(x)^2 + 1/4*cos(x)*sin(x) + 1/4*x`

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.04

$$\int x \cos(x) \sin(x) dx = \frac{x \sin^2(x)}{4} - \frac{x \cos^2(x)}{4} + \frac{\sin(x) \cos(x)}{4}$$

input `integrate(x*cos(x)*sin(x),x)`

output `x*sin(x)**2/4 - x*cos(x)**2/4 + sin(x)*cos(x)/4`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.61

$$\int x \cos(x) \sin(x) dx = -\frac{1}{4} x \cos(2x) + \frac{1}{8} \sin(2x)$$

input `integrate(x*cos(x)*sin(x),x, algorithm="maxima")`

output `-1/4*x*cos(2*x) + 1/8*sin(2*x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.61

$$\int x \cos(x) \sin(x) dx = -\frac{1}{4} x \cos(2x) + \frac{1}{8} \sin(2x)$$

input `integrate(x*cos(x)*sin(x),x, algorithm="giac")`

output `-1/4*x*cos(2*x) + 1/8*sin(2*x)`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

$$\int x \cos(x) \sin(x) dx = \frac{\sin(2x)}{8} + \frac{x(2\sin(x)^2 - 1)}{4}$$

input `int(x*cos(x)*sin(x),x)`

output `sin(2*x)/8 + (x*(2*sin(x)^2 - 1))/4`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int x \cos(x) \sin(x) dx = -\frac{\cos(x)^2 x}{4} + \frac{\cos(x) \sin(x)}{4} + \frac{\sin(x)^2 x}{4}$$

input `int(x*cos(x)*sin(x),x)`

output `(- cos(x)**2*x + cos(x)*sin(x) + sin(x)**2*x)/4`

3.28 $\int \sin^2(x) dx$

Optimal result	227
Mathematica [A] (verified)	227
Rubi [A] (verified)	228
Maple [A] (verified)	229
Fricas [A] (verification not implemented)	229
Sympy [A] (verification not implemented)	230
Maxima [A] (verification not implemented)	230
Giac [A] (verification not implemented)	230
Mupad [B] (verification not implemented)	231
Reduce [B] (verification not implemented)	231

Optimal result

Integrand size = 4, antiderivative size = 14

$$\int \sin^2(x) dx = \frac{x}{2} - \frac{1}{2} \cos(x) \sin(x)$$

output `1/2*x-1/2*cos(x)*sin(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \sin^2(x) dx = \frac{x}{2} - \frac{1}{4} \sin(2x)$$

input `Integrate[Sin[x]^2,x]`

output `x/2 - Sin[2*x]/4`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sin^2(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin(x)^2 dx \\ & \quad \downarrow \text{3115} \\ & \frac{\int 1 dx}{2} - \frac{1}{2} \sin(x) \cos(x) \\ & \quad \downarrow \text{24} \\ & \frac{x}{2} - \frac{1}{2} \sin(x) \cos(x) \end{aligned}$$

input `Int[Sin[x]^2,x]`

output `x/2 - (Cos[x]*Sin[x])/2`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.79

method	result	size
default	$\frac{x}{2} - \frac{\cos(x)\sin(x)}{2}$	11
risch	$\frac{x}{2} - \frac{\sin(2x)}{4}$	11
parallelrisc	$\frac{x}{2} - \frac{\sin(2x)}{4}$	11
meijerg	$\frac{\sqrt{\pi} \left(\frac{2x}{\sqrt{\pi}} - \frac{\sin(2x)}{\sqrt{\pi}} \right)}{4}$	22
orering	$x \sin(x)^2 - \frac{\cos(x)\sin(x)}{2} + \frac{x(2\cos(x)^2 - 2\sin(x)^2)}{4}$	30
norman	$\frac{\tan(\frac{x}{2})^3 + x \tan(\frac{x}{2})^2 + \frac{x}{2} + \frac{x \tan(\frac{x}{2})^4}{2} - \tan(\frac{x}{2})}{(1 + \tan(\frac{x}{2})^2)^2}$	45

input

```
int(sin(x)^2,x,method=_RETURNVERBOSE)
```

output

```
1/2*x-1/2*cos(x)*sin(x)
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \sin^2(x) dx = -\frac{1}{2} \cos(x) \sin(x) + \frac{1}{2} x$$

input

```
integrate(sin(x)^2,x, algorithm="fricas")
```

output

```
-1/2*cos(x)*sin(x) + 1/2*x
```

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \sin^2(x) dx = \frac{x}{2} - \frac{\sin(x) \cos(x)}{2}$$

input `integrate(sin(x)**2,x)`

output `x/2 - sin(x)*cos(x)/2`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \sin^2(x) dx = \frac{1}{2}x - \frac{1}{4} \sin(2x)$$

input `integrate(sin(x)^2,x, algorithm="maxima")`

output `1/2*x - 1/4*sin(2*x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \sin^2(x) dx = \frac{1}{2}x - \frac{1}{4} \sin(2x)$$

input `integrate(sin(x)^2,x, algorithm="giac")`

output `1/2*x - 1/4*sin(2*x)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \sin^2(x) dx = \frac{x}{2} - \frac{\sin(2x)}{4}$$

input `int(sin(x)^2,x)`

output `x/2 - sin(2*x)/4`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \sin^2(x) dx = -\frac{\cos(x) \sin(x)}{2} + \frac{x}{2}$$

input `int(sin(x)^2,x)`

output `(- cos(x)*sin(x) + x)/2`

3.29 $\int \sin^3(x) dx$

Optimal result	232
Mathematica [A] (verified)	232
Rubi [A] (verified)	233
Maple [A] (verified)	234
Fricas [A] (verification not implemented)	234
Sympy [A] (verification not implemented)	235
Maxima [A] (verification not implemented)	235
Giac [A] (verification not implemented)	235
Mupad [B] (verification not implemented)	236
Reduce [B] (verification not implemented)	236

Optimal result

Integrand size = 4, antiderivative size = 13

$$\int \sin^3(x) dx = -\cos(x) + \frac{\cos^3(x)}{3}$$

output

```
-cos(x)+1/3*cos(x)^3
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \sin^3(x) dx = -\frac{3 \cos(x)}{4} + \frac{1}{12} \cos(3x)$$

input

```
Integrate[Sin[x]^3,x]
```

output

```
(-3*Cos[x])/4 + Cos[3*x]/12
```

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3113, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sin^3(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin(x)^3 dx \\ & \quad \downarrow \text{3113} \\ & - \int (1 - \cos^2(x)) d \cos(x) \\ & \quad \downarrow \text{2009} \\ & \frac{\cos^3(x)}{3} - \cos(x) \end{aligned}$$

input `Int[Sin[x]^3,x]`

output `-Cos[x] + Cos[x]^3/3`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113

```
Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
and[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x]
&& IGtQ[(n - 1)/2, 0]
```

Maple [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

method	result	size
default	$-\frac{(2+\sin(x)^2)\cos(x)}{3}$	11
risch	$-\frac{3\cos(x)}{4} + \frac{\cos(3x)}{12}$	12
parallelrisch	$-\frac{2}{3} - \frac{3\cos(x)}{4} + \frac{\cos(3x)}{12}$	13
orering	$-\sin(x)^2\cos(x) - \frac{2\cos(x)^3}{3}$	16
norman	$\frac{-4\tan(\frac{x}{2})^2 - \frac{4}{3}}{(1+\tan(\frac{x}{2})^2)^3}$	22

input

```
int(sin(x)^3,x,method=_RETURNVERBOSE)
```

output

```
-1/3*(2+sin(x)^2)*cos(x)
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \sin^3(x) dx = \frac{1}{3} \cos(x)^3 - \cos(x)$$

input

```
integrate(sin(x)^3,x, algorithm="fricas")
```

output

```
1/3*cos(x)^3 - cos(x)
```

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.62

$$\int \sin^3(x) dx = \frac{\cos^3(x)}{3} - \cos(x)$$

input `integrate(sin(x)**3,x)`

output `cos(x)**3/3 - cos(x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \sin^3(x) dx = \frac{1}{3} \cos(x)^3 - \cos(x)$$

input `integrate(sin(x)^3,x, algorithm="maxima")`

output `1/3*cos(x)^3 - cos(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \sin^3(x) dx = \frac{1}{3} \cos(x)^3 - \cos(x)$$

input `integrate(sin(x)^3,x, algorithm="giac")`

output `1/3*cos(x)^3 - cos(x)`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \sin^3(x) dx = \frac{\cos(x) (\cos(x)^2 - 3)}{3}$$

input `int(sin(x)^3,x)`

output `(cos(x)*(cos(x)^2 - 3))/3`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.08

$$\int \sin^3(x) dx = -\frac{\cos(x) \sin(x)^2}{3} - \frac{2 \cos(x)}{3} + \frac{2}{3}$$

input `int(sin(x)^3,x)`

output `(- cos(x)*sin(x)**2 - 2*cos(x) + 2)/3`

3.30 $\int \sin^4(x) dx$

Optimal result	237
Mathematica [A] (verified)	237
Rubi [A] (verified)	238
Maple [A] (verified)	239
Fricas [A] (verification not implemented)	240
Sympy [A] (verification not implemented)	240
Maxima [A] (verification not implemented)	240
Giac [A] (verification not implemented)	241
Mupad [B] (verification not implemented)	241
Reduce [B] (verification not implemented)	241

Optimal result

Integrand size = 4, antiderivative size = 24

$$\int \sin^4(x) dx = \frac{3x}{8} - \frac{3}{8} \cos(x) \sin(x) - \frac{1}{4} \cos(x) \sin^3(x)$$

output `3/8*x-3/8*cos(x)*sin(x)-1/4*cos(x)*sin(x)^3`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \sin^4(x) dx = \frac{3x}{8} - \frac{1}{4} \sin(2x) + \frac{1}{32} \sin(4x)$$

input `Integrate[Sin[x]^4,x]`

output `(3*x)/8 - Sin[2*x]/4 + Sin[4*x]/32`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.21, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$, Rules used = {3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^4(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x)^4 dx \\
 & \quad \downarrow \text{3115} \\
 & \frac{3}{4} \int \sin^2(x) dx - \frac{1}{4} \sin^3(x) \cos(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{4} \int \sin(x)^2 dx - \frac{1}{4} \sin^3(x) \cos(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{3}{4} \left(\frac{\int 1 dx}{2} - \frac{1}{2} \sin(x) \cos(x) \right) - \frac{1}{4} \sin^3(x) \cos(x) \\
 & \quad \downarrow \text{24} \\
 & \frac{3}{4} \left(\frac{x}{2} - \frac{1}{2} \sin(x) \cos(x) \right) - \frac{1}{4} \sin^3(x) \cos(x)
 \end{aligned}$$

input `Int [Sin [x] ^4, x]`

output `-1/4*(Cos [x]*Sin [x]^3) + (3*(x/2 - (Cos [x]*Sin [x])/2))/4`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.71

method	result
risch	$\frac{3x}{8} + \frac{\sin(4x)}{32} - \frac{\sin(2x)}{4}$
parallelrisc	$\frac{3x}{8} + \frac{\sin(4x)}{32} - \frac{\sin(2x)}{4}$
default	$-\frac{(\sin(x)^3 + \frac{3\sin(x)}{2})\cos(x)}{4} + \frac{3x}{8}$
orering	$x \sin(x)^4 - \frac{5 \cos(x) \sin(x)^3}{8} + \frac{5x(-4 \sin(x)^4 + 12 \sin(x)^2 \cos(x)^2)}{16} - \frac{3 \cos(x)^3 \sin(x)}{8} + \frac{x(40 \sin(x)^4 - 192 \sin(x)^2)}{64}$
norman	$\frac{3x}{8} - \frac{11 \tan(\frac{x}{2})^3}{4} + \frac{11 \tan(\frac{x}{2})^5}{4} + \frac{3 \tan(\frac{x}{2})^7}{4} + \frac{3x \tan(\frac{x}{2})^2}{2} + \frac{9x \tan(\frac{x}{2})^4}{4} + \frac{3x \tan(\frac{x}{2})^6}{2} + \frac{3x \tan(\frac{x}{2})^8}{8} - \frac{3 \tan(\frac{x}{2})}{4}$ $(1 + \tan(\frac{x}{2})^2)^4$

input `int(sin(x)^4,x,method=_RETURNVERBOSE)`

output `3/8*x+1/32*sin(4*x)-1/4*sin(2*x)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.79

$$\int \sin^4(x) dx = \frac{1}{8} (2 \cos(x)^3 - 5 \cos(x)) \sin(x) + \frac{3}{8} x$$

input `integrate(sin(x)^4,x, algorithm="fricas")`output `1/8*(2*cos(x)^3 - 5*cos(x))*sin(x) + 3/8*x`**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \sin^4(x) dx = \frac{3x}{8} - \frac{\sin^3(x) \cos(x)}{4} - \frac{3 \sin(x) \cos(x)}{8}$$

input `integrate(sin(x)**4,x)`output `3*x/8 - sin(x)**3*cos(x)/4 - 3*sin(x)*cos(x)/8`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int \sin^4(x) dx = \frac{3}{8} x + \frac{1}{32} \sin(4x) - \frac{1}{4} \sin(2x)$$

input `integrate(sin(x)^4,x, algorithm="maxima")`output `3/8*x + 1/32*sin(4*x) - 1/4*sin(2*x)`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int \sin^4(x) dx = \frac{3}{8}x + \frac{1}{32}\sin(4x) - \frac{1}{4}\sin(2x)$$

input `integrate(sin(x)^4,x, algorithm="giac")`

output `3/8*x + 1/32*sin(4*x) - 1/4*sin(2*x)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int \sin^4(x) dx = \frac{3x}{8} - \frac{\sin(2x)}{4} + \frac{\sin(4x)}{32}$$

input `int(sin(x)^4,x)`

output `(3*x)/8 - sin(2*x)/4 + sin(4*x)/32`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.75

$$\int \sin^4(x) dx = -\frac{\cos(x)\sin(x)^3}{4} - \frac{3\cos(x)\sin(x)}{8} + \frac{3x}{8}$$

input `int(sin(x)^4,x)`

output `(- 2*cos(x)*sin(x)**3 - 3*cos(x)*sin(x) + 3*x)/8`

3.31 $\int \sin^5(x) dx$

Optimal result	242
Mathematica [A] (verified)	242
Rubi [A] (verified)	243
Maple [A] (verified)	244
Fricas [A] (verification not implemented)	244
Sympy [A] (verification not implemented)	245
Maxima [A] (verification not implemented)	245
Giac [A] (verification not implemented)	245
Mupad [B] (verification not implemented)	246
Reduce [B] (verification not implemented)	246

Optimal result

Integrand size = 4, antiderivative size = 21

$$\int \sin^5(x) dx = -\cos(x) + \frac{2 \cos^3(x)}{3} - \frac{\cos^5(x)}{5}$$

output

```
-cos(x)+2/3*cos(x)^3-1/5*cos(x)^5
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \sin^5(x) dx = -\frac{5 \cos(x)}{8} + \frac{5}{48} \cos(3x) - \frac{1}{80} \cos(5x)$$

input

```
Integrate[Sin[x]^5,x]
```

output

```
(-5*Cos[x])/8 + (5*Cos[3*x])/48 - Cos[5*x]/80
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3113, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sin^5(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin(x)^5 dx \\ & \quad \downarrow \text{3113} \\ & - \int (\cos^4(x) - 2 \cos^2(x) + 1) d \cos(x) \\ & \quad \downarrow \text{2009} \\ & -\frac{1}{5} \cos^5(x) + \frac{2 \cos^3(x)}{3} - \cos(x) \end{aligned}$$

input `Int[Sin[x]^5,x]`

output `-Cos[x] + (2*Cos[x]^3)/3 - Cos[x]^5/5`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp and[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

Maple [A] (verified)

Time = 0.51 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

method	result	size
default	$-\frac{\left(\frac{8}{3} + \sin(x)^4 + \frac{4 \sin(x)^2}{3}\right) \cos(x)}{5}$	17
risch	$-\frac{5 \cos(x)}{8} - \frac{\cos(5x)}{80} + \frac{5 \cos(3x)}{48}$	18
parallelrisch	$-\frac{8}{15} - \frac{5 \cos(x)}{8} + \frac{5 \cos(3x)}{48} - \frac{\cos(5x)}{80}$	19
orering	$-\sin(x)^4 \cos(x) - \frac{4 \sin(x)^2 \cos(x)^3}{3} - \frac{8 \cos(x)^5}{15}$	26
norman	$\frac{-\frac{32 \tan\left(\frac{x}{2}\right)^4}{3} - \frac{16 \tan\left(\frac{x}{2}\right)^2}{3} - \frac{16}{15}}{\left(1 + \tan\left(\frac{x}{2}\right)^2\right)^5}$	30

input `int(sin(x)^5,x,method=_RETURNVERBOSE)`

output `-1/5*(8/3+sin(x)^4+4/3*sin(x)^2)*cos(x)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \sin^5(x) dx = -\frac{1}{5} \cos(x)^5 + \frac{2}{3} \cos(x)^3 - \cos(x)$$

input `integrate(sin(x)^5,x, algorithm="fricas")`

output `-1/5*cos(x)^5 + 2/3*cos(x)^3 - cos(x)`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \sin^5(x) dx = -\frac{\cos^5(x)}{5} + \frac{2\cos^3(x)}{3} - \cos(x)$$

input `integrate(sin(x)**5,x)`output `-cos(x)**5/5 + 2*cos(x)**3/3 - cos(x)`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \sin^5(x) dx = -\frac{1}{5} \cos(x)^5 + \frac{2}{3} \cos(x)^3 - \cos(x)$$

input `integrate(sin(x)^5,x, algorithm="maxima")`output `-1/5*cos(x)^5 + 2/3*cos(x)^3 - cos(x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \sin^5(x) dx = -\frac{1}{5} \cos(x)^5 + \frac{2}{3} \cos(x)^3 - \cos(x)$$

input `integrate(sin(x)^5,x, algorithm="giac")`output `-1/5*cos(x)^5 + 2/3*cos(x)^3 - cos(x)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \sin^5(x) dx = -\frac{\cos(x)^5}{5} + \frac{2\cos(x)^3}{3} - \cos(x)$$

input `int(sin(x)^5,x)`

output `(2*cos(x)^3)/3 - cos(x) - cos(x)^5/5`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int \sin^5(x) dx = -\frac{\cos(x)\sin(x)^4}{5} - \frac{4\cos(x)\sin(x)^2}{15} - \frac{8\cos(x)}{15} + \frac{8}{15}$$

input `int(sin(x)^5,x)`

output `(- 3*cos(x)*sin(x)**4 - 4*cos(x)*sin(x)**2 - 8*cos(x) + 8)/15`

3.32 $\int \sin^6(x) dx$

Optimal result	247
Mathematica [A] (verified)	247
Rubi [A] (verified)	248
Maple [A] (verified)	249
Fricas [A] (verification not implemented)	250
Sympy [A] (verification not implemented)	250
Maxima [A] (verification not implemented)	250
Giac [A] (verification not implemented)	251
Mupad [B] (verification not implemented)	251
Reduce [B] (verification not implemented)	251

Optimal result

Integrand size = 4, antiderivative size = 34

$$\int \sin^6(x) dx = \frac{5x}{16} - \frac{5}{16} \cos(x) \sin(x) - \frac{5}{24} \cos(x) \sin^3(x) - \frac{1}{6} \cos(x) \sin^5(x)$$

output `5/16*x-5/16*cos(x)*sin(x)-5/24*cos(x)*sin(x)^3-1/6*cos(x)*sin(x)^5`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.88

$$\int \sin^6(x) dx = \frac{5x}{16} - \frac{15}{64} \sin(2x) + \frac{3}{64} \sin(4x) - \frac{1}{192} \sin(6x)$$

input `Integrate[Sin[x]^6,x]`

output `(5*x)/16 - (15*Sin[2*x])/64 + (3*Sin[4*x])/64 - Sin[6*x]/192`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.29, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 1.750$, Rules used = {3042, 3115, 3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^6(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x)^6 dx \\
 & \quad \downarrow \text{3115} \\
 & \frac{5}{6} \int \sin^4(x) dx - \frac{1}{6} \sin^5(x) \cos(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{6} \int \sin(x)^4 dx - \frac{1}{6} \sin^5(x) \cos(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{5}{6} \left(\frac{3}{4} \int \sin^2(x) dx - \frac{1}{4} \sin^3(x) \cos(x) \right) - \frac{1}{6} \sin^5(x) \cos(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{6} \left(\frac{3}{4} \int \sin(x)^2 dx - \frac{1}{4} \sin^3(x) \cos(x) \right) - \frac{1}{6} \sin^5(x) \cos(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{5}{6} \left(\frac{3}{4} \left(\frac{\int 1 dx}{2} - \frac{1}{2} \sin(x) \cos(x) \right) - \frac{1}{4} \sin^3(x) \cos(x) \right) - \frac{1}{6} \sin^5(x) \cos(x) \\
 & \quad \downarrow \text{24} \\
 & \frac{5}{6} \left(\frac{3}{4} \left(\frac{x}{2} - \frac{1}{2} \sin(x) \cos(x) \right) - \frac{1}{4} \sin^3(x) \cos(x) \right) - \frac{1}{6} \sin^5(x) \cos(x)
 \end{aligned}$$

input

Int [Sin [x]^6, x]

output
$$-1/6*(\text{Cos}[x]*\text{Sin}[x]^5) + (5*(-1/4*(\text{Cos}[x]*\text{Sin}[x]^3) + (3*(x/2 - (\text{Cos}[x]*\text{Sin}[x])/2))/4))/6$$

Defintions of rubi rules used

rule 24
$$\text{Int}[a_, x_Symbol] \text{ :> } \text{Simp}[a*x, x] \text{ /; } \text{FreeQ}[a, x]$$

rule 3042
$$\text{Int}[u_, x_Symbol] \text{ :> } \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /; } \text{FunctionOfTrigOfLinearQ}[u, x]$$

rule 3115
$$\text{Int}[(b_)*\text{sin}[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] \text{ :> } \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n-1)}/(d*n), x] + \text{Simp}[b^2*((n-1)/n) \text{ Int}[(b*\text{Sin}[c + d*x])^{(n-2)}, x], x] \text{ /; } \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$$

Maple [A] (verified)

Time = 0.78 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.68

method	result
risch	$\frac{5x}{16} - \frac{\sin(6x)}{192} + \frac{3 \sin(4x)}{64} - \frac{15 \sin(2x)}{64}$
parallelrisch	$\frac{5x}{16} - \frac{\sin(6x)}{192} + \frac{3 \sin(4x)}{64} - \frac{15 \sin(2x)}{64}$
default	$-\frac{\left(\sin(x)^5 + \frac{5 \sin(x)^3}{4} + \frac{15 \sin(x)}{8}\right) \cos(x)}{6} + \frac{5x}{16}$
norman	$\frac{\frac{5x}{16} - \frac{85 \tan(\frac{x}{2})^3}{24} - \frac{33 \tan(\frac{x}{2})^5}{4} + \frac{33 \tan(\frac{x}{2})^7}{4} + \frac{85 \tan(\frac{x}{2})^9}{24} + \frac{5 \tan(\frac{x}{2})^{11}}{8} + \frac{15x \tan(\frac{x}{2})^2}{8} + \frac{75x \tan(\frac{x}{2})^4}{16} + \frac{25x \tan(\frac{x}{2})^6}{4} + \frac{75x \tan(\frac{x}{2})^8}{16}}{(1 + \tan(\frac{x}{2})^2)^6}$
orering	$x \sin(x)^6 - \frac{11 \cos(x) \sin(x)^5}{16} + \frac{49x(-6 \sin(x)^6 + 30 \sin(x)^4 \cos(x)^2)}{144} - \frac{5 \sin(x)^3 \cos(x)^3}{6} + \frac{7x(96 \sin(x)^6 - 840 \sin(x)^4 \cos(x)^2 + 252 \sin(x)^2 \cos(x)^4 - 35 \cos(x)^6)}{144}$

input
$$\text{int}(\sin(x)^6, x, \text{method}=_RETURNVERBOSE)$$

output
$$5/16*x - 1/192*\sin(6*x) + 3/64*\sin(4*x) - 15/64*\sin(2*x)$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.74

$$\int \sin^6(x) dx = -\frac{1}{48} (8 \cos(x)^5 - 26 \cos(x)^3 + 33 \cos(x)) \sin(x) + \frac{5}{16} x$$

input `integrate(sin(x)^6,x, algorithm="fricas")`output `-1/48*(8*cos(x)^5 - 26*cos(x)^3 + 33*cos(x))*sin(x) + 5/16*x`**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.06

$$\int \sin^6(x) dx = \frac{5x}{16} - \frac{\sin^5(x) \cos(x)}{6} - \frac{5 \sin^3(x) \cos(x)}{24} - \frac{5 \sin(x) \cos(x)}{16}$$

input `integrate(sin(x)**6,x)`output `5*x/16 - sin(x)**5*cos(x)/6 - 5*sin(x)**3*cos(x)/24 - 5*sin(x)*cos(x)/16`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.71

$$\int \sin^6(x) dx = \frac{1}{48} \sin(2x)^3 + \frac{5}{16} x + \frac{3}{64} \sin(4x) - \frac{1}{4} \sin(2x)$$

input `integrate(sin(x)^6,x, algorithm="maxima")`output `1/48*sin(2*x)^3 + 5/16*x + 3/64*sin(4*x) - 1/4*sin(2*x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.65

$$\int \sin^6(x) dx = \frac{5}{16} x - \frac{1}{192} \sin(6x) + \frac{3}{64} \sin(4x) - \frac{15}{64} \sin(2x)$$

input `integrate(sin(x)^6,x, algorithm="giac")`output `5/16*x - 1/192*sin(6*x) + 3/64*sin(4*x) - 15/64*sin(2*x)`**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.65

$$\int \sin^6(x) dx = \frac{5x}{16} - \frac{15 \sin(2x)}{64} + \frac{3 \sin(4x)}{64} - \frac{\sin(6x)}{192}$$

input `int(sin(x)^6,x)`output `(5*x)/16 - (15*sin(2*x))/64 + (3*sin(4*x))/64 - sin(6*x)/192`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.76

$$\int \sin^6(x) dx = -\frac{\cos(x) \sin(x)^5}{6} - \frac{5 \cos(x) \sin(x)^3}{24} - \frac{5 \cos(x) \sin(x)}{16} + \frac{5x}{16}$$

input `int(sin(x)^6,x)`output `(- 8*cos(x)*sin(x)**5 - 10*cos(x)*sin(x)**3 - 15*cos(x)*sin(x) + 15*x)/48`

3.33 $\int x \sin^2(x) dx$

Optimal result	252
Mathematica [A] (verified)	252
Rubi [A] (verified)	253
Maple [A] (verified)	254
Fricas [A] (verification not implemented)	254
Sympy [A] (verification not implemented)	255
Maxima [A] (verification not implemented)	255
Giac [A] (verification not implemented)	255
Mupad [B] (verification not implemented)	256
Reduce [B] (verification not implemented)	256

Optimal result

Integrand size = 6, antiderivative size = 25

$$\int x \sin^2(x) dx = \frac{x^2}{4} - \frac{1}{2}x \cos(x) \sin(x) + \frac{\sin^2(x)}{4}$$

output

```
1/4*x^2-1/2*x*cos(x)*sin(x)+1/4*sin(x)^2
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int x \sin^2(x) dx = \frac{x^2}{4} - \frac{1}{8} \cos(2x) - \frac{1}{4} x \sin(2x)$$

input

```
Integrate[x*Sin[x]^2,x]
```

output

```
x^2/4 - Cos[2*x]/8 - (x*Sin[2*x])/4
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 3791, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x \sin^2(x) dx \\ & \quad \downarrow \text{3042} \\ & \int x \sin(x)^2 dx \\ & \quad \downarrow \text{3791} \\ & \frac{\int x dx}{2} + \frac{\sin^2(x)}{4} - \frac{1}{2} x \sin(x) \cos(x) \\ & \quad \downarrow \text{15} \\ & \frac{x^2}{4} + \frac{\sin^2(x)}{4} - \frac{1}{2} x \sin(x) \cos(x) \end{aligned}$$

input `Int[x*Sin[x]^2,x]`

output `x^2/4 - (x*Cos[x]*Sin[x])/2 + Sin[x]^2/4`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3791

```
Int[((c_.) + (d_.)*(x_))*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :=
Simp[d*((b*SIN[e + f*x])^n/(f^2*n^2)), x] + (-Simp[b*(c + d*x)*Cos[e + f*x]
]*((b*SIN[e + f*x])^(n - 1)/(f*n)), x] + Simp[b^2*((n - 1)/n Int[(c + d*
x)*(b*SIN[e + f*x])^(n - 2), x], x)] /; FreeQ[{b, c, d, e, f}, x] && GtQ[n,
1]
```

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

method	result	size
risch	$\frac{x^2}{4} - \frac{\cos(2x)}{8} - \frac{x \sin(2x)}{4}$	20
default	$x \left(\frac{x}{2} - \frac{\cos(x) \sin(x)}{2} \right) - \frac{x^2}{4} + \frac{\sin(x)^2}{4}$	25
meijerg	$\frac{\sqrt{\pi} \left(\frac{2x^2+1}{2\sqrt{\pi}} - \frac{\cos(2x)}{2\sqrt{\pi}} - \frac{x \sin(2x)}{\sqrt{\pi}} \right)}{4}$	38
orering	$\frac{(2x^2+3) \sin(x)^2}{4} - \frac{\sin(x)^2}{2} - x \cos(x) \sin(x) + \frac{x(4 \cos(x) \sin(x) - 2x \sin(x)^2 + 2x \cos(x)^2)}{8}$	52
norman	$\frac{\tan\left(\frac{x}{2}\right)^2 + \tan\left(\frac{x}{2}\right)^3 x + \frac{x^2}{4} - x \tan\left(\frac{x}{2}\right) + \frac{x^2 \tan\left(\frac{x}{2}\right)^2}{2} + \frac{x^2 \tan\left(\frac{x}{2}\right)^4}{4}}{\left(1 + \tan\left(\frac{x}{2}\right)^2\right)^2}$	61

input

```
int(x*sin(x)^2,x,method=_RETURNVERBOSE)
```

output

```
1/4*x^2-1/8*cos(2*x)-1/4*x*sin(2*x)
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int x \sin^2(x) dx = -\frac{1}{2} x \cos(x) \sin(x) + \frac{1}{4} x^2 - \frac{1}{4} \cos(x)^2$$

input

```
integrate(x*sin(x)^2,x, algorithm="fricas")
```

output

```
-1/2*x*cos(x)*sin(x) + 1/4*x^2 - 1/4*cos(x)^2
```

Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.44

$$\int x \sin^2(x) dx = \frac{x^2 \sin^2(x)}{4} + \frac{x^2 \cos^2(x)}{4} - \frac{x \sin(x) \cos(x)}{2} - \frac{\cos^2(x)}{4}$$

input `integrate(x*sin(x)**2,x)`

output `x**2*sin(x)**2/4 + x**2*cos(x)**2/4 - x*sin(x)*cos(x)/2 - cos(x)**2/4`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int x \sin^2(x) dx = \frac{1}{4} x^2 - \frac{1}{4} x \sin(2x) - \frac{1}{8} \cos(2x)$$

input `integrate(x*sin(x)^2,x, algorithm="maxima")`

output `1/4*x^2 - 1/4*x*sin(2*x) - 1/8*cos(2*x)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int x \sin^2(x) dx = \frac{1}{4} x^2 - \frac{1}{4} x \sin(2x) - \frac{1}{8} \cos(2x)$$

input `integrate(x*sin(x)^2,x, algorithm="giac")`

output `1/4*x^2 - 1/4*x*sin(2*x) - 1/8*cos(2*x)`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int x \sin^2(x) dx = \frac{\sin(x)^2}{4} - \frac{x \sin(2x)}{4} + \frac{x^2}{4}$$

input

`int(x*sin(x)^2,x)`

output

`sin(x)^2/4 - (x*sin(2*x))/4 + x^2/4`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int x \sin^2(x) dx = -\frac{\cos(x) \sin(x) x}{2} + \frac{\sin(x)^2}{4} + \frac{x^2}{4} - \frac{1}{2}$$

input

`int(x*sin(x)^2,x)`

output

`(- 2*cos(x)*sin(x)*x + sin(x)**2 + x**2 - 2)/4`

3.34 $\int x \sin^3(x) dx$

Optimal result	257
Mathematica [A] (verified)	257
Rubi [A] (verified)	258
Maple [A] (verified)	259
Fricas [A] (verification not implemented)	260
Sympy [A] (verification not implemented)	260
Maxima [A] (verification not implemented)	260
Giac [A] (verification not implemented)	261
Mupad [B] (verification not implemented)	261
Reduce [B] (verification not implemented)	261

Optimal result

Integrand size = 6, antiderivative size = 33

$$\int x \sin^3(x) dx = -\frac{2}{3}x \cos(x) + \frac{2 \sin(x)}{3} - \frac{1}{3}x \cos(x) \sin^2(x) + \frac{\sin^3(x)}{9}$$

output

```
-2/3*x*cos(x)+2/3*sin(x)-1/3*x*cos(x)*sin(x)^2+1/9*sin(x)^3
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.94

$$\int x \sin^3(x) dx = -\frac{3}{4}x \cos(x) + \frac{1}{12}x \cos(3x) + \frac{3 \sin(x)}{4} - \frac{1}{36} \sin(3x)$$

input

```
Integrate[x*Sin[x]^3,x]
```

output

```
(-3*x*Cos[x])/4 + (x*Cos[3*x])/12 + (3*Sin[x])/4 - Sin[3*x]/36
```

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.97, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 3791, 3042, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \sin^3(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x \sin(x)^3 dx \\
 & \quad \downarrow \text{3791} \\
 & \frac{2}{3} \int x \sin(x) dx + \frac{\sin^3(x)}{9} - \frac{1}{3} x \sin^2(x) \cos(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{2}{3} \int x \sin(x) dx + \frac{\sin^3(x)}{9} - \frac{1}{3} x \sin^2(x) \cos(x) \\
 & \quad \downarrow \text{3777} \\
 & \frac{2}{3} \left(\int \cos(x) dx - x \cos(x) \right) + \frac{\sin^3(x)}{9} - \frac{1}{3} x \sin^2(x) \cos(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{2}{3} \left(\int \sin \left(x + \frac{\pi}{2} \right) dx - x \cos(x) \right) + \frac{\sin^3(x)}{9} - \frac{1}{3} x \sin^2(x) \cos(x) \\
 & \quad \downarrow \text{3117} \\
 & \frac{\sin^3(x)}{9} - \frac{1}{3} x \sin^2(x) \cos(x) + \frac{2}{3} (\sin(x) - x \cos(x))
 \end{aligned}$$

input

Int [x*Sin [x]^3, x]

output

-1/3*(x*Cos [x]*Sin [x]^2) + Sin [x]^3/9 + (2*(-(x*Cos [x]) + Sin [x]))/3

Definitions of rubi rules used

rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinear}$
 $\text{Q}[u, x]$

rule 3117 $\text{Int}[\sin[\text{Pi}/2 + (c_.) + (d_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[\text{Sin}[c + d*x]/d, x] \text{ ;}$
 $\text{FreeQ}[\{c, d\}, x]$

rule 3777 $\text{Int}[((c_.) + (d_.)*(x_))^m*\sin[(e_.) + (f_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[($
 $-(c + d*x)^m*(\text{Cos}[e + f*x]/f), x] + \text{Simp}[d*(m/f) \text{ Int}[(c + d*x)^{m-1}*C$
 $\text{os}[e + f*x], x], x] \text{ ; FreeQ}[\{c, d, e, f\}, x] \ \&\& \ \text{GtQ}[m, 0]$

rule 3791 $\text{Int}[((c_.) + (d_.)*(x_))*((b_.)*\sin[(e_.) + (f_.)*(x_)])^n, x_Symbol] \rightarrow$
 $\text{Simp}[d*((b*\text{Sin}[e + f*x])^n/(f^2*n^2)), x] + (-\text{Simp}[b*(c + d*x)*\text{Cos}[e + f*x]$
 $]*(b*\text{Sin}[e + f*x])^{n-1}/(f*n), x] + \text{Simp}[b^2*((n-1)/n) \text{ Int}[(c + d*$
 $x)*(b*\text{Sin}[e + f*x])^{n-2}, x], x] \text{ ; FreeQ}[\{b, c, d, e, f\}, x] \ \&\& \ \text{GtQ}[n,$
 $1]$

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.70

method	result
default	$-\frac{x(2+\sin(x)^2)\cos(x)}{3} + \frac{\sin(x)^3}{9} + \frac{2\sin(x)}{3}$
risch	$-\frac{3x\cos(x)}{4} + \frac{3\sin(x)}{4} + \frac{x\cos(3x)}{12} - \frac{\sin(3x)}{36}$
parallelrisch	$-\frac{3x\cos(x)}{4} + \frac{3\sin(x)}{4} + \frac{x\cos(3x)}{12} - \frac{\sin(3x)}{36}$
norman	$\frac{-\frac{2x}{3} + \frac{32\tan(\frac{x}{2})^3}{9} + \frac{4\tan(\frac{x}{2})^5}{3} - 2x\tan(\frac{x}{2})^2 + 2x\tan(\frac{x}{2})^4 + \frac{2x\tan(\frac{x}{2})^6}{3} + \frac{4\tan(\frac{x}{2})}{3}}{(1+\tan(\frac{x}{2})^2)^3}$
orering	$\frac{4(5x^2+2)\sin(x)^3}{9x^2} - \frac{2(5x^2+4)(\sin(x)^3+3x\cos(x)\sin(x)^2)}{9x^2} + \frac{8\sin(x)^2\cos(x)}{3} - \frac{4x\sin(x)^3}{3} + \frac{8x\cos(x)^2\sin(x)}{3} - 2\sin(x)$

input $\text{int}(x*\sin(x)^3, x, \text{method}=_RETURNVERBOSE)$

output `-1/3*x*(2+sin(x)^2)*cos(x)+1/9*sin(x)^3+2/3*sin(x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.70

$$\int x \sin^3(x) dx = \frac{1}{3} x \cos(x)^3 - x \cos(x) - \frac{1}{9} (\cos(x)^2 - 7) \sin(x)$$

input `integrate(x*sin(x)^3,x, algorithm="fricas")`

output `1/3*x*cos(x)^3 - x*cos(x) - 1/9*(cos(x)^2 - 7)*sin(x)`

Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.18

$$\int x \sin^3(x) dx = -x \sin^2(x) \cos(x) - \frac{2x \cos^3(x)}{3} + \frac{7 \sin^3(x)}{9} + \frac{2 \sin(x) \cos^2(x)}{3}$$

input `integrate(x*sin(x)**3,x)`

output `-x*sin(x)**2*cos(x) - 2*x*cos(x)**3/3 + 7*sin(x)**3/9 + 2*sin(x)*cos(x)**2/3`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.70

$$\int x \sin^3(x) dx = \frac{1}{12} x \cos(3x) - \frac{3}{4} x \cos(x) - \frac{1}{36} \sin(3x) + \frac{3}{4} \sin(x)$$

input `integrate(x*sin(x)^3,x, algorithm="maxima")`

output $1/12*x*cos(3*x) - 3/4*x*cos(x) - 1/36*sin(3*x) + 3/4*sin(x)$

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.70

$$\int x \sin^3(x) dx = \frac{1}{12} x \cos(3x) - \frac{3}{4} x \cos(x) - \frac{1}{36} \sin(3x) + \frac{3}{4} \sin(x)$$

input `integrate(x*sin(x)^3,x, algorithm="giac")`

output $1/12*x*cos(3*x) - 3/4*x*cos(x) - 1/36*sin(3*x) + 3/4*sin(x)$

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.76

$$\int x \sin^3(x) dx = \frac{x \cos(x)^3}{3} - \frac{\sin(x) \cos(x)^2}{9} - x \cos(x) + \frac{7 \sin(x)}{9}$$

input `int(x*sin(x)^3,x)`

output $(7*\sin(x))/9 + (x*\cos(x)^3)/3 - (\cos(x)^2*\sin(x))/9 - x*\cos(x)$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.76

$$\int x \sin^3(x) dx = -\frac{\cos(x) \sin(x)^2 x}{3} - \frac{2 \cos(x) x}{3} + \frac{\sin(x)^3}{9} + \frac{2 \sin(x)}{3}$$

input `int(x*sin(x)^3,x)`

output $(-3*\cos(x)*\sin(x)**2*x - 6*\cos(x)*x + \sin(x)**3 + 6*\sin(x))/9$

3.35 $\int x^2 \sin^2(x) dx$

Optimal result	262
Mathematica [A] (verified)	262
Rubi [A] (verified)	263
Maple [C] (verified)	264
Fricas [A] (verification not implemented)	265
Sympy [A] (verification not implemented)	265
Maxima [A] (verification not implemented)	266
Giac [A] (verification not implemented)	266
Mupad [B] (verification not implemented)	267
Reduce [B] (verification not implemented)	267

Optimal result

Integrand size = 8, antiderivative size = 41

$$\int x^2 \sin^2(x) dx = -\frac{x}{4} + \frac{x^3}{6} + \frac{1}{4} \cos(x) \sin(x) - \frac{1}{2} x^2 \cos(x) \sin(x) + \frac{1}{2} x \sin^2(x)$$

output

```
-1/4*x+1/6*x^3+1/4*cos(x)*sin(x)-1/2*x^2*cos(x)*sin(x)+1/2*x*sin(x)^2
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.71

$$\int x^2 \sin^2(x) dx = \frac{1}{24} (4x^3 - 6x \cos(2x) + (3 - 6x^2) \sin(2x))$$

input

```
Integrate[x^2*Sin[x]^2,x]
```

output

```
(4*x^3 - 6*x*Cos[2*x] + (3 - 6*x^2)*Sin[2*x])/24
```

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.12, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3792, 15, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \sin^2(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x^2 \sin(x)^2 dx \\
 & \quad \downarrow \text{3792} \\
 & \frac{\int x^2 dx}{2} - \frac{1}{2} \int \sin^2(x) dx - \frac{1}{2} x^2 \sin(x) \cos(x) + \frac{1}{2} x \sin^2(x) \\
 & \quad \downarrow \text{15} \\
 & -\frac{1}{2} \int \sin^2(x) dx + \frac{x^3}{6} - \frac{1}{2} x^2 \sin(x) \cos(x) + \frac{1}{2} x \sin^2(x) \\
 & \quad \downarrow \text{3042} \\
 & -\frac{1}{2} \int \sin(x)^2 dx + \frac{x^3}{6} - \frac{1}{2} x^2 \sin(x) \cos(x) + \frac{1}{2} x \sin^2(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{1}{2} \left(\frac{1}{2} \sin(x) \cos(x) - \frac{\int 1 dx}{2} \right) + \frac{x^3}{6} - \frac{1}{2} x^2 \sin(x) \cos(x) + \frac{1}{2} x \sin^2(x) \\
 & \quad \downarrow \text{24} \\
 & \frac{x^3}{6} - \frac{1}{2} x^2 \sin(x) \cos(x) + \frac{1}{2} x \sin^2(x) + \frac{1}{2} \left(\frac{1}{2} \sin(x) \cos(x) - \frac{x}{2} \right)
 \end{aligned}$$

input `Int[x^2*Sin[x]^2,x]`

output `x^3/6 - (x^2*Cos[x]*Sin[x])/2 + (x*Sin[x]^2)/2 + (-1/2*x + (Cos[x]*Sin[x])/2)/2`

Defintions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 3792 `Int[((c_.) + (d_.)*(x_))^(m_)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[d*m*(c + d*x)^(m - 1)*((b*Sin[e + f*x])^n/(f^2*n^2)), x] + (-Simp[b*(c + d*x)^m*Cos[e + f*x]*((b*Sin[e + f*x])^(n - 1)/(f*n)), x] + Simp[b^2*((n - 1)/n) Int[(c + d*x)^m*(b*Sin[e + f*x])^(n - 2), x], x] - Simp[d^2*m*((m - 1)/(f^2*n^2)) Int[(c + d*x)^(m - 2)*(b*Sin[e + f*x])^n, x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 1]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3.

Time = 0.18 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.46

method	result
meijerg	$\frac{x^5 \operatorname{hypergeom}\left(\left[1, \frac{5}{2}\right], \left[\frac{3}{2}, 2, \frac{7}{2}\right], -x^2\right)}{5}$
risch	$\frac{x^3}{6} - \frac{x \cos(2x)}{4} - \frac{(2x^2-1) \sin(2x)}{8}$
default	$x^2 \left(\frac{x}{2} - \frac{\cos(x) \sin(x)}{2} \right) - \frac{x \cos(x)^2}{2} + \frac{\cos(x) \sin(x)}{4} + \frac{x}{4} - \frac{x^3}{3}$
norman	$\frac{x^2 \tan\left(\frac{x}{2}\right)^3 - \frac{x}{4} + \frac{x^3}{6} - \frac{\tan\left(\frac{x}{2}\right)^3}{2} + \frac{3x \tan\left(\frac{x}{2}\right)^2}{2} - \frac{x \tan\left(\frac{x}{2}\right)^4}{4} - x^2 \tan\left(\frac{x}{2}\right) + \frac{x^3 \tan\left(\frac{x}{2}\right)^2}{3} + \frac{x^3 \tan\left(\frac{x}{2}\right)^4}{6} + \frac{\tan\left(\frac{x}{2}\right)}{2}}{\left(1 + \tan\left(\frac{x}{2}\right)\right)^2}$
orering	$\frac{(x^4+3x^2-3) \sin(x)^2}{3x} - \frac{(14x^2-15)(2x \sin(x)^2+2x^2 \cos(x) \sin(x))}{24x^2} + \frac{(2x^2-3)(2 \sin(x)^2+8x \cos(x) \sin(x)-2x^2 \sin(x)^2+2x^2 \cos(x)^2)}{24x}$

input `int(x^2*sin(x)^2,x,method=_RETURNVERBOSE)`

output `1/5*x^5*hypergeom([1, 5/2], [3/2, 2, 7/2], -x^2)`

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.71

$$\int x^2 \sin^2(x) dx = \frac{1}{6} x^3 - \frac{1}{2} x \cos(x)^2 - \frac{1}{4} (2x^2 - 1) \cos(x) \sin(x) + \frac{1}{4} x$$

input `integrate(x^2*sin(x)^2,x, algorithm="fricas")`

output `1/6*x^3 - 1/2*x*cos(x)^2 - 1/4*(2*x^2 - 1)*cos(x)*sin(x) + 1/4*x`

Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.37

$$\int x^2 \sin^2(x) dx = \frac{x^3 \sin^2(x)}{6} + \frac{x^3 \cos^2(x)}{6} - \frac{x^2 \sin(x) \cos(x)}{2} + \frac{x \sin^2(x)}{4} - \frac{x \cos^2(x)}{4} + \frac{\sin(x) \cos(x)}{4}$$

input `integrate(x**2*sin(x)**2,x)`

output `x**3*sin(x)**2/6 + x**3*cos(x)**2/6 - x**2*sin(x)*cos(x)/2 + x*sin(x)**2/4
- x*cos(x)**2/4 + sin(x)*cos(x)/4`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.63

$$\int x^2 \sin^2(x) dx = \frac{1}{6} x^3 - \frac{1}{4} x \cos(2x) - \frac{1}{8} (2x^2 - 1) \sin(2x)$$

input `integrate(x^2*sin(x)^2,x, algorithm="maxima")`

output `1/6*x^3 - 1/4*x*cos(2*x) - 1/8*(2*x^2 - 1)*sin(2*x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.63

$$\int x^2 \sin^2(x) dx = \frac{1}{6} x^3 - \frac{1}{4} x \cos(2x) - \frac{1}{8} (2x^2 - 1) \sin(2x)$$

input `integrate(x^2*sin(x)^2,x, algorithm="giac")`

output `1/6*x^3 - 1/4*x*cos(2*x) - 1/8*(2*x^2 - 1)*sin(2*x)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.68

$$\int x^2 \sin^2(x) dx = \frac{\sin(2x)}{8} - \frac{x \cos(2x)}{4} - \frac{x^2 \sin(2x)}{4} + \frac{x^3}{6}$$

input `int(x^2*sin(x)^2,x)`output `sin(2*x)/8 - (x*cos(2*x))/4 - (x^2*sin(2*x))/4 + x^3/6`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.76

$$\int x^2 \sin^2(x) dx = -\frac{\cos(x) \sin(x) x^2}{2} + \frac{\cos(x) \sin(x)}{4} + \frac{\sin(x)^2 x}{2} + \frac{x^3}{6} - \frac{x}{4}$$

input `int(x^2*sin(x)^2,x)`output `(- 6*cos(x)*sin(x)*x**2 + 3*cos(x)*sin(x) + 6*sin(x)**2*x + 2*x**3 - 3*x) /12`

3.36 $\int \cos^2(x) dx$

Optimal result	268
Mathematica [A] (verified)	268
Rubi [A] (verified)	269
Maple [A] (verified)	270
Fricas [A] (verification not implemented)	270
Sympy [A] (verification not implemented)	271
Maxima [A] (verification not implemented)	271
Giac [A] (verification not implemented)	271
Mupad [B] (verification not implemented)	272
Reduce [B] (verification not implemented)	272

Optimal result

Integrand size = 4, antiderivative size = 14

$$\int \cos^2(x) dx = \frac{x}{2} + \frac{1}{2} \cos(x) \sin(x)$$

output `1/2*x+1/2*cos(x)*sin(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \cos^2(x) dx = \frac{x}{2} + \frac{1}{4} \sin(2x)$$

input `Integrate[Cos[x]^2,x]`

output `x/2 + Sin[2*x]/4`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cos^2(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin\left(x + \frac{\pi}{2}\right)^2 dx \\ & \quad \downarrow \text{3115} \\ & \frac{\int 1 dx}{2} + \frac{1}{2} \sin(x) \cos(x) \\ & \quad \downarrow \text{24} \\ & \frac{x}{2} + \frac{1}{2} \sin(x) \cos(x) \end{aligned}$$

input `Int[Cos[x]^2,x]`

output `x/2 + (Cos[x]*Sin[x])/2`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.79

method	result	size
default	$\frac{x}{2} + \frac{\cos(x)\sin(x)}{2}$	11
risch	$\frac{x}{2} + \frac{\sin(2x)}{4}$	11
parallelrisch	$\frac{x}{2} + \frac{\sin(2x)}{4}$	11
orering	$x \cos(x)^2 + \frac{\cos(x)\sin(x)}{2} + \frac{x(-2\cos(x)^2 + 2\sin(x)^2)}{4}$	30
norman	$\frac{x \tan(\frac{x}{2})^2 + \frac{x}{2} - \tan(\frac{x}{2})^3 + \frac{x \tan(\frac{x}{2})^4}{2} + \tan(\frac{x}{2})}{(1 + \tan(\frac{x}{2})^2)^2}$	45

input `int(cos(x)^2,x,method=_RETURNVERBOSE)`

output `1/2*x+1/2*cos(x)*sin(x)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \cos^2(x) dx = \frac{1}{2} \cos(x) \sin(x) + \frac{1}{2} x$$

input `integrate(cos(x)^2,x, algorithm="fricas")`

output `1/2*cos(x)*sin(x) + 1/2*x`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \cos^2(x) dx = \frac{x}{2} + \frac{\sin(x) \cos(x)}{2}$$

input `integrate(cos(x)**2,x)`

output `x/2 + sin(x)*cos(x)/2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \cos^2(x) dx = \frac{1}{2}x + \frac{1}{4} \sin(2x)$$

input `integrate(cos(x)^2,x, algorithm="maxima")`

output `1/2*x + 1/4*sin(2*x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \cos^2(x) dx = \frac{1}{2}x + \frac{1}{4} \sin(2x)$$

input `integrate(cos(x)^2,x, algorithm="giac")`

output `1/2*x + 1/4*sin(2*x)`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \cos^2(x) dx = \frac{x}{2} + \frac{\sin(2x)}{4}$$

input `int(cos(x)^2,x)`

output `x/2 + sin(2*x)/4`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \cos^2(x) dx = \frac{\cos(x)\sin(x)}{2} + \frac{x}{2}$$

input `int(cos(x)^2,x)`

output `(cos(x)*sin(x) + x)/2`

3.37 $\int \cos^3(x) dx$

Optimal result	273
Mathematica [A] (verified)	273
Rubi [A] (verified)	274
Maple [A] (verified)	275
Fricas [A] (verification not implemented)	275
Sympy [A] (verification not implemented)	276
Maxima [A] (verification not implemented)	276
Giac [A] (verification not implemented)	276
Mupad [B] (verification not implemented)	277
Reduce [B] (verification not implemented)	277

Optimal result

Integrand size = 4, antiderivative size = 11

$$\int \cos^3(x) dx = \sin(x) - \frac{\sin^3(x)}{3}$$

output

```
sin(x)-1/3*sin(x)^3
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \cos^3(x) dx = \sin(x) - \frac{\sin^3(x)}{3}$$

input

```
Integrate[Cos[x]^3,x]
```

output

```
Sin[x] - Sin[x]^3/3
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3113, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \cos^3(x) dx \\
 \downarrow \text{3042} \\
 \int \sin\left(x + \frac{\pi}{2}\right)^3 dx \\
 \downarrow \text{3113} \\
 - \int (1 - \sin^2(x)) d(-\sin(x)) \\
 \downarrow \text{2009} \\
 \sin(x) - \frac{\sin^3(x)}{3}
 \end{array}$$

input `Int[Cos[x]^3,x]`

output `Sin[x] - Sin[x]^3/3`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113

```
Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
and[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x]
&& IGtQ[(n - 1)/2, 0]
```

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

method	result	size
default	$\frac{(2+\cos(x)^2)\sin(x)}{3}$	11
risch	$\frac{3\sin(x)}{4} + \frac{\sin(3x)}{12}$	12
parallelrisch	$\frac{3\sin(x)}{4} + \frac{\sin(3x)}{12}$	12
orering	$\sin(x)\cos(x)^2 + \frac{2\sin(x)^3}{3}$	15

input

```
int(cos(x)^3,x,method=_RETURNVERBOSE)
```

output

```
1/3*(2+cos(x)^2)*sin(x)
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

$$\int \cos^3(x) dx = \frac{1}{3} (\cos(x)^2 + 2) \sin(x)$$

input

```
integrate(cos(x)^3,x, algorithm="fricas")
```

output

```
1/3*(cos(x)^2 + 2)*sin(x)
```

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int \cos^3(x) dx = -\frac{\sin^3(x)}{3} + \sin(x)$$

input `integrate(cos(x)**3,x)`

output `-sin(x)**3/3 + sin(x)`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \cos^3(x) dx = -\frac{1}{3} \sin(x)^3 + \sin(x)$$

input `integrate(cos(x)^3,x, algorithm="maxima")`

output `-1/3*sin(x)^3 + sin(x)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \cos^3(x) dx = -\frac{1}{3} \sin(x)^3 + \sin(x)$$

input `integrate(cos(x)^3,x, algorithm="giac")`

output `-1/3*sin(x)^3 + sin(x)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \cos^3(x) dx = \sin(x) - \frac{\sin(x)^3}{3}$$

input `int(cos(x)^3,x)`

output `sin(x) - sin(x)^3/3`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

$$\int \cos^3(x) dx = \frac{\sin(x) (-\sin(x)^2 + 3)}{3}$$

input `int(cos(x)^3,x)`

output `(sin(x)*(- sin(x)**2 + 3))/3`

3.38 $\int \cos^4(x) dx$

Optimal result	278
Mathematica [A] (verified)	278
Rubi [A] (verified)	279
Maple [A] (verified)	280
Fricas [A] (verification not implemented)	281
Sympy [A] (verification not implemented)	281
Maxima [A] (verification not implemented)	281
Giac [A] (verification not implemented)	282
Mupad [B] (verification not implemented)	282
Reduce [B] (verification not implemented)	282

Optimal result

Integrand size = 4, antiderivative size = 24

$$\int \cos^4(x) dx = \frac{3x}{8} + \frac{3}{8} \cos(x) \sin(x) + \frac{1}{4} \cos^3(x) \sin(x)$$

output `3/8*x+3/8*cos(x)*sin(x)+1/4*cos(x)^3*sin(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \cos^4(x) dx = \frac{3x}{8} + \frac{1}{4} \sin(2x) + \frac{1}{32} \sin(4x)$$

input `Integrate[Cos[x]^4,x]`

output `(3*x)/8 + Sin[2*x]/4 + Sin[4*x]/32`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.21, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$, Rules used = {3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^4(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(x + \frac{\pi}{2}\right)^4 dx \\
 & \quad \downarrow \text{3115} \\
 & \frac{3}{4} \int \cos^2(x) dx + \frac{1}{4} \sin(x) \cos^3(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{4} \int \sin\left(x + \frac{\pi}{2}\right)^2 dx + \frac{1}{4} \sin(x) \cos^3(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{3}{4} \left(\frac{\int 1 dx}{2} + \frac{1}{2} \sin(x) \cos(x) \right) + \frac{1}{4} \sin(x) \cos^3(x) \\
 & \quad \downarrow \text{24} \\
 & \frac{1}{4} \sin(x) \cos^3(x) + \frac{3}{4} \left(\frac{x}{2} + \frac{1}{2} \sin(x) \cos(x) \right)
 \end{aligned}$$

input `Int[Cos[x]^4,x]`

output `(Cos[x]^3*Sin[x])/4 + (3*(x/2 + (Cos[x]*Sin[x])/2))/4`

Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.71

method	result
risch	$\frac{3x}{8} + \frac{\sin(4x)}{32} + \frac{\sin(2x)}{4}$
parallelrisch	$\frac{3x}{8} + \frac{\sin(4x)}{32} + \frac{\sin(2x)}{4}$
default	$\frac{(\cos(x)^3 + \frac{3\cos(x)}{2})\sin(x)}{4} + \frac{3x}{8}$
orering	$x \cos(x)^4 + \frac{5\cos(x)^3 \sin(x)}{8} + \frac{5x(12\sin(x)^2 \cos(x)^2 - 4\cos(x)^4)}{16} + \frac{3\cos(x)\sin(x)^3}{8} + \frac{x(-192\sin(x)^2 \cos(x)^2 + 40)}{64}$
norman	$\frac{\frac{3x}{8} - \frac{3\tan(\frac{x}{2})^3}{4} + \frac{3\tan(\frac{x}{2})^5}{4} - \frac{5\tan(\frac{x}{2})^7}{4} + \frac{3x\tan(\frac{x}{2})^2}{2} + \frac{9x\tan(\frac{x}{2})^4}{4} + \frac{3x\tan(\frac{x}{2})^6}{2} + \frac{3x\tan(\frac{x}{2})^8}{8} + \frac{5\tan(\frac{x}{2})}{4}}{(1+\tan(\frac{x}{2})^2)^4}$

input `int(cos(x)^4,x,method=_RETURNVERBOSE)`

output `3/8*x+1/32*sin(4*x)+1/4*sin(2*x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.79

$$\int \cos^4(x) dx = \frac{1}{8} (2 \cos(x)^3 + 3 \cos(x)) \sin(x) + \frac{3}{8} x$$

input `integrate(cos(x)^4,x, algorithm="fricas")`

output `1/8*(2*cos(x)^3 + 3*cos(x))*sin(x) + 3/8*x`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \cos^4(x) dx = \frac{3x}{8} + \frac{\sin(x) \cos^3(x)}{4} + \frac{3 \sin(x) \cos(x)}{8}$$

input `integrate(cos(x)**4,x)`

output `3*x/8 + sin(x)*cos(x)**3/4 + 3*sin(x)*cos(x)/8`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int \cos^4(x) dx = \frac{3}{8} x + \frac{1}{32} \sin(4x) + \frac{1}{4} \sin(2x)$$

input `integrate(cos(x)^4,x, algorithm="maxima")`

output `3/8*x + 1/32*sin(4*x) + 1/4*sin(2*x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int \cos^4(x) dx = \frac{3}{8}x + \frac{1}{32}\sin(4x) + \frac{1}{4}\sin(2x)$$

input `integrate(cos(x)^4,x, algorithm="giac")`output `3/8*x + 1/32*sin(4*x) + 1/4*sin(2*x)`**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int \cos^4(x) dx = \frac{3x}{8} + \frac{\sin(2x)}{4} + \frac{\sin(4x)}{32}$$

input `int(cos(x)^4,x)`output `(3*x)/8 + sin(2*x)/4 + sin(4*x)/32`**Reduce [B] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.75

$$\int \cos^4(x) dx = -\frac{\cos(x)\sin(x)^3}{4} + \frac{5\cos(x)\sin(x)}{8} + \frac{3x}{8}$$

input `int(cos(x)^4,x)`output `(- 2*cos(x)*sin(x)**3 + 5*cos(x)*sin(x) + 3*x)/8`

3.39 $\int (a^2 - x^2)^{5/2} dx$

Optimal result	283
Mathematica [A] (verified)	283
Rubi [A] (verified)	284
Maple [A] (verified)	285
Fricas [A] (verification not implemented)	286
Sympy [C] (verification not implemented)	286
Maxima [A] (verification not implemented)	287
Giac [A] (verification not implemented)	287
Mupad [B] (verification not implemented)	287
Reduce [B] (verification not implemented)	288

Optimal result

Integrand size = 13, antiderivative size = 84

$$\int (a^2 - x^2)^{5/2} dx = \frac{5}{16} a^4 x \sqrt{a^2 - x^2} + \frac{5}{24} a^2 x (a^2 - x^2)^{3/2} + \frac{1}{6} x (a^2 - x^2)^{5/2} + \frac{5}{16} a^6 \arctan\left(\frac{x}{\sqrt{a^2 - x^2}}\right)$$

output

```
5/24*a^2*x*(a^2-x^2)^(3/2)+1/6*x*(a^2-x^2)^(5/2)+5/16*a^6*arctan(x/(a^2-x^2)^(1/2))+5/16*a^4*x*(a^2-x^2)^(1/2)
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.73

$$\int (a^2 - x^2)^{5/2} dx = \frac{1}{48} \sqrt{a^2 - x^2} (33a^4 x - 26a^2 x^3 + 8x^5) + \frac{5}{16} a^6 \arctan\left(\frac{x}{\sqrt{a^2 - x^2}}\right)$$

input

```
Integrate[(a^2 - x^2)^(5/2), x]
```

output

```
(Sqrt[a^2 - x^2]*(33*a^4*x - 26*a^2*x^3 + 8*x^5))/48 + (5*a^6*ArcTan[x/Sqrt[a^2 - x^2]])/16
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.12, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {211, 211, 211, 224, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a^2 - x^2)^{5/2} dx$$

$$\downarrow 211$$

$$\frac{5}{6}a^2 \int (a^2 - x^2)^{3/2} dx + \frac{1}{6}x(a^2 - x^2)^{5/2}$$

$$\downarrow 211$$

$$\frac{5}{6}a^2 \left(\frac{3}{4}a^2 \int \sqrt{a^2 - x^2} dx + \frac{1}{4}x(a^2 - x^2)^{3/2} \right) + \frac{1}{6}x(a^2 - x^2)^{5/2}$$

$$\downarrow 211$$

$$\frac{5}{6}a^2 \left(\frac{3}{4}a^2 \left(\frac{1}{2}a^2 \int \frac{1}{\sqrt{a^2 - x^2}} dx + \frac{1}{2}x\sqrt{a^2 - x^2} \right) + \frac{1}{4}x(a^2 - x^2)^{3/2} \right) + \frac{1}{6}x(a^2 - x^2)^{5/2}$$

$$\downarrow 224$$

$$\frac{5}{6}a^2 \left(\frac{3}{4}a^2 \left(\frac{1}{2}a^2 \int \frac{1}{\frac{x^2}{a^2 - x^2} + 1} d\frac{x}{\sqrt{a^2 - x^2}} + \frac{1}{2}x\sqrt{a^2 - x^2} \right) + \frac{1}{4}x(a^2 - x^2)^{3/2} \right) + \frac{1}{6}x(a^2 - x^2)^{5/2}$$

$$\downarrow 216$$

$$\frac{5}{6}a^2 \left(\frac{3}{4}a^2 \left(\frac{1}{2}a^2 \arctan \left(\frac{x}{\sqrt{a^2 - x^2}} \right) + \frac{1}{2}x\sqrt{a^2 - x^2} \right) + \frac{1}{4}x(a^2 - x^2)^{3/2} \right) + \frac{1}{6}x(a^2 - x^2)^{5/2}$$

input `Int[(a^2 - x^2)^(5/2), x]`

output `(x*(a^2 - x^2)^(5/2))/6 + (5*a^2*((x*(a^2 - x^2)^(3/2))/4 + (3*a^2*((x*Sqr
t[a^2 - x^2])/2 + (a^2*ArcTan[x/Sqrt[a^2 - x^2]]/2))/4))/6`

Defintions of rubi rules used

rule 211 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[x*((a + b*x^2)^p/(2*p + 1)), x] + Simp[2*a*(p/(2*p + 1)) Int[(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.64

method	result	size
risch	$\frac{x(33a^4 - 26a^2x^2 + 8x^4)\sqrt{a^2 - x^2}}{48} + \frac{5a^6 \arctan\left(\frac{x}{\sqrt{a^2 - x^2}}\right)}{16}$	54
pseudoelliptic	$-\frac{5 \arctan\left(\frac{\sqrt{a^2 - x^2}}{x}\right)a^6}{16} + \frac{11\sqrt{a^2 - x^2} \left(a^4 - \frac{26}{33}a^2x^2 + \frac{8}{33}x^4\right)x}{16}$	54
default	$\frac{x(a^2 - x^2)^{\frac{5}{2}}}{6} + \frac{5a^2 \left(\frac{(a^2 - x^2)^{\frac{3}{2}}x}{4} + \frac{3a^2 \left(\frac{x\sqrt{a^2 - x^2}}{2} + \frac{a^2 \arctan\left(\frac{x}{\sqrt{a^2 - x^2}}\right)}{2} \right)}{4} \right)}{6}$	75

input `int((a^2-x^2)^(5/2), x, method=_RETURNVERBOSE)`

output `1/48*x*(33*a^4-26*a^2*x^2+8*x^4)*(a^2-x^2)^(1/2)+5/16*a^6*arctan(x/(a^2-x^2)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.71

$$\int (a^2 - x^2)^{5/2} dx = -\frac{5}{8} a^6 \arctan\left(-\frac{a - \sqrt{a^2 - x^2}}{x}\right) + \frac{1}{48} (33 a^4 x - 26 a^2 x^3 + 8 x^5) \sqrt{a^2 - x^2}$$

input `integrate((a^2-x^2)^(5/2),x, algorithm="fricas")`

output `-5/8*a^6*arctan(-(a - sqrt(a^2 - x^2))/x) + 1/48*(33*a^4*x - 26*a^2*x^3 + 8*x^5)*sqrt(a^2 - x^2)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 2.26 (sec) , antiderivative size = 180, normalized size of antiderivative = 2.14

$$\int (a^2 - x^2)^{5/2} dx = \begin{cases} -\frac{5ia^6 \operatorname{acosh}\left(\frac{x}{a}\right)}{16} - \frac{11ia^5 x}{16\sqrt{-1+\frac{x^2}{a^2}}} + \frac{59ia^3 x^3}{48\sqrt{-1+\frac{x^2}{a^2}}} - \frac{17iax^5}{24\sqrt{-1+\frac{x^2}{a^2}}} + \frac{ix^7}{6a\sqrt{-1+\frac{x^2}{a^2}}} & \text{for } \left|\frac{x^2}{a^2}\right| > 1 \\ \frac{5a^6 \operatorname{asin}\left(\frac{x}{a}\right)}{16} + \frac{11a^5 x \sqrt{1-\frac{x^2}{a^2}}}{16} - \frac{13a^3 x^3 \sqrt{1-\frac{x^2}{a^2}}}{24} + \frac{ax^5 \sqrt{1-\frac{x^2}{a^2}}}{6} & \text{otherwise} \end{cases}$$

input `integrate((a**2-x**2)**(5/2),x)`

output `Piecewise((-5*I*a**6*acosh(x/a)/16 - 11*I*a**5*x/(16*sqrt(-1 + x**2/a**2)) + 59*I*a**3*x**3/(48*sqrt(-1 + x**2/a**2)) - 17*I*a*x**5/(24*sqrt(-1 + x**2/a**2)) + I*x**7/(6*a*sqrt(-1 + x**2/a**2)), Abs(x**2/a**2) > 1), (5*a**6*asin(x/a)/16 + 11*a**5*x*sqrt(1 - x**2/a**2)/16 - 13*a**3*x**3*sqrt(1 - x**2/a**2)/24 + a*x**5*sqrt(1 - x**2/a**2)/6, True))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.71

$$\int (a^2 - x^2)^{5/2} dx = \frac{5}{16} a^6 \arcsin\left(\frac{x}{a}\right) + \frac{5}{16} \sqrt{a^2 - x^2} a^4 x + \frac{5}{24} (a^2 - x^2)^{3/2} a^2 x + \frac{1}{6} (a^2 - x^2)^{5/2} x$$

input `integrate((a^2-x^2)^(5/2),x, algorithm="maxima")`output `5/16*a^6*arcsin(x/a) + 5/16*sqrt(a^2 - x^2)*a^4*x + 5/24*(a^2 - x^2)^(3/2)*a^2*x + 1/6*(a^2 - x^2)^(5/2)*x`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.60

$$\int (a^2 - x^2)^{5/2} dx = \frac{5}{16} a^6 \arcsin\left(\frac{x}{a}\right) \operatorname{sgn}(a) + \frac{1}{48} (33 a^4 - 2 (13 a^2 - 4 x^2) x^2) \sqrt{a^2 - x^2} x$$

input `integrate((a^2-x^2)^(5/2),x, algorithm="giac")`output `5/16*a^6*arcsin(x/a)*sgn(a) + 1/48*(33*a^4 - 2*(13*a^2 - 4*x^2)*x^2)*sqrt(a^2 - x^2)*x`**Mupad [B] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.44

$$\int (a^2 - x^2)^{5/2} dx = \frac{x (a^2 - x^2)^{5/2} {}_2F_1\left(-\frac{5}{2}, \frac{1}{2}; \frac{3}{2}; \frac{x^2}{a^2}\right)}{\left(1 - \frac{x^2}{a^2}\right)^{5/2}}$$

input `int((a^2 - x^2)^(5/2),x)`

output $(x*(a^2 - x^2)^{(5/2)}*\text{hypergeom}([-5/2, 1/2], 3/2, x^2/a^2))/(1 - x^2/a^2)^{(5/2)}$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.73

$$\int (a^2 - x^2)^{5/2} dx = \frac{5a \sin\left(\frac{x}{a}\right) a^6}{16} + \frac{11\sqrt{a^2 - x^2} a^4 x}{16} - \frac{13\sqrt{a^2 - x^2} a^2 x^3}{24} + \frac{\sqrt{a^2 - x^2} x^5}{6}$$

input $\text{int}((a^2-x^2)^{(5/2)}, x)$

output $(15*\text{asin}(x/a)*a**6 + 33*\text{sqrt}(a**2 - x**2)*a**4*x - 26*\text{sqrt}(a**2 - x**2)*a**2*x**3 + 8*\text{sqrt}(a**2 - x**2)*x**5)/48$

3.40 $\int \frac{x^5}{\sqrt{5+x^2}} dx$

Optimal result	289
Mathematica [A] (verified)	289
Rubi [A] (verified)	290
Maple [A] (verified)	291
Fricas [A] (verification not implemented)	292
Sympy [A] (verification not implemented)	292
Maxima [A] (verification not implemented)	292
Giac [A] (verification not implemented)	293
Mupad [B] (verification not implemented)	293
Reduce [B] (verification not implemented)	293

Optimal result

Integrand size = 13, antiderivative size = 38

$$\int \frac{x^5}{\sqrt{5+x^2}} dx = 25\sqrt{5+x^2} - \frac{10}{3}(5+x^2)^{3/2} + \frac{1}{5}(5+x^2)^{5/2}$$

output `-10/3*(x^2+5)^(3/2)+1/5*(x^2+5)^(5/2)+25*(x^2+5)^(1/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.66

$$\int \frac{x^5}{\sqrt{5+x^2}} dx = \frac{1}{15}\sqrt{5+x^2}(200 - 20x^2 + 3x^4)$$

input `Integrate[x^5/Sqrt[5 + x^2],x]`

output `(Sqrt[5 + x^2]*(200 - 20*x^2 + 3*x^4))/15`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.11, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^5}{\sqrt{x^2+5}} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{2} \int \frac{x^4}{\sqrt{x^2+5}} dx^2 \\ & \quad \downarrow \text{53} \\ & \frac{1}{2} \int \left((x^2+5)^{3/2} - 10\sqrt{x^2+5} + \frac{25}{\sqrt{x^2+5}} \right) dx^2 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2} \left(\frac{2}{5} (x^2+5)^{5/2} - \frac{20}{3} (x^2+5)^{3/2} + 50\sqrt{x^2+5} \right) \end{aligned}$$

input `Int[x^5/Sqrt[5 + x^2], x]`

output `(50*Sqrt[5 + x^2] - (20*(5 + x^2)^(3/2))/3 + (2*(5 + x^2)^(5/2))/5)/2`

Defintions of rubi rules used

rule 53

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && IGtQ[m, 0] && ( !IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0])
|| LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

rule 243 $\text{Int}[(x_)^{(m_)}*((a_)+(b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{(m-1)/2}*(a+b*x)^p, x], x, x^2], x] /; \text{FreeQ}\{a, b, m, p\}, x \ \&\& \ \text{IntegerQ}[(m-1)/2]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.55

method	result	size
trager	$\sqrt{x^2+5} \left(\frac{1}{5}x^4 - \frac{4}{3}x^2 + \frac{40}{3} \right)$	21
gospers	$\frac{\sqrt{x^2+5} (3x^4 - 20x^2 + 200)}{15}$	22
risch	$\frac{\sqrt{x^2+5} (3x^4 - 20x^2 + 200)}{15}$	22
pseudoelliptic	$\frac{\sqrt{x^2+5} (3x^4 - 20x^2 + 200)}{15}$	22
orering	$\frac{\sqrt{x^2+5} (3x^4 - 20x^2 + 200)}{15}$	22
default	$\frac{x^4\sqrt{x^2+5}}{5} - \frac{4x^2\sqrt{x^2+5}}{3} + \frac{40\sqrt{x^2+5}}{3}$	35
meijerg	$\frac{25\sqrt{5} \left(-\frac{16\sqrt{\pi}}{15} + \frac{\sqrt{\pi} \left(\frac{6}{25}x^4 - \frac{8}{5}x^2 + 16 \right) \sqrt{1+\frac{x^2}{5}} \right)}{2\sqrt{\pi}}$	41

input `int(x^5/(x^2+5)^(1/2),x,method=_RETURNVERBOSE)`

output $(x^2+5)^{(1/2)}*(1/5*x^4-4/3*x^2+40/3)$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.55

$$\int \frac{x^5}{\sqrt{5+x^2}} dx = \frac{1}{15} (3x^4 - 20x^2 + 200)\sqrt{x^2+5}$$

input `integrate(x^5/(x^2+5)^(1/2),x, algorithm="fricas")`output `1/15*(3*x^4 - 20*x^2 + 200)*sqrt(x^2 + 5)`**Sympy [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.03

$$\int \frac{x^5}{\sqrt{5+x^2}} dx = \frac{x^4\sqrt{x^2+5}}{5} - \frac{4x^2\sqrt{x^2+5}}{3} + \frac{40\sqrt{x^2+5}}{3}$$

input `integrate(x**5/(x**2+5)**(1/2),x)`output `x**4*sqrt(x**2 + 5)/5 - 4*x**2*sqrt(x**2 + 5)/3 + 40*sqrt(x**2 + 5)/3`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.89

$$\int \frac{x^5}{\sqrt{5+x^2}} dx = \frac{1}{5} \sqrt{x^2+5}x^4 - \frac{4}{3} \sqrt{x^2+5}x^2 + \frac{40}{3} \sqrt{x^2+5}$$

input `integrate(x^5/(x^2+5)^(1/2),x, algorithm="maxima")`output `1/5*sqrt(x^2 + 5)*x^4 - 4/3*sqrt(x^2 + 5)*x^2 + 40/3*sqrt(x^2 + 5)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.74

$$\int \frac{x^5}{\sqrt{5+x^2}} dx = \frac{1}{5} (x^2+5)^{\frac{5}{2}} - \frac{10}{3} (x^2+5)^{\frac{3}{2}} + 25\sqrt{x^2+5}$$

input `integrate(x^5/(x^2+5)^(1/2),x, algorithm="giac")`

output `1/5*(x^2 + 5)^(5/2) - 10/3*(x^2 + 5)^(3/2) + 25*sqrt(x^2 + 5)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.53

$$\int \frac{x^5}{\sqrt{5+x^2}} dx = \sqrt{x^2+5} \left(\frac{x^4}{5} - \frac{4x^2}{3} + \frac{40}{3} \right)$$

input `int(x^5/(x^2 + 5)^(1/2),x)`

output `(x^2 + 5)^(1/2)*(x^4/5 - (4*x^2)/3 + 40/3)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.53

$$\int \frac{x^5}{\sqrt{5+x^2}} dx = \frac{\sqrt{x^2+5}(3x^4 - 20x^2 + 200)}{15}$$

input `int(x^5/(x^2+5)^(1/2),x)`

output `(sqrt(x**2 + 5)*(3*x**4 - 20*x**2 + 200))/15`

3.41 $\int \frac{t^3}{\sqrt{4+t^3}} dt$

Optimal result	294
Mathematica [C] (verified)	295
Rubi [A] (verified)	295
Maple [C] (verified)	296
Fricas [A] (verification not implemented)	297
Sympy [A] (verification not implemented)	298
Maxima [F]	298
Giac [F]	298
Mupad [B] (verification not implemented)	299
Reduce [F]	299

Optimal result

Integrand size = 13, antiderivative size = 172

$$\int \frac{t^3}{\sqrt{4+t^3}} dt = \frac{2}{5}t\sqrt{4+t^3} + \frac{8 \cdot 2^{2/3} \sqrt{2+\sqrt{3}} (2^{2/3}+t) \sqrt{\frac{2\sqrt[3]{2}-2^{2/3}t+t^2}{(2^{2/3}(1+\sqrt{3})+t)^2}} \operatorname{EllipticF}\left(\arcsin\left(\frac{2^{2/3}(1-\sqrt{3})+t}{2^{2/3}(1+\sqrt{3})+t}\right), -7-4\sqrt{3}\right)}{5\sqrt[4]{3} \sqrt{\frac{2^{2/3}+t}{(2^{2/3}(1+\sqrt{3})+t)^2}} \sqrt{4+t^3}}$$

output

```
2/5*t*(t^3+4)^(1/2)-8/15*2^(2/3)*(2^(2/3)+t)*EllipticF((t+2^(2/3)*(1-3^(1/2)))/(t+2^(2/3)*(1+3^(1/2))),I*3^(1/2)+2*I)*(1/2*6^(1/2)+1/2*2^(1/2))*((2*2^(1/3)-2^(2/3)*t+t^2)/(t+2^(2/3)*(1+3^(1/2)))^2)^(1/2)*3^(3/4)/(t^3+4)^(1/2)/((2^(2/3)+t)/(t+2^(2/3)*(1+3^(1/2))))^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 9.73 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.20

$$\int \frac{t^3}{\sqrt{4+t^3}} dt = \frac{2}{5}t \left(\sqrt{4+t^3} - 2 \operatorname{Hypergeometric2F1} \left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, -\frac{t^3}{4} \right) \right)$$

input `Integrate[t^3/Sqrt[4 + t^3],t]`

output `(2*t*(Sqrt[4 + t^3] - 2*Hypergeometric2F1[1/3, 1/2, 4/3, -1/4*t^3]))/5`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {843, 759}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{t^3}{\sqrt{t^3+4}} dt \\ & \quad \downarrow \text{843} \\ & \frac{2}{5}t\sqrt{t^3+4} - \frac{8}{5} \int \frac{1}{\sqrt{t^3+4}} dt \\ & \quad \downarrow \text{759} \\ & \frac{2}{5}t\sqrt{t^3+4} - \\ & \frac{8 \cdot 2^{2/3} \sqrt{2+\sqrt{3}} (t+2^{2/3}) \sqrt{\frac{t^2-2^{2/3}t+2\sqrt[3]{2}}{(t+2^{2/3})(1+\sqrt{3})^2}} \operatorname{EllipticF} \left(\arcsin \left(\frac{t+2^{2/3}(1-\sqrt{3})}{t+2^{2/3}(1+\sqrt{3})} \right), -7-4\sqrt{3} \right)}{5\sqrt[4]{3} \sqrt{\frac{t+2^{2/3}}{(t+2^{2/3})(1+\sqrt{3})^2}} \sqrt{t^3+4}} \end{aligned}$$

input `Int[t^3/Sqrt[4 + t^3],t]`

output

```
(2*t*Sqrt[4 + t^3])/5 - (8*2^(2/3)*Sqrt[2 + Sqrt[3]]*(2^(2/3) + t)*Sqrt[(2
*2^(1/3) - 2^(2/3)*t + t^2)/(2^(2/3)*(1 + Sqrt[3]) + t)^2]*EllipticF[ArcSi
n[(2^(2/3)*(1 - Sqrt[3]) + t)/(2^(2/3)*(1 + Sqrt[3]) + t)], -7 - 4*Sqrt[3]
])/((5*3^(1/4)*Sqrt[(2^(2/3) + t)/(2^(2/3)*(1 + Sqrt[3]) + t)^2]*Sqrt[4 + t
^3])
```

Defintions of rubi rules used

rule 759

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[2*Sqrt[2 + Sqrt[3]]*(s + r*x)*(Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 + Sqrt[3])*s + r*x)^2]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[s*
((s + r*x)/((1 + Sqrt[3])*s + r*x)^2)])*EllipticF[ArcSin[((1 - Sqrt[3])*s
+ r*x)/((1 + Sqrt[3])*s + r*x)], -7 - 4*Sqrt[3]], x] /; FreeQ[{a, b}, x] &
& PosQ[a]
```

rule 843

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n
- 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Simp[
a*c^n*(m - n + 1)/(b*(m + n*p + 1)) Int[(c*x)^(m - n)*(a + b*x^n)^p, x]
, x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*
p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4.

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.10

method	result
meijerg	$\frac{t^4 \operatorname{hypergeom}\left(\left[\frac{1}{2}, \frac{4}{3}\right], \left[\frac{7}{3}\right], -\frac{t^3}{4}\right)}{8}$
default	$\frac{2t\sqrt{t^3+4}}{5} + \frac{8i\sqrt{3}2^{\frac{2}{3}}\sqrt{i\left(t-\frac{2^{\frac{2}{3}}}{2}-\frac{i\sqrt{3}2^{\frac{2}{3}}}{2}\right)}\sqrt{3}2^{\frac{1}{3}}\sqrt{\frac{2^{\frac{2}{3}}+t}{\frac{3}{2}2^{\frac{2}{3}}+\frac{i\sqrt{3}2^{\frac{2}{3}}}{2}}}\sqrt{-i\left(t-\frac{2^{\frac{2}{3}}}{2}+\frac{i\sqrt{3}2^{\frac{2}{3}}}{2}\right)}\sqrt{3}2^{\frac{1}{3}}\operatorname{EllipticF}\left(\frac{\sqrt{6}\sqrt{i\left(t-\frac{2^{\frac{2}{3}}}{2}-\frac{i\sqrt{3}2^{\frac{2}{3}}}{2}\right)}}{6}\right)}{15\sqrt{t^3+4}}$
risch	$\frac{2t\sqrt{t^3+4}}{5} + \frac{8i\sqrt{3}2^{\frac{2}{3}}\sqrt{i\left(t-\frac{2^{\frac{2}{3}}}{2}-\frac{i\sqrt{3}2^{\frac{2}{3}}}{2}\right)}\sqrt{3}2^{\frac{1}{3}}\sqrt{\frac{2^{\frac{2}{3}}+t}{\frac{3}{2}2^{\frac{2}{3}}+\frac{i\sqrt{3}2^{\frac{2}{3}}}{2}}}\sqrt{-i\left(t-\frac{2^{\frac{2}{3}}}{2}+\frac{i\sqrt{3}2^{\frac{2}{3}}}{2}\right)}\sqrt{3}2^{\frac{1}{3}}\operatorname{EllipticF}\left(\frac{\sqrt{6}\sqrt{i\left(t-\frac{2^{\frac{2}{3}}}{2}-\frac{i\sqrt{3}2^{\frac{2}{3}}}{2}\right)}}{6}\right)}{15\sqrt{t^3+4}}$
elliptic	$\frac{2t\sqrt{t^3+4}}{5} + \frac{8i\sqrt{3}2^{\frac{2}{3}}\sqrt{i\left(t-\frac{2^{\frac{2}{3}}}{2}-\frac{i\sqrt{3}2^{\frac{2}{3}}}{2}\right)}\sqrt{3}2^{\frac{1}{3}}\sqrt{\frac{2^{\frac{2}{3}}+t}{\frac{3}{2}2^{\frac{2}{3}}+\frac{i\sqrt{3}2^{\frac{2}{3}}}{2}}}\sqrt{-i\left(t-\frac{2^{\frac{2}{3}}}{2}+\frac{i\sqrt{3}2^{\frac{2}{3}}}{2}\right)}\sqrt{3}2^{\frac{1}{3}}\operatorname{EllipticF}\left(\frac{\sqrt{6}\sqrt{i\left(t-\frac{2^{\frac{2}{3}}}{2}-\frac{i\sqrt{3}2^{\frac{2}{3}}}{2}\right)}}{6}\right)}{15\sqrt{t^3+4}}$

input `int(t^3/(t^3+4)^(1/2), t, method=_RETURNVERBOSE)`

output `1/8*t^4*hypergeom([1/2, 4/3], [7/3], -1/4*t^3)`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.10

$$\int \frac{t^3}{\sqrt{4+t^3}} dt = \frac{2}{5}\sqrt{t^3+4}t - \frac{16}{5}\operatorname{weierstrassPInverse}(0, -16, t)$$

input `integrate(t^3/(t^3+4)^(1/2), t, algorithm="fricas")`

output `2/5*sqrt(t^3 + 4)*t - 16/5*weierstrassPInverse(0, -16, t)`

Sympy [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.18

$$\int \frac{t^3}{\sqrt{4+t^3}} dt = \frac{t^4 \Gamma\left(\frac{4}{3}\right) {}_2F_1\left(\frac{1}{2}, \frac{4}{3} \mid \frac{7}{3} \mid \frac{t^3 e^{i\pi}}{4}\right)}{6\Gamma\left(\frac{7}{3}\right)}$$

input `integrate(t**3/(t**3+4)**(1/2),t)`output `t**4*gamma(4/3)*hyper((1/2, 4/3), (7/3,), t**3*exp_polar(I*pi)/4)/(6*gamma(7/3))`**Maxima [F]**

$$\int \frac{t^3}{\sqrt{4+t^3}} dt = \int \frac{t^3}{\sqrt{t^3+4}} dt$$

input `integrate(t^3/(t^3+4)^(1/2),t, algorithm="maxima")`output `integrate(t^3/sqrt(t^3 + 4), t)`**Giac [F]**

$$\int \frac{t^3}{\sqrt{4+t^3}} dt = \int \frac{t^3}{\sqrt{t^3+4}} dt$$

input `integrate(t^3/(t^3+4)^(1/2),t, algorithm="giac")`output `integrate(t^3/sqrt(t^3 + 4), t)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 301, normalized size of antiderivative = 1.75

$$\int \frac{t^3}{\sqrt{4+t^3}} dt = \frac{2t\sqrt{t^3+4}}{5} - \frac{16 \sqrt{-\frac{t-2^{2/3}(\frac{1}{2}+\frac{\sqrt{3}1i}{2})}{2^{2/3}+2^{2/3}(\frac{1}{2}+\frac{\sqrt{3}1i}{2})}} \sqrt{-\frac{t+2^{2/3}(-\frac{1}{2}+\frac{\sqrt{3}1i}{2})}{2^{2/3}-2^{2/3}(-\frac{1}{2}+\frac{\sqrt{3}1i}{2})}} \sqrt{\frac{t+2^{2/3}}{2^{2/3}+2^{2/3}(\frac{1}{2}+\frac{\sqrt{3}1i}{2})}} \left(2^{2/3}+2^{2/3}(\frac{1}{2}+\frac{\sqrt{3}1i}{2})\right)}{5 \sqrt{t^3 + \left(2^{2/3} + 2^{2/3} \left(-\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right) - 2^{2/3} \left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)\right) t^2 + \left(2 \cdot 2^{1/3} \left(-\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right) - 2 \cdot 2^{1/3} \left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)\right)}}$$

input `int(t^3/(t^3 + 4)^(1/2),t)`

output

```
(2*t*(t^3 + 4)^(1/2))/5 - (16*(-(t - 2^(2/3))*((3^(1/2)*1i)/2 + 1/2))/(2^(2/3) + 2^(2/3)*((3^(1/2)*1i)/2 + 1/2)))^(1/2)*(-(t + 2^(2/3))*((3^(1/2)*1i)/2 - 1/2))/(2^(2/3) - 2^(2/3)*((3^(1/2)*1i)/2 - 1/2)))^(1/2)*((t + 2^(2/3))/(2^(2/3) + 2^(2/3)*((3^(1/2)*1i)/2 + 1/2)))^(1/2)*(2^(2/3) + 2^(2/3)*((3^(1/2)*1i)/2 + 1/2))*ellipticF(asin(((t + 2^(2/3))/(2^(2/3) + 2^(2/3)*((3^(1/2)*1i)/2 + 1/2)))^(1/2)), (2^(2/3) + 2^(2/3)*((3^(1/2)*1i)/2 + 1/2))/(2^(2/3) - 2^(2/3)*((3^(1/2)*1i)/2 - 1/2)))/(5*(t^2*(2^(2/3) + 2^(2/3)*((3^(1/2)*1i)/2 - 1/2) - 2^(2/3)*((3^(1/2)*1i)/2 + 1/2)) - 4*((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2) + t^3 - t*(2*2^(1/3)*((3^(1/2)*1i)/2 + 1/2) - 2*2^(1/3)*((3^(1/2)*1i)/2 - 1/2) + 2*2^(1/3)*((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2)))^(1/2))
```

Reduce [F]

$$\int \frac{t^3}{\sqrt{4+t^3}} dt = \frac{2\sqrt{t^3+4}t}{5} - \frac{8\left(\int \frac{\sqrt{t^3+4}}{t^3+4} dt\right)}{5}$$

input `int(t^3/(t^3+4)^(1/2),t)`output `(2*(sqrt(t**3 + 4)*t - 4*int(sqrt(t**3 + 4)/(t**3 + 4),t)))/5`

3.42 $\int \tan^2(x) dx$

Optimal result	300
Mathematica [A] (verified)	300
Rubi [A] (verified)	301
Maple [A] (verified)	302
Fricas [A] (verification not implemented)	302
Sympy [B] (verification not implemented)	303
Maxima [A] (verification not implemented)	303
Giac [A] (verification not implemented)	303
Mupad [B] (verification not implemented)	304
Reduce [B] (verification not implemented)	304

Optimal result

Integrand size = 4, antiderivative size = 6

$$\int \tan^2(x) dx = -x + \tan(x)$$

output `-x+tan(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.33

$$\int \tan^2(x) dx = -\arctan(\tan(x)) + \tan(x)$$

input `Integrate[Tan[x]^2,x]`

output `-ArcTan[Tan[x]] + Tan[x]`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \tan^2(x) dx \\ \downarrow 3042 \\ \int \tan(x)^2 dx \\ \downarrow 3954 \\ \tan(x) - \int 1 dx \\ \downarrow 24 \\ \tan(x) - x \end{array}$$

input `Int [Tan [x]^2,x]`

output `-x + Tan [x]`

Defintions of rubi rules used

rule 24 `Int [a_, x_Symbol] :> Simp [a*x, x] /; FreeQ [a, x]`

rule 3042 `Int [u_, x_Symbol] :> Int [DeactivateTrig [u, x], x] /; FunctionOfTrigOfLinear Q [u, x]`

rule 3954

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d
*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x]
, x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
```

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.17

method	result	size
norman	$-x + \tan(x)$	7
parallelrisc	$-x + \tan(x)$	7
derivativedivides	$\tan(x) - \arctan(\tan(x))$	9
default	$\tan(x) - \arctan(\tan(x))$	9
risc	$-x + \frac{2i}{e^{2ix} + 1}$	17

input

```
int(tan(x)^2,x,method=_RETURNVERBOSE)
```

output

```
-x+tan(x)
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \tan^2(x) dx = -x + \tan(x)$$

input

```
integrate(tan(x)^2,x, algorithm="fricas")
```

output

```
-x + tan(x)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7 vs. $2(3) = 6$.

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.17

$$\int \tan^2(x) dx = -x + \frac{\sin(x)}{\cos(x)}$$

input `integrate(tan(x)**2,x)`

output `-x + sin(x)/cos(x)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \tan^2(x) dx = -x + \tan(x)$$

input `integrate(tan(x)^2,x, algorithm="maxima")`

output `-x + tan(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \tan^2(x) dx = -x + \tan(x)$$

input `integrate(tan(x)^2,x, algorithm="giac")`

output `-x + tan(x)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \tan^2(x) dx = \tan(x) - x$$

input `int(tan(x)^2,x)`

output `tan(x) - x`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \tan^2(x) dx = \tan(x) - x$$

input `int(tan(x)^2,x)`

output `tan(x) - x`

3.43 $\int \tan^4(x) dx$

Optimal result	305
Mathematica [A] (verified)	305
Rubi [A] (verified)	306
Maple [A] (verified)	307
Fricas [A] (verification not implemented)	308
Sympy [A] (verification not implemented)	308
Maxima [A] (verification not implemented)	308
Giac [A] (verification not implemented)	309
Mupad [B] (verification not implemented)	309
Reduce [B] (verification not implemented)	309

Optimal result

Integrand size = 4, antiderivative size = 14

$$\int \tan^4(x) dx = x - \tan(x) + \frac{\tan^3(x)}{3}$$

output

```
x-tan(x)+1/3*tan(x)^3
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \tan^4(x) dx = \arctan(\tan(x)) - \tan(x) + \frac{\tan^3(x)}{3}$$

input

```
Integrate[Tan[x]^4,x]
```

output

```
ArcTan[Tan[x]] - Tan[x] + Tan[x]^3/3
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$, Rules used = {3042, 3954, 3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^4(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x)^4 dx \\
 & \quad \downarrow \text{3954} \\
 & \frac{\tan^3(x)}{3} - \int \tan^2(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{\tan^3(x)}{3} - \int \tan(x)^2 dx \\
 & \quad \downarrow \text{3954} \\
 & \int 1 dx + \frac{\tan^3(x)}{3} - \tan(x) \\
 & \quad \downarrow \text{24} \\
 & x + \frac{\tan^3(x)}{3} - \tan(x)
 \end{aligned}$$

input `Int [Tan [x] ^4, x]`

output `x - Tan [x] + Tan [x] ^3/3`

Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

method	result	size
norman	$x - \tan(x) + \frac{\tan(x)^3}{3}$	13
parallelrisc	$x - \tan(x) + \frac{\tan(x)^3}{3}$	13
derivativedivides	$\frac{\tan(x)^3}{3} - \tan(x) + \arctan(\tan(x))$	15
default	$\frac{\tan(x)^3}{3} - \tan(x) + \arctan(\tan(x))$	15
risc	$x - \frac{4i(3e^{4ix} + 3e^{2ix} + 2)}{3(e^{2ix} + 1)^3}$	31

input `int(tan(x)^4,x,method=_RETURNVERBOSE)`

output `x-tan(x)+1/3*tan(x)^3`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \tan^4(x) dx = \frac{1}{3} \tan(x)^3 + x - \tan(x)$$

input `integrate(tan(x)^4,x, algorithm="fricas")`output `1/3*tan(x)^3 + x - tan(x)`**Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.36

$$\int \tan^4(x) dx = x + \frac{\sin^3(x)}{3 \cos^3(x)} - \frac{\sin(x)}{\cos(x)}$$

input `integrate(tan(x)**4,x)`output `x + sin(x)**3/(3*cos(x)**3) - sin(x)/cos(x)`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \tan^4(x) dx = \frac{1}{3} \tan(x)^3 + x - \tan(x)$$

input `integrate(tan(x)^4,x, algorithm="maxima")`output `1/3*tan(x)^3 + x - tan(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \tan^4(x) dx = \frac{1}{3} \tan(x)^3 + x - \tan(x)$$

input `integrate(tan(x)^4,x, algorithm="giac")`

output `1/3*tan(x)^3 + x - tan(x)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \tan^4(x) dx = \frac{\tan(x)^3}{3} - \tan(x) + x$$

input `int(tan(x)^4,x)`

output `x - tan(x) + tan(x)^3/3`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \tan^4(x) dx = \frac{\tan(x)^3}{3} - \tan(x) + x$$

input `int(tan(x)^4,x)`

output `(tan(x)**3 - 3*tan(x) + 3*x)/3`

3.44 $\int \cot^2(x) dx$

Optimal result	310
Mathematica [C] (verified)	310
Rubi [A] (verified)	311
Maple [A] (verified)	312
Fricas [B] (verification not implemented)	312
Sympy [A] (verification not implemented)	313
Maxima [A] (verification not implemented)	313
Giac [B] (verification not implemented)	313
Mupad [B] (verification not implemented)	314
Reduce [B] (verification not implemented)	314

Optimal result

Integrand size = 4, antiderivative size = 8

$$\int \cot^2(x) dx = -x - \cot(x)$$

output `-x-cot(x)`

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 2.25

$$\int \cot^2(x) dx = -\cot(x) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, -\tan^2(x)\right)$$

input `Integrate[Cot[x]^2,x]`

output `-(Cot[x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[x]^2])`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cot^2(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \tan\left(x + \frac{\pi}{2}\right)^2 dx \\ & \quad \downarrow \text{3954} \\ & - \int 1 dx - \cot(x) \\ & \quad \downarrow \text{24} \\ & -x - \cot(x) \end{aligned}$$

input `Int[Cot[x]^2,x]`

output `-x - Cot[x]`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d
*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x]
, x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
```

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.12

method	result	size
parallelrisch	$-x - \cot(x)$	9
norman	$\frac{-1-x \tan(x)}{\tan(x)}$	13
derivativedivides	$-\cot(x) + \frac{\pi}{2} - \operatorname{arccot}(\cot(x))$	14
default	$-\cot(x) + \frac{\pi}{2} - \operatorname{arccot}(\cot(x))$	14
risch	$-x - \frac{2i}{e^{2ix}-1}$	17

input

```
int(cot(x)^2,x,method=_RETURNVERBOSE)
```

output

```
-x-cot(x)
```

Fricas [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 20 vs. $2(8) = 16$.

Time = 0.09 (sec) , antiderivative size = 20, normalized size of antiderivative = 2.50

$$\int \cot^2(x) dx = -\frac{x \sin(2x) + \cos(2x) + 1}{\sin(2x)}$$

input

```
integrate(cot(x)^2,x, algorithm="fricas")
```

output

```
-(x*sin(2*x) + cos(2*x) + 1)/sin(2*x)
```

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \cot^2(x) dx = -x - \frac{\cos(x)}{\sin(x)}$$

input `integrate(cot(x)**2,x)`

output `-x - cos(x)/sin(x)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.25

$$\int \cot^2(x) dx = -x - \frac{1}{\tan(x)}$$

input `integrate(cot(x)^2,x, algorithm="maxima")`

output `-x - 1/tan(x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18 vs. $2(8) = 16$.

Time = 0.12 (sec) , antiderivative size = 18, normalized size of antiderivative = 2.25

$$\int \cot^2(x) dx = -x - \frac{1}{2 \tan\left(\frac{1}{2}x\right)} + \frac{1}{2} \tan\left(\frac{1}{2}x\right)$$

input `integrate(cot(x)^2,x, algorithm="giac")`

output `-x - 1/2/tan(1/2*x) + 1/2*tan(1/2*x)`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \cot^2(x) dx = -x - \cot(x)$$

input `int(cot(x)^2,x)`

output `- x - cot(x)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \cot^2(x) dx = -\cot(x) - x$$

input `int(cot(x)^2,x)`

output `- (cot(x) + x)`

3.45 $\int \cot^4(x) dx$

Optimal result	315
Mathematica [C] (verified)	315
Rubi [A] (verified)	316
Maple [A] (verified)	317
Fricas [B] (verification not implemented)	318
Sympy [A] (verification not implemented)	318
Maxima [A] (verification not implemented)	318
Giac [B] (verification not implemented)	319
Mupad [B] (verification not implemented)	319
Reduce [B] (verification not implemented)	319

Optimal result

Integrand size = 4, antiderivative size = 12

$$\int \cot^4(x) dx = x + \cot(x) - \frac{\cot^3(x)}{3}$$

output

```
x+cot(x)-1/3*cot(x)^3
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.00 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.83

$$\int \cot^4(x) dx = -\frac{1}{3} \cot^3(x) \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, -\tan^2(x)\right)$$

input

```
Integrate[Cot[x]^4,x]
```

output

```
-1/3*(Cot[x]^3*Hypergeometric2F1[-3/2, 1, -1/2, -Tan[x]^2])
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$, Rules used = {3042, 3954, 3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^4(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan\left(x + \frac{\pi}{2}\right)^4 dx \\
 & \quad \downarrow \text{3954} \\
 & - \int \cot^2(x) dx - \frac{1}{3} \cot^3(x) \\
 & \quad \downarrow \text{3042} \\
 & - \int \tan\left(x + \frac{\pi}{2}\right)^2 dx - \frac{1}{3} \cot^3(x) \\
 & \quad \downarrow \text{3954} \\
 & \int 1 dx - \frac{1}{3} \cot^3(x) + \cot(x) \\
 & \quad \downarrow \text{24} \\
 & x - \frac{1}{3} \cot^3(x) + \cot(x)
 \end{aligned}$$

input `Int[Cot[x]^4,x]`

output `x + Cot[x] - Cot[x]^3/3`

Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

method	result	size
parallelrisc	$x + \cot(x) - \frac{\cot(x)^3}{3}$	11
derivativedivides	$-\frac{\cot(x)^3}{3} + \cot(x) - \frac{\pi}{2} + \operatorname{arccot}(\cot(x))$	16
default	$-\frac{\cot(x)^3}{3} + \cot(x) - \frac{\pi}{2} + \operatorname{arccot}(\cot(x))$	16
norman	$\frac{-\frac{1}{3} + \tan(x)^2 + x \tan(x)^3}{\tan(x)^3}$	18
risch	$x + \frac{4i(3e^{4ix} - 3e^{2ix} + 2)}{3(e^{2ix} - 1)^3}$	31

input `int(cot(x)^4,x,method=_RETURNVERBOSE)`

output `x+cot(x)-1/3*cot(x)^3`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 48 vs. $2(10) = 20$.

Time = 0.10 (sec) , antiderivative size = 48, normalized size of antiderivative = 4.00

$$\int \cot^4(x) dx = \frac{4 \cos(2x)^2 + 3(x \cos(2x) - x) \sin(2x) + 2 \cos(2x) - 2}{3(\cos(2x) - 1) \sin(2x)}$$

input `integrate(cot(x)^4,x, algorithm="fricas")`

output `1/3*(4*cos(2*x)^2 + 3*(x*cos(2*x) - x)*sin(2*x) + 2*cos(2*x) - 2)/((cos(2*x) - 1)*sin(2*x))`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.58

$$\int \cot^4(x) dx = x + \frac{\cos(x)}{\sin(x)} - \frac{\cos^3(x)}{3 \sin^3(x)}$$

input `integrate(cot(x)**4,x)`

output `x + cos(x)/sin(x) - cos(x)**3/(3*sin(x)**3)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.33

$$\int \cot^4(x) dx = x + \frac{3 \tan(x)^2 - 1}{3 \tan(x)^3}$$

input `integrate(cot(x)^4,x, algorithm="maxima")`

output `x + 1/3*(3*tan(x)^2 - 1)/tan(x)^3`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 34 vs. $2(10) = 20$.

Time = 0.11 (sec) , antiderivative size = 34, normalized size of antiderivative = 2.83

$$\int \cot^4(x) dx = \frac{1}{24} \tan\left(\frac{1}{2}x\right)^3 + x + \frac{15 \tan\left(\frac{1}{2}x\right)^2 - 1}{24 \tan\left(\frac{1}{2}x\right)^3} - \frac{5}{8} \tan\left(\frac{1}{2}x\right)$$

input `integrate(cot(x)^4,x, algorithm="giac")`

output `1/24*tan(1/2*x)^3 + x + 1/24*(15*tan(1/2*x)^2 - 1)/tan(1/2*x)^3 - 5/8*tan(1/2*x)`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \cot^4(x) dx = -\frac{\cot(x)^3}{3} + \cot(x) + x$$

input `int(cot(x)^4,x)`

output `x + cot(x) - cot(x)^3/3`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \cot^4(x) dx = -\frac{\cot(x)^3}{3} + \cot(x) + x$$

input `int(cot(x)^4,x)`

output `(- cot(x)**3 + 3*cot(x) + 3*x)/3`

3.46 $\int (2 + 3x) \sin(5x) dx$

Optimal result	320
Mathematica [A] (verified)	320
Rubi [A] (verified)	321
Maple [A] (verified)	322
Fricas [A] (verification not implemented)	323
Sympy [A] (verification not implemented)	323
Maxima [A] (verification not implemented)	323
Giac [A] (verification not implemented)	324
Mupad [B] (verification not implemented)	324
Reduce [B] (verification not implemented)	324

Optimal result

Integrand size = 10, antiderivative size = 22

$$\int (2 + 3x) \sin(5x) dx = -\frac{1}{5}(2 + 3x) \cos(5x) + \frac{3}{25} \sin(5x)$$

output `-1/5*(2+3*x)*cos(5*x)+3/25*sin(5*x)`

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.18

$$\int (2 + 3x) \sin(5x) dx = -\frac{2}{5} \cos(5x) - \frac{3}{5}x \cos(5x) + \frac{3}{25} \sin(5x)$$

input `Integrate[(2 + 3*x)*Sin[5*x],x]`

output `(-2*Cos[5*x])/5 - (3*x*Cos[5*x])/5 + (3*Sin[5*x])/25`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (3x + 2) \sin(5x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int (3x + 2) \sin(5x) dx \\
 & \quad \downarrow \text{3777} \\
 & \frac{3}{5} \int \cos(5x) dx - \frac{1}{5} (3x + 2) \cos(5x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{5} \int \sin\left(5x + \frac{\pi}{2}\right) dx - \frac{1}{5} (3x + 2) \cos(5x) \\
 & \quad \downarrow \text{3117} \\
 & \frac{3}{25} \sin(5x) - \frac{1}{5} (3x + 2) \cos(5x)
 \end{aligned}$$

input `Int[(2 + 3*x)*Sin[5*x],x]`

output `-1/5*((2 + 3*x)*Cos[5*x]) + (3*Sin[5*x])/25`

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

method	result	size
risch	$\left(-\frac{2}{5} - \frac{3x}{5}\right) \cos(5x) + \frac{3 \sin(5x)}{25}$	18
orering	$-\frac{(2+3x) \cos(5x)}{5} + \frac{3 \sin(5x)}{25}$	19
parallelrisc	$-\frac{2}{5} + \frac{(-3x-2) \cos(5x)}{5} + \frac{3 \sin(5x)}{25}$	20
derivativedivides	$-\frac{2 \cos(5x)}{5} + \frac{3 \sin(5x)}{25} - \frac{3 \cos(5x)x}{5}$	21
default	$-\frac{2 \cos(5x)}{5} + \frac{3 \sin(5x)}{25} - \frac{3 \cos(5x)x}{5}$	21
parts	$-\frac{2 \cos(5x)}{5} + \frac{3 \sin(5x)}{25} - \frac{3 \cos(5x)x}{5}$	21
norman	$\frac{-\frac{3x}{5} + \frac{3x \tan\left(\frac{5x}{2}\right)^2}{5} + \frac{6 \tan\left(\frac{5x}{2}\right)}{25} - \frac{4}{5}}{1 + \tan\left(\frac{5x}{2}\right)^2}$	32
meijerg	$\frac{2\sqrt{\pi} \left(\frac{1}{\sqrt{\pi}} - \frac{\cos(5x)}{\sqrt{\pi}}\right)}{5} + \frac{6\sqrt{\pi} \left(-\frac{5x \cos(5x)}{2\sqrt{\pi}} + \frac{\sin(5x)}{2\sqrt{\pi}}\right)}{25}$	45

input `int((2+3*x)*sin(5*x), x, method=_RETURNVERBOSE)`

output `(-2/5-3/5*x)*cos(5*x)+3/25*sin(5*x)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int (2 + 3x) \sin(5x) dx = -\frac{1}{5} (3x + 2) \cos(5x) + \frac{3}{25} \sin(5x)$$

input `integrate((2+3*x)*sin(5*x),x, algorithm="fricas")`output `-1/5*(3*x + 2)*cos(5*x) + 3/25*sin(5*x)`**Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.18

$$\int (2 + 3x) \sin(5x) dx = -\frac{3x \cos(5x)}{5} + \frac{3 \sin(5x)}{25} - \frac{2 \cos(5x)}{5}$$

input `integrate((2+3*x)*sin(5*x),x)`output `-3*x*cos(5*x)/5 + 3*sin(5*x)/25 - 2*cos(5*x)/5`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int (2 + 3x) \sin(5x) dx = -\frac{3}{5} x \cos(5x) - \frac{2}{5} \cos(5x) + \frac{3}{25} \sin(5x)$$

input `integrate((2+3*x)*sin(5*x),x, algorithm="maxima")`output `-3/5*x*cos(5*x) - 2/5*cos(5*x) + 3/25*sin(5*x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int (2 + 3x) \sin(5x) dx = -\frac{1}{5} (3x + 2) \cos(5x) + \frac{3}{25} \sin(5x)$$

input `integrate((2+3*x)*sin(5*x),x, algorithm="giac")`

output `-1/5*(3*x + 2)*cos(5*x) + 3/25*sin(5*x)`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int (2 + 3x) \sin(5x) dx = \frac{3 \sin(5x)}{25} - \frac{2 \cos(5x)}{5} - \frac{3x \cos(5x)}{5}$$

input `int(sin(5*x)*(3*x + 2),x)`

output `(3*sin(5*x))/25 - (2*cos(5*x))/5 - (3*x*cos(5*x))/5`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int (2 + 3x) \sin(5x) dx = -\frac{3 \cos(5x) x}{5} - \frac{2 \cos(5x)}{5} + \frac{3 \sin(5x)}{25}$$

input `int((2+3*x)*sin(5*x),x)`

output `(- 15*cos(5*x)*x - 10*cos(5*x) + 3*sin(5*x))/25`

3.47 $\int x\sqrt{1+x^2} dx$

Optimal result	325
Mathematica [A] (verified)	325
Rubi [A] (verified)	326
Maple [A] (verified)	327
Fricas [A] (verification not implemented)	327
Sympy [B] (verification not implemented)	328
Maxima [A] (verification not implemented)	328
Giac [A] (verification not implemented)	328
Mupad [B] (verification not implemented)	329
Reduce [B] (verification not implemented)	329

Optimal result

Integrand size = 11, antiderivative size = 13

$$\int x\sqrt{1+x^2} dx = \frac{1}{3}(1+x^2)^{3/2}$$

output `1/3*(x^2+1)^(3/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int x\sqrt{1+x^2} dx = \frac{1}{3}(1+x^2)^{3/2}$$

input `Integrate[x*Sqrt[1 + x^2],x]`

output `(1 + x^2)^(3/2)/3`

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x\sqrt{x^2+1} dx$$

$$\downarrow 241$$

$$\frac{1}{3}(x^2+1)^{3/2}$$

input

```
Int[x*Sqrt[1 + x^2],x]
```

output

```
(1 + x^2)^(3/2)/3
```

Defintions of rubi rules used

rule 241

```
Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(a + b*x^2)^(p + 1)/
(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]
```

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

method	result	size
gospers	$\frac{(x^2+1)^{\frac{3}{2}}}{3}$	10
derivativedivides	$\frac{(x^2+1)^{\frac{3}{2}}}{3}$	10
default	$\frac{(x^2+1)^{\frac{3}{2}}}{3}$	10
risch	$\frac{(x^2+1)^{\frac{3}{2}}}{3}$	10
pseudoelliptic	$\frac{(x^2+1)^{\frac{3}{2}}}{3}$	10
orering	$\frac{(x^2+1)^{\frac{3}{2}}}{3}$	10
trager	$\left(\frac{x^2}{3} + \frac{1}{3}\right) \sqrt{x^2 + 1}$	16
meijerg	$-\frac{\frac{4\sqrt{\pi}}{3} - \frac{2\sqrt{\pi}(2x^2+2)\sqrt{x^2+1}}{3}}{4\sqrt{\pi}}$	31

input `int(x*(x^2+1)^(1/2),x,method=_RETURNVERBOSE)`output `1/3*(x^2+1)^(3/2)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int x\sqrt{1+x^2} dx = \frac{1}{3} (x^2 + 1)^{\frac{3}{2}}$$

input `integrate(x*(x^2+1)^(1/2),x, algorithm="fricas")`output `1/3*(x^2 + 1)^(3/2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 22 vs. $2(8) = 16$.

Time = 0.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.69

$$\int x\sqrt{1+x^2} dx = \frac{x^2\sqrt{x^2+1}}{3} + \frac{\sqrt{x^2+1}}{3}$$

input `integrate(x*(x**2+1)**(1/2),x)`

output `x**2*sqrt(x**2 + 1)/3 + sqrt(x**2 + 1)/3`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int x\sqrt{1+x^2} dx = \frac{1}{3} (x^2 + 1)^{\frac{3}{2}}$$

input `integrate(x*(x^2+1)^(1/2),x, algorithm="maxima")`

output `1/3*(x^2 + 1)^(3/2)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int x\sqrt{1+x^2} dx = \frac{1}{3} (x^2 + 1)^{\frac{3}{2}}$$

input `integrate(x*(x^2+1)^(1/2),x, algorithm="giac")`

output `1/3*(x^2 + 1)^(3/2)`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int x\sqrt{1+x^2} dx = \frac{(x^2+1)^{3/2}}{3}$$

input `int(x*(x^2 + 1)^(1/2),x)`

output `(x^2 + 1)^(3/2)/3`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int x\sqrt{1+x^2} dx = \frac{\sqrt{x^2+1}(x^2+1)}{3}$$

input `int(x*(x^2+1)^(1/2),x)`

output `(sqrt(x**2 + 1)*(x**2 + 1))/3`

3.48 $\int x(-1 + x^2)^9 dx$

Optimal result	330
Mathematica [A] (verified)	330
Rubi [A] (verified)	331
Maple [A] (verified)	332
Fricas [B] (verification not implemented)	332
Sympy [B] (verification not implemented)	333
Maxima [A] (verification not implemented)	333
Giac [A] (verification not implemented)	333
Mupad [B] (verification not implemented)	334
Reduce [B] (verification not implemented)	334

Optimal result

Integrand size = 9, antiderivative size = 13

$$\int x(-1 + x^2)^9 dx = \frac{1}{20}(1 - x^2)^{10}$$

output `1/20*(-x^2+1)^10`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int x(-1 + x^2)^9 dx = \frac{1}{20}(-1 + x^2)^{10}$$

input `Integrate[x*(-1 + x^2)^9,x]`

output `(-1 + x^2)^10/20`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(x^2 - 1)^9 dx$$

$$\downarrow 241$$

$$\frac{1}{20}(1 - x^2)^{10}$$

input `Int[x*(-1 + x^2)^9,x]`

output `(1 - x^2)^10/20`

Defintions of rubi rules used

rule 241 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

method	result	size
default	$\frac{(x^2-1)^{10}}{20}$	10
gospers	$\frac{x^2(x^{18}-10x^{16}+45x^{14}-120x^{12}+210x^{10}-252x^8+210x^6-120x^4+45x^2-10)}{20}$	51
norman	$-\frac{1}{2}x^2 + \frac{9}{4}x^4 - 6x^6 + \frac{21}{2}x^8 - \frac{63}{5}x^{10} + \frac{21}{2}x^{12} - 6x^{14} + \frac{9}{4}x^{16} - \frac{1}{2}x^{18} + \frac{1}{20}x^{20}$	52
paralelrirsch	$-\frac{1}{2}x^2 + \frac{9}{4}x^4 - 6x^6 + \frac{21}{2}x^8 - \frac{63}{5}x^{10} + \frac{21}{2}x^{12} - 6x^{14} + \frac{9}{4}x^{16} - \frac{1}{2}x^{18} + \frac{1}{20}x^{20}$	52
risch	$\frac{1}{20}x^{20} - \frac{1}{2}x^{18} + \frac{9}{4}x^{16} - 6x^{14} + \frac{21}{2}x^{12} - \frac{63}{5}x^{10} + \frac{21}{2}x^8 - 6x^6 + \frac{9}{4}x^4 - \frac{1}{2}x^2 + \frac{1}{20}$	53
orering	$\frac{x^2(x^{18}-10x^{16}+45x^{14}-120x^{12}+210x^{10}-252x^8+210x^6-120x^4+45x^2-10)(x^2-1)^9}{20(-1+x)^9(1+x)^9}$	68

input `int(x*(x^2-1)^9,x,method=_RETURNVERBOSE)`output `1/20*(x^2-1)^10`**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 51 vs. 2(9) = 18.

Time = 0.08 (sec) , antiderivative size = 51, normalized size of antiderivative = 3.92

$$\int x(-1+x^2)^9 dx = \frac{1}{20}x^{20} - \frac{1}{2}x^{18} + \frac{9}{4}x^{16} - 6x^{14} + \frac{21}{2}x^{12} - \frac{63}{5}x^{10} + \frac{21}{2}x^8 - 6x^6 + \frac{9}{4}x^4 - \frac{1}{2}x^2$$

input `integrate(x*(x^2-1)^9,x, algorithm="fricas")`output `1/20*x^20 - 1/2*x^18 + 9/4*x^16 - 6*x^14 + 21/2*x^12 - 63/5*x^10 + 21/2*x^8 - 6*x^6 + 9/4*x^4 - 1/2*x^2`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 58 vs. $2(7) = 14$.

Time = 0.02 (sec) , antiderivative size = 58, normalized size of antiderivative = 4.46

$$\int x(-1+x^2)^9 dx = \frac{x^{20}}{20} - \frac{x^{18}}{2} + \frac{9x^{16}}{4} - 6x^{14} + \frac{21x^{12}}{2} - \frac{63x^{10}}{5} + \frac{21x^8}{2} - 6x^6 + \frac{9x^4}{4} - \frac{x^2}{2}$$

input `integrate(x*(x**2-1)**9,x)`

output `x**20/20 - x**18/2 + 9*x**16/4 - 6*x**14 + 21*x**12/2 - 63*x**10/5 + 21*x**8/2 - 6*x**6 + 9*x**4/4 - x**2/2`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int x(-1+x^2)^9 dx = \frac{1}{20} (x^2 - 1)^{10}$$

input `integrate(x*(x^2-1)^9,x, algorithm="maxima")`

output `1/20*(x^2 - 1)^10`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int x(-1+x^2)^9 dx = \frac{1}{20} (x^2 - 1)^{10}$$

input `integrate(x*(x^2-1)^9,x, algorithm="giac")`

output `1/20*(x^2 - 1)^10`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int x(-1 + x^2)^9 dx = \frac{(x^2 - 1)^{10}}{20}$$

input `int(x*(x^2 - 1)^9,x)`output `(x^2 - 1)^10/20`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 50, normalized size of antiderivative = 3.85

$$\int x(-1 + x^2)^9 dx$$

$$= \frac{x^2(x^{18} - 10x^{16} + 45x^{14} - 120x^{12} + 210x^{10} - 252x^8 + 210x^6 - 120x^4 + 45x^2 - 10)}{20}$$

input `int(x*(x^2-1)^9,x)`output `(x**2*(x**18 - 10*x**16 + 45*x**14 - 120*x**12 + 210*x**10 - 252*x**8 + 210*x**6 - 120*x**4 + 45*x**2 - 10))/20`

3.49 $\int \frac{3+2x}{(7+6x)^3} dx$

Optimal result	335
Mathematica [A] (verified)	335
Rubi [A] (verified)	336
Maple [A] (verified)	337
Fricas [A] (verification not implemented)	337
Sympy [A] (verification not implemented)	338
Maxima [A] (verification not implemented)	338
Giac [A] (verification not implemented)	338
Mupad [B] (verification not implemented)	339
Reduce [B] (verification not implemented)	339

Optimal result

Integrand size = 13, antiderivative size = 18

$$\int \frac{3+2x}{(7+6x)^3} dx = -\frac{(3+2x)^2}{8(7+6x)^2}$$

output `-1/8*(3+2*x)^2/(7+6*x)^2`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int \frac{3+2x}{(7+6x)^3} dx = -\frac{4+3x}{9(7+6x)^2}$$

input `Integrate[(3 + 2*x)/(7 + 6*x)^3,x]`

output `-1/9*(4 + 3*x)/(7 + 6*x)^2`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{2x + 3}{(6x + 7)^3} dx$$

↓ 48

$$-\frac{(2x + 3)^2}{8(6x + 7)^2}$$

input `Int[(3 + 2*x)/(7 + 6*x)^3,x]`

output `-1/8*(3 + 2*x)^2/(7 + 6*x)^2`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp
 [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
 a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

method	result	size
norman	$\frac{-\frac{x}{3}-\frac{4}{9}}{(7+6x)^2}$	14
gospers	$-\frac{3x+4}{9(7+6x)^2}$	15
risch	$\frac{-\frac{x}{3}-\frac{4}{9}}{(7+6x)^2}$	15
parallelrisch	$\frac{-12x-16}{36(7+6x)^2}$	15
orering	$-\frac{3x+4}{9(7+6x)^2}$	15
default	$-\frac{1}{18(7+6x)^2} - \frac{1}{18(7+6x)}$	20
meijerg	$\frac{3x(\frac{6x}{7}+2)}{686(1+\frac{6x}{7})^2} + \frac{x^2}{343(1+\frac{6x}{7})^2}$	29

input `int((3+2*x)/(7+6*x)^3,x,method=_RETURNVERBOSE)`output `1/(7+6*x)^2*(-1/3*x-4/9)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.06

$$\int \frac{3+2x}{(7+6x)^3} dx = -\frac{3x+4}{9(36x^2+84x+49)}$$

input `integrate((3+2*x)/(7+6*x)^3,x, algorithm="fricas")`output `-1/9*(3*x + 4)/(36*x^2 + 84*x + 49)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int \frac{3 + 2x}{(7 + 6x)^3} dx = \frac{-3x - 4}{324x^2 + 756x + 441}$$

input `integrate((3+2*x)/(7+6*x)**3,x)`output `(-3*x - 4)/(324*x**2 + 756*x + 441)`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.06

$$\int \frac{3 + 2x}{(7 + 6x)^3} dx = -\frac{3x + 4}{9(36x^2 + 84x + 49)}$$

input `integrate((3+2*x)/(7+6*x)^3,x, algorithm="maxima")`output `-1/9*(3*x + 4)/(36*x^2 + 84*x + 49)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{3 + 2x}{(7 + 6x)^3} dx = -\frac{3x + 4}{9(6x + 7)^2}$$

input `integrate((3+2*x)/(7+6*x)^3,x, algorithm="giac")`output `-1/9*(3*x + 4)/(6*x + 7)^2`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{3 + 2x}{(7 + 6x)^3} dx = -\frac{3x + 4}{9(6x + 7)^2}$$

input `int((2*x + 3)/(6*x + 7)^3,x)`

output `-(3*x + 4)/(9*(6*x + 7)^2)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{3 + 2x}{(7 + 6x)^3} dx = \frac{4x^2 - 7}{1008x^2 + 2352x + 1372}$$

input `int((3+2*x)/(7+6*x)^3,x)`

output `(4*x**2 - 7)/(28*(36*x**2 + 84*x + 49))`

3.50 $\int x^4(1+x^5)^5 dx$

Optimal result	340
Mathematica [B] (verified)	340
Rubi [A] (verified)	341
Maple [A] (verified)	342
Fricas [B] (verification not implemented)	342
Sympy [B] (verification not implemented)	343
Maxima [A] (verification not implemented)	343
Giac [A] (verification not implemented)	343
Mupad [B] (verification not implemented)	344
Reduce [B] (verification not implemented)	344

Optimal result

Integrand size = 11, antiderivative size = 11

$$\int x^4(1+x^5)^5 dx = \frac{1}{30}(1+x^5)^6$$

output `1/30*(x^5+1)^6`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 43 vs. $2(11) = 22$.

Time = 0.00 (sec) , antiderivative size = 43, normalized size of antiderivative = 3.91

$$\int x^4(1+x^5)^5 dx = \frac{x^5}{5} + \frac{x^{10}}{2} + \frac{2x^{15}}{3} + \frac{x^{20}}{2} + \frac{x^{25}}{5} + \frac{x^{30}}{30}$$

input `Integrate[x^4*(1 + x^5)^5,x]`

output `x^5/5 + x^10/2 + (2*x^15)/3 + x^20/2 + x^25/5 + x^30/30`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {793}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4(x^5 + 1)^5 dx$$

↓ 793

$$\frac{1}{30}(x^5 + 1)^6$$

input

```
Int[x^4*(1 + x^5)^5,x]
```

output

```
(1 + x^5)^6/30
```

Defintions of rubi rules used

rule 793

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(a + b*x^n)^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] && NeQ[p, -1]
```

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

method	result	size
default	$\frac{(x^5+1)^6}{30}$	10
gospers	$\frac{x^5(x^{25}+6x^{20}+15x^{15}+20x^{10}+15x^5+6)}{30}$	31
norman	$\frac{1}{5}x^{25} + \frac{1}{30}x^{30} + \frac{1}{5}x^5 + \frac{1}{2}x^{10} + \frac{2}{3}x^{15} + \frac{1}{2}x^{20}$	32
parallemrisch	$\frac{1}{5}x^{25} + \frac{1}{30}x^{30} + \frac{1}{5}x^5 + \frac{1}{2}x^{10} + \frac{2}{3}x^{15} + \frac{1}{2}x^{20}$	32
risch	$\frac{1}{30}x^{30} + \frac{1}{5}x^{25} + \frac{1}{2}x^{20} + \frac{2}{3}x^{15} + \frac{1}{2}x^{10} + \frac{1}{5}x^5 + \frac{1}{30}$	33
orering	$\frac{x^5(x^{25}+6x^{20}+15x^{15}+20x^{10}+15x^5+6)(x^5+1)^5}{30(1+x)^5(x^4-x^3+x^2-x+1)^5}$	61

input `int(x^4*(x^5+1)^5,x,method=_RETURNVERBOSE)`

output `1/30*(x^5+1)^6`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 31 vs. 2(9) = 18.

Time = 0.09 (sec) , antiderivative size = 31, normalized size of antiderivative = 2.82

$$\int x^4(1+x^5)^5 dx = \frac{1}{30}x^{30} + \frac{1}{5}x^{25} + \frac{1}{2}x^{20} + \frac{2}{3}x^{15} + \frac{1}{2}x^{10} + \frac{1}{5}x^5$$

input `integrate(x^4*(x^5+1)^5,x, algorithm="fricas")`

output `1/30*x^30 + 1/5*x^25 + 1/2*x^20 + 2/3*x^15 + 1/2*x^10 + 1/5*x^5`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 31 vs. $2(7) = 14$.

Time = 0.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 2.82

$$\int x^4(1+x^5)^5 dx = \frac{x^{30}}{30} + \frac{x^{25}}{5} + \frac{x^{20}}{2} + \frac{2x^{15}}{3} + \frac{x^{10}}{2} + \frac{x^5}{5}$$

input `integrate(x**4*(x**5+1)**5,x)`

output `x**30/30 + x**25/5 + x**20/2 + 2*x**15/3 + x**10/2 + x**5/5`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int x^4(1+x^5)^5 dx = \frac{1}{30} (x^5 + 1)^6$$

input `integrate(x^4*(x^5+1)^5,x, algorithm="maxima")`

output `1/30*(x^5 + 1)^6`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int x^4(1+x^5)^5 dx = \frac{1}{30} (x^5 + 1)^6$$

input `integrate(x^4*(x^5+1)^5,x, algorithm="giac")`

output `1/30*(x^5 + 1)^6`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 2.82

$$\int x^4(1+x^5)^5 dx = \frac{x^{30}}{30} + \frac{x^{25}}{5} + \frac{x^{20}}{2} + \frac{2x^{15}}{3} + \frac{x^{10}}{2} + \frac{x^5}{5}$$

input `int(x^4*(x^5 + 1)^5,x)`output `x^5/5 + x^10/2 + (2*x^15)/3 + x^20/2 + x^25/5 + x^30/30`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 30, normalized size of antiderivative = 2.73

$$\int x^4(1+x^5)^5 dx = \frac{x^5(x^{25} + 6x^{20} + 15x^{15} + 20x^{10} + 15x^5 + 6)}{30}$$

input `int(x^4*(x^5+1)^5,x)`output `(x**5*(x**25 + 6*x**20 + 15*x**15 + 20*x**10 + 15*x**5 + 6))/30`

3.51 $\int (1 - x)^{20} x^4 dx$

Optimal result	345
Mathematica [B] (verified)	345
Rubi [A] (verified)	346
Maple [B] (verified)	347
Fricas [B] (verification not implemented)	348
Sympy [B] (verification not implemented)	348
Maxima [B] (verification not implemented)	349
Giac [B] (verification not implemented)	349
Mupad [B] (verification not implemented)	350
Reduce [B] (verification not implemented)	351

Optimal result

Integrand size = 11, antiderivative size = 56

$$\int (1 - x)^{20} x^4 dx = -\frac{1}{21}(1 - x)^{21} + \frac{2}{11}(1 - x)^{22} - \frac{6}{23}(1 - x)^{23} + \frac{1}{6}(1 - x)^{24} - \frac{1}{25}(1 - x)^{25}$$

output

```
-1/21*(1-x)^21+2/11*(1-x)^22-6/23*(1-x)^23+1/6*(1-x)^24-1/25*(1-x)^25
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 140 vs. 2(56) = 112.

Time = 0.00 (sec) , antiderivative size = 140, normalized size of antiderivative = 2.50

$$\begin{aligned} \int (1 - x)^{20} x^4 dx = & \frac{x^5}{5} - \frac{10x^6}{3} + \frac{190x^7}{7} - \frac{285x^8}{2} + \frac{1615x^9}{3} - \frac{7752x^{10}}{5} + \frac{38760x^{11}}{11} - 6460x^{12} \\ & + 9690x^{13} - \frac{83980x^{14}}{7} + \frac{184756x^{15}}{15} - \frac{20995x^{16}}{2} + 7410x^{17} - \frac{12920x^{18}}{3} \\ & + 2040x^{19} - \frac{3876x^{20}}{5} + \frac{1615x^{21}}{7} - \frac{570x^{22}}{11} + \frac{190x^{23}}{23} - \frac{5x^{24}}{6} + \frac{x^{25}}{25} \end{aligned}$$

input

```
Integrate[(1 - x)^20*x^4,x]
```

output

$$x^5/5 - (10x^6)/3 + (190x^7)/7 - (285x^8)/2 + (1615x^9)/3 - (7752x^{10})/5 + (38760x^{11})/11 - 6460x^{12} + 9690x^{13} - (83980x^{14})/7 + (184756x^{15})/15 - (20995x^{16})/2 + 7410x^{17} - (12920x^{18})/3 + 2040x^{19} - (3876x^{20})/5 + (1615x^{21})/7 - (570x^{22})/11 + (190x^{23})/23 - (5x^{24})/6 + x^{25}/25$$

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (1-x)^{20} x^4 dx$$

$$\downarrow 49$$

$$\int ((1-x)^{24} - 4(1-x)^{23} + 6(1-x)^{22} - 4(1-x)^{21} + (1-x)^{20}) dx$$

$$\downarrow 2009$$

$$-\frac{1}{25}(1-x)^{25} + \frac{1}{6}(1-x)^{24} - \frac{6}{23}(1-x)^{23} + \frac{2}{11}(1-x)^{22} - \frac{1}{21}(1-x)^{21}$$

input

Int[(1 - x)^20*x^4,x]

output

$$-1/21*(1-x)^{21} + (2*(1-x)^{22})/11 - (6*(1-x)^{23})/23 + (1-x)^{24}/6 - (1-x)^{25}/25$$

Definitions of rubi rules used

rule 49

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 105 vs. $2(46) = 92$.

Time = 0.07 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.89

method	result
gospers	$x^5(10626x^{20} - 221375x^{19} + 2194500x^{18} - 13765500x^{17} + 61289250x^{16} - 205931880x^{15} + 541926000x^{14} - 1144066000x^{13} + 1968466500x^{12} - 2788660875x^{11} + 3272028760x^{10} - 3187041000x^9 + 2574148500x^8 - 1716099000x^7 + 936054000x^6 - 411863760x^5 + 143008250x^4 - 37855125x^3 + 7210500x^2 - 885500x + 53130)$
default	$9690x^{13} - \frac{83980}{7}x^{14} + \frac{184756}{15}x^{15} - \frac{20995}{2}x^{16} + 7410x^{17} - \frac{12920}{3}x^{18} + 2040x^{19} - \frac{3876}{5}x^{20} + \frac{1615}{7}x^{21}$
norman	$9690x^{13} - \frac{83980}{7}x^{14} + \frac{184756}{15}x^{15} - \frac{20995}{2}x^{16} + 7410x^{17} - \frac{12920}{3}x^{18} + 2040x^{19} - \frac{3876}{5}x^{20} + \frac{1615}{7}x^{21}$
risch	$9690x^{13} - \frac{83980}{7}x^{14} + \frac{184756}{15}x^{15} - \frac{20995}{2}x^{16} + 7410x^{17} - \frac{12920}{3}x^{18} + 2040x^{19} - \frac{3876}{5}x^{20} + \frac{1615}{7}x^{21}$
parallelrisch	$9690x^{13} - \frac{83980}{7}x^{14} + \frac{184756}{15}x^{15} - \frac{20995}{2}x^{16} + 7410x^{17} - \frac{12920}{3}x^{18} + 2040x^{19} - \frac{3876}{5}x^{20} + \frac{1615}{7}x^{21}$
orering	$x^5(10626x^{20} - 221375x^{19} + 2194500x^{18} - 13765500x^{17} + 61289250x^{16} - 205931880x^{15} + 541926000x^{14} - 1144066000x^{13} + 1968466500x^{12} - 2788660875x^{11} + 3272028760x^{10} - 3187041000x^9 + 2574148500x^8 - 1716099000x^7 + 936054000x^6 - 411863760x^5 + 143008250x^4 - 37855125x^3 + 7210500x^2 - 885500x + 53130)$

input

```
int((1-x)^20*x^4,x,method=_RETURNVERBOSE)
```

output

```
1/265650*x^5*(10626*x^20-221375*x^19+2194500*x^18-13765500*x^17+61289250*x
^16-205931880*x^15+541926000*x^14-1144066000*x^13+1968466500*x^12-27886608
75*x^11+3272028760*x^10-3187041000*x^9+2574148500*x^8-1716099000*x^7+93605
4000*x^6-411863760*x^5+143008250*x^4-37855125*x^3+7210500*x^2-885500*x+531
30)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 106 vs. $2(36) = 72$.

Time = 0.08 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.89

$$\int (1-x)^{20} x^4 dx = \frac{1}{25} x^{25} - \frac{5}{6} x^{24} + \frac{190}{23} x^{23} - \frac{570}{11} x^{22} + \frac{1615}{7} x^{21} - \frac{3876}{5} x^{20}$$

$$+ 2040 x^{19} - \frac{12920}{3} x^{18} + 7410 x^{17} - \frac{20995}{2} x^{16}$$

$$+ \frac{184756}{15} x^{15} - \frac{83980}{7} x^{14} + 9690 x^{13} - 6460 x^{12} + \frac{38760}{11} x^{11}$$

$$- \frac{7752}{5} x^{10} + \frac{1615}{3} x^9 - \frac{285}{2} x^8 + \frac{190}{7} x^7 - \frac{10}{3} x^6 + \frac{1}{5} x^5$$

input `integrate((1-x)^20*x^4,x, algorithm="fricas")`

output `1/25*x^25 - 5/6*x^24 + 190/23*x^23 - 570/11*x^22 + 1615/7*x^21 - 3876/5*x^20 + 2040*x^19 - 12920/3*x^18 + 7410*x^17 - 20995/2*x^16 + 184756/15*x^15 - 83980/7*x^14 + 9690*x^13 - 6460*x^12 + 38760/11*x^11 - 7752/5*x^10 + 1615/3*x^9 - 285/2*x^8 + 190/7*x^7 - 10/3*x^6 + 1/5*x^5`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 131 vs. $2(36) = 72$.

Time = 0.03 (sec) , antiderivative size = 131, normalized size of antiderivative = 2.34

$$\int (1-x)^{20} x^4 dx = \frac{x^{25}}{25} - \frac{5x^{24}}{6} + \frac{190x^{23}}{23} - \frac{570x^{22}}{11} + \frac{1615x^{21}}{7} - \frac{3876x^{20}}{5}$$

$$+ 2040x^{19} - \frac{12920x^{18}}{3} + 7410x^{17} - \frac{20995x^{16}}{2}$$

$$+ \frac{184756x^{15}}{15} - \frac{83980x^{14}}{7} + 9690x^{13} - 6460x^{12} + \frac{38760x^{11}}{11}$$

$$- \frac{7752x^{10}}{5} + \frac{1615x^9}{3} - \frac{285x^8}{2} + \frac{190x^7}{7} - \frac{10x^6}{3} + \frac{x^5}{5}$$

input `integrate((1-x)**20*x**4,x)`

output

```
x**25/25 - 5*x**24/6 + 190*x**23/23 - 570*x**22/11 + 1615*x**21/7 - 3876*x
**20/5 + 2040*x**19 - 12920*x**18/3 + 7410*x**17 - 20995*x**16/2 + 184756*
x**15/15 - 83980*x**14/7 + 9690*x**13 - 6460*x**12 + 38760*x**11/11 - 7752
*x**10/5 + 1615*x**9/3 - 285*x**8/2 + 190*x**7/7 - 10*x**6/3 + x**5/5
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 106 vs. $2(36) = 72$.

Time = 0.03 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.89

$$\int (1-x)^{20} x^4 dx = \frac{1}{25} x^{25} - \frac{5}{6} x^{24} + \frac{190}{23} x^{23} - \frac{570}{11} x^{22} + \frac{1615}{7} x^{21} - \frac{3876}{5} x^{20}$$

$$+ 2040 x^{19} - \frac{12920}{3} x^{18} + 7410 x^{17} - \frac{20995}{2} x^{16}$$

$$+ \frac{184756}{15} x^{15} - \frac{83980}{7} x^{14} + 9690 x^{13} - 6460 x^{12} + \frac{38760}{11} x^{11}$$

$$- \frac{7752}{5} x^{10} + \frac{1615}{3} x^9 - \frac{285}{2} x^8 + \frac{190}{7} x^7 - \frac{10}{3} x^6 + \frac{1}{5} x^5$$

input

```
integrate((1-x)^20*x^4,x, algorithm="maxima")
```

output

```
1/25*x^25 - 5/6*x^24 + 190/23*x^23 - 570/11*x^22 + 1615/7*x^21 - 3876/5*x^
20 + 2040*x^19 - 12920/3*x^18 + 7410*x^17 - 20995/2*x^16 + 184756/15*x^15
- 83980/7*x^14 + 9690*x^13 - 6460*x^12 + 38760/11*x^11 - 7752/5*x^10 + 161
5/3*x^9 - 285/2*x^8 + 190/7*x^7 - 10/3*x^6 + 1/5*x^5
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 106 vs. $2(36) = 72$.

Time = 0.13 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.89

$$\int (1-x)^{20} x^4 dx = \frac{1}{25} x^{25} - \frac{5}{6} x^{24} + \frac{190}{23} x^{23} - \frac{570}{11} x^{22} + \frac{1615}{7} x^{21} - \frac{3876}{5} x^{20}$$

$$+ 2040 x^{19} - \frac{12920}{3} x^{18} + 7410 x^{17} - \frac{20995}{2} x^{16}$$

$$+ \frac{184756}{15} x^{15} - \frac{83980}{7} x^{14} + 9690 x^{13} - 6460 x^{12} + \frac{38760}{11} x^{11}$$

$$- \frac{7752}{5} x^{10} + \frac{1615}{3} x^9 - \frac{285}{2} x^8 + \frac{190}{7} x^7 - \frac{10}{3} x^6 + \frac{1}{5} x^5$$

input `integrate((1-x)^20*x^4,x, algorithm="giac")`

output $1/25*x^{25} - 5/6*x^{24} + 190/23*x^{23} - 570/11*x^{22} + 1615/7*x^{21} - 3876/5*x^{20} + 2040*x^{19} - 12920/3*x^{18} + 7410*x^{17} - 20995/2*x^{16} + 184756/15*x^{15} - 83980/7*x^{14} + 9690*x^{13} - 6460*x^{12} + 38760/11*x^{11} - 7752/5*x^{10} + 1615/3*x^9 - 285/2*x^8 + 190/7*x^7 - 10/3*x^6 + 1/5*x^5$

Mupad [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.89

$$\int (1-x)^{20} x^4 dx = \frac{x^{25}}{25} - \frac{5x^{24}}{6} + \frac{190x^{23}}{23} - \frac{570x^{22}}{11} + \frac{1615x^{21}}{7} - \frac{3876x^{20}}{5} + 2040x^{19} - \frac{12920x^{18}}{3} + 7410x^{17} - \frac{20995x^{16}}{2} + \frac{184756x^{15}}{15} - \frac{83980x^{14}}{7} + 9690x^{13} - 6460x^{12} + \frac{38760x^{11}}{11} - \frac{7752x^{10}}{5} + \frac{1615x^9}{3} - \frac{285x^8}{2} + \frac{190x^7}{7} - \frac{10x^6}{3} + \frac{x^5}{5}$$

input `int(x^4*(x - 1)^20,x)`

output $x^5/5 - (10*x^6)/3 + (190*x^7)/7 - (285*x^8)/2 + (1615*x^9)/3 - (7752*x^{10})/5 + (38760*x^{11})/11 - 6460*x^{12} + 9690*x^{13} - (83980*x^{14})/7 + (184756*x^{15})/15 - (20995*x^{16})/2 + 7410*x^{17} - (12920*x^{18})/3 + 2040*x^{19} - (3876*x^{20})/5 + (1615*x^{21})/7 - (570*x^{22})/11 + (190*x^{23})/23 - (5*x^{24})/6 + x^25/25$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.88

$$\int (1-x)^{20} x^4 dx$$
$$= \frac{x^5(10626x^{20} - 221375x^{19} + 2194500x^{18} - 13765500x^{17} + 61289250x^{16} - 205931880x^{15} + 541926000x^{14} - 1144066000x^{13} + 196846500x^{12} - 2788660875x^{11} + 3272028760x^{10} - 3187041000x^9 + 2574148500x^8 - 1716099000x^7 + 936054000x^6 - 411863760x^5 + 143008250x^4 - 37855125x^3 + 7210500x^2 - 885500x + 53130)}{265650}$$

input `int((1-x)^20*x^4,x)`output `(x**5*(10626*x**20 - 221375*x**19 + 2194500*x**18 - 13765500*x**17 + 61289250*x**16 - 205931880*x**15 + 541926000*x**14 - 1144066000*x**13 + 196846500*x**12 - 2788660875*x**11 + 3272028760*x**10 - 3187041000*x**9 + 2574148500*x**8 - 1716099000*x**7 + 936054000*x**6 - 411863760*x**5 + 143008250*x**4 - 37855125*x**3 + 7210500*x**2 - 885500*x + 53130))/265650`

$$3.52 \quad \int \frac{\sin\left(\frac{1}{x}\right)}{x^2} dx$$

Optimal result	352
Mathematica [A] (verified)	352
Rubi [A] (verified)	353
Maple [A] (verified)	354
Fricas [A] (verification not implemented)	354
Sympy [A] (verification not implemented)	355
Maxima [A] (verification not implemented)	355
Giac [A] (verification not implemented)	355
Mupad [B] (verification not implemented)	356
Reduce [B] (verification not implemented)	356

Optimal result

Integrand size = 8, antiderivative size = 4

$$\int \frac{\sin\left(\frac{1}{x}\right)}{x^2} dx = \cos\left(\frac{1}{x}\right)$$

output `cos(1/x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{\sin\left(\frac{1}{x}\right)}{x^2} dx = \cos\left(\frac{1}{x}\right)$$

input `Integrate[Sin[x^(-1)]/x^2,x]`

output `Cos[x^(-1)]`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3860, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sin\left(\frac{1}{x}\right)}{x^2} dx \\ & \quad \downarrow \text{3860} \\ & - \int \sin\left(\frac{1}{x}\right) d\frac{1}{x} \\ & \quad \downarrow \text{3042} \\ & - \int \sin\left(\frac{1}{x}\right) d\frac{1}{x} \\ & \quad \downarrow \text{3118} \\ & \cos\left(\frac{1}{x}\right) \end{aligned}$$

input `Int[Sin[x^(-1)]/x^2,x]`

output `Cos[x^(-1)]`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3860

```
Int[(x_)^(m_.)*((a_.) + (b_.)*Sin[(c_.) + (d_.)*(x_)^(n_)])^(p_.), x_Symbol
] -> Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Sin[c + d*x])^
p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[
(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[
(m + 1)/n], 0]))
```

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

method	result	size
derivativdivides	$\cos\left(\frac{1}{x}\right)$	5
default	$\cos\left(\frac{1}{x}\right)$	5
risch	$\cos\left(\frac{1}{x}\right)$	5
parallelrisc	$1 + \cos\left(\frac{1}{x}\right)$	7
norman	$\frac{2}{1 + \tan\left(\frac{1}{2x}\right)^2}$	15
meijerg	$-\sqrt{\pi} \left(\frac{1}{\sqrt{\pi}} - \frac{\cos\left(\frac{1}{x}\right)}{\sqrt{\pi}} \right)$	19
orering	$-2 \sin\left(\frac{1}{x}\right) x - x^4 \left(-\frac{\cos\left(\frac{1}{x}\right)}{x^4} - \frac{2 \sin\left(\frac{1}{x}\right)}{x^3} \right)$	33

input `int(sin(1/x)/x^2,x,method=_RETURNVERBOSE)`output `cos(1/x)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{\sin\left(\frac{1}{x}\right)}{x^2} dx = \cos\left(\frac{1}{x}\right)$$

input `integrate(sin(1/x)/x^2,x, algorithm="fricas")`

output `cos(1/x)`

Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int \frac{\sin\left(\frac{1}{x}\right)}{x^2} dx = \cos\left(\frac{1}{x}\right)$$

input `integrate(sin(1/x)/x**2,x)`

output `cos(1/x)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{\sin\left(\frac{1}{x}\right)}{x^2} dx = \cos\left(\frac{1}{x}\right)$$

input `integrate(sin(1/x)/x^2,x, algorithm="maxima")`

output `cos(1/x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{\sin\left(\frac{1}{x}\right)}{x^2} dx = \cos\left(\frac{1}{x}\right)$$

input `integrate(sin(1/x)/x^2,x, algorithm="giac")`

output `cos(1/x)`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{\sin\left(\frac{1}{x}\right)}{x^2} dx = \cos\left(\frac{1}{x}\right)$$

input `int(sin(1/x)/x^2,x)`

output `cos(1/x)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{\sin\left(\frac{1}{x}\right)}{x^2} dx = \cos\left(\frac{1}{x}\right)$$

input `int(sin(1/x)/x^2,x)`

output `cos(1/x)`

3.53 $\int \sin(\sqrt[4]{-1+x}) dx$

Optimal result	357
Mathematica [A] (verified)	357
Rubi [A] (verified)	358
Maple [A] (verified)	360
Fricas [A] (verification not implemented)	360
Sympy [A] (verification not implemented)	361
Maxima [A] (verification not implemented)	361
Giac [A] (verification not implemented)	361
Mupad [B] (verification not implemented)	362
Reduce [B] (verification not implemented)	362

Optimal result

Integrand size = 8, antiderivative size = 62

$$\int \sin(\sqrt[4]{-1+x}) dx = 24\sqrt[4]{-1+x} \cos(\sqrt[4]{-1+x}) - 4(-1+x)^{3/4} \cos(\sqrt[4]{-1+x}) - 24 \sin(\sqrt[4]{-1+x}) + 12\sqrt{-1+x} \sin(\sqrt[4]{-1+x})$$

output

```
24*(-1+x)^(1/4)*cos((-1+x)^(1/4))-4*(-1+x)^(3/4)*cos((-1+x)^(1/4))-24*sin((-1+x)^(1/4))+12*sin((-1+x)^(1/4))*(-1+x)^(1/2)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.74

$$\int \sin(\sqrt[4]{-1+x}) dx = -4(-6 + \sqrt{-1+x}) \sqrt[4]{-1+x} \cos(\sqrt[4]{-1+x}) + 12(-2 + \sqrt{-1+x}) \sin(\sqrt[4]{-1+x})$$

input

```
Integrate[Sin[(-1 + x)^(1/4)], x]
```

output

```
-4*(-6 + Sqrt[-1 + x])*(-1 + x)^(1/4)*Cos[(-1 + x)^(1/4)] + 12*(-2 + Sqrt[-1 + x])*Sin[(-1 + x)^(1/4)]
```

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.08, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$, Rules used = {3842, 3042, 3777, 3042, 3777, 25, 3042, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin(\sqrt[4]{x-1}) \, dx \\
 & \quad \downarrow \text{3842} \\
 & 4 \int (x-1)^{3/4} \sin(\sqrt[4]{x-1}) \, d\sqrt[4]{x-1} \\
 & \quad \downarrow \text{3042} \\
 & 4 \int (x-1)^{3/4} \sin(\sqrt[4]{x-1}) \, d\sqrt[4]{x-1} \\
 & \quad \downarrow \text{3777} \\
 & 4 \left(3 \int \sqrt{x-1} \cos(\sqrt[4]{x-1}) \, d\sqrt[4]{x-1} - (x-1)^{3/4} \cos(\sqrt[4]{x-1}) \right) \\
 & \quad \downarrow \text{3042} \\
 & 4 \left(3 \int \sqrt{x-1} \sin\left(\sqrt[4]{x-1} + \frac{\pi}{2}\right) \, d\sqrt[4]{x-1} - (x-1)^{3/4} \cos(\sqrt[4]{x-1}) \right) \\
 & \quad \downarrow \text{3777} \\
 & 4 \left(3 \left(2 \int -\sqrt[4]{x-1} \sin(\sqrt[4]{x-1}) \, d\sqrt[4]{x-1} + \sqrt{x-1} \sin(\sqrt[4]{x-1}) \right) - (x-1)^{3/4} \cos(\sqrt[4]{x-1}) \right) \\
 & \quad \downarrow \text{25} \\
 & 4 \left(3 \left(\sqrt{x-1} \sin(\sqrt[4]{x-1}) - 2 \int \sqrt[4]{x-1} \sin(\sqrt[4]{x-1}) \, d\sqrt[4]{x-1} \right) - (x-1)^{3/4} \cos(\sqrt[4]{x-1}) \right) \\
 & \quad \downarrow \text{3042} \\
 & 4 \left(3 \left(\sqrt{x-1} \sin(\sqrt[4]{x-1}) - 2 \int \sqrt[4]{x-1} \sin(\sqrt[4]{x-1}) \, d\sqrt[4]{x-1} \right) - (x-1)^{3/4} \cos(\sqrt[4]{x-1}) \right) \\
 & \quad \downarrow \text{3777}
 \end{aligned}$$

$$4 \left(3 \left(\sqrt{x-1} \sin(\sqrt[4]{x-1}) - 2 \left(\int \cos(\sqrt[4]{x-1}) d\sqrt[4]{x-1} - \sqrt[4]{x-1} \cos(\sqrt[4]{x-1}) \right) \right) \right) - (x-1)^{3/4} \cos(\sqrt[4]{x-1})$$

↓ 3042

$$4 \left(3 \left(\sqrt{x-1} \sin(\sqrt[4]{x-1}) - 2 \left(\int \sin\left(\sqrt[4]{x-1} + \frac{\pi}{2}\right) d\sqrt[4]{x-1} - \sqrt[4]{x-1} \cos(\sqrt[4]{x-1}) \right) \right) \right) - (x-1)^{3/4} \cos(\sqrt[4]{x-1})$$

↓ 3117

$$4 \left(3 \left(\sqrt{x-1} \sin(\sqrt[4]{x-1}) - 2 \left(\sin(\sqrt[4]{x-1}) - \sqrt[4]{x-1} \cos(\sqrt[4]{x-1}) \right) \right) \right) - (x-1)^{3/4} \cos(\sqrt[4]{x-1})$$

input `Int[Sin[(-1 + x)^(1/4)], x]`

output `4*(-((-1 + x)^(3/4)*Cos[(-1 + x)^(1/4)]) + 3*(Sqrt[-1 + x]*Sin[(-1 + x)^(1/4)] - 2*(-((-1 + x)^(1/4)*Cos[(-1 + x)^(1/4)]) + Sin[(-1 + x)^(1/4)]))`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 3842

```
Int[((a_.) + (b_.)*Sin[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)]^(p_.), x_Symbol]
:> Simp[1/(n*f) Subst[Int[x^(1/n - 1)*(a + b*Sin[c + d*x])^p, x], x, (e + f*x)^n], x]
/; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[1/n]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.79

method	result
derivativedivides	$24(-1+x)^{\frac{1}{4}} \cos\left((-1+x)^{\frac{1}{4}}\right) - 4(-1+x)^{\frac{3}{4}} \cos\left((-1+x)^{\frac{1}{4}}\right) - 24 \sin\left((-1+x)^{\frac{1}{4}}\right)$
default	$24(-1+x)^{\frac{1}{4}} \cos\left((-1+x)^{\frac{1}{4}}\right) - 4(-1+x)^{\frac{3}{4}} \cos\left((-1+x)^{\frac{1}{4}}\right) - 24 \sin\left((-1+x)^{\frac{1}{4}}\right)$

input

```
int(sin((-1+x)^(1/4)),x,method=_RETURNVERBOSE)
```

output

```
24*(-1+x)^(1/4)*cos((-1+x)^(1/4))-4*(-1+x)^(3/4)*cos((-1+x)^(1/4))-24*sin((-1+x)^(1/4))+12*sin((-1+x)^(1/4))*(-1+x)^(1/2)
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.60

$$\int \sin(\sqrt[4]{-1+x}) dx = -4 \left((x-1)^{\frac{3}{4}} - 6(x-1)^{\frac{1}{4}} \right) \cos\left((x-1)^{\frac{1}{4}}\right) + 12(\sqrt{x-1} - 2) \sin\left((x-1)^{\frac{1}{4}}\right)$$

input

```
integrate(sin((-1+x)^(1/4)),x, algorithm="fricas")
```

output

```
-4*((x - 1)^(3/4) - 6*(x - 1)^(1/4))*cos((x - 1)^(1/4)) + 12*(sqrt(x - 1) - 2)*sin((x - 1)^(1/4))
```

Sympy [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.97

$$\int \sin(\sqrt[4]{-1+x}) dx = -4(x-1)^{\frac{3}{4}} \cos(\sqrt[4]{x-1}) + 24\sqrt[4]{x-1} \cos(\sqrt[4]{x-1}) \\ + 12\sqrt{x-1} \sin(\sqrt[4]{x-1}) - 24 \sin(\sqrt[4]{x-1})$$

input `integrate(sin((-1+x)**(1/4)),x)`output `-4*(x - 1)**(3/4)*cos((x - 1)**(1/4)) + 24*(x - 1)**(1/4)*cos((x - 1)**(1/4)) + 12*sqrt(x - 1)*sin((x - 1)**(1/4)) - 24*sin((x - 1)**(1/4))`**Maxima [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.60

$$\int \sin(\sqrt[4]{-1+x}) dx = -4 \left((x-1)^{\frac{3}{4}} - 6(x-1)^{\frac{1}{4}} \right) \cos \left((x-1)^{\frac{1}{4}} \right) \\ + 12(\sqrt{x-1} - 2) \sin \left((x-1)^{\frac{1}{4}} \right)$$

input `integrate(sin((-1+x)^(1/4)),x, algorithm="maxima")`output `-4*((x - 1)^(3/4) - 6*(x - 1)^(1/4))*cos((x - 1)^(1/4)) + 12*(sqrt(x - 1) - 2)*sin((x - 1)^(1/4))`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.60

$$\int \sin(\sqrt[4]{-1+x}) dx = -4 \left((x-1)^{\frac{3}{4}} - 6(x-1)^{\frac{1}{4}} \right) \cos \left((x-1)^{\frac{1}{4}} \right) \\ + 12(\sqrt{x-1} - 2) \sin \left((x-1)^{\frac{1}{4}} \right)$$

input `integrate(sin((-1+x)^(1/4)),x, algorithm="giac")`

output `-4*((x - 1)^(3/4) - 6*(x - 1)^(1/4))*cos((x - 1)^(1/4)) + 12*(sqrt(x - 1) - 2)*sin((x - 1)^(1/4))`

Mupad [B] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.66

$$\int \sin(\sqrt[4]{-1+x}) dx = 4 \cos\left((x-1)^{1/4}\right) \left(6(x-1)^{1/4} - (x-1)^{3/4}\right) + 4 \sin\left((x-1)^{1/4}\right) (3\sqrt{x-1} - 6)$$

input `int(sin((x - 1)^(1/4)),x)`

output `4*cos((x - 1)^(1/4))*(6*(x - 1)^(1/4) - (x - 1)^(3/4)) + 4*sin((x - 1)^(1/4))*(3*(x - 1)^(1/2) - 6)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.76

$$\int \sin(\sqrt[4]{-1+x}) dx = -4(x-1)^{3/4} \cos\left((x-1)^{1/4}\right) + 24(x-1)^{1/4} \cos\left((x-1)^{1/4}\right) + 12\sqrt{x-1} \sin\left((x-1)^{1/4}\right) - 24 \sin\left((x-1)^{1/4}\right)$$

input `int(sin((-1+x)^(1/4)),x)`

output `4*(-(x - 1)**(3/4)*cos((x - 1)**(1/4)) + 6*(x - 1)**(1/4)*cos((x - 1)**(1/4)) + 3*sqrt(x - 1)*sin((x - 1)**(1/4)) - 6*sin((x - 1)**(1/4)))`

3.54 $\int x \cos(x^2) \sin(x^2) dx$

Optimal result	363
Mathematica [A] (verified)	363
Rubi [A] (verified)	364
Maple [A] (verified)	365
Fricas [A] (verification not implemented)	365
Sympy [A] (verification not implemented)	366
Maxima [A] (verification not implemented)	366
Giac [A] (verification not implemented)	366
Mupad [B] (verification not implemented)	367
Reduce [B] (verification not implemented)	367

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int x \cos(x^2) \sin(x^2) dx = \frac{1}{4} \sin^2(x^2)$$

output `1/4*sin(x^2)^2`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int x \cos(x^2) \sin(x^2) dx = -\frac{1}{4} \cos^2(x^2)$$

input `Integrate[x*Cos[x^2]*Sin[x^2],x]`

output `-1/4*Cos[x^2]^2`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {3922}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \sin(x^2) \cos(x^2) dx$$

$$\downarrow 3922$$

$$\frac{1}{4} \sin^2(x^2)$$

input `Int[x*Cos[x^2]*Sin[x^2],x]`

output `Sin[x^2]^2/4`

Defintions of rubi rules used

rule 3922 `Int[Cos[(a_.) + (b_.)*(x_)^(n_.)]*(x_)^(m_.)*Sin[(a_.) + (b_.)*(x_)^(n_.)]^(p_.), x_Symbol] :> Simp[Sin[a + b*x^n]^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] && NeQ[p, -1]`

Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

method	result	size
derivativedivides	$-\frac{\cos(x^2)^2}{4}$	9
default	$-\frac{\cos(x^2)^2}{4}$	9
risch	$-\frac{\cos(2x^2)}{8}$	9
parallelrisch	$-\frac{\cos(2x^2)}{8} + \frac{1}{8}$	11
meijerg	$\frac{\sqrt{\pi} \left(\frac{1}{\sqrt{\pi}} - \frac{\cos(2x^2)}{\sqrt{\pi}} \right)}{8}$	21
norman	$\frac{\tan\left(\frac{x^2}{2}\right)^2}{\left(1 + \tan\left(\frac{x^2}{2}\right)^2\right)^2}$	22
orering	$\frac{\cos(x^2) \sin(x^2)}{16x^2} - \frac{\cos(x^2) \sin(x^2) - 2x^2 \sin(x^2)^2 + 2x^2 \cos(x^2)^2}{16x^2}$	52

input `int(x*cos(x^2)*sin(x^2),x,method=_RETURNVERBOSE)`output `-1/4*cos(x^2)^2`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int x \cos(x^2) \sin(x^2) dx = -\frac{1}{4} \cos(x^2)^2$$

input `integrate(x*cos(x^2)*sin(x^2),x, algorithm="fricas")`output `-1/4*cos(x^2)^2`

Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int x \cos(x^2) \sin(x^2) dx = \frac{\sin^2(x^2)}{4}$$

input `integrate(x*cos(x**2)*sin(x**2),x)`

output `sin(x**2)**2/4`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int x \cos(x^2) \sin(x^2) dx = -\frac{1}{4} \cos(x^2)^2$$

input `integrate(x*cos(x^2)*sin(x^2),x, algorithm="maxima")`

output `-1/4*cos(x^2)^2`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int x \cos(x^2) \sin(x^2) dx = -\frac{1}{4} \cos(x^2)^2$$

input `integrate(x*cos(x^2)*sin(x^2),x, algorithm="giac")`

output `-1/4*cos(x^2)^2`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int x \cos(x^2) \sin(x^2) dx = \frac{\sin(x^2)^2}{4}$$

input `int(x*cos(x^2)*sin(x^2),x)`

output `sin(x^2)^2/4`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int x \cos(x^2) \sin(x^2) dx = -\frac{\cos(x^2)^2}{4}$$

input `int(x*cos(x^2)*sin(x^2),x)`

output `(- cos(x**2)**2)/4`

3.55 $\int \sqrt{1 + 3 \cos^2(x)} \sin(2x) dx$

Optimal result	368
Mathematica [A] (verified)	368
Rubi [A] (verified)	369
Maple [A] (verified)	370
Fricas [A] (verification not implemented)	371
Sympy [A] (verification not implemented)	371
Maxima [A] (verification not implemented)	371
Giac [A] (verification not implemented)	372
Mupad [B] (verification not implemented)	372
Reduce [F]	372

Optimal result

Integrand size = 17, antiderivative size = 16

$$\int \sqrt{1 + 3 \cos^2(x)} \sin(2x) dx = -\frac{2}{9}(4 - 3 \sin^2(x))^{3/2}$$

output `-2/9*(4-3*sin(x)^2)^(3/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \sqrt{1 + 3 \cos^2(x)} \sin(2x) dx = -\frac{2}{9}(4 - 3 \sin^2(x))^{3/2}$$

input `Integrate[Sqrt[1 + 3*Cos[x]^2]*Sin[2*x], x]`

output `(-2*(4 - 3*Sin[x]^2)^(3/2))/9`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {3042, 4878, 27, 241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin(2x) \sqrt{3 \cos^2(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(2x) \sqrt{3 \cos(x)^2 + 1} dx \\
 & \quad \downarrow \text{4878} \\
 & \int 2 \sin(x) \sqrt{4 - 3 \sin^2(x)} d \sin(x) \\
 & \quad \downarrow \text{27} \\
 & 2 \int \sin(x) \sqrt{4 - 3 \sin^2(x)} d \sin(x) \\
 & \quad \downarrow \text{241} \\
 & -\frac{2}{9} (4 - 3 \sin^2(x))^{3/2}
 \end{aligned}$$

input `Int[Sqrt[1 + 3*Cos[x]^2]*Sin[2*x],x]`

output `(-2*(4 - 3*Sin[x]^2)^(3/2))/9`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 241 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4878 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Sin[v], x]}, d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Sin[v]/d, u/Cos[v], x], x], x, Sin[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Sin[v], x], u/Cos[v], x]`

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$-\frac{2(1+3\cos(x)^2)^{\frac{3}{2}}}{9}$	13
default	$-\frac{2(1+3\cos(x)^2)^{\frac{3}{2}}}{9}$	13

input `int(sin(2*x)*(1+3*cos(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

output `-2/9*(1+3*cos(x)^2)^(3/2)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \sqrt{1 + 3 \cos^2(x)} \sin(2x) dx = -\frac{2}{9} (3 \cos(x)^2 + 1)^{\frac{3}{2}}$$

input `integrate(sin(2*x)*(1+3*cos(x)^2)^(1/2),x, algorithm="fricas")`output `-2/9*(3*cos(x)^2 + 1)^(3/2)`**Sympy [A] (verification not implemented)**

Time = 0.67 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \sqrt{1 + 3 \cos^2(x)} \sin(2x) dx = -\frac{2(3 \cos^2(x) + 1)^{\frac{3}{2}}}{9}$$

input `integrate(sin(2*x)*(1+3*cos(x)**2)**(1/2),x)`output `-2*(3*cos(x)**2 + 1)**(3/2)/9`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \sqrt{1 + 3 \cos^2(x)} \sin(2x) dx = -\frac{2}{9} (3 \cos(x)^2 + 1)^{\frac{3}{2}}$$

input `integrate(sin(2*x)*(1+3*cos(x)^2)^(1/2),x, algorithm="maxima")`output `-2/9*(3*cos(x)^2 + 1)^(3/2)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \sqrt{1 + 3 \cos^2(x)} \sin(2x) dx = -\frac{2}{9} (3 \cos(x)^2 + 1)^{\frac{3}{2}}$$

input `integrate(sin(2*x)*(1+3*cos(x)^2)^(1/2),x, algorithm="giac")`output `-2/9*(3*cos(x)^2 + 1)^(3/2)`**Mupad [B] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \sqrt{1 + 3 \cos^2(x)} \sin(2x) dx = -\frac{2 (3 \cos(x)^2 + 1)^{3/2}}{9}$$

input `int(sin(2*x)*(3*cos(x)^2 + 1)^(1/2),x)`output `-(2*(3*cos(x)^2 + 1)^(3/2))/9`**Reduce [F]**

$$\int \sqrt{1 + 3 \cos^2(x)} \sin(2x) dx = \int \sqrt{3 \cos(x)^2 + 1} \sin(2x) dx$$

input `int(sin(2*x)*(1+3*cos(x)^2)^(1/2),x)`output `int(sqrt(3*cos(x)**2 + 1)*sin(2*x),x)`

3.56 $\int \frac{1}{2+3x} dx$

Optimal result	373
Mathematica [A] (verified)	373
Rubi [A] (verified)	374
Maple [A] (verified)	375
Fricas [A] (verification not implemented)	375
Sympy [A] (verification not implemented)	376
Maxima [A] (verification not implemented)	376
Giac [A] (verification not implemented)	376
Mupad [B] (verification not implemented)	377
Reduce [B] (verification not implemented)	377

Optimal result

Integrand size = 7, antiderivative size = 10

$$\int \frac{1}{2+3x} dx = \frac{1}{3} \log(2+3x)$$

output `1/3*ln(2+3*x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{2+3x} dx = \frac{1}{3} \log(2+3x)$$

input `Integrate[(2 + 3*x)^(-1),x]`

output `Log[2 + 3*x]/3`

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{3x+2} dx$$

↓ 16

$$\frac{1}{3} \log(3x+2)$$

input

```
Int[(2 + 3*x)^(-1),x]
```

output

```
Log[2 + 3*x]/3
```

Defintions of rubi rules used

rule 16

```
Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] :> Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]
```

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

method	result	size
parallelrisc	$\frac{\ln(\frac{2}{3}+x)}{3}$	7
default	$\frac{\ln(2+3x)}{3}$	9
norman	$\frac{\ln(2+3x)}{3}$	9
meijerg	$\frac{\ln(1+\frac{3x}{2})}{3}$	9
risc	$\frac{\ln(2+3x)}{3}$	9

input `int(1/(2+3*x),x,method=_RETURNVERBOSE)`

output `1/3*ln(2/3+x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{1}{2+3x} dx = \frac{1}{3} \log(3x+2)$$

input `integrate(1/(2+3*x),x, algorithm="fricas")`

output `1/3*log(3*x + 2)`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int \frac{1}{2+3x} dx = \frac{\log(3x+2)}{3}$$

input `integrate(1/(2+3*x),x)`

output `log(3*x + 2)/3`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{1}{2+3x} dx = \frac{1}{3} \log(3x+2)$$

input `integrate(1/(2+3*x),x, algorithm="maxima")`

output `1/3*log(3*x + 2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

$$\int \frac{1}{2+3x} dx = \frac{1}{3} \log(|3x+2|)$$

input `integrate(1/(2+3*x),x, algorithm="giac")`

output `1/3*log(abs(3*x + 2))`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.60

$$\int \frac{1}{2+3x} dx = \frac{\ln\left(x + \frac{2}{3}\right)}{3}$$

input `int(1/(3*x + 2),x)`

output `log(x + 2/3)/3`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{1}{2+3x} dx = \frac{\log(3x + 2)}{3}$$

input `int(1/(2+3*x),x)`

output `log(3*x + 2)/3`

3.57 $\int \log^2(x) dx$

Optimal result	378
Mathematica [A] (verified)	378
Rubi [A] (verified)	379
Maple [A] (verified)	380
Fricas [A] (verification not implemented)	380
Sympy [A] (verification not implemented)	381
Maxima [A] (verification not implemented)	381
Giac [A] (verification not implemented)	381
Mupad [B] (verification not implemented)	382
Reduce [B] (verification not implemented)	382

Optimal result

Integrand size = 4, antiderivative size = 15

$$\int \log^2(x) dx = 2x - 2x \log(x) + x \log^2(x)$$

output

```
2*x-2*x*ln(x)+x*ln(x)^2
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \log^2(x) dx = 2x - 2x \log(x) + x \log^2(x)$$

input

```
Integrate[Log[x]^2,x]
```

output

```
2*x - 2*x*Log[x] + x*Log[x]^2
```

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {2733, 2732}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \log^2(x) dx$$

$$\downarrow 2733$$

$$x \log^2(x) - 2 \int \log(x) dx$$

$$\downarrow 2732$$

$$x \log^2(x) - 2(x \log(x) - x)$$

input `Int [Log[x]^2, x]`

output `x*Log[x]^2 - 2*(-x + x*Log[x])`

Defintions of rubi rules used

rule 2732 `Int [Log[(c_.)*(x_)^(n_.)], x_Symbol] :> Simp[x*Log[c*x^n], x] - Simp[n*x, x] /; FreeQ[{c, n}, x]`

rule 2733 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_.), x_Symbol] :> Simp[x*(a + b*Log[c*x^n])^p, x] - Simp[b*n*p Int[(a + b*Log[c*x^n])^(p - 1), x], x] /; FreeQ[{a, b, c, n}, x] && GtQ[p, 0] && IntegerQ[2*p]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.07

method	result	size
default	$2x - 2x \ln(x) + x \ln(x)^2$	16
norman	$2x - 2x \ln(x) + x \ln(x)^2$	16
risch	$2x - 2x \ln(x) + x \ln(x)^2$	16
parallelrisch	$2x - 2x \ln(x) + x \ln(x)^2$	16
orering	$x \ln(x)^2 + x^3 \left(\frac{2}{x^2} - \frac{2 \ln(x)}{x^2} \right)$	25

input `int(ln(x)^2,x,method=_RETURNVERBOSE)`output `2*x-2*x*ln(x)+x*ln(x)^2`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \log^2(x) dx = x \log(x)^2 - 2x \log(x) + 2x$$

input `integrate(log(x)^2,x, algorithm="fricas")`output `x*log(x)^2 - 2*x*log(x) + 2*x`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \log^2(x) dx = x \log(x)^2 - 2x \log(x) + 2x$$

input `integrate(ln(x)**2,x)`

output `x*log(x)**2 - 2*x*log(x) + 2*x`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

$$\int \log^2(x) dx = (\log(x)^2 - 2 \log(x) + 2)x$$

input `integrate(log(x)^2,x, algorithm="maxima")`

output `(log(x)^2 - 2*log(x) + 2)*x`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \log^2(x) dx = x \log(x)^2 - 2x \log(x) + 2x$$

input `integrate(log(x)^2,x, algorithm="giac")`

output `x*log(x)^2 - 2*x*log(x) + 2*x`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

$$\int \log^2(x) dx = x (\ln(x)^2 - 2 \ln(x) + 2)$$

input `int(log(x)^2,x)`

output `x*(log(x)^2 - 2*log(x) + 2)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

$$\int \log^2(x) dx = x(\log(x)^2 - 2\log(x) + 2)$$

input `int(log(x)^2,x)`

output `x*(log(x)**2 - 2*log(x) + 2)`

3.58 $\int x \log(x) dx$

Optimal result	383
Mathematica [A] (verified)	383
Rubi [A] (verified)	384
Maple [A] (verified)	385
Fricas [A] (verification not implemented)	385
Sympy [A] (verification not implemented)	386
Maxima [A] (verification not implemented)	386
Giac [A] (verification not implemented)	386
Mupad [B] (verification not implemented)	387
Reduce [B] (verification not implemented)	387

Optimal result

Integrand size = 4, antiderivative size = 17

$$\int x \log(x) dx = -\frac{x^2}{4} + \frac{1}{2}x^2 \log(x)$$

output

```
-1/4*x^2+1/2*x^2*ln(x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int x \log(x) dx = -\frac{x^2}{4} + \frac{1}{2}x^2 \log(x)$$

input

```
Integrate[x*Log[x],x]
```

output

```
-1/4*x^2 + (x^2*Log[x])/2
```


Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2741}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \log(x) dx$$

↓ 2741

$$\frac{1}{2}x^2 \log(x) - \frac{x^2}{4}$$

input `Int [x*Log [x] , x]`

output `-1/4*x^2 + (x^2*Log [x])/2`

Defintions of rubi rules used

rule 2741 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((d_.)*(x_)^(m_.), x_Symbol] :>
Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])/(d*(m + 1))), x] - Simp[b*n*((d*x)^(
m + 1)/(d*(m + 1)^2)), x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2}$	14
norman	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2}$	14
risch	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2}$	14
parallelrisch	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2}$	14
parts	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2}$	14
orering	$\frac{3x^2 \ln(x)}{4} - \frac{x^2(1+\ln(x))}{4}$	18

input `int(x*ln(x),x,method=_RETURNVERBOSE)`output `-1/4*x^2+1/2*x^2*ln(x)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int x \log(x) dx = \frac{1}{2} x^2 \log(x) - \frac{1}{4} x^2$$

input `integrate(x*log(x),x, algorithm="fricas")`output `1/2*x^2*log(x) - 1/4*x^2`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

$$\int x \log(x) dx = \frac{x^2 \log(x)}{2} - \frac{x^2}{4}$$

input `integrate(x*ln(x),x)`

output `x**2*log(x)/2 - x**2/4`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int x \log(x) dx = \frac{1}{2} x^2 \log(x) - \frac{1}{4} x^2$$

input `integrate(x*log(x),x, algorithm="maxima")`

output `1/2*x^2*log(x) - 1/4*x^2`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int x \log(x) dx = \frac{1}{2} x^2 \log(x) - \frac{1}{4} x^2$$

input `integrate(x*log(x),x, algorithm="giac")`

output `1/2*x^2*log(x) - 1/4*x^2`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.53

$$\int x \log(x) dx = \frac{x^2 (\ln(x) - \frac{1}{2})}{2}$$

input `int(x*log(x),x)`

output `(x^2*(log(x) - 1/2))/2`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int x \log(x) dx = \frac{x^2(2 \log(x) - 1)}{4}$$

input `int(x*log(x),x)`

output `(x**2*(2*log(x) - 1))/4`

3.59 $\int x \log^2(x) dx$

Optimal result	388
Mathematica [A] (verified)	388
Rubi [A] (verified)	389
Maple [A] (verified)	390
Fricas [A] (verification not implemented)	390
Sympy [A] (verification not implemented)	391
Maxima [A] (verification not implemented)	391
Giac [A] (verification not implemented)	391
Mupad [B] (verification not implemented)	392
Reduce [B] (verification not implemented)	392

Optimal result

Integrand size = 6, antiderivative size = 28

$$\int x \log^2(x) dx = \frac{x^2}{4} - \frac{1}{2}x^2 \log(x) + \frac{1}{2}x^2 \log^2(x)$$

output

```
1/4*x^2-1/2*x^2*ln(x)+1/2*x^2*ln(x)^2
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int x \log^2(x) dx = \frac{x^2}{4} - \frac{1}{2}x^2 \log(x) + \frac{1}{2}x^2 \log^2(x)$$

input

```
Integrate[x*Log[x]^2,x]
```

output

```
x^2/4 - (x^2*Log[x])/2 + (x^2*Log[x]^2)/2
```

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2742, 2741}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \log^2(x) dx$$

$$\downarrow \text{2742}$$

$$\frac{1}{2}x^2 \log^2(x) - \int x \log(x) dx$$

$$\downarrow \text{2741}$$

$$\frac{x^2}{4} + \frac{1}{2}x^2 \log^2(x) - \frac{1}{2}x^2 \log(x)$$

input `Int[x*Log[x]^2,x]`

output `x^2/4 - (x^2*Log[x])/2 + (x^2*Log[x]^2)/2`

Defintions of rubi rules used

rule 2741 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((d_.)*(x_)^(m_.), x_Symbol] :> Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])/(d*(m + 1))), x] - Simp[b*n*((d*x)^(m + 1)/(d*(m + 1)^2)), x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[m, -1]`

rule 2742 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((d_.)*(x_)^(m_.), x_Symbol] :> Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])^p/(d*(m + 1))), x] - Simp[b*n*(p/(m + 1)) Int[(d*x)^m*(a + b*Log[c*x^n])^(p - 1), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[m, -1] && GtQ[p, 0]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.82

method	result	size
default	$\frac{x^2}{4} - \frac{x^2 \ln(x)}{2} + \frac{x^2 \ln(x)^2}{2}$	23
norman	$\frac{x^2}{4} - \frac{x^2 \ln(x)}{2} + \frac{x^2 \ln(x)^2}{2}$	23
risch	$\frac{x^2}{4} - \frac{x^2 \ln(x)}{2} + \frac{x^2 \ln(x)^2}{2}$	23
parallelrisch	$\frac{x^2}{4} - \frac{x^2 \ln(x)}{2} + \frac{x^2 \ln(x)^2}{2}$	23
parts	$\frac{x^2}{4} - \frac{x^2 \ln(x)}{2} + \frac{x^2 \ln(x)^2}{2}$	23
orering	$\frac{7x^2 \ln(x)^2}{8} - \frac{3x^2 (\ln(x)^2 + 2 \ln(x))}{8} + \frac{x^3 (\frac{2 \ln(x)}{x} + \frac{2}{x})}{8}$	43

input `int(x*ln(x)^2,x,method=_RETURNVERBOSE)`output `1/4*x^2-1/2*x^2*ln(x)+1/2*x^2*ln(x)^2`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int x \log^2(x) dx = \frac{1}{2} x^2 \log(x)^2 - \frac{1}{2} x^2 \log(x) + \frac{1}{4} x^2$$

input `integrate(x*log(x)^2,x, algorithm="fricas")`output `1/2*x^2*log(x)^2 - 1/2*x^2*log(x) + 1/4*x^2`

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int x \log^2(x) dx = \frac{x^2 \log(x)^2}{2} - \frac{x^2 \log(x)}{2} + \frac{x^2}{4}$$

input `integrate(x*ln(x)**2,x)`output `x**2*log(x)**2/2 - x**2*log(x)/2 + x**2/4`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.61

$$\int x \log^2(x) dx = \frac{1}{4} (2 \log(x)^2 - 2 \log(x) + 1) x^2$$

input `integrate(x*log(x)^2,x, algorithm="maxima")`output `1/4*(2*log(x)^2 - 2*log(x) + 1)*x^2`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int x \log^2(x) dx = \frac{1}{2} x^2 \log(x)^2 - \frac{1}{2} x^2 \log(x) + \frac{1}{4} x^2$$

input `integrate(x*log(x)^2,x, algorithm="giac")`output `1/2*x^2*log(x)^2 - 1/2*x^2*log(x) + 1/4*x^2`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.61

$$\int x \log^2(x) dx = \frac{x^2 (2 \ln(x)^2 - 2 \ln(x) + 1)}{4}$$

input `int(x*log(x)^2,x)`

output `(x^2*(2*log(x)^2 - 2*log(x) + 1))/4`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.61

$$\int x \log^2(x) dx = \frac{x^2 (2 \log(x)^2 - 2 \log(x) + 1)}{4}$$

input `int(x*log(x)^2,x)`

output `(x**2*(2*log(x)**2 - 2*log(x) + 1))/4`

3.60 $\int \frac{1}{1+t} dt$

Optimal result	393
Mathematica [A] (verified)	393
Rubi [A] (verified)	394
Maple [A] (verified)	394
Fricas [A] (verification not implemented)	395
Sympy [A] (verification not implemented)	395
Maxima [A] (verification not implemented)	396
Giac [A] (verification not implemented)	396
Mupad [B] (verification not implemented)	396
Reduce [B] (verification not implemented)	397

Optimal result

Integrand size = 5, antiderivative size = 4

$$\int \frac{1}{1+t} dt = \log(1+t)$$

output `ln(1+t)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+t} dt = \log(1+t)$$

input `Integrate[(1 + t)^(-1),t]`

output `Log[1 + t]`

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{t+1} dt$$

↓ 16

$$\log(t+1)$$

input `Int[(1 + t)^(-1), t]`

output `Log[1 + t]`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] :> Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

method	result	size
default	$\ln(1+t)$	5
norman	$\ln(1+t)$	5
meijerg	$\ln(1+t)$	5
risch	$\ln(1+t)$	5
parallelrisch	$\ln(1+t)$	5

input `int(1/(1+t),t,method=_RETURNVERBOSE)`

output `ln(1+t)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+t} dt = \log(t+1)$$

input `integrate(1/(1+t),t, algorithm="fricas")`

output `log(t + 1)`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int \frac{1}{1+t} dt = \log(t+1)$$

input `integrate(1/(1+t),t)`

output `log(t + 1)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+t} dt = \log(t+1)$$

input `integrate(1/(1+t),t, algorithm="maxima")`

output `log(t + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

$$\int \frac{1}{1+t} dt = \log(|t+1|)$$

input `integrate(1/(1+t),t, algorithm="giac")`

output `log(abs(t + 1))`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+t} dt = \ln(t+1)$$

input `int(1/(t + 1),t)`

output `log(t + 1)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+t} dt = \log(t+1)$$

input `int(1/(1+t),t)`

output `log(t + 1)`

3.61 $\int \cot(x) dx$

Optimal result	398
Mathematica [A] (verified)	398
Rubi [A] (verified)	399
Maple [A] (verified)	400
Fricas [B] (verification not implemented)	400
Sympy [A] (verification not implemented)	401
Maxima [A] (verification not implemented)	401
Giac [A] (verification not implemented)	401
Mupad [B] (verification not implemented)	402
Reduce [B] (verification not implemented)	402

Optimal result

Integrand size = 2, antiderivative size = 3

$$\int \cot(x) dx = \log(\sin(x))$$

output `ln(sin(x))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \cot(x) dx = \log(\sin(x))$$

input `Integrate[Cot[x], x]`

output `Log[Sin[x]]`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$, Rules used = {3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \cot(x) dx \\ \downarrow 3042 \\ \int -\tan\left(x + \frac{\pi}{2}\right) dx \\ \downarrow 25 \\ -\int \tan\left(x + \frac{\pi}{2}\right) dx \\ \downarrow 3956 \\ \log(\sin(x)) \end{array}$$

input `Int[Cot[x],x]`

output `Log[Sin[x]]`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
lookup	$\ln(\sin(x))$	4
default	$\ln(\sin(x))$	4
derivativedivides	$-\frac{\ln(\cot(x)^2+1)}{2}$	10
parallelrisc	$\ln\left(\frac{1}{\sqrt{\sec(x)^2}}\right) + \ln(\tan(x))$	12
norman	$-\frac{\ln(1+\tan(x)^2)}{2} + \ln(\tan(x))$	14
risc	$-ix + \ln(e^{2ix} - 1)$	14

input `int(cot(x),x,method=_RETURNVERBOSE)`

output `ln(sin(x))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 11 vs. $2(3) = 6$.

Time = 0.09 (sec) , antiderivative size = 11, normalized size of antiderivative = 3.67

$$\int \cot(x) dx = \frac{1}{2} \log\left(-\frac{1}{2} \cos(2x) + \frac{1}{2}\right)$$

input `integrate(cot(x),x, algorithm="fricas")`

output `1/2*log(-1/2*cos(2*x) + 1/2)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \cot(x) dx = \log(\sin(x))$$

input `integrate(cot(x),x)`

output `log(sin(x))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \cot(x) dx = \log(\sin(x))$$

input `integrate(cot(x),x, algorithm="maxima")`

output `log(sin(x))`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

$$\int \cot(x) dx = \log(|\sin(x)|)$$

input `integrate(cot(x),x, algorithm="giac")`

output `log(abs(sin(x)))`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \cot(x) dx = \ln(\sin(x))$$

input `int(cot(x),x)`

output `log(sin(x))`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 5.67

$$\int \cot(x) dx = -\log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) + \log\left(\tan\left(\frac{x}{2}\right)\right)$$

input `int(cot(x),x)`

output `- log(tan(x/2)**2 + 1) + log(tan(x/2))`

3.62 $\int x^n \log(ax) dx$

Optimal result	403
Mathematica [A] (verified)	403
Rubi [A] (verified)	404
Maple [A] (verified)	405
Fricas [A] (verification not implemented)	405
Sympy [B] (verification not implemented)	406
Maxima [A] (verification not implemented)	406
Giac [A] (verification not implemented)	406
Mupad [B] (verification not implemented)	407
Reduce [B] (verification not implemented)	407

Optimal result

Integrand size = 8, antiderivative size = 28

$$\int x^n \log(ax) dx = -\frac{x^{1+n}}{(1+n)^2} + \frac{x^{1+n} \log(ax)}{1+n}$$

output

```
-x^(1+n)/(1+n)^2+x^(1+n)*ln(a*x)/(1+n)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.75

$$\int x^n \log(ax) dx = \frac{x^{1+n}(-1 + (1+n) \log(ax))}{(1+n)^2}$$

input

```
Integrate[x^n*Log[a*x],x]
```

output

```
(x^(1+n)*(-1+(1+n)*Log[a*x]))/(1+n)^2
```

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2741}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^n \log(ax) dx$$

$$\downarrow 2741$$

$$\frac{x^{n+1} \log(ax)}{n+1} - \frac{x^{n+1}}{(n+1)^2}$$

input `Int [x^n*Log [a*x] , x]`

output `-(x^(1 + n)/(1 + n)^2) + (x^(1 + n)*Log[a*x])/(1 + n)`

Defintions of rubi rules used

rule 2741

```
Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((d_.)*(x_)^(m_.), x_Symbol] :>
Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])/(d*(m + 1))), x] - Simp[b*n*((d*x)^(
m + 1)/(d*(m + 1)^2)), x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.18

method	result
parallelsch	$\frac{x x^n \ln(ax)n+x^n \ln(ax)x-x^n x}{(1+n)^2}$
norman	$\frac{x \ln(ax)e^{n \ln(x)}}{1+n} - \frac{x e^{n \ln(x)}}{n^2+2n+1}$
oring	$\frac{x(2n+1)x^n \ln(ax)}{n^2+2n+1} - \frac{x^2 \left(\frac{x^n n \ln(ax)}{x} + \frac{x^n}{x} \right)}{n^2+2n+1}$
risch	$x \left(i\pi \operatorname{csgn}(ix) \operatorname{csgn}(iax)^2 n - i\pi \operatorname{csgn}(ix) \operatorname{csgn}(iax) \operatorname{csgn}(ia) n - i\pi \operatorname{csgn}(iax)^3 n + i\pi \operatorname{csgn}(iax)^2 \operatorname{csgn}(ia) n + i\pi \operatorname{csgn}(ix) \operatorname{csgn}(ia) \right) / (1+n)^2$

input `int(x^n*ln(a*x),x,method=_RETURNVERBOSE)`output `(x*x^n*ln(a*x)*n+x^n*ln(a*x)*x-x^n*x)/(1+n)^2`**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.14

$$\int x^n \log(ax) dx = \frac{((n+1)x \log(a) + (n+1)x \log(x) - x)x^n}{n^2 + 2n + 1}$$

input `integrate(x^n*log(a*x),x, algorithm="fricas")`output `((n + 1)*x*log(a) + (n + 1)*x*log(x) - x)*x^n/(n^2 + 2*n + 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 61 vs. $2(22) = 44$.

Time = 0.29 (sec) , antiderivative size = 61, normalized size of antiderivative = 2.18

$$\int x^n \log(ax) dx = \begin{cases} \frac{nx^n \log(ax)}{n^2+2n+1} + \frac{xx^n \log(ax)}{n^2+2n+1} - \frac{xx^n}{n^2+2n+1} & \text{for } n \neq -1 \\ \frac{\log(ax)^2}{2} & \text{otherwise} \end{cases}$$

input `integrate(x**n*ln(a*x),x)`

output `Piecewise((n*x*x**n*log(a*x)/(n**2 + 2*n + 1) + x*x**n*log(a*x)/(n**2 + 2*n + 1) - x*x**n/(n**2 + 2*n + 1), Ne(n, -1)), (log(a*x)**2/2, True))`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int x^n \log(ax) dx = \frac{x^{n+1} \log(ax)}{n+1} - \frac{x^{n+1}}{(n+1)^2}$$

input `integrate(x^n*log(a*x),x, algorithm="maxima")`

output `x^(n + 1)*log(a*x)/(n + 1) - x^(n + 1)/(n + 1)^2`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int x^n \log(ax) dx = \frac{x^{n+1} \log(ax)}{n+1} - \frac{x^{n+1}}{(n+1)^2}$$

input `integrate(x^n*log(a*x),x, algorithm="giac")`

output `x^(n + 1)*log(a*x)/(n + 1) - x^(n + 1)/(n + 1)^2`

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.36

$$\int x^n \log(ax) dx = \begin{cases} \frac{\ln(ax)^2}{2} & \text{if } n = -1 \\ \frac{x^{n+1} \left(\ln(ax) - \frac{1}{n+1} \right)}{n+1} & \text{if } n \neq -1 \end{cases}$$

input `int(x^n*log(a*x),x)`

output `piecewise(n == -1, log(a*x)^2/2, n ~= -1, (x^(n + 1)*(log(a*x) - 1/(n + 1)))/(n + 1))`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int x^n \log(ax) dx = \frac{x^n x (\log(ax) n + \log(ax) - 1)}{n^2 + 2n + 1}$$

input `int(x^n*log(a*x),x)`

output `(x**n*x*(log(a*x)*n + log(a*x) - 1))/(n**2 + 2*n + 1)`

3.63 $\int x^2 \log^2(x) dx$

Optimal result	408
Mathematica [A] (verified)	408
Rubi [A] (verified)	409
Maple [A] (verified)	410
Fricas [A] (verification not implemented)	410
Sympy [A] (verification not implemented)	411
Maxima [A] (verification not implemented)	411
Giac [A] (verification not implemented)	411
Mupad [B] (verification not implemented)	412
Reduce [B] (verification not implemented)	412

Optimal result

Integrand size = 8, antiderivative size = 28

$$\int x^2 \log^2(x) dx = \frac{2x^3}{27} - \frac{2}{9}x^3 \log(x) + \frac{1}{3}x^3 \log^2(x)$$

output

```
2/27*x^3-2/9*x^3*ln(x)+1/3*x^3*ln(x)^2
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int x^2 \log^2(x) dx = \frac{2x^3}{27} - \frac{2}{9}x^3 \log(x) + \frac{1}{3}x^3 \log^2(x)$$

input

```
Integrate[x^2*Log[x]^2,x]
```

output

```
(2*x^3)/27 - (2*x^3*Log[x])/9 + (x^3*Log[x]^2)/3
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.18, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2742, 2741}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \log^2(x) dx$$

$$\downarrow 2742$$

$$\frac{1}{3}x^3 \log^2(x) - \frac{2}{3} \int x^2 \log(x) dx$$

$$\downarrow 2741$$

$$\frac{1}{3}x^3 \log^2(x) - \frac{2}{3} \left(\frac{1}{3}x^3 \log(x) - \frac{x^3}{9} \right)$$

input

```
Int[x^2*Log[x]^2,x]
```

output

```
(x^3*Log[x]^2)/3 - (2*(-1/9*x^3 + (x^3*Log[x])/3))/3
```

Defintions of rubi rules used

rule 2741

```
Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((d_.)*(x_)^(m_.), x_Symbol] :=
Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])/(d*(m + 1))), x] - Simp[b*n*((d*x)^(
m + 1)/(d*(m + 1)^2)), x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[m, -1]
```

rule 2742

```
Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((d_.)*(x_)^(m_.), x_Symbol] :=
Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])^p/(d*(m + 1))), x] - Simp[b*n*
(p/(m + 1)) Int[(d*x)^m*(a + b*Log[c*x^n])^(p - 1), x], x] /; FreeQ[{a, b
, c, d, m, n}, x] && NeQ[m, -1] && GtQ[p, 0]
```

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.82

method	result	size
default	$\frac{2x^3}{27} - \frac{2x^3 \ln(x)}{9} + \frac{x^3 \ln(x)^2}{3}$	23
norman	$\frac{2x^3}{27} - \frac{2x^3 \ln(x)}{9} + \frac{x^3 \ln(x)^2}{3}$	23
risch	$\frac{2x^3}{27} - \frac{2x^3 \ln(x)}{9} + \frac{x^3 \ln(x)^2}{3}$	23
parallelrisch	$\frac{2x^3}{27} - \frac{2x^3 \ln(x)}{9} + \frac{x^3 \ln(x)^2}{3}$	23
parts	$\frac{2x^3}{27} - \frac{2x^3 \ln(x)}{9} + \frac{x^3 \ln(x)^2}{3}$	23
orering	$\frac{19x^3 \ln(x)^2}{27} - \frac{2x^2(2x \ln(x)^2 + 2x \ln(x))}{9} + \frac{x^3(2 \ln(x)^2 + 6 \ln(x) + 2)}{27}$	46

input `int(x^2*ln(x)^2,x,method=_RETURNVERBOSE)`output `2/27*x^3-2/9*x^3*ln(x)+1/3*x^3*ln(x)^2`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int x^2 \log^2(x) dx = \frac{1}{3} x^3 \log(x)^2 - \frac{2}{9} x^3 \log(x) + \frac{2}{27} x^3$$

input `integrate(x^2*log(x)^2,x, algorithm="fricas")`output `1/3*x^3*log(x)^2 - 2/9*x^3*log(x) + 2/27*x^3`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int x^2 \log^2(x) dx = \frac{x^3 \log(x)^2}{3} - \frac{2x^3 \log(x)}{9} + \frac{2x^3}{27}$$

input `integrate(x**2*ln(x)**2,x)`output `x**3*log(x)**2/3 - 2*x**3*log(x)/9 + 2*x**3/27`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.61

$$\int x^2 \log^2(x) dx = \frac{1}{27} (9 \log(x)^2 - 6 \log(x) + 2) x^3$$

input `integrate(x^2*log(x)^2,x, algorithm="maxima")`output `1/27*(9*log(x)^2 - 6*log(x) + 2)*x^3`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int x^2 \log^2(x) dx = \frac{1}{3} x^3 \log(x)^2 - \frac{2}{9} x^3 \log(x) + \frac{2}{27} x^3$$

input `integrate(x^2*log(x)^2,x, algorithm="giac")`output `1/3*x^3*log(x)^2 - 2/9*x^3*log(x) + 2/27*x^3`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.61

$$\int x^2 \log^2(x) dx = \frac{2x^3 \left(\frac{9\ln(x)^2}{2} - 3\ln(x) + 1 \right)}{27}$$

input `int(x^2*log(x)^2,x)`

output `(2*x^3*((9*log(x)^2)/2 - 3*log(x) + 1))/27`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.61

$$\int x^2 \log^2(x) dx = \frac{x^3(9\log(x)^2 - 6\log(x) + 2)}{27}$$

input `int(x^2*log(x)^2,x)`

output `(x**3*(9*log(x)**2 - 6*log(x) + 2))/27`

3.64 $\int \frac{1}{x \log(x)} dx$

Optimal result	413
Mathematica [A] (verified)	413
Rubi [A] (verified)	414
Maple [A] (verified)	415
Fricas [A] (verification not implemented)	415
Sympy [A] (verification not implemented)	415
Maxima [A] (verification not implemented)	416
Giac [A] (verification not implemented)	416
Mupad [B] (verification not implemented)	416
Reduce [B] (verification not implemented)	417

Optimal result

Integrand size = 8, antiderivative size = 3

$$\int \frac{1}{x \log(x)} dx = \log(\log(x))$$

output `ln(ln(x))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{x \log(x)} dx = \log(\log(x))$$

input `Integrate[1/(x*Log[x]),x]`

output `Log[Log[x]]`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2739, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{1}{x \log(x)} dx \\ \downarrow 2739 \\ \int \frac{1}{\log(x)} d\log(x) \\ \downarrow 14 \\ \log(\log(x)) \end{array}$$

input `Int[1/(x*Log[x]), x]`

output `Log[Log[x]]`

Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] :> Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2739 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)/(x_), x_Symbol] :> Simp[1/(b*n) Subst[Int[x^p, x], x, a + b*Log[c*x^n]], x] /; FreeQ[{a, b, c, n, p}, x]`

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
derivativedivides	$\ln(\ln(x))$	4
default	$\ln(\ln(x))$	4
norman	$\ln(\ln(x))$	4
risch	$\ln(\ln(x))$	4
parallelrisch	$\ln(\ln(x))$	4

input `int(1/x/ln(x),x,method=_RETURNVERBOSE)`output `ln(ln(x))`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{x \log(x)} dx = \log(\log(x))$$

input `integrate(1/x/log(x),x, algorithm="fricas")`output `log(log(x))`**Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{x \log(x)} dx = \log(\log(x))$$

input `integrate(1/x/ln(x),x)`

output `log(log(x))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{x \log(x)} dx = \log(\log(x))$$

input `integrate(1/x/log(x),x, algorithm="maxima")`

output `log(log(x))`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

$$\int \frac{1}{x \log(x)} dx = \log(|\log(x)|)$$

input `integrate(1/x/log(x),x, algorithm="giac")`

output `log(abs(log(x)))`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{x \log(x)} dx = \ln(\ln(x))$$

input `int(1/(x*log(x)),x)`

output `log(log(x))`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{x \log(x)} dx = \log(\log(x))$$

input `int(1/x/log(x),x)`

output `log(log(x))`

3.65 $\int \frac{\log(1-t)}{1-t} dt$

Optimal result	418
Mathematica [A] (verified)	418
Rubi [A] (verified)	419
Maple [A] (verified)	420
Fricas [A] (verification not implemented)	420
Sympy [A] (verification not implemented)	421
Maxima [A] (verification not implemented)	421
Giac [A] (verification not implemented)	421
Mupad [B] (verification not implemented)	422
Reduce [B] (verification not implemented)	422

Optimal result

Integrand size = 14, antiderivative size = 12

$$\int \frac{\log(1-t)}{1-t} dt = -\frac{1}{2} \log^2(1-t)$$

output

```
-1/2*ln(1-t)^2
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\log(1-t)}{1-t} dt = -\frac{1}{2} \log^2(1-t)$$

input

```
Integrate[Log[1 - t]/(1 - t),t]
```

output

```
-1/2*Log[1 - t]^2
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2837, 2738}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\log(1-t)}{1-t} dt \\ & \quad \downarrow \text{2837} \\ & - \int \frac{\log(1-t)}{1-t} d(1-t) \\ & \quad \downarrow \text{2738} \\ & -\frac{1}{2} \log^2(1-t) \end{aligned}$$

input `Int[Log[1 - t]/(1 - t),t]`

output `-1/2*Log[1 - t]^2`

Defintions of rubi rules used

rule 2738 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))/(x_), x_Symbol] := Simp[(a + b*Log[c*x^n])^2/(2*b*n), x] /; FreeQ[{a, b, c, n}, x]`

rule 2837 `Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_.)]*(b_.))^(p_.)*((f_) + (g_.)*(x_)^(q_.)), x_Symbol] := Simp[1/e Subst[Int[(f*(x/d))^q*(a + b*Log[c*x^n])^p, x], x, d + e*x], x] /; FreeQ[{a, b, c, d, e, f, g, n, p, q}, x] && EqQ[e*f - d*g, 0]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

method	result	size
derivativedivides	$-\frac{\ln(1-t)^2}{2}$	11
default	$-\frac{\ln(1-t)^2}{2}$	11
norman	$-\frac{\ln(1-t)^2}{2}$	11
risch	$-\frac{\ln(1-t)^2}{2}$	11
parts	$-\ln(-1+t)\ln(1-t) + \frac{\ln(-1+t)^2}{2}$	22

input `int(ln(1-t)/(1-t),t,method=_RETURNVERBOSE)`output `-1/2*ln(1-t)^2`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\log(1-t)}{1-t} dt = -\frac{1}{2} \log(-t+1)^2$$

input `integrate(log(1-t)/(1-t),t, algorithm="fricas")`output `-1/2*log(-t + 1)^2`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{\log(1-t)}{1-t} dt = -\frac{\log(1-t)^2}{2}$$

input `integrate(ln(1-t)/(1-t),t)`

output `-log(1 - t)**2/2`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\log(1-t)}{1-t} dt = -\frac{1}{2} \log(-t+1)^2$$

input `integrate(log(1-t)/(1-t),t, algorithm="maxima")`

output `-1/2*log(-t + 1)^2`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\log(1-t)}{1-t} dt = -\frac{1}{2} \log(-t+1)^2$$

input `integrate(log(1-t)/(1-t),t, algorithm="giac")`

output `-1/2*log(-t + 1)^2`

Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\log(1-t)}{1-t} dt = -\frac{\ln(1-t)^2}{2}$$

input `int(-log(1 - t)/(t - 1),t)`

output `-log(1 - t)^2/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\log(1-t)}{1-t} dt = -\frac{\log(-t+1)^2}{2}$$

input `int(log(1-t)/(1-t),t)`

output `(- log(- t + 1)**2)/2`

3.66 $\int \frac{\log(x)}{x\sqrt{1+\log(x)}} dx$

Optimal result	423
Mathematica [A] (verified)	423
Rubi [A] (verified)	424
Maple [A] (verified)	425
Fricas [A] (verification not implemented)	425
Sympy [A] (verification not implemented)	426
Maxima [A] (verification not implemented)	426
Giac [A] (verification not implemented)	426
Mupad [B] (verification not implemented)	427
Reduce [B] (verification not implemented)	427

Optimal result

Integrand size = 14, antiderivative size = 23

$$\int \frac{\log(x)}{x\sqrt{1+\log(x)}} dx = -2\sqrt{1+\log(x)} + \frac{2}{3}(1+\log(x))^{3/2}$$

output `2/3*(1+ln(x))^(3/2)-2*(1+ln(x))^(1/2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.70

$$\int \frac{\log(x)}{x\sqrt{1+\log(x)}} dx = \frac{2}{3}(-2+\log(x))\sqrt{1+\log(x)}$$

input `Integrate[Log[x]/(x*Sqrt[1 + Log[x]]),x]`

output `(2*(-2 + Log[x])*Sqrt[1 + Log[x]])/3`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {2812, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\log(x)}{x\sqrt{\log(x)+1}} dx$$

↓ 2812

$$\int \frac{\log(x)}{\sqrt{\log(x)+1}} d\log(x)$$

↓ 53

$$\int \left(\sqrt{\log(x)+1} - \frac{1}{\sqrt{\log(x)+1}} \right) d\log(x)$$

↓ 2009

$$\frac{2}{3}(\log(x)+1)^{3/2} - 2\sqrt{\log(x)+1}$$

input `Int[Log[x]/(x*Sqrt[1 + Log[x]]),x]`

output `-2*Sqrt[1 + Log[x]] + (2*(1 + Log[x])^(3/2))/3`

Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2812

```
Int[(((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((d_.) + Log[(c_.)*(x_)^(n_.)]*(e_.))^(q_.))/(x_), x_Symbol] := Simp[1/n Subst[Int[(a + b*x)^p*(d + e*x)^q, x], x, Log[c*x^n]], x] /; FreeQ[{a, b, c, d, e, n, p, q}, x]
```

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

method	result	size
derivativedivides	$\frac{2(1+\ln(x))^{\frac{3}{2}}}{3} - 2\sqrt{1+\ln(x)}$	18
default	$\frac{2(1+\ln(x))^{\frac{3}{2}}}{3} - 2\sqrt{1+\ln(x)}$	18

input

```
int(ln(x)/x/(1+ln(x))^(1/2),x,method=_RETURNVERBOSE)
```

output

```
2/3*(1+ln(x))^(3/2)-2*(1+ln(x))^(1/2)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.52

$$\int \frac{\log(x)}{x\sqrt{1+\log(x)}} dx = \frac{2}{3}\sqrt{\log(x)+1}(\log(x)-2)$$

input

```
integrate(log(x)/x/(1+log(x))^(1/2),x, algorithm="fricas")
```

output

```
2/3*sqrt(log(x) + 1)*(log(x) - 2)
```

Sympy [A] (verification not implemented)

Time = 0.87 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \frac{\log(x)}{x\sqrt{1+\log(x)}} dx = \frac{2(\log(x)+1)^{\frac{3}{2}}}{3} - 2\sqrt{\log(x)+1}$$

input `integrate(ln(x)/x/(1+ln(x))**(1/2),x)`output `2*(log(x) + 1)**(3/2)/3 - 2*sqrt(log(x) + 1)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{\log(x)}{x\sqrt{1+\log(x)}} dx = \frac{2}{3}(\log(x)+1)^{\frac{3}{2}} - 2\sqrt{\log(x)+1}$$

input `integrate(log(x)/x/(1+log(x))^(1/2),x, algorithm="maxima")`output `2/3*(log(x) + 1)^(3/2) - 2*sqrt(log(x) + 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{\log(x)}{x\sqrt{1+\log(x)}} dx = \frac{2}{3}(\log(x)+1)^{\frac{3}{2}} - 2\sqrt{\log(x)+1}$$

input `integrate(log(x)/x/(1+log(x))^(1/2),x, algorithm="giac")`output `2/3*(log(x) + 1)^(3/2) - 2*sqrt(log(x) + 1)`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.57

$$\int \frac{\log(x)}{x\sqrt{1+\log(x)}} dx = \sqrt{\ln(x)+1} \left(\frac{2 \ln(x)}{3} - \frac{4}{3} \right)$$

input `int(log(x)/(x*(log(x) + 1)^(1/2)),x)`

output `(log(x) + 1)^(1/2)*((2*log(x))/3 - 4/3)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.48

$$\int \frac{\log(x)}{x\sqrt{1+\log(x)}} dx = \frac{2\sqrt{\log(x)+1}(\log(x)-2)}{3}$$

input `int(log(x)/x/(1+log(x))^(1/2),x)`

output `(2*sqrt(log(x) + 1)*(log(x) - 2))/3`

3.67 $\int x^3 \log^3(x) dx$

Optimal result	428
Mathematica [A] (verified)	428
Rubi [A] (verified)	429
Maple [A] (verified)	430
Fricas [A] (verification not implemented)	430
Sympy [A] (verification not implemented)	431
Maxima [A] (verification not implemented)	431
Giac [A] (verification not implemented)	431
Mupad [B] (verification not implemented)	432
Reduce [B] (verification not implemented)	432

Optimal result

Integrand size = 8, antiderivative size = 39

$$\int x^3 \log^3(x) dx = -\frac{3x^4}{128} + \frac{3}{32}x^4 \log(x) - \frac{3}{16}x^4 \log^2(x) + \frac{1}{4}x^4 \log^3(x)$$

output

```
-3/128*x^4+3/32*x^4*ln(x)-3/16*x^4*ln(x)^2+1/4*x^4*ln(x)^3
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00

$$\int x^3 \log^3(x) dx = -\frac{3x^4}{128} + \frac{3}{32}x^4 \log(x) - \frac{3}{16}x^4 \log^2(x) + \frac{1}{4}x^4 \log^3(x)$$

input

```
Integrate[x^3*Log[x]^3,x]
```

output

```
(-3*x^4)/128 + (3*x^4*Log[x])/32 - (3*x^4*Log[x]^2)/16 + (x^4*Log[x]^3)/4
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.26, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {2742, 2742, 2741}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \log^3(x) dx$$

$$\downarrow 2742$$

$$\frac{1}{4}x^4 \log^3(x) - \frac{3}{4} \int x^3 \log^2(x) dx$$

$$\downarrow 2742$$

$$\frac{1}{4}x^4 \log^3(x) - \frac{3}{4} \left(\frac{1}{4}x^4 \log^2(x) - \frac{1}{2} \int x^3 \log(x) dx \right)$$

$$\downarrow 2741$$

$$\frac{1}{4}x^4 \log^3(x) - \frac{3}{4} \left(\frac{1}{4}x^4 \log^2(x) + \frac{1}{2} \left(\frac{x^4}{16} - \frac{1}{4}x^4 \log(x) \right) \right)$$

input `Int[x^3*Log[x]^3,x]`

output `(x^4*Log[x]^3)/4 - (3*((x^4*Log[x]^2)/4 + (x^4/16 - (x^4*Log[x])/4)/2))/4`

Defintions of rubi rules used

rule 2741

```
Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((d_.)*(x_))^(m_.), x_Symbol] :>
Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])/(d*(m + 1))), x] - Simp[b*n*((d*x)^(
m + 1)/(d*(m + 1)^2)), x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[m, -1]
```

rule 2742

```
Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((d_.)*(x_)^(m_.), x_Symbol)
] := Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])^p/(d*(m + 1))), x] - Simp[b*n*(p/(m + 1))
Int[(d*x)^m*(a + b*Log[c*x^n])^(p - 1), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[m, -1] && GtQ[p, 0]
```

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.82

method	result
default	$-\frac{3x^4}{128} + \frac{3x^4 \ln(x)}{32} - \frac{3x^4 \ln(x)^2}{16} + \frac{x^4 \ln(x)^3}{4}$
norman	$-\frac{3x^4}{128} + \frac{3x^4 \ln(x)}{32} - \frac{3x^4 \ln(x)^2}{16} + \frac{x^4 \ln(x)^3}{4}$
risch	$-\frac{3x^4}{128} + \frac{3x^4 \ln(x)}{32} - \frac{3x^4 \ln(x)^2}{16} + \frac{x^4 \ln(x)^3}{4}$
parallelrisch	$-\frac{3x^4}{128} + \frac{3x^4 \ln(x)}{32} - \frac{3x^4 \ln(x)^2}{16} + \frac{x^4 \ln(x)^3}{4}$
parts	$-\frac{3x^4}{128} + \frac{3x^4 \ln(x)}{32} - \frac{3x^4 \ln(x)^2}{16} + \frac{x^4 \ln(x)^3}{4}$
orering	$\frac{175x^4 \ln(x)^3}{256} - \frac{55x^2(3x^2 \ln(x)^3 + 3x^2 \ln(x)^2)}{256} + \frac{5x^3(6x \ln(x)^3 + 15x \ln(x)^2 + 6x \ln(x))}{128} - \frac{x^4(6 \ln(x)^3 + 33 \ln(x)^2 + 36 \ln(x) + 16)}{256}$

input

```
int(x^3*ln(x)^3,x,method=_RETURNVERBOSE)
```

output

```
-3/128*x^4+3/32*x^4*ln(x)-3/16*x^4*ln(x)^2+1/4*x^4*ln(x)^3
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.79

$$\int x^3 \log^3(x) dx = \frac{1}{4} x^4 \log(x)^3 - \frac{3}{16} x^4 \log(x)^2 + \frac{3}{32} x^4 \log(x) - \frac{3}{128} x^4$$

input

```
integrate(x^3*log(x)^3,x, algorithm="fricas")
```

output

```
1/4*x^4*log(x)^3 - 3/16*x^4*log(x)^2 + 3/32*x^4*log(x) - 3/128*x^4
```

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.95

$$\int x^3 \log^3(x) dx = \frac{x^4 \log(x)^3}{4} - \frac{3x^4 \log(x)^2}{16} + \frac{3x^4 \log(x)}{32} - \frac{3x^4}{128}$$

input `integrate(x**3*ln(x)**3,x)`output `x**4*log(x)**3/4 - 3*x**4*log(x)**2/16 + 3*x**4*log(x)/32 - 3*x**4/128`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.59

$$\int x^3 \log^3(x) dx = \frac{1}{128} (32 \log(x)^3 - 24 \log(x)^2 + 12 \log(x) - 3)x^4$$

input `integrate(x^3*log(x)^3,x, algorithm="maxima")`output `1/128*(32*log(x)^3 - 24*log(x)^2 + 12*log(x) - 3)*x^4`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.79

$$\int x^3 \log^3(x) dx = \frac{1}{4} x^4 \log(x)^3 - \frac{3}{16} x^4 \log(x)^2 + \frac{3}{32} x^4 \log(x) - \frac{3}{128} x^4$$

input `integrate(x^3*log(x)^3,x, algorithm="giac")`output `1/4*x^4*log(x)^3 - 3/16*x^4*log(x)^2 + 3/32*x^4*log(x) - 3/128*x^4`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.59

$$\int x^3 \log^3(x) dx = \frac{3x^4 \left(\frac{32 \ln(x)^3}{3} - 8 \ln(x)^2 + 4 \ln(x) - 1 \right)}{128}$$

input `int(x^3*log(x)^3,x)`output `(3*x^4*(4*log(x) - 8*log(x)^2 + (32*log(x)^3)/3 - 1))/128`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.59

$$\int x^3 \log^3(x) dx = \frac{x^4 (32 \log(x)^3 - 24 \log(x)^2 + 12 \log(x) - 3)}{128}$$

input `int(x^3*log(x)^3,x)`output `(x**4*(32*log(x)**3 - 24*log(x)**2 + 12*log(x) - 3))/128`

3.68 $\int e^{x^3} x^2 dx$

Optimal result	433
Mathematica [A] (verified)	433
Rubi [A] (verified)	434
Maple [A] (verified)	434
Fricas [A] (verification not implemented)	435
Sympy [A] (verification not implemented)	436
Maxima [A] (verification not implemented)	436
Giac [A] (verification not implemented)	436
Mupad [B] (verification not implemented)	437
Reduce [B] (verification not implemented)	437

Optimal result

Integrand size = 9, antiderivative size = 9

$$\int e^{x^3} x^2 dx = \frac{e^{x^3}}{3}$$

output `1/3*exp(x^3)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int e^{x^3} x^2 dx = \frac{e^{x^3}}{3}$$

input `Integrate[E^x^3*x^2,x]`

output `E^x^3/3`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {2638}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{x^3} x^2 dx$$

$$\downarrow 2638$$

$$\frac{e^{x^3}}{3}$$

input `Int [E^x^3*x^2,x]`

output `E^x^3/3`

Defintions of rubi rules used

rule 2638

```
Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^(n_)))*((e_.) + (f_.)*(x_)^(m_.), x_Symbol]
:> Simp[(e + f*x)^n*(F^(a + b*(c + d*x)^n)/(b*f*n*(c + d*x)^n *Log[F])), x]
;/; FreeQ[{F, a, b, c, d, e, f, n}, x] && EqQ[m, n - 1] && EqQ[d*e - c*f, 0]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

method	result	size
gospers	$\frac{e^{x^3}}{3}$	7
derivativdivides	$\frac{e^{x^3}}{3}$	7
default	$\frac{e^{x^3}}{3}$	7
norman	$\frac{e^{x^3}}{3}$	7
risch	$\frac{e^{x^3}}{3}$	7
parallelrisch	$\frac{e^{x^3}}{3}$	7
orering	$\frac{e^{x^3}}{3}$	7
meijerg	$-\frac{1}{3} + \frac{e^{x^3}}{3}$	9

input `int(exp(x^3)*x^2,x,method=_RETURNVERBOSE)`

output `1/3*exp(x^3)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.67

$$\int e^{x^3} x^2 dx = \frac{1}{3} e^{(x^3)}$$

input `integrate(exp(x^3)*x^2,x, algorithm="fricas")`

output `1/3*e^(x^3)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.56

$$\int e^{x^3} x^2 dx = \frac{e^{x^3}}{3}$$

input `integrate(exp(x**3)*x**2,x)`

output `exp(x**3)/3`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.67

$$\int e^{x^3} x^2 dx = \frac{1}{3} e^{(x^3)}$$

input `integrate(exp(x^3)*x^2,x, algorithm="maxima")`

output `1/3*e^(x^3)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.67

$$\int e^{x^3} x^2 dx = \frac{1}{3} e^{(x^3)}$$

input `integrate(exp(x^3)*x^2,x, algorithm="giac")`

output `1/3*e^(x^3)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.67

$$\int e^{x^3} x^2 dx = \frac{e^{x^3}}{3}$$

input `int(x^2*exp(x^3),x)`

output `exp(x^3)/3`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int e^{x^3} x^2 dx = \frac{e^{x^3}}{3}$$

input `int(exp(x^3)*x^2,x)`

output `e**(x**3)/3`

3.69 $\int \frac{2^{\sqrt{x}}}{\sqrt{x}} dx$

Optimal result	438
Mathematica [A] (verified)	438
Rubi [A] (verified)	439
Maple [A] (verified)	439
Fricas [A] (verification not implemented)	440
Sympy [A] (verification not implemented)	440
Maxima [A] (verification not implemented)	441
Giac [A] (verification not implemented)	441
Mupad [B] (verification not implemented)	441
Reduce [B] (verification not implemented)	442

Optimal result

Integrand size = 13, antiderivative size = 14

$$\int \frac{2^{\sqrt{x}}}{\sqrt{x}} dx = \frac{2^{1+\sqrt{x}}}{\log(2)}$$

output `2^(1+x^(1/2))/ln(2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{2^{\sqrt{x}}}{\sqrt{x}} dx = \frac{2^{1+\sqrt{x}}}{\log(2)}$$

input `Integrate[2^Sqrt[x]/Sqrt[x],x]`

output `2^(1 + Sqrt[x])/Log[2]`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {2638}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{2^{\sqrt{x}}}{\sqrt{x}} dx$$

↓ 2638

$$\frac{2^{\sqrt{x}+1}}{\log(2)}$$

input `Int [2^Sqrt [x]/Sqrt [x] ,x]`

output `2^(1 + Sqrt [x])/Log[2]`

Defintions of rubi rules used

rule 2638 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^(n_)))*((e_.) + (f_.)*(x_)^(m_.), x_Symbol] := Simp[(e + f*x)^n*(F^(a + b*(c + d*x)^n)/(b*f*n*(c + d*x)^n *Log[F])), x] /; FreeQ[{F, a, b, c, d, e, f, n}, x] && EqQ[m, n - 1] && EqQ[d*e - c*f, 0]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

method	result	size
derivativedivides	$\frac{2 \cdot 2^{\sqrt{x}}}{\ln(2)}$	12
default	$\frac{2 \cdot 2^{\sqrt{x}}}{\ln(2)}$	12
meijerg	$-\frac{2(1 - e^{\sqrt{x} \ln(2)})}{\ln(2)}$	18

input `int(2^(x^(1/2))/x^(1/2),x,method=_RETURNVERBOSE)`

output `2*2^(x^(1/2))/ln(2)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.79

$$\int \frac{2^{\sqrt{x}}}{\sqrt{x}} dx = \frac{2 \cdot 2^{(\sqrt{x})}}{\log(2)}$$

input `integrate(2^(x^(1/2))/x^(1/2),x, algorithm="fricas")`

output `2*2^sqrt(x)/log(2)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{2^{\sqrt{x}}}{\sqrt{x}} dx = \frac{2 \cdot 2^{\sqrt{x}}}{\log(2)}$$

input `integrate(2**(x**(1/2))/x**(1/2),x)`

output `2*2**(sqrt(x))/log(2)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{2^{\sqrt{x}}}{\sqrt{x}} dx = \frac{2^{\sqrt{x}+1}}{\log(2)}$$

input `integrate(2^(x^(1/2))/x^(1/2),x, algorithm="maxima")`output `2^(sqrt(x) + 1)/log(2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.79

$$\int \frac{2^{\sqrt{x}}}{\sqrt{x}} dx = \frac{2 \cdot 2^{(\sqrt{x})}}{\log(2)}$$

input `integrate(2^(x^(1/2))/x^(1/2),x, algorithm="giac")`output `2*2^sqrt(x)/log(2)`**Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.79

$$\int \frac{2^{\sqrt{x}}}{\sqrt{x}} dx = \frac{2 \cdot 2^{\sqrt{x}}}{\ln(2)}$$

input `int(2^(x^(1/2))/x^(1/2),x)`output `(2*2^(x^(1/2)))/log(2)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{2^{\sqrt{x}}}{\sqrt{x}} dx = \frac{2 \cdot 2^{\sqrt{x}}}{\log(2)}$$

input `int(2^(x^(1/2))/x^(1/2),x)`

output `(2*2**sqrt(x))/log(2)`

3.70 $\int e^{2\sin(x)} \cos(x) dx$

Optimal result	443
Mathematica [A] (verified)	443
Rubi [A] (verified)	444
Maple [A] (verified)	445
Fricas [A] (verification not implemented)	445
Sympy [A] (verification not implemented)	446
Maxima [A] (verification not implemented)	446
Giac [A] (verification not implemented)	446
Mupad [B] (verification not implemented)	447
Reduce [B] (verification not implemented)	447

Optimal result

Integrand size = 9, antiderivative size = 10

$$\int e^{2\sin(x)} \cos(x) dx = \frac{1}{2} e^{2\sin(x)}$$

output `1/2*exp(2*sin(x))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int e^{2\sin(x)} \cos(x) dx = \frac{1}{2} e^{2\sin(x)}$$

input `Integrate[E^(2*Sin[x])*Cos[x],x]`

output `E^(2*Sin[x])/2`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {4834, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{2\sin(x)} \cos(x) dx$$

↓ 4834

$$\int e^{2\sin(x)} d\sin(x)$$

↓ 2624

$$\frac{1}{2} e^{2\sin(x)}$$

input `Int[E^(2*Sin[x])*Cos[x],x]`

output `E^(2*Sin[x])/2`

Defintions of rubi rules used

rule 2624 `Int[((F_)^(v_))^(n_), x_Symbol] := Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;`
`FreeQ[{F, n}, x] && LinearQ[v, x]`

rule 4834 `Int[(u_)*(F_)[(c_)*((a_) + (b_)*(x_))], x_Symbol] := With[{d = FreeFacto`
`rs[Sin[c*(a + b*x)], x]}, Simp[d/(b*c) Subst[Int[SubstFor[1, Sin[c*(a + b`
`*x)]/d, u, x], x], x, Sin[c*(a + b*x)]/d, x] /; FunctionOfQ[Sin[c*(a + b*x`
`)]/d, u, x, True]] /; FreeQ[{a, b, c}, x] && (EqQ[F, Cos] || EqQ[F, cos])`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

method	result	size
derivativedivides	$\frac{e^{2\sin(x)}}{2}$	8
default	$\frac{e^{2\sin(x)}}{2}$	8
risch	$\frac{e^{2\sin(x)}}{2}$	8
parallelrisch	$\frac{e^{2\sin(x)}}{2}$	8
norman	$\frac{\tan\left(\frac{x}{2}\right)^2 e^{\frac{4 \tan\left(\frac{x}{2}\right)}{1+\tan\left(\frac{x}{2}\right)^2}}}{2} + \frac{e^{\frac{4 \tan\left(\frac{x}{2}\right)}{1+\tan\left(\frac{x}{2}\right)^2}}}{2}$	57

input `int(exp(2*sin(x))*cos(x),x,method=_RETURNVERBOSE)`output `1/2*exp(2*sin(x))`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int e^{2\sin(x)} \cos(x) dx = \frac{1}{2} e^{(2\sin(x))}$$

input `integrate(exp(2*sin(x))*cos(x),x, algorithm="fricas")`output `1/2*e^(2*sin(x))`

Sympy [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int e^{2\sin(x)} \cos(x) dx = \frac{e^{2\sin(x)}}{2}$$

input `integrate(exp(2*sin(x))*cos(x),x)`

output `exp(2*sin(x))/2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int e^{2\sin(x)} \cos(x) dx = \frac{1}{2} e^{(2\sin(x))}$$

input `integrate(exp(2*sin(x))*cos(x),x, algorithm="maxima")`

output `1/2*e^(2*sin(x))`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int e^{2\sin(x)} \cos(x) dx = \frac{1}{2} e^{(2\sin(x))}$$

input `integrate(exp(2*sin(x))*cos(x),x, algorithm="giac")`

output `1/2*e^(2*sin(x))`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int e^{2\sin(x)} \cos(x) dx = \frac{e^{2\sin(x)}}{2}$$

input `int(exp(2*sin(x))*cos(x),x)`

output `exp(2*sin(x))/2`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int e^{2\sin(x)} \cos(x) dx = \frac{e^{2\sin(x)}}{2}$$

input `int(exp(2*sin(x))*cos(x),x)`

output `e**(2*sin(x))/2`

3.71 $\int e^x \sin(x) dx$

Optimal result	448
Mathematica [A] (verified)	448
Rubi [A] (verified)	449
Maple [A] (verified)	449
Fricas [A] (verification not implemented)	450
Sympy [A] (verification not implemented)	450
Maxima [A] (verification not implemented)	451
Giac [A] (verification not implemented)	451
Mupad [B] (verification not implemented)	451
Reduce [B] (verification not implemented)	452

Optimal result

Integrand size = 6, antiderivative size = 19

$$\int e^x \sin(x) dx = -\frac{1}{2}e^x \cos(x) + \frac{1}{2}e^x \sin(x)$$

output `-1/2*exp(x)*cos(x)+1/2*exp(x)*sin(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int e^x \sin(x) dx = \frac{1}{2}e^x(-\cos(x) + \sin(x))$$

input `Integrate[E^x*Sin[x],x]`

output `(E^x*(-Cos[x] + Sin[x]))/2`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {4932}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^x \sin(x) dx$$

↓ 4932

$$\frac{1}{2}e^x \sin(x) - \frac{1}{2}e^x \cos(x)$$

input `Int [E^x*Sin [x] ,x]`

output `-1/2*(E^x*Cos [x]) + (E^x*Sin [x])/2`

Defintions of rubi rules used

rule 4932 `Int[(F_)^((c_.)*((a_.) + (b_.)*(x_)))*Sin[(d_.) + (e_.)*(x_)], x_Symbol] :>
Simp[b*c*Log[F]*F^(c*(a + b*x))*(Sin[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x
] - Simp[e*F^(c*(a + b*x))*(Cos[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] /; F
reeQ[{F, a, b, c, d, e}, x] && NeQ[e^2 + b^2*c^2*Log[F]^2, 0]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

method	result	size
parallelrisch	$\frac{e^x(-\cos(x)+\sin(x))}{2}$	12
default	$-\frac{e^x \cos(x)}{2} + \frac{e^x \sin(x)}{2}$	14
orering	$-\frac{e^x \cos(x)}{2} + \frac{e^x \sin(x)}{2}$	14
norman	$\frac{e^x \tan(\frac{x}{2}) + \frac{e^x \tan(\frac{x}{2})^2}{2} - \frac{e^x}{2}}{1 + \tan(\frac{x}{2})^2}$	34
risch	$-\frac{e^{(1+i)x}}{4} - \frac{ie^{(1+i)x}}{4} - \frac{e^{(1-i)x}}{4} + \frac{ie^{(1-i)x}}{4}$	36

input `int(exp(x)*sin(x),x,method=_RETURNVERBOSE)`

output `1/2*exp(x)*(-cos(x)+sin(x))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.68

$$\int e^x \sin(x) dx = -\frac{1}{2} \cos(x) e^x + \frac{1}{2} e^x \sin(x)$$

input `integrate(exp(x)*sin(x),x, algorithm="fricas")`

output `-1/2*cos(x)*e^x + 1/2*e^x*sin(x)`

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int e^x \sin(x) dx = \frac{e^x \sin(x)}{2} - \frac{e^x \cos(x)}{2}$$

input `integrate(exp(x)*sin(x),x)`

output `exp(x)*sin(x)/2 - exp(x)*cos(x)/2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int e^x \sin(x) dx = -\frac{1}{2} (\cos(x) - \sin(x))e^x$$

input `integrate(exp(x)*sin(x),x, algorithm="maxima")`

output `-1/2*(cos(x) - sin(x))*e^x`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int e^x \sin(x) dx = -\frac{1}{2} (\cos(x) - \sin(x))e^x$$

input `integrate(exp(x)*sin(x),x, algorithm="giac")`

output `-1/2*(cos(x) - sin(x))*e^x`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int e^x \sin(x) dx = -\frac{e^x (\cos(x) - \sin(x))}{2}$$

input `int(exp(x)*sin(x),x)`

output `-(exp(x)*(cos(x) - sin(x)))/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int e^x \sin(x) dx = \frac{e^x(-\cos(x) + \sin(x))}{2}$$

input `int(exp(x)*sin(x),x)`

output `(e**x*(- cos(x) + sin(x)))/2`

3.72 $\int e^x \cos(x) dx$

Optimal result	453
Mathematica [A] (verified)	453
Rubi [A] (verified)	454
Maple [A] (verified)	454
Fricas [A] (verification not implemented)	455
Sympy [A] (verification not implemented)	455
Maxima [A] (verification not implemented)	456
Giac [A] (verification not implemented)	456
Mupad [B] (verification not implemented)	456
Reduce [B] (verification not implemented)	457

Optimal result

Integrand size = 6, antiderivative size = 19

$$\int e^x \cos(x) dx = \frac{1}{2}e^x \cos(x) + \frac{1}{2}e^x \sin(x)$$

output `1/2*exp(x)*cos(x)+1/2*exp(x)*sin(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int e^x \cos(x) dx = \frac{1}{2}e^x(\cos(x) + \sin(x))$$

input `Integrate[E^x*Cos[x],x]`

output `(E^x*(Cos[x] + Sin[x]))/2`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {4933}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^x \cos(x) dx$$

$$\downarrow 4933$$

$$\frac{1}{2}e^x \sin(x) + \frac{1}{2}e^x \cos(x)$$

input `Int [E^x*Cos [x] , x]`

output `(E^x*Cos [x])/2 + (E^x*Sin [x])/2`

Defintions of rubi rules used

rule 4933 `Int [Cos [(d_.) + (e_.)*(x_)]*(F_)^((c_.)*((a_.) + (b_.)*(x_))), x_Symbol] :=
Simp [b*c*Log [F]*F^(c*(a + b*x))*(Cos [d + e*x]/(e^2 + b^2*c^2*Log [F]^2)), x
] + Simp [e*F^(c*(a + b*x))*(Sin [d + e*x]/(e^2 + b^2*c^2*Log [F]^2)), x] /; F
reeQ [{F, a, b, c, d, e}, x] && NeQ [e^2 + b^2*c^2*Log [F]^2, 0]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.53

method	result	size
parallelrisc	$\frac{e^x(\cos(x)+\sin(x))}{2}$	10
default	$\frac{e^x \cos(x)}{2} + \frac{e^x \sin(x)}{2}$	14
orering	$\frac{e^x \cos(x)}{2} + \frac{e^x \sin(x)}{2}$	14
norman	$\frac{e^x \tan(\frac{x}{2}) - \frac{e^x \tan(\frac{x}{2})^2}{2} + \frac{e^x}{2}}{1 + \tan(\frac{x}{2})^2}$	34
risc	$\frac{e^{(1+i)x}}{4} - \frac{ie^{(1+i)x}}{4} + \frac{e^{(1-i)x}}{4} + \frac{ie^{(1-i)x}}{4}$	36

input `int(exp(x)*cos(x),x,method=_RETURNVERBOSE)`

output `1/2*exp(x)*(cos(x)+sin(x))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.68

$$\int e^x \cos(x) dx = \frac{1}{2} \cos(x) e^x + \frac{1}{2} e^x \sin(x)$$

input `integrate(exp(x)*cos(x),x, algorithm="fricas")`

output `1/2*cos(x)*e^x + 1/2*e^x*sin(x)`

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int e^x \cos(x) dx = \frac{e^x \sin(x)}{2} + \frac{e^x \cos(x)}{2}$$

input `integrate(exp(x)*cos(x),x)`

output `exp(x)*sin(x)/2 + exp(x)*cos(x)/2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.47

$$\int e^x \cos(x) dx = \frac{1}{2} (\cos(x) + \sin(x))e^x$$

input `integrate(exp(x)*cos(x),x, algorithm="maxima")`

output `1/2*(cos(x) + sin(x))*e^x`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.47

$$\int e^x \cos(x) dx = \frac{1}{2} (\cos(x) + \sin(x))e^x$$

input `integrate(exp(x)*cos(x),x, algorithm="giac")`

output `1/2*(cos(x) + sin(x))*e^x`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.47

$$\int e^x \cos(x) dx = \frac{e^x (\cos(x) + \sin(x))}{2}$$

input `int(exp(x)*cos(x),x)`

output `(exp(x)*(cos(x) + sin(x)))/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.53

$$\int e^x \cos(x) dx = \frac{e^x(\cos(x) + \sin(x))}{2}$$

input `int(exp(x)*cos(x),x)`

output `(e**x*(cos(x) + sin(x)))/2`

3.73 $\int \frac{1}{1+e^x} dx$

Optimal result	458
Mathematica [A] (verified)	458
Rubi [A] (verified)	459
Maple [A] (verified)	460
Fricas [A] (verification not implemented)	461
Sympy [A] (verification not implemented)	461
Maxima [A] (verification not implemented)	461
Giac [A] (verification not implemented)	462
Mupad [B] (verification not implemented)	462
Reduce [B] (verification not implemented)	462

Optimal result

Integrand size = 7, antiderivative size = 10

$$\int \frac{1}{1+e^x} dx = x - \log(1+e^x)$$

output `x-ln(1+exp(x))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+e^x} dx = -2\operatorname{arctanh}(1+2e^x)$$

input `Integrate[(1 + E^x)^(-1), x]`

output `-2*ArcTanh[1 + 2*E^x]`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.30, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {2720, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{e^x + 1} dx \\
 & \quad \downarrow \text{2720} \\
 & \int \frac{e^{-x}}{e^x + 1} de^x \\
 & \quad \downarrow \text{47} \\
 & \int e^{-x} de^x - \int \frac{1}{1 + e^x} de^x \\
 & \quad \downarrow \text{14} \\
 & \log(e^x) - \int \frac{1}{1 + e^x} de^x \\
 & \quad \downarrow \text{16} \\
 & \log(e^x) - \log(e^x + 1)
 \end{aligned}$$

input `Int[(1 + E^x)^(-1), x]`

output `Log[E^x] - Log[1 + E^x]`

Definitions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

method	result	size
norman	$x - \ln(1 + e^x)$	10
risch	$x - \ln(1 + e^x)$	10
parallelrisch	$x - \ln(1 + e^x)$	10
derivativedivides	$-\ln(1 + e^x) + \ln(e^x)$	12
default	$-\ln(1 + e^x) + \ln(e^x)$	12

input `int(1/(1+exp(x)),x,method=_RETURNVERBOSE)`

output `x-ln(1+exp(x))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

$$\int \frac{1}{1+e^x} dx = x - \log(e^x + 1)$$

input `integrate(1/(1+exp(x)),x, algorithm="fricas")`

output `x - log(e^x + 1)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int \frac{1}{1+e^x} dx = x - \log(e^x + 1)$$

input `integrate(1/(1+exp(x)),x)`

output `x - log(exp(x) + 1)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

$$\int \frac{1}{1+e^x} dx = x - \log(e^x + 1)$$

input `integrate(1/(1+exp(x)),x, algorithm="maxima")`

output `x - log(e^x + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

$$\int \frac{1}{1+e^x} dx = x - \log(e^x + 1)$$

input `integrate(1/(1+exp(x)),x, algorithm="giac")`

output `x - log(e^x + 1)`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

$$\int \frac{1}{1+e^x} dx = x - \ln(e^x + 1)$$

input `int(1/(exp(x) + 1),x)`

output `x - log(exp(x) + 1)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+e^x} dx = -\log(e^x + 1) + x$$

input `int(1/(1+exp(x)),x)`

output `- log(e**x + 1) + x`

3.74 $\int e^x x dx$

Optimal result	463
Mathematica [A] (verified)	463
Rubi [A] (verified)	464
Maple [A] (verified)	465
Fricas [A] (verification not implemented)	465
Sympy [A] (verification not implemented)	466
Maxima [A] (verification not implemented)	466
Giac [A] (verification not implemented)	466
Mupad [B] (verification not implemented)	467
Reduce [B] (verification not implemented)	467

Optimal result

Integrand size = 5, antiderivative size = 11

$$\int e^x x dx = -e^x + e^x x$$

output `-exp(x)+exp(x)*x`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.64

$$\int e^x x dx = e^x(-1 + x)$$

input `Integrate[E^x*x,x]`

output `E^x*(-1 + x)`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int e^x x dx \\ \downarrow 2607 \\ e^x x - \int e^x dx \\ \downarrow 2624 \\ e^x x - e^x \end{array}$$

input `Int [Ex*x, x]`

output `-Ex + Ex*x`

Defintions of rubi rules used

rule 2607 `Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]`

rule 2624 `Int[((F_)^(v_))^(n_.), x_Symbol] := Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /; FreeQ[{F, n}, x] && LinearQ[v, x]`

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.64

method	result	size
gospers	$(-1 + x)e^x$	7
risch	$(-1 + x)e^x$	7
orering	$(-1 + x)e^x$	7
default	$-e^x + e^x x$	10
norman	$-e^x + e^x x$	10
parallelrisch	$-e^x + e^x x$	10
parts	$-e^x + e^x x$	10
meijerg	$1 - \frac{(-2x+2)e^x}{2}$	12

input `int(exp(x)*x,x,method=_RETURNVERBOSE)`

output `(-1+x)*exp(x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.55

$$\int e^x x dx = (x - 1)e^x$$

input `integrate(exp(x)*x,x, algorithm="fricas")`

output `(x - 1)*e^x`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.45

$$\int e^x x dx = (x - 1) e^x$$

input `integrate(exp(x)*x,x)`

output `(x - 1)*exp(x)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.55

$$\int e^x x dx = (x - 1) e^x$$

input `integrate(exp(x)*x,x, algorithm="maxima")`

output `(x - 1)*e^x`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.55

$$\int e^x x dx = (x - 1) e^x$$

input `integrate(exp(x)*x,x, algorithm="giac")`

output `(x - 1)*e^x`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.55

$$\int e^x x dx = e^x (x - 1)$$

input `int(x*exp(x),x)`

output `exp(x)*(x - 1)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.64

$$\int e^x x dx = e^x (x - 1)$$

input `int(exp(x)*x,x)`

output `e**x*(x - 1)`

3.75 $\int e^{-x} x dx$

Optimal result	468
Mathematica [A] (verified)	468
Rubi [A] (verified)	469
Maple [A] (warning: unable to verify)	470
Fricas [A] (verification not implemented)	470
Sympy [A] (verification not implemented)	471
Maxima [A] (verification not implemented)	471
Giac [A] (verification not implemented)	471
Mupad [B] (verification not implemented)	472
Reduce [B] (verification not implemented)	472

Optimal result

Integrand size = 7, antiderivative size = 16

$$\int e^{-x} x dx = -e^{-x} - e^{-x} x$$

output `-1/exp(x)-x/exp(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.69

$$\int e^{-x} x dx = e^{-x}(-1 - x)$$

input `Integrate[x/E^x,x]`

output `(-1 - x)/E^x`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int e^{-x} x dx \\ \downarrow 2607 \\ \int e^{-x} dx - e^{-x} x \\ \downarrow 2624 \\ -e^{-x} x - e^{-x} \end{array}$$

input `Int [x/Ex, x]`

output `-E(-x) - x/Ex`

Defintions of rubi rules used

rule 2607 `Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]`

rule 2624 `Int[((F_)^(v_))^(n_.), x_Symbol] := Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /; FreeQ[{F, n}, x] && LinearQ[v, x]`

Maple [A] (warning: unable to verify)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.62

method	result	size
gospers	$-(1+x)e^{-x}$	10
orering	$-(1+x)e^{-x}$	10
norman	$(-1-x)e^{-x}$	11
risch	$(-1-x)e^{-x}$	11
parallelrisch	$(-1-x)e^{-x}$	11
meijerg	$1 - \frac{(2x+2)e^{-x}}{2}$	14
default	$-e^{-x} - xe^{-x}$	15

input `int(x/exp(x),x,method=_RETURNVERBOSE)`output `-(1+x)/exp(x)`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.56

$$\int e^{-x}x dx = -(x+1)e^{(-x)}$$

input `integrate(x/exp(x),x, algorithm="fricas")`output `-(x + 1)*e^(-x)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.44

$$\int e^{-x} x dx = (-x - 1) e^{-x}$$

input `integrate(x/exp(x), x)`

output `(-x - 1)*exp(-x)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.56

$$\int e^{-x} x dx = -(x + 1) e^{(-x)}$$

input `integrate(x/exp(x), x, algorithm="maxima")`

output `-(x + 1)*e^(-x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.56

$$\int e^{-x} x dx = -(x + 1) e^{(-x)}$$

input `integrate(x/exp(x), x, algorithm="giac")`

output `-(x + 1)*e^(-x)`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.56

$$\int e^{-x} x dx = -e^{-x} (x + 1)$$

input `int(x*exp(-x),x)`

output `-exp(-x)*(x + 1)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.69

$$\int e^{-x} x dx = \frac{-x - 1}{e^x}$$

input `int(x/exp(x),x)`

output `(- (x + 1))/e**x`

3.76 $\int e^x x^2 dx$

Optimal result	473
Mathematica [A] (verified)	473
Rubi [A] (verified)	474
Maple [A] (verified)	475
Fricas [A] (verification not implemented)	475
Sympy [A] (verification not implemented)	476
Maxima [A] (verification not implemented)	476
Giac [A] (verification not implemented)	476
Mupad [B] (verification not implemented)	477
Reduce [B] (verification not implemented)	477

Optimal result

Integrand size = 7, antiderivative size = 19

$$\int e^x x^2 dx = 2e^x - 2e^x x + e^x x^2$$

output `2*exp(x)-2*exp(x)*x+exp(x)*x^2`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int e^x x^2 dx = e^x (2 - 2x + x^2)$$

input `Integrate[E^x*x^2,x]`

output `E^x*(2 - 2*x + x^2)`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {2607, 2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int e^x x^2 dx \\ & \quad \downarrow 2607 \\ & e^x x^2 - 2 \int e^x x dx \\ & \quad \downarrow 2607 \\ & e^x x^2 - 2 \left(e^x x - \int e^x dx \right) \\ & \quad \downarrow 2624 \\ & e^x x^2 - 2(e^x x - e^x) \end{aligned}$$

input `Int [E^x*x^2,x]`

output `E^x*x^2 - 2*(-E^x + E^x*x)`

Defintions of rubi rules used

rule 2607 `Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]`

rule 2624 `Int[((F_)^(v_))^(n_.), x_Symbol] := Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /; FreeQ[{F, n}, x] && LinearQ[v, x]`

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

method	result	size
gospers	$(x^2 - 2x + 2)e^x$	12
risch	$(x^2 - 2x + 2)e^x$	12
orering	$(x^2 - 2x + 2)e^x$	12
default	$2e^x - 2e^x x + e^x x^2$	17
norman	$2e^x - 2e^x x + e^x x^2$	17
meijerg	$-2 + \frac{(3x^2 - 6x + 6)e^x}{3}$	17
parallelrisch	$2e^x - 2e^x x + e^x x^2$	17
parts	$2e^x - 2e^x x + e^x x^2$	17

input `int(exp(x)*x^2,x,method=_RETURNVERBOSE)`output `(x^2-2*x+2)*exp(x)`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int e^x x^2 dx = (x^2 - 2x + 2)e^x$$

input `integrate(exp(x)*x^2,x, algorithm="fricas")`output `(x^2 - 2*x + 2)*e^x`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.53

$$\int e^x x^2 dx = (x^2 - 2x + 2) e^x$$

input `integrate(exp(x)*x**2,x)`

output `(x**2 - 2*x + 2)*exp(x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int e^x x^2 dx = (x^2 - 2x + 2) e^x$$

input `integrate(exp(x)*x^2,x, algorithm="maxima")`

output `(x^2 - 2*x + 2)*e^x`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int e^x x^2 dx = (x^2 - 2x + 2) e^x$$

input `integrate(exp(x)*x^2,x, algorithm="giac")`

output `(x^2 - 2*x + 2)*e^x`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int e^x x^2 dx = e^x (x^2 - 2x + 2)$$

input `int(x^2*exp(x),x)`

output `exp(x)*(x^2 - 2*x + 2)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int e^x x^2 dx = e^x (x^2 - 2x + 2)$$

input `int(exp(x)*x^2,x)`

output `e**x*(x**2 - 2*x + 2)`

3.77 $\int e^{-2x} x^2 dx$

Optimal result	478
Mathematica [A] (verified)	478
Rubi [A] (verified)	479
Maple [A] (verified)	480
Fricas [A] (verification not implemented)	480
Sympy [A] (verification not implemented)	481
Maxima [A] (verification not implemented)	481
Giac [A] (verification not implemented)	481
Mupad [B] (verification not implemented)	482
Reduce [B] (verification not implemented)	482

Optimal result

Integrand size = 9, antiderivative size = 32

$$\int e^{-2x} x^2 dx = -\frac{1}{4}e^{-2x} - \frac{1}{2}e^{-2x}x - \frac{1}{2}e^{-2x}x^2$$

output `-1/4/exp(2*x)-1/2*x/exp(2*x)-1/2*x^2/exp(2*x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.59

$$\int e^{-2x} x^2 dx = -\frac{1}{4}e^{-2x}(1 + 2x + 2x^2)$$

input `Integrate[x^2/E^(2*x),x]`

output `-1/4*(1 + 2*x + 2*x^2)/E^(2*x)`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2607, 2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int e^{-2x} x^2 dx \\ & \quad \downarrow \text{2607} \\ & \int e^{-2x} x dx - \frac{1}{2} e^{-2x} x^2 \\ & \quad \downarrow \text{2607} \\ & \frac{1}{2} \int e^{-2x} dx - \frac{1}{2} e^{-2x} x^2 - \frac{1}{2} e^{-2x} x \\ & \quad \downarrow \text{2624} \\ & -\frac{1}{2} e^{-2x} x^2 - \frac{1}{2} e^{-2x} x - \frac{e^{-2x}}{4} \end{aligned}$$

input `Int[x^2/E^(2*x),x]`

output `-1/4*1/E^(2*x) - x/(2*E^(2*x)) - x^2/(2*E^(2*x))`

Defintions of rubi rules used

rule 2607

```
Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x))))^(n_.)*((c_.) + (d_.)*(x))^(m
_.), x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])),
x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^
n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*
m] && !TrueQ[$UseGamma]
```


rule 2624 `Int[((F_)^(v_))^(n_), x_Symbol] := Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;`
`FreeQ[{F, n}, x] && LinearQ[v, x]`

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.50

method	result	size
risch	$(-\frac{1}{2}x^2 - \frac{1}{2}x - \frac{1}{4})e^{-2x}$	16
norman	$(-\frac{1}{2}x^2 - \frac{1}{2}x - \frac{1}{4})e^{-2x}$	18
gosper	$-\frac{(2x^2+2x+1)e^{-2x}}{4}$	19
meijerg	$\frac{1}{4} - \frac{(12x^2+12x+6)e^{-2x}}{24}$	19
parallelrisch	$\frac{(-2x^2-2x-1)e^{-2x}}{4}$	19
orering	$-\frac{(2x^2+2x+1)e^{-2x}}{4}$	19
derivativedivides	$-\frac{e^{-2x}}{4} - \frac{x e^{-2x}}{2} - \frac{x^2 e^{-2x}}{2}$	30
default	$-\frac{e^{-2x}}{4} - \frac{x e^{-2x}}{2} - \frac{x^2 e^{-2x}}{2}$	30

input `int(x^2/exp(2*x),x,method=_RETURNVERBOSE)`

output `(-1/2*x^2-1/2*x-1/4)*exp(-2*x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.50

$$\int e^{-2x} x^2 dx = -\frac{1}{4} (2x^2 + 2x + 1)e^{(-2x)}$$

input `integrate(x^2/exp(2*x),x, algorithm="fricas")`

output `-1/4*(2*x^2 + 2*x + 1)*e^(-2*x)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.53

$$\int e^{-2x} x^2 dx = \frac{(-2x^2 - 2x - 1) e^{-2x}}{4}$$

input `integrate(x**2/exp(2*x),x)`output `(-2*x**2 - 2*x - 1)*exp(-2*x)/4`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.50

$$\int e^{-2x} x^2 dx = -\frac{1}{4} (2x^2 + 2x + 1) e^{(-2x)}$$

input `integrate(x^2/exp(2*x),x, algorithm="maxima")`output `-1/4*(2*x^2 + 2*x + 1)*e^(-2*x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.50

$$\int e^{-2x} x^2 dx = -\frac{1}{4} (2x^2 + 2x + 1) e^{(-2x)}$$

input `integrate(x^2/exp(2*x),x, algorithm="giac")`output `-1/4*(2*x^2 + 2*x + 1)*e^(-2*x)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.50

$$\int e^{-2x} x^2 dx = -\frac{e^{-2x} (4x^2 + 4x + 2)}{8}$$

input `int(x^2*exp(-2*x),x)`

output `-(exp(-2*x)*(4*x + 4*x^2 + 2))/8`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.59

$$\int e^{-2x} x^2 dx = \frac{-2x^2 - 2x - 1}{4e^{2x}}$$

input `int(x^2/exp(2*x),x)`

output `(- 2*x**2 - 2*x - 1)/(4*e**(2*x))`

3.78 $\int e^{\sqrt{x}} dx$

Optimal result	483
Mathematica [A] (verified)	483
Rubi [A] (verified)	484
Maple [A] (verified)	485
Fricas [A] (verification not implemented)	485
Sympy [A] (verification not implemented)	486
Maxima [A] (verification not implemented)	486
Giac [A] (verification not implemented)	486
Mupad [B] (verification not implemented)	487
Reduce [B] (verification not implemented)	487

Optimal result

Integrand size = 7, antiderivative size = 24

$$\int e^{\sqrt{x}} dx = -2e^{\sqrt{x}} + 2e^{\sqrt{x}}\sqrt{x}$$

output `-2*exp(x^(1/2))+2*exp(x^(1/2))*x^(1/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int e^{\sqrt{x}} dx = 2e^{\sqrt{x}}(-1 + \sqrt{x})$$

input `Integrate[E^Sqrt[x], x]`

output `2*E^Sqrt[x]*(-1 + Sqrt[x])`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.04, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {2636, 2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int e^{\sqrt{x}} dx \\ & \quad \downarrow \text{2636} \\ & 2 \int e^{\sqrt{x}} \sqrt{x} d\sqrt{x} \\ & \quad \downarrow \text{2607} \\ & 2 \left(e^{\sqrt{x}} \sqrt{x} - \int e^{\sqrt{x}} d\sqrt{x} \right) \\ & \quad \downarrow \text{2624} \\ & 2 \left(e^{\sqrt{x}} \sqrt{x} - e^{\sqrt{x}} \right) \end{aligned}$$

input `Int[E^Sqrt[x],x]`

output `2*(-E^Sqrt[x] + E^Sqrt[x]*Sqrt[x])`

Defintions of rubi rules used

rule 2607

```
Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m
_.), x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])),
x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^
n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*
m] && !TrueQ[$UseGamma]
```

rule 2624 `Int[((F_)^(v_))^(n_), x_Symbol] := Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;`
`FreeQ[{F, n}, x] && LinearQ[v, x]`

rule 2636 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^(n_))), x_Symbol] := With[{k =`
`Denominator[n]}, Simp[k/d Subst[Int[x^(k - 1)*F^(a + b*x^(k*n)), x], x, (`
`c + d*x)^(1/k)], x]] /;` `FreeQ[{F, a, b, c, d}, x] && IntegerQ[2/n] && !IntegerQ[n]`

Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

method	result	size
meijerg	$2 - (-2\sqrt{x} + 2) e^{\sqrt{x}}$	16
derivativedivides	$-2 e^{\sqrt{x}} + 2 e^{\sqrt{x}} \sqrt{x}$	17
default	$-2 e^{\sqrt{x}} + 2 e^{\sqrt{x}} \sqrt{x}$	17

input `int(exp(x^(1/2)),x,method=_RETURNVERBOSE)`

output `2-(-2*x^(1/2)+2)*exp(x^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.46

$$\int e^{\sqrt{x}} dx = 2(\sqrt{x} - 1)e^{\sqrt{x}}$$

input `integrate(exp(x^(1/2)),x, algorithm="fricas")`

output `2*(sqrt(x) - 1)*e^sqrt(x)`

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int e^{\sqrt{x}} dx = 2\sqrt{x}e^{\sqrt{x}} - 2e^{\sqrt{x}}$$

input `integrate(exp(x**(1/2)),x)`

output `2*sqrt(x)*exp(sqrt(x)) - 2*exp(sqrt(x))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.46

$$\int e^{\sqrt{x}} dx = 2(\sqrt{x} - 1)e^{\sqrt{x}}$$

input `integrate(exp(x^(1/2)),x, algorithm="maxima")`

output `2*(sqrt(x) - 1)*e^sqrt(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.46

$$\int e^{\sqrt{x}} dx = 2(\sqrt{x} - 1)e^{\sqrt{x}}$$

input `integrate(exp(x^(1/2)),x, algorithm="giac")`

output `2*(sqrt(x) - 1)*e^sqrt(x)`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.46

$$\int e^{\sqrt{x}} dx = 2e^{\sqrt{x}}(\sqrt{x} - 1)$$

input `int(exp(x^(1/2)),x)`

output `2*exp(x^(1/2))*(x^(1/2) - 1)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.42

$$\int e^{\sqrt{x}} dx = 2e^{\sqrt{x}}(\sqrt{x} - 1)$$

input `int(exp(x^(1/2)),x)`

output `2*e**sqrt(x)*(sqrt(x) - 1)`

3.79 $\int e^{-x^2} x^3 dx$

Optimal result	488
Mathematica [A] (verified)	488
Rubi [A] (verified)	489
Maple [A] (warning: unable to verify)	490
Fricas [A] (verification not implemented)	490
Sympy [A] (verification not implemented)	491
Maxima [A] (verification not implemented)	491
Giac [A] (verification not implemented)	491
Mupad [B] (verification not implemented)	492
Reduce [B] (verification not implemented)	492

Optimal result

Integrand size = 11, antiderivative size = 26

$$\int e^{-x^2} x^3 dx = -\frac{e^{-x^2}}{2} - \frac{1}{2}e^{-x^2} x^2$$

output `-1/2/exp(x^2)-1/2*x^2/exp(x^2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.62

$$\int e^{-x^2} x^3 dx = -\frac{1}{2}e^{-x^2}(1 + x^2)$$

input `Integrate[x^3/E^x^2,x]`

output `-1/2*(1 + x^2)/E^x^2`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2641, 2638}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{-x^2} x^3 dx$$

$$\downarrow 2641$$

$$\int e^{-x^2} x dx - \frac{1}{2} e^{-x^2} x^2$$

$$\downarrow 2638$$

$$-\frac{1}{2} e^{-x^2} x^2 - \frac{e^{-x^2}}{2}$$

input `Int [x^3/E^x^2, x]`

output `-1/2*1/E^x^2 - x^2/(2*E^x^2)`

Defintions of rubi rules used

rule 2638 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^n*(F^(a + b*(c + d*x)^n)/(b*f*n*(c + d*x)^n *Log[F])), x] /; FreeQ[{F, a, b, c, d, e, f, n}, x] && EqQ[m, n - 1] && EqQ[d*e - c*f, 0]`

rule 2641 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^(m - n + 1)*(F^(a + b*(c + d*x)^n)/(b*d*n*Log[F])), x] - Simp[(m - n + 1)/(b*n*Log[F]) Int[(c + d*x)^(m - n)*F^(a + b*(c + d*x)^n), x], x] /; FreeQ[{F, a, b, c, d}, x] && IntegerQ[2*((m + 1)/n)] && LtQ[0, (m + 1)/n, 5] && IntegerQ[n] && (LtQ[0, n, m + 1] || LtQ[m, n, 0])`

Maple [A] (warning: unable to verify)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.54

method	result	size
gospers	$-\frac{(x^2+1)e^{-x^2}}{2}$	14
orering	$-\frac{(x^2+1)e^{-x^2}}{2}$	14
norman	$\left(-\frac{x^2}{2} - \frac{1}{2}\right) e^{-x^2}$	15
risch	$\left(-\frac{x^2}{2} - \frac{1}{2}\right) e^{-x^2}$	15
parallelrisch	$\frac{(-x^2-1)e^{-x^2}}{2}$	16
meijerg	$\frac{1}{2} - \frac{(2x^2+2)e^{-x^2}}{4}$	18
derivativedivides	$-\frac{e^{-x^2}}{2} - \frac{x^2 e^{-x^2}}{2}$	21
default	$-\frac{e^{-x^2}}{2} - \frac{x^2 e^{-x^2}}{2}$	21

input `int(x^3/exp(x^2),x,method=_RETURNVERBOSE)`output `-1/2*(x^2+1)/exp(x^2)`**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.50

$$\int e^{-x^2} x^3 dx = -\frac{1}{2} (x^2 + 1) e^{(-x^2)}$$

input `integrate(x^3/exp(x^2),x, algorithm="fricas")`output `-1/2*(x^2 + 1)*e^(-x^2)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.46

$$\int e^{-x^2} x^3 dx = \frac{(-x^2 - 1) e^{-x^2}}{2}$$

input `integrate(x**3/exp(x**2),x)`

output `(-x**2 - 1)*exp(-x**2)/2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.50

$$\int e^{-x^2} x^3 dx = -\frac{1}{2} (x^2 + 1) e^{(-x^2)}$$

input `integrate(x^3/exp(x^2),x, algorithm="maxima")`

output `-1/2*(x^2 + 1)*e^(-x^2)`

Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.50

$$\int e^{-x^2} x^3 dx = -\frac{1}{2} (x^2 + 1) e^{(-x^2)}$$

input `integrate(x^3/exp(x^2),x, algorithm="giac")`

output `-1/2*(x^2 + 1)*e^(-x^2)`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.50

$$\int e^{-x^2} x^3 dx = -\frac{e^{-x^2} (x^2 + 1)}{2}$$

input `int(x^3*exp(-x^2),x)`output `-(exp(-x^2)*(x^2 + 1))/2`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.62

$$\int e^{-x^2} x^3 dx = \frac{-x^2 - 1}{2e^{x^2}}$$

input `int(x^3/exp(x^2),x)`output `(- (x**2 + 1))/(2*e**(x**2))`

3.80 $\int e^{ax} \cos(bx) dx$

Optimal result	493
Mathematica [A] (verified)	493
Rubi [A] (verified)	494
Maple [A] (verified)	494
Fricas [A] (verification not implemented)	495
Sympy [C] (verification not implemented)	495
Maxima [A] (verification not implemented)	496
Giac [A] (verification not implemented)	497
Mupad [B] (verification not implemented)	497
Reduce [B] (verification not implemented)	497

Optimal result

Integrand size = 10, antiderivative size = 41

$$\int e^{ax} \cos(bx) dx = \frac{ae^{ax} \cos(bx)}{a^2 + b^2} + \frac{be^{ax} \sin(bx)}{a^2 + b^2}$$

output `a*exp(a*x)*cos(b*x)/(a^2+b^2)+b*exp(a*x)*sin(b*x)/(a^2+b^2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.68

$$\int e^{ax} \cos(bx) dx = \frac{e^{ax}(a \cos(bx) + b \sin(bx))}{a^2 + b^2}$$

input `Integrate[E^(a*x)*Cos[b*x],x]`

output `(E^(a*x)*(a*Cos[b*x] + b*Sin[b*x]))/(a^2 + b^2)`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {4933}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{ax} \cos(bx) dx$$

$$\downarrow 4933$$

$$\frac{be^{ax} \sin(bx)}{a^2 + b^2} + \frac{ae^{ax} \cos(bx)}{a^2 + b^2}$$

input `Int [E^(a*x)*Cos [b*x] , x]`

output `(a*E^(a*x)*Cos [b*x])/(a^2 + b^2) + (b*E^(a*x)*Sin [b*x])/(a^2 + b^2)`

Defintions of rubi rules used

rule 4933

```
Int[Cos[(d_.) + (e_.)*(x_)]*(F_)^((c_.)*((a_.) + (b_.)*(x_))), x_Symbol] :>
Simp[b*c*Log[F]*F^(c*(a + b*x))*(Cos[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x
] + Simp[e*F^(c*(a + b*x))*(Sin[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] /; F
reeQ[{F, a, b, c, d, e}, x] && NeQ[e^2 + b^2*c^2*Log[F]^2, 0]
```

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.68

method	result	size
parallelrisch	$\frac{(\cos(bx)a+b\sin(bx))e^{ax}}{a^2+b^2}$	28
default	$\frac{a e^{ax} \cos(bx)}{a^2+b^2} + \frac{b e^{ax} \sin(bx)}{a^2+b^2}$	40
risch	$\frac{e^{x(ib+a)}}{2ib+2a} + \frac{e^{x(-ib+a)}}{-2ib+2a}$	40
orering	$\frac{2a e^{ax} \cos(bx)}{a^2+b^2} - \frac{a e^{ax} \cos(bx) - e^{ax} b \sin(bx)}{a^2+b^2}$	55
norman	$\frac{\frac{a e^{ax}}{a^2+b^2} - \frac{a e^{ax} \tan\left(\frac{bx}{2}\right)^2}{a^2+b^2} + \frac{2b e^{ax} \tan\left(\frac{bx}{2}\right)}{a^2+b^2}}{1 + \tan\left(\frac{bx}{2}\right)^2}$	73

input `int(exp(a*x)*cos(b*x),x,method=_RETURNVERBOSE)`

output `1/(a^2+b^2)*(cos(b*x)*a+b*sin(b*x))*exp(a*x)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.76

$$\int e^{ax} \cos(bx) dx = \frac{a \cos(bx) e^{(ax)} + b e^{(ax)} \sin(bx)}{a^2 + b^2}$$

input `integrate(exp(a*x)*cos(b*x),x, algorithm="fricas")`

output `(a*cos(b*x)*e^(a*x) + b*e^(a*x)*sin(b*x))/(a^2 + b^2)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 136, normalized size of antiderivative = 3.32

$$\int e^{ax} \cos(bx) dx = \begin{cases} x & \text{for } a = 0 \wedge b = 0 \\ \frac{ixe^{-ibx} \sin(bx)}{2} + \frac{xe^{-ibx} \cos(bx)}{2} + \frac{e^{-ibx} \sin(bx)}{2b} & \text{for } a = -ib \\ -\frac{ixe^{ibx} \sin(bx)}{2} + \frac{xe^{ibx} \cos(bx)}{2} + \frac{e^{ibx} \sin(bx)}{2b} & \text{for } a = ib \\ \frac{ae^{ax} \cos(bx)}{a^2+b^2} + \frac{be^{ax} \sin(bx)}{a^2+b^2} & \text{otherwise} \end{cases}$$

input `integrate(exp(a*x)*cos(b*x),x)`

output `Piecewise((x, Eq(a, 0) & Eq(b, 0)), (I*x*exp(-I*b*x)*sin(b*x)/2 + x*exp(-I*b*x)*cos(b*x)/2 + exp(-I*b*x)*sin(b*x)/(2*b), Eq(a, -I*b)), (-I*x*exp(I*b*x)*sin(b*x)/2 + x*exp(I*b*x)*cos(b*x)/2 + exp(I*b*x)*sin(b*x)/(2*b), Eq(a, I*b)), (a*exp(a*x)*cos(b*x)/(a**2 + b**2) + b*exp(a*x)*sin(b*x)/(a**2 + b**2), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.66

$$\int e^{ax} \cos(bx) dx = \frac{(a \cos(bx) + b \sin(bx))e^{(ax)}}{a^2 + b^2}$$

input `integrate(exp(a*x)*cos(b*x),x, algorithm="maxima")`

output `(a*cos(b*x) + b*sin(b*x))*e^(a*x)/(a^2 + b^2)`

Giac [A] (verification not implemented)

Time = 1.25 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.88

$$\int e^{ax} \cos(bx) dx = \left(\frac{a \cos(bx)}{a^2 + b^2} + \frac{b \sin(bx)}{a^2 + b^2} \right) e^{(ax)}$$

input `integrate(exp(a*x)*cos(b*x),x, algorithm="giac")`output `(a*cos(b*x)/(a^2 + b^2) + b*sin(b*x)/(a^2 + b^2))*e^(a*x)`**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.66

$$\int e^{ax} \cos(bx) dx = \frac{e^{ax} (a \cos(bx) + b \sin(bx))}{a^2 + b^2}$$

input `int(exp(a*x)*cos(b*x),x)`output `(exp(a*x)*(a*cos(b*x) + b*sin(b*x)))/(a^2 + b^2)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.68

$$\int e^{ax} \cos(bx) dx = \frac{e^{ax} (\cos(bx) a + \sin(bx) b)}{a^2 + b^2}$$

input `int(exp(a*x)*cos(b*x),x)`output `(e**(a*x)*(cos(b*x)*a + sin(b*x)*b))/(a**2 + b**2)`

3.81 $\int e^{ax} \sin(bx) dx$

Optimal result	498
Mathematica [A] (verified)	498
Rubi [A] (verified)	499
Maple [A] (verified)	499
Fricas [A] (verification not implemented)	500
Sympy [C] (verification not implemented)	500
Maxima [A] (verification not implemented)	501
Giac [A] (verification not implemented)	502
Mupad [B] (verification not implemented)	502
Reduce [B] (verification not implemented)	502

Optimal result

Integrand size = 10, antiderivative size = 42

$$\int e^{ax} \sin(bx) dx = -\frac{be^{ax} \cos(bx)}{a^2 + b^2} + \frac{ae^{ax} \sin(bx)}{a^2 + b^2}$$

output `-b*exp(a*x)*cos(b*x)/(a^2+b^2)+a*exp(a*x)*sin(b*x)/(a^2+b^2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.69

$$\int e^{ax} \sin(bx) dx = \frac{e^{ax}(-b \cos(bx) + a \sin(bx))}{a^2 + b^2}$$

input `Integrate[E^(a*x)*Sin[b*x],x]`

output `(E^(a*x)*(-b*Cos[b*x] + a*Sine[b*x]))/(a^2 + b^2)`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {4932}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{ax} \sin(bx) dx$$

$$\downarrow 4932$$

$$\frac{ae^{ax} \sin(bx)}{a^2 + b^2} - \frac{be^{ax} \cos(bx)}{a^2 + b^2}$$

input `Int [E^(a*x)*Sin[b*x] , x]`

output `-((b*E^(a*x)*Cos[b*x])/(a^2 + b^2)) + (a*E^(a*x)*Sin[b*x])/(a^2 + b^2)`

Defintions of rubi rules used

rule 4932

```
Int[(F_)^((c_.)*((a_.) + (b_.)*(x_)))*Sin[(d_.) + (e_.)*(x_)], x_Symbol] :>
Simp[b*c*Log[F]*F^(c*(a + b*x))*(Sin[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x
] - Simp[e*F^(c*(a + b*x))*(Cos[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] /; F
reeQ[{F, a, b, c, d, e}, x] && NeQ[e^2 + b^2*c^2*Log[F]^2, 0]
```

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.69

method	result	size
parallelrisch	$\frac{(\sin(bx)a - b \cos(bx))e^{ax}}{a^2 + b^2}$	29
default	$-\frac{b e^{ax} \cos(bx)}{a^2 + b^2} + \frac{a e^{ax} \sin(bx)}{a^2 + b^2}$	41
risch	$-\frac{i e^{x(ib+a)}}{2(ib+a)} + \frac{i e^{x(-ib+a)}}{-2ib+2a}$	42
orering	$\frac{2a e^{ax} \sin(bx)}{a^2 + b^2} - \frac{a e^{ax} \sin(bx) + e^{ax} b \cos(bx)}{a^2 + b^2}$	54
norman	$\frac{\frac{b e^{ax} \tan\left(\frac{bx}{2}\right)^2}{a^2 + b^2} - \frac{b e^{ax}}{a^2 + b^2} + \frac{2a e^{ax} \tan\left(\frac{bx}{2}\right)}{a^2 + b^2}}{1 + \tan\left(\frac{bx}{2}\right)^2}$	73

input `int(exp(a*x)*sin(b*x), x, method=_RETURNVERBOSE)`

output `1/(a^2+b^2)*(sin(b*x)*a-b*cos(b*x))*exp(a*x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.79

$$\int e^{ax} \sin(bx) dx = -\frac{b \cos(bx) e^{(ax)} - a e^{(ax)} \sin(bx)}{a^2 + b^2}$$

input `integrate(exp(a*x)*sin(b*x), x, algorithm="fricas")`

output `-(b*cos(b*x)*e^(a*x) - a*e^(a*x)*sin(b*x))/(a^2 + b^2)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 139, normalized size of antiderivative = 3.31

$$\int e^{ax} \sin(bx) dx = \begin{cases} 0 & \text{for } a = 0 \wedge b = 0 \\ \frac{xe^{-ibx} \sin(bx)}{2} - \frac{ixe^{-ibx} \cos(bx)}{2} + \frac{ie^{-ibx} \sin(bx)}{2b} & \text{for } a = -ib \\ \frac{xe^{ibx} \sin(bx)}{2} + \frac{ixe^{ibx} \cos(bx)}{2} - \frac{ie^{ibx} \sin(bx)}{2b} & \text{for } a = ib \\ \frac{ae^{ax} \sin(bx)}{a^2+b^2} - \frac{be^{ax} \cos(bx)}{a^2+b^2} & \text{otherwise} \end{cases}$$

input `integrate(exp(a*x)*sin(b*x),x)`

output `Piecewise((0, Eq(a, 0) & Eq(b, 0)), (x*exp(-I*b*x)*sin(b*x)/2 - I*x*exp(-I*b*x)*cos(b*x)/2 + I*exp(-I*b*x)*sin(b*x)/(2*b), Eq(a, -I*b)), (x*exp(I*b*x)*sin(b*x)/2 + I*x*exp(I*b*x)*cos(b*x)/2 - I*exp(I*b*x)*sin(b*x)/(2*b), Eq(a, I*b)), (a*exp(a*x)*sin(b*x)/(a**2 + b**2) - b*exp(a*x)*cos(b*x)/(a**2 + b**2), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.69

$$\int e^{ax} \sin(bx) dx = -\frac{(b \cos(bx) - a \sin(bx))e^{(ax)}}{a^2 + b^2}$$

input `integrate(exp(a*x)*sin(b*x),x, algorithm="maxima")`

output `-(b*cos(b*x) - a*sin(b*x))*e^(a*x)/(a^2 + b^2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.90

$$\int e^{ax} \sin(bx) dx = -\left(\frac{b \cos(bx)}{a^2 + b^2} - \frac{a \sin(bx)}{a^2 + b^2}\right) e^{(ax)}$$

input `integrate(exp(a*x)*sin(b*x),x, algorithm="giac")`output `-(b*cos(b*x)/(a^2 + b^2) - a*sin(b*x)/(a^2 + b^2))*e^(a*x)`**Mupad [B] (verification not implemented)**

Time = 0.01 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.69

$$\int e^{ax} \sin(bx) dx = -\frac{e^{ax} (b \cos(bx) - a \sin(bx))}{a^2 + b^2}$$

input `int(exp(a*x)*sin(b*x),x)`output `-(exp(a*x)*(b*cos(b*x) - a*sin(b*x)))/(a^2 + b^2)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.69

$$\int e^{ax} \sin(bx) dx = \frac{e^{ax} (-\cos(bx) b + \sin(bx) a)}{a^2 + b^2}$$

input `int(exp(a*x)*sin(b*x),x)`output `(e**(a*x)*(-cos(b*x)*b + sin(b*x)*a))/(a**2 + b**2)`

3.82 $\int \cot^{-1}(x) dx$

Optimal result	503
Mathematica [A] (verified)	503
Rubi [A] (verified)	504
Maple [A] (verified)	505
Fricas [A] (verification not implemented)	505
Sympy [A] (verification not implemented)	506
Maxima [A] (verification not implemented)	506
Giac [A] (verification not implemented)	506
Mupad [B] (verification not implemented)	507
Reduce [B] (verification not implemented)	507

Optimal result

Integrand size = 2, antiderivative size = 15

$$\int \cot^{-1}(x) dx = x \cot^{-1}(x) + \frac{1}{2} \log(1 + x^2)$$

output `x*arccot(x)+1/2*ln(x^2+1)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \cot^{-1}(x) dx = x \cot^{-1}(x) + \frac{1}{2} \log(1 + x^2)$$

input `Integrate[ArcCot[x], x]`

output `x*ArcCot[x] + Log[1 + x^2]/2`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {5346, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{-1}(x) dx$$

$$\downarrow \text{5346}$$

$$\int \frac{x}{x^2 + 1} dx + x \cot^{-1}(x)$$

$$\downarrow \text{240}$$

$$\frac{1}{2} \log(x^2 + 1) + x \cot^{-1}(x)$$

input `Int[ArcCot[x], x]`

output `x*ArcCot[x] + Log[1 + x^2]/2`

Defintions of rubi rules used

rule 240 `Int[(x_)/((a_) + (b_)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`

rule 5346 `Int[((a_) + ArcCot[(c_)*(x_)^(n_)])*(b_)^(p_), x_Symbol] := Simp[x*(a + b*ArcCot[c*x^n])^p, x] + Simp[b*c*n*p Int[x^n*((a + b*ArcCot[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

method	result	size
lookup	$x \operatorname{arccot}(x) + \frac{\ln(x^2+1)}{2}$	14
default	$x \operatorname{arccot}(x) + \frac{\ln(x^2+1)}{2}$	14
parallelrisc	$x \operatorname{arccot}(x) + \frac{\ln(x^2+1)}{2}$	14
parts	$x \operatorname{arccot}(x) + \frac{\ln(x^2+1)}{2}$	14
risc	$\frac{ix \ln(ix+1)}{2} - \frac{ix \ln(-ix+1)}{2} + \frac{\pi x}{2} + \frac{\ln(x^2+1)}{2}$	36

input `int(arccot(x),x,method=_RETURNVERBOSE)`output `x*arccot(x)+1/2*ln(x^2+1)`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \cot^{-1}(x) dx = x \operatorname{arccot}(x) + \frac{1}{2} \log(x^2 + 1)$$

input `integrate(arccot(x),x, algorithm="fricas")`output `x*arccot(x) + 1/2*log(x^2 + 1)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

$$\int \cot^{-1}(x) dx = x \operatorname{acot}(x) + \frac{\log(x^2 + 1)}{2}$$

input `integrate(acot(x),x)`output `x*acot(x) + log(x**2 + 1)/2`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \cot^{-1}(x) dx = x \operatorname{arccot}(x) + \frac{1}{2} \log(x^2 + 1)$$

input `integrate(arccot(x),x, algorithm="maxima")`output `x*arccot(x) + 1/2*log(x^2 + 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.40

$$\int \cot^{-1}(x) dx = x \arctan\left(\frac{1}{x}\right) + \frac{1}{2} \log\left(\frac{1}{x^2} + 1\right) - \frac{1}{2} \log\left(\frac{1}{x^2}\right)$$

input `integrate(arccot(x),x, algorithm="giac")`output `x*arctan(1/x) + 1/2*log(1/x^2 + 1) - 1/2*log(x^(-2))`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \cot^{-1}(x) dx = \frac{\ln(x^2 + 1)}{2} + x \operatorname{acot}(x)$$

input `int(acot(x),x)`

output `log(x^2 + 1)/2 + x*acot(x)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \cot^{-1}(x) dx = \operatorname{acot}(x) x + \frac{\log(x^2 + 1)}{2}$$

input `int(acot(x),x)`

output `(2*acot(x)*x + log(x**2 + 1))/2`

3.83 $\int \sec^{-1}(x) dx$

Optimal result	508
Mathematica [B] (verified)	508
Rubi [A] (verified)	509
Maple [A] (verified)	510
Fricas [A] (verification not implemented)	511
Sympy [C] (verification not implemented)	511
Maxima [B] (verification not implemented)	511
Giac [B] (verification not implemented)	512
Mupad [B] (verification not implemented)	512
Reduce [F]	513

Optimal result

Integrand size = 2, antiderivative size = 19

$$\int \sec^{-1}(x) dx = x \sec^{-1}(x) - \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{x^2}}\right)$$

output `x*arcsec(x)-arctanh((1-1/x^2)^(1/2))`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 64 vs. $2(19) = 38$.

Time = 0.04 (sec) , antiderivative size = 64, normalized size of antiderivative = 3.37

$$\int \sec^{-1}(x) dx = x \sec^{-1}(x) - \frac{\sqrt{-1+x^2}\left(-\log\left(1-\frac{x}{\sqrt{-1+x^2}}\right)+\log\left(1+\frac{x}{\sqrt{-1+x^2}}\right)\right)}{2\sqrt{1-\frac{1}{x^2}}x}$$

input `Integrate[ArcSec[x], x]`

output `x*ArcSec[x] - (Sqrt[-1 + x^2]*(-Log[1 - x/Sqrt[-1 + x^2]] + Log[1 + x/Sqrt[-1 + x^2]]))/(2*Sqrt[1 - x^(-2)]*x)`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 2.000$, Rules used = {5737, 798, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^{-1}(x) dx \\
 & \quad \downarrow \text{5737} \\
 & x \sec^{-1}(x) - \int \frac{1}{\sqrt{1 - \frac{1}{x^2}}} dx \\
 & \quad \downarrow \text{798} \\
 & \frac{1}{2} \int \frac{x^2}{\sqrt{1 - \frac{1}{x^2}}} d\frac{1}{x^2} + x \sec^{-1}(x) \\
 & \quad \downarrow \text{73} \\
 & x \sec^{-1}(x) - \int \frac{1}{1 - \frac{1}{x^4}} d\sqrt{1 - \frac{1}{x^2}} \\
 & \quad \downarrow \text{219} \\
 & x \sec^{-1}(x) - \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{x^2}}\right)
 \end{aligned}$$

input `Int[ArcSec[x], x]`

output `x*ArcSec[x] - ArcTanh[Sqrt[1 - x^(-2)]]`

Definitions of rubi rules used

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
 ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
 Q[a, 0] || LtQ[b, 0])`

rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/n Subst
 [Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a,
 b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

rule 5737 `Int[ArcSec[(c_.)*(x_)], x_Symbol] := Simp[x*ArcSec[c*x], x] - Simp[1/c In
 t[1/(x*sqrt[1 - 1/(c^2*x^2)]), x], x] /; FreeQ[c, x]`

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.16

method	result	size
lookup	$x \operatorname{arcsec}(x) - \ln\left(x + x\sqrt{1 - \frac{1}{x^2}}\right)$	22
default	$x \operatorname{arcsec}(x) - \ln\left(x + x\sqrt{1 - \frac{1}{x^2}}\right)$	22
parts	$x \operatorname{arcsec}(x) - \frac{\sqrt{x^2-1} \ln(x+\sqrt{x^2-1})}{\sqrt{\frac{x^2-1}{x^2}} x}$	39

input `int(arcsec(x), x, method=_RETURNVERBOSE)`

output `x*arcsec(x)-ln(x+x*(1-1/x^2)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.74

$$\int \sec^{-1}(x) dx = (x - 2) \operatorname{arcsec}(x) + 4 \arctan\left(-x + \sqrt{x^2 - 1}\right) + \log\left(-x + \sqrt{x^2 - 1}\right)$$

input `integrate(arcsec(x),x, algorithm="fricas")`

output `(x - 2)*arcsec(x) + 4*arctan(-x + sqrt(x^2 - 1)) + log(-x + sqrt(x^2 - 1))`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.97 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \sec^{-1}(x) dx = x \operatorname{asec}(x) - \begin{cases} \operatorname{acosh}(x) & \text{for } |x^2| > 1 \\ -i \operatorname{asin}(x) & \text{otherwise} \end{cases}$$

input `integrate(asec(x),x)`

output `x*asec(x) - Piecewise((acosh(x), Abs(x**2) > 1), (-I*asin(x), True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 35 vs. $2(17) = 34$.

Time = 0.03 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.84

$$\int \sec^{-1}(x) dx = x \operatorname{arcsec}(x) - \frac{1}{2} \log\left(\sqrt{-\frac{1}{x^2} + 1} + 1\right) + \frac{1}{2} \log\left(-\sqrt{-\frac{1}{x^2} + 1} + 1\right)$$

input `integrate(arcsec(x),x, algorithm="maxima")`

output

```
x*arcsec(x) - 1/2*log(sqrt(-1/x^2 + 1) + 1) + 1/2*log(-sqrt(-1/x^2 + 1) + 1)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 37 vs. 2(17) = 34.

Time = 0.13 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.95

$$\int \sec^{-1}(x) dx = x \arccos\left(\frac{1}{x}\right) - \frac{1}{2} \log\left(\sqrt{-\frac{1}{x^2} + 1} + 1\right) + \frac{1}{2} \log\left(-\sqrt{-\frac{1}{x^2} + 1} + 1\right)$$

input

```
integrate(arcsec(x),x, algorithm="giac")
```

output

```
x*arccos(1/x) - 1/2*log(sqrt(-1/x^2 + 1) + 1) + 1/2*log(-sqrt(-1/x^2 + 1) + 1)
```

Mupad [B] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \sec^{-1}(x) dx = x \arccos\left(\frac{1}{x}\right) - \ln\left(x + \sqrt{x^2 - 1}\right) \operatorname{sign}(x)$$

input

```
int(acos(1/x),x)
```

output

```
x*acos(1/x) - log(x + (x^2 - 1)^(1/2))*sign(x)
```

Reduce [F]

$$\int \sec^{-1}(x) dx = \int \operatorname{asec}(x) dx$$

input `int(asec(x),x)`

output `int(asec(x),x)`

3.84 $\int \csc^{-1}(x) dx$

Optimal result	514
Mathematica [B] (verified)	514
Rubi [A] (verified)	515
Maple [A] (verified)	516
Fricas [B] (verification not implemented)	517
Sympy [A] (verification not implemented)	517
Maxima [B] (verification not implemented)	517
Giac [B] (verification not implemented)	518
Mupad [B] (verification not implemented)	518
Reduce [F]	519

Optimal result

Integrand size = 2, antiderivative size = 17

$$\int \csc^{-1}(x) dx = x \csc^{-1}(x) + \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{x^2}}\right)$$

output `x*arccsc(x)+arctanh((1-1/x^2)^(1/2))`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 64 vs. 2(17) = 34.

Time = 0.02 (sec) , antiderivative size = 64, normalized size of antiderivative = 3.76

$$\int \csc^{-1}(x) dx = x \csc^{-1}(x) + \frac{\sqrt{-1 + x^2} \left(-\log\left(1 - \frac{x}{\sqrt{-1+x^2}}\right) + \log\left(1 + \frac{x}{\sqrt{-1+x^2}}\right) \right)}{2\sqrt{1 - \frac{1}{x^2}}x}$$

input `Integrate[ArcCsc[x], x]`

output `x*ArcCsc[x] + (Sqrt[-1 + x^2]*(-Log[1 - x/Sqrt[-1 + x^2]] + Log[1 + x/Sqrt[-1 + x^2]]))/(2*Sqrt[1 - x^(-2)]*x)`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 2.000$, Rules used = {5738, 798, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \csc^{-1}(x) dx \\
 & \quad \downarrow \text{5738} \\
 & \int \frac{1}{\sqrt{1 - \frac{1}{x^2}}} dx + x \csc^{-1}(x) \\
 & \quad \downarrow \text{798} \\
 & x \csc^{-1}(x) - \frac{1}{2} \int \frac{x^2}{\sqrt{1 - \frac{1}{x^2}}} d\frac{1}{x^2} \\
 & \quad \downarrow \text{73} \\
 & \int \frac{1}{1 - \frac{1}{x^4}} d\sqrt{1 - \frac{1}{x^2}} + x \csc^{-1}(x) \\
 & \quad \downarrow \text{219} \\
 & \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{x^2}}\right) + x \csc^{-1}(x)
 \end{aligned}$$

input `Int[ArcCsc[x], x]`

output `x*ArcCsc[x] + ArcTanh[Sqrt[1 - x^(-2)]]`

Definitions of rubi rules used

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
 ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
 Q[a, 0] || LtQ[b, 0])`

rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/n Subst
 [Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a,
 b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

rule 5738 `Int[ArcCsc[(c_.)*(x_)], x_Symbol] := Simp[x*ArcCsc[c*x], x] + Simp[1/c In
 t[1/(x*sqrt[1 - 1/(c^2*x^2)]), x], x] /; FreeQ[c, x]`

Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.18

method	result	size
lookup	$x \operatorname{arccsc}(x) + \ln\left(x + x\sqrt{1 - \frac{1}{x^2}}\right)$	20
default	$x \operatorname{arccsc}(x) + \ln\left(x + x\sqrt{1 - \frac{1}{x^2}}\right)$	20
parts	$x \operatorname{arccsc}(x) + \frac{\sqrt{x^2-1} \ln(x+\sqrt{x^2-1})}{\sqrt{\frac{x^2-1}{x^2}} x}$	38

input `int(arccsc(x), x, method=_RETURNVERBOSE)`

output `x*arccsc(x)+ln(x+x*(1-1/x^2)^(1/2))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 35 vs. $2(15) = 30$.

Time = 0.11 (sec) , antiderivative size = 35, normalized size of antiderivative = 2.06

$$\int \csc^{-1}(x) dx = (x - 2) \operatorname{arccsc}(x) - 4 \arctan\left(-x + \sqrt{x^2 - 1}\right) - \log\left(-x + \sqrt{x^2 - 1}\right)$$

input `integrate(arccsc(x),x, algorithm="fricas")`

output `(x - 2)*arccsc(x) - 4*arctan(-x + sqrt(x^2 - 1)) - log(-x + sqrt(x^2 - 1))`

Sympy [A] (verification not implemented)

Time = 0.95 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \csc^{-1}(x) dx = x \operatorname{acsc}(x) + \begin{cases} \operatorname{acosh}(x) & \text{for } |x^2| > 1 \\ -i \operatorname{asin}(x) & \text{otherwise} \end{cases}$$

input `integrate(acsc(x),x)`

output `x*acsc(x) + Piecewise((acosh(x), Abs(x**2) > 1), (-I*asin(x), True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 35 vs. $2(15) = 30$.

Time = 0.02 (sec) , antiderivative size = 35, normalized size of antiderivative = 2.06

$$\int \csc^{-1}(x) dx = x \operatorname{arccsc}(x) + \frac{1}{2} \log\left(\sqrt{-\frac{1}{x^2} + 1} + 1\right) - \frac{1}{2} \log\left(-\sqrt{-\frac{1}{x^2} + 1} + 1\right)$$

input `integrate(arccsc(x),x, algorithm="maxima")`

output `x*arccsc(x) + 1/2*log(sqrt(-1/x^2 + 1) + 1) - 1/2*log(-sqrt(-1/x^2 + 1) + 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 37 vs. $2(15) = 30$.

Time = 0.13 (sec) , antiderivative size = 37, normalized size of antiderivative = 2.18

$$\int \csc^{-1}(x) dx = x \arcsin\left(\frac{1}{x}\right) + \frac{1}{2} \log\left(\sqrt{-\frac{1}{x^2} + 1} + 1\right) - \frac{1}{2} \log\left(-\sqrt{-\frac{1}{x^2} + 1} + 1\right)$$

input `integrate(arccsc(x),x, algorithm="giac")`

output `x*arcsin(1/x) + 1/2*log(sqrt(-1/x^2 + 1) + 1) - 1/2*log(-sqrt(-1/x^2 + 1) + 1)`

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.18

$$\int \csc^{-1}(x) dx = x \operatorname{asin}\left(\frac{1}{x}\right) + \ln\left(x + \sqrt{x^2 - 1}\right) \operatorname{sign}(x)$$

input `int(asin(1/x),x)`

output `x*asin(1/x) + log(x + (x^2 - 1)^(1/2))*sign(x)`

Reduce [F]

$$\int \csc^{-1}(x) dx = \int \operatorname{acsc}(x) dx$$

input `int(acsc(x), x)`output `int(acsc(x), x)`

3.85 $\int \arcsin(x)^2 dx$

Optimal result	520
Mathematica [A] (verified)	520
Rubi [A] (verified)	521
Maple [A] (verified)	522
Fricas [A] (verification not implemented)	522
Sympy [A] (verification not implemented)	523
Maxima [A] (verification not implemented)	523
Giac [A] (verification not implemented)	523
Mupad [B] (verification not implemented)	524
Reduce [B] (verification not implemented)	524

Optimal result

Integrand size = 4, antiderivative size = 25

$$\int \arcsin(x)^2 dx = -2x + 2\sqrt{1-x^2} \arcsin(x) + x \arcsin(x)^2$$

output `-2*x+x*arcsin(x)^2+2*arcsin(x)*(-x^2+1)^(1/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \arcsin(x)^2 dx = -2x + 2\sqrt{1-x^2} \arcsin(x) + x \arcsin(x)^2$$

input `Integrate[ArcSin[x]^2,x]`

output `-2*x + 2*Sqrt[1 - x^2]*ArcSin[x] + x*ArcSin[x]^2`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5130, 5182, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \arcsin(x)^2 dx \\ & \quad \downarrow \text{5130} \\ & x \arcsin(x)^2 - 2 \int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx \\ & \quad \downarrow \text{5182} \\ & x \arcsin(x)^2 - 2 \left(\int 1 dx - \sqrt{1-x^2} \arcsin(x) \right) \\ & \quad \downarrow \text{24} \\ & x \arcsin(x)^2 - 2 \left(x - \sqrt{1-x^2} \arcsin(x) \right) \end{aligned}$$

input `Int[ArcSin[x]^2,x]`

output `x*ArcSin[x]^2 - 2*(x - Sqrt[1 - x^2]*ArcSin[x])`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

rule 5130 `Int[((a_.) + ArcSin[(c_.)*(x_)]*(b_.))^n, x_Symbol] :> Simp[x*(a + b*ArcSin[c*x])^n, x] - Simp[b*c*n Int[x*((a + b*ArcSin[c*x])^(n - 1)/Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`

rule 5182

```
Int[((a_.) + ArcSin[(c_.)*(x_.)]*(b_.))^(n_.)*(x_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(d + e*x^2)^(p + 1)*((a + b*ArcSin[c*x])^n/(2*e*(p + 1))), x] + Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/(1 - c^2*x^2)^p] Int[(1 - c^2*x^2)^(p + 1/2)*(a + b*ArcSin[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

method	result	size
default	$-2x + x \arcsin(x)^2 + 2 \arcsin(x) \sqrt{-x^2 + 1}$	24
orering	$x \arcsin(x)^2 + \frac{2 \arcsin(x)}{\sqrt{-x^2 + 1}} + (-1 + x)(1 + x)x \left(\frac{2}{-x^2 + 1} + \frac{2 \arcsin(x)x}{(-x^2 + 1)^{\frac{3}{2}}} \right)$	55

input `int(arcsin(x)^2,x,method=_RETURNVERBOSE)`output `-2*x+x*arcsin(x)^2+2*arcsin(x)*(-x^2+1)^(1/2)`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \arcsin(x)^2 dx = x \arcsin(x)^2 + 2 \sqrt{-x^2 + 1} \arcsin(x) - 2x$$

input `integrate(arcsin(x)^2,x, algorithm="fricas")`output `x*arcsin(x)^2 + 2*sqrt(-x^2 + 1)*arcsin(x) - 2*x`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \arcsin(x)^2 dx = x \arcsin^2(x) - 2x + 2\sqrt{1-x^2} \arcsin(x)$$

input `integrate(asin(x)**2,x)`

output `x*asin(x)**2 - 2*x + 2*sqrt(1 - x**2)*asin(x)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \arcsin(x)^2 dx = x \arcsin(x)^2 + 2\sqrt{-x^2+1} \arcsin(x) - 2x$$

input `integrate(arcsin(x)^2,x, algorithm="maxima")`

output `x*arcsin(x)^2 + 2*sqrt(-x^2 + 1)*arcsin(x) - 2*x`

Giac [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \arcsin(x)^2 dx = x \arcsin(x)^2 + 2\sqrt{-x^2+1} \arcsin(x) - 2x$$

input `integrate(arcsin(x)^2,x, algorithm="giac")`

output `x*arcsin(x)^2 + 2*sqrt(-x^2 + 1)*arcsin(x) - 2*x`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \arcsin(x)^2 dx = 2 \arcsin(x) \sqrt{1-x^2} + x (\arcsin(x)^2 - 2)$$

input `int(asin(x)^2,x)`

output `2*asin(x)*(1 - x^2)^(1/2) + x*(asin(x)^2 - 2)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \arcsin(x)^2 dx = \arcsin(x)^2 x + 2\sqrt{-x^2+1} \arcsin(x) - 2x$$

input `int(asin(x)^2,x)`

output `asin(x)**2*x + 2*sqrt(- x**2 + 1)*asin(x) - 2*x`

3.86 $\int \frac{\arcsin(x)}{x^2} dx$

Optimal result	525
Mathematica [A] (verified)	525
Rubi [A] (verified)	526
Maple [A] (verified)	527
Fricas [A] (verification not implemented)	528
Sympy [A] (verification not implemented)	528
Maxima [A] (verification not implemented)	528
Giac [A] (verification not implemented)	529
Mupad [B] (verification not implemented)	529
Reduce [B] (verification not implemented)	529

Optimal result

Integrand size = 6, antiderivative size = 22

$$\int \frac{\arcsin(x)}{x^2} dx = -\frac{\arcsin(x)}{x} - \operatorname{arctanh}(\sqrt{1-x^2})$$

output

```
-arcsin(x)/x-arctanh((-x^2+1)^(1/2))
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{\arcsin(x)}{x^2} dx = -\frac{\arcsin(x)}{x} - \operatorname{arctanh}(\sqrt{1-x^2})$$

input

```
Integrate[ArcSin[x]/x^2,x]
```

output

```
-(ArcSin[x]/x) - ArcTanh[Sqrt[1 - x^2]]
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {5138, 243, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\arcsin(x)}{x^2} dx \\
 & \quad \downarrow \text{5138} \\
 & \int \frac{1}{x\sqrt{1-x^2}} dx - \frac{\arcsin(x)}{x} \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{2} \int \frac{1}{x^2\sqrt{1-x^2}} dx^2 - \frac{\arcsin(x)}{x} \\
 & \quad \downarrow \text{73} \\
 & - \int \frac{1}{1-x^4} d\sqrt{1-x^2} - \frac{\arcsin(x)}{x} \\
 & \quad \downarrow \text{219} \\
 & - \frac{\arcsin(x)}{x} - \operatorname{arctanh}(\sqrt{1-x^2})
 \end{aligned}$$

input `Int[ArcSin[x]/x^2,x]`

output `-(ArcSin[x]/x) - ArcTanh[Sqrt[1 - x^2]]`

Definitions of rubi rules used

- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`
- rule 5138 `Int[((a_.) + ArcSin[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcSin[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcSin[c*x])^(n - 1)/Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.95

method	result	size
default	$-\frac{\arcsin(x)}{x} - \operatorname{arctanh}\left(\frac{1}{\sqrt{-x^2+1}}\right)$	21
parts	$-\frac{\arcsin(x)}{x} - \operatorname{arctanh}\left(\frac{1}{\sqrt{-x^2+1}}\right)$	21

input `int(arcsin(x)/x^2,x,method=_RETURNVERBOSE)`

output `-arcsin(x)/x-arctanh(1/(-x^2+1)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.77

$$\int \frac{\arcsin(x)}{x^2} dx = -\frac{x \log(\sqrt{-x^2+1}+1) - x \log(\sqrt{-x^2+1}-1) + 2 \arcsin(x)}{2x}$$

input `integrate(arcsin(x)/x^2,x, algorithm="fricas")`output `-1/2*(x*log(sqrt(-x^2 + 1) + 1) - x*log(sqrt(-x^2 + 1) - 1) + 2*arcsin(x))
/x`**Sympy [A] (verification not implemented)**

Time = 0.92 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{\arcsin(x)}{x^2} dx = \begin{cases} -\operatorname{acosh}\left(\frac{1}{x}\right) & \text{for } \frac{1}{|x^2|} > 1 \\ i \operatorname{asin}\left(\frac{1}{x}\right) & \text{otherwise} \end{cases} - \frac{\operatorname{asin}(x)}{x}$$

input `integrate(asin(x)/x**2,x)`output `Piecewise((-acosh(1/x), 1/Abs(x**2) > 1), (I*asin(1/x), True)) - asin(x)/x`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.50

$$\int \frac{\arcsin(x)}{x^2} dx = -\frac{\arcsin(x)}{x} - \log\left(\frac{2\sqrt{-x^2+1}}{|x|} + \frac{2}{|x|}\right)$$

input `integrate(arcsin(x)/x^2,x, algorithm="maxima")`output `-arcsin(x)/x - log(2*sqrt(-x^2 + 1)/abs(x) + 2/abs(x))`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.73

$$\int \frac{\arcsin(x)}{x^2} dx = -\frac{\arcsin(x)}{x} - \frac{1}{2} \log(\sqrt{-x^2+1}+1) + \frac{1}{2} \log(-\sqrt{-x^2+1}+1)$$

input `integrate(arcsin(x)/x^2,x, algorithm="giac")`

output `-arcsin(x)/x - 1/2*log(sqrt(-x^2 + 1) + 1) + 1/2*log(-sqrt(-x^2 + 1) + 1)`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{\arcsin(x)}{x^2} dx = -\operatorname{atanh}\left(\frac{1}{\sqrt{1-x^2}}\right) - \frac{\operatorname{asin}(x)}{x}$$

input `int(asin(x)/x^2,x)`

output `- atanh(1/(1 - x^2)^(1/2)) - asin(x)/x`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

$$\int \frac{\arcsin(x)}{x^2} dx = \frac{-\operatorname{asin}(x) + \log\left(\tan\left(\frac{\operatorname{asin}(x)}{2}\right)\right) x}{x}$$

input `int(asin(x)/x^2,x)`

output `(- asin(x) + log(tan(asin(x)/2))*x)/x`

3.87 $\int \frac{1}{\sqrt{a^2-x^2}} dx$

Optimal result	530
Mathematica [A] (verified)	530
Rubi [A] (verified)	531
Maple [A] (verified)	532
Fricas [A] (verification not implemented)	532
Sympy [C] (verification not implemented)	532
Maxima [A] (verification not implemented)	533
Giac [A] (verification not implemented)	533
Mupad [B] (verification not implemented)	533
Reduce [B] (verification not implemented)	534

Optimal result

Integrand size = 13, antiderivative size = 16

$$\int \frac{1}{\sqrt{a^2-x^2}} dx = \arctan\left(\frac{x}{\sqrt{a^2-x^2}}\right)$$

output `arctan(x/(a^2-x^2)^(1/2))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{a^2-x^2}} dx = \arctan\left(\frac{x}{\sqrt{a^2-x^2}}\right)$$

input `Integrate[1/Sqrt[a^2 - x^2],x]`

output `ArcTan[x/Sqrt[a^2 - x^2]]`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {224, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx$$

$$\downarrow \text{224}$$

$$\int \frac{1}{\frac{x^2}{a^2 - x^2} + 1} d \frac{x}{\sqrt{a^2 - x^2}}$$

$$\downarrow \text{216}$$

$$\arctan \left(\frac{x}{\sqrt{a^2 - x^2}} \right)$$

input `Int[1/Sqrt[a^2 - x^2],x]`

output `ArcTan[x/Sqrt[a^2 - x^2]]`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result	size
default	$\arctan\left(\frac{x}{\sqrt{a^2-x^2}}\right)$	15
pseudoelliptic	$-\arctan\left(\frac{\sqrt{a^2-x^2}}{x}\right)$	19

input `int(1/(a^2-x^2)^(1/2),x,method=_RETURNVERBOSE)`

output `arctan(x/(a^2-x^2)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.44

$$\int \frac{1}{\sqrt{a^2-x^2}} dx = -2 \arctan\left(-\frac{a-\sqrt{a^2-x^2}}{x}\right)$$

input `integrate(1/(a^2-x^2)^(1/2),x, algorithm="fricas")`

output `-2*arctan(-(a - sqrt(a^2 - x^2))/x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.49 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.19

$$\int \frac{1}{\sqrt{a^2-x^2}} dx = \begin{cases} -i \operatorname{acosh}\left(\frac{x}{a}\right) & \text{for } \left|\frac{x^2}{a^2}\right| > 1 \\ \operatorname{asin}\left(\frac{x}{a}\right) & \text{otherwise} \end{cases}$$

input `integrate(1/(a**2-x**2)**(1/2),x)`

output `Piecewise((-I*acosh(x/a), Abs(x**2/a**2) > 1), (asin(x/a), True))`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.38

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \arcsin\left(\frac{x}{a}\right)$$

input `integrate(1/(a^2-x^2)^(1/2),x, algorithm="maxima")`

output `arcsin(x/a)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.75

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \frac{1}{2} a^2 \arcsin\left(\frac{x}{a}\right) \operatorname{sgn}(a) + \frac{1}{2} \sqrt{a^2 - x^2} x$$

input `integrate(1/(a^2-x^2)^(1/2),x, algorithm="giac")`

output `1/2*a^2*arcsin(x/a)*sgn(a) + 1/2*sqrt(a^2 - x^2)*x`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \operatorname{atan}\left(\frac{x}{\sqrt{a^2 - x^2}}\right)$$

input `int(1/(a^2 - x^2)^(1/2),x)`

output `atan(x/(a^2 - x^2)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.38

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \operatorname{asin}\left(\frac{x}{a}\right)$$

input `int(1/(a^2-x^2)^(1/2),x)`

output `asin(x/a)`

$$3.88 \quad \int \frac{1}{\sqrt{1-2x-x^2}} dx$$

Optimal result	535
Mathematica [B] (verified)	535
Rubi [A] (verified)	536
Maple [A] (verified)	537
Fricas [B] (verification not implemented)	537
Sympy [A] (verification not implemented)	537
Maxima [A] (verification not implemented)	538
Giac [B] (verification not implemented)	538
Mupad [B] (verification not implemented)	539
Reduce [B] (verification not implemented)	539

Optimal result

Integrand size = 14, antiderivative size = 10

$$\int \frac{1}{\sqrt{1-2x-x^2}} dx = \arcsin\left(\frac{1+x}{\sqrt{2}}\right)$$

output `arcsin(1/2*(1+x)*2^(1/2))`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 23 vs. 2(10) = 20.

Time = 0.05 (sec) , antiderivative size = 23, normalized size of antiderivative = 2.30

$$\int \frac{1}{\sqrt{1-2x-x^2}} dx = 2 \arctan\left(\frac{x}{-1 + \sqrt{1-2x-x^2}}\right)$$

input `Integrate[1/Sqrt[1 - 2*x - x^2],x]`

output `2*ArcTan[x/(-1 + Sqrt[1 - 2*x - x^2])]`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.70, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {1090, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{-x^2 - 2x + 1}} dx$$

↓ 1090

$$-\frac{\int \frac{1}{\sqrt{1 - \frac{1}{8}(-2x-2)^2}} d(-2x-2)}{2\sqrt{2}}$$

↓ 223

$$-\arcsin\left(\frac{-2x-2}{2\sqrt{2}}\right)$$

input `Int[1/Sqrt[1 - 2*x - x^2],x]`

output `-ArcSin[(-2 - 2*x)/(2*Sqrt[2])]`

Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 1090 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/(2*c*(-4*(c/(b^2 - 4*a*c))))^p) Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

method	result	size
default	$\arcsin\left(\frac{(1+x)\sqrt{2}}{2}\right)$	10
trager	$\text{RootOf}(_Z^2 + 1) \ln(-\text{RootOf}(_Z^2 + 1) x + \sqrt{-x^2 - 2x + 1} - \text{RootOf}(_Z^2 + 1))$	39

input `int(1/(-x^2-2*x+1)^(1/2),x,method=_RETURNVERBOSE)`

output `arcsin(1/2*(1+x)*2^(1/2))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 21 vs. 2(9) = 18.

Time = 0.09 (sec) , antiderivative size = 21, normalized size of antiderivative = 2.10

$$\int \frac{1}{\sqrt{1-2x-x^2}} dx = -2 \arctan\left(\frac{\sqrt{-x^2-2x+1}-1}{x}\right)$$

input `integrate(1/(-x^2-2*x+1)^(1/2),x, algorithm="fricas")`

output `-2*arctan((sqrt(-x^2 - 2*x + 1) - 1)/x)`

Sympy [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1-2x-x^2}} dx = \text{asin}\left(\frac{\sqrt{2}(x+1)}{2}\right)$$

input `integrate(1/(-x**2-2*x+1)**(1/2),x)`

output `asin(sqrt(2)*(x + 1)/2)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

$$\int \frac{1}{\sqrt{1-2x-x^2}} dx = -\arcsin\left(-\frac{1}{2}\sqrt{2}(x+1)\right)$$

input `integrate(1/(-x^2-2*x+1)^(1/2),x, algorithm="maxima")`

output `-arcsin(-1/2*sqrt(2)*(x + 1))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(9) = 18.

Time = 0.13 (sec) , antiderivative size = 27, normalized size of antiderivative = 2.70

$$\int \frac{1}{\sqrt{1-2x-x^2}} dx = \frac{1}{2}\sqrt{-x^2-2x+1}(x+1) + \arcsin\left(\frac{1}{2}\sqrt{2}(x+1)\right)$$

input `integrate(1/(-x^2-2*x+1)^(1/2),x, algorithm="giac")`

output `1/2*sqrt(-x^2 - 2*x + 1)*(x + 1) + arcsin(1/2*sqrt(2)*(x + 1))`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

$$\int \frac{1}{\sqrt{1-2x-x^2}} dx = \operatorname{asin}\left(\frac{\sqrt{8}(2x+2)}{8}\right)$$

input `int(1/(1 - x^2 - 2*x)^(1/2),x)`

output `asin((8^(1/2)*(2*x + 2))/8)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

$$\int \frac{1}{\sqrt{1-2x-x^2}} dx = \operatorname{asin}\left(\frac{x+1}{\sqrt{2}}\right)$$

input `int(1/(-x^2-2*x+1)^(1/2),x)`

output `asin((x + 1)/sqrt(2))`

3.89 $\int \frac{1}{a^2+x^2} dx$

Optimal result	540
Mathematica [A] (verified)	540
Rubi [A] (verified)	541
Maple [A] (verified)	541
Fricas [A] (verification not implemented)	542
Sympy [C] (verification not implemented)	542
Maxima [A] (verification not implemented)	543
Giac [A] (verification not implemented)	543
Mupad [B] (verification not implemented)	543
Reduce [B] (verification not implemented)	544

Optimal result

Integrand size = 9, antiderivative size = 10

$$\int \frac{1}{a^2+x^2} dx = \frac{\arctan\left(\frac{x}{a}\right)}{a}$$

output

```
arctan(x/a)/a
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{a^2+x^2} dx = \frac{\arctan\left(\frac{x}{a}\right)}{a}$$

input

```
Integrate[(a^2 + x^2)^(-1),x]
```

output

```
ArcTan[x/a]/a
```

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{a^2 + x^2} dx$$

↓ 216

$$\frac{\arctan\left(\frac{x}{a}\right)}{a}$$

input `Int[(a^2 + x^2)^(-1), x]`

output `ArcTan[x/a]/a`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

method	result	size
default	$\frac{\arctan\left(\frac{x}{a}\right)}{a}$	11
risch	$\frac{\arctan\left(\frac{x}{a}\right)}{a}$	11
parallelrisch	$-\frac{i \ln(-ia+x) - i \ln(ia+x)}{2a}$	27

input `int(1/(a^2+x^2),x,method=_RETURNVERBOSE)`

output `arctan(x/a)/a`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{a^2 + x^2} dx = \frac{\arctan\left(\frac{x}{a}\right)}{a}$$

input `integrate(1/(a^2+x^2),x, algorithm="fricas")`

output `arctan(x/a)/a`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 2.00

$$\int \frac{1}{a^2 + x^2} dx = \frac{-\frac{i \log(-ia+x)}{2} + \frac{i \log(ia+x)}{2}}{a}$$

input `integrate(1/(a**2+x**2),x)`

output `(-I*log(-I*a + x)/2 + I*log(I*a + x)/2)/a`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{a^2 + x^2} dx = \frac{\arctan\left(\frac{x}{a}\right)}{a}$$

input `integrate(1/(a^2+x^2),x, algorithm="maxima")`output `arctan(x/a)/a`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{a^2 + x^2} dx = \frac{\arctan\left(\frac{x}{a}\right)}{a}$$

input `integrate(1/(a^2+x^2),x, algorithm="giac")`output `arctan(x/a)/a`**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{a^2 + x^2} dx = \frac{\operatorname{atan}\left(\frac{x}{a}\right)}{a}$$

input `int(1/(a^2 + x^2),x)`output `atan(x/a)/a`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{a^2 + x^2} dx = \frac{\operatorname{atan}\left(\frac{x}{a}\right)}{a}$$

input `int(1/(a^2+x^2),x)`

output `atan(x/a)/a`

3.90 $\int \frac{1}{a+bx^2} dx$

Optimal result	545
Mathematica [A] (verified)	545
Rubi [A] (verified)	546
Maple [A] (verified)	546
Fricas [A] (verification not implemented)	547
Sympy [B] (verification not implemented)	547
Maxima [A] (verification not implemented)	548
Giac [A] (verification not implemented)	548
Mupad [B] (verification not implemented)	548
Reduce [B] (verification not implemented)	549

Optimal result

Integrand size = 9, antiderivative size = 24

$$\int \frac{1}{a+bx^2} dx = \frac{\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{b}}$$

output `arctan(x*b^(1/2)/a^(1/2))/a^(1/2)/b^(1/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{1}{a+bx^2} dx = \frac{\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{b}}$$

input `Integrate[(a + b*x^2)^(-1),x]`

output `ArcTan[(Sqrt[b]*x)/Sqrt[a]]/(Sqrt[a]*Sqrt[b])`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{a + bx^2} dx$$

↓ 218

$$\frac{\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{b}}$$

input `Int[(a + b*x^2)^(-1), x]`

output `ArcTan[(Sqrt[b]*x)/Sqrt[a]]/(Sqrt[a]*Sqrt[b])`

Defintions of rubi rules used

rule 218

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

method	result	size
default	$\frac{\arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}}$	16
risch	$-\frac{\ln(bx + \sqrt{-ab})}{2\sqrt{-ab}} + \frac{\ln(-bx + \sqrt{-ab})}{2\sqrt{-ab}}$	41

input `int(1/(b*x^2+a),x,method=_RETURNVERBOSE)`

output `1/(a*b)^(1/2)*arctan(b*x/(a*b)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 67, normalized size of antiderivative = 2.79

$$\int \frac{1}{a + bx^2} dx = \left[-\frac{\sqrt{-ab} \log\left(\frac{bx^2 - 2\sqrt{-ab}x - a}{bx^2 + a}\right)}{2ab}, \frac{\sqrt{ab} \arctan\left(\frac{\sqrt{ab}x}{a}\right)}{ab} \right]$$

input `integrate(1/(b*x^2+a),x, algorithm="fricas")`

output `[-1/2*sqrt(-a*b)*log((b*x^2 - 2*sqrt(-a*b)*x - a)/(b*x^2 + a))/(a*b), sqrt(a*b)*arctan(sqrt(a*b)*x/a)/(a*b)]`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 53 vs. 2(22) = 44.

Time = 0.06 (sec) , antiderivative size = 53, normalized size of antiderivative = 2.21

$$\int \frac{1}{a + bx^2} dx = -\frac{\sqrt{-\frac{1}{ab}} \log\left(-a\sqrt{-\frac{1}{ab}} + x\right)}{2} + \frac{\sqrt{-\frac{1}{ab}} \log\left(a\sqrt{-\frac{1}{ab}} + x\right)}{2}$$

input `integrate(1/(b*x**2+a),x)`

output `-sqrt(-1/(a*b))*log(-a*sqrt(-1/(a*b)) + x)/2 + sqrt(-1/(a*b))*log(a*sqrt(-1/(a*b)) + x)/2`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.62

$$\int \frac{1}{a + bx^2} dx = \frac{\arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}}$$

input `integrate(1/(b*x^2+a),x, algorithm="maxima")`output `arctan(b*x/sqrt(a*b))/sqrt(a*b)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.62

$$\int \frac{1}{a + bx^2} dx = \frac{\arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}}$$

input `integrate(1/(b*x^2+a),x, algorithm="giac")`output `arctan(b*x/sqrt(a*b))/sqrt(a*b)`**Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int \frac{1}{a + bx^2} dx = \frac{\operatorname{atan}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{b}}$$

input `int(1/(a + b*x^2),x)`output `atan((b^(1/2)*x)/a^(1/2))/(a^(1/2)*b^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.96

$$\int \frac{1}{a + bx^2} dx = \frac{\sqrt{b} \sqrt{a} \operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right)}{ab}$$

input `int(1/(b*x^2+a),x)`

output `(sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*sqrt(a))))/(a*b)`

3.91 $\int \frac{1}{2-x+x^2} dx$

Optimal result	550
Mathematica [A] (verified)	550
Rubi [A] (verified)	551
Maple [A] (verified)	552
Fricas [A] (verification not implemented)	552
Sympy [A] (verification not implemented)	552
Maxima [A] (verification not implemented)	553
Giac [A] (verification not implemented)	553
Mupad [B] (verification not implemented)	553
Reduce [B] (verification not implemented)	554

Optimal result

Integrand size = 10, antiderivative size = 19

$$\int \frac{1}{2-x+x^2} dx = -\frac{2 \arctan\left(\frac{1-2x}{\sqrt{7}}\right)}{\sqrt{7}}$$

output `-2/7*arctan(1/7*(1-2*x)*7^(1/2))*7^(1/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{1}{2-x+x^2} dx = \frac{2 \arctan\left(\frac{-1+2x}{\sqrt{7}}\right)}{\sqrt{7}}$$

input `Integrate[(2 - x + x^2)^(-1),x]`

output `(2*ArcTan[(-1 + 2*x)/Sqrt[7]])/Sqrt[7]`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {1083, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 - x + 2} dx$$

↓ 1083

$$-2 \int \frac{1}{-(2x - 1)^2 - 7} d(2x - 1)$$

↓ 217

$$\frac{2 \arctan\left(\frac{2x-1}{\sqrt{7}}\right)}{\sqrt{7}}$$

input `Int[(2 - x + x^2)^(-1), x]`

output `(2*ArcTan[(-1 + 2*x)/Sqrt[7]])/Sqrt[7]`

Defintions of rubi rules used

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])`

rule 1083 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`

Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

method	result	size
default	$\frac{2\sqrt{7} \arctan\left(\frac{(2x-1)\sqrt{7}}{7}\right)}{7}$	17
risch	$\frac{2\sqrt{7} \arctan\left(\frac{(2x-1)\sqrt{7}}{7}\right)}{7}$	17

input `int(1/(x^2-x+2),x,method=_RETURNVERBOSE)`output `2/7*7^(1/2)*arctan(1/7*(2*x-1)*7^(1/2))`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

$$\int \frac{1}{2-x+x^2} dx = \frac{2}{7} \sqrt{7} \arctan\left(\frac{1}{7} \sqrt{7}(2x-1)\right)$$

input `integrate(1/(x^2-x+2),x, algorithm="fricas")`output `2/7*sqrt(7)*arctan(1/7*sqrt(7)*(2*x - 1))`**Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.37

$$\int \frac{1}{2-x+x^2} dx = \frac{2\sqrt{7} \operatorname{atan}\left(\frac{2\sqrt{7}x}{7} - \frac{\sqrt{7}}{7}\right)}{7}$$

input `integrate(1/(x**2-x+2),x)`output `2*sqrt(7)*atan(2*sqrt(7)*x/7 - sqrt(7)/7)/7`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

$$\int \frac{1}{2-x+x^2} dx = \frac{2}{7} \sqrt{7} \arctan\left(\frac{1}{7} \sqrt{7}(2x-1)\right)$$

input `integrate(1/(x^2-x+2),x, algorithm="maxima")`output `2/7*sqrt(7)*arctan(1/7*sqrt(7)*(2*x - 1))`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

$$\int \frac{1}{2-x+x^2} dx = \frac{2}{7} \sqrt{7} \arctan\left(\frac{1}{7} \sqrt{7}(2x-1)\right)$$

input `integrate(1/(x^2-x+2),x, algorithm="giac")`output `2/7*sqrt(7)*arctan(1/7*sqrt(7)*(2*x - 1))`**Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

$$\int \frac{1}{2-x+x^2} dx = \frac{2\sqrt{7} \operatorname{atan}\left(\frac{\sqrt{7}(2x-1)}{7}\right)}{7}$$

input `int(1/(x^2 - x + 2),x)`output `(2*7^(1/2)*atan((7^(1/2)*(2*x - 1))/7))/7`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{1}{2-x+x^2} dx = \frac{2\sqrt{7} \operatorname{atan}\left(\frac{2x-1}{\sqrt{7}}\right)}{7}$$

input `int(1/(x^2-x+2),x)`

output `(2*sqrt(7)*atan((2*x - 1)/sqrt(7)))/7`

3.92 $\int x \arctan(x) dx$

Optimal result	555
Mathematica [A] (verified)	555
Rubi [A] (verified)	556
Maple [A] (verified)	557
Fricas [A] (verification not implemented)	558
Sympy [A] (verification not implemented)	558
Maxima [A] (verification not implemented)	558
Giac [A] (verification not implemented)	559
Mupad [B] (verification not implemented)	559
Reduce [B] (verification not implemented)	559

Optimal result

Integrand size = 4, antiderivative size = 21

$$\int x \arctan(x) dx = -\frac{x}{2} + \frac{\arctan(x)}{2} + \frac{1}{2}x^2 \arctan(x)$$

output `-1/2*x+1/2*arctan(x)+1/2*x^2*arctan(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.76

$$\int x \arctan(x) dx = \frac{1}{2}(-x + (1 + x^2) \arctan(x))$$

input `Integrate[x*ArcTan[x],x]`

output `(-x + (1 + x^2)*ArcTan[x])/2`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.95, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5361, 262, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \arctan(x) dx$$

$$\downarrow 5361$$

$$\frac{1}{2}x^2 \arctan(x) - \frac{1}{2} \int \frac{x^2}{x^2+1} dx$$

$$\downarrow 262$$

$$\frac{1}{2} \left(\int \frac{1}{x^2+1} dx - x \right) + \frac{1}{2}x^2 \arctan(x)$$

$$\downarrow 216$$

$$\frac{1}{2}x^2 \arctan(x) + \frac{1}{2}(\arctan(x) - x)$$

input

```
Int[x*ArcTan[x],x]
```

output

```
(x^2*ArcTan[x])/2 + (-x + ArcTan[x])/2
```

Defintions of rubi rules used

rule 216

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

rule 262

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)
^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/
(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b
, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c
, 2, m, p, x]
```

rule 5361

```
Int(((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :=
Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m +
1)) Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x],
x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] &
& IntegerQ[m])) && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.76

method	result	size
default	$-\frac{x}{2} + \frac{\arctan(x)}{2} + \frac{x^2 \arctan(x)}{2}$	16
meijerg	$-\frac{x}{2} + \frac{(3x^2+3) \arctan(x)}{6}$	16
parallelrisch	$-\frac{x}{2} + \frac{\arctan(x)}{2} + \frac{x^2 \arctan(x)}{2}$	16
parts	$-\frac{x}{2} + \frac{\arctan(x)}{2} + \frac{x^2 \arctan(x)}{2}$	16
orering	$\arctan(x)(x^2 + 1) + \left(-\frac{x^2}{2} - \frac{1}{2}\right) \left(\arctan(x) + \frac{x}{x^2+1}\right)$	30
risch	$-\frac{ix^2 \ln(ix+1)}{4} + \frac{ix^2 \ln(-ix+1)}{4} - \frac{x}{2} + \frac{\arctan(x)}{2}$	35

input

```
int(x*arctan(x),x,method=_RETURNVERBOSE)
```

output

```
-1/2*x+1/2*arctan(x)+1/2*x^2*arctan(x)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int x \arctan(x) dx = \frac{1}{2} (x^2 + 1) \arctan(x) - \frac{1}{2} x$$

input `integrate(x*arctan(x),x, algorithm="fricas")`output `1/2*(x^2 + 1)*arctan(x) - 1/2*x`**Sympy [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int x \arctan(x) dx = \frac{x^2 \operatorname{atan}(x)}{2} - \frac{x}{2} + \frac{\operatorname{atan}(x)}{2}$$

input `integrate(x*atan(x),x)`output `x**2*atan(x)/2 - x/2 + atan(x)/2`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int x \arctan(x) dx = \frac{1}{2} x^2 \arctan(x) - \frac{1}{2} x + \frac{1}{2} \arctan(x)$$

input `integrate(x*arctan(x),x, algorithm="maxima")`output `1/2*x^2*arctan(x) - 1/2*x + 1/2*arctan(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int x \arctan(x) dx = \frac{1}{2} x^2 \arctan(x) - \frac{1}{2} x + \frac{1}{2} \arctan(x)$$

input `integrate(x*arctan(x),x, algorithm="giac")`

output `1/2*x^2*arctan(x) - 1/2*x + 1/2*arctan(x)`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

$$\int x \arctan(x) dx = \operatorname{atan}(x) \left(\frac{x^2}{2} + \frac{1}{2} \right) - \frac{x}{2}$$

input `int(x*atan(x),x)`

output `atan(x)*(x^2/2 + 1/2) - x/2`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int x \arctan(x) dx = \frac{\operatorname{atan}(x) x^2}{2} + \frac{\operatorname{atan}(x)}{2} - \frac{x}{2}$$

input `int(x*atan(x),x)`

output `(atan(x)*x**2 + atan(x) - x)/2`

3.93 $\int x^2 \arccos(x) dx$

Optimal result	560
Mathematica [A] (verified)	560
Rubi [A] (verified)	561
Maple [A] (verified)	562
Fricas [A] (verification not implemented)	563
Sympy [A] (verification not implemented)	563
Maxima [A] (verification not implemented)	563
Giac [A] (verification not implemented)	564
Mupad [B] (verification not implemented)	564
Reduce [B] (verification not implemented)	564

Optimal result

Integrand size = 6, antiderivative size = 40

$$\int x^2 \arccos(x) dx = -\frac{1}{3}\sqrt{1-x^2} + \frac{1}{9}(1-x^2)^{3/2} + \frac{1}{3}x^3 \arccos(x)$$

output `1/9*(-x^2+1)^(3/2)+1/3*x^3*arccos(x)-1/3*(-x^2+1)^(1/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.75

$$\int x^2 \arccos(x) dx = -\frac{1}{9}\sqrt{1-x^2}(2+x^2) + \frac{1}{3}x^3 \arccos(x)$$

input `Integrate[x^2*ArcCos[x],x]`

output `-1/9*(Sqrt[1 - x^2]*(2 + x^2)) + (x^3*ArcCos[x])/3`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.08, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {5139, 243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \arccos(x) dx \\
 & \quad \downarrow \text{5139} \\
 & \frac{1}{3} \int \frac{x^3}{\sqrt{1-x^2}} dx + \frac{1}{3} x^3 \arccos(x) \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{6} \int \frac{x^2}{\sqrt{1-x^2}} dx^2 + \frac{1}{3} x^3 \arccos(x) \\
 & \quad \downarrow \text{53} \\
 & \frac{1}{6} \int \left(\frac{1}{\sqrt{1-x^2}} - \sqrt{1-x^2} \right) dx^2 + \frac{1}{3} x^3 \arccos(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{3} x^3 \arccos(x) + \frac{1}{6} \left(\frac{2}{3} (1-x^2)^{3/2} - 2\sqrt{1-x^2} \right)
 \end{aligned}$$

input `Int [x^2*ArcCos [x] , x]`

output `(-2*sqrt [1 - x^2] + (2*(1 - x^2)^(3/2))/3)/6 + (x^3*ArcCos [x])/3`

Definitions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5139 `Int[((a_.) + ArcCos[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCos[c*x])^n/(d*(m + 1))), x] + Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCos[c*x])^(n - 1)/Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.85

method	result	size
default	$\frac{x^3 \arccos(x)}{3} - \frac{\sqrt{-x^2+1} x^2}{9} - \frac{2\sqrt{-x^2+1}}{9}$	34
parts	$\frac{x^3 \arccos(x)}{3} - \frac{\sqrt{-x^2+1} x^2}{9} - \frac{2\sqrt{-x^2+1}}{9}$	34
orering	$\frac{(5x^4+2x^2-4) \arccos(x)}{9x} - \frac{(x^2+2)(-1+x)(1+x) \left(2x \arccos(x) - \frac{x^2}{\sqrt{-x^2+1}} \right)}{9x^2}$	57

input `int(x^2*arccos(x),x,method=_RETURNVERBOSE)`

output `1/3*x^3*arccos(x)-1/9*(-x^2+1)^(1/2)*x^2-2/9*(-x^2+1)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.60

$$\int x^2 \arccos(x) dx = \frac{1}{3} x^3 \arccos(x) - \frac{1}{9} (x^2 + 2) \sqrt{-x^2 + 1}$$

input `integrate(x^2*arccos(x),x, algorithm="fricas")`output `1/3*x^3*arccos(x) - 1/9*(x^2 + 2)*sqrt(-x^2 + 1)`**Sympy [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.80

$$\int x^2 \arccos(x) dx = \frac{x^3 \arccos(x)}{3} - \frac{x^2 \sqrt{1-x^2}}{9} - \frac{2\sqrt{1-x^2}}{9}$$

input `integrate(x**2*acos(x),x)`output `x**3*acos(x)/3 - x**2*sqrt(1 - x**2)/9 - 2*sqrt(1 - x**2)/9`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.82

$$\int x^2 \arccos(x) dx = \frac{1}{3} x^3 \arccos(x) - \frac{1}{9} \sqrt{-x^2 + 1} x^2 - \frac{2}{9} \sqrt{-x^2 + 1}$$

input `integrate(x^2*arccos(x),x, algorithm="maxima")`output `1/3*x^3*arccos(x) - 1/9*sqrt(-x^2 + 1)*x^2 - 2/9*sqrt(-x^2 + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.82

$$\int x^2 \arccos(x) dx = \frac{1}{3} x^3 \arccos(x) - \frac{1}{9} \sqrt{-x^2 + 1} x^2 - \frac{2}{9} \sqrt{-x^2 + 1}$$

input `integrate(x^2*arccos(x),x, algorithm="giac")`

output `1/3*x^3*arccos(x) - 1/9*sqrt(-x^2 + 1)*x^2 - 2/9*sqrt(-x^2 + 1)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.60

$$\int x^2 \arccos(x) dx = \frac{x^3 \arccos(x)}{3} - \frac{\sqrt{1-x^2}(x^2+2)}{9}$$

input `int(x^2*acos(x),x)`

output `(x^3*acos(x))/3 - ((1 - x^2)^(1/2)*(x^2 + 2))/9`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.78

$$\int x^2 \arccos(x) dx = \frac{\arccos(x) x^3}{3} - \frac{\sqrt{-x^2 + 1} x^2}{9} - \frac{2\sqrt{-x^2 + 1}}{9}$$

input `int(x^2*acos(x),x)`

output `(3*acos(x)*x**3 - sqrt(-x**2 + 1)*x**2 - 2*sqrt(-x**2 + 1))/9`

3.94 $\int x \arctan(x)^2 dx$

Optimal result	565
Mathematica [A] (verified)	565
Rubi [A] (verified)	566
Maple [A] (verified)	568
Fricas [A] (verification not implemented)	568
Sympy [A] (verification not implemented)	568
Maxima [A] (verification not implemented)	569
Giac [A] (verification not implemented)	569
Mupad [B] (verification not implemented)	570
Reduce [B] (verification not implemented)	570

Optimal result

Integrand size = 6, antiderivative size = 35

$$\int x \arctan(x)^2 dx = -x \arctan(x) + \frac{\arctan(x)^2}{2} + \frac{1}{2}x^2 \arctan(x)^2 + \frac{1}{2} \log(1 + x^2)$$

output `-x*arctan(x)+1/2*arctan(x)^2+1/2*x^2*arctan(x)^2+1/2*ln(x^2+1)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.74

$$\int x \arctan(x)^2 dx = \frac{1}{2}(-2x \arctan(x) + (1 + x^2) \arctan(x)^2 + \log(1 + x^2))$$

input `Integrate[x*ArcTan[x]^2,x]`

output `(-2*x*ArcTan[x] + (1 + x^2)*ArcTan[x]^2 + Log[1 + x^2])/2`

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {5361, 5451, 5345, 240, 5419}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \arctan(x)^2 dx \\
 & \quad \downarrow \text{5361} \\
 & \frac{1}{2}x^2 \arctan(x)^2 - \int \frac{x^2 \arctan(x)}{x^2 + 1} dx \\
 & \quad \downarrow \text{5451} \\
 & \int \frac{\arctan(x)}{x^2 + 1} dx - \int \arctan(x) dx + \frac{1}{2}x^2 \arctan(x)^2 \\
 & \quad \downarrow \text{5345} \\
 & \int \frac{\arctan(x)}{x^2 + 1} dx + \int \frac{x}{x^2 + 1} dx + \frac{1}{2}x^2 \arctan(x)^2 - x \arctan(x) \\
 & \quad \downarrow \text{240} \\
 & \int \frac{\arctan(x)}{x^2 + 1} dx + \frac{1}{2}x^2 \arctan(x)^2 - x \arctan(x) + \frac{1}{2} \log(x^2 + 1) \\
 & \quad \downarrow \text{5419} \\
 & \frac{1}{2}x^2 \arctan(x)^2 + \frac{\arctan(x)^2}{2} - x \arctan(x) + \frac{1}{2} \log(x^2 + 1)
 \end{aligned}$$

input

 $\text{Int}[x \cdot \text{ArcTan}[x]^2, x]$

output

 $-(x \cdot \text{ArcTan}[x]) + \text{ArcTan}[x]^2/2 + (x^2 \cdot \text{ArcTan}[x]^2)/2 + \text{Log}[1 + x^2]/2$

Definitions of rubi rules used

rule 240 $\text{Int}[(x_+)/((a_) + (b_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x^2, x]]/(2*b), x] /; \text{FreeQ}[\{a, b\}, x]$

rule 5345 $\text{Int}[(a_) + \text{ArcTan}[(c_)*(x_)^(n_)]*(b_)]^(p_), x_Symbol] \rightarrow \text{Simp}[x*(a + b*\text{ArcTan}[c*x^n])^p, x] - \text{Simp}[b*c*n*p \text{ Int}[x^n*((a + b*\text{ArcTan}[c*x^n])^(p-1)/(1 + c^2*x^(2*n))), x], x] /; \text{FreeQ}[\{a, b, c, n\}, x] \&\& \text{IGtQ}[p, 0] \&\& (\text{EqQ}[n, 1] \parallel \text{EqQ}[p, 1])$

rule 5361 $\text{Int}[(a_) + \text{ArcTan}[(c_)*(x_)^(n_)]*(b_)]^(p_)*(x_)^(m_), x_Symbol] \rightarrow \text{Simp}[x^(m+1)*((a + b*\text{ArcTan}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \text{ Int}[x^(m+n)*((a + b*\text{ArcTan}[c*x^n])^(p-1)/(1 + c^2*x^(2*n))), x], x] /; \text{FreeQ}[\{a, b, c, m, n\}, x] \&\& \text{IGtQ}[p, 0] \&\& (\text{EqQ}[p, 1] \parallel (\text{EqQ}[n, 1] \&\& \text{IntegerQ}[m])) \&\& \text{NeQ}[m, -1]$

rule 5419 $\text{Int}[(a_) + \text{ArcTan}[(c_)*(x_)]*(b_)]^(p_)/((d_) + (e_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTan}[c*x])^(p+1)/(b*c*d*(p+1)), x] /; \text{FreeQ}[\{a, b, c, d, e, p\}, x] \&\& \text{EqQ}[e, c^2*d] \&\& \text{NeQ}[p, -1]$

rule 5451 $\text{Int}[(a_) + \text{ArcTan}[(c_)*(x_)]*(b_)]^(p_)*((f_)*(x_)^(m_))/((d_) + (e_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[f^2/e \text{ Int}[(f*x)^(m-2)*(a + b*\text{ArcTan}[c*x])^p, x], x] - \text{Simp}[d*(f^2/e) \text{ Int}[(f*x)^(m-2)*((a + b*\text{ArcTan}[c*x])^p/(d + e*x^2)), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{GtQ}[p, 0] \&\& \text{GtQ}[m, 1]$

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.86

method	result
default	$-x \arctan(x) + \frac{\arctan(x)^2}{2} + \frac{x^2 \arctan(x)^2}{2} + \frac{\ln(x^2+1)}{2}$
parallelrisc	$-x \arctan(x) + \frac{\arctan(x)^2}{2} + \frac{x^2 \arctan(x)^2}{2} + \frac{\ln(x^2+1)}{2}$
parts	$-x \arctan(x) + \frac{\arctan(x)^2}{2} + \frac{x^2 \arctan(x)^2}{2} + \frac{\ln(x^2+1)}{2}$
risc	$-\frac{\left(\frac{x^2}{2} + \frac{1}{2}\right) \ln(ix+1)^2}{4} - \frac{(-x^2 \ln(-ix+1) - 2ix - \ln(-ix+1)) \ln(ix+1)}{4} - \frac{x^2 \ln(-ix+1)^2}{8} - \frac{\ln(-ix+1)^2}{8} - \frac{ix \ln(-ix+1)}{2}$

input `int(x*arctan(x)^2,x,method=_RETURNVERBOSE)`output `-x*arctan(x)+1/2*arctan(x)^2+1/2*x^2*arctan(x)^2+1/2*ln(x^2+1)`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.71

$$\int x \arctan(x)^2 dx = \frac{1}{2} (x^2 + 1) \arctan(x)^2 - x \arctan(x) + \frac{1}{2} \log(x^2 + 1)$$

input `integrate(x*arctan(x)^2,x, algorithm="fricas")`output `1/2*(x^2 + 1)*arctan(x)^2 - x*arctan(x) + 1/2*log(x^2 + 1)`**Sympy [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.83

$$\int x \arctan(x)^2 dx = \frac{x^2 \operatorname{atan}^2(x)}{2} - x \operatorname{atan}(x) + \frac{\log(x^2 + 1)}{2} + \frac{\operatorname{atan}^2(x)}{2}$$

input `integrate(x*atan(x)**2,x)`

output `x**2*atan(x)**2/2 - x*atan(x) + log(x**2 + 1)/2 + atan(x)**2/2`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.97

$$\int x \arctan(x)^2 dx = \frac{1}{2} x^2 \arctan(x)^2 - (x - \arctan(x)) \arctan(x) - \frac{1}{2} \arctan(x)^2 + \frac{1}{2} \log(x^2 + 1)$$

input `integrate(x*arctan(x)^2,x, algorithm="maxima")`

output `1/2*x^2*arctan(x)^2 - (x - arctan(x))*arctan(x) - 1/2*arctan(x)^2 + 1/2*log(x^2 + 1)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.83

$$\int x \arctan(x)^2 dx = \frac{1}{2} x^2 \arctan(x)^2 - x \arctan(x) + \frac{1}{2} \arctan(x)^2 + \frac{1}{2} \log(x^2 + 1)$$

input `integrate(x*arctan(x)^2,x, algorithm="giac")`

output `1/2*x^2*arctan(x)^2 - x*arctan(x) + 1/2*arctan(x)^2 + 1/2*log(x^2 + 1)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.83

$$\int x \arctan(x)^2 dx = \frac{\ln(x^2 + 1)}{2} + \frac{\operatorname{atan}(x)^2}{2} + \frac{x^2 \operatorname{atan}(x)^2}{2} - x \operatorname{atan}(x)$$

input `int(x*atan(x)^2,x)`output `log(x^2 + 1)/2 + atan(x)^2/2 + (x^2*atan(x)^2)/2 - x*atan(x)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.83

$$\int x \arctan(x)^2 dx = \frac{\operatorname{atan}(x)^2 x^2}{2} + \frac{\operatorname{atan}(x)^2}{2} - \operatorname{atan}(x) x + \frac{\log(x^2 + 1)}{2}$$

input `int(x*atan(x)^2,x)`output `(atan(x)**2*x**2 + atan(x)**2 - 2*atan(x)*x + log(x**2 + 1))/2`

3.95 $\int \arctan(\sqrt{x}) dx$

Optimal result	571
Mathematica [A] (verified)	571
Rubi [A] (verified)	572
Maple [A] (verified)	573
Fricas [A] (verification not implemented)	574
Sympy [A] (verification not implemented)	574
Maxima [A] (verification not implemented)	575
Giac [A] (verification not implemented)	575
Mupad [B] (verification not implemented)	575
Reduce [B] (verification not implemented)	576

Optimal result

Integrand size = 6, antiderivative size = 22

$$\int \arctan(\sqrt{x}) dx = -\sqrt{x} + \arctan(\sqrt{x}) + x \arctan(\sqrt{x})$$

output `arctan(x^(1/2))+x*arctan(x^(1/2))-x^(1/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \arctan(\sqrt{x}) dx = -\sqrt{x} + (1+x) \arctan(\sqrt{x})$$

input `Integrate[ArcTan[Sqrt[x]],x]`

output `-Sqrt[x] + (1+x)*ArcTan[Sqrt[x]]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.32, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {5345, 60, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \arctan(\sqrt{x}) \, dx \\
 & \quad \downarrow \text{5345} \\
 & x \arctan(\sqrt{x}) - \frac{1}{2} \int \frac{\sqrt{x}}{x+1} \, dx \\
 & \quad \downarrow \text{60} \\
 & \frac{1}{2} \left(\int \frac{1}{\sqrt{x}(x+1)} \, dx - 2\sqrt{x} \right) + x \arctan(\sqrt{x}) \\
 & \quad \downarrow \text{73} \\
 & \frac{1}{2} \left(2 \int \frac{1}{x+1} \, d\sqrt{x} - 2\sqrt{x} \right) + x \arctan(\sqrt{x}) \\
 & \quad \downarrow \text{216} \\
 & x \arctan(\sqrt{x}) + \frac{1}{2} (2 \arctan(\sqrt{x}) - 2\sqrt{x})
 \end{aligned}$$

input `Int[ArcTan[Sqrt[x]],x]`

output `x*ArcTan[Sqrt[x]] + (-2*Sqrt[x] + 2*ArcTan[Sqrt[x]])/2`

Defintions of rubi rules used

rule 60

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(
b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!Integer
Q[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinear
Q[a, b, c, d, m, n, x]
```

rule 73

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

rule 216

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

rule 5345

```
Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.), x_Symbol] := Simp[x*(a
+ b*ArcTan[c*x^n])^p, x] - Simp[b*c*n*p Int[x^n*((a + b*ArcTan[c*x^n])^(p
- 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] &&
(EqQ[n, 1] || EqQ[p, 1])
```

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

method	result	size
derivativedivides	$\arctan(\sqrt{x}) + x \arctan(\sqrt{x}) - \sqrt{x}$	17
default	$\arctan(\sqrt{x}) + x \arctan(\sqrt{x}) - \sqrt{x}$	17
parts	$\arctan(\sqrt{x}) + x \arctan(\sqrt{x}) - \sqrt{x}$	17
meijerg	$-\sqrt{x} + \frac{(3x+3)\arctan(\sqrt{x})}{3}$	18

input `int(arctan(x^(1/2)),x,method=_RETURNVERBOSE)`

output `arctan(x^(1/2))+x*arctan(x^(1/2))-x^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.64

$$\int \arctan(\sqrt{x}) dx = (x + 1) \arctan(\sqrt{x}) - \sqrt{x}$$

input `integrate(arctan(x^(1/2)),x, algorithm="fricas")`

output `(x + 1)*arctan(sqrt(x)) - sqrt(x)`

Sympy [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.86

$$\int \arctan(\sqrt{x}) dx = -\sqrt{x} + x \operatorname{atan}(\sqrt{x}) + \operatorname{atan}(\sqrt{x})$$

input `integrate(atan(x**(1/2)),x)`

output `-sqrt(x) + x*atan(sqrt(x)) + atan(sqrt(x))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \arctan(\sqrt{x}) dx = x \arctan(\sqrt{x}) - \sqrt{x} + \arctan(\sqrt{x})$$

input `integrate(arctan(x^(1/2)),x, algorithm="maxima")`output `x*arctan(sqrt(x)) - sqrt(x) + arctan(sqrt(x))`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \arctan(\sqrt{x}) dx = x \arctan(\sqrt{x}) - \sqrt{x} + \arctan(\sqrt{x})$$

input `integrate(arctan(x^(1/2)),x, algorithm="giac")`output `x*arctan(sqrt(x)) - sqrt(x) + arctan(sqrt(x))`**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \arctan(\sqrt{x}) dx = \operatorname{atan}(\sqrt{x}) + x \operatorname{atan}(\sqrt{x}) - \sqrt{x}$$

input `int(atan(x^(1/2)),x)`output `atan(x^(1/2)) + x*atan(x^(1/2)) - x^(1/2)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.59

$$\int \arctan(\sqrt{x}) dx = \operatorname{atan}(\sqrt{x}) x + \operatorname{atan}(\sqrt{x}) - \sqrt{x}$$

input `int(atan(x^(1/2)),x)`

output `atan(sqrt(x))*x + atan(sqrt(x)) - sqrt(x)`

3.96 $\int \frac{\arctan(\sqrt{x})}{\sqrt{x}(1+x)} dx$

Optimal result	577
Mathematica [A] (verified)	577
Rubi [A] (verified)	578
Maple [A] (verified)	578
Fricas [A] (verification not implemented)	579
Sympy [A] (verification not implemented)	579
Maxima [A] (verification not implemented)	579
Giac [A] (verification not implemented)	580
Mupad [B] (verification not implemented)	580
Reduce [B] (verification not implemented)	580

Optimal result

Integrand size = 17, antiderivative size = 8

$$\int \frac{\arctan(\sqrt{x})}{\sqrt{x}(1+x)} dx = \arctan(\sqrt{x})^2$$

output

```
arctan(x^(1/2))^2
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(\sqrt{x})}{\sqrt{x}(1+x)} dx = \arctan(\sqrt{x})^2$$

input

```
Integrate[ArcTan[Sqrt[x]]/(Sqrt[x]*(1+x)),x]
```

output

```
ArcTan[Sqrt[x]]^2
```

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {7237}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(\sqrt{x})}{\sqrt{x}(x+1)} dx$$

↓ 7237

$$\arctan(\sqrt{x})^2$$

input `Int[ArcTan[Sqrt[x]]/(Sqrt[x]*(1+x)),x]`

output `ArcTan[Sqrt[x]]^2`

Defintions of rubi rules used

rule 7237 `Int[(u_)*(y_)^(m_.), x_Symbol] := With[{q = DerivativeDivides[y, u, x]}, Simp[q*(y^(m+1)/(m+1)), x] /; !FalseQ[q] /; FreeQ[m, x] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$\arctan(\sqrt{x})^2$	7
default	$\arctan(\sqrt{x})^2$	7

input `int(arctan(x^(1/2))/(1+x)/x^(1/2),x,method=_RETURNVERBOSE)`

output `arctan(x^(1/2))^2`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\arctan(\sqrt{x})}{\sqrt{x}(1+x)} dx = \arctan(\sqrt{x})^2$$

input `integrate(arctan(x^(1/2))/(1+x)/x^(1/2),x, algorithm="fricas")`

output `arctan(sqrt(x))^2`

Sympy [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int \frac{\arctan(\sqrt{x})}{\sqrt{x}(1+x)} dx = \text{atan}^2(\sqrt{x})$$

input `integrate(atan(x**(1/2))/(1+x)/x**(1/2),x)`

output `atan(sqrt(x))**2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\arctan(\sqrt{x})}{\sqrt{x}(1+x)} dx = \arctan(\sqrt{x})^2$$

input `integrate(arctan(x^(1/2))/(1+x)/x^(1/2),x, algorithm="maxima")`

output `arctan(sqrt(x))^2`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\arctan(\sqrt{x})}{\sqrt{x}(1+x)} dx = \arctan(\sqrt{x})^2$$

input `integrate(arctan(x^(1/2))/(1+x)/x^(1/2),x, algorithm="giac")`

output `arctan(sqrt(x))^2`

Mupad [B] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\arctan(\sqrt{x})}{\sqrt{x}(1+x)} dx = \operatorname{atan}(\sqrt{x})^2$$

input `int(atan(x^(1/2))/(x^(1/2)*(x + 1)),x)`

output `atan(x^(1/2))^2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int \frac{\arctan(\sqrt{x})}{\sqrt{x}(1+x)} dx = \operatorname{atan}(\sqrt{x})^2$$

input `int(atan(x^(1/2))/(1+x)/x^(1/2),x)`

output `atan(sqrt(x))**2`

3.97 $\int \sqrt{1-x^2} dx$

Optimal result	582
Mathematica [A] (verified)	582
Rubi [A] (verified)	583
Maple [A] (verified)	584
Fricas [A] (verification not implemented)	584
Sympy [A] (verification not implemented)	585
Maxima [A] (verification not implemented)	585
Giac [A] (verification not implemented)	585
Mupad [B] (verification not implemented)	586
Reduce [B] (verification not implemented)	586

Optimal result

Integrand size = 11, antiderivative size = 23

$$\int \sqrt{1-x^2} dx = \frac{1}{2}x\sqrt{1-x^2} + \frac{\arcsin(x)}{2}$$

output `1/2*arcsin(x)+1/2*x*(-x^2+1)^(1/2)`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.61

$$\int \sqrt{1-x^2} dx = \frac{1}{2}x\sqrt{1-x^2} - \arctan\left(\frac{\sqrt{1-x^2}}{1+x}\right)$$

input `Integrate[Sqrt[1 - x^2],x]`

output `(x*Sqrt[1 - x^2])/2 - ArcTan[Sqrt[1 - x^2]/(1 + x)]`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {211, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{1-x^2} dx$$

$$\downarrow \text{211}$$

$$\frac{1}{2} \int \frac{1}{\sqrt{1-x^2}} dx + \frac{1}{2} \sqrt{1-x^2} x$$

$$\downarrow \text{223}$$

$$\frac{\arcsin(x)}{2} + \frac{1}{2} \sqrt{1-x^2} x$$

input `Int[Sqrt[1 - x^2], x]`

output `(x*Sqrt[1 - x^2])/2 + ArcSin[x]/2`

Defintions of rubi rules used

rule 211 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[x*((a + b*x^2)^p/(2*p + 1)), x] + Simp[2*a*(p/(2*p + 1)) Int[(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

method	result	size
default	$\frac{\arcsin(x)}{2} + \frac{x\sqrt{-x^2+1}}{2}$	18
risch	$-\frac{x(x^2-1)}{2\sqrt{-x^2+1}} + \frac{\arcsin(x)}{2}$	23
pseudoelliptic	$\frac{x\sqrt{-x^2+1}}{2} - \frac{\arctan\left(\frac{\sqrt{-x^2+1}}{x}\right)}{2}$	30
meijerg	$\frac{i(-2i\sqrt{\pi}x\sqrt{-x^2+1}-2i\sqrt{\pi}\arcsin(x))}{4\sqrt{\pi}}$	32
trager	$\frac{x\sqrt{-x^2+1}}{2} + \frac{\text{RootOf}(-Z^2+1)\ln(\text{RootOf}(-Z^2+1)\sqrt{-x^2+1}+x)}{2}$	41

input `int((-x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output `1/2*arcsin(x)+1/2*x*(-x^2+1)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.35

$$\int \sqrt{1-x^2} dx = \frac{1}{2} \sqrt{-x^2+1} x - \arctan\left(\frac{\sqrt{-x^2+1}-1}{x}\right)$$

input `integrate((-x^2+1)^(1/2),x, algorithm="fricas")`

output `1/2*sqrt(-x^2 + 1)*x - arctan((sqrt(-x^2 + 1) - 1)/x)`

Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int \sqrt{1-x^2} dx = \frac{x\sqrt{1-x^2}}{2} + \frac{\arcsin(x)}{2}$$

input `integrate((-x**2+1)**(1/2),x)`output `x*sqrt(1 - x**2)/2 + asin(x)/2`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \sqrt{1-x^2} dx = \frac{1}{2} \sqrt{-x^2+1}x + \frac{1}{2} \arcsin(x)$$

input `integrate((-x^2+1)^(1/2),x, algorithm="maxima")`output `1/2*sqrt(-x^2 + 1)*x + 1/2*arcsin(x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \sqrt{1-x^2} dx = \frac{1}{2} \sqrt{-x^2+1}x + \frac{1}{2} \arcsin(x)$$

input `integrate((-x^2+1)^(1/2),x, algorithm="giac")`output `1/2*sqrt(-x^2 + 1)*x + 1/2*arcsin(x)`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \sqrt{1-x^2} dx = \frac{\operatorname{asin}(x)}{2} + \frac{x\sqrt{1-x^2}}{2}$$

input `int((1 - x^2)^(1/2),x)`

output `asin(x)/2 + (x*(1 - x^2)^(1/2))/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.70

$$\int \sqrt{1-x^2} dx = \frac{\operatorname{asin}(x)}{2} + \frac{\sqrt{-x^2+1}x}{2}$$

input `int((-x^2+1)^(1/2),x)`

output `(asin(x) + sqrt(-x**2 + 1)*x)/2`

$$3.98 \quad \int \frac{e^{\arctan(x)} x}{(1+x^2)^{3/2}} dx$$

Optimal result	587
Mathematica [C] (verified)	587
Rubi [A] (verified)	588
Maple [A] (verified)	588
Fricas [A] (verification not implemented)	589
Sympy [A] (verification not implemented)	589
Maxima [F]	590
Giac [A] (verification not implemented)	590
Mupad [F(-1)]	590
Reduce [B] (verification not implemented)	591

Optimal result

Integrand size = 15, antiderivative size = 22

$$\int \frac{e^{\arctan(x)} x}{(1+x^2)^{3/2}} dx = -\frac{e^{\arctan(x)}(1-x)}{2\sqrt{1+x^2}}$$

output `-1/2*exp(arctan(x))*(1-x)/(x^2+1)^(1/2)`

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.01 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.68

$$\int \frac{e^{\arctan(x)} x}{(1+x^2)^{3/2}} dx = \frac{1}{2}(1-ix)^{-\frac{1}{2}+\frac{i}{2}}(1+ix)^{-\frac{1}{2}-\frac{i}{2}}(-1+x)$$

input `Integrate[(E^ArcTan[x]*x)/(1+x^2)^(3/2),x]`

output `(-1+x)/(2*(1-I*x)^(1/2-I/2)*(1+I*x)^(1/2+I/2))`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$, Rules used = {5600}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x e^{\arctan(x)}}{(x^2 + 1)^{3/2}} dx$$

↓ 5600

$$-\frac{(1-x)e^{\arctan(x)}}{2\sqrt{x^2+1}}$$

input `Int[(E^ArcTan[x]*x)/(1 + x^2)^(3/2), x]`

output `-1/2*(E^ArcTan[x]*(1 - x))/Sqrt[1 + x^2]`

Defintions of rubi rules used

rule 5600

```
Int[(E^(ArcTan[(a_.)*(x_)])*(n_.))*(x_))/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol]
:> Simp[(-(1 - a*n*x))*(E^(n*ArcTan[a*x])/(d*(n^2 + 1)*Sqrt[c + d*x^2]))
, x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && !IntegerQ[I*n]
```

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

method	result	size
gospers	$\frac{(-1+x)e^{\arctan(x)}}{2\sqrt{x^2+1}}$	16
orering	$\frac{(-1+x)e^{\arctan(x)}}{2\sqrt{x^2+1}}$	16

input `int(exp(arctan(x))*x/(x^2+1)^(3/2),x,method=_RETURNVERBOSE)`

output `1/2*(-1+x)*exp(arctan(x))/(x^2+1)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.68

$$\int \frac{e^{\arctan(x)}x}{(1+x^2)^{3/2}} dx = \frac{(x-1)e^{\arctan(x)}}{2\sqrt{x^2+1}}$$

input `integrate(exp(arctan(x))*x/(x^2+1)^(3/2),x, algorithm="fricas")`

output `1/2*(x - 1)*e^arctan(x)/sqrt(x^2 + 1)`

Sympy [A] (verification not implemented)

Time = 8.34 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.41

$$\int \frac{e^{\arctan(x)}x}{(1+x^2)^{3/2}} dx = \frac{xe^{\arctan(x)}}{2\sqrt{x^2+1}} - \frac{e^{\arctan(x)}}{2\sqrt{x^2+1}}$$

input `integrate(exp(atan(x))*x/(x**2+1)**(3/2),x)`

output `x*exp(atan(x))/(2*sqrt(x**2 + 1)) - exp(atan(x))/(2*sqrt(x**2 + 1))`

Maxima [F]

$$\int \frac{e^{\arctan(x)} x}{(1+x^2)^{3/2}} dx = \int \frac{x e^{\arctan(x)}}{(x^2+1)^{3/2}} dx$$

input `integrate(exp(arctan(x))*x/(x^2+1)^(3/2),x, algorithm="maxima")`

output `integrate(x*e^arctan(x)/(x^2 + 1)^(3/2), x)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{e^{\arctan(x)} x}{(1+x^2)^{3/2}} dx = \frac{1}{2} \left(\frac{x}{\sqrt{x^2+1}} - \frac{1}{\sqrt{x^2+1}} \right) e^{\arctan(x)}$$

input `integrate(exp(arctan(x))*x/(x^2+1)^(3/2),x, algorithm="giac")`

output `1/2*(x/sqrt(x^2 + 1) - 1/sqrt(x^2 + 1))*e^arctan(x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{e^{\arctan(x)} x}{(1+x^2)^{3/2}} dx = \int \frac{x e^{\arctan(x)}}{(x^2+1)^{3/2}} dx$$

input `int((x*exp(atan(x)))/(x^2 + 1)^(3/2),x)`

output `int((x*exp(atan(x)))/(x^2 + 1)^(3/2), x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.05

$$\int \frac{e^{\arctan(x)} x}{(1+x^2)^{3/2}} dx = \frac{e^{\arctan(x)} \sqrt{x^2+1} (x-1)}{2x^2+2}$$

input `int(exp(atan(x))*x/(x^2+1)^(3/2),x)`

output `(e**atan(x)*sqrt(x**2 + 1)*(x - 1))/(2*(x**2 + 1))`

3.99 $\int \frac{e^{\arctan(x)}}{(1+x^2)^{3/2}} dx$

Optimal result	592
Mathematica [A] (verified)	592
Rubi [A] (verified)	593
Maple [A] (verified)	593
Fricas [A] (verification not implemented)	594
Sympy [A] (verification not implemented)	594
Maxima [F]	595
Giac [A] (verification not implemented)	595
Mupad [F(-1)]	595
Reduce [B] (verification not implemented)	596

Optimal result

Integrand size = 14, antiderivative size = 20

$$\int \frac{e^{\arctan(x)}}{(1+x^2)^{3/2}} dx = \frac{e^{\arctan(x)}(1+x)}{2\sqrt{1+x^2}}$$

output `1/2*exp(arctan(x))*(1+x)/(x^2+1)^(1/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{e^{\arctan(x)}}{(1+x^2)^{3/2}} dx = \frac{e^{\arctan(x)}(1+x)}{2\sqrt{1+x^2}}$$

input `Integrate[E^ArcTan[x]/(1 + x^2)^(3/2), x]`

output `(E^ArcTan[x]*(1 + x))/(2*Sqrt[1 + x^2])`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$, Rules used = {5592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^{\arctan(x)}}{(x^2 + 1)^{3/2}} dx$$

↓ 5592

$$\frac{(x + 1)e^{\arctan(x)}}{2\sqrt{x^2 + 1}}$$

input

```
Int [E^ArcTan[x]/(1 + x^2)^(3/2), x]
```

output

```
(E^ArcTan[x]*(1 + x))/(2*Sqrt[1 + x^2])
```

Defintions of rubi rules used

rule 5592

```
Int [E^(ArcTan[(a_)*(x_)]*(n_.))/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] :>
Simp[(n + a*x)*(E^(n*ArcTan[a*x]))/(a*c*(n^2 + 1)*Sqrt[c + d*x^2]), x] /; F
reeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && !IntegerQ[I*n]
```

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

method	result	size
gospers	$\frac{e^{\arctan(x)}(1+x)}{2\sqrt{x^2+1}}$	16
orering	$\frac{e^{\arctan(x)}(1+x)}{2\sqrt{x^2+1}}$	16

input `int(exp(arctan(x))/(x^2+1)^(3/2),x,method=_RETURNVERBOSE)`

output `1/2*exp(arctan(x))*(1+x)/(x^2+1)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

$$\int \frac{e^{\arctan(x)}}{(1+x^2)^{3/2}} dx = \frac{(x+1)e^{\arctan(x)}}{2\sqrt{x^2+1}}$$

input `integrate(exp(arctan(x))/(x^2+1)^(3/2),x, algorithm="fricas")`

output `1/2*(x + 1)*e^arctan(x)/sqrt(x^2 + 1)`

Sympy [A] (verification not implemented)

Time = 8.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.55

$$\int \frac{e^{\arctan(x)}}{(1+x^2)^{3/2}} dx = \frac{x e^{\arctan(x)}}{2\sqrt{x^2+1}} + \frac{e^{\arctan(x)}}{2\sqrt{x^2+1}}$$

input `integrate(exp(atan(x))/(x**2+1)**(3/2),x)`

output `x*exp(atan(x))/(2*sqrt(x**2 + 1)) + exp(atan(x))/(2*sqrt(x**2 + 1))`

Maxima [F]

$$\int \frac{e^{\arctan(x)}}{(1+x^2)^{3/2}} dx = \int \frac{e^{\arctan(x)}}{(x^2+1)^{3/2}} dx$$

input `integrate(exp(arctan(x))/(x^2+1)^(3/2),x, algorithm="maxima")`

output `integrate(e^arctan(x)/(x^2 + 1)^(3/2), x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{e^{\arctan(x)}}{(1+x^2)^{3/2}} dx = \frac{1}{2} \left(\frac{x}{\sqrt{x^2+1}} + \frac{1}{\sqrt{x^2+1}} \right) e^{\arctan(x)}$$

input `integrate(exp(arctan(x))/(x^2+1)^(3/2),x, algorithm="giac")`

output `1/2*(x/sqrt(x^2 + 1) + 1/sqrt(x^2 + 1))*e^arctan(x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{e^{\arctan(x)}}{(1+x^2)^{3/2}} dx = \int \frac{e^{\operatorname{atan}(x)}}{(x^2+1)^{3/2}} dx$$

input `int(exp(atan(x))/(x^2 + 1)^(3/2),x)`

output `int(exp(atan(x))/(x^2 + 1)^(3/2), x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.15

$$\int \frac{e^{\arctan(x)}}{(1+x^2)^{3/2}} dx = \frac{e^{\arctan(x)} \sqrt{x^2+1} (x+1)}{2x^2+2}$$

input `int(exp(atan(x))/(x^2+1)^(3/2),x)`

output `(e**atan(x)*sqrt(x**2 + 1)*(x + 1))/(2*(x**2 + 1))`

3.100 $\int \frac{x^2}{(1+x^2)^2} dx$

Optimal result	597
Mathematica [A] (verified)	597
Rubi [A] (verified)	598
Maple [A] (verified)	599
Fricas [A] (verification not implemented)	599
Sympy [A] (verification not implemented)	599
Maxima [A] (verification not implemented)	600
Giac [A] (verification not implemented)	600
Mupad [B] (verification not implemented)	600
Reduce [B] (verification not implemented)	601

Optimal result

Integrand size = 11, antiderivative size = 19

$$\int \frac{x^2}{(1+x^2)^2} dx = -\frac{x}{2(1+x^2)} + \frac{\arctan(x)}{2}$$

output `-1/2*x/(x^2+1)+1/2*arctan(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{x^2}{(1+x^2)^2} dx = -\frac{x}{2(1+x^2)} + \frac{\arctan(x)}{2}$$

input `Integrate[x^2/(1 + x^2)^2,x]`

output `-1/2*x/(1 + x^2) + ArcTan[x]/2`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {252, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{(x^2 + 1)^2} dx$$

↓ 252

$$\frac{1}{2} \int \frac{1}{x^2 + 1} dx - \frac{x}{2(x^2 + 1)}$$

↓ 216

$$\frac{\arctan(x)}{2} - \frac{x}{2(x^2 + 1)}$$

input `Int[x^2/(1 + x^2)^2,x]`

output `-1/2*x/(1 + x^2) + ArcTan[x]/2`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 252 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(2*b*(p + 1))), x] - Simp[c^2*(m - 1)/(2*b*(p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && LtQ[p, -1] && GtQ[m, 1] && !ILtQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

method	result	size
default	$-\frac{x}{2(x^2+1)} + \frac{\arctan(x)}{2}$	16
meijerg	$-\frac{x}{2(x^2+1)} + \frac{\arctan(x)}{2}$	16
risch	$-\frac{x}{2(x^2+1)} + \frac{\arctan(x)}{2}$	16
parallelrisch	$-\frac{i \ln(x-i)x^2 - i \ln(x+i)x^2 + i \ln(x-i) - i \ln(x+i) + 2x}{4(x^2+1)}$	52

input `int(x^2/(x^2+1)^2,x,method=_RETURNVERBOSE)`output `-1/2*x/(x^2+1)+1/2*arctan(x)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{x^2}{(1+x^2)^2} dx = \frac{(x^2+1)\arctan(x) - x}{2(x^2+1)}$$

input `integrate(x^2/(x^2+1)^2,x, algorithm="fricas")`output `1/2*((x^2 + 1)*arctan(x) - x)/(x^2 + 1)`**Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int \frac{x^2}{(1+x^2)^2} dx = -\frac{x}{2x^2+2} + \frac{\operatorname{atan}(x)}{2}$$

input `integrate(x**2/(x**2+1)**2,x)`

output `-x/(2*x**2 + 2) + atan(x)/2`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{x^2}{(1+x^2)^2} dx = -\frac{x}{2(x^2+1)} + \frac{1}{2} \arctan(x)$$

input `integrate(x^2/(x^2+1)^2,x, algorithm="maxima")`

output `-1/2*x/(x^2 + 1) + 1/2*arctan(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{x^2}{(1+x^2)^2} dx = -\frac{x}{2(x^2+1)} + \frac{1}{2} \arctan(x)$$

input `integrate(x^2/(x^2+1)^2,x, algorithm="giac")`

output `-1/2*x/(x^2 + 1) + 1/2*arctan(x)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{x^2}{(1+x^2)^2} dx = \frac{\operatorname{atan}(x)}{2} - \frac{x}{2(x^2+1)}$$

input `int(x^2/(x^2 + 1)^2,x)`

output `atan(x)/2 - x/(2*(x^2 + 1))`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.16

$$\int \frac{x^2}{(1+x^2)^2} dx = \frac{\operatorname{atan}(x) x^2 + \operatorname{atan}(x) - x}{2x^2 + 2}$$

input `int(x^2/(x^2+1)^2,x)`

output `(atan(x)*x**2 + atan(x) - x)/(2*(x**2 + 1))`

3.101 $\int \frac{e^x}{1+e^{2x}} dx$

Optimal result	602
Mathematica [A] (verified)	602
Rubi [A] (verified)	603
Maple [A] (verified)	604
Fricas [A] (verification not implemented)	604
Sympy [B] (verification not implemented)	604
Maxima [A] (verification not implemented)	605
Giac [A] (verification not implemented)	605
Mupad [B] (verification not implemented)	605
Reduce [B] (verification not implemented)	606

Optimal result

Integrand size = 13, antiderivative size = 4

$$\int \frac{e^x}{1+e^{2x}} dx = \arctan(e^x)$$

output `arctan(exp(x))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{e^x}{1+e^{2x}} dx = \arctan(e^x)$$

input `Integrate[E^x/(1 + E^(2*x)),x]`

output `ArcTan[E^x]`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {2679, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^x}{e^{2x} + 1} dx$$

↓ 2679

$$\int \frac{1}{e^{2x} + 1} de^x$$

↓ 216

$$\arctan(e^x)$$

input `Int[E^x/(1 + E^(2*x)), x]`

output `ArcTan[E^x]`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 2679 `Int[((a_) + (b_.)*(F_)^((e_.)*((c_.) + (d_.)*(x_))))^(p_.)*(G_)^((h_.)*((f_.) + (g_.)*(x_))), x_Symbol] := With[{m = FullSimplify[d*e*(Log[F]/(g*h*Log[G]))}], Simp[Denominator[m]/(g*h*Log[G]) Subst[Int[x^(Denominator[m] - 1)*(a + b*F^(c*e - d*e*(f/g))*x^Numerator[m])^p, x], x, G^(h*((f + g*x)/Denominator[m]))], x] /; LtQ[m, -1] || GtQ[m, 1] /; FreeQ[{F, G, a, b, c, d, e, f, g, h, p}, x]`

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

method	result	size
default	$\arctan(e^x)$	4
risch	$\frac{i \ln(e^x+i)}{2} - \frac{i \ln(e^x-i)}{2}$	20

input `int(exp(x)/(1+exp(2*x)),x,method=_RETURNVERBOSE)`

output `arctan(exp(x))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int \frac{e^x}{1+e^{2x}} dx = \arctan(e^x)$$

input `integrate(exp(x)/(1+exp(2*x)),x, algorithm="fricas")`

output `arctan(e^x)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(3) = 6$.

Time = 0.05 (sec) , antiderivative size = 15, normalized size of antiderivative = 3.75

$$\int \frac{e^x}{1+e^{2x}} dx = \text{RootSum}(4z^2 + 1, (i \mapsto i \log(2i + e^x)))$$

input `integrate(exp(x)/(1+exp(2*x)),x)`

output `RootSum(4*_z**2 + 1, Lambda(_i, _i*log(2*_i + exp(x))))`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int \frac{e^x}{1 + e^{2x}} dx = \arctan(e^x)$$

input `integrate(exp(x)/(1+exp(2*x)),x, algorithm="maxima")`output `arctan(e^x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int \frac{e^x}{1 + e^{2x}} dx = \arctan(e^x)$$

input `integrate(exp(x)/(1+exp(2*x)),x, algorithm="giac")`output `arctan(e^x)`**Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int \frac{e^x}{1 + e^{2x}} dx = \operatorname{atan}(e^x)$$

input `int(exp(x)/(exp(2*x) + 1),x)`output `atan(exp(x))`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{e^x}{1 + e^{2x}} dx = \operatorname{atan}(e^x)$$

input `int(exp(x)/(1+exp(2*x)),x)`

output `atan(e**x)`

3.102 $\int e^{-x} \cot^{-1}(e^x) dx$

Optimal result	607
Mathematica [A] (verified)	607
Rubi [A] (verified)	608
Maple [A] (verified)	610
Fricas [A] (verification not implemented)	610
Sympy [A] (verification not implemented)	610
Maxima [A] (verification not implemented)	611
Giac [A] (verification not implemented)	611
Mupad [B] (verification not implemented)	611
Reduce [B] (verification not implemented)	612

Optimal result

Integrand size = 10, antiderivative size = 27

$$\int e^{-x} \cot^{-1}(e^x) dx = -x - e^{-x} \cot^{-1}(e^x) + \frac{1}{2} \log(1 + e^{2x})$$

output `-x-arccot(exp(x))/exp(x)+1/2*ln(1+exp(2*x))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int e^{-x} \cot^{-1}(e^x) dx = -x - e^{-x} \cot^{-1}(e^x) + \frac{1}{2} \log(1 + e^{2x})$$

input `Integrate[ArcCot[E^x]/E^x,x]`

output `-x - ArcCot[E^x]/E^x + Log[1 + E^(2*x)]/2`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.22, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {5731, 25, 2720, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{-x} \cot^{-1}(e^x) dx \\
 & \quad \downarrow \text{5731} \\
 & \int -\frac{1}{1+e^{2x}} dx - e^{-x} \cot^{-1}(e^x) \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{1}{1+e^{2x}} dx - e^{-x} \cot^{-1}(e^x) \\
 & \quad \downarrow \text{2720} \\
 & -\frac{1}{2} \int \frac{e^{-2x}}{1+e^{2x}} de^{2x} - e^{-x} \cot^{-1}(e^x) \\
 & \quad \downarrow \text{47} \\
 & \frac{1}{2} \left(\int \frac{1}{1+e^{2x}} de^{2x} - \int e^{-2x} de^{2x} \right) - e^{-x} \cot^{-1}(e^x) \\
 & \quad \downarrow \text{14} \\
 & \frac{1}{2} \left(\int \frac{1}{1+e^{2x}} de^{2x} - \log(e^{2x}) \right) - e^{-x} \cot^{-1}(e^x) \\
 & \quad \downarrow \text{16} \\
 & \frac{1}{2} (\log(e^{2x} + 1) - \log(e^{2x})) - e^{-x} \cot^{-1}(e^x)
 \end{aligned}$$

input `Int[ArcCot[E^x]/E^x, x]`

output `-(ArcCot[E^x]/E^x) + (-Log[E^(2*x)] + Log[1 + E^(2*x)])/2`

Definitions of rubi rules used

- rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`
- rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`
- rule 5731 `Int[((a_.) + ArcCot[u_]*(b_.))*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Simp[(a + b*ArcCot[u])*w, x] + Simp[b Int[SimplifyIntegrand[w*(D[u, x]/(1 + u^2)), x], x], x] /; InverseFunctionFreeQ[w, x] /; FreeQ[{a, b}, x] && InverseFunctionFreeQ[u, x] && !MatchQ[v, ((c_.) + (d_.)*x)^(m_.)] /; FreeQ[{c, d, m}, x] && FalseQ[FunctionOfLinear[v*(a + b*ArcCot[u]), x]]]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.93

method	result	size
derivativdivides	$-\operatorname{arccot}(e^x)e^{-x} - \ln(e^x) + \frac{\ln(1+e^{2x})}{2}$	25
default	$-\operatorname{arccot}(e^x)e^{-x} - \ln(e^x) + \frac{\ln(1+e^{2x})}{2}$	25
parallelrisch	$\frac{(\ln(1+e^{2x})e^x - 2e^x x - 2\operatorname{arccot}(e^x))e^{-x}}{2}$	28
risch	$-\frac{ie^{-x}\ln(1+ie^x)}{2} + \frac{\ln(1+e^{2x})}{2} - x + \frac{ie^{-x}\ln(1-ie^x)}{2} - \frac{e^{-x}\pi}{2}$	51

input `int(arccot(exp(x))/exp(x),x,method=_RETURNVERBOSE)`output `-arccot(exp(x))/exp(x)-ln(exp(x))+1/2*ln(1+exp(x)^2)`**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.04

$$\int e^{-x} \cot^{-1}(e^x) dx = -\frac{1}{2} (2xe^x - e^x \log(e^{2x} + 1) + 2 \operatorname{arccot}(e^x))e^{-x}$$

input `integrate(arccot(exp(x))/exp(x),x, algorithm="fricas")`output `-1/2*(2*x*e^x - e^x*log(e^(2*x) + 1) + 2*arccot(e^x))*e^(-x)`**Sympy [A] (verification not implemented)**

Time = 1.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int e^{-x} \cot^{-1}(e^x) dx = -x + \frac{\log(e^{2x} + 1)}{2} - e^{-x} \operatorname{acot}(e^x)$$

input `integrate(acot(exp(x))/exp(x),x)`

output `-x + log(exp(2*x) + 1)/2 - exp(-x)*acot(exp(x))`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int e^{-x} \cot^{-1}(e^x) dx = -\operatorname{arccot}(e^x) e^{(-x)} + \frac{1}{2} \log(e^{(-2x)} + 1)$$

input `integrate(arccot(exp(x))/exp(x),x, algorithm="maxima")`

output `-arccot(e^x)*e^(-x) + 1/2*log(e^(-2*x) + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int e^{-x} \cot^{-1}(e^x) dx = -\arctan(e^{(-x)}) e^{(-x)} + \frac{1}{2} \log(e^{(-2x)} + 1)$$

input `integrate(arccot(exp(x))/exp(x),x, algorithm="giac")`

output `-arctan(e^(-x))*e^(-x) + 1/2*log(e^(-2*x) + 1)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int e^{-x} \cot^{-1}(e^x) dx = \frac{\ln(e^{2x} + 1)}{2} - x - \operatorname{acot}(e^x) e^{-x}$$

input `int(acot(exp(x))*exp(-x),x)`

output `log(exp(2*x) + 1)/2 - x - acot(exp(x))*exp(-x)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.19

$$\int e^{-x} \cot^{-1}(e^x) dx = \frac{-2\operatorname{acot}(e^x) + e^x \log(e^{2x} + 1) - 2e^x x}{2e^x}$$

input `int(acot(exp(x))/exp(x),x)`

output `(- 2*acot(e**x) + e**x*log(e**(2*x) + 1) - 2*e**x*x)/(2*e**x)`

3.103 $\int \sqrt{\frac{a+x}{a-x}} dx$

Optimal result	613
Mathematica [A] (verified)	613
Rubi [A] (verified)	614
Maple [A] (verified)	615
Fricas [A] (verification not implemented)	616
Sympy [F]	616
Maxima [A] (verification not implemented)	616
Giac [A] (verification not implemented)	617
Mupad [B] (verification not implemented)	617
Reduce [B] (verification not implemented)	617

Optimal result

Integrand size = 15, antiderivative size = 42

$$\int \sqrt{\frac{a+x}{a-x}} dx = -\left((a-x)\sqrt{\frac{a+x}{a-x}}\right) + 2a \arctan\left(\sqrt{\frac{a+x}{a-x}}\right)$$

output

```
2*a*arctan(((a+x)/(a-x))^(1/2))- (a-x)*((a+x)/(a-x))^(1/2)
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.60

$$\int \sqrt{\frac{a+x}{a-x}} dx = \frac{\sqrt{\frac{a+x}{a-x}} \left((-a+x)\sqrt{a+x} + 2a\sqrt{a-x} \arctan\left(\frac{\sqrt{a+x}}{\sqrt{a-x}}\right) \right)}{\sqrt{a+x}}$$

input

```
Integrate[Sqrt[(a + x)/(a - x)],x]
```

output

```
(Sqrt[(a + x)/(a - x)]*((-a + x)*Sqrt[a + x] + 2*a*Sqrt[a - x]*ArcTan[Sqrt[a + x]/Sqrt[a - x]]))/Sqrt[a + x]
```

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.38, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2051, 252, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\frac{a+x}{a-x}} dx \\
 & \quad \downarrow \text{2051} \\
 & 4a \int \frac{a+x}{(a-x) \left(\frac{a+x}{a-x} + 1\right)^2} d\sqrt{\frac{a+x}{a-x}} \\
 & \quad \downarrow \text{252} \\
 & 4a \left(\frac{1}{2} \int \frac{1}{\frac{a+x}{a-x} + 1} d\sqrt{\frac{a+x}{a-x}} - \frac{\sqrt{\frac{a+x}{a-x}}}{2 \left(\frac{a+x}{a-x} + 1\right)} \right) \\
 & \quad \downarrow \text{216} \\
 & 4a \left(\frac{1}{2} \arctan \left(\sqrt{\frac{a+x}{a-x}} \right) - \frac{\sqrt{\frac{a+x}{a-x}}}{2 \left(\frac{a+x}{a-x} + 1\right)} \right)
 \end{aligned}$$

input `Int[Sqrt[(a + x)/(a - x)],x]`

output `4*a*(-1/2*Sqrt[(a + x)/(a - x)]/(1 + (a + x)/(a - x)) + ArcTan[Sqrt[(a + x)/(a - x)]]/2)`

Definitions of rubi rules used

rule 216 $\text{Int}[\{(a_)+(b_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*ArcTan[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /;$ FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

rule 252 $\text{Int}[\{(c_)*(x_)\}^{(m_)}*\{(a_)+(b_)*(x_)^2\}^{(p_)}, x_Symbol] \rightarrow \text{Simp}[c*(c*x)^{(m-1)}*((a+b*x^2)^{(p+1)}/(2*b*(p+1))), x] - \text{Simp}[c^2*((m-1)/(2*b*(p+1))) \text{Int}[(c*x)^{(m-2)}*(a+b*x^2)^{(p+1)}, x], x] /;$ FreeQ[{a, b, c}, x] && LtQ[p, -1] && GtQ[m, 1] && !ILtQ[(m+2*p+3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]

rule 2051 $\text{Int}[\{(e_)*\{(a_)+(b_)*(x_)^{n_}\}\}/\{(c_)+(d_)*(x_)^{n_}\}^{(p_)}, x_Symbol] \rightarrow \text{With}[\{q = \text{Denominator}[p]\}, \text{Simp}[q*e*((b*c-a*d)/n) \text{Subst}[\text{Int}[x^{(q*(p+1)-1)}*((-a)*e+c*x^q)^{(1/n-1)}/(b*e-d*x^q)^{(1/n+1)}], x], x, (e*((a+b*x^n)/(c+d*x^n)))^{(1/q)}, x]] /;$ FreeQ[{a, b, c, d, e}, x] && FractionQ[p] && IntegerQ[1/n]

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.45

method	result	size
default	$-\frac{\sqrt{\frac{a+x}{a-x}}(a-x)\left(\sqrt{a^2-x^2}-a \arctan\left(\frac{x}{\sqrt{a^2-x^2}}\right)\right)}{\sqrt{(a-x)(a+x)}}$	61
risch	$-\frac{(a-x)\sqrt{\frac{a+x}{a-x}}\sqrt{(a-x)(a+x)}}{\sqrt{-(-a+x)(a+x)}} + \frac{a \arctan\left(\frac{x}{\sqrt{a^2-x^2}}\right)\sqrt{\frac{a+x}{a-x}}\sqrt{(a-x)(a+x)}}{a+x}$	90

input `int(((a+x)/(a-x))^(1/2),x,method=_RETURNVERBOSE)`

output $-\{(a+x)/(a-x)\}^{(1/2)}*(a-x)*\{(a^2-x^2)\}^{(1/2)}-a*\arctan(x/\{(a^2-x^2)\}^{(1/2)})\}/\{(a-x)*(a+x)\}^{(1/2)}$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.90

$$\int \sqrt{\frac{a+x}{a-x}} dx = 2a \arctan\left(\sqrt{\frac{a+x}{a-x}}\right) - (a-x)\sqrt{\frac{a+x}{a-x}}$$

input `integrate(((a+x)/(a-x))^(1/2),x, algorithm="fricas")`output `2*a*arctan(sqrt((a + x)/(a - x))) - (a - x)*sqrt((a + x)/(a - x))`**Sympy [F]**

$$\int \sqrt{\frac{a+x}{a-x}} dx = \int \sqrt{\frac{a+x}{a-x}} dx$$

input `integrate(((a+x)/(a-x))**(1/2),x)`output `Integral(sqrt((a + x)/(a - x)), x)`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.17

$$\int \sqrt{\frac{a+x}{a-x}} dx = -2a \left(\frac{\sqrt{\frac{a+x}{a-x}}}{\frac{a+x}{a-x} + 1} - \arctan\left(\sqrt{\frac{a+x}{a-x}}\right) \right)$$

input `integrate(((a+x)/(a-x))^(1/2),x, algorithm="maxima")`output `-2*a*(sqrt((a + x)/(a - x))/((a + x)/(a - x) + 1) - arctan(sqrt((a + x)/(a - x))))`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.86

$$\int \sqrt{\frac{a+x}{a-x}} dx = a \arcsin\left(\frac{x}{a}\right) \operatorname{sgn}(a-x) \operatorname{sgn}(a) - \sqrt{a^2 - x^2} \operatorname{sgn}(a-x)$$

input `integrate(((a+x)/(a-x))^(1/2),x, algorithm="giac")`output `a*arcsin(x/a)*sgn(a - x)*sgn(a) - sqrt(a^2 - x^2)*sgn(a - x)`**Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.17

$$\int \sqrt{\frac{a+x}{a-x}} dx = 2a \operatorname{atan}\left(\sqrt{\frac{a+x}{a-x}}\right) - \frac{2a \sqrt{\frac{a+x}{a-x}}}{\frac{a+x}{a-x} + 1}$$

input `int(((a + x)/(a - x))^(1/2),x)`output `2*a*atan(((a + x)/(a - x))^(1/2)) - (2*a*((a + x)/(a - x))^(1/2))/((a + x)/(a - x) + 1)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.76

$$\int \sqrt{\frac{a+x}{a-x}} dx = -2a \sin\left(\frac{\sqrt{a-x}}{\sqrt{a}\sqrt{2}}\right) a - \sqrt{a+x} \sqrt{a-x}$$

input `int(((a+x)/(a-x))^(1/2),x)`output `- 2*asin(sqrt(a - x)/(sqrt(a)*sqrt(2)))*a - sqrt(a + x)*sqrt(a - x)`

3.104 $\int \sqrt{(b-x)(-a+x)} dx$

Optimal result	618
Mathematica [A] (verified)	618
Rubi [A] (verified)	619
Maple [A] (verified)	620
Fricas [A] (verification not implemented)	621
Sympy [A] (verification not implemented)	621
Maxima [F(-2)]	622
Giac [A] (verification not implemented)	622
Mupad [F(-1)]	623
Reduce [B] (verification not implemented)	623

Optimal result

Integrand size = 15, antiderivative size = 71

$$\int \sqrt{(b-x)(-a+x)} dx = -\frac{1}{4}(a+b-2x)\sqrt{-ab+(a+b)x-x^2} - \frac{1}{8}(a-b)^2 \arctan\left(\frac{a+b-2x}{2\sqrt{-ab+(a+b)x-x^2}}\right)$$

output

```
-1/8*(a-b)^2*arctan(1/2*(a+b-2*x)/(-a*b+(a+b)*x-x^2)^(1/2))-1/4*(a+b-2*x)*(-a*b+(a+b)*x-x^2)^(1/2)
```

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.06

$$\int \sqrt{(b-x)(-a+x)} dx = \frac{1}{4}\sqrt{(a-x)(-b+x)}\left(-a-b+2x + \frac{(a-b)^2 \arctan\left(\frac{\sqrt{-a+x}}{\sqrt{b-x}}\right)}{\sqrt{b-x}\sqrt{-a+x}}\right)$$

input

```
Integrate[Sqrt[(b-x)*(-a+x)],x]
```

output

$$\frac{(\text{Sqrt}[(a-x)*(-b+x)]*(-a-b+2*x+((a-b)^2*\text{ArcTan}[\text{Sqrt}[-a+x]/\text{Sqrt}[b-x]])/(\text{Sqrt}[b-x]*\text{Sqrt}[-a+x]))}{4}$$
Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {2048, 1087, 1092, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{(x-a)(b-x)} dx \\ & \quad \downarrow 2048 \\ & \int \sqrt{x(a+b)-ab-x^2} dx \\ & \quad \downarrow 1087 \\ & \frac{1}{8}(a-b)^2 \int \frac{1}{\sqrt{-x^2+(a+b)x-ab}} dx - \frac{1}{4}(a+b-2x)\sqrt{x(a+b)-ab-x^2} \\ & \quad \downarrow 1092 \\ & \frac{1}{4}(a-b)^2 \int \frac{1}{-\frac{(a+b-2x)^2}{-x^2+(a+b)x-ab} - 4} d \frac{a+b-2x}{\sqrt{-x^2+(a+b)x-ab}} - \frac{1}{4}(a+b-2x)\sqrt{x(a+b)-ab-x^2} \\ & \quad \downarrow 217 \\ & -\frac{1}{8}(a-b)^2 \arctan\left(\frac{a+b-2x}{2\sqrt{x(a+b)-ab-x^2}}\right) - \frac{1}{4}(a+b-2x)\sqrt{x(a+b)-ab-x^2} \end{aligned}$$

input

$$\text{Int}[\text{Sqrt}[(b-x)*(-a+x)],x]$$

output

$$-1/4*((a+b-2*x)*\text{Sqrt}[-(a*b)+(a+b)*x-x^2]) - ((a-b)^2*\text{ArcTan}[(a+b-2*x)/(2*\text{Sqrt}[-(a*b)+(a+b)*x-x^2]])/8$$

Definitions of rubi rules used

rule 217 $\text{Int}[(a_) + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1} * \text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /;$ FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

rule 1087 $\text{Int}[(a_) + (b_)*(x_) + (c_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[(b + 2*c*x) * ((a + b*x + c*x^2)^p / (2*c*(2*p + 1))), x] - \text{Simp}[p*((b^2 - 4*a*c) / (2*c*(2*p + 1))) \text{Int}[(a + b*x + c*x^2)^{(p-1)}, x], x] /;$ FreeQ[{a, b, c}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[3*p])

rule 1092 $\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_) + (c_)*(x_)^2], x_Symbol] \rightarrow \text{Simp}[2 \text{Subst}[\text{Int}[1/(4*c - x^2), x], x, (b + 2*c*x)/\text{Sqrt}[a + b*x + c*x^2]], x] /;$ FreeQ[{a, b, c}, x]

rule 2048 $\text{Int}[(u_)*((e_)*((a_) + (b_)*(x_)^{(n_)}) * ((c_) + (d_)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Int}[u*(a*c*e + (b*c + a*d)*e*x^n + b*d*e*x^{(2*n)})^p, x] /;$ FreeQ[{a, b, c, d, e, n, p}, x]

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.96

method	result	size
default	$-\frac{(a+b-2x)\sqrt{-ab+(a+b)x-x^2}}{4} - \frac{(4ab-(a+b)^2) \arctan\left(\frac{x-\frac{b}{2}-\frac{a}{2}}{\sqrt{-ab+(a+b)x-x^2}}\right)}{8}$	68
risch	$\frac{(b-x)(a-x)(a+b-2x)}{4\sqrt{-(-b+x)(-a+x)}} - \left(\frac{1}{4}ab - \frac{1}{8}b^2 - \frac{1}{8}a^2\right) \arctan\left(\frac{x-\frac{b}{2}-\frac{a}{2}}{\sqrt{-ab+(a+b)x-x^2}}\right)$	78

input $\text{int}((b-x)*(-a+x))^{(1/2)}, x, \text{method}=_RETURNVERBOSE)$

output $-1/4*(a+b-2*x)*(-a*b+(a+b)*x-x^2)^{(1/2)} - 1/8*(4*a*b-(a+b)^2)*\arctan((x-1/2*b-1/2*a)/(-a*b+(a+b)*x-x^2)^{(1/2)})$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.13

$$\int \sqrt{(b-x)(-a+x)} dx$$

$$= -\frac{1}{8} (a^2 - 2ab + b^2) \arctan \left(-\frac{\sqrt{-ab + (a+b)x - x^2}(a+b-2x)}{2(ab - (a+b)x + x^2)} \right)$$

$$- \frac{1}{4} \sqrt{-ab + (a+b)x - x^2}(a+b-2x)$$

input `integrate(((b-x)*(-a+x))^(1/2),x, algorithm="fricas")`output `-1/8*(a^2 - 2*a*b + b^2)*arctan(-1/2*sqrt(-a*b + (a + b)*x - x^2)*(a + b - 2*x)/(a*b - (a + b)*x + x^2)) - 1/4*sqrt(-a*b + (a + b)*x - x^2)*(a + b - 2*x)`**Sympy [A] (verification not implemented)**

Time = 1.65 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.61

$$\int \sqrt{(b-x)(-a+x)} dx = \left(-\frac{ab}{2} + \frac{\left(\frac{a}{4} + \frac{b}{4}\right)(a+b)}{2} \right) \left(\begin{array}{l} \left\{ \begin{array}{l} -i \log \left(a + b - 2x + 2i\sqrt{-ab - x^2 + x(a+b)} \right) \\ \frac{\left(-\frac{a}{2} - \frac{b}{2} + x\right) \log \left(-\frac{a}{2} - \frac{b}{2} + x\right)}{\sqrt{-\left(-\frac{a}{2} - \frac{b}{2} + x\right)^2}} \end{array} \right. \text{ for } ab - \frac{(a+b)^2}{4} \neq 0 \\ \text{otherwise} \end{array} \right)$$

$$+ \left(-\frac{a}{4} - \frac{b}{4} + \frac{x}{2} \right) \sqrt{-ab - x^2 + x(a+b)}$$

input `integrate(((b-x)*(-a+x))**(1/2),x)`output `(-a*b/2 + (a/4 + b/4)*(a + b)/2)*Piecewise((-I*log(a + b - 2*x + 2*I*sqrt(-a*b - x**2 + x*(a + b))), Ne(a*b - (a + b)**2/4, 0)), ((-a/2 - b/2 + x)*log(-a/2 - b/2 + x)/sqrt(-(-a/2 - b/2 + x)**2), True)) + (-a/4 - b/4 + x/2)*sqrt(-a*b - x**2 + x*(a + b))`

Maxima [F(-2)]

Exception generated.

$$\int \sqrt{(b-x)(-a+x)} dx = \text{Exception raised: ValueError}$$

input `integrate(((b-x)*(-a+x))^(1/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see `assume?` for more details)Is

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.86

$$\int \sqrt{(b-x)(-a+x)} dx = \frac{1}{8} (a^2 - 2ab + b^2) \arcsin\left(\frac{a+b-2x}{a-b}\right) \operatorname{sgn}(-a+b) - \frac{1}{4} \sqrt{-ab + ax + bx - x^2} (a+b-2x)$$

input `integrate(((b-x)*(-a+x))^(1/2),x, algorithm="giac")`

output `1/8*(a^2 - 2*a*b + b^2)*arcsin((a + b - 2*x)/(a - b))*sgn(-a + b) - 1/4*sqrt(-a*b + a*x + b*x - x^2)*(a + b - 2*x)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{(b-x)(-a+x)} dx = \int \sqrt{-(a-x)(b-x)} dx$$

input `int((-a - x)*(b - x)^(1/2),x)`output `int((-a - x)*(b - x)^(1/2), x)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 182, normalized size of antiderivative = 2.56

$$\int \sqrt{(b-x)(-a+x)} dx$$

$$= \frac{i \left(-\operatorname{asinh}\left(\frac{\sqrt{-a+x}i}{\sqrt{-a+b}}\right) a^3 + 3\operatorname{asinh}\left(\frac{\sqrt{-a+x}i}{\sqrt{-a+b}}\right) a^2 b - 3\operatorname{asinh}\left(\frac{\sqrt{-a+x}i}{\sqrt{-a+b}}\right) a b^2 + \operatorname{asinh}\left(\frac{\sqrt{-a+x}i}{\sqrt{-a+b}}\right) b^3 - \sqrt{b-x} \sqrt{a-b} \right)}{4a}$$

input `int(((b-x)*(-a+x))^(1/2),x)`output `(i*(-asinh((sqrt(-a+x)*i)/sqrt(-a+b))*a**3 + 3*asinh((sqrt(-a+x)*i)/sqrt(-a+b))*a**2*b - 3*asinh((sqrt(-a+x)*i)/sqrt(-a+b))*a*b**2 + asinh((sqrt(-a+x)*i)/sqrt(-a+b))*b**3 - sqrt(b-x)*sqrt(a-b)*sqrt(-a+x)*sqrt(-a+b)*a - sqrt(b-x)*sqrt(a-b)*sqrt(-a+x)*sqrt(-a+b)*b + 2*sqrt(b-x)*sqrt(a-b)*sqrt(-a+x)*sqrt(-a+b)*x))/(4*(a-b))`

3.105 $\int \frac{1}{\sqrt{(b-x)(-a+x)}} dx$

Optimal result	624
Mathematica [A] (verified)	624
Rubi [A] (verified)	625
Maple [A] (verified)	626
Fricas [A] (verification not implemented)	626
Sympy [C] (verification not implemented)	627
Maxima [F(-2)]	627
Giac [B] (verification not implemented)	628
Mupad [F(-1)]	628
Reduce [B] (verification not implemented)	628

Optimal result

Integrand size = 15, antiderivative size = 32

$$\int \frac{1}{\sqrt{(b-x)(-a+x)}} dx = -\arctan\left(\frac{a+b-2x}{2\sqrt{-ab+(a+b)x-x^2}}\right)$$

output `-arctan(1/2*(a+b-2*x)/(-a*b+(a+b)*x-x^2)^(1/2))`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.72

$$\int \frac{1}{\sqrt{(b-x)(-a+x)}} dx = \frac{2\sqrt{b-x}\sqrt{-a+x} \arctan\left(\frac{\sqrt{-a+x}}{\sqrt{b-x}}\right)}{\sqrt{(a-x)(-b+x)}}$$

input `Integrate[1/Sqrt[(b-x)*(-a+x)],x]`

output `(2*Sqrt[b-x]*Sqrt[-a+x]*ArcTan[Sqrt[-a+x]/Sqrt[b-x]])/Sqrt[(a-x)*(-b+x)]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2048, 1092, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{(x-a)(b-x)}} dx$$

↓ 2048

$$\int \frac{1}{\sqrt{x(a+b) - ab - x^2}} dx$$

↓ 1092

$$2 \int \frac{1}{-\frac{(a+b-2x)^2}{-x^2+(a+b)x-ab} - 4} d \frac{a+b-2x}{\sqrt{-x^2+(a+b)x-ab}}$$

↓ 217

$$-\arctan\left(\frac{a+b-2x}{2\sqrt{x(a+b) - ab - x^2}}\right)$$

input `Int[1/Sqrt[(b - x)*(-a + x)],x]`

output `-ArcTan[(a + b - 2*x)/(2*Sqrt[-(a*b) + (a + b)*x - x^2])]`

Defintions of rubi rules used

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])`

rule 1092

```
Int[1/Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2], x_Symbol] := Simp[2 Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x]
```

rule 2048

```
Int[(u_)*((e_)*((a_) + (b_)*(x_)^(n_))*((c_) + (d_)*(x_)^(n_)))^(p_), x_Symbol] := Int[u*(a*c*e + (b*c + a*d)*e*x^n + b*d*e*x^(2*n))^p, x] /; FreeQ[{a, b, c, d, e, n, p}, x]
```

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.88

method	result	size
default	$\arctan\left(\frac{x - \frac{b}{2} - \frac{a}{2}}{\sqrt{-ab + (a+b)x - x^2}}\right)$	28

input

```
int(1/((b-x)*(-a+x))^(1/2),x,method=_RETURNVERBOSE)
```

output

```
arctan((x-1/2*b-1/2*a)/(-a*b+(a+b)*x-x^2)^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.34

$$\int \frac{1}{\sqrt{(b-x)(-a+x)}} dx = -\arctan\left(-\frac{\sqrt{-ab + (a+b)x - x^2}(a+b-2x)}{2(ab - (a+b)x + x^2)}\right)$$

input

```
integrate(1/((b-x)*(-a+x))^(1/2),x, algorithm="fricas")
```

output

```
-arctan(-1/2*sqrt(-a*b + (a + b)*x - x^2)*(a + b - 2*x)/(a*b - (a + b)*x + x^2))
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.83 (sec) , antiderivative size = 71, normalized size of antiderivative = 2.22

$$\int \frac{1}{\sqrt{(b-x)(-a+x)}} dx$$

$$= \begin{cases} -i \log \left(a + b - 2x + 2i\sqrt{-ab - x^2 + x(a+b)} \right) & \text{for } ab - \frac{(a+b)^2}{4} \neq 0 \\ \frac{\left(-\frac{a}{2} - \frac{b}{2} + x\right) \log\left(-\frac{a}{2} - \frac{b}{2} + x\right)}{\sqrt{-\left(-\frac{a}{2} - \frac{b}{2} + x\right)^2}} & \text{otherwise} \end{cases}$$

input `integrate(1/((b-x)*(-a+x))**(1/2),x)`

output `Piecewise((-I*log(a + b - 2*x + 2*I*sqrt(-a*b - x**2 + x*(a + b))), Ne(a*b - (a + b)**2/4, 0)), ((-a/2 - b/2 + x)*log(-a/2 - b/2 + x)/sqrt(-(-a/2 - b/2 + x)**2), True))`

Maxima [F(-2)]

Exception generated.

$$\int \frac{1}{\sqrt{(b-x)(-a+x)}} dx = \text{Exception raised: ValueError}$$

input `integrate(1/((b-x)*(-a+x))^(1/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see `assume?` for more details)Is`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 61 vs. $2(28) = 56$.

Time = 0.13 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.91

$$\int \frac{1}{\sqrt{(b-x)(-a+x)}} dx = \frac{1}{8} (a^2 - 2ab + b^2) \arcsin\left(\frac{a+b-2x}{a-b}\right) \operatorname{sgn}(-a+b) - \frac{1}{4} \sqrt{-ab+ax+bx-x^2}(a+b-2x)$$

input `integrate(1/((b-x)*(-a+x))^(1/2),x, algorithm="giac")`

output `1/8*(a^2 - 2*a*b + b^2)*arcsin((a + b - 2*x)/(a - b))*sgn(-a + b) - 1/4*sqrt(-a*b + a*x + b*x - x^2)*(a + b - 2*x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{(b-x)(-a+x)}} dx = \int \frac{1}{\sqrt{-(a-x)(b-x)}} dx$$

input `int(1/(-(a - x)*(b - x))^(1/2),x)`

output `int(1/(-(a - x)*(b - x))^(1/2), x)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.62

$$\int \frac{1}{\sqrt{(b-x)(-a+x)}} dx = -2a \operatorname{sinh}\left(\frac{\sqrt{-a+x}i}{\sqrt{-a+b}}\right) i$$

input `int(1/((b-x)*(-a+x))^(1/2),x)`

output $- 2*\operatorname{asinh}(\sqrt{-a+x}*i)/\sqrt{-a+b}*i$

3.106 $\int \frac{3+5x}{-3+2x+x^2} dx$

Optimal result	630
Mathematica [A] (verified)	630
Rubi [A] (verified)	631
Maple [A] (verified)	632
Fricas [A] (verification not implemented)	632
Sympy [A] (verification not implemented)	632
Maxima [A] (verification not implemented)	633
Giac [A] (verification not implemented)	633
Mupad [B] (verification not implemented)	633
Reduce [B] (verification not implemented)	634

Optimal result

Integrand size = 16, antiderivative size = 15

$$\int \frac{3+5x}{-3+2x+x^2} dx = 2\log(1-x) + 3\log(3+x)$$

output `2*ln(1-x)+3*ln(3+x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{3+5x}{-3+2x+x^2} dx = 2\log(1-x) + 3\log(3+x)$$

input `Integrate[(3 + 5*x)/(-3 + 2*x + x^2), x]`

output `2*Log[1 - x] + 3*Log[3 + x]`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {1141, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{5x + 3}{x^2 + 2x - 3} dx$$

↓ 1141

$$\int \left(\frac{3}{x + 3} - \frac{2}{1 - x} \right) dx$$

↓ 2009

$$2 \log(1 - x) + 3 \log(x + 3)$$

input

```
Int[(3 + 5*x)/(-3 + 2*x + x^2),x]
```

output

```
2*Log[1 - x] + 3*Log[3 + x]
```

Defintions of rubi rules used

rule 1141

```
Int[((d_.) + (e_.)*(x_)^(m_.))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_
Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[1/c^p Int[ExpandIntegrand[
(d + e*x)^m*(b/2 - q/2 + c*x)^p*(b/2 + q/2 + c*x)^p, x], x], x] /; EqQ[p, -
1] || !FractionalPowerFactorQ[q] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[p,
0] && IntegerQ[m] && NiceSqrtQ[b^2 - 4*a*c]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```


Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

method	result	size
default	$3 \ln(3+x) + 2 \ln(-1+x)$	14
norman	$3 \ln(3+x) + 2 \ln(-1+x)$	14
risch	$3 \ln(3+x) + 2 \ln(-1+x)$	14
parallelrisc	$3 \ln(3+x) + 2 \ln(-1+x)$	14

input `int((3+5*x)/(x^2+2*x-3),x,method=_RETURNVERBOSE)`output `3*ln(3+x)+2*ln(-1+x)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{3+5x}{-3+2x+x^2} dx = 3 \log(x+3) + 2 \log(x-1)$$

input `integrate((3+5*x)/(x^2+2*x-3),x, algorithm="fricas")`output `3*log(x + 3) + 2*log(x - 1)`**Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

$$\int \frac{3+5x}{-3+2x+x^2} dx = 2 \log(x-1) + 3 \log(x+3)$$

input `integrate((3+5*x)/(x**2+2*x-3),x)`

output `2*log(x - 1) + 3*log(x + 3)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{3 + 5x}{-3 + 2x + x^2} dx = 3 \log(x + 3) + 2 \log(x - 1)$$

input `integrate((3+5*x)/(x^2+2*x-3),x, algorithm="maxima")`

output `3*log(x + 3) + 2*log(x - 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{3 + 5x}{-3 + 2x + x^2} dx = 3 \log(|x + 3|) + 2 \log(|x - 1|)$$

input `integrate((3+5*x)/(x^2+2*x-3),x, algorithm="giac")`

output `3*log(abs(x + 3)) + 2*log(abs(x - 1))`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{3 + 5x}{-3 + 2x + x^2} dx = 2 \ln(x - 1) + 3 \ln(x + 3)$$

input `int((5*x + 3)/(2*x + x^2 - 3),x)`

output `2*log(x - 1) + 3*log(x + 3)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{3 + 5x}{-3 + 2x + x^2} dx = 2 \log(x - 1) + 3 \log(x + 3)$$

input `int((3+5*x)/(x^2+2*x-3),x)`

output `2*log(x - 1) + 3*log(x + 3)`

3.107 $\int \frac{5+2x}{-3+2x+x^2} dx$

Optimal result	635
Mathematica [A] (verified)	635
Rubi [A] (verified)	636
Maple [A] (verified)	637
Fricas [A] (verification not implemented)	637
Sympy [A] (verification not implemented)	637
Maxima [A] (verification not implemented)	638
Giac [A] (verification not implemented)	638
Mupad [B] (verification not implemented)	638
Reduce [B] (verification not implemented)	639

Optimal result

Integrand size = 16, antiderivative size = 19

$$\int \frac{5+2x}{-3+2x+x^2} dx = \frac{7}{4} \log(1-x) + \frac{1}{4} \log(3+x)$$

output `7/4*ln(1-x)+1/4*ln(3+x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{5+2x}{-3+2x+x^2} dx = \frac{7}{4} \log(1-x) + \frac{1}{4} \log(3+x)$$

input `Integrate[(5 + 2*x)/(-3 + 2*x + x^2), x]`

output `(7*Log[1 - x])/4 + Log[3 + x]/4`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {1141, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{2x + 5}{x^2 + 2x - 3} dx$$

↓ 1141

$$\int \left(\frac{1}{4(x+3)} - \frac{7}{4(1-x)} \right) dx$$

↓ 2009

$$\frac{7}{4} \log(1-x) + \frac{1}{4} \log(x+3)$$

input

```
Int[(5 + 2*x)/(-3 + 2*x + x^2),x]
```

output

```
(7*Log[1 - x])/4 + Log[3 + x]/4
```

Defintions of rubi rules used

rule 1141

```
Int[((d_.) + (e_.)*(x_)^(m_.))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_
Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[1/c^p Int[ExpandIntegrand[
(d + e*x)^m*(b/2 - q/2 + c*x)^p*(b/2 + q/2 + c*x)^p, x], x], x] /; EqQ[p, -
1] || !FractionalPowerFactorQ[q] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[p,
0] && IntegerQ[m] && NiceSqrtQ[b^2 - 4*a*c]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

method	result	size
default	$\frac{\ln(3+x)}{4} + \frac{7\ln(-1+x)}{4}$	14
norman	$\frac{\ln(3+x)}{4} + \frac{7\ln(-1+x)}{4}$	14
risch	$\frac{\ln(3+x)}{4} + \frac{7\ln(-1+x)}{4}$	14
paralelrisch	$\frac{\ln(3+x)}{4} + \frac{7\ln(-1+x)}{4}$	14

input `int((5+2*x)/(x^2+2*x-3),x,method=_RETURNVERBOSE)`output `1/4*ln(3+x)+7/4*ln(-1+x)`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.68

$$\int \frac{5+2x}{-3+2x+x^2} dx = \frac{1}{4} \log(x+3) + \frac{7}{4} \log(x-1)$$

input `integrate((5+2*x)/(x^2+2*x-3),x, algorithm="fricas")`output `1/4*log(x + 3) + 7/4*log(x - 1)`**Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int \frac{5+2x}{-3+2x+x^2} dx = \frac{7 \log(x-1)}{4} + \frac{\log(x+3)}{4}$$

input `integrate((5+2*x)/(x**2+2*x-3),x)`

output $7*\log(x - 1)/4 + \log(x + 3)/4$

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.68

$$\int \frac{5 + 2x}{-3 + 2x + x^2} dx = \frac{1}{4} \log(x + 3) + \frac{7}{4} \log(x - 1)$$

input `integrate((5+2*x)/(x^2+2*x-3),x, algorithm="maxima")`

output $1/4*\log(x + 3) + 7/4*\log(x - 1)$

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{5 + 2x}{-3 + 2x + x^2} dx = \frac{1}{4} \log(|x + 3|) + \frac{7}{4} \log(|x - 1|)$$

input `integrate((5+2*x)/(x^2+2*x-3),x, algorithm="giac")`

output $1/4*\log(\text{abs}(x + 3)) + 7/4*\log(\text{abs}(x - 1))$

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.68

$$\int \frac{5 + 2x}{-3 + 2x + x^2} dx = \frac{7 \ln(x - 1)}{4} + \frac{\ln(x + 3)}{4}$$

input `int((2*x + 5)/(2*x + x^2 - 3),x)`

output $(7*\log(x - 1))/4 + \log(x + 3)/4$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.68

$$\int \frac{5 + 2x}{-3 + 2x + x^2} dx = \frac{7 \log(x - 1)}{4} + \frac{\log(x + 3)}{4}$$

input `int((5+2*x)/(x^2+2*x-3),x)`

output `(7*log(x - 1) + log(x + 3))/4`

3.108 $\int \frac{3x+x^3}{-3-2x+x^2} dx$

Optimal result	640
Mathematica [A] (verified)	640
Rubi [A] (verified)	641
Maple [A] (verified)	642
Fricas [A] (verification not implemented)	642
Sympy [A] (verification not implemented)	643
Maxima [A] (verification not implemented)	643
Giac [A] (verification not implemented)	643
Mupad [B] (verification not implemented)	644
Reduce [B] (verification not implemented)	644

Optimal result

Integrand size = 18, antiderivative size = 23

$$\int \frac{3x + x^3}{-3 - 2x + x^2} dx = 2x + \frac{x^2}{2} + 9 \log(3 - x) + \log(1 + x)$$

output `2*x+1/2*x^2+9*ln(3-x)+ln(1+x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{3x + x^3}{-3 - 2x + x^2} dx = 2x + \frac{x^2}{2} + 9 \log(3 - x) + \log(1 + x)$$

input `Integrate[(3*x + x^3)/(-3 - 2*x + x^2),x]`

output `2*x + x^2/2 + 9*Log[3 - x] + Log[1 + x]`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2027, 2159, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^3 + 3x}{x^2 - 2x - 3} dx \\ & \quad \downarrow \text{2027} \\ & \int \frac{x(x^2 + 3)}{x^2 - 2x - 3} dx \\ & \quad \downarrow \text{2159} \\ & \int \left(\frac{2(5x + 3)}{x^2 - 2x - 3} + x + 2 \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{x^2}{2} + 2x + 9 \log(3 - x) + \log(x + 1) \end{aligned}$$

input `Int[(3*x + x^3)/(-3 - 2*x + x^2),x]`

output `2*x + x^2/2 + 9*Log[3 - x] + Log[1 + x]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2027 `Int[(Fx_.)*((a_.)*(x_)^(r_.) + (b_.)*(x_)^(s_.))^(p_.), x_Symbol] := Int[x^(p*r)*(a + b*x^(s - r))^p*Fx, x] /; FreeQ[{a, b, r, s}, x] && IntegerQ[p] && PosQ[s - r] && !(EqQ[p, 1] && EqQ[u, 1])`

rule 2159

```
Int[(Pq_)*((d_.) + (e_.)*(x_))^(m_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*Pq*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

method	result	size
default	$2x + \frac{x^2}{2} + 9 \ln(-3 + x) + \ln(1 + x)$	20
norman	$2x + \frac{x^2}{2} + 9 \ln(-3 + x) + \ln(1 + x)$	20
risch	$2x + \frac{x^2}{2} + 9 \ln(-3 + x) + \ln(1 + x)$	20
parallelrisc	$2x + \frac{x^2}{2} + 9 \ln(-3 + x) + \ln(1 + x)$	20

input

```
int((x^3+3*x)/(x^2-2*x-3),x,method=_RETURNVERBOSE)
```

output

```
2*x+1/2*x^2+9*ln(-3+x)+ln(1+x)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{3x + x^3}{-3 - 2x + x^2} dx = \frac{1}{2} x^2 + 2x + \log(x + 1) + 9 \log(x - 3)$$

input

```
integrate((x^3+3*x)/(x^2-2*x-3),x, algorithm="fricas")
```

output

```
1/2*x^2 + 2*x + log(x + 1) + 9*log(x - 3)
```

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{3x + x^3}{-3 - 2x + x^2} dx = \frac{x^2}{2} + 2x + 9 \log(x - 3) + \log(x + 1)$$

input `integrate((x**3+3*x)/(x**2-2*x-3),x)`output `x**2/2 + 2*x + 9*log(x - 3) + log(x + 1)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{3x + x^3}{-3 - 2x + x^2} dx = \frac{1}{2} x^2 + 2x + \log(x + 1) + 9 \log(x - 3)$$

input `integrate((x^3+3*x)/(x^2-2*x-3),x, algorithm="maxima")`output `1/2*x^2 + 2*x + log(x + 1) + 9*log(x - 3)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{3x + x^3}{-3 - 2x + x^2} dx = \frac{1}{2} x^2 + 2x + \log(|x + 1|) + 9 \log(|x - 3|)$$

input `integrate((x^3+3*x)/(x^2-2*x-3),x, algorithm="giac")`output `1/2*x^2 + 2*x + log(abs(x + 1)) + 9*log(abs(x - 3))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{3x + x^3}{-3 - 2x + x^2} dx = 2x + \ln(x + 1) + 9 \ln(x - 3) + \frac{x^2}{2}$$

input `int(-(3*x + x^3)/(2*x - x^2 + 3),x)`output `2*x + log(x + 1) + 9*log(x - 3) + x^2/2`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{3x + x^3}{-3 - 2x + x^2} dx = 9 \log(x - 3) + \log(x + 1) + \frac{x^2}{2} + 2x$$

input `int((x^3+3*x)/(x^2-2*x-3),x)`output `(18*log(x - 3) + 2*log(x + 1) + x**2 + 4*x)/2`

3.109 $\int \frac{-1+5x+2x^2}{-2x+x^2+x^3} dx$

Optimal result	645
Mathematica [A] (verified)	645
Rubi [A] (verified)	646
Maple [A] (verified)	647
Fricas [A] (verification not implemented)	647
Sympy [A] (verification not implemented)	648
Maxima [A] (verification not implemented)	648
Giac [A] (verification not implemented)	648
Mupad [B] (verification not implemented)	649
Reduce [B] (verification not implemented)	649

Optimal result

Integrand size = 23, antiderivative size = 23

$$\int \frac{-1 + 5x + 2x^2}{-2x + x^2 + x^3} dx = 2 \log(1 - x) + \frac{\log(x)}{2} - \frac{1}{2} \log(2 + x)$$

output `2*ln(1-x)+1/2*ln(x)-1/2*ln(2+x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{-1 + 5x + 2x^2}{-2x + x^2 + x^3} dx = 2 \log(1 - x) + \frac{\log(x)}{2} - \frac{1}{2} \log(2 + x)$$

input `Integrate[(-1 + 5*x + 2*x^2)/(-2*x + x^2 + x^3), x]`

output `2*Log[1 - x] + Log[x]/2 - Log[2 + x]/2`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {2026, 2159, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{2x^2 + 5x - 1}{x^3 + x^2 - 2x} dx \\ & \quad \downarrow \text{2026} \\ & \int \frac{2x^2 + 5x - 1}{x(x^2 + x - 2)} dx \\ & \quad \downarrow \text{2159} \\ & \int \left(\frac{1}{2x} - \frac{1}{2(x+2)} + \frac{2}{x-1} \right) dx \\ & \quad \downarrow \text{2009} \\ & 2 \log(1-x) + \frac{\log(x)}{2} - \frac{1}{2} \log(x+2) \end{aligned}$$

input `Int[(-1 + 5*x + 2*x^2)/(-2*x + x^2 + x^3), x]`

output `2*Log[1 - x] + Log[x]/2 - Log[2 + x]/2`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2159

```
Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p
_), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*Pq*(a + b*x + c*x^2)^p, x
], x] /; FreeQ[{a, b, c, d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

method	result	size
default	$\frac{\ln(x)}{2} + 2 \ln(-1 + x) - \frac{\ln(2+x)}{2}$	18
norman	$\frac{\ln(x)}{2} + 2 \ln(-1 + x) - \frac{\ln(2+x)}{2}$	18
risch	$\frac{\ln(x)}{2} + 2 \ln(-1 + x) - \frac{\ln(2+x)}{2}$	18
parallelrisch	$\frac{\ln(x)}{2} + 2 \ln(-1 + x) - \frac{\ln(2+x)}{2}$	18

input

```
int((2*x^2+5*x-1)/(x^3+x^2-2*x),x,method=_RETURNVERBOSE)
```

output

```
1/2*ln(x)+2*ln(-1+x)-1/2*ln(2+x)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{-1 + 5x + 2x^2}{-2x + x^2 + x^3} dx = -\frac{1}{2} \log(x + 2) + 2 \log(x - 1) + \frac{1}{2} \log(x)$$

input

```
integrate((2*x^2+5*x-1)/(x^3+x^2-2*x),x, algorithm="fricas")
```

output

```
-1/2*log(x + 2) + 2*log(x - 1) + 1/2*log(x)
```


Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{-1 + 5x + 2x^2}{-2x + x^2 + x^3} dx = \frac{\log(x)}{2} + 2 \log(x - 1) - \frac{\log(x + 2)}{2}$$

input `integrate((2*x**2+5*x-1)/(x**3+x**2-2*x),x)`output `log(x)/2 + 2*log(x - 1) - log(x + 2)/2`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{-1 + 5x + 2x^2}{-2x + x^2 + x^3} dx = -\frac{1}{2} \log(x + 2) + 2 \log(x - 1) + \frac{1}{2} \log(x)$$

input `integrate((2*x^2+5*x-1)/(x^3+x^2-2*x),x, algorithm="maxima")`output `-1/2*log(x + 2) + 2*log(x - 1) + 1/2*log(x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \frac{-1 + 5x + 2x^2}{-2x + x^2 + x^3} dx = -\frac{1}{2} \log(|x + 2|) + 2 \log(|x - 1|) + \frac{1}{2} \log(|x|)$$

input `integrate((2*x^2+5*x-1)/(x^3+x^2-2*x),x, algorithm="giac")`output `-1/2*log(abs(x + 2)) + 2*log(abs(x - 1)) + 1/2*log(abs(x))`

Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{-1 + 5x + 2x^2}{-2x + x^2 + x^3} dx = 2 \ln(x - 1) + \operatorname{atanh}\left(\frac{135}{11(11x - 5)} + \frac{16}{11}\right)$$

input `int((5*x + 2*x^2 - 1)/(x^2 - 2*x + x^3),x)`

output `2*log(x - 1) + atanh(135/(11*(11*x - 5)) + 16/11)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{-1 + 5x + 2x^2}{-2x + x^2 + x^3} dx = 2 \log(x - 1) - \frac{\log(x + 2)}{2} + \frac{\log(x)}{2}$$

input `int((2*x^2+5*x-1)/(x^3+x^2-2*x),x)`

output `(4*log(x - 1) - log(x + 2) + log(x))/2`

$$3.110 \quad \int \frac{3+2x+x^2}{(-1+x)(1+x)^2} dx$$

Optimal result	650
Mathematica [A] (verified)	650
Rubi [A] (verified)	651
Maple [A] (verified)	652
Fricas [A] (verification not implemented)	652
Sympy [A] (verification not implemented)	652
Maxima [A] (verification not implemented)	653
Giac [A] (verification not implemented)	653
Mupad [B] (verification not implemented)	653
Reduce [B] (verification not implemented)	654

Optimal result

Integrand size = 19, antiderivative size = 24

$$\int \frac{3+2x+x^2}{(-1+x)(1+x)^2} dx = \frac{1}{1+x} + \frac{3}{2} \log(1-x) - \frac{1}{2} \log(1+x)$$

output `1/(1+x)+3/2*ln(1-x)-1/2*ln(1+x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{3+2x+x^2}{(-1+x)(1+x)^2} dx = \frac{1}{1+x} + \frac{3}{2} \log(-1+x) - \frac{1}{2} \log(1+x)$$

input `Integrate[(3 + 2*x + x^2)/((-1 + x)*(1 + x)^2),x]`

output `(1 + x)^(-1) + (3*Log[-1 + x])/2 - Log[1 + x]/2`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2 + 2x + 3}{(x-1)(x+1)^2} dx$$

$$\downarrow 1195$$

$$\int \left(-\frac{1}{2(x+1)} - \frac{1}{(x+1)^2} + \frac{3}{2(x-1)} \right) dx$$

$$\downarrow 2009$$

$$\frac{1}{x+1} + \frac{3}{2} \log(1-x) - \frac{1}{2} \log(x+1)$$

input

```
Int[(3 + 2*x + x^2)/((-1 + x)*(1 + x)^2), x]
```

output

```
(1 + x)^(-1) + (3*Log[1 - x])/2 - Log[1 + x]/2
```

Defintions of rubi rules used

rule 1195

```
Int[((d_.) + (e_.)*(x_)^(m_.))*((f_.) + (g_.)*(x_)^(n_.))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]
```

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.79

method	result	size
default	$\frac{3 \ln(-1+x)}{2} + \frac{1}{1+x} - \frac{\ln(1+x)}{2}$	19
norman	$\frac{3 \ln(-1+x)}{2} + \frac{1}{1+x} - \frac{\ln(1+x)}{2}$	19
risch	$\frac{3 \ln(-1+x)}{2} + \frac{1}{1+x} - \frac{\ln(1+x)}{2}$	19
parallelrisc	$\frac{3 \ln(-1+x)x - \ln(1+x)x + 2 + 3 \ln(-1+x) - \ln(1+x)}{2x+2}$	36

input `int((x^2+2*x+3)/(-1+x)/(1+x)^2,x,method=_RETURNVERBOSE)`output `3/2*ln(-1+x)+1/(1+x)-1/2*ln(1+x)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{3 + 2x + x^2}{(-1+x)(1+x)^2} dx = -\frac{(x+1)\log(x+1) - 3(x+1)\log(x-1) - 2}{2(x+1)}$$

input `integrate((x^2+2*x+3)/(-1+x)/(1+x)^2,x, algorithm="fricas")`output `-1/2*((x + 1)*log(x + 1) - 3*(x + 1)*log(x - 1) - 2)/(x + 1)`**Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.79

$$\int \frac{3 + 2x + x^2}{(-1+x)(1+x)^2} dx = \frac{3 \log(x-1)}{2} - \frac{\log(x+1)}{2} + \frac{1}{x+1}$$

input `integrate((x**2+2*x+3)/(-1+x)/(1+x)**2,x)`

output $3*\log(x - 1)/2 - \log(x + 1)/2 + 1/(x + 1)$

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.75

$$\int \frac{3 + 2x + x^2}{(-1 + x)(1 + x)^2} dx = \frac{1}{x + 1} - \frac{1}{2} \log(x + 1) + \frac{3}{2} \log(x - 1)$$

input `integrate((x^2+2*x+3)/(-1+x)/(1+x)^2,x, algorithm="maxima")`

output $1/(x + 1) - 1/2*\log(x + 1) + 3/2*\log(x - 1)$

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{3 + 2x + x^2}{(-1 + x)(1 + x)^2} dx = \frac{1}{x + 1} + \log(|x + 1|) + \frac{3}{2} \log\left(\left|-\frac{2}{x + 1} + 1\right|\right)$$

input `integrate((x^2+2*x+3)/(-1+x)/(1+x)^2,x, algorithm="giac")`

output $1/(x + 1) + \log(\text{abs}(x + 1)) + 3/2*\log(\text{abs}(-2/(x + 1) + 1))$

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.75

$$\int \frac{3 + 2x + x^2}{(-1 + x)(1 + x)^2} dx = \frac{3 \ln(x - 1)}{2} - \frac{\ln(x + 1)}{2} + \frac{1}{x + 1}$$

input `int((2*x + x^2 + 3)/((x - 1)*(x + 1)^2),x)`

output $(3*\log(x - 1))/2 - \log(x + 1)/2 + 1/(x + 1)$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.58

$$\int \frac{3 + 2x + x^2}{(-1 + x)(1 + x)^2} dx = \frac{3 \log(x - 1) x + 3 \log(x - 1) - \log(x + 1) x - \log(x + 1) - 2x}{2x + 2}$$

input `int((x^2+2*x+3)/(-1+x)/(1+x)^2,x)`

output `(3*log(x - 1)*x + 3*log(x - 1) - log(x + 1)*x - log(x + 1) - 2*x)/(2*(x + 1))`

$$3.111 \quad \int \frac{-2+2x+3x^2}{-1+x^3} dx$$

Optimal result	655
Mathematica [A] (verified)	655
Rubi [A] (verified)	656
Maple [A] (verified)	658
Fricas [A] (verification not implemented)	658
Sympy [A] (verification not implemented)	659
Maxima [A] (verification not implemented)	659
Giac [A] (verification not implemented)	659
Mupad [B] (verification not implemented)	660
Reduce [B] (verification not implemented)	660

Optimal result

Integrand size = 18, antiderivative size = 28

$$\int \frac{-2 + 2x + 3x^2}{-1 + x^3} dx = \frac{4 \arctan\left(\frac{1+2x}{\sqrt{3}}\right)}{\sqrt{3}} + \log(1 - x^3)$$

output `ln(-x^3+1)+4/3*arctan(1/3*(1+2*x)*3^(1/2))*3^(1/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{-2 + 2x + 3x^2}{-1 + x^3} dx = \frac{4 \arctan\left(\frac{1+2x}{\sqrt{3}}\right)}{\sqrt{3}} + \log(1 - x^3)$$

input `Integrate[(-2 + 2*x + 3*x^2)/(-1 + x^3), x]`

output `(4*ArcTan[(1 + 2*x)/Sqrt[3]])/Sqrt[3] + Log[1 - x^3]`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2410, 25, 792, 2019, 1083, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{3x^2 + 2x - 2}{x^3 - 1} dx \\
 & \quad \downarrow \text{2410} \\
 & \int \frac{2x - 2}{x^3 - 1} dx + 3 \int -\frac{x^2}{1 - x^3} dx \\
 & \quad \downarrow \text{25} \\
 & \int \frac{2x - 2}{x^3 - 1} dx - 3 \int \frac{x^2}{1 - x^3} dx \\
 & \quad \downarrow \text{792} \\
 & \int \frac{2x - 2}{x^3 - 1} dx + \log(1 - x^3) \\
 & \quad \downarrow \text{2019} \\
 & \int \frac{1}{\frac{x^2}{2} + \frac{x}{2} + \frac{1}{2}} dx + \log(1 - x^3) \\
 & \quad \downarrow \text{1083} \\
 & \log(1 - x^3) - 2 \int \frac{1}{-(x + \frac{1}{2})^2 - \frac{3}{4}} d\left(x + \frac{1}{2}\right) \\
 & \quad \downarrow \text{217} \\
 & \frac{4 \arctan\left(\frac{2(x + \frac{1}{2})}{\sqrt{3}}\right)}{\sqrt{3}} + \log(1 - x^3)
 \end{aligned}$$

input `Int[(-2 + 2*x + 3*x^2)/(-1 + x^3), x]`

output `(4*ArcTan[(2*(1/2 + x))/Sqrt[3]])/Sqrt[3] + Log[1 - x^3]`

Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 217 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 792 `Int[(x_)^(m_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]`
- rule 1083 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`
- rule 2019 `Int[(u_)*(Px_)^(p_)*(Qx_)^(q_), x_Symbol] := Int[u*PolynomialQuotient[Px, Qx, x]^p*Qx^(p + q), x] /; FreeQ[q, x] && PolyQ[Px, x] && PolyQ[Qx, x] && EqQ[PolynomialRemainder[Px, Qx, x], 0] && IntegerQ[p] && LtQ[p*q, 0]`
- rule 2410 `Int[(P2_)/((a_) + (b_)*(x_)^3), x_Symbol] := With[{A = Coeff[P2, x, 0], B = Coeff[P2, x, 1], C = Coeff[P2, x, 2]}, Int[(A + B*x)/(a + b*x^3), x] + Simp[C Int[x^2/(a + b*x^3), x], x] /; EqQ[a*B^3 - b*A^3, 0] || !RationalQ[a/b]] /; FreeQ[{a, b}, x] && PolyQ[P2, x, 2]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

method	result
default	$\ln(-1+x) + \ln(x^2+x+1) + \frac{4 \arctan\left(\frac{(1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{3}$
risch	$\ln(4x^2+4x+4) + \frac{4 \arctan\left(\frac{(1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{3} + \ln(-1+x)$
meijerg	$-\frac{2x \left(\ln\left(1-(x^3)^{\frac{1}{3}}\right) - \frac{\ln\left(1+(x^3)^{\frac{1}{3}}+(x^3)^{\frac{2}{3}}\right)}{2} - \sqrt{3} \arctan\left(\frac{\sqrt{3}(x^3)^{\frac{1}{3}}}{2+(x^3)^{\frac{1}{3}}}\right) \right)}{3(x^3)^{\frac{1}{3}}} + \ln(-x^3+1) + \frac{2x^2 \left(\ln\left(1-(x^3)^{\frac{1}{3}}\right) - \frac{\ln\left(1+(x^3)^{\frac{1}{3}}+(x^3)^{\frac{2}{3}}\right)}{2} - \sqrt{3} \arctan\left(\frac{\sqrt{3}(x^3)^{\frac{1}{3}}}{2+(x^3)^{\frac{1}{3}}}\right) \right)}{3(x^3)^{\frac{1}{3}}}$

input `int((3*x^2+2*x-2)/(x^3-1),x,method=_RETURNVERBOSE)`output `ln(-1+x)+ln(x^2+x+1)+4/3*arctan(1/3*(1+2*x)*3^(1/2))*3^(1/2)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{-2+2x+3x^2}{-1+x^3} dx = \frac{4}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x+1)\right) + \log(x^2+x+1) + \log(x-1)$$

input `integrate((3*x^2+2*x-2)/(x^3-1),x, algorithm="fricas")`output `4/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + log(x^2 + x + 1) + log(x - 1)`

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.11

$$\int \frac{-2 + 2x + 3x^2}{-1 + x^3} dx = \log(x - 1)$$

input `integrate((3*x**2+2*x-2)/(x**3-1),x)`output `log(x - 1)`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{-2 + 2x + 3x^2}{-1 + x^3} dx = \frac{4}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x + 1)\right) + \log(x^2 + x + 1) + \log(x - 1)$$

input `integrate((3*x^2+2*x-2)/(x^3-1),x, algorithm="maxima")`output `4/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + log(x^2 + x + 1) + log(x - 1)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

$$\int \frac{-2 + 2x + 3x^2}{-1 + x^3} dx = \frac{4}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x + 1)\right) + \log(x^2 + x + 1) + \log(|x - 1|)$$

input `integrate((3*x^2+2*x-2)/(x^3-1),x, algorithm="giac")`output `4/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + log(x^2 + x + 1) + log(abs(x - 1))`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 57, normalized size of antiderivative = 2.04

$$\int \frac{-2 + 2x + 3x^2}{-1 + x^3} dx = \ln \left(x + \frac{1}{2} - \frac{\sqrt{3}1i}{2} \right) + \ln \left(x + \frac{1}{2} + \frac{\sqrt{3}1i}{2} \right) + \ln(x - 1) \\ - \frac{\sqrt{3} \ln \left(x + \frac{1}{2} - \frac{\sqrt{3}1i}{2} \right) 2i}{3} + \frac{\sqrt{3} \ln \left(x + \frac{1}{2} + \frac{\sqrt{3}1i}{2} \right) 2i}{3}$$

input `int((2*x + 3*x^2 - 2)/(x^3 - 1),x)`output `log(x - (3^(1/2)*1i)/2 + 1/2) + log(x + (3^(1/2)*1i)/2 + 1/2) + log(x - 1) - (3^(1/2)*log(x - (3^(1/2)*1i)/2 + 1/2)*2i)/3 + (3^(1/2)*log(x + (3^(1/2)*1i)/2 + 1/2)*2i)/3`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \frac{-2 + 2x + 3x^2}{-1 + x^3} dx = \frac{4\sqrt{3} \operatorname{atan}\left(\frac{2x+1}{\sqrt{3}}\right)}{3} + \log(x^2 + x + 1) + \log(x - 1)$$

input `int((3*x^2+2*x-2)/(x^3-1),x)`output `(4*sqrt(3)*atan((2*x + 1)/sqrt(3)) + 3*log(x**2 + x + 1) + 3*log(x - 1))/3`

$$3.112 \quad \int \frac{2-x+2x^2-x^3+x^4}{(-1+x)(2+x^2)^2} dx$$

Optimal result	661
Mathematica [A] (verified)	661
Rubi [A] (verified)	662
Maple [A] (verified)	663
Fricas [A] (verification not implemented)	664
Sympy [A] (verification not implemented)	664
Maxima [A] (verification not implemented)	664
Giac [A] (verification not implemented)	665
Mupad [B] (verification not implemented)	665
Reduce [B] (verification not implemented)	666

Optimal result

Integrand size = 31, antiderivative size = 49

$$\int \frac{2-x+2x^2-x^3+x^4}{(-1+x)(2+x^2)^2} dx = \frac{1}{2(2+x^2)} - \frac{\arctan\left(\frac{x}{\sqrt{2}}\right)}{3\sqrt{2}} + \frac{1}{3}\log(1-x) + \frac{1}{3}\log(2+x^2)$$

output `1/2/(x^2+2)+1/3*ln(1-x)+1/3*ln(x^2+2)-1/6*arctan(1/2*x*2^(1/2))*2^(1/2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.24

$$\int \frac{2-x+2x^2-x^3+x^4}{(-1+x)(2+x^2)^2} dx = \frac{1}{2(3+2(-1+x)+(-1+x)^2)} - \frac{\arctan\left(\frac{x}{\sqrt{2}}\right)}{3\sqrt{2}} + \frac{1}{3}\log(3+2(-1+x)+(-1+x)^2) + \frac{1}{3}\log(-1+x)$$

input `Integrate[(2 - x + 2*x^2 - x^3 + x^4)/((-1 + x)*(2 + x^2)^2), x]`

output `1/(2*(3 + 2*(-1 + x) + (-1 + x)^2)) - ArcTan[x/Sqrt[2]]/(3*Sqrt[2]) + Log[3 + 2*(-1 + x) + (-1 + x)^2]/3 + Log[-1 + x]/3`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.129$, Rules used = {2178, 27, 2160, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^4 - x^3 + 2x^2 - x + 2}{(x-1)(x^2+2)^2} dx \\
 & \quad \downarrow \text{2178} \\
 & \frac{1}{2(x^2+2)} - \frac{1}{4} \int \frac{4(x^2-x+1)}{(1-x)(x^2+2)} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{2(x^2+2)} - \int \frac{x^2-x+1}{(1-x)(x^2+2)} dx \\
 & \quad \downarrow \text{2160} \\
 & \frac{1}{2(x^2+2)} - \int \left(\frac{1-2x}{3(x^2+2)} - \frac{1}{3(x-1)} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & -\frac{\arctan\left(\frac{x}{\sqrt{2}}\right)}{3\sqrt{2}} + \frac{1}{2(x^2+2)} + \frac{1}{3} \log(x^2+2) + \frac{1}{3} \log(1-x)
 \end{aligned}$$

input `Int[(2 - x + 2*x^2 - x^3 + x^4)/((-1 + x)*(2 + x^2)^2), x]`

output `1/(2*(2 + x^2)) - ArcTan[x/Sqrt[2]]/(3*Sqrt[2]) + Log[1 - x]/3 + Log[2 + x^2]/3`

Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2160 `Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*Pq*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

rule 2178 `Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] :> With[{Qx = PolynomialQuotient[(d + e*x)^m*Pq, a + b*x^2, x], R = Coeff[PolynomialRemainder[(d + e*x)^m*Pq, a + b*x^2, x], x, 0], S = Coeff[PolynomialRemainder[(d + e*x)^m*Pq, a + b*x^2, x], x, 1]}, Simp[(a*S - b*R*x)*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] + Simp[1/(2*a*b*(p + 1)) Int[(d + e*x)^m*(a + b*x^2)^(p + 1)*ExpandToSum[(2*a*b*(p + 1)*Qx)/(d + e*x)^m + (b*R*(2*p + 3))/(d + e*x)^m, x], x], x] /; FreeQ[{a, b, d, e}, x] && PolyQ[Pq, x] && NeQ[b*d^2 + a*e^2, 0] && LtQ[p, -1] && ILtQ[m, 0]`

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.76

method	result	size
default	$\frac{1}{2x^2+4} + \frac{\ln(x^2+2)}{3} - \frac{\arctan\left(\frac{x\sqrt{2}}{2}\right)\sqrt{2}}{6} + \frac{\ln(-1+x)}{3}$	37
risch	$\frac{1}{2x^2+4} + \frac{\ln(x^2+2)}{3} - \frac{\arctan\left(\frac{x\sqrt{2}}{2}\right)\sqrt{2}}{6} + \frac{\ln(-1+x)}{3}$	37

input `int((x^4-x^3+2*x^2-x+2)/(-1+x)/(x^2+2)^2,x,method=_RETURNVERBOSE)`

output `1/2/(x^2+2)+1/3*ln(x^2+2)-1/6*arctan(1/2*x*2^(1/2))*2^(1/2)+1/3*ln(-1+x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.04

$$\int \frac{2 - x + 2x^2 - x^3 + x^4}{(-1 + x)(2 + x^2)^2} dx$$

$$= -\frac{\sqrt{2}(x^2 + 2) \arctan\left(\frac{1}{2}\sqrt{2}x\right) - 2(x^2 + 2) \log(x^2 + 2) - 2(x^2 + 2) \log(x - 1) - 3}{6(x^2 + 2)}$$

input `integrate((x^4-x^3+2*x^2-x+2)/(-1+x)/(x^2+2)^2,x, algorithm="fricas")`output `-1/6*(sqrt(2)*(x^2 + 2)*arctan(1/2*sqrt(2)*x) - 2*(x^2 + 2)*log(x^2 + 2) - 2*(x^2 + 2)*log(x - 1) - 3)/(x^2 + 2)`**Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.29

$$\int \frac{2 - x + 2x^2 - x^3 + x^4}{(-1 + x)(2 + x^2)^2} dx = \frac{\log(x - 1)}{3} + \frac{1}{2x^2 + 4}$$

input `integrate((x**4-x**3+2*x**2-x+2)/(-1+x)/(x**2+2)**2,x)`output `log(x - 1)/3 + 1/(2*x**2 + 4)`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.73

$$\int \frac{2 - x + 2x^2 - x^3 + x^4}{(-1 + x)(2 + x^2)^2} dx = -\frac{1}{6} \sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}x\right) + \frac{1}{2(x^2 + 2)}$$

$$+ \frac{1}{3} \log(x^2 + 2) + \frac{1}{3} \log(x - 1)$$

input `integrate((x^4-x^3+2*x^2-x+2)/(-1+x)/(x^2+2)^2,x, algorithm="maxima")`

output $-1/6*\sqrt{2}*\arctan(1/2*\sqrt{2}*x) + 1/2/(x^2 + 2) + 1/3*\log(x^2 + 2) + 1/3*\log(x - 1)$

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.76

$$\int \frac{2 - x + 2x^2 - x^3 + x^4}{(-1 + x)(2 + x^2)^2} dx = -\frac{1}{6} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2} x\right) + \frac{1}{2(x^2 + 2)} + \frac{1}{3} \log(x^2 + 2) + \frac{1}{3} \log(|x - 1|)$$

input `integrate((x^4-x^3+2*x^2-x+2)/(-1+x)/(x^2+2)^2,x, algorithm="giac")`

output $-1/6*\sqrt{2}*\arctan(1/2*\sqrt{2}*x) + 1/2/(x^2 + 2) + 1/3*\log(x^2 + 2) + 1/3*\log(\text{abs}(x - 1))$

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.08

$$\int \frac{2 - x + 2x^2 - x^3 + x^4}{(-1 + x)(2 + x^2)^2} dx = \frac{\ln(x - 1)}{3} + \ln(x - \sqrt{2} i) \left(\frac{1}{3} + \frac{\sqrt{2} i}{12}\right) - \ln(x + \sqrt{2} i) \left(-\frac{1}{3} + \frac{\sqrt{2} i}{12}\right) + \frac{1}{2(x^2 + 2)}$$

input `int((2*x^2 - x - x^3 + x^4 + 2)/((x^2 + 2)^2*(x - 1)),x)`

output $\log(x - 1)/3 + \log(x - 2^{(1/2)}*1i)*((2^{(1/2)}*1i)/12 + 1/3) - \log(x + 2^{(1/2)}*1i)*((2^{(1/2)}*1i)/12 - 1/3) + 1/(2*(x^2 + 2))$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.53

$$\int \frac{2 - x + 2x^2 - x^3 + x^4}{(-1 + x)(2 + x^2)^2} dx$$

$$= \frac{-2\sqrt{2} \operatorname{atan}\left(\frac{x}{\sqrt{2}}\right) x^2 - 4\sqrt{2} \operatorname{atan}\left(\frac{x}{\sqrt{2}}\right) + 4 \log(x^2 + 2) x^2 + 8 \log(x^2 + 2) + 4 \log(x - 1) x^2 + 8 \log(x - 1)}{12x^2 + 24}$$

input `int((x^4-x^3+2*x^2-x+2)/(-1+x)/(x^2+2)^2,x)`output `(- 2*sqrt(2)*atan(x/sqrt(2))*x**2 - 4*sqrt(2)*atan(x/sqrt(2)) + 4*log(x**2 + 2)*x**2 + 8*log(x**2 + 2) + 4*log(x - 1)*x**2 + 8*log(x - 1) - 3*x**2)/(12*(x**2 + 2))`

3.113 $\int \frac{1}{\cos(x)+\sin(x)} dx$

Optimal result	667
Mathematica [C] (verified)	667
Rubi [A] (verified)	668
Maple [A] (verified)	669
Fricas [B] (verification not implemented)	669
Sympy [A] (verification not implemented)	670
Maxima [B] (verification not implemented)	670
Giac [B] (verification not implemented)	670
Mupad [B] (verification not implemented)	671
Reduce [B] (verification not implemented)	671

Optimal result

Integrand size = 7, antiderivative size = 21

$$\int \frac{1}{\cos(x) + \sin(x)} dx = -\frac{\operatorname{arctanh}\left(\frac{\cos(x)-\sin(x)}{\sqrt{2}}\right)}{\sqrt{2}}$$

output `-1/2*arctanh(1/2*(cos(x)-sin(x))*2^(1/2))*2^(1/2)`

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.14

$$\int \frac{1}{\cos(x) + \sin(x)} dx = (-1 - i)(-1)^{3/4} \operatorname{arctanh}\left(\frac{-1 + \tan\left(\frac{x}{2}\right)}{\sqrt{2}}\right)$$

input `Integrate[(Cos[x] + Sin[x])^(-1),x]`

output `(-1 - I)*(-1)^(3/4)*ArcTanh[(-1 + Tan[x/2])/Sqrt[2]]`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3553, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\sin(x) + \cos(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{\sin(x) + \cos(x)} dx \\ & \quad \downarrow \text{3553} \\ & - \int \frac{1}{2 - (\cos(x) - \sin(x))^2} d(\cos(x) - \sin(x)) \\ & \quad \downarrow \text{219} \\ & - \frac{\operatorname{arctanh}\left(\frac{\cos(x) - \sin(x)}{\sqrt{2}}\right)}{\sqrt{2}} \end{aligned}$$

input `Int[(Cos[x] + Sin[x])^(-1),x]`

output `-(ArcTanh[(Cos[x] - Sin[x])/Sqrt[2]]/Sqrt[2])`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3553

```
Int[(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x
_Symbol] :> Simp[-d^(-1) Subst[Int[1/(a^2 + b^2 - x^2), x], x, b*Cos[c +
d*x] - a*Sin[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]
```

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

method	result	size
default	$\sqrt{2} \operatorname{arctanh}\left(\frac{(2 \tan(\frac{x}{2}) - 2)\sqrt{2}}{4}\right)$	19
risch	$\frac{\sqrt{2} \ln\left(e^{ix} - \frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right)}{2} - \frac{\sqrt{2} \ln\left(e^{ix} + \frac{\sqrt{2}}{2} - \frac{i\sqrt{2}}{2}\right)}{2}$	48

input

```
int(1/(cos(x)+sin(x)),x,method=_RETURNVERBOSE)
```

output

```
2^(1/2)*arctanh(1/4*(2*tan(1/2*x)-2)*2^(1/2))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 38 vs. 2(18) = 36.

Time = 0.08 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.81

$$\int \frac{1}{\cos(x) + \sin(x)} dx = \frac{1}{4} \sqrt{2} \log \left(\frac{2(\sqrt{2} - \cos(x)) \sin(x) - 2\sqrt{2} \cos(x) + 3}{2 \cos(x) \sin(x) + 1} \right)$$

input

```
integrate(1/(cos(x)+sin(x)),x, algorithm="fricas")
```

output

```
1/4*sqrt(2)*log((2*(sqrt(2) - cos(x))*sin(x) - 2*sqrt(2)*cos(x) + 3)/(2*cos(x)*sin(x) + 1))
```

Sympy [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.86

$$\int \frac{1}{\cos(x) + \sin(x)} dx = \frac{\sqrt{2} \log\left(\tan\left(\frac{x}{2}\right) - 1 + \sqrt{2}\right)}{2} - \frac{\sqrt{2} \log\left(\tan\left(\frac{x}{2}\right) - \sqrt{2} - 1\right)}{2}$$

input `integrate(1/(cos(x)+sin(x)),x)`

output `sqrt(2)*log(tan(x/2) - 1 + sqrt(2))/2 - sqrt(2)*log(tan(x/2) - sqrt(2) - 1)/2`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. 2(18) = 36.

Time = 0.10 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.86

$$\int \frac{1}{\cos(x) + \sin(x)} dx = -\frac{1}{2} \sqrt{2} \log\left(-\frac{\sqrt{2} - \frac{\sin(x)}{\cos(x)+1} + 1}{\sqrt{2} + \frac{\sin(x)}{\cos(x)+1} - 1}\right)$$

input `integrate(1/(cos(x)+sin(x)),x, algorithm="maxima")`

output `-1/2*sqrt(2)*log(-(sqrt(2) - sin(x)/(cos(x) + 1) + 1)/(sqrt(2) + sin(x)/(cos(x) + 1) - 1))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 37 vs. 2(18) = 36.

Time = 0.14 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.76

$$\int \frac{1}{\cos(x) + \sin(x)} dx = -\frac{1}{2} \sqrt{2} \log\left(\frac{|-2\sqrt{2} + 2 \tan\left(\frac{1}{2}x\right) - 2|}{|2\sqrt{2} + 2 \tan\left(\frac{1}{2}x\right) - 2|}\right)$$

input `integrate(1/(cos(x)+sin(x)),x, algorithm="giac")`

output `-1/2*sqrt(2)*log(abs(-2*sqrt(2) + 2*tan(1/2*x) - 2)/abs(2*sqrt(2) + 2*tan(1/2*x) - 2))`

Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{1}{\cos(x) + \sin(x)} dx = -\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2}}{2} - \frac{\sqrt{2} \tan\left(\frac{x}{2}\right)}{2}\right)$$

input `int(1/(cos(x) + sin(x)),x)`

output `-2^(1/2)*atanh(2^(1/2)/2 - (2^(1/2)*tan(x/2))/2)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.29

$$\int \frac{1}{\cos(x) + \sin(x)} dx = \frac{\sqrt{2} (-\log(-\sqrt{2} + \tan\left(\frac{x}{2}\right) - 1) + \log(\sqrt{2} + \tan\left(\frac{x}{2}\right) - 1))}{2}$$

input `int(1/(cos(x)+sin(x)),x)`

output `(sqrt(2)*(- log(- sqrt(2) + tan(x/2) - 1) + log(sqrt(2) + tan(x/2) - 1)))/2`

3.114 $\int \frac{x}{4-x^2+\sqrt{4-x^2}} dx$

Optimal result	672
Mathematica [A] (verified)	672
Rubi [A] (verified)	673
Maple [A] (verified)	674
Fricas [B] (verification not implemented)	674
Sympy [B] (verification not implemented)	675
Maxima [A] (verification not implemented)	675
Giac [A] (verification not implemented)	675
Mupad [B] (verification not implemented)	676
Reduce [B] (verification not implemented)	676

Optimal result

Integrand size = 22, antiderivative size = 16

$$\int \frac{x}{4-x^2+\sqrt{4-x^2}} dx = -\log\left(1+\sqrt{4-x^2}\right)$$

output `-ln(1+(-x^2+4)^(1/2))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{x}{4-x^2+\sqrt{4-x^2}} dx = -\log\left(1+\sqrt{4-x^2}\right)$$

input `Integrate[x/(4 - x^2 + Sqrt[4 - x^2]),x]`

output `-Log[1 + Sqrt[4 - x^2]]`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.136$, Rules used = {2586, 7267, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{-x^2 + \sqrt{4-x^2} + 4} dx$$

↓ 2586

$$\frac{1}{2} \int \frac{1}{-x^2 + \sqrt{4-x^2} + 4} dx^2$$

↓ 7267

$$-\int \frac{1}{\sqrt{4-x^2} + 1} d\sqrt{4-x^2}$$

↓ 16

$$-\log(\sqrt{4-x^2} + 1)$$

input `Int[x/(4 - x^2 + Sqrt[4 - x^2]),x]`

output `-Log[1 + Sqrt[4 - x^2]]`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] :> Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 2586 `Int[(x_)^(m_.)/((c_) + (d_.)*(x_)^(n_) + (e_.)*Sqrt[(a_) + (b_.)*(x_)^(n_)]) , x_Symbol] :> Simp[1/n Subst[Int[x^((m + 1)/n - 1)/(c + d*x + e*Sqrt[a + b*x]), x], x, x^n], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[(m + 1)/n]`

rule 7267

```
Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Si
mp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])], x
] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]]
```

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.31

method	result
trager	$\ln\left(\frac{-1+\sqrt{-x^2+4}}{x^2-3}\right)$
default	$-\frac{\ln(x^2-3)}{2} + \frac{\sqrt{-(-2+x)^2-4x+8}-2\arcsin(\frac{x}{2})}{2(2+\sqrt{3})(-2+\sqrt{3})} + \frac{\sqrt{-(2+x)^2+4x+8}+2\arcsin(\frac{x}{2})}{2(2+\sqrt{3})(-2+\sqrt{3})} + \frac{-\sqrt{-(x-\sqrt{3})^2-2\sqrt{3}(x-\sqrt{3})+1}}{2}$

input

```
int(x/(4-x^2+(-x^2+4)^(1/2)),x,method=_RETURNVERBOSE)
```

output

```
ln((-1+(-x^2+4)^(1/2))/(x^2-3))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 55 vs. 2(14) = 28.

Time = 0.07 (sec) , antiderivative size = 55, normalized size of antiderivative = 3.44

$$\int \frac{x}{4-x^2+\sqrt{4-x^2}} dx = -\frac{1}{2} \log(x^2-3) + \frac{1}{2} \log\left(-\frac{x^2+3\sqrt{-x^2+4}-6}{x^2}\right) - \frac{1}{2} \log\left(-\frac{x^2+\sqrt{-x^2+4}-2}{x^2}\right)$$

input

```
integrate(x/(4-x^2+(-x^2+4)^(1/2)),x, algorithm="fricas")
```

output

```
-1/2*log(x^2 - 3) + 1/2*log(-(x^2 + 3*sqrt(-x^2 + 4) - 6)/x^2) - 1/2*log(-
(x^2 + sqrt(-x^2 + 4) - 2)/x^2)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 48 vs. $2(12) = 24$.

Time = 1.33 (sec) , antiderivative size = 48, normalized size of antiderivative = 3.00

$$\int \frac{x}{4 - x^2 + \sqrt{4 - x^2}} dx = \frac{\log(2\sqrt{4 - x^2})}{2} - \frac{\log(2\sqrt{4 - x^2} + 2)}{2} - \frac{\log(2x^2 - 2\sqrt{4 - x^2} - 8)}{2}$$

input `integrate(x/(4-x**2+(-x**2+4)**(1/2)),x)`

output `log(2*sqrt(4 - x**2))/2 - log(2*sqrt(4 - x**2) + 2)/2 - log(2*x**2 - 2*sqrt(4 - x**2) - 8)/2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{x}{4 - x^2 + \sqrt{4 - x^2}} dx = -\log(\sqrt{-x^2 + 4} + 1)$$

input `integrate(x/(4-x^2+(-x^2+4)^(1/2)),x, algorithm="maxima")`

output `-log(sqrt(-x^2 + 4) + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{x}{4 - x^2 + \sqrt{4 - x^2}} dx = -\log(\sqrt{-x^2 + 4} + 1)$$

input `integrate(x/(4-x^2+(-x^2+4)^(1/2)),x, algorithm="giac")`

output `-log(sqrt(-x^2 + 4) + 1)`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 87, normalized size of antiderivative = 5.44

$$\int \frac{x}{4 - x^2 + \sqrt{4 - x^2}} dx = -\frac{\ln(x - \sqrt{3})}{2} - \frac{\ln\left(\frac{\sqrt{3}x + \sqrt{4 - x^2} + 4i}{x + \sqrt{3}}\right)}{2} - \frac{\ln(x + \sqrt{3})}{2} - \frac{\ln\left(\frac{-\sqrt{3}x + \sqrt{4 - x^2} + 4i}{x - \sqrt{3}}\right)}{2}$$

input `int(x/((4 - x^2)^(1/2) - x^2 + 4),x)`

output `- log(x - 3^(1/2))/2 - log((3^(1/2)*x*1i + (4 - x^2)^(1/2)*1i + 4i)/(x + 3^(1/2)))/2 - log(x + 3^(1/2))/2 - log(((4 - x^2)^(1/2)*1i - 3^(1/2)*x*1i + 4i)/(x - 3^(1/2)))/2`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.81

$$\int \frac{x}{4 - x^2 + \sqrt{4 - x^2}} dx = -\log(\sqrt{-x^2 + 4} + 1)$$

input `int(x/(4-x^2+(-x^2+4)^(1/2)),x)`

output `- log(sqrt(- x**2 + 4) + 1)`

3.115 $\int \frac{3+2x}{(-2+x)(5+x)} dx$

Optimal result	677
Mathematica [A] (verified)	677
Rubi [A] (verified)	678
Maple [A] (verified)	679
Fricas [A] (verification not implemented)	679
Sympy [A] (verification not implemented)	679
Maxima [A] (verification not implemented)	680
Giac [A] (verification not implemented)	680
Mupad [B] (verification not implemented)	680
Reduce [B] (verification not implemented)	681

Optimal result

Integrand size = 16, antiderivative size = 11

$$\int \frac{3+2x}{(-2+x)(5+x)} dx = \log(2-x) + \log(5+x)$$

output `ln(2-x)+ln(5+x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \frac{3+2x}{(-2+x)(5+x)} dx = \log(-2+x) + \log(5+x)$$

input `Integrate[(3 + 2*x)/((-2 + x)*(5 + x)), x]`

output `Log[-2 + x] + Log[5 + x]`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{2x + 3}{(x - 2)(x + 5)} dx$$

↓ 86

$$\int \left(\frac{1}{x + 5} + \frac{1}{x - 2} \right) dx$$

↓ 2009

$$\log(2 - x) + \log(x + 5)$$

input `Int[(3 + 2*x)/((-2 + x)*(5 + x)),x]`

output `Log[2 - x] + Log[5 + x]`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

method	result	size
default	$\ln((-2+x)(5+x))$	9
norman	$\ln(-2+x) + \ln(5+x)$	10
risch	$\ln(x^2 + 3x - 10)$	10
parallelrisch	$\ln(-2+x) + \ln(5+x)$	10

input `int((3+2*x)/(-2+x)/(5+x),x,method=_RETURNVERBOSE)`output `ln((-2+x)*(5+x))`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \frac{3+2x}{(-2+x)(5+x)} dx = \log(x^2 + 3x - 10)$$

input `integrate((3+2*x)/(-2+x)/(5+x),x, algorithm="fricas")`output `log(x^2 + 3*x - 10)`**Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int \frac{3+2x}{(-2+x)(5+x)} dx = \log(x^2 + 3x - 10)$$

input `integrate((3+2*x)/(-2+x)/(5+x),x)`

output `log(x**2 + 3*x - 10)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \frac{3 + 2x}{(-2 + x)(5 + x)} dx = \log(x + 5) + \log(x - 2)$$

input `integrate((3+2*x)/(-2+x)/(5+x),x, algorithm="maxima")`

output `log(x + 5) + log(x - 2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{3 + 2x}{(-2 + x)(5 + x)} dx = \log(|x + 5|) + \log(|x - 2|)$$

input `integrate((3+2*x)/(-2+x)/(5+x),x, algorithm="giac")`

output `log(abs(x + 5)) + log(abs(x - 2))`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \frac{3 + 2x}{(-2 + x)(5 + x)} dx = \ln(x^2 + 3x - 10)$$

input `int((2*x + 3)/((x - 2)*(x + 5)),x)`

output `log(3*x + x^2 - 10)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \frac{3 + 2x}{(-2 + x)(5 + x)} dx = \log(x - 2) + \log(x + 5)$$

input `int((3+2*x)/(-2+x)/(5+x),x)`

output `log(x - 2) + log(x + 5)`

3.116 $\int \frac{x}{(1+x)(2+x)(3+x)} dx$

Optimal result	682
Mathematica [A] (verified)	682
Rubi [A] (verified)	683
Maple [A] (verified)	684
Fricas [A] (verification not implemented)	684
Sympy [A] (verification not implemented)	684
Maxima [A] (verification not implemented)	685
Giac [A] (verification not implemented)	685
Mupad [B] (verification not implemented)	685
Reduce [B] (verification not implemented)	686

Optimal result

Integrand size = 17, antiderivative size = 23

$$\int \frac{x}{(1+x)(2+x)(3+x)} dx = -\frac{1}{2} \log(1+x) + 2 \log(2+x) - \frac{3}{2} \log(3+x)$$

output

```
-1/2*ln(1+x)+2*ln(2+x)-3/2*ln(3+x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{x}{(1+x)(2+x)(3+x)} dx = -\frac{1}{2} \log(1+x) + 2 \log(2+x) - \frac{3}{2} \log(3+x)$$

input

```
Integrate[x/((1+x)*(2+x)*(3+x)),x]
```

output

```
-1/2*Log[1+x]+2*Log[2+x]-3/2*Log[3+x]
```

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {165, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{(x+1)(x+2)(x+3)} dx$$

$$\downarrow 165$$

$$\int \left(\frac{2}{x+2} - \frac{3}{2(x+3)} - \frac{1}{2(x+1)} \right) dx$$

$$\downarrow 2009$$

$$-\frac{1}{2} \log(x+1) + 2 \log(x+2) - \frac{3}{2} \log(x+3)$$

input

```
Int[x/((1 + x)*(2 + x)*(3 + x)),x]
```

output

```
-1/2*Log[1 + x] + 2*Log[2 + x] - (3*Log[3 + x])/2
```

Defintions of rubi rules used

rule 165

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p*(g + h*x), x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m}, x] && (IntegersQ[m, n, p] || (IGtQ[n, 0] && IGtQ[p, 0]))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

method	result	size
default	$-\frac{\ln(1+x)}{2} + 2 \ln(2+x) - \frac{3 \ln(3+x)}{2}$	20
norman	$-\frac{\ln(1+x)}{2} + 2 \ln(2+x) - \frac{3 \ln(3+x)}{2}$	20
risch	$-\frac{\ln(1+x)}{2} + 2 \ln(2+x) - \frac{3 \ln(3+x)}{2}$	20
parallelrisch	$-\frac{\ln(1+x)}{2} + 2 \ln(2+x) - \frac{3 \ln(3+x)}{2}$	20

input `int(x/(1+x)/(2+x)/(3+x),x,method=_RETURNVERBOSE)`output `-1/2*ln(1+x)+2*ln(2+x)-3/2*ln(3+x)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{x}{(1+x)(2+x)(3+x)} dx = -\frac{3}{2} \log(x+3) + 2 \log(x+2) - \frac{1}{2} \log(x+1)$$

input `integrate(x/(1+x)/(2+x)/(3+x),x, algorithm="fricas")`output `-3/2*log(x + 3) + 2*log(x + 2) - 1/2*log(x + 1)`**Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \frac{x}{(1+x)(2+x)(3+x)} dx = -\frac{\log(x+1)}{2} + 2 \log(x+2) - \frac{3 \log(x+3)}{2}$$

input `integrate(x/(1+x)/(2+x)/(3+x),x)`

output $-\log(x + 1)/2 + 2*\log(x + 2) - 3*\log(x + 3)/2$

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{x}{(1+x)(2+x)(3+x)} dx = -\frac{3}{2} \log(x+3) + 2 \log(x+2) - \frac{1}{2} \log(x+1)$$

input `integrate(x/(1+x)/(2+x)/(3+x),x, algorithm="maxima")`

output $-3/2*\log(x + 3) + 2*\log(x + 2) - 1/2*\log(x + 1)$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.96

$$\int \frac{x}{(1+x)(2+x)(3+x)} dx = -\frac{3}{2} \log(|x+3|) + 2 \log(|x+2|) - \frac{1}{2} \log(|x+1|)$$

input `integrate(x/(1+x)/(2+x)/(3+x),x, algorithm="giac")`

output $-3/2*\log(\text{abs}(x + 3)) + 2*\log(\text{abs}(x + 2)) - 1/2*\log(\text{abs}(x + 1))$

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{x}{(1+x)(2+x)(3+x)} dx = 2 \ln(x+2) - \frac{\ln(x+1)}{2} - \frac{3 \ln(x+3)}{2}$$

input `int(x/((x + 1)*(x + 2)*(x + 3)),x)`

output $2*\log(x + 2) - \log(x + 1)/2 - (3*\log(x + 3))/2$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{x}{(1+x)(2+x)(3+x)} dx = -\frac{3\log(x+3)}{2} + 2\log(x+2) - \frac{\log(x+1)}{2}$$

input `int(x/(1+x)/(2+x)/(3+x),x)`

output `(- 3*log(x + 3) + 4*log(x + 2) - log(x + 1))/2`

3.117 $\int \frac{x}{2-3x+x^3} dx$

Optimal result	687
Mathematica [A] (verified)	687
Rubi [A] (verified)	688
Maple [A] (verified)	689
Fricas [A] (verification not implemented)	689
Sympy [A] (verification not implemented)	689
Maxima [A] (verification not implemented)	690
Giac [A] (verification not implemented)	690
Mupad [B] (verification not implemented)	690
Reduce [B] (verification not implemented)	691

Optimal result

Integrand size = 12, antiderivative size = 30

$$\int \frac{x}{2-3x+x^3} dx = \frac{1}{3(1-x)} + \frac{2}{9} \log(1-x) - \frac{2}{9} \log(2+x)$$

output

```
1/3/(1-x)+2/9*ln(1-x)-2/9*ln(2+x)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93

$$\int \frac{x}{2-3x+x^3} dx = -\frac{1}{3(-1+x)} + \frac{2}{9} \log(1-x) - \frac{2}{9} \log(2+x)$$

input

```
Integrate[x/(2 - 3*x + x^3),x]
```

output

```
-1/3*1/(-1 + x) + (2*Log[1 - x])/9 - (2*Log[2 + x])/9
```


Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{x^3 - 3x + 2} dx$$

$$\downarrow \text{2462}$$

$$\int \left(-\frac{2}{9(x+2)} + \frac{2}{9(x-1)} + \frac{1}{3(x-1)^2} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{1}{3(1-x)} + \frac{2}{9} \log(1-x) - \frac{2}{9} \log(x+2)$$

input

```
Int[x/(2 - 3*x + x^3),x]
```

output

```
1/(3*(1 - x)) + (2*Log[1 - x])/9 - (2*Log[2 + x])/9
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2462

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegr
and[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ
[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0
] && RationalFunctionQ[u, x]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.70

method	result	size
default	$-\frac{1}{3(-1+x)} + \frac{2\ln(-1+x)}{9} - \frac{2\ln(2+x)}{9}$	21
norman	$-\frac{1}{3(-1+x)} + \frac{2\ln(-1+x)}{9} - \frac{2\ln(2+x)}{9}$	21
risch	$-\frac{1}{3(-1+x)} + \frac{2\ln(-1+x)}{9} - \frac{2\ln(2+x)}{9}$	21
parallelrisc	$\frac{2\ln(-1+x)x-2\ln(2+x)x-3-2\ln(-1+x)+2\ln(2+x)}{-9+9x}$	36

input `int(x/(x^3-3*x+2),x,method=_RETURNVERBOSE)`output `-1/3/(-1+x)+2/9*ln(-1+x)-2/9*ln(2+x)`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.90

$$\int \frac{x}{2-3x+x^3} dx = -\frac{2(x-1)\log(x+2) - 2(x-1)\log(x-1) + 3}{9(x-1)}$$

input `integrate(x/(x^3-3*x+2),x, algorithm="fricas")`output `-1/9*(2*(x - 1)*log(x + 2) - 2*(x - 1)*log(x - 1) + 3)/(x - 1)`**Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \frac{x}{2-3x+x^3} dx = \frac{2\log(x-1)}{9} - \frac{2\log(x+2)}{9} - \frac{1}{3x-3}$$

input `integrate(x/(x**3-3*x+2),x)`

output `2*log(x - 1)/9 - 2*log(x + 2)/9 - 1/(3*x - 3)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.67

$$\int \frac{x}{2 - 3x + x^3} dx = -\frac{1}{3(x-1)} - \frac{2}{9} \log(x+2) + \frac{2}{9} \log(x-1)$$

input `integrate(x/(x^3-3*x+2),x, algorithm="maxima")`

output `-1/3/(x - 1) - 2/9*log(x + 2) + 2/9*log(x - 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \frac{x}{2 - 3x + x^3} dx = -\frac{1}{3(x-1)} - \frac{2}{9} \log(|x+2|) + \frac{2}{9} \log(|x-1|)$$

input `integrate(x/(x^3-3*x+2),x, algorithm="giac")`

output `-1/3/(x - 1) - 2/9*log(abs(x + 2)) + 2/9*log(abs(x - 1))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.60

$$\int \frac{x}{2 - 3x + x^3} dx = -\frac{4 \operatorname{atanh}\left(\frac{2x}{3} + \frac{1}{3}\right)}{9} - \frac{1}{3(x-1)}$$

input `int(x/(x^3 - 3*x + 2),x)`

output `-(4*atanh((2*x)/3 + 1/3))/9 - 1/(3*(x - 1))`

Reduce [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.27

$$\int \frac{x}{2 - 3x + x^3} dx = \frac{2\log(x - 1)x - 2\log(x - 1) - 2\log(x + 2)x + 2\log(x + 2) - 3x}{9x - 9}$$

input `int(x/(x^3-3*x+2),x)`

output `(2*log(x - 1)*x - 2*log(x - 1) - 2*log(x + 2)*x + 2*log(x + 2) - 3*x)/(9*(x - 1))`

$$3.118 \quad \int \frac{-6+2x+x^4}{-2x+x^2+x^3} dx$$

Optimal result	692
Mathematica [A] (verified)	692
Rubi [A] (verified)	693
Maple [A] (verified)	694
Fricas [A] (verification not implemented)	694
Sympy [A] (verification not implemented)	695
Maxima [A] (verification not implemented)	695
Giac [A] (verification not implemented)	695
Mupad [B] (verification not implemented)	696
Reduce [B] (verification not implemented)	696

Optimal result

Integrand size = 21, antiderivative size = 27

$$\int \frac{-6 + 2x + x^4}{-2x + x^2 + x^3} dx = -x + \frac{x^2}{2} - \log(1 - x) + 3 \log(x) + \log(2 + x)$$

output

```
-x+1/2*x^2-ln(1-x)+3*ln(x)+ln(2+x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{-6 + 2x + x^4}{-2x + x^2 + x^3} dx = -x + \frac{x^2}{2} - \log(1 - x) + 3 \log(x) + \log(2 + x)$$

input

```
Integrate[(-6 + 2*x + x^4)/(-2*x + x^2 + x^3),x]
```

output

```
-x + x^2/2 - Log[1 - x] + 3*Log[x] + Log[2 + x]
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2026, 2159, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^4 + 2x - 6}{x^3 + x^2 - 2x} dx \\ & \quad \downarrow \text{2026} \\ & \int \frac{x^4 + 2x - 6}{x(x^2 + x - 2)} dx \\ & \quad \downarrow \text{2159} \\ & \int \left(x + \frac{1}{1-x} + \frac{1}{x+2} + \frac{3}{x} - 1 \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{x^2}{2} - x - \log(1-x) + 3\log(x) + \log(x+2) \end{aligned}$$

input `Int[(-6 + 2*x + x^4)/(-2*x + x^2 + x^3),x]`

output `-x + x^2/2 - Log[1 - x] + 3*Log[x] + Log[2 + x]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2159

```
Int[(Pq_)*((d_) + (e_)*(x_)^(m_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p
_), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*Pq*(a + b*x + c*x^2)^p, x
], x] /; FreeQ[{a, b, c, d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89

method	result	size
default	$-x + \frac{x^2}{2} + 3 \ln(x) - \ln(-1 + x) + \ln(2 + x)$	24
norman	$-x + \frac{x^2}{2} + 3 \ln(x) - \ln(-1 + x) + \ln(2 + x)$	24
risch	$-x + \frac{x^2}{2} + 3 \ln(x) - \ln(-1 + x) + \ln(2 + x)$	24
parallelrisc	$-x + \frac{x^2}{2} + 3 \ln(x) - \ln(-1 + x) + \ln(2 + x)$	24

input

```
int((x^4+2*x-6)/(x^3+x^2-2*x),x,method=_RETURNVERBOSE)
```

output

```
-x+1/2*x^2+3*ln(x)-ln(-1+x)+ln(2+x)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \frac{-6 + 2x + x^4}{-2x + x^2 + x^3} dx = \frac{1}{2} x^2 - x + \log(x + 2) - \log(x - 1) + 3 \log(x)$$

input

```
integrate((x^4+2*x-6)/(x^3+x^2-2*x),x, algorithm="fricas")
```

output

```
1/2*x^2 - x + log(x + 2) - log(x - 1) + 3*log(x)
```

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

$$\int \frac{-6 + 2x + x^4}{-2x + x^2 + x^3} dx = \frac{x^2}{2} - x + 3 \log(x) - \log(x - 1) + \log(x + 2)$$

input `integrate((x**4+2*x-6)/(x**3+x**2-2*x),x)`output `x**2/2 - x + 3*log(x) - log(x - 1) + log(x + 2)`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \frac{-6 + 2x + x^4}{-2x + x^2 + x^3} dx = \frac{1}{2} x^2 - x + \log(x + 2) - \log(x - 1) + 3 \log(x)$$

input `integrate((x^4+2*x-6)/(x^3+x^2-2*x),x, algorithm="maxima")`output `1/2*x^2 - x + log(x + 2) - log(x - 1) + 3*log(x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \frac{-6 + 2x + x^4}{-2x + x^2 + x^3} dx = \frac{1}{2} x^2 - x + \log(|x + 2|) - \log(|x - 1|) + 3 \log(|x|)$$

input `integrate((x^4+2*x-6)/(x^3+x^2-2*x),x, algorithm="giac")`output `1/2*x^2 - x + log(abs(x + 2)) - log(abs(x - 1)) + 3*log(abs(x))`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.11

$$\int \frac{-6 + 2x + x^4}{-2x + x^2 + x^3} dx = 3 \ln(x) - x + \frac{x^2}{2} + \operatorname{atan}\left(\frac{192i}{7(28x - 40)} + \frac{9i}{7}\right) 2i$$

input `int((2*x + x^4 - 6)/(x^2 - 2*x + x^3),x)`output `atan(192i/(7*(28*x - 40)) + 9i/7)*2i - x + 3*log(x) + x^2/2`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \frac{-6 + 2x + x^4}{-2x + x^2 + x^3} dx = -\log(x - 1) + \log(x + 2) + 3 \log(x) + \frac{x^2}{2} - x$$

input `int((x^4+2*x-6)/(x^3+x^2-2*x),x)`output `(- 2*log(x - 1) + 2*log(x + 2) + 6*log(x) + x**2 - 2*x)/2`

3.119 $\int \frac{7+8x^3}{(1+x)(1+2x)^3} dx$

Optimal result	697
Mathematica [A] (verified)	697
Rubi [A] (verified)	698
Maple [A] (verified)	699
Fricas [A] (verification not implemented)	699
Sympy [A] (verification not implemented)	699
Maxima [A] (verification not implemented)	700
Giac [A] (verification not implemented)	700
Mupad [B] (verification not implemented)	700
Reduce [B] (verification not implemented)	701

Optimal result

Integrand size = 20, antiderivative size = 23

$$\int \frac{7+8x^3}{(1+x)(1+2x)^3} dx = -\frac{3}{(1+2x)^2} + \frac{3}{1+2x} + \log(1+x)$$

output `-3/(1+2*x)^2+3/(1+2*x)+ln(1+x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.04

$$\int \frac{7+8x^3}{(1+x)(1+2x)^3} dx = \frac{6x + (1+2x)^2 \log(1+x)}{(1+2x)^2}$$

input `Integrate[(7 + 8*x^3)/((1 + x)*(1 + 2*x)^3), x]`

output `(6*x + (1 + 2*x)^2*Log[1 + x])/(1 + 2*x)^2`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2123, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{8x^3 + 7}{(x + 1)(2x + 1)^3} dx$$

↓ 2123

$$\int \left(-\frac{6}{(2x + 1)^2} + \frac{12}{(2x + 1)^3} + \frac{1}{x + 1} \right) dx$$

↓ 2009

$$\frac{3}{2x + 1} - \frac{3}{(2x + 1)^2} + \log(x + 1)$$

input `Int[(7 + 8*x^3)/((1 + x)*(1 + 2*x)^3), x]`

output `-3/(1 + 2*x)^2 + 3/(1 + 2*x) + Log[1 + x]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2123 `Int[(Px_)*((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && PolyQ[Px, x] && (IntegersQ[m, n] || IGtQ[m, -2])`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.70

method	result	size
norman	$\frac{6x}{(1+2x)^2} + \ln(1+x)$	16
risch	$\frac{6x}{(1+2x)^2} + \ln(1+x)$	16
default	$-\frac{3}{(1+2x)^2} + \frac{3}{1+2x} + \ln(1+x)$	24
parallelrisch	$\frac{4\ln(1+x)x^2 + 4\ln(1+x)x + \ln(1+x) + 6x}{(1+2x)^2}$	33

input `int((8*x^3+7)/(1+x)/(1+2*x)^3,x,method=_RETURNVERBOSE)`

output `6*x/(1+2*x)^2+ln(1+x)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.39

$$\int \frac{7 + 8x^3}{(1+x)(1+2x)^3} dx = \frac{(4x^2 + 4x + 1)\log(x+1) + 6x}{4x^2 + 4x + 1}$$

input `integrate((8*x^3+7)/(1+x)/(1+2*x)^3,x, algorithm="fricas")`

output `((4*x^2 + 4*x + 1)*log(x + 1) + 6*x)/(4*x^2 + 4*x + 1)`

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{7 + 8x^3}{(1+x)(1+2x)^3} dx = \frac{6x}{4x^2 + 4x + 1} + \log(x+1)$$

input `integrate((8*x**3+7)/(1+x)/(1+2*x)**3,x)`

output $6x/(4x^2 + 4x + 1) + \log(x + 1)$

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \frac{7 + 8x^3}{(1+x)(1+2x)^3} dx = \frac{6x}{4x^2 + 4x + 1} + \log(x + 1)$$

input `integrate((8*x^3+7)/(1+x)/(1+2*x)^3,x, algorithm="maxima")`

output $6x/(4x^2 + 4x + 1) + \log(x + 1)$

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.70

$$\int \frac{7 + 8x^3}{(1+x)(1+2x)^3} dx = \frac{6x}{(2x+1)^2} + \log(|x+1|)$$

input `integrate((8*x^3+7)/(1+x)/(1+2*x)^3,x, algorithm="giac")`

output $6x/(2x + 1)^2 + \log(\text{abs}(x + 1))$

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int \frac{7 + 8x^3}{(1+x)(1+2x)^3} dx = \ln(x + 1) + \frac{6x}{(2x + 1)^2}$$

input `int((8*x^3 + 7)/((2*x + 1)^3*(x + 1)),x)`

output $\log(x + 1) + (6x)/(2x + 1)^2$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.83

$$\int \frac{7 + 8x^3}{(1+x)(1+2x)^3} dx = \frac{8 \log(x+1) x^2 + 8 \log(x+1) x + 2 \log(x+1) - 12x^2 - 3}{8x^2 + 8x + 2}$$

input `int((8*x^3+7)/(1+x)/(1+2*x)^3,x)`

output `(8*log(x + 1)*x**2 + 8*log(x + 1)*x + 2*log(x + 1) - 12*x**2 - 3)/(2*(4*x*
*2 + 4*x + 1))`

3.120 $\int \frac{1+x+4x^2}{-1+x^3} dx$

Optimal result	702
Mathematica [A] (verified)	702
Rubi [A] (verified)	703
Maple [A] (verified)	704
Fricas [A] (verification not implemented)	705
Sympy [A] (verification not implemented)	705
Maxima [A] (verification not implemented)	705
Giac [A] (verification not implemented)	706
Mupad [B] (verification not implemented)	706
Reduce [B] (verification not implemented)	706

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{1+x+4x^2}{-1+x^3} dx = 2 \log(1-x) + \log(1+x+x^2)$$

output `2*ln(1-x)+ln(x^2+x+1)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{1+x+4x^2}{-1+x^3} dx = 2 \log(1-x) + \log(1+x+x^2)$$

input `Integrate[(1 + x + 4*x^2)/(-1 + x^3), x]`

output `2*Log[1 - x] + Log[1 + x + x^2]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2414, 16, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{4x^2 + x + 1}{x^3 - 1} dx \\
 & \quad \downarrow \text{2414} \\
 & -\frac{1}{3} \int -\frac{3(2x+1)}{x^2+x+1} dx - 2 \int \frac{1}{1-x} dx \\
 & \quad \downarrow \text{16} \\
 & 2 \log(1-x) - \frac{1}{3} \int -\frac{3(2x+1)}{x^2+x+1} dx \\
 & \quad \downarrow \text{27} \\
 & \int \frac{2x+1}{x^2+x+1} dx + 2 \log(1-x) \\
 & \quad \downarrow \text{1103} \\
 & \log(x^2+x+1) + 2 \log(1-x)
 \end{aligned}$$

input `Int[(1 + x + 4*x^2)/(-1 + x^3),x]`

output `2*Log[1 - x] + Log[1 + x + x^2]`

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 27 $\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] /; \text{FreeQ}[b, x]$
- rule 1103 $\text{Int}(((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$
- rule 2414 $\text{Int}[(P2_)/((a_) + (b_)*(x_)^3), x_Symbol] \rightarrow \text{With}[\{A = \text{Coeff}[P2, x, 0], B = \text{Coeff}[P2, x, 1], C = \text{Coeff}[P2, x, 2], q = (-a/b)^{(1/3)}\}, \text{Simp}[q*((A + B*q + C*q^2)/(3*a)) \text{ Int}[1/(q - x), x], x] + \text{Simp}[q/(3*a) \text{ Int}[(q*(2*A - B*q - C*q^2) + (A + B*q - 2*C*q^2)*x)/(q^2 + q*x + x^2), x], x] /; \text{NeQ}[a*B^3 - b*A^3, 0] \ \&\& \ \text{NeQ}[A + B*q + C*q^2, 0]] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PolyQ}[P2, x, 2] \ \&\& \ \text{LtQ}[a/b, 0]$

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result
default	$2 \ln(-1 + x) + \ln(x^2 + x + 1)$
norman	$2 \ln(-1 + x) + \ln(x^2 + x + 1)$
risch	$2 \ln(-1 + x) + \ln(x^2 + x + 1)$
parallelrisch	$2 \ln(-1 + x) + \ln(x^2 + x + 1)$
meijerg	$\frac{x \left(\ln\left(1 - (x^3)^{\frac{1}{3}}\right) - \frac{\ln\left(1 + (x^3)^{\frac{1}{3}} + (x^3)^{\frac{2}{3}}\right)}{2} - \sqrt{3} \arctan\left(\frac{\sqrt{3}(x^3)^{\frac{1}{3}}}{2 + (x^3)^{\frac{1}{3}}}\right) \right)}{3(x^3)^{\frac{1}{3}}} + \frac{4 \ln(-x^3 + 1)}{3} + \frac{x^2 \left(\ln\left(1 - (x^3)^{\frac{1}{3}}\right) - \frac{\ln\left(1 + (x^3)^{\frac{1}{3}} + (x^3)^{\frac{2}{3}}\right)}{2} - \sqrt{3} \arctan\left(\frac{\sqrt{3}(x^3)^{\frac{1}{3}}}{2 + (x^3)^{\frac{1}{3}}}\right) \right)}{3(x^3)^{\frac{1}{3}}}$

input $\text{int}((4*x^2+x+1)/(x^3-1), x, \text{method}=_RETURNVERBOSE)$

output `2*ln(-1+x)+ln(x^2+x+1)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1+x+4x^2}{-1+x^3} dx = \log(x^2+x+1) + 2 \log(x-1)$$

input `integrate((4*x^2+x+1)/(x^3-1),x, algorithm="fricas")`

output `log(x^2 + x + 1) + 2*log(x - 1)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1+x+4x^2}{-1+x^3} dx = 2 \log(x-1) + \log(x^2+x+1)$$

input `integrate((4*x**2+x+1)/(x**3-1),x)`

output `2*log(x - 1) + log(x**2 + x + 1)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1+x+4x^2}{-1+x^3} dx = \log(x^2+x+1) + 2 \log(x-1)$$

input `integrate((4*x^2+x+1)/(x^3-1),x, algorithm="maxima")`

output `log(x^2 + x + 1) + 2*log(x - 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{1+x+4x^2}{-1+x^3} dx = \log(x^2+x+1) + 2 \log(|x-1|)$$

input `integrate((4*x^2+x+1)/(x^3-1),x, algorithm="giac")`output `log(x^2 + x + 1) + 2*log(abs(x - 1))`**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1+x+4x^2}{-1+x^3} dx = \ln(x^2+x+1) + 2 \ln(x-1)$$

input `int((x + 4*x^2 + 1)/(x^3 - 1),x)`output `log(x + x^2 + 1) + 2*log(x - 1)`**Reduce [B] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1+x+4x^2}{-1+x^3} dx = \log(x^2+x+1) + 2 \log(x-1)$$

input `int((4*x^2+x+1)/(x^3-1),x)`output `log(x**2 + x + 1) + 2*log(x - 1)`

3.121

$$\int \frac{x^4}{4+5x^2+x^4} dx$$

Optimal result	707
Mathematica [A] (verified)	707
Rubi [A] (verified)	708
Maple [A] (verified)	709
Fricas [A] (verification not implemented)	710
Sympy [A] (verification not implemented)	710
Maxima [A] (verification not implemented)	710
Giac [A] (verification not implemented)	711
Mupad [B] (verification not implemented)	711
Reduce [B] (verification not implemented)	711

Optimal result

Integrand size = 16, antiderivative size = 18

$$\int \frac{x^4}{4+5x^2+x^4} dx = x - \frac{8}{3} \arctan\left(\frac{x}{2}\right) + \frac{\arctan(x)}{3}$$

output

```
x-8/3*arctan(1/2*x)+1/3*arctan(x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{x^4}{4+5x^2+x^4} dx = x + \frac{8}{3} \arctan\left(\frac{2}{x}\right) + \frac{\arctan(x)}{3}$$

input

```
Integrate[x^4/(4 + 5*x^2 + x^4),x]
```

output

```
x + (8*ArcTan[2/x])/3 + ArcTan[x]/3
```

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {1442, 1480, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^4}{x^4 + 5x^2 + 4} dx \\ & \quad \downarrow 1442 \\ & x - \int \frac{5x^2 + 4}{x^4 + 5x^2 + 4} dx \\ & \quad \downarrow 1480 \\ & \frac{1}{3} \int \frac{1}{x^2 + 1} dx - \frac{16}{3} \int \frac{1}{x^2 + 4} dx + x \\ & \quad \downarrow 216 \\ & -\frac{8}{3} \arctan\left(\frac{x}{2}\right) + \frac{\arctan(x)}{3} + x \end{aligned}$$

input `Int[x^4/(4 + 5*x^2 + x^4),x]`

output `x - (8*ArcTan[x/2])/3 + ArcTan[x]/3`

Defintions of rubi rules used

rule 216

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

rule 1442

```
Int[((d_)*(x_))^(m_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol]
:> Simp[d^3*(d*x)^(m - 3)*((a + b*x^2 + c*x^4)^(p + 1)/(c*(m + 4*p + 1))),
x] - Simp[d^4/(c*(m + 4*p + 1)) Int[(d*x)^(m - 4)*Simp[a*(m - 3) + b*(m +
2*p - 1)*x^2, x]*(a + b*x^2 + c*x^4)^p, x], x] /; FreeQ[{a, b, c, d, p}, x
] && NeQ[b^2 - 4*a*c, 0] && GtQ[m, 3] && NeQ[m + 4*p + 1, 0] && IntegerQ[2*
p] && (IntegerQ[p] || IntegerQ[m])
```

rule 1480

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (b_)*(x_)^2 + (c_)*(x_)^4), x_Symbol] :
> With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[(e/2 + (2*c*d - b*e)/(2*q)) Int[1/(
b/2 - q/2 + c*x^2), x], x] + Simp[(e/2 - (2*c*d - b*e)/(2*q)) Int[1/(b/2
+ q/2 + c*x^2), x], x]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0]
&& NeQ[c*d^2 - a*e^2, 0] && PosQ[b^2 - 4*a*c]
```

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.72

method	result	size
default	$x - \frac{8 \arctan(\frac{x}{2})}{3} + \frac{\arctan(x)}{3}$	13
risch	$x - \frac{8 \arctan(\frac{x}{2})}{3} + \frac{\arctan(x)}{3}$	13
parallelrisc	$x + \frac{4i \ln(x-2i)}{3} - \frac{4i \ln(x+2i)}{3} + \frac{i \ln(x+i)}{6} - \frac{i \ln(x-i)}{6}$	35

input

```
int(x^4/(x^4+5*x^2+4),x,method=_RETURNVERBOSE)
```

output

```
x-8/3*arctan(1/2*x)+1/3*arctan(x)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

$$\int \frac{x^4}{4 + 5x^2 + x^4} dx = x - \frac{8}{3} \arctan\left(\frac{1}{2}x\right) + \frac{1}{3} \arctan(x)$$

input `integrate(x^4/(x^4+5*x^2+4),x, algorithm="fricas")`output `x - 8/3*arctan(1/2*x) + 1/3*arctan(x)`**Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{x^4}{4 + 5x^2 + x^4} dx = x - \frac{8 \operatorname{atan}\left(\frac{x}{2}\right)}{3} + \frac{\operatorname{atan}(x)}{3}$$

input `integrate(x**4/(x**4+5*x**2+4),x)`output `x - 8*atan(x/2)/3 + atan(x)/3`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

$$\int \frac{x^4}{4 + 5x^2 + x^4} dx = x - \frac{8}{3} \arctan\left(\frac{1}{2}x\right) + \frac{1}{3} \arctan(x)$$

input `integrate(x^4/(x^4+5*x^2+4),x, algorithm="maxima")`output `x - 8/3*arctan(1/2*x) + 1/3*arctan(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

$$\int \frac{x^4}{4 + 5x^2 + x^4} dx = x - \frac{8}{3} \arctan\left(\frac{1}{2}x\right) + \frac{1}{3} \arctan(x)$$

input `integrate(x^4/(x^4+5*x^2+4),x, algorithm="giac")`output `x - 8/3*arctan(1/2*x) + 1/3*arctan(x)`**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

$$\int \frac{x^4}{4 + 5x^2 + x^4} dx = x - \frac{8 \operatorname{atan}\left(\frac{x}{2}\right)}{3} + \frac{\operatorname{atan}(x)}{3}$$

input `int(x^4/(5*x^2 + x^4 + 4),x)`output `x - (8*atan(x/2))/3 + atan(x)/3`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

$$\int \frac{x^4}{4 + 5x^2 + x^4} dx = -\frac{8 \operatorname{atan}\left(\frac{x}{2}\right)}{3} + \frac{\operatorname{atan}(x)}{3} + x$$

input `int(x^4/(x^4+5*x^2+4),x)`output `(- 8*atan(x/2) + atan(x) + 3*x)/3`

3.122 $\int \frac{2+x}{x+x^2} dx$

Optimal result	712
Mathematica [A] (verified)	712
Rubi [A] (verified)	713
Maple [A] (verified)	714
Fricas [A] (verification not implemented)	714
Sympy [A] (verification not implemented)	714
Maxima [A] (verification not implemented)	715
Giac [A] (verification not implemented)	715
Mupad [B] (verification not implemented)	715
Reduce [B] (verification not implemented)	716

Optimal result

Integrand size = 11, antiderivative size = 11

$$\int \frac{2+x}{x+x^2} dx = 2 \log(x) - \log(1+x)$$

output `2*ln(x)-ln(1+x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{2+x}{x+x^2} dx = 2 \log(x) - \log(1+x)$$

input `Integrate[(2 + x)/(x + x^2),x]`

output `2*Log[x] - Log[1 + x]`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {1141, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x+2}{x^2+x} dx$$

↓ 1141

$$\int \left(\frac{2}{x} + \frac{1}{-x-1} \right) dx$$

↓ 2009

$$2\log(x) - \log(x+1)$$

input

```
Int[(2 + x)/(x + x^2), x]
```

output

```
2*Log[x] - Log[1 + x]
```

Defintions of rubi rules used

rule 1141

```
Int[((d_.) + (e_.)*(x_)^(m_.))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_
Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[1/c^p Int[ExpandIntegrand[
(d + e*x)^m*(b/2 - q/2 + c*x)^p*(b/2 + q/2 + c*x)^p, x], x], x] /; EqQ[p, -
1] || !FractionalPowerFactorQ[q] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[p,
0] && IntegerQ[m] && NiceSqrtQ[b^2 - 4*a*c]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

method	result	size
default	$2 \ln(x) - \ln(1+x)$	12
norman	$2 \ln(x) - \ln(1+x)$	12
meijerg	$2 \ln(x) - \ln(1+x)$	12
risch	$2 \ln(x) - \ln(1+x)$	12
parallelrisch	$2 \ln(x) - \ln(1+x)$	12

input `int((2+x)/(x^2+x),x,method=_RETURNVERBOSE)`output `2*ln(x)-ln(1+x)`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{2+x}{x+x^2} dx = -\log(x+1) + 2 \log(x)$$

input `integrate((2+x)/(x^2+x),x, algorithm="fricas")`output `-log(x + 1) + 2*log(x)`**Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int \frac{2+x}{x+x^2} dx = 2 \log(x) - \log(x+1)$$

input `integrate((2+x)/(x**2+x),x)`

output `2*log(x) - log(x + 1)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{2+x}{x+x^2} dx = -\log(x+1) + 2\log(x)$$

input `integrate((2+x)/(x^2+x),x, algorithm="maxima")`

output `-log(x + 1) + 2*log(x)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.18

$$\int \frac{2+x}{x+x^2} dx = -\log(|x+1|) + 2\log(|x|)$$

input `integrate((2+x)/(x^2+x),x, algorithm="giac")`

output `-log(abs(x + 1)) + 2*log(abs(x))`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{2+x}{x+x^2} dx = 2\ln(x) - \ln(x+1)$$

input `int((x + 2)/(x + x^2),x)`

output `2*log(x) - log(x + 1)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{2+x}{x+x^2} dx = -\log(x+1) + 2\log(x)$$

input `int((2+x)/(x^2+x),x)`

output `- log(x + 1) + 2*log(x)`

3.123 $\int \frac{1}{x(1+x^2)^2} dx$

Optimal result	717
Mathematica [A] (verified)	717
Rubi [A] (verified)	718
Maple [A] (verified)	719
Fricas [A] (verification not implemented)	719
Sympy [A] (verification not implemented)	720
Maxima [A] (verification not implemented)	720
Giac [A] (verification not implemented)	720
Mupad [B] (verification not implemented)	721
Reduce [B] (verification not implemented)	721

Optimal result

Integrand size = 11, antiderivative size = 24

$$\int \frac{1}{x(1+x^2)^2} dx = \frac{1}{2(1+x^2)} + \log(x) - \frac{1}{2} \log(1+x^2)$$

output `1/2/(x^2+1)+ln(x)-1/2*ln(x^2+1)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1+x^2)^2} dx = \frac{1}{2(1+x^2)} + \log(x) - \frac{1}{2} \log(1+x^2)$$

input `Integrate[1/(x*(1 + x^2)^2),x]`

output `1/(2*(1 + x^2)) + Log[x] - Log[1 + x^2]/2`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {243, 54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{x(x^2+1)^2} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{2} \int \frac{1}{x^2(x^2+1)^2} dx^2 \\ & \quad \downarrow \text{54} \\ & \frac{1}{2} \int \left(\frac{1}{x^2} - \frac{1}{(x^2+1)^2} + \frac{1}{-x^2-1} \right) dx^2 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2} \left(\frac{1}{x^2+1} + \log(x^2) - \log(x^2+1) \right) \end{aligned}$$

input

```
Int[1/(x*(1 + x^2)^2),x]
```

output

```
((1 + x^2)^(-1) + Log[x^2] - Log[1 + x^2])/2
```

Defintions of rubi rules used

rule 54

```
Int[((a_) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])
```

rule 243 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.88

method	result	size
default	$\frac{1}{2x^2+2} + \ln(x) - \frac{\ln(x^2+1)}{2}$	21
norman	$\frac{1}{2x^2+2} + \ln(x) - \frac{\ln(x^2+1)}{2}$	21
risch	$\frac{1}{2x^2+2} + \ln(x) - \frac{\ln(x^2+1)}{2}$	21
meijerg	$\frac{1}{2} + \ln(x) - \frac{x^2}{2x^2+2} - \frac{\ln(x^2+1)}{2}$	27
parallelrisch	$\frac{2x^2 \ln(x) - \ln(x^2+1)x^2 + 1 + 2 \ln(x) - \ln(x^2+1)}{2x^2+2}$	42

input `int(1/x/(x^2+1)^2,x,method=_RETURNVERBOSE)`

output `1/2/(x^2+1)+ln(x)-1/2*ln(x^2+1)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.33

$$\int \frac{1}{x(1+x^2)^2} dx = -\frac{(x^2+1)\log(x^2+1) - 2(x^2+1)\log(x) - 1}{2(x^2+1)}$$

input `integrate(1/x/(x^2+1)^2,x, algorithm="fricas")`

output `-1/2*((x^2 + 1)*log(x^2 + 1) - 2*(x^2 + 1)*log(x) - 1)/(x^2 + 1)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.79

$$\int \frac{1}{x(1+x^2)^2} dx = \log(x) - \frac{\log(x^2+1)}{2} + \frac{1}{2x^2+2}$$

input `integrate(1/x/(x**2+1)**2,x)`output `log(x) - log(x**2 + 1)/2 + 1/(2*x**2 + 2)`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1+x^2)^2} dx = \frac{1}{2(x^2+1)} - \frac{1}{2} \log(x^2+1) + \frac{1}{2} \log(x^2)$$

input `integrate(1/x/(x^2+1)^2,x, algorithm="maxima")`output `1/2/(x^2 + 1) - 1/2*log(x^2 + 1) + 1/2*log(x^2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.21

$$\int \frac{1}{x(1+x^2)^2} dx = \frac{x^2+2}{2(x^2+1)} - \frac{1}{2} \log(x^2+1) + \frac{1}{2} \log(x^2)$$

input `integrate(1/x/(x^2+1)^2,x, algorithm="giac")`output `1/2*(x^2 + 2)/(x^2 + 1) - 1/2*log(x^2 + 1) + 1/2*log(x^2)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{1}{x(1+x^2)^2} dx = \ln(x) - \frac{\ln(x^2+1)}{2} + \frac{1}{2(x^2+1)}$$

input `int(1/(x*(x^2 + 1)^2),x)`output `log(x) - log(x^2 + 1)/2 + 1/(2*(x^2 + 1))`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.92

$$\int \frac{1}{x(1+x^2)^2} dx = \frac{-\log(x^2+1)x^2 - \log(x^2+1) + 2\log(x)x^2 + 2\log(x) - x^2}{2x^2+2}$$

input `int(1/x/(x^2+1)^2,x)`output `(- log(x**2 + 1)*x**2 - log(x**2 + 1) + 2*log(x)*x**2 + 2*log(x) - x**2)/
(2*(x**2 + 1))`

3.124 $\int \frac{1}{(1+x)(2+x)^2(3+x)^3} dx$

Optimal result	722
Mathematica [A] (verified)	722
Rubi [A] (verified)	723
Maple [A] (verified)	724
Fricas [B] (verification not implemented)	724
Sympy [A] (verification not implemented)	725
Maxima [A] (verification not implemented)	725
Giac [A] (verification not implemented)	726
Mupad [B] (verification not implemented)	726
Reduce [B] (verification not implemented)	727

Optimal result

Integrand size = 16, antiderivative size = 46

$$\int \frac{1}{(1+x)(2+x)^2(3+x)^3} dx = \frac{1}{2+x} + \frac{1}{4(3+x)^2} + \frac{5}{4(3+x)} + \frac{1}{8} \log(1+x) + 2 \log(2+x) - \frac{17}{8} \log(3+x)$$

output `1/(2+x)+1/4/(3+x)^2+5/4/(3+x)+1/8*ln(1+x)+2*ln(2+x)-17/8*ln(3+x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.96

$$\int \frac{1}{(1+x)(2+x)^2(3+x)^3} dx = \frac{1}{8} \left(\frac{8}{2+x} + \frac{2}{(3+x)^2} + \frac{10}{3+x} + \log(-1-x) + 16 \log(2+x) - 17 \log(3+x) \right)$$

input `Integrate[1/((1+x)*(2+x)^2*(3+x)^3),x]`

output $(8/(2 + x) + 2/(3 + x)^2 + 10/(3 + x) + \text{Log}[-1 - x] + 16*\text{Log}[2 + x] - 17*\text{Log}[3 + x])/8$

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {99, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(x+1)(x+2)^2(x+3)^3} dx$$

↓ 99

$$\int \left(\frac{2}{x+2} - \frac{17}{8(x+3)} - \frac{1}{(x+2)^2} - \frac{5}{4(x+3)^2} - \frac{1}{2(x+3)^3} + \frac{1}{8(x+1)} \right) dx$$

↓ 2009

$$\frac{1}{x+2} + \frac{5}{4(x+3)} + \frac{1}{4(x+3)^2} + \frac{1}{8} \log(x+1) + 2 \log(x+2) - \frac{17}{8} \log(x+3)$$

input `Int[1/((1 + x)*(2 + x)^2*(3 + x)^3),x]`

output $(2 + x)^{-1} + 1/(4*(3 + x)^2) + 5/(4*(3 + x)) + \text{Log}[1 + x]/8 + 2*\text{Log}[2 + x] - (17*\text{Log}[3 + x])/8$

Defintions of rubi rules used

rule 99 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, p}, x] && IntegersQ[m, n] && (IntegerQ[p] | (GtQ[m, 0] && GeQ[n, -1]))]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.85

method	result
default	$\frac{1}{2+x} + \frac{1}{4(3+x)^2} + \frac{5}{4(3+x)} + \frac{\ln(1+x)}{8} + 2 \ln(2+x) - \frac{17 \ln(3+x)}{8}$
norman	$\frac{\frac{9}{4}x^2 + \frac{25}{2}x + 17}{(2+x)(3+x)^2} + \frac{\ln(1+x)}{8} + 2 \ln(2+x) - \frac{17 \ln(3+x)}{8}$
risch	$\frac{\frac{9}{4}x^2 + \frac{25}{2}x + 17}{(2+x)(3+x)^2} + \frac{\ln(1+x)}{8} + 2 \ln(2+x) - \frac{17 \ln(3+x)}{8}$
parallelrisc	$\frac{\ln(1+x)x^3 + 16 \ln(2+x)x^3 - 17 \ln(3+x)x^3 + 136 + 8 \ln(1+x)x^2 + 128 \ln(2+x)x^2 - 136 \ln(3+x)x^2 + 21 \ln(1+x)x + 336 \ln(2+x)x - 17 \ln(3+x)x}{8(2+x)(3+x)^2}$

input `int(1/(1+x)/(2+x)^2/(3+x)^3,x,method=_RETURNVERBOSE)`

output `1/(2+x)+1/4/(3+x)^2+5/4/(3+x)+1/8*ln(1+x)+2*ln(2+x)-17/8*ln(3+x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 83 vs. $2(38) = 76$.

Time = 0.07 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.80

$$\int \frac{1}{(1+x)(2+x)^2(3+x)^3} dx$$

$$= \frac{18x^2 - 17(x^3 + 8x^2 + 21x + 18) \log(x+3) + 16(x^3 + 8x^2 + 21x + 18) \log(x+2) + (x^3 + 8x^2 + 21x + 18) \log(x+1) + 100x + 136}{8(x^3 + 8x^2 + 21x + 18)}$$

input `integrate(1/(1+x)/(2+x)^2/(3+x)^3,x, algorithm="fricas")`

output `1/8*(18*x^2 - 17*(x^3 + 8*x^2 + 21*x + 18)*log(x + 3) + 16*(x^3 + 8*x^2 + 21*x + 18)*log(x + 2) + (x^3 + 8*x^2 + 21*x + 18)*log(x + 1) + 100*x + 136)/(x^3 + 8*x^2 + 21*x + 18)`

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1+x)(2+x)^2(3+x)^3} dx = \frac{9x^2 + 50x + 68}{4x^3 + 32x^2 + 84x + 72} + \frac{\log(x+1)}{8} + 2\log(x+2) - \frac{17\log(x+3)}{8}$$

input `integrate(1/(1+x)/(2+x)**2/(3+x)**3,x)`

output `(9*x**2 + 50*x + 68)/(4*x**3 + 32*x**2 + 84*x + 72) + log(x + 1)/8 + 2*log(x + 2) - 17*log(x + 3)/8`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1+x)(2+x)^2(3+x)^3} dx = \frac{9x^2 + 50x + 68}{4(x^3 + 8x^2 + 21x + 18)} - \frac{17}{8} \log(x+3) + 2\log(x+2) + \frac{1}{8} \log(x+1)$$

input `integrate(1/(1+x)/(2+x)^2/(3+x)^3,x, algorithm="maxima")`

output `1/4*(9*x^2 + 50*x + 68)/(x^3 + 8*x^2 + 21*x + 18) - 17/8*log(x + 3) + 2*log(x + 2) + 1/8*log(x + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.13

$$\int \frac{1}{(1+x)(2+x)^2(3+x)^3} dx = \frac{1}{x+2} - \frac{\frac{7}{x+2} + 6}{4\left(\frac{1}{x+2} + 1\right)^2} + \frac{1}{8} \log\left(\left|-\frac{1}{x+2} + 1\right|\right) - \frac{17}{8} \log\left(\left|-\frac{1}{x+2} - 1\right|\right)$$

input `integrate(1/(1+x)/(2+x)^2/(3+x)^3,x, algorithm="giac")`output `1/(x + 2) - 1/4*(7/(x + 2) + 6)/(1/(x + 2) + 1)^2 + 1/8*log(abs(-1/(x + 2) + 1)) - 17/8*log(abs(-1/(x + 2) - 1))`**Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.98

$$\int \frac{1}{(1+x)(2+x)^2(3+x)^3} dx = \frac{\ln(x+1)}{8} + 2 \ln(x+2) - \frac{17 \ln(x+3)}{8} + \frac{\frac{9x^2}{4} + \frac{25x}{2} + 17}{x^3 + 8x^2 + 21x + 18}$$

input `int(1/((x + 1)*(x + 2)^2*(x + 3)^3),x)`output `log(x + 1)/8 + 2*log(x + 2) - (17*log(x + 3))/8 + ((25*x)/2 + (9*x^2)/4 + 17)/(21*x + 8*x^2 + x^3 + 18)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 121, normalized size of antiderivative = 2.63

$$\int \frac{1}{(1+x)(2+x)^2(3+x)^3} dx$$

$$= \frac{-68 \log(x+3)x^3 - 544 \log(x+3)x^2 - 1428 \log(x+3)x - 1224 \log(x+3) + 64 \log(x+2)x^3 + 512 \log(x+2)x^2 + 1344 \log(x+2)x + 1152 \log(x+2) + 4 \log(x+1)x^3 + 32 \log(x+1)x^2 + 84 \log(x+1)x + 72 \log(x+1) - 9x^3 + 211x + 382}{(32(x^3 + 8x^2 + 21x + 18))}$$

input

```
int(1/(1+x)/(2+x)^2/(3+x)^3,x)
```

output

```
( - 68*log(x + 3)*x**3 - 544*log(x + 3)*x**2 - 1428*log(x + 3)*x - 1224*log(x + 3) + 64*log(x + 2)*x**3 + 512*log(x + 2)*x**2 + 1344*log(x + 2)*x + 1152*log(x + 2) + 4*log(x + 1)*x**3 + 32*log(x + 1)*x**2 + 84*log(x + 1)*x + 72*log(x + 1) - 9*x**3 + 211*x + 382)/(32*(x**3 + 8*x**2 + 21*x + 18))
```


3.125 $\int \frac{x}{(1+x)^2} dx$

Optimal result	728
Mathematica [A] (verified)	728
Rubi [A] (verified)	729
Maple [A] (verified)	730
Fricas [A] (verification not implemented)	730
Sympy [A] (verification not implemented)	731
Maxima [A] (verification not implemented)	731
Giac [A] (verification not implemented)	731
Mupad [B] (verification not implemented)	732
Reduce [B] (verification not implemented)	732

Optimal result

Integrand size = 7, antiderivative size = 10

$$\int \frac{x}{(1+x)^2} dx = \frac{1}{1+x} + \log(1+x)$$

output

```
1/(1+x)+ln(1+x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{x}{(1+x)^2} dx = \frac{1}{1+x} + \log(1+x)$$

input

```
Integrate[x/(1 + x)^2,x]
```

output

```
(1 + x)^(-1) + Log[1 + x]
```

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{(x+1)^2} dx$$

$$\downarrow 49$$

$$\int \left(\frac{1}{x+1} - \frac{1}{(x+1)^2} \right) dx$$

$$\downarrow 2009$$

$$\frac{1}{x+1} + \log(x+1)$$

input

```
Int[x/(1 + x)^2,x]
```

output

```
(1 + x)^(-1) + Log[1 + x]
```

Defintions of rubi rules used

rule 49

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]
```

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

method	result	size
default	$\frac{1}{1+x} + \ln(1+x)$	11
norman	$\frac{1}{1+x} + \ln(1+x)$	11
risch	$\frac{1}{1+x} + \ln(1+x)$	11
meijerg	$-\frac{x}{1+x} + \ln(1+x)$	14
parallelrisch	$\frac{\ln(1+x)x+1+\ln(1+x)}{1+x}$	19

input `int(x/(1+x)^2,x,method=_RETURNVERBOSE)`output `1/(1+x)+ln(1+x)`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.60

$$\int \frac{x}{(1+x)^2} dx = \frac{(x+1)\log(x+1)+1}{x+1}$$

input `integrate(x/(1+x)^2,x, algorithm="fricas")`output `((x + 1)*log(x + 1) + 1)/(x + 1)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{x}{(1+x)^2} dx = \log(x+1) + \frac{1}{x+1}$$

input `integrate(x/(1+x)**2,x)`

output `log(x + 1) + 1/(x + 1)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{x}{(1+x)^2} dx = \frac{1}{x+1} + \log(x+1)$$

input `integrate(x/(1+x)^2,x, algorithm="maxima")`

output `1/(x + 1) + log(x + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

$$\int \frac{x}{(1+x)^2} dx = \frac{1}{x+1} + \log(|x+1|)$$

input `integrate(x/(1+x)^2,x, algorithm="giac")`

output `1/(x + 1) + log(abs(x + 1))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{x}{(1+x)^2} dx = \ln(x+1) + \frac{1}{x+1}$$

input `int(x/(x + 1)^2,x)`

output `log(x + 1) + 1/(x + 1)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 20, normalized size of antiderivative = 2.00

$$\int \frac{x}{(1+x)^2} dx = \frac{\log(x+1)x + \log(x+1) - x}{x+1}$$

input `int(x/(1+x)^2,x)`

output `(log(x + 1)*x + log(x + 1) - x)/(x + 1)`

3.126 $\int \frac{1}{-x+x^3} dx$

Optimal result	733
Mathematica [A] (verified)	733
Rubi [A] (verified)	734
Maple [A] (verified)	735
Fricas [A] (verification not implemented)	736
Sympy [A] (verification not implemented)	736
Maxima [A] (verification not implemented)	737
Giac [A] (verification not implemented)	737
Mupad [B] (verification not implemented)	737
Reduce [B] (verification not implemented)	738

Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \frac{1}{-x+x^3} dx = -\log(x) + \frac{1}{2} \log(1-x^2)$$

output

```
-ln(x)+1/2*ln(-x^2+1)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{1}{-x+x^3} dx = -\log(x) + \frac{1}{2} \log(1-x^2)$$

input

```
Integrate[(-x + x^3)^(-1),x]
```

output

```
-Log[x] + Log[1 - x^2]/2
```

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {2026, 243, 25, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^3 - x} dx \\
 & \quad \downarrow \text{2026} \\
 & \int \frac{1}{x(x^2 - 1)} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{2} \int -\frac{1}{x^2(1 - x^2)} dx^2 \\
 & \quad \downarrow \text{25} \\
 & -\frac{1}{2} \int \frac{1}{x^2(1 - x^2)} dx^2 \\
 & \quad \downarrow \text{47} \\
 & \frac{1}{2} \left(-\int \frac{1}{x^2} dx^2 - \int \frac{1}{1 - x^2} dx^2 \right) \\
 & \quad \downarrow \text{14} \\
 & \frac{1}{2} \left(-\int \frac{1}{1 - x^2} dx^2 - \log(x^2) \right) \\
 & \quad \downarrow \text{16} \\
 & \frac{1}{2} (\log(1 - x^2) - \log(x^2))
 \end{aligned}$$

input `Int[(-x + x^3)^(-1), x]`

output `(-Log[x^2] + Log[1 - x^2])/2`

Definitions of rubi rules used

- rule 14 $\text{Int}[(a_)/(x_), x_Symbol] \rightarrow \text{Simp}[a*\text{Log}[x], x] \text{ /; FreeQ}[a, x]$
- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ /; FreeQ}[\{a, b, c\}, x]$
- rule 25 $\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[Fx, x], x]$
- rule 47 $\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_))), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x), x], x] \text{ /; FreeQ}[\{a, b, c, d\}, x]$
- rule 243 $\text{Int}[(x_)^{(m_)*((a_)+(b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{((m-1)/2)*(a + b*x)^p}, x], x, x^2], x] \text{ /; FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$
- rule 2026 $\text{Int}[(Fx_)*(Px_)^{(p_)}, x_Symbol] \rightarrow \text{With}[\{r = \text{Expon}[Px, x, \text{Min}]\}, \text{Int}[x^{(p*r)*\text{ExpandToSum}[Px/x^r, x]^p*Fx, x] \text{ /; IGtQ}[r, 0]] \text{ /; PolyQ}[Px, x] \ \&\& \ \text{IntegerQ}[p] \ \&\& \ !\text{MonomialQ}[Px, x] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ !\text{PolyQ}[u, x])]$

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
risch	$-\ln(x) + \frac{\ln(x^2-1)}{2}$	14
default	$-\ln(x) + \frac{\ln(-1+x)}{2} + \frac{\ln(1+x)}{2}$	18
norman	$-\ln(x) + \frac{\ln(-1+x)}{2} + \frac{\ln(1+x)}{2}$	18
parallelrisch	$-\ln(x) + \frac{\ln(-1+x)}{2} + \frac{\ln(1+x)}{2}$	18
meijerg	$-\ln(x) - \frac{i\pi}{2} + \frac{\ln(-x^2+1)}{2}$	20

input `int(1/(x^3-x),x,method=_RETURNVERBOSE)`

output `-ln(x)+1/2*ln(x^2-1)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \frac{1}{-x + x^3} dx = \frac{1}{2} \log(x^2 - 1) - \log(x)$$

input `integrate(1/(x^3-x),x, algorithm="fricas")`

output `1/2*log(x^2 - 1) - log(x)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.59

$$\int \frac{1}{-x + x^3} dx = -\log(x) + \frac{\log(x^2 - 1)}{2}$$

input `integrate(1/(x**3-x),x)`

output `-log(x) + log(x**2 - 1)/2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{1}{-x + x^3} dx = \frac{1}{2} \log(x + 1) + \frac{1}{2} \log(x - 1) - \log(x)$$

input `integrate(1/(x^3-x),x, algorithm="maxima")`output `1/2*log(x + 1) + 1/2*log(x - 1) - log(x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

$$\int \frac{1}{-x + x^3} dx = -\frac{1}{2} \log(x^2) + \frac{1}{2} \log(|x^2 - 1|)$$

input `integrate(1/(x^3-x),x, algorithm="giac")`output `-1/2*log(x^2) + 1/2*log(abs(x^2 - 1))`**Mupad [B] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \frac{1}{-x + x^3} dx = \frac{\ln(x^2 - 1)}{2} - \ln(x)$$

input `int(-1/(x - x^3),x)`output `log(x^2 - 1)/2 - log(x)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{1}{-x + x^3} dx = \frac{\log(x - 1)}{2} + \frac{\log(x + 1)}{2} - \log(x)$$

input `int(1/(x^3-x),x)`

output `(log(x - 1) + log(x + 1) - 2*log(x))/2`

3.127 $\int \frac{x^2}{-6+x+x^2} dx$

Optimal result	739
Mathematica [A] (verified)	739
Rubi [A] (verified)	740
Maple [A] (verified)	741
Fricas [A] (verification not implemented)	741
Sympy [A] (verification not implemented)	741
Maxima [A] (verification not implemented)	742
Giac [A] (verification not implemented)	742
Mupad [B] (verification not implemented)	742
Reduce [B] (verification not implemented)	743

Optimal result

Integrand size = 12, antiderivative size = 20

$$\int \frac{x^2}{-6+x+x^2} dx = x + \frac{4}{5} \log(2-x) - \frac{9}{5} \log(3+x)$$

output

```
x+4/5*ln(2-x)-9/5*ln(3+x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{x^2}{-6+x+x^2} dx = x + \frac{4}{5} \log(2-x) - \frac{9}{5} \log(3+x)$$

input

```
Integrate[x^2/(-6 + x + x^2),x]
```

output

```
x + (4*Log[2 - x])/5 - (9*Log[3 + x])/5
```

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {1141, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{x^2 + x - 6} dx$$

$$\downarrow \text{1141}$$

$$\int \left(-\frac{9}{5(x+3)} - \frac{4}{5(2-x)} + 1 \right) dx$$

$$\downarrow \text{2009}$$

$$x + \frac{4}{5} \log(2-x) - \frac{9}{5} \log(x+3)$$

input `Int[x^2/(-6 + x + x^2),x]`

output `x + (4*Log[2 - x])/5 - (9*Log[3 + x])/5`

Defintions of rubi rules used

rule 1141 `Int[((d_.) + (e_.)*(x_))^(m_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] :> With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[1/c^p Int[ExpandIntegrand[(d + e*x)^(m*(b/2 - q/2 + c*x)^p*(b/2 + q/2 + c*x)^p, x], x], x] /; EqQ[p, -1] || !FractionalPowerFactorQ[q]] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[p, 0] && IntegerQ[m] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

method	result	size
default	$x - \frac{9 \ln(3+x)}{5} + \frac{4 \ln(-2+x)}{5}$	15
norman	$x - \frac{9 \ln(3+x)}{5} + \frac{4 \ln(-2+x)}{5}$	15
risch	$x - \frac{9 \ln(3+x)}{5} + \frac{4 \ln(-2+x)}{5}$	15
parallelrisch	$x - \frac{9 \ln(3+x)}{5} + \frac{4 \ln(-2+x)}{5}$	15

input `int(x^2/(x^2+x-6),x,method=_RETURNVERBOSE)`output `x-9/5*ln(3+x)+4/5*ln(-2+x)`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.70

$$\int \frac{x^2}{-6+x+x^2} dx = x - \frac{9}{5} \log(x+3) + \frac{4}{5} \log(x-2)$$

input `integrate(x^2/(x^2+x-6),x, algorithm="fricas")`output `x - 9/5*log(x + 3) + 4/5*log(x - 2)`**Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{x^2}{-6+x+x^2} dx = x + \frac{4 \log(x-2)}{5} - \frac{9 \log(x+3)}{5}$$

input `integrate(x**2/(x**2+x-6),x)`

output $x + 4 \cdot \log(x - 2)/5 - 9 \cdot \log(x + 3)/5$

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.70

$$\int \frac{x^2}{-6 + x + x^2} dx = x - \frac{9}{5} \log(x + 3) + \frac{4}{5} \log(x - 2)$$

input `integrate(x^2/(x^2+x-6),x, algorithm="maxima")`

output $x - 9/5 \cdot \log(x + 3) + 4/5 \cdot \log(x - 2)$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int \frac{x^2}{-6 + x + x^2} dx = x - \frac{9}{5} \log(|x + 3|) + \frac{4}{5} \log(|x - 2|)$$

input `integrate(x^2/(x^2+x-6),x, algorithm="giac")`

output $x - 9/5 \cdot \log(\text{abs}(x + 3)) + 4/5 \cdot \log(\text{abs}(x - 2))$

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.70

$$\int \frac{x^2}{-6 + x + x^2} dx = x + \frac{4 \ln(x - 2)}{5} - \frac{9 \ln(x + 3)}{5}$$

input `int(x^2/(x + x^2 - 6),x)`

output $x + (4 \cdot \log(x - 2))/5 - (9 \cdot \log(x + 3))/5$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.70

$$\int \frac{x^2}{-6 + x + x^2} dx = \frac{4 \log(x - 2)}{5} - \frac{9 \log(x + 3)}{5} + x$$

input `int(x^2/(x^2+x-6),x)`

output `(4*log(x - 2) - 9*log(x + 3) + 5*x)/5`

3.128 $\int \frac{2+x}{4-4x+x^2} dx$

Optimal result	744
Mathematica [A] (verified)	744
Rubi [A] (verified)	745
Maple [A] (verified)	746
Fricas [A] (verification not implemented)	746
Sympy [A] (verification not implemented)	747
Maxima [A] (verification not implemented)	747
Giac [A] (verification not implemented)	747
Mupad [B] (verification not implemented)	748
Reduce [B] (verification not implemented)	748

Optimal result

Integrand size = 14, antiderivative size = 16

$$\int \frac{2+x}{4-4x+x^2} dx = \frac{4}{2-x} + \log(2-x)$$

output `4/(2-x)+ln(2-x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{2+x}{4-4x+x^2} dx = -\frac{4}{-2+x} + \log(-2+x)$$

input `Integrate[(2 + x)/(4 - 4*x + x^2), x]`

output `-4/(-2 + x) + Log[-2 + x]`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {1098, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x+2}{x^2-4x+4} dx$$

↓ 1098

$$\int \frac{x+2}{(2-x)^2} dx$$

↓ 49

$$\int \left(\frac{1}{x-2} + \frac{4}{(x-2)^2} \right) dx$$

↓ 2009

$$\frac{4}{2-x} + \log(2-x)$$

input `Int[(2 + x)/(4 - 4*x + x^2),x]`

output `4/(2 - x) + Log[2 - x]`

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 1098 `Int[((d_.) + (e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Simp[1/c^p Int[(d + e*x)^m*(b/2 + c*x)^(2*p), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.81

method	result	size
default	$-\frac{4}{-2+x} + \ln(-2+x)$	13
norman	$-\frac{4}{-2+x} + \ln(-2+x)$	13
risch	$-\frac{4}{-2+x} + \ln(-2+x)$	13
meijerg	$\frac{x}{1-\frac{x}{2}} + \ln\left(1-\frac{x}{2}\right)$	17
parallelrisch	$\frac{\ln(-2+x)x-4-2\ln(-2+x)}{-2+x}$	21

input `int((2+x)/(x^2-4*x+4),x,method=_RETURNVERBOSE)`

output `-4/(-2+x)+ln(-2+x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{2+x}{4-4x+x^2} dx = \frac{(x-2)\log(x-2)-4}{x-2}$$

input `integrate((2+x)/(x^2-4*x+4),x, algorithm="fricas")`

output `((x-2)*log(x-2)-4)/(x-2)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.50

$$\int \frac{2+x}{4-4x+x^2} dx = \log(x-2) - \frac{4}{x-2}$$

input `integrate((2+x)/(x**2-4*x+4),x)`output `log(x - 2) - 4/(x - 2)`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{2+x}{4-4x+x^2} dx = -\frac{4}{x-2} + \log(x-2)$$

input `integrate((2+x)/(x^2-4*x+4),x, algorithm="maxima")`output `-4/(x - 2) + log(x - 2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.81

$$\int \frac{2+x}{4-4x+x^2} dx = -\frac{4}{x-2} + \log(|x-2|)$$

input `integrate((2+x)/(x^2-4*x+4),x, algorithm="giac")`output `-4/(x - 2) + log(abs(x - 2))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{2+x}{4-4x+x^2} dx = \ln(x-2) - \frac{4}{x-2}$$

input `int((x + 2)/(x^2 - 4*x + 4), x)`output `log(x - 2) - 4/(x - 2)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.38

$$\int \frac{2+x}{4-4x+x^2} dx = \frac{\log(x-2)x - 2\log(x-2) - 2x}{x-2}$$

input `int((2+x)/(x^2-4*x+4), x)`output `(log(x - 2)*x - 2*log(x - 2) - 2*x)/(x - 2)`

$$3.129 \quad \int \frac{1}{(4-4x+x^2)(5-4x+x^2)} dx$$

Optimal result	749
Mathematica [A] (verified)	749
Rubi [A] (verified)	750
Maple [A] (verified)	751
Fricas [A] (verification not implemented)	752
Sympy [A] (verification not implemented)	752
Maxima [A] (verification not implemented)	752
Giac [A] (verification not implemented)	753
Mupad [B] (verification not implemented)	753
Reduce [B] (verification not implemented)	753

Optimal result

Integrand size = 21, antiderivative size = 14

$$\int \frac{1}{(4-4x+x^2)(5-4x+x^2)} dx = \frac{1}{2-x} + \arctan(2-x)$$

output `1/(2-x)-arctan(-2+x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{(4-4x+x^2)(5-4x+x^2)} dx = -\frac{1}{-2+x} + \arctan(2-x)$$

input `Integrate[1/((4 - 4*x + x^2)*(5 - 4*x + x^2)),x]`

output `-(-2 + x)^(-1) + ArcTan[2 - x]`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.43, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {1294, 1117, 1083, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(x^2 - 4x + 4)(x^2 - 4x + 5)} dx \\
 & \quad \downarrow \text{1294} \\
 & \int \frac{1}{(2-x)^2(x^2 - 4x + 5)} dx \\
 & \quad \downarrow \text{1117} \\
 & \frac{1}{2-x} - \int \frac{1}{x^2 - 4x + 5} dx \\
 & \quad \downarrow \text{1083} \\
 & 2 \int \frac{1}{-(2x-4)^2 - 4} d(2x-4) + \frac{1}{2-x} \\
 & \quad \downarrow \text{217} \\
 & \frac{1}{2-x} - \arctan\left(\frac{1}{2}(2x-4)\right)
 \end{aligned}$$

input `Int[1/((4 - 4*x + x^2)*(5 - 4*x + x^2)),x]`

output `(2 - x)^(-1) - ArcTan[(-4 + 2*x)/2]`

Definitions of rubi rules used

rule 217 $\text{Int}[(a_ + (b_ \cdot x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2])^{-1} \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /;$ $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 1083 $\text{Int}[(a_ + (b_ \cdot x_) + (c_ \cdot x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[-2 \ \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4 \cdot a \cdot c - x^2, x], x], x, b + 2 \cdot c \cdot x], x] /;$ $\text{FreeQ}\{a, b, c, x\}$

rule 1117 $\text{Int}[(d_ + (e_ \cdot x_))^{m_} \cdot ((a_ + (b_ \cdot x_) + (c_ \cdot x_)^2)^{p_}), x_Symbol] \rightarrow \text{Simp}[-2 \cdot b \cdot d \cdot (d + e \cdot x)^{m+1} \cdot ((a + b \cdot x + c \cdot x^2)^{p+1} / (d^2 \cdot (m+1) \cdot (b^2 - 4 \cdot a \cdot c))), x] + \text{Simp}[b^2 \cdot ((m+2 \cdot p+3) / (d^2 \cdot (m+1) \cdot (b^2 - 4 \cdot a \cdot c))) \ \text{Int}[(d + e \cdot x)^{m+2} \cdot (a + b \cdot x + c \cdot x^2)^p, x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, p\}, x \ \&\& \ \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0] \ \&\& \ \text{NeQ}[m+2 \cdot p+3, 0] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ (\text{IntegerQ}[2 \cdot p] \ || \ (\text{IntegerQ}[m] \ \&\& \ \text{RationalQ}[p]) \ || \ \text{IntegerQ}[(m+2 \cdot p+3)/2])$

rule 1294 $\text{Int}[(a_ + (b_ \cdot x_) + (c_ \cdot x_)^2)^{p_} \cdot ((d_ + (e_ \cdot x_) + (f_ \cdot x_)^2)^{q_}), x_Symbol] \rightarrow \text{Simp}[1/c^p \ \text{Int}[(b/2 + c \cdot x)^{2 \cdot p} \cdot (d + e \cdot x + f \cdot x^2)^q, x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f, q\}, x \ \&\& \ \text{EqQ}[b^2 - 4 \cdot a \cdot c, 0] \ \&\& \ \text{IntegerQ}[p]$

Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

method	result	size
default	$-\arctan(-2+x) - \frac{1}{-2+x}$	15
risch	$-\arctan(-2+x) - \frac{1}{-2+x}$	15
parallelrisch	$\frac{i \ln(x-2-i)x - i \ln(x-2+i)x - 2i \ln(x-2-i) + 2i \ln(x-2+i) - x}{-4+2x}$	50

input $\text{int}(1/(x^2-4 \cdot x+4)/(x^2-4 \cdot x+5), x, \text{method}=_RETURNVERBOSE)$

output `-arctan(-2+x)-1/(-2+x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.21

$$\int \frac{1}{(4-4x+x^2)(5-4x+x^2)} dx = -\frac{(x-2)\arctan(x-2)+1}{x-2}$$

input `integrate(1/(x^2-4*x+4)/(x^2-4*x+5),x, algorithm="fricas")`

output `-((x - 2)*arctan(x - 2) + 1)/(x - 2)`

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{1}{(4-4x+x^2)(5-4x+x^2)} dx = -\operatorname{atan}(x-2) - \frac{1}{x-2}$$

input `integrate(1/(x**2-4*x+4)/(x**2-4*x+5),x)`

output `-atan(x - 2) - 1/(x - 2)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{(4-4x+x^2)(5-4x+x^2)} dx = -\frac{1}{x-2} - \arctan(x-2)$$

input `integrate(1/(x^2-4*x+4)/(x^2-4*x+5),x, algorithm="maxima")`

output `-1/(x - 2) - arctan(x - 2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{(4 - 4x + x^2)(5 - 4x + x^2)} dx = -\frac{1}{x - 2} - \arctan(x - 2)$$

input `integrate(1/(x^2-4*x+4)/(x^2-4*x+5),x, algorithm="giac")`output `-1/(x - 2) - arctan(x - 2)`**Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{(4 - 4x + x^2)(5 - 4x + x^2)} dx = -\operatorname{atan}(x - 2) - \frac{1}{x - 2}$$

input `int(1/((x^2 - 4*x + 4)*(x^2 - 4*x + 5)),x)`output `- atan(x - 2) - 1/(x - 2)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.79

$$\int \frac{1}{(4 - 4x + x^2)(5 - 4x + x^2)} dx = \frac{-2\operatorname{atan}(x - 2)x + 4\operatorname{atan}(x - 2) - x}{2x - 4}$$

input `int(1/(x^2-4*x+4)/(x^2-4*x+5),x)`output `(- 2*atan(x - 2)*x + 4*atan(x - 2) - x)/(2*(x - 2))`

3.130 $\int \frac{-3+x}{2x+3x^2+x^3} dx$

Optimal result	754
Mathematica [A] (verified)	754
Rubi [A] (verified)	755
Maple [A] (verified)	756
Fricas [A] (verification not implemented)	756
Sympy [A] (verification not implemented)	757
Maxima [A] (verification not implemented)	757
Giac [A] (verification not implemented)	757
Mupad [B] (verification not implemented)	758
Reduce [B] (verification not implemented)	758

Optimal result

Integrand size = 18, antiderivative size = 21

$$\int \frac{-3+x}{2x+3x^2+x^3} dx = -\frac{3 \log(x)}{2} + 4 \log(1+x) - \frac{5}{2} \log(2+x)$$

output

```
-3/2*ln(x)+4*ln(1+x)-5/2*ln(2+x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{-3+x}{2x+3x^2+x^3} dx = -\frac{3 \log(x)}{2} + 4 \log(1+x) - \frac{5}{2} \log(2+x)$$

input

```
Integrate[(-3 + x)/(2*x + 3*x^2 + x^3),x]
```

output

```
(-3*Log[x])/2 + 4*Log[1 + x] - (5*Log[2 + x])/2
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {1979, 1200, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x-3}{x^3+3x^2+2x} dx \\ & \quad \downarrow 1979 \\ & \int \frac{x-3}{x(x^2+3x+2)} dx \\ & \quad \downarrow 1200 \\ & \int \left(\frac{4}{x+1} - \frac{5}{2(x+2)} - \frac{3}{2x} \right) dx \\ & \quad \downarrow 2009 \\ & -\frac{3 \log(x)}{2} + 4 \log(x+1) - \frac{5}{2} \log(x+2) \end{aligned}$$

input `Int[(-3 + x)/(2*x + 3*x^2 + x^3), x]`

output `(-3*Log[x])/2 + 4*Log[1 + x] - (5*Log[2 + x])/2`

Defintions of rubi rules used

rule 1200

```
Int[(((d_.) + (e_.)*(x_)^(m_.))*((f_.) + (g_.)*(x_)^(n_.)))/((a_.) + (b_.)*
(x_) + (c_.)*(x_)^2), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*((f + g*
x)^n/(a + b*x + c*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && In
tegersQ[n]
```

rule 1979

```
Int[((c_.)*(x_)^(j_.) + (b_.)*(x_)^(n_.) + (a_.)*(x_)^(q_.))^(p_.)*((A_) +
(B_.)*(x_)^(r_.)), x_Symbol] := Int[x^(p*q)*(A + B*x^(n - q))*(a + b*x^(n -
q) + c*x^(2*(n - q)))^p, x] /; FreeQ[{a, b, c, A, B, n, q}, x] && EqQ[r, n
- q] && EqQ[j, 2*n - q] && IntegerQ[p] && PosQ[n - q]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.86

method	result	size
default	$-\frac{3\ln(x)}{2} + 4\ln(1+x) - \frac{5\ln(2+x)}{2}$	18
norman	$-\frac{3\ln(x)}{2} + 4\ln(1+x) - \frac{5\ln(2+x)}{2}$	18
risch	$-\frac{3\ln(x)}{2} + 4\ln(1+x) - \frac{5\ln(2+x)}{2}$	18
parallelrisch	$-\frac{3\ln(x)}{2} + 4\ln(1+x) - \frac{5\ln(2+x)}{2}$	18

input

```
int((-3+x)/(x^3+3*x^2+2*x),x,method=_RETURNVERBOSE)
```

output

```
-3/2*ln(x)+4*ln(1+x)-5/2*ln(2+x)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{-3+x}{2x+3x^2+x^3} dx = -\frac{5}{2} \log(x+2) + 4 \log(x+1) - \frac{3}{2} \log(x)$$

input

```
integrate((-3+x)/(x^3+3*x^2+2*x),x, algorithm="fricas")
```

output

```
-5/2*log(x + 2) + 4*log(x + 1) - 3/2*log(x)
```

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.95

$$\int \frac{-3+x}{2x+3x^2+x^3} dx = -\frac{3\log(x)}{2} + 4\log(x+1) - \frac{5\log(x+2)}{2}$$

input `integrate((-3+x)/(x**3+3*x**2+2*x),x)`output `-3*log(x)/2 + 4*log(x + 1) - 5*log(x + 2)/2`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{-3+x}{2x+3x^2+x^3} dx = -\frac{5}{2}\log(x+2) + 4\log(x+1) - \frac{3}{2}\log(x)$$

input `integrate((-3+x)/(x^3+3*x^2+2*x),x, algorithm="maxima")`output `-5/2*log(x + 2) + 4*log(x + 1) - 3/2*log(x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.95

$$\int \frac{-3+x}{2x+3x^2+x^3} dx = -\frac{5}{2}\log(|x+2|) + 4\log(|x+1|) - \frac{3}{2}\log(|x|)$$

input `integrate((-3+x)/(x^3+3*x^2+2*x),x, algorithm="giac")`output `-5/2*log(abs(x + 2)) + 4*log(abs(x + 1)) - 3/2*log(abs(x))`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{-3 + x}{2x + 3x^2 + x^3} dx = 4 \ln(x + 1) - \frac{5 \ln(x + 2)}{2} - \frac{3 \ln(x)}{2}$$

input `int((x - 3)/(2*x + 3*x^2 + x^3),x)`

output `4*log(x + 1) - (5*log(x + 2))/2 - (3*log(x))/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{-3 + x}{2x + 3x^2 + x^3} dx = -\frac{5 \log(x + 2)}{2} + 4 \log(x + 1) - \frac{3 \log(x)}{2}$$

input `int((-3+x)/(x^3+3*x^2+2*x),x)`

output `(- 5*log(x + 2) + 8*log(x + 1) - 3*log(x))/2`

3.131 $\int \frac{1}{(-1+x^2)^2} dx$

Optimal result	759
Mathematica [A] (verified)	759
Rubi [A] (verified)	760
Maple [C] (verified)	761
Fricas [B] (verification not implemented)	761
Sympy [A] (verification not implemented)	762
Maxima [A] (verification not implemented)	762
Giac [A] (verification not implemented)	762
Mupad [B] (verification not implemented)	763
Reduce [B] (verification not implemented)	763

Optimal result

Integrand size = 7, antiderivative size = 21

$$\int \frac{1}{(-1+x^2)^2} dx = \frac{x}{2(1-x^2)} + \frac{\operatorname{arctanh}(x)}{2}$$

output `1/2*x/(-x^2+1)+1/2*arctanh(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.29

$$\int \frac{1}{(-1+x^2)^2} dx = \frac{1}{4} \left(-\frac{2x}{-1+x^2} - \log(1-x) + \log(1+x) \right)$$

input `Integrate[(-1 + x^2)^(-2), x]`

output `((-2*x)/(-1 + x^2) - Log[1 - x] + Log[1 + x])/4`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {215, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(x^2 - 1)^2} dx$$

$$\downarrow \text{215}$$

$$\frac{x}{2(1 - x^2)} - \frac{1}{2} \int \frac{1}{x^2 - 1} dx$$

$$\downarrow \text{220}$$

$$\frac{\operatorname{arctanh}(x)}{2} + \frac{x}{2(1 - x^2)}$$

input `Int[(-1 + x^2)^(-2), x]`

output `x/(2*(1 - x^2)) + ArcTanh[x]/2`

Defintions of rubi rules used

rule 215 `Int[((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && LtQ[p, -1] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 220 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.07 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

method	result	size
meijerg	$-\frac{i\left(\frac{2ix}{-2x^2+2}+i\operatorname{arctanh}(x)\right)}{2}$	23
norman	$-\frac{x}{2(x^2-1)} - \frac{\ln(-1+x)}{4} + \frac{\ln(1+x)}{4}$	24
risch	$-\frac{x}{2(x^2-1)} - \frac{\ln(-1+x)}{4} + \frac{\ln(1+x)}{4}$	24
default	$-\frac{1}{4(-1+x)} - \frac{\ln(-1+x)}{4} - \frac{1}{4(1+x)} + \frac{\ln(1+x)}{4}$	28
parallelrisch	$-\frac{\ln(-1+x)x^2 - \ln(1+x)x^2 - \ln(-1+x) + \ln(1+x) + 2x}{4(x^2-1)}$	41

input `int(1/(x^2-1)^2,x,method=_RETURNVERBOSE)`

output `-1/2*I*(2*I*x/(-2*x^2+2)+I*arctanh(x))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 34 vs. 2(15) = 30.

Time = 0.08 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.62

$$\int \frac{1}{(-1+x^2)^2} dx = \frac{(x^2-1)\log(x+1) - (x^2-1)\log(x-1) - 2x}{4(x^2-1)}$$

input `integrate(1/(x^2-1)^2,x, algorithm="fricas")`

output `1/4*((x^2 - 1)*log(x + 1) - (x^2 - 1)*log(x - 1) - 2*x)/(x^2 - 1)`

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.95

$$\int \frac{1}{(-1+x^2)^2} dx = -\frac{x}{2x^2-2} - \frac{\log(x-1)}{4} + \frac{\log(x+1)}{4}$$

input `integrate(1/(x**2-1)**2,x)`output `-x/(2*x**2 - 2) - log(x - 1)/4 + log(x + 1)/4`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{1}{(-1+x^2)^2} dx = -\frac{x}{2(x^2-1)} + \frac{1}{4} \log(x+1) - \frac{1}{4} \log(x-1)$$

input `integrate(1/(x^2-1)^2,x, algorithm="maxima")`output `-1/2*x/(x^2 - 1) + 1/4*log(x + 1) - 1/4*log(x - 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.19

$$\int \frac{1}{(-1+x^2)^2} dx = -\frac{x}{2(x^2-1)} + \frac{1}{4} \log(|x+1|) - \frac{1}{4} \log(|x-1|)$$

input `integrate(1/(x^2-1)^2,x, algorithm="giac")`output `-1/2*x/(x^2 - 1) + 1/4*log(abs(x + 1)) - 1/4*log(abs(x - 1))`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{1}{(-1+x^2)^2} dx = \frac{\operatorname{atanh}(x)}{2} - \frac{x}{2(x^2-1)}$$

input `int(1/(x^2 - 1)^2,x)`output `atanh(x)/2 - x/(2*(x^2 - 1))`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.95

$$\int \frac{1}{(-1+x^2)^2} dx = \frac{-\log(x-1)x^2 + \log(x-1) + \log(x+1)x^2 - \log(x+1) - 2x}{4x^2 - 4}$$

input `int(1/(x^2-1)^2,x)`output `(- log(x - 1)*x**2 + log(x - 1) + log(x + 1)*x**2 - log(x + 1) - 2*x)/(4*(x**2 - 1))`

3.132 $\int \frac{1+x}{-1+x^3} dx$

Optimal result	764
Mathematica [A] (verified)	764
Rubi [A] (verified)	765
Maple [A] (verified)	766
Fricas [A] (verification not implemented)	767
Sympy [A] (verification not implemented)	767
Maxima [A] (verification not implemented)	767
Giac [A] (verification not implemented)	768
Mupad [B] (verification not implemented)	768
Reduce [B] (verification not implemented)	768

Optimal result

Integrand size = 11, antiderivative size = 22

$$\int \frac{1+x}{-1+x^3} dx = \frac{2}{3} \log(1-x) - \frac{1}{3} \log(1+x+x^2)$$

output `2/3*ln(1-x)-1/3*ln(x^2+x+1)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1+x}{-1+x^3} dx = \frac{2}{3} \log(1-x) - \frac{1}{3} \log(1+x+x^2)$$

input `Integrate[(1 + x)/(-1 + x^3),x]`

output `(2*Log[1 - x])/3 - Log[1 + x + x^2]/3`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {2400, 16, 25, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x+1}{x^3-1} dx \\
 & \quad \downarrow \text{2400} \\
 & \frac{1}{3} \int -\frac{2x+1}{x^2+x+1} dx - \frac{2}{3} \int \frac{1}{1-x} dx \\
 & \quad \downarrow \text{16} \\
 & \frac{1}{3} \int -\frac{2x+1}{x^2+x+1} dx + \frac{2}{3} \log(1-x) \\
 & \quad \downarrow \text{25} \\
 & \frac{2}{3} \log(1-x) - \frac{1}{3} \int \frac{2x+1}{x^2+x+1} dx \\
 & \quad \downarrow \text{1103} \\
 & \frac{2}{3} \log(1-x) - \frac{1}{3} \log(x^2+x+1)
 \end{aligned}$$

input `Int[(1 + x)/(-1 + x^3),x]`

output `(2*Log[1 - x])/3 - Log[1 + x + x^2]/3`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 2400 `Int[((A_) + (B_.)*(x_))/((a_) + (b_.)*(x_)^3), x_Symbol] := With[{r = Numerator[Rt[-a/b, 3]], s = Denominator[Rt[-a/b, 3]]}, Simp[r*(B*r + A*s)/(3*a*s) Int[1/(r - s*x), x], x] - Simp[r/(3*a*s) Int[(r*(B*r - 2*A*s) - s*(B*r + A*s)*x)/(r^2 + r*s*x + s^2*x^2), x], x]] /; FreeQ[{a, b, A, B}, x] && NeQ[a*B^3 - b*A^3, 0] && NegQ[a/b]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

method	result
default	$\frac{2 \ln(-1+x)}{3} - \frac{\ln(x^2+x+1)}{3}$
norman	$\frac{2 \ln(-1+x)}{3} - \frac{\ln(x^2+x+1)}{3}$
risch	$\frac{2 \ln(-1+x)}{3} - \frac{\ln(x^2+x+1)}{3}$
paralelrisch	$\frac{2 \ln(-1+x)}{3} - \frac{\ln(x^2+x+1)}{3}$
meijerg	$\frac{x \left(\ln\left(1 - (x^3)^{\frac{1}{3}}\right) - \frac{\ln\left(1 + (x^3)^{\frac{1}{3}} + (x^3)^{\frac{2}{3}}\right)}{2} - \sqrt{3} \arctan\left(\frac{\sqrt{3}(x^3)^{\frac{1}{3}}}{2 + (x^3)^{\frac{1}{3}}}\right) \right)}{3(x^3)^{\frac{1}{3}}} + \frac{x^2 \left(\ln\left(1 - (x^3)^{\frac{1}{3}}\right) - \frac{\ln\left(1 + (x^3)^{\frac{1}{3}} + (x^3)^{\frac{2}{3}}\right)}{2} + \sqrt{3} \arctan\left(\frac{\sqrt{3}(x^3)^{\frac{1}{3}}}{2 + (x^3)^{\frac{1}{3}}}\right) \right)}{3(x^3)^{\frac{2}{3}}}$

input `int((1+x)/(x^3-1),x,method=_RETURNVERBOSE)`

output `2/3*ln(-1+x)-1/3*ln(x^2+x+1)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \frac{1+x}{-1+x^3} dx = -\frac{1}{3} \log(x^2+x+1) + \frac{2}{3} \log(x-1)$$

input `integrate((1+x)/(x^3-1),x, algorithm="fricas")`

output `-1/3*log(x^2 + x + 1) + 2/3*log(x - 1)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

$$\int \frac{1+x}{-1+x^3} dx = \frac{2 \log(x-1)}{3} - \frac{\log(x^2+x+1)}{3}$$

input `integrate((1+x)/(x**3-1),x)`

output `2*log(x - 1)/3 - log(x**2 + x + 1)/3`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \frac{1+x}{-1+x^3} dx = -\frac{1}{3} \log(x^2+x+1) + \frac{2}{3} \log(x-1)$$

input `integrate((1+x)/(x^3-1),x, algorithm="maxima")`

output `-1/3*log(x^2 + x + 1) + 2/3*log(x - 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

$$\int \frac{1+x}{-1+x^3} dx = -\frac{1}{3} \log(x^2+x+1) + \frac{2}{3} \log(|x-1|)$$

input `integrate((1+x)/(x^3-1),x, algorithm="giac")`output `-1/3*log(x^2 + x + 1) + 2/3*log(abs(x - 1))`**Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \frac{1+x}{-1+x^3} dx = \frac{2 \ln(x-1)}{3} - \frac{\ln(x^2+x+1)}{3}$$

input `int((x + 1)/(x^3 - 1),x)`output `(2*log(x - 1))/3 - log(x + x^2 + 1)/3`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \frac{1+x}{-1+x^3} dx = -\frac{\log(x^2+x+1)}{3} + \frac{2 \log(x-1)}{3}$$

input `int((1+x)/(x^3-1),x)`output `(- log(x**2 + x + 1) + 2*log(x - 1))/3`

3.133

$$\int \frac{1+x^4}{x(1+x^2)^2} dx$$

Optimal result	769
Mathematica [A] (verified)	769
Rubi [A] (verified)	770
Maple [A] (verified)	771
Fricas [A] (verification not implemented)	771
Sympy [A] (verification not implemented)	772
Maxima [A] (verification not implemented)	772
Giac [A] (verification not implemented)	772
Mupad [B] (verification not implemented)	773
Reduce [B] (verification not implemented)	773

Optimal result

Integrand size = 16, antiderivative size = 10

$$\int \frac{1+x^4}{x(1+x^2)^2} dx = \frac{1}{1+x^2} + \log(x)$$

output `1/(x^2+1)+ln(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1+x^4}{x(1+x^2)^2} dx = \frac{1}{1+x^2} + \log(x)$$

input `Integrate[(1 + x^4)/(x*(1 + x^2)^2), x]`

output `(1 + x^2)^(-1) + Log[x]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.80, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {1579, 522, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^4 + 1}{x(x^2 + 1)^2} dx \\ & \quad \downarrow \text{1579} \\ & \frac{1}{2} \int \frac{x^4 + 1}{x^2(x^2 + 1)^2} dx^2 \\ & \quad \downarrow \text{522} \\ & \frac{1}{2} \int \left(\frac{1}{x^2} - \frac{2}{(x^2 + 1)^2} \right) dx^2 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2} \left(\frac{2}{x^2 + 1} + \log(x^2) \right) \end{aligned}$$

input `Int[(1 + x^4)/(x*(1 + x^2)^2),x]`

output `(2/(1 + x^2) + Log[x^2])/2`

Defintions of rubi rules used

rule 522

```
Int[((e_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] :> Int[ExpandIntegrand[(e*x)^m*(c + d*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && IGtQ[p, 0]
```

rule 1579 `Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (c_.)*(x_)^4)^(p_.), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, c, d, e, p, q}, x] && IntegerQ[(m + 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

method	result	size
default	$\frac{1}{x^2+1} + \ln(x)$	11
norman	$\frac{1}{x^2+1} + \ln(x)$	11
risch	$\frac{1}{x^2+1} + \ln(x)$	11
parallelrisch	$\frac{x^2 \ln(x)+1+\ln(x)}{x^2+1}$	19
meijerg	$-\frac{x^2}{2(x^2+1)} + \frac{1}{2} + \ln(x) - \frac{x^2}{2x^2+2}$	31

input `int((x^4+1)/x/(x^2+1)^2,x,method=_RETURNVERBOSE)`

output `1/(x^2+1)+ln(x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.80

$$\int \frac{1+x^4}{x(1+x^2)^2} dx = \frac{(x^2+1)\log(x)+1}{x^2+1}$$

input `integrate((x^4+1)/x/(x^2+1)^2,x, algorithm="fricas")`

output `((x^2 + 1)*log(x) + 1)/(x^2 + 1)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{1+x^4}{x(1+x^2)^2} dx = \log(x) + \frac{1}{x^2+1}$$

input `integrate((x**4+1)/x/(x**2+1)**2,x)`output `log(x) + 1/(x**2 + 1)`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.40

$$\int \frac{1+x^4}{x(1+x^2)^2} dx = \frac{1}{x^2+1} + \frac{1}{2} \log(x^2)$$

input `integrate((x^4+1)/x/(x^2+1)^2,x, algorithm="maxima")`output `1/(x^2 + 1) + 1/2*log(x^2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.40

$$\int \frac{1+x^4}{x(1+x^2)^2} dx = \frac{1}{x^2+1} + \frac{1}{2} \log(x^2)$$

input `integrate((x^4+1)/x/(x^2+1)^2,x, algorithm="giac")`output `1/(x^2 + 1) + 1/2*log(x^2)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1 + x^4}{x(1 + x^2)^2} dx = \ln(x) + \frac{1}{x^2 + 1}$$

input `int((x^4 + 1)/(x*(x^2 + 1)^2),x)`output `log(x) + 1/(x^2 + 1)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 22, normalized size of antiderivative = 2.20

$$\int \frac{1 + x^4}{x(1 + x^2)^2} dx = \frac{\log(x) x^2 + \log(x) - x^2}{x^2 + 1}$$

input `int((x^4+1)/x/(x^2+1)^2,x)`output `(log(x)*x**2 + log(x) - x**2)/(x**2 + 1)`

3.134 $\int \frac{1}{-2x^3+x^4} dx$

Optimal result	774
Mathematica [A] (verified)	774
Rubi [A] (verified)	775
Maple [A] (verified)	776
Fricas [A] (verification not implemented)	776
Sympy [A] (verification not implemented)	777
Maxima [A] (verification not implemented)	777
Giac [A] (verification not implemented)	777
Mupad [B] (verification not implemented)	778
Reduce [B] (verification not implemented)	778

Optimal result

Integrand size = 11, antiderivative size = 31

$$\int \frac{1}{-2x^3+x^4} dx = \frac{1}{4x^2} + \frac{1}{4x} + \frac{1}{8} \log(2-x) - \frac{\log(x)}{8}$$

output $1/4/x^2+1/4/x+1/8*\ln(2-x)-1/8*\ln(x)$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{1}{-2x^3+x^4} dx = \frac{1}{4x^2} + \frac{1}{4x} + \frac{1}{8} \log(2-x) - \frac{\log(x)}{8}$$

input `Integrate[(-2*x^3 + x^4)^(-1),x]`

output $1/(4*x^2) + 1/(4*x) + \text{Log}[2 - x]/8 - \text{Log}[x]/8$

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {2026, 54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{x^4 - 2x^3} dx \\ & \quad \downarrow \text{2026} \\ & \int \frac{1}{(x-2)x^3} dx \\ & \quad \downarrow \text{54} \\ & \int \left(-\frac{1}{2x^3} - \frac{1}{4x^2} - \frac{1}{8x} + \frac{1}{8(x-2)} \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{1}{4x^2} + \frac{1}{4x} + \frac{1}{8} \log(2-x) - \frac{\log(x)}{8} \end{aligned}$$

input

```
Int[(-2*x^3 + x^4)^(-1), x]
```

output

```
1/(4*x^2) + 1/(4*x) + Log[2 - x]/8 - Log[x]/8
```

Defintions of rubi rules used

rule 54

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])]
```

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```


rule 2026

```
Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p
*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && Integ
erQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])
```

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.68

method	result	size
norman	$\frac{\frac{1}{4} + \frac{x}{4}}{x^2} - \frac{\ln(x)}{8} + \frac{\ln(-2+x)}{8}$	21
risch	$\frac{\frac{1}{4} + \frac{x}{4}}{x^2} - \frac{\ln(x)}{8} + \frac{\ln(-2+x)}{8}$	21
default	$\frac{1}{4x^2} + \frac{1}{4x} - \frac{\ln(x)}{8} + \frac{\ln(-2+x)}{8}$	22
parallelrisc	$-\frac{x^2 \ln(x) - \ln(-2+x)x^2 - 2 - 2x}{8x^2}$	26
meijerg	$\frac{1}{4x^2} + \frac{1}{4x} - \frac{\ln(x)}{8} + \frac{\ln(2)}{8} - \frac{i\pi}{8} + \frac{\ln(1-\frac{x}{2})}{8}$	32

input

```
int(1/(x^4-2*x^3),x,method=_RETURNVERBOSE)
```

output

```
(1/4+1/4*x)/x^2-1/8*ln(x)+1/8*ln(-2+x)
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.81

$$\int \frac{1}{-2x^3 + x^4} dx = \frac{x^2 \log(x - 2) - x^2 \log(x) + 2x + 2}{8x^2}$$

input

```
integrate(1/(x^4-2*x^3),x, algorithm="fricas")
```

output

```
1/8*(x^2*log(x - 2) - x^2*log(x) + 2*x + 2)/x^2
```

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.61

$$\int \frac{1}{-2x^3 + x^4} dx = -\frac{\log(x)}{8} + \frac{\log(x-2)}{8} + \frac{x+1}{4x^2}$$

input `integrate(1/(x**4-2*x**3),x)`output `-log(x)/8 + log(x - 2)/8 + (x + 1)/(4*x**2)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.61

$$\int \frac{1}{-2x^3 + x^4} dx = \frac{x+1}{4x^2} + \frac{1}{8} \log(x-2) - \frac{1}{8} \log(x)$$

input `integrate(1/(x^4-2*x^3),x, algorithm="maxima")`output `1/4*(x + 1)/x^2 + 1/8*log(x - 2) - 1/8*log(x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.68

$$\int \frac{1}{-2x^3 + x^4} dx = \frac{x+1}{4x^2} + \frac{1}{8} \log(|x-2|) - \frac{1}{8} \log(|x|)$$

input `integrate(1/(x^4-2*x^3),x, algorithm="giac")`output `1/4*(x + 1)/x^2 + 1/8*log(abs(x - 2)) - 1/8*log(abs(x))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.52

$$\int \frac{1}{-2x^3 + x^4} dx = \frac{\frac{x}{4} + \frac{1}{4}}{x^2} - \frac{\operatorname{atanh}(x - 1)}{4}$$

input `int(-1/(2*x^3 - x^4),x)`output `(x/4 + 1/4)/x^2 - atanh(x - 1)/4`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.81

$$\int \frac{1}{-2x^3 + x^4} dx = \frac{\log(x - 2) x^2 - \log(x) x^2 + 2x + 2}{8x^2}$$

input `int(1/(x^4-2*x^3),x)`output `(log(x - 2)*x**2 - log(x)*x**2 + 2*x + 2)/(8*x**2)`

3.135 $\int \frac{1-x^3}{x(1+x^2)} dx$

Optimal result	779
Mathematica [A] (verified)	779
Rubi [A] (verified)	780
Maple [A] (verified)	781
Fricas [A] (verification not implemented)	781
Sympy [A] (verification not implemented)	781
Maxima [A] (verification not implemented)	782
Giac [A] (verification not implemented)	782
Mupad [B] (verification not implemented)	782
Reduce [B] (verification not implemented)	783

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{1-x^3}{x(1+x^2)} dx = -x + \arctan(x) + \log(x) - \frac{1}{2} \log(1+x^2)$$

output

```
-x+arctan(x)+ln(x)-1/2*ln(x^2+1)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{1-x^3}{x(1+x^2)} dx = -x + \arctan(x) + \log(x) - \frac{1}{2} \log(1+x^2)$$

input

```
Integrate[(1 - x^3)/(x*(1 + x^2)),x]
```

output

```
-x + ArcTan[x] + Log[x] - Log[1 + x^2]/2
```

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {2333, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1-x^3}{x(x^2+1)} dx$$

$$\downarrow \text{2333}$$

$$\int \left(\frac{1-x}{x^2+1} + \frac{1}{x} - 1 \right) dx$$

$$\downarrow \text{2009}$$

$$\arctan(x) - \frac{1}{2} \log(x^2+1) - x + \log(x)$$

input

```
Int[(1 - x^3)/(x*(1 + x^2)),x]
```

output

```
-x + ArcTan[x] + Log[x] - Log[1 + x^2]/2
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2333

```
Int[(Pq_)*((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[
ExpandIntegrand[(c*x)^m*Pq*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x]
&& PolyQ[Pq, x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

method	result	size
default	$-x + \arctan(x) + \ln(x) - \frac{\ln(x^2+1)}{2}$	17
meijerg	$-x + \arctan(x) + \ln(x) - \frac{\ln(x^2+1)}{2}$	17
risch	$-x + \arctan(x) + \ln(x) - \frac{\ln(x^2+1)}{2}$	17
parallelrisch	$-x + \ln(x) - \frac{\ln(x-i)}{2} - \frac{i \ln(x-i)}{2} - \frac{\ln(x+i)}{2} + \frac{i \ln(x+i)}{2}$	37

input `int((-x^3+1)/x/(x^2+1),x,method=_RETURNVERBOSE)`output `-x+arctan(x)+ln(x)-1/2*ln(x^2+1)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int \frac{1-x^3}{x(1+x^2)} dx = -x + \arctan(x) - \frac{1}{2} \log(x^2+1) + \log(x)$$

input `integrate((-x^3+1)/x/(x^2+1),x, algorithm="fricas")`output `-x + arctan(x) - 1/2*log(x^2 + 1) + log(x)`**Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int \frac{1-x^3}{x(1+x^2)} dx = -x + \log(x) - \frac{\log(x^2+1)}{2} + \operatorname{atan}(x)$$

input `integrate((-x**3+1)/x/(x**2+1),x)`

output `-x + log(x) - log(x**2 + 1)/2 + atan(x)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int \frac{1 - x^3}{x(1 + x^2)} dx = -x + \arctan(x) - \frac{1}{2} \log(x^2 + 1) + \log(x)$$

input `integrate((-x^3+1)/x/(x^2+1),x, algorithm="maxima")`

output `-x + arctan(x) - 1/2*log(x^2 + 1) + log(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

$$\int \frac{1 - x^3}{x(1 + x^2)} dx = -x + \arctan(x) - \frac{1}{2} \log(x^2 + 1) + \log(|x|)$$

input `integrate((-x^3+1)/x/(x^2+1),x, algorithm="giac")`

output `-x + arctan(x) - 1/2*log(x^2 + 1) + log(abs(x))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.33

$$\int \frac{1 - x^3}{x(1 + x^2)} dx = \ln(x) - x + \ln(x - i) \left(-\frac{1}{2} - \frac{1}{2}i\right) + \ln(x + i) \left(-\frac{1}{2} + \frac{1}{2}i\right)$$

input `int(-(x^3 - 1)/(x*(x^2 + 1)),x)`

output `log(x) - log(x - 1i)*(1/2 + 1i/2) - log(x + 1i)*(1/2 - 1i/2) - x`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int \frac{1-x^3}{x(1+x^2)} dx = \operatorname{atan}(x) - \frac{\log(x^2+1)}{2} + \log(x) - x$$

input `int((-x^3+1)/x/(x^2+1),x)`

output `(2*atan(x) - log(x**2 + 1) + 2*log(x) - 2*x)/2`

3.136 $\int \frac{1}{-1+x^4} dx$

Optimal result	784
Mathematica [A] (verified)	784
Rubi [A] (verified)	785
Maple [A] (verified)	786
Fricas [A] (verification not implemented)	786
Sympy [A] (verification not implemented)	787
Maxima [A] (verification not implemented)	787
Giac [B] (verification not implemented)	787
Mupad [B] (verification not implemented)	788
Reduce [B] (verification not implemented)	788

Optimal result

Integrand size = 7, antiderivative size = 13

$$\int \frac{1}{-1+x^4} dx = -\frac{\arctan(x)}{2} - \frac{\operatorname{arctanh}(x)}{2}$$

output `-1/2*arctan(x)-1/2*arctanh(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.92

$$\int \frac{1}{-1+x^4} dx = -\frac{\arctan(x)}{2} + \frac{1}{4} \log(1-x) - \frac{1}{4} \log(1+x)$$

input `Integrate[(-1 + x^4)^(-1), x]`

output `-1/2*ArcTan[x] + Log[1 - x]/4 - Log[1 + x]/4`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{x^4 - 1} dx \\ & \quad \downarrow \text{756} \\ & -\frac{1}{2} \int \frac{1}{1-x^2} dx - \frac{1}{2} \int \frac{1}{x^2+1} dx \\ & \quad \downarrow \text{216} \\ & -\frac{1}{2} \int \frac{1}{1-x^2} dx - \frac{\arctan(x)}{2} \\ & \quad \downarrow \text{219} \\ & -\frac{\arctan(x)}{2} - \frac{\operatorname{arctanh}(x)}{2} \end{aligned}$$

input `Int[(-1 + x^4)^(-1), x]`

output `-1/2*ArcTan[x] - ArcTanh[x]/2`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 756

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2
]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x]
+ Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a
/b, 0]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

method	result	size
default	$-\frac{\arctan(x)}{2} - \frac{\operatorname{arctanh}(x)}{2}$	10
risch	$-\frac{\arctan(x)}{2} - \frac{\ln(1+x)}{4} + \frac{\ln(-1+x)}{4}$	18
parallelrisc	$\frac{i \ln(x-i)}{4} - \frac{i \ln(x+i)}{4} - \frac{\ln(1+x)}{4} + \frac{\ln(-1+x)}{4}$	30
meijerg	$\frac{x \left(\ln \left(1 - (x^4)^{\frac{1}{4}} \right) - \ln \left(1 + (x^4)^{\frac{1}{4}} \right) - 2 \arctan \left((x^4)^{\frac{1}{4}} \right) \right)}{4(x^4)^{\frac{1}{4}}}$	38

input

```
int(1/(x^4-1),x,method=_RETURNVERBOSE)
```

output

```
-1/2*arctan(x)-1/2*arctanh(x)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.31

$$\int \frac{1}{-1+x^4} dx = -\frac{1}{2} \arctan(x) - \frac{1}{4} \log(x+1) + \frac{1}{4} \log(x-1)$$

input

```
integrate(1/(x^4-1),x, algorithm="fricas")
```

output

```
-1/2*arctan(x) - 1/4*log(x + 1) + 1/4*log(x - 1)
```

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.31

$$\int \frac{1}{-1+x^4} dx = \frac{\log(x-1)}{4} - \frac{\log(x+1)}{4} - \frac{\operatorname{atan}(x)}{2}$$

input `integrate(1/(x**4-1),x)`

output `log(x - 1)/4 - log(x + 1)/4 - atan(x)/2`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.31

$$\int \frac{1}{-1+x^4} dx = -\frac{1}{2} \arctan(x) - \frac{1}{4} \log(x+1) + \frac{1}{4} \log(x-1)$$

input `integrate(1/(x^4-1),x, algorithm="maxima")`

output `-1/2*arctan(x) - 1/4*log(x + 1) + 1/4*log(x - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. 2(9) = 18.

Time = 0.12 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.46

$$\int \frac{1}{-1+x^4} dx = -\frac{1}{2} \arctan(x) - \frac{1}{4} \log(|x+1|) + \frac{1}{4} \log(|x-1|)$$

input `integrate(1/(x^4-1),x, algorithm="giac")`

output `-1/2*arctan(x) - 1/4*log(abs(x + 1)) + 1/4*log(abs(x - 1))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int \frac{1}{-1+x^4} dx = -\frac{\operatorname{atan}(x)}{2} - \frac{\operatorname{atanh}(x)}{2}$$

input `int(1/(x^4 - 1),x)`

output `- atan(x)/2 - atanh(x)/2`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.31

$$\int \frac{1}{-1+x^4} dx = -\frac{\operatorname{atan}(x)}{2} + \frac{\log(x-1)}{4} - \frac{\log(x+1)}{4}$$

input `int(1/(x^4-1),x)`

output `(- 2*atan(x) + log(x - 1) - log(x + 1))/4`

3.137 $\int \frac{1}{1+x^4} dx$

Optimal result	789
Mathematica [A] (verified)	789
Rubi [A] (verified)	790
Maple [C] (verified)	793
Fricas [A] (verification not implemented)	793
Sympy [A] (verification not implemented)	794
Maxima [A] (verification not implemented)	794
Giac [A] (verification not implemented)	795
Mupad [B] (verification not implemented)	795
Reduce [B] (verification not implemented)	796

Optimal result

Integrand size = 7, antiderivative size = 85

$$\int \frac{1}{1+x^4} dx = -\frac{\arctan(1-\sqrt{2}x)}{2\sqrt{2}} + \frac{\arctan(1+\sqrt{2}x)}{2\sqrt{2}} - \frac{\log(1-\sqrt{2}x+x^2)}{4\sqrt{2}} + \frac{\log(1+\sqrt{2}x+x^2)}{4\sqrt{2}}$$

output

```
1/4*arctan(-1+x*2^(1/2))*2^(1/2)+1/4*arctan(1+x*2^(1/2))*2^(1/2)-1/8*ln(1+x^2-x*2^(1/2))*2^(1/2)+1/8*ln(1+x^2+x*2^(1/2))*2^(1/2)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.75

$$\int \frac{1}{1+x^4} dx = \frac{-2 \arctan(1-\sqrt{2}x) + 2 \arctan(1+\sqrt{2}x) - \log(1-\sqrt{2}x+x^2) + \log(1+\sqrt{2}x+x^2)}{4\sqrt{2}}$$

input

```
Integrate[(1 + x^4)^(-1),x]
```

output

$$(-2*\text{ArcTan}[1 - \text{Sqrt}[2]*x] + 2*\text{ArcTan}[1 + \text{Sqrt}[2]*x] - \text{Log}[1 - \text{Sqrt}[2]*x + x^2] + \text{Log}[1 + \text{Sqrt}[2]*x + x^2])/(4*\text{Sqrt}[2])$$
Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.06, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.143$, Rules used = {755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{x^4 + 1} dx \\ & \quad \downarrow 755 \\ & \frac{1}{2} \int \frac{1 - x^2}{x^4 + 1} dx + \frac{1}{2} \int \frac{x^2 + 1}{x^4 + 1} dx \\ & \quad \downarrow 1476 \\ & \frac{1}{2} \left(\frac{1}{2} \int \frac{1}{x^2 - \sqrt{2}x + 1} dx + \frac{1}{2} \int \frac{1}{x^2 + \sqrt{2}x + 1} dx \right) + \frac{1}{2} \int \frac{1 - x^2}{x^4 + 1} dx \\ & \quad \downarrow 1082 \\ & \frac{1}{2} \int \frac{1 - x^2}{x^4 + 1} dx + \frac{1}{2} \left(\frac{\int \frac{1}{-(1 - \sqrt{2}x)^2 - 1} d(1 - \sqrt{2}x)}{\sqrt{2}} - \frac{\int \frac{1}{-(\sqrt{2}x + 1)^2 - 1} d(\sqrt{2}x + 1)}{\sqrt{2}} \right) \\ & \quad \downarrow 217 \\ & \frac{1}{2} \int \frac{1 - x^2}{x^4 + 1} dx + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}x + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}x)}{\sqrt{2}} \right) \\ & \quad \downarrow 1479 \\ & \frac{1}{2} \left(-\frac{\int -\frac{\sqrt{2} - 2x}{x^2 - \sqrt{2}x + 1} dx}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}x + 1)}{x^2 + \sqrt{2}x + 1} dx}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}x + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}x)}{\sqrt{2}} \right) \\ & \quad \downarrow 25 \end{aligned}$$

$$\frac{1}{2} \left(\int \frac{\sqrt{2}-2x}{x^2-\sqrt{2}x+1} dx + \int \frac{\sqrt{2}(\sqrt{2}x+1)}{x^2+\sqrt{2}x+1} dx \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}x+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}x)}{\sqrt{2}} \right)$$

↓ 27

$$\frac{1}{2} \left(\int \frac{\sqrt{2}-2x}{x^2-\sqrt{2}x+1} dx + \frac{1}{2} \int \frac{\sqrt{2}x+1}{x^2+\sqrt{2}x+1} dx \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}x+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}x)}{\sqrt{2}} \right)$$

↓ 1103

$$\frac{1}{2} \left(\frac{\arctan(\sqrt{2}x+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}x)}{\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\log(x^2+\sqrt{2}x+1)}{2\sqrt{2}} - \frac{\log(x^2-\sqrt{2}x+1)}{2\sqrt{2}} \right)$$

input `Int[(1 + x^4)^(-1), x]`

output `(-(ArcTan[1 - Sqrt[2]*x]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*x]/Sqrt[2])/2 + (-1/2*Log[1 - Sqrt[2]*x + x^2]/Sqrt[2] + Log[1 + Sqrt[2]*x + x^2]/(2*Sqrt[2]))/2`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.08 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.26

method	result
risch	$\frac{\sum_{-R=\text{RootOf}(-Z^4+1)} \frac{\ln(x-\frac{R}{-R^3})}{-R^3}}{4}$
default	$\frac{\sqrt{2} \left(\ln\left(\frac{1+x^2+x\sqrt{2}}{1+x^2-x\sqrt{2}}\right) + 2 \arctan(1+x\sqrt{2}) + 2 \arctan(-1+x\sqrt{2}) \right)}{8}$
meijerg	$-\frac{x\sqrt{2} \ln\left(1-\sqrt{2}(x^4)^{\frac{1}{4}}+\sqrt{x^4}\right)}{8(x^4)^{\frac{1}{4}}} + \frac{x\sqrt{2} \arctan\left(\frac{\sqrt{2}(x^4)^{\frac{1}{4}}}{2-\sqrt{2}(x^4)^{\frac{1}{4}}}\right)}{4(x^4)^{\frac{1}{4}}} + \frac{x\sqrt{2} \ln\left(1+\sqrt{2}(x^4)^{\frac{1}{4}}+\sqrt{x^4}\right)}{8(x^4)^{\frac{1}{4}}} + \frac{x\sqrt{2} \arctan\left(\frac{\sqrt{2}(x^4)^{\frac{1}{4}}}{2+\sqrt{2}(x^4)^{\frac{1}{4}}}\right)}{4(x^4)^{\frac{1}{4}}}$

input `int(1/(x^4+1),x,method=_RETURNVERBOSE)`

output `1/4*sum(1/_R^3*ln(x-_R),_R=RootOf(_Z^4+1))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.71

$$\int \frac{1}{1+x^4} dx = \frac{1}{4} \sqrt{2} \arctan(\sqrt{2}x+1) + \frac{1}{4} \sqrt{2} \arctan(\sqrt{2}x-1) \\ + \frac{1}{8} \sqrt{2} \log(x^2 + \sqrt{2}x + 1) - \frac{1}{8} \sqrt{2} \log(x^2 - \sqrt{2}x + 1)$$

input `integrate(1/(x^4+1),x, algorithm="fricas")`

output `1/4*sqrt(2)*arctan(sqrt(2)*x + 1) + 1/4*sqrt(2)*arctan(sqrt(2)*x - 1) + 1/8*sqrt(2)*log(x^2 + sqrt(2)*x + 1) - 1/8*sqrt(2)*log(x^2 - sqrt(2)*x + 1)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.86

$$\int \frac{1}{1+x^4} dx = -\frac{\sqrt{2} \log(x^2 - \sqrt{2}x + 1)}{8} + \frac{\sqrt{2} \log(x^2 + \sqrt{2}x + 1)}{8} \\ + \frac{\sqrt{2} \operatorname{atan}(\sqrt{2}x - 1)}{4} + \frac{\sqrt{2} \operatorname{atan}(\sqrt{2}x + 1)}{4}$$

input `integrate(1/(x**4+1),x)`output `-sqrt(2)*log(x**2 - sqrt(2)*x + 1)/8 + sqrt(2)*log(x**2 + sqrt(2)*x + 1)/8
+ sqrt(2)*atan(sqrt(2)*x - 1)/4 + sqrt(2)*atan(sqrt(2)*x + 1)/4`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.85

$$\int \frac{1}{1+x^4} dx = \frac{1}{4} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}(2x + \sqrt{2})\right) + \frac{1}{4} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}(2x - \sqrt{2})\right) \\ + \frac{1}{8} \sqrt{2} \log(x^2 + \sqrt{2}x + 1) - \frac{1}{8} \sqrt{2} \log(x^2 - \sqrt{2}x + 1)$$

input `integrate(1/(x^4+1),x, algorithm="maxima")`output `1/4*sqrt(2)*arctan(1/2*sqrt(2)*(2*x + sqrt(2))) + 1/4*sqrt(2)*arctan(1/2*sqrt(2)*(2*x - sqrt(2))) + 1/8*sqrt(2)*log(x^2 + sqrt(2)*x + 1) - 1/8*sqrt(2)*log(x^2 - sqrt(2)*x + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.85

$$\int \frac{1}{1+x^4} dx = \frac{1}{4} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}(2x + \sqrt{2})\right) + \frac{1}{4} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}(2x - \sqrt{2})\right) \\ + \frac{1}{8} \sqrt{2} \log(x^2 + \sqrt{2}x + 1) - \frac{1}{8} \sqrt{2} \log(x^2 - \sqrt{2}x + 1)$$

input `integrate(1/(x^4+1),x, algorithm="giac")`

output `1/4*sqrt(2)*arctan(1/2*sqrt(2)*(2*x + sqrt(2))) + 1/4*sqrt(2)*arctan(1/2*sqrt(2)*(2*x - sqrt(2))) + 1/8*sqrt(2)*log(x^2 + sqrt(2)*x + 1) - 1/8*sqrt(2)*log(x^2 - sqrt(2)*x + 1)`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.39

$$\int \frac{1}{1+x^4} dx = \sqrt{2} \operatorname{atan}\left(\sqrt{2}x \left(\frac{1}{2} - \frac{1}{2}i\right)\right) \left(\frac{1}{4} + \frac{1}{4}i\right) \\ + \sqrt{2} \operatorname{atan}\left(\sqrt{2}x \left(\frac{1}{2} + \frac{1}{2}i\right)\right) \left(\frac{1}{4} - \frac{1}{4}i\right)$$

input `int(1/(x^4 + 1),x)`

output `2^(1/2)*atan(2^(1/2)*x*(1/2 - 1i/2))*(1/4 + 1i/4) + 2^(1/2)*atan(2^(1/2)*x*(1/2 + 1i/2))*(1/4 - 1i/4)`

Reduce [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.66

$$\int \frac{1}{1+x^4} dx$$

$$= \frac{\sqrt{2} \left(-2 \operatorname{atan} \left(\frac{\sqrt{2}-2x}{\sqrt{2}} \right) + 2 \operatorname{atan} \left(\frac{\sqrt{2}+2x}{\sqrt{2}} \right) - \log(-\sqrt{2}x + x^2 + 1) + \log(\sqrt{2}x + x^2 + 1) \right)}{8}$$

input

`int(1/(x^4+1),x)`

output

```
(sqrt(2)*(-2*atan((sqrt(2)-2*x)/sqrt(2))+2*atan((sqrt(2)+2*x)/sqrt(2))-log(-sqrt(2)*x+x**2+1)+log(sqrt(2)*x+x**2+1)))/8
```

3.138

$$\int \frac{x^2}{(2+2x+x^2)^2} dx$$

Optimal result	797
Mathematica [A] (verified)	797
Rubi [A] (verified)	798
Maple [A] (verified)	799
Fricas [A] (verification not implemented)	799
Sympy [A] (verification not implemented)	800
Maxima [A] (verification not implemented)	800
Giac [A] (verification not implemented)	800
Mupad [B] (verification not implemented)	801
Reduce [B] (verification not implemented)	801

Optimal result

Integrand size = 14, antiderivative size = 23

$$\int \frac{x^2}{(2+2x+x^2)^2} dx = -\frac{x(2+x)}{2(2+2x+x^2)} + \arctan(1+x)$$

output `-1/2*x*(2+x)/(x^2+2*x+2)+arctan(1+x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int \frac{x^2}{(2+2x+x^2)^2} dx = \frac{1}{2+2x+x^2} + \arctan(1+x)$$

input `Integrate[x^2/(2 + 2*x + x^2)^2,x]`

output `(2 + 2*x + x^2)^(-1) + ArcTan[1 + x]`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {1153, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{(x^2 + 2x + 2)^2} dx$$

$$\downarrow 1153$$

$$\int \frac{1}{x^2 + 2x + 2} dx - \frac{x(x + 2)}{2(x^2 + 2x + 2)}$$

$$\downarrow 1082$$

$$- \int \frac{1}{-(x + 1)^2 - 1} d(x + 1) - \frac{x(x + 2)}{2(x^2 + 2x + 2)}$$

$$\downarrow 217$$

$$\arctan(x + 1) - \frac{x(x + 2)}{2(x^2 + 2x + 2)}$$

input `Int[x^2/(2 + 2*x + x^2)^2,x]`

output `-1/2*(x*(2 + x))/(2 + 2*x + x^2) + ArcTan[1 + x]`

Defintions of rubi rules used

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])`

rule 1082

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; Fre
eQ[{a, b, c}, x]
```

rule 1153

```
Int[((d_) + (e_)*(x_)^(m_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_S
ymbol] := Simp[(d + e*x)^(m - 1)*(d*b - 2*a*e + (2*c*d - b*e)*x)*((a + b*x
+ c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c))), x] - Simp[2*(2*p + 3)*((c*d^2 -
b*d*e + a*e^2)/((p + 1)*(b^2 - 4*a*c)) Int[(d + e*x)^(m - 2)*(a + b*x +
c*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[m + 2*p + 2, 0]
&& LtQ[p, -1]
```

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.70

method	result	size
default	$\frac{1}{x^2+2x+2} + \arctan(1+x)$	16
risch	$\frac{1}{x^2+2x+2} + \arctan(1+x)$	16
parallelrisch	$-\frac{i \ln(x+1-i)x^2 - i \ln(x+1+i)x^2 + 2i \ln(x+1-i)x - 2i \ln(x+1+i)x - 2 + 2i \ln(x+1-i) - 2i \ln(x+1+i)}{2(x^2+2x+2)}$	77

input

```
int(x^2/(x^2+2*x+2)^2,x,method=_RETURNVERBOSE)
```

output

```
1/(x^2+2*x+2)+arctan(1+x)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.13

$$\int \frac{x^2}{(2+2x+x^2)^2} dx = \frac{(x^2+2x+2) \arctan(x+1) + 1}{x^2+2x+2}$$

input

```
integrate(x^2/(x^2+2*x+2)^2,x, algorithm="fricas")
```


output `((x^2 + 2*x + 2)*arctan(x + 1) + 1)/(x^2 + 2*x + 2)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.61

$$\int \frac{x^2}{(2 + 2x + x^2)^2} dx = \operatorname{atan}(x + 1) + \frac{1}{x^2 + 2x + 2}$$

input `integrate(x**2/(x**2+2*x+2)**2,x)`

output `atan(x + 1) + 1/(x**2 + 2*x + 2)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int \frac{x^2}{(2 + 2x + x^2)^2} dx = \frac{1}{x^2 + 2x + 2} + \operatorname{arctan}(x + 1)$$

input `integrate(x^2/(x^2+2*x+2)^2,x, algorithm="maxima")`

output `1/(x^2 + 2*x + 2) + arctan(x + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int \frac{x^2}{(2 + 2x + x^2)^2} dx = \frac{1}{x^2 + 2x + 2} + \operatorname{arctan}(x + 1)$$

input `integrate(x^2/(x^2+2*x+2)^2,x, algorithm="giac")`

output `1/(x^2 + 2*x + 2) + arctan(x + 1)`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int \frac{x^2}{(2 + 2x + x^2)^2} dx = \operatorname{atan}(x + 1) + \frac{1}{x^2 + 2x + 2}$$

input `int(x^2/(2*x + x^2 + 2)^2,x)`output `atan(x + 1) + 1/(2*x + x^2 + 2)`**Reduce [B] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.48

$$\int \frac{x^2}{(2 + 2x + x^2)^2} dx = \frac{\operatorname{atan}(x + 1) x^2 + 2\operatorname{atan}(x + 1) x + 2\operatorname{atan}(x + 1) + 1}{x^2 + 2x + 2}$$

input `int(x^2/(x^2+2*x+2)^2,x)`output `(atan(x + 1)*x**2 + 2*atan(x + 1)*x + 2*atan(x + 1) + 1)/(x**2 + 2*x + 2)`

$$3.139 \quad \int \frac{-1+4x^5}{(1+x+x^5)^2} dx$$

Optimal result	802
Mathematica [A] (verified)	802
Rubi [A] (verified)	803
Maple [A] (verified)	803
Fricas [A] (verification not implemented)	804
Sympy [A] (verification not implemented)	804
Maxima [A] (verification not implemented)	805
Giac [A] (verification not implemented)	805
Mupad [B] (verification not implemented)	805
Reduce [B] (verification not implemented)	806

Optimal result

Integrand size = 16, antiderivative size = 11

$$\int \frac{-1+4x^5}{(1+x+x^5)^2} dx = -\frac{x}{1+x+x^5}$$

output `-x/(x^5+x+1)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{-1+4x^5}{(1+x+x^5)^2} dx = -\frac{x}{1+x+x^5}$$

input `Integrate[(-1 + 4*x^5)/(1 + x + x^5)^2,x]`

output `-(x/(1 + x + x^5))`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.062$, Rules used = {2021}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{4x^5 - 1}{(x^5 + x + 1)^2} dx$$

\downarrow 2021
 $-\frac{x}{x^5 + x + 1}$

input `Int[(-1 + 4*x^5)/(1 + x + x^5)^2,x]`

output `-(x/(1 + x + x^5))`

Defintions of rubi rules used

rule 2021 `Int[(Pp_)*(Qq_)^(m_.), x_Symbol] := With[{p = Expon[Pp, x], q = Expon[Qq, x]}, Simp[Coeff[Pp, x, p]*x^(p - q + 1)*(Qq^(m + 1))/((p + m*q + 1)*Coeff[Qq, x, q]), x] /; NeQ[p + m*q + 1, 0] && EqQ[(p + m*q + 1)*Coeff[Qq, x, q]*Pp, Coeff[Pp, x, p]*x^(p - q)*((p - q + 1)*Qq + (m + 1)*x*D[Qq, x])] /; FreeQ[m, x] && PolyQ[Pp, x] && PolyQ[Qq, x] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

method	result	size
gospers	$-\frac{x}{x^5+x+1}$	12
norman	$-\frac{x}{x^5+x+1}$	12
risch	$-\frac{x}{x^5+x+1}$	12
parallelrisch	$-\frac{x}{x^5+x+1}$	12
orering	$-\frac{(x^2+x+1)(x^3-x^2+1)x}{(x^5+x+1)^2}$	28
default	$-\frac{-3x^2+5x-1}{7(x^3-x^2+1)} + \frac{-3x-1}{7x^2+7x+7}$	41

input `int((4*x^5-1)/(x^5+x+1)^2,x,method=_RETURNVERBOSE)`

output `-x/(x^5+x+1)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{-1 + 4x^5}{(1 + x + x^5)^2} dx = -\frac{x}{x^5 + x + 1}$$

input `integrate((4*x^5-1)/(x^5+x+1)^2,x, algorithm="fricas")`

output `-x/(x^5 + x + 1)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int \frac{-1 + 4x^5}{(1 + x + x^5)^2} dx = -\frac{x}{x^5 + x + 1}$$

input `integrate((4*x**5-1)/(x**5+x+1)**2,x)`

output `-x/(x**5 + x + 1)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{-1 + 4x^5}{(1 + x + x^5)^2} dx = -\frac{x}{x^5 + x + 1}$$

input `integrate((4*x^5-1)/(x^5+x+1)^2,x, algorithm="maxima")`

output `-x/(x^5 + x + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{-1 + 4x^5}{(1 + x + x^5)^2} dx = -\frac{x}{x^5 + x + 1}$$

input `integrate((4*x^5-1)/(x^5+x+1)^2,x, algorithm="giac")`

output `-x/(x^5 + x + 1)`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{-1 + 4x^5}{(1 + x + x^5)^2} dx = -\frac{x}{x^5 + x + 1}$$

input `int((4*x^5 - 1)/(x + x^5 + 1)^2,x)`

output `-x/(x + x^5 + 1)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.27

$$\int \frac{-1 + 4x^5}{(1 + x + x^5)^2} dx = \frac{x^5 + 1}{x^5 + x + 1}$$

input `int((4*x^5-1)/(x^5+x+1)^2,x)`

output `(x**5 + 1)/(x**5 + x + 1)`

3.140 $\int \frac{1}{5 - \cos(x) + 2 \sin(x)} dx$

Optimal result	807
Mathematica [A] (verified)	807
Rubi [A] (verified)	808
Maple [A] (verified)	809
Fricas [A] (verification not implemented)	810
Sympy [A] (verification not implemented)	810
Maxima [A] (verification not implemented)	810
Giac [A] (verification not implemented)	811
Mupad [B] (verification not implemented)	811
Reduce [B] (verification not implemented)	811

Optimal result

Integrand size = 12, antiderivative size = 45

$$\int \frac{1}{5 - \cos(x) + 2 \sin(x)} dx = \frac{x}{2\sqrt{5}} + \frac{\arctan\left(\frac{2 \cos(x) + \sin(x)}{5 + 2\sqrt{5} - \cos(x) + 2 \sin(x)}\right)}{\sqrt{5}}$$

output

$1/10*x*5^{(1/2)}+1/5*\arctan((2*\cos(x)+\sin(x))/(5-\cos(x)+2*\sin(x)+2*5^{(1/2)}))$
 $*5^{(1/2)}$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.51

$$\int \frac{1}{5 - \cos(x) + 2 \sin(x)} dx = \frac{\arctan\left(\frac{1+3 \tan(\frac{x}{2})}{\sqrt{5}}\right)}{\sqrt{5}}$$

input

`Integrate[(5 - Cos[x] + 2*Sin[x])^(-1),x]`

output

`ArcTan[(1 + 3*Tan[x/2])/Sqrt[5]]/Sqrt[5]`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.58, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3603, 1083, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{2 \sin(x) - \cos(x) + 5} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{2 \sin(x) - \cos(x) + 5} dx \\
 & \quad \downarrow \text{3603} \\
 & 2 \int \frac{1}{6 \tan^2\left(\frac{x}{2}\right) + 4 \tan\left(\frac{x}{2}\right) + 4} d \tan\left(\frac{x}{2}\right) \\
 & \quad \downarrow \text{1083} \\
 & -4 \int \frac{1}{-(12 \tan\left(\frac{x}{2}\right) + 4)^2 - 80} d\left(12 \tan\left(\frac{x}{2}\right) + 4\right) \\
 & \quad \downarrow \text{217} \\
 & \frac{\arctan\left(\frac{12 \tan\left(\frac{x}{2}\right) + 4}{4\sqrt{5}}\right)}{\sqrt{5}}
 \end{aligned}$$

input `Int[(5 - Cos[x] + 2*Sin[x])^(-1),x]`

output `ArcTan[(4 + 12*Tan[x/2])/(4*Sqrt[5])]/Sqrt[5]`

Definitions of rubi rules used

rule 217 $\text{Int}[(a_ + (b_ \cdot)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2])^{-1} \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[-a, 2])], x] \text{ ; FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 1083 $\text{Int}[(a_ + (b_ \cdot)(x_) + (c_ \cdot)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[-2 \ \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4 \cdot a \cdot c - x^2, x], x], x, b + 2 \cdot c \cdot x], x] \text{ ; FreeQ}\{a, b, c, x\}$

rule 3042 $\text{Int}[u_ , x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$

rule 3603 $\text{Int}[(\cos[(d_ \cdot) + (e_ \cdot)(x_)] \cdot (b_ \cdot) + (a_) + (c_ \cdot) \cdot \sin[(d_ \cdot) + (e_ \cdot)(x_)])^{-1}, x_Symbol] \rightarrow \text{Module}\{f = \text{FreeFactors}[\text{Tan}[(d + e \cdot x)/2], x]\}, \text{Simp}[2 \cdot (f/e) \ \text{Subst}[\text{Int}[1/(a + b + 2 \cdot c \cdot f \cdot x + (a - b) \cdot f^2 \cdot x^2), x], x, \text{Tan}[(d + e \cdot x)/2]/f], x]] \text{ ; FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{NeQ}[a^2 - b^2 - c^2, 0]$

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.44

method	result	size
default	$\frac{\sqrt{5} \arctan\left(\frac{(6 \tan(\frac{x}{2}) + 2)\sqrt{5}}{10}\right)}{5}$	20
risch	$\frac{i\sqrt{5} \ln\left(e^{ix} - 1 + 2i + \frac{4i\sqrt{5}}{5} - \frac{2\sqrt{5}}{5}\right)}{10} - \frac{i\sqrt{5} \ln\left(e^{ix} - 1 + 2i - \frac{4i\sqrt{5}}{5} + \frac{2\sqrt{5}}{5}\right)}{10}$	56

input `int(1/(5-cos(x)+2*sin(x)),x,method=_RETURNVERBOSE)`

output `1/5*5^(1/2)*arctan(1/10*(6*tan(1/2*x)+2)*5^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.80

$$\int \frac{1}{5 - \cos(x) + 2 \sin(x)} dx = \frac{1}{10} \sqrt{5} \arctan \left(-\frac{\sqrt{5} \cos(x) - 2 \sqrt{5} \sin(x) - \sqrt{5}}{2(2 \cos(x) + \sin(x))} \right)$$

input `integrate(1/(5-cos(x)+2*sin(x)),x, algorithm="fricas")`output `1/10*sqrt(5)*arctan(-1/2*(sqrt(5)*cos(x) - 2*sqrt(5)*sin(x) - sqrt(5))/(2*cos(x) + sin(x)))`**Sympy [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.87

$$\int \frac{1}{5 - \cos(x) + 2 \sin(x)} dx = \frac{\sqrt{5} \left(\operatorname{atan} \left(\frac{3\sqrt{5} \tan(\frac{x}{2})}{5} + \frac{\sqrt{5}}{5} \right) + \pi \left\lfloor \frac{\frac{x}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right)}{5}$$

input `integrate(1/(5-cos(x)+2*sin(x)),x)`output `sqrt(5)*(atan(3*sqrt(5)*tan(x/2)/5 + sqrt(5)/5) + pi*floor((x/2 - pi/2)/pi))/5`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.51

$$\int \frac{1}{5 - \cos(x) + 2 \sin(x)} dx = \frac{1}{5} \sqrt{5} \arctan \left(\frac{1}{5} \sqrt{5} \left(\frac{3 \sin(x)}{\cos(x) + 1} + 1 \right) \right)$$

input `integrate(1/(5-cos(x)+2*sin(x)),x, algorithm="maxima")`output `1/5*sqrt(5)*arctan(1/5*sqrt(5)*(3*sin(x)/(cos(x) + 1) + 1))`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.04

$$\int \frac{1}{5 - \cos(x) + 2 \sin(x)} dx$$

$$= \frac{1}{10} \sqrt{5} \left(x + 2 \arctan \left(-\frac{\sqrt{5} \sin(x) - \cos(x) - 3 \sin(x) - 1}{\sqrt{5} \cos(x) + \sqrt{5} - 3 \cos(x) + \sin(x) + 3} \right) \right)$$

input `integrate(1/(5-cos(x)+2*sin(x)),x, algorithm="giac")`output `1/10*sqrt(5)*(x + 2*arctan(-(sqrt(5)*sin(x) - cos(x) - 3*sin(x) - 1)/(sqrt(5)*cos(x) + sqrt(5) - 3*cos(x) + sin(x) + 3)))`**Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.47

$$\int \frac{1}{5 - \cos(x) + 2 \sin(x)} dx = \frac{\sqrt{20} \operatorname{atan} \left(\frac{3\sqrt{20} \tan(\frac{x}{2})}{10} + \frac{\sqrt{20}}{10} \right)}{10}$$

input `int(1/(2*sin(x) - cos(x) + 5),x)`output `(20^(1/2)*atan((3*20^(1/2)*tan(x/2))/10 + 20^(1/2)/10))/10`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.40

$$\int \frac{1}{5 - \cos(x) + 2 \sin(x)} dx = \frac{\sqrt{5} \operatorname{atan} \left(\frac{3 \tan(\frac{x}{2}) + 1}{\sqrt{5}} \right)}{5}$$

input `int(1/(5-cos(x)+2*sin(x)),x)`

output $(\sqrt{5} \cdot \operatorname{atan}((3 \cdot \tan(x/2) + 1)/\sqrt{5}))/5$

3.141 $\int \frac{1}{1+a \cos(x)} dx$

Optimal result	813
Mathematica [A] (verified)	813
Rubi [A] (verified)	814
Maple [A] (verified)	815
Fricas [A] (verification not implemented)	815
Sympy [B] (verification not implemented)	816
Maxima [F(-2)]	816
Giac [A] (verification not implemented)	817
Mupad [B] (verification not implemented)	817
Reduce [B] (verification not implemented)	818

Optimal result

Integrand size = 8, antiderivative size = 37

$$\int \frac{1}{1+a \cos(x)} dx = \frac{2 \arctan\left(\frac{\sqrt{1-a} \tan\left(\frac{x}{2}\right)}{\sqrt{1+a}}\right)}{\sqrt{1-a^2}}$$

output `2*arctan((1-a)^(1/2)*tan(1/2*x)/(1+a)^(1/2))/(-a^2+1)^(1/2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.84

$$\int \frac{1}{1+a \cos(x)} dx = \frac{2 \operatorname{arctanh}\left(\frac{(-1+a) \tan\left(\frac{x}{2}\right)}{\sqrt{-1+a^2}}\right)}{\sqrt{-1+a^2}}$$

input `Integrate[(1 + a*Cos[x])^(-1),x]`

output `(2*ArcTanh[((-1 + a)*Tan[x/2])/Sqrt[-1 + a^2]])/Sqrt[-1 + a^2]`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{a \cos(x) + 1} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{a \sin\left(x + \frac{\pi}{2}\right) + 1} dx \\ & \quad \downarrow \text{3138} \\ & 2 \int \frac{1}{(1-a) \tan^2\left(\frac{x}{2}\right) + a + 1} d \tan\left(\frac{x}{2}\right) \\ & \quad \downarrow \text{218} \\ & \frac{2 \arctan\left(\frac{\sqrt{1-a} \tan\left(\frac{x}{2}\right)}{\sqrt{a+1}}\right)}{\sqrt{1-a^2}} \end{aligned}$$

input

```
Int[(1 + a*Cos[x])^(-1),x]
```

output

```
(2*ArcTan[(Sqrt[1 - a]*Tan[x/2])/Sqrt[1 + a]])/Sqrt[1 - a^2]
```

Defintions of rubi rules used

rule 218

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 3138

```
Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] :> With[{
e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b +
(a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x]
&& NeQ[a^2 - b^2, 0]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.81

method	result	size
default	$\frac{2 \operatorname{arctanh}\left(\frac{(a-1) \tan\left(\frac{x}{2}\right)}{\sqrt{(1+a)(a-1)}}\right)}{\sqrt{(1+a)(a-1)}}$	30
risch	$\frac{\ln\left(\frac{e^{ix} + \frac{ia^2 + \sqrt{a^2-1}-i}{a\sqrt{a^2-1}}}{\sqrt{a^2-1}}\right)}{\sqrt{a^2-1}} - \frac{\ln\left(\frac{e^{ix} + \frac{-ia^2 + \sqrt{a^2-1}+i}{a\sqrt{a^2-1}}}{\sqrt{a^2-1}}\right)}{\sqrt{a^2-1}}$	87

input

```
int(1/(1+a*cos(x)),x,method=_RETURNVERBOSE)
```

output

```
2/((1+a)*(a-1))^(1/2)*arctanh((a-1)*tan(1/2*x)/((1+a)*(a-1))^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 111, normalized size of antiderivative = 3.00

$$\int \frac{1}{1 + a \cos(x)} dx = \left[\frac{\log\left(-\frac{(a^2-2) \cos(x)^2 - 2\sqrt{a^2-1}(a+\cos(x)) \sin(x) - 2a^2 - 2a \cos(x) + 1}{a^2 \cos(x)^2 + 2a \cos(x) + 1}\right)}{2\sqrt{a^2-1}}, \right. \\ \left. - \frac{\sqrt{-a^2+1} \arctan\left(\frac{\sqrt{-a^2+1}(a+\cos(x))}{(a^2-1) \sin(x)}\right)}{a^2-1} \right]$$

input

```
integrate(1/(1+a*cos(x)),x, algorithm="fricas")
```


output

```
[1/2*log(-(a^2 - 2)*cos(x)^2 - 2*sqrt(a^2 - 1)*(a + cos(x))*sin(x) - 2*a^2 - 2*a*cos(x) + 1)/(a^2*cos(x)^2 + 2*a*cos(x) + 1))/sqrt(a^2 - 1), -sqrt(-a^2 + 1)*arctan(sqrt(-a^2 + 1)*(a + cos(x))/((a^2 - 1)*sin(x)))/(a^2 - 1)
]
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 110 vs. $2(29) = 58$.

Time = 1.33 (sec) , antiderivative size = 110, normalized size of antiderivative = 2.97

$$\int \frac{1}{1 + a \cos(x)} dx = \begin{cases} \tan\left(\frac{x}{2}\right) & \text{for } a = 1 \\ -\frac{1}{\tan\left(\frac{x}{2}\right)} & \text{for } a = -1 \\ -\frac{\log\left(-\sqrt{\frac{a}{a-1} + \frac{1}{a-1}} + \tan\left(\frac{x}{2}\right)\right)}{a\sqrt{\frac{a}{a-1} + \frac{1}{a-1}} - \sqrt{\frac{a}{a-1} + \frac{1}{a-1}}} + \frac{\log\left(\sqrt{\frac{a}{a-1} + \frac{1}{a-1}} + \tan\left(\frac{x}{2}\right)\right)}{a\sqrt{\frac{a}{a-1} + \frac{1}{a-1}} - \sqrt{\frac{a}{a-1} + \frac{1}{a-1}}} & \text{otherwise} \end{cases}$$

input

```
integrate(1/(1+a*cos(x)),x)
```

output

```
Piecewise((tan(x/2), Eq(a, 1)), (-1/tan(x/2), Eq(a, -1)), (-log(-sqrt(a/(a - 1) + 1/(a - 1)) + tan(x/2))/(a*sqrt(a/(a - 1) + 1/(a - 1)) - sqrt(a/(a - 1) + 1/(a - 1))) + log(sqrt(a/(a - 1) + 1/(a - 1)) + tan(x/2))/(a*sqrt(a/(a - 1) + 1/(a - 1)) - sqrt(a/(a - 1) + 1/(a - 1))), True))
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{1}{1 + a \cos(x)} dx = \text{Exception raised: ValueError}$$

input

```
integrate(1/(1+a*cos(x)),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a^2-1.0>0)', see `assume?` for m
ore detail
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.43

$$\int \frac{1}{1+a\cos(x)} dx = -\frac{2\left(\pi\left\lfloor\frac{x}{2\pi} + \frac{1}{2}\right\rfloor \operatorname{sgn}(2a-2) + \arctan\left(\frac{a\tan(\frac{1}{2}x) - \tan(\frac{1}{2}x)}{\sqrt{-a^2+1}}\right)\right)}{\sqrt{-a^2+1}}$$

input

```
integrate(1/(1+a*cos(x)),x, algorithm="giac")
```

output

```
-2*(pi*floor(1/2*x/pi + 1/2)*sgn(2*a - 2) + arctan((a*tan(1/2*x) - tan(1/2
*x))/sqrt(-a^2 + 1)))/sqrt(-a^2 + 1)
```

Mupad [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.76

$$\int \frac{1}{1+a\cos(x)} dx = \frac{2 \operatorname{atanh}\left(\frac{\tan(\frac{x}{2})\sqrt{a-1}}{\sqrt{a+1}}\right)}{\sqrt{a-1}\sqrt{a+1}}$$

input

```
int(1/(a*cos(x) + 1),x)
```

output

```
(2*atanh((tan(x/2)*(a - 1)^(1/2))/(a + 1)^(1/2)))/((a - 1)^(1/2)*(a + 1)^(
1/2))
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.14

$$\int \frac{1}{1 + a \cos(x)} dx = \frac{2\sqrt{-a^2 + 1} \operatorname{atan}\left(\frac{\tan(\frac{x}{2})a - \tan(\frac{x}{2})}{\sqrt{-a^2 + 1}}\right)}{a^2 - 1}$$

input `int(1/(1+a*cos(x)),x)`

output `(2*sqrt(-a**2 + 1)*atan((tan(x/2)*a - tan(x/2))/sqrt(-a**2 + 1)))/(a**2 - 1)`

3.142 $\int \frac{1}{1+2\cos(x)} dx$

Optimal result	819
Mathematica [A] (verified)	819
Rubi [A] (verified)	820
Maple [A] (verified)	821
Fricas [A] (verification not implemented)	821
Sympy [A] (verification not implemented)	822
Maxima [A] (verification not implemented)	822
Giac [A] (verification not implemented)	822
Mupad [B] (verification not implemented)	823
Reduce [B] (verification not implemented)	823

Optimal result

Integrand size = 8, antiderivative size = 56

$$\int \frac{1}{1+2\cos(x)} dx = -\frac{\log(\sqrt{3}\cos(\frac{x}{2}) - \sin(\frac{x}{2}))}{\sqrt{3}} + \frac{\log(\sqrt{3}\cos(\frac{x}{2}) + \sin(\frac{x}{2}))}{\sqrt{3}}$$

output

```
-1/3*ln(-sin(1/2*x)+cos(1/2*x)*3^(1/2))*3^(1/2)+1/3*ln(sin(1/2*x)+cos(1/2*x)*3^(1/2))*3^(1/2)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.36

$$\int \frac{1}{1+2\cos(x)} dx = \frac{2\operatorname{arctanh}\left(\frac{\tan(\frac{x}{2})}{\sqrt{3}}\right)}{\sqrt{3}}$$

input

```
Integrate[(1 + 2*Cos[x])^(-1), x]
```

output

```
(2*ArcTanh[Tan[x/2]/Sqrt[3]])/Sqrt[3]
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.36, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3042, 3138, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{2 \cos(x) + 1} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{2 \sin\left(x + \frac{\pi}{2}\right) + 1} dx \\ & \quad \downarrow \text{3138} \\ & 2 \int \frac{1}{3 - \tan^2\left(\frac{x}{2}\right)} d \tan\left(\frac{x}{2}\right) \\ & \quad \downarrow \text{219} \\ & \frac{2 \operatorname{arctanh}\left(\frac{\tan\left(\frac{x}{2}\right)}{\sqrt{3}}\right)}{\sqrt{3}} \end{aligned}$$

input `Int[(1 + 2*Cos[x])^(-1), x]`

output `(2*ArcTanh[Tan[x/2]/Sqrt[3]])/Sqrt[3]`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138

```
Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{
e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b +
(a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x]
&& NeQ[a^2 - b^2, 0]
```

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.29

method	result	size
default	$\frac{2\sqrt{3} \operatorname{arctanh}\left(\frac{\tan\left(\frac{x}{2}\right)\sqrt{3}}{3}\right)}{3}$	16
risch	$\frac{\sqrt{3} \ln\left(e^{ix} + \frac{1}{2} + \frac{i\sqrt{3}}{2}\right)}{3} - \frac{\sqrt{3} \ln\left(e^{ix} + \frac{1}{2} - \frac{i\sqrt{3}}{2}\right)}{3}$	40

input

```
int(1/(1+2*cos(x)),x,method=_RETURNVERBOSE)
```

output

```
2/3*3^(1/2)*arctanh(1/3*tan(1/2*x)*3^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.89

$$\int \frac{1}{1 + 2 \cos(x)} dx$$

$$= \frac{1}{6} \sqrt{3} \log \left(-\frac{2 \cos(x)^2 - 2(\sqrt{3} \cos(x) + 2\sqrt{3}) \sin(x) - 4 \cos(x) - 7}{4 \cos(x)^2 + 4 \cos(x) + 1} \right)$$

input

```
integrate(1/(1+2*cos(x)),x, algorithm="fricas")
```

output

```
1/6*sqrt(3)*log(-(2*cos(x)^2 - 2*(sqrt(3)*cos(x) + 2*sqrt(3))*sin(x) - 4*cos(x) - 7)/(4*cos(x)^2 + 4*cos(x) + 1))
```

Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.64

$$\int \frac{1}{1+2\cos(x)} dx = -\frac{\sqrt{3}\log\left(\tan\left(\frac{x}{2}\right) - \sqrt{3}\right)}{3} + \frac{\sqrt{3}\log\left(\tan\left(\frac{x}{2}\right) + \sqrt{3}\right)}{3}$$

input `integrate(1/(1+2*cos(x)),x)`output `-sqrt(3)*log(tan(x/2) - sqrt(3))/3 + sqrt(3)*log(tan(x/2) + sqrt(3))/3`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.66

$$\int \frac{1}{1+2\cos(x)} dx = -\frac{1}{3}\sqrt{3}\log\left(-\frac{\sqrt{3} - \frac{\sin(x)}{\cos(x)+1}}{\sqrt{3} + \frac{\sin(x)}{\cos(x)+1}}\right)$$

input `integrate(1/(1+2*cos(x)),x, algorithm="maxima")`output `-1/3*sqrt(3)*log(-(sqrt(3) - sin(x)/(cos(x) + 1))/(sqrt(3) + sin(x)/(cos(x) + 1)))`**Giac [A] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.62

$$\int \frac{1}{1+2\cos(x)} dx = -\frac{1}{3}\sqrt{3}\log\left(\frac{|-2\sqrt{3} + 2\tan\left(\frac{1}{2}x\right)|}{|2\sqrt{3} + 2\tan\left(\frac{1}{2}x\right)|}\right)$$

input `integrate(1/(1+2*cos(x)),x, algorithm="giac")`output `-1/3*sqrt(3)*log(abs(-2*sqrt(3) + 2*tan(1/2*x))/abs(2*sqrt(3) + 2*tan(1/2*x)))`

Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.27

$$\int \frac{1}{1 + 2 \cos(x)} dx = \frac{2 \sqrt{3} \operatorname{atanh}\left(\frac{\sqrt{3} \tan\left(\frac{x}{2}\right)}{3}\right)}{3}$$

input `int(1/(2*cos(x) + 1),x)`output `(2*3^(1/2)*atanh((3^(1/2)*tan(x/2))/3))/3`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.45

$$\int \frac{1}{1 + 2 \cos(x)} dx = \frac{\sqrt{3} \left(-\log(-\sqrt{3} + \tan\left(\frac{x}{2}\right)) + \log(\sqrt{3} + \tan\left(\frac{x}{2}\right)) \right)}{3}$$

input `int(1/(1+2*cos(x)),x)`output `(sqrt(3)*(-log(-sqrt(3) + tan(x/2)) + log(sqrt(3) + tan(x/2))))/3`

$$3.143 \quad \int \frac{1}{1 + \frac{\cos(x)}{2}} dx$$

Optimal result	824
Mathematica [A] (verified)	824
Rubi [A] (verified)	825
Maple [A] (verified)	826
Fricas [A] (verification not implemented)	826
Sympy [A] (verification not implemented)	826
Maxima [A] (verification not implemented)	827
Giac [A] (verification not implemented)	827
Mupad [B] (verification not implemented)	828
Reduce [B] (verification not implemented)	828

Optimal result

Integrand size = 10, antiderivative size = 31

$$\int \frac{1}{1 + \frac{\cos(x)}{2}} dx = \frac{2x}{\sqrt{3}} - \frac{4 \arctan\left(\frac{\sin(x)}{2 + \sqrt{3} + \cos(x)}\right)}{\sqrt{3}}$$

output `2/3*x*3^(1/2)-4/3*arctan(sin(x)/(2+cos(x)+3^(1/2)))*3^(1/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.65

$$\int \frac{1}{1 + \frac{\cos(x)}{2}} dx = \frac{4 \arctan\left(\frac{\tan\left(\frac{x}{2}\right)}{\sqrt{3}}\right)}{\sqrt{3}}$$

input `Integrate[(1 + Cos[x]/2)^(-1),x]`

output `(4*ArcTan[Tan[x/2]/Sqrt[3]])/Sqrt[3]`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\frac{\cos(x)}{2} + 1} dx$$

↓ 3042

$$\int \frac{1}{\frac{1}{2} \sin\left(x + \frac{\pi}{2}\right) + 1} dx$$

↓ 3136

$$\frac{2x}{\sqrt{3}} - \frac{4 \arctan\left(\frac{\sin(x)}{\cos(x) + \sqrt{3} + 2}\right)}{\sqrt{3}}$$

input `Int[(1 + Cos[x]/2)^(-1),x]`

output `(2*x)/Sqrt[3] - (4*ArcTan[Sin[x]/(2 + Sqrt[3] + Cos[x])])/Sqrt[3]`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3136 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] :> With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + q + b*Sin[c + d*x]))], x]] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.52

method	result	size
default	$\frac{4\sqrt{3} \arctan\left(\frac{\tan\left(\frac{x}{2}\right)\sqrt{3}}{3}\right)}{3}$	16
risch	$\frac{2i\sqrt{3} \ln(e^{ix} + \sqrt{3} + 2)}{3} - \frac{2i\sqrt{3} \ln(e^{ix} - \sqrt{3} + 2)}{3}$	38

input `int(1/(1+1/2*cos(x)),x,method=_RETURNVERBOSE)`output `4/3*3^(1/2)*arctan(1/3*tan(1/2*x)*3^(1/2))`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

$$\int \frac{1}{1 + \frac{\cos(x)}{2}} dx = -\frac{2}{3} \sqrt{3} \arctan\left(\frac{2\sqrt{3} \cos(x) + \sqrt{3}}{3 \sin(x)}\right)$$

input `integrate(1/(1+1/2*cos(x)),x, algorithm="fricas")`output `-2/3*sqrt(3)*arctan(1/3*(2*sqrt(3)*cos(x) + sqrt(3))/sin(x))`**Sympy [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.03

$$\int \frac{1}{1 + \frac{\cos(x)}{2}} dx = \frac{4\sqrt{3} \left(\operatorname{atan}\left(\frac{\sqrt{3} \tan\left(\frac{x}{2}\right)}{3}\right) + \pi \left\lfloor \frac{\frac{x}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right)}{3}$$

input `integrate(1/(1+1/2*cos(x)),x)`

output `4*sqrt(3)*(atan(sqrt(3)*tan(x/2)/3) + pi*floor((x/2 - pi/2)/pi))/3`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.61

$$\int \frac{1}{1 + \frac{\cos(x)}{2}} dx = \frac{4}{3} \sqrt{3} \arctan \left(\frac{\sqrt{3} \sin(x)}{3(\cos(x) + 1)} \right)$$

input `integrate(1/(1+1/2*cos(x)),x, algorithm="maxima")`

output `4/3*sqrt(3)*arctan(1/3*sqrt(3)*sin(x)/(cos(x) + 1))`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.29

$$\int \frac{1}{1 + \frac{\cos(x)}{2}} dx = \frac{2}{3} \sqrt{3} \left(x + 2 \arctan \left(-\frac{\sqrt{3} \sin(x) - \sin(x)}{\sqrt{3} \cos(x) + \sqrt{3} - \cos(x) + 1} \right) \right)$$

input `integrate(1/(1+1/2*cos(x)),x, algorithm="giac")`

output `2/3*sqrt(3)*(x + 2*arctan(-(sqrt(3)*sin(x) - sin(x))/(sqrt(3)*cos(x) + sqrt(3) - cos(x) + 1)))`

Mupad [B] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.03

$$\int \frac{1}{1 + \frac{\cos(x)}{2}} dx = \frac{4\sqrt{3} \left(\frac{x}{2} - \operatorname{atan}\left(\tan\left(\frac{x}{2}\right)\right) \right)}{3} + \frac{4\sqrt{3} \operatorname{atan}\left(\frac{\sqrt{3} \tan\left(\frac{x}{2}\right)}{3}\right)}{3}$$

input `int(1/(cos(x)/2 + 1), x)`output `(4*3^(1/2)*(x/2 - atan(tan(x/2)))/3 + (4*3^(1/2)*atan((3^(1/2)*tan(x/2))/3))/3`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.45

$$\int \frac{1}{1 + \frac{\cos(x)}{2}} dx = \frac{4\sqrt{3} \operatorname{atan}\left(\frac{\tan\left(\frac{x}{2}\right)}{\sqrt{3}}\right)}{3}$$

input `int(1/(1+1/2*cos(x)), x)`output `(4*sqrt(3)*atan(tan(x/2)/sqrt(3)))/3`

3.144 $\int \frac{\sin^2(x)}{1+\sin^2(x)} dx$

Optimal result	829
Mathematica [A] (verified)	829
Rubi [A] (verified)	830
Maple [A] (verified)	831
Fricas [A] (verification not implemented)	832
Sympy [B] (verification not implemented)	832
Maxima [A] (verification not implemented)	833
Giac [A] (verification not implemented)	833
Mupad [B] (verification not implemented)	834
Reduce [B] (verification not implemented)	834

Optimal result

Integrand size = 13, antiderivative size = 36

$$\int \frac{\sin^2(x)}{1 + \sin^2(x)} dx = x - \frac{x}{\sqrt{2}} - \frac{\arctan\left(\frac{\cos(x)\sin(x)}{1+\sqrt{2}+\sin^2(x)}\right)}{\sqrt{2}}$$

output

```
x-1/2*x*2^(1/2)-1/2*arctan(cos(x)*sin(x)/(1+sin(x)^2+2^(1/2)))*2^(1/2)
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.50

$$\int \frac{\sin^2(x)}{1 + \sin^2(x)} dx = x - \frac{\arctan(\sqrt{2}\tan(x))}{\sqrt{2}}$$

input

```
Integrate[Sin[x]^2/(1 + Sin[x]^2),x]
```

output

```
x - ArcTan[Sqrt[2]*Tan[x]]/Sqrt[2]
```

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.50, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {3042, 3650, 3042, 3660, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin^2(x)}{\sin^2(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(x)^2}{\sin(x)^2 + 1} dx \\
 & \quad \downarrow \text{3650} \\
 & x - \int \frac{1}{\sin^2(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & x - \int \frac{1}{\sin(x)^2 + 1} dx \\
 & \quad \downarrow \text{3660} \\
 & x - \int \frac{1}{2 \tan^2(x) + 1} d \tan(x) \\
 & \quad \downarrow \text{216} \\
 & x - \frac{\arctan(\sqrt{2} \tan(x))}{\sqrt{2}}
 \end{aligned}$$

input

```
Int [Sin [x]^2/(1 + Sin [x]^2), x]
```

output

```
x - ArcTan [Sqrt [2]*Tan [x]]/Sqrt [2]
```

Definitions of rubi rules used

rule 216

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3650

```
Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]^2)/((a_) + (b_.)*sin[(e_.) + (f
.)*(x_)]^2), x_Symbol] := Simp[B*(x/b), x] + Simp[(A*b - a*B)/b Int[1/(a
+ b*Sin[e + f*x]^2), x], x] /; FreeQ[{a, b, e, f, A, B}, x]
```

rule 3660

```
Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(-1), x_Symbol] := With[{ff =
FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Subst[Int[1/(a + (a + b)*ff^2*x^
2), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, e, f}, x]
```

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.47

method	result	size
default	$\arctan(\tan(x)) - \frac{\sqrt{2} \arctan(\tan(x)\sqrt{2})}{2}$	17
risch	$x - \frac{i\sqrt{2} \ln(e^{2ix} - 2\sqrt{2} - 3)}{4} + \frac{i\sqrt{2} \ln(e^{2ix} + 2\sqrt{2} - 3)}{4}$	41

input

```
int(sin(x)^2/(1+sin(x)^2),x,method=_RETURNVERBOSE)
```

output

```
arctan(tan(x))-1/2*2^(1/2)*arctan(tan(x)*2^(1/2))
```


Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.92

$$\int \frac{\sin^2(x)}{1 + \sin^2(x)} dx = \frac{1}{4} \sqrt{2} \arctan \left(\frac{3 \sqrt{2} \cos(x)^2 - 2 \sqrt{2}}{4 \cos(x) \sin(x)} \right) + x$$

input `integrate(sin(x)^2/(1+sin(x)^2),x, algorithm="fricas")`

output `1/4*sqrt(2)*arctan(1/4*(3*sqrt(2)*cos(x)^2 - 2*sqrt(2))/(cos(x)*sin(x))) + x`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 248 vs. 2(36) = 72.

Time = 21.35 (sec) , antiderivative size = 248, normalized size of antiderivative = 6.89

$$\int \frac{\sin^2(x)}{1 + \sin^2(x)} dx = \frac{31988856\sqrt{2}x}{31988856\sqrt{2} + 45239074} + \frac{45239074x}{31988856\sqrt{2} + 45239074} - \frac{77227930\sqrt{3 - 2\sqrt{2}} \left(\operatorname{atan} \left(\frac{\tan(\frac{x}{2})}{\sqrt{3 - 2\sqrt{2}}} \right) + \pi \left\lfloor \frac{\frac{x}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right)}{31988856\sqrt{2} + 45239074} - \frac{54608393\sqrt{2}\sqrt{3 - 2\sqrt{2}} \left(\operatorname{atan} \left(\frac{\tan(\frac{x}{2})}{\sqrt{3 - 2\sqrt{2}}} \right) + \pi \left\lfloor \frac{\frac{x}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right)}{31988856\sqrt{2} + 45239074} - \frac{13250218\sqrt{2\sqrt{2} + 3} \left(\operatorname{atan} \left(\frac{\tan(\frac{x}{2})}{\sqrt{2\sqrt{2} + 3}} \right) + \pi \left\lfloor \frac{\frac{x}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right)}{31988856\sqrt{2} + 45239074} - \frac{9369319\sqrt{2}\sqrt{2\sqrt{2} + 3} \left(\operatorname{atan} \left(\frac{\tan(\frac{x}{2})}{\sqrt{2\sqrt{2} + 3}} \right) + \pi \left\lfloor \frac{\frac{x}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right)}{31988856\sqrt{2} + 45239074}$$

input `integrate(sin(x)**2/(1+sin(x)**2),x)`

output

```
31988856*sqrt(2)*x/(31988856*sqrt(2) + 45239074) + 45239074*x/(31988856*sqrt(2) + 45239074) - 77227930*sqrt(3 - 2*sqrt(2))*(atan(tan(x/2)/sqrt(3 - 2*sqrt(2)))) + pi*floor((x/2 - pi/2)/pi))/(31988856*sqrt(2) + 45239074) - 54608393*sqrt(2)*sqrt(3 - 2*sqrt(2))*(atan(tan(x/2)/sqrt(3 - 2*sqrt(2)))) + pi*floor((x/2 - pi/2)/pi))/(31988856*sqrt(2) + 45239074) - 13250218*sqrt(2*sqrt(2) + 3)*(atan(tan(x/2)/sqrt(2*sqrt(2) + 3)) + pi*floor((x/2 - pi/2)/pi))/(31988856*sqrt(2) + 45239074) - 9369319*sqrt(2)*sqrt(2*sqrt(2) + 3)*(atan(tan(x/2)/sqrt(2*sqrt(2) + 3)) + pi*floor((x/2 - pi/2)/pi))/(31988856*sqrt(2) + 45239074)
```

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.39

$$\int \frac{\sin^2(x)}{1 + \sin^2(x)} dx = -\frac{1}{2} \sqrt{2} \arctan(\sqrt{2} \tan(x)) + x$$

input

```
integrate(sin(x)^2/(1+sin(x)^2),x, algorithm="maxima")
```

output

```
-1/2*sqrt(2)*arctan(sqrt(2)*tan(x)) + x
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.33

$$\int \frac{\sin^2(x)}{1 + \sin^2(x)} dx = -\frac{1}{2} \sqrt{2} \left(x + \arctan \left(-\frac{\sqrt{2} \sin(2x) - 2 \sin(2x)}{\sqrt{2} \cos(2x) + \sqrt{2} - 2 \cos(2x) + 2} \right) \right) + x$$

input

```
integrate(sin(x)^2/(1+sin(x)^2),x, algorithm="giac")
```

output

```
-1/2*sqrt(2)*(x + arctan(-(sqrt(2)*sin(2*x) - 2*sin(2*x))/(sqrt(2)*cos(2*x) + sqrt(2) - 2*cos(2*x) + 2))) + x
```

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.72

$$\int \frac{\sin^2(x)}{1 + \sin^2(x)} dx = x - \frac{\sqrt{2}(x - \operatorname{atan}(\tan(x)))}{2} - \frac{\sqrt{2} \operatorname{atan}(\sqrt{2} \tan(x))}{2}$$

input `int(sin(x)^2/(sin(x)^2 + 1),x)`output `x - (2^(1/2)*(x - atan(tan(x))))/2 - (2^(1/2)*atan(2^(1/2)*tan(x)))/2`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.47

$$\int \frac{\sin^2(x)}{1 + \sin^2(x)} dx = -\frac{\sqrt{2} \operatorname{atan}\left(\frac{\tan\left(\frac{x}{2}\right)}{\sqrt{2}+1}\right)}{2} + \frac{\sqrt{2} \log(-\sqrt{2}i + \tan\left(\frac{x}{2}\right) + i) i}{4} \\ - \frac{\sqrt{2} \log(\sqrt{2}i + \tan\left(\frac{x}{2}\right) - i) i}{4} + x$$

input `int(sin(x)^2/(1+sin(x)^2),x)`output `(- 2*sqrt(2)*atan(tan(x/2)/(sqrt(2) + 1)) + sqrt(2)*log(- sqrt(2)*i + ta
n(x/2) + i)*i - sqrt(2)*log(sqrt(2)*i + tan(x/2) - i)*i + 4*x)/4`

$$3.145 \quad \int \frac{1}{b^2 \cos^2(x) + a^2 \sin^2(x)} dx$$

Optimal result	835
Mathematica [A] (verified)	835
Rubi [A] (verified)	836
Maple [A] (verified)	837
Fricas [B] (verification not implemented)	837
Sympy [B] (verification not implemented)	838
Maxima [A] (verification not implemented)	839
Giac [A] (verification not implemented)	839
Mupad [B] (verification not implemented)	839
Reduce [F]	840

Optimal result

Integrand size = 19, antiderivative size = 15

$$\int \frac{1}{b^2 \cos^2(x) + a^2 \sin^2(x)} dx = \frac{\arctan\left(\frac{a \tan(x)}{b}\right)}{ab}$$

output `arctan(a*tan(x)/b)/a/b`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{1}{b^2 \cos^2(x) + a^2 \sin^2(x)} dx = \frac{\arctan\left(\frac{a \tan(x)}{b}\right)}{ab}$$

input `Integrate[(b^2*Cos[x]^2 + a^2*Sin[x]^2)^(-1),x]`

output `ArcTan[(a*Tan[x])/b]/(a*b)`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {3042, 4889, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{a^2 \sin^2(x) + b^2 \cos^2(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{a^2 \sin(x)^2 + b^2 \cos(x)^2} dx \\ & \quad \downarrow \text{4889} \\ & \int \frac{1}{a^2 \tan^2(x) + b^2} d \tan(x) \\ & \quad \downarrow \text{218} \\ & \frac{\arctan\left(\frac{a \tan(x)}{b}\right)}{ab} \end{aligned}$$

input `Int[(b^2*Cos[x]^2 + a^2*Sin[x]^2)^(-1), x]`

output `ArcTan[(a*Tan[x])/b]/(a*b)`

Defintions of rubi rules used

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4889

```
Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, With[{d = FreeFactors
[Tan[v], x]}, Simp[d/Coefficient[v, x, 1] Subst[Int[SubstFor[1/(1 + d^2*x
^2), Tan[v]/d, u, x], x], x, Tan[v]/d], x]] /; !FalseQ[v] && FunctionOfQ[N
onfreeFactors[Tan[v], x], u, x]] /; InverseFunctionFreeQ[u, x] && !MatchQ[
u, (v_.)*((c_.)*tan[w_]^(n_.)*tan[z_]^(n_.))^ (p_.) /; FreeQ[{c, p}, x] && I
ntegerQ[n] && LinearQ[w, x] && EqQ[z, 2*w]]
```

Maple [A] (verified)

Time = 0.67 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.07

method	result	size
default	$\frac{\arctan\left(\frac{a \tan(x)}{b}\right)}{ab}$	16
paralelrisch	$\frac{i\left(\ln\left(2ia \tan\left(\frac{x}{2}\right) + \sec\left(\frac{x}{2}\right)^2 b - 2b\right) - \ln\left(-2ia \tan\left(\frac{x}{2}\right) + \sec\left(\frac{x}{2}\right)^2 b - 2b\right)\right)}{2ab}$	55
risch	$-\frac{i \ln\left(e^{2ix} - \frac{a-b}{a+b}\right)}{2ab} + \frac{i \ln\left(e^{2ix} - \frac{a+b}{a-b}\right)}{2ab}$	58

input

```
int(1/(b^2*cos(x)^2+a^2*sin(x)^2),x,method=_RETURNVERBOSE)
```

output

```
arctan(a*tan(x)/b)/a/b
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 43 vs. 2(15) = 30.

Time = 0.09 (sec) , antiderivative size = 43, normalized size of antiderivative = 2.87

$$\int \frac{1}{b^2 \cos^2(x) + a^2 \sin^2(x)} dx = -\frac{\arctan\left(\frac{(a^2+b^2) \cos(x)^2 - a^2}{2ab \cos(x) \sin(x)}\right)}{2ab}$$

input

```
integrate(1/(b^2*cos(x)^2+a^2*sin(x)^2),x, algorithm="fricas")
```

output

```
-1/2*arctan(1/2*((a^2 + b^2)*cos(x)^2 - a^2)/(a*b*cos(x)*sin(x)))/(a*b)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 71839 vs. $2(10) = 20$.

Time = 15.30 (sec) , antiderivative size = 71839, normalized size of antiderivative = 4789.27

$$\int \frac{1}{b^2 \cos^2(x) + a^2 \sin^2(x)} dx = \text{Too large to display}$$

input `integrate(1/(b**2*cos(x)**2+a**2*sin(x)**2),x)`

output

```
Piecewise((zoo*tan(x/2)/(tan(x/2)**2 - 1), Eq(a, 0) & Eq(b, 0)), ((tan(x/2)/2 - 1/(2*tan(x/2)))/a**2, Eq(b, 0)), (-2*tan(x/2)/(b**2*(tan(x/2)**2 - 1)), Eq(a, 0)), (x/(b**2*sin(x)**2 + b**2*cos(x)**2), Eq(a, b) | Eq(a, -b)), (8192*a**15*sqrt(-2*a**2/b**2 - 2*a*sqrt(a**2 - b**2)/b**2 + 1)*log(-sqrt(-2*a**2/b**2 + 2*a*sqrt(a**2 - b**2)/b**2 + 1) + tan(x/2))/(8192*a**15*b**2*sqrt(-2*a**2/b**2 - 2*a*sqrt(a**2 - b**2)/b**2 + 1)*sqrt(-2*a**2/b**2 + 2*a*sqrt(a**2 - b**2)/b**2 + 1) - 8192*a**14*b**2*sqrt(a**2 - b**2)*sqrt(-2*a**2/b**2 - 2*a*sqrt(a**2 - b**2)/b**2 + 1)*sqrt(-2*a**2/b**2 + 2*a*sqrt(a**2 - b**2)/b**2 + 1) - 30720*a**13*b**4*sqrt(-2*a**2/b**2 - 2*a*sqrt(a**2 - b**2)/b**2 + 1)*sqrt(-2*a**2/b**2 + 2*a*sqrt(a**2 - b**2)/b**2 + 1) + 26624*a**12*b**4*sqrt(a**2 - b**2)*sqrt(-2*a**2/b**2 - 2*a*sqrt(a**2 - b**2)/b**2 + 1)*sqrt(-2*a**2/b**2 + 2*a*sqrt(a**2 - b**2)/b**2 + 1) + 45568*a**11*b**6*sqrt(-2*a**2/b**2 - 2*a*sqrt(a**2 - b**2)/b**2 + 1)*sqrt(-2*a**2/b**2 + 2*a*sqrt(a**2 - b**2)/b**2 + 1) - 33280*a**10*b**6*sqrt(a**2 - b**2)*sqrt(-2*a**2/b**2 - 2*a*sqrt(a**2 - b**2)/b**2 + 1)*sqrt(-2*a**2/b**2 + 2*a*sqrt(a**2 - b**2)/b**2 + 1) - 33792*a**9*b**8*sqrt(-2*a**2/b**2 - 2*a*sqrt(a**2 - b**2)/b**2 + 1)*sqrt(-2*a**2/b**2 + 2*a*sqrt(a**2 - b**2)/b**2 + 1) + 19968*a**8*b**8*sqrt(a**2 - b**2)*sqrt(-2*a**2/b**2 - 2*a*sqrt(a**2 - b**2)/b**2 + 1)*sqrt(-2*a**2/b**2 + 2*a*sqrt(a**2 - b**2)/b**2 + 1) + 12992*a**7*b**10*sqrt(-2*a**2/b**2 - 2*a*sqrt(a**2 - b**2)/b**2 + 1...
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{1}{b^2 \cos^2(x) + a^2 \sin^2(x)} dx = \frac{\arctan\left(\frac{a \tan(x)}{b}\right)}{ab}$$

input `integrate(1/(b^2*cos(x)^2+a^2*sin(x)^2),x, algorithm="maxima")`output `arctan(a*tan(x)/b)/(a*b)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.73

$$\int \frac{1}{b^2 \cos^2(x) + a^2 \sin^2(x)} dx = \frac{\pi \lfloor \frac{x}{\pi} + \frac{1}{2} \rfloor + \arctan\left(\frac{a \tan(x)}{b}\right)}{ab}$$

input `integrate(1/(b^2*cos(x)^2+a^2*sin(x)^2),x, algorithm="giac")`output `(pi*floor(x/pi + 1/2) + arctan(a*tan(x)/b))/(a*b)`**Mupad [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{1}{b^2 \cos^2(x) + a^2 \sin^2(x)} dx = \frac{\operatorname{atan}\left(\frac{a \tan(x)}{b}\right)}{ab}$$

input `int(1/(b^2*cos(x)^2 + a^2*sin(x)^2),x)`output `atan((a*tan(x))/b)/(a*b)`

Reduce **[F]**

$$\int \frac{1}{b^2 \cos^2(x) + a^2 \sin^2(x)} dx = \int \frac{1}{\cos(x)^2 b^2 + \sin(x)^2 a^2} dx$$

input `int(1/(b^2*cos(x)^2+a^2*sin(x)^2),x)`

output `int(1/(cos(x)**2*b**2 + sin(x)**2*a**2),x)`

3.146 $\int \frac{1}{(b \cos(x) + a \sin(x))^2} dx$

Optimal result	841
Mathematica [A] (verified)	841
Rubi [A] (verified)	842
Maple [A] (verified)	843
Fricas [B] (verification not implemented)	843
Sympy [B] (verification not implemented)	844
Maxima [A] (verification not implemented)	844
Giac [A] (verification not implemented)	845
Mupad [B] (verification not implemented)	845
Reduce [B] (verification not implemented)	845

Optimal result

Integrand size = 11, antiderivative size = 17

$$\int \frac{1}{(b \cos(x) + a \sin(x))^2} dx = \frac{\sin(x)}{b(b \cos(x) + a \sin(x))}$$

output `sin(x)/b/(b*cos(x)+a*sin(x))`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{1}{(b \cos(x) + a \sin(x))^2} dx = \frac{\sin(x)}{b(b \cos(x) + a \sin(x))}$$

input `Integrate[(b*Cos[x] + a*Sin[x])^(-2),x]`

output `Sin[x]/(b*(b*Cos[x] + a*Sin[x]))`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3042, 3554}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a \sin(x) + b \cos(x))^2} dx$$

↓ 3042

$$\int \frac{1}{(a \sin(x) + b \cos(x))^2} dx$$

↓ 3554

$$\frac{\sin(x)}{b(a \sin(x) + b \cos(x))}$$

input `Int[(b*Cos[x] + a*Sin[x])^(-2),x]`

output `Sin[x]/(b*(b*Cos[x] + a*Sin[x]))`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3554 `Int[(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-2), x_Symbol] := Simp[Sin[c + d*x]/(a*d*(a*Cos[c + d*x] + b*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]`

Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$-\frac{1}{a(a \tan(x)+b)}$	14
parallelrisc	$\frac{\sin(x)}{b(b \cos(x)+a \sin(x))}$	18
norman	$\frac{-\frac{1}{a} + \frac{\tan(\frac{x}{2})^2}{a}}{-b \tan(\frac{x}{2})^2 + 2a \tan(\frac{x}{2}) + b}$	38
risc	$-\frac{2i}{(a e^{2ix} + i b e^{2ix} - a + i b)(i b + a)}$	38

input `int(1/(b*cos(x)+a*sin(x))^2,x,method=_RETURNVERBOSE)`

output `-1/a/(a*tan(x)+b)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. 2(17) = 34.

Time = 0.08 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.29

$$\int \frac{1}{(b \cos(x) + a \sin(x))^2} dx = -\frac{a \cos(x) - b \sin(x)}{(a^2 b + b^3) \cos(x) + (a^3 + a b^2) \sin(x)}$$

input `integrate(1/(b*cos(x)+a*sin(x))^2,x, algorithm="fricas")`

output `-(a*cos(x) - b*sin(x))/((a^2*b + b^3)*cos(x) + (a^3 + a*b^2)*sin(x))`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 602 vs. $2(14) = 28$.

Time = 103.86 (sec) , antiderivative size = 602, normalized size of antiderivative = 35.41

$$\int \frac{1}{(b \cos(x) + a \sin(x))^2} dx = \text{Too large to display}$$

input `integrate(1/(b*cos(x)+a*sin(x))**2,x)`

output `Piecewise((zoo*tan(x/2)/(tan(x/2)**2 - 1), Eq(a, 0) & Eq(b, 0)), (x*tan(x/2)**4/(2*b**2*sin(x)**2*tan(x/2)**4 - 4*b**2*sin(x)**2*tan(x/2)**2 + 2*b**2*sin(x)**2 + 8*b**2*sin(x)*cos(x)*tan(x/2)**3 - 8*b**2*sin(x)*cos(x)*tan(x/2) + 8*b**2*cos(x)**2*tan(x/2)**2) + 2*x*tan(x/2)**2/(2*b**2*sin(x)**2*tan(x/2)**4 - 4*b**2*sin(x)**2*tan(x/2)**2 + 2*b**2*sin(x)**2 + 8*b**2*sin(x)*cos(x)*tan(x/2)**3 - 8*b**2*sin(x)*cos(x)*tan(x/2) + 8*b**2*cos(x)**2*tan(x/2)**2) + x/(2*b**2*sin(x)**2*tan(x/2)**4 - 4*b**2*sin(x)**2*tan(x/2)**2 + 2*b**2*sin(x)**2 + 8*b**2*sin(x)*cos(x)*tan(x/2)**3 - 8*b**2*sin(x)*cos(x)*tan(x/2) + 8*b**2*cos(x)**2*tan(x/2)**2) - 2*tan(x/2)/(2*b**2*sin(x)**2*tan(x/2)**4 - 4*b**2*sin(x)**2*tan(x/2)**2 + 2*b**2*sin(x)**2 + 8*b**2*sin(x)*cos(x)*tan(x/2)**3 - 8*b**2*sin(x)*cos(x)*tan(x/2) + 8*b**2*cos(x)**2*tan(x/2)**2) - 2*tan(x/2)/(2*b**2*sin(x)**2*tan(x/2)**4 - 4*b**2*sin(x)**2*tan(x/2)**2 + 2*b**2*sin(x)**2 + 8*b**2*sin(x)*cos(x)*tan(x/2)**3 - 8*b**2*sin(x)*cos(x)*tan(x/2) + 8*b**2*cos(x)**2*tan(x/2)**2), Eq(a, b*(tan(x/2) - 1)*(tan(x/2) + 1)/(2*tan(x/2)))), ((tan(x/2)/2 - 1/(2*tan(x/2)))/a**2, Eq(b, 0)), (2*tan(x/2)/(2*a*b*tan(x/2) - b**2*tan(x/2)**2 + b**2), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \frac{1}{(b \cos(x) + a \sin(x))^2} dx = -\frac{1}{a^2 \tan(x) + ab}$$

input `integrate(1/(b*cos(x)+a*sin(x))^2,x, algorithm="maxima")`

output $-1/(a^2 \tan(x) + a b)$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \frac{1}{(b \cos(x) + a \sin(x))^2} dx = -\frac{1}{(a \tan(x) + b)a}$$

input `integrate(1/(b*cos(x)+a*sin(x))^2,x, algorithm="giac")`

output $-1/((a \tan(x) + b) a)$

Mupad [B] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.71

$$\int \frac{1}{(b \cos(x) + a \sin(x))^2} dx = \frac{2 \tan\left(\frac{x}{2}\right)}{b \left(-b \tan\left(\frac{x}{2}\right)^2 + 2 a \tan\left(\frac{x}{2}\right) + b\right)}$$

input `int(1/(b*cos(x) + a*sin(x))^2,x)`

output $(2 \tan(x/2)) / (b * (b + 2 * a * \tan(x/2) - b * \tan(x/2)^2))$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.06

$$\int \frac{1}{(b \cos(x) + a \sin(x))^2} dx = -\frac{\cos(x)}{a (\cos(x) b + \sin(x) a)}$$

input `int(1/(b*cos(x)+a*sin(x))^2,x)`

output $(-\cos(x))/(a(\cos(x)*b + \sin(x)*a))$

3.147 $\int \frac{\sin(x)}{1+\cos(x)+\sin(x)} dx$

Optimal result	847
Mathematica [A] (verified)	847
Rubi [A] (verified)	848
Maple [C] (verified)	849
Fricas [A] (verification not implemented)	850
Sympy [A] (verification not implemented)	850
Maxima [A] (verification not implemented)	851
Giac [A] (verification not implemented)	851
Mupad [B] (verification not implemented)	851
Reduce [B] (verification not implemented)	852

Optimal result

Integrand size = 11, antiderivative size = 30

$$\int \frac{\sin(x)}{1 + \cos(x) + \sin(x)} dx = \frac{x}{2} - \frac{1}{2} \log(1 + \cos(x) + \sin(x)) - \frac{1}{2} \log\left(1 + \tan\left(\frac{x}{2}\right)\right)$$

output `1/2*x-1/2*ln(1+cos(x)+sin(x))-1/2*ln(1+tan(1/2*x))`

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \frac{\sin(x)}{1 + \cos(x) + \sin(x)} dx = \frac{x}{2} - \log\left(\cos\left(\frac{x}{2}\right) + \sin\left(\frac{x}{2}\right)\right)$$

input `Integrate[Sin[x]/(1 + Cos[x] + Sin[x]),x]`

output `x/2 - Log[Cos[x/2] + Sin[x/2]]`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3616, 3042, 3603, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin(x)}{\sin(x) + \cos(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(x)}{\sin(x) + \cos(x) + 1} dx \\
 & \quad \downarrow \text{3616} \\
 & -\frac{1}{2} \int \frac{1}{\cos(x) + \sin(x) + 1} dx + \frac{x}{2} - \frac{1}{2} \log(\sin(x) + \cos(x) + 1) \\
 & \quad \downarrow \text{3042} \\
 & -\frac{1}{2} \int \frac{1}{\cos(x) + \sin(x) + 1} dx + \frac{x}{2} - \frac{1}{2} \log(\sin(x) + \cos(x) + 1) \\
 & \quad \downarrow \text{3603} \\
 & -\int \frac{1}{2 \tan\left(\frac{x}{2}\right) + 2} d \tan\left(\frac{x}{2}\right) + \frac{x}{2} - \frac{1}{2} \log(\sin(x) + \cos(x) + 1) \\
 & \quad \downarrow \text{16} \\
 & \frac{x}{2} - \frac{1}{2} \log\left(\tan\left(\frac{x}{2}\right) + 1\right) - \frac{1}{2} \log(\sin(x) + \cos(x) + 1)
 \end{aligned}$$

input `Int[Sin[x]/(1 + Cos[x] + Sin[x]),x]`

output `x/2 - Log[1 + Cos[x] + Sin[x]]/2 - Log[1 + Tan[x/2]]/2`

Definitions of rubi rules used

rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3603 $\text{Int}[(\cos[(d_)+(e_)*(x_)]*(b_)+(a_)+(c_)*\sin[(d_)+(e_)*(x_)])^(-1), x_Symbol] \rightarrow \text{Module}[\{f = \text{FreeFactors}[\text{Tan}[(d + e*x)/2], x]\}, \text{Simp}[2*(f/e) \text{ Subst}[\text{Int}[1/(a + b + 2*c*f*x + (a - b)*f^2*x^2), x], x, \text{Tan}[(d + e*x)/2]/f], x]] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{NeQ}[a^2 - b^2 - c^2, 0]$

rule 3616 $\text{Int}[(A_)+(C_)*\sin[(d_)+(e_)*(x_)]/((a_)+\cos[(d_)+(e_)*(x_)]*(b_)+(c_)*\sin[(d_)+(e_)*(x_)]), x_Symbol] \rightarrow \text{Simp}[c*C*((d + e*x)/(e*(b^2 + c^2))), x] + (-\text{Simp}[b*C*(\text{Log}[a + b*\text{Cos}[d + e*x] + c*\text{Sin}[d + e*x]]/(e*(b^2 + c^2))), x] + \text{Simp}[(A*(b^2 + c^2) - a*c*C)/(b^2 + c^2) \text{ Int}[1/(a + b*\text{Cos}[d + e*x] + c*\text{Sin}[d + e*x]), x], x]) /; \text{FreeQ}[\{a, b, c, d, e, A, C\}, x] \&\& \text{NeQ}[b^2 + c^2, 0] \&\& \text{NeQ}[A*(b^2 + c^2) - a*c*C, 0]$

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.67

method	result	size
risch	$\frac{x}{2} + \frac{ix}{2} - \ln(e^{ix} + i)$	20
parallelrisch	$\frac{x}{2} - \ln(1 + \tan(\frac{x}{2})) + \ln\left(\sqrt{\sec(\frac{x}{2})^2}\right)$	23
default	$-\ln(1 + \tan(\frac{x}{2})) + \frac{\ln(1 + \tan(\frac{x}{2})^2)}{2} + \arctan(\tan(\frac{x}{2}))$	27
norman	$\frac{\frac{x}{2} + \frac{x \tan(\frac{x}{2})^2}{2}}{1 + \tan(\frac{x}{2})^2} - \ln(1 + \tan(\frac{x}{2})) + \frac{\ln(1 + \tan(\frac{x}{2})^2)}{2}$	46

input `int(sin(x)/(1+cos(x)+sin(x)),x,method=_RETURNVERBOSE)`

output `1/2*x+1/2*I*x-ln(exp(I*x)+I)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.37

$$\int \frac{\sin(x)}{1 + \cos(x) + \sin(x)} dx = \frac{1}{2}x - \frac{1}{2} \log(\sin(x) + 1)$$

input `integrate(sin(x)/(1+cos(x)+sin(x)),x, algorithm="fricas")`

output `1/2*x - 1/2*log(sin(x) + 1)`

Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \frac{\sin(x)}{1 + \cos(x) + \sin(x)} dx = \frac{x}{2} - \log\left(\tan\left(\frac{x}{2}\right) + 1\right) + \frac{\log\left(\tan^2\left(\frac{x}{2}\right) + 1\right)}{2}$$

input `integrate(sin(x)/(1+cos(x)+sin(x)),x)`

output `x/2 - log(tan(x/2) + 1) + log(tan(x/2)**2 + 1)/2`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.37

$$\int \frac{\sin(x)}{1 + \cos(x) + \sin(x)} dx = \arctan\left(\frac{\sin(x)}{\cos(x) + 1}\right) - \log\left(\frac{\sin(x)}{\cos(x) + 1} + 1\right) + \frac{1}{2} \log\left(\frac{\sin(x)^2}{(\cos(x) + 1)^2} + 1\right)$$

input `integrate(sin(x)/(1+cos(x)+sin(x)),x, algorithm="maxima")`output `arctan(sin(x)/(cos(x) + 1)) - log(sin(x)/(cos(x) + 1) + 1) + 1/2*log(sin(x)^2/(cos(x) + 1)^2 + 1)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.83

$$\int \frac{\sin(x)}{1 + \cos(x) + \sin(x)} dx = \frac{1}{2}x + \frac{1}{2} \log\left(\tan\left(\frac{1}{2}x\right)^2 + 1\right) - \log\left(\left|\tan\left(\frac{1}{2}x\right) + 1\right|\right)$$

input `integrate(sin(x)/(1+cos(x)+sin(x)),x, algorithm="giac")`output `1/2*x + 1/2*log(tan(1/2*x)^2 + 1) - log(abs(tan(1/2*x) + 1))`**Mupad [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.13

$$\int \frac{\sin(x)}{1 + \cos(x) + \sin(x)} dx = -\ln\left(\tan\left(\frac{x}{2}\right) + 1\right) + \ln\left(\tan\left(\frac{x}{2}\right) - i\right) \left(\frac{1}{2} - \frac{1}{2}i\right) + \ln\left(\tan\left(\frac{x}{2}\right) + 1i\right) \left(\frac{1}{2} + \frac{1}{2}i\right)$$

input `int(sin(x)/(cos(x) + sin(x) + 1),x)`

output `log(tan(x/2) - 1i)*(1/2 - 1i/2) - log(tan(x/2) + 1) + log(tan(x/2) + 1i)*(1/2 + 1i/2)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \frac{\sin(x)}{1 + \cos(x) + \sin(x)} dx = -\frac{\log(\cos(x) + \sin(x) + 1)}{2} - \frac{\log(\tan(\frac{x}{2}) + 1)}{2} + \frac{x}{2}$$

input `int(sin(x)/(1+cos(x)+sin(x)),x)`

output `(- log(cos(x) + sin(x) + 1) - log(tan(x/2) + 1) + x)/2`

3.148 $\int \sqrt{3 - x^2} dx$

Optimal result	853
Mathematica [A] (verified)	853
Rubi [A] (verified)	854
Maple [A] (verified)	855
Fricas [A] (verification not implemented)	855
Sympy [A] (verification not implemented)	856
Maxima [A] (verification not implemented)	856
Giac [A] (verification not implemented)	856
Mupad [B] (verification not implemented)	857
Reduce [B] (verification not implemented)	857

Optimal result

Integrand size = 11, antiderivative size = 29

$$\int \sqrt{3 - x^2} dx = \frac{1}{2}x\sqrt{3 - x^2} + \frac{3}{2} \arcsin\left(\frac{x}{\sqrt{3}}\right)$$

output

```
3/2*arcsin(1/3*x*3^(1/2))+1/2*x*(-x^2+3)^(1/2)
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.41

$$\int \sqrt{3 - x^2} dx = \frac{1}{2}x\sqrt{3 - x^2} + 3 \arctan\left(\frac{-\sqrt{3} + x}{\sqrt{3 - x^2}}\right)$$

input

```
Integrate[Sqrt[3 - x^2], x]
```

output

```
(x*Sqrt[3 - x^2])/2 + 3*ArcTan[(-Sqrt[3] + x)/Sqrt[3 - x^2]]
```

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {211, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{3-x^2} dx$$

$$\downarrow \text{211}$$

$$\frac{3}{2} \int \frac{1}{\sqrt{3-x^2}} dx + \frac{1}{2} \sqrt{3-x^2} x$$

$$\downarrow \text{223}$$

$$\frac{3}{2} \arcsin\left(\frac{x}{\sqrt{3}}\right) + \frac{1}{2} \sqrt{3-x^2} x$$

input `Int[Sqrt[3 - x^2],x]`

output `(x*Sqrt[3 - x^2])/2 + (3*ArcSin[x/Sqrt[3]])/2`

Defintions of rubi rules used

rule 211 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[x*((a + b*x^2)^p/(2*p + 1)), x] + Simp[2*a*(p/(2*p + 1)) Int[(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.79

method	result	size
default	$\frac{3 \arcsin\left(\frac{x\sqrt{3}}{3}\right)}{2} + \frac{x\sqrt{-x^2+3}}{2}$	23
risch	$-\frac{x(x^2-3)}{2\sqrt{-x^2+3}} + \frac{3 \arcsin\left(\frac{x\sqrt{3}}{3}\right)}{2}$	28
pseudoelliptic	$\frac{x\sqrt{-x^2+3}}{2} - \frac{3 \arctan\left(\frac{\sqrt{-x^2+3}}{x}\right)}{2}$	30
meijerg	$\frac{3i \left(-\frac{2i\sqrt{\pi} x\sqrt{3} \sqrt{-\frac{x^2}{3}+1}}{3} - 2i\sqrt{\pi} \arcsin\left(\frac{x\sqrt{3}}{3}\right) \right)}{4\sqrt{\pi}}$	40
trager	$\frac{x\sqrt{-x^2+3}}{2} + \frac{3 \operatorname{RootOf}\left(-Z^2+1\right) \ln\left(\operatorname{RootOf}\left(-Z^2+1\right)\sqrt{-x^2+3}+x\right)}{2}$	41

input `int((-x^2+3)^(1/2),x,method=_RETURNVERBOSE)`

output `3/2*arcsin(1/3*x*3^(1/2))+1/2*x*(-x^2+3)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.17

$$\int \sqrt{3-x^2} dx = \frac{1}{2} \sqrt{-x^2+3x} - \frac{3}{2} \arctan\left(\frac{\sqrt{-x^2+3x}}{x^2-3}\right)$$

input `integrate((-x^2+3)^(1/2),x, algorithm="fricas")`

output `1/2*sqrt(-x^2 + 3)*x - 3/2*arctan(sqrt(-x^2 + 3)*x/(x^2 - 3))`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

$$\int \sqrt{3-x^2} dx = \frac{x\sqrt{3-x^2}}{2} + \frac{3 \operatorname{asin}\left(\frac{\sqrt{3}x}{3}\right)}{2}$$

input `integrate((-x**2+3)**(1/2),x)`output `x*sqrt(3 - x**2)/2 + 3*asin(sqrt(3)*x/3)/2`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.76

$$\int \sqrt{3-x^2} dx = \frac{1}{2} \sqrt{-x^2+3x} + \frac{3}{2} \operatorname{arcsin}\left(\frac{1}{3} \sqrt{3}x\right)$$

input `integrate((-x^2+3)^(1/2),x, algorithm="maxima")`output `1/2*sqrt(-x^2 + 3)*x + 3/2*arcsin(1/3*sqrt(3)*x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.76

$$\int \sqrt{3-x^2} dx = \frac{1}{2} \sqrt{-x^2+3x} + \frac{3}{2} \operatorname{arcsin}\left(\frac{1}{3} \sqrt{3}x\right)$$

input `integrate((-x^2+3)^(1/2),x, algorithm="giac")`output `1/2*sqrt(-x^2 + 3)*x + 3/2*arcsin(1/3*sqrt(3)*x)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.76

$$\int \sqrt{3-x^2} dx = \frac{3 \operatorname{asin}\left(\frac{\sqrt{3}x}{3}\right)}{2} + \frac{x\sqrt{3-x^2}}{2}$$

input `int((3 - x^2)^(1/2),x)`output `(3*asin((3^(1/2)*x)/3))/2 + (x*(3 - x^2)^(1/2))/2`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

$$\int \sqrt{3-x^2} dx = \frac{3 \operatorname{asin}\left(\frac{x}{\sqrt{3}}\right)}{2} + \frac{\sqrt{-x^2+3}x}{2}$$

input `int((-x^2+3)^(1/2),x)`output `(3*asin(x/sqrt(3)) + sqrt(-x**2 + 3)*x)/2`

3.149 $\int \frac{x}{\sqrt{3-x^2}} dx$

Optimal result	858
Mathematica [A] (verified)	858
Rubi [A] (verified)	859
Maple [A] (verified)	860
Fricas [A] (verification not implemented)	860
Sympy [A] (verification not implemented)	861
Maxima [A] (verification not implemented)	861
Giac [A] (verification not implemented)	861
Mupad [B] (verification not implemented)	862
Reduce [B] (verification not implemented)	862

Optimal result

Integrand size = 13, antiderivative size = 13

$$\int \frac{x}{\sqrt{3-x^2}} dx = -\sqrt{3-x^2}$$

output `-(-x^2+3)^(1/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{x}{\sqrt{3-x^2}} dx = -\sqrt{3-x^2}$$

input `Integrate[x/Sqrt[3 - x^2],x]`

output `-Sqrt[3 - x^2]`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\sqrt{3-x^2}} dx$$

↓ 241

$$-\sqrt{3-x^2}$$

input `Int[x/Sqrt[3 - x^2], x]`

output `-Sqrt[3 - x^2]`

Defintions of rubi rules used

rule 241 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
gospers	$-\sqrt{-x^2 + 3}$	12
derivativedivides	$-\sqrt{-x^2 + 3}$	12
default	$-\sqrt{-x^2 + 3}$	12
trager	$-\sqrt{-x^2 + 3}$	12
pseudoelliptic	$-\sqrt{-x^2 + 3}$	12
risch	$\frac{x^2-3}{\sqrt{-x^2+3}}$	16
orering	$\frac{x^2-3}{\sqrt{-x^2+3}}$	16
meijerg	$-\frac{\sqrt{3} \left(-2\sqrt{\pi} + 2\sqrt{\pi} \sqrt{-\frac{x^2}{3} + 1} \right)}{2\sqrt{\pi}}$	29

input `int(x/(-x^2+3)^(1/2),x,method=_RETURNVERBOSE)`output `-(-x^2+3)^(1/2)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{x}{\sqrt{3-x^2}} dx = -\sqrt{-x^2 + 3}$$

input `integrate(x/(-x^2+3)^(1/2),x, algorithm="fricas")`output `-sqrt(-x^2 + 3)`

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.62

$$\int \frac{x}{\sqrt{3-x^2}} dx = -\sqrt{3-x^2}$$

input `integrate(x/(-x**2+3)**(1/2),x)`

output `-sqrt(3 - x**2)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{x}{\sqrt{3-x^2}} dx = -\sqrt{-x^2+3}$$

input `integrate(x/(-x^2+3)^(1/2),x, algorithm="maxima")`

output `-sqrt(-x^2 + 3)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{x}{\sqrt{3-x^2}} dx = -\sqrt{-x^2+3}$$

input `integrate(x/(-x^2+3)^(1/2),x, algorithm="giac")`

output `-sqrt(-x^2 + 3)`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{x}{\sqrt{3-x^2}} dx = -\sqrt{3-x^2}$$

input `int(x/(3 - x^2)^(1/2), x)`

output `-(3 - x^2)^(1/2)`

Reduce [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \frac{x}{\sqrt{3-x^2}} dx = -\sqrt{-x^2+3}$$

input `int(x/(-x^2+3)^(1/2), x)`

output `- sqrt(- x**2 + 3)`

3.150 $\int \frac{\sqrt{3-x^2}}{x} dx$

Optimal result	863
Mathematica [A] (verified)	863
Rubi [A] (verified)	864
Maple [A] (verified)	865
Fricas [A] (verification not implemented)	866
Sympy [C] (verification not implemented)	867
Maxima [A] (verification not implemented)	867
Giac [A] (verification not implemented)	868
Mupad [B] (verification not implemented)	868
Reduce [B] (verification not implemented)	868

Optimal result

Integrand size = 15, antiderivative size = 37

$$\int \frac{\sqrt{3-x^2}}{x} dx = \sqrt{3-x^2} - \sqrt{3} \operatorname{arctanh}\left(\frac{\sqrt{3-x^2}}{\sqrt{3}}\right)$$

output `-arctanh(1/3*(-x^2+3)^(1/2)*3^(1/2))*3^(1/2)+(-x^2+3)^(1/2)`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.89

$$\int \frac{\sqrt{3-x^2}}{x} dx = \sqrt{3-x^2} - \sqrt{3} \operatorname{arctanh}\left(\sqrt{1-\frac{x^2}{3}}\right)$$

input `Integrate[Sqrt[3 - x^2]/x,x]`

output `Sqrt[3 - x^2] - Sqrt[3]*ArcTanh[Sqrt[1 - x^2/3]]`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.16, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {243, 60, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{3-x^2}}{x} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{2} \int \frac{\sqrt{3-x^2}}{x^2} dx^2 \\
 & \quad \downarrow \text{60} \\
 & \frac{1}{2} \left(3 \int \frac{1}{x^2 \sqrt{3-x^2}} dx^2 + 2\sqrt{3-x^2} \right) \\
 & \quad \downarrow \text{73} \\
 & \frac{1}{2} \left(2\sqrt{3-x^2} - 6 \int \frac{1}{3-x^4} d\sqrt{3-x^2} \right) \\
 & \quad \downarrow \text{219} \\
 & \frac{1}{2} \left(2\sqrt{3-x^2} - 2\sqrt{3} \operatorname{arctanh} \left(\frac{\sqrt{3-x^2}}{\sqrt{3}} \right) \right)
 \end{aligned}$$

input `Int[Sqrt[3 - x^2]/x,x]`

output `(2*Sqrt[3 - x^2] - 2*Sqrt[3]*ArcTanh[Sqrt[3 - x^2]/Sqrt[3]])/2`

Definitions of rubi rules used

- rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.81

method	result	size
default	$\sqrt{-x^2 + 3} - \sqrt{3} \operatorname{arctanh}\left(\frac{\sqrt{3}}{\sqrt{-x^2 + 3}}\right)$	30
pseudoelliptic	$-\operatorname{arctanh}\left(\frac{\sqrt{-x^2 + 3}\sqrt{3}}{3}\right) \sqrt{3} + \sqrt{-x^2 + 3}$	31
trager	$\sqrt{-x^2 + 3} + \operatorname{RootOf}(_Z^2 - 3) \ln\left(\frac{\sqrt{-x^2 + 3} - \operatorname{RootOf}(_Z^2 - 3)}{x}\right)$	41
meijerg	$\frac{\sqrt{3} \left(-2(2 - 2\ln(2) + 2\ln(x) - \ln(3) + i\pi)\sqrt{\pi} + 4\sqrt{\pi} - 4\sqrt{\pi} \sqrt{-\frac{x^2}{3} + 1} + 4\sqrt{\pi} \ln\left(\frac{1}{2} + \frac{\sqrt{-\frac{x^2}{3} + 1}}{2}\right) \right)}{4\sqrt{\pi}}$	71

input `int((-x^2+3)^(1/2)/x,x,method=_RETURNVERBOSE)`

output `(-x^2+3)^(1/2)-3^(1/2)*arctanh(3^(1/2)/(-x^2+3)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.08

$$\int \frac{\sqrt{3-x^2}}{x} dx = \frac{1}{2} \sqrt{3} \log\left(-\frac{x^2 + 2\sqrt{3}\sqrt{-x^2 + 3} - 6}{x^2}\right) + \sqrt{-x^2 + 3}$$

input `integrate((-x^2+3)^(1/2)/x,x, algorithm="fricas")`

output `1/2*sqrt(3)*log(-(x^2 + 2*sqrt(3)*sqrt(-x^2 + 3) - 6)/x^2) + sqrt(-x^2 + 3)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.74 (sec) , antiderivative size = 87, normalized size of antiderivative = 2.35

$$\int \frac{\sqrt{3-x^2}}{x} dx = \begin{cases} i\sqrt{x^2-3} - \sqrt{3}\log(x) + \frac{\sqrt{3}\log(x^2)}{2} + \sqrt{3}i \operatorname{asin}\left(\frac{\sqrt{3}}{x}\right) & \text{for } |x^2| > 3 \\ \sqrt{3-x^2} + \frac{\sqrt{3}\log(x^2)}{2} - \sqrt{3}\log\left(\sqrt{1-\frac{x^2}{3}}+1\right) & \text{otherwise} \end{cases}$$

input `integrate((-x**2+3)**(1/2)/x,x)`

output `Piecewise((I*sqrt(x**2 - 3) - sqrt(3)*log(x) + sqrt(3)*log(x**2)/2 + sqrt(3)*I*asin(sqrt(3)/x), Abs(x**2) > 3), (sqrt(3 - x**2) + sqrt(3)*log(x**2)/2 - sqrt(3)*log(sqrt(1 - x**2/3) + 1), True))`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.11

$$\int \frac{\sqrt{3-x^2}}{x} dx = -\sqrt{3}\log\left(\frac{2\sqrt{3}\sqrt{-x^2+3}}{|x|} + \frac{6}{|x|}\right) + \sqrt{-x^2+3}$$

input `integrate((-x^2+3)^(1/2)/x,x, algorithm="maxima")`

output `-sqrt(3)*log(2*sqrt(3)*sqrt(-x^2 + 3)/abs(x) + 6/abs(x)) + sqrt(-x^2 + 3)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.27

$$\int \frac{\sqrt{3-x^2}}{x} dx = \frac{1}{2} \sqrt{3} \log \left(\frac{\sqrt{3} - \sqrt{-x^2+3}}{\sqrt{3} + \sqrt{-x^2+3}} \right) + \sqrt{-x^2+3}$$

input `integrate((-x^2+3)^(1/2)/x,x, algorithm="giac")`output `1/2*sqrt(3)*log((sqrt(3) - sqrt(-x^2 + 3))/(sqrt(3) + sqrt(-x^2 + 3))) + sqrt(-x^2 + 3)`**Mupad [B] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.95

$$\int \frac{\sqrt{3-x^2}}{x} dx = \sqrt{3} \ln \left(\sqrt{\frac{3}{x^2} - 1} - \sqrt{3} \sqrt{\frac{1}{x^2}} \right) + \sqrt{3-x^2}$$

input `int((3 - x^2)^(1/2)/x,x)`output `3^(1/2)*log((3/x^2 - 1)^(1/2) - 3^(1/2)*(1/x^2)^(1/2)) + (3 - x^2)^(1/2)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.73

$$\int \frac{\sqrt{3-x^2}}{x} dx = \sqrt{-x^2+3} + \sqrt{3} \log \left(\tan \left(\frac{\arcsin\left(\frac{x}{\sqrt{3}}\right)}{2} \right) \right) - \sqrt{3}$$

input `int((-x^2+3)^(1/2)/x,x)`output `sqrt(-x**2 + 3) + sqrt(3)*log(tan(asin(x/sqrt(3))/2)) - sqrt(3)`

3.151 $\int \frac{\sqrt{x+x^2}}{x} dx$

Optimal result	869
Mathematica [A] (verified)	869
Rubi [A] (verified)	870
Maple [A] (verified)	871
Fricas [A] (verification not implemented)	871
Sympy [F]	872
Maxima [A] (verification not implemented)	872
Giac [A] (verification not implemented)	872
Mupad [B] (verification not implemented)	873
Reduce [B] (verification not implemented)	873

Optimal result

Integrand size = 13, antiderivative size = 22

$$\int \frac{\sqrt{x+x^2}}{x} dx = \sqrt{x+x^2} + \operatorname{arctanh}\left(\frac{x}{\sqrt{x+x^2}}\right)$$

output `arctanh(x/(x^2+x)^(1/2))+(x^2+x)^(1/2)`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.91

$$\int \frac{\sqrt{x+x^2}}{x} dx = \sqrt{x(1+x)} \left(1 - \frac{\log(-\sqrt{x} + \sqrt{1+x})}{\sqrt{x}\sqrt{1+x}} \right)$$

input `Integrate[Sqrt[x + x^2]/x,x]`

output `Sqrt[x*(1 + x)]*(1 - Log[-Sqrt[x] + Sqrt[1 + x]]/(Sqrt[x]*Sqrt[1 + x]))`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {1131, 1091, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{x^2 + x}}{x} dx$$

↓ 1131

$$\frac{1}{2} \int \frac{1}{\sqrt{x^2 + x}} dx + \sqrt{x^2 + x}$$

↓ 1091

$$\int \frac{1}{1 - \frac{x^2}{x^2 + x}} d \frac{x}{\sqrt{x^2 + x}} + \sqrt{x^2 + x}$$

↓ 219

$$\operatorname{arctanh}\left(\frac{x}{\sqrt{x^2 + x}}\right) + \sqrt{x^2 + x}$$

input `Int[Sqrt[x + x^2]/x,x]`

output `Sqrt[x + x^2] + ArcTanh[x/Sqrt[x + x^2]]`

Defintions of rubi rules used

rule 219

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

rule 1091

```
Int[1/Sqrt[(b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] :> Simp[2 Subst[Int[1/(1
- c*x^2), x], x, x/Sqrt[b*x + c*x^2]], x] /; FreeQ[{b, c}, x]
```

rule 1131

```
Int[((d_.) + (e_.)*(x_)^(m_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_S
ymbol] := Simp[(d + e*x)^(m + 1)*((a + b*x + c*x^2)^p/(e*(m + 2*p + 1))), x
] - Simp[p*((2*c*d - b*e)/(e^2*(m + 2*p + 1))) Int[(d + e*x)^(m + 1)*(a +
b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c*d^2 - b
*d*e + a*e^2, 0] && GtQ[p, 0] && (LeQ[-2, m, 0] || EqQ[m + p + 1, 0]) && Ne
Q[m + 2*p + 1, 0] && IntegerQ[2*p]
```

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

method	result	size
default	$\sqrt{x^2 + x} + \frac{\ln\left(x + \frac{1}{2} + \sqrt{x^2 + x}\right)}{2}$	22
trager	$\sqrt{x^2 + x} - \frac{\ln\left(2\sqrt{x^2 + x} - 1 - 2x\right)}{2}$	26
risch	$\frac{x(1+x)}{\sqrt{x(1+x)}} + \frac{\ln\left(x + \frac{1}{2} + \sqrt{x^2 + x}\right)}{2}$	27
meijerg	$-\frac{-2\sqrt{\pi}\sqrt{x}\sqrt{1+x} - 2\sqrt{\pi}\operatorname{arcsinh}(\sqrt{x})}{2\sqrt{\pi}}$	29
pseudoelliptic	$\sqrt{x(1+x)} - \frac{\ln\left(\frac{\sqrt{x(1+x)}-x}{x}\right)}{2} + \frac{\ln\left(\frac{\sqrt{x(1+x)}+x}{x}\right)}{2}$	43

input `int((x^2+x)^(1/2)/x,x,method=_RETURNVERBOSE)`output `(x^2+x)^(1/2)+1/2*ln(x+1/2+(x^2+x)^(1/2))`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.14

$$\int \frac{\sqrt{x+x^2}}{x} dx = \sqrt{x^2+x} - \frac{1}{2} \log\left(-2x + 2\sqrt{x^2+x} - 1\right)$$

input `integrate((x^2+x)^(1/2)/x,x, algorithm="fricas")`

output `sqrt(x^2 + x) - 1/2*log(-2*x + 2*sqrt(x^2 + x) - 1)`

Sympy [F]

$$\int \frac{\sqrt{x+x^2}}{x} dx = \int \frac{\sqrt{x(x+1)}}{x} dx$$

input `integrate((x**2+x)**(1/2)/x,x)`

output `Integral(sqrt(x*(x + 1))/x, x)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.14

$$\int \frac{\sqrt{x+x^2}}{x} dx = \sqrt{x^2+x} + \frac{1}{2} \log \left(2x + 2\sqrt{x^2+x} + 1 \right)$$

input `integrate((x^2+x)^(1/2)/x,x, algorithm="maxima")`

output `sqrt(x^2 + x) + 1/2*log(2*x + 2*sqrt(x^2 + x) + 1)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.18

$$\int \frac{\sqrt{x+x^2}}{x} dx = \sqrt{x^2+x} - \frac{1}{2} \log \left(\left| -2x + 2\sqrt{x^2+x} - 1 \right| \right)$$

input `integrate((x^2+x)^(1/2)/x,x, algorithm="giac")`

output `sqrt(x^2 + x) - 1/2*log(abs(-2*x + 2*sqrt(x^2 + x) - 1))`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.95

$$\int \frac{\sqrt{x+x^2}}{x} dx = \frac{\ln\left(x + \sqrt{x(x+1)} + \frac{1}{2}\right)}{2} + \sqrt{x^2+x}$$

input `int((x + x^2)^(1/2)/x,x)`output `log(x + (x*(x + 1))^(1/2) + 1/2)/2 + (x + x^2)^(1/2)`**Reduce [B] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \frac{\sqrt{x+x^2}}{x} dx = \sqrt{x} \sqrt{x+1} + \log(\sqrt{x+1} + \sqrt{x})$$

input `int((x^2+x)^(1/2)/x,x)`output `sqrt(x)*sqrt(x + 1) + log(sqrt(x + 1) + sqrt(x))`

3.152 $\int \sqrt{5 + x^2} dx$

Optimal result	874
Mathematica [A] (verified)	874
Rubi [A] (verified)	875
Maple [A] (verified)	876
Fricas [A] (verification not implemented)	876
Sympy [A] (verification not implemented)	877
Maxima [A] (verification not implemented)	877
Giac [A] (verification not implemented)	877
Mupad [B] (verification not implemented)	878
Reduce [B] (verification not implemented)	878

Optimal result

Integrand size = 9, antiderivative size = 27

$$\int \sqrt{5 + x^2} dx = \frac{1}{2}x\sqrt{5 + x^2} + \frac{5}{2}\operatorname{arcsinh}\left(\frac{x}{\sqrt{5}}\right)$$

output

```
5/2*arcsinh(1/5*x*5^(1/2))+1/2*x*(x^2+5)^(1/2)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.22

$$\int \sqrt{5 + x^2} dx = \frac{1}{2}x\sqrt{5 + x^2} - \frac{5}{2}\log\left(-x + \sqrt{5 + x^2}\right)$$

input

```
Integrate[Sqrt[5 + x^2],x]
```

output

```
(x*Sqrt[5 + x^2])/2 - (5*Log[-x + Sqrt[5 + x^2]])/2
```

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {211, 222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{x^2 + 5} dx$$

$$\downarrow \text{211}$$

$$\frac{5}{2} \int \frac{1}{\sqrt{x^2 + 5}} dx + \frac{1}{2} \sqrt{x^2 + 5} x$$

$$\downarrow \text{222}$$

$$\frac{5}{2} \operatorname{arcsinh}\left(\frac{x}{\sqrt{5}}\right) + \frac{1}{2} \sqrt{x^2 + 5} x$$

input `Int[Sqrt[5 + x^2], x]`

output `(x*Sqrt[5 + x^2])/2 + (5*ArcSinh[x/Sqrt[5]])/2`

Defintions of rubi rules used

rule 211 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[x*((a + b*x^2)^p/(2*p + 1)), x] + Simp[2*a*(p/(2*p + 1)) Int[(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 222 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

method	result	size
default	$\frac{5 \operatorname{arcsinh}\left(\frac{x\sqrt{5}}{5}\right)}{2} + \frac{x\sqrt{x^2+5}}{2}$	21
risch	$\frac{5 \operatorname{arcsinh}\left(\frac{x\sqrt{5}}{5}\right)}{2} + \frac{x\sqrt{x^2+5}}{2}$	21
trager	$\frac{x\sqrt{x^2+5}}{2} - \frac{5 \ln(x - \sqrt{x^2+5})}{2}$	26
meijerg	$5 \left(-\frac{2\sqrt{\pi} x \sqrt{5} \sqrt{1 + \frac{x^2}{5}}}{5} - 2\sqrt{\pi} \operatorname{arcsinh}\left(\frac{x\sqrt{5}}{5}\right) \right)$ $-\frac{\hspace{10em}}{4\sqrt{\pi}}$	37
pseudoelliptic	$\frac{x\sqrt{x^2+5}}{2} - \frac{5 \ln\left(\frac{\sqrt{x^2+5}-x}{x}\right)}{4} + \frac{5 \ln\left(\frac{\sqrt{x^2+5}+x}{x}\right)}{4}$	46

input `int((x^2+5)^(1/2),x,method=_RETURNVERBOSE)`output `5/2*arcsinh(1/5*x*5^(1/2))+1/2*x*(x^2+5)^(1/2)`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.93

$$\int \sqrt{5+x^2} dx = \frac{1}{2} \sqrt{x^2+5}x - \frac{5}{2} \log(-x + \sqrt{x^2+5})$$

input `integrate((x^2+5)^(1/2),x, algorithm="fricas")`output `1/2*sqrt(x^2 + 5)*x - 5/2*log(-x + sqrt(x^2 + 5))`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89

$$\int \sqrt{5+x^2} dx = \frac{x\sqrt{x^2+5}}{2} + \frac{5 \operatorname{asinh}\left(\frac{\sqrt{5}x}{5}\right)}{2}$$

input `integrate((x**2+5)**(1/2),x)`output `x*sqrt(x**2 + 5)/2 + 5*asinh(sqrt(5)*x/5)/2`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

$$\int \sqrt{5+x^2} dx = \frac{1}{2} \sqrt{x^2+5}x + \frac{5}{2} \operatorname{arsinh}\left(\frac{1}{5} \sqrt{5}x\right)$$

input `integrate((x^2+5)^(1/2),x, algorithm="maxima")`output `1/2*sqrt(x^2 + 5)*x + 5/2*arcsinh(1/5*sqrt(5)*x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.93

$$\int \sqrt{5+x^2} dx = \frac{1}{2} \sqrt{x^2+5}x - \frac{5}{2} \log\left(-x + \sqrt{x^2+5}\right)$$

input `integrate((x^2+5)^(1/2),x, algorithm="giac")`output `1/2*sqrt(x^2 + 5)*x - 5/2*log(-x + sqrt(x^2 + 5))`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

$$\int \sqrt{5+x^2} dx = \frac{5 \operatorname{asinh}\left(\frac{\sqrt{5}x}{5}\right)}{2} + \frac{x\sqrt{x^2+5}}{2}$$

input `int((x^2 + 5)^(1/2), x)`output `(5*asinh((5^(1/2)*x)/5))/2 + (x*(x^2 + 5)^(1/2))/2`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \sqrt{5+x^2} dx = \frac{\sqrt{x^2+5}x}{2} + \frac{5 \log\left(\frac{\sqrt{x^2+5}+x}{\sqrt{5}}\right)}{2}$$

input `int((x^2+5)^(1/2), x)`output `(sqrt(x**2 + 5)*x + 5*log((sqrt(x**2 + 5) + x)/sqrt(5)))/2`

3.153 $\int \frac{x}{\sqrt{1+x+x^2}} dx$

Optimal result	879
Mathematica [A] (verified)	879
Rubi [A] (verified)	880
Maple [A] (verified)	881
Fricas [A] (verification not implemented)	881
Sympy [A] (verification not implemented)	882
Maxima [A] (verification not implemented)	882
Giac [A] (verification not implemented)	882
Mupad [B] (verification not implemented)	883
Reduce [B] (verification not implemented)	883

Optimal result

Integrand size = 12, antiderivative size = 27

$$\int \frac{x}{\sqrt{1+x+x^2}} dx = \sqrt{1+x+x^2} - \frac{1}{2} \operatorname{arcsinh}\left(\frac{1+2x}{\sqrt{3}}\right)$$

output `-1/2*arcsinh(1/3*(1+2*x)*3^(1/2))+(x^2+x+1)^(1/2)`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.22

$$\int \frac{x}{\sqrt{1+x+x^2}} dx = \sqrt{1+x+x^2} + \frac{1}{2} \log\left(-1-2x+2\sqrt{1+x+x^2}\right)$$

input `Integrate[x/Sqrt[1 + x + x^2],x]`

output `Sqrt[1 + x + x^2] + Log[-1 - 2*x + 2*Sqrt[1 + x + x^2]]/2`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {1160, 1090, 222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x}{\sqrt{x^2 + x + 1}} dx \\
 & \quad \downarrow \text{1160} \\
 & \sqrt{x^2 + x + 1} - \frac{1}{2} \int \frac{1}{\sqrt{x^2 + x + 1}} dx \\
 & \quad \downarrow \text{1090} \\
 & \sqrt{x^2 + x + 1} - \frac{\int \frac{1}{\sqrt{\frac{1}{3}(2x+1)^2 + 1}} d(2x+1)}{2\sqrt{3}} \\
 & \quad \downarrow \text{222} \\
 & \sqrt{x^2 + x + 1} - \frac{1}{2} \operatorname{arcsinh}\left(\frac{2x+1}{\sqrt{3}}\right)
 \end{aligned}$$

input `Int[x/Sqrt[1 + x + x^2],x]`

output `Sqrt[1 + x + x^2] - ArcSinh[(1 + 2*x)/Sqrt[3]]/2`

Defintions of rubi rules used

rule 222 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

rule 1090 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p) Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

rule 1160

```
Int[((d_.) + (e_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol
] := Simp[e*((a + b*x + c*x^2)^(p + 1)/(2*c*(p + 1))), x] + Simp[(2*c*d - b
*e)/(2*c) Int[(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x]
&& NeQ[p, -1]
```

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

method	result	size
default	$\sqrt{x^2 + x + 1} - \frac{\operatorname{arcsinh}\left(\frac{2\sqrt{3}\left(x + \frac{1}{2}\right)}{3}\right)}{2}$	21
risch	$\sqrt{x^2 + x + 1} - \frac{\operatorname{arcsinh}\left(\frac{2\sqrt{3}\left(x + \frac{1}{2}\right)}{3}\right)}{2}$	21
trager	$\sqrt{x^2 + x + 1} - \frac{\ln\left(2x + 1 + 2\sqrt{x^2 + x + 1}\right)}{2}$	28

input

```
int(x/(x^2+x+1)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
(x^2+x+1)^(1/2)-1/2*arcsinh(2/3*3^(1/2)*(x+1/2))
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{x}{\sqrt{1+x+x^2}} dx = \sqrt{x^2+x+1} + \frac{1}{2} \log\left(-2x + 2\sqrt{x^2+x+1} - 1\right)$$

input

```
integrate(x/(x^2+x+1)^(1/2),x, algorithm="fricas")
```

output

```
sqrt(x^2 + x + 1) + 1/2*log(-2*x + 2*sqrt(x^2 + x + 1) - 1)
```

Sympy [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \frac{x}{\sqrt{1+x+x^2}} dx = \sqrt{x^2+x+1} - \frac{\operatorname{asinh}\left(\frac{2\sqrt{3}(x+\frac{1}{2})}{3}\right)}{2}$$

input `integrate(x/(x**2+x+1)**(1/2),x)`output `sqrt(x**2 + x + 1) - asinh(2*sqrt(3)*(x + 1/2)/3)/2`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int \frac{x}{\sqrt{1+x+x^2}} dx = \sqrt{x^2+x+1} - \frac{1}{2} \operatorname{arsinh}\left(\frac{1}{3}\sqrt{3}(2x+1)\right)$$

input `integrate(x/(x^2+x+1)^(1/2),x, algorithm="maxima")`output `sqrt(x^2 + x + 1) - 1/2*arcsinh(1/3*sqrt(3)*(2*x + 1))`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{x}{\sqrt{1+x+x^2}} dx = \sqrt{x^2+x+1} + \frac{1}{2} \log\left(-2x + 2\sqrt{x^2+x+1} - 1\right)$$

input `integrate(x/(x^2+x+1)^(1/2),x, algorithm="giac")`output `sqrt(x^2 + x + 1) + 1/2*log(-2*x + 2*sqrt(x^2 + x + 1) - 1)`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \frac{x}{\sqrt{1+x+x^2}} dx = \sqrt{x^2+x+1} - \frac{\ln\left(x + \sqrt{x^2+x+1} + \frac{1}{2}\right)}{2}$$

input `int(x/(x + x^2 + 1)^(1/2),x)`output `(x + x^2 + 1)^(1/2) - log(x + (x + x^2 + 1)^(1/2) + 1/2)/2`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.11

$$\int \frac{x}{\sqrt{1+x+x^2}} dx = \sqrt{x^2+x+1} - \frac{\log\left(\frac{2\sqrt{x^2+x+1}+2x+1}{\sqrt{3}}\right)}{2}$$

input `int(x/(x^2+x+1)^(1/2),x)`output `(2*sqrt(x**2 + x + 1) - log((2*sqrt(x**2 + x + 1) + 2*x + 1)/sqrt(3)))/2`

3.154 $\int \frac{1}{\sqrt{x+x^2}} dx$

Optimal result	884
Mathematica [B] (verified)	884
Rubi [A] (verified)	885
Maple [A] (verified)	886
Fricas [A] (verification not implemented)	886
Sympy [A] (verification not implemented)	886
Maxima [A] (verification not implemented)	887
Giac [B] (verification not implemented)	887
Mupad [B] (verification not implemented)	887
Reduce [B] (verification not implemented)	888

Optimal result

Integrand size = 9, antiderivative size = 14

$$\int \frac{1}{\sqrt{x+x^2}} dx = 2\operatorname{arctanh}\left(\frac{x}{\sqrt{x+x^2}}\right)$$

output `2*arctanh(x/(x^2+x)^(1/2))`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 39 vs. 2(14) = 28.

Time = 0.02 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.79

$$\int \frac{1}{\sqrt{x+x^2}} dx = -\frac{2\sqrt{x}\sqrt{1+x}\log(-\sqrt{x} + \sqrt{1+x})}{\sqrt{x(1+x)}}$$

input `Integrate[1/Sqrt[x + x^2],x]`

output `(-2*Sqrt[x]*Sqrt[1 + x]*Log[-Sqrt[x] + Sqrt[1 + x]])/Sqrt[x*(1 + x)]`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {1091, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x^2 + x}} dx$$

↓ 1091

$$2 \int \frac{1}{1 - \frac{x^2}{x^2+x}} d \frac{x}{\sqrt{x^2 + x}}$$

↓ 219

$$2 \operatorname{arctanh} \left(\frac{x}{\sqrt{x^2 + x}} \right)$$

input `Int[1/Sqrt[x + x^2], x]`

output `2*ArcTanh[x/Sqrt[x + x^2]]`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 1091 `Int[1/Sqrt[(b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Simp[2 Subst[Int[1/(1 - c*x^2), x], x, x/Sqrt[b*x + c*x^2]], x] /; FreeQ[{b, c}, x]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.50

method	result	size
meijerg	$2 \operatorname{arcsinh}(\sqrt{x})$	7
default	$\ln\left(x + \frac{1}{2} + \sqrt{x^2 + x}\right)$	12
pseudoelliptic	$2 \operatorname{arctanh}\left(\frac{\sqrt{x(1+x)}}{x}\right)$	15
trager	$\ln\left(1 + 2x + 2\sqrt{x^2 + x}\right)$	16

input `int(1/(x^2+x)^(1/2),x,method=_RETURNVERBOSE)`output `2*arcsinh(x^(1/2))`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.21

$$\int \frac{1}{\sqrt{x+x^2}} dx = -\log\left(-2x + 2\sqrt{x^2+x} - 1\right)$$

input `integrate(1/(x^2+x)^(1/2),x, algorithm="fricas")`output `-log(-2*x + 2*sqrt(x^2 + x) - 1)`**Sympy [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

$$\int \frac{1}{\sqrt{x+x^2}} dx = \log\left(2x + 2\sqrt{x^2+x} + 1\right)$$

input `integrate(1/(x**2+x)**(1/2),x)`

output `log(2*x + 2*sqrt(x**2 + x) + 1)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

$$\int \frac{1}{\sqrt{x+x^2}} dx = \log\left(2x + 2\sqrt{x^2+x} + 1\right)$$

input `integrate(1/(x^2+x)^(1/2),x, algorithm="maxima")`

output `log(2*x + 2*sqrt(x^2 + x) + 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 33 vs. $2(12) = 24$.

Time = 0.12 (sec) , antiderivative size = 33, normalized size of antiderivative = 2.36

$$\int \frac{1}{\sqrt{x+x^2}} dx = \frac{1}{4} \sqrt{x^2+x}(2x+1) + \frac{1}{8} \log\left(\left|-2x + 2\sqrt{x^2+x} - 1\right|\right)$$

input `integrate(1/(x^2+x)^(1/2),x, algorithm="giac")`

output `1/4*sqrt(x^2 + x)*(2*x + 1) + 1/8*log(abs(-2*x + 2*sqrt(x^2 + x) - 1))`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.79

$$\int \frac{1}{\sqrt{x+x^2}} dx = \ln\left(x + \sqrt{x(x+1)} + \frac{1}{2}\right)$$

input `int(1/(x + x^2)^(1/2),x)`

output `log(x + (x*(x + 1))^(1/2) + 1/2)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{1}{\sqrt{x+x^2}} dx = 2 \log(\sqrt{x+1} + \sqrt{x})$$

input `int(1/(x^2+x)^(1/2),x)`

output `2*log(sqrt(x + 1) + sqrt(x))`

3.155 $\int \frac{\sqrt{2-x-x^2}}{x^2} dx$

Optimal result	889
Mathematica [A] (verified)	889
Rubi [A] (verified)	890
Maple [A] (verified)	892
Fricas [A] (verification not implemented)	892
Sympy [F]	893
Maxima [A] (verification not implemented)	893
Giac [B] (verification not implemented)	894
Mupad [B] (verification not implemented)	894
Reduce [B] (verification not implemented)	895

Optimal result

Integrand size = 18, antiderivative size = 68

$$\int \frac{\sqrt{2-x-x^2}}{x^2} dx = -\frac{\sqrt{2-x-x^2}}{x} + \arcsin\left(\frac{1}{3}(-1-2x)\right) + \frac{\operatorname{arctanh}\left(\frac{4-x}{2\sqrt{2}\sqrt{2-x-x^2}}\right)}{2\sqrt{2}}$$

output `-arcsin(1/3+2/3*x)+1/4*arctanh(1/4*(4-x)*2^(1/2)/(-x^2-x+2)^(1/2))*2^(1/2)
-(-x^2-x+2)^(1/2)/x`

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.12

$$\int \frac{\sqrt{2-x-x^2}}{x^2} dx = -\frac{\sqrt{2-x-x^2}}{x} + 2 \arctan\left(\frac{\sqrt{2-x-x^2}}{2+x}\right) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{2-x-x^2}}{\sqrt{2}(-1+x)}\right)}{\sqrt{2}}$$

input `Integrate[Sqrt[2 - x - x^2]/x^2,x]`

output `-(Sqrt[2 - x - x^2]/x) + 2*ArcTan[Sqrt[2 - x - x^2]/(2 + x)] - ArcTanh[Sqr
t[2 - x - x^2]/(Sqrt[2]*(-1 + x))]/Sqrt[2]`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.06, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {1161, 25, 1269, 1090, 223, 1154, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{-x^2 - x + 2}}{x^2} dx \\
 & \quad \downarrow \text{1161} \\
 & \frac{1}{2} \int -\frac{2x+1}{x\sqrt{-x^2-x+2}} dx - \frac{\sqrt{-x^2-x+2}}{x} \\
 & \quad \downarrow \text{25} \\
 & -\frac{1}{2} \int \frac{2x+1}{x\sqrt{-x^2-x+2}} dx - \frac{\sqrt{-x^2-x+2}}{x} \\
 & \quad \downarrow \text{1269} \\
 & \frac{1}{2} \left(-2 \int \frac{1}{\sqrt{-x^2-x+2}} dx - \int \frac{1}{x\sqrt{-x^2-x+2}} dx \right) - \frac{\sqrt{-x^2-x+2}}{x} \\
 & \quad \downarrow \text{1090} \\
 & \frac{1}{2} \left(\frac{2}{3} \int \frac{1}{\sqrt{1-\frac{1}{9}(-2x-1)^2}} d(-2x-1) - \int \frac{1}{x\sqrt{-x^2-x+2}} dx \right) - \frac{\sqrt{-x^2-x+2}}{x} \\
 & \quad \downarrow \text{223} \\
 & \frac{1}{2} \left(2 \arcsin \left(\frac{1}{3}(-2x-1) \right) - \int \frac{1}{x\sqrt{-x^2-x+2}} dx \right) - \frac{\sqrt{-x^2-x+2}}{x} \\
 & \quad \downarrow \text{1154} \\
 & \frac{1}{2} \left(2 \int \frac{1}{8 - \frac{(4-x)^2}{-x^2-x+2}} d \frac{4-x}{\sqrt{-x^2-x+2}} + 2 \arcsin \left(\frac{1}{3}(-2x-1) \right) \right) - \frac{\sqrt{-x^2-x+2}}{x} \\
 & \quad \downarrow \text{219} \\
 & \frac{1}{2} \left(2 \arcsin \left(\frac{1}{3}(-2x-1) \right) + \frac{\operatorname{arctanh} \left(\frac{4-x}{2\sqrt{2}\sqrt{-x^2-x+2}} \right)}{\sqrt{2}} \right) - \frac{\sqrt{-x^2-x+2}}{x}
 \end{aligned}$$

input `Int[Sqrt[2 - x - x^2]/x^2,x]`

output `-(Sqrt[2 - x - x^2]/x) + (2*ArcSin[(-1 - 2*x)/3] + ArcTanh[(4 - x)/(2*Sqrt[2]*Sqrt[2 - x - x^2]])/Sqrt[2])/2`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 223 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 1090 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p) Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

rule 1154 `Int[1/(((d_) + (e_)*(x_))*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2]), x_Symbol] := Simp[-2 Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 1161 `Int[((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*x + c*x^2)^p/(e*(m + 1))), x] - Simp[p/(e*(m + 1)) Int[(d + e*x)^(m + 1)*(b + 2*c*x)*(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && GtQ[p, 0] && (IntegerQ[p] || LtQ[m, -1]) && NeQ[m, -1] && !ILtQ[m + 2*p + 1, 0] && IntQuadraticQ[a, b, c, d, e, m, p, x]`

rule 1269

```
Int[((d_.) + (e_.)*(x_)^(m_))*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[g/e Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Simp[(e*f - d*g)/e Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && !IGtQ[m, 0]
```

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.88

method	result
risch	$\frac{x^2+x-2}{x\sqrt{-x^2-x+2}} + \frac{\operatorname{arctanh}\left(\frac{(4-x)\sqrt{2}}{4\sqrt{-x^2-x+2}}\right)\sqrt{2}}{4} - \arcsin\left(\frac{1}{3} + \frac{2x}{3}\right)$
default	$-\frac{(-x^2-x+2)^{\frac{3}{2}}}{2x} - \frac{\sqrt{-x^2-x+2}}{4} - \arcsin\left(\frac{1}{3} + \frac{2x}{3}\right) + \frac{\operatorname{arctanh}\left(\frac{(4-x)\sqrt{2}}{4\sqrt{-x^2-x+2}}\right)\sqrt{2}}{4} + \frac{(-2x-1)\sqrt{-x^2-x+2}}{4}$
trager	$-\frac{\sqrt{-x^2-x+2}}{x} - \operatorname{RootOf}(_Z^2 + 1) \ln(-2 \operatorname{RootOf}(_Z^2 + 1) x + 2\sqrt{-x^2-x+2}) - \operatorname{RootOf}(_Z^2 + 1)$

input

```
int((-x^2-x+2)^(1/2)/x^2,x,method=_RETURNVERBOSE)
```

output

```
(x^2+x-2)/x/(-x^2-x+2)^(1/2)+1/4*arctanh(1/4*(4-x)*2^(1/2)/(-x^2-x+2)^(1/2))*2^(1/2)-arcsin(1/3+2/3*x)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.35

$$\int \frac{\sqrt{2-x-x^2}}{x^2} dx$$

$$= \frac{\sqrt{2}x \log\left(-\frac{4\sqrt{2}\sqrt{-x^2-x+2}(x-4)+7x^2+16x-32}{x^2}\right) + 8x \arctan\left(\frac{\sqrt{-x^2-x+2}(2x+1)}{2(x^2+x-2)}\right) - 8\sqrt{-x^2-x+2}}{8x}$$

input

```
integrate((-x^2-x+2)^(1/2)/x^2,x, algorithm="fricas")
```

output $1/8*(\sqrt{2}*x*\log(-4*\sqrt{2}*\sqrt{-x^2 - x + 2}*(x - 4) + 7*x^2 + 16*x - 32)/x^2) + 8*x*\arctan(1/2*\sqrt{-x^2 - x + 2}*(2*x + 1)/(x^2 + x - 2)) - 8*\sqrt{-x^2 - x + 2})/x$

Sympy [F]

$$\int \frac{\sqrt{2-x-x^2}}{x^2} dx = \int \frac{\sqrt{-(x-1)(x+2)}}{x^2} dx$$

input `integrate((-x**2-x+2)**(1/2)/x**2,x)`

output `Integral(sqrt(-(x - 1)*(x + 2))/x**2, x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.87

$$\int \frac{\sqrt{2-x-x^2}}{x^2} dx = \frac{1}{4} \sqrt{2} \log \left(\frac{2\sqrt{2}\sqrt{-x^2-x+2}}{|x|} + \frac{4}{|x|} - 1 \right) - \frac{\sqrt{-x^2-x+2}}{x} + \arcsin \left(-\frac{2}{3}x - \frac{1}{3} \right)$$

input `integrate((-x^2-x+2)^(1/2)/x^2,x, algorithm="maxima")`

output $1/4*\sqrt{2}*\log(2*\sqrt{2}*\sqrt{-x^2 - x + 2}/\text{abs}(x) + 4/\text{abs}(x) - 1) - \sqrt{-x^2 - x + 2}/x + \arcsin(-2/3*x - 1/3)$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 168 vs. $2(52) = 104$.

Time = 0.15 (sec) , antiderivative size = 168, normalized size of antiderivative = 2.47

$$\int \frac{\sqrt{2-x-x^2}}{x^2} dx = -\frac{1}{4} \sqrt{2} \log \left(\frac{\left| -4\sqrt{2} + \frac{2(2\sqrt{-x^2-x+2}-3)}{2x+1} + 6 \right|}{\left| 4\sqrt{2} + \frac{2(2\sqrt{-x^2-x+2}-3)}{2x+1} + 6 \right|} \right) + \frac{6 \left(\frac{3(2\sqrt{-x^2-x+2}-3)}{2x+1} + 1 \right)}{\frac{6(2\sqrt{-x^2-x+2}-3)}{2x+1} + \frac{(2\sqrt{-x^2-x+2}-3)^2}{(2x+1)^2} + 1} - \arcsin \left(\frac{2}{3}x + \frac{1}{3} \right)$$

input `integrate((-x^2-x+2)^(1/2)/x^2,x, algorithm="giac")`

output `-1/4*sqrt(2)*log(abs(-4*sqrt(2) + 2*(2*sqrt(-x^2 - x + 2) - 3)/(2*x + 1) + 6)/abs(4*sqrt(2) + 2*(2*sqrt(-x^2 - x + 2) - 3)/(2*x + 1) + 6)) + 6*(3*(2*sqrt(-x^2 - x + 2) - 3)/(2*x + 1) + 1)/(6*(2*sqrt(-x^2 - x + 2) - 3)/(2*x + 1) + (2*sqrt(-x^2 - x + 2) - 3)^2/(2*x + 1)^2 + 1) - arcsin(2/3*x + 1/3)`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.07

$$\int \frac{\sqrt{2-x-x^2}}{x^2} dx = \frac{\sqrt{2} \ln \left(\frac{2}{x} + \frac{\sqrt{2}\sqrt{-x^2-x+2}}{x} - \frac{1}{2} \right)}{4} - \frac{\sqrt{-x^2-x+2}}{x} + \ln \left(x \operatorname{li} + \sqrt{-x^2-x+2} + \frac{1}{2}i \right) \operatorname{li}$$

input `int((2 - x^2 - x)^(1/2)/x^2,x)`

output `log(x*i + (2 - x^2 - x)^(1/2) + i/2)*i - (2 - x^2 - x)^(1/2)/x + (2^(1/2)*log(2/x + (2^(1/2)*(2 - x^2 - x)^(1/2))/x - 1/2))/4`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.01

$$\int \frac{\sqrt{2-x-x^2}}{x^2} dx$$

$$= \frac{-4\operatorname{asin}\left(\frac{2x}{3} + \frac{1}{3}\right)x - 4\sqrt{-x^2 - x + 2} + \sqrt{2}\log\left(-2\sqrt{2} + \tan\left(\frac{\operatorname{asin}\left(\frac{2x}{3} + \frac{1}{3}\right)}{2}\right) - 3\right)x - \sqrt{2}\log\left(2\sqrt{2} + \tan\left(\frac{\operatorname{asin}\left(\frac{2x}{3} + \frac{1}{3}\right)}{2}\right) - 3\right)x}{4x}$$

input

```
int((-x^2-x+2)^(1/2)/x^2,x)
```

output

```
( - 4*asin((2*x + 1)/3)*x - 4*sqrt( - x**2 - x + 2) + sqrt(2)*log( - 2*sqrt(2) + tan(asin((2*x + 1)/3)/2) - 3)*x - sqrt(2)*log(2*sqrt(2) + tan(asin((2*x + 1)/3)/2) - 3)*x)/(4*x)
```


3.156 $\int \frac{\log(t)}{1+t} dt$

Optimal result	896
Mathematica [A] (verified)	896
Rubi [A] (verified)	897
Maple [A] (verified)	898
Fricas [F]	898
Sympy [C] (verification not implemented)	899
Maxima [A] (verification not implemented)	899
Giac [F]	900
Mupad [B] (verification not implemented)	900
Reduce [F]	900

Optimal result

Integrand size = 8, antiderivative size = 13

$$\int \frac{\log(t)}{1+t} dt = \log(t) \log(1+t) + \text{PolyLog}(2, -t)$$

output

`ln(t)*ln(1+t)+polylog(2,-t)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\log(t)}{1+t} dt = \log(t) \log(1+t) + \text{PolyLog}(2, -t)$$

input

`Integrate[Log[t]/(1 + t),t]`

output

`Log[t]*Log[1 + t] + PolyLog[2, -t]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2754, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\log(t)}{t+1} dt$$

↓ 2754

$$\log(t) \log(t+1) - \int \frac{\log(t+1)}{t} dt$$

↓ 2838

$$\text{PolyLog}(2, -t) + \log(t) \log(t+1)$$

input `Int[Log[t]/(1 + t), t]`

output `Log[t]*Log[1 + t] + PolyLog[2, -t]`

Defintions of rubi rules used

rule 2754 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)/((d_) + (e_.)*(x_)), x_Symbol] :> Simp[Log[1 + e*(x/d)]*((a + b*Log[c*x^n])^p/e), x] - Simp[b*n*(p/e) Int[Log[1 + e*(x/d)]*((a + b*Log[c*x^n])^(p - 1)/x), x], x] /; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[p, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

method	result	size
default	$\operatorname{dilog}(1+t) + \ln(t) \ln(1+t)$	13
risch	$\operatorname{dilog}(1+t) + \ln(t) \ln(1+t)$	13
parts	$\operatorname{dilog}(1+t) + \ln(t) \ln(1+t)$	13

input `int(ln(t)/(1+t),t,method=_RETURNVERBOSE)`

output `dilog(1+t)+ln(t)*ln(1+t)`

Fricas [F]

$$\int \frac{\log(t)}{1+t} dt = \int \frac{\log(t)}{t+1} dt$$

input `integrate(log(t)/(1+t),t, algorithm="fricas")`

output `integral(log(t)/(t + 1), t)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.90 (sec) , antiderivative size = 73, normalized size of antiderivative = 5.62

$$\int \frac{\log(t)}{1+t} dt = \begin{cases} -\operatorname{Li}_2(t+1) & \text{for } \frac{1}{|t+1|} < 1 \wedge |t+1| < 1 \\ i\pi \log(t+1) - \operatorname{Li}_2(t+1) & \text{for } |t+1| < 1 \\ -i\pi \log\left(\frac{1}{t+1}\right) - \operatorname{Li}_2(t+1) & \text{for } \frac{1}{|t+1|} < 1 \\ -i\pi G_{2,2}^{2,0}\left(0,0 \left| \begin{matrix} 1,1 \\ t+1 \end{matrix} \right.\right) + i\pi G_{2,2}^{0,2}\left(1,1 \left| \begin{matrix} 1,1 \\ 0,0 \end{matrix} \right| t+1\right) - \operatorname{Li}_2(t+1) & \text{otherwise} \end{cases}$$

input `integrate(ln(t)/(1+t),t)`

output `Piecewise((-polylog(2, t + 1), (Abs(t + 1) < 1) & (1/Abs(t + 1) < 1)), (I*pi*log(t + 1) - polylog(2, t + 1), Abs(t + 1) < 1), (-I*pi*log(1/(t + 1)) - polylog(2, t + 1), 1/Abs(t + 1) < 1), (-I*pi*meijerg(((), (1, 1)), ((0, 0), ()), t + 1) + I*pi*meijerg(((1, 1), ()), (((), (0, 0))), t + 1) - polylog(2, t + 1), True))`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

$$\int \frac{\log(t)}{1+t} dt = \log(t+1) \log(t) + \operatorname{Li}_2(-t)$$

input `integrate(log(t)/(1+t),t, algorithm="maxima")`

output `log(t + 1)*log(t) + dilog(-t)`

Giac [F]

$$\int \frac{\log(t)}{1+t} dt = \int \frac{\log(t)}{t+1} dt$$

input `integrate(log(t)/(1+t),t, algorithm="giac")`

output `integrate(log(t)/(t + 1), t)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\log(t)}{1+t} dt = \text{polylog}(2, -t) + \ln(t+1) \ln(t)$$

input `int(log(t)/(t + 1),t)`

output `polylog(2, -t) + log(t + 1)*log(t)`

Reduce [F]

$$\int \frac{\log(t)}{1+t} dt = - \left(\int \frac{\log(t)}{t^2+t} dt \right) + \frac{\log(t)^2}{2}$$

input `int(log(t)/(1+t),t)`

output `(- 2*int(log(t)/(t**2 + t),t) + log(t)**2)/2`

3.157 $\int \log(e^{\cos(x)}) dx$

Optimal result	901
Mathematica [A] (verified)	901
Rubi [A] (verified)	902
Maple [A] (verified)	903
Fricas [A] (verification not implemented)	904
Sympy [A] (verification not implemented)	904
Maxima [A] (verification not implemented)	904
Giac [A] (verification not implemented)	905
Mupad [B] (verification not implemented)	905
Reduce [B] (verification not implemented)	905

Optimal result

Integrand size = 5, antiderivative size = 15

$$\int \log(e^{\cos(x)}) dx = -x \cos(x) + x \log(e^{\cos(x)}) + \sin(x)$$

output `-x*cos(x)+x*ln(exp(cos(x)))+sin(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \log(e^{\cos(x)}) dx = x(-\cos(x) + \log(e^{\cos(x)})) + \sin(x)$$

input `Integrate[Log[E^Cos[x]],x]`

output `x*(-Cos[x] + Log[E^Cos[x]]) + Sin[x]`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 1.200$, Rules used = {3028, 25, 3042, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \log \left(e^{\cos(x)} \right) dx \\
 & \quad \downarrow \text{3028} \\
 & x \log \left(e^{\cos(x)} \right) - \int -x \sin(x) dx \\
 & \quad \downarrow \text{25} \\
 & \int x \sin(x) dx + x \log \left(e^{\cos(x)} \right) \\
 & \quad \downarrow \text{3042} \\
 & \int x \sin(x) dx + x \log \left(e^{\cos(x)} \right) \\
 & \quad \downarrow \text{3777} \\
 & \int \cos(x) dx - x \cos(x) + x \log \left(e^{\cos(x)} \right) \\
 & \quad \downarrow \text{3042} \\
 & \int \sin \left(x + \frac{\pi}{2} \right) dx - x \cos(x) + x \log \left(e^{\cos(x)} \right) \\
 & \quad \downarrow \text{3117} \\
 & \sin(x) - x \cos(x) + x \log \left(e^{\cos(x)} \right)
 \end{aligned}$$

input

`Int [Log [E^Cos [x]] , x]`

output

`-(x*Cos [x]) + x*Log [E^Cos [x]] + Sin [x]`

Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3028 `Int[Log[u_], x_Symbol] := Simp[x*Log[u], x] - Int[SimplifyIntegrand[x*(D[u, x]/u), x], x] /; InverseFunctionFreeQ[u, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`
- rule 3777 `Int[((c_.) + (d_.)*(x_)^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

method	result	size
default	$-x \cos(x) + x \ln(e^{\cos(x)}) + \sin(x)$	15
risch	$-x \cos(x) + x \ln(e^{\cos(x)}) + \sin(x)$	15
parallelrisch	$-x \cos(x) + x \ln(e^{\cos(x)}) + \sin(x)$	15
parts	$-x \cos(x) + x \ln(e^{\cos(x)}) + \sin(x)$	15

input `int(ln(exp(cos(x))), x, method=_RETURNVERBOSE)`

output `-x*cos(x)+x*ln(exp(cos(x)))+sin(x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.13

$$\int \log(e^{\cos(x)}) dx = \sin(x)$$

input `integrate(log(exp(cos(x))),x, algorithm="fricas")`

output `sin(x)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \log(e^{\cos(x)}) dx = x \log(e^{\cos(x)}) - x \cos(x) + \sin(x)$$

input `integrate(ln(exp(cos(x))),x)`

output `x*log(exp(cos(x))) - x*cos(x) + sin(x)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.13

$$\int \log(e^{\cos(x)}) dx = \sin(x)$$

input `integrate(log(exp(cos(x))),x, algorithm="maxima")`

output `sin(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.13

$$\int \log(e^{\cos(x)}) dx = \sin(x)$$

input `integrate(log(exp(cos(x))),x, algorithm="giac")`

output `sin(x)`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.13

$$\int \log(e^{\cos(x)}) dx = \sin(x)$$

input `int(log(exp(cos(x))),x)`

output `sin(x)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.13

$$\int \log(e^{\cos(x)}) dx = \sin(x)$$

input `int(log(exp(cos(x))),x)`

output `sin(x)`

3.158 $\int \frac{e^t}{t} dt$

Optimal result	906
Mathematica [A] (verified)	906
Rubi [A] (verified)	907
Maple [B] (verified)	907
Fricas [A] (verification not implemented)	908
Sympy [A] (verification not implemented)	908
Maxima [A] (verification not implemented)	909
Giac [A] (verification not implemented)	909
Mupad [B] (verification not implemented)	909
Reduce [B] (verification not implemented)	910

Optimal result

Integrand size = 7, antiderivative size = 2

$$\int \frac{e^t}{t} dt = \text{ExpIntegralEi}(t)$$

output `Ei(t)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{e^t}{t} dt = \text{ExpIntegralEi}(t)$$

input `Integrate[E^t/t,t]`

output `ExpIntegralEi[t]`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2609}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^t}{t} dt$$

↓ 2609

$$\text{ExpIntegralEi}(t)$$

input `Int[E^t/t,t]`

output `ExpIntegralEi[t]`

Defintions of rubi rules used

rule 2609 `Int[(F_)^((g_.)*((e_.) + (f_.)*(x_)))/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 7 vs. $2(2) = 4$.

Time = 0.05 (sec) , antiderivative size = 8, normalized size of antiderivative = 4.00

method	result	size
default	$-\text{expIntegral}_1(-t)$	8
risch	$-\text{expIntegral}_1(-t)$	8
meijerg	$\ln(t) + i\pi - \ln(-t) - \text{expIntegral}_1(-t)$	21

input `int(exp(t)/t,t,method=_RETURNVERBOSE)`

output `-Ei(1,-t)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{e^t}{t} dt = \text{Ei}(t)$$

input `integrate(exp(t)/t,t, algorithm="fricas")`

output `Ei(t)`

Sympy [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{e^t}{t} dt = \text{Ei}(t)$$

input `integrate(exp(t)/t,t)`

output `Ei(t)`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{e^t}{t} dt = \text{Ei}(t)$$

input `integrate(exp(t)/t,t, algorithm="maxima")`

output `Ei(t)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{e^t}{t} dt = \text{Ei}(t)$$

input `integrate(exp(t)/t,t, algorithm="giac")`

output `Ei(t)`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{e^t}{t} dt = \text{ei}(t)$$

input `int(exp(t)/t,t)`

output `ei(t)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{e^t}{t} dt = ei(t)$$

input `int(exp(t)/t,t)`

output `ei(t)`

3.159 $\int \frac{e^{at}}{t} dt$

Optimal result	911
Mathematica [A] (verified)	911
Rubi [A] (verified)	912
Maple [A] (verified)	912
Fricas [A] (verification not implemented)	913
Sympy [A] (verification not implemented)	913
Maxima [A] (verification not implemented)	914
Giac [A] (verification not implemented)	914
Mupad [B] (verification not implemented)	914
Reduce [B] (verification not implemented)	915

Optimal result

Integrand size = 9, antiderivative size = 4

$$\int \frac{e^{at}}{t} dt = \text{ExpIntegralEi}(at)$$

output `Ei(a*t)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{e^{at}}{t} dt = \text{ExpIntegralEi}(at)$$

input `Integrate[E^(a*t)/t,t]`

output `ExpIntegralEi[a*t]`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {2609}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^{at}}{t} dt$$

↓ 2609

$$\text{ExpIntegralEi}(at)$$

input `Int [E^(a*t)/t,t]`

output `ExpIntegralEi[a*t]`

Defintions of rubi rules used

rule 2609 `Int[(F_)^((g_.)*(e_.) + (f_.)*(x_)))/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 2.25

method	result	size
derivativedivides	$-\text{expIntegral}_1(-at)$	9
default	$-\text{expIntegral}_1(-at)$	9
risch	$-\text{expIntegral}_1(-at)$	9
meijerg	$\ln(t) + \ln(-a) - \ln(-at) - \text{expIntegral}_1(-at)$	23

input `int(exp(a*t)/t,t,method=_RETURNVERBOSE)`

output `-Ei(1,-a*t)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{e^{at}}{t} dt = \text{Ei}(at)$$

input `integrate(exp(a*t)/t,t, algorithm="fricas")`

output `Ei(a*t)`

Sympy [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int \frac{e^{at}}{t} dt = \text{Ei}(at)$$

input `integrate(exp(a*t)/t,t)`

output `Ei(a*t)`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{e^{at}}{t} dt = \text{Ei}(at)$$

input `integrate(exp(a*t)/t,t, algorithm="maxima")`

output `Ei(a*t)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{e^{at}}{t} dt = \text{Ei}(at)$$

input `integrate(exp(a*t)/t,t, algorithm="giac")`

output `Ei(a*t)`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{e^{at}}{t} dt = \text{ei}(a t)$$

input `int(exp(a*t)/t,t)`

output `ei(a*t)`

Reduce [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{e^{at}}{t} dt = ei(at)$$

input `int(exp(a*t)/t,t)`

output `ei(a*t)`

3.160 $\int \frac{e^t}{t^2} dt$

Optimal result	916
Mathematica [A] (verified)	916
Rubi [A] (verified)	917
Maple [A] (verified)	918
Fricas [A] (verification not implemented)	918
Sympy [A] (verification not implemented)	918
Maxima [A] (verification not implemented)	919
Giac [A] (verification not implemented)	919
Mupad [B] (verification not implemented)	919
Reduce [B] (verification not implemented)	920

Optimal result

Integrand size = 7, antiderivative size = 11

$$\int \frac{e^t}{t^2} dt = -\frac{e^t}{t} + \text{ExpIntegralEi}(t)$$

output

```
-exp(t)/t+Ei(t)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{e^t}{t^2} dt = -\frac{e^t}{t} + \text{ExpIntegralEi}(t)$$

input

```
Integrate[E^t/t^2,t]
```

output

```
-(E^t/t) + ExpIntegralEi[t]
```

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2608, 2609}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^t}{t^2} dt$$

↓ 2608

$$\int \frac{e^t}{t} dt - \frac{e^t}{t}$$

↓ 2609

$$\text{ExpIntegralEi}(t) - \frac{e^t}{t}$$

input

```
Int [E^t/t^2,t]
```

output

```
-(E^t/t) + ExpIntegralEi [t]
```

Defintions of rubi rules used

rule 2608

```
Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m_)), x_Symbol] := Simp[(c + d*x)^(m + 1)*((b*F^(g*(e + f*x)))^n/(d*(m + 1))), x] - Simp[f*g*n*(Log[F]/(d*(m + 1))) Int[(c + d*x)^(m + 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && LtQ[m, -1] && IntegerQ[2*m] && !TrueQ[$UseGamma]
```

rule 2609

```
Int[(F_)^((g_.)*((e_.) + (f_.)*(x_)))/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.45

method	result	size
default	$-\frac{e^t}{t} - \text{expIntegral}_1(-t)$	16
risch	$-\frac{e^t}{t} - \text{expIntegral}_1(-t)$	16
meijerg	$-\frac{1}{t} - 1 + \ln(t) + i\pi + \frac{2t+2}{2t} - \frac{e^t}{t} - \ln(-t) - \text{expIntegral}_1(-t)$	44

input `int(exp(t)/t^2,t,method=_RETURNVERBOSE)`output `-exp(t)/t-Ei(1,-t)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.18

$$\int \frac{e^t}{t^2} dt = \frac{t\text{Ei}(t) - e^t}{t}$$

input `integrate(exp(t)/t^2,t, algorithm="fricas")`output `(t*Ei(t) - e^t)/t`**Sympy [A] (verification not implemented)**

Time = 0.46 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.64

$$\int \frac{e^t}{t^2} dt = \text{Ei}(t) - \frac{e^t}{t}$$

input `integrate(exp(t)/t**2,t)`output `Ei(t) - exp(t)/t`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.45

$$\int \frac{e^t}{t^2} dt = \Gamma(-1, -t)$$

input `integrate(exp(t)/t^2,t, algorithm="maxima")`

output `gamma(-1, -t)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.18

$$\int \frac{e^t}{t^2} dt = \frac{t\text{Ei}(t) - e^t}{t}$$

input `integrate(exp(t)/t^2,t, algorithm="giac")`

output `(t*Ei(t) - e^t)/t`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.27

$$\int \frac{e^t}{t^2} dt = -\frac{e^t}{t} - \text{expint}(-t)$$

input `int(exp(t)/t^2,t)`

output `- exp(t)/t - expint(-t)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.27

$$\int \frac{e^t}{t^2} dt = \frac{ei(t)t - e^t}{t}$$

input `int(exp(t)/t^2,t)`

output `(ei(t)*t - e**t)/t`

3.161 $\int e^{\frac{1}{t}} dt$

Optimal result	921
Mathematica [A] (verified)	921
Rubi [A] (verified)	922
Maple [A] (verified)	923
Fricas [A] (verification not implemented)	923
Sympy [A] (verification not implemented)	923
Maxima [A] (verification not implemented)	924
Giac [A] (verification not implemented)	924
Mupad [B] (verification not implemented)	924
Reduce [B] (verification not implemented)	925

Optimal result

Integrand size = 5, antiderivative size = 14

$$\int e^{\frac{1}{t}} dt = e^{\frac{1}{t}} t - \text{ExpIntegralEi} \left(\frac{1}{t} \right)$$

output

```
exp(1/t)*t-Ei(1/t)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int e^{\frac{1}{t}} dt = e^{\frac{1}{t}} t - \text{ExpIntegralEi} \left(\frac{1}{t} \right)$$

input

```
Integrate[E^t^(-1),t]
```

output

```
E^t^(-1)*t - ExpIntegralEi[t^(-1)]
```

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {2635, 2639}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{\frac{1}{t}} dt$$

$$\downarrow 2635$$

$$\int \frac{e^{\frac{1}{t}}}{t} dt + e^{\frac{1}{t}} t$$

$$\downarrow 2639$$

$$e^{\frac{1}{t}} t - \text{ExpIntegralEi}\left(\frac{1}{t}\right)$$

input `Int[E^t^(-1),t]`

output `E^t^(-1)*t - ExpIntegralEi[t^(-1)]`

Defintions of rubi rules used

rule 2635 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^n)), x_Symbol] :> Simp[(c + d*x)*(F^(a + b*(c + d*x)^n)/d), x] - Simp[b*n*Log[F] Int[(c + d*x)^n*F^(a + b*(c + d*x)^n), x], x] /; FreeQ[{F, a, b, c, d}, x] && IntegerQ[2/n] && ILtQ[n, 0]`

rule 2639 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^n))/((e_.) + (f_.)*(x_)), x_Symbol] :> Simp[F^a*(ExpIntegralEi[b*(c + d*x)^n*Log[F]]/(f*n)), x] /; FreeQ[{F, a, b, c, d, e, f, n}, x] && EqQ[d*e - c*f, 0]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

method	result	size
derivativedivides	$e^{\frac{1}{t}}t + \text{expIntegral}_1\left(-\frac{1}{t}\right)$	15
default	$e^{\frac{1}{t}}t + \text{expIntegral}_1\left(-\frac{1}{t}\right)$	15
risch	$e^{\frac{1}{t}}t + \text{expIntegral}_1\left(-\frac{1}{t}\right)$	15
meijerg	$t + 1 + \ln(t) - i\pi - \frac{t(2+\frac{2}{t})}{2} + e^{\frac{1}{t}}t + \ln\left(-\frac{1}{t}\right) + \text{expIntegral}_1\left(-\frac{1}{t}\right)$	39

input `int(exp(1/t),t,method=_RETURNVERBOSE)`output `exp(1/t)*t+Ei(1,-1/t)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

$$\int e^{\frac{1}{t}} dt = te^{\frac{1}{t}} - \text{Ei}\left(\frac{1}{t}\right)$$

input `integrate(exp(1/t),t, algorithm="fricas")`output `t*e^(1/t) - Ei(1/t)`**Sympy [A] (verification not implemented)**

Time = 0.49 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int e^{\frac{1}{t}} dt = te^{\frac{1}{t}} - \text{Ei}\left(\frac{1}{t}\right)$$

input `integrate(exp(1/t),t)`

output `t*exp(1/t) - Ei(1/t)`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.64

$$\int e^{\frac{1}{t}} dt = -\Gamma\left(-1, -\frac{1}{t}\right)$$

input `integrate(exp(1/t),t, algorithm="maxima")`

output `-gamma(-1, -1/t)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.29

$$\int e^{\frac{1}{t}} dt = -t \left(\frac{\text{Ei}\left(\frac{1}{t}\right)}{t} - e^{\frac{1}{t}} \right)$$

input `integrate(exp(1/t),t, algorithm="giac")`

output `-t*(Ei(1/t)/t - e^(1/t))`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.64

$$\int e^{\frac{1}{t}} dt = t \text{expint}\left(2, -\frac{1}{t}\right)$$

input `int(exp(1/t),t)`

output `t*expint(2, -1/t)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int e^{\frac{1}{t}} dt = -ei\left(\frac{1}{t}\right) + e^{\frac{1}{t}}t$$

input `int(exp(1/t),t)`

output `- ei(1/t) + e**(1/t)*t`

3.162 $\int \frac{e^{-t}}{-1-a+t} dt$

Optimal result	926
Mathematica [A] (verified)	926
Rubi [A] (verified)	927
Maple [A] (verified)	927
Fricas [A] (verification not implemented)	928
Sympy [F]	928
Maxima [A] (verification not implemented)	928
Giac [A] (verification not implemented)	929
Mupad [B] (verification not implemented)	929
Reduce [F]	929

Optimal result

Integrand size = 14, antiderivative size = 15

$$\int \frac{e^{-t}}{-1-a+t} dt = e^{-1-a} \text{ExpIntegralEi}(1+a-t)$$

output

```
exp(-1-a)*Ei(1+a-t)
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{e^{-t}}{-1-a+t} dt = e^{-1-a} \text{ExpIntegralEi}(1+a-t)$$

input

```
Integrate[1/(E^-t*(-1 - a + t)),t]
```

output

```
E^(-1 - a)*ExpIntegralEi[1 + a - t]
```

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$, Rules used = {2609}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^{-t}}{-a+t-1} dt$$

↓ 2609

$$e^{-a-1} \text{ExpIntegralEi}(a-t+1)$$

input `Int[1/(E^t*(-1 - a + t)),t]`

output `E^(-1 - a)*ExpIntegralEi[1 + a - t]`

Defintions of rubi rules used

rule 2609 `Int[(F_)^((g_.)*((e_.) + (f_.)*(x_)))/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

method	result	size
default	$-e^{-1-a} \text{expIntegral}_1(-1-a+t)$	17
risch	$-e^{-1-a} \text{expIntegral}_1(-1-a+t)$	17

input `int(1/exp(t)/(-1-a+t),t,method=_RETURNVERBOSE)`

output `-exp(-1-a)*Ei(1,-1-a+t)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{e^{-t}}{-1-a+t} dt = \text{Ei}(a-t+1) e^{(-a-1)}$$

input `integrate(1/exp(t)/(-1-a+t),t, algorithm="fricas")`

output `Ei(a - t + 1)*e^(-a - 1)`

Sympy [F]

$$\int \frac{e^{-t}}{-1-a+t} dt = \int \frac{e^{-t}}{-a+t-1} dt$$

input `integrate(1/exp(t)/(-1-a+t),t)`

output `Integral(exp(-t)/(-a + t - 1), t)`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.07

$$\int \frac{e^{-t}}{-1-a+t} dt = -e^{(-a-1)} E_1(-a+t-1)$$

input `integrate(1/exp(t)/(-1-a+t),t, algorithm="maxima")`

output `-e^(-a - 1)*exp_integral_e(1, -a + t - 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{e^{-t}}{-1-a+t} dt = \text{Ei}(a-t+1) e^{(-a-1)}$$

input `integrate(1/exp(t)/(-1-a+t),t, algorithm="giac")`output `Ei(a - t + 1)*e^(-a - 1)`**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{e^{-t}}{-1-a+t} dt = e^{-a-1} \text{ei}(a-t+1)$$

input `int(-exp(-t)/(a - t + 1),t)`output `exp(- a - 1)*ei(a - t + 1)`**Reduce [F]**

$$\int \frac{e^{-t}}{-1-a+t} dt = - \left(\int \frac{1}{e^t a - e^t t + e^t} dt \right)$$

input `int(1/exp(t)/(-1-a+t),t)`output `- int(1/(e**t*a - e**t*t + e**t),t)`

3.163 $\int \frac{e^{t^2} t}{1+t^2} dt$

Optimal result	930
Mathematica [A] (verified)	930
Rubi [A] (verified)	931
Maple [A] (verified)	932
Fricas [A] (verification not implemented)	932
Sympy [F]	932
Maxima [A] (verification not implemented)	933
Giac [A] (verification not implemented)	933
Mupad [B] (verification not implemented)	933
Reduce [F]	934

Optimal result

Integrand size = 14, antiderivative size = 13

$$\int \frac{e^{t^2} t}{1+t^2} dt = \frac{\text{ExpIntegralEi}(1+t^2)}{2e}$$

output `1/2*Ei(t^2+1)/exp(1)`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{e^{t^2} t}{1+t^2} dt = \frac{\text{ExpIntegralEi}(1+t^2)}{2e}$$

input `Integrate[(E^t^2*t)/(1+t^2),t]`

output `ExpIntegralEi[1+t^2]/(2*E)`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {7266, 2609}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^{t^2} t}{t^2 + 1} dt$$

↓ 7266

$$\frac{1}{2} \int \frac{e^{t^2}}{t^2 + 1} dt^2$$

↓ 2609

$$\frac{\text{ExpIntegralEi}(t^2 + 1)}{2e}$$

input `Int[(E^t^2*t)/(1 + t^2),t]`

output `ExpIntegralEi[1 + t^2]/(2*E)`

Defintions of rubi rules used

rule 2609 `Int[(F_)^((g_)*((e_.) + (f_.)*(x_)))/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]`

rule 7266 `Int[(u_)*(x_)^(m_), x_Symbol] := Simp[1/(m + 1) Subst[Int[SubstFor[x^(m + 1), u, x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && FunctionOfQ[x^(m + 1), u, x]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.08

method	result	size
derivativdivides	$-\frac{e^{-1} \operatorname{expIntegral}_1(-t^2-1)}{2}$	14
default	$-\frac{e^{-1} \operatorname{expIntegral}_1(-t^2-1)}{2}$	14
risch	$-\frac{e^{-1} \operatorname{expIntegral}_1(-t^2-1)}{2}$	14

input `int(exp(t^2)*t/(t^2+1),t,method=_RETURNVERBOSE)`

output `-1/2*exp(-1)*Ei(1,-t^2-1)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \frac{e^{t^2} t}{1+t^2} dt = \frac{1}{2} \operatorname{Ei}(t^2+1) e^{(-1)}$$

input `integrate(exp(t^2)*t/(t^2+1),t, algorithm="fricas")`

output `1/2*Ei(t^2 + 1)*e^(-1)`

Sympy [F]

$$\int \frac{e^{t^2} t}{1+t^2} dt = \int \frac{te^{t^2}}{t^2+1} dt$$

input `integrate(exp(t**2)*t/(t**2+1),t)`

output `Integral(t*exp(t**2)/(t**2 + 1), t)`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{e^{t^2} t}{1+t^2} dt = -\frac{1}{2} e^{(-1)} E_1(-t^2 - 1)$$

input `integrate(exp(t^2)*t/(t^2+1),t, algorithm="maxima")`output `-1/2*e^(-1)*exp_integral_e(1, -t^2 - 1)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \frac{e^{t^2} t}{1+t^2} dt = \frac{1}{2} \text{Ei}(t^2 + 1) e^{(-1)}$$

input `integrate(exp(t^2)*t/(t^2+1),t, algorithm="giac")`output `1/2*Ei(t^2 + 1)*e^(-1)`**Mupad [B] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \frac{e^{t^2} t}{1+t^2} dt = \frac{e^{-1} \text{ei}(t^2 + 1)}{2}$$

input `int((t*exp(t^2))/(t^2 + 1),t)`output `(exp(-1)*ei(t^2 + 1))/2`

Reduce [F]

$$\int \frac{e^{t^2}t}{1+t^2} dt = \int \frac{e^{t^2}t}{t^2+1} dt$$

input `int(exp(t^2)*t/(t^2+1),t)`

output `int((e**(t**2)*t)/(t**2 + 1),t)`

3.164 $\int \frac{e^t}{(1+t)^2} dt$

Optimal result	935
Mathematica [A] (verified)	935
Rubi [A] (verified)	936
Maple [A] (verified)	937
Fricas [A] (verification not implemented)	937
Sympy [F]	937
Maxima [A] (verification not implemented)	938
Giac [B] (verification not implemented)	938
Mupad [B] (verification not implemented)	939
Reduce [F]	939

Optimal result

Integrand size = 9, antiderivative size = 19

$$\int \frac{e^t}{(1+t)^2} dt = -\frac{e^t}{1+t} + \frac{\text{ExpIntegralEi}(1+t)}{e}$$

output `-exp(t)/(1+t)+Ei(1+t)/exp(1)`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{e^t}{(1+t)^2} dt = -\frac{e^t}{1+t} + \frac{\text{ExpIntegralEi}(1+t)}{e}$$

input `Integrate[E^t/(1 + t)^2,t]`

output `-(E^t/(1 + t)) + ExpIntegralEi[1 + t]/E`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {2608, 2609}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{e^t}{(t+1)^2} dt \\ \downarrow 2608 \\ \int \frac{e^t}{t+1} dt - \frac{e^t}{t+1} \\ \downarrow 2609 \\ \frac{\text{ExpIntegralEi}(t+1)}{e} - \frac{e^t}{t+1} \end{array}$$

input `Int[E^t/(1 + t)^2,t]`

output `-(E^t/(1 + t)) + ExpIntegralEi[1 + t]/E`

Defintions of rubi rules used

rule 2608 `Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m_), x_Symbol] := Simp[(c + d*x)^(m + 1)*((b*F^(g*(e + f*x)))^n/(d*(m + 1))), x] - Simp[f*g*n*(Log[F]/(d*(m + 1))) Int[(c + d*x)^(m + 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && LtQ[m, -1] && IntegerQ[2*m] && !TrueQ[$UseGamma]`

rule 2609 `Int[(F_)^((g_.)*((e_.) + (f_.)*(x_)))/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.16

method	result	size
default	$-\frac{e^t}{1+t} - e^{-1} \operatorname{expIntegral}_1(-1-t)$	22
risch	$-\frac{e^t}{1+t} - e^{-1} \operatorname{expIntegral}_1(-1-t)$	22

input `int(exp(t)/(1+t)^2,t,method=_RETURNVERBOSE)`output `-exp(t)/(1+t)-exp(-1)*Ei(1,-1-t)`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.21

$$\int \frac{e^t}{(1+t)^2} dt = \frac{((t+1)\operatorname{Ei}(t+1) - e^{(t+1)})e^{(-1)}}{t+1}$$

input `integrate(exp(t)/(1+t)^2,t, algorithm="fricas")`output `((t + 1)*Ei(t + 1) - e^(t + 1))*e^(-1)/(t + 1)`**Sympy [F]**

$$\int \frac{e^t}{(1+t)^2} dt = \int \frac{e^t}{(t+1)^2} dt$$

input `integrate(exp(t)/(1+t)**2,t)`output `Integral(exp(t)/(t + 1)**2, t)`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

$$\int \frac{e^t}{(1+t)^2} dt = -\frac{e^{(-1)}E_2(-t-1)}{t+1}$$

input `integrate(exp(t)/(1+t)^2,t, algorithm="maxima")`

output `-e^(-1)*exp_integral_e(2, -t - 1)/(t + 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 80 vs. $2(17) = 34$.

Time = 0.12 (sec) , antiderivative size = 80, normalized size of antiderivative = 4.21

$$\int \frac{e^t}{(1+t)^2} dt = \frac{(t+1)\left(\frac{1}{t+1}-1\right)\text{Ei}\left(-\left(t+1\right)\left(\frac{1}{t+1}-1\right)+1\right) - \text{Ei}\left(-\left(t+1\right)\left(\frac{1}{t+1}-1\right)+1\right) + e^{\left(-\left(t+1\right)\left(\frac{1}{t+1}-1\right)+1\right)}}{(t+1)\left(\frac{1}{t+1}-1\right)e - e}$$

input `integrate(exp(t)/(1+t)^2,t, algorithm="giac")`

output `((t + 1)*(1/(t + 1) - 1)*Ei(-(t + 1)*(1/(t + 1) - 1) + 1) - Ei(-(t + 1)*(1/(t + 1) - 1) + 1) + e^(-(t + 1)*(1/(t + 1) - 1) + 1))/((t + 1)*(1/(t + 1) - 1)*e - e)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{e^t}{(1+t)^2} dt = \text{ei}(t+1) e^{-1} - \frac{e^t}{t+1}$$

input `int(exp(t)/(t + 1)^2,t)`

output `ei(t + 1)*exp(-1) - exp(t)/(t + 1)`

Reduce [F]

$$\int \frac{e^t}{(1+t)^2} dt = \int \frac{e^t}{t^2 + 2t + 1} dt$$

input `int(exp(t)/(1+t)^2,t)`

output `int(e**t/(t**2 + 2*t + 1),t)`

3.165 $\int e^t \log(1 + t) dt$

Optimal result	940
Mathematica [A] (verified)	940
Rubi [A] (verified)	941
Maple [A] (verified)	942
Fricas [A] (verification not implemented)	942
Sympy [F]	942
Maxima [A] (verification not implemented)	943
Giac [A] (verification not implemented)	943
Mupad [F(-1)]	943
Reduce [F]	944

Optimal result

Integrand size = 8, antiderivative size = 18

$$\int e^t \log(1 + t) dt = -\frac{\text{ExpIntegralEi}(1 + t)}{e} + e^t \log(1 + t)$$

output `-Ei(1+t)/exp(1)+exp(t)*ln(1+t)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int e^t \log(1 + t) dt = -\frac{\text{ExpIntegralEi}(1 + t)}{e} + e^t \log(1 + t)$$

input `Integrate[E^t*Log[1 + t],t]`

output `-(ExpIntegralEi[1 + t]/E) + E^t*Log[1 + t]`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3034, 2609}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^t \log(t+1) dt$$

$$\downarrow \text{3034}$$

$$e^t \log(t+1) - \int \frac{e^t}{t+1} dt$$

$$\downarrow \text{2609}$$

$$e^t \log(t+1) - \frac{\text{ExpIntegralEi}(t+1)}{e}$$

input `Int[E^t*Log[1 + t],t]`

output `-(ExpIntegralEi[1 + t]/E) + E^t*Log[1 + t]`

Defintions of rubi rules used

rule 2609 `Int[(F_)^((g_.)*(e_.) + (f_.)*(x_)))/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]`

rule 3034 `Int[Log[u]*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Simp[Log[u] w, x] - Int[SimplifyIntegrand[w*(D[u, x]/u), x], x] /; InverseFunctionFreeQ[w, x]] /; InverseFunctionFreeQ[u, x]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.06

method	result	size
risch	$e^t \ln(1+t) + e^{-1} \operatorname{expIntegral}_1(-1-t)$	19

input `int(exp(t)*ln(1+t),t,method=_RETURNVERBOSE)`

output `exp(t)*ln(1+t)+exp(-1)*Ei(1,-1-t)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.06

$$\int e^t \log(1+t) dt = (e^{(t+1)} \log(t+1) - \operatorname{Ei}(t+1))e^{(-1)}$$

input `integrate(exp(t)*log(1+t),t, algorithm="fricas")`

output `(e^(t + 1)*log(t + 1) - Ei(t + 1))*e^(-1)`

Sympy [F]

$$\int e^t \log(1+t) dt = \int e^t \log(t+1) dt$$

input `integrate(exp(t)*ln(1+t),t)`

output `Integral(exp(t)*log(t + 1), t)`

Maxima [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int e^t \log(1+t) dt = e^{(-1)} E_1(-t-1) + e^t \log(t+1)$$

input `integrate(exp(t)*log(1+t),t, algorithm="maxima")`output `e^(-1)*exp_integral_e(1, -t - 1) + e^t*log(t + 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int e^t \log(1+t) dt = -\text{Ei}(t+1) e^{(-1)} + e^t \log(t+1)$$

input `integrate(exp(t)*log(1+t),t, algorithm="giac")`output `-Ei(t + 1)*e^(-1) + e^t*log(t + 1)`**Mupad [F(-1)]**

Timed out.

$$\int e^t \log(1+t) dt = \int \ln(t+1) e^t dt$$

input `int(log(t + 1)*exp(t),t)`output `int(log(t + 1)*exp(t), t)`

Reduce [F]

$$\int e^t \log(1+t) dt = e^t \log(t+1) - \left(\int \frac{e^t}{t+1} dt \right)$$

input `int(exp(t)*log(1+t),t)`

output `e**t*log(t + 1) - int(e**t/(t + 1),t)`

3.166 $\int e^{-t}t dt$

Optimal result	945
Mathematica [A] (verified)	945
Rubi [A] (verified)	946
Maple [A] (warning: unable to verify)	947
Fricas [A] (verification not implemented)	947
Sympy [A] (verification not implemented)	948
Maxima [A] (verification not implemented)	948
Giac [A] (verification not implemented)	948
Mupad [B] (verification not implemented)	949
Reduce [B] (verification not implemented)	949

Optimal result

Integrand size = 7, antiderivative size = 16

$$\int e^{-t}t dt = -e^{-t} - e^{-t}t$$

output `-1/exp(t)-t/exp(t)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.69

$$\int e^{-t}t dt = e^{-t}(-1 - t)$$

input `Integrate[t/E^t,t]`

output `(-1 - t)/E^t`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{-t} dt$$

$$\downarrow 2607$$

$$\int e^{-t} dt - e^{-t}$$

$$\downarrow 2624$$

$$-e^{-t} - e^{-t}$$

input `Int[t/E^t,t]`

output `-E^(-t) - t/E^t`

Defintions of rubi rules used

rule 2607 `Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]`

rule 2624 `Int[((F_)^(v_))^(n_.), x_Symbol] := Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /; FreeQ[{F, n}, x] && LinearQ[v, x]`

Maple [A] (warning: unable to verify)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.62

method	result	size
gospers	$-(1+t)e^{-t}$	10
orering	$-(1+t)e^{-t}$	10
norman	$(-1-t)e^{-t}$	11
risch	$(-1-t)e^{-t}$	11
parallelrisch	$(-1-t)e^{-t}$	11
meijerg	$1 - \frac{(2t+2)e^{-t}}{2}$	14
default	$-e^{-t} - te^{-t}$	15

input `int(t/exp(t),t,method=_RETURNVERBOSE)`

output `-(1+t)/exp(t)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.56

$$\int e^{-t}t dt = -(t+1)e^{(-t)}$$

input `integrate(t/exp(t),t, algorithm="fricas")`

output `-(t + 1)*e^(-t)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.44

$$\int e^{-t}t dt = (-t - 1)e^{-t}$$

input `integrate(t/exp(t),t)`

output `(-t - 1)*exp(-t)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.56

$$\int e^{-t}t dt = -(t + 1)e^{(-t)}$$

input `integrate(t/exp(t),t, algorithm="maxima")`

output `-(t + 1)*e^(-t)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.56

$$\int e^{-t}t dt = -(t + 1)e^{(-t)}$$

input `integrate(t/exp(t),t, algorithm="giac")`

output `-(t + 1)*e^(-t)`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.56

$$\int e^{-t} dt = -e^{-t} (t + 1)$$

input `int(t*exp(-t),t)`

output `-exp(-t)*(t + 1)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.69

$$\int e^{-t} t dt = \frac{-t - 1}{e^t}$$

input `int(t/exp(t),t)`

output `(- (t + 1))/e**t`

3.167 $\int e^{-t}t^2 dt$

Optimal result	950
Mathematica [A] (verified)	950
Rubi [A] (verified)	951
Maple [A] (warning: unable to verify)	952
Fricas [A] (verification not implemented)	952
Sympy [A] (verification not implemented)	953
Maxima [A] (verification not implemented)	953
Giac [A] (verification not implemented)	953
Mupad [B] (verification not implemented)	954
Reduce [B] (verification not implemented)	954

Optimal result

Integrand size = 9, antiderivative size = 26

$$\int e^{-t}t^2 dt = -2e^{-t} - 2e^{-t}t - e^{-t}t^2$$

output `-2/exp(t)-2*t/exp(t)-t^2/exp(t)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.62

$$\int e^{-t}t^2 dt = e^{-t}(-2 - 2t - t^2)$$

input `Integrate[t^2/E^t,t]`

output `(-2 - 2*t - t^2)/E^t`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.12, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2607, 2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int e^{-t^2} dt \\ & \quad \downarrow 2607 \\ & 2 \int e^{-t} t dt - e^{-t^2} \\ & \quad \downarrow 2607 \\ & 2 \left(\int e^{-t} dt - e^{-t} t \right) - e^{-t^2} \\ & \quad \downarrow 2624 \\ & 2(-e^{-t} - e^{-t} t) - e^{-t^2} \end{aligned}$$

input `Int[t^2/E^t,t]`

output `-(t^2/E^t) + 2*(-E^(-t) - t/E^t)`

Defintions of rubi rules used

rule 2607

```
Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m
_.), x_Symbol] := Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])),
x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^
n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*
m] && !TrueQ[$UseGamma]
```

rule 2624

```
Int[((F_)^(v_))^(n_.), x_Symbol] := Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;
FreeQ[{F, n}, x] && LinearQ[v, x]
```


Maple [A] (warning: unable to verify)

Time = 0.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.58

method	result	size
gospers	$-(t^2 + 2t + 2)e^{-t}$	15
orering	$-(t^2 + 2t + 2)e^{-t}$	15
norman	$(-t^2 - 2t - 2)e^{-t}$	16
risch	$(-t^2 - 2t - 2)e^{-t}$	16
parallelrisc	$(-t^2 - 2t - 2)e^{-t}$	16
meijerg	$2 - \frac{(3t^2 + 6t + 6)e^{-t}}{3}$	19
default	$-2e^{-t} - 2te^{-t} - t^2e^{-t}$	24

input `int(t^2/exp(t),t,method=_RETURNVERBOSE)`output `-(t^2+2*t+2)/exp(t)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.54

$$\int e^{-t}t^2 dt = -(t^2 + 2t + 2)e^{(-t)}$$

input `integrate(t^2/exp(t),t, algorithm="fricas")`output `-(t^2 + 2*t + 2)*e^(-t)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.46

$$\int e^{-t^2} dt = (-t^2 - 2t - 2) e^{-t}$$

input `integrate(t**2/exp(t),t)`

output `(-t**2 - 2*t - 2)*exp(-t)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.54

$$\int e^{-t^2} dt = -(t^2 + 2t + 2)e^{(-t)}$$

input `integrate(t^2/exp(t),t, algorithm="maxima")`

output `-(t^2 + 2*t + 2)*e^(-t)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.54

$$\int e^{-t^2} dt = -(t^2 + 2t + 2)e^{(-t)}$$

input `integrate(t^2/exp(t),t, algorithm="giac")`

output `-(t^2 + 2*t + 2)*e^(-t)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.54

$$\int e^{-t} t^2 dt = -e^{-t} (t^2 + 2t + 2)$$

input `int(t^2*exp(-t),t)`

output `-exp(-t)*(2*t + t^2 + 2)`

Reduce [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.62

$$\int e^{-t} t^2 dt = \frac{-t^2 - 2t - 2}{e^t}$$

input `int(t^2/exp(t),t)`

output `(- t**2 - 2*t - 2)/e**t`

3.168 $\int e^{-t}t^3 dt$

Optimal result	955
Mathematica [A] (verified)	955
Rubi [A] (verified)	956
Maple [A] (warning: unable to verify)	957
Fricas [A] (verification not implemented)	958
Sympy [A] (verification not implemented)	958
Maxima [A] (verification not implemented)	958
Giac [A] (verification not implemented)	959
Mupad [B] (verification not implemented)	959
Reduce [B] (verification not implemented)	959

Optimal result

Integrand size = 9, antiderivative size = 36

$$\int e^{-t}t^3 dt = -6e^{-t} - 6e^{-t}t - 3e^{-t}t^2 - e^{-t}t^3$$

output `-6/exp(t)-6*t/exp(t)-3*t^2/exp(t)-t^3/exp(t)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.58

$$\int e^{-t}t^3 dt = e^{-t}(-6 - 6t - 3t^2 - t^3)$$

input `Integrate[t^3/E^t,t]`

output `(-6 - 6*t - 3*t^2 - t^3)/E^t`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.17, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {2607, 2607, 2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{-t} t^3 dt \\
 & \quad \downarrow \text{2607} \\
 & 3 \int e^{-t} t^2 dt - e^{-t} t^3 \\
 & \quad \downarrow \text{2607} \\
 & 3 \left(2 \int e^{-t} t dt - e^{-t} t^2 \right) - e^{-t} t^3 \\
 & \quad \downarrow \text{2607} \\
 & 3 \left(2 \left(\int e^{-t} dt - e^{-t} t \right) - e^{-t} t^2 \right) - e^{-t} t^3 \\
 & \quad \downarrow \text{2624} \\
 & 3 \left(2 \left(-e^{-t} t - e^{-t} \right) - e^{-t} t^2 \right) - e^{-t} t^3
 \end{aligned}$$

input

Int[t^3/E^t,t]

output

-(t^3/E^t) + 3*(-(t^2/E^t) + 2*(-E^(-t) - t/E^t))

Definitions of rubi rules used

rule 2607

```
Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m
_.), x_Symbol] := Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])),
x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^
n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*
m] && !TrueQ[$UseGamma]
```

rule 2624

```
Int[((F_)^(v_))^(n_.), x_Symbol] := Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;
FreeQ[{F, n}, x] && LinearQ[v, x]
```

Maple [A] (warning: unable to verify)

Time = 0.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.56

method	result	size
gosper	$-(t^3 + 3t^2 + 6t + 6)e^{-t}$	20
oring	$-(t^3 + 3t^2 + 6t + 6)e^{-t}$	20
norman	$(-t^3 - 3t^2 - 6t - 6)e^{-t}$	21
risch	$(-t^3 - 3t^2 - 6t - 6)e^{-t}$	21
parallelrisch	$(-t^3 - 3t^2 - 6t - 6)e^{-t}$	21
meijerg	$6 - \frac{(4t^3 + 12t^2 + 24t + 24)e^{-t}}{4}$	24
default	$-6e^{-t} - 6te^{-t} - 3t^2e^{-t} - t^3e^{-t}$	33

input

```
int(t^3/exp(t), t, method=_RETURNVERBOSE)
```

output

```
-(t^3+3*t^2+6*t+6)/exp(t)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.53

$$\int e^{-t} t^3 dt = -(t^3 + 3t^2 + 6t + 6)e^{-t}$$

input `integrate(t^3/exp(t),t, algorithm="fricas")`

output `-(t^3 + 3*t^2 + 6*t + 6)*e^(-t)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.47

$$\int e^{-t} t^3 dt = (-t^3 - 3t^2 - 6t - 6)e^{-t}$$

input `integrate(t**3/exp(t),t)`

output `(-t**3 - 3*t**2 - 6*t - 6)*exp(-t)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.53

$$\int e^{-t} t^3 dt = -(t^3 + 3t^2 + 6t + 6)e^{-t}$$

input `integrate(t^3/exp(t),t, algorithm="maxima")`

output `-(t^3 + 3*t^2 + 6*t + 6)*e^(-t)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.53

$$\int e^{-t}t^3 dt = -(t^3 + 3t^2 + 6t + 6)e^{(-t)}$$

input `integrate(t^3/exp(t),t, algorithm="giac")`

output `-(t^3 + 3*t^2 + 6*t + 6)*e^(-t)`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.53

$$\int e^{-t}t^3 dt = -e^{-t} (t^3 + 3t^2 + 6t + 6)$$

input `int(t^3*exp(-t),t)`

output `-exp(-t)*(6*t + 3*t^2 + t^3 + 6)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.58

$$\int e^{-t}t^3 dt = \frac{-t^3 - 3t^2 - 6t - 6}{e^t}$$

input `int(t^3/exp(t),t)`

output `(- t**3 - 3*t**2 - 6*t - 6)/e**t`

3.169 $\int \frac{b1 \cos(x) + a1 \sin(x)}{b \cos(x) + a \sin(x)} dx$

Optimal result	960
Mathematica [A] (verified)	960
Rubi [A] (verified)	961
Maple [A] (verified)	962
Fricas [A] (verification not implemented)	962
Sympy [C] (verification not implemented)	963
Maxima [B] (verification not implemented)	964
Giac [A] (verification not implemented)	964
Mupad [B] (verification not implemented)	965
Reduce [B] (verification not implemented)	965

Optimal result

Integrand size = 21, antiderivative size = 48

$$\int \frac{b1 \cos(x) + a1 \sin(x)}{b \cos(x) + a \sin(x)} dx = \frac{(aa1 + bb1)x}{a^2 + b^2} - \frac{(a1b - ab1) \log(b \cos(x) + a \sin(x))}{a^2 + b^2}$$

output `(a*a1+b*b1)*x/(a^2+b^2)-(-a*b1+a1*b)*ln(b*cos(x)+a*sin(x))/(a^2+b^2)`

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.81

$$\int \frac{b1 \cos(x) + a1 \sin(x)}{b \cos(x) + a \sin(x)} dx = \frac{(aa1 + bb1)x + (-a1b + ab1) \log(b \cos(x) + a \sin(x))}{a^2 + b^2}$$

input `Integrate[(b1*Cos[x] + a1*Sin[x])/(b*Cos[x] + a*Sin[x]),x]`

output `((a*a1 + b*b1)*x + (-a1*b) + a*b1)*Log[b*Cos[x] + a*Sin[x]]/(a^2 + b^2)`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {3042, 3612}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a_1 \sin(x) + b_1 \cos(x)}{a \sin(x) + b \cos(x)} dx$$

↓ 3042

$$\int \frac{a_1 \sin(x) + b_1 \cos(x)}{a \sin(x) + b \cos(x)} dx$$

↓ 3612

$$\frac{x(a a_1 + b b_1)}{a^2 + b^2} - \frac{(a_1 b - a b_1) \log(a \sin(x) + b \cos(x))}{a^2 + b^2}$$

input `Int[(b1*Cos[x] + a1*Sin[x])/(b*Cos[x] + a*Sin[x]),x]`

output `((a*a1 + b*b1)*x)/(a^2 + b^2) - ((a1*b - a*b1)*Log[b*Cos[x] + a*Sin[x]])/(a^2 + b^2)`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3612 `Int[((A_.) + cos[(d_.) + (e_.)*(x_.)]*(B_.) + (C_.)*sin[(d_.) + (e_.)*(x_.)]) / ((a_.) + cos[(d_.) + (e_.)*(x_.)]*(b_.) + (c_.)*sin[(d_.) + (e_.)*(x_.)]), x_Symbol] := Simp[(b*B + c*C)*(x/(b^2 + c^2)), x] + Simp[(c*B - b*C)*(Log[a + b*Cos[d + e*x] + c*Sin[d + e*x]]/(e*(b^2 + c^2))), x] /; FreeQ[{a, b, c, d, e, A, B, C}, x] && NeQ[b^2 + c^2, 0] && EqQ[A*(b^2 + c^2) - a*(b*B + c*C), 0]`

Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.25

method	result	si
parallelrisc	$\frac{xa a_1 + xb b_1 - \left(\ln\left(\sec\left(\frac{x}{2}\right)^2\right) - \ln\left(b \tan\left(\frac{x}{2}\right)^2 - 2a \tan\left(\frac{x}{2}\right) - b\right)\right)(a b_1 - a_1 b)}{a^2 + b^2}$	60
default	$\frac{(-a b_1 + a_1 b) \ln(1 + \tan(x)^2)}{2} + \frac{(a a_1 + b b_1) \arctan(\tan(x))}{a^2 + b^2} + \frac{(a b_1 - a_1 b) \ln(a \tan(x) + b)}{a^2 + b^2}$	60
norman	$\frac{\frac{(a a_1 + b b_1)x}{a^2 + b^2} + \frac{(a a_1 + b b_1)x \tan\left(\frac{x}{2}\right)^2}{a^2 + b^2}}{1 + \tan\left(\frac{x}{2}\right)^2} + \frac{(a b_1 - a_1 b) \ln\left(-b \tan\left(\frac{x}{2}\right)^2 + 2a \tan\left(\frac{x}{2}\right) + b\right)}{a^2 + b^2} - \frac{(a b_1 - a_1 b) \ln\left(1 + \tan\left(\frac{x}{2}\right)^2\right)}{a^2 + b^2}$	12
risc	$\frac{ix b_1}{ib+a} + \frac{x a_1}{ib+a} - \frac{2ix a b_1}{a^2+b^2} + \frac{2ix a_1 b}{a^2+b^2} + \frac{\ln\left(e^{2ix} + \frac{ib-a}{ib+a}\right) a b_1}{a^2+b^2} - \frac{\ln\left(e^{2ix} + \frac{ib-a}{ib+a}\right) a_1 b}{a^2+b^2}$	12

input `int((b1*cos(x)+a1*sin(x))/(b*cos(x)+a*sin(x)),x,method=_RETURNVERBOSE)`

output `1/(a^2+b^2)*(x*a*a1+x*b*b1-(ln(sec(1/2*x)^2)-ln(b*tan(1/2*x)^2-2*a*tan(1/2*x)-b))*(a*b1-a1*b))`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.25

$$\int \frac{b_1 \cos(x) + a_1 \sin(x)}{b \cos(x) + a \sin(x)} dx$$

$$= \frac{2(a a_1 + b b_1)x - (a_1 b - a b_1) \log(2 a b \cos(x) \sin(x) - (a^2 - b^2) \cos(x)^2 + a^2)}{2(a^2 + b^2)}$$

input `integrate((b1*cos(x)+a1*sin(x))/(b*cos(x)+a*sin(x)),x, algorithm="fricas")`

output `1/2*(2*(a*a1 + b*b1)*x - (a1*b - a*b1)*log(2*a*b*cos(x)*sin(x) - (a^2 - b^2)*cos(x)^2 + a^2))/(a^2 + b^2)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.34 (sec) , antiderivative size = 360, normalized size of antiderivative = 7.50

$$\int \frac{b_1 \cos(x) + a_1 \sin(x)}{b \cos(x) + a \sin(x)} dx$$

$$= \begin{cases} \tilde{\infty}(-a_1 \log(\cos(x)) + b_1 x) \\ \frac{a_1 x + b_1 \log(\sin(x))}{a} \\ \frac{a_1 x \sin(x)}{-2ib \sin(x) + 2b \cos(x)} + \frac{ia_1 x \cos(x)}{-2ib \sin(x) + 2b \cos(x)} - \frac{a_1 \cos(x)}{-2ib \sin(x) + 2b \cos(x)} - \frac{ib_1 x \sin(x)}{-2ib \sin(x) + 2b \cos(x)} + \frac{b_1 x \cos(x)}{-2ib \sin(x) + 2b \cos(x)} - \frac{ib_1 \cos(x)}{-2ib \sin(x) + 2b \cos(x)} \\ \frac{a_1 x \sin(x)}{2ib \sin(x) + 2b \cos(x)} - \frac{ia_1 x \cos(x)}{2ib \sin(x) + 2b \cos(x)} - \frac{a_1 \cos(x)}{2ib \sin(x) + 2b \cos(x)} + \frac{ib_1 x \sin(x)}{2ib \sin(x) + 2b \cos(x)} + \frac{b_1 x \cos(x)}{2ib \sin(x) + 2b \cos(x)} + \frac{ib_1 \cos(x)}{2ib \sin(x) + 2b \cos(x)} \\ \frac{aa_1 x}{a^2 + b^2} + \frac{ab_1 \log\left(\frac{a \sin(x)}{b} + \cos(x)\right)}{a^2 + b^2} - \frac{a_1 b \log\left(\frac{a \sin(x)}{b} + \cos(x)\right)}{a^2 + b^2} + \frac{bb_1 x}{a^2 + b^2} \end{cases}$$

input `integrate((b1*cos(x)+a1*sin(x))/(b*cos(x)+a*sin(x)),x)`

output `Piecewise((zoo*(-a1*log(cos(x)) + b1*x), Eq(a, 0) & Eq(b, 0)), ((a1*x + b1*log(sin(x)))/a, Eq(b, 0)), (a1*x*sin(x)/(-2*I*b*sin(x) + 2*b*cos(x)) + I*a1*x*cos(x)/(-2*I*b*sin(x) + 2*b*cos(x)) - a1*cos(x)/(-2*I*b*sin(x) + 2*b*cos(x)) - I*b1*x*sin(x)/(-2*I*b*sin(x) + 2*b*cos(x)) + b1*x*cos(x)/(-2*I*b*sin(x) + 2*b*cos(x)) - I*b1*cos(x)/(-2*I*b*sin(x) + 2*b*cos(x)), Eq(a, -I*b)), (a1*x*sin(x)/(2*I*b*sin(x) + 2*b*cos(x)) - I*a1*x*cos(x)/(2*I*b*sin(x) + 2*b*cos(x)) - a1*cos(x)/(2*I*b*sin(x) + 2*b*cos(x)) + I*b1*x*sin(x)/(2*I*b*sin(x) + 2*b*cos(x)) + b1*x*cos(x)/(2*I*b*sin(x) + 2*b*cos(x)) + I*b1*cos(x)/(2*I*b*sin(x) + 2*b*cos(x)), Eq(a, I*b)), (a*a1*x/(a**2 + b**2) + a*b1*log(a*sin(x)/b + cos(x))/(a**2 + b**2) - a1*b*log(a*sin(x)/b + cos(x))/(a**2 + b**2) + b*b1*x/(a**2 + b**2), True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 181 vs. $2(48) = 96$.

Time = 0.11 (sec) , antiderivative size = 181, normalized size of antiderivative = 3.77

$$\int \frac{b_1 \cos(x) + a_1 \sin(x)}{b \cos(x) + a \sin(x)} dx$$

$$= a_1 \left(\frac{2 a \arctan\left(\frac{\sin(x)}{\cos(x)+1}\right)}{a^2 + b^2} - \frac{b \log\left(-b - \frac{2 a \sin(x)}{\cos(x)+1} + \frac{b \sin(x)^2}{(\cos(x)+1)^2}\right)}{a^2 + b^2} + \frac{b \log\left(\frac{\sin(x)^2}{(\cos(x)+1)^2} + 1\right)}{a^2 + b^2} \right)$$

$$+ b_1 \left(\frac{2 b \arctan\left(\frac{\sin(x)}{\cos(x)+1}\right)}{a^2 + b^2} + \frac{a \log\left(-b - \frac{2 a \sin(x)}{\cos(x)+1} + \frac{b \sin(x)^2}{(\cos(x)+1)^2}\right)}{a^2 + b^2} - \frac{a \log\left(\frac{\sin(x)^2}{(\cos(x)+1)^2} + 1\right)}{a^2 + b^2} \right)$$

input `integrate((b1*cos(x)+a1*sin(x))/(b*cos(x)+a*sin(x)),x, algorithm="maxima")`

output `a1*(2*a*arctan(sin(x)/(cos(x) + 1))/(a^2 + b^2) - b*log(-b - 2*a*sin(x)/(cos(x) + 1) + b*sin(x)^2/(cos(x) + 1)^2)/(a^2 + b^2) + b*log(sin(x)^2/(cos(x) + 1)^2 + 1)/(a^2 + b^2)) + b1*(2*b*arctan(sin(x)/(cos(x) + 1))/(a^2 + b^2) + a*log(-b - 2*a*sin(x)/(cos(x) + 1) + b*sin(x)^2/(cos(x) + 1)^2)/(a^2 + b^2) - a*log(sin(x)^2/(cos(x) + 1)^2 + 1)/(a^2 + b^2))`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.60

$$\int \frac{b_1 \cos(x) + a_1 \sin(x)}{b \cos(x) + a \sin(x)} dx = \frac{(a a_1 + b b_1)x}{a^2 + b^2} + \frac{(a_1 b - a b_1) \log(\tan(x)^2 + 1)}{2(a^2 + b^2)}$$

$$- \frac{(a a_1 b - a^2 b_1) \log(|a \tan(x) + b|)}{a^3 + a b^2}$$

input `integrate((b1*cos(x)+a1*sin(x))/(b*cos(x)+a*sin(x)),x, algorithm="giac")`

output `(a*a1 + b*b1)*x/(a^2 + b^2) + 1/2*(a1*b - a*b1)*log(tan(x)^2 + 1)/(a^2 + b^2) - (a*a1*b - a^2*b1)*log(abs(a*tan(x) + b))/(a^3 + a*b^2)`

Mupad [B] (verification not implemented)

Time = 6.17 (sec) , antiderivative size = 2034, normalized size of antiderivative = 42.38

$$\int \frac{b1 \cos(x) + a1 \sin(x)}{b \cos(x) + a \sin(x)} dx = \text{Too large to display}$$

input `int((b1*cos(x) + a1*sin(x))/(b*cos(x) + a*sin(x)),x)`

output

```
(2*atan((tan(x/2)*((((a*a1 + b*b1)^3*(96*a^4*b + 96*a^2*b^3))/(a^2 + b^2)
^3 + (((a*a1 + b*b1)*(32*a^2*a1*b^2 - ((2*a*b1 - 2*a1*b)*(96*a^4*b + 96*a
^2*b^3)))/(2*(a^2 + b^2)) - 64*a1*b^4 + 128*a*b^3*b1 + 32*a^3*b*b1))/(a^2 +
b^2) - ((a*a1 + b*b1)*(2*a*b1 - 2*a1*b)*(96*a^4*b + 96*a^2*b^3))/(2*(a^2
+ b^2)^2))*(2*a*b1 - 2*a1*b))/(2*(a^2 + b^2)) - ((a*a1 + b*b1)*(32*b^3*b1^
2 - ((2*a*b1 - 2*a1*b)*(32*a^2*a1*b^2 - ((2*a*b1 - 2*a1*b)*(96*a^4*b + 96*
a^2*b^3)))/(2*(a^2 + b^2)) - 64*a1*b^4 + 128*a*b^3*b1 + 32*a^3*b*b1))/(2*(a
^2 + b^2)) + 64*a^2*a1^2*b - 96*a^2*b*b1^2 + 192*a*a1*b^2*b1))/(a^2 + b^2)
)*(a^4*a1^2 + 4*a1^2*b^4 - 4*a^4*b1^2 - b^4*b1^2 - 13*a^2*a1^2*b^2 + 13*a^
2*b^2*b1^2 - 18*a*a1*b^3*b1 + 18*a^3*a1*b*b1))/((a^2 + b^2)^2*(a^2*a1^2 +
4*a^2*b1^2 + 4*a1^2*b^2 + b^2*b1^2 - 6*a*a1*b*b1)^2) - (((((a*a1 + b*b1)*
(32*a^2*a1*b^2 - ((2*a*b1 - 2*a1*b)*(96*a^4*b + 96*a^2*b^3))/(2*(a^2 + b^2)
)) - 64*a1*b^4 + 128*a*b^3*b1 + 32*a^3*b*b1))/(a^2 + b^2) - ((a*a1 + b*b1)
*(2*a*b1 - 2*a1*b)*(96*a^4*b + 96*a^2*b^3))/(2*(a^2 + b^2)^2))*(a*a1 + b*b
1))/(a^2 + b^2) - 32*a1*b^2*b1^2 - 64*a1^3*b^2 + ((2*a*b1 - 2*a1*b)*(32*b^
3*b1^2 - ((2*a*b1 - 2*a1*b)*(32*a^2*a1*b^2 - ((2*a*b1 - 2*a1*b)*(96*a^4*b
+ 96*a^2*b^3)))/(2*(a^2 + b^2)) - 64*a1*b^4 + 128*a*b^3*b1 + 32*a^3*b*b1))/
(2*(a^2 + b^2)) + 64*a^2*a1^2*b - 96*a^2*b*b1^2 + 192*a*a1*b^2*b1))/(2*(a^
2 + b^2)) + 32*a*b*b1^3 - ((a*a1 + b*b1)^2*(2*a*b1 - 2*a1*b)*(96*a^4*b + 9
6*a^2*b^3))/(2*(a^2 + b^2)^3) + 64*a*a1^2*b*b1)*(12*a*a1^2*b^3 - 6*a^3*...
```

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.96

$$\int \frac{b1 \cos(x) + a1 \sin(x)}{b \cos(x) + a \sin(x)} dx$$

$$= \frac{\log(\cos(x) b + \sin(x) a) a b 1 - \log(\cos(x) b + \sin(x) a) a 1 b + a a 1 x + b b 1 x}{a^2 + b^2}$$

input `int((b1*cos(x)+a1*sin(x))/(b*cos(x)+a*sin(x)),x)`

output `(log(cos(x)*b + sin(x)*a)*a*b1 - log(cos(x)*b + sin(x)*a)*a1*b + a*a1*x + b*b1*x)/(a**2 + b**2)`

3.170 $\int \frac{1}{\log(t)} dt$

Optimal result	967
Mathematica [A] (verified)	967
Rubi [A] (verified)	968
Maple [B] (verified)	968
Fricas [A] (verification not implemented)	969
Sympy [A] (verification not implemented)	969
Maxima [A] (verification not implemented)	969
Giac [A] (verification not implemented)	970
Mupad [B] (verification not implemented)	970
Reduce [B] (verification not implemented)	970

Optimal result

Integrand size = 4, antiderivative size = 2

$$\int \frac{1}{\log(t)} dt = \text{LogIntegral}(t)$$

output `Li(t)`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{\log(t)} dt = \text{LogIntegral}(t)$$

input `Integrate[Log[t]^(-1), t]`

output `LogIntegral[t]`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2735}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\log(t)} dt$$

↓ 2735

$$\text{LogIntegral}(t)$$

input `Int [Log[t]^(-1), t]`

output `LogIntegral[t]`

Defintions of rubi rules used

rule 2735 `Int [Log[(c_.)*(x_)]^(-1), x_Symbol] :> Simp[LogIntegral[c*x]/c, x] /; FreeQ [c, x]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 8 vs. $2(2) = 4$.

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 4.50

method	result	size
default	$-\text{expIntegral}_1(-\ln(t))$	9
risch	$-\text{expIntegral}_1(-\ln(t))$	9

input `int(1/ln(t),t,method=_RETURNVERBOSE)`

output `-Ei(1,-ln(t))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{\log(t)} dt = \text{log_integral}(t)$$

input `integrate(1/log(t),t, algorithm="fricas")`

output `log_integral(t)`

Sympy [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{\log(t)} dt = \text{li}(t)$$

input `integrate(1/ln(t),t)`

output `li(t)`

Maxima [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

$$\int \frac{1}{\log(t)} dt = \text{Ei}(\log(t))$$

input `integrate(1/log(t),t, algorithm="maxima")`

output `Ei(log(t))`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

$$\int \frac{1}{\log(t)} dt = \text{Ei}(\log(t))$$

input `integrate(1/log(t),t, algorithm="giac")`

output `Ei(log(t))`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{\log(t)} dt = \text{logint}(t)$$

input `int(1/log(t),t)`

output `logint(t)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

$$\int \frac{1}{\log(t)} dt = \text{ei}(\log(t))$$

input `int(1/log(t),t)`

output `ei(log(t))`

3.171 $\int \frac{1}{\log^2(t)} dt$

Optimal result	971
Mathematica [A] (verified)	971
Rubi [A] (verified)	972
Maple [A] (verified)	973
Fricas [A] (verification not implemented)	973
Sympy [A] (verification not implemented)	973
Maxima [A] (verification not implemented)	974
Giac [A] (verification not implemented)	974
Mupad [B] (verification not implemented)	974
Reduce [B] (verification not implemented)	975

Optimal result

Integrand size = 4, antiderivative size = 10

$$\int \frac{1}{\log^2(t)} dt = -\frac{t}{\log(t)} + \text{LogIntegral}(t)$$

output `Li(t)-t/ln(t)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{\log^2(t)} dt = -\frac{t}{\log(t)} + \text{LogIntegral}(t)$$

input `Integrate[Log[t]^(-2), t]`

output `-(t/Log[t]) + LogIntegral[t]`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {2734, 2735}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{1}{\log^2(t)} dt \\ \downarrow 2734 \\ \int \frac{1}{\log(t)} dt - \frac{t}{\log(t)} \\ \downarrow 2735 \\ \text{LogIntegral}(t) - \frac{t}{\log(t)} \end{array}$$

input

```
Int [Log[t]^(-2), t]
```

output

```
-(t/Log[t]) + LogIntegral[t]
```

Defintions of rubi rules used

rule 2734

```
Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_), x_Symbol] := Simp[x*((a + b
*Log[c*x^n])^(p + 1)/(b*n*(p + 1))), x] - Simp[1/(b*n*(p + 1)) Int[(a + b
*Log[c*x^n])^(p + 1), x], x] /; FreeQ[{a, b, c, n}, x] && LtQ[p, -1] && Int
egerQ[2*p]
```

rule 2735

```
Int[Log[(c_.)*(x_)^(-1), x_Symbol] := Simp[LogIntegral[c*x]/c, x] /; FreeQ
[c, x]
```

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.70

method	result	size
default	$-\frac{t}{\ln(t)} - \text{expIntegral}_1(-\ln(t))$	17
risch	$-\frac{t}{\ln(t)} - \text{expIntegral}_1(-\ln(t))$	17

input `int(1/ln(t)^2,t,method=_RETURNVERBOSE)`output `-t/ln(t)-Ei(1,-ln(t))`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.40

$$\int \frac{1}{\log^2(t)} dt = \frac{\log(t) \log_integral(t) - t}{\log(t)}$$

input `integrate(1/log(t)^2,t, algorithm="fricas")`output `(log(t)*log_integral(t) - t)/log(t)`**Sympy [A] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int \frac{1}{\log^2(t)} dt = -\frac{t}{\log(t)} + \text{li}(t)$$

input `integrate(1/ln(t)**2,t)`output `-t/log(t) + li(t)`

Maxima [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.60

$$\int \frac{1}{\log^2(t)} dt = \Gamma(-1, -\log(t))$$

input `integrate(1/log(t)^2,t, algorithm="maxima")`output `gamma(-1, -log(t))`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

$$\int \frac{1}{\log^2(t)} dt = -\frac{t}{\log(t)} + \text{Ei}(\log(t))$$

input `integrate(1/log(t)^2,t, algorithm="giac")`output `-t/log(t) + Ei(log(t))`**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{\log^2(t)} dt = \text{logint}(t) - \frac{t}{\ln(t)}$$

input `int(1/log(t)^2,t)`output `logint(t) - t/log(t)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.50

$$\int \frac{1}{\log^2(t)} dt = \frac{ei(\log(t)) \log(t) - t}{\log(t)}$$

input `int(1/log(t)^2,t)`

output `(ei(log(t))*log(t) - t)/log(t)`

3.172 $\int \log^{-1-n}(t) dt$

Optimal result	976
Mathematica [A] (verified)	976
Rubi [A] (verified)	977
Maple [F]	978
Fricas [C] (verification not implemented)	978
Sympy [A] (verification not implemented)	978
Maxima [A] (verification not implemented)	979
Giac [F]	979
Mupad [B] (verification not implemented)	979
Reduce [F]	980

Optimal result

Integrand size = 8, antiderivative size = 22

$$\int \log^{-1-n}(t) dt = -\Gamma(-n, -\log(t))(-\log(t))^n \log^{-n}(t)$$

output `-GAMMA(-n, -ln(t))*(-ln(t))^n/(ln(t)^n)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \log^{-1-n}(t) dt = -\Gamma(-n, -\log(t))(-\log(t))^n \log^{-n}(t)$$

input `Integrate[Log[t]^(-1 - n), t]`

output `-((Gamma[-n, -Log[t]]*(-Log[t])^n)/Log[t]^n)`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2736, 2612}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \log^{-n-1}(t) dt$$

$$\downarrow 2736$$

$$\int t \log^{-n-1}(t) d \log(t)$$

$$\downarrow 2612$$

$$(-\log(t))^n \log^{-n}(t) (-\Gamma(-n, -\log(t)))$$

input `Int[Log[t]^(-1 - n),t]`

output `-((Gamma[-n, -Log[t]]*(-Log[t])^n)/Log[t]^n)`

Defintions of rubi rules used

rule 2612 `Int[(F_)^((g_)*(e_) + (f_)*(x_))*((c_) + (d_)*(x_))^(m_), x_Symbol]
:> Simp[(-F^(g*(e - c*(f/d))))*((c + d*x)^FracPart[m]/(d*((-f)*g*(Log[F]/d)
)^(IntPart[m] + 1)*((-f)*g*Log[F]*((c + d*x)/d))^FracPart[m]))*Gamma[m + 1,
((-f)*g*(Log[F]/d)*(c + d*x)], x] /; FreeQ[{F, c, d, e, f, g, m}, x] &&
!IntegerQ[m]`

rule 2736 `Int[((a_) + Log[(c_)*(x_)^(n_)])*(b_)^(p_), x_Symbol]
:> Simp[1/(n*c^(1/n)) Subst[Int[E^(x/n)*(a + b*x)^p, x], x, Log[c*x^n]], x] /; FreeQ[{a, b, c, p}, x] && IntegerQ[1/n]`

Maple [F]

$$\int \ln(t)^{-1-n} dt$$

input `int(ln(t)^(-1-n), t)`

output `int(ln(t)^(-1-n), t)`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \log^{-1-n}(t) dt = e^{(i\pi+i\pi n)}\Gamma(-n, -\log(t))$$

input `integrate(log(t)^(-1-n), t, algorithm="fricas")`

output `e^(I*pi + I*pi*n)*gamma(-n, -log(t))`

Sympy [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \log^{-1-n}(t) dt = (-\log(t))^{n+1} \log(t)^{-n-1} \Gamma(-n, -\log(t))$$

input `integrate(ln(t)**(-1-n), t)`

output `(-log(t))**(n + 1)*log(t)**(-n - 1)*uppergamma(-n, -log(t))`

Maxima [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \log^{-1-n}(t) dt = -(-\log(t))^n \log(t)^{-n} \Gamma(-n, -\log(t))$$

input `integrate(log(t)^(-1-n),t, algorithm="maxima")`output `-(-log(t))^n*log(t)^(-n)*gamma(-n, -log(t))`**Giac [F]**

$$\int \log^{-1-n}(t) dt = \int \log(t)^{-n-1} dt$$

input `integrate(log(t)^(-1-n),t, algorithm="giac")`output `integrate(log(t)^(-n - 1), t)`**Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \log^{-1-n}(t) dt = -\frac{(-\ln(t))^n \Gamma(-n, -\ln(t))}{\ln(t)^n}$$

input `int(1/log(t)^(n + 1),t)`output `-((-log(t))^n*igamma(-n, -log(t)))/log(t)^n`

Reduce [F]

$$\int \log^{-1-n}(t) dt = \int \frac{1}{\log(t)^n \log(t)} dt$$

input `int(log(t)^(-1-n),t)`

output `int(1/(log(t)**n*log(t)),t)`

3.173 $\int \frac{e^{2t}}{-1+t} dt$

Optimal result	981
Mathematica [A] (verified)	981
Rubi [A] (verified)	982
Maple [A] (verified)	982
Fricas [A] (verification not implemented)	983
Sympy [F]	983
Maxima [A] (verification not implemented)	984
Giac [A] (verification not implemented)	984
Mupad [B] (verification not implemented)	984
Reduce [F]	985

Optimal result

Integrand size = 11, antiderivative size = 12

$$\int \frac{e^{2t}}{-1+t} dt = e^2 \text{ExpIntegralEi}(-2(1-t))$$

output `exp(2)*Ei(-2+2*t)`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{e^{2t}}{-1+t} dt = e^2 \text{ExpIntegralEi}(2(-1+t))$$

input `Integrate[E^(2*t)/(-1 + t),t]`

output `E^2*ExpIntegralEi[2*(-1 + t)]`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {2609}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^{2t}}{t-1} dt$$

↓ 2609

$$e^2 \text{ExpIntegralEi}(-2(1-t))$$

input `Int[E^(2*t)/(-1 + t),t]`

output `E^2*ExpIntegralEi[-2*(1 - t)]`

Defintions of rubi rules used

rule 2609

```
Int[(F_)^((g_.)*((e_.) + (f_.)*(x_)))/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]
```

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

method	result	size
derivativedivides	$-e^2 \text{expIntegral}_1(-2t + 2)$	12
default	$-e^2 \text{expIntegral}_1(-2t + 2)$	12
risch	$-e^2 \text{expIntegral}_1(-2t + 2)$	12

input `int(exp(2*t)/(-1+t),t,method=_RETURNVERBOSE)`

output `-exp(2)*Ei(1,-2*t+2)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

$$\int \frac{e^{2t}}{-1+t} dt = \text{Ei}(2t-2) e^2$$

input `integrate(exp(2*t)/(-1+t),t, algorithm="fricas")`

output `Ei(2*t - 2)*e^2`

Sympy [F]

$$\int \frac{e^{2t}}{-1+t} dt = \int \frac{e^{2t}}{t-1} dt$$

input `integrate(exp(2*t)/(-1+t),t)`

output `Integral(exp(2*t)/(t - 1), t)`

Maxima [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

$$\int \frac{e^{2t}}{-1+t} dt = -e^2 E_1(-2t+2)$$

input `integrate(exp(2*t)/(-1+t),t, algorithm="maxima")`

output `-e^2*exp_integral_e(1, -2*t + 2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

$$\int \frac{e^{2t}}{-1+t} dt = \text{Ei}(2t-2) e^2$$

input `integrate(exp(2*t)/(-1+t),t, algorithm="giac")`

output `Ei(2*t - 2)*e^2`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

$$\int \frac{e^{2t}}{-1+t} dt = e^2 \text{ei}(2t-2)$$

input `int(exp(2*t)/(t - 1),t)`

output `exp(2)*ei(2*t - 2)`

Reduce **[F]**

$$\int \frac{e^{2t}}{-1+t} dt = \int \frac{e^{2t}}{t-1} dt$$

input `int(exp(2*t)/(-1+t),t)`

output `int(e**(2*t)/(t - 1),t)`

3.174 $\int \frac{e^{2x}}{2-3x+x^2} dx$

Optimal result	986
Mathematica [A] (verified)	986
Rubi [A] (verified)	987
Maple [A] (verified)	988
Fricas [A] (verification not implemented)	988
Sympy [F]	988
Maxima [F]	989
Giac [A] (verification not implemented)	989
Mupad [F(-1)]	989
Reduce [F]	990

Optimal result

Integrand size = 16, antiderivative size = 22

$$\int \frac{e^{2x}}{2-3x+x^2} dx = e^4 \text{ExpIntegralEi}(-4+2x) - e^2 \text{ExpIntegralEi}(-2+2x)$$

output `exp(4)*Ei(-4+2*x)-exp(2)*Ei(-2+2*x)`

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{e^{2x}}{2-3x+x^2} dx = e^4 \text{ExpIntegralEi}(-4+2x) - e^2 \text{ExpIntegralEi}(-2+2x)$$

input `Integrate[E^(2*x)/(2 - 3*x + x^2),x]`

output `E^4*ExpIntegralEi[-4 + 2*x] - E^2*ExpIntegralEi[-2 + 2*x]`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2698, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^{2x}}{x^2 - 3x + 2} dx$$

$$\downarrow \text{2698}$$

$$\int \left(-\frac{2e^{2x}}{2x-2} - \frac{2e^{2x}}{4-2x} \right) dx$$

$$\downarrow \text{2009}$$

$$e^4 \text{ExpIntegralEi}(2x-4) - e^2 \text{ExpIntegralEi}(2x-2)$$

input `Int[E^(2*x)/(2 - 3*x + x^2),x]`

output `E^4*ExpIntegralEi[-4 + 2*x] - E^2*ExpIntegralEi[-2 + 2*x]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2698 `Int[(F_)^((g_)*((d_) + (e_)*(x_)^(n_)))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] :> Int[ExpandIntegrand[F^(g*(d + e*x)^n), 1/(a + b*x + c*x^2), x], x] /; FreeQ[{F, a, b, c, d, e, g, n}, x]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.05

method	result	size
derivativedivides	$e^2 \exp \text{Integral}_1(-2x + 2) - e^4 \exp \text{Integral}_1(-2x + 4)$	23
default	$e^2 \exp \text{Integral}_1(-2x + 2) - e^4 \exp \text{Integral}_1(-2x + 4)$	23
risch	$e^2 \exp \text{Integral}_1(-2x + 2) - e^4 \exp \text{Integral}_1(-2x + 4)$	23

input `int(exp(2*x)/(x^2-3*x+2),x,method=_RETURNVERBOSE)`output `exp(2)*Ei(1,-2*x+2)-exp(4)*Ei(1,-2*x+4)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{e^{2x}}{2 - 3x + x^2} dx = \text{Ei}(2x - 4) e^4 - \text{Ei}(2x - 2) e^2$$

input `integrate(exp(2*x)/(x^2-3*x+2),x, algorithm="fricas")`output `Ei(2*x - 4)*e^4 - Ei(2*x - 2)*e^2`**Sympy [F]**

$$\int \frac{e^{2x}}{2 - 3x + x^2} dx = \int \frac{e^{2x}}{(x - 2)(x - 1)} dx$$

input `integrate(exp(2*x)/(x**2-3*x+2),x)`output `Integral(exp(2*x)/((x - 2)*(x - 1)), x)`

Maxima [F]

$$\int \frac{e^{2x}}{2 - 3x + x^2} dx = \int \frac{e^{(2x)}}{x^2 - 3x + 2} dx$$

input `integrate(exp(2*x)/(x^2-3*x+2),x, algorithm="maxima")`

output `integrate(e^(2*x)/(x^2 - 3*x + 2), x)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{e^{2x}}{2 - 3x + x^2} dx = \text{Ei}(2x - 4) e^4 - \text{Ei}(2x - 2) e^2$$

input `integrate(exp(2*x)/(x^2-3*x+2),x, algorithm="giac")`

output `Ei(2*x - 4)*e^4 - Ei(2*x - 2)*e^2`

Mupad [F(-1)]

Timed out.

$$\int \frac{e^{2x}}{2 - 3x + x^2} dx = \int \frac{e^{2x}}{x^2 - 3x + 2} dx$$

input `int(exp(2*x)/(x^2 - 3*x + 2),x)`

output `int(exp(2*x)/(x^2 - 3*x + 2), x)`

Reduce [F]

$$\int \frac{e^{2x}}{2 - 3x + x^2} dx = \int \frac{e^{2x}}{x^2 - 3x + 2} dx$$

input `int(exp(2*x)/(x^2-3*x+2),x)`

output `int(e**(2*x)/(x**2 - 3*x + 2),x)`

3.175 $\int \frac{1}{\sqrt{1+t^3}} dt$

Optimal result	991
Mathematica [C] (verified)	991
Rubi [A] (verified)	992
Maple [C] (verified)	993
Fricas [A] (verification not implemented)	993
Sympy [A] (verification not implemented)	994
Maxima [F]	994
Giac [F]	994
Mupad [B] (verification not implemented)	995
Reduce [F]	995

Optimal result

Integrand size = 9, antiderivative size = 103

$$\int \frac{1}{\sqrt{1+t^3}} dt = \frac{2\sqrt{2+\sqrt{3}}(1+t)\sqrt{\frac{1-t+t^2}{(1+\sqrt{3}+t)^2}} \operatorname{EllipticF}\left(\arcsin\left(\frac{1-\sqrt{3}+t}{1+\sqrt{3}+t}\right), -7-4\sqrt{3}\right)}{\sqrt[4]{3}\sqrt{\frac{1+t}{(1+\sqrt{3}+t)^2}}\sqrt{1+t^3}}$$

output

```
2/3*(1+t)*EllipticF((1+t-3^(1/2))/(1+t+3^(1/2)),I*3^(1/2)+2*I)*(1/2*6^(1/2)
)+1/2*2^(1/2))*((t^2-t+1)/(1+t+3^(1/2))^2)^(1/2)*3^(3/4)/(t^3+1)^(1/2)/((
+t)/(1+t+3^(1/2))^2)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.17

$$\int \frac{1}{\sqrt{1+t^3}} dt = t \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, -t^3\right)$$

input

```
Integrate[1/Sqrt[1 + t^3],t]
```


output `t*Hypergeometric2F1[1/3, 1/2, 4/3, -t^3]`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {759}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{t^3+1}} dt$$

↓ 759

$$\frac{2\sqrt{2+\sqrt{3}}(t+1)\sqrt{\frac{t^2-t+1}{(t+\sqrt{3}+1)^2}} \operatorname{EllipticF}\left(\arcsin\left(\frac{t-\sqrt{3}+1}{t+\sqrt{3}+1}\right), -7-4\sqrt{3}\right)}{\sqrt[4]{3}\sqrt{\frac{t+1}{(t+\sqrt{3}+1)^2}}\sqrt{t^3+1}}$$

input `Int[1/Sqrt[1 + t^3],t]`

output `(2*Sqrt[2 + Sqrt[3]]*(1 + t)*Sqrt[(1 - t + t^2)/(1 + Sqrt[3] + t)^2]*EllipticF[ArcSin[(1 - Sqrt[3] + t)/(1 + Sqrt[3] + t)], -7 - 4*Sqrt[3]])/(3^(1/4)*Sqrt[(1 + t)/(1 + Sqrt[3] + t)^2]*Sqrt[1 + t^3])`

Defintions of rubi rules used

rule 759 `Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[2*Sqrt[2 + Sqrt[3]]*(s + r*x)*(Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 + Sqrt[3])*s + r*x)^2]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[s*((s + r*x)/((1 + Sqrt[3])*s + r*x)^2]))*EllipticF[ArcSin[((1 - Sqrt[3])*s + r*x)/((1 + Sqrt[3])*s + r*x)], -7 - 4*Sqrt[3]], x] /; FreeQ[{a, b}, x] && PosQ[a]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4.

Time = 0.14 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.14

method	result	size
meijerg	$t \operatorname{hypergeom}\left(\left[\frac{1}{3}, \frac{1}{2}\right], \left[\frac{4}{3}\right], -t^3\right)$	14
default	$\frac{2\left(\frac{3}{2} - \frac{i\sqrt{3}}{2}\right) \sqrt{\frac{1+t}{\frac{3}{2} - \frac{i\sqrt{3}}{2}}} \sqrt{\frac{t - \frac{1}{2} - \frac{i\sqrt{3}}{2}}{-\frac{3}{2} - \frac{i\sqrt{3}}{2}}} \sqrt{\frac{t - \frac{1}{2} + \frac{i\sqrt{3}}{2}}{-\frac{3}{2} + \frac{i\sqrt{3}}{2}}} \operatorname{EllipticF}\left(\sqrt{\frac{1+t}{\frac{3}{2} - \frac{i\sqrt{3}}{2}}}, \sqrt{\frac{-\frac{3}{2} + \frac{i\sqrt{3}}{2}}{-\frac{3}{2} - \frac{i\sqrt{3}}{2}}}\right)}{\sqrt{t^3+1}}$	116
elliptic	$\frac{2\left(\frac{3}{2} - \frac{i\sqrt{3}}{2}\right) \sqrt{\frac{1+t}{\frac{3}{2} - \frac{i\sqrt{3}}{2}}} \sqrt{\frac{t - \frac{1}{2} - \frac{i\sqrt{3}}{2}}{-\frac{3}{2} - \frac{i\sqrt{3}}{2}}} \sqrt{\frac{t - \frac{1}{2} + \frac{i\sqrt{3}}{2}}{-\frac{3}{2} + \frac{i\sqrt{3}}{2}}} \operatorname{EllipticF}\left(\sqrt{\frac{1+t}{\frac{3}{2} - \frac{i\sqrt{3}}{2}}}, \sqrt{\frac{-\frac{3}{2} + \frac{i\sqrt{3}}{2}}{-\frac{3}{2} - \frac{i\sqrt{3}}{2}}}\right)}{\sqrt{t^3+1}}$	116

input `int(1/(t^3+1)^(1/2), t, method=_RETURNVERBOSE)`

output `t*hypergeom([1/3, 1/2], [4/3], -t^3)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.06

$$\int \frac{1}{\sqrt{1+t^3}} dt = 2 \operatorname{weierstrassPInverse}(0, -4, t)$$

input `integrate(1/(t^3+1)^(1/2), t, algorithm="fricas")`

output `2*weierstrassPInverse(0, -4, t)`

Sympy [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.26

$$\int \frac{1}{\sqrt{1+t^3}} dt = \frac{t\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{1}{2} \middle| \frac{4}{3} \middle| t^3 e^{i\pi}\right)}{3\Gamma\left(\frac{4}{3}\right)}$$

input `integrate(1/(t**3+1)**(1/2),t)`output `t*gamma(1/3)*hyper((1/3, 1/2), (4/3,), t**3*exp_polar(I*pi))/(3*gamma(4/3))`**Maxima [F]**

$$\int \frac{1}{\sqrt{1+t^3}} dt = \int \frac{1}{\sqrt{t^3+1}} dt$$

input `integrate(1/(t^3+1)^(1/2),t, algorithm="maxima")`output `integrate(1/sqrt(t^3 + 1), t)`**Giac [F]**

$$\int \frac{1}{\sqrt{1+t^3}} dt = \int \frac{1}{\sqrt{t^3+1}} dt$$

input `integrate(1/(t^3+1)^(1/2),t, algorithm="giac")`output `integrate(1/sqrt(t^3 + 1), t)`

Mupad [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.50

$$\int \frac{1}{\sqrt{1+t^3}} dt$$

$$= \frac{(3 + \sqrt{3} i) \sqrt{\frac{t - \frac{1}{2} + \frac{\sqrt{3} i}{2}}{-\frac{3}{2} + \frac{\sqrt{3} i}{2}}} \sqrt{\frac{t+1}{\frac{3}{2} + \frac{\sqrt{3} i}{2}}} \sqrt{\frac{\frac{1}{2} - t + \frac{\sqrt{3} i}{2}}{\frac{3}{2} + \frac{\sqrt{3} i}{2}}} F\left(\operatorname{asin}\left(\sqrt{\frac{t+1}{\frac{3}{2} + \frac{\sqrt{3} i}{2}}}\right) \middle| -\frac{\frac{3}{2} + \frac{\sqrt{3} i}{2}}{-\frac{3}{2} + \frac{\sqrt{3} i}{2}}\right)}{\sqrt{t^3 + \left(-\left(-\frac{1}{2} + \frac{\sqrt{3} i}{2}\right)\left(\frac{1}{2} + \frac{\sqrt{3} i}{2}\right) - 1\right) t - \left(-\frac{1}{2} + \frac{\sqrt{3} i}{2}\right)\left(\frac{1}{2} + \frac{\sqrt{3} i}{2}\right)}}$$

input `int(1/(t^3 + 1)^(1/2),t)`output `((3^(1/2)*1i + 3)*((t + (3^(1/2)*1i)/2 - 1/2)/((3^(1/2)*1i)/2 - 3/2))^(1/2) * ((t + 1)/((3^(1/2)*1i)/2 + 3/2))^(1/2) * (((3^(1/2)*1i)/2 - t + 1/2)/((3^(1/2)*1i)/2 + 3/2))^(1/2) * ellipticF(asin(((t + 1)/((3^(1/2)*1i)/2 + 3/2))^(1/2)), -((3^(1/2)*1i)/2 + 3/2)/((3^(1/2)*1i)/2 - 3/2))/(t^3 - t*((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2) + 1) - ((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2))^(1/2)`**Reduce [F]**

$$\int \frac{1}{\sqrt{1+t^3}} dt = \int \frac{\sqrt{t^3+1}}{t^3+1} dt$$

input `int(1/(t^3+1)^(1/2),t)`output `int(sqrt(t**3 + 1)/(t**3 + 1),t)`

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions 996
4.2 Links to plain text integration problems used in this report for each CAS . 1014

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal."}
        ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal."}
    ]
  ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order of result is higher than in optimal."}
  ]
  ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]===Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]===Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]===Plus || Head[expn]===Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```

```

    Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
    If[AppellFunctionQ[Head[expn]],
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
    If[Head[expn]===RootSum,
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
    If[Head[expn]===Integrate || Head[expn]===Int,
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
    9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

```

```

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

```

```

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

```

```

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```


Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

#   antiderivative
#   "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co

        fi;
    fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                    convert(ExpnType_result,string)," vs. order ",
                    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
    9
  end if
end proc
```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```



```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file