

Computer Algebra Independent Integration Tests

Summer 2024

0-Independent-test-suites/2-Bondarenko-Problems

Nasser M. Abbasi

May 17, 2024

Compiled on May 17, 2024 at 10:57pm

Contents

1	Introduction	3
1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	9
1.4	Performance based on number of rules Rubi used	11
1.5	Performance based on number of steps Rubi used	12
1.6	Solved integrals histogram based on leaf size of result	13
1.7	Solved integrals histogram based on CPU time used	14
1.8	Leaf size vs. CPU time used	15
1.9	list of integrals with no known antiderivative	16
1.10	List of integrals solved by CAS but has no known antiderivative	16
1.11	list of integrals solved by CAS but failed verification	16
1.12	Timing	17
1.13	Verification	17
1.14	Important notes about some of the results	18
1.15	Current tree layout of integration tests	21
1.16	Design of the test system	22
2	detailed summary tables of results	23
2.1	List of integrals sorted by grade for each CAS	24
2.2	Detailed conclusion table per each integral for all CAS systems	28
2.3	Detailed conclusion table specific for Rubi results	37
3	Listing of integrals	39
3.1	$\int \frac{1}{\sqrt{2+\cos(z)+\sin(z)}} dz$	41
3.2	$\int \frac{1}{(\sqrt{1-x}+\sqrt{1+x})^2} dx$	46
3.3	$\int \frac{1}{(1+\cos(x))^2} dx$	51
3.4	$\int \frac{\sin(x)}{\sqrt{1+x}} dx$	56
3.5	$\int \frac{1}{(\cos(x)+\sin(x))^6} dx$	62
3.6	$\int \log\left(\frac{1}{x^4} + x^4\right) dx$	69

3.7	$\int \frac{\log(1+x)}{x\sqrt{1+\sqrt{1+x}}} dx$	80
3.8	$\int \frac{\sqrt{1+\sqrt{1+x}} \log(1+x)}{x} dx$	87
3.9	$\int \frac{1}{1+\sqrt{x+\sqrt{1+x^2}}} dx$	95
3.10	$\int \frac{\sqrt{1+x}}{x+\sqrt{1+\sqrt{1+x}}} dx$	101
3.11	$\int \frac{1}{x-\sqrt{1+\sqrt{1+x}}} dx$	107
3.12	$\int \frac{x}{x+\sqrt{1-\sqrt{1+x}}} dx$	114
3.13	$\int \frac{\sqrt{x+\sqrt{1+x}}}{\sqrt{1+x(1+x^2)}} dx$	120
3.14	$\int \frac{\sqrt{x+\sqrt{1+x}}}{1+x^2} dx$	127
3.15	$\int \sqrt{1+\sqrt{x}+\sqrt{1+2\sqrt{x}+2x}} dx$	133
3.16	$\int \sqrt{\sqrt{2}+\sqrt{x}+\sqrt{2+2\sqrt{2}\sqrt{x}+2x}} dx$	138
3.17	$\int \frac{\sqrt{x+\sqrt{1+x}}}{x^2} dx$	144
3.18	$\int \sqrt{\sqrt{1+\frac{1}{x}+\frac{1}{x}}} dx$	151
3.19	$\int \frac{\sqrt{1+e^{-x}}}{-e^{-x}+e^x} dx$	159
3.20	$\int \sqrt{1+e^{-x}} \operatorname{csch}(x) dx$	165
3.21	$\int \frac{1}{(\cos(x)+\cos(3x))^5} dx$	172
3.22	$\int \frac{1}{(1+\cos(x)+\sin(x))^2} dx$	182
3.23	$\int \sqrt{1+\tanh(4x)} dx$	188
3.24	$\int \frac{\tanh(x)}{\sqrt{e^x+e^{2x}}} dx$	193
3.25	$\int \sqrt{\operatorname{sech}(x) \sinh(2x)} dx$	202
3.26	$\int \log(x^2+\sqrt{1-x^2}) dx$	209
3.27	$\int \frac{\log(1+e^x)}{1+e^{2x}} dx$	218
3.28	$\int \cosh(x) \log^2(1+\cosh^2(x)) dx$	224
3.29	$\int \cosh(x) \log^2(\cosh^2(x)+\sinh(x)) dx$	230
3.30	$\int \frac{\log(x+\sqrt{1+x})}{1+x^2} dx$	239
3.31	$\int \frac{\log^2(x+\sqrt{1+x})}{(1+x)^2} dx$	246
3.32	$\int \frac{\log(x+\sqrt{1+x})}{x} dx$	254
3.33	$\int \arctan(2 \tan(x)) dx$	262
3.34	$\int \frac{\arctan(x) \log(x)}{x} dx$	270
3.35	$\int \sqrt{1+x^2} \arctan(x)^2 dx$	275
4	Appendix	282
4.1	Listing of Grading functions	282
4.2	Links to plain text integration problems used in this report for each CAS300	

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	9
1.4	Performance based on number of rules Rubi used	11
1.5	Performance based on number of steps Rubi used	12
1.6	Solved integrals histogram based on leaf size of result	13
1.7	Solved integrals histogram based on CPU time used	14
1.8	Leaf size vs. CPU time used	15
1.9	list of integrals with no known antiderivative	16
1.10	List of integrals solved by CAS but has no known antiderivative	16
1.11	list of integrals solved by CAS but failed verification	16
1.12	Timing	17
1.13	Verification	17
1.14	Important notes about some of the results	18
1.15	Current tree layout of integration tests	21
1.16	Design of the test system	22

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [35]. This is test number [2].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Mathematica	97.14 (34)	2.86 (1)
Rubi	94.29 (33)	5.71 (2)
Maple	80.00 (28)	20.00 (7)
Fricas	71.43 (25)	28.57 (10)
Giac	48.57 (17)	51.43 (18)
Maxima	45.71 (16)	54.29 (19)
Reduce	31.43 (11)	68.57 (24)
Mupad	25.71 (9)	74.29 (26)
Sympy	25.71 (9)	74.29 (26)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	94.286	0.000	0.000	5.714
Mathematica	65.714	11.429	20.000	2.857
Maple	48.571	14.286	17.143	20.000
Fricas	42.857	25.714	2.857	28.571
Maxima	37.143	5.714	2.857	54.286
Giac	28.571	17.143	2.857	51.429
Sympy	17.143	8.571	0.000	74.286
Mupad	0.000	25.714	0.000	74.286
Reduce	0.000	31.429	0.000	68.571

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Mathematica	1	100.00	0.00	0.00
Rubi	2	100.00	0.00	0.00
Maple	7	100.00	0.00	0.00
Fricas	10	80.00	0.00	20.00
Giac	18	83.33	0.00	16.67
Maxima	19	100.00	0.00	0.00
Reduce	24	100.00	0.00	0.00
Mupad	26	0.00	100.00	0.00
Sympy	26	92.31	7.69	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.13
Reduce	0.16
Giac	0.18
Mupad	0.36
Rubi	0.46
Fricas	0.59
Mathematica	1.32
Sympy	2.17
Maple	4.99

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Reduce	78.00	1.06	59.00	1.00
Giac	113.12	1.57	55.00	1.03
Maple	134.50	1.15	73.00	0.82
Sympy	143.00	3.02	65.00	1.21
Mupad	160.00	1.47	49.00	0.94
Rubi	170.24	1.10	92.00	1.01
Mathematica	180.21	1.40	87.50	1.00
Fricas	493.56	2.60	81.00	1.38
Maxima	851.50	8.19	53.50	1.24

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram shows that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {9, 10, 12, 24}

Mathematica {7, 8, 26, 31, 35}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand,the_variable)
```

Which gives $\sin(x)^2/2$


1.15 Current tree layout of integration tests



Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	24
2.2	Detailed conclusion table per each integral for all CAS systems	28
2.3	Detailed conclusion table specific for Rubi results	37

2.1 List of integrals sorted by grade for each CAS

Rubi	24
Mma	24
Maple	25
Fricas	25
Maxima	25
Giac	26
Mupad	26
Sympy	26
Reduce	27

Rubi

A grade { 1, 2, 3, 4, 5, 6, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 }

B grade { }

C grade { }

F normal fail { 7, 8 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 2, 3, 5, 7, 8, 9, 10, 11, 12, 15, 16, 17, 18, 21, 22, 23, 24, 27, 28, 29, 32, 34, 35 }

B grade { 19, 20, 31, 33 }

C grade { 1, 4, 6, 13, 14, 25, 26 }

F normal fail { 30 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 3, 4, 10, 11, 12, 20, 21, 22, 23, 25, 27, 30, 32, 33, 34, 35 }

B grade { 2, 17, 19, 24, 26 }

C grade { 5, 6, 7, 8, 13, 14 }

F normal fail { 9, 15, 16, 18, 28, 29, 31 }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 3, 4, 5, 9, 12, 15, 16, 17, 18, 19, 22, 25, 26 }

B grade { 10, 11, 13, 14, 20, 21, 23, 24, 33 }

C grade { 6 }

F normal fail { 27, 28, 29, 30, 31, 32, 34, 35 }

F(-1) timedout fail { }

F(-2) exception fail { 7, 8 }

Maxima

A grade { 1, 3, 5, 7, 8, 10, 11, 12, 19, 20, 22, 33, 34 }

B grade { 21, 23 }

C grade { 4 }

F normal fail { 2, 6, 9, 13, 14, 15, 16, 17, 18, 24, 25, 26, 27, 28, 29, 30, 31, 32, 35 }

F(-1) timedout fail { }

F(-2) exception fail { }

Giac

A grade { 1, 3, 5, 6, 10, 11, 12, 21, 22, 23 }

B grade { 2, 17, 19, 20, 24, 26 }

C grade { 4 }

F normal fail { 7, 8, 9, 15, 18, 25, 27, 28, 29, 30, 31, 32, 33, 34, 35 }

F(-1) timedout fail { }

F(-2) exception fail { 13, 14, 16 }

Mupad

A grade { }

B grade { 1, 2, 3, 5, 6, 21, 22, 23, 26 }

C grade { }

F normal fail { }

F(-1) timedout fail { 4, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 24, 25, 27, 28, 29, 30, 31, 32, 33, 34, 35 }

F(-2) exception fail { }

Sympy

A grade { 3, 4, 6, 10, 11, 12 }

B grade { 1, 5, 22 }

C grade { }

F normal fail { 2, 9, 13, 14, 15, 16, 17, 18, 19, 20, 21, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 }

F(-1) timedout fail { 7, 8 }

F(-2) exception fail { }

Reduce

A grade { }

B grade { 1, 2, 3, 5, 6, 10, 11, 12, 17, 22, 26 }

C grade { }

F normal fail { 4, 7, 8, 9, 13, 14, 15, 16, 18, 19, 20, 21, 23, 24, 25, 27, 28, 29, 30, 31, 32, 33, 34, 35 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	77	18	20	25	61	18	17	16
N.S.	1	1.00	3.50	0.82	0.91	1.14	2.77	0.82	0.77	0.73
time (sec)	N/A	0.160	0.050	0.189	0.104	0.084	3.797	0.126	0.147	0.147

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	A	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	30	49	50	0	44	0	147	59	49
N.S.	1	0.94	1.53	1.56	0.00	1.38	0.00	4.59	1.84	1.53
time (sec)	N/A	0.190	0.116	0.025	0.000	0.070	0.000	0.143	0.154	0.529

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	16	16	23	20	14	15	14	14
N.S.	1	1.00	0.64	0.64	0.92	0.80	0.56	0.60	0.56	0.56
time (sec)	N/A	0.190	0.008	0.056	0.029	0.068	0.147	0.122	0.149	0.106

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	C	A	A	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	68	42	112	46	53	43	11	0
N.S.	1	1.00	1.17	0.72	1.93	0.79	0.91	0.74	0.19	0.00
time (sec)	N/A	0.349	0.013	0.138	0.112	0.081	0.823	0.120	0.153	0.000

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	55	26	30	56	67	838	32	67	88
N.S.	1	1.10	0.52	0.60	1.12	1.34	16.76	0.64	1.34	1.76
time (sec)	N/A	0.269	0.026	0.369	0.032	0.071	2.347	0.122	0.150	0.156

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	C	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	334	362	30	34	0	149	26	248	216	313
N.S.	1	1.08	0.09	0.10	0.00	0.45	0.08	0.74	0.65	0.94
time (sec)	N/A	0.615	0.004	0.230	0.000	0.080	1.144	0.127	0.152	0.149

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	C	A	F(-2)	F(-1)	F	F	F(-1)
verified	N/A	N/A	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	291	0	310	172	366	0	0	0	335	0
N.S.	1	0.00	1.07	0.59	1.26	0.00	0.00	0.00	1.15	0.00
time (sec)	N/A	0.000	0.186	0.222	0.115	0.000	0.000	0.000	0.187	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	C	A	F(-2)	F(-1)	F	F	F(-1)
verified	N/A	N/A	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	308	0	326	199	378	0	0	0	17	0
N.S.	1	0.00	1.06	0.65	1.23	0.00	0.00	0.00	0.06	0.00
time (sec)	N/A	0.000	0.095	0.028	0.126	0.000	0.000	0.000	0.150	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	F	F	F(-1)
verified	N/A	No	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	84	73	105	0	0	66	0	0	155	0
N.S.	1	0.87	1.25	0.00	0.00	0.79	0.00	0.00	1.85	0.00
time (sec)	N/A	0.252	0.129	0.000	0.000	0.073	0.000	0.000	29.220	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	71	41	34	51	101	65	51	46	0
N.S.	1	1.73	1.00	0.83	1.24	2.46	1.59	1.24	1.12	0.00
time (sec)	N/A	0.432	0.058	0.066	0.109	0.073	4.681	0.129	0.153	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	75	69	46	63	112	76	67	73	0
N.S.	1	1.03	0.95	0.63	0.86	1.53	1.04	0.92	1.00	0.00
time (sec)	N/A	0.356	0.082	0.146	0.111	0.077	2.396	0.380	0.148	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	103	52	60	77	110	88	79	55	0
N.S.	1	1.41	0.71	0.82	1.05	1.51	1.21	1.08	0.75	0.00
time (sec)	N/A	0.514	0.051	0.049	0.113	0.084	3.937	0.333	0.157	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	B	F	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	365	367	217	109	0	5235	0	0	25	0
N.S.	1	1.01	0.59	0.30	0.00	14.34	0.00	0.00	0.07	0.00
time (sec)	N/A	0.989	0.157	0.116	0.000	5.681	0.000	0.000	0.158	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	B	F	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	337	339	212	105	0	4535	0	0	17	0
N.S.	1	1.01	0.63	0.31	0.00	13.46	0.00	0.00	0.05	0.00
time (sec)	N/A	0.835	0.144	0.119	0.000	3.086	0.000	0.000	0.169	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	74	0	0	56	0	0	17	0
N.S.	1	1.00	0.96	0.00	0.00	0.73	0.00	0.00	0.22	0.00
time (sec)	N/A	0.245	10.027	0.000	0.000	0.358	0.000	0.000	0.172	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	118	118	112	0	0	73	0	0	20	0
N.S.	1	1.00	0.95	0.00	0.00	0.62	0.00	0.00	0.17	0.00
time (sec)	N/A	0.391	10.055	0.000	0.000	0.724	0.000	0.000	0.173	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	A	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	92	77	298	0	81	0	188	86	0
N.S.	1	1.11	0.93	3.59	0.00	0.98	0.00	2.27	1.04	0.00
time (sec)	N/A	0.307	0.144	0.042	0.000	1.654	0.000	0.363	0.174	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	96	106	89	0	0	122	0	0	15	0
N.S.	1	1.10	0.93	0.00	0.00	1.27	0.00	0.00	0.16	0.00
time (sec)	N/A	0.292	0.350	0.000	0.000	1.675	0.000	0.000	180.019	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	A	A	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	65	49	36	34	0	55	27	0
N.S.	1	1.00	2.60	1.96	1.44	1.36	0.00	2.20	1.08	0.00
time (sec)	N/A	0.241	0.111	0.071	0.110	0.072	0.000	0.147	0.161	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	B	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	57	33	35	55	0	54	20	0
N.S.	1	1.00	2.28	1.32	1.40	2.20	0.00	2.16	0.80	0.00
time (sec)	N/A	0.234	0.050	0.217	0.104	0.074	0.000	0.137	0.147	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	108	184	213	95	12209	219	0	104	59	307
N.S.	1	1.70	1.97	0.88	113.05	2.03	0.00	0.96	0.55	2.84
time (sec)	N/A	0.330	3.885	134.336	0.688	0.113	0.000	0.122	0.160	0.699

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	56	27	40	46	66	30	46	26
N.S.	1	1.00	1.93	0.93	1.38	1.59	2.28	1.03	1.59	0.90
time (sec)	N/A	0.221	0.025	0.185	0.027	0.081	0.298	0.112	0.157	0.158

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	26	20	43	129	0	21	9	19
N.S.	1	1.00	1.00	0.77	1.65	4.96	0.00	0.81	0.35	0.73
time (sec)	N/A	0.180	0.014	0.141	0.107	0.074	0.000	0.131	0.155	0.090

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F	B	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	110	126	121	366	0	501	0	470	24	0
N.S.	1	1.15	1.10	3.33	0.00	4.55	0.00	4.27	0.22	0.00
time (sec)	N/A	0.546	0.089	0.414	0.000	0.093	0.000	0.241	0.155	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	86	75	0	23	0	0	15	0
N.S.	1	1.00	2.15	1.88	0.00	0.58	0.00	0.00	0.38	0.00
time (sec)	N/A	0.437	10.899	0.316	0.000	0.070	0.000	0.000	0.167	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	A	F	B	B	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	185	349	920	469	0	270	0	301	179	608
N.S.	1	1.89	4.97	2.54	0.00	1.46	0.00	1.63	0.97	3.29
time (sec)	N/A	1.107	0.359	0.179	0.000	0.114	0.000	0.190	0.177	1.194

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	102	102	102	83	0	0	0	0	18	0
N.S.	1	1.00	1.00	0.81	0.00	0.00	0.00	0.00	0.18	0.00
time (sec)	N/A	0.316	0.015	0.058	0.000	0.000	0.000	0.000	0.159	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	159	161	122	0	0	0	0	0	342	0
N.S.	1	1.01	0.77	0.00	0.00	0.00	0.00	0.00	2.15	0.00
time (sec)	N/A	0.401	0.071	0.000	0.000	0.000	0.000	0.000	0.167	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	395	410	389	0	0	0	0	0	15	0
N.S.	1	1.04	0.98	0.00	0.00	0.00	0.00	0.00	0.04	0.00
time (sec)	N/A	0.730	0.158	0.000	0.000	0.000	0.000	0.000	0.154	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	981	983	0	722	0	0	0	0	17	0
N.S.	1	1.00	0.00	0.74	0.00	0.00	0.00	0.00	0.02	0.00
time (sec)	N/A	1.398	0.000	0.109	0.000	0.000	0.000	0.000	0.161	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	555	580	1280	0	0	0	0	0	22	0
N.S.	1	1.05	2.31	0.00	0.00	0.00	0.00	0.00	0.04	0.00
time (sec)	N/A	0.928	6.916	0.000	0.000	0.000	0.000	0.000	0.156	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	313	337	303	252	0	0	0	0	13	0
N.S.	1	1.08	0.97	0.81	0.00	0.00	0.00	0.00	0.04	0.00
time (sec)	N/A	0.569	0.071	0.135	0.000	0.000	0.000	0.000	0.160	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	83	262	113	84	220	0	0	7	0
N.S.	1	1.04	3.28	1.41	1.05	2.75	0.00	0.00	0.09	0.00
time (sec)	N/A	0.323	0.178	0.533	0.109	0.099	0.000	0.000	0.153	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	51	44	71	31	0	0	0	10	0
N.S.	1	0.89	0.77	1.25	0.54	0.00	0.00	0.00	0.18	0.00
time (sec)	N/A	0.304	0.043	0.642	0.147	0.000	0.000	0.000	0.162	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	121	134	131	178	0	0	0	0	13	0
N.S.	1	1.11	1.08	1.47	0.00	0.00	0.00	0.00	0.11	0.00
time (sec)	N/A	0.533	0.142	0.468	0.000	0.000	0.000	0.000	0.152	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [21] had the largest ratio of [1.55556000000000005]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	2	2	1.00	12	0.167
2	A	2	2	0.94	19	0.105
3	A	4	4	1.00	6	0.667
4	A	8	7	1.00	10	0.700
5	A	6	6	1.10	7	0.857
6	A	11	10	1.08	8	1.250
7	F	0	0	N/A	0.000	N/A
8	F	0	0	N/A	0.000	N/A
9	A	5	4	0.87	19	0.211
10	A	6	5	1.73	25	0.200
11	A	7	6	1.03	19	0.316
12	A	5	4	1.41	21	0.190
13	A	5	4	1.01	28	0.143
14	A	5	4	1.01	21	0.190
15	A	3	2	1.00	27	0.074
16	A	4	3	1.00	36	0.083
17	A	9	8	1.11	17	0.471
18	A	9	8	1.10	17	0.471
19	A	9	8	1.00	25	0.320
20	A	9	8	1.00	14	0.571
21	A	15	14	1.70	9	1.556

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	7	6	1.00	8	0.750
23	A	4	3	1.00	10	0.300
24	A	8	7	1.15	16	0.438
25	A	10	10	1.00	11	0.909
26	A	3	3	1.89	16	0.188
27	A	4	3	1.00	16	0.188
28	A	5	4	1.01	12	0.333
29	A	5	4	1.04	13	0.308
30	A	5	4	1.00	18	0.222
31	A	6	5	1.05	18	0.278
32	A	5	4	1.08	14	0.286
33	A	6	5	1.04	5	1.000
34	A	3	3	0.89	8	0.375
35	A	9	8	1.11	14	0.571

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \frac{1}{\sqrt{2+\cos(z)+\sin(z)}} dz$	41
3.2	$\int \frac{1}{(\sqrt{1-x}+\sqrt{1+x})^2} dx$	46
3.3	$\int \frac{1}{(1+\cos(x))^2} dx$	51
3.4	$\int \frac{\sin(x)}{\sqrt{1+x}} dx$	56
3.5	$\int \frac{1}{(\cos(x)+\sin(x))^6} dx$	62
3.6	$\int \log\left(\frac{1}{x^4} + x^4\right) dx$	69
3.7	$\int \frac{\log(1+x)}{x\sqrt{1+\sqrt{1+x}}} dx$	80
3.8	$\int \frac{\sqrt{1+\sqrt{1+x}} \log(1+x)}{x} dx$	87
3.9	$\int \frac{1}{1+\sqrt{x+\sqrt{1+x^2}}} dx$	95
3.10	$\int \frac{\sqrt{1+x}}{x+\sqrt{1+\sqrt{1+x}}} dx$	101
3.11	$\int \frac{1}{x-\sqrt{1+\sqrt{1+x}}} dx$	107
3.12	$\int \frac{x}{x+\sqrt{1-\sqrt{1+x}}} dx$	114
3.13	$\int \frac{\sqrt{x+\sqrt{1+x}}}{\sqrt{1+x}(1+x^2)} dx$	120
3.14	$\int \frac{\sqrt{x+\sqrt{1+x}}}{1+x^2} dx$	127
3.15	$\int \sqrt{1+\sqrt{x}+\sqrt{1+2\sqrt{x}+2x}} dx$	133
3.16	$\int \sqrt{\sqrt{2}+\sqrt{x}+\sqrt{2+2\sqrt{2}\sqrt{x}+2x}} dx$	138
3.17	$\int \frac{\sqrt{x+\sqrt{1+x}}}{x^2} dx$	144
3.18	$\int \sqrt{\sqrt{1+\frac{1}{x}+\frac{1}{x}}} dx$	151
3.19	$\int \frac{\sqrt{1+e^{-x}}}{-e^{-x}+e^x} dx$	159
3.20	$\int \sqrt{1+e^{-x}} \operatorname{csch}(x) dx$	165
3.21	$\int \frac{1}{(\cos(x)+\cos(3x))^5} dx$	172
3.22	$\int \frac{1}{(1+\cos(x)+\sin(x))^2} dx$	182

3.23	$\int \sqrt{1 + \tanh(4x)} dx$	188
3.24	$\int \frac{\tanh(x)}{\sqrt{e^x + e^{2x}}} dx$	193
3.25	$\int \sqrt{\operatorname{sech}(x) \sinh(2x)} dx$	202
3.26	$\int \log(x^2 + \sqrt{1 - x^2}) dx$	209
3.27	$\int \frac{\log(1+e^x)}{1+e^{2x}} dx$	218
3.28	$\int \cosh(x) \log^2(1 + \cosh^2(x)) dx$	224
3.29	$\int \cosh(x) \log^2(\cosh^2(x) + \sinh(x)) dx$	230
3.30	$\int \frac{\log(x+\sqrt{1+x})}{1+x^2} dx$	239
3.31	$\int \frac{\log^2(x+\sqrt{1+x})}{(1+x)^2} dx$	246
3.32	$\int \frac{\log(x+\sqrt{1+x})}{x} dx$	254
3.33	$\int \arctan(2 \tan(x)) dx$	262
3.34	$\int \frac{\arctan(x) \log(x)}{x} dx$	270
3.35	$\int \sqrt{1 + x^2} \arctan(x)^2 dx$	275

3.1 $\int \frac{1}{\sqrt{2} + \cos(z) + \sin(z)} dz$

Optimal result	41
Mathematica [C] (verified)	41
Rubi [A] (verified)	42
Maple [A] (verified)	43
Fricas [A] (verification not implemented)	43
Sympy [B] (verification not implemented)	44
Maxima [A] (verification not implemented)	44
Giac [A] (verification not implemented)	45
Mupad [B] (verification not implemented)	45
Reduce [B] (verification not implemented)	45

Optimal result

Integrand size = 12, antiderivative size = 22

$$\int \frac{1}{\sqrt{2} + \cos(z) + \sin(z)} dz = -\frac{1 - \sqrt{2} \sin(z)}{\cos(z) - \sin(z)}$$

output

```
(-1+sin(z)*2^(1/2))/(cos(z)-sin(z))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.05 (sec) , antiderivative size = 77, normalized size of antiderivative = 3.50

$$\int \frac{1}{\sqrt{2} + \cos(z) + \sin(z)} dz = \frac{-(((1 + 3i) + \sqrt{2}) \cos(\frac{z}{2})) + ((1 + i) - i\sqrt{2}) \sin(\frac{z}{2})}{((1 + i) + \sqrt{2}) \cos(\frac{z}{2}) + i((-1 - i) + \sqrt{2}) \sin(\frac{z}{2})}$$

input

```
Integrate[(Sqrt[2] + Cos[z] + Sin[z])^(-1), z]
```

output

```
(-(((1 + 3*I) + Sqrt[2])*Cos[z/2]) + ((1 + I) - I*Sqrt[2])*Sin[z/2])/(((1 + I) + Sqrt[2])*Cos[z/2] + I*((-1 - I) + Sqrt[2])*Sin[z/2])
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 3593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sin(z) + \cos(z) + \sqrt{2}} dz$$

↓ 3042

$$\int \frac{1}{\sin(z) + \cos(z) + \sqrt{2}} dz$$

↓ 3593

$$-\frac{1 - \sqrt{2} \sin(z)}{\cos(z) - \sin(z)}$$

input `Int[(Sqrt[2] + Cos[z] + Sin[z])^(-1),z]`

output `-((1 - Sqrt[2]*Sin[z])/(Cos[z] - Sin[z]))`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3593 `Int[(cos[(d_.) + (e_.)*(x_)]*(b_.) + (a_.) + (c_.)*sin[(d_.) + (e_.)*(x_)])^(-1), x_Symbol] := Simp[-(c - a*Sin[d + e*x])/(c*e*(c*Cos[d + e*x] - b*Sin[d + e*x])), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[a^2 - b^2 - c^2, 0]`

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

method	result	size
parallelsch	$\frac{2 \tan(\frac{z}{2})}{\tan(\frac{z}{2}) + \sqrt{2} + 1}$	18
default	$-\frac{2}{(\sqrt{2}-1)(\tan(\frac{z}{2}) + \sqrt{2} + 1)}$	21
norman	$\frac{(-2-2\sqrt{2})\tan(\frac{z}{2})+2}{\tan(\frac{z}{2})^2+2\tan(\frac{z}{2})-1}$	32
risch	$-\frac{2}{\sqrt{2+2e^{iz}+i\sqrt{2}}} + \frac{2i}{\sqrt{2+2e^{iz}+i\sqrt{2}}}$	45

input `int(1/(cos(z)+sin(z)+2^(1/2)),z,method=_RETURNVERBOSE)`

output `2*tan(1/2*z)/(tan(1/2*z)+2^(1/2)+1)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.14

$$\int \frac{1}{\sqrt{2} + \cos(z) + \sin(z)} dz = \frac{\sqrt{2} \cos(z) + \sqrt{2} \sin(z) - 2}{2(\cos(z) - \sin(z))}$$

input `integrate(1/(cos(z)+sin(z)+2^(1/2)),z, algorithm="fricas")`

output `1/2*(sqrt(2)*cos(z) + sqrt(2)*sin(z) - 2)/(cos(z) - sin(z))`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 61 vs. $2(15) = 30$.

Time = 3.80 (sec) , antiderivative size = 61, normalized size of antiderivative = 2.77

$$\int \frac{1}{\sqrt{2} + \cos(z) + \sin(z)} dz = -\frac{198}{-239 \tan\left(\frac{z}{2}\right) + 169\sqrt{2} \tan\left(\frac{z}{2}\right) - 70\sqrt{2} + 99} + \frac{140\sqrt{2}}{-239 \tan\left(\frac{z}{2}\right) + 169\sqrt{2} \tan\left(\frac{z}{2}\right) - 70\sqrt{2} + 99}$$

input `integrate(1/(cos(z)+sin(z)+2**(1/2)),z)`

output `-198/(-239*tan(z/2) + 169*sqrt(2)*tan(z/2) - 70*sqrt(2) + 99) + 140*sqrt(2)/(-239*tan(z/2) + 169*sqrt(2)*tan(z/2) - 70*sqrt(2) + 99)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{1}{\sqrt{2} + \cos(z) + \sin(z)} dz = -\frac{2}{\frac{(\sqrt{2}-1)\sin(z)}{\cos(z)+1} + 1}$$

input `integrate(1/(cos(z)+sin(z)+2^(1/2)),z, algorithm="maxima")`

output `-2/((sqrt(2) - 1)*sin(z)/(cos(z) + 1) + 1)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \frac{1}{\sqrt{2} + \cos(z) + \sin(z)} dz = -\frac{2(\sqrt{2} + 1)}{\sqrt{2} + \tan\left(\frac{1}{2}z\right) + 1}$$

input `integrate(1/(cos(z)+sin(z)+2^(1/2)),z, algorithm="giac")`output `-2*(sqrt(2) + 1)/(sqrt(2) + tan(1/2*z) + 1)`**Mupad [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \frac{1}{\sqrt{2} + \cos(z) + \sin(z)} dz = -\frac{2}{\tan\left(\frac{z}{2}\right) (\sqrt{2} - 1) + 1}$$

input `int(1/(cos(z) + sin(z) + 2^(1/2)),z)`output `-2/(tan(z/2)*(2^(1/2) - 1) + 1)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

$$\int \frac{1}{\sqrt{2} + \cos(z) + \sin(z)} dz = \frac{\sqrt{2} \cos(z) - 1}{\cos(z) - \sin(z)}$$

input `int(1/(cos(z)+sin(z)+2^(1/2)),z)`output `(sqrt(2)*cos(z) - 1)/(cos(z) - sin(z))`

3.2 $\int \frac{1}{(\sqrt{1-x} + \sqrt{1+x})^2} dx$

Optimal result	46
Mathematica [A] (verified)	46
Rubi [A] (verified)	47
Maple [B] (verified)	48
Fricas [A] (verification not implemented)	48
Sympy [F]	48
Maxima [F]	49
Giac [B] (verification not implemented)	49
Mupad [B] (verification not implemented)	50
Reduce [B] (verification not implemented)	50

Optimal result

Integrand size = 19, antiderivative size = 32

$$\int \frac{1}{(\sqrt{1-x} + \sqrt{1+x})^2} dx = -\frac{1}{2x} + \frac{\sqrt{1-x^2}}{2x} + \frac{\arcsin(x)}{2}$$

output `-1/2/x+1/2*arcsin(x)+1/2*(-x^2+1)^(1/2)/x`

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.53

$$\int \frac{1}{(\sqrt{1-x} + \sqrt{1+x})^2} dx = \frac{-1 + \sqrt{1-x^2} + 4x \arctan\left(\frac{-\sqrt{2} + \sqrt{1+x}}{\sqrt{1-x}}\right)}{2x}$$

input `Integrate[(Sqrt[1 - x] + Sqrt[1 + x])^(-2), x]`

output `(-1 + Sqrt[1 - x^2] + 4*x*ArcTan[(-Sqrt[2] + Sqrt[1 + x])/Sqrt[1 - x]])/(2*x)`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.94, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {7241, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(\sqrt{1-x} + \sqrt{x+1})^2} dx$$

↓ 7241

$$\frac{1}{4} \int \left(\frac{2}{x^2} - \frac{2\sqrt{1-x^2}}{x^2} \right) dx$$

↓ 2009

$$\frac{1}{4} \left(2 \arcsin(x) + \frac{2\sqrt{1-x^2}}{x} - \frac{2}{x} \right)$$

input `Int[(Sqrt[1 - x] + Sqrt[1 + x])^(-2), x]`

output `(-2/x + (2*Sqrt[1 - x^2])/x + 2*ArcSin[x])/4`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7241 `Int[(u_.)*((e_.)*Sqrt[(a_.) + (b_.)*(x_)^(n_.)] + (f_.)*Sqrt[(c_.) + (d_.)*(x_)^(n_.)])^(m_), x_Symbol] := Simp[(b*e^2 - d*f^2)^m Int[ExpandIntegrand[(u*x^(m*n))/(e*Sqrt[a + b*x^n] - f*Sqrt[c + d*x^n])^m, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ILtQ[m, 0] && EqQ[a*e^2 - c*f^2, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 49 vs. $2(24) = 48$.

Time = 0.02 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.56

method	result	size
default	$-\frac{1}{2x} - \frac{(-\arcsin(x)x - \sqrt{-x^2+1})\sqrt{1+x}\sqrt{1-x}}{2x\sqrt{-x^2+1}}$	50

input `int(1/((1-x)^(1/2)+(1+x)^(1/2))^2,x,method=_RETURNVERBOSE)`

output `-1/2/x-1/2*(-arcsin(x)*x-(-x^2+1)^(1/2))*(1+x)^(1/2)*(1-x)^(1/2)/x/(-x^2+1)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.38

$$\int \frac{1}{(\sqrt{1-x} + \sqrt{1+x})^2} dx = -\frac{2x \arctan\left(\frac{\sqrt{x+1}\sqrt{-x+1}-1}{x}\right) - \sqrt{x+1}\sqrt{-x+1} + 1}{2x}$$

input `integrate(1/((1-x)^(1/2)+(1+x)^(1/2))^2,x, algorithm="fricas")`

output `-1/2*(2*x*arctan((sqrt(x + 1)*sqrt(-x + 1) - 1)/x) - sqrt(x + 1)*sqrt(-x + 1) + 1)/x`

Sympy [F]

$$\int \frac{1}{(\sqrt{1-x} + \sqrt{1+x})^2} dx = \int \frac{1}{(\sqrt{1-x} + \sqrt{x+1})^2} dx$$

input `integrate(1/((1-x)**(1/2)+(1+x)**(1/2))**2,x)`

output `Integral((sqrt(1 - x) + sqrt(x + 1))**(-2), x)`

Maxima [F]

$$\int \frac{1}{(\sqrt{1-x} + \sqrt{1+x})^2} dx = \int \frac{1}{(\sqrt{x+1} + \sqrt{-x+1})^2} dx$$

input `integrate(1/((1-x)^(1/2)+(1+x)^(1/2))^2,x, algorithm="maxima")`

output `integrate((sqrt(x + 1) + sqrt(-x + 1))^-2, x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 147 vs. $2(24) = 48$.

Time = 0.14 (sec) , antiderivative size = 147, normalized size of antiderivative = 4.59

$$\int \frac{1}{(\sqrt{1-x} + \sqrt{1+x})^2} dx = \frac{1}{2} \pi + \frac{2 \left(\frac{\sqrt{2}-\sqrt{-x+1}}{\sqrt{x+1}} - \frac{\sqrt{x+1}}{\sqrt{2}-\sqrt{-x+1}} \right)}{\left(\frac{\sqrt{2}-\sqrt{-x+1}}{\sqrt{x+1}} - \frac{\sqrt{x+1}}{\sqrt{2}-\sqrt{-x+1}} \right)^2 - 4} - \frac{1}{2x} + \arctan \left(\frac{\sqrt{x+1} \left(\frac{(\sqrt{2}-\sqrt{-x+1})^2}{x+1} - 1 \right)}{2(\sqrt{2}-\sqrt{-x+1})} \right)$$

input `integrate(1/((1-x)^(1/2)+(1+x)^(1/2))^2,x, algorithm="giac")`

output `1/2*pi + 2*((sqrt(2) - sqrt(-x + 1))/sqrt(x + 1) - sqrt(x + 1)/(sqrt(2) - sqrt(-x + 1)))/(((sqrt(2) - sqrt(-x + 1))/sqrt(x + 1) - sqrt(x + 1)/(sqrt(2) - sqrt(-x + 1)))^2 - 4) - 1/2/x + arctan(1/2*sqrt(x + 1)*((sqrt(2) - sqrt(-x + 1))^2/(x + 1) - 1)/(sqrt(2) - sqrt(-x + 1)))`

Mupad [B] (verification not implemented)

Time = 0.53 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.53

$$\int \frac{1}{(\sqrt{1-x} + \sqrt{1+x})^2} dx = \frac{\left(\frac{x}{2} + \frac{1}{2}\right) \sqrt{1-x}}{x \sqrt{x+1}} - \frac{1}{2x} - 2 \operatorname{atan}\left(\frac{\sqrt{1-x}-1}{\sqrt{x+1}-1}\right)$$

input `int(1/((x + 1)^(1/2) + (1 - x)^(1/2))^2,x)`output `((x/2 + 1/2)*(1 - x)^(1/2))/(x*(x + 1)^(1/2)) - 1/(2*x) - 2*atan(((1 - x)^(1/2) - 1)/((x + 1)^(1/2) - 1))`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.84

$$\int \frac{1}{(\sqrt{1-x} + \sqrt{1+x})^2} dx = \frac{-\sqrt{1-x} \operatorname{asin}\left(\frac{\sqrt{1-x}}{\sqrt{2}}\right) - \sqrt{x+1} \operatorname{asin}\left(\frac{\sqrt{1-x}}{\sqrt{2}}\right) - \sqrt{x+1}}{\sqrt{1-x} + \sqrt{x+1}}$$

input `int(1/((1-x)^(1/2)+(1+x)^(1/2))^2,x)`output `(- (sqrt(- x + 1)*asin(sqrt(- x + 1)/sqrt(2)) + sqrt(x + 1)*asin(sqrt(- x + 1)/sqrt(2)) + sqrt(x + 1)))/(sqrt(- x + 1) + sqrt(x + 1))`

3.3 $\int \frac{1}{(1+\cos(x))^2} dx$

Optimal result	51
Mathematica [A] (verified)	51
Rubi [A] (verified)	52
Maple [A] (verified)	53
Fricas [A] (verification not implemented)	54
Sympy [A] (verification not implemented)	54
Maxima [A] (verification not implemented)	54
Giac [A] (verification not implemented)	55
Mupad [B] (verification not implemented)	55
Reduce [B] (verification not implemented)	55

Optimal result

Integrand size = 6, antiderivative size = 25

$$\int \frac{1}{(1+\cos(x))^2} dx = \frac{\sin(x)}{3(1+\cos(x))^2} + \frac{\sin(x)}{3(1+\cos(x))}$$

output `1/3*sin(x)/(1+cos(x))^2+1/3*sin(x)/(1+cos(x))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.64

$$\int \frac{1}{(1+\cos(x))^2} dx = \frac{(2+\cos(x))\sin(x)}{3(1+\cos(x))^2}$$

input `Integrate[(1 + Cos[x])^(-2),x]`

output `((2 + Cos[x])*Sin[x])/(3*(1 + Cos[x])^2)`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 3129, 3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(\cos(x) + 1)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(\sin(x + \frac{\pi}{2}) + 1)^2} dx \\
 & \quad \downarrow \text{3129} \\
 & \frac{1}{3} \int \frac{1}{\cos(x) + 1} dx + \frac{\sin(x)}{3(\cos(x) + 1)^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \int \frac{1}{\sin(x + \frac{\pi}{2}) + 1} dx + \frac{\sin(x)}{3(\cos(x) + 1)^2} \\
 & \quad \downarrow \text{3127} \\
 & \frac{\sin(x)}{3(\cos(x) + 1)} + \frac{\sin(x)}{3(\cos(x) + 1)^2}
 \end{aligned}$$

input `Int[(1 + Cos[x])^(-2), x]`

output `Sin[x]/(3*(1 + Cos[x])^2) + Sin[x]/(3*(1 + Cos[x]))`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] :=> Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3129 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :=> Simp[b*Cos[c + d*x]*((a + b*Sin[c + d*x])^n/(a*d*(2*n + 1))), x] + Simp[(n + 1)/(a*(2*n + 1)) Int[(a + b*Sin[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.64

method	result	size
default	$\frac{\tan(\frac{x}{2})^3}{6} + \frac{\tan(\frac{x}{2})}{2}$	16
norman	$\frac{\tan(\frac{x}{2})^3}{6} + \frac{\tan(\frac{x}{2})}{2}$	16
parallelrisch	$\frac{\tan(\frac{x}{2})^3}{6} + \frac{\tan(\frac{x}{2})}{2}$	16
risch	$\frac{2i(1+3e^{ix})}{3(1+e^{ix})^3}$	22

input `int(1/(1+cos(x))^2,x,method=_RETURNVERBOSE)`

output `1/6*tan(1/2*x)^3+1/2*tan(1/2*x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \frac{1}{(1 + \cos(x))^2} dx = \frac{(\cos(x) + 2) \sin(x)}{3 (\cos(x)^2 + 2 \cos(x) + 1)}$$

input `integrate(1/(1+cos(x))^2,x, algorithm="fricas")`

output `1/3*(cos(x) + 2)*sin(x)/(cos(x)^2 + 2*cos(x) + 1)`

Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.56

$$\int \frac{1}{(1 + \cos(x))^2} dx = \frac{\tan^3\left(\frac{x}{2}\right)}{6} + \frac{\tan\left(\frac{x}{2}\right)}{2}$$

input `integrate(1/(1+cos(x))**2,x)`

output `tan(x/2)**3/6 + tan(x/2)/2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \frac{1}{(1 + \cos(x))^2} dx = \frac{\sin(x)}{2(\cos(x) + 1)} + \frac{\sin(x)^3}{6(\cos(x) + 1)^3}$$

input `integrate(1/(1+cos(x))^2,x, algorithm="maxima")`

output `1/2*sin(x)/(cos(x) + 1) + 1/6*sin(x)^3/(cos(x) + 1)^3`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.60

$$\int \frac{1}{(1 + \cos(x))^2} dx = \frac{1}{6} \tan\left(\frac{1}{2}x\right)^3 + \frac{1}{2} \tan\left(\frac{1}{2}x\right)$$

input `integrate(1/(1+cos(x))^2,x, algorithm="giac")`

output `1/6*tan(1/2*x)^3 + 1/2*tan(1/2*x)`

Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.56

$$\int \frac{1}{(1 + \cos(x))^2} dx = \frac{\tan\left(\frac{x}{2}\right) \left(\tan\left(\frac{x}{2}\right)^2 + 3\right)}{6}$$

input `int(1/(cos(x) + 1)^2,x)`

output `(tan(x/2)*(tan(x/2)^2 + 3))/6`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.56

$$\int \frac{1}{(1 + \cos(x))^2} dx = \frac{\tan\left(\frac{x}{2}\right) \left(\tan\left(\frac{x}{2}\right)^2 + 3\right)}{6}$$

input `int(1/(1+cos(x))^2,x)`

output `(tan(x/2)*(tan(x/2)**2 + 3))/6`

3.4 $\int \frac{\sin(x)}{\sqrt{1+x}} dx$

Optimal result	56
Mathematica [C] (verified)	56
Rubi [A] (verified)	57
Maple [A] (verified)	59
Fricas [A] (verification not implemented)	59
Sympy [A] (verification not implemented)	59
Maxima [C] (verification not implemented)	60
Giac [C] (verification not implemented)	60
Mupad [F(-1)]	61
Reduce [F]	61

Optimal result

Integrand size = 10, antiderivative size = 58

$$\int \frac{\sin(x)}{\sqrt{1+x}} dx = \sqrt{2\pi} \cos(1) \operatorname{FresnelS}\left(\sqrt{\frac{2}{\pi}}\sqrt{1+x}\right) - \sqrt{2\pi} \operatorname{FresnelC}\left(\sqrt{\frac{2}{\pi}}\sqrt{1+x}\right) \sin(1)$$

output

```
cos(1)*FresnelS(2^(1/2)/Pi^(1/2)*(1+x)^(1/2))*2^(1/2)*Pi^(1/2)-FresnelC(2^(1/2)/Pi^(1/2)*(1+x)^(1/2))*sin(1)*2^(1/2)*Pi^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.01 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.17

$$\int \frac{\sin(x)}{\sqrt{1+x}} dx = -\frac{e^{-i}\left(\sqrt{-i(1+x)}\Gamma\left(\frac{1}{2}, -i(1+x)\right) + e^{2i}\sqrt{i(1+x)}\Gamma\left(\frac{1}{2}, i(1+x)\right)\right)}{2\sqrt{1+x}}$$

input

```
Integrate[Sin[x]/Sqrt[1 + x],x]
```

output

```
-1/2*(Sqrt[(-I)*(1 + x)]*Gamma[1/2, (-I)*(1 + x)] + E^(2*I)*Sqrt[I*(1 + x)]*Gamma[1/2, I*(1 + x)])/(E^I*Sqrt[1 + x])
```

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {3042, 3787, 3042, 3785, 3786, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin(x)}{\sqrt{x+1}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(x)}{\sqrt{x+1}} dx \\
 & \quad \downarrow \text{3787} \\
 & \cos(1) \int \frac{\sin(x+1)}{\sqrt{x+1}} dx - \sin(1) \int \frac{\cos(x+1)}{\sqrt{x+1}} dx \\
 & \quad \downarrow \text{3042} \\
 & \cos(1) \int \frac{\sin(x+1)}{\sqrt{x+1}} dx - \sin(1) \int \frac{\sin(x + \frac{\pi}{2} + 1)}{\sqrt{x+1}} dx \\
 & \quad \downarrow \text{3785} \\
 & \cos(1) \int \frac{\sin(x+1)}{\sqrt{x+1}} dx - 2 \sin(1) \int \cos(x+1) d\sqrt{x+1} \\
 & \quad \downarrow \text{3786} \\
 & 2 \cos(1) \int \sin(x+1) d\sqrt{x+1} - 2 \sin(1) \int \cos(x+1) d\sqrt{x+1} \\
 & \quad \downarrow \text{3832} \\
 & \sqrt{2\pi} \cos(1) \text{FresnelS} \left(\sqrt{\frac{2}{\pi}} \sqrt{x+1} \right) - 2 \sin(1) \int \cos(x+1) d\sqrt{x+1} \\
 & \quad \downarrow \text{3833}
 \end{aligned}$$

$$\sqrt{2\pi} \cos(1) \operatorname{FresnelS} \left(\sqrt{\frac{2}{\pi}} \sqrt{x+1} \right) - \sqrt{2\pi} \sin(1) \operatorname{FresnelC} \left(\sqrt{\frac{2}{\pi}} \sqrt{x+1} \right)$$

input `Int[Sin[x]/Sqrt[1 + x],x]`

output `Sqrt[2*Pi]*Cos[1]*FresnelS[Sqrt[2/Pi]*Sqrt[1 + x]] - Sqrt[2*Pi]*FresnelC[Sqrt[2/Pi]*Sqrt[1 + x]]*Sin[1]`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3785 `Int[sin[Pi/2 + (e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[2/d Subst[Int[Cos[f*(x^2/d)], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]`

rule 3786 `Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[2/d Subst[Int[Sin[f*(x^2/d)], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]`

rule 3787 `Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[Cos[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/Sqrt[c + d*x], x], x] + Simp[Sin[(d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/Sqrt[c + d*x], x], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && NeQ[d*e - c*f, 0]`

rule 3832 `Int[Sin[(d_.)*((e_.) + (f_.)*(x_))^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3833 `Int[Cos[(d_.)*((e_.) + (f_.)*(x_))^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.72

method	result	size
derivativedivides	$\sqrt{2} \sqrt{\pi} \left(\cos(1) \operatorname{FresnelS} \left(\frac{\sqrt{2} \sqrt{1+x}}{\sqrt{\pi}} \right) - \sin(1) \operatorname{FresnelC} \left(\frac{\sqrt{2} \sqrt{1+x}}{\sqrt{\pi}} \right) \right)$	42
default	$\sqrt{2} \sqrt{\pi} \left(\cos(1) \operatorname{FresnelS} \left(\frac{\sqrt{2} \sqrt{1+x}}{\sqrt{\pi}} \right) - \sin(1) \operatorname{FresnelC} \left(\frac{\sqrt{2} \sqrt{1+x}}{\sqrt{\pi}} \right) \right)$	42

input `int(sin(x)/(1+x)^(1/2),x,method=_RETURNVERBOSE)`

output `2^(1/2)*Pi^(1/2)*(cos(1)*FresnelS(2^(1/2)/Pi^(1/2)*(1+x)^(1/2))-sin(1)*FresnelC(2^(1/2)/Pi^(1/2)*(1+x)^(1/2)))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.79

$$\int \frac{\sin(x)}{\sqrt{1+x}} dx = \sqrt{2} \sqrt{\pi} \cos(1) S \left(\frac{\sqrt{2} \sqrt{x+1}}{\sqrt{\pi}} \right) - \sqrt{2} \sqrt{\pi} C \left(\frac{\sqrt{2} \sqrt{x+1}}{\sqrt{\pi}} \right) \sin(1)$$

input `integrate(sin(x)/(1+x)^(1/2),x, algorithm="fricas")`

output `sqrt(2)*sqrt(pi)*cos(1)*fresnel_sin(sqrt(2)*sqrt(x + 1)/sqrt(pi)) - sqrt(2)*sqrt(pi)*fresnel_cos(sqrt(2)*sqrt(x + 1)/sqrt(pi))*sin(1)`

Sympy [A] (verification not implemented)

Time = 0.82 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.91

$$\int \frac{\sin(x)}{\sqrt{1+x}} dx = \sqrt{2} \sqrt{\pi} \left(-\sin(1) C \left(\frac{\sqrt{2} \sqrt{x+1}}{\sqrt{\pi}} \right) + \cos(1) S \left(\frac{\sqrt{2} \sqrt{x+1}}{\sqrt{\pi}} \right) \right)$$

input `integrate(sin(x)/(1+x)**(1/2),x)`

output

```
sqrt(2)*sqrt(pi)*(-sin(1)*fresnelc(sqrt(2)*sqrt(x + 1)/sqrt(pi)) + cos(1)*
fresnels(sqrt(2)*sqrt(x + 1)/sqrt(pi)))
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.93

$$\int \frac{\sin(x)}{\sqrt{1+x}} dx$$

$$= \frac{1}{8} \sqrt{\pi} \left(\left((i+1) \sqrt{2} \cos(1) + (i-1) \sqrt{2} \sin(1) \right) \operatorname{erf} \left(\left(\frac{1}{2}i + \frac{1}{2} \right) \sqrt{2} \sqrt{x+1} \right) + \left((i-1) \sqrt{2} \cos(1) + (i+1) \sqrt{2} \sin(1) \right) \operatorname{erf} \left(\left(\frac{1}{2}i - \frac{1}{2} \right) \sqrt{2} \sqrt{x+1} \right) \right)$$

input

```
integrate(sin(x)/(1+x)^(1/2),x, algorithm="maxima")
```

output

```
1/8*sqrt(pi)*(((I + 1)*sqrt(2)*cos(1) + (I - 1)*sqrt(2)*sin(1))*erf((1/2*I
+ 1/2)*sqrt(2)*sqrt(x + 1)) + ((I - 1)*sqrt(2)*cos(1) + (I + 1)*sqrt(2)*s
in(1))*erf((1/2*I - 1/2)*sqrt(2)*sqrt(x + 1)) + (-(I - 1)*sqrt(2)*cos(1) -
(I + 1)*sqrt(2)*sin(1))*erf(sqrt(-I)*sqrt(x + 1)) + ((I + 1)*sqrt(2)*cos(
1) + (I - 1)*sqrt(2)*sin(1))*erf((-1)^(1/4)*sqrt(x + 1)))
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.74

$$\int \frac{\sin(x)}{\sqrt{1+x}} dx = -\left(\frac{1}{4}i + \frac{1}{4}\right) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(-\left(\frac{1}{2}i + \frac{1}{2}\right) \sqrt{2} \sqrt{x+1} \right) e^i$$

$$+ \left(\frac{1}{4}i - \frac{1}{4}\right) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{x+1} \right) e^{-i}$$

input

```
integrate(sin(x)/(1+x)^(1/2),x, algorithm="giac")
```

output

```
-(1/4*I + 1/4)*sqrt(2)*sqrt(pi)*erf(-(1/2*I + 1/2)*sqrt(2)*sqrt(x + 1))*e^
I + (1/4*I - 1/4)*sqrt(2)*sqrt(pi)*erf((1/2*I - 1/2)*sqrt(2)*sqrt(x + 1))*
e^(-I)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\sin(x)}{\sqrt{1+x}} dx = \int \frac{\sin(x)}{\sqrt{x+1}} dx$$

input

```
int(sin(x)/(x + 1)^(1/2),x)
```

output

```
int(sin(x)/(x + 1)^(1/2), x)
```

Reduce [F]

$$\int \frac{\sin(x)}{\sqrt{1+x}} dx = \int \frac{\sin(x)}{\sqrt{x+1}} dx$$

input

```
int(sin(x)/(1+x)^(1/2),x)
```

output

```
int(sin(x)/sqrt(x + 1),x)
```

3.5 $\int \frac{1}{(\cos(x)+\sin(x))^6} dx$

Optimal result	62
Mathematica [A] (verified)	62
Rubi [A] (verified)	63
Maple [C] (verified)	64
Fricas [A] (verification not implemented)	65
Sympy [B] (verification not implemented)	65
Maxima [A] (verification not implemented)	66
Giac [A] (verification not implemented)	67
Mupad [B] (verification not implemented)	67
Reduce [B] (verification not implemented)	68

Optimal result

Integrand size = 7, antiderivative size = 50

$$\int \frac{1}{(\cos(x) + \sin(x))^6} dx = -\frac{\cos(x) - \sin(x)}{10(\cos(x) + \sin(x))^5} - \frac{\cos(x) - \sin(x)}{15(\cos(x) + \sin(x))^3} + \frac{2 \sin(x)}{15(\cos(x) + \sin(x))}$$

output

```
1/10*(-cos(x)+sin(x))/(cos(x)+sin(x))^5+1/15*(-cos(x)+sin(x))/(cos(x)+sin(x))^3+2/15*sin(x)/(cos(x)+sin(x))
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.52

$$\int \frac{1}{(\cos(x) + \sin(x))^6} dx = -\frac{5 \cos(3x) - 10 \sin(x) + \sin(5x)}{30(\cos(x) + \sin(x))^5}$$

input

```
Integrate[(Cos[x] + Sin[x])^(-6),x]
```

output

```
-1/30*(5*Cos[3*x] - 10*Sin[x] + Sin[5*x])/(Cos[x] + Sin[x])^5
```

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.10, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.857$, Rules used = {3042, 3555, 3042, 3555, 3042, 3554}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(\sin(x) + \cos(x))^6} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(\sin(x) + \cos(x))^6} dx \\
 & \quad \downarrow \text{3555} \\
 & \frac{2}{5} \int \frac{1}{(\cos(x) + \sin(x))^4} dx - \frac{\cos(x) - \sin(x)}{10(\sin(x) + \cos(x))^5} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2}{5} \int \frac{1}{(\cos(x) + \sin(x))^4} dx - \frac{\cos(x) - \sin(x)}{10(\sin(x) + \cos(x))^5} \\
 & \quad \downarrow \text{3555} \\
 & \frac{2}{5} \left(\frac{1}{3} \int \frac{1}{(\cos(x) + \sin(x))^2} dx - \frac{\cos(x) - \sin(x)}{6(\sin(x) + \cos(x))^3} \right) - \frac{\cos(x) - \sin(x)}{10(\sin(x) + \cos(x))^5} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2}{5} \left(\frac{1}{3} \int \frac{1}{(\cos(x) + \sin(x))^2} dx - \frac{\cos(x) - \sin(x)}{6(\sin(x) + \cos(x))^3} \right) - \frac{\cos(x) - \sin(x)}{10(\sin(x) + \cos(x))^5} \\
 & \quad \downarrow \text{3554} \\
 & \frac{2}{5} \left(\frac{\sin(x)}{3(\sin(x) + \cos(x))} - \frac{\cos(x) - \sin(x)}{6(\sin(x) + \cos(x))^3} \right) - \frac{\cos(x) - \sin(x)}{10(\sin(x) + \cos(x))^5}
 \end{aligned}$$

input `Int[(Cos[x] + Sin[x])^(-6),x]`


```
output -1/10*(Cos[x] - Sin[x])/(Cos[x] + Sin[x])^5 + (2*(-1/6*(Cos[x] - Sin[x])/(Cos[x] + Sin[x])^3 + Sin[x]/(3*(Cos[x] + Sin[x]))))/5
```

Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3554 Int[(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-2), x_Symbol] := Simp[Sin[c + d*x]/(a*d*(a*cos[c + d*x] + b*sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]
```

```
rule 3555 Int[(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(b*cos[c + d*x] - a*sin[c + d*x])*((a*cos[c + d*x] + b*sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 + b^2))), x] + Simp[(n + 2)/((n + 1)*(a^2 + b^2)) Int[(a*cos[c + d*x] + b*sin[c + d*x])^(n + 2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0] && LtQ[n, -1] && NeQ[n, -2]
```

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.37 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.60

method	result	size
risch	$\frac{-\frac{2}{15} + \frac{4e^{4ix}}{3} + \frac{2ie^{2ix}}{3}}{(e^{2ix} + i)^5}$	30
default	$-\frac{4}{5(\tan(x)+1)^5} + \frac{2}{(\tan(x)+1)^4} - \frac{1}{\tan(x)+1} + \frac{2}{(\tan(x)+1)^2} - \frac{8}{3(\tan(x)+1)^3}$	42
norman	$\frac{-8 \tan\left(\frac{x}{2}\right)^2 - 2 \tan\left(\frac{x}{2}\right) - 2 \tan\left(\frac{x}{2}\right)^9 + 8 \tan\left(\frac{x}{2}\right)^8 - \frac{40 \tan\left(\frac{x}{2}\right)^3}{3} - \frac{40 \tan\left(\frac{x}{2}\right)^7}{3} - \frac{8 \tan\left(\frac{x}{2}\right)^6}{3} + \frac{8 \tan\left(\frac{x}{2}\right)^4}{3} + \frac{236 \tan\left(\frac{x}{2}\right)^5}{15}}{\left(\tan\left(\frac{x}{2}\right)^2 - 2 \tan\left(\frac{x}{2}\right) - 1\right)^5}$	89
parallelrisch	$\frac{-8 \tan\left(\frac{x}{2}\right)^2 - 2 \tan\left(\frac{x}{2}\right) - 2 \tan\left(\frac{x}{2}\right)^9 + 8 \tan\left(\frac{x}{2}\right)^8 - \frac{40 \tan\left(\frac{x}{2}\right)^3}{3} - \frac{40 \tan\left(\frac{x}{2}\right)^7}{3} - \frac{8 \tan\left(\frac{x}{2}\right)^6}{3} + \frac{8 \tan\left(\frac{x}{2}\right)^4}{3} + \frac{236 \tan\left(\frac{x}{2}\right)^5}{15}}{\left(\tan\left(\frac{x}{2}\right)^2 - 2 \tan\left(\frac{x}{2}\right) - 1\right)^5}$	90

input `int(1/(cos(x)+sin(x))^6,x,method=_RETURNVERBOSE)`

output `2/15*(-1+10*exp(4*I*x)+5*I*exp(2*I*x))/(exp(2*I*x)+I)^5`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.34

$$\int \frac{1}{(\cos(x) + \sin(x))^6} dx$$

$$= -\frac{8 \cos(x)^5 - 20 \cos(x)^3 - (8 \cos(x)^4 + 4 \cos(x)^2 - 7) \sin(x) + 5 \cos(x)}{30(4 \cos(x)^5 + (4 \cos(x)^4 - 8 \cos(x)^2 - 1) \sin(x) - 5 \cos(x))}$$

input `integrate(1/(cos(x)+sin(x))^6,x, algorithm="fricas")`

output `-1/30*(8*cos(x)^5 - 20*cos(x)^3 - (8*cos(x)^4 + 4*cos(x)^2 - 7)*sin(x) + 5*cos(x))/(4*cos(x)^5 + (4*cos(x)^4 - 8*cos(x)^2 - 1)*sin(x) - 5*cos(x))`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 838 vs. $2(51) = 102$.

Time = 2.35 (sec) , antiderivative size = 838, normalized size of antiderivative = 16.76

$$\int \frac{1}{(\cos(x) + \sin(x))^6} dx = \text{Too large to display}$$

input `integrate(1/(cos(x)+sin(x))**6,x)`

output

```

-30*tan(x/2)**9/(15*tan(x/2)**10 - 150*tan(x/2)**9 + 525*tan(x/2)**8 - 600
*tan(x/2)**7 - 450*tan(x/2)**6 + 1020*tan(x/2)**5 + 450*tan(x/2)**4 - 600*
tan(x/2)**3 - 525*tan(x/2)**2 - 150*tan(x/2) - 15) + 120*tan(x/2)**8/(15*t
an(x/2)**10 - 150*tan(x/2)**9 + 525*tan(x/2)**8 - 600*tan(x/2)**7 - 450*ta
n(x/2)**6 + 1020*tan(x/2)**5 + 450*tan(x/2)**4 - 600*tan(x/2)**3 - 525*tan
(x/2)**2 - 150*tan(x/2) - 15) - 200*tan(x/2)**7/(15*tan(x/2)**10 - 150*tan
(x/2)**9 + 525*tan(x/2)**8 - 600*tan(x/2)**7 - 450*tan(x/2)**6 + 1020*tan(
x/2)**5 + 450*tan(x/2)**4 - 600*tan(x/2)**3 - 525*tan(x/2)**2 - 150*tan(x/
2) - 15) - 40*tan(x/2)**6/(15*tan(x/2)**10 - 150*tan(x/2)**9 + 525*tan(x/2
)**8 - 600*tan(x/2)**7 - 450*tan(x/2)**6 + 1020*tan(x/2)**5 + 450*tan(x/2
)**4 - 600*tan(x/2)**3 - 525*tan(x/2)**2 - 150*tan(x/2) - 15) + 236*tan(x/2
)**5/(15*tan(x/2)**10 - 150*tan(x/2)**9 + 525*tan(x/2)**8 - 600*tan(x/2)**
7 - 450*tan(x/2)**6 + 1020*tan(x/2)**5 + 450*tan(x/2)**4 - 600*tan(x/2)**3
- 525*tan(x/2)**2 - 150*tan(x/2) - 15) + 40*tan(x/2)**4/(15*tan(x/2)**10
- 150*tan(x/2)**9 + 525*tan(x/2)**8 - 600*tan(x/2)**7 - 450*tan(x/2)**6 +
1020*tan(x/2)**5 + 450*tan(x/2)**4 - 600*tan(x/2)**3 - 525*tan(x/2)**2 - 1
50*tan(x/2) - 15) - 200*tan(x/2)**3/(15*tan(x/2)**10 - 150*tan(x/2)**9 + 5
25*tan(x/2)**8 - 600*tan(x/2)**7 - 450*tan(x/2)**6 + 1020*tan(x/2)**5 + 45
0*tan(x/2)**4 - 600*tan(x/2)**3 - 525*tan(x/2)**2 - 150*tan(x/2) - 15) - 1
20*tan(x/2)**2/(15*tan(x/2)**10 - 150*tan(x/2)**9 + 525*tan(x/2)**8 - 6...

```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.12

$$\int \frac{1}{(\cos(x) + \sin(x))^6} dx$$

$$= -\frac{15 \tan(x)^4 + 30 \tan(x)^3 + 40 \tan(x)^2 + 20 \tan(x) + 7}{15 (\tan(x)^5 + 5 \tan(x)^4 + 10 \tan(x)^3 + 10 \tan(x)^2 + 5 \tan(x) + 1)}$$

input

```
integrate(1/(cos(x)+sin(x))^6,x, algorithm="maxima")
```

output

```

-1/15*(15*tan(x)^4 + 30*tan(x)^3 + 40*tan(x)^2 + 20*tan(x) + 7)/(tan(x)^5
+ 5*tan(x)^4 + 10*tan(x)^3 + 10*tan(x)^2 + 5*tan(x) + 1)

```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.64

$$\int \frac{1}{(\cos(x) + \sin(x))^6} dx = -\frac{15 \tan(x)^4 + 30 \tan(x)^3 + 40 \tan(x)^2 + 20 \tan(x) + 7}{15 (\tan(x) + 1)^5}$$

input `integrate(1/(cos(x)+sin(x))^6,x, algorithm="giac")`

output `-1/15*(15*tan(x)^4 + 30*tan(x)^3 + 40*tan(x)^2 + 20*tan(x) + 7)/(tan(x) + 1)^5`

Mupad [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.76

$$\int \frac{1}{(\cos(x) + \sin(x))^6} dx = \frac{2 \tan\left(\frac{x}{2}\right) \left(15 \tan\left(\frac{x}{2}\right)^8 - 60 \tan\left(\frac{x}{2}\right)^7 + 100 \tan\left(\frac{x}{2}\right)^6 + 20 \tan\left(\frac{x}{2}\right)^5 - 118 \tan\left(\frac{x}{2}\right)^4 - 20 \tan\left(\frac{x}{2}\right)^3 + 100 \tan\left(\frac{x}{2}\right)^2 - 15 \tan\left(\frac{x}{2}\right) + 15\right)}{15 \left(-\tan\left(\frac{x}{2}\right)^2 + 2 \tan\left(\frac{x}{2}\right) + 1\right)^5}$$

input `int(1/(cos(x) + sin(x))^6,x)`

output `(2*tan(x/2)*(60*tan(x/2) + 100*tan(x/2)^2 - 20*tan(x/2)^3 - 118*tan(x/2)^4 + 20*tan(x/2)^5 + 100*tan(x/2)^6 - 60*tan(x/2)^7 + 15*tan(x/2)^8 + 15))/(15*(2*tan(x/2) - tan(x/2)^2 + 1)^5)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.34

$$\int \frac{1}{(\cos(x) + \sin(x))^6} dx$$

$$= \frac{-2 \cos(x) \sin(x)^4 - 6 \cos(x) \sin(x)^2 + 3 \cos(x) + 6 \sin(x)^5 - 10 \sin(x)^3}{60 \cos(x) \sin(x)^4 - 120 \cos(x) \sin(x)^2 - 15 \cos(x) + 60 \sin(x)^5 - 75 \sin(x)}$$

input `int(1/(cos(x)+sin(x))^6,x)`output `(- 2*cos(x)*sin(x)**4 - 6*cos(x)*sin(x)**2 + 3*cos(x) + 6*sin(x)**5 - 10*
sin(x)**3)/(15*(4*cos(x)*sin(x)**4 - 8*cos(x)*sin(x)**2 - cos(x) + 4*sin(x)
)**5 - 5*sin(x))`

3.6 $\int \log\left(\frac{1}{x^4} + x^4\right) dx$

Optimal result	69
Mathematica [C] (verified)	70
Rubi [A] (verified)	70
Maple [C] (verified)	74
Fricas [C] (verification not implemented)	75
Sympy [A] (verification not implemented)	75
Maxima [F]	76
Giac [A] (verification not implemented)	76
Mupad [B] (verification not implemented)	77
Reduce [B] (verification not implemented)	78

Optimal result

Integrand size = 8, antiderivative size = 334

$$\begin{aligned}
 \int \log\left(\frac{1}{x^4} + x^4\right) dx = & -4x - \sqrt{2 + \sqrt{2}} \arctan\left(\frac{\sqrt{2 - \sqrt{2}} - 2x}{\sqrt{2 + \sqrt{2}}}\right) \\
 & - \sqrt{2 - \sqrt{2}} \arctan\left(\frac{\sqrt{2 + \sqrt{2}} - 2x}{\sqrt{2 - \sqrt{2}}}\right) \\
 & + \sqrt{2 + \sqrt{2}} \arctan\left(\frac{\sqrt{2 - \sqrt{2}} + 2x}{\sqrt{2 + \sqrt{2}}}\right) \\
 & + \sqrt{2 - \sqrt{2}} \arctan\left(\frac{\sqrt{2 + \sqrt{2}} + 2x}{\sqrt{2 - \sqrt{2}}}\right) \\
 & - \frac{1}{2} \sqrt{2 - \sqrt{2}} \log\left(1 - \sqrt{2 - \sqrt{2}}x + x^2\right) \\
 & + \frac{1}{2} \sqrt{2 - \sqrt{2}} \log\left(1 + \sqrt{2 - \sqrt{2}}x + x^2\right) \\
 & - \frac{1}{2} \sqrt{2 + \sqrt{2}} \log\left(1 - \sqrt{2 + \sqrt{2}}x + x^2\right) \\
 & + \frac{1}{2} \sqrt{2 + \sqrt{2}} \log\left(1 + \sqrt{2 + \sqrt{2}}x + x^2\right) + x \log\left(\frac{1}{x^4} + x^4\right)
 \end{aligned}$$

output

$$\begin{aligned}
& -4*x+x*\ln(1/x^4+x^4)-\arctan((-2*x+(2+2^{(1/2)})^{(1/2)})/(2-2^{(1/2)})^{(1/2)})*(2 \\
& -2^{(1/2)})^{(1/2)}+\arctan((2*x+(2+2^{(1/2)})^{(1/2)})/(2-2^{(1/2)})^{(1/2)})*(2-2^{(1/2)} \\
& 2)^{(1/2)}-1/2*\ln(1+x^2-x*(2-2^{(1/2)})^{(1/2)})*(2-2^{(1/2)})^{(1/2)}+1/2*\ln(1+x^2 \\
& +x*(2-2^{(1/2)})^{(1/2)})*(2-2^{(1/2)})^{(1/2)}-\arctan((-2*x+(2-2^{(1/2)})^{(1/2)})/(2 \\
& +2^{(1/2)})^{(1/2)})*(2+2^{(1/2)})^{(1/2)}+\arctan((2*x+(2-2^{(1/2)})^{(1/2)})/(2+2^{(1/2)} \\
& 2)^{(1/2)})*(2+2^{(1/2)})^{(1/2)}-1/2*\ln(1+x^2-x*(2+2^{(1/2)})^{(1/2)})*(2+2^{(1/2)}) \\
& ^{(1/2)}+1/2*\ln(1+x^2+x*(2+2^{(1/2)})^{(1/2)})*(2+2^{(1/2)})^{(1/2)}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.00 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.09

$$\int \log\left(\frac{1}{x^4} + x^4\right) dx = -4x + 8x \operatorname{Hypergeometric2F1}\left(\frac{1}{8}, 1, \frac{9}{8}, -x^8\right) + x \log\left(\frac{1}{x^4} + x^4\right)$$

input

`Integrate[Log[x^(-4) + x^4],x]`

output

`-4*x + 8*x*Hypergeometric2F1[1/8, 1, 9/8, -x^8] + x*Log[x^(-4) + x^4]`

Rubi [A] (verified)

Time = 0.61 (sec) , antiderivative size = 362, normalized size of antiderivative = 1.08, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$, Rules used = {3003, 27, 913, 757, 1483, 1142, 25, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \log\left(x^4 + \frac{1}{x^4}\right) dx \\
& \quad \downarrow \text{3003} \\
& x \log\left(x^4 + \frac{1}{x^4}\right) - \int -\frac{4(1-x^8)}{x^8+1} dx
\end{aligned}$$

$$\begin{aligned}
& \downarrow 27 \\
& 4 \int \frac{1-x^8}{x^8+1} dx + x \log \left(x^4 + \frac{1}{x^4} \right) \\
& \downarrow 913 \\
& 4 \left(2 \int \frac{1}{x^8+1} dx - x \right) + x \log \left(x^4 + \frac{1}{x^4} \right) \\
& \downarrow 757 \\
& 4 \left(2 \left(\frac{\int \frac{\sqrt{2-x^2}}{x^4-\sqrt{2}x^2+1} dx}{2\sqrt{2}} + \frac{\int \frac{x^2+\sqrt{2}}{x^4+\sqrt{2}x^2+1} dx}{2\sqrt{2}} \right) - x \right) + x \log \left(x^4 + \frac{1}{x^4} \right) \\
& \downarrow 1483 \\
& 4 \left(2 \left(\frac{\int \frac{(1-\sqrt{2})x+\sqrt{2(2-\sqrt{2})}}{x^2-\sqrt{2-\sqrt{2}}x+1} dx}{2\sqrt{2-\sqrt{2}}} + \frac{\int \frac{\sqrt{2(2-\sqrt{2})}-(1-\sqrt{2})x}{x^2+\sqrt{2-\sqrt{2}}x+1} dx}{2\sqrt{2-\sqrt{2}}} + \frac{\int \frac{\sqrt{2(2+\sqrt{2})}-(1+\sqrt{2})x}{x^2-\sqrt{2+\sqrt{2}}x+1} dx}{2\sqrt{2+\sqrt{2}}} + \frac{\int \frac{(1+\sqrt{2})x+\sqrt{2(2+\sqrt{2})}}{x^2+\sqrt{2+\sqrt{2}}x+1} dx}{2\sqrt{2+\sqrt{2}}} \right) - x \right) + \\
& \quad x \log \left(x^4 + \frac{1}{x^4} \right) \\
& \downarrow 1142 \\
& 4 \left(2 \left(\frac{\frac{\frac{1}{2}\sqrt{2+\sqrt{2}} \int \frac{1}{x^2-\sqrt{2-\sqrt{2}}x+1} dx + \frac{1}{2}(1-\sqrt{2}) \int \frac{-\sqrt{2-\sqrt{2}}-2x}{x^2-\sqrt{2-\sqrt{2}}x+1} dx}{2\sqrt{2-\sqrt{2}}} + \frac{\frac{1}{2}\sqrt{2+\sqrt{2}} \int \frac{1}{x^2+\sqrt{2-\sqrt{2}}x+1} dx - \frac{1}{2}(1-\sqrt{2}) \int \frac{2x+\sqrt{2-\sqrt{2}}}{x^2+\sqrt{2-\sqrt{2}}x+1} dx}{2\sqrt{2-\sqrt{2}}}} \right) + \right. \\
& \quad \left. x \log \left(x^4 + \frac{1}{x^4} \right) \right) \\
& \downarrow 25 \\
& 4 \left(2 \left(\frac{\frac{\frac{1}{2}\sqrt{2+\sqrt{2}} \int \frac{1}{x^2-\sqrt{2-\sqrt{2}}x+1} dx - \frac{1}{2}(1-\sqrt{2}) \int \frac{\sqrt{2-\sqrt{2}}-2x}{x^2-\sqrt{2-\sqrt{2}}x+1} dx}{2\sqrt{2-\sqrt{2}}} + \frac{\frac{1}{2}\sqrt{2+\sqrt{2}} \int \frac{1}{x^2+\sqrt{2-\sqrt{2}}x+1} dx - \frac{1}{2}(1-\sqrt{2}) \int \frac{2x+\sqrt{2-\sqrt{2}}}{x^2+\sqrt{2-\sqrt{2}}x+1} dx}{2\sqrt{2-\sqrt{2}}}} \right) + \right. \\
& \quad \left. x \log \left(x^4 + \frac{1}{x^4} \right) \right) \\
& \downarrow 1083
\end{aligned}$$

$$4 \left(2 \left(\frac{-\frac{1}{2}(1-\sqrt{2}) \int \frac{\sqrt{2-\sqrt{2}}-2x}{x^2-\sqrt{2-\sqrt{2}}x+1} dx - \sqrt{2+\sqrt{2}} \int \frac{1}{-(2x-\sqrt{2-\sqrt{2}})^2-\sqrt{2}-2} d(2x-\sqrt{2-\sqrt{2}})}{2\sqrt{2-\sqrt{2}}} + \frac{-\frac{1}{2}(1-\sqrt{2}) \int \frac{2x+\sqrt{2-\sqrt{2}}}{x^2+\sqrt{2-\sqrt{2}}x+1} dx - \sqrt{2+\sqrt{2}} \int \frac{1}{(2x+\sqrt{2-\sqrt{2}})^2-\sqrt{2}-2} d(2x+\sqrt{2-\sqrt{2}})}{2\sqrt{2-\sqrt{2}}} \right) \right)$$

$$x \log \left(x^4 + \frac{1}{x^4} \right)$$

↓ 217

$$4 \left(2 \left(\frac{\frac{\arctan\left(\frac{2x-\sqrt{2-\sqrt{2}}}{\sqrt{2+\sqrt{2}}}\right) - \frac{1}{2}(1-\sqrt{2}) \int \frac{\sqrt{2-\sqrt{2}}-2x}{x^2-\sqrt{2-\sqrt{2}}x+1} dx}{2\sqrt{2-\sqrt{2}}} + \frac{\arctan\left(\frac{2x+\sqrt{2-\sqrt{2}}}{\sqrt{2+\sqrt{2}}}\right) - \frac{1}{2}(1-\sqrt{2}) \int \frac{2x+\sqrt{2-\sqrt{2}}}{x^2+\sqrt{2-\sqrt{2}}x+1} dx}{2\sqrt{2-\sqrt{2}}} + \frac{\frac{1}{2}(1+\sqrt{2}) \int \frac{\sqrt{2-\sqrt{2}}}{x^2-\sqrt{2-\sqrt{2}}x+1} dx}{2\sqrt{2-\sqrt{2}}} \right) \right)$$

$$x \log \left(x^4 + \frac{1}{x^4} \right)$$

↓ 1103

$$4 \left(2 \left(\frac{\frac{\arctan\left(\frac{2x-\sqrt{2-\sqrt{2}}}{\sqrt{2+\sqrt{2}}}\right) + \frac{1}{2}(1-\sqrt{2}) \log(x^2-\sqrt{2-\sqrt{2}}x+1)}{2\sqrt{2-\sqrt{2}}} + \frac{\arctan\left(\frac{2x+\sqrt{2-\sqrt{2}}}{\sqrt{2+\sqrt{2}}}\right) - \frac{1}{2}(1-\sqrt{2}) \log(x^2+\sqrt{2-\sqrt{2}}x+1)}{2\sqrt{2-\sqrt{2}}} + \frac{\arctan\left(\frac{2x-\sqrt{2-\sqrt{2}}}{\sqrt{2+\sqrt{2}}}\right) + \frac{1}{2}(1+\sqrt{2}) \log(x^2-\sqrt{2-\sqrt{2}}x+1)}{2\sqrt{2-\sqrt{2}}} \right) \right)$$

$$x \log \left(x^4 + \frac{1}{x^4} \right)$$

input `Int [Log[x^(-4) + x^4] , x]`

output `4*(-x + 2*((ArcTan[(-Sqrt[2 - Sqrt[2]] + 2*x)/Sqrt[2 + Sqrt[2]]] + ((1 - Sqrt[2])*Log[1 - Sqrt[2 - Sqrt[2]]*x + x^2])/2)/(2*Sqrt[2 - Sqrt[2]]) + (ArcTan[(Sqrt[2 - Sqrt[2]] + 2*x)/Sqrt[2 + Sqrt[2]]] - ((1 - Sqrt[2])*Log[1 + Sqrt[2 - Sqrt[2]]*x + x^2])/2)/(2*Sqrt[2 - Sqrt[2]]))/(2*Sqrt[2]) + ((ArcTan[(-Sqrt[2 + Sqrt[2]] + 2*x)/Sqrt[2 - Sqrt[2]]] - ((1 + Sqrt[2])*Log[1 - Sqrt[2 + Sqrt[2]]*x + x^2])/2)/(2*Sqrt[2 + Sqrt[2]]) + (ArcTan[(Sqrt[2 + Sqrt[2]] + 2*x)/Sqrt[2 - Sqrt[2]]] + ((1 + Sqrt[2])*Log[1 + Sqrt[2 + Sqrt[2]]*x + x^2])/2)/(2*Sqrt[2 + Sqrt[2]]))/(2*Sqrt[2])) + x*Log[x^(-4) + x^4]`

Defintions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27 $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ ; FreeQ}[\text{b}, \text{x}]$
- rule 217 $\text{Int}[(\text{a}_) + (\text{b}_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(-(\text{Rt}[-\text{a}, 2]*\text{Rt}[-\text{b}, 2])^{-1})*\text{ArcTan}[\text{Rt}[-\text{b}, 2]*(\text{x}/\text{Rt}[-\text{a}, 2])], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ \text{PosQ}[\text{a}/\text{b}] \ \&\& \ (\text{LtQ}[\text{a}, 0] \ || \ \text{LtQ}[\text{b}, 0])$
- rule 757 $\text{Int}[(\text{a}_) + (\text{b}_.)*(x_)^{(n_)}])^{-1}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{r} = \text{Numerator}[\text{Rt}[\text{a}/\text{b}, 4]], \text{s} = \text{Denominator}[\text{Rt}[\text{a}/\text{b}, 4]]\}, \text{Simp}[\text{r}/(2*\text{Sqrt}[2]*\text{a}) \quad \text{Int}[(\text{Sqrt}[2]*\text{r} - \text{s}*x^{(n/4)})/(\text{r}^2 - \text{Sqrt}[2]*\text{r}*x^{(n/4)} + \text{s}^2*x^{(n/2)}), \text{x}], \text{x}] + \text{Simp}[\text{r}/(2*\text{Sqrt}[2]*\text{a}) \quad \text{Int}[(\text{Sqrt}[2]*\text{r} + \text{s}*x^{(n/4)})/(\text{r}^2 + \text{Sqrt}[2]*\text{r}*x^{(n/4)} + \text{s}^2*x^{(n/2)}), \text{x}], \text{x}]] \text{ ; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ \text{IGtQ}[\text{n}/4, 1] \ \&\& \ \text{GtQ}[\text{a}/\text{b}, 0]$
- rule 913 $\text{Int}[(\text{a}_) + (\text{b}_.)*(x_)^{(n_)}])^{(p_)}*((\text{c}_) + (\text{d}_.)*(x_)^{(n_)}), \text{x_Symbol}] \rightarrow \text{Simp}[\text{d}*x*((\text{a} + \text{b}*x^n)^{(p+1})/(\text{b}*(\text{n}*(\text{p}+1) + 1))), \text{x}] - \text{Simp}[(\text{a}*d - \text{b}*c*(\text{n}*(\text{p}+1) + 1))/(\text{b}*(\text{n}*(\text{p}+1) + 1)) \quad \text{Int}[(\text{a} + \text{b}*x^n)^p, \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{n}, \text{p}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{n}*(\text{p}+1) + 1, 0]$
- rule 1083 $\text{Int}[(\text{a}_) + (\text{b}_.)*(x_) + (\text{c}_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[-2 \quad \text{Subst}[\text{Int}[1/\text{Simp}[\text{b}^2 - 4*\text{a}*c - x^2, \text{x}], \text{x}], \text{x}, \text{b} + 2*c*x], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}\}, \text{x}]$
- rule 1103 $\text{Int}[(\text{d}_) + (\text{e}_.)*(x_)]/((\text{a}_) + (\text{b}_.)*(x_) + (\text{c}_.)*(x_)^2), \text{x_Symbol}] \rightarrow \text{Simp}[\text{d}*(\text{Log}[\text{RemoveContent}[\text{a} + \text{b}*x + \text{c}*x^2, \text{x}]]/\text{b}), \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}\}, \text{x}] \ \&\& \ \text{EqQ}[\text{2}*c*d - \text{b}*e, 0]$
- rule 1142 $\text{Int}[(\text{d}_) + (\text{e}_.)*(x_)]/((\text{a}_) + (\text{b}_.)*(x_) + (\text{c}_.)*(x_)^2), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{2}*c*d - \text{b}*e)/(2*c) \quad \text{Int}[1/(\text{a} + \text{b}*x + \text{c}*x^2), \text{x}], \text{x}] + \text{Simp}[\text{e}/(2*c) \quad \text{Int}[(\text{b} + 2*c*x)/(\text{a} + \text{b}*x + \text{c}*x^2), \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}\}, \text{x}]$

rule 1483

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (b_)*(x_)^2 + (c_)*(x_)^4), x_Symbol] :
> With[{q = Rt[a/c, 2]}, With[{r = Rt[2*q - b/c, 2]}, Simp[1/(2*c*q*r) In
t[(d*r - (d - e*q)*x)/(q - r*x + x^2), x], x] + Simp[1/(2*c*q*r) Int[(d*r
+ (d - e*q)*x)/(q + r*x + x^2), x], x]]] /; FreeQ[{a, b, c, d, e}, x] && N
eQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NegQ[b^2 - 4*a*c]
```

rule 3003

```
Int[((a_) + Log[(c_)*(RFx_)^(p_)]*(b_))^(n_), x_Symbol] :> Simp[x*(a +
b*Log[c*RFx^p])^n, x] - Simp[b*n*p Int[SimplifyIntegrand[x*(a + b*Log[c*
RFx^p])^(n - 1)*(D[RFx, x]/RFx), x], x], x] /; FreeQ[{a, b, c, p}, x] && Ra
tionalFunctionQ[RFx, x] && IGtQ[n, 0]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.23 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.10

method	result	size
risch	$x \ln\left(\frac{1}{x^4} + x^4\right) - 4x + \left(\sum_{-R=\text{RootOf}(_Z^8+1)} \frac{\ln(x-_R)}{-R^7}\right)$	34
parts	$x \ln\left(\frac{1}{x^4} + x^4\right) - 4x + \left(\sum_{-R=\text{RootOf}(_Z^8+1)} \frac{\ln(x-_R)}{-R^7}\right)$	34
default	$x \ln\left(\frac{x^8+1}{x^4}\right) - 4x + \left(\sum_{-R=\text{RootOf}(_Z^8+1)} \frac{\ln(x-_R)}{-R^7}\right)$	36

input

```
int(ln(1/x^4+x^4),x,method=_RETURNVERBOSE)
```

output

```
x*ln(1/x^4+x^4)-4*x+sum(1/_R^7*ln(x-_R),_R=RootOf(_Z^8+1))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.45

$$\int \log \left(\frac{1}{x^4} + x^4 \right) dx = x \log \left(\frac{x^8 + 1}{x^4} \right) + \left(\frac{1}{2}i + \frac{1}{2} \right) \sqrt{2}(-1)^{\frac{1}{8}} \log \left(2x + (i + 1) \sqrt{2}(-1)^{\frac{1}{8}} \right) - \left(\frac{1}{2}i - \frac{1}{2} \right) \sqrt{2}(-1)^{\frac{1}{8}} \log \left(2x - (i - 1) \sqrt{2}(-1)^{\frac{1}{8}} \right) + \left(\frac{1}{2}i - \frac{1}{2} \right) \sqrt{2}(-1)^{\frac{1}{8}} \log \left(2x + (i - 1) \sqrt{2}(-1)^{\frac{1}{8}} \right) - \left(\frac{1}{2}i + \frac{1}{2} \right) \sqrt{2}(-1)^{\frac{1}{8}} \log \left(2x - (i + 1) \sqrt{2}(-1)^{\frac{1}{8}} \right) + (-1)^{\frac{1}{8}} \log \left(x + (-1)^{\frac{1}{8}} \right) + i(-1)^{\frac{1}{8}} \log \left(x + i(-1)^{\frac{1}{8}} \right) - i(-1)^{\frac{1}{8}} \log \left(x - i(-1)^{\frac{1}{8}} \right) - (-1)^{\frac{1}{8}} \log \left(x - (-1)^{\frac{1}{8}} \right) - 4x$$

input `integrate(log(1/x^4+x^4),x, algorithm="fricas")`

output `x*log((x^8 + 1)/x^4) + (1/2*I + 1/2)*sqrt(2)*(-1)^(1/8)*log(2*x + (I + 1)*sqrt(2)*(-1)^(1/8)) - (1/2*I - 1/2)*sqrt(2)*(-1)^(1/8)*log(2*x - (I - 1)*sqrt(2)*(-1)^(1/8)) + (1/2*I - 1/2)*sqrt(2)*(-1)^(1/8)*log(2*x + (I - 1)*sqrt(2)*(-1)^(1/8)) - (1/2*I + 1/2)*sqrt(2)*(-1)^(1/8)*log(2*x - (I + 1)*sqrt(2)*(-1)^(1/8)) + (-1)^(1/8)*log(x + (-1)^(1/8)) + I*(-1)^(1/8)*log(x + I*(-1)^(1/8)) - I*(-1)^(1/8)*log(x - I*(-1)^(1/8)) - (-1)^(1/8)*log(x - (-1)^(1/8)) - 4*x`

Sympy [A] (verification not implemented)

Time = 1.14 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.08

$$\int \log \left(\frac{1}{x^4} + x^4 \right) dx = x \log \left(x^4 + \frac{1}{x^4} \right) - 4x - \text{RootSum} \left(t^8 + 1, (t \mapsto t \log(-t + x)) \right)$$

input `integrate(ln(1/x**4+x**4),x)`

output `x*log(x**4 + x**(-4)) - 4*x - RootSum(_t**8 + 1, Lambda(_t, _t*log(-_t + x)))`

Maxima [F]

$$\int \log\left(\frac{1}{x^4} + x^4\right) dx = \int \log\left(x^4 + \frac{1}{x^4}\right) dx$$

input `integrate(log(1/x^4+x^4),x, algorithm="maxima")`

output `x*log(x^8 + 1) - 4*x*log(x) - 4*x + 8*integrate(1/(x^8 + 1), x)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 248, normalized size of antiderivative = 0.74

$$\begin{aligned} \int \log\left(\frac{1}{x^4} + x^4\right) dx &= x \log\left(x^4 + \frac{1}{x^4}\right) + \sqrt{\sqrt{2} + 2} \arctan\left(\frac{2x + \sqrt{-\sqrt{2} + 2}}{\sqrt{\sqrt{2} + 2}}\right) \\ &\quad + \sqrt{\sqrt{2} + 2} \arctan\left(\frac{2x - \sqrt{-\sqrt{2} + 2}}{\sqrt{\sqrt{2} + 2}}\right) \\ &\quad + \sqrt{-\sqrt{2} + 2} \arctan\left(\frac{2x + \sqrt{\sqrt{2} + 2}}{\sqrt{-\sqrt{2} + 2}}\right) \\ &\quad + \sqrt{-\sqrt{2} + 2} \arctan\left(\frac{2x - \sqrt{\sqrt{2} + 2}}{\sqrt{-\sqrt{2} + 2}}\right) \\ &\quad + \frac{1}{2} \sqrt{\sqrt{2} + 2} \log\left(x^2 + x\sqrt{\sqrt{2} + 2} + 1\right) \\ &\quad - \frac{1}{2} \sqrt{\sqrt{2} + 2} \log\left(x^2 - x\sqrt{\sqrt{2} + 2} + 1\right) \\ &\quad + \frac{1}{2} \sqrt{-\sqrt{2} + 2} \log\left(x^2 + x\sqrt{-\sqrt{2} + 2} + 1\right) \\ &\quad - \frac{1}{2} \sqrt{-\sqrt{2} + 2} \log\left(x^2 - x\sqrt{-\sqrt{2} + 2} + 1\right) - 4x \end{aligned}$$

input `integrate(log(1/x^4+x^4),x, algorithm="giac")`

output `x*log(x^4 + 1/x^4) + sqrt(sqrt(2) + 2)*arctan((2*x + sqrt(-sqrt(2) + 2))/sqrt(sqrt(2) + 2)) + sqrt(sqrt(2) + 2)*arctan((2*x - sqrt(-sqrt(2) + 2))/sqrt(sqrt(2) + 2)) + sqrt(-sqrt(2) + 2)*arctan((2*x + sqrt(sqrt(2) + 2))/sqrt(-sqrt(2) + 2)) + sqrt(-sqrt(2) + 2)*arctan((2*x - sqrt(sqrt(2) + 2))/sqrt(-sqrt(2) + 2)) + 1/2*sqrt(sqrt(2) + 2)*log(x^2 + x*sqrt(sqrt(2) + 2) + 1) - 1/2*sqrt(sqrt(2) + 2)*log(x^2 - x*sqrt(sqrt(2) + 2) + 1) + 1/2*sqrt(-sqrt(2) + 2)*log(x^2 + x*sqrt(-sqrt(2) + 2) + 1) - 1/2*sqrt(-sqrt(2) + 2)*log(x^2 - x*sqrt(-sqrt(2) + 2) + 1) - 4*x`

Mupad [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 313, normalized size of antiderivative = 0.94

$$\begin{aligned} & \int \log\left(\frac{1}{x^4} + x^4\right) dx \\ &= x \ln\left(\frac{1}{x^4} + x^4\right) - 4x + \operatorname{atan}\left(\frac{x \sqrt{-\sqrt{2} - 2} 2097152i}{2097152 \sqrt{2 - \sqrt{2}} \sqrt{-\sqrt{2} - 2} + 2097152 \sqrt{2}}\right. \\ &\quad \left. - \frac{x \sqrt{2 - \sqrt{2}} 2097152i}{2097152 \sqrt{2 - \sqrt{2}} \sqrt{-\sqrt{2} - 2} + 2097152 \sqrt{2}}\right) \left(\sqrt{-\sqrt{2} - 2} 1i - \sqrt{2 - \sqrt{2}} 1i\right) \\ &\quad - \operatorname{atan}\left(\frac{x \sqrt{\sqrt{2} - 2} 2097152i}{2097152 \sqrt{2} + 2097152 \sqrt{\sqrt{2} - 2} \sqrt{\sqrt{2} + 2}}\right. \\ &\quad \left. + \frac{x \sqrt{\sqrt{2} + 2} 2097152i}{2097152 \sqrt{2} + 2097152 \sqrt{\sqrt{2} - 2} \sqrt{\sqrt{2} + 2}}\right) \left(\sqrt{\sqrt{2} - 2} 1i + \sqrt{\sqrt{2} + 2} 1i\right) \\ &\quad + \operatorname{atan}\left(-\frac{\sqrt{2} x \sqrt{\sqrt{2} + 2}}{2} + x \sqrt{\sqrt{2} + 2} \left(\frac{1}{2} + \frac{1}{2}i\right)\right) \left(\frac{\sqrt{2} 1i}{2} - \frac{1}{2} - \frac{1}{2}i\right) \sqrt{\sqrt{2} + 2} 2i \\ &\quad - \operatorname{atan}\left(x \sqrt{\sqrt{2} + 2} \left(\frac{1}{2} - \frac{1}{2}i\right) + \frac{\sqrt{2} x \sqrt{\sqrt{2} + 2} 1i}{2}\right) \left(\frac{\sqrt{2}}{2} - \frac{1}{2} + \frac{1}{2}i\right) \sqrt{\sqrt{2} + 2} 2i \end{aligned}$$

input `int(log(1/x^4 + x^4),x)`

output

```

x*log(1/x^4 + x^4) - 4*x + atan((x*(- 2^(1/2) - 2)^(1/2)*2097152i)/(209715
2*(2 - 2^(1/2))^(1/2)*(- 2^(1/2) - 2)^(1/2) + 2097152*2^(1/2)) - (x*(2 - 2
^(1/2))^(1/2)*2097152i)/(2097152*(2 - 2^(1/2))^(1/2)*(- 2^(1/2) - 2)^(1/2)
+ 2097152*2^(1/2)))*((- 2^(1/2) - 2)^(1/2)*1i - (2 - 2^(1/2))^(1/2)*1i) -
atan((x*(2^(1/2) - 2)^(1/2)*2097152i)/(2097152*2^(1/2) + 2097152*(2^(1/2)
- 2)^(1/2)*(2^(1/2) + 2)^(1/2)) + (x*(2^(1/2) + 2)^(1/2)*2097152i)/(20971
52*2^(1/2) + 2097152*(2^(1/2) - 2)^(1/2)*(2^(1/2) + 2)^(1/2)))*((2^(1/2) -
2)^(1/2)*1i + (2^(1/2) + 2)^(1/2)*1i) + atan(x*(2^(1/2) + 2)^(1/2)*(1/2 +
1i/2) - (2^(1/2)*x*(2^(1/2) + 2)^(1/2))/2)*((2^(1/2)*1i)/2 - (1/2 + 1i/2)
)*(2^(1/2) + 2)^(1/2)*2i - atan(x*(2^(1/2) + 2)^(1/2)*(1/2 - 1i/2) + (2^(1
/2)*x*(2^(1/2) + 2)^(1/2)*1i)/2)*(2^(1/2)/2 - (1/2 - 1i/2))*(2^(1/2) + 2)^(
1/2)*2i

```

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 216, normalized size of antiderivative = 0.65

$$\begin{aligned}
\int \log\left(\frac{1}{x^4} + x^4\right) dx &= -\sqrt{\sqrt{2} + 2} \operatorname{atan}\left(\frac{\sqrt{-\sqrt{2} + 2} - 2x}{\sqrt{\sqrt{2} + 2}}\right) \\
&+ \sqrt{\sqrt{2} + 2} \operatorname{atan}\left(\frac{\sqrt{-\sqrt{2} + 2} + 2x}{\sqrt{\sqrt{2} + 2}}\right) \\
&- \sqrt{-\sqrt{2} + 2} \operatorname{atan}\left(\frac{\sqrt{\sqrt{2} + 2} - 2x}{\sqrt{-\sqrt{2} + 2}}\right) \\
&+ \sqrt{-\sqrt{2} + 2} \operatorname{atan}\left(\frac{\sqrt{\sqrt{2} + 2} + 2x}{\sqrt{-\sqrt{2} + 2}}\right) \\
&- \frac{\sqrt{-\sqrt{2} + 2} \log\left(-\sqrt{-\sqrt{2} + 2} x + x^2 + 1\right)}{2} \\
&+ \frac{\sqrt{-\sqrt{2} + 2} \log\left(\sqrt{-\sqrt{2} + 2} x + x^2 + 1\right)}{2} \\
&- \frac{\sqrt{\sqrt{2} + 2} \log\left(-\sqrt{\sqrt{2} + 2} x + x^2 + 1\right)}{2} \\
&+ \frac{\sqrt{\sqrt{2} + 2} \log\left(\sqrt{\sqrt{2} + 2} x + x^2 + 1\right)}{2} + \log\left(\frac{x^8 + 1}{x^4}\right) x - 4x
\end{aligned}$$

input `int(log(1/x^4+x^4),x)`

output
$$\begin{aligned} & (-2\sqrt{\sqrt{2} + 2} \operatorname{atan}(\frac{\sqrt{-\sqrt{2} + 2} - 2x}{\sqrt{\sqrt{2} + 2}}) + 2\sqrt{\sqrt{2} + 2} \operatorname{atan}(\frac{\sqrt{-\sqrt{2} + 2} + 2x}{\sqrt{\sqrt{2} + 2}}) - 2\sqrt{-\sqrt{2} + 2} \operatorname{atan}(\frac{\sqrt{\sqrt{2} + 2} - 2x}{\sqrt{-\sqrt{2} + 2}}) + 2\sqrt{-\sqrt{2} + 2} \operatorname{atan}(\frac{\sqrt{\sqrt{2} + 2} + 2x}{\sqrt{-\sqrt{2} + 2}}) - \sqrt{-\sqrt{2} + 2} \log(-\sqrt{-\sqrt{2} + 2}x + x^2 + 1) + \sqrt{-\sqrt{2} + 2} \log(\sqrt{-\sqrt{2} + 2}x + x^2 + 1) - \sqrt{\sqrt{2} + 2} \log(-\sqrt{\sqrt{2} + 2}x + x^2 + 1) + \sqrt{\sqrt{2} + 2} \log(\sqrt{\sqrt{2} + 2}x + x^2 + 1) + 2 \log((x^8 + 1)/x^4)x - 8x)/2 \end{aligned}$$

3.7 $\int \frac{\log(1+x)}{x\sqrt{1+\sqrt{1+x}}} dx$

Optimal result	80
Mathematica [A] (warning: unable to verify)	81
Rubi [F]	82
Maple [C] (verified)	83
Fricas [F(-2)]	83
Sympy [F(-1)]	84
Maxima [A] (verification not implemented)	84
Giac [F]	85
Mupad [F(-1)]	85
Reduce [F]	86

Optimal result

Integrand size = 21, antiderivative size = 291

$$\begin{aligned}
 \int \frac{\log(1+x)}{x\sqrt{1+\sqrt{1+x}}} dx = & -8\operatorname{arctanh}\left(\sqrt{1+\sqrt{1+x}}\right) - \frac{2\log(1+x)}{\sqrt{1+\sqrt{1+x}}} \\
 & - \sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{1+\sqrt{1+x}}}{\sqrt{2}}\right) \log(1+x) \\
 & + 2\sqrt{2}\operatorname{arctanh}\left(\frac{1}{\sqrt{2}}\right) \log\left(1 - \sqrt{1+\sqrt{1+x}}\right) \\
 & - 2\sqrt{2}\operatorname{arctanh}\left(\frac{1}{\sqrt{2}}\right) \log\left(1 + \sqrt{1+\sqrt{1+x}}\right) \\
 & + \sqrt{2}\operatorname{PolyLog}\left(2, -\frac{\sqrt{2}\left(1 - \sqrt{1+\sqrt{1+x}}\right)}{2 - \sqrt{2}}\right) \\
 & - \sqrt{2}\operatorname{PolyLog}\left(2, \frac{\sqrt{2}\left(1 - \sqrt{1+\sqrt{1+x}}\right)}{2 + \sqrt{2}}\right) \\
 & - \sqrt{2}\operatorname{PolyLog}\left(2, -\frac{\sqrt{2}\left(1 + \sqrt{1+\sqrt{1+x}}\right)}{2 - \sqrt{2}}\right) \\
 & + \sqrt{2}\operatorname{PolyLog}\left(2, \frac{\sqrt{2}\left(1 + \sqrt{1+\sqrt{1+x}}\right)}{2 + \sqrt{2}}\right)
 \end{aligned}$$

output

```
-8*arctanh((1+(1+x)^(1/2))^(1/2))-arctanh(1/2*(1+(1+x)^(1/2))^(1/2)*2^(1/2))
)*ln(1+x)*2^(1/2)+2*arctanh(1/2*2^(1/2))*ln(1-(1+(1+x)^(1/2))^(1/2))*2^(1/2)
)-2*arctanh(1/2*2^(1/2))*ln(1+(1+(1+x)^(1/2))^(1/2))*2^(1/2)+polylog(2,-2^(1/2)
)*(1-(1+(1+x)^(1/2))^(1/2))/(2-2^(1/2)))*2^(1/2)-polylog(2,2^(1/2))*(1-(1+(1+x)^(1/2))^(1/2)
)/(2+2^(1/2)))*2^(1/2)-polylog(2,-2^(1/2)*(1+(1+(1+x)^(1/2))^(1/2))/(2-2^(1/2)))*2^(1/2)
)+polylog(2,2^(1/2)*(1+(1+(1+x)^(1/2))^(1/2))/(2+2^(1/2)))*2^(1/2)-2*ln(1+x)/(1+(1+x)^(1/2))^(1/2)
```

Mathematica [A] (warning: unable to verify)

Time = 0.19 (sec) , antiderivative size = 310, normalized size of antiderivative = 1.07

$$\int \frac{\log(1+x)}{x\sqrt{1+\sqrt{1+x}}} dx$$

$$= -8\operatorname{arctanh}\left(\sqrt{1+\sqrt{1+x}}\right) - \frac{2\log(1+x)}{\sqrt{1+\sqrt{1+x}}}$$

$$+ \frac{\log(1+x)\left(\log\left(\sqrt{2}-\sqrt{1+\sqrt{1+x}}\right) - \log\left(\sqrt{2}+\sqrt{1+\sqrt{1+x}}\right)\right)}{\sqrt{2}}$$

$$+ \sqrt{2}\left(-\log(-1+\sqrt{2})\log\left(1-\sqrt{1+\sqrt{1+x}}\right)\right.$$

$$+ \log(1+\sqrt{2})\log\left(1-\sqrt{1+\sqrt{1+x}}\right) + \log(-1+\sqrt{2})\log\left(1+\sqrt{1+\sqrt{1+x}}\right)$$

$$\left. - \log(1+\sqrt{2})\log\left(1+\sqrt{1+\sqrt{1+x}}\right)\right)$$

$$- \operatorname{PolyLog}\left(2, -\left(\left(-1+\sqrt{2}\right)\left(-1+\sqrt{1+\sqrt{1+x}}\right)\right)\right)$$

$$+ \operatorname{PolyLog}\left(2, \left(1+\sqrt{2}\right)\left(-1+\sqrt{1+\sqrt{1+x}}\right)\right)$$

$$+ \operatorname{PolyLog}\left(2, \left(-1+\sqrt{2}\right)\left(1+\sqrt{1+\sqrt{1+x}}\right)\right)$$

$$- \operatorname{PolyLog}\left(2, -\left(\left(1+\sqrt{2}\right)\left(1+\sqrt{1+\sqrt{1+x}}\right)\right)\right)$$

input

```
Integrate[Log[1 + x]/(x*Sqrt[1 + Sqrt[1 + x]]), x]
```

output

```
-8*ArcTanh[Sqrt[1 + Sqrt[1 + x]]] - (2*Log[1 + x])/Sqrt[1 + Sqrt[1 + x]] +
(Log[1 + x]*(Log[Sqrt[2] - Sqrt[1 + Sqrt[1 + x]]] - Log[Sqrt[2] + Sqrt[1
+ Sqrt[1 + x]]]))/Sqrt[2] + Sqrt[2]*(-(Log[-1 + Sqrt[2]]*Log[1 - Sqrt[1 +
Sqrt[1 + x]]]) + Log[1 + Sqrt[2]]*Log[1 - Sqrt[1 + Sqrt[1 + x]]] + Log[-1
+ Sqrt[2]]*Log[1 + Sqrt[1 + Sqrt[1 + x]]] - Log[1 + Sqrt[2]]*Log[1 + Sqrt[
1 + Sqrt[1 + x]]] - PolyLog[2, -((-1 + Sqrt[2])*(-1 + Sqrt[1 + Sqrt[1 + x]
])]) + PolyLog[2, (1 + Sqrt[2])*(-1 + Sqrt[1 + Sqrt[1 + x]])] + PolyLog[2,
(-1 + Sqrt[2])*(1 + Sqrt[1 + Sqrt[1 + x]])] - PolyLog[2, -((1 + Sqrt[2])*
(1 + Sqrt[1 + Sqrt[1 + x]))])
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\log(x+1)}{x\sqrt{\sqrt{x+1}+1}} dx$$

↓ 2867

$$\int \frac{\log(x+1)}{x\sqrt{\sqrt{x+1}+1}} dx$$

input

```
Int[Log[1 + x]/(x*Sqrt[1 + Sqrt[1 + x]]),x]
```

output

```
$Aborted
```

Defintions of rubi rules used

rule 2867

```
Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_))^(n_.)]*(b_.))^(p_.)*(AFx_), x_Sy
mbol] := Unintegrable[AFx*(a + b*Log[c*(d + e*x)^n])^p, x] /; FreeQ[{a, b,
c, d, e, n, p}, x] && AlgebraicFunctionQ[AFx, x, True]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.22 (sec) , antiderivative size = 172, normalized size of antiderivative = 0.59

method	result
derivativedivides	$-\frac{2\ln(1+x)}{\sqrt{1+\sqrt{1+x}}} - 8 \operatorname{arctanh}\left(\sqrt{1+\sqrt{1+x}}\right) + 8 \left(\sum_{-\alpha=\operatorname{RootOf}(-Z^2-2)} \frac{\left(\frac{\ln(\sqrt{1+\sqrt{1+x}}-\alpha)\ln(1+x)}{2}\right)}{\dots} \right)$
default	$-\frac{2\ln(1+x)}{\sqrt{1+\sqrt{1+x}}} - 8 \operatorname{arctanh}\left(\sqrt{1+\sqrt{1+x}}\right) + 8 \left(\sum_{-\alpha=\operatorname{RootOf}(-Z^2-2)} \frac{\left(\frac{\ln(\sqrt{1+\sqrt{1+x}}-\alpha)\ln(1+x)}{2}\right)}{\dots} \right)$

input `int(ln(1+x)/x/(1+(1+x)^(1/2))^(1/2),x,method=_RETURNVERBOSE)`

output `-2*ln(1+x)/(1+(1+x)^(1/2))^(1/2)-8*arctanh((1+(1+x)^(1/2))^(1/2))+8*Sum(1/8*(1/2*ln((1+(1+x)^(1/2))^(1/2)-_alpha)*ln(1+x)-dilog((1+(1+(1+x)^(1/2))^(1/2))/(1+_alpha))-ln((1+(1+x)^(1/2))^(1/2)-_alpha)*ln((1+(1+(1+x)^(1/2))^(1/2))/(1+_alpha))-dilog((1+(1+x)^(1/2))^(1/2)-1)/(-1+_alpha))-ln((1+(1+x)^(1/2))^(1/2)-_alpha)*ln(((1+(1+x)^(1/2))^(1/2)-1)/(-1+_alpha)))*_alpha,_alpha=RootOf(-Z^2-2))`

Fricas [F(-2)]

Exception generated.

$$\int \frac{\log(1+x)}{x\sqrt{1+\sqrt{1+x}}} dx = \text{Exception raised: TypeError}$$

input `integrate(log(1+x)/x/(1+(1+x)^(1/2))^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\log(1+x)}{x\sqrt{1+\sqrt{1+x}}} dx = \text{Timed out}$$

input `integrate(ln(1+x)/x/(1+(1+x)**(1/2))**(1/2),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 366, normalized size of antiderivative = 1.26

$$\begin{aligned} & \int \frac{\log(1+x)}{x\sqrt{1+\sqrt{1+x}}} dx \\ &= \frac{1}{2} \left(\sqrt{2} \log \left(-\frac{\sqrt{2}-\sqrt{\sqrt{x+1}+1}}{\sqrt{2}+\sqrt{\sqrt{x+1}+1}} \right) - \frac{4}{\sqrt{\sqrt{x+1}+1}} \right) \log(x+1) \\ &+ \sqrt{2} \left(\log \left(\sqrt{2} + \sqrt{\sqrt{x+1}+1} \right) \log \left(-\frac{\sqrt{2}+\sqrt{\sqrt{x+1}+1}}{\sqrt{2}+1} + 1 \right) + \text{Li}_2 \left(\frac{\sqrt{2}+\sqrt{\sqrt{x+1}+1}}{\sqrt{2}+1} \right) \right) \\ &- \sqrt{2} \left(\log \left(-\sqrt{2} + \sqrt{\sqrt{x+1}+1} \right) \log \left(-\frac{\sqrt{2}-\sqrt{\sqrt{x+1}+1}}{\sqrt{2}+1} + 1 \right) + \text{Li}_2 \left(\frac{\sqrt{2}-\sqrt{\sqrt{x+1}+1}}{\sqrt{2}+1} \right) \right) \\ &+ \sqrt{2} \left(\log \left(\sqrt{2} + \sqrt{\sqrt{x+1}+1} \right) \log \left(-\frac{\sqrt{2}+\sqrt{\sqrt{x+1}+1}}{\sqrt{2}-1} + 1 \right) + \text{Li}_2 \left(\frac{\sqrt{2}+\sqrt{\sqrt{x+1}+1}}{\sqrt{2}-1} \right) \right) \\ &- \sqrt{2} \left(\log \left(-\sqrt{2} + \sqrt{\sqrt{x+1}+1} \right) \log \left(-\frac{\sqrt{2}-\sqrt{\sqrt{x+1}+1}}{\sqrt{2}-1} + 1 \right) + \text{Li}_2 \left(\frac{\sqrt{2}-\sqrt{\sqrt{x+1}+1}}{\sqrt{2}-1} \right) \right) \\ &- 4 \log \left(\sqrt{\sqrt{x+1}+1} + 1 \right) + 4 \log \left(\sqrt{\sqrt{x+1}+1} - 1 \right) \end{aligned}$$

input `integrate(log(1+x)/x/(1+(1+x)^(1/2))^(1/2),x, algorithm="maxima")`

output

```
1/2*(sqrt(2)*log(-sqrt(2) - sqrt(sqrt(x + 1) + 1))/(sqrt(2) + sqrt(sqrt(x
+ 1) + 1))) - 4/sqrt(sqrt(x + 1) + 1)*log(x + 1) + sqrt(2)*(log(sqrt(2)
+ sqrt(sqrt(x + 1) + 1))*log(-sqrt(2) + sqrt(sqrt(x + 1) + 1))/(sqrt(2) +
1) + 1) + dilog((sqrt(2) + sqrt(sqrt(x + 1) + 1))/(sqrt(2) + 1))) - sqrt(
2)*(log(-sqrt(2) + sqrt(sqrt(x + 1) + 1))*log(-sqrt(2) - sqrt(sqrt(x + 1)
+ 1))/(sqrt(2) + 1) + 1) + dilog((sqrt(2) - sqrt(sqrt(x + 1) + 1))/(sqrt(
2) + 1))) + sqrt(2)*(log(sqrt(2) + sqrt(sqrt(x + 1) + 1))*log(-sqrt(2) +
sqrt(sqrt(x + 1) + 1))/(sqrt(2) - 1) + 1) + dilog((sqrt(2) + sqrt(sqrt(x +
1) + 1))/(sqrt(2) - 1))) - sqrt(2)*(log(-sqrt(2) + sqrt(sqrt(x + 1) + 1))
*log(-sqrt(2) - sqrt(sqrt(x + 1) + 1))/(sqrt(2) - 1) + 1) + dilog((sqrt(2)
) - sqrt(sqrt(x + 1) + 1))/(sqrt(2) - 1))) - 4*log(sqrt(sqrt(x + 1) + 1) +
1) + 4*log(sqrt(sqrt(x + 1) + 1) - 1)
```

Giac [F]

$$\int \frac{\log(1+x)}{x\sqrt{1+\sqrt{1+x}}} dx = \int \frac{\log(x+1)}{x\sqrt{\sqrt{x+1}+1}} dx$$

input

```
integrate(log(1+x)/x/(1+(1+x)^(1/2))^(1/2),x, algorithm="giac")
```

output

```
integrate(log(x + 1)/(x*sqrt(sqrt(x + 1) + 1)), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\log(1+x)}{x\sqrt{1+\sqrt{1+x}}} dx = \int \frac{\ln(x+1)}{x\sqrt{\sqrt{x+1}+1}} dx$$

input

```
int(log(x + 1)/(x*((x + 1)^(1/2) + 1)^(1/2)),x)
```

output

```
int(log(x + 1)/(x*((x + 1)^(1/2) + 1)^(1/2)), x)
```

Reduce [F]

$$\int \frac{\log(1+x)}{x\sqrt{1+\sqrt{1+x}}} dx$$

$$4\sqrt{x+1}\sqrt{\sqrt{x+1}+1}\sqrt{2}\log\left(\sqrt{\sqrt{x+1}+1}-\sqrt{2}\right) - 4\sqrt{x+1}\sqrt{\sqrt{x+1}+1}\sqrt{2}\log\left(\sqrt{\sqrt{x+1}+1}+\sqrt{2}\right)$$

input `int(log(1+x)/x/(1+(1+x)^(1/2))^(1/2),x)`

output `(2*(2*sqrt(x + 1)*sqrt(sqrt(x + 1) + 1)*sqrt(2)*log(sqrt(sqrt(x + 1) + 1) - sqrt(2)) - 2*sqrt(x + 1)*sqrt(sqrt(x + 1) + 1)*sqrt(2)*log(sqrt(sqrt(x + 1) + 1) + sqrt(2)) - sqrt(x + 1)*sqrt(sqrt(x + 1) + 1)*int((sqrt(sqrt(x + 1) + 1)*log(x + 1))/(x**3 + x**2),x) - sqrt(x + 1)*sqrt(sqrt(x + 1) + 1)*int((sqrt(sqrt(x + 1) + 1)*log(x + 1))/(x**2 + x),x) - sqrt(x + 1)*sqrt(sqrt(x + 1) + 1)*int((sqrt(x + 1)*sqrt(sqrt(x + 1) + 1)*log(x + 1))/(x**3 + x**2),x) - 2*sqrt(sqrt(x + 1) + 1)*sqrt(2)*log(sqrt(sqrt(x + 1) + 1) - sqrt(2)) + 2*sqrt(sqrt(x + 1) + 1)*sqrt(2)*log(sqrt(sqrt(x + 1) + 1) + sqrt(2)) + sqrt(sqrt(x + 1) + 1)*int((sqrt(sqrt(x + 1) + 1)*log(x + 1))/(x**3 + x**2),x) + sqrt(sqrt(x + 1) + 1)*int((sqrt(sqrt(x + 1) + 1)*log(x + 1))/(x**2 + x),x) + sqrt(sqrt(x + 1) + 1)*int((sqrt(x + 1)*sqrt(sqrt(x + 1) + 1)*log(x + 1))/(x**3 + x**2),x) - 2*sqrt(x + 1)*log(x + 1))/(sqrt(sqrt(x + 1) + 1)*(sqrt(x + 1) - 1))`

3.8 $\int \frac{\sqrt{1+\sqrt{1+x}} \log(1+x)}{x} dx$

Optimal result	88
Mathematica [A] (warning: unable to verify)	89
Rubi [F]	90
Maple [C] (verified)	91
Fricas [F(-2)]	91
Sympy [F(-1)]	92
Maxima [A] (verification not implemented)	92
Giac [F]	93
Mupad [F(-1)]	93
Reduce [F]	94

Optimal result

Integrand size = 21, antiderivative size = 308

$$\begin{aligned}
 \int \frac{\sqrt{1 + \sqrt{1 + x}} \log(1 + x)}{x} dx = & -16\sqrt{1 + \sqrt{1 + x}} + 16\operatorname{arctanh}\left(\sqrt{1 + \sqrt{1 + x}}\right) \\
 & + 4\sqrt{1 + \sqrt{1 + x}} \log(1 + x) \\
 & - 2\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{1 + \sqrt{1 + x}}}{\sqrt{2}}\right) \log(1 + x) \\
 & + 4\sqrt{2}\operatorname{arctanh}\left(\frac{1}{\sqrt{2}}\right) \log\left(1 - \sqrt{1 + \sqrt{1 + x}}\right) \\
 & - 4\sqrt{2}\operatorname{arctanh}\left(\frac{1}{\sqrt{2}}\right) \log\left(1 + \sqrt{1 + \sqrt{1 + x}}\right) \\
 & + 2\sqrt{2}\operatorname{PolyLog}\left(2, -\frac{\sqrt{2}(1 - \sqrt{1 + \sqrt{1 + x}})}{2 - \sqrt{2}}\right) \\
 & - 2\sqrt{2}\operatorname{PolyLog}\left(2, \frac{\sqrt{2}(1 - \sqrt{1 + \sqrt{1 + x}})}{2 + \sqrt{2}}\right) \\
 & - 2\sqrt{2}\operatorname{PolyLog}\left(2, -\frac{\sqrt{2}(1 + \sqrt{1 + \sqrt{1 + x}})}{2 - \sqrt{2}}\right) \\
 & + 2\sqrt{2}\operatorname{PolyLog}\left(2, \frac{\sqrt{2}(1 + \sqrt{1 + \sqrt{1 + x}})}{2 + \sqrt{2}}\right)
 \end{aligned}$$

output

```

16*arctanh((1+(1+x)^(1/2))^(1/2))-2*arctanh(1/2*(1+(1+x)^(1/2))^(1/2)*2^(1/2))*ln(1+x)*2^(1/2)+4*arctanh(1/2*2^(1/2))*ln(1-(1+(1+x)^(1/2))^(1/2))*2^(1/2)-4*arctanh(1/2*2^(1/2))*ln(1+(1+(1+x)^(1/2))^(1/2))*2^(1/2)+2*polylog(2,-2^(1/2)*(1-(1+(1+x)^(1/2))^(1/2))/(2-2^(1/2)))*2^(1/2)-2*polylog(2,2^(1/2)*(1-(1+(1+x)^(1/2))^(1/2))/(2+2^(1/2)))*2^(1/2)-2*polylog(2,-2^(1/2)*(1+(1+(1+x)^(1/2))^(1/2))/(2-2^(1/2)))*2^(1/2)+2*polylog(2,2^(1/2)*(1+(1+(1+x)^(1/2))^(1/2))/(2+2^(1/2)))*2^(1/2)-16*(1+(1+x)^(1/2))^(1/2)+4*ln(1+x)*(1+(1+x)^(1/2))^(1/2)

```

Mathematica [A] (warning: unable to verify)

Time = 0.09 (sec) , antiderivative size = 326, normalized size of antiderivative = 1.06

$$\begin{aligned}
\int \frac{\sqrt{1 + \sqrt{1 + x}} \log(1 + x)}{x} dx = & -16\sqrt{1 + \sqrt{1 + x}} + 16\operatorname{arctanh}\left(\sqrt{1 + \sqrt{1 + x}}\right) \\
& + 4\sqrt{1 + \sqrt{1 + x}} \log(1 + x) \\
& + \sqrt{2} \log(1 + x) \left(\log\left(\sqrt{2} - \sqrt{1 + \sqrt{1 + x}}\right) \right. \\
& \qquad \qquad \qquad \left. - \log\left(\sqrt{2} + \sqrt{1 + \sqrt{1 + x}}\right) \right) \\
& - 2\sqrt{2} \left(\log(-1 + \sqrt{2}) \log\left(1 - \sqrt{1 + \sqrt{1 + x}}\right) \right. \\
& \qquad \qquad \qquad - \log(1 + \sqrt{2}) \log\left(1 - \sqrt{1 + \sqrt{1 + x}}\right) \\
& \qquad \qquad \qquad - \log(-1 + \sqrt{2}) \log\left(1 + \sqrt{1 + \sqrt{1 + x}}\right) \\
& + \log(1 + \sqrt{2}) \log\left(1 + \sqrt{1 + \sqrt{1 + x}}\right) + \operatorname{PolyLog}\left(2, \right. \\
& \qquad \qquad \qquad \left. - \left((-1 + \sqrt{2}) \left(-1 + \sqrt{1 + \sqrt{1 + x}} \right) \right) \right) \\
& - \operatorname{PolyLog}\left(2, (1 + \sqrt{2}) \left(-1 + \sqrt{1 + \sqrt{1 + x}} \right) \right) \\
& - \operatorname{PolyLog}\left(2, (-1 + \sqrt{2}) \left(1 + \sqrt{1 + \sqrt{1 + x}} \right) \right) \\
& \qquad \qquad \qquad + \operatorname{PolyLog}\left(2, \right. \\
& \qquad \qquad \qquad \left. - \left((1 + \sqrt{2}) \left(1 + \sqrt{1 + \sqrt{1 + x}} \right) \right) \right)
\end{aligned}$$

input

```
Integrate[(Sqrt[1 + Sqrt[1 + x]]*Log[1 + x])/x,x]
```

output

```
-16*Sqrt[1 + Sqrt[1 + x]] + 16*ArcTanh[Sqrt[1 + Sqrt[1 + x]]] + 4*Sqrt[1 +
Sqrt[1 + x]]*Log[1 + x] + Sqrt[2]*Log[1 + x]*(Log[Sqrt[2] - Sqrt[1 + Sqrt
[1 + x]]] - Log[Sqrt[2] + Sqrt[1 + Sqrt[1 + x]]]) - 2*Sqrt[2]*(Log[-1 + Sq
rt[2]]*Log[1 - Sqrt[1 + Sqrt[1 + x]]] - Log[1 + Sqrt[2]]*Log[1 - Sqrt[1 +
Sqrt[1 + x]]] - Log[-1 + Sqrt[2]]*Log[1 + Sqrt[1 + Sqrt[1 + x]]] + Log[1 +
Sqrt[2]]*Log[1 + Sqrt[1 + Sqrt[1 + x]]] + PolyLog[2, -((-1 + Sqrt[2])*(-1
+ Sqrt[1 + Sqrt[1 + x]]))] - PolyLog[2, (1 + Sqrt[2])*(-1 + Sqrt[1 + Sqrt
[1 + x]])] - PolyLog[2, (-1 + Sqrt[2])*(1 + Sqrt[1 + Sqrt[1 + x]])] + Poly
Log[2, -((1 + Sqrt[2])*(1 + Sqrt[1 + Sqrt[1 + x]])])])
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\sqrt{x+1}+1} \log(x+1)}{x} dx$$

↓ 2867

$$\int \frac{\sqrt{\sqrt{x+1}+1} \log(x+1)}{x} dx$$

input

```
Int[(Sqrt[1 + Sqrt[1 + x]]*Log[1 + x])/x,x]
```

output

```
$Aborted
```

Defintions of rubi rules used

rule 2867

```
Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_))^(n_.)]*(b_.))^(p_.)*(AFx_), x_Sy
mbol] := Unintegrable[AFx*(a + b*Log[c*(d + e*x)^n])^p, x] /; FreeQ[{a, b,
c, d, e, n, p}, x] && AlgebraicFunctionQ[AFx, x, True]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.03 (sec) , antiderivative size = 199, normalized size of antiderivative = 0.65

method	result
derivativedivides	$4 \ln(1+x) \sqrt{1+\sqrt{1+x}} - 16\sqrt{1+\sqrt{1+x}} - 8 \ln(\sqrt{1+\sqrt{1+x}} - 1) + 8 \ln(1 +$
default	$4 \ln(1+x) \sqrt{1+\sqrt{1+x}} - 16\sqrt{1+\sqrt{1+x}} - 8 \ln(\sqrt{1+\sqrt{1+x}} - 1) + 8 \ln(1 +$

input `int(ln(1+x)*(1+(1+x)^(1/2))^(1/2)/x,x,method=_RETURNVERBOSE)`

output `4*ln(1+x)*(1+(1+x)^(1/2))^(1/2)-16*(1+(1+x)^(1/2))^(1/2)-8*ln((1+(1+x)^(1/2))^(1/2)-1)+8*ln(1+(1+(1+x)^(1/2))^(1/2))+8*Sum(1/4*(1/2*ln((1+(1+x)^(1/2))^(1/2)-_alpha)*ln(1+x)-dilog((1+(1+(1+x)^(1/2))^(1/2))/(1+_alpha))-ln((1+(1+x)^(1/2))^(1/2)-_alpha)*ln((1+(1+(1+x)^(1/2))^(1/2))/(1+_alpha))-dilog(((1+(1+x)^(1/2))^(1/2)-1)/(-1+_alpha))-ln((1+(1+x)^(1/2))^(1/2)-_alpha)*ln(((1+(1+x)^(1/2))^(1/2)-1)/(-1+_alpha)))*_alpha,_alpha=RootOf(_Z^2-2))`

Fricas [F(-2)]

Exception generated.

$$\int \frac{\sqrt{1+\sqrt{1+x}} \log(1+x)}{x} dx = \text{Exception raised: TypeError}$$

input `integrate(log(1+x)*(1+(1+x)^(1/2))^(1/2)/x,x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{1 + \sqrt{1 + x}} \log(1 + x)}{x} dx = \text{Timed out}$$

input `integrate(ln(1+x)*(1+(1+x)**(1/2))**(1/2)/x,x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 378, normalized size of antiderivative = 1.23

$$\begin{aligned} & \int \frac{\sqrt{1 + \sqrt{1 + x}} \log(1 + x)}{x} dx \\ &= \left(\sqrt{2} \log \left(-\frac{\sqrt{2} - \sqrt{\sqrt{x+1}+1}}{\sqrt{2} + \sqrt{\sqrt{x+1}+1}} \right) + 4 \sqrt{\sqrt{x+1}+1} \right) \log(x+1) \\ &+ 2\sqrt{2} \left(\log \left(\sqrt{2} + \sqrt{\sqrt{x+1}+1} \right) \log \left(-\frac{\sqrt{2} + \sqrt{\sqrt{x+1}+1}}{\sqrt{2} + 1} + 1 \right) + \text{Li}_2 \left(\frac{\sqrt{2} + \sqrt{\sqrt{x+1}+1}}{\sqrt{2} + 1} \right) \right) \\ &- 2\sqrt{2} \left(\log \left(-\sqrt{2} + \sqrt{\sqrt{x+1}+1} \right) \log \left(-\frac{\sqrt{2} - \sqrt{\sqrt{x+1}+1}}{\sqrt{2} + 1} + 1 \right) + \text{Li}_2 \left(\frac{\sqrt{2} - \sqrt{\sqrt{x+1}+1}}{\sqrt{2} + 1} \right) \right) \\ &+ 2\sqrt{2} \left(\log \left(\sqrt{2} + \sqrt{\sqrt{x+1}+1} \right) \log \left(-\frac{\sqrt{2} + \sqrt{\sqrt{x+1}+1}}{\sqrt{2} - 1} + 1 \right) + \text{Li}_2 \left(\frac{\sqrt{2} + \sqrt{\sqrt{x+1}+1}}{\sqrt{2} - 1} \right) \right) \\ &- 2\sqrt{2} \left(\log \left(-\sqrt{2} + \sqrt{\sqrt{x+1}+1} \right) \log \left(-\frac{\sqrt{2} - \sqrt{\sqrt{x+1}+1}}{\sqrt{2} - 1} + 1 \right) + \text{Li}_2 \left(\frac{\sqrt{2} - \sqrt{\sqrt{x+1}+1}}{\sqrt{2} - 1} \right) \right) \\ &- 16 \sqrt{\sqrt{x+1}+1} + 8 \log \left(\sqrt{\sqrt{x+1}+1} + 1 \right) - 8 \log \left(\sqrt{\sqrt{x+1}+1} - 1 \right) \end{aligned}$$

input `integrate(log(1+x)*(1+(1+x)^(1/2))^(1/2)/x,x, algorithm="maxima")`

output

```
(sqrt(2)*log(-sqrt(2) - sqrt(sqrt(x + 1) + 1))/(sqrt(2) + sqrt(sqrt(x + 1)
) + 1))) + 4*sqrt(sqrt(x + 1) + 1)*log(x + 1) + 2*sqrt(2)*(log(sqrt(2) +
sqrt(sqrt(x + 1) + 1))*log(-sqrt(2) + sqrt(sqrt(x + 1) + 1))/(sqrt(2) + 1
) + 1) + dilog((sqrt(2) + sqrt(sqrt(x + 1) + 1))/(sqrt(2) + 1))) - 2*sqrt(
2)*(log(-sqrt(2) + sqrt(sqrt(x + 1) + 1))*log(-sqrt(2) - sqrt(sqrt(x + 1)
+ 1))/(sqrt(2) + 1) + 1) + dilog((sqrt(2) - sqrt(sqrt(x + 1) + 1))/(sqrt(
2) + 1))) + 2*sqrt(2)*(log(sqrt(2) + sqrt(sqrt(x + 1) + 1))*log(-sqrt(2)
+ sqrt(sqrt(x + 1) + 1))/(sqrt(2) - 1) + 1) + dilog((sqrt(2) + sqrt(sqrt(x
+ 1) + 1))/(sqrt(2) - 1))) - 2*sqrt(2)*(log(-sqrt(2) + sqrt(sqrt(x + 1) +
1))*log(-sqrt(2) - sqrt(sqrt(x + 1) + 1))/(sqrt(2) - 1) + 1) + dilog((sq
rt(2) - sqrt(sqrt(x + 1) + 1))/(sqrt(2) - 1))) - 16*sqrt(sqrt(x + 1) + 1)
+ 8*log(sqrt(sqrt(x + 1) + 1) + 1) - 8*log(sqrt(sqrt(x + 1) + 1) - 1)
```

Giac [F]

$$\int \frac{\sqrt{1 + \sqrt{1 + x}} \log(1 + x)}{x} dx = \int \frac{\sqrt{\sqrt{x + 1} + 1} \log(x + 1)}{x} dx$$

input

```
integrate(log(1+x)*(1+(1+x)^(1/2))^(1/2)/x,x, algorithm="giac")
```

output

```
integrate(sqrt(sqrt(x + 1) + 1)*log(x + 1)/x, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{1 + \sqrt{1 + x}} \log(1 + x)}{x} dx = \int \frac{\ln(x + 1) \sqrt{\sqrt{x + 1} + 1}}{x} dx$$

input

```
int((log(x + 1)*((x + 1)^(1/2) + 1)^(1/2))/x,x)
```

output

```
int((log(x + 1)*((x + 1)^(1/2) + 1)^(1/2))/x, x)
```

Reduce [F]

$$\int \frac{\sqrt{1 + \sqrt{1 + x}} \log(1 + x)}{x} dx = \int \frac{\sqrt{\sqrt{x + 1} + 1} \log(x + 1)}{x} dx$$

input `int(log(1+x)*(1+(1+x)^(1/2))^(1/2)/x,x)`

output `int((sqrt(sqrt(x + 1) + 1)*log(x + 1))/x,x)`

3.9 $\int \frac{1}{1+\sqrt{x+\sqrt{1+x^2}}} dx$

Optimal result	95
Mathematica [A] (verified)	95
Rubi [A] (warning: unable to verify)	96
Maple [F]	98
Fricas [A] (verification not implemented)	98
Sympy [F]	98
Maxima [F]	99
Giac [F]	99
Mupad [F(-1)]	99
Reduce [F]	100

Optimal result

Integrand size = 19, antiderivative size = 84

$$\int \frac{1}{1 + \sqrt{x + \sqrt{1 + x^2}}} dx = -\frac{1}{2(x + \sqrt{1 + x^2})} + \frac{1}{\sqrt{x + \sqrt{1 + x^2}}} + \sqrt{x + \sqrt{1 + x^2}} + \frac{1}{2} \log(x + \sqrt{1 + x^2}) - 2 \log\left(1 + \sqrt{x + \sqrt{1 + x^2}}\right)$$

output

$1/2*\ln(x+(x^2+1)^(1/2))-2*\ln(1+(x+(x^2+1)^(1/2))^(1/2))-1/2/(x+(x^2+1)^(1/2))+1/(x+(x^2+1)^(1/2))^(1/2)+(x+(x^2+1)^(1/2))^(1/2)$

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.25

$$\int \frac{1}{1 + \sqrt{x + \sqrt{1 + x^2}}} dx = \frac{1}{2} \left(\frac{-1 + 5x + 2(1 + x)\sqrt{x + \sqrt{1 + x^2}} + \sqrt{1 + x^2}(5 + 2\sqrt{x + \sqrt{1 + x^2}})}{x + \sqrt{1 + x^2}} + \log(x + \sqrt{1 + x^2}) - 4 \log\left(1 + \sqrt{x + \sqrt{1 + x^2}}\right) \right)$$

input `Integrate[(1 + Sqrt[x + Sqrt[1 + x^2]])^(-1), x]`

output `((-1 + 5*x + 2*(1 + x)*Sqrt[x + Sqrt[1 + x^2]] + Sqrt[1 + x^2]*(5 + 2*Sqrt[x + Sqrt[1 + x^2]]))/(x + Sqrt[1 + x^2]) + Log[x + Sqrt[1 + x^2]] - 4*Log[1 + Sqrt[x + Sqrt[1 + x^2]]])/2`

Rubi [A] (warning: unable to verify)

Time = 0.25 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.87, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {2542, 2361, 2123, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{\sqrt{x^2+1}+x}+1} dx \\
 & \quad \downarrow \text{2542} \\
 & \frac{1}{2} \int \frac{(x + \sqrt{x^2+1})^2 + 1}{(x + \sqrt{x^2+1})^2 (\sqrt{x + \sqrt{x^2+1}} + 1)} d(x + \sqrt{x^2+1}) \\
 & \quad \downarrow \text{2361} \\
 & \int \frac{(\sqrt{x^2+1}+x)^2 + 1}{(\sqrt{x^2+1}+x)^{3/2} (\sqrt{\sqrt{x^2+1}+x} + 1)} d\sqrt{\sqrt{x^2+1}+x} \\
 & \quad \downarrow \text{2123} \\
 & \int \left(-\frac{2}{\sqrt{\sqrt{x^2+1}+x}+1} + \frac{1}{\sqrt{\sqrt{x^2+1}+x}} - \frac{1}{\sqrt{x^2+1}+x} + \frac{1}{(\sqrt{x^2+1}+x)^{3/2}+1} \right) d\sqrt{\sqrt{x^2+1}+x} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$\sqrt{\sqrt{x^2+1}+x} + \frac{1}{\sqrt{x^2+1}+x} - \frac{1}{2(\sqrt{x^2+1}+x)^2} + \log(\sqrt{x^2+1}+x) - 2\log(\sqrt{x^2+1}+x+1)$$

input `Int[(1 + Sqrt[x + Sqrt[1 + x^2]])^(-1), x]`

output `-1/2*1/(x + Sqrt[1 + x^2])^2 + (x + Sqrt[1 + x^2])^(-1) + Sqrt[x + Sqrt[1 + x^2]] + Log[x + Sqrt[1 + x^2]] - 2*Log[1 + x + Sqrt[1 + x^2]]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2123 `Int[(Px_)*((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && PolyQ[Px, x] && (IntegersQ[m, n] || IGtQ[m, -2])`

rule 2361 `Int[(Pq_)*(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*SubstFor[x^n, Pq, x]*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && PolyQ[Pq, x^n] && IntegerQ[Simplify[(m + 1)/n]]`

rule 2542 `Int[((g_) + (h_)*((d_) + (e_)*(x_) + (f_)*Sqrt[(a_) + (c_)*(x_)^2]))^(n_))^(p_), x_Symbol] := Simp[1/(2*e) Subst[Int[(g + h*x^n)^p*((d^2 + a*f^2 - 2*d*x + x^2)/(d - x)^2), x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]`

Maple [F]

$$\int \frac{1}{1 + \sqrt{x + \sqrt{x^2 + 1}}} dx$$

input `int(1/(1+(x+(x^2+1)^(1/2))^(1/2)),x)`

output `int(1/(1+(x+(x^2+1)^(1/2))^(1/2)),x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.79

$$\int \frac{1}{1 + \sqrt{x + \sqrt{1 + x^2}}} dx = -\sqrt{x + \sqrt{x^2 + 1}}(x - \sqrt{x^2 + 1} - 1) + \frac{1}{2}x - \frac{1}{2}\sqrt{x^2 + 1} - 2 \log\left(\sqrt{x + \sqrt{x^2 + 1}} + 1\right) + \log\left(\sqrt{x + \sqrt{x^2 + 1}}\right)$$

input `integrate(1/(1+(x+(x^2+1)^(1/2))^(1/2)),x, algorithm="fricas")`

output `-sqrt(x + sqrt(x^2 + 1))*(x - sqrt(x^2 + 1) - 1) + 1/2*x - 1/2*sqrt(x^2 + 1) - 2*log(sqrt(x + sqrt(x^2 + 1)) + 1) + log(sqrt(x + sqrt(x^2 + 1)))`

Sympy [F]

$$\int \frac{1}{1 + \sqrt{x + \sqrt{1 + x^2}}} dx = \int \frac{1}{\sqrt{x + \sqrt{x^2 + 1}} + 1} dx$$

input `integrate(1/(1+(x+(x**2+1)**(1/2))**(1/2)),x)`

output `Integral(1/(sqrt(x + sqrt(x**2 + 1)) + 1), x)`

Maxima [F]

$$\int \frac{1}{1 + \sqrt{x + \sqrt{1 + x^2}}} dx = \int \frac{1}{\sqrt{x + \sqrt{x^2 + 1}} + 1} dx$$

input `integrate(1/(1+(x+(x^2+1)^(1/2))^(1/2)),x, algorithm="maxima")`

output `integrate(1/(sqrt(x + sqrt(x^2 + 1)) + 1), x)`

Giac [F]

$$\int \frac{1}{1 + \sqrt{x + \sqrt{1 + x^2}}} dx = \int \frac{1}{\sqrt{x + \sqrt{x^2 + 1}} + 1} dx$$

input `integrate(1/(1+(x+(x^2+1)^(1/2))^(1/2)),x, algorithm="giac")`

output `integrate(1/(sqrt(x + sqrt(x^2 + 1)) + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{1 + \sqrt{x + \sqrt{1 + x^2}}} dx = \int \frac{1}{\sqrt{x + \sqrt{x^2 + 1}} + 1} dx$$

input `int(1/((x + (x^2 + 1)^(1/2))^(1/2) + 1),x)`

output `int(1/((x + (x^2 + 1)^(1/2))^(1/2) + 1), x)`

Reduce [F]

$$\begin{aligned}
\int \frac{1}{1 + \sqrt{x + \sqrt{1 + x^2}}} dx &= \frac{\sqrt{\sqrt{x^2 + 1} + x} \sqrt{x^2 + 1}}{2} - \frac{\sqrt{x^2 + 1}}{2} \\
&- \frac{3 \left(\int \frac{\sqrt{\sqrt{x^2 + 1} + x}}{x^2 + 1} dx \right)}{4} - \frac{3 \left(\int \frac{\sqrt{\sqrt{x^2 + 1} + x} x^2}{x^2 + 1} dx \right)}{4} \\
&+ \frac{\left(\int \frac{\sqrt{\sqrt{x^2 + 1} + x} \sqrt{x^2 + 1}}{x^2 + 1} dx \right)}{2} - \frac{\log(\sqrt{x^2 + 1} + x - 1)}{2} \\
&+ \frac{\log(\sqrt{x^2 + 1} + x + 1)}{2} + \log\left(\sqrt{\sqrt{x^2 + 1} + x} - 1\right) \\
&- \log\left(\sqrt{\sqrt{x^2 + 1} + x} + 1\right) - \frac{\log(x)}{2} + \frac{x}{2}
\end{aligned}$$

input `int(1/(1+(x+(x^2+1)^(1/2))^(1/2)),x)`

output `(2*sqrt(sqrt(x**2 + 1) + x)*sqrt(x**2 + 1) - 2*sqrt(x**2 + 1) - 3*int(sqrt(sqrt(x**2 + 1) + x)/(x**2 + 1),x) - 3*int((sqrt(sqrt(x**2 + 1) + x)*x**2)/(x**2 + 1),x) + 2*int((sqrt(sqrt(x**2 + 1) + x)*sqrt(x**2 + 1))/(x**2 + 1),x) - 2*log(sqrt(x**2 + 1) + x - 1) + 2*log(sqrt(x**2 + 1) + x + 1) + 4*log(sqrt(sqrt(x**2 + 1) + x) - 1) - 4*log(sqrt(sqrt(x**2 + 1) + x) + 1) - 2*log(x) + 2*x)/4`

3.10 $\int \frac{\sqrt{1+x}}{x+\sqrt{1+\sqrt{1+x}}} dx$

Optimal result	101
Mathematica [A] (verified)	101
Rubi [A] (warning: unable to verify)	102
Maple [A] (verified)	103
Fricas [B] (verification not implemented)	104
Sympy [A] (verification not implemented)	104
Maxima [A] (verification not implemented)	105
Giac [A] (verification not implemented)	105
Mupad [F(-1)]	105
Reduce [B] (verification not implemented)	106

Optimal result

Integrand size = 25, antiderivative size = 41

$$\int \frac{\sqrt{1+x}}{x+\sqrt{1+\sqrt{1+x}}} dx = 2\sqrt{1+x} + \frac{8\operatorname{arctanh}\left(\frac{1+2\sqrt{1+\sqrt{1+x}}}{\sqrt{5}}\right)}{\sqrt{5}}$$

output `8/5*arctanh(1/5*(1+2*(1+(1+x)^(1/2))^(1/2))*5^(1/2))*5^(1/2)+2*(1+x)^(1/2)`

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{1+x}}{x+\sqrt{1+\sqrt{1+x}}} dx = 2\sqrt{1+x} + \frac{8\operatorname{arctanh}\left(\frac{1+2\sqrt{1+\sqrt{1+x}}}{\sqrt{5}}\right)}{\sqrt{5}}$$

input `Integrate[Sqrt[1 + x]/(x + Sqrt[1 + Sqrt[1 + x]]),x]`

output `2*Sqrt[1 + x] + (8*ArcTanh[(1 + 2*Sqrt[1 + Sqrt[1 + x]])/Sqrt[5]])/Sqrt[5]`

Rubi [A] (warning: unable to verify)

Time = 0.43 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.73, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {7267, 25, 7267, 1200, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{x+1}}{x + \sqrt{\sqrt{x+1}+1}} dx \\
 & \quad \downarrow \text{7267} \\
 & 2 \int -\frac{x+1}{-x - \sqrt{\sqrt{x+1}+1}} d\sqrt{x+1} \\
 & \quad \downarrow \text{25} \\
 & -2 \int \frac{x+1}{-x - \sqrt{\sqrt{x+1}+1}} d\sqrt{x+1} \\
 & \quad \downarrow \text{7267} \\
 & 4 \int \frac{(1 - \sqrt{\sqrt{x+1}+1})(\sqrt{\sqrt{x+1}+1}+1)^2}{-x - \sqrt{\sqrt{x+1}+1}} d\sqrt{\sqrt{x+1}+1} \\
 & \quad \downarrow \text{1200} \\
 & 4 \int \left(\sqrt{\sqrt{x+1}+1} + \frac{1}{-x - \sqrt{\sqrt{x+1}+1}} \right) d\sqrt{\sqrt{x+1}+1} \\
 & \quad \downarrow \text{2009} \\
 & 4 \left(\frac{x+1}{2} - \frac{\log(2\sqrt{\sqrt{x+1}+1} - \sqrt{5} + 1)}{\sqrt{5}} + \frac{\log(2\sqrt{\sqrt{x+1}+1} + \sqrt{5} + 1)}{\sqrt{5}} \right)
 \end{aligned}$$

input `Int[Sqrt[1 + x]/(x + Sqrt[1 + Sqrt[1 + x]]),x]`

output `4*((1 + x)/2 - Log[1 - Sqrt[5] + 2*Sqrt[1 + Sqrt[1 + x]]]/Sqrt[5] + Log[1 + Sqrt[5] + 2*Sqrt[1 + Sqrt[1 + x]]]/Sqrt[5])`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 1200 `Int[(((d_.) + (e_.)*(x_)^(m_.))*((f_.) + (g_.)*(x_)^(n_.)))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*((f + g*x)^n/(a + b*x + c*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && IntegersQ[n]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7267 `Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Simp[lst[[2]]*lst[[4] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])]], x] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$2\sqrt{1+x} + 2 + \frac{8 \operatorname{arctanh}\left(\frac{(1+2\sqrt{1+\sqrt{1+x}})\sqrt{5}}{5}\right)\sqrt{5}}{5}$	34
default	$2\sqrt{1+x} + 2 + \frac{8 \operatorname{arctanh}\left(\frac{(1+2\sqrt{1+\sqrt{1+x}})\sqrt{5}}{5}\right)\sqrt{5}}{5}$	34

input `int((1+x)^(1/2)/(x+(1+(1+x)^(1/2))^(1/2)),x,method=_RETURNVERBOSE)`

output `2*(1+x)^(1/2)+2+8/5*arctanh(1/5*(1+2*(1+(1+x)^(1/2))^(1/2))*5^(1/2))*5^(1/2)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 101 vs. $2(32) = 64$.

Time = 0.07 (sec) , antiderivative size = 101, normalized size of antiderivative = 2.46

$$\int \frac{\sqrt{1+x}}{x + \sqrt{1+\sqrt{1+x}}} dx$$

$$= \frac{4}{5} \sqrt{5} \log \left(\frac{2x^2 - \sqrt{5}(3x+1) - (\sqrt{5}(x+2) - 5x)\sqrt{x+1} + (\sqrt{5}(x+2) + (\sqrt{5}(2x-1) - 5)\sqrt{x+1} - 5x)\sqrt{x+1}}{x^2 - x - 1} \right) + 2\sqrt{x+1}$$

input `integrate((1+x)^(1/2)/(x+(1+(1+x)^(1/2))^(1/2)),x, algorithm="fricas")`

output `4/5*sqrt(5)*log((2*x^2 - sqrt(5)*(3*x + 1) - (sqrt(5)*(x + 2) - 5*x)*sqrt(x + 1) + (sqrt(5)*(x + 2) + (sqrt(5)*(2*x - 1) - 5)*sqrt(x + 1) - 5*x)*sqrt(x + 1) + 3*x + 3)/(x^2 - x - 1)) + 2*sqrt(x + 1)`

Sympy [A] (verification not implemented)

Time = 4.68 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.59

$$\int \frac{\sqrt{1+x}}{x + \sqrt{1+\sqrt{1+x}}} dx$$

$$= 2\sqrt{x+1} - \frac{4\sqrt{5} \left(-\log \left(\sqrt{\sqrt{x+1}+1} + \frac{1}{2} + \frac{\sqrt{5}}{2} \right) + \log \left(\sqrt{\sqrt{x+1}+1} - \frac{\sqrt{5}}{2} + \frac{1}{2} \right) \right)}{5} + 2$$

input `integrate((1+x)**(1/2)/(x+(1+(1+x)**(1/2))**(1/2)),x)`

output `2*sqrt(x + 1) - 4*sqrt(5)*(-log(sqrt(sqrt(x + 1) + 1) + 1/2 + sqrt(5)/2) + log(sqrt(sqrt(x + 1) + 1) - sqrt(5)/2 + 1/2))/5 + 2`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.24

$$\int \frac{\sqrt{1+x}}{x + \sqrt{1 + \sqrt{1+x}}} dx = -\frac{4}{5} \sqrt{5} \log \left(-\frac{\sqrt{5} - 2 \sqrt{\sqrt{x+1} + 1} - 1}{\sqrt{5} + 2 \sqrt{\sqrt{x+1} + 1} + 1} \right) + 2\sqrt{x+1} + 2$$

input `integrate((1+x)^(1/2)/(x+(1+(1+x)^(1/2))^(1/2)),x, algorithm="maxima")`

output `-4/5*sqrt(5)*log(-(sqrt(5) - 2*sqrt(sqrt(x + 1) + 1) - 1)/(sqrt(5) + 2*sqrt(sqrt(x + 1) + 1) + 1)) + 2*sqrt(x + 1) + 2`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.24

$$\int \frac{\sqrt{1+x}}{x + \sqrt{1 + \sqrt{1+x}}} dx = -\frac{4}{5} \sqrt{5} \log \left(-\frac{\sqrt{5} - 2 \sqrt{\sqrt{x+1} + 1} - 1}{\sqrt{5} + 2 \sqrt{\sqrt{x+1} + 1} + 1} \right) + 2\sqrt{x+1} + 2$$

input `integrate((1+x)^(1/2)/(x+(1+(1+x)^(1/2))^(1/2)),x, algorithm="giac")`

output `-4/5*sqrt(5)*log(-(sqrt(5) - 2*sqrt(sqrt(x + 1) + 1) - 1)/(sqrt(5) + 2*sqrt(sqrt(x + 1) + 1) + 1)) + 2*sqrt(x + 1) + 2`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{1+x}}{x + \sqrt{1 + \sqrt{1+x}}} dx = \int \frac{\sqrt{x+1}}{x + \sqrt{\sqrt{x+1} + 1}} dx$$

input `int((x + 1)^(1/2)/(x + ((x + 1)^(1/2) + 1)^(1/2)),x)`

output `int((x + 1)^(1/2)/(x + ((x + 1)^(1/2) + 1)^(1/2)), x)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.12

$$\int \frac{\sqrt{1+x}}{x + \sqrt{1+\sqrt{1+x}}} dx = 2\sqrt{x+1} - \frac{4\sqrt{5} \log\left(2\sqrt{\sqrt{x+1}+1} - \sqrt{5}+1\right)}{5} + \frac{4\sqrt{5} \log\left(2\sqrt{\sqrt{x+1}+1} + \sqrt{5}+1\right)}{5} + 2$$

input

```
int((1+x)^(1/2)/(x+(1+(1+x)^(1/2))^(1/2)),x)
```

output

```
(2*(5*sqrt(x + 1) - 2*sqrt(5)*log(2*sqrt(sqrt(x + 1) + 1) - sqrt(5) + 1) + 2*sqrt(5)*log(2*sqrt(sqrt(x + 1) + 1) + sqrt(5) + 1) + 5))/5
```

3.11 $\int \frac{1}{x - \sqrt{1 + \sqrt{1 + x}}} dx$

Optimal result	107
Mathematica [A] (verified)	107
Rubi [A] (verified)	108
Maple [A] (verified)	109
Fricas [B] (verification not implemented)	110
Sympy [A] (verification not implemented)	111
Maxima [A] (verification not implemented)	111
Giac [A] (verification not implemented)	112
Mupad [F(-1)]	112
Reduce [B] (verification not implemented)	113

Optimal result

Integrand size = 19, antiderivative size = 73

$$\int \frac{1}{x - \sqrt{1 + \sqrt{1 + x}}} dx = \frac{2}{5} (5 + \sqrt{5}) \log \left(1 - \sqrt{5} - 2\sqrt{1 + \sqrt{1 + x}} \right) + \frac{2}{5} (5 - \sqrt{5}) \log \left(1 + \sqrt{5} - 2\sqrt{1 + \sqrt{1 + x}} \right)$$

output

```
2/5*ln(1+5^(1/2)-2*(1+(1+x)^(1/2))^(1/2))*(5-5^(1/2))+2/5*ln(1-5^(1/2)-2*(1+(1+x)^(1/2))^(1/2))*(5+5^(1/2))
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.95

$$\int \frac{1}{x - \sqrt{1 + \sqrt{1 + x}}} dx = -\frac{2}{5} (-5 + \sqrt{5}) \log \left(1 + \sqrt{5} - 2\sqrt{1 + \sqrt{1 + x}} \right) + \frac{2}{5} (5 + \sqrt{5}) \log \left(-1 + \sqrt{5} + 2\sqrt{1 + \sqrt{1 + x}} \right)$$

input

```
Integrate[(x - Sqrt[1 + Sqrt[1 + x]])^(-1), x]
```

output

$$\frac{(-2*(-5 + \text{Sqrt}[5])*\text{Log}[1 + \text{Sqrt}[5] - 2*\text{Sqrt}[1 + \text{Sqrt}[1 + x]])/5 + (2*(5 + \text{Sqrt}[5])*\text{Log}[-1 + \text{Sqrt}[5] + 2*\text{Sqrt}[1 + \text{Sqrt}[1 + x]])/5}$$
Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.03, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {7267, 25, 7267, 25, 1141, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{x - \sqrt{\sqrt{x+1}+1}} dx \\ & \quad \downarrow \text{7267} \\ & 2 \int -\frac{\sqrt{x+1}}{\sqrt{\sqrt{x+1}+1}-x} d\sqrt{x+1} \\ & \quad \downarrow \text{25} \\ & -2 \int \frac{\sqrt{x+1}}{\sqrt{\sqrt{x+1}+1}-x} d\sqrt{x+1} \\ & \quad \downarrow \text{7267} \\ & -4 \int -\frac{1 - \sqrt{\sqrt{x+1}+1}}{\sqrt{\sqrt{x+1}+1}-x} d\sqrt{\sqrt{x+1}+1} \\ & \quad \downarrow \text{25} \\ & 4 \int \frac{1 - \sqrt{\sqrt{x+1}+1}}{\sqrt{\sqrt{x+1}+1}-x} d\sqrt{\sqrt{x+1}+1} \\ & \quad \downarrow \text{1141} \\ & -4 \int \left(\frac{5 + \sqrt{5}}{5(-2\sqrt{\sqrt{x+1}+1} - \sqrt{5} + 1)} - \frac{1 - \sqrt{5}}{-2\sqrt{5}\sqrt{\sqrt{x+1}+1} + \sqrt{5} + 5} \right) d\sqrt{\sqrt{x+1}+1} \\ & \quad \downarrow \text{2009} \end{aligned}$$

$$-4\left(-\frac{1}{10}(5 + \sqrt{5}) \log\left(-2\sqrt{\sqrt{x+1}+1} - \sqrt{5} + 1\right) - \frac{1}{10}(5 - \sqrt{5}) \log\left(-2\sqrt{\sqrt{x+1}+1} + \sqrt{5} + 1\right)\right)$$

input `Int[(x - Sqrt[1 + Sqrt[1 + x]])^(-1),x]`

output `-4*(-1/10*((5 + Sqrt[5])*Log[1 - Sqrt[5] - 2*Sqrt[1 + Sqrt[1 + x]]]) - ((5 - Sqrt[5])*Log[1 + Sqrt[5] - 2*Sqrt[1 + Sqrt[1 + x]]])/10)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 1141 `Int[((d_.) + (e_.)*(x_))^(m_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[1/c^p Int[ExpandIntegrand[(d + e*x)^m*(b/2 - q/2 + c*x)^p*(b/2 + q/2 + c*x)^p, x], x] /; EqQ[p, -1] || !FractionalPowerFactorQ[q] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[p, 0] && IntegerQ[m] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7267 `Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Simp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2]])], x] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]`

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.63

method	result
derivativedivides	$2 \ln \left(\sqrt{1+x} - \sqrt{1+\sqrt{1+x}} \right) + \frac{4\sqrt{5} \operatorname{arctanh} \left(\frac{(2\sqrt{1+\sqrt{1+x}}-1)\sqrt{5}}{5} \right)}{5}$
default	$\frac{\ln(x^2-x-1)}{2} + \frac{\sqrt{5} \operatorname{arctanh} \left(\frac{(2x-1)\sqrt{5}}{5} \right)}{5} - \ln \left(\sqrt{1+x} + \sqrt{1+\sqrt{1+x}} \right) + \frac{2 \operatorname{arctanh} \left(\frac{(1+2\sqrt{1+\sqrt{1+x}})}{5} \right)}{5}$

input `int(1/(x-(1+(1+x)^(1/2))^(1/2)),x,method=_RETURNVERBOSE)`

output `2*ln((1+x)^(1/2)-(1+(1+x)^(1/2))^(1/2))+4/5*5^(1/2)*arctanh(1/5*(2*(1+(1+x)^(1/2))^(1/2)-1)*5^(1/2))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 112 vs. $2(51) = 102$.

Time = 0.08 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.53

$$\int \frac{1}{x - \sqrt{1 + \sqrt{1+x}}} dx$$

$$= \frac{2}{5} \sqrt{5} \log \left(\frac{2x^2 + \sqrt{5}(3x+1) + (\sqrt{5}(x+2) + 5x)\sqrt{x+1} + (\sqrt{5}(x+2) + (\sqrt{5}(2x-1) + 5)\sqrt{x+1} + 5x)\sqrt{x+1} + 5}{x^2 - x - 1} \right)$$

$$+ 2 \log \left(\sqrt{x+1} - \sqrt{\sqrt{x+1} + 1} \right)$$

input `integrate(1/(x-(1+(1+x)^(1/2))^(1/2)),x, algorithm="fricas")`

output `2/5*sqrt(5)*log((2*x^2 + sqrt(5)*(3*x + 1) + (sqrt(5)*(x + 2) + 5*x)*sqrt(x + 1) + (sqrt(5)*(x + 2) + (sqrt(5)*(2*x - 1) + 5)*sqrt(x + 1) + 5*x)*sqrt(sqrt(x + 1) + 1) + 3*x + 3)/(x^2 - x - 1)) + 2*log(sqrt(x + 1) - sqrt(sqrt(x + 1) + 1))`

Sympy [A] (verification not implemented)

Time = 2.40 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.04

$$\int \frac{1}{x - \sqrt{1 + \sqrt{1 + x}}} dx$$

$$= -\frac{2\sqrt{5}\left(-\log\left(\sqrt{\sqrt{x+1}+1} - \frac{1}{2} + \frac{\sqrt{5}}{2}\right) + \log\left(\sqrt{\sqrt{x+1}+1} - \frac{\sqrt{5}}{2} - \frac{1}{2}\right)\right)}{5}$$

$$+ 2\log\left(\sqrt{x+1} - \sqrt{\sqrt{x+1}+1}\right)$$

input `integrate(1/(x-(1+(1+x)**(1/2))**(1/2)),x)`output `-2*sqrt(5)*(-log(sqrt(sqrt(x + 1) + 1) - 1/2 + sqrt(5)/2) + log(sqrt(sqrt(x + 1) + 1) - sqrt(5)/2 - 1/2))/5 + 2*log(sqrt(x + 1) - sqrt(sqrt(x + 1) + 1))`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.86

$$\int \frac{1}{x - \sqrt{1 + \sqrt{1 + x}}} dx = -\frac{2}{5}\sqrt{5}\log\left(-\frac{\sqrt{5} - 2\sqrt{\sqrt{x+1}+1} + 1}{\sqrt{5} + 2\sqrt{\sqrt{x+1}+1} - 1}\right)$$

$$+ 2\log\left(\sqrt{x+1} - \sqrt{\sqrt{x+1}+1}\right)$$

input `integrate(1/(x-(1+(1+x)^(1/2))^(1/2)),x, algorithm="maxima")`output `-2/5*sqrt(5)*log(-(sqrt(5) - 2*sqrt(sqrt(x + 1) + 1) + 1)/(sqrt(5) + 2*sqrt(sqrt(x + 1) + 1) - 1)) + 2*log(sqrt(x + 1) - sqrt(sqrt(x + 1) + 1))`

Giac [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.92

$$\int \frac{1}{x - \sqrt{1 + \sqrt{1 + x}}} dx = -\frac{2}{5} \sqrt{5} \log \left(\frac{\left| -\sqrt{5} + 2 \sqrt{\sqrt{x+1} + 1} - 1 \right|}{\left| \sqrt{5} + 2 \sqrt{\sqrt{x+1} + 1} - 1 \right|} \right) + 2 \log \left(\left| \sqrt{x+1} - \sqrt{\sqrt{x+1} + 1} \right| \right)$$

input `integrate(1/(x-(1+(1+x)^(1/2))^(1/2)),x, algorithm="giac")`

output `-2/5*sqrt(5)*log(abs(-sqrt(5) + 2*sqrt(sqrt(x + 1) + 1) - 1)/abs(sqrt(5) + 2*sqrt(sqrt(x + 1) + 1) - 1)) + 2*log(abs(sqrt(x + 1) - sqrt(sqrt(x + 1) + 1)))`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{x - \sqrt{1 + \sqrt{1 + x}}} dx = \int \frac{1}{x - \sqrt{\sqrt{x+1} + 1}} dx$$

input `int(1/(x - ((x + 1)^(1/2) + 1)^(1/2)),x)`

output `int(1/(x - ((x + 1)^(1/2) + 1)^(1/2)), x)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.00

$$\int \frac{1}{x - \sqrt{1 + \sqrt{1 + x}}} dx = -\frac{2\sqrt{5} \log\left(2\sqrt{\sqrt{x+1}+1} - \sqrt{5} - 1\right)}{5}$$

$$+ \frac{2\sqrt{5} \log\left(2\sqrt{\sqrt{x+1}+1} + \sqrt{5} - 1\right)}{5}$$

$$+ 2 \log\left(2\sqrt{\sqrt{x+1}+1} - \sqrt{5} - 1\right)$$

$$+ 2 \log\left(2\sqrt{\sqrt{x+1}+1} + \sqrt{5} - 1\right)$$

input `int(1/(x-(1+(1+x)^(1/2))^(1/2)),x)`output `(2*(- sqrt(5)*log(2*sqrt(sqrt(x + 1) + 1) - sqrt(5) - 1) + sqrt(5)*log(2*sqrt(sqrt(x + 1) + 1) + sqrt(5) - 1) + 5*log(2*sqrt(sqrt(x + 1) + 1) - sqrt(5) - 1) + 5*log(2*sqrt(sqrt(x + 1) + 1) + sqrt(5) - 1)))/5`

3.12 $\int \frac{x}{x + \sqrt{1 - \sqrt{1+x}}} dx$

Optimal result	114
Mathematica [A] (verified)	114
Rubi [A] (warning: unable to verify)	115
Maple [A] (verified)	116
Fricas [A] (verification not implemented)	117
Sympy [A] (verification not implemented)	117
Maxima [A] (verification not implemented)	118
Giac [A] (verification not implemented)	118
Mupad [F(-1)]	119
Reduce [B] (verification not implemented)	119

Optimal result

Integrand size = 21, antiderivative size = 73

$$\int \frac{x}{x + \sqrt{1 - \sqrt{1+x}}} dx = 2\sqrt{1+x} - 4\sqrt{1 - \sqrt{1+x}} + (1 - \sqrt{1+x})^2 + \frac{8\operatorname{arctanh}\left(\frac{1+2\sqrt{1-\sqrt{1+x}}}{\sqrt{5}}\right)}{\sqrt{5}}$$

output

```
8/5*arctanh(1/5*(1+2*(1-(1+x)^(1/2))^(1/2))*5^(1/2))*5^(1/2)+(1-(1+x)^(1/2))^2+2*(1+x)^(1/2)-4*(1-(1+x)^(1/2))^(1/2)
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.71

$$\int \frac{x}{x + \sqrt{1 - \sqrt{1+x}}} dx = x - 4\sqrt{1 - \sqrt{1+x}} + \frac{8\operatorname{arctanh}\left(\frac{1+2\sqrt{1-\sqrt{1+x}}}{\sqrt{5}}\right)}{\sqrt{5}}$$

input

```
Integrate[x/(x + Sqrt[1 - Sqrt[1 + x]]),x]
```

output

$$x - 4\sqrt{1 - \sqrt{1 + x}} + (8\operatorname{ArcTanh}[(1 + 2\sqrt{1 - \sqrt{1 + x}})/\sqrt{t[5]})]/\sqrt{t[5]})/\sqrt{5}$$
Rubi [A] (warning: unable to verify)

Time = 0.51 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.41, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {7267, 7267, 2159, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x}{x + \sqrt{1 - \sqrt{x+1}}} dx \\ & \quad \downarrow \text{7267} \\ & 2 \int -\frac{x\sqrt{x+1}}{-x - \sqrt{1 - \sqrt{x+1}}} d\sqrt{x+1} \\ & \quad \downarrow \text{7267} \\ & 4 \int \frac{(1-x)(x+1)(\sqrt{1 - \sqrt{x+1}} + 1)}{-x - \sqrt{1 - \sqrt{x+1}}} d\sqrt{1 - \sqrt{x+1}} \\ & \quad \downarrow \text{2159} \\ & 4 \int \left((x+1)^{3/2} - \sqrt{1 - \sqrt{x+1}} + \frac{1}{-x - \sqrt{1 - \sqrt{x+1}}} - 1 \right) d\sqrt{1 - \sqrt{x+1}} \\ & \quad \downarrow \text{2009} \\ & 4 \left(\frac{1}{4}(x+1)^2 + \frac{1}{2}(-x-1) - \sqrt{1 - \sqrt{x+1}} - \frac{\log(2\sqrt{1 - \sqrt{x+1}} - \sqrt{5} + 1)}{\sqrt{5}} + \frac{\log(2\sqrt{1 - \sqrt{x+1}} + \sqrt{5} + 1)}{\sqrt{5}} \right) \end{aligned}$$

input

$$\operatorname{Int}[x/(x + \sqrt{1 - \sqrt{1 + x}}), x]$$

output

```
4*((-1 - x)/2 + (1 + x)^2/4 - Sqrt[1 - Sqrt[1 + x]] - Log[1 - Sqrt[5] + 2*
Sqrt[1 - Sqrt[1 + x]]]/Sqrt[5] + Log[1 + Sqrt[5] + 2*Sqrt[1 - Sqrt[1 + x]]
]/Sqrt[5])
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2159

```
Int[(Pq_)*((d_) + (e_)*(x_)^(m_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p
_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*Pq*(a + b*x + c*x^2)^p, x
], x] /; FreeQ[{a, b, c, d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]
```

rule 7267

```
Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Si
mp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])]], x
] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]
```

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$(1 - \sqrt{1+x})^2 + 2\sqrt{1+x} - 2 - 4\sqrt{1 - \sqrt{1+x}} + \frac{8 \operatorname{arctanh}\left(\frac{(1+2\sqrt{1-\sqrt{1+x}})\sqrt{5}}{5}\right)\sqrt{5}}{5}$	60
default	$(1 - \sqrt{1+x})^2 + 2\sqrt{1+x} - 2 - 4\sqrt{1 - \sqrt{1+x}} + \frac{8 \operatorname{arctanh}\left(\frac{(1+2\sqrt{1-\sqrt{1+x}})\sqrt{5}}{5}\right)\sqrt{5}}{5}$	60

input

```
int(x/(x+(1-(1+x)^(1/2))^(1/2)),x,method=_RETURNVERBOSE)
```

output

```
(1-(1+x)^(1/2))^2+2*(1+x)^(1/2)-2-4*(1-(1+x)^(1/2))^(1/2)+8/5*arctanh(1/5*
(1+2*(1-(1+x)^(1/2))^(1/2))*5^(1/2))*5^(1/2)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.51

$$\int \frac{x}{x + \sqrt{1 - \sqrt{1 + x}}} dx$$

$$= \frac{4}{5} \sqrt{5} \log \left(\frac{2x^2 - \sqrt{5}(3x + 1) + (\sqrt{5}(x + 2) - 5x)\sqrt{x + 1} + (\sqrt{5}(x + 2) - (\sqrt{5}(2x - 1) - 5)\sqrt{x + 1} - 5x - 4\sqrt{-\sqrt{x + 1} + 1}}}{x^2 - x - 1} \right)$$

input `integrate(x/(x+(1-(1+x)^(1/2))^(1/2)),x, algorithm="fricas")`

output `4/5*sqrt(5)*log((2*x^2 - sqrt(5)*(3*x + 1) + (sqrt(5)*(x + 2) - 5*x)*sqrt(x + 1) + (sqrt(5)*(x + 2) - (sqrt(5)*(2*x - 1) - 5)*sqrt(x + 1) - 5*x)*sqrt(-sqrt(x + 1) + 1) + 3*x + 3)/(x^2 - x - 1)) + x - 4*sqrt(-sqrt(x + 1) + 1)`

Sympy [A] (verification not implemented)

Time = 3.94 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.21

$$\int \frac{x}{x + \sqrt{1 - \sqrt{1 + x}}} dx$$

$$= -4\sqrt{1 - \sqrt{x + 1}} + (1 - \sqrt{x + 1})^2 + 2\sqrt{x + 1}$$

$$- \frac{4\sqrt{5} \left(-\log \left(\sqrt{1 - \sqrt{x + 1}} + \frac{1}{2} + \frac{\sqrt{5}}{2} \right) + \log \left(\sqrt{1 - \sqrt{x + 1}} - \frac{\sqrt{5}}{2} + \frac{1}{2} \right) \right)}{5} - 2$$

input `integrate(x/(x+(1-(1+x)**(1/2))**(1/2)),x)`

output `-4*sqrt(1 - sqrt(x + 1)) + (1 - sqrt(x + 1))**2 + 2*sqrt(x + 1) - 4*sqrt(5)*(-log(sqrt(1 - sqrt(x + 1)) + 1/2 + sqrt(5)/2) + log(sqrt(1 - sqrt(x + 1)) - sqrt(5)/2 + 1/2))/5 - 2`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.05

$$\int \frac{x}{x + \sqrt{1 - \sqrt{1 + x}}} dx = (\sqrt{x + 1} - 1)^2 - \frac{4}{5} \sqrt{5} \log \left(-\frac{\sqrt{5} - 2\sqrt{-\sqrt{x + 1} + 1} - 1}{\sqrt{5} + 2\sqrt{-\sqrt{x + 1} + 1} + 1} \right) + 2\sqrt{x + 1} - 4\sqrt{-\sqrt{x + 1} + 1} - 2$$

input `integrate(x/(x+(1-(1+x)^(1/2))^(1/2)),x, algorithm="maxima")`

output `(sqrt(x + 1) - 1)^2 - 4/5*sqrt(5)*log(-(sqrt(5) - 2*sqrt(-sqrt(x + 1) + 1) - 1)/(sqrt(5) + 2*sqrt(-sqrt(x + 1) + 1) + 1)) + 2*sqrt(x + 1) - 4*sqrt(-sqrt(x + 1) + 1) - 2`

Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.08

$$\int \frac{x}{x + \sqrt{1 - \sqrt{1 + x}}} dx = (\sqrt{x + 1} - 1)^2 - \frac{4}{5} \sqrt{5} \log \left(\frac{|-\sqrt{5} + 2\sqrt{-\sqrt{x + 1} + 1} + 1|}{\sqrt{5} + 2\sqrt{-\sqrt{x + 1} + 1} + 1} \right) + 2\sqrt{x + 1} - 4\sqrt{-\sqrt{x + 1} + 1} - 2$$

input `integrate(x/(x+(1-(1+x)^(1/2))^(1/2)),x, algorithm="giac")`

output `(sqrt(x + 1) - 1)^2 - 4/5*sqrt(5)*log(abs(-sqrt(5) + 2*sqrt(-sqrt(x + 1) + 1) + 1)/(sqrt(5) + 2*sqrt(-sqrt(x + 1) + 1) + 1)) + 2*sqrt(x + 1) - 4*sqrt(-sqrt(x + 1) + 1) - 2`

Mupad [F(-1)]

Timed out.

$$\int \frac{x}{x + \sqrt{1 - \sqrt{1+x}}} dx = \int \frac{x}{x + \sqrt{1 - \sqrt{x+1}}} dx$$

input `int(x/(x + (1 - (x + 1)^(1/2))^(1/2)),x)`output `int(x/(x + (1 - (x + 1)^(1/2))^(1/2)), x)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.75

$$\int \frac{x}{x + \sqrt{1 - \sqrt{1+x}}} dx = -4\sqrt{-\sqrt{x+1}+1} - \frac{4\sqrt{5} \log(2\sqrt{-\sqrt{x+1}+1} - \sqrt{5}+1)}{5} + \frac{4\sqrt{5} \log(2\sqrt{-\sqrt{x+1}+1} + \sqrt{5}+1)}{5} + x$$

input `int(x/(x+(1-(1+x)^(1/2))^(1/2)),x)`output `(- 20*sqrt(- sqrt(x + 1) + 1) - 4*sqrt(5)*log(2*sqrt(- sqrt(x + 1) + 1) - sqrt(5) + 1) + 4*sqrt(5)*log(2*sqrt(- sqrt(x + 1) + 1) + sqrt(5) + 1) + 5*x)/5`

3.13 $\int \frac{\sqrt{x+\sqrt{1+x}}}{\sqrt{1+x}(1+x^2)} dx$

Optimal result	120
Mathematica [C] (verified)	121
Rubi [A] (verified)	122
Maple [C] (verified)	124
Fricas [B] (verification not implemented)	124
Sympy [F]	125
Maxima [F]	125
Giac [F(-2)]	125
Mupad [F(-1)]	126
Reduce [F]	126

Optimal result

Integrand size = 28, antiderivative size = 365

$$\int \frac{\sqrt{x+\sqrt{1+x}}}{\sqrt{1+x}(1+x^2)} dx = -\frac{i \arctan\left(\frac{2+\sqrt{1-i}-(1-2\sqrt{1-i})\sqrt{1+x}}{2\sqrt{i+\sqrt{1-i}}\sqrt{x+\sqrt{1+x}}}\right)}{2\sqrt{\frac{1-i}{i+\sqrt{1-i}}}} + \frac{i \arctan\left(\frac{2+\sqrt{1+i}-(1-2\sqrt{1+i})\sqrt{1+x}}{2\sqrt{-i+\sqrt{1+i}}\sqrt{x+\sqrt{1+x}}}\right)}{2\sqrt{-\frac{1+i}{i-\sqrt{1+i}}}} + \frac{i \operatorname{arctanh}\left(\frac{2-\sqrt{1-i}-(1+2\sqrt{1-i})\sqrt{1+x}}{2\sqrt{-i+\sqrt{1-i}}\sqrt{x+\sqrt{1+x}}}\right)}{2\sqrt{-\frac{1-i}{i-\sqrt{1-i}}}} - \frac{i \operatorname{arctanh}\left(\frac{2-\sqrt{1+i}-(1+2\sqrt{1+i})\sqrt{1+x}}{2\sqrt{i+\sqrt{1+i}}\sqrt{x+\sqrt{1+x}}}\right)}{2\sqrt{\frac{1+i}{i+\sqrt{1+i}}}}$$

output

```

1/2*I*arctanh(1/2*(2-(1-I)^(1/2)-(1+2*(1-I)^(1/2))*(1+x)^(1/2))/(-I+(1-I)^(1/2))^(1/2)/(x+(1+x)^(1/2))^(1/2))/((-1+I)/(I-(1-I)^(1/2)))^(1/2)-1/2*I*arctan(1/2*(2+(1-I)^(1/2)-(1-2*(1-I)^(1/2))*(1+x)^(1/2))/(I+(1-I)^(1/2))^(1/2)/(x+(1+x)^(1/2))^(1/2))/((1-I)/(I+(1-I)^(1/2)))^(1/2)+1/2*I*arctan(1/2*(2+(1+I)^(1/2)-(1-2*(1+I)^(1/2))*(1+x)^(1/2))/(-I+(1+I)^(1/2))^(1/2)/(x+(1+x)^(1/2))^(1/2))/((-1-I)/(I-(1+I)^(1/2)))^(1/2)-1/2*I*arctanh(1/2*(2-(1+I)^(1/2)-(1+2*(1+I)^(1/2))*(1+x)^(1/2))/(I+(1+I)^(1/2))^(1/2)/(x+(1+x)^(1/2))^(1/2))/((1+I)/(I+(1+I)^(1/2)))^(1/2)

```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.16 (sec) , antiderivative size = 217, normalized size of antiderivative = 0.59

$$\int \frac{\sqrt{x + \sqrt{1+x}}}{\sqrt{1+x}(1+x^2)} dx = -\frac{1}{2} \text{RootSum} \left[1 - 8\#1 + 40\#1^2 - 48\#1^3 + 20\#1^4 + 8\#1^5 - 4\#1^6 + \#1^8 \&, \frac{-\log\left(-\sqrt{1+x} + \sqrt{x + \sqrt{1+x}} - \#1\right) + 5 \log\left(-\sqrt{1+x} + \sqrt{x + \sqrt{1+x}} - \#1\right) \#1^2 - 5\#1^3}{-1 + 10\#1 - 18\#1^2 + 10\#1^3 - \#1^4} \right]$$

input

```
Integrate[Sqrt[x + Sqrt[1 + x]]/(Sqrt[1 + x]*(1 + x^2)),x]
```

output

```

-1/2*RootSum[1 - 8*#1 + 40*#1^2 - 48*#1^3 + 20*#1^4 + 8*#1^5 - 4*#1^6 + #1^8 & , (-Log[-Sqrt[1 + x] + Sqrt[x + Sqrt[1 + x]] - #1] + 5*Log[-Sqrt[1 + x] + Sqrt[x + Sqrt[1 + x]] - #1]*#1^2 - 5*Log[-Sqrt[1 + x] + Sqrt[x + Sqrt[1 + x]] - #1]*#1^4 + 2*Log[-Sqrt[1 + x] + Sqrt[x + Sqrt[1 + x]] - #1]*#1^5)/(-1 + 10*#1 - 18*#1^2 + 10*#1^3 + 5*#1^4 - 3*#1^5 + #1^7) & ]

```

Rubi [A] (verified)

Time = 0.99 (sec) , antiderivative size = 367, normalized size of antiderivative = 1.01, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {7267, 7292, 7279, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{x + \sqrt{x + 1}}}{\sqrt{x + 1} (x^2 + 1)} dx \\
 & \quad \downarrow \text{7267} \\
 & 2 \int \frac{\sqrt{x + \sqrt{x + 1}}}{x^2 + 1} d\sqrt{x + 1} \\
 & \quad \downarrow \text{7292} \\
 & 2 \int \frac{\sqrt{x + \sqrt{x + 1}}}{(x + 1)^2 - 2(x + 1) + 2} d\sqrt{x + 1} \\
 & \quad \downarrow \text{7279} \\
 & 2 \int \left(\frac{i\sqrt{x + \sqrt{x + 1}}}{(2 + 2i) - 2(x + 1)} + \frac{i\sqrt{x + \sqrt{x + 1}}}{2(x + 1) - (2 - 2i)} \right) d\sqrt{x + 1} \\
 & \quad \downarrow \text{2009} \\
 & 2 \left(\frac{i \arctan \left(\frac{-((1 - 2\sqrt{1 - i})\sqrt{x + 1}) + \sqrt{1 - i} + 2}{2\sqrt{i + \sqrt{1 - i}}\sqrt{x + \sqrt{x + 1}}} \right)}{4\sqrt{\frac{1 - i}{i + \sqrt{1 - i}}}} + \frac{i \arctan \left(\frac{-((1 - 2\sqrt{1 + i})\sqrt{x + 1}) + \sqrt{1 + i} + 2}{2\sqrt{\sqrt{1 + i} - i}\sqrt{x + \sqrt{x + 1}}} \right)}{4\sqrt{-\frac{1 + i}{i - \sqrt{1 + i}}}} + \frac{i \operatorname{arctanh} \left(\frac{-((1 + 2\sqrt{1 - i})\sqrt{x + 1}) + \sqrt{1 - i} + 2}{2\sqrt{\sqrt{1 - i} - i}\sqrt{x + \sqrt{x + 1}}} \right)}{4\sqrt{-\frac{1 - i}{i - \sqrt{1 - i}}}} \right)
 \end{aligned}$$

input

```
Int[Sqrt[x + Sqrt[1 + x]]/(Sqrt[1 + x]*(1 + x^2)),x]
```

output

```
2*(((1/4)*ArcTan[(2 + Sqrt[1 - I] - (1 - 2*Sqrt[1 - I])*Sqrt[1 + x])/(2
*Sqrt[1 + Sqrt[1 - I]]*Sqrt[x + Sqrt[1 + x]])]/Sqrt[(1 - I)/(1 + Sqrt[1 -
I])]) + ((1/4)*ArcTan[(2 + Sqrt[1 + I] - (1 - 2*Sqrt[1 + I])*Sqrt[1 + x])/
(2*Sqrt[-1 + Sqrt[1 + I]]*Sqrt[x + Sqrt[1 + x]])]/Sqrt[(-1 - I)/(1 - Sqrt
[1 + I])]) + ((1/4)*ArcTanh[(2 - Sqrt[1 - I] - (1 + 2*Sqrt[1 - I])*Sqrt[1 +
x])/(2*Sqrt[-1 + Sqrt[1 - I]]*Sqrt[x + Sqrt[1 + x]])]/Sqrt[(-1 + I)/(1 -
Sqrt[1 - I])]) - ((1/4)*ArcTanh[(2 - Sqrt[1 + I] - (1 + 2*Sqrt[1 + I])*Sqr
t[1 + x])/(2*Sqrt[1 + Sqrt[1 + I]]*Sqrt[x + Sqrt[1 + x]])]/Sqrt[(1 + I)/(
1 + Sqrt[1 + I])])
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 7267

```
Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Si
mp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])], x
] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]]
```

rule 7279

```
Int[(u_)/((a_.) + (b_.)*(x_)^(n_.) + (c_.)*(x_)^(n2_.)), x_Symbol] := With[
{v = RationalFunctionExpand[u/(a + b*x^n + c*x^(2*n)), x]}, Int[v, x] /; Su
mQ[v] /; FreeQ[{a, b, c}, x] && EqQ[n2, 2*n] && IGtQ[n, 0]
```

rule 7292

```
Int[u_, x_Symbol] := With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v !=
u]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.12 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.30

method	result
derivativedivides	$-\frac{\left(\sum_{R=\text{RootOf}(_Z^8-4_Z^6+8_Z^5+20_Z^4-48_Z^3+40_Z^2-8_Z+1)} \frac{(2_R^5-5_R^4+5_R^2-1) \ln(\sqrt{x+\sqrt{1+x}}-\sqrt{1-x})}{_R^7-3_R^5+5_R^4+10_R^3-18_R^2+10_R-1}\right)}{2}$
default	$-\frac{\left(\sum_{R=\text{RootOf}(_Z^8-4_Z^6+8_Z^5+20_Z^4-48_Z^3+40_Z^2-8_Z+1)} \frac{(2_R^5-5_R^4+5_R^2-1) \ln(\sqrt{x+\sqrt{1+x}}-\sqrt{1-x})}{_R^7-3_R^5+5_R^4+10_R^3-18_R^2+10_R-1}\right)}{2}$

input `int((x+(1+x)^(1/2))^(1/2)/(x^2+1)/(1+x)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/2*sum((2*_R^5-5*_R^4+5*_R^2-1)/(_R^7-3*_R^5+5*_R^4+10*_R^3-18*_R^2+10*_R-1)*ln((x+(1+x)^(1/2))^(1/2)-(1+x)^(1/2)-_R),_R=RootOf(_Z^8-4*_Z^6+8*_Z^5+20*_Z^4-48*_Z^3+40*_Z^2-8*_Z+1))`

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 5235 vs. $2(201) = 402$.

Time = 5.68 (sec) , antiderivative size = 5235, normalized size of antiderivative = 14.34

$$\int \frac{\sqrt{x + \sqrt{1+x}}}{\sqrt{1+x}(1+x^2)} dx = \text{Too large to display}$$

input `integrate((x+(1+x)^(1/2))^(1/2)/(x^2+1)/(1+x)^(1/2),x, algorithm="fricas")`

output `Too large to include`

Sympy [F]

$$\int \frac{\sqrt{x + \sqrt{1+x}}}{\sqrt{1+x}(1+x^2)} dx = \int \frac{\sqrt{x + \sqrt{x+1}}}{\sqrt{x+1}(x^2+1)} dx$$

input `integrate((x+(1+x)**(1/2))**(1/2)/(x**2+1)/(1+x)**(1/2), x)`

output `Integral(sqrt(x + sqrt(x + 1))/(sqrt(x + 1)*(x**2 + 1)), x)`

Maxima [F]

$$\int \frac{\sqrt{x + \sqrt{1+x}}}{\sqrt{1+x}(1+x^2)} dx = \int \frac{\sqrt{x + \sqrt{x+1}}}{(x^2+1)\sqrt{x+1}} dx$$

input `integrate((x+(1+x)^(1/2))^(1/2)/(x^2+1)/(1+x)^(1/2), x, algorithm="maxima")`

output `integrate(sqrt(x + sqrt(x + 1))/((x^2 + 1)*sqrt(x + 1)), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{x + \sqrt{1+x}}}{\sqrt{1+x}(1+x^2)} dx = \text{Exception raised: TypeError}$$

input `integrate((x+(1+x)^(1/2))^(1/2)/(x^2+1)/(1+x)^(1/2), x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:Invalid _EXT in replace_ext Error:
Bad Argument ValueInvalid _EXT in replace_ext Error: Bad Argument ValueDon
e`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{x + \sqrt{1+x}}}{\sqrt{1+x}(1+x^2)} dx = \int \frac{\sqrt{x + \sqrt{x+1}}}{(x^2 + 1)\sqrt{x+1}} dx$$

input `int((x + (x + 1)^(1/2))^(1/2)/((x^2 + 1)*(x + 1)^(1/2)),x)`

output `int((x + (x + 1)^(1/2))^(1/2)/((x^2 + 1)*(x + 1)^(1/2)), x)`

Reduce [F]

$$\int \frac{\sqrt{x + \sqrt{1+x}}}{\sqrt{1+x}(1+x^2)} dx = \int \frac{\sqrt{\sqrt{x+1} + x}}{\sqrt{x+1}x^2 + \sqrt{x+1}} dx$$

input `int((x+(1+x)^(1/2))^(1/2)/(x^2+1)/(1+x)^(1/2),x)`

output `int(sqrt(sqrt(x + 1) + x)/(sqrt(x + 1)*x**2 + sqrt(x + 1)),x)`

3.14 $\int \frac{\sqrt{x+\sqrt{1+x}}}{1+x^2} dx$

Optimal result	127
Mathematica [C] (verified)	128
Rubi [A] (verified)	128
Maple [C] (verified)	130
Fricas [B] (verification not implemented)	130
Sympy [F]	131
Maxima [F]	131
Giac [F(-2)]	131
Mupad [F(-1)]	132
Reduce [F]	132

Optimal result

Integrand size = 21, antiderivative size = 337

$$\int \frac{\sqrt{x+\sqrt{1+x}}}{1+x^2} dx$$

$$= \frac{1}{2}i\sqrt{i+\sqrt{1-i}} \arctan\left(\frac{2+\sqrt{1-i}-(1-2\sqrt{1-i})\sqrt{1+x}}{2\sqrt{i+\sqrt{1-i}}\sqrt{x+\sqrt{1+x}}}\right)$$

$$- \frac{1}{2}i\sqrt{-i+\sqrt{1+i}} \arctan\left(\frac{2+\sqrt{1+i}-(1-2\sqrt{1+i})\sqrt{1+x}}{2\sqrt{-i+\sqrt{1+i}}\sqrt{x+\sqrt{1+x}}}\right)$$

$$+ \frac{1}{2}i\sqrt{-i+\sqrt{1-i}} \operatorname{arctanh}\left(\frac{2-\sqrt{1-i}-(1+2\sqrt{1-i})\sqrt{1+x}}{2\sqrt{-i+\sqrt{1-i}}\sqrt{x+\sqrt{1+x}}}\right)$$

$$- \frac{1}{2}i\sqrt{i+\sqrt{1+i}} \operatorname{arctanh}\left(\frac{2-\sqrt{1+i}-(1+2\sqrt{1+i})\sqrt{1+x}}{2\sqrt{i+\sqrt{1+i}}\sqrt{x+\sqrt{1+x}}}\right)$$

output

```
1/2*I*arctanh(1/2*(2-(1-I)^(1/2)-(1+2*(1-I)^(1/2))*(1+x)^(1/2))/(-I+(1-I)^(1/2))^(1/2)/(x+(1+x)^(1/2))^(1/2))*(-I+(1-I)^(1/2))^(1/2)+1/2*I*arctan(1/2*(2+(1-I)^(1/2)-(1-2*(1-I)^(1/2))*(1+x)^(1/2))/(I+(1-I)^(1/2))^(1/2)/(x+(1+x)^(1/2))^(1/2))*(I+(1-I)^(1/2))^(1/2)-1/2*I*arctan(1/2*(2+(1+I)^(1/2)-(1-2*(1+I)^(1/2))*(1+x)^(1/2))/(-I+(1+I)^(1/2))^(1/2)/(x+(1+x)^(1/2))^(1/2))*(-I+(1+I)^(1/2))^(1/2)-1/2*I*arctanh(1/2*(2-(1+I)^(1/2)-(1+2*(1+I)^(1/2))*(1+x)^(1/2))/(I+(1+I)^(1/2))^(1/2)/(x+(1+x)^(1/2))^(1/2))*(I+(1+I)^(1/2))^(1/2)
```


Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.14 (sec) , antiderivative size = 212, normalized size of antiderivative = 0.63

$$\int \frac{\sqrt{x + \sqrt{1+x}}}{1+x^2} dx = \frac{1}{2} \text{RootSum} \left[1 - 8\#1 + 40\#1^2 - 48\#1^3 + 20\#1^4 + 8\#1^5 - 4\#1^6 \right. \\ \left. + \#1^8 \&, \frac{\log \left(-\sqrt{1+x} + \sqrt{x + \sqrt{1+x}} - \#1 \right) + 2 \log \left(-\sqrt{1+x} + \sqrt{x + \sqrt{1+x}} - \#1 \right) \#1 - 2 \log \left(-\sqrt{1+x} + \sqrt{x + \sqrt{1+x}} - \#1 \right) \#1^2 - 2 \log \left(-\sqrt{1+x} + \sqrt{x + \sqrt{1+x}} - \#1 \right) \#1^3}{-1 + 10\#1 - 18\#1^2 + 10\#1^3} \right]$$

input

```
Integrate[Sqrt[x + Sqrt[1 + x]]/(1 + x^2), x]
```

output

```
RootSum[1 - 8*#1 + 40*#1^2 - 48*#1^3 + 20*#1^4 + 8*#1^5 - 4*#1^6 + #1^8 &
, (Log[-Sqrt[1 + x] + Sqrt[x + Sqrt[1 + x]] - #1] + 2*Log[-Sqrt[1 + x] + S
qrt[x + Sqrt[1 + x]] - #1]*#1 - 2*Log[-Sqrt[1 + x] + Sqrt[x + Sqrt[1 + x]]
- #1]*#1^5 + Log[-Sqrt[1 + x] + Sqrt[x + Sqrt[1 + x]] - #1]*#1^6)/(-1 + 1
0*#1 - 18*#1^2 + 10*#1^3 + 5*#1^4 - 3*#1^5 + #1^7) & ]/2
```

Rubi [A] (verified)

Time = 0.84 (sec) , antiderivative size = 339, normalized size of antiderivative = 1.01, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {7267, 7292, 7279, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{x + \sqrt{x+1}}}{x^2 + 1} dx \\ \downarrow 7267 \\ 2 \int \frac{\sqrt{x+1} \sqrt{x + \sqrt{x+1}}}{x^2 + 1} d\sqrt{x+1} \\ \downarrow 7292$$

$$\begin{aligned}
& 2 \int \frac{\sqrt{x+1}\sqrt{x+\sqrt{x+1}}}{(x+1)^2 - 2(x+1) + 2} d\sqrt{x+1} \\
& \quad \downarrow \text{7279} \\
& 2 \int \left(\frac{i\sqrt{x+1}\sqrt{x+\sqrt{x+1}}}{(2+2i) - 2(x+1)} + \frac{i\sqrt{x+1}\sqrt{x+\sqrt{x+1}}}{2(x+1) - (2-2i)} \right) d\sqrt{x+1} \\
& \quad \downarrow \text{2009} \\
& 2 \left(\frac{1}{4} i \sqrt{i + \sqrt{1-i}} \arctan \left(\frac{-((1-2\sqrt{1-i})\sqrt{x+1}) + \sqrt{1-i} + 2}{2\sqrt{i + \sqrt{1-i}}\sqrt{x+\sqrt{x+1}}} \right) - \frac{1}{4} i \sqrt{\sqrt{1+i} - i} \arctan \left(\frac{-((1-2\sqrt{1+i})\sqrt{x+1}) + \sqrt{1+i} + 2}{2\sqrt{\sqrt{1+i} - i}\sqrt{x+\sqrt{x+1}}} \right) \right)
\end{aligned}$$

input `Int[Sqrt[x + Sqrt[1 + x]]/(1 + x^2), x]`

output

```

2*((I/4)*Sqrt[I + Sqrt[1 - I]]*ArcTan[(2 + Sqrt[1 - I] - (1 - 2*Sqrt[1 - I])
]*Sqrt[1 + x])/(2*Sqrt[I + Sqrt[1 - I]]*Sqrt[x + Sqrt[1 + x]])] - (I/4)*S
qrt[-I + Sqrt[1 + I]]*ArcTan[(2 + Sqrt[1 + I] - (1 - 2*Sqrt[1 + I])*Sqrt[1
+ x])/(2*Sqrt[-I + Sqrt[1 + I]]*Sqrt[x + Sqrt[1 + x]])] + (I/4)*Sqrt[-I +
Sqrt[1 - I]]*ArcTanh[(2 - Sqrt[1 - I] - (1 + 2*Sqrt[1 - I])*Sqrt[1 + x])/
(2*Sqrt[-I + Sqrt[1 - I]]*Sqrt[x + Sqrt[1 + x]])] - (I/4)*Sqrt[I + Sqrt[1
+ I]]*ArcTanh[(2 - Sqrt[1 + I] - (1 + 2*Sqrt[1 + I])*Sqrt[1 + x])/(2*Sqrt[
I + Sqrt[1 + I]]*Sqrt[x + Sqrt[1 + x]])]

```

Defintions of rubi rules used

rule 2009

```

Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]

```

rule 7267

```

Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Si
mp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2]])], x
] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]

```

rule 7279

```

Int[(u_)/((a_.) + (b_.)*(x_)^(n_.) + (c_.)*(x_)^(n2_.)), x_Symbol] := With[
{v = RationalFunctionExpand[u/(a + b*x^n + c*x^(2*n)), x]}, Int[v, x] /; Su
mQ[v]] /; FreeQ[{a, b, c}, x] && EqQ[n2, 2*n] && IGtQ[n, 0]

```

rule 7292

```
Int[u_, x_Symbol] :=> With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v !=
= u]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.12 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.31

method	result
derivativedivides	$\frac{\left(\sum_{R=\text{RootOf}(-Z^8-4Z^6+8Z^5+20Z^4-48Z^3+40Z^2-8Z+1)} \frac{(-R^6-2R^5+2R+1) \ln(\sqrt{x+\sqrt{1+x}}-\sqrt{1+x})}{-R^7-3R^5+5R^4+10R^3-18R^2+10R-1} \right)}{2}$
default	$\frac{\left(\sum_{R=\text{RootOf}(-Z^8-4Z^6+8Z^5+20Z^4-48Z^3+40Z^2-8Z+1)} \frac{(-R^6-2R^5+2R+1) \ln(\sqrt{x+\sqrt{1+x}}-\sqrt{1+x})}{-R^7-3R^5+5R^4+10R^3-18R^2+10R-1} \right)}{2}$

input

```
int((x+(1+x)^(1/2))^(1/2)/(x^2+1),x,method=_RETURNVERBOSE)
```

output

```
1/2*sum((-R^6-2*R^5+2*R+1)/(-R^7-3*R^5+5*R^4+10*R^3-18*R^2+10*R-1)*
ln((x+(1+x)^(1/2))^(1/2)-(1+x)^(1/2)-R),R=RootOf(-Z^8-4*_Z^6+8*_Z^5+20*_
Z^4-48*_Z^3+40*_Z^2-8*_Z+1))
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 4535 vs. $2(185) = 370$.

Time = 3.09 (sec) , antiderivative size = 4535, normalized size of antiderivative = 13.46

$$\int \frac{\sqrt{x + \sqrt{1+x}}}{1+x^2} dx = \text{Too large to display}$$

input

```
integrate((x+(1+x)^(1/2))^(1/2)/(x^2+1),x, algorithm="fricas")
```

output Too large to include

Sympy [F]

$$\int \frac{\sqrt{x + \sqrt{1+x}}}{1+x^2} dx = \int \frac{\sqrt{x + \sqrt{x+1}}}{x^2 + 1} dx$$

input `integrate((x+(1+x)**(1/2))**(1/2)/(x**2+1), x)`

output `Integral(sqrt(x + sqrt(x + 1))/(x**2 + 1), x)`

Maxima [F]

$$\int \frac{\sqrt{x + \sqrt{1+x}}}{1+x^2} dx = \int \frac{\sqrt{x + \sqrt{x+1}}}{x^2 + 1} dx$$

input `integrate((x+(1+x)^(1/2))^(1/2)/(x^2+1), x, algorithm="maxima")`

output `integrate(sqrt(x + sqrt(x + 1))/(x^2 + 1), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{x + \sqrt{1+x}}}{1+x^2} dx = \text{Exception raised: TypeError}$$

input `integrate((x+(1+x)^(1/2))^(1/2)/(x^2+1), x, algorithm="giac")`

output

```
Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Precision problem choosing root in
common_EXT, current precision 14Precision problem choosing root in common_
EXT, curr
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{x + \sqrt{1+x}}}{1+x^2} dx = \int \frac{\sqrt{x + \sqrt{x+1}}}{x^2 + 1} dx$$

input

```
int((x + (x + 1)^(1/2))^(1/2)/(x^2 + 1),x)
```

output

```
int((x + (x + 1)^(1/2))^(1/2)/(x^2 + 1), x)
```

Reduce [F]

$$\int \frac{\sqrt{x + \sqrt{1+x}}}{1+x^2} dx = \int \frac{\sqrt{\sqrt{x+1} + x}}{x^2 + 1} dx$$

input

```
int((x+(1+x)^(1/2))^(1/2)/(x^2+1),x)
```

output

```
int(sqrt(sqrt(x + 1) + x)/(x**2 + 1),x)
```

3.15 $\int \sqrt{1 + \sqrt{x} + \sqrt{1 + 2\sqrt{x} + 2x}} dx$

Optimal result	133
Mathematica [A] (verified)	133
Rubi [A] (verified)	134
Maple [F]	135
Fricas [A] (verification not implemented)	135
Sympy [F]	136
Maxima [F]	136
Giac [F]	136
Mupad [F(-1)]	137
Reduce [F]	137

Optimal result

Integrand size = 27, antiderivative size = 77

$$\int \sqrt{1 + \sqrt{x} + \sqrt{1 + 2\sqrt{x} + 2x}} dx$$

$$= \frac{2\sqrt{1 + \sqrt{x} + \sqrt{1 + 2\sqrt{x} + 2x}} \left(2 + \sqrt{x} + 6x^{3/2} - (2 - \sqrt{x}) \sqrt{1 + 2\sqrt{x} + 2x} \right)}{15\sqrt{x}}$$

output

```
2/15*(2+6*x^(3/2)+x^(1/2)-(2-x^(1/2))*(1+2*x+2*x^(1/2))^(1/2))*(1+x^(1/2)+
(1+2*x+2*x^(1/2))^(1/2))^1/2/x^(1/2)
```

Mathematica [A] (verified)

Time = 10.03 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.96

$$\int \sqrt{1 + \sqrt{x} + \sqrt{1 + 2\sqrt{x} + 2x}} dx$$

$$= \frac{2\sqrt{1 + \sqrt{x} + \sqrt{1 + 2\sqrt{x} + 2x}} \left(2 + \sqrt{x} + 6x^{3/2} + (-2 + \sqrt{x}) \sqrt{1 + 2\sqrt{x} + 2x} \right)}{15\sqrt{x}}$$

input

```
Integrate[Sqrt[1 + Sqrt[x] + Sqrt[1 + 2*Sqrt[x] + 2*x]],x]
```

output $(2\sqrt{1 + \sqrt{x}} + \sqrt{1 + 2\sqrt{x}} + 2x)(2 + \sqrt{x} + 6x^{3/2}) + (-2 + \sqrt{x})\sqrt{1 + 2\sqrt{x}} + 2x)/(15\sqrt{x})$

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.074$, Rules used = {7267, 2539}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\sqrt{x} + \sqrt{2x + 2\sqrt{x} + 1} + 1} dx$$

$$\downarrow 7267$$

$$2 \int \sqrt{x} \sqrt{\sqrt{x} + \sqrt{2x + 2\sqrt{x} + 1} + 1} d\sqrt{x}$$

$$\downarrow 2539$$

$$\frac{2\sqrt{\sqrt{x} + \sqrt{2x + 2\sqrt{x} + 1} + 1} (6x^{3/2} + \sqrt{x} - (2 - \sqrt{x}) \sqrt{2x + 2\sqrt{x} + 1} + 2)}{15\sqrt{x}}$$

input $\text{Int}[\sqrt{1 + \sqrt{x}} + \sqrt{1 + 2\sqrt{x}} + 2x], x]$

output $(2\sqrt{1 + \sqrt{x}} + \sqrt{1 + 2\sqrt{x}} + 2x)(2 + \sqrt{x} + 6x^{3/2}) - (2 - \sqrt{x})\sqrt{1 + 2\sqrt{x}} + 2x)/(15\sqrt{x})$

Definitions of rubi rules used

rule 2539

```
Int[((g_.) + (h_.)*(x_))*Sqrt[(d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_.) + (b_.)
*(x_) + (c_.)*(x_)^2]], x_Symbol] := Simp[2*((f*(5*b*c*g^2 - 2*b^2*g*h - 3*
a*c*g*h + 2*a*b*h^2) + c*f*(10*c*g^2 - b*g*h + a*h^2)*x + 9*c^2*f*g*h*x^2 +
3*c^2*f*h^2*x^3 - (e*g - d*h)*(5*c*g - 2*b*h + c*h*x)*Sqrt[a + b*x + c*x^2
])/((15*c^2*f*(g + h*x))*Sqrt[d + e*x + f*Sqrt[a + b*x + c*x^2]]), x] /; Fre
eQ[{a, b, c, d, e, f, g, h}, x] && EqQ[(e*g - d*h)^2 - f^2*(c*g^2 - b*g*h +
a*h^2), 0] && EqQ[2*e^2*g - 2*d*e*h - f^2*(2*c*g - b*h), 0]
```

rule 7267

```
Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Si
mp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])]], x
] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]
```

Maple [F]

$$\int \sqrt{1 + \sqrt{x} + \sqrt{1 + 2x + 2\sqrt{x}}} dx$$

input

```
int((1+x^(1/2)+(1+2*x+2*x^(1/2))^(1/2))^(1/2),x)
```

output

```
int((1+x^(1/2)+(1+2*x+2*x^(1/2))^(1/2))^(1/2),x)
```

Fricas [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.73

$$\int \sqrt{1 + \sqrt{x} + \sqrt{1 + 2\sqrt{x} + 2x}} dx$$

$$= \frac{2 \left(6x^2 + \sqrt{2x + 2\sqrt{x} + 1}(x - 2\sqrt{x}) + x + 2\sqrt{x} \right) \sqrt{\sqrt{2x + 2\sqrt{x} + 1} + \sqrt{x} + 1}}{15x}$$

input

```
integrate((1+x^(1/2)+(1+2*x+2*x^(1/2))^(1/2))^(1/2),x, algorithm="fricas")
```


output $2/15*(6*x^2 + \sqrt{2*x + 2*\sqrt{x} + 1}*(x - 2*\sqrt{x}) + x + 2*\sqrt{x})*\sqrt{\sqrt{2*x + 2*\sqrt{x} + 1} + \sqrt{x} + 1}/x$

Sympy [F]

$$\int \sqrt{1 + \sqrt{x} + \sqrt{1 + 2\sqrt{x} + 2x}} dx = \int \sqrt{\sqrt{x} + \sqrt{2\sqrt{x} + 2x + 1} + 1} dx$$

input `integrate((1+x**(1/2)+(1+2*x+2*x**(1/2))**(1/2))**(1/2)),x)`

output `Integral(sqrt(sqrt(x) + sqrt(2*sqrt(x) + 2*x + 1) + 1), x)`

Maxima [F]

$$\int \sqrt{1 + \sqrt{x} + \sqrt{1 + 2\sqrt{x} + 2x}} dx = \int \sqrt{\sqrt{2x + 2\sqrt{x} + 1} + \sqrt{x} + 1} dx$$

input `integrate((1+x^(1/2)+(1+2*x+2*x^(1/2))^(1/2))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(sqrt(2*x + 2*sqrt(x) + 1) + sqrt(x) + 1), x)`

Giac [F]

$$\int \sqrt{1 + \sqrt{x} + \sqrt{1 + 2\sqrt{x} + 2x}} dx = \int \sqrt{\sqrt{2x + 2\sqrt{x} + 1} + \sqrt{x} + 1} dx$$

input `integrate((1+x^(1/2)+(1+2*x+2*x^(1/2))^(1/2))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(sqrt(2*x + 2*sqrt(x) + 1) + sqrt(x) + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{1 + \sqrt{x} + \sqrt{1 + 2\sqrt{x} + 2x}} dx = \int \sqrt{\sqrt{2x + 2\sqrt{x} + 1} + \sqrt{x} + 1} dx$$

input `int(((2*x + 2*x^(1/2) + 1)^(1/2) + x^(1/2) + 1)^(1/2),x)`

output `int(((2*x + 2*x^(1/2) + 1)^(1/2) + x^(1/2) + 1)^(1/2), x)`

Reduce [F]

$$\int \sqrt{1 + \sqrt{x} + \sqrt{1 + 2\sqrt{x} + 2x}} dx = \int \sqrt{\sqrt{2\sqrt{x} + 2x + 1} + \sqrt{x} + 1} dx$$

input `int(((1+x^(1/2)+(1+2*x+2*x^(1/2))^(1/2))^(1/2))^(1/2),x)`

output `int(sqrt(sqrt(2*sqrt(x) + 2*x + 1) + sqrt(x) + 1),x)`

3.16 $\int \sqrt{\sqrt{2} + \sqrt{x} + \sqrt{2 + 2\sqrt{2}\sqrt{x} + 2x}} dx$

Optimal result	138
Mathematica [A] (verified)	138
Rubi [A] (verified)	139
Maple [F]	140
Fricas [A] (verification not implemented)	141
Sympy [F]	141
Maxima [F]	142
Giac [F(-2)]	142
Mupad [F(-1)]	142
Reduce [F]	143

Optimal result

Integrand size = 36, antiderivative size = 118

$$\int \sqrt{\sqrt{2} + \sqrt{x} + \sqrt{2 + 2\sqrt{2}\sqrt{x} + 2x}} dx$$

$$= \frac{2\sqrt{2}\sqrt{\sqrt{2} + \sqrt{x} + \sqrt{2}\sqrt{1 + \sqrt{2}\sqrt{x} + x}} \left(4 + \sqrt{2}\sqrt{x} + 3\sqrt{2}x^{3/2} - \sqrt{2}(2\sqrt{2} - \sqrt{x}) \sqrt{1 + \sqrt{2}\sqrt{x} + x} \right)}{15\sqrt{x}}$$

output

```
2/15*2^(1/2)*(4+3*x^(3/2)*2^(1/2)+2^(1/2)*x^(1/2)-2^(1/2)*(2*2^(1/2)-x^(1/2)))*(1+x+2^(1/2)*x^(1/2))^(1/2)*(2^(1/2)+x^(1/2)+2^(1/2)*(1+x+2^(1/2)*x^(1/2))^(1/2))/x^(1/2)
```

Mathematica [A] (verified)

Time = 10.06 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.95

$$\int \sqrt{\sqrt{2} + \sqrt{x} + \sqrt{2 + 2\sqrt{2}\sqrt{x} + 2x}} dx$$

$$= \frac{2\sqrt{2} \left(4 + \sqrt{2}\sqrt{x} + 3\sqrt{2}x^{3/2} + \sqrt{2}(-2\sqrt{2} + \sqrt{x}) \sqrt{1 + \sqrt{2}\sqrt{x} + x} \right) \sqrt{\sqrt{x} + \sqrt{2}} \left(1 + \sqrt{1 + \sqrt{2}\sqrt{x} + x} \right)}{15\sqrt{x}}$$

input `Integrate[Sqrt[Sqrt[2] + Sqrt[x] + Sqrt[2 + 2*Sqrt[2]*Sqrt[x] + 2*x]],x]`

output `(2*Sqrt[2]*(4 + Sqrt[2]*Sqrt[x] + 3*Sqrt[2]*x^(3/2) + Sqrt[2]*(-2*Sqrt[2] + Sqrt[x])*Sqrt[1 + Sqrt[2]*Sqrt[x] + x])*Sqrt[Sqrt[x] + Sqrt[2]*(1 + Sqrt[1 + Sqrt[2]*Sqrt[x] + x])]/(15*Sqrt[x])`

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {7267, 2540, 2539}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\sqrt{x} + \sqrt{2x + 2\sqrt{2}\sqrt{x} + 2} + \sqrt{2}} dx$$

$$\downarrow 7267$$

$$2 \int \sqrt{x} \sqrt{\sqrt{2} \left(\sqrt{x + \sqrt{2}\sqrt{x} + 1} + 1 \right) + \sqrt{x}} d\sqrt{x}$$

$$\downarrow 2540$$

$$2 \int \sqrt{x} \sqrt{\sqrt{x} + \sqrt{2}\sqrt{x + \sqrt{2}\sqrt{x} + 1} + \sqrt{2}} d\sqrt{x}$$

$$\downarrow 2539$$

$$\frac{2\sqrt{2}\sqrt{\sqrt{x} + \sqrt{2}\sqrt{x + \sqrt{2}\sqrt{x} + 1} + \sqrt{2}} \left(3\sqrt{2}x^{3/2} + \sqrt{2}\sqrt{x} - \sqrt{2}(2\sqrt{2} - \sqrt{x}) \sqrt{x + \sqrt{2}\sqrt{x} + 1} + 4 \right)}{15\sqrt{x}}$$

input `Int[Sqrt[Sqrt[2] + Sqrt[x] + Sqrt[2 + 2*Sqrt[2]*Sqrt[x] + 2*x]],x]`

output

```
(2*Sqrt[2]*Sqrt[Sqrt[2] + Sqrt[x] + Sqrt[2]*Sqrt[1 + Sqrt[2]*Sqrt[x] + x]]
*(4 + Sqrt[2]*Sqrt[x] + 3*Sqrt[2]*x^(3/2) - Sqrt[2]*(2*Sqrt[2] - Sqrt[x])*
Sqrt[1 + Sqrt[2]*Sqrt[x] + x]))/(15*Sqrt[x])
```

Defintions of rubi rules used

rule 2539

```
Int[((g_.) + (h_.)*(x_.))*Sqrt[(d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_.) + (b_.)
*(x_) + (c_.)*(x_)^2]], x_Symbol] := Simp[2*((f*(5*b*c*g^2 - 2*b^2*g*h - 3*
a*c*g*h + 2*a*b*h^2) + c*f*(10*c*g^2 - b*g*h + a*h^2)*x + 9*c^2*f*g*h*x^2 +
3*c^2*f*h^2*x^3 - (e*g - d*h)*(5*c*g - 2*b*h + c*h*x)*Sqrt[a + b*x + c*x^2
])/(15*c^2*f*(g + h*x))*Sqrt[d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; Fre
eQ[{a, b, c, d, e, f, g, h}, x] && EqQ[(e*g - d*h)^2 - f^2*(c*g^2 - b*g*h +
a*h^2), 0] && EqQ[2*e^2*g - 2*d*e*h - f^2*(2*c*g - b*h), 0]
```

rule 2540

```
Int[((u_) + (f_.)*((j_.) + (k_.)*Sqrt[v_]))^(n_.)*((g_.) + (h_.)*(x_.))^(m_.
), x_Symbol] := Int[(g + h*x)^m*(ExpandToSum[u + f*j, x] + f*k*Sqrt[ExpandT
oSum[v, x]])^n, x] /; FreeQ[{f, g, h, j, k, m, n}, x] && LinearQ[u, x] && Q
uadraticQ[v, x] && !(LinearMatchQ[u, x] && QuadraticMatchQ[v, x] && (EqQ[j
, 0] || EqQ[f, 1])) && EqQ[(Coefficient[u, x, 1]*g - h*(Coefficient[u, x, 0
] + f*j))^2 - f^2*k^2*(Coefficient[v, x, 2]*g^2 - Coefficient[v, x, 1]*g*h
+ Coefficient[v, x, 0]*h^2), 0]
```

rule 7267

```
Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Si
mp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])]], x
] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]
```

Maple [F]

$$\int \sqrt{\sqrt{2} + \sqrt{x} + \sqrt{2 + 2x + 2\sqrt{2}\sqrt{x}}} dx$$

input

```
int((2^(1/2)+x^(1/2)+(2+2*x+2*2^(1/2)*x^(1/2))^(1/2))^(1/2),x)
```

output

```
int((2^(1/2)+x^(1/2)+(2+2*x+2*2^(1/2)*x^(1/2))^(1/2))^(1/2),x)
```

Fricas [A] (verification not implemented)

Time = 0.72 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.62

$$\int \sqrt{\sqrt{2} + \sqrt{x} + \sqrt{2 + 2\sqrt{2}\sqrt{x} + 2x}} dx$$

$$= \frac{2 \left(6x^2 + (\sqrt{2}x - 4\sqrt{x})\sqrt{2\sqrt{2}\sqrt{x} + 2x + 2} + 4\sqrt{2}\sqrt{x} + 2x \right) \sqrt{\sqrt{2} + \sqrt{2\sqrt{2}\sqrt{x} + 2x + 2} + \sqrt{x}}}{15x}$$

input `integrate((2^(1/2)+x^(1/2)+(2+2*x+2*2^(1/2)*x^(1/2))^(1/2))^(1/2),x, algo
ithm="fricas")`

output `2/15*(6*x^2 + (sqrt(2)*x - 4*sqrt(x))*sqrt(2*sqrt(2)*sqrt(x) + 2*x + 2) +
4*sqrt(2)*sqrt(x) + 2*x)*sqrt(sqrt(2) + sqrt(2*sqrt(2)*sqrt(x) + 2*x + 2)
+ sqrt(x))/x`

Sympy [F]

$$\int \sqrt{\sqrt{2} + \sqrt{x} + \sqrt{2 + 2\sqrt{2}\sqrt{x} + 2x}} dx = \int \sqrt{\sqrt{x} + \sqrt{2\sqrt{2}\sqrt{x} + 2x + 2} + \sqrt{2}} dx$$

input `integrate((2**(1/2)+x**(1/2)+(2+2*x+2*2**(1/2)*x**(1/2))**(1/2))**(1/2),x)`

output `Integral(sqrt(sqrt(x) + sqrt(2*sqrt(2)*sqrt(x) + 2*x + 2) + sqrt(2)), x)`

Maxima [F]

$$\int \sqrt{\sqrt{2} + \sqrt{x} + \sqrt{2 + 2\sqrt{2}\sqrt{x} + 2x}} dx = \int \sqrt{\sqrt{2} + \sqrt{2\sqrt{2}\sqrt{x} + 2x} + 2 + \sqrt{x}} dx$$

input `integrate((2^(1/2)+x^(1/2)+(2+2*x+2*2^(1/2)*x^(1/2))^(1/2))^(1/2),x, algorith="maxima")`

output `integrate(sqrt(sqrt(2) + sqrt(2*sqrt(2)*sqrt(x) + 2*x + 2) + sqrt(x)), x)`

Giac [F(-2)]

Exception generated.

$$\int \sqrt{\sqrt{2} + \sqrt{x} + \sqrt{2 + 2\sqrt{2}\sqrt{x} + 2x}} dx = \text{Exception raised: TypeError}$$

input `integrate((2^(1/2)+x^(1/2)+(2+2*x+2*2^(1/2)*x^(1/2))^(1/2))^(1/2),x, algorith="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for the root of a polynomial with parameters. This might be wrong.The choice was done`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{\sqrt{2} + \sqrt{x} + \sqrt{2 + 2\sqrt{2}\sqrt{x} + 2x}} dx = \int \sqrt{\sqrt{2x + 2\sqrt{2}\sqrt{x} + 2} + \sqrt{2} + \sqrt{x}} dx$$

input `int(((2*x + 2*2^(1/2)*x^(1/2) + 2)^(1/2) + 2^(1/2) + x^(1/2))^(1/2),x)`

output `int(((2*x + 2*2^(1/2)*x^(1/2) + 2)^(1/2) + 2^(1/2) + x^(1/2))^(1/2), x)`

Reduce [F]

$$\int \sqrt{\sqrt{2} + \sqrt{x} + \sqrt{2 + 2\sqrt{2}\sqrt{x} + 2x}} dx = \int \sqrt{\sqrt{\sqrt{x}\sqrt{2} + x + 1}\sqrt{2} + \sqrt{x} + \sqrt{2}} dx$$

input `int((2^(1/2)+x^(1/2)+(2+2*x+2*2^(1/2)*x^(1/2))^(1/2))^(1/2),x)`

output `int(sqrt(sqrt(sqrt(x)*sqrt(2) + x + 1)*sqrt(2) + sqrt(x) + sqrt(2)),x)`

3.17 $\int \frac{\sqrt{x+\sqrt{1+x}}}{x^2} dx$

Optimal result	144
Mathematica [A] (verified)	144
Rubi [A] (verified)	145
Maple [B] (verified)	147
Fricas [A] (verification not implemented)	148
Sympy [F]	149
Maxima [F]	149
Giac [B] (verification not implemented)	149
Mupad [F(-1)]	150
Reduce [B] (verification not implemented)	150

Optimal result

Integrand size = 17, antiderivative size = 83

$$\int \frac{\sqrt{x+\sqrt{1+x}}}{x^2} dx = -\frac{\sqrt{x+\sqrt{1+x}}}{x} - \frac{1}{4} \arctan\left(\frac{3+\sqrt{1+x}}{2\sqrt{x+\sqrt{1+x}}}\right) + \frac{3}{4} \operatorname{arctanh}\left(\frac{1-3\sqrt{1+x}}{2\sqrt{x+\sqrt{1+x}}}\right)$$

output

```
-1/4*arctan(1/2*(3+(1+x)^(1/2))/(x+(1+x)^(1/2))^(1/2))+3/4*arctanh(1/2*(1-3*(1+x)^(1/2))/(x+(1+x)^(1/2))^(1/2))-(x+(1+x)^(1/2))^(1/2)/x
```

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.93

$$\int \frac{\sqrt{x+\sqrt{1+x}}}{x^2} dx = -\frac{\sqrt{x+\sqrt{1+x}}}{x} - \frac{1}{2} \arctan\left(1+\sqrt{1+x}-\sqrt{x+\sqrt{1+x}}\right) - \frac{3}{2} \operatorname{arctanh}\left(1-\sqrt{1+x}+\sqrt{x+\sqrt{1+x}}\right)$$

input

```
Integrate[Sqrt[x + Sqrt[1 + x]]/x^2,x]
```

output

$$-(\text{Sqrt}[x + \text{Sqrt}[1 + x]]/x) - \text{ArcTan}[1 + \text{Sqrt}[1 + x] - \text{Sqrt}[x + \text{Sqrt}[1 + x]]]/2 - (3*\text{ArcTanh}[1 - \text{Sqrt}[1 + x] + \text{Sqrt}[x + \text{Sqrt}[1 + x]])/2$$
Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.11, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.471$, Rules used = {7267, 1347, 27, 1366, 25, 1154, 217, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{x + \sqrt{x+1}}}{x^2} dx$$

$$\downarrow 7267$$

$$2 \int \frac{\sqrt{x+1} \sqrt{x + \sqrt{x+1}}}{x^2} d\sqrt{x+1}$$

$$\downarrow 1347$$

$$2 \left(\frac{1}{2} \int \frac{2\sqrt{x+1} + 1}{2x\sqrt{x + \sqrt{x+1}}} d\sqrt{x+1} - \frac{\sqrt{x + \sqrt{x+1}}}{2x} \right)$$

$$\downarrow 27$$

$$2 \left(-\frac{1}{4} \int -\frac{2\sqrt{x+1} + 1}{x\sqrt{x + \sqrt{x+1}}} d\sqrt{x+1} - \frac{\sqrt{x + \sqrt{x+1}}}{2x} \right)$$

$$\downarrow 1366$$

$$2 \left(\frac{1}{4} \left(-\frac{3}{2} \int \frac{1}{(1 - \sqrt{x+1}) \sqrt{x + \sqrt{x+1}}} d\sqrt{x+1} - \frac{1}{2} \int -\frac{1}{(\sqrt{x+1} + 1) \sqrt{x + \sqrt{x+1}}} d\sqrt{x+1} \right) - \frac{\sqrt{x + \sqrt{x+1}}}{2x} \right)$$

$$\downarrow 25$$

$$2 \left(\frac{1}{4} \left(\frac{1}{2} \int \frac{1}{(\sqrt{x+1} + 1) \sqrt{x + \sqrt{x+1}}} d\sqrt{x+1} - \frac{3}{2} \int \frac{1}{(1 - \sqrt{x+1}) \sqrt{x + \sqrt{x+1}}} d\sqrt{x+1} \right) - \frac{\sqrt{x + \sqrt{x+1}}}{2x} \right)$$

$$\downarrow 1154$$

$$\begin{aligned}
& 2\left(\frac{1}{4}\left(3\int\frac{1}{3-x}d\frac{1-3\sqrt{x+1}}{\sqrt{x+\sqrt{x+1}}}-\int\frac{1}{-x-5}d\left(-\frac{\sqrt{x+1}+3}{\sqrt{x+\sqrt{x+1}}}\right)\right)-\frac{\sqrt{x+\sqrt{x+1}}}{2x}\right) \\
& \quad \downarrow \text{217} \\
& 2\left(\frac{1}{4}\left(3\int\frac{1}{3-x}d\frac{1-3\sqrt{x+1}}{\sqrt{x+\sqrt{x+1}}}-\frac{1}{2}\arctan\left(\frac{\sqrt{x+1}+3}{2\sqrt{x+\sqrt{x+1}}}\right)\right)-\frac{\sqrt{x+\sqrt{x+1}}}{2x}\right) \\
& \quad \downarrow \text{219} \\
& 2\left(\frac{1}{4}\left(\frac{3}{2}\operatorname{arctanh}\left(\frac{1-3\sqrt{x+1}}{2\sqrt{x+\sqrt{x+1}}}\right)-\frac{1}{2}\arctan\left(\frac{\sqrt{x+1}+3}{2\sqrt{x+\sqrt{x+1}}}\right)\right)-\frac{\sqrt{x+\sqrt{x+1}}}{2x}\right)
\end{aligned}$$

input `Int[Sqrt[x + Sqrt[1 + x]]/x^2,x]`

output `2*(-1/2*Sqrt[x + Sqrt[1 + x]]/x + (-1/2*ArcTan[(3 + Sqrt[1 + x])/(2*Sqrt[x + Sqrt[1 + x]])] + (3*ArcTanh[(1 - 3*Sqrt[1 + x])/(2*Sqrt[x + Sqrt[1 + x]])])]/2)/4)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 1154 `Int[1/(((d_.) + (e_.)*(x_))*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]), x_Symbol] := Simp[-2 Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 1347 `Int[((g_.) + (h_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_)*((d_) + (e_.)*(x_) + (f_.)*(x_)^2)^(q_), x_Symbol] := Simp[(a*h - g*c*x)*(a + c*x^2)^(p + 1)*((d + e*x + f*x^2)^q/(2*a*c*(p + 1))), x] + Simp[2/(4*a*c*(p + 1)) Int[(a + c*x^2)^(p + 1)*(d + e*x + f*x^2)^(q - 1)*Simp[g*c*d*(2*p + 3) - a*(h*e*q) + (g*c*e*(2*p + q + 3) - a*(2*h*f*q))*x + g*c*f*(2*p + 2*q + 3)*x^2, x], x] /; FreeQ[{a, c, d, e, f, g, h}, x] && NeQ[e^2 - 4*d*f, 0] && LtQ[p, -1] && GtQ[q, 0]`

rule 1366 `Int[((g_.) + (h_.)*(x_))/(((a_) + (c_.)*(x_)^2)*Sqrt[(d_.) + (e_.)*(x_) + (f_.)*(x_)^2]), x_Symbol] := With[{q = Rt[(-a)*c, 2]}, Simp[(h/2 + c*(g/(2*q))) Int[1/((-q + c*x)*Sqrt[d + e*x + f*x^2]), x], x] + Simp[(h/2 - c*(g/(2*q))) Int[1/(q + c*x)*Sqrt[d + e*x + f*x^2]), x], x] /; FreeQ[{a, c, d, e, f, g, h}, x] && NeQ[e^2 - 4*d*f, 0] && PosQ[(-a)*c]`

rule 7267 `Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Simp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])]], x] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 297 vs. 2(59) = 118.

Time = 0.04 (sec) , antiderivative size = 298, normalized size of antiderivative = 3.59

method	result
derivativedivides	$-\frac{\left((-1+\sqrt{1+x})^2+3\sqrt{1+x}-2\right)^{\frac{3}{2}}}{2(-1+\sqrt{1+x})} + \frac{3\sqrt{(-1+\sqrt{1+x})^2+3\sqrt{1+x}-2}}{4} + \frac{\ln\left(\frac{1}{2}+\sqrt{1+x}+\sqrt{(-1+\sqrt{1+x})^2+3\sqrt{1+x}-2}\right)}{2}$
default	$-\frac{\left((-1+\sqrt{1+x})^2+3\sqrt{1+x}-2\right)^{\frac{3}{2}}}{2(-1+\sqrt{1+x})} + \frac{3\sqrt{(-1+\sqrt{1+x})^2+3\sqrt{1+x}-2}}{4} + \frac{\ln\left(\frac{1}{2}+\sqrt{1+x}+\sqrt{(-1+\sqrt{1+x})^2+3\sqrt{1+x}-2}\right)}{2}$

input `int((x+(1+x)^(1/2))^(1/2)/x^2,x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -1/2/(-1+(1+x)^{(1/2)})*((-1+(1+x)^{(1/2)})^2+3*(1+x)^{(1/2)-2})^{(3/2)}+3/4*((-1+ \\ & (1+x)^{(1/2)})^2+3*(1+x)^{(1/2)-2})^{(1/2)}+1/2*\ln(1/2+(1+x)^{(1/2)}+((-1+(1+x)^{(1/2)}) \\ & ^2+3*(1+x)^{(1/2)-2})^{(1/2)})-3/4*\operatorname{arctanh}(1/2*(-1+3*(1+x)^{(1/2)}))/((-1+(1+x) \\ & ^{(1/2)})^2+3*(1+x)^{(1/2)-2})^{(1/2)}+1/4*(1+2*(1+x)^{(1/2)})*((-1+(1+x)^{(1/2)}) \\ & ^2+3*(1+x)^{(1/2)-2})^{(1/2)}-1/2/(1+(1+x)^{(1/2)})*((1+(1+x)^{(1/2)})^2-(1+x)^{(1/2)-2})^{(3/2)} \\ & -1/4*((1+(1+x)^{(1/2)})^2-(1+x)^{(1/2)-2})^{(1/2)}-1/2*\ln(1/2+(1+x)^{(1/2)}+((1+(1+x)^{(1/2)})^2-(1+x)^{(1/2)-2})^{(1/2)}) \\ & +((1+(1+x)^{(1/2)})^2-(1+x)^{(1/2)-2})^{(1/2)})+1/4*\operatorname{arctan}(1/2*(-3-(1+x)^{(1/2)}))/((1+(1+x)^{(1/2)})^2-(1+x)^{(1/2)-2})^{(1/2)} \\ & +1/4*(1+2*(1+x)^{(1/2)})*((1+(1+x)^{(1/2)})^2-(1+x)^{(1/2)-2})^{(1/2)} \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 1.65 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.98

$$\int \frac{\sqrt{x + \sqrt{1 + x}}}{x^2} dx = \frac{x \arctan\left(\frac{2\sqrt{x+\sqrt{x+1}}(\sqrt{x+1}-3)}{x-8}\right) + 3x \log\left(\frac{2\sqrt{x+\sqrt{x+1}}(\sqrt{x+1}+1)-3x-2\sqrt{x+1}-2}{x}\right) - 4\sqrt{x + \sqrt{x + 1}}}{4x}$$

input `integrate((x+(1+x)^(1/2))^(1/2)/x^2,x, algorithm="fricas")`

output
$$\begin{aligned} & 1/4*(x*\arctan(2*\sqrt{x + \sqrt{x + 1}}*(\sqrt{x + 1} - 3)/(x - 8)) + 3*x*\log \\ & ((2*\sqrt{x + \sqrt{x + 1}}*(\sqrt{x + 1} + 1) - 3*x - 2*\sqrt{x + 1} - 2)/x) \\ & - 4*\sqrt{x + \sqrt{x + 1}})/x \end{aligned}$$

Sympy [F]

$$\int \frac{\sqrt{x + \sqrt{1+x}}}{x^2} dx = \int \frac{\sqrt{x + \sqrt{x+1}}}{x^2} dx$$

input `integrate((x+(1+x)**(1/2))**(1/2)/x**2,x)`

output `Integral(sqrt(x + sqrt(x + 1))/x**2, x)`

Maxima [F]

$$\int \frac{\sqrt{x + \sqrt{1+x}}}{x^2} dx = \int \frac{\sqrt{x + \sqrt{x+1}}}{x^2} dx$$

input `integrate((x+(1+x)^(1/2))^(1/2)/x^2,x, algorithm="maxima")`

output `integrate(sqrt(x + sqrt(x + 1))/x^2, x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 188 vs. $2(59) = 118$.

Time = 0.36 (sec) , antiderivative size = 188, normalized size of antiderivative = 2.27

$$\int \frac{\sqrt{x + \sqrt{1+x}}}{x^2} dx =$$

$$\frac{2 \left(\sqrt{x + \sqrt{x+1}} - \sqrt{x+1} \right)^3 - 3 \left(\sqrt{x + \sqrt{x+1}} - \sqrt{x+1} \right)^2 - \sqrt{x + \sqrt{x+1}} + \sqrt{x+1} + 1}{\left(\sqrt{x + \sqrt{x+1}} - \sqrt{x+1} \right)^4 - 2 \left(\sqrt{x + \sqrt{x+1}} - \sqrt{x+1} \right)^2 + 4 \sqrt{x + \sqrt{x+1}} - 4 \sqrt{x+1}}$$

$$+ \frac{1}{2} \arctan \left(\sqrt{x + \sqrt{x+1}} - \sqrt{x+1} - 1 \right) - \frac{3}{4} \log \left(\left| \sqrt{x + \sqrt{x+1}} - \sqrt{x+1} + 2 \right| \right)$$

$$+ \frac{3}{4} \log \left(\left| \sqrt{x + \sqrt{x+1}} - \sqrt{x+1} \right| \right)$$

input `integrate((x+(1+x)^(1/2))^(1/2)/x^2,x, algorithm="giac")`

output `-(2*(sqrt(x + sqrt(x + 1)) - sqrt(x + 1))^3 - 3*(sqrt(x + sqrt(x + 1)) - sqrt(x + 1))^2 - sqrt(x + sqrt(x + 1)) + sqrt(x + 1) + 1)/((sqrt(x + sqrt(x + 1)) - sqrt(x + 1))^4 - 2*(sqrt(x + sqrt(x + 1)) - sqrt(x + 1))^2 + 4*sqrt(x + sqrt(x + 1)) - 4*sqrt(x + 1)) + 1/2*arctan(sqrt(x + sqrt(x + 1)) - sqrt(x + 1) - 1) - 3/4*log(abs(sqrt(x + sqrt(x + 1)) - sqrt(x + 1) + 2)) + 3/4*log(abs(sqrt(x + sqrt(x + 1)) - sqrt(x + 1))))`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{x + \sqrt{1+x}}}{x^2} dx = \int \frac{\sqrt{x + \sqrt{x+1}}}{x^2} dx$$

input `int((x + (x + 1)^(1/2))^(1/2)/x^2,x)`

output `int((x + (x + 1)^(1/2))^(1/2)/x^2, x)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.04

$$\int \frac{\sqrt{x + \sqrt{1+x}}}{x^2} dx = \frac{10 \operatorname{atan}\left(\sqrt{\sqrt{x+1}+x} + \sqrt{x+1} + 1\right) x - 20\sqrt{\sqrt{x+1}+x} - 15 \log\left(\frac{10\sqrt{\sqrt{x+1}+x} + 10\sqrt{x+1}}{\sqrt{5}}\right) x + 15 \log\left(\frac{2\sqrt{\sqrt{x+1}+x} + 2\sqrt{x+1} - 4}{\sqrt{5}}\right) x - 16x}{20x}$$

input `int((x+(1+x)^(1/2))^(1/2)/x^2,x)`

output `(10*atan(sqrt(sqrt(x + 1) + x) + sqrt(x + 1) + 1)*x - 20*sqrt(sqrt(x + 1) + x) - 15*log((10*sqrt(sqrt(x + 1) + x) + 10*sqrt(x + 1))/sqrt(5))*x + 15*log((2*sqrt(sqrt(x + 1) + x) + 2*sqrt(x + 1) - 4)/sqrt(5))*x - 16*x)/(20*x)`

3.18 $\int \sqrt{\sqrt{1 + \frac{1}{x}} + \frac{1}{x}} dx$

Optimal result	151
Mathematica [A] (verified)	152
Rubi [A] (verified)	152
Maple [F]	155
Fricas [A] (verification not implemented)	156
Sympy [F]	156
Maxima [F]	157
Giac [F]	157
Mupad [F(-1)]	157
Reduce [F]	158

Optimal result

Integrand size = 17, antiderivative size = 96

$$\int \sqrt{\sqrt{1 + \frac{1}{x}} + \frac{1}{x}} dx = \sqrt{\sqrt{1 + \frac{1}{x}} + \frac{1}{x}} x + \frac{1}{4} \arctan \left(\frac{3 + \sqrt{1 + \frac{1}{x}}}{2\sqrt{\sqrt{1 + \frac{1}{x}} + \frac{1}{x}}} \right) - \frac{3}{4} \operatorname{arctanh} \left(\frac{1 - 3\sqrt{1 + \frac{1}{x}}}{2\sqrt{\sqrt{1 + \frac{1}{x}} + \frac{1}{x}}} \right)$$

output `1/4*arctan(1/2*(3+(1+1/x)^(1/2))/(1/x+(1+1/x)^(1/2))^(1/2))-3/4*arctanh(1/2*(1-3*(1+1/x)^(1/2))/(1/x+(1+1/x)^(1/2))^(1/2))+x*(1/x+(1+1/x)^(1/2))^(1/2)`

Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.93

$$\int \sqrt{\sqrt{1 + \frac{1}{x}} + \frac{1}{x}} dx = \frac{1}{2} \left(2\sqrt{\sqrt{1 + \frac{1}{x}} + \frac{1}{x}} x + \arctan \left(1 + \sqrt{1 + \frac{1}{x}} - \sqrt{\sqrt{1 + \frac{1}{x}} + \frac{1}{x}} \right) + 3\operatorname{arctanh} \left(1 - \sqrt{1 + \frac{1}{x}} + \sqrt{\sqrt{1 + \frac{1}{x}} + \frac{1}{x}} \right) \right)$$

input

```
Integrate[Sqrt[Sqrt[1 + x^(-1)] + x^(-1)], x]
```

output

```
(2*Sqrt[Sqrt[1 + x^(-1)] + x^(-1)]*x + ArcTan[1 + Sqrt[1 + x^(-1)] - Sqrt[Sqrt[1 + x^(-1)] + x^(-1)]] + 3*ArcTanh[1 - Sqrt[1 + x^(-1)] + Sqrt[Sqrt[1 + x^(-1)] + x^(-1)]])/2
```

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.10, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.471$, Rules used = {7268, 1347, 27, 1366, 25, 1154, 217, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\sqrt{\frac{1}{x} + 1} + \frac{1}{x}} dx$$

↓ 7268

$$-2 \int \sqrt{1 + \frac{1}{x}} \sqrt{\sqrt{1 + \frac{1}{x}} + \frac{1}{x}} x^2 d\sqrt{1 + \frac{1}{x}}$$

↓ 1347

$$-2 \left(\frac{1}{2} \int \frac{(2\sqrt{1+\frac{1}{x}+1})x}{2\sqrt{\sqrt{1+\frac{1}{x}+\frac{1}{x}}}} d\sqrt{1+\frac{1}{x}} - \frac{1}{2} \sqrt{\sqrt{\frac{1}{x}+1+\frac{1}{x}x}} \right)$$

↓ 27

$$-2 \left(-\frac{1}{4} \int -\frac{(2\sqrt{1+\frac{1}{x}+1})x}{\sqrt{\sqrt{1+\frac{1}{x}+\frac{1}{x}}}} d\sqrt{1+\frac{1}{x}} - \frac{1}{2} \sqrt{\sqrt{\frac{1}{x}+1+\frac{1}{x}x}} \right)$$

↓ 1366

$$-2 \left(\frac{1}{4} \left(-\frac{3}{2} \int \frac{1}{(1-\sqrt{1+\frac{1}{x}})\sqrt{\sqrt{1+\frac{1}{x}+\frac{1}{x}}}} d\sqrt{1+\frac{1}{x}} - \frac{1}{2} \int -\frac{1}{(\sqrt{1+\frac{1}{x}+1})\sqrt{\sqrt{1+\frac{1}{x}+\frac{1}{x}}}} d\sqrt{1+\frac{1}{x}} \right) - \frac{1}{2} \sqrt{\sqrt{\frac{1}{x}+1+\frac{1}{x}x}} \right)$$

↓ 25

$$-2 \left(\frac{1}{4} \left(\frac{1}{2} \int \frac{1}{(\sqrt{1+\frac{1}{x}+1})\sqrt{\sqrt{1+\frac{1}{x}+\frac{1}{x}}}} d\sqrt{1+\frac{1}{x}} - \frac{3}{2} \int \frac{1}{(1-\sqrt{1+\frac{1}{x}})\sqrt{\sqrt{1+\frac{1}{x}+\frac{1}{x}}}} d\sqrt{1+\frac{1}{x}} \right) - \frac{1}{2} \sqrt{\sqrt{\frac{1}{x}+1+\frac{1}{x}x}} \right)$$

↓ 1154

$$-2 \left(\frac{1}{4} \left(3 \int \frac{1}{3-\frac{1}{x}} d\frac{1-3\sqrt{1+\frac{1}{x}}}{\sqrt{\sqrt{1+\frac{1}{x}+\frac{1}{x}}}} - \int \frac{1}{-5-\frac{1}{x}} d\left(-\frac{\sqrt{1+\frac{1}{x}+3}}{\sqrt{\sqrt{1+\frac{1}{x}+\frac{1}{x}}}} \right) \right) - \frac{1}{2} \sqrt{\sqrt{\frac{1}{x}+1+\frac{1}{x}x}} \right)$$

↓ 217

$$-2 \left(\frac{1}{4} \left(3 \int \frac{1}{3-\frac{1}{x}} d\frac{1-3\sqrt{1+\frac{1}{x}}}{\sqrt{\sqrt{1+\frac{1}{x}+\frac{1}{x}}}} - \frac{1}{2} \arctan \left(\frac{\sqrt{\frac{1}{x}+1+3}}{2\sqrt{\sqrt{\frac{1}{x}+1+\frac{1}{x}x}}} \right) \right) - \frac{1}{2} \sqrt{\sqrt{\frac{1}{x}+1+\frac{1}{x}x}} \right)$$

↓ 219

$$-2 \left(\frac{1}{4} \left(\frac{3}{2} \operatorname{arctanh} \left(\frac{1-3\sqrt{\frac{1}{x}+1}}{2\sqrt{\sqrt{\frac{1}{x}+1+\frac{1}{x}x}}} \right) - \frac{1}{2} \arctan \left(\frac{\sqrt{\frac{1}{x}+1+3}}{2\sqrt{\sqrt{\frac{1}{x}+1+\frac{1}{x}x}}} \right) \right) - \frac{1}{2} \sqrt{\sqrt{\frac{1}{x}+1+\frac{1}{x}x}} \right)$$

input `Int[Sqrt[Sqrt[1 + x(-1)] + x(-1)], x]`

output `-2*(-1/2*(Sqrt[Sqrt[1 + x(-1)] + x(-1)]*x) + (-1/2*ArcTan[(3 + Sqrt[1 + x(-1)])/(2*Sqrt[Sqrt[1 + x(-1)] + x(-1)])]) + (3*ArcTanh[(1 - 3*Sqrt[1 + x(-1)])/(2*Sqrt[Sqrt[1 + x(-1)] + x(-1)])])/2)/4`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 1154 `Int[1/(((d_.) + (e_.)*(x_))*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]), x_Symbol] := Simp[-2 Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 1347

```
Int[((g_.) + (h_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_)*((d_) + (e_.)*(x_) + (f_.)*(x_)^2)^(q_), x_Symbol] := Simp[(a*h - g*c*x)*(a + c*x^2)^(p + 1)*((d + e*x + f*x^2)^q/(2*a*c*(p + 1))), x] + Simp[2/(4*a*c*(p + 1)) Int[(a + c*x^2)^(p + 1)*(d + e*x + f*x^2)^(q - 1)*Simp[g*c*d*(2*p + 3) - a*(h*e*q) + (g*c*e*(2*p + q + 3) - a*(2*h*f*q))*x + g*c*f*(2*p + 2*q + 3)*x^2, x], x] /; FreeQ[{a, c, d, e, f, g, h}, x] && NeQ[e^2 - 4*d*f, 0] && LtQ[p, -1] && GtQ[q, 0]
```

rule 1366

```
Int[((g_.) + (h_.)*(x_))/(((a_) + (c_.)*(x_)^2)*Sqrt[(d_.) + (e_.)*(x_) + (f_.)*(x_)^2]), x_Symbol] := With[{q = Rt[(-a)*c, 2]}, Simp[(h/2 + c*(g/(2*q))) Int[1/((-q + c*x)*Sqrt[d + e*x + f*x^2]), x], x] + Simp[(h/2 - c*(g/(2*q))) Int[1/((q + c*x)*Sqrt[d + e*x + f*x^2]), x], x] /; FreeQ[{a, c, d, e, f, g, h}, x] && NeQ[e^2 - 4*d*f, 0] && PosQ[(-a)*c]
```

rule 7268

```
Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfQuotientOfLinears[u, x]}, Simp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2]])], x] /; !FalseQ[lst]
```

Maple [F]

$$\int \sqrt{\frac{1}{x} + \sqrt{1 + \frac{1}{x}}} dx$$

input

```
int((1/x+(1+1/x)^(1/2))^(1/2),x)
```

output

```
int((1/x+(1+1/x)^(1/2))^(1/2),x)
```

Fricas [A] (verification not implemented)

Time = 1.68 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.27

$$\int \sqrt{\sqrt{1 + \frac{1}{x} + \frac{1}{x}}} dx = x \sqrt{\frac{x \sqrt{\frac{x+1}{x}} + 1}{x}} + \frac{1}{4} \arctan \left(\frac{2 \left(x \sqrt{\frac{x+1}{x}} - 3x \right) \sqrt{\frac{x \sqrt{\frac{x+1}{x}} + 1}}{x}}{8x - 1} \right) + \frac{3}{4} \log \left(2 \left(x \sqrt{\frac{x+1}{x}} + x \right) \sqrt{\frac{x \sqrt{\frac{x+1}{x}} + 1}{x}} + 2x \sqrt{\frac{x+1}{x}} + 2x + 3 \right)$$

input `integrate((1/x+(1+1/x)^(1/2))^(1/2),x, algorithm="fricas")`

output `x*sqrt((x*sqrt((x + 1)/x) + 1)/x) + 1/4*arctan(2*(x*sqrt((x + 1)/x) - 3*x)*sqrt((x*sqrt((x + 1)/x) + 1)/x)/(8*x - 1)) + 3/4*log(2*(x*sqrt((x + 1)/x) + x)*sqrt((x*sqrt((x + 1)/x) + 1)/x) + 2*x*sqrt((x + 1)/x) + 2*x + 3)`

Sympy [F]

$$\int \sqrt{\sqrt{1 + \frac{1}{x} + \frac{1}{x}}} dx = \int \sqrt{\sqrt{1 + \frac{1}{x} + \frac{1}{x}}} dx$$

input `integrate((1/x+(1+1/x)**(1/2))**(1/2),x)`

output `Integral(sqrt(sqrt(1 + 1/x) + 1/x), x)`

Maxima [F]

$$\int \sqrt{\sqrt{1 + \frac{1}{x}} + \frac{1}{x}} dx = \int \sqrt{\sqrt{\frac{1}{x} + 1} + \frac{1}{x}} dx$$

input `integrate((1/x+(1+1/x)^(1/2))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(sqrt(1/x + 1) + 1/x), x)`

Giac [F]

$$\int \sqrt{\sqrt{1 + \frac{1}{x}} + \frac{1}{x}} dx = \int \sqrt{\sqrt{\frac{1}{x} + 1} + \frac{1}{x}} dx$$

input `integrate((1/x+(1+1/x)^(1/2))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(sqrt(1/x + 1) + 1/x), x)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{\sqrt{1 + \frac{1}{x}} + \frac{1}{x}} dx = \int \sqrt{\sqrt{\frac{1}{x} + 1} + \frac{1}{x}} dx$$

input `int(((1/x + 1)^(1/2) + 1/x)^(1/2),x)`

output `int(((1/x + 1)^(1/2) + 1/x)^(1/2), x)`

Reduce **[F]**

$$\int \sqrt{\sqrt{1 + \frac{1}{x}} + \frac{1}{x}} dx = \int \sqrt{\frac{1}{x} + \sqrt{1 + \frac{1}{x}}} dx$$

input `int((1/x+(1+1/x)^(1/2))^(1/2),x)`

output `int((1/x+(1+1/x)^(1/2))^(1/2),x)`

3.19 $\int \frac{\sqrt{1+e^{-x}}}{-e^{-x}+e^x} dx$

Optimal result	159
Mathematica [B] (verified)	159
Rubi [A] (verified)	160
Maple [B] (verified)	162
Fricas [A] (verification not implemented)	162
Sympy [F]	163
Maxima [A] (verification not implemented)	163
Giac [B] (verification not implemented)	163
Mupad [F(-1)]	164
Reduce [F]	164

Optimal result

Integrand size = 25, antiderivative size = 25

$$\int \frac{\sqrt{1+e^{-x}}}{-e^{-x}+e^x} dx = -\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{1+e^{-x}}}{\sqrt{2}}\right)$$

output

```
-arctanh(1/2*(1+exp(-x))^(1/2)*2^(1/2))*2^(1/2)
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 65 vs. 2(25) = 50.

Time = 0.11 (sec) , antiderivative size = 65, normalized size of antiderivative = 2.60

$$\int \frac{\sqrt{1+e^{-x}}}{-e^{-x}+e^x} dx = -\frac{\sqrt{2}e^{x/2}\sqrt{1+e^{-x}}\operatorname{arctanh}\left(\frac{1-e^x+e^{x/2}\sqrt{1+e^x}}{\sqrt{2}}\right)}{\sqrt{1+e^x}}$$

input

```
Integrate[Sqrt[1 + E^(-x)]/(-E^(-x) + E^x), x]
```

output

```
-((Sqrt[2]*E^(x/2)*Sqrt[1 + E^(-x)]*ArcTanh[(1 - E^x + E^(x/2)*Sqrt[1 + E^x])/Sqrt[2]])/Sqrt[1 + E^x])
```


Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$, Rules used = {2720, 25, 1776, 1388, 946, 25, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{e^{-x} + 1}}{e^x - e^{-x}} dx \\
 & \quad \downarrow \text{2720} \\
 & \int -\frac{\sqrt{e^{-x} + 1}}{1 - e^{2x}} de^x \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{\sqrt{1 + e^{-x}}}{1 - e^{2x}} de^x \\
 & \quad \downarrow \text{1776} \\
 & -\int \frac{e^{-2x} \sqrt{1 + e^{-x}}}{-1 + e^{-2x}} de^x \\
 & \quad \downarrow \text{1388} \\
 & -\int \frac{e^{-2x}}{(-1 + e^{-x}) \sqrt{1 + e^{-x}}} de^x \\
 & \quad \downarrow \text{946} \\
 & \int -\frac{1}{(1 - e^{-x}) \sqrt{e^{-x} + 1}} de^{-x} \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{1}{(1 - e^{-x}) \sqrt{1 + e^{-x}}} de^{-x} \\
 & \quad \downarrow \text{73} \\
 & -2 \int \frac{1}{2 - e^{2x}} d\sqrt{1 + e^{-x}} \\
 & \quad \downarrow \text{219}
 \end{aligned}$$

$$-\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{e^{-x}+1}}{\sqrt{2}}\right)$$

input `Int[Sqrt[1 + E^(-x)]/(-E^(-x) + E^x), x]`

output `-(Sqrt[2]*ArcTanh[Sqrt[1 + E^(-x)]/Sqrt[2]])`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b)]^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])`

rule 946 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.
, x_Symbol] := Simp[1/n Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^n],
x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && EqQ[m - n
+ 1, 0]`

rule 1388 `Int[(u_.)*((a_) + (c_.)*(x_)^(n2_.))^(p_.)*((d_) + (e_.)*(x_)^(n_))^(q_.),
x_Symbol] := Int[u*(d + e*x^n)^(p + q)*(a/d + (c/e)*x^n)^p, x] /; FreeQ[{a,
c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[c*d^2 + a*e^2, 0] && (Integer
Q[p] || (GtQ[a, 0] && GtQ[d, 0]))`

rule 1776 `Int[((a_.) + (c_.)*(x_)^(mn2_.))^(p_.)*((d_) + (e_.)*(x_)^(n_.))^(q_.), x_Symbol] :> Int[(d + e*x^n)^q*(c + a*x^(2*n))^p/x^(2*n*p), x] /; FreeQ[{a, c, d, e, n, q}, x] && EqQ[mn2, -2*n] && IntegerQ[p]`

rule 2720 `Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 48 vs. 2(19) = 38.

Time = 0.07 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.96

method	result	size
default	$-\frac{\sqrt{(1+e^x)e^{-x}} e^x \sqrt{2} \operatorname{arctanh}\left(\frac{(1+3e^x)\sqrt{2}}{4\sqrt{e^x+e^{2x}}}\right)}{2\sqrt{(1+e^x)e^x}}$	49

input `int((1+exp(-x))^(1/2)/(-exp(-x)+exp(x)),x,method=_RETURNVERBOSE)`

output `-1/2*((1+exp(x))/exp(x))^(1/2)*exp(x)/((1+exp(x))*exp(x))^(1/2)*2^(1/2)*arctanh(1/4*(1+3*exp(x))*2^(1/2)/(exp(x)^2+exp(x))^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.36

$$\int \frac{\sqrt{1+e^{-x}}}{-e^{-x}+e^x} dx = \frac{1}{2} \sqrt{2} \log \left(\frac{2\sqrt{2}\sqrt{e^x+1}e^{(\frac{1}{2}x)} - 3e^x - 1}{e^x - 1} \right)$$

input `integrate((1+exp(-x))^(1/2)/(-exp(-x)+exp(x)),x, algorithm="fricas")`

output `1/2*sqrt(2)*log((2*sqrt(2)*sqrt(e^x + 1)*e^(1/2*x) - 3*e^x - 1)/(e^x - 1))`

Sympy [F]

$$\int \frac{\sqrt{1+e^{-x}}}{-e^{-x}+e^x} dx = \int \frac{\sqrt{1+e^{-x}}e^x}{(e^x-1)(e^x+1)} dx$$

input `integrate((1+exp(-x))**(1/2)/(-exp(-x)+exp(x)),x)`

output `Integral(sqrt(1 + exp(-x))*exp(x)/((exp(x) - 1)*(exp(x) + 1)), x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.44

$$\int \frac{\sqrt{1+e^{-x}}}{-e^{-x}+e^x} dx = \frac{1}{2} \sqrt{2} \log \left(-\frac{\sqrt{2} - \sqrt{e^{(-x)} + 1}}{\sqrt{2} + \sqrt{e^{(-x)} + 1}} \right)$$

input `integrate((1+exp(-x))^(1/2)/(-exp(-x)+exp(x)),x, algorithm="maxima")`

output `1/2*sqrt(2)*log(-(sqrt(2) - sqrt(e^(-x) + 1))/(sqrt(2) + sqrt(e^(-x) + 1)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 55 vs. $2(19) = 38$.

Time = 0.15 (sec) , antiderivative size = 55, normalized size of antiderivative = 2.20

$$\int \frac{\sqrt{1+e^{-x}}}{-e^{-x}+e^x} dx = \frac{1}{2} \sqrt{2} \log \left(\frac{\left| -2\sqrt{2} + 2\sqrt{e^{(2x)} + e^x} - 2e^x + 2 \right|}{\left| 2\sqrt{2} + 2\sqrt{e^{(2x)} + e^x} - 2e^x + 2 \right|} \right)$$

input `integrate((1+exp(-x))^(1/2)/(-exp(-x)+exp(x)),x, algorithm="giac")`

output `1/2*sqrt(2)*log(abs(-2*sqrt(2) + 2*sqrt(e^(2*x) + e^x) - 2*e^x + 2)/abs(2*sqrt(2) + 2*sqrt(e^(2*x) + e^x) - 2*e^x + 2))`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{1+e^{-x}}}{-e^{-x}+e^x} dx = -\int \frac{\sqrt{e^{-x}+1}}{e^{-x}-e^x} dx$$

input `int(-(exp(-x) + 1)^(1/2)/(exp(-x) - exp(x)),x)`

output `-int((exp(-x) + 1)^(1/2)/(exp(-x) - exp(x)), x)`

Reduce [F]

$$\int \frac{\sqrt{1+e^{-x}}}{-e^{-x}+e^x} dx = \int \frac{e^x \sqrt{e^x+1}}{e^{\frac{5x}{2}} - e^{\frac{x}{2}}} dx$$

input `int((1+exp(-x))^(1/2)/(-exp(-x)+exp(x)),x)`

output `int((e**x*sqrt(e**x + 1))/(e**((5*x)/2) - e**(x/2)),x)`

3.20 $\int \sqrt{1 + e^{-x}} \operatorname{csch}(x) dx$

Optimal result	165
Mathematica [B] (verified)	165
Rubi [A] (verified)	166
Maple [A] (verified)	168
Fricas [B] (verification not implemented)	169
Sympy [F]	169
Maxima [A] (verification not implemented)	170
Giac [B] (verification not implemented)	170
Mupad [F(-1)]	170
Reduce [F]	171

Optimal result

Integrand size = 14, antiderivative size = 25

$$\int \sqrt{1 + e^{-x}} \operatorname{csch}(x) dx = -2\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1 + e^{-x}}}{\sqrt{2}}\right)$$

output `-2*arctanh(1/2*(1+exp(-x))^(1/2)*2^(1/2))*2^(1/2)`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 57 vs. 2(25) = 50.

Time = 0.05 (sec) , antiderivative size = 57, normalized size of antiderivative = 2.28

$$\int \sqrt{1 + e^{-x}} \operatorname{csch}(x) dx = -\frac{2\sqrt{2}e^{x/2}\sqrt{1 + e^{-x}} \operatorname{arctanh}\left(\frac{\sqrt{2}e^{x/2}}{\sqrt{1+e^x}}\right)}{\sqrt{1 + e^x}}$$

input `Integrate[Sqrt[1 + E^(-x)]*Csch[x], x]`

output `(-2*Sqrt[2]*E^(x/2)*Sqrt[1 + E^(-x)]*ArcTanh[(Sqrt[2]*E^(x/2))/Sqrt[1 + E^x]])/Sqrt[1 + E^x]`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {2720, 27, 1776, 1388, 946, 25, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{e^{-x} + 1} \operatorname{csch}(x) dx \\
 & \quad \downarrow \text{2720} \\
 & \int -\frac{2\sqrt{e^{-x} + 1}}{1 - e^{2x}} de^x \\
 & \quad \downarrow \text{27} \\
 & -2 \int \frac{\sqrt{1 + e^{-x}}}{1 - e^{2x}} de^x \\
 & \quad \downarrow \text{1776} \\
 & -2 \int \frac{e^{-2x} \sqrt{1 + e^{-x}}}{-1 + e^{-2x}} de^x \\
 & \quad \downarrow \text{1388} \\
 & -2 \int \frac{e^{-2x}}{(-1 + e^{-x}) \sqrt{1 + e^{-x}}} de^x \\
 & \quad \downarrow \text{946} \\
 & 2 \int -\frac{1}{(1 - e^{-x}) \sqrt{1 + e^{-x}}} de^{-x} \\
 & \quad \downarrow \text{25} \\
 & -2 \int \frac{1}{(1 - e^{-x}) \sqrt{1 + e^{-x}}} de^{-x} \\
 & \quad \downarrow \text{73} \\
 & -4 \int \frac{1}{2 - e^{2x}} d\sqrt{1 + e^{-x}} \\
 & \quad \downarrow \text{219}
 \end{aligned}$$

$$-2\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{e^{-x}+1}}{\sqrt{2}}\right)$$

input `Int[Sqrt[1 + E^(-x)]*Csch[x],x]`

output `-2*Sqrt[2]*ArcTanh[Sqrt[1 + E^(-x)]/Sqrt[2]]`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 946 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] := Simp[1/n Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && EqQ[m - n + 1, 0]`

rule 1388 `Int[(u_)*((a_) + (c_)*(x_)^(n2_))^(p_)*((d_) + (e_)*(x_)^(n_))^(q_), x_Symbol] := Int[u*(d + e*x^n)^(p + q)*(a/d + (c/e)*x^n)^p, x] /; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[c*d^2 + a*e^2, 0] && (IntegerQ[p] || (GtQ[a, 0] && GtQ[d, 0]))`

rule 1776 `Int[((a_) + (c_)*(x_)^(mn2_))^(p_)*((d_) + (e_)*(x_)^(n_))^(q_), x_Symbol] := Int[((d + e*x^n)^q*(c + a*x^(2*n))^p)/x^(2*n*p), x] /; FreeQ[{a, c, d, e, n, q}, x] && EqQ[mn2, -2*n] && IntegerQ[p]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.32

method	result	size
default	$-2\sqrt{2} \sqrt{\frac{1}{\tanh(\frac{x}{2})+1}} \sqrt{\tanh(\frac{x}{2}) + 1} \operatorname{arctanh}\left(\sqrt{\tanh(\frac{x}{2}) + 1}\right)$	33

input `int((1+exp(-x))^(1/2)/sinh(x),x,method=_RETURNVERBOSE)`

output `-2*2^(1/2)*(1/(tanh(1/2*x)+1))^(1/2)*(tanh(1/2*x)+1)^(1/2)*arctanh((tanh(1/2*x)+1)^(1/2))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 55 vs. $2(19) = 38$.

Time = 0.07 (sec) , antiderivative size = 55, normalized size of antiderivative = 2.20

$$\int \sqrt{1 + e^{-x}} \operatorname{csch}(x) dx$$

$$= \sqrt{2} \log \left(\frac{2 (\sqrt{2} \cosh(x) + \sqrt{2} \sinh(x)) \sqrt{\frac{\cosh(x) + \sinh(x) + 1}{\cosh(x) + \sinh(x)}} - 3 \cosh(x) - 3 \sinh(x) - 1}{\cosh(x) + \sinh(x) - 1} \right)$$

input `integrate((1+exp(-x))^(1/2)/sinh(x),x, algorithm="fricas")`

output `sqrt(2)*log((2*(sqrt(2)*cosh(x) + sqrt(2)*sinh(x))*sqrt((cosh(x) + sinh(x) + 1)/(cosh(x) + sinh(x))) - 3*cosh(x) - 3*sinh(x) - 1)/(cosh(x) + sinh(x) - 1))`

Sympy [F]

$$\int \sqrt{1 + e^{-x}} \operatorname{csch}(x) dx = \int \frac{\sqrt{1 + e^{-x}}}{\sinh(x)} dx$$

input `integrate((1+exp(-x))**(1/2)/sinh(x),x)`

output `Integral(sqrt(1 + exp(-x))/sinh(x), x)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.40

$$\int \sqrt{1 + e^{-x}} \operatorname{csch}(x) dx = \sqrt{2} \log \left(-\frac{\sqrt{2} - \sqrt{e^{(-x)} + 1}}{\sqrt{2} + \sqrt{e^{(-x)} + 1}} \right)$$

input `integrate((1+exp(-x))^(1/2)/sinh(x),x, algorithm="maxima")`

output `sqrt(2)*log(-(sqrt(2) - sqrt(e^(-x) + 1))/(sqrt(2) + sqrt(e^(-x) + 1)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 54 vs. 2(19) = 38.

Time = 0.14 (sec) , antiderivative size = 54, normalized size of antiderivative = 2.16

$$\int \sqrt{1 + e^{-x}} \operatorname{csch}(x) dx = \sqrt{2} \log \left(\frac{\left| -2\sqrt{2} + 2\sqrt{e^{(2x)} + e^x} - 2e^x + 2 \right|}{\left| 2\sqrt{2} + 2\sqrt{e^{(2x)} + e^x} - 2e^x + 2 \right|} \right)$$

input `integrate((1+exp(-x))^(1/2)/sinh(x),x, algorithm="giac")`

output `sqrt(2)*log(abs(-2*sqrt(2) + 2*sqrt(e^(2*x) + e^x) - 2*e^x + 2)/abs(2*sqrt(2) + 2*sqrt(e^(2*x) + e^x) - 2*e^x + 2))`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{1 + e^{-x}} \operatorname{csch}(x) dx = \int \frac{\sqrt{e^{-x} + 1}}{\sinh(x)} dx$$

input `int((exp(-x) + 1)^(1/2)/sinh(x),x)`

output `int((exp(-x) + 1)^(1/2)/sinh(x), x)`

Reduce [F]

$$\int \sqrt{1 + e^{-x}} \operatorname{csch}(x) dx = \int \frac{\sqrt{e^x + 1}}{e^{\frac{x}{2}} \sinh(x)} dx$$

input `int((1+exp(-x))^(1/2)/sinh(x),x)`

output `int(sqrt(e**x + 1)/(e**(x/2)*sinh(x)),x)`

3.21 $\int \frac{1}{(\cos(x)+\cos(3x))^5} dx$

Optimal result	172
Mathematica [A] (verified)	173
Rubi [A] (verified)	173
Maple [A] (verified)	177
Fricas [B] (verification not implemented)	177
Sympy [F]	178
Maxima [B] (verification not implemented)	178
Giac [A] (verification not implemented)	179
Mupad [B] (verification not implemented)	180
Reduce [F]	181

Optimal result

Integrand size = 9, antiderivative size = 108

$$\int \frac{1}{(\cos(x) + \cos(3x))^5} dx = -\frac{523}{256} \operatorname{arctanh}(\sin(x)) + \frac{1483 \operatorname{arctanh}(\sqrt{2} \sin(x))}{512\sqrt{2}}$$

$$+ \frac{\sin(x)}{32(1-2\sin^2(x))^4} - \frac{17\sin(x)}{192(1-2\sin^2(x))^3}$$

$$+ \frac{203\sin(x)}{768(1-2\sin^2(x))^2} - \frac{437\sin(x)}{512(1-2\sin^2(x))}$$

$$- \frac{43}{256} \sec(x) \tan(x) - \frac{1}{128} \sec^3(x) \tan(x)$$

output

```
-523/256*arctanh(sin(x))+1/32*sin(x)/(1-2*sin(x)^2)^4-17/192*sin(x)/(1-2*
sin(x)^2)^3+203/768*sin(x)/(1-2*sin(x)^2)^2-437/512*sin(x)/(1-2*sin(x)^2)+
483/1024*arctanh(sin(x)*2^(1/2))*2^(1/2)-43/256*sec(x)*tan(x)-1/128*sec(x)
^3*tan(x)
```

Mathematica [A] (verified)

Time = 3.88 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.97

$$\int \frac{1}{(\cos(x) + \cos(3x))^5} dx$$

$$= \frac{12552 \log\left(\cos\left(\frac{x}{2}\right) - \sin\left(\frac{x}{2}\right)\right) - 12552 \log\left(\cos\left(\frac{x}{2}\right) + \sin\left(\frac{x}{2}\right)\right) - 4449\sqrt{2} \log\left(\sqrt{2} - 2\sin(x)\right) + 4449\sqrt{2} \log\left(\sqrt{2} + 2\sin(x)\right) - 12}{6144}$$

input

```
Integrate[(Cos[x] + Cos[3*x])^(-5), x]
```

output

```
(12552*Log[Cos[x/2] - Sin[x/2]] - 12552*Log[Cos[x/2] + Sin[x/2]] - 4449*Sqrt[2]*Log[Sqrt[2] - 2*Sin[x]] + 4449*Sqrt[2]*Log[Sqrt[2] + 2*Sin[x]] - 12/(Cos[x/2] - Sin[x/2])^4 - 516/(Cos[x/2] - Sin[x/2])^2 + 12/(Cos[x/2] + Sin[x/2])^4 + 516/(Cos[x/2] + Sin[x/2])^2 - 136/(Cos[x] - Sin[x])^3 - 2622/(Cos[x] - Sin[x]) + 136/(Cos[x] + Sin[x])^3 + 2622/(Cos[x] + Sin[x]) + 6*Sec[2*x]^4*(190*Sin[x] + 79*(-Sin[3*x] + Sin[5*x])))/6144
```

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.70, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 1.556$, Rules used = {3042, 4825, 27, 316, 27, 402, 402, 402, 402, 27, 402, 27, 397, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(\cos(x) + \cos(3x))^5} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{1}{(\cos(x) + \cos(3x))^5} dx$$

$$\downarrow \text{4825}$$

$$\int \frac{1}{32(1 - 2\sin^2(x))^5(1 - \sin^2(x))^3} d\sin(x)$$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{1}{32} \int \frac{1}{(1-2\sin^2(x))^5 (1-\sin^2(x))^3} d\sin(x) \\
& \downarrow 316 \\
& \frac{1}{32} \left(\frac{1}{8} \int \frac{2(3-11\sin^2(x))}{(1-2\sin^2(x))^4 (1-\sin^2(x))^3} d\sin(x) + \frac{\sin(x)}{4(1-2\sin^2(x))^4 (1-\sin^2(x))^2} \right) \\
& \downarrow 27 \\
& \frac{1}{32} \left(\frac{1}{4} \int \frac{3-11\sin^2(x)}{(1-2\sin^2(x))^4 (1-\sin^2(x))^3} d\sin(x) + \frac{\sin(x)}{4(1-2\sin^2(x))^4 (1-\sin^2(x))^2} \right) \\
& \downarrow 402 \\
& \frac{1}{32} \left(\frac{1}{4} \left(\frac{1}{6} \int \frac{45\sin^2(x)+23}{(1-2\sin^2(x))^3 (1-\sin^2(x))^3} d\sin(x) - \frac{5\sin(x)}{6(1-2\sin^2(x))^3 (1-\sin^2(x))^2} \right) + \frac{\sin(x)}{4(1-2\sin^2(x))^4 (1-\sin^2(x))^2} \right) \\
& \downarrow 402 \\
& \frac{1}{32} \left(\frac{1}{4} \left(\frac{1}{6} \left(\frac{1}{4} \int \frac{1-637\sin^2(x)}{(1-2\sin^2(x))^2 (1-\sin^2(x))^3} d\sin(x) + \frac{91\sin(x)}{4(1-2\sin^2(x))^2 (1-\sin^2(x))^2} \right) - \frac{5\sin(x)}{6(1-2\sin^2(x))^3 (1-\sin^2(x))^2} \right) \right) \\
& \downarrow 402 \\
& \frac{1}{32} \left(\frac{1}{4} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{1}{2} \int \frac{3175\sin^2(x)+637}{(1-2\sin^2(x))(1-\sin^2(x))^3} d\sin(x) - \frac{635\sin(x)}{2(1-2\sin^2(x))(1-\sin^2(x))^2} \right) + \frac{91\sin(x)}{4(1-2\sin^2(x))^3 (1-\sin^2(x))^2} \right) \right) \right) \\
& \downarrow 402 \\
& \frac{1}{32} \left(\frac{1}{4} \left(\frac{1}{6} \left(\frac{1}{4} \left(-\frac{1}{4} \int -\frac{24(953\sin^2(x)+265)}{(1-2\sin^2(x))(1-\sin^2(x))^2} d\sin(x) - \frac{953\sin(x)}{(1-\sin^2(x))^2} \right) - \frac{635\sin(x)}{2(1-2\sin^2(x))(1-\sin^2(x))^2} \right) \right) \right) \\
& \downarrow 27 \\
& \frac{1}{32} \left(\frac{1}{4} \left(\frac{1}{6} \left(\frac{1}{4} \left(6 \int \frac{953\sin^2(x)+265}{(1-2\sin^2(x))(1-\sin^2(x))^2} d\sin(x) - \frac{953\sin(x)}{(1-\sin^2(x))^2} \right) - \frac{635\sin(x)}{2(1-2\sin^2(x))(1-\sin^2(x))^2} \right) \right) \right) \\
& \downarrow 402
\end{aligned}$$

$$\frac{1}{32} \left(\frac{1}{4} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{1}{2} \left(6 \left(-\frac{1}{2} \int -\frac{4(609 \sin^2(x) + 437)}{(1 - 2 \sin^2(x))(1 - \sin^2(x))} d \sin(x) - \frac{609 \sin(x)}{1 - \sin^2(x)} \right) - \frac{953 \sin(x)}{(1 - \sin^2(x))^2} \right) - \frac{953 \sin(x)}{2(1 - \sin^2(x))} \right) \right) \right) \right)$$

↓ 27

$$\frac{1}{32} \left(\frac{1}{4} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{1}{2} \left(6 \left(2 \int \frac{609 \sin^2(x) + 437}{(1 - 2 \sin^2(x))(1 - \sin^2(x))} d \sin(x) - \frac{609 \sin(x)}{1 - \sin^2(x)} \right) - \frac{953 \sin(x)}{(1 - \sin^2(x))^2} \right) - \frac{953 \sin(x)}{2(1 - \sin^2(x))} \right) \right) \right) \right)$$

↓ 397

$$\frac{1}{32} \left(\frac{1}{4} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{1}{2} \left(6 \left(2 \left(1483 \int \frac{1}{1 - 2 \sin^2(x)} d \sin(x) - 1046 \int \frac{1}{1 - \sin^2(x)} d \sin(x) \right) - \frac{609 \sin(x)}{1 - \sin^2(x)} \right) - \frac{953 \sin(x)}{(1 - \sin^2(x))^2} \right) \right) \right) \right) \right)$$

↓ 219

$$\frac{1}{32} \left(\frac{1}{4} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{1}{2} \left(6 \left(2 \left(\frac{1483 \operatorname{arctanh}(\sqrt{2} \sin(x))}{\sqrt{2}} - 1046 \operatorname{arctanh}(\sin(x)) \right) - \frac{609 \sin(x)}{1 - \sin^2(x)} \right) - \frac{953 \sin(x)}{(1 - \sin^2(x))^2} \right) \right) \right) \right) \right)$$

input `Int[(Cos[x] + Cos[3*x])^(-5),x]`

output `(Sin[x]/(4*(1 - 2*SIN[x]^2)^4*(1 - SIN[x]^2)^2) + ((-5*SIN[x])/(6*(1 - 2*SIN[x]^2)^3*(1 - SIN[x]^2)^2) + ((91*SIN[x])/(4*(1 - 2*SIN[x]^2)^2*(1 - SIN[x]^2)^2) + ((-635*SIN[x])/(2*(1 - 2*SIN[x]^2)*(1 - SIN[x]^2)^2) + ((-953*SIN[x])/(1 - SIN[x]^2)^2 + 6*(2*(-1046*ArcTanh[SIN[x]] + (1483*ArcTanh[Sqrt[2]*Sin[x]])/Sqrt[2]) - (609*SIN[x])/(1 - SIN[x]^2))))/2)/4)/6)/4)/32`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 316 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp`
`p[(-b)*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(2*a*(p + 1)*(b*c - a*d))`
`), x] + Simp[1/(2*a*(p + 1)*(b*c - a*d)) Int[(a + b*x^2)^(p + 1)*(c + d*x`
`^2)^q*Simp[b*c + 2*(p + 1)*(b*c - a*d) + d*b*(2*(p + q + 2) + 1)*x^2, x], x`
`] /; FreeQ[{a, b, c, d, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !`
`(!IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, 2,`
`p, q, x]`

rule 397 `Int[((e_) + (f_.)*(x_)^2)/(((a_) + (b_.)*(x_)^2)*((c_) + (d_.)*(x_)^2)), x_`
`Symbol] := Simp[(b*e - a*f)/(b*c - a*d) Int[1/(a + b*x^2), x], x] - Simp[`
`(d*e - c*f)/(b*c - a*d) Int[1/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d, e`
`, f}, x]`

rule 402 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_.)*((e_) + (f_.)*(x`
`_)^2), x_Symbol] := Simp[(-b*e - a*f)*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^`
`(q + 1)/(a*2*(b*c - a*d)*(p + 1)), x] + Simp[1/(a*2*(b*c - a*d)*(p + 1))`
`Int[(a + b*x^2)^(p + 1)*(c + d*x^2)^q*Simp[c*(b*e - a*f) + e*2*(b*c - a*d)`
`*(p + 1) + d*(b*e - a*f)*(2*(p + q + 2) + 1)*x^2, x], x], x] /; FreeQ[{a, b`
`, c, d, e, f, q}, x] && LtQ[p, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear`
`Q[u, x]`

rule 4825 `Int[(cos[(m_.)*((c_.) + (d_.)*(x_))]*(a_.) + cos[(n_.)*((c_.) + (d_.)*(x_))`
`]*(b_.))^(p_), x_Symbol] := Simp[1/d Subst[Int[Simplify[TrigExpand[a*Cos[`
`m*ArcSin[x]] + b*Cos[n*ArcSin[x]]]]^p/Sqrt[1 - x^2], x], x, Sin[c + d*x]],`
`x] /; FreeQ[{a, b, c, d}, x] && ILtQ[(p - 1)/2, 0] && IntegerQ[(m - 1)/2] &`
`& IntegerQ[(n - 1)/2]`

Maple [A] (verified)

Time = 134.34 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.88

method	result
default	$-\frac{4\left(-\frac{437\sin(x)^7}{256} + \frac{3527\sin(x)^5}{1536} - \frac{3257\sin(x)^3}{3072} + \frac{331\sin(x)}{2048}\right)}{(2\sin(x)^2-1)^4} + \frac{1483\operatorname{arctanh}(\sin(x)\sqrt{2})\sqrt{2}}{1024} - \frac{1}{512(\sin(x)-1)^2} + \frac{43}{512(\sin(x)-1)}$
risch	$\frac{i(1827e^{23ix} + 3733e^{21ix} + 6115e^{19ix} + 9109e^{17ix} + 5746e^{15ix} + 2382e^{13ix} - 2382e^{11ix} - 5746e^{9ix} - 9109e^{7ix} - 6115e^{5ix} - 3733e^{3ix} - 1827e^{ix})}{1536(e^{6ix} + e^{4ix} + e^{2ix} + 1)^4}$

input `int(1/(cos(x)+cos(3*x))^5,x,method=_RETURNVERBOSE)`

output `-4*(-437/256*sin(x)^7+3527/1536*sin(x)^5-3257/3072*sin(x)^3+331/2048*sin(x))/((2*sin(x)^2-1)^4)+1483/1024*arctanh(sin(x)*2^(1/2))*2^(1/2)-1/512/(sin(x)-1)^2+43/512/(sin(x)-1)+523/512*ln(sin(x)-1)+1/512/(sin(x)+1)^2+43/512/(sin(x)+1)-523/512*ln(sin(x)+1)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 219 vs. 2(88) = 176.

Time = 0.11 (sec) , antiderivative size = 219, normalized size of antiderivative = 2.03

$$\int \frac{1}{(\cos(x) + \cos(3x))^5} dx$$

$$= \frac{4449 (16 \sqrt{2} \cos(x)^{12} - 32 \sqrt{2} \cos(x)^{10} + 24 \sqrt{2} \cos(x)^8 - 8 \sqrt{2} \cos(x)^6 + \sqrt{2} \cos(x)^4) \log\left(-\frac{2 \cos(x)^2 - 2}{2 \cos(x)}\right)}{1536(e^{6ix} + e^{4ix} + e^{2ix} + 1)^4}$$

input `integrate(1/(cos(x)+cos(3*x))^5,x, algorithm="fricas")`

output

```
1/6144*(4449*(16*sqrt(2)*cos(x)^12 - 32*sqrt(2)*cos(x)^10 + 24*sqrt(2)*cos
(x)^8 - 8*sqrt(2)*cos(x)^6 + sqrt(2)*cos(x)^4)*log(-(2*cos(x)^2 - 2*sqrt(2)
)*sin(x) - 3)/(2*cos(x)^2 - 1)) - 6276*(16*cos(x)^12 - 32*cos(x)^10 + 24*c
os(x)^8 - 8*cos(x)^6 + cos(x)^4)*log(sin(x) + 1) + 6276*(16*cos(x)^12 - 32
*cos(x)^10 + 24*cos(x)^8 - 8*cos(x)^6 + cos(x)^4)*log(-sin(x) + 1) - 4*(14
616*cos(x)^10 - 25420*cos(x)^8 + 15570*cos(x)^6 - 3677*cos(x)^4 + 162*cos(
x)^2 + 12)*sin(x))/(16*cos(x)^12 - 32*cos(x)^10 + 24*cos(x)^8 - 8*cos(x)^6
+ cos(x)^4)
```

Sympy [F]

$$\int \frac{1}{(\cos(x) + \cos(3x))^5} dx = \int \frac{1}{(\cos(x) + \cos(3x))^5} dx$$

input

```
integrate(1/(cos(x)+cos(3*x))**5,x)
```

output

```
Integral((cos(x) + cos(3*x))**(-5), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12209 vs. 2(88) = 176.

Time = 0.69 (sec) , antiderivative size = 12209, normalized size of antiderivative = 113.05

$$\int \frac{1}{(\cos(x) + \cos(3x))^5} dx = \text{Too large to display}$$

input

```
integrate(1/(cos(x)+cos(3*x))^5,x, algorithm="maxima")
```

output

```

-1/12288*(8*(1827*sin(23*x) + 3733*sin(21*x) + 6115*sin(19*x) + 9109*sin(17*x) + 5746*sin(15*x) + 2382*sin(13*x) - 2382*sin(11*x) - 5746*sin(9*x) - 9109*sin(7*x) - 6115*sin(5*x) - 3733*sin(3*x) - 1827*sin(x))*cos(24*x) - 14616*(4*sin(22*x) + 10*sin(20*x) + 20*sin(18*x) + 31*sin(16*x) + 40*sin(14*x) + 44*sin(12*x) + 40*sin(10*x) + 31*sin(8*x) + 20*sin(6*x) + 10*sin(4*x) + 4*sin(2*x))*cos(23*x) + 32*(3733*sin(21*x) + 6115*sin(19*x) + 9109*sin(17*x) + 5746*sin(15*x) + 2382*sin(13*x) - 2382*sin(11*x) - 5746*sin(9*x) - 9109*sin(7*x) - 6115*sin(5*x) - 3733*sin(3*x) - 1827*sin(x))*cos(22*x) - 29864*(10*sin(20*x) + 20*sin(18*x) + 31*sin(16*x) + 40*sin(14*x) + 44*sin(12*x) + 40*sin(10*x) + 31*sin(8*x) + 20*sin(6*x) + 10*sin(4*x) + 4*sin(2*x))*cos(21*x) + 80*(6115*sin(19*x) + 9109*sin(17*x) + 5746*sin(15*x) + 2382*sin(13*x) - 2382*sin(11*x) - 5746*sin(9*x) - 9109*sin(7*x) - 6115*sin(5*x) - 3733*sin(3*x) - 1827*sin(x))*cos(20*x) - 48920*(20*sin(18*x) + 31*sin(16*x) + 40*sin(14*x) + 44*sin(12*x) + 40*sin(10*x) + 31*sin(8*x) + 20*sin(6*x) + 10*sin(4*x) + 4*sin(2*x))*cos(19*x) + 160*(9109*sin(17*x) + 5746*sin(15*x) + 2382*sin(13*x) - 2382*sin(11*x) - 5746*sin(9*x) - 9109*sin(7*x) - 6115*sin(5*x) - 3733*sin(3*x) - 1827*sin(x))*cos(18*x) - 72872*(31*sin(16*x) + 40*sin(14*x) + 44*sin(12*x) + 40*sin(10*x) + 31*sin(8*x) + 20*sin(6*x) + 10*sin(4*x) + 4*sin(2*x))*cos(17*x) + 248*(5746*sin(15*x) + 2382*sin(13*x) - 2382*sin(11*x) - 5746*sin(9*x) - 9109*sin(7*x) - 6115*sin(5*x)...

```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.96

$$\begin{aligned}
& \int \frac{1}{(\cos(x) + \cos(3x))^5} dx \\
&= -\frac{1483}{2048} \sqrt{2} \log \left(\frac{|-2\sqrt{2} + 4 \sin(x)|}{|2\sqrt{2} + 4 \sin(x)|} \right) + \frac{43 \sin(x)^3 - 45 \sin(x)}{256 (\sin(x)^2 - 1)^2} \\
&+ \frac{10488 \sin(x)^7 - 14108 \sin(x)^5 + 6514 \sin(x)^3 - 993 \sin(x)}{1536 (2 \sin(x)^2 - 1)^4} \\
&- \frac{523}{512} \log(\sin(x) + 1) + \frac{523}{512} \log(-\sin(x) + 1)
\end{aligned}$$

input

```
integrate(1/(cos(x)+cos(3*x))^5,x, algorithm="giac")
```

output

```
-1483/2048*sqrt(2)*log(abs(-2*sqrt(2) + 4*sin(x))/abs(2*sqrt(2) + 4*sin(x)
)) + 1/256*(43*sin(x)^3 - 45*sin(x))/(sin(x)^2 - 1)^2 + 1/1536*(10488*sin(
x)^7 - 14108*sin(x)^5 + 6514*sin(x)^3 - 993*sin(x))/(2*sin(x)^2 - 1)^4 - 5
23/512*log(sin(x) + 1) + 523/512*log(-sin(x) + 1)
```

Mupad [B] (verification not implemented)

Time = 0.70 (sec) , antiderivative size = 307, normalized size of antiderivative = 2.84

$$\int \frac{1}{(\cos(x) + \cos(3x))^5} dx =$$

$$\frac{11492 \sin(3x) + 18218 \sin(5x) + 12230 \sin(7x) + 7466 \sin(9x) + 3654 \sin(11x) + 276144 \operatorname{atanh}(\sin(x/2)/\cos(x/2))}{(122880 \cos(2x) + 95232 \cos(4x) + 61440 \cos(6x) + 30720 \cos(8x) + 12288 \cos(10x) + 3072 \cos(12x) + 67584)}$$

input

```
int(1/(cos(3*x) + cos(x))^5,x)
```

output

```
-(11492*sin(3*x) + 18218*sin(5*x) + 12230*sin(7*x) + 7466*sin(9*x) + 3654*
sin(11*x) + 276144*atanh(sin(x/2)/cos(x/2)) + 4764*sin(x) + 502080*cos(2*x
)*atanh(sin(x/2)/cos(x/2)) + 389112*cos(4*x)*atanh(sin(x/2)/cos(x/2)) + 25
1040*cos(6*x)*atanh(sin(x/2)/cos(x/2)) + 125520*cos(8*x)*atanh(sin(x/2)/co
s(x/2)) + 50208*cos(10*x)*atanh(sin(x/2)/cos(x/2)) + 12552*cos(12*x)*atanh
(sin(x/2)/cos(x/2)) - 97878*2^(1/2)*atanh(2^(1/2)*sin(x)) - 177960*2^(1/2)
*atanh(2^(1/2)*sin(x))*cos(2*x) - 137919*2^(1/2)*atanh(2^(1/2)*sin(x))*cos
(4*x) - 88980*2^(1/2)*atanh(2^(1/2)*sin(x))*cos(6*x) - 44490*2^(1/2)*atanh
(2^(1/2)*sin(x))*cos(8*x) - 17796*2^(1/2)*atanh(2^(1/2)*sin(x))*cos(10*x)
- 4449*2^(1/2)*atanh(2^(1/2)*sin(x))*cos(12*x))/(122880*cos(2*x) + 95232*c
os(4*x) + 61440*cos(6*x) + 30720*cos(8*x) + 12288*cos(10*x) + 3072*cos(12*
x) + 67584)
```

Reduce [F]

$$\int \frac{1}{(\cos(x) + \cos(3x))^5} dx$$

$$= \int \frac{1}{\cos(3x)^5 + 5 \cos(3x)^4 \cos(x) + 10 \cos(3x)^3 \cos(x)^2 + 10 \cos(3x)^2 \cos(x)^3 + 5 \cos(3x) \cos(x)^4 + \cos(x)^5} dx$$

input `int(1/(cos(x)+cos(3*x))^5,x)`

output `int(1/(cos(3*x)**5 + 5*cos(3*x)**4*cos(x) + 10*cos(3*x)**3*cos(x)**2 + 10*cos(3*x)**2*cos(x)**3 + 5*cos(3*x)*cos(x)**4 + cos(x)**5),x)`

3.22 $\int \frac{1}{(1+\cos(x)+\sin(x))^2} dx$

Optimal result	182
Mathematica [A] (verified)	182
Rubi [A] (verified)	183
Maple [A] (verified)	184
Fricas [A] (verification not implemented)	185
Sympy [B] (verification not implemented)	185
Maxima [A] (verification not implemented)	186
Giac [A] (verification not implemented)	186
Mupad [B] (verification not implemented)	186
Reduce [B] (verification not implemented)	187

Optimal result

Integrand size = 8, antiderivative size = 29

$$\int \frac{1}{(1 + \cos(x) + \sin(x))^2} dx = -\log\left(1 + \tan\left(\frac{x}{2}\right)\right) - \frac{\cos(x) - \sin(x)}{1 + \cos(x) + \sin(x)}$$

output `-ln(1+tan(1/2*x))+(-cos(x)+sin(x))/(1+cos(x)+sin(x))`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.93

$$\int \frac{1}{(1 + \cos(x) + \sin(x))^2} dx = \log\left(\cos\left(\frac{x}{2}\right)\right) - \log\left(\cos\left(\frac{x}{2}\right) + \sin\left(\frac{x}{2}\right)\right) + \frac{\sin\left(\frac{x}{2}\right)}{\cos\left(\frac{x}{2}\right) + \sin\left(\frac{x}{2}\right)} + \frac{1}{2} \tan\left(\frac{x}{2}\right)$$

input `Integrate[(1 + Cos[x] + Sin[x])^(-2), x]`

output `Log[Cos[x/2]] - Log[Cos[x/2] + Sin[x/2]] + Sin[x/2]/(Cos[x/2] + Sin[x/2]) + Tan[x/2]/2`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3608, 25, 3042, 3603, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(\sin(x) + \cos(x) + 1)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(\sin(x) + \cos(x) + 1)^2} dx \\
 & \quad \downarrow \text{3608} \\
 & \int -\frac{1}{\cos(x) + \sin(x) + 1} dx - \frac{\cos(x) - \sin(x)}{\sin(x) + \cos(x) + 1} \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{1}{\cos(x) + \sin(x) + 1} dx - \frac{\cos(x) - \sin(x)}{\sin(x) + \cos(x) + 1} \\
 & \quad \downarrow \text{3042} \\
 & -\int \frac{1}{\cos(x) + \sin(x) + 1} dx - \frac{\cos(x) - \sin(x)}{\sin(x) + \cos(x) + 1} \\
 & \quad \downarrow \text{3603} \\
 & -2 \int \frac{1}{2 \tan\left(\frac{x}{2}\right) + 2} d \tan\left(\frac{x}{2}\right) - \frac{\cos(x) - \sin(x)}{\sin(x) + \cos(x) + 1} \\
 & \quad \downarrow \text{16} \\
 & -\log\left(\tan\left(\frac{x}{2}\right) + 1\right) - \frac{\cos(x) - \sin(x)}{\sin(x) + \cos(x) + 1}
 \end{aligned}$$

input `Int[(1 + Cos[x] + Sin[x])^(-2), x]`

output `-Log[1 + Tan[x/2]] - (Cos[x] - Sin[x])/(1 + Cos[x] + Sin[x])`

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ ; FreeQ}[\{a, b, c\}, x]$
- rule 25 $\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[Fx, x], x]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$
- rule 3603 $\text{Int}[(\cos[(d_)+(e_)*(x_)]*(b_)+(a_)+(c_)*\sin[(d_)+(e_)*(x_)]^{(-1)}, x_Symbol] \rightarrow \text{Module}[\{f = \text{FreeFactors}[\text{Tan}[(d + e*x)/2], x]\}, \text{Simp}[2*(f/e) \text{ Subst}[\text{Int}[1/(a + b + 2*c*f*x + (a - b)*f^2*x^2), x], x, \text{Tan}[(d + e*x)/2]/f], x]] \text{ ; FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2 - c^2, 0]$
- rule 3608 $\text{Int}[(\cos[(d_)+(e_)*(x_)]*(b_)+(a_)+(c_)*\sin[(d_)+(e_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[((-c)*\text{Cos}[d + e*x] + b*\text{Sin}[d + e*x])*((a + b*\text{Cos}[d + e*x] + c*\text{Sin}[d + e*x])^{(n + 1)}/(e*(n + 1)*(a^2 - b^2 - c^2))), x] + \text{Simp}[1/((n + 1)*(a^2 - b^2 - c^2)) \text{ Int}[(a*(n + 1) - b*(n + 2)*\text{Cos}[d + e*x] - c*(n + 2)*\text{Sin}[d + e*x])*(a + b*\text{Cos}[d + e*x] + c*\text{Sin}[d + e*x])^{(n + 1)}, x], x] \text{ ; FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2 - c^2, 0] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{NeQ}[n, -3/2]$

Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

method	result	size
default	$\frac{\tan(\frac{x}{2})}{2} - \frac{1}{1+\tan(\frac{x}{2})} - \ln(1 + \tan(\frac{x}{2}))$	27
norman	$\frac{\tan(\frac{x}{2})^2}{2} - \frac{3}{1+\tan(\frac{x}{2})} - \ln(1 + \tan(\frac{x}{2}))$	30
parallelrisch	$\frac{(-2-2\tan(\frac{x}{2}))\ln(1+\tan(\frac{x}{2}))+\tan(\frac{x}{2})^2-3}{2+2\tan(\frac{x}{2})}$	36
risch	$\frac{(-1+i)(e^{ix}+1+i)}{e^{2ix}+i+e^{ix}+ie^{ix}} + \ln(1 + e^{ix}) - \ln(e^{ix} + i)$	57

input `int(1/(1+cos(x)+sin(x))^2,x,method=_RETURNVERBOSE)`

output `1/2*tan(1/2*x)-1/(1+tan(1/2*x))-ln(1+tan(1/2*x))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.59

$$\int \frac{1}{(1 + \cos(x) + \sin(x))^2} dx$$

$$= \frac{(\cos(x) + \sin(x) + 1) \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) - (\cos(x) + \sin(x) + 1) \log(\sin(x) + 1) - 2 \cos(x) + 2 \sin(x)}{2(\cos(x) + \sin(x) + 1)}$$

input `integrate(1/(1+cos(x)+sin(x))^2,x, algorithm="fricas")`

output `1/2*((cos(x) + sin(x) + 1)*log(1/2*cos(x) + 1/2) - (cos(x) + sin(x) + 1)*log(sin(x) + 1) - 2*cos(x) + 2*sin(x))/(cos(x) + sin(x) + 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 66 vs. $2(22) = 44$.

Time = 0.30 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.28

$$\int \frac{1}{(1 + \cos(x) + \sin(x))^2} dx = -\frac{2 \log\left(\tan\left(\frac{x}{2}\right) + 1\right) \tan\left(\frac{x}{2}\right)}{2 \tan\left(\frac{x}{2}\right) + 2} - \frac{2 \log\left(\tan\left(\frac{x}{2}\right) + 1\right)}{2 \tan\left(\frac{x}{2}\right) + 2}$$

$$+ \frac{\tan^2\left(\frac{x}{2}\right)}{2 \tan\left(\frac{x}{2}\right) + 2} - \frac{3}{2 \tan\left(\frac{x}{2}\right) + 2}$$

input `integrate(1/(1+cos(x)+sin(x))**2,x)`

output `-2*log(tan(x/2) + 1)*tan(x/2)/(2*tan(x/2) + 2) - 2*log(tan(x/2) + 1)/(2*tan(x/2) + 2) + tan(x/2)**2/(2*tan(x/2) + 2) - 3/(2*tan(x/2) + 2)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.38

$$\int \frac{1}{(1 + \cos(x) + \sin(x))^2} dx = \frac{\sin(x)}{2(\cos(x) + 1)} - \frac{1}{\frac{\sin(x)}{\cos(x)+1} + 1} - \log\left(\frac{\sin(x)}{\cos(x) + 1} + 1\right)$$

input `integrate(1/(1+cos(x)+sin(x))^2,x, algorithm="maxima")`output `1/2*sin(x)/(cos(x) + 1) - 1/(sin(x)/(cos(x) + 1) + 1) - log(sin(x)/(cos(x) + 1) + 1)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.03

$$\int \frac{1}{(1 + \cos(x) + \sin(x))^2} dx = \frac{\tan\left(\frac{1}{2}x\right)}{\tan\left(\frac{1}{2}x\right) + 1} - \log\left(\left|\tan\left(\frac{1}{2}x\right) + 1\right|\right) + \frac{1}{2}\tan\left(\frac{1}{2}x\right)$$

input `integrate(1/(1+cos(x)+sin(x))^2,x, algorithm="giac")`output `tan(1/2*x)/(tan(1/2*x) + 1) - log(abs(tan(1/2*x) + 1)) + 1/2*tan(1/2*x)`**Mupad [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.90

$$\int \frac{1}{(1 + \cos(x) + \sin(x))^2} dx = \frac{\tan\left(\frac{x}{2}\right)}{2} - \ln\left(\tan\left(\frac{x}{2}\right) + 1\right) - \frac{1}{\tan\left(\frac{x}{2}\right) + 1}$$

input `int(1/(cos(x) + sin(x) + 1)^2,x)`output `tan(x/2)/2 - log(tan(x/2) + 1) - 1/(tan(x/2) + 1)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.59

$$\int \frac{1}{(1 + \cos(x) + \sin(x))^2} dx$$

$$= \frac{-2 \log(\tan(\frac{x}{2}) + 1) \tan(\frac{x}{2}) - 2 \log(\tan(\frac{x}{2}) + 1) + \tan(\frac{x}{2})^2 + 3 \tan(\frac{x}{2})}{2 \tan(\frac{x}{2}) + 2}$$

input

```
int(1/(1+cos(x)+sin(x))^2,x)
```

output

```
( - 2*log(tan(x/2) + 1)*tan(x/2) - 2*log(tan(x/2) + 1) + tan(x/2)**2 + 3*tan(x/2))/(2*(tan(x/2) + 1))
```

3.23 $\int \sqrt{1 + \tanh(4x)} dx$

Optimal result	188
Mathematica [A] (verified)	188
Rubi [A] (verified)	189
Maple [A] (verified)	190
Fricas [B] (verification not implemented)	190
Sympy [F]	191
Maxima [B] (verification not implemented)	191
Giac [A] (verification not implemented)	192
Mupad [B] (verification not implemented)	192
Reduce [F]	192

Optimal result

Integrand size = 10, antiderivative size = 26

$$\int \sqrt{1 + \tanh(4x)} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(4x)}}{\sqrt{2}}\right)}{2\sqrt{2}}$$

output `1/4*arctanh(1/2*(1+tanh(4*x))^(1/2)*2^(1/2))*2^(1/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \sqrt{1 + \tanh(4x)} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(4x)}}{\sqrt{2}}\right)}{2\sqrt{2}}$$

input `Integrate[Sqrt[1 + Tanh[4*x]], x]`

output `ArcTanh[Sqrt[1 + Tanh[4*x]]/Sqrt[2]]/(2*Sqrt[2])`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{\tanh(4x) + 1} dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{1 - i \tan(4ix)} dx \\ & \quad \downarrow \text{3961} \\ & \frac{1}{2} \int \frac{1}{1 - \tanh(4x)} d\sqrt{\tanh(4x) + 1} \\ & \quad \downarrow \text{219} \\ & \frac{\operatorname{arctanh}\left(\frac{\sqrt{\tanh(4x)+1}}{\sqrt{2}}\right)}{2\sqrt{2}} \end{aligned}$$

input `Int[Sqrt[1 + Tanh[4*x]], x]`

output `ArcTanh[Sqrt[1 + Tanh[4*x]]/Sqrt[2]]/(2*Sqrt[2])`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3961

```
Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d)
  Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]], x] /; FreeQ[{a
, b, c, d}, x] && EqQ[a^2 + b^2, 0]
```

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.77

method	result	size
derivativedivides	$\frac{\operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(4x)}\sqrt{2}}{2}\right)\sqrt{2}}{4}$	20
default	$\frac{\operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(4x)}\sqrt{2}}{2}\right)\sqrt{2}}{4}$	20

input

```
int((1+tanh(4*x))^(1/2),x,method=_RETURNVERBOSE)
```

output

```
1/4*arctanh(1/2*(1+tanh(4*x))^(1/2)*2^(1/2))*2^(1/2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 129 vs. 2(19) = 38.

Time = 0.07 (sec) , antiderivative size = 129, normalized size of antiderivative = 4.96

$$\int \sqrt{1 + \tanh(4x)} dx = \frac{1}{8} \sqrt{2} \log \left(\frac{-2 \cosh(4x)^2 - 4 \cosh(4x) \sinh(4x) - 2 \sinh(4x)^2 + \sqrt{2}(\sqrt{2} \cosh(4x)^3 + 3\sqrt{2} \cosh(4x) \sinh(4x)^2 + \sqrt{2} \sinh(4x)^3 + (3\sqrt{2} \cosh(4x)^2 + \sqrt{2}) \sinh(4x) - 1}{\sqrt{\cosh(4x)^2 + 2 \cosh(4x) \sinh(4x) + \sinh(4x)^2 + 1}} \right)$$

input

```
integrate((1+tanh(4*x))^(1/2),x, algorithm="fricas")
```

output

```
1/8*sqrt(2)*log(-2*cosh(4*x)^2 - 4*cosh(4*x)*sinh(4*x) - 2*sinh(4*x)^2 - s
qrt(2)*(sqrt(2)*cosh(4*x)^3 + 3*sqrt(2)*cosh(4*x)*sinh(4*x)^2 + sqrt(2)*si
nh(4*x)^3 + (3*sqrt(2)*cosh(4*x)^2 + sqrt(2))*sinh(4*x) + sqrt(2)*cosh(4*x
))/sqrt(cosh(4*x)^2 + 2*cosh(4*x)*sinh(4*x) + sinh(4*x)^2 + 1) - 1)
```

Sympy [F]

$$\int \sqrt{1 + \tanh(4x)} dx = \int \sqrt{\tanh(4x) + 1} dx$$

input

```
integrate((1+tanh(4*x))**(1/2),x)
```

output

```
Integral(sqrt(tanh(4*x) + 1), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 43 vs. 2(19) = 38.

Time = 0.11 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.65

$$\int \sqrt{1 + \tanh(4x)} dx = -\frac{1}{8} \sqrt{2} \log \left(-\frac{\sqrt{2} - \frac{\sqrt{2}}{\sqrt{e^{(-8x)}+1}}}{\sqrt{2} + \frac{\sqrt{2}}{\sqrt{e^{(-8x)}+1}}} \right)$$

input

```
integrate((1+tanh(4*x))^(1/2),x, algorithm="maxima")
```

output

```
-1/8*sqrt(2)*log(-(sqrt(2) - sqrt(2)/sqrt(e^(-8*x) + 1))/(sqrt(2) + sqrt(2)
)/sqrt(e^(-8*x) + 1)))
```


Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.81

$$\int \sqrt{1 + \tanh(4x)} dx = -\frac{1}{4} \sqrt{2} \log \left(\sqrt{e^{(8x)} + 1} - e^{(4x)} \right)$$

input `integrate((1+tanh(4*x))^(1/2),x, algorithm="giac")`output `-1/4*sqrt(2)*log(sqrt(e^(8*x) + 1) - e^(4*x))`**Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.73

$$\int \sqrt{1 + \tanh(4x)} dx = \frac{\sqrt{2} \operatorname{atanh} \left(\frac{\sqrt{2} \sqrt{\tanh(4x)+1}}{2} \right)}{4}$$

input `int((tanh(4*x) + 1)^(1/2),x)`output `(2^(1/2)*atanh((2^(1/2)*(tanh(4*x) + 1)^(1/2))/2))/4`**Reduce [F]**

$$\int \sqrt{1 + \tanh(4x)} dx = \int \sqrt{\tanh(4x) + 1} dx$$

input `int((1+tanh(4*x))^(1/2),x)`output `int(sqrt(tanh(4*x) + 1),x)`

3.24 $\int \frac{\tanh(x)}{\sqrt{e^x + e^{2x}}} dx$

Optimal result	193
Mathematica [A] (verified)	193
Rubi [A] (warning: unable to verify)	194
Maple [B] (verified)	196
Fricas [B] (verification not implemented)	197
Sympy [F]	198
Maxima [F]	199
Giac [B] (verification not implemented)	199
Mupad [F(-1)]	200
Reduce [F]	200

Optimal result

Integrand size = 16, antiderivative size = 110

$$\int \frac{\tanh(x)}{\sqrt{e^x + e^{2x}}} dx = 2e^{-x}\sqrt{e^x + e^{2x}} - \frac{\arctan\left(\frac{i-(1-2i)e^x}{2\sqrt{1+i}\sqrt{e^x + e^{2x}}}\right)}{\sqrt{1+i}} + \frac{\arctan\left(\frac{i+(1+2i)e^x}{2\sqrt{1-i}\sqrt{e^x + e^{2x}}}\right)}{\sqrt{1-i}}$$

output

```
arctan(1/2*(I+(1+2*I)*exp(x))/(1-I)^(1/2)/(exp(x)+exp(2*x))^(1/2))/(1-I)^(1/2)-arctan(1/2*(I+(-1+2*I)*exp(x))/(1+I)^(1/2)/(exp(x)+exp(2*x))^(1/2))/(1+I)^(1/2)+2*(exp(x)+exp(2*x))^(1/2)/exp(x)
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.10

$$\int \frac{\tanh(x)}{\sqrt{e^x + e^{2x}}} dx = \frac{2 + 2e^x - (1 - i)^{3/2}e^{x/2}\sqrt{1 + e^x}\operatorname{arctanh}\left(\frac{\sqrt{1-ie^{x/2}}}{\sqrt{1+e^x}}\right) - (1 + i)^{3/2}e^{x/2}\sqrt{1 + e^x}\operatorname{arctanh}\left(\frac{\sqrt{1+ie^{x/2}}}{\sqrt{1+e^x}}\right)}{\sqrt{e^x(1 + e^x)}}$$

input

```
Integrate[Tanh[x]/Sqrt[E^x + E^(2*x)], x]
```

output

$$(2 + 2E^x - (1 - I)^{(3/2)} * E^{(x/2)} * \text{Sqrt}[1 + E^x] * \text{ArcTanh}[(\text{Sqrt}[1 - I] * E^{(x/2)}) / \text{Sqrt}[1 + E^x]] - (1 + I)^{(3/2)} * E^{(x/2)} * \text{Sqrt}[1 + E^x] * \text{ArcTanh}[(\text{Sqrt}[1 + I] * E^{(x/2)}) / \text{Sqrt}[1 + E^x]]) / \text{Sqrt}[E^x * (1 + E^x)]$$
Rubi [A] (warning: unable to verify)

Time = 0.55 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.15, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {2720, 25, 2467, 2003, 2035, 2247, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tanh(x)}{\sqrt{e^x + e^{2x}}} dx \\ & \quad \downarrow \text{2720} \\ & \int -\frac{e^{-x}(1 - e^{2x})}{(e^{2x} + 1)\sqrt{e^x + e^{2x}}} de^x \\ & \quad \downarrow \text{25} \\ & -\int \frac{e^{-x}(1 - e^{2x})}{(1 + e^{2x})\sqrt{e^x + e^{2x}}} de^x \\ & \quad \downarrow \text{2467} \\ & -\frac{\sqrt{e^x}\sqrt{e^x + 1} \int \frac{1 - e^{2x}}{(e^x)^{3/2}\sqrt{1 + e^x}(1 + e^{2x})} de^x}{\sqrt{e^x + e^{2x}}} \\ & \quad \downarrow \text{2003} \\ & -\frac{\sqrt{e^x}\sqrt{e^x + 1} \int \frac{(1 - e^x)\sqrt{1 + e^x}}{(e^x)^{3/2}(1 + e^{2x})} de^x}{\sqrt{e^x + e^{2x}}} \\ & \quad \downarrow \text{2035} \\ & -\frac{2\sqrt{e^x}\sqrt{e^x + 1} \int \frac{e^{-2x}(1 - e^{2x})\sqrt{1 + e^{2x}}}{1 + e^{4x}} d\sqrt{e^x}}{\sqrt{e^x + e^{2x}}} \\ & \quad \downarrow \text{2247} \end{aligned}$$

$$\frac{2\sqrt{e^x}\sqrt{e^x+1} \int \left(\frac{\sqrt{1+e^{2x}}(-1-e^{2x})}{1+e^{4x}} + e^{-2x}\sqrt{1+e^{2x}} \right) d\sqrt{e^x}}{\sqrt{e^x+e^{2x}}}$$

↓ 2009

$$\frac{2\sqrt{e^x}\sqrt{e^x+1} \left(\frac{1}{2}(1-i)^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{1-i}\sqrt{e^x}}{\sqrt{e^{2x}+1}} \right) + \frac{1}{2}(1+i)^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{1+i}\sqrt{e^x}}{\sqrt{e^{2x}+1}} \right) - e^{-x}\sqrt{e^{2x}+1} \right)}{\sqrt{e^x+e^{2x}}}$$

input `Int [Tanh [x] / Sqrt [E^x + E^(2*x)], x]`

output `(-2*Sqrt [E^x] * Sqrt [1 + E^x] * (-Sqrt [1 + E^(2*x)] / E^x) + ((1 - I)^(3/2) * ArcTanh [(Sqrt [1 - I] * Sqrt [E^x]) / Sqrt [1 + E^(2*x)])] / 2 + ((1 + I)^(3/2) * ArcTanh [(Sqrt [1 + I] * Sqrt [E^x]) / Sqrt [1 + E^(2*x)])] / 2) / Sqrt [E^x + E^(2*x)]`

Defintions of rubi rules used

rule 25 `Int [- (Fx_), x_Symbol] := Simp [Identity [-1] Int [Fx, x], x]`

rule 2003 `Int [(u_) * ((c_) + (d_) * (x_))^(n_) * ((a_) + (b_) * (x_)^2)^(p_), x_Symbol] :> Int [u * (c + d*x)^(n + p) * (a/c + (b/d)*x)^p, x] /; FreeQ[{a, b, c, d, n, p}, x] && EqQ[b*c^2 + a*d^2, 0] && (IntegerQ[p] || (GtQ[a, 0] && GtQ[c, 0] && !IntegerQ[n]))`

rule 2009 `Int [u_, x_Symbol] := Simp [IntSum [u, x], x] /; SumQ [u]`

rule 2035 `Int [(Fx_) * (x_)^(m_), x_Symbol] := With [{k = Denominator [m]}, Simp [k Subst [Int [x^(k*(m + 1) - 1) * SubstPower [Fx, x, k], x], x, x^(1/k)], x] /; FractionQ [m] && AlgebraicFunctionQ [Fx, x]`

rule 2247

```
Int[(Px_)*((f_)*(x_)^(m_))*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)
^4)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(f*x)^m*(d + e*x^2)^q*(a + c
*x^4)^p, x], x] /; FreeQ[{a, c, d, e, f, m, q}, x] && PolyQ[Px, x] && IntegerQ[p]
```

rule 2467

```
Int[(Fx_)*(Px_)^(p_), x_Symbol] := With[{r = Expon[Px, x, Min]}, Simp[Px^FracPart[p]/(x^(r*FracPart[p]))*ExpandToSum[Px/x^r, x]^FracPart[p]) Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x], x] /; IGtQ[r, 0] /; FreeQ[p, x] && PolyQ[Px, x] && !IntegerQ[p] && !MonomialQ[Px, x] && !PolyQ[Fx, x]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 365 vs. $2(81) = 162$.

Time = 0.41 (sec) , antiderivative size = 366, normalized size of antiderivative = 3.33

method	result
default	$\frac{\sqrt{2} \left(\sqrt{\tanh\left(\frac{x}{2}\right)+1} \sqrt{2\sqrt{2}+2} \sqrt{2} \sqrt{2\sqrt{2}-2} \ln\left(\tanh\left(\frac{x}{2}\right)+1-\sqrt{\tanh\left(\frac{x}{2}\right)+1} \sqrt{2\sqrt{2}+2+\sqrt{2}}\right)-\sqrt{\tanh\left(\frac{x}{2}\right)+1} \sqrt{2\sqrt{2}+2} \sqrt{2} \sqrt{2\sqrt{2}-2} \right)}{\dots}$

input

```
int(tanh(x)/(exp(x)+exp(2*x))^(1/2), x, method=_RETURNVERBOSE)
```

output

```
-1/4*2^(1/2)*((tanh(1/2*x)+1)^(1/2)*(2*2^(1/2)+2)^(1/2)*2^(1/2)*(2*2^(1/2)
-2)^(1/2)*ln(tanh(1/2*x)+1-(tanh(1/2*x)+1)^(1/2)*(2*2^(1/2)+2)^(1/2)+2^(1/
2))-(tanh(1/2*x)+1)^(1/2)*(2*2^(1/2)+2)^(1/2)*2^(1/2)*(2*2^(1/2)-2)^(1/2)*
ln(tanh(1/2*x)+1+(tanh(1/2*x)+1)^(1/2)*(2*2^(1/2)+2)^(1/2)+2^(1/2))-(tanh(
1/2*x)+1)^(1/2)*(2*2^(1/2)+2)^(1/2)*(2*2^(1/2)-2)^(1/2)*ln(tanh(1/2*x)+1-(
tanh(1/2*x)+1)^(1/2)*(2*2^(1/2)+2)^(1/2)+2^(1/2)))+(tanh(1/2*x)+1)^(1/2)*(2
*2^(1/2)+2)^(1/2)*(2*2^(1/2)-2)^(1/2)*ln(tanh(1/2*x)+1+(tanh(1/2*x)+1)^(1/
2)*(2*2^(1/2)+2)^(1/2)+2^(1/2)))+4*(tanh(1/2*x)+1)^(1/2)*arctan((2*(tanh(1/
2*x)+1)^(1/2)-(2*2^(1/2)+2)^(1/2))/(2*2^(1/2)-2)^(1/2))+4*(tanh(1/2*x)+1)^(
1/2)*arctan((2*(tanh(1/2*x)+1)^(1/2)+(2*2^(1/2)+2)^(1/2))/(2*2^(1/2)-2)^(
1/2))+8*(2*2^(1/2)-2)^(1/2))/(2*2^(1/2)-2)^(1/2)/(tanh(1/2*x)-1)/((tanh(1/
2*x)+1)/(tanh(1/2*x)-1)^2)^(1/2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 501 vs. $2(67) = 134$.

Time = 0.09 (sec) , antiderivative size = 501, normalized size of antiderivative = 4.55

$$\int \frac{\tanh(x)}{\sqrt{e^x + e^{2x}}} dx = \text{Too large to display}$$

input

```
integrate(tanh(x)/(exp(x)+exp(2*x))^(1/2),x, algorithm="fricas")
```

output

```

1/2*(2*sqrt(sqrt(2) + 1)*(cosh(x) + sinh(x))*arctan((2*(sqrt(2) + 1)*sqrt(
sqrt(2) - 1) + sqrt(2) + 2)*sqrt(sqrt(2) + 1)*sqrt((cosh(x) + sinh(x) + 1)
/(cosh(x) - sinh(x))) - ((sqrt(2) + 2)*cosh(x) + (sqrt(2) + 2)*sinh(x) + (
2*(sqrt(2) + 1)*cosh(x) + 2*(sqrt(2) + 1)*sinh(x) + sqrt(2) + 1)*sqrt(sqrt(
2) - 1) + sqrt(2))*sqrt(sqrt(2) + 1)) - 2*sqrt(sqrt(2) + 1)*(cosh(x) + si
nh(x))*arctan((2*(sqrt(2) + 1)*sqrt(sqrt(2) - 1) - sqrt(2) - 2)*sqrt(sqrt(
2) + 1)*sqrt((cosh(x) + sinh(x) + 1)/(cosh(x) - sinh(x))) + ((sqrt(2) + 2)
*cosh(x) + (sqrt(2) + 2)*sinh(x) - (2*(sqrt(2) + 1)*cosh(x) + 2*(sqrt(2) +
1)*sinh(x) + sqrt(2) + 1)*sqrt(sqrt(2) - 1) + sqrt(2))*sqrt(sqrt(2) + 1))
+ sqrt(sqrt(2) - 1)*(cosh(x) + sinh(x))*log(2*cosh(x)^2 + (4*cosh(x) + 1)
*sinh(x) + 2*sinh(x)^2 + (sqrt(2)*cosh(x) + sqrt(2)*sinh(x) - sqrt(2) - 2)
*sqrt(sqrt(2) - 1) - (sqrt(2)*sqrt(sqrt(2) - 1) + 2*cosh(x) + 2*sinh(x))*s
qrt((cosh(x) + sinh(x) + 1)/(cosh(x) - sinh(x))) + sqrt(2) + cosh(x) + 1)
- sqrt(sqrt(2) - 1)*(cosh(x) + sinh(x))*log(2*cosh(x)^2 + (4*cosh(x) + 1)*
sinh(x) + 2*sinh(x)^2 - (sqrt(2)*cosh(x) + sqrt(2)*sinh(x) - sqrt(2) - 2)*
sqrt(sqrt(2) - 1) + (sqrt(2)*sqrt(sqrt(2) - 1) - 2*cosh(x) - 2*sinh(x))*sq
rt((cosh(x) + sinh(x) + 1)/(cosh(x) - sinh(x))) + sqrt(2) + cosh(x) + 1) +
4*sqrt((cosh(x) + sinh(x) + 1)/(cosh(x) - sinh(x))) + 4*cosh(x) + 4*sinh(
x))/(cosh(x) + sinh(x))

```

Sympy [F]

$$\int \frac{\tanh(x)}{\sqrt{e^x + e^{2x}}} dx = \int \frac{\tanh(x)}{\sqrt{(e^x + 1)e^x}} dx$$

input

```
integrate(tanh(x)/(exp(x)+exp(2*x))**(1/2), x)
```

output

```
Integral(tanh(x)/sqrt((exp(x) + 1)*exp(x)), x)
```

Maxima [F]

$$\int \frac{\tanh(x)}{\sqrt{e^x + e^{2x}}} dx = \int \frac{\tanh(x)}{\sqrt{e^{(2x)} + e^x}} dx$$

input `integrate(tanh(x)/(exp(x)+exp(2*x))^(1/2),x, algorithm="maxima")`

output `integrate(tanh(x)/sqrt(e^(2*x) + e^x), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 470 vs. $2(67) = 134$.

Time = 0.24 (sec) , antiderivative size = 470, normalized size of antiderivative = 4.27

$$\begin{aligned} & \int \frac{\tanh(x)}{\sqrt{e^x + e^{2x}}} dx = \\ & -\frac{1}{2} \sqrt{\sqrt{2} - 1} \log \left(\left(65 \sqrt{2} \left(\sqrt{e^{(2x)} + e^x} - e^x \right) + 13 \sqrt{2} \sqrt{13 \sqrt{2} - 17} - 13 \sqrt{2} - 17 \sqrt{13 \sqrt{2} - 17} - 85 \right. \right. \\ & \left. \left. + \left(13 \sqrt{2} \left(\sqrt{e^{(2x)} + e^x} - e^x \right) + 65 \sqrt{2} + 7 \sqrt{13 \sqrt{2} - 17} - 17 \sqrt{e^{(2x)} + e^x} + 17 e^x - 85 \right)^2 \right) \right) \\ & + \frac{1}{2} \sqrt{\sqrt{2} - 1} \log \left(\left(65 \sqrt{2} \left(\sqrt{e^{(2x)} + e^x} - e^x \right) - 13 \sqrt{2} \sqrt{13 \sqrt{2} - 17} - 13 \sqrt{2} + 17 \sqrt{13 \sqrt{2} - 17} - 85 \right. \right. \\ & \left. \left. + \left(13 \sqrt{2} \left(\sqrt{e^{(2x)} + e^x} - e^x \right) + 65 \sqrt{2} - 7 \sqrt{13 \sqrt{2} - 17} - 17 \sqrt{e^{(2x)} + e^x} + 17 e^x - 85 \right)^2 \right) \right) \\ & + \frac{\arctan\left(\frac{1}{5}\right) + \arctan\left(\frac{1}{7} \left(11 \sqrt{2} \sqrt{13 \sqrt{2} - 17} + 14 \sqrt{2} + 16 \sqrt{13 \sqrt{2} - 17} + 14 \right) \left(\sqrt{e^{(2x)} + e^x} - e^x \right) - \sqrt{\sqrt{2} - 1}\right)}{\sqrt{\sqrt{2} - 1}} \\ & - \frac{\arctan\left(\frac{1}{5}\right) + \arctan\left(-\frac{1}{7} \left(11 \sqrt{2} \sqrt{13 \sqrt{2} - 17} - 14 \sqrt{2} + 16 \sqrt{13 \sqrt{2} - 17} - 14 \right) \left(\sqrt{e^{(2x)} + e^x} - e^x \right) - \sqrt{\sqrt{2} - 1}\right)}{\sqrt{\sqrt{2} - 1}} \\ & + \frac{2}{\sqrt{e^{(2x)} + e^x} - e^x} \end{aligned}$$

input `integrate(tanh(x)/(exp(x)+exp(2*x))^(1/2),x, algorithm="giac")`

output

```
-1/2*sqrt(sqrt(2) - 1)*log((65*sqrt(2)*(sqrt(e^(2*x) + e^x) - e^x) + 13*sqrt(2)*sqrt(13*sqrt(2) - 17) - 13*sqrt(2) - 17*sqrt(13*sqrt(2) - 17) - 85*sqrt(e^(2*x) + e^x) + 85*e^x + 17)^2 + (13*sqrt(2)*(sqrt(e^(2*x) + e^x) - e^x) + 65*sqrt(2) + 7*sqrt(13*sqrt(2) - 17) - 17*sqrt(e^(2*x) + e^x) + 17*e^x - 85)^2) + 1/2*sqrt(sqrt(2) - 1)*log((65*sqrt(2)*(sqrt(e^(2*x) + e^x) - e^x) - 13*sqrt(2)*sqrt(13*sqrt(2) - 17) - 13*sqrt(2) + 17*sqrt(13*sqrt(2) - 17) - 85*sqrt(e^(2*x) + e^x) + 85*e^x + 17)^2 + (13*sqrt(2)*(sqrt(e^(2*x) + e^x) - e^x) + 65*sqrt(2) - 7*sqrt(13*sqrt(2) - 17) - 17*sqrt(e^(2*x) + e^x) + 17*e^x - 85)^2) + (arctan(1/5) + arctan(1/7*(11*sqrt(2)*sqrt(13*sqrt(2) - 17) + 14*sqrt(2) + 16*sqrt(13*sqrt(2) - 17) + 14)*(sqrt(e^(2*x) + e^x) - e^x) - 5/7*sqrt(2)*sqrt(13*sqrt(2) - 17) - sqrt(2) - 6/7*sqrt(13*sqrt(2) - 17) - 1))/sqrt(sqrt(2) - 1) - (arctan(1/5) + arctan(-1/7*(11*sqrt(2)*sqrt(13*sqrt(2) - 17) - 14*sqrt(2) + 16*sqrt(13*sqrt(2) - 17) - 14)*(sqrt(e^(2*x) + e^x) - e^x) + 5/7*sqrt(2)*sqrt(13*sqrt(2) - 17) - sqrt(2) + 6/7*sqrt(13*sqrt(2) - 17) - 1))/sqrt(sqrt(2) - 1) + 2/(sqrt(e^(2*x) + e^x) - e^x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\tanh(x)}{\sqrt{e^x + e^{2x}}} dx = \int \frac{\tanh(x)}{\sqrt{e^{2x} + e^x}} dx$$

input

```
int(tanh(x)/(exp(2*x) + exp(x))^(1/2), x)
```

output

```
int(tanh(x)/(exp(2*x) + exp(x))^(1/2), x)
```

Reduce [F]

$$\int \frac{\tanh(x)}{\sqrt{e^x + e^{2x}}} dx = \int \frac{\sqrt{e^x + 1} \tanh(x)}{e^{\frac{3x}{2}} + e^{\frac{x}{2}}} dx$$

input

```
int(tanh(x)/(exp(x)+exp(2*x))^(1/2), x)
```

output `int((sqrt(e**x + 1)*tanh(x))/(e**((3*x)/2) + e**(x/2)),x)`

3.25 $\int \sqrt{\operatorname{sech}(x) \sinh(2x)} dx$

Optimal result	202
Mathematica [C] (verified)	202
Rubi [A] (verified)	203
Maple [A] (verified)	205
Fricas [A] (verification not implemented)	206
Sympy [F]	206
Maxima [F]	207
Giac [F]	207
Mupad [F(-1)]	207
Reduce [F]	208

Optimal result

Integrand size = 11, antiderivative size = 40

$$\int \sqrt{\operatorname{sech}(x) \sinh(2x)} dx = \frac{2i\sqrt{2}E\left(\frac{\pi}{4} - \frac{ix}{2} \mid 2\right) \sqrt{\sinh(x)}}{\sqrt{i \sinh(x)}}$$

output

```
2*I*(sin(1/4*Pi+1/2*I*x)^2)^(1/2)/sin(1/4*Pi+1/2*I*x)*EllipticE(cos(1/4*Pi
+1/2*I*x),2^(1/2))*2^(1/2)*sinh(x)^(1/2)/(I*sinh(x))^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.90 (sec) , antiderivative size = 86, normalized size of antiderivative = 2.15

$$\int \sqrt{\operatorname{sech}(x) \sinh(2x)} dx$$

$$= \frac{2}{3} \left(-3 + \operatorname{Hypergeometric2F1} \left(\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, \tanh^2 \left(\frac{x}{2} \right) \right) \sqrt{\operatorname{sech}^2 \left(\frac{x}{2} \right)} \right.$$

$$\left. + 4 \operatorname{Hypergeometric2F1} \left(\frac{3}{4}, \frac{3}{2}, \frac{7}{4}, \tanh^2 \left(\frac{x}{2} \right) \right) \sqrt{\operatorname{sech}^2 \left(\frac{x}{2} \right)} \sqrt{\operatorname{sech}(x) \sinh(2x)} \tanh \left(\frac{x}{2} \right) \right)$$

input

```
Integrate[Sqrt[Sech[x]*Sinh[2*x]],x]
```

output

```
(2*(-3 + Hypergeometric2F1[1/2, 3/4, 7/4, Tanh[x/2]^2]*Sqrt[Sech[x/2]^2] +
4*Hypergeometric2F1[3/4, 3/2, 7/4, Tanh[x/2]^2]*Sqrt[Sech[x/2]^2])*Sqrt[S
ech[x]*Sinh[2*x]]*Tanh[x/2])/3
```

Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.909$, Rules used = {3042, 4898, 3042, 4900, 3042, 4709, 3042, 4797, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\sinh(2x)\operatorname{sech}(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{-i \sin(2ix) \sec(ix)} dx \\
 & \quad \downarrow \text{4898} \\
 & \frac{\sqrt{\sinh(2x)\operatorname{sech}(x)} \int \sqrt{i \operatorname{sech}(x) \sinh(2x)} dx}{\sqrt{i \sinh(2x)\operatorname{sech}(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sinh(2x)\operatorname{sech}(x)} \int \sqrt{\sec(ix) \sin(2ix)} dx}{\sqrt{i \sinh(2x)\operatorname{sech}(x)}} \\
 & \quad \downarrow \text{4900} \\
 & \frac{\sqrt{\sinh(2x)\operatorname{sech}(x)} \int \sqrt{\operatorname{sech}(x)} \sqrt{i \sinh(2x)} dx}{\sqrt{i \sinh(2x)} \sqrt{\operatorname{sech}(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sinh(2x)\operatorname{sech}(x)} \int \sqrt{\sec(ix)} \sqrt{\sin(2ix)} dx}{\sqrt{i \sinh(2x)} \sqrt{\operatorname{sech}(x)}} \\
 & \quad \downarrow \text{4709}
 \end{aligned}$$

$$\begin{aligned}
& \frac{\sqrt{\cosh(x)}\sqrt{\sinh(2x)\operatorname{sech}(x)} \int \frac{\sqrt{i\sinh(2x)}}{\sqrt{\cosh(x)}} dx}{\sqrt{i\sinh(2x)}} \\
& \quad \downarrow \text{3042} \\
& \frac{\sqrt{\cosh(x)}\sqrt{\sinh(2x)\operatorname{sech}(x)} \int \frac{\sqrt{\sin(2ix)}}{\sqrt{\cos(ix)}} dx}{\sqrt{i\sinh(2x)}} \\
& \quad \downarrow \text{4797} \\
& \frac{\sqrt{\sinh(2x)\operatorname{sech}(x)} \int \sqrt{i\sinh(x)} dx}{\sqrt{i\sinh(x)}} \\
& \quad \downarrow \text{3042} \\
& \frac{\sqrt{\sinh(2x)\operatorname{sech}(x)} \int \sqrt{\sin(ix)} dx}{\sqrt{i\sinh(x)}} \\
& \quad \downarrow \text{3119} \\
& \frac{2iE\left(\frac{\pi}{4} - \frac{ix}{2} \mid 2\right) \sqrt{\sinh(2x)\operatorname{sech}(x)}}{\sqrt{i\sinh(x)}}
\end{aligned}$$

input `Int[Sqrt[Sech[x]*Sinh[2*x]],x]`

output `((2*I)*EllipticE[Pi/4 - (I/2)*x, 2]*Sqrt[Sech[x]*Sinh[2*x]])/Sqrt[I*Sinh[x]]`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4709 `Int[(u_)*((c_)*sec[(a_) + (b_)*(x_)])^(m_), x_Symbol] := Simp[(c*Sec[a + b*x])^m*(c*cos[a + b*x])^m Int[ActivateTrig[u]/(c*cos[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownSineIntegrandQ[u, x]`

rule 4797 `Int[(cos[(a_) + (b_)*(x_)]*(e_))^(m_)*((g_)*sin[(c_) + (d_)*(x_)])^(p_), x_Symbol] := Simp[(g*sin[c + d*x])^p/((e*cos[a + b*x])^p*sin[a + b*x]^p) Int[(e*cos[a + b*x])^(m + p)*sin[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, e, g, m, p}, x] && EqQ[b*c - a*d, 0] && EqQ[d/b, 2] && !IntegerQ[p]`

rule 4898 `Int[(u_)*((a_)*(v_))^(p_), x_Symbol] := With[{uu = ActivateTrig[u], vv = ActivateTrig[v]}, Simp[a^IntPart[p]*((a*vv)^FracPart[p]/vv^FracPart[p]) Int[uu*vv^p, x], x] /; FreeQ[{a, p}, x] && !IntegerQ[p] && !InertTrigFreeQ[v]`

rule 4900 `Int[(u_)*((v_)^(m_)*(w_)^(n_))^(p_), x_Symbol] := With[{uu = ActivateTrig[u], vv = ActivateTrig[v], ww = ActivateTrig[w]}, Simp[(vv^m*ww^n)^FracPart[p]/(vv^(m*FracPart[p])*ww^(n*FracPart[p])) Int[uu*vv^(m*p)*ww^(n*p), x], x] /; FreeQ[{m, n, p}, x] && !IntegerQ[p] && (!InertTrigFreeQ[v] || !InertTrigFreeQ[w])`

Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.88

method	result
default	$\frac{2\sqrt{-i(\sinh(x)+i)}\sqrt{-i(-\sinh(x)+i)}\sqrt{i\sinh(x)}\left(2\operatorname{EllipticE}\left(\sqrt{1-i\sinh(x)},\frac{\sqrt{2}}{2}\right)-\operatorname{EllipticF}\left(\sqrt{1-i\sinh(x)},\frac{\sqrt{2}}{2}\right)\right)}{\cosh(x)\sqrt{\sinh(x)}}$
risch	$2\sqrt{e^{-x}(-1+e^{2x})} + \frac{\left(-\frac{4(-1+e^{2x})}{\sqrt{e^x(-1+e^{2x})}} + \frac{2\sqrt{1+e^x}\sqrt{-2e^x+2}\sqrt{-e^x}\left(-2\operatorname{EllipticE}\left(\sqrt{1+e^x},\frac{\sqrt{2}}{2}\right)+\operatorname{EllipticF}\left(\sqrt{1+e^x},\frac{\sqrt{2}}{2}\right)\right)\right)}{\sqrt{e^{3x}-e^x}}\sqrt{e^{-x}}}{-1+e^{2x}}$

input `int((sinh(2*x)/cosh(x))^(1/2),x,method=_RETURNVERBOSE)`

output

```
2*(-I*(sinh(x)+I))^(1/2)*(-I*(-sinh(x)+I))^(1/2)*(I*sinh(x))^(1/2)*(2*EllipticE((1-I*sinh(x))^(1/2),1/2*2^(1/2))-EllipticF((1-I*sinh(x))^(1/2),1/2*2^(1/2)))/cosh(x)/sinh(x)^(1/2)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.58

$$\int \sqrt{\operatorname{sech}(x) \sinh(2x)} dx = -2\sqrt{2}\sqrt{\sinh(x)} - 4 \operatorname{weierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cosh(x) + \sinh(x)))$$

input

```
integrate((sinh(2*x)/cosh(x))^(1/2),x, algorithm="fricas")
```

output

```
-2*sqrt(2)*sqrt(sinh(x)) - 4*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cosh(x) + sinh(x)))
```

Sympy [F]

$$\int \sqrt{\operatorname{sech}(x) \sinh(2x)} dx = \int \sqrt{\frac{\sinh(2x)}{\cosh(x)}} dx$$

input

```
integrate((sinh(2*x)/cosh(x))**(1/2),x)
```

output

```
Integral(sqrt(sinh(2*x)/cosh(x)), x)
```

Maxima [F]

$$\int \sqrt{\operatorname{sech}(x) \sinh(2x)} dx = \int \sqrt{\frac{\sinh(2x)}{\cosh(x)}} dx$$

input `integrate((sinh(2*x)/cosh(x))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(sinh(2*x)/cosh(x)), x)`

Giac [F]

$$\int \sqrt{\operatorname{sech}(x) \sinh(2x)} dx = \int \sqrt{\frac{\sinh(2x)}{\cosh(x)}} dx$$

input `integrate((sinh(2*x)/cosh(x))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(sinh(2*x)/cosh(x)), x)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{\operatorname{sech}(x) \sinh(2x)} dx = \int \sqrt{\frac{\sinh(2x)}{\cosh(x)}} dx$$

input `int((sinh(2*x)/cosh(x))^(1/2),x)`

output `int((sinh(2*x)/cosh(x))^(1/2), x)`

Reduce [F]

$$\int \sqrt{\operatorname{sech}(x) \sinh(2x)} dx = \int \frac{\sqrt{\sinh(2x)} \sqrt{\cosh(x)}}{\cosh(x)} dx$$

input `int((sinh(2*x)/cosh(x))^(1/2),x)`

output `int((sqrt(sinh(2*x))*sqrt(cosh(x)))/cosh(x),x)`

3.26 $\int \log(x^2 + \sqrt{1-x^2}) dx$

Optimal result	209
Mathematica [C] (warning: unable to verify)	210
Rubi [A] (verified)	211
Maple [B] (verified)	212
Fricas [A] (verification not implemented)	213
Sympy [F]	214
Maxima [F]	214
Giac [B] (verification not implemented)	215
Mupad [B] (verification not implemented)	216
Reduce [B] (verification not implemented)	217

Optimal result

Integrand size = 16, antiderivative size = 185

$$\begin{aligned} \int \log(x^2 + \sqrt{1-x^2}) dx = & -2x - \arcsin(x) + \sqrt{\frac{1}{2}(1+\sqrt{5})} \arctan\left(\sqrt{\frac{2}{1+\sqrt{5}}}x\right) \\ & + \sqrt{\frac{1}{2}(1+\sqrt{5})} \arctan\left(\frac{\sqrt{\frac{1}{2}(1+\sqrt{5})}x}{\sqrt{1-x^2}}\right) \\ & + \sqrt{\frac{1}{2}(-1+\sqrt{5})} \operatorname{arctanh}\left(\sqrt{\frac{2}{-1+\sqrt{5}}}x\right) \\ & - \sqrt{\frac{1}{2}(-1+\sqrt{5})} \operatorname{arctanh}\left(\frac{\sqrt{\frac{1}{2}(-1+\sqrt{5})}x}{\sqrt{1-x^2}}\right) \\ & + x \log(x^2 + \sqrt{1-x^2}) \end{aligned}$$

output

```
-2*x-arcsin(x)+x*ln(x^2+(-x^2+1)^(1/2))+1/2*arctanh(x*2^(1/2)/(5^(1/2)-1)^(1/2))*(-2+2*5^(1/2))^(1/2)-1/2*arctanh(1/2*x*(-2+2*5^(1/2))^(1/2)/(-x^2+1)^(1/2))*(-2+2*5^(1/2))^(1/2)+1/2*arctan(x*2^(1/2)/(5^(1/2)+1)^(1/2))*(2+2*5^(1/2))^(1/2)+1/2*arctan(1/2*x*(2+2*5^(1/2))^(1/2)/(-x^2+1)^(1/2))*(2+2*5^(1/2))^(1/2)
```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.36 (sec) , antiderivative size = 920, normalized size of antiderivative = 4.97

$$\int \log(x^2 + \sqrt{1-x^2}) dx = \text{Too large to display}$$

input `Integrate[Log[x^2 + Sqrt[1 - x^2]],x]`

output

```
(-8*Sqrt[5]*x - 4*Sqrt[5]*ArcSin[x] + 5*Sqrt[2*(-1 + Sqrt[5])]*ArcTan[Sqrt[2/(1 + Sqrt[5])]*x] + Sqrt[10*(-1 + Sqrt[5])]*ArcTan[Sqrt[2/(1 + Sqrt[5])]*x] - (-5 + Sqrt[5])*Sqrt[2*(1 + Sqrt[5])]*ArcTanh[Sqrt[2/(-1 + Sqrt[5])]*x] - 5*Sqrt[2 + Sqrt[5]]*Log[-Sqrt[2*(-1 + Sqrt[5])]] + 2*x] + 3*Sqrt[5*(2 + Sqrt[5])]*Log[-Sqrt[2*(-1 + Sqrt[5])]] + 2*x] + 5*Sqrt[2 + Sqrt[5]]*Log[Sqrt[2*(-1 + Sqrt[5])]] + 2*x] - 3*Sqrt[5*(2 + Sqrt[5])]*Log[Sqrt[2*(-1 + Sqrt[5])]] + 2*x] - (5*I)*Sqrt[-2 + Sqrt[5]]*Log[(-I)*Sqrt[2*(1 + Sqrt[5])]] + 2*x] - (3*I)*Sqrt[5*(-2 + Sqrt[5])]*Log[(-I)*Sqrt[2*(1 + Sqrt[5])]] + 2*x] + (5*I)*Sqrt[-2 + Sqrt[5]]*Log[I*Sqrt[2*(1 + Sqrt[5])]] + 2*x] + (3*I)*Sqrt[5*(-2 + Sqrt[5])]*Log[I*Sqrt[2*(1 + Sqrt[5])]] + 2*x] + 4*Sqrt[5]*x*Log[x^2 + Sqrt[1 - x^2]] + (5*I)*Sqrt[-2 + Sqrt[5]]*Log[4 - (2*I)*Sqrt[2*(1 + Sqrt[5])]*x + 2*Sqrt[2*(3 + Sqrt[5])]*Sqrt[1 - x^2]] + (3*I)*Sqrt[5*(-2 + Sqrt[5])]*Log[4 - (2*I)*Sqrt[2*(1 + Sqrt[5])]*x + 2*Sqrt[2*(3 + Sqrt[5])]*Sqrt[1 - x^2]] - (5*I)*Sqrt[-2 + Sqrt[5]]*Log[4 + (2*I)*Sqrt[2*(1 + Sqrt[5])]*x + 2*Sqrt[2*(3 + Sqrt[5])]*Sqrt[1 - x^2]] - (3*I)*Sqrt[5*(-2 + Sqrt[5])]*Log[4 + (2*I)*Sqrt[2*(1 + Sqrt[5])]*x + 2*Sqrt[2*(3 + Sqrt[5])]*Sqrt[1 - x^2]] - 5*Sqrt[2 + Sqrt[5]]*Log[2*(2 + Sqrt[2*(-1 + Sqrt[5])]*x + Sqrt[2]*Sqrt[(-3 + Sqrt[5])*(-1 + x^2)])] + 3*Sqrt[5*(2 + Sqrt[5])]*Log[2*(2 + Sqrt[2*(-1 + Sqrt[5])]*x + Sqrt[2]*Sqrt[(-3 + Sqrt[5])*(-1 + x^2)])] + 5*Sqrt[2 + Sqrt[5]]*Log[4 - 2*Sqrt[2*(-1 + Sqrt[5])]*x + 2*Sqrt[2]*Sqrt[(-...
```

Rubi [A] (verified)

Time = 1.11 (sec) , antiderivative size = 349, normalized size of antiderivative = 1.89, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3028, 7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \log \left(x^2 + \sqrt{1-x^2} \right) dx \\
 & \quad \downarrow \text{3028} \\
 & x \log \left(x^2 + \sqrt{1-x^2} \right) - \int \frac{x^2 \left(2 - \frac{1}{\sqrt{1-x^2}} \right)}{x^2 + \sqrt{1-x^2}} dx \\
 & \quad \downarrow \text{7293} \\
 & x \log \left(x^2 + \sqrt{1-x^2} \right) - \int \left(\frac{2x^2}{x^2 + \sqrt{1-x^2}} - \frac{x^2}{\sqrt{1-x^2}x^2 - x^2 + 1} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & -\arcsin(x) + 2\sqrt{\frac{1}{5}(2+\sqrt{5})} \arctan \left(\frac{\sqrt{\frac{1}{2}(1+\sqrt{5})}x}{\sqrt{1-x^2}} \right) - \\
 & \sqrt{\frac{1}{10}(1+\sqrt{5})} \arctan \left(\frac{\sqrt{\frac{1}{2}(1+\sqrt{5})}x}{\sqrt{1-x^2}} \right) + 2\sqrt{\frac{1}{5}(2+\sqrt{5})} \arctan \left(\sqrt{\frac{2}{1+\sqrt{5}}}x \right) - \\
 & \sqrt{\frac{1}{10}(1+\sqrt{5})} \arctan \left(\sqrt{\frac{2}{1+\sqrt{5}}}x \right) - \sqrt{\frac{1}{10}(\sqrt{5}-1)} \operatorname{arctanh} \left(\frac{\sqrt{\frac{1}{2}(\sqrt{5}-1)}x}{\sqrt{1-x^2}} \right) - \\
 & 2\sqrt{\frac{1}{5}(\sqrt{5}-2)} \operatorname{arctanh} \left(\frac{\sqrt{\frac{1}{2}(\sqrt{5}-1)}x}{\sqrt{1-x^2}} \right) + \sqrt{\frac{1}{10}(\sqrt{5}-1)} \operatorname{arctanh} \left(\sqrt{\frac{2}{\sqrt{5}-1}}x \right) + \\
 & 2\sqrt{\frac{1}{5}(\sqrt{5}-2)} \operatorname{arctanh} \left(\sqrt{\frac{2}{\sqrt{5}-1}}x \right) + x \log \left(x^2 + \sqrt{1-x^2} \right) - 2x
 \end{aligned}$$

input `Int[Log[x^2 + Sqrt[1 - x^2]],x]`

output

```
-2*x - ArcSin[x] - Sqrt[(1 + Sqrt[5])/10]*ArcTan[Sqrt[2/(1 + Sqrt[5])]*x]
+ 2*Sqrt[(2 + Sqrt[5])/5]*ArcTan[Sqrt[2/(1 + Sqrt[5])]*x] - Sqrt[(1 + Sqrt
[5])/10]*ArcTan[(Sqrt[(1 + Sqrt[5])/2]*x)/Sqrt[1 - x^2]] + 2*Sqrt[(2 + Sqr
t[5])/5]*ArcTan[(Sqrt[(1 + Sqrt[5])/2]*x)/Sqrt[1 - x^2]] + 2*Sqrt[(-2 + Sq
rt[5])/5]*ArcTanh[Sqrt[2/(-1 + Sqrt[5])]*x] + Sqrt[(-1 + Sqrt[5])/10]*ArcT
anh[Sqrt[2/(-1 + Sqrt[5])]*x] - 2*Sqrt[(-2 + Sqrt[5])/5]*ArcTanh[(Sqrt[(-1
+ Sqrt[5])/2]*x)/Sqrt[1 - x^2]] - Sqrt[(-1 + Sqrt[5])/10]*ArcTanh[(Sqrt[(-
1 + Sqrt[5])/2]*x)/Sqrt[1 - x^2]] + x*Log[x^2 + Sqrt[1 - x^2]]
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 3028

```
Int[Log[u_], x_Symbol] := Simp[x*Log[u], x] - Int[SimplifyIntegrand[x*(D[u,
x]/u), x], x] /; InverseFunctionFreeQ[u, x]
```

rule 7293

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 468 vs. 2(138) = 276.

Time = 0.18 (sec) , antiderivative size = 469, normalized size of antiderivative = 2.54

method	result
parts	$x \ln(x^2 + \sqrt{-x^2 + 1}) - \frac{\sqrt{5}(\sqrt{5}+1) \arctan\left(\frac{2x}{\sqrt{2+2\sqrt{5}}}\right)}{5\sqrt{2+2\sqrt{5}}} + \frac{(\sqrt{5}-1)\sqrt{5} \operatorname{arctanh}\left(\frac{2x}{\sqrt{-2+2\sqrt{5}}}\right)}{5\sqrt{-2+2\sqrt{5}}} + \arcsin(x) - \frac{(\sqrt{5}-1)\sqrt{5} \operatorname{arctanh}\left(\frac{2x}{\sqrt{-2+2\sqrt{5}}}\right)}{5\sqrt{-2+2\sqrt{5}}}$
default	$x \ln(x^2 + \sqrt{-x^2 + 1}) - \frac{\sqrt{5}(\sqrt{5}+1) \arctan\left(\frac{2x}{\sqrt{2+2\sqrt{5}}}\right)}{5\sqrt{2+2\sqrt{5}}} + \frac{(\sqrt{5}-1)\sqrt{5} \operatorname{arctanh}\left(\frac{2x}{\sqrt{-2+2\sqrt{5}}}\right)}{5\sqrt{-2+2\sqrt{5}}} - 2x + \frac{2(3+\sqrt{5})\sqrt{5}}{5}$

input

```
int(ln(x^2+(-x^2+1)^(1/2)),x,method=_RETURNVERBOSE)
```

output

```

x*ln(x^2+(-x^2+1)^(1/2))-1/5*5^(1/2)*(5^(1/2)+1)/(2+2*5^(1/2))^(1/2)*arctan
n(2*x/(2+2*5^(1/2))^(1/2))+1/5*(5^(1/2)-1)*5^(1/2)/(-2+2*5^(1/2))^(1/2)*ar
ctanh(2*x/(-2+2*5^(1/2))^(1/2))+arcsin(x)-1/10*(5^(1/2)-3)*5^(1/2)/(-2+5^(
1/2))^(1/2)*arctanh(((x^2+1)^(1/2)-1)/x/(-2+5^(1/2))^(1/2))+1/10*(3+5^(1/
2))*5^(1/2)/(2+5^(1/2))^(1/2)*arctan(((x^2+1)^(1/2)-1)/x/(2+5^(1/2))^(1/2
))+1/10*(5^(1/2)-1)*5^(1/2)/(-2+5^(1/2))^(1/2)*arctan(((x^2+1)^(1/2)-1)/x
/(-2+5^(1/2))^(1/2))-1/10*5^(1/2)*(5^(1/2)+1)/(2+5^(1/2))^(1/2)*arctanh(((
-x^2+1)^(1/2)-1)/x/(2+5^(1/2))^(1/2))+2/5*(-2+5^(1/2))^(1/2)*5^(1/2)*arcta
nh(((x^2+1)^(1/2)-1)/x/(-2+5^(1/2))^(1/2))-2/5*(2+5^(1/2))^(1/2)*5^(1/2)*
arctan(((x^2+1)^(1/2)-1)/x/(2+5^(1/2))^(1/2))+4*arctan(((x^2+1)^(1/2)-1)
/x)-2/5*5^(1/2)/(-2+5^(1/2))^(1/2)*arctan(((x^2+1)^(1/2)-1)/x/(-2+5^(1/2)
)^(1/2))-2/5*5^(1/2)/(2+5^(1/2))^(1/2)*arctanh(((x^2+1)^(1/2)-1)/x/(2+5^(
1/2))^(1/2))-2*x+2/5*(3+5^(1/2))*5^(1/2)/(2+2*5^(1/2))^(1/2)*arctan(2*x/(2
+2*5^(1/2))^(1/2))-2/5*(5^(1/2)-3)*5^(1/2)/(-2+2*5^(1/2))^(1/2)*arctanh(2*
x/(-2+2*5^(1/2))^(1/2))

```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.46

$$\begin{aligned}
& \int \log(x^2 + \sqrt{1-x^2}) dx \\
&= x \log(x^2 + \sqrt{-x^2+1}) + \sqrt{\frac{1}{2}\sqrt{5} + \frac{1}{2}} \arctan\left(\frac{1}{2}(\sqrt{5}x - x)\sqrt{\frac{1}{2}\sqrt{5} + \frac{1}{2}}\right) \\
&\quad - \sqrt{\frac{1}{2}\sqrt{5} + \frac{1}{2}} \arctan\left(\frac{\sqrt{-x^2+1}(\sqrt{5}-1)\sqrt{\frac{1}{2}\sqrt{5} + \frac{1}{2}}}{2x}\right) \\
&\quad + \frac{1}{2}\sqrt{\frac{1}{2}\sqrt{5} - \frac{1}{2}} \log\left(x + \sqrt{\frac{1}{2}\sqrt{5} - \frac{1}{2}}\right) - \frac{1}{2}\sqrt{\frac{1}{2}\sqrt{5} - \frac{1}{2}} \log\left(x - \sqrt{\frac{1}{2}\sqrt{5} - \frac{1}{2}}\right) \\
&\quad + \frac{1}{2}\sqrt{\frac{1}{2}\sqrt{5} - \frac{1}{2}} \log\left(-\frac{x^2 + (\sqrt{-x^2+1}x - x)\sqrt{\frac{1}{2}\sqrt{5} - \frac{1}{2}} + \sqrt{-x^2+1} - 1}{x^2}\right) \\
&\quad - \frac{1}{2}\sqrt{\frac{1}{2}\sqrt{5} - \frac{1}{2}} \log\left(-\frac{x^2 - (\sqrt{-x^2+1}x - x)\sqrt{\frac{1}{2}\sqrt{5} - \frac{1}{2}} + \sqrt{-x^2+1} - 1}{x^2}\right) \\
&\quad - 2x + 2 \arctan\left(\frac{\sqrt{-x^2+1} - 1}{x}\right)
\end{aligned}$$

input `integrate(log(x^2+(-x^2+1)^(1/2)),x, algorithm="fricas")`

output `x*log(x^2 + sqrt(-x^2 + 1)) + sqrt(1/2*sqrt(5) + 1/2)*arctan(1/2*(sqrt(5)*
x - x)*sqrt(1/2*sqrt(5) + 1/2)) - sqrt(1/2*sqrt(5) + 1/2)*arctan(1/2*sqrt(
-x^2 + 1)*(sqrt(5) - 1)*sqrt(1/2*sqrt(5) + 1/2)/x) + 1/2*sqrt(1/2*sqrt(5)
- 1/2)*log(x + sqrt(1/2*sqrt(5) - 1/2)) - 1/2*sqrt(1/2*sqrt(5) - 1/2)*log(
x - sqrt(1/2*sqrt(5) - 1/2)) + 1/2*sqrt(1/2*sqrt(5) - 1/2)*log(-(x^2 + (sq
rt(-x^2 + 1)*x - x)*sqrt(1/2*sqrt(5) - 1/2) + sqrt(-x^2 + 1) - 1)/x^2) - 1
/2*sqrt(1/2*sqrt(5) - 1/2)*log(-(x^2 - (sqrt(-x^2 + 1)*x - x)*sqrt(1/2*sq
rt(5) - 1/2) + sqrt(-x^2 + 1) - 1)/x^2) - 2*x + 2*arctan((sqrt(-x^2 + 1) -
1)/x)`

Sympy [F]

$$\int \log(x^2 + \sqrt{1-x^2}) dx = \int \log(x^2 + \sqrt{1-x^2}) dx$$

input `integrate(ln(x**2+(-x**2+1)**(1/2)),x)`

output `Integral(log(x**2 + sqrt(1 - x**2)), x)`

Maxima [F]

$$\int \log(x^2 + \sqrt{1-x^2}) dx = \int \log(x^2 + \sqrt{-x^2+1}) dx$$

input `integrate(log(x^2+(-x^2+1)^(1/2)),x, algorithm="maxima")`

output `x*log(x^2 + sqrt(x + 1)*sqrt(-x + 1)) - x - integrate((x^4 - 2*x^2)/(x^4 -
x^2 + (x^2 - 1)*e^(1/2*log(x + 1) + 1/2*log(-x + 1))), x) + 1/2*log(x + 1
) - 1/2*log(-x + 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 301 vs. $2(138) = 276$.

Time = 0.19 (sec) , antiderivative size = 301, normalized size of antiderivative = 1.63

$$\begin{aligned}
 & \int \log(x^2 + \sqrt{1-x^2}) dx \\
 &= x \log(x^2 + \sqrt{-x^2+1}) - \frac{1}{2} \pi \operatorname{sgn}(x) + \frac{1}{2} \sqrt{2\sqrt{5}+2} \arctan\left(\frac{x}{\sqrt{\frac{1}{2}\sqrt{5}+\frac{1}{2}}}\right) \\
 &\quad - \frac{1}{2} \sqrt{2\sqrt{5}+2} \arctan\left(-\frac{\frac{x}{\sqrt{-x^2+1}-1} - \frac{\sqrt{-x^2+1}-1}{x}}{\sqrt{2\sqrt{5}+2}}\right) \\
 &\quad + \frac{1}{4} \sqrt{2\sqrt{5}-2} \log\left(\left|x + \sqrt{\frac{1}{2}\sqrt{5}-\frac{1}{2}}\right|\right) - \frac{1}{4} \sqrt{2\sqrt{5}-2} \log\left(\left|x - \sqrt{\frac{1}{2}\sqrt{5}-\frac{1}{2}}\right|\right) \\
 &\quad - \frac{1}{4} \sqrt{2\sqrt{5}-2} \log\left(\left|\sqrt{2\sqrt{5}-2} - \frac{x}{\sqrt{-x^2+1}-1} + \frac{\sqrt{-x^2+1}-1}{x}\right|\right) \\
 &\quad + \frac{1}{4} \sqrt{2\sqrt{5}-2} \log\left(\left|-\sqrt{2\sqrt{5}-2} - \frac{x}{\sqrt{-x^2+1}-1} + \frac{\sqrt{-x^2+1}-1}{x}\right|\right) \\
 &\quad - 2x - \arctan\left(-\frac{x\left(\frac{(\sqrt{-x^2+1}-1)^2}{x^2} - 1\right)}{2(\sqrt{-x^2+1}-1)}\right)
 \end{aligned}$$

input `integrate(log(x^2+(-x^2+1)^(1/2)),x, algorithm="giac")`

output `x*log(x^2 + sqrt(-x^2 + 1)) - 1/2*pi*sgn(x) + 1/2*sqrt(2*sqrt(5) + 2)*arctan(x/sqrt(1/2*sqrt(5) + 1/2)) - 1/2*sqrt(2*sqrt(5) + 2)*arctan(-(x/(sqrt(-x^2 + 1) - 1) - (sqrt(-x^2 + 1) - 1)/x)/sqrt(2*sqrt(5) + 2)) + 1/4*sqrt(2*sqrt(5) - 2)*log(abs(x + sqrt(1/2*sqrt(5) - 1/2))) - 1/4*sqrt(2*sqrt(5) - 2)*log(abs(x - sqrt(1/2*sqrt(5) - 1/2))) - 1/4*sqrt(2*sqrt(5) - 2)*log(abs(sqrt(2*sqrt(5) - 2) - x/(sqrt(-x^2 + 1) - 1) + (sqrt(-x^2 + 1) - 1)/x)) + 1/4*sqrt(2*sqrt(5) - 2)*log(abs(-sqrt(2*sqrt(5) - 2) - x/(sqrt(-x^2 + 1) - 1) + (sqrt(-x^2 + 1) - 1)/x)) - 2*x - arctan(-1/2*x*((sqrt(-x^2 + 1) - 1)^2/x^2 - 1)/(sqrt(-x^2 + 1) - 1))`

Mupad [B] (verification not implemented)

Time = 1.19 (sec) , antiderivative size = 608, normalized size of antiderivative = 3.29

$$\int \log(x^2 + \sqrt{1-x^2}) dx = \text{Too large to display}$$

input `int(log(x^2 + (1 - x^2)^(1/2)),x)`

output

```
x*log(x^2 + (1 - x^2)^(1/2)) - asin(x) - 2*x + (log(x - (2^(1/2))*(5^(1/2)
- 1)^(1/2))/2)*(5^(1/2)/2 - 5/2))/(2*(5^(1/2)/2 - 1/2)^(1/2) + 4*(5^(1/2)/
2 - 1/2)^(3/2)) - (log(x + (2^(1/2))*(5^(1/2) - 1)^(1/2))/2)*(5^(1/2)/2 - 5
/2))/(2*(5^(1/2)/2 - 1/2)^(1/2) + 4*(5^(1/2)/2 - 1/2)^(3/2)) - (log(x - (2
^(1/2))*(- 5^(1/2) - 1)^(1/2))/2)*(5^(1/2)/2 + 5/2))/(2*(- 5^(1/2)/2 - 1/2)
^(1/2) + 4*(- 5^(1/2)/2 - 1/2)^(3/2)) + (log(x + (2^(1/2))*(- 5^(1/2) - 1)^(
1/2))/2)*(5^(1/2)/2 + 5/2))/(2*(- 5^(1/2)/2 - 1/2)^(1/2) + 4*(- 5^(1/2)/2
- 1/2)^(3/2)) + (log((((x*(5^(1/2)/2 - 1/2)^(1/2) + 1)*i)/(3/2 - 5^(1/2)
/2)^(1/2) + (1 - x^2)^(1/2)*i)/(x + (5^(1/2)/2 - 1/2)^(1/2)))*((3*5^(1/2)
)/2 - 5/2))/((2*(5^(1/2)/2 - 1/2)^(1/2) + 4*(5^(1/2)/2 - 1/2)^(3/2))*((3/2
- 5^(1/2)/2)^(1/2)) - (log((((x*(- 5^(1/2)/2 - 1/2)^(1/2) + 1)*i)/(5^(1/2)
)/2 + 3/2)^(1/2) + (1 - x^2)^(1/2)*i)/(x + (- 5^(1/2)/2 - 1/2)^(1/2)))*((
3*5^(1/2))/2 + 5/2))/((2*(- 5^(1/2)/2 - 1/2)^(1/2) + 4*(- 5^(1/2)/2 - 1/2)
^(3/2))*((5^(1/2)/2 + 3/2)^(1/2)) - (log((((x*(5^(1/2)/2 - 1/2)^(1/2) - 1)*
i)/(3/2 - 5^(1/2)/2)^(1/2) - (1 - x^2)^(1/2)*i)/(x - (5^(1/2)/2 - 1/2)^(
1/2)))*((3*5^(1/2))/2 - 5/2))/((2*(5^(1/2)/2 - 1/2)^(1/2) + 4*(5^(1/2)/2 -
1/2)^(3/2))*((3/2 - 5^(1/2)/2)^(1/2)) + (log((((x*(- 5^(1/2)/2 - 1/2)^(1/2)
) - 1)*i)/(5^(1/2)/2 + 3/2)^(1/2) - (1 - x^2)^(1/2)*i)/(x - (- 5^(1/2)/2
- 1/2)^(1/2)))*((3*5^(1/2))/2 + 5/2))/((2*(- 5^(1/2)/2 - 1/2)^(1/2) + 4*(-
5^(1/2)/2 - 1/2)^(3/2))*((5^(1/2)/2 + 3/2)^(1/2)))
```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 179, normalized size of antiderivative = 0.97

$$\begin{aligned}
\int \log(x^2 + \sqrt{1-x^2}) dx = & -\operatorname{asin}(x) + \sqrt{\sqrt{5}-2}\sqrt{5} \operatorname{atan}\left(\frac{\tan\left(\frac{\operatorname{asin}(x)}{2}\right)}{\sqrt{\sqrt{5}-2}}\right) \\
& + 3\sqrt{\sqrt{5}-2} \operatorname{atan}\left(\frac{\tan\left(\frac{\operatorname{asin}(x)}{2}\right)}{\sqrt{\sqrt{5}-2}}\right) \\
& + \frac{\sqrt{\sqrt{5}+2}\sqrt{5} \log\left(-\sqrt{\sqrt{5}+2} + \tan\left(\frac{\operatorname{asin}(x)}{2}\right)\right)}{2} \\
& - \frac{\sqrt{\sqrt{5}+2}\sqrt{5} \log\left(\sqrt{\sqrt{5}+2} + \tan\left(\frac{\operatorname{asin}(x)}{2}\right)\right)}{2} \\
& - \frac{3\sqrt{\sqrt{5}+2} \log\left(-\sqrt{\sqrt{5}+2} + \tan\left(\frac{\operatorname{asin}(x)}{2}\right)\right)}{2} \\
& + \frac{3\sqrt{\sqrt{5}+2} \log\left(\sqrt{\sqrt{5}+2} + \tan\left(\frac{\operatorname{asin}(x)}{2}\right)\right)}{2} \\
& + \log\left(\frac{-\tan\left(\frac{\operatorname{asin}(x)}{2}\right)^4 + 4\tan\left(\frac{\operatorname{asin}(x)}{2}\right)^2 + 1}{\tan\left(\frac{\operatorname{asin}(x)}{2}\right)^4 + 2\tan\left(\frac{\operatorname{asin}(x)}{2}\right)^2 + 1}\right) x - 2x
\end{aligned}$$

input `int(log(x^2+(-x^2+1)^(1/2)),x)`output `(- 2*asin(x) + 2*sqrt(sqrt(5) - 2)*sqrt(5)*atan(tan(asin(x)/2)/sqrt(sqrt(5) - 2)) + 6*sqrt(sqrt(5) - 2)*atan(tan(asin(x)/2)/sqrt(sqrt(5) - 2)) + sqrt(sqrt(5) + 2)*sqrt(5)*log(- sqrt(sqrt(5) + 2) + tan(asin(x)/2)) - sqrt(sqrt(5) + 2)*sqrt(5)*log(sqrt(sqrt(5) + 2) + tan(asin(x)/2)) - 3*sqrt(sqrt(5) + 2)*log(- sqrt(sqrt(5) + 2) + tan(asin(x)/2)) + 3*sqrt(sqrt(5) + 2)*log(sqrt(sqrt(5) + 2) + tan(asin(x)/2)) + 2*log((- tan(asin(x)/2)**4 + 4*tan(asin(x)/2)**2 + 1)/(tan(asin(x)/2)**4 + 2*tan(asin(x)/2)**2 + 1))*x - 4*x)/2`

3.27 $\int \frac{\log(1+e^x)}{1+e^{2x}} dx$

Optimal result	218
Mathematica [A] (verified)	219
Rubi [A] (verified)	219
Maple [A] (verified)	221
Fricas [F]	221
Sympy [F]	221
Maxima [F]	222
Giac [F]	222
Mupad [F(-1)]	222
Reduce [F]	223

Optimal result

Integrand size = 16, antiderivative size = 102

$$\begin{aligned} \int \frac{\log(1+e^x)}{1+e^{2x}} dx = & -\frac{1}{2} \log\left(\left(\frac{1}{2} - \frac{i}{2}\right)(i - e^x)\right) \log(1+e^x) \\ & - \frac{1}{2} \log\left(\left(-\frac{1}{2} - \frac{i}{2}\right)(i + e^x)\right) \log(1+e^x) \\ & - \text{PolyLog}\left(2, -e^x\right) - \frac{1}{2} \text{PolyLog}\left(2, \left(\frac{1}{2} - \frac{i}{2}\right)(1+e^x)\right) \\ & - \frac{1}{2} \text{PolyLog}\left(2, \left(\frac{1}{2} + \frac{i}{2}\right)(1+e^x)\right) \end{aligned}$$

output

```
-1/2*ln((1/2-1/2*I)*(I-exp(x)))*ln(1+exp(x))-1/2*ln((-1/2-1/2*I)*(I+exp(x)))*ln(1+exp(x))-polylog(2,-exp(x))-1/2*polylog(2,(1/2-1/2*I)*(1+exp(x)))-1/2*polylog(2,(1/2+1/2*I)*(1+exp(x)))
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.00

$$\begin{aligned} \int \frac{\log(1+e^x)}{1+e^{2x}} dx &= -\frac{1}{2} \log\left(\left(\frac{1}{2} - \frac{i}{2}\right)(i - e^x)\right) \log(1+e^x) \\ &\quad - \frac{1}{2} \log\left(\left(-\frac{1}{2} - \frac{i}{2}\right)(i + e^x)\right) \log(1+e^x) \\ &\quad - \text{PolyLog}(2, -e^x) - \frac{1}{2} \text{PolyLog}\left(2, \left(\frac{1}{2} - \frac{i}{2}\right)(1+e^x)\right) \\ &\quad - \frac{1}{2} \text{PolyLog}\left(2, \left(\frac{1}{2} + \frac{i}{2}\right)(1+e^x)\right) \end{aligned}$$

input `Integrate[Log[1 + E^x]/(1 + E^(2*x)), x]`

output `-1/2*(Log[(1/2 - I/2)*(I - E^x)]*Log[1 + E^x]) - (Log[(-1/2 - I/2)*(I + E^x)]*Log[1 + E^x])/2 - PolyLog[2, -E^x] - PolyLog[2, (1/2 - I/2)*(1 + E^x)]/2 - PolyLog[2, (1/2 + I/2)*(1 + E^x)]/2`

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {2720, 2863, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} &\int \frac{\log(e^x + 1)}{e^{2x} + 1} dx \\ &\quad \downarrow \text{2720} \\ &\int \frac{e^{-x} \log(e^x + 1)}{e^{2x} + 1} de^x \\ &\quad \downarrow \text{2863} \\ &\int \left(e^{-x} \log(e^x + 1) - \frac{e^x \log(e^x + 1)}{e^{2x} + 1} \right) de^x \end{aligned}$$

$$\begin{aligned} & \downarrow \text{2009} \\ & -\text{PolyLog}(2, -e^x) - \frac{1}{2}\text{PolyLog}\left(2, \left(\frac{1}{2} - \frac{i}{2}\right)(1 + e^x)\right) - \frac{1}{2}\text{PolyLog}\left(2, \left(\frac{1}{2} + \frac{i}{2}\right)(1 + e^x)\right) - \\ & \frac{1}{2}\log\left(\left(\frac{1}{2} - \frac{i}{2}\right)(-e^x + i)\right)\log(e^x + 1) - \frac{1}{2}\log\left(\left(-\frac{1}{2} - \frac{i}{2}\right)(e^x + i)\right)\log(e^x + 1) \end{aligned}$$

input `Int[Log[1 + E^x]/(1 + E^(2*x)),x]`

output `-1/2*(Log[(1/2 - I/2)*(1 - E^x)]*Log[1 + E^x]) - (Log[(-1/2 - I/2)*(1 + E^x)]*Log[1 + E^x])/2 - PolyLog[2, -E^x] - PolyLog[2, (1/2 - I/2)*(1 + E^x)]/2 - PolyLog[2, (1/2 + I/2)*(1 + E^x)]/2`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :=> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2720 `Int[u_, x_Symbol] :=> With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 2863 `Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_))^(n_.)]*(b_.))^(p_.)*((h_.)*(x_))^(m_.)*((f_) + (g_.)*(x_)^(r_.))^(q_.), x_Symbol] :=> Int[ExpandIntegrand[(a + b*Log[c*(d + e*x)^n])^p, (h*x)^m*(f + g*x^r)^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n, p, q, r}, x] && IntegerQ[m] && IntegerQ[q]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.81

method	result
default	$-\frac{\ln(1+e^x) \ln\left(\frac{1}{2} - \frac{e^x}{2} + \frac{i(1+e^x)}{2}\right)}{2} - \frac{\ln(1+e^x) \ln\left(\frac{1}{2} - \frac{e^x}{2} - \frac{i(1+e^x)}{2}\right)}{2} - \frac{\operatorname{dilog}\left(\frac{1}{2} - \frac{e^x}{2} + \frac{i(1+e^x)}{2}\right)}{2} - \frac{\operatorname{dilog}\left(\frac{1}{2} - \frac{e^x}{2} - \frac{i(1+e^x)}{2}\right)}{2}$
risch	$-\frac{\ln(1+e^x) \ln\left(\frac{1}{2} - \frac{e^x}{2} + \frac{i(1+e^x)}{2}\right)}{2} - \frac{\ln(1+e^x) \ln\left(\frac{1}{2} - \frac{e^x}{2} - \frac{i(1+e^x)}{2}\right)}{2} - \frac{\operatorname{dilog}\left(\frac{1}{2} - \frac{e^x}{2} + \frac{i(1+e^x)}{2}\right)}{2} - \frac{\operatorname{dilog}\left(\frac{1}{2} - \frac{e^x}{2} - \frac{i(1+e^x)}{2}\right)}{2}$

input `int(ln(1+exp(x))/(1+exp(2*x)),x,method=_RETURNVERBOSE)`

output `-1/2*ln(1+exp(x))*ln(1/2-1/2*exp(x)+1/2*I*(1+exp(x)))-1/2*ln(1+exp(x))*ln(1/2-1/2*exp(x)-1/2*I*(1+exp(x)))-1/2*dilog(1/2-1/2*exp(x)+1/2*I*(1+exp(x)))-1/2*dilog(1/2-1/2*exp(x)-1/2*I*(1+exp(x)))-dilog(1+exp(x))`

Fricas [F]

$$\int \frac{\log(1+e^x)}{1+e^{2x}} dx = \int \frac{\log(e^x+1)}{e^{(2x)}+1} dx$$

input `integrate(log(1+exp(x))/(1+exp(2*x)),x, algorithm="fricas")`

output `integral(log(e^x + 1)/(e^(2*x) + 1), x)`

Sympy [F]

$$\int \frac{\log(1+e^x)}{1+e^{2x}} dx = \int \frac{\log(e^x+1)}{e^{2x}+1} dx$$

input `integrate(ln(1+exp(x))/(1+exp(2*x)),x)`

output `Integral(log(exp(x) + 1)/(exp(2*x) + 1), x)`

Maxima [F]

$$\int \frac{\log(1 + e^x)}{1 + e^{2x}} dx = \int \frac{\log(e^x + 1)}{e^{(2x)} + 1} dx$$

input `integrate(log(1+exp(x))/(1+exp(2*x)),x, algorithm="maxima")`

output `integrate(log(e^x + 1)/(e^(2*x) + 1), x)`

Giac [F]

$$\int \frac{\log(1 + e^x)}{1 + e^{2x}} dx = \int \frac{\log(e^x + 1)}{e^{(2x)} + 1} dx$$

input `integrate(log(1+exp(x))/(1+exp(2*x)),x, algorithm="giac")`

output `integrate(log(e^x + 1)/(e^(2*x) + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\log(1 + e^x)}{1 + e^{2x}} dx = \int \frac{\ln(e^x + 1)}{e^{2x} + 1} dx$$

input `int(log(exp(x) + 1)/(exp(2*x) + 1),x)`

output `int(log(exp(x) + 1)/(exp(2*x) + 1), x)`

Reduce [F]

$$\int \frac{\log(1 + e^x)}{1 + e^{2x}} dx = \int \frac{\log(e^x + 1)}{e^{2x} + 1} dx$$

input `int(log(1+exp(x))/(1+exp(2*x)),x)`

output `int(log(e**x + 1)/(e**(2*x) + 1),x)`

3.28 $\int \cosh(x) \log^2(1 + \cosh^2(x)) dx$

Optimal result	224
Mathematica [A] (verified)	225
Rubi [A] (verified)	225
Maple [F]	227
Fricas [F]	227
Sympy [F]	227
Maxima [F]	228
Giac [F]	228
Mupad [F(-1)]	229
Reduce [F]	229

Optimal result

Integrand size = 12, antiderivative size = 159

$$\begin{aligned} \int \cosh(x) \log^2(1 + \cosh^2(x)) dx = & -8\sqrt{2} \arctan\left(\frac{\sinh(x)}{\sqrt{2}}\right) + 4i\sqrt{2} \arctan\left(\frac{\sinh(x)}{\sqrt{2}}\right)^2 \\ & + 8\sqrt{2} \arctan\left(\frac{\sinh(x)}{\sqrt{2}}\right) \log\left(\frac{2\sqrt{2}}{\sqrt{2} + i \sinh(x)}\right) \\ & + 4\sqrt{2} \arctan\left(\frac{\sinh(x)}{\sqrt{2}}\right) \log(2 + \sinh^2(x)) \\ & + 4i\sqrt{2} \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{2}}{\sqrt{2} + i \sinh(x)}\right) \\ & + 8 \sinh(x) - 4 \log(2 + \sinh^2(x)) \sinh(x) \\ & + \log^2(2 + \sinh^2(x)) \sinh(x) \end{aligned}$$

output

```
8*sinh(x)-4*ln(2+sinh(x)^2)*sinh(x)+ln(2+sinh(x)^2)^2*sinh(x)-8*arctan(1/2
*sinh(x)*2^(1/2))*2^(1/2)+4*I*arctan(1/2*sinh(x)*2^(1/2))^2*2^(1/2)+4*arct
an(1/2*sinh(x)*2^(1/2))*ln(2+sinh(x)^2)*2^(1/2)+8*arctan(1/2*sinh(x)*2^(1/
2))*ln(2*2^(1/2)/(I*sinh(x)+2^(1/2)))*2^(1/2)+4*I*polylog(2,1-2*2^(1/2)/(I
*sinh(x)+2^(1/2)))*2^(1/2)
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.77

$$\int \cosh(x) \log^2(1 + \cosh^2(x)) dx = 4\sqrt{2} \arctan\left(\frac{\sinh(x)}{\sqrt{2}}\right) \left(-2 + i \arctan\left(\frac{\sinh(x)}{\sqrt{2}}\right)\right) + 2 \log\left(\frac{4i}{2i - \sqrt{2} \sinh(x)}\right) + \log(2 + \sinh^2(x)) + 4i\sqrt{2} \text{PolyLog}\left(2, \frac{2i + \sqrt{2} \sinh(x)}{-2i + \sqrt{2} \sinh(x)}\right) + (8 - 4 \log(2 + \sinh^2(x)) + \log^2(2 + \sinh^2(x))) \sinh(x)$$

input `Integrate[Cosh[x]*Log[1 + Cosh[x]^2],x]`

output `4*Sqrt[2]*ArcTan[Sinh[x]/Sqrt[2]]*(-2 + I*ArcTan[Sinh[x]/Sqrt[2]] + 2*Log[(4*I)/(2*I - Sqrt[2]*Sinh[x])] + Log[2 + Sinh[x]^2]) + (4*I)*Sqrt[2]*PolyLog[2, (2*I + Sqrt[2]*Sinh[x])/(-2*I + Sqrt[2]*Sinh[x])] + (8 - 4*Log[2 + Sinh[x]^2] + Log[2 + Sinh[x]^2]^2)*Sinh[x]`

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.01, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {4858, 2900, 2926, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(x) \log^2(\cosh^2(x) + 1) dx$$

↓ 4858

$$\int \log^2(\sinh^2(x) + 2) d\sinh(x)$$

↓ 2900

$$\sinh(x) \log^2(\sinh^2(x) + 2) - 4 \int \frac{\log(\sinh^2(x) + 2) \sinh^2(x)}{\sinh^2(x) + 2} d \sinh(x)$$

↓ 2926

$$\sinh(x) \log^2(\sinh^2(x) + 2) - 4 \int \left(\log(\sinh^2(x) + 2) - \frac{2 \log(\sinh^2(x) + 2)}{\sinh^2(x) + 2} \right) d \sinh(x)$$

↓ 2009

$$4 \left(-i\sqrt{2} \arctan\left(\frac{\sinh(x)}{\sqrt{2}}\right)^2 + 2\sqrt{2} \arctan\left(\frac{\sinh(x)}{\sqrt{2}}\right) - \sqrt{2} \arctan\left(\frac{\sinh(x)}{\sqrt{2}}\right) \log(\sinh^2(x) + 2) - 2\sqrt{2} \arctan\left(\frac{\sinh(x)}{\sqrt{2}}\right) \right) + \sinh(x) \log^2(\sinh^2(x) + 2) -$$

input `Int[Cosh[x]*Log[1 + Cosh[x]^2]^2,x]`

output `Log[2 + Sinh[x]^2]^2*Sinh[x] - 4*(2*Sqrt[2]*ArcTan[Sinh[x]/Sqrt[2]] - I*Sqrt[2]*ArcTan[Sinh[x]/Sqrt[2]]^2 - 2*Sqrt[2]*ArcTan[Sinh[x]/Sqrt[2]]*Log[(2*Sqrt[2])/(Sqrt[2] + I*Sinh[x])]) - Sqrt[2]*ArcTan[Sinh[x]/Sqrt[2]]*Log[2 + Sinh[x]^2] - I*Sqrt[2]*PolyLog[2, 1 - (2*Sqrt[2])/(Sqrt[2] + I*Sinh[x])]) - 2*Sinh[x] + Log[2 + Sinh[x]^2]*Sinh[x]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2900 `Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_))^(p_.)]*(b_.))^(q_), x_Symbol] := Simp[x*(a + b*Log[c*(d + e*x^n)^p])^q, x] - Simp[b*e*n*p*q Int[x^n*((a + b*Log[c*(d + e*x^n)^p])^(q - 1)/(d + e*x^n)), x], x] /; FreeQ[{a, b, c, d, e, n, p}, x] && IGtQ[q, 0] && (EqQ[q, 1] || IntegerQ[n])`

rule 2926 `Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_))^(p_.)]*(b_.))^(q_.)*(x_)^(m_.)*((f_) + (g_.)*(x_)^(s_))^(r_.), x_Symbol] := Int[ExpandIntegrand[(a + b*Log[c*(d + e*x^n)^p])^q, x^m*(f + g*x^s)^r, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p, q, r, s}, x] && IGtQ[q, 0] && IntegerQ[m] && IntegerQ[r] && IntegerQ[s]`

rule 4858

```
Int[Cosh[(c_.)*((a_.) + (b_.)*(x_))]*(u_), x_Symbol] := With[{d = FreeFactors[Sinh[c*(a + b*x)], x]}, Simp[d/(b*c) Subst[Int[SubstFor[1, Sinh[c*(a + b*x)]]/d, u, x], x], x, Sinh[c*(a + b*x)]/d, x] /; FunctionOfQ[Sinh[c*(a + b*x)]/d, u, x] /; FreeQ[{a, b, c}, x]
```

Maple [F]

$$\int \cosh(x) \ln(1 + \cosh(x)^2)^2 dx$$

input

```
int(cosh(x)*ln(1+cosh(x)^2)^2,x)
```

output

```
int(cosh(x)*ln(1+cosh(x)^2)^2,x)
```

Fricas [F]

$$\int \cosh(x) \log^2(1 + \cosh^2(x)) dx = \int \cosh(x) \log(\cosh(x)^2 + 1)^2 dx$$

input

```
integrate(cosh(x)*log(1+cosh(x)^2)^2,x, algorithm="fricas")
```

output

```
integral(cosh(x)*log(cosh(x)^2 + 1)^2, x)
```

Sympy [F]

$$\int \cosh(x) \log^2(1 + \cosh^2(x)) dx = \int \log(\cosh^2(x) + 1)^2 \cosh(x) dx$$

input

```
integrate(cosh(x)*ln(1+cosh(x)**2)**2,x)
```

output

```
Integral(log(cosh(x)**2 + 1)**2*cosh(x), x)
```

Maxima [F]

$$\int \cosh(x) \log^2(1 + \cosh^2(x)) dx = \int \cosh(x) \log(\cosh(x)^2 + 1)^2 dx$$

input `integrate(cosh(x)*log(1+cosh(x)^2)^2,x, algorithm="maxima")`

output

```
1/2*(e^(2*x) - 1)*e^(-x)*log(e^(4*x) + 6*e^(2*x) + 1)^2 - 2*(e^(-x) + inte
grate((e^(2*x) + 6)*e^x/(e^(4*x) + 6*e^(2*x) + 1), x))*log(2)^2 + 2*(e^x -
integrate((6*e^(2*x) + 1)*e^x/(e^(4*x) + 6*e^(2*x) + 1), x))*log(2)^2 + 1
4*integrate(e^(3*x)/(e^(4*x) + 6*e^(2*x) + 1), x)*log(2)^2 + 14*integrate(
e^x/(e^(4*x) + 6*e^(2*x) + 1), x)*log(2)^2 + 4*integrate(x*e^(6*x)/(e^(5*x
) + 6*e^(3*x) + e^x), x)*log(2) + 28*integrate(x*e^(4*x)/(e^(5*x) + 6*e^(3
*x) + e^x), x)*log(2) + 28*integrate(x*e^(2*x)/(e^(5*x) + 6*e^(3*x) + e^x
), x)*log(2) - 2*integrate(e^(6*x)*log(e^(4*x) + 6*e^(2*x) + 1)/(e^(5*x) +
6*e^(3*x) + e^x), x)*log(2) - 14*integrate(e^(4*x)*log(e^(4*x) + 6*e^(2*x)
+ 1)/(e^(5*x) + 6*e^(3*x) + e^x), x)*log(2) - 14*integrate(e^(2*x)*log(e^
(4*x) + 6*e^(2*x) + 1)/(e^(5*x) + 6*e^(3*x) + e^x), x)*log(2) + 4*integrat
e(x/(e^(5*x) + 6*e^(3*x) + e^x), x)*log(2) - 2*integrate(log(e^(4*x) + 6*e
^(2*x) + 1)/(e^(5*x) + 6*e^(3*x) + e^x), x)*log(2) + 2*integrate(x^2*e^(6*
x)/(e^(5*x) + 6*e^(3*x) + e^x), x) + 14*integrate(x^2*e^(4*x)/(e^(5*x) + 6
*e^(3*x) + e^x), x) + 14*integrate(x^2*e^(2*x)/(e^(5*x) + 6*e^(3*x) + e^x)
, x) - 2*integrate(x*e^(6*x)*log(e^(4*x) + 6*e^(2*x) + 1)/(e^(5*x) + 6*e^(
3*x) + e^x), x) - 14*integrate(x*e^(4*x)*log(e^(4*x) + 6*e^(2*x) + 1)/(e^(
5*x) + 6*e^(3*x) + e^x), x) - 14*integrate(x*e^(2*x)*log(e^(4*x) + 6*e^(2*
x) + 1)/(e^(5*x) + 6*e^(3*x) + e^x), x) + 2*integrate(x^2/(e^(5*x) + 6*e^(
3*x) + e^x), x) - 2*integrate(x*log(e^(4*x) + 6*e^(2*x) + 1)/(e^(5*x) + ...
```

Giac [F]

$$\int \cosh(x) \log^2(1 + \cosh^2(x)) dx = \int \cosh(x) \log(\cosh(x)^2 + 1)^2 dx$$

input `integrate(cosh(x)*log(1+cosh(x)^2)^2,x, algorithm="giac")`

output `integrate(cosh(x)*log(cosh(x)^2 + 1)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \cosh(x) \log^2(1 + \cosh^2(x)) dx = \int \ln(\cosh(x)^2 + 1)^2 \cosh(x) dx$$

input `int(log(cosh(x)^2 + 1)^2*cosh(x), x)`output `int(log(cosh(x)^2 + 1)^2*cosh(x), x)`**Reduce [F]**

$$\int \cosh(x) \log^2(1 + \cosh^2(x)) dx$$

$$48e^x \sqrt{2} \operatorname{atan}\left(\frac{e^x}{\sqrt{2}+1}\right) - 64e^x \operatorname{atan}\left(\frac{e^x}{\sqrt{2}+1}\right) + e^{2x} \log\left(\frac{e^{4x}+6e^{2x}+1}{4e^{2x}}\right)^2 - 4e^{2x} \log\left(\frac{e^{4x}+6e^{2x}+1}{4e^{2x}}\right) + 8e^{2x} - 24e^x \sqrt{2}$$

input `int(cosh(x)*log(1+cosh(x)^2)^2,x)`output `(48*e**x*sqrt(2)*atan(e**x/(sqrt(2) + 1)) - 64*e**x*atan(e**x/(sqrt(2) + 1)) + e**(2*x)*log((e**(4*x) + 6*e**(2*x) + 1)/(4*e**(2*x)))**2 - 4*e**(2*x)*log((e**(4*x) + 6*e**(2*x) + 1)/(4*e**(2*x))) + 8*e**(2*x) - 24*e**x*sqrt(2)*log(e**x - sqrt(2)*i + i)*i + 24*e**x*sqrt(2)*log(e**x + sqrt(2)*i - i)*i - 32*e**x*int(log((e**(4*x) + 6*e**(2*x) + 1)/(4*e**(2*x)))/(e**(5*x) + 6*e**(3*x) + e**x),x) - 160*e**x*int((e**x*log((e**(4*x) + 6*e**(2*x) + 1)/(4*e**(2*x))))/(e**(4*x) + 6*e**(2*x) + 1),x) - 32*e**x*log(e**x - sqrt(2)*i + i)*i + 32*e**x*log(e**x + sqrt(2)*i - i)*i - log((e**(4*x) + 6*e**(2*x) + 1)/(4*e**(2*x)))**2 - 28*log((e**(4*x) + 6*e**(2*x) + 1)/(4*e**(2*x))) + 56)/(2*e**x)`

3.29 $\int \cosh(x) \log^2 (\cosh^2(x) + \sinh(x)) \, dx$

Optimal result	231
Mathematica [A] (verified)	233
Rubi [A] (verified)	234
Maple [F]	236
Fricas [F]	236
Sympy [F]	236
Maxima [F]	237
Giac [F]	237
Mupad [F(-1)]	238
Reduce [F]	238

Optimal result

Integrand size = 13, antiderivative size = 395

$$\begin{aligned}
\int \cosh(x) \log^2(\cosh^2(x) + \sinh(x)) \, dx = & -4\sqrt{3} \arctan\left(\frac{1 + 2\sinh(x)}{\sqrt{3}}\right) \\
& - \frac{1}{2}(1 - i\sqrt{3}) \log^2(1 - i\sqrt{3} + 2\sinh(x)) - (1 \\
& + i\sqrt{3}) \log\left(\frac{i(1 - i\sqrt{3} + 2\sinh(x))}{2\sqrt{3}}\right) \log(1 \\
& \qquad \qquad \qquad + i\sqrt{3} + 2\sinh(x)) \\
& - \frac{1}{2}(1 + i\sqrt{3}) \log^2(1 + i\sqrt{3} + 2\sinh(x)) \\
& - (1 - i\sqrt{3}) \log(1 - i\sqrt{3} \\
& + 2\sinh(x)) \log\left(-\frac{i(1 + i\sqrt{3} + 2\sinh(x))}{2\sqrt{3}}\right) \\
& - 2 \log(1 + \sinh(x) + \sinh^2(x)) \\
& + (1 - i\sqrt{3}) \log(1 - i\sqrt{3} + 2\sinh(x)) \log(1 \\
& \qquad \qquad \qquad + \sinh(x) + \sinh^2(x)) \\
& + (1 + i\sqrt{3}) \log(1 + i\sqrt{3} + 2\sinh(x)) \log(1 \\
& \qquad \qquad \qquad + \sinh(x) + \sinh^2(x)) \\
& - (1 + i\sqrt{3}) \operatorname{PolyLog}\left(2, \right. \\
& \qquad \qquad \qquad \left. -\frac{i - \sqrt{3} + 2i\sinh(x)}{2\sqrt{3}}\right) - (1 \\
& - i\sqrt{3}) \operatorname{PolyLog}\left(2, \frac{i + \sqrt{3} + 2i\sinh(x)}{2\sqrt{3}}\right) \\
& + 8\sinh(x) \\
& - 4 \log(1 + \sinh(x) + \sinh^2(x)) \sinh(x) \\
& + \log^2(1 + \sinh(x) + \sinh^2(x)) \sinh(x)
\end{aligned}$$

output

```

-2*ln(1+sinh(x)+sinh(x)^2)+8*sinh(x)-4*ln(1+sinh(x)+sinh(x)^2)*sinh(x)+ln(
1+sinh(x)+sinh(x)^2)^2*sinh(x)+ln(1+sinh(x)+sinh(x)^2)*ln(1+2*sinh(x)-I*3^
(1/2))*(1-I*3^(1/2))-1/2*ln(1+2*sinh(x)-I*3^(1/2))^2*(1-I*3^(1/2))-ln(1+2*
sinh(x)-I*3^(1/2))*ln(-1/6*I*(1+2*sinh(x)+I*3^(1/2))*3^(1/2))*(1-I*3^(1/2)
)-polylog(2,1/6*(I+2*I*sinh(x)+3^(1/2))*3^(1/2))*(1-I*3^(1/2))+ln(1+sinh(x)
)+sinh(x)^2)*ln(1+2*sinh(x)+I*3^(1/2))*(1+I*3^(1/2))-1/2*ln(1+2*sinh(x)+I*
3^(1/2))^2*(1+I*3^(1/2))-ln(1+2*sinh(x)+I*3^(1/2))*ln(1/6*I*(1+2*sinh(x)-I
*3^(1/2))*3^(1/2))*(1+I*3^(1/2))-polylog(2,1/6*(-I-2*I*sinh(x)+3^(1/2))*3^
(1/2))*(1+I*3^(1/2))-4*arctan(1/3*(1+2*sinh(x))*3^(1/2))*3^(1/2)

```

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 389, normalized size of antiderivative = 0.98

$$\begin{aligned}
\int \cosh(x) \log^2(\cosh^2(x) + \sinh(x)) dx = & -4\sqrt{3} \arctan\left(\frac{1 + 2\sinh(x)}{\sqrt{3}}\right) + i\left(i\right. \\
& + \sqrt{3}\left.) \log\left(\frac{-i + \sqrt{3} - 2i\sinh(x)}{2\sqrt{3}}\right) \log\left(1\right. \right. \\
& \left. \left. - i\sqrt{3} + 2\sinh(x)\right)\right) \\
& + \frac{1}{2}i(i + \sqrt{3}) \log^2\left(1 - i\sqrt{3} + 2\sinh(x)\right) - \left(1\right. \\
& \left. + i\sqrt{3}\right) \log\left(\frac{i + \sqrt{3} + 2i\sinh(x)}{2\sqrt{3}}\right) \log\left(1\right. \\
& \left. + i\sqrt{3} + 2\sinh(x)\right) \\
& - \frac{1}{2}\left(1 + i\sqrt{3}\right) \log^2\left(1 + i\sqrt{3} + 2\sinh(x)\right) \\
& - 2 \log\left(1 + \sinh(x) + \sinh^2(x)\right) \\
& + \left(1 - i\sqrt{3}\right) \log\left(1 - i\sqrt{3} + 2\sinh(x)\right) \log\left(1\right. \\
& \left. + \sinh(x) + \sinh^2(x)\right) \\
& + \left(1 + i\sqrt{3}\right) \log\left(1 + i\sqrt{3} + 2\sinh(x)\right) \log\left(1\right. \\
& \left. + \sinh(x) + \sinh^2(x)\right) - \left(1\right. \\
& \left. + i\sqrt{3}\right) \text{PolyLog}\left(2, \frac{-i + \sqrt{3} - 2i\sinh(x)}{2\sqrt{3}}\right) \\
& + i\left(i\right. \\
& \left. + \sqrt{3}\right) \text{PolyLog}\left(2, \frac{i + \sqrt{3} + 2i\sinh(x)}{2\sqrt{3}}\right) \\
& + 8\sinh(x) \\
& - 4 \log\left(1 + \sinh(x) + \sinh^2(x)\right) \sinh(x) \\
& + \log^2\left(1 + \sinh(x) + \sinh^2(x)\right) \sinh(x)
\end{aligned}$$

input

Integrate[Cosh[x]*Log[Cosh[x]^2 + Sinh[x]]^2,x]

output

```

-4*Sqrt[3]*ArcTan[(1 + 2*Sinh[x])/Sqrt[3]] + I*(I + Sqrt[3])*Log[(-I + Sqr
t[3] - (2*I)*Sinh[x])/(2*Sqrt[3])] *Log[1 - I*Sqrt[3] + 2*Sinh[x]] + (I/2)*
(I + Sqrt[3])*Log[1 - I*Sqrt[3] + 2*Sinh[x]]^2 - (1 + I*Sqrt[3])*Log[(I +
Sqrt[3] + (2*I)*Sinh[x])/(2*Sqrt[3])] *Log[1 + I*Sqrt[3] + 2*Sinh[x]] - ((1
+ I*Sqrt[3])*Log[1 + I*Sqrt[3] + 2*Sinh[x]]^2)/2 - 2*Log[1 + Sinh[x] + Si
nh[x]^2] + (1 - I*Sqrt[3])*Log[1 - I*Sqrt[3] + 2*Sinh[x]] *Log[1 + Sinh[x]
+ Sinh[x]^2] + (1 + I*Sqrt[3])*Log[1 + I*Sqrt[3] + 2*Sinh[x]] *Log[1 + Sinh
[x] + Sinh[x]^2] - (1 + I*Sqrt[3])*PolyLog[2, (-I + Sqrt[3] - (2*I)*Sinh[x
])/(2*Sqrt[3])] + I*(I + Sqrt[3])*PolyLog[2, (I + Sqrt[3] + (2*I)*Sinh[x])
/(2*Sqrt[3])] + 8*Sinh[x] - 4*Log[1 + Sinh[x] + Sinh[x]^2]*Sinh[x] + Log[1
+ Sinh[x] + Sinh[x]^2]^2*Sinh[x]

```

Rubi [A] (verified)

Time = 0.73 (sec) , antiderivative size = 410, normalized size of antiderivative = 1.04, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {4858, 3003, 3008, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(x) \log^2(\sinh(x) + \cosh^2(x)) \, dx \\
 & \quad \downarrow \text{4858} \\
 & \int \log^2(\sinh^2(x) + \sinh(x) + 1) \, d\sinh(x) \\
 & \quad \downarrow \text{3003} \\
 & 2 \int \frac{\sinh(x) \log^2(\sinh^2(x) + \sinh(x) + 1) - \log(\sinh^2(x) + \sinh(x) + 1) \sinh(x)(2\sinh(x) + 1)}{\sinh^2(x) + \sinh(x) + 1} \, d\sinh(x) \\
 & \quad \downarrow \text{3008} \\
 & 2 \int \left(2 \log(\sinh^2(x) + \sinh(x) + 1) - \frac{\log(\sinh^2(x) + \sinh(x) + 1) (\sinh(x) + 2)}{\sinh^2(x) + \sinh(x) + 1} \right) \, d\sinh(x) \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$\frac{\sinh(x) \log^2(\sinh^2(x) + \sinh(x) + 1) - 2 \left(2\sqrt{3} \arctan\left(\frac{2\sinh(x) + 1}{\sqrt{3}}\right) + \frac{1}{2}(1 + i\sqrt{3}) \operatorname{PolyLog}\left(2, -\frac{2i\sinh(x) - \sqrt{3} + i}{2\sqrt{3}}\right) + \frac{1}{2}(1 - i\sqrt{3}) \operatorname{PolyLog}\left(2, \frac{2i\sinh(x) + \sqrt{3} + i}{2\sqrt{3}}\right) \right)}{1}$$

input `Int[Cosh[x]*Log[Cosh[x]^2 + Sinh[x]]^2,x]`

output `Log[1 + Sinh[x] + Sinh[x]^2]^2*Sinh[x] - 2*(2*Sqrt[3]*ArcTan[(1 + 2*Sinh[x])/Sqrt[3]] + ((1 - I*Sqrt[3])*Log[1 - I*Sqrt[3] + 2*Sinh[x]]^2)/4 + ((1 + I*Sqrt[3])*Log[((I/2)*(1 - I*Sqrt[3] + 2*Sinh[x]))/Sqrt[3]]*Log[1 + I*Sqrt[3] + 2*Sinh[x]])/2 + ((1 + I*Sqrt[3])*Log[1 + I*Sqrt[3] + 2*Sinh[x]]^2)/4 + ((1 - I*Sqrt[3])*Log[1 - I*Sqrt[3] + 2*Sinh[x]]*Log[((-1/2*I)*(1 + I*Sqrt[3] + 2*Sinh[x]))/Sqrt[3]])/2 + Log[1 + Sinh[x] + Sinh[x]^2] - ((1 - I*Sqrt[3])*Log[1 - I*Sqrt[3] + 2*Sinh[x]]*Log[1 + Sinh[x] + Sinh[x]^2])/2 - ((1 + I*Sqrt[3])*Log[1 + I*Sqrt[3] + 2*Sinh[x]]*Log[1 + Sinh[x] + Sinh[x]^2])/2 + ((1 + I*Sqrt[3])*PolyLog[2, -1/2*(I - Sqrt[3] + (2*I)*Sinh[x])/Sqrt[3]])/2 + ((1 - I*Sqrt[3])*PolyLog[2, (I + Sqrt[3] + (2*I)*Sinh[x])/(2*Sqrt[3])])/2 - 4*Sinh[x] + 2*Log[1 + Sinh[x] + Sinh[x]^2]*Sinh[x]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3003 `Int[((a_.) + Log[(c_.)*(RFx_)^(p_.)]*(b_.))^(n_.), x_Symbol] := Simp[x*(a + b*Log[c*RFx^p])^n, x] - Simp[b*n*p Int[SimplifyIntegrand[x*(a + b*Log[c*RFx^p])^(n - 1)*(D[RFx, x]/RFx), x], x] /; FreeQ[{a, b, c, p}, x] && RationalFunctionQ[RFx, x] && IGtQ[n, 0]`

rule 3008 `Int[((a_.) + Log[(c_.)*(RFx_)^(p_.)]*(b_.))^(n_.)*(RGx_), x_Symbol] := With[{u = ExpandIntegrand[(a + b*Log[c*RFx^p])^n, RGx, x]}, Int[u, x] /; SumQ[u]] /; FreeQ[{a, b, c, p}, x] && RationalFunctionQ[RFx, x] && RationalFunctionQ[RGx, x] && IGtQ[n, 0]`

rule 4858

```
Int[Cosh[(c_.)*((a_.) + (b_.)*(x_))]*(u_), x_Symbol] := With[{d = FreeFactors[Sinh[c*(a + b*x)], x]}, Simp[d/(b*c) Subst[Int[SubstFor[1, Sinh[c*(a + b*x)]]/d, u, x], x], x, Sinh[c*(a + b*x)]/d, x] /; FunctionOfQ[Sinh[c*(a + b*x)]/d, u, x] /; FreeQ[{a, b, c}, x]
```

Maple [F]

$$\int \cosh(x) \ln(\cosh(x)^2 + \sinh(x))^2 dx$$

input

```
int(cosh(x)*ln(cosh(x)^2+sinh(x))^2,x)
```

output

```
int(cosh(x)*ln(cosh(x)^2+sinh(x))^2,x)
```

Fricas [F]

$$\int \cosh(x) \log^2(\cosh^2(x) + \sinh(x)) dx = \int \cosh(x) \log(\cosh(x)^2 + \sinh(x))^2 dx$$

input

```
integrate(cosh(x)*log(cosh(x)^2+sinh(x))^2,x, algorithm="fricas")
```

output

```
integral(cosh(x)*log(cosh(x)^2 + sinh(x))^2, x)
```

Sympy [F]

$$\int \cosh(x) \log^2(\cosh^2(x) + \sinh(x)) dx = \int \log(\sinh(x) + \cosh^2(x))^2 \cosh(x) dx$$

input

```
integrate(cosh(x)*ln(cosh(x)**2+sinh(x))**2,x)
```

output

```
Integral(log(sinh(x) + cosh(x)**2)**2*cosh(x), x)
```

Maxima [F]

$$\int \cosh(x) \log^2(\cosh^2(x) + \sinh(x)) dx = \int \cosh(x) \log(\cosh(x)^2 + \sinh(x))^2 dx$$

input `integrate(cosh(x)*log(cosh(x)^2+sinh(x))^2,x, algorithm="maxima")`

output

```
1/2*(e^(2*x) - 1)*e^(-x)*log(e^(4*x) + 2*e^(3*x) + 2*e^(2*x) - 2*e^x + 1)^
2 + 2*(2*x - e^(-x) - integrate((2*e^(3*x) + 5*e^(2*x) + 6*e^x - 2)*e^x/(e
^(4*x) + 2*e^(3*x) + 2*e^(2*x) - 2*e^x + 1), x))*log(2)^2 - 4*(x - integra
te((e^(3*x) + 2*e^(2*x) + 2*e^x - 2)*e^x/(e^(4*x) + 2*e^(3*x) + 2*e^(2*x)
- 2*e^x + 1), x))*log(2)^2 + 2*(e^x - integrate((2*e^(3*x) + 2*e^(2*x) - 2
*e^x + 1)*e^x/(e^(4*x) + 2*e^(3*x) + 2*e^(2*x) - 2*e^x + 1), x))*log(2)^2
+ 4*integrate(e^(4*x)/(e^(4*x) + 2*e^(3*x) + 2*e^(2*x) - 2*e^x + 1), x)*lo
g(2)^2 + 6*integrate(e^(3*x)/(e^(4*x) + 2*e^(3*x) + 2*e^(2*x) - 2*e^x + 1)
, x)*log(2)^2 + 6*integrate(e^x/(e^(4*x) + 2*e^(3*x) + 2*e^(2*x) - 2*e^x +
1), x)*log(2)^2 + 4*integrate(x*e^(6*x)/(e^(5*x) + 2*e^(4*x) + 2*e^(3*x)
- 2*e^(2*x) + e^x), x)*log(2) + 8*integrate(x*e^(5*x)/(e^(5*x) + 2*e^(4*x)
+ 2*e^(3*x) - 2*e^(2*x) + e^x), x)*log(2) + 12*integrate(x*e^(4*x)/(e^(5*
x) + 2*e^(4*x) + 2*e^(3*x) - 2*e^(2*x) + e^x), x)*log(2) + 12*integrate(x*
e^(2*x)/(e^(5*x) + 2*e^(4*x) + 2*e^(3*x) - 2*e^(2*x) + e^x), x)*log(2) - 8
*integrate(x*e^x/(e^(5*x) + 2*e^(4*x) + 2*e^(3*x) - 2*e^(2*x) + e^x), x)*l
og(2) - 2*integrate(e^(6*x)*log(e^(4*x) + 2*e^(3*x) + 2*e^(2*x) - 2*e^x +
1)/(e^(5*x) + 2*e^(4*x) + 2*e^(3*x) - 2*e^(2*x) + e^x), x)*log(2) - 4*inte
grate(e^(5*x)*log(e^(4*x) + 2*e^(3*x) + 2*e^(2*x) - 2*e^x + 1)/(e^(5*x) +
2*e^(4*x) + 2*e^(3*x) - 2*e^(2*x) + e^x), x)*log(2) - 6*integrate(e^(4*x)*
log(e^(4*x) + 2*e^(3*x) + 2*e^(2*x) - 2*e^x + 1)/(e^(5*x) + 2*e^(4*x) + ...
```

Giac [F]

$$\int \cosh(x) \log^2(\cosh^2(x) + \sinh(x)) dx = \int \cosh(x) \log(\cosh(x)^2 + \sinh(x))^2 dx$$

input `integrate(cosh(x)*log(cosh(x)^2+sinh(x))^2,x, algorithm="giac")`

output `integrate(cosh(x)*log(cosh(x)^2 + sinh(x))^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \cosh(x) \log^2(\cosh^2(x) + \sinh(x)) dx = \int \cosh(x) \ln(\cosh(x)^2 + \sinh(x))^2 dx$$

input `int(cosh(x)*log(sinh(x) + cosh(x)^2)^2,x)`

output `int(cosh(x)*log(sinh(x) + cosh(x)^2)^2, x)`

Reduce [F]

$$\int \cosh(x) \log^2(\cosh^2(x) + \sinh(x)) dx = \int \cosh(x) \log(\cosh(x)^2 + \sinh(x))^2 dx$$

input `int(cosh(x)*log(cosh(x)^2+sinh(x))^2,x)`

output `int(cosh(x)*log(cosh(x)**2 + sinh(x))**2,x)`

$$3.30 \quad \int \frac{\log(x + \sqrt{1+x})}{1+x^2} dx$$

Optimal result	239
Mathematica [F]	240
Rubi [A] (verified)	241
Maple [A] (verified)	243
Fricas [F]	244
Sympy [F]	244
Maxima [F]	244
Giac [F]	245
Mupad [F(-1)]	245
Reduce [F]	245

Optimal result

Integrand size = 18, antiderivative size = 981

$$\int \frac{\log(x + \sqrt{1+x})}{1+x^2} dx = \text{Too large to display}$$

output

```

1/2*I*ln((1+I)^(1/2)+(1+x)^(1/2))*ln((1+5^(1/2)+2*(1+x)^(1/2))/(1-2*(1+I)^(1/2)+5^(1/2)))+1/2*I*ln((1-I)^(1/2)-(1+x)^(1/2))*ln(x+(1+x)^(1/2))-1/2*I*ln((1+I)^(1/2)-(1+x)^(1/2))*ln(x+(1+x)^(1/2))-1/2*I*ln((1-I)^(1/2)-(1+x)^(1/2))*ln((1+5^(1/2)+2*(1+x)^(1/2))/(1+2*(1-I)^(1/2)+5^(1/2)))+1/2*I*ln((1+I)^(1/2)-(1+x)^(1/2))*ln((1+5^(1/2)+2*(1+x)^(1/2))/(1+2*(1+I)^(1/2)+5^(1/2))) -1/2*I*ln((1-I)^(1/2)+(1+x)^(1/2))*ln((1-5^(1/2)+2*(1+x)^(1/2))/(1-2*(1-I)^(1/2)-5^(1/2)))+1/2*I*polylog(2,2*((1+I)^(1/2)-(1+x)^(1/2))/(1+2*(1+I)^(1/2)-5^(1/2)))-1/2*I*polylog(2,-2*((1-I)^(1/2)+(1+x)^(1/2))/(1-2*(1-I)^(1/2)+5^(1/2)))-1/2*I*ln((1-I)^(1/2)-(1+x)^(1/2))*ln((1-5^(1/2)+2*(1+x)^(1/2))/(1+2*(1-I)^(1/2)-5^(1/2)))-1/2*I*polylog(2,-2*((1-I)^(1/2)+(1+x)^(1/2))/(1-2*(1-I)^(1/2)-5^(1/2)))+1/2*I*polylog(2,-2*((1+I)^(1/2)+(1+x)^(1/2))/(1-2*(1+I)^(1/2)-5^(1/2)))-1/2*I*ln((1-I)^(1/2)+(1+x)^(1/2))*ln((1+5^(1/2)+2*(1+x)^(1/2))/(1-2*(1-I)^(1/2)+5^(1/2)))+1/2*I*ln(x+(1+x)^(1/2))*ln((1-I)^(1/2)+(1+x)^(1/2))+1/2*I*ln((1+I)^(1/2)+(1+x)^(1/2))*ln((1-5^(1/2)+2*(1+x)^(1/2))/(1-2*(1+I)^(1/2)-5^(1/2)))+1/2*I*ln((1+I)^(1/2)-(1+x)^(1/2))*ln((1-5^(1/2)+2*(1+x)^(1/2))/(1+2*(1+I)^(1/2)-5^(1/2)))-1/2*I*polylog(2,2*((1-I)^(1/2)-(1+x)^(1/2))/(1+2*(1-I)^(1/2)+5^(1/2)))+1/2*I*polylog(2,2*((1+I)^(1/2)-(1+x)^(1/2))/(1+2*(1+I)^(1/2)+5^(1/2)))+1/2*I*polylog(2,-2*((1+I)^(1/2)+(1+x)^(1/2))/(1-2*(1+I)^(1/2)+5^(1/2)))-1/2*I*polylog(2,2*((1-I)^(1/2)-(1+x)^(1/2))/(1+2*(1-I)^(1/2)-5^(1/2)))-1/2*I*ln(x+(1+x)^(1/2))*ln((1...

```

Mathematica [F]

$$\int \frac{\log(x + \sqrt{1+x})}{1+x^2} dx = \int \frac{\log(x + \sqrt{1+x})}{1+x^2} dx$$

input

```
Integrate[Log[x + Sqrt[1 + x]]/(1 + x^2), x]
```

output

```
Integrate[Log[x + Sqrt[1 + x]]/(1 + x^2), x]
```

Rubi [A] (verified)

Time = 1.40 (sec) , antiderivative size = 983, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3010, 7292, 3008, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\log(x + \sqrt{x+1})}{x^2 + 1} dx \\
 & \quad \downarrow \text{3010} \\
 & 2 \int \frac{\sqrt{x+1} \log(x + \sqrt{x+1})}{x^2 + 1} d\sqrt{x+1} \\
 & \quad \downarrow \text{7292} \\
 & 2 \int \frac{\sqrt{x+1} \log(x + \sqrt{x+1})}{(x+1)^2 - 2(x+1) + 2} d\sqrt{x+1} \\
 & \quad \downarrow \text{3008} \\
 & 2 \int \left(\frac{i\sqrt{x+1} \log(x + \sqrt{x+1})}{(2+2i) - 2(x+1)} + \frac{i\sqrt{x+1} \log(x + \sqrt{x+1})}{2(x+1) - (2-2i)} \right) d\sqrt{x+1} \\
 & \quad \downarrow \text{2009} \\
 & 2 \left(\frac{1}{4} i \log(\sqrt{1-i} - \sqrt{x+1}) \log(x + \sqrt{x+1}) - \frac{1}{4} i \log(\sqrt{1+i} - \sqrt{x+1}) \log(x + \sqrt{x+1}) + \frac{1}{4} i \log(\sqrt{x+1}) \right)
 \end{aligned}$$

input `Int[Log[x + Sqrt[1 + x]]/(1 + x^2),x]`

output

```

2*((I/4)*Log[Sqrt[1 - I] - Sqrt[1 + x]]*Log[x + Sqrt[1 + x]] - (I/4)*Log[S
qrt[1 + I] - Sqrt[1 + x]]*Log[x + Sqrt[1 + x]] + (I/4)*Log[Sqrt[1 - I] + S
qrt[1 + x]]*Log[x + Sqrt[1 + x]] - (I/4)*Log[Sqrt[1 + I] + Sqrt[1 + x]]*Lo
g[x + Sqrt[1 + x]] - (I/4)*Log[Sqrt[1 - I] + Sqrt[1 + x]]*Log[(1 - Sqrt[5]
+ 2*Sqrt[1 + x])/(1 - 2*Sqrt[1 - I] - Sqrt[5])] - (I/4)*Log[Sqrt[1 - I] -
Sqrt[1 + x]]*Log[(1 - Sqrt[5] + 2*Sqrt[1 + x])/(1 + 2*Sqrt[1 - I] - Sqrt[
5])] + (I/4)*Log[Sqrt[1 + I] + Sqrt[1 + x]]*Log[(1 - Sqrt[5] + 2*Sqrt[1 +
x])/(1 - 2*Sqrt[1 + I] - Sqrt[5])] + (I/4)*Log[Sqrt[1 + I] - Sqrt[1 + x]]*
Log[(1 - Sqrt[5] + 2*Sqrt[1 + x])/(1 + 2*Sqrt[1 + I] - Sqrt[5])] - (I/4)*L
og[Sqrt[1 - I] + Sqrt[1 + x]]*Log[(1 + Sqrt[5] + 2*Sqrt[1 + x])/(1 - 2*Sqr
t[1 - I] + Sqrt[5])] - (I/4)*Log[Sqrt[1 - I] - Sqrt[1 + x]]*Log[(1 + Sqrt[
5] + 2*Sqrt[1 + x])/(1 + 2*Sqrt[1 - I] + Sqrt[5])] + (I/4)*Log[Sqrt[1 + I]
+ Sqrt[1 + x]]*Log[(1 + Sqrt[5] + 2*Sqrt[1 + x])/(1 - 2*Sqrt[1 + I] + Sqr
t[5])] + (I/4)*Log[Sqrt[1 + I] - Sqrt[1 + x]]*Log[(1 + Sqrt[5] + 2*Sqrt[1
+ x])/(1 + 2*Sqrt[1 + I] + Sqrt[5])] - (I/4)*PolyLog[2, (2*(Sqrt[1 - I] -
Sqrt[1 + x]))/(1 + 2*Sqrt[1 - I] - Sqrt[5])] - (I/4)*PolyLog[2, (2*(Sqrt[1
- I] - Sqrt[1 + x]))/(1 + 2*Sqrt[1 - I] + Sqrt[5])] + (I/4)*PolyLog[2, (2
*(Sqrt[1 + I] - Sqrt[1 + x]))/(1 + 2*Sqrt[1 + I] - Sqrt[5])] + (I/4)*PolyL
og[2, (2*(Sqrt[1 + I] - Sqrt[1 + x]))/(1 + 2*Sqrt[1 + I] + Sqrt[5])] - (I/
4)*PolyLog[2, (-2*(Sqrt[1 - I] + Sqrt[1 + x]))/(1 - 2*Sqrt[1 - I] - Sqr...

```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 3008

```
Int[((a_.) + Log[(c_.)*(RFx_)^(p_.)]*(b_.))^(n_.)*(RGx_), x_Symbol] := With
[{u = ExpandIntegrand[(a + b*Log[c*RFx^p])^n, RGx, x]}, Int[u, x] /; SumQ[u
]] /; FreeQ[{a, b, c, p}, x] && RationalFunctionQ[RFx, x] && RationalFunci
onQ[RGx, x] && IGtQ[n, 0]
```

rule 3010

```
Int[((a_.) + Log[u_]*(b_.))*(RFx_), x_Symbol] := With[{lst = SubstForFracti
onalPowerOfLinear[RFx*(a + b*Log[u]), x]}, Simp[lst[[2]]*lst[[4]] Subst[I
nt[lst[[1]], x], x, lst[[3]]^(1/lst[[2])], x] /; !FalseQ[lst]] /; FreeQ[{
a, b}, x] && RationalFunctionQ[RFx, x]
```

rule 7292

```
Int[u_, x_Symbol] := With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v != u]
```

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 722, normalized size of antiderivative = 0.74

method	result
derivativedivides	$\frac{i \left(\ln(\sqrt{1+x}-\sqrt{1-i}) \ln(x+\sqrt{1+x}) - \ln(\sqrt{1+x}-\sqrt{1-i}) \ln\left(\frac{1-\sqrt{5}+2\sqrt{1+x}}{1+2\sqrt{1-i}-\sqrt{5}}\right) - \ln(\sqrt{1+x}-\sqrt{1-i}) \ln\left(\frac{1+\sqrt{5}+2\sqrt{1+x}}{1+2\sqrt{1-i}+\sqrt{5}}\right) \right) - \text{dilog}\left(\frac{1-\sqrt{5}+2\sqrt{1+x}}{1+2\sqrt{1-i}-\sqrt{5}}\right) - \text{dilog}\left(\frac{1+\sqrt{5}+2\sqrt{1+x}}{1+2\sqrt{1-i}+\sqrt{5}}\right)}{2}$
default	$\frac{i \left(\ln(\sqrt{1+x}-\sqrt{1-i}) \ln(x+\sqrt{1+x}) - \ln(\sqrt{1+x}-\sqrt{1-i}) \ln\left(\frac{1-\sqrt{5}+2\sqrt{1+x}}{1+2\sqrt{1-i}-\sqrt{5}}\right) - \ln(\sqrt{1+x}-\sqrt{1-i}) \ln\left(\frac{1+\sqrt{5}+2\sqrt{1+x}}{1+2\sqrt{1-i}+\sqrt{5}}\right) \right) - \text{dilog}\left(\frac{1-\sqrt{5}+2\sqrt{1+x}}{1+2\sqrt{1-i}-\sqrt{5}}\right) - \text{dilog}\left(\frac{1+\sqrt{5}+2\sqrt{1+x}}{1+2\sqrt{1-i}+\sqrt{5}}\right)}{2}$
parts	$\frac{i \left(\ln(\sqrt{1+x}-\sqrt{1-i}) \ln(x+\sqrt{1+x}) - \ln(\sqrt{1+x}-\sqrt{1-i}) \ln\left(\frac{1-\sqrt{5}+2\sqrt{1+x}}{1+2\sqrt{1-i}-\sqrt{5}}\right) - \ln(\sqrt{1+x}-\sqrt{1-i}) \ln\left(\frac{1+\sqrt{5}+2\sqrt{1+x}}{1+2\sqrt{1-i}+\sqrt{5}}\right) \right) - \text{dilog}\left(\frac{1-\sqrt{5}+2\sqrt{1+x}}{1+2\sqrt{1-i}-\sqrt{5}}\right) - \text{dilog}\left(\frac{1+\sqrt{5}+2\sqrt{1+x}}{1+2\sqrt{1-i}+\sqrt{5}}\right)}{2}$

input

```
int(ln(x+(1+x)^(1/2))/(x^2+1), x, method=_RETURNVERBOSE)
```

output

```
1/2*I*(ln((1+x)^(1/2)-(1-I)^(1/2))*ln(x+(1+x)^(1/2))-ln((1+x)^(1/2)-(1-I)^(1/2))*ln((1-5^(1/2)+2*(1+x)^(1/2))/(1+2*(1-I)^(1/2)-5^(1/2)))-ln((1+x)^(1/2)-(1-I)^(1/2))*ln((1+5^(1/2)+2*(1+x)^(1/2))/(1+2*(1-I)^(1/2)+5^(1/2)))-dilog((1-5^(1/2)+2*(1+x)^(1/2))/(1+2*(1-I)^(1/2)-5^(1/2)))-dilog((1+5^(1/2)+2*(1+x)^(1/2))/(1+2*(1-I)^(1/2)+5^(1/2)))+1/2*I*(ln((1-I)^(1/2)+(1+x)^(1/2))*ln(x+(1+x)^(1/2))-ln((1-I)^(1/2)+(1+x)^(1/2))*ln((1-5^(1/2)+2*(1+x)^(1/2))/(1-2*(1-I)^(1/2)-5^(1/2)))-ln((1-I)^(1/2)+(1+x)^(1/2))*ln((1+5^(1/2)+2*(1+x)^(1/2))/(1-2*(1-I)^(1/2)+5^(1/2)))-dilog((1-5^(1/2)+2*(1+x)^(1/2))/(1-2*(1-I)^(1/2)-5^(1/2)))-dilog((1+5^(1/2)+2*(1+x)^(1/2))/(1-2*(1-I)^(1/2)+5^(1/2)))-1/2*I*(ln((1+x)^(1/2)-(1+I)^(1/2))*ln(x+(1+x)^(1/2))-ln((1+x)^(1/2)-(1+I)^(1/2))*ln((1-5^(1/2)+2*(1+x)^(1/2))/(1+2*(1+I)^(1/2)-5^(1/2)))-ln((1+x)^(1/2)-(1+I)^(1/2))*ln((1+5^(1/2)+2*(1+x)^(1/2))/(1+2*(1+I)^(1/2)+5^(1/2)))-dilog((1-5^(1/2)+2*(1+x)^(1/2))/(1+2*(1+I)^(1/2)-5^(1/2)))-dilog((1+5^(1/2)+2*(1+x)^(1/2))/(1+2*(1+I)^(1/2)+5^(1/2)))-1/2*I*(ln((1+I)^(1/2)+(1+x)^(1/2))*ln(x+(1+x)^(1/2))-ln((1+I)^(1/2)+(1+x)^(1/2))*ln((1-5^(1/2)+2*(1+x)^(1/2))/(1-2*(1+I)^(1/2)-5^(1/2)))-ln((1+I)^(1/2)+(1+x)^(1/2))*ln((1+5^(1/2)+2*(1+x)^(1/2))/(1-2*(1+I)^(1/2)+5^(1/2)))-dilog((1-5^(1/2)+2*(1+x)^(1/2))/(1-2*(1+I)^(1/2)-5^(1/2)))-dilog((1+5^(1/2)+2*(1+x)^(1/2))/(1-2*(1+I)^(1/2)+5^(1/2)))))
```

Fricas [F]

$$\int \frac{\log(x + \sqrt{1+x})}{1+x^2} dx = \int \frac{\log(x + \sqrt{x+1})}{x^2+1} dx$$

input `integrate(log(x+(1+x)^(1/2))/(x^2+1),x, algorithm="fricas")`

output `integral(log(x + sqrt(x + 1))/(x^2 + 1), x)`

Sympy [F]

$$\int \frac{\log(x + \sqrt{1+x})}{1+x^2} dx = \int \frac{\log(x + \sqrt{x+1})}{x^2+1} dx$$

input `integrate(ln(x+(1+x)**(1/2))/(x**2+1),x)`

output `Integral(log(x + sqrt(x + 1))/(x**2 + 1), x)`

Maxima [F]

$$\int \frac{\log(x + \sqrt{1+x})}{1+x^2} dx = \int \frac{\log(x + \sqrt{x+1})}{x^2+1} dx$$

input `integrate(log(x+(1+x)^(1/2))/(x^2+1),x, algorithm="maxima")`

output `integrate(log(x + sqrt(x + 1))/(x^2 + 1), x)`

Giac [F]

$$\int \frac{\log(x + \sqrt{1+x})}{1+x^2} dx = \int \frac{\log(x + \sqrt{x+1})}{x^2+1} dx$$

input `integrate(log(x+(1+x)^(1/2))/(x^2+1),x, algorithm="giac")`

output `integrate(log(x + sqrt(x + 1))/(x^2 + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\log(x + \sqrt{1+x})}{1+x^2} dx = \int \frac{\ln(x + \sqrt{x+1})}{x^2+1} dx$$

input `int(log(x + (x + 1)^(1/2))/(x^2 + 1),x)`

output `int(log(x + (x + 1)^(1/2))/(x^2 + 1), x)`

Reduce [F]

$$\int \frac{\log(x + \sqrt{1+x})}{1+x^2} dx = \int \frac{\log(\sqrt{x+1} + x)}{x^2+1} dx$$

input `int(log(x+(1+x)^(1/2))/(x^2+1),x)`

output `int(log(sqrt(x + 1) + x)/(x**2 + 1),x)`

3.31
$$\int \frac{\log^2(x+\sqrt{1+x})}{(1+x)^2} dx$$

Optimal result	247
Mathematica [B] (warning: unable to verify)	248
Rubi [A] (verified)	249
Maple [F]	251
Fricas [F]	252
Sympy [F]	252
Maxima [F]	252
Giac [F]	253
Mupad [F(-1)]	253
Reduce [F]	253

Optimal result

Integrand size = 18, antiderivative size = 555

$$\begin{aligned}
\int \frac{\log^2(x + \sqrt{1+x})}{(1+x)^2} dx = & \log(1+x) + \frac{2\log(x + \sqrt{1+x})}{\sqrt{1+x}} \\
& - 6\log(\sqrt{1+x})\log(x + \sqrt{1+x}) - \frac{\log^2(x + \sqrt{1+x})}{1+x} \\
& - (1 + \sqrt{5})\log(1 - \sqrt{5} + 2\sqrt{1+x}) \\
& + 6\log\left(\frac{1}{2}(-1 + \sqrt{5})\right)\log(1 - \sqrt{5} + 2\sqrt{1+x}) \\
& + (3 + \sqrt{5})\log(x + \sqrt{1+x})\log(1 - \sqrt{5} + 2\sqrt{1+x}) \\
& - \frac{1}{2}(3 + \sqrt{5})\log^2(1 - \sqrt{5} + 2\sqrt{1+x}) \\
& - (1 - \sqrt{5})\log(1 + \sqrt{5} + 2\sqrt{1+x}) \\
& + (3 - \sqrt{5})\log(x + \sqrt{1+x})\log(1 + \sqrt{5} + 2\sqrt{1+x}) \\
& - (3 - \sqrt{5})\log\left(-\frac{1 - \sqrt{5} + 2\sqrt{1+x}}{2\sqrt{5}}\right)\log(1 + \sqrt{5} \\
& \quad + 2\sqrt{1+x}) - \frac{1}{2}(3 - \sqrt{5})\log^2(1 + \sqrt{5} + 2\sqrt{1+x}) - (3 \\
& \quad + \sqrt{5})\log(1 - \sqrt{5} + 2\sqrt{1+x})\log\left(\frac{1 + \sqrt{5} + 2\sqrt{1+x}}{2\sqrt{5}}\right) \\
& + 6\log(\sqrt{1+x})\log\left(1 + \frac{2\sqrt{1+x}}{1 + \sqrt{5}}\right) \\
& + 6\text{PolyLog}\left(2, -\frac{2\sqrt{1+x}}{1 + \sqrt{5}}\right) \\
& - (3 + \sqrt{5})\text{PolyLog}\left(2, -\frac{1 - \sqrt{5} + 2\sqrt{1+x}}{2\sqrt{5}}\right) \\
& - (3 - \sqrt{5})\text{PolyLog}\left(2, \frac{1 + \sqrt{5} + 2\sqrt{1+x}}{2\sqrt{5}}\right) \\
& - 6\text{PolyLog}\left(2, 1 + \frac{2\sqrt{1+x}}{1 - \sqrt{5}}\right)
\end{aligned}$$

output

```

ln(1+x)-3*ln(1+x)*ln(x+(1+x)^(1/2))-ln(x+(1+x)^(1/2))^2/(1+x)+6*ln(1/2*5^(
1/2)-1/2)*ln(1-5^(1/2)+2*(1+x)^(1/2))+3*ln(1+x)*ln(1+2*(1+x)^(1/2)/(5^(1/2
)+1))+6*polylog(2,-2*(1+x)^(1/2)/(5^(1/2)+1))-6*polylog(2,1+2*(1+x)^(1/2)/
(-5^(1/2)+1))-ln(1+5^(1/2)+2*(1+x)^(1/2))*(-5^(1/2)+1)+ln(x+(1+x)^(1/2))*l
n(1+5^(1/2)+2*(1+x)^(1/2))*(3-5^(1/2))-ln(1/10*(-1+5^(1/2)-2*(1+x)^(1/2))*
5^(1/2))*ln(1+5^(1/2)+2*(1+x)^(1/2))*(3-5^(1/2))-1/2*ln(1+5^(1/2)+2*(1+x)^(
1/2))^2*(3-5^(1/2))-polylog(2,1/10*(1+5^(1/2)+2*(1+x)^(1/2))*5^(1/2))*(3-
5^(1/2))-ln(1-5^(1/2)+2*(1+x)^(1/2))*(5^(1/2)+1)+ln(x+(1+x)^(1/2))*ln(1-5^(
1/2)+2*(1+x)^(1/2))*(3+5^(1/2))-1/2*ln(1-5^(1/2)+2*(1+x)^(1/2))^2*(3+5^(1
/2))-ln(1-5^(1/2)+2*(1+x)^(1/2))*ln(1/10*(1+5^(1/2)+2*(1+x)^(1/2))*5^(1/2
))*(3+5^(1/2))-polylog(2,1/10*(-1+5^(1/2)-2*(1+x)^(1/2))*5^(1/2))*(3+5^(1/2
))+2*ln(x+(1+x)^(1/2))/(1+x)^(1/2)

```

Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 1280 vs. 2(555) = 1110.

Time = 6.92 (sec) , antiderivative size = 1280, normalized size of antiderivative = 2.31

$$\int \frac{\log^2(x + \sqrt{1+x})}{(1+x)^2} dx = \text{Too large to display}$$

input

```
Integrate[Log[x + Sqrt[1 + x]]^2/(1 + x)^2,x]
```

output

```
(2*Log[1 + x])/(-1 + Sqrt[5]) - (2*Log[1 + x])/(1 + Sqrt[5]) - (4*Log[-1 +
Sqrt[5] - 2*Sqrt[1 + x]])/(-1 + Sqrt[5]) + (Log[100]*Log[1/2 - Sqrt[5]/2
+ Sqrt[1 + x]])/Sqrt[5] - 6*Log[(2*Sqrt[1 + x])/(-1 + Sqrt[5])]*Log[1/2 -
Sqrt[5]/2 + Sqrt[1 + x]] + 3*Log[1 + x]*Log[1/2 - Sqrt[5]/2 + Sqrt[1 + x]]
- 3*Log[-1 + Sqrt[5] - 2*Sqrt[1 + x]]*Log[1/2 - Sqrt[5]/2 + Sqrt[1 + x]]
- Sqrt[5]*Log[-1 + Sqrt[5] - 2*Sqrt[1 + x]]*Log[1/2 - Sqrt[5]/2 + Sqrt[1 +
x]] + (3*Log[1/2 - Sqrt[5]/2 + Sqrt[1 + x]]^2)/2 + (Sqrt[5]*Log[1/2 - Sqr
t[5]/2 + Sqrt[1 + x]]^2)/2 + (Log[8]*Log[(1 + Sqrt[5])/2 + Sqrt[1 + x]])/(
2*Sqrt[5]) - 3*Log[-1 + Sqrt[5] - 2*Sqrt[1 + x]]*Log[(1 + Sqrt[5])/2 + Sqr
t[1 + x]] - Sqrt[5]*Log[-1 + Sqrt[5] - 2*Sqrt[1 + x]]*Log[(1 + Sqrt[5])/2
+ Sqrt[1 + x]] + (3*Log[(1 + Sqrt[5])/2 + Sqrt[1 + x]]^2)/2 - Log[(1 + Sqr
t[5])/2 + Sqrt[1 + x]]^2/Sqrt[5] + (2*Log[x + Sqrt[1 + x]])/Sqrt[1 + x] -
3*Log[1 + x]*Log[x + Sqrt[1 + x]] + 3*Log[-1 + Sqrt[5] - 2*Sqrt[1 + x]]*Lo
g[x + Sqrt[1 + x]] + Sqrt[5]*Log[-1 + Sqrt[5] - 2*Sqrt[1 + x]]*Log[x + Sqr
t[1 + x]] - Log[x + Sqrt[1 + x]]^2/(1 + x) + (4*Log[1 + Sqrt[5] + 2*Sqrt[1
+ x]])/(1 + Sqrt[5]) - 3*Log[1/2 - Sqrt[5]/2 + Sqrt[1 + x]]*Log[1 + Sqrt[
5] + 2*Sqrt[1 + x]] + Sqrt[5]*Log[1/2 - Sqrt[5]/2 + Sqrt[1 + x]]*Log[1 + S
qrt[5] + 2*Sqrt[1 + x]] - 3*Log[(1 + Sqrt[5])/2 + Sqrt[1 + x]]*Log[1 + Sqr
t[5] + 2*Sqrt[1 + x]] + (7*Log[(1 + Sqrt[5])/2 + Sqrt[1 + x]]*Log[1 + Sqrt
[5] + 2*Sqrt[1 + x]])/(2*Sqrt[5]) + 3*Log[x + Sqrt[1 + x]]*Log[1 + Sqrt...
```

Rubi [A] (verified)

Time = 0.93 (sec) , antiderivative size = 580, normalized size of antiderivative = 1.05, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {7267, 3005, 25, 3008, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\log^2(x + \sqrt{x+1})}{(x+1)^2} dx$$

$$\downarrow 7267$$

$$2 \int \frac{\log^2(x + \sqrt{x+1})}{(x+1)^{3/2}} d\sqrt{x+1}$$

$$\downarrow 3005$$

$$2 \left(\int -\frac{(2\sqrt{x+1}+1)\log(x+\sqrt{x+1})}{(x+1)(-x-\sqrt{x+1})} d\sqrt{x+1} - \frac{\log^2(x+\sqrt{x+1})}{2(x+1)} \right)$$

↓ 25

$$2 \left(- \int \frac{(2\sqrt{x+1}+1)\log(x+\sqrt{x+1})}{(x+1)(-x-\sqrt{x+1})} d\sqrt{x+1} - \frac{\log^2(x+\sqrt{x+1})}{2(x+1)} \right)$$

↓ 3008

$$2 \left(- \int \left(\frac{3\log(x+\sqrt{x+1})}{\sqrt{x+1}} + \frac{\log(x+\sqrt{x+1})}{x+1} + \frac{(-3\sqrt{x+1}-4)\log(x+\sqrt{x+1})}{x+\sqrt{x+1}} \right) d\sqrt{x+1} - \frac{\log^2(x+\sqrt{x+1})}{2(x+1)} \right)$$

↓ 2009

$$2 \left(3 \operatorname{PolyLog} \left(2, -\frac{2\sqrt{x+1}}{1+\sqrt{5}} \right) - \frac{1}{2} (3+\sqrt{5}) \operatorname{PolyLog} \left(2, -\frac{2\sqrt{x+1}-\sqrt{5}+1}{2\sqrt{5}} \right) - \frac{1}{2} (3-\sqrt{5}) \operatorname{PolyLog} \left(2, \frac{2\sqrt{x+1}+\sqrt{5}+1}{2\sqrt{5}} \right) \right)$$

input `Int[Log[x + Sqrt[1 + x]]^2/(1 + x)^2,x]`

output

```
2*(Log[Sqrt[1 + x]] + Log[x + Sqrt[1 + x]]/Sqrt[1 + x] - 3*Log[Sqrt[1 + x]]
)*Log[x + Sqrt[1 + x]] - Log[x + Sqrt[1 + x]]^2/(2*(1 + x)) - ((1 + Sqrt[5]
))*Log[1 - Sqrt[5] + 2*Sqrt[1 + x]]/2 + 3*Log[(-1 + Sqrt[5])/2]*Log[1 - S
qrt[5] + 2*Sqrt[1 + x]] + ((3 + Sqrt[5])*Log[x + Sqrt[1 + x]]*Log[1 - Sqrt
[5] + 2*Sqrt[1 + x]])/2 - ((3 + Sqrt[5])*Log[1 - Sqrt[5] + 2*Sqrt[1 + x]]^
2)/4 - ((1 - Sqrt[5])*Log[1 + Sqrt[5] + 2*Sqrt[1 + x]])/2 + ((3 - Sqrt[5]
)*Log[x + Sqrt[1 + x]]*Log[1 + Sqrt[5] + 2*Sqrt[1 + x]])/2 - ((3 - Sqrt[5]
)*Log[-1/2*(1 - Sqrt[5] + 2*Sqrt[1 + x])/Sqrt[5]]*Log[1 + Sqrt[5] + 2*Sqrt[
1 + x]])/2 - ((3 - Sqrt[5])*Log[1 + Sqrt[5] + 2*Sqrt[1 + x]]^2)/4 - ((3 +
Sqrt[5])*Log[1 - Sqrt[5] + 2*Sqrt[1 + x]]*Log[(1 + Sqrt[5] + 2*Sqrt[1 + x]
)/(2*Sqrt[5]])/2 + 3*Log[Sqrt[1 + x]]*Log[1 + (2*Sqrt[1 + x])/(1 + Sqrt[5
])] + 3*PolyLog[2, (-2*Sqrt[1 + x])/(1 + Sqrt[5])] - ((3 + Sqrt[5])*PolyLo
g[2, -1/2*(1 - Sqrt[5] + 2*Sqrt[1 + x])/Sqrt[5]])/2 - ((3 - Sqrt[5])*PolyLo
g[2, (1 + Sqrt[5] + 2*Sqrt[1 + x])/(2*Sqrt[5]])/2 - 3*PolyLog[2, 1 + (2*
Sqrt[1 + x])/(1 - Sqrt[5])])
```

Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3005 `Int[((a_.) + Log[(c_.)*(Rfx_)^(p_.)]*(b_.))^(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*Log[c*Rfx^p])^n/(e*(m + 1))), x] - Simp[b*n*(p/(e*(m + 1))) Int[SimplifyIntegrand[(d + e*x)^(m + 1)*(a + b*Log[c*Rfx^p])^(n - 1)*(D[Rfx, x]/Rfx), x], x], x] /; FreeQ[{a, b, c, d, e, m, p}, x] && RationalFunctionQ[Rfx, x] && IGtQ[n, 0] && (EqQ[n, 1] || IntegerQ[m]) && NeQ[m, -1]`
- rule 3008 `Int[((a_.) + Log[(c_.)*(Rfx_)^(p_.)]*(b_.))^(n_.)*(RGx_), x_Symbol] := With[{u = ExpandIntegrand[(a + b*Log[c*Rfx^p])^n, RGx, x]}, Int[u, x] /; SumQ[u]] /; FreeQ[{a, b, c, p}, x] && RationalFunctionQ[Rfx, x] && RationalFunctionQ[RGx, x] && IGtQ[n, 0]`
- rule 7267 `Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Simp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])], x] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]]`

Maple **[F]**

$$\int \frac{\ln(x + \sqrt{1+x})^2}{(1+x)^2} dx$$

input `int(ln(x+(1+x)^(1/2))^2/(1+x)^2,x)`

output `int(ln(x+(1+x)^(1/2))^2/(1+x)^2,x)`

Fricas [F]

$$\int \frac{\log^2(x + \sqrt{1+x})}{(1+x)^2} dx = \int \frac{\log(x + \sqrt{x+1})^2}{(x+1)^2} dx$$

input `integrate(log(x+(1+x)^(1/2))^2/(1+x)^2,x, algorithm="fricas")`

output `integral(log(x + sqrt(x + 1))^2/(x^2 + 2*x + 1), x)`

Sympy [F]

$$\int \frac{\log^2(x + \sqrt{1+x})}{(1+x)^2} dx = \int \frac{\log(x + \sqrt{x+1})^2}{(x+1)^2} dx$$

input `integrate(ln(x+(1+x)**(1/2))**2/(1+x)**2,x)`

output `Integral(log(x + sqrt(x + 1))**2/(x + 1)**2, x)`

Maxima [F]

$$\int \frac{\log^2(x + \sqrt{1+x})}{(1+x)^2} dx = \int \frac{\log(x + \sqrt{x+1})^2}{(x+1)^2} dx$$

input `integrate(log(x+(1+x)^(1/2))^2/(1+x)^2,x, algorithm="maxima")`

output `-log(x + sqrt(x + 1))^2/(x + 1) + integrate((2*x + sqrt(x + 1) + 2)*log(x + sqrt(x + 1))/(x^3 + 2*x^2 + (x^2 + 2*x + 1)*sqrt(x + 1) + x), x)`

Giac [F]

$$\int \frac{\log^2(x + \sqrt{1+x})}{(1+x)^2} dx = \int \frac{\log(x + \sqrt{x+1})^2}{(x+1)^2} dx$$

input `integrate(log(x+(1+x)^(1/2))^2/(1+x)^2,x, algorithm="giac")`

output `integrate(log(x + sqrt(x + 1))^2/(x + 1)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\log^2(x + \sqrt{1+x})}{(1+x)^2} dx = \int \frac{\ln(x + \sqrt{x+1})^2}{(x+1)^2} dx$$

input `int(log(x + (x + 1)^(1/2))^2/(x + 1)^2,x)`

output `int(log(x + (x + 1)^(1/2))^2/(x + 1)^2, x)`

Reduce [F]

$$\int \frac{\log^2(x + \sqrt{1+x})}{(1+x)^2} dx = \int \frac{\log(\sqrt{x+1} + x)^2}{x^2 + 2x + 1} dx$$

input `int(log(x+(1+x)^(1/2))^2/(1+x)^2,x)`

output `int(log(sqrt(x + 1) + x)**2/(x**2 + 2*x + 1),x)`

3.32
$$\int \frac{\log(x+\sqrt{1+x})}{x} dx$$

Optimal result	255
Mathematica [A] (verified)	256
Rubi [A] (verified)	257
Maple [A] (verified)	259
Fricas [F]	259
Sympy [F]	260
Maxima [F]	260
Giac [F]	260
Mupad [F(-1)]	261
Reduce [F]	261

Optimal result

Integrand size = 14, antiderivative size = 313

$$\begin{aligned}
\int \frac{\log(x + \sqrt{1+x})}{x} dx &= \log(-1 + \sqrt{1+x}) \log(x + \sqrt{1+x}) \\
&+ \log(1 + \sqrt{1+x}) \log(x + \sqrt{1+x}) \\
&- \log(-1 + \sqrt{1+x}) \log\left(\frac{1 - \sqrt{5} + 2\sqrt{1+x}}{3 - \sqrt{5}}\right) \\
&- \log(1 + \sqrt{1+x}) \log\left(-\frac{1 - \sqrt{5} + 2\sqrt{1+x}}{1 + \sqrt{5}}\right) \\
&- \log(1 + \sqrt{1+x}) \log\left(-\frac{1 + \sqrt{5} + 2\sqrt{1+x}}{1 - \sqrt{5}}\right) \\
&- \log(-1 + \sqrt{1+x}) \log\left(\frac{1 + \sqrt{5} + 2\sqrt{1+x}}{3 + \sqrt{5}}\right) \\
&- \text{PolyLog}\left(2, \frac{2(1 - \sqrt{1+x})}{3 - \sqrt{5}}\right) \\
&- \text{PolyLog}\left(2, \frac{2(1 - \sqrt{1+x})}{3 + \sqrt{5}}\right) \\
&- \text{PolyLog}\left(2, \frac{2(1 + \sqrt{1+x})}{1 - \sqrt{5}}\right) \\
&- \text{PolyLog}\left(2, \frac{2(1 + \sqrt{1+x})}{1 + \sqrt{5}}\right)
\end{aligned}$$

output

```

ln(-1+(1+x)^(1/2))*ln(x+(1+x)^(1/2))+ln(1+(1+x)^(1/2))*ln(x+(1+x)^(1/2))-1
n(-1+(1+x)^(1/2))*ln((1-5^(1/2)+2*(1+x)^(1/2))/(3-5^(1/2)))-ln(1+(1+x)^(1/
2))*ln((-1+5^(1/2)-2*(1+x)^(1/2))/(5^(1/2)+1))-ln(1+(1+x)^(1/2))*ln((-1-5^
(1/2)-2*(1+x)^(1/2))/(-5^(1/2)+1))-ln(-1+(1+x)^(1/2))*ln((1+5^(1/2)+2*(1+x
)^(1/2))/(3+5^(1/2)))-polylog(2,2*(1-(1+x)^(1/2))/(3-5^(1/2)))-polylog(2,2
*(1-(1+x)^(1/2))/(3+5^(1/2)))-polylog(2,2*(1+(1+x)^(1/2))/(-5^(1/2)+1))-po
lylog(2,2*(1+(1+x)^(1/2))/(5^(1/2)+1))

```


Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 303, normalized size of antiderivative = 0.97

$$\begin{aligned}
\int \frac{\log(x + \sqrt{1+x})}{x} dx = & \log(1 - \sqrt{1+x}) \log(x + \sqrt{1+x}) \\
& + \log(1 + \sqrt{1+x}) \log(x + \sqrt{1+x}) \\
& - \log\left(\frac{1}{2}(3 - \sqrt{5})\right) \log(1 - \sqrt{5} + 2\sqrt{1+x}) \\
& - \log\left(\frac{1}{2}(1 + \sqrt{5})\right) \log(1 - \sqrt{5} + 2\sqrt{1+x}) \\
& - \log\left(\frac{1}{2}(3 + \sqrt{5})\right) \log(1 + \sqrt{5} + 2\sqrt{1+x}) \\
& - \log(1 + \sqrt{1+x}) \log\left(-\frac{1 + \sqrt{5} + 2\sqrt{1+x}}{1 - \sqrt{5}}\right) \\
& - \text{PolyLog}\left(2, \frac{2(1 + \sqrt{1+x})}{1 - \sqrt{5}}\right) \\
& + \text{PolyLog}\left(2, \frac{1 - \sqrt{5} + 2\sqrt{1+x}}{3 - \sqrt{5}}\right) \\
& + \text{PolyLog}\left(2, -\frac{1 - \sqrt{5} + 2\sqrt{1+x}}{1 + \sqrt{5}}\right) \\
& + \text{PolyLog}\left(2, \frac{1 + \sqrt{5} + 2\sqrt{1+x}}{3 + \sqrt{5}}\right)
\end{aligned}$$

input `Integrate[Log[x + Sqrt[1 + x]]/x,x]`

output

```

Log[1 - Sqrt[1 + x]]*Log[x + Sqrt[1 + x]] + Log[1 + Sqrt[1 + x]]*Log[x + S
qrt[1 + x]] - Log[(3 - Sqrt[5])/2]*Log[1 - Sqrt[5] + 2*Sqrt[1 + x]] - Log[
(1 + Sqrt[5])/2]*Log[1 - Sqrt[5] + 2*Sqrt[1 + x]] - Log[(3 + Sqrt[5])/2]*L
og[1 + Sqrt[5] + 2*Sqrt[1 + x]] - Log[1 + Sqrt[1 + x]]*Log[-((1 + Sqrt[5]
+ 2*Sqrt[1 + x])/(1 - Sqrt[5]))] - PolyLog[2, (2*(1 + Sqrt[1 + x]))/(1 - S
qrt[5])] + PolyLog[2, (1 - Sqrt[5] + 2*Sqrt[1 + x])/(3 - Sqrt[5])] + PolyL
og[2, -((1 - Sqrt[5] + 2*Sqrt[1 + x])/(1 + Sqrt[5]))] + PolyLog[2, (1 + Sq
rt[5] + 2*Sqrt[1 + x])/(3 + Sqrt[5])]

```

Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 337, normalized size of antiderivative = 1.08, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3010, 25, 3008, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\log(x + \sqrt{x+1})}{x} dx \\
 & \quad \downarrow \text{3010} \\
 & 2 \int \frac{\sqrt{x+1} \log(x + \sqrt{x+1})}{x} d\sqrt{x+1} \\
 & \quad \downarrow \text{25} \\
 & -2 \int -\frac{\sqrt{x+1} \log(x + \sqrt{x+1})}{x} d\sqrt{x+1} \\
 & \quad \downarrow \text{3008} \\
 & -2 \int \left(-\frac{\log(x + \sqrt{x+1})}{2(\sqrt{x+1} - 1)} - \frac{\log(x + \sqrt{x+1})}{2(\sqrt{x+1} + 1)} \right) d\sqrt{x+1} \\
 & \quad \downarrow \text{2009} \\
 & 2 \left(-\frac{1}{2} \text{PolyLog} \left(2, \frac{2(1 - \sqrt{x+1})}{3 - \sqrt{5}} \right) - \frac{1}{2} \text{PolyLog} \left(2, \frac{2(1 - \sqrt{x+1})}{3 + \sqrt{5}} \right) - \frac{1}{2} \text{PolyLog} \left(2, \frac{2(\sqrt{x+1} + 1)}{1 - \sqrt{5}} \right) - \frac{1}{2} \right)
 \end{aligned}$$

input `Int[Log[x + Sqrt[1 + x]]/x,x]`

output

```
2*((Log[-1 + Sqrt[1 + x]]*Log[x + Sqrt[1 + x]])/2 + (Log[1 + Sqrt[1 + x]]*
Log[x + Sqrt[1 + x]])/2 - (Log[-1 + Sqrt[1 + x]]*Log[(1 - Sqrt[5] + 2*Sqrt
[1 + x])/(3 - Sqrt[5])])/2 - (Log[1 + Sqrt[1 + x]]*Log[-((1 - Sqrt[5] + 2*
Sqrt[1 + x])/(1 + Sqrt[5]))])/2 - (Log[1 + Sqrt[1 + x]]*Log[-((1 + Sqrt[5]
+ 2*Sqrt[1 + x])/(1 - Sqrt[5]))])/2 - (Log[-1 + Sqrt[1 + x]]*Log[(1 + Sqr
t[5] + 2*Sqrt[1 + x])/(3 + Sqrt[5])])/2 - PolyLog[2, (2*(1 - Sqrt[1 + x]))
/(3 - Sqrt[5])]/2 - PolyLog[2, (2*(1 - Sqrt[1 + x]))/(3 + Sqrt[5])]/2 - Po
lyLog[2, (2*(1 + Sqrt[1 + x]))/(1 - Sqrt[5])]/2 - PolyLog[2, (2*(1 + Sqrt[
1 + x]))/(1 + Sqrt[5])]/2)
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 3008

```
Int[((a_.) + Log[(c_.)*(RFx_)^(p_.)]*(b_.))^(n_.)*(RGx_), x_Symbol] := With
[{u = ExpandIntegrand[(a + b*Log[c*RFx^p])^n, RGx, x]}, Int[u, x] /; SumQ[u
]] /; FreeQ[{a, b, c, p}, x] && RationalFunctionQ[RFx, x] && RationalFunci
onQ[RGx, x] && IGtQ[n, 0]
```

rule 3010

```
Int[((a_.) + Log[u]*(b_.))*(RFx_), x_Symbol] := With[{lst = SubstForFracti
onalPowerOfLinear[RFx*(a + b*Log[u]), x]}, Simp[lst[[2]]*lst[[4]] Subst[I
nt[lst[[1]], x], x, lst[[3]]^(1/lst[[2])], x] /; !FalseQ[lst]] /; FreeQ[{
a, b}, x] && RationalFunctionQ[RFx, x]
```

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 252, normalized size of antiderivative = 0.81

method	result
derivativedivides	$\ln(-1 + \sqrt{1+x}) \ln(x + \sqrt{1+x}) - \ln(-1 + \sqrt{1+x}) \ln\left(\frac{-1 + \sqrt{5} - 2\sqrt{1+x}}{\sqrt{5}-3}\right) - \ln(-1 + \sqrt{1+x}) \ln\left(\frac{-1 + \sqrt{5} - 2\sqrt{1+x}}{\sqrt{5}-3}\right) - \ln(-1 + \sqrt{1+x}) \ln\left(\frac{-1 + \sqrt{5} - 2\sqrt{1+x}}{\sqrt{5}-3}\right) - \ln(-1 + \sqrt{1+x}) \ln\left(\frac{-1 + \sqrt{5} - 2\sqrt{1+x}}{\sqrt{5}-3}\right)$
default	$\ln(-1 + \sqrt{1+x}) \ln(x + \sqrt{1+x}) - \ln(-1 + \sqrt{1+x}) \ln\left(\frac{-1 + \sqrt{5} - 2\sqrt{1+x}}{\sqrt{5}-3}\right) - \ln(-1 + \sqrt{1+x}) \ln\left(\frac{-1 + \sqrt{5} - 2\sqrt{1+x}}{\sqrt{5}-3}\right) - \ln(-1 + \sqrt{1+x}) \ln\left(\frac{-1 + \sqrt{5} - 2\sqrt{1+x}}{\sqrt{5}-3}\right) - \ln(-1 + \sqrt{1+x}) \ln\left(\frac{-1 + \sqrt{5} - 2\sqrt{1+x}}{\sqrt{5}-3}\right)$
parts	$\ln(x) \ln(x + \sqrt{1+x}) - \ln\left(\sqrt{1+x} - \frac{\sqrt{5}}{2} + \frac{1}{2}\right) \ln(x) + \operatorname{dilog}\left(\frac{1 + \sqrt{1+x}}{\frac{1}{2} + \frac{\sqrt{5}}{2}}\right) + \ln\left(\sqrt{1+x} - \frac{\sqrt{5}}{2} + \frac{1}{2}\right)$

input `int(ln(x+(1+x)^(1/2))/x,x,method=_RETURNVERBOSE)`

output `ln(-1+(1+x)^(1/2))*ln(x+(1+x)^(1/2))-ln(-1+(1+x)^(1/2))*ln((-1+5^(1/2)-2*(1+x)^(1/2))/(5^(1/2)-3))-ln(-1+(1+x)^(1/2))*ln((1+5^(1/2)+2*(1+x)^(1/2))/(3+5^(1/2)))-dilog((-1+5^(1/2)-2*(1+x)^(1/2))/(5^(1/2)-3))-dilog((1+5^(1/2)+2*(1+x)^(1/2))/(3+5^(1/2)))+ln(1+(1+x)^(1/2))*ln(x+(1+x)^(1/2))-ln(1+(1+x)^(1/2))*ln((-1+5^(1/2)-2*(1+x)^(1/2))/(5^(1/2)+1))-ln(1+(1+x)^(1/2))*ln((1+5^(1/2)+2*(1+x)^(1/2))/(5^(1/2)-1))-dilog((-1+5^(1/2)-2*(1+x)^(1/2))/(5^(1/2)+1))-dilog((1+5^(1/2)+2*(1+x)^(1/2))/(5^(1/2)-1))`

Fricas [F]

$$\int \frac{\log(x + \sqrt{1+x})}{x} dx = \int \frac{\log(x + \sqrt{x+1})}{x} dx$$

input `integrate(log(x+(1+x)^(1/2))/x,x, algorithm="fricas")`

output `integral(log(x + sqrt(x + 1))/x, x)`

Sympy [F]

$$\int \frac{\log(x + \sqrt{1+x})}{x} dx = \int \frac{\log(x + \sqrt{x+1})}{x} dx$$

input `integrate(ln(x+(1+x)**(1/2))/x,x)`

output `Integral(log(x + sqrt(x + 1))/x, x)`

Maxima [F]

$$\int \frac{\log(x + \sqrt{1+x})}{x} dx = \int \frac{\log(x + \sqrt{x+1})}{x} dx$$

input `integrate(log(x+(1+x)^(1/2))/x,x, algorithm="maxima")`

output `integrate(log(x + sqrt(x + 1))/x, x)`

Giac [F]

$$\int \frac{\log(x + \sqrt{1+x})}{x} dx = \int \frac{\log(x + \sqrt{x+1})}{x} dx$$

input `integrate(log(x+(1+x)^(1/2))/x,x, algorithm="giac")`

output `integrate(log(x + sqrt(x + 1))/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\log(x + \sqrt{1+x})}{x} dx = \int \frac{\ln(x + \sqrt{x+1})}{x} dx$$

input `int(log(x + (x + 1)^(1/2))/x,x)`output `int(log(x + (x + 1)^(1/2))/x, x)`**Reduce [F]**

$$\int \frac{\log(x + \sqrt{1+x})}{x} dx = \int \frac{\log(\sqrt{x+1} + x)}{x} dx$$

input `int(log(x+(1+x)^(1/2))/x,x)`output `int(log(sqrt(x + 1) + x)/x,x)`

3.33 $\int \arctan(2 \tan(x)) dx$

Optimal result	262
Mathematica [B] (verified)	263
Rubi [A] (verified)	264
Maple [A] (verified)	266
Fricas [B] (verification not implemented)	267
Sympy [F]	268
Maxima [A] (verification not implemented)	268
Giac [F]	269
Mupad [F(-1)]	269
Reduce [F]	269

Optimal result

Integrand size = 5, antiderivative size = 80

$$\int \arctan(2 \tan(x)) dx = x \arctan(2 \tan(x)) + \frac{1}{2}ix \log(1 - 3e^{2ix}) - \frac{1}{2}ix \log\left(1 - \frac{1}{3}e^{2ix}\right) - \frac{1}{4} \text{PolyLog}\left(2, \frac{1}{3}e^{2ix}\right) + \frac{1}{4} \text{PolyLog}(2, 3e^{2ix})$$

output `x*arctan(2*tan(x))+1/2*I*x*ln(1-3*exp(2*I*x))-1/2*I*x*ln(1-1/3*exp(2*I*x))-1/4*polylog(2,1/3*exp(2*I*x))+1/4*polylog(2,3*exp(2*I*x))`

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 262 vs. 2(80) = 160.

Time = 0.18 (sec) , antiderivative size = 262, normalized size of antiderivative = 3.28

$$\begin{aligned}
 \int \arctan(2 \tan(x)) dx &= x \arctan(2 \tan(x)) \\
 &- \frac{1}{4}i \left(4ix \arctan\left(\frac{\cot(x)}{2}\right) + 2i \arccos\left(\frac{5}{3}\right) \arctan(2 \tan(x)) \right. \\
 &\quad + \left(\arccos\left(\frac{5}{3}\right) + 2 \arctan\left(\frac{\cot(x)}{2}\right) \right. \\
 &\quad \left. \left. + 2 \arctan(2 \tan(x)) \right) \log\left(\frac{2i\sqrt{\frac{2}{3}}e^{-ix}}{\sqrt{-5+3\cos(2x)}}\right) \right. \\
 &\quad + \left(\arccos\left(\frac{5}{3}\right) - 2 \arctan\left(\frac{\cot(x)}{2}\right) \right. \\
 &\quad \left. \left. - 2 \arctan(2 \tan(x)) \right) \log\left(\frac{2i\sqrt{\frac{2}{3}}e^{ix}}{\sqrt{-5+3\cos(2x)}}\right) \right. \\
 &\quad - \left(\arccos\left(\frac{5}{3}\right) - 2 \arctan(2 \tan(x)) \right) \log\left(\frac{4i-4\tan(x)}{i+2\tan(x)}\right) \\
 &\quad - \left(\arccos\left(\frac{5}{3}\right) + 2 \arctan(2 \tan(x)) \right) \log\left(\frac{4(i+\tan(x))}{3i+6\tan(x)}\right) \\
 &\quad + i \left(-\text{PolyLog}\left(2, \frac{-3i+6\tan(x)}{i+2\tan(x)}\right) \right. \\
 &\quad \left. \left. + \text{PolyLog}\left(2, \frac{-i+2\tan(x)}{3i+6\tan(x)}\right) \right) \right)
 \end{aligned}$$

input

```
Integrate[ArcTan[2*Tan[x]], x]
```


output

```
x*ArcTan[2*Tan[x]] - (I/4)*((4*I)*x*ArcTan[Cot[x]/2] + (2*I)*ArcCos[5/3]*ArcTan[2*Tan[x]] + (ArcCos[5/3] + 2*ArcTan[Cot[x]/2] + 2*ArcTan[2*Tan[x]])*Log[((2*I)*Sqrt[2/3])/(E^(I*x)*Sqrt[-5 + 3*Cos[2*x]])] + (ArcCos[5/3] - 2*ArcTan[Cot[x]/2] - 2*ArcTan[2*Tan[x]])*Log[((2*I)*Sqrt[2/3]*E^(I*x))/Sqrt[-5 + 3*Cos[2*x]]) - (ArcCos[5/3] - 2*ArcTan[2*Tan[x]])*Log[(4*I - 4*Tan[x])/(I + 2*Tan[x])] - (ArcCos[5/3] + 2*ArcTan[2*Tan[x]])*Log[(4*(I + Tan[x]))/(3*I + 6*Tan[x])] + I*(-PolyLog[2, (-3*I + 6*Tan[x])/(I + 2*Tan[x])] + PolyLog[2, (-I + 2*Tan[x])/(3*I + 6*Tan[x])])])
```

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.04, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {5690, 25, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \arctan(2 \tan(x)) \, dx \\
 & \quad \downarrow \text{5690} \\
 & -3 \int -\frac{e^{2ix} x}{1 - 3e^{2ix}} \, dx - \int \frac{e^{2ix} x}{3 - e^{2ix}} \, dx + x \arctan(2 \tan(x)) \\
 & \quad \downarrow \text{25} \\
 & 3 \int \frac{e^{2ix} x}{1 - 3e^{2ix}} \, dx - \int \frac{e^{2ix} x}{3 - e^{2ix}} \, dx + x \arctan(2 \tan(x)) \\
 & \quad \downarrow \text{2620} \\
 & 3 \left(\frac{1}{6} ix \log(1 - 3e^{2ix}) - \frac{1}{6} i \int \log(1 - 3e^{2ix}) \, dx \right) + \frac{1}{2} i \int \log\left(1 - \frac{1}{3} e^{2ix}\right) \, dx + \\
 & \quad x \arctan(2 \tan(x)) - \frac{1}{2} ix \log\left(1 - \frac{1}{3} e^{2ix}\right) \\
 & \quad \downarrow \text{2715} \\
 & 3 \left(\frac{1}{6} ix \log(1 - 3e^{2ix}) - \frac{1}{12} \int e^{-2ix} \log(1 - 3e^{2ix}) \, de^{2ix} \right) + \frac{1}{4} \int e^{-2ix} \log\left(1 - \frac{1}{3} e^{2ix}\right) \, de^{2ix} + \\
 & \quad x \arctan(2 \tan(x)) - \frac{1}{2} ix \log\left(1 - \frac{1}{3} e^{2ix}\right)
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow 2838 \\
 x \arctan(2 \tan(x)) - \frac{1}{4} \operatorname{PolyLog}\left(2, \frac{1}{3} e^{2ix}\right) + \\
 3 \left(\frac{1}{12} \operatorname{PolyLog}(2, 3e^{2ix}) + \frac{1}{6} ix \log(1 - 3e^{2ix}) \right) - \frac{1}{2} ix \log\left(1 - \frac{1}{3} e^{2ix}\right)
 \end{array}$$

input `Int[ArcTan[2*Tan[x]], x]`

output `x*ArcTan[2*Tan[x]] - (I/2)*x*Log[1 - E^((2*I)*x)/3] - PolyLog[2, E^((2*I)*x)/3]/4 + 3*((I/6)*x*Log[1 - 3E^((2*I)*x)] + PolyLog[2, 3E^((2*I)*x)]/12)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 5690

```
Int[ArcTan[(c_.) + (d_.)*Tan[(a_.) + (b_.)*(x_)]], x_Symbol] := Simp[x*ArcTan[c + d*Tan[a + b*x]], x] + (Simp[b*(1 - I*c - d) Int[x*(E^(2*I*a + 2*I*b*x))/(1 - I*c + d + (1 - I*c - d)*E^(2*I*a + 2*I*b*x))], x], x] - Simp[b*(1 + I*c + d) Int[x*(E^(2*I*a + 2*I*b*x))/(1 + I*c - d + (1 + I*c + d)*E^(2*I*a + 2*I*b*x))], x], x) /; FreeQ[{a, b, c, d}, x] && NeQ[(c + I*d)^2, -1]
```

Maple [A] (verified)

Time = 0.53 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.41

method	result
derivativedivides	$\arctan(2 \tan(x)) \arctan(\tan(x)) + \frac{i \arctan(\tan(x)) \ln\left(1 - \frac{3(1+i \tan(x))^2}{1+\tan(x)^2}\right)}{2} + \frac{\text{polylog}\left(2, \frac{3(1+i \tan(x))}{1+\tan(x)^2}\right)}{4}$
default	$\arctan(2 \tan(x)) \arctan(\tan(x)) + \frac{i \arctan(\tan(x)) \ln\left(1 - \frac{3(1+i \tan(x))^2}{1+\tan(x)^2}\right)}{2} + \frac{\text{polylog}\left(2, \frac{3(1+i \tan(x))}{1+\tan(x)^2}\right)}{4}$
risch	$\frac{ix \ln(1-\sqrt{3}e^{ix})}{2} - \frac{ix \ln(e^{2ix}-\frac{1}{3})}{2} + \frac{ix \ln(1+\sqrt{3}e^{ix})}{2} - \frac{ix \ln(3)}{2} - \frac{\pi x}{2} - \frac{\pi \text{csgn}(i(e^{2ix}-3)) \text{csgn}\left(\frac{i(e^{2ix}-3)}{e^{2ix}+1}\right)}{4}$

input

```
int(arctan(2*tan(x)), x, method=_RETURNVERBOSE)
```

output

```
arctan(2*tan(x))*arctan(tan(x))+1/2*I*arctan(tan(x))*ln(1-3*(1+I*tan(x))^2/(1+tan(x)^2))+1/4*polylog(2,3*(1+I*tan(x))^2/(1+tan(x)^2))-1/2*I*arctan(tan(x))*ln(1-1/3*(1+I*tan(x))^2/(1+tan(x)^2))-1/4*polylog(2,1/3*(1+I*tan(x))^2/(1+tan(x)^2))
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 220 vs. $2(50) = 100$.

Time = 0.10 (sec) , antiderivative size = 220, normalized size of antiderivative = 2.75

$$\int \arctan(2 \tan(x)) dx = x \arctan(2 \tan(x))$$

$$\begin{aligned}
& - \frac{1}{4} i x \log \left(\frac{2(2 \tan(x)^2 + 3i \tan(x) - 1)}{\tan(x)^2 + 1} \right) \\
& + \frac{1}{4} i x \log \left(\frac{2(2 \tan(x)^2 + i \tan(x) + 1)}{3(\tan(x)^2 + 1)} \right) \\
& - \frac{1}{4} i x \log \left(\frac{2(2 \tan(x)^2 - i \tan(x) + 1)}{3(\tan(x)^2 + 1)} \right) \\
& + \frac{1}{4} i x \log \left(\frac{2(2 \tan(x)^2 - 3i \tan(x) - 1)}{\tan(x)^2 + 1} \right) \\
& + \frac{1}{8} \operatorname{Li}_2 \left(-\frac{2(2 \tan(x)^2 + 3i \tan(x) - 1)}{\tan(x)^2 + 1} + 1 \right) \\
& - \frac{1}{8} \operatorname{Li}_2 \left(-\frac{2(2 \tan(x)^2 + i \tan(x) + 1)}{3(\tan(x)^2 + 1)} + 1 \right) \\
& - \frac{1}{8} \operatorname{Li}_2 \left(-\frac{2(2 \tan(x)^2 - i \tan(x) + 1)}{3(\tan(x)^2 + 1)} + 1 \right) \\
& + \frac{1}{8} \operatorname{Li}_2 \left(-\frac{2(2 \tan(x)^2 - 3i \tan(x) - 1)}{\tan(x)^2 + 1} + 1 \right)
\end{aligned}$$

input

```
integrate(arctan(2*tan(x)),x, algorithm="fricas")
```

output

```
x*arctan(2*tan(x)) - 1/4*I*x*log(2*(2*tan(x)^2 + 3*I*tan(x) - 1)/(tan(x)^2 + 1)) + 1/4*I*x*log(2/3*(2*tan(x)^2 + I*tan(x) + 1)/(tan(x)^2 + 1)) - 1/4*I*x*log(2/3*(2*tan(x)^2 - I*tan(x) + 1)/(tan(x)^2 + 1)) + 1/4*I*x*log(2*(2*tan(x)^2 - 3*I*tan(x) - 1)/(tan(x)^2 + 1)) + 1/8*dilog(-2*(2*tan(x)^2 + 3*I*tan(x) - 1)/(tan(x)^2 + 1) + 1) - 1/8*dilog(-2/3*(2*tan(x)^2 + I*tan(x) + 1)/(tan(x)^2 + 1) + 1) - 1/8*dilog(-2/3*(2*tan(x)^2 - I*tan(x) + 1)/(tan(x)^2 + 1) + 1) + 1/8*dilog(-2*(2*tan(x)^2 - 3*I*tan(x) - 1)/(tan(x)^2 + 1) + 1)
```

Sympy [F]

$$\int \arctan(2 \tan(x)) dx = \int \operatorname{atan}(2 \tan(x)) dx$$

input

```
integrate(atan(2*tan(x)),x)
```

output

```
Integral(atan(2*tan(x)), x)
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.05

$$\begin{aligned} \int \arctan(2 \tan(x)) dx &= x \arctan(2 \tan(x)) - \frac{1}{8} \log(4 \tan(x)^2 + 4) \log(4 \tan(x)^2 + 1) \\ &\quad + \frac{1}{8} \log(4 \tan(x)^2 + 1) \log\left(\frac{4}{9} \tan(x)^2 + \frac{4}{9}\right) \\ &\quad - \frac{1}{4} \operatorname{Li}_2(2i \tan(x) - 1) + \frac{1}{4} \operatorname{Li}_2\left(\frac{2}{3}i \tan(x) + \frac{1}{3}\right) \\ &\quad + \frac{1}{4} \operatorname{Li}_2\left(-\frac{2}{3}i \tan(x) + \frac{1}{3}\right) - \frac{1}{4} \operatorname{Li}_2(-2i \tan(x) - 1) \end{aligned}$$

input

```
integrate(arctan(2*tan(x)),x, algorithm="maxima")
```

output

```
x*arctan(2*tan(x)) - 1/8*log(4*tan(x)^2 + 4)*log(4*tan(x)^2 + 1) + 1/8*log
(4*tan(x)^2 + 1)*log(4/9*tan(x)^2 + 4/9) - 1/4*dilog(2*I*tan(x) - 1) + 1/4
*dilog(2/3*I*tan(x) + 1/3) + 1/4*dilog(-2/3*I*tan(x) + 1/3) - 1/4*dilog(-2
*I*tan(x) - 1)
```

Giac [F]

$$\int \arctan(2 \tan(x)) dx = \int \arctan(2 \tan(x)) dx$$

input

```
integrate(arctan(2*tan(x)),x, algorithm="giac")
```

output

```
integrate(arctan(2*tan(x)), x)
```

Mupad [F(-1)]

Timed out.

$$\int \arctan(2 \tan(x)) dx = \int \operatorname{atan}(2 \tan(x)) dx$$

input

```
int(atan(2*tan(x)),x)
```

output

```
int(atan(2*tan(x)), x)
```

Reduce [F]

$$\int \arctan(2 \tan(x)) dx = \int \operatorname{atan}(2 \tan(x)) dx$$

input

```
int(atan(2*tan(x)),x)
```

output

```
int(atan(2*tan(x)),x)
```

3.34 $\int \frac{\arctan(x) \log(x)}{x} dx$

Optimal result	270
Mathematica [A] (verified)	270
Rubi [A] (verified)	271
Maple [A] (verified)	272
Fricas [F]	273
Sympy [F]	273
Maxima [A] (verification not implemented)	273
Giac [F]	274
Mupad [F(-1)]	274
Reduce [F]	274

Optimal result

Integrand size = 8, antiderivative size = 57

$$\int \frac{\arctan(x) \log(x)}{x} dx = \frac{1}{2}i \log(x) \text{PolyLog}(2, -ix) - \frac{1}{2}i \log(x) \text{PolyLog}(2, ix) - \frac{1}{2}i \text{PolyLog}(3, -ix) + \frac{1}{2}i \text{PolyLog}(3, ix)$$

output `1/2*I*ln(x)*polylog(2,-I*x)-1/2*I*ln(x)*polylog(2,I*x)-1/2*I*polylog(3,-I*x)+1/2*I*polylog(3,I*x)`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.77

$$\int \frac{\arctan(x) \log(x)}{x} dx = \frac{1}{2}i(\log(x) \text{PolyLog}(2, -ix) - \log(x) \text{PolyLog}(2, ix) - \text{PolyLog}(3, -ix) + \text{PolyLog}(3, ix))$$

input `Integrate[(ArcTan[x]*Log[x])/x,x]`

output `(I/2)*(Log[x]*PolyLog[2, (-I)*x] - Log[x]*PolyLog[2, I*x] - PolyLog[3, (-I)*x] + PolyLog[3, I*x])`

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.89, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {5540, 2821, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\arctan(x) \log(x)}{x} dx \\
 & \quad \downarrow \text{5540} \\
 & \frac{1}{2}i \int \frac{\log(1-ix) \log(x)}{x} dx - \frac{1}{2}i \int \frac{\log(ix+1) \log(x)}{x} dx \\
 & \quad \downarrow \text{2821} \\
 & \frac{1}{2}i \left(\int \frac{\text{PolyLog}(2, ix)}{x} dx - \text{PolyLog}(2, ix) \log(x) \right) - \\
 & \frac{1}{2}i \left(\int \frac{\text{PolyLog}(2, -ix)}{x} dx - \text{PolyLog}(2, -ix) \log(x) \right) \\
 & \quad \downarrow \text{7143} \\
 & \frac{1}{2}i(\text{PolyLog}(3, ix) - \text{PolyLog}(2, ix) \log(x)) - \frac{1}{2}i(\text{PolyLog}(3, -ix) - \text{PolyLog}(2, -ix) \log(x))
 \end{aligned}$$

input

```
Int[(ArcTan[x]*Log[x])/x,x]
```

output

```
(-1/2*I)*(-Log[x]*PolyLog[2, (-I)*x]) + PolyLog[3, (-I)*x] + (I/2)*(-Log[x]*PolyLog[2, I*x]) + PolyLog[3, I*x]
```


Definitions of rubi rules used

rule 2821

```
Int[(Log[(d_.)*((e_) + (f_.)*(x_)^(m_.))]*((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_.))/(x_), x_Symbol] := Simp[(-PolyLog[2, (-d)*f*x^m]*((a + b*Log[c*x^n])^p/m), x] + Simp[b*n*(p/m) Int[PolyLog[2, (-d)*f*x^m]*((a + b*Log[c*x^n])^(p - 1)/x), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && IGtQ[p, 0] && EqQ[d*e, 1]
```

rule 5540

```
Int[(ArcTan[(c_.)*(x_)^(n_.)]*Log[(d_.)*(x_)^(m_.)]/(x_), x_Symbol] := Simp[I/2 Int[Log[d*x^m]*(Log[1 - I*c*x^n]/x), x], x] - Simp[I/2 Int[Log[d*x^m]*(Log[1 + I*c*x^n]/x), x], x] /; FreeQ[{c, d, m, n}, x]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_)^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.25

method	result
risch	$\frac{i \ln(x)^2 \ln(-i(x+i))}{4} - \frac{i \ln(x)^2 \ln(-ix+1)}{4} - \frac{i \ln(x) \operatorname{polylog}(2, ix)}{2} + \frac{i \operatorname{polylog}(3, ix)}{2} + \frac{i \ln(x) \operatorname{polylog}(2, -ix)}{2} - \frac{i \operatorname{polylog}(3, -ix)}{2}$

input

```
int(arctan(x)*ln(x)/x,x,method=_RETURNVERBOSE)
```

output

```
1/4*I*ln(x)^2*ln(-I*(x+I))-1/4*I*ln(x)^2*ln(1-I*x)-1/2*I*ln(x)*polylog(2,I*x)+1/2*I*polylog(3,I*x)+1/2*I*ln(x)*polylog(2,-I*x)-1/2*I*polylog(3,-I*x)
```

Fricas [F]

$$\int \frac{\arctan(x) \log(x)}{x} dx = \int \frac{\arctan(x) \log(x)}{x} dx$$

input `integrate(arctan(x)*log(x)/x,x, algorithm="fricas")`

output `integral(arctan(x)*log(x)/x, x)`

Sympy [F]

$$\int \frac{\arctan(x) \log(x)}{x} dx = \int \frac{\log(x) \operatorname{atan}(x)}{x} dx$$

input `integrate(atan(x)*ln(x)/x,x)`

output `Integral(log(x)*atan(x)/x, x)`

Maxima [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.54

$$\int \frac{\arctan(x) \log(x)}{x} dx = -\frac{1}{2}i \operatorname{Li}_2(ix) \log(x) + \frac{1}{2}i \operatorname{Li}_2(-ix) \log(x) \\ + \frac{1}{2}i \operatorname{Li}_3(ix) - \frac{1}{2}i \operatorname{Li}_3(-ix)$$

input `integrate(arctan(x)*log(x)/x,x, algorithm="maxima")`

output `-1/2*I*dilog(I*x)*log(x) + 1/2*I*dilog(-I*x)*log(x) + 1/2*I*polylog(3, I*x) - 1/2*I*polylog(3, -I*x)`

Giac [F]

$$\int \frac{\arctan(x) \log(x)}{x} dx = \int \frac{\arctan(x) \log(x)}{x} dx$$

input `integrate(arctan(x)*log(x)/x,x, algorithm="giac")`

output `integrate(arctan(x)*log(x)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan(x) \log(x)}{x} dx = \int \frac{\operatorname{atan}(x) \ln(x)}{x} dx$$

input `int((atan(x)*log(x))/x,x)`

output `int((atan(x)*log(x))/x, x)`

Reduce [F]

$$\int \frac{\arctan(x) \log(x)}{x} dx = \int \frac{\operatorname{atan}(x) \log(x)}{x} dx$$

input `int(atan(x)*log(x)/x,x)`

output `int((atan(x)*log(x))/x,x)`

3.35 $\int \sqrt{1+x^2} \arctan(x)^2 dx$

Optimal result	275
Mathematica [A] (warning: unable to verify)	276
Rubi [A] (verified)	276
Maple [A] (verified)	279
Fricas [F]	280
Sympy [F]	280
Maxima [F]	280
Giac [F]	281
Mupad [F(-1)]	281
Reduce [F]	281

Optimal result

Integrand size = 14, antiderivative size = 121

$$\begin{aligned} \int \sqrt{1+x^2} \arctan(x)^2 dx = & \operatorname{arcsinh}(x) - \sqrt{1+x^2} \arctan(x) + \frac{1}{2} x \sqrt{1+x^2} \arctan(x)^2 \\ & - i \arctan(e^{i \arctan(x)}) \arctan(x)^2 \\ & + i \arctan(x) \operatorname{PolyLog}(2, -ie^{i \arctan(x)}) \\ & - i \arctan(x) \operatorname{PolyLog}(2, ie^{i \arctan(x)}) \\ & - \operatorname{PolyLog}(3, -ie^{i \arctan(x)}) + \operatorname{PolyLog}(3, ie^{i \arctan(x)}) \end{aligned}$$

output

```
arcsinh(x)-I*arctan((1+I*x)/(x^2+1)^(1/2))*arctan(x)^2+I*arctan(x)*polylog
(2,-I*(1+I*x)/(x^2+1)^(1/2))-I*arctan(x)*polylog(2,I*(1+I*x)/(x^2+1)^(1/2)
)-polylog(3,-I*(1+I*x)/(x^2+1)^(1/2))+polylog(3,I*(1+I*x)/(x^2+1)^(1/2))-a
rctan(x)*(x^2+1)^(1/2)+1/2*x*arctan(x)^2*(x^2+1)^(1/2)
```

Mathematica [A] (warning: unable to verify)

Time = 0.14 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.08

$$\int \sqrt{1+x^2} \arctan(x)^2 dx = \coth^{-1}\left(\frac{x}{\sqrt{1+x^2}}\right) - \sqrt{1+x^2} \arctan(x) \\ + \frac{1}{2}x\sqrt{1+x^2} \arctan(x)^2 - i \arctan(e^{i \arctan(x)}) \arctan(x)^2 \\ + i \arctan(x) \operatorname{PolyLog}(2, -ie^{i \arctan(x)}) \\ - i \arctan(x) \operatorname{PolyLog}(2, ie^{i \arctan(x)}) \\ - \operatorname{PolyLog}(3, -ie^{i \arctan(x)}) + \operatorname{PolyLog}(3, ie^{i \arctan(x)})$$

input `Integrate[Sqrt[1 + x^2]*ArcTan[x]^2,x]`output `ArcCoth[x/Sqrt[1 + x^2]] - Sqrt[1 + x^2]*ArcTan[x] + (x*Sqrt[1 + x^2]*ArcTan[x]^2)/2 - I*ArcTan[E^(I*ArcTan[x])]*ArcTan[x]^2 + I*ArcTan[x]*PolyLog[2, (-I)*E^(I*ArcTan[x])] - I*ArcTan[x]*PolyLog[2, I*E^(I*ArcTan[x])] - PolyLog[3, (-I)*E^(I*ArcTan[x])] + PolyLog[3, I*E^(I*ArcTan[x])]`**Rubi [A] (verified)**Time = 0.53 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.11, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {5415, 222, 5423, 3042, 4669, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{x^2+1} \arctan(x)^2 dx \\ \downarrow 5415 \\ \frac{1}{2} \int \frac{\arctan(x)^2}{\sqrt{x^2+1}} dx + \int \frac{1}{\sqrt{x^2+1}} dx + \frac{1}{2}x\sqrt{x^2+1} \arctan(x)^2 - \sqrt{x^2+1} \arctan(x) \\ \downarrow 222 \\ \frac{1}{2} \int \frac{\arctan(x)^2}{\sqrt{x^2+1}} dx + \operatorname{arcsinh}(x) + \frac{1}{2}x\sqrt{x^2+1} \arctan(x)^2 - \sqrt{x^2+1} \arctan(x)$$

↓ 5423

$$\frac{1}{2} \int \sqrt{x^2 + 1} \arctan(x)^2 d \arctan(x) + \operatorname{arcsinh}(x) + \frac{1}{2} x \sqrt{x^2 + 1} \arctan(x)^2 - \sqrt{x^2 + 1} \arctan(x)$$

↓ 3042

$$\frac{1}{2} \int \arctan(x)^2 \csc \left(\arctan(x) + \frac{\pi}{2} \right) d \arctan(x) + \operatorname{arcsinh}(x) + \frac{1}{2} x \sqrt{x^2 + 1} \arctan(x)^2 - \sqrt{x^2 + 1} \arctan(x)$$

↓ 4669

$$\frac{1}{2} \left(-2 \int \arctan(x) \log \left(1 - i e^{i \arctan(x)} \right) d \arctan(x) + 2 \int \arctan(x) \log \left(1 + i e^{i \arctan(x)} \right) d \arctan(x) - 2i \operatorname{arcsinh}(x) + \frac{1}{2} x \sqrt{x^2 + 1} \arctan(x)^2 - \sqrt{x^2 + 1} \arctan(x) \right)$$

↓ 3011

$$\frac{1}{2} \left(2 \left(i \arctan(x) \operatorname{PolyLog} \left(2, -i e^{i \arctan(x)} \right) - i \int \operatorname{PolyLog} \left(2, -i e^{i \arctan(x)} \right) d \arctan(x) \right) - 2 \left(i \arctan(x) \operatorname{PolyLog} \left(2, i e^{i \arctan(x)} \right) - i \int \operatorname{PolyLog} \left(2, i e^{i \arctan(x)} \right) d \arctan(x) \right) - 2i \operatorname{arcsinh}(x) + \frac{1}{2} x \sqrt{x^2 + 1} \arctan(x)^2 - \sqrt{x^2 + 1} \arctan(x) \right)$$

↓ 2720

$$\frac{1}{2} \left(2 \left(i \arctan(x) \operatorname{PolyLog} \left(2, -i e^{i \arctan(x)} \right) - \int e^{-i \arctan(x)} \operatorname{PolyLog} \left(2, -i e^{i \arctan(x)} \right) d e^{i \arctan(x)} \right) - 2 \left(i \arctan(x) \operatorname{PolyLog} \left(2, i e^{i \arctan(x)} \right) - \int e^{i \arctan(x)} \operatorname{PolyLog} \left(2, i e^{i \arctan(x)} \right) d e^{i \arctan(x)} \right) - 2i \operatorname{arcsinh}(x) + \frac{1}{2} x \sqrt{x^2 + 1} \arctan(x)^2 - \sqrt{x^2 + 1} \arctan(x) \right)$$

↓ 7143

$$\frac{1}{2} \left(2 \left(i \arctan(x) \operatorname{PolyLog} \left(2, -i e^{i \arctan(x)} \right) - \operatorname{arcsinh}(x) + \operatorname{PolyLog} \left(3, -i e^{i \arctan(x)} \right) \right) - 2 \left(i \arctan(x) \operatorname{PolyLog} \left(2, i e^{i \arctan(x)} \right) - \operatorname{arcsinh}(x) + \operatorname{PolyLog} \left(3, i e^{i \arctan(x)} \right) \right) - \frac{1}{2} x \sqrt{x^2 + 1} \arctan(x)^2 - \sqrt{x^2 + 1} \arctan(x) \right)$$

input

Int[Sqrt[1 + x^2]*ArcTan[x]^2,x]

output

```
ArcSinh[x] - Sqrt[1 + x^2]*ArcTan[x] + (x*Sqrt[1 + x^2]*ArcTan[x]^2)/2 + (
(-2*I)*ArcTan[E^(I*ArcTan[x])]*ArcTan[x]^2 + 2*(I*ArcTan[x]*PolyLog[2, (-I
)*E^(I*ArcTan[x])]) - PolyLog[3, (-I)*E^(I*ArcTan[x])]) - 2*(I*ArcTan[x]*Po
lyLog[2, I*E^(I*ArcTan[x])]) - PolyLog[3, I*E^(I*ArcTan[x])]))/2
```

Defintions of rubi rules used

rule 222

```
Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt
[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^(c_)*((a_) + (b_)*(x_)))^(n_)]*((f_) + (g_)
*(x_)^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4669

```
Int[csc[(e_) + Pi*(k_) + (f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol
] := Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*k*Pi)*E^(I*(e + f*x))]/f), x] + (-Si
mp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 - E^(I*k*Pi)*E^(I*(e + f*x))], x],
x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 + E^(I*k*Pi)*E^(I*(e + f*x
))], x], x]) /; FreeQ[{c, d, e, f}, x] && IntegerQ[2*k] && IGtQ[m, 0]
```

rule 5415

```
Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^(p_)*((d_) + (e_.)*(x_)^2)^(q_.), x_
Symbol] := Simp[(-b)*p*(d + e*x^2)^q*((a + b*ArcTan[c*x])^(p - 1)/(2*c*q*(2
*q + 1))), x] + (Simp[x*(d + e*x^2)^q*((a + b*ArcTan[c*x])^p/(2*q + 1)), x]
+ Simp[2*d*(q/(2*q + 1)) Int[(d + e*x^2)^(q - 1)*(a + b*ArcTan[c*x])^p,
x], x] + Simp[b^2*d*p*((p - 1)/(2*q*(2*q + 1))) Int[(d + e*x^2)^(q - 1)*(
a + b*ArcTan[c*x])^(p - 2), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[e,
c^2*d] && GtQ[q, 0] && GtQ[p, 1]
```

rule 5423

```
Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^(p_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_S
ymbol] := Simp[1/(c*Sqrt[d]) Subst[Int[(a + b*x)^p*Sec[x], x], x, ArcTan[
c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[e, c^2*d] && IGtQ[p, 0] && Gt
Q[d, 0]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_)^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.47

method	result
default	$\frac{(x \arctan(x) - 2) \arctan(x) \sqrt{x^2 + 1}}{2} - i \left(i \arctan(x)^2 \ln \left(1 - \frac{i(x+1)}{\sqrt{x^2+1}} \right) - i \arctan(x)^2 \ln \left(1 + \frac{i(x+1)}{\sqrt{x^2+1}} \right) + 2 \arctan(x) \operatorname{polylog} \left(2, \frac{i(x+1)}{\sqrt{x^2+1}} \right) \right)$

input

```
int(arctan(x)^2*(x^2+1)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
1/2*(x*arctan(x)-2)*arctan(x)*(x^2+1)^(1/2)-1/2*I*(I*arctan(x)^2*ln(1-I*(I
*x+1)/(x^2+1)^(1/2))-I*arctan(x)^2*ln(1+I*(I*x+1)/(x^2+1)^(1/2))+2*arctan(
x)*polylog(2,I*(I*x+1)/(x^2+1)^(1/2))-2*arctan(x)*polylog(2,-I*(I*x+1)/(x^
2+1)^(1/2))+2*I*polylog(3,I*(I*x+1)/(x^2+1)^(1/2))-2*I*polylog(3,-I*(I*x+1
)/(x^2+1)^(1/2))+4*arctan((I*x+1)/(x^2+1)^(1/2)))
```


Fricas [F]

$$\int \sqrt{1+x^2} \arctan(x)^2 dx = \int \sqrt{x^2+1} \arctan(x)^2 dx$$

input `integrate(arctan(x)^2*(x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(sqrt(x^2 + 1)*arctan(x)^2, x)`

Sympy [F]

$$\int \sqrt{1+x^2} \arctan(x)^2 dx = \int \sqrt{x^2+1} \operatorname{atan}^2(x) dx$$

input `integrate(atan(x)**2*(x**2+1)**(1/2),x)`

output `Integral(sqrt(x**2 + 1)*atan(x)**2, x)`

Maxima [F]

$$\int \sqrt{1+x^2} \arctan(x)^2 dx = \int \sqrt{x^2+1} \arctan(x)^2 dx$$

input `integrate(arctan(x)^2*(x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(x^2 + 1)*arctan(x)^2, x)`

Giac [F]

$$\int \sqrt{1+x^2} \arctan(x)^2 dx = \int \sqrt{x^2+1} \arctan(x)^2 dx$$

input `integrate(arctan(x)^2*(x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(x^2 + 1)*arctan(x)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{1+x^2} \arctan(x)^2 dx = \int \operatorname{atan}(x)^2 \sqrt{x^2+1} dx$$

input `int(atan(x)^2*(x^2 + 1)^(1/2),x)`

output `int(atan(x)^2*(x^2 + 1)^(1/2), x)`

Reduce [F]

$$\int \sqrt{1+x^2} \arctan(x)^2 dx = \int \sqrt{x^2+1} \operatorname{atan}(x)^2 dx$$

input `int(atan(x)^2*(x^2+1)^(1/2),x)`

output `int(sqrt(x**2 + 1)*atan(x)**2,x)`

CHAPTER 4

APPENDIX

4.1	Listing of Grading functions	282
4.2	Links to plain text integration problems used in this report for each CAS .	300

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
  ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "
  ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]===Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]===Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]===Plus || Head[expn]===Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```


Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022  add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue
fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result    := ExpnType(result);
      ExpnType_optimal   := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
#     is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

#      antiderivative
#      "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

        end if
    else #result contains complex but optimal is not
        if debug then
            print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
    fi;
else # result do not contain complex
    # this assumes optimal do not as well. No check is needed here.
    if debug then
        print("result do not contain complex, this assumes optimal do not as well
    fi;

```



```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
            fi;
        fi;
    else #ExpnType(result) > ExpnType(optimal)
        if debug then
            print("ExpnType(result) > ExpnType(optimal)");
        fi;
        return "C",cat("Result contains higher order function than in optimal. Order ",
                        convert(ExpnType_result,string)," vs. order ",
                        convert(ExpnType_optimal,string),".");
    fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9
end proc

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```



```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file