

Computer Algebra Independent Integration Tests

Summer 2024

0-Independent-test-suites/4-Calculus-Textbook-Problems

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 1:12am

Contents

1	Introduction	5
1.1	Listing of CAS systems tested	6
1.2	Results	7
1.3	Time and leaf size Performance	11
1.4	Performance based on number of rules Rubi used	13
1.5	Performance based on number of steps Rubi used	14
1.6	Solved integrals histogram based on leaf size of result	15
1.7	Solved integrals histogram based on CPU time used	16
1.8	Leaf size vs. CPU time used	17
1.9	list of integrals with no known antiderivative	18
1.10	List of integrals solved by CAS but has no known antiderivative	18
1.11	list of integrals solved by CAS but failed verification	18
1.12	Timing	19
1.13	Verification	19
1.14	Important notes about some of the results	20
1.15	Current tree layout of integration tests	23
1.16	Design of the test system	24
2	detailed summary tables of results	25
2.1	List of integrals sorted by grade for each CAS	26
2.2	Detailed conclusion table per each integral for all CAS systems	30
2.3	Detailed conclusion table specific for Rubi results	50
3	Listing of integrals	53
3.1	$\int \frac{1-3x^4}{(-2+x)(1+x^2)^2} dx$	56
3.2	$\int \frac{-9-9x+2x^2}{-9x+x^3} dx$	62
3.3	$\int \frac{1+2x^2+x^5}{-x+x^3} dx$	67
3.4	$\int \frac{3+2x^2}{(-1+x)^2x} dx$	72
3.5	$\int \frac{-1+2x^2}{(-1+4x)(1+x^2)} dx$	77
3.6	$\int \frac{-3+2x-3x^2+x^3}{1+x^2} dx$	82

3.7	$\int \frac{x+10x^2+6x^3+x^4}{10+6x+x^2} dx$	87
3.8	$\int \frac{1}{-18+27x-7x^2-3x^3+x^4} dx$	92
3.9	$\int \frac{1+x^3}{-2+x} dx$	97
3.10	$\int \frac{3x-4x^2+3x^3}{1+x^2} dx$	102
3.11	$\int \frac{5+3x}{1-x-x^2+x^3} dx$	107
3.12	$\int \frac{-1-x-x^3+x^4}{-x^2+x^3} dx$	112
3.13	$\int \frac{2+x+x^2+x^3}{2+3x^2+x^4} dx$	117
3.14	$\int \frac{-4+8x-4x^2+4x^3-x^4+x^5}{(2+x^2)^3} dx$	122
3.15	$\int \frac{-1-3x+x^2}{-2x+x^2+x^3} dx$	128
3.16	$\int \frac{3-x+3x^2-2x^3+x^4}{3x-2x^2+x^3} dx$	133
3.17	$\int \frac{-1+x+x^3}{(1+x^2)^2} dx$	138
3.18	$\int \frac{1+2x-x^2+8x^3+x^4}{(x+x^2)(1+x^3)} dx$	143
3.19	$\int \frac{15-5x+x^2+x^3}{(5+x^2)(3+2x+x^2)} dx$	149
3.20	$\int \frac{-3+25x+23x^2+32x^3+15x^4+7x^5+x^6}{(1+x^2)^2(2+x+x^2)^2} dx$	155
3.21	$\int \frac{1}{(1+x^2)(4+x^2)} dx$	161
3.22	$\int \frac{a+bx^3}{1+x^2} dx$	166
3.23	$\int \frac{x+x^2}{(4+x)(-4+x^2)} dx$	171
3.24	$\int \frac{4+x^2}{(1+x^2)(2+x^2)} dx$	176
3.25	$\int \frac{5-4x+3x^2+x^4}{(-1+x)^2(1+x^2)} dx$	181
3.26	$\int \frac{1+x^4}{2+x^2} dx$	186
3.27	$\int \frac{2+2x+x^4}{x^4+x^5} dx$	191
3.28	$\int \frac{-1-5x+2x^2}{2-x-2x^2+x^3} dx$	196
3.29	$\int \frac{2+x+x^3}{1+2x^2+x^4} dx$	201
3.30	$\int \frac{1+2x+x^2+x^3}{1+2x^2+x^4} dx$	207
3.31	$\int \frac{3+4x}{(1+x^2)(2+x^2)} dx$	213
3.32	$\int \frac{2+x}{(1+x^2)(4+x^2)} dx$	219
3.33	$\int \frac{2-x+x^3}{-7-6x+x^2} dx$	225
3.34	$\int \frac{-1+x^5}{-1+x^2} dx$	230
3.35	$\int \frac{5+2x-x^2+x^3}{1+x+x^2} dx$	235
3.36	$\int \frac{-3+x-2x^3+x^4}{10-8x+2x^2} dx$	240
3.37	$\int \frac{1+2x+3x^2+x^3}{(-3+x)(-2+x)(-1+x)} dx$	245
3.38	$\int \frac{2-7x+x^2-x^3+x^4}{-24-14x+x^2+x^3} dx$	250
3.39	$\int \frac{2+x^2}{(-1+x)^2x(1+x)} dx$	255
3.40	$\int \frac{3+x^2+x^3}{(2+x^2)^2} dx$	260
3.41	$\int \frac{-35+70x-4x^2+2x^3}{(26-10x+x^2)(17-2x+x^2)} dx$	266

3.42	$\int \frac{2+x^2}{(-5+x)(-3+x)(4+x)} dx$	272
3.43	$\int \frac{x^4}{(-1+x)(2+x^2)} dx$	277
3.44	$\int \frac{-1+7x+2x^2}{-1-x+x^2+x^3} dx$	283
3.45	$\int \frac{1+2x}{-1+3x-3x^2+x^3} dx$	288
3.46	$\int \frac{5-5x+7x^2+x^3}{(-1+x)^2(1+x)^3} dx$	293
3.47	$\int \frac{1+3x+3x^2}{1+2x+2x^2+x^3} dx$	298
3.48	$\int \frac{-1+2x+x^2}{-2x+3x^2+2x^3} dx$	303
3.49	$\int \frac{1+4x-2x^2+x^4}{1-x-x^2+x^3} dx$	308
3.50	$\int \frac{4-x+2x^2}{4x+x^3} dx$	313
3.51	$\int \frac{1+x^2+x^3}{(-1+x)x(1+x^2)^3(1+x+x^2)} dx$	318
3.52	$\int \frac{1-3x+2x^2-x^3}{(1+x^2)^2} dx$	325
3.53	$\int \frac{1-3x+2x^2-x^3}{x(1+x^2)^2} dx$	331
3.54	$\int \frac{1-x-x^2+x^3+x^4}{-x+x^3} dx$	337
3.55	$\int \frac{2-4x^2+x^3}{(1+x^2)(2+x^2)} dx$	342
3.56	$\int \frac{1+x^2+x^4}{(1+x^2)(4+x^2)^2} dx$	347
3.57	$\int \frac{1+x^2+x^3}{2x^2+x^3+x^4} dx$	352
3.58	$\int \frac{1-12x+x^2+x^3}{-12+x+x^2} dx$	357
3.59	$\int \frac{-3+5x+6x^2}{-3x+2x^2+x^3} dx$	362
3.60	$\int \frac{-2+3x+5x^2}{2x^2+x^3} dx$	367
3.61	$\int \frac{18-2x-4x^2}{-6+x+4x^2+x^3} dx$	372
3.62	$\int \frac{1+x-2x^2+x^3}{4+5x^2+x^4} dx$	377
3.63	$\int \frac{-32+5x-27x^2+4x^3}{-70-299x-286x^2+50x^3-13x^4+30x^5} dx$	383
3.64	$\int \frac{8-13x^2-7x^3+12x^5}{4-20x+41x^2-80x^3+116x^4-80x^5+100x^6} dx$	389
3.65	$\int \frac{9+x^4}{x^2(9+x^2)} dx$	395
3.66	$\int \frac{2x+x^4}{1+x^2} dx$	400
3.67	$\int \frac{-x+x^3}{(-1+x)^2(1+x^2)} dx$	405
3.68	$\int \frac{2+5x+3x^2+2x^3}{1+x+x^2} dx$	410
3.69	$\int \frac{3-4x-5x^2+3x^3}{x^3(-1+x+x^2)} dx$	415
3.70	$\int \frac{4+8x+5x^2+2x^3}{(2+2x+x^2)^2} dx$	421
3.71	$\int \frac{(-1+x)^4 x^4}{1+x^2} dx$	427
3.72	$\int \frac{-20x+4x^2}{9-10x^2+x^4} dx$	432
3.73	$\int \frac{-1+x+4x^3}{(-1+x)x^2(1+x^2)} dx$	438
3.74	$\int \frac{1-3x+2x^2-4x^3+x^4}{(1+x^2)^3} dx$	443
3.75	$\int \frac{1-3x+2x^2-4x^3+x^4}{1+3x^2+3x^4+x^6} dx$	448

3.76	$\int \frac{1+x+2x^2+2x^3}{x^2+x^3+x^4} dx$	454
4	Appendix	459
4.1	Listing of Grading functions	459
4.2	Links to plain text integration problems used in this report for each CAS477	

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	6
1.2	Results	7
1.3	Time and leaf size Performance	11
1.4	Performance based on number of rules Rubi used	13
1.5	Performance based on number of steps Rubi used	14
1.6	Solved integrals histogram based on leaf size of result	15
1.7	Solved integrals histogram based on CPU time used	16
1.8	Leaf size vs. CPU time used	17
1.9	list of integrals with no known antiderivative	18
1.10	List of integrals solved by CAS but has no known antiderivative	18
1.11	list of integrals solved by CAS but failed verification	18
1.12	Timing	19
1.13	Verification	19
1.14	Important notes about some of the results	20
1.15	Current tree layout of integration tests	23
1.16	Design of the test system	24

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [76]. This is test number [4].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (76)	0.00 (0)
Mathematica	100.00 (76)	0.00 (0)
Maple	100.00 (76)	0.00 (0)
Fricas	100.00 (76)	0.00 (0)
Mupad	100.00 (76)	0.00 (0)
Giac	100.00 (76)	0.00 (0)
Maxima	100.00 (76)	0.00 (0)
Reduce	100.00 (76)	0.00 (0)
Sympy	100.00 (76)	0.00 (0)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Mathematica	100.000	0.000	0.000	0.000
Maple	100.000	0.000	0.000	0.000
Maxima	100.000	0.000	0.000	0.000
Fricas	97.368	2.632	0.000	0.000
Sympy	97.368	1.316	1.316	0.000
Giac	96.053	3.947	0.000	0.000
Mupad	0.000	100.000	0.000	0.000
Reduce	0.000	100.000	0.000	0.000

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	0	0.00	0.00	0.00
Maple	0	0.00	0.00	0.00
Mupad	0	0.00	0.00	0.00
Giac	0	0.00	0.00	0.00
Maxima	0	0.00	0.00	0.00
Reduce	0	0.00	0.00	0.00
Sympy	0	0.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Mathematica	0.01
Mupad	0.06
Sympy	0.06
Fricas	0.07
Maxima	0.07
Maple	0.09
Giac	0.12
Reduce	0.15
Rubi	0.21

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Maple	24.21	0.86	21.50	0.85
Maxima	24.21	0.85	21.00	0.82
Sympy	25.80	0.86	20.00	0.82
Giac	26.00	0.94	23.00	0.85
Mupad	28.18	0.96	22.00	0.87
Mathematica	29.17	1.01	25.00	1.00
Fricas	29.45	1.00	23.00	0.88
Rubi	29.89	1.01	25.50	1.00
Reduce	35.21	1.15	23.00	0.88

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram shows that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand,the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	26
2.2	Detailed conclusion table per each integral for all CAS systems	30
2.3	Detailed conclusion table specific for Rubi results	50

2.1 List of integrals sorted by grade for each CAS

Rubi	26
Mma	26
Maple	27
Fricas	27
Maxima	28
Giac	28
Mupad	28
Sympy	29
Reduce	29

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76 }
}

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76 }
}

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 13, 14, 15, 16, 17, 18, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76 }

B grade { 11, 20 }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Maxima

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76 }
}

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Giac

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 68, 69, 70, 71, 72, 73, 74, 75, 76 }
}

B grade { 11, 25, 67 }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76 }
}

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Sympy

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76 }

B grade { 11 }

C grade { 22 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Reduce

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76 }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	48	57	33	33	47	37	34	64	38
N.S.	1	1.12	1.33	0.77	0.77	1.09	0.86	0.79	1.49	0.88
time (sec)	N/A	0.247	0.016	0.099	0.100	0.073	0.076	0.123	0.153	0.030

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	16	15	15	14	18	15	21
N.S.	1	1.00	1.00	0.94	0.88	0.88	0.82	1.06	0.88	1.24
time (sec)	N/A	0.197	0.004	0.088	0.023	0.076	0.056	0.116	0.151	0.038

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	25	22	21	21	20	24	21	30
N.S.	1	1.00	1.00	0.88	0.84	0.84	0.80	0.96	0.84	1.20
time (sec)	N/A	0.203	0.005	0.093	0.032	0.067	0.053	0.117	0.152	0.077

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	20	19	18	24	14	28	30	18
N.S.	1	1.00	0.91	0.86	0.82	1.09	0.64	1.27	1.36	0.82
time (sec)	N/A	0.155	0.006	0.089	0.030	0.067	0.042	0.114	0.158	0.023

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	38	22	21	21	26	22	21	25
N.S.	1	1.00	1.41	0.81	0.78	0.78	0.96	0.81	0.78	0.93
time (sec)	N/A	0.196	0.008	0.105	0.104	0.074	0.063	0.122	0.163	0.126

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	21	18	17	17	15	17	17	17
N.S.	1	1.00	1.00	0.86	0.81	0.81	0.71	0.81	0.81	0.81
time (sec)	N/A	0.165	0.004	0.065	0.100	0.066	0.028	0.117	0.154	0.103

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	27	24	23	23	22	23	23	23
N.S.	1	1.00	1.00	0.89	0.85	0.85	0.81	0.85	0.85	0.85
time (sec)	N/A	0.202	0.005	0.219	0.103	0.067	0.043	0.124	0.159	0.051

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	39	26	25	25	26	29	25	25
N.S.	1	1.00	1.00	0.67	0.64	0.64	0.67	0.74	0.64	0.64
time (sec)	N/A	0.182	0.004	0.040	0.023	0.073	0.110	0.114	0.151	0.091

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	23	19	18	18	17	19	18	18
N.S.	1	1.00	1.05	0.86	0.82	0.82	0.77	0.86	0.82	0.82
time (sec)	N/A	0.164	0.003	0.076	0.023	0.063	0.027	0.120	0.148	0.016

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	13	14	13	13	13
N.S.	1	1.00	1.00	0.93	0.87	0.87	0.93	0.87	0.87	0.87
time (sec)	N/A	0.186	0.004	0.079	0.104	0.065	0.033	0.116	0.154	0.020

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	24	21	20	26	17	22	35	10
N.S.	1	1.00	2.00	1.75	1.67	2.17	1.42	1.83	2.92	0.83
time (sec)	N/A	0.163	0.009	0.036	0.024	0.063	0.041	0.122	0.152	0.047

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	25	22	21	22	19	23	22	19
N.S.	1	1.00	1.00	0.88	0.84	0.88	0.76	0.92	0.88	0.76
time (sec)	N/A	0.203	0.005	0.095	0.028	0.071	0.040	0.117	0.159	0.025

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	13	12	11	11	10	11	11	11
N.S.	1	1.00	1.00	0.92	0.85	0.85	0.77	0.85	0.85	0.85
time (sec)	N/A	0.174	0.006	0.050	0.101	0.068	0.046	0.121	0.157	0.082

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	35	31	35	55	36	30	85	35
N.S.	1	1.00	1.00	0.89	1.00	1.57	1.03	0.86	2.43	1.00
time (sec)	N/A	0.200	0.011	0.089	0.106	0.088	0.059	0.121	0.156	0.025

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	23	18	17	17	17	20	17	17
N.S.	1	1.00	1.00	0.78	0.74	0.74	0.74	0.87	0.74	0.74
time (sec)	N/A	0.195	0.005	0.033	0.023	0.073	0.057	0.125	0.159	0.105

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	23	20	19	19	19	20	19	19
N.S.	1	1.00	1.00	0.87	0.83	0.83	0.83	0.87	0.83	0.83
time (sec)	N/A	0.216	0.006	0.036	0.029	0.068	0.049	0.121	0.147	0.082

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	28	25	24	23	32	20	23	41	25
N.S.	1	0.97	0.86	0.83	0.79	1.10	0.69	0.79	1.41	0.86
time (sec)	N/A	0.164	0.009	0.083	0.101	0.065	0.047	0.121	0.155	0.019

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	44	44	42	41	58	49	43	88	55
N.S.	1	1.00	1.00	0.95	0.93	1.32	1.11	0.98	2.00	1.25
time (sec)	N/A	0.399	0.017	0.106	0.108	0.075	0.091	0.120	0.156	0.075

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	46	39	38	38	51	38	36	88
N.S.	1	1.00	1.00	0.85	0.83	0.83	1.11	0.83	0.78	1.91
time (sec)	N/A	0.310	0.013	0.304	0.101	0.077	0.100	0.118	0.155	0.090

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	33	34	44	72	41	44	129	56
N.S.	1	1.00	1.00	1.03	1.33	2.18	1.24	1.33	3.91	1.70
time (sec)	N/A	0.340	0.015	0.278	0.029	0.068	0.071	0.119	0.150	0.037

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	12	11	11	10	11	11	11
N.S.	1	1.00	1.00	0.71	0.65	0.65	0.59	0.65	0.65	0.65
time (sec)	N/A	0.132	0.005	0.097	0.106	0.073	0.066	0.122	0.158	0.070

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	22	21	20	20	34	20	20	20
N.S.	1	1.00	0.92	0.88	0.83	0.83	1.42	0.83	0.83	0.83
time (sec)	N/A	0.169	0.007	0.085	0.103	0.069	0.073	0.121	0.157	0.068

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	23	18	17	17	17	20	17	19
N.S.	1	1.00	1.53	1.20	1.13	1.13	1.13	1.33	1.13	1.27
time (sec)	N/A	0.205	0.005	0.083	0.100	0.074	0.065	0.119	0.158	0.077

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	18	17	17	19	17	16	17
N.S.	1	1.00	1.00	0.90	0.85	0.85	0.95	0.85	0.80	0.85
time (sec)	N/A	0.135	0.007	0.095	0.102	0.081	0.066	0.119	0.159	0.032

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	33	28	27	44	29	60	56	35
N.S.	1	1.00	0.89	0.76	0.73	1.19	0.78	1.62	1.51	0.95
time (sec)	N/A	0.198	0.017	0.123	0.109	0.072	0.068	0.120	0.158	0.028

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	26	22	21	21	26	21	20	21
N.S.	1	1.00	1.00	0.85	0.81	0.81	1.00	0.81	0.77	0.81
time (sec)	N/A	0.155	0.006	0.080	0.103	0.068	0.042	0.116	0.153	0.018

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	12	11	10	16	10	11	16	10
N.S.	1	1.00	1.00	0.92	0.83	1.33	0.83	0.92	1.33	0.83
time (sec)	N/A	0.189	0.004	0.083	0.023	0.067	0.033	0.118	0.155	0.070

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	21	18	17	17	15	20	17	21
N.S.	1	1.00	1.00	0.86	0.81	0.81	0.71	0.95	0.81	1.00
time (sec)	N/A	0.177	0.004	0.042	0.023	0.071	0.068	0.121	0.153	0.106

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	22	21	20	34	17	20	41	20
N.S.	1	1.00	1.00	0.95	0.91	1.55	0.77	0.91	1.86	0.91
time (sec)	N/A	0.176	0.007	0.044	0.101	0.070	0.043	0.120	0.156	0.039

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	24	21	20	32	19	20	41	22
N.S.	1	1.00	1.00	0.88	0.83	1.33	0.79	0.83	1.71	0.92
time (sec)	N/A	0.174	0.007	0.049	0.105	0.068	0.048	0.122	0.150	0.037

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	38	36	34	33	33	39	33	32	56
N.S.	1	1.06	1.00	0.94	0.92	0.92	1.08	0.92	0.89	1.56
time (sec)	N/A	0.170	0.011	0.091	0.104	0.076	0.085	0.119	0.147	0.091

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	45	37	28	27	27	29	27	27	37
N.S.	1	1.22	1.00	0.76	0.73	0.73	0.78	0.73	0.73	1.00
time (sec)	N/A	0.166	0.006	0.097	0.101	0.073	0.082	0.109	0.160	0.030

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	29	22	21	21	22	23	21	21
N.S.	1	1.00	1.00	0.76	0.72	0.72	0.76	0.79	0.72	0.72
time (sec)	N/A	0.181	0.004	0.092	0.026	0.067	0.055	0.125	0.160	0.083

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	19	16	15	15	14	16	15	15
N.S.	1	1.00	1.00	0.84	0.79	0.79	0.74	0.84	0.79	0.79
time (sec)	N/A	0.162	0.003	0.082	0.027	0.062	0.036	0.122	0.152	0.020

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	41	35	34	34	46	34	33	36
N.S.	1	1.00	1.00	0.85	0.83	0.83	1.12	0.83	0.80	0.88
time (sec)	N/A	0.186	0.010	0.163	0.103	0.065	0.051	0.121	0.156	0.075

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	39	32	31	31	34	31	31	31
N.S.	1	1.00	0.95	0.78	0.76	0.76	0.83	0.76	0.76	0.76
time (sec)	N/A	0.190	0.005	0.133	0.104	0.066	0.048	0.121	0.142	0.025

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	30	24	21	20	20	24	23	20	20
N.S.	1	1.00	0.80	0.70	0.67	0.67	0.80	0.77	0.67	0.67
time (sec)	N/A	0.208	0.008	0.101	0.025	0.080	0.062	0.114	0.156	0.025

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	35	28	27	27	31	30	27	27
N.S.	1	1.00	1.00	0.80	0.77	0.77	0.89	0.86	0.77	0.77
time (sec)	N/A	0.194	0.005	0.046	0.026	0.076	0.059	0.121	0.154	0.028

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	32	25	24	34	27	34	47	26
N.S.	1	1.00	0.94	0.74	0.71	1.00	0.79	1.00	1.38	0.76
time (sec)	N/A	0.197	0.010	0.085	0.033	0.093	0.059	0.116	0.154	0.020

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	43	42	35	33	44	36	33	63	39
N.S.	1	0.96	0.93	0.78	0.73	0.98	0.80	0.73	1.40	0.87
time (sec)	N/A	0.171	0.013	0.085	0.106	0.074	0.054	0.119	0.150	0.072

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	49	38	37	37	46	37	37	41
N.S.	1	1.00	1.00	0.78	0.76	0.76	0.94	0.76	0.76	0.84
time (sec)	N/A	0.320	0.009	0.465	0.101	0.073	0.096	0.116	0.148	0.104

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	29	20	19	19	24	22	19	19
N.S.	1	1.00	1.00	0.69	0.66	0.66	0.83	0.76	0.66	0.66
time (sec)	N/A	0.194	0.005	0.086	0.026	0.077	0.059	0.123	0.160	0.114

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	52	43	34	33	33	41	34	32	50
N.S.	1	1.13	0.93	0.74	0.72	0.72	0.89	0.74	0.70	1.09
time (sec)	N/A	0.221	0.012	0.082	0.120	0.072	0.062	0.119	0.149	0.053

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	14	15	14	17	10	15	23	14
N.S.	1	1.00	0.88	0.94	0.88	1.06	0.62	0.94	1.44	0.88
time (sec)	N/A	0.169	0.006	0.033	0.038	0.080	0.037	0.116	0.147	0.026

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	14	12	17	17	14	12	20	12
N.S.	1	1.00	0.78	0.67	0.94	0.94	0.78	0.67	1.11	0.67
time (sec)	N/A	0.143	0.003	0.024	0.029	0.066	0.033	0.119	0.146	0.065

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	16	23	23	17	30	20	15
N.S.	1	1.00	1.00	1.07	1.53	1.53	1.13	2.00	1.33	1.00
time (sec)	N/A	0.175	0.008	0.078	0.025	0.062	0.044	0.117	0.159	0.076

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	31	27	28	28	3	29	27	57
N.S.	1	1.00	1.00	0.87	0.90	0.90	0.10	0.94	0.87	1.84
time (sec)	N/A	0.189	0.011	0.044	0.101	0.070	0.063	0.115	0.154	0.203

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	25	18	19	19	19	22	19	19
N.S.	1	1.00	1.00	0.72	0.76	0.76	0.76	0.88	0.76	0.76
time (sec)	N/A	0.205	0.004	0.040	0.028	0.073	0.061	0.125	0.157	0.067

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	30	29	25	24	36	20	26	44	22
N.S.	1	1.36	1.32	1.14	1.09	1.64	0.91	1.18	2.00	1.00
time (sec)	N/A	0.183	0.011	0.040	0.028	0.067	0.046	0.114	0.148	0.024

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	23	18	17	17	17	18	17	21
N.S.	1	1.00	1.00	0.78	0.74	0.74	0.74	0.78	0.74	0.91
time (sec)	N/A	0.199	0.004	0.084	0.101	0.079	0.057	0.121	0.142	0.080

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	103	103	93	72	77	136	88	74	204	96
N.S.	1	1.00	0.90	0.70	0.75	1.32	0.85	0.72	1.98	0.93
time (sec)	N/A	0.549	0.029	0.222	0.101	0.078	0.248	0.129	0.143	0.078

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	30	27	25	36	24	25	49	32
N.S.	1	1.00	0.88	0.79	0.74	1.06	0.71	0.74	1.44	0.94
time (sec)	N/A	0.172	0.008	0.089	0.106	0.065	0.065	0.121	0.169	0.021

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	33	33	28	29	44	27	30	58	33
N.S.	1	0.87	0.87	0.74	0.76	1.16	0.71	0.79	1.53	0.87
time (sec)	N/A	0.205	0.012	0.100	0.102	0.069	0.060	0.120	0.153	0.068

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	25	20	23	19	17	26	23	19
N.S.	1	1.00	1.00	0.80	0.92	0.76	0.68	1.04	0.92	0.76
time (sec)	N/A	0.209	0.005	0.088	0.026	0.064	0.038	0.120	0.156	0.018

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	36	32	31	31	36	31	30	56
N.S.	1	1.00	1.00	0.89	0.86	0.86	1.00	0.86	0.83	1.56
time (sec)	N/A	0.278	0.010	0.093	0.109	0.077	0.097	0.126	0.157	0.061

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	29	22	21	33	22	21	40	23
N.S.	1	1.00	1.00	0.76	0.72	1.14	0.76	0.72	1.38	0.79
time (sec)	N/A	0.262	0.011	0.108	0.104	0.077	0.076	0.119	0.172	0.025

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	46	34	35	39	46	36	38	49
N.S.	1	1.00	1.00	0.74	0.76	0.85	1.00	0.78	0.83	1.07
time (sec)	N/A	0.231	0.016	0.045	0.109	0.082	0.070	0.125	0.171	0.063

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	26	19	18	18	17	20	18	14
N.S.	1	1.00	1.00	0.73	0.69	0.69	0.65	0.77	0.69	0.54
time (sec)	N/A	0.173	0.004	0.100	0.026	0.071	0.035	0.121	0.156	0.026

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	16	15	15	15	18	15	15
N.S.	1	1.00	1.00	0.94	0.88	0.88	0.88	1.06	0.88	0.88
time (sec)	N/A	0.205	0.004	0.044	0.031	0.077	0.056	0.114	0.154	0.087

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	15	14	18	14	16	18	14
N.S.	1	1.00	1.00	1.07	1.00	1.29	1.00	1.14	1.29	1.00
time (sec)	N/A	0.176	0.003	0.084	0.023	0.078	0.046	0.124	0.153	0.041

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	25	18	17	17	17	20	17	17
N.S.	1	1.00	1.32	0.95	0.89	0.89	0.89	1.05	0.89	0.89
time (sec)	N/A	0.178	0.005	0.039	0.024	0.078	0.053	0.120	0.153	0.072

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	23	18	17	17	19	17	17	33
N.S.	1	1.00	1.00	0.78	0.74	0.74	0.83	0.74	0.74	1.43
time (sec)	N/A	0.190	0.006	0.053	0.110	0.078	0.080	0.121	0.156	0.162

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	57	51	50	50	68	53	49	58
N.S.	1	1.00	0.90	0.81	0.79	0.79	1.08	0.84	0.78	0.92
time (sec)	N/A	0.250	0.017	0.066	0.120	0.082	0.169	0.124	0.153	0.140

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	86	67	54	59	103	65	59	169	71
N.S.	1	1.00	0.78	0.63	0.69	1.20	0.76	0.69	1.97	0.83
time (sec)	N/A	0.260	0.029	0.072	0.105	0.080	0.095	0.125	0.157	0.071

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	13	19	12	13	19	13
N.S.	1	1.00	1.00	0.82	0.76	1.12	0.71	0.76	1.12	0.76
time (sec)	N/A	0.157	0.005	0.095	0.100	0.073	0.040	0.117	0.151	0.018

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	19	18	17	17	15	17	17	17
N.S.	1	1.00	1.00	0.95	0.89	0.89	0.79	0.89	0.89	0.89
time (sec)	N/A	0.192	0.004	0.082	0.105	0.073	0.041	0.132	0.150	0.078

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	9	9	9	8	7	7	7	28	7	19
N.S.	1	1.00	1.00	0.89	0.78	0.78	0.78	3.11	0.78	2.11
time (sec)	N/A	0.222	0.004	0.095	0.102	0.077	0.059	0.125	0.156	0.025

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	12	13	12	12	12	12	12	12
N.S.	1	1.00	1.00	1.08	1.00	1.00	1.00	1.00	1.00	1.00
time (sec)	N/A	0.160	0.004	0.149	0.028	0.064	0.041	0.123	0.152	0.017

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	58	41	51	66	99	55	76	48
N.S.	1	1.00	0.89	0.63	0.78	1.02	1.52	0.85	1.17	0.74
time (sec)	N/A	0.240	0.021	0.132	0.103	0.081	0.239	0.117	0.141	0.060

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	28	29	28	46	24	28	71	28
N.S.	1	1.00	1.00	1.04	1.00	1.64	0.86	1.00	2.54	1.00
time (sec)	N/A	0.200	0.006	0.134	0.105	0.072	0.049	0.121	0.154	0.070

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	32	27	26	26	29	26	26	26
N.S.	1	1.00	1.00	0.84	0.81	0.81	0.91	0.81	0.81	0.81
time (sec)	N/A	0.206	0.017	0.122	0.104	0.068	0.042	0.126	0.157	0.014

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	47	39	24	23	23	26	27	23	23
N.S.	1	1.52	1.26	0.77	0.74	0.74	0.84	0.87	0.74	0.74
time (sec)	N/A	0.262	0.005	0.046	0.024	0.083	0.093	0.125	0.155	0.035

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	24	23	22	26	19	23	26	30
N.S.	1	1.00	1.00	0.96	0.92	1.08	0.79	0.96	1.08	1.25
time (sec)	N/A	0.210	0.006	0.100	0.107	0.075	0.065	0.120	0.159	0.027

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	23	23	19	24	35	20	19	40	23
N.S.	1	0.82	0.82	0.68	0.86	1.25	0.71	0.68	1.43	0.82
time (sec)	N/A	0.185	0.007	0.080	0.101	0.064	0.060	0.118	0.156	0.119

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	23	19	24	35	20	19	40	23
N.S.	1	1.00	1.00	0.83	1.04	1.52	0.87	0.83	1.74	1.00
time (sec)	N/A	0.214	0.004	0.050	0.101	0.064	0.054	0.127	0.147	0.031

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	13	14	13	15	10	13	15	13
N.S.	1	1.00	1.00	1.08	1.00	1.15	0.77	1.00	1.15	1.00
time (sec)	N/A	0.208	0.005	0.043	0.027	0.064	0.039	0.118	0.159	0.039

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [72] had the largest ratio of [.363636000000000015]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	4	4	1.12	20	0.200
2	A	3	3	1.00	20	0.150
3	A	3	3	1.00	20	0.150
4	A	2	2	1.00	16	0.125
5	A	2	2	1.00	22	0.091
6	A	2	2	1.00	21	0.095
7	A	3	3	1.00	26	0.115
8	A	2	2	1.00	20	0.100
9	A	2	2	1.00	11	0.182
10	A	3	3	1.00	22	0.136
11	A	2	2	1.00	21	0.095
12	A	3	3	1.00	25	0.120
13	A	4	4	1.00	22	0.182
14	A	6	6	1.00	31	0.194
15	A	3	3	1.00	21	0.143
16	A	3	3	1.00	33	0.091
17	A	4	4	0.97	14	0.286
18	A	3	3	1.00	33	0.091
19	A	2	2	1.00	29	0.069
20	A	2	2	1.00	44	0.045
21	A	2	2	1.00	15	0.133

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	2	2	1.00	15	0.133
23	A	3	3	1.00	18	0.167
24	A	2	2	1.00	20	0.100
25	A	2	2	1.00	26	0.077
26	A	2	2	1.00	13	0.154
27	A	3	3	1.00	18	0.167
28	A	2	2	1.00	26	0.077
29	A	6	6	1.00	19	0.316
30	A	6	6	1.00	24	0.250
31	A	7	6	1.06	20	0.300
32	A	7	6	1.22	18	0.333
33	A	2	2	1.00	19	0.105
34	A	2	2	1.00	13	0.154
35	A	2	2	1.00	22	0.091
36	A	2	2	1.00	24	0.083
37	A	2	2	1.00	29	0.069
38	A	2	2	1.00	30	0.067
39	A	2	2	1.00	19	0.105
40	A	5	5	0.96	16	0.312
41	A	2	2	1.00	36	0.056
42	A	2	2	1.00	21	0.095
43	A	4	4	1.13	16	0.250
44	A	2	2	1.00	24	0.083
45	A	2	2	1.00	21	0.095
46	A	2	2	1.00	24	0.083
47	A	2	2	1.00	26	0.077
48	A	3	3	1.00	25	0.120
49	A	2	2	1.36	29	0.069
50	A	3	3	1.00	20	0.150
51	A	2	2	1.00	32	0.062
52	A	5	5	1.00	23	0.217
53	A	4	4	0.87	26	0.154

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
54	A	3	3	1.00	26	0.115
55	A	2	2	1.00	25	0.080
56	A	2	2	1.00	23	0.087
57	A	3	3	1.00	23	0.130
58	A	2	2	1.00	20	0.100
59	A	3	3	1.00	25	0.120
60	A	3	3	1.00	22	0.136
61	A	2	2	1.00	24	0.083
62	A	5	5	1.00	24	0.208
63	A	2	2	1.00	43	0.047
64	A	2	2	1.00	50	0.040
65	A	2	2	1.00	16	0.125
66	A	3	3	1.00	15	0.200
67	A	4	4	1.00	20	0.200
68	A	2	2	1.00	24	0.083
69	A	2	2	1.00	27	0.074
70	A	8	7	1.00	26	0.269
71	A	3	3	1.00	16	0.188
72	A	9	8	1.52	22	0.364
73	A	2	2	1.00	24	0.083
74	A	5	5	0.82	26	0.192
75	A	6	6	1.00	36	0.167
76	A	3	3	1.00	26	0.115

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \frac{1-3x^4}{(-2+x)(1+x^2)^2} dx$	56
3.2	$\int \frac{-9-9x+2x^2}{-9x+x^3} dx$	62
3.3	$\int \frac{1+2x^2+x^5}{-x+x^3} dx$	67
3.4	$\int \frac{3+2x^2}{(-1+x)^2x} dx$	72
3.5	$\int \frac{-1+2x^2}{(-1+4x)(1+x^2)} dx$	77
3.6	$\int \frac{-3+2x-3x^2+x^3}{1+x^2} dx$	82
3.7	$\int \frac{x+10x^2+6x^3+x^4}{10+6x+x^2} dx$	87
3.8	$\int \frac{1}{-18+27x-7x^2-3x^3+x^4} dx$	92
3.9	$\int \frac{1+x^3}{-2+x} dx$	97
3.10	$\int \frac{3x-4x^2+3x^3}{1+x^2} dx$	102
3.11	$\int \frac{5+3x}{1-x-x^2+x^3} dx$	107
3.12	$\int \frac{-1-x-x^3+x^4}{-x^2+x^3} dx$	112
3.13	$\int \frac{2+x+x^2+x^3}{2+3x^2+x^4} dx$	117
3.14	$\int \frac{-4+8x-4x^2+4x^3-x^4+x^5}{(2+x^2)^3} dx$	122
3.15	$\int \frac{-1-3x+x^2}{-2x+x^2+x^3} dx$	128
3.16	$\int \frac{3-x+3x^2-2x^3+x^4}{3x-2x^2+x^3} dx$	133
3.17	$\int \frac{-1+x+x^3}{(1+x^2)^2} dx$	138
3.18	$\int \frac{1+2x-x^2+8x^3+x^4}{(x+x^2)(1+x^3)} dx$	143
3.19	$\int \frac{15-5x+x^2+x^3}{(5+x^2)(3+2x+x^2)} dx$	149
3.20	$\int \frac{-3+25x+23x^2+32x^3+15x^4+7x^5+x^6}{(1+x^2)^2(2+x+x^2)^2} dx$	155
3.21	$\int \frac{1}{(1+x^2)(4+x^2)} dx$	161
3.22	$\int \frac{a+bx^3}{1+x^2} dx$	166
3.23	$\int \frac{x+x^2}{(4+x)(-4+x^2)} dx$	171
3.24	$\int \frac{4+x^2}{(1+x^2)(2+x^2)} dx$	176

3.25	$\int \frac{5-4x+3x^2+x^4}{(-1+x)^2(1+x^2)} dx$	181
3.26	$\int \frac{1+x^4}{2+x^2} dx$	186
3.27	$\int \frac{2+2x+x^4}{x^4+x^5} dx$	191
3.28	$\int \frac{-1-5x+2x^2}{2-x-2x^2+x^3} dx$	196
3.29	$\int \frac{2+x+x^3}{1+2x^2+x^4} dx$	201
3.30	$\int \frac{1+2x+x^2+x^3}{1+2x^2+x^4} dx$	207
3.31	$\int \frac{3+4x}{(1+x^2)(2+x^2)} dx$	213
3.32	$\int \frac{2+x}{(1+x^2)(4+x^2)} dx$	219
3.33	$\int \frac{2-x+x^3}{-7-6x+x^2} dx$	225
3.34	$\int \frac{-1+x^5}{-1+x^2} dx$	230
3.35	$\int \frac{5+2x-x^2+x^3}{1+x+x^2} dx$	235
3.36	$\int \frac{-3+x-2x^3+x^4}{10-8x+2x^2} dx$	240
3.37	$\int \frac{1+2x+3x^2+x^3}{(-3+x)(-2+x)(-1+x)} dx$	245
3.38	$\int \frac{2-7x+x^2-x^3+x^4}{-24-14x+x^2+x^3} dx$	250
3.39	$\int \frac{2+x^2}{(-1+x)^2x(1+x)} dx$	255
3.40	$\int \frac{3+x^2+x^3}{(2+x^2)^2} dx$	260
3.41	$\int \frac{-35+70x-4x^2+2x^3}{(26-10x+x^2)(17-2x+x^2)} dx$	266
3.42	$\int \frac{2+x^2}{(-5+x)(-3+x)(4+x)} dx$	272
3.43	$\int \frac{x^4}{(-1+x)(2+x^2)} dx$	277
3.44	$\int \frac{-1+7x+2x^2}{-1-x+x^2+x^3} dx$	283
3.45	$\int \frac{1+2x}{-1+3x-3x^2+x^3} dx$	288
3.46	$\int \frac{5-5x+7x^2+x^3}{(-1+x)^2(1+x)^3} dx$	293
3.47	$\int \frac{1+3x+3x^2}{1+2x+2x^2+x^3} dx$	298
3.48	$\int \frac{-1+2x+x^2}{-2x+3x^2+2x^3} dx$	303
3.49	$\int \frac{1+4x-2x^2+x^4}{1-x-x^2+x^3} dx$	308
3.50	$\int \frac{4-x+2x^2}{4x+x^3} dx$	313
3.51	$\int \frac{1+x^2+x^3}{(-1+x)x(1+x^2)^3(1+x+x^2)} dx$	318
3.52	$\int \frac{1-3x+2x^2-x^3}{(1+x^2)^2} dx$	325
3.53	$\int \frac{1-3x+2x^2-x^3}{x(1+x^2)^2} dx$	331
3.54	$\int \frac{1-x-x^2+x^3+x^4}{-x+x^3} dx$	337
3.55	$\int \frac{2-4x^2+x^3}{(1+x^2)(2+x^2)} dx$	342
3.56	$\int \frac{1+x^2+x^4}{(1+x^2)(4+x^2)^2} dx$	347
3.57	$\int \frac{1+x^2+x^3}{2x^2+x^3+x^4} dx$	352
3.58	$\int \frac{1-12x+x^2+x^3}{-12+x+x^2} dx$	357

3.59	$\int \frac{-3+5x+6x^2}{-3x+2x^2+x^3} dx$	362
3.60	$\int \frac{-2+3x+5x^2}{2x^2+x^3} dx$	367
3.61	$\int \frac{18-2x-4x^2}{-6+x+4x^2+x^3} dx$	372
3.62	$\int \frac{1+x-2x^2+x^3}{4+5x^2+x^4} dx$	377
3.63	$\int \frac{-32+5x-27x^2+4x^3}{-70-299x-286x^2+50x^3-13x^4+30x^5} dx$	383
3.64	$\int \frac{8-13x^2-7x^3+12x^5}{4-20x+41x^2-80x^3+116x^4-80x^5+100x^6} dx$	389
3.65	$\int \frac{9+x^4}{x^2(9+x^2)} dx$	395
3.66	$\int \frac{2x+x^4}{1+x^2} dx$	400
3.67	$\int \frac{-x+x^3}{(-1+x)^2(1+x^2)} dx$	405
3.68	$\int \frac{2+5x+3x^2+2x^3}{1+x+x^2} dx$	410
3.69	$\int \frac{3-4x-5x^2+3x^3}{x^3(-1+x+x^2)} dx$	415
3.70	$\int \frac{4+8x+5x^2+2x^3}{(2+2x+x^2)^2} dx$	421
3.71	$\int \frac{(-1+x)^4 x^4}{1+x^2} dx$	427
3.72	$\int \frac{-20x+4x^2}{9-10x^2+x^4} dx$	432
3.73	$\int \frac{-1+x+4x^3}{(-1+x)x^2(1+x^2)} dx$	438
3.74	$\int \frac{1-3x+2x^2-4x^3+x^4}{(1+x^2)^3} dx$	443
3.75	$\int \frac{1-3x+2x^2-4x^3+x^4}{1+3x^2+3x^4+x^6} dx$	448
3.76	$\int \frac{1+x+2x^2+2x^3}{x^2+x^3+x^4} dx$	454

3.1 $\int \frac{1-3x^4}{(-2+x)(1+x^2)^2} dx$

Optimal result	56
Mathematica [A] (verified)	56
Rubi [A] (verified)	57
Maple [A] (verified)	58
Fricas [A] (verification not implemented)	59
Sympy [A] (verification not implemented)	59
Maxima [A] (verification not implemented)	59
Giac [A] (verification not implemented)	60
Mupad [B] (verification not implemented)	60
Reduce [B] (verification not implemented)	61

Optimal result

Integrand size = 20, antiderivative size = 43

$$\int \frac{1-3x^4}{(-2+x)(1+x^2)^2} dx = -\frac{1-2x}{5(1+x^2)} - \frac{46 \arctan(x)}{25} - \frac{47}{25} \log(2-x) - \frac{14}{25} \log(1+x^2)$$

output `-(1-2*x)/(5*x^2+5)-46/25*arctan(x)-47/25*ln(2-x)-14/25*ln(x^2+1)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.33

$$\int \frac{1-3x^4}{(-2+x)(1+x^2)^2} dx = \frac{3+2(-2+x)}{5(5+4(-2+x)+(-2+x)^2)} - \frac{46 \arctan(x)}{25} - \frac{14}{25} \log(5+4(-2+x)+(-2+x)^2) - \frac{47}{25} \log(-2+x)$$

input `Integrate[(1 - 3*x^4)/((-2 + x)*(1 + x^2)^2), x]`

output `(3 + 2*(-2 + x))/(5*(5 + 4*(-2 + x) + (-2 + x)^2)) - (46*ArcTan[x])/25 - (14*Log[5 + 4*(-2 + x) + (-2 + x)^2])/25 - (47*Log[-2 + x])/25`

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.12, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2178, 27, 2160, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1 - 3x^4}{(x - 2)(x^2 + 1)^2} dx \\
 & \quad \downarrow \text{2178} \\
 & -\frac{1}{2} \int \frac{2(-15x^2 + 2x + 9)}{5(2 - x)(x^2 + 1)} dx - \frac{1 - 2x}{5(x^2 + 1)} \\
 & \quad \downarrow \text{27} \\
 & -\frac{1}{5} \int \frac{-15x^2 + 2x + 9}{(2 - x)(x^2 + 1)} dx - \frac{1 - 2x}{5(x^2 + 1)} \\
 & \quad \downarrow \text{2160} \\
 & -\frac{1}{5} \int \left(\frac{2(14x + 23)}{5(x^2 + 1)} + \frac{47}{5(x - 2)} \right) dx - \frac{1 - 2x}{5(x^2 + 1)} \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{5} \left(-\frac{46 \arctan(x)}{5} - \frac{14}{5} \log(x^2 + 1) - \frac{47}{5} \log(2 - x) \right) - \frac{1 - 2x}{5(x^2 + 1)}
 \end{aligned}$$

input `Int[(1 - 3*x^4)/((-2 + x)*(1 + x^2)^2), x]`

output `-1/5*(1 - 2*x)/(1 + x^2) + ((-46*ArcTan[x])/5 - (47*Log[2 - x])/5 - (14*Log[1 + x^2])/5)/5`

Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2160 `Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*Pq*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

rule 2178 `Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{Qx = PolynomialQuotient[(d + e*x)^m*Pq, a + b*x^2, x], R = Coeff[PolynomialRemainder[(d + e*x)^m*Pq, a + b*x^2, x], x, 0], S = Coeff[PolynomialRemainder[(d + e*x)^m*Pq, a + b*x^2, x], x, 1]}, Simp[(a*S - b*R*x)*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] + Simp[1/(2*a*b*(p + 1)) Int[(d + e*x)^m*(a + b*x^2)^(p + 1)*ExpandToSum[(2*a*b*(p + 1)*Qx)/(d + e*x)^m + (b*R*(2*p + 3))/(d + e*x)^m, x], x], x] /; FreeQ[{a, b, d, e}, x] && PolyQ[Pq, x] && NeQ[b*d^2 + a*e^2, 0] && LtQ[p, -1] && ILtQ[m, 0]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.77

method	result
risch	$\frac{2x - \frac{1}{5}}{x^2 + 1} - \frac{14 \ln(x^2 + 1)}{25} - \frac{46 \arctan(x)}{25} - \frac{47 \ln(-2 + x)}{25}$
default	$-\frac{2(-5x + \frac{5}{2})}{25(x^2 + 1)} - \frac{14 \ln(x^2 + 1)}{25} - \frac{46 \arctan(x)}{25} - \frac{47 \ln(-2 + x)}{25}$
paralelrisch	$-\frac{-23i \ln(x-i)x^2 + 23i \ln(x+i)x^2 + 47 \ln(-2+x)x^2 + 14 \ln(x-i)x^2 + 14 \ln(x+i)x^2 + 5 - 23i \ln(x-i) + 23i \ln(x+i) + 47 \ln(-2+x)}{25(x^2 + 1)}$

input `int((-3*x^4+1)/(-2+x)/(x^2+1)^2,x,method=_RETURNVERBOSE)`

output `(2/5*x-1/5)/(x^2+1)-14/25*ln(x^2+1)-46/25*arctan(x)-47/25*ln(-2+x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.09

$$\int \frac{1 - 3x^4}{(-2 + x)(1 + x^2)^2} dx$$

$$= -\frac{46(x^2 + 1) \arctan(x) + 14(x^2 + 1) \log(x^2 + 1) + 47(x^2 + 1) \log(x - 2) - 10x + 5}{25(x^2 + 1)}$$

input `integrate((-3*x^4+1)/(-2+x)/(x^2+1)^2,x, algorithm="fricas")`output `-1/25*(46*(x^2 + 1)*arctan(x) + 14*(x^2 + 1)*log(x^2 + 1) + 47*(x^2 + 1)*log(x - 2) - 10*x + 5)/(x^2 + 1)`**Sympy [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.86

$$\int \frac{1 - 3x^4}{(-2 + x)(1 + x^2)^2} dx = -\frac{1 - 2x}{5x^2 + 5} - \frac{47 \log(x - 2)}{25} - \frac{14 \log(x^2 + 1)}{25} - \frac{46 \operatorname{atan}(x)}{25}$$

input `integrate((-3*x**4+1)/(-2+x)/(x**2+1)**2,x)`output `-(1 - 2*x)/(5*x**2 + 5) - 47*log(x - 2)/25 - 14*log(x**2 + 1)/25 - 46*atan(x)/25`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.77

$$\int \frac{1 - 3x^4}{(-2 + x)(1 + x^2)^2} dx = \frac{2x - 1}{5(x^2 + 1)} - \frac{46}{25} \arctan(x) - \frac{14}{25} \log(x^2 + 1) - \frac{47}{25} \log(x - 2)$$

input `integrate((-3*x^4+1)/(-2+x)/(x^2+1)^2,x, algorithm="maxima")`

output $\frac{1}{5} \cdot \frac{2x - 1}{x^2 + 1} - \frac{46}{25} \arctan(x) - \frac{14}{25} \log(x^2 + 1) - \frac{47}{25} \log(x - 2)$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.79

$$\int \frac{1 - 3x^4}{(-2 + x)(1 + x^2)^2} dx = \frac{2x - 1}{5(x^2 + 1)} - \frac{46}{25} \arctan(x) - \frac{14}{25} \log(x^2 + 1) - \frac{47}{25} \log(|x - 2|)$$

input `integrate((-3*x^4+1)/(-2+x)/(x^2+1)^2,x, algorithm="giac")`

output $\frac{1}{5} \cdot \frac{2x - 1}{x^2 + 1} - \frac{46}{25} \arctan(x) - \frac{14}{25} \log(x^2 + 1) - \frac{47}{25} \log(\text{abs}(x - 2))$

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.88

$$\int \frac{1 - 3x^4}{(-2 + x)(1 + x^2)^2} dx = \frac{\frac{2x}{5} - \frac{1}{5}}{x^2 + 1} - \frac{47 \ln(x - 2)}{25} + \ln(x - i) \left(-\frac{14}{25} + \frac{23}{25}i \right) + \ln(x + i) \left(-\frac{14}{25} - \frac{23}{25}i \right)$$

input `int(-(3*x^4 - 1)/((x^2 + 1)^2*(x - 2)),x)`

output $\left(\frac{2x}{5} - \frac{1}{5}\right)/(x^2 + 1) - \log(x - 1i) \cdot (14/25 - 23i/25) - \log(x + 1i) \cdot (14/25 + 23i/25) - (47 \cdot \log(x - 2))/25$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.49

$$\int \frac{1 - 3x^4}{(-2 + x)(1 + x^2)^2} dx$$

$$= \frac{-46 \operatorname{atan}(x) x^2 - 46 \operatorname{atan}(x) - 14 \log(x^2 + 1) x^2 - 14 \log(x^2 + 1) - 47 \log(x - 2) x^2 - 47 \log(x - 2) + 5x^2 + 10x}{25x^2 + 25}$$

input `int((-3*x^4+1)/(-2+x)/(x^2+1)^2,x)`output `(- 46*atan(x)*x**2 - 46*atan(x) - 14*log(x**2 + 1)*x**2 - 14*log(x**2 + 1) - 47*log(x - 2)*x**2 - 47*log(x - 2) + 5*x**2 + 10*x)/(25*(x**2 + 1))`

3.2 $\int \frac{-9-9x+2x^2}{-9x+x^3} dx$

Optimal result	62
Mathematica [A] (verified)	62
Rubi [A] (verified)	63
Maple [A] (verified)	64
Fricas [A] (verification not implemented)	64
Sympy [A] (verification not implemented)	65
Maxima [A] (verification not implemented)	65
Giac [A] (verification not implemented)	65
Mupad [B] (verification not implemented)	66
Reduce [B] (verification not implemented)	66

Optimal result

Integrand size = 20, antiderivative size = 17

$$\int \frac{-9 - 9x + 2x^2}{-9x + x^3} dx = -\log(3 - x) + \log(x) + 2\log(3 + x)$$

output

```
-ln(3-x)+ln(x)+2*ln(3+x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{-9 - 9x + 2x^2}{-9x + x^3} dx = -\log(3 - x) + \log(x) + 2\log(3 + x)$$

input

```
Integrate[(-9 - 9*x + 2*x^2)/(-9*x + x^3), x]
```

output

```
-Log[3 - x] + Log[x] + 2*Log[3 + x]
```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {2026, 2333, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{2x^2 - 9x - 9}{x^3 - 9x} dx \\ & \quad \downarrow \text{2026} \\ & \int \frac{2x^2 - 9x - 9}{x(x^2 - 9)} dx \\ & \quad \downarrow \text{2333} \\ & \int \left(\frac{1}{x} + \frac{2}{x+3} + \frac{1}{3-x} \right) dx \\ & \quad \downarrow \text{2009} \\ & -\log(3-x) + \log(x) + 2\log(x+3) \end{aligned}$$

input `Int[(-9 - 9*x + 2*x^2)/(-9*x + x^3),x]`

output `-Log[3 - x] + Log[x] + 2*Log[3 + x]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2333

```
Int[(Pq_)*((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[
ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x]
&& PolyQ[Pq, x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

method	result	size
default	$-\ln(-3+x) + 2\ln(3+x) + \ln(x)$	16
norman	$-\ln(-3+x) + 2\ln(3+x) + \ln(x)$	16
risch	$-\ln(-3+x) + 2\ln(3+x) + \ln(x)$	16
parallelrisc	$-\ln(-3+x) + 2\ln(3+x) + \ln(x)$	16
meijerg	$\ln(x) - \ln(3) + \frac{i\pi}{2} + \frac{\ln\left(1-\frac{x^2}{9}\right)}{2} + 3 \operatorname{arctanh}\left(\frac{x}{3}\right)$	28

input

```
int((2*x^2-9*x-9)/(x^3-9*x),x,method=_RETURNVERBOSE)
```

output

```
-ln(-3+x)+2*ln(3+x)+ln(x)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{-9 - 9x + 2x^2}{-9x + x^3} dx = 2 \log(x + 3) - \log(x - 3) + \log(x)$$

input

```
integrate((2*x^2-9*x-9)/(x^3-9*x),x, algorithm="fricas")
```

output

```
2*log(x + 3) - log(x - 3) + log(x)
```

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \frac{-9 - 9x + 2x^2}{-9x + x^3} dx = \log(x) - \log(x - 3) + 2 \log(x + 3)$$

input `integrate((2*x**2-9*x-9)/(x**3-9*x),x)`output `log(x) - log(x - 3) + 2*log(x + 3)`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{-9 - 9x + 2x^2}{-9x + x^3} dx = 2 \log(x + 3) - \log(x - 3) + \log(x)$$

input `integrate((2*x^2-9*x-9)/(x^3-9*x),x, algorithm="maxima")`output `2*log(x + 3) - log(x - 3) + log(x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.06

$$\int \frac{-9 - 9x + 2x^2}{-9x + x^3} dx = 2 \log(|x + 3|) - \log(|x - 3|) + \log(|x|)$$

input `integrate((2*x^2-9*x-9)/(x^3-9*x),x, algorithm="giac")`output `2*log(abs(x + 3)) - log(abs(x - 3)) + log(abs(x))`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.24

$$\int \frac{-9 - 9x + 2x^2}{-9x + x^3} dx = 2 \ln(x + 3) - 2 \operatorname{atanh}\left(\frac{1296}{18x + 162} - 7\right)$$

input `int((9*x - 2*x^2 + 9)/(9*x - x^3),x)`

output `2*log(x + 3) - 2*atanh(1296/(18*x + 162) - 7)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{-9 - 9x + 2x^2}{-9x + x^3} dx = -\log(x - 3) + 2 \log(x + 3) + \log(x)$$

input `int((2*x^2-9*x-9)/(x^3-9*x),x)`

output `- log(x - 3) + 2*log(x + 3) + log(x)`

3.3 $\int \frac{1+2x^2+x^5}{-x+x^3} dx$

Optimal result	67
Mathematica [A] (verified)	67
Rubi [A] (verified)	68
Maple [A] (verified)	69
Fricas [A] (verification not implemented)	69
Sympy [A] (verification not implemented)	70
Maxima [A] (verification not implemented)	70
Giac [A] (verification not implemented)	70
Mupad [B] (verification not implemented)	71
Reduce [B] (verification not implemented)	71

Optimal result

Integrand size = 20, antiderivative size = 25

$$\int \frac{1 + 2x^2 + x^5}{-x + x^3} dx = x + \frac{x^3}{3} + 2 \log(1 - x) - \log(x) + \log(1 + x)$$

output

```
x+1/3*x^3+2*ln(1-x)-ln(x)+ln(1+x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{1 + 2x^2 + x^5}{-x + x^3} dx = x + \frac{x^3}{3} + 2 \log(1 - x) - \log(x) + \log(1 + x)$$

input

```
Integrate[(1 + 2*x^2 + x^5)/(-x + x^3),x]
```

output

```
x + x^3/3 + 2*Log[1 - x] - Log[x] + Log[1 + x]
```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {2026, 2333, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^5 + 2x^2 + 1}{x^3 - x} dx \\ & \quad \downarrow \text{2026} \\ & \int \frac{x^5 + 2x^2 + 1}{x(x^2 - 1)} dx \\ & \quad \downarrow \text{2333} \\ & \int \left(x^2 + \frac{2}{x-1} + \frac{1}{x+1} - \frac{1}{x} + 1 \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{x^3}{3} + x + 2 \log(1-x) - \log(x) + \log(x+1) \end{aligned}$$

input `Int[(1 + 2*x^2 + x^5)/(-x + x^3),x]`

output `x + x^3/3 + 2*Log[1 - x] - Log[x] + Log[1 + x]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2333

```
Int[(Pq_)*((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[
ExpandIntegrand[(c*x)^m*Pq*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x]
&& PolyQ[Pq, x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

method	result	size
default	$\frac{x^3}{3} + x - \ln(x) + 2 \ln(-1 + x) + \ln(1 + x)$	22
norman	$\frac{x^3}{3} + x - \ln(x) + 2 \ln(-1 + x) + \ln(1 + x)$	22
risch	$\frac{x^3}{3} + x - \ln(x) + 2 \ln(-1 + x) + \ln(1 + x)$	22
parallelrisc	$\frac{x^3}{3} + x - \ln(x) + 2 \ln(-1 + x) + \ln(1 + x)$	22
meijerg	$-\ln(x) - \frac{i\pi}{2} + \frac{3 \ln(-x^2+1)}{2} + \frac{i \left(-\frac{2ix(5x^2+15)}{15} + 2i \operatorname{arctanh}(x) \right)}{2}$	40

input

```
int((x^5+2*x^2+1)/(x^3-x),x,method=_RETURNVERBOSE)
```

output

```
1/3*x^3+x-ln(x)+2*ln(-1+x)+ln(1+x)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int \frac{1 + 2x^2 + x^5}{-x + x^3} dx = \frac{1}{3}x^3 + x + \log(x + 1) + 2 \log(x - 1) - \log(x)$$

input

```
integrate((x^5+2*x^2+1)/(x^3-x),x, algorithm="fricas")
```

output

```
1/3*x^3 + x + log(x + 1) + 2*log(x - 1) - log(x)
```

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \frac{1 + 2x^2 + x^5}{-x + x^3} dx = \frac{x^3}{3} + x - \log(x) + 2 \log(x - 1) + \log(x + 1)$$

input `integrate((x**5+2*x**2+1)/(x**3-x),x)`output `x**3/3 + x - log(x) + 2*log(x - 1) + log(x + 1)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int \frac{1 + 2x^2 + x^5}{-x + x^3} dx = \frac{1}{3} x^3 + x + \log(x + 1) + 2 \log(x - 1) - \log(x)$$

input `integrate((x^5+2*x^2+1)/(x^3-x),x, algorithm="maxima")`output `1/3*x^3 + x + log(x + 1) + 2*log(x - 1) - log(x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{1 + 2x^2 + x^5}{-x + x^3} dx = \frac{1}{3} x^3 + x + \log(|x + 1|) + 2 \log(|x - 1|) - \log(|x|)$$

input `integrate((x^5+2*x^2+1)/(x^3-x),x, algorithm="giac")`output `1/3*x^3 + x + log(abs(x + 1)) + 2*log(abs(x - 1)) - log(abs(x))`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.20

$$\int \frac{1 + 2x^2 + x^5}{-x + x^3} dx = x + 2 \ln(x - 1) + \frac{x^3}{3} + \operatorname{atan}\left(\frac{48i}{11(22x - 2)} + \frac{13i}{11}\right) 2i$$

input `int(-(2*x^2 + x^5 + 1)/(x - x^3),x)`

output `x + 2*log(x - 1) + atan(48i/(11*(22*x - 2)) + 13i/11)*2i + x^3/3`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int \frac{1 + 2x^2 + x^5}{-x + x^3} dx = 2 \log(x - 1) + \log(x + 1) - \log(x) + \frac{x^3}{3} + x$$

input `int((x^5+2*x^2+1)/(x^3-x),x)`

output `(6*log(x - 1) + 3*log(x + 1) - 3*log(x) + x**3 + 3*x)/3`

3.4 $\int \frac{3+2x^2}{(-1+x)^2x} dx$

Optimal result	72
Mathematica [A] (verified)	72
Rubi [A] (verified)	73
Maple [A] (verified)	74
Fricas [A] (verification not implemented)	74
Sympy [A] (verification not implemented)	75
Maxima [A] (verification not implemented)	75
Giac [A] (verification not implemented)	75
Mupad [B] (verification not implemented)	76
Reduce [B] (verification not implemented)	76

Optimal result

Integrand size = 16, antiderivative size = 22

$$\int \frac{3+2x^2}{(-1+x)^2x} dx = \frac{5}{1-x} - \log(1-x) + 3 \log(x)$$

output `5/(1-x)-ln(1-x)+3*ln(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{3+2x^2}{(-1+x)^2x} dx = -\frac{5}{-1+x} - \log(1-x) + 3 \log(x)$$

input `Integrate[(3 + 2*x^2)/((-1 + x)^2*x), x]`

output `-5/(-1 + x) - Log[1 - x] + 3*Log[x]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {522, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{2x^2 + 3}{(x-1)^2 x} dx$$

$$\downarrow 522$$

$$\int \left(\frac{3}{x} + \frac{5}{(x-1)^2} + \frac{1}{1-x} \right) dx$$

$$\downarrow 2009$$

$$\frac{5}{1-x} - \log(1-x) + 3 \log(x)$$

input `Int[(3 + 2*x^2)/((-1 + x)^2*x),x]`

output `5/(1 - x) - Log[1 - x] + 3*Log[x]`

Defintions of rubi rules used

rule 522 `Int[((e_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] :> Int[ExpandIntegrand[(e*x)^m*(c + d*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.86

method	result	size
default	$3 \ln(x) - \frac{5}{-1+x} - \ln(-1+x)$	19
norman	$3 \ln(x) - \frac{5}{-1+x} - \ln(-1+x)$	19
risch	$3 \ln(x) - \frac{5}{-1+x} - \ln(-1+x)$	19
parallelrisch	$\frac{3x \ln(x) - \ln(-1+x)x - 5 - 3 \ln(x) + \ln(-1+x)}{-1+x}$	29
meijerg	$\frac{2x}{1-x} - \ln(1-x) + 3 + 3 \ln(x) + 3i\pi + \frac{6x}{-2x+2}$	39

input `int((2*x^2+3)/(-1+x)^2/x,x,method=_RETURNVERBOSE)`output `3*ln(x)-5/(-1+x)-ln(-1+x)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{3 + 2x^2}{(-1+x)^2x} dx = -\frac{(x-1)\log(x-1) - 3(x-1)\log(x) + 5}{x-1}$$

input `integrate((2*x^2+3)/(-1+x)^2/x,x, algorithm="fricas")`output `-((x - 1)*log(x - 1) - 3*(x - 1)*log(x) + 5)/(x - 1)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.64

$$\int \frac{3 + 2x^2}{(-1 + x)^2 x} dx = 3 \log(x) - \log(x - 1) - \frac{5}{x - 1}$$

input `integrate((2*x**2+3)/(-1+x)**2/x,x)`output `3*log(x) - log(x - 1) - 5/(x - 1)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \frac{3 + 2x^2}{(-1 + x)^2 x} dx = -\frac{5}{x - 1} - \log(x - 1) + 3 \log(x)$$

input `integrate((2*x^2+3)/(-1+x)^2/x,x, algorithm="maxima")`output `-5/(x - 1) - log(x - 1) + 3*log(x)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.27

$$\int \frac{3 + 2x^2}{(-1 + x)^2 x} dx = -\frac{5}{x - 1} + 2 \log(|x - 1|) + 3 \log\left(\left|-\frac{1}{x - 1} - 1\right|\right)$$

input `integrate((2*x^2+3)/(-1+x)^2/x,x, algorithm="giac")`output `-5/(x - 1) + 2*log(abs(x - 1)) + 3*log(abs(-1/(x - 1) - 1))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \frac{3 + 2x^2}{(-1 + x)^2 x} dx = 3 \ln(x) - \ln(x - 1) - \frac{5}{x - 1}$$

input `int((2*x^2 + 3)/(x*(x - 1)^2),x)`

output `3*log(x) - log(x - 1) - 5/(x - 1)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.36

$$\int \frac{3 + 2x^2}{(-1 + x)^2 x} dx = \frac{-\log(x - 1)x + \log(x - 1) + 3 \log(x)x - 3 \log(x) - 5x}{x - 1}$$

input `int((2*x^2+3)/(-1+x)^2/x,x)`

output `(- log(x - 1)*x + log(x - 1) + 3*log(x)*x - 3*log(x) - 5*x)/(x - 1)`

3.5 $\int \frac{-1+2x^2}{(-1+4x)(1+x^2)} dx$

Optimal result	77
Mathematica [A] (verified)	77
Rubi [A] (verified)	78
Maple [A] (verified)	79
Fricas [A] (verification not implemented)	79
Sympy [A] (verification not implemented)	79
Maxima [A] (verification not implemented)	80
Giac [A] (verification not implemented)	80
Mupad [B] (verification not implemented)	80
Reduce [B] (verification not implemented)	81

Optimal result

Integrand size = 22, antiderivative size = 27

$$\int \frac{-1+2x^2}{(-1+4x)(1+x^2)} dx = \frac{3 \arctan(x)}{17} - \frac{7}{34} \log(1-4x) + \frac{6}{17} \log(1+x^2)$$

output `3/17*arctan(x)-7/34*ln(1-4*x)+6/17*ln(x^2+1)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.41

$$\int \frac{-1+2x^2}{(-1+4x)(1+x^2)} dx = \frac{3 \arctan(x)}{17} - \frac{7}{34} \log(-1+4x) + \frac{6}{17} \log(17+2(-1+4x)+(-1+4x)^2)$$

input `Integrate[(-1 + 2*x^2)/((-1 + 4*x)*(1 + x^2)),x]`

output `(3*ArcTan[x])/17 - (7*Log[-1 + 4*x])/34 + (6*Log[17 + 2*(-1 + 4*x) + (-1 + 4*x)^2])/17`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {2160, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{2x^2 - 1}{(4x - 1)(x^2 + 1)} dx$$

↓ 2160

$$\int \left(\frac{3(4x + 1)}{17(x^2 + 1)} - \frac{14}{17(4x - 1)} \right) dx$$

↓ 2009

$$\frac{3 \arctan(x)}{17} + \frac{6}{17} \log(x^2 + 1) - \frac{7}{34} \log(1 - 4x)$$

input

```
Int[(-1 + 2*x^2)/((-1 + 4*x)*(1 + x^2)),x]
```

output

```
(3*ArcTan[x])/17 - (7*Log[1 - 4*x])/34 + (6*Log[1 + x^2])/17
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2160

```
Int[(Pq_)*((d_) + (e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol]
:= Int[ExpandIntegrand[(d + e*x)^m*Pq*(a + b*x^2)^p, x], x] /; FreeQ[{a, b,
d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

method	result	size
default	$-\frac{7 \ln(-1+4x)}{34} + \frac{6 \ln(x^2+1)}{17} + \frac{3 \arctan(x)}{17}$	22
risch	$-\frac{7 \ln(-1+4x)}{34} + \frac{6 \ln(x^2+1)}{17} + \frac{3 \arctan(x)}{17}$	22
parallelrisc	$-\frac{7 \ln(x-\frac{1}{4})}{34} + \frac{6 \ln(x-i)}{17} - \frac{3i \ln(x-i)}{34} + \frac{6 \ln(x+i)}{17} + \frac{3i \ln(x+i)}{34}$	38

input `int((2*x^2-1)/(-1+4*x)/(x^2+1),x,method=_RETURNVERBOSE)`

output `-7/34*ln(-1+4*x)+6/17*ln(x^2+1)+3/17*arctan(x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int \frac{-1 + 2x^2}{(-1 + 4x)(1 + x^2)} dx = \frac{3}{17} \arctan(x) + \frac{6}{17} \log(x^2 + 1) - \frac{7}{34} \log(4x - 1)$$

input `integrate((2*x^2-1)/(-1+4*x)/(x^2+1),x, algorithm="fricas")`

output `3/17*arctan(x) + 6/17*log(x^2 + 1) - 7/34*log(4*x - 1)`

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \frac{-1 + 2x^2}{(-1 + 4x)(1 + x^2)} dx = -\frac{7 \log(x - \frac{1}{4})}{34} + \frac{6 \log(x^2 + 1)}{17} + \frac{3 \operatorname{atan}(x)}{17}$$

input `integrate((2*x**2-1)/(-1+4*x)/(x**2+1),x)`

output `-7*log(x - 1/4)/34 + 6*log(x**2 + 1)/17 + 3*atan(x)/17`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int \frac{-1 + 2x^2}{(-1 + 4x)(1 + x^2)} dx = \frac{3}{17} \arctan(x) + \frac{6}{17} \log(x^2 + 1) - \frac{7}{34} \log(4x - 1)$$

input `integrate((2*x^2-1)/(-1+4*x)/(x^2+1),x, algorithm="maxima")`

output `3/17*arctan(x) + 6/17*log(x^2 + 1) - 7/34*log(4*x - 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int \frac{-1 + 2x^2}{(-1 + 4x)(1 + x^2)} dx = \frac{3}{17} \arctan(x) + \frac{6}{17} \log(x^2 + 1) - \frac{7}{34} \log(|4x - 1|)$$

input `integrate((2*x^2-1)/(-1+4*x)/(x^2+1),x, algorithm="giac")`

output `3/17*arctan(x) + 6/17*log(x^2 + 1) - 7/34*log(abs(4*x - 1))`

Mupad [B] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.93

$$\int \frac{-1 + 2x^2}{(-1 + 4x)(1 + x^2)} dx = -\frac{7 \ln\left(x - \frac{1}{4}\right)}{34} + \ln(x - i) \left(\frac{6}{17} - \frac{3}{34}i\right) + \ln(x + i) \left(\frac{6}{17} + \frac{3}{34}i\right)$$

input `int((2*x^2 - 1)/((4*x - 1)*(x^2 + 1)),x)`

output `log(x - 1i)*(6/17 - 3i/34) - (7*log(x - 1/4))/34 + log(x + 1i)*(6/17 + 3i/34)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int \frac{-1 + 2x^2}{(-1 + 4x)(1 + x^2)} dx = \frac{3 \operatorname{atan}(x)}{17} + \frac{6 \log(x^2 + 1)}{17} - \frac{7 \log(4x - 1)}{34}$$

input `int((2*x^2-1)/(-1+4*x)/(x^2+1),x)`

output `(6*atan(x) + 12*log(x**2 + 1) - 7*log(4*x - 1))/34`

3.6 $\int \frac{-3+2x-3x^2+x^3}{1+x^2} dx$

Optimal result	82
Mathematica [A] (verified)	82
Rubi [A] (verified)	83
Maple [A] (verified)	84
Fricas [A] (verification not implemented)	84
Sympy [A] (verification not implemented)	85
Maxima [A] (verification not implemented)	85
Giac [A] (verification not implemented)	85
Mupad [B] (verification not implemented)	86
Reduce [B] (verification not implemented)	86

Optimal result

Integrand size = 21, antiderivative size = 21

$$\int \frac{-3 + 2x - 3x^2 + x^3}{1 + x^2} dx = -3x + \frac{x^2}{2} + \frac{1}{2} \log(1 + x^2)$$

output

```
-3*x+1/2*x^2+1/2*ln(x^2+1)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{-3 + 2x - 3x^2 + x^3}{1 + x^2} dx = -3x + \frac{x^2}{2} + \frac{1}{2} \log(1 + x^2)$$

input

```
Integrate[(-3 + 2*x - 3*x^2 + x^3)/(1 + x^2),x]
```

output

```
-3*x + x^2/2 + Log[1 + x^2]/2
```

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2341, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3 - 3x^2 + 2x - 3}{x^2 + 1} dx$$

↓ 2341

$$\int \left(\frac{x}{x^2 + 1} + x - 3 \right) dx$$

↓ 2009

$$\frac{x^2}{2} + \frac{1}{2} \log(x^2 + 1) - 3x$$

input `Int[(-3 + 2*x - 3*x^2 + x^3)/(1 + x^2),x]`

output `-3*x + x^2/2 + Log[1 + x^2]/2`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2341 `Int[(Pq_)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[Pq*(a + b*x^2)^p, x], x] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.86

method	result	size
default	$-3x + \frac{x^2}{2} + \frac{\ln(x^2+1)}{2}$	18
norman	$-3x + \frac{x^2}{2} + \frac{\ln(x^2+1)}{2}$	18
meijerg	$-3x + \frac{x^2}{2} + \frac{\ln(x^2+1)}{2}$	18
risch	$-3x + \frac{x^2}{2} + \frac{\ln(x^2+1)}{2}$	18
parallelrisc	$-3x + \frac{x^2}{2} + \frac{\ln(x^2+1)}{2}$	18

input `int((x^3-3*x^2+2*x-3)/(x^2+1),x,method=_RETURNVERBOSE)`

output `-3*x+1/2*x^2+1/2*ln(x^2+1)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{-3 + 2x - 3x^2 + x^3}{1 + x^2} dx = \frac{1}{2}x^2 - 3x + \frac{1}{2} \log(x^2 + 1)$$

input `integrate((x^3-3*x^2+2*x-3)/(x^2+1),x, algorithm="fricas")`

output `1/2*x^2 - 3*x + 1/2*log(x^2 + 1)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int \frac{-3 + 2x - 3x^2 + x^3}{1 + x^2} dx = \frac{x^2}{2} - 3x + \frac{\log(x^2 + 1)}{2}$$

input `integrate((x**3-3*x**2+2*x-3)/(x**2+1),x)`output `x**2/2 - 3*x + log(x**2 + 1)/2`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{-3 + 2x - 3x^2 + x^3}{1 + x^2} dx = \frac{1}{2}x^2 - 3x + \frac{1}{2}\log(x^2 + 1)$$

input `integrate((x^3-3*x^2+2*x-3)/(x^2+1),x, algorithm="maxima")`output `1/2*x^2 - 3*x + 1/2*log(x^2 + 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{-3 + 2x - 3x^2 + x^3}{1 + x^2} dx = \frac{1}{2}x^2 - 3x + \frac{1}{2}\log(x^2 + 1)$$

input `integrate((x^3-3*x^2+2*x-3)/(x^2+1),x, algorithm="giac")`output `1/2*x^2 - 3*x + 1/2*log(x^2 + 1)`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{-3 + 2x - 3x^2 + x^3}{1 + x^2} dx = \frac{\ln(x^2 + 1)}{2} - 3x + \frac{x^2}{2}$$

input `int((2*x - 3*x^2 + x^3 - 3)/(x^2 + 1),x)`output `log(x^2 + 1)/2 - 3*x + x^2/2`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{-3 + 2x - 3x^2 + x^3}{1 + x^2} dx = \frac{\log(x^2 + 1)}{2} + \frac{x^2}{2} - 3x$$

input `int((x^3-3*x^2+2*x-3)/(x^2+1),x)`output `(log(x**2 + 1) + x**2 - 6*x)/2`

3.7 $\int \frac{x+10x^2+6x^3+x^4}{10+6x+x^2} dx$

Optimal result	87
Mathematica [A] (verified)	87
Rubi [A] (verified)	88
Maple [A] (verified)	89
Fricas [A] (verification not implemented)	89
Sympy [A] (verification not implemented)	90
Maxima [A] (verification not implemented)	90
Giac [A] (verification not implemented)	90
Mupad [B] (verification not implemented)	91
Reduce [B] (verification not implemented)	91

Optimal result

Integrand size = 26, antiderivative size = 27

$$\int \frac{x + 10x^2 + 6x^3 + x^4}{10 + 6x + x^2} dx = \frac{x^3}{3} - 3 \arctan(3 + x) + \frac{1}{2} \log(10 + 6x + x^2)$$

output `1/3*x^3-3*arctan(3+x)+1/2*ln(x^2+6*x+10)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{x + 10x^2 + 6x^3 + x^4}{10 + 6x + x^2} dx = \frac{x^3}{3} - 3 \arctan(3 + x) + \frac{1}{2} \log(10 + 6x + x^2)$$

input `Integrate[(x + 10*x^2 + 6*x^3 + x^4)/(10 + 6*x + x^2),x]`

output `x^3/3 - 3*ArcTan[3 + x] + Log[10 + 6*x + x^2]/2`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$, Rules used = {2029, 2159, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4 + 6x^3 + 10x^2 + x}{x^2 + 6x + 10} dx$$

↓ 2029

$$\int \frac{x(x^3 + 6x^2 + 10x + 1)}{x^2 + 6x + 10} dx$$

↓ 2159

$$\int \left(x^2 + \frac{x}{x^2 + 6x + 10} \right) dx$$

↓ 2009

$$-3 \arctan(x + 3) + \frac{x^3}{3} + \frac{1}{2} \log(x^2 + 6x + 10)$$

input `Int[(x + 10*x^2 + 6*x^3 + x^4)/(10 + 6*x + x^2),x]`

output `x^3/3 - 3*ArcTan[3 + x] + Log[10 + 6*x + x^2]/2`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2029 `Int[(Fx_.)*((d_.)*(x_)^(q_.) + (a_.)*(x_)^(r_.) + (b_.)*(x_)^(s_.) + (c_.)*(x_)^(t_.))^(p_.), x_Symbol] := Int[x^(p*r)*(a + b*x^(s - r) + c*x^(t - r) + d*x^(q - r))^p*Fx, x] /; FreeQ[{a, b, c, d, r, s, t, q}, x] && IntegerQ[p] && PosQ[s - r] && PosQ[t - r] && PosQ[q - r] && !(EqQ[p, 1] && EqQ[u, 1])`

rule 2159

```
Int[(Pq_)*((d_.) + (e_.)*(x_))^(m_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*Pq*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89

method	result	size
default	$\frac{x^3}{3} - 3 \arctan(3 + x) + \frac{\ln(x^2+6x+10)}{2}$	24
risch	$\frac{x^3}{3} - 3 \arctan(3 + x) + \frac{\ln(x^2+6x+10)}{2}$	24
parallelrisc	$\frac{x^3}{3} + \frac{\ln(x+3-i)}{2} + \frac{3i \ln(x+3-i)}{2} + \frac{\ln(x+3+i)}{2} - \frac{3i \ln(x+3+i)}{2}$	41

input

```
int((x^4+6*x^3+10*x^2+x)/(x^2+6*x+10),x,method=_RETURNVERBOSE)
```

output

```
1/3*x^3-3*arctan(3+x)+1/2*ln(x^2+6*x+10)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \frac{x + 10x^2 + 6x^3 + x^4}{10 + 6x + x^2} dx = \frac{1}{3} x^3 - 3 \arctan(x + 3) + \frac{1}{2} \log(x^2 + 6x + 10)$$

input

```
integrate((x^4+6*x^3+10*x^2+x)/(x^2+6*x+10),x, algorithm="fricas")
```

output

```
1/3*x^3 - 3*arctan(x + 3) + 1/2*log(x^2 + 6*x + 10)
```

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int \frac{x + 10x^2 + 6x^3 + x^4}{10 + 6x + x^2} dx = \frac{x^3}{3} + \frac{\log(x^2 + 6x + 10)}{2} - 3 \operatorname{atan}(x + 3)$$

input `integrate((x**4+6*x**3+10*x**2+x)/(x**2+6*x+10),x)`output `x**3/3 + log(x**2 + 6*x + 10)/2 - 3*atan(x + 3)`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \frac{x + 10x^2 + 6x^3 + x^4}{10 + 6x + x^2} dx = \frac{1}{3} x^3 - 3 \arctan(x + 3) + \frac{1}{2} \log(x^2 + 6x + 10)$$

input `integrate((x^4+6*x^3+10*x^2+x)/(x^2+6*x+10),x, algorithm="maxima")`output `1/3*x^3 - 3*arctan(x + 3) + 1/2*log(x^2 + 6*x + 10)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \frac{x + 10x^2 + 6x^3 + x^4}{10 + 6x + x^2} dx = \frac{1}{3} x^3 - 3 \arctan(x + 3) + \frac{1}{2} \log(x^2 + 6x + 10)$$

input `integrate((x^4+6*x^3+10*x^2+x)/(x^2+6*x+10),x, algorithm="giac")`output `1/3*x^3 - 3*arctan(x + 3) + 1/2*log(x^2 + 6*x + 10)`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \frac{x + 10x^2 + 6x^3 + x^4}{10 + 6x + x^2} dx = \frac{\ln(x^2 + 6x + 10)}{2} - 3 \operatorname{atan}(x + 3) + \frac{x^3}{3}$$

input `int((x + 10*x^2 + 6*x^3 + x^4)/(6*x + x^2 + 10),x)`output `log(6*x + x^2 + 10)/2 - 3*atan(x + 3) + x^3/3`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \frac{x + 10x^2 + 6x^3 + x^4}{10 + 6x + x^2} dx = -3 \operatorname{atan}(x + 3) + \frac{\log(x^2 + 6x + 10)}{2} + \frac{x^3}{3}$$

input `int((x^4+6*x^3+10*x^2+x)/(x^2+6*x+10),x)`output `(- 18*atan(x + 3) + 3*log(x**2 + 6*x + 10) + 2*x**3)/6`

$$3.8 \quad \int \frac{1}{-18+27x-7x^2-3x^3+x^4} dx$$

Optimal result	92
Mathematica [A] (verified)	92
Rubi [A] (verified)	93
Maple [A] (verified)	94
Fricas [A] (verification not implemented)	94
Sympy [A] (verification not implemented)	95
Maxima [A] (verification not implemented)	95
Giac [A] (verification not implemented)	95
Mupad [B] (verification not implemented)	96
Reduce [B] (verification not implemented)	96

Optimal result

Integrand size = 20, antiderivative size = 39

$$\int \frac{1}{-18+27x-7x^2-3x^3+x^4} dx = \frac{1}{8} \log(1-x) - \frac{1}{5} \log(2-x) + \frac{1}{12} \log(3-x) - \frac{1}{120} \log(3+x)$$

output `1/8*ln(1-x)-1/5*ln(2-x)+1/12*ln(3-x)-1/120*ln(3+x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00

$$\int \frac{1}{-18+27x-7x^2-3x^3+x^4} dx = \frac{1}{8} \log(1-x) - \frac{1}{5} \log(2-x) + \frac{1}{12} \log(3-x) - \frac{1}{120} \log(3+x)$$

input `Integrate[(-18 + 27*x - 7*x^2 - 3*x^3 + x^4)^(-1),x]`

output `Log[1 - x]/8 - Log[2 - x]/5 + Log[3 - x]/12 - Log[3 + x]/120`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^4 - 3x^3 - 7x^2 + 27x - 18} dx$$

↓ 2462

$$\int \left(-\frac{1}{5(x-2)} + \frac{1}{8(x-1)} - \frac{1}{120(x+3)} + \frac{1}{12(x-3)} \right) dx$$

↓ 2009

$$\frac{1}{8} \log(1-x) - \frac{1}{5} \log(2-x) + \frac{1}{12} \log(3-x) - \frac{1}{120} \log(x+3)$$

input

```
Int[(-18 + 27*x - 7*x^2 - 3*x^3 + x^4)^(-1), x]
```

output

```
Log[1 - x]/8 - Log[2 - x]/5 + Log[3 - x]/12 - Log[3 + x]/120
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2462

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegr
and[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ
[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && !LtQ[p, 0
] && RationalFunctionQ[u, x]
```

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.67

method	result	size
default	$\frac{\ln(-3+x)}{12} - \frac{\ln(3+x)}{120} + \frac{\ln(-1+x)}{8} - \frac{\ln(-2+x)}{5}$	26
norman	$\frac{\ln(-3+x)}{12} - \frac{\ln(3+x)}{120} + \frac{\ln(-1+x)}{8} - \frac{\ln(-2+x)}{5}$	26
risch	$\frac{\ln(-3+x)}{12} - \frac{\ln(3+x)}{120} + \frac{\ln(-1+x)}{8} - \frac{\ln(-2+x)}{5}$	26
parallelrisc	$\frac{\ln(-3+x)}{12} - \frac{\ln(3+x)}{120} + \frac{\ln(-1+x)}{8} - \frac{\ln(-2+x)}{5}$	26

input `int(1/(x^4-3*x^3-7*x^2+27*x-18),x,method=_RETURNVERBOSE)`

output `1/12*ln(-3+x)-1/120*ln(3+x)+1/8*ln(-1+x)-1/5*ln(-2+x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.64

$$\int \frac{1}{-18 + 27x - 7x^2 - 3x^3 + x^4} dx = -\frac{1}{120} \log(x + 3) + \frac{1}{8} \log(x - 1) - \frac{1}{5} \log(x - 2) + \frac{1}{12} \log(x - 3)$$

input `integrate(1/(x^4-3*x^3-7*x^2+27*x-18),x, algorithm="fricas")`

output `-1/120*log(x + 3) + 1/8*log(x - 1) - 1/5*log(x - 2) + 1/12*log(x - 3)`

Sympy [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.67

$$\int \frac{1}{-18 + 27x - 7x^2 - 3x^3 + x^4} dx = \frac{\log(x-3)}{12} - \frac{\log(x-2)}{5} + \frac{\log(x-1)}{8} - \frac{\log(x+3)}{120}$$

input `integrate(1/(x**4-3*x**3-7*x**2+27*x-18),x)`output `log(x - 3)/12 - log(x - 2)/5 + log(x - 1)/8 - log(x + 3)/120`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.64

$$\int \frac{1}{-18 + 27x - 7x^2 - 3x^3 + x^4} dx = -\frac{1}{120} \log(x+3) + \frac{1}{8} \log(x-1) - \frac{1}{5} \log(x-2) + \frac{1}{12} \log(x-3)$$

input `integrate(1/(x^4-3*x^3-7*x^2+27*x-18),x, algorithm="maxima")`output `-1/120*log(x + 3) + 1/8*log(x - 1) - 1/5*log(x - 2) + 1/12*log(x - 3)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.74

$$\int \frac{1}{-18 + 27x - 7x^2 - 3x^3 + x^4} dx = -\frac{1}{120} \log(|x+3|) + \frac{1}{8} \log(|x-1|) - \frac{1}{5} \log(|x-2|) + \frac{1}{12} \log(|x-3|)$$

input `integrate(1/(x^4-3*x^3-7*x^2+27*x-18),x, algorithm="giac")`

output $-1/120*\log(\text{abs}(x + 3)) + 1/8*\log(\text{abs}(x - 1)) - 1/5*\log(\text{abs}(x - 2)) + 1/12*\log(\text{abs}(x - 3))$

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.64

$$\int \frac{1}{-18 + 27x - 7x^2 - 3x^3 + x^4} dx = \frac{\ln(x - 1)}{8} - \frac{\ln(x - 2)}{5} + \frac{\ln(x - 3)}{12} - \frac{\ln(x + 3)}{120}$$

input $\text{int}(-1/(7*x^2 - 27*x + 3*x^3 - x^4 + 18), x)$

output $\log(x - 1)/8 - \log(x - 2)/5 + \log(x - 3)/12 - \log(x + 3)/120$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.64

$$\int \frac{1}{-18 + 27x - 7x^2 - 3x^3 + x^4} dx = \frac{\log(x - 3)}{12} - \frac{\log(x - 2)}{5} + \frac{\log(x - 1)}{8} - \frac{\log(x + 3)}{120}$$

input $\text{int}(1/(x^4-3*x^3-7*x^2+27*x-18), x)$

output $(10*\log(x - 3) - 24*\log(x - 2) + 15*\log(x - 1) - \log(x + 3))/120$

3.9 $\int \frac{1+x^3}{-2+x} dx$

Optimal result	97
Mathematica [A] (verified)	97
Rubi [A] (verified)	98
Maple [A] (verified)	99
Fricas [A] (verification not implemented)	99
Sympy [A] (verification not implemented)	100
Maxima [A] (verification not implemented)	100
Giac [A] (verification not implemented)	100
Mupad [B] (verification not implemented)	101
Reduce [B] (verification not implemented)	101

Optimal result

Integrand size = 11, antiderivative size = 22

$$\int \frac{1+x^3}{-2+x} dx = 4x + x^2 + \frac{x^3}{3} + 9 \log(2-x)$$

output `4*x+x^2+1/3*x^3+9*ln(2-x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.05

$$\int \frac{1+x^3}{-2+x} dx = -\frac{44}{3} + 4x + x^2 + \frac{x^3}{3} + 9 \log(-2+x)$$

input `Integrate[(1 + x^3)/(-2 + x),x]`

output `-44/3 + 4*x + x^2 + x^3/3 + 9*Log[-2 + x]`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2389, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3 + 1}{x - 2} dx$$

$$\downarrow \text{2389}$$

$$\int \left(x^2 + 2x + \frac{9}{x - 2} + 4 \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{x^3}{3} + x^2 + 4x + 9 \log(2 - x)$$

input

```
Int[(1 + x^3)/(-2 + x),x]
```

output

```
4*x + x^2 + x^3/3 + 9*Log[2 - x]
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2389

```
Int[(Pq_)*((a_) + (b_.)*(x_)^(n_.))^(p_.), x_Symbol] :> Int[ExpandIntegrand
[Pq*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, n}, x] && PolyQ[Pq, x] && (IGtQ[p
, 0] || EqQ[n, 1])
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.86

method	result	size
default	$\frac{x^3}{3} + x^2 + 4x + 9 \ln(-2 + x)$	19
norman	$\frac{x^3}{3} + x^2 + 4x + 9 \ln(-2 + x)$	19
risch	$\frac{x^3}{3} + x^2 + 4x + 9 \ln(-2 + x)$	19
parallelrisch	$\frac{x^3}{3} + x^2 + 4x + 9 \ln(-2 + x)$	19
meijerg	$9 \ln\left(1 - \frac{x}{2}\right) + \frac{x(x^2+3x+12)}{3}$	21

input `int((x^3+1)/(-2+x),x,method=_RETURNVERBOSE)`output `1/3*x^3+x^2+4*x+9*ln(-2+x)`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \frac{1+x^3}{-2+x} dx = \frac{1}{3}x^3 + x^2 + 4x + 9 \log(x-2)$$

input `integrate((x^3+1)/(-2+x),x, algorithm="fricas")`output `1/3*x^3 + x^2 + 4*x + 9*log(x - 2)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

$$\int \frac{1+x^3}{-2+x} dx = \frac{x^3}{3} + x^2 + 4x + 9 \log(x-2)$$

input `integrate((x**3+1)/(-2+x),x)`output `x**3/3 + x**2 + 4*x + 9*log(x - 2)`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \frac{1+x^3}{-2+x} dx = \frac{1}{3} x^3 + x^2 + 4x + 9 \log(x-2)$$

input `integrate((x^3+1)/(-2+x),x, algorithm="maxima")`output `1/3*x^3 + x^2 + 4*x + 9*log(x - 2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.86

$$\int \frac{1+x^3}{-2+x} dx = \frac{1}{3} x^3 + x^2 + 4x + 9 \log(|x-2|)$$

input `integrate((x^3+1)/(-2+x),x, algorithm="giac")`output `1/3*x^3 + x^2 + 4*x + 9*log(abs(x - 2))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \frac{1+x^3}{-2+x} dx = 4x + 9 \ln(x-2) + x^2 + \frac{x^3}{3}$$

input `int((x^3 + 1)/(x - 2), x)`

output `4*x + 9*log(x - 2) + x^2 + x^3/3`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \frac{1+x^3}{-2+x} dx = 9 \log(x-2) + \frac{x^3}{3} + x^2 + 4x$$

input `int((x^3+1)/(-2+x), x)`

output `(27*log(x - 2) + x**3 + 3*x**2 + 12*x)/3`

3.10 $\int \frac{3x-4x^2+3x^3}{1+x^2} dx$

Optimal result	102
Mathematica [A] (verified)	102
Rubi [A] (verified)	103
Maple [A] (verified)	104
Fricas [A] (verification not implemented)	104
Sympy [A] (verification not implemented)	105
Maxima [A] (verification not implemented)	105
Giac [A] (verification not implemented)	105
Mupad [B] (verification not implemented)	106
Reduce [B] (verification not implemented)	106

Optimal result

Integrand size = 22, antiderivative size = 15

$$\int \frac{3x - 4x^2 + 3x^3}{1 + x^2} dx = -4x + \frac{3x^2}{2} + 4 \arctan(x)$$

output

```
-4*x+3/2*x^2+4*arctan(x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{3x - 4x^2 + 3x^3}{1 + x^2} dx = -4x + \frac{3x^2}{2} + 4 \arctan(x)$$

input

```
Integrate[(3*x - 4*x^2 + 3*x^3)/(1 + x^2), x]
```

output

```
-4*x + (3*x^2)/2 + 4*ArcTan[x]
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.136$, Rules used = {2028, 2333, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{3x^3 - 4x^2 + 3x}{x^2 + 1} dx$$

$$\downarrow \text{2028}$$

$$\int \frac{x(3x^2 - 4x + 3)}{x^2 + 1} dx$$

$$\downarrow \text{2333}$$

$$\int \left(\frac{4}{x^2 + 1} + 3x - 4 \right) dx$$

$$\downarrow \text{2009}$$

$$4 \arctan(x) + \frac{3x^2}{2} - 4x$$

input `Int[(3*x - 4*x^2 + 3*x^3)/(1 + x^2), x]`

output `-4*x + (3*x^2)/2 + 4*ArcTan[x]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2028 `Int[(Fx_.)*((a_.)*(x_)^(r_.) + (b_.)*(x_)^(s_.) + (c_.)*(x_)^(t_.))^(p_.), x_Symbol] := Int[x^(p*r)*(a + b*x^(s - r) + c*x^(t - r))^p*Fx, x] /; FreeQ[{a, b, c, r, s, t}, x] && IntegerQ[p] && PosQ[s - r] && PosQ[t - r] && !(EqQ[p, 1] && EqQ[u, 1])`

rule 2333

```
Int[(Pq_)*((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[
ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x]
&& PolyQ[Pq, x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

method	result	size
default	$-4x + \frac{3x^2}{2} + 4 \arctan(x)$	14
meijerg	$-4x + \frac{3x^2}{2} + 4 \arctan(x)$	14
risch	$-4x + \frac{3x^2}{2} + 4 \arctan(x)$	14
parallelrisch	$-4x + \frac{3x^2}{2} + 2i \ln(x + i) - 2i \ln(x - i)$	26

input

```
int((3*x^3-4*x^2+3*x)/(x^2+1),x,method=_RETURNVERBOSE)
```

output

```
-4*x+3/2*x^2+4*arctan(x)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{3x - 4x^2 + 3x^3}{1 + x^2} dx = \frac{3}{2}x^2 - 4x + 4 \arctan(x)$$

input

```
integrate((3*x^3-4*x^2+3*x)/(x^2+1),x, algorithm="fricas")
```

output

```
3/2*x^2 - 4*x + 4*arctan(x)
```

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{3x - 4x^2 + 3x^3}{1 + x^2} dx = \frac{3x^2}{2} - 4x + 4 \operatorname{atan}(x)$$

input `integrate((3*x**3-4*x**2+3*x)/(x**2+1),x)`

output `3*x**2/2 - 4*x + 4*atan(x)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{3x - 4x^2 + 3x^3}{1 + x^2} dx = \frac{3}{2}x^2 - 4x + 4 \arctan(x)$$

input `integrate((3*x^3-4*x^2+3*x)/(x^2+1),x, algorithm="maxima")`

output `3/2*x^2 - 4*x + 4*arctan(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{3x - 4x^2 + 3x^3}{1 + x^2} dx = \frac{3}{2}x^2 - 4x + 4 \arctan(x)$$

input `integrate((3*x^3-4*x^2+3*x)/(x^2+1),x, algorithm="giac")`

output `3/2*x^2 - 4*x + 4*arctan(x)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{3x - 4x^2 + 3x^3}{1 + x^2} dx = 4 \operatorname{atan}(x) - 4x + \frac{3x^2}{2}$$

input `int((3*x - 4*x^2 + 3*x^3)/(x^2 + 1),x)`

output `4*atan(x) - 4*x + (3*x^2)/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{3x - 4x^2 + 3x^3}{1 + x^2} dx = 4 \operatorname{atan}(x) + \frac{3x^2}{2} - 4x$$

input `int((3*x^3-4*x^2+3*x)/(x^2+1),x)`

output `(8*atan(x) + 3*x**2 - 8*x)/2`

3.11 $\int \frac{5+3x}{1-x-x^2+x^3} dx$

Optimal result	107
Mathematica [A] (verified)	107
Rubi [A] (verified)	108
Maple [A] (verified)	109
Fricas [B] (verification not implemented)	109
Sympy [B] (verification not implemented)	110
Maxima [A] (verification not implemented)	110
Giac [B] (verification not implemented)	110
Mupad [B] (verification not implemented)	111
Reduce [B] (verification not implemented)	111

Optimal result

Integrand size = 21, antiderivative size = 12

$$\int \frac{5+3x}{1-x-x^2+x^3} dx = \frac{4}{1-x} + \operatorname{arctanh}(x)$$

output `4/(1-x)+arctanh(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 2.00

$$\int \frac{5+3x}{1-x-x^2+x^3} dx = -\frac{4}{-1+x} - \frac{1}{2} \log(-1+x) + \frac{1}{2} \log(1+x)$$

input `Integrate[(5 + 3*x)/(1 - x - x^2 + x^3), x]`

output `-4/(-1 + x) - Log[-1 + x]/2 + Log[1 + x]/2`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{3x + 5}{x^3 - x^2 - x + 1} dx$$

$$\downarrow \text{2462}$$

$$\int \left(\frac{1}{1 - x^2} + \frac{4}{(x - 1)^2} \right) dx$$

$$\downarrow \text{2009}$$

$$\operatorname{arctanh}(x) + \frac{4}{1 - x}$$

input

```
Int[(5 + 3*x)/(1 - x - x^2 + x^3),x]
```

output

```
4/(1 - x) + ArcTanh[x]
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2462

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegr
and[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ
[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0
] && RationalFunctionQ[u, x]
```

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.75

method	result	size
default	$-\frac{4}{-1+x} - \frac{\ln(-1+x)}{2} + \frac{\ln(1+x)}{2}$	21
norman	$-\frac{4}{-1+x} - \frac{\ln(-1+x)}{2} + \frac{\ln(1+x)}{2}$	21
risch	$-\frac{4}{-1+x} - \frac{\ln(-1+x)}{2} + \frac{\ln(1+x)}{2}$	21
parallelrisch	$-\frac{\ln(-1+x)x - \ln(1+x)x + 8 - \ln(-1+x) + \ln(1+x)}{2(-1+x)}$	33

input `int((5+3*x)/(x^3-x^2-x+1),x,method=_RETURNVERBOSE)`

output `-4/(-1+x)-1/2*ln(-1+x)+1/2*ln(1+x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 26 vs. 2(10) = 20.

Time = 0.06 (sec) , antiderivative size = 26, normalized size of antiderivative = 2.17

$$\int \frac{5+3x}{1-x-x^2+x^3} dx = \frac{(x-1)\log(x+1) - (x-1)\log(x-1) - 8}{2(x-1)}$$

input `integrate((5+3*x)/(x^3-x^2-x+1),x, algorithm="fricas")`

output `1/2*((x - 1)*log(x + 1) - (x - 1)*log(x - 1) - 8)/(x - 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17 vs. $2(7) = 14$.

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.42

$$\int \frac{5 + 3x}{1 - x - x^2 + x^3} dx = -\frac{\log(x - 1)}{2} + \frac{\log(x + 1)}{2} - \frac{4}{x - 1}$$

input `integrate((5+3*x)/(x**3-x**2-x+1),x)`

output `-log(x - 1)/2 + log(x + 1)/2 - 4/(x - 1)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.67

$$\int \frac{5 + 3x}{1 - x - x^2 + x^3} dx = -\frac{4}{x - 1} + \frac{1}{2} \log(x + 1) - \frac{1}{2} \log(x - 1)$$

input `integrate((5+3*x)/(x^3-x^2-x+1),x, algorithm="maxima")`

output `-4/(x - 1) + 1/2*log(x + 1) - 1/2*log(x - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 22 vs. $2(10) = 20$.

Time = 0.12 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.83

$$\int \frac{5 + 3x}{1 - x - x^2 + x^3} dx = -\frac{4}{x - 1} + \frac{1}{2} \log(|x + 1|) - \frac{1}{2} \log(|x - 1|)$$

input `integrate((5+3*x)/(x^3-x^2-x+1),x, algorithm="giac")`

output `-4/(x - 1) + 1/2*log(abs(x + 1)) - 1/2*log(abs(x - 1))`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{5 + 3x}{1 - x - x^2 + x^3} dx = \operatorname{atanh}(x) - \frac{4}{x - 1}$$

input `int(-(3*x + 5)/(x + x^2 - x^3 - 1),x)`output `atanh(x) - 4/(x - 1)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 35, normalized size of antiderivative = 2.92

$$\int \frac{5 + 3x}{1 - x - x^2 + x^3} dx = \frac{-\log(x - 1)x + \log(x - 1) + \log(x + 1)x - \log(x + 1) - 8x}{2x - 2}$$

input `int((5+3*x)/(x^3-x^2-x+1),x)`output `(- log(x - 1)*x + log(x - 1) + log(x + 1)*x - log(x + 1) - 8*x)/(2*(x - 1))`

$$3.12 \quad \int \frac{-1-x-x^3+x^4}{-x^2+x^3} dx$$

Optimal result	112
Mathematica [A] (verified)	112
Rubi [A] (verified)	113
Maple [A] (verified)	114
Fricas [A] (verification not implemented)	114
Sympy [A] (verification not implemented)	115
Maxima [A] (verification not implemented)	115
Giac [A] (verification not implemented)	115
Mupad [B] (verification not implemented)	116
Reduce [B] (verification not implemented)	116

Optimal result

Integrand size = 25, antiderivative size = 25

$$\int \frac{-1-x-x^3+x^4}{-x^2+x^3} dx = -\frac{1}{x} + \frac{x^2}{2} - 2\log(1-x) + 2\log(x)$$

output

```
-1/x+1/2*x^2-2*ln(1-x)+2*ln(x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{-1-x-x^3+x^4}{-x^2+x^3} dx = -\frac{1}{x} + \frac{x^2}{2} - 2\log(1-x) + 2\log(x)$$

input

```
Integrate[(-1 - x - x^3 + x^4)/(-x^2 + x^3), x]
```

output

```
-x^(-1) + x^2/2 - 2*Log[1 - x] + 2*Log[x]
```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2026, 2123, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4 - x^3 - x - 1}{x^3 - x^2} dx$$

↓ 2026

$$\int \frac{x^4 - x^3 - x - 1}{(x - 1)x^2} dx$$

↓ 2123

$$\int \left(\frac{1}{x^2} + x - \frac{2}{x - 1} + \frac{2}{x} \right) dx$$

↓ 2009

$$\frac{x^2}{2} - \frac{1}{x} - 2 \log(1 - x) + 2 \log(x)$$

input `Int[(-1 - x - x^3 + x^4)/(-x^2 + x^3), x]`

output `-x^(-1) + x^2/2 - 2*Log[1 - x] + 2*Log[x]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2123

```
Int[(Px_)*((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol]
:> Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c,
d, m, n}, x] && PolyQ[Px, x] && (IntegersQ[m, n] || IGtQ[m, -2])
```

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

method	result	size
default	$\frac{x^2}{2} - \frac{1}{x} + 2 \ln(x) - 2 \ln(-1 + x)$	22
risch	$\frac{x^2}{2} - \frac{1}{x} + 2 \ln(x) - 2 \ln(-1 + x)$	22
norman	$\frac{-1 + \frac{x^3}{2}}{x} + 2 \ln(x) - 2 \ln(-1 + x)$	23
parallelrisc	$\frac{x^3 + 4x \ln(x) - 4 \ln(-1 + x)x - 2}{2x}$	23
meijerg	$-\frac{1}{x} + 2 \ln(x) + 2i\pi + \frac{x(3x+6)}{6} - x - 2 \ln(1 - x)$	34

input

```
int((x^4-x^3-x-1)/(x^3-x^2),x,method=_RETURNVERBOSE)
```

output

```
1/2*x^2-1/x+2*ln(x)-2*ln(-1+x)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \frac{-1 - x - x^3 + x^4}{-x^2 + x^3} dx = \frac{x^3 - 4x \log(x - 1) + 4x \log(x) - 2}{2x}$$

input

```
integrate((x^4-x^3-x-1)/(x^3-x^2),x, algorithm="fricas")
```

output

```
1/2*(x^3 - 4*x*log(x - 1) + 4*x*log(x) - 2)/x
```

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{-1 - x - x^3 + x^4}{-x^2 + x^3} dx = \frac{x^2}{2} + 2 \log(x) - 2 \log(x - 1) - \frac{1}{x}$$

input `integrate((x**4-x**3-x-1)/(x**3-x**2),x)`output `x**2/2 + 2*log(x) - 2*log(x - 1) - 1/x`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int \frac{-1 - x - x^3 + x^4}{-x^2 + x^3} dx = \frac{1}{2} x^2 - \frac{1}{x} - 2 \log(x - 1) + 2 \log(x)$$

input `integrate((x^4-x^3-x-1)/(x^3-x^2),x, algorithm="maxima")`output `1/2*x^2 - 1/x - 2*log(x - 1) + 2*log(x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \frac{-1 - x - x^3 + x^4}{-x^2 + x^3} dx = \frac{1}{2} x^2 - \frac{1}{x} - 2 \log(|x - 1|) + 2 \log(|x|)$$

input `integrate((x^4-x^3-x-1)/(x^3-x^2),x, algorithm="giac")`output `1/2*x^2 - 1/x - 2*log(abs(x - 1)) + 2*log(abs(x))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{-1 - x - x^3 + x^4}{-x^2 + x^3} dx = 4 \operatorname{atanh}(2x - 1) - \frac{1}{x} + \frac{x^2}{2}$$

input `int((x + x^3 - x^4 + 1)/(x^2 - x^3), x)`output `4*atanh(2*x - 1) - 1/x + x^2/2`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \frac{-1 - x - x^3 + x^4}{-x^2 + x^3} dx = \frac{-4 \log(x - 1) x + 4 \log(x) x + x^3 - 2}{2x}$$

input `int((x^4-x^3-x-1)/(x^3-x^2), x)`output `(- 4*log(x - 1)*x + 4*log(x)*x + x**3 - 2)/(2*x)`

3.13 $\int \frac{2+x+x^2+x^3}{2+3x^2+x^4} dx$

Optimal result	117
Mathematica [A] (verified)	117
Rubi [A] (verified)	118
Maple [A] (verified)	119
Fricas [A] (verification not implemented)	120
Sympy [A] (verification not implemented)	120
Maxima [A] (verification not implemented)	120
Giac [A] (verification not implemented)	121
Mupad [B] (verification not implemented)	121
Reduce [B] (verification not implemented)	121

Optimal result

Integrand size = 22, antiderivative size = 13

$$\int \frac{2+x+x^2+x^3}{2+3x^2+x^4} dx = \arctan(x) + \frac{1}{2} \log(2+x^2)$$

output

```
arctan(x)+1/2*ln(x^2+2)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{2+x+x^2+x^3}{2+3x^2+x^4} dx = \arctan(x) + \frac{1}{2} \log(2+x^2)$$

input

```
Integrate[(2 + x + x^2 + x^3)/(2 + 3*x^2 + x^4), x]
```

output

```
ArcTan[x] + Log[2 + x^2]/2
```

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2202, 1387, 216, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3 + x^2 + x + 2}{x^4 + 3x^2 + 2} dx \\
 & \quad \downarrow \text{2202} \\
 & \int \frac{x(x^2 + 1)}{x^4 + 3x^2 + 2} dx + \int \frac{x^2 + 2}{x^4 + 3x^2 + 2} dx \\
 & \quad \downarrow \text{1387} \\
 & \int \frac{1}{x^2 + 1} dx + \int \frac{x}{x^2 + 2} dx \\
 & \quad \downarrow \text{216} \\
 & \int \frac{x}{x^2 + 2} dx + \arctan(x) \\
 & \quad \downarrow \text{240} \\
 & \arctan(x) + \frac{1}{2} \log(x^2 + 2)
 \end{aligned}$$

input `Int[(2 + x + x^2 + x^3)/(2 + 3*x^2 + x^4), x]`

output `ArcTan[x] + Log[2 + x^2]/2`

Definitions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 240 `Int[(x_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`

rule 1387 `Int[(u_.)*((a_) + (c_.)*(x_)^(n2_.) + (b_.)*(x_)^(n_.))^p_.*((d_) + (e_.)*(x_)^(n_.))^q_.], x_Symbol] := Int[u*(d + e*x^n)^(p + q)*(a/d + (c/e)*x^n)^p, x] /; FreeQ[{a, b, c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[c*d^2 - b*d*e + a*e^2, 0] && (IntegerQ[p] || (GtQ[a, 0] && GtQ[d, 0] && LtQ[c, 0]))`

rule 2202 `Int[(Pn_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^p_], x_Symbol] := Module[{n = Expon[Pn, x], k}, Int[Sum[Coeff[Pn, x, 2*k]*x^(2*k), {k, 0, n/2}]*(a + b*x^2 + c*x^4)^p, x] + Int[x*Sum[Coeff[Pn, x, 2*k + 1]*x^(2*k), {k, 0, (n - 1)/2}]*(a + b*x^2 + c*x^4)^p, x]] /; FreeQ[{a, b, c, p}, x] && PolyQ[Pn, x] && !PolyQ[Pn, x^2]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
default	$\arctan(x) + \frac{\ln(x^2+2)}{2}$	12
risch	$\arctan(x) + \frac{\ln(x^2+2)}{2}$	12
parallelrisch	$\frac{i \ln(x+i)}{2} - \frac{i \ln(x-i)}{2} + \frac{\ln(x^2+2)}{2}$	26

input `int((x^3+x^2+x+2)/(x^4+3*x^2+2),x,method=_RETURNVERBOSE)`

output `arctan(x)+1/2*ln(x^2+2)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{2 + x + x^2 + x^3}{2 + 3x^2 + x^4} dx = \arctan(x) + \frac{1}{2} \log(x^2 + 2)$$

input `integrate((x^3+x^2+x+2)/(x^4+3*x^2+2),x, algorithm="fricas")`output `arctan(x) + 1/2*log(x^2 + 2)`**Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \frac{2 + x + x^2 + x^3}{2 + 3x^2 + x^4} dx = \frac{\log(x^2 + 2)}{2} + \operatorname{atan}(x)$$

input `integrate((x**3+x**2+x+2)/(x**4+3*x**2+2),x)`output `log(x**2 + 2)/2 + atan(x)`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{2 + x + x^2 + x^3}{2 + 3x^2 + x^4} dx = \arctan(x) + \frac{1}{2} \log(x^2 + 2)$$

input `integrate((x^3+x^2+x+2)/(x^4+3*x^2+2),x, algorithm="maxima")`output `arctan(x) + 1/2*log(x^2 + 2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{2 + x + x^2 + x^3}{2 + 3x^2 + x^4} dx = \arctan(x) + \frac{1}{2} \log(x^2 + 2)$$

input `integrate((x^3+x^2+x+2)/(x^4+3*x^2+2),x, algorithm="giac")`

output `arctan(x) + 1/2*log(x^2 + 2)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{2 + x + x^2 + x^3}{2 + 3x^2 + x^4} dx = \frac{\ln(x^2 + 2)}{2} + \operatorname{atan}(x)$$

input `int((x + x^2 + x^3 + 2)/(3*x^2 + x^4 + 2),x)`

output `log(x^2 + 2)/2 + atan(x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{2 + x + x^2 + x^3}{2 + 3x^2 + x^4} dx = \operatorname{atan}(x) + \frac{\log(x^2 + 2)}{2}$$

input `int((x^3+x^2+x+2)/(x^4+3*x^2+2),x)`

output `(2*atan(x) + log(x**2 + 2))/2`

$$3.14 \quad \int \frac{-4+8x-4x^2+4x^3-x^4+x^5}{(2+x^2)^3} dx$$

Optimal result	122
Mathematica [A] (verified)	122
Rubi [A] (verified)	123
Maple [A] (verified)	125
Fricas [A] (verification not implemented)	125
Sympy [A] (verification not implemented)	126
Maxima [A] (verification not implemented)	126
Giac [A] (verification not implemented)	126
Mupad [B] (verification not implemented)	127
Reduce [B] (verification not implemented)	127

Optimal result

Integrand size = 31, antiderivative size = 35

$$\int \frac{-4 + 8x - 4x^2 + 4x^3 - x^4 + x^5}{(2 + x^2)^3} dx = -\frac{1}{(2 + x^2)^2} - \frac{\arctan\left(\frac{x}{\sqrt{2}}\right)}{\sqrt{2}} + \frac{1}{2} \log(2 + x^2)$$

output `-1/(x^2+2)^2-1/2*arctan(1/2*x*2^(1/2))*2^(1/2)+1/2*ln(x^2+2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{-4 + 8x - 4x^2 + 4x^3 - x^4 + x^5}{(2 + x^2)^3} dx = -\frac{1}{(2 + x^2)^2} - \frac{\arctan\left(\frac{x}{\sqrt{2}}\right)}{\sqrt{2}} + \frac{1}{2} \log(2 + x^2)$$

input `Integrate[(-4 + 8*x - 4*x^2 + 4*x^3 - x^4 + x^5)/(2 + x^2)^3,x]`

output `-(2 + x^2)^(-2) - ArcTan[x/Sqrt[2]]/Sqrt[2] + Log[2 + x^2]/2`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.194$, Rules used = {2345, 27, 2019, 452, 216, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^5 - x^4 + 4x^3 - 4x^2 + 8x - 4}{(x^2 + 2)^3} dx \\
 & \quad \downarrow \text{2345} \\
 & -\frac{1}{8} \int \frac{8(-x^3 + x^2 - 2x + 2)}{(x^2 + 2)^2} dx - \frac{1}{(x^2 + 2)^2} \\
 & \quad \downarrow \text{27} \\
 & - \int \frac{-x^3 + x^2 - 2x + 2}{(x^2 + 2)^2} dx - \frac{1}{(x^2 + 2)^2} \\
 & \quad \downarrow \text{2019} \\
 & - \int \frac{1 - x}{x^2 + 2} dx - \frac{1}{(x^2 + 2)^2} \\
 & \quad \downarrow \text{452} \\
 & - \int \frac{1}{x^2 + 2} dx + \int \frac{x}{x^2 + 2} dx - \frac{1}{(x^2 + 2)^2} \\
 & \quad \downarrow \text{216} \\
 & \int \frac{x}{x^2 + 2} dx - \frac{\arctan\left(\frac{x}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{1}{(x^2 + 2)^2} \\
 & \quad \downarrow \text{240} \\
 & -\frac{\arctan\left(\frac{x}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{1}{(x^2 + 2)^2} + \frac{1}{2} \log(x^2 + 2)
 \end{aligned}$$

input

```
Int[(-4 + 8*x - 4*x^2 + 4*x^3 - x^4 + x^5)/(2 + x^2)^3,x]
```

output $-(2 + x^2)^{-2} - \text{ArcTan}[x/\text{Sqrt}[2]]/\text{Sqrt}[2] + \text{Log}[2 + x^2]/2$

Defintions of rubi rules used

rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$

rule 216 $\text{Int}[(a_*) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

rule 240 $\text{Int}[(x_)/((a_*) + (b_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x^2, x]]/(2*b), x] /; \text{FreeQ}\{a, b\}, x]$

rule 452 $\text{Int}[(c_*) + (d_*)(x_)/((a_*) + (b_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[c \text{ Int}[1/(a + b*x^2), x], x] + \text{Simp}[d \text{ Int}[x/(a + b*x^2), x], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[b*c^2 + a*d^2, 0]$

rule 2019 $\text{Int}[(u_*)(Px_)^{(p_*)}(Qx_)^{(q_*)}, x_Symbol] \rightarrow \text{Int}[u*\text{PolynomialQuotient}[Px, Qx, x]^p*Qx^{(p+q)}, x] /; \text{FreeQ}[q, x] \ \&\& \ \text{PolyQ}[Px, x] \ \&\& \ \text{PolyQ}[Qx, x] \ \&\& \ \text{EqQ}[\text{PolynomialRemainder}[Px, Qx, x], 0] \ \&\& \ \text{IntegerQ}[p] \ \&\& \ \text{LtQ}[p*q, 0]$

rule 2345 $\text{Int}[(Pq_*)((a_*) + (b_*)(x_)^2)^{(p_*)}, x_Symbol] \rightarrow \text{With}\{Q = \text{PolynomialQuotient}[Pq, a + b*x^2, x], f = \text{Coeff}[\text{PolynomialRemainder}[Pq, a + b*x^2, x], x, 0], g = \text{Coeff}[\text{PolynomialRemainder}[Pq, a + b*x^2, x], x, 1]\}, \text{Simp}[(a*g - b*f*x)*((a + b*x^2)^{(p+1})/(2*a*b*(p+1))), x] + \text{Simp}[1/(2*a*(p+1)) \text{Int}[(a + b*x^2)^{(p+1)}*\text{ExpandToSum}[2*a*(p+1)*Q + f*(2*p+3), x], x], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PolyQ}[Pq, x] \ \&\& \ \text{LtQ}[p, -1]$

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.89

method	result
default	$-\frac{1}{(x^2+2)^2} - \frac{\arctan\left(\frac{x\sqrt{2}}{2}\right)\sqrt{2}}{2} + \frac{\ln(x^2+2)}{2}$
risch	$-\frac{1}{(x^2+2)^2} - \frac{\arctan\left(\frac{x\sqrt{2}}{2}\right)\sqrt{2}}{2} + \frac{\ln(x^2+2)}{2}$
meijerg	$-\frac{\sqrt{2}\left(\frac{x\sqrt{2}\left(\frac{3x^2+5}{2}\right) + \frac{3\arctan\left(\frac{x\sqrt{2}}{2}\right)}{2}\right)}{8} - \frac{x^2\left(\frac{9x^2+6}{2}\right)}{24\left(\frac{x^2+1}{2}\right)^2} + \frac{\ln\left(\frac{x^2+1}{2}\right)}{2} - \frac{\sqrt{2}\left(\frac{x\sqrt{2}\left(\frac{25x^2+15}{2}\right) + \frac{3\arctan\left(\frac{x\sqrt{2}}{2}\right)}{2}\right)}{8} + \frac{1}{8\left(\frac{x^2+1}{2}\right)}$

input `int((x^5-x^4+4*x^3-4*x^2+8*x-4)/(x^2+2)^3,x,method=_RETURNVERBOSE)`

output `-1/(x^2+2)^2-1/2*arctan(1/2*x*2^(1/2))*2^(1/2)+1/2*ln(x^2+2)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.57

$$\int \frac{-4 + 8x - 4x^2 + 4x^3 - x^4 + x^5}{(2 + x^2)^3} dx$$

$$= -\frac{\sqrt{2}(x^4 + 4x^2 + 4) \arctan\left(\frac{1}{2}\sqrt{2}x\right) - (x^4 + 4x^2 + 4) \log(x^2 + 2) + 2}{2(x^4 + 4x^2 + 4)}$$

input `integrate((x^5-x^4+4*x^3-4*x^2+8*x-4)/(x^2+2)^3,x, algorithm="fricas")`

output `-1/2*(sqrt(2)*(x^4 + 4*x^2 + 4)*arctan(1/2*sqrt(2)*x) - (x^4 + 4*x^2 + 4)*log(x^2 + 2))/(x^4 + 4*x^2 + 4)`

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.03

$$\int \frac{-4 + 8x - 4x^2 + 4x^3 - x^4 + x^5}{(2 + x^2)^3} dx = \frac{\log(x^2 + 2)}{2} - \frac{\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}x}{2}\right)}{2} - \frac{1}{x^4 + 4x^2 + 4}$$

input `integrate((x**5-x**4+4*x**3-4*x**2+8*x-4)/(x**2+2)**3,x)`output `log(x**2 + 2)/2 - sqrt(2)*atan(sqrt(2)*x/2)/2 - 1/(x**4 + 4*x**2 + 4)`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{-4 + 8x - 4x^2 + 4x^3 - x^4 + x^5}{(2 + x^2)^3} dx = -\frac{1}{2} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}x\right) - \frac{1}{x^4 + 4x^2 + 4} + \frac{1}{2} \log(x^2 + 2)$$

input `integrate((x^5-x^4+4*x^3-4*x^2+8*x-4)/(x^2+2)^3,x, algorithm="maxima")`output `-1/2*sqrt(2)*arctan(1/2*sqrt(2)*x) - 1/(x^4 + 4*x^2 + 4) + 1/2*log(x^2 + 2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.86

$$\int \frac{-4 + 8x - 4x^2 + 4x^3 - x^4 + x^5}{(2 + x^2)^3} dx = -\frac{1}{2} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}x\right) - \frac{1}{(x^2 + 2)^2} + \frac{1}{2} \log(x^2 + 2)$$

input `integrate((x^5-x^4+4*x^3-4*x^2+8*x-4)/(x^2+2)^3,x, algorithm="giac")`

output $-1/2*\text{sqrt}(2)*\text{arctan}(1/2*\text{sqrt}(2)*x) - 1/(x^2 + 2)^2 + 1/2*\log(x^2 + 2)$

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{-4 + 8x - 4x^2 + 4x^3 - x^4 + x^5}{(2 + x^2)^3} dx = \frac{\ln(x^2 + 2)}{2} - \frac{\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}x}{2}\right)}{2} - \frac{1}{x^4 + 4x^2 + 4}$$

input $\text{int}((8*x - 4*x^2 + 4*x^3 - x^4 + x^5 - 4)/(x^2 + 2)^3, x)$

output $\log(x^2 + 2)/2 - (2^{(1/2)}*\text{atan}(2^{(1/2)}*x/2))/2 - 1/(4*x^2 + x^4 + 4)$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 85, normalized size of antiderivative = 2.43

$$\int \frac{-4 + 8x - 4x^2 + 4x^3 - x^4 + x^5}{(2 + x^2)^3} dx$$

$$= \frac{-\sqrt{2} \operatorname{atan}\left(\frac{x}{\sqrt{2}}\right) x^4 - 4\sqrt{2} \operatorname{atan}\left(\frac{x}{\sqrt{2}}\right) x^2 - 4\sqrt{2} \operatorname{atan}\left(\frac{x}{\sqrt{2}}\right) + \log(x^2 + 2) x^4 + 4 \log(x^2 + 2) x^2 + 4 \log(x^2 + 2) - 2}{2x^4 + 8x^2 + 8}$$

input $\text{int}((x^5-x^4+4*x^3-4*x^2+8*x-4)/(x^2+2)^3, x)$

output $(- \text{sqrt}(2)*\text{atan}(x/\text{sqrt}(2))*x**4 - 4*\text{sqrt}(2)*\text{atan}(x/\text{sqrt}(2))*x**2 - 4*\text{sqrt}(2)*\text{atan}(x/\text{sqrt}(2)) + \log(x**2 + 2)*x**4 + 4*\log(x**2 + 2)*x**2 + 4*\log(x**2 + 2) - 2)/(2*(x**4 + 4*x**2 + 4))$

3.15 $\int \frac{-1-3x+x^2}{-2x+x^2+x^3} dx$

Optimal result	128
Mathematica [A] (verified)	128
Rubi [A] (verified)	129
Maple [A] (verified)	130
Fricas [A] (verification not implemented)	130
Sympy [A] (verification not implemented)	131
Maxima [A] (verification not implemented)	131
Giac [A] (verification not implemented)	131
Mupad [B] (verification not implemented)	132
Reduce [B] (verification not implemented)	132

Optimal result

Integrand size = 21, antiderivative size = 23

$$\int \frac{-1-3x+x^2}{-2x+x^2+x^3} dx = -\log(1-x) + \frac{\log(x)}{2} + \frac{3}{2}\log(2+x)$$

output

```
-ln(1-x)+1/2*ln(x)+3/2*ln(2+x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{-1-3x+x^2}{-2x+x^2+x^3} dx = -\log(1-x) + \frac{\log(x)}{2} + \frac{3}{2}\log(2+x)$$

input

```
Integrate[(-1 - 3*x + x^2)/(-2*x + x^2 + x^3), x]
```

output

```
-Log[1 - x] + Log[x]/2 + (3*Log[2 + x])/2
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2026, 2159, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^2 - 3x - 1}{x^3 + x^2 - 2x} dx \\ & \quad \downarrow \text{2026} \\ & \int \frac{x^2 - 3x - 1}{x(x^2 + x - 2)} dx \\ & \quad \downarrow \text{2159} \\ & \int \left(\frac{1}{2x} + \frac{3}{2(x+2)} + \frac{1}{1-x} \right) dx \\ & \quad \downarrow \text{2009} \\ & -\log(1-x) + \frac{\log(x)}{2} + \frac{3}{2}\log(x+2) \end{aligned}$$

input `Int[(-1 - 3*x + x^2)/(-2*x + x^2 + x^3),x]`

output `-Log[1 - x] + Log[x]/2 + (3*Log[2 + x])/2`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2159 `Int[(Pq_)*((d_.) + (e_)*(x_))^(m_)*((a_.) + (b_)*(x_) + (c_)*(x_)^2)^(p_.), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*Pq*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

method	result	size
default	$\frac{\ln(x)}{2} - \ln(-1 + x) + \frac{3\ln(2+x)}{2}$	18
norman	$\frac{\ln(x)}{2} - \ln(-1 + x) + \frac{3\ln(2+x)}{2}$	18
risch	$\frac{\ln(x)}{2} - \ln(-1 + x) + \frac{3\ln(2+x)}{2}$	18
parallelrisch	$\frac{\ln(x)}{2} - \ln(-1 + x) + \frac{3\ln(2+x)}{2}$	18

input `int((x^2-3*x-1)/(x^3+x^2-2*x),x,method=_RETURNVERBOSE)`

output `1/2*ln(x)-ln(-1+x)+3/2*ln(2+x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{-1 - 3x + x^2}{-2x + x^2 + x^3} dx = \frac{3}{2} \log(x + 2) - \log(x - 1) + \frac{1}{2} \log(x)$$

input `integrate((x^2-3*x-1)/(x^3+x^2-2*x),x, algorithm="fricas")`

output `3/2*log(x + 2) - log(x - 1) + 1/2*log(x)`

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{-1 - 3x + x^2}{-2x + x^2 + x^3} dx = \frac{\log(x)}{2} - \log(x - 1) + \frac{3\log(x + 2)}{2}$$

input `integrate((x**2-3*x-1)/(x**3+x**2-2*x),x)`output `log(x)/2 - log(x - 1) + 3*log(x + 2)/2`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{-1 - 3x + x^2}{-2x + x^2 + x^3} dx = \frac{3}{2} \log(x + 2) - \log(x - 1) + \frac{1}{2} \log(x)$$

input `integrate((x^2-3*x-1)/(x^3+x^2-2*x),x, algorithm="maxima")`output `3/2*log(x + 2) - log(x - 1) + 1/2*log(x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \frac{-1 - 3x + x^2}{-2x + x^2 + x^3} dx = \frac{3}{2} \log(|x + 2|) - \log(|x - 1|) + \frac{1}{2} \log(|x|)$$

input `integrate((x^2-3*x-1)/(x^3+x^2-2*x),x, algorithm="giac")`output `3/2*log(abs(x + 2)) - log(abs(x - 1)) + 1/2*log(abs(x))`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{-1 - 3x + x^2}{-2x + x^2 + x^3} dx = \frac{3 \ln(x + 2)}{2} - \ln(x - 1) + \frac{\ln(x)}{2}$$

input `int(-(3*x - x^2 + 1)/(x^2 - 2*x + x^3),x)`

output `(3*log(x + 2))/2 - log(x - 1) + log(x)/2`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{-1 - 3x + x^2}{-2x + x^2 + x^3} dx = -\log(x - 1) + \frac{3 \log(x + 2)}{2} + \frac{\log(x)}{2}$$

input `int((x^2-3*x-1)/(x^3+x^2-2*x),x)`

output `(- 2*log(x - 1) + 3*log(x + 2) + log(x))/2`

$$3.16 \quad \int \frac{3-x+3x^2-2x^3+x^4}{3x-2x^2+x^3} dx$$

Optimal result	133
Mathematica [A] (verified)	133
Rubi [A] (verified)	134
Maple [A] (verified)	135
Fricas [A] (verification not implemented)	135
Sympy [A] (verification not implemented)	136
Maxima [A] (verification not implemented)	136
Giac [A] (verification not implemented)	136
Mupad [B] (verification not implemented)	137
Reduce [B] (verification not implemented)	137

Optimal result

Integrand size = 33, antiderivative size = 23

$$\int \frac{3-x+3x^2-2x^3+x^4}{3x-2x^2+x^3} dx = \frac{x^2}{2} + \log(x) - \frac{1}{2} \log(3-2x+x^2)$$

output

```
1/2*x^2+ln(x)-1/2*ln(x^2-2*x+3)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{3-x+3x^2-2x^3+x^4}{3x-2x^2+x^3} dx = \frac{x^2}{2} + \log(x) - \frac{1}{2} \log(3-2x+x^2)$$

input

```
Integrate[(3 - x + 3*x^2 - 2*x^3 + x^4)/(3*x - 2*x^2 + x^3), x]
```

output

```
x^2/2 + Log[x] - Log[3 - 2*x + x^2]/2
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {2026, 2159, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4 - 2x^3 + 3x^2 - x + 3}{x^3 - 2x^2 + 3x} dx$$

↓ 2026

$$\int \frac{x^4 - 2x^3 + 3x^2 - x + 3}{x(x^2 - 2x + 3)} dx$$

↓ 2159

$$\int \left(\frac{1-x}{x^2 - 2x + 3} + x + \frac{1}{x} \right) dx$$

↓ 2009

$$\frac{x^2}{2} - \frac{1}{2} \log(x^2 - 2x + 3) + \log(x)$$

input `Int[(3 - x + 3*x^2 - 2*x^3 + x^4)/(3*x - 2*x^2 + x^3),x]`

output `x^2/2 + Log[x] - Log[3 - 2*x + x^2]/2`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2159

```
Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p
_), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*Pq*(a + b*x + c*x^2)^p, x
], x] /; FreeQ[{a, b, c, d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

method	result	size
default	$\frac{x^2}{2} + \ln(x) - \frac{\ln(x^2-2x+3)}{2}$	20
norman	$\frac{x^2}{2} + \ln(x) - \frac{\ln(x^2-2x+3)}{2}$	20
risch	$\frac{x^2}{2} + \ln(x) - \frac{\ln(x^2-2x+3)}{2}$	20
parallelrisch	$\frac{x^2}{2} + \ln(x) - \frac{\ln(x^2-2x+3)}{2}$	20

input

```
int((x^4-2*x^3+3*x^2-x+3)/(x^3-2*x^2+3*x),x,method=_RETURNVERBOSE)
```

output

```
1/2*x^2+ln(x)-1/2*ln(x^2-2*x+3)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{3 - x + 3x^2 - 2x^3 + x^4}{3x - 2x^2 + x^3} dx = \frac{1}{2}x^2 - \frac{1}{2}\log(x^2 - 2x + 3) + \log(x)$$

input

```
integrate((x^4-2*x^3+3*x^2-x+3)/(x^3-2*x^2+3*x),x, algorithm="fricas")
```

output

```
1/2*x^2 - 1/2*log(x^2 - 2*x + 3) + log(x)
```


Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{3 - x + 3x^2 - 2x^3 + x^4}{3x - 2x^2 + x^3} dx = \frac{x^2}{2} + \log(x) - \frac{\log(x^2 - 2x + 3)}{2}$$

input `integrate((x**4-2*x**3+3*x**2-x+3)/(x**3-2*x**2+3*x),x)`output `x**2/2 + log(x) - log(x**2 - 2*x + 3)/2`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{3 - x + 3x^2 - 2x^3 + x^4}{3x - 2x^2 + x^3} dx = \frac{1}{2}x^2 - \frac{1}{2}\log(x^2 - 2x + 3) + \log(x)$$

input `integrate((x^4-2*x^3+3*x^2-x+3)/(x^3-2*x^2+3*x),x, algorithm="maxima")`output `1/2*x^2 - 1/2*log(x^2 - 2*x + 3) + log(x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \frac{3 - x + 3x^2 - 2x^3 + x^4}{3x - 2x^2 + x^3} dx = \frac{1}{2}x^2 - \frac{1}{2}\log(x^2 - 2x + 3) + \log(|x|)$$

input `integrate((x^4-2*x^3+3*x^2-x+3)/(x^3-2*x^2+3*x),x, algorithm="giac")`output `1/2*x^2 - 1/2*log(x^2 - 2*x + 3) + log(abs(x))`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{3 - x + 3x^2 - 2x^3 + x^4}{3x - 2x^2 + x^3} dx = \ln(x) - \frac{\ln(x^2 - 2x + 3)}{2} + \frac{x^2}{2}$$

input `int((3*x^2 - x - 2*x^3 + x^4 + 3)/(3*x - 2*x^2 + x^3),x)`output `log(x) - log(x^2 - 2*x + 3)/2 + x^2/2`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{3 - x + 3x^2 - 2x^3 + x^4}{3x - 2x^2 + x^3} dx = -\frac{\log(x^2 - 2x + 3)}{2} + \log(x) + \frac{x^2}{2}$$

input `int((x^4-2*x^3+3*x^2-x+3)/(x^3-2*x^2+3*x),x)`output `(- log(x**2 - 2*x + 3) + 2*log(x) + x**2)/2`

$$3.17 \quad \int \frac{-1+x+x^3}{(1+x^2)^2} dx$$

Optimal result	138
Mathematica [A] (verified)	138
Rubi [A] (verified)	139
Maple [A] (verified)	140
Fricas [A] (verification not implemented)	141
Sympy [A] (verification not implemented)	141
Maxima [A] (verification not implemented)	141
Giac [A] (verification not implemented)	142
Mupad [B] (verification not implemented)	142
Reduce [B] (verification not implemented)	142

Optimal result

Integrand size = 14, antiderivative size = 29

$$\int \frac{-1+x+x^3}{(1+x^2)^2} dx = -\frac{x}{2(1+x^2)} - \frac{\arctan(x)}{2} + \frac{1}{2} \log(1+x^2)$$

output `-x/(2*x^2+2)-1/2*arctan(x)+1/2*ln(x^2+1)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.86

$$\int \frac{-1+x+x^3}{(1+x^2)^2} dx = \frac{1}{2} \left(-\frac{x}{1+x^2} - \arctan(x) + \log(1+x^2) \right)$$

input `Integrate[(-1 + x + x^3)/(1 + x^2)^2,x]`

output `(-(x/(1 + x^2)) - ArcTan[x] + Log[1 + x^2])/2`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.97, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2345, 452, 216, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^3 + x - 1}{(x^2 + 1)^2} dx \\ & \quad \downarrow \text{2345} \\ & -\frac{1}{2} \int \frac{1 - 2x}{x^2 + 1} dx - \frac{x}{2(x^2 + 1)} \\ & \quad \downarrow \text{452} \\ & \frac{1}{2} \left(2 \int \frac{x}{x^2 + 1} dx - \int \frac{1}{x^2 + 1} dx \right) - \frac{x}{2(x^2 + 1)} \\ & \quad \downarrow \text{216} \\ & \frac{1}{2} \left(2 \int \frac{x}{x^2 + 1} dx - \arctan(x) \right) - \frac{x}{2(x^2 + 1)} \\ & \quad \downarrow \text{240} \\ & \frac{1}{2} (\log(x^2 + 1) - \arctan(x)) - \frac{x}{2(x^2 + 1)} \end{aligned}$$

input

```
Int[(-1 + x + x^3)/(1 + x^2)^2,x]
```

output

```
-1/2*x/(1 + x^2) + (-ArcTan[x] + Log[1 + x^2])/2
```

Definitions of rubi rules used

rule 216 $\text{Int}[(a_ + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))* \text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

rule 240 $\text{Int}[(x_)/((a_ + (b_.)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x^2, x]]/(2*b), x] /; \text{FreeQ}\{a, b\}, x]$

rule 452 $\text{Int}(((c_ + (d_.)*(x_))/((a_ + (b_.)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[c \ \text{Int}[1/(a + b*x^2), x], x] + \text{Simp}[d \ \text{Int}[x/(a + b*x^2), x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[b*c^2 + a*d^2, 0]$

rule 2345 $\text{Int}[(Pq_)*((a_ + (b_.)*(x_)^2)^{p_}), x_Symbol] \rightarrow \text{With}\{Q = \text{PolynomialQuotient}[Pq, a + b*x^2, x], f = \text{Coeff}[\text{PolynomialRemainder}[Pq, a + b*x^2, x], x, 0], g = \text{Coeff}[\text{PolynomialRemainder}[Pq, a + b*x^2, x], x, 1]\}, \text{Simp}[(a*g - b*f*x)*((a + b*x^2)^{(p + 1})/(2*a*b*(p + 1))), x] + \text{Simp}[1/(2*a*(p + 1)) \ \text{Int}[(a + b*x^2)^{(p + 1)}*\text{ExpandToSum}[2*a*(p + 1)*Q + f*(2*p + 3), x], x], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{PolyQ}[Pq, x] \ \&\& \ \text{LtQ}[p, -1]$

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

method	result	size
default	$-\frac{x}{2(x^2+1)} + \frac{\ln(x^2+1)}{2} - \frac{\arctan(x)}{2}$	24
risch	$-\frac{x}{2(x^2+1)} + \frac{\ln(x^2+1)}{2} - \frac{\arctan(x)}{2}$	24
meijerg	$-\frac{x}{2x^2+2} - \frac{\arctan(x)}{2} + \frac{\ln(x^2+1)}{2}$	26
paralelrisch	$\frac{i \ln(x-i)x^2 - i \ln(x+i)x^2 + 2 \ln(x-i)x^2 + 2 \ln(x+i)x^2 + i \ln(x-i) - i \ln(x+i) + 2 \ln(x-i) + 2 \ln(x+i) - 2x}{4x^2+4}$	86

input $\text{int}((x^3+x-1)/(x^2+1)^2, x, \text{method}=_RETURNVERBOSE)$

output `-1/2*x/(x^2+1)+1/2*ln(x^2+1)-1/2*arctan(x)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.10

$$\int \frac{-1 + x + x^3}{(1 + x^2)^2} dx = -\frac{(x^2 + 1) \arctan(x) - (x^2 + 1) \log(x^2 + 1) + x}{2(x^2 + 1)}$$

input `integrate((x^3+x-1)/(x^2+1)^2,x, algorithm="fricas")`

output `-1/2*((x^2 + 1)*arctan(x) - (x^2 + 1)*log(x^2 + 1) + x)/(x^2 + 1)`

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.69

$$\int \frac{-1 + x + x^3}{(1 + x^2)^2} dx = -\frac{x}{2x^2 + 2} + \frac{\log(x^2 + 1)}{2} - \frac{\operatorname{atan}(x)}{2}$$

input `integrate((x**3+x-1)/(x**2+1)**2,x)`

output `-x/(2*x**2 + 2) + log(x**2 + 1)/2 - atan(x)/2`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.79

$$\int \frac{-1 + x + x^3}{(1 + x^2)^2} dx = -\frac{x}{2(x^2 + 1)} - \frac{1}{2} \arctan(x) + \frac{1}{2} \log(x^2 + 1)$$

input `integrate((x^3+x-1)/(x^2+1)^2,x, algorithm="maxima")`

output `-1/2*x/(x^2 + 1) - 1/2*arctan(x) + 1/2*log(x^2 + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.79

$$\int \frac{-1 + x + x^3}{(1 + x^2)^2} dx = -\frac{x}{2(x^2 + 1)} - \frac{1}{2} \arctan(x) + \frac{1}{2} \log(x^2 + 1)$$

input `integrate((x^3+x-1)/(x^2+1)^2,x, algorithm="giac")`

output `-1/2*x/(x^2 + 1) - 1/2*arctan(x) + 1/2*log(x^2 + 1)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.86

$$\int \frac{-1 + x + x^3}{(1 + x^2)^2} dx = \frac{\ln(x^2 + 1)}{2} - \frac{\operatorname{atan}(x)}{2} - \frac{x}{2(x^2 + 1)}$$

input `int((x + x^3 - 1)/(x^2 + 1)^2,x)`

output `log(x^2 + 1)/2 - atan(x)/2 - x/(2*(x^2 + 1))`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.41

$$\int \frac{-1 + x + x^3}{(1 + x^2)^2} dx = \frac{-\operatorname{atan}(x) x^2 - \operatorname{atan}(x) + \log(x^2 + 1) x^2 + \log(x^2 + 1) - x}{2x^2 + 2}$$

input `int((x^3+x-1)/(x^2+1)^2,x)`

output `(- atan(x)*x**2 - atan(x) + log(x**2 + 1)*x**2 + log(x**2 + 1) - x)/(2*(x**2 + 1))`

$$3.18 \quad \int \frac{1+2x-x^2+8x^3+x^4}{(x+x^2)(1+x^3)} dx$$

Optimal result	143
Mathematica [A] (verified)	143
Rubi [A] (verified)	144
Maple [A] (verified)	145
Fricas [A] (verification not implemented)	146
Sympy [A] (verification not implemented)	146
Maxima [A] (verification not implemented)	147
Giac [A] (verification not implemented)	147
Mupad [B] (verification not implemented)	147
Reduce [B] (verification not implemented)	148

Optimal result

Integrand size = 33, antiderivative size = 44

$$\int \frac{1+2x-x^2+8x^3+x^4}{(x+x^2)(1+x^3)} dx = -\frac{3}{1+x} - \frac{2 \arctan\left(\frac{1-2x}{\sqrt{3}}\right)}{\sqrt{3}} + \log(x) - 2 \log(1+x) + \log(1-x+x^2)$$

output

```
-3/(1+x)-2/3*arctan(1/3*(1-2*x)*3^(1/2))*3^(1/2)+ln(x)-2*ln(1+x)+ln(x^2-x+1)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00

$$\int \frac{1+2x-x^2+8x^3+x^4}{(x+x^2)(1+x^3)} dx = -\frac{3}{1+x} + \frac{2 \arctan\left(\frac{-1+2x}{\sqrt{3}}\right)}{\sqrt{3}} + \log(x) - 2 \log(1+x) + \log(1-x+x^2)$$

input

```
Integrate[(1 + 2*x - x^2 + 8*x^3 + x^4)/((x + x^2)*(1 + x^3)), x]
```


output

$$-3/(1+x) + (2*\text{ArcTan}[(-1+2*x)/\text{Sqrt}[3]])/\text{Sqrt}[3] + \text{Log}[x] - 2*\text{Log}[1+x] + \text{Log}[1-x+x^2]$$

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {2026, 7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^4 + 8x^3 - x^2 + 2x + 1}{(x^2 + x)(x^3 + 1)} dx \\ & \quad \downarrow \text{2026} \\ & \int \frac{x^4 + 8x^3 - x^2 + 2x + 1}{x(x+1)(x^3 + 1)} dx \\ & \quad \downarrow \text{7276} \\ & \int \left(\frac{2x}{x^2 - x + 1} - \frac{2}{x+1} + \frac{3}{(x+1)^2} + \frac{1}{x} \right) dx \\ & \quad \downarrow \text{2009} \\ & -\frac{2 \arctan\left(\frac{1-2x}{\sqrt{3}}\right)}{\sqrt{3}} + \log(x^2 - x + 1) - \frac{3}{x+1} + \log(x) - 2\log(x+1) \end{aligned}$$

input

$$\text{Int}[(1 + 2*x - x^2 + 8*x^3 + x^4)/((x + x^2)*(1 + x^3)), x]$$

output

$$-3/(1+x) - (2*\text{ArcTan}[(1-2*x)/\text{Sqrt}[3]])/\text{Sqrt}[3] + \text{Log}[x] - 2*\text{Log}[1+x] + \text{Log}[1-x+x^2]$$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.95

method	result	size
default	$\ln(x) - \frac{3}{1+x} - 2 \ln(1+x) + \ln(x^2 - x + 1) + \frac{2\sqrt{3} \arctan\left(\frac{(2x-1)\sqrt{3}}{3}\right)}{3}$	42
risch	$-\frac{3}{1+x} + \frac{2\sqrt{3} \arctan\left(\frac{(2x-1)\sqrt{3}}{3}\right)}{3} + \ln(4x^2 - 4x + 4) + \ln(x) - 2 \ln(1+x)$	44

input `int((x^4+8*x^3-x^2+2*x+1)/(x^2+x)/(x^3+1), x, method=_RETURNVERBOSE)`

output `ln(x)-3/(1+x)-2*ln(1+x)+ln(x^2-x+1)+2/3*3^(1/2)*arctan(1/3*(2*x-1)*3^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.32

$$\int \frac{1 + 2x - x^2 + 8x^3 + x^4}{(x + x^2)(1 + x^3)} dx$$

$$= \frac{2\sqrt{3}(x+1)\arctan\left(\frac{1}{3}\sqrt{3}(2x-1)\right) + 3(x+1)\log(x^2-x+1) - 6(x+1)\log(x+1) + 3(x+1)\log(x)}{3(x+1)}$$

input `integrate((x^4+8*x^3-x^2+2*x+1)/(x^2+x)/(x^3+1),x, algorithm="fricas")`

output `1/3*(2*sqrt(3)*(x + 1)*arctan(1/3*sqrt(3)*(2*x - 1)) + 3*(x + 1)*log(x^2 - x + 1) - 6*(x + 1)*log(x + 1) + 3*(x + 1)*log(x) - 9)/(x + 1)`

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.11

$$\int \frac{1 + 2x - x^2 + 8x^3 + x^4}{(x + x^2)(1 + x^3)} dx = \log(x) - 2\log(x + 1) + \log(x^2 - x + 1)$$

$$+ \frac{2\sqrt{3}\operatorname{atan}\left(\frac{2\sqrt{3}x}{3} - \frac{\sqrt{3}}{3}\right)}{3} - \frac{3}{x + 1}$$

input `integrate((x**4+8*x**3-x**2+2*x+1)/(x**2+x)/(x**3+1),x)`

output `log(x) - 2*log(x + 1) + log(x**2 - x + 1) + 2*sqrt(3)*atan(2*sqrt(3)*x/3 - sqrt(3)/3)/3 - 3/(x + 1)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.93

$$\int \frac{1 + 2x - x^2 + 8x^3 + x^4}{(x + x^2)(1 + x^3)} dx = \frac{2}{3} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3}(2x - 1) \right) - \frac{3}{x + 1} + \log(x^2 - x + 1) - 2 \log(x + 1) + \log(x)$$

input `integrate((x^4+8*x^3-x^2+2*x+1)/(x^2+x)/(x^3+1),x, algorithm="maxima")`output `2/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - 1)) - 3/(x + 1) + log(x^2 - x + 1) - 2*log(x + 1) + log(x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.98

$$\int \frac{1 + 2x - x^2 + 8x^3 + x^4}{(x + x^2)(1 + x^3)} dx = \frac{2}{3} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3}(2x - 1) \right) - \frac{3}{x + 1} + \log(x^2 - x + 1) - 2 \log(|x + 1|) + \log(|x|)$$

input `integrate((x^4+8*x^3-x^2+2*x+1)/(x^2+x)/(x^3+1),x, algorithm="giac")`output `2/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - 1)) - 3/(x + 1) + log(x^2 - x + 1) - 2*log(abs(x + 1)) + log(abs(x))`**Mupad [B] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.25

$$\int \frac{1 + 2x - x^2 + 8x^3 + x^4}{(x + x^2)(1 + x^3)} dx = \ln(x) - 2 \ln(x + 1) - \frac{3}{x + 1} - \ln \left(x - \frac{1}{2} - \frac{\sqrt{3} \text{li}}{2} \right) \left(-1 + \frac{\sqrt{3} \text{li}}{3} \right) + \ln \left(x - \frac{1}{2} + \frac{\sqrt{3} \text{li}}{2} \right) \left(1 + \frac{\sqrt{3} \text{li}}{3} \right)$$

input `int((2*x - x^2 + 8*x^3 + x^4 + 1)/((x^3 + 1)*(x + x^2)),x)`

output `log(x) - 2*log(x + 1) - 3/(x + 1) - log(x - (3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/3 - 1) + log(x + (3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/3 + 1)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 88, normalized size of antiderivative = 2.00

$$\int \frac{1 + 2x - x^2 + 8x^3 + x^4}{(x + x^2)(1 + x^3)} dx$$

$$= \frac{2\sqrt{3} \operatorname{atan}\left(\frac{2x-1}{\sqrt{3}}\right) x + 2\sqrt{3} \operatorname{atan}\left(\frac{2x-1}{\sqrt{3}}\right) + 3 \log(x^2 - x + 1) x + 3 \log(x^2 - x + 1) - 6 \log(x + 1) x - 6 \log(x + 1)}{3x + 3}$$

input `int((x^4+8*x^3-x^2+2*x+1)/(x^2+x)/(x^3+1),x)`

output `(2*sqrt(3)*atan((2*x - 1)/sqrt(3))*x + 2*sqrt(3)*atan((2*x - 1)/sqrt(3)) + 3*log(x**2 - x + 1)*x + 3*log(x**2 - x + 1) - 6*log(x + 1)*x - 6*log(x + 1) + 3*log(x)*x + 3*log(x) + 9*x)/(3*(x + 1))`

$$3.19 \quad \int \frac{15-5x+x^2+x^3}{(5+x^2)(3+2x+x^2)} dx$$

Optimal result	149
Mathematica [A] (verified)	149
Rubi [A] (verified)	150
Maple [A] (verified)	151
Fricas [A] (verification not implemented)	151
Sympy [A] (verification not implemented)	152
Maxima [A] (verification not implemented)	152
Giac [A] (verification not implemented)	153
Mupad [B] (verification not implemented)	153
Reduce [B] (verification not implemented)	154

Optimal result

Integrand size = 29, antiderivative size = 46

$$\int \frac{15-5x+x^2+x^3}{(5+x^2)(3+2x+x^2)} dx = -\sqrt{5} \arctan\left(\frac{x}{\sqrt{5}}\right) + \frac{5 \arctan\left(\frac{1+x}{\sqrt{2}}\right)}{\sqrt{2}} + \frac{1}{2} \log(3+2x+x^2)$$

output

```
-5^(1/2)*arctan(1/5*x*5^(1/2))+5/2*arctan(1/2*(1+x)*2^(1/2))*2^(1/2)+1/2*ln(x^2+2*x+3)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

$$\int \frac{15-5x+x^2+x^3}{(5+x^2)(3+2x+x^2)} dx = -\sqrt{5} \arctan\left(\frac{x}{\sqrt{5}}\right) + \frac{5 \arctan\left(\frac{1+x}{\sqrt{2}}\right)}{\sqrt{2}} + \frac{1}{2} \log(3+2x+x^2)$$

input

```
Integrate[(15 - 5*x + x^2 + x^3)/((5 + x^2)*(3 + 2*x + x^2)),x]
```

output

```
-(Sqrt[5]*ArcTan[x/Sqrt[5]]) + (5*ArcTan[(1 + x)/Sqrt[2]])/Sqrt[2] + Log[3 + 2*x + x^2]/2
```

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.069$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3 + x^2 - 5x + 15}{(x^2 + 5)(x^2 + 2x + 3)} dx$$

$$\downarrow 7276$$

$$\int \left(\frac{x + 6}{x^2 + 2x + 3} - \frac{5}{x^2 + 5} \right) dx$$

$$\downarrow 2009$$

$$-\sqrt{5} \arctan\left(\frac{x}{\sqrt{5}}\right) + \frac{5 \arctan\left(\frac{x+1}{\sqrt{2}}\right)}{\sqrt{2}} + \frac{1}{2} \log(x^2 + 2x + 3)$$

input `Int[(15 - 5*x + x^2 + x^3)/((5 + x^2)*(3 + 2*x + x^2)),x]`

output `-(Sqrt[5]*ArcTan[x/Sqrt[5]]) + (5*ArcTan[(1 + x)/Sqrt[2]])/Sqrt[2] + Log[3 + 2*x + x^2]/2`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] :> With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]`

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.85

method	result	size
risch	$-\sqrt{5} \arctan\left(\frac{x\sqrt{5}}{5}\right) + \frac{5 \arctan\left(\frac{(1+x)\sqrt{2}}{2}\right)\sqrt{2}}{2} + \frac{\ln(x^2+2x+3)}{2}$	39
default	$\frac{\ln(x^2+2x+3)}{2} + \frac{5\sqrt{2} \arctan\left(\frac{(2x+2)\sqrt{2}}{4}\right)}{2} - \sqrt{5} \arctan\left(\frac{x\sqrt{5}}{5}\right)$	41

input `int((x^3+x^2-5*x+15)/(x^2+5)/(x^2+2*x+3),x,method=_RETURNVERBOSE)`

output `-5^(1/2)*arctan(1/5*x*5^(1/2))+5/2*arctan(1/2*(1+x)*2^(1/2))*2^(1/2)+1/2*ln(x^2+2*x+3)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.83

$$\int \frac{15 - 5x + x^2 + x^3}{(5 + x^2)(3 + 2x + x^2)} dx = \frac{5}{2} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}(x + 1)\right) - \sqrt{5} \arctan\left(\frac{1}{5} \sqrt{5}x\right) + \frac{1}{2} \log(x^2 + 2x + 3)$$

input `integrate((x^3+x^2-5*x+15)/(x^2+5)/(x^2+2*x+3),x, algorithm="fricas")`

output `5/2*sqrt(2)*arctan(1/2*sqrt(2)*(x + 1)) - sqrt(5)*arctan(1/5*sqrt(5)*x) + 1/2*log(x^2 + 2*x + 3)`

Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.11

$$\int \frac{15 - 5x + x^2 + x^3}{(5 + x^2)(3 + 2x + x^2)} dx = \frac{\log(x^2 + 2x + 3)}{2} - \sqrt{5} \operatorname{atan}\left(\frac{\sqrt{5}x}{5}\right) + \frac{5\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}x}{2} + \frac{\sqrt{2}}{2}\right)}{2}$$

input `integrate((x**3+x**2-5*x+15)/(x**2+5)/(x**2+2*x+3),x)`output `log(x**2 + 2*x + 3)/2 - sqrt(5)*atan(sqrt(5)*x/5) + 5*sqrt(2)*atan(sqrt(2)*x/2 + sqrt(2)/2)/2`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.83

$$\int \frac{15 - 5x + x^2 + x^3}{(5 + x^2)(3 + 2x + x^2)} dx = \frac{5}{2} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}(x + 1)\right) - \sqrt{5} \arctan\left(\frac{1}{5} \sqrt{5}x\right) + \frac{1}{2} \log(x^2 + 2x + 3)$$

input `integrate((x^3+x^2-5*x+15)/(x^2+5)/(x^2+2*x+3),x, algorithm="maxima")`output `5/2*sqrt(2)*arctan(1/2*sqrt(2)*(x + 1)) - sqrt(5)*arctan(1/5*sqrt(5)*x) + 1/2*log(x^2 + 2*x + 3)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.83

$$\int \frac{15 - 5x + x^2 + x^3}{(5 + x^2)(3 + 2x + x^2)} dx = \frac{5}{2} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}(x + 1)\right) - \sqrt{5} \arctan\left(\frac{1}{5} \sqrt{5}x\right) + \frac{1}{2} \log(x^2 + 2x + 3)$$

input `integrate((x^3+x^2-5*x+15)/(x^2+5)/(x^2+2*x+3),x, algorithm="giac")`

output `5/2*sqrt(2)*arctan(1/2*sqrt(2)*(x + 1)) - sqrt(5)*arctan(1/5*sqrt(5)*x) + 1/2*log(x^2 + 2*x + 3)`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.91

$$\int \frac{15 - 5x + x^2 + x^3}{(5 + x^2)(3 + 2x + x^2)} dx = \frac{\ln(x + 1 - \sqrt{2}1i)}{2} + \frac{\ln(x + 1 + \sqrt{2}1i)}{2} + \sqrt{5} \operatorname{atan}\left(\frac{2000\sqrt{5}}{2000x + 1120} - \frac{224\sqrt{5}x}{2000x + 1120}\right) - \frac{\sqrt{2} \ln(x + 1 - \sqrt{2}1i) 5i}{4} + \frac{\sqrt{2} \ln(x + 1 + \sqrt{2}1i) 5i}{4}$$

input `int((x^2 - 5*x + x^3 + 15)/((x^2 + 5)*(2*x + x^2 + 3)),x)`

output `log(x - 2^(1/2)*1i + 1)/2 + log(x + 2^(1/2)*1i + 1)/2 + 5^(1/2)*atan((2000*5^(1/2))/(2000*x + 1120) - (224*5^(1/2)*x)/(2000*x + 1120)) - (2^(1/2)*log(x - 2^(1/2)*1i + 1)*5i)/4 + (2^(1/2)*log(x + 2^(1/2)*1i + 1)*5i)/4`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.78

$$\int \frac{15 - 5x + x^2 + x^3}{(5 + x^2)(3 + 2x + x^2)} dx = \frac{5\sqrt{2} \operatorname{atan}\left(\frac{x+1}{\sqrt{2}}\right)}{2} - \sqrt{5} \operatorname{atan}\left(\frac{x}{\sqrt{5}}\right) + \frac{\log(x^2 + 2x + 3)}{2}$$

input `int((x^3+x^2-5*x+15)/(x^2+5)/(x^2+2*x+3),x)`

output `(5*sqrt(2)*atan((x + 1)/sqrt(2)) - 2*sqrt(5)*atan(x/sqrt(5)) + log(x**2 + 2*x + 3))/2`

$$3.20 \quad \int \frac{-3+25x+23x^2+32x^3+15x^4+7x^5+x^6}{(1+x^2)^2(2+x+x^2)^2} dx$$

Optimal result	155
Mathematica [A] (verified)	155
Rubi [A] (verified)	156
Maple [A] (verified)	157
Fricas [B] (verification not implemented)	157
Sympy [A] (verification not implemented)	158
Maxima [A] (verification not implemented)	158
Giac [A] (verification not implemented)	159
Mupad [B] (verification not implemented)	159
Reduce [B] (verification not implemented)	160

Optimal result

Integrand size = 44, antiderivative size = 33

$$\int \frac{-3 + 25x + 23x^2 + 32x^3 + 15x^4 + 7x^5 + x^6}{(1 + x^2)^2 (2 + x + x^2)^2} dx$$

$$= -\frac{3}{1 + x^2} + \frac{1}{2 + x + x^2} + \log(1 + x^2) - \log(2 + x + x^2)$$

output

```
-3/(x^2+1)+1/(x^2+x+2)+ln(x^2+1)-ln(x^2+x+2)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int \frac{-3 + 25x + 23x^2 + 32x^3 + 15x^4 + 7x^5 + x^6}{(1 + x^2)^2 (2 + x + x^2)^2} dx$$

$$= -\frac{3}{1 + x^2} + \frac{1}{2 + x + x^2} + \log(1 + x^2) - \log(2 + x + x^2)$$

input

```
Integrate[(-3 + 25*x + 23*x^2 + 32*x^3 + 15*x^4 + 7*x^5 + x^6)/((1 + x^2)^2*(2 + x + x^2)^2), x]
```

output $-3/(1 + x^2) + (2 + x + x^2)^{-1} + \text{Log}[1 + x^2] - \text{Log}[2 + x + x^2]$

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.045$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^6 + 7x^5 + 15x^4 + 32x^3 + 23x^2 + 25x - 3}{(x^2 + 1)^2 (x^2 + x + 2)^2} dx$$

↓ 7293

$$\int \left(\frac{-2x - 1}{x^2 + x + 2} + \frac{-2x - 1}{(x^2 + x + 2)^2} + \frac{2x}{x^2 + 1} + \frac{6x}{(x^2 + 1)^2} \right) dx$$

↓ 2009

$$-\frac{3}{x^2 + 1} + \frac{1}{x^2 + x + 2} + \log(x^2 + 1) - \log(x^2 + x + 2)$$

input `Int[(-3 + 25*x + 23*x^2 + 32*x^3 + 15*x^4 + 7*x^5 + x^6)/((1 + x^2)^2*(2 + x + x^2)^2),x]`

output $-3/(1 + x^2) + (2 + x + x^2)^{-1} + \text{Log}[1 + x^2] - \text{Log}[2 + x + x^2]$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.03

method	result
default	$-\frac{3}{x^2+1} + \frac{1}{x^2+x+2} + \ln(x^2+1) - \ln(x^2+x+2)$
norman	$\frac{-2x^2-3x-5}{(x^2+x+2)(x^2+1)} - \ln(x^2+x+2) + \ln(x^2+1)$
risch	$\frac{-2x^2-3x-5}{(x^2+x+2)(x^2+1)} - \ln(x^2+x+2) + \ln(x^2+1)$
parallelrisch	$\frac{\ln(x^2+1)x^4 - \ln(x^2+x+2)x^4 - 5 + \ln(x^2+1)x^3 - \ln(x^2+x+2)x^3 + 3\ln(x^2+1)x^2 - 3\ln(x^2+x+2)x^2 + \ln(x^2+1)x - \ln(x^2+x+2)}{(x^2+1)(x^2+x+2)}$

input `int((x^6+7*x^5+15*x^4+32*x^3+23*x^2+25*x-3)/(x^2+1)^2/(x^2+x+2)^2,x,method =_RETURNVERBOSE)`

output `-3/(x^2+1)+1/(x^2+x+2)+ln(x^2+1)-ln(x^2+x+2)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 72 vs. 2(33) = 66.

Time = 0.07 (sec) , antiderivative size = 72, normalized size of antiderivative = 2.18

$$\int \frac{-3 + 25x + 23x^2 + 32x^3 + 15x^4 + 7x^5 + x^6}{(1+x^2)^2(2+x+x^2)^2} dx =$$

$$\frac{2x^2 + (x^4 + x^3 + 3x^2 + x + 2) \log(x^2 + x + 2) - (x^4 + x^3 + 3x^2 + x + 2) \log(x^2 + 1) + 3x + 5}{x^4 + x^3 + 3x^2 + x + 2}$$

input `integrate((x^6+7*x^5+15*x^4+32*x^3+23*x^2+25*x-3)/(x^2+1)^2/(x^2+x+2)^2,x,algorithm="fricas")`

output `-(2*x^2 + (x^4 + x^3 + 3*x^2 + x + 2)*log(x^2 + x + 2) - (x^4 + x^3 + 3*x^2 + x + 2)*log(x^2 + 1) + 3*x + 5)/(x^4 + x^3 + 3*x^2 + x + 2)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.24

$$\int \frac{-3 + 25x + 23x^2 + 32x^3 + 15x^4 + 7x^5 + x^6}{(1 + x^2)^2 (2 + x + x^2)^2} dx$$

$$= \frac{-2x^2 - 3x - 5}{x^4 + x^3 + 3x^2 + x + 2} + \log(x^2 + 1) - \log(x^2 + x + 2)$$

input `integrate((x**6+7*x**5+15*x**4+32*x**3+23*x**2+25*x-3)/(x**2+1)**2/(x**2+x+2)**2,x)`

output `(-2*x**2 - 3*x - 5)/(x**4 + x**3 + 3*x**2 + x + 2) + log(x**2 + 1) - log(x**2 + x + 2)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.33

$$\int \frac{-3 + 25x + 23x^2 + 32x^3 + 15x^4 + 7x^5 + x^6}{(1 + x^2)^2 (2 + x + x^2)^2} dx$$

$$= -\frac{2x^2 + 3x + 5}{x^4 + x^3 + 3x^2 + x + 2} - \log(x^2 + x + 2) + \log(x^2 + 1)$$

input `integrate((x^6+7*x^5+15*x^4+32*x^3+23*x^2+25*x-3)/(x^2+1)^2/(x^2+x+2)^2,x,algorithm="maxima")`

output `-(2*x^2 + 3*x + 5)/(x^4 + x^3 + 3*x^2 + x + 2) - log(x^2 + x + 2) + log(x^2 + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.33

$$\int \frac{-3 + 25x + 23x^2 + 32x^3 + 15x^4 + 7x^5 + x^6}{(1 + x^2)^2 (2 + x + x^2)^2} dx$$

$$= -\frac{2x^2 + 3x + 5}{x^4 + x^3 + 3x^2 + x + 2} - \log(x^2 + x + 2) + \log(x^2 + 1)$$

input

```
integrate((x^6+7*x^5+15*x^4+32*x^3+23*x^2+25*x-3)/(x^2+1)^2/(x^2+x+2)^2,x,
algorithm="giac")
```

output

```
-(2*x^2 + 3*x + 5)/(x^4 + x^3 + 3*x^2 + x + 2) - log(x^2 + x + 2) + log(x^2 + 1)
```

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.70

$$\int \frac{-3 + 25x + 23x^2 + 32x^3 + 15x^4 + 7x^5 + x^6}{(1 + x^2)^2 (2 + x + x^2)^2} dx$$

$$= -\frac{2x^2 + 3x + 5}{x^4 + x^3 + 3x^2 + x + 2} + \operatorname{atan}\left(\frac{\frac{x224i}{11} + \frac{224i}{11}}{44x^2 + 16x + 60} - \frac{3}{11}i\right) 2i$$

input

```
int((25*x + 23*x^2 + 32*x^3 + 15*x^4 + 7*x^5 + x^6 - 3)/((x^2 + 1)^2*(x + x^2 + 2)^2),x)
```

output

```
atan(((x*224i)/11 + 224i/11)/(16*x + 44*x^2 + 60) - 3i/11)*2i - (3*x + 2*x^2 + 5)/(x + 3*x^2 + x^3 + x^4 + 2)
```


Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 129, normalized size of antiderivative = 3.91

$$\int \frac{-3 + 25x + 23x^2 + 32x^3 + 15x^4 + 7x^5 + x^6}{(1 + x^2)^2 (2 + x + x^2)^2} dx$$

$$= \frac{-\log(x^2 + x + 2) x^4 - \log(x^2 + x + 2) x^3 - 3 \log(x^2 + x + 2) x^2 - \log(x^2 + x + 2) x - 2 \log(x^2 + x + 2)}{x^4 + x^3 + 3x^2 + 2x + 2}$$

input

```
int((x^6+7*x^5+15*x^4+32*x^3+23*x^2+25*x-3)/(x^2+1)^2/(x^2+x+2)^2,x)
```

output

```
( - log(x**2 + x + 2)*x**4 - log(x**2 + x + 2)*x**3 - 3*log(x**2 + x + 2)*
x**2 - log(x**2 + x + 2)*x - 2*log(x**2 + x + 2) + log(x**2 + 1)*x**4 + lo
g(x**2 + 1)*x**3 + 3*log(x**2 + 1)*x**2 + log(x**2 + 1)*x + 2*log(x**2 + 1
) - 2*x**2 - 3*x - 5)/(x**4 + x**3 + 3*x**2 + x + 2)
```

3.21 $\int \frac{1}{(1+x^2)(4+x^2)} dx$

Optimal result	161
Mathematica [A] (verified)	161
Rubi [A] (verified)	162
Maple [A] (verified)	163
Fricas [A] (verification not implemented)	163
Sympy [A] (verification not implemented)	163
Maxima [A] (verification not implemented)	164
Giac [A] (verification not implemented)	164
Mupad [B] (verification not implemented)	164
Reduce [B] (verification not implemented)	165

Optimal result

Integrand size = 15, antiderivative size = 17

$$\int \frac{1}{(1+x^2)(4+x^2)} dx = -\frac{1}{6} \arctan\left(\frac{x}{2}\right) + \frac{\arctan(x)}{3}$$

output `-1/6*arctan(1/2*x)+1/3*arctan(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1+x^2)(4+x^2)} dx = \frac{1}{6} \arctan\left(\frac{2}{x}\right) + \frac{\arctan(x)}{3}$$

input `Integrate[1/((1 + x^2)*(4 + x^2)),x]`

output `ArcTan[2/x]/6 + ArcTan[x]/3`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {303, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(x^2 + 1)(x^2 + 4)} dx$$

↓ 303

$$\frac{1}{3} \int \frac{1}{x^2 + 1} dx - \frac{1}{3} \int \frac{1}{x^2 + 4} dx$$

↓ 216

$$\frac{\arctan(x)}{3} - \frac{1}{6} \arctan\left(\frac{x}{2}\right)$$

input `Int[1/((1 + x^2)*(4 + x^2)),x]`

output `-1/6*ArcTan[x/2] + ArcTan[x]/3`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 303 `Int[1/(((a_) + (b_)*(x_)^2)*((c_) + (d_)*(x_)^2)), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x^2), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

method	result	size
default	$-\frac{\arctan(\frac{x}{2})}{6} + \frac{\arctan(x)}{3}$	12
risch	$-\frac{\arctan(\frac{x}{2})}{6} + \frac{\arctan(x)}{3}$	12
parallelrisc	$\frac{i \ln(x+i)}{6} - \frac{i \ln(x-i)}{6} + \frac{i \ln(x-2i)}{12} - \frac{i \ln(x+2i)}{12}$	34

input `int(1/(x^2+1)/(x^2+4),x,method=_RETURNVERBOSE)`output `-1/6*arctan(1/2*x)+1/3*arctan(x)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int \frac{1}{(1+x^2)(4+x^2)} dx = -\frac{1}{6} \arctan\left(\frac{1}{2}x\right) + \frac{1}{3} \arctan(x)$$

input `integrate(1/(x^2+1)/(x^2+4),x, algorithm="fricas")`output `-1/6*arctan(1/2*x) + 1/3*arctan(x)`**Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.59

$$\int \frac{1}{(1+x^2)(4+x^2)} dx = -\frac{\operatorname{atan}\left(\frac{x}{2}\right)}{6} + \frac{\operatorname{atan}(x)}{3}$$

input `integrate(1/(x**2+1)/(x**2+4),x)`

output `-atan(x/2)/6 + atan(x)/3`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int \frac{1}{(1+x^2)(4+x^2)} dx = -\frac{1}{6} \arctan\left(\frac{1}{2}x\right) + \frac{1}{3} \arctan(x)$$

input `integrate(1/(x^2+1)/(x^2+4),x, algorithm="maxima")`

output `-1/6*arctan(1/2*x) + 1/3*arctan(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int \frac{1}{(1+x^2)(4+x^2)} dx = -\frac{1}{6} \arctan\left(\frac{1}{2}x\right) + \frac{1}{3} \arctan(x)$$

input `integrate(1/(x^2+1)/(x^2+4),x, algorithm="giac")`

output `-1/6*arctan(1/2*x) + 1/3*arctan(x)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int \frac{1}{(1+x^2)(4+x^2)} dx = \frac{\operatorname{atan}(x)}{3} - \frac{\operatorname{atan}\left(\frac{x}{2}\right)}{6}$$

input `int(1/((x^2 + 1)*(x^2 + 4)),x)`

output `atan(x)/3 - atan(x/2)/6`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int \frac{1}{(1+x^2)(4+x^2)} dx = -\frac{\operatorname{atan}\left(\frac{x}{2}\right)}{6} + \frac{\operatorname{atan}(x)}{3}$$

input `int(1/(x^2+1)/(x^2+4),x)`

output `(- atan(x/2) + 2*atan(x))/6`

3.22 $\int \frac{a+bx^3}{1+x^2} dx$

Optimal result	166
Mathematica [A] (verified)	166
Rubi [A] (verified)	167
Maple [A] (verified)	168
Fricas [A] (verification not implemented)	168
Sympy [C] (verification not implemented)	168
Maxima [A] (verification not implemented)	169
Giac [A] (verification not implemented)	169
Mupad [B] (verification not implemented)	170
Reduce [B] (verification not implemented)	170

Optimal result

Integrand size = 15, antiderivative size = 24

$$\int \frac{a + bx^3}{1 + x^2} dx = \frac{bx^2}{2} + a \arctan(x) - \frac{1}{2}b \log(1 + x^2)$$

output `1/2*b*x^2+a*arctan(x)-1/2*b*ln(x^2+1)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{a + bx^3}{1 + x^2} dx = a \arctan(x) + \frac{1}{2}b(x^2 - \log(1 + x^2))$$

input `Integrate[(a + b*x^3)/(1 + x^2),x]`

output `a*ArcTan[x] + (b*(x^2 - Log[1 + x^2]))/2`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2341, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + bx^3}{x^2 + 1} dx$$

$$\downarrow \text{2341}$$

$$\int \left(\frac{a - bx}{x^2 + 1} + bx \right) dx$$

$$\downarrow \text{2009}$$

$$a \arctan(x) + \frac{bx^2}{2} - \frac{1}{2}b \log(x^2 + 1)$$

input

```
Int[(a + b*x^3)/(1 + x^2),x]
```

output

```
(b*x^2)/2 + a*ArcTan[x] - (b*Log[1 + x^2])/2
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2341

```
Int[(Pq_)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[Pq*(a + b*x^2)^p, x], x] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && IGtQ[p, -2]
```


Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.88

method	result	size
default	$\frac{bx^2}{2} + a \arctan(x) - \frac{b \ln(x^2+1)}{2}$	21
meijerg	$\frac{b(x^2 - \ln(x^2+1))}{2} + a \arctan(x)$	21
risch	$\frac{bx^2}{2} + a \arctan(x) - \frac{b \ln(x^2+1)}{2}$	21
parallelrisch	$\frac{bx^2}{2} - \frac{\ln(x-i)b}{2} - \frac{i \ln(x-i)a}{2} - \frac{\ln(x+i)b}{2} + \frac{i \ln(x+i)a}{2}$	42

input `int((b*x^3+a)/(x^2+1),x,method=_RETURNVERBOSE)`

output `1/2*b*x^2+a*arctan(x)-1/2*b*ln(x^2+1)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{a + bx^3}{1 + x^2} dx = \frac{1}{2} bx^2 + a \arctan(x) - \frac{1}{2} b \log(x^2 + 1)$$

input `integrate((b*x^3+a)/(x^2+1),x, algorithm="fricas")`

output `1/2*b*x^2 + a*arctan(x) - 1/2*b*log(x^2 + 1)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.07 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.42

$$\int \frac{a + bx^3}{1 + x^2} dx = \frac{bx^2}{2} + \left(-\frac{ia}{2} - \frac{b}{2}\right) \log(x - i) + \left(\frac{ia}{2} - \frac{b}{2}\right) \log(x + i)$$

input `integrate((b*x**3+a)/(x**2+1),x)`

output `b*x**2/2 + (-I*a/2 - b/2)*log(x - I) + (I*a/2 - b/2)*log(x + I)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{a + bx^3}{1 + x^2} dx = \frac{1}{2} bx^2 + a \arctan(x) - \frac{1}{2} b \log(x^2 + 1)$$

input `integrate((b*x^3+a)/(x^2+1),x, algorithm="maxima")`

output `1/2*b*x^2 + a*arctan(x) - 1/2*b*log(x^2 + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{a + bx^3}{1 + x^2} dx = \frac{1}{2} bx^2 + a \arctan(x) - \frac{1}{2} b \log(x^2 + 1)$$

input `integrate((b*x^3+a)/(x^2+1),x, algorithm="giac")`

output `1/2*b*x^2 + a*arctan(x) - 1/2*b*log(x^2 + 1)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{a + bx^3}{1 + x^2} dx = \frac{bx^2}{2} - \frac{b \ln(x^2 + 1)}{2} + a \operatorname{atan}(x)$$

input `int((a + b*x^3)/(x^2 + 1),x)`

output `(b*x^2)/2 - (b*log(x^2 + 1))/2 + a*atan(x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{a + bx^3}{1 + x^2} dx = \operatorname{atan}(x) a - \frac{\log(x^2 + 1) b}{2} + \frac{bx^2}{2}$$

input `int((b*x^3+a)/(x^2+1),x)`

output `(2*atan(x)*a - log(x**2 + 1)*b + b*x**2)/2`

3.23 $\int \frac{x+x^2}{(4+x)(-4+x^2)} dx$

Optimal result	171
Mathematica [A] (verified)	171
Rubi [A] (verified)	172
Maple [A] (verified)	173
Fricas [A] (verification not implemented)	173
Sympy [A] (verification not implemented)	174
Maxima [A] (verification not implemented)	174
Giac [A] (verification not implemented)	174
Mupad [B] (verification not implemented)	175
Reduce [B] (verification not implemented)	175

Optimal result

Integrand size = 18, antiderivative size = 15

$$\int \frac{x+x^2}{(4+x)(-4+x^2)} dx = -\frac{1}{2} \operatorname{arctanh}\left(\frac{x}{2}\right) + \log(4+x)$$

output

```
-1/2*arctanh(1/2*x)+ln(4+x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.53

$$\int \frac{x+x^2}{(4+x)(-4+x^2)} dx = \frac{1}{4} \log(2-x) - \frac{1}{4} \log(2+x) + \log(4+x)$$

input

```
Integrate[(x + x^2)/((4 + x)*(-4 + x^2)), x]
```

output

```
Log[2 - x]/4 - Log[2 + x]/4 + Log[4 + x]
```

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2027, 2160, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^2 + x}{(x + 4)(x^2 - 4)} dx \\ & \quad \downarrow \text{2027} \\ & \int \frac{x(x + 1)}{(x + 4)(x^2 - 4)} dx \\ & \quad \downarrow \text{2160} \\ & \int \left(\frac{1}{x^2 - 4} + \frac{1}{x + 4} \right) dx \\ & \quad \downarrow \text{2009} \\ & \log(x + 4) - \frac{1}{2} \operatorname{arctanh}\left(\frac{x}{2}\right) \end{aligned}$$

input `Int[(x + x^2)/((4 + x)*(-4 + x^2)),x]`

output `-1/2*ArcTanh[x/2] + Log[4 + x]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2027 `Int[(F*x_.)*((a_.)*(x_)^(r_.) + (b_.)*(x_)^(s_.))^(p_.), x_Symbol] := Int[x^(p*r)*(a + b*x^(s - r))^p*Fx, x] /; FreeQ[{a, b, r, s}, x] && IntegerQ[p] && PosQ[s - r] && !(EqQ[p, 1] && EqQ[u, 1])`

rule 2160

```
Int[(Pq_)*((d_) + (e_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol]
:> Int[ExpandIntegrand[(d + e*x)^m*Pq*(a + b*x^2)^p, x], x] /; FreeQ[{a, b,
d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.20

method	result	size
default	$\ln(x+4) - \frac{\ln(2+x)}{4} + \frac{\ln(-2+x)}{4}$	18
norman	$\ln(x+4) - \frac{\ln(2+x)}{4} + \frac{\ln(-2+x)}{4}$	18
risch	$\ln(x+4) - \frac{\ln(2+x)}{4} + \frac{\ln(-2+x)}{4}$	18
parallelrisch	$\ln(x+4) - \frac{\ln(2+x)}{4} + \frac{\ln(-2+x)}{4}$	18

input

```
int((x^2+x)/(x+4)/(x^2-4),x,method=_RETURNVERBOSE)
```

output

```
ln(x+4)-1/4*ln(2+x)+1/4*ln(-2+x)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{x + x^2}{(4 + x)(-4 + x^2)} dx = \log(x + 4) - \frac{1}{4} \log(x + 2) + \frac{1}{4} \log(x - 2)$$

input

```
integrate((x^2+x)/(4+x)/(x^2-4),x, algorithm="fricas")
```

output

```
log(x + 4) - 1/4*log(x + 2) + 1/4*log(x - 2)
```

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{x + x^2}{(4 + x)(-4 + x^2)} dx = \frac{\log(x - 2)}{4} - \frac{\log(x + 2)}{4} + \log(x + 4)$$

input `integrate((x**2+x)/(4+x)/(x**2-4),x)`output `log(x - 2)/4 - log(x + 2)/4 + log(x + 4)`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{x + x^2}{(4 + x)(-4 + x^2)} dx = \log(x + 4) - \frac{1}{4} \log(x + 2) + \frac{1}{4} \log(x - 2)$$

input `integrate((x^2+x)/(4+x)/(x^2-4),x, algorithm="maxima")`output `log(x + 4) - 1/4*log(x + 2) + 1/4*log(x - 2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.33

$$\int \frac{x + x^2}{(4 + x)(-4 + x^2)} dx = \log(|x + 4|) - \frac{1}{4} \log(|x + 2|) + \frac{1}{4} \log(|x - 2|)$$

input `integrate((x^2+x)/(4+x)/(x^2-4),x, algorithm="giac")`output `log(abs(x + 4)) - 1/4*log(abs(x + 2)) + 1/4*log(abs(x - 2))`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.27

$$\int \frac{x + x^2}{(4 + x)(-4 + x^2)} dx = \ln(x + 4) + \frac{\operatorname{atanh}\left(\frac{90}{7(21x+48)} - \frac{8}{7}\right)}{2}$$

input `int((x + x^2)/((x^2 - 4)*(x + 4)),x)`output `log(x + 4) + atanh(90/(7*(21*x + 48)) - 8/7)/2`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{x + x^2}{(4 + x)(-4 + x^2)} dx = \frac{\log(x - 2)}{4} + \log(x + 4) - \frac{\log(x + 2)}{4}$$

input `int((x^2+x)/(4+x)/(x^2-4),x)`output `(log(x - 2) + 4*log(x + 4) - log(x + 2))/4`

3.24 $\int \frac{4+x^2}{(1+x^2)(2+x^2)} dx$

Optimal result	176
Mathematica [A] (verified)	176
Rubi [A] (verified)	177
Maple [A] (verified)	178
Fricas [A] (verification not implemented)	178
Sympy [A] (verification not implemented)	178
Maxima [A] (verification not implemented)	179
Giac [A] (verification not implemented)	179
Mupad [B] (verification not implemented)	179
Reduce [B] (verification not implemented)	180

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{4+x^2}{(1+x^2)(2+x^2)} dx = 3 \arctan(x) - \sqrt{2} \arctan\left(\frac{x}{\sqrt{2}}\right)$$

output `3*arctan(x)-arctan(1/2*x*2^(1/2))*2^(1/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{4+x^2}{(1+x^2)(2+x^2)} dx = 3 \arctan(x) - \sqrt{2} \arctan\left(\frac{x}{\sqrt{2}}\right)$$

input `Integrate[(4 + x^2)/((1 + x^2)*(2 + x^2)), x]`

output `3*ArcTan[x] - Sqrt[2]*ArcTan[x/Sqrt[2]]`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {397, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2 + 4}{(x^2 + 1)(x^2 + 2)} dx$$

$$\downarrow 397$$

$$3 \int \frac{1}{x^2 + 1} dx - 2 \int \frac{1}{x^2 + 2} dx$$

$$\downarrow 216$$

$$3 \arctan(x) - \sqrt{2} \arctan\left(\frac{x}{\sqrt{2}}\right)$$

input `Int[(4 + x^2)/((1 + x^2)*(2 + x^2)),x]`

output `3*ArcTan[x] - Sqrt[2]*ArcTan[x/Sqrt[2]]`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 397 `Int[((e_) + (f_.)*(x_)^2)/(((a_) + (b_.)*(x_)^2)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Simp[(b*e - a*f)/(b*c - a*d) Int[1/(a + b*x^2), x], x] - Simp[(d*e - c*f)/(b*c - a*d) Int[1/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d, e, f}, x]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

method	result	size
default	$3 \arctan(x) - \arctan\left(\frac{x\sqrt{2}}{2}\right) \sqrt{2}$	18
risch	$3 \arctan(x) - \arctan\left(\frac{x\sqrt{2}}{2}\right) \sqrt{2}$	18

input `int((x^2+4)/(x^2+1)/(x^2+2),x,method=_RETURNVERBOSE)`

output `3*arctan(x)-arctan(1/2*x*2^(1/2))*2^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{4 + x^2}{(1 + x^2)(2 + x^2)} dx = -\sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}x\right) + 3 \arctan(x)$$

input `integrate((x^2+4)/(x^2+1)/(x^2+2),x, algorithm="fricas")`

output `-sqrt(2)*arctan(1/2*sqrt(2)*x) + 3*arctan(x)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{4 + x^2}{(1 + x^2)(2 + x^2)} dx = 3 \operatorname{atan}(x) - \sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}x}{2}\right)$$

input `integrate((x**2+4)/(x**2+1)/(x**2+2),x)`

output `3*atan(x) - sqrt(2)*atan(sqrt(2)*x/2)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{4 + x^2}{(1 + x^2)(2 + x^2)} dx = -\sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}x\right) + 3 \arctan(x)$$

input `integrate((x^2+4)/(x^2+1)/(x^2+2),x, algorithm="maxima")`output `-sqrt(2)*arctan(1/2*sqrt(2)*x) + 3*arctan(x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{4 + x^2}{(1 + x^2)(2 + x^2)} dx = -\sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}x\right) + 3 \arctan(x)$$

input `integrate((x^2+4)/(x^2+1)/(x^2+2),x, algorithm="giac")`output `-sqrt(2)*arctan(1/2*sqrt(2)*x) + 3*arctan(x)`**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{4 + x^2}{(1 + x^2)(2 + x^2)} dx = 3 \operatorname{atan}(x) - \sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}x}{2}\right)$$

input `int((x^2 + 4)/((x^2 + 1)*(x^2 + 2)),x)`output `3*atan(x) - 2^(1/2)*atan((2^(1/2)*x)/2)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int \frac{4 + x^2}{(1 + x^2)(2 + x^2)} dx = -\sqrt{2} \operatorname{atan}\left(\frac{x}{\sqrt{2}}\right) + 3\operatorname{atan}(x)$$

input `int((x^2+4)/(x^2+1)/(x^2+2),x)`

output `- sqrt(2)*atan(x/sqrt(2)) + 3*atan(x)`

$$3.25 \quad \int \frac{5-4x+3x^2+x^4}{(-1+x)^2(1+x^2)} dx$$

Optimal result	181
Mathematica [A] (verified)	181
Rubi [A] (verified)	182
Maple [A] (verified)	183
Fricas [A] (verification not implemented)	183
Sympy [A] (verification not implemented)	184
Maxima [A] (verification not implemented)	184
Giac [B] (verification not implemented)	184
Mupad [B] (verification not implemented)	185
Reduce [B] (verification not implemented)	185

Optimal result

Integrand size = 26, antiderivative size = 37

$$\int \frac{5-4x+3x^2+x^4}{(-1+x)^2(1+x^2)} dx = \frac{5}{2(1-x)} + x + 2 \arctan(x) + \frac{1}{2} \log(1-x) + \frac{3}{4} \log(1+x^2)$$

output `5/(2-2*x)+x+2*arctan(x)+1/2*ln(1-x)+3/4*ln(x^2+1)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.89

$$\int \frac{5-4x+3x^2+x^4}{(-1+x)^2(1+x^2)} dx = \frac{5}{2-2x} + x + 2 \arctan(x) + \frac{1}{2} \log(-1+x) + \frac{3}{4} \log(1+x^2)$$

input `Integrate[(5 - 4*x + 3*x^2 + x^4)/((-1 + x)^2*(1 + x^2)),x]`

output `5/(2 - 2*x) + x + 2*ArcTan[x] + Log[-1 + x]/2 + (3*Log[1 + x^2])/4`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {2160, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4 + 3x^2 - 4x + 5}{(x-1)^2(x^2+1)} dx$$

$$\downarrow \text{2160}$$

$$\int \left(\frac{3x+4}{2(x^2+1)} + \frac{1}{2(x-1)} + \frac{5}{2(x-1)^2} + 1 \right) dx$$

$$\downarrow \text{2009}$$

$$2 \arctan(x) + \frac{3}{4} \log(x^2+1) + x + \frac{5}{2(1-x)} + \frac{1}{2} \log(1-x)$$

input

```
Int[(5 - 4*x + 3*x^2 + x^4)/((-1 + x)^2*(1 + x^2)),x]
```

output

```
5/(2*(1 - x)) + x + 2*ArcTan[x] + Log[1 - x]/2 + (3*Log[1 + x^2])/4
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2160

```
Int[(Pq_)*((d_) + (e_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol]
:> Int[ExpandIntegrand[(d + e*x)^m*Pq*(a + b*x^2)^p, x], x] /; FreeQ[{a, b,
d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.76

method	result
default	$x + \frac{3\ln(x^2+1)}{4} + 2 \arctan(x) - \frac{5}{2(-1+x)} + \frac{\ln(-1+x)}{2}$
risch	$x + \frac{3\ln(x^2+1)}{4} + 2 \arctan(x) - \frac{5}{2(-1+x)} + \frac{\ln(-1+x)}{2}$
parallelrisc	$-\frac{4i \ln(x-i)x - 4i \ln(x+i)x - 4i \ln(x-i) + 4i \ln(x+i) - 2 \ln(-1+x)x - 3 \ln(x-i)x - 3 \ln(x+i)x - 4x^2 + 14 + 2 \ln(-1+x) + 3 \ln(x-i)}{4(-1+x)}$

input `int((x^4+3*x^2-4*x+5)/(-1+x)^2/(x^2+1),x,method=_RETURNVERBOSE)`

output `x+3/4*ln(x^2+1)+2*arctan(x)-5/2/(-1+x)+1/2*ln(-1+x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.19

$$\int \frac{5 - 4x + 3x^2 + x^4}{(-1 + x)^2 (1 + x^2)} dx$$

$$= \frac{4x^2 + 8(x - 1) \arctan(x) + 3(x - 1) \log(x^2 + 1) + 2(x - 1) \log(x - 1) - 4x - 10}{4(x - 1)}$$

input `integrate((x^4+3*x^2-4*x+5)/(-1+x)^2/(x^2+1),x, algorithm="fricas")`

output `1/4*(4*x^2 + 8*(x - 1)*arctan(x) + 3*(x - 1)*log(x^2 + 1) + 2*(x - 1)*log(x - 1) - 4*x - 10)/(x - 1)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.78

$$\int \frac{5 - 4x + 3x^2 + x^4}{(-1 + x)^2 (1 + x^2)} dx = x + \frac{\log(x - 1)}{2} + \frac{3 \log(x^2 + 1)}{4} + 2 \operatorname{atan}(x) - \frac{5}{2x - 2}$$

input `integrate((x**4+3*x**2-4*x+5)/(-1+x)**2/(x**2+1),x)`

output `x + log(x - 1)/2 + 3*log(x**2 + 1)/4 + 2*atan(x) - 5/(2*x - 2)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.73

$$\int \frac{5 - 4x + 3x^2 + x^4}{(-1 + x)^2 (1 + x^2)} dx = x - \frac{5}{2(x - 1)} + 2 \arctan(x) + \frac{3}{4} \log(x^2 + 1) + \frac{1}{2} \log(x - 1)$$

input `integrate((x^4+3*x^2-4*x+5)/(-1+x)^2/(x^2+1),x, algorithm="maxima")`

output `x - 5/2/(x - 1) + 2*arctan(x) + 3/4*log(x^2 + 1) + 1/2*log(x - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 60 vs. $2(29) = 58$.

Time = 0.12 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.62

$$\int \frac{5 - 4x + 3x^2 + x^4}{(-1 + x)^2 (1 + x^2)} dx = \frac{1}{2} \pi - 2 \pi \left[\frac{\pi + 4 \arctan(x)}{4 \pi} + \frac{1}{2} \right] + x - \frac{5}{2(x - 1)} + 2 \arctan(x) + \frac{3}{4} \log \left(\frac{2}{x - 1} + \frac{2}{(x - 1)^2} + 1 \right) + 2 \log(|x - 1|) - 1$$

input `integrate((x^4+3*x^2-4*x+5)/(-1+x)^2/(x^2+1),x, algorithm="giac")`

output `1/2*pi - 2*pi*floor(1/4*(pi + 4*arctan(x))/pi + 1/2) + x - 5/2/(x - 1) + 2*arctan(x) + 3/4*log(2/(x - 1) + 2/(x - 1)^2 + 1) + 2*log(abs(x - 1)) - 1`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.95

$$\int \frac{5 - 4x + 3x^2 + x^4}{(-1 + x)^2 (1 + x^2)} dx = x + \frac{\ln(x - 1)}{2} - \frac{5}{2(x - 1)} + \ln(x - i) \left(\frac{3}{4} - i\right) + \ln(x + i) \left(\frac{3}{4} + i\right)$$

input `int((3*x^2 - 4*x + x^4 + 5)/((x^2 + 1)*(x - 1)^2),x)`

output `x + log(x - 1)/2 + log(x - 1i)*(3/4 - 1i) + log(x + 1i)*(3/4 + 1i) - 5/(2*(x - 1))`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.51

$$\int \frac{5 - 4x + 3x^2 + x^4}{(-1 + x)^2 (1 + x^2)} dx = \frac{8 \operatorname{atan}(x) x - 8 \operatorname{atan}(x) + 3 \log(x^2 + 1) x - 3 \log(x^2 + 1) + 2 \log(x - 1) x - 2 \log(x - 1) + 4x^2 - 14x}{4x - 4}$$

input `int((x^4+3*x^2-4*x+5)/(-1+x)^2/(x^2+1),x)`

output `(8*atan(x)*x - 8*atan(x) + 3*log(x**2 + 1)*x - 3*log(x**2 + 1) + 2*log(x - 1)*x - 2*log(x - 1) + 4*x**2 - 14*x)/(4*(x - 1))`

3.26 $\int \frac{1+x^4}{2+x^2} dx$

Optimal result	186
Mathematica [A] (verified)	186
Rubi [A] (verified)	187
Maple [A] (verified)	188
Fricas [A] (verification not implemented)	188
Sympy [A] (verification not implemented)	188
Maxima [A] (verification not implemented)	189
Giac [A] (verification not implemented)	189
Mupad [B] (verification not implemented)	189
Reduce [B] (verification not implemented)	190

Optimal result

Integrand size = 13, antiderivative size = 26

$$\int \frac{1+x^4}{2+x^2} dx = -2x + \frac{x^3}{3} + \frac{5 \arctan\left(\frac{x}{\sqrt{2}}\right)}{\sqrt{2}}$$

output `-2*x+1/3*x^3+5/2*arctan(1/2*x*2^(1/2))*2^(1/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{1+x^4}{2+x^2} dx = -2x + \frac{x^3}{3} + \frac{5 \arctan\left(\frac{x}{\sqrt{2}}\right)}{\sqrt{2}}$$

input `Integrate[(1 + x^4)/(2 + x^2), x]`

output `-2*x + x^3/3 + (5*ArcTan[x/Sqrt[2]])/Sqrt[2]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {1468, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4 + 1}{x^2 + 2} dx$$

$$\downarrow 1468$$

$$\int \left(x^2 + \frac{5}{x^2 + 2} - 2 \right) dx$$

$$\downarrow 2009$$

$$\frac{5 \arctan\left(\frac{x}{\sqrt{2}}\right)}{\sqrt{2}} + \frac{x^3}{3} - 2x$$

input `Int[(1 + x^4)/(2 + x^2),x]`

output `-2*x + x^3/3 + (5*ArcTan[x/Sqrt[2]])/Sqrt[2]`

Defintions of rubi rules used

rule 1468 `Int[((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (c_.)*(x_)^4)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && IGtQ[p, 0] && IGtQ[q, -2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

method	result	size
default	$-2x + \frac{x^3}{3} + \frac{5 \arctan\left(\frac{x\sqrt{2}}{2}\right)\sqrt{2}}{2}$	22
risch	$-2x + \frac{x^3}{3} + \frac{5 \arctan\left(\frac{x\sqrt{2}}{2}\right)\sqrt{2}}{2}$	22
meijerg	$\frac{\arctan\left(\frac{x\sqrt{2}}{2}\right)\sqrt{2}}{2} + \sqrt{2} \left(-\frac{x\sqrt{2}\left(-\frac{5x^2}{2}+15\right)}{15} + 2 \arctan\left(\frac{x\sqrt{2}}{2}\right) \right)$	41

input `int((x^4+1)/(x^2+2),x,method=_RETURNVERBOSE)`output `-2*x+1/3*x^3+5/2*arctan(1/2*x*2^(1/2))*2^(1/2)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.81

$$\int \frac{1+x^4}{2+x^2} dx = \frac{1}{3}x^3 + \frac{5}{2}\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}x\right) - 2x$$

input `integrate((x^4+1)/(x^2+2),x, algorithm="fricas")`output `1/3*x^3 + 5/2*sqrt(2)*arctan(1/2*sqrt(2)*x) - 2*x`**Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{1+x^4}{2+x^2} dx = \frac{x^3}{3} - 2x + \frac{5\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}x}{2}\right)}{2}$$

input `integrate((x**4+1)/(x**2+2),x)`

output `x**3/3 - 2*x + 5*sqrt(2)*atan(sqrt(2)*x/2)/2`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.81

$$\int \frac{1+x^4}{2+x^2} dx = \frac{1}{3}x^3 + \frac{5}{2}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}x\right) - 2x$$

input `integrate((x^4+1)/(x^2+2),x, algorithm="maxima")`

output `1/3*x^3 + 5/2*sqrt(2)*arctan(1/2*sqrt(2)*x) - 2*x`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.81

$$\int \frac{1+x^4}{2+x^2} dx = \frac{1}{3}x^3 + \frac{5}{2}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}x\right) - 2x$$

input `integrate((x^4+1)/(x^2+2),x, algorithm="giac")`

output `1/3*x^3 + 5/2*sqrt(2)*arctan(1/2*sqrt(2)*x) - 2*x`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.81

$$\int \frac{1+x^4}{2+x^2} dx = \frac{5\sqrt{2}\operatorname{atan}\left(\frac{\sqrt{2}x}{2}\right)}{2} - 2x + \frac{x^3}{3}$$

input `int((x^4 + 1)/(x^2 + 2),x)`

output $(5\sqrt{2}\operatorname{atan}(\sqrt{2}x)/2 - 2x + x^3/3)$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.77

$$\int \frac{1+x^4}{2+x^2} dx = \frac{5\sqrt{2}\operatorname{atan}\left(\frac{x}{\sqrt{2}}\right)}{2} + \frac{x^3}{3} - 2x$$

input $\operatorname{int}((x^4+1)/(x^2+2), x)$

output $(15\sqrt{2}\operatorname{atan}(x/\sqrt{2}) + 2x^3 - 12x)/6$

3.27 $\int \frac{2+2x+x^4}{x^4+x^5} dx$

Optimal result	191
Mathematica [A] (verified)	191
Rubi [A] (verified)	192
Maple [A] (verified)	193
Fricas [A] (verification not implemented)	193
Sympy [A] (verification not implemented)	194
Maxima [A] (verification not implemented)	194
Giac [A] (verification not implemented)	194
Mupad [B] (verification not implemented)	195
Reduce [B] (verification not implemented)	195

Optimal result

Integrand size = 18, antiderivative size = 12

$$\int \frac{2 + 2x + x^4}{x^4 + x^5} dx = -\frac{2}{3x^3} + \log(1 + x)$$

output

```
-2/3/x^3+ln(1+x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{2 + 2x + x^4}{x^4 + x^5} dx = -\frac{2}{3x^3} + \log(1 + x)$$

input

```
Integrate[(2 + 2*x + x^4)/(x^4 + x^5), x]
```

output

```
-2/(3*x^3) + Log[1 + x]
```


Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2026, 2123, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^4 + 2x + 2}{x^5 + x^4} dx \\ & \quad \downarrow \text{2026} \\ & \int \frac{x^4 + 2x + 2}{x^4(x+1)} dx \\ & \quad \downarrow \text{2123} \\ & \int \left(\frac{2}{x^4} + \frac{1}{x+1} \right) dx \\ & \quad \downarrow \text{2009} \\ & \log(x+1) - \frac{2}{3x^3} \end{aligned}$$

input `Int[(2 + 2*x + x^4)/(x^4 + x^5),x]`

output `-2/(3*x^3) + Log[1 + x]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2123

```
Int[(Px_)*((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.), x_Symbol]
:> Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c,
d, m, n}, x] && PolyQ[Px, x] && (IntegersQ[m, n] || IGtQ[m, -2])
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

method	result	size
default	$-\frac{2}{3x^3} + \ln(1+x)$	11
norman	$-\frac{2}{3x^3} + \ln(1+x)$	11
meijerg	$-\frac{2}{3x^3} + \ln(1+x)$	11
risch	$-\frac{2}{3x^3} + \ln(1+x)$	11
parallelrisc	$\frac{3\ln(1+x)x^3-2}{3x^3}$	17

input

```
int((x^4+2*x+2)/(x^5+x^4),x,method=_RETURNVERBOSE)
```

output

```
-2/3/x^3+ln(1+x)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.33

$$\int \frac{2 + 2x + x^4}{x^4 + x^5} dx = \frac{3x^3 \log(x+1) - 2}{3x^3}$$

input

```
integrate((x^4+2*x+2)/(x^5+x^4),x, algorithm="fricas")
```

output

```
1/3*(3*x^3*log(x + 1) - 2)/x^3
```

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{2 + 2x + x^4}{x^4 + x^5} dx = \log(x + 1) - \frac{2}{3x^3}$$

input `integrate((x**4+2*x+2)/(x**5+x**4),x)`output `log(x + 1) - 2/(3*x**3)`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{2 + 2x + x^4}{x^4 + x^5} dx = -\frac{2}{3x^3} + \log(x + 1)$$

input `integrate((x^4+2*x+2)/(x^5+x^4),x, algorithm="maxima")`output `-2/3/x^3 + log(x + 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

$$\int \frac{2 + 2x + x^4}{x^4 + x^5} dx = -\frac{2}{3x^3} + \log(|x + 1|)$$

input `integrate((x^4+2*x+2)/(x^5+x^4),x, algorithm="giac")`output `-2/3/x^3 + log(abs(x + 1))`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{2 + 2x + x^4}{x^4 + x^5} dx = \ln(x + 1) - \frac{2}{3x^3}$$

input `int((2*x + x^4 + 2)/(x^4 + x^5),x)`

output `log(x + 1) - 2/(3*x^3)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.33

$$\int \frac{2 + 2x + x^4}{x^4 + x^5} dx = \frac{3 \log(x + 1) x^3 - 2}{3x^3}$$

input `int((x^4+2*x+2)/(x^5+x^4),x)`

output `(3*log(x + 1)*x**3 - 2)/(3*x**3)`

3.28 $\int \frac{-1-5x+2x^2}{2-x-2x^2+x^3} dx$

Optimal result	196
Mathematica [A] (verified)	196
Rubi [A] (verified)	197
Maple [A] (verified)	198
Fricas [A] (verification not implemented)	198
Sympy [A] (verification not implemented)	198
Maxima [A] (verification not implemented)	199
Giac [A] (verification not implemented)	199
Mupad [B] (verification not implemented)	199
Reduce [B] (verification not implemented)	200

Optimal result

Integrand size = 26, antiderivative size = 21

$$\int \frac{-1-5x+2x^2}{2-x-2x^2+x^3} dx = 2\log(1-x) - \log(2-x) + \log(1+x)$$

output `2*ln(1-x)-ln(2-x)+ln(1+x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{-1-5x+2x^2}{2-x-2x^2+x^3} dx = 2\log(1-x) - \log(2-x) + \log(1+x)$$

input `Integrate[(-1 - 5*x + 2*x^2)/(2 - x - 2*x^2 + x^3),x]`

output `2*Log[1 - x] - Log[2 - x] + Log[1 + x]`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{2x^2 - 5x - 1}{x^3 - 2x^2 - x + 2} dx$$

$$\downarrow \text{2462}$$

$$\int \left(\frac{2}{x-1} + \frac{1}{x+1} + \frac{1}{2-x} \right) dx$$

$$\downarrow \text{2009}$$

$$2 \log(1-x) - \log(2-x) + \log(x+1)$$

input `Int[(-1 - 5*x + 2*x^2)/(2 - x - 2*x^2 + x^3),x]`

output `2*Log[1 - x] - Log[2 - x] + Log[1 + x]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2462 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegr and[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ [Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0] && RationalFunctionQ[u, x]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.86

method	result	size
default	$2 \ln(-1+x) - \ln(-2+x) + \ln(1+x)$	18
norman	$2 \ln(-1+x) - \ln(-2+x) + \ln(1+x)$	18
risch	$2 \ln(-1+x) - \ln(-2+x) + \ln(1+x)$	18
parallelrisch	$2 \ln(-1+x) - \ln(-2+x) + \ln(1+x)$	18

input `int((2*x^2-5*x-1)/(x^3-2*x^2-x+2),x,method=_RETURNVERBOSE)`

output `2*ln(-1+x)-ln(-2+x)+ln(1+x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{-1 - 5x + 2x^2}{2 - x - 2x^2 + x^3} dx = \log(x + 1) + 2 \log(x - 1) - \log(x - 2)$$

input `integrate((2*x^2-5*x-1)/(x^3-2*x^2-x+2),x, algorithm="fricas")`

output `log(x + 1) + 2*log(x - 1) - log(x - 2)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int \frac{-1 - 5x + 2x^2}{2 - x - 2x^2 + x^3} dx = -\log(x - 2) + 2 \log(x - 1) + \log(x + 1)$$

input `integrate((2*x**2-5*x-1)/(x**3-2*x**2-x+2),x)`

output `-log(x - 2) + 2*log(x - 1) + log(x + 1)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{-1 - 5x + 2x^2}{2 - x - 2x^2 + x^3} dx = \log(x + 1) + 2 \log(x - 1) - \log(x - 2)$$

input `integrate((2*x^2-5*x-1)/(x^3-2*x^2-x+2),x, algorithm="maxima")`

output `log(x + 1) + 2*log(x - 1) - log(x - 2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.95

$$\int \frac{-1 - 5x + 2x^2}{2 - x - 2x^2 + x^3} dx = \log(|x + 1|) + 2 \log(|x - 1|) - \log(|x - 2|)$$

input `integrate((2*x^2-5*x-1)/(x^3-2*x^2-x+2),x, algorithm="giac")`

output `log(abs(x + 1)) + 2*log(abs(x - 1)) - log(abs(x - 2))`

Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{-1 - 5x + 2x^2}{2 - x - 2x^2 + x^3} dx = 2 \ln(x - 1) - 2 \operatorname{atanh}\left(\frac{144}{11(22x - 50)} + \frac{13}{11}\right)$$

input `int((5*x - 2*x^2 + 1)/(x + 2*x^2 - x^3 - 2),x)`

output `2*log(x - 1) - 2*atanh(144/(11*(22*x - 50)) + 13/11)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{-1 - 5x + 2x^2}{2 - x - 2x^2 + x^3} dx = -\log(x - 2) + 2\log(x - 1) + \log(x + 1)$$

input `int((2*x^2-5*x-1)/(x^3-2*x^2-x+2),x)`

output `- log(x - 2) + 2*log(x - 1) + log(x + 1)`

3.29 $\int \frac{2+x+x^3}{1+2x^2+x^4} dx$

Optimal result	201
Mathematica [A] (verified)	201
Rubi [A] (verified)	202
Maple [A] (verified)	204
Fricas [A] (verification not implemented)	204
Sympy [A] (verification not implemented)	204
Maxima [A] (verification not implemented)	205
Giac [A] (verification not implemented)	205
Mupad [B] (verification not implemented)	205
Reduce [B] (verification not implemented)	206

Optimal result

Integrand size = 19, antiderivative size = 22

$$\int \frac{2+x+x^3}{1+2x^2+x^4} dx = \frac{x}{1+x^2} + \arctan(x) + \frac{1}{2} \log(1+x^2)$$

output `x/(x^2+1)+arctan(x)+1/2*ln(x^2+1)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{2+x+x^3}{1+2x^2+x^4} dx = \frac{x}{1+x^2} + \arctan(x) + \frac{1}{2} \log(1+x^2)$$

input `Integrate[(2 + x + x^3)/(1 + 2*x^2 + x^4), x]`

output `x/(1 + x^2) + ArcTan[x] + Log[1 + x^2]/2`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {1380, 2345, 27, 452, 216, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3 + x + 2}{x^4 + 2x^2 + 1} dx \\
 & \quad \downarrow \text{1380} \\
 & \int \frac{x^3 + x + 2}{(x^2 + 1)^2} dx \\
 & \quad \downarrow \text{2345} \\
 & \frac{x}{x^2 + 1} - \frac{1}{2} \int -\frac{2(x + 1)}{x^2 + 1} dx \\
 & \quad \downarrow \text{27} \\
 & \int \frac{x + 1}{x^2 + 1} dx + \frac{x}{x^2 + 1} \\
 & \quad \downarrow \text{452} \\
 & \int \frac{1}{x^2 + 1} dx + \int \frac{x}{x^2 + 1} dx + \frac{x}{x^2 + 1} \\
 & \quad \downarrow \text{216} \\
 & \int \frac{x}{x^2 + 1} dx + \arctan(x) + \frac{x}{x^2 + 1} \\
 & \quad \downarrow \text{240} \\
 & \arctan(x) + \frac{x}{x^2 + 1} + \frac{1}{2} \log(x^2 + 1)
 \end{aligned}$$

input

 $\text{Int}[(2 + x + x^3)/(1 + 2*x^2 + x^4), x]$

output

 $x/(1 + x^2) + \text{ArcTan}[x] + \text{Log}[1 + x^2]/2$

Definitions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 240 `Int[(x_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`
- rule 452 `Int[((c_) + (d_.)*(x_))/((a_) + (b_.)*(x_)^2), x_Symbol] := Simp[c Int[1/(a + b*x^2), x], x] + Simp[d Int[x/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c^2 + a*d^2, 0]`
- rule 1380 `Int[(u_)*((a_) + (c_.)*(x_)^(n2_.) + (b_.)*(x_)^(n_.))^p_., x_Symbol] := Simp[1/c^p Int[u*(b/2 + c*x^n)^(2*p), x], x] /; FreeQ[{a, b, c, n, p}, x] && EqQ[n2, 2*n] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p]`
- rule 2345 `Int[(Pq_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{Q = PolynomialQuotient[Pq, a + b*x^2, x], f = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 0], g = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 1]}, Simp[(a*g - b*f*x)*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] + Simp[1/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1)*ExpandToSum[2*a*(p + 1)*Q + f*(2*p + 3), x], x], x] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && LtQ[p, -1]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.95

method	result	size
default	$\frac{x}{x^2+1} + \arctan(x) + \frac{\ln(x^2+1)}{2}$	21
risch	$\frac{x}{x^2+1} + \arctan(x) + \frac{\ln(x^2+1)}{2}$	21
parallelrisch	$-\frac{i \ln(x-i)x^2 - i \ln(x+i)x^2 - \ln(x-i)x^2 - \ln(x+i)x^2 + i \ln(x-i) - i \ln(x+i) - \ln(x-i) - \ln(x+i) - 2x}{2(x^2+1)}$	86

input `int((x^3+x+2)/(x^4+2*x^2+1),x,method=_RETURNVERBOSE)`output `x/(x^2+1)+arctan(x)+1/2*ln(x^2+1)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.55

$$\int \frac{2+x+x^3}{1+2x^2+x^4} dx = \frac{2(x^2+1)\arctan(x) + (x^2+1)\log(x^2+1) + 2x}{2(x^2+1)}$$

input `integrate((x^3+x+2)/(x^4+2*x^2+1),x, algorithm="fricas")`output `1/2*(2*(x^2+1)*arctan(x) + (x^2+1)*log(x^2+1) + 2*x)/(x^2+1)`**Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

$$\int \frac{2+x+x^3}{1+2x^2+x^4} dx = \frac{x}{x^2+1} + \frac{\log(x^2+1)}{2} + \operatorname{atan}(x)$$

input `integrate((x**3+x+2)/(x**4+2*x**2+1),x)`

output `x/(x**2 + 1) + log(x**2 + 1)/2 + atan(x)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{2 + x + x^3}{1 + 2x^2 + x^4} dx = \frac{x}{x^2 + 1} + \arctan(x) + \frac{1}{2} \log(x^2 + 1)$$

input `integrate((x^3+x+2)/(x^4+2*x^2+1),x, algorithm="maxima")`

output `x/(x^2 + 1) + arctan(x) + 1/2*log(x^2 + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{2 + x + x^3}{1 + 2x^2 + x^4} dx = \frac{x}{x^2 + 1} + \arctan(x) + \frac{1}{2} \log(x^2 + 1)$$

input `integrate((x^3+x+2)/(x^4+2*x^2+1),x, algorithm="giac")`

output `x/(x^2 + 1) + arctan(x) + 1/2*log(x^2 + 1)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{2 + x + x^3}{1 + 2x^2 + x^4} dx = \frac{\ln(x^2 + 1)}{2} + \operatorname{atan}(x) + \frac{x}{x^2 + 1}$$

input `int((x + x^3 + 2)/(2*x^2 + x^4 + 1),x)`

output `log(x^2 + 1)/2 + atan(x) + x/(x^2 + 1)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.86

$$\int \frac{2 + x + x^3}{1 + 2x^2 + x^4} dx = \frac{2\operatorname{atan}(x)x^2 + 2\operatorname{atan}(x) + \log(x^2 + 1)x^2 + \log(x^2 + 1) + 2x}{2x^2 + 2}$$

input `int((x^3+x+2)/(x^4+2*x^2+1),x)`

output `(2*atan(x)*x**2 + 2*atan(x) + log(x**2 + 1)*x**2 + log(x**2 + 1) + 2*x)/(2*(x**2 + 1))`

3.30 $\int \frac{1+2x+x^2+x^3}{1+2x^2+x^4} dx$

Optimal result	207
Mathematica [A] (verified)	207
Rubi [A] (verified)	208
Maple [A] (verified)	210
Fricas [A] (verification not implemented)	210
Sympy [A] (verification not implemented)	210
Maxima [A] (verification not implemented)	211
Giac [A] (verification not implemented)	211
Mupad [B] (verification not implemented)	211
Reduce [B] (verification not implemented)	212

Optimal result

Integrand size = 24, antiderivative size = 24

$$\int \frac{1+2x+x^2+x^3}{1+2x^2+x^4} dx = -\frac{1}{2(1+x^2)} + \arctan(x) + \frac{1}{2} \log(1+x^2)$$

output `-1/(2*x^2+2)+arctan(x)+1/2*ln(x^2+1)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{1+2x+x^2+x^3}{1+2x^2+x^4} dx = -\frac{1}{2(1+x^2)} + \arctan(x) + \frac{1}{2} \log(1+x^2)$$

input `Integrate[(1 + 2*x + x^2 + x^3)/(1 + 2*x^2 + x^4),x]`

output `-1/2*1/(1 + x^2) + ArcTan[x] + Log[1 + x^2]/2`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {1380, 2345, 27, 452, 216, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3 + x^2 + 2x + 1}{x^4 + 2x^2 + 1} dx \\
 & \quad \downarrow \text{1380} \\
 & \int \frac{x^3 + x^2 + 2x + 1}{(x^2 + 1)^2} dx \\
 & \quad \downarrow \text{2345} \\
 & -\frac{1}{2} \int -\frac{2(x+1)}{x^2+1} dx - \frac{1}{2(x^2+1)} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{x+1}{x^2+1} dx - \frac{1}{2(x^2+1)} \\
 & \quad \downarrow \text{452} \\
 & \int \frac{1}{x^2+1} dx + \int \frac{x}{x^2+1} dx - \frac{1}{2(x^2+1)} \\
 & \quad \downarrow \text{216} \\
 & \int \frac{x}{x^2+1} dx + \arctan(x) - \frac{1}{2(x^2+1)} \\
 & \quad \downarrow \text{240} \\
 & \arctan(x) - \frac{1}{2(x^2+1)} + \frac{1}{2} \log(x^2+1)
 \end{aligned}$$

input

 $\text{Int}[(1 + 2*x + x^2 + x^3)/(1 + 2*x^2 + x^4), x]$

output

 $-1/2*1/(1 + x^2) + \text{ArcTan}[x] + \text{Log}[1 + x^2]/2$

Definitions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 240 `Int[(x_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`
- rule 452 `Int[((c_) + (d_.)*(x_))/((a_) + (b_.)*(x_)^2), x_Symbol] := Simp[c Int[1/(a + b*x^2), x], x] + Simp[d Int[x/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c^2 + a*d^2, 0]`
- rule 1380 `Int[(u_)*((a_) + (c_.)*(x_)^(n2_.) + (b_.)*(x_)^(n_.))^p_., x_Symbol] := Simp[1/c^p Int[u*(b/2 + c*x^n)^(2*p), x], x] /; FreeQ[{a, b, c, n, p}, x] && EqQ[n2, 2*n] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p]`
- rule 2345 `Int[(Pq_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{Q = PolynomialQuotient[Pq, a + b*x^2, x], f = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 0], g = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 1]}, Simp[(a*g - b*f*x)*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] + Simp[1/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1)*ExpandToSum[2*a*(p + 1)*Q + f*(2*p + 3), x], x], x] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && LtQ[p, -1]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.88

method	result	size
default	$-\frac{1}{2(x^2+1)} + \frac{\ln(x^2+1)}{2} + \arctan(x)$	21
risch	$-\frac{1}{2(x^2+1)} + \frac{\ln(x^2+1)}{2} + \arctan(x)$	21
parallelrisch	$-\frac{i \ln(x-i)x^2 - i \ln(x+i)x^2 - \ln(x-i)x^2 - \ln(x+i)x^2 + 1 + i \ln(x-i) - i \ln(x+i) - \ln(x-i) - \ln(x+i)}{2(x^2+1)}$	84

input `int((x^3+x^2+2*x+1)/(x^4+2*x^2+1),x,method=_RETURNVERBOSE)`output `-1/2/(x^2+1)+1/2*ln(x^2+1)+arctan(x)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.33

$$\int \frac{1 + 2x + x^2 + x^3}{1 + 2x^2 + x^4} dx = \frac{2(x^2 + 1) \arctan(x) + (x^2 + 1) \log(x^2 + 1) - 1}{2(x^2 + 1)}$$

input `integrate((x^3+x^2+2*x+1)/(x^4+2*x^2+1),x, algorithm="fricas")`output `1/2*(2*(x^2 + 1)*arctan(x) + (x^2 + 1)*log(x^2 + 1) - 1)/(x^2 + 1)`**Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.79

$$\int \frac{1 + 2x + x^2 + x^3}{1 + 2x^2 + x^4} dx = \frac{\log(x^2 + 1)}{2} + \operatorname{atan}(x) - \frac{1}{2x^2 + 2}$$

input `integrate((x**3+x**2+2*x+1)/(x**4+2*x**2+1),x)`

output `log(x**2 + 1)/2 + atan(x) - 1/(2*x**2 + 2)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{1 + 2x + x^2 + x^3}{1 + 2x^2 + x^4} dx = -\frac{1}{2(x^2 + 1)} + \arctan(x) + \frac{1}{2} \log(x^2 + 1)$$

input `integrate((x^3+x^2+2*x+1)/(x^4+2*x^2+1),x, algorithm="maxima")`

output `-1/2/(x^2 + 1) + arctan(x) + 1/2*log(x^2 + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{1 + 2x + x^2 + x^3}{1 + 2x^2 + x^4} dx = -\frac{1}{2(x^2 + 1)} + \arctan(x) + \frac{1}{2} \log(x^2 + 1)$$

input `integrate((x^3+x^2+2*x+1)/(x^4+2*x^2+1),x, algorithm="giac")`

output `-1/2/(x^2 + 1) + arctan(x) + 1/2*log(x^2 + 1)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1 + 2x + x^2 + x^3}{1 + 2x^2 + x^4} dx = \frac{\ln(x^2 + 1)}{2} + \operatorname{atan}(x) - \frac{1}{2(x^2 + 1)}$$

input `int((2*x + x^2 + x^3 + 1)/(2*x^2 + x^4 + 1),x)`

output `log(x^2 + 1)/2 + atan(x) - 1/(2*(x^2 + 1))`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.71

$$\int \frac{1 + 2x + x^2 + x^3}{1 + 2x^2 + x^4} dx = \frac{2\operatorname{atan}(x)x^2 + 2\operatorname{atan}(x) + \log(x^2 + 1)x^2 + \log(x^2 + 1) + x^2}{2x^2 + 2}$$

input `int((x^3+x^2+2*x+1)/(x^4+2*x^2+1),x)`

output `(2*atan(x)*x**2 + 2*atan(x) + log(x**2 + 1)*x**2 + log(x**2 + 1) + x**2)/(2*(x**2 + 1))`

3.31 $\int \frac{3+4x}{(1+x^2)(2+x^2)} dx$

Optimal result	213
Mathematica [A] (verified)	213
Rubi [A] (verified)	214
Maple [A] (verified)	216
Fricas [A] (verification not implemented)	216
Sympy [A] (verification not implemented)	216
Maxima [A] (verification not implemented)	217
Giac [A] (verification not implemented)	217
Mupad [B] (verification not implemented)	218
Reduce [B] (verification not implemented)	218

Optimal result

Integrand size = 20, antiderivative size = 36

$$\int \frac{3+4x}{(1+x^2)(2+x^2)} dx = 3 \arctan(x) - \frac{3 \arctan\left(\frac{x}{\sqrt{2}}\right)}{\sqrt{2}} + 2 \log(1+x^2) - 2 \log(2+x^2)$$

output `3*arctan(x)-3/2*arctan(1/2*x*2^(1/2))*2^(1/2)+2*ln(x^2+1)-2*ln(x^2+2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00

$$\int \frac{3+4x}{(1+x^2)(2+x^2)} dx = 3 \arctan(x) - \frac{3 \arctan\left(\frac{x}{\sqrt{2}}\right)}{\sqrt{2}} + 2 \log(1+x^2) - 2 \log(2+x^2)$$

input `Integrate[(3 + 4*x)/((1 + x^2)*(2 + x^2)),x]`

output `3*ArcTan[x] - (3*ArcTan[x/Sqrt[2]])/Sqrt[2] + 2*Log[1 + x^2] - 2*Log[2 + x^2]`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.06, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {1343, 303, 216, 353, 47, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{4x + 3}{(x^2 + 1)(x^2 + 2)} dx \\
 & \quad \downarrow \text{1343} \\
 & 3 \int \frac{1}{(x^2 + 1)(x^2 + 2)} dx + 4 \int \frac{x}{(x^2 + 1)(x^2 + 2)} dx \\
 & \quad \downarrow \text{303} \\
 & 3 \left(\int \frac{1}{x^2 + 1} dx - \int \frac{1}{x^2 + 2} dx \right) + 4 \int \frac{x}{(x^2 + 1)(x^2 + 2)} dx \\
 & \quad \downarrow \text{216} \\
 & 4 \int \frac{x}{(x^2 + 1)(x^2 + 2)} dx + 3 \left(\arctan(x) - \frac{\arctan\left(\frac{x}{\sqrt{2}}\right)}{\sqrt{2}} \right) \\
 & \quad \downarrow \text{353} \\
 & 2 \int \frac{1}{(x^2 + 1)(x^2 + 2)} dx^2 + 3 \left(\arctan(x) - \frac{\arctan\left(\frac{x}{\sqrt{2}}\right)}{\sqrt{2}} \right) \\
 & \quad \downarrow \text{47} \\
 & 2 \left(\int \frac{1}{x^2 + 1} dx^2 - \int \frac{1}{x^2 + 2} dx^2 \right) + 3 \left(\arctan(x) - \frac{\arctan\left(\frac{x}{\sqrt{2}}\right)}{\sqrt{2}} \right) \\
 & \quad \downarrow \text{16} \\
 & 3 \left(\arctan(x) - \frac{\arctan\left(\frac{x}{\sqrt{2}}\right)}{\sqrt{2}} \right) + 2(\log(x^2 + 1) - \log(x^2 + 2))
 \end{aligned}$$

input

```
Int[(3 + 4*x)/((1 + x^2)*(2 + x^2)), x]
```

output $3*(\text{ArcTan}[x] - \text{ArcTan}[x/\text{Sqrt}[2]]/\text{Sqrt}[2]) + 2*(\text{Log}[1 + x^2] - \text{Log}[2 + x^2])$

Defintions of rubi rules used

rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 47 $\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_))), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x), x], x] /; \text{FreeQ}[\{a, b, c, d\}, x]$

rule 216 $\text{Int}[(a_)+(b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b] \&\& (\text{GtQ}[a, 0] \parallel \text{GtQ}[b, 0])$

rule 303 $\text{Int}[1/(((a_)+(b_)*(x_)^2))*((c_)+(d_)*(x_)^2)), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x^2), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0]$

rule 353 $\text{Int}[(x_)*((a_)+(b_)*(x_)^2)^{(p_)}*((c_)+(d_)*(x_)^2)^{(q_)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; \text{FreeQ}[\{a, b, c, d, p, q\}, x] \&\& \text{NeQ}[b*c - a*d, 0]$

rule 1343 $\text{Int}[(g_)+(h_)*(x_))*((a_)+(c_)*(x_)^2)^{(p_)}*((d_)+(f_)*(x_)^2)^{(q_)}, x_Symbol] \rightarrow \text{Simp}[g \text{ Int}[(a + c*x^2)^p*(d + f*x^2)^q, x], x] + \text{Simp}[h \text{ Int}[x*(a + c*x^2)^p*(d + f*x^2)^q, x], x] /; \text{FreeQ}[\{a, c, d, f, g, h, p, q\}, x]$

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.94

method	result	size
default	$3 \arctan(x) - \frac{3 \arctan\left(\frac{x\sqrt{2}}{2}\right)\sqrt{2}}{2} + 2 \ln(x^2 + 1) - 2 \ln(x^2 + 2)$	34
risch	$3 \arctan(x) - \frac{3 \arctan\left(\frac{x\sqrt{2}}{2}\right)\sqrt{2}}{2} + 2 \ln(x^2 + 1) - 2 \ln(x^2 + 2)$	34

input `int((4*x+3)/(x^2+1)/(x^2+2),x,method=_RETURNVERBOSE)`

output `3*arctan(x)-3/2*arctan(1/2*x*2^(1/2))*2^(1/2)+2*ln(x^2+1)-2*ln(x^2+2)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.92

$$\int \frac{3 + 4x}{(1 + x^2)(2 + x^2)} dx = -\frac{3}{2} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}x\right) + 3 \arctan(x) - 2 \log(x^2 + 2) + 2 \log(x^2 + 1)$$

input `integrate((3+4*x)/(x^2+1)/(x^2+2),x, algorithm="fricas")`

output `-3/2*sqrt(2)*arctan(1/2*sqrt(2)*x) + 3*arctan(x) - 2*log(x^2 + 2) + 2*log(x^2 + 1)`

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.08

$$\int \frac{3 + 4x}{(1 + x^2)(2 + x^2)} dx = 2 \log(x^2 + 1) - 2 \log(x^2 + 2) + 3 \operatorname{atan}(x) - \frac{3\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}x}{2}\right)}{2}$$

input `integrate((3+4*x)/(x**2+1)/(x**2+2),x)`

output $2*\log(x**2 + 1) - 2*\log(x**2 + 2) + 3*\operatorname{atan}(x) - 3*\sqrt{2}*\operatorname{atan}(\sqrt{2}*x/2)/2$

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.92

$$\int \frac{3 + 4x}{(1 + x^2)(2 + x^2)} dx = -\frac{3}{2} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}x\right) + 3 \arctan(x) - 2 \log(x^2 + 2) + 2 \log(x^2 + 1)$$

input `integrate((3+4*x)/(x^2+1)/(x^2+2),x, algorithm="maxima")`

output $-3/2*\sqrt{2}*\arctan(1/2*\sqrt{2}*x) + 3*\arctan(x) - 2*\log(x^2 + 2) + 2*\log(x^2 + 1)$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.92

$$\int \frac{3 + 4x}{(1 + x^2)(2 + x^2)} dx = -\frac{3}{2} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}x\right) + 3 \arctan(x) - 2 \log(x^2 + 2) + 2 \log(x^2 + 1)$$

input `integrate((3+4*x)/(x^2+1)/(x^2+2),x, algorithm="giac")`

output $-3/2*\sqrt{2}*\arctan(1/2*\sqrt{2}*x) + 3*\arctan(x) - 2*\log(x^2 + 2) + 2*\log(x^2 + 1)$

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.56

$$\int \frac{3 + 4x}{(1 + x^2)(2 + x^2)} dx = \ln(x - i) \left(2 - \frac{3}{2}i\right) + \ln(x + i) \left(2 + \frac{3}{2}i\right) \\ + \ln(x - \sqrt{2}i) \left(-2 + \frac{\sqrt{2}3i}{4}\right) - \ln(x + \sqrt{2}i) \left(2 + \frac{\sqrt{2}3i}{4}\right)$$

input `int((4*x + 3)/((x^2 + 1)*(x^2 + 2)),x)`output `log(x - 1i)*(2 - 3i/2) + log(x + 1i)*(2 + 3i/2) + log(x - 2^(1/2)*1i)*((2^(1/2)*3i)/4 - 2) - log(x + 2^(1/2)*1i)*((2^(1/2)*3i)/4 + 2)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.89

$$\int \frac{3 + 4x}{(1 + x^2)(2 + x^2)} dx = -\frac{3\sqrt{2} \operatorname{atan}\left(\frac{x}{\sqrt{2}}\right)}{2} + 3\operatorname{atan}(x) - 2\log(x^2 + 2) + 2\log(x^2 + 1)$$

input `int((3+4*x)/(x^2+1)/(x^2+2),x)`output `(- 3*sqrt(2)*atan(x/sqrt(2)) + 6*atan(x) - 4*log(x**2 + 2) + 4*log(x**2 + 1))/2`

3.32 $\int \frac{2+x}{(1+x^2)(4+x^2)} dx$

Optimal result	219
Mathematica [A] (verified)	219
Rubi [A] (verified)	220
Maple [A] (verified)	222
Fricas [A] (verification not implemented)	222
Sympy [A] (verification not implemented)	222
Maxima [A] (verification not implemented)	223
Giac [A] (verification not implemented)	223
Mupad [B] (verification not implemented)	224
Reduce [B] (verification not implemented)	224

Optimal result

Integrand size = 18, antiderivative size = 37

$$\int \frac{2+x}{(1+x^2)(4+x^2)} dx = -\frac{1}{3} \arctan\left(\frac{x}{2}\right) + \frac{2 \arctan(x)}{3} + \frac{1}{6} \log(1+x^2) - \frac{1}{6} \log(4+x^2)$$

output `-1/3*arctan(1/2*x)+2/3*arctan(x)+1/6*ln(x^2+1)-1/6*ln(x^2+4)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00

$$\int \frac{2+x}{(1+x^2)(4+x^2)} dx = -\frac{1}{3} \arctan\left(\frac{x}{2}\right) + \frac{2 \arctan(x)}{3} + \frac{1}{6} \log(1+x^2) - \frac{1}{6} \log(4+x^2)$$

input `Integrate[(2 + x)/((1 + x^2)*(4 + x^2)),x]`

output `-1/3*ArcTan[x/2] + (2*ArcTan[x])/3 + Log[1 + x^2]/6 - Log[4 + x^2]/6`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.22, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {1343, 303, 216, 353, 47, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x+2}{(x^2+1)(x^2+4)} dx \\
 & \quad \downarrow \text{1343} \\
 & 2 \int \frac{1}{(x^2+1)(x^2+4)} dx + \int \frac{x}{(x^2+1)(x^2+4)} dx \\
 & \quad \downarrow \text{303} \\
 & 2 \left(\frac{1}{3} \int \frac{1}{x^2+1} dx - \frac{1}{3} \int \frac{1}{x^2+4} dx \right) + \int \frac{x}{(x^2+1)(x^2+4)} dx \\
 & \quad \downarrow \text{216} \\
 & \int \frac{x}{(x^2+1)(x^2+4)} dx + 2 \left(\frac{\arctan(x)}{3} - \frac{1}{6} \arctan\left(\frac{x}{2}\right) \right) \\
 & \quad \downarrow \text{353} \\
 & \frac{1}{2} \int \frac{1}{(x^2+1)(x^2+4)} dx^2 + 2 \left(\frac{\arctan(x)}{3} - \frac{1}{6} \arctan\left(\frac{x}{2}\right) \right) \\
 & \quad \downarrow \text{47} \\
 & \frac{1}{2} \left(\frac{1}{3} \int \frac{1}{x^2+1} dx^2 - \frac{1}{3} \int \frac{1}{x^2+4} dx^2 \right) + 2 \left(\frac{\arctan(x)}{3} - \frac{1}{6} \arctan\left(\frac{x}{2}\right) \right) \\
 & \quad \downarrow \text{16} \\
 & 2 \left(\frac{\arctan(x)}{3} - \frac{1}{6} \arctan\left(\frac{x}{2}\right) \right) + \frac{1}{2} \left(\frac{1}{3} \log(x^2+1) - \frac{1}{3} \log(x^2+4) \right)
 \end{aligned}$$

input `Int[(2 + x)/((1 + x^2)*(4 + x^2)), x]`

output `2*(-1/6*ArcTan[x/2] + ArcTan[x]/3) + (Log[1 + x^2]/3 - Log[4 + x^2]/3)/2`

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}\{a, b, c\}, x]$
- rule 47 $\text{Int}[1/(((a_)+(b_)(x_))*((c_)+(d_)(x_))), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x), x], x] /; \text{FreeQ}\{a, b, c, d\}, x]$
- rule 216 $\text{Int}(((a_) + (b_)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*ArcTan[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x] \&\& \text{PosQ}[a/b] \&\& (\text{GtQ}[a, 0] || \text{GtQ}[b, 0])]$
- rule 303 $\text{Int}[1/(((a_) + (b_)(x_)^2)*((c_) + (d_)(x_)^2)), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x^2), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x^2), x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0]$
- rule 353 $\text{Int}[(x_)*((a_) + (b_)(x_)^2)^{(p_)}*((c_) + (d_)(x_)^2)^{(q_)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; \text{FreeQ}\{a, b, c, d, p, q\}, x] \&\& \text{NeQ}[b*c - a*d, 0]$
- rule 1343 $\text{Int}(((g_) + (h_)(x_))*((a_) + (c_)(x_)^2)^{(p_)}*((d_) + (f_)(x_)^2)^{(q_)}, x_Symbol] \rightarrow \text{Simp}[g \text{ Int}[(a + c*x^2)^p*(d + f*x^2)^q, x], x] + \text{Simp}[h \text{ Int}[x*(a + c*x^2)^p*(d + f*x^2)^q, x], x] /; \text{FreeQ}\{a, c, d, f, g, h, p, q\}, x]$

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.76

method	result	size
default	$-\frac{\arctan(\frac{x}{2})}{3} + \frac{2\arctan(x)}{3} + \frac{\ln(x^2+1)}{6} - \frac{\ln(x^2+4)}{6}$	28
risch	$-\frac{\arctan(\frac{x}{2})}{3} + \frac{2\arctan(x)}{3} + \frac{\ln(x^2+1)}{6} - \frac{\ln(x^2+4)}{6}$	28
parallelrisc	$\frac{\ln(x-i)}{6} - \frac{i\ln(x-i)}{3} - \frac{\ln(x-2i)}{6} + \frac{i\ln(x-2i)}{6} + \frac{\ln(x+i)}{6} + \frac{i\ln(x+i)}{3} - \frac{\ln(x+2i)}{6} - \frac{i\ln(x+2i)}{6}$	62

input `int((2+x)/(x^2+1)/(x^2+4),x,method=_RETURNVERBOSE)`output `-1/3*arctan(1/2*x)+2/3*arctan(x)+1/6*ln(x^2+1)-1/6*ln(x^2+4)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.73

$$\int \frac{2+x}{(1+x^2)(4+x^2)} dx = -\frac{1}{3} \arctan\left(\frac{1}{2}x\right) + \frac{2}{3} \arctan(x) - \frac{1}{6} \log(x^2+4) + \frac{1}{6} \log(x^2+1)$$

input `integrate((2+x)/(x^2+1)/(x^2+4),x, algorithm="fricas")`output `-1/3*arctan(1/2*x) + 2/3*arctan(x) - 1/6*log(x^2 + 4) + 1/6*log(x^2 + 1)`**Sympy [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.78

$$\int \frac{2+x}{(1+x^2)(4+x^2)} dx = \frac{\log(x^2+1)}{6} - \frac{\log(x^2+4)}{6} - \frac{\operatorname{atan}\left(\frac{x}{2}\right)}{3} + \frac{2\operatorname{atan}(x)}{3}$$

input `integrate((2+x)/(x**2+1)/(x**2+4),x)`

output $\log(x^2 + 1)/6 - \log(x^2 + 4)/6 - \operatorname{atan}(x/2)/3 + 2*\operatorname{atan}(x)/3$

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.73

$$\int \frac{2+x}{(1+x^2)(4+x^2)} dx = -\frac{1}{3} \arctan\left(\frac{1}{2}x\right) + \frac{2}{3} \arctan(x) - \frac{1}{6} \log(x^2 + 4) + \frac{1}{6} \log(x^2 + 1)$$

input `integrate((2+x)/(x^2+1)/(x^2+4),x, algorithm="maxima")`

output $-1/3*\arctan(1/2*x) + 2/3*\arctan(x) - 1/6*\log(x^2 + 4) + 1/6*\log(x^2 + 1)$

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.73

$$\int \frac{2+x}{(1+x^2)(4+x^2)} dx = -\frac{1}{3} \arctan\left(\frac{1}{2}x\right) + \frac{2}{3} \arctan(x) - \frac{1}{6} \log(x^2 + 4) + \frac{1}{6} \log(x^2 + 1)$$

input `integrate((2+x)/(x^2+1)/(x^2+4),x, algorithm="giac")`

output $-1/3*\arctan(1/2*x) + 2/3*\arctan(x) - 1/6*\log(x^2 + 4) + 1/6*\log(x^2 + 1)$

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00

$$\int \frac{2+x}{(1+x^2)(4+x^2)} dx = \ln(x-i) \left(\frac{1}{6} - \frac{1}{3}i \right) + \ln(x+1i) \left(\frac{1}{6} + \frac{1}{3}i \right) \\ + \ln(x-2i) \left(-\frac{1}{6} + \frac{1}{6}i \right) + \ln(x+2i) \left(-\frac{1}{6} - \frac{1}{6}i \right)$$

input `int((x + 2)/((x^2 + 1)*(x^2 + 4)),x)`output `log(x - 1i)*(1/6 - 1i/3) + log(x + 1i)*(1/6 + 1i/3) - log(x - 2i)*(1/6 - 1i/6) - log(x + 2i)*(1/6 + 1i/6)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.73

$$\int \frac{2+x}{(1+x^2)(4+x^2)} dx = -\frac{\operatorname{atan}\left(\frac{x}{2}\right)}{3} + \frac{2\operatorname{atan}(x)}{3} - \frac{\log(x^2+4)}{6} + \frac{\log(x^2+1)}{6}$$

input `int((2+x)/(x^2+1)/(x^2+4),x)`output `(- 2*atan(x/2) + 4*atan(x) - log(x**2 + 4) + log(x**2 + 1))/6`

3.33 $\int \frac{2-x+x^3}{-7-6x+x^2} dx$

Optimal result	225
Mathematica [A] (verified)	225
Rubi [A] (verified)	226
Maple [A] (verified)	227
Fricas [A] (verification not implemented)	227
Sympy [A] (verification not implemented)	227
Maxima [A] (verification not implemented)	228
Giac [A] (verification not implemented)	228
Mupad [B] (verification not implemented)	228
Reduce [B] (verification not implemented)	229

Optimal result

Integrand size = 19, antiderivative size = 29

$$\int \frac{2-x+x^3}{-7-6x+x^2} dx = 6x + \frac{x^2}{2} + \frac{169}{4} \log(7-x) - \frac{1}{4} \log(1+x)$$

output `6*x+1/2*x^2+169/4*ln(7-x)-1/4*ln(1+x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{2-x+x^3}{-7-6x+x^2} dx = 6x + \frac{x^2}{2} + \frac{169}{4} \log(7-x) - \frac{1}{4} \log(1+x)$$

input `Integrate[(2 - x + x^3)/(-7 - 6*x + x^2),x]`

output `6*x + x^2/2 + (169*Log[7 - x])/4 - Log[1 + x]/4`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2188, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3 - x + 2}{x^2 - 6x - 7} dx$$

↓ 2188

$$\int \left(\frac{2(21x + 22)}{x^2 - 6x - 7} + x + 6 \right) dx$$

↓ 2009

$$\frac{x^2}{2} + 6x + \frac{169}{4} \log(7 - x) - \frac{1}{4} \log(x + 1)$$

input

```
Int[(2 - x + x^3)/(-7 - 6*x + x^2),x]
```

output

```
6*x + x^2/2 + (169*Log[7 - x])/4 - Log[1 + x]/4
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2188

```
Int[(Pq_)*((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^p, x_Symbol] := Int[Expand
Integrand[Pq*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c}, x] && PolyQ[Pq,
x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.76

method	result	size
default	$\frac{x^2}{2} + 6x + \frac{169 \ln(x-7)}{4} - \frac{\ln(1+x)}{4}$	22
norman	$\frac{x^2}{2} + 6x + \frac{169 \ln(x-7)}{4} - \frac{\ln(1+x)}{4}$	22
risch	$\frac{x^2}{2} + 6x + \frac{169 \ln(x-7)}{4} - \frac{\ln(1+x)}{4}$	22
parallelrisch	$\frac{x^2}{2} + 6x + \frac{169 \ln(x-7)}{4} - \frac{\ln(1+x)}{4}$	22

input `int((x^3-x+2)/(x^2-6*x-7),x,method=_RETURNVERBOSE)`output `1/2*x^2+6*x+169/4*ln(x-7)-1/4*ln(1+x)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

$$\int \frac{2-x+x^3}{-7-6x+x^2} dx = \frac{1}{2}x^2 + 6x - \frac{1}{4} \log(x+1) + \frac{169}{4} \log(x-7)$$

input `integrate((x^3-x+2)/(x^2-6*x-7),x, algorithm="fricas")`output `1/2*x^2 + 6*x - 1/4*log(x + 1) + 169/4*log(x - 7)`**Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.76

$$\int \frac{2-x+x^3}{-7-6x+x^2} dx = \frac{x^2}{2} + 6x + \frac{169 \log(x-7)}{4} - \frac{\log(x+1)}{4}$$

input `integrate((x**3-x+2)/(x**2-6*x-7),x)`

output `x**2/2 + 6*x + 169*log(x - 7)/4 - log(x + 1)/4`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

$$\int \frac{2 - x + x^3}{-7 - 6x + x^2} dx = \frac{1}{2} x^2 + 6x - \frac{1}{4} \log(x + 1) + \frac{169}{4} \log(x - 7)$$

input `integrate((x^3-x+2)/(x^2-6*x-7),x, algorithm="maxima")`

output `1/2*x^2 + 6*x - 1/4*log(x + 1) + 169/4*log(x - 7)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.79

$$\int \frac{2 - x + x^3}{-7 - 6x + x^2} dx = \frac{1}{2} x^2 + 6x - \frac{1}{4} \log(|x + 1|) + \frac{169}{4} \log(|x - 7|)$$

input `integrate((x^3-x+2)/(x^2-6*x-7),x, algorithm="giac")`

output `1/2*x^2 + 6*x - 1/4*log(abs(x + 1)) + 169/4*log(abs(x - 7))`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

$$\int \frac{2 - x + x^3}{-7 - 6x + x^2} dx = 6x - \frac{\ln(x + 1)}{4} + \frac{169 \ln(x - 7)}{4} + \frac{x^2}{2}$$

input `int(-(x^3 - x + 2)/(6*x - x^2 + 7),x)`

output `6*x - log(x + 1)/4 + (169*log(x - 7))/4 + x^2/2`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

$$\int \frac{2 - x + x^3}{-7 - 6x + x^2} dx = \frac{169 \log(x - 7)}{4} - \frac{\log(x + 1)}{4} + \frac{x^2}{2} + 6x$$

input `int((x^3-x+2)/(x^2-6*x-7),x)`

output `(169*log(x - 7) - log(x + 1) + 2*x**2 + 24*x)/4`

3.34 $\int \frac{-1+x^5}{-1+x^2} dx$

Optimal result	230
Mathematica [A] (verified)	230
Rubi [A] (verified)	231
Maple [A] (verified)	232
Fricas [A] (verification not implemented)	232
Sympy [A] (verification not implemented)	233
Maxima [A] (verification not implemented)	233
Giac [A] (verification not implemented)	233
Mupad [B] (verification not implemented)	234
Reduce [B] (verification not implemented)	234

Optimal result

Integrand size = 13, antiderivative size = 19

$$\int \frac{-1+x^5}{-1+x^2} dx = \frac{x^2}{2} + \frac{x^4}{4} + \log(1+x)$$

output `1/2*x^2+1/4*x^4+ln(1+x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{-1+x^5}{-1+x^2} dx = \frac{x^2}{2} + \frac{x^4}{4} + \log(1+x)$$

input `Integrate[(-1 + x^5)/(-1 + x^2),x]`

output `x^2/2 + x^4/4 + Log[1 + x]`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {2341, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^5 - 1}{x^2 - 1} dx$$

$$\downarrow \text{2341}$$

$$\int \left(x^3 - \frac{1-x}{x^2-1} + x \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{x^4}{4} + \frac{x^2}{2} + \log(x+1)$$

input `Int[(-1 + x^5)/(-1 + x^2),x]`

output `x^2/2 + x^4/4 + Log[1 + x]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2341 `Int[(Pq_)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[Pq*(a + b*x^2)^p, x], x] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

method	result	size
default	$\frac{x^2}{2} + \frac{x^4}{4} + \ln(1+x)$	16
norman	$\frac{x^2}{2} + \frac{x^4}{4} + \ln(1+x)$	16
parallelrisc	$\frac{x^2}{2} + \frac{x^4}{4} + \ln(1+x)$	16
risc	$\frac{x^4}{4} + \frac{x^2}{2} + \frac{1}{4} + \ln(1+x)$	17
meijerg	$\operatorname{arctanh}(x) + \frac{x^2(3x^2+6)}{12} + \frac{\ln(-x^2+1)}{2}$	26

input `int((x^5-1)/(x^2-1),x,method=_RETURNVERBOSE)`

output `1/2*x^2+1/4*x^4+ln(1+x)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{-1+x^5}{-1+x^2} dx = \frac{1}{4}x^4 + \frac{1}{2}x^2 + \log(x+1)$$

input `integrate((x^5-1)/(x^2-1),x, algorithm="fricas")`

output `1/4*x^4 + 1/2*x^2 + log(x + 1)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int \frac{-1 + x^5}{-1 + x^2} dx = \frac{x^4}{4} + \frac{x^2}{2} + \log(x + 1)$$

input `integrate((x**5-1)/(x**2-1),x)`output `x**4/4 + x**2/2 + log(x + 1)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{-1 + x^5}{-1 + x^2} dx = \frac{1}{4} x^4 + \frac{1}{2} x^2 + \log(x + 1)$$

input `integrate((x^5-1)/(x^2-1),x, algorithm="maxima")`output `1/4*x^4 + 1/2*x^2 + log(x + 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

$$\int \frac{-1 + x^5}{-1 + x^2} dx = \frac{1}{4} x^4 + \frac{1}{2} x^2 + \log(|x + 1|)$$

input `integrate((x^5-1)/(x^2-1),x, algorithm="giac")`output `1/4*x^4 + 1/2*x^2 + log(abs(x + 1))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{-1 + x^5}{-1 + x^2} dx = \ln(x + 1) + \frac{x^2}{2} + \frac{x^4}{4}$$

input `int((x^5 - 1)/(x^2 - 1),x)`

output `log(x + 1) + x^2/2 + x^4/4`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{-1 + x^5}{-1 + x^2} dx = \log(x + 1) + \frac{x^4}{4} + \frac{x^2}{2}$$

input `int((x^5-1)/(x^2-1),x)`

output `(4*log(x + 1) + x**4 + 2*x**2)/4`

3.35 $\int \frac{5+2x-x^2+x^3}{1+x+x^2} dx$

Optimal result	235
Mathematica [A] (verified)	235
Rubi [A] (verified)	236
Maple [A] (verified)	237
Fricas [A] (verification not implemented)	237
Sympy [A] (verification not implemented)	237
Maxima [A] (verification not implemented)	238
Giac [A] (verification not implemented)	238
Mupad [B] (verification not implemented)	239
Reduce [B] (verification not implemented)	239

Optimal result

Integrand size = 22, antiderivative size = 41

$$\int \frac{5+2x-x^2+x^3}{1+x+x^2} dx = -2x + \frac{x^2}{2} + \frac{11 \arctan\left(\frac{1+2x}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{3}{2} \log(1+x+x^2)$$

output `-2*x+1/2*x^2+11/3*arctan(1/3*(1+2*x)*3^(1/2))*3^(1/2)+3/2*ln(x^2+x+1)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \frac{5+2x-x^2+x^3}{1+x+x^2} dx = -2x + \frac{x^2}{2} + \frac{11 \arctan\left(\frac{1+2x}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{3}{2} \log(1+x+x^2)$$

input `Integrate[(5 + 2*x - x^2 + x^3)/(1 + x + x^2), x]`

output `-2*x + x^2/2 + (11*ArcTan[(1 + 2*x)/Sqrt[3]])/Sqrt[3] + (3*Log[1 + x + x^2])/2`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {2188, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3 - x^2 + 2x + 5}{x^2 + x + 1} dx$$

↓ 2188

$$\int \left(\frac{3x + 7}{x^2 + x + 1} + x - 2 \right) dx$$

↓ 2009

$$\frac{11 \arctan\left(\frac{2x+1}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{x^2}{2} + \frac{3}{2} \log(x^2 + x + 1) - 2x$$

input `Int[(5 + 2*x - x^2 + x^3)/(1 + x + x^2), x]`

output `-2*x + x^2/2 + (11*ArcTan[(1 + 2*x)/Sqrt[3]])/Sqrt[3] + (3*Log[1 + x + x^2])/2`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2188 `Int[(Pq_)*((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[Pq*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.85

method	result	size
default	$-2x + \frac{x^2}{2} + \frac{11 \arctan\left(\frac{(1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{3} + \frac{3 \ln(x^2+x+1)}{2}$	35
risch	$-2x + \frac{x^2}{2} + \frac{3 \ln(4x^2+4x+4)}{2} + \frac{11 \arctan\left(\frac{(1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{3}$	39

input `int((x^3-x^2+2*x+5)/(x^2+x+1),x,method=_RETURNVERBOSE)`output `-2*x+1/2*x^2+11/3*arctan(1/3*(1+2*x)*3^(1/2))*3^(1/2)+3/2*ln(x^2+x+1)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.83

$$\int \frac{5 + 2x - x^2 + x^3}{1 + x + x^2} dx = \frac{1}{2}x^2 + \frac{11}{3}\sqrt{3}\arctan\left(\frac{1}{3}\sqrt{3}(2x+1)\right) - 2x + \frac{3}{2}\log(x^2+x+1)$$

input `integrate((x^3-x^2+2*x+5)/(x^2+x+1),x, algorithm="fricas")`output `1/2*x^2 + 11/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) - 2*x + 3/2*log(x^2 + x + 1)`**Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.12

$$\int \frac{5 + 2x - x^2 + x^3}{1 + x + x^2} dx = \frac{x^2}{2} - 2x + \frac{3 \log(x^2 + x + 1)}{2} + \frac{11\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}x}{3} + \frac{\sqrt{3}}{3}\right)}{3}$$

input `integrate((x**3-x**2+2*x+5)/(x**2+x+1),x)`

output `x**2/2 - 2*x + 3*log(x**2 + x + 1)/2 + 11*sqrt(3)*atan(2*sqrt(3)*x/3 + sqrt(3)/3)/3`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.83

$$\int \frac{5 + 2x - x^2 + x^3}{1 + x + x^2} dx = \frac{1}{2} x^2 + \frac{11}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x + 1)\right) - 2x + \frac{3}{2} \log(x^2 + x + 1)$$

input `integrate((x^3-x^2+2*x+5)/(x^2+x+1),x, algorithm="maxima")`

output `1/2*x^2 + 11/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) - 2*x + 3/2*log(x^2 + x + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.83

$$\int \frac{5 + 2x - x^2 + x^3}{1 + x + x^2} dx = \frac{1}{2} x^2 + \frac{11}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x + 1)\right) - 2x + \frac{3}{2} \log(x^2 + x + 1)$$

input `integrate((x^3-x^2+2*x+5)/(x^2+x+1),x, algorithm="giac")`

output `1/2*x^2 + 11/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) - 2*x + 3/2*log(x^2 + x + 1)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.88

$$\int \frac{5 + 2x - x^2 + x^3}{1 + x + x^2} dx = \frac{3 \ln(x^2 + x + 1)}{2} - 2x + \frac{11\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}x + \sqrt{3}}{3}\right)}{3} + \frac{x^2}{2}$$

input `int((2*x - x^2 + x^3 + 5)/(x + x^2 + 1),x)`output `(3*log(x + x^2 + 1))/2 - 2*x + (11*3^(1/2)*atan((2*3^(1/2)*x)/3 + 3^(1/2)/3))/3 + x^2/2`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.80

$$\int \frac{5 + 2x - x^2 + x^3}{1 + x + x^2} dx = \frac{11\sqrt{3} \operatorname{atan}\left(\frac{2x+1}{\sqrt{3}}\right)}{3} + \frac{3 \log(x^2 + x + 1)}{2} + \frac{x^2}{2} - 2x$$

input `int((x^3-x^2+2*x+5)/(x^2+x+1),x)`output `(22*sqrt(3)*atan((2*x + 1)/sqrt(3)) + 9*log(x**2 + x + 1) + 3*x**2 - 12*x)/6`

$$3.36 \quad \int \frac{-3+x-2x^3+x^4}{10-8x+2x^2} dx$$

Optimal result	240
Mathematica [A] (verified)	240
Rubi [A] (verified)	241
Maple [A] (verified)	242
Fricas [A] (verification not implemented)	242
Sympy [A] (verification not implemented)	242
Maxima [A] (verification not implemented)	243
Giac [A] (verification not implemented)	243
Mupad [B] (verification not implemented)	243
Reduce [B] (verification not implemented)	244

Optimal result

Integrand size = 24, antiderivative size = 41

$$\int \frac{-3+x-2x^3+x^4}{10-8x+2x^2} dx = \frac{3x}{2} + \frac{x^2}{2} + \frac{x^3}{6} + 6 \arctan(2-x) + \frac{3}{4} \log(5-4x+x^2)$$

output `3/2*x+1/2*x^2+1/6*x^3-6*arctan(-2+x)+3/4*ln(x^2-4*x+5)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.95

$$\int \frac{-3+x-2x^3+x^4}{10-8x+2x^2} dx = \frac{1}{2} \left(3x + x^2 + \frac{x^3}{3} + 12 \arctan(2-x) + \frac{3}{2} \log(5-4x+x^2) \right)$$

input `Integrate[(-3 + x - 2*x^3 + x^4)/(10 - 8*x + 2*x^2), x]`

output `(3*x + x^2 + x^3/3 + 12*ArcTan[2 - x] + (3*Log[5 - 4*x + x^2])/2)/2`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2188, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4 - 2x^3 + x - 3}{2x^2 - 8x + 10} dx$$

↓ 2188

$$\int \left(\frac{x^2}{2} - \frac{3(6-x)}{2x^2 - 8x + 10} + x + \frac{3}{2} \right) dx$$

↓ 2009

$$6 \arctan(2-x) + \frac{x^3}{6} + \frac{x^2}{2} + \frac{3}{4} \log(x^2 - 4x + 5) + \frac{3x}{2}$$

input

```
Int[(-3 + x - 2*x^3 + x^4)/(10 - 8*x + 2*x^2),x]
```

output

```
(3*x)/2 + x^2/2 + x^3/6 + 6*ArcTan[2 - x] + (3*Log[5 - 4*x + x^2])/4
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2188

```
Int[(Pq_)*((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Int[Expand
Integrand[Pq*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c}, x] && PolyQ[Pq,
x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.78

method	result	size
default	$\frac{3x}{2} + \frac{x^2}{2} + \frac{x^3}{6} - 6 \arctan(-2 + x) + \frac{3 \ln(x^2 - 4x + 5)}{4}$	32
risch	$\frac{3x}{2} + \frac{x^2}{2} + \frac{x^3}{6} - 6 \arctan(-2 + x) + \frac{3 \ln(x^2 - 4x + 5)}{4}$	32
parallelrisc	$\frac{x^3}{6} + \frac{x^2}{2} + \frac{3x}{2} + \frac{3 \ln(x-2-i)}{4} + 3i \ln(x-2-i) + \frac{3 \ln(x-2+i)}{4} - 3i \ln(x-2+i)$	49

input `int((x^4-2*x^3+x-3)/(2*x^2-8*x+10),x,method=_RETURNVERBOSE)`

output `3/2*x+1/2*x^2+1/6*x^3-6*arctan(-2+x)+3/4*ln(x^2-4*x+5)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.76

$$\int \frac{-3 + x - 2x^3 + x^4}{10 - 8x + 2x^2} dx = \frac{1}{6}x^3 + \frac{1}{2}x^2 + \frac{3}{2}x - 6 \arctan(x - 2) + \frac{3}{4} \log(x^2 - 4x + 5)$$

input `integrate((x^4-2*x^3+x-3)/(2*x^2-8*x+10),x, algorithm="fricas")`

output `1/6*x^3 + 1/2*x^2 + 3/2*x - 6*arctan(x - 2) + 3/4*log(x^2 - 4*x + 5)`

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.83

$$\int \frac{-3 + x - 2x^3 + x^4}{10 - 8x + 2x^2} dx = \frac{x^3}{6} + \frac{x^2}{2} + \frac{3x}{2} + \frac{3 \log(x^2 - 4x + 5)}{4} - 6 \operatorname{atan}(x - 2)$$

input `integrate((x**4-2*x**3+x-3)/(2*x**2-8*x+10),x)`

output `x**3/6 + x**2/2 + 3*x/2 + 3*log(x**2 - 4*x + 5)/4 - 6*atan(x - 2)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.76

$$\int \frac{-3 + x - 2x^3 + x^4}{10 - 8x + 2x^2} dx = \frac{1}{6}x^3 + \frac{1}{2}x^2 + \frac{3}{2}x - 6 \arctan(x - 2) + \frac{3}{4} \log(x^2 - 4x + 5)$$

input `integrate((x^4-2*x^3+x-3)/(2*x^2-8*x+10),x, algorithm="maxima")`output `1/6*x^3 + 1/2*x^2 + 3/2*x - 6*arctan(x - 2) + 3/4*log(x^2 - 4*x + 5)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.76

$$\int \frac{-3 + x - 2x^3 + x^4}{10 - 8x + 2x^2} dx = \frac{1}{6}x^3 + \frac{1}{2}x^2 + \frac{3}{2}x - 6 \arctan(x - 2) + \frac{3}{4} \log(x^2 - 4x + 5)$$

input `integrate((x^4-2*x^3+x-3)/(2*x^2-8*x+10),x, algorithm="giac")`output `1/6*x^3 + 1/2*x^2 + 3/2*x - 6*arctan(x - 2) + 3/4*log(x^2 - 4*x + 5)`**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.76

$$\int \frac{-3 + x - 2x^3 + x^4}{10 - 8x + 2x^2} dx = \frac{3x}{2} - 6 \operatorname{atan}(x - 2) + \frac{3 \ln(x^2 - 4x + 5)}{4} + \frac{x^2}{2} + \frac{x^3}{6}$$

input `int((x - 2*x^3 + x^4 - 3)/(2*x^2 - 8*x + 10),x)`output `(3*x)/2 - 6*atan(x - 2) + (3*log(x^2 - 4*x + 5))/4 + x^2/2 + x^3/6`

Reduce [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.76

$$\int \frac{-3 + x - 2x^3 + x^4}{10 - 8x + 2x^2} dx = -6\operatorname{atan}(x - 2) + \frac{3\log(x^2 - 4x + 5)}{4} + \frac{x^3}{6} + \frac{x^2}{2} + \frac{3x}{2}$$

input `int((x^4-2*x^3+x-3)/(2*x^2-8*x+10),x)`

output `(- 72*atan(x - 2) + 9*log(x**2 - 4*x + 5) + 2*x**3 + 6*x**2 + 18*x)/12`

$$3.37 \quad \int \frac{1+2x+3x^2+x^3}{(-3+x)(-2+x)(-1+x)} dx$$

Optimal result	245
Mathematica [A] (verified)	245
Rubi [A] (verified)	246
Maple [A] (verified)	247
Fricas [A] (verification not implemented)	247
Sympy [A] (verification not implemented)	247
Maxima [A] (verification not implemented)	248
Giac [A] (verification not implemented)	248
Mupad [B] (verification not implemented)	248
Reduce [B] (verification not implemented)	249

Optimal result

Integrand size = 29, antiderivative size = 30

$$\int \frac{1+2x+3x^2+x^3}{(-3+x)(-2+x)(-1+x)} dx = x + \frac{7}{2} \log(1-x) - 25 \log(2-x) + \frac{61}{2} \log(3-x)$$

output `x+7/2*ln(1-x)-25*ln(2-x)+61/2*ln(3-x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.80

$$\int \frac{1+2x+3x^2+x^3}{(-3+x)(-2+x)(-1+x)} dx = x + \frac{61}{2} \log(-3+x) - 25 \log(-2+x) + \frac{7}{2} \log(-1+x)$$

input `Integrate[(1 + 2*x + 3*x^2 + x^3)/((-3 + x)*(-2 + x)*(-1 + x)),x]`

output `x + (61*Log[-3 + x])/2 - 25*Log[-2 + x] + (7*Log[-1 + x])/2`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.069$, Rules used = {2115, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3 + 3x^2 + 2x + 1}{(x-3)(x-2)(x-1)} dx$$

$$\downarrow \text{2115}$$

$$\int \left(-\frac{25}{x-2} + \frac{7}{2(x-1)} + \frac{61}{2(x-3)} + 1 \right) dx$$

$$\downarrow \text{2009}$$

$$x + \frac{7}{2} \log(1-x) - 25 \log(2-x) + \frac{61}{2} \log(3-x)$$

input

```
Int[(1 + 2*x + 3*x^2 + x^3)/((-3 + x)*(-2 + x)*(-1 + x)),x]
```

output

```
x + (7*Log[1 - x])/2 - 25*Log[2 - x] + (61*Log[3 - x])/2
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2115

```
Int[(Px_)*((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_Symbol] := Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && IntegersQ[m, n]
```

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.70

method	result	size
default	$x + \frac{61 \ln(-3+x)}{2} + \frac{7 \ln(-1+x)}{2} - 25 \ln(-2+x)$	21
norman	$x + \frac{61 \ln(-3+x)}{2} + \frac{7 \ln(-1+x)}{2} - 25 \ln(-2+x)$	21
risch	$x + \frac{61 \ln(-3+x)}{2} + \frac{7 \ln(-1+x)}{2} - 25 \ln(-2+x)$	21
parallelsch	$x + \frac{61 \ln(-3+x)}{2} + \frac{7 \ln(-1+x)}{2} - 25 \ln(-2+x)$	21

input `int((x^3+3*x^2+2*x+1)/(-3+x)/(-2+x)/(-1+x),x,method=_RETURNVERBOSE)`output `x+61/2*ln(-3+x)+7/2*ln(-1+x)-25*ln(-2+x)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.67

$$\int \frac{1 + 2x + 3x^2 + x^3}{(-3+x)(-2+x)(-1+x)} dx = x + \frac{7}{2} \log(x-1) - 25 \log(x-2) + \frac{61}{2} \log(x-3)$$

input `integrate((x^3+3*x^2+2*x+1)/(-3+x)/(-2+x)/(-1+x),x, algorithm="fricas")`output `x + 7/2*log(x - 1) - 25*log(x - 2) + 61/2*log(x - 3)`**Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.80

$$\int \frac{1 + 2x + 3x^2 + x^3}{(-3+x)(-2+x)(-1+x)} dx = x + \frac{61 \log(x-3)}{2} - 25 \log(x-2) + \frac{7 \log(x-1)}{2}$$

input `integrate((x**3+3*x**2+2*x+1)/(-3+x)/(-2+x)/(-1+x),x)`

output $x + 61 \cdot \log(x - 3)/2 - 25 \cdot \log(x - 2) + 7 \cdot \log(x - 1)/2$

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.67

$$\int \frac{1 + 2x + 3x^2 + x^3}{(-3 + x)(-2 + x)(-1 + x)} dx = x + \frac{7}{2} \log(x - 1) - 25 \log(x - 2) + \frac{61}{2} \log(x - 3)$$

input `integrate((x^3+3*x^2+2*x+1)/(-3+x)/(-2+x)/(-1+x),x, algorithm="maxima")`

output $x + 7/2 \cdot \log(x - 1) - 25 \cdot \log(x - 2) + 61/2 \cdot \log(x - 3)$

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.77

$$\int \frac{1 + 2x + 3x^2 + x^3}{(-3 + x)(-2 + x)(-1 + x)} dx = x + \frac{7}{2} \log(|x - 1|) - 25 \log(|x - 2|) + \frac{61}{2} \log(|x - 3|)$$

input `integrate((x^3+3*x^2+2*x+1)/(-3+x)/(-2+x)/(-1+x),x, algorithm="giac")`

output $x + 7/2 \cdot \log(\text{abs}(x - 1)) - 25 \cdot \log(\text{abs}(x - 2)) + 61/2 \cdot \log(\text{abs}(x - 3))$

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.67

$$\int \frac{1 + 2x + 3x^2 + x^3}{(-3 + x)(-2 + x)(-1 + x)} dx = x + \frac{7 \ln(x - 1)}{2} - 25 \ln(x - 2) + \frac{61 \ln(x - 3)}{2}$$

input `int((2*x + 3*x^2 + x^3 + 1)/((x - 1)*(x - 2)*(x - 3)),x)`

output $x + (7 \cdot \log(x - 1))/2 - 25 \cdot \log(x - 2) + (61 \cdot \log(x - 3))/2$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.67

$$\int \frac{1 + 2x + 3x^2 + x^3}{(-3 + x)(-2 + x)(-1 + x)} dx = \frac{61 \log(x - 3)}{2} - 25 \log(x - 2) + \frac{7 \log(x - 1)}{2} + x$$

input `int((x^3+3*x^2+2*x+1)/(-3+x)/(-2+x)/(-1+x),x)`

output `(61*log(x - 3) - 50*log(x - 2) + 7*log(x - 1) + 2*x)/2`

$$3.38 \quad \int \frac{2-7x+x^2-x^3+x^4}{-24-14x+x^2+x^3} dx$$

Optimal result	250
Mathematica [A] (verified)	250
Rubi [A] (verified)	251
Maple [A] (verified)	252
Fricas [A] (verification not implemented)	252
Sympy [A] (verification not implemented)	252
Maxima [A] (verification not implemented)	253
Giac [A] (verification not implemented)	253
Mupad [B] (verification not implemented)	254
Reduce [B] (verification not implemented)	254

Optimal result

Integrand size = 30, antiderivative size = 35

$$\int \frac{2-7x+x^2-x^3+x^4}{-24-14x+x^2+x^3} dx = -2x + \frac{x^2}{2} + \frac{13}{3} \log(4-x) - \frac{22}{3} \log(2+x) + 20 \log(3+x)$$

output

```
-2*x+1/2*x^2+13/3*ln(4-x)-22/3*ln(2+x)+20*ln(3+x)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{2-7x+x^2-x^3+x^4}{-24-14x+x^2+x^3} dx = -2x + \frac{x^2}{2} + \frac{13}{3} \log(4-x) - \frac{22}{3} \log(2+x) + 20 \log(3+x)$$

input

```
Integrate[(2 - 7*x + x^2 - x^3 + x^4)/(-24 - 14*x + x^2 + x^3),x]
```

output

```
-2*x + x^2/2 + (13*Log[4 - x])/3 - (22*Log[2 + x])/3 + 20*Log[3 + x]
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$, Rules used = {2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4 - x^3 + x^2 - 7x + 2}{x^3 + x^2 - 14x - 24} dx$$

$$\downarrow \text{2462}$$

$$\int \left(x + \frac{13}{3(x-4)} - \frac{22}{3(x+2)} + \frac{20}{x+3} - 2 \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{x^2}{2} - 2x + \frac{13}{3} \log(4-x) - \frac{22}{3} \log(x+2) + 20 \log(x+3)$$

input

```
Int[(2 - 7*x + x^2 - x^3 + x^4)/(-24 - 14*x + x^2 + x^3),x]
```

output

```
-2*x + x^2/2 + (13*Log[4 - x])/3 - (22*Log[2 + x])/3 + 20*Log[3 + x]
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2462

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegr
and[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ
[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0
] && RationalFunctionQ[u, x]
```

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.80

method	result	size
default	$-2x + \frac{x^2}{2} + 20 \ln(3+x) - \frac{22 \ln(2+x)}{3} + \frac{13 \ln(x-4)}{3}$	28
norman	$-2x + \frac{x^2}{2} + 20 \ln(3+x) - \frac{22 \ln(2+x)}{3} + \frac{13 \ln(x-4)}{3}$	28
risch	$-2x + \frac{x^2}{2} + 20 \ln(3+x) - \frac{22 \ln(2+x)}{3} + \frac{13 \ln(x-4)}{3}$	28
parallelrisc	$-2x + \frac{x^2}{2} + 20 \ln(3+x) - \frac{22 \ln(2+x)}{3} + \frac{13 \ln(x-4)}{3}$	28

input `int((x^4-x^3+x^2-7*x+2)/(x^3+x^2-14*x-24),x,method=_RETURNVERBOSE)`

output `-2*x+1/2*x^2+20*ln(3+x)-22/3*ln(2+x)+13/3*ln(x-4)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.77

$$\int \frac{2 - 7x + x^2 - x^3 + x^4}{-24 - 14x + x^2 + x^3} dx = \frac{1}{2} x^2 - 2x + 20 \log(x+3) - \frac{22}{3} \log(x+2) + \frac{13}{3} \log(x-4)$$

input `integrate((x^4-x^3+x^2-7*x+2)/(x^3+x^2-14*x-24),x, algorithm="fricas")`

output `1/2*x^2 - 2*x + 20*log(x + 3) - 22/3*log(x + 2) + 13/3*log(x - 4)`

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.89

$$\int \frac{2 - 7x + x^2 - x^3 + x^4}{-24 - 14x + x^2 + x^3} dx = \frac{x^2}{2} - 2x + \frac{13 \log(x-4)}{3} - \frac{22 \log(x+2)}{3} + 20 \log(x+3)$$

input `integrate((x**4-x**3+x**2-7*x+2)/(x**3+x**2-14*x-24),x)`

output `x**2/2 - 2*x + 13*log(x - 4)/3 - 22*log(x + 2)/3 + 20*log(x + 3)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.77

$$\int \frac{2 - 7x + x^2 - x^3 + x^4}{-24 - 14x + x^2 + x^3} dx = \frac{1}{2} x^2 - 2x + 20 \log(x+3) - \frac{22}{3} \log(x+2) + \frac{13}{3} \log(x-4)$$

input `integrate((x^4-x^3+x^2-7*x+2)/(x^3+x^2-14*x-24),x, algorithm="maxima")`

output `1/2*x^2 - 2*x + 20*log(x + 3) - 22/3*log(x + 2) + 13/3*log(x - 4)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.86

$$\int \frac{2 - 7x + x^2 - x^3 + x^4}{-24 - 14x + x^2 + x^3} dx = \frac{1}{2} x^2 - 2x + 20 \log(|x+3|) - \frac{22}{3} \log(|x+2|) + \frac{13}{3} \log(|x-4|)$$

input `integrate((x^4-x^3+x^2-7*x+2)/(x^3+x^2-14*x-24),x, algorithm="giac")`

output `1/2*x^2 - 2*x + 20*log(abs(x + 3)) - 22/3*log(abs(x + 2)) + 13/3*log(abs(x - 4))`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.77

$$\int \frac{2 - 7x + x^2 - x^3 + x^4}{-24 - 14x + x^2 + x^3} dx = 20 \ln(x + 3) - \frac{22 \ln(x + 2)}{3} - 2x + \frac{13 \ln(x - 4)}{3} + \frac{x^2}{2}$$

input `int(-(x^2 - 7*x - x^3 + x^4 + 2)/(14*x - x^2 - x^3 + 24),x)`output `20*log(x + 3) - (22*log(x + 2))/3 - 2*x + (13*log(x - 4))/3 + x^2/2`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.77

$$\int \frac{2 - 7x + x^2 - x^3 + x^4}{-24 - 14x + x^2 + x^3} dx = \frac{13 \log(x - 4)}{3} + 20 \log(x + 3) - \frac{22 \log(x + 2)}{3} + \frac{x^2}{2} - 2x$$

input `int((x^4-x^3+x^2-7*x+2)/(x^3+x^2-14*x-24),x)`output `(26*log(x - 4) + 120*log(x + 3) - 44*log(x + 2) + 3*x**2 - 12*x)/6`

$$3.39 \quad \int \frac{2+x^2}{(-1+x)^2 x(1+x)} dx$$

Optimal result	255
Mathematica [A] (verified)	255
Rubi [A] (verified)	256
Maple [A] (verified)	257
Fricas [A] (verification not implemented)	257
Sympy [A] (verification not implemented)	258
Maxima [A] (verification not implemented)	258
Giac [A] (verification not implemented)	258
Mupad [B] (verification not implemented)	259
Reduce [B] (verification not implemented)	259

Optimal result

Integrand size = 19, antiderivative size = 34

$$\int \frac{2+x^2}{(-1+x)^2 x(1+x)} dx = \frac{3}{2(1-x)} - \frac{5}{4} \log(1-x) + 2 \log(x) - \frac{3}{4} \log(1+x)$$

output `3/(2-2*x)-5/4*ln(1-x)+2*ln(x)-3/4*ln(1+x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.94

$$\int \frac{2+x^2}{(-1+x)^2 x(1+x)} dx = -\frac{3}{2(-1+x)} - \frac{5}{4} \log(1-x) + 2 \log(x) - \frac{3}{4} \log(1+x)$$

input `Integrate[(2 + x^2)/((-1 + x)^2*x*(1 + x)),x]`

output `-3/(2*(-1 + x)) - (5*Log[1 - x])/4 + 2*Log[x] - (3*Log[1 + x])/4`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2115, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2 + 2}{(x - 1)^2 x (x + 1)} dx$$

$$\downarrow \text{2115}$$

$$\int \left(\frac{2}{x} - \frac{3}{4(x + 1)} - \frac{5}{4(x - 1)} + \frac{3}{2(x - 1)^2} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{3}{2(1 - x)} - \frac{5}{4} \log(1 - x) + 2 \log(x) - \frac{3}{4} \log(x + 1)$$

input `Int[(2 + x^2)/((-1 + x)^2*x*(1 + x)),x]`

output `3/(2*(1 - x)) - (5*Log[1 - x])/4 + 2*Log[x] - (3*Log[1 + x])/4`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2115 `Int[(Px_)*((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && IntegersQ[m, n]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.74

method	result	size
default	$2 \ln(x) - \frac{3}{2(-1+x)} - \frac{5 \ln(-1+x)}{4} - \frac{3 \ln(1+x)}{4}$	25
norman	$2 \ln(x) - \frac{3}{2(-1+x)} - \frac{5 \ln(-1+x)}{4} - \frac{3 \ln(1+x)}{4}$	25
risch	$2 \ln(x) - \frac{3}{2(-1+x)} - \frac{5 \ln(-1+x)}{4} - \frac{3 \ln(1+x)}{4}$	25
parallelrisc	$\frac{8x \ln(x) - 5 \ln(-1+x)x - 3 \ln(1+x)x - 6 - 8 \ln(x) + 5 \ln(-1+x) + 3 \ln(1+x)}{-4 + 4x}$	45

input `int((x^2+2)/(-1+x)^2/x/(1+x),x,method=_RETURNVERBOSE)`output `2*ln(x)-3/2/(-1+x)-5/4*ln(-1+x)-3/4*ln(1+x)`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00

$$\int \frac{2+x^2}{(-1+x)^2 x(1+x)} dx$$

$$= -\frac{3(x-1)\log(x+1) + 5(x-1)\log(x-1) - 8(x-1)\log(x) + 6}{4(x-1)}$$

input `integrate((x^2+2)/(-1+x)^2/x/(1+x),x, algorithm="fricas")`output `-1/4*(3*(x - 1)*log(x + 1) + 5*(x - 1)*log(x - 1) - 8*(x - 1)*log(x) + 6)/
(x - 1)`

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.79

$$\int \frac{2+x^2}{(-1+x)^2 x(1+x)} dx = 2 \log(x) - \frac{5 \log(x-1)}{4} - \frac{3 \log(x+1)}{4} - \frac{3}{2x-2}$$

input `integrate((x**2+2)/(-1+x)**2/x/(1+x),x)`output `2*log(x) - 5*log(x - 1)/4 - 3*log(x + 1)/4 - 3/(2*x - 2)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.71

$$\int \frac{2+x^2}{(-1+x)^2 x(1+x)} dx = -\frac{3}{2(x-1)} - \frac{3}{4} \log(x+1) - \frac{5}{4} \log(x-1) + 2 \log(x)$$

input `integrate((x^2+2)/(-1+x)^2/x/(1+x),x, algorithm="maxima")`output `-3/2/(x - 1) - 3/4*log(x + 1) - 5/4*log(x - 1) + 2*log(x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00

$$\int \frac{2+x^2}{(-1+x)^2 x(1+x)} dx = -\frac{3}{2(x-1)} + 2 \log\left(\left|-\frac{1}{x-1} - 1\right|\right) - \frac{3}{4} \log\left(\left|-\frac{2}{x-1} - 1\right|\right)$$

input `integrate((x^2+2)/(-1+x)^2/x/(1+x),x, algorithm="giac")`output `-3/2/(x - 1) + 2*log(abs(-1/(x - 1) - 1)) - 3/4*log(abs(-2/(x - 1) - 1))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.76

$$\int \frac{2 + x^2}{(-1 + x)^2 x (1 + x)} dx = 2 \ln(x) - \frac{3 \ln(x + 1)}{4} - \frac{5 \ln(x - 1)}{4} - \frac{3}{2(x - 1)}$$

input `int((x^2 + 2)/(x*(x - 1)^2*(x + 1)),x)`output `2*log(x) - (3*log(x + 1))/4 - (5*log(x - 1))/4 - 3/(2*(x - 1))`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.38

$$\int \frac{2 + x^2}{(-1 + x)^2 x (1 + x)} dx = \frac{-5 \log(x - 1) x + 5 \log(x - 1) - 3 \log(x + 1) x + 3 \log(x + 1) + 8 \log(x) x - 8 \log(x) - 6x}{4x - 4}$$

input `int((x^2+2)/(-1+x)^2/x/(1+x),x)`output `(- 5*log(x - 1)*x + 5*log(x - 1) - 3*log(x + 1)*x + 3*log(x + 1) + 8*log(x)*x - 8*log(x) - 6*x)/(4*(x - 1))`

3.40 $\int \frac{3+x^2+x^3}{(2+x^2)^2} dx$

Optimal result	260
Mathematica [A] (verified)	260
Rubi [A] (verified)	261
Maple [A] (verified)	262
Fricas [A] (verification not implemented)	263
Sympy [A] (verification not implemented)	263
Maxima [A] (verification not implemented)	264
Giac [A] (verification not implemented)	264
Mupad [B] (verification not implemented)	264
Reduce [B] (verification not implemented)	265

Optimal result

Integrand size = 16, antiderivative size = 45

$$\int \frac{3+x^2+x^3}{(2+x^2)^2} dx = \frac{(1-2x)x}{4(2+x^2)} + \frac{5 \arctan\left(\frac{x}{\sqrt{2}}\right)}{4\sqrt{2}} + \frac{1}{2} \log(2+x^2)$$

output `(1-2*x)*x/(4*x^2+8)+5/8*arctan(1/2*x*2^(1/2))*2^(1/2)+1/2*ln(x^2+2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.93

$$\int \frac{3+x^2+x^3}{(2+x^2)^2} dx = \frac{4+x}{4(2+x^2)} + \frac{5 \arctan\left(\frac{x}{\sqrt{2}}\right)}{4\sqrt{2}} + \frac{1}{2} \log(2+x^2)$$

input `Integrate[(3 + x^2 + x^3)/(2 + x^2)^2,x]`

output `(4 + x)/(4*(2 + x^2)) + (5*ArcTan[x/Sqrt[2]])/(4*Sqrt[2]) + Log[2 + x^2]/2`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.96, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {2345, 25, 452, 216, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3 + x^2 + 3}{(x^2 + 2)^2} dx \\
 & \quad \downarrow \text{2345} \\
 & \frac{x + 4}{4(x^2 + 2)} - \frac{1}{4} \int -\frac{4x + 5}{x^2 + 2} dx \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{4} \int \frac{4x + 5}{x^2 + 2} dx + \frac{x + 4}{4(x^2 + 2)} \\
 & \quad \downarrow \text{452} \\
 & \frac{1}{4} \left(5 \int \frac{1}{x^2 + 2} dx + 4 \int \frac{x}{x^2 + 2} dx \right) + \frac{x + 4}{4(x^2 + 2)} \\
 & \quad \downarrow \text{216} \\
 & \frac{1}{4} \left(4 \int \frac{x}{x^2 + 2} dx + \frac{5 \arctan\left(\frac{x}{\sqrt{2}}\right)}{\sqrt{2}} \right) + \frac{x + 4}{4(x^2 + 2)} \\
 & \quad \downarrow \text{240} \\
 & \frac{1}{4} \left(\frac{5 \arctan\left(\frac{x}{\sqrt{2}}\right)}{\sqrt{2}} + 2 \log(x^2 + 2) \right) + \frac{x + 4}{4(x^2 + 2)}
 \end{aligned}$$

input `Int[(3 + x^2 + x^3)/(2 + x^2)^2,x]`

output `(4 + x)/(4*(2 + x^2)) + ((5*ArcTan[x/Sqrt[2]])/Sqrt[2] + 2*Log[2 + x^2])/4`

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 216 $\text{Int}[(\text{a}_) + (\text{b}_) * (\text{x}_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2] * \text{Rt}[\text{b}, 2])) * \text{ArcTan}[\text{Rt}[\text{b}, 2] * (\text{x}/\text{Rt}[\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{PosQ}[\text{a}/\text{b}] \&\& (\text{GtQ}[\text{a}, 0] \parallel \text{GtQ}[\text{b}, 0])$
- rule 240 $\text{Int}[(\text{x}_)/((\text{a}_) + (\text{b}_) * (\text{x}_)^2), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[\text{a} + \text{b} * \text{x}^2, \text{x}]] / (2 * \text{b}), \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}]$
- rule 452 $\text{Int}[(\text{c}_) + (\text{d}_) * (\text{x}_)] / ((\text{a}_) + (\text{b}_) * (\text{x}_)^2), \text{x_Symbol}] \rightarrow \text{Simp}[\text{c} \quad \text{Int}[1/(\text{a} + \text{b} * \text{x}^2), \text{x}], \text{x}] + \text{Simp}[\text{d} \quad \text{Int}[\text{x}/(\text{a} + \text{b} * \text{x}^2), \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \&\& \text{NeQ}[\text{b} * \text{c}^2 + \text{a} * \text{d}^2, 0]$
- rule 2345 $\text{Int}[(\text{Pq}_) * ((\text{a}_) + (\text{b}_) * (\text{x}_)^2)^{\text{p}_}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{Q} = \text{PolynomialQuotient}[\text{Pq}, \text{a} + \text{b} * \text{x}^2, \text{x}], \text{f} = \text{Coeff}[\text{PolynomialRemainder}[\text{Pq}, \text{a} + \text{b} * \text{x}^2, \text{x}], \text{x}, 0], \text{g} = \text{Coeff}[\text{PolynomialRemainder}[\text{Pq}, \text{a} + \text{b} * \text{x}^2, \text{x}], \text{x}, 1]\}, \text{Simp}[(\text{a} * \text{g} - \text{b} * \text{f} * \text{x}) * ((\text{a} + \text{b} * \text{x}^2)^{\text{p} + 1} / (2 * \text{a} * \text{b} * (\text{p} + 1))), \text{x}] + \text{Simp}[1 / (2 * \text{a} * (\text{p} + 1)) \quad \text{Int}[(\text{a} + \text{b} * \text{x}^2)^{\text{p} + 1} * \text{ExpandToSum}[2 * \text{a} * (\text{p} + 1) * \text{Q} + \text{f} * (2 * \text{p} + 3), \text{x}], \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{PolyQ}[\text{Pq}, \text{x}] \&\& \text{LtQ}[\text{p}, -1]$

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.78

method	result	size
default	$\frac{x+1}{x^2+2} + \frac{\ln(x^2+2)}{2} + \frac{5 \arctan\left(\frac{x\sqrt{2}}{2}\right)\sqrt{2}}{8}$	35
risch	$\frac{x+1}{x^2+2} + \frac{\ln(x^2+2)}{2} + \frac{5 \arctan\left(\frac{x\sqrt{2}}{2}\right)\sqrt{2}}{8}$	35
meijerg	$\frac{3\sqrt{2}\left(\frac{x\sqrt{2}}{x^2+2} + \arctan\left(\frac{x\sqrt{2}}{2}\right)\right)}{8} - \frac{x^2}{4\left(\frac{x^2}{2}+1\right)} + \frac{\ln\left(\frac{x^2}{2}+1\right)}{2} + \frac{\sqrt{2}\left(-\frac{x\sqrt{2}}{2\left(\frac{x^2}{2}+1\right)} + \arctan\left(\frac{x\sqrt{2}}{2}\right)\right)}{4}$	79

input `int((x^3+x^2+3)/(x^2+2)^2,x,method=_RETURNVERBOSE)`

output `(1/4*x+1)/(x^2+2)+1/2*ln(x^2+2)+5/8*arctan(1/2*x*2^(1/2))*2^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.98

$$\int \frac{3 + x^2 + x^3}{(2 + x^2)^2} dx = \frac{5\sqrt{2}(x^2 + 2) \arctan\left(\frac{1}{2}\sqrt{2}x\right) + 4(x^2 + 2) \log(x^2 + 2) + 2x + 8}{8(x^2 + 2)}$$

input `integrate((x^3+x^2+3)/(x^2+2)^2,x, algorithm="fricas")`

output `1/8*(5*sqrt(2)*(x^2 + 2)*arctan(1/2*sqrt(2)*x) + 4*(x^2 + 2)*log(x^2 + 2) + 2*x + 8)/(x^2 + 2)`

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.80

$$\int \frac{3 + x^2 + x^3}{(2 + x^2)^2} dx = \frac{x + 4}{4x^2 + 8} + \frac{\log(x^2 + 2)}{2} + \frac{5\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}x}{2}\right)}{8}$$

input `integrate((x**3+x**2+3)/(x**2+2)**2,x)`

output `(x + 4)/(4*x**2 + 8) + log(x**2 + 2)/2 + 5*sqrt(2)*atan(sqrt(2)*x/2)/8`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.73

$$\int \frac{3 + x^2 + x^3}{(2 + x^2)^2} dx = \frac{5}{8} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2} x\right) + \frac{x + 4}{4(x^2 + 2)} + \frac{1}{2} \log(x^2 + 2)$$

input `integrate((x^3+x^2+3)/(x^2+2)^2,x, algorithm="maxima")`output `5/8*sqrt(2)*arctan(1/2*sqrt(2)*x) + 1/4*(x + 4)/(x^2 + 2) + 1/2*log(x^2 + 2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.73

$$\int \frac{3 + x^2 + x^3}{(2 + x^2)^2} dx = \frac{5}{8} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2} x\right) + \frac{x + 4}{4(x^2 + 2)} + \frac{1}{2} \log(x^2 + 2)$$

input `integrate((x^3+x^2+3)/(x^2+2)^2,x, algorithm="giac")`output `5/8*sqrt(2)*arctan(1/2*sqrt(2)*x) + 1/4*(x + 4)/(x^2 + 2) + 1/2*log(x^2 + 2)`**Mupad [B] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.87

$$\int \frac{3 + x^2 + x^3}{(2 + x^2)^2} dx = \frac{\ln(x^2 + 2)}{2} + \frac{5\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}x}{2}\right)}{8} + \frac{x}{4(x^2 + 2)} + \frac{1}{x^2 + 2}$$

input `int((x^2 + x^3 + 3)/(x^2 + 2)^2,x)`output `log(x^2 + 2)/2 + (5*2^(1/2)*atan((2^(1/2)*x)/2))/8 + x/(4*(x^2 + 2)) + 1/(x^2 + 2)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.40

$$\int \frac{3 + x^2 + x^3}{(2 + x^2)^2} dx$$

$$= \frac{5\sqrt{2} \operatorname{atan}\left(\frac{x}{\sqrt{2}}\right) x^2 + 10\sqrt{2} \operatorname{atan}\left(\frac{x}{\sqrt{2}}\right) + 4 \log(x^2 + 2) x^2 + 8 \log(x^2 + 2) - 4x^2 + 2x}{8x^2 + 16}$$

input `int((x^3+x^2+3)/(x^2+2)^2,x)`output `(5*sqrt(2)*atan(x/sqrt(2))*x**2 + 10*sqrt(2)*atan(x/sqrt(2)) + 4*log(x**2 + 2)*x**2 + 8*log(x**2 + 2) - 4*x**2 + 2*x)/(8*(x**2 + 2))`

3.41 $\int \frac{-35+70x-4x^2+2x^3}{(26-10x+x^2)(17-2x+x^2)} dx$

Optimal result	266
Mathematica [A] (verified)	267
Rubi [A] (verified)	267
Maple [A] (verified)	268
Fricas [A] (verification not implemented)	269
Sympy [A] (verification not implemented)	269
Maxima [A] (verification not implemented)	270
Giac [A] (verification not implemented)	270
Mupad [B] (verification not implemented)	271
Reduce [B] (verification not implemented)	271

Optimal result

Integrand size = 36, antiderivative size = 49

$$\int \frac{-35 + 70x - 4x^2 + 2x^3}{(26 - 10x + x^2)(17 - 2x + x^2)} dx = -\frac{15033 \arctan(5 - x)}{1025} - \frac{4607 \arctan\left(\frac{1}{4}(-1 + x)\right)}{4100} + \frac{1003 \log(26 - 10x + x^2)}{1025} + \frac{22 \log(17 - 2x + x^2)}{1025}$$

output `15033/1025*arctan(-5+x)-4607/4100*arctan(-1/4+1/4*x)+1003/1025*ln(x^2-10*x+26)+22/1025*ln(x^2-2*x+17)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00

$$\int \frac{-35 + 70x - 4x^2 + 2x^3}{(26 - 10x + x^2)(17 - 2x + x^2)} dx = -\frac{15033 \arctan(5 - x)}{1025} - \frac{4607 \arctan\left(\frac{1}{4}(-1 + x)\right)}{4100} + \frac{1003 \log(26 - 10x + x^2)}{1025} + \frac{22 \log(17 - 2x + x^2)}{1025}$$

input

```
Integrate[(-35 + 70*x - 4*x^2 + 2*x^3)/((26 - 10*x + x^2)*(17 - 2*x + x^2)),x]
```

output

```
(-15033*ArcTan[5 - x])/1025 - (4607*ArcTan[(-1 + x)/4])/4100 + (1003*Log[26 - 10*x + x^2])/1025 + (22*Log[17 - 2*x + x^2])/1025
```

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {7279, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{2x^3 - 4x^2 + 70x - 35}{(x^2 - 10x + 26)(x^2 - 2x + 17)} dx$$

$$\downarrow \text{7279}$$

$$\int \left(\frac{44x - 4651}{1025(x^2 - 2x + 17)} + \frac{2006x + 5003}{1025(x^2 - 10x + 26)} \right) dx$$

$$\downarrow \text{2009}$$

$$-\frac{15033 \arctan(5 - x)}{1025} - \frac{4607 \arctan\left(\frac{x-1}{4}\right)}{4100} + \frac{1003 \log(x^2 - 10x + 26)}{1025} + \frac{22 \log(x^2 - 2x + 17)}{1025}$$

input `Int[(-35 + 70*x - 4*x^2 + 2*x^3)/((26 - 10*x + x^2)*(17 - 2*x + x^2)),x]`

output `(-15033*ArcTan[5 - x])/1025 - (4607*ArcTan[(-1 + x)/4])/4100 + (1003*Log[26 - 10*x + x^2])/1025 + (22*Log[17 - 2*x + x^2])/1025`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7279 `Int[(u_)/((a_.) + (b_.)*(x_)^(n_.) + (c_.)*(x_)^(n2_.)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n + c*x^(2*n)), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b, c}, x] && EqQ[n2, 2*n] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.78

method	result
default	$\frac{15033 \arctan(x-5)}{1025} - \frac{4607 \arctan(-\frac{1}{4} + \frac{x}{4})}{4100} + \frac{1003 \ln(x^2-10x+26)}{1025} + \frac{22 \ln(x^2-2x+17)}{1025}$
risch	$\frac{15033 \arctan(x-5)}{1025} - \frac{4607 \arctan(-\frac{1}{4} + \frac{x}{4})}{4100} + \frac{1003 \ln(x^2-10x+26)}{1025} + \frac{22 \ln(x^2-2x+17)}{1025}$
paralelrisch	$\frac{1003 \ln(x-5-i)}{1025} - \frac{15033i \ln(x-5-i)}{2050} + \frac{1003 \ln(x-5+i)}{1025} + \frac{15033i \ln(x-5+i)}{2050} + \frac{22 \ln(x-1-4i)}{1025} + \frac{4607i \ln(x-1-4i)}{8200}$

input `int((2*x^3-4*x^2+70*x-35)/(x^2-10*x+26)/(x^2-2*x+17),x,method=_RETURNVERBOSE)`

output `15033/1025*arctan(x-5)-4607/4100*arctan(-1/4+1/4*x)+1003/1025*ln(x^2-10*x+26)+22/1025*ln(x^2-2*x+17)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.76

$$\int \frac{-35 + 70x - 4x^2 + 2x^3}{(26 - 10x + x^2)(17 - 2x + x^2)} dx = \frac{15033}{1025} \arctan(x - 5) - \frac{4607}{4100} \arctan\left(\frac{1}{4}x - \frac{1}{4}\right) + \frac{22}{1025} \log(x^2 - 2x + 17) + \frac{1003}{1025} \log(x^2 - 10x + 26)$$

input `integrate((2*x^3-4*x^2+70*x-35)/(x^2-10*x+26)/(x^2-2*x+17),x, algorithm="fricas")`

output `15033/1025*arctan(x - 5) - 4607/4100*arctan(1/4*x - 1/4) + 22/1025*log(x^2 - 2*x + 17) + 1003/1025*log(x^2 - 10*x + 26)`

Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.94

$$\int \frac{-35 + 70x - 4x^2 + 2x^3}{(26 - 10x + x^2)(17 - 2x + x^2)} dx = \frac{1003 \log(x^2 - 10x + 26)}{1025} + \frac{22 \log(x^2 - 2x + 17)}{1025} - \frac{4607 \operatorname{atan}\left(\frac{x}{4} - \frac{1}{4}\right)}{4100} + \frac{15033 \operatorname{atan}(x - 5)}{1025}$$

input `integrate((2*x**3-4*x**2+70*x-35)/(x**2-10*x+26)/(x**2-2*x+17),x)`

output `1003*log(x**2 - 10*x + 26)/1025 + 22*log(x**2 - 2*x + 17)/1025 - 4607*atan(x/4 - 1/4)/4100 + 15033*atan(x - 5)/1025`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.76

$$\int \frac{-35 + 70x - 4x^2 + 2x^3}{(26 - 10x + x^2)(17 - 2x + x^2)} dx = \frac{15033}{1025} \arctan(x - 5) - \frac{4607}{4100} \arctan\left(\frac{1}{4}x - \frac{1}{4}\right) + \frac{22}{1025} \log(x^2 - 2x + 17) + \frac{1003}{1025} \log(x^2 - 10x + 26)$$

input `integrate((2*x^3-4*x^2+70*x-35)/(x^2-10*x+26)/(x^2-2*x+17),x, algorithm="maxima")`

output `15033/1025*arctan(x - 5) - 4607/4100*arctan(1/4*x - 1/4) + 22/1025*log(x^2 - 2*x + 17) + 1003/1025*log(x^2 - 10*x + 26)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.76

$$\int \frac{-35 + 70x - 4x^2 + 2x^3}{(26 - 10x + x^2)(17 - 2x + x^2)} dx = \frac{15033}{1025} \arctan(x - 5) - \frac{4607}{4100} \arctan\left(\frac{1}{4}x - \frac{1}{4}\right) + \frac{22}{1025} \log(x^2 - 2x + 17) + \frac{1003}{1025} \log(x^2 - 10x + 26)$$

input `integrate((2*x^3-4*x^2+70*x-35)/(x^2-10*x+26)/(x^2-2*x+17),x, algorithm="giac")`

output `15033/1025*arctan(x - 5) - 4607/4100*arctan(1/4*x - 1/4) + 22/1025*log(x^2 - 2*x + 17) + 1003/1025*log(x^2 - 10*x + 26)`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.84

$$\int \frac{-35 + 70x - 4x^2 + 2x^3}{(26 - 10x + x^2)(17 - 2x + x^2)} dx = \ln(x - 1 - 4i) \left(\frac{22}{1025} + \frac{4607}{8200}i \right) \\ + \ln(x - 1 + 4i) \left(\frac{22}{1025} - \frac{4607}{8200}i \right) \\ + \ln(x - 5 - i) \left(\frac{1003}{1025} - \frac{15033}{2050}i \right) \\ + \ln(x - 5 + i) \left(\frac{1003}{1025} + \frac{15033}{2050}i \right)$$

input `int((70*x - 4*x^2 + 2*x^3 - 35)/((x^2 - 2*x + 17)*(x^2 - 10*x + 26)),x)`

output `log(x - (1 + 4i))*(22/1025 + 4607i/8200) + log(x - (1 - 4i))*(22/1025 - 4607i/8200) + log(x - (5 + i))*(1003/1025 - 15033i/2050) + log(x - (5 - i))*(1003/1025 + 15033i/2050)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.76

$$\int \frac{-35 + 70x - 4x^2 + 2x^3}{(26 - 10x + x^2)(17 - 2x + x^2)} dx = \frac{15033 \operatorname{atan}(x - 5)}{1025} - \frac{4607 \operatorname{atan}\left(\frac{x}{4} - \frac{1}{4}\right)}{4100} \\ + \frac{1003 \log(x^2 - 10x + 26)}{1025} \\ + \frac{22 \log(x^2 - 2x + 17)}{1025}$$

input `int((2*x^3-4*x^2+70*x-35)/(x^2-10*x+26)/(x^2-2*x+17),x)`

output `(60132*atan(x - 5) - 4607*atan((x - 1)/4) + 4012*log(x**2 - 10*x + 26) + 88*log(x**2 - 2*x + 17))/4100`

$$3.42 \quad \int \frac{2+x^2}{(-5+x)(-3+x)(4+x)} dx$$

Optimal result	272
Mathematica [A] (verified)	272
Rubi [A] (verified)	273
Maple [A] (verified)	274
Fricas [A] (verification not implemented)	274
Sympy [A] (verification not implemented)	274
Maxima [A] (verification not implemented)	275
Giac [A] (verification not implemented)	275
Mupad [B] (verification not implemented)	275
Reduce [B] (verification not implemented)	276

Optimal result

Integrand size = 21, antiderivative size = 29

$$\int \frac{2+x^2}{(-5+x)(-3+x)(4+x)} dx = -\frac{11}{14} \log(3-x) + \frac{3}{2} \log(5-x) + \frac{2}{7} \log(4+x)$$

output `-11/14*ln(3-x)+3/2*ln(5-x)+2/7*ln(4+x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{2+x^2}{(-5+x)(-3+x)(4+x)} dx = -\frac{11}{14} \log(3-x) + \frac{3}{2} \log(5-x) + \frac{2}{7} \log(4+x)$$

input `Integrate[(2 + x^2)/((-5 + x)*(-3 + x)*(4 + x)),x]`

output `(-11*Log[3 - x])/14 + (3*Log[5 - x])/2 + (2*Log[4 + x])/7`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2115, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2 + 2}{(x - 5)(x - 3)(x + 4)} dx$$

$$\downarrow \text{2115}$$

$$\int \left(-\frac{11}{14(x - 3)} + \frac{2}{7(x + 4)} + \frac{3}{2(x - 5)} \right) dx$$

$$\downarrow \text{2009}$$

$$-\frac{11}{14} \log(3 - x) + \frac{3}{2} \log(5 - x) + \frac{2}{7} \log(x + 4)$$

input

```
Int[(2 + x^2)/((-5 + x)*(-3 + x)*(4 + x)),x]
```

output

```
(-11*Log[3 - x])/14 + (3*Log[5 - x])/2 + (2*Log[4 + x])/7
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2115

```
Int[(Px_)*((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_Symbol] := Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && IntegersQ[m, n]
```

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.69

method	result	size
default	$-\frac{11 \ln(-3+x)}{14} + \frac{3 \ln(x-5)}{2} + \frac{2 \ln(x+4)}{7}$	20
norman	$-\frac{11 \ln(-3+x)}{14} + \frac{3 \ln(x-5)}{2} + \frac{2 \ln(x+4)}{7}$	20
risch	$-\frac{11 \ln(-3+x)}{14} + \frac{3 \ln(x-5)}{2} + \frac{2 \ln(x+4)}{7}$	20
parallelrisc	$-\frac{11 \ln(-3+x)}{14} + \frac{3 \ln(x-5)}{2} + \frac{2 \ln(x+4)}{7}$	20

input `int((x^2+2)/(x-5)/(-3+x)/(x+4),x,method=_RETURNVERBOSE)`output `-11/14*ln(-3+x)+3/2*ln(x-5)+2/7*ln(x+4)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.66

$$\int \frac{2+x^2}{(-5+x)(-3+x)(4+x)} dx = \frac{2}{7} \log(x+4) - \frac{11}{14} \log(x-3) + \frac{3}{2} \log(x-5)$$

input `integrate((x^2+2)/(-5+x)/(-3+x)/(4+x),x, algorithm="fricas")`output `2/7*log(x + 4) - 11/14*log(x - 3) + 3/2*log(x - 5)`**Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

$$\int \frac{2+x^2}{(-5+x)(-3+x)(4+x)} dx = \frac{3 \log(x-5)}{2} - \frac{11 \log(x-3)}{14} + \frac{2 \log(x+4)}{7}$$

input `integrate((x**2+2)/(-5+x)/(-3+x)/(4+x),x)`

output $3*\log(x - 5)/2 - 11*\log(x - 3)/14 + 2*\log(x + 4)/7$

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.66

$$\int \frac{2 + x^2}{(-5 + x)(-3 + x)(4 + x)} dx = \frac{2}{7} \log(x + 4) - \frac{11}{14} \log(x - 3) + \frac{3}{2} \log(x - 5)$$

input `integrate((x^2+2)/(-5+x)/(-3+x)/(4+x),x, algorithm="maxima")`

output $2/7*\log(x + 4) - 11/14*\log(x - 3) + 3/2*\log(x - 5)$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.76

$$\int \frac{2 + x^2}{(-5 + x)(-3 + x)(4 + x)} dx = \frac{2}{7} \log(|x + 4|) - \frac{11}{14} \log(|x - 3|) + \frac{3}{2} \log(|x - 5|)$$

input `integrate((x^2+2)/(-5+x)/(-3+x)/(4+x),x, algorithm="giac")`

output $2/7*\log(\text{abs}(x + 4)) - 11/14*\log(\text{abs}(x - 3)) + 3/2*\log(\text{abs}(x - 5))$

Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.66

$$\int \frac{2 + x^2}{(-5 + x)(-3 + x)(4 + x)} dx = \frac{2 \ln(x + 4)}{7} - \frac{11 \ln(x - 3)}{14} + \frac{3 \ln(x - 5)}{2}$$

input `int((x^2 + 2)/((x - 3)*(x + 4)*(x - 5)),x)`

output $(2*\log(x + 4))/7 - (11*\log(x - 3))/14 + (3*\log(x - 5))/2$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.66

$$\int \frac{2 + x^2}{(-5 + x)(-3 + x)(4 + x)} dx = \frac{3 \log(x - 5)}{2} - \frac{11 \log(x - 3)}{14} + \frac{2 \log(x + 4)}{7}$$

input `int((x^2+2)/(-5+x)/(-3+x)/(4+x),x)`

output `(21*log(x - 5) - 11*log(x - 3) + 4*log(x + 4))/14`

3.43 $\int \frac{x^4}{(-1+x)(2+x^2)} dx$

Optimal result	277
Mathematica [A] (verified)	277
Rubi [A] (verified)	278
Maple [A] (verified)	279
Fricas [A] (verification not implemented)	280
Sympy [A] (verification not implemented)	280
Maxima [A] (verification not implemented)	280
Giac [A] (verification not implemented)	281
Mupad [B] (verification not implemented)	281
Reduce [B] (verification not implemented)	282

Optimal result

Integrand size = 16, antiderivative size = 46

$$\int \frac{x^4}{(-1+x)(2+x^2)} dx = x + \frac{x^2}{2} - \frac{2}{3}\sqrt{2} \arctan\left(\frac{x}{\sqrt{2}}\right) + \frac{1}{3} \log(1-x) - \frac{2}{3} \log(2+x^2)$$

output

```
x+1/2*x^2-2/3*arctan(1/2*x*2^(1/2))*2^(1/2)+1/3*ln(1-x)-2/3*ln(x^2+2)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.93

$$\int \frac{x^4}{(-1+x)(2+x^2)} dx = \frac{1}{6} \left(-9 + 6x + 3x^2 - 4\sqrt{2} \arctan\left(\frac{x}{\sqrt{2}}\right) + 2 \log(-1+x) - 4 \log(2+x^2) \right)$$

input

```
Integrate[x^4/((-1+x)*(2+x^2)),x]
```

output

```
(-9 + 6*x + 3*x^2 - 4*Sqrt[2]*ArcTan[x/Sqrt[2]] + 2*Log[-1 + x] - 4*Log[2 + x^2])/6
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.13, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {604, 27, 2160, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^4}{(x-1)(x^2+2)} dx \\
 & \quad \downarrow \text{604} \\
 & \frac{1}{2} \int \frac{2(-2x^3 + 3x^2 - 4x + 2)}{(1-x)(x^2+2)} dx + \frac{1}{2}(1-x)^2 \\
 & \quad \downarrow \text{27} \\
 & \int \frac{-2x^3 + 3x^2 - 4x + 2}{(1-x)(x^2+2)} dx + \frac{1}{2}(1-x)^2 \\
 & \quad \downarrow \text{2160} \\
 & \int \left(-\frac{4(x+1)}{3(x^2+2)} + \frac{1}{3(x-1)} + 2 \right) dx + \frac{1}{2}(1-x)^2 \\
 & \quad \downarrow \text{2009} \\
 & -\frac{2}{3}\sqrt{2} \arctan\left(\frac{x}{\sqrt{2}}\right) - \frac{2}{3} \log(x^2+2) + \frac{1}{2}(1-x)^2 + 2x + \frac{1}{3} \log(1-x)
 \end{aligned}$$

input `Int[x^4/((-1 + x)*(2 + x^2)),x]`

output `(1 - x)^2/2 + 2*x - (2*sqrt[2]*ArcTan[x/sqrt[2]])/3 + Log[1 - x]/3 - (2*Log[2 + x^2])/3`

Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 604 `Int[(x_)^(m_)*((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(c + d*x)^(m + n - 1)*((a + b*x^2)^(p + 1)/(b*d^(m - 1)*(m + n + 2*p + 1))), x] + Simp[1/(b*d^m*(m + n + 2*p + 1)) Int[(c + d*x)^n*(a + b*x^2)^p*ExpandToSum[b*d^m*(m + n + 2*p + 1)*x^m - b*(m + n + 2*p + 1)*(c + d*x)^m - (c + d*x)^(m - 2)*(a*d^2*(m + n - 1) - b*c^2*(m + n + 2*p + 1) - 2*b*c*d*(m + n + p)*x), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && IGtQ[m, 1] && NeQ[m + n + 2*p + 1, 0] && IntegerQ[2*p]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2160 `Int[(P_q)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*P_q*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, d, e, m}, x] && PolyQ[P_q, x] && IGtQ[p, -2]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.74

method	result	size
default	$x + \frac{x^2}{2} - \frac{2 \ln(x^2+2)}{3} - \frac{2 \arctan\left(\frac{x\sqrt{2}}{2}\right)\sqrt{2}}{3} + \frac{\ln(-1+x)}{3}$	34
risch	$x + \frac{x^2}{2} - \frac{2 \ln(x^2+2)}{3} - \frac{2 \arctan\left(\frac{x\sqrt{2}}{2}\right)\sqrt{2}}{3} + \frac{\ln(-1+x)}{3}$	34

input `int(x^4/(-1+x)/(x^2+2),x,method=_RETURNVERBOSE)`

output `x+1/2*x^2-2/3*ln(x^2+2)-2/3*arctan(1/2*x*2^(1/2))*2^(1/2)+1/3*ln(-1+x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.72

$$\int \frac{x^4}{(-1+x)(2+x^2)} dx = \frac{1}{2}x^2 - \frac{2}{3}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}x\right) + x - \frac{2}{3}\log(x^2+2) + \frac{1}{3}\log(x-1)$$

input `integrate(x^4/(-1+x)/(x^2+2),x, algorithm="fricas")`output `1/2*x^2 - 2/3*sqrt(2)*arctan(1/2*sqrt(2)*x) + x - 2/3*log(x^2 + 2) + 1/3*log(x - 1)`**Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.89

$$\int \frac{x^4}{(-1+x)(2+x^2)} dx = \frac{x^2}{2} + x + \frac{\log(x-1)}{3} - \frac{2\log(x^2+2)}{3} - \frac{2\sqrt{2}\operatorname{atan}\left(\frac{\sqrt{2}x}{2}\right)}{3}$$

input `integrate(x**4/(-1+x)/(x**2+2),x)`output `x**2/2 + x + log(x - 1)/3 - 2*log(x**2 + 2)/3 - 2*sqrt(2)*atan(sqrt(2)*x/2)/3`**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.72

$$\int \frac{x^4}{(-1+x)(2+x^2)} dx = \frac{1}{2}x^2 - \frac{2}{3}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}x\right) + x - \frac{2}{3}\log(x^2+2) + \frac{1}{3}\log(x-1)$$

input `integrate(x^4/(-1+x)/(x^2+2),x, algorithm="maxima")`

output $\frac{1}{2}x^2 - \frac{2}{3}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}x\right) + x - \frac{2}{3}\log(x^2 + 2) + \frac{1}{3}\log(x - 1)$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.74

$$\int \frac{x^4}{(-1+x)(2+x^2)} dx = \frac{1}{2}x^2 - \frac{2}{3}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}x\right) + x - \frac{2}{3}\log(x^2 + 2) + \frac{1}{3}\log(|x - 1|)$$

input `integrate(x^4/(-1+x)/(x^2+2),x, algorithm="giac")`

output $\frac{1}{2}x^2 - \frac{2}{3}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}x\right) + x - \frac{2}{3}\log(x^2 + 2) + \frac{1}{3}\log(\text{abs}(x - 1))$

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.09

$$\int \frac{x^4}{(-1+x)(2+x^2)} dx = x + \frac{\ln(x-1)}{3} + \ln(x - \sqrt{2}1i) \left(-\frac{2}{3} + \frac{\sqrt{2}1i}{3} \right) - \ln(x + \sqrt{2}1i) \left(\frac{2}{3} + \frac{\sqrt{2}1i}{3} \right) + \frac{x^2}{2}$$

input `int(x^4/((x^2 + 2)*(x - 1)),x)`

output $x + \log(x - 1)/3 + \log(x - 2^{(1/2)}1i) * ((2^{(1/2)}1i)/3 - 2/3) - \log(x + 2^{(1/2)}1i) * ((2^{(1/2)}1i)/3 + 2/3) + x^2/2$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.70

$$\int \frac{x^4}{(-1+x)(2+x^2)} dx = -\frac{2\sqrt{2} \operatorname{atan}\left(\frac{x}{\sqrt{2}}\right)}{3} - \frac{2 \log(x^2 + 2)}{3} + \frac{\log(x - 1)}{3} + \frac{x^2}{2} + x$$

input `int(x^4/(-1+x)/(x^2+2),x)`

output `(- 4*sqrt(2)*atan(x/sqrt(2)) - 4*log(x**2 + 2) + 2*log(x - 1) + 3*x**2 + 6*x)/6`

$$3.44 \quad \int \frac{-1+7x+2x^2}{-1-x+x^2+x^3} dx$$

Optimal result	283
Mathematica [A] (verified)	283
Rubi [A] (verified)	284
Maple [A] (verified)	285
Fricas [A] (verification not implemented)	285
Sympy [A] (verification not implemented)	285
Maxima [A] (verification not implemented)	286
Giac [A] (verification not implemented)	286
Mupad [B] (verification not implemented)	286
Reduce [B] (verification not implemented)	287

Optimal result

Integrand size = 24, antiderivative size = 16

$$\int \frac{-1+7x+2x^2}{-1-x+x^2+x^3} dx = -\frac{3}{1+x} + 2 \log(1-x)$$

output

```
-3/(1+x)+2*ln(1-x)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{-1+7x+2x^2}{-1-x+x^2+x^3} dx = -\frac{3}{1+x} + 2 \log(-1+x)$$

input

```
Integrate[(-1 + 7*x + 2*x^2)/(-1 - x + x^2 + x^3),x]
```

output

```
-3/(1 + x) + 2*Log[-1 + x]
```

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{2x^2 + 7x - 1}{x^3 + x^2 - x - 1} dx$$

$$\downarrow \text{2462}$$

$$\int \left(\frac{3}{(x+1)^2} + \frac{2}{x-1} \right) dx$$

$$\downarrow \text{2009}$$

$$2 \log(1-x) - \frac{3}{x+1}$$

input

```
Int[(-1 + 7*x + 2*x^2)/(-1 - x + x^2 + x^3),x]
```

output

```
-3/(1 + x) + 2*Log[1 - x]
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2462

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegr
and[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ
[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0
] && RationalFunctionQ[u, x]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result	size
default	$-\frac{3}{1+x} + 2 \ln(-1+x)$	15
norman	$-\frac{3}{1+x} + 2 \ln(-1+x)$	15
risch	$-\frac{3}{1+x} + 2 \ln(-1+x)$	15
parallelrisch	$\frac{2 \ln(-1+x)x - 3 + 2 \ln(-1+x)}{1+x}$	22

input `int((2*x^2+7*x-1)/(x^3+x^2-x-1),x,method=_RETURNVERBOSE)`

output `-3/(1+x)+2*ln(-1+x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

$$\int \frac{-1 + 7x + 2x^2}{-1 - x + x^2 + x^3} dx = \frac{2(x+1) \log(x-1) - 3}{x+1}$$

input `integrate((2*x^2+7*x-1)/(x^3+x^2-x-1),x, algorithm="fricas")`

output `(2*(x + 1)*log(x - 1) - 3)/(x + 1)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.62

$$\int \frac{-1 + 7x + 2x^2}{-1 - x + x^2 + x^3} dx = 2 \log(x-1) - \frac{3}{x+1}$$

input `integrate((2*x**2+7*x-1)/(x**3+x**2-x-1),x)`

output `2*log(x - 1) - 3/(x + 1)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{-1 + 7x + 2x^2}{-1 - x + x^2 + x^3} dx = -\frac{3}{x + 1} + 2 \log(x - 1)$$

input `integrate((2*x^2+7*x-1)/(x^3+x^2-x-1),x, algorithm="maxima")`

output `-3/(x + 1) + 2*log(x - 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{-1 + 7x + 2x^2}{-1 - x + x^2 + x^3} dx = -\frac{3}{x + 1} + 2 \log(|x - 1|)$$

input `integrate((2*x^2+7*x-1)/(x^3+x^2-x-1),x, algorithm="giac")`

output `-3/(x + 1) + 2*log(abs(x - 1))`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{-1 + 7x + 2x^2}{-1 - x + x^2 + x^3} dx = 2 \ln(x - 1) - \frac{3}{x + 1}$$

input `int(-(7*x + 2*x^2 - 1)/(x - x^2 - x^3 + 1),x)`

output `2*log(x - 1) - 3/(x + 1)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.44

$$\int \frac{-1 + 7x + 2x^2}{-1 - x + x^2 + x^3} dx = \frac{2 \log(x - 1) x + 2 \log(x - 1) + 3x}{x + 1}$$

input `int((2*x^2+7*x-1)/(x^3+x^2-x-1),x)`

output `(2*log(x - 1)*x + 2*log(x - 1) + 3*x)/(x + 1)`

$$3.45 \quad \int \frac{1+2x}{-1+3x-3x^2+x^3} dx$$

Optimal result	288
Mathematica [A] (verified)	288
Rubi [A] (verified)	289
Maple [A] (verified)	290
Fricas [A] (verification not implemented)	290
Sympy [A] (verification not implemented)	291
Maxima [A] (verification not implemented)	291
Giac [A] (verification not implemented)	291
Mupad [B] (verification not implemented)	292
Reduce [B] (verification not implemented)	292

Optimal result

Integrand size = 21, antiderivative size = 18

$$\int \frac{1+2x}{-1+3x-3x^2+x^3} dx = -\frac{(1+2x)^2}{6(1-x)^2}$$

output

```
-1/6*(1+2*x)^2/(1-x)^2
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{1+2x}{-1+3x-3x^2+x^3} dx = \frac{1-4x}{2(-1+x)^2}$$

input

```
Integrate[(1 + 2*x)/(-1 + 3*x - 3*x^2 + x^3),x]
```

output

```
(1 - 4*x)/(2*(-1 + x)^2)
```

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2007, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{2x + 1}{x^3 - 3x^2 + 3x - 1} dx$$

↓ 2007

$$\int \frac{2x + 1}{(x - 1)^3} dx$$

↓ 48

$$-\frac{(2x + 1)^2}{6(1 - x)^2}$$

input

```
Int[(1 + 2*x)/(-1 + 3*x - 3*x^2 + x^3),x]
```

output

```
-1/6*(1 + 2*x)^2/(1 - x)^2
```

Defintions of rubi rules used

rule 48

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

rule 2007

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{a = Rt[Coeff[Px, x, 0], Expon[Px,
x]], b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Int[u*(a + b*x)^(Ex
pon[Px, x]*p), x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; IntegerQ[p] && Pol
yQ[Px, x] && GtQ[Expon[Px, x], 1] && NeQ[Coeff[Px, x, 0], 0]
```

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

method	result	size
norman	$\frac{-2x + \frac{1}{2}}{(-1+x)^2}$	12
default	$-\frac{3}{2(-1+x)^2} - \frac{2}{-1+x}$	16
risch	$\frac{-2x + \frac{1}{2}}{x^2 - 2x + 1}$	17
gospers	$-\frac{-1+4x}{2(x^2-2x+1)}$	18
paralelrisch	$\frac{1-4x}{2x^2-4x+2}$	18
orering	$-\frac{(-1+x)(-1+4x)}{2(x^3-3x^2+3x-1)}$	26

input `int((1+2*x)/(x^3-3*x^2+3*x-1),x,method=_RETURNVERBOSE)`

output `(-2*x+1/2)/(-1+x)^2`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

$$\int \frac{1+2x}{-1+3x-3x^2+x^3} dx = -\frac{4x-1}{2(x^2-2x+1)}$$

input `integrate((1+2*x)/(x^3-3*x^2+3*x-1),x, algorithm="fricas")`

output `-1/2*(4*x - 1)/(x^2 - 2*x + 1)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{1 + 2x}{-1 + 3x - 3x^2 + x^3} dx = \frac{1 - 4x}{2x^2 - 4x + 2}$$

input `integrate((1+2*x)/(x**3-3*x**2+3*x-1),x)`output `(1 - 4*x)/(2*x**2 - 4*x + 2)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

$$\int \frac{1 + 2x}{-1 + 3x - 3x^2 + x^3} dx = -\frac{4x - 1}{2(x^2 - 2x + 1)}$$

input `integrate((1+2*x)/(x^3-3*x^2+3*x-1),x, algorithm="maxima")`output `-1/2*(4*x - 1)/(x^2 - 2*x + 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

$$\int \frac{1 + 2x}{-1 + 3x - 3x^2 + x^3} dx = -\frac{4x - 1}{2(x - 1)^2}$$

input `integrate((1+2*x)/(x^3-3*x^2+3*x-1),x, algorithm="giac")`output `-1/2*(4*x - 1)/(x - 1)^2`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

$$\int \frac{1 + 2x}{-1 + 3x - 3x^2 + x^3} dx = -\frac{4x - 1}{2(x - 1)^2}$$

input `int((2*x + 1)/(3*x - 3*x^2 + x^3 - 1),x)`output `-(4*x - 1)/(2*(x - 1)^2)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{1 + 2x}{-1 + 3x - 3x^2 + x^3} dx = \frac{-2x^2 - 1}{2x^2 - 4x + 2}$$

input `int((1+2*x)/(x^3-3*x^2+3*x-1),x)`output `(- 2*x**2 - 1)/(2*(x**2 - 2*x + 1))`

$$3.46 \quad \int \frac{5-5x+7x^2+x^3}{(-1+x)^2(1+x)^3} dx$$

Optimal result	293
Mathematica [A] (verified)	293
Rubi [A] (verified)	294
Maple [A] (verified)	295
Fricas [A] (verification not implemented)	295
Sympy [A] (verification not implemented)	296
Maxima [A] (verification not implemented)	296
Giac [A] (verification not implemented)	296
Mupad [B] (verification not implemented)	297
Reduce [B] (verification not implemented)	297

Optimal result

Integrand size = 24, antiderivative size = 15

$$\int \frac{5-5x+7x^2+x^3}{(-1+x)^2(1+x)^3} dx = \frac{1}{1-x} - \frac{2}{(1+x)^2}$$

output

```
1/(1-x)-2/(1+x)^2
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{5-5x+7x^2+x^3}{(-1+x)^2(1+x)^3} dx = -\frac{1}{-1+x} - \frac{2}{(1+x)^2}$$

input

```
Integrate[(5 - 5*x + 7*x^2 + x^3)/((-1 + x)^2*(1 + x)^3),x]
```

output

```
-(-1 + x)^(-1) - 2/(1 + x)^2
```

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2123, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3 + 7x^2 - 5x + 5}{(x-1)^2(x+1)^3} dx$$

$$\downarrow \text{2123}$$

$$\int \left(\frac{4}{(x+1)^3} + \frac{1}{(x-1)^2} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{1}{1-x} - \frac{2}{(x+1)^2}$$

input `Int[(5 - 5*x + 7*x^2 + x^3)/((-1 + x)^2*(1 + x)^3),x]`

output `(1 - x)^(-1) - 2/(1 + x)^2`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2123 `Int[(Px_)*((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && PolyQ[Px, x] && (IntegersQ[m, n] || IGtQ[m, -2])`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.07

method	result	size
default	$-\frac{2}{(1+x)^2} - \frac{1}{-1+x}$	16
gosper	$-\frac{x^2+4x-1}{(-1+x)(1+x)^2}$	21
orering	$-\frac{x^2+4x-1}{(-1+x)(1+x)^2}$	21
norman	$\frac{-x^2-4x+1}{(-1+x)(1+x)^2}$	22
risch	$\frac{-x^2-4x+1}{(-1+x)(1+x)^2}$	22
parallelrisc	$\frac{-x^2-4x+1}{(-1+x)(1+x)^2}$	22

input `int((x^3+7*x^2-5*x+5)/(-1+x)^2/(1+x)^3,x,method=_RETURNVERBOSE)`

output `-2/(1+x)^2-1/(-1+x)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.53

$$\int \frac{5 - 5x + 7x^2 + x^3}{(-1+x)^2(1+x)^3} dx = -\frac{x^2 + 4x - 1}{x^3 + x^2 - x - 1}$$

input `integrate((x^3+7*x^2-5*x+5)/(-1+x)^2/(1+x)^3,x, algorithm="fricas")`

output `-(x^2 + 4*x - 1)/(x^3 + x^2 - x - 1)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{5 - 5x + 7x^2 + x^3}{(-1 + x)^2(1 + x)^3} dx = \frac{-x^2 - 4x + 1}{x^3 + x^2 - x - 1}$$

input `integrate((x**3+7*x**2-5*x+5)/(-1+x)**2/(1+x)**3,x)`output `(-x**2 - 4*x + 1)/(x**3 + x**2 - x - 1)`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.53

$$\int \frac{5 - 5x + 7x^2 + x^3}{(-1 + x)^2(1 + x)^3} dx = -\frac{x^2 + 4x - 1}{x^3 + x^2 - x - 1}$$

input `integrate((x^3+7*x^2-5*x+5)/(-1+x)^2/(1+x)^3,x, algorithm="maxima")`output `-(x^2 + 4*x - 1)/(x^3 + x^2 - x - 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 30, normalized size of antiderivative = 2.00

$$\int \frac{5 - 5x + 7x^2 + x^3}{(-1 + x)^2(1 + x)^3} dx = -\frac{1}{x - 1} + \frac{\frac{4}{x-1} + 1}{2\left(\frac{2}{x-1} + 1\right)^2}$$

input `integrate((x^3+7*x^2-5*x+5)/(-1+x)^2/(1+x)^3,x, algorithm="giac")`output `-1/(x - 1) + 1/2*(4/(x - 1) + 1)/(2/(x - 1) + 1)^2`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{5 - 5x + 7x^2 + x^3}{(-1 + x)^2(1 + x)^3} dx = -\frac{1}{x - 1} - \frac{2}{(x + 1)^2}$$

input `int((7*x^2 - 5*x + x^3 + 5)/((x - 1)^2*(x + 1)^3),x)`output `- 1/(x - 1) - 2/(x + 1)^2`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.33

$$\int \frac{5 - 5x + 7x^2 + x^3}{(-1 + x)^2(1 + x)^3} dx = \frac{x(x^2 - 5)}{x^3 + x^2 - x - 1}$$

input `int((x^3+7*x^2-5*x+5)/(-1+x)^2/(1+x)^3,x)`output `(x*(x**2 - 5))/(x**3 + x**2 - x - 1)`

$$3.47 \quad \int \frac{1+3x+3x^2}{1+2x+2x^2+x^3} dx$$

Optimal result	298
Mathematica [A] (verified)	298
Rubi [A] (verified)	299
Maple [A] (verified)	300
Fricas [A] (verification not implemented)	300
Sympy [A] (verification not implemented)	300
Maxima [A] (verification not implemented)	301
Giac [A] (verification not implemented)	301
Mupad [B] (verification not implemented)	301
Reduce [B] (verification not implemented)	302

Optimal result

Integrand size = 26, antiderivative size = 31

$$\int \frac{1+3x+3x^2}{1+2x+2x^2+x^3} dx = -\frac{2 \arctan\left(\frac{1+2x}{\sqrt{3}}\right)}{\sqrt{3}} + \log(1+x) + \log(1+x+x^2)$$

output `-2/3*arctan(1/3*(1+2*x)*3^(1/2))*3^(1/2)+ln(1+x)+ln(x^2+x+1)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{1+3x+3x^2}{1+2x+2x^2+x^3} dx = -\frac{2 \arctan\left(\frac{1+2x}{\sqrt{3}}\right)}{\sqrt{3}} + \log(1+x) + \log(1+x+x^2)$$

input `Integrate[(1 + 3*x + 3*x^2)/(1 + 2*x + 2*x^2 + x^3),x]`

output `(-2*ArcTan[(1 + 2*x)/Sqrt[3]])/Sqrt[3] + Log[1 + x] + Log[1 + x + x^2]`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{3x^2 + 3x + 1}{x^3 + 2x^2 + 2x + 1} dx$$

$$\downarrow 2462$$

$$\int \left(\frac{2x}{x^2 + x + 1} + \frac{1}{x + 1} \right) dx$$

$$\downarrow 2009$$

$$-\frac{2 \arctan\left(\frac{2x+1}{\sqrt{3}}\right)}{\sqrt{3}} + \log(x^2 + x + 1) + \log(x + 1)$$

input

```
Int[(1 + 3*x + 3*x^2)/(1 + 2*x + 2*x^2 + x^3),x]
```

output

```
(-2*ArcTan[(1 + 2*x)/Sqrt[3]])/Sqrt[3] + Log[1 + x] + Log[1 + x + x^2]
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2462

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegr
and[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ
[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0
] && RationalFunctionQ[u, x]
```

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.87

method	result	size
risch	$\ln(1+x) - \frac{2\sqrt{3} \arctan\left(\frac{2\sqrt{3}\left(x+\frac{1}{2}\right)}{3}\right)}{3} + \ln(x^2+x+1)$	27
default	$-\frac{2 \arctan\left(\frac{(1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{3} + \ln(1+x) + \ln(x^2+x+1)$	29

input `int((3*x^2+3*x+1)/(x^3+2*x^2+2*x+1),x,method=_RETURNVERBOSE)`

output `ln(1+x)-2/3*3^(1/2)*arctan(2/3*3^(1/2)*(x+1/2))+ln(x^2+x+1)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.90

$$\int \frac{1+3x+3x^2}{1+2x+2x^2+x^3} dx = -\frac{2}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x+1)\right) + \log(x^2+x+1) + \log(x+1)$$

input `integrate((3*x^2+3*x+1)/(x^3+2*x^2+2*x+1),x, algorithm="fricas")`

output `-2/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + log(x^2 + x + 1) + log(x + 1)`

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.10

$$\int \frac{1+3x+3x^2}{1+2x+2x^2+x^3} dx = \log(x+1)$$

input `integrate((3*x**2+3*x+1)/(x**3+2*x**2+2*x+1),x)`

output `log(x + 1)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.90

$$\int \frac{1 + 3x + 3x^2}{1 + 2x + 2x^2 + x^3} dx = -\frac{2}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x + 1)\right) + \log(x^2 + x + 1) + \log(x + 1)$$

input `integrate((3*x^2+3*x+1)/(x^3+2*x^2+2*x+1),x, algorithm="maxima")`output `-2/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + log(x^2 + x + 1) + log(x + 1)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int \frac{1 + 3x + 3x^2}{1 + 2x + 2x^2 + x^3} dx = -\frac{2}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x + 1)\right) + \log(x^2 + x + 1) + \log(|x + 1|)$$

input `integrate((3*x^2+3*x+1)/(x^3+2*x^2+2*x+1),x, algorithm="giac")`output `-2/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + log(x^2 + x + 1) + log(abs(x + 1))`**Mupad [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.84

$$\int \frac{1 + 3x + 3x^2}{1 + 2x + 2x^2 + x^3} dx = \ln\left(x + \frac{1}{2} - \frac{\sqrt{3} \text{li}}{2}\right) + \ln\left(x + \frac{1}{2} + \frac{\sqrt{3} \text{li}}{2}\right) + \ln(x + 1) + \frac{\sqrt{3} \ln\left(x + \frac{1}{2} - \frac{\sqrt{3} \text{li}}{2}\right) \text{li}}{3} - \frac{\sqrt{3} \ln\left(x + \frac{1}{2} + \frac{\sqrt{3} \text{li}}{2}\right) \text{li}}{3}$$

input `int((3*x + 3*x^2 + 1)/(2*x + 2*x^2 + x^3 + 1),x)`

output

$$\log(x - (3^{1/2})i)/2 + 1/2) + \log(x + (3^{1/2})i)/2 + 1/2) + \log(x + 1) + (3^{1/2})\log(x - (3^{1/2})i)/2 + 1/2)/3 - (3^{1/2})\log(x + (3^{1/2})i)/2 + 1/2)/3$$
Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.87

$$\int \frac{1 + 3x + 3x^2}{1 + 2x + 2x^2 + x^3} dx = -\frac{2\sqrt{3} \operatorname{atan}\left(\frac{2x+1}{\sqrt{3}}\right)}{3} + \log(x^2 + x + 1) + \log(x + 1)$$

input

$$\operatorname{int}((3x^2+3x+1)/(x^3+2x^2+2x+1), x)$$

output

$$(-2\sqrt{3}\operatorname{atan}((2x+1)/\sqrt{3}) + 3\log(x^2 + x + 1) + 3\log(x + 1))/3$$

$$3.48 \quad \int \frac{-1+2x+x^2}{-2x+3x^2+2x^3} dx$$

Optimal result	303
Mathematica [A] (verified)	303
Rubi [A] (verified)	304
Maple [A] (verified)	305
Fricas [A] (verification not implemented)	305
Sympy [A] (verification not implemented)	306
Maxima [A] (verification not implemented)	306
Giac [A] (verification not implemented)	306
Mupad [B] (verification not implemented)	307
Reduce [B] (verification not implemented)	307

Optimal result

Integrand size = 25, antiderivative size = 25

$$\int \frac{-1+2x+x^2}{-2x+3x^2+2x^3} dx = \frac{1}{10} \log(1-2x) + \frac{\log(x)}{2} - \frac{1}{10} \log(2+x)$$

output

```
1/10*ln(1-2*x)+1/2*ln(x)-1/10*ln(2+x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{-1+2x+x^2}{-2x+3x^2+2x^3} dx = \frac{1}{10} \log(1-2x) + \frac{\log(x)}{2} - \frac{1}{10} \log(2+x)$$

input

```
Integrate[(-1 + 2*x + x^2)/(-2*x + 3*x^2 + 2*x^3),x]
```

output

```
Log[1 - 2*x]/10 + Log[x]/2 - Log[2 + x]/10
```


Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2026, 2159, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2 + 2x - 1}{2x^3 + 3x^2 - 2x} dx$$

↓ 2026

$$\int \frac{x^2 + 2x - 1}{x(2x^2 + 3x - 2)} dx$$

↓ 2159

$$\int \left(-\frac{1}{10(x+2)} + \frac{1}{5(2x-1)} + \frac{1}{2x} \right) dx$$

↓ 2009

$$\frac{1}{10} \log(1-2x) + \frac{\log(x)}{2} - \frac{1}{10} \log(x+2)$$

input `Int[(-1 + 2*x + x^2)/(-2*x + 3*x^2 + 2*x^3), x]`

output `Log[1 - 2*x]/10 + Log[x]/2 - Log[2 + x]/10`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2159

```
Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*Pq*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.72

method	result	size
parallelsch	$\frac{\ln(x)}{2} - \frac{\ln(2+x)}{10} + \frac{\ln(x-\frac{1}{2})}{10}$	18
default	$\frac{\ln(x)}{2} - \frac{\ln(2+x)}{10} + \frac{\ln(2x-1)}{10}$	20
norman	$\frac{\ln(x)}{2} - \frac{\ln(2+x)}{10} + \frac{\ln(2x-1)}{10}$	20
risch	$\frac{\ln(x)}{2} - \frac{\ln(2+x)}{10} + \frac{\ln(2x-1)}{10}$	20

input

```
int((x^2+2*x-1)/(2*x^3+3*x^2-2*x),x,method=_RETURNVERBOSE)
```

output

```
1/2*ln(x)-1/10*ln(2+x)+1/10*ln(x-1/2)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{-1 + 2x + x^2}{-2x + 3x^2 + 2x^3} dx = \frac{1}{10} \log(2x - 1) - \frac{1}{10} \log(x + 2) + \frac{1}{2} \log(x)$$

input

```
integrate((x^2+2*x-1)/(2*x^3+3*x^2-2*x),x, algorithm="fricas")
```

output

```
1/10*log(2*x - 1) - 1/10*log(x + 2) + 1/2*log(x)
```

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{-1 + 2x + x^2}{-2x + 3x^2 + 2x^3} dx = \frac{\log(x)}{2} + \frac{\log(x - \frac{1}{2})}{10} - \frac{\log(x + 2)}{10}$$

input `integrate((x**2+2*x-1)/(2*x**3+3*x**2-2*x),x)`output `log(x)/2 + log(x - 1/2)/10 - log(x + 2)/10`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{-1 + 2x + x^2}{-2x + 3x^2 + 2x^3} dx = \frac{1}{10} \log(2x - 1) - \frac{1}{10} \log(x + 2) + \frac{1}{2} \log(x)$$

input `integrate((x^2+2*x-1)/(2*x^3+3*x^2-2*x),x, algorithm="maxima")`output `1/10*log(2*x - 1) - 1/10*log(x + 2) + 1/2*log(x)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \frac{-1 + 2x + x^2}{-2x + 3x^2 + 2x^3} dx = \frac{1}{10} \log(|2x - 1|) - \frac{1}{10} \log(|x + 2|) + \frac{1}{2} \log(|x|)$$

input `integrate((x^2+2*x-1)/(2*x^3+3*x^2-2*x),x, algorithm="giac")`output `1/10*log(abs(2*x - 1)) - 1/10*log(abs(x + 2)) + 1/2*log(abs(x))`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{-1 + 2x + x^2}{-2x + 3x^2 + 2x^3} dx = \frac{\operatorname{atanh}\left(\frac{24}{145\left(\frac{29x}{100} - \frac{11}{50}\right) + \frac{35}{29}}\right)}{5} + \frac{\ln(x)}{2}$$

input `int((2*x + x^2 - 1)/(3*x^2 - 2*x + 2*x^3), x)`output `atanh(24/(145*((29*x)/100 - 11/50)) + 35/29)/5 + log(x)/2`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{-1 + 2x + x^2}{-2x + 3x^2 + 2x^3} dx = \frac{\log(2x - 1)}{10} - \frac{\log(x + 2)}{10} + \frac{\log(x)}{2}$$

input `int((x^2+2*x-1)/(2*x^3+3*x^2-2*x), x)`output `(log(2*x - 1) - log(x + 2) + 5*log(x))/10`

$$3.49 \quad \int \frac{1+4x-2x^2+x^4}{1-x-x^2+x^3} dx$$

Optimal result	308
Mathematica [A] (verified)	308
Rubi [A] (verified)	309
Maple [A] (verified)	310
Fricas [A] (verification not implemented)	310
Sympy [A] (verification not implemented)	311
Maxima [A] (verification not implemented)	311
Giac [A] (verification not implemented)	311
Mupad [B] (verification not implemented)	312
Reduce [B] (verification not implemented)	312

Optimal result

Integrand size = 29, antiderivative size = 22

$$\int \frac{1+4x-2x^2+x^4}{1-x-x^2+x^3} dx = \frac{2}{1-x} + x + \frac{x^2}{2} - 2\operatorname{arctanh}(x)$$

output

```
2/(1-x)+x+1/2*x^2-2*arctanh(x)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.32

$$\int \frac{1+4x-2x^2+x^4}{1-x-x^2+x^3} dx = -\frac{2}{-1+x} + \frac{1}{2}(1+x)^2 + \log(1-x) - \log(1+x)$$

input

```
Integrate[(1 + 4*x - 2*x^2 + x^4)/(1 - x - x^2 + x^3),x]
```

output

```
-2/(-1 + x) + (1 + x)^2/2 + Log[1 - x] - Log[1 + x]
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.36, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.069$, Rules used = {2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4 - 2x^2 + 4x + 1}{x^3 - x^2 - x + 1} dx$$

$$\downarrow \text{2462}$$

$$\int \left(x + \frac{1}{-x-1} + \frac{1}{x-1} + \frac{2}{(x-1)^2} + 1 \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{x^2}{2} + x + \frac{2}{1-x} + \log(1-x) - \log(x+1)$$

input

```
Int[(1 + 4*x - 2*x^2 + x^4)/(1 - x - x^2 + x^3),x]
```

output

```
2/(1 - x) + x + x^2/2 + Log[1 - x] - Log[1 + x]
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2462

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegr
and[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ
[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0
] && RationalFunctionQ[u, x]
```

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.14

method	result	size
default	$x + \frac{x^2}{2} + \ln(-1+x) - \frac{2}{-1+x} - \ln(1+x)$	25
risch	$x + \frac{x^2}{2} + \ln(-1+x) - \frac{2}{-1+x} - \ln(1+x)$	25
norman	$\frac{\frac{1}{2}x^2 + \frac{1}{2}x^3 - 3}{-1+x} - \ln(1+x) + \ln(-1+x)$	30
parallelrisc	$\frac{x^3 + 2\ln(-1+x)x - 2\ln(1+x)x + x^2 - 6 - 2\ln(-1+x) + 2\ln(1+x)}{-2+2x}$	42

input `int((x^4-2*x^2+4*x+1)/(x^3-x^2-x+1),x,method=_RETURNVERBOSE)`

output `x+1/2*x^2+ln(-1+x)-2/(-1+x)-ln(1+x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.64

$$\int \frac{1 + 4x - 2x^2 + x^4}{1 - x - x^2 + x^3} dx$$

$$= \frac{x^3 + x^2 - 2(x-1)\log(x+1) + 2(x-1)\log(x-1) - 2x - 4}{2(x-1)}$$

input `integrate((x^4-2*x^2+4*x+1)/(x^3-x^2-x+1),x, algorithm="fricas")`

output `1/2*(x^3 + x^2 - 2*(x - 1)*log(x + 1) + 2*(x - 1)*log(x - 1) - 2*x - 4)/(x - 1)`

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{1 + 4x - 2x^2 + x^4}{1 - x - x^2 + x^3} dx = \frac{x^2}{2} + x + \log(x - 1) - \log(x + 1) - \frac{2}{x - 1}$$

input `integrate((x**4-2*x**2+4*x+1)/(x**3-x**2-x+1),x)`output `x**2/2 + x + log(x - 1) - log(x + 1) - 2/(x - 1)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{1 + 4x - 2x^2 + x^4}{1 - x - x^2 + x^3} dx = \frac{1}{2}x^2 + x - \frac{2}{x - 1} - \log(x + 1) + \log(x - 1)$$

input `integrate((x^4-2*x^2+4*x+1)/(x^3-x^2-x+1),x, algorithm="maxima")`output `1/2*x^2 + x - 2/(x - 1) - log(x + 1) + log(x - 1)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.18

$$\int \frac{1 + 4x - 2x^2 + x^4}{1 - x - x^2 + x^3} dx = \frac{1}{2}x^2 + x - \frac{2}{x - 1} - \log(|x + 1|) + \log(|x - 1|)$$

input `integrate((x^4-2*x^2+4*x+1)/(x^3-x^2-x+1),x, algorithm="giac")`output `1/2*x^2 + x - 2/(x - 1) - log(abs(x + 1)) + log(abs(x - 1))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1 + 4x - 2x^2 + x^4}{1 - x - x^2 + x^3} dx = x - \frac{2}{x - 1} + \frac{x^2}{2} + \operatorname{atan}(x \operatorname{li} 2) 2i$$

input `int(-(4*x - 2*x^2 + x^4 + 1)/(x + x^2 - x^3 - 1),x)`output `x + atan(x*1i)*2i - 2/(x - 1) + x^2/2`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 44, normalized size of antiderivative = 2.00

$$\int \frac{1 + 4x - 2x^2 + x^4}{1 - x - x^2 + x^3} dx$$

$$= \frac{2 \log(x - 1) x - 2 \log(x - 1) - 2 \log(x + 1) x + 2 \log(x + 1) + x^3 + x^2 - 6x}{2x - 2}$$

input `int((x^4-2*x^2+4*x+1)/(x^3-x^2-x+1),x)`output `(2*log(x - 1)*x - 2*log(x - 1) - 2*log(x + 1)*x + 2*log(x + 1) + x**3 + x**2 - 6*x)/(2*(x - 1))`

3.50 $\int \frac{4-x+2x^2}{4x+x^3} dx$

Optimal result	313
Mathematica [A] (verified)	313
Rubi [A] (verified)	314
Maple [A] (verified)	315
Fricas [A] (verification not implemented)	315
Sympy [A] (verification not implemented)	316
Maxima [A] (verification not implemented)	316
Giac [A] (verification not implemented)	316
Mupad [B] (verification not implemented)	317
Reduce [B] (verification not implemented)	317

Optimal result

Integrand size = 20, antiderivative size = 23

$$\int \frac{4-x+2x^2}{4x+x^3} dx = -\frac{1}{2} \arctan\left(\frac{x}{2}\right) + \log(x) + \frac{1}{2} \log(4+x^2)$$

output

```
-1/2*arctan(1/2*x)+ln(x)+1/2*ln(x^2+4)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{4-x+2x^2}{4x+x^3} dx = -\frac{1}{2} \arctan\left(\frac{x}{2}\right) + \log(x) + \frac{1}{2} \log(4+x^2)$$

input

```
Integrate[(4 - x + 2*x^2)/(4*x + x^3), x]
```

output

```
-1/2*ArcTan[x/2] + Log[x] + Log[4 + x^2]/2
```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {2026, 2333, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{2x^2 - x + 4}{x^3 + 4x} dx \\ & \quad \downarrow \text{2026} \\ & \int \frac{2x^2 - x + 4}{x(x^2 + 4)} dx \\ & \quad \downarrow \text{2333} \\ & \int \left(\frac{x-1}{x^2+4} + \frac{1}{x} \right) dx \\ & \quad \downarrow \text{2009} \\ & -\frac{1}{2} \arctan\left(\frac{x}{2}\right) + \frac{1}{2} \log(x^2 + 4) + \log(x) \end{aligned}$$

input `Int[(4 - x + 2*x^2)/(4*x + x^3),x]`

output `-1/2*ArcTan[x/2] + Log[x] + Log[4 + x^2]/2`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2333

```
Int[(Pq_)*((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[
ExpandIntegrand[(c*x)^m*Pq*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x]
&& PolyQ[Pq, x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

method	result	size
default	$-\frac{\arctan(\frac{x}{2})}{2} + \ln(x) + \frac{\ln(x^2+4)}{2}$	18
risch	$-\frac{\arctan(\frac{x}{2})}{2} + \ln(x) + \frac{\ln(x^2+4)}{2}$	18
meijerg	$\ln(x) - \ln(2) + \frac{\ln(\frac{x^2}{4}+1)}{2} - \frac{\arctan(\frac{x}{2})}{2}$	24
parallelrisc	$\ln(x) + \frac{\ln(x-2i)}{2} + \frac{i \ln(x-2i)}{4} + \frac{\ln(x+2i)}{2} - \frac{i \ln(x+2i)}{4}$	34

input

```
int((2*x^2-x+4)/(x^3+4*x),x,method=_RETURNVERBOSE)
```

output

```
-1/2*arctan(1/2*x)+ln(x)+1/2*ln(x^2+4)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{4-x+2x^2}{4x+x^3} dx = -\frac{1}{2} \arctan\left(\frac{1}{2}x\right) + \frac{1}{2} \log(x^2+4) + \log(x)$$

input

```
integrate((2*x^2-x+4)/(x^3+4*x),x, algorithm="fricas")
```

output

```
-1/2*arctan(1/2*x) + 1/2*log(x^2 + 4) + log(x)
```

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{4 - x + 2x^2}{4x + x^3} dx = \log(x) + \frac{\log(x^2 + 4)}{2} - \frac{\operatorname{atan}\left(\frac{x}{2}\right)}{2}$$

input `integrate((2*x**2-x+4)/(x**3+4*x),x)`output `log(x) + log(x**2 + 4)/2 - atan(x/2)/2`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{4 - x + 2x^2}{4x + x^3} dx = -\frac{1}{2} \arctan\left(\frac{1}{2}x\right) + \frac{1}{2} \log(x^2 + 4) + \log(x)$$

input `integrate((2*x^2-x+4)/(x^3+4*x),x, algorithm="maxima")`output `-1/2*arctan(1/2*x) + 1/2*log(x^2 + 4) + log(x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

$$\int \frac{4 - x + 2x^2}{4x + x^3} dx = -\frac{1}{2} \arctan\left(\frac{1}{2}x\right) + \frac{1}{2} \log(x^2 + 4) + \log(|x|)$$

input `integrate((2*x^2-x+4)/(x^3+4*x),x, algorithm="giac")`output `-1/2*arctan(1/2*x) + 1/2*log(x^2 + 4) + log(abs(x))`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{4 - x + 2x^2}{4x + x^3} dx = \ln(x) + \ln(x - 2i) \left(\frac{1}{2} + \frac{1}{4}i \right) + \ln(x + 2i) \left(\frac{1}{2} - \frac{1}{4}i \right)$$

input `int((2*x^2 - x + 4)/(4*x + x^3),x)`

output `log(x - 2i)*(1/2 + 1i/4) + log(x + 2i)*(1/2 - 1i/4) + log(x)`

Reduce [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{4 - x + 2x^2}{4x + x^3} dx = -\frac{\operatorname{atan}\left(\frac{x}{2}\right)}{2} + \frac{\log(x^2 + 4)}{2} + \log(x)$$

input `int((2*x^2-x+4)/(x^3+4*x),x)`

output `(- atan(x/2) + log(x**2 + 4) + 2*log(x))/2`

3.51 $\int \frac{1+x^2+x^3}{(-1+x)x(1+x^2)^3(1+x+x^2)} dx$

Optimal result	318
Mathematica [A] (verified)	319
Rubi [A] (verified)	319
Maple [A] (verified)	320
Fricas [A] (verification not implemented)	321
Sympy [A] (verification not implemented)	321
Maxima [A] (verification not implemented)	322
Giac [A] (verification not implemented)	322
Mupad [B] (verification not implemented)	323
Reduce [B] (verification not implemented)	323

Optimal result

Integrand size = 32, antiderivative size = 103

$$\int \frac{1+x^2+x^3}{(-1+x)x(1+x^2)^3(1+x+x^2)} dx = \frac{1+x}{8(1+x^2)^2} - \frac{3(1-x)}{8(1+x^2)} + \frac{3x}{16(1+x^2)} + \frac{7 \arctan(x)}{16} - \frac{\arctan\left(\frac{1+2x}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{1}{8} \log(1-x) - \log(x) + \frac{15}{16} \log(1+x^2) - \frac{1}{2} \log(1+x+x^2)$$

output

```
1/8*(1+x)/(x^2+1)^2-3*(1-x)/(8*x^2+8)+3*x/(16*x^2+16)+7/16*arctan(x)-1/3*arctan(1/3*(1+2*x)*3^(1/2))*3^(1/2)+1/8*ln(1-x)-ln(x)+15/16*ln(x^2+1)-1/2*ln(x^2+x+1)
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.90

$$\int \frac{1 + x^2 + x^3}{(-1 + x)x(1 + x^2)^3(1 + x + x^2)} dx = \frac{1}{48} \left(\frac{6(1 + x)}{(1 + x^2)^2} + \frac{9(-2 + 3x)}{1 + x^2} + 21 \arctan(x) \right. \\ \left. - 16\sqrt{3} \arctan\left(\frac{1 + 2x}{\sqrt{3}}\right) + 20 \log(1 - x) \right. \\ \left. - 48 \log(x) + 45 \log(1 + x^2) \right. \\ \left. - 10 \log(1 + x + x^2) - 14 \log(1 - x^3) \right)$$

input

```
Integrate[(1 + x^2 + x^3)/((-1 + x)*x*(1 + x^2)^3*(1 + x + x^2)), x]
```

output

```
((6*(1 + x))/(1 + x^2)^2 + (9*(-2 + 3*x))/(1 + x^2) + 21*ArcTan[x] - 16*Sqrt[3]*ArcTan[(1 + 2*x)/Sqrt[3]] + 20*Log[1 - x] - 48*Log[x] + 45*Log[1 + x^2] - 10*Log[1 + x + x^2] - 14*Log[1 - x^3])/48
```

Rubi [A] (verified)

Time = 0.55 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.062$, Rules used = {7279, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3 + x^2 + 1}{(x - 1)x(x^2 + 1)^3(x^2 + x + 1)} dx \\ \downarrow 7279 \\ \int \left(\frac{-x - 1}{x^2 + x + 1} + \frac{15x - 1}{8(x^2 + 1)} + \frac{3(x + 1)}{4(x^2 + 1)^2} + \frac{1 - x}{2(x^2 + 1)^3} + \frac{1}{8(x - 1)} - \frac{1}{x} \right) dx \\ \downarrow 2009$$

$$\frac{7 \arctan(x)}{16} - \frac{\arctan\left(\frac{2x+1}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{3(1-x)}{8(x^2+1)} + \frac{3x}{16(x^2+1)} + \frac{x+1}{8(x^2+1)^2} + \frac{15}{16} \log(x^2+1) - \frac{1}{2} \log(x^2+x+1) + \frac{1}{8} \log(1-x) - \log(x)$$

input `Int[(1 + x^2 + x^3)/((-1 + x)*x*(1 + x^2)^3*(1 + x + x^2)),x]`

output `(1 + x)/(8*(1 + x^2)^2) - (3*(1 - x))/(8*(1 + x^2)) + (3*x)/(16*(1 + x^2)) + (7*ArcTan[x])/16 - ArcTan[(1 + 2*x)/Sqrt[3]]/Sqrt[3] + Log[1 - x]/8 - Log[x] + (15*Log[1 + x^2])/16 - Log[1 + x + x^2]/2`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7279 `Int[(u_)/((a_.) + (b_.)*(x_)^(n_.) + (c_.)*(x_)^(n2_.)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n + c*x^(2*n)), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b, c}, x] && EqQ[n2, 2*n] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.70

method	result
risch	$\frac{\frac{9}{16}x^3 - \frac{3}{8}x^2 + \frac{11}{16}x - \frac{1}{4}}{(x^2+1)^2} - \ln(x) + \frac{\ln(-1+x)}{8} + \frac{15 \ln(49x^2+49)}{16} + \frac{7 \arctan(x)}{16} - \frac{\sqrt{3} \arctan\left(\frac{2\sqrt{3}\left(x+\frac{1}{2}\right)}{3}\right)}{3} - \frac{\ln(x^2+x+1)}{2}$
default	$-\ln(x) + \frac{\frac{9}{2}x^3 - 3x^2 + \frac{11}{2}x - 2}{8(x^2+1)^2} + \frac{15 \ln(x^2+1)}{16} + \frac{7 \arctan(x)}{16} + \frac{\ln(-1+x)}{8} - \frac{\ln(x^2+x+1)}{2} - \frac{\arctan\left(\frac{(1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{3}$

input `int((x^3+x^2+1)/(-1+x)/x/(x^2+1)^3/(x^2+x+1),x,method=_RETURNVERBOSE)`

output

```
(9/16*x^3-3/8*x^2+11/16*x-1/4)/(x^2+1)^2-ln(x)+1/8*ln(-1+x)+15/16*ln(49*x^2+49)+7/16*arctan(x)-1/3*3^(1/2)*arctan(2/3*3^(1/2)*(x+1/2))-1/2*ln(x^2+x+1)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.32

$$\int \frac{1+x^2+x^3}{(-1+x)x(1+x^2)^3(1+x+x^2)} dx$$

$$= \frac{27x^3 - 16\sqrt{3}(x^4 + 2x^2 + 1) \arctan\left(\frac{1}{3}\sqrt{3}(2x+1)\right) - 18x^2 + 21(x^4 + 2x^2 + 1) \arctan(x) - 24(x^4 + 2x^2 + 1) \log(x^2 + x + 1) + 45(x^4 + 2x^2 + 1) \log(x^2 + 1) + 6(x^4 + 2x^2 + 1) \log(x - 1) - 48(x^4 + 2x^2 + 1) \log(x) + 33x - 12}{16(x^4 + 2x^2 + 1)}$$

input

```
integrate((x^3+x^2+1)/(-1+x)/x/(x^2+1)^3/(x^2+x+1),x, algorithm="fricas")
```

output

```
1/48*(27*x^3 - 16*sqrt(3)*(x^4 + 2*x^2 + 1)*arctan(1/3*sqrt(3)*(2*x + 1)) - 18*x^2 + 21*(x^4 + 2*x^2 + 1)*arctan(x) - 24*(x^4 + 2*x^2 + 1)*log(x^2 + x + 1) + 45*(x^4 + 2*x^2 + 1)*log(x^2 + 1) + 6*(x^4 + 2*x^2 + 1)*log(x - 1) - 48*(x^4 + 2*x^2 + 1)*log(x) + 33*x - 12)/(x^4 + 2*x^2 + 1)
```

Sympy [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.85

$$\int \frac{1+x^2+x^3}{(-1+x)x(1+x^2)^3(1+x+x^2)} dx = -\log(x) + \frac{\log(x-1)}{8} + \frac{15 \log(x^2+1)}{16}$$

$$- \frac{\log(x^2+x+1)}{2} + \frac{7 \operatorname{atan}(x)}{16}$$

$$- \frac{\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}x}{3} + \frac{\sqrt{3}}{3}\right)}{3}$$

$$+ \frac{9x^3 - 6x^2 + 11x - 4}{16x^4 + 32x^2 + 16}$$

input

```
integrate((x**3+x**2+1)/(-1+x)/x/(x**2+1)**3/(x**2+x+1),x)
```

output

```
-log(x) + log(x - 1)/8 + 15*log(x**2 + 1)/16 - log(x**2 + x + 1)/2 + 7*atan(x)/16 - sqrt(3)*atan(2*sqrt(3)*x/3 + sqrt(3)/3)/3 + (9*x**3 - 6*x**2 + 11*x - 4)/(16*x**4 + 32*x**2 + 16)
```

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.75

$$\int \frac{1 + x^2 + x^3}{(-1 + x)x(1 + x^2)^3(1 + x + x^2)} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x + 1)\right) + \frac{9x^3 - 6x^2 + 11x - 4}{16(x^4 + 2x^2 + 1)} + \frac{7}{16} \arctan(x) - \frac{1}{2} \log(x^2 + x + 1) + \frac{15}{16} \log(x^2 + 1) + \frac{1}{8} \log(x - 1) - \log(x)$$

input

```
integrate((x^3+x^2+1)/(-1+x)/x/(x^2+1)^3/(x^2+x+1),x, algorithm="maxima")
```

output

```
-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + 1/16*(9*x^3 - 6*x^2 + 11*x - 4)/(x^4 + 2*x^2 + 1) + 7/16*arctan(x) - 1/2*log(x^2 + x + 1) + 15/16*log(x^2 + 1) + 1/8*log(x - 1) - log(x)
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.72

$$\int \frac{1 + x^2 + x^3}{(-1 + x)x(1 + x^2)^3(1 + x + x^2)} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x + 1)\right) + \frac{9x^3 - 6x^2 + 11x - 4}{16(x^2 + 1)^2} + \frac{7}{16} \arctan(x) - \frac{1}{2} \log(x^2 + x + 1) + \frac{15}{16} \log(x^2 + 1) + \frac{1}{8} \log(|x - 1|) - \log(|x|)$$

input

```
integrate((x^3+x^2+1)/(-1+x)/x/(x^2+1)^3/(x^2+x+1),x, algorithm="giac")
```

output

```
-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + 1/16*(9*x^3 - 6*x^2 + 11*x -
4)/(x^2 + 1)^2 + 7/16*arctan(x) - 1/2*log(x^2 + x + 1) + 15/16*log(x^2 + 1
) + 1/8*log(abs(x - 1)) - log(abs(x))
```

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.93

$$\int \frac{1 + x^2 + x^3}{(-1 + x)x(1 + x^2)^3(1 + x + x^2)} dx = \frac{\ln(x - 1)}{8} - \ln(x) + \ln\left(x + \frac{1}{2} - \frac{\sqrt{3}i}{2}\right) \left(-\frac{1}{2} + \frac{\sqrt{3}i}{6}\right) - \ln\left(x + \frac{1}{2} + \frac{\sqrt{3}i}{2}\right) \left(\frac{1}{2} + \frac{\sqrt{3}i}{6}\right) + \frac{\frac{9x^3}{16} - \frac{3x^2}{8} + \frac{11x}{16} - \frac{1}{4}}{x^4 + 2x^2 + 1} + \ln(x - i) \left(\frac{15}{16} - \frac{7i}{32}\right) + \ln(x + i) \left(\frac{15}{16} + \frac{7i}{32}\right)$$

input

```
int((x^2 + x^3 + 1)/(x*(x^2 + 1)^3*(x - 1)*(x + x^2 + 1)),x)
```

output

```
log(x - 1)/8 + log(x - 1i)*(15/16 - 7i/32) + log(x + 1i)*(15/16 + 7i/32) -
log(x) + log(x - (3^(1/2)*1i)/2 + 1/2)*((3^(1/2)*1i)/6 - 1/2) - log(x + (
3^(1/2)*1i)/2 + 1/2)*((3^(1/2)*1i)/6 + 1/2) + ((11*x)/16 - (3*x^2)/8 + (9*
x^3)/16 - 1/4)/(2*x^2 + x^4 + 1)
```

Reduce [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.98

$$\int \frac{1 + x^2 + x^3}{(-1 + x)x(1 + x^2)^3(1 + x + x^2)} dx = \frac{-16\sqrt{3} \operatorname{atan}\left(\frac{2x+1}{\sqrt{3}}\right) x^4 - 32\sqrt{3} \operatorname{atan}\left(\frac{2x+1}{\sqrt{3}}\right) x^2 - 16\sqrt{3} \operatorname{atan}\left(\frac{2x+1}{\sqrt{3}}\right) + 21 \operatorname{atan}(x) x^4 + 42 \operatorname{atan}(x) x^2 + 21 \operatorname{atan}(x)}{(-1 + x)x(1 + x^2)^3(1 + x + x^2)}$$

input `int((x^3+x^2+1)/(-1+x)/x/(x^2+1)^3/(x^2+x+1),x)`

output `(- 16*sqrt(3)*atan((2*x + 1)/sqrt(3))*x**4 - 32*sqrt(3)*atan((2*x + 1)/sqrt(3))*x**2 - 16*sqrt(3)*atan((2*x + 1)/sqrt(3)) + 21*atan(x)*x**4 + 42*atan(x)*x**2 + 21*atan(x) - 24*log(x**2 + x + 1)*x**4 - 48*log(x**2 + x + 1)*x**2 - 24*log(x**2 + x + 1) + 45*log(x**2 + 1)*x**4 + 90*log(x**2 + 1)*x**2 + 45*log(x**2 + 1) + 6*log(x - 1)*x**4 + 12*log(x - 1)*x**2 + 6*log(x - 1) - 48*log(x)*x**4 - 96*log(x)*x**2 - 48*log(x) + 9*x**4 + 27*x**3 + 33*x - 3)/(48*(x**4 + 2*x**2 + 1))`

$$3.52 \quad \int \frac{1-3x+2x^2-x^3}{(1+x^2)^2} dx$$

Optimal result	325
Mathematica [A] (verified)	325
Rubi [A] (verified)	326
Maple [A] (verified)	327
Fricas [A] (verification not implemented)	328
Sympy [A] (verification not implemented)	328
Maxima [A] (verification not implemented)	329
Giac [A] (verification not implemented)	329
Mupad [B] (verification not implemented)	329
Reduce [B] (verification not implemented)	330

Optimal result

Integrand size = 23, antiderivative size = 34

$$\int \frac{1-3x+2x^2-x^3}{(1+x^2)^2} dx = -\frac{x(1+2x)}{2(1+x^2)} + \frac{3 \arctan(x)}{2} - \frac{1}{2} \log(1+x^2)$$

output `-x*(1+2*x)/(2*x^2+2)+3/2*arctan(x)-1/2*ln(x^2+1)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.88

$$\int \frac{1-3x+2x^2-x^3}{(1+x^2)^2} dx = \frac{1}{2} \left(\frac{2-x}{1+x^2} + 3 \arctan(x) - \log(1+x^2) \right)$$

input `Integrate[(1 - 3*x + 2*x^2 - x^3)/(1 + x^2)^2,x]`

output `((2 - x)/(1 + x^2) + 3*ArcTan[x] - Log[1 + x^2])/2`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$, Rules used = {2345, 25, 452, 216, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{-x^3 + 2x^2 - 3x + 1}{(x^2 + 1)^2} dx$$

$$\downarrow \text{2345}$$

$$\frac{2-x}{2(x^2+1)} - \frac{1}{2} \int -\frac{3-2x}{x^2+1} dx$$

$$\downarrow \text{25}$$

$$\frac{1}{2} \int \frac{3-2x}{x^2+1} dx + \frac{2-x}{2(x^2+1)}$$

$$\downarrow \text{452}$$

$$\frac{1}{2} \left(3 \int \frac{1}{x^2+1} dx - 2 \int \frac{x}{x^2+1} dx \right) + \frac{2-x}{2(x^2+1)}$$

$$\downarrow \text{216}$$

$$\frac{1}{2} \left(3 \arctan(x) - 2 \int \frac{x}{x^2+1} dx \right) + \frac{2-x}{2(x^2+1)}$$

$$\downarrow \text{240}$$

$$\frac{1}{2} (3 \arctan(x) - \log(x^2+1)) + \frac{2-x}{2(x^2+1)}$$

input

 $\text{Int}[(1 - 3*x + 2*x^2 - x^3)/(1 + x^2)^2, x]$

output

 $(2 - x)/(2*(1 + x^2)) + (3*\text{ArcTan}[x] - \text{Log}[1 + x^2])/2$

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 216 $\text{Int}[(\text{a}_) + (\text{b}_) * (\text{x}_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2] * \text{Rt}[\text{b}, 2])) * \text{ArcTan}[\text{Rt}[\text{b}, 2] * (\text{x}/\text{Rt}[\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{PosQ}[\text{a}/\text{b}] \&\& (\text{GtQ}[\text{a}, 0] \parallel \text{GtQ}[\text{b}, 0])$
- rule 240 $\text{Int}[(\text{x}_)/((\text{a}_) + (\text{b}_) * (\text{x}_)^2), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[\text{a} + \text{b} * \text{x}^2, \text{x}]]/(2 * \text{b}), \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}]$
- rule 452 $\text{Int}[(\text{c}_) + (\text{d}_) * (\text{x}_)]/((\text{a}_) + (\text{b}_) * (\text{x}_)^2), \text{x_Symbol}] \rightarrow \text{Simp}[\text{c} \quad \text{Int}[1/(\text{a} + \text{b} * \text{x}^2), \text{x}], \text{x}] + \text{Simp}[\text{d} \quad \text{Int}[\text{x}/(\text{a} + \text{b} * \text{x}^2), \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \&\& \text{NeQ}[\text{b} * \text{c}^2 + \text{a} * \text{d}^2, 0]$
- rule 2345 $\text{Int}[(\text{Pq}_) * ((\text{a}_) + (\text{b}_) * (\text{x}_)^2)^{\text{p}_}], \text{x_Symbol}] \rightarrow \text{With}[\{\text{Q} = \text{PolynomialQuotient}[\text{Pq}, \text{a} + \text{b} * \text{x}^2, \text{x}], \text{f} = \text{Coeff}[\text{PolynomialRemainder}[\text{Pq}, \text{a} + \text{b} * \text{x}^2, \text{x}], \text{x}, 0], \text{g} = \text{Coeff}[\text{PolynomialRemainder}[\text{Pq}, \text{a} + \text{b} * \text{x}^2, \text{x}], \text{x}, 1]\}, \text{Simp}[(\text{a} * \text{g} - \text{b} * \text{f} * \text{x}) * ((\text{a} + \text{b} * \text{x}^2)^{\text{p} + 1}) / (2 * \text{a} * \text{b} * (\text{p} + 1)), \text{x}] + \text{Simp}[1 / (2 * \text{a} * (\text{p} + 1)) \quad \text{Int}[(\text{a} + \text{b} * \text{x}^2)^{\text{p} + 1} * \text{ExpandToSum}[2 * \text{a} * (\text{p} + 1) * \text{Q} + \text{f} * (2 * \text{p} + 3), \text{x}], \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{PolyQ}[\text{Pq}, \text{x}] \&\& \text{LtQ}[\text{p}, -1]$

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.79

method	result	size
risch	$\frac{1 - \frac{\pi}{2}}{x^2 + 1} - \frac{\ln(x^2 + 1)}{2} + \frac{3 \arctan(x)}{2}$	27
default	$-\frac{\frac{\pi}{2} - 1}{x^2 + 1} - \frac{\ln(x^2 + 1)}{2} + \frac{3 \arctan(x)}{2}$	28
meijerg	$-\frac{x^2}{x^2 + 1} - \frac{\ln(x^2 + 1)}{2} - \frac{x}{x^2 + 1} + \frac{3 \arctan(x)}{2} + \frac{x}{2x^2 + 2}$	47
parallelrisch	$-\frac{3i \ln(x-i)x^2 - 3i \ln(x+i)x^2 + 2 \ln(x-i)x^2 + 2 \ln(x+i)x^2 - 4 + 3i \ln(x-i) - 3i \ln(x+i) + 2 \ln(x-i) + 2 \ln(x+i) + 2x}{4(x^2 + 1)}$	87

input `int((-x^3+2*x^2-3*x+1)/(x^2+1)^2,x,method=_RETURNVERBOSE)`

output `(1-1/2*x)/(x^2+1)-1/2*ln(x^2+1)+3/2*arctan(x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.06

$$\int \frac{1 - 3x + 2x^2 - x^3}{(1 + x^2)^2} dx = \frac{3(x^2 + 1) \arctan(x) - (x^2 + 1) \log(x^2 + 1) - x + 2}{2(x^2 + 1)}$$

input `integrate((-x^3+2*x^2-3*x+1)/(x^2+1)^2,x, algorithm="fricas")`

output `1/2*(3*(x^2 + 1)*arctan(x) - (x^2 + 1)*log(x^2 + 1) - x + 2)/(x^2 + 1)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.71

$$\int \frac{1 - 3x + 2x^2 - x^3}{(1 + x^2)^2} dx = -\frac{x - 2}{2x^2 + 2} - \frac{\log(x^2 + 1)}{2} + \frac{3 \operatorname{atan}(x)}{2}$$

input `integrate((-x**3+2*x**2-3*x+1)/(x**2+1)**2,x)`

output `-(x - 2)/(2*x**2 + 2) - log(x**2 + 1)/2 + 3*atan(x)/2`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.74

$$\int \frac{1 - 3x + 2x^2 - x^3}{(1 + x^2)^2} dx = -\frac{x - 2}{2(x^2 + 1)} + \frac{3}{2} \arctan(x) - \frac{1}{2} \log(x^2 + 1)$$

input `integrate((-x^3+2*x^2-3*x+1)/(x^2+1)^2,x, algorithm="maxima")`output `-1/2*(x - 2)/(x^2 + 1) + 3/2*arctan(x) - 1/2*log(x^2 + 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.74

$$\int \frac{1 - 3x + 2x^2 - x^3}{(1 + x^2)^2} dx = -\frac{x - 2}{2(x^2 + 1)} + \frac{3}{2} \arctan(x) - \frac{1}{2} \log(x^2 + 1)$$

input `integrate((-x^3+2*x^2-3*x+1)/(x^2+1)^2,x, algorithm="giac")`output `-1/2*(x - 2)/(x^2 + 1) + 3/2*arctan(x) - 1/2*log(x^2 + 1)`**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.94

$$\int \frac{1 - 3x + 2x^2 - x^3}{(1 + x^2)^2} dx = \frac{3 \operatorname{atan}(x)}{2} - \frac{\ln(x^2 + 1)}{2} - \frac{x}{2(x^2 + 1)} + \frac{1}{x^2 + 1}$$

input `int(-(3*x - 2*x^2 + x^3 - 1)/(x^2 + 1)^2,x)`output `(3*atan(x))/2 - log(x^2 + 1)/2 - x/(2*(x^2 + 1)) + 1/(x^2 + 1)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.44

$$\int \frac{1 - 3x + 2x^2 - x^3}{(1 + x^2)^2} dx$$

$$= \frac{3\operatorname{atan}(x)x^2 + 3\operatorname{atan}(x) - \log(x^2 + 1)x^2 - \log(x^2 + 1) - 2x^2 - x}{2x^2 + 2}$$

input `int((-x^3+2*x^2-3*x+1)/(x^2+1)^2,x)`output `(3*atan(x)*x**2 + 3*atan(x) - log(x**2 + 1)*x**2 - log(x**2 + 1) - 2*x**2 - x)/(2*(x**2 + 1))`

3.53

$$\int \frac{1-3x+2x^2-x^3}{x(1+x^2)^2} dx$$

Optimal result	331
Mathematica [A] (verified)	331
Rubi [A] (verified)	332
Maple [A] (verified)	333
Fricas [A] (verification not implemented)	334
Sympy [A] (verification not implemented)	334
Maxima [A] (verification not implemented)	334
Giac [A] (verification not implemented)	335
Mupad [B] (verification not implemented)	335
Reduce [B] (verification not implemented)	335

Optimal result

Integrand size = 26, antiderivative size = 38

$$\int \frac{1-3x+2x^2-x^3}{x(1+x^2)^2} dx = -\frac{1}{2(1+x^2)} - \frac{x}{1+x^2} - 2 \arctan(x) + \log(x) - \frac{1}{2} \log(1+x^2)$$

output `-1/(2*x^2+2)-x/(x^2+1)-2*arctan(x)+ln(x)-1/2*ln(x^2+1)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.87

$$\int \frac{1-3x+2x^2-x^3}{x(1+x^2)^2} dx = \frac{-1-2x}{2(1+x^2)} - 2 \arctan(x) + \log(x) - \frac{1}{2} \log(1+x^2)$$

input `Integrate[(1 - 3*x + 2*x^2 - x^3)/(x*(1 + x^2)^2),x]`

output `(-1 - 2*x)/(2*(1 + x^2)) - 2*ArcTan[x] + Log[x] - Log[1 + x^2]/2`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.87, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {2336, 27, 523, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{-x^3 + 2x^2 - 3x + 1}{x(x^2 + 1)^2} dx \\
 & \quad \downarrow \text{2336} \\
 & -\frac{1}{2} \int -\frac{2(1 - 2x)}{x(x^2 + 1)} dx - \frac{2x + 1}{2(x^2 + 1)} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{1 - 2x}{x(x^2 + 1)} dx - \frac{2x + 1}{2(x^2 + 1)} \\
 & \quad \downarrow \text{523} \\
 & \int \left(\frac{-x - 2}{x^2 + 1} + \frac{1}{x} \right) dx - \frac{2x + 1}{2(x^2 + 1)} \\
 & \quad \downarrow \text{2009} \\
 & -2 \arctan(x) - \frac{2x + 1}{2(x^2 + 1)} - \frac{1}{2} \log(x^2 + 1) + \log(x)
 \end{aligned}$$

input `Int[(1 - 3*x + 2*x^2 - x^3)/(x*(1 + x^2)^2),x]`

output `-1/2*(1 + 2*x)/(1 + x^2) - 2*ArcTan[x] + Log[x] - Log[1 + x^2]/2`

Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 523 `Int[((x_)^(m_)*((c_) + (d_)*(x_)))/((a_) + (b_)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[x^m*((c + d*x)/(a + b*x^2)), x], x] /; FreeQ[{a, b, c, d}, x] && IntegerQ[m]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2336 `Int[(Pq_)*((c_)*(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{Q = PolynomialQuotient[(c*x)^m*Pq, a + b*x^2, x], f = Coeff[PolynomialRemainder[(c*x)^m*Pq, a + b*x^2, x], x, 0], g = Coeff[PolynomialRemainder[(c*x)^m*Pq, a + b*x^2, x], x, 1]}, Simp[(a*g - b*f*x)*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] + Simp[1/(2*a*(p + 1)) Int[(c*x)^m*(a + b*x^2)^(p + 1)*ExpandToSum[(2*a*(p + 1)*Q)/(c*x)^m + (f*(2*p + 3))/(c*x)^m, x], x], x] /; FreeQ[{a, b, c}, x] && PolyQ[Pq, x] && LtQ[p, -1] && ILtQ[m, 0]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.74

method	result
default	$\ln(x) - \frac{x+\frac{1}{2}}{x^2+1} - \frac{\ln(x^2+1)}{2} - 2 \arctan(x)$
risch	$\frac{-x-\frac{1}{2}}{x^2+1} + \ln(x) - \frac{\ln(4x^2+4)}{2} - 2 \arctan(x)$
meijerg	$-\frac{2x}{2x^2+2} - 2 \arctan(x) + \frac{x^2}{x^2+1} + \frac{1}{2} + \ln(x) - \frac{x^2}{2x^2+2} - \frac{\ln(x^2+1)}{2}$
paralelrisch	$\frac{2i \ln(x-i)x^2 - 2i \ln(x+i)x^2 + 2x^2 \ln(x) - \ln(x-i)x^2 - \ln(x+i)x^2 - 1 + 2i \ln(x-i) - 2i \ln(x+i) + 2 \ln(x) - \ln(x-i) - \ln(x+i) - 2x}{2x^2+2}$

input `int((-x^3+2*x^2-3*x+1)/x/(x^2+1)^2,x,method=_RETURNVERBOSE)`

output `ln(x)-(x+1/2)/(x^2+1)-1/2*ln(x^2+1)-2*arctan(x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.16

$$\int \frac{1 - 3x + 2x^2 - x^3}{x(1 + x^2)^2} dx$$

$$= -\frac{4(x^2 + 1) \arctan(x) + (x^2 + 1) \log(x^2 + 1) - 2(x^2 + 1) \log(x) + 2x + 1}{2(x^2 + 1)}$$

input `integrate((-x^3+2*x^2-3*x+1)/x/(x^2+1)^2,x, algorithm="fricas")`output `-1/2*(4*(x^2 + 1)*arctan(x) + (x^2 + 1)*log(x^2 + 1) - 2*(x^2 + 1)*log(x) + 2*x + 1)/(x^2 + 1)`**Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.71

$$\int \frac{1 - 3x + 2x^2 - x^3}{x(1 + x^2)^2} dx = -\frac{2x + 1}{2x^2 + 2} + \log(x) - \frac{\log(x^2 + 1)}{2} - 2 \operatorname{atan}(x)$$

input `integrate((-x**3+2*x**2-3*x+1)/x/(x**2+1)**2,x)`output `-(2*x + 1)/(2*x**2 + 2) + log(x) - log(x**2 + 1)/2 - 2*atan(x)`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.76

$$\int \frac{1 - 3x + 2x^2 - x^3}{x(1 + x^2)^2} dx = -\frac{2x + 1}{2(x^2 + 1)} - 2 \arctan(x) - \frac{1}{2} \log(x^2 + 1) + \log(x)$$

input `integrate((-x^3+2*x^2-3*x+1)/x/(x^2+1)^2,x, algorithm="maxima")`output `-1/2*(2*x + 1)/(x^2 + 1) - 2*arctan(x) - 1/2*log(x^2 + 1) + log(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.79

$$\int \frac{1 - 3x + 2x^2 - x^3}{x(1+x^2)^2} dx = -\frac{2x+1}{2(x^2+1)} - 2 \arctan(x) - \frac{1}{2} \log(x^2+1) + \log(|x|)$$

input `integrate((-x^3+2*x^2-3*x+1)/x/(x^2+1)^2,x, algorithm="giac")`output `-1/2*(2*x + 1)/(x^2 + 1) - 2*arctan(x) - 1/2*log(x^2 + 1) + log(abs(x))`**Mupad [B] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.87

$$\int \frac{1 - 3x + 2x^2 - x^3}{x(1+x^2)^2} dx = \ln(x) - \frac{x + \frac{1}{2}}{x^2 + 1} + \ln(x-i) \left(-\frac{1}{2} + i\right) + \ln(x+i) \left(-\frac{1}{2} - i\right)$$

input `int(-(3*x - 2*x^2 + x^3 - 1)/(x*(x^2 + 1)^2),x)`output `log(x) - log(x + 1i)*(1/2 + 1i) - log(x - 1i)*(1/2 - 1i) - (x + 1/2)/(x^2 + 1)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.53

$$\int \frac{1 - 3x + 2x^2 - x^3}{x(1+x^2)^2} dx = \frac{-4\operatorname{atan}(x)x^2 - 4\operatorname{atan}(x) - \log(x^2+1)x^2 - \log(x^2+1) + 2\log(x)x^2 + 2\log(x) + x^2 - 2x}{2x^2 + 2}$$

input `int((-x^3+2*x^2-3*x+1)/x/(x^2+1)^2,x)`

output $(-4*\operatorname{atan}(x)*x^{**2} - 4*\operatorname{atan}(x) - \log(x^{**2} + 1)*x^{**2} - \log(x^{**2} + 1) + 2*\log(x)*x^{**2} + 2*\log(x) + x^{**2} - 2*x)/(2*(x^{**2} + 1))$

3.54

$$\int \frac{1-x-x^2+x^3+x^4}{-x+x^3} dx$$

Optimal result	337
Mathematica [A] (verified)	337
Rubi [A] (verified)	338
Maple [A] (verified)	339
Fricas [A] (verification not implemented)	339
Sympy [A] (verification not implemented)	340
Maxima [A] (verification not implemented)	340
Giac [A] (verification not implemented)	340
Mupad [B] (verification not implemented)	341
Reduce [B] (verification not implemented)	341

Optimal result

Integrand size = 26, antiderivative size = 25

$$\int \frac{1-x-x^2+x^3+x^4}{-x+x^3} dx = x + \frac{x^2}{2} - \log(x) + \frac{1}{2} \log(1-x^2)$$

output

```
x+1/2*x^2-ln(x)+1/2*ln(-x^2+1)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{1-x-x^2+x^3+x^4}{-x+x^3} dx = x + \frac{x^2}{2} - \log(x) + \frac{1}{2} \log(1-x^2)$$

input

```
Integrate[(1 - x - x^2 + x^3 + x^4)/(-x + x^3),x]
```

output

```
x + x^2/2 - Log[x] + Log[1 - x^2]/2
```

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$, Rules used = {2026, 2333, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4 + x^3 - x^2 - x + 1}{x^3 - x} dx$$

$$\downarrow \text{2026}$$

$$\int \frac{x^4 + x^3 - x^2 - x + 1}{x(x^2 - 1)} dx$$

$$\downarrow \text{2333}$$

$$\int \left(\frac{x}{x^2 - 1} + x - \frac{1}{x} + 1 \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{x^2}{2} + \frac{1}{2} \log(1 - x^2) + x - \log(x)$$

input `Int[(1 - x - x^2 + x^3 + x^4)/(-x + x^3),x]`

output `x + x^2/2 - Log[x] + Log[1 - x^2]/2`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2333

```
Int[(Pq_)*((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[
ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x]
&& PolyQ[Pq, x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

method	result	size
risch	$x + \frac{x^2}{2} - \ln(x) + \frac{\ln(x^2-1)}{2}$	20
default	$x + \frac{x^2}{2} - \ln(x) + \frac{\ln(-1+x)}{2} + \frac{\ln(1+x)}{2}$	24
norman	$x + \frac{x^2}{2} - \ln(x) + \frac{\ln(-1+x)}{2} + \frac{\ln(1+x)}{2}$	24
parallelrisch	$x + \frac{x^2}{2} - \ln(x) + \frac{\ln(-1+x)}{2} + \frac{\ln(1+x)}{2}$	24
meijerg	$-\ln(x) - \frac{i\pi}{2} + \frac{\ln(-x^2+1)}{2} + \frac{x^2}{2} - \frac{i(2ix-2i \operatorname{arctanh}(x))}{2} + \operatorname{arctanh}(x)$	40

input

```
int((x^4+x^3-x^2-x+1)/(x^3-x),x,method=_RETURNVERBOSE)
```

output

```
x+1/2*x^2-ln(x)+1/2*ln(x^2-1)
```

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{1-x-x^2+x^3+x^4}{-x+x^3} dx = \frac{1}{2}x^2 + x + \frac{1}{2} \log(x^2-1) - \log(x)$$

input

```
integrate((x^4+x^3-x^2-x+1)/(x^3-x),x, algorithm="fricas")
```

output

```
1/2*x^2 + x + 1/2*log(x^2 - 1) - log(x)
```

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.68

$$\int \frac{1 - x - x^2 + x^3 + x^4}{-x + x^3} dx = \frac{x^2}{2} + x - \log(x) + \frac{\log(x^2 - 1)}{2}$$

input `integrate((x**4+x**3-x**2-x+1)/(x**3-x),x)`output `x**2/2 + x - log(x) + log(x**2 - 1)/2`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \frac{1 - x - x^2 + x^3 + x^4}{-x + x^3} dx = \frac{1}{2} x^2 + x + \frac{1}{2} \log(x + 1) + \frac{1}{2} \log(x - 1) - \log(x)$$

input `integrate((x^4+x^3-x^2-x+1)/(x^3-x),x, algorithm="maxima")`output `1/2*x^2 + x + 1/2*log(x + 1) + 1/2*log(x - 1) - log(x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04

$$\int \frac{1 - x - x^2 + x^3 + x^4}{-x + x^3} dx = \frac{1}{2} x^2 + x + \frac{1}{2} \log(|x + 1|) + \frac{1}{2} \log(|x - 1|) - \log(|x|)$$

input `integrate((x^4+x^3-x^2-x+1)/(x^3-x),x, algorithm="giac")`output `1/2*x^2 + x + 1/2*log(abs(x + 1)) + 1/2*log(abs(x - 1)) - log(abs(x))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{1 - x - x^2 + x^3 + x^4}{-x + x^3} dx = x + \frac{\ln(x^2 - 1)}{2} - \ln(x) + \frac{x^2}{2}$$

input `int(-(x^3 - x^2 - x + x^4 + 1)/(x - x^3),x)`output `x + log(x^2 - 1)/2 - log(x) + x^2/2`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \frac{1 - x - x^2 + x^3 + x^4}{-x + x^3} dx = \frac{\log(x - 1)}{2} + \frac{\log(x + 1)}{2} - \log(x) + \frac{x^2}{2} + x$$

input `int((x^4+x^3-x^2-x+1)/(x^3-x),x)`output `(log(x - 1) + log(x + 1) - 2*log(x) + x**2 + 2*x)/2`

$$3.55 \quad \int \frac{2-4x^2+x^3}{(1+x^2)(2+x^2)} dx$$

Optimal result	342
Mathematica [A] (verified)	342
Rubi [A] (verified)	343
Maple [A] (verified)	344
Fricas [A] (verification not implemented)	344
Sympy [A] (verification not implemented)	345
Maxima [A] (verification not implemented)	345
Giac [A] (verification not implemented)	345
Mupad [B] (verification not implemented)	346
Reduce [B] (verification not implemented)	346

Optimal result

Integrand size = 25, antiderivative size = 36

$$\int \frac{2-4x^2+x^3}{(1+x^2)(2+x^2)} dx = 6 \arctan(x) - 5\sqrt{2} \arctan\left(\frac{x}{\sqrt{2}}\right) - \frac{1}{2} \log(1+x^2) + \log(2+x^2)$$

output `6*arctan(x)-5*arctan(1/2*x*2^(1/2))*2^(1/2)-1/2*ln(x^2+1)+ln(x^2+2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00

$$\int \frac{2-4x^2+x^3}{(1+x^2)(2+x^2)} dx = 6 \arctan(x) - 5\sqrt{2} \arctan\left(\frac{x}{\sqrt{2}}\right) - \frac{1}{2} \log(1+x^2) + \log(2+x^2)$$

input `Integrate[(2 - 4*x^2 + x^3)/((1 + x^2)*(2 + x^2)),x]`

output `6*ArcTan[x] - 5*Sqrt[2]*ArcTan[x/Sqrt[2]] - Log[1 + x^2]/2 + Log[2 + x^2]`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3 - 4x^2 + 2}{(x^2 + 1)(x^2 + 2)} dx$$

$$\downarrow 7276$$

$$\int \left(\frac{6 - x}{x^2 + 1} + \frac{2(x - 5)}{x^2 + 2} \right) dx$$

$$\downarrow 2009$$

$$6 \arctan(x) - 5\sqrt{2} \arctan\left(\frac{x}{\sqrt{2}}\right) - \frac{1}{2} \log(x^2 + 1) + \log(x^2 + 2)$$

input `Int[(2 - 4*x^2 + x^3)/((1 + x^2)*(2 + x^2)),x]`

output `6*ArcTan[x] - 5*sqrt[2]*ArcTan[x/sqrt[2]] - Log[1 + x^2]/2 + Log[2 + x^2]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] :> With[{v = RationalFunctionE
x
pand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]`

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.89

method	result	size
default	$6 \arctan(x) - 5 \arctan\left(\frac{x\sqrt{2}}{2}\right) \sqrt{2} - \frac{\ln(x^2+1)}{2} + \ln(x^2+2)$	32
risch	$6 \arctan(x) - 5 \arctan\left(\frac{x\sqrt{2}}{2}\right) \sqrt{2} - \frac{\ln(x^2+1)}{2} + \ln(x^2+2)$	32

input `int((x^3-4*x^2+2)/(x^2+1)/(x^2+2),x,method=_RETURNVERBOSE)`

output `6*arctan(x)-5*arctan(1/2*x*2^(1/2))*2^(1/2)-1/2*ln(x^2+1)+ln(x^2+2)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.86

$$\int \frac{2 - 4x^2 + x^3}{(1 + x^2)(2 + x^2)} dx = -5\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}x\right) + 6 \arctan(x) + \log(x^2 + 2) - \frac{1}{2} \log(x^2 + 1)$$

input `integrate((x^3-4*x^2+2)/(x^2+1)/(x^2+2),x, algorithm="fricas")`

output `-5*sqrt(2)*arctan(1/2*sqrt(2)*x) + 6*arctan(x) + log(x^2 + 2) - 1/2*log(x^2 + 1)`

Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00

$$\int \frac{2 - 4x^2 + x^3}{(1 + x^2)(2 + x^2)} dx = -\frac{\log(x^2 + 1)}{2} + \log(x^2 + 2) + 6 \operatorname{atan}(x) - 5\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}x}{2}\right)$$

input `integrate((x**3-4*x**2+2)/(x**2+1)/(x**2+2),x)`output `-log(x**2 + 1)/2 + log(x**2 + 2) + 6*atan(x) - 5*sqrt(2)*atan(sqrt(2)*x/2)`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.86

$$\int \frac{2 - 4x^2 + x^3}{(1 + x^2)(2 + x^2)} dx = -5\sqrt{2} \operatorname{arctan}\left(\frac{1}{2}\sqrt{2}x\right) + 6 \operatorname{arctan}(x) + \log(x^2 + 2) - \frac{1}{2} \log(x^2 + 1)$$

input `integrate((x^3-4*x^2+2)/(x^2+1)/(x^2+2),x, algorithm="maxima")`output `-5*sqrt(2)*arctan(1/2*sqrt(2)*x) + 6*arctan(x) + log(x^2 + 2) - 1/2*log(x^2 + 1)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.86

$$\int \frac{2 - 4x^2 + x^3}{(1 + x^2)(2 + x^2)} dx = -5\sqrt{2} \operatorname{arctan}\left(\frac{1}{2}\sqrt{2}x\right) + 6 \operatorname{arctan}(x) + \log(x^2 + 2) - \frac{1}{2} \log(x^2 + 1)$$

input `integrate((x^3-4*x^2+2)/(x^2+1)/(x^2+2),x, algorithm="giac")`

output `-5*sqrt(2)*arctan(1/2*sqrt(2)*x) + 6*arctan(x) + log(x^2 + 2) - 1/2*log(x^2 + 1)`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.56

$$\int \frac{2 - 4x^2 + x^3}{(1 + x^2)(2 + x^2)} dx = \ln(x - i) \left(-\frac{1}{2} - 3i \right) + \ln(x + 1i) \left(-\frac{1}{2} + 3i \right) \\ + \ln(x - \sqrt{2}1i) \left(1 + \frac{\sqrt{2}5i}{2} \right) - \ln(x + \sqrt{2}1i) \left(-1 + \frac{\sqrt{2}5i}{2} \right)$$

input `int((x^3 - 4*x^2 + 2)/((x^2 + 1)*(x^2 + 2)),x)`

output `log(x - 2^(1/2)*1i)*((2^(1/2)*5i)/2 + 1) - log(x + 1i)*(1/2 - 3i) - log(x - 1i)*(1/2 + 3i) - log(x + 2^(1/2)*1i)*((2^(1/2)*5i)/2 - 1)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.83

$$\int \frac{2 - 4x^2 + x^3}{(1 + x^2)(2 + x^2)} dx = -5\sqrt{2} \operatorname{atan}\left(\frac{x}{\sqrt{2}}\right) + 6\operatorname{atan}(x) + \log(x^2 + 2) - \frac{\log(x^2 + 1)}{2}$$

input `int((x^3-4*x^2+2)/(x^2+1)/(x^2+2),x)`

output `(- 10*sqrt(2)*atan(x/sqrt(2)) + 12*atan(x) + 2*log(x**2 + 2) - log(x**2 + 1))/2`

$$3.56 \quad \int \frac{1+x^2+x^4}{(1+x^2)(4+x^2)^2} dx$$

Optimal result	347
Mathematica [A] (verified)	347
Rubi [A] (verified)	348
Maple [A] (verified)	349
Fricas [A] (verification not implemented)	349
Sympy [A] (verification not implemented)	349
Maxima [A] (verification not implemented)	350
Giac [A] (verification not implemented)	350
Mupad [B] (verification not implemented)	350
Reduce [B] (verification not implemented)	351

Optimal result

Integrand size = 23, antiderivative size = 29

$$\int \frac{1+x^2+x^4}{(1+x^2)(4+x^2)^2} dx = -\frac{13x}{24(4+x^2)} + \frac{25}{144} \arctan\left(\frac{x}{2}\right) + \frac{\arctan(x)}{9}$$

output `-13*x/(24*x^2+96)+25/144*arctan(1/2*x)+1/9*arctan(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{1+x^2+x^4}{(1+x^2)(4+x^2)^2} dx = -\frac{13x}{24(4+x^2)} + \frac{25}{144} \arctan\left(\frac{x}{2}\right) + \frac{\arctan(x)}{9}$$

input `Integrate[(1 + x^2 + x^4)/((1 + x^2)*(4 + x^2)^2), x]`

output `(-13*x)/(24*(4 + x^2)) + (25*ArcTan[x/2])/144 + ArcTan[x]/9`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4 + x^2 + 1}{(x^2 + 1)(x^2 + 4)^2} dx$$

↓ 7276

$$\int \left(\frac{8}{9(x^2 + 4)} - \frac{13}{3(x^2 + 4)^2} + \frac{1}{9(x^2 + 1)} \right) dx$$

↓ 2009

$$\frac{25}{144} \arctan\left(\frac{x}{2}\right) + \frac{\arctan(x)}{9} - \frac{13x}{24(x^2 + 4)}$$

input `Int[(1 + x^2 + x^4)/((1 + x^2)*(4 + x^2)^2), x]`

output `(-13*x)/(24*(4 + x^2)) + (25*ArcTan[x/2])/144 + ArcTan[x]/9`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] :> With[{v = RationalFunctionE
x
pand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]`

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.76

method	result
default	$\frac{\arctan(x)}{9} - \frac{13x}{24(x^2+4)} + \frac{25 \arctan(\frac{x}{2})}{144}$
risch	$\frac{\arctan(x)}{9} - \frac{13x}{24(x^2+4)} + \frac{25 \arctan(\frac{x}{2})}{144}$
parallelrisch	$-\frac{16i \ln(x-i)x^2 + 25i \ln(x-2i)x^2 - 16i \ln(x+i)x^2 - 25i \ln(x+2i)x^2 + 64i \ln(x-i) + 100i \ln(x-2i) - 64i \ln(x+i) - 100i \ln(x+2i)}{288(x^2+4)}$

input `int((x^4+x^2+1)/(x^2+1)/(x^2+4)^2,x,method=_RETURNVERBOSE)`output `1/9*arctan(x)-13/24*x/(x^2+4)+25/144*arctan(1/2*x)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.14

$$\int \frac{1+x^2+x^4}{(1+x^2)(4+x^2)^2} dx = \frac{25(x^2+4)\arctan(\frac{1}{2}x) + 16(x^2+4)\arctan(x) - 78x}{144(x^2+4)}$$

input `integrate((x^4+x^2+1)/(x^2+1)/(x^2+4)^2,x, algorithm="fricas")`output `1/144*(25*(x^2 + 4)*arctan(1/2*x) + 16*(x^2 + 4)*arctan(x) - 78*x)/(x^2 + 4)`**Sympy [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.76

$$\int \frac{1+x^2+x^4}{(1+x^2)(4+x^2)^2} dx = -\frac{13x}{24x^2+96} + \frac{25 \operatorname{atan}(\frac{x}{2})}{144} + \frac{\operatorname{atan}(x)}{9}$$

input `integrate((x**4+x**2+1)/(x**2+1)/(x**2+4)**2,x)`

output `-13*x/(24*x**2 + 96) + 25*atan(x/2)/144 + atan(x)/9`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

$$\int \frac{1 + x^2 + x^4}{(1 + x^2)(4 + x^2)^2} dx = -\frac{13x}{24(x^2 + 4)} + \frac{25}{144} \arctan\left(\frac{1}{2}x\right) + \frac{1}{9} \arctan(x)$$

input `integrate((x^4+x^2+1)/(x^2+1)/(x^2+4)^2,x, algorithm="maxima")`

output `-13/24*x/(x^2 + 4) + 25/144*arctan(1/2*x) + 1/9*arctan(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

$$\int \frac{1 + x^2 + x^4}{(1 + x^2)(4 + x^2)^2} dx = -\frac{13x}{24(x^2 + 4)} + \frac{25}{144} \arctan\left(\frac{1}{2}x\right) + \frac{1}{9} \arctan(x)$$

input `integrate((x^4+x^2+1)/(x^2+1)/(x^2+4)^2,x, algorithm="giac")`

output `-13/24*x/(x^2 + 4) + 25/144*arctan(1/2*x) + 1/9*arctan(x)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.79

$$\int \frac{1 + x^2 + x^4}{(1 + x^2)(4 + x^2)^2} dx = \frac{25 \operatorname{atan}\left(\frac{x}{2}\right)}{144} + \frac{\operatorname{atan}(x)}{9} - \frac{13x}{24(x^2 + 4)}$$

input `int((x^2 + x^4 + 1)/((x^2 + 1)*(x^2 + 4)^2), x)`

output $(25*\operatorname{atan}(x/2))/144 + \operatorname{atan}(x)/9 - (13*x)/(24*(x^2 + 4))$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.38

$$\int \frac{1 + x^2 + x^4}{(1 + x^2)(4 + x^2)^2} dx$$

$$= \frac{25\operatorname{atan}\left(\frac{x}{2}\right)x^2 + 100\operatorname{atan}\left(\frac{x}{2}\right) + 16\operatorname{atan}(x)x^2 + 64\operatorname{atan}(x) - 78x}{144x^2 + 576}$$

input $\operatorname{int}((x^4+x^2+1)/(x^2+1)/(x^2+4)^2,x)$

output $(25*\operatorname{atan}(x/2)*x**2 + 100*\operatorname{atan}(x/2) + 16*\operatorname{atan}(x)*x**2 + 64*\operatorname{atan}(x) - 78*x)/(144*(x**2 + 4))$

3.57 $\int \frac{1+x^2+x^3}{2x^2+x^3+x^4} dx$

Optimal result	352
Mathematica [A] (verified)	352
Rubi [A] (verified)	353
Maple [A] (verified)	354
Fricas [A] (verification not implemented)	354
Sympy [A] (verification not implemented)	355
Maxima [A] (verification not implemented)	355
Giac [A] (verification not implemented)	355
Mupad [B] (verification not implemented)	356
Reduce [B] (verification not implemented)	356

Optimal result

Integrand size = 23, antiderivative size = 46

$$\int \frac{1+x^2+x^3}{2x^2+x^3+x^4} dx = -\frac{1}{2x} + \frac{\arctan\left(\frac{1+2x}{\sqrt{7}}\right)}{4\sqrt{7}} - \frac{\log(x)}{4} + \frac{5}{8} \log(2+x+x^2)$$

output `-1/2/x+1/28*arctan(1/7*(1+2*x)*7^(1/2))*7^(1/2)-1/4*ln(x)+5/8*ln(x^2+x+2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

$$\int \frac{1+x^2+x^3}{2x^2+x^3+x^4} dx = -\frac{1}{2x} + \frac{\arctan\left(\frac{1+2x}{\sqrt{7}}\right)}{4\sqrt{7}} - \frac{\log(x)}{4} + \frac{5}{8} \log(2+x+x^2)$$

input `Integrate[(1 + x^2 + x^3)/(2*x^2 + x^3 + x^4), x]`

output `-1/2*1/x + ArcTan[(1 + 2*x)/Sqrt[7]]/(4*Sqrt[7]) - Log[x]/4 + (5*Log[2 + x + x^2])/8`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {2026, 2159, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3 + x^2 + 1}{x^4 + x^3 + 2x^2} dx$$

↓ 2026

$$\int \frac{x^3 + x^2 + 1}{x^2(x^2 + x + 2)} dx$$

↓ 2159

$$\int \left(\frac{5x + 3}{4(x^2 + x + 2)} + \frac{1}{2x^2} - \frac{1}{4x} \right) dx$$

↓ 2009

$$\frac{\arctan\left(\frac{2x+1}{\sqrt{7}}\right)}{4\sqrt{7}} + \frac{5}{8} \log(x^2 + x + 2) - \frac{1}{2x} - \frac{\log(x)}{4}$$

input `Int[(1 + x^2 + x^3)/(2*x^2 + x^3 + x^4),x]`

output `-1/2*1/x + ArcTan[(1 + 2*x)/Sqrt[7]]/(4*Sqrt[7]) - Log[x]/4 + (5*Log[2 + x + x^2])/8`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2159

```
Int[(Pq_)*((d_.) + (e_)*(x_))^(m_)*((a_.) + (b_)*(x_) + (c_)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.74

method	result	size
risch	$-\frac{1}{2x} - \frac{\ln(x)}{4} + \frac{5 \ln(x^2+x+2)}{8} + \frac{\sqrt{7} \arctan\left(\frac{2(x+\frac{1}{2})\sqrt{7}}{7}\right)}{28}$	34
default	$-\frac{1}{2x} + \frac{\arctan\left(\frac{(1+2x)\sqrt{7}}{7}\right)\sqrt{7}}{28} - \frac{\ln(x)}{4} + \frac{5 \ln(x^2+x+2)}{8}$	36

input

```
int((x^3+x^2+1)/(x^4+x^3+2*x^2), x, method=_RETURNVERBOSE)
```

output

```
-1/2/x-1/4*ln(x)+5/8*ln(x^2+x+2)+1/28*7^(1/2)*arctan(2/7*(x+1/2)*7^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.85

$$\int \frac{1+x^2+x^3}{2x^2+x^3+x^4} dx = \frac{2\sqrt{7}x \arctan\left(\frac{1}{7}\sqrt{7}(2x+1)\right) + 35x \log(x^2+x+2) - 14x \log(x) - 28}{56x}$$

input

```
integrate((x^3+x^2+1)/(x^4+x^3+2*x^2), x, algorithm="fricas")
```

output

```
1/56*(2*sqrt(7)*x*arctan(1/7*sqrt(7)*(2*x + 1)) + 35*x*log(x^2 + x + 2) - 14*x*log(x) - 28)/x
```

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

$$\int \frac{1+x^2+x^3}{2x^2+x^3+x^4} dx = -\frac{\log(x)}{4} + \frac{5 \log(x^2+x+2)}{8} + \frac{\sqrt{7} \operatorname{atan}\left(\frac{2\sqrt{7}x}{7} + \frac{\sqrt{7}}{7}\right)}{28} - \frac{1}{2x}$$

input `integrate((x**3+x**2+1)/(x**4+x**3+2*x**2),x)`output `-log(x)/4 + 5*log(x**2 + x + 2)/8 + sqrt(7)*atan(2*sqrt(7)*x/7 + sqrt(7)/7)/28 - 1/(2*x)`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.76

$$\int \frac{1+x^2+x^3}{2x^2+x^3+x^4} dx = \frac{1}{28} \sqrt{7} \arctan\left(\frac{1}{7} \sqrt{7}(2x+1)\right) - \frac{1}{2x} + \frac{5}{8} \log(x^2+x+2) - \frac{1}{4} \log(x)$$

input `integrate((x^3+x^2+1)/(x^4+x^3+2*x^2),x, algorithm="maxima")`output `1/28*sqrt(7)*arctan(1/7*sqrt(7)*(2*x + 1)) - 1/2/x + 5/8*log(x^2 + x + 2) - 1/4*log(x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.78

$$\int \frac{1+x^2+x^3}{2x^2+x^3+x^4} dx = \frac{1}{28} \sqrt{7} \arctan\left(\frac{1}{7} \sqrt{7}(2x+1)\right) - \frac{1}{2x} + \frac{5}{8} \log(x^2+x+2) - \frac{1}{4} \log(|x|)$$

input `integrate((x^3+x^2+1)/(x^4+x^3+2*x^2),x, algorithm="giac")`

output `1/28*sqrt(7)*arctan(1/7*sqrt(7)*(2*x + 1)) - 1/2/x + 5/8*log(x^2 + x + 2) - 1/4*log(abs(x))`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.07

$$\int \frac{1+x^2+x^3}{2x^2+x^3+x^4} dx = -\frac{\ln(x)}{4} - \ln\left(x + \frac{1}{2} - \frac{\sqrt{7} \operatorname{li}}{2}\right) \left(-\frac{5}{8} + \frac{\sqrt{7} \operatorname{li}}{56}\right) + \ln\left(x + \frac{1}{2} + \frac{\sqrt{7} \operatorname{li}}{2}\right) \left(\frac{5}{8} + \frac{\sqrt{7} \operatorname{li}}{56}\right) - \frac{1}{2x}$$

input `int((x^2 + x^3 + 1)/(2*x^2 + x^3 + x^4),x)`

output `log(x + (7^(1/2)*1i)/2 + 1/2)*((7^(1/2)*1i)/56 + 5/8) - log(x - (7^(1/2)*1i)/2 + 1/2)*((7^(1/2)*1i)/56 - 5/8) - log(x)/4 - 1/(2*x)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.83

$$\int \frac{1+x^2+x^3}{2x^2+x^3+x^4} dx = \frac{2\sqrt{7} \operatorname{atan}\left(\frac{2x+1}{\sqrt{7}}\right) x + 35 \log(x^2+x+2) x - 14 \log(x) x - 28}{56x}$$

input `int((x^3+x^2+1)/(x^4+x^3+2*x^2),x)`

output `(2*sqrt(7)*atan((2*x + 1)/sqrt(7))*x + 35*log(x**2 + x + 2)*x - 14*log(x)*x - 28)/(56*x)`

$$3.58 \quad \int \frac{1-12x+x^2+x^3}{-12+x+x^2} dx$$

Optimal result	357
Mathematica [A] (verified)	357
Rubi [A] (verified)	358
Maple [A] (verified)	359
Fricas [A] (verification not implemented)	359
Sympy [A] (verification not implemented)	359
Maxima [A] (verification not implemented)	360
Giac [A] (verification not implemented)	360
Mupad [B] (verification not implemented)	360
Reduce [B] (verification not implemented)	361

Optimal result

Integrand size = 20, antiderivative size = 26

$$\int \frac{1-12x+x^2+x^3}{-12+x+x^2} dx = \frac{x^2}{2} + \frac{1}{7} \log(3-x) - \frac{1}{7} \log(4+x)$$

output `1/2*x^2+1/7*ln(3-x)-1/7*ln(4+x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{1-12x+x^2+x^3}{-12+x+x^2} dx = \frac{x^2}{2} + \frac{1}{7} \log(3-x) - \frac{1}{7} \log(4+x)$$

input `Integrate[(1 - 12*x + x^2 + x^3)/(-12 + x + x^2),x]`

output `x^2/2 + Log[3 - x]/7 - Log[4 + x]/7`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2188, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3 + x^2 - 12x + 1}{x^2 + x - 12} dx$$

$$\downarrow \text{2188}$$

$$\int \left(\frac{1}{x^2 + x - 12} + x \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{x^2}{2} + \frac{1}{7} \log(3 - x) - \frac{1}{7} \log(x + 4)$$

input `Int[(1 - 12*x + x^2 + x^3)/(-12 + x + x^2),x]`

output `x^2/2 + Log[3 - x]/7 - Log[4 + x]/7`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2188 `Int[(Pq_)*((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[Pq*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.73

method	result	size
default	$\frac{x^2}{2} + \frac{\ln(-3+x)}{7} - \frac{\ln(x+4)}{7}$	19
norman	$\frac{x^2}{2} + \frac{\ln(-3+x)}{7} - \frac{\ln(x+4)}{7}$	19
risch	$\frac{x^2}{2} + \frac{\ln(-3+x)}{7} - \frac{\ln(x+4)}{7}$	19
parallelrisch	$\frac{x^2}{2} + \frac{\ln(-3+x)}{7} - \frac{\ln(x+4)}{7}$	19

input `int((x^3+x^2-12*x+1)/(x^2+x-12),x,method=_RETURNVERBOSE)`output `1/2*x^2+1/7*ln(-3+x)-1/7*ln(x+4)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.69

$$\int \frac{1 - 12x + x^2 + x^3}{-12 + x + x^2} dx = \frac{1}{2} x^2 - \frac{1}{7} \log(x + 4) + \frac{1}{7} \log(x - 3)$$

input `integrate((x^3+x^2-12*x+1)/(x^2+x-12),x, algorithm="fricas")`output `1/2*x^2 - 1/7*log(x + 4) + 1/7*log(x - 3)`**Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.65

$$\int \frac{1 - 12x + x^2 + x^3}{-12 + x + x^2} dx = \frac{x^2}{2} + \frac{\log(x - 3)}{7} - \frac{\log(x + 4)}{7}$$

input `integrate((x**3+x**2-12*x+1)/(x**2+x-12),x)`

output `x**2/2 + log(x - 3)/7 - log(x + 4)/7`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.69

$$\int \frac{1 - 12x + x^2 + x^3}{-12 + x + x^2} dx = \frac{1}{2} x^2 - \frac{1}{7} \log(x + 4) + \frac{1}{7} \log(x - 3)$$

input `integrate((x^3+x^2-12*x+1)/(x^2+x-12),x, algorithm="maxima")`

output `1/2*x^2 - 1/7*log(x + 4) + 1/7*log(x - 3)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.77

$$\int \frac{1 - 12x + x^2 + x^3}{-12 + x + x^2} dx = \frac{1}{2} x^2 - \frac{1}{7} \log(|x + 4|) + \frac{1}{7} \log(|x - 3|)$$

input `integrate((x^3+x^2-12*x+1)/(x^2+x-12),x, algorithm="giac")`

output `1/2*x^2 - 1/7*log(abs(x + 4)) + 1/7*log(abs(x - 3))`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.54

$$\int \frac{1 - 12x + x^2 + x^3}{-12 + x + x^2} dx = \frac{x^2}{2} - \frac{2 \operatorname{atanh}\left(\frac{2x}{7} + \frac{1}{7}\right)}{7}$$

input `int((x^2 - 12*x + x^3 + 1)/(x + x^2 - 12),x)`

output `x^2/2 - (2*atanh((2*x)/7 + 1/7))/7`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.69

$$\int \frac{1 - 12x + x^2 + x^3}{-12 + x + x^2} dx = \frac{\log(x - 3)}{7} - \frac{\log(x + 4)}{7} + \frac{x^2}{2}$$

input `int((x^3+x^2-12*x+1)/(x^2+x-12),x)`

output `(2*log(x - 3) - 2*log(x + 4) + 7*x**2)/14`

$$3.59 \quad \int \frac{-3+5x+6x^2}{-3x+2x^2+x^3} dx$$

Optimal result	362
Mathematica [A] (verified)	362
Rubi [A] (verified)	363
Maple [A] (verified)	364
Fricas [A] (verification not implemented)	364
Sympy [A] (verification not implemented)	365
Maxima [A] (verification not implemented)	365
Giac [A] (verification not implemented)	365
Mupad [B] (verification not implemented)	366
Reduce [B] (verification not implemented)	366

Optimal result

Integrand size = 25, antiderivative size = 17

$$\int \frac{-3 + 5x + 6x^2}{-3x + 2x^2 + x^3} dx = 2 \log(1 - x) + \log(x) + 3 \log(3 + x)$$

output `2*ln(1-x)+ln(x)+3*ln(3+x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{-3 + 5x + 6x^2}{-3x + 2x^2 + x^3} dx = 2 \log(1 - x) + \log(x) + 3 \log(3 + x)$$

input `Integrate[(-3 + 5*x + 6*x^2)/(-3*x + 2*x^2 + x^3),x]`

output `2*Log[1 - x] + Log[x] + 3*Log[3 + x]`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2026, 2159, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{6x^2 + 5x - 3}{x^3 + 2x^2 - 3x} dx$$

↓ 2026

$$\int \frac{6x^2 + 5x - 3}{x(x^2 + 2x - 3)} dx$$

↓ 2159

$$\int \left(\frac{1}{x} + \frac{3}{x+3} + \frac{2}{x-1} \right) dx$$

↓ 2009

$$2 \log(1-x) + \log(x) + 3 \log(x+3)$$

input `Int[(-3 + 5*x + 6*x^2)/(-3*x + 2*x^2 + x^3), x]`

output `2*Log[1 - x] + Log[x] + 3*Log[3 + x]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2159

```
Int[(Pq_)*((d_.) + (e_)*(x_))^(m_)*((a_.) + (b_)*(x_) + (c_)*(x_)^2)^(p_.), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

method	result	size
default	$3 \ln(3 + x) + \ln(x) + 2 \ln(-1 + x)$	16
norman	$3 \ln(3 + x) + \ln(x) + 2 \ln(-1 + x)$	16
risch	$3 \ln(3 + x) + \ln(x) + 2 \ln(-1 + x)$	16
parallelrisk	$3 \ln(3 + x) + \ln(x) + 2 \ln(-1 + x)$	16

input

```
int((6*x^2+5*x-3)/(x^3+2*x^2-3*x),x,method=_RETURNVERBOSE)
```

output

```
3*ln(3+x)+ln(x)+2*ln(-1+x)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{-3 + 5x + 6x^2}{-3x + 2x^2 + x^3} dx = 3 \log(x + 3) + 2 \log(x - 1) + \log(x)$$

input

```
integrate((6*x^2+5*x-3)/(x^3+2*x^2-3*x),x, algorithm="fricas")
```

output

```
3*log(x + 3) + 2*log(x - 1) + log(x)
```

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{-3 + 5x + 6x^2}{-3x + 2x^2 + x^3} dx = \log(x) + 2 \log(x - 1) + 3 \log(x + 3)$$

input `integrate((6*x**2+5*x-3)/(x**3+2*x**2-3*x),x)`output `log(x) + 2*log(x - 1) + 3*log(x + 3)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{-3 + 5x + 6x^2}{-3x + 2x^2 + x^3} dx = 3 \log(x + 3) + 2 \log(x - 1) + \log(x)$$

input `integrate((6*x^2+5*x-3)/(x^3+2*x^2-3*x),x, algorithm="maxima")`output `3*log(x + 3) + 2*log(x - 1) + log(x)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.06

$$\int \frac{-3 + 5x + 6x^2}{-3x + 2x^2 + x^3} dx = 3 \log(|x + 3|) + 2 \log(|x - 1|) + \log(|x|)$$

input `integrate((6*x^2+5*x-3)/(x^3+2*x^2-3*x),x, algorithm="giac")`output `3*log(abs(x + 3)) + 2*log(abs(x - 1)) + log(abs(x))`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{-3 + 5x + 6x^2}{-3x + 2x^2 + x^3} dx = 2 \ln(x - 1) + 3 \ln(x + 3) + \ln(x)$$

input `int((5*x + 6*x^2 - 3)/(2*x^2 - 3*x + x^3),x)`

output `2*log(x - 1) + 3*log(x + 3) + log(x)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{-3 + 5x + 6x^2}{-3x + 2x^2 + x^3} dx = 2 \log(x - 1) + 3 \log(x + 3) + \log(x)$$

input `int((6*x^2+5*x-3)/(x^3+2*x^2-3*x),x)`

output `2*log(x - 1) + 3*log(x + 3) + log(x)`

3.60

$$\int \frac{-2+3x+5x^2}{2x^2+x^3} dx$$

Optimal result	367
Mathematica [A] (verified)	367
Rubi [A] (verified)	368
Maple [A] (verified)	369
Fricas [A] (verification not implemented)	369
Sympy [A] (verification not implemented)	370
Maxima [A] (verification not implemented)	370
Giac [A] (verification not implemented)	370
Mupad [B] (verification not implemented)	371
Reduce [B] (verification not implemented)	371

Optimal result

Integrand size = 22, antiderivative size = 14

$$\int \frac{-2+3x+5x^2}{2x^2+x^3} dx = \frac{1}{x} + 2 \log(x) + 3 \log(2+x)$$

output `1/x+2*ln(x)+3*ln(2+x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{-2+3x+5x^2}{2x^2+x^3} dx = \frac{1}{x} + 2 \log(x) + 3 \log(2+x)$$

input `Integrate[(-2 + 3*x + 5*x^2)/(2*x^2 + x^3), x]`

output `x^(-1) + 2*Log[x] + 3*Log[2 + x]`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.136$, Rules used = {2026, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{5x^2 + 3x - 2}{x^3 + 2x^2} dx$$

↓ 2026

$$\int \frac{5x^2 + 3x - 2}{x^2(x + 2)} dx$$

↓ 1195

$$\int \left(-\frac{1}{x^2} + \frac{3}{x + 2} + \frac{2}{x} \right) dx$$

↓ 2009

$$\frac{1}{x} + 2 \log(x) + 3 \log(x + 2)$$

input `Int[(-2 + 3*x + 5*x^2)/(2*x^2 + x^3), x]`

output `x^(-1) + 2*Log[x] + 3*Log[2 + x]`

Defintions of rubi rules used

rule 1195 `Int[((d_.) + (e_.)*(x_))^(m_.)*((f_.) + (g_.)*(x_))^(n_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026

```
Int[(Fx_)*(Px_)^(p_), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p
*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && Integ
erQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

method	result	size
default	$\frac{1}{x} + 2 \ln(x) + 3 \ln(2 + x)$	15
norman	$\frac{1}{x} + 2 \ln(x) + 3 \ln(2 + x)$	15
risch	$\frac{1}{x} + 2 \ln(x) + 3 \ln(2 + x)$	15
parallelrisch	$\frac{2x \ln(x) + 3 \ln(2+x)x + 1}{x}$	19
meijerg	$\frac{1}{x} + 2 \ln(x) - 2 \ln(2) + 3 \ln\left(1 + \frac{x}{2}\right)$	21

input

```
int((5*x^2+3*x-2)/(x^3+2*x^2),x,method=_RETURNVERBOSE)
```

output

```
1/x+2*ln(x)+3*ln(2+x)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.29

$$\int \frac{-2 + 3x + 5x^2}{2x^2 + x^3} dx = \frac{3x \log(x + 2) + 2x \log(x) + 1}{x}$$

input

```
integrate((5*x^2+3*x-2)/(x^3+2*x^2),x, algorithm="fricas")
```

output

```
(3*x*log(x + 2) + 2*x*log(x) + 1)/x
```

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{-2 + 3x + 5x^2}{2x^2 + x^3} dx = 2 \log(x) + 3 \log(x + 2) + \frac{1}{x}$$

input `integrate((5*x**2+3*x-2)/(x**3+2*x**2),x)`output `2*log(x) + 3*log(x + 2) + 1/x`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{-2 + 3x + 5x^2}{2x^2 + x^3} dx = \frac{1}{x} + 3 \log(x + 2) + 2 \log(x)$$

input `integrate((5*x^2+3*x-2)/(x^3+2*x^2),x, algorithm="maxima")`output `1/x + 3*log(x + 2) + 2*log(x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{-2 + 3x + 5x^2}{2x^2 + x^3} dx = \frac{1}{x} + 3 \log(|x + 2|) + 2 \log(|x|)$$

input `integrate((5*x^2+3*x-2)/(x^3+2*x^2),x, algorithm="giac")`output `1/x + 3*log(abs(x + 2)) + 2*log(abs(x))`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{-2 + 3x + 5x^2}{2x^2 + x^3} dx = 3 \ln(x + 2) + 2 \ln(x) + \frac{1}{x}$$

input `int((3*x + 5*x^2 - 2)/(2*x^2 + x^3),x)`

output `3*log(x + 2) + 2*log(x) + 1/x`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.29

$$\int \frac{-2 + 3x + 5x^2}{2x^2 + x^3} dx = \frac{3 \log(x + 2) x + 2 \log(x) x + 1}{x}$$

input `int((5*x^2+3*x-2)/(x^3+2*x^2),x)`

output `(3*log(x + 2)*x + 2*log(x)*x + 1)/x`

3.61 $\int \frac{18-2x-4x^2}{-6+x+4x^2+x^3} dx$

Optimal result	372
Mathematica [A] (verified)	372
Rubi [A] (verified)	373
Maple [A] (verified)	374
Fricas [A] (verification not implemented)	374
Sympy [A] (verification not implemented)	374
Maxima [A] (verification not implemented)	375
Giac [A] (verification not implemented)	375
Mupad [B] (verification not implemented)	375
Reduce [B] (verification not implemented)	376

Optimal result

Integrand size = 24, antiderivative size = 19

$$\int \frac{18 - 2x - 4x^2}{-6 + x + 4x^2 + x^3} dx = \log(1 - x) - 2 \log(2 + x) - 3 \log(3 + x)$$

output `ln(1-x)-2*ln(2+x)-3*ln(3+x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.32

$$\int \frac{18 - 2x - 4x^2}{-6 + x + 4x^2 + x^3} dx = -2 \left(-\frac{1}{2} \log(1 - x) + \log(2 + x) + \frac{3}{2} \log(3 + x) \right)$$

input `Integrate[(18 - 2*x - 4*x^2)/(-6 + x + 4*x^2 + x^3),x]`

output `-2*(-1/2*Log[1 - x] + Log[2 + x] + (3*Log[3 + x])/2)`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{-4x^2 - 2x + 18}{x^3 + 4x^2 + x - 6} dx$$

$$\downarrow \text{2462}$$

$$\int \left(-\frac{2}{x+2} - \frac{3}{x+3} + \frac{1}{x-1} \right) dx$$

$$\downarrow \text{2009}$$

$$\log(1-x) - 2\log(x+2) - 3\log(x+3)$$

input

```
Int[(18 - 2*x - 4*x^2)/(-6 + x + 4*x^2 + x^3),x]
```

output

```
Log[1 - x] - 2*Log[2 + x] - 3*Log[3 + x]
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2462

```
Int[(u_.)*(Px_)^(p_), x_Symbol] :> With[{Qx = Factor[Px]}, Int[ExpandIntegr
and[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ
[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0
] && RationalFunctionQ[u, x]
```

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.95

method	result	size
default	$-3 \ln(3+x) + \ln(-1+x) - 2 \ln(2+x)$	18
norman	$-3 \ln(3+x) + \ln(-1+x) - 2 \ln(2+x)$	18
risch	$-3 \ln(3+x) + \ln(-1+x) - 2 \ln(2+x)$	18
parallelrisch	$-3 \ln(3+x) + \ln(-1+x) - 2 \ln(2+x)$	18

input `int((-4*x^2-2*x+18)/(x^3+4*x^2+x-6),x,method=_RETURNVERBOSE)`

output `-3*ln(3+x)+ln(-1+x)-2*ln(2+x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{18 - 2x - 4x^2}{-6 + x + 4x^2 + x^3} dx = -3 \log(x + 3) - 2 \log(x + 2) + \log(x - 1)$$

input `integrate((-4*x^2-2*x+18)/(x^3+4*x^2+x-6),x, algorithm="fricas")`

output `-3*log(x + 3) - 2*log(x + 2) + log(x - 1)`

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{18 - 2x - 4x^2}{-6 + x + 4x^2 + x^3} dx = \log(x - 1) - 2 \log(x + 2) - 3 \log(x + 3)$$

input `integrate((-4*x**2-2*x+18)/(x**3+4*x**2+x-6),x)`

output $\log(x - 1) - 2*\log(x + 2) - 3*\log(x + 3)$

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{18 - 2x - 4x^2}{-6 + x + 4x^2 + x^3} dx = -3 \log(x + 3) - 2 \log(x + 2) + \log(x - 1)$$

input `integrate((-4*x^2-2*x+18)/(x^3+4*x^2+x-6),x, algorithm="maxima")`

output $-3*\log(x + 3) - 2*\log(x + 2) + \log(x - 1)$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.05

$$\int \frac{18 - 2x - 4x^2}{-6 + x + 4x^2 + x^3} dx = -3 \log(|x + 3|) - 2 \log(|x + 2|) + \log(|x - 1|)$$

input `integrate((-4*x^2-2*x+18)/(x^3+4*x^2+x-6),x, algorithm="giac")`

output $-3*\log(\text{abs}(x + 3)) - 2*\log(\text{abs}(x + 2)) + \log(\text{abs}(x - 1))$

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{18 - 2x - 4x^2}{-6 + x + 4x^2 + x^3} dx = \ln(x - 1) - 2 \ln(x + 2) - 3 \ln(x + 3)$$

input `int(-(2*x + 4*x^2 - 18)/(x + 4*x^2 + x^3 - 6),x)`

output $\log(x - 1) - 2*\log(x + 2) - 3*\log(x + 3)$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{18 - 2x - 4x^2}{-6 + x + 4x^2 + x^3} dx = \log(x - 1) - 3 \log(x + 3) - 2 \log(x + 2)$$

input `int((-4*x^2-2*x+18)/(x^3+4*x^2+x-6),x)`

output `log(x - 1) - 3*log(x + 3) - 2*log(x + 2)`

$$3.62 \quad \int \frac{1+x-2x^2+x^3}{4+5x^2+x^4} dx$$

Optimal result	377
Mathematica [A] (verified)	377
Rubi [A] (verified)	378
Maple [A] (verified)	379
Fricas [A] (verification not implemented)	380
Sympy [A] (verification not implemented)	380
Maxima [A] (verification not implemented)	381
Giac [A] (verification not implemented)	381
Mupad [B] (verification not implemented)	381
Reduce [B] (verification not implemented)	382

Optimal result

Integrand size = 24, antiderivative size = 23

$$\int \frac{1+x-2x^2+x^3}{4+5x^2+x^4} dx = -\frac{3}{2} \arctan\left(\frac{x}{2}\right) + \arctan(x) + \frac{1}{2} \log(4+x^2)$$

output

```
-3/2*arctan(1/2*x)+arctan(x)+1/2*ln(x^2+4)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{1+x-2x^2+x^3}{4+5x^2+x^4} dx = -\frac{3}{2} \arctan\left(\frac{x}{2}\right) + \arctan(x) + \frac{1}{2} \log(4+x^2)$$

input

```
Integrate[(1 + x - 2*x^2 + x^3)/(4 + 5*x^2 + x^4),x]
```

output

```
(-3*ArcTan[x/2])/2 + ArcTan[x] + Log[4 + x^2]/2
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$, Rules used = {2202, 1387, 240, 1480, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3 - 2x^2 + x + 1}{x^4 + 5x^2 + 4} dx \\
 & \quad \downarrow \text{2202} \\
 & \int \frac{1 - 2x^2}{x^4 + 5x^2 + 4} dx + \int \frac{x(x^2 + 1)}{x^4 + 5x^2 + 4} dx \\
 & \quad \downarrow \text{1387} \\
 & \int \frac{x}{x^2 + 4} dx + \int \frac{1 - 2x^2}{x^4 + 5x^2 + 4} dx \\
 & \quad \downarrow \text{240} \\
 & \int \frac{1 - 2x^2}{x^4 + 5x^2 + 4} dx + \frac{1}{2} \log(x^2 + 4) \\
 & \quad \downarrow \text{1480} \\
 & \int \frac{1}{x^2 + 1} dx - 3 \int \frac{1}{x^2 + 4} dx + \frac{1}{2} \log(x^2 + 4) \\
 & \quad \downarrow \text{216} \\
 & -\frac{3}{2} \arctan\left(\frac{x}{2}\right) + \arctan(x) + \frac{1}{2} \log(x^2 + 4)
 \end{aligned}$$

input `Int[(1 + x - 2*x^2 + x^3)/(4 + 5*x^2 + x^4), x]`

output `(-3*ArcTan[x/2])/2 + ArcTan[x] + Log[4 + x^2]/2`

Definitions of rubi rules used

rule 216 $\text{Int}[(a_+) + (b_+)(x_+)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*A$
 $\text{rcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /;$ $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a$
 $, 0] \ || \ \text{GtQ}[b, 0])$

rule 240 $\text{Int}[(x_+)/((a_+) + (b_+)(x_+)^2), x_Symbol] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x$
 $^2, x]]/(2*b), x] /;$ $\text{FreeQ}\{a, b, x\}$

rule 1387 $\text{Int}[(u_+)((a_+) + (c_+)(x_+)^{n2_+}) + (b_+)(x_+)^{n_+})^{p_+}((d_+) + (e_+)$
 $(x_+)^{n_+})^{q_+}, x_Symbol] \rightarrow \text{Int}[u*(d + e*x^n)^{(p+q)}*(a/d + (c/e)*x^n)^$
 $p, x] /;$ $\text{FreeQ}\{a, b, c, d, e, n, p, q, x\} \ \&\& \ \text{EqQ}[n2, 2*n] \ \&\& \ \text{EqQ}[c*d^2 -$
 $b*d*e + a*e^2, 0] \ \&\& \ (\text{IntegerQ}[p] \ || \ (\text{GtQ}[a, 0] \ \&\& \ \text{GtQ}[d, 0] \ \&\& \ \text{LtQ}[c, 0]))$

rule 1480 $\text{Int}[(d_+) + (e_+)(x_+)^2)/((a_+) + (b_+)(x_+)^2 + (c_+)(x_+)^4), x_Symbol] :$
 $> \text{With}\{q = \text{Rt}[b^2 - 4*a*c, 2]\}, \text{Simp}[(e/2 + (2*c*d - b*e)/(2*q)) \ \text{Int}[1/($
 $b/2 - q/2 + c*x^2), x], x] + \text{Simp}[(e/2 - (2*c*d - b*e)/(2*q)) \ \text{Int}[1/(b/2$
 $+ q/2 + c*x^2), x], x]] /;$ $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0]$
 $\ \&\& \ \text{NeQ}[c*d^2 - a*e^2, 0] \ \&\& \ \text{PosQ}[b^2 - 4*a*c]$

rule 2202 $\text{Int}[(Pn_+)((a_+) + (b_+)(x_+)^2 + (c_+)(x_+)^4)^{p_+}, x_Symbol] \rightarrow \text{Module}\{n$
 $= \text{Expon}[Pn, x], k\}, \text{Int}[\text{Sum}[\text{Coeff}[Pn, x, 2*k]*x^{(2*k)}, \{k, 0, n/2\}]* (a + b$
 $*x^2 + c*x^4)^p, x] + \text{Int}[x*\text{Sum}[\text{Coeff}[Pn, x, 2*k + 1]*x^{(2*k)}, \{k, 0, (n -$
 $1)/2\}]* (a + b*x^2 + c*x^4)^p, x]] /;$ $\text{FreeQ}\{a, b, c, p, x\} \ \&\& \ \text{PolyQ}[Pn, x]$
 $\ \&\& \ !\text{PolyQ}[Pn, x^2]$

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

method	result	size
default	$-\frac{3 \arctan(\frac{x}{2})}{2} + \arctan(x) + \frac{\ln(x^2+4)}{2}$	18
risch	$-\frac{3 \arctan(\frac{x}{2})}{2} + \arctan(x) + \frac{\ln(x^2+4)}{2}$	18
parallelrisc	$-\frac{i \ln(x-i)}{2} + \frac{\ln(x-2i)}{2} + \frac{3i \ln(x-2i)}{4} + \frac{i \ln(x+i)}{2} + \frac{\ln(x+2i)}{2} - \frac{3i \ln(x+2i)}{4}$	48

input `int((x^3-2*x^2+x+1)/(x^4+5*x^2+4),x,method=_RETURNVERBOSE)`

output `-3/2*arctan(1/2*x)+arctan(x)+1/2*ln(x^2+4)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{1+x-2x^2+x^3}{4+5x^2+x^4} dx = -\frac{3}{2} \arctan\left(\frac{1}{2}x\right) + \arctan(x) + \frac{1}{2} \log(x^2+4)$$

input `integrate((x^3-2*x^2+x+1)/(x^4+5*x^2+4),x, algorithm="fricas")`

output `-3/2*arctan(1/2*x) + arctan(x) + 1/2*log(x^2 + 4)`

Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{1+x-2x^2+x^3}{4+5x^2+x^4} dx = \frac{\log(x^2+4)}{2} - \frac{3 \operatorname{atan}\left(\frac{x}{2}\right)}{2} + \operatorname{atan}(x)$$

input `integrate((x**3-2*x**2+x+1)/(x**4+5*x**2+4),x)`

output `log(x**2 + 4)/2 - 3*atan(x/2)/2 + atan(x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{1+x-2x^2+x^3}{4+5x^2+x^4} dx = -\frac{3}{2} \arctan\left(\frac{1}{2}x\right) + \arctan(x) + \frac{1}{2} \log(x^2+4)$$

input `integrate((x^3-2*x^2+x+1)/(x^4+5*x^2+4),x, algorithm="maxima")`output `-3/2*arctan(1/2*x) + arctan(x) + 1/2*log(x^2 + 4)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{1+x-2x^2+x^3}{4+5x^2+x^4} dx = -\frac{3}{2} \arctan\left(\frac{1}{2}x\right) + \arctan(x) + \frac{1}{2} \log(x^2+4)$$

input `integrate((x^3-2*x^2+x+1)/(x^4+5*x^2+4),x, algorithm="giac")`output `-3/2*arctan(1/2*x) + arctan(x) + 1/2*log(x^2 + 4)`**Mupad [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.43

$$\int \frac{1+x-2x^2+x^3}{4+5x^2+x^4} dx = -\operatorname{atan}\left(\frac{1305}{4(144x-162)} + \frac{9}{8}\right) + \ln(x-2i) \left(\frac{1}{2} + \frac{3}{4}i\right) + \ln(x+2i) \left(\frac{1}{2} - \frac{3}{4}i\right)$$

input `int((x - 2*x^2 + x^3 + 1)/(5*x^2 + x^4 + 4),x)`output `log(x - 2i)*(1/2 + 3i/4) + log(x + 2i)*(1/2 - 3i/4) - atan(1305/(4*(144*x - 162))) + 9/8)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{1 + x - 2x^2 + x^3}{4 + 5x^2 + x^4} dx = -\frac{3\operatorname{atan}\left(\frac{x}{2}\right)}{2} + \operatorname{atan}(x) + \frac{\log(x^2 + 4)}{2}$$

input `int((x^3-2*x^2+x+1)/(x^4+5*x^2+4),x)`

output `(- 3*atan(x/2) + 2*atan(x) + log(x**2 + 4))/2`

3.63 $\int \frac{-32+5x-27x^2+4x^3}{-70-299x-286x^2+50x^3-13x^4+30x^5} dx$

Optimal result	383
Mathematica [A] (verified)	383
Rubi [A] (verified)	384
Maple [A] (verified)	385
Fricas [A] (verification not implemented)	385
Sympy [A] (verification not implemented)	386
Maxima [A] (verification not implemented)	386
Giac [A] (verification not implemented)	387
Mupad [B] (verification not implemented)	388
Reduce [B] (verification not implemented)	388

Optimal result

Integrand size = 43, antiderivative size = 63

$$\int \frac{-32 + 5x - 27x^2 + 4x^3}{-70 - 299x - 286x^2 + 50x^3 - 13x^4 + 30x^5} dx$$

$$= \frac{3988 \arctan\left(\frac{1+2x}{\sqrt{19}}\right)}{13685\sqrt{19}} - \frac{3146 \log(7 - 3x)}{80155} - \frac{334}{323} \log(1 + 2x)$$

$$+ \frac{4822 \log(2 + 5x)}{4879} + \frac{11049 \log(5 + x + x^2)}{260015}$$

output

```
3988/260015*arctan(1/19*(1+2*x)*19^(1/2))*19^(1/2)-3146/80155*ln(7-3*x)-334/323*ln(1+2*x)+4822/4879*ln(2+5*x)+11049/260015*ln(x^2+x+5)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.90

$$\int \frac{-32 + 5x - 27x^2 + 4x^3}{-70 - 299x - 286x^2 + 50x^3 - 13x^4 + 30x^5} dx$$

$$= \frac{163508\sqrt{19} \arctan\left(\frac{1+2x}{\sqrt{19}}\right) - 418418 \log(7 - 3x) - 11023670 \log(1 + 2x) + 10536070 \log(2 + 5x) + 4536070 \log(5 + x + x^2)}{10660615}$$

input `Integrate[(-32 + 5*x - 27*x^2 + 4*x^3)/(-70 - 299*x - 286*x^2 + 50*x^3 - 13*x^4 + 30*x^5),x]`

output `(163508*sqrt[19]*ArcTan[(1 + 2*x)/sqrt[19]] - 418418*Log[7 - 3*x] - 11023670*Log[1 + 2*x] + 10536070*Log[2 + 5*x] + 453009*Log[5 + x + x^2])/10660615`

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.047$, Rules used = {2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{4x^3 - 27x^2 + 5x - 32}{30x^5 - 13x^4 + 50x^3 - 286x^2 - 299x - 70} dx$$

↓ 2462

$$\int \left(\frac{22098x + 48935}{260015(x^2 + x + 5)} - \frac{668}{323(2x + 1)} - \frac{9438}{80155(3x - 7)} + \frac{24110}{4879(5x + 2)} \right) dx$$

↓ 2009

$$\frac{3988 \arctan\left(\frac{2x+1}{\sqrt{19}}\right)}{13685\sqrt{19}} + \frac{11049 \log(x^2 + x + 5)}{260015} - \frac{3146 \log(7 - 3x)}{80155} - \frac{334}{323} \log(2x + 1) + \frac{4822 \log(5x + 2)}{4879}$$

input `Int[(-32 + 5*x - 27*x^2 + 4*x^3)/(-70 - 299*x - 286*x^2 + 50*x^3 - 13*x^4 + 30*x^5),x]`

output `(3988*ArcTan[(1 + 2*x)/sqrt[19]]/(13685*sqrt[19]) - (3146*Log[7 - 3*x])/80155 - (334*Log[1 + 2*x])/323 + (4822*Log[2 + 5*x])/4879 + (11049*Log[5 + x + x^2])/260015`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2462 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegr
and[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ
[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0
] && RationalFunctionQ[u, x]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.81

method	result
default	$\frac{11049 \ln(x^2+x+5)}{260015} + \frac{3988 \arctan\left(\frac{(1+2x)\sqrt{19}}{19}\right)\sqrt{19}}{260015} + \frac{4822 \ln(2+5x)}{4879} - \frac{3146 \ln(3x-7)}{80155} - \frac{334 \ln(1+2x)}{323}$
risch	$-\frac{3146 \ln(3x-7)}{80155} + \frac{4822 \ln(2+5x)}{4879} + \frac{11049 \ln(15904144x^2+15904144x+79520720)}{260015} + \frac{3988\sqrt{19} \arctan\left(\frac{(3988x+1994)\sqrt{19}}{37886}\right)}{260015}$

input `int((4*x^3-27*x^2+5*x-32)/(30*x^5-13*x^4+50*x^3-286*x^2-299*x-70),x,method
=_RETURNVERBOSE)`

output `11049/260015*ln(x^2+x+5)+3988/260015*arctan(1/19*(1+2*x)*19^(1/2))*19^(1/2
) + 4822/4879*ln(2+5*x) - 3146/80155*ln(3*x-7) - 334/323*ln(1+2*x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.79

$$\int \frac{-32 + 5x - 27x^2 + 4x^3}{-70 - 299x - 286x^2 + 50x^3 - 13x^4 + 30x^5} dx$$

$$= \frac{3988}{260015} \sqrt{19} \arctan\left(\frac{1}{19} \sqrt{19}(2x+1)\right) + \frac{11049}{260015} \log(x^2+x+5)$$

$$+ \frac{4822}{4879} \log(5x+2) - \frac{3146}{80155} \log(3x-7) - \frac{334}{323} \log(2x+1)$$

input `integrate((4*x^3-27*x^2+5*x-32)/(30*x^5-13*x^4+50*x^3-286*x^2-299*x-70),x,
algorithm="fricas")`

output `3988/260015*sqrt(19)*arctan(1/19*sqrt(19)*(2*x + 1)) + 11049/260015*log(x^2 + x + 5) + 4822/4879*log(5*x + 2) - 3146/80155*log(3*x - 7) - 334/323*log(2*x + 1)`

Sympy [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.08

$$\int \frac{-32 + 5x - 27x^2 + 4x^3}{-70 - 299x - 286x^2 + 50x^3 - 13x^4 + 30x^5} dx$$

$$= -\frac{3146 \log\left(x - \frac{7}{3}\right)}{80155} + \frac{4822 \log\left(x + \frac{2}{5}\right)}{4879} - \frac{334 \log\left(x + \frac{1}{2}\right)}{323}$$

$$+ \frac{11049 \log(x^2 + x + 5)}{260015} + \frac{3988\sqrt{19} \operatorname{atan}\left(\frac{2\sqrt{19}x}{19} + \frac{\sqrt{19}}{19}\right)}{260015}$$

input `integrate((4*x**3-27*x**2+5*x-32)/(30*x**5-13*x**4+50*x**3-286*x**2-299*x-70),x)`

output `-3146*log(x - 7/3)/80155 + 4822*log(x + 2/5)/4879 - 334*log(x + 1/2)/323 + 11049*log(x**2 + x + 5)/260015 + 3988*sqrt(19)*atan(2*sqrt(19)*x/19 + sqrt(19)/19)/260015`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.79

$$\int \frac{-32 + 5x - 27x^2 + 4x^3}{-70 - 299x - 286x^2 + 50x^3 - 13x^4 + 30x^5} dx$$

$$= \frac{3988}{260015} \sqrt{19} \arctan\left(\frac{1}{19} \sqrt{19}(2x + 1)\right) + \frac{11049}{260015} \log(x^2 + x + 5)$$

$$+ \frac{4822}{4879} \log(5x + 2) - \frac{3146}{80155} \log(3x - 7) - \frac{334}{323} \log(2x + 1)$$

input `integrate((4*x^3-27*x^2+5*x-32)/(30*x^5-13*x^4+50*x^3-286*x^2-299*x-70),x,
algorithm="maxima")`

output `3988/260015*sqrt(19)*arctan(1/19*sqrt(19)*(2*x + 1)) + 11049/260015*log(x^2 + x + 5) + 4822/4879*log(5*x + 2) - 3146/80155*log(3*x - 7) - 334/323*log(2*x + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.84

$$\int \frac{-32 + 5x - 27x^2 + 4x^3}{-70 - 299x - 286x^2 + 50x^3 - 13x^4 + 30x^5} dx$$

$$= \frac{3988}{260015} \sqrt{19} \arctan\left(\frac{1}{19} \sqrt{19}(2x + 1)\right) + \frac{11049}{260015} \log(x^2 + x + 5)$$

$$+ \frac{4822}{4879} \log(|5x + 2|) - \frac{3146}{80155} \log(|3x - 7|) - \frac{334}{323} \log(|2x + 1|)$$

input `integrate((4*x^3-27*x^2+5*x-32)/(30*x^5-13*x^4+50*x^3-286*x^2-299*x-70),x,
algorithm="giac")`

output `3988/260015*sqrt(19)*arctan(1/19*sqrt(19)*(2*x + 1)) + 11049/260015*log(x^2 + x + 5) + 4822/4879*log(abs(5*x + 2)) - 3146/80155*log(abs(3*x - 7)) - 334/323*log(abs(2*x + 1))`

Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.92

$$\int \frac{-32 + 5x - 27x^2 + 4x^3}{-70 - 299x - 286x^2 + 50x^3 - 13x^4 + 30x^5} dx$$

$$= \frac{4822 \ln\left(x + \frac{2}{5}\right)}{4879} - \frac{334 \ln\left(x + \frac{1}{2}\right)}{323} - \frac{3146 \ln\left(x - \frac{7}{3}\right)}{80155}$$

$$- \ln\left(x + \frac{1}{2} - \frac{\sqrt{19}i}{2}\right) \left(-\frac{11049}{260015} + \frac{\sqrt{19}1994i}{260015}\right)$$

$$+ \ln\left(x + \frac{1}{2} + \frac{\sqrt{19}i}{2}\right) \left(\frac{11049}{260015} + \frac{\sqrt{19}1994i}{260015}\right)$$

input

```
int(-(5*x - 27*x^2 + 4*x^3 - 32)/(299*x + 286*x^2 - 50*x^3 + 13*x^4 - 30*x^5 + 70),x)
```

output

```
(4822*log(x + 2/5))/4879 - (334*log(x + 1/2))/323 - (3146*log(x - 7/3))/80155 - log(x - (19^(1/2)*1i)/2 + 1/2)*((19^(1/2)*1994i)/260015 - 11049/260015) + log(x + (19^(1/2)*1i)/2 + 1/2)*((19^(1/2)*1994i)/260015 + 11049/260015)
```

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.78

$$\int \frac{-32 + 5x - 27x^2 + 4x^3}{-70 - 299x - 286x^2 + 50x^3 - 13x^4 + 30x^5} dx$$

$$= \frac{3988\sqrt{19} \operatorname{atan}\left(\frac{2x+1}{\sqrt{19}}\right)}{260015} + \frac{11049 \log(x^2 + x + 5)}{260015}$$

$$+ \frac{4822 \log(5x + 2)}{4879} - \frac{3146 \log(3x - 7)}{80155} - \frac{334 \log(2x + 1)}{323}$$

input

```
int((4*x^3-27*x^2+5*x-32)/(30*x^5-13*x^4+50*x^3-286*x^2-299*x-70),x)
```

output

```
(163508*sqrt(19)*atan((2*x + 1)/sqrt(19)) + 453009*log(x**2 + x + 5) + 10536070*log(5*x + 2) - 418418*log(3*x - 7) - 11023670*log(2*x + 1))/10660615
```

3.64 $\int \frac{8-13x^2-7x^3+12x^5}{4-20x+41x^2-80x^3+116x^4-80x^5+100x^6} dx$

Optimal result	389
Mathematica [A] (verified)	389
Rubi [A] (verified)	390
Maple [A] (verified)	391
Fricas [A] (verification not implemented)	391
Sympy [A] (verification not implemented)	392
Maxima [A] (verification not implemented)	392
Giac [A] (verification not implemented)	393
Mupad [B] (verification not implemented)	393
Reduce [B] (verification not implemented)	394

Optimal result

Integrand size = 50, antiderivative size = 86

$$\int \frac{8 - 13x^2 - 7x^3 + 12x^5}{4 - 20x + 41x^2 - 80x^3 + 116x^4 - 80x^5 + 100x^6} dx$$

$$= \frac{5828}{9075(2 - 5x)} - \frac{313 + 502x}{1452(1 + 2x^2)} - \frac{251 \arctan(\sqrt{2}x)}{726\sqrt{2}}$$

$$+ \frac{272\sqrt{2} \arctan(\sqrt{2}x)}{1331} - \frac{59096 \log(2 - 5x)}{99825} + \frac{2843 \log(1 + 2x^2)}{7986}$$

output 5828/(18150-45375*x)-(313+502*x)/(2904*x^2+1452)+503/15972*arctan(x*2^(1/2))*2^(1/2)-59096/99825*ln(2-5*x)+2843/7986*ln(2*x^2+1)

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.78

$$\int \frac{8 - 13x^2 - 7x^3 + 12x^5}{4 - 20x + 41x^2 - 80x^3 + 116x^4 - 80x^5 + 100x^6} dx$$

$$= \frac{-\frac{33(2554+4675x+36458x^2)}{-2+5x-4x^2+10x^3} + 12575\sqrt{2} \arctan(\sqrt{2}x) - 236384 \log(2 - 5x) + 142150 \log(1 + 2x^2)}{399300}$$

input `Integrate[(8 - 13*x^2 - 7*x^3 + 12*x^5)/(4 - 20*x + 41*x^2 - 80*x^3 + 116*x^4 - 80*x^5 + 100*x^6),x]`

output `((-33*(2554 + 4675*x + 36458*x^2))/(-2 + 5*x - 4*x^2 + 10*x^3) + 12575*
Sqrt[2]*ArcTan[Sqrt[2]*x] - 236384*Log[2 - 5*x] + 142150*Log[1 + 2*x^2])/3993
00`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{12x^5 - 7x^3 - 13x^2 + 8}{100x^6 - 80x^5 + 116x^4 - 80x^3 + 41x^2 - 20x + 4} dx$$

↓ 2462

$$\int \left(\frac{313x - 251}{363(2x^2 + 1)^2} + \frac{2(2843x + 816)}{3993(2x^2 + 1)} - \frac{59096}{19965(5x - 2)} + \frac{5828}{1815(5x - 2)^2} \right) dx$$

↓ 2009

$$\frac{272\sqrt{2} \arctan(\sqrt{2}x)}{1331} - \frac{251 \arctan(\sqrt{2}x)}{726\sqrt{2}} - \frac{502x + 313}{1452(2x^2 + 1)} + \frac{2843 \log(2x^2 + 1)}{7986} +$$

$$\frac{5828}{9075(2 - 5x)} - \frac{59096 \log(2 - 5x)}{99825}$$

input `Int[(8 - 13*x^2 - 7*x^3 + 12*x^5)/(4 - 20*x + 41*x^2 - 80*x^3 + 116*x^4 - 80*x^5 + 100*x^6),x]`

output `5828/(9075*(2 - 5*x)) - (313 + 502*x)/(1452*(1 + 2*x^2)) - (251*ArcTan[Sqr
t[2]*x])/(726*Sqrt[2]) + (272*Sqrt[2]*ArcTan[Sqrt[2]*x])/1331 - (59096*Log
[2 - 5*x])/99825 + (2843*Log[1 + 2*x^2])/7986`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2462 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegr
and[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ
[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0
] && RationalFunctionQ[u, x]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.63

method	result	size
default	$\frac{-\frac{2761x}{4} - \frac{3443}{8}}{3993x^2 + \frac{3993}{2}} + \frac{2843 \ln(2x^2+1)}{7986} + \frac{503 \arctan(x\sqrt{2})\sqrt{2}}{15972} - \frac{5828}{9075(5x-2)} - \frac{59096 \ln(5x-2)}{99825}$	54
risch	$\frac{-\frac{18229}{60500}x^2 - \frac{17}{440}x - \frac{1277}{60500}}{x^3 - \frac{2}{5}x^2 + \frac{1}{2}x - \frac{1}{5}} - \frac{59096 \ln(5x-2)}{99825} + \frac{2843 \ln(\frac{253009}{2} + 253009x^2)}{7986} + \frac{503 \arctan(x\sqrt{2})\sqrt{2}}{15972}$	57

input `int((12*x^5-7*x^3-13*x^2+8)/(100*x^6-80*x^5+116*x^4-80*x^3+41*x^2-20*x+4),
x,method=_RETURNVERBOSE)`

output `1/3993*(-2761/4*x-3443/8)/(x^2+1/2)+2843/7986*ln(2*x^2+1)+503/15972*arctan
(x*2^(1/2))*2^(1/2)-5828/9075/(5*x-2)-59096/99825*ln(5*x-2)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.20

$$\int \frac{8 - 13x^2 - 7x^3 + 12x^5}{4 - 20x + 41x^2 - 80x^3 + 116x^4 - 80x^5 + 100x^6} dx$$

$$= \frac{12575 \sqrt{2}(10x^3 - 4x^2 + 5x - 2) \arctan(\sqrt{2}x) - 1203114x^2 + 142150(10x^3 - 4x^2 + 5x - 2) \log(2x^2)}{399300(10x^3 - 4x^2 + 5x - 2)}$$

input `integrate((12*x^5-7*x^3-13*x^2+8)/(100*x^6-80*x^5+116*x^4-80*x^3+41*x^2-20*x+4),x, algorithm="fricas")`

output `1/399300*(12575*sqrt(2)*(10*x^3 - 4*x^2 + 5*x - 2)*arctan(sqrt(2)*x) - 1203114*x^2 + 142150*(10*x^3 - 4*x^2 + 5*x - 2)*log(2*x^2 + 1) - 236384*(10*x^3 - 4*x^2 + 5*x - 2)*log(5*x - 2) - 154275*x - 84282)/(10*x^3 - 4*x^2 + 5*x - 2)`

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.76

$$\int \frac{8 - 13x^2 - 7x^3 + 12x^5}{4 - 20x + 41x^2 - 80x^3 + 116x^4 - 80x^5 + 100x^6} dx$$

$$= \frac{-36458x^2 - 4675x - 2554}{121000x^3 - 48400x^2 + 60500x - 24200} - \frac{59096 \log\left(x - \frac{2}{5}\right)}{99825}$$

$$+ \frac{2843 \log\left(x^2 + \frac{1}{2}\right)}{7986} + \frac{503\sqrt{2} \operatorname{atan}\left(\sqrt{2}x\right)}{15972}$$

input `integrate((12*x**5-7*x**3-13*x**2+8)/(100*x**6-80*x**5+116*x**4-80*x**3+41*x**2-20*x+4),x)`

output `(-36458*x**2 - 4675*x - 2554)/(121000*x**3 - 48400*x**2 + 60500*x - 24200) - 59096*log(x - 2/5)/99825 + 2843*log(x**2 + 1/2)/7986 + 503*sqrt(2)*atan(sqrt(2)*x)/15972`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.69

$$\int \frac{8 - 13x^2 - 7x^3 + 12x^5}{4 - 20x + 41x^2 - 80x^3 + 116x^4 - 80x^5 + 100x^6} dx$$

$$= \frac{503}{15972} \sqrt{2} \arctan\left(\sqrt{2}x\right) - \frac{36458x^2 + 4675x + 2554}{12100(10x^3 - 4x^2 + 5x - 2)}$$

$$+ \frac{2843}{7986} \log(2x^2 + 1) - \frac{59096}{99825} \log(5x - 2)$$

input `integrate((12*x^5-7*x^3-13*x^2+8)/(100*x^6-80*x^5+116*x^4-80*x^3+41*x^2-20*x+4),x, algorithm="maxima")`

output `503/15972*sqrt(2)*arctan(sqrt(2)*x) - 1/12100*(36458*x^2 + 4675*x + 2554)/(10*x^3 - 4*x^2 + 5*x - 2) + 2843/7986*log(2*x^2 + 1) - 59096/99825*log(5*x - 2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.69

$$\begin{aligned} & \int \frac{8 - 13x^2 - 7x^3 + 12x^5}{4 - 20x + 41x^2 - 80x^3 + 116x^4 - 80x^5 + 100x^6} dx \\ &= \frac{503}{15972} \sqrt{2} \arctan(\sqrt{2}x) - \frac{36458x^2 + 4675x + 2554}{12100(2x^2 + 1)(5x - 2)} \\ & \quad + \frac{2843}{7986} \log(2x^2 + 1) - \frac{59096}{99825} \log(|5x - 2|) \end{aligned}$$

input `integrate((12*x^5-7*x^3-13*x^2+8)/(100*x^6-80*x^5+116*x^4-80*x^3+41*x^2-20*x+4),x, algorithm="giac")`

output `503/15972*sqrt(2)*arctan(sqrt(2)*x) - 1/12100*(36458*x^2 + 4675*x + 2554)/((2*x^2 + 1)*(5*x - 2)) + 2843/7986*log(2*x^2 + 1) - 59096/99825*log(abs(5*x - 2))`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.83

$$\begin{aligned} & \int \frac{8 - 13x^2 - 7x^3 + 12x^5}{4 - 20x + 41x^2 - 80x^3 + 116x^4 - 80x^5 + 100x^6} dx \\ &= -\frac{59096 \ln\left(x - \frac{2}{5}\right)}{99825} - \frac{\frac{18229x^2}{60500} + \frac{17x}{440} + \frac{1277}{60500}}{x^3 - \frac{2x^2}{5} + \frac{x}{2} - \frac{1}{5}} \\ & \quad - \ln\left(x - \frac{\sqrt{2}1i}{2}\right) \left(-\frac{2843}{7986} + \frac{\sqrt{2}503i}{31944}\right) + \ln\left(x + \frac{\sqrt{2}1i}{2}\right) \left(\frac{2843}{7986} + \frac{\sqrt{2}503i}{31944}\right) \end{aligned}$$

input

```
int(-(13*x^2 + 7*x^3 - 12*x^5 - 8)/(41*x^2 - 20*x - 80*x^3 + 116*x^4 - 80*
x^5 + 100*x^6 + 4),x)
```

output

```
log(x + (2^(1/2)*1i)/2)*((2^(1/2)*503i)/31944 + 2843/7986) - ((17*x)/440 +
(18229*x^2)/60500 + 1277/60500)/(x/2 - (2*x^2)/5 + x^3 - 1/5) - log(x - (
2^(1/2)*1i)/2)*((2^(1/2)*503i)/31944 - 2843/7986) - (59096*log(x - 2/5))/9
9825
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.97

$$\int \frac{8 - 13x^2 - 7x^3 + 12x^5}{4 - 20x + 41x^2 - 80x^3 + 116x^4 - 80x^5 + 100x^6} dx$$

$$= \frac{251500\sqrt{2} \operatorname{atan}\left(\frac{2x}{\sqrt{2}}\right) x^3 - 100600\sqrt{2} \operatorname{atan}\left(\frac{2x}{\sqrt{2}}\right) x^2 + 125750\sqrt{2} \operatorname{atan}\left(\frac{2x}{\sqrt{2}}\right) x - 50300\sqrt{2} \operatorname{atan}\left(\frac{2x}{\sqrt{2}}\right) - 4727680 \log(5x - 2) x^3 + 1891072 \log(5x - 2) x^2 - 2363840 \log(5x - 2) x + 945536 \log(5x - 2) + 2843000 \log(2x^2 + 1) x^3 - 1137200 \log(2x^2 + 1) x^2 + 1421500 \log(2x^2 + 1) x - 568600 \log(2x^2 + 1) - 6015570 x^3 - 3316335 x + 1034550}{(798600(10x^3 - 4x^2 + 5x - 2))}$$

input

```
int((12*x^5-7*x^3-13*x^2+8)/(100*x^6-80*x^5+116*x^4-80*x^3+41*x^2-20*x+4),
x)
```

output

```
(251500*sqrt(2)*atan((2*x)/sqrt(2))*x**3 - 100600*sqrt(2)*atan((2*x)/sqrt(
2))*x**2 + 125750*sqrt(2)*atan((2*x)/sqrt(2))*x - 50300*sqrt(2)*atan((2*x)
/sqrt(2)) - 4727680*log(5*x - 2)*x**3 + 1891072*log(5*x - 2)*x**2 - 236384
0*log(5*x - 2)*x + 945536*log(5*x - 2) + 2843000*log(2*x**2 + 1)*x**3 - 11
37200*log(2*x**2 + 1)*x**2 + 1421500*log(2*x**2 + 1)*x - 568600*log(2*x**2
+ 1) - 6015570*x**3 - 3316335*x + 1034550)/(798600*(10*x**3 - 4*x**2 + 5*
x - 2))
```

3.65 $\int \frac{9+x^4}{x^2(9+x^2)} dx$

Optimal result	395
Mathematica [A] (verified)	395
Rubi [A] (verified)	396
Maple [A] (verified)	397
Fricas [A] (verification not implemented)	397
Sympy [A] (verification not implemented)	397
Maxima [A] (verification not implemented)	398
Giac [A] (verification not implemented)	398
Mupad [B] (verification not implemented)	398
Reduce [B] (verification not implemented)	399

Optimal result

Integrand size = 16, antiderivative size = 17

$$\int \frac{9+x^4}{x^2(9+x^2)} dx = -\frac{1}{x} + x - \frac{10}{3} \arctan\left(\frac{x}{3}\right)$$

output `-1/x+x-10/3*arctan(1/3*x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{9+x^4}{x^2(9+x^2)} dx = -\frac{1}{x} + x - \frac{10}{3} \arctan\left(\frac{x}{3}\right)$$

input `Integrate[(9 + x^4)/(x^2*(9 + x^2)),x]`

output `-x^(-1) + x - (10*ArcTan[x/3])/3`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {1585, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4 + 9}{x^2(x^2 + 9)} dx$$

$$\downarrow 1585$$

$$\int \left(\frac{1}{x^2} - \frac{10}{x^2 + 9} + 1 \right) dx$$

$$\downarrow 2009$$

$$-\frac{10}{3} \arctan\left(\frac{x}{3}\right) + x - \frac{1}{x}$$

input `Int[(9 + x^4)/(x^2*(9 + x^2)),x]`

output `-x^(-1) + x - (10*ArcTan[x/3])/3`

Defintions of rubi rules used

rule 1585 `Int[((f_)*(x_))^(m_)*((d_)+(e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] :> Int[ExpandIntegrand[(f*x)^m*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a, c, d, e, f, m, q}, x] && IGtQ[p, 0] && IGtQ[q, -2]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$-\frac{1}{x} + x - \frac{10 \arctan\left(\frac{x}{3}\right)}{3}$	14
meijerg	$-\frac{1}{x} + x - \frac{10 \arctan\left(\frac{x}{3}\right)}{3}$	14
risch	$-\frac{1}{x} + x - \frac{10 \arctan\left(\frac{x}{3}\right)}{3}$	14
parallelrisc	$\frac{5i \ln(x-3i)x - 5i \ln(x+3i)x + 3x^2 - 3}{3x}$	31

input `int((x^4+9)/x^2/(x^2+9),x,method=_RETURNVERBOSE)`output `-1/x+x-10/3*arctan(1/3*x)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{9 + x^4}{x^2(9 + x^2)} dx = \frac{3x^2 - 10x \arctan\left(\frac{1}{3}x\right) - 3}{3x}$$

input `integrate((x^4+9)/x^2/(x^2+9),x, algorithm="fricas")`output `1/3*(3*x^2 - 10*x*arctan(1/3*x) - 3)/x`**Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

$$\int \frac{9 + x^4}{x^2(9 + x^2)} dx = x - \frac{10 \operatorname{atan}\left(\frac{x}{3}\right)}{3} - \frac{1}{x}$$

input `integrate((x**4+9)/x**2/(x**2+9),x)`

output `x - 10*atan(x/3)/3 - 1/x`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \frac{9 + x^4}{x^2(9 + x^2)} dx = x - \frac{1}{x} - \frac{10}{3} \arctan\left(\frac{1}{3}x\right)$$

input `integrate((x^4+9)/x^2/(x^2+9),x, algorithm="maxima")`

output `x - 1/x - 10/3*arctan(1/3*x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \frac{9 + x^4}{x^2(9 + x^2)} dx = x - \frac{1}{x} - \frac{10}{3} \arctan\left(\frac{1}{3}x\right)$$

input `integrate((x^4+9)/x^2/(x^2+9),x, algorithm="giac")`

output `x - 1/x - 10/3*arctan(1/3*x)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \frac{9 + x^4}{x^2(9 + x^2)} dx = x - \frac{10 \operatorname{atan}\left(\frac{x}{3}\right)}{3} - \frac{1}{x}$$

input `int((x^4 + 9)/(x^2*(x^2 + 9)),x)`

output `x - (10*atan(x/3))/3 - 1/x`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{9 + x^4}{x^2(9 + x^2)} dx = \frac{-10 \operatorname{atan}\left(\frac{x}{3}\right) x + 3x^2 - 3}{3x}$$

input `int((x^4+9)/x^2/(x^2+9),x)`

output `(- 10*atan(x/3)*x + 3*x**2 - 3)/(3*x)`

3.66 $\int \frac{2x+x^4}{1+x^2} dx$

Optimal result	400
Mathematica [A] (verified)	400
Rubi [A] (verified)	401
Maple [A] (verified)	402
Fricas [A] (verification not implemented)	402
Sympy [A] (verification not implemented)	403
Maxima [A] (verification not implemented)	403
Giac [A] (verification not implemented)	403
Mupad [B] (verification not implemented)	404
Reduce [B] (verification not implemented)	404

Optimal result

Integrand size = 15, antiderivative size = 19

$$\int \frac{2x + x^4}{1 + x^2} dx = -x + \frac{x^3}{3} + \arctan(x) + \log(1 + x^2)$$

output `-x+1/3*x^3+arctan(x)+ln(x^2+1)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{2x + x^4}{1 + x^2} dx = -x + \frac{x^3}{3} + \arctan(x) + \log(1 + x^2)$$

input `Integrate[(2*x + x^4)/(1 + x^2),x]`

output `-x + x^3/3 + ArcTan[x] + Log[1 + x^2]`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2027, 2333, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^4 + 2x}{x^2 + 1} dx \\ & \quad \downarrow \text{2027} \\ & \int \frac{x(x^3 + 2)}{x^2 + 1} dx \\ & \quad \downarrow \text{2333} \\ & \int \left(x^2 + \frac{2x + 1}{x^2 + 1} - 1 \right) dx \\ & \quad \downarrow \text{2009} \\ & \arctan(x) + \frac{x^3}{3} + \log(x^2 + 1) - x \end{aligned}$$

input `Int[(2*x + x^4)/(1 + x^2),x]`

output `-x + x^3/3 + ArcTan[x] + Log[1 + x^2]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2027 `Int[(Fx_.)*((a_.)*(x_)^(r_.) + (b_.)*(x_)^(s_.))^(p_.), x_Symbol] := Int[x^(p*r)*(a + b*x^(s - r))^p*Fx, x] /; FreeQ[{a, b, r, s}, x] && IntegerQ[p] && PosQ[s - r] && !(EqQ[p, 1] && EqQ[u, 1])`

rule 2333

```
Int[(Pq_)*((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[
ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x]
&& PolyQ[Pq, x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.95

method	result	size
default	$-x + \frac{x^3}{3} + \arctan(x) + \ln(x^2 + 1)$	18
risch	$-x + \frac{x^3}{3} + \arctan(x) + \ln(x^2 + 1)$	18
meijerg	$-\frac{x(-5x^2+15)}{15} + \arctan(x) + \ln(x^2 + 1)$	20
parallelrisch	$\frac{x^3}{3} - x + \ln(x - i) - \frac{i \ln(x-i)}{2} + \ln(x + i) + \frac{i \ln(x+i)}{2}$	36

input

```
int((x^4+2*x)/(x^2+1),x,method=_RETURNVERBOSE)
```

output

```
-x+1/3*x^3+arctan(x)+ln(x^2+1)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{2x + x^4}{1 + x^2} dx = \frac{1}{3} x^3 - x + \arctan(x) + \log(x^2 + 1)$$

input

```
integrate((x^4+2*x)/(x^2+1),x, algorithm="fricas")
```

output

```
1/3*x^3 - x + arctan(x) + log(x^2 + 1)
```

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{2x + x^4}{1 + x^2} dx = \frac{x^3}{3} - x + \log(x^2 + 1) + \operatorname{atan}(x)$$

input `integrate((x**4+2*x)/(x**2+1),x)`output `x**3/3 - x + log(x**2 + 1) + atan(x)`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{2x + x^4}{1 + x^2} dx = \frac{1}{3} x^3 - x + \arctan(x) + \log(x^2 + 1)$$

input `integrate((x^4+2*x)/(x^2+1),x, algorithm="maxima")`output `1/3*x^3 - x + arctan(x) + log(x^2 + 1)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{2x + x^4}{1 + x^2} dx = \frac{1}{3} x^3 - x + \arctan(x) + \log(x^2 + 1)$$

input `integrate((x^4+2*x)/(x^2+1),x, algorithm="giac")`output `1/3*x^3 - x + arctan(x) + log(x^2 + 1)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{2x + x^4}{1 + x^2} dx = \ln(x^2 + 1) - x + \operatorname{atan}(x) + \frac{x^3}{3}$$

input `int((2*x + x^4)/(x^2 + 1),x)`

output `log(x^2 + 1) - x + atan(x) + x^3/3`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{2x + x^4}{1 + x^2} dx = \operatorname{atan}(x) + \log(x^2 + 1) + \frac{x^3}{3} - x$$

input `int((x^4+2*x)/(x^2+1),x)`

output `(3*atan(x) + 3*log(x**2 + 1) + x**3 - 3*x)/3`

$$3.67 \quad \int \frac{-x+x^3}{(-1+x)^2(1+x^2)} dx$$

Optimal result	405
Mathematica [A] (verified)	405
Rubi [A] (verified)	406
Maple [A] (verified)	407
Fricas [A] (verification not implemented)	408
Sympy [A] (verification not implemented)	408
Maxima [A] (verification not implemented)	408
Giac [B] (verification not implemented)	409
Mupad [B] (verification not implemented)	409
Reduce [B] (verification not implemented)	409

Optimal result

Integrand size = 20, antiderivative size = 9

$$\int \frac{-x+x^3}{(-1+x)^2(1+x^2)} dx = \arctan(x) + \log(1-x)$$

output

```
arctan(x)+ln(1-x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \frac{-x+x^3}{(-1+x)^2(1+x^2)} dx = \arctan(x) + \log(1-x)$$

input

```
Integrate[(-x + x^3)/((-1 + x)^2*(1 + x^2)), x]
```

output

```
ArcTan[x] + Log[1 - x]
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2019, 2027, 2160, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3 - x}{(x - 1)^2 (x^2 + 1)} dx \\
 & \quad \downarrow \text{2019} \\
 & \int \frac{x^2 + x}{(x - 1)(x^2 + 1)} dx \\
 & \quad \downarrow \text{2027} \\
 & \int \frac{x(x + 1)}{(x - 1)(x^2 + 1)} dx \\
 & \quad \downarrow \text{2160} \\
 & \int \left(\frac{1}{x^2 + 1} + \frac{1}{x - 1} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \arctan(x) + \log(1 - x)
 \end{aligned}$$

input `Int[(-x + x^3)/((-1 + x)^2*(1 + x^2)),x]`

output `ArcTan[x] + Log[1 - x]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2019 `Int[(u_)*(Px_)^(p_)*(Qx_)^(q_), x_Symbol] := Int[u*PolynomialQuotient[Px, Qx, x]^p*Qx^(p + q), x] /; FreeQ[q, x] && PolyQ[Px, x] && PolyQ[Qx, x] && EqQ[PolynomialRemainder[Px, Qx, x], 0] && IntegerQ[p] && LtQ[p*q, 0]`

rule 2027 `Int[(Fx_)*((a_)*(x_)^(r_) + (b_)*(x_)^(s_))^(p_), x_Symbol] := Int[x^(p*r)*(a + b*x^(s - r))^p*Fx, x] /; FreeQ[{a, b, r, s}, x] && IntegerQ[p] && PosQ[s - r] && !(EqQ[p, 1] && EqQ[u, 1])`

rule 2160 `Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*Pq*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

method	result	size
default	$\arctan(x) + \ln(-1 + x)$	8
risch	$\arctan(x) + \ln(-1 + x)$	8
parallelrisch	$\frac{i \ln(x+i)}{2} - \frac{i \ln(x-i)}{2} + \ln(-1 + x)$	22

input `int((x^3-x)/(-1+x)^2/(x^2+1),x,method=_RETURNVERBOSE)`

output `arctan(x)+ln(-1+x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \frac{-x + x^3}{(-1 + x)^2 (1 + x^2)} dx = \arctan(x) + \log(x - 1)$$

input `integrate((x^3-x)/(-1+x)^2/(x^2+1),x, algorithm="fricas")`output `arctan(x) + log(x - 1)`**Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \frac{-x + x^3}{(-1 + x)^2 (1 + x^2)} dx = \log(x - 1) + \operatorname{atan}(x)$$

input `integrate((x**3-x)/(-1+x)**2/(x**2+1),x)`output `log(x - 1) + atan(x)`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \frac{-x + x^3}{(-1 + x)^2 (1 + x^2)} dx = \arctan(x) + \log(x - 1)$$

input `integrate((x^3-x)/(-1+x)^2/(x^2+1),x, algorithm="maxima")`output `arctan(x) + log(x - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 28 vs. $2(9) = 18$.

Time = 0.12 (sec) , antiderivative size = 28, normalized size of antiderivative = 3.11

$$\int \frac{-x + x^3}{(-1 + x)^2 (1 + x^2)} dx = \frac{1}{4} \pi - \pi \left[\frac{\pi + 4 \arctan(x)}{4 \pi} + \frac{1}{2} \right] + \arctan(x) + \log(|x - 1|)$$

input `integrate((x^3-x)/(-1+x)^2/(x^2+1),x, algorithm="giac")`

output `1/4*pi - pi*floor(1/4*(pi + 4*arctan(x))/pi + 1/2) + arctan(x) + log(abs(x - 1))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 2.11

$$\int \frac{-x + x^3}{(-1 + x)^2 (1 + x^2)} dx = \ln(x - 1) - \operatorname{atan}\left(\frac{5}{4x + 2} - \frac{1}{2}\right)$$

input `int(-(x - x^3)/((x^2 + 1)*(x - 1)^2),x)`

output `log(x - 1) - atan(5/(4*x + 2) - 1/2)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \frac{-x + x^3}{(-1 + x)^2 (1 + x^2)} dx = \operatorname{atan}(x) + \log(x - 1)$$

input `int((x^3-x)/(-1+x)^2/(x^2+1),x)`

output `atan(x) + log(x - 1)`

$$3.68 \quad \int \frac{2+5x+3x^2+2x^3}{1+x+x^2} dx$$

Optimal result	410
Mathematica [A] (verified)	410
Rubi [A] (verified)	411
Maple [A] (verified)	412
Fricas [A] (verification not implemented)	412
Sympy [A] (verification not implemented)	412
Maxima [A] (verification not implemented)	413
Giac [A] (verification not implemented)	413
Mupad [B] (verification not implemented)	413
Reduce [B] (verification not implemented)	414

Optimal result

Integrand size = 24, antiderivative size = 12

$$\int \frac{2 + 5x + 3x^2 + 2x^3}{1 + x + x^2} dx = x + x^2 + \log(1 + x + x^2)$$

output `x+x^2+ln(x^2+x+1)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{2 + 5x + 3x^2 + 2x^3}{1 + x + x^2} dx = x + x^2 + \log(1 + x + x^2)$$

input `Integrate[(2 + 5*x + 3*x^2 + 2*x^3)/(1 + x + x^2),x]`

output `x + x^2 + Log[1 + x + x^2]`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2188, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{2x^3 + 3x^2 + 5x + 2}{x^2 + x + 1} dx$$

↓ 2188

$$\int \left(\frac{2x + 1}{x^2 + x + 1} + 2x + 1 \right) dx$$

↓ 2009

$$x^2 + \log(x^2 + x + 1) + x$$

input

```
Int[(2 + 5*x + 3*x^2 + 2*x^3)/(1 + x + x^2), x]
```

output

```
x + x^2 + Log[1 + x + x^2]
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2188

```
Int[(Pq)*((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Int[Expand
Integrand[Pq*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c}, x] && PolyQ[Pq,
x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

method	result	size
default	$x + x^2 + \ln(x^2 + x + 1)$	13
norman	$x + x^2 + \ln(x^2 + x + 1)$	13
risch	$x + x^2 + \ln(x^2 + x + 1)$	13
parallelrisch	$x + x^2 + \ln(x^2 + x + 1)$	13

input `int((2*x^3+3*x^2+5*x+2)/(x^2+x+1),x,method=_RETURNVERBOSE)`

output `x+x^2+ln(x^2+x+1)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{2 + 5x + 3x^2 + 2x^3}{1 + x + x^2} dx = x^2 + x + \log(x^2 + x + 1)$$

input `integrate((2*x^3+3*x^2+5*x+2)/(x^2+x+1),x, algorithm="fricas")`

output `x^2 + x + log(x^2 + x + 1)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{2 + 5x + 3x^2 + 2x^3}{1 + x + x^2} dx = x^2 + x + \log(x^2 + x + 1)$$

input `integrate((2*x**3+3*x**2+5*x+2)/(x**2+x+1),x)`

output `x**2 + x + log(x**2 + x + 1)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{2 + 5x + 3x^2 + 2x^3}{1 + x + x^2} dx = x^2 + x + \log(x^2 + x + 1)$$

input `integrate((2*x^3+3*x^2+5*x+2)/(x^2+x+1),x, algorithm="maxima")`

output `x^2 + x + log(x^2 + x + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{2 + 5x + 3x^2 + 2x^3}{1 + x + x^2} dx = x^2 + x + \log(x^2 + x + 1)$$

input `integrate((2*x^3+3*x^2+5*x+2)/(x^2+x+1),x, algorithm="giac")`

output `x^2 + x + log(x^2 + x + 1)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{2 + 5x + 3x^2 + 2x^3}{1 + x + x^2} dx = x + \ln(x^2 + x + 1) + x^2$$

input `int((5*x + 3*x^2 + 2*x^3 + 2)/(x + x^2 + 1),x)`

output `x + log(x + x^2 + 1) + x^2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{2 + 5x + 3x^2 + 2x^3}{1 + x + x^2} dx = \log(x^2 + x + 1) + x^2 + x$$

input `int((2*x^3+3*x^2+5*x+2)/(x^2+x+1),x)`

output `log(x**2 + x + 1) + x**2 + x`

$$3.69 \quad \int \frac{3-4x-5x^2+3x^3}{x^3(-1+x+x^2)} dx$$

Optimal result	415
Mathematica [A] (verified)	415
Rubi [A] (verified)	416
Maple [A] (verified)	417
Fricas [A] (verification not implemented)	417
Sympy [A] (verification not implemented)	418
Maxima [A] (verification not implemented)	418
Giac [A] (verification not implemented)	419
Mupad [B] (verification not implemented)	419
Reduce [B] (verification not implemented)	420

Optimal result

Integrand size = 27, antiderivative size = 65

$$\int \frac{3-4x-5x^2+3x^3}{x^3(-1+x+x^2)} dx = \frac{3}{2x^2} - \frac{1}{x} + 3 \log(x) - \frac{1}{10} (15 - \sqrt{5}) \log(1 - \sqrt{5} + 2x) - \frac{1}{10} (15 + \sqrt{5}) \log(1 + \sqrt{5} + 2x)$$

output

```
3/2/x^2-1/x+3*ln(x)-1/10*(15-5^(1/2))*ln(1-5^(1/2)+2*x)-1/10*(15+5^(1/2))*ln(1+5^(1/2)+2*x)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.89

$$\int \frac{3-4x-5x^2+3x^3}{x^3(-1+x+x^2)} dx = \frac{1}{10} \left(\frac{15}{x^2} - \frac{10}{x} + (-15 + \sqrt{5}) \log(-1 + \sqrt{5} - 2x) + 30 \log(x) - (15 + \sqrt{5}) \log(1 + \sqrt{5} + 2x) \right)$$

input

```
Integrate[(3 - 4*x - 5*x^2 + 3*x^3)/(x^3*(-1 + x + x^2)),x]
```


output

$$(15/x^2 - 10/x + (-15 + \text{Sqrt}[5])\text{Log}[-1 + \text{Sqrt}[5] - 2*x] + 30*\text{Log}[x] - (15 + \text{Sqrt}[5])\text{Log}[1 + \text{Sqrt}[5] + 2*x])/10$$
Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.074$, Rules used = {2159, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{3x^3 - 5x^2 - 4x + 3}{x^3(x^2 + x - 1)} dx$$

↓ 2159

$$\int \left(-\frac{3}{x^3} + \frac{-3x - 1}{x^2 + x - 1} + \frac{1}{x^2} + \frac{3}{x} \right) dx$$

↓ 2009

$$\frac{3}{2x^2} - \frac{1}{x} + 3 \log(x) - \frac{1}{10} (15 - \sqrt{5}) \log(2x - \sqrt{5} + 1) - \frac{1}{10} (15 + \sqrt{5}) \log(2x + \sqrt{5} + 1)$$

input

$$\text{Int}[(3 - 4*x - 5*x^2 + 3*x^3)/(x^3*(-1 + x + x^2)),x]$$

output

$$3/(2*x^2) - x^{(-1)} + 3*\text{Log}[x] - ((15 - \text{Sqrt}[5])*\text{Log}[1 - \text{Sqrt}[5] + 2*x])/10 - ((15 + \text{Sqrt}[5])*\text{Log}[1 + \text{Sqrt}[5] + 2*x])/10$$
Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x_Symbol] \text{ :> } \text{Simp}[\text{IntSum}[u, x], x] \text{ /; } \text{SumQ}[u]$$

rule 2159

$$\text{Int}[(Pq_)*((d_) + (e_)*(x_))^{(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^{(p_)}], x_Symbol] \text{ :> } \text{Int}[\text{ExpandIntegrand}[(d + e*x)^m*Pq*(a + b*x + c*x^2)^p, x], x] \text{ /; } \text{FreeQ}\{a, b, c, d, e, m\}, x \text{ \&\& } \text{PolyQ}[Pq, x] \text{ \&\& } \text{IGtQ}[p, -2]$$

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.63

method	result	size
default	$-\frac{3 \ln(x^2+x-1)}{2} - \frac{\sqrt{5} \operatorname{arctanh}\left(\frac{(1+2x)\sqrt{5}}{5}\right)}{5} - \frac{1}{x} + \frac{3}{2x^2} + 3 \ln(x)$	41
risch	$\frac{-x+\frac{3}{2}}{x^2} - \frac{3 \ln(1-\sqrt{5}+2x)}{2} + \frac{\ln(1-\sqrt{5}+2x)\sqrt{5}}{10} - \frac{3 \ln(1+\sqrt{5}+2x)}{2} - \frac{\ln(1+\sqrt{5}+2x)\sqrt{5}}{10} + 3 \ln(x)$	69

input `int((3*x^3-5*x^2-4*x+3)/x^3/(x^2+x-1),x,method=_RETURNVERBOSE)`

output `-3/2*ln(x^2+x-1)-1/5*5^(1/2)*arctanh(1/5*(1+2*x)*5^(1/2))-1/x+3/2/x^2+3*ln(x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.02

$$\int \frac{3 - 4x - 5x^2 + 3x^3}{x^3(-1 + x + x^2)} dx$$

$$= \frac{\sqrt{5}x^2 \log\left(\frac{2x^2 - \sqrt{5}(2x+1) + 2x+3}{x^2+x-1}\right) - 15x^2 \log(x^2 + x - 1) + 30x^2 \log(x) - 10x + 15}{10x^2}$$

input `integrate((3*x^3-5*x^2-4*x+3)/x^3/(x^2+x-1),x, algorithm="fricas")`

output `1/10*(sqrt(5)*x^2*log((2*x^2 - sqrt(5)*(2*x + 1) + 2*x + 3)/(x^2 + x - 1)) - 15*x^2*log(x^2 + x - 1) + 30*x^2*log(x) - 10*x + 15)/x^2`

Sympy [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.52

$$\int \frac{3 - 4x - 5x^2 + 3x^3}{x^3(-1 + x + x^2)} dx = 3 \log(x) + \left(-\frac{3}{2} + \frac{\sqrt{5}}{10} \right) \log \left(x - \frac{405}{202} - \frac{35\sqrt{5}}{202} + \frac{110 \left(-\frac{3}{2} + \frac{\sqrt{5}}{10} \right)^2}{101} \right) + \left(-\frac{3}{2} - \frac{\sqrt{5}}{10} \right) \log \left(x - \frac{405}{202} + \frac{35\sqrt{5}}{202} + \frac{110 \left(-\frac{3}{2} - \frac{\sqrt{5}}{10} \right)^2}{101} \right) + \frac{3 - 2x}{2x^2}$$

input `integrate((3*x**3-5*x**2-4*x+3)/x**3/(x**2+x-1),x)`output `3*log(x) + (-3/2 + sqrt(5)/10)*log(x - 405/202 - 35*sqrt(5)/202 + 110*(-3/2 + sqrt(5)/10)**2/101) + (-3/2 - sqrt(5)/10)*log(x - 405/202 + 35*sqrt(5)/202 + 110*(-3/2 - sqrt(5)/10)**2/101) + (3 - 2*x)/(2*x**2)`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.78

$$\int \frac{3 - 4x - 5x^2 + 3x^3}{x^3(-1 + x + x^2)} dx = \frac{1}{10} \sqrt{5} \log \left(\frac{2x - \sqrt{5} + 1}{2x + \sqrt{5} + 1} \right) - \frac{2x - 3}{2x^2} - \frac{3}{2} \log(x^2 + x - 1) + 3 \log(x)$$

input `integrate((3*x^3-5*x^2-4*x+3)/x^3/(x^2+x-1),x, algorithm="maxima")`

output $1/10*\sqrt{5}*\log((2*x - \sqrt{5} + 1)/(2*x + \sqrt{5} + 1)) - 1/2*(2*x - 3)/x^2 - 3/2*\log(x^2 + x - 1) + 3*\log(x)$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.85

$$\int \frac{3 - 4x - 5x^2 + 3x^3}{x^3(-1 + x + x^2)} dx = \frac{1}{10} \sqrt{5} \log \left(\frac{|2x - \sqrt{5} + 1|}{|2x + \sqrt{5} + 1|} \right) - \frac{2x - 3}{2x^2} - \frac{3}{2} \log(|x^2 + x - 1|) + 3 \log(|x|)$$

input `integrate((3*x^3-5*x^2-4*x+3)/x^3/(x^2+x-1),x, algorithm="giac")`

output $1/10*\sqrt{5}*\log(\text{abs}(2*x - \sqrt{5} + 1)/\text{abs}(2*x + \sqrt{5} + 1)) - 1/2*(2*x - 3)/x^2 - 3/2*\log(\text{abs}(x^2 + x - 1)) + 3*\log(\text{abs}(x))$

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.74

$$\int \frac{3 - 4x - 5x^2 + 3x^3}{x^3(-1 + x + x^2)} dx = 3 \ln(x) - \frac{x - \frac{3}{2}}{x^2} + \ln \left(x - \frac{\sqrt{5}}{2} + \frac{1}{2} \right) \left(\frac{\sqrt{5}}{10} - \frac{3}{2} \right) - \ln \left(x + \frac{\sqrt{5}}{2} + \frac{1}{2} \right) \left(\frac{\sqrt{5}}{10} + \frac{3}{2} \right)$$

input `int(-(4*x + 5*x^2 - 3*x^3 - 3)/(x^3*(x + x^2 - 1)),x)`

output $3*\log(x) - (x - 3/2)/x^2 + \log(x - 5^{(1/2)}/2 + 1/2)*(5^{(1/2)}/10 - 3/2) - 1*\log(x + 5^{(1/2)}/2 + 1/2)*(5^{(1/2)}/10 + 3/2)$

Reduce [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.17

$$\int \frac{3 - 4x - 5x^2 + 3x^3}{x^3(-1 + x + x^2)} dx$$

$$= \frac{\sqrt{5} \log(-\sqrt{5} + 2x + 1) x^2 - \sqrt{5} \log(\sqrt{5} + 2x + 1) x^2 - 15 \log(-\sqrt{5} + 2x + 1) x^2 - 15 \log(\sqrt{5} + 2x + 1) x^2 + 30 \log(x) x^2 - 10x + 15}{10x^2}$$

input `int((3*x^3-5*x^2-4*x+3)/x^3/(x^2+x-1),x)`

output `(sqrt(5)*log(-sqrt(5)+2*x+1)*x**2 - sqrt(5)*log(sqrt(5)+2*x+1)*x**2 - 15*log(-sqrt(5)+2*x+1)*x**2 - 15*log(sqrt(5)+2*x+1)*x**2 + 30*log(x)*x**2 - 10*x + 15)/(10*x**2)`

$$3.70 \quad \int \frac{4+8x+5x^2+2x^3}{(2+2x+x^2)^2} dx$$

Optimal result	421
Mathematica [A] (verified)	421
Rubi [A] (verified)	422
Maple [A] (verified)	424
Fricas [A] (verification not implemented)	424
Sympy [A] (verification not implemented)	425
Maxima [A] (verification not implemented)	425
Giac [A] (verification not implemented)	425
Mupad [B] (verification not implemented)	426
Reduce [B] (verification not implemented)	426

Optimal result

Integrand size = 26, antiderivative size = 28

$$\int \frac{4 + 8x + 5x^2 + 2x^3}{(2 + 2x + x^2)^2} dx = -\frac{1}{2 + 2x + x^2} - \arctan(1 + x) + \log(2 + 2x + x^2)$$

output `-1/(x^2+2*x+2)-arctan(1+x)+ln(x^2+2*x+2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{4 + 8x + 5x^2 + 2x^3}{(2 + 2x + x^2)^2} dx = -\frac{1}{2 + 2x + x^2} - \arctan(1 + x) + \log(2 + 2x + x^2)$$

input `Integrate[(4 + 8*x + 5*x^2 + 2*x^3)/(2 + 2*x + x^2)^2,x]`

output `-(2 + 2*x + x^2)^(-1) - ArcTan[1 + x] + Log[2 + 2*x + x^2]`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.269$, Rules used = {2191, 27, 1142, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{2x^3 + 5x^2 + 8x + 4}{(x^2 + 2x + 2)^2} dx \\
 & \quad \downarrow \text{2191} \\
 & \frac{1}{4} \int \frac{4(2x + 1)}{x^2 + 2x + 2} dx - \frac{1}{x^2 + 2x + 2} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{2x + 1}{x^2 + 2x + 2} dx - \frac{1}{x^2 + 2x + 2} \\
 & \quad \downarrow \text{1142} \\
 & - \int \frac{1}{x^2 + 2x + 2} dx + \int \frac{2(x + 1)}{x^2 + 2x + 2} dx - \frac{1}{x^2 + 2x + 2} \\
 & \quad \downarrow \text{27} \\
 & - \int \frac{1}{x^2 + 2x + 2} dx + 2 \int \frac{x + 1}{x^2 + 2x + 2} dx - \frac{1}{x^2 + 2x + 2} \\
 & \quad \downarrow \text{1082} \\
 & 2 \int \frac{x + 1}{x^2 + 2x + 2} dx + \int \frac{1}{-(x + 1)^2 - 1} d(x + 1) - \frac{1}{x^2 + 2x + 2} \\
 & \quad \downarrow \text{217} \\
 & 2 \int \frac{x + 1}{x^2 + 2x + 2} dx - \arctan(x + 1) - \frac{1}{x^2 + 2x + 2} \\
 & \quad \downarrow \text{1103} \\
 & - \arctan(x + 1) - \frac{1}{x^2 + 2x + 2} + \log(x^2 + 2x + 2)
 \end{aligned}$$

input

```
Int[(4 + 8*x + 5*x^2 + 2*x^3)/(2 + 2*x + x^2)^2,x]
```

output $-(2 + 2x + x^2)^{-1} - \text{ArcTan}[1 + x] + \text{Log}[2 + 2x + x^2]$

Defintions of rubi rules used

rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$

rule 217 $\text{Int}[(a_*) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 1082 $\text{Int}[(a_*) + (b_*)(x_) + (c_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1103 $\text{Int}[(d_*) + (e_*)(x_)]/((a_*) + (b_*)(x_) + (c_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$

rule 1142 $\text{Int}[(d_*) + (e_*)(x_)]/((a_*) + (b_*)(x_) + (c_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \ \text{Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \ \text{Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

rule 2191 $\text{Int}[(Pq_*)*((a_*) + (b_*)(x_) + (c_*)(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{With}[\{Q = \text{PolynomialQuotient}[Pq, a + b*x + c*x^2, x], f = \text{Coeff}[\text{PolynomialRemainder}[Pq, a + b*x + c*x^2, x], x, 0], g = \text{Coeff}[\text{PolynomialRemainder}[Pq, a + b*x + c*x^2, x], x, 1]\}, \text{Simp}[(b*f - 2*a*g + (2*c*f - b*g)*x)*((a + b*x + c*x^2)^{(p + 1})/((p + 1)*(b^2 - 4*a*c))), x] + \text{Simp}[1/((p + 1)*(b^2 - 4*a*c)) \ \text{Int}[(a + b*x + c*x^2)^{(p + 1)}*\text{ExpandToSum}[(p + 1)*(b^2 - 4*a*c)*Q - (2*p + 3)*(2*c*f - b*g), x], x], x]] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{PolyQ}[Pq, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{LtQ}[p, -1]$

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

method	result
default	$-\frac{1}{x^2+2x+2} - \arctan(1+x) + \ln(x^2+2x+2)$
risch	$-\frac{1}{x^2+2x+2} - \arctan(1+x) + \ln(x^2+2x+2)$
parallelrisc	$\frac{-2i \ln(x+1+i)+2i \ln(x+1-i)+2i \ln(x+1-i)x+i \ln(x+1-i)x^2+2 \ln(x+1-i)x^2+2 \ln(x+1+i)x^2-2-2i \ln(x+1+i)x-i \ln(x+1+i)x^2}{2x^2+4x+4}$

input `int((2*x^3+5*x^2+8*x+4)/(x^2+2*x+2)^2,x,method=_RETURNVERBOSE)`

output `-1/(x^2+2*x+2)-arctan(1+x)+ln(x^2+2*x+2)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.64

$$\int \frac{4 + 8x + 5x^2 + 2x^3}{(2 + 2x + x^2)^2} dx$$

$$= -\frac{(x^2 + 2x + 2) \arctan(x + 1) - (x^2 + 2x + 2) \log(x^2 + 2x + 2) + 1}{x^2 + 2x + 2}$$

input `integrate((2*x^3+5*x^2+8*x+4)/(x^2+2*x+2)^2,x, algorithm="fricas")`

output `-((x^2 + 2*x + 2)*arctan(x + 1) - (x^2 + 2*x + 2)*log(x^2 + 2*x + 2) + 1)/(x^2 + 2*x + 2)`

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.86

$$\int \frac{4 + 8x + 5x^2 + 2x^3}{(2 + 2x + x^2)^2} dx = \log(x^2 + 2x + 2) - \operatorname{atan}(x + 1) - \frac{1}{x^2 + 2x + 2}$$

input `integrate((2*x**3+5*x**2+8*x+4)/(x**2+2*x+2)**2,x)`output `log(x**2 + 2*x + 2) - atan(x + 1) - 1/(x**2 + 2*x + 2)`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{4 + 8x + 5x^2 + 2x^3}{(2 + 2x + x^2)^2} dx = -\frac{1}{x^2 + 2x + 2} - \arctan(x + 1) + \log(x^2 + 2x + 2)$$

input `integrate((2*x^3+5*x^2+8*x+4)/(x^2+2*x+2)^2,x, algorithm="maxima")`output `-1/(x^2 + 2*x + 2) - arctan(x + 1) + log(x^2 + 2*x + 2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{4 + 8x + 5x^2 + 2x^3}{(2 + 2x + x^2)^2} dx = -\frac{1}{x^2 + 2x + 2} - \arctan(x + 1) + \log(x^2 + 2x + 2)$$

input `integrate((2*x^3+5*x^2+8*x+4)/(x^2+2*x+2)^2,x, algorithm="giac")`output `-1/(x^2 + 2*x + 2) - arctan(x + 1) + log(x^2 + 2*x + 2)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{4 + 8x + 5x^2 + 2x^3}{(2 + 2x + x^2)^2} dx = \ln(x^2 + 2x + 2) - \operatorname{atan}(x + 1) - \frac{1}{x^2 + 2x + 2}$$

input `int((8*x + 5*x^2 + 2*x^3 + 4)/(2*x + x^2 + 2)^2,x)`output `log(2*x + x^2 + 2) - atan(x + 1) - 1/(2*x + x^2 + 2)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 71, normalized size of antiderivative = 2.54

$$\int \frac{4 + 8x + 5x^2 + 2x^3}{(2 + 2x + x^2)^2} dx$$

$$= \frac{-\operatorname{atan}(x + 1)x^2 - 2\operatorname{atan}(x + 1)x - 2\operatorname{atan}(x + 1) + \log(x^2 + 2x + 2)x^2 + 2\log(x^2 + 2x + 2)x + 2\log(x^2 + 2x + 2)}{x^2 + 2x + 2}$$

input `int((2*x^3+5*x^2+8*x+4)/(x^2+2*x+2)^2,x)`output `(- atan(x + 1)*x**2 - 2*atan(x + 1)*x - 2*atan(x + 1) + log(x**2 + 2*x + 2)*x**2 + 2*log(x**2 + 2*x + 2)*x + 2*log(x**2 + 2*x + 2) - 1)/(x**2 + 2*x + 2)`

3.71 $\int \frac{(-1+x)^4 x^4}{1+x^2} dx$

Optimal result	427
Mathematica [A] (verified)	427
Rubi [A] (verified)	428
Maple [A] (verified)	429
Fricas [A] (verification not implemented)	429
Sympy [A] (verification not implemented)	430
Maxima [A] (verification not implemented)	430
Giac [A] (verification not implemented)	430
Mupad [B] (verification not implemented)	431
Reduce [B] (verification not implemented)	431

Optimal result

Integrand size = 16, antiderivative size = 32

$$\int \frac{(-1+x)^4 x^4}{1+x^2} dx = 4x - \frac{4x^3}{3} + x^5 - \frac{2x^6}{3} + \frac{x^7}{7} - 4 \arctan(x)$$

output `4*x-4/3*x^3+x^5-2/3*x^6+1/7*x^7-4*arctan(x)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

$$\int \frac{(-1+x)^4 x^4}{1+x^2} dx = 4x - \frac{4x^3}{3} + x^5 - \frac{2x^6}{3} + \frac{x^7}{7} - 4 \arctan(x)$$

input `Integrate[((-1 + x)^4*x^4)/(1 + x^2), x]`

output `4*x - (4*x^3)/3 + x^5 - (2*x^6)/3 + x^7/7 - 4*ArcTan[x]`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {525, 2333, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(x-1)^4 x^4}{x^2+1} dx$$

↓ 525

$$\int \frac{x^4(-4x^3 + 5x^2 - 4x + 1)}{x^2+1} dx + \frac{x^7}{7}$$

↓ 2333

$$\int \left(-4x^5 + 5x^4 - 4x^2 - \frac{4}{x^2+1} + 4 \right) dx + \frac{x^7}{7}$$

↓ 2009

$$-4 \arctan(x) + \frac{x^7}{7} - \frac{2x^6}{3} + x^5 - \frac{4x^3}{3} + 4x$$

input `Int[((-1 + x)^4*x^4)/(1 + x^2),x]`

output `4*x - (4*x^3)/3 + x^5 - (2*x^6)/3 + x^7/7 - 4*ArcTan[x]`

Defintions of rubi rules used

rule 525 `Int[((x_)^(m_.)*((c_) + (d_.)*(x_))^(n_))/((a_) + (b_.)*(x_)^2), x_Symbol]
:> Simp[d^n*(x^(m + n - 1)/(b*(m + n - 1))), x] + Simp[1/b Int[x^m*(ExpandToSum[b*(c + d*x)^n - b*d^n*x^n - a*d^n*x^(n - 2), x]/(a + b*x^2)), x], x]
/; FreeQ[{a, b, c, d}, x] && IGtQ[n, 1] && IGtQ[m, -2] && NeQ[m + n - 1, 0]
]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2333

```
Int[(Pq_)*((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[
ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x]
&& PolyQ[Pq, x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.84

method	result
default	$4x - \frac{4x^3}{3} + x^5 - \frac{2x^6}{3} + \frac{x^7}{7} - 4 \arctan(x)$
risch	$4x - \frac{4x^3}{3} + x^5 - \frac{2x^6}{3} + \frac{x^7}{7} - 4 \arctan(x)$
parallelrisch	$\frac{x^7}{7} - \frac{2x^6}{3} + x^5 - \frac{4x^3}{3} + 4x + 2i \ln(x - i) - 2i \ln(x + i)$
meijerg	$-\frac{x(-5x^2+15)}{15} - 4 \arctan(x) + \frac{x^2(-3x^2+6)}{3} + \frac{2x(21x^4-35x^2+105)}{35} - \frac{x^2(4x^4-6x^2+12)}{6} - \frac{x(-45x^6+63x^4)}{315}$

input

```
int((-1+x)^4*x^4/(x^2+1),x,method=_RETURNVERBOSE)
```

output

```
4*x-4/3*x^3+x^5-2/3*x^6+1/7*x^7-4*arctan(x)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.81

$$\int \frac{(-1+x)^4 x^4}{1+x^2} dx = \frac{1}{7} x^7 - \frac{2}{3} x^6 + x^5 - \frac{4}{3} x^3 + 4x - 4 \arctan(x)$$

input

```
integrate((-1+x)^4*x^4/(x^2+1),x, algorithm="fricas")
```

output

```
1/7*x^7 - 2/3*x^6 + x^5 - 4/3*x^3 + 4*x - 4*arctan(x)
```

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.91

$$\int \frac{(-1+x)^4 x^4}{1+x^2} dx = \frac{x^7}{7} - \frac{2x^6}{3} + x^5 - \frac{4x^3}{3} + 4x - 4 \operatorname{atan}(x)$$

input `integrate((-1+x)**4*x**4/(x**2+1),x)`output `x**7/7 - 2*x**6/3 + x**5 - 4*x**3/3 + 4*x - 4*atan(x)`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.81

$$\int \frac{(-1+x)^4 x^4}{1+x^2} dx = \frac{1}{7} x^7 - \frac{2}{3} x^6 + x^5 - \frac{4}{3} x^3 + 4x - 4 \operatorname{arctan}(x)$$

input `integrate((-1+x)^4*x^4/(x^2+1),x, algorithm="maxima")`output `1/7*x^7 - 2/3*x^6 + x^5 - 4/3*x^3 + 4*x - 4*arctan(x)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.81

$$\int \frac{(-1+x)^4 x^4}{1+x^2} dx = \frac{1}{7} x^7 - \frac{2}{3} x^6 + x^5 - \frac{4}{3} x^3 + 4x - 4 \operatorname{arctan}(x)$$

input `integrate((-1+x)^4*x^4/(x^2+1),x, algorithm="giac")`output `1/7*x^7 - 2/3*x^6 + x^5 - 4/3*x^3 + 4*x - 4*arctan(x)`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.81

$$\int \frac{(-1+x)^4 x^4}{1+x^2} dx = 4x - 4 \operatorname{atan}(x) - \frac{4x^3}{3} + x^5 - \frac{2x^6}{3} + \frac{x^7}{7}$$

input `int((x^4*(x - 1)^4)/(x^2 + 1),x)`output `4*x - 4*atan(x) - (4*x^3)/3 + x^5 - (2*x^6)/3 + x^7/7`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.81

$$\int \frac{(-1+x)^4 x^4}{1+x^2} dx = -4 \operatorname{atan}(x) + \frac{x^7}{7} - \frac{2x^6}{3} + x^5 - \frac{4x^3}{3} + 4x$$

input `int((-1+x)^4*x^4/(x^2+1),x)`output `(- 84*atan(x) + 3*x**7 - 14*x**6 + 21*x**5 - 28*x**3 + 84*x)/21`

3.72 $\int \frac{-20x+4x^2}{9-10x^2+x^4} dx$

Optimal result	432
Mathematica [A] (verified)	432
Rubi [A] (verified)	433
Maple [A] (verified)	435
Fricas [A] (verification not implemented)	435
Sympy [A] (verification not implemented)	436
Maxima [A] (verification not implemented)	436
Giac [A] (verification not implemented)	436
Mupad [B] (verification not implemented)	437
Reduce [B] (verification not implemented)	437

Optimal result

Integrand size = 22, antiderivative size = 31

$$\int \frac{-20x + 4x^2}{9 - 10x^2 + x^4} dx = \log(1 - x) - \frac{1}{2} \log(3 - x) + \frac{3}{2} \log(1 + x) - 2 \log(3 + x)$$

output `ln(1-x)-1/2*ln(3-x)+3/2*ln(1+x)-2*ln(3+x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.26

$$\int \frac{-20x + 4x^2}{9 - 10x^2 + x^4} dx = 4 \left(\frac{1}{4} \log(1 - x) - \frac{1}{8} \log(3 - x) + \frac{3}{8} \log(1 + x) - \frac{1}{2} \log(3 + x) \right)$$

input `Integrate[(-20*x + 4*x^2)/(9 - 10*x^2 + x^4),x]`

output `4*(Log[1 - x]/4 - Log[3 - x]/8 + (3*Log[1 + x])/8 - Log[3 + x]/2)`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.52, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {2027, 2193, 27, 1432, 1081, 1450, 220, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{4x^2 - 20x}{x^4 - 10x^2 + 9} dx \\
 & \quad \downarrow \text{2027} \\
 & \int \frac{x(4x - 20)}{x^4 - 10x^2 + 9} dx \\
 & \quad \downarrow \text{2193} \\
 & \int -\frac{20x}{x^4 - 10x^2 + 9} dx + \int \frac{4x^2}{x^4 - 10x^2 + 9} dx \\
 & \quad \downarrow \text{27} \\
 & 4 \int \frac{x^2}{x^4 - 10x^2 + 9} dx - 20 \int \frac{x}{x^4 - 10x^2 + 9} dx \\
 & \quad \downarrow \text{1432} \\
 & 4 \int \frac{x^2}{x^4 - 10x^2 + 9} dx - 10 \int \frac{1}{x^4 - 10x^2 + 9} dx^2 \\
 & \quad \downarrow \text{1081} \\
 & 4 \int \frac{x^2}{x^4 - 10x^2 + 9} dx - 10 \int \left(\frac{1}{8(1-x^2)} - \frac{1}{8(9-x^2)} \right) dx^2 \\
 & \quad \downarrow \text{1450} \\
 & 4 \left(\frac{9}{8} \int \frac{1}{x^2-9} dx - \frac{1}{8} \int \frac{1}{x^2-1} dx \right) - 10 \int \left(\frac{1}{8(1-x^2)} - \frac{1}{8(9-x^2)} \right) dx^2 \\
 & \quad \downarrow \text{220} \\
 & 4 \left(\frac{\operatorname{arctanh}(x)}{8} - \frac{3}{8} \operatorname{arctanh}\left(\frac{x}{3}\right) \right) - 10 \int \left(\frac{1}{8(1-x^2)} - \frac{1}{8(9-x^2)} \right) dx^2 \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$4\left(\frac{\operatorname{arctanh}(x)}{8} - \frac{3}{8}\operatorname{arctanh}\left(\frac{x}{3}\right)\right) - 10\left(\frac{1}{8}\log(9 - x^2) - \frac{1}{8}\log(1 - x^2)\right)$$

input `Int[(-20*x + 4*x^2)/(9 - 10*x^2 + x^4),x]`

output `4*((-3*ArcTanh[x/3])/8 + ArcTanh[x]/8) - 10*(-1/8*Log[1 - x^2] + Log[9 - x^2])/8)`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 1081 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c Int[ExpandIntegrand[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x] /; FreeQ[{a, b, c}, x] && NiceSqrtQ[b^2 - 4*a*c]`

rule 1432 `Int[(x_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, p}, x]`

rule 1450 `Int[((d_.)*(x_))^(m_)/((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[(d^2/2)*(b/q + 1) Int[(d*x)^(m - 2)/(b/2 + q/2 + c*x^2), x], x] - Simp[(d^2/2)*(b/q - 1) Int[(d*x)^(m - 2)/(b/2 - q/2 + c*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - 4*a*c, 0] && GeQ[m, 2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2027 `Int[(Fx_.)*((a_.)*(x_)^(r_.) + (b_.)*(x_)^(s_.))^(p_.), x_Symbol] := Int[x^(p*r)*(a + b*x^(s - r))^p*Fx, x] /; FreeQ[{a, b, r, s}, x] && IntegerQ[p] && PosQ[s - r] && !(EqQ[p, 1] && EqQ[u, 1])`

rule 2193 `Int[(Pq_)*((d_.)*(x_)^(m_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := Module[{q = Expon[Pq, x], k}, Int[Sum[Coeff[Pq, x, 2*k]*x^(2*k), {k, 0, q/2 + 1}](d*x)^m*(a + b*x^2 + c*x^4)^p, x] + Simp[1/d Int[Sum[Coeff[Pq, x, 2*k + 1]*x^(2*k), {k, 0, (q + 1)/2}](d*x)^(m + 1)*(a + b*x^2 + c*x^4)^p, x], x]] /; FreeQ[{a, b, c, d, m, p}, x] && PolyQ[Pq, x] && !PolyQ[Pq, x^2]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.77

method	result	size
default	$-\frac{\ln(-3+x)}{2} - 2 \ln(3+x) + \ln(-1+x) + \frac{3 \ln(1+x)}{2}$	24
norman	$-\frac{\ln(-3+x)}{2} - 2 \ln(3+x) + \ln(-1+x) + \frac{3 \ln(1+x)}{2}$	24
risch	$-\frac{\ln(-3+x)}{2} - 2 \ln(3+x) + \ln(-1+x) + \frac{3 \ln(1+x)}{2}$	24
parallelrisch	$-\frac{\ln(-3+x)}{2} - 2 \ln(3+x) + \ln(-1+x) + \frac{3 \ln(1+x)}{2}$	24

input `int((4*x^2-20*x)/(x^4-10*x^2+9),x,method=_RETURNVERBOSE)`

output `-1/2*ln(-3+x)-2*ln(3+x)+ln(-1+x)+3/2*ln(1+x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

$$\int \frac{-20x + 4x^2}{9 - 10x^2 + x^4} dx = -2 \log(x + 3) + \frac{3}{2} \log(x + 1) + \log(x - 1) - \frac{1}{2} \log(x - 3)$$

input `integrate((4*x^2-20*x)/(x^4-10*x^2+9),x, algorithm="fricas")`

output $-2*\log(x + 3) + 3/2*\log(x + 1) + \log(x - 1) - 1/2*\log(x - 3)$

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int \frac{-20x + 4x^2}{9 - 10x^2 + x^4} dx = -\frac{\log(x - 3)}{2} + \log(x - 1) + \frac{3 \log(x + 1)}{2} - 2 \log(x + 3)$$

input `integrate((4*x**2-20*x)/(x**4-10*x**2+9),x)`

output $-\log(x - 3)/2 + \log(x - 1) + 3*\log(x + 1)/2 - 2*\log(x + 3)$

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

$$\int \frac{-20x + 4x^2}{9 - 10x^2 + x^4} dx = -2 \log(x + 3) + \frac{3}{2} \log(x + 1) + \log(x - 1) - \frac{1}{2} \log(x - 3)$$

input `integrate((4*x^2-20*x)/(x^4-10*x^2+9),x, algorithm="maxima")`

output $-2*\log(x + 3) + 3/2*\log(x + 1) + \log(x - 1) - 1/2*\log(x - 3)$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.87

$$\int \frac{-20x + 4x^2}{9 - 10x^2 + x^4} dx = -2 \log(|x + 3|) + \frac{3}{2} \log(|x + 1|) + \log(|x - 1|) - \frac{1}{2} \log(|x - 3|)$$

input `integrate((4*x^2-20*x)/(x^4-10*x^2+9),x, algorithm="giac")`

output $-2*\log(\text{abs}(x + 3)) + 3/2*\log(\text{abs}(x + 1)) + \log(\text{abs}(x - 1)) - 1/2*\log(\text{abs}(x - 3))$

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

$$\int \frac{-20x + 4x^2}{9 - 10x^2 + x^4} dx = \ln(x - 1) + \frac{3 \ln(x + 1)}{2} - \frac{\ln(x - 3)}{2} - 2 \ln(x + 3)$$

input $\text{int}(-(20*x - 4*x^2)/(x^4 - 10*x^2 + 9), x)$

output $\log(x - 1) + (3*\log(x + 1))/2 - \log(x - 3)/2 - 2*\log(x + 3)$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

$$\int \frac{-20x + 4x^2}{9 - 10x^2 + x^4} dx = -\frac{\log(x - 3)}{2} + \log(x - 1) - 2\log(x + 3) + \frac{3\log(x + 1)}{2}$$

input $\text{int}((4*x^2-20*x)/(x^4-10*x^2+9), x)$

output $(-\log(x - 3) + 2*\log(x - 1) - 4*\log(x + 3) + 3*\log(x + 1))/2$

3.73 $\int \frac{-1+x+4x^3}{(-1+x)x^2(1+x^2)} dx$

Optimal result	438
Mathematica [A] (verified)	438
Rubi [A] (verified)	439
Maple [A] (verified)	440
Fricas [A] (verification not implemented)	440
Sympy [A] (verification not implemented)	440
Maxima [A] (verification not implemented)	441
Giac [A] (verification not implemented)	441
Mupad [B] (verification not implemented)	441
Reduce [B] (verification not implemented)	442

Optimal result

Integrand size = 24, antiderivative size = 24

$$\int \frac{-1+x+4x^3}{(-1+x)x^2(1+x^2)} dx = -\frac{1}{x} + \arctan(x) + 2 \log(1-x) - \log(1+x^2)$$

output

```
-1/x+arctan(x)+2*ln(1-x)-ln(x^2+1)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{-1+x+4x^3}{(-1+x)x^2(1+x^2)} dx = -\frac{1}{x} + \arctan(x) + 2 \log(1-x) - \log(1+x^2)$$

input

```
Integrate[(-1 + x + 4*x^3)/((-1 + x)*x^2*(1 + x^2)),x]
```

output

```
-x^(-1) + ArcTan[x] + 2*Log[1 - x] - Log[1 + x^2]
```

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2353, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{4x^3 + x - 1}{(x-1)x^2(x^2+1)} dx$$

↓ 2353

$$\int \left(\frac{1-2x}{x^2+1} + \frac{1}{x^2} + \frac{2}{x-1} \right) dx$$

↓ 2009

$$\arctan(x) - \log(x^2+1) - \frac{1}{x} + 2\log(1-x)$$

input

```
Int[(-1 + x + 4*x^3)/((-1 + x)*x^2*(1 + x^2)),x]
```

output

```
-x^(-1) + ArcTan[x] + 2*Log[1 - x] - Log[1 + x^2]
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2353

```
Int[(Px_)*((e_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(e*x)^m*(c + d*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && PolyQ[Px, x] && (IntegerQ[p] || (IntegerQ[2*p] && IntegerQ[m] && ILtQ[n, 0]))
```


Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.96

method	result	size
default	$-\frac{1}{x} - \ln(x^2 + 1) + \arctan(x) + 2 \ln(-1 + x)$	23
risch	$-\frac{1}{x} - \ln(x^2 + 1) + \arctan(x) + 2 \ln(-1 + x)$	23
parallelrisc	$\frac{-i \ln(x-i)x+i \ln(x+i)x+4 \ln(-1+x)x-2 \ln(x-i)x-2 \ln(x+i)x-2}{2x}$	49

input `int((4*x^3+x-1)/(-1+x)/x^2/(x^2+1),x,method=_RETURNVERBOSE)`

output `-1/x-ln(x^2+1)+arctan(x)+2*ln(-1+x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{-1 + x + 4x^3}{(-1 + x)x^2(1 + x^2)} dx = \frac{x \arctan(x) - x \log(x^2 + 1) + 2x \log(x - 1) - 1}{x}$$

input `integrate((4*x^3+x-1)/(-1+x)/x^2/(x^2+1),x, algorithm="fricas")`

output `(x*arctan(x) - x*log(x^2 + 1) + 2*x*log(x - 1) - 1)/x`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.79

$$\int \frac{-1 + x + 4x^3}{(-1 + x)x^2(1 + x^2)} dx = 2 \log(x - 1) - \log(x^2 + 1) + \operatorname{atan}(x) - \frac{1}{x}$$

input `integrate((4*x**3+x-1)/(-1+x)/x**2/(x**2+1),x)`

output `2*log(x - 1) - log(x**2 + 1) + atan(x) - 1/x`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{-1 + x + 4x^3}{(-1 + x)x^2(1 + x^2)} dx = -\frac{1}{x} + \arctan(x) - \log(x^2 + 1) + 2 \log(x - 1)$$

input `integrate((4*x^3+x-1)/(-1+x)/x^2/(x^2+1),x, algorithm="maxima")`output `-1/x + arctan(x) - log(x^2 + 1) + 2*log(x - 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.96

$$\int \frac{-1 + x + 4x^3}{(-1 + x)x^2(1 + x^2)} dx = -\frac{1}{x} + \arctan(x) - \log(x^2 + 1) + 2 \log(|x - 1|)$$

input `integrate((4*x^3+x-1)/(-1+x)/x^2/(x^2+1),x, algorithm="giac")`output `-1/x + arctan(x) - log(x^2 + 1) + 2*log(abs(x - 1))`**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.25

$$\int \frac{-1 + x + 4x^3}{(-1 + x)x^2(1 + x^2)} dx = 2 \ln(x - 1) - \frac{1}{x} + \ln(x - i) \left(-1 - \frac{1}{2}i\right) + \ln(x + i) \left(-1 + \frac{1}{2}i\right)$$

input `int((x + 4*x^3 - 1)/(x^2*(x^2 + 1)*(x - 1)),x)`output `2*log(x - 1) - log(x - 1i)*(1 + 1i/2) - log(x + 1i)*(1 - 1i/2) - 1/x`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{-1 + x + 4x^3}{(-1 + x)x^2(1 + x^2)} dx = \frac{\operatorname{atan}(x)x - \log(x^2 + 1)x + 2\log(x - 1)x - 1}{x}$$

input `int((4*x^3+x-1)/(-1+x)/x^2/(x^2+1),x)`

output `(atan(x)*x - log(x**2 + 1)*x + 2*log(x - 1)*x - 1)/x`

$$3.74 \quad \int \frac{1-3x+2x^2-4x^3+x^4}{(1+x^2)^3} dx$$

Optimal result	443
Mathematica [A] (verified)	443
Rubi [A] (verified)	444
Maple [A] (verified)	445
Fricas [A] (verification not implemented)	446
Sympy [A] (verification not implemented)	446
Maxima [A] (verification not implemented)	446
Giac [A] (verification not implemented)	447
Mupad [B] (verification not implemented)	447
Reduce [B] (verification not implemented)	447

Optimal result

Integrand size = 26, antiderivative size = 28

$$\int \frac{1-3x+2x^2-4x^3+x^4}{(1+x^2)^3} dx = \frac{x^2}{4(1+x^2)^2} + \frac{7}{4(1+x^2)} + \arctan(x)$$

output `1/4*x^2/(x^2+1)^2+7/(4*x^2+4)+arctan(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.82

$$\int \frac{1-3x+2x^2-4x^3+x^4}{(1+x^2)^3} dx = -\frac{1}{4(1+x^2)^2} + \frac{2}{1+x^2} + \arctan(x)$$

input `Integrate[(1 - 3*x + 2*x^2 - 4*x^3 + x^4)/(1 + x^2)^3,x]`

output `-1/4*1/(1 + x^2)^2 + 2/(1 + x^2) + ArcTan[x]`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.82, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.192$, Rules used = {2345, 27, 2345, 27, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^4 - 4x^3 + 2x^2 - 3x + 1}{(x^2 + 1)^3} dx \\
 & \quad \downarrow \text{2345} \\
 & -\frac{1}{4} \int -\frac{4(x^2 - 4x + 1)}{(x^2 + 1)^2} dx - \frac{1}{4(x^2 + 1)^2} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{x^2 - 4x + 1}{(x^2 + 1)^2} dx - \frac{1}{4(x^2 + 1)^2} \\
 & \quad \downarrow \text{2345} \\
 & -\frac{1}{2} \int -\frac{2}{x^2 + 1} dx + \frac{2}{x^2 + 1} - \frac{1}{4(x^2 + 1)^2} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{1}{x^2 + 1} dx + \frac{2}{x^2 + 1} - \frac{1}{4(x^2 + 1)^2} \\
 & \quad \downarrow \text{216} \\
 & \arctan(x) + \frac{2}{x^2 + 1} - \frac{1}{4(x^2 + 1)^2}
 \end{aligned}$$

input `Int[(1 - 3*x + 2*x^2 - 4*x^3 + x^4)/(1 + x^2)^3,x]`

output `-1/4*1/(1 + x^2)^2 + 2/(1 + x^2) + ArcTan[x]`

Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 2345 `Int[(Pq_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{Q = PolynomialQuotient[Pq, a + b*x^2, x], f = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 0], g = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 1]}, Simp[(a*g - b*f*x)*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] + Simp[1/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1)*ExpandToSum[2*a*(p + 1)*Q + f*(2*p + 3), x], x] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && LtQ[p, -1]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.68

method	result	size
default	$\frac{2x^2 + \frac{7}{4}}{(x^2 + 1)^2} + \arctan(x)$	19
risch	$\frac{2x^2 + \frac{7}{4}}{(x^2 + 1)^2} + \arctan(x)$	19
parallelrisch	$-\frac{2i \ln(x-i)x^4 - 2i \ln(x+i)x^4 - 3 + 4i \ln(x-i)x^2 - 4i \ln(x+i)x^2 + 4x^4 + 2i \ln(x-i) - 2i \ln(x+i)}{4(x^2 + 1)^2}$	77
meijerg	$\frac{x(3x^2 + 5)}{8(x^2 + 1)^2} + \arctan(x) - \frac{x(25x^2 + 15)}{40(x^2 + 1)^2} - \frac{x^4}{(x^2 + 1)^2} - \frac{x(-3x^2 + 3)}{12(x^2 + 1)^2} - \frac{3x^2(x^2 + 2)}{4(x^2 + 1)^2}$	84

input `int((x^4-4*x^3+2*x^2-3*x+1)/(x^2+1)^3,x,method=_RETURNVERBOSE)`

output `(2*x^2+7/4)/(x^2+1)^2+arctan(x)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.25

$$\int \frac{1 - 3x + 2x^2 - 4x^3 + x^4}{(1 + x^2)^3} dx = \frac{8x^2 + 4(x^4 + 2x^2 + 1) \arctan(x) + 7}{4(x^4 + 2x^2 + 1)}$$

input `integrate((x^4-4*x^3+2*x^2-3*x+1)/(x^2+1)^3,x, algorithm="fricas")`

output `1/4*(8*x^2 + 4*(x^4 + 2*x^2 + 1)*arctan(x) + 7)/(x^4 + 2*x^2 + 1)`

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.71

$$\int \frac{1 - 3x + 2x^2 - 4x^3 + x^4}{(1 + x^2)^3} dx = \frac{8x^2 + 7}{4x^4 + 8x^2 + 4} + \operatorname{atan}(x)$$

input `integrate((x**4-4*x**3+2*x**2-3*x+1)/(x**2+1)**3,x)`

output `(8*x**2 + 7)/(4*x**4 + 8*x**2 + 4) + atan(x)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.86

$$\int \frac{1 - 3x + 2x^2 - 4x^3 + x^4}{(1 + x^2)^3} dx = \frac{8x^2 + 7}{4(x^4 + 2x^2 + 1)} + \arctan(x)$$

input `integrate((x^4-4*x^3+2*x^2-3*x+1)/(x^2+1)^3,x, algorithm="maxima")`

output `1/4*(8*x^2 + 7)/(x^4 + 2*x^2 + 1) + arctan(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.68

$$\int \frac{1 - 3x + 2x^2 - 4x^3 + x^4}{(1 + x^2)^3} dx = \frac{8x^2 + 7}{4(x^2 + 1)^2} + \arctan(x)$$

input `integrate((x^4-4*x^3+2*x^2-3*x+1)/(x^2+1)^3,x, algorithm="giac")`output `1/4*(8*x^2 + 7)/(x^2 + 1)^2 + arctan(x)`**Mupad [B] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.82

$$\int \frac{1 - 3x + 2x^2 - 4x^3 + x^4}{(1 + x^2)^3} dx = \operatorname{atan}(x) + \frac{2x^2 + \frac{7}{4}}{x^4 + 2x^2 + 1}$$

input `int((2*x^2 - 3*x - 4*x^3 + x^4 + 1)/(x^2 + 1)^3,x)`output `atan(x) + (2*x^2 + 7/4)/(2*x^2 + x^4 + 1)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.43

$$\int \frac{1 - 3x + 2x^2 - 4x^3 + x^4}{(1 + x^2)^3} dx = \frac{4\operatorname{atan}(x)x^4 + 8\operatorname{atan}(x)x^2 + 4\operatorname{atan}(x) - 4x^4 + 3}{4x^4 + 8x^2 + 4}$$

input `int((x^4-4*x^3+2*x^2-3*x+1)/(x^2+1)^3,x)`output `(4*atan(x)*x**4 + 8*atan(x)*x**2 + 4*atan(x) - 4*x**4 + 3)/(4*(x**4 + 2*x**2 + 1))`

$$3.75 \quad \int \frac{1-3x+2x^2-4x^3+x^4}{1+3x^2+3x^4+x^6} dx$$

Optimal result	448
Mathematica [A] (verified)	448
Rubi [A] (verified)	449
Maple [A] (verified)	450
Fricas [A] (verification not implemented)	451
Sympy [A] (verification not implemented)	451
Maxima [A] (verification not implemented)	451
Giac [A] (verification not implemented)	452
Mupad [B] (verification not implemented)	452
Reduce [B] (verification not implemented)	452

Optimal result

Integrand size = 36, antiderivative size = 23

$$\int \frac{1-3x+2x^2-4x^3+x^4}{1+3x^2+3x^4+x^6} dx = -\frac{1}{4(1+x^2)^2} + \frac{2}{1+x^2} + \arctan(x)$$

output `-1/4/(x^2+1)^2+2/(x^2+1)+arctan(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{1-3x+2x^2-4x^3+x^4}{1+3x^2+3x^4+x^6} dx = -\frac{1}{4(1+x^2)^2} + \frac{2}{1+x^2} + \arctan(x)$$

input `Integrate[(1 - 3*x + 2*x^2 - 4*x^3 + x^4)/(1 + 3*x^2 + 3*x^4 + x^6),x]`

output `-1/4*1/(1 + x^2)^2 + 2/(1 + x^2) + ArcTan[x]`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2070, 2345, 27, 2345, 27, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^4 - 4x^3 + 2x^2 - 3x + 1}{x^6 + 3x^4 + 3x^2 + 1} dx \\
 & \quad \downarrow \text{2070} \\
 & \int \frac{x^4 - 4x^3 + 2x^2 - 3x + 1}{(x^2 + 1)^3} dx \\
 & \quad \downarrow \text{2345} \\
 & -\frac{1}{4} \int -\frac{4(x^2 - 4x + 1)}{(x^2 + 1)^2} dx - \frac{1}{4(x^2 + 1)^2} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{x^2 - 4x + 1}{(x^2 + 1)^2} dx - \frac{1}{4(x^2 + 1)^2} \\
 & \quad \downarrow \text{2345} \\
 & -\frac{1}{2} \int -\frac{2}{x^2 + 1} dx + \frac{2}{x^2 + 1} - \frac{1}{4(x^2 + 1)^2} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{1}{x^2 + 1} dx + \frac{2}{x^2 + 1} - \frac{1}{4(x^2 + 1)^2} \\
 & \quad \downarrow \text{216} \\
 & \arctan(x) + \frac{2}{x^2 + 1} - \frac{1}{4(x^2 + 1)^2}
 \end{aligned}$$

input

$$\text{Int}[(1 - 3*x + 2*x^2 - 4*x^3 + x^4)/(1 + 3*x^2 + 3*x^4 + x^6), x]$$

output

$$-1/4*1/(1 + x^2)^2 + 2/(1 + x^2) + \text{ArcTan}[x]$$

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 2070 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{a = Rt[Coeff[Px, x^2, 0], Expon[Px, x^2]], b = Rt[Coeff[Px, x^2, Expon[Px, x^2]], Expon[Px, x^2]]}, Int[u*(a + b*x^2)^(Expon[Px, x^2]*p), x] /; EqQ[Px, (a + b*x^2)^Expon[Px, x^2]] /; IntegerQ[p] && PolyQ[Px, x^2] && GtQ[Expon[Px, x^2], 1] && NeQ[Coeff[Px, x^2, 0], 0]`

rule 2345 `Int[(Pq_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{Q = PolynomialQuotient[Pq, a + b*x^2, x], f = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 0], g = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 1]}, Simp[(a*g - b*f*x)*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] + Simp[1/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1)*ExpandToSum[2*a*(p + 1)*Q + f*(2*p + 3), x], x], x] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && LtQ[p, -1]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

method	result	size
default	$\frac{2x^2 + \frac{7}{4}}{(x^2 + 1)^2} + \arctan(x)$	19
risch	$\frac{2x^2 + \frac{7}{4}}{x^4 + 2x^2 + 1} + \arctan(x)$	24
parallelrisc	$-\frac{2i \ln(x-i)x^4 - 2i \ln(x+i)x^4 - 3 + 4i \ln(x-i)x^2 - 4i \ln(x+i)x^2 + 4x^4 + 2i \ln(x-i) - 2i \ln(x+i)}{4(x^4 + 2x^2 + 1)}$	82

input `int((x^4-4*x^3+2*x^2-3*x+1)/(x^6+3*x^4+3*x^2+1),x,method=_RETURNVERBOSE)`

output $(2x^2+7/4)/(x^2+1)^2+\arctan(x)$

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.52

$$\int \frac{1 - 3x + 2x^2 - 4x^3 + x^4}{1 + 3x^2 + 3x^4 + x^6} dx = \frac{8x^2 + 4(x^4 + 2x^2 + 1)\arctan(x) + 7}{4(x^4 + 2x^2 + 1)}$$

input `integrate((x^4-4*x^3+2*x^2-3*x+1)/(x^6+3*x^4+3*x^2+1),x, algorithm="fricas")`

output $1/4*(8x^2 + 4*(x^4 + 2x^2 + 1)*\arctan(x) + 7)/(x^4 + 2x^2 + 1)$

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \frac{1 - 3x + 2x^2 - 4x^3 + x^4}{1 + 3x^2 + 3x^4 + x^6} dx = \frac{8x^2 + 7}{4x^4 + 8x^2 + 4} + \operatorname{atan}(x)$$

input `integrate((x**4-4*x**3+2*x**2-3*x+1)/(x**6+3*x**4+3*x**2+1),x)`

output $(8x^2 + 7)/(4x^4 + 8x^2 + 4) + \operatorname{atan}(x)$

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.04

$$\int \frac{1 - 3x + 2x^2 - 4x^3 + x^4}{1 + 3x^2 + 3x^4 + x^6} dx = \frac{8x^2 + 7}{4(x^4 + 2x^2 + 1)} + \arctan(x)$$

input `integrate((x^4-4*x^3+2*x^2-3*x+1)/(x^6+3*x^4+3*x^2+1),x, algorithm="maxima")`

output $1/4*(8*x^2 + 7)/(x^4 + 2*x^2 + 1) + \arctan(x)$

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{1 - 3x + 2x^2 - 4x^3 + x^4}{1 + 3x^2 + 3x^4 + x^6} dx = \frac{8x^2 + 7}{4(x^2 + 1)^2} + \arctan(x)$$

input `integrate((x^4-4*x^3+2*x^2-3*x+1)/(x^6+3*x^4+3*x^2+1),x, algorithm="giac")`

output $1/4*(8*x^2 + 7)/(x^2 + 1)^2 + \arctan(x)$

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{1 - 3x + 2x^2 - 4x^3 + x^4}{1 + 3x^2 + 3x^4 + x^6} dx = \operatorname{atan}(x) + \frac{2x^2 + \frac{7}{4}}{x^4 + 2x^2 + 1}$$

input `int((2*x^2 - 3*x - 4*x^3 + x^4 + 1)/(3*x^2 + 3*x^4 + x^6 + 1),x)`

output $\operatorname{atan}(x) + (2*x^2 + 7/4)/(2*x^2 + x^4 + 1)$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.74

$$\int \frac{1 - 3x + 2x^2 - 4x^3 + x^4}{1 + 3x^2 + 3x^4 + x^6} dx = \frac{4\operatorname{atan}(x)x^4 + 8\operatorname{atan}(x)x^2 + 4\operatorname{atan}(x) - 4x^4 + 3}{4x^4 + 8x^2 + 4}$$

input `int((x^4-4*x^3+2*x^2-3*x+1)/(x^6+3*x^4+3*x^2+1),x)`

output $(4*\operatorname{atan}(x)*x^{**4} + 8*\operatorname{atan}(x)*x^{**2} + 4*\operatorname{atan}(x) - 4*x^{**4} + 3)/(4*(x^{**4} + 2*x^{**2} + 1))$

3.76

$$\int \frac{1+x+2x^2+2x^3}{x^2+x^3+x^4} dx$$

Optimal result	454
Mathematica [A] (verified)	454
Rubi [A] (verified)	455
Maple [A] (verified)	456
Fricas [A] (verification not implemented)	456
Sympy [A] (verification not implemented)	457
Maxima [A] (verification not implemented)	457
Giac [A] (verification not implemented)	457
Mupad [B] (verification not implemented)	458
Reduce [B] (verification not implemented)	458

Optimal result

Integrand size = 26, antiderivative size = 13

$$\int \frac{1+x+2x^2+2x^3}{x^2+x^3+x^4} dx = -\frac{1}{x} + \log(1+x+x^2)$$

output `-1/x+ln(x^2+x+1)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{1+x+2x^2+2x^3}{x^2+x^3+x^4} dx = -\frac{1}{x} + \log(1+x+x^2)$$

input `Integrate[(1 + x + 2*x^2 + 2*x^3)/(x^2 + x^3 + x^4),x]`

output `-x^(-1) + Log[1 + x + x^2]`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$, Rules used = {2026, 2159, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{2x^3 + 2x^2 + x + 1}{x^4 + x^3 + x^2} dx$$

$$\downarrow \text{2026}$$

$$\int \frac{2x^3 + 2x^2 + x + 1}{x^2(x^2 + x + 1)} dx$$

$$\downarrow \text{2159}$$

$$\int \left(\frac{2x + 1}{x^2 + x + 1} + \frac{1}{x^2} \right) dx$$

$$\downarrow \text{2009}$$

$$\log(x^2 + x + 1) - \frac{1}{x}$$

input `Int[(1 + x + 2*x^2 + 2*x^3)/(x^2 + x^3 + x^4), x]`

output `-x^(-1) + Log[1 + x + x^2]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2159

```
Int[(Pq_)*((d_) + (e_)*(x_)^(m_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p
_), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*Pq*(a + b*x + c*x^2)^p, x
], x] /; FreeQ[{a, b, c, d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]
```

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.08

method	result	size
default	$-\frac{1}{x} + \ln(x^2 + x + 1)$	14
norman	$-\frac{1}{x} + \ln(x^2 + x + 1)$	14
risch	$-\frac{1}{x} + \ln(x^2 + x + 1)$	14
parallelrisch	$\frac{\ln(x^2+x+1)x-1}{x}$	16

input

```
int((2*x^3+2*x^2+x+1)/(x^4+x^3+x^2),x,method=_RETURNVERBOSE)
```

output

```
-1/x+ln(x^2+x+1)
```

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{1 + x + 2x^2 + 2x^3}{x^2 + x^3 + x^4} dx = \frac{x \log(x^2 + x + 1) - 1}{x}$$

input

```
integrate((2*x^3+2*x^2+x+1)/(x^4+x^3+x^2),x, algorithm="fricas")
```

output

```
(x*log(x^2 + x + 1) - 1)/x
```

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \frac{1+x+2x^2+2x^3}{x^2+x^3+x^4} dx = \log(x^2+x+1) - \frac{1}{x}$$

input `integrate((2*x**3+2*x**2+x+1)/(x**4+x**3+x**2),x)`output `log(x**2 + x + 1) - 1/x`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{1+x+2x^2+2x^3}{x^2+x^3+x^4} dx = -\frac{1}{x} + \log(x^2+x+1)$$

input `integrate((2*x^3+2*x^2+x+1)/(x^4+x^3+x^2),x, algorithm="maxima")`output `-1/x + log(x^2 + x + 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{1+x+2x^2+2x^3}{x^2+x^3+x^4} dx = -\frac{1}{x} + \log(x^2+x+1)$$

input `integrate((2*x^3+2*x^2+x+1)/(x^4+x^3+x^2),x, algorithm="giac")`output `-1/x + log(x^2 + x + 1)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{1 + x + 2x^2 + 2x^3}{x^2 + x^3 + x^4} dx = \ln(x^2 + x + 1) - \frac{1}{x}$$

input `int((x + 2*x^2 + 2*x^3 + 1)/(x^2 + x^3 + x^4),x)`output `log(x + x^2 + 1) - 1/x`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{1 + x + 2x^2 + 2x^3}{x^2 + x^3 + x^4} dx = \frac{\log(x^2 + x + 1)x - 1}{x}$$

input `int((2*x^3+2*x^2+x+1)/(x^4+x^3+x^2),x)`output `(log(x**2 + x + 1)*x - 1)/x`

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions	459
4.2 Links to plain text integration problems used in this report for each CAS .	477

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
  ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "
  ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]===Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]===Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]===Plus || Head[expn]===Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```

```

    Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
    If[AppellFunctionQ[Head[expn]],
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
      If[Head[expn]===RootSum,
        Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
        If[Head[expn]===Integrate || Head[expn]===Int,
          Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
          9]]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

```

```

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

```

```

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

```

```

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022  add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result    := ExpnType(result);
      ExpnType_optimal   := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
#     is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```



```

# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```
    if leaf_count_result<=2*leaf_count_optimal then
      if debug then
        print("leaf_count_result<=2*leaf_count_optimal");
      fi;
      return "A"," ";
    else
      if debug then
        print("leaf_count_result>2*leaf_count_optimal");
      fi;
      return "B",cat("Leaf count of result is larger than twice the leaf count of
                    convert(leaf_count_result,string)," $ vs. $2(",
                    convert(leaf_count_optimal,string),")=",convert(2*leaf_co
      fi;
    fi;
  else #ExpnType(result) > ExpnType(optimal)
    if debug then
      print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                  convert(ExpnType_result,string)," vs. order ",
                  convert(ExpnType_optimal,string),".");
  fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```



```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file