

Computer Algebra Independent Integration Tests

Summer 2024

0-Independent-test-suites/9-Moses-Problems

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 4:56am

Contents

1	Introduction	6
1.1	Listing of CAS systems tested	7
1.2	Results	8
1.3	Time and leaf size Performance	12
1.4	Performance based on number of rules Rubi used	14
1.5	Performance based on number of steps Rubi used	15
1.6	Solved integrals histogram based on leaf size of result	16
1.7	Solved integrals histogram based on CPU time used	17
1.8	Leaf size vs. CPU time used	18
1.9	list of integrals with no known antiderivative	19
1.10	List of integrals solved by CAS but has no known antiderivative	19
1.11	list of integrals solved by CAS but failed verification	19
1.12	Timing	20
1.13	Verification	20
1.14	Important notes about some of the results	21
1.15	Current tree layout of integration tests	24
1.16	Design of the test system	25
2	detailed summary tables of results	26
2.1	List of integrals sorted by grade for each CAS	27
2.2	Detailed conclusion table per each integral for all CAS systems	32
2.3	Detailed conclusion table specific for Rubi results	61
3	Listing of integrals	65
3.1	$\int \cot^4(x) dx$	69
3.2	$\int \frac{1}{x^4(1+x^2)} dx$	74
3.3	$\int \frac{x+x^2}{\sqrt{x}} dx$	79
3.4	$\int \cos(x) dx$	84
3.5	$\int e^{x^2} x dx$	89
3.6	$\int \sec^2(x) \tan(x) dx$	94
3.7	$\int x\sqrt{1+x^2} dx$	99

3.8	$\int e^x \sin(x) dx$	104
3.9	$\int \cot(x) \csc^3(x) dx$	109
3.10	$\int \sin(e^x) dx$	114
3.11	$\int \frac{\sin(y)}{y} dy$	119
3.12	$\int (e^x + \sin(x)) dx$	124
3.13	$\int (e^{x^2} + 2e^{x^2} x^2) dx$	129
3.14	$\int (e^x + x)^2 dx$	134
3.15	$\int (2e^x + e^{2x} + x^2) dx$	139
3.16	$\int \cos(x) \sin(x) dx$	144
3.17	$\int e^{x^2} x dx$	149
3.18	$\int x\sqrt{1+x^2} dx$	154
3.19	$\int \frac{e^x}{1+e^x} dx$	159
3.20	$\int x^{3/2} dx$	164
3.21	$\int \cos(3+2x) dx$	169
3.22	$\int 2e^{2x} yz dx$	174
3.23	$\int e^x \cos^2(e^x) \sin(e^x) dx$	179
3.24	$\int x\sqrt{1+x} dx$	184
3.25	$\int \frac{1}{-1+x^4} dx$	189
3.26	$\int \frac{e^x}{2+3e^{2x}} dx$	194
3.27	$\int \frac{e^{2x}}{A+Be^{4x}} dx$	199
3.28	$\int \frac{e^{1+x}}{1+e^x} dx$	204
3.29	$\int (10e)^x dx$	209
3.30	$\int x^3 \sin(x^2) dx$	214
3.31	$\int \frac{x^7}{1+x^{12}} dx$	220
3.32	$\int x^{3a} \sin(x^{2a}) dx$	229
3.33	$\int \cos(\sqrt{x}) dx$	234
3.34	$\int x\sqrt{1+x} dx$	239
3.35	$\int \frac{1}{\sqrt[3]{x+\sqrt{x}}} dx$	244
3.36	$\int \sqrt{\frac{1+x}{3+2x}} dx$	249
3.37	$\int \frac{x^4}{(1-x^2)^{5/2}} dx$	255
3.38	$\int \sqrt{x}(1+x)^{5/2} dx$	261
3.39	$\int \frac{x^4}{(1-x^2)^{5/2}} dx$	267
3.40	$\int \frac{\sqrt{A^2+B^2-B^2y^2}}{1-y^2} dy$	273
3.41	$\int \sin^2(x) dx$	280
3.42	$\int \csc(x) \sqrt{A^2+B^2 \sin^2(x)} dx$	285
3.43	$\int \frac{1}{1+\cos(x)} dx$	292
3.44	$\int e^x x dx$	297

3.45	$\int \frac{e^x x}{(1+x)^2} dx$	302
3.46	$\int e^{x^2} (1 + 2x^2) dx$	307
3.47	$\int e^{x^2} dx$	312
3.48	$\int \frac{e^x}{x} dx$	317
3.49	$\int \frac{x}{1+x^3} dx$	322
3.50	$\int \frac{1}{-1+x^6} dx$	328
3.51	$\int \frac{1}{A^4 - A^2 B^2 + (-A^2 + B^2)x^2} dx$	335
3.52	$\int x \log(x) dx$	340
3.53	$\int x^2 \arcsin(x) dx$	345
3.54	$\int \frac{1}{1+2x+x^2} dx$	350
3.55	$\int \frac{\log(x)}{(1+\log(x))^2} dx$	355
3.56	$\int \frac{1}{x(1+\log^2(x))} dx$	360
3.57	$\int \frac{1}{\log(x)} dx$	365
3.58	$\int x(\cos(x) + \sin(x)) dx$	369
3.59	$\int e^{-x}(e^x + x) dx$	374
3.60	$\int (1 + e^x)^2 x dx$	379
3.61	$\int x \cos(x) dx$	384
3.62	$\int \cos(\sqrt{x}) dx$	389
3.63	$\int x \cos(x) dx$	394
3.64	$\int x \log^2(x) dx$	399
3.65	$\int \cos(x) (1 + \sin^3(x)) dx$	404
3.66	$\int \frac{1}{x(1+\log^2(x))} dx$	409
3.67	$\int \frac{1}{\sqrt{1-x^2}(1+\arcsin(x)^2)} dx$	414
3.68	$\int \frac{\sin(x)}{\cos(x)+\sin(x)} dx$	419
3.69	$\int -\frac{\sqrt{A^2+B^2(1-y^2)}}{1-y^2} dy$	425
3.70	$\int \frac{(-A^2-B^2)\cos^2(z)}{B\left(1-\frac{(A^2+B^2)\sin^2(z)}{B^2}\right)} dz$	433
3.71	$\int \frac{-A^2-B^2}{B(1+w^2)^2\left(1-\frac{(A^2+B^2)w^2}{B^2(1+w^2)}\right)} dw$	441
3.72	$\int -\frac{B(A^2+B^2)}{(1+w^2)(B^2-A^2w^2)} dw$	450
3.73	$\int \frac{x^4}{(1-x^2)^{5/2}} dx$	457
3.74	$\int \tan^4(y) dy$	463
3.75	$\int \frac{z^4}{1+z^2} dz$	468
3.76	$\int e^{x^2} (1 + 2x^2) dx$	473
3.77	$\int \frac{e^{x^2}(1+4x^2+x^3+5x^4+2x^6)}{(1+x^2)^2} dx$	478
3.78	$\int e^{-1-x} dx$	483

3.79	$\int \left(\frac{1}{x} + x\right) \log(x) dx$	488
3.80	$\int \frac{x}{1+x^4} dx$	493
3.81	$\int \frac{x^5}{1+x^4} dx$	498
3.82	$\int \frac{1}{1+\tan^2(x)} dx$	503
3.83	$\int \frac{x^4}{(1-x^2)^{5/2}} dx$	508
3.84	$\int -\frac{x^2}{(1-x^2)^{3/2}} dx$	514
3.85	$\int e^x \sin(x) dx$	519
3.86	$\int \frac{1}{x} dx$	524
3.87	$\int \frac{\sec(2t)}{1+\sec^2(t)+3 \tan(t)} dt$	528
3.88	$\int \cos^2(x) dx$	534
3.89	$\int \frac{1+x^2}{\sqrt{x}} dx$	539
3.90	$\int \frac{x}{\sqrt{5+2x+x^2}} dx$	544
3.91	$\int \cos(x) \sin^2(x) dx$	549
3.92	$\int \frac{e^x}{1+e^x} dx$	554
3.93	$\int \frac{e^{2x}}{1+e^x} dx$	559
3.94	$\int \frac{1}{1-\cos(x)} dx$	564
3.95	$\int \sec^2(x) \tan(x) dx$	569
3.96	$\int x \log(x) dx$	574
3.97	$\int \cos(x) \sin(x) dx$	579
3.98	$\int \frac{1+x}{\sqrt{2x-x^2}} dx$	584
3.99	$\int \frac{2e^x}{2+3e^{2x}} dx$	589
3.100	$\int \frac{x^4}{(1-x^2)^{5/2}} dx$	594
3.101	$\int \frac{e^{6x}}{1+e^{4x}} dx$	600
3.102	$\int \log(2 + 3x^2) dx$	605
3.103	$\int \frac{1}{r\sqrt{-a^2+2Hr^2}} dx$	610
3.104	$\int \frac{1}{r\sqrt{-a^2-e^2+2Hr^2}} dx$	615
3.105	$\int \frac{1}{r\sqrt{-a^2+2Hr^2-2Kr^4}} dx$	620
3.106	$\int \frac{1}{r\sqrt{-a^2-e^2+2Hr^2-2Kr^4}} dx$	625
3.107	$\int \frac{1}{r\sqrt{-a^2-2Kr+2Hr^2}} dx$	630
3.108	$\int \frac{1}{r\sqrt{-a^2-e^2-2Kr+2Hr^2}} dx$	635
3.109	$\int \frac{r}{\sqrt{-a^2+2er^2}} dx$	640
3.110	$\int \frac{r}{\sqrt{-a^2-e^2+2er^2}} dx$	645
3.111	$\int \frac{r}{\sqrt{-a^2+2er^2-2Kr^4}} dx$	650
3.112	$\int \frac{r}{\sqrt{-a^2-e^2+2er^2-2Kr^4}} dx$	655
3.113	$\int \frac{r}{\sqrt{-a^2-e^2-2Kr+2Hr^2}} dx$	660

- 4.1 Listing of Grading functions 665
- 4.2 Links to plain text integration problems used in this report for each CAS683

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	7
1.2	Results	8
1.3	Time and leaf size Performance	12
1.4	Performance based on number of rules Rubi used	14
1.5	Performance based on number of steps Rubi used	15
1.6	Solved integrals histogram based on leaf size of result	16
1.7	Solved integrals histogram based on CPU time used	17
1.8	Leaf size vs. CPU time used	18
1.9	list of integrals with no known antiderivative	19
1.10	List of integrals solved by CAS but has no known antiderivative	19
1.11	list of integrals solved by CAS but failed verification	19
1.12	Timing	20
1.13	Verification	20
1.14	Important notes about some of the results	21
1.15	Current tree layout of integration tests	24
1.16	Design of the test system	25

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [**113**]. This is test number [9].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (113)	0.00 (0)
Mathematica	100.00 (113)	0.00 (0)
Maple	100.00 (113)	0.00 (0)
Fricas	99.12 (112)	0.88 (1)
Giac	98.23 (111)	1.77 (2)
Maxima	98.23 (111)	1.77 (2)
Reduce	96.46 (109)	3.54 (4)
Sympy	94.69 (107)	5.31 (6)
Mupad	93.81 (106)	6.19 (7)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	97.345	2.655	0.000	0.000
Mathematica	93.805	5.310	0.885	0.000
Maple	92.920	3.540	3.540	0.000
Giac	88.496	8.850	0.885	1.770
Fricas	87.611	11.504	0.000	0.885
Maxima	86.726	8.850	2.655	1.770
Sympy	80.531	11.504	2.655	5.310
Mupad	0.000	93.805	0.000	6.195
Reduce	0.000	96.460	0.000	3.540

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Maple	0	0.00	0.00	0.00
Fricas	1	100.00	0.00	0.00
Giac	2	100.00	0.00	0.00
Maxima	2	100.00	0.00	0.00
Reduce	4	100.00	0.00	0.00
Sympy	6	100.00	0.00	0.00
Mupad	7	0.00	100.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Mathematica	0.03
Maxima	0.05
Mupad	0.06
Fricas	0.07
Maple	0.11
Giac	0.12
Reduce	0.15
Rubi	0.17
Sympy	1.24

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Maple	19.90	1.07	13.00	0.88
Rubi	21.19	1.06	17.00	1.00
Giac	23.58	1.15	14.00	0.85
Maxima	23.66	1.16	14.00	0.88
Mathematica	24.39	1.13	16.00	1.00
Fricas	25.05	1.17	14.00	0.91
Reduce	25.58	1.21	13.00	1.00
Mupad	28.25	1.65	12.00	0.83
Sympy	30.84	1.68	15.00	0.83

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {23}

Mathematica {}

Maple {79}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand,the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer. the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	27
2.2	Detailed conclusion table per each integral for all CAS systems	32
2.3	Detailed conclusion table specific for Rubi results	61

2.1 List of integrals sorted by grade for each CAS

Rubi	27
Mma	27
Maple	28
Fricas	28
Maxima	29
Giac	29
Mupad	30
Sympy	30
Reduce	30

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113 }

B grade { 70, 71, 72 }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113 }

B grade { 31, 42, 70, 71, 72, 84 }

C grade { 1 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 33, 34, 35, 37, 38, 39, 40, 41, 43, 44, 45, 46, 47, 49, 50, 51, 52, 53, 54, 55, 56, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 69, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113 }

B grade { 36, 48, 57, 70 }

C grade { 32, 42, 68, 71 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { 2, 3, 4, 5, 6, 7, 8, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 33, 34, 35, 36, 38, 41, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 71, 72, 75, 76, 77, 78, 79, 80, 81, 82, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113 }

B grade { 1, 9, 37, 39, 40, 42, 69, 70, 73, 74, 83, 84, 100 }

C grade { }

F normal fail { 32 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maxima

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 33, 34, 35, 37, 39, 41, 43, 44, 45, 46, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113 }

B grade { 36, 38, 40, 42, 68, 69, 70, 71, 72, 87 }

C grade { 10, 11, 47 }

F normal fail { 32, 67 }

F(-1) timedout fail { }

F(-2) exception fail { }

Giac

A grade { 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 26, 27, 28, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 44, 46, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113 }

B grade { 1, 25, 36, 40, 43, 45, 69, 70, 71, 72 }

C grade { 47 }

F normal fail { 32, 42 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 33, 34, 35, 36, 37, 39, 41, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113 }

C grade { }

F normal fail { }

F(-1) timedout fail { 10, 11, 32, 38, 40, 42, 69 }

F(-2) exception fail { }

Sympy

A grade { 1, 2, 3, 4, 5, 6, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 41, 43, 44, 45, 46, 47, 48, 49, 52, 53, 54, 55, 57, 58, 59, 60, 61, 62, 63, 64, 65, 67, 68, 70, 74, 75, 76, 77, 78, 79, 80, 81, 85, 86, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113 }

B grade { 7, 18, 37, 39, 50, 51, 56, 66, 73, 82, 83, 84, 100 }

C grade { 38, 71, 72 }

F normal fail { 35, 36, 40, 42, 69, 87 }

F(-1) timedout fail { }

F(-2) exception fail { }

Reduce

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 33, 34, 35, 36, 37, 38, 39, 40, 41, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113 }

C grade { }

F normal fail { 10, 32, 42, 87 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	22	11	16	48	19	34	10	10
N.S.	1	1.00	1.83	0.92	1.33	4.00	1.58	2.83	0.83	0.83
time (sec)	N/A	0.188	0.003	0.035	0.102	0.075	0.029	0.120	0.147	0.001

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	13	12	15	19	14	15	19	12
N.S.	1	1.00	1.00	0.92	1.15	1.46	1.08	1.15	1.46	0.92
time (sec)	N/A	0.125	0.004	0.094	0.102	0.065	0.046	0.119	0.169	0.094

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	14	11	11	14	15	11	10	10
N.S.	1	1.00	0.74	0.58	0.58	0.74	0.79	0.58	0.53	0.53
time (sec)	N/A	0.128	0.008	0.060	0.023	0.066	0.071	0.116	0.143	0.019

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2	2	2	3	2	2	2	2	2	2
N.S.	1	1.00	1.00	1.50	1.00	1.00	1.00	1.00	1.00	1.00
time (sec)	N/A	0.133	0.004	0.065	0.025	0.071	0.035	0.114	0.151	0.017

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	9	9	9	7	6	6	5	6	7	6
N.S.	1	1.00	1.00	0.78	0.67	0.67	0.56	0.67	0.78	0.67
time (sec)	N/A	0.129	0.009	0.026	0.024	0.067	0.028	0.122	0.150	0.001

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	6	7	6	6	6
N.S.	1	1.00	1.00	0.88	0.75	0.75	0.88	0.75	0.75	0.75
time (sec)	N/A	0.146	0.004	0.116	0.023	0.077	0.021	0.114	0.154	0.012

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	13	10	9	9	22	9	13	9
N.S.	1	1.00	1.00	0.77	0.69	0.69	1.69	0.69	1.00	0.69
time (sec)	N/A	0.118	0.001	0.082	0.023	0.079	0.061	0.123	0.156	0.015

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	14	12	11	13	15	11	12	11
N.S.	1	1.00	0.74	0.63	0.58	0.68	0.79	0.58	0.63	0.58
time (sec)	N/A	0.146	0.007	0.123	0.026	0.080	0.090	0.121	0.156	0.010

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	14	8	6	6	6
N.S.	1	1.00	1.00	0.88	0.75	1.75	1.00	0.75	0.75	0.75
time (sec)	N/A	0.158	0.004	0.121	0.028	0.066	0.021	0.118	0.152	0.028

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	4	4	4	4	15	3	3	3	6	0
N.S.	1	1.00	1.00	1.00	3.75	0.75	0.75	0.75	1.50	0.00
time (sec)	N/A	0.169	0.012	0.108	0.062	0.077	0.300	0.121	0.154	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2	2	2	3	13	2	2	2	2	0
N.S.	1	1.00	1.00	1.50	6.50	1.00	1.00	1.00	1.00	0.00
time (sec)	N/A	0.149	0.012	0.133	0.057	0.086	0.327	0.112	0.146	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	8	7	7	5	7	8	7
N.S.	1	1.00	1.00	1.00	0.88	0.88	0.62	0.88	1.00	0.88
time (sec)	N/A	0.130	0.007	0.144	0.030	0.090	0.027	0.118	0.143	0.023

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	7	7	7	7	6	6	5	6	7	6
N.S.	1	1.00	1.00	1.00	0.86	0.86	0.71	0.86	1.00	0.86
time (sec)	N/A	0.141	0.023	0.036	0.028	0.080	0.029	0.118	0.156	0.093

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	26	21	19	19	20	19	24	21
N.S.	1	1.00	0.93	0.75	0.68	0.68	0.71	0.68	0.86	0.75
time (sec)	N/A	0.187	0.027	0.082	0.030	0.083	0.034	0.118	0.149	0.097

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	22	17	16	16	15	16	18	16
N.S.	1	1.00	1.00	0.77	0.73	0.73	0.68	0.73	0.82	0.73
time (sec)	N/A	0.142	0.008	0.063	0.024	0.077	0.037	0.118	0.167	0.025

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	6	5	6	6	6
N.S.	1	1.00	1.00	0.88	0.75	0.75	0.62	0.75	0.75	0.75
time (sec)	N/A	0.147	0.001	0.146	0.025	0.071	0.039	0.110	0.163	0.009

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	9	9	9	7	6	6	5	6	7	6
N.S.	1	1.00	1.00	0.78	0.67	0.67	0.56	0.67	0.78	0.67
time (sec)	N/A	0.136	0.000	0.027	0.029	0.074	0.031	0.113	0.169	0.001

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	13	10	9	9	22	9	13	9
N.S.	1	1.00	1.00	0.77	0.69	0.69	1.69	0.69	1.00	0.69
time (sec)	N/A	0.121	0.000	0.075	0.025	0.074	0.082	0.114	0.158	0.001

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	6	6	6	6	5	5	5	5	6	5
N.S.	1	1.00	1.00	1.00	0.83	0.83	0.83	0.83	1.00	0.83
time (sec)	N/A	0.147	0.008	0.023	0.024	0.076	0.036	0.118	0.149	0.015

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	9	9	9	6	5	5	7	5	7	5
N.S.	1	1.00	1.00	0.67	0.56	0.56	0.78	0.56	0.78	0.56
time (sec)	N/A	0.118	0.000	0.033	0.025	0.065	0.040	0.119	0.152	0.026

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	10	9	8	8	7	8	8	8
N.S.	1	1.00	1.00	0.90	0.80	0.80	0.70	0.80	0.80	0.80
time (sec)	N/A	0.152	0.016	0.150	0.024	0.083	0.065	0.118	0.150	0.038

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	8	7	7	7	7	8	7
N.S.	1	1.00	1.00	1.00	0.88	0.88	0.88	0.88	1.00	0.88
time (sec)	N/A	0.134	0.000	0.030	0.026	0.066	0.031	0.124	0.155	0.010

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	9	19	8	7	7	8	7	8	7
N.S.	1	0.90	1.90	0.80	0.70	0.70	0.80	0.70	0.80	0.70
time (sec)	N/A	0.193	0.019	0.150	0.031	0.078	0.234	0.118	0.153	0.128

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	18	13	15	15	34	15	14	12
N.S.	1	1.00	0.78	0.57	0.65	0.65	1.48	0.65	0.61	0.52
time (sec)	N/A	0.143	0.009	0.076	0.024	0.064	0.481	0.121	0.152	0.017

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	25	10	17	17	17	19	17	9
N.S.	1	1.00	1.92	0.77	1.31	1.31	1.31	1.46	1.31	0.69
time (sec)	N/A	0.133	0.003	0.093	0.108	0.071	0.070	0.125	0.150	0.085

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	18	14	13	13	15	13	14	13
N.S.	1	1.00	1.00	0.78	0.72	0.72	0.83	0.72	0.78	0.72
time (sec)	N/A	0.162	0.023	0.036	0.103	0.071	0.074	0.114	0.149	0.046

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	31	20	19	76	22	19	71	19
N.S.	1	1.00	1.00	0.65	0.61	2.45	0.71	0.61	2.29	0.61
time (sec)	N/A	0.174	0.024	0.059	0.103	0.078	0.078	0.121	0.155	0.126

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	9	8	11	8	8	8	8
N.S.	1	1.00	1.00	1.12	1.00	1.38	1.00	1.00	1.00	1.00
time (sec)	N/A	0.176	0.010	0.030	0.036	0.069	0.053	0.118	0.154	0.103

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	12	15	14	14	10	13	13	12
N.S.	1	1.00	1.00	1.25	1.17	1.17	0.83	1.08	1.08	1.00
time (sec)	N/A	0.143	0.003	0.067	0.028	0.069	0.034	0.118	0.156	0.035

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	18	20	17	16	16	15	16	16	16
N.S.	1	0.90	1.00	0.85	0.80	0.80	0.75	0.80	0.80	0.80
time (sec)	N/A	0.219	0.003	0.220	0.031	0.087	0.114	0.118	0.157	0.100

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	55	260	39	40	40	46	40	353	52
N.S.	1	1.12	5.31	0.80	0.82	0.82	0.94	0.82	7.20	1.06
time (sec)	N/A	0.196	0.071	0.092	0.107	0.076	0.071	0.126	0.169	0.069

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	115	142	41	0	0	54	0	130	0
N.S.	1	1.00	1.23	0.36	0.00	0.00	0.47	0.00	1.13	0.00
time (sec)	N/A	0.251	0.193	0.278	0.000	0.000	1.386	0.000	0.162	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	21	22	17	16	16	20	16	13	16
N.S.	1	0.95	1.00	0.77	0.73	0.73	0.91	0.73	0.59	0.73
time (sec)	N/A	0.216	0.017	0.057	0.034	0.083	0.111	0.120	0.150	0.119

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	18	13	15	15	34	15	14	12
N.S.	1	1.00	0.78	0.57	0.65	0.65	1.48	0.65	0.61	0.52
time (sec)	N/A	0.138	0.002	0.072	0.025	0.077	0.482	0.122	0.154	0.001

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	36	33	25	24	24	0	24	23	24
N.S.	1	1.12	1.03	0.78	0.75	0.75	0.00	0.75	0.72	0.75
time (sec)	N/A	0.164	0.021	0.060	0.023	0.071	0.000	0.115	0.152	0.001

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	A	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	44	75	75	80	55	0	61	32	57
N.S.	1	1.00	1.70	1.70	1.82	1.25	0.00	1.39	0.73	1.30
time (sec)	N/A	0.156	0.087	0.142	0.112	0.071	0.000	0.130	0.146	0.108

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	44	30	44	63	105	29	55	91
N.S.	1	1.00	1.26	0.86	1.26	1.80	3.00	0.83	1.57	2.60
time (sec)	N/A	0.146	0.064	0.190	0.114	0.077	0.333	0.132	0.155	0.087

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	C	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	85	52	44	113	44	190	90	50	0
N.S.	1	1.13	0.69	0.59	1.51	0.59	2.53	1.20	0.67	0.00
time (sec)	N/A	0.159	0.057	0.081	0.024	0.071	18.274	0.139	0.140	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	44	30	44	63	105	29	55	91
N.S.	1	1.00	1.26	0.86	1.26	1.80	3.00	0.83	1.57	2.60
time (sec)	N/A	0.139	0.000	0.112	0.100	0.082	0.412	0.111	0.145	0.002

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	78	90	114	145	0	295	126	0
N.S.	1	1.00	1.53	1.76	2.24	2.84	0.00	5.78	2.47	0.00
time (sec)	N/A	0.181	0.712	0.399	0.112	0.094	0.000	0.194	0.179	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	11	10	10	10	10	10	10
N.S.	1	1.00	1.00	0.79	0.71	0.71	0.71	0.71	0.71	0.71
time (sec)	N/A	0.151	0.001	0.146	0.030	0.080	0.018	0.121	0.146	0.084

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	C	B	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	57	99	149	116	244	0	0	20	0
N.S.	1	1.16	2.02	3.04	2.37	4.98	0.00	0.00	0.41	0.00
time (sec)	N/A	0.265	0.101	0.521	0.114	0.164	0.000	0.000	0.145	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	9	9	6	5	9	9	3	30	4	4
N.S.	1	1.00	0.67	0.56	1.00	1.00	0.33	3.33	0.44	0.44
time (sec)	N/A	0.153	0.005	0.039	0.028	0.074	0.076	0.123	0.146	0.097

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	7	7	6	6	5	6	7	6
N.S.	1	1.00	0.64	0.64	0.55	0.55	0.45	0.55	0.64	0.55
time (sec)	N/A	0.150	0.001	0.023	0.028	0.077	0.029	0.120	0.154	0.009

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	9	9	9	9	8	8	5	30	9	8
N.S.	1	1.00	1.00	1.00	0.89	0.89	0.56	3.33	1.00	0.89
time (sec)	N/A	0.159	0.044	0.062	0.031	0.061	0.038	0.118	0.151	0.048

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	7	7	7	7	6	6	5	6	7	6
N.S.	1	1.00	1.00	1.00	0.86	0.86	0.71	0.86	1.00	0.86
time (sec)	N/A	0.167	0.001	0.032	0.030	0.070	0.030	0.128	0.164	0.018

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	C	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	11	8	9	7	8	9	9	7
N.S.	1	1.00	1.00	0.73	0.82	0.64	0.73	0.82	0.82	0.64
time (sec)	N/A	0.134	0.010	0.017	0.026	0.071	0.069	0.130	0.149	0.001

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2	2	2	8	2	2	2	2	2	2
N.S.	1	1.00	1.00	4.00	1.00	1.00	1.00	1.00	1.00	1.00
time (sec)	N/A	0.138	0.001	0.028	0.054	0.075	0.309	0.119	0.156	0.005

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	45	40	33	34	34	41	35	33	46
N.S.	1	1.10	0.98	0.80	0.83	0.83	1.00	0.85	0.80	1.12
time (sec)	N/A	0.185	0.007	0.088	0.106	0.072	0.051	0.125	0.162	0.143

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	79	75	66	65	65	83	67	63	88
N.S.	1	1.68	1.60	1.40	1.38	1.38	1.77	1.43	1.34	1.87
time (sec)	N/A	0.227	0.010	0.088	0.107	0.073	0.113	0.119	0.153	0.049

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	21	30	39	27	70	41	33	21
N.S.	1	1.00	1.00	1.43	1.86	1.29	3.33	1.95	1.57	1.00
time (sec)	N/A	0.138	0.003	0.145	0.030	0.069	0.136	0.119	0.155	0.153

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	13	13	12	13	11	9
N.S.	1	1.00	1.00	0.82	0.76	0.76	0.71	0.76	0.65	0.53
time (sec)	N/A	0.139	0.000	0.027	0.021	0.072	0.033	0.124	0.159	0.020

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	43	29	34	33	24	32	38	31	24
N.S.	1	1.08	0.72	0.85	0.82	0.60	0.80	0.95	0.78	0.60
time (sec)	N/A	0.188	0.009	0.049	0.101	0.084	0.084	0.123	0.157	0.001

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	7	7	7	8	7	7	5	7	7	7
N.S.	1	1.00	1.00	1.14	1.00	1.00	0.71	1.00	1.00	1.00
time (sec)	N/A	0.130	0.001	0.084	0.022	0.070	0.027	0.123	0.142	0.011

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	9	8	8	5	8	8	8
N.S.	1	1.00	1.00	1.12	1.00	1.00	0.62	1.00	1.00	1.00
time (sec)	N/A	0.183	0.011	0.083	0.027	0.064	0.034	0.122	0.147	0.148

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	3	3	3	4	3	3	15	3	3	3
N.S.	1	1.00	1.00	1.33	1.00	1.00	5.00	1.00	1.00	1.00
time (sec)	N/A	0.152	0.018	0.069	0.103	0.071	0.072	0.122	0.145	0.186

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2	2	2	9	3	2	2	3	3	2
N.S.	1	1.00	1.00	4.50	1.50	1.00	1.00	1.50	1.50	1.00
time (sec)	N/A	0.131	0.003	0.010	0.048	0.064	0.172	0.119	0.158	0.005

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	15	14	14	15	14	14	14
N.S.	1	1.00	1.00	1.07	1.00	1.00	1.07	1.00	1.00	1.00
time (sec)	N/A	0.162	0.003	0.258	0.024	0.073	0.057	0.121	0.156	0.033

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	13	13	11	14	8	11	16	15
N.S.	1	1.00	0.76	0.76	0.65	0.82	0.47	0.65	0.94	0.88
time (sec)	N/A	0.198	0.004	0.053	0.031	0.066	0.030	0.124	0.159	0.031

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	29	25	24	24	26	24	32	28
N.S.	1	1.00	0.76	0.66	0.63	0.63	0.68	0.63	0.84	0.74
time (sec)	N/A	0.185	0.037	0.067	0.025	0.078	0.037	0.117	0.152	0.096

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	7	7	7	8	7	7	7	7	7	7
N.S.	1	1.00	1.00	1.14	1.00	1.00	1.00	1.00	1.00	1.00
time (sec)	N/A	0.184	0.002	0.115	0.025	0.076	0.065	0.120	0.158	0.010

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	21	22	17	16	16	20	16	13	16
N.S.	1	0.95	1.00	0.77	0.73	0.73	0.91	0.73	0.59	0.73
time (sec)	N/A	0.215	0.004	0.055	0.024	0.083	0.085	0.115	0.160	0.001

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	7	7	7	8	7	7	7	7	7	7
N.S.	1	1.00	1.00	1.14	1.00	1.00	1.00	1.00	1.00	1.00
time (sec)	N/A	0.184	0.001	0.088	0.025	0.073	0.056	0.124	0.141	0.001

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	28	23	17	22	22	22	17	17
N.S.	1	1.00	1.00	0.82	0.61	0.79	0.79	0.79	0.61	0.61
time (sec)	N/A	0.167	0.000	0.069	0.023	0.072	0.040	0.124	0.160	0.017

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	11	10	9	15	8	9	10	9
N.S.	1	1.00	1.00	0.91	0.82	1.36	0.73	0.82	0.91	0.82
time (sec)	N/A	0.175	0.002	0.687	0.027	0.075	0.106	0.123	0.157	0.018

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	3	3	3	4	3	3	15	3	3	3
N.S.	1	1.00	1.00	1.33	1.00	1.00	5.00	1.00	1.00	1.00
time (sec)	N/A	0.148	0.002	0.059	0.104	0.070	0.057	0.125	0.146	0.001

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	3	3	3	4	0	3	3	3	3	43
N.S.	1	1.00	1.00	1.33	0.00	1.00	1.00	1.00	1.00	14.33
time (sec)	N/A	0.197	0.067	0.079	0.000	0.078	0.112	0.123	0.147	1.791

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	20	53	15	12	21	12	13
N.S.	1	1.00	1.00	1.25	3.31	0.94	0.75	1.31	0.75	0.81
time (sec)	N/A	0.235	0.036	0.182	0.112	0.086	0.052	0.118	0.154	0.014

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	73	89	115	144	0	295	126	0
N.S.	1	1.00	1.38	1.68	2.17	2.72	0.00	5.57	2.38	0.00
time (sec)	N/A	0.221	0.123	0.197	0.114	0.085	0.000	0.189	0.158	0.000

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	B	B	B	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	46	35	74	69	67	202	83	74	360
N.S.	1	2.88	2.19	4.62	4.31	4.19	12.62	5.19	4.62	22.50
time (sec)	N/A	0.270	0.175	0.548	0.102	0.095	101.856	0.126	0.158	0.274

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	B	C	B	A	C	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	44	35	70	68	26	422	82	33	352
N.S.	1	2.75	2.19	4.38	4.25	1.62	26.38	5.12	2.06	22.00
time (sec)	N/A	0.306	0.013	0.121	0.106	0.076	0.894	0.123	0.156	0.134

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	B	A	B	A	C	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	44	35	29	65	26	422	79	33	352
N.S.	1	2.75	2.19	1.81	4.06	1.62	26.38	4.94	2.06	22.00
time (sec)	N/A	0.166	0.009	0.163	0.101	0.077	0.894	0.116	0.151	0.031

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	44	30	44	63	105	29	55	91
N.S.	1	1.00	1.26	0.86	1.26	1.80	3.00	0.83	1.57	2.60
time (sec)	N/A	0.146	0.001	0.141	0.102	0.069	0.354	0.128	0.157	0.001

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	16	13	12	26	19	12	39	12
N.S.	1	1.00	1.14	0.93	0.86	1.86	1.36	0.86	2.79	0.86
time (sec)	N/A	0.194	0.005	0.255	0.102	0.081	0.021	0.118	0.173	0.041

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	13	12	11	11	8	11	11	11
N.S.	1	1.00	1.00	0.92	0.85	0.85	0.62	0.85	0.85	0.85
time (sec)	N/A	0.140	0.004	0.085	0.107	0.064	0.034	0.117	0.153	0.016

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	7	7	7	7	6	6	5	6	7	6
N.S.	1	1.00	1.00	1.00	0.86	0.86	0.71	0.86	1.00	0.86
time (sec)	N/A	0.170	0.001	0.032	0.025	0.071	0.032	0.119	0.150	0.001

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	19	24	23	23	20	30	25	24
N.S.	1	1.00	0.79	1.00	0.96	0.96	0.83	1.25	1.04	1.00
time (sec)	N/A	0.450	0.239	0.095	0.140	0.073	0.049	0.119	0.143	0.133

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	9	9	9	9	8	8	7	8	10	8
N.S.	1	1.00	1.00	1.00	0.89	0.89	0.78	0.89	1.11	0.89
time (sec)	N/A	0.139	0.009	0.031	0.023	0.061	0.025	0.117	0.149	0.018

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	25	20	24	19	19	19	19	19
N.S.	1	1.00	1.00	0.80	0.96	0.76	0.76	0.76	0.76	0.76
time (sec)	N/A	0.195	0.002	0.052	0.027	0.071	0.037	0.114	0.154	0.122

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	6	5	6	29	6
N.S.	1	1.00	1.00	0.88	0.75	0.75	0.62	0.75	3.62	0.75
time (sec)	N/A	0.129	0.003	0.086	0.101	0.065	0.039	0.123	0.148	0.095

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	14	16	13	12	12	10	12	34	12
N.S.	1	0.88	1.00	0.81	0.75	0.75	0.62	0.75	2.12	0.75
time (sec)	N/A	0.138	0.004	0.082	0.113	0.065	0.036	0.118	0.150	0.011

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	11	16	20	36	16	21	10
N.S.	1	1.00	1.00	0.79	1.14	1.43	2.57	1.14	1.50	0.71
time (sec)	N/A	0.190	0.002	0.049	0.101	0.073	0.165	0.121	0.152	0.108

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	44	30	44	63	105	29	55	91
N.S.	1	1.00	1.26	0.86	1.26	1.80	3.00	0.83	1.57	2.60
time (sec)	N/A	0.144	0.000	0.131	0.102	0.070	0.333	0.126	0.151	0.001

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	35	16	15	45	34	21	26	37
N.S.	1	1.00	2.06	0.94	0.88	2.65	2.00	1.24	1.53	2.18
time (sec)	N/A	0.133	0.046	0.144	0.101	0.090	0.178	0.117	0.151	0.093

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	14	12	11	13	15	11	12	11
N.S.	1	1.00	0.74	0.63	0.58	0.68	0.79	0.58	0.63	0.58
time (sec)	N/A	0.152	0.007	0.130	0.025	0.073	0.085	0.123	0.150	0.001

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2	2	2	3	2	2	2	3	2	2
N.S.	1	1.00	1.00	1.50	1.00	1.00	1.00	1.50	1.00	1.00
time (sec)	N/A	0.114	0.000	0.013	0.025	0.065	0.039	0.110	0.154	0.005

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	40	73	31	256	71	0	33	207	32
N.S.	1	0.89	1.62	0.69	5.69	1.58	0.00	0.73	4.60	0.71
time (sec)	N/A	0.282	0.450	1.006	0.182	0.112	0.000	0.131	0.190	0.382

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	11	10	10	10	16	10	10
N.S.	1	1.00	1.00	0.79	0.71	0.71	0.71	1.14	0.71	0.71
time (sec)	N/A	0.159	0.001	0.049	0.024	0.076	0.022	0.125	0.152	0.001

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	14	11	11	10	14	11	9	10
N.S.	1	1.00	0.82	0.65	0.65	0.59	0.82	0.65	0.53	0.59
time (sec)	N/A	0.136	0.008	0.042	0.022	0.063	0.076	0.120	0.156	0.012

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	25	31	20	19	27	19	27	29	27
N.S.	1	1.09	1.35	0.87	0.83	1.17	0.83	1.17	1.26	1.17
time (sec)	N/A	0.152	0.040	0.286	0.105	0.069	0.248	0.127	0.151	0.115

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	10	5	6	6	6
N.S.	1	1.00	1.00	0.88	0.75	1.25	0.62	0.75	0.75	0.75
time (sec)	N/A	0.158	0.001	0.274	0.026	0.074	0.035	0.121	0.157	0.015

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	6	6	6	6	5	5	5	5	6	5
N.S.	1	1.00	1.00	1.00	0.83	0.83	0.83	0.83	1.00	0.83
time (sec)	N/A	0.146	0.001	0.017	0.027	0.065	0.028	0.113	0.153	0.001

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	12	11	10	10	8	10	12	10
N.S.	1	1.00	1.00	0.92	0.83	0.83	0.67	0.83	1.00	0.83
time (sec)	N/A	0.170	0.013	0.025	0.023	0.070	0.033	0.111	0.154	0.027

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	8	7	10	10	7	8	8	6
N.S.	1	1.00	0.67	0.58	0.83	0.83	0.58	0.67	0.67	0.50
time (sec)	N/A	0.166	0.034	0.076	0.028	0.069	0.156	0.111	0.156	0.114

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	6	7	6	6	6
N.S.	1	1.00	1.00	0.88	0.75	0.75	0.88	0.75	0.75	0.75
time (sec)	N/A	0.157	0.001	0.102	0.030	0.078	0.024	0.115	0.153	0.001

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	13	13	12	13	11	9
N.S.	1	1.00	1.00	0.82	0.76	0.76	0.71	0.76	0.65	0.53
time (sec)	N/A	0.141	0.000	0.022	0.022	0.066	0.031	0.117	0.153	0.001

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	6	5	6	6	6
N.S.	1	1.00	1.00	0.88	0.75	0.75	0.62	0.75	0.75	0.75
time (sec)	N/A	0.154	0.001	0.115	0.023	0.078	0.035	0.112	0.159	0.001

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	28	47	21	22	34	15	20	31	20
N.S.	1	1.17	1.96	0.88	0.92	1.42	0.62	0.83	1.29	0.83
time (sec)	N/A	0.154	0.037	0.299	0.101	0.071	0.330	0.131	0.155	0.155

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	14	13	12	15	13	14	13
N.S.	1	1.00	1.00	0.70	0.65	0.60	0.75	0.65	0.70	0.65
time (sec)	N/A	0.158	0.004	0.033	0.108	0.072	0.047	0.118	0.152	0.046

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	44	30	44	63	105	29	55	91
N.S.	1	1.00	1.26	0.86	1.26	1.80	3.00	0.83	1.57	2.60
time (sec)	N/A	0.143	0.001	0.136	0.103	0.073	0.339	0.123	0.153	0.001

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	18	18	15	14	14	24	14	42	14
N.S.	1	0.90	0.90	0.75	0.70	0.70	1.20	0.70	2.10	0.70
time (sec)	N/A	0.158	0.021	0.039	0.106	0.074	0.048	0.116	0.158	0.040

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	40	33	27	26	26	31	26	26	26
N.S.	1	1.21	1.00	0.82	0.79	0.79	0.94	0.79	0.79	0.79
time (sec)	N/A	0.154	0.009	0.033	0.105	0.078	0.052	0.119	0.154	0.015

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	21	20	19	31	15	19	31	19
N.S.	1	1.00	1.00	0.95	0.90	1.48	0.71	0.90	1.48	0.90
time (sec)	N/A	0.142	0.000	0.036	0.023	0.062	0.016	0.124	0.154	0.001

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	26	25	24	40	19	24	39	24
N.S.	1	1.00	1.00	0.96	0.92	1.54	0.73	0.92	1.50	0.92
time (sec)	N/A	0.144	0.000	0.033	0.023	0.060	0.018	0.111	0.153	0.001

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	27	26	25	43	22	25	43	25
N.S.	1	1.00	1.00	0.96	0.93	1.59	0.81	0.93	1.59	0.93
time (sec)	N/A	0.146	0.000	0.041	0.024	0.061	0.016	0.117	0.151	0.001

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	32	31	30	52	26	30	51	30
N.S.	1	1.00	1.00	0.97	0.94	1.62	0.81	0.94	1.59	0.94
time (sec)	N/A	0.151	0.000	0.047	0.023	0.066	0.017	0.120	0.157	0.001

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	25	24	23	41	20	23	39	23
N.S.	1	1.00	1.04	1.00	0.96	1.71	0.83	0.96	1.62	0.96
time (sec)	N/A	0.149	0.000	0.040	0.029	0.062	0.015	0.117	0.156	0.002

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	30	29	28	50	24	28	47	28
N.S.	1	1.00	1.03	1.00	0.97	1.72	0.83	0.97	1.62	0.97
time (sec)	N/A	0.154	0.000	0.042	0.024	0.063	0.017	0.114	0.151	0.001

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	19	19	18	18	17	18	29	18
N.S.	1	1.00	1.00	1.00	0.95	0.95	0.89	0.95	1.53	0.95
time (sec)	N/A	0.136	0.000	0.042	0.026	0.063	0.014	0.111	0.144	0.001

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	24	24	23	23	20	23	37	23
N.S.	1	1.00	1.00	1.00	0.96	0.96	0.83	0.96	1.54	0.96
time (sec)	N/A	0.141	0.000	0.040	0.029	0.062	0.013	0.122	0.153	0.001

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	25	25	24	44	24	24	41	24
N.S.	1	1.00	1.00	1.00	0.96	1.76	0.96	0.96	1.64	0.96
time (sec)	N/A	0.144	0.000	0.055	0.031	0.067	0.014	0.116	0.148	0.001

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	30	30	30	29	52	27	29	49	29
N.S.	1	1.00	1.00	1.00	0.97	1.73	0.90	0.97	1.63	0.97
time (sec)	N/A	0.147	0.000	0.049	0.023	0.063	0.014	0.122	0.153	0.002

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	28	27	26	26	24	26	45	26
N.S.	1	1.00	1.04	1.00	0.96	0.96	0.89	0.96	1.67	0.96
time (sec)	N/A	0.141	0.000	0.040	0.024	0.060	0.013	0.121	0.149	0.001

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. The column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [1] had the largest ratio of [1.2500000000000000]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	5	5	1.00	4	1.250
2	A	3	3	1.00	11	0.273
3	A	3	3	1.00	11	0.273
4	A	2	2	1.00	2	1.000
5	A	1	1	1.00	7	0.143
6	A	4	3	1.00	7	0.429
7	A	1	1	1.00	11	0.091
8	A	1	1	1.00	6	0.167
9	A	5	4	1.00	7	0.571
10	A	4	3	1.00	4	0.750
11	A	2	2	1.00	6	0.333
12	A	1	1	1.00	6	0.167
13	A	1	1	1.00	16	0.062
14	A	2	2	1.00	7	0.286
15	A	1	1	1.00	14	0.071
16	A	4	3	1.00	5	0.600
17	A	1	1	1.00	7	0.143
18	A	1	1	1.00	11	0.091
19	A	3	2	1.00	11	0.182
20	A	1	1	1.00	5	0.200
21	A	2	2	1.00	6	0.333

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	2	2	1.00	9	0.222
23	A	5	4	0.90	14	0.286
24	A	2	2	1.00	9	0.222
25	A	3	3	1.00	7	0.429
26	A	3	2	1.00	15	0.133
27	A	3	2	1.00	17	0.118
28	A	4	3	1.00	13	0.231
29	A	1	1	1.00	5	0.200
30	A	6	5	0.90	8	0.625
31	A	9	8	1.12	11	0.727
32	A	2	2	1.00	12	0.167
33	A	7	6	0.95	6	1.000
34	A	2	2	1.00	9	0.222
35	A	5	4	1.12	13	0.308
36	A	5	4	1.00	15	0.267
37	A	3	3	1.00	15	0.200
38	A	7	6	1.13	13	0.462
39	A	3	3	1.00	15	0.200
40	A	6	5	1.00	29	0.172
41	A	3	3	1.00	4	0.750
42	A	8	7	1.16	19	0.368
43	A	2	2	1.00	6	0.333
44	A	2	2	1.00	5	0.400
45	A	1	1	1.00	10	0.100
46	A	2	2	1.00	13	0.154
47	A	1	1	1.00	5	0.200
48	A	1	1	1.00	7	0.143
49	A	8	7	1.10	9	0.778
50	A	9	8	1.68	7	1.143
51	A	1	1	1.00	27	0.037
52	A	1	1	1.00	4	0.250
53	A	5	4	1.08	6	0.667

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
54	A	2	2	1.00	10	0.200
55	A	2	2	1.00	9	0.222
56	A	3	2	1.00	12	0.167
57	A	1	1	1.00	4	0.250
58	A	2	2	1.00	7	0.286
59	A	2	2	1.00	11	0.182
60	A	2	2	1.00	9	0.222
61	A	5	5	1.00	4	1.250
62	A	7	6	0.95	6	1.000
63	A	5	5	1.00	4	1.250
64	A	2	2	1.00	6	0.333
65	A	4	3	1.00	9	0.333
66	A	3	2	1.00	12	0.167
67	A	3	2	1.00	20	0.100
68	A	4	4	1.00	10	0.400
69	A	8	7	1.00	30	0.233
70	B	8	7	2.88	39	0.179
71	B	6	6	2.75	48	0.125
72	B	5	5	2.75	31	0.161
73	A	3	3	1.00	15	0.200
74	A	5	5	1.00	4	1.250
75	A	2	2	1.00	11	0.182
76	A	2	2	1.00	13	0.154
77	A	2	2	1.00	33	0.061
78	A	1	1	1.00	7	0.143
79	A	3	3	1.00	8	0.375
80	A	3	2	1.00	9	0.222
81	A	4	3	0.88	11	0.273
82	A	5	5	1.00	8	0.625
83	A	3	3	1.00	15	0.200
84	A	3	3	1.00	16	0.188
85	A	1	1	1.00	6	0.167

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
86	A	1	1	1.00	3	0.333
87	A	5	4	0.89	17	0.235
88	A	3	3	1.00	4	0.750
89	A	2	2	1.00	11	0.182
90	A	4	3	1.09	14	0.214
91	A	4	3	1.00	7	0.429
92	A	3	2	1.00	11	0.182
93	A	4	3	1.00	13	0.231
94	A	2	2	1.00	8	0.250
95	A	4	3	1.00	7	0.429
96	A	1	1	1.00	4	0.250
97	A	4	3	1.00	5	0.600
98	A	4	3	1.17	17	0.176
99	A	4	3	1.00	16	0.188
100	A	3	3	1.00	15	0.200
101	A	4	3	0.90	15	0.200
102	A	3	3	1.21	8	0.375
103	A	1	1	1.00	20	0.050
104	A	1	1	1.00	25	0.040
105	A	1	1	1.00	26	0.038
106	A	1	1	1.00	31	0.032
107	A	1	1	1.00	24	0.042
108	A	1	1	1.00	29	0.034
109	A	1	1	1.00	18	0.056
110	A	1	1	1.00	23	0.043
111	A	1	1	1.00	24	0.042
112	A	1	1	1.00	29	0.034
113	A	1	1	1.00	27	0.037

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \cot^4(x) dx$	69
3.2	$\int \frac{1}{x^4(1+x^2)} dx$	74
3.3	$\int \frac{x+x^2}{\sqrt{x}} dx$	79
3.4	$\int \cos(x) dx$	84
3.5	$\int e^{x^2} x dx$	89
3.6	$\int \sec^2(x) \tan(x) dx$	94
3.7	$\int x\sqrt{1+x^2} dx$	99
3.8	$\int e^x \sin(x) dx$	104
3.9	$\int \cot(x) \csc^3(x) dx$	109
3.10	$\int \sin(e^x) dx$	114
3.11	$\int \frac{\sin(y)}{y} dy$	119
3.12	$\int (e^x + \sin(x)) dx$	124
3.13	$\int (e^{x^2} + 2e^{x^2} x^2) dx$	129
3.14	$\int (e^x + x)^2 dx$	134
3.15	$\int (2e^x + e^{2x} + x^2) dx$	139
3.16	$\int \cos(x) \sin(x) dx$	144
3.17	$\int e^{x^2} x dx$	149
3.18	$\int x\sqrt{1+x^2} dx$	154
3.19	$\int \frac{e^x}{1+e^x} dx$	159
3.20	$\int x^{3/2} dx$	164
3.21	$\int \cos(3+2x) dx$	169
3.22	$\int 2e^{2x} yz dx$	174
3.23	$\int e^x \cos^2(e^x) \sin(e^x) dx$	179
3.24	$\int x\sqrt{1+x} dx$	184
3.25	$\int \frac{1}{-1+x^4} dx$	189
3.26	$\int \frac{e^x}{2+3e^{2x}} dx$	194

3.27	$\int \frac{e^{2x}}{A+Be^{4x}} dx$	199
3.28	$\int \frac{e^{1+x}}{1+e^x} dx$	204
3.29	$\int (10e)^x dx$	209
3.30	$\int x^3 \sin(x^2) dx$	214
3.31	$\int \frac{x^7}{1+x^{12}} dx$	220
3.32	$\int x^{3a} \sin(x^{2a}) dx$	229
3.33	$\int \cos(\sqrt{x}) dx$	234
3.34	$\int x\sqrt{1+x} dx$	239
3.35	$\int \frac{1}{\sqrt[3]{x+\sqrt{x}}} dx$	244
3.36	$\int \sqrt{\frac{1+x}{3+2x}} dx$	249
3.37	$\int \frac{x^4}{(1-x^2)^{5/2}} dx$	255
3.38	$\int \sqrt{x}(1+x)^{5/2} dx$	261
3.39	$\int \frac{x^4}{(1-x^2)^{5/2}} dx$	267
3.40	$\int \frac{\sqrt{A^2+B^2-B^2y^2}}{1-y^2} dy$	273
3.41	$\int \sin^2(x) dx$	280
3.42	$\int \csc(x) \sqrt{A^2+B^2 \sin^2(x)} dx$	285
3.43	$\int \frac{1}{1+\cos(x)} dx$	292
3.44	$\int e^x x dx$	297
3.45	$\int \frac{e^x x}{(1+x)^2} dx$	302
3.46	$\int e^{x^2} (1+2x^2) dx$	307
3.47	$\int e^{x^2} dx$	312
3.48	$\int \frac{e^x}{x} dx$	317
3.49	$\int \frac{x}{1+x^3} dx$	322
3.50	$\int \frac{1}{-1+x^6} dx$	328
3.51	$\int \frac{1}{A^4-A^2B^2+(-A^2+B^2)x^2} dx$	335
3.52	$\int x \log(x) dx$	340
3.53	$\int x^2 \arcsin(x) dx$	345
3.54	$\int \frac{1}{1+2x+x^2} dx$	350
3.55	$\int \frac{\log(x)}{(1+\log(x))^2} dx$	355
3.56	$\int \frac{1}{x(1+\log^2(x))} dx$	360
3.57	$\int \frac{1}{\log(x)} dx$	365
3.58	$\int x(\cos(x) + \sin(x)) dx$	369
3.59	$\int e^{-x}(e^x + x) dx$	374
3.60	$\int (1+e^x)^2 x dx$	379
3.61	$\int x \cos(x) dx$	384
3.62	$\int \cos(\sqrt{x}) dx$	389
3.63	$\int x \cos(x) dx$	394

3.64	$\int x \log^2(x) dx$	399
3.65	$\int \cos(x) (1 + \sin^3(x)) dx$	404
3.66	$\int \frac{1}{x(1+\log^2(x))} dx$	409
3.67	$\int \frac{1}{\sqrt{1-x^2(1+\arcsin(x)^2)}} dx$	414
3.68	$\int \frac{\sin(x)}{\cos(x)+\sin(x)} dx$	419
3.69	$\int -\frac{\sqrt{A^2+B^2(1-y^2)}}{1-y^2} dy$	425
3.70	$\int \frac{(-A^2-B^2)\cos^2(z)}{B\left(1-\frac{(A^2+B^2)\sin^2(z)}{B^2}\right)} dz$	433
3.71	$\int \frac{-A^2-B^2}{B(1+w^2)^2\left(1-\frac{(A^2+B^2)w^2}{B^2(1+w^2)}\right)} dw$	441
3.72	$\int -\frac{B(A^2+B^2)}{(1+w^2)(B^2-A^2w^2)} dw$	450
3.73	$\int \frac{x^4}{(1-x^2)^{5/2}} dx$	457
3.74	$\int \tan^4(y) dy$	463
3.75	$\int \frac{z^4}{1+z^2} dz$	468
3.76	$\int e^{x^2}(1+2x^2) dx$	473
3.77	$\int \frac{e^{x^2}(1+4x^2+x^3+5x^4+2x^6)}{(1+x^2)^2} dx$	478
3.78	$\int e^{-1-x} dx$	483
3.79	$\int \left(\frac{1}{x} + x\right) \log(x) dx$	488
3.80	$\int \frac{x}{1+x^4} dx$	493
3.81	$\int \frac{x^5}{1+x^4} dx$	498
3.82	$\int \frac{1}{1+\tan^2(x)} dx$	503
3.83	$\int \frac{x^4}{(1-x^2)^{5/2}} dx$	508
3.84	$\int -\frac{x^2}{(1-x^2)^{3/2}} dx$	514
3.85	$\int e^x \sin(x) dx$	519
3.86	$\int \frac{1}{x} dx$	524
3.87	$\int \frac{\sec(2t)}{1+\sec^2(t)+3\tan(t)} dt$	528
3.88	$\int \cos^2(x) dx$	534
3.89	$\int \frac{1+x^2}{\sqrt{x}} dx$	539
3.90	$\int \frac{x}{\sqrt{5+2x+x^2}} dx$	544
3.91	$\int \cos(x) \sin^2(x) dx$	549
3.92	$\int \frac{e^x}{1+e^x} dx$	554
3.93	$\int \frac{e^{2x}}{1+e^x} dx$	559
3.94	$\int \frac{1}{1-\cos(x)} dx$	564
3.95	$\int \sec^2(x) \tan(x) dx$	569
3.96	$\int x \log(x) dx$	574
3.97	$\int \cos(x) \sin(x) dx$	579

3.98	$\int \frac{1+x}{\sqrt{2x-x^2}} dx$	584
3.99	$\int \frac{2e^x}{2+3e^{2x}} dx$	589
3.100	$\int \frac{x^4}{(1-x^2)^{5/2}} dx$	594
3.101	$\int \frac{e^{6x}}{1+e^{4x}} dx$	600
3.102	$\int \log(2+3x^2) dx$	605
3.103	$\int \frac{1}{r\sqrt{-a^2+2Hr^2}} dx$	610
3.104	$\int \frac{1}{r\sqrt{-a^2-e^2+2Hr^2}} dx$	615
3.105	$\int \frac{1}{r\sqrt{-a^2+2Hr^2-2Kr^4}} dx$	620
3.106	$\int \frac{1}{r\sqrt{-a^2-e^2+2Hr^2-2Kr^4}} dx$	625
3.107	$\int \frac{1}{r\sqrt{-a^2-2Kr+2Hr^2}} dx$	630
3.108	$\int \frac{1}{r\sqrt{-a^2-e^2-2Kr+2Hr^2}} dx$	635
3.109	$\int \frac{r}{\sqrt{-a^2+2er^2}} dx$	640
3.110	$\int \frac{r}{\sqrt{-a^2-e^2+2er^2}} dx$	645
3.111	$\int \frac{r}{\sqrt{-a^2+2er^2-2Kr^4}} dx$	650
3.112	$\int \frac{r}{\sqrt{-a^2-e^2+2er^2-2Kr^4}} dx$	655
3.113	$\int \frac{r}{\sqrt{-a^2-e^2-2Kr+2Hr^2}} dx$	660

3.1 $\int \cot^4(x) dx$

Optimal result	69
Mathematica [C] (verified)	69
Rubi [A] (verified)	70
Maple [A] (verified)	71
Fricas [B] (verification not implemented)	72
Sympy [A] (verification not implemented)	72
Maxima [A] (verification not implemented)	72
Giac [B] (verification not implemented)	73
Mupad [B] (verification not implemented)	73
Reduce [B] (verification not implemented)	73

Optimal result

Integrand size = 4, antiderivative size = 12

$$\int \cot^4(x) dx = x + \cot(x) - \frac{\cot^3(x)}{3}$$

output `x+cot(x)-1/3*cot(x)^3`

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.00 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.83

$$\int \cot^4(x) dx = -\frac{1}{3} \cot^3(x) \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, -\tan^2(x)\right)$$

input `Integrate[Cot[x]^4,x]`

output `-1/3*(Cot[x]^3*Hypergeometric2F1[-3/2, 1, -1/2, -Tan[x]^2])`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$, Rules used = {3042, 3954, 3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^4(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan\left(x + \frac{\pi}{2}\right)^4 dx \\
 & \quad \downarrow \text{3954} \\
 & - \int \cot^2(x) dx - \frac{1}{3} \cot^3(x) \\
 & \quad \downarrow \text{3042} \\
 & - \int \tan\left(x + \frac{\pi}{2}\right)^2 dx - \frac{1}{3} \cot^3(x) \\
 & \quad \downarrow \text{3954} \\
 & \int 1 dx - \frac{1}{3} \cot^3(x) + \cot(x) \\
 & \quad \downarrow \text{24} \\
 & x - \frac{1}{3} \cot^3(x) + \cot(x)
 \end{aligned}$$

input `Int[Cot[x]^4,x]`

output `x + Cot[x] - Cot[x]^3/3`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

method	result	size
parallelrisch	$x + \cot(x) - \frac{\cot(x)^3}{3}$	11
derivativedivides	$-\frac{\cot(x)^3}{3} + \cot(x) - \frac{\pi}{2} + \operatorname{arccot}(\cot(x))$	16
default	$-\frac{\cot(x)^3}{3} + \cot(x) - \frac{\pi}{2} + \operatorname{arccot}(\cot(x))$	16
norman	$\frac{-\frac{1}{3} + \tan(x)^2 + x \tan(x)^3}{\tan(x)^3}$	18
risch	$x + \frac{4i(3e^{4ix} - 3e^{2ix} + 2)}{3(e^{2ix} - 1)^3}$	31

input `int(cot(x)^4,x,method=_RETURNVERBOSE)`

output `x+cot(x)-1/3*cot(x)^3`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 48 vs. $2(10) = 20$.

Time = 0.07 (sec) , antiderivative size = 48, normalized size of antiderivative = 4.00

$$\int \cot^4(x) dx = \frac{4 \cos(2x)^2 + 3(x \cos(2x) - x) \sin(2x) + 2 \cos(2x) - 2}{3(\cos(2x) - 1) \sin(2x)}$$

input `integrate(cot(x)^4,x, algorithm="fricas")`

output `1/3*(4*cos(2*x)^2 + 3*(x*cos(2*x) - x)*sin(2*x) + 2*cos(2*x) - 2)/((cos(2*x) - 1)*sin(2*x))`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.58

$$\int \cot^4(x) dx = x + \frac{\cos(x)}{\sin(x)} - \frac{\cos^3(x)}{3 \sin^3(x)}$$

input `integrate(cot(x)**4,x)`

output `x + cos(x)/sin(x) - cos(x)**3/(3*sin(x)**3)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.33

$$\int \cot^4(x) dx = x + \frac{3 \tan(x)^2 - 1}{3 \tan(x)^3}$$

input `integrate(cot(x)^4,x, algorithm="maxima")`

output `x + 1/3*(3*tan(x)^2 - 1)/tan(x)^3`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 34 vs. $2(10) = 20$.

Time = 0.12 (sec) , antiderivative size = 34, normalized size of antiderivative = 2.83

$$\int \cot^4(x) dx = \frac{1}{24} \tan\left(\frac{1}{2}x\right)^3 + x + \frac{15 \tan\left(\frac{1}{2}x\right)^2 - 1}{24 \tan\left(\frac{1}{2}x\right)^3} - \frac{5}{8} \tan\left(\frac{1}{2}x\right)$$

input `integrate(cot(x)^4,x, algorithm="giac")`

output `1/24*tan(1/2*x)^3 + x + 1/24*(15*tan(1/2*x)^2 - 1)/tan(1/2*x)^3 - 5/8*tan(1/2*x)`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \cot^4(x) dx = -\frac{\cot(x)^3}{3} + \cot(x) + x$$

input `int(cot(x)^4,x)`

output `x + cot(x) - cot(x)^3/3`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \cot^4(x) dx = -\frac{\cot(x)^3}{3} + \cot(x) + x$$

input `int(cot(x)^4,x)`

output `(- cot(x)**3 + 3*cot(x) + 3*x)/3`

3.2 $\int \frac{1}{x^4(1+x^2)} dx$

Optimal result	74
Mathematica [A] (verified)	74
Rubi [A] (verified)	75
Maple [A] (verified)	76
Fricas [A] (verification not implemented)	76
Sympy [A] (verification not implemented)	77
Maxima [A] (verification not implemented)	77
Giac [A] (verification not implemented)	77
Mupad [B] (verification not implemented)	78
Reduce [B] (verification not implemented)	78

Optimal result

Integrand size = 11, antiderivative size = 13

$$\int \frac{1}{x^4(1+x^2)} dx = -\frac{1}{3x^3} + \frac{1}{x} + \arctan(x)$$

output `-1/3/x^3+1/x+arctan(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^4(1+x^2)} dx = -\frac{1}{3x^3} + \frac{1}{x} + \arctan(x)$$

input `Integrate[1/(x^4*(1 + x^2)),x]`

output `-1/3*1/x^3 + x^(-1) + ArcTan[x]`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {264, 264, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{x^4(x^2+1)} dx \\ & \quad \downarrow 264 \\ & - \int \frac{1}{x^2(x^2+1)} dx - \frac{1}{3x^3} \\ & \quad \downarrow 264 \\ & \int \frac{1}{x^2+1} dx - \frac{1}{3x^3} + \frac{1}{x} \\ & \quad \downarrow 216 \\ & \arctan(x) - \frac{1}{3x^3} + \frac{1}{x} \end{aligned}$$

input `Int[1/(x^4*(1 + x^2)),x]`

output `-1/3*1/x^3 + x^(-1) + ArcTan[x]`

Defintions of rubi rules used

rule 216

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

rule 264

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(
m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*((m + 2*p + 3)/(a*c
^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p
}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]
```

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
default	$-\frac{1}{3x^3} + \frac{1}{x} + \arctan(x)$	12
meijerg	$-\frac{1}{3x^3} + \frac{1}{x} + \arctan(x)$	12
risch	$\frac{x^2 - \frac{1}{3}}{x^3} + \arctan(x)$	13
parallelrisc	$-\frac{3i \ln(x-i)x^3 - 3i \ln(x+i)x^3 + 2 - 6x^2}{6x^3}$	35

input

```
int(1/x^4/(x^2+1),x,method=_RETURNVERBOSE)
```

output

```
-1/3/x^3+1/x+arctan(x)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.46

$$\int \frac{1}{x^4(1+x^2)} dx = \frac{3x^3 \arctan(x) + 3x^2 - 1}{3x^3}$$

input

```
integrate(1/x^4/(x^2+1),x, algorithm="fricas")
```

output

```
1/3*(3*x^3*arctan(x) + 3*x^2 - 1)/x^3
```

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.08

$$\int \frac{1}{x^4(1+x^2)} dx = \operatorname{atan}(x) + \frac{3x^2 - 1}{3x^3}$$

input `integrate(1/x**4/(x**2+1),x)`output `atan(x) + (3*x**2 - 1)/(3*x**3)`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{1}{x^4(1+x^2)} dx = \frac{3x^2 - 1}{3x^3} + \arctan(x)$$

input `integrate(1/x^4/(x^2+1),x, algorithm="maxima")`output `1/3*(3*x^2 - 1)/x^3 + arctan(x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{1}{x^4(1+x^2)} dx = \frac{3x^2 - 1}{3x^3} + \arctan(x)$$

input `integrate(1/x^4/(x^2+1),x, algorithm="giac")`output `1/3*(3*x^2 - 1)/x^3 + arctan(x)`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

$$\int \frac{1}{x^4(1+x^2)} dx = \operatorname{atan}(x) + \frac{x^2 - \frac{1}{3}}{x^3}$$

input `int(1/(x^4*(x^2 + 1)),x)`

output `atan(x) + (x^2 - 1/3)/x^3`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.46

$$\int \frac{1}{x^4(1+x^2)} dx = \frac{3\operatorname{atan}(x)x^3 + 3x^2 - 1}{3x^3}$$

input `int(1/x^4/(x^2+1),x)`

output `(3*atan(x)*x**3 + 3*x**2 - 1)/(3*x**3)`

3.3 $\int \frac{x+x^2}{\sqrt{x}} dx$

Optimal result	79
Mathematica [A] (verified)	79
Rubi [A] (verified)	80
Maple [A] (verified)	81
Fricas [A] (verification not implemented)	81
Sympy [A] (verification not implemented)	82
Maxima [A] (verification not implemented)	82
Giac [A] (verification not implemented)	82
Mupad [B] (verification not implemented)	83
Reduce [B] (verification not implemented)	83

Optimal result

Integrand size = 11, antiderivative size = 19

$$\int \frac{x+x^2}{\sqrt{x}} dx = \frac{2x^{3/2}}{3} + \frac{2x^{5/2}}{5}$$

output `2/3*x^(3/2)+2/5*x^(5/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int \frac{x+x^2}{\sqrt{x}} dx = \frac{2}{15}x^{3/2}(5+3x)$$

input `Integrate[(x + x^2)/Sqrt[x],x]`

output `(2*x^(3/2)*(5 + 3*x))/15`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {9, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2 + x}{\sqrt{x}} dx$$

↓ 9

$$\int \sqrt{x}(x + 1) dx$$

↓ 53

$$\int (x^{3/2} + \sqrt{x}) dx$$

↓ 2009

$$\frac{2x^{5/2}}{5} + \frac{2x^{3/2}}{3}$$

input `Int[(x + x^2)/Sqrt[x],x]`

output `(2*x^(3/2))/3 + (2*x^(5/2))/5`

Defintions of rubi rules used

rule 9 `Int[(u_.)*(Px_)^(p_.)*((e_.)*(x_))^(m_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Simp[1/e^(p*r) Int[u*(e*x)^(m + p*r)*ExpandToSum[Px/x^r, x]^p, x], x] /; IGtQ[r, 0] /; FreeQ[{e, m}, x] && PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x]`

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0])
|| LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

method	result	size
gospers	$\frac{2x^{\frac{3}{2}}(5+3x)}{15}$	11
trager	$\frac{2x^{\frac{3}{2}}(5+3x)}{15}$	11
risch	$\frac{2x^{\frac{3}{2}}(5+3x)}{15}$	11
derivativedivides	$\frac{2x^{\frac{3}{2}}}{3} + \frac{2x^{\frac{5}{2}}}{5}$	12
default	$\frac{2x^{\frac{3}{2}}}{3} + \frac{2x^{\frac{5}{2}}}{5}$	12
orering	$\frac{2\sqrt{x}(5+3x)(x^2+x)}{15(1+x)}$	21

input `int((x^2+x)/x^(1/2),x,method=_RETURNVERBOSE)`

output `2/15*x^(3/2)*(5+3*x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int \frac{x + x^2}{\sqrt{x}} dx = \frac{2}{15} (3x^2 + 5x)\sqrt{x}$$

input `integrate((x^2+x)/x^(1/2),x, algorithm="fricas")`

output `2/15*(3*x^2 + 5*x)*sqrt(x)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{x + x^2}{\sqrt{x}} dx = \frac{2x^{\frac{5}{2}}}{5} + \frac{2x^{\frac{3}{2}}}{3}$$

input `integrate((x**2+x)/x**(1/2),x)`

output `2*x**(5/2)/5 + 2*x**(3/2)/3`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int \frac{x + x^2}{\sqrt{x}} dx = \frac{2}{5} x^{\frac{5}{2}} + \frac{2}{3} x^{\frac{3}{2}}$$

input `integrate((x^2+x)/x^(1/2),x, algorithm="maxima")`

output `2/5*x^(5/2) + 2/3*x^(3/2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int \frac{x + x^2}{\sqrt{x}} dx = \frac{2}{5} x^{\frac{5}{2}} + \frac{2}{3} x^{\frac{3}{2}}$$

input `integrate((x^2+x)/x^(1/2),x, algorithm="giac")`

output `2/5*x^(5/2) + 2/3*x^(3/2)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.53

$$\int \frac{x + x^2}{\sqrt{x}} dx = \frac{2x^{3/2}(3x + 5)}{15}$$

input `int((x + x^2)/x^(1/2),x)`output `(2*x^(3/2)*(3*x + 5))/15`**Reduce [B] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.53

$$\int \frac{x + x^2}{\sqrt{x}} dx = \frac{2\sqrt{x}x(3x + 5)}{15}$$

input `int((x^2+x)/x^(1/2),x)`output `(2*sqrt(x)*x*(3*x + 5))/15`

3.4 $\int \cos(x) dx$

Optimal result	84
Mathematica [A] (verified)	84
Rubi [A] (verified)	85
Maple [A] (verified)	86
Fricas [A] (verification not implemented)	86
Sympy [A] (verification not implemented)	87
Maxima [A] (verification not implemented)	87
Giac [A] (verification not implemented)	87
Mupad [B] (verification not implemented)	88
Reduce [B] (verification not implemented)	88

Optimal result

Integrand size = 2, antiderivative size = 2

$$\int \cos(x) dx = \sin(x)$$

output `sin(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cos(x) dx = \sin(x)$$

input `Integrate[Cos[x], x]`

output `Sin[x]`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos(x) dx$$

$$\downarrow 3042$$

$$\int \sin\left(x + \frac{\pi}{2}\right) dx$$

$$\downarrow 3117$$

$$\sin(x)$$

input `Int[Cos[x], x]`

output `Sin[x]`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] :> Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
lookup	$\sin(x)$	3
default	$\sin(x)$	3
meijerg	$\sin(x)$	3
risch	$\sin(x)$	3
parallelrisc	$\sin(x)$	3
orering	$\sin(x)$	3
norman	$\frac{2 \tan(\frac{x}{2})}{1 + \tan(\frac{x}{2})^2}$	17

input `int(cos(x),x,method=_RETURNVERBOSE)`

output `sin(x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cos(x) dx = \sin(x)$$

input `integrate(cos(x),x, algorithm="fricas")`

output `sin(x)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cos(x) dx = \sin(x)$$

input `integrate(cos(x),x)`

output `sin(x)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cos(x) dx = \sin(x)$$

input `integrate(cos(x),x, algorithm="maxima")`

output `sin(x)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cos(x) dx = \sin(x)$$

input `integrate(cos(x),x, algorithm="giac")`

output `sin(x)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cos(x) dx = \sin(x)$$

input `int(cos(x),x)`

output `sin(x)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cos(x) dx = \sin(x)$$

input `int(cos(x),x)`

output `sin(x)`

3.5 $\int e^{x^2} x dx$

Optimal result	89
Mathematica [A] (verified)	89
Rubi [A] (verified)	90
Maple [A] (verified)	90
Fricas [A] (verification not implemented)	91
Sympy [A] (verification not implemented)	92
Maxima [A] (verification not implemented)	92
Giac [A] (verification not implemented)	92
Mupad [B] (verification not implemented)	93
Reduce [B] (verification not implemented)	93

Optimal result

Integrand size = 7, antiderivative size = 9

$$\int e^{x^2} x dx = \frac{e^{x^2}}{2}$$

output `1/2*exp(x^2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int e^{x^2} x dx = \frac{e^{x^2}}{2}$$

input `Integrate[E^x^2*x,x]`

output `E^x^2/2`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2638}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{x^2} x dx$$

$$\downarrow 2638$$

$$\frac{e^{x^2}}{2}$$

input `Int [E^x^2*x, x]`

output `E^x^2/2`

Defintions of rubi rules used

rule 2638

```
Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^(n_)))*((e_.) + (f_.)*(x_)^(m_.), x_Symbol] := Simp[(e + f*x)^n*(F^(a + b*(c + d*x)^n)/(b*f*n*(c + d*x)^n *Log[F])), x] /; FreeQ[{F, a, b, c, d, e, f, n}, x] && EqQ[m, n - 1] && EqQ[d*e - c*f, 0]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

method	result	size
gosper	$\frac{e^{x^2}}{2}$	7
derivativedivides	$\frac{e^{x^2}}{2}$	7
default	$\frac{e^{x^2}}{2}$	7
norman	$\frac{e^{x^2}}{2}$	7
risch	$\frac{e^{x^2}}{2}$	7
parallelrisch	$\frac{e^{x^2}}{2}$	7
orering	$\frac{e^{x^2}}{2}$	7
meijerg	$-\frac{1}{2} + \frac{e^{x^2}}{2}$	9
parts	$\frac{\operatorname{erfi}(x)\sqrt{\pi}x}{2} - \frac{\sqrt{\pi}\left(x \operatorname{erfi}(x) - \frac{e^{x^2}}{\sqrt{\pi}}\right)}{2}$	29

input `int(exp(x^2)*x,x,method=_RETURNVERBOSE)`

output `1/2*exp(x^2)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.67

$$\int e^{x^2} x dx = \frac{1}{2} e^{(x^2)}$$

input `integrate(exp(x^2)*x,x, algorithm="fricas")`

output `1/2*e^(x^2)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.56

$$\int e^{x^2} x dx = \frac{e^{x^2}}{2}$$

input `integrate(exp(x**2)*x,x)`

output `exp(x**2)/2`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.67

$$\int e^{x^2} x dx = \frac{1}{2} e^{(x^2)}$$

input `integrate(exp(x^2)*x,x, algorithm="maxima")`

output `1/2*e^(x^2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.67

$$\int e^{x^2} x dx = \frac{1}{2} e^{(x^2)}$$

input `integrate(exp(x^2)*x,x, algorithm="giac")`

output `1/2*e^(x^2)`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.67

$$\int e^{x^2} x dx = \frac{e^{x^2}}{2}$$

input `int(x*exp(x^2),x)`

output `exp(x^2)/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int e^{x^2} x dx = \frac{e^{x^2}}{2}$$

input `int(exp(x^2)*x,x)`

output `e**(x**2)/2`

3.6 $\int \sec^2(x) \tan(x) dx$

Optimal result	94
Mathematica [A] (verified)	94
Rubi [A] (verified)	95
Maple [A] (verified)	96
Fricas [A] (verification not implemented)	96
Sympy [A] (verification not implemented)	97
Maxima [A] (verification not implemented)	97
Giac [A] (verification not implemented)	97
Mupad [B] (verification not implemented)	98
Reduce [B] (verification not implemented)	98

Optimal result

Integrand size = 7, antiderivative size = 8

$$\int \sec^2(x) \tan(x) dx = \frac{\sec^2(x)}{2}$$

output `1/2*sec(x)^2`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \sec^2(x) \tan(x) dx = \frac{\sec^2(x)}{2}$$

input `Integrate[Sec[x]^2*Tan[x],x]`

output `Sec[x]^2/2`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3086, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \tan(x) \sec^2(x) dx \\ \downarrow 3042 \\ \int \tan(x) \sec(x)^2 dx \\ \downarrow 3086 \\ \int \sec(x) d\sec(x) \\ \downarrow 15 \\ \frac{\sec^2(x)}{2} \end{array}$$

input `Int[Sec[x]^2*Tan[x],x]`

output `Sec[x]^2/2`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086

```
Int[((a_.)*sec[(e_.) + (f_.)*(x_.)]^(m_.))*((b_.)*tan[(e_.) + (f_.)*(x_.)]^(n_.), x_Symbol] :> Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])
```

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$\frac{\sec(x)^2}{2}$	7
default	$\frac{\sec(x)^2}{2}$	7
risch	$\frac{2e^{2ix}}{(e^{2ix}+1)^2}$	17

input

```
int(sec(x)^2*tan(x), x, method=_RETURNVERBOSE)
```

output

```
1/2*sec(x)^2
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^2(x) \tan(x) dx = \frac{1}{2 \cos(x)^2}$$

input

```
integrate(sec(x)^2*tan(x), x, algorithm="fricas")
```

output

```
1/2/cos(x)^2
```

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int \sec^2(x) \tan(x) dx = \frac{1}{2 \cos^2(x)}$$

input `integrate(sec(x)**2*tan(x),x)`

output `1/(2*cos(x)**2)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^2(x) \tan(x) dx = \frac{1}{2} \tan(x)^2$$

input `integrate(sec(x)^2*tan(x),x, algorithm="maxima")`

output `1/2*tan(x)^2`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^2(x) \tan(x) dx = \frac{1}{2 \cos(x)^2}$$

input `integrate(sec(x)^2*tan(x),x, algorithm="giac")`

output `1/2/cos(x)^2`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^2(x) \tan(x) dx = \frac{\tan(x)^2}{2}$$

input `int(tan(x)/cos(x)^2,x)`

output `tan(x)^2/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^2(x) \tan(x) dx = \frac{\sec(x)^2}{2}$$

input `int(sec(x)^2*tan(x),x)`

output `sec(x)**2/2`

3.7 $\int x\sqrt{1+x^2} dx$

Optimal result	99
Mathematica [A] (verified)	99
Rubi [A] (verified)	100
Maple [A] (verified)	101
Fricas [A] (verification not implemented)	101
Sympy [B] (verification not implemented)	102
Maxima [A] (verification not implemented)	102
Giac [A] (verification not implemented)	102
Mupad [B] (verification not implemented)	103
Reduce [B] (verification not implemented)	103

Optimal result

Integrand size = 11, antiderivative size = 13

$$\int x\sqrt{1+x^2} dx = \frac{1}{3}(1+x^2)^{3/2}$$

output `1/3*(x^2+1)^(3/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int x\sqrt{1+x^2} dx = \frac{1}{3}(1+x^2)^{3/2}$$

input `Integrate[x*Sqrt[1 + x^2],x]`

output `(1 + x^2)^(3/2)/3`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x\sqrt{x^2+1} dx$$

$$\downarrow 241$$

$$\frac{1}{3}(x^2+1)^{3/2}$$

input

```
Int[x*Sqrt[1 + x^2],x]
```

output

```
(1 + x^2)^(3/2)/3
```

Defintions of rubi rules used

rule 241

```
Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(a + b*x^2)^(p + 1)/
(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

method	result	size
gospers	$\frac{(x^2+1)^{\frac{3}{2}}}{3}$	10
derivativedivides	$\frac{(x^2+1)^{\frac{3}{2}}}{3}$	10
default	$\frac{(x^2+1)^{\frac{3}{2}}}{3}$	10
risch	$\frac{(x^2+1)^{\frac{3}{2}}}{3}$	10
pseudoelliptic	$\frac{(x^2+1)^{\frac{3}{2}}}{3}$	10
orering	$\frac{(x^2+1)^{\frac{3}{2}}}{3}$	10
trager	$\left(\frac{x^2}{3} + \frac{1}{3}\right) \sqrt{x^2 + 1}$	16
meijerg	$-\frac{\frac{4\sqrt{\pi}}{3} - \frac{2\sqrt{\pi}(2x^2+2)\sqrt{x^2+1}}{3}}{4\sqrt{\pi}}$	31

input `int(x*(x^2+1)^(1/2),x,method=_RETURNVERBOSE)`output `1/3*(x^2+1)^(3/2)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int x\sqrt{1+x^2} dx = \frac{1}{3} (x^2 + 1)^{\frac{3}{2}}$$

input `integrate(x*(x^2+1)^(1/2),x, algorithm="fricas")`output `1/3*(x^2 + 1)^(3/2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 22 vs. $2(8) = 16$.

Time = 0.06 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.69

$$\int x\sqrt{1+x^2} dx = \frac{x^2\sqrt{x^2+1}}{3} + \frac{\sqrt{x^2+1}}{3}$$

input `integrate(x*(x**2+1)**(1/2),x)`

output `x**2*sqrt(x**2 + 1)/3 + sqrt(x**2 + 1)/3`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int x\sqrt{1+x^2} dx = \frac{1}{3} (x^2 + 1)^{\frac{3}{2}}$$

input `integrate(x*(x^2+1)^(1/2),x, algorithm="maxima")`

output `1/3*(x^2 + 1)^(3/2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int x\sqrt{1+x^2} dx = \frac{1}{3} (x^2 + 1)^{\frac{3}{2}}$$

input `integrate(x*(x^2+1)^(1/2),x, algorithm="giac")`

output `1/3*(x^2 + 1)^(3/2)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int x\sqrt{1+x^2} dx = \frac{(x^2+1)^{3/2}}{3}$$

input `int(x*(x^2 + 1)^(1/2),x)`

output `(x^2 + 1)^(3/2)/3`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int x\sqrt{1+x^2} dx = \frac{\sqrt{x^2+1}(x^2+1)}{3}$$

input `int(x*(x^2+1)^(1/2),x)`

output `(sqrt(x**2 + 1)*(x**2 + 1))/3`

3.8 $\int e^x \sin(x) dx$

Optimal result	104
Mathematica [A] (verified)	104
Rubi [A] (verified)	105
Maple [A] (verified)	105
Fricas [A] (verification not implemented)	106
Sympy [A] (verification not implemented)	106
Maxima [A] (verification not implemented)	107
Giac [A] (verification not implemented)	107
Mupad [B] (verification not implemented)	107
Reduce [B] (verification not implemented)	108

Optimal result

Integrand size = 6, antiderivative size = 19

$$\int e^x \sin(x) dx = -\frac{1}{2}e^x \cos(x) + \frac{1}{2}e^x \sin(x)$$

output `-1/2*exp(x)*cos(x)+1/2*exp(x)*sin(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int e^x \sin(x) dx = \frac{1}{2}e^x(-\cos(x) + \sin(x))$$

input `Integrate[E^x*Sin[x],x]`

output `(E^x*(-Cos[x] + Sin[x]))/2`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {4932}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^x \sin(x) dx$$

↓ 4932

$$\frac{1}{2}e^x \sin(x) - \frac{1}{2}e^x \cos(x)$$

input

```
Int[E^x*Sin[x],x]
```

output

```
-1/2*(E^x*Cos[x]) + (E^x*Sin[x])/2
```

Defintions of rubi rules used

rule 4932

```
Int[(F_)^((c_.)*((a_.) + (b_.)*(x_)))*Sin[(d_.) + (e_.)*(x_)], x_Symbol] :>
Simp[b*c*Log[F]*F^(c*(a + b*x))*(Sin[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x
] - Simp[e*F^(c*(a + b*x))*(Cos[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] /; F
reeQ[{F, a, b, c, d, e}, x] && NeQ[e^2 + b^2*c^2*Log[F]^2, 0]
```

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

method	result	size
parallelrisch	$\frac{e^x(-\cos(x)+\sin(x))}{2}$	12
default	$-\frac{e^x \cos(x)}{2} + \frac{e^x \sin(x)}{2}$	14
orering	$-\frac{e^x \cos(x)}{2} + \frac{e^x \sin(x)}{2}$	14
norman	$\frac{e^x \tan(\frac{x}{2}) + \frac{e^x \tan(\frac{x}{2})^2}{2} - \frac{e^x}{2}}{1 + \tan(\frac{x}{2})^2}$	34
risch	$-\frac{e^{(1+i)x}}{4} - \frac{ie^{(1+i)x}}{4} - \frac{e^{(1-i)x}}{4} + \frac{ie^{(1-i)x}}{4}$	36

input `int(exp(x)*sin(x),x,method=_RETURNVERBOSE)`

output `1/2*exp(x)*(-cos(x)+sin(x))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.68

$$\int e^x \sin(x) dx = -\frac{1}{2} \cos(x) e^x + \frac{1}{2} e^x \sin(x)$$

input `integrate(exp(x)*sin(x),x, algorithm="fricas")`

output `-1/2*cos(x)*e^x + 1/2*e^x*sin(x)`

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int e^x \sin(x) dx = \frac{e^x \sin(x)}{2} - \frac{e^x \cos(x)}{2}$$

input `integrate(exp(x)*sin(x),x)`

output `exp(x)*sin(x)/2 - exp(x)*cos(x)/2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int e^x \sin(x) dx = -\frac{1}{2} (\cos(x) - \sin(x))e^x$$

input `integrate(exp(x)*sin(x),x, algorithm="maxima")`

output `-1/2*(cos(x) - sin(x))*e^x`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int e^x \sin(x) dx = -\frac{1}{2} (\cos(x) - \sin(x))e^x$$

input `integrate(exp(x)*sin(x),x, algorithm="giac")`

output `-1/2*(cos(x) - sin(x))*e^x`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int e^x \sin(x) dx = -\frac{e^x (\cos(x) - \sin(x))}{2}$$

input `int(exp(x)*sin(x),x)`

output `-(exp(x)*(cos(x) - sin(x)))/2`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int e^x \sin(x) dx = \frac{e^x(-\cos(x) + \sin(x))}{2}$$

input `int(exp(x)*sin(x),x)`

output `(e**x*(- cos(x) + sin(x)))/2`

3.9 $\int \cot(x) \csc^3(x) dx$

Optimal result	109
Mathematica [A] (verified)	109
Rubi [A] (verified)	110
Maple [A] (verified)	111
Fricas [B] (verification not implemented)	112
Sympy [A] (verification not implemented)	112
Maxima [A] (verification not implemented)	112
Giac [A] (verification not implemented)	113
Mupad [B] (verification not implemented)	113
Reduce [B] (verification not implemented)	113

Optimal result

Integrand size = 7, antiderivative size = 8

$$\int \cot(x) \csc^3(x) dx = -\frac{1}{3} \csc^3(x)$$

output `-1/3*csc(x)^3`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \cot(x) \csc^3(x) dx = -\frac{1}{3} \csc^3(x)$$

input `Integrate[Cot[x]*Csc[x]^3,x]`

output `-1/3*Csc[x]^3`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 25, 3086, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot(x) \csc^3(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan\left(x - \frac{\pi}{2}\right) \left(-\sec\left(x - \frac{\pi}{2}\right)\right)^3 dx \\
 & \quad \downarrow \text{25} \\
 & - \int \sec\left(x - \frac{\pi}{2}\right)^3 \tan\left(x - \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{3086} \\
 & - \int \csc^2(x) d \csc(x) \\
 & \quad \downarrow \text{15} \\
 & -\frac{1}{3} \csc^3(x)
 \end{aligned}$$

input `Int [Cot [x] *Csc [x] ^3, x]`

output `-1/3*Csc [x] ^3`

Definitions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)]^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_.)), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$-\frac{\csc(x)^3}{3}$	7
default	$-\frac{\csc(x)^3}{3}$	7
parallelrisc	$-\frac{\sec(\frac{x}{2})^3 \csc(\frac{x}{2})^3}{24}$	15
risc	$\frac{8ie^{3ix}}{3(e^{2ix}-1)^3}$	18
norman	$\frac{-\frac{1}{24} - \frac{\tan(\frac{x}{2})^2}{8} - \frac{\tan(\frac{x}{2})^4}{8} - \frac{\tan(\frac{x}{2})^6}{24}}{\tan(\frac{x}{2})^3}$	34

input `int(cos(x)*csc(x)^2/sin(x)^2,x,method=_RETURNVERBOSE)`

output `-1/3*csc(x)^3`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 14 vs. $2(6) = 12$.

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.75

$$\int \cot(x) \csc^3(x) dx = \frac{1}{3 (\cos(x)^2 - 1) \sin(x)}$$

input `integrate(cos(x)*csc(x)^2/sin(x)^2,x, algorithm="fricas")`

output `1/3/((cos(x)^2 - 1)*sin(x))`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \cot(x) \csc^3(x) dx = -\frac{1}{3 \sin^3(x)}$$

input `integrate(cos(x)*csc(x)**2/sin(x)**2,x)`

output `-1/(3*sin(x)**3)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cot(x) \csc^3(x) dx = -\frac{1}{3 \sin(x)^3}$$

input `integrate(cos(x)*csc(x)^2/sin(x)^2,x, algorithm="maxima")`

output `-1/3/sin(x)^3`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cot(x) \csc^3(x) dx = -\frac{1}{3 \sin(x)^3}$$

input `integrate(cos(x)*csc(x)^2/sin(x)^2,x, algorithm="giac")`

output `-1/3/sin(x)^3`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cot(x) \csc^3(x) dx = -\frac{1}{3 \sin(x)^3}$$

input `int(cos(x)/sin(x)^4,x)`

output `-1/(3*sin(x)^3)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cot(x) \csc^3(x) dx = -\frac{1}{3 \sin(x)^3}$$

input `int(cos(x)*csc(x)^2/sin(x)^2,x)`

output `(- 1)/(3*sin(x)**3)`

3.10 $\int \sin(e^x) dx$

Optimal result	114
Mathematica [A] (verified)	114
Rubi [A] (verified)	115
Maple [A] (verified)	116
Fricas [A] (verification not implemented)	116
Sympy [A] (verification not implemented)	117
Maxima [C] (verification not implemented)	117
Giac [A] (verification not implemented)	117
Mupad [F(-1)]	118
Reduce [F]	118

Optimal result

Integrand size = 4, antiderivative size = 4

$$\int \sin(e^x) dx = \text{Si}(e^x)$$

output `Si(exp(x))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \sin(e^x) dx = \text{Si}(e^x)$$

input `Integrate[Sin[E^x], x]`

output `SinIntegral[E^x]`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {2720, 3042, 3780}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \sin(e^x) dx \\ \downarrow 2720 \\ \int e^{-x} \sin(e^x) de^x \\ \downarrow 3042 \\ \int e^{-x} \sin(e^x) de^x \\ \downarrow 3780 \\ \text{Si}(e^x) \end{array}$$

input `Int[Sin[E^x],x]`

output `SinIntegral[E^x]`

Defintions of rubi rules used

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 3780

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[SinInte
gral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]
```

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

method	result	size
derivativedivides	$\text{Si}(e^x)$	4
default	$\text{Si}(e^x)$	4
risch	$-\frac{\pi \operatorname{csgn}(e^x)}{2} + \text{Si}(e^x)$	11

```
input int(sin(exp(x)),x,method=_RETURNVERBOSE)
```

```
output Si(exp(x))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int \sin(e^x) dx = \text{Si}(e^x)$$

```
input integrate(sin(exp(x)),x, algorithm="fricas")
```

```
output sin_integral(e^x)
```

Sympy [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int \sin(e^x) dx = \text{Si}(e^x)$$

input `integrate(sin(exp(x)),x)`

output `Si(exp(x))`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.06 (sec) , antiderivative size = 15, normalized size of antiderivative = 3.75

$$\int \sin(e^x) dx = -\frac{1}{2}i \text{Ei}(i e^x) + \frac{1}{2}i \text{Ei}(-i e^x)$$

input `integrate(sin(exp(x)),x, algorithm="maxima")`

output `-1/2*I*Ei(I*e^x) + 1/2*I*Ei(-I*e^x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int \sin(e^x) dx = \text{Si}(e^x)$$

input `integrate(sin(exp(x)),x, algorithm="giac")`

output `sin_integral(e^x)`

Mupad [F(-1)]

Timed out.

$$\int \sin(e^x) dx = \text{sinint}(e^x)$$

input `int(sin(exp(x)),x)`output `sinint(exp(x))`**Reduce [F]**

$$\int \sin(e^x) dx = \int \sin(e^x) dx$$

input `int(sin(exp(x)),x)`output `int(sin(e**x),x)`

3.11 $\int \frac{\sin(y)}{y} dy$

Optimal result	119
Mathematica [A] (verified)	119
Rubi [A] (verified)	120
Maple [A] (verified)	121
Fricas [A] (verification not implemented)	121
Sympy [A] (verification not implemented)	121
Maxima [C] (verification not implemented)	122
Giac [A] (verification not implemented)	122
Mupad [F(-1)]	122
Reduce [B] (verification not implemented)	123

Optimal result

Integrand size = 6, antiderivative size = 2

$$\int \frac{\sin(y)}{y} dy = \text{Si}(y)$$

output

Si(y)

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{\sin(y)}{y} dy = \text{Si}(y)$$

input

Integrate[Sin[y]/y,y]

output

SinIntegral[y]

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3780}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sin(y)}{y} dy$$

↓ 3042

$$\int \frac{\sin(y)}{y} dy$$

↓ 3780

$$\text{Si}(y)$$

input

```
Int[Sin[y]/y,y]
```

output

```
SinIntegral[y]
```

Defintions of rubi rules used

rule 3042

```
Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 3780

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]
```

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
default	$\text{Si}(y)$	3
meijerg	$\text{Si}(y)$	3
risch	$-\frac{\pi \operatorname{csgn}(y)}{2} + \text{Si}(y)$	9

input `int(sin(y)/y,y,method=_RETURNVERBOSE)`

output `Si(y)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{\sin(y)}{y} dy = \text{Si}(y)$$

input `integrate(sin(y)/y,y, algorithm="fricas")`

output `sin_integral(y)`

Sympy [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{\sin(y)}{y} dy = \text{Si}(y)$$

input `integrate(sin(y)/y,y)`

output `Si(y)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.06 (sec) , antiderivative size = 13, normalized size of antiderivative = 6.50

$$\int \frac{\sin(y)}{y} dy = -\frac{1}{2}i \operatorname{Ei}(iy) + \frac{1}{2}i \operatorname{Ei}(-iy)$$

input `integrate(sin(y)/y,y, algorithm="maxima")`

output `-1/2*I*Ei(I*y) + 1/2*I*Ei(-I*y)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{\sin(y)}{y} dy = \operatorname{Si}(y)$$

input `integrate(sin(y)/y,y, algorithm="giac")`

output `sin_integral(y)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sin(y)}{y} dy = \operatorname{sinint}(y)$$

input `int(sin(y)/y,y)`

output `sinint(y)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{\sin(y)}{y} dy = si(y)$$

input `int(sin(y)/y,y)`

output `si(y)`

3.12 $\int (e^x + \sin(x)) dx$

Optimal result	124
Mathematica [A] (verified)	124
Rubi [A] (verified)	125
Maple [A] (verified)	126
Fricas [A] (verification not implemented)	126
Sympy [A] (verification not implemented)	127
Maxima [A] (verification not implemented)	127
Giac [A] (verification not implemented)	127
Mupad [B] (verification not implemented)	128
Reduce [B] (verification not implemented)	128

Optimal result

Integrand size = 6, antiderivative size = 8

$$\int (e^x + \sin(x)) dx = e^x - \cos(x)$$

output `exp(x)-cos(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int (e^x + \sin(x)) dx = e^x - \cos(x)$$

input `Integrate[E^x + Sin[x],x]`

output `E^x - Cos[x]`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e^x + \sin(x)) dx$$

↓ 2009

$$e^x - \cos(x)$$

input `Int[E^x + Sin[x], x]`

output `E^x - Cos[x]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

method	result	size
default	$e^x - \cos(x)$	8
risch	$e^x - \cos(x)$	8
parts	$e^x - \cos(x)$	8
orering	$e^x - \cos(x)$	8
parallelrisc	$e^x - \cos(x) - 1$	9
meijerg	$-1 + e^x + \sqrt{\pi} \left(\frac{1}{\sqrt{\pi}} - \frac{\cos(x)}{\sqrt{\pi}} \right)$	20
norman	$\frac{e^x \tan(\frac{x}{2})^2 - 2 + e^x}{1 + \tan(\frac{x}{2})^2}$	25

input `int(exp(x)+sin(x),x,method=_RETURNVERBOSE)`output `exp(x)-cos(x)`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int (e^x + \sin(x)) dx = -\cos(x) + e^x$$

input `integrate(exp(x)+sin(x),x, algorithm="fricas")`output `-cos(x) + e^x`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int (e^x + \sin(x)) dx = e^x - \cos(x)$$

input `integrate(exp(x)+sin(x),x)`

output `exp(x) - cos(x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int (e^x + \sin(x)) dx = -\cos(x) + e^x$$

input `integrate(exp(x)+sin(x),x, algorithm="maxima")`

output `-cos(x) + e^x`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int (e^x + \sin(x)) dx = -\cos(x) + e^x$$

input `integrate(exp(x)+sin(x),x, algorithm="giac")`

output `-cos(x) + e^x`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int (e^x + \sin(x)) dx = e^x - \cos(x)$$

input `int(exp(x) + sin(x),x)`

output `exp(x) - cos(x)`

Reduce [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int (e^x + \sin(x)) dx = -\cos(x) + e^x$$

input `int(exp(x)+sin(x),x)`

output `- cos(x) + e**x`

3.13 $\int \left(e^{x^2} + 2e^{x^2} x^2 \right) dx$

Optimal result	129
Mathematica [A] (verified)	129
Rubi [A] (verified)	130
Maple [A] (verified)	131
Fricas [A] (verification not implemented)	131
Sympy [A] (verification not implemented)	132
Maxima [A] (verification not implemented)	132
Giac [A] (verification not implemented)	132
Mupad [B] (verification not implemented)	133
Reduce [B] (verification not implemented)	133

Optimal result

Integrand size = 16, antiderivative size = 7

$$\int \left(e^{x^2} + 2e^{x^2} x^2 \right) dx = e^{x^2} x$$

output `exp(x^2)*x`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \left(e^{x^2} + 2e^{x^2} x^2 \right) dx = e^{x^2} x$$

input `Integrate[E^x^2 + 2*E^x^2*x^2,x]`

output `E^x^2*x`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.062$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (2e^{x^2}x^2 + e^{x^2}) dx$$

$$\downarrow \text{2009}$$

$$e^{x^2}x$$

input `Int[E^x^2 + 2*E^x^2*x^2,x]`

output `E^x^2*x`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

method	result	size
gospers	$e^{x^2} x$	7
default	$e^{x^2} x$	7
norman	$e^{x^2} x$	7
risch	$e^{x^2} x$	7
parallelrisch	$e^{x^2} x$	7
orering	$\frac{x(2e^{x^2}x^2 + e^{x^2})}{2x^2 + 1}$	26
meijerg	$i\left(-ie^{x^2}x + \frac{i\operatorname{erfi}(x)\sqrt{\pi}}{2}\right) + \frac{\operatorname{erfi}(x)\sqrt{\pi}}{2}$	29
parts	$\operatorname{erfi}(x)\sqrt{\pi}x^2 + \frac{\operatorname{erfi}(x)\sqrt{\pi}}{2} - 2\sqrt{\pi}\left(\frac{x^2\operatorname{erfi}(x)}{2} - \frac{\frac{e^{x^2}x - \operatorname{erfi}(x)\sqrt{\pi}}{4}}{\sqrt{\pi}}\right)$	51

input `int(2*exp(x^2)*x^2+exp(x^2),x,method=_RETURNVERBOSE)`output `exp(x^2)*x`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int (e^{x^2} + 2e^{x^2}x^2) dx = xe^{(x^2)}$$

input `integrate(exp(x^2)+2*exp(x^2)*x^2,x, algorithm="fricas")`output `x*e^(x^2)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.71

$$\int \left(e^{x^2} + 2e^{x^2} x^2 \right) dx = x e^{x^2}$$

input `integrate(exp(x**2)+2*exp(x**2)*x**2,x)`

output `x*exp(x**2)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int \left(e^{x^2} + 2e^{x^2} x^2 \right) dx = x e^{(x^2)}$$

input `integrate(exp(x^2)+2*exp(x^2)*x^2,x, algorithm="maxima")`

output `x*e^(x^2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int \left(e^{x^2} + 2e^{x^2} x^2 \right) dx = x e^{(x^2)}$$

input `integrate(exp(x^2)+2*exp(x^2)*x^2,x, algorithm="giac")`

output `x*e^(x^2)`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int \left(e^{x^2} + 2e^{x^2} x^2 \right) dx = x e^{x^2}$$

input `int(exp(x^2) + 2*x^2*exp(x^2),x)`

output `x*exp(x^2)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \left(e^{x^2} + 2e^{x^2} x^2 \right) dx = e^{x^2} x$$

input `int(exp(x^2)+2*exp(x^2)*x^2,x)`

output `e**(x**2)*x`

3.14 $\int (e^x + x)^2 dx$

Optimal result	134
Mathematica [A] (verified)	134
Rubi [A] (verified)	135
Maple [A] (verified)	136
Fricas [A] (verification not implemented)	136
Sympy [A] (verification not implemented)	137
Maxima [A] (verification not implemented)	137
Giac [A] (verification not implemented)	137
Mupad [B] (verification not implemented)	138
Reduce [B] (verification not implemented)	138

Optimal result

Integrand size = 7, antiderivative size = 28

$$\int (e^x + x)^2 dx = -2e^x + \frac{e^{2x}}{2} + 2e^x x + \frac{x^3}{3}$$

output

```
-2*exp(x)+1/2*exp(2*x)+2*exp(x)*x+1/3*x^3
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int (e^x + x)^2 dx = \frac{e^{2x}}{2} + \frac{x^3}{3} + e^x(-2 + 2x)$$

input

```
Integrate[(E^x + x)^2, x]
```

output

```
E^(2*x)/2 + x^3/3 + E^x*(-2 + 2*x)
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (x + e^x)^2 dx$$

$$\downarrow 7293$$

$$\int (x^2 + 2e^x x + e^{2x}) dx$$

$$\downarrow 2009$$

$$\frac{x^3}{3} + 2e^x x - 2e^x + \frac{e^{2x}}{2}$$

input

```
Int[(E^x + x)^2, x]
```

output

```
-2*E^x + E^(2*x)/2 + 2*E^x*x + x^3/3
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 7293

```
Int[u_, x_Symbol] :> With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]
```


Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.75

method	result
risch	$\frac{x^3}{3} + (-2 + 2x)e^x + \frac{e^{2x}}{2}$
default	$-2e^x + \frac{e^{2x}}{2} + 2e^x x + \frac{x^3}{3}$
norman	$-2e^x + \frac{e^{2x}}{2} + 2e^x x + \frac{x^3}{3}$
parallelrisch	$-2e^x + \frac{e^{2x}}{2} + 2e^x x + \frac{x^3}{3}$
parts	$-2e^x + \frac{e^{2x}}{2} + 2e^x x + \frac{x^3}{3}$
orering	$\frac{(2x^4 - 21x^2 + 39x - 15)(e^x + x)^2}{6(-1+x)^3} - \frac{(6x^4 - 25x^3 + 30x^2 + 6x - 12)(e^x + x)(1 + e^x)}{6(-1+x)^3} + \frac{x(2x^2 - 9x + 12)(2(1 + e^x)^2 + 2(e^x + x)e^x)}{12(-1+x)^2}$

input `int((exp(x)+x)^2,x,method=_RETURNVERBOSE)`output `1/3*x^3+(-2+2*x)*exp(x)+1/2*exp(2*x)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.68

$$\int (e^x + x)^2 dx = \frac{1}{3} x^3 + 2(x - 1)e^x + \frac{1}{2} e^{(2x)}$$

input `integrate((exp(x)+x)^2,x, algorithm="fricas")`output `1/3*x^3 + 2*(x - 1)*e^x + 1/2*e^(2*x)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.71

$$\int (e^x + x)^2 dx = \frac{x^3}{3} + \frac{(4x - 4)e^x}{2} + \frac{e^{2x}}{2}$$

input `integrate((exp(x)+x)**2,x)`

output `x**3/3 + (4*x - 4)*exp(x)/2 + exp(2*x)/2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.68

$$\int (e^x + x)^2 dx = \frac{1}{3} x^3 + 2(x - 1)e^x + \frac{1}{2} e^{(2x)}$$

input `integrate((exp(x)+x)^2,x, algorithm="maxima")`

output `1/3*x^3 + 2*(x - 1)*e^x + 1/2*e^(2*x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.68

$$\int (e^x + x)^2 dx = \frac{1}{3} x^3 + 2(x - 1)e^x + \frac{1}{2} e^{(2x)}$$

input `integrate((exp(x)+x)^2,x, algorithm="giac")`

output `1/3*x^3 + 2*(x - 1)*e^x + 1/2*e^(2*x)`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.75

$$\int (e^x + x)^2 dx = \frac{e^{2x}}{2} - 2e^x + 2xe^x + \frac{x^3}{3}$$

input `int((x + exp(x))^2,x)`

output `exp(2*x)/2 - 2*exp(x) + 2*x*exp(x) + x^3/3`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.86

$$\int (e^x + x)^2 dx = \frac{e^{2x}}{2} + 2e^x x - 2e^x + \frac{x^3}{3}$$

input `int((exp(x)+x)^2,x)`

output `(3*e**(2*x) + 12*e**x*x - 12*e**x + 2*x**3)/6`

3.15 $\int (2e^x + e^{2x} + x^2) dx$

Optimal result	139
Mathematica [A] (verified)	139
Rubi [A] (verified)	140
Maple [A] (verified)	141
Fricas [A] (verification not implemented)	141
Sympy [A] (verification not implemented)	142
Maxima [A] (verification not implemented)	142
Giac [A] (verification not implemented)	142
Mupad [B] (verification not implemented)	143
Reduce [B] (verification not implemented)	143

Optimal result

Integrand size = 14, antiderivative size = 22

$$\int (2e^x + e^{2x} + x^2) dx = 2e^x + \frac{e^{2x}}{2} + \frac{x^3}{3}$$

output `2*exp(x)+1/2*exp(2*x)+1/3*x^3`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int (2e^x + e^{2x} + x^2) dx = 2e^x + \frac{e^{2x}}{2} + \frac{x^3}{3}$$

input `Integrate[2*E^x + E^(2*x) + x^2,x]`

output `2*E^x + E^(2*x)/2 + x^3/3`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (x^2 + 2e^x + e^{2x}) dx$$

$$\downarrow \text{2009}$$

$$\frac{x^3}{3} + 2e^x + \frac{e^{2x}}{2}$$

input `Int [2*E^x + E^(2*x) + x^2,x]`

output `2*E^x + E^(2*x)/2 + x^3/3`

Defintions of rubi rules used

rule 2009 `Int [u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

method	result	size
default	$2e^x + \frac{e^{2x}}{2} + \frac{x^3}{3}$	17
norman	$2e^x + \frac{e^{2x}}{2} + \frac{x^3}{3}$	17
risch	$2e^x + \frac{e^{2x}}{2} + \frac{x^3}{3}$	17
parallelrisch	$2e^x + \frac{e^{2x}}{2} + \frac{x^3}{3}$	17
parts	$2e^x + \frac{e^{2x}}{2} + \frac{x^3}{3}$	17
orering	$\frac{(2x^3-21x+9)(2e^x+e^{2x}+x^2)}{6x^2-18x+6} - \frac{(2x^3-7x^2+2)(2e^x+2e^{2x}+2x)}{4(x^2-3x+1)} + \frac{x(2x^2-9x+6)(2e^x+4e^{2x}+2)}{12x^2-36x+12}$	109

input `int(2*exp(x)+exp(2*x)+x^2,x,method=_RETURNVERBOSE)`output `2*exp(x)+1/2*exp(2*x)+1/3*x^3`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int (2e^x + e^{2x} + x^2) dx = \frac{1}{3}x^3 + \frac{1}{2}e^{(2x)} + 2e^x$$

input `integrate(2*exp(x)+exp(2*x)+x^2,x, algorithm="fricas")`output `1/3*x^3 + 1/2*e^(2*x) + 2*e^x`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.68

$$\int (2e^x + e^{2x} + x^2) dx = \frac{x^3}{3} + \frac{e^{2x}}{2} + 2e^x$$

input `integrate(2*exp(x)+exp(2*x)+x**2,x)`output `x**3/3 + exp(2*x)/2 + 2*exp(x)`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int (2e^x + e^{2x} + x^2) dx = \frac{1}{3} x^3 + \frac{1}{2} e^{(2x)} + 2 e^x$$

input `integrate(2*exp(x)+exp(2*x)+x^2,x, algorithm="maxima")`output `1/3*x^3 + 1/2*e^(2*x) + 2*e^x`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int (2e^x + e^{2x} + x^2) dx = \frac{1}{3} x^3 + \frac{1}{2} e^{(2x)} + 2 e^x$$

input `integrate(2*exp(x)+exp(2*x)+x^2,x, algorithm="giac")`output `1/3*x^3 + 1/2*e^(2*x) + 2*e^x`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int (2e^x + e^{2x} + x^2) dx = \frac{e^{2x}}{2} + 2e^x + \frac{x^3}{3}$$

input `int(exp(2*x) + 2*exp(x) + x^2,x)`

output `exp(2*x)/2 + 2*exp(x) + x^3/3`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int (2e^x + e^{2x} + x^2) dx = \frac{e^{2x}}{2} + 2e^x + \frac{x^3}{3}$$

input `int(2*exp(x)+exp(2*x)+x^2,x)`

output `(3*e**(2*x) + 12*e**x + 2*x**3)/6`

3.16 $\int \cos(x) \sin(x) dx$

Optimal result	144
Mathematica [A] (verified)	144
Rubi [A] (verified)	145
Maple [A] (verified)	146
Fricas [A] (verification not implemented)	146
Sympy [A] (verification not implemented)	147
Maxima [A] (verification not implemented)	147
Giac [A] (verification not implemented)	147
Mupad [B] (verification not implemented)	148
Reduce [B] (verification not implemented)	148

Optimal result

Integrand size = 5, antiderivative size = 8

$$\int \cos(x) \sin(x) dx = \frac{\sin^2(x)}{2}$$

output `1/2*sin(x)^2`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \cos(x) \sin(x) dx = -\frac{1}{2} \cos^2(x)$$

input `Integrate[Cos[x]*Sin[x],x]`

output `-1/2*Cos[x]^2`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3042, 3044, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \sin(x) \cos(x) dx \\ \downarrow \text{3042} \\ \int \sin(x) \cos(x) dx \\ \downarrow \text{3044} \\ \int \sin(x) d \sin(x) \\ \downarrow \text{15} \\ \frac{\sin^2(x)}{2} \end{array}$$

input `Int[Cos[x]*Sin[x],x]`

output `Sin[x]^2/2`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3044

```
Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)]^(m_.), x_
Symbol] := Simp[1/(a*f) Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a
*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(I
ntegerQ[(m - 1)/2] && LtQ[0, m, n])
```

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$\frac{\sin(x)^2}{2}$	7
default	$\frac{\sin(x)^2}{2}$	7
risch	$-\frac{\cos(2x)}{4}$	7
parallelrisch	$\frac{1}{4} - \frac{\cos(2x)}{4}$	9
orering	$-\frac{\cos(x)^2}{4} + \frac{\sin(x)^2}{4}$	14
norman	$\frac{2 \tan(\frac{x}{2})^2}{(1 + \tan(\frac{x}{2})^2)^2}$	19
meijerg	$\frac{\sqrt{\pi} \left(\frac{1}{\sqrt{\pi}} - \frac{\cos(2x)}{\sqrt{\pi}} \right)}{4}$	19

input

```
int(cos(x)*sin(x), x, method=_RETURNVERBOSE)
```

output

```
1/2*sin(x)^2
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos(x) \sin(x) dx = -\frac{1}{2} \cos(x)^2$$

input

```
integrate(cos(x)*sin(x), x, algorithm="fricas")
```

output `-1/2*cos(x)^2`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int \cos(x) \sin(x) dx = \frac{\sin^2(x)}{2}$$

input `integrate(cos(x)*sin(x),x)`

output `sin(x)**2/2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos(x) \sin(x) dx = -\frac{1}{2} \cos(x)^2$$

input `integrate(cos(x)*sin(x),x, algorithm="maxima")`

output `-1/2*cos(x)^2`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos(x) \sin(x) dx = -\frac{1}{2} \cos(x)^2$$

input `integrate(cos(x)*sin(x),x, algorithm="giac")`

output `-1/2*cos(x)^2`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos(x) \sin(x) dx = \frac{\sin(x)^2}{2}$$

input `int(cos(x)*sin(x),x)`

output `sin(x)^2/2`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos(x) \sin(x) dx = -\frac{\cos(x)^2}{2}$$

input `int(cos(x)*sin(x),x)`

output `(- cos(x)**2)/2`

3.17 $\int e^{x^2} x dx$

Optimal result	149
Mathematica [A] (verified)	149
Rubi [A] (verified)	150
Maple [A] (verified)	150
Fricas [A] (verification not implemented)	151
Sympy [A] (verification not implemented)	152
Maxima [A] (verification not implemented)	152
Giac [A] (verification not implemented)	152
Mupad [B] (verification not implemented)	153
Reduce [B] (verification not implemented)	153

Optimal result

Integrand size = 7, antiderivative size = 9

$$\int e^{x^2} x dx = \frac{e^{x^2}}{2}$$

output `1/2*exp(x^2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int e^{x^2} x dx = \frac{e^{x^2}}{2}$$

input `Integrate[E^x^2*x,x]`

output `E^x^2/2`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2638}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{x^2} x dx$$

$$\downarrow 2638$$

$$\frac{e^{x^2}}{2}$$

input `Int [E^x^2*x, x]`

output `E^x^2/2`

Defintions of rubi rules used

rule 2638

```
Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^(n_)))*((e_.) + (f_.)*(x_)^(m_.), x_Symbol] := Simp[(e + f*x)^n*(F^(a + b*(c + d*x)^n)/(b*f*n*(c + d*x)^n *Log[F])), x] /; FreeQ[{F, a, b, c, d, e, f, n}, x] && EqQ[m, n - 1] && EqQ[d*e - c*f, 0]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

method	result	size
gosper	$\frac{e^{x^2}}{2}$	7
derivativeldivides	$\frac{e^{x^2}}{2}$	7
default	$\frac{e^{x^2}}{2}$	7
norman	$\frac{e^{x^2}}{2}$	7
risch	$\frac{e^{x^2}}{2}$	7
parallelrisch	$\frac{e^{x^2}}{2}$	7
orering	$\frac{e^{x^2}}{2}$	7
meijerg	$-\frac{1}{2} + \frac{e^{x^2}}{2}$	9
parts	$\frac{\operatorname{erfi}(x)\sqrt{\pi}x}{2} - \frac{\sqrt{\pi}\left(x \operatorname{erfi}(x) - \frac{e^{x^2}}{\sqrt{\pi}}\right)}{2}$	29

input `int(exp(x^2)*x,x,method=_RETURNVERBOSE)`

output `1/2*exp(x^2)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.67

$$\int e^{x^2} x dx = \frac{1}{2} e^{(x^2)}$$

input `integrate(exp(x^2)*x,x, algorithm="fricas")`

output `1/2*e^(x^2)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.56

$$\int e^{x^2} x dx = \frac{e^{x^2}}{2}$$

input `integrate(exp(x**2)*x,x)`

output `exp(x**2)/2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.67

$$\int e^{x^2} x dx = \frac{1}{2} e^{(x^2)}$$

input `integrate(exp(x^2)*x,x, algorithm="maxima")`

output `1/2*e^(x^2)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.67

$$\int e^{x^2} x dx = \frac{1}{2} e^{(x^2)}$$

input `integrate(exp(x^2)*x,x, algorithm="giac")`

output `1/2*e^(x^2)`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.67

$$\int e^{x^2} x dx = \frac{e^{x^2}}{2}$$

input `int(x*exp(x^2),x)`

output `exp(x^2)/2`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int e^{x^2} x dx = \frac{e^{x^2}}{2}$$

input `int(exp(x^2)*x,x)`

output `e**(x**2)/2`

3.18 $\int x\sqrt{1+x^2} dx$

Optimal result	154
Mathematica [A] (verified)	154
Rubi [A] (verified)	155
Maple [A] (verified)	156
Fricas [A] (verification not implemented)	156
Sympy [B] (verification not implemented)	157
Maxima [A] (verification not implemented)	157
Giac [A] (verification not implemented)	157
Mupad [B] (verification not implemented)	158
Reduce [B] (verification not implemented)	158

Optimal result

Integrand size = 11, antiderivative size = 13

$$\int x\sqrt{1+x^2} dx = \frac{1}{3}(1+x^2)^{3/2}$$

output `1/3*(x^2+1)^(3/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int x\sqrt{1+x^2} dx = \frac{1}{3}(1+x^2)^{3/2}$$

input `Integrate[x*Sqrt[1 + x^2],x]`

output `(1 + x^2)^(3/2)/3`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x\sqrt{x^2+1} dx$$

$$\downarrow 241$$

$$\frac{1}{3}(x^2+1)^{3/2}$$

input

```
Int[x*Sqrt[1 + x^2],x]
```

output

```
(1 + x^2)^(3/2)/3
```

Defintions of rubi rules used

rule 241

```
Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(a + b*x^2)^(p + 1)/
(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

method	result	size
gospers	$\frac{(x^2+1)^{\frac{3}{2}}}{3}$	10
derivativedivides	$\frac{(x^2+1)^{\frac{3}{2}}}{3}$	10
default	$\frac{(x^2+1)^{\frac{3}{2}}}{3}$	10
risch	$\frac{(x^2+1)^{\frac{3}{2}}}{3}$	10
pseudoelliptic	$\frac{(x^2+1)^{\frac{3}{2}}}{3}$	10
orering	$\frac{(x^2+1)^{\frac{3}{2}}}{3}$	10
trager	$\left(\frac{x^2}{3} + \frac{1}{3}\right) \sqrt{x^2 + 1}$	16
meijerg	$-\frac{\frac{4\sqrt{\pi}}{3} - \frac{2\sqrt{\pi}(2x^2+2)\sqrt{x^2+1}}{3}}{4\sqrt{\pi}}$	31

input `int(x*(x^2+1)^(1/2),x,method=_RETURNVERBOSE)`output `1/3*(x^2+1)^(3/2)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int x\sqrt{1+x^2} dx = \frac{1}{3}(x^2+1)^{\frac{3}{2}}$$

input `integrate(x*(x^2+1)^(1/2),x, algorithm="fricas")`output `1/3*(x^2 + 1)^(3/2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 22 vs. $2(8) = 16$.

Time = 0.08 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.69

$$\int x\sqrt{1+x^2} dx = \frac{x^2\sqrt{x^2+1}}{3} + \frac{\sqrt{x^2+1}}{3}$$

input `integrate(x*(x**2+1)**(1/2),x)`

output `x**2*sqrt(x**2 + 1)/3 + sqrt(x**2 + 1)/3`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int x\sqrt{1+x^2} dx = \frac{1}{3} (x^2 + 1)^{\frac{3}{2}}$$

input `integrate(x*(x^2+1)^(1/2),x, algorithm="maxima")`

output `1/3*(x^2 + 1)^(3/2)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int x\sqrt{1+x^2} dx = \frac{1}{3} (x^2 + 1)^{\frac{3}{2}}$$

input `integrate(x*(x^2+1)^(1/2),x, algorithm="giac")`

output `1/3*(x^2 + 1)^(3/2)`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int x\sqrt{1+x^2} dx = \frac{(x^2+1)^{3/2}}{3}$$

input `int(x*(x^2 + 1)^(1/2),x)`

output `(x^2 + 1)^(3/2)/3`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int x\sqrt{1+x^2} dx = \frac{\sqrt{x^2+1}(x^2+1)}{3}$$

input `int(x*(x^2+1)^(1/2),x)`

output `(sqrt(x**2 + 1)*(x**2 + 1))/3`

3.19 $\int \frac{e^x}{1+e^x} dx$

Optimal result	159
Mathematica [A] (verified)	159
Rubi [A] (verified)	160
Maple [A] (verified)	161
Fricas [A] (verification not implemented)	161
Sympy [A] (verification not implemented)	161
Maxima [A] (verification not implemented)	162
Giac [A] (verification not implemented)	162
Mupad [B] (verification not implemented)	162
Reduce [B] (verification not implemented)	163

Optimal result

Integrand size = 11, antiderivative size = 6

$$\int \frac{e^x}{1+e^x} dx = \log(1+e^x)$$

output `ln(1+exp(x))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{e^x}{1+e^x} dx = \log(1+e^x)$$

input `Integrate[E^x/(1 + E^x),x]`

output `Log[1 + E^x]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2676, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^x}{e^x + 1} dx$$

↓ 2676

$$\int \frac{1}{e^x + 1} de^x$$

↓ 16

$$\log(e^x + 1)$$

input `Int[E^x/(1 + E^x),x]`

output `Log[1 + E^x]`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 2676 `Int[((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)*((a_) + (b_.)*(F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.))^(p_.), x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[(a + b*x)^p, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n, p}, x]`

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

method	result	size
derivativedivides	$\ln(1 + e^x)$	6
default	$\ln(1 + e^x)$	6
norman	$\ln(1 + e^x)$	6
risch	$\ln(1 + e^x)$	6
parallelrisch	$\ln(1 + e^x)$	6

input `int(1/(1+exp(x))*exp(x),x,method=_RETURNVERBOSE)`

output `ln(1+exp(x))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int \frac{e^x}{1 + e^x} dx = \log(e^x + 1)$$

input `integrate(exp(x)/(1+exp(x)),x, algorithm="fricas")`

output `log(e^x + 1)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int \frac{e^x}{1 + e^x} dx = \log(e^x + 1)$$

input `integrate(exp(x)/(1+exp(x)),x)`

output `log(exp(x) + 1)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int \frac{e^x}{1 + e^x} dx = \log(e^x + 1)$$

input `integrate(exp(x)/(1+exp(x)),x, algorithm="maxima")`

output `log(e^x + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int \frac{e^x}{1 + e^x} dx = \log(e^x + 1)$$

input `integrate(exp(x)/(1+exp(x)),x, algorithm="giac")`

output `log(e^x + 1)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int \frac{e^x}{1 + e^x} dx = \ln(e^x + 1)$$

input `int(exp(x)/(exp(x) + 1),x)`

output `log(exp(x) + 1)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{e^x}{1 + e^x} dx = \log(e^x + 1)$$

input `int(exp(x)/(1+exp(x)), x)`

output `log(e**x + 1)`

3.20 $\int x^{3/2} dx$

Optimal result	164
Mathematica [A] (verified)	164
Rubi [A] (verified)	165
Maple [A] (verified)	166
Fricas [A] (verification not implemented)	166
Sympy [A] (verification not implemented)	167
Maxima [A] (verification not implemented)	167
Giac [A] (verification not implemented)	167
Mupad [B] (verification not implemented)	168
Reduce [B] (verification not implemented)	168

Optimal result

Integrand size = 5, antiderivative size = 9

$$\int x^{3/2} dx = \frac{2x^{5/2}}{5}$$

output `2/5*x^(5/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int x^{3/2} dx = \frac{2x^{5/2}}{5}$$

input `Integrate[x^(3/2),x]`

output `(2*x^(5/2))/5`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{3/2} dx$$

↓ 15

$$\frac{2x^{5/2}}{5}$$

input `Int[x^(3/2),x]`

output `(2*x^(5/2))/5`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.67

method	result	size
gospers	$\frac{2x^{\frac{5}{2}}}{5}$	6
derivativedivides	$\frac{2x^{\frac{5}{2}}}{5}$	6
default	$\frac{2x^{\frac{5}{2}}}{5}$	6
trager	$\frac{2x^{\frac{5}{2}}}{5}$	6
risch	$\frac{2x^{\frac{5}{2}}}{5}$	6
orering	$\frac{2x^{\frac{5}{2}}}{5}$	6

input `int(x^(3/2),x,method=_RETURNVERBOSE)`

output `2/5*x^(5/2)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.56

$$\int x^{3/2} dx = \frac{2}{5} x^{5/2}$$

input `integrate(x^(3/2),x, algorithm="fricas")`

output `2/5*x^(5/2)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int x^{3/2} dx = \frac{2x^{5/2}}{5}$$

input `integrate(x**(3/2),x)`

output `2*x**(5/2)/5`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.56

$$\int x^{3/2} dx = \frac{2}{5} x^{5/2}$$

input `integrate(x^(3/2),x, algorithm="maxima")`

output `2/5*x^(5/2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.56

$$\int x^{3/2} dx = \frac{2}{5} x^{5/2}$$

input `integrate(x^(3/2),x, algorithm="giac")`

output `2/5*x^(5/2)`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.56

$$\int x^{3/2} dx = \frac{2x^{5/2}}{5}$$

input `int(x^(3/2),x)`

output `(2*x^(5/2))/5`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int x^{3/2} dx = \frac{2\sqrt{x} x^2}{5}$$

input `int(x^(3/2),x)`

output `(2*sqrt(x)*x**2)/5`

3.21 $\int \cos(3 + 2x) dx$

Optimal result	169
Mathematica [A] (verified)	169
Rubi [A] (verified)	170
Maple [A] (verified)	171
Fricas [A] (verification not implemented)	171
Sympy [A] (verification not implemented)	172
Maxima [A] (verification not implemented)	172
Giac [A] (verification not implemented)	172
Mupad [B] (verification not implemented)	173
Reduce [B] (verification not implemented)	173

Optimal result

Integrand size = 6, antiderivative size = 10

$$\int \cos(3 + 2x) dx = \frac{1}{2} \sin(3 + 2x)$$

output `1/2*sin(3+2*x)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \cos(3 + 2x) dx = \frac{1}{2} \sin(3 + 2x)$$

input `Integrate[Cos[3 + 2*x],x]`

output `Sin[3 + 2*x]/2`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos(2x + 3) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(2x + \frac{\pi}{2} + 3\right) dx$$

$$\downarrow \text{3117}$$

$$\frac{1}{2} \sin(2x + 3)$$

input `Int[Cos[3 + 2*x], x]`

output `Sin[3 + 2*x]/2`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

method	result	size
derivativedivides	$\frac{\sin(3+2x)}{2}$	9
default	$\frac{\sin(3+2x)}{2}$	9
risch	$\frac{\sin(3+2x)}{2}$	9
parallelrisch	$\frac{\sin(3+2x)}{2}$	9
orering	$\frac{\sin(3+2x)}{2}$	9
norman	$\frac{\tan(\frac{3}{2}+x)}{1+\tan(\frac{3}{2}+x)^2}$	16
meijerg	$\frac{\cos(3)\sin(2x)}{2} - \frac{\sin(3)\sqrt{\pi}\left(\frac{1}{\sqrt{\pi}} - \frac{\cos(2x)}{\sqrt{\pi}}\right)}{2}$	30

input `int(cos(3+2*x),x,method=_RETURNVERBOSE)`output `1/2*sin(3+2*x)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \cos(3 + 2x) dx = \frac{1}{2} \sin(2x + 3)$$

input `integrate(cos(3+2*x),x, algorithm="fricas")`output `1/2*sin(2*x + 3)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int \cos(3 + 2x) dx = \frac{\sin(2x + 3)}{2}$$

input `integrate(cos(3+2*x), x)`

output `sin(2*x + 3)/2`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \cos(3 + 2x) dx = \frac{1}{2} \sin(2x + 3)$$

input `integrate(cos(3+2*x), x, algorithm="maxima")`

output `1/2*sin(2*x + 3)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \cos(3 + 2x) dx = \frac{1}{2} \sin(2x + 3)$$

input `integrate(cos(3+2*x), x, algorithm="giac")`

output `1/2*sin(2*x + 3)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \cos(3 + 2x) dx = \frac{\sin(2x + 3)}{2}$$

input `int(cos(2*x + 3),x)`

output `sin(2*x + 3)/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \cos(3 + 2x) dx = \frac{\sin(2x + 3)}{2}$$

input `int(cos(3+2*x),x)`

output `sin(2*x + 3)/2`

3.22 $\int 2e^{2x}yz dx$

Optimal result	174
Mathematica [A] (verified)	174
Rubi [A] (verified)	175
Maple [A] (verified)	176
Fricas [A] (verification not implemented)	176
Sympy [A] (verification not implemented)	177
Maxima [A] (verification not implemented)	177
Giac [A] (verification not implemented)	177
Mupad [B] (verification not implemented)	178
Reduce [B] (verification not implemented)	178

Optimal result

Integrand size = 9, antiderivative size = 8

$$\int 2e^{2x}yz dx = e^{2x}yz$$

output `exp(2*x)*y*z`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int 2e^{2x}yz dx = e^{2x}yz$$

input `Integrate[2*E^(2*x)*y*z,x]`

output `E^(2*x)*y*z`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {27, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int 2e^{2x}yz \, dx$$

$$\downarrow 27$$

$$2yz \int e^{2x} \, dx$$

$$\downarrow 2624$$

$$e^{2x}yz$$

input `Int [2*E^(2*x)*y*z, x]`

output `E^(2*x)*y*z`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2624 `Int[((F_)^(v_))^(n_), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /; FreeQ[{F, n}, x] && LinearQ[v, x]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

method	result	size
gospers	$e^{2x}yz$	8
derivativeldivides	$e^{2x}yz$	8
default	$e^{2x}yz$	8
norman	$e^{2x}yz$	8
risch	$e^{2x}yz$	8
parallelrisch	$e^{2x}yz$	8
parts	$e^{2x}yz$	8
orering	$e^{2x}yz$	8
meijerg	$-yz(1 - e^{2x})$	13

input `int(2*exp(2*x)*y*z,x,method=_RETURNVERBOSE)`

output `exp(2*x)*y*z`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int 2e^{2x}yz dx = yze^{(2x)}$$

input `integrate(2*exp(2*x)*y*z,x, algorithm="fricas")`

output `y*z*e^(2*x)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int 2e^{2x}yz dx = yze^{2x}$$

input `integrate(2*exp(2*x)*y*z,x)`

output `y*z*exp(2*x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int 2e^{2x}yz dx = yze^{(2x)}$$

input `integrate(2*exp(2*x)*y*z,x, algorithm="maxima")`

output `y*z*e^(2*x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int 2e^{2x}yz dx = yze^{(2x)}$$

input `integrate(2*exp(2*x)*y*z,x, algorithm="giac")`

output `y*z*e^(2*x)`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int 2e^{2x}yz \, dx = yz e^{2x}$$

input `int(2*y*z*exp(2*x), x)`

output `y*z*exp(2*x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int 2e^{2x}yz \, dx = e^{2x}yz$$

input `int(2*exp(2*x)*y*z, x)`

output `e**(2*x)*y*z`

3.23 $\int e^x \cos^2(e^x) \sin(e^x) dx$

Optimal result	179
Mathematica [A] (verified)	179
Rubi [A] (warning: unable to verify)	180
Maple [A] (verified)	181
Fricas [A] (verification not implemented)	182
Sympy [A] (verification not implemented)	182
Maxima [A] (verification not implemented)	182
Giac [A] (verification not implemented)	183
Mupad [B] (verification not implemented)	183
Reduce [B] (verification not implemented)	183

Optimal result

Integrand size = 14, antiderivative size = 10

$$\int e^x \cos^2(e^x) \sin(e^x) dx = -\frac{1}{3} \cos^3(e^x)$$

output `-1/3*cos(exp(x))^3`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.90

$$\int e^x \cos^2(e^x) \sin(e^x) dx = -\frac{1}{4} \cos(e^x) - \frac{1}{12} \cos(3e^x)$$

input `Integrate[E^x*Cos[E^x]^2*Sin[E^x],x]`

output `-1/4*Cos[E^x] - Cos[3*E^x]/12`

Rubi [A] (warning: unable to verify)

Time = 0.19 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2720, 3042, 3045, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int e^x \sin(e^x) \cos^2(e^x) dx \\ & \quad \downarrow \text{2720} \\ & \int \sin(e^x) \cos^2(e^x) de^x \\ & \quad \downarrow \text{3042} \\ & \int \sin(e^x) \cos(e^x)^2 de^x \\ & \quad \downarrow \text{3045} \\ & - \int e^{2x} d \cos(e^x) \\ & \quad \downarrow \text{15} \\ & -\frac{e^{3x}}{3} \end{aligned}$$

input

 $\text{Int}[E^x \cdot \text{Cos}[E^x]^2 \cdot \text{Sin}[E^x], x]$

output

 $-1/3 \cdot E^{(3 \cdot x)}$

Definitions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3045 `Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[-(a*f)^(-1) Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])`

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

method	result	size
derivativedivides	$-\frac{\cos(e^x)^3}{3}$	8
default	$-\frac{\cos(e^x)^3}{3}$	8
risch	$-\frac{\cos(e^x)}{4} - \frac{\cos(3e^x)}{12}$	14
parallelrisch	$-\frac{\cos(e^x)}{4} - \frac{\cos(3e^x)}{12} + \frac{1}{3}$	15
norman	$\frac{-2 \tan\left(\frac{e^x}{2}\right)^4 - \frac{2}{3}}{\left(1 + \tan\left(\frac{e^x}{2}\right)^2\right)^3}$	24

input `int(exp(x)*cos(exp(x))^2*sin(exp(x)),x,method=_RETURNVERBOSE)`

output `-1/3*cos(exp(x))^3`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int e^x \cos^2(e^x) \sin(e^x) dx = -\frac{1}{3} \cos(e^x)^3$$

input `integrate(exp(x)*cos(exp(x))^2*sin(exp(x)),x, algorithm="fricas")`

output `-1/3*cos(e^x)^3`

Sympy [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int e^x \cos^2(e^x) \sin(e^x) dx = -\frac{\cos^3(e^x)}{3}$$

input `integrate(exp(x)*cos(exp(x))**2*sin(exp(x)),x)`

output `-cos(exp(x))**3/3`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int e^x \cos^2(e^x) \sin(e^x) dx = -\frac{1}{3} \cos(e^x)^3$$

input `integrate(exp(x)*cos(exp(x))^2*sin(exp(x)),x, algorithm="maxima")`

output `-1/3*cos(e^x)^3`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int e^x \cos^2(e^x) \sin(e^x) dx = -\frac{1}{3} \cos(e^x)^3$$

input `integrate(exp(x)*cos(exp(x))^2*sin(exp(x)),x, algorithm="giac")`output `-1/3*cos(e^x)^3`**Mupad [B] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int e^x \cos^2(e^x) \sin(e^x) dx = -\frac{\cos(e^x)^3}{3}$$

input `int(cos(exp(x))^2*sin(exp(x))*exp(x),x)`output `-cos(exp(x))^3/3`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int e^x \cos^2(e^x) \sin(e^x) dx = -\frac{\cos(e^x)^3}{3}$$

input `int(exp(x)*cos(exp(x))^2*sin(exp(x)),x)`output `(- cos(e**x)**3)/3`

3.24 $\int x\sqrt{1+x} dx$

Optimal result	184
Mathematica [A] (verified)	184
Rubi [A] (verified)	185
Maple [A] (verified)	186
Fricas [A] (verification not implemented)	186
Sympy [A] (verification not implemented)	187
Maxima [A] (verification not implemented)	187
Giac [A] (verification not implemented)	187
Mupad [B] (verification not implemented)	188
Reduce [B] (verification not implemented)	188

Optimal result

Integrand size = 9, antiderivative size = 23

$$\int x\sqrt{1+x} dx = -\frac{2}{3}(1+x)^{3/2} + \frac{2}{5}(1+x)^{5/2}$$

output `-2/3*(1+x)^(3/2)+2/5*(1+x)^(5/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

$$\int x\sqrt{1+x} dx = \frac{2}{15}(1+x)^{3/2}(-5+3(1+x))$$

input `Integrate[x*Sqrt[1 + x],x]`

output `(2*(1 + x)^(3/2)*(-5 + 3*(1 + x)))/15`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x\sqrt{x+1} dx$$

↓ 53

$$\int \left((x+1)^{3/2} - \sqrt{x+1} \right) dx$$

↓ 2009

$$\frac{2}{5}(x+1)^{5/2} - \frac{2}{3}(x+1)^{3/2}$$

input `Int[x*Sqrt[1 + x],x]`

output `(-2*(1 + x)^(3/2))/3 + (2*(1 + x)^(5/2))/5`

Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.57

method	result	size
gosper	$\frac{2(1+x)^{\frac{3}{2}}(-2+3x)}{15}$	13
orering	$\frac{2(1+x)^{\frac{3}{2}}(-2+3x)}{15}$	13
derivativedivides	$-\frac{2(1+x)^{\frac{3}{2}}}{3} + \frac{2(1+x)^{\frac{5}{2}}}{5}$	16
default	$-\frac{2(1+x)^{\frac{3}{2}}}{3} + \frac{2(1+x)^{\frac{5}{2}}}{5}$	16
risch	$\frac{2(3x^2+x-2)\sqrt{1+x}}{15}$	16
trager	$\left(\frac{2}{5}x^2 + \frac{2}{15}x - \frac{4}{15}\right)\sqrt{1+x}$	17
meijerg	$-\frac{8\sqrt{\pi}}{15} + \frac{4\sqrt{\pi}(1+x)^{\frac{3}{2}}(2-3x)}{15 \cdot 2\sqrt{\pi}}$	27

input `int(x*(1+x)^(1/2),x,method=_RETURNVERBOSE)`output `2/15*(1+x)^(3/2)*(-2+3*x)`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int x\sqrt{1+x} dx = \frac{2}{15} (3x^2 + x - 2)\sqrt{x+1}$$

input `integrate(x*(1+x)^(1/2),x, algorithm="fricas")`output `2/15*(3*x^2 + x - 2)*sqrt(x + 1)`

Sympy [A] (verification not implemented)

Time = 0.48 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.48

$$\int x\sqrt{1+x} dx = \frac{2x^2\sqrt{x+1}}{5} + \frac{2x\sqrt{x+1}}{15} - \frac{4\sqrt{x+1}}{15}$$

input `integrate(x*(1+x)**(1/2),x)`output `2*x**2*sqrt(x + 1)/5 + 2*x*sqrt(x + 1)/15 - 4*sqrt(x + 1)/15`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int x\sqrt{1+x} dx = \frac{2}{5}(x+1)^{\frac{5}{2}} - \frac{2}{3}(x+1)^{\frac{3}{2}}$$

input `integrate(x*(1+x)^(1/2),x, algorithm="maxima")`output `2/5*(x + 1)^(5/2) - 2/3*(x + 1)^(3/2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int x\sqrt{1+x} dx = \frac{2}{5}(x+1)^{\frac{5}{2}} - \frac{2}{3}(x+1)^{\frac{3}{2}}$$

input `integrate(x*(1+x)^(1/2),x, algorithm="giac")`output `2/5*(x + 1)^(5/2) - 2/3*(x + 1)^(3/2)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.52

$$\int x\sqrt{1+x} dx = \frac{2(3x-2)(x+1)^{3/2}}{15}$$

input `int(x*(x + 1)^(1/2),x)`

output `(2*(3*x - 2)*(x + 1)^(3/2))/15`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.61

$$\int x\sqrt{1+x} dx = \frac{2\sqrt{x+1}(3x^2+x-2)}{15}$$

input `int(x*(1+x)^(1/2),x)`

output `(2*sqrt(x + 1)*(3*x**2 + x - 2))/15`

3.25 $\int \frac{1}{-1+x^4} dx$

Optimal result	189
Mathematica [A] (verified)	189
Rubi [A] (verified)	190
Maple [A] (verified)	191
Fricas [A] (verification not implemented)	191
Sympy [A] (verification not implemented)	192
Maxima [A] (verification not implemented)	192
Giac [B] (verification not implemented)	192
Mupad [B] (verification not implemented)	193
Reduce [B] (verification not implemented)	193

Optimal result

Integrand size = 7, antiderivative size = 13

$$\int \frac{1}{-1+x^4} dx = -\frac{\arctan(x)}{2} - \frac{\operatorname{arctanh}(x)}{2}$$

output

```
-1/2*arctan(x)-1/2*arctanh(x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.92

$$\int \frac{1}{-1+x^4} dx = -\frac{\arctan(x)}{2} + \frac{1}{4} \log(1-x) - \frac{1}{4} \log(1+x)$$

input

```
Integrate[(-1 + x^4)^(-1),x]
```

output

```
-1/2*ArcTan[x] + Log[1 - x]/4 - Log[1 + x]/4
```

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{x^4 - 1} dx \\ & \quad \downarrow \text{756} \\ & -\frac{1}{2} \int \frac{1}{1-x^2} dx - \frac{1}{2} \int \frac{1}{x^2+1} dx \\ & \quad \downarrow \text{216} \\ & -\frac{1}{2} \int \frac{1}{1-x^2} dx - \frac{\arctan(x)}{2} \\ & \quad \downarrow \text{219} \\ & -\frac{\arctan(x)}{2} - \frac{\operatorname{arctanh}(x)}{2} \end{aligned}$$

input `Int[(-1 + x^4)^(-1), x]`

output `-1/2*ArcTan[x] - ArcTanh[x]/2`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 756

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2
]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x]
+ Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a
/b, 0]
```

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

method	result	size
default	$-\frac{\arctan(x)}{2} - \frac{\operatorname{arctanh}(x)}{2}$	10
risch	$-\frac{\arctan(x)}{2} - \frac{\ln(1+x)}{4} + \frac{\ln(-1+x)}{4}$	18
parallelrisc	$\frac{i \ln(x-i)}{4} - \frac{i \ln(x+i)}{4} - \frac{\ln(1+x)}{4} + \frac{\ln(-1+x)}{4}$	30
meijerg	$\frac{x \left(\ln \left(1 - (x^4)^{\frac{1}{4}} \right) - \ln \left(1 + (x^4)^{\frac{1}{4}} \right) - 2 \arctan \left((x^4)^{\frac{1}{4}} \right) \right)}{4(x^4)^{\frac{1}{4}}}$	38

input

```
int(1/(x^4-1),x,method=_RETURNVERBOSE)
```

output

```
-1/2*arctan(x)-1/2*arctanh(x)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.31

$$\int \frac{1}{-1+x^4} dx = -\frac{1}{2} \arctan(x) - \frac{1}{4} \log(x+1) + \frac{1}{4} \log(x-1)$$

input

```
integrate(1/(x^4-1),x, algorithm="fricas")
```

output

```
-1/2*arctan(x) - 1/4*log(x + 1) + 1/4*log(x - 1)
```


Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.31

$$\int \frac{1}{-1+x^4} dx = \frac{\log(x-1)}{4} - \frac{\log(x+1)}{4} - \frac{\operatorname{atan}(x)}{2}$$

input `integrate(1/(x**4-1),x)`

output `log(x - 1)/4 - log(x + 1)/4 - atan(x)/2`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.31

$$\int \frac{1}{-1+x^4} dx = -\frac{1}{2} \arctan(x) - \frac{1}{4} \log(x+1) + \frac{1}{4} \log(x-1)$$

input `integrate(1/(x^4-1),x, algorithm="maxima")`

output `-1/2*arctan(x) - 1/4*log(x + 1) + 1/4*log(x - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. 2(9) = 18.

Time = 0.12 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.46

$$\int \frac{1}{-1+x^4} dx = -\frac{1}{2} \arctan(x) - \frac{1}{4} \log(|x+1|) + \frac{1}{4} \log(|x-1|)$$

input `integrate(1/(x^4-1),x, algorithm="giac")`

output `-1/2*arctan(x) - 1/4*log(abs(x + 1)) + 1/4*log(abs(x - 1))`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int \frac{1}{-1+x^4} dx = -\frac{\operatorname{atan}(x)}{2} - \frac{\operatorname{atanh}(x)}{2}$$

input `int(1/(x^4 - 1),x)`

output `- atan(x)/2 - atanh(x)/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.31

$$\int \frac{1}{-1+x^4} dx = -\frac{\operatorname{atan}(x)}{2} + \frac{\log(x-1)}{4} - \frac{\log(x+1)}{4}$$

input `int(1/(x^4-1),x)`

output `(- 2*atan(x) + log(x - 1) - log(x + 1))/4`

3.26 $\int \frac{e^x}{2+3e^{2x}} dx$

Optimal result	194
Mathematica [A] (verified)	194
Rubi [A] (verified)	195
Maple [A] (verified)	196
Fricas [A] (verification not implemented)	196
Sympy [A] (verification not implemented)	196
Maxima [A] (verification not implemented)	197
Giac [A] (verification not implemented)	197
Mupad [B] (verification not implemented)	197
Reduce [B] (verification not implemented)	198

Optimal result

Integrand size = 15, antiderivative size = 18

$$\int \frac{e^x}{2+3e^{2x}} dx = \frac{\arctan\left(\sqrt{\frac{3}{2}}e^x\right)}{\sqrt{6}}$$

output `1/6*arctan(1/2*exp(x)*6^(1/2))*6^(1/2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{e^x}{2+3e^{2x}} dx = \frac{\arctan\left(\sqrt{\frac{3}{2}}e^x\right)}{\sqrt{6}}$$

input `Integrate[E^x/(2 + 3*E^(2*x)), x]`

output `ArcTan[Sqrt[3/2]*E^x]/Sqrt[6]`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2679, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^x}{3e^{2x} + 2} dx$$

↓ 2679

$$\int \frac{1}{3e^{2x} + 2} de^x$$

↓ 216

$$\frac{\arctan\left(\sqrt{\frac{3}{2}}e^x\right)}{\sqrt{6}}$$

input `Int[E^x/(2 + 3*E^(2*x)),x]`

output `ArcTan[Sqrt[3/2]*E^x]/Sqrt[6]`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 2679 `Int[((a_) + (b_.)*(F_)^((e_.)*((c_.) + (d_.)*(x_))))^(p_.)*(G_)^((h_.)*((f_.) + (g_.)*(x_))), x_Symbol] := With[{m = FullSimplify[d*e*(Log[F]/(g*h*Log[G]))]}, Simp[Denominator[m]/(g*h*Log[G]) Subst[Int[x^(Denominator[m] - 1)]*(a + b*F^(c*e - d*e*(f/g))*x^Numerator[m])^p, x], x, G^(h*((f + g*x)/Denominator[m]))], x] /; LtQ[m, -1] || GtQ[m, 1] /; FreeQ[{F, G, a, b, c, d, e, f, g, h, p}, x]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

method	result	size
default	$\frac{\arctan\left(\frac{e^x\sqrt{6}}{2}\right)\sqrt{6}}{6}$	14
risch	$\frac{i\sqrt{6}\ln\left(e^x + \frac{i\sqrt{6}}{3}\right)}{12} - \frac{i\sqrt{6}\ln\left(e^x - \frac{i\sqrt{6}}{3}\right)}{12}$	34

input `int(exp(x)/(2+3*exp(2*x)),x,method=_RETURNVERBOSE)`output `1/6*arctan(1/2*exp(x)*6^(1/2))*6^(1/2)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.72

$$\int \frac{e^x}{2 + 3e^{2x}} dx = \frac{1}{6} \sqrt{6} \arctan\left(\frac{1}{2} \sqrt{6} e^x\right)$$

input `integrate(exp(x)/(2+3*exp(2*x)),x, algorithm="fricas")`output `1/6*sqrt(6)*arctan(1/2*sqrt(6)*e^x)`**Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int \frac{e^x}{2 + 3e^{2x}} dx = \text{RootSum}(24z^2 + 1, (i \mapsto i \log(4i + e^x)))$$

input `integrate(exp(x)/(2+3*exp(2*x)),x)`output `RootSum(24*_z**2 + 1, Lambda(_i, _i*log(4*_i + exp(x))))`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.72

$$\int \frac{e^x}{2 + 3e^{2x}} dx = \frac{1}{6} \sqrt{6} \arctan \left(\frac{1}{2} \sqrt{6} e^x \right)$$

input `integrate(exp(x)/(2+3*exp(2*x)),x, algorithm="maxima")`output `1/6*sqrt(6)*arctan(1/2*sqrt(6)*e^x)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.72

$$\int \frac{e^x}{2 + 3e^{2x}} dx = \frac{1}{6} \sqrt{6} \arctan \left(\frac{1}{2} \sqrt{6} e^x \right)$$

input `integrate(exp(x)/(2+3*exp(2*x)),x, algorithm="giac")`output `1/6*sqrt(6)*arctan(1/2*sqrt(6)*e^x)`**Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.72

$$\int \frac{e^x}{2 + 3e^{2x}} dx = \frac{\sqrt{6} \operatorname{atan} \left(\frac{\sqrt{6} e^x}{2} \right)}{6}$$

input `int(exp(x)/(3*exp(2*x) + 2),x)`output `(6^(1/2)*atan((6^(1/2)*exp(x))/2))/6`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{e^x}{2 + 3e^{2x}} dx = \frac{\sqrt{6} \operatorname{atan}\left(\frac{3e^x}{\sqrt{6}}\right)}{6}$$

input `int(exp(x)/(2+3*exp(2*x)),x)`

output `(sqrt(6)*atan((3*e**x)/sqrt(6)))/6`

3.27 $\int \frac{e^{2x}}{A+Be^{4x}} dx$

Optimal result	199
Mathematica [A] (verified)	199
Rubi [A] (verified)	200
Maple [A] (verified)	201
Fricas [A] (verification not implemented)	201
Sympy [A] (verification not implemented)	201
Maxima [A] (verification not implemented)	202
Giac [A] (verification not implemented)	202
Mupad [B] (verification not implemented)	202
Reduce [B] (verification not implemented)	203

Optimal result

Integrand size = 17, antiderivative size = 31

$$\int \frac{e^{2x}}{A + Be^{4x}} dx = \frac{\arctan\left(\frac{\sqrt{B}e^{2x}}{\sqrt{A}}\right)}{2\sqrt{A}\sqrt{B}}$$

output `1/2*arctan(exp(2*x)*B^(1/2)/A^(1/2))/A^(1/2)/B^(1/2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{e^{2x}}{A + Be^{4x}} dx = \frac{\arctan\left(\frac{\sqrt{B}e^{2x}}{\sqrt{A}}\right)}{2\sqrt{A}\sqrt{B}}$$

input `Integrate[E^(2*x)/(A + B*E^(4*x)), x]`

output `ArcTan[(Sqrt[B]*E^(2*x))/Sqrt[A]]/(2*Sqrt[A]*Sqrt[B])`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2679, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^{2x}}{A + Be^{4x}} dx$$

↓ 2679

$$\frac{1}{2} \int \frac{1}{A + Be^{4x}} de^{2x}$$

↓ 218

$$\frac{\arctan\left(\frac{\sqrt{B}e^{2x}}{\sqrt{A}}\right)}{2\sqrt{A}\sqrt{B}}$$

input `Int [E^(2*x)/(A + B*E^(4*x)), x]`

output `ArcTan[(Sqrt[B]*E^(2*x))/Sqrt[A]]/(2*Sqrt[A]*Sqrt[B])`

Defintions of rubi rules used

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 2679 `Int[((a_) + (b_)*(F_)^(e_)*((c_) + (d_)*(x_)))^(p_)*(G_)^(h_)*((f_) + (g_)*(x_)), x_Symbol] := With[{m = FullSimplify[d*e*(Log[F]/(g*h*Log[G]))]}, Simp[Denominator[m]/(g*h*Log[G]) Subst[Int[x^(Denominator[m] - 1)*(a + b*F^(c*e - d*e*(f/g))*x^Numerator[m])^p, x], x, G^(h*((f + g*x)/Denominator[m]))], x] /; LtQ[m, -1] || GtQ[m, 1]] /; FreeQ[{F, G, a, b, c, d, e, f, g, h, p}, x]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.65

method	result	size
default	$\frac{\arctan\left(\frac{B e^{2x}}{\sqrt{AB}}\right)}{2\sqrt{AB}}$	20
risch	$-\frac{\ln\left(e^{2x} - \frac{A}{\sqrt{-AB}}\right)}{4\sqrt{-AB}} + \frac{\ln\left(e^{2x} + \frac{A}{\sqrt{-AB}}\right)}{4\sqrt{-AB}}$	47

input `int(exp(2*x)/(A+B*exp(4*x)),x,method=_RETURNVERBOSE)`output `1/2/(A*B)^(1/2)*arctan(B*exp(x)^2/(A*B)^(1/2))`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 76, normalized size of antiderivative = 2.45

$$\int \frac{e^{2x}}{A + B e^{4x}} dx = \left[-\frac{\sqrt{-AB} \log\left(\frac{B e^{(4x)} - 2\sqrt{-AB} e^{(2x)} - A}{B e^{(4x)} + A}\right)}{4AB}, -\frac{\sqrt{AB} \arctan\left(\frac{\sqrt{AB} e^{(-2x)}}{B}\right)}{2AB} \right]$$

input `integrate(exp(2*x)/(A+B*exp(4*x)),x, algorithm="fricas")`output `[-1/4*sqrt(-A*B)*log((B*e^(4*x) - 2*sqrt(-A*B)*e^(2*x) - A)/(B*e^(4*x) + A))/ (A*B), -1/2*sqrt(A*B)*arctan(sqrt(A*B)*e^(-2*x)/B)/(A*B)]`**Sympy [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.71

$$\int \frac{e^{2x}}{A + B e^{4x}} dx = \text{RootSum}(16z^2 AB + 1, (i \mapsto i \log(4iA + e^{2x})))$$

input `integrate(exp(2*x)/(A+B*exp(4*x)),x)`

output `RootSum(16*_z**2*A*B + 1, Lambda(_i, _i*log(4*_i*A + exp(2*x))))`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.61

$$\int \frac{e^{2x}}{A + Be^{4x}} dx = \frac{\arctan\left(\frac{Be^{2x}}{\sqrt{AB}}\right)}{2\sqrt{AB}}$$

input `integrate(exp(2*x)/(A+B*exp(4*x)),x, algorithm="maxima")`

output `1/2*arctan(B*e^(2*x)/sqrt(A*B))/sqrt(A*B)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.61

$$\int \frac{e^{2x}}{A + Be^{4x}} dx = \frac{\arctan\left(\frac{Be^{2x}}{\sqrt{AB}}\right)}{2\sqrt{AB}}$$

input `integrate(exp(2*x)/(A+B*exp(4*x)),x, algorithm="giac")`

output `1/2*arctan(B*e^(2*x)/sqrt(A*B))/sqrt(A*B)`

Mupad [B] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.61

$$\int \frac{e^{2x}}{A + Be^{4x}} dx = \frac{\operatorname{atan}\left(\frac{Be^{2x}}{\sqrt{AB}}\right)}{2\sqrt{AB}}$$

input `int(exp(2*x)/(A + B*exp(4*x)),x)`

output `atan((B*exp(2*x))/(A*B)^(1/2))/(2*(A*B)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 71, normalized size of antiderivative = 2.29

$$\int \frac{e^{2x}}{A + Be^{4x}} dx = -\frac{\sqrt{b}\sqrt{a} \left(\operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}-2e^x\sqrt{b}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) + \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}+2e^x\sqrt{b}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) \right)}{2ab}$$

input `int(exp(2*x)/(A+B*exp(4*x)),x)`

output `(- sqrt(b)*sqrt(a)*(atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*e**x*sqrt(b))/(b**
*(1/4)*a**(1/4)*sqrt(2))) + atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*e**x*sqrt(
b))/(b**(1/4)*a**(1/4)*sqrt(2))))/(2*a*b)`

3.28 $\int \frac{e^{1+x}}{1+e^x} dx$

Optimal result	204
Mathematica [A] (verified)	204
Rubi [A] (verified)	205
Maple [A] (verified)	206
Fricas [A] (verification not implemented)	206
Sympy [A] (verification not implemented)	207
Maxima [A] (verification not implemented)	207
Giac [A] (verification not implemented)	207
Mupad [B] (verification not implemented)	208
Reduce [B] (verification not implemented)	208

Optimal result

Integrand size = 13, antiderivative size = 8

$$\int \frac{e^{1+x}}{1+e^x} dx = e \log(1+e^x)$$

output

```
exp(1)*ln(1+exp(x))
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{e^{1+x}}{1+e^x} dx = e \log(1+e^x)$$

input

```
Integrate[E^(1 + x)/(1 + E^x),x]
```

output

```
E*Log[1 + E^x]
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {2677, 2676, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{e^{x+1}}{e^x + 1} dx \\ & \quad \downarrow \text{2677} \\ & e \int \frac{e^x}{1 + e^x} dx \\ & \quad \downarrow \text{2676} \\ & e \int \frac{1}{1 + e^x} de^x \\ & \quad \downarrow \text{16} \\ & e \log(e^x + 1) \end{aligned}$$

input `Int[E^(1 + x)/(1 + E^x),x]`

output `E*Log[1 + E^x]`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] :> Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 2676 `Int[((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)*((a_) + (b_.)*(F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.))^p, x_Symbol] :> Simp[1/(d*e*n*Log[F]) Subst[Int[(a + b*x)^p, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n, p}, x]`

rule 2677

```
Int[((a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))^(p_))*((G_)^((
h_)*((f_) + (g_)*(x_)))^(m_)), x_Symbol] :-> Simp[(G^(h*(f + g*x)))^m/(F
^(e*(c + d*x)))^n Int[(F^(e*(c + d*x)))^n*(a + b*(F^(e*(c + d*x)))^n)^p,
x], x] /; FreeQ[{F, G, a, b, c, d, e, f, g, h, m, n, p}, x] && EqQ[d*e*n*Lo
g[F], g*h*m*Log[G]]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.12

method	result	size
default	$e \ln(1 + e^x)$	9
norman	$e \ln(1 + e^x)$	9
risch	$e \ln(1 + e^x)$	9

input

```
int(exp(1+x)/(1+exp(x)),x,method=_RETURNVERBOSE)
```

output

```
exp(1)*ln(1+exp(x))
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.38

$$\int \frac{e^{1+x}}{1 + e^x} dx = e \log(e + e^{(x+1)})$$

input

```
integrate(exp(1+x)/(1+exp(x)),x, algorithm="fricas")
```

output

```
e*log(e + e^(x + 1))
```

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{e^{1+x}}{1+e^x} dx = e \log(e^x + 1)$$

input `integrate(exp(1+x)/(1+exp(x)),x)`

output `E*log(exp(x) + 1)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{e^{1+x}}{1+e^x} dx = e \log(e^x + 1)$$

input `integrate(exp(1+x)/(1+exp(x)),x, algorithm="maxima")`

output `e*log(e^x + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{e^{1+x}}{1+e^x} dx = e \log(e^x + 1)$$

input `integrate(exp(1+x)/(1+exp(x)),x, algorithm="giac")`

output `e*log(e^x + 1)`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{e^{1+x}}{1+e^x} dx = e \ln(e^x + 1)$$

input `int(exp(x + 1)/(exp(x) + 1),x)`

output `exp(1)*log(exp(x) + 1)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{e^{1+x}}{1+e^x} dx = \log(e^x + 1) e$$

input `int(exp(1+x)/(1+exp(x)),x)`

output `log(e**x + 1)*e`

3.29 $\int (10e)^x dx$

Optimal result	209
Mathematica [A] (verified)	209
Rubi [A] (verified)	210
Maple [A] (verified)	211
Fricas [A] (verification not implemented)	211
Sympy [A] (verification not implemented)	212
Maxima [A] (verification not implemented)	212
Giac [A] (verification not implemented)	212
Mupad [B] (verification not implemented)	213
Reduce [B] (verification not implemented)	213

Optimal result

Integrand size = 5, antiderivative size = 12

$$\int (10e)^x dx = \frac{(10e)^x}{1 + \log(10)}$$

output

```
(10*exp(1))^x/(1+ln(10))
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int (10e)^x dx = \frac{(10e)^x}{\log(10e)}$$

input

```
Integrate[(10*E)^x,x]
```

output

```
(10*E)^x/Log[10*E]
```

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (10e)^x dx$$

$$\downarrow 2624$$

$$\frac{(10e)^x}{1 + \log(10)}$$

input

```
Int[(10*E)^x,x]
```

output

```
(10*E)^x/(1 + Log[10])
```

Defintions of rubi rules used

rule 2624

```
Int[((F_)^(v_))^(n_), x_Symbol] := Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;
FreeQ[{F, n}, x] && LinearQ[v, x]
```

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.25

method	result	size
gospers	$\frac{(10e)^x}{\ln(10e)}$	15
parallelrisch	$\frac{(10e)^x}{\ln(10e)}$	15
norman	$\frac{e^x \ln(10e)}{1+\ln(10)}$	16
orering	$\frac{(10e)^x}{1+\ln(2)+\ln(5)}$	16
risch	$\frac{5^x 2^x e^x}{1+\ln(2)+\ln(5)}$	18
meijerg	$-\frac{1-e^{x(1+\ln(10))}}{1+\ln(10)}$	20

input `int((10*exp(1))^x,x,method=_RETURNVERBOSE)`output `1/ln(10*exp(1))*(10*exp(1))^x`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int (10e)^x dx = \frac{e^{(x \log(10)+x)}}{\log(10) + 1}$$

input `integrate((10*exp(1))^x,x, algorithm="fricas")`output `e^(x*log(10) + x)/(log(10) + 1)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int (10e)^x dx = \frac{(10e)^x}{1 + \log(10)}$$

input `integrate((10*exp(1))**x,x)`

output `(10*E)**x/(1 + log(10))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int (10e)^x dx = \frac{(10e)^x}{\log(10e)}$$

input `integrate((10*exp(1))^x,x, algorithm="maxima")`

output `(10*e)^x/log(10*e)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

$$\int (10e)^x dx = \frac{(10e)^x}{\log(10) + 1}$$

input `integrate((10*exp(1))^x,x, algorithm="giac")`

output `(10*e)^x/(log(10) + 1)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int (10e)^x dx = \frac{10^x e^x}{\ln(10) + 1}$$

input `int((10*exp(1))^x,x)`

output `(10^x*exp(x))/(log(10) + 1)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

$$\int (10e)^x dx = \frac{e^x 10^x}{\log(10) + 1}$$

input `int((10*exp(1))^x,x)`

output `(e**x*10**x)/(log(10) + 1)`

3.30 $\int x^3 \sin(x^2) dx$

Optimal result	214
Mathematica [A] (verified)	214
Rubi [A] (verified)	215
Maple [A] (verified)	216
Fricas [A] (verification not implemented)	217
Sympy [A] (verification not implemented)	218
Maxima [A] (verification not implemented)	218
Giac [A] (verification not implemented)	218
Mupad [B] (verification not implemented)	219
Reduce [B] (verification not implemented)	219

Optimal result

Integrand size = 8, antiderivative size = 20

$$\int x^3 \sin(x^2) dx = -\frac{1}{2}x^2 \cos(x^2) + \frac{\sin(x^2)}{2}$$

output `-1/2*x^2*cos(x^2)+1/2*sin(x^2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int x^3 \sin(x^2) dx = -\frac{1}{2}x^2 \cos(x^2) + \frac{\sin(x^2)}{2}$$

input `Integrate[x^3*Sin[x^2],x]`

output `-1/2*(x^2*Cos[x^2]) + Sin[x^2]/2`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {3860, 3042, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \sin(x^2) dx \\
 & \quad \downarrow \text{3860} \\
 & \frac{1}{2} \int x^2 \sin(x^2) dx^2 \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int x^2 \sin(x^2) dx^2 \\
 & \quad \downarrow \text{3777} \\
 & \frac{1}{2} \left(\int \cos(x^2) dx^2 - x^2 \cos(x^2) \right) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \left(\int \sin\left(x^2 + \frac{\pi}{2}\right) dx^2 - x^2 \cos(x^2) \right) \\
 & \quad \downarrow \text{3117} \\
 & \frac{1}{2} (\sin(x^2) - x^2 \cos(x^2))
 \end{aligned}$$

input

Int[x^3*Sin[x^2],x]

output

(-(x^2*Cos[x^2]) + Sin[x^2])/2

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 3860 `Int[(x_)^(m_.)*((a_.) + (b_.)*Sin[(c_.) + (d_.)*(x_)^(n_)])^(p_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Sin[c + d*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))`

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

method	result	size
derivativedivides	$-\frac{x^2 \cos(x^2)}{2} + \frac{\sin(x^2)}{2}$	17
default	$-\frac{x^2 \cos(x^2)}{2} + \frac{\sin(x^2)}{2}$	17
risch	$-\frac{x^2 \cos(x^2)}{2} + \frac{\sin(x^2)}{2}$	17
parallelrisc	$-\frac{x^2 \cos(x^2)}{2} + \frac{\sin(x^2)}{2}$	17
meijerg	$\sqrt{\pi} \left(-\frac{x^2 \cos(x^2)}{2\sqrt{\pi}} + \frac{\sin(x^2)}{2\sqrt{\pi}} \right)$	27
orering	$\frac{5 \sin(x^2)}{4} - \frac{3x^2 \sin(x^2) + 2x^4 \cos(x^2)}{4x^2}$	32
norman	$\frac{-\frac{x^2}{2} + \frac{x^2 \tan\left(\frac{x^2}{2}\right)^2}{2} + \tan\left(\frac{x^2}{2}\right)}{1 + \tan\left(\frac{x^2}{2}\right)^2}$	39
parts	$\frac{\sqrt{2} \sqrt{\pi} \operatorname{FresnelS}\left(\frac{\sqrt{2}x}{\sqrt{\pi}}\right) x^3}{2} - \frac{3\pi^2 \left(\frac{2 \operatorname{FresnelS}\left(\frac{\sqrt{2}x}{\sqrt{\pi}}\right) \sqrt{2} x^3}{3\pi^{\frac{3}{2}}} + \frac{2x^2 \cos(x^2)}{3\pi^2} - \frac{2 \sin(x^2)}{3\pi^2} \right)}{4}$	69

input `int(x^3*sin(x^2),x,method=_RETURNVERBOSE)`

output `-1/2*x^2*cos(x^2)+1/2*sin(x^2)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int x^3 \sin(x^2) dx = -\frac{1}{2} x^2 \cos(x^2) + \frac{1}{2} \sin(x^2)$$

input `integrate(x^3*sin(x^2),x, algorithm="fricas")`

output `-1/2*x^2*cos(x^2) + 1/2*sin(x^2)`

Sympy [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

$$\int x^3 \sin(x^2) dx = -\frac{x^2 \cos(x^2)}{2} + \frac{\sin(x^2)}{2}$$

input `integrate(x**3*sin(x**2),x)`output `-x**2*cos(x**2)/2 + sin(x**2)/2`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int x^3 \sin(x^2) dx = -\frac{1}{2} x^2 \cos(x^2) + \frac{1}{2} \sin(x^2)$$

input `integrate(x^3*sin(x^2),x, algorithm="maxima")`output `-1/2*x^2*cos(x^2) + 1/2*sin(x^2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int x^3 \sin(x^2) dx = -\frac{1}{2} x^2 \cos(x^2) + \frac{1}{2} \sin(x^2)$$

input `integrate(x^3*sin(x^2),x, algorithm="giac")`output `-1/2*x^2*cos(x^2) + 1/2*sin(x^2)`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int x^3 \sin(x^2) dx = \frac{\sin(x^2)}{2} - \frac{x^2 \cos(x^2)}{2}$$

input `int(x^3*sin(x^2),x)`

output `sin(x^2)/2 - (x^2*cos(x^2))/2`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int x^3 \sin(x^2) dx = -\frac{\cos(x^2) x^2}{2} + \frac{\sin(x^2)}{2}$$

input `int(x^3*sin(x^2),x)`

output `(- cos(x**2)*x**2 + sin(x**2))/2`

3.31 $\int \frac{x^7}{1+x^{12}} dx$

Optimal result	220
Mathematica [B] (verified)	221
Rubi [A] (verified)	221
Maple [A] (verified)	224
Fricas [A] (verification not implemented)	224
Sympy [A] (verification not implemented)	225
Maxima [A] (verification not implemented)	225
Giac [A] (verification not implemented)	225
Mupad [B] (verification not implemented)	226
Reduce [B] (verification not implemented)	227

Optimal result

Integrand size = 11, antiderivative size = 49

$$\int \frac{x^7}{1+x^{12}} dx = -\frac{\arctan\left(\frac{1-2x^4}{\sqrt{3}}\right)}{4\sqrt{3}} - \frac{1}{12} \log(1+x^4) + \frac{1}{24} \log(1-x^4+x^8)$$

output

`-1/12*ln(x^4+1)+1/24*ln(x^8-x^4+1)-1/12*arctan(1/3*(-2*x^4+1)*3^(1/2))*3^(1/2)`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 260 vs. $2(49) = 98$.

Time = 0.07 (sec) , antiderivative size = 260, normalized size of antiderivative = 5.31

$$\int \frac{x^7}{1+x^{12}} dx = \frac{1}{24} \left(2\sqrt{3} \arctan \left(\frac{1+\sqrt{3}-2\sqrt{2}x}{1-\sqrt{3}} \right) - 2\sqrt{3} \arctan \left(\frac{1-\sqrt{3}+2\sqrt{2}x}{1+\sqrt{3}} \right) \right. \\ \left. + 2\sqrt{3} \arctan \left(\frac{-1+\sqrt{3}+2\sqrt{2}x}{1+\sqrt{3}} \right) - 2\sqrt{3} \arctan \left(\frac{1+\sqrt{3}+2\sqrt{2}x}{-1+\sqrt{3}} \right) \right. \\ \left. - 2 \log \left(1 - \sqrt{2}x + x^2 \right) - 2 \log \left(1 + \sqrt{2}x + x^2 \right) \right. \\ \left. + \log \left(2 + \sqrt{2}x - \sqrt{6}x + 2x^2 \right) + \log \left(2 + \sqrt{2}(-1 + \sqrt{3})x + 2x^2 \right) \right. \\ \left. + \log \left(2 - (\sqrt{2} + \sqrt{6})x + 2x^2 \right) + \log \left(2 + (\sqrt{2} + \sqrt{6})x + 2x^2 \right) \right)$$

input

```
Integrate[x^7/(1 + x^12),x]
```

output

```
(2*Sqrt[3]*ArcTan[(1 + Sqrt[3] - 2*Sqrt[2]*x)/(1 - Sqrt[3])] - 2*Sqrt[3]*ArcTan[(1 - Sqrt[3] + 2*Sqrt[2]*x)/(1 + Sqrt[3])] + 2*Sqrt[3]*ArcTan[(-1 + Sqrt[3] + 2*Sqrt[2]*x)/(1 + Sqrt[3])] - 2*Sqrt[3]*ArcTan[(1 + Sqrt[3] + 2*Sqrt[2]*x)/(-1 + Sqrt[3])] - 2*Log[1 - Sqrt[2]*x + x^2] - 2*Log[1 + Sqrt[2]*x + x^2] + Log[2 + Sqrt[2]*x - Sqrt[6]*x + 2*x^2] + Log[2 + Sqrt[2]*(-1 + Sqrt[3])*x + 2*x^2] + Log[2 - (Sqrt[2] + Sqrt[6])*x + 2*x^2] + Log[2 + (Sqrt[2] + Sqrt[6])*x + 2*x^2])/24
```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.12, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.727$, Rules used = {807, 821, 16, 1142, 25, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{x^7}{x^{12} + 1} dx \\
& \quad \downarrow \text{807} \\
& \frac{1}{4} \int \frac{x^4}{x^{12} + 1} dx^4 \\
& \quad \downarrow \text{821} \\
& \frac{1}{4} \left(\frac{1}{3} \int \frac{x^4 + 1}{x^8 - x^4 + 1} dx^4 - \frac{1}{3} \int \frac{1}{x^4 + 1} dx^4 \right) \\
& \quad \downarrow \text{16} \\
& \frac{1}{4} \left(\frac{1}{3} \int \frac{x^4 + 1}{x^8 - x^4 + 1} dx^4 - \frac{1}{3} \log(x^4 + 1) \right) \\
& \quad \downarrow \text{1142} \\
& \frac{1}{4} \left(\frac{1}{3} \left(\frac{3}{2} \int \frac{1}{x^8 - x^4 + 1} dx^4 + \frac{1}{2} \int -\frac{1 - 2x^4}{x^8 - x^4 + 1} dx^4 \right) - \frac{1}{3} \log(x^4 + 1) \right) \\
& \quad \downarrow \text{25} \\
& \frac{1}{4} \left(\frac{1}{3} \left(\frac{3}{2} \int \frac{1}{x^8 - x^4 + 1} dx^4 - \frac{1}{2} \int \frac{1 - 2x^4}{x^8 - x^4 + 1} dx^4 \right) - \frac{1}{3} \log(x^4 + 1) \right) \\
& \quad \downarrow \text{1083} \\
& \frac{1}{4} \left(\frac{1}{3} \left(-3 \int \frac{1}{-x^8 - 3} d(2x^4 - 1) - \frac{1}{2} \int \frac{1 - 2x^4}{x^8 - x^4 + 1} dx^4 \right) - \frac{1}{3} \log(x^4 + 1) \right) \\
& \quad \downarrow \text{217} \\
& \frac{1}{4} \left(\frac{1}{3} \left(\sqrt{3} \arctan\left(\frac{2x^4 - 1}{\sqrt{3}}\right) - \frac{1}{2} \int \frac{1 - 2x^4}{x^8 - x^4 + 1} dx^4 \right) - \frac{1}{3} \log(x^4 + 1) \right) \\
& \quad \downarrow \text{1103} \\
& \frac{1}{4} \left(\frac{1}{3} \left(\sqrt{3} \arctan\left(\frac{2x^4 - 1}{\sqrt{3}}\right) + \frac{1}{2} \log(x^8 - x^4 + 1) \right) - \frac{1}{3} \log(x^4 + 1) \right)
\end{aligned}$$

input `Int[x^7/(1 + x^12),x]`

output `(-1/3*Log[1 + x^4] + (Sqrt[3]*ArcTan[(-1 + 2*x^4)/Sqrt[3]] + Log[1 - x^4 + x^8]/2)/3)/4`

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 25 $\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[Fx, x], x]$
- rule 217 $\text{Int}[(a_)+(b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b] \& \& (\text{LtQ}[a, 0] \parallel \text{LtQ}[b, 0])]$
- rule 807 $\text{Int}[(x_)^{(m_)}*((a_)+(b_)*(x_)^{(n_))}^{(p_)}, x_Symbol] \rightarrow \text{With}[\{k = \text{GCD}[m + 1, n]\}, \text{Simp}[1/k \text{ Subst}[\text{Int}[x^{(m+1)/k-1}*(a + b*x^{(n/k)})^p, x], x, x^k], x] /; k \neq 1] /; \text{FreeQ}[\{a, b, p\}, x] \&\& \text{IGtQ}[n, 0] \&\& \text{IntegerQ}[m]$
- rule 821 $\text{Int}[(x_)/((a_)+(b_)*(x_)^3), x_Symbol] \rightarrow \text{Simp}[-(3*\text{Rt}[a, 3]*\text{Rt}[b, 3])^{(-1)} \text{ Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]*\text{Rt}[b, 3]) \text{ Int}[(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$
- rule 1083 $\text{Int}[(a_)+(b_)*(x_)+(c_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[-2 \text{ Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{EqQ}[2*c*d - b*e, 0]$
- rule 1142 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \text{ Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \text{ Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.80

method	result	size
risch	$-\frac{\ln(x^4+1)}{12} + \frac{\ln(x^8-x^4+1)}{24} + \frac{\sqrt{3} \arctan\left(\frac{2(x^4-\frac{1}{2})\sqrt{3}}{3}\right)}{12}$	39
default	$-\frac{\ln(x^4+1)}{12} + \frac{\ln(x^8-x^4+1)}{24} + \frac{\arctan\left(\frac{(2x^4-1)\sqrt{3}}{3}\right)\sqrt{3}}{12}$	41
meijerg	$-\frac{x^8 \ln\left(1+(x^{12})^{\frac{1}{3}}\right)}{12(x^{12})^{\frac{2}{3}}} + \frac{x^8 \ln\left(1-(x^{12})^{\frac{1}{3}}+(x^{12})^{\frac{2}{3}}\right)}{24(x^{12})^{\frac{2}{3}}} + \frac{x^8 \sqrt{3} \arctan\left(\frac{\sqrt{3}(x^{12})^{\frac{1}{3}}}{2-(x^{12})^{\frac{1}{3}}}\right)}{12(x^{12})^{\frac{2}{3}}}$	80

input `int(x^7/(x^12+1),x,method=_RETURNVERBOSE)`output
$$-1/12*\ln(x^4+1)+1/24*\ln(x^8-x^4+1)+1/12*3^{(1/2)}*\arctan(2/3*(x^4-1/2)*3^{(1/2)})$$
Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.82

$$\int \frac{x^7}{1+x^{12}} dx = \frac{1}{12} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x^4-1)\right) + \frac{1}{24} \log(x^8-x^4+1) - \frac{1}{12} \log(x^4+1)$$

input `integrate(x^7/(x^12+1),x, algorithm="fricas")`output
$$1/12*\sqrt{3}*\arctan(1/3*\sqrt{3}*(2*x^4-1)) + 1/24*\log(x^8-x^4+1) - 1/12*\log(x^4+1)$$

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.94

$$\int \frac{x^7}{1+x^{12}} dx = -\frac{\log(x^4+1)}{12} + \frac{\log(x^8-x^4+1)}{24} + \frac{\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}x^4}{3} - \frac{\sqrt{3}}{3}\right)}{12}$$

input `integrate(x**7/(x**12+1),x)`output `-log(x**4 + 1)/12 + log(x**8 - x**4 + 1)/24 + sqrt(3)*atan(2*sqrt(3)*x**4/
3 - sqrt(3)/3)/12`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.82

$$\int \frac{x^7}{1+x^{12}} dx = \frac{1}{12} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x^4-1)\right) + \frac{1}{24} \log(x^8-x^4+1) - \frac{1}{12} \log(x^4+1)$$

input `integrate(x^7/(x^12+1),x, algorithm="maxima")`output `1/12*sqrt(3)*arctan(1/3*sqrt(3)*(2*x^4 - 1)) + 1/24*log(x^8 - x^4 + 1) - 1
/12*log(x^4 + 1)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.82

$$\int \frac{x^7}{1+x^{12}} dx = \frac{1}{12} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x^4-1)\right) + \frac{1}{24} \log(x^8-x^4+1) - \frac{1}{12} \log(x^4+1)$$

input `integrate(x^7/(x^12+1),x, algorithm="giac")`output `1/12*sqrt(3)*arctan(1/3*sqrt(3)*(2*x^4 - 1)) + 1/24*log(x^8 - x^4 + 1) - 1
/12*log(x^4 + 1)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.06

$$\int \frac{x^7}{1+x^{12}} dx = -\frac{\ln(x^4+1)}{12} - \ln\left(x^4 - \frac{\sqrt{3}i}{2} - \frac{1}{2}\right) \left(-\frac{1}{24} + \frac{\sqrt{3}i}{24}\right) \\ + \ln\left(x^4 + \frac{\sqrt{3}i}{2} - \frac{1}{2}\right) \left(\frac{1}{24} + \frac{\sqrt{3}i}{24}\right)$$

input `int(x^7/(x^12 + 1),x)`output `log((3^(1/2)*1i)/2 + x^4 - 1/2)*((3^(1/2)*1i)/24 + 1/24) - log(x^4 - (3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/24 - 1/24) - log(x^4 + 1)/12`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 353, normalized size of antiderivative = 7.20

$$\begin{aligned}
\int \frac{x^7}{1+x^{12}} dx = & -\frac{\sqrt{-\sqrt{3}+2}\sqrt{6}\operatorname{atan}\left(\frac{\sqrt{6}+\sqrt{2}-4x}{2\sqrt{-\sqrt{3}+2}}\right)}{24} - \frac{\sqrt{-\sqrt{3}+2}\sqrt{2}\operatorname{atan}\left(\frac{\sqrt{6}+\sqrt{2}-4x}{2\sqrt{-\sqrt{3}+2}}\right)}{8} \\
& - \frac{\sqrt{-\sqrt{3}+2}\sqrt{6}\operatorname{atan}\left(\frac{\sqrt{6}+\sqrt{2}+4x}{2\sqrt{-\sqrt{3}+2}}\right)}{24} - \frac{\sqrt{-\sqrt{3}+2}\sqrt{2}\operatorname{atan}\left(\frac{\sqrt{6}+\sqrt{2}+4x}{2\sqrt{-\sqrt{3}+2}}\right)}{8} \\
& + \frac{\sqrt{-\sqrt{3}+2}\sqrt{6}\operatorname{atan}\left(\frac{2\sqrt{-\sqrt{3}+2}-4x}{\sqrt{6}+\sqrt{2}}\right)}{24} \\
& + \frac{\sqrt{-\sqrt{3}+2}\sqrt{2}\operatorname{atan}\left(\frac{2\sqrt{-\sqrt{3}+2}-4x}{\sqrt{6}+\sqrt{2}}\right)}{8} \\
& + \frac{\sqrt{-\sqrt{3}+2}\sqrt{6}\operatorname{atan}\left(\frac{2\sqrt{-\sqrt{3}+2}+4x}{\sqrt{6}+\sqrt{2}}\right)}{24} \\
& + \frac{\sqrt{-\sqrt{3}+2}\sqrt{2}\operatorname{atan}\left(\frac{2\sqrt{-\sqrt{3}+2}+4x}{\sqrt{6}+\sqrt{2}}\right)}{8} \\
& + \frac{\log\left(-\sqrt{-\sqrt{3}+2}x+x^2+1\right)}{24} - \frac{\log\left(-\sqrt{2}x+x^2+1\right)}{12} \\
& + \frac{\log\left(\sqrt{-\sqrt{3}+2}x+x^2+1\right)}{24} - \frac{\log\left(\sqrt{2}x+x^2+1\right)}{12} \\
& + \frac{\log\left(-\frac{\sqrt{6}x}{2}-\frac{\sqrt{2}x}{2}+x^2+1\right)}{24} + \frac{\log\left(\frac{\sqrt{6}x}{2}+\frac{\sqrt{2}x}{2}+x^2+1\right)}{24}
\end{aligned}$$

input `int(x^7/(x^12+1),x)`

output

```
( - sqrt( - sqrt(3) + 2)*sqrt(6)*atan((sqrt(6) + sqrt(2) - 4*x)/(2*sqrt( -
sqrt(3) + 2))) - 3*sqrt( - sqrt(3) + 2)*sqrt(2)*atan((sqrt(6) + sqrt(2) -
4*x)/(2*sqrt( - sqrt(3) + 2))) - sqrt( - sqrt(3) + 2)*sqrt(6)*atan((sqrt(
6) + sqrt(2) + 4*x)/(2*sqrt( - sqrt(3) + 2))) - 3*sqrt( - sqrt(3) + 2)*sqr
t(2)*atan((sqrt(6) + sqrt(2) + 4*x)/(2*sqrt( - sqrt(3) + 2))) + sqrt( - sq
rt(3) + 2)*sqrt(6)*atan((2*sqrt( - sqrt(3) + 2) - 4*x)/(sqrt(6) + sqrt(2))
) + 3*sqrt( - sqrt(3) + 2)*sqrt(2)*atan((2*sqrt( - sqrt(3) + 2) - 4*x)/(sq
rt(6) + sqrt(2))) + sqrt( - sqrt(3) + 2)*sqrt(6)*atan((2*sqrt( - sqrt(3) +
2) + 4*x)/(sqrt(6) + sqrt(2))) + 3*sqrt( - sqrt(3) + 2)*sqrt(2)*atan((2*s
qrt( - sqrt(3) + 2) + 4*x)/(sqrt(6) + sqrt(2))) + log( - sqrt( - sqrt(3) +
2)*x + x**2 + 1) - 2*log( - sqrt(2)*x + x**2 + 1) + log(sqrt( - sqrt(3) +
2)*x + x**2 + 1) - 2*log(sqrt(2)*x + x**2 + 1) + log((- sqrt(6)*x - sqrt
(2)*x + 2*x**2 + 2)/2) + log((sqrt(6)*x + sqrt(2)*x + 2*x**2 + 2)/2))/24
```

3.32 $\int x^{3a} \sin(x^{2a}) dx$

Optimal result	229
Mathematica [A] (verified)	229
Rubi [A] (verified)	230
Maple [C] (verified)	231
Fricas [F]	231
Sympy [A] (verification not implemented)	232
Maxima [F]	232
Giac [F]	232
Mupad [F(-1)]	233
Reduce [F]	233

Optimal result

Integrand size = 12, antiderivative size = 115

$$\int x^{3a} \sin(x^{2a}) dx = \frac{ix^{1+3a}(-ix^{2a})^{-\frac{1+3a}{2a}} \Gamma(\frac{1}{2}(3 + \frac{1}{a}), -ix^{2a})}{4a} - \frac{ix^{1+3a}(ix^{2a})^{-\frac{1+3a}{2a}} \Gamma(\frac{1}{2}(3 + \frac{1}{a}), ix^{2a})}{4a}$$

output

```
1/4*I*x^(1+3*a)*GAMMA(3/2+1/2/a,-I*x^(2*a))/a/((-I*x^(2*a))^(1/2*(1+3*a)/a))-1/4*I*x^(1+3*a)*GAMMA(3/2+1/2/a,I*x^(2*a))/a/((I*x^(2*a))^(1/2*(1+3*a)/a))
```

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.23

$$\int x^{3a} \sin(x^{2a}) dx = \frac{x^{1+a}(x^{4a})^{-\frac{1+a}{2a}} \left(4a(x^{4a})^{\frac{1+a}{2a}} \cos(x^{2a}) + (1+a)(ix^{2a})^{\frac{1+a}{2a}} \Gamma(\frac{1+a}{2a}, -ix^{2a}) + (1+a)(-ix^{2a})^{\frac{1+a}{2a}} \Gamma(\frac{1+a}{2a}, ix^{2a}) \right)}{8a^2}$$

input

```
Integrate[x^(3*a)*Sin[x^(2*a)],x]
```

output

```
-1/8*(x^(1 + a)*(4*a*(x^(4*a))^((1 + a)/(2*a))*Cos[x^(2*a)] + (1 + a)*(I*x
^(2*a))^((1 + a)/(2*a))*Gamma[(1 + a)/(2*a), (-I)*x^(2*a)] + (1 + a)*((-I)
*x^(2*a))^((1 + a)/(2*a))*Gamma[(1 + a)/(2*a), I*x^(2*a)]))/(a^2*(x^(4*a))
^((1 + a)/(2*a)))
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3904, 2648}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{3a} \sin(x^{2a}) dx$$

↓ 3904

$$\frac{1}{2}i \int e^{-ix^{2a}} x^{3a} dx - \frac{1}{2}i \int e^{ix^{2a}} x^{3a} dx$$

↓ 2648

$$\frac{ix^{3a+1}(-ix^{2a})^{-\frac{3a+1}{2a}} \Gamma(\frac{1}{2}(3 + \frac{1}{a}), -ix^{2a})}{4a} - \frac{ix^{3a+1}(ix^{2a})^{-\frac{3a+1}{2a}} \Gamma(\frac{1}{2}(3 + \frac{1}{a}), ix^{2a})}{4a}$$

input

```
Int[x^(3*a)*Sin[x^(2*a)],x]
```

output

```
((I/4)*x^(1 + 3*a)*Gamma[(3 + a^(-1))/2, (-I)*x^(2*a)]/(a*((-I)*x^(2*a))^
((1 + 3*a)/(2*a))) - ((I/4)*x^(1 + 3*a)*Gamma[(3 + a^(-1))/2, I*x^(2*a)]/
(a*(I*x^(2*a))^((1 + 3*a)/(2*a))))
```

Defintions of rubi rules used

rule 2648

```
Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^(n_)))*((e_.) + (f_.)*(x_)^(m_
.), x_Symbol] := Simp[(-F^a)*((e + f*x)^(m + 1)/(f*n*((-b)*(c + d*x)^n*Log[
F])^((m + 1)/n)))*Gamma[(m + 1)/n, (-b)*(c + d*x)^n*Log[F]], x] /; FreeQ[{F
, a, b, c, d, e, f, m, n}, x] && EqQ[d*e - c*f, 0]
```

rule 3904

```
Int[((e_.)*(x_))^(m_.)*Sin[(c_.) + (d_.)*(x_)^(n_)], x_Symbol] := Simp[I/2
  Int[(e*x)^m*E^((-c)*I - d*I*x^n), x], x] - Simp[I/2  Int[(e*x)^m*E^(c*I
+ d*I*x^n), x], x] /; FreeQ[{c, d, e, m, n}, x]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4.

Time = 0.28 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.36

method	result	size
meijerg	$\frac{x^{5a+1} \operatorname{hypergeom}\left(\left[\frac{5}{4} + \frac{1}{4a}\right], \left[\frac{3}{2}, \frac{9}{4} + \frac{1}{4a}\right], -\frac{x^{4a}}{4}\right)}{5a+1}$	41

input

```
int(x^(3*a)*sin(x^(2*a)),x,method=_RETURNVERBOSE)
```

output

```
1/(5*a+1)*x^(5*a+1)*hypergeom([5/4+1/4/a], [3/2, 9/4+1/4/a], -1/4*x^(4*a))
```

Fricas [F]

$$\int x^{3a} \sin(x^{2a}) dx = \int x^{3a} \sin(x^{2a}) dx$$

input

```
integrate(x^(3*a)*sin(x^(2*a)),x, algorithm="fricas")
```

output

```
integral(x^(3*a)*sin(x^(2*a)), x)
```


Sympy [A] (verification not implemented)

Time = 1.39 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.47

$$\int x^{3a} \sin(x^{2a}) dx = \frac{x^{5a+1} \Gamma\left(\frac{5}{4} + \frac{1}{4a}\right) {}_1F_2\left(\frac{5}{4} + \frac{1}{4a} \mid -\frac{x^{4a}}{4}\right)}{4a \Gamma\left(\frac{9}{4} + \frac{1}{4a}\right)}$$

input `integrate(x**(3*a)*sin(x**(2*a)),x)`output `x**(5*a + 1)*gamma(5/4 + 1/(4*a))*hyper((5/4 + 1/(4*a)), (3/2, 9/4 + 1/(4*a)), -x**(4*a)/4)/(4*a*gamma(9/4 + 1/(4*a)))`**Maxima [F]**

$$\int x^{3a} \sin(x^{2a}) dx = \int x^{3a} \sin(x^{2a}) dx$$

input `integrate(x^(3*a)*sin(x^(2*a)),x, algorithm="maxima")`output `-1/2*(x*x^a*cos(x^(2*a)) - (a + 1)*integrate(x^a*cos(x^(2*a)), x))/a`**Giac [F]**

$$\int x^{3a} \sin(x^{2a}) dx = \int x^{3a} \sin(x^{2a}) dx$$

input `integrate(x^(3*a)*sin(x^(2*a)),x, algorithm="giac")`output `integrate(x^(3*a)*sin(x^(2*a)), x)`

Mupad [F(-1)]

Timed out.

$$\int x^{3a} \sin(x^{2a}) dx = \int x^{3a} \sin(x^{2a}) dx$$

input `int(x^(3*a)*sin(x^(2*a)),x)`output `int(x^(3*a)*sin(x^(2*a)), x)`**Reduce [F]**

$$\int x^{3a} \sin(x^{2a}) dx$$

$$= \frac{-x^a x + \left(\int \frac{x^a}{\tan\left(\frac{x^{2a}}{2}\right)^2 + 1} dx \right) \tan\left(\frac{x^{2a}}{2}\right)^2 a + \left(\int \frac{x^a}{\tan\left(\frac{x^{2a}}{2}\right)^2 + 1} dx \right) \tan\left(\frac{x^{2a}}{2}\right)^2 + \left(\int \frac{x^a}{\tan\left(\frac{x^{2a}}{2}\right)^2 + 1} dx \right) a + \int \frac{x^a}{\tan\left(\frac{x^{2a}}{2}\right)^2 + 1} dx}{a \left(\tan\left(\frac{x^{2a}}{2}\right)^2 + 1 \right)}$$

input `int(x^(3*a)*sin(x^(2*a)),x)`output `(- x**a*x + int(x**a/(tan(x**(2*a)/2)**2 + 1),x)*tan(x**(2*a)/2)**2*a + int(x**a/(tan(x**(2*a)/2)**2 + 1),x)*tan(x**(2*a)/2)**2 + int(x**a/(tan(x**(2*a)/2)**2 + 1),x)*a + int(x**a/(tan(x**(2*a)/2)**2 + 1),x))/(a*(tan(x**(2*a)/2)**2 + 1))`

3.33 $\int \cos(\sqrt{x}) dx$

Optimal result	234
Mathematica [A] (verified)	234
Rubi [A] (verified)	235
Maple [A] (verified)	236
Fricas [A] (verification not implemented)	237
Sympy [A] (verification not implemented)	237
Maxima [A] (verification not implemented)	237
Giac [A] (verification not implemented)	238
Mupad [B] (verification not implemented)	238
Reduce [B] (verification not implemented)	238

Optimal result

Integrand size = 6, antiderivative size = 22

$$\int \cos(\sqrt{x}) dx = 2 \cos(\sqrt{x}) + 2\sqrt{x} \sin(\sqrt{x})$$

output `2*cos(x^(1/2))+2*sin(x^(1/2))*x^(1/2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \cos(\sqrt{x}) dx = 2 \cos(\sqrt{x}) + 2\sqrt{x} \sin(\sqrt{x})$$

input `Integrate[Cos[Sqrt[x]],x]`

output `2*Cos[Sqrt[x]] + 2*Sqrt[x]*Sin[Sqrt[x]]`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.95, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3843, 3042, 3777, 25, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(\sqrt{x}) \, dx \\
 & \quad \downarrow \text{3843} \\
 & 2 \int \sqrt{x} \cos(\sqrt{x}) \, d\sqrt{x} \\
 & \quad \downarrow \text{3042} \\
 & 2 \int \sqrt{x} \sin\left(\sqrt{x} + \frac{\pi}{2}\right) \, d\sqrt{x} \\
 & \quad \downarrow \text{3777} \\
 & 2 \left(\int -\sin(\sqrt{x}) \, d\sqrt{x} + \sqrt{x} \sin(\sqrt{x}) \right) \\
 & \quad \downarrow \text{25} \\
 & 2 \left(\sqrt{x} \sin(\sqrt{x}) - \int \sin(\sqrt{x}) \, d\sqrt{x} \right) \\
 & \quad \downarrow \text{3042} \\
 & 2 \left(\sqrt{x} \sin(\sqrt{x}) - \int \sin(\sqrt{x}) \, d\sqrt{x} \right) \\
 & \quad \downarrow \text{3118} \\
 & 2(\sqrt{x} \sin(\sqrt{x}) + \cos(\sqrt{x}))
 \end{aligned}$$

input

Int[Cos[Sqrt[x]], x]

output

2*(Cos[Sqrt[x]] + Sqrt[x]*Sin[Sqrt[x]])

Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`
- rule 3777 `Int[((c_.) + (d_.)*(x_)^(m_.))*sin[(e_.) + (f_.)*(x_)], x_Symbol] :> Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`
- rule 3843 `Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_)^(n_)])*(b_.))^(p_.), x_Symbol] :> Simp[1/(n*f) Subst[Int[x^(1/n - 1)*(a + b*Cos[c + d*x])^p, x], x, (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[1/n]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

method	result	size
derivativedivides	$2 \cos(\sqrt{x}) + 2 \sin(\sqrt{x}) \sqrt{x}$	17
default	$2 \cos(\sqrt{x}) + 2 \sin(\sqrt{x}) \sqrt{x}$	17
meijerg	$4\sqrt{\pi} \left(-\frac{1}{2\sqrt{\pi}} + \frac{\cos(\sqrt{x})}{2\sqrt{\pi}} + \frac{\sqrt{x} \sin(\sqrt{x})}{2\sqrt{\pi}} \right)$	33

input `int(cos(x^(1/2)),x,method=_RETURNVERBOSE)`

output `2*cos(x^(1/2))+2*sin(x^(1/2))*x^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \cos(\sqrt{x}) dx = 2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

input `integrate(cos(x^(1/2)),x, algorithm="fricas")`

output `2*sqrt(x)*sin(sqrt(x)) + 2*cos(sqrt(x))`

Sympy [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \cos(\sqrt{x}) dx = 2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

input `integrate(cos(x**(1/2)),x)`

output `2*sqrt(x)*sin(sqrt(x)) + 2*cos(sqrt(x))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \cos(\sqrt{x}) dx = 2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

input `integrate(cos(x^(1/2)),x, algorithm="maxima")`

output `2*sqrt(x)*sin(sqrt(x)) + 2*cos(sqrt(x))`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \cos(\sqrt{x}) dx = 2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

input `integrate(cos(x^(1/2)),x, algorithm="giac")`

output `2*sqrt(x)*sin(sqrt(x)) + 2*cos(sqrt(x))`

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \cos(\sqrt{x}) dx = 2 \cos(\sqrt{x}) + 2\sqrt{x} \sin(\sqrt{x})$$

input `int(cos(x^(1/2)),x)`

output `2*cos(x^(1/2)) + 2*x^(1/2)*sin(x^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.59

$$\int \cos(\sqrt{x}) dx = 2 \cos(\sqrt{x}) + 2\sqrt{x} \sin(\sqrt{x})$$

input `int(cos(x^(1/2)),x)`

output `2*(cos(sqrt(x)) + sqrt(x)*sin(sqrt(x)))`

3.34 $\int x\sqrt{1+x} dx$

Optimal result	239
Mathematica [A] (verified)	239
Rubi [A] (verified)	240
Maple [A] (verified)	241
Fricas [A] (verification not implemented)	241
Sympy [A] (verification not implemented)	242
Maxima [A] (verification not implemented)	242
Giac [A] (verification not implemented)	242
Mupad [B] (verification not implemented)	243
Reduce [B] (verification not implemented)	243

Optimal result

Integrand size = 9, antiderivative size = 23

$$\int x\sqrt{1+x} dx = -\frac{2}{3}(1+x)^{3/2} + \frac{2}{5}(1+x)^{5/2}$$

output `-2/3*(1+x)^(3/2)+2/5*(1+x)^(5/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

$$\int x\sqrt{1+x} dx = \frac{2}{15}(1+x)^{3/2}(-5+3(1+x))$$

input `Integrate[x*Sqrt[1 + x],x]`

output `(2*(1 + x)^(3/2)*(-5 + 3*(1 + x)))/15`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x\sqrt{x+1} dx$$

↓ 53

$$\int \left((x+1)^{3/2} - \sqrt{x+1} \right) dx$$

↓ 2009

$$\frac{2}{5}(x+1)^{5/2} - \frac{2}{3}(x+1)^{3/2}$$

input `Int[x*Sqrt[1 + x],x]`

output `(-2*(1 + x)^(3/2))/3 + (2*(1 + x)^(5/2))/5`

Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.57

method	result	size
gospers	$\frac{2(1+x)^{\frac{3}{2}}(-2+3x)}{15}$	13
orering	$\frac{2(1+x)^{\frac{3}{2}}(-2+3x)}{15}$	13
derivativedivides	$-\frac{2(1+x)^{\frac{3}{2}}}{3} + \frac{2(1+x)^{\frac{5}{2}}}{5}$	16
default	$-\frac{2(1+x)^{\frac{3}{2}}}{3} + \frac{2(1+x)^{\frac{5}{2}}}{5}$	16
risch	$\frac{2(3x^2+x-2)\sqrt{1+x}}{15}$	16
trager	$\left(\frac{2}{5}x^2 + \frac{2}{15}x - \frac{4}{15}\right)\sqrt{1+x}$	17
meijerg	$-\frac{8\sqrt{\pi}}{15} + \frac{4\sqrt{\pi}(1+x)^{\frac{3}{2}}(2-3x)}{2\sqrt{\pi}}$	27

input `int(x*(1+x)^(1/2),x,method=_RETURNVERBOSE)`output `2/15*(1+x)^(3/2)*(-2+3*x)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int x\sqrt{1+x} dx = \frac{2}{15} (3x^2 + x - 2)\sqrt{x+1}$$

input `integrate(x*(1+x)^(1/2),x, algorithm="fricas")`output `2/15*(3*x^2 + x - 2)*sqrt(x + 1)`

Sympy [A] (verification not implemented)

Time = 0.48 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.48

$$\int x\sqrt{1+x} dx = \frac{2x^2\sqrt{x+1}}{5} + \frac{2x\sqrt{x+1}}{15} - \frac{4\sqrt{x+1}}{15}$$

input `integrate(x*(1+x)**(1/2),x)`

output `2*x**2*sqrt(x + 1)/5 + 2*x*sqrt(x + 1)/15 - 4*sqrt(x + 1)/15`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int x\sqrt{1+x} dx = \frac{2}{5}(x+1)^{\frac{5}{2}} - \frac{2}{3}(x+1)^{\frac{3}{2}}$$

input `integrate(x*(1+x)^(1/2),x, algorithm="maxima")`

output `2/5*(x + 1)^(5/2) - 2/3*(x + 1)^(3/2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int x\sqrt{1+x} dx = \frac{2}{5}(x+1)^{\frac{5}{2}} - \frac{2}{3}(x+1)^{\frac{3}{2}}$$

input `integrate(x*(1+x)^(1/2),x, algorithm="giac")`

output `2/5*(x + 1)^(5/2) - 2/3*(x + 1)^(3/2)`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.52

$$\int x\sqrt{1+x} dx = \frac{2(3x-2)(x+1)^{3/2}}{15}$$

input `int(x*(x + 1)^(1/2),x)`

output `(2*(3*x - 2)*(x + 1)^(3/2))/15`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.61

$$\int x\sqrt{1+x} dx = \frac{2\sqrt{x+1}(3x^2+x-2)}{15}$$

input `int(x*(1+x)^(1/2),x)`

output `(2*sqrt(x + 1)*(3*x**2 + x - 2))/15`

3.35 $\int \frac{1}{\sqrt[3]{x} + \sqrt{x}} dx$

Optimal result	244
Mathematica [A] (verified)	244
Rubi [A] (verified)	245
Maple [A] (verified)	246
Fricas [A] (verification not implemented)	247
Sympy [F]	247
Maxima [A] (verification not implemented)	247
Giac [A] (verification not implemented)	248
Mupad [B] (verification not implemented)	248
Reduce [B] (verification not implemented)	248

Optimal result

Integrand size = 13, antiderivative size = 32

$$\int \frac{1}{\sqrt[3]{x} + \sqrt{x}} dx = 6\sqrt[6]{x} - 3\sqrt[3]{x} + 2\sqrt{x} - 6 \log(1 + \sqrt[6]{x})$$

output `6*x^(1/6)-3*x^(1/3)-6*ln(1+x^(1/6))+2*x^(1/2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.03

$$\int \frac{1}{\sqrt[3]{x} + \sqrt{x}} dx = (6 - 3\sqrt[6]{x} + 2\sqrt[3]{x}) \sqrt[6]{x} - 6 \log(1 + \sqrt[6]{x})$$

input `Integrate[(x^(1/3) + Sqrt[x])^(-1), x]`

output `(6 - 3*x^(1/6) + 2*x^(1/3))*x^(1/6) - 6*Log[1 + x^(1/6)]`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.12, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {2027, 798, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{x} + \sqrt[3]{x}} dx \\
 & \quad \downarrow \text{2027} \\
 & \int \frac{1}{(\sqrt[6]{x} + 1) \sqrt[3]{x}} dx \\
 & \quad \downarrow \text{798} \\
 & 6 \int \frac{\sqrt{x}}{\sqrt[6]{x} + 1} d\sqrt[6]{x} \\
 & \quad \downarrow \text{49} \\
 & 6 \int \left(\sqrt[3]{x} - \sqrt[6]{x} + \frac{1}{-\sqrt[6]{x} - 1} + 1 \right) d\sqrt[6]{x} \\
 & \quad \downarrow \text{2009} \\
 & 6 \left(\frac{\sqrt{x}}{3} - \frac{\sqrt[3]{x}}{2} + \sqrt[6]{x} - \log(\sqrt[6]{x} + 1) \right)
 \end{aligned}$$

input `Int[(x^(1/3) + Sqrt[x])^(-1),x]`

output `6*(x^(1/6) - x^(1/3)/2 + Sqrt[x]/3 - Log[1 + x^(1/6)])`

Definitions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/n Subst
[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a,
b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2027 `Int[(Fx_.)*((a_.)*(x_)^(r_.) + (b_.)*(x_)^(s_.))^(p_.), x_Symbol] := Int[x^(
(p*r)*(a + b*x^(s - r))^p*Fx, x] /; FreeQ[{a, b, r, s}, x] && IntegerQ[p] &
&& PosQ[s - r] && !(EqQ[p, 1] && EqQ[u, 1])`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.78

method	result
derivativedivides	$6x^{\frac{1}{6}} - 3x^{\frac{1}{3}} - 6 \ln(1 + x^{\frac{1}{6}}) + 2\sqrt{x}$
meijerg	$\frac{x^{\frac{1}{6}}(4x^{\frac{1}{3}} - 6x^{\frac{1}{6}} + 12)}{2} - 6 \ln(1 + x^{\frac{1}{6}})$
default	$-\ln(x^{\frac{1}{3}} + x^{\frac{1}{6}} + 1) + 2 \ln(x^{\frac{1}{6}} - 1) + \ln(x^{\frac{1}{3}} - x^{\frac{1}{6}} + 1) - 2 \ln(1 + x^{\frac{1}{6}}) + 2\sqrt{x} + \ln$

input `int(1/(x^(1/3)+x^(1/2)),x,method=_RETURNVERBOSE)`

output `6*x^(1/6)-3*x^(1/3)-6*ln(1+x^(1/6))+2*x^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt[3]{x} + \sqrt{x}} dx = 2\sqrt{x} - 3x^{\frac{1}{3}} + 6x^{\frac{1}{6}} - 6 \log(x^{\frac{1}{6}} + 1)$$

input `integrate(1/(x^(1/3)+x^(1/2)),x, algorithm="fricas")`output `2*sqrt(x) - 3*x^(1/3) + 6*x^(1/6) - 6*log(x^(1/6) + 1)`**Sympy [F]**

$$\int \frac{1}{\sqrt[3]{x} + \sqrt{x}} dx = \int \frac{1}{\sqrt[3]{x} + \sqrt{x}} dx$$

input `integrate(1/(x**(1/3)+x**(1/2)),x)`output `Integral(1/(x**(1/3) + sqrt(x)), x)`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt[3]{x} + \sqrt{x}} dx = 2\sqrt{x} - 3x^{\frac{1}{3}} + 6x^{\frac{1}{6}} - 6 \log(x^{\frac{1}{6}} + 1)$$

input `integrate(1/(x^(1/3)+x^(1/2)),x, algorithm="maxima")`output `2*sqrt(x) - 3*x^(1/3) + 6*x^(1/6) - 6*log(x^(1/6) + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt[3]{x} + \sqrt{x}} dx = 2\sqrt{x} - 3x^{\frac{1}{3}} + 6x^{\frac{1}{6}} - 6\log\left(x^{\frac{1}{6}} + 1\right)$$

input `integrate(1/(x^(1/3)+x^(1/2)),x, algorithm="giac")`output `2*sqrt(x) - 3*x^(1/3) + 6*x^(1/6) - 6*log(x^(1/6) + 1)`**Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt[3]{x} + \sqrt{x}} dx = 2\sqrt{x} - 6\ln\left(x^{1/6} + 1\right) - 3x^{1/3} + 6x^{1/6}$$

input `int(1/(x^(1/2) + x^(1/3)),x)`output `2*x^(1/2) - 6*log(x^(1/6) + 1) - 3*x^(1/3) + 6*x^(1/6)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.72

$$\int \frac{1}{\sqrt[3]{x} + \sqrt{x}} dx = 6x^{\frac{1}{6}} - 3x^{\frac{1}{3}} + 2\sqrt{x} - 6\log\left(x^{\frac{1}{6}} + 1\right)$$

input `int(1/(x^(1/3)+x^(1/2)),x)`output `6*x**(1/6) - 3*x**(1/3) + 2*sqrt(x) - 6*log(x**(1/6) + 1)`

3.36 $\int \sqrt{\frac{1+x}{3+2x}} dx$

Optimal result	249
Mathematica [A] (verified)	249
Rubi [A] (verified)	250
Maple [B] (verified)	251
Fricas [A] (verification not implemented)	252
Sympy [F]	252
Maxima [B] (verification not implemented)	253
Giac [B] (verification not implemented)	253
Mupad [B] (verification not implemented)	254
Reduce [B] (verification not implemented)	254

Optimal result

Integrand size = 15, antiderivative size = 44

$$\int \sqrt{\frac{1+x}{3+2x}} dx = \frac{1}{2} \sqrt{1+x} \sqrt{3+2x} - \frac{\operatorname{arcsinh}(\sqrt{2}\sqrt{1+x})}{2\sqrt{2}}$$

output

```
-1/4*arcsinh(2^(1/2)*(1+x)^(1/2))*2^(1/2)+1/2*(1+x)^(1/2)*(3+2*x)^(1/2)
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.70

$$\int \sqrt{\frac{1+x}{3+2x}} dx = \frac{\sqrt{\frac{1+x}{3+2x}} \left(2\sqrt{1+x}(3+2x) - \sqrt{6+4x} \operatorname{arctanh}\left(\frac{\sqrt{3+2x}}{\sqrt{2}\sqrt{1+x}}\right) \right)}{4\sqrt{1+x}}$$

input

```
Integrate[Sqrt[(1 + x)/(3 + 2*x)], x]
```

output

```
(Sqrt[(1 + x)/(3 + 2*x)]*(2*Sqrt[1 + x]*(3 + 2*x) - Sqrt[6 + 4*x]*ArcTanh[
Sqrt[3 + 2*x]/(Sqrt[2]*Sqrt[1 + x])]))/(4*Sqrt[1 + x])
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {2050, 60, 64, 222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\frac{x+1}{2x+3}} dx \\
 & \quad \downarrow \text{2050} \\
 & \int \frac{\sqrt{x+1}}{\sqrt{2x+3}} dx \\
 & \quad \downarrow \text{60} \\
 & \frac{1}{2}\sqrt{x+1}\sqrt{2x+3} - \frac{1}{4} \int \frac{1}{\sqrt{x+1}\sqrt{2x+3}} dx \\
 & \quad \downarrow \text{64} \\
 & \frac{1}{2}\sqrt{x+1}\sqrt{2x+3} - \frac{1}{2} \int \frac{1}{\sqrt{2(x+1)+1}} d\sqrt{x+1} \\
 & \quad \downarrow \text{222} \\
 & \frac{1}{2}\sqrt{x+1}\sqrt{2x+3} - \frac{\operatorname{arcsinh}(\sqrt{2}\sqrt{x+1})}{2\sqrt{2}}
 \end{aligned}$$

input `Int[Sqrt[(1 + x)/(3 + 2*x)],x]`

output `(Sqrt[1 + x]*Sqrt[3 + 2*x])/2 - ArcSinh[Sqrt[2]*Sqrt[1 + x]]/(2*Sqrt[2])`

Defintions of rubi rules used

```
rule 60 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(
b*(m + n + 1))) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && ( !Integer
Q[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinear
Q[a, b, c, d, m, n, x]
```

```
rule 64 Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp
[2/b Subst[Int[1/Sqrt[c - a*(d/b) + d*(x^2/b)], x], x, Sqrt[a + b*x]], x]
/; FreeQ[{a, b, c, d}, x] && GtQ[c - a*(d/b), 0] && ( !GtQ[a - c*(b/d), 0]
|| PosQ[b])
```

```
rule 222 Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt
[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]
```

```
rule 2050 Int[(u_.)*(((e_.)*((a_.) + (b_.)*(x_)^(n_.)))/((c_.) + (d_.)*(x_)^(n_.)))^(p
_), x_Symbol] := Int[u*((a*e + b*e*x^n)^p/(c + d*x^n)^p), x] /; FreeQ[{a, b,
c, d, e, n, p}, x] && GtQ[b*d*e, 0] && GtQ[c - a*(d/b), 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 74 vs. 2(30) = 60.

Time = 0.14 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.70

method	result
default	$-\frac{\sqrt{\frac{1+x}{3+2x}}(3+2x)\left(\ln\left(\frac{5\sqrt{2}}{4}+x\sqrt{2+\sqrt{2x^2+5x+3}}\right)\sqrt{2-4\sqrt{2x^2+5x+3}}\right)}{8\sqrt{(3+2x)(1+x)}}$
risch	$\frac{(3+2x)\sqrt{\frac{1+x}{3+2x}}}{2} - \frac{\ln\left(\frac{(\frac{5}{2}+2x)\sqrt{2}}{2} + \sqrt{2x^2+5x+3}\right)\sqrt{2}\sqrt{\frac{1+x}{3+2x}}\sqrt{(3+2x)(1+x)}}{8(1+x)}$
trager	$3\left(\frac{1}{2} + \frac{x}{3}\right)\sqrt{-\frac{-1-x}{3+2x}} - \frac{\text{RootOf}(_Z^2-2)\ln\left(4\text{RootOf}(_Z^2-2)x+8\sqrt{-\frac{-1-x}{3+2x}}x+5\text{RootOf}(_Z^2-2)+12\sqrt{-\frac{-1-x}{3+2x}}\right)}{8}$

input `int(((1+x)/(3+2*x))^(1/2),x,method=_RETURNVERBOSE)`

output `-1/8*((1+x)/(3+2*x))^(1/2)*(3+2*x)*(ln(5/4*2^(1/2)+x*2^(1/2)+(2*x^2+5*x+3)^(1/2))*2^(1/2)-4*(2*x^2+5*x+3)^(1/2))/((3+2*x)*(1+x))^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.25

$$\int \sqrt{\frac{1+x}{3+2x}} dx = \frac{1}{2} (2x+3) \sqrt{\frac{x+1}{2x+3}} + \frac{1}{8} \sqrt{2} \log \left(2\sqrt{2}(2x+3) \sqrt{\frac{x+1}{2x+3}} - 4x - 5 \right)$$

input `integrate(((1+x)/(3+2*x))^(1/2),x, algorithm="fricas")`

output `1/2*(2*x + 3)*sqrt((x + 1)/(2*x + 3)) + 1/8*sqrt(2)*log(2*sqrt(2)*(2*x + 3)*sqrt((x + 1)/(2*x + 3)) - 4*x - 5)`

Sympy [F]

$$\int \sqrt{\frac{1+x}{3+2x}} dx = \int \sqrt{\frac{x+1}{2x+3}} dx$$

input `integrate(((1+x)/(3+2*x))**(1/2),x)`

output `Integral(sqrt((x + 1)/(2*x + 3)), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 80 vs. $2(30) = 60$.

Time = 0.11 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.82

$$\int \sqrt{\frac{1+x}{3+2x}} dx = \frac{1}{8} \sqrt{2} \log \left(-\frac{\sqrt{2} - 2 \sqrt{\frac{x+1}{2x+3}}}{\sqrt{2} + 2 \sqrt{\frac{x+1}{2x+3}}} \right) - \frac{\sqrt{\frac{x+1}{2x+3}}}{2 \left(\frac{2(x+1)}{2x+3} - 1 \right)}$$

input `integrate(((1+x)/(3+2*x))^(1/2),x, algorithm="maxima")`

output `1/8*sqrt(2)*log(-(sqrt(2) - 2*sqrt((x + 1)/(2*x + 3)))/(sqrt(2) + 2*sqrt((x + 1)/(2*x + 3)))) - 1/2*sqrt((x + 1)/(2*x + 3))/(2*(x + 1)/(2*x + 3) - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 61 vs. $2(30) = 60$.

Time = 0.13 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.39

$$\int \sqrt{\frac{1+x}{3+2x}} dx = \frac{1}{8} \sqrt{2} \log \left(\left| -2 \sqrt{2} \left(\sqrt{2x} - \sqrt{2x^2 + 5x + 3} \right) - 5 \right| \right) \operatorname{sgn}(2x + 3) + \frac{1}{2} \sqrt{2x^2 + 5x + 3} \operatorname{sgn}(2x + 3)$$

input `integrate(((1+x)/(3+2*x))^(1/2),x, algorithm="giac")`

output `1/8*sqrt(2)*log(abs(-2*sqrt(2)*(sqrt(2)*x - sqrt(2*x^2 + 5*x + 3)) - 5))*sgn(2*x + 3) + 1/2*sqrt(2*x^2 + 5*x + 3)*sgn(2*x + 3)`

Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.30

$$\int \sqrt{\frac{1+x}{3+2x}} dx = -\frac{\sqrt{2} \operatorname{atanh}\left(\sqrt{2} \sqrt{\frac{x+1}{2x+3}}\right)}{4} - \frac{\sqrt{\frac{x+1}{2x+3}}}{2\left(\frac{2x+2}{2x+3} - 1\right)}$$

input `int(((x + 1)/(2*x + 3))^(1/2),x)`output `-(2^(1/2)*atanh(2^(1/2)*((x + 1)/(2*x + 3))^(1/2)))/4 - ((x + 1)/(2*x + 3))^(1/2)/(2*((2*x + 2)/(2*x + 3) - 1))`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.73

$$\int \sqrt{\frac{1+x}{3+2x}} dx = \frac{\sqrt{x+1} \sqrt{2x+3}}{2} - \frac{\sqrt{2} \log(\sqrt{2x+3} + \sqrt{x+1} \sqrt{2})}{4}$$

input `int(((1+x)/(3+2*x))^(1/2),x)`output `(2*sqrt(x + 1)*sqrt(2*x + 3) - sqrt(2)*log(sqrt(2*x + 3) + sqrt(x + 1)*sqrt(2)))/4`

$$3.37 \quad \int \frac{x^4}{(1-x^2)^{5/2}} dx$$

Optimal result	255
Mathematica [A] (verified)	255
Rubi [A] (verified)	256
Maple [A] (verified)	257
Fricas [B] (verification not implemented)	257
Sympy [B] (verification not implemented)	258
Maxima [A] (verification not implemented)	258
Giac [A] (verification not implemented)	259
Mupad [B] (verification not implemented)	259
Reduce [B] (verification not implemented)	259

Optimal result

Integrand size = 15, antiderivative size = 35

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{x^3}{3(1-x^2)^{3/2}} - \frac{x}{\sqrt{1-x^2}} + \arcsin(x)$$

output `1/3*x^3/(-x^2+1)^(3/2)+arcsin(x)-x/(-x^2+1)^(1/2)`

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.26

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{x(-3+4x^2)}{3(1-x^2)^{3/2}} + 2 \arctan\left(\frac{x}{-1+\sqrt{1-x^2}}\right)$$

input `Integrate[x^4/(1-x^2)^(5/2),x]`

output `(x*(-3+4*x^2))/(3*(1-x^2)^(3/2))+2*ArcTan[x/(-1+Sqrt[1-x^2])]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {252, 252, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^4}{(1-x^2)^{5/2}} dx \\
 & \quad \downarrow \text{252} \\
 & \frac{x^3}{3(1-x^2)^{3/2}} - \int \frac{x^2}{(1-x^2)^{3/2}} dx \\
 & \quad \downarrow \text{252} \\
 & \int \frac{1}{\sqrt{1-x^2}} dx - \frac{x}{\sqrt{1-x^2}} + \frac{x^3}{3(1-x^2)^{3/2}} \\
 & \quad \downarrow \text{223} \\
 & \arcsin(x) - \frac{x}{\sqrt{1-x^2}} + \frac{x^3}{3(1-x^2)^{3/2}}
 \end{aligned}$$

input `Int[x^4/(1 - x^2)^(5/2),x]`

output `x^3/(3*(1 - x^2)^(3/2)) - x/Sqrt[1 - x^2] + ArcSin[x]`

Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 252

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(2*b*(p + 1))), x] - Simp[c^2*(m - 1)/(2*b*(p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && LtQ[p, -1] && GtQ[m, 1] && !ILtQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]
```

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.86

method	result	size
default	$\frac{x^3}{3(-x^2+1)^{\frac{3}{2}}} + \arcsin(x) - \frac{x}{\sqrt{-x^2+1}}$	30
risch	$-\frac{x(4x^2-3)}{3(x^2-1)\sqrt{-x^2+1}} + \arcsin(x)$	30
meijerg	$-\frac{2i \left(-\frac{i\sqrt{\pi}x(-20x^2+15)}{10(-x^2+1)^{\frac{3}{2}}} + \frac{3i\sqrt{\pi}\arcsin(x)}{2} \right)}{3\sqrt{\pi}}$	39
pseudoelliptic	$\frac{(x^2-1)\sqrt{-x^2+1}\arctan\left(\frac{\sqrt{-x^2+1}}{x}\right) + \frac{4x^3}{3} - x}{(-x^2+1)^{\frac{3}{2}}}$	49
trager	$\frac{(4x^2-3)x\sqrt{-x^2+1}}{3(x^2-1)^2} + \text{RootOf}(_Z^2 + 1) \ln(\text{RootOf}(_Z^2 + 1)\sqrt{-x^2+1} + x)$	54

input `int(x^4/(-x^2+1)^(5/2),x,method=_RETURNVERBOSE)`output `1/3*x^3/(-x^2+1)^(3/2)+arcsin(x)-1/(-x^2+1)^(1/2)*x`**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 63 vs. 2(29) = 58.

Time = 0.08 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.80

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = -\frac{6(x^4 - 2x^2 + 1) \arctan\left(\frac{\sqrt{-x^2+1}-1}{x}\right) - (4x^3 - 3x)\sqrt{-x^2+1}}{3(x^4 - 2x^2 + 1)}$$

input `integrate(x^4/(-x^2+1)^(5/2),x, algorithm="fricas")`

output `-1/3*(6*(x^4 - 2*x^2 + 1)*arctan((sqrt(-x^2 + 1) - 1)/x) - (4*x^3 - 3*x)*sqrt(-x^2 + 1))/(x^4 - 2*x^2 + 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 105 vs. $2(26) = 52$.

Time = 0.33 (sec) , antiderivative size = 105, normalized size of antiderivative = 3.00

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{3x^4 \operatorname{asin}(x)}{3x^4 - 6x^2 + 3} + \frac{4x^3 \sqrt{1-x^2}}{3x^4 - 6x^2 + 3} - \frac{6x^2 \operatorname{asin}(x)}{3x^4 - 6x^2 + 3} - \frac{3x \sqrt{1-x^2}}{3x^4 - 6x^2 + 3} + \frac{3 \operatorname{asin}(x)}{3x^4 - 6x^2 + 3}$$

input `integrate(x**4/(-x**2+1)**(5/2),x)`

output `3*x**4*asin(x)/(3*x**4 - 6*x**2 + 3) + 4*x**3*sqrt(1 - x**2)/(3*x**4 - 6*x**2 + 3) - 6*x**2*asin(x)/(3*x**4 - 6*x**2 + 3) - 3*x*sqrt(1 - x**2)/(3*x**4 - 6*x**2 + 3) + 3*asin(x)/(3*x**4 - 6*x**2 + 3)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.26

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{1}{3} x \left(\frac{3x^2}{(-x^2+1)^{3/2}} - \frac{2}{(-x^2+1)^{3/2}} \right) - \frac{x}{3\sqrt{-x^2+1}} + \operatorname{arcsin}(x)$$

input `integrate(x^4/(-x^2+1)^(5/2),x, algorithm="maxima")`

output `1/3*x*(3*x^2/(-x^2 + 1)^(3/2) - 2/(-x^2 + 1)^(3/2)) - 1/3*x/sqrt(-x^2 + 1) + arcsin(x)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.83

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{(4x^2-3)\sqrt{-x^2+1}x}{3(x^2-1)^2} + \arcsin(x)$$

input `integrate(x^4/(-x^2+1)^(5/2),x, algorithm="giac")`output `1/3*(4*x^2 - 3)*sqrt(-x^2 + 1)*x/(x^2 - 1)^2 + arcsin(x)`**Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 91, normalized size of antiderivative = 2.60

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \arcsin(x) + \frac{3\sqrt{1-x^2}}{4(x-1)} + \frac{3\sqrt{1-x^2}}{4(x+1)} - \sqrt{1-x^2} \left(\frac{1}{12(x-1)} - \frac{1}{12(x-1)^2} \right) - \sqrt{1-x^2} \left(\frac{1}{12(x+1)} + \frac{1}{12(x+1)^2} \right)$$

input `int(x^4/(1 - x^2)^(5/2),x)`output `asin(x) + (3*(1 - x^2)^(1/2))/(4*(x - 1)) + (3*(1 - x^2)^(1/2))/(4*(x + 1)) - (1 - x^2)^(1/2)*(1/(12*(x - 1)) - 1/(12*(x - 1)^2)) - (1 - x^2)^(1/2)*(1/(12*(x + 1)) + 1/(12*(x + 1)^2))`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.57

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{3\sqrt{-x^2+1} \arcsin(x) x^2 - 3\sqrt{-x^2+1} \arcsin(x) - 4x^3 + 3x}{3\sqrt{-x^2+1} (x^2 - 1)}$$

input `int(x^4/(-x^2+1)^(5/2),x)`

output
$$\frac{(3\sqrt{-x^2 + 1})\operatorname{asin}(x)x^{**2} - 3\sqrt{-x^2 + 1}\operatorname{asin}(x) - 4x^{**3} + 3x}{(3\sqrt{-x^2 + 1})(x^{**2} - 1)}$$

3.38 $\int \sqrt{x}(1+x)^{5/2} dx$

Optimal result	261
Mathematica [A] (verified)	261
Rubi [A] (verified)	262
Maple [A] (verified)	263
Fricas [A] (verification not implemented)	264
Sympy [C] (verification not implemented)	264
Maxima [B] (verification not implemented)	265
Giac [A] (verification not implemented)	265
Mupad [F(-1)]	266
Reduce [B] (verification not implemented)	266

Optimal result

Integrand size = 13, antiderivative size = 75

$$\int \sqrt{x}(1+x)^{5/2} dx = \frac{5}{64}\sqrt{x}\sqrt{1+x} + \frac{5}{32}x^{3/2}\sqrt{1+x} + \frac{5}{24}x^{3/2}(1+x)^{3/2} + \frac{1}{4}x^{3/2}(1+x)^{5/2} - \frac{5\operatorname{arcsinh}(\sqrt{x})}{64}$$

output $5/24*x^{(3/2)}*(1+x)^{(3/2)}+1/4*x^{(3/2)}*(1+x)^{(5/2)}-5/64*\operatorname{arcsinh}(x^{(1/2)})+5/32*x^{(3/2)}*(1+x)^{(1/2)}+5/64*x^{(1/2)}*(1+x)^{(1/2)}$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.69

$$\int \sqrt{x}(1+x)^{5/2} dx = \frac{1}{192}\sqrt{x}\sqrt{1+x}(15 + 118x + 136x^2 + 48x^3) + \frac{5}{64}\log\left(-\sqrt{x} + \sqrt{1+x}\right)$$

input $\operatorname{Integrate}[\operatorname{Sqrt}[x]*(1+x)^{(5/2)},x]$

output

```
(Sqrt[x]*Sqrt[1 + x]*(15 + 118*x + 136*x^2 + 48*x^3))/192 + (5*Log[-Sqrt[x]
] + Sqrt[1 + x])/64
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.13, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$, Rules used = {60, 60, 60, 60, 63, 222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{x}(x+1)^{5/2} dx$$

$$\downarrow 60$$

$$\frac{5}{8} \int \sqrt{x}(x+1)^{3/2} dx + \frac{1}{4} x^{3/2}(x+1)^{5/2}$$

$$\downarrow 60$$

$$\frac{5}{8} \left(\frac{1}{2} \int \sqrt{x}\sqrt{x+1} dx + \frac{1}{3} x^{3/2}(x+1)^{3/2} \right) + \frac{1}{4} x^{3/2}(x+1)^{5/2}$$

$$\downarrow 60$$

$$\frac{5}{8} \left(\frac{1}{2} \left(\frac{1}{4} \int \frac{\sqrt{x}}{\sqrt{x+1}} dx + \frac{1}{2} \sqrt{x+1} x^{3/2} \right) + \frac{1}{3} x^{3/2}(x+1)^{3/2} \right) + \frac{1}{4} x^{3/2}(x+1)^{5/2}$$

$$\downarrow 60$$

$$\frac{5}{8} \left(\frac{1}{2} \left(\frac{1}{4} \left(\sqrt{x}\sqrt{x+1} - \frac{1}{2} \int \frac{1}{\sqrt{x}\sqrt{x+1}} dx \right) + \frac{1}{2} \sqrt{x+1} x^{3/2} \right) + \frac{1}{3} x^{3/2}(x+1)^{3/2} \right) + \frac{1}{4} x^{3/2}(x+1)^{5/2}$$

$$\downarrow 63$$

$$\frac{5}{8} \left(\frac{1}{2} \left(\frac{1}{4} \left(\sqrt{x}\sqrt{x+1} - \int \frac{1}{\sqrt{x+1}} d\sqrt{x} \right) + \frac{1}{2} \sqrt{x+1} x^{3/2} \right) + \frac{1}{3} x^{3/2}(x+1)^{3/2} \right) + \frac{1}{4} x^{3/2}(x+1)^{5/2}$$

$$\downarrow 222$$

$$\frac{5}{8} \left(\frac{1}{2} \left(\frac{1}{4} \left(\sqrt{x}\sqrt{x+1} - \operatorname{arcsinh}(\sqrt{x}) \right) + \frac{1}{2} \sqrt{x+1} x^{3/2} \right) + \frac{1}{3} x^{3/2}(x+1)^{3/2} \right) + \frac{1}{4} x^{3/2}(x+1)^{5/2}$$

input `Int[Sqrt[x]*(1 + x)^(5/2),x]`

output `(x^(3/2)*(1 + x)^(5/2))/4 + (5*((x^(3/2)*(1 + x)^(3/2))/3 + ((x^(3/2)*Sqrt[1 + x])/2 + (Sqrt[x]*Sqrt[1 + x] - ArcSinh[Sqrt[x]])/4)/2))/8`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 63 `Int[1/(Sqrt[(b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2/b Subst[Int[1/Sqrt[c + d*(x^2/b)], x], x, Sqrt[b*x]], x] /; FreeQ[{b, c, d}, x] && GtQ[c, 0]`

rule 222 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.59

method	result	size
meijerg	$-\frac{15 \left(-\frac{\sqrt{\pi} \sqrt{x} (48x^3 + 136x^2 + 118x + 15) \sqrt{1+x}}{360} + \frac{\sqrt{\pi} \operatorname{arcsinh}(\sqrt{x})}{24} \right)}{8\sqrt{\pi}}$	44
risch	$\frac{(48x^3 + 136x^2 + 118x + 15)\sqrt{x}\sqrt{1+x}}{192} - \frac{5\sqrt{x(1+x)} \ln\left(x + \frac{1}{2} + \sqrt{x^2+x}\right)}{128\sqrt{1+x}\sqrt{x}}$	55
default	$\frac{\sqrt{x}(1+x)^{\frac{7}{2}}}{4} - \frac{\sqrt{x}(1+x)^{\frac{5}{2}}}{24} - \frac{5\sqrt{x}(1+x)^{\frac{3}{2}}}{96} - \frac{5\sqrt{x}\sqrt{1+x}}{64} - \frac{5\sqrt{x(1+x)} \ln\left(x + \frac{1}{2} + \sqrt{x^2+x}\right)}{128\sqrt{1+x}\sqrt{x}}$	70

input `int(x^(1/2)*(1+x)^(5/2),x,method=_RETURNVERBOSE)`

output `-15/8/Pi^(1/2)*(-1/360*Pi^(1/2)*x^(1/2)*(48*x^3+136*x^2+118*x+15)*(1+x)^(1/2)+1/24*Pi^(1/2)*arcsinh(x^(1/2)))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.59

$$\int \sqrt{x}(1+x)^{5/2} dx = \frac{1}{192} (48x^3 + 136x^2 + 118x + 15)\sqrt{x+1}\sqrt{x} + \frac{5}{128} \log(2\sqrt{x+1}\sqrt{x} - 2x - 1)$$

input `integrate(x^(1/2)*(1+x)^(5/2),x, algorithm="fricas")`

output `1/192*(48*x^3 + 136*x^2 + 118*x + 15)*sqrt(x + 1)*sqrt(x) + 5/128*log(2*sqrt(x + 1)*sqrt(x) - 2*x - 1)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 18.27 (sec) , antiderivative size = 190, normalized size of antiderivative = 2.53

$$\int \sqrt{x}(1+x)^{5/2} dx = \begin{cases} -\frac{5 \operatorname{acosh}(\sqrt{x+1})}{64} + \frac{(x+1)^{\frac{9}{2}}}{4\sqrt{x}} - \frac{7(x+1)^{\frac{7}{2}}}{24\sqrt{x}} - \frac{(x+1)^{\frac{5}{2}}}{96\sqrt{x}} - \frac{5(x+1)^{\frac{3}{2}}}{192\sqrt{x}} + \frac{5\sqrt{x+1}}{64\sqrt{x}} & \text{for } |x+1| > 1 \\ \frac{5i \operatorname{asin}(\sqrt{x+1})}{64} - \frac{i(x+1)^{\frac{9}{2}}}{4\sqrt{-x}} + \frac{7i(x+1)^{\frac{7}{2}}}{24\sqrt{-x}} + \frac{i(x+1)^{\frac{5}{2}}}{96\sqrt{-x}} + \frac{5i(x+1)^{\frac{3}{2}}}{192\sqrt{-x}} - \frac{5i\sqrt{x+1}}{64\sqrt{-x}} & \text{otherwise} \end{cases}$$

input `integrate(x**(1/2)*(1+x)**(5/2),x)`

output

```
Piecewise((-5*acosh(sqrt(x + 1))/64 + (x + 1)**(9/2)/(4*sqrt(x)) - 7*(x + 1)**(7/2)/(24*sqrt(x)) - (x + 1)**(5/2)/(96*sqrt(x)) - 5*(x + 1)**(3/2)/(192*sqrt(x)) + 5*sqrt(x + 1)/(64*sqrt(x)), Abs(x + 1) > 1), (5*I*asin(sqrt(x + 1))/64 - I*(x + 1)**(9/2)/(4*sqrt(-x)) + 7*I*(x + 1)**(7/2)/(24*sqrt(-x)) + I*(x + 1)**(5/2)/(96*sqrt(-x)) + 5*I*(x + 1)**(3/2)/(192*sqrt(-x)) - 5*I*sqrt(x + 1)/(64*sqrt(-x)), True))
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 113 vs. $2(47) = 94$.

Time = 0.02 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.51

$$\int \sqrt{x}(1+x)^{5/2} dx = \frac{\frac{15(x+1)^{7/2}}{x^{7/2}} + \frac{73(x+1)^{5/2}}{x^{5/2}} - \frac{55(x+1)^{3/2}}{x^{3/2}} + \frac{15\sqrt{x+1}}{\sqrt{x}}}{192 \left(\frac{(x+1)^4}{x^4} - \frac{4(x+1)^3}{x^3} + \frac{6(x+1)^2}{x^2} - \frac{4(x+1)}{x} + 1 \right)} - \frac{5}{128} \log \left(\frac{\sqrt{x+1}}{\sqrt{x}} + 1 \right) + \frac{5}{128} \log \left(\frac{\sqrt{x+1}}{\sqrt{x}} - 1 \right)$$

input

```
integrate(x^(1/2)*(1+x)^(5/2),x, algorithm="maxima")
```

output

```
1/192*(15*(x + 1)^(7/2)/x^(7/2) + 73*(x + 1)^(5/2)/x^(5/2) - 55*(x + 1)^(3/2)/x^(3/2) + 15*sqrt(x + 1)/sqrt(x))/(x + 1)^4/x^4 - 4*(x + 1)^3/x^3 + 6*(x + 1)^2/x^2 - 4*(x + 1)/x + 1) - 5/128*log(sqrt(x + 1)/sqrt(x) + 1) + 5/128*log(sqrt(x + 1)/sqrt(x) - 1)
```

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.20

$$\int \sqrt{x}(1+x)^{5/2} dx = \frac{1}{192} (2(4(6x-19)(x+1)+163)(x+1)-279)\sqrt{x+1}\sqrt{x} + \frac{1}{8} (2(4x-9)(x+1)+33)\sqrt{x+1}\sqrt{x} + \frac{3}{4} (2x-3)\sqrt{x+1}\sqrt{x} + \sqrt{x+1}\sqrt{x} + \frac{5}{64} \log(\sqrt{x+1}-\sqrt{x})$$

input `integrate(x^(1/2)*(1+x)^(5/2),x, algorithm="giac")`

output `1/192*(2*(4*(6*x - 19)*(x + 1) + 163)*(x + 1) - 279)*sqrt(x + 1)*sqrt(x) + 1/8*(2*(4*x - 9)*(x + 1) + 33)*sqrt(x + 1)*sqrt(x) + 3/4*(2*x - 3)*sqrt(x + 1)*sqrt(x) + sqrt(x + 1)*sqrt(x) + 5/64*log(sqrt(x + 1) - sqrt(x))`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{x}(1+x)^{5/2} dx = \int \sqrt{x}(x+1)^{5/2} dx$$

input `int(x^(1/2)*(x + 1)^(5/2),x)`

output `int(x^(1/2)*(x + 1)^(5/2), x)`

Reduce [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.67

$$\int \sqrt{x}(1+x)^{5/2} dx = \frac{\sqrt{x}\sqrt{x+1}x^3}{4} + \frac{17\sqrt{x}\sqrt{x+1}x^2}{24} + \frac{59\sqrt{x}\sqrt{x+1}x}{96} + \frac{5\sqrt{x}\sqrt{x+1}}{64} - \frac{5\log(\sqrt{x+1} + \sqrt{x})}{64}$$

input `int(x^(1/2)*(1+x)^(5/2),x)`

output `(48*sqrt(x)*sqrt(x + 1)*x**3 + 136*sqrt(x)*sqrt(x + 1)*x**2 + 118*sqrt(x)*sqrt(x + 1)*x + 15*sqrt(x)*sqrt(x + 1) - 15*log(sqrt(x + 1) + sqrt(x)))/192`

$$3.39 \quad \int \frac{x^4}{(1-x^2)^{5/2}} dx$$

Optimal result	267
Mathematica [A] (verified)	267
Rubi [A] (verified)	268
Maple [A] (verified)	269
Fricas [B] (verification not implemented)	269
Sympy [B] (verification not implemented)	270
Maxima [A] (verification not implemented)	270
Giac [A] (verification not implemented)	271
Mupad [B] (verification not implemented)	271
Reduce [B] (verification not implemented)	271

Optimal result

Integrand size = 15, antiderivative size = 35

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{x^3}{3(1-x^2)^{3/2}} - \frac{x}{\sqrt{1-x^2}} + \arcsin(x)$$

output `1/3*x^3/(-x^2+1)^(3/2)+arcsin(x)-x/(-x^2+1)^(1/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.26

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{x(-3+4x^2)}{3(1-x^2)^{3/2}} + 2 \arctan\left(\frac{x}{-1+\sqrt{1-x^2}}\right)$$

input `Integrate[x^4/(1-x^2)^(5/2),x]`

output `(x*(-3+4*x^2))/(3*(1-x^2)^(3/2))+2*ArcTan[x/(-1+Sqrt[1-x^2])]`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {252, 252, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx$$

$$\downarrow \text{252}$$

$$\frac{x^3}{3(1-x^2)^{3/2}} - \int \frac{x^2}{(1-x^2)^{3/2}} dx$$

$$\downarrow \text{252}$$

$$\int \frac{1}{\sqrt{1-x^2}} dx - \frac{x}{\sqrt{1-x^2}} + \frac{x^3}{3(1-x^2)^{3/2}}$$

$$\downarrow \text{223}$$

$$\arcsin(x) - \frac{x}{\sqrt{1-x^2}} + \frac{x^3}{3(1-x^2)^{3/2}}$$

input `Int[x^4/(1 - x^2)^(5/2),x]`

output `x^3/(3*(1 - x^2)^(3/2)) - x/Sqrt[1 - x^2] + ArcSin[x]`

Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 252

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(2*b*(p + 1))), x] - Simp[c^2*(m - 1)/(2*b*(p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && LtQ[p, -1] && GtQ[m, 1] && !ILtQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]
```

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.86

method	result	size
default	$\frac{x^3}{3(-x^2+1)^{\frac{3}{2}}} + \arcsin(x) - \frac{x}{\sqrt{-x^2+1}}$	30
risch	$-\frac{x(4x^2-3)}{3(x^2-1)\sqrt{-x^2+1}} + \arcsin(x)$	30
meijerg	$-\frac{2i \left(-\frac{i\sqrt{\pi}x(-20x^2+15)}{10(-x^2+1)^{\frac{3}{2}}} + \frac{3i\sqrt{\pi}\arcsin(x)}{2} \right)}{3\sqrt{\pi}}$	39
pseudoelliptic	$\frac{(x^2-1)\sqrt{-x^2+1}\arctan\left(\frac{\sqrt{-x^2+1}}{x}\right) + \frac{4x^3}{3} - x}{(-x^2+1)^{\frac{3}{2}}}$	49
trager	$\frac{(4x^2-3)x\sqrt{-x^2+1}}{3(x^2-1)^2} + \text{RootOf}(_Z^2 + 1) \ln(\text{RootOf}(_Z^2 + 1)\sqrt{-x^2+1} + x)$	54

input

```
int(x^4/(-x^2+1)^(5/2),x,method=_RETURNVERBOSE)
```

output

```
1/3*x^3/(-x^2+1)^(3/2)+arcsin(x)-1/(-x^2+1)^(1/2)*x
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 63 vs. 2(29) = 58.

Time = 0.08 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.80

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = -\frac{6(x^4 - 2x^2 + 1) \arctan\left(\frac{\sqrt{-x^2+1}-1}{x}\right) - (4x^3 - 3x)\sqrt{-x^2+1}}{3(x^4 - 2x^2 + 1)}$$

input `integrate(x^4/(-x^2+1)^(5/2),x, algorithm="fricas")`

output `-1/3*(6*(x^4 - 2*x^2 + 1)*arctan((sqrt(-x^2 + 1) - 1)/x) - (4*x^3 - 3*x)*sqrt(-x^2 + 1))/(x^4 - 2*x^2 + 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 105 vs. $2(26) = 52$.

Time = 0.41 (sec) , antiderivative size = 105, normalized size of antiderivative = 3.00

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{3x^4 \operatorname{asin}(x)}{3x^4 - 6x^2 + 3} + \frac{4x^3 \sqrt{1-x^2}}{3x^4 - 6x^2 + 3} - \frac{6x^2 \operatorname{asin}(x)}{3x^4 - 6x^2 + 3} - \frac{3x \sqrt{1-x^2}}{3x^4 - 6x^2 + 3} + \frac{3 \operatorname{asin}(x)}{3x^4 - 6x^2 + 3}$$

input `integrate(x**4/(-x**2+1)**(5/2),x)`

output `3*x**4*asin(x)/(3*x**4 - 6*x**2 + 3) + 4*x**3*sqrt(1 - x**2)/(3*x**4 - 6*x**2 + 3) - 6*x**2*asin(x)/(3*x**4 - 6*x**2 + 3) - 3*x*sqrt(1 - x**2)/(3*x**4 - 6*x**2 + 3) + 3*asin(x)/(3*x**4 - 6*x**2 + 3)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.26

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{1}{3} x \left(\frac{3x^2}{(-x^2+1)^{3/2}} - \frac{2}{(-x^2+1)^{3/2}} \right) - \frac{x}{3\sqrt{-x^2+1}} + \operatorname{arcsin}(x)$$

input `integrate(x^4/(-x^2+1)^(5/2),x, algorithm="maxima")`

output `1/3*x*(3*x^2/(-x^2 + 1)^(3/2) - 2/(-x^2 + 1)^(3/2)) - 1/3*x/sqrt(-x^2 + 1) + arcsin(x)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.83

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{(4x^2-3)\sqrt{-x^2+1}x}{3(x^2-1)^2} + \arcsin(x)$$

input `integrate(x^4/(-x^2+1)^(5/2),x, algorithm="giac")`output `1/3*(4*x^2 - 3)*sqrt(-x^2 + 1)*x/(x^2 - 1)^2 + arcsin(x)`**Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 91, normalized size of antiderivative = 2.60

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \arcsin(x) + \frac{3\sqrt{1-x^2}}{4(x-1)} + \frac{3\sqrt{1-x^2}}{4(x+1)} - \sqrt{1-x^2} \left(\frac{1}{12(x-1)} - \frac{1}{12(x-1)^2} \right) - \sqrt{1-x^2} \left(\frac{1}{12(x+1)} + \frac{1}{12(x+1)^2} \right)$$

input `int(x^4/(1 - x^2)^(5/2),x)`output `asin(x) + (3*(1 - x^2)^(1/2))/(4*(x - 1)) + (3*(1 - x^2)^(1/2))/(4*(x + 1)) - (1 - x^2)^(1/2)*(1/(12*(x - 1)) - 1/(12*(x - 1)^2)) - (1 - x^2)^(1/2)*(1/(12*(x + 1)) + 1/(12*(x + 1)^2))`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.57

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{3\sqrt{-x^2+1} \arcsin(x) x^2 - 3\sqrt{-x^2+1} \arcsin(x) - 4x^3 + 3x}{3\sqrt{-x^2+1} (x^2 - 1)}$$

input `int(x^4/(-x^2+1)^(5/2),x)`

output
$$\frac{(3\sqrt{-x^2 + 1})\operatorname{asin}(x)x^{**2} - 3\sqrt{-x^2 + 1}\operatorname{asin}(x) - 4x^{**3} + 3x}{(3\sqrt{-x^2 + 1})(x^{**2} - 1)}$$

3.40 $\int \frac{\sqrt{A^2+B^2-B^2y^2}}{1-y^2} dy$

Optimal result	273
Mathematica [A] (verified)	273
Rubi [A] (verified)	274
Maple [A] (verified)	276
Fricas [B] (verification not implemented)	276
Sympy [F]	277
Maxima [B] (verification not implemented)	277
Giac [B] (verification not implemented)	278
Mupad [F(-1)]	278
Reduce [B] (verification not implemented)	279

Optimal result

Integrand size = 29, antiderivative size = 51

$$\int \frac{\sqrt{A^2 + B^2 - B^2y^2}}{1 - y^2} dy = B \arctan\left(\frac{By}{\sqrt{A^2 + B^2 - B^2y^2}}\right) + A \operatorname{arctanh}\left(\frac{Ay}{\sqrt{A^2 + B^2 - B^2y^2}}\right)$$

output `B*arctan(B*y/(-B^2*y^2+A^2+B^2)^(1/2))+A*arctanh(A*y/(-B^2*y^2+A^2+B^2)^(1/2))`

Mathematica [A] (verified)

Time = 0.71 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.53

$$\int \frac{\sqrt{A^2 + B^2 - B^2y^2}}{1 - y^2} dy = -2B \arctan\left(\frac{-\frac{\sqrt{A^2+B^2}}{B} + \frac{\sqrt{A^2+B^2-B^2y^2}}{B}}{y}\right) + A \operatorname{arctanh}\left(\frac{\sqrt{A^2 + B^2 - B^2y^2}}{Ay}\right)$$

input `Integrate[Sqrt[A^2 + B^2 - B^2*y^2]/(1 - y^2),y]`

output

```
-2*B*ArcTan[(-(Sqrt[A^2 + B^2]/B) + Sqrt[A^2 + B^2 - B^2*y^2]/B)/y] + A*ArcTanh[Sqrt[A^2 + B^2 - B^2*y^2]/(A*y)]
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.172$, Rules used = {301, 224, 216, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{A^2 - B^2 y^2 + B^2}}{1 - y^2} dy$$

↓ 301

$$A^2 \int \frac{1}{(1 - y^2) \sqrt{A^2 + B^2 - B^2 y^2}} dy + B^2 \int \frac{1}{\sqrt{A^2 + B^2 - B^2 y^2}} dy$$

↓ 224

$$A^2 \int \frac{1}{(1 - y^2) \sqrt{A^2 + B^2 - B^2 y^2}} dy + B^2 \int \frac{1}{\frac{B^2 y^2}{A^2 + B^2 - B^2 y^2} + 1} d \frac{y}{\sqrt{A^2 + B^2 - B^2 y^2}}$$

↓ 216

$$A^2 \int \frac{1}{(1 - y^2) \sqrt{A^2 + B^2 - B^2 y^2}} dy + B \arctan \left(\frac{By}{\sqrt{A^2 - B^2 y^2 + B^2}} \right)$$

↓ 291

$$A^2 \int \frac{1}{1 - \frac{A^2 y^2}{A^2 + B^2 - B^2 y^2}} d \frac{y}{\sqrt{A^2 + B^2 - B^2 y^2}} + B \arctan \left(\frac{By}{\sqrt{A^2 - B^2 y^2 + B^2}} \right)$$

↓ 219

$$B \arctan \left(\frac{By}{\sqrt{A^2 - B^2 y^2 + B^2}} \right) + A \operatorname{arctanh} \left(\frac{Ay}{\sqrt{A^2 - B^2 y^2 + B^2}} \right)$$

input

```
Int[Sqrt[A^2 + B^2 - B^2*y^2]/(1 - y^2), y]
```

output $B \cdot \text{ArcTan}[(B \cdot y) / \sqrt{A^2 + B^2 - B^2 \cdot y^2}] + A \cdot \text{ArcTanh}[(A \cdot y) / \sqrt{A^2 + B^2 - B^2 \cdot y^2}]$

Defintions of rubi rules used

rule 216 $\text{Int}[(a_) + (b_.) \cdot (x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1 / (\text{Rt}[a, 2] \cdot \text{Rt}[b, 2])) \cdot \text{ArcTan}[\text{Rt}[b, 2] \cdot (x / \text{Rt}[a, 2])], x] /;$ FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

rule 219 $\text{Int}[(a_) + (b_.) \cdot (x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1 / (\text{Rt}[a, 2] \cdot \text{Rt}[-b, 2])) \cdot \text{ArcTanh}[\text{Rt}[-b, 2] \cdot (x / \text{Rt}[a, 2])], x] /;$ FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

rule 224 $\text{Int}[1 / \sqrt{(a_) + (b_.) \cdot (x_)^2}, x_Symbol] \rightarrow \text{Subst}[\text{Int}[1 / (1 - b \cdot x^2), x], x, x / \sqrt{a + b \cdot x^2}] /;$ FreeQ[{a, b}, x] && !GtQ[a, 0]

rule 291 $\text{Int}[1 / (\sqrt{(a_) + (b_.) \cdot (x_)^2} \cdot ((c_) + (d_.) \cdot (x_)^2)), x_Symbol] \rightarrow \text{Subst}[\text{Int}[1 / (c - (b \cdot c - a \cdot d) \cdot x^2), x], x, x / \sqrt{a + b \cdot x^2}] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b \cdot c - a \cdot d, 0]

rule 301 $\text{Int}[(a_) + (b_.) \cdot (x_)^2)^{p_.)} / ((c_) + (d_.) \cdot (x_)^2), x_Symbol] \rightarrow \text{Simp}[b / d \cdot \text{Int}[(a + b \cdot x^2)^{p-1}, x], x] - \text{Simp}[(b \cdot c - a \cdot d) / d \cdot \text{Int}[(a + b \cdot x^2)^{p-1} / (c + d \cdot x^2), x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b \cdot c - a \cdot d, 0] && GtQ[p, 0] && (EqQ[p, 1/2] || EqQ[Denominator[p], 4] || (EqQ[p, 2/3] && EqQ[b \cdot c + 3 \cdot a \cdot d, 0]))

Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.76

method	result
pseudoelliptic	$-\frac{A \ln\left(\frac{Ay - \sqrt{-B^2y^2 + A^2 + B^2}}{y}\right)}{2} + \frac{A \ln\left(\frac{Ay + \sqrt{-B^2y^2 + A^2 + B^2}}{y}\right)}{2} - B \arctan\left(\frac{\sqrt{-B^2y^2 + A^2 + B^2}}{By}\right)$
default	$\frac{\sqrt{-B^2(1+y)^2 + 2B^2(1+y) + A^2}}{2} + \frac{B^2 \arctan\left(\frac{\sqrt{B^2}y}{\sqrt{-B^2(1+y)^2 + 2B^2(1+y) + A^2}}\right)}{2\sqrt{B^2}} - \frac{A^2 \ln\left(\frac{2A^2 + 2B^2(1+y) + 2\sqrt{A^2}\sqrt{-B^2(1+y)^2 + 2B^2(1+y) + A^2}}{1+y}\right)}{2\sqrt{A^2}}$

input `int((-B^2*y^2+A^2+B^2)^(1/2)/(-y^2+1),y,method=_RETURNVERBOSE)`

output `-1/2*A*ln((A*y-(-B^2*y^2+A^2+B^2)^(1/2))/y)+1/2*A*ln((A*y+(-B^2*y^2+A^2+B^2)^(1/2))/y)-B*arctan(1/B/y*(-B^2*y^2+A^2+B^2)^(1/2))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 145 vs. 2(47) = 94.

Time = 0.09 (sec) , antiderivative size = 145, normalized size of antiderivative = 2.84

$$\begin{aligned} & \int \frac{\sqrt{A^2 + B^2 - B^2y^2}}{1 - y^2} dy \\ &= -B \arctan\left(\frac{\sqrt{-B^2y^2 + A^2 + B^2}By}{B^2y^2 - A^2 - B^2}\right) \\ & \quad + \frac{1}{4} A \log\left(-\frac{(A^2 - B^2)y^2 + 2\sqrt{-B^2y^2 + A^2 + B^2}Ay + A^2 + B^2}{y^2}\right) \\ & \quad - \frac{1}{4} A \log\left(-\frac{(A^2 - B^2)y^2 - 2\sqrt{-B^2y^2 + A^2 + B^2}Ay + A^2 + B^2}{y^2}\right) \end{aligned}$$

input `integrate((-B^2*y^2+A^2+B^2)^(1/2)/(-y^2+1),y, algorithm="fricas")`

output

```
-B*arctan(sqrt(-B^2*y^2 + A^2 + B^2)*B*y/(B^2*y^2 - A^2 - B^2)) + 1/4*A*log(-((A^2 - B^2)*y^2 + 2*sqrt(-B^2*y^2 + A^2 + B^2)*A*y + A^2 + B^2)/y^2) - 1/4*A*log(-((A^2 - B^2)*y^2 - 2*sqrt(-B^2*y^2 + A^2 + B^2)*A*y + A^2 + B^2)/y^2)
```

Sympy [F]

$$\int \frac{\sqrt{A^2 + B^2 - B^2 y^2}}{1 - y^2} dy = - \int \frac{\sqrt{A^2 - B^2 y^2 + B^2}}{y^2 - 1} dy$$

input

```
integrate((-B**2*y**2+A**2+B**2)**(1/2)/(-y**2+1),y)
```

output

```
-Integral(sqrt(A**2 - B**2*y**2 + B**2)/(y**2 - 1), y)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 114 vs. 2(47) = 94.

Time = 0.11 (sec) , antiderivative size = 114, normalized size of antiderivative = 2.24

$$\int \frac{\sqrt{A^2 + B^2 - B^2 y^2}}{1 - y^2} dy = B \arcsin \left(\frac{B^2 y}{\sqrt{A^2 B^2 + B^4}} \right) - \frac{1}{2} A \log \left(B^2 + \frac{A^2}{y+1} + \frac{\sqrt{-B^2 y^2 + A^2 + B^2 A}}{y+1} \right) + \frac{1}{2} A \log \left(-B^2 + \frac{2 A^2}{|2y-2|} + \frac{2 \sqrt{-B^2 y^2 + A^2 + B^2 A}}{|2y-2|} \right)$$

input

```
integrate((-B^2*y^2+A^2+B^2)^(1/2)/(-y^2+1),y, algorithm="maxima")
```

output

```
B*arcsin(B^2*y/sqrt(A^2*B^2 + B^4)) - 1/2*A*log(B^2 + A^2/(y + 1) + sqrt(-B^2*y^2 + A^2 + B^2)*A/(y + 1)) + 1/2*A*log(-B^2 + 2*A^2/abs(2*y - 2) + 2*sqrt(-B^2*y^2 + A^2 + B^2)*A/abs(2*y - 2))
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 295 vs. $2(47) = 94$.

Time = 0.19 (sec) , antiderivative size = 295, normalized size of antiderivative = 5.78

$$\int \frac{\sqrt{A^2 + B^2 - B^2 y^2}}{1 - y^2} dy$$

$$= - \frac{\left(\pi \operatorname{sgn}(y) - 2 \arctan \left(- \frac{B^2 y \left(\frac{(\sqrt{A^2 + B^2} B + \sqrt{-B^2 y^2 + A^2 + B^2} |B|)^2}{B^4 y^2} - 1 \right)}{2 (\sqrt{A^2 + B^2} B + \sqrt{-B^2 y^2 + A^2 + B^2} |B|)} \right) \right) B^2}{2 |B|}$$

$$+ \frac{AB \log \left(\left| - \left(\frac{B^2 y}{\sqrt{A^2 + B^2} B + \sqrt{-B^2 y^2 + A^2 + B^2} |B|} - \frac{\sqrt{A^2 + B^2} B + \sqrt{-B^2 y^2 + A^2 + B^2} |B|}{B^2 y} \right) B + 2A \right| \right)}{2 |B|}$$

$$- \frac{AB \log \left(\left| - \left(\frac{B^2 y}{\sqrt{A^2 + B^2} B + \sqrt{-B^2 y^2 + A^2 + B^2} |B|} - \frac{\sqrt{A^2 + B^2} B + \sqrt{-B^2 y^2 + A^2 + B^2} |B|}{B^2 y} \right) B - 2A \right| \right)}{2 |B|}$$

input `integrate((-B^2*y^2+A^2+B^2)^(1/2)/(-y^2+1),y, algorithm="giac")`

output `-1/2*(pi*sgn(y) - 2*arctan(-1/2*B^2*y*((sqrt(A^2 + B^2)*B + sqrt(-B^2*y^2 + A^2 + B^2)*abs(B))^2/(B^4*y^2) - 1)/(sqrt(A^2 + B^2)*B + sqrt(-B^2*y^2 + A^2 + B^2)*abs(B)))*B^2/abs(B) + 1/2*A*B*log(abs(-(B^2*y/(sqrt(A^2 + B^2)*B + sqrt(-B^2*y^2 + A^2 + B^2)*abs(B)))/(B^2*y))*B + 2*A))/abs(B) - 1/2*A*B*log(abs(-(B^2*y/(sqrt(A^2 + B^2)*B + sqrt(-B^2*y^2 + A^2 + B^2)*abs(B)) - (sqrt(A^2 + B^2)*B + sqrt(-B^2*y^2 + A^2 + B^2)*abs(B))/(B^2*y))*B - 2*A))/abs(B)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{A^2 + B^2 - B^2 y^2}}{1 - y^2} dy$$

$$= \begin{cases} - \int \frac{\sqrt{-B^2 y^2}}{y^2 - 1} dy & \text{if } A^2 + B^2 = 0 \\ - \ln(2y \sqrt{-B^2} + 2 \sqrt{A^2 - B^2 y^2 + B^2}) \sqrt{-B^2} - \operatorname{atan} \left(\frac{y \sqrt{A^2}}{\sqrt{A^2 - B^2 y^2 + B^2}} \right) \sqrt{A^2} \operatorname{li} & \text{if } A^2 + B^2 \neq 0 \end{cases}$$

input `int(-(A^2 + B^2 - B^2*y^2)^(1/2)/(y^2 - 1),y)`

output `piecewise(A^2 + B^2 == 0, -int((-B^2*y^2)^(1/2)/(y^2 - 1), y), A^2 + B^2 ~
= 0, - atan((y*(A^2)^(1/2)*1i)/(A^2 + B^2 - B^2*y^2)^(1/2))*(A^2)^(1/2)*1i
- log(2*y*(-B^2)^(1/2) + 2*(A^2 + B^2 - B^2*y^2)^(1/2))*(-B^2)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 126, normalized size of antiderivative = 2.47

$$\int \frac{\sqrt{A^2 + B^2 - B^2 y^2}}{1 - y^2} dy = \operatorname{asin}\left(\frac{by}{\sqrt{a^2 + b^2}}\right) b - \operatorname{atan}\left(\frac{\tan\left(\frac{\operatorname{asin}\left(\frac{by}{\sqrt{a^2 + b^2}}\right)}{2}\right) bi}{\sqrt{a^2 + b^2} - a}\right) ai$$

$$+ \frac{\log\left(-\sqrt{a^2 + b^2} + \tan\left(\frac{\operatorname{asin}\left(\frac{by}{\sqrt{a^2 + b^2}}\right)}{2}\right) b - a\right) a}{2}$$

$$- \frac{\log\left(\sqrt{a^2 + b^2} + \tan\left(\frac{\operatorname{asin}\left(\frac{by}{\sqrt{a^2 + b^2}}\right)}{2}\right) b + a\right) a}{2}$$

input `int((-B^2*y^2+A^2+B^2)^(1/2)/(-y^2+1),y)`

output `(2*asin((b*y)/sqrt(a**2 + b**2))*b - 2*atan((tan(asin((b*y)/sqrt(a**2 + b*
*2))/2)*b*i)/(sqrt(a**2 + b**2) - a))*a*i + log(- sqrt(a**2 + b**2) + tan
(asin((b*y)/sqrt(a**2 + b**2))/2)*b - a)*a - log(sqrt(a**2 + b**2) + tan(a
sin((b*y)/sqrt(a**2 + b**2))/2)*b + a)*a)/2`

3.41 $\int \sin^2(x) dx$

Optimal result	280
Mathematica [A] (verified)	280
Rubi [A] (verified)	281
Maple [A] (verified)	282
Fricas [A] (verification not implemented)	282
Sympy [A] (verification not implemented)	283
Maxima [A] (verification not implemented)	283
Giac [A] (verification not implemented)	283
Mupad [B] (verification not implemented)	284
Reduce [B] (verification not implemented)	284

Optimal result

Integrand size = 4, antiderivative size = 14

$$\int \sin^2(x) dx = \frac{x}{2} - \frac{1}{2} \cos(x) \sin(x)$$

output `1/2*x-1/2*cos(x)*sin(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \sin^2(x) dx = \frac{x}{2} - \frac{1}{4} \sin(2x)$$

input `Integrate[Sin[x]^2,x]`

output `x/2 - Sin[2*x]/4`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sin^2(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin(x)^2 dx \\ & \quad \downarrow \text{3115} \\ & \frac{\int 1 dx}{2} - \frac{1}{2} \sin(x) \cos(x) \\ & \quad \downarrow \text{24} \\ & \frac{x}{2} - \frac{1}{2} \sin(x) \cos(x) \end{aligned}$$

input `Int[Sin[x]^2,x]`

output `x/2 - (Cos[x]*Sin[x])/2`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.79

method	result	size
default	$\frac{x}{2} - \frac{\cos(x)\sin(x)}{2}$	11
risch	$\frac{x}{2} - \frac{\sin(2x)}{4}$	11
parallelrisch	$\frac{x}{2} - \frac{\sin(2x)}{4}$	11
meijerg	$\frac{\sqrt{\pi} \left(\frac{2x}{\sqrt{\pi}} - \frac{\sin(2x)}{\sqrt{\pi}} \right)}{4}$	22
orering	$x \sin(x)^2 - \frac{\cos(x)\sin(x)}{2} + \frac{x(2\cos(x)^2 - 2\sin(x)^2)}{4}$	30
norman	$\frac{\tan(\frac{x}{2})^3 + x \tan(\frac{x}{2})^2 + \frac{x}{2} + \frac{x \tan(\frac{x}{2})^4}{2} - \tan(\frac{x}{2})}{(1 + \tan(\frac{x}{2})^2)^2}$	45

input

```
int(sin(x)^2,x,method=_RETURNVERBOSE)
```

output

```
1/2*x-1/2*cos(x)*sin(x)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \sin^2(x) dx = -\frac{1}{2} \cos(x) \sin(x) + \frac{1}{2} x$$

input

```
integrate(sin(x)^2,x, algorithm="fricas")
```

output

```
-1/2*cos(x)*sin(x) + 1/2*x
```

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \sin^2(x) dx = \frac{x}{2} - \frac{\sin(x) \cos(x)}{2}$$

input `integrate(sin(x)**2,x)`

output `x/2 - sin(x)*cos(x)/2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \sin^2(x) dx = \frac{1}{2}x - \frac{1}{4} \sin(2x)$$

input `integrate(sin(x)^2,x, algorithm="maxima")`

output `1/2*x - 1/4*sin(2*x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \sin^2(x) dx = \frac{1}{2}x - \frac{1}{4} \sin(2x)$$

input `integrate(sin(x)^2,x, algorithm="giac")`

output `1/2*x - 1/4*sin(2*x)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \sin^2(x) dx = \frac{x}{2} - \frac{\sin(2x)}{4}$$

input `int(sin(x)^2,x)`

output `x/2 - sin(2*x)/4`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \sin^2(x) dx = -\frac{\cos(x) \sin(x)}{2} + \frac{x}{2}$$

input `int(sin(x)^2,x)`

output `(- cos(x)*sin(x) + x)/2`

3.42 $\int \csc(x) \sqrt{A^2 + B^2 \sin^2(x)} dx$

Optimal result	285
Mathematica [B] (verified)	285
Rubi [A] (verified)	286
Maple [C] (verified)	288
Fricas [B] (verification not implemented)	289
Sympy [F]	290
Maxima [B] (verification not implemented)	290
Giac [F]	291
Mupad [F(-1)]	291
Reduce [F]	291

Optimal result

Integrand size = 19, antiderivative size = 49

$$\int \csc(x) \sqrt{A^2 + B^2 \sin^2(x)} dx = -B \arctan \left(\frac{B \cos(x)}{\sqrt{A^2 + B^2 \sin^2(x)}} \right) - A \operatorname{arctanh} \left(\frac{A \cos(x)}{\sqrt{A^2 + B^2 \sin^2(x)}} \right)$$

output

`-B*arctan(B*cos(x)/(A^2+B^2*sin(x)^2)^(1/2))-A*arctanh(A*cos(x)/(A^2+B^2*sin(x)^2)^(1/2))`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 99 vs. 2(49) = 98.

Time = 0.10 (sec) , antiderivative size = 99, normalized size of antiderivative = 2.02

$$\int \csc(x) \sqrt{A^2 + B^2 \sin^2(x)} dx = -\sqrt{A^2} \operatorname{arctanh} \left(\frac{\sqrt{2} \sqrt{A^2} \cos(x)}{\sqrt{2A^2 + B^2 - B^2 \cos(2x)}} \right) + \sqrt{-B^2} \log \left(\sqrt{2} \sqrt{-B^2} \cos(x) + \sqrt{2A^2 + B^2 - B^2 \cos(2x)} \right)$$

input `Integrate[Csc[x]*Sqrt[A^2 + B^2*Sin[x]^2],x]`

output `-(Sqrt[A^2]*ArcTanh[(Sqrt[2]*Sqrt[A^2]*Cos[x])/Sqrt[2*A^2 + B^2 - B^2*Cos[2*x]]) + Sqrt[-B^2]*Log[Sqrt[2]*Sqrt[-B^2]*Cos[x] + Sqrt[2*A^2 + B^2 - B^2*Cos[2*x]]]`

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.16, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {3042, 3665, 301, 224, 216, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \csc(x) \sqrt{A^2 + B^2 \sin^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{A^2 + B^2 \sin^2(x)}}{\sin(x)} dx \\
 & \quad \downarrow \text{3665} \\
 & - \int \frac{\sqrt{A^2 + B^2 - B^2 \cos^2(x)}}{1 - \cos^2(x)} d \cos(x) \\
 & \quad \downarrow \text{301} \\
 & A^2 \left(- \int \frac{1}{(1 - \cos^2(x)) \sqrt{A^2 + B^2 - B^2 \cos^2(x)}} d \cos(x) \right) - \\
 & \quad B^2 \int \frac{1}{\sqrt{A^2 + B^2 - B^2 \cos^2(x)}} d \cos(x) \\
 & \quad \downarrow \text{224} \\
 & A^2 \left(- \int \frac{1}{(1 - \cos^2(x)) \sqrt{A^2 + B^2 - B^2 \cos^2(x)}} d \cos(x) \right) - \\
 & \quad B^2 \int \frac{1}{\frac{B^2 \cos^2(x)}{A^2 + B^2 - B^2 \cos^2(x)} + 1} d \frac{\cos(x)}{\sqrt{A^2 + B^2 - B^2 \cos^2(x)}}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 216 \\
& A^2 \left(- \int \frac{1}{(1 - \cos^2(x)) \sqrt{A^2 + B^2 - B^2 \cos^2(x)}} d \cos(x) \right) - \\
& \quad B \arctan \left(\frac{B \cos(x)}{\sqrt{A^2 - B^2 \cos^2(x) + B^2}} \right) \\
& \downarrow 291 \\
& A^2 \left(- \int \frac{1}{1 - \frac{A^2 \cos^2(x)}{A^2 + B^2 - B^2 \cos^2(x)}} d \frac{\cos(x)}{\sqrt{A^2 + B^2 - B^2 \cos^2(x)}} \right) - \\
& \quad B \arctan \left(\frac{B \cos(x)}{\sqrt{A^2 - B^2 \cos^2(x) + B^2}} \right) \\
& \downarrow 219 \\
& -B \arctan \left(\frac{B \cos(x)}{\sqrt{A^2 - B^2 \cos^2(x) + B^2}} \right) - A \operatorname{arctanh} \left(\frac{A \cos(x)}{\sqrt{A^2 - B^2 \cos^2(x) + B^2}} \right)
\end{aligned}$$

input `Int[Csc[x]*Sqrt[A^2 + B^2*Sin[x]^2],x]`

output `-(B*ArcTan[(B*Cos[x])/Sqrt[A^2 + B^2 - B^2*Cos[x]^2]]) - A*ArcTanh[(A*Cos[x])/Sqrt[A^2 + B^2 - B^2*Cos[x]^2]]`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

rule 291 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

rule 301 `Int[((a_) + (b_.)*(x_)^2)^(p_.)/((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[b/d Int[(a + b*x^2)^(p - 1), x], x] - Simp[(b*c - a*d)/d Int[(a + b*x^2)^(p - 1)/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[p, 0] && (EqQ[p, 1/2] || EqQ[Denominator[p], 4] || (EqQ[p, 2/3] && EqQ[b*c + 3*a*d, 0]))`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3665 `Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_.), x_Symbol] := With[{ff = FreeFactors[Cos[e + f*x], x]}, Simp[-ff/f Subst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b - b*ff^2*x^2)^p, x], x, Cos[e + f*x]/ff], x] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.52 (sec) , antiderivative size = 149, normalized size of antiderivative = 3.04

method	result
default	$-\frac{\sqrt{(A^2+B^2 \sin(x)^2) \cos(x)^2} \left(A \operatorname{csgn}(A) \ln \left(-\frac{A^2 \sin(x)^2 - B^2 \sin(x)^2 - 2 \operatorname{csgn}(A) A \sqrt{(A^2+B^2 \sin(x)^2) \cos(x)^2 - 2A^2}}{\sin(x)^2} \right) - B \operatorname{csgn}(B) \right)}{2 \cos(x) \sqrt{A^2+B^2 \sin(x)^2}}$

input `int((A^2+B^2*sin(x)^2)^(1/2)/sin(x),x,method=_RETURNVERBOSE)`

output

```
-1/2*((A^2+B^2*sin(x)^2)*cos(x)^2)^(1/2)*(A*csgn(A)*ln(-(A^2*sin(x)^2-B^2*
sin(x)^2-2*csgn(A)*A*((A^2+B^2*sin(x)^2)*cos(x)^2)^(1/2)-2*A^2)/sin(x)^2)-
B*csgn(B)*arctan(1/2*csgn(B)/B*(2*B^2*sin(x)^2+A^2-B^2)/((A^2+B^2*sin(x)^2
)*cos(x)^2)^(1/2)))/cos(x)/(A^2+B^2*sin(x)^2)^(1/2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 244 vs. $2(45) = 90$.

Time = 0.16 (sec) , antiderivative size = 244, normalized size of antiderivative = 4.98

$$\int \csc(x) \sqrt{A^2 + B^2 \sin^2(x)} dx$$

$$= \frac{1}{2} B \arctan \left(-\frac{(A^4 + 2A^2B^2 + B^4) \cos(x) \sin(x) - 2(2B^3 \cos(x)^3 - (A^2B + B^3) \cos(x)) \sqrt{-B^2 \cos(x)^2 + A^2 + B^2}}{4B^4 \cos(x)^4 + A^4 + 2A^2B^2 + B^4 - (A^4 + 6A^2B^2 + 5B^4) \cos(x)^2} \right) - \frac{1}{2} B \arctan \left(\frac{\sin(x)}{\cos(x)} \right) - \frac{1}{2} A \log \left(-B^2 \cos(x)^2 + AB \cos(x) \sin(x) + A^2 + B^2 + \sqrt{-B^2 \cos(x)^2 + A^2 + B^2} (A \cos(x) + B \sin(x)) \right) + \frac{1}{2} A \log \left(-B^2 \cos(x)^2 - AB \cos(x) \sin(x) + A^2 + B^2 - \sqrt{-B^2 \cos(x)^2 + A^2 + B^2} (A \cos(x) - B \sin(x)) \right)$$

input

```
integrate((A^2+B^2*sin(x)^2)^(1/2)/sin(x),x, algorithm="fricas")
```

output

```
1/2*B*arctan(-((A^4 + 2*A^2*B^2 + B^4)*cos(x)*sin(x) - 2*(2*B^3*cos(x)^3 -
(A^2*B + B^3)*cos(x))*sqrt(-B^2*cos(x)^2 + A^2 + B^2))/(4*B^4*cos(x)^4 +
A^4 + 2*A^2*B^2 + B^4 - (A^4 + 6*A^2*B^2 + 5*B^4)*cos(x)^2)) - 1/2*B*arcta
n(sin(x)/cos(x)) - 1/2*A*log(-B^2*cos(x)^2 + A*B*cos(x)*sin(x) + A^2 + B^2
+ sqrt(-B^2*cos(x)^2 + A^2 + B^2)*(A*cos(x) + B*sin(x))) + 1/2*A*log(-B^2
*cos(x)^2 - A*B*cos(x)*sin(x) + A^2 + B^2 - sqrt(-B^2*cos(x)^2 + A^2 + B^2
)*(A*cos(x) - B*sin(x)))
```

Sympy [F]

$$\int \csc(x) \sqrt{A^2 + B^2 \sin^2(x)} dx = \int \frac{\sqrt{A^2 + B^2 \sin^2(x)}}{\sin(x)} dx$$

input `integrate((A**2+B**2*sin(x)**2)**(1/2)/sin(x),x)`

output `Integral(sqrt(A**2 + B**2*sin(x)**2)/sin(x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 116 vs. $2(45) = 90$.

Time = 0.11 (sec) , antiderivative size = 116, normalized size of antiderivative = 2.37

$$\int \csc(x) \sqrt{A^2 + B^2 \sin^2(x)} dx = -B \arcsin\left(\frac{B^2 \cos(x)}{\sqrt{A^2 B^2 + B^4}}\right) - \frac{1}{2} A \log\left(B^2 - \frac{A^2}{\cos(x) - 1} - \frac{\sqrt{-B^2 \cos(x)^2 + A^2 + B^2 A}}{\cos(x) - 1}\right) + \frac{1}{2} A \log\left(-B^2 + \frac{A^2}{\cos(x) + 1} + \frac{\sqrt{-B^2 \cos(x)^2 + A^2 + B^2 A}}{\cos(x) + 1}\right)$$

input `integrate((A^2+B^2*sin(x)^2)^(1/2)/sin(x),x, algorithm="maxima")`

output `-B*arcsin(B^2*cos(x)/sqrt(A^2*B^2 + B^4)) - 1/2*A*log(B^2 - A^2/(cos(x) - 1) - sqrt(-B^2*cos(x)^2 + A^2 + B^2)*A/(cos(x) - 1)) + 1/2*A*log(-B^2 + A^2/(cos(x) + 1) + sqrt(-B^2*cos(x)^2 + A^2 + B^2)*A/(cos(x) + 1))`

Giac [F]

$$\int \csc(x) \sqrt{A^2 + B^2 \sin^2(x)} dx = \int \frac{\sqrt{B^2 \sin(x)^2 + A^2}}{\sin(x)} dx$$

input `integrate((A^2+B^2*sin(x)^2)^(1/2)/sin(x),x, algorithm="giac")`

output `integrate(sqrt(B^2*sin(x)^2 + A^2)/sin(x), x)`

Mupad [F(-1)]

Timed out.

$$\int \csc(x) \sqrt{A^2 + B^2 \sin^2(x)} dx = \int \frac{\sqrt{A^2 + B^2 \sin(x)^2}}{\sin(x)} dx$$

input `int((B^2*sin(x)^2 + A^2)^(1/2)/sin(x),x)`

output `int((B^2*sin(x)^2 + A^2)^(1/2)/sin(x), x)`

Reduce [F]

$$\int \csc(x) \sqrt{A^2 + B^2 \sin^2(x)} dx = \int \frac{\sqrt{\sin(x)^2 b^2 + a^2}}{\sin(x)} dx$$

input `int((A^2+B^2*sin(x)^2)^(1/2)/sin(x),x)`

output `int(sqrt(sin(x)**2*b**2 + a**2)/sin(x),x)`

3.43 $\int \frac{1}{1+\cos(x)} dx$

Optimal result	292
Mathematica [A] (verified)	292
Rubi [A] (verified)	293
Maple [A] (verified)	294
Fricas [A] (verification not implemented)	294
Sympy [A] (verification not implemented)	294
Maxima [A] (verification not implemented)	295
Giac [B] (verification not implemented)	295
Mupad [B] (verification not implemented)	296
Reduce [B] (verification not implemented)	296

Optimal result

Integrand size = 6, antiderivative size = 9

$$\int \frac{1}{1+\cos(x)} dx = \frac{\sin(x)}{1+\cos(x)}$$

output `sin(x)/(1+cos(x))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.67

$$\int \frac{1}{1+\cos(x)} dx = \tan\left(\frac{x}{2}\right)$$

input `Integrate[(1 + Cos[x])^(-1),x]`

output `Tan[x/2]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\cos(x) + 1} dx$$

↓ 3042

$$\int \frac{1}{\sin\left(x + \frac{\pi}{2}\right) + 1} dx$$

↓ 3127

$$\frac{\sin(x)}{\cos(x) + 1}$$

input `Int[(1 + Cos[x])^(-1), x]`

output `Sin[x]/(1 + Cos[x])`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3127 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.56

method	result	size
default	$\tan\left(\frac{x}{2}\right)$	5
norman	$\tan\left(\frac{x}{2}\right)$	5
parallelrisk	$\tan\left(\frac{x}{2}\right)$	5
risch	$\frac{2i}{1+e^{ix}}$	13

input `int(1/(1+cos(x)),x,method=_RETURNVERBOSE)`

output `tan(1/2*x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + \cos(x)} dx = \frac{\sin(x)}{\cos(x) + 1}$$

input `integrate(1/(1+cos(x)),x, algorithm="fricas")`

output `sin(x)/(cos(x) + 1)`

Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.33

$$\int \frac{1}{1 + \cos(x)} dx = \tan\left(\frac{x}{2}\right)$$

input `integrate(1/(1+cos(x)),x)`

output `tan(x/2)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + \cos(x)} dx = \frac{\sin(x)}{\cos(x) + 1}$$

input `integrate(1/(1+cos(x)),x, algorithm="maxima")`

output `sin(x)/(cos(x) + 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 30 vs. 2(9) = 18.

Time = 0.12 (sec) , antiderivative size = 30, normalized size of antiderivative = 3.33

$$\int \frac{1}{1 + \cos(x)} dx = -\frac{2 \tan\left(\frac{1}{2}x\right)}{(x^2 + 1)\left(\frac{x^2-1}{x^2+1} - 1\right)}$$

input `integrate(1/(1+cos(x)),x, algorithm="giac")`

output `-2*tan(1/2*x)/((x^2 + 1)*((x^2 - 1)/(x^2 + 1) - 1))`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 4, normalized size of antiderivative = 0.44

$$\int \frac{1}{1 + \cos(x)} dx = \tan\left(\frac{x}{2}\right)$$

input `int(1/(cos(x) + 1),x)`

output `tan(x/2)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 4, normalized size of antiderivative = 0.44

$$\int \frac{1}{1 + \cos(x)} dx = \tan\left(\frac{x}{2}\right)$$

input `int(1/(1+cos(x)),x)`

output `tan(x/2)`

3.44 $\int e^x x dx$

Optimal result	297
Mathematica [A] (verified)	297
Rubi [A] (verified)	298
Maple [A] (verified)	299
Fricas [A] (verification not implemented)	299
Sympy [A] (verification not implemented)	300
Maxima [A] (verification not implemented)	300
Giac [A] (verification not implemented)	300
Mupad [B] (verification not implemented)	301
Reduce [B] (verification not implemented)	301

Optimal result

Integrand size = 5, antiderivative size = 11

$$\int e^x x dx = -e^x + e^x x$$

output `-exp(x)+exp(x)*x`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.64

$$\int e^x x dx = e^x(-1 + x)$$

input `Integrate[E^x*x,x]`

output `E^x*(-1 + x)`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int e^x x \, dx \\ \downarrow 2607 \\ e^x x - \int e^x \, dx \\ \downarrow 2624 \\ e^x x - e^x \end{array}$$

input `Int [E^x*x, x]`

output `-E^x + E^x*x`

Defintions of rubi rules used

rule 2607 `Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]`

rule 2624 `Int[((F_)^(v_))^(n_.), x_Symbol] := Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /; FreeQ[{F, n}, x] && LinearQ[v, x]`

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.64

method	result	size
gospers	$(-1 + x)e^x$	7
risch	$(-1 + x)e^x$	7
orering	$(-1 + x)e^x$	7
default	$-e^x + e^x x$	10
norman	$-e^x + e^x x$	10
parallelrisch	$-e^x + e^x x$	10
parts	$-e^x + e^x x$	10
meijerg	$1 - \frac{(-2x+2)e^x}{2}$	12

input `int(exp(x)*x,x,method=_RETURNVERBOSE)`

output `(-1+x)*exp(x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.55

$$\int e^x x dx = (x - 1)e^x$$

input `integrate(exp(x)*x,x, algorithm="fricas")`

output `(x - 1)*e^x`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.45

$$\int e^x x dx = (x - 1) e^x$$

input `integrate(exp(x)*x,x)`

output `(x - 1)*exp(x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.55

$$\int e^x x dx = (x - 1) e^x$$

input `integrate(exp(x)*x,x, algorithm="maxima")`

output `(x - 1)*e^x`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.55

$$\int e^x x dx = (x - 1) e^x$$

input `integrate(exp(x)*x,x, algorithm="giac")`

output `(x - 1)*e^x`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.55

$$\int e^x x dx = e^x (x - 1)$$

input `int(x*exp(x),x)`

output `exp(x)*(x - 1)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.64

$$\int e^x x dx = e^x (x - 1)$$

input `int(exp(x)*x,x)`

output `e**x*(x - 1)`

3.45 $\int \frac{e^x x}{(1+x)^2} dx$

Optimal result	302
Mathematica [A] (verified)	302
Rubi [A] (verified)	303
Maple [A] (verified)	303
Fricas [A] (verification not implemented)	304
Sympy [A] (verification not implemented)	304
Maxima [A] (verification not implemented)	305
Giac [B] (verification not implemented)	305
Mupad [B] (verification not implemented)	306
Reduce [B] (verification not implemented)	306

Optimal result

Integrand size = 10, antiderivative size = 9

$$\int \frac{e^x x}{(1+x)^2} dx = \frac{e^x}{1+x}$$

output `exp(x)/(1+x)`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \frac{e^x x}{(1+x)^2} dx = \frac{e^x}{1+x}$$

input `Integrate[(E^x*x)/(1 + x)^2,x]`

output `E^x/(1 + x)`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2627}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^x x}{(x+1)^2} dx$$

↓ 2627

$$\frac{e^x}{x+1}$$

input `Int[(E^x*x)/(1 + x)^2,x]`

output `E^x/(1 + x)`

Defintions of rubi rules used

rule 2627 `Int[(F_)^(v_)*((d_.) + (e_.)*(x_))^(m_.)*((f_.) + (g_.)*(x_)), x_Symbol] :>
Simp[g*(d + e*x)^(m + 1)*(F^v/(D[v, x]*e*Log[F])), x] /; FreeQ[{F, d, e, f, g, m}, x] && LinearQ[v, x] && EqQ[e*g*(m + 1) - D[v, x]*(e*f - d*g)*Log[F], 0]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

method	result	size
gosper	$\frac{e^x}{1+x}$	9
default	$\frac{e^x}{1+x}$	9
norman	$\frac{e^x}{1+x}$	9
risch	$\frac{e^x}{1+x}$	9
parallelrisc	$\frac{e^x}{1+x}$	9
orering	$\frac{e^x}{1+x}$	9

input `int(exp(x)*x/(1+x)^2,x,method=_RETURNVERBOSE)`

output `exp(x)/(1+x)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

$$\int \frac{e^x x}{(1+x)^2} dx = \frac{e^x}{x+1}$$

input `integrate(exp(x)*x/(1+x)^2,x, algorithm="fricas")`

output `e^x/(x + 1)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.56

$$\int \frac{e^x x}{(1+x)^2} dx = \frac{e^x}{x+1}$$

input `integrate(exp(x)*x/(1+x)**2,x)`

output `exp(x)/(x + 1)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

$$\int \frac{e^x x}{(1+x)^2} dx = \frac{e^x}{x+1}$$

input `integrate(exp(x)*x/(1+x)^2,x, algorithm="maxima")`

output `e^x/(x + 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 30 vs. 2(8) = 16.

Time = 0.12 (sec) , antiderivative size = 30, normalized size of antiderivative = 3.33

$$\int \frac{e^x x}{(1+x)^2} dx = -\frac{e^{-(x+1)\left(\frac{1}{x+1}-1\right)}}{(x+1)\left(\frac{1}{x+1}-1\right)-1}$$

input `integrate(exp(x)*x/(1+x)^2,x, algorithm="giac")`

output `-e^(-(x + 1)*(1/(x + 1) - 1))/((x + 1)*(1/(x + 1) - 1) - 1)`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

$$\int \frac{e^x x}{(1+x)^2} dx = \frac{e^x}{x+1}$$

input `int((x*exp(x))/(x + 1)^2,x)`

output `exp(x)/(x + 1)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \frac{e^x x}{(1+x)^2} dx = \frac{e^x}{x+1}$$

input `int(exp(x)*x/(1+x)^2,x)`

output `e**x/(x + 1)`

3.46 $\int e^{x^2}(1 + 2x^2) dx$

Optimal result	307
Mathematica [A] (verified)	307
Rubi [A] (verified)	308
Maple [A] (verified)	309
Fricas [A] (verification not implemented)	309
Sympy [A] (verification not implemented)	310
Maxima [A] (verification not implemented)	310
Giac [A] (verification not implemented)	310
Mupad [B] (verification not implemented)	311
Reduce [B] (verification not implemented)	311

Optimal result

Integrand size = 13, antiderivative size = 7

$$\int e^{x^2}(1 + 2x^2) dx = e^{x^2}x$$

output `exp(x^2)*x`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int e^{x^2}(1 + 2x^2) dx = e^{x^2}x$$

input `Integrate[E^x^2*(1 + 2*x^2),x]`

output `E^x^2*x`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {2656, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{x^2} (2x^2 + 1) dx$$

$$\downarrow 2656$$

$$\int (2e^{x^2} x^2 + e^{x^2}) dx$$

$$\downarrow 2009$$

$$e^{x^2} x$$

input `Int[E^x^2*(1 + 2*x^2),x]`

output `E^x^2*x`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2656 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*(Px_), x_Symbol] := Int[ExpandLinearProduct[F^(a + b*(c + d*x)^n), Px, c, d, x], x] /; FreeQ[{F, a, b, c, d, n}, x] && PolynomialQ[Px, x]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

method	result	size
gospers	$e^{x^2} x$	7
default	$e^{x^2} x$	7
norman	$e^{x^2} x$	7
risch	$e^{x^2} x$	7
parallelrisch	$e^{x^2} x$	7
orering	$e^{x^2} x$	7
meijerg	$i \left(-ie^{x^2} x + \frac{i \operatorname{erfi}(x)\sqrt{\pi}}{2} \right) + \frac{\operatorname{erfi}(x)\sqrt{\pi}}{2}$	29
parts	$\sqrt{\pi} \operatorname{erfi}(x) x^2 + \frac{\operatorname{erfi}(x)\sqrt{\pi}}{2} - 2\sqrt{\pi} \left(\frac{x^2 \operatorname{erfi}(x)}{2} - \frac{\frac{e^{x^2} x - \operatorname{erfi}(x)\sqrt{\pi}}{4}}{\sqrt{\pi}} \right)$	51

input `int(exp(x^2)*(2*x^2+1),x,method=_RETURNVERBOSE)`output `exp(x^2)*x`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int e^{x^2} (1 + 2x^2) dx = x e^{(x^2)}$$

input `integrate(exp(x^2)*(2*x^2+1),x, algorithm="fricas")`output `x*e^(x^2)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.71

$$\int e^{x^2} (1 + 2x^2) dx = xe^{x^2}$$

input `integrate(exp(x**2)*(2*x**2+1),x)`

output `x*exp(x**2)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int e^{x^2} (1 + 2x^2) dx = xe^{(x^2)}$$

input `integrate(exp(x^2)*(2*x^2+1),x, algorithm="maxima")`

output `x*e^(x^2)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int e^{x^2} (1 + 2x^2) dx = xe^{(x^2)}$$

input `integrate(exp(x^2)*(2*x^2+1),x, algorithm="giac")`

output `x*e^(x^2)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int e^{x^2} (1 + 2x^2) dx = x e^{x^2}$$

input `int(exp(x^2)*(2*x^2 + 1),x)`

output `x*exp(x^2)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int e^{x^2} (1 + 2x^2) dx = e^{x^2} x$$

input `int(exp(x^2)*(2*x^2+1),x)`

output `e**(x**2)*x`

3.47 $\int e^{x^2} dx$

Optimal result	312
Mathematica [A] (verified)	312
Rubi [A] (verified)	313
Maple [A] (verified)	313
Fricas [A] (verification not implemented)	314
Sympy [A] (verification not implemented)	314
Maxima [C] (verification not implemented)	315
Giac [C] (verification not implemented)	315
Mupad [B] (verification not implemented)	315
Reduce [B] (verification not implemented)	316

Optimal result

Integrand size = 5, antiderivative size = 11

$$\int e^{x^2} dx = \frac{1}{2}\sqrt{\pi}\operatorname{erfi}(x)$$

output `1/2*erfi(x)*Pi^(1/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int e^{x^2} dx = \frac{1}{2}\sqrt{\pi}\operatorname{erfi}(x)$$

input `Integrate[E^x^2,x]`

output `(Sqrt[Pi]*Erfi[x])/2`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2633}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{x^2} dx$$

↓ 2633

$$\frac{1}{2}\sqrt{\pi}\operatorname{erfi}(x)$$

input `Int [E^x^2,x]`

output `(Sqrt [Pi]*Erfi [x])/2`

Defintions of rubi rules used

rule 2633 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))2), x_Symbol] :> Simp[F^a*Sqrt [Pi]*(Erfi[(c + d*x)*Rt [b*Log[F], 2]]/(2*d*Rt [b*Log[F], 2])), x] /; FreeQ[{ F, a, b, c, d}, x] && PosQ[b]`

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

method	result	size
default	$\frac{\operatorname{erfi}(x)\sqrt{\pi}}{2}$	8
meijerg	$\frac{\operatorname{erfi}(x)\sqrt{\pi}}{2}$	8
risch	$\frac{\operatorname{erfi}(x)\sqrt{\pi}}{2}$	8

input `int(exp(x^2),x,method=_RETURNVERBOSE)`

output `1/2*erfi(x)*Pi^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.64

$$\int e^{x^2} dx = \frac{1}{2} \sqrt{\pi} \operatorname{erfi}(x)$$

input `integrate(exp(x^2),x, algorithm="fricas")`

output `1/2*sqrt(pi)*erfi(x)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int e^{x^2} dx = \frac{\sqrt{\pi} \operatorname{erfi}(x)}{2}$$

input `integrate(exp(x**2),x)`

output `sqrt(pi)*erfi(x)/2`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int e^{x^2} dx = -\frac{1}{2}i\sqrt{\pi} \operatorname{erf}(ix)$$

input `integrate(exp(x^2),x, algorithm="maxima")`

output `-1/2*I*sqrt(pi)*erf(I*x)`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.13 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int e^{x^2} dx = \frac{1}{2}i\sqrt{\pi} \operatorname{erf}(-ix)$$

input `integrate(exp(x^2),x, algorithm="giac")`

output `1/2*I*sqrt(pi)*erf(-I*x)`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.64

$$\int e^{x^2} dx = \frac{\sqrt{\pi} \operatorname{erfi}(x)}{2}$$

input `int(exp(x^2),x)`

output `(pi^(1/2)*erfi(x))/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int e^{x^2} dx = -\frac{\sqrt{\pi} \operatorname{erf}(ix) i}{2}$$

input `int(exp(x^2),x)`

output `(- sqrt(pi)*erf(i*x)*i)/2`

3.48 $\int \frac{e^x}{x} dx$

Optimal result	317
Mathematica [A] (verified)	317
Rubi [A] (verified)	318
Maple [B] (verified)	318
Fricas [A] (verification not implemented)	319
Sympy [A] (verification not implemented)	319
Maxima [A] (verification not implemented)	320
Giac [A] (verification not implemented)	320
Mupad [B] (verification not implemented)	320
Reduce [B] (verification not implemented)	321

Optimal result

Integrand size = 7, antiderivative size = 2

$$\int \frac{e^x}{x} dx = \text{ExpIntegralEi}(x)$$

output Ei(x)

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{e^x}{x} dx = \text{ExpIntegralEi}(x)$$

input Integrate[E^x/x,x]

output ExpIntegralEi[x]

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2609}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^x}{x} dx$$

↓ 2609

$$\text{ExpIntegralEi}(x)$$

input `Int [E^x/x, x]`

output `ExpIntegralEi [x]`

Defintions of rubi rules used

rule 2609 `Int[(F_)^((g_.)*((e_.) + (f_.)*(x_)))/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 7 vs. $2(2) = 4$.

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 4.00

method	result	size
default	$-\text{expIntegral}_1(-x)$	8
risch	$-\text{expIntegral}_1(-x)$	8
meijerg	$\ln(x) + i\pi - \ln(-x) - \text{expIntegral}_1(-x)$	21

input `int(exp(x)/x,x,method=_RETURNVERBOSE)`

output `-Ei(1,-x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{e^x}{x} dx = \text{Ei}(x)$$

input `integrate(exp(x)/x,x, algorithm="fricas")`

output `Ei(x)`

Sympy [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{e^x}{x} dx = \text{Ei}(x)$$

input `integrate(exp(x)/x,x)`

output `Ei(x)`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{e^x}{x} dx = \text{Ei}(x)$$

input `integrate(exp(x)/x,x, algorithm="maxima")`

output `Ei(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{e^x}{x} dx = \text{Ei}(x)$$

input `integrate(exp(x)/x,x, algorithm="giac")`

output `Ei(x)`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{e^x}{x} dx = \text{ei}(x)$$

input `int(exp(x)/x,x)`

output `ei(x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{e^x}{x} dx = ei(x)$$

input `int(exp(x)/x,x)`

output `ei(x)`

3.49 $\int \frac{x}{1+x^3} dx$

Optimal result	322
Mathematica [A] (verified)	322
Rubi [A] (verified)	323
Maple [A] (verified)	325
Fricas [A] (verification not implemented)	325
Sympy [A] (verification not implemented)	326
Maxima [A] (verification not implemented)	326
Giac [A] (verification not implemented)	326
Mupad [B] (verification not implemented)	327
Reduce [B] (verification not implemented)	327

Optimal result

Integrand size = 9, antiderivative size = 41

$$\int \frac{x}{1+x^3} dx = -\frac{\arctan\left(\frac{1-2x}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{3}\log(1+x) + \frac{1}{6}\log(1-x+x^2)$$

output `-1/3*ln(1+x)+1/6*ln(x^2-x+1)-1/3*arctan(1/3*(1-2*x)*3^(1/2))*3^(1/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.98

$$\int \frac{x}{1+x^3} dx = \frac{\arctan\left(\frac{-1+2x}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{3}\log(1+x) + \frac{1}{6}\log(1-x+x^2)$$

input `Integrate[x/(1 + x^3), x]`

output `ArcTan[(-1 + 2*x)/Sqrt[3]]/Sqrt[3] - Log[1 + x]/3 + Log[1 - x + x^2]/6`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.10, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.778$, Rules used = {821, 16, 1142, 25, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x}{x^3 + 1} dx \\
 & \quad \downarrow \text{821} \\
 & \frac{1}{3} \int \frac{x+1}{x^2-x+1} dx - \frac{1}{3} \int \frac{1}{x+1} dx \\
 & \quad \downarrow \text{16} \\
 & \frac{1}{3} \int \frac{x+1}{x^2-x+1} dx - \frac{1}{3} \log(x+1) \\
 & \quad \downarrow \text{1142} \\
 & \frac{1}{3} \left(\frac{3}{2} \int \frac{1}{x^2-x+1} dx + \frac{1}{2} \int -\frac{1-2x}{x^2-x+1} dx \right) - \frac{1}{3} \log(x+1) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{3} \left(\frac{3}{2} \int \frac{1}{x^2-x+1} dx - \frac{1}{2} \int \frac{1-2x}{x^2-x+1} dx \right) - \frac{1}{3} \log(x+1) \\
 & \quad \downarrow \text{1083} \\
 & \frac{1}{3} \left(-\frac{1}{2} \int \frac{1-2x}{x^2-x+1} dx - 3 \int \frac{1}{-(2x-1)^2-3} d(2x-1) \right) - \frac{1}{3} \log(x+1) \\
 & \quad \downarrow \text{217} \\
 & \frac{1}{3} \left(\sqrt{3} \arctan \left(\frac{2x-1}{\sqrt{3}} \right) - \frac{1}{2} \int \frac{1-2x}{x^2-x+1} dx \right) - \frac{1}{3} \log(x+1) \\
 & \quad \downarrow \text{1103} \\
 & \frac{1}{3} \left(\sqrt{3} \arctan \left(\frac{2x-1}{\sqrt{3}} \right) + \frac{1}{2} \log(x^2-x+1) \right) - \frac{1}{3} \log(x+1)
 \end{aligned}$$

input

Int[x/(1 + x^3), x]

output
$$-1/3 \cdot \text{Log}[1 + x] + (\text{Sqrt}[3] \cdot \text{ArcTan}[(-1 + 2x)/\text{Sqrt}[3]] + \text{Log}[1 - x + x^2]/2)/3$$

Defintions of rubi rules used

rule 16
$$\text{Int}[(c_)/((a_)+(b_)(x_)), x_Symbol] \rightarrow \text{Simp}[c \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$$

rule 25
$$\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[Fx, x], x]$$

rule 217
$$\text{Int}(((a_) + (b_)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-(\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2]))^{(-1)} \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \& \ \& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$$

rule 821
$$\text{Int}[(x_)/((a_) + (b_)(x_)^3), x_Symbol] \rightarrow \text{Simp}[-(3 \cdot \text{Rt}[a, 3] \cdot \text{Rt}[b, 3])^{(-1)} \quad \text{Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3] \cdot x), x], x] + \text{Simp}[1/(3 \cdot \text{Rt}[a, 3] \cdot \text{Rt}[b, 3]) \quad \text{Int}[(\text{Rt}[a, 3] + \text{Rt}[b, 3] \cdot x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3] \cdot \text{Rt}[b, 3] \cdot x + \text{Rt}[b, 3]^2 \cdot x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$$

rule 1083
$$\text{Int}(((a_) + (b_)(x_) + (c_)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[-2 \quad \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4 \cdot a \cdot c - x^2, x], x], x, b + 2 \cdot c \cdot x], x] /; \text{FreeQ}[\{a, b, c\}, x]$$

rule 1103
$$\text{Int}(((d_) + (e_)(x_))/((a_) + (b_)(x_) + (c_)(x_)^2), x_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$$

rule 1142
$$\text{Int}(((d_) + (e_)(x_))/((a_) + (b_)(x_) + (c_)(x_)^2), x_Symbol] \rightarrow \text{Simp}[(2 \cdot c \cdot d - b \cdot e)/(2 \cdot c) \quad \text{Int}[1/(a + b \cdot x + c \cdot x^2), x], x] + \text{Simp}[e/(2 \cdot c) \quad \text{Int}[(b + 2 \cdot c \cdot x)/(a + b \cdot x + c \cdot x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$$

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.80

method	result	size
risch	$-\frac{\ln(1+x)}{3} + \frac{\ln(x^2-x+1)}{6} + \frac{\sqrt{3} \arctan\left(\frac{2\left(x-\frac{1}{2}\right)\sqrt{3}}{3}\right)}{3}$	33
default	$-\frac{\ln(1+x)}{3} + \frac{\ln(x^2-x+1)}{6} + \frac{\sqrt{3} \arctan\left(\frac{(2x-1)\sqrt{3}}{3}\right)}{3}$	35
meijerg	$-\frac{x^2 \ln\left(1+(x^3)^{\frac{1}{3}}\right)}{3(x^3)^{\frac{2}{3}}} + \frac{x^2 \ln\left(1-(x^3)^{\frac{1}{3}}+(x^3)^{\frac{2}{3}}\right)}{6(x^3)^{\frac{2}{3}}} + \frac{x^2 \sqrt{3} \arctan\left(\frac{\sqrt{3}(x^3)^{\frac{1}{3}}}{2-(x^3)^{\frac{1}{3}}}\right)}{3(x^3)^{\frac{2}{3}}}$	80

input `int(x/(x^3+1),x,method=_RETURNVERBOSE)`

output `-1/3*ln(1+x)+1/6*ln(x^2-x+1)+1/3*3^(1/2)*arctan(2/3*(x-1/2)*3^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.83

$$\int \frac{x}{1+x^3} dx = \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x-1)\right) + \frac{1}{6} \log(x^2-x+1) - \frac{1}{3} \log(x+1)$$

input `integrate(x/(x^3+1),x, algorithm="fricas")`

output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - 1)) + 1/6*log(x^2 - x + 1) - 1/3*log(x + 1)`

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \frac{x}{1+x^3} dx = -\frac{\log(x+1)}{3} + \frac{\log(x^2-x+1)}{6} + \frac{\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}x - \sqrt{3}}{3}\right)}{3}$$

input `integrate(x/(x**3+1),x)`output `-log(x + 1)/3 + log(x**2 - x + 1)/6 + sqrt(3)*atan(2*sqrt(3)*x/3 - sqrt(3)/3)/3`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.83

$$\int \frac{x}{1+x^3} dx = \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x-1)\right) + \frac{1}{6} \log(x^2-x+1) - \frac{1}{3} \log(x+1)$$

input `integrate(x/(x^3+1),x, algorithm="maxima")`output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - 1)) + 1/6*log(x^2 - x + 1) - 1/3*log(x + 1)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.85

$$\int \frac{x}{1+x^3} dx = \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x-1)\right) + \frac{1}{6} \log(x^2-x+1) - \frac{1}{3} \log(|x+1|)$$

input `integrate(x/(x^3+1),x, algorithm="giac")`output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - 1)) + 1/6*log(x^2 - x + 1) - 1/3*log(abs(x + 1))`

Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.12

$$\int \frac{x}{1+x^3} dx = -\frac{\ln(x+1)}{3} - \ln\left(x - \frac{1}{2} - \frac{\sqrt{3}1i}{2}\right) \left(-\frac{1}{6} + \frac{\sqrt{3}1i}{6}\right) \\ + \ln\left(x - \frac{1}{2} + \frac{\sqrt{3}1i}{2}\right) \left(\frac{1}{6} + \frac{\sqrt{3}1i}{6}\right)$$

input `int(x/(x^3 + 1),x)`output `log(x + (3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/6 + 1/6) - log(x - (3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/6 - 1/6) - log(x + 1)/3`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.80

$$\int \frac{x}{1+x^3} dx = \frac{\sqrt{3} \operatorname{atan}\left(\frac{2x-1}{\sqrt{3}}\right)}{3} + \frac{\log(x^2 - x + 1)}{6} - \frac{\log(x + 1)}{3}$$

input `int(x/(x^3+1),x)`output `(2*sqrt(3)*atan((2*x - 1)/sqrt(3)) + log(x**2 - x + 1) - 2*log(x + 1))/6`

3.50 $\int \frac{1}{-1+x^6} dx$

Optimal result	328
Mathematica [A] (verified)	328
Rubi [A] (verified)	329
Maple [A] (verified)	331
Fricas [A] (verification not implemented)	332
Sympy [B] (verification not implemented)	332
Maxima [A] (verification not implemented)	333
Giac [A] (verification not implemented)	333
Mupad [B] (verification not implemented)	334
Reduce [B] (verification not implemented)	334

Optimal result

Integrand size = 7, antiderivative size = 47

$$\int \frac{1}{-1+x^6} dx = -\frac{\arctan\left(\frac{\sqrt{3}x}{1-x^2}\right)}{2\sqrt{3}} - \frac{\operatorname{arctanh}(x)}{3} - \frac{1}{6}\operatorname{arctanh}\left(\frac{x}{1+x^2}\right)$$

output `-1/3*arctanh(x)-1/6*arctanh(x/(x^2+1))-1/6*arctan(x*3^(1/2)/(-x^2+1))*3^(1/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.60

$$\int \frac{1}{-1+x^6} dx = \frac{1}{12} \left(-2\sqrt{3} \arctan\left(\frac{-1+2x}{\sqrt{3}}\right) - 2\sqrt{3} \arctan\left(\frac{1+2x}{\sqrt{3}}\right) + 2\log(1-x) - 2\log(1+x) + \log(1-x+x^2) - \log(1+x+x^2) \right)$$

input `Integrate[(-1 + x^6)^(-1), x]`

output `(-2*Sqrt[3]*ArcTan[(-1 + 2*x)/Sqrt[3]] - 2*Sqrt[3]*ArcTan[(1 + 2*x)/Sqrt[3]] + 2*Log[1 - x] - 2*Log[1 + x] + Log[1 - x + x^2] - Log[1 + x + x^2])/12`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.68, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.143$, Rules used = {754, 27, 219, 1142, 25, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^6 - 1} dx \\
 & \quad \downarrow 754 \\
 & -\frac{1}{3} \int \frac{1}{1 - x^2} dx - \frac{1}{3} \int \frac{2 - x}{2(x^2 - x + 1)} dx - \frac{1}{3} \int \frac{x + 2}{2(x^2 + x + 1)} dx \\
 & \quad \downarrow 27 \\
 & -\frac{1}{3} \int \frac{1}{1 - x^2} dx - \frac{1}{6} \int \frac{2 - x}{x^2 - x + 1} dx - \frac{1}{6} \int \frac{x + 2}{x^2 + x + 1} dx \\
 & \quad \downarrow 219 \\
 & -\frac{1}{6} \int \frac{2 - x}{x^2 - x + 1} dx - \frac{1}{6} \int \frac{x + 2}{x^2 + x + 1} dx - \frac{\operatorname{arctanh}(x)}{3} \\
 & \quad \downarrow 1142 \\
 & \frac{1}{6} \left(\frac{1}{2} \int -\frac{1 - 2x}{x^2 - x + 1} dx - \frac{3}{2} \int \frac{1}{x^2 - x + 1} dx \right) + \\
 & \frac{1}{6} \left(-\frac{3}{2} \int \frac{1}{x^2 + x + 1} dx - \frac{1}{2} \int \frac{2x + 1}{x^2 + x + 1} dx \right) - \frac{\operatorname{arctanh}(x)}{3} \\
 & \quad \downarrow 25 \\
 & \frac{1}{6} \left(-\frac{3}{2} \int \frac{1}{x^2 - x + 1} dx - \frac{1}{2} \int \frac{1 - 2x}{x^2 - x + 1} dx \right) + \\
 & \frac{1}{6} \left(-\frac{3}{2} \int \frac{1}{x^2 + x + 1} dx - \frac{1}{2} \int \frac{2x + 1}{x^2 + x + 1} dx \right) - \frac{\operatorname{arctanh}(x)}{3} \\
 & \quad \downarrow 1083 \\
 & \frac{1}{6} \left(3 \int \frac{1}{-(2x - 1)^2 - 3} d(2x - 1) - \frac{1}{2} \int \frac{1 - 2x}{x^2 - x + 1} dx \right) + \\
 & \frac{1}{6} \left(3 \int \frac{1}{-(2x + 1)^2 - 3} d(2x + 1) - \frac{1}{2} \int \frac{2x + 1}{x^2 + x + 1} dx \right) - \frac{\operatorname{arctanh}(x)}{3} \\
 & \quad \downarrow 217
 \end{aligned}$$

$$\frac{1}{6} \left(-\frac{1}{2} \int \frac{1-2x}{x^2-x+1} dx - \sqrt{3} \arctan \left(\frac{2x-1}{\sqrt{3}} \right) \right) +$$

$$\frac{1}{6} \left(-\frac{1}{2} \int \frac{2x+1}{x^2+x+1} dx - \sqrt{3} \arctan \left(\frac{2x+1}{\sqrt{3}} \right) \right) - \frac{\operatorname{arctanh}(x)}{3}$$

↓ 1103

$$\frac{1}{6} \left(\frac{1}{2} \log(x^2-x+1) - \sqrt{3} \arctan \left(\frac{2x-1}{\sqrt{3}} \right) \right) +$$

$$\frac{1}{6} \left(-\sqrt{3} \arctan \left(\frac{2x+1}{\sqrt{3}} \right) - \frac{1}{2} \log(x^2+x+1) \right) - \frac{\operatorname{arctanh}(x)}{3}$$

input `Int[(-1 + x^6)^(-1), x]`

output `-1/3*ArcTanh[x] + (-Sqrt[3]*ArcTan[(-1 + 2*x)/Sqrt[3]]) + Log[1 - x + x^2
]/2)/6 + (-Sqrt[3]*ArcTan[(1 + 2*x)/Sqrt[3]]) - Log[1 + x + x^2]/2)/6`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])`

```
rule 754 Int[((a_) + (b_.)*(x_)^(n_))^(n_)-1, x_Symbol] := Module[{r = Numerator[Rt[-a/b, n]], s = Denominator[Rt[-a/b, n]], k, u}, Simp[u = Int[(r - s*Cos[(2*k*Pi)/n]*x)/(r^2 - 2*r*s*Cos[(2*k*Pi)/n]*x + s^2*x^2), x] + Int[(r + s*Cos[(2*k*Pi)/n]*x)/(r^2 + 2*r*s*Cos[(2*k*Pi)/n]*x + s^2*x^2), x]; 2*(r^2/(a*n)) Int[1/(r^2 - s^2*x^2), x] + 2*(r/(a*n)) Sum[u, {k, 1, (n - 2)/4}], x] /; FreeQ[{a, b}, x] && IGtQ[(n - 2)/4, 0] && NegQ[a/b]
```

```
rule 1083 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(n_)-1, x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]
```

```
rule 1103 Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

```
rule 1142 Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]
```

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.40

method	result
default	$\frac{\ln(-1+x)}{6} - \frac{\ln(x^2+x+1)}{12} - \frac{\arctan\left(\frac{(1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{6} - \frac{\ln(1+x)}{6} + \frac{\ln(x^2-x+1)}{12} - \frac{\sqrt{3} \arctan\left(\frac{(2x-1)\sqrt{3}}{3}\right)}{6}$
risch	$-\frac{\ln(4x^2+4x+4)}{12} - \frac{\arctan\left(\frac{(1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{6} - \frac{\ln(1+x)}{6} + \frac{\ln(-1+x)}{6} + \frac{\ln(x^2-x+1)}{12} - \frac{\sqrt{3} \arctan\left(\frac{2(x-\frac{1}{2})\sqrt{3}}{3}\right)}{6}$
meijerg	$\frac{x \left(\ln\left(1-(x^6)^{\frac{1}{6}}\right) - \ln\left(1+(x^6)^{\frac{1}{6}}\right) + \frac{\ln\left(1-(x^6)^{\frac{1}{6}}+(x^6)^{\frac{1}{3}}\right)}{2} - \sqrt{3} \arctan\left(\frac{\sqrt{3}(x^6)^{\frac{1}{6}}}{2-(x^6)^{\frac{1}{6}}}\right) - \frac{\ln\left(1+(x^6)^{\frac{1}{6}}+(x^6)^{\frac{1}{3}}\right)}{2} - \sqrt{3} \arctan\left(\frac{\sqrt{3}(x^6)^{\frac{1}{6}}}{2+(x^6)^{\frac{1}{6}}}\right) \right)}{6(x^6)^{\frac{1}{6}}}$

```
input int(1/(x^6-1),x,method=_RETURNVERBOSE)
```

output $1/6*\ln(-1+x)-1/12*\ln(x^2+x+1)-1/6*\arctan(1/3*(1+2*x)*3^{(1/2)})*3^{(1/2)}-1/6*\ln(1+x)+1/12*\ln(x^2-x+1)-1/6*3^{(1/2)}*\arctan(1/3*(2*x-1)*3^{(1/2)})$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.38

$$\int \frac{1}{-1+x^6} dx = -\frac{1}{6} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x+1)\right) - \frac{1}{6} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x-1)\right) - \frac{1}{12} \log(x^2+x+1) + \frac{1}{12} \log(x^2-x+1) - \frac{1}{6} \log(x+1) + \frac{1}{6} \log(x-1)$$

input `integrate(1/(x^6-1),x, algorithm="fricas")`

output $-1/6*\sqrt{3}*\arctan(1/3*\sqrt{3}*(2*x + 1)) - 1/6*\sqrt{3}*\arctan(1/3*\sqrt{3}*(2*x - 1)) - 1/12*\log(x^2 + x + 1) + 1/12*\log(x^2 - x + 1) - 1/6*\log(x + 1) + 1/6*\log(x - 1)$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 83 vs. $2(36) = 72$.

Time = 0.11 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.77

$$\int \frac{1}{-1+x^6} dx = \frac{\log(x-1)}{6} - \frac{\log(x+1)}{6} + \frac{\log(x^2-x+1)}{12} - \frac{\log(x^2+x+1)}{12} - \frac{\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}x - \sqrt{3}}{3}\right)}{6} - \frac{\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}x + \sqrt{3}}{3}\right)}{6}$$

input `integrate(1/(x**6-1),x)`

output $\log(x - 1)/6 - \log(x + 1)/6 + \log(x**2 - x + 1)/12 - \log(x**2 + x + 1)/12 - \sqrt{3}*\operatorname{atan}(2*\sqrt{3}*x/3 - \sqrt{3}/3)/6 - \sqrt{3}*\operatorname{atan}(2*\sqrt{3}*x/3 + \sqrt{3}/3)/6$

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.38

$$\int \frac{1}{-1+x^6} dx = -\frac{1}{6} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x+1)\right) - \frac{1}{6} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x-1)\right) \\ - \frac{1}{12} \log(x^2+x+1) + \frac{1}{12} \log(x^2-x+1) - \frac{1}{6} \log(x+1) + \frac{1}{6} \log(x-1)$$

input `integrate(1/(x^6-1),x, algorithm="maxima")`output `-1/6*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) - 1/6*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - 1)) - 1/12*log(x^2 + x + 1) + 1/12*log(x^2 - x + 1) - 1/6*log(x + 1) + 1/6*log(x - 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.43

$$\int \frac{1}{-1+x^6} dx = -\frac{1}{6} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x+1)\right) \\ - \frac{1}{6} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x-1)\right) - \frac{1}{12} \log(x^2+x+1) \\ + \frac{1}{12} \log(x^2-x+1) - \frac{1}{6} \log(|x+1|) + \frac{1}{6} \log(|x-1|)$$

input `integrate(1/(x^6-1),x, algorithm="giac")`output `-1/6*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) - 1/6*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - 1)) - 1/12*log(x^2 + x + 1) + 1/12*log(x^2 - x + 1) - 1/6*log(abs(x + 1)) + 1/6*log(abs(x - 1))`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.87

$$\int \frac{1}{-1+x^6} dx = -\frac{\operatorname{atanh}(x)}{3} - \operatorname{atan}\left(\frac{x \operatorname{li}}{1+\sqrt{3} \operatorname{li}} + \frac{\sqrt{3}x}{1+\sqrt{3} \operatorname{li}}\right) \left(\frac{\sqrt{3}}{6} + \frac{1}{6}i\right) \\ - \operatorname{atan}\left(\frac{x \operatorname{li}}{-1+\sqrt{3} \operatorname{li}} - \frac{\sqrt{3}x}{-1+\sqrt{3} \operatorname{li}}\right) \left(\frac{\sqrt{3}}{6} - \frac{1}{6}i\right)$$

input `int(1/(x^6 - 1),x)`output `- atanh(x)/3 - atan((x*1i)/(3^(1/2)*1i + 1) + (3^(1/2)*x)/(3^(1/2)*1i + 1))*(3^(1/2)/6 + 1i/6) - atan((x*1i)/(3^(1/2)*1i - 1) - (3^(1/2)*x)/(3^(1/2)*1i - 1))*(3^(1/2)/6 - 1i/6)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.34

$$\int \frac{1}{-1+x^6} dx = -\frac{\sqrt{3} \operatorname{atan}\left(\frac{2x-1}{\sqrt{3}}\right)}{6} - \frac{\sqrt{3} \operatorname{atan}\left(\frac{2x+1}{\sqrt{3}}\right)}{6} + \frac{\log(x^2 - x + 1)}{12} \\ - \frac{\log(x^2 + x + 1)}{12} + \frac{\log(x - 1)}{6} - \frac{\log(x + 1)}{6}$$

input `int(1/(x^6-1),x)`output `(- 2*sqrt(3)*atan((2*x - 1)/sqrt(3)) - 2*sqrt(3)*atan((2*x + 1)/sqrt(3)) + log(x**2 - x + 1) - log(x**2 + x + 1) + 2*log(x - 1) - 2*log(x + 1))/12`

3.51 $\int \frac{1}{A^4 - A^2 B^2 + (-A^2 + B^2)x^2} dx$

Optimal result	335
Mathematica [A] (verified)	335
Rubi [A] (verified)	336
Maple [A] (verified)	336
Fricas [A] (verification not implemented)	337
Sympy [B] (verification not implemented)	337
Maxima [A] (verification not implemented)	338
Giac [A] (verification not implemented)	338
Mupad [B] (verification not implemented)	338
Reduce [B] (verification not implemented)	339

Optimal result

Integrand size = 27, antiderivative size = 21

$$\int \frac{1}{A^4 - A^2 B^2 + (-A^2 + B^2)x^2} dx = \frac{\operatorname{arctanh}\left(\frac{x}{A}\right)}{A(A^2 - B^2)}$$

output `arctanh(x/A)/A/(A^2-B^2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{1}{A^4 - A^2 B^2 + (-A^2 + B^2)x^2} dx = \frac{\operatorname{arctanh}\left(\frac{x}{A}\right)}{A(A^2 - B^2)}$$

input `Integrate[(A^4 - A^2*B^2 + (-A^2 + B^2)*x^2)^(-1),x]`

output `ArcTanh[x/A]/(A*(A^2 - B^2))`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.037$, Rules used = {221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{A^4 + x^2(B^2 - A^2) - A^2B^2} dx$$

$$\downarrow \text{221}$$

$$\frac{\operatorname{arctanh}\left(\frac{x}{A}\right)}{A(A^2 - B^2)}$$

input `Int[(A^4 - A^2*B^2 + (-A^2 + B^2)*x^2)^(-1), x]`

output `ArcTanh[x/A]/(A*(A^2 - B^2))`

Defintions of rubi rules used

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.43

method	result	size
parallelrisch	$-\frac{\ln(A-x)+\ln(A+x)}{2A(A^2-B^2)}$	30
default	$\frac{\frac{\ln(A+x)}{2A} - \frac{\ln(A-x)}{2A}}{A^2-B^2}$	34
norman	$-\frac{\ln(A-x)}{2A(A^2-B^2)} + \frac{\ln(A+x)}{2A(A^2-B^2)}$	44
risch	$-\frac{\ln(A-x)}{2A(A^2-B^2)} + \frac{\ln(A+x)}{2A(A^2-B^2)}$	44

input `int(1/(A^4-A^2*B^2+(-A^2+B^2)*x^2),x,method=_RETURNVERBOSE)`

output `1/2*(-ln(A-x)+ln(A+x))/A/(A^2-B^2)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.29

$$\int \frac{1}{A^4 - A^2B^2 + (-A^2 + B^2)x^2} dx = \frac{\log(A+x) - \log(-A+x)}{2(A^3 - AB^2)}$$

input `integrate(1/(A^4-A^2*B^2+(-A^2+B^2)*x^2),x, algorithm="fricas")`

output `1/2*(log(A + x) - log(-A + x))/(A^3 - A*B^2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 70 vs. 2(12) = 24.

Time = 0.14 (sec) , antiderivative size = 70, normalized size of antiderivative = 3.33

$$\int \frac{1}{A^4 - A^2B^2 + (-A^2 + B^2)x^2} dx = -\frac{\log\left(-\frac{A^3}{(A-B)(A+B)} + \frac{AB^2}{(A-B)(A+B)} + x\right)}{2A(A-B)(A+B)} + \frac{\log\left(\frac{A^3}{(A-B)(A+B)} - \frac{AB^2}{(A-B)(A+B)} + x\right)}{2A(A-B)(A+B)}$$

input `integrate(1/(A**4-A**2*B**2+(-A**2+B**2)*x**2),x)`

output `-log(-A**3/((A - B)*(A + B)) + A*B**2/((A - B)*(A + B)) + x)/(2*A*(A - B)*(A + B)) + log(A**3/((A - B)*(A + B)) - A*B**2/((A - B)*(A + B)) + x)/(2*A*(A - B)*(A + B))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.86

$$\int \frac{1}{A^4 - A^2 B^2 + (-A^2 + B^2) x^2} dx = \frac{\log(A + x)}{2(A^3 - AB^2)} - \frac{\log(-A + x)}{2(A^3 - AB^2)}$$

input `integrate(1/(A^4-A^2*B^2+(-A^2+B^2)*x^2),x, algorithm="maxima")`output `1/2*log(A + x)/(A^3 - A*B^2) - 1/2*log(-A + x)/(A^3 - A*B^2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.95

$$\int \frac{1}{A^4 - A^2 B^2 + (-A^2 + B^2) x^2} dx = \frac{\log(|A + x|)}{2(A^3 - AB^2)} - \frac{\log(|-A + x|)}{2(A^3 - AB^2)}$$

input `integrate(1/(A^4-A^2*B^2+(-A^2+B^2)*x^2),x, algorithm="giac")`output `1/2*log(abs(A + x))/(A^3 - A*B^2) - 1/2*log(abs(-A + x))/(A^3 - A*B^2)`**Mupad [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{1}{A^4 - A^2 B^2 + (-A^2 + B^2) x^2} dx = -\frac{\operatorname{atanh}\left(\frac{x}{A}\right)}{A B^2 - A^3}$$

input `int(-1/(x^2*(A^2 - B^2) - A^4 + A^2*B^2),x)`output `-atanh(x/A)/(A*B^2 - A^3)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.57

$$\int \frac{1}{A^4 - A^2 B^2 + (-A^2 + B^2) x^2} dx = \frac{\log(-a - x) - \log(a - x)}{2a(a^2 - b^2)}$$

input `int(1/(A^4-A^2*B^2+(-A^2+B^2)*x^2),x)`

output `(log(-a-x)-log(a-x))/(2*a*(a**2-b**2))`

3.52 $\int x \log(x) dx$

Optimal result	340
Mathematica [A] (verified)	340
Rubi [A] (verified)	341
Maple [A] (verified)	342
Fricas [A] (verification not implemented)	342
Sympy [A] (verification not implemented)	343
Maxima [A] (verification not implemented)	343
Giac [A] (verification not implemented)	343
Mupad [B] (verification not implemented)	344
Reduce [B] (verification not implemented)	344

Optimal result

Integrand size = 4, antiderivative size = 17

$$\int x \log(x) dx = -\frac{x^2}{4} + \frac{1}{2}x^2 \log(x)$$

output

```
-1/4*x^2+1/2*x^2*ln(x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int x \log(x) dx = -\frac{x^2}{4} + \frac{1}{2}x^2 \log(x)$$

input

```
Integrate[x*Log[x],x]
```

output

```
-1/4*x^2 + (x^2*Log[x])/2
```

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2741}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \log(x) dx$$

↓ 2741

$$\frac{1}{2}x^2 \log(x) - \frac{x^2}{4}$$

input `Int [x*Log [x] , x]`

output `-1/4*x^2 + (x^2*Log [x])/2`

Defintions of rubi rules used

rule 2741 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((d_.)*(x_)^(m_.), x_Symbol] :>
Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])/(d*(m + 1))), x] - Simp[b*n*((d*x)^(
m + 1)/(d*(m + 1)^2)), x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2}$	14
norman	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2}$	14
risch	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2}$	14
parallelrisch	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2}$	14
parts	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2}$	14
orering	$\frac{3x^2 \ln(x)}{4} - \frac{x^2(1+\ln(x))}{4}$	18

input `int(x*ln(x),x,method=_RETURNVERBOSE)`output `-1/4*x^2+1/2*x^2*ln(x)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int x \log(x) dx = \frac{1}{2} x^2 \log(x) - \frac{1}{4} x^2$$

input `integrate(x*log(x),x, algorithm="fricas")`output `1/2*x^2*log(x) - 1/4*x^2`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

$$\int x \log(x) dx = \frac{x^2 \log(x)}{2} - \frac{x^2}{4}$$

input `integrate(x*ln(x),x)`

output `x**2*log(x)/2 - x**2/4`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int x \log(x) dx = \frac{1}{2} x^2 \log(x) - \frac{1}{4} x^2$$

input `integrate(x*log(x),x, algorithm="maxima")`

output `1/2*x^2*log(x) - 1/4*x^2`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int x \log(x) dx = \frac{1}{2} x^2 \log(x) - \frac{1}{4} x^2$$

input `integrate(x*log(x),x, algorithm="giac")`

output `1/2*x^2*log(x) - 1/4*x^2`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.53

$$\int x \log(x) dx = \frac{x^2 (\ln(x) - \frac{1}{2})}{2}$$

input `int(x*log(x),x)`

output `(x^2*(log(x) - 1/2))/2`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int x \log(x) dx = \frac{x^2(2 \log(x) - 1)}{4}$$

input `int(x*log(x),x)`

output `(x**2*(2*log(x) - 1))/4`

3.53 $\int x^2 \arcsin(x) dx$

Optimal result	345
Mathematica [A] (verified)	345
Rubi [A] (verified)	346
Maple [A] (verified)	347
Fricas [A] (verification not implemented)	348
Sympy [A] (verification not implemented)	348
Maxima [A] (verification not implemented)	348
Giac [A] (verification not implemented)	349
Mupad [B] (verification not implemented)	349
Reduce [B] (verification not implemented)	349

Optimal result

Integrand size = 6, antiderivative size = 40

$$\int x^2 \arcsin(x) dx = \frac{\sqrt{1-x^2}}{3} - \frac{1}{9}(1-x^2)^{3/2} + \frac{1}{3}x^3 \arcsin(x)$$

output

```
-1/9*(-x^2+1)^(3/2)+1/3*x^3*arcsin(x)+1/3*(-x^2+1)^(1/2)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.72

$$\int x^2 \arcsin(x) dx = \frac{1}{9} \left(\sqrt{1-x^2} (2+x^2) + 3x^3 \arcsin(x) \right)$$

input

```
Integrate[x^2*ArcSin[x],x]
```

output

```
(Sqrt[1 - x^2]*(2 + x^2) + 3*x^3*ArcSin[x])/9
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.08, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {5138, 243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \arcsin(x) dx \\
 & \quad \downarrow \text{5138} \\
 & \frac{1}{3} x^3 \arcsin(x) - \frac{1}{3} \int \frac{x^3}{\sqrt{1-x^2}} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{3} x^3 \arcsin(x) - \frac{1}{6} \int \frac{x^2}{\sqrt{1-x^2}} dx^2 \\
 & \quad \downarrow \text{53} \\
 & \frac{1}{3} x^3 \arcsin(x) - \frac{1}{6} \int \left(\frac{1}{\sqrt{1-x^2}} - \sqrt{1-x^2} \right) dx^2 \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{3} x^3 \arcsin(x) + \frac{1}{6} \left(2\sqrt{1-x^2} - \frac{2}{3}(1-x^2)^{3/2} \right)
 \end{aligned}$$

input `Int [x^2*ArcSin[x] , x]`

output `(2*sqrt [1 - x^2] - (2*(1 - x^2)^(3/2))/3)/6 + (x^3*ArcSin[x])/3`

Definitions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5138 `Int[((a_.) + ArcSin[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcSin[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcSin[c*x])^(n - 1)/Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.85

method	result	size
default	$\frac{x^3 \arcsin(x)}{3} + \frac{\sqrt{-x^2+1} x^2}{9} + \frac{2\sqrt{-x^2+1}}{9}$	34
parts	$\frac{x^3 \arcsin(x)}{3} + \frac{\sqrt{-x^2+1} x^2}{9} + \frac{2\sqrt{-x^2+1}}{9}$	34
orering	$\frac{(5x^4+2x^2-4) \arcsin(x)}{9x} - \frac{(x^2+2)(-1+x)(1+x) \left(\frac{x^2}{\sqrt{-x^2+1}} + 2 \arcsin(x)x \right)}{9x^2}$	56

input `int(arcsin(x)*x^2,x,method=_RETURNVERBOSE)`

output `1/3*x^3*arcsin(x)+1/9*(-x^2+1)^(1/2)*x^2+2/9*(-x^2+1)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.60

$$\int x^2 \arcsin(x) dx = \frac{1}{3} x^3 \arcsin(x) + \frac{1}{9} (x^2 + 2) \sqrt{-x^2 + 1}$$

input `integrate(x^2*arcsin(x),x, algorithm="fricas")`output `1/3*x^3*arcsin(x) + 1/9*(x^2 + 2)*sqrt(-x^2 + 1)`**Sympy [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.80

$$\int x^2 \arcsin(x) dx = \frac{x^3 \arcsin(x)}{3} + \frac{x^2 \sqrt{1-x^2}}{9} + \frac{2\sqrt{1-x^2}}{9}$$

input `integrate(x**2*asin(x),x)`output `x**3*asin(x)/3 + x**2*sqrt(1 - x**2)/9 + 2*sqrt(1 - x**2)/9`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.82

$$\int x^2 \arcsin(x) dx = \frac{1}{3} x^3 \arcsin(x) + \frac{1}{9} \sqrt{-x^2 + 1} x^2 + \frac{2}{9} \sqrt{-x^2 + 1}$$

input `integrate(x^2*arcsin(x),x, algorithm="maxima")`output `1/3*x^3*arcsin(x) + 1/9*sqrt(-x^2 + 1)*x^2 + 2/9*sqrt(-x^2 + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.95

$$\int x^2 \arcsin(x) dx = \frac{1}{3} (x^2 - 1)x \arcsin(x) + \frac{1}{3} x \arcsin(x) - \frac{1}{9} (-x^2 + 1)^{\frac{3}{2}} + \frac{1}{3} \sqrt{-x^2 + 1}$$

input `integrate(x^2*arcsin(x),x, algorithm="giac")`

output `1/3*(x^2 - 1)*x*arcsin(x) + 1/3*x*arcsin(x) - 1/9*(-x^2 + 1)^(3/2) + 1/3*sqrt(-x^2 + 1)`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.60

$$\int x^2 \arcsin(x) dx = \frac{x^3 \arcsin(x)}{3} + \frac{\sqrt{1-x^2}(x^2+2)}{9}$$

input `int(x^2*asin(x),x)`

output `(x^3*asin(x))/3 + ((1 - x^2)^(1/2)*(x^2 + 2))/9`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.78

$$\int x^2 \arcsin(x) dx = \frac{\arcsin(x) x^3}{3} + \frac{\sqrt{-x^2 + 1} x^2}{9} + \frac{2\sqrt{-x^2 + 1}}{9}$$

input `int(x^2*asin(x),x)`

output `(3*asin(x)*x**3 + sqrt(-x**2 + 1)*x**2 + 2*sqrt(-x**2 + 1))/9`

3.54 $\int \frac{1}{1+2x+x^2} dx$

Optimal result	350
Mathematica [A] (verified)	350
Rubi [A] (verified)	351
Maple [A] (verified)	352
Fricas [A] (verification not implemented)	352
Sympy [A] (verification not implemented)	353
Maxima [A] (verification not implemented)	353
Giac [A] (verification not implemented)	353
Mupad [B] (verification not implemented)	354
Reduce [B] (verification not implemented)	354

Optimal result

Integrand size = 10, antiderivative size = 7

$$\int \frac{1}{1+2x+x^2} dx = -\frac{1}{1+x}$$

output `-1/(1+x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+2x+x^2} dx = -\frac{1}{1+x}$$

input `Integrate[(1 + 2*x + x^2)^(-1),x]`

output `-(1 + x)^(-1)`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {1077, 17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 + 2x + 1} dx$$

↓ 1077

$$\int \frac{1}{(x + 1)^2} dx$$

↓ 17

$$-\frac{1}{x + 1}$$

input `Int[(1 + 2*x + x^2)^(-1),x]`

output `-(1 + x)^(-1)`

Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1))], x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

rule 1077 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Simp[1/c^p Int[(b/2 + c*x)^(2*p), x], x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.14

method	result	size
gospers	$-\frac{1}{1+x}$	8
default	$-\frac{1}{1+x}$	8
norman	$-\frac{1}{1+x}$	8
meijerg	$\frac{x}{1+x}$	8
risch	$-\frac{1}{1+x}$	8
parallelrisch	$-\frac{1}{1+x}$	8
orering	$\frac{-1-x}{x^2+2x+1}$	17

input `int(1/(x^2+2*x+1),x,method=_RETURNVERBOSE)`

output `-1/(1+x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+2x+x^2} dx = -\frac{1}{x+1}$$

input `integrate(1/(x^2+2*x+1),x, algorithm="fricas")`

output `-1/(x + 1)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.71

$$\int \frac{1}{1 + 2x + x^2} dx = -\frac{1}{x + 1}$$

input `integrate(1/(x**2+2*x+1),x)`

output `-1/(x + 1)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + 2x + x^2} dx = -\frac{1}{x + 1}$$

input `integrate(1/(x^2+2*x+1),x, algorithm="maxima")`

output `-1/(x + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + 2x + x^2} dx = -\frac{1}{x + 1}$$

input `integrate(1/(x^2+2*x+1),x, algorithm="giac")`

output `-1/(x + 1)`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + 2x + x^2} dx = -\frac{1}{x + 1}$$

input `int(1/(2*x + x^2 + 1), x)`

output `-1/(x + 1)`

Reduce [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + 2x + x^2} dx = \frac{x}{x + 1}$$

input `int(1/(x^2+2*x+1), x)`

output `x/(x + 1)`

3.55 $\int \frac{\log(x)}{(1+\log(x))^2} dx$

Optimal result	355
Mathematica [A] (verified)	355
Rubi [A] (verified)	356
Maple [A] (verified)	357
Fricas [A] (verification not implemented)	357
Sympy [A] (verification not implemented)	357
Maxima [A] (verification not implemented)	358
Giac [A] (verification not implemented)	358
Mupad [B] (verification not implemented)	358
Reduce [B] (verification not implemented)	359

Optimal result

Integrand size = 9, antiderivative size = 8

$$\int \frac{\log(x)}{(1 + \log(x))^2} dx = \frac{x}{1 + \log(x)}$$

output `x/(1+ln(x))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{\log(x)}{(1 + \log(x))^2} dx = \frac{x}{1 + \log(x)}$$

input `Integrate[Log[x]/(1 + Log[x])^2,x]`

output `x/(1 + Log[x])`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {2807, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\log(x)}{(\log(x) + 1)^2} dx$$

↓ 2807

$$\int \left(\frac{1}{\log(x) + 1} - \frac{1}{(\log(x) + 1)^2} \right) dx$$

↓ 2009

$$\frac{x}{\log(x) + 1}$$

input

```
Int[Log[x]/(1 + Log[x])^2,x]
```

output

```
x/(1 + Log[x])
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2807

```
Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^p_.*(Log[(c_.)*(x_)^(n_.)]*(e_.) + (d_.)^(q_.), x_Symbol] := Int[ExpandIntegrand[(a + b*Log[c*x^n])^p*(d + e*Log[c*x^n])^q, x], x] /; FreeQ[{a, b, c, d, e, n}, x] && IntegerQ[p] && IntegerQ[q]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.12

method	result	size
default	$\frac{x}{1+\ln(x)}$	9
norman	$\frac{x}{1+\ln(x)}$	9
risch	$\frac{x}{1+\ln(x)}$	9
parallelrisch	$\frac{x}{1+\ln(x)}$	9

input `int(ln(x)/(1+ln(x))^2,x,method=_RETURNVERBOSE)`

output `x/(1+ln(x))`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{\log(x)}{(1 + \log(x))^2} dx = \frac{x}{\log(x) + 1}$$

input `integrate(log(x)/(1+log(x))^2,x, algorithm="fricas")`

output `x/(log(x) + 1)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int \frac{\log(x)}{(1 + \log(x))^2} dx = \frac{x}{\log(x) + 1}$$

input `integrate(ln(x)/(1+ln(x))**2,x)`

output $x/(\log(x) + 1)$

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{\log(x)}{(1 + \log(x))^2} dx = \frac{x}{\log(x) + 1}$$

input `integrate(log(x)/(1+log(x))^2,x, algorithm="maxima")`

output $x/(\log(x) + 1)$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{\log(x)}{(1 + \log(x))^2} dx = \frac{x}{\log(x) + 1}$$

input `integrate(log(x)/(1+log(x))^2,x, algorithm="giac")`

output $x/(\log(x) + 1)$

Mupad [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{\log(x)}{(1 + \log(x))^2} dx = \frac{x}{\ln(x) + 1}$$

input `int(log(x)/(log(x) + 1)^2,x)`

output $x/(\log(x) + 1)$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{\log(x)}{(1 + \log(x))^2} dx = \frac{x}{\log(x) + 1}$$

input `int(log(x)/(1+log(x))^2,x)`

output `x/(log(x) + 1)`

$$3.56 \quad \int \frac{1}{x(1+\log^2(x))} dx$$

Optimal result	360
Mathematica [A] (verified)	360
Rubi [A] (verified)	361
Maple [A] (verified)	362
Fricas [A] (verification not implemented)	362
Sympy [B] (verification not implemented)	362
Maxima [A] (verification not implemented)	363
Giac [A] (verification not implemented)	363
Mupad [B] (verification not implemented)	363
Reduce [B] (verification not implemented)	364

Optimal result

Integrand size = 12, antiderivative size = 3

$$\int \frac{1}{x(1+\log^2(x))} dx = \arctan(\log(x))$$

output `arctan(ln(x))`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1+\log^2(x))} dx = \arctan(\log(x))$$

input `Integrate[1/(x*(1 + Log[x]^2)),x]`

output `ArcTan[Log[x]]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3039, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x (\log^2(x) + 1)} dx$$

↓ 3039

$$\int \frac{1}{\log^2(x) + 1} d\log(x)$$

↓ 216

$$\arctan(\log(x))$$

input `Int[1/(x*(1 + Log[x]^2)),x]`

output `ArcTan[Log[x]]`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3039 `Int[u_, x_Symbol] := With[{lst = FunctionOfLog[Cancel[x*u], x]}, Simp[1/lst[[3]] Subst[Int[lst[[1]], x], x, Log[lst[[2]]]], x] /; !FalseQ[lst] /; NonsumQ[u]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
derivativedivides	$\arctan(\ln(x))$	4
default	$\arctan(\ln(x))$	4
risch	$\frac{i \ln(\ln(x)+i)}{2} - \frac{i \ln(\ln(x)-i)}{2}$	20
parallelrisc	$\frac{i \ln(\ln(x)+i)}{2} - \frac{i \ln(\ln(x)-i)}{2}$	20

input `int(1/x/(1+ln(x)^2),x,method=_RETURNVERBOSE)`

output `arctan(ln(x))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1+\log^2(x))} dx = \arctan(\log(x))$$

input `integrate(1/x/(1+log(x)^2),x, algorithm="fricas")`

output `arctan(log(x))`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(3) = 6$.

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 5.00

$$\int \frac{1}{x(1+\log^2(x))} dx = \text{RootSum}(4z^2 + 1, (i \mapsto i \log(2i + \log(x))))$$

input `integrate(1/x/(1+ln(x)**2),x)`

output `RootSum(4*_z**2 + 1, Lambda(_i, _i*log(2*_i + log(x))))`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1+\log^2(x))} dx = \arctan(\log(x))$$

input `integrate(1/x/(1+log(x)^2),x, algorithm="maxima")`

output `arctan(log(x))`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1+\log^2(x))} dx = \arctan(\log(x))$$

input `integrate(1/x/(1+log(x)^2),x, algorithm="giac")`

output `arctan(log(x))`

Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1+\log^2(x))} dx = \operatorname{atan}(\ln(x))$$

input `int(1/(x*(log(x)^2 + 1)),x)`

output `atan(log(x))`

Reduce [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1+\log^2(x))} dx = \text{atan}(\log(x))$$

input `int(1/x/(1+log(x)^2),x)`

output `atan(log(x))`

3.57 $\int \frac{1}{\log(x)} dx$

Optimal result	365
Mathematica [A] (verified)	365
Rubi [A] (verified)	366
Maple [B] (verified)	366
Fricas [A] (verification not implemented)	367
Sympy [A] (verification not implemented)	367
Maxima [A] (verification not implemented)	367
Giac [A] (verification not implemented)	368
Mupad [B] (verification not implemented)	368
Reduce [B] (verification not implemented)	368

Optimal result

Integrand size = 4, antiderivative size = 2

$$\int \frac{1}{\log(x)} dx = \text{LogIntegral}(x)$$

output `Li(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{\log(x)} dx = \text{LogIntegral}(x)$$

input `Integrate[Log[x]^(-1),x]`

output `LogIntegral[x]`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2735}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\log(x)} dx$$

↓ 2735

$$\text{LogIntegral}(x)$$

input `Int [Log[x]^(-1) , x]`

output `LogIntegral [x]`

Defintions of rubi rules used

rule 2735 `Int [Log[(c_.)*(x_)]^(-1), x_Symbol] :> Simp[LogIntegral[c*x]/c, x] /; FreeQ [c, x]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 8 vs. $2(2) = 4$.

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 4.50

method	result	size
default	$-\text{expIntegral}_1(-\ln(x))$	9
risch	$-\text{expIntegral}_1(-\ln(x))$	9

input `int(1/ln(x),x,method=_RETURNVERBOSE)`

output `-Ei(1,-ln(x))`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{\log(x)} dx = \log_integral(x)$$

input `integrate(1/log(x),x, algorithm="fricas")`

output `log_integral(x)`

Sympy [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{\log(x)} dx = \text{li}(x)$$

input `integrate(1/ln(x),x)`

output `li(x)`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

$$\int \frac{1}{\log(x)} dx = \text{Ei}(\log(x))$$

input `integrate(1/log(x),x, algorithm="maxima")`

output `Ei(log(x))`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

$$\int \frac{1}{\log(x)} dx = \text{Ei}(\log(x))$$

input `integrate(1/log(x),x, algorithm="giac")`

output `Ei(log(x))`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{\log(x)} dx = \text{logint}(x)$$

input `int(1/log(x),x)`

output `logint(x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

$$\int \frac{1}{\log(x)} dx = \text{ei}(\log(x))$$

input `int(1/log(x),x)`

output `ei(log(x))`

3.58 $\int x(\cos(x) + \sin(x)) dx$

Optimal result	369
Mathematica [A] (verified)	369
Rubi [A] (verified)	370
Maple [A] (verified)	371
Fricas [A] (verification not implemented)	371
Sympy [A] (verification not implemented)	372
Maxima [A] (verification not implemented)	372
Giac [A] (verification not implemented)	372
Mupad [B] (verification not implemented)	373
Reduce [B] (verification not implemented)	373

Optimal result

Integrand size = 7, antiderivative size = 14

$$\int x(\cos(x) + \sin(x)) dx = \cos(x) - x \cos(x) + \sin(x) + x \sin(x)$$

output

```
cos(x)-x*cos(x)+sin(x)+x*sin(x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int x(\cos(x) + \sin(x)) dx = \cos(x) - x \cos(x) + \sin(x) + x \sin(x)$$

input

```
Integrate[x*(Cos[x] + Sin[x]),x]
```

output

```
Cos[x] - x*Cos[x] + Sin[x] + x*Sin[x]
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2010, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(\sin(x) + \cos(x)) dx$$

$$\downarrow \text{2010}$$

$$\int (x \sin(x) + x \cos(x)) dx$$

$$\downarrow \text{2009}$$

$$x \sin(x) + \sin(x) - x \cos(x) + \cos(x)$$

input `Int[x*(Cos[x] + Sin[x]),x]`

output `Cos[x] - x*Cos[x] + Sin[x] + x*Sin[x]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2010 `Int[(u_)*((c_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_) + (b_.)*(v_)] /; FreeQ[{a, b}, x] && InverseFunctionQ[v]`

Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

method	result	size
default	$\cos(x) - x \cos(x) + \sin(x) + x \sin(x)$	15
parts	$\cos(x) - x \cos(x) + \sin(x) + x \sin(x)$	15
risch	$(1 - x) \cos(x) + (1 + x) \sin(x)$	16
orering	$\cos(x) + \sin(x) - x(\cos(x) - \sin(x))$	16
parallelrisc	$(1 - x) \cos(x) + 1 + (1 + x) \sin(x)$	17
norman	$\frac{x \tan(\frac{x}{2})^2 - x + 2x \tan(\frac{x}{2}) + 2 \tan(\frac{x}{2}) + 2}{1 + \tan(\frac{x}{2})^2}$	38
meijerg	$2\sqrt{\pi} \left(-\frac{1}{2\sqrt{\pi}} + \frac{\cos(x)}{2\sqrt{\pi}} + \frac{x \sin(x)}{2\sqrt{\pi}} \right) + 2\sqrt{\pi} \left(-\frac{x \cos(x)}{2\sqrt{\pi}} + \frac{\sin(x)}{2\sqrt{\pi}} \right)$	49

input `int(x*(cos(x)+sin(x)),x,method=_RETURNVERBOSE)`output `cos(x)-x*cos(x)+sin(x)+x*sin(x)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int x(\cos(x) + \sin(x)) dx = -(x - 1) \cos(x) + (x + 1) \sin(x)$$

input `integrate(x*(cos(x)+sin(x)),x, algorithm="fricas")`output `-(x - 1)*cos(x) + (x + 1)*sin(x)`

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

$$\int x(\cos(x) + \sin(x)) dx = x \sin(x) - x \cos(x) + \sin(x) + \cos(x)$$

input `integrate(x*(cos(x)+sin(x)),x)`

output `x*sin(x) - x*cos(x) + sin(x) + cos(x)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int x(\cos(x) + \sin(x)) dx = -x \cos(x) + x \sin(x) + \cos(x) + \sin(x)$$

input `integrate(x*(cos(x)+sin(x)),x, algorithm="maxima")`

output `-x*cos(x) + x*sin(x) + cos(x) + sin(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int x(\cos(x) + \sin(x)) dx = -x \cos(x) + x \sin(x) + \cos(x) + \sin(x)$$

input `integrate(x*(cos(x)+sin(x)),x, algorithm="giac")`

output `-x*cos(x) + x*sin(x) + cos(x) + sin(x)`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int x(\cos(x) + \sin(x)) dx = \cos(x) + \sin(x) - x \cos(x) + x \sin(x)$$

input `int(x*(cos(x) + sin(x)),x)`

output `cos(x) + sin(x) - x*cos(x) + x*sin(x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int x(\cos(x) + \sin(x)) dx = -\cos(x)x + \cos(x) + \sin(x)x + \sin(x)$$

input `int(x*(cos(x)+sin(x)),x)`

output `- cos(x)*x + cos(x) + sin(x)*x + sin(x)`

3.59 $\int e^{-x}(e^x + x) dx$

Optimal result	374
Mathematica [A] (verified)	374
Rubi [A] (verified)	375
Maple [A] (verified)	376
Fricas [A] (verification not implemented)	376
Sympy [A] (verification not implemented)	377
Maxima [A] (verification not implemented)	377
Giac [A] (verification not implemented)	377
Mupad [B] (verification not implemented)	378
Reduce [B] (verification not implemented)	378

Optimal result

Integrand size = 11, antiderivative size = 17

$$\int e^{-x}(e^x + x) dx = -e^{-x} + x - e^{-x}x$$

output `-1/exp(x)+x-x/exp(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int e^{-x}(e^x + x) dx = e^{-x}(-1 - x) + x$$

input `Integrate[(E^x + x)/E^x,x]`

output `(-1 - x)/E^x + x`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{-x}(x + e^x) dx$$

$$\downarrow 7293$$

$$\int (e^{-x}x + 1) dx$$

$$\downarrow 2009$$

$$-e^{-x}x + x - e^{-x}$$

input `Int[(E^-x + x)/E^x,x]`

output `-E^(-x) + x - x/E^x`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] :> With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

method	result	size
risch	$x + (-1 - x)e^{-x}$	13
norman	$(-1 + e^x x - x)e^{-x}$	15
parallelrisch	$(-1 + e^x x - x)e^{-x}$	15
default	$-e^{-x}x - e^{-x} + x$	16
parts	$-e^{-x}x - e^{-x} + x$	16
orering	$(-1 + x)(e^x + x)e^{-x} + \frac{(x^2+1)((1+e^x)e^{-x} - (e^x+x)e^{-x})}{-1+x}$	45

input `int((exp(x)+x)/exp(x),x,method=_RETURNVERBOSE)`output `x+(-1-x)*exp(-x)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int e^{-x}(e^x + x) dx = (xe^x - x - 1)e^{(-x)}$$

input `integrate((exp(x)+x)/exp(x),x, algorithm="fricas")`output `(x*e^x - x - 1)*e^(-x)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.47

$$\int e^{-x}(e^x + x) dx = x + (-x - 1)e^{-x}$$

input `integrate((exp(x)+x)/exp(x),x)`

output `x + (-x - 1)*exp(-x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int e^{-x}(e^x + x) dx = -(x + 1)e^{(-x)} + x$$

input `integrate((exp(x)+x)/exp(x),x, algorithm="maxima")`

output `-(x + 1)*e^(-x) + x`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int e^{-x}(e^x + x) dx = -(x + 1)e^{(-x)} + x$$

input `integrate((exp(x)+x)/exp(x),x, algorithm="giac")`

output `-(x + 1)*e^(-x) + x`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int e^{-x}(e^x + x) dx = x - e^{-x} - x e^{-x}$$

input `int(exp(-x)*(x + exp(x)),x)`

output `x - exp(-x) - x*exp(-x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

$$\int e^{-x}(e^x + x) dx = \frac{e^x x - x - 1}{e^x}$$

input `int((exp(x)+x)/exp(x),x)`

output `(e**x*x - x - 1)/e**x`

3.60 $\int (1 + e^x)^2 x dx$

Optimal result	379
Mathematica [A] (verified)	379
Rubi [A] (verified)	380
Maple [A] (verified)	381
Fricas [A] (verification not implemented)	381
Sympy [A] (verification not implemented)	382
Maxima [A] (verification not implemented)	382
Giac [A] (verification not implemented)	382
Mupad [B] (verification not implemented)	383
Reduce [B] (verification not implemented)	383

Optimal result

Integrand size = 9, antiderivative size = 38

$$\int (1 + e^x)^2 x dx = -2e^x - \frac{e^{2x}}{4} + 2e^x x + \frac{1}{2}e^{2x} x + \frac{x^2}{2}$$

output

```
-2*exp(x)-1/4*exp(2*x)+2*exp(x)*x+1/2*exp(2*x)*x+1/2*x^2
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.76

$$\int (1 + e^x)^2 x dx = \frac{1}{4}(8e^x(-1 + x) + 2x^2 + e^{2x}(-1 + 2x))$$

input

```
Integrate[(1 + E^x)^2*x,x]
```

output

```
(8*E^x*(-1 + x) + 2*x^2 + E^(2*x)*(-1 + 2*x))/4
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {2614, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e^x + 1)^2 x dx$$

↓ 2614

$$\int (2e^x x + e^{2x} x + x) dx$$

↓ 2009

$$\frac{x^2}{2} + 2e^x x + \frac{1}{2}e^{2x} x - 2e^x - \frac{e^{2x}}{4}$$

input

```
Int[(1 + E^x)^2*x, x]
```

output

```
-2*E^x - E^(2*x)/4 + 2*E^x*x + (E^(2*x)*x)/2 + x^2/2
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2614

```
Int[((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))^(p_)*((c_) + (d_)*(x_))^(m_), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*(F^(g*(e + f*x)))^n)^p, x], x] /; FreeQ[{F, a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]
```

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.66

method	result
risch	$\frac{x^2}{2} + \left(-\frac{1}{4} + \frac{x}{2}\right) e^{2x} + (-2 + 2x) e^x$
default	$-2 e^x - \frac{e^{2x}}{4} + 2 e^x x + \frac{e^{2x} x}{2} + \frac{x^2}{2}$
norman	$-2 e^x - \frac{e^{2x}}{4} + 2 e^x x + \frac{e^{2x} x}{2} + \frac{x^2}{2}$
parallelrisch	$-2 e^x - \frac{e^{2x}}{4} + 2 e^x x + \frac{e^{2x} x}{2} + \frac{x^2}{2}$
parts	$-2 e^x - \frac{e^{2x}}{4} + 2 e^x x + \frac{e^{2x} x}{2} + \frac{x^2}{2}$
orering	$\frac{(2x^3+3x^2-12x-9)(1+e^x)^2}{4x} - \frac{(3x^2-5x-9)(2(1+e^x)x e^x+(1+e^x)^2)}{4x} + \left(-\frac{3}{4} + \frac{x}{4}\right) (2 e^{2x} x + 4(1+e^x) e^x +$

input `int((1+exp(x))^2*x,x,method=_RETURNVERBOSE)`output `1/2*x^2+(-1/4+1/2*x)*exp(x)^2+(-2+2*x)*exp(x)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.63

$$\int (1 + e^x)^2 x dx = \frac{1}{2} x^2 + \frac{1}{4} (2x - 1) e^{(2x)} + 2(x - 1) e^x$$

input `integrate((1+exp(x))^2*x,x,algorithm="fricas")`output `1/2*x^2 + 1/4*(2*x - 1)*e^(2*x) + 2*(x - 1)*e^x`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.68

$$\int (1 + e^x)^2 x dx = \frac{x^2}{2} + \frac{(2x - 1)e^{2x}}{4} + \frac{(8x - 8)e^x}{4}$$

input `integrate((1+exp(x))**2*x,x)`output `x**2/2 + (2*x - 1)*exp(2*x)/4 + (8*x - 8)*exp(x)/4`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.63

$$\int (1 + e^x)^2 x dx = \frac{1}{2} x^2 + \frac{1}{4} (2x - 1)e^{(2x)} + 2(x - 1)e^x$$

input `integrate((1+exp(x))^2*x,x, algorithm="maxima")`output `1/2*x^2 + 1/4*(2*x - 1)*e^(2*x) + 2*(x - 1)*e^x`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.63

$$\int (1 + e^x)^2 x dx = \frac{1}{2} x^2 + \frac{1}{4} (2x - 1)e^{(2x)} + 2(x - 1)e^x$$

input `integrate((1+exp(x))^2*x,x, algorithm="giac")`output `1/2*x^2 + 1/4*(2*x - 1)*e^(2*x) + 2*(x - 1)*e^x`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.74

$$\int (1 + e^x)^2 x dx = \frac{x e^{2x}}{2} - 2 e^x - \frac{e^{2x}}{4} + 2 x e^x + \frac{x^2}{2}$$

input `int(x*(exp(x) + 1)^2,x)`output `(x*exp(2*x))/2 - 2*exp(x) - exp(2*x)/4 + 2*x*exp(x) + x^2/2`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int (1 + e^x)^2 x dx = \frac{e^{2x}x}{2} - \frac{e^{2x}}{4} + 2e^x x - 2e^x + \frac{x^2}{2}$$

input `int((1+exp(x))^2*x,x)`output `(2*e**(2*x)*x - e**(2*x) + 8*e**x*x - 8*e**x + 2*x**2)/4`

3.61 $\int x \cos(x) dx$

Optimal result	384
Mathematica [A] (verified)	384
Rubi [A] (verified)	385
Maple [A] (verified)	386
Fricas [A] (verification not implemented)	387
Sympy [A] (verification not implemented)	387
Maxima [A] (verification not implemented)	387
Giac [A] (verification not implemented)	388
Mupad [B] (verification not implemented)	388
Reduce [B] (verification not implemented)	388

Optimal result

Integrand size = 4, antiderivative size = 7

$$\int x \cos(x) dx = \cos(x) + x \sin(x)$$

output `cos(x)+x*sin(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int x \cos(x) dx = \cos(x) + x \sin(x)$$

input `Integrate[x*Cos[x],x]`

output `Cos[x] + x*Sin[x]`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$, Rules used = {3042, 3777, 25, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \cos(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x \sin\left(x + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{3777} \\
 & \int -\sin(x) dx + x \sin(x) \\
 & \quad \downarrow \text{25} \\
 & x \sin(x) - \int \sin(x) dx \\
 & \quad \downarrow \text{3042} \\
 & x \sin(x) - \int \sin(x) dx \\
 & \quad \downarrow \text{3118} \\
 & x \sin(x) + \cos(x)
 \end{aligned}$$

input

Int [x*Cos [x] , x]

output

Cos [x] + x*Sin [x]

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_)^(m_.))*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.14

method	result	size
default	$\cos(x) + x \sin(x)$	8
risch	$\cos(x) + x \sin(x)$	8
parts	$\cos(x) + x \sin(x)$	8
orering	$\cos(x) + x \sin(x)$	8
parallelrisc	$1 + \cos(x) + x \sin(x)$	9
norman	$\frac{2x \tan(\frac{x}{2}) + 2}{1 + \tan(\frac{x}{2})^2}$	21
meijerg	$2\sqrt{\pi} \left(-\frac{1}{2\sqrt{\pi}} + \frac{\cos(x)}{2\sqrt{\pi}} + \frac{x \sin(x)}{2\sqrt{\pi}} \right)$	27

input `int(x*cos(x), x, method=_RETURNVERBOSE)`

output `cos(x)+x*sin(x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int x \cos(x) dx = x \sin(x) + \cos(x)$$

input `integrate(x*cos(x),x, algorithm="fricas")`

output `x*sin(x) + cos(x)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int x \cos(x) dx = x \sin(x) + \cos(x)$$

input `integrate(x*cos(x),x)`

output `x*sin(x) + cos(x)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int x \cos(x) dx = x \sin(x) + \cos(x)$$

input `integrate(x*cos(x),x, algorithm="maxima")`

output `x*sin(x) + cos(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int x \cos(x) dx = x \sin(x) + \cos(x)$$

input `integrate(x*cos(x),x, algorithm="giac")`

output `x*sin(x) + cos(x)`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int x \cos(x) dx = \cos(x) + x \sin(x)$$

input `int(x*cos(x),x)`

output `cos(x) + x*sin(x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int x \cos(x) dx = \cos(x) + \sin(x)x$$

input `int(x*cos(x),x)`

output `cos(x) + sin(x)*x`

3.62 $\int \cos(\sqrt{x}) dx$

Optimal result	389
Mathematica [A] (verified)	389
Rubi [A] (verified)	390
Maple [A] (verified)	391
Fricas [A] (verification not implemented)	392
Sympy [A] (verification not implemented)	392
Maxima [A] (verification not implemented)	392
Giac [A] (verification not implemented)	393
Mupad [B] (verification not implemented)	393
Reduce [B] (verification not implemented)	393

Optimal result

Integrand size = 6, antiderivative size = 22

$$\int \cos(\sqrt{x}) dx = 2 \cos(\sqrt{x}) + 2\sqrt{x} \sin(\sqrt{x})$$

output `2*cos(x^(1/2))+2*sin(x^(1/2))*x^(1/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \cos(\sqrt{x}) dx = 2 \cos(\sqrt{x}) + 2\sqrt{x} \sin(\sqrt{x})$$

input `Integrate[Cos[Sqrt[x]],x]`

output `2*Cos[Sqrt[x]] + 2*Sqrt[x]*Sin[Sqrt[x]]`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.95, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3843, 3042, 3777, 25, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(\sqrt{x}) \, dx \\
 & \quad \downarrow \text{3843} \\
 & 2 \int \sqrt{x} \cos(\sqrt{x}) \, d\sqrt{x} \\
 & \quad \downarrow \text{3042} \\
 & 2 \int \sqrt{x} \sin\left(\sqrt{x} + \frac{\pi}{2}\right) \, d\sqrt{x} \\
 & \quad \downarrow \text{3777} \\
 & 2 \left(\int -\sin(\sqrt{x}) \, d\sqrt{x} + \sqrt{x} \sin(\sqrt{x}) \right) \\
 & \quad \downarrow \text{25} \\
 & 2 \left(\sqrt{x} \sin(\sqrt{x}) - \int \sin(\sqrt{x}) \, d\sqrt{x} \right) \\
 & \quad \downarrow \text{3042} \\
 & 2 \left(\sqrt{x} \sin(\sqrt{x}) - \int \sin(\sqrt{x}) \, d\sqrt{x} \right) \\
 & \quad \downarrow \text{3118} \\
 & 2(\sqrt{x} \sin(\sqrt{x}) + \cos(\sqrt{x}))
 \end{aligned}$$

input `Int[Cos[Sqrt[x]], x]`

output `2*(Cos[Sqrt[x]] + Sqrt[x]*Sin[Sqrt[x]])`

Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`
- rule 3777 `Int[((c_.) + (d_.)*(x_)^(m_.))*sin[(e_.) + (f_.)*(x_)], x_Symbol] :> Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`
- rule 3843 `Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_)^(n_)])*(b_.))^(p_.), x_Symbol] :> Simp[1/(n*f) Subst[Int[x^(1/n - 1)*(a + b*Cos[c + d*x]]^p, x], x, (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[1/n]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

method	result	size
derivativedivides	$2 \cos(\sqrt{x}) + 2 \sin(\sqrt{x}) \sqrt{x}$	17
default	$2 \cos(\sqrt{x}) + 2 \sin(\sqrt{x}) \sqrt{x}$	17
meijerg	$4\sqrt{\pi} \left(-\frac{1}{2\sqrt{\pi}} + \frac{\cos(\sqrt{x})}{2\sqrt{\pi}} + \frac{\sqrt{x} \sin(\sqrt{x})}{2\sqrt{\pi}} \right)$	33

input `int(cos(x^(1/2)),x,method=_RETURNVERBOSE)`

output `2*cos(x^(1/2))+2*sin(x^(1/2))*x^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \cos(\sqrt{x}) dx = 2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

input `integrate(cos(x^(1/2)),x, algorithm="fricas")`

output `2*sqrt(x)*sin(sqrt(x)) + 2*cos(sqrt(x))`

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \cos(\sqrt{x}) dx = 2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

input `integrate(cos(x**(1/2)),x)`

output `2*sqrt(x)*sin(sqrt(x)) + 2*cos(sqrt(x))`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \cos(\sqrt{x}) dx = 2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

input `integrate(cos(x^(1/2)),x, algorithm="maxima")`

output `2*sqrt(x)*sin(sqrt(x)) + 2*cos(sqrt(x))`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \cos(\sqrt{x}) dx = 2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

input `integrate(cos(x^(1/2)),x, algorithm="giac")`

output `2*sqrt(x)*sin(sqrt(x)) + 2*cos(sqrt(x))`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \cos(\sqrt{x}) dx = 2 \cos(\sqrt{x}) + 2\sqrt{x} \sin(\sqrt{x})$$

input `int(cos(x^(1/2)),x)`

output `2*cos(x^(1/2)) + 2*x^(1/2)*sin(x^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.59

$$\int \cos(\sqrt{x}) dx = 2 \cos(\sqrt{x}) + 2\sqrt{x} \sin(\sqrt{x})$$

input `int(cos(x^(1/2)),x)`

output `2*(cos(sqrt(x)) + sqrt(x)*sin(sqrt(x)))`

3.63 $\int x \cos(x) dx$

Optimal result	394
Mathematica [A] (verified)	394
Rubi [A] (verified)	395
Maple [A] (verified)	396
Fricas [A] (verification not implemented)	397
Sympy [A] (verification not implemented)	397
Maxima [A] (verification not implemented)	397
Giac [A] (verification not implemented)	398
Mupad [B] (verification not implemented)	398
Reduce [B] (verification not implemented)	398

Optimal result

Integrand size = 4, antiderivative size = 7

$$\int x \cos(x) dx = \cos(x) + x \sin(x)$$

output `cos(x)+x*sin(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int x \cos(x) dx = \cos(x) + x \sin(x)$$

input `Integrate[x*Cos[x],x]`

output `Cos[x] + x*Sin[x]`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$, Rules used = {3042, 3777, 25, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \cos(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x \sin\left(x + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{3777} \\
 & \int -\sin(x) dx + x \sin(x) \\
 & \quad \downarrow \text{25} \\
 & x \sin(x) - \int \sin(x) dx \\
 & \quad \downarrow \text{3042} \\
 & x \sin(x) - \int \sin(x) dx \\
 & \quad \downarrow \text{3118} \\
 & x \sin(x) + \cos(x)
 \end{aligned}$$

input

Int [x*Cos [x] , x]

output

Cos [x] + x*Sin [x]

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_)^(m_.))*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.14

method	result	size
default	$\cos(x) + x \sin(x)$	8
risch	$\cos(x) + x \sin(x)$	8
parts	$\cos(x) + x \sin(x)$	8
orering	$\cos(x) + x \sin(x)$	8
parallelrisc	$1 + \cos(x) + x \sin(x)$	9
norman	$\frac{2x \tan(\frac{x}{2}) + 2}{1 + \tan(\frac{x}{2})^2}$	21
meijerg	$2\sqrt{\pi} \left(-\frac{1}{2\sqrt{\pi}} + \frac{\cos(x)}{2\sqrt{\pi}} + \frac{x \sin(x)}{2\sqrt{\pi}} \right)$	27

input `int(x*cos(x), x, method=_RETURNVERBOSE)`

output `cos(x)+x*sin(x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int x \cos(x) dx = x \sin(x) + \cos(x)$$

input `integrate(x*cos(x),x, algorithm="fricas")`

output `x*sin(x) + cos(x)`

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int x \cos(x) dx = x \sin(x) + \cos(x)$$

input `integrate(x*cos(x),x)`

output `x*sin(x) + cos(x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int x \cos(x) dx = x \sin(x) + \cos(x)$$

input `integrate(x*cos(x),x, algorithm="maxima")`

output `x*sin(x) + cos(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int x \cos(x) dx = x \sin(x) + \cos(x)$$

input `integrate(x*cos(x),x, algorithm="giac")`

output `x*sin(x) + cos(x)`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int x \cos(x) dx = \cos(x) + x \sin(x)$$

input `int(x*cos(x),x)`

output `cos(x) + x*sin(x)`

Reduce [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int x \cos(x) dx = \cos(x) + \sin(x)x$$

input `int(x*cos(x),x)`

output `cos(x) + sin(x)*x`

3.64 $\int x \log^2(x) dx$

Optimal result	399
Mathematica [A] (verified)	399
Rubi [A] (verified)	400
Maple [A] (verified)	401
Fricas [A] (verification not implemented)	401
Sympy [A] (verification not implemented)	402
Maxima [A] (verification not implemented)	402
Giac [A] (verification not implemented)	402
Mupad [B] (verification not implemented)	403
Reduce [B] (verification not implemented)	403

Optimal result

Integrand size = 6, antiderivative size = 28

$$\int x \log^2(x) dx = \frac{x^2}{4} - \frac{1}{2}x^2 \log(x) + \frac{1}{2}x^2 \log^2(x)$$

output

```
1/4*x^2-1/2*x^2*ln(x)+1/2*x^2*ln(x)^2
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int x \log^2(x) dx = \frac{x^2}{4} - \frac{1}{2}x^2 \log(x) + \frac{1}{2}x^2 \log^2(x)$$

input

```
Integrate[x*Log[x]^2,x]
```

output

```
x^2/4 - (x^2*Log[x])/2 + (x^2*Log[x]^2)/2
```


Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2742, 2741}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \log^2(x) dx$$

$$\downarrow 2742$$

$$\frac{1}{2}x^2 \log^2(x) - \int x \log(x) dx$$

$$\downarrow 2741$$

$$\frac{x^2}{4} + \frac{1}{2}x^2 \log^2(x) - \frac{1}{2}x^2 \log(x)$$

input `Int[x*Log[x]^2,x]`

output `x^2/4 - (x^2*Log[x])/2 + (x^2*Log[x]^2)/2`

Defintions of rubi rules used

rule 2741 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((d_.)*(x_)^(m_.), x_Symbol] :> Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])/(d*(m + 1))), x] - Simp[b*n*((d*x)^(m + 1)/(d*(m + 1)^2)), x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[m, -1]`

rule 2742 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((d_.)*(x_)^(m_.), x_Symbol] :> Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])^p/(d*(m + 1))), x] - Simp[b*n*(p/(m + 1)) Int[(d*x)^m*(a + b*Log[c*x^n])^(p - 1), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[m, -1] && GtQ[p, 0]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.82

method	result	size
default	$\frac{x^2}{4} - \frac{x^2 \ln(x)}{2} + \frac{x^2 \ln(x)^2}{2}$	23
norman	$\frac{x^2}{4} - \frac{x^2 \ln(x)}{2} + \frac{x^2 \ln(x)^2}{2}$	23
risch	$\frac{x^2}{4} - \frac{x^2 \ln(x)}{2} + \frac{x^2 \ln(x)^2}{2}$	23
parallelrisc	$\frac{x^2}{4} - \frac{x^2 \ln(x)}{2} + \frac{x^2 \ln(x)^2}{2}$	23
parts	$\frac{x^2}{4} - \frac{x^2 \ln(x)}{2} + \frac{x^2 \ln(x)^2}{2}$	23
orering	$\frac{7x^2 \ln(x)^2}{8} - \frac{3x^2 (\ln(x)^2 + 2 \ln(x))}{8} + \frac{x^3 (\frac{2 \ln(x)}{x} + \frac{2}{x})}{8}$	43

input `int(x*ln(x)^2,x,method=_RETURNVERBOSE)`output `1/4*x^2-1/2*x^2*ln(x)+1/2*x^2*ln(x)^2`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int x \log^2(x) dx = \frac{1}{2} x^2 \log(x)^2 - \frac{1}{2} x^2 \log(x) + \frac{1}{4} x^2$$

input `integrate(x*log(x)^2,x, algorithm="fricas")`output `1/2*x^2*log(x)^2 - 1/2*x^2*log(x) + 1/4*x^2`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int x \log^2(x) dx = \frac{x^2 \log(x)^2}{2} - \frac{x^2 \log(x)}{2} + \frac{x^2}{4}$$

input `integrate(x*ln(x)**2,x)`output `x**2*log(x)**2/2 - x**2*log(x)/2 + x**2/4`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.61

$$\int x \log^2(x) dx = \frac{1}{4} (2 \log(x)^2 - 2 \log(x) + 1) x^2$$

input `integrate(x*log(x)^2,x, algorithm="maxima")`output `1/4*(2*log(x)^2 - 2*log(x) + 1)*x^2`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int x \log^2(x) dx = \frac{1}{2} x^2 \log(x)^2 - \frac{1}{2} x^2 \log(x) + \frac{1}{4} x^2$$

input `integrate(x*log(x)^2,x, algorithm="giac")`output `1/2*x^2*log(x)^2 - 1/2*x^2*log(x) + 1/4*x^2`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.61

$$\int x \log^2(x) dx = \frac{x^2 (2 \ln(x)^2 - 2 \ln(x) + 1)}{4}$$

input `int(x*log(x)^2,x)`

output `(x^2*(2*log(x)^2 - 2*log(x) + 1))/4`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.61

$$\int x \log^2(x) dx = \frac{x^2 (2 \log(x)^2 - 2 \log(x) + 1)}{4}$$

input `int(x*log(x)^2,x)`

output `(x**2*(2*log(x)**2 - 2*log(x) + 1))/4`

3.65 $\int \cos(x) (1 + \sin^3(x)) dx$

Optimal result	404
Mathematica [A] (verified)	404
Rubi [A] (verified)	405
Maple [A] (verified)	406
Fricas [A] (verification not implemented)	406
Sympy [A] (verification not implemented)	407
Maxima [A] (verification not implemented)	407
Giac [A] (verification not implemented)	407
Mupad [B] (verification not implemented)	408
Reduce [B] (verification not implemented)	408

Optimal result

Integrand size = 9, antiderivative size = 11

$$\int \cos(x) (1 + \sin^3(x)) dx = \sin(x) + \frac{\sin^4(x)}{4}$$

output

```
sin(x)+1/4*sin(x)^4
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \cos(x) (1 + \sin^3(x)) dx = \sin(x) + \frac{\sin^4(x)}{4}$$

input

```
Integrate[Cos[x]*(1 + Sin[x]^3),x]
```

output

```
Sin[x] + Sin[x]^4/4
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3702, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (\sin^3(x) + 1) \cos(x) dx \\ & \quad \downarrow \text{3042} \\ & \int (\sin(x)^3 + 1) \cos(x) dx \\ & \quad \downarrow \text{3702} \\ & \int (\sin^3(x) + 1) d \sin(x) \\ & \quad \downarrow \text{2009} \\ & \frac{\sin^4(x)}{4} + \sin(x) \end{aligned}$$

input `Int[Cos[x]*(1 + Sin[x]^3),x]`

output `Sin[x] + Sin[x]^4/4`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3702

```
Int[cos[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*((c_.)*sin[(e_.) + (f_.)*(x_)]^(n_.))^(p_.), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Simp[ff/f Subst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b*(c*ff*x)^n)^p, x], x, Sin[e + f*x]/ff], x]] /; FreeQ[{a, b, c, e, f, n, p}, x] && IntegerQ[(m - 1)/2] && (EqQ[n, 4] || GtQ[m, 0] || IGtQ[p, 0] || IntegersQ[m, p])
```

Maple [A] (verified)

Time = 0.69 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

method	result	size
derivativedivides	$\sin(x) + \frac{\sin(x)^4}{4}$	10
default	$\sin(x) + \frac{\sin(x)^4}{4}$	10
risch	$\sin(x) + \frac{\cos(4x)}{32} - \frac{\cos(2x)}{8}$	16
parallelrisc	$\frac{3}{32} + \sin(x) + \frac{\cos(4x)}{32} - \frac{\cos(2x)}{8}$	17
orering	$\sin(x) (1 + \sin(x)^3) - \frac{3 \sin(x)^2 \cos(x)^2}{16} - \frac{27 \sin(x)^4}{32} - \frac{3 \cos(x)^4}{32}$	33

input

```
int(cos(x)*(1+sin(x)^3),x,method=_RETURNVERBOSE)
```

output

```
sin(x)+1/4*sin(x)^4
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.36

$$\int \cos(x) (1 + \sin^3(x)) dx = \frac{1}{4} \cos(x)^4 - \frac{1}{2} \cos(x)^2 + \sin(x)$$

input

```
integrate(cos(x)*(1+sin(x)^3),x, algorithm="fricas")
```

output

```
1/4*cos(x)^4 - 1/2*cos(x)^2 + sin(x)
```

Sympy [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int \cos(x) (1 + \sin^3(x)) dx = \frac{\sin^4(x)}{4} + \sin(x)$$

input `integrate(cos(x)*(1+sin(x)**3),x)`

output `sin(x)**4/4 + sin(x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \cos(x) (1 + \sin^3(x)) dx = \frac{1}{4} \sin(x)^4 + \sin(x)$$

input `integrate(cos(x)*(1+sin(x)^3),x, algorithm="maxima")`

output `1/4*sin(x)^4 + sin(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \cos(x) (1 + \sin^3(x)) dx = \frac{1}{4} \sin(x)^4 + \sin(x)$$

input `integrate(cos(x)*(1+sin(x)^3),x, algorithm="giac")`

output `1/4*sin(x)^4 + sin(x)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \cos(x) (1 + \sin^3(x)) dx = \frac{\sin(x)^4}{4} + \sin(x)$$

input `int(cos(x)*(sin(x)^3 + 1),x)`

output `sin(x) + sin(x)^4/4`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

$$\int \cos(x) (1 + \sin^3(x)) dx = \frac{\sin(x) (\sin(x)^3 + 4)}{4}$$

input `int(cos(x)*(1+sin(x)^3),x)`

output `(sin(x)*(sin(x)**3 + 4))/4`

$$3.66 \quad \int \frac{1}{x(1+\log^2(x))} dx$$

Optimal result	409
Mathematica [A] (verified)	409
Rubi [A] (verified)	410
Maple [A] (verified)	411
Fricas [A] (verification not implemented)	411
Sympy [B] (verification not implemented)	411
Maxima [A] (verification not implemented)	412
Giac [A] (verification not implemented)	412
Mupad [B] (verification not implemented)	412
Reduce [B] (verification not implemented)	413

Optimal result

Integrand size = 12, antiderivative size = 3

$$\int \frac{1}{x(1+\log^2(x))} dx = \arctan(\log(x))$$

output `arctan(ln(x))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1+\log^2(x))} dx = \arctan(\log(x))$$

input `Integrate[1/(x*(1 + Log[x]^2)),x]`

output `ArcTan[Log[x]]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3039, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x (\log^2(x) + 1)} dx$$

↓ 3039

$$\int \frac{1}{\log^2(x) + 1} d\log(x)$$

↓ 216

$$\arctan(\log(x))$$

input `Int[1/(x*(1 + Log[x]^2)),x]`

output `ArcTan[Log[x]]`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3039 `Int[u_, x_Symbol] := With[{lst = FunctionOfLog[Cancel[x*u], x]}, Simp[1/lst[[3]] Subst[Int[lst[[1]], x], x, Log[lst[[2]]]], x] /; !FalseQ[lst] /; NonsumQ[u]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
derivativdivides	$\arctan(\ln(x))$	4
default	$\arctan(\ln(x))$	4
risch	$\frac{i \ln(\ln(x)+i)}{2} - \frac{i \ln(\ln(x)-i)}{2}$	20
parallelrisch	$\frac{i \ln(\ln(x)+i)}{2} - \frac{i \ln(\ln(x)-i)}{2}$	20

input `int(1/x/(1+ln(x)^2),x,method=_RETURNVERBOSE)`

output `arctan(ln(x))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1+\log^2(x))} dx = \arctan(\log(x))$$

input `integrate(1/x/(1+log(x)^2),x, algorithm="fricas")`

output `arctan(log(x))`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(3) = 6$.

Time = 0.06 (sec) , antiderivative size = 15, normalized size of antiderivative = 5.00

$$\int \frac{1}{x(1+\log^2(x))} dx = \text{RootSum}(4z^2 + 1, (i \mapsto i \log(2i + \log(x))))$$

input `integrate(1/x/(1+ln(x)**2),x)`

output `RootSum(4*_z**2 + 1, Lambda(_i, _i*log(2*_i + log(x))))`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1+\log^2(x))} dx = \arctan(\log(x))$$

input `integrate(1/x/(1+log(x)^2),x, algorithm="maxima")`

output `arctan(log(x))`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1+\log^2(x))} dx = \arctan(\log(x))$$

input `integrate(1/x/(1+log(x)^2),x, algorithm="giac")`

output `arctan(log(x))`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1+\log^2(x))} dx = \operatorname{atan}(\ln(x))$$

input `int(1/(x*(log(x)^2 + 1)),x)`

output `atan(log(x))`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1+\log^2(x))} dx = \text{atan}(\log(x))$$

input `int(1/x/(1+log(x)^2),x)`

output `atan(log(x))`

3.67 $\int \frac{1}{\sqrt{1-x^2}(1+\arcsin(x)^2)} dx$

Optimal result	414
Mathematica [A] (verified)	414
Rubi [A] (verified)	415
Maple [A] (verified)	416
Fricas [A] (verification not implemented)	416
Sympy [A] (verification not implemented)	416
Maxima [F]	417
Giac [A] (verification not implemented)	417
Mupad [B] (verification not implemented)	417
Reduce [B] (verification not implemented)	418

Optimal result

Integrand size = 20, antiderivative size = 3

$$\int \frac{1}{\sqrt{1-x^2}(1+\arcsin(x)^2)} dx = \arctan(\arcsin(x))$$

output `arctan(arcsin(x))`

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1-x^2}(1+\arcsin(x)^2)} dx = \arctan(\arcsin(x))$$

input `Integrate[1/(Sqrt[1 - x^2]*(1 + ArcSin[x]^2)),x]`

output `ArcTan[ArcSin[x]]`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {7247, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{1-x^2}(\arcsin(x)^2+1)} dx$$

↓ 7247

$$\int \frac{1}{\arcsin(x)^2+1} d\arcsin(x)$$

↓ 216

$$\arctan(\arcsin(x))$$

input `Int[1/(Sqrt[1 - x^2]*(1 + ArcSin[x]^2)),x]`

output `ArcTan[ArcSin[x]]`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 7247 `Int[(u_.)*((a_.) + (b_.)*(y_)^(n_))^(p_), x_Symbol] := With[{q = Derivative Divides[y, u, x]}, Simp[q Subst[Int[(a + b*x^n)^p, x], x, y], x] /; !FalseQ[q]] /; FreeQ[{a, b, n, p}, x]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
derivativdivides	$\arctan(\arcsin(x))$	4
default	$\arctan(\arcsin(x))$	4

input `int(1/(1+arcsin(x)^2)/(-x^2+1)^(1/2),x,method=_RETURNVERBOSE)`output `arctan(arcsin(x))`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1-x^2}(1+\arcsin(x)^2)} dx = \arctan(\arcsin(x))$$

input `integrate(1/(1+arcsin(x)^2)/(-x^2+1)^(1/2),x, algorithm="fricas")`output `arctan(arcsin(x))`**Sympy [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1-x^2}(1+\arcsin(x)^2)} dx = \operatorname{atan}(\operatorname{asin}(x))$$

input `integrate(1/(1+asin(x)**2)/(-x**2+1)**(1/2),x)`output `atan(asin(x))`

Maxima [F]

$$\int \frac{1}{\sqrt{1-x^2}(1+\arcsin(x)^2)} dx = \int \frac{1}{\sqrt{-x^2+1}(\arcsin(x)^2+1)} dx$$

input `integrate(1/(1+arcsin(x)^2)/(-x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(1/(sqrt(-x^2 + 1)*(arcsin(x)^2 + 1)), x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1-x^2}(1+\arcsin(x)^2)} dx = \arctan(\arcsin(x))$$

input `integrate(1/(1+arcsin(x)^2)/(-x^2+1)^(1/2),x, algorithm="giac")`

output `arctan(arcsin(x))`

Mupad [B] (verification not implemented)

Time = 1.79 (sec) , antiderivative size = 43, normalized size of antiderivative = 14.33

$$\int \frac{1}{\sqrt{1-x^2}(1+\arcsin(x)^2)} dx = \frac{\ln\left(\frac{-1+\arcsin(x) \text{ li}}{\sqrt{1-x^2}}\right) \text{ li}}{2} - \frac{\ln\left(\frac{1+\arcsin(x) \text{ li}}{\sqrt{1-x^2}}\right) \text{ li}}{2}$$

input `int(1/((1 - x^2)^(1/2)*(asin(x)^2 + 1)),x)`

output `(log((asin(x)*li - 1)/(1 - x^2)^(1/2))*li)/2 - (log((asin(x)*li + 1)/(1 - x^2)^(1/2))*li)/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1-x^2}(1+\arcsin(x)^2)} dx = \text{atan}(\text{asin}(x))$$

input `int(1/(1+asin(x)^2)/(-x^2+1)^(1/2),x)`

output `atan(asin(x))`

3.68 $\int \frac{\sin(x)}{\cos(x)+\sin(x)} dx$

Optimal result	419
Mathematica [A] (verified)	419
Rubi [A] (verified)	420
Maple [C] (verified)	421
Fricas [A] (verification not implemented)	422
Sympy [A] (verification not implemented)	422
Maxima [B] (verification not implemented)	422
Giac [A] (verification not implemented)	423
Mupad [B] (verification not implemented)	423
Reduce [B] (verification not implemented)	424

Optimal result

Integrand size = 10, antiderivative size = 16

$$\int \frac{\sin(x)}{\cos(x) + \sin(x)} dx = \frac{x}{2} - \frac{1}{2} \log(\cos(x) + \sin(x))$$

output `1/2*x-1/2*ln(cos(x)+sin(x))`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\sin(x)}{\cos(x) + \sin(x)} dx = \frac{x}{2} - \frac{1}{2} \log(\cos(x) + \sin(x))$$

input `Integrate[Sin[x]/(Cos[x] + Sin[x]),x]`

output `x/2 - Log[Cos[x] + Sin[x]]/2`

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3576, 3042, 3612}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin(x)}{\sin(x) + \cos(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(x)}{\sin(x) + \cos(x)} dx \\
 & \quad \downarrow \text{3576} \\
 & \frac{x}{2} - \frac{1}{2} \int \frac{\cos(x) - \sin(x)}{\cos(x) + \sin(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{x}{2} - \frac{1}{2} \int \frac{\cos(x) - \sin(x)}{\cos(x) + \sin(x)} dx \\
 & \quad \downarrow \text{3612} \\
 & \frac{x}{2} - \frac{1}{2} \log(\sin(x) + \cos(x))
 \end{aligned}$$

input `Int[Sin[x]/(Cos[x] + Sin[x]),x]`

output `x/2 - Log[Cos[x] + Sin[x]]/2`

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3576 `Int[sin[(c_.) + (d_.)*(x_)]/(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[b*(x/(a^2 + b^2)), x] - Simp[a/(a^2 + b^2) Int[(b*Cos[c + d*x] - a*Sin[c + d*x])/(a*Cos[c + d*x] + b*Sin[c + d*x]), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]`

rule 3612 `Int[((A_.) + cos[(d_.) + (e_.)*(x_)]*(B_.) + (C_.)*sin[(d_.) + (e_.)*(x_)]) / ((a_.) + cos[(d_.) + (e_.)*(x_)]*(b_.) + (c_.)*sin[(d_.) + (e_.)*(x_)]), x_Symbol] := Simp[(b*B + c*C)*(x/(b^2 + c^2)), x] + Simp[(c*B - b*C)*(Log[a + b*Cos[d + e*x] + c*Sin[d + e*x]]/(e*(b^2 + c^2))), x] /; FreeQ[{a, b, c, d, e, A, B, C}, x] && NeQ[b^2 + c^2, 0] && EqQ[A*(b^2 + c^2) - a*(b*B + c*C), 0]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.25

method	result	size
risch	$\frac{x}{2} + \frac{ix}{2} - \frac{\ln(e^{2ix} + i)}{2}$	20
default	$-\frac{\ln(\tan(x)+1)}{2} + \frac{\ln(1+\tan(x)^2)}{4} + \frac{\arctan(\tan(x))}{2}$	23
parallelrisch	$\frac{x}{2} + \ln\left(\sqrt{\sec\left(\frac{x}{2}\right)^2}\right) + \ln\left(\frac{1}{\sqrt{\tan\left(\frac{x}{2}\right)^2 - 2\tan\left(\frac{x}{2}\right) - 1}}\right)$	31
norman	$\frac{x}{2} + \frac{x \tan\left(\frac{x}{2}\right)^2}{1 + \tan\left(\frac{x}{2}\right)^2} + \frac{\ln\left(1 + \tan\left(\frac{x}{2}\right)^2\right)}{2} - \frac{\ln\left(\tan\left(\frac{x}{2}\right)^2 - 2\tan\left(\frac{x}{2}\right) - 1\right)}{2}$	54

input `int(sin(x)/(cos(x)+sin(x)),x,method=_RETURNVERBOSE)`

output `1/2*x+1/2*I*x-1/2*ln(exp(2*I*x)+I)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{\sin(x)}{\cos(x) + \sin(x)} dx = \frac{1}{2}x - \frac{1}{4} \log(2 \cos(x) \sin(x) + 1)$$

input `integrate(sin(x)/(cos(x)+sin(x)),x, algorithm="fricas")`

output `1/2*x - 1/4*log(2*cos(x)*sin(x) + 1)`

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{\sin(x)}{\cos(x) + \sin(x)} dx = \frac{x}{2} - \frac{\log(\sin(x) + \cos(x))}{2}$$

input `integrate(sin(x)/(cos(x)+sin(x)),x)`

output `x/2 - log(sin(x) + cos(x))/2`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 53 vs. 2(12) = 24.

Time = 0.11 (sec) , antiderivative size = 53, normalized size of antiderivative = 3.31

$$\int \frac{\sin(x)}{\cos(x) + \sin(x)} dx = \arctan\left(\frac{\sin(x)}{\cos(x) + 1}\right) - \frac{1}{2} \log\left(-\frac{2 \sin(x)}{\cos(x) + 1} + \frac{\sin(x)^2}{(\cos(x) + 1)^2} - 1\right) + \frac{1}{2} \log\left(\frac{\sin(x)^2}{(\cos(x) + 1)^2} + 1\right)$$

input `integrate(sin(x)/(cos(x)+sin(x)),x, algorithm="maxima")`

output `arctan(sin(x)/(cos(x) + 1)) - 1/2*log(-2*sin(x)/(cos(x) + 1) + sin(x)^2/(cos(x) + 1)^2 - 1) + 1/2*log(sin(x)^2/(cos(x) + 1)^2 + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.31

$$\int \frac{\sin(x)}{\cos(x) + \sin(x)} dx = \frac{1}{2}x + \frac{1}{4} \log(\tan(x)^2 + 1) - \frac{1}{2} \log(|\tan(x) + 1|)$$

input `integrate(sin(x)/(cos(x)+sin(x)),x, algorithm="giac")`

output `1/2*x + 1/4*log(tan(x)^2 + 1) - 1/2*log(abs(tan(x) + 1))`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.81

$$\int \frac{\sin(x)}{\cos(x) + \sin(x)} dx = \frac{x}{2} - \frac{\ln(\cos(x - \frac{\pi}{4}))}{2}$$

input `int(sin(x)/(cos(x) + sin(x)),x)`

output `x/2 - log(cos(x - pi/4))/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{\sin(x)}{\cos(x) + \sin(x)} dx = -\frac{\log(\cos(x) + \sin(x))}{2} + \frac{x}{2}$$

input `int(sin(x)/(cos(x)+sin(x)),x)`

output `(- log(cos(x) + sin(x)) + x)/2`

$$3.69 \quad \int -\frac{\sqrt{A^2+B^2(1-y^2)}}{1-y^2} dy$$

Optimal result	425
Mathematica [A] (verified)	425
Rubi [A] (verified)	426
Maple [A] (verified)	428
Fricas [B] (verification not implemented)	429
Sympy [F]	429
Maxima [B] (verification not implemented)	430
Giac [B] (verification not implemented)	430
Mupad [F(-1)]	431
Reduce [B] (verification not implemented)	432

Optimal result

Integrand size = 30, antiderivative size = 53

$$\int -\frac{\sqrt{A^2+B^2(1-y^2)}}{1-y^2} dy = -B \arctan\left(\frac{By}{\sqrt{A^2+B^2-B^2y^2}}\right) - A \operatorname{arctanh}\left(\frac{Ay}{\sqrt{A^2+B^2-B^2y^2}}\right)$$

output

```
-B*arctan(B*y/(-B^2*y^2+A^2+B^2)^(1/2))-A*arctanh(A*y/(-B^2*y^2+A^2+B^2)^(1/2))
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.38

$$\int -\frac{\sqrt{A^2+B^2(1-y^2)}}{1-y^2} dy = 2B \arctan\left(\frac{By}{\sqrt{A^2+B^2}-\sqrt{A^2+B^2-B^2y^2}}\right) - A \operatorname{arctanh}\left(\frac{\sqrt{A^2+B^2-B^2y^2}}{Ay}\right)$$

input

```
Integrate[-(Sqrt[A^2 + B^2*(1 - y^2)]/(1 - y^2)),y]
```

output

```
2*B*ArcTan[(B*y)/(Sqrt[A^2 + B^2] - Sqrt[A^2 + B^2 - B^2*y^2])] - A*ArcTan
h[Sqrt[A^2 + B^2 - B^2*y^2]/(A*y)]
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.233$, Rules used = {25, 2074, 301, 224, 216, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int -\frac{\sqrt{A^2 + B^2(1 - y^2)}}{1 - y^2} dy \\
 & \quad \downarrow \text{25} \\
 & - \int \frac{\sqrt{A^2 + B^2(1 - y^2)}}{1 - y^2} dy \\
 & \quad \downarrow \text{2074} \\
 & - \int \frac{\sqrt{A^2 + B^2 - B^2y^2}}{1 - y^2} dy \\
 & \quad \downarrow \text{301} \\
 & A^2 \left(- \int \frac{1}{(1 - y^2)\sqrt{A^2 + B^2 - B^2y^2}} dy \right) - B^2 \int \frac{1}{\sqrt{A^2 + B^2 - B^2y^2}} dy \\
 & \quad \downarrow \text{224} \\
 & A^2 \left(- \int \frac{1}{(1 - y^2)\sqrt{A^2 + B^2 - B^2y^2}} dy \right) - B^2 \int \frac{1}{\frac{B^2y^2}{A^2 + B^2 - B^2y^2} + 1} d\frac{y}{\sqrt{A^2 + B^2 - B^2y^2}} \\
 & \quad \downarrow \text{216} \\
 & A^2 \left(- \int \frac{1}{(1 - y^2)\sqrt{A^2 + B^2 - B^2y^2}} dy \right) - B \arctan \left(\frac{By}{\sqrt{A^2 - B^2y^2 + B^2}} \right) \\
 & \quad \downarrow \text{291}
 \end{aligned}$$

$$A^2 \left(- \int \frac{1}{1 - \frac{A^2 y^2}{A^2 + B^2 - B^2 y^2}} d \frac{y}{\sqrt{A^2 + B^2 - B^2 y^2}} \right) - B \arctan \left(\frac{By}{\sqrt{A^2 - B^2 y^2 + B^2}} \right)$$

↓ 219

$$-B \arctan \left(\frac{By}{\sqrt{A^2 - B^2 y^2 + B^2}} \right) - A \operatorname{arctanh} \left(\frac{Ay}{\sqrt{A^2 - B^2 y^2 + B^2}} \right)$$

input `Int[-(Sqrt[A^2 + B^2*(1 - y^2)]/(1 - y^2)),y]`

output `-(B*ArcTan[(B*y)/Sqrt[A^2 + B^2 - B^2*y^2]]) - A*ArcTanh[(A*y)/Sqrt[A^2 + B^2 - B^2*y^2]]`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

rule 291 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

rule 301 `Int[((a_) + (b_.)*(x_)^2)^(p_.)/((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[b/d Int[(a + b*x^2)^(p - 1), x], x] - Simp[(b*c - a*d)/d Int[(a + b*x^2)^(p - 1)/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[p, 0] && (EqQ[p, 1/2] || EqQ[Denominator[p], 4] || (EqQ[p, 2/3] && EqQ[b*c + 3*a*d, 0]))`

rule 2074 `Int[(u_)^(p_.)*(v_)^(q_.), x_Symbol] := Int[ExpandToSum[u, x]^p*ExpandToSum[v, x]^q, x] /; FreeQ[{p, q}, x] && BinomialQ[{u, v}, x] && EqQ[BinomialDegree[u, x] - BinomialDegree[v, x], 0] && !BinomialMatchQ[{u, v}, x]`

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.68

method	result
pseudoelliptic	$\frac{A \ln\left(\frac{Ay - \sqrt{-B^2y^2 + A^2 + B^2}}{y}\right)}{2} - \frac{A \ln\left(\frac{Ay + \sqrt{-B^2y^2 + A^2 + B^2}}{y}\right)}{2} + B \arctan\left(\frac{\sqrt{-B^2y^2 + A^2 + B^2}}{By}\right)$
default	$-\frac{\sqrt{-B^2(1+y)^2 + 2B^2(1+y) + A^2}}{2} - \frac{B^2 \arctan\left(\frac{\sqrt{B^2}y}{\sqrt{-B^2(1+y)^2 + 2B^2(1+y) + A^2}}\right)}{2\sqrt{B^2}} + \frac{A^2 \ln\left(\frac{2A^2 + 2B^2(1+y) + 2\sqrt{A^2}\sqrt{-B^2(1+y)^2 + 2B^2(1+y) + A^2}}{1+y}\right)}{2\sqrt{A^2}}$

input `int(-(A^2+B^2*(-y^2+1))^(1/2)/(-y^2+1), y, method=_RETURNVERBOSE)`

output `1/2*A*ln((A*y-(-B^2*y^2+A^2+B^2)^(1/2))/y)-1/2*A*ln((A*y+(-B^2*y^2+A^2+B^2)^(1/2))/y)+B*arctan(1/B/y*(-B^2*y^2+A^2+B^2)^(1/2))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 144 vs. $2(49) = 98$.

Time = 0.08 (sec) , antiderivative size = 144, normalized size of antiderivative = 2.72

$$\begin{aligned} & \int -\frac{\sqrt{A^2 + B^2(1 - y^2)}}{1 - y^2} dy \\ &= B \arctan\left(\frac{\sqrt{-B^2y^2 + A^2 + B^2By}}{B^2y^2 - A^2 - B^2}\right) \\ &\quad - \frac{1}{4} A \log\left(-\frac{(A^2 - B^2)y^2 + 2\sqrt{-B^2y^2 + A^2 + B^2}Ay + A^2 + B^2}{y^2}\right) \\ &\quad + \frac{1}{4} A \log\left(-\frac{(A^2 - B^2)y^2 - 2\sqrt{-B^2y^2 + A^2 + B^2}Ay + A^2 + B^2}{y^2}\right) \end{aligned}$$

input `integrate(-(A^2+B^2*(-y^2+1))^(1/2)/(-y^2+1),y, algorithm="fricas")`

output `B*arctan(sqrt(-B^2*y^2 + A^2 + B^2)*B*y/(B^2*y^2 - A^2 - B^2)) - 1/4*A*log(-((A^2 - B^2)*y^2 + 2*sqrt(-B^2*y^2 + A^2 + B^2)*A*y + A^2 + B^2)/y^2) + 1/4*A*log(-((A^2 - B^2)*y^2 - 2*sqrt(-B^2*y^2 + A^2 + B^2)*A*y + A^2 + B^2)/y^2)`

Sympy [F]

$$\int -\frac{\sqrt{A^2 + B^2(1 - y^2)}}{1 - y^2} dy = \int \frac{\sqrt{A^2 - B^2y^2 + B^2}}{(y - 1)(y + 1)} dy$$

input `integrate(-(A**2+B**2*(-y**2+1))**(1/2)/(-y**2+1),y)`

output `Integral(sqrt(A**2 - B**2*y**2 + B**2)/((y - 1)*(y + 1)), y)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 115 vs. $2(49) = 98$.

Time = 0.11 (sec) , antiderivative size = 115, normalized size of antiderivative = 2.17

$$\int -\frac{\sqrt{A^2 + B^2(1 - y^2)}}{1 - y^2} dy = -B \arcsin\left(\frac{B^2 y}{\sqrt{A^2 B^2 + B^4}}\right) + \frac{1}{2} A \log\left(B^2 + \frac{A^2}{y + 1} + \frac{\sqrt{-B^2 y^2 + A^2 + B^2 A}}{y + 1}\right) - \frac{1}{2} A \log\left(-B^2 + \frac{2 A^2}{|2y - 2|} + \frac{2\sqrt{-B^2 y^2 + A^2 + B^2 A}}{|2y - 2|}\right)$$

input `integrate(-(A^2+B^2*(-y^2+1))^(1/2)/(-y^2+1),y, algorithm="maxima")`

output `-B*arcsin(B^2*y/sqrt(A^2*B^2 + B^4)) + 1/2*A*log(B^2 + A^2/(y + 1) + sqrt(-B^2*y^2 + A^2 + B^2)*A/(y + 1)) - 1/2*A*log(-B^2 + 2*A^2/abs(2*y - 2) + 2*sqrt(-B^2*y^2 + A^2 + B^2)*A/abs(2*y - 2))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 295 vs. $2(49) = 98$.

Time = 0.19 (sec) , antiderivative size = 295, normalized size of antiderivative = 5.57

$$\int -\frac{\sqrt{A^2 + B^2(1 - y^2)}}{1 - y^2} dy = \frac{\left(\pi \operatorname{sgn}(y) - 2 \arctan\left(\frac{B^2 y \left(\frac{(\sqrt{A^2 + B^2 B + \sqrt{-B^2 y^2 + A^2 + B^2 |B|}})^2}{B^4 y^2} - 1\right)}{2(\sqrt{A^2 + B^2 B + \sqrt{-B^2 y^2 + A^2 + B^2 |B|}})}\right)\right) B^2}{2 |B|} - \frac{AB \log\left(\left|-\left(\frac{B^2 y}{\sqrt{A^2 + B^2 B + \sqrt{-B^2 y^2 + A^2 + B^2 |B|}}}-\frac{\sqrt{A^2 + B^2 B + \sqrt{-B^2 y^2 + A^2 + B^2 |B|}}}{B^2 y}\right) B + 2 A\right|\right)}{2 |B|} + \frac{AB \log\left(\left|-\left(\frac{B^2 y}{\sqrt{A^2 + B^2 B + \sqrt{-B^2 y^2 + A^2 + B^2 |B|}}}-\frac{\sqrt{A^2 + B^2 B + \sqrt{-B^2 y^2 + A^2 + B^2 |B|}}}{B^2 y}\right) B - 2 A\right|\right)}{2 |B|}$$

input `integrate(-(A^2+B^2*(-y^2+1))^(1/2)/(-y^2+1),y, algorithm="giac")`

output `1/2*(pi*sgn(y) - 2*arctan(-1/2*B^2*y*((sqrt(A^2 + B^2)*B + sqrt(-B^2*y^2 + A^2 + B^2)*abs(B)))^2/(B^4*y^2) - 1)/(sqrt(A^2 + B^2)*B + sqrt(-B^2*y^2 + A^2 + B^2)*abs(B)))*B^2/abs(B) - 1/2*A*B*log(abs(-(B^2*y/(sqrt(A^2 + B^2)*B + sqrt(-B^2*y^2 + A^2 + B^2)*abs(B)) - (sqrt(A^2 + B^2)*B + sqrt(-B^2*y^2 + A^2 + B^2)*abs(B))/(B^2*y))*B + 2*A))/abs(B) + 1/2*A*B*log(abs(-(B^2*y/(sqrt(A^2 + B^2)*B + sqrt(-B^2*y^2 + A^2 + B^2)*abs(B)) - (sqrt(A^2 + B^2)*B + sqrt(-B^2*y^2 + A^2 + B^2)*abs(B))/(B^2*y))*B - 2*A))/abs(B)`

Mupad [F(-1)]

Timed out.

$$\int -\frac{\sqrt{A^2 + B^2(1 - y^2)}}{1 - y^2} dy$$

$$= \begin{cases} \int \frac{\sqrt{-B^2 y^2}}{y^2 - 1} dy & \text{if } A^2 + B^2 = 0 \\ \ln(2y\sqrt{-B^2} + 2\sqrt{A^2 - B^2 y^2 + B^2})\sqrt{-B^2} + \operatorname{atan}\left(\frac{y\sqrt{A^2} \operatorname{li}}{\sqrt{A^2 - B^2 y^2 + B^2}}\right)\sqrt{A^2} \operatorname{li} & \text{if } A^2 + B^2 \neq 0 \end{cases}$$

input `int((A^2 - B^2*(y^2 - 1))^(1/2)/(y^2 - 1),y)`

output `piecewise(A^2 + B^2 == 0, int((-B^2*y^2)^(1/2)/(y^2 - 1), y), A^2 + B^2 ~= 0, atan((y*(A^2)^(1/2)*1i)/(A^2 + B^2 - B^2*y^2)^(1/2))*(A^2)^(1/2)*1i + log(2*y*(-B^2)^(1/2) + 2*(A^2 + B^2 - B^2*y^2)^(1/2))*(-B^2)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 126, normalized size of antiderivative = 2.38

$$\int -\frac{\sqrt{A^2 + B^2(1-y^2)}}{1-y^2} dy = -a \sin\left(\frac{by}{\sqrt{a^2+b^2}}\right) b + a \operatorname{atan}\left(\frac{\tan\left(\frac{\operatorname{asin}\left(\frac{by}{\sqrt{a^2+b^2}}\right)}{2}\right) bi}{\sqrt{a^2+b^2}-a}\right) ai$$

$$- \frac{\log\left(-\sqrt{a^2+b^2} + \tan\left(\frac{\operatorname{asin}\left(\frac{by}{\sqrt{a^2+b^2}}\right)}{2}\right) b - a\right) a}{2}$$

$$+ \frac{\log\left(\sqrt{a^2+b^2} + \tan\left(\frac{\operatorname{asin}\left(\frac{by}{\sqrt{a^2+b^2}}\right)}{2}\right) b + a\right) a}{2}$$

input `int(-(A^2+B^2*(-y^2+1))^(1/2)/(-y^2+1),y)`output `(- 2*asin((b*y)/sqrt(a**2 + b**2))*b + 2*atan((tan(asin((b*y)/sqrt(a**2 + b**2)))/2)*b*i)/(sqrt(a**2 + b**2) - a)*a*i - log(- sqrt(a**2 + b**2) + tan(asin((b*y)/sqrt(a**2 + b**2)))/2)*b - a)*a + log(sqrt(a**2 + b**2) + tan(asin((b*y)/sqrt(a**2 + b**2)))/2)*b + a)*a)/2`

3.70
$$\int \frac{(-A^2 - B^2) \cos^2(z)}{B \left(1 - \frac{(A^2 + B^2) \sin^2(z)}{B^2}\right)} dz$$

Optimal result	433
Mathematica [B] (verified)	433
Rubi [B] (verified)	434
Maple [B] (verified)	436
Fricas [B] (verification not implemented)	437
Sympy [A] (verification not implemented)	437
Maxima [B] (verification not implemented)	438
Giac [B] (verification not implemented)	438
Mupad [B] (verification not implemented)	439
Reduce [B] (verification not implemented)	440

Optimal result

Integrand size = 39, antiderivative size = 16

$$\int \frac{(-A^2 - B^2) \cos^2(z)}{B \left(1 - \frac{(A^2 + B^2) \sin^2(z)}{B^2}\right)} dz = -Bz - A \operatorname{arctanh}\left(\frac{A \tan(z)}{B}\right)$$

output `-B*z-A*arctanh(A*tan(z)/B)`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 35 vs. 2(16) = 32.

Time = 0.17 (sec) , antiderivative size = 35, normalized size of antiderivative = 2.19

$$\int \frac{(-A^2 - B^2) \cos^2(z)}{B \left(1 - \frac{(A^2 + B^2) \sin^2(z)}{B^2}\right)} dz = -\frac{B(A^2 + B^2) \left(Bz + A \operatorname{arctanh}\left(\frac{A \tan(z)}{B}\right)\right)}{A^2 B + B^3}$$

input `Integrate[((-A^2 - B^2)*Cos[z]^2)/(B*(1 - ((A^2 + B^2)*Sin[z]^2)/B^2)),z]`

output

$$-\left(\frac{B(A^2 + B^2)(Bz + A \operatorname{ArcTanh}[(A \tan z)/B])}{A^2 B + B^3}\right)$$

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 46 vs. $2(16) = 32$.

Time = 0.27 (sec) , antiderivative size = 46, normalized size of antiderivative = 2.88, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.179$, Rules used = {27, 3042, 3670, 27, 303, 216, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(-A^2 - B^2) \cos^2(z)}{B \left(1 - \frac{(A^2 + B^2) \sin^2(z)}{B^2}\right)} dz \\ & \quad \downarrow \text{27} \\ & \frac{(A^2 + B^2) \int \frac{\cos^2(z)}{1 - \frac{(A^2 + B^2) \sin^2(z)}{B^2}} dz}{B} \\ & \quad \downarrow \text{3042} \\ & \frac{(A^2 + B^2) \int \frac{\cos(z)^2}{1 - \frac{(A^2 + B^2) \sin(z)^2}{B^2}} dz}{B} \\ & \quad \downarrow \text{3670} \\ & \frac{(A^2 + B^2) \int \frac{B^2}{(\tan^2(z) + 1)(B^2 - A^2 \tan^2(z))} d \tan(z)}{B} \\ & \quad \downarrow \text{27} \\ & -B(A^2 + B^2) \int \frac{1}{(\tan^2(z) + 1)(B^2 - A^2 \tan^2(z))} d \tan(z) \\ & \quad \downarrow \text{303} \\ & -B(A^2 + B^2) \left(\frac{A^2 \int \frac{1}{B^2 - A^2 \tan^2(z)} d \tan(z)}{A^2 + B^2} + \frac{\int \frac{1}{\tan^2(z) + 1} d \tan(z)}{A^2 + B^2} \right) \\ & \quad \downarrow \text{216} \end{aligned}$$

$$-B(A^2 + B^2) \left(\frac{A^2 \int \frac{1}{B^2 - A^2 \tan^2(z)} d \tan(z)}{A^2 + B^2} + \frac{\arctan(\tan(z))}{A^2 + B^2} \right)$$

↓ 221

$$-B(A^2 + B^2) \left(\frac{\arctan(\tan(z))}{A^2 + B^2} + \frac{A \operatorname{arctanh}\left(\frac{A \tan(z)}{B}\right)}{B(A^2 + B^2)} \right)$$

input `Int[((-A^2 - B^2)*Cos[z]^2)/(B*(1 - ((A^2 + B^2)*Sin[z]^2)/B^2)),z]`

output `-(B*(A^2 + B^2)*(ArcTan[Tan[z]]/(A^2 + B^2) + (A*ArcTanh[(A*Tan[z])/B])/(B*(A^2 + B^2))))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 303 `Int[1/(((a_) + (b_.)*(x_)^2)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x^2), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3670

```
Int[cos[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(
p_.), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Su
bst[Int[(a + (a + b)*ff^2*x^2)^p/(1 + ff^2*x^2)^(m/2 + p + 1), x], x, Tan[e
+ f*x]/ff], x]] /; FreeQ[{a, b, e, f}, x] && IntegerQ[m/2] && IntegerQ[p]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 73 vs. 2(16) = 32.

Time = 0.55 (sec) , antiderivative size = 74, normalized size of antiderivative = 4.62

method	result
default	$(-A^2 - B^2) B \left(\frac{\arctan(\tan(z))}{A^2+B^2} + \frac{A \ln(A \tan(z)+B)}{2B(A^2+B^2)} - \frac{A \ln(A \tan(z)-B)}{2B(A^2+B^2)} \right)$
parallelsch	$\frac{(-A^2 - B^2) \left(2Bz + A \left(\ln \left(2A \tan\left(\frac{z}{2}\right) + 2B - \sec\left(\frac{z}{2}\right)^2 B \right) - \ln \left(2A \tan\left(\frac{z}{2}\right) - 2B + \sec\left(\frac{z}{2}\right)^2 B \right) \right) \right)}{2A^2 + 2B^2}$
norman	$\frac{-Bz - 2Bz \tan\left(\frac{z}{2}\right)^2 - Bz \tan\left(\frac{z}{2}\right)^4}{\left(1 + \tan\left(\frac{z}{2}\right)^2\right)^2} - \frac{A \ln\left(-B \tan\left(\frac{z}{2}\right)^2 + 2A \tan\left(\frac{z}{2}\right) + B\right)}{2} + \frac{A \ln\left(B \tan\left(\frac{z}{2}\right)^2 + 2A \tan\left(\frac{z}{2}\right) - B\right)}{2}$
risch	$-\frac{Bz A^2}{A^2+B^2} - \frac{B^3 z}{A^2+B^2} + \frac{A^3 \ln\left(e^{2iz} - \frac{iB+A}{-iB+A}\right)}{2A^2+2B^2} + \frac{A \ln\left(e^{2iz} - \frac{iB+A}{-iB+A}\right) B^2}{2A^2+2B^2} - \frac{A^3 \ln\left(e^{2iz} - \frac{-iB+A}{iB+A}\right)}{2(A^2+B^2)} - \frac{A \ln\left(e^{2iz} - \frac{-iB+A}{iB+A}\right)}{2(A^2+B^2)}$

input

```
int((-A^2-B^2)*cos(z)^2/B/(1-(A^2+B^2)*sin(z)^2/B^2),z,method=_RETURNVERBO
SE)
```

output

```
(-A^2-B^2)*B*(1/(A^2+B^2)*arctan(tan(z))+1/2*A/B/(A^2+B^2)*ln(A*tan(z)+B)-
1/2*A/B/(A^2+B^2)*ln(A*tan(z)-B))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 67 vs. $2(16) = 32$.

Time = 0.09 (sec) , antiderivative size = 67, normalized size of antiderivative = 4.19

$$\int \frac{(-A^2 - B^2) \cos^2(z)}{B \left(1 - \frac{(A^2+B^2) \sin^2(z)}{B^2}\right)} dz = -Bz$$

$$- \frac{1}{4} A \log(2AB \cos(z) \sin(z) - (A^2 - B^2) \cos^2(z) + A^2)$$

$$+ \frac{1}{4} A \log(-2AB \cos(z) \sin(z) - (A^2 - B^2) \cos^2(z) + A^2)$$

input `integrate((-A^2-B^2)*cos(z)^2/B/(1-(A^2+B^2)*sin(z)^2/B^2),z, algorithm="fricas")`

output `-B*z - 1/4*A*log(2*A*B*cos(z)*sin(z) - (A^2 - B^2)*cos(z)^2 + A^2) + 1/4*A*log(-2*A*B*cos(z)*sin(z) - (A^2 - B^2)*cos(z)^2 + A^2)`

Sympy [A] (verification not implemented)

Time = 101.86 (sec) , antiderivative size = 202, normalized size of antiderivative = 12.62

$$\int \frac{(-A^2 - B^2) \cos^2(z)}{B \left(1 - \frac{(A^2+B^2) \sin^2(z)}{B^2}\right)} dz$$

$$= \frac{(-A^2 - B^2) \left(\begin{array}{l} z \\ \frac{z \sin^2(z)}{2} + \frac{z \cos^2(z)}{2} + \frac{\sin(z) \cos(z)}{2} \\ \frac{AB \log\left(-\frac{A}{B} + \tan\left(\frac{z}{2}\right) - \frac{\sqrt{A^2+B^2}}{B}\right)}{2A^2+2B^2} + \frac{AB \log\left(-\frac{A}{B} + \tan\left(\frac{z}{2}\right) + \frac{\sqrt{A^2+B^2}}{B}\right)}{2A^2+2B^2} - \frac{AB \log\left(\frac{A}{B} + \tan\left(\frac{z}{2}\right) - \frac{\sqrt{A^2+B^2}}{B}\right)}{2A^2+2B^2} \end{array} \right)}{B}$$

input `integrate((-A**2-B**2)*cos(z)**2/B/(1-(A**2+B**2)*sin(z)**2/B**2),z)`

output

```
(-A**2 - B**2)*Piecewise((z, Eq(A, 0) & Eq(B, 0)), (z*sin(z)**2/2 + z*cos(z)**2/2 + sin(z)*cos(z)/2, Eq(A, I*B) | Eq(A, -I*B)), (A*B*log(-A/B + tan(z/2) - sqrt(A**2 + B**2)/B)/(2*A**2 + 2*B**2) + A*B*log(-A/B + tan(z/2) + sqrt(A**2 + B**2)/B)/(2*A**2 + 2*B**2) - A*B*log(A/B + tan(z/2) - sqrt(A**2 + B**2)/B)/(2*A**2 + 2*B**2) - A*B*log(A/B + tan(z/2) + sqrt(A**2 + B**2)/B)/(2*A**2 + 2*B**2) + 2*B**2*z/(2*A**2 + 2*B**2), True))/B
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 69 vs. $2(16) = 32$.

Time = 0.10 (sec) , antiderivative size = 69, normalized size of antiderivative = 4.31

$$\int \frac{(-A^2 - B^2) \cos^2(z)}{B \left(1 - \frac{(A^2 + B^2) \sin^2(z)}{B^2}\right)} dz$$

$$= - \frac{(A^2 + B^2) \left(\frac{2B^2z}{A^2 + B^2} + \frac{AB \log(A \tan(z) + B)}{A^2 + B^2} - \frac{AB \log(A \tan(z) - B)}{A^2 + B^2} \right)}{2B}$$

input

```
integrate((-A^2-B^2)*cos(z)^2/B/(1-(A^2+B^2)*sin(z)^2/B^2),z, algorithm="maxima")
```

output

```
-1/2*(A^2 + B^2)*(2*B^2*z/(A^2 + B^2) + A*B*log(A*tan(z) + B)/(A^2 + B^2) - A*B*log(A*tan(z) - B)/(A^2 + B^2))/B
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 83 vs. $2(16) = 32$.

Time = 0.13 (sec) , antiderivative size = 83, normalized size of antiderivative = 5.19

$$\int \frac{(-A^2 - B^2) \cos^2(z)}{B \left(1 - \frac{(A^2 + B^2) \sin^2(z)}{B^2}\right)} dz$$

$$= - \frac{\left(\frac{A^3 B \log(|A \tan(z) + B|)}{A^4 + A^2 B^2} - \frac{A^3 B \log(|A \tan(z) - B|)}{A^4 + A^2 B^2} + \frac{2B^2z}{A^2 + B^2} \right) (A^2 + B^2)}{2B}$$

input `integrate((-A^2-B^2)*cos(z)^2/B/(1-(A^2+B^2)*sin(z)^2/B^2),z, algorithm="giac")`

output `-1/2*(A^3*B*log(abs(A*tan(z) + B))/(A^4 + A^2*B^2) - A^3*B*log(abs(A*tan(z) - B))/(A^4 + A^2*B^2) + 2*B^2*z/(A^2 + B^2))*(A^2 + B^2)/B`

Mupad [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 360, normalized size of antiderivative = 22.50

$$\int \frac{(-A^2 - B^2) \cos^2(z)}{B \left(1 - \frac{(A^2+B^2) \sin^2(z)}{B^2}\right)} dz = -A \operatorname{atanh} \left(\frac{2 A^{13} \tan(z)}{2 A^{12} B + 6 A^{10} B^3 + 6 A^8 B^5 + 2 A^6 B^7} \right. \\ \left. + \frac{2 A^7 B^6 \tan(z)}{2 A^{12} B + 6 A^{10} B^3 + 6 A^8 B^5 + 2 A^6 B^7} \right. \\ \left. + \frac{6 A^9 B^4 \tan(z)}{2 A^{12} B + 6 A^{10} B^3 + 6 A^8 B^5 + 2 A^6 B^7} \right. \\ \left. + \frac{6 A^{11} B^2 \tan(z)}{2 A^{12} B + 6 A^{10} B^3 + 6 A^8 B^5 + 2 A^6 B^7} \right) \\ - B \operatorname{atan} \left(\frac{2 A^4 B^9 \tan(z)}{2 A^{10} B^3 + 6 A^8 B^5 + 6 A^6 B^7 + 2 A^4 B^9} \right. \\ \left. + \frac{6 A^6 B^7 \tan(z)}{2 A^{10} B^3 + 6 A^8 B^5 + 6 A^6 B^7 + 2 A^4 B^9} \right. \\ \left. + \frac{6 A^8 B^5 \tan(z)}{2 A^{10} B^3 + 6 A^8 B^5 + 6 A^6 B^7 + 2 A^4 B^9} \right. \\ \left. + \frac{2 A^{10} B^3 \tan(z)}{2 A^{10} B^3 + 6 A^8 B^5 + 6 A^6 B^7 + 2 A^4 B^9} \right)$$

input `int((cos(z)^2*(A^2 + B^2))/(B*((sin(z)^2*(A^2 + B^2))/B^2 - 1)),z)`

output

```
- A*atanh((2*A^13*tan(z))/(2*A^12*B + 2*A^6*B^7 + 6*A^8*B^5 + 6*A^10*B^3)
+ (2*A^7*B^6*tan(z))/(2*A^12*B + 2*A^6*B^7 + 6*A^8*B^5 + 6*A^10*B^3) + (6*
A^9*B^4*tan(z))/(2*A^12*B + 2*A^6*B^7 + 6*A^8*B^5 + 6*A^10*B^3) + (6*A^11*
B^2*tan(z))/(2*A^12*B + 2*A^6*B^7 + 6*A^8*B^5 + 6*A^10*B^3)) - B*atan((2*A
^4*B^9*tan(z))/(2*A^4*B^9 + 6*A^6*B^7 + 6*A^8*B^5 + 2*A^10*B^3) + (6*A^6*B
^7*tan(z))/(2*A^4*B^9 + 6*A^6*B^7 + 6*A^8*B^5 + 2*A^10*B^3) + (6*A^8*B^5*t
an(z))/(2*A^4*B^9 + 6*A^6*B^7 + 6*A^8*B^5 + 2*A^10*B^3) + (2*A^10*B^3*tan(
z))/(2*A^4*B^9 + 6*A^6*B^7 + 6*A^8*B^5 + 2*A^10*B^3))
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 74, normalized size of antiderivative = 4.62

$$\int \frac{(-A^2 - B^2) \cos^2(z)}{B \left(1 - \frac{(A^2 + B^2) \sin^2(z)}{B^2}\right)} dz = \operatorname{atan}\left(\frac{\tan\left(\frac{z}{2}\right) b i}{\sqrt{a^2 + b^2} - a}\right) a i$$

$$- \frac{\log(-\sqrt{a^2 + b^2} + \tan\left(\frac{z}{2}\right) b - a) a}{2}$$

$$+ \frac{\log(\sqrt{a^2 + b^2} + \tan\left(\frac{z}{2}\right) b + a) a}{2} - b z$$

input

```
int((-A^2-B^2)*cos(z)^2/B/(1-(A^2+B^2)*sin(z)^2/B^2),z)
```

output

```
(2*atan((tan(z/2)*b*i)/(sqrt(a**2 + b**2) - a))*a*i - log(- sqrt(a**2 + b
**2) + tan(z/2)*b - a)*a + log(sqrt(a**2 + b**2) + tan(z/2)*b + a)*a - 2*b
*z)/2
```

3.71
$$\int \frac{-A^2 - B^2}{B(1+w^2)^2 \left(1 - \frac{(A^2+B^2)w^2}{B^2(1+w^2)}\right)} dw$$

Optimal result	441
Mathematica [B] (verified)	441
Rubi [B] (verified)	442
Maple [C] (verified)	444
Fricas [A] (verification not implemented)	444
Sympy [C] (verification not implemented)	445
Maxima [B] (verification not implemented)	446
Giac [B] (verification not implemented)	447
Mupad [B] (verification not implemented)	448
Reduce [B] (verification not implemented)	449

Optimal result

Integrand size = 48, antiderivative size = 16

$$\int \frac{-A^2 - B^2}{B(1+w^2)^2 \left(1 - \frac{(A^2+B^2)w^2}{B^2(1+w^2)}\right)} dw = -B \arctan(w) - A \operatorname{arctanh}\left(\frac{Aw}{B}\right)$$

output

`-B*arctan(w)-A*arctanh(A*w/B)`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 35 vs. 2(16) = 32.

Time = 0.01 (sec) , antiderivative size = 35, normalized size of antiderivative = 2.19

$$\int \frac{-A^2 - B^2}{B(1+w^2)^2 \left(1 - \frac{(A^2+B^2)w^2}{B^2(1+w^2)}\right)} dw = -\frac{B(A^2 + B^2) (B \arctan(w) + A \operatorname{arctanh}\left(\frac{Aw}{B}\right))}{A^2 B + B^3}$$

input

`Integrate[(-A^2 - B^2)/(B*(1 + w^2)^2*(1 - ((A^2 + B^2)*w^2)/(B^2*(1 + w^2))))],w]`

output $-\left(\frac{B(A^2 + B^2)(B \operatorname{ArcTan}[w] + A \operatorname{ArcTanh}[(A*w)/B])}{A^2*B + B^3}\right)$

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 44 vs. $2(16) = 32$.

Time = 0.31 (sec) , antiderivative size = 44, normalized size of antiderivative = 2.75, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {27, 7239, 27, 303, 216, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{-A^2 - B^2}{B(w^2 + 1)^2 \left(1 - \frac{w^2(A^2 + B^2)}{B^2(w^2 + 1)}\right)} dw \\
 & \quad \downarrow 27 \\
 & \frac{(A^2 + B^2) \int \frac{1}{(w^2 + 1)^2 \left(1 - \frac{(A^2 + B^2)w^2}{B^2(w^2 + 1)}\right)} dw}{B} \\
 & \quad \downarrow 7239 \\
 & -\frac{(A^2 + B^2) \int \frac{B^2}{(w^2 + 1)(B^2 - A^2 w^2)} dw}{B} \\
 & \quad \downarrow 27 \\
 & -B(A^2 + B^2) \int \frac{1}{(w^2 + 1)(B^2 - A^2 w^2)} dw \\
 & \quad \downarrow 303 \\
 & -B(A^2 + B^2) \left(\frac{A^2 \int \frac{1}{B^2 - A^2 w^2} dw}{A^2 + B^2} + \frac{\int \frac{1}{w^2 + 1} dw}{A^2 + B^2} \right) \\
 & \quad \downarrow 216 \\
 & -B(A^2 + B^2) \left(\frac{A^2 \int \frac{1}{B^2 - A^2 w^2} dw}{A^2 + B^2} + \frac{\arctan(w)}{A^2 + B^2} \right) \\
 & \quad \downarrow 221
 \end{aligned}$$

$$-B(A^2 + B^2) \left(\frac{\arctan(w)}{A^2 + B^2} + \frac{A \operatorname{arctanh}\left(\frac{Aw}{B}\right)}{B(A^2 + B^2)} \right)$$

input

```
Int[(-A^2 - B^2)/(B*(1 + w^2)^2*(1 - ((A^2 + B^2)*w^2)/(B^2*(1 + w^2))),w
]
```

output

```
-(B*(A^2 + B^2)*(ArcTan[w]/(A^2 + B^2) + (A*ArcTanh[(A*w)/B])/(B*(A^2 + B^2))))
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 216

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

rule 221

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

rule 303

```
Int[1/(((a_) + (b_.)*(x_)^2)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x^2), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]
```

rule 7239

```
Int[u_, x_Symbol] := With[{v = SimplifyIntegrand[u, x]}, Int[v, x] /; SimplifyIntegrandQ[v, u, x]]
```

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 70, normalized size of antiderivative = 4.38

method	result
parallelrisch	$\frac{(-A^2 - B^2)(-iB^2 \ln(-i+w) + iB^2 \ln(w+i) - AB \ln(Aw-B) + AB \ln(Aw+B))}{2B(A^2 + B^2)}$
default	$(-A^2 - B^2) B \left(-\frac{A \ln(Aw-B)}{2B(A^2 + B^2)} + \frac{A \ln(Aw+B)}{2B(A^2 + B^2)} + \frac{\arctan(w)}{A^2 + B^2} \right)$
risch	$-\frac{A^3 \ln(-Aw-B)}{2(A^2 + B^2)} - \frac{A \ln(-Aw-B)B^2}{2(A^2 + B^2)} + \frac{A^3 \ln(-Aw+B)}{2A^2 + 2B^2} + \frac{A \ln(-Aw+B)B^2}{2A^2 + 2B^2} - \left(\sum_{R=\text{RootOf}((A^4 + 2A^2B^2 + B^4))} \right)$

input `int((-A^2-B^2)/B/(w^2+1)^2/(1-(A^2+B^2)*w^2/B^2/(w^2+1)),w,method=_RETURNV
ERBOSE)`

output `1/2*(-A^2-B^2)/B*(-I*B^2*ln(-I+w)+I*B^2*ln(w+I)-A*B*ln(A*w-B)+A*B*ln(A*w+B
))/ (A^2+B^2)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.62

$$\int \frac{-A^2 - B^2}{B(1 + w^2)^2 \left(1 - \frac{(A^2 + B^2)w^2}{B^2(1 + w^2)}\right)} dw = -B \arctan(w) - \frac{1}{2} A \log(Aw + B) + \frac{1}{2} A \log(Aw - B)$$

input `integrate((-A^2-B^2)/B/(w^2+1)^2/(1-(A^2+B^2)*w^2/B^2/(w^2+1)),w, algorithm
m="fricas")`

output `-B*arctan(w) - 1/2*A*log(A*w + B) + 1/2*A*log(A*w - B)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.89 (sec) , antiderivative size = 422, normalized size of antiderivative = 26.38

$$\int \frac{-A^2 - B^2}{B(1+w^2)^2 \left(1 - \frac{(A^2+B^2)w^2}{B^2(1+w^2)}\right)} dw$$

$$= (A^2B + B^3) \left(\begin{aligned} & - \frac{A \log \left(w + \frac{-\frac{A^9}{B(A^2+B^2)^3} - \frac{A^7B}{(A^2+B^2)^3} + \frac{A^5B^3}{(A^2+B^2)^3} + \frac{A^5}{B(A^2+B^2)} + \frac{A^3B^5}{(A^2+B^2)^3} + \frac{AB^3}{A^2+B^2}}{A^2} \right)}{2B(A^2+B^2)} \\ & + \frac{A \log \left(w + \frac{\frac{A^9}{B(A^2+B^2)^3} + \frac{A^7B}{(A^2+B^2)^3} - \frac{A^5B^3}{(A^2+B^2)^3} - \frac{A^5}{B(A^2+B^2)} - \frac{A^3B^5}{(A^2+B^2)^3} - \frac{AB^3}{A^2+B^2}}{A^2} \right)}{2B(A^2+B^2)} \\ & + \frac{i \log \left(w + \frac{-\frac{iA^6B^2}{(A^2+B^2)^3} - \frac{iA^4B^4}{(A^2+B^2)^3} - \frac{iA^4}{A^2+B^2} + \frac{iA^2B^6}{(A^2+B^2)^3} + \frac{iB^8}{(A^2+B^2)^3} - \frac{iB^4}{A^2+B^2}}{A^2} \right)}{2(A^2+B^2)} \\ & - \frac{i \log \left(w + \frac{\frac{iA^6B^2}{(A^2+B^2)^3} + \frac{iA^4B^4}{(A^2+B^2)^3} + \frac{iA^4}{A^2+B^2} - \frac{iA^2B^6}{(A^2+B^2)^3} - \frac{iB^8}{(A^2+B^2)^3} + \frac{iB^4}{A^2+B^2}}{A^2} \right)}{2(A^2+B^2)} \end{aligned} \right)$$

input `integrate((-A**2-B**2)/B/(w**2+1)**2/(1-(A**2+B**2)*w**2/B**2/(w**2+1)),w)`

output

```
(A**2*B + B**3)*(-A*log(w + (-A**9/(B*(A**2 + B**2)**3) - A**7*B/(A**2 + B**2)**3 + A**5*B**3/(A**2 + B**2)**3 + A**5/(B*(A**2 + B**2))) + A**3*B**5/(A**2 + B**2)**3 + A*B**3/(A**2 + B**2))/A**2)/(2*B*(A**2 + B**2)) + A*log(w + (A**9/(B*(A**2 + B**2)**3) + A**7*B/(A**2 + B**2)**3 - A**5*B**3/(A**2 + B**2)**3 - A**5/(B*(A**2 + B**2))) - A**3*B**5/(A**2 + B**2)**3 - A*B**3/(A**2 + B**2))/A**2)/(2*B*(A**2 + B**2)) + I*log(w + (-I*A**6*B**2/(A**2 + B**2)**3 - I*A**4*B**4/(A**2 + B**2)**3 - I*A**4/(A**2 + B**2) + I*A**2*B**6/(A**2 + B**2)**3 + I*B**8/(A**2 + B**2)**3 - I*B**4/(A**2 + B**2)))/A**2)/(2*(A**2 + B**2)) - I*log(w + (I*A**6*B**2/(A**2 + B**2)**3 + I*A**4*B**4/(A**2 + B**2)**3 + I*A**4/(A**2 + B**2) - I*A**2*B**6/(A**2 + B**2)**3 - I*B**8/(A**2 + B**2)**3 + I*B**4/(A**2 + B**2)))/A**2)/(2*(A**2 + B**2)))
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 68 vs. $2(16) = 32$.

Time = 0.11 (sec) , antiderivative size = 68, normalized size of antiderivative = 4.25

$$\int \frac{-A^2 - B^2}{B(1+w^2)^2 \left(1 - \frac{(A^2+B^2)w^2}{B^2(1+w^2)}\right)} dw$$

$$= -\frac{(A^2 + B^2) \left(\frac{2B^2 \arctan(w)}{A^2+B^2} + \frac{AB \log(Aw+B)}{A^2+B^2} - \frac{AB \log(Aw-B)}{A^2+B^2} \right)}{2B}$$

input

```
integrate((-A^2-B^2)/B/(w^2+1)^2/(1-(A^2+B^2)*w^2/B^2/(w^2+1)),w, algorithm m="maxima")
```

output

```
-1/2*(A^2 + B^2)*(2*B^2*arctan(w)/(A^2 + B^2) + A*B*log(A*w + B)/(A^2 + B^2) - A*B*log(A*w - B)/(A^2 + B^2))/B
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 82 vs. $2(16) = 32$.

Time = 0.12 (sec) , antiderivative size = 82, normalized size of antiderivative = 5.12

$$\int \frac{-A^2 - B^2}{B(1+w^2)^2 \left(1 - \frac{(A^2+B^2)w^2}{B^2(1+w^2)}\right)} dw$$

$$= -\frac{\left(\frac{A^3 B \log(|Aw+B|)}{A^4+A^2 B^2} - \frac{A^3 B \log(|Aw-B|)}{A^4+A^2 B^2} + \frac{2 B^2 \arctan(w)}{A^2+B^2}\right)(A^2 + B^2)}{2 B}$$

input

```
integrate((-A^2-B^2)/B/(w^2+1)^2/(1-(A^2+B^2)*w^2/B^2/(w^2+1)),w, algorithm
m="giac")
```

output

```
-1/2*(A^3*B*log(abs(A*w + B))/(A^4 + A^2*B^2) - A^3*B*log(abs(A*w - B))/(A
^4 + A^2*B^2) + 2*B^2*arctan(w)/(A^2 + B^2))*(A^2 + B^2)/B
```


Mupad [B] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 352, normalized size of antiderivative = 22.00

$$\begin{aligned}
& \int \frac{-A^2 - B^2}{B(1+w^2)^2 \left(1 - \frac{(A^2+B^2)w^2}{B^2(1+w^2)}\right)} dw \\
&= -A \operatorname{atanh} \left(\frac{2A^{13}w}{2A^{12}B + 6A^{10}B^3 + 6A^8B^5 + 2A^6B^7} \right. \\
&\quad \left. + \frac{2A^7B^6w}{2A^{12}B + 6A^{10}B^3 + 6A^8B^5 + 2A^6B^7} \right. \\
&\quad \left. + \frac{6A^9B^4w}{2A^{12}B + 6A^{10}B^3 + 6A^8B^5 + 2A^6B^7} \right. \\
&\quad \left. + \frac{6A^{11}B^2w}{2A^{12}B + 6A^{10}B^3 + 6A^8B^5 + 2A^6B^7} \right) \\
&\quad - B \operatorname{atan} \left(\frac{2A^4B^9w}{2A^{10}B^3 + 6A^8B^5 + 6A^6B^7 + 2A^4B^9} \right. \\
&\quad \left. + \frac{6A^6B^7w}{2A^{10}B^3 + 6A^8B^5 + 6A^6B^7 + 2A^4B^9} \right. \\
&\quad \left. + \frac{6A^8B^5w}{2A^{10}B^3 + 6A^8B^5 + 6A^6B^7 + 2A^4B^9} \right. \\
&\quad \left. + \frac{2A^{10}B^3w}{2A^{10}B^3 + 6A^8B^5 + 6A^6B^7 + 2A^4B^9} \right)
\end{aligned}$$

input `int((A^2 + B^2)/(B*(w^2 + 1)^2*((w^2*(A^2 + B^2))/(B^2*(w^2 + 1)) - 1)),w)`

output `- A*atanh((2*A^13*w)/(2*A^12*B + 2*A^6*B^7 + 6*A^8*B^5 + 6*A^10*B^3) + (2*A^7*B^6*w)/(2*A^12*B + 2*A^6*B^7 + 6*A^8*B^5 + 6*A^10*B^3) + (6*A^9*B^4*w)/(2*A^12*B + 2*A^6*B^7 + 6*A^8*B^5 + 6*A^10*B^3) + (6*A^11*B^2*w)/(2*A^12*B + 2*A^6*B^7 + 6*A^8*B^5 + 6*A^10*B^3)) - B*atan((2*A^4*B^9*w)/(2*A^4*B^9 + 6*A^6*B^7 + 6*A^8*B^5 + 2*A^10*B^3) + (6*A^6*B^7*w)/(2*A^4*B^9 + 6*A^6*B^7 + 6*A^8*B^5 + 2*A^10*B^3) + (6*A^8*B^5*w)/(2*A^4*B^9 + 6*A^6*B^7 + 6*A^8*B^5 + 2*A^10*B^3) + (2*A^10*B^3*w)/(2*A^4*B^9 + 6*A^6*B^7 + 6*A^8*B^5 + 2*A^10*B^3))`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 33, normalized size of antiderivative = 2.06

$$\int \frac{-A^2 - B^2}{B(1+w^2)^2 \left(1 - \frac{(A^2+B^2)w^2}{B^2(1+w^2)}\right)} dw = -atan(w)b + \frac{\log(a^2w - ab)a}{2} - \frac{\log(a^2w + ab)a}{2}$$

input `int((-A^2-B^2)/B/(w^2+1)^2/(1-(A^2+B^2)*w^2/B^2/(w^2+1)),w)`

output `(- 2*atan(w)*b + log(a**2*w - a*b)*a - log(a**2*w + a*b)*a)/2`

3.72
$$\int -\frac{B(A^2+B^2)}{(1+w^2)(B^2-A^2w^2)} dw$$

Optimal result	450
Mathematica [B] (verified)	450
Rubi [B] (verified)	451
Maple [A] (verified)	452
Fricas [A] (verification not implemented)	453
Sympy [C] (verification not implemented)	453
Maxima [B] (verification not implemented)	454
Giac [B] (verification not implemented)	455
Mupad [B] (verification not implemented)	455
Reduce [B] (verification not implemented)	456

Optimal result

Integrand size = 31, antiderivative size = 16

$$\int -\frac{B(A^2 + B^2)}{(1 + w^2)(B^2 - A^2w^2)} dw = -B \arctan(w) - A \operatorname{arctanh}\left(\frac{Aw}{B}\right)$$

output

`-B*arctan(w)-A*arctanh(A*w/B)`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 35 vs. 2(16) = 32.

Time = 0.01 (sec) , antiderivative size = 35, normalized size of antiderivative = 2.19

$$\int -\frac{B(A^2 + B^2)}{(1 + w^2)(B^2 - A^2w^2)} dw = -\frac{B(A^2 + B^2) (B \arctan(w) + A \operatorname{arctanh}\left(\frac{Aw}{B}\right))}{A^2B + B^3}$$

input

`Integrate[-((B*(A^2 + B^2))/((1 + w^2)*(B^2 - A^2*w^2))),w]`

output

`-((B*(A^2 + B^2)*(B*ArcTan[w] + A*ArcTanh[(A*w)/B]))/(A^2*B + B^3))`

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 44 vs. $2(16) = 32$.

Time = 0.17 (sec) , antiderivative size = 44, normalized size of antiderivative = 2.75, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.161$, Rules used = {25, 27, 303, 216, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int -\frac{B(A^2 + B^2)}{(w^2 + 1)(B^2 - A^2w^2)} dw \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{B(A^2 + B^2)}{(w^2 + 1)(B^2 - A^2w^2)} dw \\
 & \quad \downarrow \text{27} \\
 & -B(A^2 + B^2) \int \frac{1}{(w^2 + 1)(B^2 - A^2w^2)} dw \\
 & \quad \downarrow \text{303} \\
 & -B(A^2 + B^2) \left(\frac{A^2 \int \frac{1}{B^2 - A^2w^2} dw}{A^2 + B^2} + \frac{\int \frac{1}{w^2 + 1} dw}{A^2 + B^2} \right) \\
 & \quad \downarrow \text{216} \\
 & -B(A^2 + B^2) \left(\frac{A^2 \int \frac{1}{B^2 - A^2w^2} dw}{A^2 + B^2} + \frac{\arctan(w)}{A^2 + B^2} \right) \\
 & \quad \downarrow \text{221} \\
 & -B(A^2 + B^2) \left(\frac{\arctan(w)}{A^2 + B^2} + \frac{A \operatorname{arctanh}\left(\frac{Aw}{B}\right)}{B(A^2 + B^2)} \right)
 \end{aligned}$$

input `Int[-((B*(A^2 + B^2))/((1 + w^2)*(B^2 - A^2*w^2))),w]`

output `-(B*(A^2 + B^2)*(ArcTan[w]/(A^2 + B^2) + (A*ArcTanh[(A*w)/B])/(B*(A^2 + B^2))))`

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27 $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ ; FreeQ}[\text{b}, \text{x}]$
- rule 216 $\text{Int}[(\text{a}_) + (\text{b}_.)*(\text{x}_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2]*\text{Rt}[\text{b}, 2]))*\text{ArcTan}[\text{Rt}[\text{b}, 2]*(\text{x}/\text{Rt}[\text{a}, 2])], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ \text{PosQ}[\text{a}/\text{b}] \ \&\& \ (\text{GtQ}[\text{a}, 0] \ || \ \text{GtQ}[\text{b}, 0])$
- rule 221 $\text{Int}[(\text{a}_) + (\text{b}_.)*(\text{x}_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-\text{a}/\text{b}, 2]/\text{a})*\text{ArcTanh}[\text{x}/\text{Rt}[-\text{a}/\text{b}, 2]], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ \text{NegQ}[\text{a}/\text{b}]$
- rule 303 $\text{Int}[1/((\text{a}_) + (\text{b}_.)*(\text{x}_)^2)*((\text{c}_) + (\text{d}_.)*(\text{x}_)^2)), \text{x_Symbol}] \rightarrow \text{Simp}[\text{b}/(\text{b}*c - \text{a}*d) \quad \text{Int}[1/(\text{a} + \text{b}*x^2), \text{x}], \text{x}] - \text{Simp}[\text{d}/(\text{b}*c - \text{a}*d) \quad \text{Int}[1/(\text{c} + \text{d}*x^2), \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0]$

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.81

method	result	size
risch	$-B \arctan(w) + \frac{A \ln(-Aw+B)}{2} - \frac{A \ln(-Aw-B)}{2}$	29
parallelrisch	$\frac{i \ln(-i+w)B}{2} - \frac{i \ln(w+i)B}{2} + \frac{A \ln(Aw-B)}{2} - \frac{A \ln(Aw+B)}{2}$	40
default	$-(A^2 + B^2) B \left(-\frac{A \ln(Aw-B)}{2B(A^2+B^2)} + \frac{A \ln(Aw+B)}{2B(A^2+B^2)} + \frac{\arctan(w)}{A^2+B^2} \right)$	68

input $\text{int}(-B*(A^2+B^2)/(w^2+1)/(-A^2*w^2+B^2), w, \text{method}=_RETURNVERBOSE)$

output $-B*\arctan(w)+1/2*A*\ln(-A*w+B)-1/2*A*\ln(-A*w-B)$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.62

$$\int -\frac{B(A^2 + B^2)}{(1 + w^2)(B^2 - A^2w^2)} dw = -B \arctan(w) - \frac{1}{2} A \log(Aw + B) + \frac{1}{2} A \log(Aw - B)$$

input `integrate(-B*(A^2+B^2)/(w^2+1)/(-A^2*w^2+B^2),w, algorithm="fricas")`

output `-B*arctan(w) - 1/2*A*log(A*w + B) + 1/2*A*log(A*w - B)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.89 (sec) , antiderivative size = 422, normalized size of antiderivative = 26.38

$$\int -\frac{B(A^2 + B^2)}{(1 + w^2)(B^2 - A^2w^2)} dw$$

$$= (A^2B + B^3) \left(\begin{aligned} & \frac{A \log \left(w + \frac{-\frac{A^9}{B(A^2+B^2)^3} - \frac{A^7B}{(A^2+B^2)^3} + \frac{A^5B^3}{(A^2+B^2)^3} + \frac{A^5}{B(A^2+B^2)} + \frac{A^3B^5}{(A^2+B^2)^3} + \frac{AB^3}{A^2+B^2}}{A^2} \right)}{2B(A^2 + B^2)} \\ & + \frac{A \log \left(w + \frac{\frac{A^9}{B(A^2+B^2)^3} + \frac{A^7B}{(A^2+B^2)^3} - \frac{A^5B^3}{(A^2+B^2)^3} - \frac{A^5}{B(A^2+B^2)} - \frac{A^3B^5}{(A^2+B^2)^3} - \frac{AB^3}{A^2+B^2}}{A^2} \right)}{2B(A^2 + B^2)} \\ & + \frac{i \log \left(w + \frac{-\frac{iA^6B^2}{(A^2+B^2)^3} - \frac{iA^4B^4}{(A^2+B^2)^3} - \frac{iA^4}{A^2+B^2} + \frac{iA^2B^6}{(A^2+B^2)^3} + \frac{iB^8}{(A^2+B^2)^3} - \frac{iB^4}{A^2+B^2}}{A^2} \right)}{2(A^2 + B^2)} \\ & - \frac{i \log \left(w + \frac{\frac{iA^6B^2}{(A^2+B^2)^3} + \frac{iA^4B^4}{(A^2+B^2)^3} + \frac{iA^4}{A^2+B^2} - \frac{iA^2B^6}{(A^2+B^2)^3} - \frac{iB^8}{(A^2+B^2)^3} + \frac{iB^4}{A^2+B^2}}{A^2} \right)}{2(A^2 + B^2)} \end{aligned} \right)$$

input `integrate(-B*(A**2+B**2)/(w**2+1)/(-A**2*w**2+B**2),w)`

output `(A**2*B + B**3)*(-A*log(w + (-A**9/(B*(A**2 + B**2)**3) - A**7*B/(A**2 + B**2)**3 + A**5*B**3/(A**2 + B**2)**3 + A**5/(B*(A**2 + B**2)) + A**3*B**5/(A**2 + B**2)**3 + A*B**3/(A**2 + B**2))/A**2)/(2*B*(A**2 + B**2)) + A*log(w + (A**9/(B*(A**2 + B**2)**3) + A**7*B/(A**2 + B**2)**3 - A**5*B**3/(A**2 + B**2)**3 - A**5/(B*(A**2 + B**2)) - A**3*B**5/(A**2 + B**2)**3 - A*B**3/(A**2 + B**2))/A**2)/(2*B*(A**2 + B**2)) + I*log(w + (-I*A**6*B**2/(A**2 + B**2)**3 - I*A**4*B**4/(A**2 + B**2)**3 - I*A**4/(A**2 + B**2) + I*A**2*B**6/(A**2 + B**2)**3 + I*B**8/(A**2 + B**2)**3 - I*B**4/(A**2 + B**2))/A**2)/(2*(A**2 + B**2)) - I*log(w + (I*A**6*B**2/(A**2 + B**2)**3 + I*A**4*B**4/(A**2 + B**2)**3 + I*A**4/(A**2 + B**2) - I*A**2*B**6/(A**2 + B**2)**3 - I*B**8/(A**2 + B**2)**3 + I*B**4/(A**2 + B**2))/A**2)/(2*(A**2 + B**2)))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 65 vs. $2(16) = 32$.

Time = 0.10 (sec) , antiderivative size = 65, normalized size of antiderivative = 4.06

$$\int -\frac{B(A^2 + B^2)}{(1 + w^2)(B^2 - A^2w^2)} dw$$

$$= -\frac{1}{2} (A^2 + B^2) B \left(\frac{A \log(Aw + B)}{A^2B + B^3} - \frac{A \log(Aw - B)}{A^2B + B^3} + \frac{2 \arctan(w)}{A^2 + B^2} \right)$$

input `integrate(-B*(A^2+B^2)/(w^2+1)/(-A^2*w^2+B^2),w, algorithm="maxima")`

output `-1/2*(A^2 + B^2)*B*(A*log(A*w + B)/(A^2*B + B^3) - A*log(A*w - B)/(A^2*B + B^3) + 2*arctan(w)/(A^2 + B^2))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 79 vs. $2(16) = 32$.

Time = 0.12 (sec) , antiderivative size = 79, normalized size of antiderivative = 4.94

$$\int -\frac{B(A^2 + B^2)}{(1 + w^2)(B^2 - A^2w^2)} dw$$

$$= -\frac{1}{2} \left(\frac{A^3 \log(|Aw + B|)}{A^4B + A^2B^3} - \frac{A^3 \log(|Aw - B|)}{A^4B + A^2B^3} + \frac{2 \arctan(w)}{A^2 + B^2} \right) (A^2 + B^2)B$$

input `integrate(-B*(A^2+B^2)/(w^2+1)/(-A^2*w^2+B^2),w, algorithm="giac")`

output `-1/2*(A^3*log(abs(A*w + B))/(A^4*B + A^2*B^3) - A^3*log(abs(A*w - B))/(A^4*B + A^2*B^3) + 2*arctan(w)/(A^2 + B^2))*(A^2 + B^2)*B`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 352, normalized size of antiderivative = 22.00

$$\int -\frac{B(A^2 + B^2)}{(1 + w^2)(B^2 - A^2w^2)} dw = -A \operatorname{atanh} \left(\frac{2A^{13}w}{2A^{12}B + 6A^{10}B^3 + 6A^8B^5 + 2A^6B^7} + \frac{2A^7B^6w}{2A^{12}B + 6A^{10}B^3 + 6A^8B^5 + 2A^6B^7} + \frac{6A^9B^4w}{2A^{12}B + 6A^{10}B^3 + 6A^8B^5 + 2A^6B^7} + \frac{6A^{11}B^2w}{2A^{12}B + 6A^{10}B^3 + 6A^8B^5 + 2A^6B^7} \right) - B \operatorname{atan} \left(\frac{2A^4B^9w}{2A^{10}B^3 + 6A^8B^5 + 6A^6B^7 + 2A^4B^9} + \frac{6A^6B^7w}{2A^{10}B^3 + 6A^8B^5 + 6A^6B^7 + 2A^4B^9} + \frac{6A^8B^5w}{2A^{10}B^3 + 6A^8B^5 + 6A^6B^7 + 2A^4B^9} + \frac{2A^{10}B^3w}{2A^{10}B^3 + 6A^8B^5 + 6A^6B^7 + 2A^4B^9} \right)$$

input `int(-(B*(A^2 + B^2))/((w^2 + 1)*(B^2 - A^2*w^2)),w)`

output

```
- A*atanh((2*A^13*w)/(2*A^12*B + 2*A^6*B^7 + 6*A^8*B^5 + 6*A^10*B^3) + (2*
A^7*B^6*w)/(2*A^12*B + 2*A^6*B^7 + 6*A^8*B^5 + 6*A^10*B^3) + (6*A^9*B^4*w)
/(2*A^12*B + 2*A^6*B^7 + 6*A^8*B^5 + 6*A^10*B^3) + (6*A^11*B^2*w)/(2*A^12*
B + 2*A^6*B^7 + 6*A^8*B^5 + 6*A^10*B^3)) - B*atan((2*A^4*B^9*w)/(2*A^4*B^9
+ 6*A^6*B^7 + 6*A^8*B^5 + 2*A^10*B^3) + (6*A^6*B^7*w)/(2*A^4*B^9 + 6*A^6*
B^7 + 6*A^8*B^5 + 2*A^10*B^3) + (6*A^8*B^5*w)/(2*A^4*B^9 + 6*A^6*B^7 + 6*A
^8*B^5 + 2*A^10*B^3) + (2*A^10*B^3*w)/(2*A^4*B^9 + 6*A^6*B^7 + 6*A^8*B^5 +
2*A^10*B^3))
```

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 33, normalized size of antiderivative = 2.06

$$\int -\frac{B(A^2 + B^2)}{(1 + w^2)(B^2 - A^2w^2)} dw = -atan(w)b + \frac{\log(a^2w - ab)a}{2} - \frac{\log(a^2w + ab)a}{2}$$

input

```
int(-B*(A^2+B^2)/(w^2+1)/(-A^2*w^2+B^2),w)
```

output

```
( - 2*atan(w)*b + log(a**2*w - a*b)*a - log(a**2*w + a*b)*a)/2
```

3.73 $\int \frac{x^4}{(1-x^2)^{5/2}} dx$

Optimal result	457
Mathematica [A] (verified)	457
Rubi [A] (verified)	458
Maple [A] (verified)	459
Fricas [B] (verification not implemented)	459
Sympy [B] (verification not implemented)	460
Maxima [A] (verification not implemented)	460
Giac [A] (verification not implemented)	461
Mupad [B] (verification not implemented)	461
Reduce [B] (verification not implemented)	461

Optimal result

Integrand size = 15, antiderivative size = 35

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{x^3}{3(1-x^2)^{3/2}} - \frac{x}{\sqrt{1-x^2}} + \arcsin(x)$$

output

```
1/3*x^3/(-x^2+1)^(3/2)+arcsin(x)-x/(-x^2+1)^(1/2)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.26

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{x(-3+4x^2)}{3(1-x^2)^{3/2}} + 2 \arctan\left(\frac{x}{-1+\sqrt{1-x^2}}\right)$$

input

```
Integrate[x^4/(1-x^2)^(5/2),x]
```

output

```
(x*(-3+4*x^2))/(3*(1-x^2)^(3/2))+2*ArcTan[x/(-1+Sqrt[1-x^2])]
```

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {252, 252, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx$$

$$\downarrow 252$$

$$\frac{x^3}{3(1-x^2)^{3/2}} - \int \frac{x^2}{(1-x^2)^{3/2}} dx$$

$$\downarrow 252$$

$$\int \frac{1}{\sqrt{1-x^2}} dx - \frac{x}{\sqrt{1-x^2}} + \frac{x^3}{3(1-x^2)^{3/2}}$$

$$\downarrow 223$$

$$\arcsin(x) - \frac{x}{\sqrt{1-x^2}} + \frac{x^3}{3(1-x^2)^{3/2}}$$

input `Int[x^4/(1 - x^2)^(5/2),x]`

output `x^3/(3*(1 - x^2)^(3/2)) - x/Sqrt[1 - x^2] + ArcSin[x]`

Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 252

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(2*b*(p + 1))), x] - Simp[c^2*(m - 1)/(2*b*(p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && LtQ[p, -1] && GtQ[m, 1] && !ILtQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]
```

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.86

method	result	size
default	$\frac{x^3}{3(-x^2+1)^{\frac{3}{2}}} + \arcsin(x) - \frac{x}{\sqrt{-x^2+1}}$	30
risch	$-\frac{x(4x^2-3)}{3(x^2-1)\sqrt{-x^2+1}} + \arcsin(x)$	30
meijerg	$-\frac{2i \left(-\frac{i\sqrt{\pi}x(-20x^2+15)}{10(-x^2+1)^{\frac{3}{2}}} + \frac{3i\sqrt{\pi}\arcsin(x)}{2} \right)}{3\sqrt{\pi}}$	39
pseudoelliptic	$\frac{(x^2-1)\sqrt{-x^2+1}\arctan\left(\frac{\sqrt{-x^2+1}}{x}\right) + \frac{4x^3}{3} - x}{(-x^2+1)^{\frac{3}{2}}}$	49
trager	$\frac{(4x^2-3)x\sqrt{-x^2+1}}{3(x^2-1)^2} + \text{RootOf}(_Z^2 + 1) \ln(\text{RootOf}(_Z^2 + 1)\sqrt{-x^2+1} + x)$	54

input `int(x^4/(-x^2+1)^(5/2),x,method=_RETURNVERBOSE)`output `1/3*x^3/(-x^2+1)^(3/2)+arcsin(x)-1/(-x^2+1)^(1/2)*x`**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 63 vs. 2(29) = 58.

Time = 0.07 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.80

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = -\frac{6(x^4 - 2x^2 + 1) \arctan\left(\frac{\sqrt{-x^2+1}-1}{x}\right) - (4x^3 - 3x)\sqrt{-x^2+1}}{3(x^4 - 2x^2 + 1)}$$

input `integrate(x^4/(-x^2+1)^(5/2),x, algorithm="fricas")`

output `-1/3*(6*(x^4 - 2*x^2 + 1)*arctan((sqrt(-x^2 + 1) - 1)/x) - (4*x^3 - 3*x)*sqrt(-x^2 + 1))/(x^4 - 2*x^2 + 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 105 vs. $2(26) = 52$.

Time = 0.35 (sec) , antiderivative size = 105, normalized size of antiderivative = 3.00

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{3x^4 \operatorname{asin}(x)}{3x^4 - 6x^2 + 3} + \frac{4x^3 \sqrt{1-x^2}}{3x^4 - 6x^2 + 3} - \frac{6x^2 \operatorname{asin}(x)}{3x^4 - 6x^2 + 3} - \frac{3x \sqrt{1-x^2}}{3x^4 - 6x^2 + 3} + \frac{3 \operatorname{asin}(x)}{3x^4 - 6x^2 + 3}$$

input `integrate(x**4/(-x**2+1)**(5/2),x)`

output `3*x**4*asin(x)/(3*x**4 - 6*x**2 + 3) + 4*x**3*sqrt(1 - x**2)/(3*x**4 - 6*x**2 + 3) - 6*x**2*asin(x)/(3*x**4 - 6*x**2 + 3) - 3*x*sqrt(1 - x**2)/(3*x**4 - 6*x**2 + 3) + 3*asin(x)/(3*x**4 - 6*x**2 + 3)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.26

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{1}{3} x \left(\frac{3x^2}{(-x^2+1)^{3/2}} - \frac{2}{(-x^2+1)^{3/2}} \right) - \frac{x}{3\sqrt{-x^2+1}} + \operatorname{arcsin}(x)$$

input `integrate(x^4/(-x^2+1)^(5/2),x, algorithm="maxima")`

output `1/3*x*(3*x^2/(-x^2 + 1)^(3/2) - 2/(-x^2 + 1)^(3/2)) - 1/3*x/sqrt(-x^2 + 1) + arcsin(x)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.83

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{(4x^2-3)\sqrt{-x^2+1}x}{3(x^2-1)^2} + \arcsin(x)$$

input `integrate(x^4/(-x^2+1)^(5/2),x, algorithm="giac")`output `1/3*(4*x^2 - 3)*sqrt(-x^2 + 1)*x/(x^2 - 1)^2 + arcsin(x)`**Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 91, normalized size of antiderivative = 2.60

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \arcsin(x) + \frac{3\sqrt{1-x^2}}{4(x-1)} + \frac{3\sqrt{1-x^2}}{4(x+1)} - \sqrt{1-x^2} \left(\frac{1}{12(x-1)} - \frac{1}{12(x-1)^2} \right) - \sqrt{1-x^2} \left(\frac{1}{12(x+1)} + \frac{1}{12(x+1)^2} \right)$$

input `int(x^4/(1 - x^2)^(5/2),x)`output `asin(x) + (3*(1 - x^2)^(1/2))/(4*(x - 1)) + (3*(1 - x^2)^(1/2))/(4*(x + 1)) - (1 - x^2)^(1/2)*(1/(12*(x - 1)) - 1/(12*(x - 1)^2)) - (1 - x^2)^(1/2)*(1/(12*(x + 1)) + 1/(12*(x + 1)^2))`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.57

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{3\sqrt{-x^2+1} \arcsin(x) x^2 - 3\sqrt{-x^2+1} \arcsin(x) - 4x^3 + 3x}{3\sqrt{-x^2+1} (x^2 - 1)}$$

input `int(x^4/(-x^2+1)^(5/2),x)`

output
$$\frac{(3\sqrt{-x^2 + 1})\operatorname{asin}(x)x^2 - 3\sqrt{-x^2 + 1}\operatorname{asin}(x) - 4x^3 + 3x}{(3\sqrt{-x^2 + 1})(x^2 - 1)}$$

3.74 $\int \tan^4(y) dy$

Optimal result	463
Mathematica [A] (verified)	463
Rubi [A] (verified)	464
Maple [A] (verified)	465
Fricas [B] (verification not implemented)	466
Sympy [A] (verification not implemented)	466
Maxima [A] (verification not implemented)	466
Giac [A] (verification not implemented)	467
Mupad [B] (verification not implemented)	467
Reduce [B] (verification not implemented)	467

Optimal result

Integrand size = 4, antiderivative size = 14

$$\int \tan^4(y) dy = y - \tan(y) + \frac{\tan^3(y)}{3}$$

output `y-tan(y)+1/3*tan(y)^3`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \tan^4(y) dy = \arctan(\tan(y)) - \tan(y) + \frac{\tan^3(y)}{3}$$

input `Integrate[Tan[y]^4,y]`

output `ArcTan[Tan[y]] - Tan[y] + Tan[y]^3/3`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$, Rules used = {3042, 3954, 3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^4(y) dy \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(y)^4 dy \\
 & \quad \downarrow \text{3954} \\
 & \frac{\tan^3(y)}{3} - \int \tan^2(y) dy \\
 & \quad \downarrow \text{3042} \\
 & \frac{\tan^3(y)}{3} - \int \tan(y)^2 dy \\
 & \quad \downarrow \text{3954} \\
 & \int 1 dy + \frac{\tan^3(y)}{3} - \tan(y) \\
 & \quad \downarrow \text{24} \\
 & y + \frac{\tan^3(y)}{3} - \tan(y)
 \end{aligned}$$

input `Int [Tan[y]^4, y]`

output `y - Tan[y] + Tan[y]^3/3`

Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

method	result	size
default	$y - \tan(y) + \frac{\tan(y)^3}{3}$	13
risch	$y - \frac{4i(3e^{4iy} + 3e^{2iy} + 2)}{3(e^{2iy} + 1)^3}$	31
norman	$\frac{y \tan(\frac{y}{2})^6 - y - \frac{20 \tan(\frac{y}{2})^3}{3} + 2 \tan(\frac{y}{2})^5 + 3y \tan(\frac{y}{2})^2 - 3y \tan(\frac{y}{2})^4 + 2 \tan(\frac{y}{2})}{(\tan(\frac{y}{2})^2 - 1)^3}$	64
parallelrisch	$\frac{3y \tan(\frac{y}{2})^6 - 9y \tan(\frac{y}{2})^4 + 6 \tan(\frac{y}{2})^5 + 9y \tan(\frac{y}{2})^2 - 20 \tan(\frac{y}{2})^3 - 3y + 6 \tan(\frac{y}{2})}{3(\tan(\frac{y}{2}) - 1)^3 (\tan(\frac{y}{2}) + 1)^3}$	72

input `int(sin(y)^4/cos(y)^4,y,method=_RETURNVERBOSE)`

output `y-tan(y)+1/3*tan(y)^3`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 26 vs. $2(12) = 24$.

Time = 0.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.86

$$\int \tan^4(y) dy = \frac{3y \cos(y)^3 - (4 \cos(y)^2 - 1) \sin(y)}{3 \cos(y)^3}$$

input `integrate(sin(y)^4/cos(y)^4,y, algorithm="fricas")`

output `1/3*(3*y*cos(y)^3 - (4*cos(y)^2 - 1)*sin(y))/cos(y)^3`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.36

$$\int \tan^4(y) dy = y + \frac{\sin^3(y)}{3 \cos^3(y)} - \frac{\sin(y)}{\cos(y)}$$

input `integrate(sin(y)**4/cos(y)**4,y)`

output `y + sin(y)**3/(3*cos(y)**3) - sin(y)/cos(y)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \tan^4(y) dy = \frac{1}{3} \tan(y)^3 + y - \tan(y)$$

input `integrate(sin(y)^4/cos(y)^4,y, algorithm="maxima")`

output `1/3*tan(y)^3 + y - tan(y)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \tan^4(y) dy = \frac{1}{3} \tan(y)^3 + y - \tan(y)$$

input `integrate(sin(y)^4/cos(y)^4,y, algorithm="giac")`output `1/3*tan(y)^3 + y - tan(y)`**Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \tan^4(y) dy = \frac{\tan(y)^3}{3} - \tan(y) + y$$

input `int(sin(y)^4/cos(y)^4,y)`output `y - tan(y) + tan(y)^3/3`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.79

$$\int \tan^4(y) dy = \frac{3 \cos(y) \sin(y)^2 y - 3 \cos(y) y - 4 \sin(y)^3 + 3 \sin(y)}{3 \cos(y) (\sin(y)^2 - 1)}$$

input `int(sin(y)^4/cos(y)^4,y)`output `(3*cos(y)*sin(y)**2*y - 3*cos(y)*y - 4*sin(y)**3 + 3*sin(y))/(3*cos(y)*(sin(y)**2 - 1))`

3.75 $\int \frac{z^4}{1+z^2} dz$

Optimal result	468
Mathematica [A] (verified)	468
Rubi [A] (verified)	469
Maple [A] (verified)	470
Fricas [A] (verification not implemented)	470
Sympy [A] (verification not implemented)	470
Maxima [A] (verification not implemented)	471
Giac [A] (verification not implemented)	471
Mupad [B] (verification not implemented)	471
Reduce [B] (verification not implemented)	472

Optimal result

Integrand size = 11, antiderivative size = 13

$$\int \frac{z^4}{1+z^2} dz = -z + \frac{z^3}{3} + \arctan(z)$$

output

```
-z+1/3*z^3+arctan(z)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{z^4}{1+z^2} dz = -z + \frac{z^3}{3} + \arctan(z)$$

input

```
Integrate[z^4/(1+z^2),z]
```

output

```
-z + z^3/3 + ArcTan[z]
```

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{z^4}{z^2 + 1} dz$$

$$\downarrow \text{254}$$

$$\int \left(z^2 + \frac{1}{z^2 + 1} - 1 \right) dz$$

$$\downarrow \text{2009}$$

$$\arctan(z) + \frac{z^3}{3} - z$$

input `Int[z^4/(1 + z^2),z]`

output `-z + z^3/3 + ArcTan[z]`

Defintions of rubi rules used

rule 254 `Int[(x_)^(m_)/((a_) + (b_.)*(x_)^2), x_Symbol] :> Int[PolynomialDivide[x^m, a + b*x^2, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 3]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
default	$-z + \frac{z^3}{3} + \arctan(z)$	12
risch	$-z + \frac{z^3}{3} + \arctan(z)$	12
meijerg	$-\frac{z(-5z^2+15)}{15} + \arctan(z)$	14
parallelrisc	$\frac{z^3}{3} - z + \frac{i \ln(z+i)}{2} - \frac{i \ln(z-i)}{2}$	26

input `int(z^4/(z^2+1),z,method=_RETURNVERBOSE)`output `-z+1/3*z^3+arctan(z)`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{z^4}{1+z^2} dz = \frac{1}{3} z^3 - z + \arctan(z)$$

input `integrate(z^4/(z^2+1),z, algorithm="fricas")`output `1/3*z^3 - z + arctan(z)`**Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.62

$$\int \frac{z^4}{1+z^2} dz = \frac{z^3}{3} - z + \operatorname{atan}(z)$$

input `integrate(z**4/(z**2+1),z)`

output `z**3/3 - z + atan(z)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{z^4}{1+z^2} dz = \frac{1}{3}z^3 - z + \arctan(z)$$

input `integrate(z^4/(z^2+1),z, algorithm="maxima")`

output `1/3*z^3 - z + arctan(z)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{z^4}{1+z^2} dz = \frac{1}{3}z^3 - z + \arctan(z)$$

input `integrate(z^4/(z^2+1),z, algorithm="giac")`

output `1/3*z^3 - z + arctan(z)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{z^4}{1+z^2} dz = \operatorname{atan}(z) - z + \frac{z^3}{3}$$

input `int(z^4/(z^2 + 1),z)`

output `atan(z) - z + z^3/3`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{z^4}{1+z^2} dz = \operatorname{atan}(z) + \frac{z^3}{3} - z$$

input `int(z^4/(z^2+1),z)`

output `(3*atan(z) + z**3 - 3*z)/3`

3.76 $\int e^{x^2}(1 + 2x^2) dx$

Optimal result	473
Mathematica [A] (verified)	473
Rubi [A] (verified)	474
Maple [A] (verified)	475
Fricas [A] (verification not implemented)	475
Sympy [A] (verification not implemented)	476
Maxima [A] (verification not implemented)	476
Giac [A] (verification not implemented)	476
Mupad [B] (verification not implemented)	477
Reduce [B] (verification not implemented)	477

Optimal result

Integrand size = 13, antiderivative size = 7

$$\int e^{x^2}(1 + 2x^2) dx = e^{x^2}x$$

output `exp(x^2)*x`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int e^{x^2}(1 + 2x^2) dx = e^{x^2}x$$

input `Integrate[E^x^2*(1 + 2*x^2),x]`

output `E^x^2*x`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {2656, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{x^2} (2x^2 + 1) dx$$

$$\downarrow \text{2656}$$

$$\int (2e^{x^2} x^2 + e^{x^2}) dx$$

$$\downarrow \text{2009}$$

$$e^{x^2} x$$

input `Int[E^x^2*(1 + 2*x^2),x]`

output `E^x^2*x`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2656 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*(Px_), x_Symbol] := Int[ExpandLinearProduct[F^(a + b*(c + d*x)^n), Px, c, d, x], x] /; FreeQ[{F, a, b, c, d, n}, x] && PolynomialQ[Px, x]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

method	result	size
gospers	$e^{x^2} x$	7
default	$e^{x^2} x$	7
norman	$e^{x^2} x$	7
risch	$e^{x^2} x$	7
parallelrisch	$e^{x^2} x$	7
orering	$e^{x^2} x$	7
meijerg	$i \left(-ie^{x^2} x + \frac{i \operatorname{erfi}(x)\sqrt{\pi}}{2} \right) + \frac{\operatorname{erfi}(x)\sqrt{\pi}}{2}$	29
parts	$\operatorname{erfi}(x) \sqrt{\pi} x^2 + \frac{\operatorname{erfi}(x)\sqrt{\pi}}{2} - 2\sqrt{\pi} \left(\frac{x^2 \operatorname{erfi}(x)}{2} - \frac{\frac{e^{x^2} x - \operatorname{erfi}(x)\sqrt{\pi}}{4}}{\sqrt{\pi}} \right)$	51

input `int(exp(x^2)*(2*x^2+1),x,method=_RETURNVERBOSE)`output `exp(x^2)*x`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int e^{x^2} (1 + 2x^2) dx = x e^{(x^2)}$$

input `integrate(exp(x^2)*(2*x^2+1),x, algorithm="fricas")`output `x*e^(x^2)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.71

$$\int e^{x^2} (1 + 2x^2) dx = xe^{x^2}$$

input `integrate(exp(x**2)*(2*x**2+1),x)`

output `x*exp(x**2)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int e^{x^2} (1 + 2x^2) dx = xe^{(x^2)}$$

input `integrate(exp(x^2)*(2*x^2+1),x, algorithm="maxima")`

output `x*e^(x^2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int e^{x^2} (1 + 2x^2) dx = xe^{(x^2)}$$

input `integrate(exp(x^2)*(2*x^2+1),x, algorithm="giac")`

output `x*e^(x^2)`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int e^{x^2} (1 + 2x^2) dx = x e^{x^2}$$

input `int(exp(x^2)*(2*x^2 + 1),x)`

output `x*exp(x^2)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int e^{x^2} (1 + 2x^2) dx = e^{x^2} x$$

input `int(exp(x^2)*(2*x^2+1),x)`

output `e**(x**2)*x`

$$3.77 \quad \int \frac{e^{x^2}(1+4x^2+x^3+5x^4+2x^6)}{(1+x^2)^2} dx$$

Optimal result	478
Mathematica [A] (verified)	478
Rubi [A] (verified)	479
Maple [A] (verified)	480
Fricas [A] (verification not implemented)	480
Sympy [A] (verification not implemented)	481
Maxima [A] (verification not implemented)	481
Giac [A] (verification not implemented)	481
Mupad [B] (verification not implemented)	482
Reduce [B] (verification not implemented)	482

Optimal result

Integrand size = 33, antiderivative size = 24

$$\int \frac{e^{x^2}(1+4x^2+x^3+5x^4+2x^6)}{(1+x^2)^2} dx = e^{x^2}x + \frac{e^{x^2}}{2(1+x^2)}$$

output `exp(x^2)*x+1/2*exp(x^2)/(x^2+1)`

Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.79

$$\int \frac{e^{x^2}(1+4x^2+x^3+5x^4+2x^6)}{(1+x^2)^2} dx = e^{x^2} \left(x + \frac{1}{2(1+x^2)} \right)$$

input `Integrate[(E^x^2*(1+4*x^2+x^3+5*x^4+2*x^6))/(1+x^2)^2,x]`

output `E^x^2*(x+1/(2*(1+x^2)))`

Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.061$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^{x^2} (2x^6 + 5x^4 + x^3 + 4x^2 + 1)}{(x^2 + 1)^2} dx$$

$$\downarrow \text{7293}$$

$$\int \left(2e^{x^2} x^2 + \frac{e^{x^2} x}{x^2 + 1} - \frac{e^{x^2} x}{(x^2 + 1)^2} + e^{x^2} \right) dx$$

$$\downarrow \text{2009}$$

$$e^{x^2} x + \frac{e^{x^2}}{2(x^2 + 1)}$$

input `Int[(E^x^2*(1 + 4*x^2 + x^3 + 5*x^4 + 2*x^6))/(1 + x^2)^2,x]`

output `E^x^2*x + E^x^2/(2*(1 + x^2))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

method	result	size
gospers	$\frac{(2x^3+2x+1)e^{x^2}}{2x^2+2}$	24
risch	$\frac{(2x^3+2x+1)e^{x^2}}{2x^2+2}$	24
orering	$\frac{(2x^3+2x+1)e^{x^2}}{2x^2+2}$	24
norman	$\frac{e^{x^2}x + e^{x^2}x^3 + \frac{e^{x^2}}{2}}{x^2+1}$	30
parallelrisch	$\frac{2e^{x^2}x^3 + 2e^{x^2}x + e^{x^2}}{2x^2+2}$	31

input `int(exp(x^2)*(2*x^6+5*x^4+x^3+4*x^2+1)/(x^2+1)^2,x,method=_RETURNVERBOSE)`

output `1/2*(2*x^3+2*x+1)*exp(x^2)/(x^2+1)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.96

$$\int \frac{e^{x^2}(1 + 4x^2 + x^3 + 5x^4 + 2x^6)}{(1 + x^2)^2} dx = \frac{(2x^3 + 2x + 1)e^{(x^2)}}{2(x^2 + 1)}$$

input `integrate(exp(x^2)*(2*x^6+5*x^4+x^3+4*x^2+1)/(x^2+1)^2,x, algorithm="fricas")`

output `1/2*(2*x^3 + 2*x + 1)*e^(x^2)/(x^2 + 1)`

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{e^{x^2}(1 + 4x^2 + x^3 + 5x^4 + 2x^6)}{(1 + x^2)^2} dx = \frac{(2x^3 + 2x + 1)e^{x^2}}{2x^2 + 2}$$

input `integrate(exp(x**2)*(2*x**6+5*x**4+x**3+4*x**2+1)/(x**2+1)**2,x)`output `(2*x**3 + 2*x + 1)*exp(x**2)/(2*x**2 + 2)`**Maxima [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.96

$$\int \frac{e^{x^2}(1 + 4x^2 + x^3 + 5x^4 + 2x^6)}{(1 + x^2)^2} dx = \frac{(2x^3 + 2x + 1)e^{(x^2)}}{2(x^2 + 1)}$$

input `integrate(exp(x^2)*(2*x^6+5*x^4+x^3+4*x^2+1)/(x^2+1)^2,x, algorithm="maxima")`output `1/2*(2*x^3 + 2*x + 1)*e^(x^2)/(x^2 + 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.25

$$\int \frac{e^{x^2}(1 + 4x^2 + x^3 + 5x^4 + 2x^6)}{(1 + x^2)^2} dx = \frac{2x^3e^{(x^2)} + 2xe^{(x^2)} + e^{(x^2)}}{2(x^2 + 1)}$$

input `integrate(exp(x^2)*(2*x^6+5*x^4+x^3+4*x^2+1)/(x^2+1)^2,x, algorithm="giac")`output `1/2*(2*x^3*e^(x^2) + 2*x*e^(x^2) + e^(x^2))/(x^2 + 1)`

Mupad [B] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{e^{x^2}(1 + 4x^2 + x^3 + 5x^4 + 2x^6)}{(1 + x^2)^2} dx = \frac{e^{x^2}(2x^3 + 2x + 1)}{2(x^2 + 1)}$$

input `int((exp(x^2)*(4*x^2 + x^3 + 5*x^4 + 2*x^6 + 1))/(x^2 + 1)^2,x)`output `(exp(x^2)*(2*x + 2*x^3 + 1))/(2*(x^2 + 1))`**Reduce [B] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.04

$$\int \frac{e^{x^2}(1 + 4x^2 + x^3 + 5x^4 + 2x^6)}{(1 + x^2)^2} dx = \frac{e^{x^2}(2x^3 + 2x + 1)}{2x^2 + 2}$$

input `int(exp(x^2)*(2*x^6+5*x^4+x^3+4*x^2+1)/(x^2+1)^2,x)`output `(e**(x**2)*(2*x**3 + 2*x + 1))/(2*(x**2 + 1))`

3.78 $\int e^{-1-x} dx$

Optimal result	483
Mathematica [A] (verified)	483
Rubi [A] (verified)	484
Maple [A] (verified)	485
Fricas [A] (verification not implemented)	485
Sympy [A] (verification not implemented)	486
Maxima [A] (verification not implemented)	486
Giac [A] (verification not implemented)	486
Mupad [B] (verification not implemented)	487
Reduce [B] (verification not implemented)	487

Optimal result

Integrand size = 7, antiderivative size = 9

$$\int e^{-1-x} dx = -e^{-1-x}$$

output `-exp(-1-x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int e^{-1-x} dx = -e^{-1-x}$$

input `Integrate[E^(-1 - x), x]`

output `-E^(-1 - x)`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{-x-1} dx$$

$$\downarrow 2624$$

$$-e^{-x-1}$$

input `Int [E^(-1 - x), x]`

output `-E^(-1 - x)`

Defintions of rubi rules used

rule 2624 `Int[((F_)^(v_))^(n_), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;`
`FreeQ[{F, n}, x] && LinearQ[v, x]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

method	result	size
gospers	$-e^{-1-x}$	9
derivativedivides	$-e^{-1-x}$	9
default	$-e^{-1-x}$	9
norman	$-e^{-1-x}$	9
risch	$-e^{-1-x}$	9
parallelrisc	$-e^{-1-x}$	9
orering	$-e^{-1-x}$	9
meijerg	$e^{-1}(1 - e^{-x})$	12

input `int(exp(-1-x),x,method=_RETURNVERBOSE)`

output `-exp(-1-x)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

$$\int e^{-1-x} dx = -e^{(-x-1)}$$

input `integrate(exp(-1-x),x, algorithm="fricas")`

output `-e^(-x - 1)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int e^{-1-x} dx = -e^{-x-1}$$

input `integrate(exp(-1-x), x)`

output `-exp(-x - 1)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

$$\int e^{-1-x} dx = -e^{(-x-1)}$$

input `integrate(exp(-1-x), x, algorithm="maxima")`

output `-e^(-x - 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

$$\int e^{-1-x} dx = -e^{(-x-1)}$$

input `integrate(exp(-1-x), x, algorithm="giac")`

output `-e^(-x - 1)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

$$\int e^{-1-x} dx = -e^{-x-1}$$

input `int(exp(- x - 1),x)`

output `-exp(- x - 1)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.11

$$\int e^{-1-x} dx = -\frac{1}{e^x e}$$

input `int(exp(-1-x),x)`

output `(- 1)/(e**x*e)`

3.79 $\int \left(\frac{1}{x} + x\right) \log(x) dx$

Optimal result	488
Mathematica [A] (verified)	488
Rubi [A] (verified)	489
Maple [A] (warning: unable to verify)	490
Fricas [A] (verification not implemented)	490
Sympy [A] (verification not implemented)	491
Maxima [A] (verification not implemented)	491
Giac [A] (verification not implemented)	491
Mupad [B] (verification not implemented)	492
Reduce [B] (verification not implemented)	492

Optimal result

Integrand size = 8, antiderivative size = 25

$$\int \left(\frac{1}{x} + x\right) \log(x) dx = -\frac{x^2}{4} + \frac{1}{2}x^2 \log(x) + \frac{\log^2(x)}{2}$$

output

```
-1/4*x^2+1/2*x^2*ln(x)+1/2*ln(x)^2
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \left(\frac{1}{x} + x\right) \log(x) dx = -\frac{x^2}{4} + \frac{1}{2}x^2 \log(x) + \frac{\log^2(x)}{2}$$

input

```
Integrate[(x^(-1) + x)*Log[x],x]
```

output

```
-1/4*x^2 + (x^2*Log[x])/2 + Log[x]^2/2
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {2027, 2793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \left(x + \frac{1}{x} \right) \log(x) dx \\ & \quad \downarrow \text{2027} \\ & \int \frac{(x^2 + 1) \log(x)}{x} dx \\ & \quad \downarrow \text{2793} \\ & \int \left(x \log(x) + \frac{\log(x)}{x} \right) dx \\ & \quad \downarrow \text{2009} \\ & -\frac{x^2}{4} + \frac{1}{2}x^2 \log(x) + \frac{\log^2(x)}{2} \end{aligned}$$

input `Int[(x^(-1) + x)*Log[x],x]`

output `-1/4*x^2 + (x^2*Log[x])/2 + Log[x]^2/2`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2027 `Int[(Fx_.)*((a_.)*(x_)^(r_.) + (b_.)*(x_)^(s_.))^(p_.), x_Symbol] := Int[x^(p*r)*(a + b*x^(s - r))^p*Fx, x] /; FreeQ[{a, b, r, s}, x] && IntegerQ[p] && PosQ[s - r] && !(EqQ[p, 1] && EqQ[u, 1])`

rule 2793

```
Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((f_.)*(x_)^(m_.))*((d_) + (e_.)*
(x_)^(r_.))^(q_.), x_Symbol] := With[{u = ExpandIntegrand[a + b*Log[c*x^n],
(f*x)^m*(d + e*x^r)^q, x]}, Int[u, x] /; SumQ[u]] /; FreeQ[{a, b, c, d, e,
f, m, n, q, r}, x] && IntegerQ[q] && (GtQ[q, 0] || (IntegerQ[m] && Integer
Q[r]))
```

Maple [A] (warning: unable to verify)

Time = 0.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

method	result	size
default	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2} + \frac{\ln(x)^2}{2}$	20
norman	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2} + \frac{\ln(x)^2}{2}$	20
risch	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2} + \frac{\ln(x)^2}{2}$	20
parallelrisch	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2} + \frac{\ln(x)^2}{2}$	20
parts	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2} + \frac{\ln(x)^2}{2}$	20

input

```
int((1/x+x)*ln(x),x,method=_RETURNVERBOSE)
```

output

```
-1/4*x^2+1/2*x^2*ln(x)+1/2*ln(x)^2
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \left(\frac{1}{x} + x \right) \log(x) dx = \frac{1}{2} x^2 \log(x) - \frac{1}{4} x^2 + \frac{1}{2} \log(x)^2$$

input

```
integrate((1/x+x)*log(x),x, algorithm="fricas")
```

output

```
1/2*x^2*log(x) - 1/4*x^2 + 1/2*log(x)^2
```

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \left(\frac{1}{x} + x \right) \log(x) dx = \frac{x^2 \log(x)}{2} - \frac{x^2}{4} + \frac{\log(x)^2}{2}$$

input `integrate((1/x+x)*ln(x),x)`output `x**2*log(x)/2 - x**2/4 + log(x)**2/2`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \left(\frac{1}{x} + x \right) \log(x) dx = -\frac{1}{4} x^2 + \frac{1}{2} (x^2 + 2 \log(x)) \log(x) - \frac{1}{2} \log(x)^2$$

input `integrate((1/x+x)*log(x),x, algorithm="maxima")`output `-1/4*x^2 + 1/2*(x^2 + 2*log(x))*log(x) - 1/2*log(x)^2`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \left(\frac{1}{x} + x \right) \log(x) dx = \frac{1}{2} x^2 \log(x) - \frac{1}{4} x^2 + \frac{1}{2} \log(x)^2$$

input `integrate((1/x+x)*log(x),x, algorithm="giac")`output `1/2*x^2*log(x) - 1/4*x^2 + 1/2*log(x)^2`

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \left(\frac{1}{x} + x \right) \log(x) dx = \frac{x^2 \ln(x)}{2} - \frac{x^2}{4} + \frac{\ln(x)^2}{2}$$

input `int(log(x)*(x + 1/x),x)`output `(x^2*log(x))/2 + log(x)^2/2 - x^2/4`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \left(\frac{1}{x} + x \right) \log(x) dx = \frac{\log(x)^2}{2} + \frac{\log(x) x^2}{2} - \frac{x^2}{4}$$

input `int((1/x+x)*log(x),x)`output `(2*log(x)**2 + 2*log(x)*x**2 - x**2)/4`

3.80 $\int \frac{x}{1+x^4} dx$

Optimal result	493
Mathematica [A] (verified)	493
Rubi [A] (verified)	494
Maple [A] (verified)	495
Fricas [A] (verification not implemented)	495
Sympy [A] (verification not implemented)	495
Maxima [A] (verification not implemented)	496
Giac [A] (verification not implemented)	496
Mupad [B] (verification not implemented)	496
Reduce [B] (verification not implemented)	497

Optimal result

Integrand size = 9, antiderivative size = 8

$$\int \frac{x}{1+x^4} dx = \frac{\arctan(x^2)}{2}$$

output `1/2*arctan(x^2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{x}{1+x^4} dx = \frac{\arctan(x^2)}{2}$$

input `Integrate[x/(1 + x^4), x]`

output `ArcTan[x^2]/2`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {807, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{x^4 + 1} dx$$

↓ 807

$$\frac{1}{2} \int \frac{1}{x^4 + 1} dx^2$$

↓ 216

$$\frac{\arctan(x^2)}{2}$$

input

```
Int[x/(1 + x^4), x]
```

output

```
ArcTan[x^2]/2
```

Defintions of rubi rules used

rule 216

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

rule 807

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]
```

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
default	$\frac{\arctan(x^2)}{2}$	7
meijerg	$\frac{\arctan(x^2)}{2}$	7
risch	$\frac{\arctan(x^2)}{2}$	7
parallelrisc	$\frac{i \ln(x^2+i)}{4} - \frac{i \ln(x^2-i)}{4}$	22

input `int(x/(x^4+1),x,method=_RETURNVERBOSE)`

output `1/2*arctan(x^2)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{x}{1+x^4} dx = \frac{1}{2} \arctan(x^2)$$

input `integrate(x/(x^4+1),x, algorithm="fricas")`

output `1/2*arctan(x^2)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int \frac{x}{1+x^4} dx = \frac{\text{atan}(x^2)}{2}$$

input `integrate(x/(x**4+1),x)`

output `atan(x**2)/2`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{x}{1+x^4} dx = \frac{1}{2} \arctan(x^2)$$

input `integrate(x/(x^4+1),x, algorithm="maxima")`

output `1/2*arctan(x^2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{x}{1+x^4} dx = \frac{1}{2} \arctan(x^2)$$

input `integrate(x/(x^4+1),x, algorithm="giac")`

output `1/2*arctan(x^2)`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{x}{1+x^4} dx = \frac{\operatorname{atan}(x^2)}{2}$$

input `int(x/(x^4 + 1),x)`

output `atan(x^2)/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 29, normalized size of antiderivative = 3.62

$$\int \frac{x}{1+x^4} dx = -\frac{\operatorname{atan}\left(\frac{\sqrt{2}-2x}{\sqrt{2}}\right)}{2} - \frac{\operatorname{atan}\left(\frac{\sqrt{2}+2x}{\sqrt{2}}\right)}{2}$$

input `int(x/(x^4+1),x)`output `(- (atan((sqrt(2) - 2*x)/sqrt(2)) + atan((sqrt(2) + 2*x)/sqrt(2))))/2`

3.81 $\int \frac{x^5}{1+x^4} dx$

Optimal result	498
Mathematica [A] (verified)	498
Rubi [A] (verified)	499
Maple [A] (verified)	500
Fricas [A] (verification not implemented)	500
Sympy [A] (verification not implemented)	501
Maxima [A] (verification not implemented)	501
Giac [A] (verification not implemented)	501
Mupad [B] (verification not implemented)	502
Reduce [B] (verification not implemented)	502

Optimal result

Integrand size = 11, antiderivative size = 16

$$\int \frac{x^5}{1+x^4} dx = \frac{x^2}{2} - \frac{\arctan(x^2)}{2}$$

output `1/2*x^2-1/2*arctan(x^2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{x^5}{1+x^4} dx = \frac{x^2}{2} - \frac{\arctan(x^2)}{2}$$

input `Integrate[x^5/(1 + x^4),x]`

output `x^2/2 - ArcTan[x^2]/2`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {807, 262, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^5}{x^4 + 1} dx \\ & \quad \downarrow \text{807} \\ & \frac{1}{2} \int \frac{x^4}{x^4 + 1} dx^2 \\ & \quad \downarrow \text{262} \\ & \frac{1}{2} \left(x^2 - \int \frac{1}{x^4 + 1} dx^2 \right) \\ & \quad \downarrow \text{216} \\ & \frac{1}{2} (x^2 - \arctan(x^2)) \end{aligned}$$

input `Int[x^5/(1 + x^4), x]`

output `(x^2 - ArcTan[x^2])/2`

Defintions of rubi rules used

rule 216

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

rule 262 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.81

method	result	size
default	$\frac{x^2}{2} - \frac{\arctan(x^2)}{2}$	13
meijerg	$\frac{x^2}{2} - \frac{\arctan(x^2)}{2}$	13
risch	$\frac{x^2}{2} - \frac{\arctan(x^2)}{2}$	13
parallelrisch	$\frac{x^2}{2} + \frac{i \ln(x^2 - i)}{4} - \frac{i \ln(x^2 + i)}{4}$	27

input `int(x^5/(x^4+1),x,method=_RETURNVERBOSE)`

output `1/2*x^2-1/2*arctan(x^2)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{x^5}{1+x^4} dx = \frac{1}{2}x^2 - \frac{1}{2}\arctan(x^2)$$

input `integrate(x^5/(x^4+1),x, algorithm="fricas")`

output `1/2*x^2 - 1/2*arctan(x^2)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.62

$$\int \frac{x^5}{1+x^4} dx = \frac{x^2}{2} - \frac{\operatorname{atan}(x^2)}{2}$$

input `integrate(x**5/(x**4+1),x)`

output `x**2/2 - atan(x**2)/2`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{x^5}{1+x^4} dx = \frac{1}{2} x^2 - \frac{1}{2} \operatorname{arctan}(x^2)$$

input `integrate(x^5/(x^4+1),x, algorithm="maxima")`

output `1/2*x^2 - 1/2*arctan(x^2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{x^5}{1+x^4} dx = \frac{1}{2} x^2 - \frac{1}{2} \operatorname{arctan}(x^2)$$

input `integrate(x^5/(x^4+1),x, algorithm="giac")`

output `1/2*x^2 - 1/2*arctan(x^2)`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{x^5}{1+x^4} dx = \frac{x^2}{2} - \frac{\operatorname{atan}(x^2)}{2}$$

input `int(x^5/(x^4 + 1),x)`output `x^2/2 - atan(x^2)/2`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 34, normalized size of antiderivative = 2.12

$$\int \frac{x^5}{1+x^4} dx = \frac{\operatorname{atan}\left(\frac{\sqrt{2}-2x}{\sqrt{2}}\right)}{2} + \frac{\operatorname{atan}\left(\frac{\sqrt{2}+2x}{\sqrt{2}}\right)}{2} + \frac{x^2}{2}$$

input `int(x^5/(x^4+1),x)`output `(atan((sqrt(2) - 2*x)/sqrt(2)) + atan((sqrt(2) + 2*x)/sqrt(2)) + x**2)/2`

3.82 $\int \frac{1}{1+\tan^2(x)} dx$

Optimal result	503
Mathematica [A] (verified)	503
Rubi [A] (verified)	504
Maple [A] (verified)	505
Fricas [A] (verification not implemented)	506
Sympy [B] (verification not implemented)	506
Maxima [A] (verification not implemented)	506
Giac [A] (verification not implemented)	507
Mupad [B] (verification not implemented)	507
Reduce [B] (verification not implemented)	507

Optimal result

Integrand size = 8, antiderivative size = 14

$$\int \frac{1}{1 + \tan^2(x)} dx = \frac{x}{2} + \frac{1}{2} \cos(x) \sin(x)$$

output `1/2*x+1/2*cos(x)*sin(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + \tan^2(x)} dx = \frac{x}{2} + \frac{1}{4} \sin(2x)$$

input `Integrate[(1 + Tan[x]^2)^(-1),x]`

output `x/2 + Sin[2*x]/4`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {3042, 4140, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\tan^2(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\tan(x)^2 + 1} dx \\
 & \quad \downarrow \text{4140} \\
 & \int \cos^2(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(x + \frac{\pi}{2}\right)^2 dx \\
 & \quad \downarrow \text{3115} \\
 & \frac{\int 1 dx}{2} + \frac{1}{2} \sin(x) \cos(x) \\
 & \quad \downarrow \text{24} \\
 & \frac{x}{2} + \frac{1}{2} \sin(x) \cos(x)
 \end{aligned}$$

input `Int[(1 + Tan[x]^2)^(-1),x]`

output `x/2 + (Cos[x]*Sin[x])/2`

Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4140 `Int[(u_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)]^2)^(p_), x_Symbol] := Int[ActivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a, b]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.79

method	result	size
risch	$\frac{x}{2} + \frac{\sin(2x)}{4}$	11
derivativedivides	$\frac{\tan(x)}{2+2\tan(x)^2} + \frac{\arctan(\tan(x))}{2}$	19
default	$\frac{\tan(x)}{2+2\tan(x)^2} + \frac{\arctan(\tan(x))}{2}$	19
parallelrisch	$\frac{x \tan(x)^2 + x + \tan(x)}{2+2\tan(x)^2}$	21
norman	$\frac{\frac{x}{2} + \frac{x \tan(x)^2}{2} + \frac{\tan(x)}{2}}{1+\tan(x)^2}$	25

input `int(1/(1+tan(x)^2), x, method=_RETURNVERBOSE)`

output `1/2*x+1/4*sin(2*x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.43

$$\int \frac{1}{1 + \tan^2(x)} dx = \frac{x \tan(x)^2 + x + \tan(x)}{2(\tan(x)^2 + 1)}$$

input `integrate(1/(1+tan(x)^2),x, algorithm="fricas")`

output `1/2*(x*tan(x)^2 + x + tan(x))/(tan(x)^2 + 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 36 vs. 2(10) = 20.

Time = 0.17 (sec) , antiderivative size = 36, normalized size of antiderivative = 2.57

$$\int \frac{1}{1 + \tan^2(x)} dx = \frac{x \tan^2(x)}{2 \tan^2(x) + 2} + \frac{x}{2 \tan^2(x) + 2} + \frac{\tan(x)}{2 \tan^2(x) + 2}$$

input `integrate(1/(1+tan(x)**2),x)`

output `x*tan(x)**2/(2*tan(x)**2 + 2) + x/(2*tan(x)**2 + 2) + tan(x)/(2*tan(x)**2 + 2)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{1}{1 + \tan^2(x)} dx = \frac{1}{2} x + \frac{\tan(x)}{2(\tan(x)^2 + 1)}$$

input `integrate(1/(1+tan(x)^2),x, algorithm="maxima")`

output `1/2*x + 1/2*tan(x)/(tan(x)^2 + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{1}{1 + \tan^2(x)} dx = \frac{1}{2}x + \frac{\tan(x)}{2(\tan(x)^2 + 1)}$$

input `integrate(1/(1+tan(x)^2),x, algorithm="giac")`

output `1/2*x + 1/2*tan(x)/(tan(x)^2 + 1)`

Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{1}{1 + \tan^2(x)} dx = \frac{x}{2} + \frac{\sin(2x)}{4}$$

input `int(1/(tan(x)^2 + 1),x)`

output `x/2 + sin(2*x)/4`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.50

$$\int \frac{1}{1 + \tan^2(x)} dx = \frac{\tan(x)^2 x + \tan(x) + x}{2 \tan(x)^2 + 2}$$

input `int(1/(1+tan(x)^2),x)`

output `(tan(x)**2*x + tan(x) + x)/(2*(tan(x)**2 + 1))`

3.83 $\int \frac{x^4}{(1-x^2)^{5/2}} dx$

Optimal result	508
Mathematica [A] (verified)	508
Rubi [A] (verified)	509
Maple [A] (verified)	510
Fricas [B] (verification not implemented)	510
Sympy [B] (verification not implemented)	511
Maxima [A] (verification not implemented)	511
Giac [A] (verification not implemented)	512
Mupad [B] (verification not implemented)	512
Reduce [B] (verification not implemented)	512

Optimal result

Integrand size = 15, antiderivative size = 35

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{x^3}{3(1-x^2)^{3/2}} - \frac{x}{\sqrt{1-x^2}} + \arcsin(x)$$

output `1/3*x^3/(-x^2+1)^(3/2)+arcsin(x)-x/(-x^2+1)^(1/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.26

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{x(-3+4x^2)}{3(1-x^2)^{3/2}} + 2 \arctan\left(\frac{x}{-1+\sqrt{1-x^2}}\right)$$

input `Integrate[x^4/(1-x^2)^(5/2),x]`

output `(x*(-3+4*x^2))/(3*(1-x^2)^(3/2))+2*ArcTan[x/(-1+Sqrt[1-x^2])]`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {252, 252, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx$$

$$\downarrow 252$$

$$\frac{x^3}{3(1-x^2)^{3/2}} - \int \frac{x^2}{(1-x^2)^{3/2}} dx$$

$$\downarrow 252$$

$$\int \frac{1}{\sqrt{1-x^2}} dx - \frac{x}{\sqrt{1-x^2}} + \frac{x^3}{3(1-x^2)^{3/2}}$$

$$\downarrow 223$$

$$\arcsin(x) - \frac{x}{\sqrt{1-x^2}} + \frac{x^3}{3(1-x^2)^{3/2}}$$

input `Int[x^4/(1 - x^2)^(5/2),x]`

output `x^3/(3*(1 - x^2)^(3/2)) - x/Sqrt[1 - x^2] + ArcSin[x]`

Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 252

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(2*b*(p + 1))), x] - Simp[c^2*(m - 1)/(2*b*(p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && LtQ[p, -1] && GtQ[m, 1] && !ILtQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]
```

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.86

method	result	size
default	$\frac{x^3}{3(-x^2+1)^{\frac{3}{2}}} + \arcsin(x) - \frac{x}{\sqrt{-x^2+1}}$	30
risch	$-\frac{x(4x^2-3)}{3(x^2-1)\sqrt{-x^2+1}} + \arcsin(x)$	30
meijerg	$-\frac{2i \left(-\frac{i\sqrt{\pi}x(-20x^2+15)}{10(-x^2+1)^{\frac{3}{2}}} + \frac{3i\sqrt{\pi}\arcsin(x)}{2} \right)}{3\sqrt{\pi}}$	39
pseudoelliptic	$\frac{(x^2-1)\sqrt{-x^2+1}\arctan\left(\frac{\sqrt{-x^2+1}}{x}\right) + \frac{4x^3}{3} - x}{(-x^2+1)^{\frac{3}{2}}}$	49
trager	$\frac{(4x^2-3)x\sqrt{-x^2+1}}{3(x^2-1)^2} + \text{RootOf}(_Z^2 + 1) \ln(\text{RootOf}(_Z^2 + 1)\sqrt{-x^2+1} + x)$	54

input `int(x^4/(-x^2+1)^(5/2),x,method=_RETURNVERBOSE)`output `1/3*x^3/(-x^2+1)^(3/2)+arcsin(x)-1/(-x^2+1)^(1/2)*x`**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 63 vs. 2(29) = 58.

Time = 0.07 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.80

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = -\frac{6(x^4 - 2x^2 + 1) \arctan\left(\frac{\sqrt{-x^2+1}-1}{x}\right) - (4x^3 - 3x)\sqrt{-x^2+1}}{3(x^4 - 2x^2 + 1)}$$

input `integrate(x^4/(-x^2+1)^(5/2),x, algorithm="fricas")`

output `-1/3*(6*(x^4 - 2*x^2 + 1)*arctan((sqrt(-x^2 + 1) - 1)/x) - (4*x^3 - 3*x)*sqrt(-x^2 + 1))/(x^4 - 2*x^2 + 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 105 vs. $2(26) = 52$.

Time = 0.33 (sec) , antiderivative size = 105, normalized size of antiderivative = 3.00

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{3x^4 \operatorname{asin}(x)}{3x^4 - 6x^2 + 3} + \frac{4x^3 \sqrt{1-x^2}}{3x^4 - 6x^2 + 3} - \frac{6x^2 \operatorname{asin}(x)}{3x^4 - 6x^2 + 3} - \frac{3x \sqrt{1-x^2}}{3x^4 - 6x^2 + 3} + \frac{3 \operatorname{asin}(x)}{3x^4 - 6x^2 + 3}$$

input `integrate(x**4/(-x**2+1)**(5/2),x)`

output `3*x**4*asin(x)/(3*x**4 - 6*x**2 + 3) + 4*x**3*sqrt(1 - x**2)/(3*x**4 - 6*x**2 + 3) - 6*x**2*asin(x)/(3*x**4 - 6*x**2 + 3) - 3*x*sqrt(1 - x**2)/(3*x**4 - 6*x**2 + 3) + 3*asin(x)/(3*x**4 - 6*x**2 + 3)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.26

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{1}{3} x \left(\frac{3x^2}{(-x^2+1)^{3/2}} - \frac{2}{(-x^2+1)^{3/2}} \right) - \frac{x}{3\sqrt{-x^2+1}} + \operatorname{arcsin}(x)$$

input `integrate(x^4/(-x^2+1)^(5/2),x, algorithm="maxima")`

output `1/3*x*(3*x^2/(-x^2 + 1)^(3/2) - 2/(-x^2 + 1)^(3/2)) - 1/3*x/sqrt(-x^2 + 1) + arcsin(x)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.83

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{(4x^2-3)\sqrt{-x^2+1}x}{3(x^2-1)^2} + \arcsin(x)$$

input `integrate(x^4/(-x^2+1)^(5/2),x, algorithm="giac")`output `1/3*(4*x^2 - 3)*sqrt(-x^2 + 1)*x/(x^2 - 1)^2 + arcsin(x)`**Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 91, normalized size of antiderivative = 2.60

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \arcsin(x) + \frac{3\sqrt{1-x^2}}{4(x-1)} + \frac{3\sqrt{1-x^2}}{4(x+1)} - \sqrt{1-x^2} \left(\frac{1}{12(x-1)} - \frac{1}{12(x-1)^2} \right) - \sqrt{1-x^2} \left(\frac{1}{12(x+1)} + \frac{1}{12(x+1)^2} \right)$$

input `int(x^4/(1 - x^2)^(5/2),x)`output `asin(x) + (3*(1 - x^2)^(1/2))/(4*(x - 1)) + (3*(1 - x^2)^(1/2))/(4*(x + 1)) - (1 - x^2)^(1/2)*(1/(12*(x - 1)) - 1/(12*(x - 1)^2)) - (1 - x^2)^(1/2)*(1/(12*(x + 1)) + 1/(12*(x + 1)^2))`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.57

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{3\sqrt{-x^2+1} \arcsin(x) x^2 - 3\sqrt{-x^2+1} \arcsin(x) - 4x^3 + 3x}{3\sqrt{-x^2+1} (x^2 - 1)}$$

input `int(x^4/(-x^2+1)^(5/2),x)`

output
$$\frac{(3\sqrt{-x^2 + 1})\operatorname{asin}(x)x^2 - 3\sqrt{-x^2 + 1}\operatorname{asin}(x) - 4x^3 + 3x}{(3\sqrt{-x^2 + 1})(x^2 - 1)}$$

$$3.84 \quad \int -\frac{x^2}{(1-x^2)^{3/2}} dx$$

Optimal result	514
Mathematica [B] (verified)	514
Rubi [A] (verified)	515
Maple [A] (verified)	516
Fricas [B] (verification not implemented)	516
Sympy [B] (verification not implemented)	517
Maxima [A] (verification not implemented)	517
Giac [A] (verification not implemented)	518
Mupad [B] (verification not implemented)	518
Reduce [B] (verification not implemented)	518

Optimal result

Integrand size = 16, antiderivative size = 17

$$\int -\frac{x^2}{(1-x^2)^{3/2}} dx = -\frac{x}{\sqrt{1-x^2}} + \arcsin(x)$$

output `arcsin(x)-x/(-x^2+1)^(1/2)`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 35 vs. 2(17) = 34.

Time = 0.05 (sec) , antiderivative size = 35, normalized size of antiderivative = 2.06

$$\int -\frac{x^2}{(1-x^2)^{3/2}} dx = -\frac{x}{\sqrt{1-x^2}} + 2 \arctan\left(\frac{x}{-1 + \sqrt{1-x^2}}\right)$$

input `Integrate[-(x^2/(1-x^2)^(3/2)),x]`

output `-(x/Sqrt[1-x^2]) + 2*ArcTan[x/(-1+Sqrt[1-x^2])]`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {25, 252, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int -\frac{x^2}{(1-x^2)^{3/2}} dx$$

↓ 25

$$-\int \frac{x^2}{(1-x^2)^{3/2}} dx$$

↓ 252

$$\int \frac{1}{\sqrt{1-x^2}} dx - \frac{x}{\sqrt{1-x^2}}$$

↓ 223

$$\arcsin(x) - \frac{x}{\sqrt{1-x^2}}$$

input `Int[-(x^2/(1 - x^2)^(3/2)),x]`

output `-(x/Sqrt[1 - x^2]) + ArcSin[x]`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 252

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(2*b*(p + 1))), x] - Simp[c^2*(m - 1)/(2*b*(p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && LtQ[p, -1] && GtQ[m, 1] && !ILtQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]
```

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

method	result	size
default	$\arcsin(x) - \frac{x}{\sqrt{-x^2+1}}$	16
risch	$\arcsin(x) - \frac{x}{\sqrt{-x^2+1}}$	16
meijerg	$-\frac{i\left(-\frac{i\sqrt{\pi}x}{\sqrt{-x^2+1}} + i\sqrt{\pi} \arcsin(x)\right)}{\sqrt{\pi}}$	32
pseudoelliptic	$-\frac{\arctan\left(\frac{\sqrt{-x^2+1}}{x}\right)\sqrt{-x^2+1}+x}{\sqrt{-x^2+1}}$	38
trager	$\frac{x\sqrt{-x^2+1}}{x^2-1} + \text{RootOf}(_Z^2 + 1) \ln(\text{RootOf}(_Z^2 + 1) \sqrt{-x^2 + 1} + x)$	46

input

```
int(-x^2/(-x^2+1)^(3/2),x,method=_RETURNVERBOSE)
```

output

```
arcsin(x)-1/(-x^2+1)^(1/2)*x
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 45 vs. 2(15) = 30.

Time = 0.09 (sec) , antiderivative size = 45, normalized size of antiderivative = 2.65

$$\int -\frac{x^2}{(1-x^2)^{3/2}} dx = -\frac{2(x^2-1)\arctan\left(\frac{\sqrt{-x^2+1}-1}{x}\right) - \sqrt{-x^2+1}x}{x^2-1}$$

input

```
integrate(-x^2/(-x^2+1)^(3/2),x, algorithm="fricas")
```

output $-(2*(x^2 - 1)*\arctan((\sqrt{-x^2 + 1} - 1)/x) - \sqrt{-x^2 + 1}*x)/(x^2 - 1)$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 34 vs. $2(12) = 24$.

Time = 0.18 (sec) , antiderivative size = 34, normalized size of antiderivative = 2.00

$$\int -\frac{x^2}{(1-x^2)^{3/2}} dx = \frac{x^2 \operatorname{asin}(x)}{x^2 - 1} + \frac{x\sqrt{1-x^2}}{x^2 - 1} - \frac{\operatorname{asin}(x)}{x^2 - 1}$$

input `integrate(-x**2/(-x**2+1)**(3/2),x)`

output $x**2*\operatorname{asin}(x)/(x**2 - 1) + x*\sqrt{1 - x**2}/(x**2 - 1) - \operatorname{asin}(x)/(x**2 - 1)$

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int -\frac{x^2}{(1-x^2)^{3/2}} dx = -\frac{x}{\sqrt{-x^2 + 1}} + \operatorname{arcsin}(x)$$

input `integrate(-x^2/(-x^2+1)^(3/2),x, algorithm="maxima")`

output $-x/\sqrt{-x^2 + 1} + \operatorname{arcsin}(x)$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.24

$$\int -\frac{x^2}{(1-x^2)^{3/2}} dx = \frac{\sqrt{-x^2+1}x}{x^2-1} + \arcsin(x)$$

input `integrate(-x^2/(-x^2+1)^(3/2),x, algorithm="giac")`output `sqrt(-x^2 + 1)*x/(x^2 - 1) + arcsin(x)`**Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 2.18

$$\int -\frac{x^2}{(1-x^2)^{3/2}} dx = \operatorname{asin}(x) + \frac{\sqrt{1-x^2}}{2(x-1)} + \frac{\sqrt{1-x^2}}{2(x+1)}$$

input `int(-x^2/(1 - x^2)^(3/2),x)`output `asin(x) + (1 - x^2)^(1/2)/(2*(x - 1)) + (1 - x^2)^(1/2)/(2*(x + 1))`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.53

$$\int -\frac{x^2}{(1-x^2)^{3/2}} dx = \frac{\sqrt{-x^2+1} \operatorname{asin}(x) - x}{\sqrt{-x^2+1}}$$

input `int(-x^2/(-x^2+1)^(3/2),x)`output `(sqrt(-x**2 + 1)*asin(x) - x)/sqrt(-x**2 + 1)`

3.85 $\int e^x \sin(x) dx$

Optimal result	519
Mathematica [A] (verified)	519
Rubi [A] (verified)	520
Maple [A] (verified)	520
Fricas [A] (verification not implemented)	521
Sympy [A] (verification not implemented)	521
Maxima [A] (verification not implemented)	522
Giac [A] (verification not implemented)	522
Mupad [B] (verification not implemented)	522
Reduce [B] (verification not implemented)	523

Optimal result

Integrand size = 6, antiderivative size = 19

$$\int e^x \sin(x) dx = -\frac{1}{2}e^x \cos(x) + \frac{1}{2}e^x \sin(x)$$

output `-1/2*exp(x)*cos(x)+1/2*exp(x)*sin(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int e^x \sin(x) dx = \frac{1}{2}e^x(-\cos(x) + \sin(x))$$

input `Integrate[E^x*Sin[x],x]`

output `(E^x*(-Cos[x] + Sin[x]))/2`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {4932}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^x \sin(x) dx$$

↓ 4932

$$\frac{1}{2}e^x \sin(x) - \frac{1}{2}e^x \cos(x)$$

input

```
Int[E^x*Sin[x],x]
```

output

```
-1/2*(E^x*Cos[x]) + (E^x*Sin[x])/2
```

Defintions of rubi rules used

rule 4932

```
Int[(F_)^((c_.)*((a_.) + (b_.)*(x_)))*Sin[(d_.) + (e_.)*(x_)], x_Symbol] :>
Simp[b*c*Log[F]*F^(c*(a + b*x))*(Sin[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x
] - Simp[e*F^(c*(a + b*x))*(Cos[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] /; F
reeQ[{F, a, b, c, d, e}, x] && NeQ[e^2 + b^2*c^2*Log[F]^2, 0]
```

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

method	result	size
parallelrisch	$\frac{e^x(-\cos(x)+\sin(x))}{2}$	12
default	$-\frac{e^x \cos(x)}{2} + \frac{e^x \sin(x)}{2}$	14
orering	$-\frac{e^x \cos(x)}{2} + \frac{e^x \sin(x)}{2}$	14
norman	$\frac{e^x \tan(\frac{x}{2}) + \frac{e^x \tan(\frac{x}{2})^2}{2} - \frac{e^x}{2}}{1 + \tan(\frac{x}{2})^2}$	34
risch	$-\frac{e^{(1+i)x}}{4} - \frac{ie^{(1+i)x}}{4} - \frac{e^{(1-i)x}}{4} + \frac{ie^{(1-i)x}}{4}$	36

input `int(exp(x)*sin(x),x,method=_RETURNVERBOSE)`

output `1/2*exp(x)*(-cos(x)+sin(x))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.68

$$\int e^x \sin(x) dx = -\frac{1}{2} \cos(x) e^x + \frac{1}{2} e^x \sin(x)$$

input `integrate(exp(x)*sin(x),x, algorithm="fricas")`

output `-1/2*cos(x)*e^x + 1/2*e^x*sin(x)`

Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int e^x \sin(x) dx = \frac{e^x \sin(x)}{2} - \frac{e^x \cos(x)}{2}$$

input `integrate(exp(x)*sin(x),x)`

output `exp(x)*sin(x)/2 - exp(x)*cos(x)/2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int e^x \sin(x) dx = -\frac{1}{2} (\cos(x) - \sin(x))e^x$$

input `integrate(exp(x)*sin(x),x, algorithm="maxima")`

output `-1/2*(cos(x) - sin(x))*e^x`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int e^x \sin(x) dx = -\frac{1}{2} (\cos(x) - \sin(x))e^x$$

input `integrate(exp(x)*sin(x),x, algorithm="giac")`

output `-1/2*(cos(x) - sin(x))*e^x`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int e^x \sin(x) dx = -\frac{e^x (\cos(x) - \sin(x))}{2}$$

input `int(exp(x)*sin(x),x)`

output `-(exp(x)*(cos(x) - sin(x)))/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int e^x \sin(x) dx = \frac{e^x(-\cos(x) + \sin(x))}{2}$$

input `int(exp(x)*sin(x),x)`

output `(e**x*(- cos(x) + sin(x)))/2`

3.86 $\int \frac{1}{x} dx$

Optimal result	524
Mathematica [A] (verified)	524
Rubi [A] (verified)	525
Maple [A] (verified)	525
Fricas [A] (verification not implemented)	526
Sympy [A] (verification not implemented)	526
Maxima [A] (verification not implemented)	526
Giac [A] (verification not implemented)	527
Mupad [B] (verification not implemented)	527
Reduce [B] (verification not implemented)	527

Optimal result

Integrand size = 3, antiderivative size = 2

$$\int \frac{1}{x} dx = \log(x)$$

output `ln(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{x} dx = \log(x)$$

input `Integrate[x^(-1), x]`

output `Log[x]`

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x} dx$$

↓ 14

$$\log(x)$$

input `Int [x(-1), x]`

output `Log [x]`

Defintions of rubi rules used

rule 14 `Int [(a_.)/(x_), x_Symbol] :> Simp[a*Log[x], x] /; FreeQ[a, x]`

Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
default	$\ln(x)$	3
norman	$\ln(x)$	3
risch	$\ln(x)$	3
parallelrisc	$\ln(x)$	3

input `int(1/x,x,method=_RETURNVERBOSE)`

output `ln(x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{x} dx = \log(x)$$

input `integrate(1/x,x, algorithm="fricas")`

output `log(x)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{x} dx = \log(x)$$

input `integrate(1/x,x)`

output `log(x)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{x} dx = \log(x)$$

input `integrate(1/x,x, algorithm="maxima")`

output `log(x)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

$$\int \frac{1}{x} dx = \log(|x|)$$

input `integrate(1/x,x, algorithm="giac")`

output `log(abs(x))`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{x} dx = \ln(x)$$

input `int(1/x,x)`

output `log(x)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{x} dx = \log(x)$$

input `int(1/x,x)`

output `log(x)`

3.87 $\int \frac{\sec(2t)}{1+\sec^2(t)+3 \tan(t)} dt$

Optimal result	528
Mathematica [A] (verified)	528
Rubi [A] (verified)	529
Maple [A] (verified)	530
Fricas [A] (verification not implemented)	531
Sympy [F]	531
Maxima [B] (verification not implemented)	531
Giac [A] (verification not implemented)	532
Mupad [B] (verification not implemented)	532
Reduce [F]	533

Optimal result

Integrand size = 17, antiderivative size = 45

$$\int \frac{\sec(2t)}{1 + \sec^2(t) + 3 \tan(t)} dt = -\frac{1}{12} \log(\cos(t) - \sin(t)) - \frac{1}{4} \log(\cos(t) + \sin(t)) + \frac{1}{3} \log(2 \cos(t) + \sin(t)) - \frac{1}{2(1 + \tan(t))}$$

output -1/12*ln(cos(t)-sin(t))-1/4*ln(cos(t)+sin(t))+1/3*ln(2*cos(t)+sin(t))-1/2/(1+tan(t))

Mathematica [A] (verified)

Time = 0.45 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.62

$$\int \frac{\sec(2t)}{1 + \sec^2(t) + 3 \tan(t)} dt = \frac{\cos(t)(\log(\cos(t) - \sin(t)) + 3 \log(\cos(t) + \sin(t)) - 4 \log(2 \cos(t) + \sin(t))) + (-6 + \log(\cos(t) - \sin(t)))}{12(\cos(t) + \sin(t))}$$

input Integrate[Sec[2*t]/(1 + Sec[t]^2 + 3*Tan[t]),t]

output

```
-1/12*(Cos[t]*(Log[Cos[t] - Sin[t]] + 3*Log[Cos[t] + Sin[t]] - 4*Log[2*Cos[t] + Sin[t]]) + (-6 + Log[Cos[t] - Sin[t]] + 3*Log[Cos[t] + Sin[t]] - 4*Log[2*Cos[t] + Sin[t]])*Sin[t])/(Cos[t] + Sin[t])
```

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.89, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {3042, 4889, 1141, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(2t)}{3 \tan(t) + \sec^2(t) + 1} dt$$

↓ 3042

$$\int \frac{\sec(2t)}{3 \tan(t) + \sec(t)^2 + 1} dt$$

↓ 4889

$$\int \frac{1}{(\tan(t) + 1)^2 (-\tan^2(t) - \tan(t) + 2)} d \tan(t)$$

↓ 1141

$$-\int \left(\frac{1}{4(\tan(t) + 1)} - \frac{1}{3(\tan(t) + 2)} - \frac{1}{2(\tan(t) + 1)^2} - \frac{1}{12(1 - \tan(t))} \right) d \tan(t)$$

↓ 2009

$$-\frac{1}{2(\tan(t) + 1)} - \frac{1}{12} \log(1 - \tan(t)) - \frac{1}{4} \log(\tan(t) + 1) + \frac{1}{3} \log(\tan(t) + 2)$$

input

```
Int[Sec[2*t]/(1 + Sec[t]^2 + 3*Tan[t]),t]
```

output

```
-1/12*Log[1 - Tan[t]] - Log[1 + Tan[t]]/4 + Log[2 + Tan[t]]/3 - 1/(2*(1 + Tan[t]))
```

Definitions of rubi rules used

rule 1141

```
Int[((d_.) + (e_.)*(x_))^(m_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_
Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[1/c^p Int[ExpandIntegrand[
(d + e*x)^m*(b/2 - q/2 + c*x)^p*(b/2 + q/2 + c*x)^p, x], x] /; EqQ[p, -
1] || !FractionalPowerFactorQ[q]] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[p,
0] && IntegerQ[m] && NiceSqrtQ[b^2 - 4*a*c]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4889

```
Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, With[{d = FreeFactors
[Tan[v], x]}, Simp[d/Coefficient[v, x, 1] Subst[Int[SubstFor[1/(1 + d^2*x
^2), Tan[v]/d, u, x], x], x, Tan[v]/d], x]] /; !FalseQ[v] && FunctionOfQ[N
onfreeFactors[Tan[v], x], u, x]] /; InverseFunctionFreeQ[u, x] && !MatchQ[
u, (v_.)*((c_.)*tan[w_]^(n_.)*tan[z_]^(n_.))^p /; FreeQ[{c, p}, x] && I
ntegerQ[n] && LinearQ[w, x] && EqQ[z, 2*w]]
```

Maple [A] (verified)

Time = 1.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.69

method	result	size
default	$-\frac{\ln(\tan(t)-1)}{12} + \frac{\ln(\tan(t)+2)}{3} - \frac{1}{2(1+\tan(t))} - \frac{\ln(1+\tan(t))}{4}$	31
risch	$-\frac{1}{2(e^{2it}+i)} - \frac{\ln(e^{2it}-i)}{12} + \frac{\ln(e^{2it}+\frac{3}{5}+\frac{4i}{5})}{3} - \frac{\ln(e^{2it}+i)}{4}$	48

input

```
int(sec(2*t)/(1+sec(t)^2+3*tan(t)),t,method=_RETURNVERBOSE)
```

output

```
-1/12*ln(tan(t)-1)+1/3*ln(tan(t)+2)-1/2/(1+tan(t))-1/4*ln(1+tan(t))
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.58

$$\int \frac{\sec(2t)}{1 + \sec^2(t) + 3 \tan(t)} dt$$

$$= \frac{4(\cos(t) + \sin(t)) \log\left(\frac{3}{4} \cos^2(t) + \cos(t) \sin(t) + \frac{1}{4}\right) - 3(\cos(t) + \sin(t)) \log(2 \cos(t) \sin(t) + 1) - (\cos(t) + \sin(t)) \log(-2 \cos(t) \sin(t) + 1) - 6 \cos(t) + 6 \sin(t)}{24(\cos(t) + \sin(t))}$$

input `integrate(sec(2*t)/(1+sec(t)^2+3*tan(t)),t, algorithm="fricas")`

output `1/24*(4*(cos(t) + sin(t))*log(3/4*cos(t)^2 + cos(t)*sin(t) + 1/4) - 3*(cos(t) + sin(t))*log(2*cos(t)*sin(t) + 1) - (cos(t) + sin(t))*log(-2*cos(t)*sin(t) + 1) - 6*cos(t) + 6*sin(t))/(cos(t) + sin(t))`

Sympy [F]

$$\int \frac{\sec(2t)}{1 + \sec^2(t) + 3 \tan(t)} dt = \int \frac{\sec(2t)}{3 \tan(t) + \sec^2(t) + 1} dt$$

input `integrate(sec(2*t)/(1+sec(t)**2+3*tan(t)),t)`

output `Integral(sec(2*t)/(3*tan(t) + sec(t)**2 + 1), t)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 256 vs. 2(37) = 74.

Time = 0.18 (sec) , antiderivative size = 256, normalized size of antiderivative = 5.69

$$\int \frac{\sec(2t)}{1 + \sec^2(t) + 3 \tan(t)} dt$$

$$= \frac{3(\cos(2t)^2 + \sin(2t)^2 + 2 \sin(2t) + 1) \log(953674316406250(3 \cos(2t) + \sin(2t) + 4) \cos(4t) + 23)}{\dots}$$

input `integrate(sec(2*t)/(1+sec(t)^2+3*tan(t)),t, algorithm="maxima")`

output `1/48*(3*(cos(2*t)^2 + sin(2*t)^2 + 2*sin(2*t) + 1)*log(953674316406250*(3*cos(2*t) + sin(2*t) + 4)*cos(4*t) + 2384185791015625*cos(4*t)^2 + 953674316406250*cos(2*t)^2 - 953674316406250*(cos(2*t) - 3*sin(2*t) + 3)*sin(4*t) + 2384185791015625*sin(4*t)^2 + 953674316406250*sin(2*t)^2 + 2861022949218750*cos(2*t) - 953674316406250*sin(2*t) + 2384185791015625) - 6*(cos(2*t)^2 + sin(2*t)^2 + 2*sin(2*t) + 1)*log(cos(2*t)^2 + sin(2*t)^2 + 2*sin(2*t) + 1) + 5*(cos(2*t)^2 + sin(2*t)^2 + 2*sin(2*t) + 1)*log(1/5*(5*cos(2*t)^2 + 5*sin(2*t)^2 + 6*cos(2*t) + 8*sin(2*t) + 5)/(cos(2*t)^2 + sin(2*t)^2 - 2*sin(2*t) + 1)) - 24*cos(2*t))/(cos(2*t)^2 + sin(2*t)^2 + 2*sin(2*t) + 1)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.73

$$\int \frac{\sec(2t)}{1 + \sec^2(t) + 3 \tan(t)} dt = -\frac{1}{2(\tan(t) + 1)} + \frac{1}{3} \log(|\tan(t) + 2|) - \frac{1}{4} \log(|\tan(t) + 1|) - \frac{1}{12} \log(|\tan(t) - 1|)$$

input `integrate(sec(2*t)/(1+sec(t)^2+3*tan(t)),t, algorithm="giac")`

output `-1/2/(tan(t) + 1) + 1/3*log(abs(tan(t) + 2)) - 1/4*log(abs(tan(t) + 1)) - 1/12*log(abs(tan(t) - 1))`

Mupad [B] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.71

$$\int \frac{\sec(2t)}{1 + \sec^2(t) + 3 \tan(t)} dt = \frac{\ln(\tan(t) + 2)}{3} - \frac{\ln(\tan(t) + 1)}{4} - \frac{\ln(\tan(t) - 1)}{12} - \frac{1}{2(\tan(t) + 1)}$$

input `int(1/(cos(2*t)*(3*tan(t) + 1/cos(t)^2 + 1)),t)`

output $\log(\tan(t) + 2)/3 - \log(\tan(t) + 1)/4 - \log(\tan(t) - 1)/12 - 1/(2*(\tan(t) + 1))$

Reduce [F]

$$\int \frac{\sec(2t)}{1 + \sec^2(t) + 3 \tan(t)} dt = -\frac{\left(\int \frac{\sin(t)^2}{3 \cos(2t) \cos(t) \sin(t) - \cos(2t) \sin(t)^2 + 2 \cos(2t)} dt\right)}{2} - \frac{\left(\int \frac{\sin(t)^2}{3 \cos(t) \sin(t) - \sin(t)^2 + 2}\right)}{2} - \frac{3\left(\int \frac{\cos(t) \sin(t)}{3 \cos(2t) \cos(t) \sin(t) - \cos(2t) \sin(t)^2 + 2 \cos(2t)} dt\right)}{2} - \frac{3\left(\int \frac{\cos(t) \sin(t)}{3 \cos(t) \sin(t) - \sin(t)^2 + 2}\right)}{2} + \frac{3 \log\left(\tan\left(\frac{t}{2}\right)^2 + 1\right)}{10} + \frac{\log(-\sqrt{5} + 2 \tan\left(\frac{t}{2}\right) - 1)}{5} - \frac{\log(-\sqrt{2} + \tan\left(\frac{t}{2}\right) - 1)}{2} + \frac{\log(\sqrt{5} + 2 \tan\left(\frac{t}{2}\right) - 1)}{5} - \frac{\log(\sqrt{2} + \tan\left(\frac{t}{2}\right) - 1)}{2} - \frac{\log(\tan(t) - 1)}{4} + \frac{\log(\tan(t) + 1)}{4} + \frac{2t}{5}$$

input `int(sec(2*t)/(1+sec(t)^2+3*tan(t)),t)`

output `(- 10*int(sin(t)**2/(3*cos(2*t)*cos(t)*sin(t) - cos(2*t)*sin(t)**2 + 2*cos(2*t)),t) - 10*int(sin(t)**2/(3*cos(t)*sin(t) - sin(t)**2 + 2),t) - 30*int((cos(t)*sin(t))/(3*cos(2*t)*cos(t)*sin(t) - cos(2*t)*sin(t)**2 + 2*cos(2*t)),t) - 30*int((cos(t)*sin(t))/(3*cos(t)*sin(t) - sin(t)**2 + 2),t) + 6*log(tan(t/2)**2 + 1) + 4*log(-sqrt(5) + 2*tan(t/2) - 1) - 10*log(-sqrt(2) + tan(t/2) - 1) + 4*log(sqrt(5) + 2*tan(t/2) - 1) - 10*log(sqrt(2) + tan(t/2) - 1) - 5*log(tan(t) - 1) + 5*log(tan(t) + 1) + 8*t)/20`

3.88 $\int \cos^2(x) dx$

Optimal result	534
Mathematica [A] (verified)	534
Rubi [A] (verified)	535
Maple [A] (verified)	536
Fricas [A] (verification not implemented)	536
Sympy [A] (verification not implemented)	537
Maxima [A] (verification not implemented)	537
Giac [A] (verification not implemented)	537
Mupad [B] (verification not implemented)	538
Reduce [B] (verification not implemented)	538

Optimal result

Integrand size = 4, antiderivative size = 14

$$\int \cos^2(x) dx = \frac{x}{2} + \frac{1}{2} \cos(x) \sin(x)$$

output `1/2*x+1/2*cos(x)*sin(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \cos^2(x) dx = \frac{x}{2} + \frac{1}{4} \sin(2x)$$

input `Integrate[Cos[x]^2,x]`

output `x/2 + Sin[2*x]/4`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cos^2(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin\left(x + \frac{\pi}{2}\right)^2 dx \\ & \quad \downarrow \text{3115} \\ & \frac{\int 1 dx}{2} + \frac{1}{2} \sin(x) \cos(x) \\ & \quad \downarrow \text{24} \\ & \frac{x}{2} + \frac{1}{2} \sin(x) \cos(x) \end{aligned}$$

input `Int[Cos[x]^2,x]`

output `x/2 + (Cos[x]*Sin[x])/2`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.79

method	result	size
default	$\frac{x}{2} + \frac{\cos(x)\sin(x)}{2}$	11
risch	$\frac{x}{2} + \frac{\sin(2x)}{4}$	11
parallelrisc	$\frac{x}{2} + \frac{\sin(2x)}{4}$	11
norman	$\frac{x \tan(\frac{x}{2})^2 + \frac{x}{2} - \tan(\frac{x}{2})^3 + \frac{x \tan(\frac{x}{2})^4}{2} + \tan(\frac{x}{2})}{(1 + \tan(\frac{x}{2})^2)^2}$	45

input `int(1/sec(x)^2,x,method=_RETURNVERBOSE)`output `1/2*x+1/2*cos(x)*sin(x)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \cos^2(x) dx = \frac{1}{2} \cos(x) \sin(x) + \frac{1}{2} x$$

input `integrate(1/sec(x)^2,x, algorithm="fricas")`output `1/2*cos(x)*sin(x) + 1/2*x`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \cos^2(x) dx = \frac{x}{2} + \frac{\sin(x) \cos(x)}{2}$$

input `integrate(1/sec(x)**2,x)`output `x/2 + sin(x)*cos(x)/2`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \cos^2(x) dx = \frac{1}{2}x + \frac{1}{4} \sin(2x)$$

input `integrate(1/sec(x)^2,x, algorithm="maxima")`output `1/2*x + 1/4*sin(2*x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \cos^2(x) dx = \frac{1}{2}x + \frac{\tan(x)}{2(\tan(x)^2 + 1)}$$

input `integrate(1/sec(x)^2,x, algorithm="giac")`output `1/2*x + 1/2*tan(x)/(tan(x)^2 + 1)`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \cos^2(x) dx = \frac{x}{2} + \frac{\sin(2x)}{4}$$

input `int(cos(x)^2,x)`

output `x/2 + sin(2*x)/4`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \cos^2(x) dx = \frac{\cos(x) \sin(x)}{2} + \frac{x}{2}$$

input `int(1/sec(x)^2,x)`

output `(cos(x)*sin(x) + x)/2`

3.89 $\int \frac{1+x^2}{\sqrt{x}} dx$

Optimal result	539
Mathematica [A] (verified)	539
Rubi [A] (verified)	540
Maple [A] (verified)	541
Fricas [A] (verification not implemented)	541
Sympy [A] (verification not implemented)	542
Maxima [A] (verification not implemented)	542
Giac [A] (verification not implemented)	542
Mupad [B] (verification not implemented)	543
Reduce [B] (verification not implemented)	543

Optimal result

Integrand size = 11, antiderivative size = 17

$$\int \frac{1+x^2}{\sqrt{x}} dx = 2\sqrt{x} + \frac{2x^{5/2}}{5}$$

output `2/5*x^(5/2)+2*x^(1/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \frac{1+x^2}{\sqrt{x}} dx = \frac{2}{5}\sqrt{x}(5+x^2)$$

input `Integrate[(1 + x^2)/Sqrt[x],x]`

output `(2*Sqrt[x]*(5 + x^2))/5`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2 + 1}{\sqrt{x}} dx$$

↓ 244

$$\int \left(x^{3/2} + \frac{1}{\sqrt{x}} \right) dx$$

↓ 2009

$$\frac{2x^{5/2}}{5} + 2\sqrt{x}$$

input `Int[(1 + x^2)/Sqrt[x],x]`

output `2*Sqrt[x] + (2*x^(5/2))/5`

Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] :> Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

method	result	size
gospers	$\frac{2\sqrt{x}(x^2+5)}{5}$	11
risch	$\frac{2\sqrt{x}(x^2+5)}{5}$	11
orering	$\frac{2\sqrt{x}(x^2+5)}{5}$	11
derivativdivides	$\frac{2x^{\frac{5}{2}}}{5} + 2\sqrt{x}$	12
default	$\frac{2x^{\frac{5}{2}}}{5} + 2\sqrt{x}$	12
trager	$\left(\frac{2x^2}{5} + 2\right)\sqrt{x}$	12

input `int((x^2+1)/x^(1/2),x,method=_RETURNVERBOSE)`output `2/5*x^(1/2)*(x^2+5)`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.59

$$\int \frac{1+x^2}{\sqrt{x}} dx = \frac{2}{5}(x^2+5)\sqrt{x}$$

input `integrate((x^2+1)/x^(1/2),x, algorithm="fricas")`output `2/5*(x^2 + 5)*sqrt(x)`

Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \frac{1+x^2}{\sqrt{x}} dx = \frac{2x^{\frac{5}{2}}}{5} + 2\sqrt{x}$$

input `integrate((x**2+1)/x**(1/2),x)`

output `2*x**(5/2)/5 + 2*sqrt(x)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int \frac{1+x^2}{\sqrt{x}} dx = \frac{2}{5} x^{\frac{5}{2}} + 2\sqrt{x}$$

input `integrate((x^2+1)/x^(1/2),x, algorithm="maxima")`

output `2/5*x^(5/2) + 2*sqrt(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int \frac{1+x^2}{\sqrt{x}} dx = \frac{2}{5} x^{\frac{5}{2}} + 2\sqrt{x}$$

input `integrate((x^2+1)/x^(1/2),x, algorithm="giac")`

output `2/5*x^(5/2) + 2*sqrt(x)`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.59

$$\int \frac{1+x^2}{\sqrt{x}} dx = \frac{2\sqrt{x}(x^2+5)}{5}$$

input `int((x^2 + 1)/x^(1/2),x)`

output `(2*x^(1/2)*(x^2 + 5))/5`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.53

$$\int \frac{1+x^2}{\sqrt{x}} dx = \frac{2\sqrt{x}(x^2+5)}{5}$$

input `int((x^2+1)/x^(1/2),x)`

output `(2*sqrt(x)*(x**2 + 5))/5`

3.90 $\int \frac{x}{\sqrt{5+2x+x^2}} dx$

Optimal result	544
Mathematica [A] (verified)	544
Rubi [A] (verified)	545
Maple [A] (verified)	546
Fricas [A] (verification not implemented)	546
Sympy [A] (verification not implemented)	547
Maxima [A] (verification not implemented)	547
Giac [A] (verification not implemented)	547
Mupad [B] (verification not implemented)	548
Reduce [B] (verification not implemented)	548

Optimal result

Integrand size = 14, antiderivative size = 23

$$\int \frac{x}{\sqrt{5+2x+x^2}} dx = \sqrt{5+2x+x^2} - \operatorname{arcsinh}\left(\frac{1+x}{2}\right)$$

output `-arcsinh(1/2+1/2*x)+(x^2+2*x+5)^(1/2)`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.35

$$\int \frac{x}{\sqrt{5+2x+x^2}} dx = \sqrt{5+2x+x^2} + \log\left(-1-x+\sqrt{5+2x+x^2}\right)$$

input `Integrate[x/Sqrt[5 + 2*x + x^2],x]`

output `Sqrt[5 + 2*x + x^2] + Log[-1 - x + Sqrt[5 + 2*x + x^2]]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {1160, 1090, 222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x}{\sqrt{x^2 + 2x + 5}} dx \\ & \quad \downarrow \text{1160} \\ & \sqrt{x^2 + 2x + 5} - \int \frac{1}{\sqrt{x^2 + 2x + 5}} dx \\ & \quad \downarrow \text{1090} \\ & \sqrt{x^2 + 2x + 5} - \frac{1}{4} \int \frac{1}{\sqrt{\frac{1}{16}(2x + 2)^2 + 1}} d(2x + 2) \\ & \quad \downarrow \text{222} \\ & \sqrt{x^2 + 2x + 5} - \operatorname{arcsinh}\left(\frac{1}{4}(2x + 2)\right) \end{aligned}$$

input `Int[x/Sqrt[5 + 2*x + x^2],x]`

output `Sqrt[5 + 2*x + x^2] - ArcSinh[(2 + 2*x)/4]`

Defintions of rubi rules used

rule 222 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

rule 1090 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/(2*c*(-4*(c/(b^2 - 4*a*c))))^p) Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

rule 1160

```
Int[((d_.) + (e_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol
] := Simp[e*((a + b*x + c*x^2)^(p + 1)/(2*c*(p + 1))), x] + Simp[(2*c*d - b
*e)/(2*c) Int[(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x]
&& NeQ[p, -1]
```

Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

method	result	size
default	$-\operatorname{arcsinh}\left(\frac{1}{2} + \frac{x}{2}\right) + \sqrt{x^2 + 2x + 5}$	20
risch	$-\operatorname{arcsinh}\left(\frac{1}{2} + \frac{x}{2}\right) + \sqrt{x^2 + 2x + 5}$	20
trager	$\sqrt{x^2 + 2x + 5} - \ln(1 + x + \sqrt{x^2 + 2x + 5})$	28

input

```
int(x/(x^2+2*x+5)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
-arcsinh(1/2+1/2*x)+(x^2+2*x+5)^(1/2)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{x}{\sqrt{5 + 2x + x^2}} dx = \sqrt{x^2 + 2x + 5} + \log(-x + \sqrt{x^2 + 2x + 5} - 1)$$

input

```
integrate(x/(x^2+2*x+5)^(1/2),x, algorithm="fricas")
```

output

```
sqrt(x^2 + 2*x + 5) + log(-x + sqrt(x^2 + 2*x + 5) - 1)
```

Sympy [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{x}{\sqrt{5+2x+x^2}} dx = \sqrt{x^2+2x+5} - \operatorname{asinh}\left(\frac{x}{2} + \frac{1}{2}\right)$$

input `integrate(x/(x**2+2*x+5)**(1/2),x)`output `sqrt(x**2 + 2*x + 5) - asinh(x/2 + 1/2)`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{x}{\sqrt{5+2x+x^2}} dx = \sqrt{x^2+2x+5} - \operatorname{arsinh}\left(\frac{1}{2}x + \frac{1}{2}\right)$$

input `integrate(x/(x^2+2*x+5)^(1/2),x, algorithm="maxima")`output `sqrt(x^2 + 2*x + 5) - arcsinh(1/2*x + 1/2)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{x}{\sqrt{5+2x+x^2}} dx = \sqrt{x^2+2x+5} + \log\left(-x + \sqrt{x^2+2x+5} - 1\right)$$

input `integrate(x/(x^2+2*x+5)^(1/2),x, algorithm="giac")`output `sqrt(x^2 + 2*x + 5) + log(-x + sqrt(x^2 + 2*x + 5) - 1)`

Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{x}{\sqrt{5+2x+x^2}} dx = \sqrt{x^2+2x+5} - \ln(x + \sqrt{x^2+2x+5} + 1)$$

input `int(x/(2*x + x^2 + 5)^(1/2),x)`output `(2*x + x^2 + 5)^(1/2) - log(x + (2*x + x^2 + 5)^(1/2) + 1)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.26

$$\int \frac{x}{\sqrt{5+2x+x^2}} dx = \sqrt{x^2+2x+5} - \log\left(\frac{\sqrt{x^2+2x+5}}{2} + \frac{x}{2} + \frac{1}{2}\right)$$

input `int(x/(x^2+2*x+5)^(1/2),x)`output `sqrt(x**2 + 2*x + 5) - log((sqrt(x**2 + 2*x + 5) + x + 1)/2)`

3.91 $\int \cos(x) \sin^2(x) dx$

Optimal result	549
Mathematica [A] (verified)	549
Rubi [A] (verified)	550
Maple [A] (verified)	551
Fricas [A] (verification not implemented)	551
Sympy [A] (verification not implemented)	552
Maxima [A] (verification not implemented)	552
Giac [A] (verification not implemented)	552
Mupad [B] (verification not implemented)	553
Reduce [B] (verification not implemented)	553

Optimal result

Integrand size = 7, antiderivative size = 8

$$\int \cos(x) \sin^2(x) dx = \frac{\sin^3(x)}{3}$$

output `1/3*sin(x)^3`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \cos(x) \sin^2(x) dx = \frac{\sin^3(x)}{3}$$

input `Integrate[Cos[x]*Sin[x]^2,x]`

output `Sin[x]^3/3`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3044, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sin^2(x) \cos(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin(x)^2 \cos(x) dx \\ & \quad \downarrow \text{3044} \\ & \int \sin^2(x) d \sin(x) \\ & \quad \downarrow \text{15} \\ & \frac{\sin^3(x)}{3} \end{aligned}$$

input `Int[Cos[x]*Sin[x]^2,x]`

output `Sin[x]^3/3`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3044

```
Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)]^(m_.), x_
Symbol] := Simp[1/(a*f) Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a
*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(I
ntegerQ[(m - 1)/2] && LtQ[0, m, n])
```

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativdivides	$\frac{\sin(x)^3}{3}$	7
default	$\frac{\sin(x)^3}{3}$	7
orering	$\frac{\sin(x)^3}{3}$	7
risch	$\frac{\sin(x)}{4} - \frac{\sin(3x)}{12}$	12
parallelrisch	$\frac{\sin(x)}{4} - \frac{\sin(3x)}{12}$	12
norman	$\frac{8 \tan(\frac{x}{2})^3}{3(1 + \tan(\frac{x}{2})^2)^3}$	19

input

```
int(sin(x)^2*cos(x), x, method=_RETURNVERBOSE)
```

output

```
1/3*sin(x)^3
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.25

$$\int \cos(x) \sin^2(x) dx = -\frac{1}{3} (\cos(x)^2 - 1) \sin(x)$$

input

```
integrate(cos(x)*sin(x)^2,x, algorithm="fricas")
```

output

```
-1/3*(cos(x)^2 - 1)*sin(x)
```


Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int \cos(x) \sin^2(x) dx = \frac{\sin^3(x)}{3}$$

input `integrate(cos(x)*sin(x)**2,x)`

output `sin(x)**3/3`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos(x) \sin^2(x) dx = \frac{1}{3} \sin(x)^3$$

input `integrate(cos(x)*sin(x)^2,x, algorithm="maxima")`

output `1/3*sin(x)^3`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos(x) \sin^2(x) dx = \frac{1}{3} \sin(x)^3$$

input `integrate(cos(x)*sin(x)^2,x, algorithm="giac")`

output `1/3*sin(x)^3`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos(x) \sin^2(x) dx = \frac{\sin(x)^3}{3}$$

input `int(cos(x)*sin(x)^2,x)`

output `sin(x)^3/3`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos(x) \sin^2(x) dx = \frac{\sin(x)^3}{3}$$

input `int(cos(x)*sin(x)^2,x)`

output `sin(x)**3/3`

3.92 $\int \frac{e^x}{1+e^x} dx$

Optimal result	554
Mathematica [A] (verified)	554
Rubi [A] (verified)	555
Maple [A] (verified)	556
Fricas [A] (verification not implemented)	556
Sympy [A] (verification not implemented)	556
Maxima [A] (verification not implemented)	557
Giac [A] (verification not implemented)	557
Mupad [B] (verification not implemented)	557
Reduce [B] (verification not implemented)	558

Optimal result

Integrand size = 11, antiderivative size = 6

$$\int \frac{e^x}{1+e^x} dx = \log(1+e^x)$$

output `ln(1+exp(x))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{e^x}{1+e^x} dx = \log(1+e^x)$$

input `Integrate[E^x/(1 + E^x), x]`

output `Log[1 + E^x]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2676, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^x}{e^x + 1} dx$$

↓ 2676

$$\int \frac{1}{e^x + 1} de^x$$

↓ 16

$$\log(e^x + 1)$$

input `Int[E^x/(1 + E^x),x]`

output `Log[1 + E^x]`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 2676 `Int[((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)*((a_) + (b_.)*(F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.))^(p_.), x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[(a + b*x)^p, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n, p}, x]`

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

method	result	size
derivativedivides	$\ln(1 + e^x)$	6
default	$\ln(1 + e^x)$	6
norman	$\ln(1 + e^x)$	6
risch	$\ln(1 + e^x)$	6
parallelrisch	$\ln(1 + e^x)$	6

input `int(1/(1+exp(x))*exp(x),x,method=_RETURNVERBOSE)`output `ln(1+exp(x))`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int \frac{e^x}{1 + e^x} dx = \log(e^x + 1)$$

input `integrate(exp(x)/(1+exp(x)),x, algorithm="fricas")`output `log(e^x + 1)`**Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int \frac{e^x}{1 + e^x} dx = \log(e^x + 1)$$

input `integrate(exp(x)/(1+exp(x)),x)`

output `log(exp(x) + 1)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int \frac{e^x}{1 + e^x} dx = \log(e^x + 1)$$

input `integrate(exp(x)/(1+exp(x)),x, algorithm="maxima")`

output `log(e^x + 1)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int \frac{e^x}{1 + e^x} dx = \log(e^x + 1)$$

input `integrate(exp(x)/(1+exp(x)),x, algorithm="giac")`

output `log(e^x + 1)`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int \frac{e^x}{1 + e^x} dx = \ln(e^x + 1)$$

input `int(exp(x)/(exp(x) + 1),x)`

output `log(exp(x) + 1)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{e^x}{1 + e^x} dx = \log(e^x + 1)$$

input `int(exp(x)/(1+exp(x)), x)`

output `log(e**x + 1)`

3.93 $\int \frac{e^{2x}}{1+e^x} dx$

Optimal result	559
Mathematica [A] (verified)	559
Rubi [A] (verified)	560
Maple [A] (verified)	561
Fricas [A] (verification not implemented)	561
Sympy [A] (verification not implemented)	562
Maxima [A] (verification not implemented)	562
Giac [A] (verification not implemented)	562
Mupad [B] (verification not implemented)	563
Reduce [B] (verification not implemented)	563

Optimal result

Integrand size = 13, antiderivative size = 12

$$\int \frac{e^{2x}}{1+e^x} dx = e^x - \log(1+e^x)$$

output

```
exp(x)-ln(1+exp(x))
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{e^{2x}}{1+e^x} dx = e^x - \log(1+e^x)$$

input

```
Integrate[E^(2*x)/(1 + E^x),x]
```

output

```
E^x - Log[1 + E^x]
```


Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {2678, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{e^{2x}}{e^x + 1} dx \\ & \quad \downarrow \text{2678} \\ & \int \frac{e^x}{e^x + 1} de^x \\ & \quad \downarrow \text{49} \\ & \int \left(\frac{1}{-e^x - 1} + 1 \right) de^x \\ & \quad \downarrow \text{2009} \\ & e^x - \log(e^x + 1) \end{aligned}$$

input `Int[E^(2*x)/(1 + E^x), x]`

output `E^x - Log[1 + E^x]`

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2678

```
Int[((a_) + (b_)*(F_)^((e_)*((c_) + (d_)*(x_)))^(p_)*(G_)^((h_)*((f_
.) + (g_)*(x_))), x_Symbol] := With[{m = FullSimplify[g*h*(Log[G]/(d*e*Log
[F]))]}, Simp[Denominator[m]*(G^(f*h - c*g*(h/d))/(d*e*Log[F])) Subst[Int
[x^(Numerator[m] - 1)*(a + b*x^Denominator[m])^p, x], x, F^(e*((c + d*x)/De
nominator[m]))], x] /; LeQ[m, -1] || GeQ[m, 1]] /; FreeQ[{F, G, a, b, c, d,
e, f, g, h, p}, x]
```

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

method	result	size
default	$e^x - \ln(1 + e^x)$	11
norman	$e^x - \ln(1 + e^x)$	11
risch	$e^x - \ln(1 + e^x)$	11

input

```
int(exp(2*x)/(1+exp(x)),x,method=_RETURNVERBOSE)
```

output

```
exp(x)-ln(1+exp(x))
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{e^{2x}}{1 + e^x} dx = e^x - \log(e^x + 1)$$

input

```
integrate(exp(2*x)/(1+exp(x)),x, algorithm="fricas")
```

output

```
e^x - log(e^x + 1)
```

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{e^{2x}}{1+e^x} dx = e^x - \log(e^x + 1)$$

input `integrate(exp(2*x)/(1+exp(x)),x)`

output `exp(x) - log(exp(x) + 1)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{e^{2x}}{1+e^x} dx = e^x - \log(e^x + 1)$$

input `integrate(exp(2*x)/(1+exp(x)),x, algorithm="maxima")`

output `e^x - log(e^x + 1)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{e^{2x}}{1+e^x} dx = e^x - \log(e^x + 1)$$

input `integrate(exp(2*x)/(1+exp(x)),x, algorithm="giac")`

output `e^x - log(e^x + 1)`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{e^{2x}}{1 + e^x} dx = e^x - \ln(e^x + 1)$$

input `int(exp(2*x)/(exp(x) + 1),x)`

output `exp(x) - log(exp(x) + 1)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{e^{2x}}{1 + e^x} dx = e^x - \log(e^x + 1)$$

input `int(exp(2*x)/(1+exp(x)),x)`

output `e**x - log(e**x + 1)`

3.94 $\int \frac{1}{1-\cos(x)} dx$

Optimal result	564
Mathematica [A] (verified)	564
Rubi [A] (verified)	565
Maple [A] (verified)	566
Fricas [A] (verification not implemented)	566
Sympy [A] (verification not implemented)	566
Maxima [A] (verification not implemented)	567
Giac [A] (verification not implemented)	567
Mupad [B] (verification not implemented)	567
Reduce [B] (verification not implemented)	568

Optimal result

Integrand size = 8, antiderivative size = 12

$$\int \frac{1}{1-\cos(x)} dx = -\frac{\sin(x)}{1-\cos(x)}$$

output `-sin(x)/(1-cos(x))`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1}{1-\cos(x)} dx = -\cot\left(\frac{x}{2}\right)$$

input `Integrate[(1 - Cos[x])^(-1),x]`

output `-Cot[x/2]`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{1 - \cos(x)} dx$$

↓ 3042

$$\int \frac{1}{1 - \sin\left(x + \frac{\pi}{2}\right)} dx$$

↓ 3127

$$-\frac{\sin(x)}{1 - \cos(x)}$$

input `Int[(1 - Cos[x])^(-1), x]`

output `-(Sin[x]/(1 - Cos[x]))`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.58

method	result	size
parallelrisch	$-\cot\left(\frac{x}{2}\right)$	7
default	$-\frac{1}{\tan\left(\frac{x}{2}\right)}$	9
norman	$-\frac{1}{\tan\left(\frac{x}{2}\right)}$	9
risch	$-\frac{2i}{e^{ix}-1}$	13

input `int(1/(1-cos(x)),x,method=_RETURNVERBOSE)`output `-cot(1/2*x)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{1}{1 - \cos(x)} dx = -\frac{\cos(x) + 1}{\sin(x)}$$

input `integrate(1/(1-cos(x)),x, algorithm="fricas")`output `-(cos(x) + 1)/sin(x)`**Sympy [A] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.58

$$\int \frac{1}{1 - \cos(x)} dx = -\frac{1}{\tan\left(\frac{x}{2}\right)}$$

input `integrate(1/(1-cos(x)),x)`

output `-1/tan(x/2)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{1}{1 - \cos(x)} dx = -\frac{\cos(x) + 1}{\sin(x)}$$

input `integrate(1/(1-cos(x)),x, algorithm="maxima")`

output `-(cos(x) + 1)/sin(x)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1}{1 - \cos(x)} dx = -\frac{1}{\tan\left(\frac{1}{2}x\right)}$$

input `integrate(1/(1-cos(x)),x, algorithm="giac")`

output `-1/tan(1/2*x)`

Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.50

$$\int \frac{1}{1 - \cos(x)} dx = -\cot\left(\frac{x}{2}\right)$$

input `int(-1/(cos(x) - 1),x)`

output `-cot(x/2)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1}{1 - \cos(x)} dx = -\frac{1}{\tan\left(\frac{x}{2}\right)}$$

input `int(1/(1-cos(x)),x)`

output `(- 1)/tan(x/2)`

3.95 $\int \sec^2(x) \tan(x) dx$

Optimal result	569
Mathematica [A] (verified)	569
Rubi [A] (verified)	570
Maple [A] (verified)	571
Fricas [A] (verification not implemented)	571
Sympy [A] (verification not implemented)	572
Maxima [A] (verification not implemented)	572
Giac [A] (verification not implemented)	572
Mupad [B] (verification not implemented)	573
Reduce [B] (verification not implemented)	573

Optimal result

Integrand size = 7, antiderivative size = 8

$$\int \sec^2(x) \tan(x) dx = \frac{\sec^2(x)}{2}$$

output `1/2*sec(x)^2`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \sec^2(x) \tan(x) dx = \frac{\sec^2(x)}{2}$$

input `Integrate[Sec[x]^2*Tan[x],x]`

output `Sec[x]^2/2`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3086, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tan(x) \sec^2(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \tan(x) \sec(x)^2 dx \\ & \quad \downarrow \text{3086} \\ & \int \sec(x) d\sec(x) \\ & \quad \downarrow \text{15} \\ & \frac{\sec^2(x)}{2} \end{aligned}$$

input `Int[Sec[x]^2*Tan[x],x]`

output `Sec[x]^2/2`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086

```
Int[((a_.)*sec[(e_.) + (f_.)*(x_.)]^(m_.))*((b_.)*tan[(e_.) + (f_.)*(x_.)]^(n_.), x_Symbol] :> Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])
```

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$\frac{\sec(x)^2}{2}$	7
default	$\frac{\sec(x)^2}{2}$	7
risch	$\frac{2e^{2ix}}{(e^{2ix}+1)^2}$	17

input

```
int(sec(x)^2*tan(x), x, method=_RETURNVERBOSE)
```

output

```
1/2*sec(x)^2
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^2(x) \tan(x) dx = \frac{1}{2 \cos(x)^2}$$

input

```
integrate(sec(x)^2*tan(x), x, algorithm="fricas")
```

output

```
1/2/cos(x)^2
```

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int \sec^2(x) \tan(x) dx = \frac{1}{2 \cos^2(x)}$$

input `integrate(sec(x)**2*tan(x),x)`

output `1/(2*cos(x)**2)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^2(x) \tan(x) dx = \frac{1}{2} \tan(x)^2$$

input `integrate(sec(x)^2*tan(x),x, algorithm="maxima")`

output `1/2*tan(x)^2`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^2(x) \tan(x) dx = \frac{1}{2 \cos(x)^2}$$

input `integrate(sec(x)^2*tan(x),x, algorithm="giac")`

output `1/2/cos(x)^2`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^2(x) \tan(x) dx = \frac{\tan(x)^2}{2}$$

input `int(tan(x)/cos(x)^2,x)`

output `tan(x)^2/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^2(x) \tan(x) dx = \frac{\sec(x)^2}{2}$$

input `int(sec(x)^2*tan(x),x)`

output `sec(x)**2/2`

3.96 $\int x \log(x) dx$

Optimal result	574
Mathematica [A] (verified)	574
Rubi [A] (verified)	575
Maple [A] (verified)	576
Fricas [A] (verification not implemented)	576
Sympy [A] (verification not implemented)	577
Maxima [A] (verification not implemented)	577
Giac [A] (verification not implemented)	577
Mupad [B] (verification not implemented)	578
Reduce [B] (verification not implemented)	578

Optimal result

Integrand size = 4, antiderivative size = 17

$$\int x \log(x) dx = -\frac{x^2}{4} + \frac{1}{2}x^2 \log(x)$$

output

```
-1/4*x^2+1/2*x^2*ln(x)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int x \log(x) dx = -\frac{x^2}{4} + \frac{1}{2}x^2 \log(x)$$

input

```
Integrate[x*Log[x],x]
```

output

```
-1/4*x^2 + (x^2*Log[x])/2
```

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2741}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \log(x) dx$$

↓ 2741

$$\frac{1}{2}x^2 \log(x) - \frac{x^2}{4}$$

input `Int [x*Log [x] , x]`

output `-1/4*x^2 + (x^2*Log [x])/2`

Defintions of rubi rules used

rule 2741 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((d_.)*(x_)^(m_.), x_Symbol] :>
Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])/(d*(m + 1))), x] - Simp[b*n*((d*x)^(
m + 1)/(d*(m + 1)^2)), x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2}$	14
norman	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2}$	14
risch	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2}$	14
parallelrisch	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2}$	14
parts	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2}$	14
orering	$\frac{3x^2 \ln(x)}{4} - \frac{x^2(1+\ln(x))}{4}$	18

input `int(x*ln(x),x,method=_RETURNVERBOSE)`output `-1/4*x^2+1/2*x^2*ln(x)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int x \log(x) dx = \frac{1}{2} x^2 \log(x) - \frac{1}{4} x^2$$

input `integrate(x*log(x),x, algorithm="fricas")`output `1/2*x^2*log(x) - 1/4*x^2`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

$$\int x \log(x) dx = \frac{x^2 \log(x)}{2} - \frac{x^2}{4}$$

input `integrate(x*ln(x),x)`

output `x**2*log(x)/2 - x**2/4`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int x \log(x) dx = \frac{1}{2} x^2 \log(x) - \frac{1}{4} x^2$$

input `integrate(x*log(x),x, algorithm="maxima")`

output `1/2*x^2*log(x) - 1/4*x^2`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int x \log(x) dx = \frac{1}{2} x^2 \log(x) - \frac{1}{4} x^2$$

input `integrate(x*log(x),x, algorithm="giac")`

output `1/2*x^2*log(x) - 1/4*x^2`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.53

$$\int x \log(x) dx = \frac{x^2 (\ln(x) - \frac{1}{2})}{2}$$

input `int(x*log(x),x)`

output `(x^2*(log(x) - 1/2))/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int x \log(x) dx = \frac{x^2(2 \log(x) - 1)}{4}$$

input `int(x*log(x),x)`

output `(x**2*(2*log(x) - 1))/4`

3.97 $\int \cos(x) \sin(x) dx$

Optimal result	579
Mathematica [A] (verified)	579
Rubi [A] (verified)	580
Maple [A] (verified)	581
Fricas [A] (verification not implemented)	581
Sympy [A] (verification not implemented)	582
Maxima [A] (verification not implemented)	582
Giac [A] (verification not implemented)	582
Mupad [B] (verification not implemented)	583
Reduce [B] (verification not implemented)	583

Optimal result

Integrand size = 5, antiderivative size = 8

$$\int \cos(x) \sin(x) dx = \frac{\sin^2(x)}{2}$$

output `1/2*sin(x)^2`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \cos(x) \sin(x) dx = -\frac{1}{2} \cos^2(x)$$

input `Integrate[Cos[x]*Sin[x],x]`

output `-1/2*Cos[x]^2`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3042, 3044, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \sin(x) \cos(x) dx \\ \downarrow \text{3042} \\ \int \sin(x) \cos(x) dx \\ \downarrow \text{3044} \\ \int \sin(x) d \sin(x) \\ \downarrow \text{15} \\ \frac{\sin^2(x)}{2} \end{array}$$

input `Int[Cos[x]*Sin[x],x]`

output `Sin[x]^2/2`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3044

```
Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)]^(m_.), x_
Symbol] := Simp[1/(a*f) Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a
*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(I
ntegerQ[(m - 1)/2] && LtQ[0, m, n])
```

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativdivides	$\frac{\sin(x)^2}{2}$	7
default	$\frac{\sin(x)^2}{2}$	7
risch	$-\frac{\cos(2x)}{4}$	7
parallelrisch	$\frac{1}{4} - \frac{\cos(2x)}{4}$	9
oring	$\frac{\sin(x)^2}{4} - \frac{\cos(x)^2}{4}$	14
norman	$\frac{2 \tan(\frac{x}{2})^2}{(1 + \tan(\frac{x}{2})^2)^2}$	19
meijerg	$\frac{\sqrt{\pi} \left(\frac{1}{\sqrt{\pi}} - \frac{\cos(2x)}{\sqrt{\pi}} \right)}{4}$	19

input

```
int(cos(x)*sin(x), x, method=_RETURNVERBOSE)
```

output

```
1/2*sin(x)^2
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos(x) \sin(x) dx = -\frac{1}{2} \cos(x)^2$$

input

```
integrate(cos(x)*sin(x), x, algorithm="fricas")
```

output `-1/2*cos(x)^2`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int \cos(x) \sin(x) dx = \frac{\sin^2(x)}{2}$$

input `integrate(cos(x)*sin(x),x)`

output `sin(x)**2/2`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos(x) \sin(x) dx = -\frac{1}{2} \cos(x)^2$$

input `integrate(cos(x)*sin(x),x, algorithm="maxima")`

output `-1/2*cos(x)^2`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos(x) \sin(x) dx = -\frac{1}{2} \cos(x)^2$$

input `integrate(cos(x)*sin(x),x, algorithm="giac")`

output `-1/2*cos(x)^2`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos(x) \sin(x) dx = \frac{\sin(x)^2}{2}$$

input `int(cos(x)*sin(x),x)`

output `sin(x)^2/2`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos(x) \sin(x) dx = -\frac{\cos(x)^2}{2}$$

input `int(cos(x)*sin(x),x)`

output `(- cos(x)**2)/2`

3.98 $\int \frac{1+x}{\sqrt{2x-x^2}} dx$

Optimal result	584
Mathematica [A] (verified)	584
Rubi [A] (verified)	585
Maple [A] (verified)	586
Fricas [A] (verification not implemented)	586
Sympy [A] (verification not implemented)	587
Maxima [A] (verification not implemented)	587
Giac [A] (verification not implemented)	587
Mupad [B] (verification not implemented)	588
Reduce [B] (verification not implemented)	588

Optimal result

Integrand size = 17, antiderivative size = 24

$$\int \frac{1+x}{\sqrt{2x-x^2}} dx = -\sqrt{2x-x^2} - 2 \arcsin(1-x)$$

output `2*arcsin(-1+x)-(-x^2+2*x)^(1/2)`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.96

$$\int \frac{1+x}{\sqrt{2x-x^2}} dx = \frac{(-2+x)x - 4\sqrt{-2+x}\sqrt{x} \log(\sqrt{-2+x} - \sqrt{x})}{\sqrt{-((-2+x)x)}}$$

input `Integrate[(1 + x)/Sqrt[2*x - x^2], x]`

output `((-2 + x)*x - 4*Sqrt[-2 + x]*Sqrt[x]*Log[Sqrt[-2 + x] - Sqrt[x]])/Sqrt[-((-2 + x)*x)]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.17, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {1160, 1090, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x+1}{\sqrt{2x-x^2}} dx \\ & \quad \downarrow \text{1160} \\ & 2 \int \frac{1}{\sqrt{2x-x^2}} dx - \sqrt{2x-x^2} \\ & \quad \downarrow \text{1090} \\ & - \int \frac{1}{\sqrt{1-\frac{1}{4}(2-2x)^2}} d(2-2x) - \sqrt{2x-x^2} \\ & \quad \downarrow \text{223} \\ & -2 \arcsin\left(\frac{1}{2}(2-2x)\right) - \sqrt{2x-x^2} \end{aligned}$$

input `Int[(1 + x)/Sqrt[2*x - x^2],x]`

output `-Sqrt[2*x - x^2] - 2*ArcSin[(2 - 2*x)/2]`

Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 1090 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p) Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

rule 1160

```
Int[((d_.) + (e_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol
] :> Simp[e*((a + b*x + c*x^2)^(p + 1)/(2*c*(p + 1))), x] + Simp[(2*c*d - b
*e)/(2*c) Int[(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x]
&& NeQ[p, -1]
```

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.88

method	result	size
default	$2 \arcsin(-1 + x) - \sqrt{-x^2 + 2x}$	21
risch	$\frac{x(-2+x)}{\sqrt{-x(-2+x)}} + 2 \arcsin(-1 + x)$	21
pseudoelliptic	$-\sqrt{-x(-2+x)} - 4 \arctan\left(\frac{\sqrt{-x(-2+x)}}{x}\right)$	27
trager	$-\sqrt{-x^2 + 2x} + 2 \operatorname{RootOf}(_Z^2 + 1) \ln(\operatorname{RootOf}(_Z^2 + 1) \sqrt{-x^2 + 2x} + x - 1)$	45
meijerg	$2 \arcsin\left(\frac{\sqrt{2}\sqrt{x}}{2}\right) + \frac{2i\left(\frac{i\sqrt{\pi}\sqrt{x}\sqrt{2}\sqrt{1-\frac{x}{2}}}{2} - i\sqrt{\pi} \arcsin\left(\frac{\sqrt{2}\sqrt{x}}{2}\right)\right)}{\sqrt{\pi}}$	54

```
input int((1+x)/(-x^2+2*x)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 2*arcsin(-1+x)-(-x^2+2*x)^(1/2)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.42

$$\int \frac{1+x}{\sqrt{2x-x^2}} dx = -\sqrt{-x^2+2x} - 4 \arctan\left(\frac{\sqrt{-x^2+2x}}{x-2}\right)$$

```
input integrate((1+x)/(-x^2+2*x)^(1/2),x, algorithm="fricas")
```

```
output -sqrt(-x^2 + 2*x) - 4*arctan(sqrt(-x^2 + 2*x)/(x - 2))
```

Sympy [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.62

$$\int \frac{1+x}{\sqrt{2x-x^2}} dx = -\sqrt{-x^2+2x} + 2 \operatorname{asin}(x-1)$$

input `integrate((1+x)/(-x**2+2*x)**(1/2),x)`output `-sqrt(-x**2 + 2*x) + 2*asin(x - 1)`**Maxima [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1+x}{\sqrt{2x-x^2}} dx = -\sqrt{-x^2+2x} - 2 \operatorname{arcsin}(-x+1)$$

input `integrate((1+x)/(-x^2+2*x)^(1/2),x, algorithm="maxima")`output `-sqrt(-x^2 + 2*x) - 2*arcsin(-x + 1)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{1+x}{\sqrt{2x-x^2}} dx = -\sqrt{-x^2+2x} + 2 \operatorname{arcsin}(x-1)$$

input `integrate((1+x)/(-x^2+2*x)^(1/2),x, algorithm="giac")`output `-sqrt(-x^2 + 2*x) + 2*arcsin(x - 1)`

Mupad [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{1+x}{\sqrt{2x-x^2}} dx = 2 \operatorname{asin}(x-1) - \sqrt{2x-x^2}$$

input `int((x + 1)/(2*x - x^2)^(1/2),x)`output `2*asin(x - 1) - (2*x - x^2)^(1/2)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.29

$$\int \frac{1+x}{\sqrt{2x-x^2}} dx = -\sqrt{x}\sqrt{-x+2} - 4 \log\left(\frac{\sqrt{-x+2} + \sqrt{x}i}{\sqrt{2}}\right) i$$

input `int((1+x)/(-x^2+2*x)^(1/2),x)`output `- sqrt(x)*sqrt(- x + 2) - 4*log((sqrt(- x + 2) + sqrt(x)*i)/sqrt(2))*i`

3.99 $\int \frac{2e^x}{2+3e^{2x}} dx$

Optimal result	589
Mathematica [A] (verified)	589
Rubi [A] (verified)	590
Maple [A] (verified)	591
Fricas [A] (verification not implemented)	591
Sympy [A] (verification not implemented)	592
Maxima [A] (verification not implemented)	592
Giac [A] (verification not implemented)	592
Mupad [B] (verification not implemented)	593
Reduce [B] (verification not implemented)	593

Optimal result

Integrand size = 16, antiderivative size = 20

$$\int \frac{2e^x}{2+3e^{2x}} dx = \sqrt{\frac{2}{3}} \arctan \left(\sqrt{\frac{3}{2}} e^x \right)$$

output `1/3*arctan(1/2*exp(x)*6^(1/2))*6^(1/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{2e^x}{2+3e^{2x}} dx = \sqrt{\frac{2}{3}} \arctan \left(\sqrt{\frac{3}{2}} e^x \right)$$

input `Integrate[(2*E^x)/(2 + 3*E^(2*x)),x]`

output `Sqrt[2/3]*ArcTan[Sqrt[3/2]*E^x]`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {27, 2679, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{2e^x}{3e^{2x} + 2} dx \\ & \quad \downarrow \text{27} \\ & 2 \int \frac{e^x}{2 + 3e^{2x}} dx \\ & \quad \downarrow \text{2679} \\ & 2 \int \frac{1}{2 + 3e^{2x}} de^x \\ & \quad \downarrow \text{216} \\ & \sqrt{\frac{2}{3}} \arctan \left(\sqrt{\frac{3}{2}} e^x \right) \end{aligned}$$

input `Int[(2*E^x)/(2 + 3*E^(2*x)),x]`

output `Sqrt[2/3]*ArcTan[Sqrt[3/2]*E^x]`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] :> Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 2679

```
Int[((a_) + (b_)*(F_)^((e_)*((c_) + (d_)*(x_)))^(p_)*(G_)^((h_)*((f_
.) + (g_)*(x_))), x_Symbol] := With[{m = FullSimplify[d*e*(Log[F]/(g*h*Log
[G]))]}, Simp[Denominator[m]/(g*h*Log[G]) Subst[Int[x^(Denominator[m] - 1
)*(a + b*F^(c*e - d*e*(f/g))*x^Numerator[m])^p, x], x, G^(h*((f + g*x)/Deno
minator[m]))], x] /; LtQ[m, -1] || GtQ[m, 1]] /; FreeQ[{F, G, a, b, c, d, e
, f, g, h, p}, x]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.70

method	result	size
default	$\frac{\arctan\left(\frac{e^x\sqrt{6}}{2}\right)\sqrt{6}}{3}$	14
risch	$\frac{i\sqrt{6}\ln\left(e^x + \frac{i\sqrt{6}}{3}\right)}{6} - \frac{i\sqrt{6}\ln\left(e^x - \frac{i\sqrt{6}}{3}\right)}{6}$	34

input

```
int(2*exp(x)/(2+3*exp(2*x)),x,method=_RETURNVERBOSE)
```

output

```
1/3*arctan(1/2*exp(x)*6^(1/2))*6^(1/2)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.60

$$\int \frac{2e^x}{2 + 3e^{2x}} dx = \sqrt{\frac{2}{3}} \arctan\left(\frac{3}{2} \sqrt{\frac{2}{3}} e^x\right)$$

input

```
integrate(2*exp(x)/(2+3*exp(2*x)),x, algorithm="fricas")
```

output

```
sqrt(2/3)*arctan(3/2*sqrt(2/3)*e^x)
```


Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

$$\int \frac{2e^x}{2 + 3e^{2x}} dx = \text{RootSum}(6z^2 + 1, (i \mapsto i \log(2i + e^x)))$$

input `integrate(2*exp(x)/(2+3*exp(2*x)),x)`output `RootSum(6*_z**2 + 1, Lambda(_i, _i*log(2*_i + exp(x))))`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.65

$$\int \frac{2e^x}{2 + 3e^{2x}} dx = \frac{1}{3} \sqrt{6} \arctan\left(\frac{1}{2} \sqrt{6} e^x\right)$$

input `integrate(2*exp(x)/(2+3*exp(2*x)),x, algorithm="maxima")`output `1/3*sqrt(6)*arctan(1/2*sqrt(6)*e^x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.65

$$\int \frac{2e^x}{2 + 3e^{2x}} dx = \frac{1}{3} \sqrt{6} \arctan\left(\frac{1}{2} \sqrt{6} e^x\right)$$

input `integrate(2*exp(x)/(2+3*exp(2*x)),x, algorithm="giac")`output `1/3*sqrt(6)*arctan(1/2*sqrt(6)*e^x)`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.65

$$\int \frac{2e^x}{2 + 3e^{2x}} dx = \frac{\sqrt{6} \operatorname{atan}\left(\frac{\sqrt{6}e^x}{2}\right)}{3}$$

input `int((2*exp(x))/(3*exp(2*x) + 2),x)`output `(6^(1/2)*atan((6^(1/2)*exp(x))/2))/3`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.70

$$\int \frac{2e^x}{2 + 3e^{2x}} dx = \frac{\sqrt{6} \operatorname{atan}\left(\frac{3e^x}{\sqrt{6}}\right)}{3}$$

input `int(2*exp(x)/(2+3*exp(2*x)),x)`output `(sqrt(6)*atan((3*e**x)/sqrt(6)))/3`

$$3.100 \quad \int \frac{x^4}{(1-x^2)^{5/2}} dx$$

Optimal result	594
Mathematica [A] (verified)	594
Rubi [A] (verified)	595
Maple [A] (verified)	596
Fricas [B] (verification not implemented)	596
Sympy [B] (verification not implemented)	597
Maxima [A] (verification not implemented)	597
Giac [A] (verification not implemented)	598
Mupad [B] (verification not implemented)	598
Reduce [B] (verification not implemented)	598

Optimal result

Integrand size = 15, antiderivative size = 35

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{x^3}{3(1-x^2)^{3/2}} - \frac{x}{\sqrt{1-x^2}} + \arcsin(x)$$

output `1/3*x^3/(-x^2+1)^(3/2)+arcsin(x)-x/(-x^2+1)^(1/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.26

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{x(-3+4x^2)}{3(1-x^2)^{3/2}} + 2 \arctan\left(\frac{x}{-1+\sqrt{1-x^2}}\right)$$

input `Integrate[x^4/(1-x^2)^(5/2),x]`

output `(x*(-3+4*x^2))/(3*(1-x^2)^(3/2))+2*ArcTan[x/(-1+Sqrt[1-x^2])]`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {252, 252, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^4}{(1-x^2)^{5/2}} dx \\ & \quad \downarrow \text{252} \\ & \frac{x^3}{3(1-x^2)^{3/2}} - \int \frac{x^2}{(1-x^2)^{3/2}} dx \\ & \quad \downarrow \text{252} \\ & \int \frac{1}{\sqrt{1-x^2}} dx - \frac{x}{\sqrt{1-x^2}} + \frac{x^3}{3(1-x^2)^{3/2}} \\ & \quad \downarrow \text{223} \\ & \arcsin(x) - \frac{x}{\sqrt{1-x^2}} + \frac{x^3}{3(1-x^2)^{3/2}} \end{aligned}$$

input `Int[x^4/(1 - x^2)^(5/2),x]`

output `x^3/(3*(1 - x^2)^(3/2)) - x/Sqrt[1 - x^2] + ArcSin[x]`

Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 252

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(2*b*(p + 1))), x] - Simp[c^2*(m - 1)/(2*b*(p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && LtQ[p, -1] && GtQ[m, 1] && !ILtQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]
```

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.86

method	result	size
default	$\frac{x^3}{3(-x^2+1)^{\frac{3}{2}}} + \arcsin(x) - \frac{x}{\sqrt{-x^2+1}}$	30
risch	$-\frac{x(4x^2-3)}{3(x^2-1)\sqrt{-x^2+1}} + \arcsin(x)$	30
meijerg	$-\frac{2i \left(-\frac{i\sqrt{\pi}x(-20x^2+15)}{10(-x^2+1)^{\frac{3}{2}}} + \frac{3i\sqrt{\pi}\arcsin(x)}{2} \right)}{3\sqrt{\pi}}$	39
pseudoelliptic	$\frac{(x^2-1)\sqrt{-x^2+1}\arctan\left(\frac{\sqrt{-x^2+1}}{x}\right) + \frac{4x^3}{3} - x}{(-x^2+1)^{\frac{3}{2}}}$	49
trager	$\frac{(4x^2-3)x\sqrt{-x^2+1}}{3(x^2-1)^2} + \text{RootOf}(_Z^2 + 1) \ln(\text{RootOf}(_Z^2 + 1)\sqrt{-x^2+1} + x)$	54

input `int(x^4/(-x^2+1)^(5/2),x,method=_RETURNVERBOSE)`output `1/3*x^3/(-x^2+1)^(3/2)+arcsin(x)-1/(-x^2+1)^(1/2)*x`**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 63 vs. 2(29) = 58.

Time = 0.07 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.80

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = -\frac{6(x^4 - 2x^2 + 1) \arctan\left(\frac{\sqrt{-x^2+1}-1}{x}\right) - (4x^3 - 3x)\sqrt{-x^2+1}}{3(x^4 - 2x^2 + 1)}$$

input `integrate(x^4/(-x^2+1)^(5/2),x, algorithm="fricas")`

output `-1/3*(6*(x^4 - 2*x^2 + 1)*arctan((sqrt(-x^2 + 1) - 1)/x) - (4*x^3 - 3*x)*sqrt(-x^2 + 1))/(x^4 - 2*x^2 + 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 105 vs. $2(26) = 52$.

Time = 0.34 (sec) , antiderivative size = 105, normalized size of antiderivative = 3.00

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{3x^4 \operatorname{asin}(x)}{3x^4 - 6x^2 + 3} + \frac{4x^3 \sqrt{1-x^2}}{3x^4 - 6x^2 + 3} - \frac{6x^2 \operatorname{asin}(x)}{3x^4 - 6x^2 + 3} - \frac{3x \sqrt{1-x^2}}{3x^4 - 6x^2 + 3} + \frac{3 \operatorname{asin}(x)}{3x^4 - 6x^2 + 3}$$

input `integrate(x**4/(-x**2+1)**(5/2),x)`

output `3*x**4*asin(x)/(3*x**4 - 6*x**2 + 3) + 4*x**3*sqrt(1 - x**2)/(3*x**4 - 6*x**2 + 3) - 6*x**2*asin(x)/(3*x**4 - 6*x**2 + 3) - 3*x*sqrt(1 - x**2)/(3*x**4 - 6*x**2 + 3) + 3*asin(x)/(3*x**4 - 6*x**2 + 3)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.26

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{1}{3} x \left(\frac{3x^2}{(-x^2+1)^{3/2}} - \frac{2}{(-x^2+1)^{3/2}} \right) - \frac{x}{3\sqrt{-x^2+1}} + \operatorname{arcsin}(x)$$

input `integrate(x^4/(-x^2+1)^(5/2),x, algorithm="maxima")`

output `1/3*x*(3*x^2/(-x^2 + 1)^(3/2) - 2/(-x^2 + 1)^(3/2)) - 1/3*x/sqrt(-x^2 + 1) + arcsin(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.83

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{(4x^2-3)\sqrt{-x^2+1}x}{3(x^2-1)^2} + \arcsin(x)$$

input `integrate(x^4/(-x^2+1)^(5/2),x, algorithm="giac")`output `1/3*(4*x^2 - 3)*sqrt(-x^2 + 1)*x/(x^2 - 1)^2 + arcsin(x)`**Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 91, normalized size of antiderivative = 2.60

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \arcsin(x) + \frac{3\sqrt{1-x^2}}{4(x-1)} + \frac{3\sqrt{1-x^2}}{4(x+1)} - \sqrt{1-x^2} \left(\frac{1}{12(x-1)} - \frac{1}{12(x-1)^2} \right) - \sqrt{1-x^2} \left(\frac{1}{12(x+1)} + \frac{1}{12(x+1)^2} \right)$$

input `int(x^4/(1 - x^2)^(5/2),x)`output `asin(x) + (3*(1 - x^2)^(1/2))/(4*(x - 1)) + (3*(1 - x^2)^(1/2))/(4*(x + 1)) - (1 - x^2)^(1/2)*(1/(12*(x - 1)) - 1/(12*(x - 1)^2)) - (1 - x^2)^(1/2)*(1/(12*(x + 1)) + 1/(12*(x + 1)^2))`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.57

$$\int \frac{x^4}{(1-x^2)^{5/2}} dx = \frac{3\sqrt{-x^2+1} \arcsin(x) x^2 - 3\sqrt{-x^2+1} \arcsin(x) - 4x^3 + 3x}{3\sqrt{-x^2+1} (x^2 - 1)}$$

input `int(x^4/(-x^2+1)^(5/2),x)`

output
$$\frac{(3\sqrt{-x^2 + 1})\operatorname{asin}(x)x^{**2} - 3\sqrt{-x^2 + 1}\operatorname{asin}(x) - 4x^{**3} + 3x}{(3\sqrt{-x^2 + 1})(x^{**2} - 1)}$$

3.101 $\int \frac{e^{6x}}{1+e^{4x}} dx$

Optimal result	600
Mathematica [A] (verified)	600
Rubi [A] (verified)	601
Maple [A] (verified)	602
Fricas [A] (verification not implemented)	602
Sympy [A] (verification not implemented)	603
Maxima [A] (verification not implemented)	603
Giac [A] (verification not implemented)	603
Mupad [B] (verification not implemented)	604
Reduce [B] (verification not implemented)	604

Optimal result

Integrand size = 15, antiderivative size = 20

$$\int \frac{e^{6x}}{1+e^{4x}} dx = \frac{e^{2x}}{2} - \frac{1}{2} \arctan(e^{2x})$$

output `1/2*exp(2*x)-1/2*arctan(exp(2*x))`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{e^{6x}}{1+e^{4x}} dx = \frac{1}{2}(e^{2x} - \arctan(e^{2x}))$$

input `Integrate[E^(6*x)/(1 + E^(4*x)),x]`

output `(E^(2*x) - ArcTan[E^(2*x)])/2`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2678, 262, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{e^{6x}}{e^{4x} + 1} dx \\ & \quad \downarrow \text{2678} \\ & \frac{1}{2} \int \frac{e^{4x}}{1 + e^{4x}} de^{2x} \\ & \quad \downarrow \text{262} \\ & \frac{1}{2} \left(e^{2x} - \int \frac{1}{1 + e^{4x}} de^{2x} \right) \\ & \quad \downarrow \text{216} \\ & \frac{1}{2} (e^{2x} - \arctan(e^{2x})) \end{aligned}$$

input `Int[E^(6*x)/(1 + E^(4*x)),x]`

output `(E^(2*x) - ArcTan[E^(2*x)])/2`

Defintions of rubi rules used

rule 216

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

rule 262

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)
^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/
(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b
, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c
, 2, m, p, x]
```

rule 2678

```
Int[((a_) + (b_.)*(F_)^((e_.)*((c_.) + (d_.)*(x_))))^(p_.)*(G_)^((h_.)*((f_
.) + (g_.)*(x_))), x_Symbol] := With[{m = FullSimplify[g*h*(Log[G]/(d*e*Log
[F]))}], Simp[Denominator[m]*(G^(f*h - c*g*(h/d))/(d*e*Log[F])) Subst[Int
[x^(Numerator[m] - 1)*(a + b*x^Denominator[m])^p, x], x, F^(e*((c + d*x)/De
nominator[m]))], x] /; LeQ[m, -1] || GeQ[m, 1]] /; FreeQ[{F, G, a, b, c, d,
e, f, g, h, p}, x]
```

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

method	result	size
default	$\frac{e^{2x}}{2} - \frac{\arctan(e^{2x})}{2}$	15
risch	$\frac{e^{2x}}{2} + \frac{i \ln(e^{2x} - i)}{4} - \frac{i \ln(e^{2x} + i)}{4}$	30

input

```
int(exp(6*x)/(1+exp(4*x)),x,method=_RETURNVERBOSE)
```

output

```
1/2*exp(x)^2-1/2*arctan(exp(x)^2)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.70

$$\int \frac{e^{6x}}{1 + e^{4x}} dx = -\frac{1}{2} \arctan(e^{(2x)}) + \frac{1}{2} e^{(2x)}$$

input

```
integrate(exp(6*x)/(1+exp(4*x)),x, algorithm="fricas")
```

output `-1/2*arctan(e^(2*x)) + 1/2*e^(2*x)`

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.20

$$\int \frac{e^{6x}}{1 + e^{4x}} dx = \frac{e^{2x}}{2} + \text{RootSum}(16z^2 + 1, (i \mapsto i \log(-4i + e^{2x})))$$

input `integrate(exp(6*x)/(1+exp(4*x)),x)`

output `exp(2*x)/2 + RootSum(16*_z**2 + 1, Lambda(_i, _i*log(-4*_i + exp(2*x))))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.70

$$\int \frac{e^{6x}}{1 + e^{4x}} dx = -\frac{1}{2} \arctan(e^{(2x)}) + \frac{1}{2} e^{(2x)}$$

input `integrate(exp(6*x)/(1+exp(4*x)),x, algorithm="maxima")`

output `-1/2*arctan(e^(2*x)) + 1/2*e^(2*x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.70

$$\int \frac{e^{6x}}{1 + e^{4x}} dx = -\frac{1}{2} \arctan(e^{(2x)}) + \frac{1}{2} e^{(2x)}$$

input `integrate(exp(6*x)/(1+exp(4*x)),x, algorithm="giac")`

output `-1/2*arctan(e^(2*x)) + 1/2*e^(2*x)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.70

$$\int \frac{e^{6x}}{1 + e^{4x}} dx = \frac{e^{2x}}{2} - \frac{\operatorname{atan}(e^{2x})}{2}$$

input `int(exp(6*x)/(exp(4*x) + 1),x)`output `exp(2*x)/2 - atan(exp(2*x))/2`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 42, normalized size of antiderivative = 2.10

$$\int \frac{e^{6x}}{1 + e^{4x}} dx = -\frac{\operatorname{atan}\left(\frac{2e^x - \sqrt{2}}{\sqrt{2}}\right)}{2} + \frac{\operatorname{atan}\left(\frac{2e^x + \sqrt{2}}{\sqrt{2}}\right)}{2} + \frac{e^{2x}}{2}$$

input `int(exp(6*x)/(1+exp(4*x)),x)`output `(- atan((2*e**x - sqrt(2))/sqrt(2)) + atan((2*e**x + sqrt(2))/sqrt(2)) + e**(2*x))/2`

3.102 $\int \log(2 + 3x^2) dx$

Optimal result	605
Mathematica [A] (verified)	605
Rubi [A] (verified)	606
Maple [A] (verified)	607
Fricas [A] (verification not implemented)	607
Sympy [A] (verification not implemented)	608
Maxima [A] (verification not implemented)	608
Giac [A] (verification not implemented)	608
Mupad [B] (verification not implemented)	609
Reduce [B] (verification not implemented)	609

Optimal result

Integrand size = 8, antiderivative size = 33

$$\int \log(2 + 3x^2) dx = -2x + 2\sqrt{\frac{2}{3}} \arctan\left(\sqrt{\frac{3}{2}}x\right) + x \log(2 + 3x^2)$$

output

```
-2*x+x*ln(3*x^2+2)+2/3*arctan(1/2*x*6^(1/2))*6^(1/2)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int \log(2 + 3x^2) dx = -2x + 2\sqrt{\frac{2}{3}} \arctan\left(\sqrt{\frac{3}{2}}x\right) + x \log(2 + 3x^2)$$

input

```
Integrate[Log[2 + 3*x^2], x]
```

output

```
-2*x + 2*Sqrt[2/3]*ArcTan[Sqrt[3/2]*x] + x*Log[2 + 3*x^2]
```

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.21, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {2898, 262, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \log(3x^2 + 2) dx \\ & \quad \downarrow \text{2898} \\ & x \log(3x^2 + 2) - 6 \int \frac{x^2}{3x^2 + 2} dx \\ & \quad \downarrow \text{262} \\ & x \log(3x^2 + 2) - 6 \left(\frac{x}{3} - \frac{2}{3} \int \frac{1}{3x^2 + 2} dx \right) \\ & \quad \downarrow \text{216} \\ & x \log(3x^2 + 2) - 6 \left(\frac{x}{3} - \frac{1}{3} \sqrt{\frac{2}{3}} \arctan \left(\sqrt{\frac{3}{2}} x \right) \right) \end{aligned}$$

input `Int[Log[2 + 3*x^2], x]`

output `-6*(x/3 - (Sqrt[2/3]*ArcTan[Sqrt[3/2]*x])/3) + x*Log[2 + 3*x^2]`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 262

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)
^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/
(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b
, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c
, 2, m, p, x]
```

rule 2898

```
Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_))^(p_.)], x_Symbol] := Simp[x*Log[c*(d
+ e*x^n)^p], x] - Simp[e*n*p Int[x^n/(d + e*x^n), x], x] /; FreeQ[{c, d,
e, n, p}, x]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.82

method	result	size
default	$-2x + x \ln(3x^2 + 2) + \frac{2 \arctan\left(\frac{x\sqrt{6}}{2}\right)\sqrt{6}}{3}$	27
risch	$-2x + x \ln(3x^2 + 2) + \frac{2 \arctan\left(\frac{x\sqrt{6}}{2}\right)\sqrt{6}}{3}$	27
parts	$-2x + x \ln(3x^2 + 2) + \frac{2 \arctan\left(\frac{x\sqrt{6}}{2}\right)\sqrt{6}}{3}$	27

input `int(ln(3*x^2+2),x,method=_RETURNVERBOSE)`output `-2*x+x*ln(3*x^2+2)+2/3*arctan(1/2*x*6^(1/2))*6^(1/2)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.79

$$\int \log(2 + 3x^2) dx = x \log(3x^2 + 2) + 2 \sqrt{\frac{2}{3}} \arctan\left(\frac{3}{2} \sqrt{\frac{2}{3}} x\right) - 2x$$

input `integrate(log(3*x^2+2),x, algorithm="fricas")`

output `x*log(3*x^2 + 2) + 2*sqrt(2/3)*arctan(3/2*sqrt(2/3)*x) - 2*x`

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.94

$$\int \log(2 + 3x^2) dx = x \log(3x^2 + 2) - 2x + \frac{2\sqrt{6} \operatorname{atan}\left(\frac{\sqrt{6}x}{2}\right)}{3}$$

input `integrate(ln(3*x**2+2),x)`

output `x*log(3*x**2 + 2) - 2*x + 2*sqrt(6)*atan(sqrt(6)*x/2)/3`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.79

$$\int \log(2 + 3x^2) dx = x \log(3x^2 + 2) + \frac{2}{3} \sqrt{6} \arctan\left(\frac{1}{2} \sqrt{6}x\right) - 2x$$

input `integrate(log(3*x^2+2),x, algorithm="maxima")`

output `x*log(3*x^2 + 2) + 2/3*sqrt(6)*arctan(1/2*sqrt(6)*x) - 2*x`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.79

$$\int \log(2 + 3x^2) dx = x \log(3x^2 + 2) + \frac{2}{3} \sqrt{6} \arctan\left(\frac{1}{2} \sqrt{6}x\right) - 2x$$

input `integrate(log(3*x^2+2),x, algorithm="giac")`

output $x \log(3x^2 + 2) + \frac{2}{3} \sqrt{6} \arctan\left(\frac{1}{2} \sqrt{6} x\right) - 2x$

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.79

$$\int \log(2 + 3x^2) dx = \frac{2\sqrt{6} \operatorname{atan}\left(\frac{\sqrt{6}x}{2}\right)}{3} - 2x + x \ln(3x^2 + 2)$$

input `int(log(3*x^2 + 2),x)`

output $(2 \cdot 6^{(1/2)} \cdot \operatorname{atan}((6^{(1/2)} \cdot x)/2))/3 - 2 \cdot x + x \cdot \log(3 \cdot x^2 + 2)$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.79

$$\int \log(2 + 3x^2) dx = \frac{2\sqrt{6} \operatorname{atan}\left(\frac{3x}{\sqrt{6}}\right)}{3} + \log(3x^2 + 2) x - 2x$$

input `int(log(3*x^2+2),x)`

output $(2 \cdot \sqrt{6} \cdot \operatorname{atan}((3 \cdot x)/\sqrt{6}) + 3 \cdot \log(3 \cdot x^2 + 2) \cdot x - 6 \cdot x)/3$

3.103 $\int \frac{1}{r\sqrt{-a^2+2Hr^2}} dx$

Optimal result	610
Mathematica [A] (verified)	610
Rubi [A] (verified)	611
Maple [A] (verified)	611
Fricas [A] (verification not implemented)	612
Sympy [A] (verification not implemented)	612
Maxima [A] (verification not implemented)	613
Giac [A] (verification not implemented)	613
Mupad [B] (verification not implemented)	613
Reduce [B] (verification not implemented)	614

Optimal result

Integrand size = 20, antiderivative size = 21

$$\int \frac{1}{r\sqrt{-a^2+2Hr^2}} dx = \frac{x}{r\sqrt{-a^2+2Hr^2}}$$

output $x/r/(2*H*r^2-a^2)^{(1/2)}$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{1}{r\sqrt{-a^2+2Hr^2}} dx = \frac{x}{r\sqrt{-a^2+2Hr^2}}$$

input `Integrate[1/(r*Sqrt[-a^2 + 2*H*r^2]),x]`

output $x/(r*\text{Sqrt}[-a^2 + 2*H*r^2])$

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$, Rules used = {24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{r\sqrt{2Hr^2 - a^2}} dx$$

↓ 24

$$\frac{x}{r\sqrt{2Hr^2 - a^2}}$$

input `Int[1/(r*Sqrt[-a^2 + 2*H*r^2]),x]`

output `x/(r*Sqrt[-a^2 + 2*H*r^2])`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.95

method	result	size
default	$\frac{x}{r\sqrt{2Hr^2 - a^2}}$	20
norman	$\frac{x}{r\sqrt{2Hr^2 - a^2}}$	20
parallelrisch	$\frac{x}{r\sqrt{2Hr^2 - a^2}}$	20
orering	$\frac{x}{r\sqrt{2Hr^2 - a^2}}$	20

input `int(1/r/(2*H*r^2-a^2)^(1/2),x,method=_RETURNVERBOSE)`

output `x/r/(2*H*r^2-a^2)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.48

$$\int \frac{1}{r\sqrt{-a^2 + 2Hr^2}} dx = \frac{\sqrt{2Hr^2 - a^2}x}{2Hr^3 - a^2r}$$

input `integrate(1/r/(2*H*r^2-a^2)^(1/2),x, algorithm="fricas")`

output `sqrt(2*H*r^2 - a^2)*x/(2*H*r^3 - a^2*r)`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int \frac{1}{r\sqrt{-a^2 + 2Hr^2}} dx = \frac{x}{r\sqrt{2Hr^2 - a^2}}$$

input `integrate(1/r/(2*H*r**2-a**2)**(1/2),x)`

output `x/(r*sqrt(2*H*r**2 - a**2))`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{1}{r\sqrt{-a^2 + 2Hr^2}} dx = \frac{x}{\sqrt{2Hr^2 - a^2}r}$$

input `integrate(1/r/(2*H*r^2-a^2)^(1/2),x, algorithm="maxima")`output `x/(sqrt(2*H*r^2 - a^2)*r)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{1}{r\sqrt{-a^2 + 2Hr^2}} dx = \frac{x}{\sqrt{2Hr^2 - a^2}r}$$

input `integrate(1/r/(2*H*r^2-a^2)^(1/2),x, algorithm="giac")`output `x/(sqrt(2*H*r^2 - a^2)*r)`**Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{1}{r\sqrt{-a^2 + 2Hr^2}} dx = \frac{x}{r\sqrt{2Hr^2 - a^2}}$$

input `int(1/(r*(2*H*r^2 - a^2)^(1/2)),x)`output `x/(r*(2*H*r^2 - a^2)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.48

$$\int \frac{1}{r\sqrt{-a^2 + 2Hr^2}} dx = -\frac{\sqrt{2hr^2 - a^2} x}{r(-2hr^2 + a^2)}$$

input `int(1/r/(2*H*r^2-a^2)^(1/2),x)`

output `(- sqrt(- a**2 + 2*h*r**2)*x)/(r*(a**2 - 2*h*r**2))`

3.104 $\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2}} dx$

Optimal result	615
Mathematica [A] (verified)	615
Rubi [A] (verified)	616
Maple [A] (verified)	616
Fricas [A] (verification not implemented)	617
Sympy [A] (verification not implemented)	617
Maxima [A] (verification not implemented)	618
Giac [A] (verification not implemented)	618
Mupad [B] (verification not implemented)	618
Reduce [B] (verification not implemented)	619

Optimal result

Integrand size = 25, antiderivative size = 26

$$\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2}} dx = \frac{x}{r\sqrt{-a^2 - e^2 + 2Hr^2}}$$

output $x/r/(2*H*r^2 - a^2 - e^2)^{(1/2)}$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2}} dx = \frac{x}{r\sqrt{-a^2 - e^2 + 2Hr^2}}$$

input `Integrate[1/(r*Sqrt[-a^2 - e^2 + 2*H*r^2]),x]`

output $x/(r*\text{Sqrt}[-a^2 - e^2 + 2*H*r^2])$

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2}} dx$$

↓ 24

$$\frac{x}{r\sqrt{-a^2 - e^2 + 2Hr^2}}$$

input `Int[1/(r*Sqrt[-a^2 - e^2 + 2*H*r^2]),x]`

output `x/(r*Sqrt[-a^2 - e^2 + 2*H*r^2])`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.96

method	result	size
default	$\frac{x}{r\sqrt{2Hr^2 - a^2 - e^2}}$	25
norman	$\frac{x}{r\sqrt{2Hr^2 - a^2 - e^2}}$	25
parallelrisc	$\frac{x}{r\sqrt{2Hr^2 - a^2 - e^2}}$	25
orering	$\frac{x}{r\sqrt{2Hr^2 - a^2 - e^2}}$	25

input `int(1/r/(2*H*r^2-a^2-e^2)^(1/2),x,method=_RETURNVERBOSE)`

output `x/r/(2*H*r^2-a^2-e^2)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.54

$$\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2}} dx = \frac{\sqrt{2Hr^2 - a^2 - e^2}x}{2Hr^3 - (a^2 + e^2)r}$$

input `integrate(1/r/(2*H*r^2-a^2-e^2)^(1/2),x, algorithm="fricas")`

output `sqrt(2*H*r^2 - a^2 - e^2)*x/(2*H*r^3 - (a^2 + e^2)*r)`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.73

$$\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2}} dx = \frac{x}{r\sqrt{2Hr^2 - a^2 - e^2}}$$

input `integrate(1/r/(2*H*r**2-a**2-e**2)**(1/2),x)`

output `x/(r*sqrt(2*H*r**2 - a**2 - e**2))`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

$$\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2}} dx = \frac{x}{\sqrt{2Hr^2 - a^2 - e^2}r}$$

input `integrate(1/r/(2*H*r^2-a^2-e^2)^(1/2),x, algorithm="maxima")`output `x/(sqrt(2*H*r^2 - a^2 - e^2)*r)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

$$\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2}} dx = \frac{x}{\sqrt{2Hr^2 - a^2 - e^2}r}$$

input `integrate(1/r/(2*H*r^2-a^2-e^2)^(1/2),x, algorithm="giac")`output `x/(sqrt(2*H*r^2 - a^2 - e^2)*r)`**Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

$$\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2}} dx = \frac{x}{r\sqrt{-a^2 - e^2 + 2Hr^2}}$$

input `int(1/(r*(2*H*r^2 - a^2 - e^2)^(1/2)),x)`output `x/(r*(2*H*r^2 - a^2 - e^2)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.50

$$\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2}} dx = -\frac{\sqrt{2hr^2 - a^2 - e^2} x}{r(-2hr^2 + a^2 + e^2)}$$

input `int(1/r/(2*H*r^2-a^2-e^2)^(1/2),x)`

output `(- sqrt(- a**2 - e**2 + 2*h*r**2)*x)/(r*(a**2 + e**2 - 2*h*r**2))`

$$3.105 \quad \int \frac{1}{r\sqrt{-a^2+2Hr^2-2Kr^4}} dx$$

Optimal result	620
Mathematica [A] (verified)	620
Rubi [A] (verified)	621
Maple [A] (verified)	621
Fricas [A] (verification not implemented)	622
Sympy [A] (verification not implemented)	622
Maxima [A] (verification not implemented)	623
Giac [A] (verification not implemented)	623
Mupad [B] (verification not implemented)	623
Reduce [B] (verification not implemented)	624

Optimal result

Integrand size = 26, antiderivative size = 27

$$\int \frac{1}{r\sqrt{-a^2+2Hr^2-2Kr^4}} dx = \frac{x}{r\sqrt{-a^2+2Hr^2-2Kr^4}}$$

output `x/r/(-2*K*r^4+2*H*r^2-a^2)^(1/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{1}{r\sqrt{-a^2+2Hr^2-2Kr^4}} dx = \frac{x}{r\sqrt{-a^2+2Hr^2-2Kr^4}}$$

input `Integrate[1/(r*Sqrt[-a^2 + 2*H*r^2 - 2*K*r^4]),x]`

output `x/(r*Sqrt[-a^2 + 2*H*r^2 - 2*K*r^4])`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.038$, Rules used = {24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{r\sqrt{-a^2 + 2Hr^2 - 2Kr^4}} dx$$

\downarrow 24
 x

$$\frac{x}{r\sqrt{-a^2 + 2Hr^2 - 2Kr^4}}$$

input `Int[1/(r*Sqrt[-a^2 + 2*H*r^2 - 2*K*r^4]),x]`

output `x/(r*Sqrt[-a^2 + 2*H*r^2 - 2*K*r^4])`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

method	result	size
default	$\frac{x}{r\sqrt{-2Kr^4+2Hr^2-a^2}}$	26
norman	$\frac{x}{r\sqrt{-2Kr^4+2Hr^2-a^2}}$	26
parallelrisch	$\frac{x}{r\sqrt{-2Kr^4+2Hr^2-a^2}}$	26
orering	$\frac{x}{r\sqrt{-2Kr^4+2Hr^2-a^2}}$	26

input `int(1/r/(-2*K*r^4+2*H*r^2-a^2)^(1/2),x,method=_RETURNVERBOSE)`

output `x/r/(-2*K*r^4+2*H*r^2-a^2)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.59

$$\int \frac{1}{r\sqrt{-a^2 + 2Hr^2 - 2Kr^4}} dx = -\frac{\sqrt{-2Kr^4 + 2Hr^2 - a^2}x}{2Kr^5 - 2Hr^3 + a^2r}$$

input `integrate(1/r/(-2*K*r^4+2*H*r^2-a^2)^(1/2),x, algorithm="fricas")`

output `-sqrt(-2*K*r^4 + 2*H*r^2 - a^2)*x/(2*K*r^5 - 2*H*r^3 + a^2*r)`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int \frac{1}{r\sqrt{-a^2 + 2Hr^2 - 2Kr^4}} dx = \frac{x}{r\sqrt{2Hr^2 - 2Kr^4 - a^2}}$$

input `integrate(1/r/(-2*K*r**4+2*H*r**2-a**2)**(1/2),x)`

output `x/(r*sqrt(2*H*r**2 - 2*K*r**4 - a**2))`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.93

$$\int \frac{1}{r\sqrt{-a^2 + 2Hr^2 - 2Kr^4}} dx = \frac{x}{\sqrt{-2Kr^4 + 2Hr^2 - a^2}r}$$

input `integrate(1/r/(-2*K*r^4+2*H*r^2-a^2)^(1/2),x, algorithm="maxima")`output `x/(sqrt(-2*K*r^4 + 2*H*r^2 - a^2)*r)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.93

$$\int \frac{1}{r\sqrt{-a^2 + 2Hr^2 - 2Kr^4}} dx = \frac{x}{\sqrt{-2Kr^4 + 2Hr^2 - a^2}r}$$

input `integrate(1/r/(-2*K*r^4+2*H*r^2-a^2)^(1/2),x, algorithm="giac")`output `x/(sqrt(-2*K*r^4 + 2*H*r^2 - a^2)*r)`**Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.93

$$\int \frac{1}{r\sqrt{-a^2 + 2Hr^2 - 2Kr^4}} dx = \frac{x}{r\sqrt{-a^2 - 2Kr^4 + 2Hr^2}}$$

input `int(1/(r*(2*H*r^2 - 2*K*r^4 - a^2)^(1/2)),x)`output `x/(r*(2*H*r^2 - 2*K*r^4 - a^2)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.59

$$\int \frac{1}{r\sqrt{-a^2 + 2Hr^2 - 2Kr^4}} dx = -\frac{\sqrt{-2kr^4 + 2hr^2 - a^2} x}{r(2kr^4 - 2hr^2 + a^2)}$$

input `int(1/r/(-2*K*r^4+2*H*r^2-a^2)^(1/2),x)`

output `(- sqrt(- a**2 + 2*h*r**2 - 2*k*r**4)*x)/(r*(a**2 - 2*h*r**2 + 2*k*r**4))`

3.106 $\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2 - 2Kr^4}} dx$

Optimal result	625
Mathematica [A] (verified)	625
Rubi [A] (verified)	626
Maple [A] (verified)	626
Fricas [A] (verification not implemented)	627
Sympy [A] (verification not implemented)	627
Maxima [A] (verification not implemented)	628
Giac [A] (verification not implemented)	628
Mupad [B] (verification not implemented)	628
Reduce [B] (verification not implemented)	629

Optimal result

Integrand size = 31, antiderivative size = 32

$$\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2 - 2Kr^4}} dx = \frac{x}{r\sqrt{-a^2 - e^2 + 2Hr^2 - 2Kr^4}}$$

output `x/r/(-2*K*r^4+2*H*r^2-a^2-e^2)^(1/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

$$\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2 - 2Kr^4}} dx = \frac{x}{r\sqrt{-a^2 - e^2 + 2Hr^2 - 2Kr^4}}$$

input `Integrate[1/(r*Sqrt[-a^2 - e^2 + 2*H*r^2 - 2*K*r^4]),x]`

output `x/(r*Sqrt[-a^2 - e^2 + 2*H*r^2 - 2*K*r^4])`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.032$, Rules used = {24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2 - 2Kr^4}} dx$$

\downarrow 24
 x

$$\frac{x}{r\sqrt{-a^2 - e^2 + 2Hr^2 - 2Kr^4}}$$

input `Int[1/(r*Sqrt[-a^2 - e^2 + 2*H*r^2 - 2*K*r^4]),x]`

output `x/(r*Sqrt[-a^2 - e^2 + 2*H*r^2 - 2*K*r^4])`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.97

method	result	size
default	$\frac{x}{r\sqrt{-2K r^4+2H r^2-a^2-e^2}}$	31
norman	$\frac{x}{r\sqrt{-2K r^4+2H r^2-a^2-e^2}}$	31
parallelrisc	$\frac{x}{r\sqrt{-2K r^4+2H r^2-a^2-e^2}}$	31
orering	$\frac{x}{r\sqrt{-2K r^4+2H r^2-a^2-e^2}}$	31

input `int(1/r/(-2*K*r^4+2*H*r^2-a^2-e^2)^(1/2),x,method=_RETURNVERBOSE)`

output `x/r/(-2*K*r^4+2*H*r^2-a^2-e^2)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.62

$$\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2 - 2Kr^4}} dx = -\frac{\sqrt{-2Kr^4 + 2Hr^2 - a^2 - e^2}x}{2Kr^5 - 2Hr^3 + (a^2 + e^2)r}$$

input `integrate(1/r/(-2*K*r^4+2*H*r^2-a^2-e^2)^(1/2),x, algorithm="fricas")`

output `-sqrt(-2*K*r^4 + 2*H*r^2 - a^2 - e^2)*x/(2*K*r^5 - 2*H*r^3 + (a^2 + e^2)*r)`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.81

$$\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2 - 2Kr^4}} dx = \frac{x}{r\sqrt{2Hr^2 - 2Kr^4 - a^2 - e^2}}$$

input `integrate(1/r/(-2*K*r**4+2*H*r**2-a**2-e**2)**(1/2),x)`

output `x/(r*sqrt(2*H*r**2 - 2*K*r**4 - a**2 - e**2))`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.94

$$\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2 - 2Kr^4}} dx = \frac{x}{\sqrt{-2Kr^4 + 2Hr^2 - a^2 - e^2}r}$$

input `integrate(1/r/(-2*K*r^4+2*H*r^2-a^2-e^2)^(1/2),x, algorithm="maxima")`output `x/(sqrt(-2*K*r^4 + 2*H*r^2 - a^2 - e^2)*r)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.94

$$\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2 - 2Kr^4}} dx = \frac{x}{\sqrt{-2Kr^4 + 2Hr^2 - a^2 - e^2}r}$$

input `integrate(1/r/(-2*K*r^4+2*H*r^2-a^2-e^2)^(1/2),x, algorithm="giac")`output `x/(sqrt(-2*K*r^4 + 2*H*r^2 - a^2 - e^2)*r)`**Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.94

$$\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2 - 2Kr^4}} dx = \frac{x}{r\sqrt{-a^2 - e^2 - 2Kr^4 + 2Hr^2}}$$

input `int(1/(r*(2*H*r^2 - 2*K*r^4 - a^2 - e^2)^(1/2)),x)`output `x/(r*(2*H*r^2 - 2*K*r^4 - a^2 - e^2)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.59

$$\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2 - 2Kr^4}} dx = -\frac{\sqrt{-2kr^4 + 2hr^2 - a^2 - e^2} x}{r(2kr^4 - 2hr^2 + a^2 + e^2)}$$

input `int(1/r/(-2*K*r^4+2*H*r^2-a^2-e^2)^(1/2),x)`output `(- sqrt(- a**2 - e**2 + 2*h*r**2 - 2*k*r**4)*x)/(r*(a**2 + e**2 - 2*h*r**2 + 2*k*r**4))`

3.107 $\int \frac{1}{r\sqrt{-a^2-2Kr+2Hr^2}} dx$

Optimal result	630
Mathematica [A] (verified)	630
Rubi [A] (verified)	631
Maple [A] (verified)	631
Fricas [A] (verification not implemented)	632
Sympy [A] (verification not implemented)	632
Maxima [A] (verification not implemented)	633
Giac [A] (verification not implemented)	633
Mupad [B] (verification not implemented)	633
Reduce [B] (verification not implemented)	634

Optimal result

Integrand size = 24, antiderivative size = 24

$$\int \frac{1}{r\sqrt{-a^2-2Kr+2Hr^2}} dx = \frac{x}{r\sqrt{-a^2-2r(K-Hr)}}$$

output `x/r/(-a^2-2*r*(-H*r+K))^(1/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.04

$$\int \frac{1}{r\sqrt{-a^2-2Kr+2Hr^2}} dx = \frac{x}{r\sqrt{-a^2-2Kr+2Hr^2}}$$

input `Integrate[1/(r*Sqrt[-a^2 - 2*K*r + 2*H*r^2]),x]`

output `x/(r*Sqrt[-a^2 - 2*K*r + 2*H*r^2])`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$, Rules used = {24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{r\sqrt{-a^2 + 2Hr^2 - 2Kr}} dx$$

$$\downarrow 24$$

$$\frac{x}{r\sqrt{-a^2 - 2r(K - Hr)}}$$

input `Int[1/(r*Sqrt[-a^2 - 2*K*r + 2*H*r^2]),x]`

output `x/(r*Sqrt[-a^2 - 2*r*(K - H*r)])`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

method	result	size
default	$\frac{x}{r\sqrt{2Hr^2 - 2Kr - a^2}}$	24
norman	$\frac{x}{r\sqrt{2Hr^2 - 2Kr - a^2}}$	24
parallelrisch	$\frac{x}{r\sqrt{2Hr^2 - 2Kr - a^2}}$	24
orering	$\frac{x}{r\sqrt{2Hr^2 - 2Kr - a^2}}$	24

input `int(1/r/(2*H*r^2-2*K*r-a^2)^(1/2),x,method=_RETURNVERBOSE)`

output `1/r/(2*H*r^2-2*K*r-a^2)^(1/2)*x`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.71

$$\int \frac{1}{r\sqrt{-a^2 - 2Kr + 2Hr^2}} dx = \frac{\sqrt{2Hr^2 - a^2 - 2Krx}}{2Hr^3 - a^2r - 2Kr^2}$$

input `integrate(1/r/(2*H*r^2-2*K*r-a^2)^(1/2),x, algorithm="fricas")`

output `sqrt(2*H*r^2 - a^2 - 2*K*r)*x/(2*H*r^3 - a^2*r - 2*K*r^2)`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{1}{r\sqrt{-a^2 - 2Kr + 2Hr^2}} dx = \frac{x}{r\sqrt{2Hr^2 - 2Kr - a^2}}$$

input `integrate(1/r/(2*H*r**2-2*K*r-a**2)**(1/2),x)`

output `x/(r*sqrt(2*H*r**2 - 2*K*r - a**2))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.96

$$\int \frac{1}{r\sqrt{-a^2 - 2Kr + 2Hr^2}} dx = \frac{x}{\sqrt{2Hr^2 - a^2 - 2Krr}}$$

input `integrate(1/r/(2*H*r^2-2*K*r-a^2)^(1/2),x, algorithm="maxima")`output `x/(sqrt(2*H*r^2 - a^2 - 2*K*r)*r)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.96

$$\int \frac{1}{r\sqrt{-a^2 - 2Kr + 2Hr^2}} dx = \frac{x}{\sqrt{2Hr^2 - a^2 - 2Krr}}$$

input `integrate(1/r/(2*H*r^2-2*K*r-a^2)^(1/2),x, algorithm="giac")`output `x/(sqrt(2*H*r^2 - a^2 - 2*K*r)*r)`**Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.96

$$\int \frac{1}{r\sqrt{-a^2 - 2Kr + 2Hr^2}} dx = \frac{x}{r\sqrt{-a^2 + 2Hr^2 - 2Kr}}$$

input `int(1/(r*(2*H*r^2 - 2*K*r - a^2)^(1/2)),x)`output `x/(r*(2*H*r^2 - 2*K*r - a^2)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.62

$$\int \frac{1}{r\sqrt{-a^2 - 2Kr + 2Hr^2}} dx = -\frac{\sqrt{2hr^2 - a^2 - 2kr} x}{r(-2hr^2 + a^2 + 2kr)}$$

input `int(1/r/(2*H*r^2-2*K*r-a^2)^(1/2),x)`

output `(- sqrt(- a**2 + 2*h*r**2 - 2*k*r)*x)/(r*(a**2 - 2*h*r**2 + 2*k*r))`

3.108 $\int \frac{1}{r\sqrt{-a^2 - e^2 - 2Kr + 2Hr^2}} dx$

Optimal result	635
Mathematica [A] (verified)	635
Rubi [A] (verified)	636
Maple [A] (verified)	636
Fricas [A] (verification not implemented)	637
Sympy [A] (verification not implemented)	637
Maxima [A] (verification not implemented)	638
Giac [A] (verification not implemented)	638
Mupad [B] (verification not implemented)	638
Reduce [B] (verification not implemented)	639

Optimal result

Integrand size = 29, antiderivative size = 29

$$\int \frac{1}{r\sqrt{-a^2 - e^2 - 2Kr + 2Hr^2}} dx = \frac{x}{r\sqrt{-a^2 - e^2 - 2r(K - Hr)}}$$

output `x/r/(2*H*r^2-2*K*r-a^2-e^2)^(1/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.03

$$\int \frac{1}{r\sqrt{-a^2 - e^2 - 2Kr + 2Hr^2}} dx = \frac{x}{r\sqrt{-a^2 - e^2 - 2Kr + 2Hr^2}}$$

input `Integrate[1/(r*Sqrt[-a^2 - e^2 - 2*K*r + 2*H*r^2]),x]`

output `x/(r*Sqrt[-a^2 - e^2 - 2*K*r + 2*H*r^2])`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.034$, Rules used = {24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{r\sqrt{-a^2 - e^2 + 2Hr^2 - 2Kr}} dx$$

\downarrow 24
 x

$$\frac{x}{r\sqrt{-a^2 - e^2 - 2r(K - Hr)}}$$

input `Int[1/(r*Sqrt[-a^2 - e^2 - 2*K*r + 2*H*r^2]),x]`

output `x/(r*Sqrt[-a^2 - e^2 - 2*r*(K - H*r)])`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

method	result	size
default	$\frac{x}{r\sqrt{2Hr^2 - 2Kr - a^2 - e^2}}$	29
norman	$\frac{x}{r\sqrt{2Hr^2 - 2Kr - a^2 - e^2}}$	29
parallelrisc	$\frac{x}{r\sqrt{2Hr^2 - 2Kr - a^2 - e^2}}$	29
orering	$\frac{x}{r\sqrt{2Hr^2 - 2Kr - a^2 - e^2}}$	29

input `int(1/r/(2*H*r^2-2*K*r-a^2-e^2)^(1/2),x,method=_RETURNVERBOSE)`

output `x/r/(2*H*r^2-2*K*r-a^2-e^2)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.72

$$\int \frac{1}{r\sqrt{-a^2 - e^2 - 2Kr + 2Hr^2}} dx = \frac{\sqrt{2Hr^2 - a^2 - e^2 - 2Krx}}{2Hr^3 - 2Kr^2 - (a^2 + e^2)r}$$

input `integrate(1/r/(2*H*r^2-2*K*r-a^2-e^2)^(1/2),x, algorithm="fricas")`

output `sqrt(2*H*r^2 - a^2 - e^2 - 2*K*r)*x/(2*H*r^3 - 2*K*r^2 - (a^2 + e^2)*r)`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

$$\int \frac{1}{r\sqrt{-a^2 - e^2 - 2Kr + 2Hr^2}} dx = \frac{x}{r\sqrt{2Hr^2 - 2Kr - a^2 - e^2}}$$

input `integrate(1/r/(2*H*r**2-2*K*r-a**2-e**2)**(1/2),x)`

output `x/(r*sqrt(2*H*r**2 - 2*K*r - a**2 - e**2))`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.97

$$\int \frac{1}{r\sqrt{-a^2 - e^2 - 2Kr + 2Hr^2}} dx = \frac{x}{\sqrt{2Hr^2 - a^2 - e^2 - 2Krr}}$$

input `integrate(1/r/(2*H*r^2-2*K*r-a^2-e^2)^(1/2),x, algorithm="maxima")`output `x/(sqrt(2*H*r^2 - a^2 - e^2 - 2*K*r)*r)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.97

$$\int \frac{1}{r\sqrt{-a^2 - e^2 - 2Kr + 2Hr^2}} dx = \frac{x}{\sqrt{2Hr^2 - a^2 - e^2 - 2Krr}}$$

input `integrate(1/r/(2*H*r^2-2*K*r-a^2-e^2)^(1/2),x, algorithm="giac")`output `x/(sqrt(2*H*r^2 - a^2 - e^2 - 2*K*r)*r)`**Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.97

$$\int \frac{1}{r\sqrt{-a^2 - e^2 - 2Kr + 2Hr^2}} dx = \frac{x}{r\sqrt{-a^2 - e^2 + 2Hr^2 - 2Kr}}$$

input `int(1/(r*(2*H*r^2 - 2*K*r - a^2 - e^2)^(1/2)),x)`output `x/(r*(2*H*r^2 - 2*K*r - a^2 - e^2)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.62

$$\int \frac{1}{r\sqrt{-a^2 - e^2 - 2Kr + 2Hr^2}} dx = -\frac{\sqrt{2hr^2 - a^2 - e^2 - 2kr} x}{r(-2hr^2 + a^2 + e^2 + 2kr)}$$

input `int(1/r/(2*H*r^2-2*K*r-a^2-e^2)^(1/2),x)`

output `(- sqrt(- a**2 - e**2 + 2*h*r**2 - 2*k*r)*x)/(r*(a**2 + e**2 - 2*h*r**2 + 2*k*r))`

3.109 $\int \frac{r}{\sqrt{-a^2+2er^2}} dx$

Optimal result	640
Mathematica [A] (verified)	640
Rubi [A] (verified)	641
Maple [A] (verified)	641
Fricas [A] (verification not implemented)	642
Sympy [A] (verification not implemented)	642
Maxima [A] (verification not implemented)	643
Giac [A] (verification not implemented)	643
Mupad [B] (verification not implemented)	643
Reduce [B] (verification not implemented)	644

Optimal result

Integrand size = 18, antiderivative size = 19

$$\int \frac{r}{\sqrt{-a^2+2er^2}} dx = \frac{rx}{\sqrt{-a^2+2er^2}}$$

output `r*x/(-a^2+2*exp(1)*r^2)^(1/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{r}{\sqrt{-a^2+2er^2}} dx = \frac{rx}{\sqrt{-a^2+2er^2}}$$

input `Integrate[r/Sqrt[-a^2 + 2*E*r^2],x]`

output `(r*x)/Sqrt[-a^2 + 2*E*r^2]`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{r}{\sqrt{2er^2 - a^2}} dx$$

↓ 24

$$\frac{rx}{\sqrt{2er^2 - a^2}}$$

input `Int[r/Sqrt[-a^2 + 2*E*r^2],x]`

output `(r*x)/Sqrt[-a^2 + 2*E*r^2]`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

method	result	size
default	$\frac{rx}{\sqrt{-a^2+2er^2}}$	19
norman	$\frac{rx}{\sqrt{-a^2+2er^2}}$	19
parallelrisch	$\frac{rx}{\sqrt{-a^2+2er^2}}$	19
orering	$\frac{rx}{\sqrt{-a^2+2er^2}}$	19

input `int(r/(-a^2+2*exp(1)*r^2)^(1/2),x,method=_RETURNVERBOSE)`

output `r*x/(-a^2+2*exp(1)*r^2)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.95

$$\int \frac{r}{\sqrt{-a^2 + 2er^2}} dx = \frac{rx}{\sqrt{2r^2e - a^2}}$$

input `integrate(r/(-a^2+2*exp(1)*r^2)^(1/2),x, algorithm="fricas")`

output `r*x/sqrt(2*r^2*e - a^2)`

Sympy [A] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{r}{\sqrt{-a^2 + 2er^2}} dx = \frac{rx}{\sqrt{-a^2 + 2er^2}}$$

input `integrate(r/(-a**2+2*exp(1)*r**2)**(1/2),x)`

output `r*x/sqrt(-a**2 + 2*E*r**2)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.95

$$\int \frac{r}{\sqrt{-a^2 + 2er^2}} dx = \frac{rx}{\sqrt{2r^2e - a^2}}$$

input `integrate(r/(-a^2+2*exp(1)*r^2)^(1/2),x, algorithm="maxima")`output `r*x/sqrt(2*r^2*e - a^2)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.95

$$\int \frac{r}{\sqrt{-a^2 + 2er^2}} dx = \frac{rx}{\sqrt{2r^2e - a^2}}$$

input `integrate(r/(-a^2+2*exp(1)*r^2)^(1/2),x, algorithm="giac")`output `r*x/sqrt(2*r^2*e - a^2)`**Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.95

$$\int \frac{r}{\sqrt{-a^2 + 2er^2}} dx = \frac{rx}{\sqrt{2r^2e - a^2}}$$

input `int(r/(2*r^2*exp(1) - a^2)^(1/2),x)`output `(r*x)/(2*r^2*exp(1) - a^2)^(1/2)`

Reduce [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.53

$$\int \frac{r}{\sqrt{-a^2 + 2er^2}} dx = -\frac{\sqrt{2er^2 - a^2} rx}{-2er^2 + a^2}$$

input `int(r/(-a^2+2*exp(1)*r^2)^(1/2),x)`

output `(- sqrt(- a**2 + 2*e*r**2)*r*x)/(a**2 - 2*e*r**2)`

$$3.110 \quad \int \frac{r}{\sqrt{-a^2 - e^2 + 2er^2}} dx$$

Optimal result	645
Mathematica [A] (verified)	645
Rubi [A] (verified)	646
Maple [A] (verified)	646
Fricas [A] (verification not implemented)	647
Sympy [A] (verification not implemented)	647
Maxima [A] (verification not implemented)	648
Giac [A] (verification not implemented)	648
Mupad [B] (verification not implemented)	648
Reduce [B] (verification not implemented)	649

Optimal result

Integrand size = 23, antiderivative size = 24

$$\int \frac{r}{\sqrt{-a^2 - e^2 + 2er^2}} dx = \frac{rx}{\sqrt{-a^2 - e^2 + 2er^2}}$$

output `r*x/(-a^2-e^2+2*exp(1)*r^2)^(1/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{r}{\sqrt{-a^2 - e^2 + 2er^2}} dx = \frac{rx}{\sqrt{-a^2 - e^2 + 2er^2}}$$

input `Integrate[r/Sqrt[-a^2 - e^2 + 2*E*r^2],x]`

output `(r*x)/Sqrt[-a^2 - e^2 + 2*E*r^2]`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{r}{\sqrt{-a^2 - e^2 + 2er^2}} dx$$

\downarrow 24
 $\frac{rx}{\sqrt{-a^2 - e^2 + 2er^2}}$

input `Int[r/Sqrt[-a^2 - e^2 + 2*E*r^2],x]`

output `(r*x)/Sqrt[-a^2 - e^2 + 2*E*r^2]`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

method	result	size
default	$\frac{rx}{\sqrt{-a^2 - e^2 + 2er^2}}$	24
norman	$\frac{rx}{\sqrt{-a^2 - e^2 + 2er^2}}$	24
parallelrisc	$\frac{rx}{\sqrt{-a^2 - e^2 + 2er^2}}$	24
orering	$\frac{rx}{\sqrt{-a^2 - e^2 + 2er^2}}$	24

input `int(r/(-a^2-e^2+2*exp(1)*r^2)^(1/2),x,method=_RETURNVERBOSE)`

output `r*x/(-a^2-e^2+2*exp(1)*r^2)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.96

$$\int \frac{r}{\sqrt{-a^2 - e^2 + 2er^2}} dx = \frac{rx}{\sqrt{2r^2e - a^2 - e^2}}$$

input `integrate(r/(-a^2-e^2+2*exp(1)*r^2)^(1/2),x, algorithm="fricas")`

output `r*x/sqrt(2*r^2*e - a^2 - e^2)`

Sympy [A] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{r}{\sqrt{-a^2 - e^2 + 2er^2}} dx = \frac{rx}{\sqrt{-a^2 - e^2 + 2er^2}}$$

input `integrate(r/(-a**2-e**2+2*exp(1)*r**2)**(1/2),x)`

output `r*x/sqrt(-a**2 - e**2 + 2*E*r**2)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.96

$$\int \frac{r}{\sqrt{-a^2 - e^2 + 2er^2}} dx = \frac{rx}{\sqrt{2r^2e - a^2 - e^2}}$$

input `integrate(r/(-a^2-e^2+2*exp(1)*r^2)^(1/2),x, algorithm="maxima")`output `r*x/sqrt(2*r^2*e - a^2 - e^2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.96

$$\int \frac{r}{\sqrt{-a^2 - e^2 + 2er^2}} dx = \frac{rx}{\sqrt{2r^2e - a^2 - e^2}}$$

input `integrate(r/(-a^2-e^2+2*exp(1)*r^2)^(1/2),x, algorithm="giac")`output `r*x/sqrt(2*r^2*e - a^2 - e^2)`**Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.96

$$\int \frac{r}{\sqrt{-a^2 - e^2 + 2er^2}} dx = \frac{rx}{\sqrt{-a^2 - e^2 + 2er^2}}$$

input `int(r/(2*r^2*exp(1) - a^2 - e^2)^(1/2),x)`output `(r*x)/(2*r^2*exp(1) - a^2 - e^2)^(1/2)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.54

$$\int \frac{r}{\sqrt{-a^2 - e^2 + 2er^2}} dx = -\frac{\sqrt{2er^2 - a^2 - e^2}rx}{-2er^2 + a^2 + e^2}$$

input `int(r/(-a^2-e^2+2*exp(1)*r^2)^(1/2),x)`

output `(- sqrt(- a**2 - e**2 + 2*e*r**2)*r*x)/(a**2 + e**2 - 2*e*r**2)`

$$3.111 \quad \int \frac{r}{\sqrt{-a^2 + 2er^2 - 2Kr^4}} dx$$

Optimal result	650
Mathematica [A] (verified)	650
Rubi [A] (verified)	651
Maple [A] (verified)	651
Fricas [A] (verification not implemented)	652
Sympy [A] (verification not implemented)	652
Maxima [A] (verification not implemented)	653
Giac [A] (verification not implemented)	653
Mupad [B] (verification not implemented)	653
Reduce [B] (verification not implemented)	654

Optimal result

Integrand size = 24, antiderivative size = 25

$$\int \frac{r}{\sqrt{-a^2 + 2er^2 - 2Kr^4}} dx = \frac{rx}{\sqrt{-a^2 + 2er^2 - 2Kr^4}}$$

output `r*x/(-a^2+2*exp(1)*r^2-2*K*r^4)^(1/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{r}{\sqrt{-a^2 + 2er^2 - 2Kr^4}} dx = \frac{rx}{\sqrt{-a^2 + 2er^2 - 2Kr^4}}$$

input `Integrate[r/Sqrt[-a^2 + 2*E*r^2 - 2*K*r^4],x]`

output `(r*x)/Sqrt[-a^2 + 2*E*r^2 - 2*K*r^4]`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$, Rules used = {24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{r}{\sqrt{-a^2 - 2Kr^4 + 2er^2}} dx$$

↓ 24

$$\frac{rx}{\sqrt{-a^2 - 2Kr^4 + 2er^2}}$$

input `Int[r/Sqrt[-a^2 + 2*E*r^2 - 2*K*r^4],x]`

output `(r*x)/Sqrt[-a^2 + 2*E*r^2 - 2*K*r^4]`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

method	result	size
default	$\frac{rx}{\sqrt{-a^2+2er^2-2Kr^4}}$	25
norman	$\frac{rx}{\sqrt{-a^2+2er^2-2Kr^4}}$	25
parallelrisc	$\frac{rx}{\sqrt{-a^2+2er^2-2Kr^4}}$	25
orering	$\frac{rx}{\sqrt{-a^2+2er^2-2Kr^4}}$	25

input `int(r/(-a^2+2*exp(1)*r^2-2*K*r^4)^(1/2),x,method=_RETURNVERBOSE)`

output `r*x/(-a^2+2*exp(1)*r^2-2*K*r^4)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.76

$$\int \frac{r}{\sqrt{-a^2 + 2er^2 - 2Kr^4}} dx = -\frac{\sqrt{-2Kr^4 + 2r^2e - a^2}rx}{2Kr^4 - 2r^2e + a^2}$$

input `integrate(r/(-a^2+2*exp(1)*r^2-2*K*r^4)^(1/2),x, algorithm="fricas")`

output `-sqrt(-2*K*r^4 + 2*r^2*e - a^2)*r*x/(2*K*r^4 - 2*r^2*e + a^2)`

Sympy [A] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{r}{\sqrt{-a^2 + 2er^2 - 2Kr^4}} dx = \frac{rx}{\sqrt{-2Kr^4 - a^2 + 2er^2}}$$

input `integrate(r/(-a**2+2*exp(1)*r**2-2*K*r**4)**(1/2),x)`

output `r*x/sqrt(-2*K*r**4 - a**2 + 2*E*r**2)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{r}{\sqrt{-a^2 + 2er^2 - 2Kr^4}} dx = \frac{rx}{\sqrt{-2Kr^4 + 2r^2e - a^2}}$$

input `integrate(r/(-a^2+2*exp(1)*r^2-2*K*r^4)^(1/2),x, algorithm="maxima")`output `r*x/sqrt(-2*K*r^4 + 2*r^2*e - a^2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{r}{\sqrt{-a^2 + 2er^2 - 2Kr^4}} dx = \frac{rx}{\sqrt{-2Kr^4 + 2r^2e - a^2}}$$

input `integrate(r/(-a^2+2*exp(1)*r^2-2*K*r^4)^(1/2),x, algorithm="giac")`output `r*x/sqrt(-2*K*r^4 + 2*r^2*e - a^2)`**Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{r}{\sqrt{-a^2 + 2er^2 - 2Kr^4}} dx = \frac{rx}{\sqrt{-a^2 - 2Kr^4 + 2er^2}}$$

input `int(r/(2*r^2*exp(1) - 2*K*r^4 - a^2)^(1/2),x)`output `(r*x)/(2*r^2*exp(1) - 2*K*r^4 - a^2)^(1/2)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.64

$$\int \frac{r}{\sqrt{-a^2 + 2er^2 - 2Kr^4}} dx = -\frac{\sqrt{-2kr^4 + 2er^2 - a^2} rx}{2kr^4 - 2er^2 + a^2}$$

input `int(r/(-a^2+2*exp(1)*r^2-2*K*r^4)^(1/2),x)`

output `(- sqrt(- a**2 + 2*e*r**2 - 2*k*r**4)*r*x)/(a**2 - 2*e*r**2 + 2*k*r**4)`

3.112 $\int \frac{r}{\sqrt{-a^2 - e^2 + 2er^2 - 2Kr^4}} dx$

Optimal result	655
Mathematica [A] (verified)	655
Rubi [A] (verified)	656
Maple [A] (verified)	656
Fricas [A] (verification not implemented)	657
Sympy [A] (verification not implemented)	657
Maxima [A] (verification not implemented)	658
Giac [A] (verification not implemented)	658
Mupad [B] (verification not implemented)	658
Reduce [B] (verification not implemented)	659

Optimal result

Integrand size = 29, antiderivative size = 30

$$\int \frac{r}{\sqrt{-a^2 - e^2 + 2er^2 - 2Kr^4}} dx = \frac{rx}{\sqrt{-a^2 - e^2 + 2er^2 - 2Kr^4}}$$

output

```
r*x/(-a^2-e^2+2*exp(1)*r^2-2*K*r^4)^(1/2)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{r}{\sqrt{-a^2 - e^2 + 2er^2 - 2Kr^4}} dx = \frac{rx}{\sqrt{-a^2 - e^2 + 2er^2 - 2Kr^4}}$$

input

```
Integrate[r/Sqrt[-a^2 - e^2 + 2*E*r^2 - 2*K*r^4],x]
```

output

```
(r*x)/Sqrt[-a^2 - e^2 + 2*E*r^2 - 2*K*r^4]
```


Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.034$, Rules used = {24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{r}{\sqrt{-a^2 - e^2 - 2Kr^4 + 2er^2}} dx$$

\downarrow 24
 rx

$$\frac{rx}{\sqrt{-a^2 - e^2 - 2Kr^4 + 2er^2}}$$

input `Int[r/Sqrt[-a^2 - e^2 + 2*E*r^2 - 2*K*r^4], x]`

output `(r*x)/Sqrt[-a^2 - e^2 + 2*E*r^2 - 2*K*r^4]`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

method	result	size
default	$\frac{rx}{\sqrt{-a^2 - e^2 + 2er^2 - 2Kr^4}}$	30
norman	$\frac{rx}{\sqrt{-a^2 - e^2 + 2er^2 - 2Kr^4}}$	30
parallelrisc	$\frac{rx}{\sqrt{-a^2 - e^2 + 2er^2 - 2Kr^4}}$	30
orering	$\frac{rx}{\sqrt{-a^2 - e^2 + 2er^2 - 2Kr^4}}$	30

input `int(r/(-a^2-e^2+2*exp(1)*r^2-2*K*r^4)^(1/2),x,method=_RETURNVERBOSE)`

output `r*x/(-a^2-e^2+2*exp(1)*r^2-2*K*r^4)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.73

$$\int \frac{r}{\sqrt{-a^2 - e^2 + 2er^2 - 2Kr^4}} dx = -\frac{\sqrt{-2Kr^4 + 2r^2e - a^2 - e^2}rx}{2Kr^4 - 2r^2e + a^2 + e^2}$$

input `integrate(r/(-a^2-e^2+2*exp(1)*r^2-2*K*r^4)^(1/2),x, algorithm="fricas")`

output `-sqrt(-2*K*r^4 + 2*r^2*e - a^2 - e^2)*r*x/(2*K*r^4 - 2*r^2*e + a^2 + e^2)`

Sympy [A] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.90

$$\int \frac{r}{\sqrt{-a^2 - e^2 + 2er^2 - 2Kr^4}} dx = \frac{rx}{\sqrt{-2Kr^4 - a^2 - e^2 + 2er^2}}$$

input `integrate(r/(-a**2-e**2+2*exp(1)*r**2-2*K*r**4)**(1/2),x)`

output `r*x/sqrt(-2*K*r**4 - a**2 - e**2 + 2*E*r**2)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.97

$$\int \frac{r}{\sqrt{-a^2 - e^2 + 2er^2 - 2Kr^4}} dx = \frac{rx}{\sqrt{-2Kr^4 + 2r^2e - a^2 - e^2}}$$

input `integrate(r/(-a^2-e^2+2*exp(1)*r^2-2*K*r^4)^(1/2),x, algorithm="maxima")`output `r*x/sqrt(-2*K*r^4 + 2*r^2*e - a^2 - e^2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.97

$$\int \frac{r}{\sqrt{-a^2 - e^2 + 2er^2 - 2Kr^4}} dx = \frac{rx}{\sqrt{-2Kr^4 + 2r^2e - a^2 - e^2}}$$

input `integrate(r/(-a^2-e^2+2*exp(1)*r^2-2*K*r^4)^(1/2),x, algorithm="giac")`output `r*x/sqrt(-2*K*r^4 + 2*r^2*e - a^2 - e^2)`**Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.97

$$\int \frac{r}{\sqrt{-a^2 - e^2 + 2er^2 - 2Kr^4}} dx = \frac{rx}{\sqrt{-a^2 - e^2 - 2Kr^4 + 2er^2}}$$

input `int(r/(2*r^2*exp(1) - 2*K*r^4 - a^2 - e^2)^(1/2),x)`output `(r*x)/(2*r^2*exp(1) - 2*K*r^4 - a^2 - e^2)^(1/2)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.63

$$\int \frac{r}{\sqrt{-a^2 - e^2 + 2er^2 - 2Kr^4}} dx = -\frac{\sqrt{-2kr^4 + 2er^2 - a^2 - e^2}rx}{2kr^4 - 2er^2 + a^2 + e^2}$$

input `int(r/(-a^2-e^2+2*exp(1)*r^2-2*K*r^4)^(1/2),x)`

output `(- sqrt(- a**2 - e**2 + 2*e*r**2 - 2*k*r**4)*r*x)/(a**2 + e**2 - 2*e*r**2 + 2*k*r**4)`

$$3.113 \quad \int \frac{r}{\sqrt{-a^2 - e^2 - 2Kr + 2Hr^2}} dx$$

Optimal result	660
Mathematica [A] (verified)	660
Rubi [A] (verified)	661
Maple [A] (verified)	661
Fricas [A] (verification not implemented)	662
Sympy [A] (verification not implemented)	662
Maxima [A] (verification not implemented)	663
Giac [A] (verification not implemented)	663
Mupad [B] (verification not implemented)	663
Reduce [B] (verification not implemented)	664

Optimal result

Integrand size = 27, antiderivative size = 27

$$\int \frac{r}{\sqrt{-a^2 - e^2 - 2Kr + 2Hr^2}} dx = \frac{rx}{\sqrt{-a^2 - e^2 - 2r(K - Hr)}}$$

output `r*x/(2*H*r^2-2*K*r-a^2-e^2)^(1/2)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.04

$$\int \frac{r}{\sqrt{-a^2 - e^2 - 2Kr + 2Hr^2}} dx = \frac{rx}{\sqrt{-a^2 - e^2 - 2Kr + 2Hr^2}}$$

input `Integrate[r/Sqrt[-a^2 - e^2 - 2*K*r + 2*H*r^2],x]`

output `(r*x)/Sqrt[-a^2 - e^2 - 2*K*r + 2*H*r^2]`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.037$, Rules used = {24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{r}{\sqrt{-a^2 - e^2 + 2Hr^2 - 2Kr}} dx$$

↓ 24

$$\frac{rx}{\sqrt{-a^2 - e^2 - 2r(K - Hr)}}$$

input `Int[r/Sqrt[-a^2 - e^2 - 2*K*r + 2*H*r^2],x]`

output `(r*x)/Sqrt[-a^2 - e^2 - 2*r*(K - H*r)]`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

method	result	size
default	$\frac{rx}{\sqrt{2Hr^2 - 2Kr - a^2 - e^2}}$	27
norman	$\frac{rx}{\sqrt{2Hr^2 - 2Kr - a^2 - e^2}}$	27
parallelrisc	$\frac{rx}{\sqrt{2Hr^2 - 2Kr - a^2 - e^2}}$	27
orering	$\frac{rx}{\sqrt{2Hr^2 - 2Kr - a^2 - e^2}}$	27

input `int(r/(2*H*r^2-2*K*r-a^2-e^2)^(1/2),x,method=_RETURNVERBOSE)`

output `r*x/(2*H*r^2-2*K*r-a^2-e^2)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \frac{r}{\sqrt{-a^2 - e^2 - 2Kr + 2Hr^2}} dx = \frac{rx}{\sqrt{2Hr^2 - a^2 - e^2 - 2Kr}}$$

input `integrate(r/(2*H*r^2-2*K*r-a^2-e^2)^(1/2),x, algorithm="fricas")`

output `r*x/sqrt(2*H*r^2 - a^2 - e^2 - 2*K*r)`

Sympy [A] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89

$$\int \frac{r}{\sqrt{-a^2 - e^2 - 2Kr + 2Hr^2}} dx = \frac{rx}{\sqrt{2Hr^2 - 2Kr - a^2 - e^2}}$$

input `integrate(r/(2*H*r**2-2*K*r-a**2-e**2)**(1/2),x)`

output `r*x/sqrt(2*H*r**2 - 2*K*r - a**2 - e**2)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \frac{r}{\sqrt{-a^2 - e^2 - 2Kr + 2Hr^2}} dx = \frac{rx}{\sqrt{2Hr^2 - a^2 - e^2 - 2Kr}}$$

input `integrate(r/(2*H*r^2-2*K*r-a^2-e^2)^(1/2),x, algorithm="maxima")`output `r*x/sqrt(2*H*r^2 - a^2 - e^2 - 2*K*r)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \frac{r}{\sqrt{-a^2 - e^2 - 2Kr + 2Hr^2}} dx = \frac{rx}{\sqrt{2Hr^2 - a^2 - e^2 - 2Kr}}$$

input `integrate(r/(2*H*r^2-2*K*r-a^2-e^2)^(1/2),x, algorithm="giac")`output `r*x/sqrt(2*H*r^2 - a^2 - e^2 - 2*K*r)`**Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \frac{r}{\sqrt{-a^2 - e^2 - 2Kr + 2Hr^2}} dx = \frac{rx}{\sqrt{-a^2 - e^2 + 2Hr^2 - 2Kr}}$$

input `int(r/(2*H*r^2 - 2*K*r - a^2 - e^2)^(1/2),x)`output `(r*x)/(2*H*r^2 - 2*K*r - a^2 - e^2)^(1/2)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.67

$$\int \frac{r}{\sqrt{-a^2 - e^2 - 2Kr + 2Hr^2}} dx = -\frac{\sqrt{2hr^2 - a^2 - e^2 - 2kr} rx}{-2hr^2 + a^2 + e^2 + 2kr}$$

input `int(r/(2*H*r^2-2*K*r-a^2-e^2)^(1/2),x)`output `(- sqrt(- a**2 - e**2 + 2*h*r**2 - 2*k*r)*r*x)/(a**2 + e**2 - 2*h*r**2 + 2*k*r)`

CHAPTER 4

APPENDIX

4.1	Listing of Grading functions	665
4.2	Links to plain text integration problems used in this report for each CAS .	683

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
  ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "
  ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType, expn]],
  If[Head[expn]===Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]===Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
        1,
        Max[ExpnType[expn[[1]], 2]],
      Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]],
    If[Head[expn]===Plus || Head[expn]===Times,
      Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3, ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
    If[HypergeometricFunctionQ[Head[expn]],

```


Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue
fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
            fi;
        fi;
    else #ExpnType(result) > ExpnType(optimal)
        if debug then
            print("ExpnType(result) > ExpnType(optimal)");
        fi;
        return "C",cat("Result contains higher order function than in optimal. Order ",
                        convert(ExpnType_result,string)," vs. order ",
                        convert(ExpnType_optimal,string),".");
    fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```



```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```



```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file