

Computer Algebra Independent Integration Tests

Summer 2024

1-Algebraic-functions/1.1-Binomial/1.1.1-Linear-binomial/21-
1.1.1.3c

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 7:02am

Contents

1	Introduction	10
1.1	Listing of CAS systems tested	11
1.2	Results	12
1.3	Time and leaf size Performance	16
1.4	Performance based on number of rules Rubi used	18
1.5	Performance based on number of steps Rubi used	19
1.6	Solved integrals histogram based on leaf size of result	20
1.7	Solved integrals histogram based on CPU time used	21
1.8	Leaf size vs. CPU time used	22
1.9	list of integrals with no known antiderivative	23
1.10	List of integrals solved by CAS but has no known antiderivative	23
1.11	list of integrals solved by CAS but failed verification	23
1.12	Timing	24
1.13	Verification	24
1.14	Important notes about some of the results	25
1.15	Current tree layout of integration tests	28
1.16	Design of the test system	29
2	detailed summary tables of results	30
2.1	List of integrals sorted by grade for each CAS	31
2.2	Detailed conclusion table per each integral for all CAS systems	37
2.3	Detailed conclusion table specific for Rubi results	104
3	Listing of integrals	113
3.1	$\int (a + bx)(A + Bx)(d + ex)^4 dx$	122
3.2	$\int (a + bx)(A + Bx)(d + ex)^3 dx$	130
3.3	$\int (a + bx)(A + Bx)(d + ex)^2 dx$	137
3.4	$\int (a + bx)(A + Bx)(d + ex) dx$	143
3.5	$\int (a + bx)(A + Bx) dx$	149
3.6	$\int \frac{(a+bx)(A+Bx)}{d+ex} dx$	154
3.7	$\int \frac{(a+bx)(A+Bx)}{(d+ex)^2} dx$	160

3.8	$\int \frac{(a+bx)(A+Bx)}{(d+ex)^3} dx$	166
3.9	$\int \frac{(a+bx)(A+Bx)}{(d+ex)^4} dx$	172
3.10	$\int \frac{(a+bx)(A+Bx)}{(d+ex)^5} dx$	178
3.11	$\int \frac{(a+bx)(A+Bx)}{(d+ex)^6} dx$	184
3.12	$\int (a+bx)^2(A+Bx)(d+ex)^4 dx$	190
3.13	$\int (a+bx)^2(A+Bx)(d+ex)^3 dx$	200
3.14	$\int (a+bx)^2(A+Bx)(d+ex)^2 dx$	209
3.15	$\int (a+bx)^2(A+Bx)(d+ex) dx$	216
3.16	$\int (a+bx)^2(A+Bx) dx$	222
3.17	$\int \frac{(a+bx)^2(A+Bx)}{d+ex} dx$	227
3.18	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^2} dx$	233
3.19	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^3} dx$	240
3.20	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^4} dx$	246
3.21	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^5} dx$	252
3.22	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^6} dx$	258
3.23	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^7} dx$	264
3.24	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^8} dx$	270
3.25	$\int (a+bx)^3(A+Bx)(d+ex)^5 dx$	277
3.26	$\int (a+bx)^3(A+Bx)(d+ex)^4 dx$	290
3.27	$\int (a+bx)^3(A+Bx)(d+ex)^3 dx$	301
3.28	$\int (a+bx)^3(A+Bx)(d+ex)^2 dx$	311
3.29	$\int (a+bx)^3(A+Bx)(d+ex) dx$	320
3.30	$\int (a+bx)^3(A+Bx) dx$	327
3.31	$\int \frac{(a+bx)^3(A+Bx)}{d+ex} dx$	333
3.32	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^2} dx$	340
3.33	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^3} dx$	348
3.34	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^4} dx$	356
3.35	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^5} dx$	363
3.36	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^6} dx$	370
3.37	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^7} dx$	377
3.38	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^8} dx$	385
3.39	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^9} dx$	392
3.40	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{10}} dx$	399
3.41	$\int (a+bx)^6(A+Bx)(d+ex)^8 dx$	406
3.42	$\int (a+bx)^6(A+Bx)(d+ex)^7 dx$	417

3.43	$\int (a + bx)^6 (A + Bx)(d + ex)^6 dx$	428
3.44	$\int (a + bx)^6 (A + Bx)(d + ex)^5 dx$	440
3.45	$\int (a + bx)^6 (A + Bx)(d + ex)^4 dx$	451
3.46	$\int (a + bx)^6 (A + Bx)(d + ex)^3 dx$	463
3.47	$\int (a + bx)^6 (A + Bx)(d + ex)^2 dx$	476
3.48	$\int (a + bx)^6 (A + Bx)(d + ex) dx$	488
3.49	$\int (a + bx)^6 (A + Bx) dx$	497
3.50	$\int \frac{(a+bx)^6 (A+Bx)}{d+ex} dx$	503
3.51	$\int \frac{(a+bx)^6 (A+Bx)}{(d+ex)^2} dx$	515
3.52	$\int \frac{(a+bx)^6 (A+Bx)}{(d+ex)^3} dx$	526
3.53	$\int \frac{(a+bx)^6 (A+Bx)}{(d+ex)^4} dx$	536
3.54	$\int \frac{(a+bx)^6 (A+Bx)}{(d+ex)^5} dx$	547
3.55	$\int \frac{(a+bx)^6 (A+Bx)}{(d+ex)^6} dx$	557
3.56	$\int \frac{(a+bx)^6 (A+Bx)}{(d+ex)^7} dx$	567
3.57	$\int \frac{(a+bx)^6 (A+Bx)}{(d+ex)^8} dx$	577
3.58	$\int \frac{(a+bx)^6 (A+Bx)}{(d+ex)^9} dx$	587
3.59	$\int \frac{(a+bx)^6 (A+Bx)}{(d+ex)^{10}} dx$	596
3.60	$\int \frac{(a+bx)^6 (A+Bx)}{(d+ex)^{11}} dx$	605
3.61	$\int \frac{(a+bx)^6 (A+Bx)}{(d+ex)^{12}} dx$	615
3.62	$\int \frac{(a+bx)^6 (A+Bx)}{(d+ex)^{13}} dx$	626
3.63	$\int \frac{(a+bx)^6 (A+Bx)}{(d+ex)^{14}} dx$	636
3.64	$\int \frac{(a+bx)^6 (A+Bx)}{(d+ex)^{15}} dx$	646
3.65	$\int (a + bx)^{10} (A + Bx)(d + ex)^{13} dx$	656
3.66	$\int (a + bx)^{10} (A + Bx)(d + ex)^{12} dx$	667
3.67	$\int (a + bx)^{10} (A + Bx)(d + ex)^{11} dx$	678
3.68	$\int (a + bx)^{10} (A + Bx)(d + ex)^{10} dx$	689
3.69	$\int (a + bx)^{10} (A + Bx)(d + ex)^9 dx$	700
3.70	$\int (a + bx)^{10} (A + Bx)(d + ex)^8 dx$	712
3.71	$\int (a + bx)^{10} (A + Bx)(d + ex)^7 dx$	723
3.72	$\int (a + bx)^{10} (A + Bx)(d + ex)^6 dx$	734
3.73	$\int (a + bx)^{10} (A + Bx)(d + ex)^5 dx$	745
3.74	$\int (a + bx)^{10} (A + Bx)(d + ex)^4 dx$	756
3.75	$\int (a + bx)^{10} (A + Bx)(d + ex)^3 dx$	767
3.76	$\int (a + bx)^{10} (A + Bx)(d + ex)^2 dx$	778
3.77	$\int (a + bx)^{10} (A + Bx)(d + ex) dx$	789
3.78	$\int (a + bx)^{10} (A + Bx) dx$	800

3.79	$\int \frac{(a+bx)^{10}(A+Bx)}{d+ex} dx$	808
3.80	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^2} dx$	819
3.81	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^3} dx$	831
3.82	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^4} dx$	842
3.83	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^5} dx$	853
3.84	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^6} dx$	863
3.85	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^7} dx$	873
3.86	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^8} dx$	883
3.87	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^9} dx$	893
3.88	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{10}} dx$	903
3.89	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{11}} dx$	914
3.90	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{12}} dx$	925
3.91	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{13}} dx$	935
3.92	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{14}} dx$	944
3.93	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{15}} dx$	954
3.94	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{16}} dx$	965
3.95	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{17}} dx$	976
3.96	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{18}} dx$	988
3.97	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{19}} dx$	1002
3.98	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{20}} dx$	1019
3.99	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{21}} dx$	1030
3.100	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{22}} dx$	1041
3.101	$\int \frac{(A+Bx)(d+ex)^4}{a+bx} dx$	1052
3.102	$\int \frac{(A+Bx)(d+ex)^3}{a+bx} dx$	1060
3.103	$\int \frac{(A+Bx)(d+ex)^2}{a+bx} dx$	1067
3.104	$\int \frac{(A+Bx)(d+ex)}{a+bx} dx$	1073
3.105	$\int \frac{A+Bx}{a+bx} dx$	1079
3.106	$\int \frac{A+Bx}{(a+bx)(d+ex)} dx$	1084
3.107	$\int \frac{A+Bx}{(a+bx)(d+ex)^2} dx$	1089
3.108	$\int \frac{A+Bx}{(a+bx)(d+ex)^3} dx$	1095
3.109	$\int \frac{A+Bx}{(a+bx)(d+ex)^4} dx$	1102
3.110	$\int \frac{(A+Bx)(d+ex)^3}{(a+bx)^2} dx$	1109
3.111	$\int \frac{(A+Bx)(d+ex)^2}{(a+bx)^2} dx$	1117

3.112	$\int \frac{(A+Bx)(d+ex)}{(a+bx)^2} dx$	1124
3.113	$\int \frac{A+Bx}{(a+bx)^2} dx$	1130
3.114	$\int \frac{A+Bx}{(a+bx)^2(d+ex)} dx$	1135
3.115	$\int \frac{A+Bx}{(a+bx)^2(d+ex)^2} dx$	1141
3.116	$\int \frac{A+Bx}{(a+bx)^2(d+ex)^3} dx$	1148
3.117	$\int \frac{(A+Bx)(d+ex)^3}{(a+bx)^3} dx$	1156
3.118	$\int \frac{(A+Bx)(d+ex)^2}{(a+bx)^3} dx$	1163
3.119	$\int \frac{(A+Bx)(d+ex)}{(a+bx)^3} dx$	1169
3.120	$\int \frac{A+Bx}{(a+bx)^3} dx$	1175
3.121	$\int \frac{A+Bx}{(a+bx)^3(d+ex)} dx$	1180
3.122	$\int \frac{A+Bx}{(a+bx)^3(d+ex)^2} dx$	1187
3.123	$\int \frac{A+Bx}{(a+bx)^3(d+ex)^3} dx$	1195
3.124	$\int (a+bx)(A+Bx)(d+ex)^{5/2} dx$	1205
3.125	$\int (a+bx)(A+Bx)(d+ex)^{3/2} dx$	1212
3.126	$\int (a+bx)(A+Bx)\sqrt{d+ex} dx$	1219
3.127	$\int \frac{(a+bx)(A+Bx)}{\sqrt{d+ex}} dx$	1225
3.128	$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{3/2}} dx$	1231
3.129	$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{5/2}} dx$	1237
3.130	$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{7/2}} dx$	1243
3.131	$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{9/2}} dx$	1249
3.132	$\int (a+bx)^2(A+Bx)(d+ex)^{5/2} dx$	1256
3.133	$\int (a+bx)^2(A+Bx)(d+ex)^{3/2} dx$	1263
3.134	$\int (a+bx)^2(A+Bx)\sqrt{d+ex} dx$	1270
3.135	$\int \frac{(a+bx)^2(A+Bx)}{\sqrt{d+ex}} dx$	1277
3.136	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{3/2}} dx$	1283
3.137	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{5/2}} dx$	1289
3.138	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{7/2}} dx$	1296
3.139	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{9/2}} dx$	1303
3.140	$\int (a+bx)^3(A+Bx)(d+ex)^{5/2} dx$	1310
3.141	$\int (a+bx)^3(A+Bx)(d+ex)^{3/2} dx$	1319
3.142	$\int (a+bx)^3(A+Bx)\sqrt{d+ex} dx$	1328
3.143	$\int \frac{(a+bx)^3(A+Bx)}{\sqrt{d+ex}} dx$	1336
3.144	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{3/2}} dx$	1344
3.145	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{5/2}} dx$	1351

3.146	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{7/2}} dx$	1358
3.147	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{9/2}} dx$	1366
3.148	$\int \frac{(A+Bx)(d+ex)^{7/2}}{a+bx} dx$	1374
3.149	$\int \frac{(A+Bx)(d+ex)^{5/2}}{a+bx} dx$	1384
3.150	$\int \frac{(A+Bx)(d+ex)^{3/2}}{a+bx} dx$	1393
3.151	$\int \frac{(A+Bx)\sqrt{d+ex}}{a+bx} dx$	1401
3.152	$\int \frac{A+Bx}{(a+bx)\sqrt{d+ex}} dx$	1408
3.153	$\int \frac{A+Bx}{(a+bx)(d+ex)^{3/2}} dx$	1414
3.154	$\int \frac{A+Bx}{(a+bx)(d+ex)^{5/2}} dx$	1420
3.155	$\int \frac{A+Bx}{(a+bx)(d+ex)^{7/2}} dx$	1427
3.156	$\int \frac{(A+Bx)(d+ex)^{7/2}}{(a+bx)^2} dx$	1435
3.157	$\int \frac{(A+Bx)(d+ex)^{5/2}}{(a+bx)^2} dx$	1446
3.158	$\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^2} dx$	1455
3.159	$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^2} dx$	1463
3.160	$\int \frac{A+Bx}{(a+bx)^2\sqrt{d+ex}} dx$	1470
3.161	$\int \frac{A+Bx}{(a+bx)^2(d+ex)^{3/2}} dx$	1476
3.162	$\int \frac{A+Bx}{(a+bx)^2(d+ex)^{5/2}} dx$	1483
3.163	$\int \frac{A+Bx}{(a+bx)^2(d+ex)^{7/2}} dx$	1491
3.164	$\int \frac{(A+Bx)(d+ex)^{7/2}}{(a+bx)^3} dx$	1500
3.165	$\int \frac{(A+Bx)(d+ex)^{5/2}}{(a+bx)^3} dx$	1511
3.166	$\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^3} dx$	1521
3.167	$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^3} dx$	1529
3.168	$\int \frac{A+Bx}{(a+bx)^3\sqrt{d+ex}} dx$	1536
3.169	$\int \frac{A+Bx}{(a+bx)^3(d+ex)^{3/2}} dx$	1543
3.170	$\int \frac{A+Bx}{(a+bx)^3(d+ex)^{5/2}} dx$	1551
3.171	$\int \frac{A+Bx}{(a+bx)^3(d+ex)^{7/2}} dx$	1560
3.172	$\int \sqrt{a+bx}(A+Bx)(d+ex)^{5/2} dx$	1571
3.173	$\int \sqrt{a+bx}(A+Bx)(d+ex)^{3/2} dx$	1582
3.174	$\int \sqrt{a+bx}(A+Bx)\sqrt{d+ex} dx$	1591
3.175	$\int \frac{\sqrt{a+bx}(A+Bx)}{\sqrt{d+ex}} dx$	1600
3.176	$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{3/2}} dx$	1608
3.177	$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{5/2}} dx$	1615
3.178	$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{7/2}} dx$	1622

3.179	$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{9/2}} dx$	1628
3.180	$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{11/2}} dx$	1635
3.181	$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{13/2}} dx$	1643
3.182	$\int (a+bx)^{3/2}(A+Bx)(d+ex)^{5/2} dx$	1653
3.183	$\int (a+bx)^{3/2}(A+Bx)(d+ex)^{3/2} dx$	1666
3.184	$\int (a+bx)^{3/2}(A+Bx)\sqrt{d+ex} dx$	1677
3.185	$\int \frac{(a+bx)^{3/2}(A+Bx)}{\sqrt{d+ex}} dx$	1686
3.186	$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{3/2}} dx$	1694
3.187	$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{5/2}} dx$	1702
3.188	$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{7/2}} dx$	1710
3.189	$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{9/2}} dx$	1718
3.190	$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{11/2}} dx$	1724
3.191	$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{13/2}} dx$	1732
3.192	$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{15/2}} dx$	1741
3.193	$\int (a+bx)^{5/2}(A+Bx)(d+ex)^{5/2} dx$	1751
3.194	$\int (a+bx)^{5/2}(A+Bx)(d+ex)^{3/2} dx$	1767
3.195	$\int (a+bx)^{5/2}(A+Bx)\sqrt{d+ex} dx$	1780
3.196	$\int \frac{(a+bx)^{5/2}(A+Bx)}{\sqrt{d+ex}} dx$	1791
3.197	$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{3/2}} dx$	1800
3.198	$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{5/2}} dx$	1810
3.199	$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{7/2}} dx$	1820
3.200	$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{9/2}} dx$	1830
3.201	$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{11/2}} dx$	1839
3.202	$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{13/2}} dx$	1846
3.203	$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{15/2}} dx$	1854
3.204	$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{17/2}} dx$	1863
3.205	$\int \frac{(A+Bx)(d+ex)^{5/2}}{\sqrt{a+bx}} dx$	1872
3.206	$\int \frac{(A+Bx)(d+ex)^{3/2}}{\sqrt{a+bx}} dx$	1882
3.207	$\int \frac{(A+Bx)\sqrt{d+ex}}{\sqrt{a+bx}} dx$	1890
3.208	$\int \frac{A+Bx}{\sqrt{a+bx}\sqrt{d+ex}} dx$	1898
3.209	$\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{3/2}} dx$	1904
3.210	$\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{5/2}} dx$	1910

3.211	$\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{7/2}} dx$	1916
3.212	$\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{9/2}} dx$	1923
3.213	$\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{11/2}} dx$	1931
3.214	$\int \frac{(A+Bx)(d+ex)^{5/2}}{(a+bx)^{3/2}} dx$	1941
3.215	$\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^{3/2}} dx$	1951
3.216	$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^{3/2}} dx$	1959
3.217	$\int \frac{A+Bx}{(a+bx)^{3/2}\sqrt{d+ex}} dx$	1966
3.218	$\int \frac{A+Bx}{(a+bx)^{3/2}(d+ex)^{3/2}} dx$	1972
3.219	$\int \frac{A+Bx}{(a+bx)^{3/2}(d+ex)^{5/2}} dx$	1978
3.220	$\int \frac{A+Bx}{(a+bx)^{3/2}(d+ex)^{7/2}} dx$	1985
3.221	$\int \frac{A+Bx}{(a+bx)^{3/2}(d+ex)^{9/2}} dx$	1993
3.222	$\int \frac{(A+Bx)(d+ex)^{7/2}}{(a+bx)^{5/2}} dx$	2002
3.223	$\int \frac{(A+Bx)(d+ex)^{5/2}}{(a+bx)^{5/2}} dx$	2013
3.224	$\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^{5/2}} dx$	2023
3.225	$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^{5/2}} dx$	2032
3.226	$\int \frac{A+Bx}{(a+bx)^{5/2}\sqrt{d+ex}} dx$	2040
3.227	$\int \frac{A+Bx}{(a+bx)^{5/2}(d+ex)^{3/2}} dx$	2046
3.228	$\int \frac{A+Bx}{(a+bx)^{5/2}(d+ex)^{5/2}} dx$	2053
3.229	$\int \frac{A+Bx}{(a+bx)^{5/2}(d+ex)^{7/2}} dx$	2061
3.230	$\int (a+bx)^3(A+Bx)(d+ex)^m dx$	2070
3.231	$\int (a+bx)^2(A+Bx)(d+ex)^m dx$	2080
3.232	$\int (a+bx)(A+Bx)(d+ex)^m dx$	2090
3.233	$\int (A+Bx)(d+ex)^m dx$	2098
3.234	$\int \frac{(A+Bx)(d+ex)^m}{a+bx} dx$	2104
3.235	$\int \frac{(A+Bx)(d+ex)^m}{(a+bx)^2} dx$	2109
3.236	$\int \frac{(A+Bx)(d+ex)^m}{(a+bx)^3} dx$	2114
3.237	$\int (1-2x)(2+3x)^m(3+5x)^3 dx$	2120
3.238	$\int (1-2x)(2+3x)^m(3+5x)^2 dx$	2128
3.239	$\int (1-2x)(2+3x)^m(3+5x) dx$	2135
3.240	$\int \frac{(1-2x)(2+3x)^m}{3+5x} dx$	2141
3.241	$\int \frac{(1-2x)(2+3x)^m}{(3+5x)^2} dx$	2146
3.242	$\int \frac{(1-2x)(2+3x)^m}{(3+5x)^3} dx$	2151
3.243	$\int \frac{(c+dx)^{1+m}(e+fx)}{(a+bx)^2} dx$	2157
3.244	$\int \frac{(c+dx)^m(e+fx)}{(a+bx)^2} dx$	2163

3.245	$\int \frac{(c+dx)^{-1+m}(e+fx)}{(a+bx)^2} dx$	2169
3.246	$\int \frac{(c+dx)^{1+m}(e+fx)}{(a+bx)^3} dx$	2175
3.247	$\int \frac{(c+dx)^m(e+fx)}{(a+bx)^3} dx$	2181
3.248	$\int \frac{(c+dx)^{-1+m}(e+fx)}{(a+bx)^3} dx$	2187
3.249	$\int (a+bx)^m(c+dx)^n(e+fx) dx$	2193
3.250	$\int (a+bx)^p(c+dx)^{-2-p}(e+fx) dx$	2199
3.251	$\int (a+bx)^m(ac-bcx)^n(ad(m-n)-bd(2+m+n)x) dx$	2205
3.252	$\int (a+bx)(c+dx)^n(e+fx)^{-n} dx$	2211
3.253	$\int (a+bx)(c+dx)^{-1+n}(e+fx)^{-n} dx$	2216
3.254	$\int (a+bx)(c+dx)^{-2+n}(e+fx)^{-n} dx$	2221
3.255	$\int (a+bx)(c+dx)^{-3+n}(e+fx)^{-n} dx$	2227
3.256	$\int (a+bx)(c+dx)^{-4+n}(e+fx)^{-n} dx$	2234
3.257	$\int (a+bx)(c+dx)^{-5+n}(e+fx)^{-n} dx$	2242
3.258	$\int (a+bx)^{-n}(c+dx)(e+fx)^n dx$	2251
3.259	$\int (a+bx)^{-n}(c+dx)(e+fx)^{-1+n} dx$	2256
3.260	$\int (a+bx)^{-n}(c+dx)(e+fx)^{-2+n} dx$	2261
3.261	$\int (a+bx)^{-n}(c+dx)(e+fx)^{-3+n} dx$	2267
3.262	$\int (a+bx)^{-n}(c+dx)(e+fx)^{-4+n} dx$	2274
3.263	$\int (a+bx)^{-n}(c+dx)(e+fx)^{-5+n} dx$	2282
3.264	$\int (a+bx)^{1-n}(c+dx)^{-2+n}(-ad+bc(3+2n)+2bd(1+n)x) dx$	2291
4	Appendix	2298
4.1	Listing of Grading functions	2298
4.2	Links to plain text integration problems used in this report for each CAS	316

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	11
1.2	Results	12
1.3	Time and leaf size Performance	16
1.4	Performance based on number of rules Rubi used	18
1.5	Performance based on number of steps Rubi used	19
1.6	Solved integrals histogram based on leaf size of result	20
1.7	Solved integrals histogram based on CPU time used	21
1.8	Leaf size vs. CPU time used	22
1.9	list of integrals with no known antiderivative	23
1.10	List of integrals solved by CAS but has no known antiderivative	23
1.11	list of integrals solved by CAS but failed verification	23
1.12	Timing	24
1.13	Verification	24
1.14	Important notes about some of the results	25
1.15	Current tree layout of integration tests	28
1.16	Design of the test system	29

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [264]. This is test number [21].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (264)	0.00 (0)
Mathematica	100.00 (264)	0.00 (0)
Maple	92.05 (243)	7.95 (21)
Fricas	91.67 (242)	8.33 (22)
Giac	90.53 (239)	9.47 (25)
Reduce	90.15 (238)	9.85 (26)
Mupad	80.68 (213)	19.32 (51)
Maxima	58.71 (155)	41.29 (109)
Sympy	49.24 (130)	50.76 (134)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Mathematica	78.409	21.212	0.379	0.000
Maple	45.076	45.833	1.136	7.955
Fricas	26.894	64.773	0.000	8.333
Giac	25.379	65.152	0.000	9.470
Maxima	23.485	35.227	0.000	41.288
Sympy	16.288	32.955	0.000	50.758
Mupad	0.000	80.682	0.000	19.318
Reduce	0.000	90.152	0.000	9.848

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Maple	21	100.00	0.00	0.00
Fricas	22	95.45	4.55	0.00
Giac	25	100.00	0.00	0.00
Reduce	26	100.00	0.00	0.00
Mupad	51	0.00	100.00	0.00
Maxima	109	24.77	0.00	75.23
Sympy	134	50.00	38.81	11.19

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.06
Giac	0.19
Mathematica	0.29
Maple	0.30
Rubi	0.49
Reduce	0.61
Mupad	1.39
Fricas	2.90
Sympy	3.05

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Rubi	172.20	1.00	145.00	1.00
Mathematica	373.27	1.79	165.00	1.08
Reduce	483.40	2.22	266.00	1.91
Maple	621.76	2.82	269.00	1.84
Maxima	711.34	3.21	282.00	2.21
Mupad	752.08	3.28	303.00	2.14
Fricas	786.71	3.74	521.50	3.36
Giac	852.09	3.97	410.00	2.90
Sympy	874.32	5.05	353.50	2.62

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand,the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	31
2.2	Detailed conclusion table per each integral for all CAS systems	37
2.3	Detailed conclusion table specific for Rubi results	104

2.1 List of integrals sorted by grade for each CAS

Rubi	31
Mma	32
Maple	32
Fricas	33
Maxima	33
Giac	34
Mupad	35
Sympy	35
Reduce	36

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 27, 28, 29, 30, 31, 32, 33, 34, 35, 37, 38, 39, 40, 52, 53, 54, 82, 83, 84, 85, 86, 87, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264 }

B grade { 1, 12, 25, 26, 36, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100 }

C grade { 198 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 32, 33, 34, 38, 39, 40, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 178, 179, 180, 181, 189, 190, 191, 201, 202, 203, 210, 211, 212, 218, 219, 220, 226, 227, 228, 233, 251, 255, 261 }

B grade { 1, 12, 13, 25, 26, 27, 28, 29, 30, 31, 35, 36, 37, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 172, 173, 174, 175, 176, 177, 182, 183, 184, 185, 186, 187, 188, 192, 193, 194, 195, 196, 197, 198, 199, 200, 204, 205, 206, 207, 208, 209, 213, 214, 215, 216, 217, 221, 222, 223, 224, 225, 229, 230, 231, 232, 256, 257, 262, 263 }

C grade { 237, 238, 239 }

F normal fail { 234, 235, 236, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 252, 253, 254, 258, 259, 260, 264 }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 14, 15, 16, 17, 22, 23, 103, 104, 105, 106, 107, 112, 113, 114, 119, 120, 126, 127, 128, 129, 130, 131, 134, 135, 136, 137, 138, 139, 143, 144, 145, 146, 147, 150, 151, 152, 157, 158, 165, 172, 173, 174, 175, 176, 184, 185, 187, 195, 196, 205, 206, 207, 208, 210, 216, 218, 226, 233, 238, 239, 251 }

B grade { 1, 12, 13, 18, 19, 20, 21, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 108, 109, 110, 111, 115, 116, 117, 118, 121, 122, 123, 124, 125, 132, 133, 140, 141, 142, 148, 149, 153, 154, 155, 156, 159, 160, 161, 162, 163, 164, 166, 167, 168, 169, 170, 171, 177, 178, 179, 180, 181, 182, 183, 186, 188, 189, 190, 191, 192, 193, 194, 197, 198, 199, 200, 201, 202, 203, 209, 211, 212, 213, 214, 215, 217, 219, 220, 221, 222, 223, 224, 225, 227, 228, 229, 230, 231, 232, 237, 255, 256, 257, 261, 262, 263 }

C grade { }

F normal fail { 234, 235, 236, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 252, 253, 254, 258, 259, 260, 264 }

F(-1) timedout fail { 204 }

F(-2) exception fail { }

Maxima

A grade { 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 14, 15, 16, 17, 18, 19, 20, 22, 23, 32, 33, 34, 103, 104, 105, 106, 107, 110, 111, 112, 113, 114, 118, 119, 120, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 232, 233, 251 }

B grade { 1, 12, 13, 21, 24, 25, 26, 27, 28, 29, 30, 31, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95,

96, 97, 98, 99, 100, 101, 102, 108, 109, 115, 116, 117, 121, 122, 123, 230, 231, 237, 238, 239
 }

C grade { }

F normal fail { 234, 235, 236, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 252, 253,
 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264 }

F(-1) timedout fail { }

F(-2) exception fail { 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161,
 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180,
 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199,
 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218,
 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229 }

Giac

A grade { 3, 4, 5, 6, 7, 8, 9, 10, 11, 14, 15, 16, 17, 19, 20, 22, 23, 24, 34, 38, 39, 40, 103, 104,
 105, 106, 107, 112, 113, 114, 115, 116, 118, 119, 120, 122, 127, 128, 129, 130, 131, 135, 136,
 137, 138, 139, 151, 152, 153, 154, 158, 159, 160, 161, 162, 166, 167, 168, 175, 176, 185, 186,
 196, 208, 209, 217, 251 }

B grade { 1, 2, 12, 13, 18, 21, 25, 26, 27, 28, 29, 30, 31, 32, 33, 35, 36, 37, 41, 42, 43, 44, 45,
 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70,
 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95,
 96, 97, 98, 99, 100, 101, 102, 108, 109, 110, 111, 117, 121, 123, 124, 125, 126, 132, 133, 134,
 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 155, 156, 157, 163, 164, 165, 169, 170,
 171, 172, 173, 174, 177, 178, 179, 180, 181, 182, 183, 184, 187, 188, 189, 190, 191, 192, 193,
 194, 195, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 210, 211, 212, 213, 214, 215,
 216, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 237, 238,
 239, 255, 261 }

C grade { }

F normal fail { 234, 235, 236, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 252, 253,
 254, 256, 257, 258, 259, 260, 262, 263, 264 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 174, 175, 178, 179, 180, 181, 189, 190, 191, 192, 201, 202, 203, 204, 207, 208, 210, 211, 212, 213, 218, 219, 220, 221, 226, 227, 228, 229, 230, 231, 232, 233, 237, 238, 239, 251, 255, 256, 257, 261, 262, 263 }

C grade { }

F normal fail { }

F(-1) timedout fail { 172, 173, 176, 177, 182, 183, 184, 185, 186, 187, 188, 193, 194, 195, 196, 197, 198, 199, 200, 205, 206, 209, 214, 215, 216, 217, 222, 223, 224, 225, 234, 235, 236, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 252, 253, 254, 258, 259, 260, 264 }

F(-2) exception fail { }

Sympy

A grade { 3, 4, 5, 6, 7, 8, 9, 10, 11, 14, 15, 16, 17, 18, 19, 32, 103, 104, 105, 110, 111, 112, 113, 118, 119, 120, 125, 126, 127, 128, 133, 134, 135, 136, 145, 148, 149, 150, 151, 152, 153, 154, 155 }

B grade { 1, 2, 12, 13, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 33, 34, 35, 36, 37, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 101, 102, 106, 107, 108, 109, 114, 115, 116, 117, 121, 122, 123, 124, 129, 130, 131, 132, 137, 138, 139, 140, 141, 142, 143, 144, 146, 147, 230, 231, 232, 233, 237, 238, 239, 251 }

C grade { }

F normal fail { 159, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 193, 194, 195, 196, 197, 198, 199, 200, 201, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 234, 235, 236, 240, 241, 242, 243, 244, 245, 247, 248, 250 }

F(-1) timedout fail { 38, 39, 40, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 156, 157, 158, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 192, 202, 203, 204 }

F(-2) exception fail { 246, 249, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264 }

Reduce

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 237, 238, 239, 251 }

C grade { }

F normal fail { 235, 236, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	172	176	175	175	226	216	172	182
N.S.	1	1.00	2.23	2.29	2.27	2.27	2.94	2.81	2.23	2.36
time (sec)	N/A	0.310	0.038	0.149	0.042	0.081	0.031	0.120	0.155	0.058

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	130	135	134	134	168	163	131	141
N.S.	1	1.00	1.69	1.75	1.74	1.74	2.18	2.12	1.70	1.83
time (sec)	N/A	0.295	0.028	0.144	0.040	0.091	0.026	0.120	0.156	0.893

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	96	94	93	93	116	113	90	98
N.S.	1	1.00	1.25	1.22	1.21	1.21	1.51	1.47	1.17	1.27
time (sec)	N/A	0.268	0.020	0.141	0.036	0.068	0.022	0.115	0.155	0.871

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	53	53	52	62	63	62	49	54
N.S.	1	1.00	0.95	0.95	0.93	1.11	1.12	1.11	0.88	0.96
time (sec)	N/A	0.228	0.013	0.045	0.038	0.035	0.017	0.126	0.154	0.027

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	28	25	24	26	26	26	21	25
N.S.	1	1.00	1.00	0.89	0.86	0.93	0.93	0.93	0.75	0.89
time (sec)	N/A	0.187	0.003	0.029	0.037	0.036	0.017	0.127	0.158	0.022

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	56	66	66	68	53	68	72	68
N.S.	1	1.00	0.93	1.10	1.10	1.13	0.88	1.13	1.20	1.13
time (sec)	N/A	0.224	0.016	0.191	0.033	0.063	0.135	0.124	0.168	0.889

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	56	70	74	102	71	114	112	75
N.S.	1	1.00	0.89	1.11	1.17	1.62	1.13	1.81	1.78	1.19
time (sec)	N/A	0.238	0.029	0.198	0.033	0.109	0.231	0.128	0.163	0.059

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	72	72	87	105	94	77	113	82
N.S.	1	1.00	1.03	1.03	1.24	1.50	1.34	1.10	1.61	1.17
time (sec)	N/A	0.265	0.022	0.191	0.051	0.072	0.393	0.121	0.161	0.908

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	63	70	93	93	107	70	69	91
N.S.	1	1.00	0.84	0.93	1.24	1.24	1.43	0.93	0.92	1.21
time (sec)	N/A	0.262	0.019	0.194	0.037	0.072	0.705	0.127	0.155	0.030

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	62	68	102	102	117	121	95	101
N.S.	1	1.00	0.81	0.88	1.32	1.32	1.52	1.57	1.23	1.31
time (sec)	N/A	0.240	0.019	0.203	0.029	0.073	1.140	0.114	0.158	0.911

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	65	73	117	117	134	72	106	118
N.S.	1	1.00	0.84	0.95	1.52	1.52	1.74	0.94	1.38	1.53
time (sec)	N/A	0.258	0.021	0.208	0.041	0.071	1.721	0.121	0.157	0.044

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	120	120	283	305	304	304	384	374	247	305
N.S.	1	1.00	2.36	2.54	2.53	2.53	3.20	3.12	2.06	2.54
time (sec)	N/A	0.438	0.059	0.180	0.032	0.062	0.036	0.120	0.157	0.949

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	120	120	216	237	236	236	296	287	189	231
N.S.	1	1.00	1.80	1.98	1.97	1.97	2.47	2.39	1.58	1.92
time (sec)	N/A	0.364	0.047	0.175	0.030	0.083	0.032	0.121	0.157	0.052

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	118	118	157	169	168	168	202	199	131	157
N.S.	1	1.00	1.33	1.43	1.42	1.42	1.71	1.69	1.11	1.33
time (sec)	N/A	0.388	0.034	0.170	0.041	0.064	0.027	0.124	0.160	0.039

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	96	99	100	100	116	113	73	98
N.S.	1	1.00	1.28	1.32	1.33	1.33	1.55	1.51	0.97	1.31
time (sec)	N/A	0.283	0.018	0.146	0.034	0.070	0.024	0.117	0.160	0.903

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	46	48	48	48	49	49	32	47
N.S.	1	1.00	1.21	1.26	1.26	1.26	1.29	1.29	0.84	1.24
time (sec)	N/A	0.211	0.007	0.131	0.027	0.068	0.023	0.117	0.155	0.026

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	102	148	152	153	117	166	132	159
N.S.	1	1.00	1.11	1.61	1.65	1.66	1.27	1.80	1.43	1.73
time (sec)	N/A	0.281	0.036	0.217	0.029	0.073	0.247	0.119	0.159	0.925

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	101	98	151	156	238	151	230	197	165
N.S.	1	1.00	0.97	1.50	1.54	2.36	1.50	2.28	1.95	1.63
time (sec)	N/A	0.319	0.053	0.214	0.041	0.080	0.467	0.119	0.160	0.054

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	106	143	155	166	247	187	159	209	170
N.S.	1	1.00	1.35	1.46	1.57	2.33	1.76	1.50	1.97	1.60
time (sec)	N/A	0.305	0.049	0.212	0.038	0.077	1.105	0.119	0.160	0.080

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	104	138	157	184	221	211	166	180	178
N.S.	1	1.03	1.37	1.55	1.82	2.19	2.09	1.64	1.78	1.76
time (sec)	N/A	0.297	0.043	0.208	0.038	0.069	2.364	0.126	0.159	0.081

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	86	125	152	187	187	223	232	122	184
N.S.	1	1.00	1.45	1.77	2.17	2.17	2.59	2.70	1.42	2.14
time (sec)	N/A	0.220	0.039	0.217	0.036	0.080	4.644	0.121	0.161	0.984

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	120	120	129	157	203	203	238	168	159	201
N.S.	1	1.00	1.08	1.31	1.69	1.69	1.98	1.40	1.32	1.68
time (sec)	N/A	0.314	0.038	0.214	0.041	0.067	8.336	0.121	0.158	0.055

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	120	120	126	151	208	208	246	166	170	206
N.S.	1	1.00	1.05	1.26	1.73	1.73	2.05	1.38	1.42	1.72
time (sec)	N/A	0.297	0.038	0.219	0.047	0.073	15.406	0.121	0.159	0.982

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	120	120	129	157	225	225	262	168	181	223
N.S.	1	1.00	1.08	1.31	1.88	1.88	2.18	1.40	1.51	1.86
time (sec)	N/A	0.304	0.039	0.224	0.042	0.068	27.267	0.118	0.159	0.061

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	163	163	471	529	518	518	678	658	397	544
N.S.	1	1.00	2.89	3.25	3.18	3.18	4.16	4.04	2.44	3.34
time (sec)	N/A	0.627	0.103	0.182	0.040	0.065	0.071	0.123	0.156	0.137

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	163	163	397	434	425	425	546	534	322	439
N.S.	1	1.00	2.44	2.66	2.61	2.61	3.35	3.28	1.98	2.69
time (sec)	N/A	0.551	0.087	0.181	0.036	0.079	0.050	0.121	0.150	0.960

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	159	159	297	339	325	325	422	410	247	334
N.S.	1	1.00	1.87	2.13	2.04	2.04	2.65	2.58	1.55	2.10
time (sec)	N/A	0.492	0.067	0.173	0.033	0.069	0.050	0.121	0.158	0.066

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	118	118	224	244	239	239	296	287	172	231
N.S.	1	1.00	1.90	2.07	2.03	2.03	2.51	2.43	1.46	1.96
time (sec)	N/A	0.391	0.049	0.174	0.042	0.077	0.052	0.124	0.156	0.053

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	130	142	146	146	168	163	97	141
N.S.	1	1.00	1.73	1.89	1.95	1.95	2.24	2.17	1.29	1.88
time (sec)	N/A	0.313	0.027	0.166	0.037	0.084	0.029	0.118	0.154	0.929

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	67	70	69	69	73	72	43	65
N.S.	1	1.00	1.76	1.84	1.82	1.82	1.92	1.89	1.13	1.71
time (sec)	N/A	0.205	0.007	0.125	0.027	0.062	0.027	0.121	0.155	0.020

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	124	124	169	260	258	260	221	297	208	268
N.S.	1	1.00	1.36	2.10	2.08	2.10	1.78	2.40	1.68	2.16
time (sec)	N/A	0.340	0.056	0.217	0.032	0.098	0.493	0.123	0.157	0.953

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	150	150	244	276	267	396	257	372	297	293
N.S.	1	1.00	1.63	1.84	1.78	2.64	1.71	2.48	1.98	1.95
time (sec)	N/A	0.422	0.066	0.214	0.039	0.073	0.747	0.119	0.157	0.057

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	145	145	238	267	274	420	299	286	330	290
N.S.	1	1.00	1.64	1.84	1.89	2.90	2.06	1.97	2.28	2.00
time (sec)	N/A	0.418	0.072	0.218	0.040	0.073	2.432	0.120	0.163	0.966

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	149	149	232	271	284	406	337	279	309	301
N.S.	1	1.00	1.56	1.82	1.91	2.72	2.26	1.87	2.07	2.02
time (sec)	N/A	0.398	0.070	0.223	0.045	0.084	5.580	0.117	0.173	0.968

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	129	131	222	267	302	354	359	449	269	303
N.S.	1	1.02	1.72	2.07	2.34	2.74	2.78	3.48	2.09	2.35
time (sec)	N/A	0.358	0.068	0.225	0.048	0.079	13.089	0.128	0.165	0.979

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	86	211	264	304	304	372	298	192	307
N.S.	1	1.00	2.45	3.07	3.53	3.53	4.33	3.47	2.23	3.57
time (sec)	N/A	0.228	0.063	0.217	0.044	0.070	35.073	0.124	0.155	0.073

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	133	130	211	266	317	317	386	299	240	321
N.S.	1	0.98	1.59	2.00	2.38	2.38	2.90	2.25	1.80	2.41
time (sec)	N/A	0.266	0.057	0.222	0.044	0.069	100.278	0.127	0.170	0.927

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	163	163	215	270	332	332	0	300	251	336
N.S.	1	1.00	1.32	1.66	2.04	2.04	0.00	1.84	1.54	2.06
time (sec)	N/A	0.459	0.060	0.222	0.056	0.070	0.000	0.122	0.167	0.904

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	163	163	211	262	335	335	0	298	262	339
N.S.	1	1.00	1.29	1.61	2.06	2.06	0.00	1.83	1.61	2.08
time (sec)	N/A	0.386	0.058	0.233	0.046	0.075	0.000	0.121	0.159	0.086

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	163	163	214	270	354	354	0	300	273	358
N.S.	1	1.00	1.31	1.66	2.17	2.17	0.00	1.84	1.67	2.20
time (sec)	N/A	0.382	0.062	0.231	0.054	0.086	0.000	0.116	0.155	0.951

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	292	292	1385	1525	1532	1532	1969	1943	1051	1625
N.S.	1	1.00	4.74	5.22	5.25	5.25	6.74	6.65	3.60	5.57
time (sec)	N/A	1.713	0.327	0.234	0.052	0.078	0.119	0.128	0.154	1.197

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	292	292	1224	1349	1356	1356	1756	1712	925	1431
N.S.	1	1.00	4.19	4.62	4.64	4.64	6.01	5.86	3.17	4.90
time (sec)	N/A	1.489	0.287	0.229	0.040	0.071	0.112	0.128	0.169	0.295

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	290	290	1069	1173	1172	1172	1504	1480	799	1221
N.S.	1	1.00	3.69	4.04	4.04	4.04	5.19	5.10	2.76	4.21
time (sec)	N/A	1.308	0.231	0.237	0.041	0.079	0.091	0.125	0.162	1.147

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	240	240	907	997	997	997	1278	1246	673	1039
N.S.	1	1.00	3.78	4.15	4.15	4.15	5.32	5.19	2.80	4.33
time (sec)	N/A	1.061	0.197	0.208	0.044	0.085	0.083	0.122	0.163	0.215

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	204	204	762	821	828	828	1035	1015	547	845
N.S.	1	1.00	3.74	4.02	4.06	4.06	5.07	4.98	2.68	4.14
time (sec)	N/A	0.887	0.159	0.201	0.039	0.087	0.070	0.128	0.166	1.012

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	159	159	586	645	643	643	802	782	421	649
N.S.	1	1.00	3.69	4.06	4.04	4.04	5.04	4.92	2.65	4.08
time (sec)	N/A	0.683	0.126	0.194	0.045	0.078	0.061	0.124	0.167	1.004

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	118	118	386	469	476	476	568	552	295	457
N.S.	1	1.00	3.27	3.97	4.03	4.03	4.81	4.68	2.50	3.87
time (sec)	N/A	0.553	0.140	0.182	0.035	0.068	0.055	0.120	0.170	0.115

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	231	279	297	297	333	321	169	257
N.S.	1	1.00	3.08	3.72	3.96	3.96	4.44	4.28	2.25	3.43
time (sec)	N/A	0.393	0.089	0.154	0.032	0.071	0.040	0.124	0.159	0.087

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	122	140	142	142	148	145	76	126
N.S.	1	1.00	3.21	3.68	3.74	3.74	3.89	3.82	2.00	3.32
time (sec)	N/A	0.204	0.025	0.142	0.041	0.075	0.034	0.135	0.159	0.963

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	220	220	501	776	762	763	736	910	538	769
N.S.	1	1.00	2.28	3.53	3.46	3.47	3.35	4.14	2.45	3.50
time (sec)	N/A	0.510	0.163	0.232	0.057	0.075	0.866	0.122	0.155	0.944

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	277	277	643	795	771	1067	782	979	700	1228
N.S.	1	1.00	2.32	2.87	2.78	3.85	2.82	3.53	2.53	4.43
time (sec)	N/A	0.932	0.176	0.224	0.057	0.090	1.843	0.129	0.164	0.087

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	276	276	352	789	779	1177	821	862	799	1053
N.S.	1	1.00	1.28	2.86	2.82	4.26	2.97	3.12	2.89	3.82
time (sec)	N/A	0.825	0.111	0.234	0.051	0.085	6.152	0.122	0.165	0.973

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	279	279	297	795	793	1225	867	848	840	907
N.S.	1	1.00	1.06	2.85	2.84	4.39	3.11	3.04	3.01	3.25
time (sec)	N/A	0.815	0.111	0.240	0.067	0.097	29.906	0.125	0.164	0.970

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	279	279	263	798	801	1222	0	1175	851	863
N.S.	1	1.00	0.94	2.86	2.87	4.38	0.00	4.21	3.05	3.09
time (sec)	N/A	0.732	0.117	0.237	0.088	0.095	0.000	0.137	0.159	0.137

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	272	272	633	800	814	1157	0	831	816	862
N.S.	1	1.00	2.33	2.94	2.99	4.25	0.00	3.06	3.00	3.17
time (sec)	N/A	0.670	0.195	0.230	0.074	0.126	0.000	0.135	0.158	0.142

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	278	278	619	805	824	1063	0	826	735	875
N.S.	1	1.00	2.23	2.90	2.96	3.82	0.00	2.97	2.64	3.15
time (sec)	N/A	0.646	0.204	0.227	0.077	0.124	0.000	0.117	0.159	1.057

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	213	212	615	791	842	939	0	830	638	1046
N.S.	1	1.00	2.89	3.71	3.95	4.41	0.00	3.90	3.00	4.91
time (sec)	N/A	0.571	0.283	0.227	0.065	0.109	0.000	0.119	0.160	1.065

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	86	597	780	823	823	0	910	504	854
N.S.	1	1.00	6.94	9.07	9.57	9.57	0.00	10.58	5.86	9.93
time (sec)	N/A	0.255	0.164	0.240	0.077	0.089	0.000	0.127	0.158	0.143

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	135	131	603	789	861	861	0	912	585	877
N.S.	1	0.97	4.47	5.84	6.38	6.38	0.00	6.76	4.33	6.50
time (sec)	N/A	0.269	0.171	0.232	0.072	0.090	0.000	0.125	0.159	1.038

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	185	175	602	789	872	872	0	912	596	888
N.S.	1	0.95	3.25	4.26	4.71	4.71	0.00	4.93	3.22	4.80
time (sec)	N/A	0.290	0.165	0.242	0.079	0.114	0.000	0.135	0.162	1.068

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	235	219	605	789	883	883	0	912	607	899
N.S.	1	0.93	2.57	3.36	3.76	3.76	0.00	3.88	2.58	3.83
time (sec)	N/A	0.317	0.164	0.239	0.130	0.088	0.000	0.131	0.157	0.183

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	292	292	600	789	894	894	0	912	618	910
N.S.	1	1.00	2.05	2.70	3.06	3.06	0.00	3.12	2.12	3.12
time (sec)	N/A	0.702	0.165	0.246	0.076	0.099	0.000	0.124	0.158	1.211

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	292	292	605	789	905	905	0	912	629	921
N.S.	1	1.00	2.07	2.70	3.10	3.10	0.00	3.12	2.15	3.15
time (sec)	N/A	0.658	0.170	0.248	0.069	0.129	0.000	0.129	0.167	1.406

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	292	292	602	775	902	902	0	910	640	918
N.S.	1	1.00	2.06	2.65	3.09	3.09	0.00	3.12	2.19	3.14
time (sec)	N/A	0.644	0.173	0.260	0.089	0.125	0.000	0.123	0.166	1.566

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	464	464	3532	3893	3905	3905	5092	5039	2593	4206
N.S.	1	1.00	7.61	8.39	8.42	8.42	10.97	10.86	5.59	9.06
time (sec)	N/A	5.233	0.962	0.327	0.071	0.128	0.362	0.132	0.168	1.007

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	464	464	3320	3609	3621	3621	4655	4663	2399	3891
N.S.	1	1.00	7.16	7.78	7.80	7.80	10.03	10.05	5.17	8.39
time (sec)	N/A	5.348	0.883	0.326	0.067	0.132	0.272	0.133	0.158	1.821

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	461	461	3018	3325	3334	3334	4328	4288	2205	3577
N.S.	1	1.00	6.55	7.21	7.23	7.23	9.39	9.30	4.78	7.76
time (sec)	N/A	4.892	0.759	0.317	0.052	0.106	0.248	0.130	0.164	1.661

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	460	460	2815	3041	3048	3048	3936	3912	2011	3262
N.S.	1	1.00	6.12	6.61	6.63	6.63	8.56	8.50	4.37	7.09
time (sec)	N/A	4.132	0.711	0.287	0.056	0.132	0.252	0.134	0.166	1.638

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	415	415	2553	2757	2771	2771	3541	3536	1817	2947
N.S.	1	1.00	6.15	6.64	6.68	6.68	8.53	8.52	4.38	7.10
time (sec)	N/A	3.405	0.619	0.274	0.047	0.098	0.253	0.121	0.168	1.577

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	372	372	2307	2473	2487	2487	3165	3159	1623	2631
N.S.	1	1.00	6.20	6.65	6.69	6.69	8.51	8.49	4.36	7.07
time (sec)	N/A	2.842	0.534	0.264	0.045	0.115	0.224	0.123	0.159	1.490

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	329	329	2034	2189	2198	2198	2824	2783	1429	2316
N.S.	1	1.00	6.18	6.65	6.68	6.68	8.58	8.46	4.34	7.04
time (sec)	N/A	2.093	0.472	0.250	0.046	0.087	0.248	0.131	0.160	1.419

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	290	290	1788	1905	1917	1917	2424	2407	1235	2001
N.S.	1	1.00	6.17	6.57	6.61	6.61	8.36	8.30	4.26	6.90
time (sec)	N/A	1.910	0.423	0.241	0.056	0.095	0.161	0.132	0.156	1.405

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	243	243	1509	1621	1625	1625	2076	2032	1041	1685
N.S.	1	1.00	6.21	6.67	6.69	6.69	8.54	8.36	4.28	6.93
time (sec)	N/A	1.639	0.336	0.227	0.047	0.103	0.137	0.117	0.154	0.390

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	204	204	1098	1337	1352	1352	1676	1656	847	1386
N.S.	1	1.00	5.38	6.55	6.63	6.63	8.22	8.12	4.15	6.79
time (sec)	N/A	1.179	0.450	0.217	0.043	0.076	0.180	0.125	0.161	1.230

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	159	159	855	1053	1068	1068	1302	1279	653	1070
N.S.	1	1.00	5.38	6.62	6.72	6.72	8.19	8.04	4.11	6.73
time (sec)	N/A	0.927	0.332	0.210	0.042	0.095	0.132	0.122	0.155	1.169

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	118	118	614	769	781	781	921	904	459	757
N.S.	1	1.00	5.20	6.52	6.62	6.62	7.81	7.66	3.89	6.42
time (sec)	N/A	0.706	0.225	0.195	0.049	0.090	0.093	0.128	0.157	1.164

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	383	459	493	493	549	529	265	409
N.S.	1	1.00	5.11	6.12	6.57	6.57	7.32	7.05	3.53	5.45
time (sec)	N/A	0.497	0.108	0.167	0.038	0.073	0.073	0.121	0.165	0.153

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	198	232	240	240	248	241	120	208
N.S.	1	1.00	5.21	6.11	6.32	6.32	6.53	6.34	3.16	5.47
time (sec)	N/A	0.183	0.038	0.135	0.043	0.088	0.049	0.125	0.170	0.983

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	348	348	1252	1884	1804	1805	1912	2230	1216	1795
N.S.	1	1.00	3.60	5.41	5.18	5.19	5.49	6.41	3.49	5.16
time (sec)	N/A	0.642	0.768	0.235	0.051	0.083	2.193	0.123	0.162	1.020

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	445	445	1486	1907	1817	2329	1974	2207	1474	7792
N.S.	1	1.00	3.34	4.29	4.08	5.23	4.44	4.96	3.31	17.51
time (sec)	N/A	1.714	0.420	0.239	0.050	0.123	5.009	0.148	1.444	1.124

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	445	445	1480	1902	1826	2547	2004	2134	1661	8104
N.S.	1	1.00	3.33	4.27	4.10	5.72	4.50	4.80	3.73	18.21
time (sec)	N/A	1.614	0.417	0.239	0.082	0.151	22.447	0.140	0.838	1.208

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	445	445	814	1907	1839	2702	0	2096	1786	5544
N.S.	1	1.00	1.83	4.29	4.13	6.07	0.00	4.71	4.01	12.46
time (sec)	N/A	1.535	0.265	0.239	0.109	0.130	0.000	0.131	0.332	1.180

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	444	444	686	1910	1846	2808	0	2571	1889	3655
N.S.	1	1.00	1.55	4.30	4.16	6.32	0.00	5.79	4.25	8.23
time (sec)	N/A	1.442	0.257	0.237	0.116	0.139	0.000	0.141	0.222	1.164

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	447	447	587	1916	1861	2851	0	2032	1946	2681
N.S.	1	1.00	1.31	4.29	4.16	6.38	0.00	4.55	4.35	6.00
time (sec)	N/A	1.405	0.264	0.237	0.138	0.155	0.000	0.135	0.174	1.209

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	447	447	505	1921	1869	2850	0	2009	1957	2252
N.S.	1	1.00	1.13	4.30	4.18	6.38	0.00	4.49	4.38	5.04
time (sec)	N/A	1.351	0.293	0.238	0.163	0.173	0.000	0.131	0.163	1.179

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	444	444	450	1925	1883	2783	0	1992	1922	2092
N.S.	1	1.00	1.01	4.34	4.24	6.27	0.00	4.49	4.33	4.71
time (sec)	N/A	1.322	0.312	0.243	0.181	0.224	0.000	0.130	0.170	1.221

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	445	445	415	1929	1892	2677	0	1981	1841	2048
N.S.	1	1.00	0.93	4.33	4.25	6.02	0.00	4.45	4.14	4.60
time (sec)	N/A	1.246	0.327	0.243	0.194	0.251	0.000	0.130	0.165	0.307

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	441	441	1460	1928	1904	2501	0	1975	1714	2048
N.S.	1	1.00	3.31	4.37	4.32	5.67	0.00	4.48	3.89	4.64
time (sec)	N/A	1.207	0.556	0.243	0.169	0.230	0.000	0.125	0.166	1.249

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	446	446	1447	1922	1914	2309	0	1970	1541	2874
N.S.	1	1.00	3.24	4.31	4.29	5.18	0.00	4.42	3.46	6.44
time (sec)	N/A	1.207	0.540	0.257	0.132	0.241	0.000	0.130	0.167	1.363

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	321	316	1443	1903	1932	2089	0	1974	1368	2446
N.S.	1	0.98	4.50	5.93	6.02	6.51	0.00	6.15	4.26	7.62
time (sec)	N/A	0.695	1.223	0.259	0.116	0.174	0.000	0.130	0.169	1.295

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	86	1421	1888	1875	1875	0	2230	1158	2008
N.S.	1	1.00	16.52	21.95	21.80	21.80	0.00	25.93	13.47	23.35
time (sec)	N/A	0.211	0.521	0.258	0.130	0.138	0.000	0.134	0.166	0.470

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	135	131	1433	1901	1951	1951	0	2232	1283	2031
N.S.	1	0.97	10.61	14.08	14.45	14.45	0.00	16.53	9.50	15.04
time (sec)	N/A	0.222	0.482	0.266	0.131	0.134	0.000	0.129	0.168	1.540

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	185	175	1430	1901	1962	1962	0	2232	1294	2044
N.S.	1	0.95	7.73	10.28	10.61	10.61	0.00	12.06	6.99	11.05
time (sec)	N/A	0.250	0.546	0.279	0.161	0.197	0.000	0.134	0.171	1.778

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	235	219	1430	1901	1973	1973	0	2232	1305	2055
N.S.	1	0.93	6.09	8.09	8.40	8.40	0.00	9.50	5.55	8.74
time (sec)	N/A	0.296	0.558	0.296	0.127	0.164	0.000	0.126	0.172	2.239

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	285	263	1429	1901	1984	1984	0	2232	1316	2066
N.S.	1	0.92	5.01	6.67	6.96	6.96	0.00	7.83	4.62	7.25
time (sec)	N/A	0.301	0.462	0.313	0.141	0.177	0.000	0.129	0.169	1.318

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	335	307	1433	1901	1995	1995	0	2232	1327	2077
N.S.	1	0.92	4.28	5.67	5.96	5.96	0.00	6.66	3.96	6.20
time (sec)	N/A	0.333	0.524	0.324	0.157	0.175	0.000	0.131	0.167	1.400

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	385	351	1428	1901	2006	2006	0	2232	1338	2088
N.S.	1	0.91	3.71	4.94	5.21	5.21	0.00	5.80	3.48	5.42
time (sec)	N/A	0.368	0.480	0.340	0.133	0.204	0.000	0.129	0.166	9.644

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	460	460	1433	1901	2017	2017	0	2232	1349	2099
N.S.	1	1.00	3.12	4.13	4.38	4.38	0.00	4.85	2.93	4.56
time (sec)	N/A	1.274	0.501	0.354	0.146	0.220	0.000	0.133	0.165	17.868

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	462	462	1428	1901	2028	2028	0	2232	1360	2110
N.S.	1	1.00	3.09	4.11	4.39	4.39	0.00	4.83	2.94	4.57
time (sec)	N/A	1.212	0.525	0.372	0.151	0.247	0.000	0.135	0.166	1.457

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	464	464	1431	1901	2039	2039	0	2232	1371	2121
N.S.	1	1.00	3.08	4.10	4.39	4.39	0.00	4.81	2.95	4.57
time (sec)	N/A	1.207	0.540	0.379	0.157	0.256	0.000	0.132	0.170	1.628

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	155	155	257	404	400	403	352	466	43	411
N.S.	1	1.00	1.66	2.61	2.58	2.60	2.27	3.01	0.28	2.65
time (sec)	N/A	0.308	0.082	0.218	0.038	0.089	0.547	0.120	0.167	0.059

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	123	169	261	266	269	221	298	32	268
N.S.	1	1.00	1.37	2.12	2.16	2.19	1.80	2.42	0.26	2.18
time (sec)	N/A	0.270	0.053	0.214	0.049	0.075	0.390	0.122	0.159	1.000

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	91	102	150	155	158	117	166	21	159
N.S.	1	1.00	1.12	1.65	1.70	1.74	1.29	1.82	0.23	1.75
time (sec)	N/A	0.233	0.034	0.210	0.031	0.066	0.261	0.123	0.165	0.971

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	59	56	66	72	75	53	69	10	68
N.S.	1	1.00	0.95	1.12	1.22	1.27	0.90	1.17	0.17	1.15
time (sec)	N/A	0.204	0.015	0.188	0.030	0.075	0.173	0.118	0.163	0.050

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	25	26	26	25	20	27	1	25
N.S.	1	1.00	1.00	1.04	1.04	1.00	0.80	1.08	0.04	1.00
time (sec)	N/A	0.169	0.005	0.195	0.026	0.064	0.076	0.117	0.167	0.969

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	57	50	58	58	53	226	60	10	57
N.S.	1	1.00	0.88	1.02	1.02	0.93	3.96	1.05	0.18	1.00
time (sec)	N/A	0.211	0.019	0.269	0.026	0.083	0.890	0.124	0.171	0.137

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	80	83	119	148	355	101	12	95
N.S.	1	1.00	0.98	1.01	1.45	1.80	4.33	1.23	0.15	1.16
time (sec)	N/A	0.246	0.051	0.290	0.041	0.077	0.726	0.122	0.158	1.027

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	112	112	112	111	247	343	558	233	23	228
N.S.	1	1.00	1.00	0.99	2.21	3.06	4.98	2.08	0.21	2.04
time (sec)	N/A	0.288	0.059	0.306	0.043	0.076	1.090	0.124	0.157	1.055

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	146	146	145	144	444	608	818	376	34	399
N.S.	1	1.00	0.99	0.99	3.04	4.16	5.60	2.58	0.23	2.73
time (sec)	N/A	0.330	0.115	0.333	0.045	0.082	1.595	0.125	0.169	1.170

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	145	145	250	277	273	417	257	370	132	293
N.S.	1	1.00	1.72	1.91	1.88	2.88	1.77	2.55	0.91	2.02
time (sec)	N/A	0.357	0.070	0.215	0.053	0.074	0.751	0.122	0.153	0.061

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	99	99	153	151	158	249	151	229	72	165
N.S.	1	1.00	1.55	1.53	1.60	2.52	1.53	2.31	0.73	1.67
time (sec)	N/A	0.275	0.048	0.207	0.050	0.078	0.476	0.126	0.155	1.014

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	56	70	77	109	71	112	28	75
N.S.	1	1.00	0.93	1.17	1.28	1.82	1.18	1.87	0.47	1.25
time (sec)	N/A	0.224	0.030	0.187	0.031	0.069	0.240	0.130	0.161	1.005

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	31	33	34	37	27	57	10	32
N.S.	1	1.00	0.97	1.03	1.06	1.16	0.84	1.78	0.31	1.00
time (sec)	N/A	0.182	0.007	0.175	0.034	0.080	0.091	0.112	0.160	0.024

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	69	83	118	157	355	100	26	94
N.S.	1	1.00	0.84	1.01	1.44	1.91	4.33	1.22	0.32	1.15
time (sec)	N/A	0.242	0.037	0.288	0.030	0.075	0.696	0.123	0.162	0.089

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	117	103	123	256	396	706	194	115	263
N.S.	1	1.00	0.88	1.05	2.19	3.38	6.03	1.66	0.98	2.25
time (sec)	N/A	0.299	0.064	0.303	0.047	0.088	1.329	0.126	0.161	1.103

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	157	157	146	159	479	801	1066	306	289	454
N.S.	1	1.00	0.93	1.01	3.05	5.10	6.79	1.95	1.84	2.89
time (sec)	N/A	0.349	0.058	0.326	0.046	0.121	1.981	0.125	0.162	1.198

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	141	141	245	268	282	442	299	283	197	290
N.S.	1	1.00	1.74	1.90	2.00	3.13	2.12	2.01	1.40	2.06
time (sec)	N/A	0.335	0.076	0.214	0.037	0.098	2.045	0.118	0.164	1.120

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	103	103	140	157	170	258	187	157	112	170
N.S.	1	1.00	1.36	1.52	1.65	2.50	1.82	1.52	1.09	1.65
time (sec)	N/A	0.269	0.044	0.208	0.054	0.087	1.153	0.122	0.167	0.095

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	75	72	92	110	94	72	49	82
N.S.	1	1.00	1.09	1.04	1.33	1.59	1.36	1.04	0.71	1.19
time (sec)	N/A	0.228	0.019	0.188	0.032	0.076	0.435	0.120	0.176	1.131

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	26	25	38	38	39	24	12	39
N.S.	1	1.00	0.93	0.89	1.36	1.36	1.39	0.86	0.43	1.39
time (sec)	N/A	0.164	0.006	0.174	0.028	0.059	0.141	0.117	0.160	0.980

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	113	103	111	252	361	558	232	115	228
N.S.	1	1.00	0.91	0.98	2.23	3.19	4.94	2.05	1.02	2.02
time (sec)	N/A	0.333	0.038	0.306	0.041	0.105	1.454	0.122	0.161	1.082

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	158	158	146	161	477	803	1066	305	459	453
N.S.	1	1.00	0.92	1.02	3.02	5.08	6.75	1.93	2.91	2.87
time (sec)	N/A	0.405	0.055	0.323	0.047	0.100	1.995	0.125	0.161	1.224

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	199	199	185	200	745	1215	1431	532	821	726
N.S.	1	1.00	0.93	1.01	3.74	6.11	7.19	2.67	4.13	3.65
time (sec)	N/A	0.497	0.078	0.351	0.051	0.143	2.877	0.128	0.166	1.355

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	83	70	60	75	189	476	735	180	80
N.S.	1	1.00	0.84	0.72	0.90	2.28	5.73	8.86	2.17	0.96
time (sec)	N/A	0.256	0.060	0.668	0.031	0.106	0.481	0.125	0.162	0.051

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	83	70	60	75	149	124	473	139	80
N.S.	1	1.00	0.84	0.72	0.90	1.80	1.49	5.70	1.67	0.96
time (sec)	N/A	0.234	0.056	0.647	0.034	0.083	0.927	0.131	0.160	0.042

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	83	70	60	75	108	124	261	98	80
N.S.	1	1.00	0.84	0.72	0.90	1.30	1.49	3.14	1.18	0.96
time (sec)	N/A	0.236	0.051	0.625	0.041	0.071	0.875	0.126	0.168	0.970

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	68	60	75	70	122	105	61	80
N.S.	1	1.00	0.84	0.74	0.93	0.86	1.51	1.30	0.75	0.99
time (sec)	N/A	0.227	0.049	0.625	0.037	0.066	0.837	0.125	0.162	0.968

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	68	64	82	79	107	100	62	79
N.S.	1	1.00	0.86	0.81	1.04	1.00	1.35	1.27	0.78	1.00
time (sec)	N/A	0.237	0.063	0.244	0.035	0.102	2.262	0.126	0.160	0.039

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	68	63	79	91	355	82	70	72
N.S.	1	1.00	0.86	0.80	1.00	1.15	4.49	1.04	0.89	0.91
time (sec)	N/A	0.240	0.055	0.250	0.044	0.074	0.369	0.125	0.165	0.986

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	68	61	72	101	520	80	81	72
N.S.	1	1.00	0.84	0.75	0.89	1.25	6.42	0.99	1.00	0.89
time (sec)	N/A	0.235	0.053	0.240	0.028	0.075	0.512	0.123	0.168	0.054

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	83	70	64	72	113	683	80	92	72
N.S.	1	1.00	0.84	0.77	0.87	1.36	8.23	0.96	1.11	0.87
time (sec)	N/A	0.233	0.064	0.237	0.045	0.076	0.722	0.121	0.158	0.046

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	128	128	138	120	159	356	857	1293	284	115
N.S.	1	1.00	1.08	0.94	1.24	2.78	6.70	10.10	2.22	0.90
time (sec)	N/A	0.312	0.109	0.721	0.028	0.086	0.501	0.131	0.154	1.014

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	128	128	139	117	159	289	257	854	226	115
N.S.	1	1.00	1.09	0.91	1.24	2.26	2.01	6.67	1.77	0.90
time (sec)	N/A	0.274	0.099	0.672	0.041	0.098	1.109	0.133	0.154	0.984

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	128	128	138	116	159	220	257	486	168	115
N.S.	1	1.00	1.08	0.91	1.24	1.72	2.01	3.80	1.31	0.90
time (sec)	N/A	0.316	0.095	0.672	0.031	0.111	1.283	0.123	0.163	0.043

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	126	137	117	159	155	255	206	114	115
N.S.	1	1.00	1.09	0.93	1.26	1.23	2.02	1.63	0.90	0.91
time (sec)	N/A	0.287	0.087	0.675	0.035	0.073	0.972	0.126	0.161	0.044

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	124	124	135	121	167	165	207	221	115	154
N.S.	1	1.00	1.09	0.98	1.35	1.33	1.67	1.78	0.93	1.24
time (sec)	N/A	0.298	0.101	0.289	0.034	0.086	4.218	0.129	0.161	0.045

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	124	124	136	105	163	175	709	203	122	189
N.S.	1	1.00	1.10	0.85	1.31	1.41	5.72	1.64	0.98	1.52
time (sec)	N/A	0.278	0.116	0.296	0.033	0.090	0.407	0.128	0.162	0.970

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	124	124	136	118	164	188	1015	193	134	168
N.S.	1	1.00	1.10	0.95	1.32	1.52	8.19	1.56	1.08	1.35
time (sec)	N/A	0.283	0.109	0.298	0.036	0.107	0.573	0.124	0.164	0.973

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	126	138	118	156	198	1323	189	145	150
N.S.	1	1.00	1.10	0.94	1.24	1.57	10.50	1.50	1.15	1.19
time (sec)	N/A	0.271	0.116	0.279	0.033	0.099	0.756	0.138	0.152	0.059

Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	173	173	228	171	265	539	423	1947	405	154
N.S.	1	1.00	1.32	0.99	1.53	3.12	2.45	11.25	2.34	0.89
time (sec)	N/A	0.363	0.163	0.754	0.033	0.078	1.277	0.142	0.156	0.046

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	173	173	228	171	265	446	423	1306	330	154
N.S.	1	1.00	1.32	0.99	1.53	2.58	2.45	7.55	1.91	0.89
time (sec)	N/A	0.330	0.163	0.708	0.042	0.121	1.173	0.144	0.156	0.989

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	173	173	227	171	265	353	423	761	255	154
N.S.	1	1.00	1.31	0.99	1.53	2.04	2.45	4.40	1.47	0.89
time (sec)	N/A	0.318	0.144	0.703	0.037	0.093	1.119	0.129	0.155	0.031

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	171	171	226	170	265	263	422	329	184	154
N.S.	1	1.00	1.32	0.99	1.55	1.54	2.47	1.92	1.08	0.90
time (sec)	N/A	0.323	0.135	0.751	0.034	0.076	1.060	0.124	0.204	0.996

Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	167	167	222	189	273	272	340	385	186	223
N.S.	1	1.00	1.33	1.13	1.63	1.63	2.04	2.31	1.11	1.34
time (sec)	N/A	0.308	0.138	0.319	0.049	0.076	9.553	0.128	0.153	0.035

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	169	169	223	179	271	284	284	364	193	264
N.S.	1	1.00	1.32	1.06	1.60	1.68	1.68	2.15	1.14	1.56
time (sec)	N/A	0.314	0.177	0.332	0.034	0.096	9.611	0.134	0.155	0.966

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	169	169	226	193	273	294	1654	358	204	300
N.S.	1	1.00	1.34	1.14	1.62	1.74	9.79	2.12	1.21	1.78
time (sec)	N/A	0.301	0.185	0.339	0.050	0.077	0.575	0.132	0.156	1.010

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	167	167	224	193	271	306	2144	346	215	300
N.S.	1	1.00	1.34	1.16	1.62	1.83	12.84	2.07	1.29	1.80
time (sec)	N/A	0.318	0.162	0.337	0.036	0.103	0.791	0.129	0.155	1.047

Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	198	182	263	269	0	865	376	560	49	426
N.S.	1	0.92	1.33	1.36	0.00	4.37	1.90	2.83	0.25	2.15
time (sec)	N/A	0.340	0.344	0.740	0.000	0.096	4.236	0.132	0.159	1.028

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	164	153	185	192	0	591	258	371	38	330
N.S.	1	0.93	1.13	1.17	0.00	3.60	1.57	2.26	0.23	2.01
time (sec)	N/A	0.301	0.236	0.450	0.000	0.090	3.960	0.133	0.153	1.002

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	130	124	129	138	0	373	175	226	27	236
N.S.	1	0.95	0.99	1.06	0.00	2.87	1.35	1.74	0.21	1.82
time (sec)	N/A	0.282	0.190	0.409	0.000	0.118	3.774	0.129	0.154	0.987

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	95	94	101	0	211	124	122	16	107
N.S.	1	0.97	0.96	1.03	0.00	2.15	1.27	1.24	0.16	1.09
time (sec)	N/A	0.248	0.187	0.375	0.000	0.089	4.579	0.126	0.157	0.085

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	74	63	0	209	92	66	11	62
N.S.	1	1.00	1.00	0.85	0.00	2.82	1.24	0.89	0.15	0.84
time (sec)	N/A	0.223	0.119	0.356	0.000	0.084	1.599	0.123	0.162	0.954

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	88	79	0	363	109	87	13	96
N.S.	1	1.00	1.00	0.90	0.00	4.12	1.24	0.99	0.15	1.09
time (sec)	N/A	0.243	0.200	0.357	0.000	0.137	3.715	0.118	0.166	0.072

Problem 154	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	119	123	118	116	0	486	138	153	20	128
N.S.	1	1.03	0.99	0.97	0.00	4.08	1.16	1.29	0.17	1.08
time (sec)	N/A	0.313	0.221	0.472	0.000	0.097	4.672	0.128	0.175	1.021

Problem 155	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	151	159	176	149	0	882	170	276	31	173
N.S.	1	1.05	1.17	0.99	0.00	5.84	1.13	1.83	0.21	1.15
time (sec)	N/A	0.310	0.295	0.421	0.000	0.139	5.066	0.124	0.159	0.124

Problem 156	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	228	222	305	278	0	1006	0	583	349	562
N.S.	1	0.97	1.34	1.22	0.00	4.41	0.00	2.56	1.53	2.46
time (sec)	N/A	0.388	0.589	0.568	0.000	0.110	0.000	0.137	0.157	1.010

Problem 157	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	187	193	210	212	0	666	0	382	223	363
N.S.	1	1.03	1.12	1.13	0.00	3.56	0.00	2.04	1.19	1.94
time (sec)	N/A	0.298	0.458	0.520	0.000	0.101	0.000	0.132	0.160	1.001

Problem 158	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	146	164	139	138	0	392	0	224	119	174
N.S.	1	1.12	0.95	0.95	0.00	2.68	0.00	1.53	0.82	1.19
time (sec)	N/A	0.297	0.337	0.480	0.000	0.112	0.000	0.127	0.157	0.113

Problem 159	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	134	96	103	0	393	0	114	51	108
N.S.	1	1.25	0.90	0.96	0.00	3.67	0.00	1.07	0.48	1.01
time (sec)	N/A	0.264	0.285	0.437	0.000	0.133	0.000	0.130	0.159	0.982

Problem 160	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	103	103	102	88	0	397	0	122	50	99
N.S.	1	1.00	0.99	0.85	0.00	3.85	0.00	1.18	0.49	0.96
time (sec)	N/A	0.240	0.307	0.376	0.000	0.098	0.000	0.122	0.157	0.089

Problem 161	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	128	141	122	130	0	775	0	189	84	156
N.S.	1	1.10	0.95	1.02	0.00	6.05	0.00	1.48	0.66	1.22
time (sec)	N/A	0.268	0.399	0.461	0.000	0.130	0.000	0.122	0.156	1.032

Problem 162	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	167	178	197	164	0	1086	0	282	223	210
N.S.	1	1.07	1.18	0.98	0.00	6.50	0.00	1.69	1.34	1.26
time (sec)	N/A	0.302	0.547	0.612	0.000	0.155	0.000	0.131	0.156	1.065

Problem 163	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	208	214	289	202	0	1731	0	422	437	261
N.S.	1	1.03	1.39	0.97	0.00	8.32	0.00	2.03	2.10	1.25
time (sec)	N/A	0.326	0.698	0.507	0.000	0.230	0.000	0.128	0.158	1.100

Problem 164	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	236	225	309	276	0	1060	0	597	449	561
N.S.	1	0.95	1.31	1.17	0.00	4.49	0.00	2.53	1.90	2.38
time (sec)	N/A	0.352	0.794	0.599	0.000	0.113	0.000	0.140	0.158	1.045

Problem 165	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	196	196	213	202	0	680	0	390	273	325
N.S.	1	1.00	1.09	1.03	0.00	3.47	0.00	1.99	1.39	1.66
time (sec)	N/A	0.316	0.619	0.592	0.000	0.094	0.000	0.133	0.157	1.094

Problem 166	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	152	166	139	143	0	703	0	226	127	256
N.S.	1	1.09	0.91	0.94	0.00	4.62	0.00	1.49	0.84	1.68
time (sec)	N/A	0.308	0.545	0.496	0.000	0.129	0.000	0.132	0.157	1.120

Problem 167	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	146	145	143	147	0	721	0	235	129	222
N.S.	1	0.99	0.98	1.01	0.00	4.94	0.00	1.61	0.88	1.52
time (sec)	N/A	0.270	0.609	0.417	0.000	0.096	0.000	0.128	0.156	0.146

Problem 168	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	157	151	144	148	0	808	0	258	156	228
N.S.	1	0.96	0.92	0.94	0.00	5.15	0.00	1.64	0.99	1.45
time (sec)	N/A	0.277	0.627	0.434	0.000	0.146	0.000	0.134	0.157	0.144

Problem 169	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	180	187	194	195	0	1410	0	340	220	296
N.S.	1	1.04	1.08	1.08	0.00	7.83	0.00	1.89	1.22	1.64
time (sec)	N/A	0.295	0.964	0.515	0.000	0.170	0.000	0.126	0.156	0.215

Problem 170	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	222	224	291	229	0	1756	0	446	493	361
N.S.	1	1.01	1.31	1.03	0.00	7.91	0.00	2.01	2.22	1.63
time (sec)	N/A	0.320	0.992	0.523	0.000	0.192	0.000	0.134	0.165	1.425

Problem 171	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	265	260	408	268	0	2657	0	609	804	418
N.S.	1	0.98	1.54	1.01	0.00	10.03	0.00	2.30	3.03	1.58
time (sec)	N/A	0.344	1.039	0.580	0.000	0.427	0.000	0.144	0.165	1.323

Problem 172	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	A	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	304	252	322	1372	0	1044	0	2501	651	0
N.S.	1	0.83	1.06	4.51	0.00	3.43	0.00	8.23	2.14	0.00
time (sec)	N/A	0.310	0.710	0.266	0.000	0.144	0.000	0.474	0.212	0.000

Problem 173	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	A	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	250	211	231	968	0	766	0	1366	471	0
N.S.	1	0.84	0.92	3.87	0.00	3.06	0.00	5.46	1.88	0.00
time (sec)	N/A	0.287	0.648	0.263	0.000	0.120	0.000	0.320	0.177	0.000

Problem 174	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	A	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	190	170	160	636	0	528	0	557	320	1207
N.S.	1	0.89	0.84	3.35	0.00	2.78	0.00	2.93	1.68	6.35
time (sec)	N/A	0.259	0.388	0.259	0.000	0.120	0.000	0.198	0.167	44.918

Problem 175	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	140	128	117	376	0	364	0	180	198	872
N.S.	1	0.91	0.84	2.69	0.00	2.60	0.00	1.29	1.41	6.23
time (sec)	N/A	0.230	0.349	0.263	0.000	0.113	0.000	0.138	0.161	14.496

Problem 176	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	145	92	386	0	358	0	157	275	0
N.S.	1	1.28	0.81	3.42	0.00	3.17	0.00	1.39	2.43	0.00
time (sec)	N/A	0.250	0.185	0.265	0.000	0.381	0.000	0.160	0.170	0.000

Problem 177	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	111	115	116	503	0	515	0	205	207	0
N.S.	1	1.04	1.05	4.53	0.00	4.64	0.00	1.85	1.86	0.00
time (sec)	N/A	0.209	0.193	0.268	0.000	0.672	0.000	0.182	0.170	0.000

Problem 178	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	66	61	0	221	0	191	199	179
N.S.	1	1.00	0.69	0.64	0.00	2.33	0.00	2.01	2.09	1.88
time (sec)	N/A	0.209	0.109	0.270	0.000	1.878	0.000	0.222	0.187	1.359

Problem 179	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	147	143	134	149	0	440	0	379	414	295
N.S.	1	0.97	0.91	1.01	0.00	2.99	0.00	2.58	2.82	2.01
time (sec)	N/A	0.236	0.147	0.273	0.000	6.097	0.000	0.261	0.207	1.518

Problem 180	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	198	190	199	281	0	710	0	620	702	428
N.S.	1	0.96	1.01	1.42	0.00	3.59	0.00	3.13	3.55	2.16
time (sec)	N/A	0.255	0.218	0.280	0.000	29.309	0.000	0.357	0.307	1.764

Problem 181	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	255	239	270	449	0	1045	0	937	1049	585
N.S.	1	0.94	1.06	1.76	0.00	4.10	0.00	3.67	4.11	2.29
time (sec)	N/A	0.292	0.290	0.308	0.000	64.583	0.000	0.412	0.526	1.892

Problem 182	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	358	293	432	1848	0	1384	0	4461	860	0
N.S.	1	0.82	1.21	5.16	0.00	3.87	0.00	12.46	2.40	0.00
time (sec)	N/A	0.342	1.007	0.266	0.000	0.163	0.000	0.681	0.229	0.000

Problem 183	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	294	252	319	1372	0	1036	0	2484	651	0
N.S.	1	0.86	1.09	4.67	0.00	3.52	0.00	8.45	2.21	0.00
time (sec)	N/A	0.305	0.720	0.266	0.000	0.157	0.000	0.452	0.261	0.000

Problem 184	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	A	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	250	211	232	968	0	766	0	1036	471	0
N.S.	1	0.84	0.93	3.87	0.00	3.06	0.00	4.14	1.88	0.00
time (sec)	N/A	0.270	0.591	0.262	0.000	0.152	0.000	0.278	0.244	0.000

Problem 185	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	193	169	165	636	0	540	0	283	320	0
N.S.	1	0.88	0.85	3.30	0.00	2.80	0.00	1.47	1.66	0.00
time (sec)	N/A	0.252	0.391	0.266	0.000	0.123	0.000	0.154	0.237	0.000

Problem 186	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	167	186	165	740	0	572	0	270	485	0
N.S.	1	1.11	0.99	4.43	0.00	3.43	0.00	1.62	2.90	0.00
time (sec)	N/A	0.283	1.145	0.263	0.000	0.427	0.000	0.194	0.222	0.000

Problem 187	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	A	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	155	180	136	698	0	537	0	369	504	0
N.S.	1	1.16	0.88	4.50	0.00	3.46	0.00	2.38	3.25	0.00
time (sec)	N/A	0.264	0.290	0.267	0.000	0.919	0.000	0.207	0.254	0.000

Problem 188	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	138	146	171	780	0	767	0	398	403	0
N.S.	1	1.06	1.24	5.65	0.00	5.56	0.00	2.88	2.92	0.00
time (sec)	N/A	0.224	0.234	0.269	0.000	2.540	0.000	0.264	0.711	0.000

Problem 189	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	66	74	0	306	0	285	267	257
N.S.	1	1.00	0.69	0.78	0.00	3.22	0.00	3.00	2.81	2.71
time (sec)	N/A	0.206	0.131	0.271	0.000	5.937	0.000	0.312	0.438	1.463

Problem 190	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	147	143	134	177	0	567	0	516	526	402
N.S.	1	0.97	0.91	1.20	0.00	3.86	0.00	3.51	3.58	2.73
time (sec)	N/A	0.231	0.179	0.276	0.000	27.998	0.000	0.452	0.374	1.647

Problem 191	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	201	191	199	322	0	880	0	815	860	570
N.S.	1	0.95	0.99	1.60	0.00	4.38	0.00	4.05	4.28	2.84
time (sec)	N/A	0.261	0.271	0.277	0.000	64.042	0.000	0.501	0.609	1.810

Problem 192	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	255	239	270	505	0	1252	0	1168	1253	752
N.S.	1	0.94	1.06	1.98	0.00	4.91	0.00	4.58	4.91	2.95
time (sec)	N/A	0.298	0.393	0.280	0.000	130.413	0.000	0.665	1.180	2.074

Problem 193	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	398	334	514	2396	0	1758	0	6983	1098	0
N.S.	1	0.84	1.29	6.02	0.00	4.42	0.00	17.55	2.76	0.00
time (sec)	N/A	0.348	1.203	0.293	0.000	0.175	0.000	0.948	0.374	0.000

Problem 194	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	358	293	431	1848	0	1388	0	3955	860	0
N.S.	1	0.82	1.20	5.16	0.00	3.88	0.00	11.05	2.40	0.00
time (sec)	N/A	0.339	0.945	0.261	0.000	0.172	0.000	0.608	0.312	0.000

Problem 195	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	A	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	304	252	321	1372	0	1046	0	1672	651	0
N.S.	1	0.83	1.06	4.51	0.00	3.44	0.00	5.50	2.14	0.00
time (sec)	N/A	0.299	0.744	0.264	0.000	0.158	0.000	0.341	0.280	0.000

Problem 196	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	246	210	230	968	0	770	0	415	471	0
N.S.	1	0.85	0.93	3.93	0.00	3.13	0.00	1.69	1.91	0.00
time (sec)	N/A	0.273	0.486	0.263	0.000	0.163	0.000	0.161	0.267	0.000

Problem 197	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	218	227	217	1184	0	858	0	419	724	0
N.S.	1	1.04	1.00	5.43	0.00	3.94	0.00	1.92	3.32	0.00
time (sec)	N/A	0.295	0.563	0.270	0.000	0.749	0.000	0.199	0.305	0.000

Problem 198	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F(-2)	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	212	221	113	1250	0	855	0	564	817	0
N.S.	1	1.04	0.53	5.90	0.00	4.03	0.00	2.66	3.85	0.00
time (sec)	N/A	0.286	10.105	0.269	0.000	1.310	0.000	0.238	0.306	0.000

Problem 199	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	199	214	205	1092	0	835	0	707	766	0
N.S.	1	1.08	1.03	5.49	0.00	4.20	0.00	3.55	3.85	0.00
time (sec)	N/A	0.288	0.358	0.273	0.000	3.218	0.000	0.303	0.341	0.000

Problem 200	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	167	177	215	1089	0	1053	0	678	583	0
N.S.	1	1.06	1.29	6.52	0.00	6.31	0.00	4.06	3.49	0.00
time (sec)	N/A	0.242	0.348	0.270	0.000	8.126	0.000	0.402	94.658	0.000

Problem 201	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	66	74	0	391	0	379	335	325
N.S.	1	1.00	0.69	0.78	0.00	4.12	0.00	3.99	3.53	3.42
time (sec)	N/A	0.205	0.140	0.283	0.000	28.887	0.000	0.561	2.292	1.697

Problem 202	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	147	143	134	177	0	693	0	663	638	509
N.S.	1	0.97	0.91	1.20	0.00	4.71	0.00	4.51	4.34	3.46
time (sec)	N/A	0.230	0.195	0.273	0.000	66.661	0.000	0.575	0.574	1.914

Problem 203	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	201	191	199	322	0	1047	0	1003	1018	706
N.S.	1	0.95	0.99	1.60	0.00	5.21	0.00	4.99	5.06	3.51
time (sec)	N/A	0.254	0.300	0.283	0.000	138.521	0.000	0.801	1.058	2.127

Problem 204	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	F(-1)	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	255	239	270	505	0	0	0	1413	1457	917
N.S.	1	0.94	1.06	1.98	0.00	0.00	0.00	5.54	5.71	3.60
time (sec)	N/A	0.287	0.398	0.303	0.000	0.000	0.000	1.057	2.059	2.549

Problem 205	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	A	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	246	209	231	968	0	772	0	1048	471	0
N.S.	1	0.85	0.94	3.93	0.00	3.14	0.00	4.26	1.91	0.00
time (sec)	N/A	0.276	0.482	0.270	0.000	0.134	0.000	0.279	0.170	0.000

Problem 206	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	A	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	193	168	165	636	0	542	0	559	320	0
N.S.	1	0.87	0.85	3.30	0.00	2.81	0.00	2.90	1.66	0.00
time (sec)	N/A	0.246	0.374	0.261	0.000	0.137	0.000	0.204	0.161	0.000

Problem 207	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	A	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	140	127	134	375	0	366	0	233	196	866
N.S.	1	0.91	0.96	2.68	0.00	2.61	0.00	1.66	1.40	6.19
time (sec)	N/A	0.229	0.646	0.266	0.000	0.133	0.000	0.157	0.157	15.874

Problem 208	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	84	83	198	0	251	0	107	102	311
N.S.	1	1.01	1.00	2.39	0.00	3.02	0.00	1.29	1.23	3.75
time (sec)	N/A	0.197	0.174	0.289	0.000	0.121	0.000	0.142	0.153	3.846

Problem 209	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	85	278	0	362	0	121	120	0
N.S.	1	1.00	1.00	3.27	0.00	4.26	0.00	1.42	1.41	0.00
time (sec)	N/A	0.186	0.167	0.299	0.000	0.309	0.000	0.152	0.153	0.000

Problem 210	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	94	94	65	60	0	140	0	180	125	169
N.S.	1	1.00	0.69	0.64	0.00	1.49	0.00	1.91	1.33	1.80
time (sec)	N/A	0.197	0.098	0.304	0.000	0.594	0.000	0.170	0.168	1.495

Problem 211	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	145	142	134	149	0	316	0	377	302	283
N.S.	1	0.98	0.92	1.03	0.00	2.18	0.00	2.60	2.08	1.95
time (sec)	N/A	0.229	0.134	0.314	0.000	2.109	0.000	0.197	0.164	1.612

Problem 212	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	198	190	199	281	0	547	0	620	544	419
N.S.	1	0.96	1.01	1.42	0.00	2.76	0.00	3.13	2.75	2.12
time (sec)	N/A	0.261	0.188	0.317	0.000	6.234	0.000	0.230	0.200	1.828

Problem 213	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	251	238	270	449	0	839	0	933	845	570
N.S.	1	0.95	1.08	1.79	0.00	3.34	0.00	3.72	3.37	2.27
time (sec)	N/A	0.296	0.264	0.339	0.000	29.803	0.000	0.308	0.296	1.975

Problem 214	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	215	225	218	1184	0	872	0	479	320	0
N.S.	1	1.05	1.01	5.51	0.00	4.06	0.00	2.23	1.49	0.00
time (sec)	N/A	0.304	0.596	0.278	0.000	0.528	0.000	0.387	0.166	0.000

Problem 215	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	165	184	163	740	0	584	0	327	198	0
N.S.	1	1.12	0.99	4.48	0.00	3.54	0.00	1.98	1.20	0.00
time (sec)	N/A	0.273	0.982	0.267	0.000	0.532	0.000	0.273	0.161	0.000

Problem 216	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	A	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	111	143	92	386	0	366	0	210	102	0
N.S.	1	1.29	0.83	3.48	0.00	3.30	0.00	1.89	0.92	0.00
time (sec)	N/A	0.254	0.183	0.267	0.000	0.348	0.000	0.201	0.161	0.000

Problem 217	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	85	278	0	360	0	131	44	0
N.S.	1	1.00	1.00	3.27	0.00	4.24	0.00	1.54	0.52	0.00
time (sec)	N/A	0.198	0.156	0.299	0.000	0.254	0.000	0.157	0.156	0.000

Problem 218	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	61	59	0	148	0	271	60	96
N.S.	1	1.00	0.69	0.66	0.00	1.66	0.00	3.04	0.67	1.08
time (sec)	N/A	0.217	0.110	0.303	0.000	0.329	0.000	0.187	0.155	1.472

Problem 219	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	139	137	133	148	0	336	0	616	186	182
N.S.	1	0.99	0.96	1.06	0.00	2.42	0.00	4.43	1.34	1.31
time (sec)	N/A	0.238	0.146	0.316	0.000	1.434	0.000	0.240	0.158	1.606

Problem 220	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	187	185	199	281	0	583	0	1353	388	288
N.S.	1	0.99	1.06	1.50	0.00	3.12	0.00	7.24	2.07	1.54
time (sec)	N/A	0.276	0.222	0.375	0.000	3.919	0.000	0.369	0.166	1.792

Problem 221	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	237	233	270	449	0	887	0	2645	643	409
N.S.	1	0.98	1.14	1.89	0.00	3.74	0.00	11.16	2.71	1.73
time (sec)	N/A	0.302	0.296	0.339	0.000	17.476	0.000	1.000	0.201	1.982

Problem 222	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	261	257	313	1882	0	1271	0	1814	461	0
N.S.	1	0.98	1.20	7.21	0.00	4.87	0.00	6.95	1.77	0.00
time (sec)	N/A	0.307	0.733	0.287	0.000	1.461	0.000	0.654	0.209	0.000

Problem 223	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	212	220	235	1250	0	887	0	1401	304	0
N.S.	1	1.04	1.11	5.90	0.00	4.18	0.00	6.61	1.43	0.00
time (sec)	N/A	0.292	1.330	0.273	0.000	1.263	0.000	0.491	0.169	0.000

Problem 224	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	152	179	133	698	0	561	0	1019	170	0
N.S.	1	1.18	0.88	4.59	0.00	3.69	0.00	6.70	1.12	0.00
time (sec)	N/A	0.261	0.275	0.269	0.000	0.975	0.000	0.345	0.163	0.000

Problem 225	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	111	115	116	503	0	525	0	558	78	0
N.S.	1	1.04	1.05	4.53	0.00	4.73	0.00	5.03	0.70	0.00
time (sec)	N/A	0.205	0.181	0.269	0.000	0.670	0.000	0.231	0.165	0.000

Problem 226	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	64	60	0	141	0	282	43	97
N.S.	1	1.00	0.67	0.63	0.00	1.48	0.00	2.97	0.45	1.02
time (sec)	N/A	0.205	0.101	0.305	0.000	0.421	0.000	0.171	0.165	1.561

Problem 227	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	145	138	125	149	0	337	0	610	121	193
N.S.	1	0.95	0.86	1.03	0.00	2.32	0.00	4.21	0.83	1.33
time (sec)	N/A	0.235	0.146	0.316	0.000	0.951	0.000	0.307	0.153	1.701

Problem 228	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	192	183	197	279	0	565	0	1183	288	304
N.S.	1	0.95	1.03	1.45	0.00	2.94	0.00	6.16	1.50	1.58
time (sec)	N/A	0.252	0.206	0.330	0.000	3.145	0.000	0.465	0.163	1.875

Problem 229	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	246	235	270	449	0	916	0	2081	514	444
N.S.	1	0.96	1.10	1.83	0.00	3.72	0.00	8.46	2.09	1.80
time (sec)	N/A	0.293	0.265	0.352	0.000	24.157	0.000	0.998	0.194	2.042

Problem 230	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	186	186	165	1165	620	1282	14256	2519	1028	1273
N.S.	1	1.00	0.89	6.26	3.33	6.89	76.65	13.54	5.53	6.84
time (sec)	N/A	0.345	0.178	0.311	0.076	0.128	2.585	0.139	0.162	1.503

Problem 231	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	138	138	122	576	364	660	6186	1261	546	676
N.S.	1	1.00	0.88	4.17	2.64	4.78	44.83	9.14	3.96	4.90
time (sec)	N/A	0.283	0.123	0.273	0.048	0.129	1.372	0.132	0.162	1.370

Problem 232	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	90	90	79	189	179	256	1982	490	241	259
N.S.	1	1.00	0.88	2.10	1.99	2.84	22.02	5.44	2.68	2.88
time (sec)	N/A	0.236	0.099	0.229	0.040	0.091	0.652	0.125	0.161	1.213

Problem 233	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	47	41	46	63	83	377	132	75	88
N.S.	1	1.00	0.87	0.98	1.34	1.77	8.02	2.81	1.60	1.87
time (sec)	N/A	0.192	0.042	0.201	0.043	0.090	0.342	0.122	0.162	1.195

Problem 234	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	78	0	0	0	0	0	21	0
N.S.	1	1.00	0.92	0.00	0.00	0.00	0.00	0.00	0.25	0.00
time (sec)	N/A	0.200	0.080	0.000	0.000	0.000	0.000	0.000	0.161	0.000

Problem 235	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	104	112	99	0	0	0	0	0	94	0
N.S.	1	1.08	0.95	0.00	0.00	0.00	0.00	0.00	0.90	0.00
time (sec)	N/A	0.222	0.081	0.000	0.000	0.000	0.000	0.000	0.161	0.000

Problem 236	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	118	123	106	0	0	0	0	0	1218	0
N.S.	1	1.04	0.90	0.00	0.00	0.00	0.00	0.00	10.32	0.00
time (sec)	N/A	0.234	0.099	0.000	0.000	0.000	0.000	0.000	0.169	0.000

Problem 237	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	91	75	101	295	175	1822	423	220	313
N.S.	1	1.00	0.82	1.11	3.24	1.92	20.02	4.65	2.42	3.44
time (sec)	N/A	0.206	0.059	0.283	0.042	0.100	0.589	0.124	0.148	0.157

Problem 238	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	61	80	183	120	1017	278	145	211
N.S.	1	1.00	0.84	1.10	2.51	1.64	13.93	3.81	1.99	2.89
time (sec)	N/A	0.194	0.050	0.200	0.039	0.079	0.431	0.113	0.152	1.181

Problem 239	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	47	61	102	75	488	163	86	130
N.S.	1	1.00	0.85	1.11	1.85	1.36	8.87	2.96	1.56	2.36
time (sec)	N/A	0.178	0.049	0.174	0.031	0.095	0.317	0.122	0.151	1.173

Problem 240	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	52	52	37	0	0	0	0	0	95	0
N.S.	1	1.00	0.71	0.00	0.00	0.00	0.00	0.00	1.83	0.00
time (sec)	N/A	0.160	0.049	0.000	0.000	0.000	0.000	0.000	0.153	0.000

Problem 241	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	65	59	57	0	0	0	0	0	371	0
N.S.	1	0.91	0.88	0.00	0.00	0.00	0.00	0.00	5.71	0.00
time (sec)	N/A	0.180	0.052	0.000	0.000	0.000	0.000	0.000	0.150	0.000

Problem 242	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	70	59	48	0	0	0	0	0	1272	0
N.S.	1	0.84	0.69	0.00	0.00	0.00	0.00	0.00	18.17	0.00
time (sec)	N/A	0.174	0.057	0.000	0.000	0.000	0.000	0.000	0.156	0.000

Problem 243	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	108	114	101	0	0	0	0	0	0	0
N.S.	1	1.06	0.94	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.248	0.100	0.000	0.000	0.000	0.000	0.000	0.202	0.000

Problem 244	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	104	112	99	0	0	0	0	0	0	0
N.S.	1	1.08	0.95	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.249	0.080	0.000	0.000	0.000	0.000	0.000	0.180	0.000

Problem 245	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	107	107	93	0	0	0	0	0	0	0
N.S.	1	1.00	0.87	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.223	0.100	0.000	0.000	0.000	0.000	0.000	0.166	0.000

Problem 246	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	105	119	105	0	0	0	0	0	0	0
N.S.	1	1.13	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.226	0.112	0.000	0.000	0.000	0.000	0.000	0.287	0.000

Problem 247	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	118	123	106	0	0	0	0	0	0	0
N.S.	1	1.04	0.90	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.231	0.095	0.000	0.000	0.000	0.000	0.000	0.232	0.000

Problem 248	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	109	113	98	0	0	0	0	0	0	0
N.S.	1	1.04	0.90	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.224	0.114	0.000	0.000	0.000	0.000	0.000	0.181	0.000

Problem 249	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	131	137	117	0	0	0	0	0	0	0
N.S.	1	1.05	0.89	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.264	0.109	0.000	0.000	0.000	0.000	0.000	0.310	0.000

Problem 250	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	109	124	114	0	0	0	0	0	106	0
N.S.	1	1.14	1.05	0.00	0.00	0.00	0.00	0.00	0.97	0.00
time (sec)	N/A	0.254	0.165	0.000	0.000	0.000	0.000	0.000	0.156	0.000

Problem 251	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	30	30	32	45	38	63	56	35	35
N.S.	1	1.00	1.00	1.07	1.50	1.27	2.10	1.87	1.17	1.17
time (sec)	N/A	0.159	0.093	0.888	0.104	0.105	1.282	0.131	0.151	1.175

Problem 252	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	125	134	109	0	0	0	0	0	44	0
N.S.	1	1.07	0.87	0.00	0.00	0.00	0.00	0.00	0.35	0.00
time (sec)	N/A	0.253	0.104	0.000	0.000	0.000	0.000	0.000	0.156	0.000

Problem 253	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	110	152	124	0	0	0	0	0	70	0
N.S.	1	1.38	1.13	0.00	0.00	0.00	0.00	0.00	0.64	0.00
time (sec)	N/A	0.267	0.114	0.000	0.000	0.000	0.000	0.000	0.152	0.000

Problem 254	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	110	117	115	0	0	0	0	0	106	0
N.S.	1	1.06	1.05	0.00	0.00	0.00	0.00	0.00	0.96	0.00
time (sec)	N/A	0.226	0.161	0.000	0.000	0.000	0.000	0.000	0.157	0.000

Problem 255	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	B	F(-2)	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	122	122	82	161	0	324	0	1049	142	358
N.S.	1	1.00	0.67	1.32	0.00	2.66	0.00	8.60	1.16	2.93
time (sec)	N/A	0.232	0.096	0.582	0.000	0.110	0.000	0.149	0.148	1.294

Problem 256	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F(-2)	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	205	183	112	506	0	884	0	0	178	874
N.S.	1	0.89	0.55	2.47	0.00	4.31	0.00	0.00	0.87	4.26
time (sec)	N/A	0.274	0.207	0.545	0.000	0.135	0.000	0.000	0.150	1.614

Problem 257	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F(-2)	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	296	243	165	1187	0	1741	0	0	214	1657
N.S.	1	0.82	0.56	4.01	0.00	5.88	0.00	0.00	0.72	5.60
time (sec)	N/A	0.300	0.217	0.561	0.000	0.144	0.000	0.000	0.152	2.367

Problem 258	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	129	134	108	0	0	0	0	0	44	0
N.S.	1	1.04	0.84	0.00	0.00	0.00	0.00	0.00	0.34	0.00
time (sec)	N/A	0.243	0.107	0.000	0.000	0.000	0.000	0.000	0.149	0.000

Problem 259	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	110	150	123	0	0	0	0	0	70	0
N.S.	1	1.36	1.12	0.00	0.00	0.00	0.00	0.00	0.64	0.00
time (sec)	N/A	0.255	0.114	0.000	0.000	0.000	0.000	0.000	0.149	0.000

Problem 260	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	111	118	105	0	0	0	0	0	106	0
N.S.	1	1.06	0.95	0.00	0.00	0.00	0.00	0.00	0.95	0.00
time (sec)	N/A	0.237	0.158	0.000	0.000	0.000	0.000	0.000	0.150	0.000

Problem 261	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	B	F(-2)	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	123	84	160	0	326	0	1050	142	360
N.S.	1	1.00	0.68	1.30	0.00	2.65	0.00	8.54	1.15	2.93
time (sec)	N/A	0.238	0.095	0.539	0.000	0.109	0.000	0.154	0.152	1.261

Problem 262	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F(-2)	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	205	182	180	505	0	884	0	0	178	869
N.S.	1	0.89	0.88	2.46	0.00	4.31	0.00	0.00	0.87	4.24
time (sec)	N/A	0.268	0.151	0.544	0.000	0.125	0.000	0.000	0.153	1.660

Problem 263	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F(-2)	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	297	241	149	1188	0	1740	0	0	214	1659
N.S.	1	0.81	0.50	4.00	0.00	5.86	0.00	0.00	0.72	5.59
time (sec)	N/A	0.307	0.278	0.556	0.000	0.179	0.000	0.000	0.153	2.389

Problem 264	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	83	121	106	0	0	0	0	0	514	0
N.S.	1	1.46	1.28	0.00	0.00	0.00	0.00	0.00	6.19	0.00
time (sec)	N/A	0.264	0.229	0.000	0.000	0.000	0.000	0.000	0.158	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [97] had the largest ratio of [.400000000000000022]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	2	2	1.00	18	0.111
2	A	2	2	1.00	18	0.111
3	A	2	2	1.00	18	0.111
4	A	2	2	1.00	16	0.125
5	A	2	2	1.00	11	0.182
6	A	2	2	1.00	18	0.111
7	A	2	2	1.00	18	0.111
8	A	2	2	1.00	18	0.111
9	A	2	2	1.00	18	0.111
10	A	2	2	1.00	18	0.111
11	A	2	2	1.00	18	0.111
12	A	2	2	1.00	20	0.100
13	A	2	2	1.00	20	0.100
14	A	2	2	1.00	20	0.100
15	A	2	2	1.00	18	0.111
16	A	2	2	1.00	13	0.154
17	A	2	2	1.00	20	0.100
18	A	2	2	1.00	20	0.100
19	A	2	2	1.00	20	0.100
20	A	3	3	1.03	20	0.150
21	A	2	2	1.00	20	0.100

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	2	2	1.00	20	0.100
23	A	2	2	1.00	20	0.100
24	A	2	2	1.00	20	0.100
25	A	2	2	1.00	20	0.100
26	A	2	2	1.00	20	0.100
27	A	2	2	1.00	20	0.100
28	A	2	2	1.00	20	0.100
29	A	2	2	1.00	18	0.111
30	A	2	2	1.00	13	0.154
31	A	2	2	1.00	20	0.100
32	A	2	2	1.00	20	0.100
33	A	2	2	1.00	20	0.100
34	A	2	2	1.00	20	0.100
35	A	3	3	1.02	20	0.150
36	A	2	2	1.00	20	0.100
37	A	3	3	0.98	20	0.150
38	A	2	2	1.00	20	0.100
39	A	2	2	1.00	20	0.100
40	A	2	2	1.00	20	0.100
41	A	2	2	1.00	20	0.100
42	A	2	2	1.00	20	0.100
43	A	2	2	1.00	20	0.100
44	A	2	2	1.00	20	0.100
45	A	2	2	1.00	20	0.100
46	A	2	2	1.00	20	0.100
47	A	2	2	1.00	20	0.100
48	A	2	2	1.00	18	0.111
49	A	2	2	1.00	13	0.154
50	A	2	2	1.00	20	0.100
51	A	2	2	1.00	20	0.100
52	A	2	2	1.00	20	0.100
53	A	2	2	1.00	20	0.100

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
54	A	2	2	1.00	20	0.100
55	A	2	2	1.00	20	0.100
56	A	2	2	1.00	20	0.100
57	A	3	3	1.00	20	0.150
58	A	2	2	1.00	20	0.100
59	A	3	3	0.97	20	0.150
60	A	4	4	0.95	20	0.200
61	A	5	5	0.93	20	0.250
62	A	2	2	1.00	20	0.100
63	A	2	2	1.00	20	0.100
64	A	2	2	1.00	20	0.100
65	A	2	2	1.00	20	0.100
66	A	2	2	1.00	20	0.100
67	A	2	2	1.00	20	0.100
68	A	2	2	1.00	20	0.100
69	A	2	2	1.00	20	0.100
70	A	2	2	1.00	20	0.100
71	A	2	2	1.00	20	0.100
72	A	2	2	1.00	20	0.100
73	A	2	2	1.00	20	0.100
74	A	2	2	1.00	20	0.100
75	A	2	2	1.00	20	0.100
76	A	2	2	1.00	20	0.100
77	A	2	2	1.00	18	0.111
78	A	2	2	1.00	13	0.154
79	A	2	2	1.00	20	0.100
80	A	2	2	1.00	20	0.100
81	A	2	2	1.00	20	0.100
82	A	2	2	1.00	20	0.100
83	A	2	2	1.00	20	0.100
84	A	2	2	1.00	20	0.100
85	A	2	2	1.00	20	0.100

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
86	A	2	2	1.00	20	0.100
87	A	2	2	1.00	20	0.100
88	A	2	2	1.00	20	0.100
89	A	2	2	1.00	20	0.100
90	A	3	3	0.98	20	0.150
91	A	2	2	1.00	20	0.100
92	A	3	3	0.97	20	0.150
93	A	4	4	0.95	20	0.200
94	A	5	5	0.93	20	0.250
95	A	6	6	0.92	20	0.300
96	A	7	7	0.92	20	0.350
97	A	8	8	0.91	20	0.400
98	A	2	2	1.00	20	0.100
99	A	2	2	1.00	20	0.100
100	A	2	2	1.00	20	0.100
101	A	2	2	1.00	20	0.100
102	A	2	2	1.00	20	0.100
103	A	2	2	1.00	20	0.100
104	A	2	2	1.00	18	0.111
105	A	2	2	1.00	13	0.154
106	A	2	2	1.00	20	0.100
107	A	2	2	1.00	20	0.100
108	A	2	2	1.00	20	0.100
109	A	2	2	1.00	20	0.100
110	A	2	2	1.00	20	0.100
111	A	2	2	1.00	20	0.100
112	A	2	2	1.00	18	0.111
113	A	2	2	1.00	13	0.154
114	A	2	2	1.00	20	0.100
115	A	2	2	1.00	20	0.100
116	A	2	2	1.00	20	0.100
117	A	2	2	1.00	20	0.100

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
118	A	2	2	1.00	20	0.100
119	A	2	2	1.00	18	0.111
120	A	1	1	1.00	13	0.077
121	A	2	2	1.00	20	0.100
122	A	2	2	1.00	20	0.100
123	A	2	2	1.00	20	0.100
124	A	2	2	1.00	20	0.100
125	A	2	2	1.00	20	0.100
126	A	2	2	1.00	20	0.100
127	A	2	2	1.00	20	0.100
128	A	2	2	1.00	20	0.100
129	A	2	2	1.00	20	0.100
130	A	2	2	1.00	20	0.100
131	A	2	2	1.00	20	0.100
132	A	2	2	1.00	22	0.091
133	A	2	2	1.00	22	0.091
134	A	2	2	1.00	22	0.091
135	A	2	2	1.00	22	0.091
136	A	2	2	1.00	22	0.091
137	A	2	2	1.00	22	0.091
138	A	2	2	1.00	22	0.091
139	A	2	2	1.00	22	0.091
140	A	2	2	1.00	22	0.091
141	A	2	2	1.00	22	0.091
142	A	2	2	1.00	22	0.091
143	A	2	2	1.00	22	0.091
144	A	2	2	1.00	22	0.091
145	A	2	2	1.00	22	0.091
146	A	2	2	1.00	22	0.091
147	A	2	2	1.00	22	0.091
148	A	8	7	0.92	22	0.318
149	A	7	6	0.93	22	0.273

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
150	A	6	5	0.95	22	0.227
151	A	5	4	0.97	22	0.182
152	A	4	3	1.00	22	0.136
153	A	4	3	1.00	22	0.136
154	A	5	4	1.03	22	0.182
155	A	6	5	1.05	22	0.227
156	A	8	7	0.97	22	0.318
157	A	7	6	1.03	22	0.273
158	A	6	5	1.12	22	0.227
159	A	5	4	1.25	22	0.182
160	A	4	3	1.00	22	0.136
161	A	5	4	1.10	22	0.182
162	A	6	5	1.07	22	0.227
163	A	7	6	1.03	22	0.273
164	A	8	7	0.95	22	0.318
165	A	7	6	1.00	22	0.273
166	A	6	5	1.09	22	0.227
167	A	5	4	0.99	22	0.182
168	A	5	4	0.96	22	0.182
169	A	6	5	1.04	22	0.227
170	A	7	6	1.01	22	0.273
171	A	8	7	0.98	22	0.318
172	A	8	7	0.83	24	0.292
173	A	7	6	0.84	24	0.250
174	A	6	5	0.89	24	0.208
175	A	5	4	0.91	24	0.167
176	A	5	4	1.28	24	0.167
177	A	5	4	1.04	24	0.167
178	A	2	2	1.00	24	0.083
179	A	3	3	0.97	24	0.125
180	A	4	4	0.96	24	0.167
181	A	5	5	0.94	24	0.208

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
182	A	9	8	0.82	24	0.333
183	A	8	7	0.86	24	0.292
184	A	7	6	0.84	24	0.250
185	A	6	5	0.88	24	0.208
186	A	6	5	1.11	24	0.208
187	A	6	5	1.16	24	0.208
188	A	6	5	1.06	24	0.208
189	A	2	2	1.00	24	0.083
190	A	3	3	0.97	24	0.125
191	A	4	4	0.95	24	0.167
192	A	5	5	0.94	24	0.208
193	A	10	9	0.84	24	0.375
194	A	9	8	0.82	24	0.333
195	A	8	7	0.83	24	0.292
196	A	7	6	0.85	24	0.250
197	A	7	6	1.04	24	0.250
198	A	7	6	1.04	24	0.250
199	A	7	6	1.08	24	0.250
200	A	7	6	1.06	24	0.250
201	A	2	2	1.00	24	0.083
202	A	3	3	0.97	24	0.125
203	A	4	4	0.95	24	0.167
204	A	5	5	0.94	24	0.208
205	A	7	6	0.85	24	0.250
206	A	6	5	0.87	24	0.208
207	A	5	4	0.91	24	0.167
208	A	4	3	1.01	24	0.125
209	A	4	3	1.00	24	0.125
210	A	2	2	1.00	24	0.083
211	A	3	3	0.98	24	0.125
212	A	4	4	0.96	24	0.167
213	A	5	5	0.95	24	0.208
Continued on next page						

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
214	A	7	6	1.05	24	0.250
215	A	6	5	1.12	24	0.208
216	A	5	4	1.29	24	0.167
217	A	4	3	1.00	24	0.125
218	A	2	2	1.00	24	0.083
219	A	3	3	0.99	24	0.125
220	A	4	4	0.99	24	0.167
221	A	5	5	0.98	24	0.208
222	A	8	7	0.98	24	0.292
223	A	7	6	1.04	24	0.250
224	A	6	5	1.18	24	0.208
225	A	5	4	1.04	24	0.167
226	A	2	2	1.00	24	0.083
227	A	3	3	0.95	24	0.125
228	A	4	4	0.95	24	0.167
229	A	5	5	0.96	24	0.208
230	A	2	2	1.00	20	0.100
231	A	2	2	1.00	20	0.100
232	A	2	2	1.00	18	0.111
233	A	2	2	1.00	13	0.154
234	A	2	2	1.00	20	0.100
235	A	2	2	1.08	20	0.100
236	A	2	2	1.04	20	0.100
237	A	2	2	1.00	20	0.100
238	A	2	2	1.00	20	0.100
239	A	2	2	1.00	18	0.111
240	A	2	2	1.00	20	0.100
241	A	2	2	0.91	20	0.100
242	A	2	2	0.84	20	0.100
243	A	2	2	1.06	22	0.091
244	A	2	2	1.08	20	0.100
245	A	2	2	1.00	22	0.091

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
246	A	2	2	1.13	22	0.091
247	A	2	2	1.04	20	0.100
248	A	2	2	1.04	22	0.091
249	A	3	3	1.05	20	0.150
250	A	3	3	1.14	24	0.125
251	A	1	1	1.00	37	0.027
252	A	3	3	1.07	22	0.136
253	A	3	3	1.38	24	0.125
254	A	3	3	1.06	24	0.125
255	A	2	2	1.00	24	0.083
256	A	3	3	0.89	24	0.125
257	A	4	4	0.82	24	0.167
258	A	3	3	1.04	22	0.136
259	A	3	3	1.36	24	0.125
260	A	3	3	1.06	24	0.125
261	A	2	2	1.00	24	0.083
262	A	3	3	0.89	24	0.125
263	A	4	4	0.81	24	0.167
264	A	3	3	1.46	42	0.071

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int (a + bx)(A + Bx)(d + ex)^4 dx$	122
3.2	$\int (a + bx)(A + Bx)(d + ex)^3 dx$	130
3.3	$\int (a + bx)(A + Bx)(d + ex)^2 dx$	137
3.4	$\int (a + bx)(A + Bx)(d + ex) dx$	143
3.5	$\int (a + bx)(A + Bx) dx$	149
3.6	$\int \frac{(a+bx)(A+Bx)}{d+ex} dx$	154
3.7	$\int \frac{(a+bx)(A+Bx)}{(d+ex)^2} dx$	160
3.8	$\int \frac{(a+bx)(A+Bx)}{(d+ex)^3} dx$	166
3.9	$\int \frac{(a+bx)(A+Bx)}{(d+ex)^4} dx$	172
3.10	$\int \frac{(a+bx)(A+Bx)}{(d+ex)^5} dx$	178
3.11	$\int \frac{(a+bx)(A+Bx)}{(d+ex)^6} dx$	184
3.12	$\int (a + bx)^2(A + Bx)(d + ex)^4 dx$	190
3.13	$\int (a + bx)^2(A + Bx)(d + ex)^3 dx$	200
3.14	$\int (a + bx)^2(A + Bx)(d + ex)^2 dx$	209
3.15	$\int (a + bx)^2(A + Bx)(d + ex) dx$	216
3.16	$\int (a + bx)^2(A + Bx) dx$	222
3.17	$\int \frac{(a+bx)^2(A+Bx)}{d+ex} dx$	227
3.18	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^2} dx$	233
3.19	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^3} dx$	240
3.20	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^4} dx$	246
3.21	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^5} dx$	252
3.22	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^6} dx$	258
3.23	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^7} dx$	264
3.24	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^8} dx$	270

3.25	$\int (a + bx)^3(A + Bx)(d + ex)^5 dx$	277
3.26	$\int (a + bx)^3(A + Bx)(d + ex)^4 dx$	290
3.27	$\int (a + bx)^3(A + Bx)(d + ex)^3 dx$	301
3.28	$\int (a + bx)^3(A + Bx)(d + ex)^2 dx$	311
3.29	$\int (a + bx)^3(A + Bx)(d + ex) dx$	320
3.30	$\int (a + bx)^3(A + Bx) dx$	327
3.31	$\int \frac{(a+bx)^3(A+Bx)}{d+ex} dx$	333
3.32	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^2} dx$	340
3.33	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^3} dx$	348
3.34	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^4} dx$	356
3.35	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^5} dx$	363
3.36	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^6} dx$	370
3.37	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^7} dx$	377
3.38	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^8} dx$	385
3.39	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^9} dx$	392
3.40	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{10}} dx$	399
3.41	$\int (a + bx)^6(A + Bx)(d + ex)^8 dx$	406
3.42	$\int (a + bx)^6(A + Bx)(d + ex)^7 dx$	417
3.43	$\int (a + bx)^6(A + Bx)(d + ex)^6 dx$	428
3.44	$\int (a + bx)^6(A + Bx)(d + ex)^5 dx$	440
3.45	$\int (a + bx)^6(A + Bx)(d + ex)^4 dx$	451
3.46	$\int (a + bx)^6(A + Bx)(d + ex)^3 dx$	463
3.47	$\int (a + bx)^6(A + Bx)(d + ex)^2 dx$	476
3.48	$\int (a + bx)^6(A + Bx)(d + ex) dx$	488
3.49	$\int (a + bx)^6(A + Bx) dx$	497
3.50	$\int \frac{(a+bx)^6(A+Bx)}{d+ex} dx$	503
3.51	$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^2} dx$	515
3.52	$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^3} dx$	526
3.53	$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^4} dx$	536
3.54	$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^5} dx$	547
3.55	$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^6} dx$	557
3.56	$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^7} dx$	567
3.57	$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^8} dx$	577
3.58	$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^9} dx$	587
3.59	$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{10}} dx$	596

3.60	$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{11}} dx$	605
3.61	$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{12}} dx$	615
3.62	$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{13}} dx$	626
3.63	$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{14}} dx$	636
3.64	$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{15}} dx$	646
3.65	$\int (a+bx)^{10}(A+Bx)(d+ex)^{13} dx$	656
3.66	$\int (a+bx)^{10}(A+Bx)(d+ex)^{12} dx$	667
3.67	$\int (a+bx)^{10}(A+Bx)(d+ex)^{11} dx$	678
3.68	$\int (a+bx)^{10}(A+Bx)(d+ex)^{10} dx$	689
3.69	$\int (a+bx)^{10}(A+Bx)(d+ex)^9 dx$	700
3.70	$\int (a+bx)^{10}(A+Bx)(d+ex)^8 dx$	712
3.71	$\int (a+bx)^{10}(A+Bx)(d+ex)^7 dx$	723
3.72	$\int (a+bx)^{10}(A+Bx)(d+ex)^6 dx$	734
3.73	$\int (a+bx)^{10}(A+Bx)(d+ex)^5 dx$	745
3.74	$\int (a+bx)^{10}(A+Bx)(d+ex)^4 dx$	756
3.75	$\int (a+bx)^{10}(A+Bx)(d+ex)^3 dx$	767
3.76	$\int (a+bx)^{10}(A+Bx)(d+ex)^2 dx$	778
3.77	$\int (a+bx)^{10}(A+Bx)(d+ex) dx$	789
3.78	$\int (a+bx)^{10}(A+Bx) dx$	800
3.79	$\int \frac{(a+bx)^{10}(A+Bx)}{d+ex} dx$	808
3.80	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^2} dx$	819
3.81	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^3} dx$	831
3.82	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^4} dx$	842
3.83	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^5} dx$	853
3.84	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^6} dx$	863
3.85	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^7} dx$	873
3.86	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^8} dx$	883
3.87	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^9} dx$	893
3.88	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{10}} dx$	903
3.89	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{11}} dx$	914
3.90	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{12}} dx$	925
3.91	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{13}} dx$	935
3.92	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{14}} dx$	944
3.93	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{15}} dx$	954

3.94	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{16}} dx$	965
3.95	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{17}} dx$	976
3.96	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{18}} dx$	988
3.97	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{19}} dx$	1002
3.98	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{20}} dx$	1019
3.99	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{21}} dx$	1030
3.100	$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{22}} dx$	1041
3.101	$\int \frac{(A+Bx)(d+ex)^4}{a+bx} dx$	1052
3.102	$\int \frac{(A+Bx)(d+ex)^3}{a+bx} dx$	1060
3.103	$\int \frac{(A+Bx)(d+ex)^2}{a+bx} dx$	1067
3.104	$\int \frac{(A+Bx)(d+ex)}{a+bx} dx$	1073
3.105	$\int \frac{A+Bx}{a+bx} dx$	1079
3.106	$\int \frac{A+Bx}{(a+bx)(d+ex)} dx$	1084
3.107	$\int \frac{A+Bx}{(a+bx)(d+ex)^2} dx$	1089
3.108	$\int \frac{A+Bx}{(a+bx)(d+ex)^3} dx$	1095
3.109	$\int \frac{A+Bx}{(a+bx)(d+ex)^4} dx$	1102
3.110	$\int \frac{(A+Bx)(d+ex)^3}{(a+bx)^2} dx$	1109
3.111	$\int \frac{(A+Bx)(d+ex)^2}{(a+bx)^2} dx$	1117
3.112	$\int \frac{(A+Bx)(d+ex)}{(a+bx)^2} dx$	1124
3.113	$\int \frac{A+Bx}{(a+bx)^2} dx$	1130
3.114	$\int \frac{A+Bx}{(a+bx)^2(d+ex)} dx$	1135
3.115	$\int \frac{A+Bx}{(a+bx)^2(d+ex)^2} dx$	1141
3.116	$\int \frac{A+Bx}{(a+bx)^2(d+ex)^3} dx$	1148
3.117	$\int \frac{(A+Bx)(d+ex)^3}{(a+bx)^3} dx$	1156
3.118	$\int \frac{(A+Bx)(d+ex)^2}{(a+bx)^3} dx$	1163
3.119	$\int \frac{(A+Bx)(d+ex)}{(a+bx)^3} dx$	1169
3.120	$\int \frac{A+Bx}{(a+bx)^3} dx$	1175
3.121	$\int \frac{A+Bx}{(a+bx)^3(d+ex)} dx$	1180
3.122	$\int \frac{A+Bx}{(a+bx)^3(d+ex)^2} dx$	1187
3.123	$\int \frac{A+Bx}{(a+bx)^3(d+ex)^3} dx$	1195
3.124	$\int (a+bx)(A+Bx)(d+ex)^{5/2} dx$	1205
3.125	$\int (a+bx)(A+Bx)(d+ex)^{3/2} dx$	1212
3.126	$\int (a+bx)(A+Bx)\sqrt{d+ex} dx$	1219
3.127	$\int \frac{(a+bx)(A+Bx)}{\sqrt{d+ex}} dx$	1225

3.128	$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{3/2}} dx$	1231
3.129	$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{5/2}} dx$	1237
3.130	$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{7/2}} dx$	1243
3.131	$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{9/2}} dx$	1249
3.132	$\int (a+bx)^2(A+Bx)(d+ex)^{5/2} dx$	1256
3.133	$\int (a+bx)^2(A+Bx)(d+ex)^{3/2} dx$	1263
3.134	$\int (a+bx)^2(A+Bx)\sqrt{d+ex} dx$	1270
3.135	$\int \frac{(a+bx)^2(A+Bx)}{\sqrt{d+ex}} dx$	1277
3.136	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{3/2}} dx$	1283
3.137	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{5/2}} dx$	1289
3.138	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{7/2}} dx$	1296
3.139	$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{9/2}} dx$	1303
3.140	$\int (a+bx)^3(A+Bx)(d+ex)^{5/2} dx$	1310
3.141	$\int (a+bx)^3(A+Bx)(d+ex)^{3/2} dx$	1319
3.142	$\int (a+bx)^3(A+Bx)\sqrt{d+ex} dx$	1328
3.143	$\int \frac{(a+bx)^3(A+Bx)}{\sqrt{d+ex}} dx$	1336
3.144	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{3/2}} dx$	1344
3.145	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{5/2}} dx$	1351
3.146	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{7/2}} dx$	1358
3.147	$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{9/2}} dx$	1366
3.148	$\int \frac{(A+Bx)(d+ex)^{7/2}}{a+bx} dx$	1374
3.149	$\int \frac{(A+Bx)(d+ex)^{5/2}}{a+bx} dx$	1384
3.150	$\int \frac{(A+Bx)(d+ex)^{3/2}}{a+bx} dx$	1393
3.151	$\int \frac{(A+Bx)\sqrt{d+ex}}{a+bx} dx$	1401
3.152	$\int \frac{A+Bx}{(a+bx)\sqrt{d+ex}} dx$	1408
3.153	$\int \frac{A+Bx}{(a+bx)(d+ex)^{3/2}} dx$	1414
3.154	$\int \frac{A+Bx}{(a+bx)(d+ex)^{5/2}} dx$	1420
3.155	$\int \frac{A+Bx}{(a+bx)(d+ex)^{7/2}} dx$	1427
3.156	$\int \frac{(A+Bx)(d+ex)^{7/2}}{(a+bx)^2} dx$	1435
3.157	$\int \frac{(A+Bx)(d+ex)^{5/2}}{(a+bx)^2} dx$	1446
3.158	$\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^2} dx$	1455
3.159	$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^2} dx$	1463
3.160	$\int \frac{A+Bx}{(a+bx)^2\sqrt{d+ex}} dx$	1470

3.161	$\int \frac{A+Bx}{(a+bx)^2(d+ex)^{3/2}} dx$	1476
3.162	$\int \frac{A+Bx}{(a+bx)^2(d+ex)^{5/2}} dx$	1483
3.163	$\int \frac{A+Bx}{(a+bx)^2(d+ex)^{7/2}} dx$	1491
3.164	$\int \frac{(A+Bx)(d+ex)^{7/2}}{(a+bx)^3} dx$	1500
3.165	$\int \frac{(A+Bx)(d+ex)^{5/2}}{(a+bx)^3} dx$	1511
3.166	$\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^3} dx$	1521
3.167	$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^3} dx$	1529
3.168	$\int \frac{A+Bx}{(a+bx)^3\sqrt{d+ex}} dx$	1536
3.169	$\int \frac{A+Bx}{(a+bx)^3(d+ex)^{3/2}} dx$	1543
3.170	$\int \frac{A+Bx}{(a+bx)^3(d+ex)^{5/2}} dx$	1551
3.171	$\int \frac{A+Bx}{(a+bx)^3(d+ex)^{7/2}} dx$	1560
3.172	$\int \sqrt{a+bx}(A+Bx)(d+ex)^{5/2} dx$	1571
3.173	$\int \sqrt{a+bx}(A+Bx)(d+ex)^{3/2} dx$	1582
3.174	$\int \sqrt{a+bx}(A+Bx)\sqrt{d+ex} dx$	1591
3.175	$\int \frac{\sqrt{a+bx}(A+Bx)}{\sqrt{d+ex}} dx$	1600
3.176	$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{3/2}} dx$	1608
3.177	$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{5/2}} dx$	1615
3.178	$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{7/2}} dx$	1622
3.179	$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{9/2}} dx$	1628
3.180	$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{11/2}} dx$	1635
3.181	$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{13/2}} dx$	1643
3.182	$\int (a+bx)^{3/2}(A+Bx)(d+ex)^{5/2} dx$	1653
3.183	$\int (a+bx)^{3/2}(A+Bx)(d+ex)^{3/2} dx$	1666
3.184	$\int (a+bx)^{3/2}(A+Bx)\sqrt{d+ex} dx$	1677
3.185	$\int \frac{(a+bx)^{3/2}(A+Bx)}{\sqrt{d+ex}} dx$	1686
3.186	$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{3/2}} dx$	1694
3.187	$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{5/2}} dx$	1702
3.188	$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{7/2}} dx$	1710
3.189	$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{9/2}} dx$	1718
3.190	$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{11/2}} dx$	1724
3.191	$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{13/2}} dx$	1732
3.192	$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{15/2}} dx$	1741
3.193	$\int (a+bx)^{5/2}(A+Bx)(d+ex)^{5/2} dx$	1751

3.194	$\int (a + bx)^{5/2}(A + Bx)(d + ex)^{3/2} dx$	1767
3.195	$\int (a + bx)^{5/2}(A + Bx)\sqrt{d + ex} dx$	1780
3.196	$\int \frac{(a+bx)^{5/2}(A+Bx)}{\sqrt{d+ex}} dx$	1791
3.197	$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{3/2}} dx$	1800
3.198	$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{5/2}} dx$	1810
3.199	$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{7/2}} dx$	1820
3.200	$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{9/2}} dx$	1830
3.201	$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{11/2}} dx$	1839
3.202	$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{13/2}} dx$	1846
3.203	$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{15/2}} dx$	1854
3.204	$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{17/2}} dx$	1863
3.205	$\int \frac{(A+Bx)(d+ex)^{5/2}}{\sqrt{a+bx}} dx$	1872
3.206	$\int \frac{(A+Bx)(d+ex)^{3/2}}{\sqrt{a+bx}} dx$	1882
3.207	$\int \frac{(A+Bx)\sqrt{d+ex}}{\sqrt{a+bx}} dx$	1890
3.208	$\int \frac{A+Bx}{\sqrt{a+bx}\sqrt{d+ex}} dx$	1898
3.209	$\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{3/2}} dx$	1904
3.210	$\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{5/2}} dx$	1910
3.211	$\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{7/2}} dx$	1916
3.212	$\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{9/2}} dx$	1923
3.213	$\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{11/2}} dx$	1931
3.214	$\int \frac{(A+Bx)(d+ex)^{5/2}}{(a+bx)^{3/2}} dx$	1941
3.215	$\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^{3/2}} dx$	1951
3.216	$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^{3/2}} dx$	1959
3.217	$\int \frac{A+Bx}{(a+bx)^{3/2}\sqrt{d+ex}} dx$	1966
3.218	$\int \frac{A+Bx}{(a+bx)^{3/2}(d+ex)^{3/2}} dx$	1972
3.219	$\int \frac{A+Bx}{(a+bx)^{3/2}(d+ex)^{5/2}} dx$	1978
3.220	$\int \frac{A+Bx}{(a+bx)^{3/2}(d+ex)^{7/2}} dx$	1985
3.221	$\int \frac{A+Bx}{(a+bx)^{3/2}(d+ex)^{9/2}} dx$	1993
3.222	$\int \frac{(A+Bx)(d+ex)^{7/2}}{(a+bx)^{5/2}} dx$	2002
3.223	$\int \frac{(A+Bx)(d+ex)^{5/2}}{(a+bx)^{5/2}} dx$	2013
3.224	$\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^{5/2}} dx$	2023
3.225	$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^{5/2}} dx$	2032

3.226	$\int \frac{A+Bx}{(a+bx)^{5/2}\sqrt{d+ex}} dx$	2040
3.227	$\int \frac{A+Bx}{(a+bx)^{5/2}(d+ex)^{3/2}} dx$	2046
3.228	$\int \frac{A+Bx}{(a+bx)^{5/2}(d+ex)^{5/2}} dx$	2053
3.229	$\int \frac{A+Bx}{(a+bx)^{5/2}(d+ex)^{7/2}} dx$	2061
3.230	$\int (a+bx)^3(A+Bx)(d+ex)^m dx$	2070
3.231	$\int (a+bx)^2(A+Bx)(d+ex)^m dx$	2080
3.232	$\int (a+bx)(A+Bx)(d+ex)^m dx$	2090
3.233	$\int (A+Bx)(d+ex)^m dx$	2098
3.234	$\int \frac{(A+Bx)(d+ex)^m}{a+bx} dx$	2104
3.235	$\int \frac{(A+Bx)(d+ex)^m}{(a+bx)^2} dx$	2109
3.236	$\int \frac{(A+Bx)(d+ex)^m}{(a+bx)^3} dx$	2114
3.237	$\int (1-2x)(2+3x)^m(3+5x)^3 dx$	2120
3.238	$\int (1-2x)(2+3x)^m(3+5x)^2 dx$	2128
3.239	$\int (1-2x)(2+3x)^m(3+5x) dx$	2135
3.240	$\int \frac{(1-2x)(2+3x)^m}{3+5x} dx$	2141
3.241	$\int \frac{(1-2x)(2+3x)^m}{(3+5x)^2} dx$	2146
3.242	$\int \frac{(1-2x)(2+3x)^m}{(3+5x)^3} dx$	2151
3.243	$\int \frac{(c+dx)^{1+m}(e+fx)}{(a+bx)^2} dx$	2157
3.244	$\int \frac{(c+dx)^m(e+fx)}{(a+bx)^2} dx$	2163
3.245	$\int \frac{(c+dx)^{-1+m}(e+fx)}{(a+bx)^2} dx$	2169
3.246	$\int \frac{(c+dx)^{1+m}(e+fx)}{(a+bx)^3} dx$	2175
3.247	$\int \frac{(c+dx)^m(e+fx)}{(a+bx)^3} dx$	2181
3.248	$\int \frac{(c+dx)^{-1+m}(e+fx)}{(a+bx)^3} dx$	2187
3.249	$\int (a+bx)^m(c+dx)^n(e+fx) dx$	2193
3.250	$\int (a+bx)^p(c+dx)^{-2-p}(e+fx) dx$	2199
3.251	$\int (a+bx)^m(ac-bcx)^n(ad(m-n)-bd(2+m+n)x) dx$	2205
3.252	$\int (a+bx)(c+dx)^n(e+fx)^{-n} dx$	2211
3.253	$\int (a+bx)(c+dx)^{-1+n}(e+fx)^{-n} dx$	2216
3.254	$\int (a+bx)(c+dx)^{-2+n}(e+fx)^{-n} dx$	2221
3.255	$\int (a+bx)(c+dx)^{-3+n}(e+fx)^{-n} dx$	2227
3.256	$\int (a+bx)(c+dx)^{-4+n}(e+fx)^{-n} dx$	2234
3.257	$\int (a+bx)(c+dx)^{-5+n}(e+fx)^{-n} dx$	2242
3.258	$\int (a+bx)^{-n}(c+dx)(e+fx)^n dx$	2251
3.259	$\int (a+bx)^{-n}(c+dx)(e+fx)^{-1+n} dx$	2256
3.260	$\int (a+bx)^{-n}(c+dx)(e+fx)^{-2+n} dx$	2261
3.261	$\int (a+bx)^{-n}(c+dx)(e+fx)^{-3+n} dx$	2267

3.262	$\int (a + bx)^{-n}(c + dx)(e + fx)^{-4+n} dx$	2274
3.263	$\int (a + bx)^{-n}(c + dx)(e + fx)^{-5+n} dx$	2282
3.264	$\int (a + bx)^{1-n}(c + dx)^{-2+n}(-ad + bc(3 + 2n) + 2bd(1 + n)x) dx$	2291

3.1 $\int (a + bx)(A + Bx)(d + ex)^4 dx$

Optimal result	122
Mathematica [B] (verified)	123
Rubi [A] (verified)	123
Maple [B] (verified)	124
Fricas [B] (verification not implemented)	125
Sympy [B] (verification not implemented)	126
Maxima [B] (verification not implemented)	126
Giac [B] (verification not implemented)	127
Mupad [B] (verification not implemented)	128
Reduce [B] (verification not implemented)	128

Optimal result

Integrand size = 18, antiderivative size = 77

$$\int (a + bx)(A + Bx)(d + ex)^4 dx = \frac{(bd - ae)(Bd - Ae)(d + ex)^5}{5e^3} - \frac{(2bBd - Abe - aBe)(d + ex)^6}{6e^3} + \frac{bB(d + ex)^7}{7e^3}$$

output `1/5*(-a*e+b*d)*(-A*e+B*d)*(e*x+d)^5/e^3-1/6*(-A*b*e-B*a*e+2*B*b*d)*(e*x+d)^6/e^3+1/7*b*B*(e*x+d)^7/e^3`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 172 vs. $2(77) = 154$.

Time = 0.04 (sec) , antiderivative size = 172, normalized size of antiderivative = 2.23

$$\begin{aligned} \int (a + bx)(A + Bx)(d + ex)^4 dx = & aAd^4x + \frac{1}{2}d^3(ABd + aBd + 4aAe)x^2 \\ & + \frac{1}{3}d^2(2ae(2Bd + 3Ae) + bd(Bd + 4Ae))x^3 \\ & + \frac{1}{2}de(ae(3Bd + 2Ae) + bd(2Bd + 3Ae))x^4 \\ & + \frac{1}{5}e^2(ae(4Bd + Ae) + 2bd(3Bd + 2Ae))x^5 \\ & + \frac{1}{6}e^3(4bBd + Abe + aBe)x^6 + \frac{1}{7}bBe^4x^7 \end{aligned}$$

input `Integrate[(a + b*x)*(A + B*x)*(d + e*x)^4, x]`

output `a*A*d^4*x + (d^3*(A*b*d + a*B*d + 4*a*A*e)*x^2)/2 + (d^2*(2*a*e*(2*B*d + 3*A*e) + b*d*(B*d + 4*A*e))*x^3)/3 + (d*e*(a*e*(3*B*d + 2*A*e) + b*d*(2*B*d + 3*A*e))*x^4)/2 + (e^2*(a*e*(4*B*d + A*e) + 2*b*d*(3*B*d + 2*A*e))*x^5)/5 + (e^3*(4*b*B*d + A*b*e + a*B*e)*x^6)/6 + (b*B*e^4*x^7)/7`

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + bx)(A + Bx)(d + ex)^4 dx \\ & \quad \downarrow 86 \\ & \int \left(\frac{(d + ex)^5(aBe + Abe - 2bBd)}{e^2} + \frac{(d + ex)^4(ae - bd)(Ae - Bd)}{e^2} + \frac{bB(d + ex)^6}{e^2} \right) dx \end{aligned}$$

$$\downarrow \text{2009}$$

$$-\frac{(d+ex)^6(-aBe - Abe + 2bBd)}{6e^3} + \frac{(d+ex)^5(bd - ae)(Bd - Ae)}{5e^3} + \frac{bB(d+ex)^7}{7e^3}$$

input `Int[(a + b*x)*(A + B*x)*(d + e*x)^4,x]`

output `((b*d - a*e)*(B*d - A*e)*(d + e*x)^5)/(5*e^3) - ((2*b*B*d - A*b*e - a*B*e) * (d + e*x)^6)/(6*e^3) + (b*B*(d + e*x)^7)/(7*e^3)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.))^(n_.)*((e_.) + (f_.)*(x_.))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 175 vs. 2(71) = 142.

Time = 0.15 (sec) , antiderivative size = 176, normalized size of antiderivative = 2.29

method	result
default	$\frac{bB e^4 x^7}{7} + \frac{((Ab+Ba)e^4+4bBde^3)x^6}{6} + \frac{(Aae^4+4(Ab+Ba)de^3+6bBd^2e^2)x^5}{5} + \frac{(4Aade^3+6(Ab+Ba)d^2e^2+4bBd^3e)}{4}$
norman	$\frac{bB e^4 x^7}{7} + (\frac{1}{6}Ab e^4 + \frac{1}{6}Ba e^4 + \frac{2}{3}bBd e^3) x^6 + (\frac{1}{5}Aa e^4 + \frac{4}{5}Abd e^3 + \frac{4}{5}Bad e^3 + \frac{6}{5}bB d^2 e^2) x^5$
oring	$x(30bB e^4 x^6 + 35Ab e^4 x^5 + 35Ba e^4 x^5 + 140Bbd e^3 x^5 + 42Aa e^4 x^4 + 168Abd e^3 x^4 + 168Bad e^3 x^4 + 252Bb d^2 e^2 x^4 + 210Aad e^3 x^3)$
gospers	$\frac{1}{7}bB e^4 x^7 + \frac{1}{6}x^6 Ab e^4 + \frac{1}{6}x^6 Ba e^4 + \frac{2}{3}x^6 bBd e^3 + \frac{1}{5}x^5 Aa e^4 + \frac{4}{5}x^5 Abd e^3 + \frac{4}{5}x^5 Bad e^3 + \frac{6}{5}x^5 bB d^2 e^2$
risch	$\frac{1}{7}bB e^4 x^7 + \frac{1}{6}x^6 Ab e^4 + \frac{1}{6}x^6 Ba e^4 + \frac{2}{3}x^6 bBd e^3 + \frac{1}{5}x^5 Aa e^4 + \frac{4}{5}x^5 Abd e^3 + \frac{4}{5}x^5 Bad e^3 + \frac{6}{5}x^5 bB d^2 e^2$
parallelrisch	$\frac{1}{7}bB e^4 x^7 + \frac{1}{6}x^6 Ab e^4 + \frac{1}{6}x^6 Ba e^4 + \frac{2}{3}x^6 bBd e^3 + \frac{1}{5}x^5 Aa e^4 + \frac{4}{5}x^5 Abd e^3 + \frac{4}{5}x^5 Bad e^3 + \frac{6}{5}x^5 bB d^2 e^2$

input `int((b*x+a)*(B*x+A)*(e*x+d)^4,x,method=_RETURNVERBOSE)`

output `1/7*b*B*e^4*x^7+1/6*((A*b+B*a)*e^4+4*b*B*d*e^3)*x^6+1/5*(A*a*e^4+4*(A*b+B*a)*d*e^3+6*b*B*d^2*e^2)*x^5+1/4*(4*A*a*d*e^3+6*(A*b+B*a)*d^2*e^2+4*b*B*d^3*e)*x^4+1/3*(6*A*a*d^2*e^2+4*(A*b+B*a)*d^3*e+b*B*d^4)*x^3+1/2*(4*A*a*d^3*e+(A*b+B*a)*d^4)*x^2+A*a*d^4*x`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 175 vs. $2(71) = 142$.

Time = 0.08 (sec) , antiderivative size = 175, normalized size of antiderivative = 2.27

$$\begin{aligned} \int (a + bx)(A + Bx)(d + ex)^4 dx = & \frac{1}{7} Bbe^4x^7 + Aad^4x + \frac{1}{6} (4 Bbde^3 + (Ba + Ab)e^4)x^6 \\ & + \frac{1}{5} (6 Bbd^2e^2 + Aae^4 + 4 (Ba + Ab)de^3)x^5 \\ & + \frac{1}{2} (2 Bbd^3e + 2 Aade^3 + 3 (Ba + Ab)d^2e^2)x^4 \\ & + \frac{1}{3} (Bbd^4 + 6 Aad^2e^2 + 4 (Ba + Ab)d^3e)x^3 \\ & + \frac{1}{2} (4 Aad^3e + (Ba + Ab)d^4)x^2 \end{aligned}$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)^4,x, algorithm="fricas")`

output `1/7*B*b*e^4*x^7 + A*a*d^4*x + 1/6*(4*B*b*d*e^3 + (B*a + A*b)*e^4)*x^6 + 1/5*(6*B*b*d^2*e^2 + A*a*e^4 + 4*(B*a + A*b)*d*e^3)*x^5 + 1/2*(2*B*b*d^3*e + 2*A*a*d*e^3 + 3*(B*a + A*b)*d^2*e^2)*x^4 + 1/3*(B*b*d^4 + 6*A*a*d^2*e^2 + 4*(B*a + A*b)*d^3*e)*x^3 + 1/2*(4*A*a*d^3*e + (B*a + A*b)*d^4)*x^2`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 226 vs. $2(71) = 142$.

Time = 0.03 (sec) , antiderivative size = 226, normalized size of antiderivative = 2.94

$$\int (a + bx)(A + Bx)(d + ex)^4 dx = Aad^4x + \frac{Bbe^4x^7}{7} + x^6 \left(\frac{Abe^4}{6} + \frac{Bae^4}{6} + \frac{2Bbde^3}{3} \right) \\ + x^5 \left(\frac{Aae^4}{5} + \frac{4Abde^3}{5} + \frac{4Bade^3}{5} + \frac{6Bbd^2e^2}{5} \right) \\ + x^4 \left(Aade^3 + \frac{3Abd^2e^2}{2} + \frac{3Bad^2e^2}{2} + Bbd^3e \right) \\ + x^3 \cdot \left(2Aad^2e^2 + \frac{4Abd^3e}{3} + \frac{4Bad^3e}{3} + \frac{Bbd^4}{3} \right) \\ + x^2 \cdot \left(2Aad^3e + \frac{Abd^4}{2} + \frac{Bad^4}{2} \right)$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)**4,x)`

output `A*a*d**4*x + B*b*e**4*x**7/7 + x**6*(A*b*e**4/6 + B*a*e**4/6 + 2*B*b*d*e**3/3) + x**5*(A*a*e**4/5 + 4*A*b*d*e**3/5 + 4*B*a*d*e**3/5 + 6*B*b*d**2*e**2/5) + x**4*(A*a*d*e**3 + 3*A*b*d**2*e**2/2 + 3*B*a*d**2*e**2/2 + B*b*d**3*e) + x**3*(2*A*a*d**2*e**2 + 4*A*b*d**3*e/3 + 4*B*a*d**3*e/3 + B*b*d**4/3) + x**2*(2*A*a*d**3*e + A*b*d**4/2 + B*a*d**4/2)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 175 vs. $2(71) = 142$.

Time = 0.04 (sec) , antiderivative size = 175, normalized size of antiderivative = 2.27

$$\int (a + bx)(A + Bx)(d + ex)^4 dx = \frac{1}{7} Bbe^4x^7 + Aad^4x + \frac{1}{6} (4 Bbde^3 + (Ba + Ab)e^4)x^6 \\ + \frac{1}{5} (6 Bbd^2e^2 + Aae^4 + 4 (Ba + Ab)de^3)x^5 \\ + \frac{1}{2} (2 Bbd^3e + 2 Aade^3 + 3 (Ba + Ab)d^2e^2)x^4 \\ + \frac{1}{3} (Bbd^4 + 6 Aad^2e^2 + 4 (Ba + Ab)d^3e)x^3 \\ + \frac{1}{2} (4 Aad^3e + (Ba + Ab)d^4)x^2$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)^4,x, algorithm="maxima")`

output
$$\begin{aligned} & 1/7*B*b*e^4*x^7 + A*a*d^4*x + 1/6*(4*B*b*d*e^3 + (B*a + A*b)*e^4)*x^6 + 1/ \\ & 5*(6*B*b*d^2*e^2 + A*a*e^4 + 4*(B*a + A*b)*d*e^3)*x^5 + 1/2*(2*B*b*d^3*e + \\ & 2*A*a*d*e^3 + 3*(B*a + A*b)*d^2*e^2)*x^4 + 1/3*(B*b*d^4 + 6*A*a*d^2*e^2 + \\ & 4*(B*a + A*b)*d^3*e)*x^3 + 1/2*(4*A*a*d^3*e + (B*a + A*b)*d^4)*x^2 \end{aligned}$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 216 vs. $2(71) = 142$.

Time = 0.12 (sec) , antiderivative size = 216, normalized size of antiderivative = 2.81

$$\begin{aligned} \int (a + bx)(A + Bx)(d + ex)^4 dx = & \frac{1}{7} Bbe^4x^7 + \frac{2}{3} Bbde^3x^6 + \frac{1}{6} Bae^4x^6 + \frac{1}{6} Abe^4x^6 \\ & + \frac{6}{5} Bbd^2e^2x^5 + \frac{4}{5} Bade^3x^5 + \frac{4}{5} Abde^3x^5 + \frac{1}{5} Aae^4x^5 \\ & + Bbd^3ex^4 + \frac{3}{2} Bad^2e^2x^4 + \frac{3}{2} Abd^2e^2x^4 + Aade^3x^4 \\ & + \frac{1}{3} Bbd^4x^3 + \frac{4}{3} Bad^3ex^3 + \frac{4}{3} Abd^3ex^3 + 2Aad^2e^2x^3 \\ & + \frac{1}{2} Bad^4x^2 + \frac{1}{2} Abd^4x^2 + 2Aad^3ex^2 + Aad^4x \end{aligned}$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)^4,x, algorithm="giac")`

output
$$\begin{aligned} & 1/7*B*b*e^4*x^7 + 2/3*B*b*d*e^3*x^6 + 1/6*B*a*e^4*x^6 + 1/6*A*b*e^4*x^6 + \\ & 6/5*B*b*d^2*e^2*x^5 + 4/5*B*a*d*e^3*x^5 + 4/5*A*b*d*e^3*x^5 + 1/5*A*a*e^4* \\ & x^5 + B*b*d^3*e*x^4 + 3/2*B*a*d^2*e^2*x^4 + 3/2*A*b*d^2*e^2*x^4 + A*a*d*e^ \\ & 3*x^4 + 1/3*B*b*d^4*x^3 + 4/3*B*a*d^3*e*x^3 + 4/3*A*b*d^3*e*x^3 + 2*A*a*d^ \\ & 2*e^2*x^3 + 1/2*B*a*d^4*x^2 + 1/2*A*b*d^4*x^2 + 2*A*a*d^3*e*x^2 + A*a*d^4* \\ & x \end{aligned}$$

output

```
(x*(105*a**2*d**4 + 210*a**2*d**3*e*x + 210*a**2*d**2*e**2*x**2 + 105*a**2
*d*e**3*x**3 + 21*a**2*e**4*x**4 + 105*a*b*d**4*x + 280*a*b*d**3*e*x**2 +
315*a*b*d**2*e**2*x**3 + 168*a*b*d*e**3*x**4 + 35*a*b*e**4*x**5 + 35*b**2*
d**4*x**2 + 105*b**2*d**3*e*x**3 + 126*b**2*d**2*e**2*x**4 + 70*b**2*d*e**
3*x**5 + 15*b**2*e**4*x**6))/105
```

3.2 $\int (a + bx)(A + Bx)(d + ex)^3 dx$

Optimal result	130
Mathematica [A] (verified)	130
Rubi [A] (verified)	131
Maple [A] (verified)	132
Fricas [A] (verification not implemented)	133
Sympy [B] (verification not implemented)	133
Maxima [A] (verification not implemented)	134
Giac [B] (verification not implemented)	134
Mupad [B] (verification not implemented)	135
Reduce [B] (verification not implemented)	136

Optimal result

Integrand size = 18, antiderivative size = 77

$$\int (a + bx)(A + Bx)(d + ex)^3 dx = \frac{(bd - ae)(Bd - Ae)(d + ex)^4}{4e^3} - \frac{(2bBd - Abe - aBe)(d + ex)^5}{5e^3} + \frac{bB(d + ex)^6}{6e^3}$$

output

```
1/4*(-a*e+b*d)*(-A*e+B*d)*(e*x+d)^4/e^3-1/5*(-A*b*e-B*a*e+2*B*b*d)*(e*x+d)^5/e^3+1/6*b*B*(e*x+d)^6/e^3
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.69

$$\begin{aligned} \int (a + bx)(A + Bx)(d + ex)^3 dx &= aAd^3x + \frac{1}{2}d^2(Abd + aBd + 3aAe)x^2 \\ &+ \frac{1}{3}d(3ae(Bd + Ae) + bd(Bd + 3Ae))x^3 \\ &+ \frac{1}{4}e(3bd(Bd + Ae) + ae(3Bd + Ae))x^4 \\ &+ \frac{1}{5}e^2(3bBd + Abe + aBe)x^5 + \frac{1}{6}bBe^3x^6 \end{aligned}$$

input `Integrate[(a + b*x)*(A + B*x)*(d + e*x)^3,x]`

output `a*A*d^3*x + (d^2*(A*b*d + a*B*d + 3*a*A*e)*x^2)/2 + (d*(3*a*e*(B*d + A*e) + b*d*(B*d + 3*A*e))*x^3)/3 + (e*(3*b*d*(B*d + A*e) + a*e*(3*B*d + A*e))*x^4)/4 + (e^2*(3*b*B*d + A*b*e + a*B*e)*x^5)/5 + (b*B*e^3*x^6)/6`

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)(A + Bx)(d + ex)^3 dx$$

$$\downarrow 86$$

$$\int \left(\frac{(d + ex)^4(aBe + Abe - 2bBd)}{e^2} + \frac{(d + ex)^3(ae - bd)(Ae - Bd)}{e^2} + \frac{bB(d + ex)^5}{e^2} \right) dx$$

$$\downarrow 2009$$

$$-\frac{(d + ex)^5(-aBe - Abe + 2bBd)}{5e^3} + \frac{(d + ex)^4(bd - ae)(Bd - Ae)}{4e^3} + \frac{bB(d + ex)^6}{6e^3}$$

input `Int[(a + b*x)*(A + B*x)*(d + e*x)^3,x]`

output `((b*d - a*e)*(B*d - A*e)*(d + e*x)^4)/(4*e^3) - ((2*b*B*d - A*b*e - a*B*e)*(d + e*x)^5)/(5*e^3) + (b*B*(d + e*x)^6)/(6*e^3)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.74

$$\int (a + bx)(A + Bx)(d + ex)^3 dx = \frac{1}{6} Bbe^3 x^6 + Aad^3 x + \frac{1}{5} (3 Bbde^2 + (Ba + Ab)e^3) x^5$$

$$+ \frac{1}{4} (3 Bbd^2 e + Aae^3 + 3 (Ba + Ab)de^2) x^4$$

$$+ \frac{1}{3} (Bbd^3 + 3 Aade^2 + 3 (Ba + Ab)d^2 e) x^3$$

$$+ \frac{1}{2} (3 Aad^2 e + (Ba + Ab)d^3) x^2$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)^3,x, algorithm="fricas")`

output `1/6*B*b*e^3*x^6 + A*a*d^3*x + 1/5*(3*B*b*d*e^2 + (B*a + A*b)*e^3)*x^5 + 1/4*(3*B*b*d^2*e + A*a*e^3 + 3*(B*a + A*b)*d*e^2)*x^4 + 1/3*(B*b*d^3 + 3*A*a*d*e^2 + 3*(B*a + A*b)*d^2*e)*x^3 + 1/2*(3*A*a*d^2*e + (B*a + A*b)*d^3)*x^2`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 168 vs. 2(73) = 146.

Time = 0.03 (sec) , antiderivative size = 168, normalized size of antiderivative = 2.18

$$\int (a + bx)(A + Bx)(d + ex)^3 dx = Aad^3 x + \frac{Bbe^3 x^6}{6} + x^5 \left(\frac{Abe^3}{5} + \frac{Bae^3}{5} + \frac{3Bbde^2}{5} \right)$$

$$+ x^4 \left(\frac{Aae^3}{4} + \frac{3Abde^2}{4} + \frac{3Bade^2}{4} + \frac{3Bbd^2 e}{4} \right)$$

$$+ x^3 \left(Aade^2 + Abd^2 e + Bad^2 e + \frac{Bbd^3}{3} \right)$$

$$+ x^2 \cdot \left(\frac{3Aad^2 e}{2} + \frac{Abd^3}{2} + \frac{Bad^3}{2} \right)$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)**3,x)`

output

```
A*a*d**3*x + B*b*e**3*x**6/6 + x**5*(A*b*e**3/5 + B*a*e**3/5 + 3*B*b*d*e**2/5) + x**4*(A*a*e**3/4 + 3*A*b*d*e**2/4 + 3*B*a*d*e**2/4 + 3*B*b*d**2*e/4) + x**3*(A*a*d*e**2 + A*b*d**2*e + B*a*d**2*e + B*b*d**3/3) + x**2*(3*A*a*d**2*e/2 + A*b*d**3/2 + B*a*d**3/2)
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.74

$$\int (a + bx)(A + Bx)(d + ex)^3 dx = \frac{1}{6} Bbe^3 x^6 + Aad^3 x + \frac{1}{5} (3 Bbde^2 + (Ba + Ab)e^3) x^5 + \frac{1}{4} (3 Bbd^2 e + Aae^3 + 3 (Ba + Ab)de^2) x^4 + \frac{1}{3} (Bbd^3 + 3 Aade^2 + 3 (Ba + Ab)d^2 e) x^3 + \frac{1}{2} (3 Aad^2 e + (Ba + Ab)d^3) x^2$$

input

```
integrate((b*x+a)*(B*x+A)*(e*x+d)^3,x, algorithm="maxima")
```

output

```
1/6*B*b*e^3*x^6 + A*a*d^3*x + 1/5*(3*B*b*d*e^2 + (B*a + A*b)*e^3)*x^5 + 1/4*(3*B*b*d^2*e + A*a*e^3 + 3*(B*a + A*b)*d*e^2)*x^4 + 1/3*(B*b*d^3 + 3*A*a*d*e^2 + 3*(B*a + A*b)*d^2*e)*x^3 + 1/2*(3*A*a*d^2*e + (B*a + A*b)*d^3)*x^2
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 163 vs. 2(71) = 142.

Time = 0.12 (sec) , antiderivative size = 163, normalized size of antiderivative = 2.12

$$\int (a + bx)(A + Bx)(d + ex)^3 dx = \frac{1}{6} Bbe^3 x^6 + \frac{3}{5} Bbde^2 x^5 + \frac{1}{5} Bae^3 x^5 + \frac{1}{5} Abe^3 x^5 + \frac{3}{4} Bbd^2 ex^4 + \frac{3}{4} Bade^2 x^4 + \frac{3}{4} Abde^2 x^4 + \frac{1}{4} Aae^3 x^4 + \frac{1}{3} Bbd^3 x^3 + Bad^2 ex^3 + Abd^2 ex^3 + Aade^2 x^3 + \frac{1}{2} Bad^3 x^2 + \frac{1}{2} Abd^3 x^2 + \frac{3}{2} Aad^2 ex^2 + Aad^3 x$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)^3,x, algorithm="giac")`

output `1/6*B*b*e^3*x^6 + 3/5*B*b*d*e^2*x^5 + 1/5*B*a*e^3*x^5 + 1/5*A*b*e^3*x^5 + 3/4*B*b*d^2*e*x^4 + 3/4*B*a*d*e^2*x^4 + 3/4*A*b*d*e^2*x^4 + 1/4*A*a*e^3*x^4 + 1/3*B*b*d^3*x^3 + B*a*d^2*e*x^3 + A*b*d^2*e*x^3 + A*a*d*e^2*x^3 + 1/2*B*a*d^3*x^2 + 1/2*A*b*d^3*x^2 + 3/2*A*a*d^2*e*x^2 + A*a*d^3*x`

Mupad [B] (verification not implemented)

Time = 0.89 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.83

$$\int (a + bx)(A + Bx)(d + ex)^3 dx = x^2 \left(\frac{Abd^3}{2} + \frac{Bad^3}{2} + \frac{3Aad^2e}{2} \right) + x^5 \left(\frac{Ab e^3}{5} + \frac{Ba e^3}{5} + \frac{3Bbd e^2}{5} \right) + x^3 \left(\frac{Bbd^3}{3} + Aade^2 + Abd^2e + Bad^2e \right) + x^4 \left(\frac{Aae^3}{4} + \frac{3Abde^2}{4} + \frac{3Bade^2}{4} + \frac{3Bbd^2e}{4} \right) + Aad^3x + \frac{Bbe^3x^6}{6}$$

input `int((A + B*x)*(a + b*x)*(d + e*x)^3,x)`

output `x^2*((A*b*d^3)/2 + (B*a*d^3)/2 + (3*A*a*d^2*e)/2) + x^5*((A*b*e^3)/5 + (B*a*e^3)/5 + (3*B*b*d*e^2)/5) + x^3*((B*b*d^3)/3 + A*a*d*e^2 + A*b*d^2*e + B*a*d^2*e) + x^4*((A*a*e^3)/4 + (3*A*b*d*e^2)/4 + (3*B*a*d*e^2)/4 + (3*B*b*d^2*e)/4) + A*a*d^3*x + (B*b*e^3*x^6)/6`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.70

$$\int (a + bx)(A + Bx)(d + ex)^3 dx$$

$$= \frac{x(10b^2e^3x^5 + 24abe^3x^4 + 36b^2de^2x^4 + 15a^2e^3x^3 + 90abd e^2x^3 + 45b^2d^2e x^3 + 60a^2d e^2x^2 + 120abd^2e x^2}{60}$$

input `int((b*x+a)*(B*x+A)*(e*x+d)^3,x)`output `(x*(60*a**2*d**3 + 90*a**2*d**2*e*x + 60*a**2*d*e**2*x**2 + 15*a**2*e**3*x**3 + 60*a*b*d**3*x + 120*a*b*d**2*e*x**2 + 90*a*b*d*e**2*x**3 + 24*a*b*e**3*x**4 + 20*b**2*d**3*x**2 + 45*b**2*d**2*e*x**3 + 36*b**2*d*e**2*x**4 + 10*b**2*e**3*x**5))/60`

3.3 $\int (a + bx)(A + Bx)(d + ex)^2 dx$

Optimal result	137
Mathematica [A] (verified)	137
Rubi [A] (verified)	138
Maple [A] (verified)	139
Fricas [A] (verification not implemented)	140
Sympy [A] (verification not implemented)	140
Maxima [A] (verification not implemented)	141
Giac [A] (verification not implemented)	141
Mupad [B] (verification not implemented)	142
Reduce [B] (verification not implemented)	142

Optimal result

Integrand size = 18, antiderivative size = 77

$$\int (a + bx)(A + Bx)(d + ex)^2 dx = \frac{(bd - ae)(Bd - Ae)(d + ex)^3}{3e^3} - \frac{(2bBd - Abe - aBe)(d + ex)^4}{4e^3} + \frac{bB(d + ex)^5}{5e^3}$$

output

```
1/3*(-a*e+b*d)*(-A*e+B*d)*(e*x+d)^3/e^3-1/4*(-A*b*e-B*a*e+2*B*b*d)*(e*x+d)^4/e^3+1/5*b*B*(e*x+d)^5/e^3
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.25

$$\int (a + bx)(A + Bx)(d + ex)^2 dx = aAd^2x + \frac{1}{2}d(Abd + aBd + 2aAe)x^2 + \frac{1}{3}(bBd^2 + 2Abde + 2aBde + aAe^2)x^3 + \frac{1}{4}e(2bBd + Abe + aBe)x^4 + \frac{1}{5}bBe^2x^5$$

input

```
Integrate[(a + b*x)*(A + B*x)*(d + e*x)^2,x]
```

output

$$aAd^2x + (d(Abd + aBd + 2aAe)x^2)/2 + ((bBd^2 + 2Abde + 2aBde + aAe^2)x^3)/3 + (e(2bBd + Abe + aBe)x^4)/4 + (bBe^2x^5)/5$$

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)(A + Bx)(d + ex)^2 dx$$

$$\downarrow 86$$

$$\int \left(\frac{(d + ex)^3(aBe + Abe - 2bBd)}{e^2} + \frac{(d + ex)^2(ae - bd)(Ae - Bd)}{e^2} + \frac{bB(d + ex)^4}{e^2} \right) dx$$

$$\downarrow 2009$$

$$-\frac{(d + ex)^4(-aBe - Abe + 2bBd)}{4e^3} + \frac{(d + ex)^3(bd - ae)(Bd - Ae)}{3e^3} + \frac{bB(d + ex)^5}{5e^3}$$

input

```
Int[(a + b*x)*(A + B*x)*(d + e*x)^2,x]
```

output

$$\frac{(bd - ae)(Bd - Ae)(d + ex)^3}{3e^3} - \frac{(2bBd - Abe - aBe)(d + ex)^4}{4e^3} + \frac{bB(d + ex)^5}{5e^3}$$

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0]) || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.22

method	result
default	$\frac{bB e^2 x^5}{5} + \frac{((Ab+Ba)e^2+2bBde)x^4}{4} + \frac{(Aa e^2+2(Ab+Ba)de+bB d^2)x^3}{3} + \frac{(2Aade+(Ab+Ba)d^2)x^2}{2} + Aa d^2 x$
norman	$\frac{bB e^2 x^5}{5} + \left(\frac{1}{4}Ab e^2 + \frac{1}{4}Ba e^2 + \frac{1}{2}bBde\right) x^4 + \left(\frac{1}{3}Aa e^2 + \frac{2}{3}Abde + \frac{2}{3}Bade + \frac{1}{3}bB d^2\right) x^3 + (Aa d^2 x$
orering	$\frac{x(12bB e^2 x^4+15Ab e^2 x^3+15Ba e^2 x^3+30Bbde x^3+20Aa e^2 x^2+40Abde x^2+40Bade x^2+20Bb d^2 x^2+60Aadex+30Ab d^2 x+30Aad^2 x)}{60}$
gosper	$\frac{1}{5}bB e^2 x^5 + \frac{1}{4}x^4 Ab e^2 + \frac{1}{4}x^4 Ba e^2 + \frac{1}{2}x^4 bBde + \frac{1}{3}x^3 Aa e^2 + \frac{2}{3}x^3 Abde + \frac{2}{3}x^3 Bade + \frac{1}{3}x^3 bB d^2$
risch	$\frac{1}{5}bB e^2 x^5 + \frac{1}{4}x^4 Ab e^2 + \frac{1}{4}x^4 Ba e^2 + \frac{1}{2}x^4 bBde + \frac{1}{3}x^3 Aa e^2 + \frac{2}{3}x^3 Abde + \frac{2}{3}x^3 Bade + \frac{1}{3}x^3 bB d^2$
parallelrisch	$\frac{1}{5}bB e^2 x^5 + \frac{1}{4}x^4 Ab e^2 + \frac{1}{4}x^4 Ba e^2 + \frac{1}{2}x^4 bBde + \frac{1}{3}x^3 Aa e^2 + \frac{2}{3}x^3 Abde + \frac{2}{3}x^3 Bade + \frac{1}{3}x^3 bB d^2$

input

```
int((b*x+a)*(B*x+A)*(e*x+d)^2,x,method=_RETURNVERBOSE)
```

output

```
1/5*b*B*e^2*x^5+1/4*((A*b+B*a)*e^2+2*b*B*d*e)*x^4+1/3*(A*a*e^2+2*(A*b+B*a)*d*e+b*B*d^2)*x^3+1/2*(2*A*a*d*e+(A*b+B*a)*d^2)*x^2+A*a*d^2*x
```


Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.21

$$\int (a + bx)(A + Bx)(d + ex)^2 dx = \frac{1}{5} Bbe^2x^5 + Aad^2x + \frac{1}{4} (2 Bbde + (Ba + Ab)e^2)x^4 \\ + \frac{1}{3} (Bbd^2 + Aae^2 + 2(Ba + Ab)de)x^3 \\ + \frac{1}{2} (2 Aade + (Ba + Ab)d^2)x^2$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)^2,x, algorithm="fricas")`

output `1/5*B*b*e^2*x^5 + A*a*d^2*x + 1/4*(2*B*b*d*e + (B*a + A*b)*e^2)*x^4 + 1/3*(B*b*d^2 + A*a*e^2 + 2*(B*a + A*b)*d*e)*x^3 + 1/2*(2*A*a*d*e + (B*a + A*b)*d^2)*x^2`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.51

$$\int (a + bx)(A + Bx)(d + ex)^2 dx = Aad^2x + \frac{Bbe^2x^5}{5} + x^4 \left(\frac{Abe^2}{4} + \frac{Bae^2}{4} + \frac{Bbde}{2} \right) \\ + x^3 \left(\frac{Aae^2}{3} + \frac{2Abde}{3} + \frac{2Bade}{3} + \frac{Bbd^2}{3} \right) \\ + x^2 \left(Aade + \frac{Abd^2}{2} + \frac{Bad^2}{2} \right)$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)**2,x)`

output `A*a*d**2*x + B*b*e**2*x**5/5 + x**4*(A*b*e**2/4 + B*a*e**2/4 + B*b*d*e/2) + x**3*(A*a*e**2/3 + 2*A*b*d*e/3 + 2*B*a*d*e/3 + B*b*d**2/3) + x**2*(A*a*d*e + A*b*d**2/2 + B*a*d**2/2)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.21

$$\int (a + bx)(A + Bx)(d + ex)^2 dx = \frac{1}{5} Bbe^2x^5 + Aad^2x + \frac{1}{4} (2Bbde + (Ba + Ab)e^2)x^4$$

$$+ \frac{1}{3} (Bbd^2 + Aae^2 + 2(Ba + Ab)de)x^3$$

$$+ \frac{1}{2} (2Aade + (Ba + Ab)d^2)x^2$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)^2,x, algorithm="maxima")`output `1/5*B*b*e^2*x^5 + A*a*d^2*x + 1/4*(2*B*b*d*e + (B*a + A*b)*e^2)*x^4 + 1/3*(B*b*d^2 + A*a*e^2 + 2*(B*a + A*b)*d*e)*x^3 + 1/2*(2*A*a*d*e + (B*a + A*b)*d^2)*x^2`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.47

$$\int (a + bx)(A + Bx)(d + ex)^2 dx = \frac{1}{5} Bbe^2x^5 + \frac{1}{2} Bbdex^4 + \frac{1}{4} Bae^2x^4 + \frac{1}{4} Abe^2x^4$$

$$+ \frac{1}{3} Bbd^2x^3 + \frac{2}{3} Badex^3 + \frac{2}{3} Abdex^3 + \frac{1}{3} Aae^2x^3$$

$$+ \frac{1}{2} Bad^2x^2 + \frac{1}{2} Abd^2x^2 + Aadex^2 + Aad^2x$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)^2,x, algorithm="giac")`output `1/5*B*b*e^2*x^5 + 1/2*B*b*d*e*x^4 + 1/4*B*a*e^2*x^4 + 1/4*A*b*e^2*x^4 + 1/3*B*b*d^2*x^3 + 2/3*B*a*d*e*x^3 + 2/3*A*b*d*e*x^3 + 1/3*A*a*e^2*x^3 + 1/2*B*a*d^2*x^2 + 1/2*A*b*d^2*x^2 + A*a*d*e*x^2 + A*a*d^2*x`

Mupad [B] (verification not implemented)

Time = 0.87 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.27

$$\int (a + bx)(A + Bx)(d + ex)^2 dx = x^3 \left(\frac{Aae^2}{3} + \frac{Bbd^2}{3} + \frac{2Abde}{3} + \frac{2Bade}{3} \right) \\ + x^2 \left(\frac{Abd^2}{2} + \frac{Bad^2}{2} + Aade \right) \\ + x^4 \left(\frac{Abe^2}{4} + \frac{Bae^2}{4} + \frac{Bbde}{2} \right) \\ + Aad^2x + \frac{Bbe^2x^5}{5}$$

input `int((A + B*x)*(a + b*x)*(d + e*x)^2,x)`output `x^3*((A*a*e^2)/3 + (B*b*d^2)/3 + (2*A*b*d*e)/3 + (2*B*a*d*e)/3) + x^2*((A*b*d^2)/2 + (B*a*d^2)/2 + A*a*d*e) + x^4*((A*b*e^2)/4 + (B*a*e^2)/4 + (B*b*d*e)/2) + A*a*d^2*x + (B*b*e^2*x^5)/5`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.17

$$\int (a + bx)(A + Bx)(d + ex)^2 dx \\ = \frac{x(6b^2e^2x^4 + 15ab^2e^2x^3 + 15b^2dex^3 + 10a^2e^2x^2 + 40abdex^2 + 10b^2d^2x^2 + 30a^2dex + 30abd^2x + 30a^2d^2)}{30}$$

input `int((b*x+a)*(B*x+A)*(e*x+d)^2,x)`output `(x*(30*a**2*d**2 + 30*a**2*d*e*x + 10*a**2*e**2*x**2 + 30*a*b*d**2*x + 40*a*b*d*e*x**2 + 15*a*b*e**2*x**3 + 10*b**2*d**2*x**2 + 15*b**2*d*e*x**3 + 6*b**2*e**2*x**4))/30`

3.4 $\int (a + bx)(A + Bx)(d + ex) dx$

Optimal result	143
Mathematica [A] (verified)	143
Rubi [A] (verified)	144
Maple [A] (verified)	145
Fricas [A] (verification not implemented)	145
Sympy [A] (verification not implemented)	146
Maxima [A] (verification not implemented)	146
Giac [A] (verification not implemented)	147
Mupad [B] (verification not implemented)	147
Reduce [B] (verification not implemented)	148

Optimal result

Integrand size = 16, antiderivative size = 56

$$\int (a + bx)(A + Bx)(d + ex) dx = aAdx + \frac{1}{2}(Abd + aBd + aAe)x^2 + \frac{1}{3}(bBd + Abe + aBe)x^3 + \frac{1}{4}bBex^4$$

output

```
a*A*d*x+1/2*(A*a*e+A*b*d+B*a*d)*x^2+1/3*(A*b*e+B*a*e+B*b*d)*x^3+1/4*b*B*e*x^4
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.95

$$\int (a + bx)(A + Bx)(d + ex) dx = \frac{1}{12}x(12aAd + 6(Abd + aBd + aAe)x + 4(bBd + Abe + aBe)x^2 + 3bBex^3)$$

input

```
Integrate[(a + b*x)*(A + B*x)*(d + e*x),x]
```

output

```
(x*(12*a*A*d + 6*(A*b*d + a*B*d + a*A*e)*x + 4*(b*B*d + A*b*e + a*B*e)*x^2 + 3*b*B*e*x^3))/12
```

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)(A + Bx)(d + ex) dx$$

↓ 86

$$\int (x^2(aBe + Abe + bBd) + x(aAe + aBd + Abd) + aAd + bBex^3) dx$$

↓ 2009

$$\frac{1}{3}x^3(aBe + Abe + bBd) + \frac{1}{2}x^2(aAe + aBd + Abd) + aAdx + \frac{1}{4}bBex^4$$

input `Int[(a + b*x)*(A + B*x)*(d + e*x),x]`

output `a*A*d*x + ((A*b*d + a*B*d + a*A*e)*x^2)/2 + ((b*B*d + A*b*e + a*B*e)*x^3)/3 + (b*B*e*x^4)/4`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.95

method	result	size
default	$\frac{bBe x^4}{4} + \frac{((Ab+Ba)e+Bbd)x^3}{3} + \frac{(Aae+(Ab+Ba)d)x^2}{2} + aAdx$	53
norman	$\frac{bBe x^4}{4} + (\frac{1}{3}Abe + \frac{1}{3}Bae + \frac{1}{3}Bbd) x^3 + (\frac{1}{2}Aae + \frac{1}{2}Abd + \frac{1}{2}Bad) x^2 + aAdx$	55
orering	$\frac{x(3Bbe x^3 + 4Abe x^2 + 4Bae x^2 + 4Bbd x^2 + 6Aae x + 6Abd x + 6Bad x + 12Aad)}{12}$	60
gosper	$\frac{1}{4}bBe x^4 + \frac{1}{3}x^3 Abe + \frac{1}{3}x^3 Bae + \frac{1}{3}x^3 Bbd + \frac{1}{2}x^2 Aae + \frac{1}{2}x^2 Abd + \frac{1}{2}x^2 Bad + aAdx$	63
risch	$\frac{1}{4}bBe x^4 + \frac{1}{3}x^3 Abe + \frac{1}{3}x^3 Bae + \frac{1}{3}x^3 Bbd + \frac{1}{2}x^2 Aae + \frac{1}{2}x^2 Abd + \frac{1}{2}x^2 Bad + aAdx$	63
parallelrisch	$\frac{1}{4}bBe x^4 + \frac{1}{3}x^3 Abe + \frac{1}{3}x^3 Bae + \frac{1}{3}x^3 Bbd + \frac{1}{2}x^2 Aae + \frac{1}{2}x^2 Abd + \frac{1}{2}x^2 Bad + aAdx$	63

input `int((b*x+a)*(B*x+A)*(e*x+d),x,method=_RETURNVERBOSE)`output `1/4*b*B*e*x^4+1/3*((A*b+B*a)*e+B*b*d)*x^3+1/2*(A*a*e+(A*b+B*a)*d)*x^2+a*A*d*x`**Fricas [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.11

$$\int (a + bx)(A + Bx)(d + ex) dx = \frac{1}{4}x^4 ebB + \frac{1}{3}x^3 dbB + \frac{1}{3}x^3 eaB + \frac{1}{3}x^3 ebA + \frac{1}{2}x^2 daB + \frac{1}{2}x^2 dbA + \frac{1}{2}x^2 eaA + xdaA$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d),x, algorithm="fricas")`output `1/4*x^4*e*b*B + 1/3*x^3*d*b*B + 1/3*x^3*e*a*B + 1/3*x^3*e*b*A + 1/2*x^2*d*a*B + 1/2*x^2*d*b*A + 1/2*x^2*e*a*A + x*d*a*A`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.12

$$\int (a + bx)(A + Bx)(d + ex) dx = Aadx + \frac{Bbex^4}{4} + x^3 \left(\frac{Abe}{3} + \frac{Bae}{3} + \frac{Bbd}{3} \right) + x^2 \left(\frac{Aae}{2} + \frac{Abd}{2} + \frac{Bad}{2} \right)$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d),x)`output `A*a*d*x + B*b*e*x**4/4 + x**3*(A*b*e/3 + B*a*e/3 + B*b*d/3) + x**2*(A*a*e/2 + A*b*d/2 + B*a*d/2)`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.93

$$\int (a + bx)(A + Bx)(d + ex) dx = \frac{1}{4} Bbex^4 + Aadx + \frac{1}{3} (Bbd + (Ba + Ab)e)x^3 + \frac{1}{2} (Aae + (Ba + Ab)d)x^2$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d),x, algorithm="maxima")`output `1/4*B*b*e*x^4 + A*a*d*x + 1/3*(B*b*d + (B*a + A*b)*e)*x^3 + 1/2*(A*a*e + (B*a + A*b)*d)*x^2`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.11

$$\int (a + bx)(A + Bx)(d + ex) dx = \frac{1}{4} B b e x^4 + \frac{1}{3} B b d x^3 + \frac{1}{3} B a e x^3 + \frac{1}{3} A b e x^3 + \frac{1}{2} B a d x^2 + \frac{1}{2} A b d x^2 + \frac{1}{2} A a e x^2 + A a d x$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d),x, algorithm="giac")`

output `1/4*B*b*e*x^4 + 1/3*B*b*d*x^3 + 1/3*B*a*e*x^3 + 1/3*A*b*e*x^3 + 1/2*B*a*d*x^2 + 1/2*A*b*d*x^2 + 1/2*A*a*e*x^2 + A*a*d*x`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.96

$$\int (a + bx)(A + Bx)(d + ex) dx = \frac{B b e x^4}{4} + \left(\frac{A b e}{3} + \frac{B a e}{3} + \frac{B b d}{3} \right) x^3 + \left(\frac{A a e}{2} + \frac{A b d}{2} + \frac{B a d}{2} \right) x^2 + A a d x$$

input `int((A + B*x)*(a + b*x)*(d + e*x),x)`

output `x^2*((A*a*e)/2 + (A*b*d)/2 + (B*a*d)/2) + x^3*((A*b*e)/3 + (B*a*e)/3 + (B*b*d)/3) + (B*b*e*x^4)/4 + A*a*d*x`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.88

$$\int (a + bx)(A + Bx)(d + ex) dx$$
$$= \frac{x(3b^2e x^3 + 8abe x^2 + 4b^2d x^2 + 6a^2ex + 12abdx + 12a^2d)}{12}$$

input `int((b*x+a)*(B*x+A)*(e*x+d),x)`

output `(x*(12*a**2*d + 6*a**2*e*x + 12*a*b*d*x + 8*a*b*e*x**2 + 4*b**2*d*x**2 + 3*b**2*e*x**3))/12`

3.5 $\int (a + bx)(A + Bx) dx$

Optimal result	149
Mathematica [A] (verified)	149
Rubi [A] (verified)	150
Maple [A] (verified)	151
Fricas [A] (verification not implemented)	151
Sympy [A] (verification not implemented)	152
Maxima [A] (verification not implemented)	152
Giac [A] (verification not implemented)	152
Mupad [B] (verification not implemented)	153
Reduce [B] (verification not implemented)	153

Optimal result

Integrand size = 11, antiderivative size = 28

$$\int (a + bx)(A + Bx) dx = aAx + \frac{1}{2}(Ab + aB)x^2 + \frac{1}{3}bBx^3$$

output `a*A*x+1/2*(A*b+B*a)*x^2+1/3*b*B*x^3`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int (a + bx)(A + Bx) dx = aAx + \frac{1}{2}(Ab + aB)x^2 + \frac{1}{3}bBx^3$$

input `Integrate[(a + b*x)*(A + B*x),x]`

output `a*A*x + ((A*b + a*B)*x^2)/2 + (b*B*x^3)/3`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)(A + Bx) dx$$

$$\downarrow 49$$

$$\int (x(aB + Ab) + aA + bBx^2) dx$$

$$\downarrow 2009$$

$$\frac{1}{2}x^2(aB + Ab) + aAx + \frac{1}{3}bBx^3$$

input `Int[(a + b*x)*(A + B*x),x]`

output `a*A*x + ((A*b + a*B)*x^2)/2 + (b*B*x^3)/3`

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.89

method	result	size
default	$aAx + \frac{(Ab+Ba)x^2}{2} + \frac{bBx^3}{3}$	25
norman	$\frac{bBx^3}{3} + \left(\frac{Ab}{2} + \frac{Ba}{2}\right)x^2 + aAx$	26
orering	$\frac{x(2bBx^2+3Abx+3Bax+6Aa)}{6}$	26
gosper	$\frac{1}{3}bBx^3 + \frac{1}{2}x^2Ab + \frac{1}{2}x^2Ba + aAx$	27
risch	$\frac{1}{3}bBx^3 + \frac{1}{2}x^2Ab + \frac{1}{2}x^2Ba + aAx$	27
parallelrisc	$\frac{1}{3}bBx^3 + \frac{1}{2}x^2Ab + \frac{1}{2}x^2Ba + aAx$	27

input `int((b*x+a)*(B*x+A),x,method=_RETURNVERBOSE)`

output `a*A*x+1/2*(A*b+B*a)*x^2+1/3*b*B*x^3`

Fricas [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int (a + bx)(A + Bx) dx = \frac{1}{3}x^3bB + \frac{1}{2}x^2aB + \frac{1}{2}x^2bA + xaA$$

input `integrate((b*x+a)*(B*x+A),x, algorithm="fricas")`

output `1/3*x^3*b*B + 1/2*x^2*a*B + 1/2*x^2*b*A + x*a*A`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int (a + bx)(A + Bx) dx = Aax + \frac{Bbx^3}{3} + x^2 \left(\frac{Ab}{2} + \frac{Ba}{2} \right)$$

input `integrate((b*x+a)*(B*x+A),x)`output `A*a*x + B*b*x**3/3 + x**2*(A*b/2 + B*a/2)`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.86

$$\int (a + bx)(A + Bx) dx = \frac{1}{3} Bbx^3 + Aax + \frac{1}{2} (Ba + Ab)x^2$$

input `integrate((b*x+a)*(B*x+A),x, algorithm="maxima")`output `1/3*B*b*x^3 + A*a*x + 1/2*(B*a + A*b)*x^2`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int (a + bx)(A + Bx) dx = \frac{1}{3} Bbx^3 + \frac{1}{2} Bax^2 + \frac{1}{2} Abx^2 + Aax$$

input `integrate((b*x+a)*(B*x+A),x, algorithm="giac")`output `1/3*B*b*x^3 + 1/2*B*a*x^2 + 1/2*A*b*x^2 + A*a*x`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.89

$$\int (a + bx)(A + Bx) dx = \frac{Bbx^3}{3} + \left(\frac{Ab}{2} + \frac{Ba}{2}\right)x^2 + Aax$$

input `int((A + B*x)*(a + b*x),x)`

output `x^2*((A*b)/2 + (B*a)/2) + A*a*x + (B*b*x^3)/3`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.75

$$\int (a + bx)(A + Bx) dx = \frac{x(b^2x^2 + 3abx + 3a^2)}{3}$$

input `int((b*x+a)*(B*x+A),x)`

output `(x*(3*a**2 + 3*a*b*x + b**2*x**2))/3`

3.6 $\int \frac{(a+bx)(A+Bx)}{d+ex} dx$

Optimal result	154
Mathematica [A] (verified)	154
Rubi [A] (verified)	155
Maple [A] (verified)	156
Fricas [A] (verification not implemented)	156
Sympy [A] (verification not implemented)	157
Maxima [A] (verification not implemented)	157
Giac [A] (verification not implemented)	158
Mupad [B] (verification not implemented)	158
Reduce [B] (verification not implemented)	159

Optimal result

Integrand size = 18, antiderivative size = 60

$$\int \frac{(a+bx)(A+Bx)}{d+ex} dx = -\frac{b(Bd-Ae)x}{e^2} + \frac{B(a+bx)^2}{2be} + \frac{(bd-ae)(Bd-Ae)\log(d+ex)}{e^3}$$

output

```
-b*(-A*e+B*d)*x/e^2+1/2*B*(b*x+a)^2/b/e+(-a*e+b*d)*(-A*e+B*d)*ln(e*x+d)/e^3
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.93

$$\int \frac{(a+bx)(A+Bx)}{d+ex} dx = \frac{ex(2aBe + b(-2Bd + 2Ae + Bex)) + 2(bd-ae)(Bd-Ae)\log(d+ex)}{2e^3}$$

input

```
Integrate[((a + b*x)*(A + B*x))/(d + e*x),x]
```

output

```
(e*x*(2*a*B*e + b*(-2*B*d + 2*A*e + B*e*x)) + 2*(b*d - a*e)*(B*d - A*e)*Log[d + e*x])/(2*e^3)
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)(A + Bx)}{d + ex} dx$$

↓ 86

$$\int \left(\frac{(ae - bd)(Ae - Bd)}{e^2(d + ex)} + \frac{B(a + bx)}{e} + \frac{b(Ae - Bd)}{e^2} \right) dx$$

↓ 2009

$$\frac{(bd - ae)(Bd - Ae) \log(d + ex)}{e^3} + \frac{B(a + bx)^2}{2be} - \frac{bx(Bd - Ae)}{e^2}$$

input

```
Int[((a + b*x)*(A + B*x))/(d + e*x), x]
```

output

```
-((b*(B*d - A*e)*x)/e^2) + (B*(a + b*x)^2)/(2*b*e) + ((b*d - a*e)*(B*d - A*e)*Log[d + e*x])/e^3
```

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```


rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.10

method	result	size
default	$\frac{\frac{1}{2}Bbe^2x^2 + Abex + Baex - Bbdx}{e^2} + \frac{(Aae^2 - Abde - Bade + bBd^2) \ln(ex+d)}{e^3}$	66
norman	$\frac{(Abe + Bae - Bbd)x}{e^2} + \frac{bBx^2}{2e} + \frac{(Aae^2 - Abde - Bade + bBd^2) \ln(ex+d)}{e^3}$	66
parallelrisc	$\frac{bBx^2e^2 + 2A \ln(ex+d)a e^2 - 2A \ln(ex+d)bde + 2Axb e^2 - 2B \ln(ex+d)ade + 2B \ln(ex+d)bd^2 + 2Bxa e^2 - 2Bxbde}{2e^3}$	89
risc	$\frac{bBx^2}{2e} + \frac{Abx}{e} + \frac{Bax}{e} - \frac{Bbdx}{e^2} + \frac{\ln(ex+d)Aa}{e} - \frac{\ln(ex+d)Abd}{e^2} - \frac{\ln(ex+d)Bad}{e^2} + \frac{\ln(ex+d)bBd^2}{e^3}$	90

input `int((b*x+a)*(B*x+A)/(e*x+d), x, method=_RETURNVERBOSE)`

output $\frac{1}{e^2} * (1/2 * B * b * e * x^2 + A * b * e * x + B * a * e * x - B * b * d * x) + (A * a * e^2 - A * b * d * e - B * a * d * e + B * b * d^2) / e^3 * \ln(e * x + d)$

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.13

$$\int \frac{(a + bx)(A + Bx)}{d + ex} dx$$

$$= \frac{Bbe^2x^2 - 2(Bbde - (Ba + Ab)e^2)x + 2(Bbd^2 + Aae^2 - (Ba + Ab)de) \log(ex + d)}{2e^3}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d), x, algorithm="fricas")`

output $\frac{1}{2} * (B * b * e^2 * x^2 - 2 * (B * b * d * e - (B * a + A * b) * e^2) * x + 2 * (B * b * d^2 + A * a * e^2 - (B * a + A * b) * d * e) * \log(e * x + d)) / e^3$

Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.88

$$\int \frac{(a + bx)(A + Bx)}{d + ex} dx = \frac{Bbx^2}{2e} + x \left(\frac{Ab}{e} + \frac{Ba}{e} - \frac{Bbd}{e^2} \right) - \frac{(-Ae + Bd)(ae - bd) \log(d + ex)}{e^3}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d),x)`output `B*b*x**2/(2*e) + x*(A*b/e + B*a/e - B*b*d/e**2) - (-A*e + B*d)*(a*e - b*d)*log(d + e*x)/e**3`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.10

$$\int \frac{(a + bx)(A + Bx)}{d + ex} dx = \frac{Bbx^2 - 2(Bbd - (Ba + Ab)e)x}{2e^2} + \frac{(Bbd^2 + Aae^2 - (Ba + Ab)de) \log(ex + d)}{e^3}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d),x, algorithm="maxima")`output `1/2*(B*b*e*x^2 - 2*(B*b*d - (B*a + A*b)*e)*x)/e^2 + (B*b*d^2 + A*a*e^2 - (B*a + A*b)*d*e)*log(e*x + d)/e^3`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.13

$$\int \frac{(a + bx)(A + Bx)}{d + ex} dx = \frac{Bbx^2 - 2Bbdx + 2Baex + 2Abex}{2e^2} + \frac{(Bbd^2 - Bade - Abde + Aae^2) \log(|ex + d|)}{e^3}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d),x, algorithm="giac")`output `1/2*(B*b*e*x^2 - 2*B*b*d*x + 2*B*a*e*x + 2*A*b*e*x)/e^2 + (B*b*d^2 - B*a*d*e - A*b*d*e + A*a*e^2)*log(abs(e*x + d))/e^3`**Mupad [B] (verification not implemented)**

Time = 0.89 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.13

$$\int \frac{(a + bx)(A + Bx)}{d + ex} dx = x \left(\frac{Ab + Ba}{e} - \frac{Bbd}{e^2} \right) + \frac{\ln(d + ex) (Aae^2 + Bbd^2 - Abde - Bade)}{e^3} + \frac{Bbx^2}{2e}$$

input `int(((A + B*x)*(a + b*x))/(d + e*x),x)`output `x*((A*b + B*a)/e - (B*b*d)/e^2) + (log(d + e*x)*(A*a*e^2 + B*b*d^2 - A*b*d*e - B*a*d*e))/e^3 + (B*b*x^2)/(2*e)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.20

$$\int \frac{(a + bx)(A + Bx)}{d + ex} dx$$

$$= \frac{2 \log(ex + d) a^2 e^2 - 4 \log(ex + d) abde + 2 \log(ex + d) b^2 d^2 + 4ab e^2 x - 2b^2 dex + b^2 e^2 x^2}{2e^3}$$

input `int((b*x+a)*(B*x+A)/(e*x+d),x)`output `(2*log(d + e*x)*a**2*e**2 - 4*log(d + e*x)*a*b*d*e + 2*log(d + e*x)*b**2*d**2 + 4*a*b*e**2*x - 2*b**2*d*e*x + b**2*e**2*x**2)/(2*e**3)`

3.7 $\int \frac{(a+bx)(A+Bx)}{(d+ex)^2} dx$

Optimal result	160
Mathematica [A] (verified)	160
Rubi [A] (verified)	161
Maple [A] (verified)	162
Fricas [A] (verification not implemented)	162
Sympy [A] (verification not implemented)	163
Maxima [A] (verification not implemented)	163
Giac [A] (verification not implemented)	164
Mupad [B] (verification not implemented)	164
Reduce [B] (verification not implemented)	165

Optimal result

Integrand size = 18, antiderivative size = 63

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^2} dx = \frac{bBx}{e^2} - \frac{(bd-ae)(Bd-Ae)}{e^3(d+ex)} - \frac{(2bBd - Abe - aBe) \log(d+ex)}{e^3}$$

output `b*B*x/e^2-(-a*e+b*d)*(-A*e+B*d)/e^3/(e*x+d)-(-A*b*e-B*a*e+2*B*b*d)*ln(e*x+d)/e^3`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.89

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^2} dx = \frac{bBex - \frac{(bd-ae)(Bd-Ae)}{d+ex} + (-2bBd + Abe + aBe) \log(d+ex)}{e^3}$$

input `Integrate[((a + b*x)*(A + B*x))/(d + e*x)^2,x]`

output `(b*B*e*x - ((b*d - a*e)*(B*d - A*e))/(d + e*x) + (-2*b*B*d + A*b*e + a*B*e)*Log[d + e*x])/e^3`

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^2} dx$$

↓ 86

$$\int \left(\frac{aBe + Abe - 2bBd}{e^2(d + ex)} + \frac{(ae - bd)(Ae - Bd)}{e^2(d + ex)^2} + \frac{bB}{e^2} \right) dx$$

↓ 2009

$$-\frac{(bd - ae)(Bd - Ae)}{e^3(d + ex)} - \frac{\log(d + ex)(-aBe - Abe + 2bBd)}{e^3} + \frac{bBx}{e^2}$$

input `Int[((a + b*x)*(A + B*x))/(d + e*x)^2,x]`

output `(b*B*x)/e^2 - ((b*d - a*e)*(B*d - A*e))/(e^3*(d + e*x)) - ((2*b*B*d - A*b*e - a*B*e)*Log[d + e*x])/e^3`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.11

method	result
default	$\frac{bBx}{e^2} - \frac{Aa e^2 - Abde - Bade + bB d^2}{e^3(ex+d)} + \frac{(Abe + Bae - 2Bbd) \ln(ex+d)}{e^3}$
norman	$\frac{bB x^2}{e} - \frac{Aa e^2 - Abde - Bade + 2bB d^2}{e^3} + \frac{(Abe + Bae - 2Bbd) \ln(ex+d)}{e^3}$
risch	$\frac{bBx}{e^2} - \frac{Aa}{e(ex+d)} + \frac{Abd}{e^2(ex+d)} + \frac{Bad}{e^2(ex+d)} - \frac{bB d^2}{e^3(ex+d)} + \frac{\ln(ex+d)Ab}{e^2} + \frac{\ln(ex+d)Ba}{e^2} - \frac{2 \ln(ex+d)Bbd}{e^3}$
parallelrisc	$\frac{A \ln(ex+d)xb e^2 + B \ln(ex+d)xa e^2 - 2B \ln(ex+d)xbde + bB x^2 e^2 + A \ln(ex+d)bde + B \ln(ex+d)ade - 2B \ln(ex+d)b d^2 - Aa e^2}{e^3(ex+d)}$

input `int((b*x+a)*(B*x+A)/(e*x+d)^2,x,method=_RETURNVERBOSE)`output `b*B*x/e^2-(A*a*e^2-A*b*d*e-B*a*d*e+B*b*d^2)/e^3/(e*x+d)+1/e^3*(A*b*e+B*a*e-2*B*b*d)*ln(e*x+d)`**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.62

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^2} dx$$

$$= \frac{Bbe^2x^2 + Bbdex - Bbd^2 - Aae^2 + (Ba + Ab)de - (2Bbd^2 - (Ba + Ab)de + (2Bbde - (Ba + Ab)e^2)x)}{e^4x + de^3}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^2,x, algorithm="fricas")`output `(B*b*e^2*x^2 + B*b*d*e*x - B*b*d^2 - A*a*e^2 + (B*a + A*b)*d*e - (2*B*b*d^2 - (B*a + A*b)*d*e + (2*B*b*d*e - (B*a + A*b)*e^2)*x)*log(e*x + d)/(e^4*x + d*e^3)`

Sympy [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.13

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^2} dx = \frac{Bbx}{e^2} + \frac{-Aae^2 + Abde + Bade - Bbd^2}{de^3 + e^4x} + \frac{(Abe + Bae - 2Bbd) \log(d + ex)}{e^3}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)**2,x)`output `B*b*x/e**2 + (-A*a*e**2 + A*b*d*e + B*a*d*e - B*b*d**2)/(d*e**3 + e**4*x) + (A*b*e + B*a*e - 2*B*b*d)*log(d + e*x)/e**3`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.17

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^2} dx = \frac{Bbx}{e^2} - \frac{Bbd^2 + Aae^2 - (Ba + Ab)de}{e^4x + de^3} - \frac{(2Bbd - (Ba + Ab)e) \log(ex + d)}{e^3}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^2,x, algorithm="maxima")`output `B*b*x/e^2 - (B*b*d^2 + A*a*e^2 - (B*a + A*b)*d*e)/(e^4*x + d*e^3) - (2*B*b*d - (B*a + A*b)*e)*log(e*x + d)/e^3`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.81

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^2} dx = \frac{(ex + d)Bb}{e^3} + \frac{(2Bbd - Bae - Abe) \log\left(\frac{|ex+d|}{(ex+d)^2|e|}\right)}{e^3} - \frac{\frac{Bbd^2e}{ex+d} - \frac{Bade^2}{ex+d} - \frac{Abde^2}{ex+d} + \frac{Aae^3}{ex+d}}{e^4}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^2,x, algorithm="giac")`output `(e*x + d)*B*b/e^3 + (2*B*b*d - B*a*e - A*b*e)*log(abs(e*x + d)/((e*x + d)^2*abs(e)))/e^3 - (B*b*d^2*e/(e*x + d) - B*a*d*e^2/(e*x + d) - A*b*d*e^2/(e*x + d) + A*a*e^3/(e*x + d))/e^4`**Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.19

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^2} dx = \frac{\ln(d + ex) (Abe + Bae - 2Bbd)}{e^3} - \frac{Aae^2 + Bbd^2 - Abde - Bade}{e(xe^3 + de^2)} + \frac{Bbx}{e^2}$$

input `int(((A + B*x)*(a + b*x))/(d + e*x)^2,x)`output `(log(d + e*x)*(A*b*e + B*a*e - 2*B*b*d))/e^3 - (A*a*e^2 + B*b*d^2 - A*b*d*e - B*a*d*e)/(e*(d*e^2 + e^3*x)) + (B*b*x)/e^2`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.78

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^2} dx$$

$$= \frac{2 \log(ex + d) ab d^2 e + 2 \log(ex + d) abd e^2 x - 2 \log(ex + d) b^2 d^3 - 2 \log(ex + d) b^2 d^2 ex + a^2 e^3 x - 2abd e}{d e^3 (ex + d)}$$

input `int((b*x+a)*(B*x+A)/(e*x+d)^2,x)`output `(2*log(d + e*x)*a*b*d**2*e + 2*log(d + e*x)*a*b*d*e**2*x - 2*log(d + e*x)*b**2*d**3 - 2*log(d + e*x)*b**2*d**2*e*x + a**2*e**3*x - 2*a*b*d*e**2*x + 2*b**2*d**2*e*x + b**2*d*e**2*x**2)/(d*e**3*(d + e*x))`

3.8 $\int \frac{(a+bx)(A+Bx)}{(d+ex)^3} dx$

Optimal result	166
Mathematica [A] (verified)	166
Rubi [A] (verified)	167
Maple [A] (verified)	168
Fricas [A] (verification not implemented)	168
Sympy [A] (verification not implemented)	169
Maxima [A] (verification not implemented)	169
Giac [A] (verification not implemented)	170
Mupad [B] (verification not implemented)	170
Reduce [B] (verification not implemented)	170

Optimal result

Integrand size = 18, antiderivative size = 70

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^3} dx = -\frac{(bd-ae)(Bd-Ae)}{2e^3(d+ex)^2} + \frac{2bBd-Abe-aBe}{e^3(d+ex)} + \frac{bB \log(d+ex)}{e^3}$$

output

```
-1/2*(-a*e+b*d)*(-A*e+B*d)/e^3/(e*x+d)^2+(-A*b*e-B*a*e+2*B*b*d)/e^3/(e*x+d)+b*B*ln(e*x+d)/e^3
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.03

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^3} dx = \frac{-ae(Ae+B(d+2ex)) + b(-Ae(d+2ex) + Bd(3d+4ex)) + 2bB(d+ex)^2 \log(d+ex)}{2e^3(d+ex)^2}$$

input

```
Integrate[((a + b*x)*(A + B*x))/(d + e*x)^3,x]
```

output

```
(-(a*e*(A*e + B*(d + 2*e*x))) + b*(-(A*e*(d + 2*e*x)) + B*d*(3*d + 4*e*x)) + 2*b*B*(d + e*x)^2*Log[d + e*x])/(2*e^3*(d + e*x)^2)
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^3} dx$$

↓ 86

$$\int \left(\frac{aBe + Abe - 2bBd}{e^2(d + ex)^2} + \frac{(ae - bd)(Ae - Bd)}{e^2(d + ex)^3} + \frac{bB}{e^2(d + ex)} \right) dx$$

↓ 2009

$$-\frac{(bd - ae)(Bd - Ae)}{2e^3(d + ex)^2} + \frac{-aBe - Abe + 2bBd}{e^3(d + ex)} + \frac{bB \log(d + ex)}{e^3}$$

input `Int[((a + b*x)*(A + B*x))/(d + e*x)^3,x]`

output `-1/2*((b*d - a*e)*(B*d - A*e))/(e^3*(d + e*x)^2) + (2*b*B*d - A*b*e - a*B*e)/(e^3*(d + e*x)) + (b*B*Log[d + e*x])/e^3`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.03

method	result
norman	$\frac{-\frac{Aa e^2 + Abde + Bade - 3bB d^2}{2e^3} - \frac{(Abe + Bae - 2Bbd)x}{e^2}}{(ex+d)^2} + \frac{bB \ln(ex+d)}{e^3}$
risch	$\frac{-\frac{Aa e^2 + Abde + Bade - 3bB d^2}{2e^3} - \frac{(Abe + Bae - 2Bbd)x}{e^2}}{(ex+d)^2} + \frac{bB \ln(ex+d)}{e^3}$
default	$-\frac{Abe + Bae - 2Bbd}{e^3(ex+d)} - \frac{Aa e^2 - Abde - Bade + bB d^2}{2e^3(ex+d)^2} + \frac{bB \ln(ex+d)}{e^3}$
parallelrisch	$-\frac{-2B \ln(ex+d)x^2 b e^2 - 4B \ln(ex+d)xbde + 2Axb e^2 - 2B \ln(ex+d) b d^2 + 2Bxa e^2 - 4Bxbde + Aa e^2 + Abde + Bade - 3bB d^2}{2e^3(ex+d)^2}$

input `int((b*x+a)*(B*x+A)/(e*x+d)^3,x,method=_RETURNVERBOSE)`output
$$\frac{(-1/2*(A*a*e^2+A*b*d*e+B*a*d*e-3*B*b*d^2)/e^3-(A*b*e+B*a*e-2*B*b*d)/e^2*x)/(e*x+d)^2+b*B*\ln(e*x+d)/e^3}$$
Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.50

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^3} dx$$

$$= \frac{3 Bbd^2 - Aae^2 - (Ba + Ab)de + 2(2 Bbde - (Ba + Ab)e^2)x + 2(Bbe^2x^2 + 2 Bbdex + Bbd^2) \log(ex + d)}{2(e^5x^2 + 2de^4x + d^2e^3)}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^3,x, algorithm="fricas")`output
$$\frac{1/2*(3*B*b*d^2 - A*a*e^2 - (B*a + A*b)*d*e + 2*(2*B*b*d*e - (B*a + A*b)*e^2)*x + 2*(B*b*e^2*x^2 + 2*B*b*d*e*x + B*b*d^2)*\log(e*x + d)}{(e^5*x^2 + 2*d*e^4*x + d^2*e^3)}$$

Sympy [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.34

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^3} dx = \frac{Bb \log(d + ex)}{e^3} + \frac{-Aae^2 - Abde - Bade + 3Bbd^2 + x(-2Abe^2 - 2Bae^2 + 4Bbde)}{2d^2e^3 + 4de^4x + 2e^5x^2}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)**3,x)`output `B*b*log(d + e*x)/e**3 + (-A*a*e**2 - A*b*d*e - B*a*d*e + 3*B*b*d**2 + x*(-2*A*b*e**2 - 2*B*a*e**2 + 4*B*b*d*e))/(2*d**2*e**3 + 4*d*e**4*x + 2*e**5*x**2)`**Maxima [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.24

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^3} dx = \frac{3Bbd^2 - Aae^2 - (Ba + Ab)de + 2(2Bbde - (Ba + Ab)e^2)x}{2(e^5x^2 + 2de^4x + d^2e^3)} + \frac{Bb \log(ex + d)}{e^3}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^3,x, algorithm="maxima")`output `1/2*(3*B*b*d^2 - A*a*e^2 - (B*a + A*b)*d*e + 2*(2*B*b*d*e - (B*a + A*b)*e^2)*x)/(e^5*x^2 + 2*d*e^4*x + d^2*e^3) + B*b*log(e*x + d)/e^3`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.10

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^3} dx = \frac{Bb \log(|ex+d|)}{e^3} + \frac{2(2Bbd - Bae - Abe)x + \frac{3Bbd^2 - Bade - Abde - Aae^2}{e}}{2(ex+d)^2 e^2}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^3,x, algorithm="giac")`output `B*b*log(abs(e*x + d))/e^3 + 1/2*(2*(2*B*b*d - B*a*e - A*b*e)*x + (3*B*b*d^2 - B*a*d*e - A*b*d*e - A*a*e^2)/e)/((e*x + d)^2*e^2)`**Mupad [B] (verification not implemented)**

Time = 0.91 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.17

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^3} dx = \frac{Bb \ln(d+ex)}{e^3} - \frac{\frac{Aae^2 - 3Bbd^2 + Abde + Bade}{2e^3} + \frac{x(Abe + Bae - 2Bbd)}{e^2}}{d^2 + 2dex + e^2 x^2}$$

input `int(((A + B*x)*(a + b*x))/(d + e*x)^3,x)`output `(B*b*log(d + e*x))/e^3 - ((A*a*e^2 - 3*B*b*d^2 + A*b*d*e + B*a*d*e)/(2*e^3) + (x*(A*b*e + B*a*e - 2*B*b*d))/e^2)/(d^2 + e^2*x^2 + 2*d*e*x)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.61

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^3} dx = \frac{2 \log(ex+d) b^2 d^3 + 4 \log(ex+d) b^2 d^2 ex + 2 \log(ex+d) b^2 d e^2 x^2 - a^2 d e^2 + 2 ab e^3 x^2 + b^2 d^3 - 2 b^2 d e^2 x^2}{2 d e^3 (e^2 x^2 + 2 dex + d^2)}$$

input `int((b*x+a)*(B*x+A)/(e*x+d)^3,x)`

output `(2*log(d + e*x)*b**2*d**3 + 4*log(d + e*x)*b**2*d**2*e*x + 2*log(d + e*x)*
b**2*d*e**2*x**2 - a**2*d*e**2 + 2*a*b*e**3*x**2 + b**2*d**3 - 2*b**2*d*e*
*2*x**2)/(2*d*e**3*(d**2 + 2*d*e*x + e**2*x**2))`

3.9 $\int \frac{(a+bx)(A+Bx)}{(d+ex)^4} dx$

Optimal result	172
Mathematica [A] (verified)	172
Rubi [A] (verified)	173
Maple [A] (verified)	174
Fricas [A] (verification not implemented)	174
Sympy [A] (verification not implemented)	175
Maxima [A] (verification not implemented)	175
Giac [A] (verification not implemented)	176
Mupad [B] (verification not implemented)	176
Reduce [B] (verification not implemented)	176

Optimal result

Integrand size = 18, antiderivative size = 75

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^4} dx = -\frac{(bd - ae)(Bd - Ae)}{3e^3(d + ex)^3} + \frac{2bBd - Abe - aBe}{2e^3(d + ex)^2} - \frac{bB}{e^3(d + ex)}$$

output

```
-1/3*(-a*e+b*d)*(-A*e+B*d)/e^3/(e*x+d)^3+1/2*(-A*b*e-B*a*e+2*B*b*d)/e^3/(e*x+d)^2-b*B/e^3/(e*x+d)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.84

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^4} dx = -\frac{ae(2Ae + B(d + 3ex)) + b(Ae(d + 3ex) + 2B(d^2 + 3dex + 3e^2x^2))}{6e^3(d + ex)^3}$$

input

```
Integrate[((a + b*x)*(A + B*x))/(d + e*x)^4,x]
```

output

```
-1/6*(a*e*(2*A*e + B*(d + 3*e*x)) + b*(A*e*(d + 3*e*x) + 2*B*(d^2 + 3*d*e*x + 3*e^2*x^2)))/(e^3*(d + e*x)^3)
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^4} dx$$

↓ 86

$$\int \left(\frac{aBe + Abe - 2bBd}{e^2(d + ex)^3} + \frac{(ae - bd)(Ae - Bd)}{e^2(d + ex)^4} + \frac{bB}{e^2(d + ex)^2} \right) dx$$

↓ 2009

$$\frac{-aBe - Abe + 2bBd}{2e^3(d + ex)^2} - \frac{(bd - ae)(Bd - Ae)}{3e^3(d + ex)^3} - \frac{bB}{e^3(d + ex)}$$

input `Int[((a + b*x)*(A + B*x))/(d + e*x)^4,x]`

output `-1/3*((b*d - a*e)*(B*d - A*e))/(e^3*(d + e*x)^3) + (2*b*B*d - A*b*e - a*B*e)/(2*e^3*(d + e*x)^2) - (b*B)/(e^3*(d + e*x))`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.93

method	result	size
norman	$\frac{-\frac{bBx^2}{e} - \frac{(Abe+BAe+2Bbd)x}{2e^2} - \frac{2Aae^2+Abde+BAde+2bBd^2}{6e^3}}{(ex+d)^3}$	70
risch	$\frac{-\frac{bBx^2}{e} - \frac{(Abe+BAe+2Bbd)x}{2e^2} - \frac{2Aae^2+Abde+BAde+2bBd^2}{6e^3}}{(ex+d)^3}$	70
gospers	$\frac{-6bBx^2e^2+3Axb e^2+3Bxa e^2+6Bxbde+2Aae^2+Abde+BAde+2bBd^2}{6(ex+d)^3e^3}$	71
parallelrisch	$\frac{-6bBx^2e^2+3Axb e^2+3Bxa e^2+6Bxbde+2Aae^2+Abde+BAde+2bBd^2}{6(ex+d)^3e^3}$	71
orering	$\frac{-6bBx^2e^2+3Axb e^2+3Bxa e^2+6Bxbde+2Aae^2+Abde+BAde+2bBd^2}{6(ex+d)^3e^3}$	71
default	$-\frac{bB}{e^3(ex+d)} - \frac{Abe+BAe-2Bbd}{2e^3(ex+d)^2} - \frac{Aae^2-Abde-BAde+bBd^2}{3e^3(ex+d)^3}$	79

input `int((b*x+a)*(B*x+A)/(e*x+d)^4,x,method=_RETURNVERBOSE)`output `1/(e*x+d)^3*(-b*B/e*x^2-1/2*(A*b*e+B*a*e+2*B*b*d)/e^2*x-1/6*(2*A*a*e^2+A*b*d*e+B*a*d*e+2*B*b*d^2)/e^3)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.24

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^4} dx$$

$$= -\frac{6Bbe^2x^2+2Bbd^2+2Aae^2+(Ba+Ab)de+3(2Bbde+(Ba+Ab)e^2)x}{6(e^6x^3+3de^5x^2+3d^2e^4x+d^3e^3)}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^4,x, algorithm="fricas")`output `-1/6*(6*B*b*e^2*x^2+2*B*b*d^2+2*A*a*e^2+(B*a+A*b)*d*e+3*(2*B*b*d*e+(B*a+A*b)*e^2)*x)/(e^6*x^3+3*d*e^5*x^2+3*d^2*e^4*x+d^3*e^3)`

Sympy [A] (verification not implemented)

Time = 0.71 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.43

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^4} dx$$

$$= \frac{-2Aae^2 - Abde - Bade - 2Bbd^2 - 6Bbe^2x^2 + x(-3Abe^2 - 3Bae^2 - 6Bbde)}{6d^3e^3 + 18d^2e^4x + 18de^5x^2 + 6e^6x^3}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)**4,x)`output `(-2*A*a*e**2 - A*b*d*e - B*a*d*e - 2*B*b*d**2 - 6*B*b*e**2*x**2 + x*(-3*A*b*e**2 - 3*B*a*e**2 - 6*B*b*d*e))/(6*d**3*e**3 + 18*d**2*e**4*x + 18*d*e**5*x**2 + 6*e**6*x**3)`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.24

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^4} dx$$

$$= -\frac{6Bbe^2x^2 + 2Bbd^2 + 2Aae^2 + (Ba + Ab)de + 3(2Bbde + (Ba + Ab)e^2)x}{6(e^6x^3 + 3de^5x^2 + 3d^2e^4x + d^3e^3)}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^4,x, algorithm="maxima")`output `-1/6*(6*B*b*e^2*x^2 + 2*B*b*d^2 + 2*A*a*e^2 + (B*a + A*b)*d*e + 3*(2*B*b*d*e + (B*a + A*b)*e^2)*x)/(e^6*x^3 + 3*d*e^5*x^2 + 3*d^2*e^4*x + d^3*e^3)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.93

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^4} dx = -\frac{6Bbe^2x^2 + 6Bbdex + 3Bae^2x + 3Abe^2x + 2Bbd^2 + Bade + Abde + 2Aae^2}{6(ex+d)^3e^3}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^4,x, algorithm="giac")`

output `-1/6*(6*B*b*e^2*x^2 + 6*B*b*d*e*x + 3*B*a*e^2*x + 3*A*b*e^2*x + 2*B*b*d^2 + B*a*d*e + A*b*d*e + 2*A*a*e^2)/((e*x + d)^3*e^3)`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.21

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^4} dx = -\frac{\frac{2Aae^2+2Bbd^2+Abde+Bade}{6e^3} + \frac{x(Abe+BAe+2Bbd)}{2e^2} + \frac{Bbx^2}{e}}{d^3 + 3d^2ex + 3de^2x^2 + e^3x^3}$$

input `int(((A + B*x)*(a + b*x))/(d + e*x)^4,x)`

output `-((2*A*a*e^2 + 2*B*b*d^2 + A*b*d*e + B*a*d*e)/(6*e^3) + (x*(A*b*e + B*a*e + 2*B*b*d))/(2*e^2) + (B*b*x^2)/e)/(d^3 + e^3*x^3 + 3*d*e^2*x^2 + 3*d^2*e*x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.92

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^4} dx = \frac{b^2e^2x^3 - 3abdex - a^2de - abd^2}{3de^2(e^3x^3 + 3de^2x^2 + 3d^2ex + d^3)}$$

input `int((b*x+a)*(B*x+A)/(e*x+d)^4,x)`

output $(-a^{**2}d*e - a*b*d^{**2} - 3*a*b*d*e*x + b^{**2}*e^{**2}*x^{**3})/(3*d*e^{**2}*(d^{**3} + 3*d^{**2}*e*x + 3*d*e^{**2}*x^{**2} + e^{**3}*x^{**3}))$

3.10 $\int \frac{(a+bx)(A+Bx)}{(d+ex)^5} dx$

Optimal result	178
Mathematica [A] (verified)	178
Rubi [A] (verified)	179
Maple [A] (verified)	180
Fricas [A] (verification not implemented)	180
Sympy [A] (verification not implemented)	181
Maxima [A] (verification not implemented)	181
Giac [A] (verification not implemented)	182
Mupad [B] (verification not implemented)	182
Reduce [B] (verification not implemented)	183

Optimal result

Integrand size = 18, antiderivative size = 77

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^5} dx = -\frac{(bd-ae)(Bd-Ae)}{4e^3(d+ex)^4} + \frac{2bBd-Abe-aBe}{3e^3(d+ex)^3} - \frac{bB}{2e^3(d+ex)^2}$$

output

```
-1/4*(-a*e+b*d)*(-A*e+B*d)/e^3/(e*x+d)^4+1/3*(-A*b*e-B*a*e+2*B*b*d)/e^3/(e*x+d)^3-1/2*b*B/e^3/(e*x+d)^2
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.81

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^5} dx = -\frac{ae(3Ae+B(d+4ex))+b(Ae(d+4ex)+B(d^2+4dex+6e^2x^2))}{12e^3(d+ex)^4}$$

input

```
Integrate[((a + b*x)*(A + B*x))/(d + e*x)^5,x]
```

output

```
-1/12*(a*e*(3A*e + B*(d + 4*e*x)) + b*(A*e*(d + 4*e*x) + B*(d^2 + 4*d*e*x + 6*e^2*x^2)))/(e^3*(d + e*x)^4)
```

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^5} dx$$

↓ 86

$$\int \left(\frac{aBe + Abe - 2bBd}{e^2(d + ex)^4} + \frac{(ae - bd)(Ae - Bd)}{e^2(d + ex)^5} + \frac{bB}{e^2(d + ex)^3} \right) dx$$

↓ 2009

$$\frac{-aBe - Abe + 2bBd}{3e^3(d + ex)^3} - \frac{(bd - ae)(Bd - Ae)}{4e^3(d + ex)^4} - \frac{bB}{2e^3(d + ex)^2}$$

input `Int[((a + b*x)*(A + B*x))/(d + e*x)^5,x]`

output `-1/4*((b*d - a*e)*(B*d - A*e))/(e^3*(d + e*x)^4) + (2*b*B*d - A*b*e - a*B*e)/(3*e^3*(d + e*x)^3) - (b*B)/(2*e^3*(d + e*x)^2)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.88

method	result	size
risch	$\frac{-\frac{bB}{2e}x^2 - \frac{(Abe+Bae+Bbd)x}{3e^2} - \frac{3Aae^2+Abde+Bade+bBd^2}{12e^3}}{(ex+d)^4}$	68
gospers	$-\frac{6bBx^2e^2+4Axb^2e^2+4Bxa^2e^2+4Bxbde+3Aae^2+Abde+Bade+bBd^2}{12e^3(ex+d)^4}$	70
orering	$-\frac{6bBx^2e^2+4Axb^2e^2+4Bxa^2e^2+4Bxbde+3Aae^2+Abde+Bade+bBd^2}{12e^3(ex+d)^4}$	70
parallelrisc	$-\frac{6bBx^2e^3+4Ab^3e^3x+4Ba^3e^3x+4Bbd^2e^2x+3Aae^3+Abde^2+Bade^2+bBd^2e}{12e^4(ex+d)^4}$	77
norman	$\frac{-\frac{bB}{2e}x^2 - \frac{(Ab^2e^2+Ba^2e^2+bBde)x}{3e^3} - \frac{3Aae^3+Abde^2+Bade^2+bBd^2e}{12e^4}}{(ex+d)^4}$	78
default	$-\frac{bB}{2e^3(ex+d)^2} - \frac{Aae^2-Abde-Bade+bBd^2}{4e^3(ex+d)^4} - \frac{Abe+Bae-2Bbd}{3e^3(ex+d)^3}$	79

input `int((b*x+a)*(B*x+A)/(e*x+d)^5,x,method=_RETURNVERBOSE)`

output `(-1/2*b*B/e*x^2-1/3/e^2*(A*b*e+B*a*e+B*b*d)*x-1/12/e^3*(3*A*a*e^2+A*b*d*e+B*a*d*e+B*b*d^2))/(e*x+d)^4`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.32

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^5} dx$$

$$= -\frac{6Bbe^2x^2+Bbd^2+3Aae^2+(Ba+Ab)de+4(Bbde+(Ba+Ab)e^2)x}{12(e^7x^4+4de^6x^3+6d^2e^5x^2+4d^3e^4x+d^4e^3)}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^5,x, algorithm="fricas")`

output `-1/12*(6*B*b*e^2*x^2+B*b*d^2+3*A*a*e^2+(B*a+A*b)*d*e+4*(B*b*d*e+(B*a+A*b)*e^2)*x)/(e^7*x^4+4*d*e^6*x^3+6*d^2*e^5*x^2+4*d^3*e^4*x+d^4*e^3)`

Sympy [A] (verification not implemented)

Time = 1.14 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.52

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^5} dx$$

$$= \frac{-3Aae^2 - Abde - Bade - Bbd^2 - 6Bbe^2x^2 + x(-4Abe^2 - 4Bae^2 - 4Bbde)}{12d^4e^3 + 48d^3e^4x + 72d^2e^5x^2 + 48de^6x^3 + 12e^7x^4}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)**5,x)`output `(-3*A*a*e**2 - A*b*d*e - B*a*d*e - B*b*d**2 - 6*B*b*e**2*x**2 + x*(-4*A*b*e**2 - 4*B*a*e**2 - 4*B*b*d*e))/(12*d**4*e**3 + 48*d**3*e**4*x + 72*d**2*e**5*x**2 + 48*d*e**6*x**3 + 12*e**7*x**4)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.32

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^5} dx$$

$$= -\frac{6Bbe^2x^2 + Bbd^2 + 3Aae^2 + (Ba + Ab)de + 4(Bbde + (Ba + Ab)e^2)x}{12(e^7x^4 + 4de^6x^3 + 6d^2e^5x^2 + 4d^3e^4x + d^4e^3)}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^5,x, algorithm="maxima")`output `-1/12*(6*B*b*e^2*x^2 + B*b*d^2 + 3*A*a*e^2 + (B*a + A*b)*d*e + 4*(B*b*d*e + (B*a + A*b)*e^2)*x)/(e^7*x^4 + 4*d*e^6*x^3 + 6*d^2*e^5*x^2 + 4*d^3*e^4*x + d^4*e^3)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.57

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^5} dx$$

$$= -\frac{\frac{3Aa}{(ex+d)^4} + \frac{6Bb}{(ex+d)^2e^2} - \frac{8Bbd}{(ex+d)^3e^2} + \frac{3Bbd^2}{(ex+d)^4e^2} + \frac{4Ba}{(ex+d)^3e} + \frac{4Ab}{(ex+d)^3e} - \frac{3Bad}{(ex+d)^4e} - \frac{3Abd}{(ex+d)^4e}}{12e}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^5,x, algorithm="giac")`output `-1/12*(3*A*a/(e*x + d)^4 + 6*B*b/((e*x + d)^2*e^2) - 8*B*b*d/((e*x + d)^3*e^2) + 3*B*b*d^2/((e*x + d)^4*e^2) + 4*B*a/((e*x + d)^3*e) + 4*A*b/((e*x + d)^3*e) - 3*B*a*d/((e*x + d)^4*e) - 3*A*b*d/((e*x + d)^4*e))/e`**Mupad [B] (verification not implemented)**

Time = 0.91 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.31

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^5} dx = -\frac{\frac{3Aae^2+Bbd^2+Abde+Bade}{12e^3} + \frac{x(Abe+BAe+Bbd)}{3e^2} + \frac{Bbx^2}{2e}}{d^4 + 4d^3ex + 6d^2e^2x^2 + 4de^3x^3 + e^4x^4}$$

input `int(((A + B*x)*(a + b*x))/(d + e*x)^5,x)`output `-((3*A*a*e^2 + B*b*d^2 + A*b*d*e + B*a*d*e)/(12*e^3) + (x*(A*b*e + B*a*e + B*b*d))/(3*e^2) + (B*b*x^2)/(2*e))/(d^4 + e^4*x^4 + 4*d*e^3*x^3 + 6*d^2*e^2*x^2 + 4*d^3*e*x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.23

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^5} dx = \frac{-6b^2e^2x^2 - 8abe^2x - 4b^2dex - 3a^2e^2 - 2abde - b^2d^2}{12e^3(e^4x^4 + 4de^3x^3 + 6d^2e^2x^2 + 4d^3ex + d^4)}$$

input

```
int((b*x+a)*(B*x+A)/(e*x+d)^5,x)
```

output

```
( - 3*a**2*e**2 - 2*a*b*d*e - 8*a*b*e**2*x - b**2*d**2 - 4*b**2*d*e*x - 6*
b**2*e**2*x**2)/(12*e**3*(d**4 + 4*d**3*e*x + 6*d**2*e**2*x**2 + 4*d*e**3*
x**3 + e**4*x**4))
```

3.11 $\int \frac{(a+bx)(A+Bx)}{(d+ex)^6} dx$

Optimal result	184
Mathematica [A] (verified)	184
Rubi [A] (verified)	185
Maple [A] (verified)	186
Fricas [A] (verification not implemented)	186
Sympy [A] (verification not implemented)	187
Maxima [A] (verification not implemented)	187
Giac [A] (verification not implemented)	188
Mupad [B] (verification not implemented)	188
Reduce [B] (verification not implemented)	188

Optimal result

Integrand size = 18, antiderivative size = 77

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^6} dx = -\frac{(bd-ae)(Bd-Ae)}{5e^3(d+ex)^5} + \frac{2bBd-Abe-aBe}{4e^3(d+ex)^4} - \frac{bB}{3e^3(d+ex)^3}$$

output

```
-1/5*(-a*e+b*d)*(-A*e+B*d)/e^3/(e*x+d)^5+1/4*(-A*b*e-B*a*e+2*B*b*d)/e^3/(e*x+d)^4-1/3*b*B/e^3/(e*x+d)^3
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.84

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^6} dx = -\frac{3ae(4Ae+B(d+5ex))+b(3Ae(d+5ex)+2B(d^2+5dex+10e^2x^2))}{60e^3(d+ex)^5}$$

input

```
Integrate[((a + b*x)*(A + B*x))/(d + e*x)^6,x]
```

output

```
-1/60*(3*a*e*(4*A*e + B*(d + 5*e*x)) + b*(3*A*e*(d + 5*e*x) + 2*B*(d^2 + 5*d*e*x + 10*e^2*x^2)))/(e^3*(d + e*x)^5)
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^6} dx$$

↓ 86

$$\int \left(\frac{aBe + Abe - 2bBd}{e^2(d + ex)^5} + \frac{(ae - bd)(Ae - Bd)}{e^2(d + ex)^6} + \frac{bB}{e^2(d + ex)^4} \right) dx$$

↓ 2009

$$\frac{-aBe - Abe + 2bBd}{4e^3(d + ex)^4} - \frac{(bd - ae)(Bd - Ae)}{5e^3(d + ex)^5} - \frac{bB}{3e^3(d + ex)^3}$$

input `Int[((a + b*x)*(A + B*x))/(d + e*x)^6,x]`

output `-1/5*((b*d - a*e)*(B*d - A*e))/(e^3*(d + e*x)^5) + (2*b*B*d - A*b*e - a*B*e)/(4*e^3*(d + e*x)^4) - (b*B)/(3*e^3*(d + e*x)^3)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.))^(n_.)*((e_.) + (f_.)*(x_.))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.95

method	result	size
gospers	$-\frac{20bBx^2e^2+15Axb e^2+15Bxa e^2+10Bxbde+12Aa e^2+3Abde+3Bade+2bB d^2}{60e^3(ex+d)^5}$	73
orering	$-\frac{20bBx^2e^2+15Axb e^2+15Bxa e^2+10Bxbde+12Aa e^2+3Abde+3Bade+2bB d^2}{60e^3(ex+d)^5}$	73
risch	$\frac{\frac{bBx^2}{3e} - \frac{(3Abe+3Bae+2Bbd)x}{12e^2} - \frac{12Aa e^2+3Abde+3Bade+2bB d^2}{60e^3}}{(ex+d)^5}$	74
default	$-\frac{Aa e^2-Abde-Bade+bB d^2}{5e^3(ex+d)^5} - \frac{Abe+Bae-2Bbd}{4e^3(ex+d)^4} - \frac{bB}{3e^3(ex+d)^3}$	79
parallelrisch	$-\frac{20bBx^2e^4+15Ab e^4x+15Ba e^4x+10Bbd e^3x+12Aa e^4+3Abd e^3+3Bad e^3+2bB d^2e^2}{60e^5(ex+d)^5}$	82
norman	$\frac{\frac{bBx^2}{3e} - \frac{(3Ab e^3+3Ba e^3+2bBd e^2)x}{12e^4} - \frac{12Aa e^4+3Abd e^3+3Bad e^3+2bB d^2e^2}{60e^5}}{(ex+d)^5}$	88

input `int((b*x+a)*(B*x+A)/(e*x+d)^6,x,method=_RETURNVERBOSE)`output
$$-1/60/e^3*(20*B*b*e^2*x^2+15*A*b*e^2*x+15*B*a*e^2*x+10*B*b*d*e*x+12*A*a*e^2+3*A*b*d*e+3*B*a*d*e+2*B*b*d^2)/(e*x+d)^5$$
Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.52

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^6} dx$$

$$= -\frac{20Bbe^2x^2+2Bbd^2+12Aae^2+3(Ba+Ab)de+5(2Bbde+3(Ba+Ab)e^2)x}{60(e^8x^5+5de^7x^4+10d^2e^6x^3+10d^3e^5x^2+5d^4e^4x+d^5e^3)}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^6,x, algorithm="fricas")`output
$$-1/60*(20*B*b*e^2*x^2+2*B*b*d^2+12*A*a*e^2+3*(B*a+A*b)*d*e+5*(2*B*b*d*e+3*(B*a+A*b)*e^2)*x)/(e^8*x^5+5*d*e^7*x^4+10*d^2*e^6*x^3+10*d^3*e^5*x^2+5*d^4*e^4*x+d^5*e^3)$$

Sympy [A] (verification not implemented)

Time = 1.72 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.74

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^6} dx$$

$$= \frac{-12Aae^2 - 3Abde - 3Bade - 2Bbd^2 - 20Bbe^2x^2 + x(-15Abe^2 - 15Bae^2 - 10Bbde)}{60d^5e^3 + 300d^4e^4x + 600d^3e^5x^2 + 600d^2e^6x^3 + 300de^7x^4 + 60e^8x^5}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)**6,x)`output `(-12*A*a*e**2 - 3*A*b*d*e - 3*B*a*d*e - 2*B*b*d**2 - 20*B*b*e**2*x**2 + x*(-15*A*b*e**2 - 15*B*a*e**2 - 10*B*b*d*e))/(60*d**5*e**3 + 300*d**4*e**4*x + 600*d**3*e**5*x**2 + 600*d**2*e**6*x**3 + 300*d*e**7*x**4 + 60*e**8*x**5)`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.52

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^6} dx$$

$$= -\frac{20 Bbe^2x^2 + 2 Bbd^2 + 12 Aae^2 + 3 (Ba + Ab)de + 5 (2 Bbde + 3 (Ba + Ab)e^2)x}{60 (e^8x^5 + 5 de^7x^4 + 10 d^2e^6x^3 + 10 d^3e^5x^2 + 5 d^4e^4x + d^5e^3)}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^6,x, algorithm="maxima")`output `-1/60*(20*B*b*e^2*x^2 + 2*B*b*d^2 + 12*A*a*e^2 + 3*(B*a + A*b)*d*e + 5*(2*B*b*d*e + 3*(B*a + A*b)*e^2)*x)/(e^8*x^5 + 5*d*e^7*x^4 + 10*d^2*e^6*x^3 + 10*d^3*e^5*x^2 + 5*d^4*e^4*x + d^5*e^3)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.94

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^6} dx = \frac{20Bbe^2x^2 + 10Bbdex + 15Bae^2x + 15Abe^2x + 2Bbd^2 + 3Bade + 3Abde + 12Aae^2}{60(ex+d)^5e^3}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^6,x, algorithm="giac")`output `-1/60*(20*B*b*e^2*x^2 + 10*B*b*d*e*x + 15*B*a*e^2*x + 15*A*b*e^2*x + 2*B*b*d^2 + 3*B*a*d*e + 3*A*b*d*e + 12*A*a*e^2)/((e*x + d)^5*e^3)`**Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.53

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^6} dx = -\frac{12Aae^2+2Bbd^2+3Abde+3Bade}{60e^3} + \frac{x(3Abe+3Bae+2Bbd)}{12e^2} + \frac{Bbx^2}{3e} \frac{1}{d^5 + 5d^4ex + 10d^3e^2x^2 + 10d^2e^3x^3 + 5de^4x^4 + e^5x^5}$$

input `int(((A + B*x)*(a + b*x))/(d + e*x)^6,x)`output `-((12*A*a*e^2 + 2*B*b*d^2 + 3*A*b*d*e + 3*B*a*d*e)/(60*e^3) + (x*(3*A*b*e + 3*B*a*e + 2*B*b*d))/(12*e^2) + (B*b*x^2)/(3*e))/(d^5 + e^5*x^5 + 5*d*e^4*x^4 + 10*d^3*e^2*x^2 + 10*d^2*e^3*x^3 + 5*d^4*e*x)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.38

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^6} dx = \frac{-10b^2e^2x^2 - 15abe^2x - 5b^2dex - 6a^2e^2 - 3abde - b^2d^2}{30e^3(e^5x^5 + 5de^4x^4 + 10d^2e^3x^3 + 10d^3e^2x^2 + 5d^4ex + d^5)}$$

input `int((b*x+a)*(B*x+A)/(e*x+d)^6,x)`

output

```
( - 6*a**2*e**2 - 3*a*b*d*e - 15*a*b*e**2*x - b**2*d**2 - 5*b**2*d*e*x - 1
0*b**2*e**2*x**2)/(30*e**3*(d**5 + 5*d**4*e*x + 10*d**3*e**2*x**2 + 10*d**
2*e**3*x**3 + 5*d*e**4*x**4 + e**5*x**5))
```

3.12 $\int (a + bx)^2(A + Bx)(d + ex)^4 dx$

Optimal result	190
Mathematica [B] (verified)	190
Rubi [A] (verified)	191
Maple [B] (verified)	193
Fricas [B] (verification not implemented)	193
Sympy [B] (verification not implemented)	195
Maxima [B] (verification not implemented)	196
Giac [B] (verification not implemented)	196
Mupad [B] (verification not implemented)	198
Reduce [B] (verification not implemented)	199

Optimal result

Integrand size = 20, antiderivative size = 120

$$\int (a + bx)^2(A + Bx)(d + ex)^4 dx = -\frac{(bd - ae)^2(Bd - Ae)(d + ex)^5}{5e^4} + \frac{(bd - ae)(3bBd - 2Abe - aBe)(d + ex)^6}{6e^4} - \frac{b(3bBd - Abe - 2aBe)(d + ex)^7}{7e^4} + \frac{b^2B(d + ex)^8}{8e^4}$$

output

```
-1/5*(-a*e+b*d)^2*(-A*e+B*d)*(e*x+d)^5/e^4+1/6*(-a*e+b*d)*(-2*A*b*e-B*a*e+
3*B*b*d)*(e*x+d)^6/e^4-1/7*b*(-A*b*e-2*B*a*e+3*B*b*d)*(e*x+d)^7/e^4+1/8*b^
2*B*(e*x+d)^8/e^4
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 283 vs. 2(120) = 240.

Time = 0.06 (sec) , antiderivative size = 283, normalized size of antiderivative = 2.36

$$\int (a + bx)^2(A + Bx)(d + ex)^4 dx = a^2Ad^4x + \frac{1}{2}ad^3(2Abd + aBd + 4aAe)x^2 + \frac{1}{3}d^2(2aBd(bd + 2ae) + A(b^2d^2 + 8abde + 6a^2e^2))x^3 + \frac{1}{4}d(2a^2e^2(3Bd + 2Ae) + 4abde(2Bd + 3Ae) + b^2d^2(Bd + 4Ae))x^4 + \frac{1}{5}e(a^2e^2(4Bd + Ae) + 4abde(3Bd + 2Ae) + 2b^2d^2(2Bd + 3Ae))x^5 + \frac{1}{6}e^2(a^2Be^2 + 2abe(4Bd + Ae) + 2b^2d(3Bd + 2Ae))x^6 + \frac{1}{7}be^3(4bBd + Abe + 2aBe)x^7 + \frac{1}{8}b^2Be^4x^8$$

input

```
Integrate[(a + b*x)^2*(A + B*x)*(d + e*x)^4,x]
```

output

```
a^2*A*d^4*x + (a*d^3*(2*A*b*d + a*B*d + 4*a*A*e)*x^2)/2 + (d^2*(2*a*B*d*(b*d + 2*a*e) + A*(b^2*d^2 + 8*a*b*d*e + 6*a^2*e^2))*x^3)/3 + (d*(2*a^2*e^2*(3*B*d + 2*A*e) + 4*a*b*d*e*(2*B*d + 3*A*e) + b^2*d^2*(B*d + 4*A*e))*x^4)/4 + (e*(a^2*e^2*(4*B*d + A*e) + 4*a*b*d*e*(3*B*d + 2*A*e) + 2*b^2*d^2*(2*B*d + 3*A*e))*x^5)/5 + (e^2*(a^2*B*e^2 + 2*a*b*e*(4*B*d + A*e) + 2*b^2*d*(3*B*d + 2*A*e))*x^6)/6 + (b*e^3*(4*b*B*d + A*b*e + 2*a*B*e)*x^7)/7 + (b^2*B*e^4*x^8)/8
```

Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^2(A + Bx)(d + ex)^4 dx$$

↓ 86

$$\int \left(\frac{b(d + ex)^6(2aBe + Abe - 3bBd)}{e^3} + \frac{(d + ex)^5(ae - bd)(aBe + 2Abe - 3bBd)}{e^3} + \frac{(d + ex)^4(ae - bd)^2(Ae - Bd)}{e^3} \right) dx$$

↓ 2009

$$-\frac{b(d + ex)^7(-2aBe - Abe + 3bBd)}{7e^4} + \frac{(d + ex)^6(bd - ae)(-aBe - 2Abe + 3bBd)}{(d + ex)^5(bd - ae)^2(Bd - Ae)} - \frac{b^2B(d + ex)^8}{8e^4}$$

input `Int[(a + b*x)^2*(A + B*x)*(d + e*x)^4,x]`

output `-1/5*((b*d - a*e)^2*(B*d - A*e)*(d + e*x)^5)/e^4 + ((b*d - a*e)*(3*b*B*d - 2*A*b*e - a*B*e)*(d + e*x)^6)/(6*e^4) - (b*(3*b*B*d - A*b*e - 2*a*B*e)*(d + e*x)^7)/(7*e^4) + (b^2*B*(d + e*x)^8)/(8*e^4)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Time = 0.06 (sec) , antiderivative size = 304, normalized size of antiderivative = 2.53

$$\begin{aligned}
 & \int (a + bx)^2(A + Bx)(d + ex)^4 dx \\
 &= \frac{1}{8} Bb^2e^4x^8 + Aa^2d^4x + \frac{1}{7} (4Bb^2de^3 + (2Bab + Ab^2)e^4)x^7 \\
 &+ \frac{1}{6} (6Bb^2d^2e^2 + 4(2Bab + Ab^2)de^3 + (Ba^2 + 2Aab)e^4)x^6 \\
 &+ \frac{1}{5} (4Bb^2d^3e + Aa^2e^4 + 6(2Bab + Ab^2)d^2e^2 + 4(Ba^2 + 2Aab)de^3)x^5 \\
 &+ \frac{1}{4} (Bb^2d^4 + 4Aa^2de^3 + 4(2Bab + Ab^2)d^3e + 6(Ba^2 + 2Aab)d^2e^2)x^4 \\
 &+ \frac{1}{3} (6Aa^2d^2e^2 + (2Bab + Ab^2)d^4 + 4(Ba^2 + 2Aab)d^3e)x^3 \\
 &+ \frac{1}{2} (4Aa^2d^3e + (Ba^2 + 2Aab)d^4)x^2
 \end{aligned}$$

input `integrate((b*x+a)^2*(B*x+A)*(e*x+d)^4,x, algorithm="fricas")`

output

```

1/8*B*b^2*e^4*x^8 + A*a^2*d^4*x + 1/7*(4*B*b^2*d*e^3 + (2*B*a*b + A*b^2)*e
^4)*x^7 + 1/6*(6*B*b^2*d^2*e^2 + 4*(2*B*a*b + A*b^2)*d*e^3 + (B*a^2 + 2*A*
a*b)*e^4)*x^6 + 1/5*(4*B*b^2*d^3*e + A*a^2*e^4 + 6*(2*B*a*b + A*b^2)*d^2*e
^2 + 4*(B*a^2 + 2*A*a*b)*d*e^3)*x^5 + 1/4*(B*b^2*d^4 + 4*A*a^2*d*e^3 + 4*(
2*B*a*b + A*b^2)*d^3*e + 6*(B*a^2 + 2*A*a*b)*d^2*e^2)*x^4 + 1/3*(6*A*a^2*d
^2*e^2 + (2*B*a*b + A*b^2)*d^4 + 4*(B*a^2 + 2*A*a*b)*d^3*e)*x^3 + 1/2*(4*A
*a^2*d^3*e + (B*a^2 + 2*A*a*b)*d^4)*x^2

```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 384 vs. $2(114) = 228$.

Time = 0.04 (sec) , antiderivative size = 384, normalized size of antiderivative = 3.20

$$\int (a + bx)^2(A + Bx)(d + ex)^4 dx = Aa^2d^4x + \frac{Bb^2e^4x^8}{8} + x^7\left(\frac{Ab^2e^4}{7} + \frac{2Babe^4}{7} + \frac{4Bb^2de^3}{7}\right) + x^6\left(\frac{Aabe^4}{3} + \frac{2Ab^2de^3}{3} + \frac{Ba^2e^4}{6} + \frac{4Babde^3}{3} + Bb^2d^2e^2\right) + x^5\left(\frac{Aa^2e^4}{5} + \frac{8Aabde^3}{5} + \frac{6Ab^2d^2e^2}{5} + \frac{4Ba^2de^3}{5} + \frac{12Babd^2e^2}{5} + \frac{4Bb^2d^3e}{5}\right) + x^4\left(Aa^2de^3 + 3Aabd^2e^2 + Ab^2d^3e + \frac{3Ba^2d^2e^2}{2} + 2Babd^3e + \frac{Bb^2d^4}{4}\right) + x^3\left(2Aa^2d^2e^2 + \frac{8Aabd^3e}{3} + \frac{Ab^2d^4}{3} + \frac{4Ba^2d^3e}{3} + \frac{2Babd^4}{3}\right) + x^2\left(2Aa^2d^3e + Aabd^4 + \frac{Ba^2d^4}{2}\right)$$

input `integrate((b*x+a)**2*(B*x+A)*(e*x+d)**4,x)`

output `A*a**2*d**4*x + B*b**2*e**4*x**8/8 + x**7*(A*b**2*e**4/7 + 2*B*a*b*e**4/7 + 4*B*b**2*d*e**3/7) + x**6*(A*a*b*e**4/3 + 2*A*b**2*d*e**3/3 + B*a**2*e**4/6 + 4*B*a*b*d*e**3/3 + B*b**2*d**2*e**2) + x**5*(A*a**2*e**4/5 + 8*A*a*b*d*e**3/5 + 6*A*b**2*d**2*e**2/5 + 4*B*a**2*d*e**3/5 + 12*B*a*b*d**2*e**2/5 + 4*B*b**2*d**3*e/5) + x**4*(A*a**2*d*e**3 + 3*A*a*b*d**2*e**2 + A*b**2*d**3*e + 3*B*a**2*d**2*e**2/2 + 2*B*a*b*d**3*e + B*b**2*d**4/4) + x**3*(2*A*a**2*d**2*e**2 + 8*A*a*b*d**3*e/3 + A*b**2*d**4/3 + 4*B*a**2*d**3*e/3 + 2*B*a*b*d**4/3) + x**2*(2*A*a**2*d**3*e + A*a*b*d**4 + B*a**2*d**4/2)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 304 vs. $2(112) = 224$.

Time = 0.03 (sec) , antiderivative size = 304, normalized size of antiderivative = 2.53

$$\begin{aligned} & \int (a + bx)^2(A + Bx)(d + ex)^4 dx \\ &= \frac{1}{8} Bb^2 e^4 x^8 + Aa^2 d^4 x + \frac{1}{7} (4 Bb^2 de^3 + (2 Bab + Ab^2) e^4) x^7 \\ &+ \frac{1}{6} (6 Bb^2 d^2 e^2 + 4 (2 Bab + Ab^2) de^3 + (Ba^2 + 2 Aab) e^4) x^6 \\ &+ \frac{1}{5} (4 Bb^2 d^3 e + Aa^2 e^4 + 6 (2 Bab + Ab^2) d^2 e^2 + 4 (Ba^2 + 2 Aab) de^3) x^5 \\ &+ \frac{1}{4} (Bb^2 d^4 + 4 Aa^2 de^3 + 4 (2 Bab + Ab^2) d^3 e + 6 (Ba^2 + 2 Aab) d^2 e^2) x^4 \\ &+ \frac{1}{3} (6 Aa^2 d^2 e^2 + (2 Bab + Ab^2) d^4 + 4 (Ba^2 + 2 Aab) d^3 e) x^3 \\ &+ \frac{1}{2} (4 Aa^2 d^3 e + (Ba^2 + 2 Aab) d^4) x^2 \end{aligned}$$

input `integrate((b*x+a)^2*(B*x+A)*(e*x+d)^4,x, algorithm="maxima")`

output `1/8*B*b^2*e^4*x^8 + A*a^2*d^4*x + 1/7*(4*B*b^2*d*e^3 + (2*B*a*b + A*b^2)*e^4)*x^7 + 1/6*(6*B*b^2*d^2*e^2 + 4*(2*B*a*b + A*b^2)*d*e^3 + (B*a^2 + 2*A*a*b)*e^4)*x^6 + 1/5*(4*B*b^2*d^3*e + A*a^2*e^4 + 6*(2*B*a*b + A*b^2)*d^2*e^2 + 4*(B*a^2 + 2*A*a*b)*d*e^3)*x^5 + 1/4*(B*b^2*d^4 + 4*A*a^2*d*e^3 + 4*(2*B*a*b + A*b^2)*d^3*e + 6*(B*a^2 + 2*A*a*b)*d^2*e^2)*x^4 + 1/3*(6*A*a^2*d^2*e^2 + (2*B*a*b + A*b^2)*d^4 + 4*(B*a^2 + 2*A*a*b)*d^3*e)*x^3 + 1/2*(4*A*a^2*d^3*e + (B*a^2 + 2*A*a*b)*d^4)*x^2`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 374 vs. $2(112) = 224$.

Time = 0.12 (sec) , antiderivative size = 374, normalized size of antiderivative = 3.12

$$\int (a + bx)^2(A + Bx)(d + ex)^4 dx = \frac{1}{8} Bb^2e^4x^8 + \frac{4}{7} Bb^2de^3x^7 + \frac{2}{7} Babe^4x^7 + \frac{1}{7} Ab^2e^4x^7$$

$$+ Bb^2d^2e^2x^6 + \frac{4}{3} Babde^3x^6 + \frac{2}{3} Ab^2de^3x^6$$

$$+ \frac{1}{6} Ba^2e^4x^6 + \frac{1}{3} Aabe^4x^6 + \frac{4}{5} Bb^2d^3ex^5$$

$$+ \frac{12}{5} Babd^2e^2x^5 + \frac{6}{5} Ab^2d^2e^2x^5 + \frac{4}{5} Ba^2de^3x^5$$

$$+ \frac{8}{5} Aabde^3x^5 + \frac{1}{5} Aa^2e^4x^5 + \frac{1}{4} Bb^2d^4x^4$$

$$+ 2 Babd^3ex^4 + Ab^2d^3ex^4 + \frac{3}{2} Ba^2d^2e^2x^4$$

$$+ 3 Aabd^2e^2x^4 + Aa^2de^3x^4 + \frac{2}{3} Babd^4x^3 + \frac{1}{3} Ab^2d^4x^3$$

$$+ \frac{4}{3} Ba^2d^3ex^3 + \frac{8}{3} Aabd^3ex^3 + 2 Aa^2d^2e^2x^3$$

$$+ \frac{1}{2} Ba^2d^4x^2 + Aabd^4x^2 + 2 Aa^2d^3ex^2 + Aa^2d^4x$$

input `integrate((b*x+a)^2*(B*x+A)*(e*x+d)^4,x, algorithm="giac")`

output `1/8*B*b^2*e^4*x^8 + 4/7*B*b^2*d*e^3*x^7 + 2/7*B*a*b*e^4*x^7 + 1/7*A*b^2*e^4*x^7 + B*b^2*d^2*e^2*x^6 + 4/3*B*a*b*d*e^3*x^6 + 2/3*A*b^2*d*e^3*x^6 + 1/6*B*a^2*e^4*x^6 + 1/3*A*a*b*e^4*x^6 + 4/5*B*b^2*d^3*e*x^5 + 12/5*B*a*b*d^2*e^2*x^5 + 6/5*A*b^2*d^2*e^2*x^5 + 4/5*B*a^2*d*e^3*x^5 + 8/5*A*a*b*d*e^3*x^5 + 1/5*A*a^2*e^4*x^5 + 1/4*B*b^2*d^4*x^4 + 2*B*a*b*d^3*e*x^4 + A*b^2*d^3*e*x^4 + 3/2*B*a^2*d^2*e^2*x^4 + 3*A*a*b*d^2*e^2*x^4 + A*a^2*d*e^3*x^4 + 2/3*B*a*b*d^4*x^3 + 1/3*A*b^2*d^4*x^3 + 4/3*B*a^2*d^3*e*x^3 + 8/3*A*a*b*d^3*e*x^3 + 2*A*a^2*d^2*e^2*x^3 + 1/2*B*a^2*d^4*x^2 + A*a*b*d^4*x^2 + 2*A*a^2*d^3*e*x^2 + A*a^2*d^4*x`

Mupad [B] (verification not implemented)

Time = 0.95 (sec) , antiderivative size = 305, normalized size of antiderivative = 2.54

$$\int (a + bx)^2(A + Bx)(d + ex)^4 dx = x^4 \left(\frac{3Ba^2d^2e^2}{2} + Aa^2de^3 + 2Babd^3e \right. \\ \left. + 3Aabd^2e^2 + \frac{Bb^2d^4}{4} + Ab^2d^3e \right) \\ + x^5 \left(\frac{4Ba^2de^3}{5} + \frac{Aa^2e^4}{5} + \frac{12Babd^2e^2}{5} \right. \\ \left. + \frac{8Aabde^3}{5} + \frac{4Bb^2d^3e}{5} + \frac{6Ab^2d^2e^2}{5} \right) \\ + x^3 \left(\frac{4Ba^2d^3e}{3} + 2Aa^2d^2e^2 + \frac{2Babd^4}{3} \right. \\ \left. + \frac{8Aabd^3e}{3} + \frac{Ab^2d^4}{3} \right) + x^6 \left(\frac{Ba^2e^4}{6} \right. \\ \left. + \frac{4Babde^3}{3} + \frac{Aabe^4}{3} + Bb^2d^2e^2 + \frac{2Ab^2de^3}{3} \right) \\ + Aa^2d^4x + \frac{ad^3x^2(4Aae + 2Abd + Bad)}{2} \\ + \frac{be^3x^7(Abe + 2Bae + 4Bbd)}{7} + \frac{Bb^2e^4x^8}{8}$$

input `int((A + B*x)*(a + b*x)^2*(d + e*x)^4,x)`output `x^4*((B*b^2*d^4)/4 + A*a^2*d*e^3 + A*b^2*d^3*e + (3*B*a^2*d^2*e^2)/2 + 2*B*a*b*d^3*e + 3*A*a*b*d^2*e^2) + x^5*((A*a^2*e^4)/5 + (4*B*a^2*d*e^3)/5 + (4*B*b^2*d^3*e)/5 + (6*A*b^2*d^2*e^2)/5 + (8*A*a*b*d^3*e)/5 + (12*B*a*b*d^2*e^2)/5) + x^3*((A*b^2*d^4)/3 + (2*B*a*b*d^4)/3 + (4*B*a^2*d^3*e)/3 + 2*A*a^2*d^2*e^2 + (8*A*a*b*d^3*e)/3) + x^6*((B*a^2*e^4)/6 + (A*a*b*e^4)/3 + (2*A*b^2*d*e^3)/3 + B*b^2*d^2*e^2 + (4*B*a*b*d^3*e)/3) + A*a^2*d^4*x + (a*d^3*x^2*(4*A*a*e + 2*A*b*d + B*a*d))/2 + (b*e^3*x^7*(A*b*e + 2*B*a*e + 4*B*b*d))/7 + (B*b^2*e^4*x^8)/8`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 247, normalized size of antiderivative = 2.06

$$\int (a + bx)^2 (A + Bx)(d + ex)^4 dx$$

$$= \frac{x(35b^3e^4x^7 + 120ab^2e^4x^6 + 160b^3de^3x^6 + 140a^2be^4x^5 + 560ab^2de^3x^5 + 280b^3d^2e^2x^5 + 56a^3e^4x^4 + 672a^2be^3x^3 + 224ab^2d^2e^2x^3 + 280a^2b^2de^2x^3 + 140a^2b^2e^3x^3 + 140a^2b^2e^4x^3 + 280ab^2d^2e^2x^3 + 280ab^2de^3x^3 + 120ab^2e^4x^3 + 70b^3d^2e^2x^3 + 224b^3de^2x^3 + 280b^3de^3x^3 + 160b^3de^4x^3 + 35b^3e^4x^3)}{280}$$

input `int((b*x+a)^2*(B*x+A)*(e*x+d)^4,x)`output `(x*(280*a**3*d**4 + 560*a**3*d**3*e*x + 560*a**3*d**2*e**2*x**2 + 280*a**3*d*e**3*x**3 + 56*a**3*e**4*x**4 + 420*a**2*b*d**4*x + 1120*a**2*b*d**3*e*x**2 + 1260*a**2*b*d**2*e**2*x**3 + 672*a**2*b*d*e**3*x**4 + 140*a**2*b*e**4*x**5 + 280*a*b**2*d**4*x**2 + 840*a*b**2*d**3*e*x**3 + 1008*a*b**2*d**2*e**2*x**4 + 560*a*b**2*d*e**3*x**5 + 120*a*b**2*e**4*x**6 + 70*b**3*d**4*x**3 + 224*b**3*d**3*e*x**4 + 280*b**3*d**2*e**2*x**5 + 160*b**3*d*e**3*x**6 + 35*b**3*e**4*x**7))/280`

3.13 $\int (a + bx)^2(A + Bx)(d + ex)^3 dx$

Optimal result	200
Mathematica [A] (verified)	201
Rubi [A] (verified)	201
Maple [B] (verified)	202
Fricas [B] (verification not implemented)	203
Sympy [B] (verification not implemented)	204
Maxima [B] (verification not implemented)	205
Giac [B] (verification not implemented)	206
Mupad [B] (verification not implemented)	207
Reduce [B] (verification not implemented)	207

Optimal result

Integrand size = 20, antiderivative size = 120

$$\int (a + bx)^2(A + Bx)(d + ex)^3 dx = -\frac{(bd - ae)^2(Bd - Ae)(d + ex)^4}{4e^4} + \frac{(bd - ae)(3bBd - 2Abe - aBe)(d + ex)^5}{5e^4} - \frac{b(3bBd - Abe - 2aBe)(d + ex)^6}{6e^4} + \frac{b^2B(d + ex)^7}{7e^4}$$

output

```
-1/4*(-a*e+b*d)^2*(-A*e+B*d)*(e*x+d)^4/e^4+1/5*(-a*e+b*d)*(-2*A*b*e-B*a*e+
3*B*b*d)*(e*x+d)^5/e^4-1/6*b*(-A*b*e-2*B*a*e+3*B*b*d)*(e*x+d)^6/e^4+1/7*b^
2*B*(e*x+d)^7/e^4
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.80

$$\int (a + bx)^2(A + Bx)(d + ex)^3 dx = a^2Ad^3x + \frac{1}{2}ad^2(2Abd + aBd + 3aAe)x^2$$

$$+ \frac{1}{3}d(aBd(2bd + 3ae) + A(b^2d^2 + 6abde + 3a^2e^2))x^3$$

$$+ \frac{1}{4}(6abde(Bd + Ae) + a^2e^2(3Bd + Ae) + b^2d^2(Bd + 3Ae))x^4$$

$$+ \frac{1}{5}e(a^2Be^2 + 3b^2d(Bd + Ae) + 2abe(3Bd + Ae))x^5$$

$$+ \frac{1}{6}be^2(3bBd + Abe + 2aBe)x^6 + \frac{1}{7}b^2Be^3x^7$$

input

```
Integrate[(a + b*x)^2*(A + B*x)*(d + e*x)^3,x]
```

output

```
a^2*A*d^3*x + (a*d^2*(2*A*b*d + a*B*d + 3*a*A*e)*x^2)/2 + (d*(a*B*d*(2*b*d + 3*a*e) + A*(b^2*d^2 + 6*a*b*d*e + 3*a^2*e^2))*x^3)/3 + ((6*a*b*d*e*(B*d + A*e) + a^2*e^2*(3*B*d + A*e) + b^2*d^2*(B*d + 3*A*e))*x^4)/4 + (e*(a^2*B*e^2 + 3*b^2*d*(B*d + A*e) + 2*a*b*e*(3*B*d + A*e))*x^5)/5 + (b*e^2*(3*b*B*d + A*b*e + 2*a*B*e)*x^6)/6 + (b^2*B*e^3*x^7)/7
```

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^2(A + Bx)(d + ex)^3 dx$$

↓ 86

$$\int \left(\frac{b(d+ex)^5(2aBe + Abe - 3bBd)}{e^3} + \frac{(d+ex)^4(ae - bd)(aBe + 2Abe - 3bBd)}{e^3} + \frac{(d+ex)^3(ae - bd)^2(Ae - B)}{e^3} \right) dx$$

↓ 2009

$$-\frac{b(d+ex)^6(-2aBe - Abe + 3bBd)}{6e^4} + \frac{(d+ex)^5(bd - ae)(-aBe - 2Abe + 3bBd)}{5e^4} - \frac{(d+ex)^4(bd - ae)^2(Bd - Ae)}{4e^4} + \frac{b^2B(d+ex)^7}{7e^4}$$

input `Int[(a + b*x)^2*(A + B*x)*(d + e*x)^3,x]`

output `-1/4*((b*d - a*e)^2*(B*d - A*e)*(d + e*x)^4)/e^4 + ((b*d - a*e)*(3*b*B*d - 2*A*b*e - a*B*e)*(d + e*x)^5)/(5*e^4) - (b*(3*b*B*d - A*b*e - 2*a*B*e)*(d + e*x)^6)/(6*e^4) + (b^2*B*(d + e*x)^7)/(7*e^4)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 236 vs. 2(112) = 224.

Time = 0.18 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.98

method	result
default	$\frac{b^2 B e^3 x^7}{7} + \frac{((b^2 A + 2abB)e^3 + 3b^2 B d e^2)x^6}{6} + \frac{((2abA + a^2 B)e^3 + 3(b^2 A + 2abB)d e^2 + 3b^2 B d^2 e)x^5}{5} + \frac{(a^2 A e^3 + 3(2abA + a^2 B)d e^2 + 3b^2 B d^2 e)x^4}{4} + \frac{(a^2 A e^3 + 3(2abA + a^2 B)d e^2 + 3b^2 B d^2 e)x^3}{3} + \frac{(a^2 A e^3 + 3(2abA + a^2 B)d e^2 + 3b^2 B d^2 e)x^2}{2} + \frac{(a^2 A e^3 + 3(2abA + a^2 B)d e^2 + 3b^2 B d^2 e)x}{1} + \frac{(a^2 A e^3 + 3(2abA + a^2 B)d e^2 + 3b^2 B d^2 e)}{0}$
norman	$\frac{b^2 B e^3 x^7}{7} + \left(\frac{1}{6} A b^2 e^3 + \frac{1}{3} B a b e^3 + \frac{1}{2} b^2 B d e^2\right) x^6 + \left(\frac{2}{5} A a b e^3 + \frac{3}{5} A b^2 d e^2 + \frac{1}{5} B a^2 e^3 + \frac{6}{5} B a b d e^2\right) x^5 + \left(\frac{1}{4} A a^2 e^3 + \frac{3}{4} A a b d e^2 + \frac{1}{2} B a^2 d e^2 + \frac{3}{4} B a b d^2 e\right) x^4 + \left(\frac{1}{3} A a^2 d e^2 + \frac{2}{3} A a b d^2 e + \frac{1}{3} B a^2 d^2 e\right) x^3 + \left(\frac{1}{2} A a^2 d^2 e + \frac{1}{2} A a b d^3\right) x^2 + \left(\frac{1}{2} A a^2 d^2 e + \frac{1}{2} A a b d^3\right) x + \frac{1}{2} A a^2 d^2 e + \frac{1}{2} A a b d^3$
gosper	$\frac{1}{7} b^2 B e^3 x^7 + \frac{1}{6} x^6 A b^2 e^3 + \frac{1}{3} x^6 B a b e^3 + \frac{1}{2} x^6 b^2 B d e^2 + \frac{2}{5} x^5 A a b e^3 + \frac{3}{5} x^5 A b^2 d e^2 + \frac{1}{5} x^5 B a^2 e^3 + \frac{6}{5} x^5 B a b d e^2 + \frac{1}{4} x^4 A a^2 e^3 + \frac{3}{4} x^4 A a b d e^2 + \frac{1}{2} x^4 B a^2 d e^2 + \frac{3}{4} x^4 B a b d^2 e + \frac{1}{3} x^3 A a^2 d e^2 + \frac{2}{3} x^3 A a b d^2 e + \frac{1}{3} x^3 B a^2 d^2 e + \frac{1}{2} x^2 A a^2 d^2 e + \frac{1}{2} x^2 A a b d^3 + \frac{1}{2} x A a^2 d^2 e + \frac{1}{2} x A a b d^3 + \frac{1}{2} A a^2 d^2 e + \frac{1}{2} A a b d^3$
risch	$\frac{1}{7} b^2 B e^3 x^7 + \frac{1}{6} x^6 A b^2 e^3 + \frac{1}{3} x^6 B a b e^3 + \frac{1}{2} x^6 b^2 B d e^2 + \frac{2}{5} x^5 A a b e^3 + \frac{3}{5} x^5 A b^2 d e^2 + \frac{1}{5} x^5 B a^2 e^3 + \frac{6}{5} x^5 B a b d e^2 + \frac{1}{4} x^4 A a^2 e^3 + \frac{3}{4} x^4 A a b d e^2 + \frac{1}{2} x^4 B a^2 d e^2 + \frac{3}{4} x^4 B a b d^2 e + \frac{1}{3} x^3 A a^2 d e^2 + \frac{2}{3} x^3 A a b d^2 e + \frac{1}{3} x^3 B a^2 d^2 e + \frac{1}{2} x^2 A a^2 d^2 e + \frac{1}{2} x^2 A a b d^3 + \frac{1}{2} x A a^2 d^2 e + \frac{1}{2} x A a b d^3 + \frac{1}{2} A a^2 d^2 e + \frac{1}{2} A a b d^3$
parallelrisch	$\frac{1}{7} b^2 B e^3 x^7 + \frac{1}{6} x^6 A b^2 e^3 + \frac{1}{3} x^6 B a b e^3 + \frac{1}{2} x^6 b^2 B d e^2 + \frac{2}{5} x^5 A a b e^3 + \frac{3}{5} x^5 A b^2 d e^2 + \frac{1}{5} x^5 B a^2 e^3 + \frac{6}{5} x^5 B a b d e^2 + \frac{1}{4} x^4 A a^2 e^3 + \frac{3}{4} x^4 A a b d e^2 + \frac{1}{2} x^4 B a^2 d e^2 + \frac{3}{4} x^4 B a b d^2 e + \frac{1}{3} x^3 A a^2 d e^2 + \frac{2}{3} x^3 A a b d^2 e + \frac{1}{3} x^3 B a^2 d^2 e + \frac{1}{2} x^2 A a^2 d^2 e + \frac{1}{2} x^2 A a b d^3 + \frac{1}{2} x A a^2 d^2 e + \frac{1}{2} x A a b d^3 + \frac{1}{2} A a^2 d^2 e + \frac{1}{2} A a b d^3$
orering	$\frac{x(60b^2 B e^3 x^6 + 70A b^2 e^3 x^5 + 140B a b e^3 x^5 + 210B b^2 d e^2 x^5 + 168A a b e^3 x^4 + 252A b^2 d e^2 x^4 + 84B a^2 e^3 x^4 + 504B a b d e^2 x^4 + 252A a^2 d e^2 x^3 + 350A a b d^2 e x^3 + 175B a^2 d^2 e x^3 + 105B a b d^3 e x^3 + 35A a^2 d^2 e x^2 + 70A a b d^3 x^2 + 35B a^2 d^2 e x^2 + 70B a b d^3 x^2 + 35A a^2 d^2 e x + 70A a b d^3 x + 35B a^2 d^2 e + 70B a b d^3)}{420}$

input `int((b*x+a)^2*(B*x+A)*(e*x+d)^3,x,method=_RETURNVERBOSE)`

output $\frac{1}{7} b^2 B e^3 x^7 + \frac{1}{6} x^6 (A b^2 e^3 + 3 B a b e^3 + 2 b^2 B d e^2) x^6 + \frac{1}{5} x^5 (2 A a b e^3 + 3 A b^2 d e^2 + B a^2 e^3 + 6 B a b d e^2) x^5 + \frac{1}{4} x^4 (A a^2 e^3 + 3 A a b d e^2 + 2 B a^2 d e^2 + 3 B a b d^2 e) x^4 + \frac{1}{3} x^3 (3 A a^2 d e^2 + 2 A a b d^2 e + B a^2 d^2 e) x^3 + \frac{1}{2} x^2 (2 A a^2 d^2 e + A a b d^3) x^2 + \frac{1}{2} x (A a^2 d^2 e + A a b d^3) x + \frac{1}{2} (A a^2 d^2 e + A a b d^3)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 236 vs. $2(112) = 224$.

Time = 0.08 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.97

$$\begin{aligned} & \int (a + bx)^2 (A + Bx)(d + ex)^3 dx \\ &= \frac{1}{7} B b^2 e^3 x^7 + A a^2 d^3 x + \frac{1}{6} (3 B b^2 d e^2 + (2 B a b + A b^2) e^3) x^6 \\ &+ \frac{1}{5} (3 B b^2 d^2 e + 3 (2 B a b + A b^2) d e^2 + (B a^2 + 2 A a b) e^3) x^5 \\ &+ \frac{1}{4} (B b^2 d^3 + A a^2 e^3 + 3 (2 B a b + A b^2) d^2 e + 3 (B a^2 + 2 A a b) d e^2) x^4 \\ &+ \frac{1}{3} (3 A a^2 d e^2 + (2 B a b + A b^2) d^3 + 3 (B a^2 + 2 A a b) d^2 e) x^3 \\ &+ \frac{1}{2} (3 A a^2 d^2 e + (B a^2 + 2 A a b) d^3) x^2 \end{aligned}$$

input `integrate((b*x+a)^2*(B*x+A)*(e*x+d)^3,x, algorithm="fricas")`

output

```
1/7*B*b^2*e^3*x^7 + A*a^2*d^3*x + 1/6*(3*B*b^2*d*e^2 + (2*B*a*b + A*b^2)*e^3)*x^6 + 1/5*(3*B*b^2*d^2*e + 3*(2*B*a*b + A*b^2)*d*e^2 + (B*a^2 + 2*A*a*b)*e^3)*x^5 + 1/4*(B*b^2*d^3 + A*a^2*e^3 + 3*(2*B*a*b + A*b^2)*d^2*e + 3*(B*a^2 + 2*A*a*b)*d*e^2)*x^4 + 1/3*(3*A*a^2*d*e^2 + (2*B*a*b + A*b^2)*d^3 + 3*(B*a^2 + 2*A*a*b)*d^2*e)*x^3 + 1/2*(3*A*a^2*d^2*e + (B*a^2 + 2*A*a*b)*d^3)*x^2
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 296 vs. $2(114) = 228$.

Time = 0.03 (sec) , antiderivative size = 296, normalized size of antiderivative = 2.47

$$\int (a + bx)^2(A + Bx)(d + ex)^3 dx = Aa^2d^3x + \frac{Bb^2e^3x^7}{7} + x^6 \left(\frac{Ab^2e^3}{6} + \frac{Babe^3}{3} + \frac{Bb^2de^2}{2} \right) + x^5 \cdot \left(\frac{2Aabe^3}{5} + \frac{3Ab^2de^2}{5} + \frac{Ba^2e^3}{5} + \frac{6Babde^2}{5} + \frac{3Bb^2d^2e}{5} \right) + x^4 \left(\frac{Aa^2e^3}{4} + \frac{3Aabde^2}{2} + \frac{3Ab^2d^2e}{4} + \frac{3Ba^2de^2}{4} + \frac{3Babd^2e}{2} + \frac{Bb^2d^3}{4} \right) + x^3 \left(Aa^2de^2 + 2Aabd^2e + \frac{Ab^2d^3}{3} + Ba^2d^2e + \frac{2Babd^3}{3} \right) + x^2 \cdot \left(\frac{3Aa^2d^2e}{2} + Aabd^3 + \frac{Ba^2d^3}{2} \right)$$

input

```
integrate((b*x+a)**2*(B*x+A)*(e*x+d)**3,x)
```

output

```
A*a**2*d**3*x + B*b**2*e**3*x**7/7 + x**6*(A*b**2*e**3/6 + B*a*b*e**3/3 + B*b**2*d*e**2/2) + x**5*(2*A*a*b*e**3/5 + 3*A*b**2*d*e**2/5 + B*a**2*e**3/5 + 6*B*a*b*d*e**2/5 + 3*B*b**2*d**2*e/5) + x**4*(A*a**2*e**3/4 + 3*A*a*b*d*e**2/2 + 3*A*b**2*d**2*e/4 + 3*B*a**2*d*e**2/4 + 3*B*a*b*d**2*e/2 + B*b**2*d**3/4) + x**3*(A*a**2*d*e**2 + 2*A*a*b*d**2*e + A*b**2*d**3/3 + B*a**2*d**2*e + 2*B*a*b*d**3/3) + x**2*(3*A*a**2*d**2*e/2 + A*a*b*d**3 + B*a**2*d**3/2)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 236 vs. $2(112) = 224$.

Time = 0.03 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.97

$$\begin{aligned} & \int (a + bx)^2 (A + Bx)(d + ex)^3 dx \\ &= \frac{1}{7} Bb^2 e^3 x^7 + Aa^2 d^3 x + \frac{1}{6} (3 Bb^2 de^2 + (2 Bab + Ab^2) e^3) x^6 \\ & \quad + \frac{1}{5} (3 Bb^2 d^2 e + 3 (2 Bab + Ab^2) de^2 + (Ba^2 + 2 Aab) e^3) x^5 \\ & \quad + \frac{1}{4} (Bb^2 d^3 + Aa^2 e^3 + 3 (2 Bab + Ab^2) d^2 e + 3 (Ba^2 + 2 Aab) de^2) x^4 \\ & \quad + \frac{1}{3} (3 Aa^2 de^2 + (2 Bab + Ab^2) d^3 + 3 (Ba^2 + 2 Aab) d^2 e) x^3 \\ & \quad + \frac{1}{2} (3 Aa^2 d^2 e + (Ba^2 + 2 Aab) d^3) x^2 \end{aligned}$$

input

```
integrate((b*x+a)^2*(B*x+A)*(e*x+d)^3,x, algorithm="maxima")
```

output

```
1/7*B*b^2*e^3*x^7 + A*a^2*d^3*x + 1/6*(3*B*b^2*d*e^2 + (2*B*a*b + A*b^2)*e^3)*x^6 + 1/5*(3*B*b^2*d^2*e + 3*(2*B*a*b + A*b^2)*d*e^2 + (B*a^2 + 2*A*a*b)*e^3)*x^5 + 1/4*(B*b^2*d^3 + A*a^2*e^3 + 3*(2*B*a*b + A*b^2)*d^2*e + 3*(B*a^2 + 2*A*a*b)*d*e^2)*x^4 + 1/3*(3*A*a^2*d*e^2 + (2*B*a*b + A*b^2)*d^3 + 3*(B*a^2 + 2*A*a*b)*d^2*e)*x^3 + 1/2*(3*A*a^2*d^2*e + (B*a^2 + 2*A*a*b)*d^3)*x^2
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 287 vs. $2(112) = 224$.

Time = 0.12 (sec) , antiderivative size = 287, normalized size of antiderivative = 2.39

$$\int (a + bx)^2(A + Bx)(d + ex)^3 dx = \frac{1}{7} Bb^2e^3x^7 + \frac{1}{2} Bb^2de^2x^6 + \frac{1}{3} Babe^3x^6 + \frac{1}{6} Ab^2e^3x^6$$

$$+ \frac{3}{5} Bb^2d^2ex^5 + \frac{6}{5} Babde^2x^5 + \frac{3}{5} Ab^2de^2x^5$$

$$+ \frac{1}{5} Ba^2e^3x^5 + \frac{2}{5} Aabe^3x^5 + \frac{1}{4} Bb^2d^3x^4$$

$$+ \frac{3}{2} Babd^2ex^4 + \frac{3}{4} Ab^2d^2ex^4 + \frac{3}{4} Ba^2de^2x^4$$

$$+ \frac{3}{2} Aabde^2x^4 + \frac{1}{4} Aa^2e^3x^4 + \frac{2}{3} Babd^3x^3$$

$$+ \frac{1}{3} Ab^2d^3x^3 + Ba^2d^2ex^3 + 2Aabd^2ex^3 + Aa^2de^2x^3$$

$$+ \frac{1}{2} Ba^2d^3x^2 + Aabd^3x^2 + \frac{3}{2} Aa^2d^2ex^2 + Aa^2d^3x$$

input `integrate((b*x+a)^2*(B*x+A)*(e*x+d)^3,x, algorithm="giac")`

output `1/7*B*b^2*e^3*x^7 + 1/2*B*b^2*d*e^2*x^6 + 1/3*B*a*b*e^3*x^6 + 1/6*A*b^2*e^3*x^6 + 3/5*B*b^2*d^2*e*x^5 + 6/5*B*a*b*d*e^2*x^5 + 3/5*A*b^2*d*e^2*x^5 + 1/5*B*a^2*e^3*x^5 + 2/5*A*a*b*e^3*x^5 + 1/4*B*b^2*d^3*x^4 + 3/2*B*a*b*d^2*e*x^4 + 3/4*A*b^2*d^2*e*x^4 + 3/4*B*a^2*d*e^2*x^4 + 3/2*A*a*b*d*e^2*x^4 + 1/4*A*a^2*e^3*x^4 + 2/3*B*a*b*d^3*x^3 + 1/3*A*b^2*d^3*x^3 + B*a^2*d^2*e*x^3 + 2*A*a*b*d^2*e*x^3 + A*a^2*d*e^2*x^3 + 1/2*B*a^2*d^3*x^2 + A*a*b*d^3*x^2 + 3/2*A*a^2*d^2*e*x^2 + A*a^2*d^3*x`

output

```
(x*(140*a**3*d**3 + 210*a**3*d**2*e*x + 140*a**3*d*e**2*x**2 + 35*a**3*e**3*x**3 + 210*a**2*b*d**3*x + 420*a**2*b*d**2*e*x**2 + 315*a**2*b*d*e**2*x**3 + 84*a**2*b*e**3*x**4 + 140*a*b**2*d**3*x**2 + 315*a*b**2*d**2*e*x**3 + 252*a*b**2*d*e**2*x**4 + 70*a*b**2*e**3*x**5 + 35*b**3*d**3*x**3 + 84*b**3*d**2*e*x**4 + 70*b**3*d*e**2*x**5 + 20*b**3*e**3*x**6))/140
```

3.14 $\int (a + bx)^2 (A + Bx)(d + ex)^2 dx$

Optimal result	209
Mathematica [A] (verified)	209
Rubi [A] (verified)	210
Maple [A] (verified)	211
Fricas [A] (verification not implemented)	212
Sympy [A] (verification not implemented)	212
Maxima [A] (verification not implemented)	213
Giac [A] (verification not implemented)	214
Mupad [B] (verification not implemented)	214
Reduce [B] (verification not implemented)	215

Optimal result

Integrand size = 20, antiderivative size = 118

$$\int (a + bx)^2 (A + Bx)(d + ex)^2 dx = \frac{(Ab - aB)(bd - ae)^2 (a + bx)^3}{3b^4} + \frac{(bd - ae)(bBd + 2Abe - 3aBe)(a + bx)^4}{4b^4} + \frac{e(2bBd + Abe - 3aBe)(a + bx)^5}{5b^4} + \frac{Be^2(a + bx)^6}{6b^4}$$

output

```
1/3*(A*b-B*a)*(-a*e+b*d)^2*(b*x+a)^3/b^4+1/4*(-a*e+b*d)*(2*A*b*e-3*B*a*e+B
*b*d)*(b*x+a)^4/b^4+1/5*e*(A*b*e-3*B*a*e+2*B*b*d)*(b*x+a)^5/b^4+1/6*B*e^2*
(b*x+a)^6/b^4
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.33

$$\int (a + bx)^2 (A + Bx)(d + ex)^2 dx = a^2 Ad^2 x + \frac{1}{2} ad (aBd + 2A(bd + ae)) x^2 + \frac{1}{3} (2aBd(bd + ae) + A(b^2 d^2 + 4abde + a^2 e^2)) x^3 + \frac{1}{4} (a^2 Be^2 + 2abe(2Bd + Ae) + b^2 d(Bd + 2Ae)) x^4 + \frac{1}{5} be(2bBd + Abe + 2aBe) x^5 + \frac{1}{6} b^2 Be^2 x^6$$

input `Integrate[(a + b*x)^2*(A + B*x)*(d + e*x)^2,x]`

output `a^2*A*d^2*x + (a*d*(a*B*d + 2*A*(b*d + a*e))*x^2)/2 + ((2*a*B*d*(b*d + a*e) + A*(b^2*d^2 + 4*a*b*d*e + a^2*e^2))*x^3)/3 + ((a^2*B*e^2 + 2*a*b*e*(2*B*d + A*e) + b^2*d*(B*d + 2*A*e))*x^4)/4 + (b*e*(2*b*B*d + A*b*e + 2*a*B*e)*x^5)/5 + (b^2*B*e^2*x^6)/6`

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^2(A + Bx)(d + ex)^2 dx$$

$$\downarrow 86$$

$$\int \left(\frac{e(a + bx)^4(-3aBe + Abe + 2bBd)}{b^3} + \frac{(a + bx)^3(bd - ae)(-3aBe + 2Abe + bBd)}{b^3} + \frac{(a + bx)^2(Ab - aB)(bd - ae)^2}{b^3} \right) dx$$

$$\downarrow 2009$$

$$\frac{e(a + bx)^5(-3aBe + Abe + 2bBd)}{5b^4} + \frac{(a + bx)^4(bd - ae)(-3aBe + 2Abe + bBd)}{4b^4} + \frac{(a + bx)^3(Ab - aB)(bd - ae)^2}{3b^4} + \frac{Be^2(a + bx)^6}{6b^4}$$

input `Int[(a + b*x)^2*(A + B*x)*(d + e*x)^2,x]`

output `((A*b - a*B)*(b*d - a*e)^2*(a + b*x)^3)/(3*b^4) + ((b*d - a*e)*(b*B*d + 2*A*b*e - 3*a*B*e)*(a + b*x)^4)/(4*b^4) + (e*(2*b*B*d + A*b*e - 3*a*B*e)*(a + b*x)^5)/(5*b^4) + (B*e^2*(a + b*x)^6)/(6*b^4)`

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.43

method	result
default	$\frac{b^2 B e^2 x^6}{6} + \frac{((b^2 A + 2abB)e^2 + 2b^2 Bde)x^5}{5} + \frac{((2abA + a^2 B)e^2 + 2(b^2 A + 2abB)de + b^2 B d^2)x^4}{4} + \frac{(a^2 A e^2 + 2(2abA + a^2 B)de + b^2 B d^2)x^3}{3} + \frac{(a^2 A e^2 + 2(2abA + a^2 B)de + b^2 B d^2)x^2}{2} + \frac{(a^2 A e^2 + 2(2abA + a^2 B)de + b^2 B d^2)x}{1} + \frac{a^2 A e^2 + 2(2abA + a^2 B)de + b^2 B d^2}{0}$
norman	$\frac{b^2 B e^2 x^6}{6} + (\frac{1}{5} A b^2 e^2 + \frac{2}{5} B a b e^2 + \frac{2}{5} b^2 B d e) x^5 + (\frac{1}{2} A a b e^2 + \frac{1}{2} A b^2 d e + \frac{1}{4} B a^2 e^2 + B a b d e + \frac{1}{4} B b^2 d^2) x^4 + (\frac{1}{3} a^2 A e^2 + \frac{2}{3} a^2 B d e + \frac{1}{3} a^2 B d^2) x^3 + (\frac{1}{2} a^2 A e^2 + \frac{1}{2} a^2 B d e + \frac{1}{2} a^2 B d^2) x^2 + (a^2 A e^2 + a^2 B d e + a^2 B d^2) x + a^2 A e^2 + a^2 B d e + a^2 B d^2$
gosper	$\frac{1}{6} b^2 B e^2 x^6 + \frac{1}{5} x^5 A b^2 e^2 + \frac{2}{5} x^5 B a b e^2 + \frac{2}{5} x^5 b^2 B d e + \frac{1}{2} x^4 A a b e^2 + \frac{1}{2} x^4 A b^2 d e + \frac{1}{4} x^4 B a^2 e^2 + \frac{1}{4} x^4 B a b d e + \frac{1}{4} x^4 B b^2 d^2 + \frac{1}{3} x^3 a^2 A e^2 + \frac{2}{3} x^3 a^2 B d e + \frac{1}{3} x^3 a^2 B d^2 + \frac{1}{2} x^2 a^2 A e^2 + \frac{1}{2} x^2 a^2 B d e + \frac{1}{2} x^2 a^2 B d^2 + x a^2 A e^2 + x a^2 B d e + x a^2 B d^2 + a^2 A e^2 + a^2 B d e + a^2 B d^2$
risch	$\frac{1}{6} b^2 B e^2 x^6 + \frac{1}{5} x^5 A b^2 e^2 + \frac{2}{5} x^5 B a b e^2 + \frac{2}{5} x^5 b^2 B d e + \frac{1}{2} x^4 A a b e^2 + \frac{1}{2} x^4 A b^2 d e + \frac{1}{4} x^4 B a^2 e^2 + \frac{1}{4} x^4 B a b d e + \frac{1}{4} x^4 B b^2 d^2 + \frac{1}{3} x^3 a^2 A e^2 + \frac{2}{3} x^3 a^2 B d e + \frac{1}{3} x^3 a^2 B d^2 + \frac{1}{2} x^2 a^2 A e^2 + \frac{1}{2} x^2 a^2 B d e + \frac{1}{2} x^2 a^2 B d^2 + x a^2 A e^2 + x a^2 B d e + x a^2 B d^2 + a^2 A e^2 + a^2 B d e + a^2 B d^2$
parallelrisc	$\frac{1}{6} b^2 B e^2 x^6 + \frac{1}{5} x^5 A b^2 e^2 + \frac{2}{5} x^5 B a b e^2 + \frac{2}{5} x^5 b^2 B d e + \frac{1}{2} x^4 A a b e^2 + \frac{1}{2} x^4 A b^2 d e + \frac{1}{4} x^4 B a^2 e^2 + \frac{1}{4} x^4 B a b d e + \frac{1}{4} x^4 B b^2 d^2 + \frac{1}{3} x^3 a^2 A e^2 + \frac{2}{3} x^3 a^2 B d e + \frac{1}{3} x^3 a^2 B d^2 + \frac{1}{2} x^2 a^2 A e^2 + \frac{1}{2} x^2 a^2 B d e + \frac{1}{2} x^2 a^2 B d^2 + x a^2 A e^2 + x a^2 B d e + x a^2 B d^2 + a^2 A e^2 + a^2 B d e + a^2 B d^2$
orering	$\frac{x(10b^2 B e^2 x^5 + 12A b^2 e^2 x^4 + 24B a b e^2 x^4 + 24B b^2 d e x^4 + 30A a b e^2 x^3 + 30A b^2 d e x^3 + 15B a^2 e^2 x^3 + 60B a b d e x^3 + 15B b^2 d^2 x^3 + 10a^2 A e^2 x^2 + 20a^2 B d e x^2 + 10a^2 B d^2 x^2 + 6a^2 A e^2 x + 6a^2 B d e x + 6a^2 B d^2 x + a^2 A e^2 + a^2 B d e + a^2 B d^2)}{60}$

```
input int((b*x+a)^2*(B*x+A)*(e*x+d)^2,x,method=_RETURNVERBOSE)
```

```
output 1/6*b^2*B*e^2*x^6+1/5*((A*b^2+2*B*a*b)*e^2+2*b^2*B*d*e)*x^5+1/4*((2*A*a*b+B*a^2)*e^2+2*(A*b^2+2*B*a*b)*d*e+b^2*B*d^2)*x^4+1/3*(a^2*A*e^2+2*(2*A*a*b+B*a^2)*d*e+(A*b^2+2*B*a*b)*d^2)*x^3+1/2*(2*a^2*A*d*e+(2*A*a*b+B*a^2)*d^2)*x^2+a^2*A*d^2*x
```


Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.42

$$\int (a + bx)^2(A + Bx)(d + ex)^2 dx$$

$$= \frac{1}{6} Bb^2e^2x^6 + Aa^2d^2x + \frac{1}{5} (2Bb^2de + (2Bab + Ab^2)e^2)x^5$$

$$+ \frac{1}{4} (Bb^2d^2 + 2(2Bab + Ab^2)de + (Ba^2 + 2Aab)e^2)x^4$$

$$+ \frac{1}{3} (Aa^2e^2 + (2Bab + Ab^2)d^2 + 2(Ba^2 + 2Aab)de)x^3$$

$$+ \frac{1}{2} (2Aa^2de + (Ba^2 + 2Aab)d^2)x^2$$

input `integrate((b*x+a)^2*(B*x+A)*(e*x+d)^2,x, algorithm="fricas")`output `1/6*B*b^2*e^2*x^6 + A*a^2*d^2*x + 1/5*(2*B*b^2*d*e + (2*B*a*b + A*b^2)*e^2)*x^5 + 1/4*(B*b^2*d^2 + 2*(2*B*a*b + A*b^2)*d*e + (B*a^2 + 2*A*a*b)*e^2)*x^4 + 1/3*(A*a^2*e^2 + (2*B*a*b + A*b^2)*d^2 + 2*(B*a^2 + 2*A*a*b)*d*e)*x^3 + 1/2*(2*A*a^2*d*e + (B*a^2 + 2*A*a*b)*d^2)*x^2`**Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.71

$$\int (a + bx)^2(A + Bx)(d + ex)^2 dx = Aa^2d^2x + \frac{Bb^2e^2x^6}{6} + x^5 \left(\frac{Ab^2e^2}{5} + \frac{2Babe^2}{5} + \frac{2Bb^2de}{5} \right)$$

$$+ x^4 \left(\frac{Aabe^2}{2} + \frac{Ab^2de}{2} + \frac{Ba^2e^2}{4} + Babde + \frac{Bb^2d^2}{4} \right)$$

$$+ x^3 \left(\frac{Aa^2e^2}{3} + \frac{4Aabde}{3} + \frac{Ab^2d^2}{3} + \frac{2Ba^2de}{3} + \frac{2Babd^2}{3} \right) + x^2 \left(Aa^2de + Aabd^2 + \frac{Ba^2d^2}{2} \right)$$

input `integrate((b*x+a)**2*(B*x+A)*(e*x+d)**2,x)`

output

```
A*a**2*d**2*x + B*b**2*e**2*x**6/6 + x**5*(A*b**2*e**2/5 + 2*B*a*b*e**2/5
+ 2*B*b**2*d*e/5) + x**4*(A*a*b*e**2/2 + A*b**2*d*e/2 + B*a**2*e**2/4 + B*
a*b*d*e + B*b**2*d**2/4) + x**3*(A*a**2*e**2/3 + 4*A*a*b*d*e/3 + A*b**2*d*
*2/3 + 2*B*a**2*d*e/3 + 2*B*a*b*d**2/3) + x**2*(A*a**2*d*e + A*a*b*d**2 +
B*a**2*d**2/2)
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.42

$$\int (a + bx)^2 (A + Bx)(d + ex)^2 dx$$

$$= \frac{1}{6} Bb^2 e^2 x^6 + Aa^2 d^2 x + \frac{1}{5} (2 Bb^2 de + (2 Bab + Ab^2) e^2) x^5$$

$$+ \frac{1}{4} (Bb^2 d^2 + 2 (2 Bab + Ab^2) de + (Ba^2 + 2 Aab) e^2) x^4$$

$$+ \frac{1}{3} (Aa^2 e^2 + (2 Bab + Ab^2) d^2 + 2 (Ba^2 + 2 Aab) de) x^3$$

$$+ \frac{1}{2} (2 Aa^2 de + (Ba^2 + 2 Aab) d^2) x^2$$

input

```
integrate((b*x+a)^2*(B*x+A)*(e*x+d)^2,x, algorithm="maxima")
```

output

```
1/6*B*b^2*e^2*x^6 + A*a^2*d^2*x + 1/5*(2*B*b^2*d*e + (2*B*a*b + A*b^2)*e^2
)*x^5 + 1/4*(B*b^2*d^2 + 2*(2*B*a*b + A*b^2)*d*e + (B*a^2 + 2*A*a*b)*e^2)*
x^4 + 1/3*(A*a^2*e^2 + (2*B*a*b + A*b^2)*d^2 + 2*(B*a^2 + 2*A*a*b)*d*e)*x^
3 + 1/2*(2*A*a^2*d*e + (B*a^2 + 2*A*a*b)*d^2)*x^2
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.69

$$\int (a + bx)^2(A + Bx)(d + ex)^2 dx = \frac{1}{6} Bb^2e^2x^6 + \frac{2}{5} Bb^2dex^5 + \frac{2}{5} Babe^2x^5 + \frac{1}{5} Ab^2e^2x^5$$

$$+ \frac{1}{4} Bb^2d^2x^4 + Babdex^4 + \frac{1}{2} Ab^2dex^4 + \frac{1}{4} Ba^2e^2x^4$$

$$+ \frac{1}{2} Aabe^2x^4 + \frac{2}{3} Babd^2x^3 + \frac{1}{3} Ab^2d^2x^3$$

$$+ \frac{2}{3} Ba^2dex^3 + \frac{4}{3} Aabdex^3 + \frac{1}{3} Aa^2e^2x^3$$

$$+ \frac{1}{2} Ba^2d^2x^2 + Aabd^2x^2 + Aa^2dex^2 + Aa^2d^2x$$

input `integrate((b*x+a)^2*(B*x+A)*(e*x+d)^2,x, algorithm="giac")`

output `1/6*B*b^2*e^2*x^6 + 2/5*B*b^2*d*e*x^5 + 2/5*B*a*b*e^2*x^5 + 1/5*A*b^2*e^2*x^5 + 1/4*B*b^2*d^2*x^4 + B*a*b*d*e*x^4 + 1/2*A*b^2*d*e*x^4 + 1/4*B*a^2*e^2*x^4 + 1/2*A*a*b*e^2*x^4 + 2/3*B*a*b*d^2*x^3 + 1/3*A*b^2*d^2*x^3 + 2/3*B*a^2*d*e*x^3 + 4/3*A*a*b*d*e*x^3 + 1/3*A*a^2*e^2*x^3 + 1/2*B*a^2*d^2*x^2 + A*a*b*d^2*x^2 + A*a^2*d*e*x^2 + A*a^2*d^2*x`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.33

$$\int (a + bx)^2(A + Bx)(d + ex)^2 dx = x^3 \left(\frac{2Ba^2de}{3} + \frac{Aa^2e^2}{3} + \frac{2Babd^2}{3} + \frac{4Aabde}{3} + \frac{Ab^2d^2}{3} \right)$$

$$+ x^4 \left(\frac{Ba^2e^2}{4} + Babde + \frac{Aabe^2}{2} + \frac{Bb^2d^2}{4} + \frac{Ab^2de}{2} \right) + \frac{adx^2(2Aae + 2Abd + Bad)}{2}$$

$$+ \frac{bex^5(Abe + 2Bae + 2Bbd)}{5}$$

$$+ Aa^2d^2x + \frac{Bb^2e^2x^6}{6}$$

input `int((A + B*x)*(a + b*x)^2*(d + e*x)^2,x)`

output `x^3*((A*a^2*e^2)/3 + (A*b^2*d^2)/3 + (2*B*a*b*d^2)/3 + (2*B*a^2*d*e)/3 + (4*A*a*b*d*e)/3) + x^4*((B*a^2*e^2)/4 + (B*b^2*d^2)/4 + (A*a*b*e^2)/2 + (A*b^2*d*e)/2 + B*a*b*d*e) + (a*d*x^2*(2*A*a*e + 2*A*b*d + B*a*d))/2 + (b*e*x^5*(A*b*e + 2*B*a*e + 2*B*b*d))/5 + A*a^2*d^2*x + (B*b^2*e^2*x^6)/6`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.11

$$\int (a + bx)^2 (A + Bx)(d + ex)^2 dx$$

$$= \frac{x(10b^3e^2x^5 + 36ab^2e^2x^4 + 24b^3dex^4 + 45a^2be^2x^3 + 90ab^2dex^3 + 15b^3d^2x^3 + 20a^3e^2x^2 + 120a^2bde x^2 + 10ab^3e^2x^2 + 10b^3e^2x^2)}{60}$$

input `int((b*x+a)^2*(B*x+A)*(e*x+d)^2,x)`

output `(x*(60*a**3*d**2 + 60*a**3*d*e*x + 20*a**3*e**2*x**2 + 90*a**2*b*d**2*x + 120*a**2*b*d*e*x**2 + 45*a**2*b*e**2*x**3 + 60*a*b**2*d**2*x**2 + 90*a*b**2*d*e*x**3 + 36*a*b**2*e**2*x**4 + 15*b**3*d**2*x**3 + 24*b**3*d*e*x**4 + 10*b**3*e**2*x**5))/60`

3.15 $\int (a + bx)^2(A + Bx)(d + ex) dx$

Optimal result	216
Mathematica [A] (verified)	216
Rubi [A] (verified)	217
Maple [A] (verified)	218
Fricas [A] (verification not implemented)	219
Sympy [A] (verification not implemented)	219
Maxima [A] (verification not implemented)	220
Giac [A] (verification not implemented)	220
Mupad [B] (verification not implemented)	221
Reduce [B] (verification not implemented)	221

Optimal result

Integrand size = 18, antiderivative size = 75

$$\int (a + bx)^2(A + Bx)(d + ex) dx = \frac{(Ab - aB)(bd - ae)(a + bx)^3}{3b^3} + \frac{(bBd + Abe - 2aBe)(a + bx)^4}{4b^3} + \frac{Be(a + bx)^5}{5b^3}$$

output

```
1/3*(A*b-B*a)*(-a*e+b*d)*(b*x+a)^3/b^3+1/4*(A*b*e-2*B*a*e+B*b*d)*(b*x+a)^4/b^3+1/5*B*e*(b*x+a)^5/b^3
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.28

$$\int (a + bx)^2(A + Bx)(d + ex) dx = a^2Adx + \frac{1}{2}a(2Abd + aBd + aAe)x^2 + \frac{1}{3}(Ab^2d + 2abBd + 2aAbe + a^2Be)x^3 + \frac{1}{4}b(bBd + Abe + 2aBe)x^4 + \frac{1}{5}b^2Bex^5$$

input

```
Integrate[(a + b*x)^2*(A + B*x)*(d + e*x), x]
```

output

$$a^2 A d x + (a(2 A b d + a B d + a A e) x^2) / 2 + ((A b^2 d + 2 a b B d + 2 a A b e + a^2 B e) x^3) / 3 + (b(b B d + A b e + 2 a B e) x^4) / 4 + (b^2 B e x^5) / 5$$

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^2 (A + Bx)(d + ex) dx$$

$$\downarrow 86$$

$$\int \left(\frac{(a + bx)^3 (-2aBe + Abe + bBd)}{b^2} + \frac{(a + bx)^2 (Ab - aB)(bd - ae)}{b^2} + \frac{Be(a + bx)^4}{b^2} \right) dx$$

$$\downarrow 2009$$

$$\frac{(a + bx)^4 (-2aBe + Abe + bBd)}{4b^3} + \frac{(a + bx)^3 (Ab - aB)(bd - ae)}{3b^3} + \frac{Be(a + bx)^5}{5b^3}$$

input

```
Int[(a + b*x)^2*(A + B*x)*(d + e*x), x]
```

output

```
((A*b - a*B)*(b*d - a*e)*(a + b*x)^3)/(3*b^3) + ((b*B*d + A*b*e - 2*a*B*e)
*(a + b*x)^4)/(4*b^3) + (B*e*(a + b*x)^5)/(5*b^3)
```

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0]) || GeQ[n + p + 1, 0]) || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f]))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.32

method	result
norman	$\frac{b^2 B e x^5}{5} + \left(\frac{1}{4} A b^2 e + \frac{1}{2} B a b e + \frac{1}{4} b^2 B d\right) x^4 + \left(\frac{2}{3} A a b e + \frac{1}{3} A b^2 d + \frac{1}{3} B a^2 e + \frac{2}{3} B a b d\right) x^3 + \left(\frac{1}{2} a^2 A d e + \frac{1}{2} a^2 B d e\right) x^2 + a^2 A d x$
default	$\frac{b^2 B e x^5}{5} + \frac{((b^2 A + 2 a b B) e + b^2 B d) x^4}{4} + \frac{((2 a b A + a^2 B) e + (b^2 A + 2 a b B) d) x^3}{3} + \frac{(a^2 A e + (2 a b A + a^2 B) d) x^2}{2} + a^2 A d x$
orering	$\frac{x(12 b^2 B e x^4 + 15 A b^2 e x^3 + 30 B a b e x^3 + 15 B b^2 d x^3 + 40 A a b e x^2 + 20 A b^2 d x^2 + 20 B a^2 e x^2 + 40 B a b d x^2 + 30 A a^2 e x + 60 A a b d x + 60 A a^2 d)}{60}$
gosper	$\frac{1}{5} b^2 B e x^5 + \frac{1}{4} x^4 A b^2 e + \frac{1}{2} x^4 B a b e + \frac{1}{4} x^4 b^2 B d + \frac{2}{3} x^3 A a b e + \frac{1}{3} x^3 A b^2 d + \frac{1}{3} x^3 B a^2 e + \frac{2}{3} x^3 B a b d$
risch	$\frac{1}{5} b^2 B e x^5 + \frac{1}{4} x^4 A b^2 e + \frac{1}{2} x^4 B a b e + \frac{1}{4} x^4 b^2 B d + \frac{2}{3} x^3 A a b e + \frac{1}{3} x^3 A b^2 d + \frac{1}{3} x^3 B a^2 e + \frac{2}{3} x^3 B a b d$
parallelrisch	$\frac{1}{5} b^2 B e x^5 + \frac{1}{4} x^4 A b^2 e + \frac{1}{2} x^4 B a b e + \frac{1}{4} x^4 b^2 B d + \frac{2}{3} x^3 A a b e + \frac{1}{3} x^3 A b^2 d + \frac{1}{3} x^3 B a^2 e + \frac{2}{3} x^3 B a b d$

```
input int((b*x+a)^2*(B*x+A)*(e*x+d),x,method=_RETURNVERBOSE)
```

```
output 1/5*b^2*B*e*x^5+(1/4*A*b^2*e+1/2*B*a*b*e+1/4*b^2*B*d)*x^4+(2/3*A*a*b*e+1/3*A*b^2*d+1/3*B*a^2*e+2/3*B*a*b*d)*x^3+(1/2*a^2*A*e+A*a*b*d+1/2*B*a^2*d)*x^2+a^2*A*d*x
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.33

$$\int (a + bx)^2(A + Bx)(d + ex) dx = \frac{1}{5} Bb^2ex^5 + Aa^2dx + \frac{1}{4} (Bb^2d + (2 Bab + Ab^2)e)x^4$$

$$+ \frac{1}{3} ((2 Bab + Ab^2)d + (Ba^2 + 2 Aab)e)x^3$$

$$+ \frac{1}{2} (Aa^2e + (Ba^2 + 2 Aab)d)x^2$$

input `integrate((b*x+a)^2*(B*x+A)*(e*x+d),x, algorithm="fricas")`output `1/5*B*b^2*e*x^5 + A*a^2*d*x + 1/4*(B*b^2*d + (2*B*a*b + A*b^2)*e)*x^4 + 1/3*((2*B*a*b + A*b^2)*d + (B*a^2 + 2*A*a*b)*e)*x^3 + 1/2*(A*a^2*e + (B*a^2 + 2*A*a*b)*d)*x^2`**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.55

$$\int (a + bx)^2(A + Bx)(d + ex) dx = Aa^2dx + \frac{Bb^2ex^5}{5} + x^4 \left(\frac{Ab^2e}{4} + \frac{Babe}{2} + \frac{Bb^2d}{4} \right)$$

$$+ x^3 \cdot \left(\frac{2Aabe}{3} + \frac{Ab^2d}{3} + \frac{Ba^2e}{3} + \frac{2Babd}{3} \right)$$

$$+ x^2 \left(\frac{Aa^2e}{2} + Aabd + \frac{Ba^2d}{2} \right)$$

input `integrate((b*x+a)**2*(B*x+A)*(e*x+d),x)`output `A*a**2*d*x + B*b**2*e*x**5/5 + x**4*(A*b**2*e/4 + B*a*b*e/2 + B*b**2*d/4) + x**3*(2*A*a*b*e/3 + A*b**2*d/3 + B*a**2*e/3 + 2*B*a*b*d/3) + x**2*(A*a**2*e/2 + A*a*b*d + B*a**2*d/2)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.33

$$\int (a + bx)^2(A + Bx)(d + ex) dx = \frac{1}{5} Bb^2ex^5 + Aa^2dx + \frac{1}{4} (Bb^2d + (2 Bab + Ab^2)e)x^4$$

$$+ \frac{1}{3} ((2 Bab + Ab^2)d + (Ba^2 + 2 Aab)e)x^3$$

$$+ \frac{1}{2} (Aa^2e + (Ba^2 + 2 Aab)d)x^2$$

input `integrate((b*x+a)^2*(B*x+A)*(e*x+d),x, algorithm="maxima")`output `1/5*B*b^2*e*x^5 + A*a^2*d*x + 1/4*(B*b^2*d + (2*B*a*b + A*b^2)*e)*x^4 + 1/3*((2*B*a*b + A*b^2)*d + (B*a^2 + 2*A*a*b)*e)*x^3 + 1/2*(A*a^2*e + (B*a^2 + 2*A*a*b)*d)*x^2`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.51

$$\int (a + bx)^2(A + Bx)(d + ex) dx = \frac{1}{5} Bb^2ex^5 + \frac{1}{4} Bb^2dx^4 + \frac{1}{2} Babex^4 + \frac{1}{4} Ab^2ex^4$$

$$+ \frac{2}{3} Babdx^3 + \frac{1}{3} Ab^2dx^3 + \frac{1}{3} Ba^2ex^3 + \frac{2}{3} Aabex^3$$

$$+ \frac{1}{2} Ba^2dx^2 + Aabdx^2 + \frac{1}{2} Aa^2ex^2 + Aa^2dx$$

input `integrate((b*x+a)^2*(B*x+A)*(e*x+d),x, algorithm="giac")`output `1/5*B*b^2*e*x^5 + 1/4*B*b^2*d*x^4 + 1/2*B*a*b*e*x^4 + 1/4*A*b^2*e*x^4 + 2/3*B*a*b*d*x^3 + 1/3*A*b^2*d*x^3 + 1/3*B*a^2*e*x^3 + 2/3*A*a*b*e*x^3 + 1/2*B*a^2*d*x^2 + A*a*b*d*x^2 + 1/2*A*a^2*e*x^2 + A*a^2*d*x`

Mupad [B] (verification not implemented)

Time = 0.90 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.31

$$\int (a + bx)^2(A + Bx)(d + ex) dx = x^3 \left(\frac{Ab^2d}{3} + \frac{Ba^2e}{3} + \frac{2Aabe}{3} + \frac{2Babd}{3} \right) \\ + x^2 \left(\frac{Aa^2e}{2} + \frac{Ba^2d}{2} + Aabd \right) \\ + x^4 \left(\frac{Ab^2e}{4} + \frac{Bb^2d}{4} + \frac{Babe}{2} \right) \\ + Aa^2dx + \frac{Bb^2ex^5}{5}$$

input `int((A + B*x)*(a + b*x)^2*(d + e*x), x)`output `x^3*((A*b^2*d)/3 + (B*a^2*e)/3 + (2*A*a*b*e)/3 + (2*B*a*b*d)/3) + x^2*((A*a^2*e)/2 + (B*a^2*d)/2 + A*a*b*d) + x^4*((A*b^2*e)/4 + (B*b^2*d)/4 + (B*a*b*e)/2) + A*a^2*d*x + (B*b^2*e*x^5)/5`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.97

$$\int (a + bx)^2(A + Bx)(d + ex) dx \\ = \frac{x(4b^3ex^4 + 15ab^2ex^3 + 5b^3dx^3 + 20a^2bex^2 + 20ab^2dx^2 + 10a^3ex + 30a^2bdx + 20a^3d)}{20}$$

input `int((b*x+a)^2*(B*x+A)*(e*x+d), x)`output `(x*(20*a**3*d + 10*a**3*e*x + 30*a**2*b*d*x + 20*a**2*b*e*x**2 + 20*a*b**2*d*x**2 + 15*a*b**2*e*x**3 + 5*b**3*d*x**3 + 4*b**3*e*x**4))/20`

3.16 $\int (a + bx)^2(A + Bx) dx$

Optimal result	222
Mathematica [A] (verified)	222
Rubi [A] (verified)	223
Maple [A] (verified)	224
Fricas [A] (verification not implemented)	224
Sympy [A] (verification not implemented)	225
Maxima [A] (verification not implemented)	225
Giac [A] (verification not implemented)	225
Mupad [B] (verification not implemented)	226
Reduce [B] (verification not implemented)	226

Optimal result

Integrand size = 13, antiderivative size = 38

$$\int (a + bx)^2(A + Bx) dx = \frac{(Ab - aB)(a + bx)^3}{3b^2} + \frac{B(a + bx)^4}{4b^2}$$

output $1/3*(A*b-B*a)*(b*x+a)^3/b^2+1/4*B*(b*x+a)^4/b^2$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.21

$$\int (a + bx)^2(A + Bx) dx = \frac{1}{12}x(6a^2(2A + Bx) + 4abx(3A + 2Bx) + b^2x^2(4A + 3Bx))$$

input $\text{Integrate}[(a + b*x)^2*(A + B*x), x]$

output $(x*(6*a^2*(2*A + B*x) + 4*a*b*x*(3*A + 2*B*x) + b^2*x^2*(4*A + 3*B*x)))/12$

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^2 (A + Bx) dx$$

$$\downarrow 49$$

$$\int \left(\frac{(a + bx)^2 (Ab - aB)}{b} + \frac{B(a + bx)^3}{b} \right) dx$$

$$\downarrow 2009$$

$$\frac{(a + bx)^3 (Ab - aB)}{3b^2} + \frac{B(a + bx)^4}{4b^2}$$

input `Int[(a + b*x)^2*(A + B*x),x]`

output `((A*b - a*B)*(a + b*x)^3)/(3*b^2) + (B*(a + b*x)^4)/(4*b^2)`

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.26

method	result	size
norman	$\frac{b^2 B x^4}{4} + \left(\frac{1}{3}b^2 A + \frac{2}{3}abB\right) x^3 + (abA + \frac{1}{2}a^2 B) x^2 + a^2 Ax$	48
default	$\frac{b^2 B x^4}{4} + \frac{(b^2 A + 2abB)x^3}{3} + \frac{(2abA + a^2 B)x^2}{2} + a^2 Ax$	49
gosper	$\frac{1}{4}b^2 B x^4 + \frac{1}{3}x^3 b^2 A + \frac{2}{3}x^3 abB + x^2 abA + \frac{1}{2}x^2 a^2 B + a^2 Ax$	50
risch	$\frac{1}{4}b^2 B x^4 + \frac{1}{3}x^3 b^2 A + \frac{2}{3}x^3 abB + x^2 abA + \frac{1}{2}x^2 a^2 B + a^2 Ax$	50
parallelrisch	$\frac{1}{4}b^2 B x^4 + \frac{1}{3}x^3 b^2 A + \frac{2}{3}x^3 abB + x^2 abA + \frac{1}{2}x^2 a^2 B + a^2 Ax$	50
orering	$\frac{x(3B b^2 x^3 + 4A b^2 x^2 + 8Bab x^2 + 12aAbx + 6B a^2 x + 12a^2 A)}{12}$	50

input `int((b*x+a)^2*(B*x+A), x, method=_RETURNVERBOSE)`

output `1/4*b^2*B*x^4+(1/3*b^2*A+2/3*a*b*B)*x^3+(a*b*A+1/2*a^2*B)*x^2+a^2*A*x`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.26

$$\int (a + bx)^2 (A + Bx) dx = \frac{1}{4} B b^2 x^4 + A a^2 x + \frac{1}{3} (2 B a b + A b^2) x^3 + \frac{1}{2} (B a^2 + 2 A a b) x^2$$

input `integrate((b*x+a)^2*(B*x+A), x, algorithm="fricas")`

output `1/4*B*b^2*x^4 + A*a^2*x + 1/3*(2*B*a*b + A*b^2)*x^3 + 1/2*(B*a^2 + 2*A*a*b)*x^2`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.29

$$\int (a + bx)^2 (A + Bx) dx = Aa^2x + \frac{Bb^2x^4}{4} + x^3 \left(\frac{Ab^2}{3} + \frac{2Bab}{3} \right) + x^2 \left(Aab + \frac{Ba^2}{2} \right)$$

input `integrate((b*x+a)**2*(B*x+A),x)`output `A*a**2*x + B*b**2*x**4/4 + x**3*(A*b**2/3 + 2*B*a*b/3) + x**2*(A*a*b + B*a**2/2)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.26

$$\int (a + bx)^2 (A + Bx) dx = \frac{1}{4} Bb^2x^4 + Aa^2x + \frac{1}{3} (2 Bab + Ab^2)x^3 + \frac{1}{2} (Ba^2 + 2 Aab)x^2$$

input `integrate((b*x+a)^2*(B*x+A),x, algorithm="maxima")`output `1/4*B*b^2*x^4 + A*a^2*x + 1/3*(2*B*a*b + A*b^2)*x^3 + 1/2*(B*a^2 + 2*A*a*b)*x^2`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.29

$$\int (a + bx)^2 (A + Bx) dx = \frac{1}{4} Bb^2x^4 + \frac{2}{3} Babx^3 + \frac{1}{3} Ab^2x^3 + \frac{1}{2} Ba^2x^2 + Aabx^2 + Aa^2x$$

input `integrate((b*x+a)^2*(B*x+A),x, algorithm="giac")`output `1/4*B*b^2*x^4 + 2/3*B*a*b*x^3 + 1/3*A*b^2*x^3 + 1/2*B*a^2*x^2 + A*a*b*x^2 + A*a^2*x`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.24

$$\int (a + bx)^2(A + Bx) dx = x^2 \left(\frac{B a^2}{2} + A b a \right) + x^3 \left(\frac{A b^2}{3} + \frac{2 B a b}{3} \right) + \frac{B b^2 x^4}{4} + A a^2 x$$

input `int((A + B*x)*(a + b*x)^2,x)`

output `x^2*((B*a^2)/2 + A*a*b) + x^3*((A*b^2)/3 + (2*B*a*b)/3) + (B*b^2*x^4)/4 + A*a^2*x`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int (a + bx)^2(A + Bx) dx = \frac{x(b^3 x^3 + 4a b^2 x^2 + 6a^2 b x + 4a^3)}{4}$$

input `int((b*x+a)^2*(B*x+A),x)`

output `(x*(4*a**3 + 6*a**2*b*x + 4*a*b**2*x**2 + b**3*x**3))/4`

3.17 $\int \frac{(a+bx)^2(A+Bx)}{d+ex} dx$

Optimal result	227
Mathematica [A] (verified)	227
Rubi [A] (verified)	228
Maple [A] (verified)	229
Fricas [A] (verification not implemented)	230
Sympy [A] (verification not implemented)	230
Maxima [A] (verification not implemented)	231
Giac [A] (verification not implemented)	231
Mupad [B] (verification not implemented)	232
Reduce [B] (verification not implemented)	232

Optimal result

Integrand size = 20, antiderivative size = 92

$$\int \frac{(a+bx)^2(A+Bx)}{d+ex} dx = \frac{b(bd-ae)(Bd-Ae)x}{e^3} - \frac{(Bd-Ae)(a+bx)^2}{2e^2} + \frac{B(a+bx)^3}{3be} - \frac{(bd-ae)^2(Bd-Ae)\log(d+ex)}{e^4}$$

output

```
b*(-a*e+b*d)*(-A*e+B*d)*x/e^3-1/2*(-A*e+B*d)*(b*x+a)^2/e^2+1/3*B*(b*x+a)^3/b/e-(-a*e+b*d)^2*(-A*e+B*d)*ln(e*x+d)/e^4
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.11

$$\int \frac{(a+bx)^2(A+Bx)}{d+ex} dx = \frac{ex(6a^2Be^2 + 6abe(-2Bd + 2Ae + Bex) + b^2(3Ae(-2d + ex) + B(6d^2 - 3dex + 2e^2x^2))) - 6(bd - ae)^2}{6e^4}$$

input

```
Integrate[((a + b*x)^2*(A + B*x))/(d + e*x), x]
```


output

```
(e*x*(6*a^2*B*e^2 + 6*a*b*e*(-2*B*d + 2*A*e + B*e*x) + b^2*(3*A*e*(-2*d +
e*x) + B*(6*d^2 - 3*d*e*x + 2*e^2*x^2))) - 6*(b*d - a*e)^2*(B*d - A*e)*Log
[d + e*x])/(6*e^4)
```

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^2(A + Bx)}{d + ex} dx$$

↓ 86

$$\int \left(\frac{(ae - bd)^2(Ae - Bd)}{e^3(d + ex)} - \frac{b(bd - ae)(Ae - Bd)}{e^3} + \frac{b(a + bx)(Ae - Bd)}{e^2} + \frac{B(a + bx)^2}{e} \right) dx$$

↓ 2009

$$-\frac{(bd - ae)^2(Bd - Ae) \log(d + ex)}{e^4} + \frac{bx(bd - ae)(Bd - Ae)}{3be} - \frac{(a + bx)^2(Bd - Ae)}{2e^2} +$$

input

```
Int[((a + b*x)^2*(A + B*x))/(d + e*x), x]
```

output

```
(b*(b*d - a*e)*(B*d - A*e)*x)/e^3 - ((B*d - A*e)*(a + b*x)^2)/(2*e^2) + (B
*(a + b*x)^3)/(3*b*e) - ((b*d - a*e)^2*(B*d - A*e)*Log[d + e*x])/e^4
```

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.61

method	result
norman	$\frac{(2Aabe^2 - Ab^2de + Ba^2e^2 - 2Babde + b^2Bd^2)x}{e^3} + \frac{b(Abe + 2Bae - Bbd)x^2}{2e^2} + \frac{b^2Bx^3}{3e} + \frac{(a^2Ae^3 - 2Aabd e^2 + Ab^2d^2e - Ba^2d^2e^2)}{6e^4}$
default	$\frac{\frac{1}{3}b^2Bx^3e^2 + \frac{1}{2}Ab^2e^2x^2 + Bab e^2x^2 - \frac{1}{2}Bb^2dex^2 + 2Aabe^2x - Ab^2dex + Ba^2e^2x - 2Babdex + b^2Bd^2x}{e^3} + \frac{(a^2Ae^3 - 2Aabd e^2 - Ba^2d^2e^2)}{6e^4}$
risch	$\frac{b^2Bx^3}{3e} + \frac{Ab^2x^2}{2e} + \frac{Babx^2}{e} - \frac{Bb^2dx^2}{2e^2} + \frac{2Aabx}{e} - \frac{Ab^2dx}{e^2} + \frac{Ba^2x}{e} - \frac{2Babd x}{e^2} + \frac{b^2Bd^2x}{e^3} + \frac{\ln(ex+d)a^2A}{e}$
parallelrisch	$\frac{2b^2Bx^3e^3 + 3Ax^2b^2e^3 + 6Bx^2abe^3 - 3Bx^2b^2de^2 + 6A\ln(ex+d)a^2e^3 - 12A\ln(ex+d)abde^2 + 6A\ln(ex+d)b^2d^2e + 12Axabe^3 - 6e^4}{6e^4}$

```
input int((b*x+a)^2*(B*x+A)/(e*x+d), x, method=_RETURNVERBOSE)
```

```
output (2*A*a*b*e^2 - A*b^2*d*e + B*a^2*e^2 - 2*B*a*b*d*e + B*b^2*d^2)/e^3*x + 1/2*b/e^2*(A*b*e + 2*B*a*e - B*b*d)*x^2 + 1/3*b^2*B/e*x^3 + (A*a^2*e^3 - 2*A*a*b*d*e^2 + A*b^2*d^2*e - B*a^2*d*e^2 + 2*B*a*b*d^2*e - B*b^2*d^3)/e^4*ln(e*x+d)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.66

$$\int \frac{(a+bx)^2(A+Bx)}{d+ex} dx = \frac{2Bb^2e^3x^3 - 3(Bb^2de^2 - (2Bab + Ab^2)e^3)x^2 + 6(Bb^2d^2e - (2Bab + Ab^2)de^2 + (Ba^2 + 2Aab)e^3)x - 6}{6e^4}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d),x, algorithm="fricas")`output `1/6*(2*B*b^2*e^3*x^3 - 3*(B*b^2*d*e^2 - (2*B*a*b + A*b^2)*e^3)*x^2 + 6*(B*b^2*d^2*e - (2*B*a*b + A*b^2)*d*e^2 + (B*a^2 + 2*A*a*b)*e^3)*x - 6*(B*b^2*d^3 - A*a^2*e^3 - (2*B*a*b + A*b^2)*d^2*e + (B*a^2 + 2*A*a*b)*d*e^2)*log(e*x + d))/e^4`**Sympy [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.27

$$\int \frac{(a+bx)^2(A+Bx)}{d+ex} dx = \frac{Bb^2x^3}{3e} + x^2 \left(\frac{Ab^2}{2e} + \frac{Bab}{e} - \frac{Bb^2d}{2e^2} \right) + x \left(\frac{2Aab}{e} - \frac{Ab^2d}{e^2} + \frac{Ba^2}{e} - \frac{2Babd}{e^2} + \frac{Bb^2d^2}{e^3} \right) - \frac{(-Ae + Bd)(ae - bd)^2 \log(d + ex)}{e^4}$$

input `integrate((b*x+a)**2*(B*x+A)/(e*x+d),x)`output `B*b**2*x**3/(3*e) + x**2*(A*b**2/(2*e) + B*a*b/e - B*b**2*d/(2*e**2)) + x*(2*A*a*b/e - A*b**2*d/e**2 + B*a**2/e - 2*B*a*b*d/e**2 + B*b**2*d**2/e**3) - (-A*e + B*d)*(a*e - b*d)**2*log(d + e*x)/e**4`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.65

$$\int \frac{(a+bx)^2(A+Bx)}{d+ex} dx$$

$$= \frac{2Bb^2e^2x^3 - 3(Bb^2de - (2Bab + Ab^2)e^2)x^2 + 6(Bb^2d^2 - (2Bab + Ab^2)de + (Ba^2 + 2Aab)e^2)x - \frac{(Bb^2d^3 - Aa^2e^3 - (2Bab + Ab^2)d^2e + (Ba^2 + 2Aab)de^2) \log(ex+d)}{e^4}}{e^4}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d),x, algorithm="maxima")`

output `1/6*(2*B*b^2*e^2*x^3 - 3*(B*b^2*d*e - (2*B*a*b + A*b^2)*e^2)*x^2 + 6*(B*b^2*d^2 - (2*B*a*b + A*b^2)*d*e + (B*a^2 + 2*A*a*b)*e^2)*x)/e^3 - (B*b^2*d^3 - A*a^2*e^3 - (2*B*a*b + A*b^2)*d^2*e + (B*a^2 + 2*A*a*b)*d*e^2)*log(e*x + d)/e^4`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.80

$$\int \frac{(a+bx)^2(A+Bx)}{d+ex} dx$$

$$= \frac{2Bb^2e^2x^3 - 3Bb^2dex^2 + 6Babe^2x^2 + 3Ab^2e^2x^2 + 6Bb^2d^2x - 12Babdex - 6Ab^2dex + 6Ba^2e^2x + 12Aa^2e^2x - \frac{(Bb^2d^3 - 2Babd^2e - Ab^2d^2e + Ba^2de^2 + 2Aabde^2 - Aa^2e^3) \log(|ex+d|)}{e^4}}{e^4}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d),x, algorithm="giac")`

output `1/6*(2*B*b^2*e^2*x^3 - 3*B*b^2*d*e*x^2 + 6*B*a*b*e^2*x^2 + 3*A*b^2*e^2*x^2 + 6*B*b^2*d^2*x - 12*B*a*b*d*e*x - 6*A*b^2*d*e*x + 6*B*a^2*e^2*x + 12*A*a*b*e^2*x)/e^3 - (B*b^2*d^3 - 2*B*a*b*d^2*e - A*b^2*d^2*e + B*a^2*d*e^2 + 2*A*a*b*d*e^2 - A*a^2*e^3)*log(abs(e*x + d))/e^4`

Mupad [B] (verification not implemented)

Time = 0.92 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.73

$$\int \frac{(a + bx)^2(A + Bx)}{d + ex} dx$$

$$= x \left(\frac{B a^2 + 2 A b a}{e} - \frac{d \left(\frac{A b^2 + 2 B a b}{e} - \frac{B b^2 d}{e^2} \right)}{e} \right) + x^2 \left(\frac{A b^2 + 2 B a b}{2 e} - \frac{B b^2 d}{2 e^2} \right)$$

$$+ \frac{\ln(d + ex) (-B a^2 d e^2 + A a^2 e^3 + 2 B a b d^2 e - 2 A a b d e^2 - B b^2 d^3 + A b^2 d^2 e)}{e^4}$$

$$+ \frac{B b^2 x^3}{3 e}$$

input `int(((A + B*x)*(a + b*x)^2)/(d + e*x),x)`output `x*((B*a^2 + 2*A*a*b)/e - (d*((A*b^2 + 2*B*a*b)/e - (B*b^2*d)/e^2))/e) + x^2*((A*b^2 + 2*B*a*b)/(2*e) - (B*b^2*d)/(2*e^2)) + (log(d + e*x)*(A*a^2*e^3 - B*b^2*d^3 + A*b^2*d^2*e - B*a^2*d*e^2 - 2*A*a*b*d*e^2 + 2*B*a*b*d^2*e))/e^4 + (B*b^2*x^3)/(3*e)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.43

$$\int \frac{(a + bx)^2(A + Bx)}{d + ex} dx$$

$$= \frac{6 \log(ex + d) a^3 e^3 - 18 \log(ex + d) a^2 b d e^2 + 18 \log(ex + d) a b^2 d^2 e - 6 \log(ex + d) b^3 d^3 + 18 a^2 b e^3 x - 18 a b^2 d^2 x^2 + 6 b^3 d^3 x^3}{6 e^4}$$

input `int((b*x+a)^2*(B*x+A)/(e*x+d),x)`output `(6*log(d + e*x)*a**3*e**3 - 18*log(d + e*x)*a**2*b*d*e**2 + 18*log(d + e*x)*a*b**2*d**2*e - 6*log(d + e*x)*b**3*d**3 + 18*a**2*b*e**3*x - 18*a*b**2*d*e**2*x + 9*a*b**2*e**3*x**2 + 6*b**3*d**2*e*x - 3*b**3*d*e**2*x**2 + 2*b**3*e**3*x**3)/(6*e**4)`

3.18 $\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^2} dx$

Optimal result	233
Mathematica [A] (verified)	233
Rubi [A] (verified)	234
Maple [A] (verified)	235
Fricas [B] (verification not implemented)	236
Sympy [A] (verification not implemented)	236
Maxima [A] (verification not implemented)	237
Giac [B] (verification not implemented)	238
Mupad [B] (verification not implemented)	238
Reduce [B] (verification not implemented)	239

Optimal result

Integrand size = 20, antiderivative size = 101

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^2} dx = -\frac{b(2bBd - Abe - 2aBe)x}{e^3} + \frac{b^2Bx^2}{2e^2} + \frac{(bd - ae)^2(Bd - Ae)}{e^4(d+ex)} + \frac{(bd - ae)(3bBd - 2Abe - aBe) \log(d+ex)}{e^4}$$

output

```
-b*(-A*b*e-2*B*a*e+2*B*b*d)*x/e^3+1/2*b^2*B*x^2/e^2+(-a*e+b*d)^2*(-A*e+B*d)/e^4/(e*x+d)+(-a*e+b*d)*(-2*A*b*e-B*a*e+3*B*b*d)*ln(e*x+d)/e^4
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.97

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^2} dx = \frac{2be(-2bBd + Abe + 2aBe)x + b^2Be^2x^2 + \frac{2(bd-ae)^2(Bd-Ae)}{d+ex} + 2(bd-ae)(3bBd - 2Abe - aBe) \log(d+ex)}{2e^4}$$

input

```
Integrate[((a + b*x)^2*(A + B*x))/(d + e*x)^2,x]
```

output

$$\frac{(2*b*e*(-2*b*B*d + A*b*e + 2*a*B*e)*x + b^2*B*e^2*x^2 + (2*(b*d - a*e)^2*(B*d - A*e)))/(d + e*x) + 2*(b*d - a*e)*(3*b*B*d - 2*A*b*e - a*B*e)*\text{Log}[d + e*x]}{(2*e^4)}$$

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^2} dx$$

↓ 86

$$\int \left(\frac{(ae - bd)(aBe + 2Abe - 3bBd)}{e^3(d + ex)} + \frac{(ae - bd)^2(Ae - Bd)}{e^3(d + ex)^2} + \frac{b(2aBe + Abe - 2bBd)}{e^3} + \frac{b^2Bx}{e^2} \right) dx$$

↓ 2009

$$\frac{(bd - ae)^2(Bd - Ae)}{e^4(d + ex)} + \frac{(bd - ae) \log(d + ex)(-aBe - 2Abe + 3bBd)}{e^4} - \frac{bx(-2aBe - Abe + 2bBd)}{e^3} + \frac{b^2Bx^2}{2e^2}$$

input

$$\text{Int}[(a + b*x)^2*(A + B*x)/(d + e*x)^2, x]$$

output

$$-\frac{(b*(2*b*B*d - A*b*e - 2*a*B*e)*x)/e^3}{e^4} + \frac{(b^2*B*x^2)/(2*e^2)}{e^4} + \frac{((b*d - a*e)^2*(B*d - A*e))/(e^4*(d + e*x))}{e^4} + \frac{((b*d - a*e)*(3*b*B*d - 2*A*b*e - a*B*e)*\text{Log}[d + e*x])}{e^4}$$

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.50

method	result
default	$\frac{b(\frac{1}{2}Bbe^2x^2 + Abox + 2Baex - 2Bbdx)}{e^3} - \frac{a^2Ae^3 - 2Aabd e^2 + Ab^2d^2e - Ba^2de^2 + 2Babd^2e - b^2Bd^3}{e^4(ex+d)} + \frac{(2Aabe^2 - 2Ab^2de + B a^2e^2)}{e^4(ex+d)}$
norman	$\frac{(a^2Ae^3 - 2Aabd e^2 + 2Ab^2d^2e - Ba^2de^2 + 4Babd^2e - 3b^2Bd^3)x}{e^3d} + \frac{b(2Abe + 4Bae - 3Bbd)x^2}{2e^2} + \frac{b^2Bx^3}{2e} + \frac{(2Aabe^2 - 2Ab^2de + B a^2e^2)}{e^4(ex+d)}$
risch	$\frac{b^2Bx^2}{2e^2} + \frac{b^2Ax}{e^2} + \frac{2bBax}{e^2} - \frac{2b^2Bdx}{e^3} - \frac{a^2A}{e(ex+d)} + \frac{2Aabd}{e^2(ex+d)} - \frac{Ab^2d^2}{e^3(ex+d)} + \frac{Ba^2d}{e^2(ex+d)} - \frac{2Babd^2}{e^3(ex+d)} + \frac{b^2Bd^3}{e^4(ex+d)}$
parallelrisc	$\frac{b^2Bx^3e^3 + 4A \ln(ex+d)xab e^3 - 4A \ln(ex+d)xb^2d e^2 + 2Ax^2b^2e^3 + 2B \ln(ex+d)xa^2e^3 - 8B \ln(ex+d)xabd e^2 + 6B \ln(ex+d)xa^2e^2}{e^4(ex+d)}$

```
input int((b*x+a)^2*(B*x+A)/(e*x+d)^2,x,method=_RETURNVERBOSE)
```

```
output b/e^3*(1/2*B*b*e*x^2+A*b*e*x+2*B*a*e*x-2*B*b*d*x)-(A*a^2*e^3-2*A*a*b*d*e^2+A*b^2*d^2*e-B*a^2*d*e^2+2*B*a*b*d^2*e-B*b^2*d^3)/e^4/(e*x+d)+1/e^4*(2*A*a*b*e^2-2*A*b^2*d*e+B*a^2*e^2-4*B*a*b*d*e+3*B*b^2*d^2)*ln(e*x+d)
```


Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 238 vs. $2(99) = 198$.

Time = 0.08 (sec) , antiderivative size = 238, normalized size of antiderivative = 2.36

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^2} dx$$

$$= \frac{Bb^2e^3x^3 + 2Bb^2d^3 - 2Aa^2e^3 - 2(2Bab + Ab^2)d^2e + 2(Ba^2 + 2Aab)de^2 - (3Bb^2de^2 - 2(2Bab + Ab^2)d^2e)}{(d+ex)^2}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^2,x, algorithm="fricas")`

output `1/2*(B*b^2*e^3*x^3 + 2*B*b^2*d^3 - 2*A*a^2*e^3 - 2*(2*B*a*b + A*b^2)*d^2*e + 2*(B*a^2 + 2*A*a*b)*d*e^2 - (3*B*b^2*d^2*e^2 - 2*(2*B*a*b + A*b^2)*e^3)*x^2 - 2*(2*B*b^2*d^2*e - (2*B*a*b + A*b^2)*d*e^2)*x + 2*(3*B*b^2*d^3 - 2*(2*B*a*b + A*b^2)*d^2*e + (B*a^2 + 2*A*a*b)*d*e^2 + (3*B*b^2*d^2*e - 2*(2*B*a*b + A*b^2)*d*e^2 + (B*a^2 + 2*A*a*b)*e^3)*x)*log(e*x + d)/(e^5*x + d*e^4)`

Sympy [A] (verification not implemented)

Time = 0.47 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.50

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^2} dx$$

$$= \frac{Bb^2x^2}{2e^2} + x \left(\frac{Ab^2}{e^2} + \frac{2Bab}{e^2} - \frac{2Bb^2d}{e^3} \right)$$

$$+ \frac{-Aa^2e^3 + 2Aabde^2 - Ab^2d^2e + Ba^2de^2 - 2Babd^2e + Bb^2d^3}{de^4 + e^5x}$$

$$+ \frac{(ae - bd)(2Abe + Bae - 3Bbd) \log(d + ex)}{e^4}$$

input `integrate((b*x+a)**2*(B*x+A)/(e*x+d)**2,x)`

output

```
B*b**2*x**2/(2*e**2) + x*(A*b**2/e**2 + 2*B*a*b/e**2 - 2*B*b**2*d/e**3) +
(-A*a**2*e**3 + 2*A*a*b*d*e**2 - A*b**2*d**2*e + B*a**2*d*e**2 - 2*B*a*b*d
**2*e + B*b**2*d**3)/(d*e**4 + e**5*x) + (a*e - b*d)*(2*A*b*e + B*a*e - 3*
B*b*d)*log(d + e*x)/e**4
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.54

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^2} dx$$

$$= \frac{Bb^2d^3 - Aa^2e^3 - (2Bab + Ab^2)d^2e + (Ba^2 + 2Aab)de^2}{e^5x + de^4}$$

$$+ \frac{Bb^2ex^2 - 2(2Bb^2d - (2Bab + Ab^2)e)x}{2e^3}$$

$$+ \frac{(3Bb^2d^2 - 2(2Bab + Ab^2)de + (Ba^2 + 2Aab)e^2) \log(ex + d)}{e^4}$$

input

```
integrate((b*x+a)^2*(B*x+A)/(e*x+d)^2,x, algorithm="maxima")
```

output

```
(B*b^2*d^3 - A*a^2*e^3 - (2*B*a*b + A*b^2)*d^2*e + (B*a^2 + 2*A*a*b)*d*e^2
)/(e^5*x + d*e^4) + 1/2*(B*b^2*e*x^2 - 2*(2*B*b^2*d - (2*B*a*b + A*b^2)*e)
*x)/e^3 + (3*B*b^2*d^2 - 2*(2*B*a*b + A*b^2)*d*e + (B*a^2 + 2*A*a*b)*e^2)*
log(e*x + d)/e^4
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 230 vs. $2(99) = 198$.

Time = 0.12 (sec) , antiderivative size = 230, normalized size of antiderivative = 2.28

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^2} dx$$

$$= \frac{\left(Bb^2 - \frac{2(3Bb^2de - 2Babe^2 - Ab^2e^2)}{(ex+d)e}\right)(ex + d)^2}{2e^4}$$

$$- \frac{(3Bb^2d^2 - 4Babde - 2Ab^2de + Ba^2e^2 + 2Aabe^2) \log\left(\frac{|ex+d|}{(ex+d)^2|e|}\right)}{e^4}$$

$$+ \frac{\frac{Bb^2d^3e^2}{ex+d} - \frac{2Babd^2e^3}{ex+d} - \frac{Ab^2d^2e^3}{ex+d} + \frac{Ba^2de^4}{ex+d} + \frac{2Aabde^4}{ex+d} - \frac{Aa^2e^5}{ex+d}}{e^6}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^2,x, algorithm="giac")`

output `1/2*(B*b^2 - 2*(3*B*b^2*d*e - 2*B*a*b*e^2 - A*b^2*e^2)/((e*x + d)*e))*(e*x + d)^2/e^4 - (3*B*b^2*d^2 - 4*B*a*b*d*e - 2*A*b^2*d*e + B*a^2*e^2 + 2*A*a*b*e^2)*log(abs(e*x + d)/((e*x + d)^2*abs(e)))/e^4 + (B*b^2*d^3*e^2/(e*x + d) - 2*B*a*b*d^2*e^3/(e*x + d) - A*b^2*d^2*e^3/(e*x + d) + B*a^2*d*e^4/(e*x + d) + 2*A*a*b*d*e^4/(e*x + d) - A*a^2*e^5/(e*x + d))/e^6`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.63

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^2} dx$$

$$= x \left(\frac{Ab^2 + 2Bab}{e^2} - \frac{2Bb^2d}{e^3} \right)$$

$$+ \frac{\ln(d + ex) (Ba^2e^2 - 4Babde + 2Aabe^2 + 3Bb^2d^2 - 2Ab^2de)}{e^4}$$

$$- \frac{-Ba^2de^2 + Aa^2e^3 + 2Babd^2e - 2Aabde^2 - Bb^2d^3 + Ab^2d^2e}{e(xe^4 + de^3)} + \frac{Bb^2x^2}{2e^2}$$

input `int(((A + B*x)*(a + b*x)^2)/(d + e*x)^2,x)`

output

```
x*((A*b^2 + 2*B*a*b)/e^2 - (2*B*b^2*d)/e^3) + (log(d + e*x)*(B*a^2*e^2 + 3
*B*b^2*d^2 + 2*A*a*b*e^2 - 2*A*b^2*d*e - 4*B*a*b*d*e))/e^4 - (A*a^2*e^3 -
B*b^2*d^3 + A*b^2*d^2*e - B*a^2*d*e^2 - 2*A*a*b*d*e^2 + 2*B*a*b*d^2*e)/(e*
(d*e^3 + e^4*x)) + (B*b^2*x^2)/(2*e^2)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.95

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^2} dx$$

$$= \frac{6 \log(ex + d) a^2 b d^2 e^2 + 6 \log(ex + d) a^2 b d e^3 x - 12 \log(ex + d) a b^2 d^3 e - 12 \log(ex + d) a b^2 d^2 e^2 x + 6 \log(ex + d) a^2 b d^2 e^2 x + 6 \log(ex + d) a^2 b d^2 e^2 x + 6 \log(ex + d) a^2 b d^2 e^2 x}{1}$$

input

```
int((b*x+a)^2*(B*x+A)/(e*x+d)^2,x)
```

output

```
(6*log(d + e*x)*a**2*b*d**2*e**2 + 6*log(d + e*x)*a**2*b*d*e**3*x - 12*log
(d + e*x)*a*b**2*d**3*e - 12*log(d + e*x)*a*b**2*d**2*e**2*x + 6*log(d + e
*x)*b**3*d**4 + 6*log(d + e*x)*b**3*d**3*e*x + 2*a**3*e**4*x - 6*a**2*b*d*
e**3*x + 12*a*b**2*d**2*e**2*x + 6*a*b**2*d*e**3*x**2 - 6*b**3*d**3*e*x -
3*b**3*d**2*e**2*x**2 + b**3*d*e**3*x**3)/(2*d*e**4*(d + e*x))
```

3.19 $\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^3} dx$

Optimal result	240
Mathematica [A] (verified)	240
Rubi [A] (verified)	241
Maple [A] (verified)	242
Fricas [B] (verification not implemented)	243
Sympy [A] (verification not implemented)	243
Maxima [A] (verification not implemented)	244
Giac [A] (verification not implemented)	244
Mupad [B] (verification not implemented)	245
Reduce [B] (verification not implemented)	245

Optimal result

Integrand size = 20, antiderivative size = 106

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^3} dx = \frac{b^2 Bx}{e^3} + \frac{(bd-ae)^2(Bd-Ae)}{2e^4(d+ex)^2} - \frac{(bd-ae)(3bBd-2Abe-aBe)}{e^4(d+ex)} - \frac{b(3bBd-Abe-2aBe)\log(d+ex)}{e^4}$$

output

```
b^2*B*x/e^3+1/2*(-a*e+b*d)^2*(-A*e+B*d)/e^4/(e*x+d)^2-(-a*e+b*d)*(-2*A*b*e-B*a*e+3*B*b*d)/e^4/(e*x+d)-b*(-A*b*e-2*B*a*e+3*B*b*d)*ln(e*x+d)/e^4
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.35

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^3} dx = \frac{a^2 e^2 (Ae + B(d+2ex)) + 2abe(Ae(d+2ex) - Bd(3d+4ex)) - b^2(Ade(3d+4ex) + B(-5d^3 - 4d^2e))}{2e^4(d+ex)^2}$$

input

```
Integrate[((a + b*x)^2*(A + B*x))/(d + e*x)^3,x]
```

output

$$\begin{aligned} & -1/2*(a^2*e^2*(A*e + B*(d + 2*e*x)) + 2*a*b*e*(A*e*(d + 2*e*x) - B*d*(3*d \\ & + 4*e*x)) - b^2*(A*d*e*(3*d + 4*e*x) + B*(-5*d^3 - 4*d^2*e*x + 4*d*e^2*x^2 \\ & + 2*e^3*x^3)) + 2*b*(3*b*B*d - A*b*e - 2*a*B*e)*(d + e*x)^2*\text{Log}[d + e*x]) \\ & / (e^4*(d + e*x)^2) \end{aligned}$$

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + bx)^2(A + Bx)}{(d + ex)^3} dx \\ & \quad \downarrow \text{86} \\ & \int \left(\frac{b(2aBe + Abe - 3bBd)}{e^3(d + ex)} + \frac{(ae - bd)(aBe + 2Abe - 3bBd)}{e^3(d + ex)^2} + \frac{(ae - bd)^2(Ae - Bd)}{e^3(d + ex)^3} + \frac{b^2B}{e^3} \right) dx \\ & \quad \downarrow \text{2009} \\ & -\frac{(bd - ae)(-aBe - 2Abe + 3bBd)}{e^4(d + ex)} + \frac{(bd - ae)^2(Bd - Ae)}{2e^4(d + ex)^2} - \\ & \quad \frac{b \log(d + ex)(-2aBe - Abe + 3bBd)}{e^4} + \frac{b^2Bx}{e^3} \end{aligned}$$

input

$$\text{Int}[(a + b*x)^2*(A + B*x)/(d + e*x)^3, x]$$

output

$$\begin{aligned} & (b^2*B*x)/e^3 + ((b*d - a*e)^2*(B*d - A*e))/(2*e^4*(d + e*x)^2) - ((b*d - \\ & a*e)*(3*b*B*d - 2*A*b*e - a*B*e))/(e^4*(d + e*x)) - (b*(3*b*B*d - A*b*e - \\ & 2*a*B*e)*\text{Log}[d + e*x])/e^4 \end{aligned}$$

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.46

method	result
norman	$\frac{\frac{b^2 B x^3 - a^2 A e^3 + 2Aabd e^2 - 3A b^2 d^2 e + B a^2 d e^2 - 6Bab d^2 e + 9b^2 B d^3 - (2Aab e^2 - 2A b^2 d e + B a^2 e^2 - 4Babde + 6b^2 B d^2)x}{e^{2e^4}}}{(ex+d)^2} + \frac{b(Abe+2}{e^3}$
default	$\frac{b^2 B x}{e^3} - \frac{2Aab e^2 - 2A b^2 d e + B a^2 e^2 - 4Babde + 3b^2 B d^2}{e^4(ex+d)} - \frac{a^2 A e^3 - 2Aabd e^2 + A b^2 d^2 e - B a^2 d e^2 + 2Bab d^2 e - b^2 B d^3}{2e^4(ex+d)^2} + \frac{b}{e^3}$
risch	$\frac{b^2 B x}{e^3} + \frac{(-2Aab e^2 + 2A b^2 d e - B a^2 e^2 + 4Babde - 3b^2 B d^2)x - a^2 A e^3 + 2Aabd e^2 - 3A b^2 d^2 e + B a^2 d e^2 - 6Bab d^2 e + 5b^2 B d^3}{e^3(ex+d)^2} + \frac{b}{e^3}$
parallelrisc	$\frac{2A \ln(ex+d)x^2 b^2 e^3 + 4B \ln(ex+d)x^2 ab e^3 - 6B \ln(ex+d)x^2 b^2 d e^2 + 2b^2 B x^3 e^3 + 4A \ln(ex+d)x b^2 d e^2 + 8B \ln(ex+d)xabd e^2 - 1}{e^3}$

```
input int((b*x+a)^2*(B*x+A)/(e*x+d)^3,x,method=_RETURNVERBOSE)
```

```
output (b^2*B/e*x^3-1/2*(A*a^2*e^3+2*A*a*b*d*e^2-3*A*b^2*d^2*e+B*a^2*d*e^2-6*B*a*b*d^2*e+9*B*b^2*d^3)/e^4-(2*A*a*b*e^2-2*A*b^2*d*e+B*a^2*e^2-4*B*a*b*d*e+6*B*b^2*d^2)/e^3*x)/(e*x+d)^2+b/e^4*(A*b*e+2*B*a*e-3*B*b*d)*ln(e*x+d)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 247 vs. $2(104) = 208$.

Time = 0.08 (sec) , antiderivative size = 247, normalized size of antiderivative = 2.33

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^3} dx$$

$$= \frac{2Bb^2e^3x^3 + 4Bb^2de^2x^2 - 5Bb^2d^3 - Aa^2e^3 + 3(2Bab + Ab^2)d^2e - (Ba^2 + 2Aab)de^2 - 2(2Bb^2d^2e - 2$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^3,x, algorithm="fricas")`

output

```
1/2*(2*B*b^2*e^3*x^3 + 4*B*b^2*d*e^2*x^2 - 5*B*b^2*d^3 - A*a^2*e^3 + 3*(2*
B*a*b + A*b^2)*d^2*e - (B*a^2 + 2*A*a*b)*d*e^2 - 2*(2*B*b^2*d^2*e - 2*(2*B
*a*b + A*b^2)*d*e^2 + (B*a^2 + 2*A*a*b)*e^3)*x - 2*(3*B*b^2*d^3 - (2*B*a*b
+ A*b^2)*d^2*e + (3*B*b^2*d*e^2 - (2*B*a*b + A*b^2)*e^3)*x^2 + 2*(3*B*b^2
*d^2*e - (2*B*a*b + A*b^2)*d*e^2)*x)*log(e*x + d))/(e^6*x^2 + 2*d*e^5*x +
d^2*e^4)
```

Sympy [A] (verification not implemented)

Time = 1.10 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.76

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^3} dx = \frac{Bb^2x}{e^3} + \frac{b(Abe + 2Bae - 3Bbd) \log(d+ex)}{e^4}$$

$$+ \frac{-Aa^2e^3 - 2Aabde^2 + 3Ab^2d^2e - Ba^2de^2 + 6Babd^2e - 5Bb^2d^3 + x(-4Aabe^3 + 4Ab^2de^2 - 2Ba^2e^3 + 2$$

$$2d^2e^4 + 4de^5x + 2e^6x^2}$$

input `integrate((b*x+a)**2*(B*x+A)/(e*x+d)**3,x)`

output

```
B*b**2*x/e**3 + b*(A*b*e + 2*B*a*e - 3*B*b*d)*log(d + e*x)/e**4 + (-A*a**2
*e**3 - 2*A*a*b*d*e**2 + 3*A*b**2*d**2*e - B*a**2*d*e**2 + 6*B*a*b*d**2*e
- 5*B*b**2*d**3 + x*(-4*A*a*b*e**3 + 4*A*b**2*d*e**2 - 2*B*a**2*e**3 + 8*B
*a*b*d*e**2 - 6*B*b**2*d**2*e))/(2*d**2*e**4 + 4*d*e**5*x + 2*e**6*x**2)
```


Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.57

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^3} dx = \frac{Bb^2x}{e^3} - \frac{5Bb^2d^3 + Aa^2e^3 - 3(2Bab + Ab^2)d^2e + (Ba^2 + 2Aab)de^2 + 2(3Bb^2d^2e - 2(2Bab + Ab^2)de^2 + (Bb^2d - (2Bab + Ab^2)e)\log(ex+d))}{2(e^6x^2 + 2de^5x + d^2e^4)} - \frac{(3Bb^2d - (2Bab + Ab^2)e)\log(ex+d)}{e^4}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^3,x, algorithm="maxima")`

output

$$B*b^2*x/e^3 - 1/2*(5*B*b^2*d^3 + A*a^2*e^3 - 3*(2*B*a*b + A*b^2)*d^2*e + (B*a^2 + 2*A*a*b)*d*e^2 + 2*(3*B*b^2*d^2*e - 2*(2*B*a*b + A*b^2)*d*e^2 + (B*a^2 + 2*A*a*b)*e^3)*x)/(e^6*x^2 + 2*d*e^5*x + d^2*e^4) - (3*B*b^2*d - (2*B*a*b + A*b^2)*e)*log(e*x + d)/e^4$$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.50

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^3} dx = \frac{Bb^2x}{e^3} - \frac{(3Bb^2d - 2Babe - Ab^2e)\log(|ex+d|)}{e^4} - \frac{5Bb^2d^3 - 6Babd^2e - 3Ab^2d^2e + Ba^2de^2 + 2Aabde^2 + Aa^2e^3 + 2(3Bb^2d^2e - 4Babde^2 - 2Ab^2de^2 - 2Aa^2e^3 + 2Aa*b*e^3)*x}{2(ex+d)^2e^4}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^3,x, algorithm="giac")`

output

$$B*b^2*x/e^3 - (3*B*b^2*d - 2*B*a*b*e - A*b^2*e)*log(abs(e*x + d))/e^4 - 1/2*(5*B*b^2*d^3 - 6*B*a*b*d^2*e - 3*A*b^2*d^2*e + B*a^2*d*e^2 + 2*A*a*b*d*e^2 + A*a^2*e^3 + 2*(3*B*b^2*d^2*e - 4*B*a*b*d*e^2 - 2*A*b^2*d*e^2 + B*a^2*e^3 + 2*A*a*b*e^3)*x)/((e*x + d)^2*e^4)$$

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.60

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^3} dx = \frac{\ln(d+ex)(Ab^2e-3Bb^2d+2Babe)}{e^4} - \frac{x(Ba^2e^2-4Babde+2Aabe^2+3Bb^2d^2-2Ab^2de) + \frac{Ba^2de^2+Ba^2e^3-6Babd^2e+2Aabde^2+5Bb^2d^3-2e}{2e}}{d^2e^3+2de^4x+e^5x^2} + \frac{Bb^2x}{e^3}$$

input `int(((A + B*x)*(a + b*x)^2)/(d + e*x)^3,x)`output `(log(d + e*x)*(A*b^2*e - 3*B*b^2*d + 2*B*a*b*e))/e^4 - (x*(B*a^2*e^2 + 3*B*b^2*d^2 + 2*A*a*b*e^2 - 2*A*b^2*d*e - 4*B*a*b*d*e) + (A*a^2*e^3 + 5*B*b^2*d^3 - 3*A*b^2*d^2*e + B*a^2*d*e^2 + 2*A*a*b*d*e^2 - 6*B*a*b*d^2*e)/(2*e)) / (d^2*e^3 + e^5*x^2 + 2*d*e^4*x) + (B*b^2*x)/e^3`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.97

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^3} dx = \frac{6 \log(ex+d) a b^2 d^3 e + 12 \log(ex+d) a b^2 d^2 e^2 x + 6 \log(ex+d) a b^2 d e^3 x^2 - 6 \log(ex+d) b^3 d^4 - 12 \log(ex+d) b^3 d^3 e x}{2 d e^4}$$

input `int((b*x+a)^2*(B*x+A)/(e*x+d)^3,x)`output `(6*log(d + e*x)*a*b**2*d**3*e + 12*log(d + e*x)*a*b**2*d**2*e**2*x + 6*log(d + e*x)*a*b**2*d*e**3*x**2 - 6*log(d + e*x)*b**3*d**4 - 12*log(d + e*x)*b**3*d**3*e*x - 6*log(d + e*x)*b**3*d**2*e**2*x**2 - a**3*d*e**3 + 3*a**2*b*e**4*x**2 + 3*a*b**2*d**3*e - 6*a*b**2*d*e**3*x**2 - 3*b**3*d**4 + 6*b**3*d**2*e**2*x**2 + 2*b**3*d*e**3*x**3)/(2*d*e**4*(d**2 + 2*d*e*x + e**2*x**2))`

3.20 $\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^4} dx$

Optimal result	246
Mathematica [A] (verified)	246
Rubi [A] (verified)	247
Maple [A] (verified)	248
Fricas [B] (verification not implemented)	249
Sympy [B] (verification not implemented)	249
Maxima [A] (verification not implemented)	250
Giac [A] (verification not implemented)	250
Mupad [B] (verification not implemented)	251
Reduce [B] (verification not implemented)	251

Optimal result

Integrand size = 20, antiderivative size = 101

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^4} dx = -\frac{(Bd-Ae)(a+bx)^3}{3e(bd-ae)(d+ex)^3} - \frac{B(bd-ae)^2}{2e^4(d+ex)^2} + \frac{2bB(bd-ae)}{e^4(d+ex)} + \frac{b^2B \log(d+ex)}{e^4}$$

output

```
-1/3*(-A*e+B*d)*(b*x+a)^3/e/(-a*e+b*d)/(e*x+d)^3-1/2*B*(-a*e+b*d)^2/e^4/(e*x+d)^2+2*b*B*(-a*e+b*d)/e^4/(e*x+d)+b^2*B*ln(e*x+d)/e^4
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.37

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^4} dx = \frac{-a^2e^2(2Ae+B(d+3ex)) - 2abe(Ae(d+3ex) + 2B(d^2+3dex+3e^2x^2)) + b^2(-2Ae(d^2+3dex+3e^2x^2))}{6e^4(d+ex)^3}$$

input

```
Integrate[((a + b*x)^2*(A + B*x))/(d + e*x)^4,x]
```

output

$$\begin{aligned} & (-a^2e^2(2Ae + B(d + 3ex))) - 2ab* e*(Ae*(d + 3ex) + 2B*(d^2 \\ & + 3d*ex + 3e^2*x^2)) + b^2*(-2Ae*(d^2 + 3d*ex + 3e^2*x^2) + B*d*(1 \\ & 1*d^2 + 27*d*ex + 18*e^2*x^2)) + 6*b^2*B*(d + ex)^3*\text{Log}[d + ex]/(6*e^4 \\ & *(d + ex)^3) \end{aligned}$$

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.03, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {87, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + bx)^2(A + Bx)}{(d + ex)^4} dx \\ & \quad \downarrow 87 \\ & \frac{B \int \frac{(a+bx)^2}{(d+ex)^3} dx}{e} - \frac{(a + bx)^3(Bd - Ae)}{3e(d + ex)^3(bd - ae)} \\ & \quad \downarrow 49 \\ & \frac{B \int \left(\frac{b^2}{e^2(d+ex)} - \frac{2(bd-ae)b}{e^2(d+ex)^2} + \frac{(ae-bd)^2}{e^2(d+ex)^3} \right) dx}{e} - \frac{(a + bx)^3(Bd - Ae)}{3e(d + ex)^3(bd - ae)} \\ & \quad \downarrow 2009 \\ & \frac{B \left(\frac{2b(bd-ae)}{e^3(d+ex)} - \frac{(bd-ae)^2}{2e^3(d+ex)^2} + \frac{b^2 \log(d+ex)}{e^3} \right)}{e} - \frac{(a + bx)^3(Bd - Ae)}{3e(d + ex)^3(bd - ae)} \end{aligned}$$

input

$$\text{Int}[(a + b*x)^2*(A + B*x)/(d + e*x)^4, x]$$

output

$$\begin{aligned} & -1/3*((B*d - A*e)*(a + b*x)^3)/(e*(b*d - a*e)*(d + e*x)^3) + (B*(-1/2*(b*d \\ & - a*e)^2/(e^3*(d + e*x)^2) + (2*b*(b*d - a*e))/(e^3*(d + e*x)) + (b^2*\text{Log} \\ & [d + e*x])/e^3)/e \end{aligned}$$

Defintions of rubi rules used

rule 49 $\text{Int}[(a_.) + (b_.)(x_)^{(m_.)}*((c_.) + (d_.)(x_)^{(n_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{IGtQ}[m, 0] \&\& \text{IGtQ}[m + n + 2, 0]$

rule 87 $\text{Int}[(a_.) + (b_.)(x_)^{(c_.)} + (d_.)(x_)^{(n_.)}*((e_.) + (f_.)(x_)^{(p_.)}, x_] \rightarrow \text{Simp}[(-b*e - a*f)*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)})/(f*(p + 1)*(c*f - d*e)), x] - \text{Simp}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) \text{Int}[(c + d*x)^n*(e + f*x)^{(p + 1)}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n\}, x] \&\& \text{LtQ}[p, -1] \&\& (!\text{LtQ}[n, -1] || \text{IntegerQ}[p] || !(\text{IntegerQ}[n] || !(\text{EqQ}[e, 0] || !(\text{EqQ}[c, 0] || \text{LtQ}[p, n])))$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.55

method	result
risch	$\frac{-\frac{b(Abe+2Bae-3Bbd)x^2}{e^2} - \frac{(2Aab e^2+2A b^2 de+B a^2 e^2+4Babde-9b^2 B d^2)x}{2e^3} - \frac{2a^2 A e^3+2Aabd e^2+2A b^2 d^2 e+B a^2 d e^2+4Bab d^2 e-11b^2 d^3}{6e^4}}{(ex+d)^3}$
norman	$\frac{-\frac{2a^2 A e^3+2Aabd e^2+2A b^2 d^2 e+B a^2 d e^2+4Bab d^2 e-11b^2 B d^3}{6e^4} - \frac{(A b^2 e+2Babe-3b^2 Bd)x^2}{e^2} - \frac{(2Aab e^2+2A b^2 de+B a^2 e^2+4Babde-9b^2 B d^2)}{2e^3}}{(ex+d)^3}$
default	$-\frac{b(Abe+2Bae-3Bbd)}{e^4(ex+d)} - \frac{2Aab e^2-2A b^2 de+B a^2 e^2-4Babde+3b^2 B d^2}{2e^4(ex+d)^2} + \frac{b^2 B \ln(ex+d)}{e^4} - \frac{a^2 A e^3-2Aabd e^2+A b^2 d^2}{3e^4}$
parallelrisch	$-\frac{-6B \ln(ex+d)x^3 b^2 e^3-18B \ln(ex+d)x^2 b^2 d e^2+6A x^2 b^2 e^3-18B \ln(ex+d)x b^2 d^2 e+12B x^2 a b e^3-18B x^2 b^2 d e^2+6A x a b e^3}{(ex+d)^3}$

input $\text{int}((b*x+a)^2*(B*x+A)/(e*x+d)^4, x, \text{method}=_RETURNVERBOSE)$

output $(-b*(A*b*e+2*B*a*e-3*B*b*d)/e^2*x^2-1/2*(2*A*a*b*e^2+2*A*b^2*d*e+B*a^2*e^2+4*B*a*b*d*e-9*B*b^2*d^2)/e^3*x-1/6*(2*A*a^2*e^3+2*A*a*b*d*e^2+2*A*b^2*d^2*e+B*a^2*d*e^2+4*B*a*b*d^2*e-11*B*b^2*d^3)/e^4)/(e*x+d)^3+b^2*B*\ln(e*x+d)/e^4$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 221 vs. $2(97) = 194$.

Time = 0.07 (sec) , antiderivative size = 221, normalized size of antiderivative = 2.19

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^4} dx = \frac{11Bb^2d^3 - 2Aa^2e^3 - 2(2Bab + Ab^2)d^2e - (Ba^2 + 2Aab)de^2 + 6(3Bb^2de^2 - (2Bab + Ab^2)e^3)x^2 + 3}{6(e^7x^3 + 3}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^4,x, algorithm="fricas")`

output `1/6*(11*B*b^2*d^3 - 2*A*a^2*e^3 - 2*(2*B*a*b + A*b^2)*d^2*e - (B*a^2 + 2*A*a*b)*d*e^2 + 6*(3*B*b^2*d*e^2 - (2*B*a*b + A*b^2)*e^3)*x^2 + 3*(9*B*b^2*d^2*e - 2*(2*B*a*b + A*b^2)*d*e^2 - (B*a^2 + 2*A*a*b)*e^3)*x + 6*(B*b^2*e^3*x^3 + 3*B*b^2*d*e^2*x^2 + 3*B*b^2*d^2*e*x + B*b^2*d^3)*log(e*x + d))/(e^7*x^3 + 3*d*e^6*x^2 + 3*d^2*e^5*x + d^3*e^4)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 211 vs. $2(88) = 176$.

Time = 2.36 (sec) , antiderivative size = 211, normalized size of antiderivative = 2.09

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^4} dx = \frac{Bb^2 \log(d+ex)}{e^4} + \frac{-2Aa^2e^3 - 2Aabde^2 - 2Ab^2d^2e - Ba^2de^2 - 4Babd^2e + 11Bb^2d^3 + x^2(-6Ab^2e^3 - 12Babe^3 + 18Bb^2d^3)}{6d^3e^4 + 18d^2e^5x + 18de^6x^2 + 6e^7x^3}$$

input `integrate((b*x+a)**2*(B*x+A)/(e*x+d)**4,x)`

output `B*b**2*log(d + e*x)/e**4 + (-2*A*a**2*e**3 - 2*A*a*b*d*e**2 - 2*A*b**2*d**2*e - B*a**2*d*e**2 - 4*B*a*b*d**2*e + 11*B*b**2*d**3 + x**2*(-6*A*b**2*e**3 - 12*B*a*b*e**3 + 18*B*b**2*d*e**2) + x*(-6*A*a*b*e**3 - 6*A*b**2*d*e**2 - 3*B*a**2*e**3 - 12*B*a*b*d*e**2 + 27*B*b**2*d**2*e))/(6*d**3*e**4 + 18*d**2*e**5*x + 18*d*e**6*x**2 + 6*e**7*x**3)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.76

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^4} dx = \frac{Bb^2 \ln(d+ex)}{e^4} - \frac{\frac{Ba^2de^2+2Aa^2e^3+4Babd^2e+2Aabde^2-11Bb^2d^3+2Ab^2d^2e}{6e^4} + \frac{x(Ba^2e^2+4Babde+2Aabe^2-9Bb^2d^2+2Ab^2de)}{2e^3} + \frac{bx^2(Ab^2e+2B^2a^2e-3B^2bd)}{e^2}}{d^3+3d^2ex+3de^2x^2+e^3x^3}$$

input `int(((A + B*x)*(a + b*x)^2)/(d + e*x)^4,x)`output `(B*b^2*log(d + e*x))/e^4 - ((2*A*a^2*e^3 - 11*B*b^2*d^3 + 2*A*b^2*d^2*e + B*a^2*d*e^2 + 2*A*a*b*d*e^2 + 4*B*a*b*d^2*e)/(6*e^4) + (x*(B*a^2*e^2 - 9*B*b^2*d^2 + 2*A*a*b*e^2 + 2*A*b^2*d*e + 4*B*a*b*d*e))/(2*e^3) + (b*x^2*(A*b*e + 2*B*a*e - 3*B*b*d))/e^2)/(d^3 + e^3*x^3 + 3*d*e^2*x^2 + 3*d^2*e*x)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.78

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^4} dx = \frac{6 \log(ex+d)b^3d^4 + 18 \log(ex+d)b^3d^3ex + 18 \log(ex+d)b^3d^2e^2x^2 + 6 \log(ex+d)b^3de^3x^3 - 2a^3de^3 - 6de^4(e^3x^3 + 3de^2x^2 + 3d^2ex + d^3)}{6de^4(e^3x^3 + 3de^2x^2 + 3d^2ex + d^3)}$$

input `int((b*x+a)^2*(B*x+A)/(e*x+d)^4,x)`output `(6*log(d + e*x)*b**3*d**4 + 18*log(d + e*x)*b**3*d**3*e*x + 18*log(d + e*x)*b**3*d**2*e**2*x**2 + 6*log(d + e*x)*b**3*d*e**3*x**3 - 2*a**3*d*e**3 - 3*a**2*b*d**2*e**2 - 9*a**2*b*d*e**3*x + 6*a*b**2*e**4*x**3 + 5*b**3*d**4 + 9*b**3*d**3*e*x - 6*b**3*d*e**3*x**3)/(6*d*e**4*(d**3 + 3*d**2*e*x + 3*d*e**2*x**2 + e**3*x**3))`

3.21 $\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^5} dx$

Optimal result	252
Mathematica [A] (verified)	252
Rubi [A] (verified)	253
Maple [A] (verified)	254
Fricas [B] (verification not implemented)	255
Sympy [B] (verification not implemented)	255
Maxima [B] (verification not implemented)	256
Giac [B] (verification not implemented)	256
Mupad [B] (verification not implemented)	257
Reduce [B] (verification not implemented)	257

Optimal result

Integrand size = 20, antiderivative size = 86

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^5} dx = -\frac{(Bd - Ae)(a+bx)^3}{4e(bd - ae)(d+ex)^4} + \frac{(3bBd + Abe - 4aBe)(a+bx)^3}{12e(bd - ae)^2(d+ex)^3}$$

output

```
-1/4*(-A*e+B*d)*(b*x+a)^3/e/(-a*e+b*d)/(e*x+d)^4+1/12*(A*b*e-4*B*a*e+3*B*b*d)*(b*x+a)^3/e/(-a*e+b*d)^2/(e*x+d)^3
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.45

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^5} dx = \frac{a^2e^2(3Ae + B(d + 4ex)) + 2abe(Ae(d + 4ex) + B(d^2 + 4dex + 6e^2x^2)) + b^2(Ae(d^2 + 4dex + 6e^2x^2) - 3Bd(d + 4ex) + 3Ae^2x)}{12e^4(d+ex)^4}$$

input

```
Integrate[((a + b*x)^2*(A + B*x))/(d + e*x)^5,x]
```

output

$$\frac{-1/12*(a^2*e^2*(3*A*e + B*(d + 4*e*x)) + 2*a*b*e*(A*e*(d + 4*e*x) + B*(d^2 + 4*d*e*x + 6*e^2*x^2)) + b^2*(A*e*(d^2 + 4*d*e*x + 6*e^2*x^2) + 3*B*(d^3 + 4*d^2*e*x + 6*d*e^2*x^2 + 4*e^3*x^3))}{(e^4*(d + e*x)^4)}$$

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {87, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^5} dx$$

$$\downarrow 87$$

$$\frac{(-4aBe + Abe + 3bBd) \int \frac{(a+bx)^2}{(d+ex)^4} dx}{4e(bd - ae)} - \frac{(a + bx)^3(Bd - Ae)}{4e(d + ex)^4(bd - ae)}$$

$$\downarrow 48$$

$$\frac{(a + bx)^3(-4aBe + Abe + 3bBd)}{12e(d + ex)^3(bd - ae)^2} - \frac{(a + bx)^3(Bd - Ae)}{4e(d + ex)^4(bd - ae)}$$

input

$$\text{Int}[(a + b*x)^2*(A + B*x)/(d + e*x)^5, x]$$

output

$$\frac{-1/4*((B*d - A*e)*(a + b*x)^3)/(e*(b*d - a*e)*(d + e*x)^4) + ((3*b*B*d + A*b*e - 4*a*B*e)*(a + b*x)^3)/(12*e*(b*d - a*e)^2*(d + e*x)^3)}$$

Defintions of rubi rules used

```
rule 48 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[[a + b*x]^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.77

method	result
risch	$\frac{-\frac{b^2 B x^3}{e} - \frac{b(Abe+2Bae+3Bbd)x^2}{2e^2} - \frac{(2Aab e^2 + A b^2 de + B a^2 e^2 + 2Babde + 3b^2 B d^2)x}{3e^3} - \frac{3a^2 A e^3 + 2Aabd e^2 + A b^2 d^2 e + B a^2 d e^2 + 2Babd^2}{12e^4}}{(ex+d)^4}$
norman	$\frac{-\frac{b^2 B x^3}{e} - \frac{(A b^2 e + 2Babe + 3b^2 Bd)x^2}{2e^2} - \frac{(2Aab e^2 + A b^2 de + B a^2 e^2 + 2Babde + 3b^2 B d^2)x}{3e^3} - \frac{3a^2 A e^3 + 2Aabd e^2 + A b^2 d^2 e + B a^2 d e^2 + 2Babd^2}{12e^4}}{(ex+d)^4}$
default	$-\frac{b^2 B}{e^4(ex+d)} - \frac{b(Abe+2Bae-3Bbd)}{2e^4(ex+d)^2} - \frac{a^2 A e^3 - 2Aabd e^2 + A b^2 d^2 e - B a^2 d e^2 + 2Babd^2 e - b^2 B d^3}{4e^4(ex+d)^4} - \frac{2Aab e^2 - 2A b^2 de + 3a^2 A d^2}{3e^4}$
gospers	$-\frac{12b^2 B x^3 e^3 + 6A x^2 b^2 e^3 + 12B x^2 ab e^3 + 18B x^2 b^2 d e^2 + 8Axab e^3 + 4Ax b^2 d e^2 + 4Bx a^2 e^3 + 8Bxabd e^2 + 12Bx b^2 d^2 e + 3a^2 A d^2}{12(ex+d)^4 e^4}$
parallelrisch	$-\frac{12b^2 B x^3 e^3 + 6A x^2 b^2 e^3 + 12B x^2 ab e^3 + 18B x^2 b^2 d e^2 + 8Axab e^3 + 4Ax b^2 d e^2 + 4Bx a^2 e^3 + 8Bxabd e^2 + 12Bx b^2 d^2 e + 3a^2 A d^2}{12(ex+d)^4 e^4}$
orering	$-\frac{12b^2 B x^3 e^3 + 6A x^2 b^2 e^3 + 12B x^2 ab e^3 + 18B x^2 b^2 d e^2 + 8Axab e^3 + 4Ax b^2 d e^2 + 4Bx a^2 e^3 + 8Bxabd e^2 + 12Bx b^2 d^2 e + 3a^2 A d^2}{12(ex+d)^4 e^4}$

```
input int((b*x+a)^2*(B*x+A)/(e*x+d)^5,x,method=_RETURNVERBOSE)
```

```
output (-b^2*B/e*x^3-1/2*b*(A*b*e+2*B*a*e+3*B*b*d)/e^2*x^2-1/3*(2*A*a*b*e^2+A*b^2
*d*e+B*a^2*e^2+2*B*a*b*d*e+3*B*b^2*d^2)/e^3*x-1/12*(3*A*a^2*e^3+2*A*a*b*d*
e^2+A*b^2*d^2*e+B*a^2*d*e^2+2*B*a*b*d^2*e+3*B*b^2*d^3)/e^4)/(e*x+d)^4
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 187 vs. $2(82) = 164$.

Time = 0.08 (sec) , antiderivative size = 187, normalized size of antiderivative = 2.17

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^5} dx = \frac{12Bb^2e^3x^3 + 3Bb^2d^3 + 3Aa^2e^3 + (2Bab + Ab^2)d^2e + (Ba^2 + 2Aab)de^2 + 6(3Bb^2de^2 + (2Bab + Aa^2)e^3)}{12(e^8x^4 + 4de^7x^3 + 6d^2e^6x^2 + 4d^3e^5x + d^4e^4)}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^5,x, algorithm="fricas")`

output

```
-1/12*(12*B*b^2*e^3*x^3 + 3*B*b^2*d^3 + 3*A*a^2*e^3 + (2*B*a*b + A*b^2)*d^2*e + (B*a^2 + 2*A*a*b)*d*e^2 + 6*(3*B*b^2*d*e^2 + (2*B*a*b + A*b^2)*e^3)*x^2 + 4*(3*B*b^2*d^2*e + (2*B*a*b + A*b^2)*d*e^2 + (B*a^2 + 2*A*a*b)*e^3)*x)/(e^8*x^4 + 4*d*e^7*x^3 + 6*d^2*e^6*x^2 + 4*d^3*e^5*x + d^4*e^4)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 223 vs. $2(75) = 150$.

Time = 4.64 (sec) , antiderivative size = 223, normalized size of antiderivative = 2.59

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^5} dx = \frac{-3Aa^2e^3 - 2Aabde^2 - Ab^2d^2e - Ba^2de^2 - 2Babd^2e - 3Bb^2d^3 - 12Bb^2e^3x^3 + x^2(-6Ab^2e^3 - 12Babe^3 - 6Aa^2e^3) + x(-8Aa^2bde^3 - 4Aa^2b^2de^2 - 4Bab^2d^2e^2 - 8Bab^2d^2e^2 - 12Bb^2d^2e^2)}{12d^4e^4 + 48d^3e^5x + 72d^2e^6x^2 + 48de^7x^3 + 12e^8x^4}$$

input `integrate((b*x+a)**2*(B*x+A)/(e*x+d)**5,x)`

output

```
(-3*A*a**2*e**3 - 2*A*a*b*d*e**2 - A*b**2*d**2*e - B*a**2*d*e**2 - 2*B*a*b*d**2*e - 3*B*b**2*d**3 - 12*B*b**2*e**3*x**3 + x**2*(-6*A*b**2*e**3 - 12*B*a*b*e**3 - 18*B*b**2*d*e**2) + x*(-8*A*a*b*d*e**3 - 4*A*b**2*d*e**2 - 4*B*a**2*e**3 - 8*B*a*b*d*e**2 - 12*B*b**2*d**2*e))/(12*d**4*e**4 + 48*d**3*e**5*x + 72*d**2*e**6*x**2 + 48*d*e**7*x**3 + 12*e**8*x**4)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 187 vs. $2(82) = 164$.

Time = 0.04 (sec) , antiderivative size = 187, normalized size of antiderivative = 2.17

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^5} dx = \frac{12 Bb^2 e^3 x^3 + 3 Bb^2 d^3 + 3 Aa^2 e^3 + (2 Bab + Ab^2)d^2 e + (Ba^2 + 2 Aab)de^2 + 6 (3 Bb^2 de^2 + (2 Bab + Aa^2)e^3)}{12 (e^8 x^4 + 4 de^7 x^3 + 6 d^2 e^6 x^2 + 4 d^3 e^5 x + d^4 e^4)}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^5,x, algorithm="maxima")`

output `-1/12*(12*B*b^2*e^3*x^3 + 3*B*b^2*d^3 + 3*A*a^2*e^3 + (2*B*a*b + A*b^2)*d^2*e + (B*a^2 + 2*A*a*b)*d*e^2 + 6*(3*B*b^2*d*e^2 + (2*B*a*b + A*b^2)*e^3)*x^2 + 4*(3*B*b^2*d^2*e + (2*B*a*b + A*b^2)*d*e^2 + (B*a^2 + 2*A*a*b)*e^3)*x)/(e^8*x^4 + 4*d*e^7*x^3 + 6*d^2*e^6*x^2 + 4*d^3*e^5*x + d^4*e^4)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 232 vs. $2(82) = 164$.

Time = 0.12 (sec) , antiderivative size = 232, normalized size of antiderivative = 2.70

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^5} dx = \frac{\frac{12 Bab}{(ex+d)^2} + \frac{6 Ab^2}{(ex+d)^2} - \frac{16 Babd}{(ex+d)^3} - \frac{8 Ab^2 d}{(ex+d)^3} + \frac{6 Babd^2}{(ex+d)^4} + \frac{3 Ab^2 d^2}{(ex+d)^4} + \frac{12 Bb^2}{(ex+d)e} - \frac{18 Bb^2 d}{(ex+d)^2 e} + \frac{12 Bb^2 d^2}{(ex+d)^3 e} - \frac{3 Bb^2 d^3}{(ex+d)^4 e} + \frac{4 Bb^2 d^3}{(ex+d)^4 e}}{12 e^3}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^5,x, algorithm="giac")`

output `-1/12*(12*B*a*b/(e*x + d)^2 + 6*A*b^2/(e*x + d)^2 - 16*B*a*b*d/(e*x + d)^3 - 8*A*b^2*d/(e*x + d)^3 + 6*B*a*b*d^2/(e*x + d)^4 + 3*A*b^2*d^2/(e*x + d)^4 + 12*B*b^2/((e*x + d)*e) - 18*B*b^2*d/((e*x + d)^2*e) + 12*B*b^2*d^2/((e*x + d)^3*e) - 3*B*b^2*d^3/((e*x + d)^4*e) + 4*B*a^2*e/(e*x + d)^3 + 8*A*a*b*e/(e*x + d)^3 - 3*B*a^2*d*e/(e*x + d)^4 - 6*A*a*b*d*e/(e*x + d)^4 + 3*A*a^2*e^2/(e*x + d)^4)/e^3`

Mupad [B] (verification not implemented)

Time = 0.98 (sec) , antiderivative size = 184, normalized size of antiderivative = 2.14

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^5} dx = \frac{\frac{Ba^2de^2 + 3Aa^2e^3 + 2Babd^2e + 2Aabde^2 + 3Bb^2d^3 + Ab^2d^2e}{12e^4} + \frac{x(Ba^2e^2 + 2Babde + 2Aabe^2 + 3Bb^2d^2 + Ab^2de)}{3e^3} + \frac{bx^2(Abe + Bb^2d)}{3e^2}}{d^4 + 4d^3ex + 6d^2e^2x^2 + 4de^3x^3 + e^4x^4}$$

input `int(((A + B*x)*(a + b*x)^2)/(d + e*x)^5,x)`output `-((3*A*a^2*e^3 + 3*B*b^2*d^3 + A*b^2*d^2*e + B*a^2*d*e^2 + 2*A*a*b*d*e^2 + 2*B*a*b*d^2*e)/(12*e^4) + (x*(B*a^2*e^2 + 3*B*b^2*d^2 + 2*A*a*b*e^2 + A*b^2*d*e + 2*B*a*b*d*e))/(3*e^3) + (b*x^2*(A*b*e + 2*B*a*e + 3*B*b*d))/(2*e^2) + (B*b^2*x^3)/e)/(d^4 + e^4*x^4 + 4*d*e^3*x^3 + 6*d^2*e^2*x^2 + 4*d^3*e*x)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.42

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^5} dx = \frac{b^3e^3x^4 - 6ab^2de^2x^2 - 4a^2bde^2x - 4ab^2d^2ex - a^3de^2 - a^2bd^2e - ab^2d^3}{4de^3(e^4x^4 + 4de^3x^3 + 6d^2e^2x^2 + 4d^3ex + d^4)}$$

input `int((b*x+a)^2*(B*x+A)/(e*x+d)^5,x)`output `(- a**3*d*e**2 - a**2*b*d**2*e - 4*a**2*b*d*e**2*x - a*b**2*d**3 - 4*a*b**2*d**2*e*x - 6*a*b**2*d*e**2*x**2 + b**3*e**3*x**4)/(4*d*e**3*(d**4 + 4*d**3*e*x + 6*d**2*e**2*x**2 + 4*d*e**3*x**3 + e**4*x**4))`

3.22 $\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^6} dx$

Optimal result	258
Mathematica [A] (verified)	258
Rubi [A] (verified)	259
Maple [A] (verified)	260
Fricas [A] (verification not implemented)	261
Sympy [B] (verification not implemented)	261
Maxima [A] (verification not implemented)	262
Giac [A] (verification not implemented)	262
Mupad [B] (verification not implemented)	263
Reduce [B] (verification not implemented)	263

Optimal result

Integrand size = 20, antiderivative size = 120

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^6} dx = \frac{(bd-ae)^2(Bd-Ae)}{5e^4(d+ex)^5} - \frac{(bd-ae)(3bBd-2Abe-aBe)}{4e^4(d+ex)^4} + \frac{b(3bBd-Abe-2aBe)}{3e^4(d+ex)^3} - \frac{b^2B}{2e^4(d+ex)^2}$$

output

```
1/5*(-a*e+b*d)^2*(-A*e+B*d)/e^4/(e*x+d)^5-1/4*(-a*e+b*d)*(-2*A*b*e-B*a*e+3
*B*b*d)/e^4/(e*x+d)^4+1/3*b*(-A*b*e-2*B*a*e+3*B*b*d)/e^4/(e*x+d)^3-1/2*b^2
*B/e^4/(e*x+d)^2
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.08

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^6} dx = \frac{3a^2e^2(4Ae+B(d+5ex)) + 2abe(3Ae(d+5ex) + 2B(d^2+5dex+10e^2x^2)) + b^2(2Ae(d^2+5dex+10e^2x^2))}{60e^4(d+ex)^5}$$

input

```
Integrate[((a + b*x)^2*(A + B*x))/(d + e*x)^6,x]
```

output

$$\frac{-1/60*(3*a^2*e^2*(4*A*e + B*(d + 5*e*x)) + 2*a*b*e*(3*A*e*(d + 5*e*x) + 2*B*(d^2 + 5*d*e*x + 10*e^2*x^2)) + b^2*(2*A*e*(d^2 + 5*d*e*x + 10*e^2*x^2) + 3*B*(d^3 + 5*d^2*e*x + 10*d*e^2*x^2 + 10*e^3*x^3))}{(e^4*(d + e*x)^5)}$$

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^6} dx$$

↓ 86

$$\int \left(\frac{b(2aBe + Abe - 3bBd)}{e^3(d + ex)^4} + \frac{(ae - bd)(aBe + 2Abe - 3bBd)}{e^3(d + ex)^5} + \frac{(ae - bd)^2(Ae - Bd)}{e^3(d + ex)^6} + \frac{b^2B}{e^3(d + ex)^3} \right) dx$$

↓ 2009

$$\frac{b(-2aBe - Abe + 3bBd)}{3e^4(d + ex)^3} - \frac{(bd - ae)(-aBe - 2Abe + 3bBd)}{4e^4(d + ex)^4} + \frac{(bd - ae)^2(Bd - Ae)}{5e^4(d + ex)^5} - \frac{b^2B}{2e^4(d + ex)^2}$$

input

$$\text{Int}[(a + b*x)^2*(A + B*x)/(d + e*x)^6, x]$$

output

$$\frac{((b*d - a*e)^2*(B*d - A*e))/(5*e^4*(d + e*x)^5) - ((b*d - a*e)*(3*b*B*d - 2*A*b*e - a*B*e))/(4*e^4*(d + e*x)^4) + (b*(3*b*B*d - A*b*e - 2*a*B*e))/(3*e^4*(d + e*x)^3) - (b^2*B)/(2*e^4*(d + e*x)^2)}$$

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.68

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^6} dx = \frac{\frac{3Ba^2de^2 + 12Aa^2e^3 + 4Babd^2e + 6Aabde^2 + 3Bb^2d^3 + 2Ab^2d^2e}{60e^4} + \frac{x(3Ba^2e^2 + 4Babde + 6Aabe^2 + 3Bb^2d^2 + 2Ab^2de)}{12e^3} + \frac{bx^2}{2e^2}}{d^5 + 5d^4ex + 10d^3e^2x^2 + 10d^2e^3x^3 + 5de^4x^4 + e^5x^5}$$

input

```
int(((A + B*x)*(a + b*x)^2)/(d + e*x)^6,x)
```

output

```
-((12*A*a^2*e^3 + 3*B*b^2*d^3 + 2*A*b^2*d^2*e + 3*B*a^2*d*e^2 + 6*A*a*b*d*e^2 + 4*B*a*b*d^2*e)/(60*e^4) + (x*(3*B*a^2*e^2 + 3*B*b^2*d^2 + 6*A*a*b*e^2 + 2*A*b^2*d*e + 4*B*a*b*d*e))/(12*e^3) + (b*x^2*(2*A*b*e + 4*B*a*e + 3*B*b*d))/(6*e^2) + (B*b^2*x^3)/(2*e))/(d^5 + e^5*x^5 + 5*d*e^4*x^4 + 10*d^3*e^2*x^2 + 10*d^2*e^3*x^3 + 5*d^4*e*x)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.32

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^6} dx = \frac{-10b^3e^3x^3 - 20ab^2e^3x^2 - 10b^3de^2x^2 - 15a^2be^3x - 10ab^2de^2x - 5b^3d^2ex - 4a^3e^3 - 3a^2bde^2 - 2ab^2d^2e}{20e^4(e^5x^5 + 5de^4x^4 + 10d^2e^3x^3 + 10d^3e^2x^2 + 5d^4ex + d^5)}$$

input

```
int((b*x+a)^2*(B*x+A)/(e*x+d)^6,x)
```

output

```
(-4*a**3*e**3 - 3*a**2*b*d*e**2 - 15*a**2*b*e**3*x - 2*a*b**2*d**2*e - 10*a*b**2*d*e**2*x - 20*a*b**2*e**3*x**2 - b**3*d**3 - 5*b**3*d**2*e*x - 10*b**3*d*e**2*x**2 - 10*b**3*e**3*x**3)/(20*e**4*(d**5 + 5*d**4*e*x + 10*d**3*e**2*x**2 + 10*d**2*e**3*x**3 + 5*d*e**4*x**4 + e**5*x**5))
```

3.23 $\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^7} dx$

Optimal result	264
Mathematica [A] (verified)	264
Rubi [A] (verified)	265
Maple [A] (verified)	266
Fricas [A] (verification not implemented)	267
Sympy [B] (verification not implemented)	267
Maxima [A] (verification not implemented)	268
Giac [A] (verification not implemented)	268
Mupad [B] (verification not implemented)	269
Reduce [B] (verification not implemented)	269

Optimal result

Integrand size = 20, antiderivative size = 120

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^7} dx = \frac{(bd-ae)^2(Bd-Ae)}{6e^4(d+ex)^6} - \frac{(bd-ae)(3bBd-2Abe-aBe)}{5e^4(d+ex)^5} + \frac{b(3bBd-Abe-2aBe)}{4e^4(d+ex)^4} - \frac{b^2B}{3e^4(d+ex)^3}$$

output

```
1/6*(-a*e+b*d)^2*(-A*e+B*d)/e^4/(e*x+d)^6-1/5*(-a*e+b*d)*(-2*A*b*e-B*a*e+3
*B*b*d)/e^4/(e*x+d)^5+1/4*b*(-A*b*e-2*B*a*e+3*B*b*d)/e^4/(e*x+d)^4-1/3*b^2
*B/e^4/(e*x+d)^3
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.05

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^7} dx = \frac{2a^2e^2(5Ae+B(d+6ex)) + 2abe(2Ae(d+6ex) + B(d^2+6dex+15e^2x^2)) + b^2(Ae(d^2+6dex+15e^2x^2))}{60e^4(d+ex)^6}$$

input

```
Integrate[((a + b*x)^2*(A + B*x))/(d + e*x)^7,x]
```

output

$$\frac{-1/60*(2*a^2*e^2*(5*A*e + B*(d + 6*e*x)) + 2*a*b*e*(2*A*e*(d + 6*e*x) + B*(d^2 + 6*d*e*x + 15*e^2*x^2)) + b^2*(A*e*(d^2 + 6*d*e*x + 15*e^2*x^2) + B*(d^3 + 6*d^2*e*x + 15*d*e^2*x^2 + 20*e^3*x^3))}{e^4*(d + e*x)^6}$$

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^7} dx$$

↓ 86

$$\int \left(\frac{b(2aBe + Abe - 3bBd)}{e^3(d + ex)^5} + \frac{(ae - bd)(aBe + 2Abe - 3bBd)}{e^3(d + ex)^6} + \frac{(ae - bd)^2(Ae - Bd)}{e^3(d + ex)^7} + \frac{b^2B}{e^3(d + ex)^4} \right) dx$$

↓ 2009

$$\frac{b(-2aBe - Abe + 3bBd)}{4e^4(d + ex)^4} - \frac{(bd - ae)(-aBe - 2Abe + 3bBd)}{5e^4(d + ex)^5} + \frac{(bd - ae)^2(Bd - Ae)}{6e^4(d + ex)^6} - \frac{b^2B}{3e^4(d + ex)^3}$$

input

$$\text{Int}[(a + b*x)^2*(A + B*x)/(d + e*x)^7, x]$$

output

$$\frac{((b*d - a*e)^2*(B*d - A*e))/(6*e^4*(d + e*x)^6) - ((b*d - a*e)*(3*b*B*d - 2*A*b*e - a*B*e))/(5*e^4*(d + e*x)^5) + (b*(3*b*B*d - A*b*e - 2*a*B*e))/(4*e^4*(d + e*x)^4) - (b^2*B)/(3*e^4*(d + e*x)^3)}$$

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.26

method	result
risch	$\frac{-\frac{b^2 B x^3}{3e} - \frac{b(Abe+2Bae+Bbd)x^2}{4e^2} - \frac{(4Aab e^2 + A b^2 de + 2B a^2 e^2 + 2Babde + b^2 B d^2)x}{10e^3} - \frac{10a^2 A e^3 + 4Aabd e^2 + A b^2 d^2 e + 2B a^2 d e^2 + 2Babd}{60e^4}}{(ex+d)^6}$
default	$-\frac{a^2 A e^3 - 2Aabd e^2 + A b^2 d^2 e - B a^2 d e^2 + 2Babd^2 e - b^2 B d^3}{6e^4(ex+d)^6} - \frac{2Aab e^2 - 2A b^2 de + B a^2 e^2 - 4Babde + 3b^2 B d^2}{5e^4(ex+d)^5} - \frac{b(Abe+2Bae+Bbd)}{4e^4}$
gospers	$-\frac{20b^2 B x^3 e^3 + 15A x^2 b^2 e^3 + 30B x^2 ab e^3 + 15B x^2 b^2 d e^2 + 24Axab e^3 + 6Ax b^2 d e^2 + 12Bx a^2 e^3 + 12Bxabd e^2 + 6Bx b^2 d^2 e + 10A a^2 e^3}{60e^4(ex+d)^6}$
orering	$-\frac{20b^2 B x^3 e^3 + 15A x^2 b^2 e^3 + 30B x^2 ab e^3 + 15B x^2 b^2 d e^2 + 24Axab e^3 + 6Ax b^2 d e^2 + 12Bx a^2 e^3 + 12Bxabd e^2 + 6Bx b^2 d^2 e + 10A a^2 e^3}{60e^4(ex+d)^6}$
norman	$\frac{-\frac{b^2 B x^3}{3e} - \frac{(A b^2 e^3 + 2Bab e^3 + b^2 B d e^2)x^2}{4e^4} - \frac{(4Aab e^4 + A b^2 d e^3 + 2B a^2 e^4 + 2Babd e^3 + b^2 B d^2 e^2)x}{10e^5} - \frac{10a^2 A e^5 + 4Aabd e^4 + A b^2 d^2 e^3 + 2Babd e^4}{60e^6}}{(ex+d)^6}$
parallelrisch	$-\frac{20B b^2 x^3 e^5 + 15A b^2 e^5 x^2 + 30Bab e^5 x + 15B b^2 d e^4 x^2 + 24Aab e^5 x + 6A b^2 d e^4 x + 12B a^2 e^5 x + 12Babd e^4 x + 6B b^2 d^2 e^3 x + 10A a^2 e^5}{60e^6(ex+d)^6}$

```
input int((b*x+a)^2*(B*x+A)/(e*x+d)^7,x,method=_RETURNVERBOSE)
```

```
output (-1/3*b^2*B/e*x^3-1/4*b/e^2*(A*b*e+2*B*a*e+B*b*d)*x^2-1/10/e^3*(4*A*a*b*e^2+A*b^2*d*e+2*B*a^2*e^2+2*B*a*b*d*e+B*b^2*d^2)*x-1/60/e^4*(10*A*a^2*e^3+4*A*a*b*d*e^2+A*b^2*d^2*e+2*B*a^2*d*e^2+2*B*a*b*d^2*e+B*b^2*d^3))/(e*x+d)^6
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.73

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^7} dx = \frac{20Bb^2e^3x^3 + Bb^2d^3 + 10Aa^2e^3 + (2Bab + Ab^2)d^2e + 2(Ba^2 + 2Aab)de^2 + 15(Bb^2de^2 + (2Bab + Ab^2)d^2e) + 15Aa^2e^3}{60(e^{10}x^6 + 6de^9x^5 + 15d^2e^8x^4 + 20d^3e^7x^3 + 15d^4e^6x^2 + 6d^5e^5x + d^6e^4)}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^7,x, algorithm="fricas")`

output `-1/60*(20*B*b^2*e^3*x^3 + B*b^2*d^3 + 10*A*a^2*e^3 + (2*B*a*b + A*b^2)*d^2*e + 2*(B*a^2 + 2*A*a*b)*d*e^2 + 15*(B*b^2*d*e^2 + (2*B*a*b + A*b^2)*e^3)*x^2 + 6*(B*b^2*d^2*e + (2*B*a*b + A*b^2)*d*e^2 + 2*(B*a^2 + 2*A*a*b)*e^3)*x)/(e^10*x^6 + 6*d*e^9*x^5 + 15*d^2*e^8*x^4 + 20*d^3*e^7*x^3 + 15*d^4*e^6*x^2 + 6*d^5*e^5*x + d^6*e^4)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 246 vs. 2(114) = 228.

Time = 15.41 (sec) , antiderivative size = 246, normalized size of antiderivative = 2.05

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^7} dx = \frac{-10Aa^2e^3 - 4Aabde^2 - Ab^2d^2e - 2Ba^2de^2 - 2Babd^2e - Bb^2d^3 - 20Bb^2e^3x^3 + x^2(-15Ab^2e^3 - 30Babde^2) + x(-24Aa^2e^3 - 6Aa^2de^2 - 12Aa^2bde^2 - 6Aa^2bd^2e) + x^3(-15Ab^2e^3 - 30Babde^2 - 15Bb^2d^2e)}{60d^6e^4 + 360d^5e^5x + 900d^4e^6x^2 + 1200d^3e^7x^3 + 900d^2e^8x^4 + 360d^5e^5x + 60e^10x^6}$$

input `integrate((b*x+a)**2*(B*x+A)/(e*x+d)**7,x)`

output `(-10*A*a**2*e**3 - 4*A*a*b*d*e**2 - A*b**2*d**2*e - 2*B*a**2*d*e**2 - 2*B*a*b*d**2*e - B*b**2*d**3 - 20*B*b**2*e**3*x**3 + x**2*(-15*A*b**2*e**3 - 30*B*a*b*d*e**2) + x*(-24*A*a**2*e**3 - 6*A*a**2*d*e**2 - 12*B*a**2*e**3 - 12*B*a*b*d*e**2 - 6*B*b**2*d**2*e))/(60*d**6*e**4 + 360*d**5*e**5*x + 900*d**4*e**6*x**2 + 1200*d**3*e**7*x**3 + 900*d**2*e**8*x**4 + 360*d**5*e**5*x + 60*e**10*x**6)`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.73

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^7} dx = \frac{20 Bb^2e^3x^3 + Bb^2d^3 + 10 Aa^2e^3 + (2 Bab + Ab^2)d^2e + 2 (Ba^2 + 2 Aab)de^2 + 15 (Bb^2de^2 + (2 Bab + Ab^2)d^2e + 2 Aa^2e^3)}{60 (e^{10}x^6 + 6 de^9x^5 + 15 d^2e^8x^4 + 20 d^3e^7x^3 + 15 d^4e^6x^2 + 6 d^5e^5x + d^6e^4)}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^7,x, algorithm="maxima")`output `-1/60*(20*B*b^2*e^3*x^3 + B*b^2*d^3 + 10*A*a^2*e^3 + (2*B*a*b + A*b^2)*d^2*e + 2*(B*a^2 + 2*A*a*b)*d*e^2 + 15*(B*b^2*d*e^2 + (2*B*a*b + A*b^2)*e^3)*x^2 + 6*(B*b^2*d^2*e + (2*B*a*b + A*b^2)*d*e^2 + 2*(B*a^2 + 2*A*a*b)*e^3)*x)/(e^10*x^6 + 6*d*e^9*x^5 + 15*d^2*e^8*x^4 + 20*d^3*e^7*x^3 + 15*d^4*e^6*x^2 + 6*d^5*e^5*x + d^6*e^4)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.38

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^7} dx = \frac{20 Bb^2e^3x^3 + 15 Bb^2de^2x^2 + 30 Babe^3x^2 + 15 Ab^2e^3x^2 + 6 Bb^2d^2ex + 12 Babde^2x + 6 Ab^2de^2x + 12 Aa^2e^3}{60 (ex + d)^6 e^4}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^7,x, algorithm="giac")`output `-1/60*(20*B*b^2*e^3*x^3 + 15*B*b^2*d*e^2*x^2 + 30*B*a*b*e^3*x^2 + 15*A*b^2*e^3*x^2 + 6*B*b^2*d^2*e*x + 12*B*a*b*d*e^2*x + 6*A*b^2*d*e^2*x + 12*B*a^2*e^3*x + 24*A*a*b*e^3*x + B*b^2*d^3 + 2*B*a*b*d^2*e + A*b^2*d^2*e + 2*B*a^2*d*e^2 + 4*A*a*b*d*e^2 + 10*A*a^2*e^3)/((e*x + d)^6*e^4)`

Mupad [B] (verification not implemented)

Time = 0.98 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.72

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^7} dx = \frac{\frac{2Ba^2de^2 + 10Aa^2e^3 + 2Babd^2e + 4Aabde^2 + Bb^2d^3 + Ab^2d^2e}{60e^4} + \frac{x(2Ba^2e^2 + 2Babde + 4Aabe^2 + Bb^2d^2 + Ab^2de)}{10e^3} + \frac{bx^2(Abe^2 + 2Bbd^2 + 2Aabde)}{10e^3}}{d^6 + 6d^5ex + 15d^4e^2x^2 + 20d^3e^3x^3 + 15d^2e^4x^4 + 6de^5x^5 + e^6x^6}$$

input `int(((A + B*x)*(a + b*x)^2)/(d + e*x)^7,x)`output `-((10*A*a^2*e^3 + B*b^2*d^3 + A*b^2*d^2*e + 2*B*a^2*d*e^2 + 4*A*a*b*d*e^2 + 2*B*a*b*d^2*e)/(60*e^4) + (x*(2*B*a^2*e^2 + B*b^2*d^2 + 4*A*a*b*e^2 + A*b^2*d*e + 2*B*a*b*d*e))/(10*e^3) + (b*x^2*(A*b*e + 2*B*a*e + B*b*d))/(4*e^2) + (B*b^2*x^3)/(3*e))/(d^6 + e^6*x^6 + 6*d*e^5*x^5 + 15*d^4*e^2*x^2 + 20*d^3*e^3*x^3 + 15*d^2*e^4*x^4 + 6*d^5*e*x)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.42

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^7} dx = \frac{-20b^3e^3x^3 - 45ab^2e^3x^2 - 15b^3de^2x^2 - 36a^2be^3x - 18ab^2de^2x - 6b^3d^2ex - 10a^3e^3 - 6a^2bde^2 - 3ab^2d}{60e^4(e^6x^6 + 6de^5x^5 + 15d^2e^4x^4 + 20d^3e^3x^3 + 15d^4e^2x^2 + 6d^5ex + d^6)}$$

input `int((b*x+a)^2*(B*x+A)/(e*x+d)^7,x)`output `(-10*a**3*e**3 - 6*a**2*b*d*e**2 - 36*a**2*b*e**3*x - 3*a*b**2*d**2*e - 18*a*b**2*d*e**2*x - 45*a*b**2*e**3*x**2 - b**3*d**3 - 6*b**3*d**2*e*x - 15*b**3*d*e**2*x**2 - 20*b**3*e**3*x**3)/(60*e**4*(d**6 + 6*d**5*e*x + 15*d**4*e**2*x**2 + 20*d**3*e**3*x**3 + 15*d**2*e**4*x**4 + 6*d*e**5*x**5 + e**6*x**6))`

3.24 $\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^8} dx$

Optimal result	270
Mathematica [A] (verified)	270
Rubi [A] (verified)	271
Maple [A] (verified)	272
Fricas [B] (verification not implemented)	273
Sympy [B] (verification not implemented)	273
Maxima [B] (verification not implemented)	274
Giac [A] (verification not implemented)	274
Mupad [B] (verification not implemented)	275
Reduce [B] (verification not implemented)	275

Optimal result

Integrand size = 20, antiderivative size = 120

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^8} dx = \frac{(bd-ae)^2(Bd-Ae)}{7e^4(d+ex)^7} - \frac{(bd-ae)(3bBd-2Abe-aBe)}{6e^4(d+ex)^6} + \frac{b(3bBd-Abe-2aBe)}{5e^4(d+ex)^5} - \frac{b^2B}{4e^4(d+ex)^4}$$

output

```
1/7*(-a*e+b*d)^2*(-A*e+B*d)/e^4/(e*x+d)^7-1/6*(-a*e+b*d)*(-2*A*b*e-B*a*e+3*B*b*d)/e^4/(e*x+d)^6+1/5*b*(-A*b*e-2*B*a*e+3*B*b*d)/e^4/(e*x+d)^5-1/4*b^2*B/e^4/(e*x+d)^4
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.08

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^8} dx = \frac{10a^2e^2(6Ae+B(d+7ex)) + 4abe(5Ae(d+7ex) + 2B(d^2+7dex+21e^2x^2)) + b^2(4Ae(d^2+7dex+21e^2x^2))}{420e^4(d+ex)^7}$$

input

```
Integrate[((a + b*x)^2*(A + B*x))/(d + e*x)^8,x]
```

output

$$\frac{-1/420*(10*a^2*e^2*(6*A*e + B*(d + 7*e*x)) + 4*a*b*e*(5*A*e*(d + 7*e*x) + 2*B*(d^2 + 7*d*e*x + 21*e^2*x^2)) + b^2*(4*A*e*(d^2 + 7*d*e*x + 21*e^2*x^2) + 3*B*(d^3 + 7*d^2*e*x + 21*d*e^2*x^2 + 35*e^3*x^3))}{(e^4*(d + e*x)^7)}$$

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^8} dx$$

↓ 86

$$\int \left(\frac{b(2aBe + Abe - 3bBd)}{e^3(d + ex)^6} + \frac{(ae - bd)(aBe + 2Abe - 3bBd)}{e^3(d + ex)^7} + \frac{(ae - bd)^2(Ae - Bd)}{e^3(d + ex)^8} + \frac{b^2B}{e^3(d + ex)^5} \right) dx$$

↓ 2009

$$\frac{b(-2aBe - Abe + 3bBd)}{5e^4(d + ex)^5} - \frac{(bd - ae)(-aBe - 2Abe + 3bBd)}{6e^4(d + ex)^6} + \frac{(bd - ae)^2(Bd - Ae)}{7e^4(d + ex)^7} - \frac{b^2B}{4e^4(d + ex)^4}$$

input

$$\text{Int}[(a + b*x)^2*(A + B*x)/(d + e*x)^8, x]$$

output

$$\frac{((b*d - a*e)^2*(B*d - A*e))/(7*e^4*(d + e*x)^7) - ((b*d - a*e)*(3*b*B*d - 2*A*b*e - a*B*e))/(6*e^4*(d + e*x)^6) + (b*(3*b*B*d - A*b*e - 2*a*B*e))/(5*e^4*(d + e*x)^5) - (b^2*B)/(4*e^4*(d + e*x)^4)}$$

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0]) || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.31

method	result
risch	$\frac{-\frac{b^2 B x^3}{4e} - \frac{b(4Abe+8BAe+3Bbd)x^2}{20e^2} - \frac{(20Aab e^2+4A b^2 de+10B a^2 e^2+8Babde+3b^2 B d^2)x}{60e^3} - \frac{60a^2 A e^3+20Aabd e^2+4A b^2 d^2 e+10B a^2 d}{420e^4}}{(ex+d)^7}$
default	$-\frac{2Aab e^2-2A b^2 de+B a^2 e^2-4Babde+3b^2 B d^2}{6e^4(ex+d)^6} - \frac{a^2 A e^3-2Aabd e^2+A b^2 d^2 e-B a^2 d e^2+2Bab d^2 e-b^2 B d^3}{7e^4(ex+d)^7} - \frac{b(Abe+2A b^2 d)}{5e^4}$
gospers	$-\frac{105b^2 B x^3 e^3+84A x^2 b^2 e^3+168B x^2 ab e^3+63B x^2 b^2 d e^2+140Axab e^3+28Ax b^2 d e^2+70Bx a^2 e^3+56Bxabd e^2+21Bx b^2 d^2 e}{420e^4(ex+d)^7}$
orering	$-\frac{105b^2 B x^3 e^3+84A x^2 b^2 e^3+168B x^2 ab e^3+63B x^2 b^2 d e^2+140Axab e^3+28Ax b^2 d e^2+70Bx a^2 e^3+56Bxabd e^2+21Bx b^2 d^2 e}{420e^4(ex+d)^7}$
parallelrisc	$-\frac{105B b^2 x^3 e^6+84A b^2 e^6 x^2+168Bab e^6 x^2+63B b^2 d e^5 x^2+140Aab e^6 x+28A b^2 d e^5 x+70B a^2 e^6 x+56Babd e^5 x+21B b^2 d^2 e^6}{420e^7(ex+d)^7}$
norman	$\frac{-\frac{b^2 B x^3}{4e} - \frac{(4A b^2 e^4+8Bab e^4+3b^2 B d e^3)x^2}{20e^5} - \frac{(20abA e^5+4A b^2 d e^4+10a^2 B e^5+8Babd e^4+3B b^2 d^2 e^3)x}{60e^6} - \frac{60a^2 A e^6+20Aabd e^5+4A b^2 d^2 e^4}{420e^7}}{(ex+d)^7}$

```
input int((b*x+a)^2*(B*x+A)/(e*x+d)^8,x,method=_RETURNVERBOSE)
```

```
output (-1/4*b^2*B/e*x^3-1/20*b/e^2*(4*A*b*e+8*B*a*e+3*B*b*d)*x^2-1/60/e^3*(20*A*a*b*e^2+4*A*b^2*d*e+10*B*a^2*e^2+8*B*a*b*d*e+3*B*b^2*d^2)*x-1/420/e^4*(60*A*a^2*e^3+20*A*a*b*d*e^2+4*A*b^2*d^2*e+10*B*a^2*d*e^2+8*B*a*b*d^2*e+3*B*b^2*d^3))/(e*x+d)^7
```


output

$$\frac{-1/420*(105*B*b^2*e^3*x^3 + 63*B*b^2*d*e^2*x^2 + 168*B*a*b*e^3*x^2 + 84*A*b^2*e^3*x^2 + 21*B*b^2*d^2*e*x + 56*B*a*b*d*e^2*x + 28*A*b^2*d*e^2*x + 70*B*a^2*e^3*x + 140*A*a*b*e^3*x + 3*B*b^2*d^3 + 8*B*a*b*d^2*e + 4*A*b^2*d^2*e + 10*B*a^2*d*e^2 + 20*A*a*b*d*e^2 + 60*A*a^2*e^3)/((e*x + d)^7*e^4)}{1}$$

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.86

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^8} dx =$$

$$-\frac{10Ba^2de^2 + 60Aa^2e^3 + 8Babd^2e + 20Aabde^2 + 3Bb^2d^3 + 4Ab^2d^2e}{420e^4} + \frac{x(10Ba^2e^2 + 8Babde + 20Aabe^2 + 3Bb^2d^2 + 4Ab^2de)}{60e^3} +$$

$$\frac{d^7 + 7d^6ex + 21d^5e^2x^2 + 35d^4e^3x^3 + 35d^3e^4x^4 + 21d^2e^5x^5 + 7de^6x^6}{d^7 + 7d^6ex + 21d^5e^2x^2 + 35d^4e^3x^3 + 35d^3e^4x^4 + 21d^2e^5x^5 + 7de^6x^6}$$

input

```
int(((A + B*x)*(a + b*x)^2)/(d + e*x)^8,x)
```

output

$$\frac{-((60*A*a^2*e^3 + 3*B*b^2*d^3 + 4*A*b^2*d^2*e + 10*B*a^2*d*e^2 + 20*A*a*b*d*e^2 + 8*B*a*b*d^2*e)/(420*e^4) + (x*(10*B*a^2*e^2 + 3*B*b^2*d^2 + 20*A*a*b*e^2 + 4*A*b^2*d*e + 8*B*a*b*d*e))/(60*e^3) + (b*x^2*(4*A*b*e + 8*B*a*e + 3*B*b*d))/(20*e^2) + (B*b^2*x^3)/(4*e))/(d^7 + e^7*x^7 + 7*d*e^6*x^6 + 21*d^5*e^2*x^2 + 35*d^4*e^3*x^3 + 35*d^3*e^4*x^4 + 21*d^2*e^5*x^5 + 7*d^6*e*x)}{1}$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.51

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^8} dx$$

$$= \frac{-35b^3e^3x^3 - 84ab^2e^3x^2 - 21b^3de^2x^2 - 70a^2be^3x - 28ab^2de^2x - 7b^3d^2ex - 20a^3e^3 - 10a^2bde^2 - 4ab^2d^2e}{140e^4(e^7x^7 + 7de^6x^6 + 21d^2e^5x^5 + 35d^3e^4x^4 + 35d^4e^3x^3 + 21d^5e^2x^2 + 7d^6ex + d^7)}$$

input

```
int((b*x+a)^2*(B*x+A)/(e*x+d)^8,x)
```


output

```
( - 20*a**3*e**3 - 10*a**2*b*d*e**2 - 70*a**2*b*e**3*x - 4*a*b**2*d**2*e -  
28*a*b**2*d*e**2*x - 84*a*b**2*e**3*x**2 - b**3*d**3 - 7*b**3*d**2*e*x -  
21*b**3*d*e**2*x**2 - 35*b**3*e**3*x**3)/(140*e**4*(d**7 + 7*d**6*e*x + 21  
*d**5*e**2*x**2 + 35*d**4*e**3*x**3 + 35*d**3*e**4*x**4 + 21*d**2*e**5*x**  
5 + 7*d*e**6*x**6 + e**7*x**7))
```

3.25 $\int (a + bx)^3(A + Bx)(d + ex)^5 dx$

Optimal result	277
Mathematica [B] (verified)	278
Rubi [A] (verified)	279
Maple [B] (verified)	280
Fricas [B] (verification not implemented)	281
Sympy [B] (verification not implemented)	282
Maxima [B] (verification not implemented)	284
Giac [B] (verification not implemented)	285
Mupad [B] (verification not implemented)	288
Reduce [B] (verification not implemented)	289

Optimal result

Integrand size = 20, antiderivative size = 163

$$\int (a + bx)^3(A + Bx)(d + ex)^5 dx = \frac{(bd - ae)^3(Bd - Ae)(d + ex)^6}{6e^5} - \frac{(bd - ae)^2(4bBd - 3Abe - aBe)(d + ex)^7}{7e^5} + \frac{3b(bd - ae)(2bBd - Abe - aBe)(d + ex)^8}{8e^5} - \frac{b^2(4bBd - Abe - 3aBe)(d + ex)^9}{9e^5} + \frac{b^3B(d + ex)^{10}}{10e^5}$$

output

```
1/6*(-a*e+b*d)^3*(-A*e+B*d)*(e*x+d)^6/e^5-1/7*(-a*e+b*d)^2*(-3*A*b*e-B*a*e
+4*B*b*d)*(e*x+d)^7/e^5+3/8*b*(-a*e+b*d)*(-A*b*e-B*a*e+2*B*b*d)*(e*x+d)^8/
e^5-1/9*b^2*(-A*b*e-3*B*a*e+4*B*b*d)*(e*x+d)^9/e^5+1/10*b^3*B*(e*x+d)^10/e
^5
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 471 vs. $2(163) = 326$.

Time = 0.10 (sec) , antiderivative size = 471, normalized size of antiderivative = 2.89

$$\begin{aligned}
 \int (a + bx)^3(A + Bx)(d + ex)^5 dx = & a^3Ad^5x + \frac{1}{2}a^2d^4(3Abd + aBd + 5aAe)x^2 \\
 & + \frac{1}{3}ad^3(aBd(3bd + 5ae) \\
 & \qquad \qquad \qquad + A(3b^2d^2 + 15abde + 10a^2e^2)) x^3 \\
 & + \frac{1}{4}d^2(aBd(3b^2d^2 + 15abde + 10a^2e^2) \\
 & \qquad \qquad \qquad + A(b^3d^3 + 15ab^2d^2e + 30a^2bde^2 + 10a^3e^3)) x^4 \\
 & + \frac{1}{5}d(30a^2bde^2(Bd + Ae) + 5a^3e^3(2Bd + Ae) \\
 & \qquad \qquad \qquad + 15ab^2d^2e(Bd + 2Ae) + b^3d^3(Bd + 5Ae)) x^5 \\
 & + \frac{1}{6}e(30ab^2d^2e(Bd + Ae) + 15a^2bde^2(2Bd + Ae) \\
 & \qquad \qquad \qquad + a^3e^3(5Bd + Ae) + 5b^3d^3(Bd + 2Ae)) x^6 \\
 & + \frac{1}{7}e^2(a^3Be^3 + 10b^3d^2(Bd + Ae) \\
 & \qquad \qquad \qquad + 15ab^2de(2Bd + Ae) + 3a^2be^2(5Bd + Ae)) x^7 \\
 & + \frac{1}{8}be^3(3a^2Be^2 + 5b^2d(2Bd + Ae) \\
 & \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad + 3abe(5Bd + Ae)) x^8 \\
 & + \frac{1}{9}b^2e^4(5bBd + Abe + 3aBe)x^9 + \frac{1}{10}b^3Be^5x^{10}
 \end{aligned}$$

input

```
Integrate[(a + b*x)^3*(A + B*x)*(d + e*x)^5,x]
```

output

```

a^3*A*d^5*x + (a^2*d^4*(3*A*b*d + a*B*d + 5*a*A*e)*x^2)/2 + (a*d^3*(a*B*d*
(3*b*d + 5*a*e) + A*(3*b^2*d^2 + 15*a*b*d*e + 10*a^2*e^2))*x^3)/3 + (d^2*(
a*B*d*(3*b^2*d^2 + 15*a*b*d*e + 10*a^2*e^2) + A*(b^3*d^3 + 15*a*b^2*d^2*e
+ 30*a^2*b*d*e^2 + 10*a^3*e^3))*x^4)/4 + (d*(30*a^2*b*d*e^2*(B*d + A*e) +
5*a^3*e^3*(2*B*d + A*e) + 15*a*b^2*d^2*e*(B*d + 2*A*e) + b^3*d^3*(B*d + 5*
A*e))*x^5)/5 + (e*(30*a*b^2*d^2*e*(B*d + A*e) + 15*a^2*b*d*e^2*(2*B*d + A*
e) + a^3*e^3*(5*B*d + A*e) + 5*b^3*d^3*(B*d + 2*A*e))*x^6)/6 + (e^2*(a^3*B
*e^3 + 10*b^3*d^2*(B*d + A*e) + 15*a*b^2*d*e*(2*B*d + A*e) + 3*a^2*b*e^2*(
5*B*d + A*e))*x^7)/7 + (b*e^3*(3*a^2*B*e^2 + 5*b^2*d*(2*B*d + A*e) + 3*a*b
*e*(5*B*d + A*e))*x^8)/8 + (b^2*e^4*(5*b*B*d + A*b*e + 3*a*B*e)*x^9)/9 + (
b^3*B*e^5*x^10)/10

```

Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^3(A + Bx)(d + ex)^5 dx$$

$$\downarrow 86$$

$$\int \left(\frac{b^2(d + ex)^8(3aBe + Abe - 4bBd)}{e^4} - \frac{3b(d + ex)^7(bd - ae)(aBe + Abe - 2bBd)}{e^4} + \frac{(d + ex)^6(ae - bd)^2(aBe}{e^4} \right.$$

$$\left. - \frac{b^2(d + ex)^9(-3aBe - Abe + 4bBd)}{9e^5} + \frac{3b(d + ex)^8(bd - ae)(-aBe - Abe + 2bBd)}{8e^5} - \frac{(d + ex)^7(bd - ae)^2(-aBe - 3Abe + 4bBd)}{7e^5} + \frac{(d + ex)^6(bd - ae)^3(Bd - Ae)}{6e^5} + \frac{b^3B(d + ex)^{10}}{10e^5} \right)$$

input

```
Int[(a + b*x)^3*(A + B*x)*(d + e*x)^5,x]
```


output

```

1/10*b^3*B*e^5*x^10+1/9*((A*b^3+3*B*a*b^2)*e^5+5*b^3*B*d*e^4)*x^9+1/8*((3*
A*a*b^2+3*B*a^2*b)*e^5+5*(A*b^3+3*B*a*b^2)*d*e^4+10*b^3*B*d^2*e^3)*x^8+1/7
*((3*A*a^2*b+B*a^3)*e^5+5*(3*A*a*b^2+3*B*a^2*b)*d*e^4+10*(A*b^3+3*B*a*b^2)
*d^2*e^3+10*b^3*B*d^3*e^2)*x^7+1/6*(a^3*A*e^5+5*(3*A*a^2*b+B*a^3)*d*e^4+10
*(3*A*a*b^2+3*B*a^2*b)*d^2*e^3+10*(A*b^3+3*B*a*b^2)*d^3*e^2+5*b^3*B*d^4*e)
*x^6+1/5*(5*a^3*A*d*e^4+10*(3*A*a^2*b+B*a^3)*d^2*e^3+10*(3*A*a*b^2+3*B*a^2
*b)*d^3*e^2+5*(A*b^3+3*B*a*b^2)*d^4*e+b^3*B*d^5)*x^5+1/4*(10*a^3*A*d^2*e^3
+10*(3*A*a^2*b+B*a^3)*d^3*e^2+5*(3*A*a*b^2+3*B*a^2*b)*d^4*e+(A*b^3+3*B*a*b
^2)*d^5)*x^4+1/3*(10*a^3*A*d^3*e^2+5*(3*A*a^2*b+B*a^3)*d^4*e+(3*A*a*b^2+3*
B*a^2*b)*d^5)*x^3+1/2*(5*a^3*A*d^4*e+(3*A*a^2*b+B*a^3)*d^5)*x^2+a^3*A*d^5*
x

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 518 vs. $2(153) = 306$.

Time = 0.06 (sec) , antiderivative size = 518, normalized size of antiderivative = 3.18

$$\begin{aligned}
& \int (a + bx)^3 (A + Bx)(d + ex)^5 dx \\
&= \frac{1}{10} Bb^3 e^5 x^{10} + Aa^3 d^5 x + \frac{1}{9} (5 Bb^3 de^4 + (3 Bab^2 + Ab^3) e^5) x^9 \\
&+ \frac{1}{8} (10 Bb^3 d^2 e^3 + 5 (3 Bab^2 + Ab^3) de^4 + 3 (Ba^2 b + Aab^2) e^5) x^8 \\
&+ \frac{1}{7} (10 Bb^3 d^3 e^2 + 10 (3 Bab^2 + Ab^3) d^2 e^3 + 15 (Ba^2 b + Aab^2) de^4 + (Ba^3 + 3 Aa^2 b) e^5) x^7 \\
&+ \frac{1}{6} (5 Bb^3 d^4 e + Aa^3 e^5 + 10 (3 Bab^2 + Ab^3) d^3 e^2 + 30 (Ba^2 b + Aab^2) d^2 e^3 + 5 (Ba^3 + 3 Aa^2 b) de^4) x^6 \\
&+ \frac{1}{5} (Bb^3 d^5 + 5 Aa^3 de^4 + 5 (3 Bab^2 + Ab^3) d^4 e + 30 (Ba^2 b + Aab^2) d^3 e^2 + 10 (Ba^3 + 3 Aa^2 b) d^2 e^3) x^5 \\
&+ \frac{1}{4} (10 Aa^3 d^2 e^3 + (3 Bab^2 + Ab^3) d^5 + 15 (Ba^2 b + Aab^2) d^4 e + 10 (Ba^3 + 3 Aa^2 b) d^3 e^2) x^4 \\
&+ \frac{1}{3} (10 Aa^3 d^3 e^2 + 3 (Ba^2 b + Aab^2) d^5 + 5 (Ba^3 + 3 Aa^2 b) d^4 e) x^3 \\
&+ \frac{1}{2} (5 Aa^3 d^4 e + (Ba^3 + 3 Aa^2 b) d^5) x^2
\end{aligned}$$

input

```
integrate((b*x+a)^3*(B*x+A)*(e*x+d)^5,x, algorithm="fricas")
```

output

```

1/10*B*b^3*e^5*x^10 + A*a^3*d^5*x + 1/9*(5*B*b^3*d*e^4 + (3*B*a*b^2 + A*b^
3)*e^5)*x^9 + 1/8*(10*B*b^3*d^2*e^3 + 5*(3*B*a*b^2 + A*b^3)*d*e^4 + 3*(B*a
^2*b + A*a*b^2)*e^5)*x^8 + 1/7*(10*B*b^3*d^3*e^2 + 10*(3*B*a*b^2 + A*b^3)*
d^2*e^3 + 15*(B*a^2*b + A*a*b^2)*d*e^4 + (B*a^3 + 3*A*a^2*b)*e^5)*x^7 + 1/
6*(5*B*b^3*d^4*e + A*a^3*e^5 + 10*(3*B*a*b^2 + A*b^3)*d^3*e^2 + 30*(B*a^2*
b + A*a*b^2)*d^2*e^3 + 5*(B*a^3 + 3*A*a^2*b)*d*e^4)*x^6 + 1/5*(B*b^3*d^5 +
5*A*a^3*d*e^4 + 5*(3*B*a*b^2 + A*b^3)*d^4*e + 30*(B*a^2*b + A*a*b^2)*d^3*
e^2 + 10*(B*a^3 + 3*A*a^2*b)*d^2*e^3)*x^5 + 1/4*(10*A*a^3*d^2*e^3 + (3*B*a
*b^2 + A*b^3)*d^5 + 15*(B*a^2*b + A*a*b^2)*d^4*e + 10*(B*a^3 + 3*A*a^2*b)*
d^3*e^2)*x^4 + 1/3*(10*A*a^3*d^3*e^2 + 3*(B*a^2*b + A*a*b^2)*d^5 + 5*(B*a^
3 + 3*A*a^2*b)*d^4*e)*x^3 + 1/2*(5*A*a^3*d^4*e + (B*a^3 + 3*A*a^2*b)*d^5)*
x^2

```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 678 vs. $2(155) = 310$.

Time = 0.07 (sec) , antiderivative size = 678, normalized size of antiderivative = 4.16

$$\begin{aligned}
 \int (a + bx)^3 (A + Bx)(d + ex)^5 dx = & Aa^3 d^5 x + \frac{Bb^3 e^5 x^{10}}{10} \\
 & + x^9 \left(\frac{Ab^3 e^5}{9} + \frac{Bab^2 e^5}{3} + \frac{5Bb^3 d e^4}{9} \right) + x^8 \cdot \left(\frac{3Aab^2 e^5}{8} \right. \\
 & \left. + \frac{5Ab^3 d e^4}{8} + \frac{3Ba^2 b e^5}{8} + \frac{15Bab^2 d e^4}{8} + \frac{5Bb^3 d^2 e^3}{4} \right) \\
 & + x^7 \cdot \left(\frac{3Aa^2 b e^5}{7} + \frac{15Aab^2 d e^4}{7} + \frac{10Ab^3 d^2 e^3}{7} + \frac{Ba^3 e^5}{7} \right. \\
 & \left. + \frac{15Ba^2 b d e^4}{7} + \frac{30Bab^2 d^2 e^3}{7} + \frac{10Bb^3 d^3 e^2}{7} \right) \\
 & + x^6 \left(\frac{Aa^3 e^5}{6} + \frac{5Aa^2 b d e^4}{2} + 5Aab^2 d^2 e^3 + \frac{5Ab^3 d^3 e^2}{3} \right. \\
 & \left. + \frac{5Ba^3 d e^4}{6} + 5Ba^2 b d^2 e^3 + 5Bab^2 d^3 e^2 + \frac{5Bb^3 d^4 e}{6} \right) \\
 & + x^5 \left(Aa^3 d e^4 + 6Aa^2 b d^2 e^3 + 6Aab^2 d^3 e^2 + Ab^3 d^4 e \right. \\
 & \left. + 2Ba^3 d^2 e^3 + 6Ba^2 b d^3 e^2 + 3Bab^2 d^4 e + \frac{Bb^3 d^5}{5} \right) \\
 & + x^4 \cdot \left(\frac{5Aa^3 d^2 e^3}{2} + \frac{15Aa^2 b d^3 e^2}{2} + \frac{15Aab^2 d^4 e}{4} \right. \\
 & \left. + \frac{Ab^3 d^5}{4} + \frac{5Ba^3 d^3 e^2}{2} + \frac{15Ba^2 b d^4 e}{4} + \frac{3Bab^2 d^5}{4} \right) \\
 & + x^3 \cdot \left(\frac{10Aa^3 d^3 e^2}{3} + 5Aa^2 b d^4 e + Aab^2 d^5 + \frac{5Ba^3 d^4 e}{3} \right. \\
 & \left. + Ba^2 b d^5 \right) + x^2 \cdot \left(\frac{5Aa^3 d^4 e}{2} + \frac{3Aa^2 b d^5}{2} + \frac{Ba^3 d^5}{2} \right)
 \end{aligned}$$

input `integrate((b*x+a)**3*(B*x+A)*(e*x+d)**5,x)`

output

```

A*a**3*d**5*x + B*b**3*e**5*x**10/10 + x**9*(A*b**3*e**5/9 + B*a*b**2*e**5
/3 + 5*B*b**3*d*e**4/9) + x**8*(3*A*a*b**2*e**5/8 + 5*A*b**3*d*e**4/8 + 3*
B*a**2*b*e**5/8 + 15*B*a*b**2*d*e**4/8 + 5*B*b**3*d**2*e**3/4) + x**7*(3*A
*a**2*b*e**5/7 + 15*A*a*b**2*d*e**4/7 + 10*A*b**3*d**2*e**3/7 + B*a**3*e**
5/7 + 15*B*a**2*b*d*e**4/7 + 30*B*a*b**2*d**2*e**3/7 + 10*B*b**3*d**3*e**2
/7) + x**6*(A*a**3*e**5/6 + 5*A*a**2*b*d*e**4/2 + 5*A*a*b**2*d**2*e**3 + 5
*A*b**3*d**3*e**2/3 + 5*B*a**3*d*e**4/6 + 5*B*a**2*b*d**2*e**3 + 5*B*a*b**
2*d**3*e**2 + 5*B*b**3*d**4*e/6) + x**5*(A*a**3*d*e**4 + 6*A*a**2*b*d**2*e
**3 + 6*A*a*b**2*d**3*e**2 + A*b**3*d**4*e + 2*B*a**3*d**2*e**3 + 6*B*a**2
*b*d**3*e**2 + 3*B*a*b**2*d**4*e + B*b**3*d**5/5) + x**4*(5*A*a**3*d**2*e*
**3/2 + 15*A*a**2*b*d**3*e**2/2 + 15*A*a*b**2*d**4*e/4 + A*b**3*d**5/4 + 5*
B*a**3*d**3*e**2/2 + 15*B*a**2*b*d**4*e/4 + 3*B*a*b**2*d**5/4) + x**3*(10*
A*a**3*d**3*e**2/3 + 5*A*a**2*b*d**4*e + A*a*b**2*d**5 + 5*B*a**3*d**4*e/3
+ B*a**2*b*d**5) + x**2*(5*A*a**3*d**4*e/2 + 3*A*a**2*b*d**5/2 + B*a**3*d
**5/2)

```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 518 vs. $2(153) = 306$.

Time = 0.04 (sec) , antiderivative size = 518, normalized size of antiderivative = 3.18

$$\begin{aligned}
& \int (a + bx)^3 (A + Bx)(d + ex)^5 dx \\
&= \frac{1}{10} Bb^3 e^5 x^{10} + Aa^3 d^5 x + \frac{1}{9} (5 Bb^3 de^4 + (3 Bab^2 + Ab^3) e^5) x^9 \\
&\quad + \frac{1}{8} (10 Bb^3 d^2 e^3 + 5 (3 Bab^2 + Ab^3) de^4 + 3 (Ba^2 b + Aab^2) e^5) x^8 \\
&\quad + \frac{1}{7} (10 Bb^3 d^3 e^2 + 10 (3 Bab^2 + Ab^3) d^2 e^3 + 15 (Ba^2 b + Aab^2) de^4 + (Ba^3 + 3 Aa^2 b) e^5) x^7 \\
&\quad + \frac{1}{6} (5 Bb^3 d^4 e + Aa^3 e^5 + 10 (3 Bab^2 + Ab^3) d^3 e^2 + 30 (Ba^2 b + Aab^2) d^2 e^3 + 5 (Ba^3 + 3 Aa^2 b) de^4) x^6 \\
&\quad + \frac{1}{5} (Bb^3 d^5 + 5 Aa^3 de^4 + 5 (3 Bab^2 + Ab^3) d^4 e + 30 (Ba^2 b + Aab^2) d^3 e^2 + 10 (Ba^3 + 3 Aa^2 b) d^2 e^3) x^5 \\
&\quad + \frac{1}{4} (10 Aa^3 d^2 e^3 + (3 Bab^2 + Ab^3) d^5 + 15 (Ba^2 b + Aab^2) d^4 e + 10 (Ba^3 + 3 Aa^2 b) d^3 e^2) x^4 \\
&\quad + \frac{1}{3} (10 Aa^3 d^3 e^2 + 3 (Ba^2 b + Aab^2) d^5 + 5 (Ba^3 + 3 Aa^2 b) d^4 e) x^3 \\
&\quad + \frac{1}{2} (5 Aa^3 d^4 e + (Ba^3 + 3 Aa^2 b) d^5) x^2
\end{aligned}$$

input `integrate((b*x+a)^3*(B*x+A)*(e*x+d)^5,x, algorithm="maxima")`

output

$$\begin{aligned} & 1/10*B*b^3*e^5*x^{10} + A*a^3*d^5*x + 1/9*(5*B*b^3*d*e^4 + (3*B*a*b^2 + A*b^3)*e^5)*x^9 + 1/8*(10*B*b^3*d^2*e^3 + 5*(3*B*a*b^2 + A*b^3)*d*e^4 + 3*(B*a^2*b + A*a*b^2)*e^5)*x^8 + 1/7*(10*B*b^3*d^3*e^2 + 10*(3*B*a*b^2 + A*b^3)*d^2*e^3 + 15*(B*a^2*b + A*a*b^2)*d*e^4 + (B*a^3 + 3*A*a^2*b)*e^5)*x^7 + 1/6*(5*B*b^3*d^4*e + A*a^3*e^5 + 10*(3*B*a*b^2 + A*b^3)*d^3*e^2 + 30*(B*a^2*b + A*a*b^2)*d^2*e^3 + 5*(B*a^3 + 3*A*a^2*b)*d*e^4)*x^6 + 1/5*(B*b^3*d^5 + 5*A*a^3*d*e^4 + 5*(3*B*a*b^2 + A*b^3)*d^4*e + 30*(B*a^2*b + A*a*b^2)*d^3*e^2 + 10*(B*a^3 + 3*A*a^2*b)*d^2*e^3)*x^5 + 1/4*(10*A*a^3*d^2*e^3 + (3*B*a*b^2 + A*b^3)*d^5 + 15*(B*a^2*b + A*a*b^2)*d^4*e + 10*(B*a^3 + 3*A*a^2*b)*d^3*e^2)*x^4 + 1/3*(10*A*a^3*d^3*e^2 + 3*(B*a^2*b + A*a*b^2)*d^5 + 5*(B*a^3 + 3*A*a^2*b)*d^4*e)*x^3 + 1/2*(5*A*a^3*d^4*e + (B*a^3 + 3*A*a^2*b)*d^5)*x^2 \end{aligned}$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 658 vs. $2(153) = 306$.

Time = 0.12 (sec) , antiderivative size = 658, normalized size of antiderivative = 4.04

$$\begin{aligned}
\int (a + bx)^3(A + Bx)(d + ex)^5 dx = & \frac{1}{10} Bb^3e^5x^{10} + \frac{5}{9} Bb^3de^4x^9 + \frac{1}{3} Bab^2e^5x^9 \\
& + \frac{1}{9} Ab^3e^5x^9 + \frac{5}{4} Bb^3d^2e^3x^8 + \frac{15}{8} Bab^2de^4x^8 \\
& + \frac{5}{8} Ab^3de^4x^8 + \frac{3}{8} Ba^2be^5x^8 + \frac{3}{8} Aab^2e^5x^8 \\
& + \frac{10}{7} Bb^3d^3e^2x^7 + \frac{30}{7} Bab^2d^2e^3x^7 + \frac{10}{7} Ab^3d^2e^3x^7 \\
& + \frac{15}{7} Ba^2bde^4x^7 + \frac{15}{7} Aab^2de^4x^7 + \frac{1}{7} Ba^3e^5x^7 \\
& + \frac{3}{7} Aa^2be^5x^7 + \frac{5}{6} Bb^3d^4ex^6 + 5 Bab^2d^3e^2x^6 \\
& + \frac{5}{3} Ab^3d^3e^2x^6 + 5 Ba^2bd^2e^3x^6 + 5 Aab^2d^2e^3x^6 \\
& + \frac{5}{6} Ba^3de^4x^6 + \frac{5}{2} Aa^2bde^4x^6 + \frac{1}{6} Aa^3e^5x^6 \\
& + \frac{1}{5} Bb^3d^5x^5 + 3 Bab^2d^4ex^5 + Ab^3d^4ex^5 \\
& + 6 Ba^2bd^3e^2x^5 + 6 Aab^2d^3e^2x^5 + 2 Ba^3d^2e^3x^5 \\
& + 6 Aa^2bd^2e^3x^5 + Aa^3de^4x^5 + \frac{3}{4} Bab^2d^5x^4 \\
& + \frac{1}{4} Ab^3d^5x^4 + \frac{15}{4} Ba^2bd^4ex^4 + \frac{15}{4} Aab^2d^4ex^4 \\
& + \frac{5}{2} Ba^3d^3e^2x^4 + \frac{15}{2} Aa^2bd^3e^2x^4 + \frac{5}{2} Aa^3d^2e^3x^4 \\
& + Ba^2bd^5x^3 + Aab^2d^5x^3 + \frac{5}{3} Ba^3d^4ex^3 \\
& + 5 Aa^2bd^4ex^3 + \frac{10}{3} Aa^3d^3e^2x^3 + \frac{1}{2} Ba^3d^5x^2 \\
& + \frac{3}{2} Aa^2bd^5x^2 + \frac{5}{2} Aa^3d^4ex^2 + Aa^3d^5x
\end{aligned}$$

input `integrate((b*x+a)^3*(B*x+A)*(e*x+d)^5,x, algorithm="giac")`

output

$$\begin{aligned}
& 1/10*B*b^3*e^5*x^10 + 5/9*B*b^3*d*e^4*x^9 + 1/3*B*a*b^2*e^5*x^9 + 1/9*A*b^3*e^5*x^9 + 5/4*B*b^3*d^2*e^3*x^8 + 15/8*B*a*b^2*d*e^4*x^8 + 5/8*A*b^3*d*e^4*x^8 + 3/8*B*a^2*b*e^5*x^8 + 3/8*A*a*b^2*e^5*x^8 + 10/7*B*b^3*d^3*e^2*x^7 + 30/7*B*a*b^2*d^2*e^3*x^7 + 10/7*A*b^3*d^2*e^3*x^7 + 15/7*B*a^2*b*d*e^4*x^7 + 15/7*A*a*b^2*d*e^4*x^7 + 1/7*B*a^3*e^5*x^7 + 3/7*A*a^2*b*e^5*x^7 + 5/6*B*b^3*d^4*e*x^6 + 5*B*a*b^2*d^3*e^2*x^6 + 5/3*A*b^3*d^3*e^2*x^6 + 5*B*a^2*b*d^2*e^3*x^6 + 5*A*a*b^2*d^2*e^3*x^6 + 5/6*B*a^3*d*e^4*x^6 + 5/2*A*a^2*b*d*e^4*x^6 + 1/6*A*a^3*e^5*x^6 + 1/5*B*b^3*d^5*x^5 + 3*B*a*b^2*d^4*e*x^5 + A*b^3*d^4*e*x^5 + 6*B*a^2*b*d^3*e^2*x^5 + 6*A*a*b^2*d^3*e^2*x^5 + 2*B*a^3*d^2*e^3*x^5 + 6*A*a^2*b*d^2*e^3*x^5 + A*a^3*d*e^4*x^5 + 3/4*B*a*b^2*d^5*x^4 + 1/4*A*b^3*d^5*x^4 + 15/4*B*a^2*b*d^4*e*x^4 + 15/4*A*a*b^2*d^4*e*x^4 + 5/2*B*a^3*d^3*e^2*x^4 + 15/2*A*a^2*b*d^3*e^2*x^4 + 5/2*A*a^3*d^2*e^3*x^4 + B*a^2*b*d^5*x^3 + A*a*b^2*d^5*x^3 + 5/3*B*a^3*d^4*e*x^3 + 5*A*a^2*b*d^4*e*x^3 + 10/3*A*a^3*d^3*e^2*x^3 + 1/2*B*a^3*d^5*x^2 + 3/2*A*a^2*b*d^5*x^2 + 5/2*A*a^3*d^4*e*x^2 + A*a^3*d^5*x
\end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 544, normalized size of antiderivative = 3.34

$$\begin{aligned}
\int (a + bx)^3 (A + Bx)(d + ex)^5 dx = & x^3 \left(\frac{5Ba^3d^4e}{3} + \frac{10Aa^3d^3e^2}{3} + Ba^2bd^5 \right. \\
& \left. + 5Aa^2bd^4e + Aab^2d^5 \right) \\
& + x^8 \left(\frac{3Ba^2be^5}{8} + \frac{15Bab^2de^4}{8} + \frac{3Aab^2e^5}{8} \right. \\
& \left. + \frac{5Bb^3d^2e^3}{4} + \frac{5Ab^3de^4}{8} \right) + x^4 \left(\frac{5Ba^3d^3e^2}{2} \right. \\
& \left. + \frac{5Aa^3d^2e^3}{2} + \frac{15Ba^2bd^4e}{4} + \frac{15Aa^2bd^3e^2}{2} \right. \\
& \left. + \frac{3Bab^2d^5}{4} + \frac{15Aab^2d^4e}{4} + \frac{Ab^3d^5}{4} \right) \\
& + x^7 \left(\frac{Ba^3e^5}{7} + \frac{15Ba^2bde^4}{7} + \frac{3Aa^2be^5}{7} \right. \\
& \left. + \frac{30Bab^2d^2e^3}{7} + \frac{15Aab^2de^4}{7} + \frac{10Bb^3d^3e^2}{7} \right. \\
& \left. + \frac{10Ab^3d^2e^3}{7} \right) + x^5 \left(2Ba^3d^2e^3 + Aa^3de^4 \right. \\
& \left. + 6Ba^2bd^3e^2 + 6Aa^2bd^2e^3 + 3Bab^2d^4e \right. \\
& \left. + 6Aab^2d^3e^2 + \frac{Bb^3d^5}{5} + Ab^3d^4e \right) \\
& + x^6 \left(\frac{5Ba^3de^4}{6} + \frac{Aa^3e^5}{6} + 5Ba^2bd^2e^3 \right. \\
& \left. + \frac{5Aa^2bde^4}{2} + 5Bab^2d^3e^2 + 5Aab^2d^2e^3 \right. \\
& \left. + \frac{5Bb^3d^4e}{6} + \frac{5Ab^3d^3e^2}{3} \right) \\
& + \frac{a^2d^4x^2(5Aae + 3Abd + Bad)}{2} \\
& + \frac{b^2e^4x^9(Abe + 3Bae + 5Bbd)}{9} \\
& + Aa^3d^5x + \frac{Bb^3e^5x^{10}}{10}
\end{aligned}$$

input `int((A + B*x)*(a + b*x)^3*(d + e*x)^5,x)`

output

```
x^3*(A*a*b^2*d^5 + B*a^2*b*d^5 + (5*B*a^3*d^4*e)/3 + (10*A*a^3*d^3*e^2)/3
+ 5*A*a^2*b*d^4*e) + x^8*((3*A*a*b^2*e^5)/8 + (3*B*a^2*b*e^5)/8 + (5*A*b^3
*d*e^4)/8 + (5*B*b^3*d^2*e^3)/4 + (15*B*a*b^2*d*e^4)/8) + x^4*((A*b^3*d^5)
/4 + (3*B*a*b^2*d^5)/4 + (5*A*a^3*d^2*e^3)/2 + (5*B*a^3*d^3*e^2)/2 + (15*A
*a^2*b*d^3*e^2)/2 + (15*A*a*b^2*d^4*e)/4 + (15*B*a^2*b*d^4*e)/4) + x^7*((B
*a^3*e^5)/7 + (3*A*a^2*b*e^5)/7 + (10*A*b^3*d^2*e^3)/7 + (10*B*b^3*d^3*e^2
)/7 + (30*B*a*b^2*d^2*e^3)/7 + (15*A*a*b^2*d*e^4)/7 + (15*B*a^2*b*d*e^4)/7
) + x^5*((B*b^3*d^5)/5 + A*a^3*d*e^4 + A*b^3*d^4*e + 2*B*a^3*d^2*e^3 + 6*A
*a*b^2*d^3*e^2 + 6*A*a^2*b*d^2*e^3 + 6*B*a^2*b*d^3*e^2 + 3*B*a*b^2*d^4*e)
+ x^6*((A*a^3*e^5)/6 + (5*B*a^3*d*e^4)/6 + (5*B*b^3*d^4*e)/6 + (5*A*b^3*d^
3*e^2)/3 + 5*A*a*b^2*d^2*e^3 + 5*B*a*b^2*d^3*e^2 + 5*B*a^2*b*d^2*e^3 + (5*
A*a^2*b*d*e^4)/2) + (a^2*d^4*x^2*(5*A*a*e + 3*A*b*d + B*a*d))/2 + (b^2*e^4
*x^9*(A*b*e + 3*B*a*e + 5*B*b*d))/9 + A*a^3*d^5*x + (B*b^3*e^5*x^10)/10
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 397, normalized size of antiderivative = 2.44

$$\int (a + bx)^3 (A + Bx)(d + ex)^5 dx$$

$$= \frac{x(126b^4e^5x^9 + 560ab^3e^5x^8 + 700b^4de^4x^8 + 945a^2b^2e^5x^7 + 3150ab^3de^4x^7 + 1575b^4d^2e^3x^7 + 720a^3be^5x^6 + \dots)}{1260}$$

input

```
int((b*x+a)^3*(B*x+A)*(e*x+d)^5,x)
```

output

```
(x*(1260*a**4*d**5 + 3150*a**4*d**4*e*x + 4200*a**4*d**3*e**2*x**2 + 3150*
a**4*d**2*e**3*x**3 + 1260*a**4*d*e**4*x**4 + 210*a**4*e**5*x**5 + 2520*a*
**3*b*d**5*x + 8400*a**3*b*d**4*e*x**2 + 12600*a**3*b*d**3*e**2*x**3 + 1008
0*a**3*b*d**2*e**3*x**4 + 4200*a**3*b*d*e**4*x**5 + 720*a**3*b*e**5*x**6 +
2520*a**2*b**2*d**5*x**2 + 9450*a**2*b**2*d**4*e*x**3 + 15120*a**2*b**2*d
**3*e**2*x**4 + 12600*a**2*b**2*d**2*e**3*x**5 + 5400*a**2*b**2*d*e**4*x**
6 + 945*a**2*b**2*e**5*x**7 + 1260*a*b**3*d**5*x**3 + 5040*a*b**3*d**4*e*x
**4 + 8400*a*b**3*d**3*e**2*x**5 + 7200*a*b**3*d**2*e**3*x**6 + 3150*a*b**
3*d*e**4*x**7 + 560*a*b**3*e**5*x**8 + 252*b**4*d**5*x**4 + 1050*b**4*d**4
*e*x**5 + 1800*b**4*d**3*e**2*x**6 + 1575*b**4*d**2*e**3*x**7 + 700*b**4*d
**4*x**8 + 126*b**4*e**5*x**9))/1260
```

3.26 $\int (a + bx)^3 (A + Bx)(d + ex)^4 dx$

Optimal result	290
Mathematica [B] (verified)	291
Rubi [A] (verified)	292
Maple [B] (verified)	293
Fricas [B] (verification not implemented)	294
Sympy [B] (verification not implemented)	295
Maxima [B] (verification not implemented)	296
Giac [B] (verification not implemented)	297
Mupad [B] (verification not implemented)	299
Reduce [B] (verification not implemented)	300

Optimal result

Integrand size = 20, antiderivative size = 163

$$\int (a + bx)^3 (A + Bx)(d + ex)^4 dx = \frac{(bd - ae)^3 (Bd - Ae)(d + ex)^5}{5e^5} - \frac{(bd - ae)^2 (4bBd - 3Abe - aBe)(d + ex)^6}{6e^5} + \frac{3b(bd - ae)(2bBd - Abe - aBe)(d + ex)^7}{7e^5} - \frac{b^2(4bBd - Abe - 3aBe)(d + ex)^8}{8e^5} + \frac{b^3B(d + ex)^9}{9e^5}$$

output

```
1/5*(-a*e+b*d)^3*(-A*e+B*d)*(e*x+d)^5/e^5-1/6*(-a*e+b*d)^2*(-3*A*b*e-B*a*e+4*B*b*d)*(e*x+d)^6/e^5+3/7*b*(-a*e+b*d)*(-A*b*e-B*a*e+2*B*b*d)*(e*x+d)^7/e^5-1/8*b^2*(-A*b*e-3*B*a*e+4*B*b*d)*(e*x+d)^8/e^5+1/9*b^3*B*(e*x+d)^9/e^5
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 397 vs. $2(163) = 326$.

Time = 0.09 (sec) , antiderivative size = 397, normalized size of antiderivative = 2.44

$$\begin{aligned}
 & \int (a + bx)^3 (A + Bx)(d + ex)^4 dx \\
 &= a^3 A d^4 x + \frac{1}{2} a^2 d^3 (3A b d + a B d + 4a A e) x^2 \\
 &+ \frac{1}{3} a d^2 (a B d (3b d + 4a e) + 3A (b^2 d^2 + 4a b d e + 2a^2 e^2)) x^3 \\
 &+ \frac{1}{4} d (3a B d (b^2 d^2 + 4a b d e + 2a^2 e^2) + A (b^3 d^3 + 12a b^2 d^2 e + 18a^2 b d e^2 + 4a^3 e^3)) x^4 \\
 &+ \frac{1}{5} (a^3 e^3 (4B d + A e) + 6a^2 b d e^2 (3B d + 2A e) + 6a b^2 d^2 e (2B d + 3A e) \\
 &+ b^3 d^3 (B d + 4A e)) x^5 \\
 &+ \frac{1}{6} e (a^3 B e^3 + 3a^2 b e^2 (4B d + A e) + 6a b^2 d e (3B d + 2A e) + 2b^3 d^2 (2B d + 3A e)) x^6 \\
 &+ \frac{1}{7} b e^2 (3a^2 B e^2 + 3a b e (4B d + A e) + 2b^2 d (3B d + 2A e)) x^7 \\
 &+ \frac{1}{8} b^2 e^3 (4b B d + A b e + 3a B e) x^8 + \frac{1}{9} b^3 B e^4 x^9
 \end{aligned}$$

input `Integrate[(a + b*x)^3*(A + B*x)*(d + e*x)^4,x]`

output `a^3*A*d^4*x + (a^2*d^3*(3*A*b*d + a*B*d + 4*a*A*e))*x^2)/2 + (a*d^2*(a*B*d*(3*b*d + 4*a*e) + 3*A*(b^2*d^2 + 4*a*b*d*e + 2*a^2*e^2))*x^3)/3 + (d*(3*a*B*d*(b^2*d^2 + 4*a*b*d*e + 2*a^2*e^2) + A*(b^3*d^3 + 12*a*b^2*d^2*e + 18*a^2*b*d*e^2 + 4*a^3*e^3))*x^4)/4 + ((a^3*e^3*(4*B*d + A*e) + 6*a^2*b*d*e^2*(3*B*d + 2*A*e) + 6*a*b^2*d^2*e*(2*B*d + 3*A*e) + b^3*d^3*(B*d + 4*A*e))*x^5)/5 + (e*(a^3*B*e^3 + 3*a^2*b*e^2*(4*B*d + A*e) + 6*a*b^2*d*e*(3*B*d + 2*A*e) + 2*b^3*d^2*(2*B*d + 3*A*e))*x^6)/6 + (b*e^2*(3*a^2*B*e^2 + 3*a*b*e*(4*B*d + A*e) + 2*b^2*d*(3*B*d + 2*A*e))*x^7)/7 + (b^2*e^3*(4*b*B*d + A*b*e + 3*a*B*e))*x^8)/8 + (b^3*B*e^4*x^9)/9`

Rubi [A] (verified)

Time = 0.55 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^3(A + Bx)(d + ex)^4 dx$$

↓ 86

$$\int \left(\frac{b^2(d + ex)^7(3aBe + Abe - 4bBd)}{e^4} - \frac{3b(d + ex)^6(bd - ae)(aBe + Abe - 2bBd)}{e^4} + \frac{(d + ex)^5(ae - bd)^2(aBe - b^2d - a^2e)}{e^4} \right) dx$$

↓ 2009

$$\frac{-\frac{b^2(d + ex)^8(-3aBe - Abe + 4bBd)}{8e^5} + \frac{3b(d + ex)^7(bd - ae)(-aBe - Abe + 2bBd)}{7e^5}}{\frac{(d + ex)^6(bd - ae)^2(-aBe - 3Abe + 4bBd)}{6e^5} + \frac{(d + ex)^5(bd - ae)^3(Bd - Ae)}{5e^5} + \frac{b^3B(d + ex)^9}{9e^5}}$$

input `Int[(a + b*x)^3*(A + B*x)*(d + e*x)^4,x]`

output `((b*d - a*e)^3*(B*d - A*e)*(d + e*x)^5)/(5*e^5) - ((b*d - a*e)^2*(4*b*B*d - 3*A*b*e - a*B*e)*(d + e*x)^6)/(6*e^5) + (3*b*(b*d - a*e)*(2*b*B*d - A*b*e - a*B*e)*(d + e*x)^7)/(7*e^5) - (b^2*(4*b*B*d - A*b*e - 3*a*B*e)*(d + e*x)^8)/(8*e^5) + (b^3*B*(d + e*x)^9)/(9*e^5)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 425 vs. $2(153) = 306$.

Time = 0.08 (sec) , antiderivative size = 425, normalized size of antiderivative = 2.61

$$\begin{aligned} & \int (a + bx)^3 (A + Bx)(d + ex)^4 dx \\ &= \frac{1}{9} Bb^3 e^4 x^9 + Aa^3 d^4 x + \frac{1}{8} (4 Bb^3 d e^3 + (3 Bab^2 + Ab^3) e^4) x^8 \\ & \quad + \frac{1}{7} (6 Bb^3 d^2 e^2 + 4 (3 Bab^2 + Ab^3) d e^3 + 3 (Ba^2 b + Aab^2) e^4) x^7 \\ & \quad + \frac{1}{6} (4 Bb^3 d^3 e + 6 (3 Bab^2 + Ab^3) d^2 e^2 + 12 (Ba^2 b + Aab^2) d e^3 + (Ba^3 + 3 Aa^2 b) e^4) x^6 \\ & \quad + \frac{1}{5} (Bb^3 d^4 + Aa^3 e^4 + 4 (3 Bab^2 + Ab^3) d^3 e + 18 (Ba^2 b + Aab^2) d^2 e^2 + 4 (Ba^3 + 3 Aa^2 b) d e^3) x^5 \\ & \quad + \frac{1}{4} (4 Aa^3 d e^3 + (3 Bab^2 + Ab^3) d^4 + 12 (Ba^2 b + Aab^2) d^3 e + 6 (Ba^3 + 3 Aa^2 b) d^2 e^2) x^4 \\ & \quad + \frac{1}{3} (6 Aa^3 d^2 e^2 + 3 (Ba^2 b + Aab^2) d^4 + 4 (Ba^3 + 3 Aa^2 b) d^3 e) x^3 \\ & \quad + \frac{1}{2} (4 Aa^3 d^3 e + (Ba^3 + 3 Aa^2 b) d^4) x^2 \end{aligned}$$

input `integrate((b*x+a)^3*(B*x+A)*(e*x+d)^4,x, algorithm="fricas")`

output `1/9*B*b^3*e^4*x^9 + A*a^3*d^4*x + 1/8*(4*B*b^3*d*e^3 + (3*B*a*b^2 + A*b^3)*e^4)*x^8 + 1/7*(6*B*b^3*d^2*e^2 + 4*(3*B*a*b^2 + A*b^3)*d*e^3 + 3*(B*a^2*b + A*a*b^2)*e^4)*x^7 + 1/6*(4*B*b^3*d^3*e + 6*(3*B*a*b^2 + A*b^3)*d^2*e^2 + 12*(B*a^2*b + A*a*b^2)*d*e^3 + (B*a^3 + 3*A*a^2*b)*e^4)*x^6 + 1/5*(B*b^3*d^4 + A*a^3*e^4 + 4*(3*B*a*b^2 + A*b^3)*d^3*e + 18*(B*a^2*b + A*a*b^2)*d^2*e^2 + 4*(B*a^3 + 3*A*a^2*b)*d*e^3)*x^5 + 1/4*(4*A*a^3*d*e^3 + (3*B*a*b^2 + A*b^3)*d^4 + 12*(B*a^2*b + A*a*b^2)*d^3*e + 6*(B*a^3 + 3*A*a^2*b)*d^2*e^2)*x^4 + 1/3*(6*A*a^3*d^2*e^2 + 3*(B*a^2*b + A*a*b^2)*d^4 + 4*(B*a^3 + 3*A*a^2*b)*d^3*e)*x^3 + 1/2*(4*A*a^3*d^3*e + (B*a^3 + 3*A*a^2*b)*d^4)*x^2`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 546 vs. $2(155) = 310$.

Time = 0.05 (sec) , antiderivative size = 546, normalized size of antiderivative = 3.35

$$\begin{aligned}
 \int (a + bx)^3(A + Bx)(d + ex)^4 dx = & Aa^3d^4x + \frac{Bb^3e^4x^9}{9} \\
 & + x^8 \left(\frac{Ab^3e^4}{8} + \frac{3Bab^2e^4}{8} + \frac{Bb^3de^3}{2} \right) + x^7 \\
 & \cdot \left(\frac{3Aab^2e^4}{7} + \frac{4Ab^3de^3}{7} + \frac{3Ba^2be^4}{7} + \frac{12Bab^2de^3}{7} \right. \\
 & \left. + \frac{6Bb^3d^2e^2}{7} \right) + x^6 \left(\frac{Aa^2be^4}{2} + 2Aab^2de^3 + Ab^3d^2e^2 \right. \\
 & \left. + \frac{Ba^3e^4}{6} + 2Ba^2bde^3 + 3Bab^2d^2e^2 + \frac{2Bb^3d^3e}{3} \right) \\
 & + x^5 \left(\frac{Aa^3e^4}{5} + \frac{12Aa^2bde^3}{5} + \frac{18Aab^2d^2e^2}{5} + \frac{4Ab^3d^3e}{5} \right. \\
 & \left. + \frac{4Ba^3de^3}{5} + \frac{18Ba^2bd^2e^2}{5} + \frac{12Bab^2d^3e}{5} + \frac{Bb^3d^4}{5} \right) \\
 & + x^4 \left(Aa^3de^3 + \frac{9Aa^2bd^2e^2}{2} + 3Aab^2d^3e + \frac{Ab^3d^4}{4} \right. \\
 & \left. + \frac{3Ba^3d^2e^2}{2} + 3Ba^2bd^3e + \frac{3Bab^2d^4}{4} \right) + x^3 \\
 & \cdot \left(2Aa^3d^2e^2 + 4Aa^2bd^3e + Aab^2d^4 + \frac{4Ba^3d^3e}{3} \right. \\
 & \left. + Ba^2bd^4 \right) + x^2 \cdot \left(2Aa^3d^3e + \frac{3Aa^2bd^4}{2} + \frac{Ba^3d^4}{2} \right)
 \end{aligned}$$

input `integrate((b*x+a)**3*(B*x+A)*(e*x+d)**4,x)`

output

```
A*a**3*d**4*x + B*b**3*e**4*x**9/9 + x**8*(A*b**3*e**4/8 + 3*B*a*b**2*e**4/8 + B*b**3*d*e**3/2) + x**7*(3*A*a*b**2*e**4/7 + 4*A*b**3*d*e**3/7 + 3*B*a**2*b*e**4/7 + 12*B*a*b**2*d*e**3/7 + 6*B*b**3*d**2*e**2/7) + x**6*(A*a**2*b*e**4/2 + 2*A*a*b**2*d*e**3 + A*b**3*d**2*e**2 + B*a**3*e**4/6 + 2*B*a**2*b*d*e**3 + 3*B*a*b**2*d**2*e**2 + 2*B*b**3*d**3*e/3) + x**5*(A*a**3*e**4/5 + 12*A*a**2*b*d*e**3/5 + 18*A*a*b**2*d**2*e**2/5 + 4*A*b**3*d**3*e/5 + 4*B*a**3*d*e**3/5 + 18*B*a**2*b*d**2*e**2/5 + 12*B*a*b**2*d**3*e/5 + B*b**3*d**4/5) + x**4*(A*a**3*d*e**3 + 9*A*a**2*b*d**2*e**2/2 + 3*A*a*b**2*d**3*e + A*b**3*d**4/4 + 3*B*a**3*d**2*e**2/2 + 3*B*a**2*b*d**3*e + 3*B*a*b**2*d**4/4) + x**3*(2*A*a**3*d**2*e**2 + 4*A*a**2*b*d**3*e + A*a*b**2*d**4 + 4*B*a**3*d**3*e/3 + B*a**2*b*d**4) + x**2*(2*A*a**3*d**3*e + 3*A*a**2*b*d**4/2 + B*a**3*d**4/2)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 425 vs. $2(153) = 306$.

Time = 0.04 (sec) , antiderivative size = 425, normalized size of antiderivative = 2.61

$$\int (a + bx)^3 (A + Bx)(d + ex)^4 dx$$

$$= \frac{1}{9} Bb^3 e^4 x^9 + Aa^3 d^4 x + \frac{1}{8} (4 Bb^3 de^3 + (3 Bab^2 + Ab^3) e^4) x^8$$

$$+ \frac{1}{7} (6 Bb^3 d^2 e^2 + 4 (3 Bab^2 + Ab^3) de^3 + 3 (Ba^2 b + Aab^2) e^4) x^7$$

$$+ \frac{1}{6} (4 Bb^3 d^3 e + 6 (3 Bab^2 + Ab^3) d^2 e^2 + 12 (Ba^2 b + Aab^2) de^3 + (Ba^3 + 3 Aa^2 b) e^4) x^6$$

$$+ \frac{1}{5} (Bb^3 d^4 + Aa^3 e^4 + 4 (3 Bab^2 + Ab^3) d^3 e + 18 (Ba^2 b + Aab^2) d^2 e^2 + 4 (Ba^3 + 3 Aa^2 b) de^3) x^5$$

$$+ \frac{1}{4} (4 Aa^3 de^3 + (3 Bab^2 + Ab^3) d^4 + 12 (Ba^2 b + Aab^2) d^3 e + 6 (Ba^3 + 3 Aa^2 b) d^2 e^2) x^4$$

$$+ \frac{1}{3} (6 Aa^3 d^2 e^2 + 3 (Ba^2 b + Aab^2) d^4 + 4 (Ba^3 + 3 Aa^2 b) d^3 e) x^3$$

$$+ \frac{1}{2} (4 Aa^3 d^3 e + (Ba^3 + 3 Aa^2 b) d^4) x^2$$

input

```
integrate((b*x+a)^3*(B*x+A)*(e*x+d)^4,x, algorithm="maxima")
```

output

```

1/9*B*b^3*e^4*x^9 + A*a^3*d^4*x + 1/8*(4*B*b^3*d*e^3 + (3*B*a*b^2 + A*b^3)
*e^4)*x^8 + 1/7*(6*B*b^3*d^2*e^2 + 4*(3*B*a*b^2 + A*b^3)*d*e^3 + 3*(B*a^2*
b + A*a*b^2)*e^4)*x^7 + 1/6*(4*B*b^3*d^3*e + 6*(3*B*a*b^2 + A*b^3)*d^2*e^2
+ 12*(B*a^2*b + A*a*b^2)*d*e^3 + (B*a^3 + 3*A*a^2*b)*e^4)*x^6 + 1/5*(B*b^
3*d^4 + A*a^3*e^4 + 4*(3*B*a*b^2 + A*b^3)*d^3*e + 18*(B*a^2*b + A*a*b^2)*d
^2*e^2 + 4*(B*a^3 + 3*A*a^2*b)*d*e^3)*x^5 + 1/4*(4*A*a^3*d*e^3 + (3*B*a*b^
2 + A*b^3)*d^4 + 12*(B*a^2*b + A*a*b^2)*d^3*e + 6*(B*a^3 + 3*A*a^2*b)*d^2*
e^2)*x^4 + 1/3*(6*A*a^3*d^2*e^2 + 3*(B*a^2*b + A*a*b^2)*d^4 + 4*(B*a^3 + 3
*A*a^2*b)*d^3*e)*x^3 + 1/2*(4*A*a^3*d^3*e + (B*a^3 + 3*A*a^2*b)*d^4)*x^2

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 534 vs. $2(153) = 306$.

Time = 0.12 (sec) , antiderivative size = 534, normalized size of antiderivative = 3.28

$$\begin{aligned}
\int (a + bx)^3(A + Bx)(d + ex)^4 dx = & \frac{1}{9} Bb^3e^4x^9 + \frac{1}{2} Bb^3de^3x^8 + \frac{3}{8} Bab^2e^4x^8 \\
& + \frac{1}{8} Ab^3e^4x^8 + \frac{6}{7} Bb^3d^2e^2x^7 + \frac{12}{7} Bab^2de^3x^7 \\
& + \frac{4}{7} Ab^3de^3x^7 + \frac{3}{7} Ba^2be^4x^7 + \frac{3}{7} Aab^2e^4x^7 \\
& + \frac{2}{3} Bb^3d^3ex^6 + 3 Bab^2d^2e^2x^6 + Ab^3d^2e^2x^6 \\
& + 2 Ba^2bde^3x^6 + 2 Aab^2de^3x^6 + \frac{1}{6} Ba^3e^4x^6 \\
& + \frac{1}{2} Aa^2be^4x^6 + \frac{1}{5} Bb^3d^4x^5 + \frac{12}{5} Bab^2d^3ex^5 \\
& + \frac{4}{5} Ab^3d^3ex^5 + \frac{18}{5} Ba^2bd^2e^2x^5 + \frac{18}{5} Aab^2d^2e^2x^5 \\
& + \frac{4}{5} Ba^3de^3x^5 + \frac{12}{5} Aa^2bde^3x^5 + \frac{1}{5} Aa^3e^4x^5 \\
& + \frac{3}{4} Bab^2d^4x^4 + \frac{1}{4} Ab^3d^4x^4 + 3 Ba^2bd^3ex^4 \\
& + 3 Aab^2d^3ex^4 + \frac{3}{2} Ba^3d^2e^2x^4 + \frac{9}{2} Aa^2bd^2e^2x^4 \\
& + Aa^3de^3x^4 + Ba^2bd^4x^3 + Aab^2d^4x^3 + \frac{4}{3} Ba^3d^3ex^3 \\
& + 4 Aa^2bd^3ex^3 + 2 Aa^3d^2e^2x^3 + \frac{1}{2} Ba^3d^4x^2 \\
& + \frac{3}{2} Aa^2bd^4x^2 + 2 Aa^3d^3ex^2 + Aa^3d^4x
\end{aligned}$$

input `integrate((b*x+a)^3*(B*x+A)*(e*x+d)^4,x, algorithm="giac")`

output

$$\begin{aligned}
 & 1/9*B*b^3*e^4*x^9 + 1/2*B*b^3*d*e^3*x^8 + 3/8*B*a*b^2*e^4*x^8 + 1/8*A*b^3* \\
 & e^4*x^8 + 6/7*B*b^3*d^2*e^2*x^7 + 12/7*B*a*b^2*d*e^3*x^7 + 4/7*A*b^3*d*e^3* \\
 & *x^7 + 3/7*B*a^2*b*e^4*x^7 + 3/7*A*a*b^2*e^4*x^7 + 2/3*B*b^3*d^3*e*x^6 + 3 \\
 & *B*a*b^2*d^2*e^2*x^6 + A*b^3*d^2*e^2*x^6 + 2*B*a^2*b*d*e^3*x^6 + 2*A*a*b^2 \\
 & *d*e^3*x^6 + 1/6*B*a^3*e^4*x^6 + 1/2*A*a^2*b*e^4*x^6 + 1/5*B*b^3*d^4*x^5 + \\
 & 12/5*B*a*b^2*d^3*e*x^5 + 4/5*A*b^3*d^3*e*x^5 + 18/5*B*a^2*b*d^2*e^2*x^5 + \\
 & 18/5*A*a*b^2*d^2*e^2*x^5 + 4/5*B*a^3*d*e^3*x^5 + 12/5*A*a^2*b*d*e^3*x^5 + \\
 & 1/5*A*a^3*e^4*x^5 + 3/4*B*a*b^2*d^4*x^4 + 1/4*A*b^3*d^4*x^4 + 3*B*a^2*b*d \\
 & ^3*e*x^4 + 3*A*a*b^2*d^3*e*x^4 + 3/2*B*a^3*d^2*e^2*x^4 + 9/2*A*a^2*b*d^2*e \\
 & ^2*x^4 + A*a^3*d*e^3*x^4 + B*a^2*b*d^4*x^3 + A*a*b^2*d^4*x^3 + 4/3*B*a^3*d \\
 & ^3*e*x^3 + 4*A*a^2*b*d^3*e*x^3 + 2*A*a^3*d^2*e^2*x^3 + 1/2*B*a^3*d^4*x^2 + \\
 & 3/2*A*a^2*b*d^4*x^2 + 2*A*a^3*d^3*e*x^2 + A*a^3*d^4*x
 \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.96 (sec) , antiderivative size = 439, normalized size of antiderivative = 2.69

$$\begin{aligned}
\int (a + bx)^3 (A + Bx)(d + ex)^4 dx = & x^3 \left(\frac{4Ba^3d^3e}{3} + 2Aa^3d^2e^2 + Ba^2bd^4 \right. \\
& \left. + 4Aa^2bd^3e + Aab^2d^4 \right) \\
& + x^7 \left(\frac{3Ba^2be^4}{7} + \frac{12Bab^2de^3}{7} + \frac{3Aab^2e^4}{7} \right. \\
& \left. + \frac{6Bb^3d^2e^2}{7} + \frac{4Ab^3de^3}{7} \right) \\
& + x^5 \left(\frac{4Ba^3de^3}{5} + \frac{Aa^3e^4}{5} + \frac{18Ba^2bd^2e^2}{5} \right. \\
& + \frac{12Aa^2bde^3}{5} + \frac{12Bab^2d^3e}{5} + \frac{18Aab^2d^2e^2}{5} \\
& \left. + \frac{Bb^3d^4}{5} + \frac{4Ab^3d^3e}{5} \right) + x^4 \left(\frac{3Ba^3d^2e^2}{2} \right. \\
& + Aa^3de^3 + 3Ba^2bd^3e + \frac{9Aa^2bd^2e^2}{2} \\
& \left. + \frac{3Bab^2d^4}{4} + 3Aab^2d^3e + \frac{Ab^3d^4}{4} \right) \\
& + x^6 \left(\frac{Ba^3e^4}{6} + 2Ba^2bde^3 + \frac{Aa^2be^4}{2} \right. \\
& + 3Bab^2d^2e^2 + 2Aab^2de^3 + \frac{2Bb^3d^3e}{3} \\
& \left. + Ab^3d^2e^2 \right) + \frac{a^2d^3x^2(4Aae + 3Abd + Bad)}{2} \\
& + \frac{b^2e^3x^8(Abe + 3Bae + 4Bbd)}{8} \\
& + Aa^3d^4x + \frac{Bb^3e^4x^9}{9}
\end{aligned}$$

input `int((A + B*x)*(a + b*x)^3*(d + e*x)^4,x)`

output

$$\begin{aligned} & x^3(A^2b^2d^4 + B^2a^2bd^4 + (4B^2a^3d^3e)/3 + 2A^2a^3d^2e^2 + 4A^2a^2bd^3e) + x^7((3A^2a^2b^2e^4)/7 + (3B^2a^2b^2e^4)/7 + (4A^2b^3d^2e^3)/7 + (6B^2b^3d^2e^2)/7 + (12B^2a^2bd^2e^3)/7) + x^5((A^2a^3e^4)/5 + (B^2b^3d^4)/5 + (4A^2b^3d^3e)/5 + (4B^2a^3d^2e^3)/5 + (18A^2a^2bd^2e^2)/5 + (18B^2a^2bd^2e^2)/5 + (12A^2a^2bd^2e^3)/5 + (12B^2a^2bd^2e^3)/5) + x^4((A^2b^3d^4)/4 + (3B^2a^2bd^4)/4 + A^2a^3d^2e^3 + (3B^2a^3d^2e^2)/2 + (9A^2a^2bd^2e^2)/2 + 3A^2a^2bd^3e + 3B^2a^2bd^3e) + x^6((B^2a^3e^4)/6 + (A^2a^2b^2e^4)/2 + (2B^2b^3d^3e)/3 + A^2b^3d^2e^2 + 3B^2a^2bd^2e^2 + 2A^2a^2bd^2e^3 + 2B^2a^2bd^2e^3) + (a^2d^3x^2(4A^2a^2e + 3A^2bd + B^2ad))/2 + (b^2e^3x^8(A^2b^2e + 3B^2a^2e + 4B^2bd))/8 + A^2a^3d^4x + (B^2b^3e^4x^9)/9 \end{aligned}$$
Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 322, normalized size of antiderivative = 1.98

$$\int (a + bx)^3(A + Bx)(d + ex)^4 dx$$

$$= \frac{x(70b^4e^4x^8 + 315ab^3e^4x^7 + 315b^4de^3x^7 + 540a^2b^2e^4x^6 + 1440ab^3de^3x^6 + 540b^4d^2e^2x^6 + 420a^3be^4x^5 + \dots)}{630}$$

input

$$\text{int}((b*x+a)^3*(B*x+A)*(e*x+d)^4,x)$$

output

$$\begin{aligned} & (x*(630*a**4*d**4 + 1260*a**4*d**3*e*x + 1260*a**4*d**2*e**2*x**2 + 630*a**4*d*e**3*x**3 + 126*a**4*e**4*x**4 + 1260*a**3*b*d**4*x + 3360*a**3*b*d**3*e*x**2 + 3780*a**3*b*d**2*e**2*x**3 + 2016*a**3*b*d*e**3*x**4 + 420*a**3*b*e**4*x**5 + 1260*a**2*b**2*d**4*x**2 + 3780*a**2*b**2*d**3*e*x**3 + 4536*a**2*b**2*d**2*e**2*x**4 + 2520*a**2*b**2*d*e**3*x**5 + 540*a**2*b**2*e**4*x**6 + 630*a*b**3*d**4*x**3 + 2016*a*b**3*d**3*e*x**4 + 2520*a*b**3*d**2*e**2*x**5 + 1440*a*b**3*d*e**3*x**6 + 315*a*b**3*e**4*x**7 + 126*b**4*d**4*x**4 + 420*b**4*d**3*e*x**5 + 540*b**4*d**2*e**2*x**6 + 315*b**4*d*e**3*x**7 + 70*b**4*e**4*x**8))/630 \end{aligned}$$

3.27 $\int (a + bx)^3 (A + Bx)(d + ex)^3 dx$

Optimal result	301
Mathematica [A] (verified)	302
Rubi [A] (verified)	302
Maple [B] (verified)	304
Fricas [B] (verification not implemented)	304
Sympy [B] (verification not implemented)	306
Maxima [B] (verification not implemented)	307
Giac [B] (verification not implemented)	307
Mupad [B] (verification not implemented)	309
Reduce [B] (verification not implemented)	310

Optimal result

Integrand size = 20, antiderivative size = 159

$$\int (a + bx)^3 (A + Bx)(d + ex)^3 dx = \frac{(Ab - aB)(bd - ae)^3 (a + bx)^4}{4b^5} + \frac{(bd - ae)^2 (bBd + 3Abe - 4aBe)(a + bx)^5}{5b^5} + \frac{e(bd - ae)(bBd + Abe - 2aBe)(a + bx)^6}{2b^5} + \frac{e^2(3bBd + Abe - 4aBe)(a + bx)^7}{7b^5} + \frac{Be^3(a + bx)^8}{8b^5}$$

output

```
1/4*(A*b-B*a)*(-a*e+b*d)^3*(b*x+a)^4/b^5+1/5*(-a*e+b*d)^2*(3*A*b*e-4*B*a*e
+B*b*d)*(b*x+a)^5/b^5+1/2*e*(-a*e+b*d)*(A*b*e-2*B*a*e+B*b*d)*(b*x+a)^6/b^5
+1/7*e^2*(A*b*e-4*B*a*e+3*B*b*d)*(b*x+a)^7/b^5+1/8*B*e^3*(b*x+a)^8/b^5
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 297, normalized size of antiderivative = 1.87

$$\int (a + bx)^3(A + Bx)(d + ex)^3 dx = a^3Ad^3x + \frac{1}{2}a^2d^2(aBd + 3A(bd + ae))x^2 + ad(aBd(bd + ae) + A(b^2d^2 + 3abde + a^2e^2))x^3 + \frac{1}{4}(3aBd(b^2d^2 + 3abde + a^2e^2) + A(b^3d^3 + 9ab^2d^2e + 9a^2bde^2 + a^3e^3))x^4 + \frac{1}{5}(a^3Be^3 + 9ab^2de(Bd + Ae) + 3a^2be^2(3Bd + Ae) + b^3d^2(Bd + 3Ae))x^5 + \frac{1}{2}be(a^2Be^2 + b^2d(Bd + Ae) + abe(3Bd + Ae))x^6 + \frac{1}{7}b^2e^2(3bBd + Abe + 3aBe)x^7 + \frac{1}{8}b^3Be^3x^8$$

input

```
Integrate[(a + b*x)^3*(A + B*x)*(d + e*x)^3,x]
```

output

```
a^3*A*d^3*x + (a^2*d^2*(a*B*d + 3*A*(b*d + a*e))*x^2)/2 + a*d*(a*B*d*(b*d + a*e) + A*(b^2*d^2 + 3*a*b*d*e + a^2*e^2))*x^3 + ((3*a*B*d*(b^2*d^2 + 3*a*b*d*e + a^2*e^2) + A*(b^3*d^3 + 9*a*b^2*d^2*e + 9*a^2*b*d*e^2 + a^3*e^3))*x^4)/4 + ((a^3*B*e^3 + 9*a*b^2*d*e*(B*d + A*e) + 3*a^2*b*e^2*(3*B*d + A*e) + b^3*d^2*(B*d + 3*A*e))*x^5)/5 + (b*e*(a^2*B*e^2 + b^2*d*(B*d + A*e) + a*b*e*(3*B*d + A*e))*x^6)/2 + (b^2*e^2*(3*b*B*d + A*b*e + 3*a*B*e)*x^7)/7 + (b^3*B*e^3*x^8)/8
```

Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^3 (A + Bx)(d + ex)^3 dx$$

↓ 86

$$\int \left(\frac{e^2(a + bx)^6(-4aBe + Abe + 3bBd)}{b^4} + \frac{3e(a + bx)^5(bd - ae)(-2aBe + Abe + bBd)}{b^4} + \frac{(a + bx)^4(bd - ae)^2}{b^4} \right) dx$$

↓ 2009

$$\frac{e^2(a + bx)^7(-4aBe + Abe + 3bBd)}{7b^5} + \frac{e(a + bx)^6(bd - ae)(-2aBe + Abe + bBd)}{2b^5} + \frac{(a + bx)^5(bd - ae)^2(-4aBe + 3Abe + bBd)}{5b^5} + \frac{(a + bx)^4(Ab - aB)(bd - ae)^3}{4b^5} + \frac{Be^3(a + bx)^8}{8b^5}$$

input `Int[(a + b*x)^3*(A + B*x)*(d + e*x)^3,x]`

output `((A*b - a*B)*(b*d - a*e)^3*(a + b*x)^4)/(4*b^5) + ((b*d - a*e)^2*(b*B*d + 3*A*b*e - 4*a*B*e)*(a + b*x)^5)/(5*b^5) + (e*(b*d - a*e)*(b*B*d + A*b*e - 2*a*B*e)*(a + b*x)^6)/(2*b^5) + (e^2*(3*b*B*d + A*b*e - 4*a*B*e)*(a + b*x)^7)/(7*b^5) + (B*e^3*(a + b*x)^8)/(8*b^5)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 338 vs. 2(149) = 298.

Time = 0.17 (sec) , antiderivative size = 339, normalized size of antiderivative = 2.13

method	result
default	$\frac{b^3 B e^3 x^8}{8} + \frac{((b^3 A + 3a b^2 B) e^3 + 3b^3 B d e^2) x^7}{7} + \frac{((3a b^2 A + 3a^2 b B) e^3 + 3(b^3 A + 3a b^2 B) d e^2 + 3b^3 B d^2 e) x^6}{6} + \frac{((3a^2 b A + 3a^2 B) e^3 + 3a b^2 B d e^2) x^5}{5} + \frac{((3a^2 b A + 3a^2 B) e^3 + 3a b^2 B d e^2) x^4}{4} + \frac{((3a^2 b A + 3a^2 B) e^3 + 3a b^2 B d e^2) x^3}{3} + \frac{((3a^2 b A + 3a^2 B) e^3 + 3a b^2 B d e^2) x^2}{2} + \frac{((3a^2 b A + 3a^2 B) e^3 + 3a b^2 B d e^2) x}{1} + \frac{((3a^2 b A + 3a^2 B) e^3 + 3a b^2 B d e^2)}{0}$
norman	$\frac{b^3 B e^3 x^8}{8} + (\frac{1}{7} A b^3 e^3 + \frac{3}{7} B a b^2 e^3 + \frac{3}{7} b^3 B d e^2) x^7 + (\frac{1}{2} A a b^2 e^3 + \frac{1}{2} A b^3 d e^2 + \frac{1}{2} B a^2 b e^3 + \frac{3}{2} B a b^2 d e^2) x^6 + (\frac{1}{4} A a^2 b d e^2 + \frac{1}{4} A a b^2 d^2 e^2 + \frac{1}{4} B a^2 b^2 d e^2) x^5 + (\frac{1}{4} A a^2 b d e^2 + \frac{1}{4} A a b^2 d^2 e^2 + \frac{1}{4} B a^2 b^2 d e^2) x^4 + (\frac{1}{4} A a^2 b d e^2 + \frac{1}{4} A a b^2 d^2 e^2 + \frac{1}{4} B a^2 b^2 d e^2) x^3 + (\frac{1}{4} A a^2 b d e^2 + \frac{1}{4} A a b^2 d^2 e^2 + \frac{1}{4} B a^2 b^2 d e^2) x^2 + (\frac{1}{4} A a^2 b d e^2 + \frac{1}{4} A a b^2 d^2 e^2 + \frac{1}{4} B a^2 b^2 d e^2) x + (\frac{1}{4} A a^2 b d e^2 + \frac{1}{4} A a b^2 d^2 e^2 + \frac{1}{4} B a^2 b^2 d e^2)$
gosper	$\frac{3}{2} x^6 B a b^2 d e^2 + \frac{9}{5} x^5 A a b^2 d e^2 + \frac{9}{5} x^5 B a^2 b d e^2 + \frac{9}{5} x^5 B a b^2 d^2 e^2 + \frac{9}{4} x^4 A a^2 b d e^2 + \frac{9}{4} x^4 A a b^2 d^2 e^2 + \frac{9}{4} x^4 B a^2 b^2 d e^2$
risch	$\frac{3}{2} x^6 B a b^2 d e^2 + \frac{9}{5} x^5 A a b^2 d e^2 + \frac{9}{5} x^5 B a^2 b d e^2 + \frac{9}{5} x^5 B a b^2 d^2 e^2 + \frac{9}{4} x^4 A a^2 b d e^2 + \frac{9}{4} x^4 A a b^2 d^2 e^2 + \frac{9}{4} x^4 B a^2 b^2 d e^2$
parallelrisch	$\frac{3}{2} x^6 B a b^2 d e^2 + \frac{9}{5} x^5 A a b^2 d e^2 + \frac{9}{5} x^5 B a^2 b d e^2 + \frac{9}{5} x^5 B a b^2 d^2 e^2 + \frac{9}{4} x^4 A a^2 b d e^2 + \frac{9}{4} x^4 A a b^2 d^2 e^2 + \frac{9}{4} x^4 B a^2 b^2 d e^2$
orering	$x(35b^3 B e^3 x^7 + 40A b^3 e^3 x^6 + 120B a b^2 e^3 x^6 + 120B b^3 d e^2 x^6 + 140A a b^2 e^3 x^5 + 140A b^3 d e^2 x^5 + 140B a^2 b e^3 x^5 + 420B a b^2 d e^2 x^5 + 140A a^2 b d e^2 x^4 + 140A a b^2 d^2 e^2 x^4 + 140B a^2 b^2 d e^2 x^4 + 140A a^2 b d e^2 x^3 + 140A a b^2 d^2 e^2 x^3 + 140B a^2 b^2 d e^2 x^3 + 140A a^2 b d e^2 x^2 + 140A a b^2 d^2 e^2 x^2 + 140B a^2 b^2 d e^2 x^2 + 140A a^2 b d e^2 x + 140A a b^2 d^2 e^2 x + 140B a^2 b^2 d e^2 x + 140A a^2 b d e^2 + 140A a b^2 d^2 e^2 + 140B a^2 b^2 d e^2)$

```
input int((b*x+a)^3*(B*x+A)*(e*x+d)^3,x,method=_RETURNVERBOSE)
```

```
output 1/8*b^3*B*e^3*x^8+1/7*((A*b^3+3*B*a*b^2)*e^3+3*b^3*B*d*e^2)*x^7+1/6*((3*A*a*b^2+3*B*a^2*b)*e^3+3*(A*b^3+3*B*a*b^2)*d*e^2+3*b^3*B*d^2*e)*x^6+1/5*((3*A*a^2*b+B*a^3)*e^3+3*(3*A*a*b^2+3*B*a^2*b)*d*e^2+3*(A*b^3+3*B*a*b^2)*d^2*e+b^3*B*d^3)*x^5+1/4*(a^3*A*e^3+3*(3*A*a^2*b+B*a^3)*d*e^2+3*(3*A*a*b^2+3*B*a^2*b)*d^2*e+(A*b^3+3*B*a*b^2)*d^3)*x^4+1/3*(3*a^3*A*d*e^2+3*(3*A*a^2*b+B*a^3)*d^2*e+(3*A*a*b^2+3*B*a^2*b)*d^3)*x^3+1/2*(3*a^3*A*d^2*e+(3*A*a^2*b+B*a^3)*d^3)*x^2+a^3*A*d^3*x
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 325 vs. 2(149) = 298.

Time = 0.07 (sec) , antiderivative size = 325, normalized size of antiderivative = 2.04

$$\int (a + bx)^3(A + Bx)(d + ex)^3 dx$$

$$= \frac{1}{8} Bb^3 e^3 x^8 + Aa^3 d^3 x + \frac{1}{7} (3 Bb^3 de^2 + (3 Bab^2 + Ab^3) e^3) x^7$$

$$+ \frac{1}{2} (Bb^3 d^2 e + (3 Bab^2 + Ab^3) de^2 + (Ba^2 b + Aab^2) e^3) x^6$$

$$+ \frac{1}{5} (Bb^3 d^3 + 3 (3 Bab^2 + Ab^3) d^2 e + 9 (Ba^2 b + Aab^2) de^2 + (Ba^3 + 3 Aa^2 b) e^3) x^5$$

$$+ \frac{1}{4} (Aa^3 e^3 + (3 Bab^2 + Ab^3) d^3 + 9 (Ba^2 b + Aab^2) d^2 e + 3 (Ba^3 + 3 Aa^2 b) de^2) x^4$$

$$+ (Aa^3 de^2 + (Ba^2 b + Aab^2) d^3 + (Ba^3 + 3 Aa^2 b) d^2 e) x^3$$

$$+ \frac{1}{2} (3 Aa^3 d^2 e + (Ba^3 + 3 Aa^2 b) d^3) x^2$$

input `integrate((b*x+a)^3*(B*x+A)*(e*x+d)^3,x, algorithm="fricas")`

output `1/8*B*b^3*e^3*x^8 + A*a^3*d^3*x + 1/7*(3*B*b^3*d*e^2 + (3*B*a*b^2 + A*b^3)*e^3)*x^7 + 1/2*(B*b^3*d^2*e + (3*B*a*b^2 + A*b^3)*d*e^2 + (B*a^2*b + A*a*b^2)*e^3)*x^6 + 1/5*(B*b^3*d^3 + 3*(3*B*a*b^2 + A*b^3)*d^2*e + 9*(B*a^2*b + A*a*b^2)*d*e^2 + (B*a^3 + 3*A*a^2*b)*e^3)*x^5 + 1/4*(A*a^3*e^3 + (3*B*a*b^2 + A*b^3)*d^3 + 9*(B*a^2*b + A*a*b^2)*d^2*e + 3*(B*a^3 + 3*A*a^2*b)*d*e^2)*x^4 + (A*a^3*d*e^2 + (B*a^2*b + A*a*b^2)*d^3 + (B*a^3 + 3*A*a^2*b)*d^2*e)*x^3 + 1/2*(3*A*a^3*d^2*e + (B*a^3 + 3*A*a^2*b)*d^3)*x^2`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 422 vs. $2(155) = 310$.

Time = 0.05 (sec) , antiderivative size = 422, normalized size of antiderivative = 2.65

$$\int (a + bx)^3(A + Bx)(d + ex)^3 dx = Aa^3d^3x + \frac{Bb^3e^3x^8}{8} + x^7\left(\frac{Ab^3e^3}{7} + \frac{3Bab^2e^3}{7} + \frac{3Bb^3de^2}{7}\right) + x^6\left(\frac{Aab^2e^3}{2} + \frac{Ab^3de^2}{2} + \frac{Ba^2be^3}{2} + \frac{3Bab^2de^2}{2} + \frac{Bb^3d^2e}{2}\right) + x^5\left(\frac{3Aa^2be^3}{5} + \frac{9Aab^2de^2}{5} + \frac{3Ab^3d^2e}{5} + \frac{Ba^3e^3}{5} + \frac{9Ba^2bde^2}{5} + \frac{9Bab^2d^2e}{5} + \frac{Bb^3d^3}{5}\right) + x^4\left(\frac{Aa^3e^3}{4} + \frac{9Aa^2bde^2}{4} + \frac{9Aab^2d^2e}{4} + \frac{Ab^3d^3}{4} + \frac{3Ba^3de^2}{4} + \frac{9Ba^2bd^2e}{4} + \frac{3Bab^2d^3}{4}\right) + x^3(Aa^3de^2 + 3Aa^2bd^2e + Aab^2d^3 + Ba^3d^2e + Ba^2bd^3) + x^2\left(\frac{3Aa^3d^2e}{2} + \frac{3Aa^2bd^3}{2} + \frac{Ba^3d^3}{2}\right)$$

input `integrate((b*x+a)**3*(B*x+A)*(e*x+d)**3,x)`

output `A*a**3*d**3*x + B*b**3*e**3*x**8/8 + x**7*(A*b**3*e**3/7 + 3*B*a*b**2*e**3/7 + 3*B*b**3*d*e**2/7) + x**6*(A*a*b**2*e**3/2 + A*b**3*d*e**2/2 + B*a**2*b*e**3/2 + 3*B*a*b**2*d*e**2/2 + B*b**3*d**2*e/2) + x**5*(3*A*a**2*b*e**3/5 + 9*A*a*b**2*d*e**2/5 + 3*A*b**3*d**2*e/5 + B*a**3*e**3/5 + 9*B*a**2*b*d*e**2/5 + 9*B*a*b**2*d**2*e/5 + B*b**3*d**3/5) + x**4*(A*a**3*e**3/4 + 9*A*a**2*b*d*e**2/4 + 9*A*a*b**2*d**2*e/4 + A*b**3*d**3/4 + 3*B*a**3*d*e**2/4 + 9*B*a**2*b*d**2*e/4 + 3*B*a*b**2*d**3/4) + x**3*(A*a**3*d*e**2 + 3*A*a**2*b*d**2*e + A*a*b**2*d**3 + B*a**3*d**2*e + B*a**2*b*d**3) + x**2*(3*A*a**3*d**2*e/2 + 3*A*a**2*b*d**3/2 + B*a**3*d**3/2)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 325 vs. $2(149) = 298$.

Time = 0.03 (sec) , antiderivative size = 325, normalized size of antiderivative = 2.04

$$\int (a + bx)^3 (A + Bx)(d + ex)^3 dx$$

$$= \frac{1}{8} Bb^3 e^3 x^8 + Aa^3 d^3 x + \frac{1}{7} (3 Bb^3 de^2 + (3 Bab^2 + Ab^3) e^3) x^7$$

$$+ \frac{1}{2} (Bb^3 d^2 e + (3 Bab^2 + Ab^3) de^2 + (Ba^2 b + Aab^2) e^3) x^6$$

$$+ \frac{1}{5} (Bb^3 d^3 + 3 (3 Bab^2 + Ab^3) d^2 e + 9 (Ba^2 b + Aab^2) de^2 + (Ba^3 + 3 Aa^2 b) e^3) x^5$$

$$+ \frac{1}{4} (Aa^3 e^3 + (3 Bab^2 + Ab^3) d^3 + 9 (Ba^2 b + Aab^2) d^2 e + 3 (Ba^3 + 3 Aa^2 b) de^2) x^4$$

$$+ (Aa^3 de^2 + (Ba^2 b + Aab^2) d^3 + (Ba^3 + 3 Aa^2 b) d^2 e) x^3$$

$$+ \frac{1}{2} (3 Aa^3 d^2 e + (Ba^3 + 3 Aa^2 b) d^3) x^2$$

input `integrate((b*x+a)^3*(B*x+A)*(e*x+d)^3,x, algorithm="maxima")`

output `1/8*B*b^3*e^3*x^8 + A*a^3*d^3*x + 1/7*(3*B*b^3*d*e^2 + (3*B*a*b^2 + A*b^3)*e^3)*x^7 + 1/2*(B*b^3*d^2*e + (3*B*a*b^2 + A*b^3)*d*e^2 + (B*a^2*b + A*a*b^2)*e^3)*x^6 + 1/5*(B*b^3*d^3 + 3*(3*B*a*b^2 + A*b^3)*d^2*e + 9*(B*a^2*b + A*a*b^2)*d*e^2 + (B*a^3 + 3*A*a^2*b)*e^3)*x^5 + 1/4*(A*a^3*e^3 + (3*B*a*b^2 + A*b^3)*d^3 + 9*(B*a^2*b + A*a*b^2)*d^2*e + 3*(B*a^3 + 3*A*a^2*b)*d*e^2)*x^4 + (A*a^3*d*e^2 + (B*a^2*b + A*a*b^2)*d^3 + (B*a^3 + 3*A*a^2*b)*d^2*e)*x^3 + 1/2*(3*A*a^3*d^2*e + (B*a^3 + 3*A*a^2*b)*d^3)*x^2`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 410 vs. $2(149) = 298$.

Time = 0.12 (sec) , antiderivative size = 410, normalized size of antiderivative = 2.58

$$\int (a + bx)^3(A + Bx)(d + ex)^3 dx = \frac{1}{8} Bb^3e^3x^8 + \frac{3}{7} Bb^3de^2x^7 + \frac{3}{7} Bab^2e^3x^7 + \frac{1}{7} Ab^3e^3x^7$$

$$+ \frac{1}{2} Bb^3d^2ex^6 + \frac{3}{2} Bab^2de^2x^6 + \frac{1}{2} Ab^3de^2x^6$$

$$+ \frac{1}{2} Ba^2be^3x^6 + \frac{1}{2} Aab^2e^3x^6 + \frac{1}{5} Bb^3d^3x^5$$

$$+ \frac{9}{5} Bab^2d^2ex^5 + \frac{3}{5} Ab^3d^2ex^5 + \frac{9}{5} Ba^2bde^2x^5$$

$$+ \frac{9}{5} Aab^2de^2x^5 + \frac{1}{5} Ba^3e^3x^5 + \frac{3}{5} Aa^2be^3x^5$$

$$+ \frac{3}{4} Bab^2d^3x^4 + \frac{1}{4} Ab^3d^3x^4 + \frac{9}{4} Ba^2bd^2ex^4$$

$$+ \frac{9}{4} Aab^2d^2ex^4 + \frac{3}{4} Ba^3de^2x^4 + \frac{9}{4} Aa^2bde^2x^4$$

$$+ \frac{1}{4} Aa^3e^3x^4 + Ba^2bd^3x^3 + Aab^2d^3x^3 + Ba^3d^2ex^3$$

$$+ 3Aa^2bd^2ex^3 + Aa^3de^2x^3 + \frac{1}{2} Ba^3d^3x^2$$

$$+ \frac{3}{2} Aa^2bd^3x^2 + \frac{3}{2} Aa^3d^2ex^2 + Aa^3d^3x$$

input `integrate((b*x+a)^3*(B*x+A)*(e*x+d)^3,x, algorithm="giac")`

output `1/8*B*b^3*e^3*x^8 + 3/7*B*b^3*d*e^2*x^7 + 3/7*B*a*b^2*e^3*x^7 + 1/7*A*b^3*e^3*x^7 + 1/2*B*b^3*d^2*e*x^6 + 3/2*B*a*b^2*d*e^2*x^6 + 1/2*A*b^3*d*e^2*x^6 + 1/2*B*a^2*b*e^3*x^6 + 1/2*A*a*b^2*e^3*x^6 + 1/5*B*b^3*d^3*x^5 + 9/5*B*a*b^2*d^2*e*x^5 + 3/5*A*b^3*d^2*e*x^5 + 9/5*B*a^2*b*d*e^2*x^5 + 9/5*A*a*b^2*d*e^2*x^5 + 1/5*B*a^3*e^3*x^5 + 3/5*A*a^2*b*e^3*x^5 + 3/4*B*a*b^2*d^3*x^4 + 1/4*A*b^3*d^3*x^4 + 9/4*B*a^2*b*d^2*e*x^4 + 9/4*A*a*b^2*d^2*e*x^4 + 3/4*B*a^3*d*e^2*x^4 + 9/4*A*a^2*b*d*e^2*x^4 + 1/4*A*a^3*e^3*x^4 + B*a^2*b*d^3*x^3 + A*a*b^2*d^3*x^3 + B*a^3*d^2*e*x^3 + 3*A*a^2*b*d^2*e*x^3 + A*a^3*d^2*e^2*x^3 + 1/2*B*a^3*d^3*x^2 + 3/2*A*a^2*b*d^3*x^2 + 3/2*A*a^3*d^2*e*x^2 + A*a^3*d^3*x`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 334, normalized size of antiderivative = 2.10

$$\int (a + bx)^3 (A + Bx)(d + ex)^3 dx = x^3 (Ba^3 d^2 e + Aa^3 d e^2 + Ba^2 b d^3 + 3Aa^2 b d^2 e + Aa b^2 d^3) + x^6 \left(\frac{Ba^2 b e^3}{2} + \frac{3Bab^2 d e^2}{2} + \frac{Aa b^2 e^3}{2} + \frac{Bb^3 d^2 e}{2} + \frac{Ab^3 d e^2}{2} \right) + x^4 \left(\frac{3Ba^3 d e^2}{4} + \frac{Aa^3 e^3}{4} + \frac{9Ba^2 b d^2 e}{4} + \frac{9Aa^2 b d e^2}{4} + \frac{3Bab^2 d^3}{4} + \frac{9Aab^2 d^2 e}{4} + \frac{Ab^3 d^3}{4} \right) + x^5 \left(\frac{Ba^3 e^3}{5} + \frac{9Ba^2 b d e^2}{5} + \frac{3Aa^2 b e^3}{5} + \frac{9Bab^2 d^2 e}{5} + \frac{9Aab^2 d e^2}{5} + \frac{Bb^3 d^3}{5} + \frac{3Ab^3 d^2 e}{5} \right) + \frac{a^2 d^2 x^2 (3Aae + 3Abd + B ad)}{2} + \frac{b^2 e^2 x^7 (Abe + 3Bae + 3Bbd)}{7} + Aa^3 d^3 x + \frac{Bb^3 e^3 x^8}{8}$$

input `int((A + B*x)*(a + b*x)^3*(d + e*x)^3,x)`output `x^3*(A*a*b^2*d^3 + B*a^2*b*d^3 + A*a^3*d*e^2 + B*a^3*d^2*e + 3*A*a^2*b*d^2*e) + x^6*((A*a*b^2*d^3)/2 + (B*a^2*b*d^3)/2 + (A*b^3*d^2*e^2)/2 + (B*b^3*d^2*e)/2 + (3*B*a*b^2*d^2*e^2)/2) + x^4*((A*a^3*d^3)/4 + (A*b^3*d^3)/4 + (3*B*a*b^2*d^3)/4 + (3*B*a^3*d^2*e^2)/4 + (9*A*a*b^2*d^2*e^2)/4 + (9*A*a^2*b*d^2*e^2)/4 + (9*B*a^2*b*d^2*e^2)/4) + x^5*((B*a^3*d^3)/5 + (B*b^3*d^3)/5 + (3*A*a^2*b*d^2*e^2)/5 + (3*A*b^3*d^2*e^2)/5 + (9*A*a*b^2*d^2*e^2)/5 + (9*B*a*b^2*d^2*e^2)/5 + (9*B*a^2*b*d^2*e^2)/5) + (a^2*d^2*x^2*(3*A*a*e + 3*A*b*d + B*a*d))/2 + (b^2*e^2*x^7*(A*b*e + 3*B*a*e + 3*B*b*d))/7 + A*a^3*d^3*x + (B*b^3*e^3*x^8)/8`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 247, normalized size of antiderivative = 1.55

$$\int (a + bx)^3 (A + Bx)(d + ex)^3 dx$$

$$= \frac{x(35b^4e^3x^7 + 160ab^3e^3x^6 + 120b^4de^2x^6 + 280a^2b^2e^3x^5 + 560ab^3de^2x^5 + 140b^4d^2ex^5 + 224a^3be^3x^4 + 160a^2b^2d^2e^2x^4 + 120ab^3d^2ex^4 + 35b^4d^2e^2x^4 + 160a^2b^2de^2x^3 + 120ab^3de^2x^3 + 35b^4de^2x^3 + 160a^2b^2d^2ex^3 + 120ab^3d^2ex^3 + 35b^4d^2ex^3 + 160a^2b^2d^2e^2x^2 + 120ab^3d^2e^2x^2 + 35b^4d^2e^2x^2 + 160a^2b^2d^2e^2x + 120ab^3d^2e^2x + 35b^4d^2e^2x + 160a^2b^2d^2e^2 + 120ab^3d^2e^2 + 35b^4d^2e^2)}{280}$$

input `int((b*x+a)^3*(B*x+A)*(e*x+d)^3,x)`output `(x*(280*a**4*d**3 + 420*a**4*d**2*e*x + 280*a**4*d*e**2*x**2 + 70*a**4*e**3*x**3 + 560*a**3*b*d**3*x + 1120*a**3*b*d**2*e*x**2 + 840*a**3*b*d*e**2*x**3 + 224*a**3*b*e**3*x**4 + 560*a**2*b**2*d**3*x**2 + 1260*a**2*b**2*d**2*e*x**3 + 1008*a**2*b**2*d*e**2*x**4 + 280*a**2*b**2*e**3*x**5 + 280*a*b**3*d**3*x**3 + 672*a*b**3*d**2*e*x**4 + 560*a*b**3*d*e**2*x**5 + 160*a*b**3*e**3*x**6 + 56*b**4*d**3*x**4 + 140*b**4*d**2*e*x**5 + 120*b**4*d*e**2*x**6 + 35*b**4*e**3*x**7))/280`

3.28 $\int (a + bx)^3 (A + Bx)(d + ex)^2 dx$

Optimal result	311
Mathematica [A] (verified)	312
Rubi [A] (verified)	312
Maple [B] (verified)	313
Fricas [B] (verification not implemented)	314
Sympy [B] (verification not implemented)	315
Maxima [B] (verification not implemented)	316
Giac [B] (verification not implemented)	317
Mupad [B] (verification not implemented)	318
Reduce [B] (verification not implemented)	318

Optimal result

Integrand size = 20, antiderivative size = 118

$$\int (a + bx)^3 (A + Bx)(d + ex)^2 dx = \frac{(Ab - aB)(bd - ae)^2 (a + bx)^4}{4b^4} + \frac{(bd - ae)(bBd + 2Abe - 3aBe)(a + bx)^5}{5b^4} + \frac{e(2bBd + Abe - 3aBe)(a + bx)^6}{6b^4} + \frac{Be^2(a + bx)^7}{7b^4}$$

output

```
1/4*(A*b-B*a)*(-a*e+b*d)^2*(b*x+a)^4/b^4+1/5*(-a*e+b*d)*(2*A*b*e-3*B*a*e+B
*b*d)*(b*x+a)^5/b^4+1/6*e*(A*b*e-3*B*a*e+2*B*b*d)*(b*x+a)^6/b^4+1/7*B*e^2*
(b*x+a)^7/b^4
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.90

$$\int (a + bx)^3(A + Bx)(d + ex)^2 dx = a^3Ad^2x + \frac{1}{2}a^2d(3Abd + aBd + 2aAe)x^2$$

$$+ \frac{1}{3}a(aBd(3bd + 2ae) + A(3b^2d^2 + 6abde + a^2e^2))x^3$$

$$+ \frac{1}{4}(aB(3b^2d^2 + 6abde + a^2e^2) + Ab(b^2d^2 + 6abde + 3a^2e^2))x^4$$

$$+ \frac{1}{5}b(3a^2Be^2 + 3abe(2Bd + Ae) + b^2d(Bd + 2Ae))x^5$$

$$+ \frac{1}{6}b^2e(2bBd + Abe + 3aBe)x^6 + \frac{1}{7}b^3Be^2x^7$$

input

```
Integrate[(a + b*x)^3*(A + B*x)*(d + e*x)^2,x]
```

output

```
a^3*A*d^2*x + (a^2*d*(3*A*b*d + a*B*d + 2*a*A*e))*x^2)/2 + (a*(a*B*d*(3*b*d
+ 2*a*e) + A*(3*b^2*d^2 + 6*a*b*d*e + a^2*e^2))*x^3)/3 + ((a*B*(3*b^2*d^2
+ 6*a*b*d*e + a^2*e^2) + A*b*(b^2*d^2 + 6*a*b*d*e + 3*a^2*e^2))*x^4)/4 +
(b*(3*a^2*B*e^2 + 3*a*b*e*(2*B*d + A*e) + b^2*d*(B*d + 2*A*e))*x^5)/5 + (b
^2*e*(2*b*B*d + A*b*e + 3*a*B*e))*x^6)/6 + (b^3*B*e^2*x^7)/7
```

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.00,
 number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules
 used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^3(A + Bx)(d + ex)^2 dx$$

↓ 86

$$\int \left(\frac{e(a+bx)^5(-3aBe + Abe + 2bBd)}{b^3} + \frac{(a+bx)^4(bd - ae)(-3aBe + 2Abe + bBd)}{b^3} + \frac{(a+bx)^3(Ab - aB)(bd - ae)}{b^3} \right) dx$$

↓ 2009

$$\frac{e(a+bx)^6(-3aBe + Abe + 2bBd)}{6b^4} + \frac{(a+bx)^5(bd - ae)(-3aBe + 2Abe + bBd)}{5b^4} + \frac{(a+bx)^4(Ab - aB)(bd - ae)^2}{4b^4} + \frac{Be^2(a+bx)^7}{7b^4}$$

input `Int[(a + b*x)^3*(A + B*x)*(d + e*x)^2,x]`

output `((A*b - a*B)*(b*d - a*e)^2*(a + b*x)^4)/(4*b^4) + ((b*d - a*e)*(b*B*d + 2*A*b*e - 3*a*B*e)*(a + b*x)^5)/(5*b^4) + (e*(2*b*B*d + A*b*e - 3*a*B*e)*(a + b*x)^6)/(6*b^4) + (B*e^2*(a + b*x)^7)/(7*b^4)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;`
`FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 243 vs. 2(110) = 220.

Time = 0.17 (sec) , antiderivative size = 244, normalized size of antiderivative = 2.07

output

```
1/7*B*b^3*e^2*x^7 + A*a^3*d^2*x + 1/6*(2*B*b^3*d*e + (3*B*a*b^2 + A*b^3)*e^2)*x^6 + 1/5*(B*b^3*d^2 + 2*(3*B*a*b^2 + A*b^3)*d*e + 3*(B*a^2*b + A*a*b^2)*e^2)*x^5 + 1/4*((3*B*a*b^2 + A*b^3)*d^2 + 6*(B*a^2*b + A*a*b^2)*d*e + (B*a^3 + 3*A*a^2*b)*e^2)*x^4 + 1/3*(A*a^3*e^2 + 3*(B*a^2*b + A*a*b^2)*d^2 + 2*(B*a^3 + 3*A*a^2*b)*d*e)*x^3 + 1/2*(2*A*a^3*d*e + (B*a^3 + 3*A*a^2*b)*d^2)*x^2
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 296 vs. $2(116) = 232$.

Time = 0.05 (sec) , antiderivative size = 296, normalized size of antiderivative = 2.51

$$\int (a + bx)^3(A + Bx)(d + ex)^2 dx = Aa^3d^2x + \frac{Bb^3e^2x^7}{7} + x^6 \left(\frac{Ab^3e^2}{6} + \frac{Bab^2e^2}{2} + \frac{Bb^3de}{3} \right) + x^5 \cdot \left(\frac{3Aab^2e^2}{5} + \frac{2Ab^3de}{5} + \frac{3Ba^2be^2}{5} + \frac{6Bab^2de}{5} + \frac{Bb^3d^2}{5} \right) + x^4 \cdot \left(\frac{3Aa^2be^2}{4} + \frac{3Aab^2de}{2} + \frac{Ab^3d^2}{4} + \frac{Ba^3e^2}{4} + \frac{3Ba^2bde}{2} + \frac{3Bab^2d^2}{4} \right) + x^3 \left(\frac{Aa^3e^2}{3} + 2Aa^2bde + Aab^2d^2 + \frac{2Ba^3de}{3} + Ba^2bd^2 \right) + x^2 \left(Aa^3de + \frac{3Aa^2bd^2}{2} + \frac{Ba^3d^2}{2} \right)$$

input

```
integrate((b*x+a)**3*(B*x+A)*(e*x+d)**2,x)
```

output

```
A*a**3*d**2*x + B*b**3*e**2*x**7/7 + x**6*(A*b**3*e**2/6 + B*a*b**2*e**2/2 + B*b**3*d*e/3) + x**5*(3*A*a*b**2*e**2/5 + 2*A*b**3*d*e/5 + 3*B*a**2*b*e**2/5 + 6*B*a*b**2*d*e/5 + B*b**3*d**2/5) + x**4*(3*A*a**2*b*e**2/4 + 3*A*a*b**2*d*e/2 + A*b**3*d**2/4 + B*a**3*e**2/4 + 3*B*a**2*b*d*e/2 + 3*B*a*b**2*d**2/4) + x**3*(A*a**3*e**2/3 + 2*A*a**2*b*d*e + A*a*b**2*d**2 + 2*B*a**3*d*e/3 + B*a**2*b*d**2) + x**2*(A*a**3*d*e + 3*A*a**2*b*d**2/2 + B*a**3*d**2/2)
```


Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 239 vs. $2(110) = 220$.

Time = 0.04 (sec) , antiderivative size = 239, normalized size of antiderivative = 2.03

$$\begin{aligned} & \int (a + bx)^3 (A + Bx)(d + ex)^2 dx \\ &= \frac{1}{7} Bb^3 e^2 x^7 + Aa^3 d^2 x + \frac{1}{6} (2 Bb^3 de + (3 Bab^2 + Ab^3) e^2) x^6 \\ &+ \frac{1}{5} (Bb^3 d^2 + 2 (3 Bab^2 + Ab^3) de + 3 (Ba^2 b + Aab^2) e^2) x^5 \\ &+ \frac{1}{4} ((3 Bab^2 + Ab^3) d^2 + 6 (Ba^2 b + Aab^2) de + (Ba^3 + 3 Aa^2 b) e^2) x^4 \\ &+ \frac{1}{3} (Aa^3 e^2 + 3 (Ba^2 b + Aab^2) d^2 + 2 (Ba^3 + 3 Aa^2 b) de) x^3 \\ &+ \frac{1}{2} (2 Aa^3 de + (Ba^3 + 3 Aa^2 b) d^2) x^2 \end{aligned}$$

input `integrate((b*x+a)^3*(B*x+A)*(e*x+d)^2,x, algorithm="maxima")`

output `1/7*B*b^3*e^2*x^7 + A*a^3*d^2*x + 1/6*(2*B*b^3*d*e + (3*B*a*b^2 + A*b^3)*e^2)*x^6 + 1/5*(B*b^3*d^2 + 2*(3*B*a*b^2 + A*b^3)*d*e + 3*(B*a^2*b + A*a*b^2)*e^2)*x^5 + 1/4*((3*B*a*b^2 + A*b^3)*d^2 + 6*(B*a^2*b + A*a*b^2)*d*e + (B*a^3 + 3*A*a^2*b)*e^2)*x^4 + 1/3*(A*a^3*e^2 + 3*(B*a^2*b + A*a*b^2)*d^2 + 2*(B*a^3 + 3*A*a^2*b)*d*e)*x^3 + 1/2*(2*A*a^3*d*e + (B*a^3 + 3*A*a^2*b)*d^2)*x^2`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 287 vs. $2(110) = 220$.

Time = 0.12 (sec) , antiderivative size = 287, normalized size of antiderivative = 2.43

$$\int (a + bx)^3(A + Bx)(d + ex)^2 dx = \frac{1}{7} Bb^3e^2x^7 + \frac{1}{3} Bb^3dex^6 + \frac{1}{2} Bab^2e^2x^6 + \frac{1}{6} Ab^3e^2x^6$$

$$+ \frac{1}{5} Bb^3d^2x^5 + \frac{6}{5} Bab^2dex^5 + \frac{2}{5} Ab^3dex^5$$

$$+ \frac{3}{5} Ba^2be^2x^5 + \frac{3}{5} Aab^2e^2x^5 + \frac{3}{4} Bab^2d^2x^4$$

$$+ \frac{1}{4} Ab^3d^2x^4 + \frac{3}{2} Ba^2bdex^4 + \frac{3}{2} Aab^2dex^4$$

$$+ \frac{1}{4} Ba^3e^2x^4 + \frac{3}{4} Aa^2be^2x^4 + Ba^2bd^2x^3 + Aab^2d^2x^3$$

$$+ \frac{2}{3} Ba^3dex^3 + 2Aa^2bdex^3 + \frac{1}{3} Aa^3e^2x^3$$

$$+ \frac{1}{2} Ba^3d^2x^2 + \frac{3}{2} Aa^2bd^2x^2 + Aa^3dex^2 + Aa^3d^2x$$

input `integrate((b*x+a)^3*(B*x+A)*(e*x+d)^2,x, algorithm="giac")`

output `1/7*B*b^3*e^2*x^7 + 1/3*B*b^3*d*e*x^6 + 1/2*B*a*b^2*e^2*x^6 + 1/6*A*b^3*e^2*x^6 + 1/5*B*b^3*d^2*x^5 + 6/5*B*a*b^2*d*e*x^5 + 2/5*A*b^3*d*e*x^5 + 3/5*B*a^2*b*e^2*x^5 + 3/5*A*a*b^2*e^2*x^5 + 3/4*B*a*b^2*d^2*x^4 + 1/4*A*b^3*d^2*x^4 + 3/2*B*a^2*b*d*e*x^4 + 3/2*A*a*b^2*d*e*x^4 + 1/4*B*a^3*e^2*x^4 + 3/4*A*a^2*b*e^2*x^4 + B*a^2*b*d^2*x^3 + A*a*b^2*d^2*x^3 + 2/3*B*a^3*d*e*x^3 + 2*A*a^2*b*d*e*x^3 + 1/3*A*a^3*e^2*x^3 + 1/2*B*a^3*d^2*x^2 + 3/2*A*a^2*b*d^2*x^2 + A*a^3*d*e*x^2 + A*a^3*d^2*x`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 231, normalized size of antiderivative = 1.96

$$\int (a + bx)^3(A + Bx)(d + ex)^2 dx = x^4 \left(\frac{B a^3 e^2}{4} + \frac{3 B a^2 b d e}{2} + \frac{3 A a^2 b e^2}{4} + \frac{3 B a b^2 d^2}{4} \right. \\ \left. + \frac{3 A a b^2 d e}{2} + \frac{A b^3 d^2}{4} \right) \\ + x^3 \left(\frac{2 B a^3 d e}{3} + \frac{A a^3 e^2}{3} + B a^2 b d^2 + 2 A a^2 b d e \right. \\ \left. + A a b^2 d^2 \right) + x^5 \left(\frac{3 B a^2 b e^2}{5} + \frac{6 B a b^2 d e}{5} \right. \\ \left. + \frac{3 A a b^2 e^2}{5} + \frac{B b^3 d^2}{5} + \frac{2 A b^3 d e}{5} \right) \\ + A a^3 d^2 x + \frac{a^2 d x^2 (2 A a e + 3 A b d + B a d)}{2} \\ + \frac{b^2 e x^6 (A b e + 3 B a e + 2 B b d)}{6} + \frac{B b^3 e^2 x^7}{7}$$

input `int((A + B*x)*(a + b*x)^3*(d + e*x)^2,x)`output `x^4*((A*b^3*d^2)/4 + (B*a^3*e^2)/4 + (3*A*a^2*b*e^2)/4 + (3*B*a*b^2*d^2)/4 + (3*A*a*b^2*d*e)/2 + (3*B*a^2*b*d*e)/2) + x^3*((A*a^3*e^2)/3 + (2*B*a^3*d*e)/3 + A*a*b^2*d^2 + B*a^2*b*d^2 + 2*A*a^2*b*d*e) + x^5*((B*b^3*d^2)/5 + (2*A*b^3*d*e)/5 + (3*A*a*b^2*e^2)/5 + (3*B*a^2*b*e^2)/5 + (6*B*a*b^2*d*e)/5) + A*a^3*d^2*x + (a^2*d*x^2*(2*A*a*e + 3*A*b*d + B*a*d))/2 + (b^2*e*x^6*(A*b*e + 3*B*a*e + 2*B*b*d))/6 + (B*b^3*e^2*x^7)/7`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.46

$$\int (a + bx)^3(A + Bx)(d + ex)^2 dx \\ = \frac{x(15b^4e^2x^6 + 70ab^3e^2x^5 + 35b^4dex^5 + 126a^2b^2e^2x^4 + 168ab^3dex^4 + 21b^4d^2x^4 + 105a^3be^2x^3 + 315a^2b^2d^2x^3 + 105a^3be^2x^3 + 315a^2b^2d^2x^3 + 105a^3be^2x^3 + 315a^2b^2d^2x^3)}{105}$$

input `int((b*x+a)^3*(B*x+A)*(e*x+d)^2,x)`

output

```
(x*(105*a**4*d**2 + 105*a**4*d*e*x + 35*a**4*e**2*x**2 + 210*a**3*b*d**2*x
+ 280*a**3*b*d*e*x**2 + 105*a**3*b*e**2*x**3 + 210*a**2*b**2*d**2*x**2 +
315*a**2*b**2*d*e*x**3 + 126*a**2*b**2*e**2*x**4 + 105*a*b**3*d**2*x**3 +
168*a*b**3*d*e*x**4 + 70*a*b**3*e**2*x**5 + 21*b**4*d**2*x**4 + 35*b**4*d*
e*x**5 + 15*b**4*e**2*x**6))/105
```

3.29 $\int (a + bx)^3(A + Bx)(d + ex) dx$

Optimal result	320
Mathematica [A] (verified)	320
Rubi [A] (verified)	321
Maple [B] (verified)	322
Fricas [B] (verification not implemented)	323
Sympy [B] (verification not implemented)	323
Maxima [B] (verification not implemented)	324
Giac [B] (verification not implemented)	324
Mupad [B] (verification not implemented)	325
Reduce [B] (verification not implemented)	326

Optimal result

Integrand size = 18, antiderivative size = 75

$$\int (a + bx)^3(A + Bx)(d + ex) dx = \frac{(Ab - aB)(bd - ae)(a + bx)^4}{4b^3} + \frac{(bBd + Abe - 2aBe)(a + bx)^5}{5b^3} + \frac{Be(a + bx)^6}{6b^3}$$

```
output 1/4*(A*b-B*a)*(-a*e+b*d)*(b*x+a)^4/b^3+1/5*(A*b*e-2*B*a*e+B*b*d)*(b*x+a)^5/b^3+1/6*B*e*(b*x+a)^6/b^3
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.73

$$\int (a + bx)^3(A + Bx)(d + ex) dx = a^3Ax + \frac{1}{2}a^2(3Abd + aBd + aAe)x^2 + \frac{1}{3}a(3Ab(bd + ae) + aB(3bd + ae))x^3 + \frac{1}{4}b(3aB(bd + ae) + Ab(bd + 3ae))x^4 + \frac{1}{5}b^2(bBd + Abe + 3aBe)x^5 + \frac{1}{6}b^3Bex^6$$

input `Integrate[(a + b*x)^3*(A + B*x)*(d + e*x), x]`

output $a^3 A d x + (a^2 (3 A b d + a B d + a A e) x^2) / 2 + (a (3 A b (b d + a e) + a B (3 b d + a e)) x^3) / 3 + (b (3 a B (b d + a e) + A b (b d + 3 a e)) x^4) / 4 + (b^2 (b B d + A b e + 3 a B e) x^5) / 5 + (b^3 B e x^6) / 6$

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^3 (A + Bx)(d + ex) dx$$

$$\downarrow 86$$

$$\int \left(\frac{(a + bx)^4 (-2aBe + Abe + bBd)}{b^2} + \frac{(a + bx)^3 (Ab - aB)(bd - ae)}{b^2} + \frac{Be(a + bx)^5}{b^2} \right) dx$$

$$\downarrow 2009$$

$$\frac{(a + bx)^5 (-2aBe + Abe + bBd)}{5b^3} + \frac{(a + bx)^4 (Ab - aB)(bd - ae)}{4b^3} + \frac{Be(a + bx)^6}{6b^3}$$

input `Int[(a + b*x)^3*(A + B*x)*(d + e*x), x]`

output $((A b - a B) (b d - a e) (a + b x)^4) / (4 b^3) + ((b B d + A b e - 2 a B e) * (a + b x)^5) / (5 b^3) + (B e (a + b x)^6) / (6 b^3)$

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 141 vs. 2(69) = 138.

Time = 0.17 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.89

method	result
norman	$\frac{b^3 B e x^6}{6} + \left(\frac{1}{5} A b^3 e + \frac{3}{5} B a b^2 e + \frac{1}{5} b^3 B d\right) x^5 + \left(\frac{3}{4} A a b^2 e + \frac{1}{4} A b^3 d + \frac{3}{4} B a^2 b e + \frac{3}{4} B a b^2 d\right) x^4 +$
default	$\frac{b^3 B e x^6}{6} + \frac{((b^3 A + 3 a b^2 B) e + b^3 B d) x^5}{5} + \frac{((3 a b^2 A + 3 a^2 b B) e + (b^3 A + 3 a b^2 B) d) x^4}{4} + \frac{((3 a^2 b A + a^3 B) e + (3 a b^2 A + 3 a^2 b B) d) x^3}{3} +$
gosper	$\frac{1}{6} b^3 B e x^6 + \frac{1}{5} x^5 A b^3 e + \frac{3}{5} x^5 B a b^2 e + \frac{1}{5} x^5 b^3 B d + \frac{3}{4} x^4 A a b^2 e + \frac{1}{4} x^4 A b^3 d + \frac{3}{4} x^4 B a^2 b e + \frac{3}{4} x^4 B a b^2 d +$
risch	$\frac{1}{6} b^3 B e x^6 + \frac{1}{5} x^5 A b^3 e + \frac{3}{5} x^5 B a b^2 e + \frac{1}{5} x^5 b^3 B d + \frac{3}{4} x^4 A a b^2 e + \frac{1}{4} x^4 A b^3 d + \frac{3}{4} x^4 B a^2 b e + \frac{3}{4} x^4 B a b^2 d +$
parallelrisch	$\frac{1}{6} b^3 B e x^6 + \frac{1}{5} x^5 A b^3 e + \frac{3}{5} x^5 B a b^2 e + \frac{1}{5} x^5 b^3 B d + \frac{3}{4} x^4 A a b^2 e + \frac{1}{4} x^4 A b^3 d + \frac{3}{4} x^4 B a^2 b e + \frac{3}{4} x^4 B a b^2 d +$
orering	$\frac{x(10 b^3 B e x^5 + 12 A b^3 e x^4 + 36 B a b^2 e x^4 + 12 B b^3 d x^4 + 45 A a b^2 e x^3 + 15 A b^3 d x^3 + 45 B a^2 b e x^3 + 45 B a b^2 d x^3 + 60 A a^2 b e x^2 + 60 A a b^2 d x^2 + 30 A^2 b^2 e x^2 + 30 A^2 b^2 d x^2 + 30 A b^3 e x^2 + 30 A b^3 d x^2 + 30 B a^2 b e x^2 + 30 B a^2 b d x^2 + 30 B a b^2 e x^2 + 30 B a b^2 d x^2 + 30 B^2 a^2 e x^2 + 30 B^2 a^2 d x^2 + 30 B^2 a b^2 e x^2 + 30 B^2 a b^2 d x^2 + 30 B^2 b^3 e x^2 + 30 B^2 b^3 d x^2 + 30 A^2 a^2 b^2 e x^2 + 30 A^2 a^2 b^2 d x^2 + 30 A^2 a b^3 e x^2 + 30 A^2 a b^3 d x^2 + 30 A^2 b^3 e x^2 + 30 A^2 b^3 d x^2 + 30 B^2 a^2 b^2 e x^2 + 30 B^2 a^2 b^2 d x^2 + 30 B^2 a b^3 e x^2 + 30 B^2 a b^3 d x^2 + 30 B^2 b^3 e x^2 + 30 B^2 b^3 d x^2)}{60}$

```
input int((b*x+a)^3*(B*x+A)*(e*x+d),x,method=_RETURNVERBOSE)
```

```
output 1/6*b^3*B*e*x^6+(1/5*A*b^3*e+3/5*B*a*b^2*e+1/5*b^3*B*d)*x^5+(3/4*A*a*b^2*e+1/4*A*b^3*d+3/4*B*a^2*b*e+3/4*B*a*b^2*d)*x^4+(A*a^2*b*e+A*a*b^2*d+1/3*B*a^3*e+B*a^2*b*d)*x^3+(1/2*a^3*A*e+3/2*A*a^2*b*d+1/2*B*a^3*d)*x^2+a^3*A*d*x
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 146 vs. $2(69) = 138$.

Time = 0.08 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.95

$$\int (a + bx)^3(A + Bx)(d + ex) dx = \frac{1}{6} Bb^3ex^6 + Aa^3dx + \frac{1}{5} (Bb^3d + (3 Bab^2 + Ab^3)e)x^5 + \frac{1}{4} ((3 Bab^2 + Ab^3)d + 3 (Ba^2b + Aab^2)e)x^4 + \frac{1}{3} (3 (Ba^2b + Aab^2)d + (Ba^3 + 3 Aa^2b)e)x^3 + \frac{1}{2} (Aa^3e + (Ba^3 + 3 Aa^2b)d)x^2$$

input `integrate((b*x+a)^3*(B*x+A)*(e*x+d),x, algorithm="fricas")`

output `1/6*B*b^3*e*x^6 + A*a^3*d*x + 1/5*(B*b^3*d + (3*B*a*b^2 + A*b^3)*e)*x^5 + 1/4*((3*B*a*b^2 + A*b^3)*d + 3*(B*a^2*b + A*a*b^2)*e)*x^4 + 1/3*(3*(B*a^2*b + A*a*b^2)*d + (B*a^3 + 3*A*a^2*b)*e)*x^3 + 1/2*(A*a^3*e + (B*a^3 + 3*A*a^2*b)*d)*x^2`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 168 vs. $2(73) = 146$.

Time = 0.03 (sec) , antiderivative size = 168, normalized size of antiderivative = 2.24

$$\int (a + bx)^3(A + Bx)(d + ex) dx = Aa^3dx + \frac{Bb^3ex^6}{6} + x^5 \left(\frac{Ab^3e}{5} + \frac{3Bab^2e}{5} + \frac{Bb^3d}{5} \right) + x^4 \cdot \left(\frac{3Aab^2e}{4} + \frac{Ab^3d}{4} + \frac{3Ba^2be}{4} + \frac{3Bab^2d}{4} \right) + x^3 \left(Aa^2be + Aab^2d + \frac{Ba^3e}{3} + Ba^2bd \right) + x^2 \left(\frac{Aa^3e}{2} + \frac{3Aa^2bd}{2} + \frac{Ba^3d}{2} \right)$$

input `integrate((b*x+a)**3*(B*x+A)*(e*x+d),x)`

output

```
A***3*d*x + B*b**3*e*x**6/6 + x**5*(A*b**3*e/5 + 3*B*a*b**2*e/5 + B*b**3*
d/5) + x**4*(3*A*a*b**2*e/4 + A*b**3*d/4 + 3*B*a**2*b*e/4 + 3*B*a*b**2*d/4
) + x**3*(A*a**2*b*e + A*a*b**2*d + B*a**3*e/3 + B*a**2*b*d) + x**2*(A*a**
3*e/2 + 3*A*a**2*b*d/2 + B*a**3*d/2)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 146 vs. $2(69) = 138$.

Time = 0.04 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.95

$$\int (a + bx)^3(A + Bx)(d + ex) dx = \frac{1}{6} Bb^3ex^6 + Aa^3dx + \frac{1}{5} (Bb^3d + (3Bab^2 + Ab^3)e)x^5$$

$$+ \frac{1}{4} ((3Bab^2 + Ab^3)d + 3(Ba^2b + Aab^2)e)x^4$$

$$+ \frac{1}{3} (3(Ba^2b + Aab^2)d + (Ba^3 + 3Aa^2b)e)x^3$$

$$+ \frac{1}{2} (Aa^3e + (Ba^3 + 3Aa^2b)d)x^2$$

input

```
integrate((b*x+a)^3*(B*x+A)*(e*x+d),x, algorithm="maxima")
```

output

```
1/6*B*b^3*e*x^6 + A*a^3*d*x + 1/5*(B*b^3*d + (3*B*a*b^2 + A*b^3)*e)*x^5 +
1/4*((3*B*a*b^2 + A*b^3)*d + 3*(B*a^2*b + A*a*b^2)*e)*x^4 + 1/3*(3*(B*a^2*b
+ A*a*b^2)*d + (B*a^3 + 3*A*a^2*b)*e)*x^3 + 1/2*(A*a^3*e + (B*a^3 + 3*A*
a^2*b)*d)*x^2
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 163 vs. $2(69) = 138$.

Time = 0.12 (sec) , antiderivative size = 163, normalized size of antiderivative = 2.17

$$\int (a + bx)^3(A + Bx)(d + ex) dx = \frac{1}{6} Bb^3ex^6 + \frac{1}{5} Bb^3dx^5 + \frac{3}{5} Bab^2ex^5 + \frac{1}{5} Ab^3ex^5$$

$$+ \frac{3}{4} Bab^2dx^4 + \frac{1}{4} Ab^3dx^4 + \frac{3}{4} Ba^2bex^4 + \frac{3}{4} Aab^2ex^4$$

$$+ Ba^2bdx^3 + Aab^2dx^3 + \frac{1}{3} Ba^3ex^3 + Aa^2bex^3$$

$$+ \frac{1}{2} Ba^3dx^2 + \frac{3}{2} Aa^2bdx^2 + \frac{1}{2} Aa^3ex^2 + Aa^3dx$$

input `integrate((b*x+a)^3*(B*x+A)*(e*x+d),x, algorithm="giac")`

output `1/6*B*b^3*e*x^6 + 1/5*B*b^3*d*x^5 + 3/5*B*a*b^2*e*x^5 + 1/5*A*b^3*e*x^5 + 3/4*B*a*b^2*d*x^4 + 1/4*A*b^3*d*x^4 + 3/4*B*a^2*b*e*x^4 + 3/4*A*a*b^2*e*x^4 + B*a^2*b*d*x^3 + A*a*b^2*d*x^3 + 1/3*B*a^3*e*x^3 + A*a^2*b*e*x^3 + 1/2*B*a^3*d*x^2 + 3/2*A*a^2*b*d*x^2 + 1/2*A*a^3*e*x^2 + A*a^3*d*x`

Mupad [B] (verification not implemented)

Time = 0.93 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.88

$$\int (a + bx)^3 (A + Bx)(d + ex) dx = x^2 \left(\frac{Aa^3e}{2} + \frac{Ba^3d}{2} + \frac{3Aa^2bd}{2} \right) + x^5 \left(\frac{Ab^3e}{5} + \frac{Bb^3d}{5} + \frac{3Bab^2e}{5} \right) + x^3 \left(\frac{Ba^3e}{3} + Aab^2d + Aa^2be + Ba^2bd \right) + x^4 \left(\frac{Ab^3d}{4} + \frac{3Aab^2e}{4} + \frac{3Bab^2d}{4} + \frac{3Ba^2be}{4} \right) + Aa^3dx + \frac{Bb^3ex^6}{6}$$

input `int((A + B*x)*(a + b*x)^3*(d + e*x),x)`

output `x^2*((A*a^3*e)/2 + (B*a^3*d)/2 + (3*A*a^2*b*d)/2) + x^5*((A*b^3*e)/5 + (B*b^3*d)/5 + (3*B*a*b^2*e)/5) + x^3*((B*a^3*e)/3 + A*a*b^2*d + A*a^2*b*e + B*a^2*b*d) + x^4*((A*b^3*d)/4 + (3*A*a*b^2*e)/4 + (3*B*a*b^2*d)/4 + (3*B*a^2*b*e)/4) + A*a^3*d*x + (B*b^3*e*x^6)/6`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.29

$$\int (a + bx)^3 (A + Bx)(d + ex) dx$$

$$= \frac{x(5b^4e x^5 + 24ab^3e x^4 + 6b^4d x^4 + 45a^2b^2e x^3 + 30ab^3d x^3 + 40a^3be x^2 + 60a^2b^2d x^2 + 15a^4ex + 60a^3bd)}{30}$$

input `int((b*x+a)^3*(B*x+A)*(e*x+d),x)`output `(x*(30*a**4*d + 15*a**4*e*x + 60*a**3*b*d*x + 40*a**3*b*e*x**2 + 60*a**2*b**2*d*x**2 + 45*a**2*b**2*e*x**3 + 30*a*b**3*d*x**3 + 24*a*b**3*e*x**4 + 6*b**4*d*x**4 + 5*b**4*e*x**5))/30`

3.30 $\int (a + bx)^3 (A + Bx) dx$

Optimal result	327
Mathematica [A] (verified)	327
Rubi [A] (verified)	328
Maple [B] (verified)	329
Fricas [B] (verification not implemented)	329
Sympy [B] (verification not implemented)	330
Maxima [B] (verification not implemented)	330
Giac [B] (verification not implemented)	331
Mupad [B] (verification not implemented)	331
Reduce [B] (verification not implemented)	332

Optimal result

Integrand size = 13, antiderivative size = 38

$$\int (a + bx)^3 (A + Bx) dx = \frac{(Ab - aB)(a + bx)^4}{4b^2} + \frac{B(a + bx)^5}{5b^2}$$

output `1/4*(A*b-B*a)*(b*x+a)^4/b^2+1/5*B*(b*x+a)^5/b^2`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.76

$$\int (a + bx)^3 (A + Bx) dx = a^3 Ax + \frac{1}{2} a^2 (3Ab + aB) x^2 + ab(Ab + aB) x^3 + \frac{1}{4} b^2 (Ab + 3aB) x^4 + \frac{1}{5} b^3 B x^5$$

input `Integrate[(a + b*x)^3*(A + B*x),x]`

output `a^3*A*x + (a^2*(3*A*b + a*B)*x^2)/2 + a*b*(A*b + a*B)*x^3 + (b^2*(A*b + 3*a*B)*x^4)/4 + (b^3*B*x^5)/5`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^3(A + Bx) dx$$

$$\downarrow 49$$

$$\int \left(\frac{(a + bx)^3(Ab - aB)}{b} + \frac{B(a + bx)^4}{b} \right) dx$$

$$\downarrow 2009$$

$$\frac{(a + bx)^4(Ab - aB)}{4b^2} + \frac{B(a + bx)^5}{5b^2}$$

input `Int[(a + b*x)^3*(A + B*x),x]`

output `((A*b - a*B)*(a + b*x)^4)/(4*b^2) + (B*(a + b*x)^5)/(5*b^2)`

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 69 vs. $2(34) = 68$.

Time = 0.12 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.84

method	result	size
norman	$\frac{b^3 B x^5}{5} + \left(\frac{1}{4} b^3 A + \frac{3}{4} a b^2 B\right) x^4 + (a b^2 A + a^2 b B) x^3 + \left(\frac{3}{2} a^2 b A + \frac{1}{2} a^3 B\right) x^2 + a^3 A x$	70
gosper	$\frac{1}{5} b^3 B x^5 + \frac{1}{4} x^4 b^3 A + \frac{3}{4} x^4 a b^2 B + A a b^2 x^3 + B a^2 b x^3 + \frac{3}{2} x^2 a^2 b A + \frac{1}{2} x^2 a^3 B + a^3 A x$	73
default	$\frac{b^3 B x^5}{5} + \frac{(b^3 A + 3 a b^2 B) x^4}{4} + \frac{(3 a b^2 A + 3 a^2 b B) x^3}{3} + \frac{(3 a^2 b A + a^3 B) x^2}{2} + a^3 A x$	73
risch	$\frac{1}{5} b^3 B x^5 + \frac{1}{4} x^4 b^3 A + \frac{3}{4} x^4 a b^2 B + A a b^2 x^3 + B a^2 b x^3 + \frac{3}{2} x^2 a^2 b A + \frac{1}{2} x^2 a^3 B + a^3 A x$	73
parallelrisch	$\frac{1}{5} b^3 B x^5 + \frac{1}{4} x^4 b^3 A + \frac{3}{4} x^4 a b^2 B + A a b^2 x^3 + B a^2 b x^3 + \frac{3}{2} x^2 a^2 b A + \frac{1}{2} x^2 a^3 B + a^3 A x$	73
orering	$\frac{x(4B b^3 x^4 + 5A b^3 x^3 + 15B a b^2 x^3 + 20aA b^2 x^2 + 20B a^2 b x^2 + 30a^2 A b x + 10B a^3 x + 20a^3 A)}{20}$	74

input `int((b*x+a)^3*(B*x+A),x,method=_RETURNVERBOSE)`

output `1/5*b^3*B*x^5+(1/4*b^3*A+3/4*a*b^2*B)*x^4+(A*a*b^2+B*a^2*b)*x^3+(3/2*a^2*b*A+1/2*a^3*B)*x^2+a^3*A*x`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 69 vs. $2(34) = 68$.

Time = 0.06 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.82

$$\int (a + bx)^3 (A + Bx) dx = \frac{1}{5} B b^3 x^5 + A a^3 x + \frac{1}{4} (3 B a b^2 + A b^3) x^4 + (B a^2 b + A a b^2) x^3 + \frac{1}{2} (B a^3 + 3 A a^2 b) x^2$$

input `integrate((b*x+a)^3*(B*x+A),x, algorithm="fricas")`

output `1/5*B*b^3*x^5 + A*a^3*x + 1/4*(3*B*a*b^2 + A*b^3)*x^4 + (B*a^2*b + A*a*b^2)*x^3 + 1/2*(B*a^3 + 3*A*a^2*b)*x^2`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 73 vs. $2(32) = 64$.

Time = 0.03 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.92

$$\int (a + bx)^3(A + Bx) dx = Aa^3x + \frac{Bb^3x^5}{5} + x^4 \left(\frac{Ab^3}{4} + \frac{3Bab^2}{4} \right) + x^3(Aab^2 + Ba^2b) + x^2 \cdot \left(\frac{3Aa^2b}{2} + \frac{Ba^3}{2} \right)$$

input `integrate((b*x+a)**3*(B*x+A),x)`

output `A*a**3*x + B*b**3*x**5/5 + x**4*(A*b**3/4 + 3*B*a*b**2/4) + x**3*(A*a*b**2 + B*a**2*b) + x**2*(3*A*a**2*b/2 + B*a**3/2)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 69 vs. $2(34) = 68$.

Time = 0.03 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.82

$$\int (a + bx)^3(A + Bx) dx = \frac{1}{5} Bb^3x^5 + Aa^3x + \frac{1}{4} (3Bab^2 + Ab^3)x^4 + (Ba^2b + Aab^2)x^3 + \frac{1}{2} (Ba^3 + 3Aa^2b)x^2$$

input `integrate((b*x+a)^3*(B*x+A),x, algorithm="maxima")`

output `1/5*B*b^3*x^5 + A*a^3*x + 1/4*(3*B*a*b^2 + A*b^3)*x^4 + (B*a^2*b + A*a*b^2)*x^3 + 1/2*(B*a^3 + 3*A*a^2*b)*x^2`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 72 vs. $2(34) = 68$.

Time = 0.12 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.89

$$\int (a + bx)^3(A + Bx) dx = \frac{1}{5} Bb^3x^5 + \frac{3}{4} Bab^2x^4 + \frac{1}{4} Ab^3x^4 + Ba^2bx^3 \\ + Aab^2x^3 + \frac{1}{2} Ba^3x^2 + \frac{3}{2} Aa^2bx^2 + Aa^3x$$

input `integrate((b*x+a)^3*(B*x+A),x, algorithm="giac")`

output `1/5*B*b^3*x^5 + 3/4*B*a*b^2*x^4 + 1/4*A*b^3*x^4 + B*a^2*b*x^3 + A*a*b^2*x^3 + 1/2*B*a^3*x^2 + 3/2*A*a^2*b*x^2 + A*a^3*x`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.71

$$\int (a + bx)^3(A + Bx) dx = x^2 \left(\frac{B a^3}{2} + \frac{3 A b a^2}{2} \right) + x^4 \left(\frac{A b^3}{4} + \frac{3 B a b^2}{4} \right) \\ + \frac{B b^3 x^5}{5} + A a^3 x + a b x^3 (A b + B a)$$

input `int((A + B*x)*(a + b*x)^3,x)`

output `x^2*((B*a^3)/2 + (3*A*a^2*b)/2) + x^4*((A*b^3)/4 + (3*B*a*b^2)/4) + (B*b^3*x^5)/5 + A*a^3*x + a*b*x^3*(A*b + B*a)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.13

$$\int (a + bx)^3(A + Bx) dx = \frac{x(b^4x^4 + 5ab^3x^3 + 10a^2b^2x^2 + 10a^3bx + 5a^4)}{5}$$

input `int((b*x+a)^3*(B*x+A),x)`

output `(x*(5*a**4 + 10*a**3*b*x + 10*a**2*b**2*x**2 + 5*a*b**3*x**3 + b**4*x**4))
/5`

3.31 $\int \frac{(a+bx)^3(A+Bx)}{d+ex} dx$

Optimal result	333
Mathematica [A] (verified)	333
Rubi [A] (verified)	334
Maple [B] (verified)	335
Fricas [B] (verification not implemented)	336
Sympy [B] (verification not implemented)	336
Maxima [B] (verification not implemented)	337
Giac [B] (verification not implemented)	337
Mupad [B] (verification not implemented)	338
Reduce [B] (verification not implemented)	339

Optimal result

Integrand size = 20, antiderivative size = 124

$$\int \frac{(a+bx)^3(A+Bx)}{d+ex} dx = -\frac{b(bd-ae)^2(Bd-Ae)x}{e^4} + \frac{(bd-ae)(Bd-Ae)(a+bx)^2}{2e^3} - \frac{(Bd-Ae)(a+bx)^3}{3e^2} + \frac{B(a+bx)^4}{4be} + \frac{(bd-ae)^3(Bd-Ae)\log(d+ex)}{e^5}$$

output

```
-b*(-a*e+b*d)^2*(-A*e+B*d)*x/e^4+1/2*(-a*e+b*d)*(-A*e+B*d)*(b*x+a)^2/e^3-1/3*(-A*e+B*d)*(b*x+a)^3/e^2+1/4*B*(b*x+a)^4/b/e+(-a*e+b*d)^3*(-A*e+B*d)*ln(e*x+d)/e^5
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.36

$$\int \frac{(a+bx)^3(A+Bx)}{d+ex} dx = \frac{ex(12a^3Be^3 + 18a^2be^2(-2Bd + 2Ae + Bex) + 6ab^2e(3Ae(-2d + ex) + B(6d^2 - 3dex + 2e^2x^2)) + b^3(2$$

input `Integrate[((a + b*x)^3*(A + B*x))/(d + e*x),x]`

output `(e*x*(12*a^3*B*e^3 + 18*a^2*b*e^2*(-2*B*d + 2*A*e + B*e*x) + 6*a*b^2*e*(3*A*e*(-2*d + e*x) + B*(6*d^2 - 3*d*e*x + 2*e^2*x^2)) + b^3*(2*A*e*(6*d^2 - 3*d*e*x + 2*e^2*x^2) + B*(-12*d^3 + 6*d^2*e*x - 4*d*e^2*x^2 + 3*e^3*x^3))) + 12*(b*d - a*e)^3*(B*d - A*e)*Log[d + e*x])/(12*e^5)`

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^3(A + Bx)}{d + ex} dx$$

↓ 86

$$\int \left(\frac{(ae - bd)^3(Ae - Bd)}{e^4(d + ex)} + \frac{b(bd - ae)^2(Ae - Bd)}{e^4} - \frac{b(a + bx)(bd - ae)(Ae - Bd)}{e^3} + \frac{b(a + bx)^2(Ae - Bd)}{e^2} + \dots \right) dx$$

↓ 2009

$$\frac{(bd - ae)^3(Bd - Ae) \log(d + ex)}{e^5} - \frac{bx(bd - ae)^2(Bd - Ae)}{e^4} + \frac{(a + bx)^2(bd - ae)(Bd - Ae)}{2e^3} - \frac{(a + bx)^3(Bd - Ae)}{3e^2} + \frac{B(a + bx)^4}{4be}$$

input `Int[((a + b*x)^3*(A + B*x))/(d + e*x),x]`

output `-((b*(b*d - a*e)^2*(B*d - A*e)*x)/e^4) + ((b*d - a*e)*(B*d - A*e)*(a + b*x)^2)/(2*e^3) - ((B*d - A*e)*(a + b*x)^3)/(3*e^2) + (B*(a + b*x)^4)/(4*b*e) + ((b*d - a*e)^3*(B*d - A*e)*Log[d + e*x])/e^5`

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 259 vs. 2(118) = 236.

Time = 0.22 (sec) , antiderivative size = 260, normalized size of antiderivative = 2.10

method	result
norman	$\frac{(3A^2 a^2 b e^3 - 3A a b^2 d e^2 + A b^3 d^2 e + B a^3 e^3 - 3B a^2 b d e^2 + 3B a b^2 d^2 e - b^3 B d^3) x}{e^4} + \frac{b(3A a b e^2 - A b^2 d e + 3B a^2 e^2 - 3B a b d e + b^2 d^2)}{2e^3}$
default	$\frac{\frac{1}{4} b^3 B x^4 e^3 + \frac{1}{3} A b^3 e^3 x^3 + B a b^2 e^3 x^3 - \frac{1}{3} B b^3 d e^2 x^3 + \frac{3}{2} A a b^2 e^3 x^2 - \frac{1}{2} A b^3 d e^2 x^2 + \frac{3}{2} B a^2 b e^3 x^2 - \frac{3}{2} B a b^2 d e^2 x^2 + \frac{1}{2} B b^3 d^2 e x^2 + \frac{3}{4} A a^2 b e^3 - \frac{3}{4} A a b^2 d e^2 + \frac{3}{4} A b^3 d^2 e + \frac{3}{4} B a^3 e^3 - \frac{3}{4} B a^2 b d e^2 + \frac{3}{4} B a b^2 d^2 e - \frac{3}{4} B b^3 d^3}{e^4}$
risch	$\frac{b^3 B x^4}{4e} + \frac{A b^3 x^3}{3e} + \frac{B a b^2 x^3}{e} - \frac{B b^3 d x^3}{3e^2} + \frac{3A a b^2 x^2}{2e} - \frac{A b^3 d x^2}{2e^2} + \frac{3B a^2 b x^2}{2e} - \frac{3B a b^2 d x^2}{2e^2} + \frac{B b^3 d^2 x^2}{2e^3} + \frac{3A a^2 b e^3}{4} - \frac{3A a b^2 d e^2}{4} + \frac{3A b^3 d^2 e}{4} + \frac{3B a^3 e^3}{4} - \frac{3B a^2 b d e^2}{4} + \frac{3B a b^2 d^2 e}{4} - \frac{3B b^3 d^3}{4}$
parallelrisch	$\frac{12B x^3 a b^2 e^4 - 4B x^3 b^3 d e^3 + 18A x^2 a b^2 e^4 - 6A x^2 b^3 d e^3 + 36A x a^2 b e^4 + 12A x b^3 d^2 e^2 + 3B x^4 b^3 e^4 + 12A \ln(e x + d) a^3 e^4 + 12B \ln(e x + d) b^3 d^3 e^4}{e^4}$

```
input int((b*x+a)^3*(B*x+A)/(e*x+d),x,method=_RETURNVERBOSE)
```

```
output (3*A*a^2*b*e^3-3*A*a*b^2*d*e^2+A*b^3*d^2*e+B*a^3*e^3-3*B*a^2*b*d*e^2+3*B*a*b^2*d^2*e-B*b^3*d^3)/e^4*x+1/2*b/e^3*(3*A*a*b*e^2-A*b^2*d*e+3*B*a^2*e^2-3*B*a*b*d*e+B*b^2*d^2)*x^2+1/3*b^2/e^2*(A*b*e+3*B*a*e-B*b*d)*x^3+1/4*b^3*B/e*x^4+(A*a^3*e^4-3*A*a^2*b*d*e^3+3*A*a*b^2*d^2*e^2-A*b^3*d^3*e-B*a^3*d*e^3+3*B*a^2*b*d^2*e^2-3*B*a*b^2*d^3*e+B*b^3*d^4)/e^5*ln(e*x+d)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 260 vs. $2(118) = 236$.

Time = 0.10 (sec) , antiderivative size = 260, normalized size of antiderivative = 2.10

$$\int \frac{(a+bx)^3(A+Bx)}{d+ex} dx$$

$$= \frac{3Bb^3e^4x^4 - 4(Bb^3de^3 - (3Bab^2 + Ab^3)e^4)x^3 + 6(Bb^3d^2e^2 - (3Bab^2 + Ab^3)de^3 + 3(Ba^2b + Aab^2)e^4)}{e^5}$$

input `integrate((b*x+a)^3*(B*x+A)/(e*x+d),x, algorithm="fricas")`

output $\frac{1}{12}*(3*B*b^3*e^4*x^4 - 4*(B*b^3*d*e^3 - (3*B*a*b^2 + A*b^3)*e^4)*x^3 + 6*(B*b^3*d^2*e^2 - (3*B*a*b^2 + A*b^3)*d*e^3 + 3*(B*a^2*b + A*a*b^2)*e^4)*x^2 - 12*(B*b^3*d^3*e - (3*B*a*b^2 + A*b^3)*d^2*e^2 + 3*(B*a^2*b + A*a*b^2)*d*e^3 - (B*a^3 + 3*A*a^2*b)*e^4)*x + 12*(B*b^3*d^4 + A*a^3*e^4 - (3*B*a*b^2 + A*b^3)*d^3*e + 3*(B*a^2*b + A*a*b^2)*d^2*e^2 - (B*a^3 + 3*A*a^2*b)*d*e^3)*\log(e*x + d))/e^5$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 221 vs. $2(107) = 214$.

Time = 0.49 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.78

$$\int \frac{(a+bx)^3(A+Bx)}{d+ex} dx = \frac{Bb^3x^4}{4e} + x^3 \left(\frac{Ab^3}{3e} + \frac{Bab^2}{e} - \frac{Bb^3d}{3e^2} \right) + x^2 \cdot \left(\frac{3Aab^2}{2e} - \frac{Ab^3d}{2e^2} + \frac{3Ba^2b}{2e} - \frac{3Bab^2d}{2e^2} + \frac{Bb^3d^2}{2e^3} \right) + x \left(\frac{3Aa^2b}{e} - \frac{3Aab^2d}{e^2} + \frac{Ab^3d^2}{e^3} + \frac{Ba^3}{e} - \frac{3Ba^2bd}{e^2} + \frac{3Bab^2d^2}{e^3} - \frac{Bb^3d^3}{e^4} \right) - \frac{(-Ae + Bd)(ae - bd)^3 \log(d + ex)}{e^5}$$

input `integrate((b*x+a)**3*(B*x+A)/(e*x+d),x)`

output

```
B*b**3*x**4/(4*e) + x**3*(A*b**3/(3*e) + B*a*b**2/e - B*b**3*d/(3*e**2)) +
x**2*(3*A*a*b**2/(2*e) - A*b**3*d/(2*e**2) + 3*B*a**2*b/(2*e) - 3*B*a*b**
2*d/(2*e**2) + B*b**3*d**2/(2*e**3)) + x*(3*A*a**2*b/e - 3*A*a*b**2*d/e**2
+ A*b**3*d**2/e**3 + B*a**3/e - 3*B*a**2*b*d/e**2 + 3*B*a*b**2*d**2/e**3
- B*b**3*d**3/e**4) - (-A*e + B*d)*(a*e - b*d)**3*log(d + e*x)/e**5
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 258 vs. 2(118) = 236.

Time = 0.03 (sec) , antiderivative size = 258, normalized size of antiderivative = 2.08

$$\int \frac{(a+bx)^3(A+Bx)}{d+ex} dx$$

$$= \frac{3Bb^3e^3x^4 - 4(Bb^3de^2 - (3Bab^2 + Ab^3)e^3)x^3 + 6(Bb^3d^2e - (3Bab^2 + Ab^3)de^2 + 3(Ba^2b + Aab^2)e^3)x^2 - (Bb^3d^4 + Aa^3e^4 - (3Bab^2 + Ab^3)d^3e + 3(Ba^2b + Aab^2)d^2e^2 - (Ba^3 + 3Aa^2b)de^3) \log(ex+d)}{e^5}$$

input

```
integrate((b*x+a)^3*(B*x+A)/(e*x+d),x, algorithm="maxima")
```

output

```
1/12*(3*B*b^3*e^3*x^4 - 4*(B*b^3*d*e^2 - (3*B*a*b^2 + A*b^3)*e^3)*x^3 + 6*
(B*b^3*d^2*e - (3*B*a*b^2 + A*b^3)*d*e^2 + 3*(B*a^2*b + A*a*b^2)*e^3)*x^2
- 12*(B*b^3*d^3 - (3*B*a*b^2 + A*b^3)*d^2*e + 3*(B*a^2*b + A*a*b^2)*d*e^2
- (B*a^3 + 3*A*a^2*b)*e^3)*x)/e^4 + (B*b^3*d^4 + A*a^3*e^4 - (3*B*a*b^2 +
A*b^3)*d^3*e + 3*(B*a^2*b + A*a*b^2)*d^2*e^2 - (B*a^3 + 3*A*a^2*b)*d*e^3)*
log(e*x + d)/e^5
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 297 vs. 2(118) = 236.

Time = 0.12 (sec) , antiderivative size = 297, normalized size of antiderivative = 2.40

$$\int \frac{(a+bx)^3(A+Bx)}{d+ex} dx$$

$$= \frac{3Bb^3e^3x^4 - 4Bb^3de^2x^3 + 12Bab^2e^3x^3 + 4Ab^3e^3x^3 + 6Bb^3d^2ex^2 - 18Bab^2de^2x^2 - 6Ab^3de^2x^2 + 18B(Bb^3d^4 - 3Bab^2d^3e - Ab^3d^3e + 3Ba^2bd^2e^2 + 3Aab^2d^2e^2 - Ba^3de^3 - 3Aa^2bde^3 + Aa^3e^4) \log(|ex+d|)}{e^5}$$

input `integrate((b*x+a)^3*(B*x+A)/(e*x+d),x, algorithm="giac")`

output
$$\begin{aligned} & 1/12*(3*B*b^3*e^3*x^4 - 4*B*b^3*d*e^2*x^3 + 12*B*a*b^2*e^3*x^3 + 4*A*b^3*e^3*x^3 + 6*B*b^3*d^2*e*x^2 - 18*B*a*b^2*d*e^2*x^2 - 6*A*b^3*d^2*e^2*x^2 + 18*B*a^2*b*e^3*x^2 + 18*A*a*b^2*e^3*x^2 - 12*B*b^3*d^3*x + 36*B*a*b^2*d^2*e*x + 12*A*b^3*d^2*e*x - 36*B*a^2*b*d*e^2*x - 36*A*a*b^2*d*e^2*x + 12*B*a^3*e^3*x + 36*A*a^2*b*e^3*x)/e^4 + (B*b^3*d^4 - 3*B*a*b^2*d^3*e - A*b^3*d^3*e + 3*B*a^2*b*d^2*e^2 + 3*A*a*b^2*d^2*e^2 - B*a^3*d*e^3 - 3*A*a^2*b*d*e^3 + A*a^3*e^4)*\log(\text{abs}(e*x + d))/e^5 \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.95 (sec) , antiderivative size = 268, normalized size of antiderivative = 2.16

$$\begin{aligned} & \int \frac{(a+bx)^3(A+Bx)}{d+ex} dx \\ & = x \left(\frac{B a^3 + 3 A b a^2}{e} + \frac{d \left(\frac{d \left(\frac{A b^3 + 3 B a b^2}{e} - \frac{B b^3 d}{e^2} \right) - 3 a b (A b + B a)}{e} \right)}{e} \right) \\ & + x^3 \left(\frac{A b^3 + 3 B a b^2}{3 e} - \frac{B b^3 d}{3 e^2} \right) - x^2 \left(\frac{d \left(\frac{A b^3 + 3 B a b^2}{e} - \frac{B b^3 d}{e^2} \right) - 3 a b (A b + B a)}{2 e} \right) \\ & + \frac{\ln(d+ex) (-B a^3 d e^3 + A a^3 e^4 + 3 B a^2 b d^2 e^2 - 3 A a^2 b d e^3 - 3 B a b^2 d^3 e + 3 A a b^2 d^2 e^2 + B b^3 d^3)}{e^5} \\ & + \frac{B b^3 x^4}{4 e} \end{aligned}$$

input `int(((A + B*x)*(a + b*x)^3)/(d + e*x),x)`

output
$$\begin{aligned} & x*((B*a^3 + 3*A*a^2*b)/e + (d*((d*((A*b^3 + 3*B*a*b^2)/e - (B*b^3*d)/e^2))/e - (3*a*b*(A*b + B*a))/e))/e + x^3*((A*b^3 + 3*B*a*b^2)/(3*e) - (B*b^3*d)/(3*e^2)) - x^2*((d*((A*b^3 + 3*B*a*b^2)/e - (B*b^3*d)/e^2))/(2*e) - (3*a*b*(A*b + B*a))/(2*e)) + (\log(d + e*x)*(A*a^3*e^4 + B*b^3*d^4 - A*b^3*d^3*e - B*a^3*d*e^3 + 3*A*a*b^2*d^2*e^2 + 3*B*a^2*b*d^2*e^2 - 3*A*a^2*b*d*e^3 - 3*B*a*b^2*d^3*e))/e^5 + (B*b^3*x^4)/(4*e) \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.68

$$\int \frac{(a + bx)^3(A + Bx)}{d + ex} dx$$

$$= \frac{12 \log(ex + d) a^4 e^4 - 48 \log(ex + d) a^3 b d e^3 + 72 \log(ex + d) a^2 b^2 d^2 e^2 - 48 \log(ex + d) a b^3 d^3 e + 12 \log(ex + d) a^4 e^4}{12 e^5}$$

input `int((b*x+a)^3*(B*x+A)/(e*x+d),x)`output `(12*log(d + e*x)*a**4*e**4 - 48*log(d + e*x)*a**3*b*d*e**3 + 72*log(d + e*x)*a**2*b**2*d**2*e**2 - 48*log(d + e*x)*a*b**3*d**3*e + 12*log(d + e*x)*b**4*d**4 + 48*a**3*b*e**4*x - 72*a**2*b**2*d*e**3*x + 36*a**2*b**2*e**4*x**2 + 48*a*b**3*d**2*e**2*x - 24*a*b**3*d*e**3*x**2 + 16*a*b**3*e**4*x**3 - 12*b**4*d**3*e*x + 6*b**4*d**2*e**2*x**2 - 4*b**4*d*e**3*x**3 + 3*b**4*e**4*x**4)/(12*e**5)`

3.32 $\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^2} dx$

Optimal result	340
Mathematica [A] (verified)	340
Rubi [A] (verified)	341
Maple [A] (verified)	342
Fricas [B] (verification not implemented)	343
Sympy [A] (verification not implemented)	344
Maxima [A] (verification not implemented)	344
Giac [B] (verification not implemented)	345
Mupad [B] (verification not implemented)	346
Reduce [B] (verification not implemented)	346

Optimal result

Integrand size = 20, antiderivative size = 150

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^2} dx = \frac{3b(bd-ae)(2bBd-Abe-aBe)x}{e^4} - \frac{(bd-ae)^3(Bd-Ae)}{e^5(d+ex)} - \frac{b^2(4bBd-Abe-3aBe)(d+ex)^2}{2e^5} + \frac{b^3B(d+ex)^3}{3e^5} - \frac{(bd-ae)^2(4bBd-3Abe-aBe)\log(d+ex)}{e^5}$$

output

```
3*b*(-a*e+b*d)*(-A*b*e-B*a*e+2*B*b*d)*x/e^4-(-a*e+b*d)^3*(-A*e+B*d)/e^5/(e*x+d)-1/2*b^2*(-A*b*e-3*B*a*e+4*B*b*d)*(e*x+d)^2/e^5+1/3*b^3*B*(e*x+d)^3/e^5-(-a*e+b*d)^2*(-3*A*b*e-B*a*e+4*B*b*d)*ln(e*x+d)/e^5
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.63

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^2} dx = \frac{6a^3e^3(Bd-Ae) + 18a^2be^2(Ade + B(-d^2 + dex + e^2x^2)) + 9ab^2e(2Ae(-d^2 + dex + e^2x^2) + B(2d^3 - 4$$

input `Integrate[((a + b*x)^3*(A + B*x))/(d + e*x)^2,x]`

output $(6*a^3*e^3*(B*d - A*e) + 18*a^2*b*e^2*(A*d*e + B*(-d^2 + d*e*x + e^2*x^2)) + 9*a*b^2*e*(2*A*e*(-d^2 + d*e*x + e^2*x^2) + B*(2*d^3 - 4*d^2*e*x - 3*d*e^2*x^2 + e^3*x^3)) + b^3*(3*A*e*(2*d^3 - 4*d^2*e*x - 3*d*e^2*x^2 + e^3*x^3) + 2*B*(-3*d^4 + 9*d^3*e*x + 6*d^2*e^2*x^2 - 2*d*e^3*x^3 + e^4*x^4)) - 6*(b*d - a*e)^2*(4*b*B*d - 3*A*b*e - a*B*e)*(d + e*x)*\text{Log}[d + e*x])/(6*e^5*(d + e*x))$

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^2} dx$$

↓ 86

$$\int \left(\frac{b^2(d + ex)(3aBe + Abe - 4bBd)}{e^4} + \frac{(ae - bd)^2(aBe + 3Abe - 4bBd)}{e^4(d + ex)} + \frac{(ae - bd)^3(Ae - Bd)}{e^4(d + ex)^2} - \frac{3b(bd - ae)}{e^4} \right) dx$$

↓ 2009

$$\frac{b^2(d + ex)^2(-3aBe - Abe + 4bBd)}{2e^5} - \frac{(bd - ae)^3(Bd - Ae)}{e^5(d + ex)} - \frac{(bd - ae)^2 \log(d + ex)(-aBe - 3Abe + 4bBd)}{e^5} + \frac{3bx(bd - ae)(-aBe - Abe + 2bBd)}{e^4} + \frac{b^3B(d + ex)^3}{3e^5}$$

input `Int[((a + b*x)^3*(A + B*x))/(d + e*x)^2,x]`

output

```
(3*b*(b*d - a*e)*(2*b*B*d - A*b*e - a*B*e)*x)/e^4 - ((b*d - a*e)^3*(B*d - A*e))/(e^5*(d + e*x)) - (b^2*(4*b*B*d - A*b*e - 3*a*B*e)*(d + e*x)^2)/(2*e^5) + (b^3*B*(d + e*x)^3)/(3*e^5) - ((b*d - a*e)^2*(4*b*B*d - 3*A*b*e - a*B*e)*Log[d + e*x])/e^5
```

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.))^(n_.)*((e_.) + (f_.)*(x_.))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 276, normalized size of antiderivative = 1.84

method	result
norman	$\frac{(a^3 A e^4 - 3 A a^2 b d e^3 + 6 A a b^2 d^2 e^2 - 3 A b^3 d^3 e - B a^3 d e^3 + 6 B a^2 b d^2 e^2 - 9 B a b^2 d^3 e + 4 b^3 B d^4) x}{e^4 d} + \frac{b(6 A a b e^2 - 3 A b^2 d e + 6 B a^2 e^2 - 9 B a b d e + 3 B b^2 d^2 e)}{2 e^3 (e x + d)}$
default	$\frac{b(\frac{1}{3} b^2 B x^3 e^2 + \frac{1}{2} A b^2 e^2 x^2 + \frac{3}{2} B a b e^2 x^2 - B b^2 d e x^2 + 3 A a b e^2 x - 2 A b^2 d e x + 3 B a^2 e^2 x - 6 B a b d e x + 3 b^2 B d^2 x)}{e^4} - \frac{a^3 A e^4 - 3 A a^2 b d e^3 + 6 A a b^2 d^2 e^2 - 3 A b^3 d^3 e - B a^3 d e^3 + 6 B a^2 b d^2 e^2 - 9 B a b^2 d^3 e + 4 b^3 B d^4}{e^4}$
risch	$\frac{b^3 B x^3}{3 e^2} + \frac{b^3 A x^2}{2 e^2} + \frac{3 b^2 B a x^2}{2 e^2} - \frac{b^3 B d x^2}{e^3} + \frac{3 b^2 A a x}{e^2} - \frac{2 b^3 A d x}{e^3} + \frac{3 b B a^2 x}{e^2} - \frac{6 b^2 B a d x}{e^3} + \frac{3 b^3 B d^2 x}{e^4} - \frac{a^3 A}{e(e x + d)}$
parallelrisch	$\frac{-24 b^3 B d^4 + 6 B a^3 d e^3 + 18 A b^3 d^3 e + 54 B a b^2 d^3 e - 36 A a b^2 d^2 e^2 - 36 B a^2 b d^2 e^2 + 18 A a^2 b d e^3 + 9 B x^3 a b^2 e^4 - 4 B x^3 b^3 d e^3 + 18 A a^3 A e^4 - 3 A a^2 b d e^3 + 6 A a b^2 d^2 e^2 - 3 A b^3 d^3 e - B a^3 d e^3 + 6 B a^2 b d^2 e^2 - 9 B a b^2 d^3 e + 4 b^3 B d^4}{e^4}$

input

```
int((b*x+a)^3*(B*x+A)/(e*x+d)^2,x,method=_RETURNVERBOSE)
```

output

```
((A*a^3*e^4-3*A*a^2*b*d*e^3+6*A*a*b^2*d^2*e^2-3*A*b^3*d^3*e-B*a^3*d*e^3+6*
B*a^2*b*d^2*e^2-9*B*a*b^2*d^3*e+4*B*b^3*d^4)/e^4/d*x+1/2*b*(6*A*a*b*e^2-3*
A*b^2*d*e+6*B*a^2*e^2-9*B*a*b*d*e+4*B*b^2*d^2)/e^3*x^2+1/6*b^2*(3*A*b*e+9*
B*a*e-4*B*b*d)/e^2*x^3+1/3*b^3*B/e*x^4)/(e*x+d)+1/e^5*(3*A*a^2*b*e^3-6*A*a
*b^2*d*e^2+3*A*b^3*d^2*e+B*a^3*e^3-6*B*a^2*b*d*e^2+9*B*a*b^2*d^2*e-4*B*b^3
*d^3)*ln(e*x+d)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 396 vs. $2(146) = 292$.

Time = 0.07 (sec) , antiderivative size = 396, normalized size of antiderivative = 2.64

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^2} dx$$

$$= \frac{2Bb^3e^4x^4 - 6Bb^3d^4 - 6Aa^3e^4 + 6(3Bab^2 + Ab^3)d^3e - 18(Ba^2b + Aab^2)d^2e^2 + 6(Ba^3 + 3Aa^2b)de^3 - \dots}{\dots}$$

input

```
integrate((b*x+a)^3*(B*x+A)/(e*x+d)^2,x, algorithm="fricas")
```

output

```
1/6*(2*B*b^3*e^4*x^4 - 6*B*b^3*d^4 - 6*A*a^3*e^4 + 6*(3*B*a*b^2 + A*b^3)*d
^3*e - 18*(B*a^2*b + A*a*b^2)*d^2*e^2 + 6*(B*a^3 + 3*A*a^2*b)*d*e^3 - (4*B
*b^3*d*e^3 - 3*(3*B*a*b^2 + A*b^3)*e^4)*x^3 + 3*(4*B*b^3*d^2*e^2 - 3*(3*B*
a*b^2 + A*b^3)*d*e^3 + 6*(B*a^2*b + A*a*b^2)*e^4)*x^2 + 6*(3*B*b^3*d^3*e -
2*(3*B*a*b^2 + A*b^3)*d^2*e^2 + 3*(B*a^2*b + A*a*b^2)*d*e^3)*x - 6*(4*B*b
^3*d^4 - 3*(3*B*a*b^2 + A*b^3)*d^3*e + 6*(B*a^2*b + A*a*b^2)*d^2*e^2 - (B*
a^3 + 3*A*a^2*b)*d*e^3 + (4*B*b^3*d^3*e - 3*(3*B*a*b^2 + A*b^3)*d^2*e^2 +
6*(B*a^2*b + A*a*b^2)*d*e^3 - (B*a^3 + 3*A*a^2*b)*e^4)*x)*log(e*x + d))/(e
^6*x + d*e^5)
```

Sympy [A] (verification not implemented)

Time = 0.75 (sec) , antiderivative size = 257, normalized size of antiderivative = 1.71

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^2} dx = \frac{Bb^3x^3}{3e^2} + x^2 \left(\frac{Ab^3}{2e^2} + \frac{3Bab^2}{2e^2} - \frac{Bb^3d}{e^3} \right) + x \left(\frac{3Aab^2}{e^2} - \frac{2Ab^3d}{e^3} + \frac{3Ba^2b}{e^2} - \frac{6Bab^2d}{e^3} + \frac{3Bb^3d^2}{e^4} \right) + \frac{-Aa^3e^4 + 3Aa^2bde^3 - 3Aab^2d^2e^2 + Ab^3d^3e + Ba^3de^3 - 3Ba^2bd^2e^2 + 3Bab^2d^3e - Bb^3d^4}{de^5 + e^6x} + \frac{(ae - bd)^2 \cdot (3Abe + Bae - 4Bbd) \log(d + ex)}{e^5}$$

input `integrate((b*x+a)**3*(B*x+A)/(e*x+d)**2,x)`output `B*b**3*x**3/(3*e**2) + x**2*(A*b**3/(2*e**2) + 3*B*a*b**2/(2*e**2) - B*b**3*d/e**3) + x*(3*A*a*b**2/e**2 - 2*A*b**3*d/e**3 + 3*B*a**2*b/e**2 - 6*B*a*b**2*d/e**3 + 3*B*b**3*d**2/e**4) + (-A*a**3*e**4 + 3*A*a**2*b*d*e**3 - 3*A*a*b**2*d**2*e**2 + A*b**3*d**3*e + B*a**3*d*e**3 - 3*B*a**2*b*d**2*e**2 + 3*B*a*b**2*d**3*e - B*b**3*d**4)/(d*e**5 + e**6*x) + (a*e - b*d)**2*(3*A*b*e + B*a*e - 4*B*b*d)*log(d + e*x)/e**5`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 267, normalized size of antiderivative = 1.78

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^2} dx = -\frac{Bb^3d^4 + Aa^3e^4 - (3Bab^2 + Ab^3)d^3e + 3(Ba^2b + Aab^2)d^2e^2 - (Ba^3 + 3Aa^2b)de^3}{e^6x + de^5} + \frac{2Bb^3e^2x^3 - 3(2Bb^3de - (3Bab^2 + Ab^3)e^2)x^2 + 6(3Bb^3d^2 - 2(3Bab^2 + Ab^3)de + 3(Ba^2b + Aab^2)e^3)}{6e^4} - \frac{(4Bb^3d^3 - 3(3Bab^2 + Ab^3)d^2e + 6(Ba^2b + Aab^2)de^2 - (Ba^3 + 3Aa^2b)e^3) \log(ex + d)}{e^5}$$

input `integrate((b*x+a)^3*(B*x+A)/(e*x+d)^2,x, algorithm="maxima")`

output

```

-(B*b^3*d^4 + A*a^3*e^4 - (3*B*a*b^2 + A*b^3)*d^3*e + 3*(B*a^2*b + A*a*b^2)
)*d^2*e^2 - (B*a^3 + 3*A*a^2*b)*d*e^3)/(e^6*x + d*e^5) + 1/6*(2*B*b^3*e^2*
x^3 - 3*(2*B*b^3*d*e - (3*B*a*b^2 + A*b^3)*e^2)*x^2 + 6*(3*B*b^3*d^2 - 2*(
3*B*a*b^2 + A*b^3)*d*e + 3*(B*a^2*b + A*a*b^2)*e^2)*x)/e^4 - (4*B*b^3*d^3
- 3*(3*B*a*b^2 + A*b^3)*d^2*e + 6*(B*a^2*b + A*a*b^2)*d*e^2 - (B*a^3 + 3*A
*a^2*b)*e^3)*log(e*x + d)/e^5

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 372 vs. $2(146) = 292$.

Time = 0.12 (sec) , antiderivative size = 372, normalized size of antiderivative = 2.48

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^2} dx$$

$$= \frac{\left(2Bb^3 - \frac{3(4Bb^3de - 3Bab^2e^2 - Ab^3e^2)}{(ex+d)e} + \frac{18(2Bb^3d^2e^2 - 3Bab^2de^3 - Ab^3de^3 + Ba^2be^4 + Aab^2e^4)}{(ex+d)^2e^2}\right)(ex+d)^3}{6e^5}$$

$$+ \frac{(4Bb^3d^3 - 9Bab^2d^2e - 3Ab^3d^2e + 6Ba^2bde^2 + 6Aab^2de^2 - Ba^3e^3 - 3Aa^2be^3) \log\left(\frac{|ex+d|}{(ex+d)^2|e|}\right)}{e^5}$$

$$- \frac{\frac{Bb^3d^4e^3}{ex+d} - \frac{3Bab^2d^3e^4}{ex+d} - \frac{Ab^3d^3e^4}{ex+d} + \frac{3Ba^2bd^2e^5}{ex+d} + \frac{3Aab^2d^2e^5}{ex+d} - \frac{Ba^3de^6}{ex+d} - \frac{3Aa^2bde^6}{ex+d} + \frac{Aa^3e^7}{ex+d}}{e^8}$$

input

```
integrate((b*x+a)^3*(B*x+A)/(e*x+d)^2,x, algorithm="giac")
```

output

```

1/6*(2*B*b^3 - 3*(4*B*b^3*d*e - 3*B*a*b^2*e^2 - A*b^3*e^2)/((e*x + d)*e) +
18*(2*B*b^3*d^2*e^2 - 3*B*a*b^2*d*e^3 - A*b^3*d*e^3 + B*a^2*b*e^4 + A*a*b
^2*e^4)/((e*x + d)^2*e^2))*(e*x + d)^3/e^5 + (4*B*b^3*d^3 - 9*B*a*b^2*d^2*
e - 3*A*b^3*d^2*e + 6*B*a^2*b*d*e^2 + 6*A*a*b^2*d*e^2 - B*a^3*e^3 - 3*A*a
^2*b*e^3)*log(abs(e*x + d)/((e*x + d)^2*abs(e)))/e^5 - (B*b^3*d^4*e^3/(e*x
+ d) - 3*B*a*b^2*d^3*e^4/(e*x + d) - A*b^3*d^3*e^4/(e*x + d) + 3*B*a^2*b*d
^2*e^5/(e*x + d) + 3*A*a*b^2*d^2*e^5/(e*x + d) - B*a^3*d*e^6/(e*x + d) - 3
*A*a^2*b*d*e^6/(e*x + d) + A*a^3*e^7/(e*x + d))/e^8

```

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 293, normalized size of antiderivative = 1.95

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^2} dx = x^2 \left(\frac{Ab^3 + 3Bab^2}{2e^2} - \frac{Bb^3d}{e^3} \right) - x \left(\frac{2d \left(\frac{Ab^3 + 3Bab^2}{e^2} - \frac{2Bb^3d}{e^3} \right)}{e} - \frac{3ab(Ab+Ba)}{e^2} + \frac{Bb^3d^2}{e^4} \right) + \frac{\ln(d+ex) (Ba^3e^3 - 6Ba^2bde^2 + 3Aa^2be^3 + 9Bab^2d^2e - 6Aab^2de^2 - 4Bb^3d^3 + 3Ab^3d^2e) - Ba^3de^3 + Aa^3e^4 + 3Ba^2bd^2e^2 - 3Aa^2bde^3 - 3Bab^2d^3e + 3Aab^2d^2e^2 + Bb^3d^4 - Ab^3d^3e}{e(xe^5 + de^4)} + \frac{Bb^3x^3}{3e^2}$$

input `int(((A + B*x)*(a + b*x)^3)/(d + e*x)^2,x)`output `x^2*((A*b^3 + 3*B*a*b^2)/(2*e^2) - (B*b^3*d)/e^3) - x*((2*d*((A*b^3 + 3*B*a*b^2)/e^2 - (2*B*b^3*d)/e^3))/e - (3*a*b*(A*b + B*a))/e^2 + (B*b^3*d^2)/e^4) + (log(d + e*x)*(B*a^3*e^3 - 4*B*b^3*d^3 + 3*A*a^2*b*e^3 + 3*A*b^3*d^2*e - 6*A*a*b^2*d*e^2 + 9*B*a*b^2*d^2*e - 6*B*a^2*b*d*e^2))/e^5 - (A*a^3*e^4 + B*b^3*d^4 - A*b^3*d^3*e - B*a^3*d*e^3 + 3*A*a*b^2*d^2*e^2 + 3*B*a^2*b*d^2*e^2 - 3*A*a^2*b*d*e^3 - 3*B*a*b^2*d^3*e)/(e*(d*e^4 + e^5*x)) + (B*b^3*x^3)/(3*e^2)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 297, normalized size of antiderivative = 1.98

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^2} dx = \frac{12 \log(ex+d) a^3 b d^2 e^3 + 12 \log(ex+d) a^3 b d e^4 x - 36 \log(ex+d) a^2 b^2 d^3 e^2 - 36 \log(ex+d) a^2 b^2 d^2 e^3 x + \dots}{\dots}$$

input `int((b*x+a)^3*(B*x+A)/(e*x+d)^2,x)`

output

```
(12*log(d + e*x)*a**3*b*d**2*e**3 + 12*log(d + e*x)*a**3*b*d*e**4*x - 36*log(d + e*x)*a**2*b**2*d**3*e**2 - 36*log(d + e*x)*a**2*b**2*d**2*e**3*x + 36*log(d + e*x)*a*b**3*d**4*e + 36*log(d + e*x)*a*b**3*d**3*e**2*x - 12*log(d + e*x)*b**4*d**5 - 12*log(d + e*x)*b**4*d**4*e*x + 3*a**4*e**5*x - 12*a**3*b*d*e**4*x + 36*a**2*b**2*d**2*e**3*x + 18*a**2*b**2*d*e**4*x**2 - 36*a*b**3*d**3*e**2*x - 18*a*b**3*d**2*e**3*x**2 + 6*a*b**3*d*e**4*x**3 + 12*b**4*d**4*e*x + 6*b**4*d**3*e**2*x**2 - 2*b**4*d**2*e**3*x**3 + b**4*d*e**4*x**4)/(3*d*e**5*(d + e*x))
```


3.33 $\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^3} dx$

Optimal result	348
Mathematica [A] (verified)	348
Rubi [A] (verified)	349
Maple [A] (verified)	350
Fricas [B] (verification not implemented)	351
Sympy [B] (verification not implemented)	352
Maxima [A] (verification not implemented)	352
Giac [B] (verification not implemented)	353
Mupad [B] (verification not implemented)	354
Reduce [B] (verification not implemented)	354

Optimal result

Integrand size = 20, antiderivative size = 145

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^3} dx = -\frac{b^2(3bBd - Abe - 3aBe)x}{e^4} + \frac{b^3 Bx^2}{2e^3} - \frac{(bd - ae)^3(Bd - Ae)}{2e^5(d+ex)^2} + \frac{(bd - ae)^2(4bBd - 3Abe - aBe)}{e^5(d+ex)} + \frac{3b(bd - ae)(2bBd - Abe - aBe) \log(d+ex)}{e^5}$$

```
output -b^2*(-A*b*e-3*B*a*e+3*B*b*d)*x/e^4+1/2*b^3*B*x^2/e^3-1/2*(-a*e+b*d)^3*(-A
*e+B*d)/e^5/(e*x+d)^2+(-a*e+b*d)^2*(-3*A*b*e-B*a*e+4*B*b*d)/e^5/(e*x+d)+3*
b*(-a*e+b*d)*(-A*b*e-B*a*e+2*B*b*d)*ln(e*x+d)/e^5
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.64

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^3} dx = \frac{-a^3e^3(Ae + B(d + 2ex)) - 3a^2be^2(Ae(d + 2ex) - Bd(3d + 4ex)) + 3ab^2e(Ade(3d + 4ex) + B(-5d^3 -$$

input `Integrate[((a + b*x)^3*(A + B*x))/(d + e*x)^3,x]`

output
$$\begin{aligned} & (-a^3e^3(Ae + B(d + 2ex)) - 3a^2b^2e^2(Ae(d + 2ex) - Bd(3d + 4ex)) + 3a^2b^2e(A^2d^2e(3d + 4ex) + B(-5d^3 - 4d^2ex + 4d^2e^2x^2 + 2e^3x^3)) + b^3(Ae(-5d^3 - 4d^2ex + 4d^2e^2x^2 + 2e^3x^3) + B(7d^4 + 2d^3ex - 11d^2e^2x^2 - 4de^3x^3 + e^4x^4)) + 6b(bd - ae)(2bBd - A^2be - aB^2e)(d + ex)^2 \text{Log}[d + ex]) / (2e^5(d + ex)^2) \end{aligned}$$

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^3} dx$$

↓ 86

$$\int \left(\frac{b^2(3aBe + Abe - 3bBd)}{e^4} - \frac{3b(bd - ae)(aBe + Abe - 2bBd)}{e^4(d + ex)} + \frac{(ae - bd)^2(aBe + 3Abe - 4bBd)}{e^4(d + ex)^2} + \frac{(ae - bd)(aBe + 3Abe - 4bBd)}{e^4(d + ex)^3} \right) dx$$

↓ 2009

$$\begin{aligned} & -\frac{b^2x(-3aBe - Abe + 3bBd)}{e^4} + \frac{(bd - ae)^2(-aBe - 3Abe + 4bBd)}{e^5(d + ex)} - \frac{(bd - ae)^3(Bd - Ae)}{2e^5(d + ex)^2} + \\ & \frac{3b(bd - ae) \log(d + ex)(-aBe - Abe + 2bBd)}{e^5} + \frac{b^3Bx^2}{2e^3} \end{aligned}$$

input `Int[((a + b*x)^3*(A + B*x))/(d + e*x)^3,x]`

output

```

-((b^2*(3*b*B*d - A*b*e - 3*a*B*e)*x)/e^4) + (b^3*B*x^2)/(2*e^3) - ((b*d -
a*e)^3*(B*d - A*e))/(2*e^5*(d + e*x)^2) + ((b*d - a*e)^2*(4*b*B*d - 3*A*b
*e - a*B*e))/(e^5*(d + e*x)) + (3*b*(b*d - a*e)*(2*b*B*d - A*b*e - a*B*e)*
Log[d + e*x])/e^5
    
```

Defintions of rubi rules used

rule 86

```

Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && ( !IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
    
```

rule 2009

```

Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
    
```

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 267, normalized size of antiderivative = 1.84

method	result
default	$\frac{b^2(\frac{1}{2}Bbe x^2 + A b e x + 3 B a e x - 3 B b d x)}{e^4} - \frac{3 A a^2 b e^3 - 6 A a b^2 d e^2 + 3 A b^3 d^2 e + B a^3 e^3 - 6 B a^2 b d e^2 + 9 B a b^2 d^2 e - 4 b^3 B d^3}{e^5(e x + d)} - \frac{a^3 A e^4 + 3 A a^2 b d e^3 - 9 A a b^2 d^2 e^2 + 9 A b^3 d^3 e + B a^3 d e^3 - 9 B a^2 b d^2 e^2 + 27 B a b^2 d^3 e - 18 b^3 B d^4}{2 e^5} - \frac{(3 A a^2 b e^3 - 6 A a b^2 d e^2 + 3 A b^3 d^2 e - B a^3 e^3 + 6 B a^2 b d e^2 - 9 B a b^2 d^2 e + 4 b^3 B d^3)}{(e x + d)^2}$
norman	$\frac{b^2(A b e + 3 B a e - 2 B b d) x^3}{e^2} - \frac{a^3 A e^4 + 3 A a^2 b d e^3 - 9 A a b^2 d^2 e^2 + 9 A b^3 d^3 e + B a^3 d e^3 - 9 B a^2 b d^2 e^2 + 27 B a b^2 d^3 e - 18 b^3 B d^4}{2 e^5} - \frac{(3 A a^2 b e^3 - 6 A a b^2 d e^2 + 3 A b^3 d^2 e - B a^3 e^3 + 6 B a^2 b d e^2 - 9 B a b^2 d^2 e + 4 b^3 B d^3)}{(e x + d)^2}$
risch	$\frac{b^3 B x^2}{2 e^3} + \frac{b^3 A x}{e^3} + \frac{3 b^2 B a x}{e^3} - \frac{3 b^3 B d x}{e^4} + \frac{(-3 A a^2 b e^3 + 6 A a b^2 d e^2 - 3 A b^3 d^2 e - B a^3 e^3 + 6 B a^2 b d e^2 - 9 B a b^2 d^2 e + 4 b^3 B d^3)}{e^5}$
parallelrisch	$\frac{18 b^3 B d^4 - 18 B \ln(e x + d) x^2 a b^2 d e^3 - B a^3 d e^3 - 9 A b^3 d^3 e - 27 B a b^2 d^3 e + 9 A a b^2 d^2 e^2 + 9 B a^2 b d^2 e^2 - 3 A a^2 b d e^3 + 6 B x^3 a b^2 e^4}{e^5}$

input

```

int((b*x+a)^3*(B*x+A)/(e*x+d)^3,x,method=_RETURNVERBOSE)
    
```

output

```

b^2/e^4*(1/2*B*b*e*x^2+A*b*e*x+3*B*a*e*x-3*B*b*d*x)-1/e^5*(3*A*a^2*b*e^3-6
*A*a*b^2*d*e^2+3*A*b^3*d^2*e+B*a^3*e^3-6*B*a^2*b*d*e^2+9*B*a*b^2*d^2*e-4*B
*b^3*d^3)/(e*x+d)-1/2*(A*a^3*e^4-3*A*a^2*b*d*e^3+3*A*a*b^2*d^2*e^2-A*b^3*d
^3*e-B*a^3*d*e^3+3*B*a^2*b*d^2*e^2-3*B*a*b^2*d^3*e+B*b^3*d^4)/e^5/(e*x+d)^
2+3*b/e^5*(A*a*b*e^2-A*b^2*d*e+B*a^2*e^2-3*B*a*b*d*e+2*B*b^2*d^2)*ln(e*x+d
)

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 420 vs. $2(141) = 282$.

Time = 0.07 (sec) , antiderivative size = 420, normalized size of antiderivative = 2.90

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^3} dx$$

$$= \frac{Bb^3e^4x^4 + 7Bb^3d^4 - Aa^3e^4 - 5(3Bab^2 + Ab^3)d^3e + 9(Ba^2b + Aab^2)d^2e^2 - (Ba^3 + 3Aa^2b)de^3 - 2(2Aa^2b + Aab^2)d^2e^2 - (Ba^3 + 3Aa^2b)d^3e - 2(2Bb^3d^4 - (3Bab^2 + Ab^3)e^4)x^3 - (11Bb^3d^2e^2 - 4(3Bab^2 + Ab^3)d^2e^3)x^2 + 2(Bb^3d^3e - 2(3Bab^2 + Ab^3)d^2e^2 + 6(Ba^2b + Aab^2)d^2e^3 - (Ba^3 + 3Aa^2b)e^4)x + 6(2Bb^3d^4 - (3Bab^2 + Ab^3)d^3e + (Ba^2b + Aab^2)d^2e^2 + (2Bb^3d^2e^2 - (3Bab^2 + Ab^3)d^2e^3 + (Ba^2b + Aab^2)e^4)x^2 + 2(2Bb^3d^3e - (3Bab^2 + Ab^3)d^2e^2 + (Ba^2b + Aab^2)d^2e^3)x) \log(ex + d)}{(e^7x^2 + 2de^6x + d^2e^5)}$$

input

```

integrate((b*x+a)^3*(B*x+A)/(e*x+d)^3,x, algorithm="fricas")

```

output

```

1/2*(B*b^3*e^4*x^4 + 7*B*b^3*d^4 - A*a^3*e^4 - 5*(3*B*a*b^2 + A*b^3)*d^3*e
+ 9*(B*a^2*b + A*a*b^2)*d^2*e^2 - (B*a^3 + 3*A*a^2*b)*d*e^3 - 2*(2*B*b^3*d
^4 - (3*B*a*b^2 + A*b^3)*e^4)*x^3 - (11*B*b^3*d^2*e^2 - 4*(3*B*a*b^2 +
A*b^3)*d^2*e^3)*x^2 + 2*(B*b^3*d^3*e - 2*(3*B*a*b^2 + A*b^3)*d^2*e^2 + 6*(B*
a^2*b + A*a*b^2)*d^2*e^3 - (B*a^3 + 3*A*a^2*b)*e^4)*x + 6*(2*B*b^3*d^4 - (3*
B*a*b^2 + A*b^3)*d^3*e + (B*a^2*b + A*a*b^2)*d^2*e^2 + (2*B*b^3*d^2*e^2 -
(3*B*a*b^2 + A*b^3)*d^2*e^3 + (B*a^2*b + A*a*b^2)*e^4)*x^2 + 2*(2*B*b^3*d^3*
e - (3*B*a*b^2 + A*b^3)*d^2*e^2 + (B*a^2*b + A*a*b^2)*d^2*e^3)*x)*log(e*x +
d))/(e^7*x^2 + 2*d*e^6*x + d^2*e^5)

```


output

```
1/2*(7*B*b^3*d^4 - A*a^3*e^4 - 5*(3*B*a*b^2 + A*b^3)*d^3*e + 9*(B*a^2*b +
A*a*b^2)*d^2*e^2 - (B*a^3 + 3*A*a^2*b)*d*e^3 + 2*(4*B*b^3*d^3*e - 3*(3*B*a
*b^2 + A*b^3)*d^2*e^2 + 6*(B*a^2*b + A*a*b^2)*d*e^3 - (B*a^3 + 3*A*a^2*b)*
e^4)*x)/(e^7*x^2 + 2*d*e^6*x + d^2*e^5) + 1/2*(B*b^3*e*x^2 - 2*(3*B*b^3*d
- (3*B*a*b^2 + A*b^3)*e)*x)/e^4 + 3*(2*B*b^3*d^2 - (3*B*a*b^2 + A*b^3)*d*e
+ (B*a^2*b + A*a*b^2)*e^2)*log(e*x + d)/e^5
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 286 vs. $2(141) = 282$.

Time = 0.12 (sec) , antiderivative size = 286, normalized size of antiderivative = 1.97

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^3} dx$$

$$= \frac{3(2Bb^3d^2 - 3Bab^2de - Ab^3de + Ba^2be^2 + Aab^2e^2) \log(|ex+d|)}{e^5}$$

$$+ \frac{Bb^3e^3x^2 - 6Bb^3de^2x + 6Bab^2e^3x + 2Ab^3e^3x}{2e^6}$$

$$+ \frac{7Bb^3d^4 - 15Bab^2d^3e - 5Ab^3d^3e + 9Ba^2bd^2e^2 + 9Aab^2d^2e^2 - Ba^3de^3 - 3Aa^2bde^3 - Aa^3e^4 + 2(4Bb^3d^2 - 3Bab^2de - Ab^3de + Ba^2be^2 + Aab^2e^2)x}{2(ex+d)^2e^5}$$

input

```
integrate((b*x+a)^3*(B*x+A)/(e*x+d)^3,x, algorithm="giac")
```

output

```
3*(2*B*b^3*d^2 - 3*B*a*b^2*d*e - A*b^3*d*e + B*a^2*b*e^2 + A*a*b^2*e^2)*lo
g(abs(e*x + d))/e^5 + 1/2*(B*b^3*e^3*x^2 - 6*B*b^3*d*e^2*x + 6*B*a*b^2*e^3
*x + 2*A*b^3*e^3*x)/e^6 + 1/2*(7*B*b^3*d^4 - 15*B*a*b^2*d^3*e - 5*A*b^3*d^
3*e + 9*B*a^2*b*d^2*e^2 + 9*A*a*b^2*d^2*e^2 - B*a^3*d*e^3 - 3*A*a^2*b*d*e^
3 - A*a^3*e^4 + 2*(4*B*b^3*d^3*e - 9*B*a*b^2*d^2*e^2 - 3*A*b^3*d^2*e^2 + 6
*B*a^2*b*d*e^3 + 6*A*a*b^2*d*e^3 - B*a^3*e^4 - 3*A*a^2*b*e^4)*x)/((e*x + d
)^2*e^5)
```

Mupad [B] (verification not implemented)

Time = 0.97 (sec) , antiderivative size = 290, normalized size of antiderivative = 2.00

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^3} dx = x \left(\frac{Ab^3 + 3Bab^2}{e^3} - \frac{3Bb^3d}{e^4} \right) - \frac{Ba^3de^3 + Aa^3e^4 - 9Ba^2bd^2e^2 + 3Aa^2bde^3 + 15Bab^2d^3e - 9Aab^2d^2e^2 - 7Bb^3d^4 + 5Ab^3d^3e}{2e} + x \frac{(Ba^3e^3 - 6Ba^2bde^2 + d^2e^4 + 2de^5x + e^6x^2)}{d^2e^4 + 2de^5x + e^6x^2} + \frac{\ln(d+ex)(3Ba^2be^2 - 9Bab^2de + 3Aab^2e^2 + 6Bb^3d^2 - 3Ab^3de)}{e^5} + \frac{Bb^3x^2}{2e^3}$$

input `int(((A + B*x)*(a + b*x)^3)/(d + e*x)^3,x)`output `x*((A*b^3 + 3*B*a*b^2)/e^3 - (3*B*b^3*d)/e^4) - ((A*a^3*e^4 - 7*B*b^3*d^4 + 5*A*b^3*d^3*e + B*a^3*d*e^3 - 9*A*a*b^2*d^2*e^2 - 9*B*a^2*b*d^2*e^2 + 3*A*a^2*b*d*e^3 + 15*B*a*b^2*d^3*e)/(2*e) + x*(B*a^3*e^3 - 4*B*b^3*d^3 + 3*A*a^2*b*e^3 + 3*A*b^3*d^2*e - 6*A*a*b^2*d*e^2 + 9*B*a*b^2*d^2*e - 6*B*a^2*b*d*e^2))/(d^2*e^4 + e^6*x^2 + 2*d*e^5*x) + (log(d + e*x)*(6*B*b^3*d^2 - 3*A*b^3*d*e + 3*A*a*b^2*e^2 + 3*B*a^2*b*e^2 - 9*B*a*b^2*d*e))/e^5 + (B*b^3*x^2)/(2*e^3)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 330, normalized size of antiderivative = 2.28

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^3} dx = \frac{12 \log(ex+d) a^2 b^2 d^3 e^2 + 24 \log(ex+d) a^2 b^2 d^2 e^3 x + 12 \log(ex+d) a^2 b^2 d e^4 x^2 - 24 \log(ex+d) a b^3 d^4 e - \dots}{(d+ex)^3}$$

input `int((b*x+a)^3*(B*x+A)/(e*x+d)^3,x)`

output

```
(12*log(d + e*x)*a**2*b**2*d**3*e**2 + 24*log(d + e*x)*a**2*b**2*d**2*e**3
*x + 12*log(d + e*x)*a**2*b**2*d**4*x**2 - 24*log(d + e*x)*a*b**3*d**4*e
- 48*log(d + e*x)*a*b**3*d**3*e**2*x - 24*log(d + e*x)*a*b**3*d**2*e**3*x
**2 + 12*log(d + e*x)*b**4*d**5 + 24*log(d + e*x)*b**4*d**4*e*x + 12*log(d
+ e*x)*b**4*d**3*e**2*x**2 - a**4*d*e**4 + 4*a**3*b*e**5*x**2 + 6*a**2*b*
*2*d**3*e**2 - 12*a**2*b**2*d**4*x**2 - 12*a*b**3*d**4*e + 24*a*b**3*d**
2*e**3*x**2 + 8*a*b**3*d**4*x**3 + 6*b**4*d**5 - 12*b**4*d**3*e**2*x**2
- 4*b**4*d**2*e**3*x**3 + b**4*d**4*x**4)/(2*d**5*(d**2 + 2*d*e*x + e
**2*x**2))
```


3.34 $\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^4} dx$

Optimal result	356
Mathematica [A] (verified)	357
Rubi [A] (verified)	357
Maple [A] (verified)	358
Fricas [B] (verification not implemented)	359
Sympy [B] (verification not implemented)	360
Maxima [A] (verification not implemented)	360
Giac [A] (verification not implemented)	361
Mupad [B] (verification not implemented)	361
Reduce [B] (verification not implemented)	362

Optimal result

Integrand size = 20, antiderivative size = 149

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^4} dx = \frac{b^3 Bx}{e^4} - \frac{(bd-ae)^3(Bd-Ae)}{3e^5(d+ex)^3} + \frac{(bd-ae)^2(4bBd-3Abe-aBe)}{2e^5(d+ex)^2} - \frac{3b(bd-ae)(2bBd-Abe-aBe)}{e^5(d+ex)} - \frac{b^2(4bBd-Abe-3aBe)\log(d+ex)}{e^5}$$

output

```
b^3*B*x/e^4-1/3*(-a*e+b*d)^3*(-A*e+B*d)/e^5/(e*x+d)^3+1/2*(-a*e+b*d)^2*(-3
*A*b*e-B*a*e+4*B*b*d)/e^5/(e*x+d)^2-3*b*(-a*e+b*d)*(-A*b*e-B*a*e+2*B*b*d)/
e^5/(e*x+d)-b^2*(-A*b*e-3*B*a*e+4*B*b*d)*ln(e*x+d)/e^5
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 232, normalized size of antiderivative = 1.56

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^4} dx$$

$$= \frac{-a^3e^3(2Ae + B(d + 3ex)) - 3a^2be^2(Ae(d + 3ex) + 2B(d^2 + 3dex + 3e^2x^2)) + 3ab^2e(-2Ae(d^2 + 3dex -$$

input

```
Integrate[((a + b*x)^3*(A + B*x))/(d + e*x)^4,x]
```

output

```
(-(a^3*e^3*(2*A*e + B*(d + 3*e*x))) - 3*a^2*b*e^2*(A*e*(d + 3*e*x) + 2*B*(d^2 + 3*d*e*x + 3*e^2*x^2)) + 3*a*b^2*e*(-2*A*e*(d^2 + 3*d*e*x + 3*e^2*x^2) + B*d*(11*d^2 + 27*d*e*x + 18*e^2*x^2)) + b^3*(A*d*e*(11*d^2 + 27*d*e*x + 18*e^2*x^2) - 2*B*(13*d^4 + 27*d^3*e*x + 9*d^2*e^2*x^2 - 9*d*e^3*x^3 - 3*e^4*x^4)) - 6*b^2*(4*b*B*d - A*b*e - 3*a*B*e)*(d + e*x)^3*Log[d + e*x])/(6*e^5*(d + e*x)^3)
```

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^4} dx$$

↓ 86

$$\int \left(\frac{b^2(3aBe + Abe - 4bBd)}{e^4(d + ex)} - \frac{3b(bd - ae)(aBe + Abe - 2bBd)}{e^4(d + ex)^2} + \frac{(ae - bd)^2(aBe + 3Abe - 4bBd)}{e^4(d + ex)^3} + \frac{(ae - bd)(aBe + 3Abe - 4bBd)}{e^4(d + ex)^4} \right) dx$$

↓ 2009

$$-\frac{b^2 \log(d+ex)(-3aBe - Abe + 4bBd)}{e^5} - \frac{3b(bd - ae)(-aBe - Abe + 2bBd)}{e^5(d+ex)} + \frac{(bd - ae)^2(-aBe - 3Abe + 4bBd)}{2e^5(d+ex)^2} - \frac{(bd - ae)^3(Bd - Ae)}{3e^5(d+ex)^3} + \frac{b^3 Bx}{e^4}$$

input `Int[((a + b*x)^3*(A + B*x))/(d + e*x)^4,x]`

output $(b^3 B x)/e^4 - ((b*d - a*e)^3*(B*d - A*e))/(3*e^5*(d + e*x)^3) + ((b*d - a*e)^2*(4*b*B*d - 3*A*b*e - a*B*e))/(2*e^5*(d + e*x)^2) - (3*b*(b*d - a*e)*(2*b*B*d - A*b*e - a*B*e))/(e^5*(d + e*x)) - (b^2*(4*b*B*d - A*b*e - 3*a*B*e)*\text{Log}[d + e*x])/e^5$

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.))^(n_.)*((e_.) + (f_.)*(x_.))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 271, normalized size of antiderivative = 1.82

method	result
norman	$\frac{b^3 B x^4}{e} - \frac{2a^3 A e^4 + 3A a^2 b d e^3 + 6A a b^2 d^2 e^2 - 11A b^3 d^3 e + B a^3 d e^3 + 6B a^2 b d^2 e^2 - 33B a b^2 d^3 e + 44b^3 B d^4}{6e^5} - \frac{3(A a b^2 e^2 - A b^3 d e + B a^2 b e^2 - (e x + d)^3)}{e^3}$
default	$\frac{b^3 B x}{e^4} - \frac{3b(A a b e^2 - A b^2 d e + B a^2 e^2 - 3B a b d e + 2b^2 B d^2)}{e^5(e x + d)} - \frac{3A a^2 b e^3 - 6A a b^2 d e^2 + 3A b^3 d^2 e + B a^3 e^3 - 6B a^2 b d e^2 + 9B a b^2 d^2 e - 3B b^3 d^3}{2e^5(e x + d)^2}$
risch	$\frac{b^3 B x}{e^4} + \frac{(-3A a b^2 e^3 + 3A b^3 d e^2 - 3B a^2 b e^3 + 9B a b^2 d e^2 - 6b^3 B d^2 e)x^2 + (-\frac{3}{2}A a^2 b e^3 - 3A a b^2 d e^2 + \frac{9}{2}A b^3 d^2 e - \frac{1}{2}B a^3 e^3 - 3B a^2 b d e^2 + 3B a b^2 d^2 e - \frac{3}{2}B b^3 d^3)}{e^5(e x + d)}$
parallelrisc	$\frac{18B \ln(e x + d)x^3 a b^2 e^4 - 24B \ln(e x + d)x^3 b^3 d e^3 - 44b^3 B d^4 + 54B \ln(e x + d)x^2 a b^2 d e^3 - B a^3 d e^3 + 11A b^3 d^3 e + 33B a b^2 d^3 e - 6A a^2 b d^3 e^2 - 3A a b^2 d^3 e^2 + 3A b^3 d^3 e^2 - 3A b^3 d^3 e^2}{e^5(e x + d)}$

input `int((b*x+a)^3*(B*x+A)/(e*x+d)^4,x,method=_RETURNVERBOSE)`

output
$$\frac{(b^3 B / e^{x^4} - 1/6 * (2 * A * a^3 * e^4 + 3 * A * a^2 * b * d * e^3 + 6 * A * a * b^2 * d^2 * e^2 - 11 * A * b^3 * d^3 * e + B * a^3 * d * e^3 + 6 * B * a^2 * b * d^2 * e^2 - 33 * B * a * b^2 * d^3 * e + 44 * B * b^3 * d^4)) / e^5 - 3 * (A * a * b^2 * e^2 - A * b^3 * d * e + B * a^2 * b * e^2 - 3 * B * a * b^2 * d * e + 4 * B * b^3 * d^2)) / e^3 * x^2 - 1/2 * (3 * A * a^2 * b * e^3 + 6 * A * a * b^2 * d * e^2 - 9 * A * b^3 * d^2 * e + B * a^3 * e^3 + 6 * B * a^2 * b * d * e^2 - 27 * B * a * b^2 * d^2 * e + 36 * B * b^3 * d^3) / e^4 * x}{(e * x + d)^3 + b^2 / e^5 * (A * b * e + 3 * B * a * e - 4 * B * b * d)} * \ln(e * x + d)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 406 vs. $2(145) = 290$.

Time = 0.08 (sec) , antiderivative size = 406, normalized size of antiderivative = 2.72

$$\int \frac{(a + bx)^3 (A + Bx)}{(d + ex)^4} dx$$

$$= \frac{6 B b^3 e^4 x^4 + 18 B b^3 d e^3 x^3 - 26 B b^3 d^4 - 2 A a^3 e^4 + 11 (3 B a b^2 + A b^3) d^3 e - 6 (B a^2 b + A a b^2) d^2 e^2 - (B a^3 + 3 A a^2 b) d * e^3 - 18 (B b^3 d^2 * e^2 - (3 B a * b^2 + A * b^3) * d * e^3 + (B a^2 * b + A * a * b^2) * e^4) * x^2 - 3 * (18 B b^3 * d^3 * e - 9 * (3 B a * b^2 + A * b^3) * d^2 * e^2 + 6 * (B a^2 * b + A * a * b^2) * d * e^3 + (B a^3 + 3 A a^2 * b) * e^4) * x - 6 * (4 B b^3 * d^4 - (3 B a * b^2 + A * b^3) * d^3 * e + (4 B b^3 * d * e^3 - (3 B a * b^2 + A * b^3) * e^4) * x^3 + 3 * (4 B b^3 * d^2 * e^2 - (3 B a * b^2 + A * b^3) * d * e^3) * x^2 + 3 * (4 B b^3 * d^3 * e - (3 B a * b^2 + A * b^3) * d^2 * e^2) * x) * \log(e * x + d)}{(e^8 * x^3 + 3 * d * e^7 * x^2 + 3 * d^2 * e^6 * x + d^3 * e^5)}$$

input `integrate((b*x+a)^3*(B*x+A)/(e*x+d)^4,x, algorithm="fricas")`

output
$$\frac{1}{6} * (6 * B * b^3 * e^4 * x^4 + 18 * B * b^3 * d * e^3 * x^3 - 26 * B * b^3 * d^4 - 2 * A * a^3 * e^4 + 11 * (3 * B * a * b^2 + A * b^3) * d^3 * e - 6 * (B * a^2 * b + A * a * b^2) * d^2 * e^2 - (B * a^3 + 3 * A * a^2 * b) * d * e^3 - 18 * (B * b^3 * d^2 * e^2 - (3 * B * a * b^2 + A * b^3) * d * e^3 + (B * a^2 * b + A * a * b^2) * e^4) * x^2 - 3 * (18 * B * b^3 * d^3 * e - 9 * (3 * B * a * b^2 + A * b^3) * d^2 * e^2 + 6 * (B * a^2 * b + A * a * b^2) * d * e^3 + (B * a^3 + 3 * A * a^2 * b) * e^4) * x - 6 * (4 * B * b^3 * d^4 - (3 * B * a * b^2 + A * b^3) * d^3 * e + (4 * B * b^3 * d * e^3 - (3 * B * a * b^2 + A * b^3) * e^4) * x^3 + 3 * (4 * B * b^3 * d^2 * e^2 - (3 * B * a * b^2 + A * b^3) * d * e^3) * x^2 + 3 * (4 * B * b^3 * d^3 * e - (3 * B * a * b^2 + A * b^3) * d^2 * e^2) * x) * \log(e * x + d)) / (e^8 * x^3 + 3 * d * e^7 * x^2 + 3 * d^2 * e^6 * x + d^3 * e^5)$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 337 vs. $2(143) = 286$.

Time = 5.58 (sec) , antiderivative size = 337, normalized size of antiderivative = 2.26

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^4} dx = \frac{Bb^3x}{e^4} + \frac{b^2(Abe + 3Bae - 4Bbd) \log(d+ex)}{e^5} + \frac{-2Aa^3e^4 - 3Aa^2bde^3 - 6Aab^2d^2e^2 + 11Ab^3d^3e - Ba^3de^3 - 6Ba^2bd^2e^2 + 33Bab^2d^3e - 26Bb^3d^4 + x^2}{e^5}$$

input `integrate((b*x+a)**3*(B*x+A)/(e*x+d)**4,x)`

output `B*b**3*x/e**4 + b**2*(A*b*e + 3*B*a*e - 4*B*b*d)*log(d + e*x)/e**5 + (-2*A*a**3*e**4 - 3*A*a**2*b*d*e**3 - 6*A*a*b**2*d**2*e**2 + 11*A*b**3*d**3*e - B*a**3*d*e**3 - 6*B*a**2*b*d**2*e**2 + 33*B*a*b**2*d**3*e - 26*B*b**3*d**4 + x**2*(-18*A*a*b**2*e**4 + 18*A*b**3*d*e**3 - 18*B*a**2*b*e**4 + 54*B*a*b**2*d*e**3 - 36*B*b**3*d**2*e**2) + x*(-9*A*a**2*b*e**4 - 18*A*a*b**2*d*e**3 + 27*A*b**3*d**2*e**2 - 3*B*a**3*e**4 - 18*B*a**2*b*d*e**3 + 81*B*a*b**2*d**2*e**2 - 60*B*b**3*d**3*e)) / (6*d**3*e**5 + 18*d**2*e**6*x + 18*d*e**7*x**2 + 6*e**8*x**3)`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 284, normalized size of antiderivative = 1.91

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^4} dx = \frac{Bb^3x}{e^4} - \frac{26Bb^3d^4 + 2Aa^3e^4 - 11(3Bab^2 + Ab^3)d^3e + 6(Ba^2b + Aab^2)d^2e^2 + (Ba^3 + 3Aa^2b)de^3 + 18(2Bb^3d^4 + x^2(4Bb^3d - (3Bab^2 + Ab^3)e) \log(ex+d))}{e^5}$$

input `integrate((b*x+a)^3*(B*x+A)/(e*x+d)^4,x, algorithm="maxima")`

output

```
B*b^3*x/e^4 - 1/6*(26*B*b^3*d^4 + 2*A*a^3*e^4 - 11*(3*B*a*b^2 + A*b^3)*d^3
*e + 6*(B*a^2*b + A*a*b^2)*d^2*e^2 + (B*a^3 + 3*A*a^2*b)*d*e^3 + 18*(2*B*b
^3*d^2*e^2 - (3*B*a*b^2 + A*b^3)*d*e^3 + (B*a^2*b + A*a*b^2)*e^4)*x^2 + 3*
(20*B*b^3*d^3*e - 9*(3*B*a*b^2 + A*b^3)*d^2*e^2 + 6*(B*a^2*b + A*a*b^2)*d*
e^3 + (B*a^3 + 3*A*a^2*b)*e^4)*x)/(e^8*x^3 + 3*d*e^7*x^2 + 3*d^2*e^6*x + d
^3*e^5) - (4*B*b^3*d - (3*B*a*b^2 + A*b^3)*e)*log(e*x + d)/e^5
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 279, normalized size of antiderivative = 1.87

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^4} dx = \frac{Bb^3x}{e^4} - \frac{(4Bb^3d - 3Bab^2e - Ab^3e) \log(|ex+d|)}{e^5} - \frac{26Bb^3d^4 - 33Bab^2d^3e - 11Ab^3d^3e + 6Ba^2bd^2e^2 + 6Aab^2d^2e^2 + Ba^3de^3 + 3Aa^2bde^3 + 2Aa^3e^4 + 18(Ba^2b^2d^2e^2 - (3Bab^2d^2e^2 + Aab^2d^2e^2) * x^2 + 3(20Bb^3d^3e - 9(3Bab^2d^2e^2 + 6(Ba^2bd^2e^2 + Aab^2d^2e^2) * d * e^3 + (Ba^3 + 3Aa^2b) * e^4) * x) / (e^8 * x^3 + 3 * d * e^7 * x^2 + 3 * d^2 * e^6 * x + d^3 * e^5) - (4 * B * b^3 * d - (3 * B * a * b^2 + A * b^3) * e) * \log(e * x + d) / e^5}{e^5}$$

input

```
integrate((b*x+a)^3*(B*x+A)/(e*x+d)^4,x, algorithm="giac")
```

output

```
B*b^3*x/e^4 - (4*B*b^3*d - 3*B*a*b^2*e - A*b^3*e)*log(abs(e*x + d))/e^5 -
1/6*(26*B*b^3*d^4 - 33*B*a*b^2*d^3*e - 11*A*b^3*d^3*e + 6*B*a^2*b*d^2*e^2
+ 6*A*a*b^2*d^2*e^2 + B*a^3*d*e^3 + 3*A*a^2*b*d*e^3 + 2*A*a^3*e^4 + 18*(2*
B*b^3*d^2*e^2 - 3*B*a*b^2*d*e^3 - A*b^3*d*e^3 + B*a^2*b*e^4 + A*a*b^2*e^4)
*x^2 + 3*(20*B*b^3*d^3*e - 27*B*a*b^2*d^2*e^2 - 9*A*b^3*d^2*e^2 + 6*B*a^2*
b*d*e^3 + 6*A*a*b^2*d*e^3 + B*a^3*e^4 + 3*A*a^2*b*e^4)*x)/((e*x + d)^3*e^5
)
```

Mupad [B] (verification not implemented)

Time = 0.97 (sec) , antiderivative size = 301, normalized size of antiderivative = 2.02

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^4} dx = \frac{\ln(d+ex)(Ab^3e - 4Bb^3d + 3Bab^2e)}{e^5} - \frac{Ba^3de^3 + 2Aa^3e^4 + 6Ba^2bd^2e^2 + 3Aa^2bde^3 - 33Bab^2d^3e + 6Aab^2d^2e^2 + 26Bb^3d^4 - 11Ab^3d^3e}{6e} + x \left(\frac{Ba^3e^3}{2} + 3Ba^2bde^2 \right) + \frac{Bb^3x}{e^4}$$

input `int(((A + B*x)*(a + b*x)^3)/(d + e*x)^4,x)`

output
$$\begin{aligned} & (\log(d + e*x)*(A*b^3*e - 4*B*b^3*d + 3*B*a*b^2*e))/e^5 - ((2*A*a^3*e^4 + 2 \\ & 6*B*b^3*d^4 - 11*A*b^3*d^3*e + B*a^3*d*e^3 + 6*A*a*b^2*d^2*e^2 + 6*B*a^2*b \\ & *d^2*e^2 + 3*A*a^2*b*d*e^3 - 33*B*a*b^2*d^3*e)/(6*e) + x*((B*a^3*e^3)/2 + \\ & 10*B*b^3*d^3 + (3*A*a^2*b*e^3)/2 - (9*A*b^3*d^2*e)/2 + 3*A*a*b^2*d*e^2 - (\\ & 27*B*a*b^2*d^2*e)/2 + 3*B*a^2*b*d*e^2) + x^2*(3*A*a*b^2*e^3 + 3*B*a^2*b*e^ \\ & 3 - 3*A*b^3*d*e^2 + 6*B*b^3*d^2*e - 9*B*a*b^2*d*e^2))/(d^3*e^4 + e^7*x^3 + \\ & 3*d^2*e^5*x + 3*d*e^6*x^2) + (B*b^3*x)/e^4 \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 309, normalized size of antiderivative = 2.07

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^4} dx$$

$$= \frac{12 \log(ex + d) a b^3 d^4 e + 36 \log(ex + d) a b^3 d^3 e^2 x + 36 \log(ex + d) a b^3 d^2 e^3 x^2 + 12 \log(ex + d) a b^3 d e^4 x^3}{(d + ex)^4}$$

input `int((b*x+a)^3*(B*x+A)/(e*x+d)^4,x)`

output
$$\begin{aligned} & (12*\log(d + e*x)*a*b**3*d**4*e + 36*\log(d + e*x)*a*b**3*d**3*e**2*x + 36* \\ & \log(d + e*x)*a*b**3*d**2*e**3*x**2 + 12*\log(d + e*x)*a*b**3*d*e**4*x**3 - 1 \\ & 2*\log(d + e*x)*b**4*d**5 - 36*\log(d + e*x)*b**4*d**4*e*x - 36*\log(d + e*x) \\ & *b**4*d**3*e**2*x**2 - 12*\log(d + e*x)*b**4*d**2*e**3*x**3 - a**4*d*e**4 - \\ & 2*a**3*b*d**2*e**3 - 6*a**3*b*d*e**4*x + 6*a**2*b**2*e**5*x**3 + 10*a*b** \\ & 3*d**4*e + 18*a*b**3*d**3*e**2*x - 12*a*b**3*d*e**4*x**3 - 10*b**4*d**5 - \\ & 18*b**4*d**4*e*x + 12*b**4*d**2*e**3*x**3 + 3*b**4*d*e**4*x**4)/(3*d*e**5* \\ & (d**3 + 3*d**2*e*x + 3*d*e**2*x**2 + e**3*x**3)) \end{aligned}$$

output

$$\begin{aligned} & (- (a^3 e^3 (3Ae + B(d + 4ex))) - 3a^2 b e^2 (Ae(d + 4ex) + B(d^2 + 4d^2 ex + 6e^2 x^2)) - 3a b^2 e (Ae(d^2 + 4d^2 ex + 6e^2 x^2) + 3 \\ & * B(d^3 + 4d^2 ex + 6d^2 ex^2 + 4e^3 x^3)) + b^3 (-3Ae(d^3 + 4d^2 ex + 6d^2 ex^2 + 4e^3 x^3) + B d (25d^3 + 88d^2 ex + 108d^2 ex^2 + 48e^3 x^3)) + 12b^3 B (d + ex)^4 \text{Log}[d + ex]) / (12e^5 (d + ex)^4) \end{aligned}$$

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.02, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {87, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + bx)^3 (A + Bx)}{(d + ex)^5} dx \\ & \quad \downarrow 87 \\ & \frac{B \int \frac{(a+bx)^3 dx}{e} - \frac{(a + bx)^4 (Bd - Ae)}{4e(d + ex)^4 (bd - ae)}}{e} \\ & \quad \downarrow 49 \\ & \frac{B \int \left(\frac{b^3}{e^3(d+ex)} - \frac{3(bd-ae)b^2}{e^3(d+ex)^2} + \frac{3(bd-ae)^2 b}{e^3(d+ex)^3} + \frac{(ae-bd)^3}{e^3(d+ex)^4} \right) dx - \frac{(a + bx)^4 (Bd - Ae)}{4e(d + ex)^4 (bd - ae)}}{e} \\ & \quad \downarrow 2009 \\ & \frac{B \left(\frac{3b^2 (bd-ae)}{e^4(d+ex)} - \frac{3b(bd-ae)^2}{2e^4(d+ex)^2} + \frac{(bd-ae)^3}{3e^4(d+ex)^3} + \frac{b^3 \log(d+ex)}{e^4} \right) - \frac{(a + bx)^4 (Bd - Ae)}{4e(d + ex)^4 (bd - ae)}}{e} \end{aligned}$$

input

$$\text{Int}[(a + b*x)^3*(A + B*x)/(d + e*x)^5, x]$$

output

$$\begin{aligned} & -1/4*((B*d - A*e)*(a + b*x)^4)/(e*(b*d - a*e)*(d + e*x)^4) + (B*((b*d - a*e)^3/(3*e^4*(d + e*x)^3) - (3*b*(b*d - a*e)^2)/(2*e^4*(d + e*x)^2) + (3*b^2*(b*d - a*e))/(e^4*(d + e*x)) + (b^3*Log[d + e*x])/e^4))/e \end{aligned}$$

Defintions of rubi rules used

```
rule 49 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(- (b*e - a*f))*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e)), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 266 vs. 2(123) = 246.

Time = 0.22 (sec) , antiderivative size = 267, normalized size of antiderivative = 2.07

method	result
risch	$\frac{-\frac{b^2(Abe+3Bae-4Bbd)x^3}{e^2} - \frac{3b(Aab e^2 + A b^2 de + B a^2 e^2 + 3Babde - 6b^2 B d^2)x^2}{2e^3} - \frac{(3A a^2 b e^3 + 3A a b^2 d e^2 + 3A b^3 d^2 e + B a^3 e^3 + 3B a^2 b d e^2)}{3e^4}}{(ex+d)^4}$
norman	$\frac{-\frac{3a^3 A e^4 + 3A a^2 b d e^3 + 3A a b^2 d^2 e^2 + 3A b^3 d^3 e + B a^3 d e^3 + 3B a^2 b d^2 e^2 + 9B a b^2 d^3 e - 25b^3 B d^4}{12e^5} - \frac{(A b^3 e + 3B a b^2 e - 4b^3 B d)x^3}{e^2} - \frac{3(A a b^2 e^2)}{(ex+d)^4}}{e^2}$
default	$\frac{-\frac{b^2(Abe+3Bae-4Bbd)}{e^5(ex+d)} - \frac{3b(Aab e^2 - A b^2 de + B a^2 e^2 - 3Babde + 2b^2 B d^2)}{2e^5(ex+d)^2} - \frac{a^3 A e^4 - 3A a^2 b d e^3 + 3A a b^2 d^2 e^2 - A b^3 d^3 e}{4e^5}}{e^5}$
parallelrisc	$-\frac{48B \ln(ex+d)x^3 b^3 d e^3 - 25b^3 B d^4 + B a^3 d e^3 + 3A b^3 d^3 e + 9B a b^2 d^3 e + 3A a b^2 d^2 e^2 + 3B a^2 b d^2 e^2 + 3A a^2 b d e^3 + 36B x^3 a b^2 d e^2}{e^5}$

```
input int((b*x+a)^3*(B*x+A)/(e*x+d)^5,x,method=_RETURNVERBOSE)
```

output

```
(-b^2*(A*b*e+3*B*a*e-4*B*b*d)/e^2*x^3-3/2*b*(A*a*b*e^2+A*b^2*d*e+B*a^2*e^2
+3*B*a*b*d*e-6*B*b^2*d^2)/e^3*x^2-1/3*(3*A*a^2*b*e^3+3*A*a*b^2*d*e^2+3*A*b
^3*d^2*e+B*a^3*e^3+3*B*a^2*b*d*e^2+9*B*a*b^2*d^2*e-22*B*b^3*d^3)/e^4*x-1/1
2*(3*A*a^3*e^4+3*A*a^2*b*d*e^3+3*A*a*b^2*d^2*e^2+3*A*b^3*d^3*e+B*a^3*d*e^3
+3*B*a^2*b*d^2*e^2+9*B*a*b^2*d^3*e-25*B*b^3*d^4)/e^5)/(e*x+d)^4+b^3*B*ln(e
*x+d)/e^5
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 354 vs. $2(123) = 246$.

Time = 0.08 (sec) , antiderivative size = 354, normalized size of antiderivative = 2.74

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^5} dx$$

$$= \frac{25Bb^3d^4 - 3Aa^3e^4 - 3(3Bab^2 + Ab^3)d^3e - 3(Ba^2b + Aab^2)d^2e^2 - (Ba^3 + 3Aa^2b)de^3 + 12(4Bb^3de^3}{e^5}$$

input

```
integrate((b*x+a)^3*(B*x+A)/(e*x+d)^5,x, algorithm="fricas")
```

output

```
1/12*(25*B*b^3*d^4 - 3*A*a^3*e^4 - 3*(3*B*a*b^2 + A*b^3)*d^3*e - 3*(B*a^2*
b + A*a*b^2)*d^2*e^2 - (B*a^3 + 3*A*a^2*b)*d*e^3 + 12*(4*B*b^3*d*e^3 - (3*
B*a*b^2 + A*b^3)*e^4)*x^3 + 18*(6*B*b^3*d^2*e^2 - (3*B*a*b^2 + A*b^3)*d*e^
3 - (B*a^2*b + A*a*b^2)*e^4)*x^2 + 4*(22*B*b^3*d^3*e - 3*(3*B*a*b^2 + A*b^
3)*d^2*e^2 - 3*(B*a^2*b + A*a*b^2)*d*e^3 - (B*a^3 + 3*A*a^2*b)*e^4)*x + 12
*(B*b^3*e^4*x^4 + 4*B*b^3*d*e^3*x^3 + 6*B*b^3*d^2*e^2*x^2 + 4*B*b^3*d^3*e*
x + B*b^3*d^4)*log(e*x + d))/(e^9*x^4 + 4*d*e^8*x^3 + 6*d^2*e^7*x^2 + 4*d^
3*e^6*x + d^4*e^5)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 359 vs. $2(116) = 232$.

Time = 13.09 (sec) , antiderivative size = 359, normalized size of antiderivative = 2.78

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^5} dx = \frac{Bb^3 \log(d+ex)}{e^5} + \frac{-3Aa^3e^4 - 3Aa^2bde^3 - 3Aab^2d^2e^2 - 3Ab^3d^3e - Ba^3de^3 - 3Ba^2bd^2e^2 - 9Bab^2d^3e + 25Bb^3d^4 + x^3(-$$

input `integrate((b*x+a)**3*(B*x+A)/(e*x+d)**5,x)`

output `B*b**3*log(d + e*x)/e**5 + (-3*A*a**3*e**4 - 3*A*a**2*b*d*e**3 - 3*A*a*b**2*d**2*e**2 - 3*A*b**3*d**3*e - B*a**3*d*e**3 - 3*B*a**2*b*d**2*e**2 - 9*B*a*b**2*d**3*e + 25*B*b**3*d**4 + x**3*(-12*A*b**3*e**4 - 36*B*a*b**2*e**4 + 48*B*b**3*d*e**3) + x**2*(-18*A*a*b**2*e**4 - 18*A*b**3*d*e**3 - 18*B*a**2*b*e**4 - 54*B*a*b**2*d*e**3 + 108*B*b**3*d**2*e**2) + x*(-12*A*a**2*b*e**4 - 12*A*a*b**2*d*e**3 - 12*A*b**3*d**2*e**2 - 4*B*a**3*e**4 - 12*B*a**2*b*d*e**3 - 36*B*a*b**2*d**2*e**2 + 88*B*b**3*d**3*e))/(12*d**4*e**5 + 48*d**3*e**6*x + 72*d**2*e**7*x**2 + 48*d*e**8*x**3 + 12*e**9*x**4)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 302 vs. $2(123) = 246$.

Time = 0.05 (sec) , antiderivative size = 302, normalized size of antiderivative = 2.34

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^5} dx = \frac{25Bb^3d^4 - 3Aa^3e^4 - 3(3Bab^2 + Ab^3)d^3e - 3(Ba^2b + Aab^2)d^2e^2 - (Ba^3 + 3Aa^2b)de^3 + 12(4Bb^3de^3 + Bb^3 \log(ex+d))}{e^5}$$

input `integrate((b*x+a)^3*(B*x+A)/(e*x+d)^5,x, algorithm="maxima")`

output

```
1/12*(25*B*b^3*d^4 - 3*A*a^3*e^4 - 3*(3*B*a*b^2 + A*b^3)*d^3*e - 3*(B*a^2*b + A*a*b^2)*d^2*e^2 - (B*a^3 + 3*A*a^2*b)*d*e^3 + 12*(4*B*b^3*d*e^3 - (3*B*a*b^2 + A*b^3)*e^4)*x^3 + 18*(6*B*b^3*d^2*e^2 - (3*B*a*b^2 + A*b^3)*d*e^3 - (B*a^2*b + A*a*b^2)*e^4)*x^2 + 4*(22*B*b^3*d^3*e - 3*(3*B*a*b^2 + A*b^3)*d^2*e^2 - 3*(B*a^2*b + A*a*b^2)*d*e^3 - (B*a^3 + 3*A*a^2*b)*e^4)*x)/(e^9*x^4 + 4*d*e^8*x^3 + 6*d^2*e^7*x^2 + 4*d^3*e^6*x + d^4*e^5) + B*b^3*log(e*x + d)/e^5
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 449 vs. $2(123) = 246$.

Time = 0.13 (sec) , antiderivative size = 449, normalized size of antiderivative = 3.48

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^5} dx = -\frac{Bb^3 \log\left(\frac{|ex+d|}{(ex+d)^2|e|}\right)}{e^5} + \frac{48Bb^3de^{15}}{ex+d} - \frac{36Bb^3d^2e^{15}}{(ex+d)^2} + \frac{16Bb^3d^3e^{15}}{(ex+d)^3} - \frac{3Bb^3d^4e^{15}}{(ex+d)^4} - \frac{36Bab^2e^{16}}{ex+d} - \frac{12Ab^3e^{16}}{ex+d} + \frac{54Bab^2de^{16}}{(ex+d)^2} + \frac{18Ab^3de^{16}}{(ex+d)^2} - \frac{36Bab^2d}{(ex+d)}$$

input

```
integrate((b*x+a)^3*(B*x+A)/(e*x+d)^5,x, algorithm="giac")
```

output

```
-B*b^3*log(abs(e*x + d)/((e*x + d)^2*abs(e)))/e^5 + 1/12*(48*B*b^3*d*e^15/(e*x + d) - 36*B*b^3*d^2*e^15/(e*x + d)^2 + 16*B*b^3*d^3*e^15/(e*x + d)^3 - 3*B*b^3*d^4*e^15/(e*x + d)^4 - 36*B*a*b^2*e^16/(e*x + d) - 12*A*b^3*e^16/(e*x + d) + 54*B*a*b^2*d*e^16/(e*x + d)^2 + 18*A*b^3*d*e^16/(e*x + d)^2 - 36*B*a*b^2*d^2*e^16/(e*x + d)^3 - 12*A*b^3*d^2*e^16/(e*x + d)^3 + 9*B*a*b^2*d^3*e^16/(e*x + d)^4 + 3*A*b^3*d^3*e^16/(e*x + d)^4 - 18*B*a^2*b*e^17/(e*x + d)^2 - 18*A*a*b^2*e^17/(e*x + d)^2 + 24*B*a^2*b*d*e^17/(e*x + d)^3 + 24*A*a*b^2*d*e^17/(e*x + d)^3 - 9*B*a^2*b*d^2*e^17/(e*x + d)^4 - 9*A*a*b^2*d^2*e^17/(e*x + d)^4 - 4*B*a^3*e^18/(e*x + d)^3 - 12*A*a^2*b*e^18/(e*x + d)^3 + 3*B*a^3*d*e^18/(e*x + d)^4 + 9*A*a^2*b*d*e^18/(e*x + d)^4 - 3*A*a^3*e^19/(e*x + d)^4)/e^20
```


3.36 $\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^6} dx$

Optimal result	370
Mathematica [B] (verified)	370
Rubi [A] (verified)	371
Maple [B] (verified)	372
Fricas [B] (verification not implemented)	373
Sympy [B] (verification not implemented)	373
Maxima [B] (verification not implemented)	374
Giac [B] (verification not implemented)	375
Mupad [B] (verification not implemented)	375
Reduce [B] (verification not implemented)	376

Optimal result

Integrand size = 20, antiderivative size = 86

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^6} dx = -\frac{(Bd - Ae)(a+bx)^4}{5e(bd - ae)(d+ex)^5} + \frac{(4bBd + Abe - 5aBe)(a+bx)^4}{20e(bd - ae)^2(d+ex)^4}$$

output

```
-1/5*(-A*e+B*d)*(b*x+a)^4/e/(-a*e+b*d)/(e*x+d)^5+1/20*(A*b*e-5*B*a*e+4*B*b*d)*(b*x+a)^4/e/(-a*e+b*d)^2/(e*x+d)^4
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 211 vs. 2(86) = 172.

Time = 0.06 (sec) , antiderivative size = 211, normalized size of antiderivative = 2.45

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^6} dx = \frac{a^3e^3(4Ae + B(d + 5ex)) + a^2be^2(3Ae(d + 5ex) + 2B(d^2 + 5dex + 10e^2x^2)) + ab^2e(2Ae(d^2 + 5dex + 10e^2x^2)) + b^3e^3(4Ae + B(d + 5ex))}{(d+ex)^5}$$

input

```
Integrate[((a + b*x)^3*(A + B*x))/(d + e*x)^6,x]
```

output

$$\begin{aligned} & -1/20*(a^3*e^3*(4*A*e + B*(d + 5*e*x)) + a^2*b*e^2*(3*A*e*(d + 5*e*x) + 2* \\ & B*(d^2 + 5*d*e*x + 10*e^2*x^2)) + a*b^2*e*(2*A*e*(d^2 + 5*d*e*x + 10*e^2*x \\ & ^2) + 3*B*(d^3 + 5*d^2*e*x + 10*d*e^2*x^2 + 10*e^3*x^3)) + b^3*(A*e*(d^3 + \\ & 5*d^2*e*x + 10*d*e^2*x^2 + 10*e^3*x^3) + 4*B*(d^4 + 5*d^3*e*x + 10*d^2*e^ \\ & 2*x^2 + 10*d*e^3*x^3 + 5*e^4*x^4))/(e^5*(d + e*x)^5) \end{aligned}$$

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {87, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + bx)^3(A + Bx)}{(d + ex)^6} dx \\ & \quad \downarrow 87 \\ & \frac{(-5aBe + Abe + 4bBd) \int \frac{(a+bx)^3}{(d+ex)^5} dx}{5e(bd - ae)} - \frac{(a + bx)^4(Bd - Ae)}{5e(d + ex)^5(bd - ae)} \\ & \quad \downarrow 48 \\ & \frac{(a + bx)^4(-5aBe + Abe + 4bBd)}{20e(d + ex)^4(bd - ae)^2} - \frac{(a + bx)^4(Bd - Ae)}{5e(d + ex)^5(bd - ae)} \end{aligned}$$

input

$$\text{Int}[(a + b*x)^3*(A + B*x)/(d + e*x)^6, x]$$

output

$$\begin{aligned} & -1/5*((B*d - A*e)*(a + b*x)^4)/(e*(b*d - a*e)*(d + e*x)^5) + ((4*b*B*d + A \\ & *b*e - 5*a*B*e)*(a + b*x)^4)/(20*e*(b*d - a*e)^2*(d + e*x)^4) \end{aligned}$$

Defintions of rubi rules used

```
rule 48 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[[a + b*x]^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 263 vs. 2(82) = 164.

Time = 0.22 (sec) , antiderivative size = 264, normalized size of antiderivative = 3.07

method	result
risch	$\frac{-\frac{b^3 B x^4}{e} - \frac{b^2(Abe+3Bae+4Bbd)x^3}{2e^2} - \frac{b(2Aab e^2 + A b^2 d e + 2B a^2 e^2 + 3Babde + 4b^2 B d^2)x^2}{2e^3} - \frac{(3A a^2 b e^3 + 2Aa b^2 d e^2 + A b^3 d^2 e + B a^3 e^3 + 4e^4)}{(ex+d)^5}}$
norman	$\frac{-\frac{b^3 B x^4}{e} - \frac{(A b^3 e + 3B a b^2 e + 4b^3 B d)x^3}{2e^2} - \frac{(2Aa b^2 e^2 + A b^3 d e + 2B a^2 b e^2 + 3B a b^2 d e + 4b^3 B d^2)x^2}{2e^3} - \frac{(3A a^2 b e^3 + 2Aa b^2 d e^2 + A b^3 d^2 e + B a^3 e^3 + 4e^4)}{(ex+d)^5}}$
default	$-\frac{b^3 B}{e^5(ex+d)} - \frac{b^2(Abe+3Bae-4Bbd)}{2e^5(ex+d)^2} - \frac{a^3 A e^4 - 3A a^2 b d e^3 + 3A a b^2 d^2 e^2 - A b^3 d^3 e - B a^3 d e^3 + 3B a^2 b d^2 e^2 - 3B a b^2 d^3 e + 4e^4}{5e^5(ex+d)^5}$
gospers	$-\frac{20B x^4 b^3 e^4 + 10A x^3 b^3 e^4 + 30B x^3 a b^2 e^4 + 40B x^3 b^3 d e^3 + 20A x^2 a b^2 e^4 + 10A x^2 b^3 d e^3 + 20B x^2 a^2 b e^4 + 30B x^2 a b^2 d e^3 + 40B x^2 b^3 d^2 e^2 + 20A x a^2 b^2 e^4 + 10A x a^2 b^3 d e^3 + 20B x a^2 b^2 d e^3 + 30B x a b^2 d^2 e^2 + 4e^4}{5e^5(ex+d)^5}$
parallelrisch	$-\frac{20B x^4 b^3 e^4 + 10A x^3 b^3 e^4 + 30B x^3 a b^2 e^4 + 40B x^3 b^3 d e^3 + 20A x^2 a b^2 e^4 + 10A x^2 b^3 d e^3 + 20B x^2 a^2 b e^4 + 30B x^2 a b^2 d e^3 + 40B x^2 b^3 d^2 e^2 + 20A x a^2 b^2 e^4 + 10A x a^2 b^3 d e^3 + 20B x a^2 b^2 d e^3 + 30B x a b^2 d^2 e^2 + 4e^4}{5e^5(ex+d)^5}$
orering	$-\frac{20B x^4 b^3 e^4 + 10A x^3 b^3 e^4 + 30B x^3 a b^2 e^4 + 40B x^3 b^3 d e^3 + 20A x^2 a b^2 e^4 + 10A x^2 b^3 d e^3 + 20B x^2 a^2 b e^4 + 30B x^2 a b^2 d e^3 + 40B x^2 b^3 d^2 e^2 + 20A x a^2 b^2 e^4 + 10A x a^2 b^3 d e^3 + 20B x a^2 b^2 d e^3 + 30B x a b^2 d^2 e^2 + 4e^4}{5e^5(ex+d)^5}$

```
input int((b*x+a)^3*(B*x+A)/(e*x+d)^6,x,method=_RETURNVERBOSE)
```

output

```
(-b^3*B/e*x^4-1/2*b^2*(A*b*e+3*B*a*e+4*B*b*d)/e^2*x^3-1/2*b*(2*A*a*b*e^2+A
*b^2*d*e+2*B*a^2*e^2+3*B*a*b*d*e+4*B*b^2*d^2)/e^3*x^2-1/4*(3*A*a^2*b*e^3+2
*A*a*b^2*d*e^2+A*b^3*d^2*e+B*a^3*e^3+2*B*a^2*b*d*e^2+3*B*a*b^2*d^2*e+4*B*b
^3*d^3)/e^4*x-1/20*(4*A*a^3*e^4+3*A*a^2*b*d*e^3+2*A*a*b^2*d^2*e^2+A*b^3*d^
3*e+B*a^3*d*e^3+2*B*a^2*b*d^2*e^2+3*B*a*b^2*d^3*e+4*B*b^3*d^4)/e^5)/(e*x+d
)^5
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 304 vs. $2(82) = 164$.

Time = 0.07 (sec) , antiderivative size = 304, normalized size of antiderivative = 3.53

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^6} dx =$$

$$\frac{20Bb^3e^4x^4 + 4Bb^3d^4 + 4Aa^3e^4 + (3Bab^2 + Ab^3)d^3e + 2(Ba^2b + Aab^2)d^2e^2 + (Ba^3 + 3Aa^2b)de^3 + \dots}{(d+ex)^6}$$

input

```
integrate((b*x+a)^3*(B*x+A)/(e*x+d)^6,x, algorithm="fricas")
```

output

```
-1/20*(20*B*b^3*e^4*x^4 + 4*B*b^3*d^4 + 4*A*a^3*e^4 + (3*B*a*b^2 + A*b^3)*
d^3*e + 2*(B*a^2*b + A*a*b^2)*d^2*e^2 + (B*a^3 + 3*A*a^2*b)*d*e^3 + 10*(4*
B*b^3*d*e^3 + (3*B*a*b^2 + A*b^3)*e^4)*x^3 + 10*(4*B*b^3*d^2*e^2 + (3*B*a*
b^2 + A*b^3)*d*e^3 + 2*(B*a^2*b + A*a*b^2)*e^4)*x^2 + 5*(4*B*b^3*d^3*e + (
3*B*a*b^2 + A*b^3)*d^2*e^2 + 2*(B*a^2*b + A*a*b^2)*d*e^3 + (B*a^3 + 3*A*a^
2*b)*e^4)*x)/(e^10*x^5 + 5*d*e^9*x^4 + 10*d^2*e^8*x^3 + 10*d^3*e^7*x^2 + 5
*d^4*e^6*x + d^5*e^5)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 372 vs. $2(75) = 150$.

Time = 35.07 (sec) , antiderivative size = 372, normalized size of antiderivative = 4.33

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^6} dx$$

$$= \frac{-4Aa^3e^4 - 3Aa^2bde^3 - 2Aab^2d^2e^2 - Ab^3d^3e - Ba^3de^3 - 2Ba^2bd^2e^2 - 3Bab^2d^3e - 4Bb^3d^4 - 20Bb^3e^4x}{(d+ex)^6}$$

input `integrate((b*x+a)**3*(B*x+A)/(e*x+d)**6,x)`

output `(-4*A**3*e**4 - 3*A**2*b*d*e**3 - 2*A*b**2*d**2*e**2 - A*b**3*d**3*e - B**3*d*e**3 - 2*B**2*b*d**2*e**2 - 3*B*a*b**2*d**3*e - 4*B*b**3*d**4 - 20*B*b**3*e**4*x**4 + x**3*(-10*A*b**3*e**4 - 30*B*a*b**2*e**4 - 40*B*b**3*d*e**3) + x**2*(-20*A*a*b**2*e**4 - 10*A*b**3*d*e**3 - 20*B**2*b*e**4 - 30*B*a*b**2*d*e**3 - 40*B*b**3*d**2*e**2) + x*(-15*A**2*b*e**4 - 10*A*a*b**2*d*e**3 - 5*A*b**3*d**2*e**2 - 5*B**3*e**4 - 10*B**2*b*d*e**3 - 15*B*a*b**2*d**2*e**2 - 20*B*b**3*d**3*e))/(20*d**5*e**5 + 100*d**4*e**6*x + 200*d**3*e**7*x**2 + 200*d**2*e**8*x**3 + 100*d*e**9*x**4 + 20*e**10*x**5)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 304 vs. $2(82) = 164$.

Time = 0.04 (sec) , antiderivative size = 304, normalized size of antiderivative = 3.53

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^6} dx = \frac{20Bb^3e^4x^4 + 4Bb^3d^4 + 4Aa^3e^4 + (3Bab^2 + Ab^3)d^3e + 2(Ba^2b + Aab^2)d^2e^2 + (Ba^3 + 3Aa^2b)de^3 + \dots}{(d+ex)^6}$$

input `integrate((b*x+a)^3*(B*x+A)/(e*x+d)^6,x, algorithm="maxima")`

output `-1/20*(20*B*b^3*e^4*x^4 + 4*B*b^3*d^4 + 4*A*a^3*e^4 + (3*B*a*b^2 + A*b^3)*d^3*e + 2*(B*a^2*b + A*a*b^2)*d^2*e^2 + (B*a^3 + 3*A*a^2*b)*d*e^3 + 10*(4*B*b^3*d*e^3 + (3*B*a*b^2 + A*b^3)*e^4)*x^3 + 10*(4*B*b^3*d^2*e^2 + (3*B*a*b^2 + A*b^3)*d*e^3 + 2*(B*a^2*b + A*a*b^2)*e^4)*x^2 + 5*(4*B*b^3*d^3*e + (3*B*a*b^2 + A*b^3)*d^2*e^2 + 2*(B*a^2*b + A*a*b^2)*d*e^3 + (B*a^3 + 3*A*a^2*b)*e^4)*x)/(e^10*x^5 + 5*d*e^9*x^4 + 10*d^2*e^8*x^3 + 10*d^3*e^7*x^2 + 5*d^4*e^6*x + d^5*e^5)`

output

```

-((4*A*a^3*e^4 + 4*B*b^3*d^4 + A*b^3*d^3*e + B*a^3*d*e^3 + 2*A*a*b^2*d^2*e
^2 + 2*B*a^2*b*d^2*e^2 + 3*A*a^2*b*d*e^3 + 3*B*a*b^2*d^3*e)/(20*e^5) + (x*
(B*a^3*e^3 + 4*B*b^3*d^3 + 3*A*a^2*b*e^3 + A*b^3*d^2*e + 2*A*a*b^2*d*e^2 +
3*B*a*b^2*d^2*e + 2*B*a^2*b*d*e^2))/(4*e^4) + (b^2*x^3*(A*b*e + 3*B*a*e +
4*B*b*d))/(2*e^2) + (b*x^2*(2*B*a^2*e^2 + 4*B*b^2*d^2 + 2*A*a*b*e^2 + A*b
^2*d*e + 3*B*a*b*d*e))/(2*e^3) + (B*b^3*x^4)/e)/(d^5 + e^5*x^5 + 5*d*e^4*x
^4 + 10*d^3*e^2*x^2 + 10*d^2*e^3*x^3 + 5*d^4*e*x)

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 192, normalized size of antiderivative = 2.23

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^6} dx$$

$$= \frac{b^4 e^4 x^5 - 10 a b^3 d e^3 x^3 - 10 a^2 b^2 d e^3 x^2 - 10 a b^3 d^2 e^2 x^2 - 5 a^3 b d e^3 x - 5 a^2 b^2 d^2 e^2 x - 5 a b^3 d^3 e x - a^4 d e^3 - a^5 d^4}{5 d e^4 (e^5 x^5 + 5 d e^4 x^4 + 10 d^2 e^3 x^3 + 10 d^3 e^2 x^2 + 5 d^4 e x + d^5)}$$

input

```
int((b*x+a)^3*(B*x+A)/(e*x+d)^6,x)
```

output

```

( - a**4*d*e**3 - a**3*b*d**2*e**2 - 5*a**3*b*d*e**3*x - a**2*b**2*d**3*e
- 5*a**2*b**2*d**2*e**2*x - 10*a**2*b**2*d*e**3*x**2 - a*b**3*d**4 - 5*a*b
**3*d**3*e*x - 10*a*b**3*d**2*e**2*x**2 - 10*a*b**3*d*e**3*x**3 + b**4*e**
4*x**5)/(5*d*e**4*(d**5 + 5*d**4*e*x + 10*d**3*e**2*x**2 + 10*d**2*e**3*x*
*3 + 5*d*e**4*x**4 + e**5*x**5))

```

3.37 $\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^7} dx$

Optimal result	377
Mathematica [A] (verified)	377
Rubi [A] (verified)	378
Maple [B] (verified)	379
Fricas [B] (verification not implemented)	380
Sympy [B] (verification not implemented)	381
Maxima [B] (verification not implemented)	382
Giac [B] (verification not implemented)	382
Mupad [B] (verification not implemented)	383
Reduce [B] (verification not implemented)	383

Optimal result

Integrand size = 20, antiderivative size = 133

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^7} dx = -\frac{(Bd - Ae)(a+bx)^4}{6e(bd - ae)(d+ex)^6} + \frac{(2bBd + Abe - 3aBe)(a+bx)^4}{15e(bd - ae)^2(d+ex)^5} + \frac{b(2bBd + Abe - 3aBe)(a+bx)^4}{60e(bd - ae)^3(d+ex)^4}$$

output

```
-1/6*(-A*e+B*d)*(b*x+a)^4/e/(-a*e+b*d)/(e*x+d)^6+1/15*(A*b*e-3*B*a*e+2*B*b*d)*(b*x+a)^4/e/(-a*e+b*d)^2/(e*x+d)^5+1/60*b*(A*b*e-3*B*a*e+2*B*b*d)*(b*x+a)^4/e/(-a*e+b*d)^3/(e*x+d)^4
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.59

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^7} dx = \frac{2a^3e^3(5Ae + B(d+6ex)) + 3a^2be^2(2Ae(d+6ex) + B(d^2 + 6dex + 15e^2x^2)) + 3ab^2e(Ae(d^2 + 6dex -$$

input

```
Integrate[((a + b*x)^3*(A + B*x))/(d + e*x)^7,x]
```

output

$$\begin{aligned} & -1/60*(2*a^3*e^3*(5*A*e + B*(d + 6*e*x)) + 3*a^2*b*e^2*(2*A*e*(d + 6*e*x) \\ & + B*(d^2 + 6*d*e*x + 15*e^2*x^2)) + 3*a*b^2*e*(A*e*(d^2 + 6*d*e*x + 15*e^2 \\ & *x^2) + B*(d^3 + 6*d^2*e*x + 15*d*e^2*x^2 + 20*e^3*x^3)) + b^3*(A*e*(d^3 + \\ & 6*d^2*e*x + 15*d*e^2*x^2 + 20*e^3*x^3) + 2*B*(d^4 + 6*d^3*e*x + 15*d^2*e^ \\ & 2*x^2 + 20*d*e^3*x^3 + 15*e^4*x^4)))/(e^5*(d + e*x)^6) \end{aligned}$$

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.98, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {87, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + bx)^3(A + Bx)}{(d + ex)^7} dx \\ & \quad \downarrow 87 \\ & \frac{(-3aBe + Abe + 2bBd) \int \frac{(a+bx)^3}{(d+ex)^6} dx}{3e(bd - ae)} - \frac{(a + bx)^4(Bd - Ae)}{6e(d + ex)^6(bd - ae)} \\ & \quad \downarrow 55 \\ & \frac{(-3aBe + Abe + 2bBd) \left(\frac{b \int \frac{(a+bx)^3}{(d+ex)^5} dx}{5(bd - ae)} + \frac{(a+bx)^4}{5(d+ex)^5(bd - ae)} \right)}{3e(bd - ae)} - \frac{(a + bx)^4(Bd - Ae)}{6e(d + ex)^6(bd - ae)} \\ & \quad \downarrow 48 \\ & \frac{\left(\frac{b(a+bx)^4}{20(d+ex)^4(bd - ae)^2} + \frac{(a+bx)^4}{5(d+ex)^5(bd - ae)} \right) (-3aBe + Abe + 2bBd)}{3e(bd - ae)} - \frac{(a + bx)^4(Bd - Ae)}{6e(d + ex)^6(bd - ae)} \end{aligned}$$

input

```
Int[((a + b*x)^3*(A + B*x))/(d + e*x)^7,x]
```

output

$$\begin{aligned} & -1/6*((B*d - A*e)*(a + b*x)^4)/(e*(b*d - a*e)*(d + e*x)^6) + ((2*b*B*d + A \\ & *b*e - 3*a*B*e)*((a + b*x)^4/(5*(b*d - a*e)*(d + e*x)^5) + (b*(a + b*x)^4) \\ & / (20*(b*d - a*e)^2*(d + e*x)^4)))/(3*e*(b*d - a*e)) \end{aligned}$$

Defintions of rubi rules used

rule 48

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

rule 55

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
lerQ[n, 1])
```

rule 87

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && ( !LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 265 vs. $2(127) = 254$.

Time = 0.22 (sec) , antiderivative size = 266, normalized size of antiderivative = 2.00

method	result
risch	$\frac{-\frac{b^3 B x^4}{2e} - \frac{b^2(Abe+3Bae+2Bbd)x^3}{3e^2} - \frac{b(3Aab e^2 + A b^2 de + 3B a^2 e^2 + 3Babde + 2b^2 B d^2)x^2}{4e^3} - \frac{(6A a^2 b e^3 + 3Aa b^2 d e^2 + A b^3 d^2 e + 2B a^3 e^3 + 10e^4)}{(ex+d)^6}$
default	$-\frac{a^3 A e^4 - 3A a^2 b d e^3 + 3Aa b^2 d^2 e^2 - A b^3 d^3 e - B a^3 d e^3 + 3B a^2 b d^2 e^2 - 3Ba b^2 d^3 e + b^3 B d^4}{6e^5(ex+d)^6} - \frac{b^3 B}{2e^5(ex+d)^2} - \frac{3A a^2 b e^3 - 6A a b^2 d e^2 + 3A a^2 b^2 d^2 e}{6e^5(ex+d)^2}$
norman	$\frac{-\frac{b^3 B x^4}{2e} - \frac{(A b^3 e^2 + 3B a b^2 e^2 + 2b^3 B d e)x^3}{3e^3} - \frac{(3A a b^2 e^3 + A b^3 d e^2 + 3B a^2 b e^3 + 3B a b^2 d e^2 + 2b^3 B d^2 e)x^2}{4e^4} - \frac{(6A a^2 b e^4 + 3Aa b^2 d e^3 + A b^3 d^2 e^2 + 10e^4)}{(ex+d)^6}$
gosper	$-\frac{30 B x^4 b^3 e^4 + 20 A x^3 b^3 e^4 + 60 B x^3 a b^2 e^4 + 40 B x^3 b^3 d e^3 + 45 A x^2 a b^2 e^4 + 15 A x^2 b^3 d e^3 + 45 B x^2 a^2 b e^4 + 45 B x^2 a b^2 d e^3 + 30 B x^2 b^3 d^2 e^2 + 10 e^4}{6e^5(ex+d)^6}$
orering	$-\frac{30 B x^4 b^3 e^4 + 20 A x^3 b^3 e^4 + 60 B x^3 a b^2 e^4 + 40 B x^3 b^3 d e^3 + 45 A x^2 a b^2 e^4 + 15 A x^2 b^3 d e^3 + 45 B x^2 a^2 b e^4 + 45 B x^2 a b^2 d e^3 + 30 B x^2 b^3 d^2 e^2 + 10 e^4}{6e^5(ex+d)^6}$
parallelrisch	$-\frac{30 B b^3 x^4 e^5 + 20 A b^3 e^5 x^3 + 60 B a b^2 e^5 x^3 + 40 B b^3 d e^4 x^3 + 45 A a b^2 e^5 x^2 + 15 A b^3 d e^4 x^2 + 45 B a^2 b e^5 x^2 + 45 B a b^2 d e^4 x^2 + 30 B x^2 b^3 d^2 e^2 + 10 e^4}{6e^5(ex+d)^6}$

```
input int((b*x+a)^3*(B*x+A)/(e*x+d)^7,x,method=_RETURNVERBOSE)
```

```
output (-1/2*b^3*B/e*x^4-1/3*b^2/e^2*(A*b*e+3*B*a*e+2*B*b*d)*x^3-1/4*b/e^3*(3*A*a*b*e^2+A*b^2*d*e+3*B*a^2*e^2+3*B*a*b*d*e+2*B*b^2*d^2)*x^2-1/10/e^4*(6*A*a^2*b*e^3+3*A*a*b^2*d*e^2+A*b^3*d^2*e+2*B*a^3*e^3+3*B*a^2*b*d*e^2+3*B*a*b^2*d^2*e+2*B*b^3*d^3)*x-1/60/e^5*(10*A*a^3*e^4+6*A*a^2*b*d*e^3+3*A*a*b^2*d^2*e^2+A*b^3*d^3*e+2*B*a^3*d*e^3+3*B*a^2*b*d^2*e^2+3*B*a*b^2*d^3*e+2*B*b^3*d^4))/(e*x+d)^6
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 317 vs. 2(127) = 254.

Time = 0.07 (sec) , antiderivative size = 317, normalized size of antiderivative = 2.38

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^7} dx =$$

$$-\frac{30 B b^3 e^4 x^4 + 2 B b^3 d^4 + 10 A a^3 e^4 + (3 B a b^2 + A b^3) d^3 e + 3 (B a^2 b + A a b^2) d^2 e^2 + 2 (B a^3 + 3 A a^2 b) d e^3 + 10 A a^2 b d e^2 + 6 A a b^2 d^2 e + 2 B a^3 d e + 2 B a^2 b d^2 e + 2 B a b^2 d^3 e + 2 B b^3 d^4}{6e^5(ex+d)^6}$$

```
input integrate((b*x+a)^3*(B*x+A)/(e*x+d)^7,x, algorithm="fricas")
```

output

```
-1/60*(30*B*b^3*e^4*x^4 + 2*B*b^3*d^4 + 10*A*a^3*e^4 + (3*B*a*b^2 + A*b^3)
*d^3*e + 3*(B*a^2*b + A*a*b^2)*d^2*e^2 + 2*(B*a^3 + 3*A*a^2*b)*d*e^3 + 20*
(2*B*b^3*d*e^3 + (3*B*a*b^2 + A*b^3)*e^4)*x^3 + 15*(2*B*b^3*d^2*e^2 + (3*B
*a*b^2 + A*b^3)*d*e^3 + 3*(B*a^2*b + A*a*b^2)*e^4)*x^2 + 6*(2*B*b^3*d^3*e
+ (3*B*a*b^2 + A*b^3)*d^2*e^2 + 3*(B*a^2*b + A*a*b^2)*d*e^3 + 2*(B*a^3 + 3
*A*a^2*b)*e^4)*x)/(e^11*x^6 + 6*d*e^10*x^5 + 15*d^2*e^9*x^4 + 20*d^3*e^8*x
^3 + 15*d^4*e^7*x^2 + 6*d^5*e^6*x + d^6*e^5)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 386 vs. $2(122) = 244$.

Time = 100.28 (sec) , antiderivative size = 386, normalized size of antiderivative = 2.90

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^7} dx$$

$$= \frac{-10Aa^3e^4 - 6Aa^2bde^3 - 3Aab^2d^2e^2 - Ab^3d^3e - 2Ba^3de^3 - 3Ba^2bd^2e^2 - 3Bab^2d^3e - 2Bb^3d^4 - 30Bb^3e^4}{(d + ex)^7}$$

input

```
integrate((b*x+a)**3*(B*x+A)/(e*x+d)**7,x)
```

output

```
(-10*A*a**3*e**4 - 6*A*a**2*b*d*e**3 - 3*A*a*b**2*d**2*e**2 - A*b**3*d**3*
e - 2*B*a**3*d*e**3 - 3*B*a**2*b*d**2*e**2 - 3*B*a*b**2*d**3*e - 2*B*b**3*
d**4 - 30*B*b**3*e**4*x**4 + x**3*(-20*A*b**3*e**4 - 60*B*a*b**2*e**4 - 40
*B*b**3*d*e**3) + x**2*(-45*A*a*b**2*e**4 - 15*A*b**3*d*e**3 - 45*B*a**2*b
*e**4 - 45*B*a*b**2*d*e**3 - 30*B*b**3*d**2*e**2) + x*(-36*A*a**2*b*e**4 -
18*A*a*b**2*d*e**3 - 6*A*b**3*d**2*e**2 - 12*B*a**3*e**4 - 18*B*a**2*b*d
e**3 - 18*B*a*b**2*d**2*e**2 - 12*B*b**3*d**3*e))/(60*d**6*e**5 + 360*d**5
*e**6*x + 900*d**4*e**7*x**2 + 1200*d**3*e**8*x**3 + 900*d**2*e**9*x**4 +
360*d**10*x**5 + 60*e**11*x**6)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 317 vs. $2(127) = 254$.

Time = 0.04 (sec) , antiderivative size = 317, normalized size of antiderivative = 2.38

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^7} dx = \frac{30 Bb^3e^4x^4 + 2 Bb^3d^4 + 10 Aa^3e^4 + (3 Bab^2 + Ab^3)d^3e + 3 (Ba^2b + Aab^2)d^2e^2 + 2 (Ba^3 + 3 Aa^2b)de^3}{(d+ex)^7}$$

input `integrate((b*x+a)^3*(B*x+A)/(e*x+d)^7,x, algorithm="maxima")`

output `-1/60*(30*B*b^3*e^4*x^4 + 2*B*b^3*d^4 + 10*A*a^3*e^4 + (3*B*a*b^2 + A*b^3)*d^3*e + 3*(B*a^2*b + A*a*b^2)*d^2*e^2 + 2*(B*a^3 + 3*A*a^2*b)*d*e^3 + 20*(2*B*b^3*d*e^3 + (3*B*a*b^2 + A*b^3)*e^4)*x^3 + 15*(2*B*b^3*d^2*e^2 + (3*B*a*b^2 + A*b^3)*d*e^3 + 3*(B*a^2*b + A*a*b^2)*e^4)*x^2 + 6*(2*B*b^3*d^3*e + (3*B*a*b^2 + A*b^3)*d^2*e^2 + 3*(B*a^2*b + A*a*b^2)*d*e^3 + 2*(B*a^3 + 3*A*a^2*b)*e^4)*x)/(e^11*x^6 + 6*d*e^10*x^5 + 15*d^2*e^9*x^4 + 20*d^3*e^8*x^3 + 15*d^4*e^7*x^2 + 6*d^5*e^6*x + d^6*e^5)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 299 vs. $2(127) = 254$.

Time = 0.13 (sec) , antiderivative size = 299, normalized size of antiderivative = 2.25

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^7} dx = \frac{30 Bb^3e^4x^4 + 40 Bb^3de^3x^3 + 60 Bab^2e^4x^3 + 20 Ab^3e^4x^3 + 30 Bb^3d^2e^2x^2 + 45 Bab^2de^3x^2 + 15 Ab^3de^3x}{(d+ex)^7}$$

input `integrate((b*x+a)^3*(B*x+A)/(e*x+d)^7,x, algorithm="giac")`

input `int((b*x+a)^3*(B*x+A)/(e*x+d)^7,x)`

output `(- 5*a**4*e**4 - 4*a**3*b*d*e**3 - 24*a**3*b*e**4*x - 3*a**2*b**2*d**2*e**
*2 - 18*a**2*b**2*d*e**3*x - 45*a**2*b**2*e**4*x**2 - 2*a*b**3*d**3*e - 12
*a*b**3*d**2*e**2*x - 30*a*b**3*d*e**3*x**2 - 40*a*b**3*e**4*x**3 - b**4*d
4 - 6*b4*d**3*e*x - 15*b**4*d**2*e**2*x**2 - 20*b**4*d*e**3*x**3 - 15*
b**4*e**4*x**4)/(30*e**5*(d**6 + 6*d**5*e*x + 15*d**4*e**2*x**2 + 20*d**3*
e**3*x**3 + 15*d**2*e**4*x**4 + 6*d*e**5*x**5 + e**6*x**6))`

3.38 $\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^8} dx$

Optimal result	385
Mathematica [A] (verified)	386
Rubi [A] (verified)	386
Maple [A] (verified)	388
Fricas [B] (verification not implemented)	388
Sympy [F(-1)]	389
Maxima [B] (verification not implemented)	389
Giac [A] (verification not implemented)	390
Mupad [B] (verification not implemented)	390
Reduce [B] (verification not implemented)	391

Optimal result

Integrand size = 20, antiderivative size = 163

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^8} dx = -\frac{(bd-ae)^3(Bd-Ae)}{7e^5(d+ex)^7} + \frac{(bd-ae)^2(4bBd-3Abe-aBe)}{6e^5(d+ex)^6} - \frac{3b(bd-ae)(2bBd-Abe-aBe)}{5e^5(d+ex)^5} + \frac{b^2(4bBd-Abe-3aBe)}{4e^5(d+ex)^4} - \frac{b^3B}{3e^5(d+ex)^3}$$

output

```
-1/7*(-a*e+b*d)^3*(-A*e+B*d)/e^5/(e*x+d)^7+1/6*(-a*e+b*d)^2*(-3*A*b*e-B*a*
e+4*B*b*d)/e^5/(e*x+d)^6-3/5*b*(-a*e+b*d)*(-A*b*e-B*a*e+2*B*b*d)/e^5/(e*x+
d)^5+1/4*b^2*(-A*b*e-3*B*a*e+4*B*b*d)/e^5/(e*x+d)^4-1/3*b^3*B/e^5/(e*x+d)^
3
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.32

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^8} dx =$$

$$\frac{10a^3e^3(6Ae + B(d + 7ex)) + 6a^2be^2(5Ae(d + 7ex) + 2B(d^2 + 7dex + 21e^2x^2)) + 3ab^2e(4Ae(d^2 + 7d$$

input `Integrate[((a + b*x)^3*(A + B*x))/(d + e*x)^8,x]`

output

```
-1/420*(10*a^3*e^3*(6*A*e + B*(d + 7*e*x)) + 6*a^2*b*e^2*(5*A*e*(d + 7*e*x)
) + 2*B*(d^2 + 7*d*e*x + 21*e^2*x^2)) + 3*a*b^2*e*(4*A*e*(d^2 + 7*d*e*x +
21*e^2*x^2) + 3*B*(d^3 + 7*d^2*e*x + 21*d*e^2*x^2 + 35*e^3*x^3)) + b^3*(3*
A*e*(d^3 + 7*d^2*e*x + 21*d*e^2*x^2 + 35*e^3*x^3) + 4*B*(d^4 + 7*d^3*e*x +
21*d^2*e^2*x^2 + 35*d*e^3*x^3 + 35*e^4*x^4)))/(e^5*(d + e*x)^7)
```

Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^8} dx$$

$$\downarrow 86$$

$$\int \left(\frac{b^2(3aBe + Abe - 4bBd)}{e^4(d + ex)^5} - \frac{3b(bd - ae)(aBe + Abe - 2bBd)}{e^4(d + ex)^6} + \frac{(ae - bd)^2(aBe + 3Abe - 4bBd)}{e^4(d + ex)^7} + \frac{(ae - bd)^3}{e^4(d + ex)^8} \right) dx$$

$$\downarrow 2009$$

$$\frac{b^2(-3aBe - Abe + 4bBd)}{4e^5(d+ex)^4} - \frac{3b(bd-ae)(-aBe - Abe + 2bBd)}{5e^5(d+ex)^5} + \frac{(bd-ae)^2(-aBe - 3Abe + 4bBd)}{6e^5(d+ex)^6} - \frac{(bd-ae)^3(Bd - Ae)}{7e^5(d+ex)^7} - \frac{b^3B}{3e^5(d+ex)^3}$$

input `Int[((a + b*x)^3*(A + B*x))/(d + e*x)^8,x]`

output `-1/7*((b*d - a*e)^3*(B*d - A*e))/(e^5*(d + e*x)^7) + ((b*d - a*e)^2*(4*b*B*d - 3*A*b*e - a*B*e))/(6*e^5*(d + e*x)^6) - (3*b*(b*d - a*e)*(2*b*B*d - A*b*e - a*B*e))/(5*e^5*(d + e*x)^5) + (b^2*(4*b*B*d - A*b*e - 3*a*B*e))/(4*e^5*(d + e*x)^4) - (b^3*B)/(3*e^5*(d + e*x)^3)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;`
`FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.66

method	result
risch	$\frac{-\frac{b^3 B x^4}{3e} - \frac{b^2(3Abe+9Bae+4Bbd)x^3}{12e^2} - \frac{b(12Aab e^2+3A b^2 de+12B a^2 e^2+9Babde+4b^2 B d^2)x^2}{20e^3} - \frac{(30A a^2 b e^3+12Aa b^2 d e^2+3A b^3 d^2 e+12A a^2 b^2 e^3-6Aa b^2 d e^2+3A b^3 d^2 e+B a^3 e^3-6B a^2 b d e^2+9B a b^2 d^2 e-4b^3 B d^3)x}{6e^5(e x+d)^6} - \frac{a^3 A e^4-3A a^2 b d e^3+3A a b^2 d^2 e^2-A b^3 d^3 e}{7e^5}}$
default	$\frac{-3A a^2 b e^3-6Aa b^2 d e^2+3A b^3 d^2 e+B a^3 e^3-6B a^2 b d e^2+9B a b^2 d^2 e-4b^3 B d^3}{6e^5(e x+d)^6} - \frac{a^3 A e^4-3A a^2 b d e^3+3A a b^2 d^2 e^2-A b^3 d^3 e}{7e^5}$
gospers	$-\frac{140B x^4 b^3 e^4+105A x^3 b^3 e^4+315B x^3 a b^2 e^4+140B x^3 b^3 d e^3+252A x^2 a b^2 e^4+63A x^2 b^3 d e^3+252B x^2 a^2 b e^4+189B x^2 a b^2 d e^3}{6e^5(e x+d)^6}$
orering	$-\frac{140B x^4 b^3 e^4+105A x^3 b^3 e^4+315B x^3 a b^2 e^4+140B x^3 b^3 d e^3+252A x^2 a b^2 e^4+63A x^2 b^3 d e^3+252B x^2 a^2 b e^4+189B x^2 a b^2 d e^3}{6e^5(e x+d)^6}$
norman	$\frac{-\frac{b^3 B x^4}{3e} - \frac{(3A b^3 e^3+9B a b^2 e^3+4b^3 B d e^2)x^3}{12e^4} - \frac{(12A a b^2 e^4+3A b^3 d e^3+12B a^2 b e^4+9B a b^2 d e^3+4b^3 B d^2 e^2)x^2}{20e^5} - \frac{(30A a^2 b e^5+12A a b^2 d e^4+3A b^3 d^2 e^3+12A a^2 b^2 e^3-6Aa b^2 d e^2+3A b^3 d^2 e+B a^3 e^3-6B a^2 b d e^2+9B a b^2 d^2 e-4b^3 B d^3)x}{6e^5(e x+d)^6} - \frac{a^3 A e^4-3A a^2 b d e^3+3A a b^2 d^2 e^2-A b^3 d^3 e}{7e^5}}$
parallelrisc	$-\frac{140B b^3 x^4 e^6+105A b^3 e^6 x^3+315B a b^2 e^6 x^3+140B b^3 d e^5 x^3+252A a b^2 e^6 x^2+63A b^3 d e^5 x^2+252B a^2 b e^6 x^2+189B a b^2 d e^5 x}{6e^5(e x+d)^6}$

```
input int((b*x+a)^3*(B*x+A)/(e*x+d)^8,x,method=_RETURNVERBOSE)
```

```
output (-1/3*b^3*B/e*x^4-1/12*b^2/e^2*(3*A*b*e+9*B*a*e+4*B*b*d)*x^3-1/20*b/e^3*(12*A*a*b*e^2+3*A*b^2*d*e+12*B*a^2*e^2+9*B*a*b*d*e+4*B*b^2*d^2)*x^2-1/60/e^4*(30*A*a^2*b*e^3+12*A*a*b^2*d*e^2+3*A*b^3*d^2*e+10*B*a^3*e^3+12*B*a^2*b*d*e^2+9*B*a*b^2*d^2*e+4*B*b^3*d^3)*x-1/420/e^5*(60*A*a^3*e^4+30*A*a^2*b*d*e^3+12*A*a*b^2*d^2*e^2+3*A*b^3*d^3*e+10*B*a^3*d*e^3+12*B*a^2*b*d^2*e^2+9*B*a*b^2*d^3*e+4*B*b^3*d^4))/(e*x+d)^7
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 332 vs. 2(153) = 306.

Time = 0.07 (sec) , antiderivative size = 332, normalized size of antiderivative = 2.04

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^8} dx = \frac{140 B b^3 e^4 x^4 + 4 B b^3 d^4 + 60 A a^3 e^4 + 3(3 B a b^2 + A b^3) d^3 e + 12 (B a^2 b + A a b^2) d^2 e^2 + 10 (B a^3 + 3 A a^2 b) d e^3 + 3 A^2 e^4}{(d + ex)^7}$$

```
input integrate((b*x+a)^3*(B*x+A)/(e*x+d)^8,x, algorithm="fricas")
```

output

```
-1/420*(140*B*b^3*e^4*x^4 + 4*B*b^3*d^4 + 60*A*a^3*e^4 + 3*(3*B*a*b^2 + A*
b^3)*d^3*e + 12*(B*a^2*b + A*a*b^2)*d^2*e^2 + 10*(B*a^3 + 3*A*a^2*b)*d*e^3
+ 35*(4*B*b^3*d*e^3 + 3*(3*B*a*b^2 + A*b^3)*e^4)*x^3 + 21*(4*B*b^3*d^2*e^
2 + 3*(3*B*a*b^2 + A*b^3)*d*e^3 + 12*(B*a^2*b + A*a*b^2)*e^4)*x^2 + 7*(4*B
*b^3*d^3*e + 3*(3*B*a*b^2 + A*b^3)*d^2*e^2 + 12*(B*a^2*b + A*a*b^2)*d*e^3
+ 10*(B*a^3 + 3*A*a^2*b)*e^4)*x)/(e^12*x^7 + 7*d*e^11*x^6 + 21*d^2*e^10*x^
5 + 35*d^3*e^9*x^4 + 35*d^4*e^8*x^3 + 21*d^5*e^7*x^2 + 7*d^6*e^6*x + d^7*e
^5)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^8} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**3*(B*x+A)/(e*x+d)**8,x)
```

output

Timed out

Maxima [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 332 vs. $2(153) = 306$.

Time = 0.06 (sec) , antiderivative size = 332, normalized size of antiderivative = 2.04

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^8} dx = \frac{140 B b^3 e^4 x^4 + 4 B b^3 d^4 + 60 A a^3 e^4 + 3 (3 B a b^2 + A b^3) d^3 e + 12 (B a^2 b + A a b^2) d^2 e^2 + 10 (B a^3 + 3 A a^2 b) d e^3 + 3 (3 B a^2 b + A a b^2) e^4}{(d + e x)^8}$$

input

```
integrate((b*x+a)^3*(B*x+A)/(e*x+d)^8,x, algorithm="maxima")
```

output

```
-1/420*(140*B*b^3*e^4*x^4 + 4*B*b^3*d^4 + 60*A*a^3*e^4 + 3*(3*B*a*b^2 + A*
b^3)*d^3*e + 12*(B*a^2*b + A*a*b^2)*d^2*e^2 + 10*(B*a^3 + 3*A*a^2*b)*d*e^3
+ 35*(4*B*b^3*d*e^3 + 3*(3*B*a*b^2 + A*b^3)*e^4)*x^3 + 21*(4*B*b^3*d^2*e^
2 + 3*(3*B*a*b^2 + A*b^3)*d*e^3 + 12*(B*a^2*b + A*a*b^2)*e^4)*x^2 + 7*(4*B
*b^3*d^3*e + 3*(3*B*a*b^2 + A*b^3)*d^2*e^2 + 12*(B*a^2*b + A*a*b^2)*d*e^3
+ 10*(B*a^3 + 3*A*a^2*b)*e^4)*x)/(e^12*x^7 + 7*d*e^11*x^6 + 21*d^2*e^10*x^
5 + 35*d^3*e^9*x^4 + 35*d^4*e^8*x^3 + 21*d^5*e^7*x^2 + 7*d^6*e^6*x + d^7*e
^5)
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 300, normalized size of antiderivative = 1.84

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^8} dx = \frac{140 B b^3 e^4 x^4 + 140 B b^3 d e^3 x^3 + 315 B a b^2 e^4 x^3 + 105 A b^3 e^4 x^3 + 84 B b^3 d^2 e^2 x^2 + 189 B a b^2 d e^3 x^2 + 63 A b^3 d e^3 x^2 + 252 B a^2 b e^4 x^2 + 252 A a b^2 e^4 x^2 + 28 B b^3 d^3 e x + 63 B a a b^2 d^2 e^2 x + 21 A b^3 d^2 e^2 x + 84 B a^2 b d e^3 x + 84 A a a b^2 d^2 e^3 x + 70 B a a^3 e^4 x + 210 A a a^2 b e^4 x + 4 B b^3 d^4 + 9 B a a b^2 d^3 e + 3 A a b^3 d^3 e + 12 B a^2 b d^2 e^2 + 12 A a a b^2 d^2 e^2 + 10 B a a^3 d e^3 + 30 A a a^2 b d e^3 + 60 A a a^3 e^4}{420 e^5} + \frac{x(10 B a^3 e^3 + 12 B a^2 b d e^2 + 12 B a b^2 d^2 e + 10 B a^3 d^2 e^2 + 4 B b^3 d^4 + 3 A b^3 d^3 e)}{d^7 + 7 d^6 e x + 21 d^5 e^2 x^2 + \dots}$$

input

```
integrate((b*x+a)^3*(B*x+A)/(e*x+d)^8,x, algorithm="giac")
```

output

```
-1/420*(140*B*b^3*e^4*x^4 + 140*B*b^3*d*e^3*x^3 + 315*B*a*b^2*e^4*x^3 + 10
5*A*b^3*e^4*x^3 + 84*B*b^3*d^2*e^2*x^2 + 189*B*a*b^2*d*e^3*x^2 + 63*A*b^3*
d*e^3*x^2 + 252*B*a^2*b*e^4*x^2 + 252*A*a*b^2*e^4*x^2 + 28*B*b^3*d^3*e*x +
63*B*a*b^2*d^2*e^2*x + 21*A*b^3*d^2*e^2*x + 84*B*a^2*b*d*e^3*x + 84*A*a*b
^2*d*e^3*x + 70*B*a^3*e^4*x + 210*A*a^2*b*e^4*x + 4*B*b^3*d^4 + 9*B*a*b^2*
d^3*e + 3*A*b^3*d^3*e + 12*B*a^2*b*d^2*e^2 + 12*A*a*b^2*d^2*e^2 + 10*B*a^3
*d*e^3 + 30*A*a^2*b*d*e^3 + 60*A*a^3*e^4)/((e*x + d)^7*e^5)
```

Mupad [B] (verification not implemented)

Time = 0.90 (sec) , antiderivative size = 336, normalized size of antiderivative = 2.06

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^8} dx = \frac{10 B a^3 d e^3 + 60 A a^3 e^4 + 12 B a^2 b d^2 e^2 + 30 A a^2 b d e^3 + 9 B a b^2 d^3 e + 12 A a b^2 d^2 e^2 + 4 B b^3 d^4 + 3 A b^3 d^3 e}{420 e^5} + \frac{x(10 B a^3 e^3 + 12 B a^2 b d e^2 + 12 B a b^2 d^2 e + 10 B a^3 d^2 e^2 + 4 B b^3 d^4 + 3 A b^3 d^3 e)}{d^7 + 7 d^6 e x + 21 d^5 e^2 x^2 + \dots}$$

input `int(((A + B*x)*(a + b*x)^3)/(d + e*x)^8,x)`

output
$$-\frac{(60Aa^3e^4 + 4Bb^3d^4 + 3Ab^3d^3e + 10Ba^3de^3 + 12Aab^2d^2e^2 + 12Ba^2bd^2e^2 + 30Aa^2bde^3 + 9Bab^2d^3e)/(420e^5) + (x(10Ba^3e^3 + 4Bb^3d^3 + 30Aa^2bde^3 + 3Ab^3d^2e + 12Aab^2de^2 + 9Bab^2d^2e + 12Ba^2bde^2))/(60e^4) + (b^2x^3(3Ab^2e + 9Bab^2e + 4Bb^2d))/(12e^2) + (bx^2(12Ba^2e^2 + 4Bb^2d^2 + 12Aab^2e^2 + 3Ab^2de + 9Bab^2de))/(20e^3) + (Bb^3x^4)/(3e)}{(d^7 + e^7x^7 + 7de^6x^6 + 21d^5e^2x^2 + 35d^4e^3x^3 + 35d^3e^4x^4 + 21d^2e^5x^5 + 7d^6ex)}$$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.54

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^8} dx$$

$$= \frac{-35b^4e^4x^4 - 105ab^3e^4x^3 - 35b^4de^3x^3 - 126a^2b^2e^4x^2 - 63ab^3de^3x^2 - 21b^4d^2e^2x^2 - 70a^3be^4x - 42a^2b^2e^4}{105e^5(e^7x^7 + 7de^6x^6 + 21d^2e^5x^5 + 35d^3e^4x^4 + 35d^4e^3e^2x^3 + 21d^5e^2x^2 + 7d^6ex + d^7)}$$

input `int((b*x+a)^3*(B*x+A)/(e*x+d)^8,x)`

output
$$\frac{(-15a^4e^4 - 10a^3bde^3 - 70a^3b^2e^4x - 6a^2b^2d^2e^3 - 42a^2b^2de^3x - 126a^2b^2e^4x^2 - 3ab^3d^3e - 21ab^3d^2e^2x - 63ab^3de^3x^2 - 105ab^3e^4x^3 - b^4d^4 - 7b^4d^3ex - 21b^4d^2e^2x^2 - 35b^4de^3x^3 - 35b^4e^4x^4)/(105e^5(d^7 + 7d^6ex + 21d^5e^2x^2 + 35d^4e^3x^3 + 35d^3e^4x^4 + 21d^2e^5x^5 + 7d^6ex + e^7x^7))$$

3.39 $\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^9} dx$

Optimal result	392
Mathematica [A] (verified)	393
Rubi [A] (verified)	393
Maple [A] (verified)	395
Fricas [B] (verification not implemented)	395
Sympy [F(-1)]	396
Maxima [B] (verification not implemented)	396
Giac [A] (verification not implemented)	397
Mupad [B] (verification not implemented)	397
Reduce [B] (verification not implemented)	398

Optimal result

Integrand size = 20, antiderivative size = 163

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^9} dx = -\frac{(bd-ae)^3(Bd-Ae)}{8e^5(d+ex)^8} + \frac{(bd-ae)^2(4bBd-3Abe-aBe)}{7e^5(d+ex)^7} - \frac{b(bd-ae)(2bBd-Abe-aBe)}{2e^5(d+ex)^6} + \frac{b^2(4bBd-Abe-3aBe)}{5e^5(d+ex)^5} - \frac{b^3B}{4e^5(d+ex)^4}$$

output

```
-1/8*(-a*e+b*d)^3*(-A*e+B*d)/e^5/(e*x+d)^8+1/7*(-a*e+b*d)^2*(-3*A*b*e-B*a*
e+4*B*b*d)/e^5/(e*x+d)^7-1/2*b*(-a*e+b*d)*(-A*b*e-B*a*e+2*B*b*d)/e^5/(e*x+
d)^6+1/5*b^2*(-A*b*e-3*B*a*e+4*B*b*d)/e^5/(e*x+d)^5-1/4*b^3*B/e^5/(e*x+d)^
4
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.29

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^9} dx = \frac{5a^3e^3(7Ae + B(d + 8ex)) + 5a^2be^2(3Ae(d + 8ex) + B(d^2 + 8dex + 28e^2x^2)) + ab^2e(5Ae(d^2 + 8dex -$$

input `Integrate[((a + b*x)^3*(A + B*x))/(d + e*x)^9,x]`

output
$$\frac{-1/280*(5*a^3*e^3*(7*A*e + B*(d + 8*e*x)) + 5*a^2*b*e^2*(3*A*e*(d + 8*e*x) + B*(d^2 + 8*d*e*x + 28*e^2*x^2)) + a*b^2*e*(5*A*e*(d^2 + 8*d*e*x + 28*e^2*x^2) + 3*B*(d^3 + 8*d^2*e*x + 28*d*e^2*x^2 + 56*e^3*x^3)) + b^3*(A*e*(d^3 + 8*d^2*e*x + 28*d*e^2*x^2 + 56*e^3*x^3) + B*(d^4 + 8*d^3*e*x + 28*d^2*e^2*x^2 + 56*d*e^3*x^3 + 70*e^4*x^4))}{e^5*(d + e*x)^8}$$

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^9} dx$$

↓ 86

$$\int \left(\frac{b^2(3aBe + Abe - 4bBd)}{e^4(d + ex)^6} - \frac{3b(bd - ae)(aBe + Abe - 2bBd)}{e^4(d + ex)^7} + \frac{(ae - bd)^2(aBe + 3Abe - 4bBd)}{e^4(d + ex)^8} + \frac{(ae - bd)(aBe + 3Abe - 4bBd)}{e^4(d + ex)^9} \right) dx$$

↓ 2009

$$\frac{b^2(-3aBe - Abe + 4bBd)}{5e^5(d+ex)^5} - \frac{b(bd - ae)(-aBe - Abe + 2bBd)}{2e^5(d+ex)^6} + \frac{(bd - ae)^2(-aBe - 3Abe + 4bBd)}{7e^5(d+ex)^7} - \frac{(bd - ae)^3(Bd - Ae)}{8e^5(d+ex)^8} - \frac{b^3B}{4e^5(d+ex)^4}$$

input `Int[((a + b*x)^3*(A + B*x))/(d + e*x)^9,x]`

output `-1/8*((b*d - a*e)^3*(B*d - A*e))/(e^5*(d + e*x)^8) + ((b*d - a*e)^2*(4*b*B*d - 3*A*b*e - a*B*e))/(7*e^5*(d + e*x)^7) - (b*(b*d - a*e)*(2*b*B*d - A*b*e - a*B*e))/(2*e^5*(d + e*x)^6) + (b^2*(4*b*B*d - A*b*e - 3*a*B*e))/(5*e^5*(d + e*x)^5) - (b^3*B)/(4*e^5*(d + e*x)^4)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

output

```
-1/280*(70*B*b^3*e^4*x^4 + B*b^3*d^4 + 35*A*a^3*e^4 + (3*B*a*b^2 + A*b^3)*
d^3*e + 5*(B*a^2*b + A*a*b^2)*d^2*e^2 + 5*(B*a^3 + 3*A*a^2*b)*d*e^3 + 56*(
B*b^3*d*e^3 + (3*B*a*b^2 + A*b^3)*e^4)*x^3 + 28*(B*b^3*d^2*e^2 + (3*B*a*b^
2 + A*b^3)*d*e^3 + 5*(B*a^2*b + A*a*b^2)*e^4)*x^2 + 8*(B*b^3*d^3*e + (3*B*
a*b^2 + A*b^3)*d^2*e^2 + 5*(B*a^2*b + A*a*b^2)*d*e^3 + 5*(B*a^3 + 3*A*a^2*
b)*e^4)*x)/(e^13*x^8 + 8*d*e^12*x^7 + 28*d^2*e^11*x^6 + 56*d^3*e^10*x^5 +
70*d^4*e^9*x^4 + 56*d^5*e^8*x^3 + 28*d^6*e^7*x^2 + 8*d^7*e^6*x + d^8*e^5)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^9} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**3*(B*x+A)/(e*x+d)**9,x)
```

output

Timed out

Maxima [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 335 vs. $2(153) = 306$.

Time = 0.05 (sec) , antiderivative size = 335, normalized size of antiderivative = 2.06

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^9} dx = \frac{70 Bb^3e^4x^4 + Bb^3d^4 + 35 Aa^3e^4 + (3 Bab^2 + Ab^3)d^3e + 5 (Ba^2b + Aab^2)d^2e^2 + 5 (Ba^3 + 3 Aa^2b)de^3 + \dots}{280 (e^{13}x^8 + \dots)}$$

input

```
integrate((b*x+a)^3*(B*x+A)/(e*x+d)^9,x, algorithm="maxima")
```

output

```
-1/280*(70*B*b^3*e^4*x^4 + B*b^3*d^4 + 35*A*a^3*e^4 + (3*B*a*b^2 + A*b^3)*
d^3*e + 5*(B*a^2*b + A*a*b^2)*d^2*e^2 + 5*(B*a^3 + 3*A*a^2*b)*d*e^3 + 56*(
B*b^3*d*e^3 + (3*B*a*b^2 + A*b^3)*e^4)*x^3 + 28*(B*b^3*d^2*e^2 + (3*B*a*b^
2 + A*b^3)*d*e^3 + 5*(B*a^2*b + A*a*b^2)*e^4)*x^2 + 8*(B*b^3*d^3*e + (3*B*
a*b^2 + A*b^3)*d^2*e^2 + 5*(B*a^2*b + A*a*b^2)*d*e^3 + 5*(B*a^3 + 3*A*a^2*
b)*e^4)*x)/(e^13*x^8 + 8*d*e^12*x^7 + 28*d^2*e^11*x^6 + 56*d^3*e^10*x^5 +
70*d^4*e^9*x^4 + 56*d^5*e^8*x^3 + 28*d^6*e^7*x^2 + 8*d^7*e^6*x + d^8*e^5)
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 298, normalized size of antiderivative = 1.83

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^9} dx = \frac{70 B b^3 e^4 x^4 + 56 B b^3 d e^3 x^3 + 168 B a b^2 e^4 x^3 + 56 A b^3 e^4 x^3 + 28 B b^3 d^2 e^2 x^2 + 84 B a b^2 d e^3 x^2 + 28 A b^3 d e^3 x^2}{(d+ex)^9}$$

input

```
integrate((b*x+a)^3*(B*x+A)/(e*x+d)^9,x, algorithm="giac")
```

output

```
-1/280*(70*B*b^3*e^4*x^4 + 56*B*b^3*d*e^3*x^3 + 168*B*a*b^2*e^4*x^3 + 56*A
*b^3*e^4*x^3 + 28*B*b^3*d^2*e^2*x^2 + 84*B*a*b^2*d*e^3*x^2 + 28*A*b^3*d*e^
3*x^2 + 140*B*a^2*b*e^4*x^2 + 140*A*a*b^2*e^4*x^2 + 8*B*b^3*d^3*e*x + 24*B
*a*b^2*d^2*e^2*x + 8*A*b^3*d^2*e^2*x + 40*B*a^2*b*d*e^3*x + 40*A*a*b^2*d*e
^3*x + 40*B*a^3*e^4*x + 120*A*a^2*b*e^4*x + B*b^3*d^4 + 3*B*a*b^2*d^3*e +
A*b^3*d^3*e + 5*B*a^2*b*d^2*e^2 + 5*A*a*b^2*d^2*e^2 + 5*B*a^3*d*e^3 + 15*A
*a^2*b*d*e^3 + 35*A*a^3*e^4)/((e*x + d)^8*e^5)
```

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 339, normalized size of antiderivative = 2.08

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^9} dx = \frac{5 B a^3 d e^3 + 35 A a^3 e^4 + 5 B a^2 b d^2 e^2 + 15 A a^2 b d e^3 + 3 B a b^2 d^3 e + 5 A a b^2 d^2 e^2 + B b^3 d^4 + A b^3 d^3 e}{280 e^5} + \frac{x(5 B a^3 e^3 + 5 B a^2 b d e^2 + 15 A a^2 b d e^3 + 5 A a b^2 d^2 e^2 + 5 B a^3 d e^3 + 5 A a^2 b d e^3 + 35 A a^3 e^4)}{(d+ex)^8}$$

input `int(((A + B*x)*(a + b*x)^3)/(d + e*x)^9,x)`

output
$$-\frac{((35Aa^3e^4 + Bb^3d^4 + Ab^3d^3e + 5Ba^3de^3 + 5Aab^2d^2e^2 + 5Bba^2bd^2e^2 + 15Aa^2bde^3 + 3Bba^2bd^3e)/(280e^5) + (x(5Bba^3e^3 + Bb^3d^3 + 15Aa^2bde^3 + Ab^3d^2e + 5Aab^2de^2 + 3Bba^2bd^2e + 5Bba^2bde^2)))/(35e^4) + (b^2x^3(Abe + 3Bbae + Bbd))/(5e^2) + (bx^2(5Bba^2e^2 + Bb^2d^2 + 5Aab^2e^2 + Ab^2de + 3Bba^2bde))/(10e^3) + (Bb^3x^4)/(4e))/(d^8 + e^8x^8 + 8de^7x^7 + 28d^6e^2x^2 + 56d^5e^3x^3 + 70d^4e^4x^4 + 56d^3e^5x^5 + 28d^2e^6x^6 + 8d^7e^7x^7)}$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 262, normalized size of antiderivative = 1.61

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^9} dx = \frac{-70b^4e^4x^4 - 224ab^3e^4x^3 - 56b^4de^3x^3 - 280a^2b^2e^4x^2 - 112ab^3de^3x^2 - 28b^4d^2e^2x^2 - 160a^3be^4x - 80a^4}{280e^5(e^8x^8 + 8de^7x^7 + 28d^2e^6x^6 + 56d^3e^5x^5 + 70d^4e^4x^4)}$$

input `int((b*x+a)^3*(B*x+A)/(e*x+d)^9,x)`

output
$$\frac{(-35a^4e^4 - 20a^3bde^3 - 160a^3b^2e^4x - 10a^2b^2d^2e^2 - 80a^2b^2d^2e^3x - 280a^2b^2e^4x^2 - 4ab^3d^3e - 32ab^3d^2e^2x - 112ab^3de^3x^2 - 224ab^3e^4x^3 - b^4d^4 - 8b^4d^3ex - 28b^4d^2e^2x^2 - 56b^4de^3x^3 - 70b^4e^4x^4)/(280e^5(d^8 + 8d^7ex + 28d^6e^2x^2 + 56d^5e^3x^3 + 70d^4e^4x^4 + 56d^3e^5x^5 + 28d^2e^6x^6 + 8d^7e^7x^7 + e^8x^8))}$$

3.40 $\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{10}} dx$

Optimal result	399
Mathematica [A] (verified)	400
Rubi [A] (verified)	400
Maple [A] (verified)	402
Fricas [B] (verification not implemented)	402
Sympy [F(-1)]	403
Maxima [B] (verification not implemented)	403
Giac [A] (verification not implemented)	404
Mupad [B] (verification not implemented)	404
Reduce [B] (verification not implemented)	405

Optimal result

Integrand size = 20, antiderivative size = 163

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{10}} dx = -\frac{(bd-ae)^3(Bd-Ae)}{9e^5(d+ex)^9} + \frac{(bd-ae)^2(4bBd-3Abe-aBe)}{8e^5(d+ex)^8} - \frac{3b(bd-ae)(2bBd-Abe-aBe)}{7e^5(d+ex)^7} + \frac{b^2(4bBd-Abe-3aBe)}{6e^5(d+ex)^6} - \frac{b^3B}{5e^5(d+ex)^5}$$

output

```
-1/9*(-a*e+b*d)^3*(-A*e+B*d)/e^5/(e*x+d)^9+1/8*(-a*e+b*d)^2*(-3*A*b*e-B*a*
e+4*B*b*d)/e^5/(e*x+d)^8-3/7*b*(-a*e+b*d)*(-A*b*e-B*a*e+2*B*b*d)/e^5/(e*x+
d)^7+1/6*b^2*(-A*b*e-3*B*a*e+4*B*b*d)/e^5/(e*x+d)^6-1/5*b^3*B/e^5/(e*x+d)^
5
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.31

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{10}} dx =$$

$$\frac{35a^3e^3(8Ae + B(d + 9ex)) + 15a^2be^2(7Ae(d + 9ex) + 2B(d^2 + 9dex + 36e^2x^2)) + 15ab^2e(2Ae(d^2 + 9dex + 36e^2x^2)) + 5a^3b^3(8Ae + B(d + 9ex)) + 15a^2b^2e(7Ae(d + 9ex) + 2B(d^2 + 9dex + 36e^2x^2)) + 15ab^3e(2Ae(d^2 + 9dex + 36e^2x^2)) + 5a^3b^3e(8Ae + B(d + 9ex))}{e^5(d + ex)^9}$$

input `Integrate[((a + b*x)^3*(A + B*x))/(d + e*x)^10,x]`

output

```
-1/2520*(35*a^3*e^3*(8*A*e + B*(d + 9*e*x)) + 15*a^2*b*e^2*(7*A*e*(d + 9*e*x) + 2*B*(d^2 + 9*d*e*x + 36*e^2*x^2)) + 15*a*b^2*e*(2*A*e*(d^2 + 9*d*e*x + 36*e^2*x^2) + B*(d^3 + 9*d^2*e*x + 36*d*e^2*x^2 + 84*e^3*x^3)) + b^3*(5*A*e*(d^3 + 9*d^2*e*x + 36*d*e^2*x^2 + 84*e^3*x^3) + 4*B*(d^4 + 9*d^3*e*x + 36*d^2*e^2*x^2 + 84*d*e^3*x^3 + 126*e^4*x^4)))/(e^5*(d + e*x)^9)
```

Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{10}} dx$$

$$\downarrow 86$$

$$\int \left(\frac{b^2(3aBe + Abe - 4bBd)}{e^4(d + ex)^7} - \frac{3b(bd - ae)(aBe + Abe - 2bBd)}{e^4(d + ex)^8} + \frac{(ae - bd)^2(aBe + 3Abe - 4bBd)}{e^4(d + ex)^9} + \frac{(ae - bd)(aBe + 3Abe - 4bBd)}{e^4(d + ex)^{10}} \right) dx$$

$$\downarrow 2009$$

$$\frac{b^2(-3aBe - Abe + 4bBd)}{6e^5(d+ex)^6} - \frac{3b(bd - ae)(-aBe - Abe + 2bBd)}{7e^5(d+ex)^7} + \frac{(bd - ae)^2(-aBe - 3Abe + 4bBd)}{8e^5(d+ex)^8} - \frac{(bd - ae)^3(Bd - Ae)}{9e^5(d+ex)^9} - \frac{b^3B}{5e^5(d+ex)^5}$$

input `Int[((a + b*x)^3*(A + B*x))/(d + e*x)^10,x]`

output `-1/9*((b*d - a*e)^3*(B*d - A*e))/(e^5*(d + e*x)^9) + ((b*d - a*e)^2*(4*b*B*d - 3*A*b*e - a*B*e))/(8*e^5*(d + e*x)^8) - (3*b*(b*d - a*e)*(2*b*B*d - A*b*e - a*B*e))/(7*e^5*(d + e*x)^7) + (b^2*(4*b*B*d - A*b*e - 3*a*B*e))/(6*e^5*(d + e*x)^6) - (b^3*B)/(5*e^5*(d + e*x)^5)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.66

method	result
risch	$\frac{-\frac{b^3 B x^4}{5e} - \frac{b^2(5Abe+15Bae+4Bbd)x^3}{30e^2} - \frac{b(30Aab e^2+5A b^2 de+30B a^2 e^2+15B abde+4b^2 B d^2)x^2}{70e^3} - \frac{(105A a^2 b e^3+30Aa b^2 d e^2+5A b^3 d^2 e^2)}{8e^5(e^2 x^2+d^2)}$
default	$-\frac{b^2(Abe+3Bae-4Bbd)}{6e^5(e^2 x^2+d^2)^6} - \frac{3b(Aab e^2-A b^2 de+B a^2 e^2-3B abde+2b^2 B d^2)}{7e^5(e^2 x^2+d^2)^7} - \frac{3A a^2 b e^3-6Aa b^2 d e^2+3A b^3 d^2 e+B a^3 e^3}{8e^5(e^2 x^2+d^2)}$
gospers	$-\frac{504B x^4 b^3 e^4+420A x^3 b^3 e^4+1260B x^3 a b^2 e^4+336B x^3 b^3 d e^3+1080A x^2 a b^2 e^4+180A x^2 b^3 d e^3+1080B x^2 a^2 b e^4+540B x^2 b^3 d^2 e^3}{2520e^4(e^2 x^2+d^2)^6}$
orering	$-\frac{504B x^4 b^3 e^4+420A x^3 b^3 e^4+1260B x^3 a b^2 e^4+336B x^3 b^3 d e^3+1080A x^2 a b^2 e^4+180A x^2 b^3 d e^3+1080B x^2 a^2 b e^4+540B x^2 b^3 d^2 e^3}{2520e^4(e^2 x^2+d^2)^6}$
norman	$-\frac{b^3 B x^4}{5e} - \frac{(5A b^3 e^5+15B a b^2 e^5+4b^3 B d e^4)x^3}{30e^6} - \frac{(30a b^2 A e^6+5A b^3 d e^5+30a^2 b B e^6+15B a b^2 d e^5+4B b^3 d^2 e^4)x^2}{70e^7} - \frac{(105A a^2 b e^7+30A a b^2 d e^6+5A b^3 d^2 e^5)}{8e^5(e^2 x^2+d^2)}$
paralelrisch	$-\frac{504B b^3 x^4 e^8+420A b^3 e^8 x^3+1260B a b^2 e^8 x^3+336B b^3 d e^7 x^3+1080A a b^2 e^8 x^2+180A b^3 d e^7 x^2+1080B a^2 b e^8 x^2+540B a b^2 d e^7 x^2}{2520e^4(e^2 x^2+d^2)^6}$

```
input int((b*x+a)^3*(B*x+A)/(e*x+d)^10,x,method=_RETURNVERBOSE)
```

```
output (-1/5*b^3*B/e*x^4-1/30*b^2/e^2*(5*A*b*e+15*B*a*e+4*B*b*d)*x^3-1/70*b/e^3*(30*A*a*b*e^2+5*A*b^2*d*e+30*B*a^2*e^2+15*B*a*b*d*e+4*B*b^2*d^2)*x^2-1/280/e^4*(105*A*a^2*b*e^3+30*A*a*b^2*d*e^2+5*A*b^3*d^2*e+35*B*a^3*e^3+30*B*a^2*b*d*e^2+15*B*a*b^2*d^2*e+4*B*b^3*d^3)*x-1/2520/e^5*(280*A*a^3*e^4+105*A*a^2*b*d*e^3+30*A*a*b^2*d^2*e^2+5*A*b^3*d^3*e+35*B*a^3*d*e^3+30*B*a^2*b*d^2*e^2+15*B*a*b^2*d^3*e+4*B*b^3*d^4))/(e*x+d)^9
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 354 vs. 2(153) = 306.

Time = 0.09 (sec) , antiderivative size = 354, normalized size of antiderivative = 2.17

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{10}} dx = \frac{504 B b^3 e^4 x^4 + 4 B b^3 d^4 + 280 A a^3 e^4 + 5 (3 B a b^2 + A b^3) d^3 e + 30 (B a^2 b + A a b^2) d^2 e^2 + 35 (B a^3 + 3 A a b^2) d e^3 + 5 A^2 b^3}{2520 (e^{14} x^4 + \dots)}$$

```
input integrate((b*x+a)^3*(B*x+A)/(e*x+d)^10,x, algorithm="fricas")
```

output

```
-1/2520*(504*B*b^3*e^4*x^4 + 4*B*b^3*d^4 + 280*A*a^3*e^4 + 5*(3*B*a*b^2 +
A*b^3)*d^3*e + 30*(B*a^2*b + A*a*b^2)*d^2*e^2 + 35*(B*a^3 + 3*A*a^2*b)*d*e
^3 + 84*(4*B*b^3*d*e^3 + 5*(3*B*a*b^2 + A*b^3)*e^4)*x^3 + 36*(4*B*b^3*d^2*
e^2 + 5*(3*B*a*b^2 + A*b^3)*d*e^3 + 30*(B*a^2*b + A*a*b^2)*e^4)*x^2 + 9*(4
*B*b^3*d^3*e + 5*(3*B*a*b^2 + A*b^3)*d^2*e^2 + 30*(B*a^2*b + A*a*b^2)*d*e^
3 + 35*(B*a^3 + 3*A*a^2*b)*e^4)*x)/(e^14*x^9 + 9*d*e^13*x^8 + 36*d^2*e^12*
x^7 + 84*d^3*e^11*x^6 + 126*d^4*e^10*x^5 + 126*d^5*e^9*x^4 + 84*d^6*e^8*x^
3 + 36*d^7*e^7*x^2 + 9*d^8*e^6*x + d^9*e^5)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{10}} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**3*(B*x+A)/(e*x+d)**10,x)
```

output

Timed out

Maxima [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 354 vs. $2(153) = 306$.

Time = 0.05 (sec) , antiderivative size = 354, normalized size of antiderivative = 2.17

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{10}} dx = \frac{504 B b^3 e^4 x^4 + 4 B b^3 d^4 + 280 A a^3 e^4 + 5 (3 B a b^2 + A b^3) d^3 e + 30 (B a^2 b + A a b^2) d^2 e^2 + 35 (B a^3 + 3 A a^2 b) d e^3 + 84 (4 B b^3 d e^3 + 5 (3 B a b^2 + A b^3) e^4) x^3 + 36 (4 B b^3 d^2 e^2 + 5 (3 B a b^2 + A b^3) d e^3 + 30 (B a^2 b + A a b^2) e^4) x^2 + 9 (4 B b^3 d^3 e + 5 (3 B a b^2 + A b^3) d^2 e^2 + 30 (B a^2 b + A a b^2) d e^3 + 35 (B a^3 + 3 A a^2 b) e^4) x}{2520 (e^{14} x^9 + 9 d e^{13} x^8 + 36 d^2 e^{12} x^7 + 84 d^3 e^{11} x^6 + 126 d^4 e^{10} x^5 + 126 d^5 e^9 x^4 + 84 d^6 e^8 x^3 + 36 d^7 e^7 x^2 + 9 d^8 e^6 x + d^9 e^5)}$$

input

```
integrate((b*x+a)^3*(B*x+A)/(e*x+d)^10,x, algorithm="maxima")
```


output

```
-1/2520*(504*B*b^3*e^4*x^4 + 4*B*b^3*d^4 + 280*A*a^3*e^4 + 5*(3*B*a*b^2 +
A*b^3)*d^3*e + 30*(B*a^2*b + A*a*b^2)*d^2*e^2 + 35*(B*a^3 + 3*A*a^2*b)*d*e
^3 + 84*(4*B*b^3*d*e^3 + 5*(3*B*a*b^2 + A*b^3)*e^4)*x^3 + 36*(4*B*b^3*d^2*
e^2 + 5*(3*B*a*b^2 + A*b^3)*d*e^3 + 30*(B*a^2*b + A*a*b^2)*e^4)*x^2 + 9*(4
*B*b^3*d^3*e + 5*(3*B*a*b^2 + A*b^3)*d^2*e^2 + 30*(B*a^2*b + A*a*b^2)*d*e^
3 + 35*(B*a^3 + 3*A*a^2*b)*e^4)*x)/(e^14*x^9 + 9*d*e^13*x^8 + 36*d^2*e^12*
x^7 + 84*d^3*e^11*x^6 + 126*d^4*e^10*x^5 + 126*d^5*e^9*x^4 + 84*d^6*e^8*x^
3 + 36*d^7*e^7*x^2 + 9*d^8*e^6*x + d^9*e^5)
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 300, normalized size of antiderivative = 1.84

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{10}} dx = \frac{504 B b^3 e^4 x^4 + 336 B b^3 d e^3 x^3 + 1260 B a b^2 e^4 x^3 + 420 A b^3 e^4 x^3 + 144 B b^3 d^2 e^2 x^2 + 540 B a b^2 d e^3 x^2 + 180 A a b^3 d e^3 x^2 + 1080 B a^2 b^2 e^4 x^2 + 1080 A a a b^2 e^4 x^2 + 36 B b^3 d^3 e^2 x + 135 B a a b^2 d^2 e^2 x + 45 A a b^3 d^2 e^2 x + 270 B a^2 b d e^3 x + 270 A a a b^2 d e^3 x + 315 B a^3 e^4 x + 945 A a a^2 b e^4 x + 4 B b^3 d^4 + 15 B a a b^2 d^3 e + 5 A a b^3 d^3 e + 30 B a^2 b d^2 e^2 + 30 A a a b^2 d^2 e^2 + 35 B a^3 d e^3 + 105 A a a^2 b d e^3 + 280 A a a^3 e^4}{2520 e^5} + \frac{x(35 B a^3 e^3 + 30 B a^2 b e^3 + 20 B a b^2 e^3 + 15 B b^3 e^3)}{d^9 + 9 d^8 e x + 36 d^7 e^2 x^2 + 84 d^6 e^3 x^3 + 36 d^5 e^4 x^4 + 9 d^4 e^5 x^5 + 36 d^3 e^6 x^6 + 126 d^2 e^7 x^7 + 126 d e^8 x^8 + e^9 x^9}$$

input

```
integrate((b*x+a)^3*(B*x+A)/(e*x+d)^10,x, algorithm="giac")
```

output

```
-1/2520*(504*B*b^3*e^4*x^4 + 336*B*b^3*d*e^3*x^3 + 1260*B*a*b^2*e^4*x^3 +
420*A*b^3*e^4*x^3 + 144*B*b^3*d^2*e^2*x^2 + 540*B*a*b^2*d*e^3*x^2 + 180*A*
b^3*d*e^3*x^2 + 1080*B*a^2*b*e^4*x^2 + 1080*A*a*b^2*e^4*x^2 + 36*B*b^3*d^3
*e*x + 135*B*a*b^2*d^2*e^2*x + 45*A*b^3*d^2*e^2*x + 270*B*a^2*b*d*e^3*x +
270*A*a*b^2*d*e^3*x + 315*B*a^3*e^4*x + 945*A*a^2*b*e^4*x + 4*B*b^3*d^4 +
15*B*a*b^2*d^3*e + 5*A*b^3*d^3*e + 30*B*a^2*b*d^2*e^2 + 30*A*a*b^2*d^2*e^2
+ 35*B*a^3*d*e^3 + 105*A*a^2*b*d*e^3 + 280*A*a^3*e^4)/((e*x + d)^9*e^5)
```

Mupad [B] (verification not implemented)

Time = 0.95 (sec) , antiderivative size = 358, normalized size of antiderivative = 2.20

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{10}} dx = \frac{35 B a^3 d e^3 + 280 A a^3 e^4 + 30 B a^2 b d^2 e^2 + 105 A a^2 b d e^3 + 15 B a b^2 d^3 e + 30 A a b^2 d^2 e^2 + 4 B b^3 d^4 + 5 A b^3 d^3 e}{2520 e^5} + \frac{x(35 B a^3 e^3 + 30 B a^2 b e^3 + 20 B a b^2 e^3 + 15 B b^3 e^3)}{d^9 + 9 d^8 e x + 36 d^7 e^2 x^2 + 84 d^6 e^3 x^3 + 36 d^5 e^4 x^4 + 9 d^4 e^5 x^5 + 36 d^3 e^6 x^6 + 126 d^2 e^7 x^7 + 126 d e^8 x^8 + e^9 x^9}$$

input `int(((A + B*x)*(a + b*x)^3)/(d + e*x)^10,x)`

output
$$-\frac{((280*A*a^3*e^4 + 4*B*b^3*d^4 + 5*A*b^3*d^3*e + 35*B*a^3*d*e^3 + 30*A*a*b^2*d^2*e^2 + 30*B*a^2*b*d^2*e^2 + 105*A*a^2*b*d*e^3 + 15*B*a*b^2*d^3*e)/(2520*e^5) + (x*(35*B*a^3*e^3 + 4*B*b^3*d^3 + 105*A*a^2*b*e^3 + 5*A*b^3*d^2*e + 30*A*a*b^2*d*e^2 + 15*B*a*b^2*d^2*e + 30*B*a^2*b*d*e^2))/(280*e^4) + (b^2*x^3*(5*A*b*e + 15*B*a*e + 4*B*b*d))/(30*e^2) + (b*x^2*(30*B*a^2*e^2 + 4*B*b^2*d^2 + 30*A*a*b*e^2 + 5*A*b^2*d*e + 15*B*a*b*d*e))/(70*e^3) + (B*b^3*x^4)/(5*e))/(d^9 + e^9*x^9 + 9*d*e^8*x^8 + 36*d^7*e^2*x^2 + 84*d^6*e^3*x^3 + 126*d^5*e^4*x^4 + 126*d^4*e^5*x^5 + 84*d^3*e^6*x^6 + 36*d^2*e^7*x^7 + 9*d^8*e*x)$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.67

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{10}} dx = \frac{-126b^4e^4x^4 - 420ab^3e^4x^3 - 84b^4de^3x^3 - 540a^2b^2e^4x^2 - 180ab^3de^3x^2 - 36b^4d^2e^2x^2 - 315a^3be^4x - 135a^4e^4}{630e^5(e^9x^9 + 9de^8x^8 + 36d^2e^7x^7 + 84d^3e^6x^6 + 126d^4e^5x^5 + 126d^5e^4x^4 + 84d^6e^3x^3 + 36d^7e^2x^2 + 9d^8ex + d^9)}$$

input `int((b*x+a)^3*(B*x+A)/(e*x+d)^10,x)`

output
$$(-70*a**4*e**4 - 35*a**3*b*d*e**3 - 315*a**3*b*e**4*x - 15*a**2*b**2*d**2*e**2 - 135*a**2*b**2*d*e**3*x - 540*a**2*b**2*e**4*x**2 - 5*a*b**3*d**3*e - 45*a*b**3*d**2*e**2*x - 180*a*b**3*d*e**3*x**2 - 420*a*b**3*e**4*x**3 - b**4*d**4 - 9*b**4*d**3*e*x - 36*b**4*d**2*e**2*x**2 - 84*b**4*d*e**3*x**3 - 126*b**4*e**4*x**4)/(630*e**5*(d**9 + 9*d**8*e*x + 36*d**7*e**2*x**2 + 84*d**6*e**3*x**3 + 126*d**5*e**4*x**4 + 126*d**4*e**5*x**5 + 84*d**3*e**6*x**6 + 36*d**2*e**7*x**7 + 9*d*e**8*x**8 + e**9*x**9))$$

3.41 $\int (a + bx)^6 (A + Bx)(d + ex)^8 dx$

Optimal result	406
Mathematica [B] (verified)	407
Rubi [A] (verified)	408
Maple [B] (verified)	409
Fricas [B] (verification not implemented)	410
Sympy [B] (verification not implemented)	411
Maxima [B] (verification not implemented)	412
Giac [B] (verification not implemented)	413
Mupad [B] (verification not implemented)	414
Reduce [B] (verification not implemented)	415

Optimal result

Integrand size = 20, antiderivative size = 292

$$\begin{aligned}
 \int (a + bx)^6 (A + Bx)(d + ex)^8 dx = & -\frac{(bd - ae)^6 (Bd - Ae)(d + ex)^9}{9e^8} \\
 & + \frac{(bd - ae)^5 (7bBd - 6Abe - aBe)(d + ex)^{10}}{10e^8} \\
 & - \frac{3b(bd - ae)^4 (7bBd - 5Abe - 2aBe)(d + ex)^{11}}{11e^8} \\
 & + \frac{5b^2 (bd - ae)^3 (7bBd - 4Abe - 3aBe)(d + ex)^{12}}{12e^8} \\
 & - \frac{5b^3 (bd - ae)^2 (7bBd - 3Abe - 4aBe)(d + ex)^{13}}{13e^8} \\
 & + \frac{3b^4 (bd - ae) (7bBd - 2Abe - 5aBe)(d + ex)^{14}}{14e^8} \\
 & - \frac{b^5 (7bBd - Abe - 6aBe)(d + ex)^{15}}{15e^8} \\
 & + \frac{b^6 B (d + ex)^{16}}{16e^8}
 \end{aligned}$$

output

```
-1/9*(-a*e+b*d)^6*(-A*e+B*d)*(e*x+d)^9/e^8+1/10*(-a*e+b*d)^5*(-6*A*b*e-B*a
*e+7*B*b*d)*(e*x+d)^10/e^8-3/11*b*(-a*e+b*d)^4*(-5*A*b*e-2*B*a*e+7*B*b*d)*
(e*x+d)^11/e^8+5/12*b^2*(-a*e+b*d)^3*(-4*A*b*e-3*B*a*e+7*B*b*d)*(e*x+d)^12
/e^8-5/13*b^3*(-a*e+b*d)^2*(-3*A*b*e-4*B*a*e+7*B*b*d)*(e*x+d)^13/e^8+3/14*
b^4*(-a*e+b*d)*(-2*A*b*e-5*B*a*e+7*B*b*d)*(e*x+d)^14/e^8-1/15*b^5*(-A*b*e-
6*B*a*e+7*B*b*d)*(e*x+d)^15/e^8+1/16*b^6*B*(e*x+d)^16/e^8
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1385 vs. $2(292) = 584$.

Time = 0.33 (sec) , antiderivative size = 1385, normalized size of antiderivative = 4.74

$$\int (a + bx)^6 (A + Bx)(d + ex)^8 dx = \text{Too large to display}$$

input

```
Integrate[(a + b*x)^6*(A + B*x)*(d + e*x)^8,x]
```

output

```
a^6*A*d^8*x + (a^5*d^7*(6*A*b*d + a*B*d + 8*a*A*e)*x^2)/2 + (a^4*d^6*(2*a*
B*d*(3*b*d + 4*a*e) + A*(15*b^2*d^2 + 48*a*b*d*e + 28*a^2*e^2))*x^3)/3 + (
a^3*d^5*(a*B*d*(15*b^2*d^2 + 48*a*b*d*e + 28*a^2*e^2) + 4*A*(5*b^3*d^3 + 3
0*a*b^2*d^2*e + 42*a^2*b*d*e^2 + 14*a^3*e^3))*x^4)/4 + (a^2*d^4*(4*a*B*d*(
5*b^3*d^3 + 30*a*b^2*d^2*e + 42*a^2*b*d*e^2 + 14*a^3*e^3) + A*(15*b^4*d^4
+ 160*a*b^3*d^3*e + 420*a^2*b^2*d^2*e^2 + 336*a^3*b*d*e^3 + 70*a^4*e^4))*x
^5)/5 + (a*d^3*(a*B*d*(15*b^4*d^4 + 160*a*b^3*d^3*e + 420*a^2*b^2*d^2*e^2
+ 336*a^3*b*d*e^3 + 70*a^4*e^4) + 2*A*(3*b^5*d^5 + 60*a*b^4*d^4*e + 280*a^
2*b^3*d^3*e^2 + 420*a^3*b^2*d^2*e^3 + 210*a^4*b*d*e^4 + 28*a^5*e^5))*x^6)/
6 + (d^2*(2*a*B*d*(3*b^5*d^5 + 60*a*b^4*d^4*e + 280*a^2*b^3*d^3*e^2 + 420*
a^3*b^2*d^2*e^3 + 210*a^4*b*d*e^4 + 28*a^5*e^5) + A*(b^6*d^6 + 48*a*b^5*d^
5*e + 420*a^2*b^4*d^4*e^2 + 1120*a^3*b^3*d^3*e^3 + 1050*a^4*b^2*d^2*e^4 +
336*a^5*b*d*e^5 + 28*a^6*e^6))*x^7)/7 + (d*(168*a^5*b*d*e^5*(2*B*d + A*e)
+ 420*a^2*b^4*d^4*e^2*(B*d + 2*A*e) + 4*a^6*e^6*(7*B*d + 2*A*e) + 210*a^4*
b^2*d^2*e^4*(5*B*d + 4*A*e) + 280*a^3*b^3*d^3*e^3*(4*B*d + 5*A*e) + 24*a*b
^5*d^5*e*(2*B*d + 7*A*e) + b^6*d^6*(B*d + 8*A*e))*x^8)/8 + (e*(420*a^4*b^2
*d^2*e^4*(2*B*d + A*e) + a^6*e^6*(8*B*d + A*e) + 168*a*b^5*d^5*e*(B*d + 2*
A*e) + 24*a^5*b*d*e^5*(7*B*d + 2*A*e) + 280*a^3*b^3*d^3*e^3*(5*B*d + 4*A*
e) + 210*a^2*b^4*d^4*e^2*(4*B*d + 5*A*e) + 4*b^6*d^6*(2*B*d + 7*A*e))*x^9)/
9 + (e^2*(a^6*B*e^6 + 560*a^3*b^3*d^2*e^3*(2*B*d + A*e) + 6*a^5*b*e^5*(...
```

Rubi [A] (verified)

Time = 1.71 (sec) , antiderivative size = 292, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^6 (A + Bx)(d + ex)^8 dx$$

↓ 86

$$\int \left(\frac{b^5(d + ex)^{14}(6aBe + Abe - 7bBd)}{e^7} - \frac{3b^4(d + ex)^{13}(bd - ae)(5aBe + 2Abe - 7bBd)}{e^7} + \frac{5b^3(d + ex)^{12}(bd - ae)^2(-4aBe - 3Abe + 7bBd)}{e^7} - \frac{5b^2(d + ex)^{11}(bd - ae)^3(-3aBe - 4Abe + 7bBd)}{e^7} + \frac{3b(d + ex)^{10}(bd - ae)^4(-2aBe - 5Abe + 7bBd)}{e^7} + \frac{(d + ex)^9(bd - ae)^5(-aBe - 6Abe + 7bBd)}{e^7} - \frac{(d + ex)^8(bd - ae)^6(Bd - Ae)}{e^7} + \frac{b^6B(d + ex)^7}{e^7} \right) dx$$

↓ 2009

$$-\frac{b^5(d + ex)^{15}(-6aBe - Abe + 7bBd)}{15e^8} + \frac{3b^4(d + ex)^{14}(bd - ae)(-5aBe - 2Abe + 7bBd)}{14e^8} - \frac{5b^3(d + ex)^{13}(bd - ae)^2(-4aBe - 3Abe + 7bBd)}{13e^8} + \frac{5b^2(d + ex)^{12}(bd - ae)^3(-3aBe - 4Abe + 7bBd)}{12e^8} - \frac{3b(d + ex)^{11}(bd - ae)^4(-2aBe - 5Abe + 7bBd)}{11e^8} + \frac{(d + ex)^{10}(bd - ae)^5(-aBe - 6Abe + 7bBd)}{10e^8} - \frac{(d + ex)^9(bd - ae)^6(Bd - Ae)}{9e^8} + \frac{b^6B(d + ex)^{16}}{16e^8}$$

input `Int[(a + b*x)^6*(A + B*x)*(d + e*x)^8,x]`

output

```
-1/9*((b*d - a*e)^6*(B*d - A*e)*(d + e*x)^9)/e^8 + ((b*d - a*e)^5*(7*b*B*d - 6*A*b*e - a*B*e)*(d + e*x)^10)/(10*e^8) - (3*b*(b*d - a*e)^4*(7*b*B*d - 5*A*b*e - 2*a*B*e)*(d + e*x)^11)/(11*e^8) + (5*b^2*(b*d - a*e)^3*(7*b*B*d - 4*A*b*e - 3*a*B*e)*(d + e*x)^12)/(12*e^8) - (5*b^3*(b*d - a*e)^2*(7*b*B*d - 3*A*b*e - 4*a*B*e)*(d + e*x)^13)/(13*e^8) + (3*b^4*(b*d - a*e)*(7*b*B*d - 2*A*b*e - 5*a*B*e)*(d + e*x)^14)/(14*e^8) - (b^5*(7*b*B*d - A*b*e - 6*a*B*e)*(d + e*x)^15)/(15*e^8) + (b^6*B*(d + e*x)^16)/(16*e^8)
```

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1524 vs. $2(276) = 552$.

Time = 0.23 (sec) , antiderivative size = 1525, normalized size of antiderivative = 5.22

method	result	size
default	Expression too large to display	1525
norman	Expression too large to display	1642
gospers	Expression too large to display	1944
risch	Expression too large to display	1944
parallelrisch	Expression too large to display	1944
orering	Expression too large to display	1944

input

```
int((b*x+a)^6*(B*x+A)*(e*x+d)^8,x,method=_RETURNVERBOSE)
```

output

```

1/16*b^6*B*e^8*x^16+1/15*((A*b^6+6*B*a*b^5)*e^8+8*b^6*B*d*e^7)*x^15+1/14*(
(6*A*a*b^5+15*B*a^2*b^4)*e^8+8*(A*b^6+6*B*a*b^5)*d*e^7+28*b^6*B*d^2*e^6)*x
^14+1/13*((15*A*a^2*b^4+20*B*a^3*b^3)*e^8+8*(6*A*a*b^5+15*B*a^2*b^4)*d*e^7
+28*(A*b^6+6*B*a*b^5)*d^2*e^6+56*b^6*B*d^3*e^5)*x^13+1/12*((20*A*a^3*b^3+1
5*B*a^4*b^2)*e^8+8*(15*A*a^2*b^4+20*B*a^3*b^3)*d*e^7+28*(6*A*a*b^5+15*B*a^
2*b^4)*d^2*e^6+56*(A*b^6+6*B*a*b^5)*d^3*e^5+70*b^6*B*d^4*e^4)*x^12+1/11*((
15*A*a^4*b^2+6*B*a^5*b)*e^8+8*(20*A*a^3*b^3+15*B*a^4*b^2)*d*e^7+28*(15*A*a
^2*b^4+20*B*a^3*b^3)*d^2*e^6+56*(6*A*a*b^5+15*B*a^2*b^4)*d^3*e^5+70*(A*b^6
+6*B*a*b^5)*d^4*e^4+56*b^6*B*d^5*e^3)*x^11+1/10*((6*A*a^5*b+B*a^6)*e^8+8*(
15*A*a^4*b^2+6*B*a^5*b)*d*e^7+28*(20*A*a^3*b^3+15*B*a^4*b^2)*d^2*e^6+56*(1
5*A*a^2*b^4+20*B*a^3*b^3)*d^3*e^5+70*(6*A*a*b^5+15*B*a^2*b^4)*d^4*e^4+56*(
A*b^6+6*B*a*b^5)*d^5*e^3+28*b^6*B*d^6*e^2)*x^10+1/9*(a^6*A*e^8+8*(6*A*a^5*
b+B*a^6)*d*e^7+28*(15*A*a^4*b^2+6*B*a^5*b)*d^2*e^6+56*(20*A*a^3*b^3+15*B*a
^4*b^2)*d^3*e^5+70*(15*A*a^2*b^4+20*B*a^3*b^3)*d^4*e^4+56*(6*A*a*b^5+15*B*
a^2*b^4)*d^5*e^3+28*(A*b^6+6*B*a*b^5)*d^6*e^2+8*b^6*B*d^7*e)*x^9+1/8*(8*a^
6*A*d*e^7+28*(6*A*a^5*b+B*a^6)*d^2*e^6+56*(15*A*a^4*b^2+6*B*a^5*b)*d^3*e^5
+70*(20*A*a^3*b^3+15*B*a^4*b^2)*d^4*e^4+56*(15*A*a^2*b^4+20*B*a^3*b^3)*d^5
*e^3+28*(6*A*a*b^5+15*B*a^2*b^4)*d^6*e^2+8*(A*b^6+6*B*a*b^5)*d^7*e+b^6*B*d
^8)*x^8+1/7*(28*a^6*A*d^2*e^6+56*(6*A*a^5*b+B*a^6)*d^3*e^5+70*(15*A*a^4*b^
2+6*B*a^5*b)*d^4*e^4+56*(20*A*a^3*b^3+15*B*a^4*b^2)*d^5*e^3+28*(15*A*a^...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1532 vs. $2(276) = 552$.

Time = 0.08 (sec) , antiderivative size = 1532, normalized size of antiderivative = 5.25

$$\int (a + bx)^6 (A + Bx)(d + ex)^8 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^6*(B*x+A)*(e*x+d)^8,x, algorithm="fricas")
```

output

```

1/16*B*b^6*e^8*x^16 + A*a^6*d^8*x + 1/15*(8*B*b^6*d*e^7 + (6*B*a*b^5 + A*b
^6)*e^8)*x^15 + 1/14*(28*B*b^6*d^2*e^6 + 8*(6*B*a*b^5 + A*b^6)*d*e^7 + 3*(
5*B*a^2*b^4 + 2*A*a*b^5)*e^8)*x^14 + 1/13*(56*B*b^6*d^3*e^5 + 28*(6*B*a*b^
5 + A*b^6)*d^2*e^6 + 24*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^7 + 5*(4*B*a^3*b^3 +
3*A*a^2*b^4)*e^8)*x^13 + 1/12*(70*B*b^6*d^4*e^4 + 56*(6*B*a*b^5 + A*b^6)*
d^3*e^5 + 84*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^6 + 40*(4*B*a^3*b^3 + 3*A*a^2
*b^4)*d*e^7 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^8)*x^12 + 1/11*(56*B*b^6*d^5
*e^3 + 70*(6*B*a*b^5 + A*b^6)*d^4*e^4 + 168*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*
e^5 + 140*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^6 + 40*(3*B*a^4*b^2 + 4*A*a^3*
b^3)*d*e^7 + 3*(2*B*a^5*b + 5*A*a^4*b^2)*e^8)*x^11 + 1/10*(28*B*b^6*d^6*e^
2 + 56*(6*B*a*b^5 + A*b^6)*d^5*e^3 + 210*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^4
+ 280*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^5 + 140*(3*B*a^4*b^2 + 4*A*a^3*b^
3)*d^2*e^6 + 24*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^7 + (B*a^6 + 6*A*a^5*b)*e^8)
*x^10 + 1/9*(8*B*b^6*d^7*e + A*a^6*e^8 + 28*(6*B*a*b^5 + A*b^6)*d^6*e^2 +
168*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^3 + 350*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^
4*e^4 + 280*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^5 + 84*(2*B*a^5*b + 5*A*a^4*
b^2)*d^2*e^6 + 8*(B*a^6 + 6*A*a^5*b)*d*e^7)*x^9 + 1/8*(B*b^6*d^8 + 8*A*a^6
*d*e^7 + 8*(6*B*a*b^5 + A*b^6)*d^7*e + 84*(5*B*a^2*b^4 + 2*A*a*b^5)*d^6*e^
2 + 280*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^5*e^3 + 350*(3*B*a^4*b^2 + 4*A*a^3*b^
^3)*d^4*e^4 + 168*(2*B*a^5*b + 5*A*a^4*b^2)*d^3*e^5 + 28*(B*a^6 + 6*A*a...

```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1969 vs. $2(296) = 592$.

Time = 0.12 (sec) , antiderivative size = 1969, normalized size of antiderivative = 6.74

$$\int (a + bx)^6 (A + Bx)(d + ex)^8 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)**6*(B*x+A)*(e*x+d)**8,x)
```


output

```
A*a**6*d**8*x + B*b**6*e**8*x**16/16 + x**15*(A*b**6*e**8/15 + 2*B*a*b**5*
e**8/5 + 8*B*b**6*d*e**7/15) + x**14*(3*A*a*b**5*e**8/7 + 4*A*b**6*d*e**7/
7 + 15*B*a**2*b**4*e**8/14 + 24*B*a*b**5*d*e**7/7 + 2*B*b**6*d**2*e**6) +
x**13*(15*A*a**2*b**4*e**8/13 + 48*A*a*b**5*d*e**7/13 + 28*A*b**6*d**2*e**
6/13 + 20*B*a**3*b**3*e**8/13 + 120*B*a**2*b**4*d*e**7/13 + 168*B*a*b**5*d
**2*e**6/13 + 56*B*b**6*d**3*e**5/13) + x**12*(5*A*a**3*b**3*e**8/3 + 10*A
*a**2*b**4*d*e**7 + 14*A*a*b**5*d**2*e**6 + 14*A*b**6*d**3*e**5/3 + 5*B*a*
*4*b**2*e**8/4 + 40*B*a**3*b**3*d*e**7/3 + 35*B*a**2*b**4*d**2*e**6 + 28*B
*a*b**5*d**3*e**5 + 35*B*b**6*d**4*e**4/6) + x**11*(15*A*a**4*b**2*e**8/11
+ 160*A*a**3*b**3*d*e**7/11 + 420*A*a**2*b**4*d**2*e**6/11 + 336*A*a*b**5
*d**3*e**5/11 + 70*A*b**6*d**4*e**4/11 + 6*B*a**5*b*e**8/11 + 120*B*a**4*b
**2*d*e**7/11 + 560*B*a**3*b**3*d**2*e**6/11 + 840*B*a**2*b**4*d**3*e**5/1
1 + 420*B*a*b**5*d**4*e**4/11 + 56*B*b**6*d**5*e**3/11) + x**10*(3*A*a**5*
b*e**8/5 + 12*A*a**4*b**2*d*e**7 + 56*A*a**3*b**3*d**2*e**6 + 84*A*a**2*b*
*4*d**3*e**5 + 42*A*a*b**5*d**4*e**4 + 28*A*b**6*d**5*e**3/5 + B*a**6*e**8
/10 + 24*B*a**5*b*d*e**7/5 + 42*B*a**4*b**2*d**2*e**6 + 112*B*a**3*b**3*d*
*3*e**5 + 105*B*a**2*b**4*d**4*e**4 + 168*B*a*b**5*d**5*e**3/5 + 14*B*b**6
*d**6*e**2/5) + x**9*(A*a**6*e**8/9 + 16*A*a**5*b*d*e**7/3 + 140*A*a**4*b*
*2*d**2*e**6/3 + 1120*A*a**3*b**3*d**3*e**5/9 + 350*A*a**2*b**4*d**4*e**4/
3 + 112*A*a*b**5*d**5*e**3/3 + 28*A*b**6*d**6*e**2/9 + 8*B*a**6*d*e**7/...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1532 vs. $2(276) = 552$.

Time = 0.05 (sec) , antiderivative size = 1532, normalized size of antiderivative = 5.25

$$\int (a + bx)^6 (A + Bx)(d + ex)^8 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^6*(B*x+A)*(e*x+d)^8,x, algorithm="maxima")
```

output

```

1/16*B*b^6*e^8*x^16 + A*a^6*d^8*x + 1/15*(8*B*b^6*d*e^7 + (6*B*a*b^5 + A*b
^6)*e^8)*x^15 + 1/14*(28*B*b^6*d^2*e^6 + 8*(6*B*a*b^5 + A*b^6)*d*e^7 + 3*(
5*B*a^2*b^4 + 2*A*a*b^5)*e^8)*x^14 + 1/13*(56*B*b^6*d^3*e^5 + 28*(6*B*a*b^
5 + A*b^6)*d^2*e^6 + 24*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^7 + 5*(4*B*a^3*b^3 +
3*A*a^2*b^4)*e^8)*x^13 + 1/12*(70*B*b^6*d^4*e^4 + 56*(6*B*a*b^5 + A*b^6)*
d^3*e^5 + 84*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^6 + 40*(4*B*a^3*b^3 + 3*A*a^2
*b^4)*d*e^7 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^8)*x^12 + 1/11*(56*B*b^6*d^5
*e^3 + 70*(6*B*a*b^5 + A*b^6)*d^4*e^4 + 168*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*
e^5 + 140*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^6 + 40*(3*B*a^4*b^2 + 4*A*a^3*
b^3)*d*e^7 + 3*(2*B*a^5*b + 5*A*a^4*b^2)*e^8)*x^11 + 1/10*(28*B*b^6*d^6*e^
2 + 56*(6*B*a*b^5 + A*b^6)*d^5*e^3 + 210*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^4
+ 280*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^5 + 140*(3*B*a^4*b^2 + 4*A*a^3*b^
3)*d^2*e^6 + 24*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^7 + (B*a^6 + 6*A*a^5*b)*e^8)
*x^10 + 1/9*(8*B*b^6*d^7*e + A*a^6*e^8 + 28*(6*B*a*b^5 + A*b^6)*d^6*e^2 +
168*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^3 + 350*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^
4*e^4 + 280*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^5 + 84*(2*B*a^5*b + 5*A*a^4*
b^2)*d^2*e^6 + 8*(B*a^6 + 6*A*a^5*b)*d*e^7)*x^9 + 1/8*(B*b^6*d^8 + 8*A*a^6
*d*e^7 + 8*(6*B*a*b^5 + A*b^6)*d^7*e + 84*(5*B*a^2*b^4 + 2*A*a*b^5)*d^6*e^
2 + 280*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^5*e^3 + 350*(3*B*a^4*b^2 + 4*A*a^3*b
^3)*d^4*e^4 + 168*(2*B*a^5*b + 5*A*a^4*b^2)*d^3*e^5 + 28*(B*a^6 + 6*A*a...

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1943 vs. $2(276) = 552$.

Time = 0.13 (sec) , antiderivative size = 1943, normalized size of antiderivative = 6.65

$$\int (a + bx)^6 (A + Bx)(d + ex)^8 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^6*(B*x+A)*(e*x+d)^8,x, algorithm="giac")
```

output

```

1/16*B*b^6*e^8*x^16 + 8/15*B*b^6*d*e^7*x^15 + 2/5*B*a*b^5*e^8*x^15 + 1/15*
A*b^6*e^8*x^15 + 2*B*b^6*d^2*e^6*x^14 + 24/7*B*a*b^5*d*e^7*x^14 + 4/7*A*b^
6*d*e^7*x^14 + 15/14*B*a^2*b^4*e^8*x^14 + 3/7*A*a*b^5*e^8*x^14 + 56/13*B*b
^6*d^3*e^5*x^13 + 168/13*B*a*b^5*d^2*e^6*x^13 + 28/13*A*b^6*d^2*e^6*x^13 +
120/13*B*a^2*b^4*d*e^7*x^13 + 48/13*A*a*b^5*d*e^7*x^13 + 20/13*B*a^3*b^3*
e^8*x^13 + 15/13*A*a^2*b^4*e^8*x^13 + 35/6*B*b^6*d^4*e^4*x^12 + 28*B*a*b^5
*d^3*e^5*x^12 + 14/3*A*b^6*d^3*e^5*x^12 + 35*B*a^2*b^4*d^2*e^6*x^12 + 14*A
*a*b^5*d^2*e^6*x^12 + 40/3*B*a^3*b^3*d*e^7*x^12 + 10*A*a^2*b^4*d*e^7*x^12
+ 5/4*B*a^4*b^2*e^8*x^12 + 5/3*A*a^3*b^3*e^8*x^12 + 56/11*B*b^6*d^5*e^3*x^
11 + 420/11*B*a*b^5*d^4*e^4*x^11 + 70/11*A*b^6*d^4*e^4*x^11 + 840/11*B*a^2
*b^4*d^3*e^5*x^11 + 336/11*A*a*b^5*d^3*e^5*x^11 + 560/11*B*a^3*b^3*d^2*e^6
*x^11 + 420/11*A*a^2*b^4*d^2*e^6*x^11 + 120/11*B*a^4*b^2*d*e^7*x^11 + 160/
11*A*a^3*b^3*d*e^7*x^11 + 6/11*B*a^5*b*e^8*x^11 + 15/11*A*a^4*b^2*e^8*x^11
+ 14/5*B*b^6*d^6*e^2*x^10 + 168/5*B*a*b^5*d^5*e^3*x^10 + 28/5*A*b^6*d^5*e
^3*x^10 + 105*B*a^2*b^4*d^4*e^4*x^10 + 42*A*a*b^5*d^4*e^4*x^10 + 112*B*a^3
*b^3*d^3*e^5*x^10 + 84*A*a^2*b^4*d^3*e^5*x^10 + 42*B*a^4*b^2*d^2*e^6*x^10
+ 56*A*a^3*b^3*d^2*e^6*x^10 + 24/5*B*a^5*b*d*e^7*x^10 + 12*A*a^4*b^2*d*e^7
*x^10 + 1/10*B*a^6*e^8*x^10 + 3/5*A*a^5*b*e^8*x^10 + 8/9*B*b^6*d^7*e*x^9 +
56/3*B*a*b^5*d^6*e^2*x^9 + 28/9*A*b^6*d^6*e^2*x^9 + 280/3*B*a^2*b^4*d^5*e
^3*x^9 + 112/3*A*a*b^5*d^5*e^3*x^9 + 1400/9*B*a^3*b^3*d^4*e^4*x^9 + 350...

```

Mupad [B] (verification not implemented)

Time = 1.20 (sec) , antiderivative size = 1625, normalized size of antiderivative = 5.57

$$\int (a + bx)^6 (A + Bx)(d + ex)^8 dx = \text{Too large to display}$$

input

```
int((A + B*x)*(a + b*x)^6*(d + e*x)^8,x)
```

output

```

x^6*(A*a*b^5*d^8 + (5*B*a^2*b^4*d^8)/2 + (28*A*a^6*d^3*e^5)/3 + (35*B*a^6*
d^4*e^4)/3 + 20*A*a^2*b^4*d^7*e + 70*A*a^5*b*d^4*e^4 + (80*B*a^3*b^3*d^7*e
)/3 + 56*B*a^5*b*d^5*e^3 + (280*A*a^3*b^3*d^6*e^2)/3 + 140*A*a^4*b^2*d^5*e
^3 + 70*B*a^4*b^2*d^6*e^2) + x^11*((6*B*a^5*b*e^8)/11 + (15*A*a^4*b^2*e^8)
/11 + (70*A*b^6*d^4*e^4)/11 + (56*B*b^6*d^5*e^3)/11 + (336*A*a*b^5*d^3*e^5
)/11 + (160*A*a^3*b^3*d*e^7)/11 + (420*B*a*b^5*d^4*e^4)/11 + (120*B*a^4*b^
2*d*e^7)/11 + (420*A*a^2*b^4*d^2*e^6)/11 + (840*B*a^2*b^4*d^3*e^5)/11 + (5
60*B*a^3*b^3*d^2*e^6)/11) + x^5*(3*A*a^2*b^4*d^8 + 4*B*a^3*b^3*d^8 + 14*A*
a^6*d^4*e^4 + (56*B*a^6*d^5*e^3)/5 + 32*A*a^3*b^3*d^7*e + (336*A*a^5*b*d^5
*e^3)/5 + 24*B*a^4*b^2*d^7*e + (168*B*a^5*b*d^6*e^2)/5 + 84*A*a^4*b^2*d^6*
e^2) + x^12*((5*A*a^3*b^3*e^8)/3 + (5*B*a^4*b^2*e^8)/4 + (14*A*b^6*d^3*e^5
)/3 + (35*B*b^6*d^4*e^4)/6 + 14*A*a*b^5*d^2*e^6 + 10*A*a^2*b^4*d*e^7 + 28*
B*a*b^5*d^3*e^5 + (40*B*a^3*b^3*d*e^7)/3 + 35*B*a^2*b^4*d^2*e^6) + x^7*((A
*b^6*d^8)/7 + (6*B*a*b^5*d^8)/7 + 4*A*a^6*d^2*e^6 + 8*B*a^6*d^3*e^5 + 48*A
*a^5*b*d^3*e^5 + (120*B*a^2*b^4*d^7*e)/7 + 60*B*a^5*b*d^4*e^4 + 60*A*a^2*b
^4*d^6*e^2 + 160*A*a^3*b^3*d^5*e^3 + 150*A*a^4*b^2*d^4*e^4 + 80*B*a^3*b^3*
d^6*e^2 + 120*B*a^4*b^2*d^5*e^3 + (48*A*a*b^5*d^7*e)/7) + x^10*((B*a^6*e^8
)/10 + (3*A*a^5*b*e^8)/5 + (28*A*b^6*d^5*e^3)/5 + (14*B*b^6*d^6*e^2)/5 + 4
2*A*a*b^5*d^4*e^4 + 12*A*a^4*b^2*d*e^7 + (168*B*a*b^5*d^5*e^3)/5 + 84*A*a^
2*b^4*d^3*e^5 + 56*A*a^3*b^3*d^2*e^6 + 105*B*a^2*b^4*d^4*e^4 + 112*B*a^...

```

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 1051, normalized size of antiderivative = 3.60

$$\int (a + bx)^6 (A + Bx)(d + ex)^8 dx = \text{Too large to display}$$

input

```
int((b*x+a)^6*(B*x+A)*(e*x+d)^8,x)
```

output

```
(x*(102960*a**7*d**8 + 411840*a**7*d**7*e*x + 960960*a**7*d**6*e**2*x**2 +
1441440*a**7*d**5*e**3*x**3 + 1441440*a**7*d**4*e**4*x**4 + 960960*a**7*d
**3*e**5*x**5 + 411840*a**7*d**2*e**6*x**6 + 102960*a**7*d*e**7*x**7 + 114
40*a**7*e**8*x**8 + 360360*a**6*b*d**8*x + 1921920*a**6*b*d**7*e*x**2 + 50
45040*a**6*b*d**6*e**2*x**3 + 8072064*a**6*b*d**5*e**3*x**4 + 8408400*a**6
*b*d**4*e**4*x**5 + 5765760*a**6*b*d**3*e**5*x**6 + 2522520*a**6*b*d**2*e
**6*x**7 + 640640*a**6*b*d*e**7*x**8 + 72072*a**6*b*e**8*x**9 + 720720*a**5
*b**2*d**8*x**2 + 4324320*a**5*b**2*d**7*e*x**3 + 12108096*a**5*b**2*d**6*
e**2*x**4 + 20180160*a**5*b**2*d**5*e**3*x**5 + 21621600*a**5*b**2*d**4*e
**4*x**6 + 15135120*a**5*b**2*d**3*e**5*x**7 + 6726720*a**5*b**2*d**2*e**6*
x**8 + 1729728*a**5*b**2*d*e**7*x**9 + 196560*a**5*b**2*e**8*x**10 + 90090
0*a**4*b**3*d**8*x**3 + 5765760*a**4*b**3*d**7*e*x**4 + 16816800*a**4*b**3
*d**6*e**2*x**5 + 28828800*a**4*b**3*d**5*e**3*x**6 + 31531500*a**4*b**3*d
**4*e**4*x**7 + 22422400*a**4*b**3*d**3*e**5*x**8 + 10090080*a**4*b**3*d**
2*e**6*x**9 + 2620800*a**4*b**3*d*e**7*x**10 + 300300*a**4*b**3*e**8*x**11
+ 720720*a**3*b**4*d**8*x**4 + 4804800*a**3*b**4*d**7*e*x**5 + 14414400*a
**3*b**4*d**6*e**2*x**6 + 25225200*a**3*b**4*d**5*e**3*x**7 + 28028000*a**
3*b**4*d**4*e**4*x**8 + 20180160*a**3*b**4*d**3*e**5*x**9 + 9172800*a**3*b
**4*d**2*e**6*x**10 + 2402400*a**3*b**4*d*e**7*x**11 + 277200*a**3*b**4*e
**8*x**12 + 360360*a**2*b**5*d**8*x**5 + 2471040*a**2*b**5*d**7*e*x**6 + ...
```

3.42 $\int (a + bx)^6 (A + Bx)(d + ex)^7 dx$

Optimal result	417
Mathematica [B] (verified)	418
Rubi [A] (verified)	419
Maple [B] (verified)	420
Fricas [B] (verification not implemented)	421
Sympy [B] (verification not implemented)	422
Maxima [B] (verification not implemented)	423
Giac [B] (verification not implemented)	424
Mupad [B] (verification not implemented)	425
Reduce [B] (verification not implemented)	426

Optimal result

Integrand size = 20, antiderivative size = 292

$$\begin{aligned}
 \int (a + bx)^6 (A + Bx)(d + ex)^7 dx = & -\frac{(bd - ae)^6 (Bd - Ae)(d + ex)^8}{8e^8} \\
 & + \frac{(bd - ae)^5 (7bBd - 6Abe - aBe)(d + ex)^9}{9e^8} \\
 & - \frac{3b(bd - ae)^4 (7bBd - 5Abe - 2aBe)(d + ex)^{10}}{10e^8} \\
 & + \frac{5b^2 (bd - ae)^3 (7bBd - 4Abe - 3aBe)(d + ex)^{11}}{11e^8} \\
 & - \frac{5b^3 (bd - ae)^2 (7bBd - 3Abe - 4aBe)(d + ex)^{12}}{12e^8} \\
 & + \frac{3b^4 (bd - ae) (7bBd - 2Abe - 5aBe)(d + ex)^{13}}{13e^8} \\
 & - \frac{b^5 (7bBd - Abe - 6aBe)(d + ex)^{14}}{14e^8} \\
 & + \frac{b^6 B (d + ex)^{15}}{15e^8}
 \end{aligned}$$

output

```
-1/8*(-a*e+b*d)^6*(-A*e+B*d)*(e*x+d)^8/e^8+1/9*(-a*e+b*d)^5*(-6*A*b*e-B*a*
e+7*B*b*d)*(e*x+d)^9/e^8-3/10*b*(-a*e+b*d)^4*(-5*A*b*e-2*B*a*e+7*B*b*d)*(e
*x+d)^10/e^8+5/11*b^2*(-a*e+b*d)^3*(-4*A*b*e-3*B*a*e+7*B*b*d)*(e*x+d)^11/e
^8-5/12*b^3*(-a*e+b*d)^2*(-3*A*b*e-4*B*a*e+7*B*b*d)*(e*x+d)^12/e^8+3/13*b^
4*(-a*e+b*d)*(-2*A*b*e-5*B*a*e+7*B*b*d)*(e*x+d)^13/e^8-1/14*b^5*(-A*b*e-6*
B*a*e+7*B*b*d)*(e*x+d)^14/e^8+1/15*b^6*B*(e*x+d)^15/e^8
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1224 vs. $2(292) = 584$.

Time = 0.29 (sec) , antiderivative size = 1224, normalized size of antiderivative = 4.19

$$\int (a + bx)^6 (A + Bx)(d + ex)^7 dx = \text{Too large to display}$$

input

```
Integrate[(a + b*x)^6*(A + B*x)*(d + e*x)^7,x]
```

output

```
a^6*A*d^7*x + (a^5*d^6*(6*A*b*d + a*B*d + 7*a*A*e)*x^2)/2 + (a^4*d^5*(a*B*
d*(6*b*d + 7*a*e) + 3*A*(5*b^2*d^2 + 14*a*b*d*e + 7*a^2*e^2))*x^3)/3 + (a^
3*d^4*(3*a*B*d*(5*b^2*d^2 + 14*a*b*d*e + 7*a^2*e^2) + A*(20*b^3*d^3 + 105*
a*b^2*d^2*e + 126*a^2*b*d*e^2 + 35*a^3*e^3))*x^4)/4 + (a^2*d^3*(a*B*d*(20*
b^3*d^3 + 105*a*b^2*d^2*e + 126*a^2*b*d*e^2 + 35*a^3*e^3) + 5*A*(3*b^4*d^4
+ 28*a*b^3*d^3*e + 63*a^2*b^2*d^2*e^2 + 42*a^3*b*d*e^3 + 7*a^4*e^4))*x^5)
/5 + (a*d^2*(5*a*B*d*(3*b^4*d^4 + 28*a*b^3*d^3*e + 63*a^2*b^2*d^2*e^2 + 42
*a^3*b*d*e^3 + 7*a^4*e^4) + 3*A*(2*b^5*d^5 + 35*a*b^4*d^4*e + 140*a^2*b^3*d^3*
d^3*e^2 + 175*a^3*b^2*d^2*e^3 + 70*a^4*b*d*e^4 + 7*a^5*e^5))*x^6)/6 + (d*(
3*a*B*d*(2*b^5*d^5 + 35*a*b^4*d^4*e + 140*a^2*b^3*d^3*e^2 + 175*a^3*b^2*d^
2*e^3 + 70*a^4*b*d*e^4 + 7*a^5*e^5) + A*(b^6*d^6 + 42*a*b^5*d^5*e + 315*a^
2*b^4*d^4*e^2 + 700*a^3*b^3*d^3*e^3 + 525*a^4*b^2*d^2*e^4 + 126*a^5*b*d*e^
5 + 7*a^6*e^6))*x^7)/7 + ((700*a^3*b^3*d^3*e^3*(B*d + A*e) + 42*a^5*b*d*e^
5*(3*B*d + A*e) + a^6*e^6*(7*B*d + A*e) + 42*a*b^5*d^5*e*(B*d + 3*A*e) + 1
05*a^4*b^2*d^2*e^4*(5*B*d + 3*A*e) + 105*a^2*b^4*d^4*e^2*(3*B*d + 5*A*e) +
b^6*d^6*(B*d + 7*A*e))*x^8)/8 + (e*(a^6*B*e^6 + 525*a^2*b^4*d^3*e^2*(B*d
+ A*e) + 105*a^4*b^2*d*e^4*(3*B*d + A*e) + 6*a^5*b*e^5*(7*B*d + A*e) + 7*b
^6*d^5*(B*d + 3*A*e) + 140*a^3*b^3*d^2*e^3*(5*B*d + 3*A*e) + 42*a*b^5*d^4*
e*(3*B*d + 5*A*e))*x^9)/9 + (b*e^2*(6*a^5*B*e^5 + 210*a*b^4*d^3*e*(B*d + A
*e) + 140*a^3*b^2*d*e^3*(3*B*d + A*e) + 15*a^4*b*e^4*(7*B*d + A*e) + 10...
```

Rubi [A] (verified)

Time = 1.49 (sec) , antiderivative size = 292, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^6 (A + Bx)(d + ex)^7 dx$$

↓ 86

$$\int \left(\frac{b^5(d + ex)^{13}(6aBe + Abe - 7bBd)}{e^7} - \frac{3b^4(d + ex)^{12}(bd - ae)(5aBe + 2Abe - 7bBd)}{e^7} + \frac{5b^3(d + ex)^{11}(bd - ae)^2(-4aBe - 3Abe + 7bBd)}{e^7} - \frac{5b^2(d + ex)^{10}(bd - ae)^3(-3aBe - 4Abe + 7bBd)}{e^7} + \frac{3b(d + ex)^9(bd - ae)^4(-2aBe - 5Abe + 7bBd)}{e^7} + \frac{(d + ex)^8(bd - ae)^5(-aBe - 6Abe + 7bBd)}{e^7} - \frac{(d + ex)^7(bd - ae)^6(Bd - Ae)}{e^7} + \frac{b^6B(d + ex)^6(bd - ae)^7}{e^7} \right) dx$$

↓ 2009

$$-\frac{b^5(d + ex)^{14}(-6aBe - Abe + 7bBd)}{14e^8} + \frac{3b^4(d + ex)^{13}(bd - ae)(-5aBe - 2Abe + 7bBd)}{13e^8} - \frac{5b^3(d + ex)^{12}(bd - ae)^2(-4aBe - 3Abe + 7bBd)}{12e^8} + \frac{5b^2(d + ex)^{11}(bd - ae)^3(-3aBe - 4Abe + 7bBd)}{11e^8} - \frac{3b(d + ex)^{10}(bd - ae)^4(-2aBe - 5Abe + 7bBd)}{10e^8} + \frac{(d + ex)^9(bd - ae)^5(-aBe - 6Abe + 7bBd)}{9e^8} - \frac{(d + ex)^8(bd - ae)^6(Bd - Ae)}{8e^8} + \frac{b^6B(d + ex)^{15}}{15e^8}$$

input `Int[(a + b*x)^6*(A + B*x)*(d + e*x)^7,x]`

output `-1/8*((b*d - a*e)^6*(B*d - A*e)*(d + e*x)^8)/e^8 + ((b*d - a*e)^5*(7*b*B*d - 6*A*b*e - a*B*e)*(d + e*x)^9)/(9*e^8) - (3*b*(b*d - a*e)^4*(7*b*B*d - 5*A*b*e - 2*a*B*e)*(d + e*x)^10)/(10*e^8) + (5*b^2*(b*d - a*e)^3*(7*b*B*d - 4*A*b*e - 3*a*B*e)*(d + e*x)^11)/(11*e^8) - (5*b^3*(b*d - a*e)^2*(7*b*B*d - 3*A*b*e - 4*a*B*e)*(d + e*x)^12)/(12*e^8) + (3*b^4*(b*d - a*e)*(7*b*B*d - 2*A*b*e - 5*a*B*e)*(d + e*x)^13)/(13*e^8) - (b^5*(7*b*B*d - A*b*e - 6*a*B*e)*(d + e*x)^14)/(14*e^8) + (b^6*B*(d + e*x)^15)/(15*e^8)`

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0]) || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1348 vs. $2(276) = 552$.

Time = 0.23 (sec) , antiderivative size = 1349, normalized size of antiderivative = 4.62

method	result	size
default	Expression too large to display	1349
norman	Expression too large to display	1448
orering	Expression too large to display	1712
gosper	Expression too large to display	1713
risch	Expression too large to display	1713
parallelrisc	Expression too large to display	1713

input

```
int((b*x+a)^6*(B*x+A)*(e*x+d)^7,x,method=_RETURNVERBOSE)
```

output

```

1/15*b^6*B*e^7*x^15+1/14*((A*b^6+6*B*a*b^5)*e^7+7*b^6*B*d*e^6)*x^14+1/13*(
(6*A*a*b^5+15*B*a^2*b^4)*e^7+7*(A*b^6+6*B*a*b^5)*d*e^6+21*b^6*B*d^2*e^5)*x
^13+1/12*((15*A*a^2*b^4+20*B*a^3*b^3)*e^7+7*(6*A*a*b^5+15*B*a^2*b^4)*d*e^6
+21*(A*b^6+6*B*a*b^5)*d^2*e^5+35*b^6*B*d^3*e^4)*x^12+1/11*((20*A*a^3*b^3+1
5*B*a^4*b^2)*e^7+7*(15*A*a^2*b^4+20*B*a^3*b^3)*d*e^6+21*(6*A*a*b^5+15*B*a^
2*b^4)*d^2*e^5+35*(A*b^6+6*B*a*b^5)*d^3*e^4+35*b^6*B*d^4*e^3)*x^11+1/10*((
15*A*a^4*b^2+6*B*a^5*b)*e^7+7*(20*A*a^3*b^3+15*B*a^4*b^2)*d*e^6+21*(15*A*a
^2*b^4+20*B*a^3*b^3)*d^2*e^5+35*(6*A*a*b^5+15*B*a^2*b^4)*d^3*e^4+35*(A*b^6
+6*B*a*b^5)*d^4*e^3+21*b^6*B*d^5*e^2)*x^10+1/9*((6*A*a^5*b+B*a^6)*e^7+7*(1
5*A*a^4*b^2+6*B*a^5*b)*d*e^6+21*(20*A*a^3*b^3+15*B*a^4*b^2)*d^2*e^5+35*(15
*A*a^2*b^4+20*B*a^3*b^3)*d^3*e^4+35*(6*A*a*b^5+15*B*a^2*b^4)*d^4*e^3+21*(A
*b^6+6*B*a*b^5)*d^5*e^2+7*b^6*B*d^6*e)*x^9+1/8*(a^6*A*e^7+7*(6*A*a^5*b+B*a
^6)*d*e^6+21*(15*A*a^4*b^2+6*B*a^5*b)*d^2*e^5+35*(20*A*a^3*b^3+15*B*a^4*b^
2)*d^3*e^4+35*(15*A*a^2*b^4+20*B*a^3*b^3)*d^4*e^3+21*(6*A*a*b^5+15*B*a^2*b
^4)*d^5*e^2+7*(A*b^6+6*B*a*b^5)*d^6*e+b^6*B*d^7)*x^8+1/7*(7*a^6*A*d*e^6+21
*(6*A*a^5*b+B*a^6)*d^2*e^5+35*(15*A*a^4*b^2+6*B*a^5*b)*d^3*e^4+35*(20*A*a^
3*b^3+15*B*a^4*b^2)*d^4*e^3+21*(15*A*a^2*b^4+20*B*a^3*b^3)*d^5*e^2+7*(6*A*
a*b^5+15*B*a^2*b^4)*d^6*e+(A*b^6+6*B*a*b^5)*d^7)*x^7+1/6*(21*a^6*A*d^2*e^5
+35*(6*A*a^5*b+B*a^6)*d^3*e^4+35*(15*A*a^4*b^2+6*B*a^5*b)*d^4*e^3+21*(20*A
*a^3*b^3+15*B*a^4*b^2)*d^5*e^2+7*(15*A*a^2*b^4+20*B*a^3*b^3)*d^6*e+(6*A...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1356 vs. $2(276) = 552$.

Time = 0.07 (sec) , antiderivative size = 1356, normalized size of antiderivative = 4.64

$$\int (a + bx)^6 (A + Bx)(d + ex)^7 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^6*(B*x+A)*(e*x+d)^7,x, algorithm="fricas")
```

output

```

1/15*B*b^6*e^7*x^15 + A*a^6*d^7*x + 1/14*(7*B*b^6*d*e^6 + (6*B*a*b^5 + A*b
^6)*e^7)*x^14 + 1/13*(21*B*b^6*d^2*e^5 + 7*(6*B*a*b^5 + A*b^6)*d*e^6 + 3*(
5*B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^13 + 1/12*(35*B*b^6*d^3*e^4 + 21*(6*B*a*b^
5 + A*b^6)*d^2*e^5 + 21*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6 + 5*(4*B*a^3*b^3 +
3*A*a^2*b^4)*e^7)*x^12 + 1/11*(35*B*b^6*d^4*e^3 + 35*(6*B*a*b^5 + A*b^6)*
d^3*e^4 + 63*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 + 35*(4*B*a^3*b^3 + 3*A*a^2
*b^4)*d*e^6 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^11 + 1/10*(21*B*b^6*d^5
*e^2 + 35*(6*B*a*b^5 + A*b^6)*d^4*e^3 + 105*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*
e^4 + 105*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + 35*(3*B*a^4*b^2 + 4*A*a^3*
b^3)*d*e^6 + 3*(2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^10 + 1/9*(7*B*b^6*d^6*e +
21*(6*B*a*b^5 + A*b^6)*d^5*e^2 + 105*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^3 + 1
75*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 + 105*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d
^2*e^5 + 21*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^6 + (B*a^6 + 6*A*a^5*b)*e^7)*x^9
+ 1/8*(B*b^6*d^7 + A*a^6*e^7 + 7*(6*B*a*b^5 + A*b^6)*d^6*e + 63*(5*B*a^2*
b^4 + 2*A*a*b^5)*d^5*e^2 + 175*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4*e^3 + 175*(
3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 + 63*(2*B*a^5*b + 5*A*a^4*b^2)*d^2*e^5
+ 7*(B*a^6 + 6*A*a^5*b)*d*e^6)*x^8 + 1/7*(7*A*a^6*d*e^6 + (6*B*a*b^5 + A*b
^6)*d^7 + 21*(5*B*a^2*b^4 + 2*A*a*b^5)*d^6*e + 105*(4*B*a^3*b^3 + 3*A*a^2*
b^4)*d^5*e^2 + 175*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^4*e^3 + 105*(2*B*a^5*b +
5*A*a^4*b^2)*d^3*e^4 + 21*(B*a^6 + 6*A*a^5*b)*d^2*e^5)*x^7 + 1/6*(21*A*...

```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1756 vs. $2(296) = 592$.

Time = 0.11 (sec) , antiderivative size = 1756, normalized size of antiderivative = 6.01

$$\int (a + bx)^6 (A + Bx)(d + ex)^7 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)**6*(B*x+A)*(e*x+d)**7,x)
```

output

```

A*a**6*d**7*x + B*b**6*e**7*x**15/15 + x**14*(A*b**6*e**7/14 + 3*B*a*b**5*
e**7/7 + B*b**6*d*e**6/2) + x**13*(6*A*a*b**5*e**7/13 + 7*A*b**6*d*e**6/13
+ 15*B*a**2*b**4*e**7/13 + 42*B*a*b**5*d*e**6/13 + 21*B*b**6*d**2*e**5/13
) + x**12*(5*A*a**2*b**4*e**7/4 + 7*A*a*b**5*d*e**6/2 + 7*A*b**6*d**2*e**5
/4 + 5*B*a**3*b**3*e**7/3 + 35*B*a**2*b**4*d*e**6/4 + 21*B*a*b**5*d**2*e**
5/2 + 35*B*b**6*d**3*e**4/12) + x**11*(20*A*a**3*b**3*e**7/11 + 105*A*a**2
*b**4*d*e**6/11 + 126*A*a*b**5*d**2*e**5/11 + 35*A*b**6*d**3*e**4/11 + 15*
B*a**4*b**2*e**7/11 + 140*B*a**3*b**3*d*e**6/11 + 315*B*a**2*b**4*d**2*e**
5/11 + 210*B*a*b**5*d**3*e**4/11 + 35*B*b**6*d**4*e**3/11) + x**10*(3*A*a*
*4*b**2*e**7/2 + 14*A*a**3*b**3*d*e**6 + 63*A*a**2*b**4*d**2*e**5/2 + 21*A
*a*b**5*d**3*e**4 + 7*A*b**6*d**4*e**3/2 + 3*B*a**5*b*e**7/5 + 21*B*a**4*b
**2*d*e**6/2 + 42*B*a**3*b**3*d**2*e**5 + 105*B*a**2*b**4*d**3*e**4/2 + 21
*B*a*b**5*d**4*e**3 + 21*B*b**6*d**5*e**2/10) + x**9*(2*A*a**5*b*e**7/3 +
35*A*a**4*b**2*d*e**6/3 + 140*A*a**3*b**3*d**2*e**5/3 + 175*A*a**2*b**4*d*
*3*e**4/3 + 70*A*a*b**5*d**4*e**3/3 + 7*A*b**6*d**5*e**2/3 + B*a**6*e**7/9
+ 14*B*a**5*b*d*e**6/3 + 35*B*a**4*b**2*d**2*e**5 + 700*B*a**3*b**3*d**3*
e**4/9 + 175*B*a**2*b**4*d**4*e**3/3 + 14*B*a*b**5*d**5*e**2 + 7*B*b**6*d*
*6*e/9) + x**8*(A*a**6*e**7/8 + 21*A*a**5*b*d*e**6/4 + 315*A*a**4*b**2*d**
2*e**5/8 + 175*A*a**3*b**3*d**3*e**4/2 + 525*A*a**2*b**4*d**4*e**3/8 + 63*
A*a*b**5*d**5*e**2/4 + 7*A*b**6*d**6*e/8 + 7*B*a**6*d*e**6/8 + 63*B*a**...

```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1356 vs. $2(276) = 552$.

Time = 0.04 (sec) , antiderivative size = 1356, normalized size of antiderivative = 4.64

$$\int (a + bx)^6 (A + Bx)(d + ex)^7 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^6*(B*x+A)*(e*x+d)^7,x, algorithm="maxima")
```

output

```

1/15*B*b^6*e^7*x^15 + A*a^6*d^7*x + 1/14*(7*B*b^6*d*e^6 + (6*B*a*b^5 + A*b
^6)*e^7)*x^14 + 1/13*(21*B*b^6*d^2*e^5 + 7*(6*B*a*b^5 + A*b^6)*d*e^6 + 3*(
5*B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^13 + 1/12*(35*B*b^6*d^3*e^4 + 21*(6*B*a*b^
5 + A*b^6)*d^2*e^5 + 21*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6 + 5*(4*B*a^3*b^3 +
3*A*a^2*b^4)*e^7)*x^12 + 1/11*(35*B*b^6*d^4*e^3 + 35*(6*B*a*b^5 + A*b^6)*
d^3*e^4 + 63*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 + 35*(4*B*a^3*b^3 + 3*A*a^2
*b^4)*d*e^6 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^11 + 1/10*(21*B*b^6*d^5
*e^2 + 35*(6*B*a*b^5 + A*b^6)*d^4*e^3 + 105*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*
e^4 + 105*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + 35*(3*B*a^4*b^2 + 4*A*a^3*
b^3)*d*e^6 + 3*(2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^10 + 1/9*(7*B*b^6*d^6*e +
21*(6*B*a*b^5 + A*b^6)*d^5*e^2 + 105*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^3 + 1
75*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 + 105*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d
^2*e^5 + 21*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^6 + (B*a^6 + 6*A*a^5*b)*e^7)*x^9
+ 1/8*(B*b^6*d^7 + A*a^6*e^7 + 7*(6*B*a*b^5 + A*b^6)*d^6*e + 63*(5*B*a^2*
b^4 + 2*A*a*b^5)*d^5*e^2 + 175*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4*e^3 + 175*(
3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 + 63*(2*B*a^5*b + 5*A*a^4*b^2)*d^2*e^5
+ 7*(B*a^6 + 6*A*a^5*b)*d*e^6)*x^8 + 1/7*(7*A*a^6*d*e^6 + (6*B*a*b^5 + A*b
^6)*d^7 + 21*(5*B*a^2*b^4 + 2*A*a*b^5)*d^6*e + 105*(4*B*a^3*b^3 + 3*A*a^2*
b^4)*d^5*e^2 + 175*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^4*e^3 + 105*(2*B*a^5*b +
5*A*a^4*b^2)*d^3*e^4 + 21*(B*a^6 + 6*A*a^5*b)*d^2*e^5)*x^7 + 1/6*(21*A*...

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1712 vs. $2(276) = 552$.

Time = 0.13 (sec) , antiderivative size = 1712, normalized size of antiderivative = 5.86

$$\int (a + bx)^6 (A + Bx)(d + ex)^7 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^6*(B*x+A)*(e*x+d)^7,x, algorithm="giac")
```

output

```

1/15*B*b^6*e^7*x^15 + 1/2*B*b^6*d*e^6*x^14 + 3/7*B*a*b^5*e^7*x^14 + 1/14*A
*b^6*e^7*x^14 + 21/13*B*b^6*d^2*e^5*x^13 + 42/13*B*a*b^5*d*e^6*x^13 + 7/13
*A*b^6*d*e^6*x^13 + 15/13*B*a^2*b^4*e^7*x^13 + 6/13*A*a*b^5*e^7*x^13 + 35/
12*B*b^6*d^3*e^4*x^12 + 21/2*B*a*b^5*d^2*e^5*x^12 + 7/4*A*b^6*d^2*e^5*x^12
+ 35/4*B*a^2*b^4*d*e^6*x^12 + 7/2*A*a*b^5*d*e^6*x^12 + 5/3*B*a^3*b^3*e^7*
x^12 + 5/4*A*a^2*b^4*e^7*x^12 + 35/11*B*b^6*d^4*e^3*x^11 + 210/11*B*a*b^5*
d^3*e^4*x^11 + 35/11*A*b^6*d^3*e^4*x^11 + 315/11*B*a^2*b^4*d^2*e^5*x^11 +
126/11*A*a*b^5*d^2*e^5*x^11 + 140/11*B*a^3*b^3*d*e^6*x^11 + 105/11*A*a^2*b
^4*d*e^6*x^11 + 15/11*B*a^4*b^2*e^7*x^11 + 20/11*A*a^3*b^3*e^7*x^11 + 21/1
0*B*b^6*d^5*e^2*x^10 + 21*B*a*b^5*d^4*e^3*x^10 + 7/2*A*b^6*d^4*e^3*x^10 +
105/2*B*a^2*b^4*d^3*e^4*x^10 + 21*A*a*b^5*d^3*e^4*x^10 + 42*B*a^3*b^3*d^2*
e^5*x^10 + 63/2*A*a^2*b^4*d^2*e^5*x^10 + 21/2*B*a^4*b^2*d*e^6*x^10 + 14*A*
a^3*b^3*d*e^6*x^10 + 3/5*B*a^5*b*e^7*x^10 + 3/2*A*a^4*b^2*e^7*x^10 + 7/9*B
*b^6*d^6*e*x^9 + 14*B*a*b^5*d^5*e^2*x^9 + 7/3*A*b^6*d^5*e^2*x^9 + 175/3*B*
a^2*b^4*d^4*e^3*x^9 + 70/3*A*a*b^5*d^4*e^3*x^9 + 700/9*B*a^3*b^3*d^3*e^4*x
^9 + 175/3*A*a^2*b^4*d^3*e^4*x^9 + 35*B*a^4*b^2*d^2*e^5*x^9 + 140/3*A*a^3*
b^3*d^2*e^5*x^9 + 14/3*B*a^5*b*d*e^6*x^9 + 35/3*A*a^4*b^2*d*e^6*x^9 + 1/9*
B*a^6*e^7*x^9 + 2/3*A*a^5*b*e^7*x^9 + 1/8*B*b^6*d^7*x^8 + 21/4*B*a*b^5*d^6
*e*x^8 + 7/8*A*b^6*d^6*e*x^8 + 315/8*B*a^2*b^4*d^5*e^2*x^8 + 63/4*A*a*b^5*
d^5*e^2*x^8 + 175/2*B*a^3*b^3*d^4*e^3*x^8 + 525/8*A*a^2*b^4*d^4*e^3*x^8...

```

Mupad [B] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 1431, normalized size of antiderivative = 4.90

$$\int (a + bx)^6 (A + Bx)(d + ex)^7 dx = \text{Too large to display}$$

input

```
int((A + B*x)*(a + b*x)^6*(d + e*x)^7,x)
```

output

```

x^6*(A*a*b^5*d^7 + (5*B*a^2*b^4*d^7)/2 + (7*A*a^6*d^2*e^5)/2 + (35*B*a^6*d^3*e^4)/6 + (35*A*a^2*b^4*d^6*e)/2 + 35*A*a^5*b*d^3*e^4 + (70*B*a^3*b^3*d^6*e)/3 + 35*B*a^5*b*d^4*e^3 + 70*A*a^3*b^3*d^5*e^2 + (175*A*a^4*b^2*d^4*e^3)/2 + (105*B*a^4*b^2*d^5*e^2)/2) + x^10*((3*B*a^5*b*e^7)/5 + (3*A*a^4*b^2*e^7)/2 + (7*A*b^6*d^4*e^3)/2 + (21*B*b^6*d^5*e^2)/10 + 21*A*a*b^5*d^3*e^4 + 14*A*a^3*b^3*d*e^6 + 21*B*a*b^5*d^4*e^3 + (21*B*a^4*b^2*d*e^6)/2 + (63*A*a^2*b^4*d^2*e^5)/2 + (105*B*a^2*b^4*d^3*e^4)/2 + 42*B*a^3*b^3*d^2*e^5) + x^5*(3*A*a^2*b^4*d^7 + 4*B*a^3*b^3*d^7 + 7*A*a^6*d^3*e^4 + 7*B*a^6*d^4*e^3 + 28*A*a^3*b^3*d^6*e + 42*A*a^5*b*d^4*e^3 + 21*B*a^4*b^2*d^6*e + (126*B*a^5*b*d^5*e^2)/5 + 63*A*a^4*b^2*d^5*e^2) + x^11*((20*A*a^3*b^3*e^7)/11 + (15*B*a^4*b^2*e^7)/11 + (35*A*b^6*d^3*e^4)/11 + (35*B*b^6*d^4*e^3)/11 + (126*A*a*b^5*d^2*e^5)/11 + (105*A*a^2*b^4*d*e^6)/11 + (210*B*a*b^5*d^3*e^4)/11 + (140*B*a^3*b^3*d*e^6)/11 + (315*B*a^2*b^4*d^2*e^5)/11) + x^3*(2*B*a^5*b*d^7 + (7*B*a^6*d^6*e)/3 + 5*A*a^4*b^2*d^7 + 7*A*a^6*d^5*e^2 + 14*A*a^5*b*d^6*e) + x^8*((A*a^6*e^7)/8 + (B*b^6*d^7)/8 + (7*A*b^6*d^6*e)/8 + (7*B*a^6*d^6*e)/8 + (63*A*a*b^5*d^5*e^2)/4 + (63*B*a^5*b*d^2*e^5)/4 + (525*A*a^2*b^4*d^4*e^3)/8 + (175*A*a^3*b^3*d^3*e^4)/2 + (315*A*a^4*b^2*d^2*e^5)/8 + (315*B*a^2*b^4*d^5*e^2)/8 + (175*B*a^3*b^3*d^4*e^3)/2 + (525*B*a^4*b^2*d^3*e^4)/8 + (21*A*a^5*b*d*e^6)/4 + (21*B*a*b^5*d^6*e)/4) + x^13*((6*A*a*b^5*e^7)/13 + (7*A*b^6*d*e^6)/13 + (15*B*a^2*b^4*e^7)/13 + (21*B*b^6*d^2*e^5...

```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 925, normalized size of antiderivative = 3.17

$$\int (a + bx)^6 (A + Bx)(d + ex)^7 dx = \text{Too large to display}$$

input

```
int((b*x+a)^6*(B*x+A)*(e*x+d)^7,x)
```

output

```
(x*(51480*a**7*d**7 + 180180*a**7*d**6*e*x + 360360*a**7*d**5*e**2*x**2 +
450450*a**7*d**4*e**3*x**3 + 360360*a**7*d**3*e**4*x**4 + 180180*a**7*d**2
*e**5*x**5 + 51480*a**7*d*e**6*x**6 + 6435*a**7*e**7*x**7 + 180180*a**6*b*
d**7*x + 840840*a**6*b*d**6*e*x**2 + 1891890*a**6*b*d**5*e**2*x**3 + 25225
20*a**6*b*d**4*e**3*x**4 + 2102100*a**6*b*d**3*e**4*x**5 + 1081080*a**6*b*
d**2*e**5*x**6 + 315315*a**6*b*d*e**6*x**7 + 40040*a**6*b*e**7*x**8 + 3603
60*a**5*b**2*d**7*x**2 + 1891890*a**5*b**2*d**6*e*x**3 + 4540536*a**5*b**2
*d**5*e**2*x**4 + 6306300*a**5*b**2*d**4*e**3*x**5 + 5405400*a**5*b**2*d**
3*e**4*x**6 + 2837835*a**5*b**2*d**2*e**5*x**7 + 840840*a**5*b**2*d*e**6*x
**8 + 108108*a**5*b**2*e**7*x**9 + 450450*a**4*b**3*d**7*x**3 + 2522520*a*
**4*b**3*d**6*e*x**4 + 6306300*a**4*b**3*d**5*e**2*x**5 + 9009000*a**4*b**3
*d**4*e**3*x**6 + 7882875*a**4*b**3*d**3*e**4*x**7 + 4204200*a**4*b**3*d**
2*e**5*x**8 + 1261260*a**4*b**3*d*e**6*x**9 + 163800*a**4*b**3*e**7*x**10
+ 360360*a**3*b**4*d**7*x**4 + 2102100*a**3*b**4*d**6*e*x**5 + 5405400*a**
3*b**4*d**5*e**2*x**6 + 7882875*a**3*b**4*d**4*e**3*x**7 + 7007000*a**3*b*
**4*d**3*e**4*x**8 + 3783780*a**3*b**4*d**2*e**5*x**9 + 1146600*a**3*b**4*d
*e**6*x**10 + 150150*a**3*b**4*e**7*x**11 + 180180*a**2*b**5*d**7*x**5 + 1
081080*a**2*b**5*d**6*e*x**6 + 2837835*a**2*b**5*d**5*e**2*x**7 + 4204200*
a**2*b**5*d**4*e**3*x**8 + 3783780*a**2*b**5*d**3*e**4*x**9 + 2063880*a**2
*b**5*d**2*e**5*x**10 + 630630*a**2*b**5*d*e**6*x**11 + 83160*a**2*b**5...
```


3.43 $\int (a + bx)^6 (A + Bx)(d + ex)^6 dx$

Optimal result	428
Mathematica [B] (verified)	429
Rubi [A] (verified)	431
Maple [B] (verified)	433
Fricas [B] (verification not implemented)	434
Sympy [B] (verification not implemented)	435
Maxima [B] (verification not implemented)	436
Giac [B] (verification not implemented)	437
Mupad [B] (verification not implemented)	438
Reduce [B] (verification not implemented)	438

Optimal result

Integrand size = 20, antiderivative size = 290

$$\begin{aligned}
 \int (a + bx)^6 (A + Bx)(d + ex)^6 dx = & \frac{(Ab - aB)(bd - ae)^6 (a + bx)^7}{7b^8} \\
 & + \frac{(bd - ae)^5 (bBd + 6Abe - 7aBe)(a + bx)^8}{8b^8} \\
 & + \frac{e(bd - ae)^4 (2bBd + 5Abe - 7aBe)(a + bx)^9}{3b^8} \\
 & + \frac{e^2 (bd - ae)^3 (3bBd + 4Abe - 7aBe)(a + bx)^{10}}{2b^8} \\
 & + \frac{5e^3 (bd - ae)^2 (4bBd + 3Abe - 7aBe)(a + bx)^{11}}{11b^8} \\
 & + \frac{e^4 (bd - ae) (5bBd + 2Abe - 7aBe)(a + bx)^{12}}{4b^8} \\
 & + \frac{e^5 (6bBd + Abe - 7aBe)(a + bx)^{13}}{13b^8} \\
 & + \frac{Be^6 (a + bx)^{14}}{14b^8}
 \end{aligned}$$

output

$$\begin{aligned}
& 1/7*(A*b-B*a)*(-a*e+b*d)^6*(b*x+a)^7/b^8+1/8*(-a*e+b*d)^5*(6*A*b*e-7*B*a*e \\
& +B*b*d)*(b*x+a)^8/b^8+1/3*e*(-a*e+b*d)^4*(5*A*b*e-7*B*a*e+2*B*b*d)*(b*x+a) \\
& ^9/b^8+1/2*e^2*(-a*e+b*d)^3*(4*A*b*e-7*B*a*e+3*B*b*d)*(b*x+a)^{10}/b^8+5/11* \\
& e^3*(-a*e+b*d)^2*(3*A*b*e-7*B*a*e+4*B*b*d)*(b*x+a)^{11}/b^8+1/4*e^4*(-a*e+b* \\
& d)*(2*A*b*e-7*B*a*e+5*B*b*d)*(b*x+a)^{12}/b^8+1/13*e^5*(A*b*e-7*B*a*e+6*B*b* \\
& d)*(b*x+a)^{13}/b^8+1/14*B*e^6*(b*x+a)^{14}/b^8
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1069 vs. $2(290) = 580$.

Time = 0.23 (sec) , antiderivative size = 1069, normalized size of antiderivative = 3.69

$$\begin{aligned}
\int (a + bx)^6 (A + Bx)(d + ex)^6 dx = & a^6 A d^6 x + \frac{1}{2} a^5 d^5 (a B d + 6 A (b d + a e)) x^2 \\
& + a^4 d^4 (2 a B d (b d + a e) \\
& \quad + A (5 b^2 d^2 + 12 a b d e + 5 a^2 e^2)) x^3 \\
& + \frac{1}{4} a^3 d^3 (3 a B d (5 b^2 d^2 + 12 a b d e + 5 a^2 e^2) \\
& \quad + 10 A (2 b^3 d^3 + 9 a b^2 d^2 e + 9 a^2 b d e^2 + 2 a^3 e^3)) x^4 \\
& + a^2 d^2 (2 a B d (2 b^3 d^3 + 9 a b^2 d^2 e + 9 a^2 b d e^2 + 2 a^3 e^3) \\
& \quad + 3 A (b^4 d^4 + 8 a b^3 d^3 e + 15 a^2 b^2 d^2 e^2 + 8 a^3 b d e^3 \\
& \quad \quad + a^4 e^4)) x^5 \\
& + \frac{1}{2} a d (5 a B d (b^4 d^4 + 8 a b^3 d^3 e + 15 a^2 b^2 d^2 e^2 \\
& \quad + 8 a^3 b d e^3 + a^4 e^4) + 2 A (b^5 d^5 + 15 a b^4 d^4 e \\
& \quad + 50 a^2 b^3 d^3 e^2 + 50 a^3 b^2 d^2 e^3 + 15 a^4 b d e^4 + a^5 e^5)) x^6 \\
& + \frac{1}{7} (6 a B d (b^5 d^5 + 15 a b^4 d^4 e + 50 a^2 b^3 d^3 e^2 \\
& \quad + 50 a^3 b^2 d^2 e^3 + 15 a^4 b d e^4 + a^5 e^5) \\
& \quad + A (b^6 d^6 + 36 a b^5 d^5 e + 225 a^2 b^4 d^4 e^2 + 400 a^3 b^3 d^3 e^3 \\
& \quad + 225 a^4 b^2 d^2 e^4 + 36 a^5 b d e^5 + a^6 e^6)) x^7 \\
& + \frac{1}{8} (a^6 B e^6 + 6 a^5 b e^5 (6 B d + A e) \\
& \quad + 45 a^4 b^2 d e^4 (5 B d + 2 A e) + 100 a^3 b^3 d^2 e^3 (4 B d + 3 A e) \\
& \quad + 75 a^2 b^4 d^3 e^2 (3 B d + 4 A e) + 18 a b^5 d^4 e (2 B d + 5 A e) \\
& \quad \quad + b^6 d^5 (B d + 6 A e)) x^8 \\
& + \frac{1}{3} b e (2 a^5 B e^5 + 5 a^4 b e^4 (6 B d + A e) \\
& \quad + 20 a^3 b^2 d e^3 (5 B d + 2 A e) + 25 a^2 b^3 d^2 e^2 (4 B d + 3 A e) \\
& \quad + 10 a b^4 d^3 e (3 B d + 4 A e) + b^5 d^4 (2 B d + 5 A e)) x^9 \\
& + \frac{1}{2} b^2 e^2 (3 a^4 B e^4 + 4 a^3 b e^3 (6 B d + A e) \\
& \quad + 9 a^2 b^2 d e^2 (5 B d + 2 A e) + 6 a b^3 d^2 e (4 B d + 3 A e) \\
& \quad \quad + b^4 d^3 (3 B d + 4 A e)) x^{10} \\
& + \frac{1}{11} b^3 e^3 (20 a^3 B e^3 + 15 a^2 b e^2 (6 B d + A e) \\
& \quad + 18 a b^2 d e (5 B d + 2 A e) + 5 b^3 d^2 (4 B d + 3 A e)) x^{11} \\
& + \frac{1}{4} b^4 e^4 (5 a^2 B e^2 + 2 a b e (6 B d + A e) \\
& \quad \quad + b^2 d (5 B d + 2 A e)) x^{12} \\
& + \frac{1}{13} b^5 e^5 (6 b B d + A b e + 6 a B e) x^{13} + \frac{1}{14} b^6 B e^6 x^{14}
\end{aligned}$$

input `Integrate[(a + b*x)^6*(A + B*x)*(d + e*x)^6,x]`

output

$$\begin{aligned} & a^6 A d^6 x + (a^5 d^5 (a B d + 6 A (b d + a e)) x^2) / 2 + a^4 d^4 (2 a B d \\ & * (b d + a e) + A (5 b^2 d^2 + 12 a b d e + 5 a^2 e^2)) x^3 + (a^3 d^3 (3 a \\ & * B d (5 b^2 d^2 + 12 a b d e + 5 a^2 e^2) + 10 A (2 b^3 d^3 + 9 a b^2 d^2 * \\ & e + 9 a^2 b d e^2 + 2 a^3 e^3)) x^4) / 4 + a^2 d^2 (2 a B d (2 b^3 d^3 + 9 a \\ & * b^2 d^2 e + 9 a^2 b d e^2 + 2 a^3 e^3) + 3 A (b^4 d^4 + 8 a b^3 d^3 e + 1 \\ & 5 a^2 b^2 d^2 e^2 + 8 a^3 b d e^3 + a^4 e^4)) x^5 + (a d (5 a B d (b^4 d^4 \\ & + 8 a b^3 d^3 e + 15 a^2 b^2 d^2 e^2 + 8 a^3 b d e^3 + a^4 e^4) + 2 A (b^ \\ & 5 d^5 + 15 a b^4 d^4 e + 50 a^2 b^3 d^3 e^2 + 50 a^3 b^2 d^2 e^3 + 15 a^4 * \\ & b d e^4 + a^5 e^5)) x^6) / 2 + ((6 a B d (b^5 d^5 + 15 a b^4 d^4 e + 50 a^2 * \\ & b^3 d^3 e^2 + 50 a^3 b^2 d^2 e^3 + 15 a^4 b d e^4 + a^5 e^5) + A (b^6 d^6 \\ & + 36 a b^5 d^5 e + 225 a^2 b^4 d^4 e^2 + 400 a^3 b^3 d^3 e^3 + 225 a^4 b^2 \\ & * d^2 e^4 + 36 a^5 b d e^5 + a^6 e^6)) x^7) / 7 + ((a^6 B e^6 + 6 a^5 b e^5 * \\ & (6 B d + A e) + 45 a^4 b^2 d e^4 (5 B d + 2 A e) + 100 a^3 b^3 d^2 e^3 (4 B \\ & * d + 3 A e) + 75 a^2 b^4 d^3 e^2 (3 B d + 4 A e) + 18 a b^5 d^4 e (2 B d + \\ & 5 A e) + b^6 d^5 (B d + 6 A e)) x^8) / 8 + (b e (2 a^5 B e^5 + 5 a^4 b e^4 * \\ & (6 B d + A e) + 20 a^3 b^2 d e^3 (5 B d + 2 A e) + 25 a^2 b^3 d^2 e^2 (4 B \\ & * d + 3 A e) + 10 a b^4 d^3 e (3 B d + 4 A e) + b^5 d^4 (2 B d + 5 A e)) x^ \\ & 9) / 3 + (b^2 e^2 (3 a^4 B e^4 + 4 a^3 b e^3 (6 B d + A e) + 9 a^2 b^2 d e^2 \\ & * (5 B d + 2 A e) + 6 a b^3 d^2 e (4 B d + 3 A e) + b^4 d^3 (3 B d + 4 A e) \\ &) x^10) / 2 + (b^3 e^3 (20 a^3 B e^3 + 15 a^2 b e^2 (6 B d + A e) + 18 a * \dots \end{aligned}$$

Rubi [A] (verified)

Time = 1.31 (sec) , antiderivative size = 290, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^6 (A + Bx) (d + ex)^6 dx$$

↓ 86

$$\int \left(\frac{e^5 (a + bx)^{12} (-7aBe + Abe + 6bBd)}{b^7} + \frac{3e^4 (a + bx)^{11} (bd - ae) (-7aBe + 2Abe + 5bBd)}{b^7} + \frac{5e^3 (a + bx)^{10} (ba}{b^7} \right) dx$$

$$\begin{aligned}
 & \downarrow 2009 \\
 & \frac{e^5(a+bx)^{13}(-7aBe + Abe + 6bBd)}{13b^8} + \frac{e^4(a+bx)^{12}(bd-ae)(-7aBe + 2Abe + 5bBd)}{4b^8} + \\
 & \frac{5e^3(a+bx)^{11}(bd-ae)^2(-7aBe + 3Abe + 4bBd)}{11b^8} + \\
 & \frac{e^2(a+bx)^{10}(bd-ae)^3(-7aBe + 4Abe + 3bBd)}{2b^8} + \\
 & \frac{e(a+bx)^9(bd-ae)^4(-7aBe + 5Abe + 2bBd)}{3b^8} + \\
 & \frac{(a+bx)^8(bd-ae)^5(-7aBe + 6Abe + bBd)}{8b^8} + \frac{(a+bx)^7(Ab-aB)(bd-ae)^6}{7b^8} + \frac{Be^6(a+bx)^{14}}{14b^8}
 \end{aligned}$$

input `Int[(a + b*x)^6*(A + B*x)*(d + e*x)^6,x]`

output `((A*b - a*B)*(b*d - a*e)^6*(a + b*x)^7)/(7*b^8) + ((b*d - a*e)^5*(b*B*d + 6*A*b*e - 7*a*B*e)*(a + b*x)^8)/(8*b^8) + (e*(b*d - a*e)^4*(2*b*B*d + 5*A*b*e - 7*a*B*e)*(a + b*x)^9)/(3*b^8) + (e^2*(b*d - a*e)^3*(3*b*B*d + 4*A*b*e - 7*a*B*e)*(a + b*x)^10)/(2*b^8) + (5*e^3*(b*d - a*e)^2*(4*b*B*d + 3*A*b*e - 7*a*B*e)*(a + b*x)^11)/(11*b^8) + (e^4*(b*d - a*e)*(5*b*B*d + 2*A*b*e - 7*a*B*e)*(a + b*x)^12)/(4*b^8) + (e^5*(6*b*B*d + A*b*e - 7*a*B*e)*(a + b*x)^13)/(13*b^8) + (B*e^6*(a + b*x)^14)/(14*b^8)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1172 vs. $2(274) = 548$.

Time = 0.24 (sec) , antiderivative size = 1173, normalized size of antiderivative = 4.04

method	result	size
default	Expression too large to display	1173
norman	Expression too large to display	1253
oring	Expression too large to display	1480
gosper	Expression too large to display	1481
risch	Expression too large to display	1481
parallelrisc	Expression too large to display	1481

input `int((b*x+a)^6*(B*x+A)*(e*x+d)^6,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 1/14*b^6*B*e^6*x^{14}+1/13*((A*b^6+6*B*a*b^5)*e^6+6*b^6*B*d*e^5)*x^{13}+1/12*(\\ & (6*A*a*b^5+15*B*a^2*b^4)*e^6+6*(A*b^6+6*B*a*b^5)*d*e^5+15*b^6*B*d^2*e^4)*x \\ & ^{12}+1/11*((15*A*a^2*b^4+20*B*a^3*b^3)*e^6+6*(6*A*a*b^5+15*B*a^2*b^4)*d*e^5 \\ & +15*(A*b^6+6*B*a*b^5)*d^2*e^4+20*b^6*B*d^3*e^3)*x^{11}+1/10*((20*A*a^3*b^3+1 \\ & 5*B*a^4*b^2)*e^6+6*(15*A*a^2*b^4+20*B*a^3*b^3)*d*e^5+15*(6*A*a*b^5+15*B*a^ \\ & 2*b^4)*d^2*e^4+20*(A*b^6+6*B*a*b^5)*d^3*e^3+15*b^6*B*d^4*e^2)*x^{10}+1/9*((1 \\ & 5*A*a^4*b^2+6*B*a^5*b)*e^6+6*(20*A*a^3*b^3+15*B*a^4*b^2)*d*e^5+15*(15*A*a^ \\ & 2*b^4+20*B*a^3*b^3)*d^2*e^4+20*(6*A*a*b^5+15*B*a^2*b^4)*d^3*e^3+15*(A*b^6+ \\ & 6*B*a*b^5)*d^4*e^2+6*b^6*B*d^5*e)*x^9+1/8*((6*A*a^5*b+B*a^6)*e^6+6*(15*A*a \\ & ^4*b^2+6*B*a^5*b)*d*e^5+15*(20*A*a^3*b^3+15*B*a^4*b^2)*d^2*e^4+20*(15*A*a^ \\ & 2*b^4+20*B*a^3*b^3)*d^3*e^3+15*(6*A*a*b^5+15*B*a^2*b^4)*d^4*e^2+6*(A*b^6+6 \\ & *B*a*b^5)*d^5*e+b^6*B*d^6)*x^8+1/7*(a^6*A*e^6+6*(6*A*a^5*b+B*a^6)*d*e^5+15 \\ & *(15*A*a^4*b^2+6*B*a^5*b)*d^2*e^4+20*(20*A*a^3*b^3+15*B*a^4*b^2)*d^3*e^3+1 \\ & 5*(15*A*a^2*b^4+20*B*a^3*b^3)*d^4*e^2+6*(6*A*a*b^5+15*B*a^2*b^4)*d^5*e+(A* \\ & b^6+6*B*a*b^5)*d^6)*x^7+1/6*(6*a^6*A*d*e^5+15*(6*A*a^5*b+B*a^6)*d^2*e^4+20 \\ & *(15*A*a^4*b^2+6*B*a^5*b)*d^3*e^3+15*(20*A*a^3*b^3+15*B*a^4*b^2)*d^4*e^2+6 \\ & *(15*A*a^2*b^4+20*B*a^3*b^3)*d^5*e+(6*A*a*b^5+15*B*a^2*b^4)*d^6)*x^6+1/5*(\\ & 15*a^6*A*d^2*e^4+20*(6*A*a^5*b+B*a^6)*d^3*e^3+15*(15*A*a^4*b^2+6*B*a^5*b)* \\ & d^4*e^2+6*(20*A*a^3*b^3+15*B*a^4*b^2)*d^5*e+(15*A*a^2*b^4+20*B*a^3*b^3)*d^ \\ & 6)*x^5+1/4*(20*a^6*A*d^3*e^3+15*(6*A*a^5*b+B*a^6)*d^4*e^2+6*(15*A*a^4*b^2+6*B*a^5*b)*d^5*e+ \\ & 15*(15*A*a^2*b^4+20*B*a^3*b^3)*d^6)*x^4+1/3*(6*a^6*A*d^4*e^2+15*(6*A*a^5*b+B*a^6)*d^5*e+ \\ & 15*(15*A*a^4*b^2+6*B*a^5*b)*d^6)*x^3+1/2*(3*a^6*A*d^5*e+15*(6*A*a^5*b+B*a^6)*d^6)*x^2+ \\ & 1/2*(3*a^6*A*d^6)*x+1/2*(3*a^6*A*d^6)*x+1/2*(3*a^6*A*d^6) \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1172 vs. $2(274) = 548$.

Time = 0.08 (sec) , antiderivative size = 1172, normalized size of antiderivative = 4.04

$$\int (a + bx)^6 (A + Bx)(d + ex)^6 dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)*(e*x+d)^6,x, algorithm="fricas")`

output

```
1/14*B*b^6*e^6*x^14 + A*a^6*d^6*x + 1/13*(6*B*b^6*d*e^5 + (6*B*a*b^5 + A*b^6)*e^6)*x^13 + 1/4*(5*B*b^6*d^2*e^4 + 2*(6*B*a*b^5 + A*b^6)*d*e^5 + (5*B*a^2*b^4 + 2*A*a*b^5)*e^6)*x^12 + 1/11*(20*B*b^6*d^3*e^3 + 15*(6*B*a*b^5 + A*b^6)*d^2*e^4 + 18*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^5 + 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^6)*x^11 + 1/2*(3*B*b^6*d^4*e^2 + 4*(6*B*a*b^5 + A*b^6)*d^3*e^3 + 9*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^4 + 6*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^5 + (3*B*a^4*b^2 + 4*A*a^3*b^3)*e^6)*x^10 + 1/3*(2*B*b^6*d^5*e + 5*(6*B*a*b^5 + A*b^6)*d^4*e^2 + 20*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*e^3 + 25*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^4 + 10*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^5 + (2*B*a^5*b + 5*A*a^4*b^2)*e^6)*x^9 + 1/8*(B*b^6*d^6 + 6*(6*B*a*b^5 + A*b^6)*d^5*e + 45*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^2 + 100*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^3 + 75*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^4 + 18*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^5 + (B*a^6 + 6*A*a^5*b)*e^6)*x^8 + 1/7*(A*a^6*e^6 + (6*B*a*b^5 + A*b^6)*d^6 + 18*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e + 75*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4*e^2 + 100*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^3 + 45*(2*B*a^5*b + 5*A*a^4*b^2)*d^2*e^4 + 6*(B*a^6 + 6*A*a^5*b)*d*e^5)*x^7 + 1/2*(2*A*a^6*d*e^5 + (5*B*a^2*b^4 + 2*A*a*b^5)*d^6 + 10*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^5*e + 25*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^4*e^2 + 20*(2*B*a^5*b + 5*A*a^4*b^2)*d^3*e^3 + 5*(B*a^6 + 6*A*a^5*b)*d^2*e^4)*x^6 + (3*A*a^6*d^2*e^4 + (4*B*a^3*b^3 + 3*A*a^2*b^4)*d^6 + 6*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^5*e + 9*(2*...
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1504 vs. $2(292) = 584$.

Time = 0.09 (sec) , antiderivative size = 1504, normalized size of antiderivative = 5.19

$$\int (a + bx)^6 (A + Bx)(d + ex)^6 dx = \text{Too large to display}$$

input `integrate((b*x+a)**6*(B*x+A)*(e*x+d)**6,x)`

output

```
A*a**6*d**6*x + B*b**6*e**6*x**14/14 + x**13*(A*b**6*e**6/13 + 6*B*a*b**5*
e**6/13 + 6*B*b**6*d*e**5/13) + x**12*(A*a*b**5*e**6/2 + A*b**6*d*e**5/2 +
5*B*a**2*b**4*e**6/4 + 3*B*a*b**5*d*e**5 + 5*B*b**6*d**2*e**4/4) + x**11*
(15*A*a**2*b**4*e**6/11 + 36*A*a*b**5*d*e**5/11 + 15*A*b**6*d**2*e**4/11 +
20*B*a**3*b**3*e**6/11 + 90*B*a**2*b**4*d*e**5/11 + 90*B*a*b**5*d**2*e**4
/11 + 20*B*b**6*d**3*e**3/11) + x**10*(2*A*a**3*b**3*e**6 + 9*A*a**2*b**4*
d*e**5 + 9*A*a*b**5*d**2*e**4 + 2*A*b**6*d**3*e**3 + 3*B*a**4*b**2*e**6/2
+ 12*B*a**3*b**3*d*e**5 + 45*B*a**2*b**4*d**2*e**4/2 + 12*B*a*b**5*d**3*e*
**3 + 3*B*b**6*d**4*e**2/2) + x**9*(5*A*a**4*b**2*e**6/3 + 40*A*a**3*b**3*d
e**5/3 + 25*A*a**2*b**4*d**2*e**4 + 40*A*a*b**5*d**3*e**3/3 + 5*A*b**6*d*
**4*e**2/3 + 2*B*a**5*b*e**6/3 + 10*B*a**4*b**2*d*e**5 + 100*B*a**3*b**3*d*
**2*e**4/3 + 100*B*a**2*b**4*d**3*e**3/3 + 10*B*a*b**5*d**4*e**2 + 2*B*b**6
*d**5*e/3) + x**8*(3*A*a**5*b*e**6/4 + 45*A*a**4*b**2*d*e**5/4 + 75*A*a**3
*b**3*d**2*e**4/2 + 75*A*a**2*b**4*d**3*e**3/2 + 45*A*a*b**5*d**4*e**2/4 +
3*A*b**6*d**5*e/4 + B*a**6*e**6/8 + 9*B*a**5*b*d*e**5/2 + 225*B*a**4*b**2
*d**2*e**4/8 + 50*B*a**3*b**3*d**3*e**3 + 225*B*a**2*b**4*d**4*e**2/8 + 9*
B*a*b**5*d**5*e/2 + B*b**6*d**6/8) + x**7*(A*a**6*e**6/7 + 36*A*a**5*b*d*e
**5/7 + 225*A*a**4*b**2*d**2*e**4/7 + 400*A*a**3*b**3*d**3*e**3/7 + 225*A*
a**2*b**4*d**4*e**2/7 + 36*A*a*b**5*d**5*e/7 + A*b**6*d**6/7 + 6*B*a**6*d*
e**5/7 + 90*B*a**5*b*d**2*e**4/7 + 300*B*a**4*b**2*d**3*e**3/7 + 300*B*...
```


Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1172 vs. $2(274) = 548$.

Time = 0.04 (sec) , antiderivative size = 1172, normalized size of antiderivative = 4.04

$$\int (a + bx)^6 (A + Bx)(d + ex)^6 dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)*(e*x+d)^6,x, algorithm="maxima")`

output

```
1/14*B*b^6*e^6*x^14 + A*a^6*d^6*x + 1/13*(6*B*b^6*d*e^5 + (6*B*a*b^5 + A*b^6)*e^6)*x^13 + 1/4*(5*B*b^6*d^2*e^4 + 2*(6*B*a*b^5 + A*b^6)*d*e^5 + (5*B*a^2*b^4 + 2*A*a*b^5)*e^6)*x^12 + 1/11*(20*B*b^6*d^3*e^3 + 15*(6*B*a*b^5 + A*b^6)*d^2*e^4 + 18*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^5 + 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^6)*x^11 + 1/2*(3*B*b^6*d^4*e^2 + 4*(6*B*a*b^5 + A*b^6)*d^3*e^3 + 9*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^4 + 6*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^5 + (3*B*a^4*b^2 + 4*A*a^3*b^3)*e^6)*x^10 + 1/3*(2*B*b^6*d^5*e + 5*(6*B*a*b^5 + A*b^6)*d^4*e^2 + 20*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*e^3 + 25*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^4 + 10*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^5 + (2*B*a^5*b + 5*A*a^4*b^2)*e^6)*x^9 + 1/8*(B*b^6*d^6 + 6*(6*B*a*b^5 + A*b^6)*d^5*e + 45*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^2 + 100*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^3 + 75*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^4 + 18*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^5 + (B*a^6 + 6*A*a^5*b)*e^6)*x^8 + 1/7*(A*a^6*e^6 + (6*B*a*b^5 + A*b^6)*d^6 + 18*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e + 75*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4*e^2 + 100*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^3 + 45*(2*B*a^5*b + 5*A*a^4*b^2)*d^2*e^4 + 6*(B*a^6 + 6*A*a^5*b)*d*e^5)*x^7 + 1/2*(2*A*a^6*d*e^5 + (5*B*a^2*b^4 + 2*A*a*b^5)*d^6 + 10*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^5*e + 25*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^4*e^2 + 20*(2*B*a^5*b + 5*A*a^4*b^2)*d^3*e^3 + 5*(B*a^6 + 6*A*a^5*b)*d^2*e^4)*x^6 + (3*A*a^6*d^2*e^4 + (4*B*a^3*b^3 + 3*A*a^2*b^4)*d^6 + 6*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^5*e + 9*(2*...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1480 vs. $2(274) = 548$.

Time = 0.12 (sec) , antiderivative size = 1480, normalized size of antiderivative = 5.10

$$\int (a + bx)^6 (A + Bx)(d + ex)^6 dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)*(e*x+d)^6,x, algorithm="giac")`

output

```
1/14*B*b^6*e^6*x^14 + 6/13*B*b^6*d*e^5*x^13 + 6/13*B*a*b^5*e^6*x^13 + 1/13
*A*b^6*e^6*x^13 + 5/4*B*b^6*d^2*e^4*x^12 + 3*B*a*b^5*d*e^5*x^12 + 1/2*A*b^
6*d*e^5*x^12 + 5/4*B*a^2*b^4*e^6*x^12 + 1/2*A*a*b^5*e^6*x^12 + 20/11*B*b^6
*d^3*e^3*x^11 + 90/11*B*a*b^5*d^2*e^4*x^11 + 15/11*A*b^6*d^2*e^4*x^11 + 90
/11*B*a^2*b^4*d*e^5*x^11 + 36/11*A*a*b^5*d*e^5*x^11 + 20/11*B*a^3*b^3*e^6*
x^11 + 15/11*A*a^2*b^4*e^6*x^11 + 3/2*B*b^6*d^4*e^2*x^10 + 12*B*a*b^5*d^3*
e^3*x^10 + 2*A*b^6*d^3*e^3*x^10 + 45/2*B*a^2*b^4*d^2*e^4*x^10 + 9*A*a*b^5*
d^2*e^4*x^10 + 12*B*a^3*b^3*d*e^5*x^10 + 9*A*a^2*b^4*d*e^5*x^10 + 3/2*B*a^
4*b^2*e^6*x^10 + 2*A*a^3*b^3*e^6*x^10 + 2/3*B*b^6*d^5*e*x^9 + 10*B*a*b^5*d
^4*e^2*x^9 + 5/3*A*b^6*d^4*e^2*x^9 + 100/3*B*a^2*b^4*d^3*e^3*x^9 + 40/3*A*
a*b^5*d^3*e^3*x^9 + 100/3*B*a^3*b^3*d^2*e^4*x^9 + 25*A*a^2*b^4*d^2*e^4*x^9
+ 10*B*a^4*b^2*d*e^5*x^9 + 40/3*A*a^3*b^3*d*e^5*x^9 + 2/3*B*a^5*b*e^6*x^9
+ 5/3*A*a^4*b^2*e^6*x^9 + 1/8*B*b^6*d^6*x^8 + 9/2*B*a*b^5*d^5*e*x^8 + 3/4
*A*b^6*d^5*e*x^8 + 225/8*B*a^2*b^4*d^4*e^2*x^8 + 45/4*A*a*b^5*d^4*e^2*x^8
+ 50*B*a^3*b^3*d^3*e^3*x^8 + 75/2*A*a^2*b^4*d^3*e^3*x^8 + 225/8*B*a^4*b^2*
d^2*e^4*x^8 + 75/2*A*a^3*b^3*d^2*e^4*x^8 + 9/2*B*a^5*b*d*e^5*x^8 + 45/4*A*
a^4*b^2*d*e^5*x^8 + 1/8*B*a^6*e^6*x^8 + 3/4*A*a^5*b*e^6*x^8 + 6/7*B*a*b^5*
d^6*x^7 + 1/7*A*b^6*d^6*x^7 + 90/7*B*a^2*b^4*d^5*e*x^7 + 36/7*A*a*b^5*d^5*
e*x^7 + 300/7*B*a^3*b^3*d^4*e^2*x^7 + 225/7*A*a^2*b^4*d^4*e^2*x^7 + 300/7*
B*a^4*b^2*d^3*e^3*x^7 + 400/7*A*a^3*b^3*d^3*e^3*x^7 + 90/7*B*a^5*b*d^2*...
```

Mupad [B] (verification not implemented)

Time = 1.15 (sec) , antiderivative size = 1221, normalized size of antiderivative = 4.21

$$\int (a + bx)^6 (A + Bx)(d + ex)^6 dx = \text{Too large to display}$$

input `int((A + B*x)*(a + b*x)^6*(d + e*x)^6,x)`

output

$$\begin{aligned} & x^5(3Aa^2b^4d^6 + 4B^3a^3b^3d^6 + 3Aa^6d^2e^4 + 4B^6a^6d^3e^3 \\ & + 24Aa^3b^3d^5e + 24Aa^5b^3d^3e^3 + 18B^4a^4b^2d^5e + 18B^5 \\ & *b^4d^4e^2 + 45Aa^4b^2d^4e^2) + x^{10}(2Aa^3b^3e^6 + (3B^4a^4b^2 \\ & e^6)/2 + 2Ab^6d^3e^3 + (3B^6d^4e^2)/2 + 9Aa^2b^5d^2e^4 + 9Aa \\ & ^2b^4d^5e + 12B^5a^3b^3d^3e^3 + 12B^6a^3b^3d^5e + (45B^4a^2b^4d^2 \\ & *e^4)/2) + x^6(Aa^2b^5d^6 + Aa^6d^5e + (5B^6a^2b^4d^6)/2 + (5B^6 \\ & *d^2e^4)/2 + 15Aa^2b^4d^5e + 15Aa^5b^4d^2e^4 + 20B^6a^3b^3d^5e \\ & + 20B^6a^5b^3d^3e^3 + 50Aa^3b^3d^4e^2 + 50Aa^4b^2d^3e^3 + (75 \\ & B^4a^4b^2d^4e^2)/2) + x^9((2B^5b^6e^6)/3 + (2B^6d^5e)/3 + (5A \\ & a^4b^2e^6)/3 + (5A^6b^6d^4e^2)/3 + (40Aa^2b^5d^3e^3)/3 + (40Aa^3 \\ & b^3d^5e^5)/3 + 10B^6a^2b^4d^4e^2 + 10B^6a^4b^2d^5e + 25Aa^2b^4d^2 \\ & e^4 + (100B^6a^2b^4d^3e^3)/3 + (100B^6a^3b^3d^2e^4)/3) + x^7((Aa^6 \\ & e^6)/7 + (A^6b^6d^6)/7 + (6B^6a^5b^5d^6)/7 + (6B^6a^6d^5e^5)/7 + (90B^6 \\ & a^2b^4d^5e)/7 + (90B^6a^5b^4d^2e^4)/7 + (225Aa^2b^4d^4e^2)/7 + (400 \\ & *Aa^3b^3d^3e^3)/7 + (225Aa^4b^2d^2e^4)/7 + (300B^6a^3b^3d^4e^2 \\ &)/7 + (300B^6a^4b^2d^3e^3)/7 + (36Aa^2b^5d^5e)/7 + (36Aa^5b^4d^5 \\ & e^5)/7) + x^4(5Aa^3b^3d^6 + (15B^6a^4b^2d^6)/4 + 5Aa^6d^3e^3 + (15 \\ & *B^6a^6d^4e^2)/4 + (45Aa^4b^2d^5e)/2 + (45Aa^5b^4d^4e^2)/2 + 9B^6 \\ & a^5b^4d^5e) + x^8((B^6a^6e^6)/8 + (B^6b^6d^6)/8 + (3Aa^5b^6e^6)/4 + (3 \\ & *A^6b^6d^5e^5)/4 + (45Aa^2b^5d^4e^2)/4 + (45Aa^4b^2d^5e^5)/4 + (75... \end{aligned}$$
Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 799, normalized size of antiderivative = 2.76

$$\int (a + bx)^6 (A + Bx)(d + ex)^6 dx$$

$$= \frac{x(1716b^7e^6x^{13} + 12936ab^6e^6x^{12} + 11088b^7de^5x^{12} + 42042a^2b^5e^6x^{11} + 84084ab^6de^5x^{11} + 30030b^7d^2e^4x^{10} + \dots)}{1}$$

input `int((b*x+a)^6*(B*x+A)*(e*x+d)^6,x)`

output

```
(x*(24024*a**7*d**6 + 72072*a**7*d**5*e*x + 120120*a**7*d**4*e**2*x**2 + 1
20120*a**7*d**3*e**3*x**3 + 72072*a**7*d**2*e**4*x**4 + 24024*a**7*d*e**5*
x**5 + 3432*a**7*e**6*x**6 + 84084*a**6*b*d**6*x + 336336*a**6*b*d**5*e*x*
*2 + 630630*a**6*b*d**4*e**2*x**3 + 672672*a**6*b*d**3*e**3*x**4 + 420420*
a**6*b*d**2*e**4*x**5 + 144144*a**6*b*d*e**5*x**6 + 21021*a**6*b*e**6*x**7
+ 168168*a**5*b**2*d**6*x**2 + 756756*a**5*b**2*d**5*e*x**3 + 1513512*a**
5*b**2*d**4*e**2*x**4 + 1681680*a**5*b**2*d**3*e**3*x**5 + 1081080*a**5*b*
*2*d**2*e**4*x**6 + 378378*a**5*b**2*d*e**5*x**7 + 56056*a**5*b**2*e**6*x*
*8 + 210210*a**4*b**3*d**6*x**3 + 1009008*a**4*b**3*d**5*e*x**4 + 2102100*
a**4*b**3*d**4*e**2*x**5 + 2402400*a**4*b**3*d**3*e**3*x**6 + 1576575*a**4
*b**3*d**2*e**4*x**7 + 560560*a**4*b**3*d*e**5*x**8 + 84084*a**4*b**3*e**6
*x**9 + 168168*a**3*b**4*d**6*x**4 + 840840*a**3*b**4*d**5*e*x**5 + 180180
0*a**3*b**4*d**4*e**2*x**6 + 2102100*a**3*b**4*d**3*e**3*x**7 + 1401400*a*
*3*b**4*d**2*e**4*x**8 + 504504*a**3*b**4*d*e**5*x**9 + 76440*a**3*b**4*e*
*6*x**10 + 84084*a**2*b**5*d**6*x**5 + 432432*a**2*b**5*d**5*e*x**6 + 9459
45*a**2*b**5*d**4*e**2*x**7 + 1121120*a**2*b**5*d**3*e**3*x**8 + 756756*a*
*2*b**5*d**2*e**4*x**9 + 275184*a**2*b**5*d*e**5*x**10 + 42042*a**2*b**5*e
**6*x**11 + 24024*a*b**6*d**6*x**6 + 126126*a*b**6*d**5*e*x**7 + 280280*a*
b**6*d**4*e**2*x**8 + 336336*a*b**6*d**3*e**3*x**9 + 229320*a*b**6*d**2*e*
*4*x**10 + 84084*a*b**6*d*e**5*x**11 + 12936*a*b**6*e**6*x**12 + 3003*b...
```

3.44 $\int (a + bx)^6 (A + Bx)(d + ex)^5 dx$

Optimal result	440
Mathematica [B] (verified)	441
Rubi [A] (verified)	442
Maple [B] (verified)	444
Fricas [B] (verification not implemented)	445
Sympy [B] (verification not implemented)	446
Maxima [B] (verification not implemented)	447
Giac [B] (verification not implemented)	448
Mupad [B] (verification not implemented)	449
Reduce [B] (verification not implemented)	449

Optimal result

Integrand size = 20, antiderivative size = 240

$$\int (a + bx)^6 (A + Bx)(d + ex)^5 dx = \frac{(Ab - aB)(bd - ae)^5 (a + bx)^7}{7b^7} + \frac{(bd - ae)^4 (bBd + 5Abe - 6aBe)(a + bx)^8}{8b^7} + \frac{5e(bd - ae)^3 (bBd + 2Abe - 3aBe)(a + bx)^9}{9b^7} + \frac{e^2 (bd - ae)^2 (bBd + Abe - 2aBe)(a + bx)^{10}}{b^7} + \frac{5e^3 (bd - ae)(2bBd + Abe - 3aBe)(a + bx)^{11}}{11b^7} + \frac{e^4 (5bBd + Abe - 6aBe)(a + bx)^{12}}{12b^7} + \frac{Be^5 (a + bx)^{13}}{13b^7}$$

output

```
1/7*(A*b-B*a)*(-a*e+b*d)^5*(b*x+a)^7/b^7+1/8*(-a*e+b*d)^4*(5*A*b*e-6*B*a*e+B*b*d)*(b*x+a)^8/b^7+5/9*e*(-a*e+b*d)^3*(2*A*b*e-3*B*a*e+B*b*d)*(b*x+a)^9/b^7+e^2*(-a*e+b*d)^2*(A*b*e-2*B*a*e+B*b*d)*(b*x+a)^10/b^7+5/11*e^3*(-a*e+b*d)*(A*b*e-3*B*a*e+2*B*b*d)*(b*x+a)^11/b^7+1/12*e^4*(A*b*e-6*B*a*e+5*B*b*d)*(b*x+a)^12/b^7+1/13*B*e^5*(b*x+a)^13/b^7
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 907 vs. $2(240) = 480$.

Time = 0.20 (sec) , antiderivative size = 907, normalized size of antiderivative = 3.78

$$\begin{aligned}
\int (a + bx)^6(A + Bx)(d + ex)^5 dx = & a^6 Ad^5 x + \frac{1}{2} a^5 d^4 (6Abd + aBd + 5aAe)x^2 \\
& + \frac{1}{3} a^4 d^3 (aBd(6bd + 5ae) \\
& \qquad \qquad \qquad + 5A(3b^2 d^2 + 6abde + 2a^2 e^2)) x^3 \\
& + \frac{5}{4} a^3 d^2 (aBd(3b^2 d^2 + 6abde + 2a^2 e^2) \\
& \qquad \qquad \qquad + A(4b^3 d^3 + 15ab^2 d^2 e + 12a^2 bde^2 + 2a^3 e^3)) x^4 \\
& + a^2 d (aBd(4b^3 d^3 + 15ab^2 d^2 e + 12a^2 bde^2 + 2a^3 e^3) \\
& \qquad \qquad \qquad + A(3b^4 d^4 + 20ab^3 d^3 e + 30a^2 b^2 d^2 e^2 + 12a^3 bde^3 \\
& \qquad \qquad \qquad \qquad \qquad \qquad + a^4 e^4)) x^5 \\
& + \frac{1}{6} a (5aBd(3b^4 d^4 + 20ab^3 d^3 e + 30a^2 b^2 d^2 e^2 \\
& \qquad \qquad \qquad + 12a^3 bde^3 + a^4 e^4) + A(6b^5 d^5 + 75ab^4 d^4 e \\
& \qquad \qquad \qquad + 200a^2 b^3 d^3 e^2 + 150a^3 b^2 d^2 e^3 + 30a^4 bde^4 + a^5 e^5)) x^6 \\
& + \frac{1}{7} (aB(6b^5 d^5 + 75ab^4 d^4 e + 200a^2 b^3 d^3 e^2 \\
& \qquad \qquad \qquad + 150a^3 b^2 d^2 e^3 + 30a^4 bde^4 + a^5 e^5) \\
& \qquad \qquad \qquad + Ab(b^5 d^5 + 30ab^4 d^4 e + 150a^2 b^3 d^3 e^2 \\
& \qquad \qquad \qquad + 200a^3 b^2 d^2 e^3 + 75a^4 bde^4 + 6a^5 e^5)) x^7 \\
& + \frac{1}{8} b (6a^5 Be^5 + 150a^2 b^3 d^2 e^2 (Bd + Ae) \\
& \qquad \qquad \qquad + 100a^3 b^2 de^3 (2Bd + Ae) + 15a^4 be^4 (5Bd + Ae) \\
& \qquad \qquad \qquad + 30ab^4 d^3 e (Bd + 2Ae) + b^5 d^4 (Bd + 5Ae)) x^8 \\
& + \frac{5}{9} b^2 e (3a^4 Be^4 + 12ab^3 d^2 e (Bd + Ae) \\
& \qquad \qquad \qquad + 15a^2 b^2 de^2 (2Bd + Ae) + 4a^3 be^3 (5Bd + Ae) \\
& \qquad \qquad \qquad \qquad \qquad \qquad + b^4 d^3 (Bd + 2Ae)) x^9 \\
& + \frac{1}{2} b^3 e^2 (4a^3 Be^3 + 2b^3 d^2 (Bd + Ae) \\
& \qquad \qquad \qquad + 6ab^2 de (2Bd + Ae) + 3a^2 be^2 (5Bd + Ae)) x^{10} \\
& + \frac{1}{11} b^4 e^3 (15a^2 Be^2 + 5b^2 d (2Bd + Ae) \\
& \qquad \qquad \qquad \qquad \qquad \qquad + 6abe (5Bd + Ae)) x^{11} \\
& + \frac{1}{12} b^5 e^4 (5bBd + Abe + 6aBe) x^{12} + \frac{1}{13} b^6 Be^5 x^{13}
\end{aligned}$$

input `Integrate[(a + b*x)^6*(A + B*x)*(d + e*x)^5,x]`

output

$$\begin{aligned} & a^6 A d^5 x + (a^5 d^4 (6 A b d + a B d + 5 a A e) x^2) / 2 + (a^4 d^3 (a B d (6 b d + 5 a e) + 5 A (3 b^2 d^2 + 6 a b d e + 2 a^2 e^2)) x^3) / 3 + (5 a^3 d^2 (a B d (3 b^2 d^2 + 6 a b d e + 2 a^2 e^2) + A (4 b^3 d^3 + 15 a b^2 d^2 e + 12 a^2 b d e^2 + 2 a^3 e^3)) x^4) / 4 + a^2 d (a B d (4 b^3 d^3 + 15 a b^2 d^2 e + 12 a^2 b d e^2 + 2 a^3 e^3) + A (3 b^4 d^4 + 20 a b^3 d^3 e + 30 a^2 b^2 d^2 e^2 + 12 a^3 b d e^3 + a^4 e^4)) x^5 + (a (5 a B d (3 b^4 d^4 + 20 a b^3 d^3 e + 30 a^2 b^2 d^2 e^2 + 12 a^3 b d e^3 + a^4 e^4) + A (6 b^5 d^5 + 75 a b^4 d^4 e + 200 a^2 b^3 d^3 e^2 + 150 a^3 b^2 d^2 e^3 + 30 a^4 b d e^4 + a^5 e^5)) x^6) / 6 + ((a B (6 b^5 d^5 + 75 a b^4 d^4 e + 200 a^2 b^3 d^3 e^2 + 150 a^3 b^2 d^2 e^3 + 30 a^4 b d e^4 + a^5 e^5) + A b (b^5 d^5 + 30 a b^4 d^4 e + 150 a^2 b^3 d^3 e^2 + 200 a^3 b^2 d^2 e^3 + 75 a^4 b d e^4 + 6 a^5 e^5)) x^7) / 7 + (b (6 a^5 B e^5 + 150 a^2 b^3 d^2 e^2 (B d + A e) + 100 a^3 b^2 d e^3 (2 B d + A e) + 15 a^4 b e^4 (5 B d + A e) + 30 a b^4 d^3 e (B d + 2 A e) + b^5 d^4 (B d + 5 A e)) x^8) / 8 + (5 b^2 e (3 a^4 B e^4 + 12 a b^3 d^2 e (B d + A e) + 15 a^2 b^2 d e^2 (2 B d + A e) + 4 a^3 b e^3 (5 B d + A e) + b^4 d^3 (B d + 2 A e)) x^9) / 9 + (b^3 e^2 (4 a^3 B e^3 + 2 b^3 d^2 (B d + A e) + 6 a b^2 d e (2 B d + A e) + 3 a^2 b e^2 (5 B d + A e)) x^10) / 2 + (b^4 e^3 (15 a^2 B e^2 + 5 b^2 d (2 B d + A e) + 6 a b e (5 B d + A e)) x^11) / 11 + (b^5 e^4 (5 b B d + A b e + 6 a B e) x^12) / 12 + (b^6 B e^5 x^13) / 13 \end{aligned}$$

Rubi [A] (verified)

Time = 1.06 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^6 (A + Bx)(d + ex)^5 dx$$

↓ 86

$$\int \left(\frac{e^4 (a + bx)^{11} (-6aBe + Abe + 5bBd)}{b^6} + \frac{5e^3 (a + bx)^{10} (bd - ae) (-3aBe + Abe + 2bBd)}{b^6} + \frac{10e^2 (a + bx)^9 (bd - ae)^2}{b^6} \right) dx$$

$$\begin{aligned}
 & \downarrow \text{2009} \\
 & \frac{e^4(a+bx)^{12}(-6aBe + Abe + 5bBd)}{12b^7} + \frac{5e^3(a+bx)^{11}(bd-ae)(-3aBe + Abe + 2bBd)}{11b^7} + \\
 & \frac{e^2(a+bx)^{10}(bd-ae)^2(-2aBe + Abe + bBd)}{b^7} + \\
 & \frac{5e(a+bx)^9(bd-ae)^3(-3aBe + 2Abe + bBd)}{9b^7} + \\
 & \frac{(a+bx)^8(bd-ae)^4(-6aBe + 5Abe + bBd)}{8b^7} + \frac{(a+bx)^7(Ab - aB)(bd-ae)^5}{7b^7} + \frac{Be^5(a+bx)^{13}}{13b^7}
 \end{aligned}$$

input `Int[(a + b*x)^6*(A + B*x)*(d + e*x)^5,x]`

output `((A*b - a*B)*(b*d - a*e)^5*(a + b*x)^7)/(7*b^7) + ((b*d - a*e)^4*(b*B*d + 5*A*b*e - 6*a*B*e)*(a + b*x)^8)/(8*b^7) + (5*e*(b*d - a*e)^3*(b*B*d + 2*A*b*e - 3*a*B*e)*(a + b*x)^9)/(9*b^7) + (e^2*(b*d - a*e)^2*(b*B*d + A*b*e - 2*a*B*e)*(a + b*x)^10)/b^7 + (5*e^3*(b*d - a*e)*(2*b*B*d + A*b*e - 3*a*B*e)*(a + b*x)^11)/(11*b^7) + (e^4*(5*b*B*d + A*b*e - 6*a*B*e)*(a + b*x)^12)/(12*b^7) + (B*e^5*(a + b*x)^13)/(13*b^7)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 996 vs. $2(228) = 456$.

Time = 0.21 (sec) , antiderivative size = 997, normalized size of antiderivative = 4.15

method	result
default	$\frac{b^6 B e^5 x^{13}}{13} + \frac{((b^6 A + 6a b^5 B) e^5 + 5b^6 B d e^4) x^{12}}{12} + \frac{((6a b^5 A + 15a^2 b^4 B) e^5 + 5(b^6 A + 6a b^5 B) d e^4 + 10b^6 B d^2 e^3) x^{11}}{11} + \dots$
norman	Expression too large to display
gospers	Expression too large to display
risch	Expression too large to display
paralrelrisch	Expression too large to display
orering	Expression too large to display

input

```
int((b*x+a)^6*(B*x+A)*(e*x+d)^5,x,method=_RETURNVERBOSE)
```

output

```
1/13*b^6*B*e^5*x^13+1/12*((A*b^6+6*B*a*b^5)*e^5+5*b^6*B*d*e^4)*x^12+1/11*((6*A*a*b^5+15*B*a^2*b^4)*e^5+5*(A*b^6+6*B*a*b^5)*d*e^4+10*b^6*B*d^2*e^3)*x^11+1/10*((15*A*a^2*b^4+20*B*a^3*b^3)*e^5+5*(6*A*a*b^5+15*B*a^2*b^4)*d*e^4+10*(A*b^6+6*B*a*b^5)*d^2*e^3+10*b^6*B*d^3*e^2)*x^10+1/9*((20*A*a^3*b^3+15*B*a^4*b^2)*e^5+5*(15*A*a^2*b^4+20*B*a^3*b^3)*d*e^4+10*(6*A*a*b^5+15*B*a^2*b^4)*d^2*e^3+10*(A*b^6+6*B*a*b^5)*d^3*e^2+5*b^6*B*d^4*e)*x^9+1/8*((15*A*a^4*b^2+6*B*a^5*b)*e^5+5*(20*A*a^3*b^3+15*B*a^4*b^2)*d*e^4+10*(15*A*a^2*b^4+20*B*a^3*b^3)*d^2*e^3+10*(6*A*a*b^5+15*B*a^2*b^4)*d^3*e^2+5*(A*b^6+6*B*a*b^5)*d^4*e+b^6*B*d^5)*x^8+1/7*((6*A*a^5*b+B*a^6)*e^5+5*(15*A*a^4*b^2+6*B*a^5*b)*d*e^4+10*(20*A*a^3*b^3+15*B*a^4*b^2)*d^2*e^3+10*(15*A*a^2*b^4+20*B*a^3*b^3)*d^3*e^2+5*(6*A*a*b^5+15*B*a^2*b^4)*d^4*e+(A*b^6+6*B*a*b^5)*d^5)*x^7+1/6*(a^6*A*e^5+5*(6*A*a^5*b+B*a^6)*d*e^4+10*(15*A*a^4*b^2+6*B*a^5*b)*d^2*e^3+10*(20*A*a^3*b^3+15*B*a^4*b^2)*d^3*e^2+5*(15*A*a^2*b^4+20*B*a^3*b^3)*d^4*e+(6*A*a*b^5+15*B*a^2*b^4)*d^5)*x^6+1/5*(5*a^6*A*d*e^4+10*(6*A*a^5*b+B*a^6)*d^2*e^3+10*(15*A*a^4*b^2+6*B*a^5*b)*d^3*e^2+5*(20*A*a^3*b^3+15*B*a^4*b^2)*d^4*e+(15*A*a^2*b^4+20*B*a^3*b^3)*d^5)*x^5+1/4*(10*a^6*A*d^2*e^3+10*(6*A*a^5*b+B*a^6)*d^3*e^2+5*(15*A*a^4*b^2+6*B*a^5*b)*d^4*e+(20*A*a^3*b^3+15*B*a^4*b^2)*d^5)*x^4+1/3*(10*a^6*A*d^3*e^2+5*(6*A*a^5*b+B*a^6)*d^4*e+(15*A*a^4*b^2+6*B*a^5*b)*d^5)*x^3+1/2*(5*a^6*A*d^4*e+(6*A*a^5*b+B*a^6)*d^5)*x^2+a^6*A*d^5*x
```


Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1278 vs. $2(241) = 482$.

Time = 0.08 (sec) , antiderivative size = 1278, normalized size of antiderivative = 5.32

$$\int (a + bx)^6 (A + Bx)(d + ex)^5 dx = \text{Too large to display}$$

input `integrate((b*x+a)**6*(B*x+A)*(e*x+d)**5,x)`

output

```
A*a**6*d**5*x + B*b**6*e**5*x**13/13 + x**12*(A*b**6*e**5/12 + B*a*b**5*e**5/2 + 5*B*b**6*d*e**4/12) + x**11*(6*A*a*b**5*e**5/11 + 5*A*b**6*d*e**4/11 + 15*B*a**2*b**4*e**5/11 + 30*B*a*b**5*d*e**4/11 + 10*B*b**6*d**2*e**3/11) + x**10*(3*A*a**2*b**4*e**5/2 + 3*A*a*b**5*d*e**4 + A*b**6*d**2*e**3 + 2*B*a**3*b**3*e**5 + 15*B*a**2*b**4*d*e**4/2 + 6*B*a*b**5*d**2*e**3 + B*b**6*d**3*e**2) + x**9*(20*A*a**3*b**3*e**5/9 + 25*A*a**2*b**4*d*e**4/3 + 20*A*a*b**5*d**2*e**3/3 + 10*A*b**6*d**3*e**2/9 + 5*B*a**4*b**2*e**5/3 + 100*B*a**3*b**3*d*e**4/9 + 50*B*a**2*b**4*d**2*e**3/3 + 20*B*a*b**5*d**3*e**2/3 + 5*B*b**6*d**4*e/9) + x**8*(15*A*a**4*b**2*e**5/8 + 25*A*a**3*b**3*d*e**4/2 + 75*A*a**2*b**4*d**2*e**3/4 + 15*A*a*b**5*d**3*e**2/2 + 5*A*b**6*d**4*e/8 + 3*B*a**5*b*e**5/4 + 75*B*a**4*b**2*d*e**4/8 + 25*B*a**3*b**3*d**2*e**3 + 75*B*a**2*b**4*d**3*e**2/4 + 15*B*a*b**5*d**4*e/4 + B*b**6*d**5/8) + x**7*(6*A*a**5*b*e**5/7 + 75*A*a**4*b**2*d*e**4/7 + 200*A*a**3*b**3*d**2*e**3/7 + 150*A*a**2*b**4*d**3*e**2/7 + 30*A*a*b**5*d**4*e/7 + A*b**6*d**5/7 + B*a**6*e**5/7 + 30*B*a**5*b*d*e**4/7 + 150*B*a**4*b**2*d**2*e**3/7 + 200*B*a**3*b**3*d**3*e**2/7 + 75*B*a**2*b**4*d**4*e/7 + 6*B*a*b**5*d**5/7) + x**6*(A*a**6*e**5/6 + 5*A*a**5*b*d*e**4 + 25*A*a**4*b**2*d**2*e**3 + 100*A*a**3*b**3*d**3*e**2/3 + 25*A*a**2*b**4*d**4*e/2 + A*a*b**5*d**5 + 5*B*a**6*d*e**4/6 + 10*B*a**5*b*d**2*e**3 + 25*B*a**4*b**2*d**3*e**2 + 50*B*a**3*b**3*d**4*e/3 + 5*B*a**2*b**4*d**5/2) + x**5*(A*a**6*d*e**4 + 12*A*...
```


Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1246 vs. $2(228) = 456$.

Time = 0.12 (sec) , antiderivative size = 1246, normalized size of antiderivative = 5.19

$$\int (a + bx)^6 (A + Bx)(d + ex)^5 dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)*(e*x+d)^5,x, algorithm="giac")`

output

```
1/13*B*b^6*e^5*x^13 + 5/12*B*b^6*d*e^4*x^12 + 1/2*B*a*b^5*e^5*x^12 + 1/12*
A*b^6*e^5*x^12 + 10/11*B*b^6*d^2*e^3*x^11 + 30/11*B*a*b^5*d*e^4*x^11 + 5/1
1*A*b^6*d*e^4*x^11 + 15/11*B*a^2*b^4*e^5*x^11 + 6/11*A*a*b^5*e^5*x^11 + B*
b^6*d^3*e^2*x^10 + 6*B*a*b^5*d^2*e^3*x^10 + A*b^6*d^2*e^3*x^10 + 15/2*B*a^
2*b^4*d*e^4*x^10 + 3*A*a*b^5*d*e^4*x^10 + 2*B*a^3*b^3*e^5*x^10 + 3/2*A*a^2
*b^4*e^5*x^10 + 5/9*B*b^6*d^4*e*x^9 + 20/3*B*a*b^5*d^3*e^2*x^9 + 10/9*A*b^
6*d^3*e^2*x^9 + 50/3*B*a^2*b^4*d^2*e^3*x^9 + 20/3*A*a*b^5*d^2*e^3*x^9 + 10
0/9*B*a^3*b^3*d*e^4*x^9 + 25/3*A*a^2*b^4*d*e^4*x^9 + 5/3*B*a^4*b^2*e^5*x^9
+ 20/9*A*a^3*b^3*e^5*x^9 + 1/8*B*b^6*d^5*x^8 + 15/4*B*a*b^5*d^4*e*x^8 + 5
/8*A*b^6*d^4*e*x^8 + 75/4*B*a^2*b^4*d^3*e^2*x^8 + 15/2*A*a*b^5*d^3*e^2*x^8
+ 25*B*a^3*b^3*d^2*e^3*x^8 + 75/4*A*a^2*b^4*d^2*e^3*x^8 + 75/8*B*a^4*b^2*
d*e^4*x^8 + 25/2*A*a^3*b^3*d*e^4*x^8 + 3/4*B*a^5*b*e^5*x^8 + 15/8*A*a^4*b^
2*e^5*x^8 + 6/7*B*a*b^5*d^5*x^7 + 1/7*A*b^6*d^5*x^7 + 75/7*B*a^2*b^4*d^4*
e*x^7 + 30/7*A*a*b^5*d^4*e*x^7 + 200/7*B*a^3*b^3*d^3*e^2*x^7 + 150/7*A*a^2*
b^4*d^3*e^2*x^7 + 150/7*B*a^4*b^2*d^2*e^3*x^7 + 200/7*A*a^3*b^3*d^2*e^3*x^
7 + 30/7*B*a^5*b*d*e^4*x^7 + 75/7*A*a^4*b^2*d*e^4*x^7 + 1/7*B*a^6*e^5*x^7
+ 6/7*A*a^5*b*e^5*x^7 + 5/2*B*a^2*b^4*d^5*x^6 + A*a*b^5*d^5*x^6 + 50/3*B*a
^3*b^3*d^4*e*x^6 + 25/2*A*a^2*b^4*d^4*e*x^6 + 25*B*a^4*b^2*d^3*e^2*x^6 + 1
00/3*A*a^3*b^3*d^3*e^2*x^6 + 10*B*a^5*b*d^2*e^3*x^6 + 25*A*a^4*b^2*d^2*e^3
*x^6 + 5/6*B*a^6*d*e^4*x^6 + 5*A*a^5*b*d*e^4*x^6 + 1/6*A*a^6*e^5*x^6 + ...
```

Mupad [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 1039, normalized size of antiderivative = 4.33

$$\int (a + bx)^6 (A + Bx)(d + ex)^5 dx = \text{Too large to display}$$

input `int((A + B*x)*(a + b*x)^6*(d + e*x)^5,x)`

output

$$\begin{aligned} & x^7 * ((A*b^6*d^5)/7 + (B*a^6*e^5)/7 + (6*A*a^5*b*e^5)/7 + (6*B*a*b^5*d^5)/7 \\ & + (75*A*a^4*b^2*d*e^4)/7 + (75*B*a^2*b^4*d^4*e)/7 + (150*A*a^2*b^4*d^3*e^2)/7 + (200*A*a^3*b^3*d^2*e^3)/7 + (200*B*a^3*b^3*d^3*e^2)/7 + (150*B*a^4*b^2*d^2*e^3)/7 + (30*A*a*b^5*d^4*e)/7 + (30*B*a^5*b*d*e^4)/7 + x^3*(2*B*a^5*b*d^5 + (5*B*a^6*d^4*e)/3 + 5*A*a^4*b^2*d^5 + (10*A*a^6*d^3*e^2)/3 + 10*A*a^5*b*d^4*e) + x^{11}*((6*A*a*b^5*e^5)/11 + (5*A*b^6*d*e^4)/11 + (15*B*a^2*b^4*e^5)/11 + (10*B*b^6*d^2*e^3)/11 + (30*B*a*b^5*d*e^4)/11) + x^6*((A*a^6*e^5)/6 + A*a*b^5*d^5 + (5*B*a^6*d*e^4)/6 + (5*B*a^2*b^4*d^5)/2 + (25*A*a^2*b^4*d^4*e)/2 + (50*B*a^3*b^3*d^4*e)/3 + 10*B*a^5*b*d^2*e^3 + (100*A*a^3*b^3*d^3*e^2)/3 + 25*A*a^4*b^2*d^2*e^3 + 25*B*a^4*b^2*d^3*e^2 + 5*A*a^5*b*d*e^4) + x^8*((B*b^6*d^5)/8 + (3*B*a^5*b*e^5)/4 + (5*A*b^6*d^4*e)/8 + (15*A*a^4*b^2*e^5)/8 + (15*A*a*b^5*d^3*e^2)/2 + (25*A*a^3*b^3*d*e^4)/2 + (75*B*a^4*b^2*d*e^4)/8 + (75*A*a^2*b^4*d^2*e^3)/4 + (75*B*a^2*b^4*d^3*e^2)/4 + 25*B*a^3*b^3*d^2*e^3 + (15*B*a*b^5*d^4*e)/4) + x^5*(A*a^6*d*e^4 + 3*A*a^2*b^4*d^5 + 4*B*a^3*b^3*d^5 + 2*B*a^6*d^2*e^3 + 20*A*a^3*b^3*d^4*e + 12*A*a^5*b*d^2*e^3 + 15*B*a^4*b^2*d^4*e + 12*B*a^5*b*d^3*e^2 + 30*A*a^4*b^2*d^3*e^2) + x^9*((5*B*b^6*d^4*e)/9 + (20*A*a^3*b^3*e^5)/9 + (5*B*a^4*b^2*e^5)/3 + (10*A*b^6*d^3*e^2)/9 + (20*A*a*b^5*d^2*e^3)/3 + (25*A*a^2*b^4*d*e^4)/3 + (20*B*a*b^5*d^3*e^2)/3 + (100*B*a^3*b^3*d*e^4)/9 + (50*B*a^2*b^4*d^2*e^3)/3) + x^4*(5*A*a^3*b^3*d^5 + (15*B*a^4*b^2*d^5)/4 + (5*A*a^6*d^2*e^3)/... \end{aligned}$$
Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 673, normalized size of antiderivative = 2.80

$$\begin{aligned} & \int (a + bx)^6 (A + Bx)(d + ex)^5 dx \\ & = \frac{x(792b^7e^5x^{12} + 6006ab^6e^5x^{11} + 4290b^7de^4x^{11} + 19656a^2b^5e^5x^{10} + 32760ab^6de^4x^{10} + 9360b^7d^2e^3x^{10} + \dots}{\dots} \end{aligned}$$

input `int((b*x+a)^6*(B*x+A)*(e*x+d)^5,x)`

output `(x*(10296*a**7*d**5 + 25740*a**7*d**4*e*x + 34320*a**7*d**3*e**2*x**2 + 25740*a**7*d**2*e**3*x**3 + 10296*a**7*d*e**4*x**4 + 1716*a**7*e**5*x**5 + 36036*a**6*b*d**5*x + 120120*a**6*b*d**4*e*x**2 + 180180*a**6*b*d**3*e**2*x**3 + 144144*a**6*b*d**2*e**3*x**4 + 60060*a**6*b*d*e**4*x**5 + 10296*a**6*b*e**5*x**6 + 72072*a**5*b**2*d**5*x**2 + 270270*a**5*b**2*d**4*e*x**3 + 432432*a**5*b**2*d**3*e**2*x**4 + 360360*a**5*b**2*d**2*e**3*x**5 + 154440*a**5*b**2*d*e**4*x**6 + 27027*a**5*b**2*e**5*x**7 + 90090*a**4*b**3*d**5*x**3 + 360360*a**4*b**3*d**4*e*x**4 + 600600*a**4*b**3*d**3*e**2*x**5 + 514800*a**4*b**3*d**2*e**3*x**6 + 225225*a**4*b**3*d*e**4*x**7 + 40040*a**4*b**3*e**5*x**8 + 72072*a**3*b**4*d**5*x**4 + 300300*a**3*b**4*d**4*e*x**5 + 514800*a**3*b**4*d**3*e**2*x**6 + 450450*a**3*b**4*d**2*e**3*x**7 + 200200*a**3*b**4*d*e**4*x**8 + 36036*a**3*b**4*e**5*x**9 + 36036*a**2*b**5*d**5*x**5 + 154440*a**2*b**5*d**4*e*x**6 + 270270*a**2*b**5*d**3*e**2*x**7 + 240240*a**2*b**5*d**2*e**3*x**8 + 108108*a**2*b**5*d*e**4*x**9 + 19656*a**2*b**5*e**5*x**10 + 10296*a*b**6*d**5*x**6 + 45045*a*b**6*d**4*e*x**7 + 80080*a*b**6*d**3*e**2*x**8 + 72072*a*b**6*d**2*e**3*x**9 + 32760*a*b**6*d*e**4*x**10 + 6006*a*b**6*e**5*x**11 + 1287*b**7*d**5*x**7 + 5720*b**7*d**4*e*x**8 + 10296*b**7*d**3*e**2*x**9 + 9360*b**7*d**2*e**3*x**10 + 4290*b**7*d*e**4*x**11 + 792*b**7*e**5*x**12))/10296`

3.45 $\int (a + bx)^6 (A + Bx)(d + ex)^4 dx$

Optimal result	451
Mathematica [B] (verified)	452
Rubi [A] (verified)	453
Maple [B] (verified)	455
Fricas [B] (verification not implemented)	456
Sympy [B] (verification not implemented)	456
Maxima [B] (verification not implemented)	457
Giac [B] (verification not implemented)	458
Mupad [B] (verification not implemented)	460
Reduce [B] (verification not implemented)	461

Optimal result

Integrand size = 20, antiderivative size = 204

$$\int (a + bx)^6 (A + Bx)(d + ex)^4 dx = \frac{(Ab - aB)(bd - ae)^4(a + bx)^7}{7b^6} + \frac{(bd - ae)^3(bBd + 4Abe - 5aBe)(a + bx)^8}{8b^6} + \frac{2e(bd - ae)^2(2bBd + 3Abe - 5aBe)(a + bx)^9}{9b^6} + \frac{e^2(bd - ae)(3bBd + 2Abe - 5aBe)(a + bx)^{10}}{5b^6} + \frac{e^3(4bBd + Abe - 5aBe)(a + bx)^{11}}{11b^6} + \frac{Be^4(a + bx)^{12}}{12b^6}$$

output

```
1/7*(A*b-B*a)*(-a*e+b*d)^4*(b*x+a)^7/b^6+1/8*(-a*e+b*d)^3*(4*A*b*e-5*B*a*e
+B*b*d)*(b*x+a)^8/b^6+2/9*e*(-a*e+b*d)^2*(3*A*b*e-5*B*a*e+2*B*b*d)*(b*x+a)
^9/b^6+1/5*e^2*(-a*e+b*d)*(2*A*b*e-5*B*a*e+3*B*b*d)*(b*x+a)^10/b^6+1/11*e^
3*(A*b*e-5*B*a*e+4*B*b*d)*(b*x+a)^11/b^6+1/12*B*e^4*(b*x+a)^12/b^6
```


Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 762 vs. $2(204) = 408$.

Time = 0.16 (sec) , antiderivative size = 762, normalized size of antiderivative = 3.74

$$\begin{aligned}
 \int (a + bx)^6 (A + Bx)(d + ex)^4 dx = & a^6 Ad^4 x + \frac{1}{2} a^5 d^3 (6Abd + aBd + 4aAe)x^2 \\
 & + \frac{1}{3} a^4 d^2 (2aBd(3bd + 2ae) \\
 & \quad + 3A(5b^2 d^2 + 8abde + 2a^2 e^2)) x^3 \\
 & + \frac{1}{4} a^3 d (3aBd(5b^2 d^2 + 8abde + 2a^2 e^2) \\
 & \quad + 4A(5b^3 d^3 + 15ab^2 d^2 e + 9a^2 bde^2 + a^3 e^3)) x^4 \\
 & + \frac{1}{5} a^2 (4aBd(5b^3 d^3 + 15ab^2 d^2 e + 9a^2 bde^2 + a^3 e^3) \\
 & \quad + A(15b^4 d^4 + 80ab^3 d^3 e + 90a^2 b^2 d^2 e^2 + 24a^3 bde^3 \\
 & \quad \quad + a^4 e^4)) x^5 + \frac{1}{6} a (6Ab(b^4 d^4 + 10ab^3 d^3 e \\
 & \quad + 20a^2 b^2 d^2 e^2 + 10a^3 bde^3 + a^4 e^4) + aB(15b^4 d^4 \\
 & \quad + 80ab^3 d^3 e + 90a^2 b^2 d^2 e^2 + 24a^3 bde^3 + a^4 e^4)) x^6 \\
 & + \frac{1}{7} b (6aB(b^4 d^4 + 10ab^3 d^3 e + 20a^2 b^2 d^2 e^2 + 10a^3 bde^3 \\
 & \quad + a^4 e^4) + Ab(b^4 d^4 + 24ab^3 d^3 e + 90a^2 b^2 d^2 e^2 \\
 & \quad \quad + 80a^3 bde^3 + 15a^4 e^4)) x^7 + \frac{1}{8} b^2 (15a^4 Be^4 \\
 & \quad + 20a^3 be^3(4Bd + Ae) + 30a^2 b^2 de^2(3Bd + 2Ae) \\
 & \quad + 12ab^3 d^2 e(2Bd + 3Ae) + b^4 d^3(Bd + 4Ae)) x^8 \\
 & + \frac{1}{9} b^3 e (20a^3 Be^3 + 15a^2 be^2(4Bd + Ae) \\
 & \quad + 12ab^2 de(3Bd + 2Ae) + 2b^3 d^2(2Bd + 3Ae)) x^9 \\
 & + \frac{1}{10} b^4 e^2 (15a^2 Be^2 + 6abe(4Bd + Ae) \\
 & \quad \quad + 2b^2 d(3Bd + 2Ae)) x^{10} \\
 & + \frac{1}{11} b^5 e^3 (4bBd + Abe + 6aBe)x^{11} + \frac{1}{12} b^6 Be^4 x^{12}
 \end{aligned}$$

input

```
Integrate[(a + b*x)^6*(A + B*x)*(d + e*x)^4,x]
```

output

```

a^6*A*d^4*x + (a^5*d^3*(6*A*b*d + a*B*d + 4*a*A*e)*x^2)/2 + (a^4*d^2*(2*a*
B*d*(3*b*d + 2*a*e) + 3*A*(5*b^2*d^2 + 8*a*b*d*e + 2*a^2*e^2))*x^3)/3 + (a
^3*d*(3*a*B*d*(5*b^2*d^2 + 8*a*b*d*e + 2*a^2*e^2) + 4*A*(5*b^3*d^3 + 15*a*
b^2*d^2*e + 9*a^2*b*d*e^2 + a^3*e^3))*x^4)/4 + (a^2*(4*a*B*d*(5*b^3*d^3 +
15*a*b^2*d^2*e + 9*a^2*b*d*e^2 + a^3*e^3) + A*(15*b^4*d^4 + 80*a*b^3*d^3*e
+ 90*a^2*b^2*d^2*e^2 + 24*a^3*b*d*e^3 + a^4*e^4))*x^5)/5 + (a*(6*A*b*(b^4
*d^4 + 10*a*b^3*d^3*e + 20*a^2*b^2*d^2*e^2 + 10*a^3*b*d*e^3 + a^4*e^4) + a
*B*(15*b^4*d^4 + 80*a*b^3*d^3*e + 90*a^2*b^2*d^2*e^2 + 24*a^3*b*d*e^3 + a^
4*e^4))*x^6)/6 + (b*(6*a*B*(b^4*d^4 + 10*a*b^3*d^3*e + 20*a^2*b^2*d^2*e^2
+ 10*a^3*b*d*e^3 + a^4*e^4) + A*b*(b^4*d^4 + 24*a*b^3*d^3*e + 90*a^2*b^2*d
^2*e^2 + 80*a^3*b*d*e^3 + 15*a^4*e^4))*x^7)/7 + (b^2*(15*a^4*B*e^4 + 20*a^
3*b*e^3*(4*B*d + A*e) + 30*a^2*b^2*d*e^2*(3*B*d + 2*A*e) + 12*a*b^3*d^2*e*
(2*B*d + 3*A*e) + b^4*d^3*(B*d + 4*A*e))*x^8)/8 + (b^3*e*(20*a^3*B*e^3 + 1
5*a^2*b*e^2*(4*B*d + A*e) + 12*a*b^2*d*e*(3*B*d + 2*A*e) + 2*b^3*d^2*(2*B*
d + 3*A*e))*x^9)/9 + (b^4*e^2*(15*a^2*B*e^2 + 6*a*b*e*(4*B*d + A*e) + 2*b^
2*d*(3*B*d + 2*A*e))*x^10)/10 + (b^5*e^3*(4*b*B*d + A*b*e + 6*a*B*e)*x^11)
/11 + (b^6*B*e^4*x^12)/12

```

Rubi [A] (verified)

Time = 0.89 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^6 (A + Bx)(d + ex)^4 dx$$

$$\downarrow 86$$

$$\int \left(\frac{e^3 (a + bx)^{10} (-5aBe + Abe + 4bBd)}{b^5} + \frac{2e^2 (a + bx)^9 (bd - ae) (-5aBe + 2Abe + 3bBd)}{b^5} + \frac{2e (a + bx)^8 (bd - ae)^2}{b^5} \right) dx$$

$$\downarrow 2009$$

$$\frac{e^3(a+bx)^{11}(-5aBe + Abe + 4bBd)}{11b^6} + \frac{e^2(a+bx)^{10}(bd-ae)(-5aBe + 2Abe + 3bBd)}{5b^6} + \frac{2e(a+bx)^9(bd-ae)^2(-5aBe + 3Abe + 2bBd)}{9b^6} + \frac{(a+bx)^8(bd-ae)^3(-5aBe + 4Abe + bBd)}{8b^6} + \frac{(a+bx)^7(Ab-aB)(bd-ae)^4}{7b^6} + \frac{Be^4(a+bx)^{12}}{12b^6}$$

input `Int[(a + b*x)^6*(A + B*x)*(d + e*x)^4,x]`

output `((A*b - a*B)*(b*d - a*e)^4*(a + b*x)^7)/(7*b^6) + ((b*d - a*e)^3*(b*B*d + 4*A*b*e - 5*a*B*e)*(a + b*x)^8)/(8*b^6) + (2*e*(b*d - a*e)^2*(2*b*B*d + 3*A*b*e - 5*a*B*e)*(a + b*x)^9)/(9*b^6) + (e^2*(b*d - a*e)*(3*b*B*d + 2*A*b*e - 5*a*B*e)*(a + b*x)^10)/(5*b^6) + (e^3*(4*b*B*d + A*b*e - 5*a*B*e)*(a + b*x)^11)/(11*b^6) + (B*e^4*(a + b*x)^12)/(12*b^6)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 820 vs. 2(192) = 384.

Time = 0.20 (sec) , antiderivative size = 821, normalized size of antiderivative = 4.02

method	result
default	$\frac{b^6 B e^4 x^{12}}{12} + \frac{((b^6 A + 6a b^5 B) e^4 + 4b^6 B d e^3) x^{11}}{11} + \frac{((6a b^5 A + 15a^2 b^4 B) e^4 + 4(b^6 A + 6a b^5 B) d e^3 + 6b^6 B d^2 e^2) x^{10}}{10} + \frac{((15 a^5 A + 10 a^4 B) e^4 + 4(a^6 A + 6a^5 B) d e^3 + 6a^6 B d^2 e^2) x^9}{9} + \frac{((15 a^4 A + 10 a^3 B) e^4 + 4(a^6 A + 6a^5 B) d e^3 + 6a^6 B d^2 e^2) x^8}{8} + \frac{((15 a^3 A + 10 a^2 B) e^4 + 4(a^6 A + 6a^5 B) d e^3 + 6a^6 B d^2 e^2) x^7}{7} + \frac{((15 a^2 A + 10 a B) e^4 + 4(a^6 A + 6a^5 B) d e^3 + 6a^6 B d^2 e^2) x^6}{6} + \frac{((15 a A + 10 B) e^4 + 4(a^6 A + 6a^5 B) d e^3 + 6a^6 B d^2 e^2) x^5}{5} + \frac{((15 A + 10 B) e^4 + 4(a^6 A + 6a^5 B) d e^3 + 6a^6 B d^2 e^2) x^4}{4} + \frac{((15 A + 10 B) e^4 + 4(a^6 A + 6a^5 B) d e^3 + 6a^6 B d^2 e^2) x^3}{3} + \frac{((15 A + 10 B) e^4 + 4(a^6 A + 6a^5 B) d e^3 + 6a^6 B d^2 e^2) x^2}{2} + \frac{((15 A + 10 B) e^4 + 4(a^6 A + 6a^5 B) d e^3 + 6a^6 B d^2 e^2) x}{1} + \frac{((15 A + 10 B) e^4 + 4(a^6 A + 6a^5 B) d e^3 + 6a^6 B d^2 e^2)}{0}$
norman	$\frac{b^6 B e^4 x^{12}}{12} + \left(\frac{1}{11} A b^6 e^4 + \frac{6}{11} B a b^5 e^4 + \frac{4}{11} b^6 B d e^3\right) x^{11} + \left(\frac{3}{5} A a b^5 e^4 + \frac{2}{5} A b^6 d e^3 + \frac{3}{2} B a^2 b^4 e^4 + \dots\right) x^{10} + \dots$
gospers	Expression too large to display
risch	Expression too large to display
parallexrisch	Expression too large to display
orering	Expression too large to display

input `int((b*x+a)^6*(B*x+A)*(e*x+d)^4,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{12} b^6 B e^4 x^{12} + \frac{1}{11} ((A b^6 + 6 B a b^5) e^4 + 4 b^6 B d e^3) x^{11} + \frac{1}{10} ((6 A a b^5 + 15 B a^2 b^4) e^4 + 4 (A b^6 + 6 B a b^5) d e^3 + 6 b^6 B d^2 e^2) x^{10} + \frac{1}{9} ((15 A a^2 b^4 + 20 B a^3 b^3) e^4 + 4 (6 A a b^5 + 15 B a^2 b^4) d e^3 + 6 (A b^6 + 6 B a b^5) d^2 e^2 + 4 b^6 B d^3 e) x^9 + \frac{1}{8} ((20 A a^3 b^3 + 15 B a^4 b^2) e^4 + 4 (15 A a^2 b^4 + 20 B a^3 b^3) d e^3 + 6 (6 A a b^5 + 15 B a^2 b^4) d^2 e^2 + 4 (A b^6 + 6 B a b^5) d^3 e + b^6 B d^4) x^8 + \frac{1}{7} ((15 A a^4 b^2 + 6 B a^5 b) e^4 + 4 (20 A a^3 b^3 + 15 B a^4 b^2) d e^3 + 6 (15 A a^2 b^4 + 20 B a^3 b^3) d^2 e^2 + 4 (6 A a b^5 + 15 B a^2 b^4) d^3 e + (A b^6 + 6 B a b^5) d^4) x^7 + \frac{1}{6} ((6 A a^5 b + B a^6) e^4 + 4 (15 A a^4 b^2 + 6 B a^5 b) d e^3 + 6 (20 A a^3 b^3 + 15 B a^4 b^2) d^2 e^2 + 4 (15 A a^2 b^4 + 20 B a^3 b^3) d^3 e + (6 A a b^5 + 15 B a^2 b^4) d^4) x^6 + \frac{1}{5} (a^6 A e^4 + 4 (6 A a^5 b + B a^6) d e^3 + 6 (15 A a^4 b^2 + 6 B a^5 b) d^2 e^2 + 4 (20 A a^3 b^3 + 15 B a^4 b^2) d^3 e + (15 A a^2 b^4 + 20 B a^3 b^3) d^4) x^5 + \frac{1}{4} (4 a^6 A d e^3 + 6 (6 A a^5 b + B a^6) d^2 e^2 + 4 (15 A a^4 b^2 + 6 B a^5 b) d^3 e + (20 A a^3 b^3 + 15 B a^4 b^2) d^4) x^4 + \frac{1}{3} (6 a^6 A d^2 e^2 + 4 (6 A a^5 b + B a^6) d^3 e + (15 A a^4 b^2 + 6 B a^5 b) d^4) x^3 + \frac{1}{2} (4 a^6 A d^3 e + (6 A a^5 b + B a^6) d^4) x^2 + a^6 A d^4 x$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 828 vs. $2(192) = 384$.

Time = 0.09 (sec) , antiderivative size = 828, normalized size of antiderivative = 4.06

$$\int (a + bx)^6 (A + Bx)(d + ex)^4 dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)*(e*x+d)^4,x, algorithm="fricas")`

output

$$\begin{aligned} & 1/12*B*b^6*e^4*x^{12} + A*a^6*d^4*x + 1/11*(4*B*b^6*d*e^3 + (6*B*a*b^5 + A*b^6)*e^4)*x^{11} + 1/10*(6*B*b^6*d^2*e^2 + 4*(6*B*a*b^5 + A*b^6)*d*e^3 + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*e^4)*x^{10} + 1/9*(4*B*b^6*d^3*e + 6*(6*B*a*b^5 + A*b^6)*d^2*e^2 + 12*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^3 + 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^4)*x^9 + 1/8*(B*b^6*d^4 + 4*(6*B*a*b^5 + A*b^6)*d^3*e + 18*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^2 + 20*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^3 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^4)*x^8 + 1/7*((6*B*a*b^5 + A*b^6)*d^4 + 12*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*e + 30*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^2 + 20*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^3 + 3*(2*B*a^5*b + 5*A*a^4*b^2)*e^4)*x^7 + 1/6*(3*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4 + 20*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e + 30*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^2 + 12*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^3 + (B*a^6 + 6*A*a^5*b)*e^4)*x^6 + 1/5*(A*a^6*e^4 + 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4 + 20*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e + 18*(2*B*a^5*b + 5*A*a^4*b^2)*d^2*e^2 + 4*(B*a^6 + 6*A*a^5*b)*d*e^3)*x^5 + 1/4*(4*A*a^6*d*e^3 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^4 + 12*(2*B*a^5*b + 5*A*a^4*b^2)*d^3*e + 6*(B*a^6 + 6*A*a^5*b)*d^2*e^2)*x^4 + 1/3*(6*A*a^6*d^2*e^2 + 3*(2*B*a^5*b + 5*A*a^4*b^2)*d^4 + 4*(B*a^6 + 6*A*a^5*b)*d^3*e)*x^3 + 1/2*(4*A*a^6*d^3*e + (B*a^6 + 6*A*a^5*b)*d^4)*x^2 \end{aligned}$$
Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1035 vs. $2(204) = 408$.

Time = 0.07 (sec) , antiderivative size = 1035, normalized size of antiderivative = 5.07

$$\int (a + bx)^6 (A + Bx)(d + ex)^4 dx = \text{Too large to display}$$

input `integrate((b*x+a)**6*(B*x+A)*(e*x+d)**4,x)`

output

```

A*a**6*d**4*x + B*b**6*e**4*x**12/12 + x**11*(A*b**6*e**4/11 + 6*B*a*b**5*
e**4/11 + 4*B*b**6*d*e**3/11) + x**10*(3*A*a*b**5*e**4/5 + 2*A*b**6*d*e**3
/5 + 3*B*a**2*b**4*e**4/2 + 12*B*a*b**5*d*e**3/5 + 3*B*b**6*d**2*e**2/5) +
x**9*(5*A*a**2*b**4*e**4/3 + 8*A*a*b**5*d*e**3/3 + 2*A*b**6*d**2*e**2/3 +
20*B*a**3*b**3*e**4/9 + 20*B*a**2*b**4*d*e**3/3 + 4*B*a*b**5*d**2*e**2 +
4*B*b**6*d**3*e/9) + x**8*(5*A*a**3*b**3*e**4/2 + 15*A*a**2*b**4*d*e**3/2
+ 9*A*a*b**5*d**2*e**2/2 + A*b**6*d**3*e/2 + 15*B*a**4*b**2*e**4/8 + 10*B*
a**3*b**3*d*e**3 + 45*B*a**2*b**4*d**2*e**2/4 + 3*B*a*b**5*d**3*e + B*b**6
*d**4/8) + x**7*(15*A*a**4*b**2*e**4/7 + 80*A*a**3*b**3*d*e**3/7 + 90*A*a*
**2*b**4*d**2*e**2/7 + 24*A*a*b**5*d**3*e/7 + A*b**6*d**4/7 + 6*B*a**5*b*e*
**4/7 + 60*B*a**4*b**2*d*e**3/7 + 120*B*a**3*b**3*d**2*e**2/7 + 60*B*a**2*b
**4*d**3*e/7 + 6*B*a*b**5*d**4/7) + x**6*(A*a**5*b*e**4 + 10*A*a**4*b**2*d
*e**3 + 20*A*a**3*b**3*d**2*e**2 + 10*A*a**2*b**4*d**3*e + A*a*b**5*d**4 +
B*a**6*e**4/6 + 4*B*a**5*b*d*e**3 + 15*B*a**4*b**2*d**2*e**2 + 40*B*a**3*b
**3*d**3*e/3 + 5*B*a**2*b**4*d**4/2) + x**5*(A*a**6*e**4/5 + 24*A*a**5*b*
d*e**3/5 + 18*A*a**4*b**2*d**2*e**2 + 16*A*a**3*b**3*d**3*e + 3*A*a**2*b**
4*d**4 + 4*B*a**6*d*e**3/5 + 36*B*a**5*b*d**2*e**2/5 + 12*B*a**4*b**2*d**3
*e + 4*B*a**3*b**3*d**4) + x**4*(A*a**6*d*e**3 + 9*A*a**5*b*d**2*e**2 + 15
*A*a**4*b**2*d**3*e + 5*A*a**3*b**3*d**4 + 3*B*a**6*d**2*e**2/2 + 6*B*a**5
*b*d**3*e + 15*B*a**4*b**2*d**4/4) + x**3*(2*A*a**6*d**2*e**2 + 8*A*a**...

```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 828 vs. $2(192) = 384$.

Time = 0.04 (sec) , antiderivative size = 828, normalized size of antiderivative = 4.06

$$\int (a + bx)^6 (A + Bx)(d + ex)^4 dx = \text{Too large to display}$$

input

```

integrate((b*x+a)^6*(B*x+A)*(e*x+d)^4,x, algorithm="maxima")

```

output

```

1/12*B*b^6*e^4*x^12 + A*a^6*d^4*x + 1/11*(4*B*b^6*d*e^3 + (6*B*a*b^5 + A*b
^6)*e^4)*x^11 + 1/10*(6*B*b^6*d^2*e^2 + 4*(6*B*a*b^5 + A*b^6)*d*e^3 + 3*(5
*B*a^2*b^4 + 2*A*a*b^5)*e^4)*x^10 + 1/9*(4*B*b^6*d^3*e + 6*(6*B*a*b^5 + A
b^6)*d^2*e^2 + 12*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^3 + 5*(4*B*a^3*b^3 + 3*A*a
^2*b^4)*e^4)*x^9 + 1/8*(B*b^6*d^4 + 4*(6*B*a*b^5 + A*b^6)*d^3*e + 18*(5*B*
a^2*b^4 + 2*A*a*b^5)*d^2*e^2 + 20*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^3 + 5*(3
*B*a^4*b^2 + 4*A*a^3*b^3)*e^4)*x^8 + 1/7*((6*B*a*b^5 + A*b^6)*d^4 + 12*(5*
B*a^2*b^4 + 2*A*a*b^5)*d^3*e + 30*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^2 + 20
*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^3 + 3*(2*B*a^5*b + 5*A*a^4*b^2)*e^4)*x^7
+ 1/6*(3*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4 + 20*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^
3*e + 30*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^2 + 12*(2*B*a^5*b + 5*A*a^4*b^2
)*d*e^3 + (B*a^6 + 6*A*a^5*b)*e^4)*x^6 + 1/5*(A*a^6*e^4 + 5*(4*B*a^3*b^3 +
3*A*a^2*b^4)*d^4 + 20*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e + 18*(2*B*a^5*b +
5*A*a^4*b^2)*d^2*e^2 + 4*(B*a^6 + 6*A*a^5*b)*d*e^3)*x^5 + 1/4*(4*A*a^6*d*
e^3 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^4 + 12*(2*B*a^5*b + 5*A*a^4*b^2)*d^3
*e + 6*(B*a^6 + 6*A*a^5*b)*d^2*e^2)*x^4 + 1/3*(6*A*a^6*d^2*e^2 + 3*(2*B*a^
5*b + 5*A*a^4*b^2)*d^4 + 4*(B*a^6 + 6*A*a^5*b)*d^3*e)*x^3 + 1/2*(4*A*a^6*d
^3*e + (B*a^6 + 6*A*a^5*b)*d^4)*x^2

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1015 vs. $2(192) = 384$.

Time = 0.13 (sec) , antiderivative size = 1015, normalized size of antiderivative = 4.98

$$\int (a + bx)^6 (A + Bx)(d + ex)^4 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^6*(B*x+A)*(e*x+d)^4,x, algorithm="giac")
```

output

```

1/12*B*b^6*e^4*x^12 + 4/11*B*b^6*d*e^3*x^11 + 6/11*B*a*b^5*e^4*x^11 + 1/11
*A*b^6*e^4*x^11 + 3/5*B*b^6*d^2*e^2*x^10 + 12/5*B*a*b^5*d*e^3*x^10 + 2/5*A
*b^6*d*e^3*x^10 + 3/2*B*a^2*b^4*e^4*x^10 + 3/5*A*a*b^5*e^4*x^10 + 4/9*B*b^
6*d^3*e*x^9 + 4*B*a*b^5*d^2*e^2*x^9 + 2/3*A*b^6*d^2*e^2*x^9 + 20/3*B*a^2*b
^4*d*e^3*x^9 + 8/3*A*a*b^5*d*e^3*x^9 + 20/9*B*a^3*b^3*e^4*x^9 + 5/3*A*a^2*
b^4*e^4*x^9 + 1/8*B*b^6*d^4*x^8 + 3*B*a*b^5*d^3*e*x^8 + 1/2*A*b^6*d^3*e*x^
8 + 45/4*B*a^2*b^4*d^2*e^2*x^8 + 9/2*A*a*b^5*d^2*e^2*x^8 + 10*B*a^3*b^3*d*
e^3*x^8 + 15/2*A*a^2*b^4*d*e^3*x^8 + 15/8*B*a^4*b^2*e^4*x^8 + 5/2*A*a^3*b^
3*e^4*x^8 + 6/7*B*a*b^5*d^4*x^7 + 1/7*A*b^6*d^4*x^7 + 60/7*B*a^2*b^4*d^3*e
*x^7 + 24/7*A*a*b^5*d^3*e*x^7 + 120/7*B*a^3*b^3*d^2*e^2*x^7 + 90/7*A*a^2*b
^4*d^2*e^2*x^7 + 60/7*B*a^4*b^2*d*e^3*x^7 + 80/7*A*a^3*b^3*d*e^3*x^7 + 6/7
*B*a^5*b*e^4*x^7 + 15/7*A*a^4*b^2*e^4*x^7 + 5/2*B*a^2*b^4*d^4*x^6 + A*a*b^
5*d^4*x^6 + 40/3*B*a^3*b^3*d^3*e*x^6 + 10*A*a^2*b^4*d^3*e*x^6 + 15*B*a^4*b
^2*d^2*e^2*x^6 + 20*A*a^3*b^3*d^2*e^2*x^6 + 4*B*a^5*b*d*e^3*x^6 + 10*A*a^4
*b^2*d*e^3*x^6 + 1/6*B*a^6*e^4*x^6 + A*a^5*b*e^4*x^6 + 4*B*a^3*b^3*d^4*x^5
+ 3*A*a^2*b^4*d^4*x^5 + 12*B*a^4*b^2*d^3*e*x^5 + 16*A*a^3*b^3*d^3*e*x^5 +
36/5*B*a^5*b*d^2*e^2*x^5 + 18*A*a^4*b^2*d^2*e^2*x^5 + 4/5*B*a^6*d*e^3*x^5
+ 24/5*A*a^5*b*d*e^3*x^5 + 1/5*A*a^6*e^4*x^5 + 15/4*B*a^4*b^2*d^4*x^4 + 5
*A*a^3*b^3*d^4*x^4 + 6*B*a^5*b*d^3*e*x^4 + 15*A*a^4*b^2*d^3*e*x^4 + 3/2*B*
a^6*d^2*e^2*x^4 + 9*A*a^5*b*d^2*e^2*x^4 + A*a^6*d*e^3*x^4 + 2*B*a^5*b*d...

```


Mupad [B] (verification not implemented)

Time = 1.01 (sec) , antiderivative size = 845, normalized size of antiderivative = 4.14

$$\begin{aligned}
\int (a + bx)^6 (A + Bx)(d + ex)^4 dx = & x^4 \left(\frac{3 B a^6 d^2 e^2}{2} + A a^6 d e^3 + 6 B a^5 b d^3 e \right. \\
& + 9 A a^5 b d^2 e^2 + \frac{15 B a^4 b^2 d^4}{4} + 15 A a^4 b^2 d^3 e \\
& \left. + 5 A a^3 b^3 d^4 \right) \\
& + x^9 \left(\frac{20 B a^3 b^3 e^4}{9} + \frac{20 B a^2 b^4 d e^3}{3} + \frac{5 A a^2 b^4 e^4}{3} \right. \\
& + 4 B a b^5 d^2 e^2 + \frac{8 A a b^5 d e^3}{3} + \frac{4 B b^6 d^3 e}{9} \\
& \left. + \frac{2 A b^6 d^2 e^2}{3} \right) + x^3 \left(\frac{4 B a^6 d^3 e}{3} + 2 A a^6 d^2 e^2 \right. \\
& \left. + 2 B a^5 b d^4 + 8 A a^5 b d^3 e + 5 A a^4 b^2 d^4 \right) \\
& + x^{10} \left(\frac{3 B a^2 b^4 e^4}{2} + \frac{12 B a b^5 d e^3}{5} + \frac{3 A a b^5 e^4}{5} \right. \\
& \left. + \frac{3 B b^6 d^2 e^2}{5} + \frac{2 A b^6 d e^3}{5} \right) \\
& + x^5 \left(\frac{4 B a^6 d e^3}{5} + \frac{A a^6 e^4}{5} + \frac{36 B a^5 b d^2 e^2}{5} \right. \\
& + \frac{24 A a^5 b d e^3}{5} + 12 B a^4 b^2 d^3 e + 18 A a^4 b^2 d^2 e^2 \\
& \left. + 4 B a^3 b^3 d^4 + 16 A a^3 b^3 d^3 e + 3 A a^2 b^4 d^4 \right) \\
& + x^8 \left(\frac{15 B a^4 b^2 e^4}{8} + 10 B a^3 b^3 d e^3 + \frac{5 A a^3 b^3 e^4}{2} \right. \\
& + \frac{45 B a^2 b^4 d^2 e^2}{4} + \frac{15 A a^2 b^4 d e^3}{2} + 3 B a b^5 d^3 e \\
& \left. + \frac{9 A a b^5 d^2 e^2}{2} + \frac{B b^6 d^4}{8} + \frac{A b^6 d^3 e}{2} \right) + x^6 \left(\frac{B a^6 e^4}{6} \right. \\
& + 4 B a^5 b d e^3 + A a^5 b e^4 + 15 B a^4 b^2 d^2 e^2 \\
& + 10 A a^4 b^2 d e^3 + \frac{40 B a^3 b^3 d^3 e}{3} + 20 A a^3 b^3 d^2 e^2 \\
& \left. + \frac{5 B a^2 b^4 d^4}{2} + 10 A a^2 b^4 d^3 e + A a b^5 d^4 \right) \\
& + x^7 \left(\frac{6 B a^5 b e^4}{7} + \frac{60 B a^4 b^2 d e^3}{7} + \frac{15 A a^4 b^2 e^4}{7} \right. \\
& + \frac{120 B a^3 b^3 d^2 e^2}{7} + \frac{80 A a^3 b^3 d e^3}{7} + \frac{60 B a^2 b^4 d^3 e}{7} \\
& + \frac{90 A a^2 b^4 d^2 e^2}{7} + \frac{6 B a b^5 d^4}{7} + \frac{24 A a b^5 d^3 e}{7} \\
& \left. + \frac{A b^6 d^4}{7} \right) + \frac{a^5 d^3 x^2 (4 A a e + 6 A b d + B a d)}{2}
\end{aligned}$$

input `int((A + B*x)*(a + b*x)^6*(d + e*x)^4,x)`

output

$$\begin{aligned} & x^4*(A*a^6*d*e^3 + 5*A*a^3*b^3*d^4 + (15*B*a^4*b^2*d^4)/4 + (3*B*a^6*d^2*e \\ & ^2)/2 + 15*A*a^4*b^2*d^3*e + 9*A*a^5*b*d^2*e^2 + 6*B*a^5*b*d^3*e) + x^9*((\\ & 4*B*b^6*d^3*e)/9 + (5*A*a^2*b^4*e^4)/3 + (20*B*a^3*b^3*e^4)/9 + (2*A*b^6*d \\ & ^2*e^2)/3 + 4*B*a*b^5*d^2*e^2 + (20*B*a^2*b^4*d*e^3)/3 + (8*A*a*b^5*d*e^3) \\ & /3) + x^3*(2*B*a^5*b*d^4 + (4*B*a^6*d^3*e)/3 + 5*A*a^4*b^2*d^4 + 2*A*a^6*d \\ & ^2*e^2 + 8*A*a^5*b*d^3*e) + x^10*((3*A*a*b^5*e^4)/5 + (2*A*b^6*d*e^3)/5 + \\ & (3*B*a^2*b^4*e^4)/2 + (3*B*b^6*d^2*e^2)/5 + (12*B*a*b^5*d*e^3)/5) + x^5*((\\ & A*a^6*e^4)/5 + (4*B*a^6*d*e^3)/5 + 3*A*a^2*b^4*d^4 + 4*B*a^3*b^3*d^4 + 16* \\ & A*a^3*b^3*d^3*e + 12*B*a^4*b^2*d^3*e + (36*B*a^5*b*d^2*e^2)/5 + 18*A*a^4*b \\ & ^2*d^2*e^2 + (24*A*a^5*b*d*e^3)/5) + x^8*((B*b^6*d^4)/8 + (A*b^6*d^3*e)/2 \\ & + (5*A*a^3*b^3*e^4)/2 + (15*B*a^4*b^2*e^4)/8 + (9*A*a*b^5*d^2*e^2)/2 + (15 \\ & *A*a^2*b^4*d*e^3)/2 + 10*B*a^3*b^3*d*e^3 + (45*B*a^2*b^4*d^2*e^2)/4 + 3*B* \\ & a*b^5*d^3*e) + x^6*((B*a^6*e^4)/6 + A*a*b^5*d^4 + A*a^5*b*e^4 + (5*B*a^2*b \\ & ^4*d^4)/2 + 10*A*a^2*b^4*d^3*e + 10*A*a^4*b^2*d*e^3 + (40*B*a^3*b^3*d^3*e) \\ & /3 + 20*A*a^3*b^3*d^2*e^2 + 15*B*a^4*b^2*d^2*e^2 + 4*B*a^5*b*d*e^3) + x^7* \\ & ((A*b^6*d^4)/7 + (6*B*a*b^5*d^4)/7 + (6*B*a^5*b*e^4)/7 + (15*A*a^4*b^2*e^4 \\ &)/7 + (80*A*a^3*b^3*d*e^3)/7 + (60*B*a^2*b^4*d^3*e)/7 + (60*B*a^4*b^2*d*e^ \\ & 3)/7 + (90*A*a^2*b^4*d^2*e^2)/7 + (120*B*a^3*b^3*d^2*e^2)/7 + (24*A*a*b^5* \\ & d^3*e)/7) + (a^5*d^3*x^2*(4*A*a*e + 6*A*b*d + B*a*d))/2 + (b^5*e^3*x^11*(A \\ & *b*e + 6*B*a*e + 4*B*b*d))/11 + A*a^6*d^4*x + (B*b^6*e^4*x^12)/12 \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 547, normalized size of antiderivative = 2.68

$$\int (a + bx)^6 (A + Bx)(d + ex)^4 dx$$

$$= \frac{x(330b^7e^4x^{11} + 2520ab^6e^4x^{10} + 1440b^7de^3x^{10} + 8316a^2b^5e^4x^9 + 11088ab^6de^3x^9 + 2376b^7d^2e^2x^9 + 1540a^2b^5d^2e^2x^8 + 11088ab^6de^3x^8 + 2376b^7d^2e^2x^8 + 1540a^2b^5d^2e^2x^7 + 11088ab^6de^3x^7 + 2376b^7d^2e^2x^7 + 1540a^2b^5d^2e^2x^6 + 11088ab^6de^3x^6 + 2376b^7d^2e^2x^6 + 1540a^2b^5d^2e^2x^5 + 11088ab^6de^3x^5 + 2376b^7d^2e^2x^5 + 1540a^2b^5d^2e^2x^4 + 11088ab^6de^3x^4 + 2376b^7d^2e^2x^4 + 1540a^2b^5d^2e^2x^3 + 11088ab^6de^3x^3 + 2376b^7d^2e^2x^3 + 1540a^2b^5d^2e^2x^2 + 11088ab^6de^3x^2 + 2376b^7d^2e^2x^2 + 1540a^2b^5d^2e^2x + 11088ab^6de^3x + 2376b^7d^2e^2x + 1540a^2b^5d^2e^2)}{11}$$

input `int((b*x+a)^6*(B*x+A)*(e*x+d)^4,x)`

output

```
(x*(3960*a**7*d**4 + 7920*a**7*d**3*e*x + 7920*a**7*d**2*e**2*x**2 + 3960*
a**7*d*e**3*x**3 + 792*a**7*e**4*x**4 + 13860*a**6*b*d**4*x + 36960*a**6*b
*d**3*e*x**2 + 41580*a**6*b*d**2*e**2*x**3 + 22176*a**6*b*d*e**3*x**4 + 46
20*a**6*b*e**4*x**5 + 27720*a**5*b**2*d**4*x**2 + 83160*a**5*b**2*d**3*e*x
**3 + 99792*a**5*b**2*d**2*e**2*x**4 + 55440*a**5*b**2*d*e**3*x**5 + 11880
*a**5*b**2*e**4*x**6 + 34650*a**4*b**3*d**4*x**3 + 110880*a**4*b**3*d**3*e
*x**4 + 138600*a**4*b**3*d**2*e**2*x**5 + 79200*a**4*b**3*d*e**3*x**6 + 17
325*a**4*b**3*e**4*x**7 + 27720*a**3*b**4*d**4*x**4 + 92400*a**3*b**4*d**3
*e*x**5 + 118800*a**3*b**4*d**2*e**2*x**6 + 69300*a**3*b**4*d*e**3*x**7 +
15400*a**3*b**4*e**4*x**8 + 13860*a**2*b**5*d**4*x**5 + 47520*a**2*b**5*d
**3*e*x**6 + 62370*a**2*b**5*d**2*e**2*x**7 + 36960*a**2*b**5*d*e**3*x**8 +
8316*a**2*b**5*e**4*x**9 + 3960*a*b**6*d**4*x**6 + 13860*a*b**6*d**3*e*x
**7 + 18480*a*b**6*d**2*e**2*x**8 + 11088*a*b**6*d*e**3*x**9 + 2520*a*b**6*
e**4*x**10 + 495*b**7*d**4*x**7 + 1760*b**7*d**3*e*x**8 + 2376*b**7*d**2*
e**2*x**9 + 1440*b**7*d*e**3*x**10 + 330*b**7*e**4*x**11))/3960
```

3.46 $\int (a + bx)^6 (A + Bx)(d + ex)^3 dx$

Optimal result	463
Mathematica [B] (verified)	464
Rubi [A] (verified)	465
Maple [B] (verified)	466
Fricas [B] (verification not implemented)	467
Sympy [B] (verification not implemented)	468
Maxima [B] (verification not implemented)	470
Giac [B] (verification not implemented)	472
Mupad [B] (verification not implemented)	473
Reduce [B] (verification not implemented)	474

Optimal result

Integrand size = 20, antiderivative size = 159

$$\int (a + bx)^6 (A + Bx)(d + ex)^3 dx = \frac{(Ab - aB)(bd - ae)^3 (a + bx)^7}{7b^5} + \frac{(bd - ae)^2 (bBd + 3Abe - 4aBe)(a + bx)^8}{8b^5} + \frac{e(bd - ae)(bBd + Abe - 2aBe)(a + bx)^9}{3b^5} + \frac{e^2(3bBd + Abe - 4aBe)(a + bx)^{10}}{10b^5} + \frac{Be^3(a + bx)^{11}}{11b^5}$$

output

```
1/7*(A*b-B*a)*(-a*e+b*d)^3*(b*x+a)^7/b^5+1/8*(-a*e+b*d)^2*(3*A*b*e-4*B*a*e
+B*b*d)*(b*x+a)^8/b^5+1/3*e*(-a*e+b*d)*(A*b*e-2*B*a*e+B*b*d)*(b*x+a)^9/b^5
+1/10*e^2*(A*b*e-4*B*a*e+3*B*b*d)*(b*x+a)^10/b^5+1/11*B*e^3*(b*x+a)^11/b^5
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 586 vs. $2(159) = 318$.

Time = 0.13 (sec) , antiderivative size = 586, normalized size of antiderivative = 3.69

$$\int (a + bx)^6 (A + Bx)(d + ex)^3 dx = a^6 Ad^3 x + \frac{1}{2} a^5 d^2 (6Abd + aBd + 3aAe) x^2$$

$$+ a^4 d (aBd(2bd + ae) + A(5b^2 d^2 + 6abde + a^2 e^2)) x^3$$

$$+ \frac{1}{4} a^3 (3aBd(5b^2 d^2 + 6abde + a^2 e^2)$$

$$+ A(20b^3 d^3 + 45ab^2 d^2 e + 18a^2 bde^2 + a^3 e^3)) x^4$$

$$+ \frac{1}{5} a^2 (aB(20b^3 d^3 + 45ab^2 d^2 e + 18a^2 bde^2 + a^3 e^3)$$

$$+ 3Ab(5b^3 d^3 + 20ab^2 d^2 e + 15a^2 bde^2 + 2a^3 e^3)) x^5$$

$$+ \frac{1}{2} ab (aB(5b^3 d^3 + 20ab^2 d^2 e + 15a^2 bde^2 + 2a^3 e^3)$$

$$+ Ab(2b^3 d^3 + 15ab^2 d^2 e + 20a^2 bde^2 + 5a^3 e^3)) x^6$$

$$+ \frac{1}{7} b^2 (3aB(2b^3 d^3 + 15ab^2 d^2 e + 20a^2 bde^2 + 5a^3 e^3)$$

$$+ Ab(b^3 d^3 + 18ab^2 d^2 e + 45a^2 bde^2 + 20a^3 e^3)) x^7$$

$$+ \frac{1}{8} b^3 (20a^3 Be^3 + 18ab^2 de(Bd + Ae)$$

$$+ 15a^2 be^2(3Bd + Ae) + b^3 d^2(Bd + 3Ae)) x^8$$

$$+ \frac{1}{3} b^4 e (5a^2 Be^2 + b^2 d(Bd + Ae)$$

$$+ 2abe(3Bd + Ae)) x^9$$

$$+ \frac{1}{10} b^5 e^2 (3bBd + Abe + 6aBe) x^{10} + \frac{1}{11} b^6 Be^3 x^{11}$$

input `Integrate[(a + b*x)^6*(A + B*x)*(d + e*x)^3,x]`

output

$$\begin{aligned}
& a^6 A d^3 x + (a^5 d^2 (6 A b d + a B d + 3 a A e) x^2) / 2 + a^4 d (a B d (2 b d + a e) + A (5 b^2 d^2 + 6 a b d e + a^2 e^2)) x^3 + (a^3 (3 a B d (5 b^2 d^2 + 6 a b d e + a^2 e^2) + A (20 b^3 d^3 + 45 a b^2 d^2 e + 18 a^2 b d e^2 + a^3 e^3)) x^4) / 4 + (a^2 (a B (20 b^3 d^3 + 45 a b^2 d^2 e + 18 a^2 b d e^2 + a^3 e^3) + 3 A b (5 b^3 d^3 + 20 a b^2 d^2 e + 15 a^2 b d e^2 + 2 a^3 e^3)) x^5) / 5 + (a b (a B (5 b^3 d^3 + 20 a b^2 d^2 e + 15 a^2 b d e^2 + 2 a^3 e^3) + A b (2 b^3 d^3 + 15 a b^2 d^2 e + 20 a^2 b d e^2 + 5 a^3 e^3)) x^6) / 2 + (b^2 (3 a B (2 b^3 d^3 + 15 a b^2 d^2 e + 20 a^2 b d e^2 + 5 a^3 e^3) + A b (b^3 d^3 + 18 a b^2 d^2 e + 45 a^2 b d e^2 + 20 a^3 e^3)) x^7) / 7 + (b^3 (20 a^3 B e^3 + 18 a b^2 d e (B d + A e) + 15 a^2 b e^2 (3 B d + A e) + b^3 d^2 (B d + 3 A e)) x^8) / 8 + (b^4 e (5 a^2 B e^2 + b^2 d (B d + A e) + 2 a b e (3 B d + A e)) x^9) / 3 + (b^5 e^2 (3 b B d + A b e + 6 a B e) x^{10}) / 10 + (b^6 B e^3 x^{11}) / 11
\end{aligned}$$
Rubi [A] (verified)

Time = 0.68 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int (a + bx)^6 (A + Bx)(d + ex)^3 dx \\
& \quad \downarrow 86 \\
& \int \left(\frac{e^2 (a + bx)^9 (-4aBe + Abe + 3bBd)}{b^4} + \frac{3e(a + bx)^8 (bd - ae)(-2aBe + Abe + bBd)}{b^4} + \frac{(a + bx)^7 (bd - ae)^2 (-2aBe + Abe + bBd)}{b^4} \right) dx \\
& \quad \downarrow 2009 \\
& \frac{e^2 (a + bx)^{10} (-4aBe + Abe + 3bBd)}{10b^5} + \frac{e(a + bx)^9 (bd - ae)(-2aBe + Abe + bBd)}{3b^5} + \frac{(a + bx)^8 (bd - ae)^2 (-4aBe + 3Abe + bBd)}{8b^5} + \frac{(a + bx)^7 (Ab - aB)(bd - ae)^3}{7b^5} + \frac{Be^3 (a + bx)^{11}}{11b^5}
\end{aligned}$$

input

$$\text{Int}[(a + b*x)^6*(A + B*x)*(d + e*x)^3,x]$$

output

```

1/11*b^6*B*e^3*x^11+1/10*((A*b^6+6*B*a*b^5)*e^3+3*b^6*B*d*e^2)*x^10+1/9*((
6*A*a*b^5+15*B*a^2*b^4)*e^3+3*(A*b^6+6*B*a*b^5)*d*e^2+3*b^6*B*d^2*e)*x^9+1
/8*((15*A*a^2*b^4+20*B*a^3*b^3)*e^3+3*(6*A*a*b^5+15*B*a^2*b^4)*d*e^2+3*(A*
b^6+6*B*a*b^5)*d^2*e+b^6*B*d^3)*x^8+1/7*((20*A*a^3*b^3+15*B*a^4*b^2)*e^3+3
*(15*A*a^2*b^4+20*B*a^3*b^3)*d*e^2+3*(6*A*a*b^5+15*B*a^2*b^4)*d^2*e+(A*b^6
+6*B*a*b^5)*d^3)*x^7+1/6*((15*A*a^4*b^2+6*B*a^5*b)*e^3+3*(20*A*a^3*b^3+15*
B*a^4*b^2)*d*e^2+3*(15*A*a^2*b^4+20*B*a^3*b^3)*d^2*e+(6*A*a*b^5+15*B*a^2*b
^4)*d^3)*x^6+1/5*((6*A*a^5*b+B*a^6)*e^3+3*(15*A*a^4*b^2+6*B*a^5*b)*d*e^2+3
*(20*A*a^3*b^3+15*B*a^4*b^2)*d^2*e+(15*A*a^2*b^4+20*B*a^3*b^3)*d^3)*x^5+1/
4*(a^6*A*e^3+3*(6*A*a^5*b+B*a^6)*d*e^2+3*(15*A*a^4*b^2+6*B*a^5*b)*d^2*e+(2
0*A*a^3*b^3+15*B*a^4*b^2)*d^3)*x^4+1/3*(3*a^6*A*d*e^2+3*(6*A*a^5*b+B*a^6)*
d^2*e+(15*A*a^4*b^2+6*B*a^5*b)*d^3)*x^3+1/2*(3*a^6*A*d^2*e+(6*A*a^5*b+B*a^
6)*d^3)*x^2+a^6*A*d^3*x

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 643 vs. $2(149) = 298$.

Time = 0.08 (sec) , antiderivative size = 643, normalized size of antiderivative = 4.04

$$\begin{aligned}
& \int (a + bx)^6 (A + Bx)(d + ex)^3 dx \\
&= \frac{1}{11} Bb^6 e^3 x^{11} + Aa^6 d^3 x + \frac{1}{10} (3 Bb^6 de^2 + (6 Bab^5 + Ab^6) e^3) x^{10} \\
&+ \frac{1}{3} (Bb^6 d^2 e + (6 Bab^5 + Ab^6) de^2 + (5 Ba^2 b^4 + 2 Aab^5) e^3) x^9 \\
&+ \frac{1}{8} (Bb^6 d^3 + 3 (6 Bab^5 + Ab^6) d^2 e + 9 (5 Ba^2 b^4 + 2 Aab^5) de^2 + 5 (4 Ba^3 b^3 + 3 Aa^2 b^4) e^3) x^8 \\
&+ \frac{1}{7} ((6 Bab^5 + Ab^6) d^3 + 9 (5 Ba^2 b^4 + 2 Aab^5) d^2 e + 15 (4 Ba^3 b^3 + 3 Aa^2 b^4) de^2 + 5 (3 Ba^4 b^2 + 4 Aa^3 b^3) e^3) x^7 \\
&+ \frac{1}{2} ((5 Ba^2 b^4 + 2 Aab^5) d^3 + 5 (4 Ba^3 b^3 + 3 Aa^2 b^4) d^2 e + 5 (3 Ba^4 b^2 + 4 Aa^3 b^3) de^2 + (2 Ba^5 b + 5 Aa^4 b^2) e^3) x^6 \\
&+ \frac{1}{5} (5 (4 Ba^3 b^3 + 3 Aa^2 b^4) d^3 + 15 (3 Ba^4 b^2 + 4 Aa^3 b^3) d^2 e + 9 (2 Ba^5 b + 5 Aa^4 b^2) de^2 + (Ba^6 + 6 Aa^5 b) e^3) x^5 \\
&+ \frac{1}{4} (Aa^6 e^3 + 5 (3 Ba^4 b^2 + 4 Aa^3 b^3) d^3 + 9 (2 Ba^5 b + 5 Aa^4 b^2) d^2 e + 3 (Ba^6 + 6 Aa^5 b) de^2) x^4 \\
&+ (Aa^6 de^2 + (2 Ba^5 b + 5 Aa^4 b^2) d^3 + (Ba^6 + 6 Aa^5 b) d^2 e) x^3 \\
&+ \frac{1}{2} (3 Aa^6 d^2 e + (Ba^6 + 6 Aa^5 b) d^3) x^2
\end{aligned}$$

input

```
integrate((b*x+a)^6*(B*x+A)*(e*x+d)^3,x, algorithm="fricas")
```


output

```

1/11*B*b^6*e^3*x^11 + A*a^6*d^3*x + 1/10*(3*B*b^6*d*e^2 + (6*B*a*b^5 + A*b
^6)*e^3)*x^10 + 1/3*(B*b^6*d^2*e + (6*B*a*b^5 + A*b^6)*d*e^2 + (5*B*a^2*b^
4 + 2*A*a*b^5)*e^3)*x^9 + 1/8*(B*b^6*d^3 + 3*(6*B*a*b^5 + A*b^6)*d^2*e + 9
*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^2 + 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^3)*x^8
+ 1/7*((6*B*a*b^5 + A*b^6)*d^3 + 9*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e + 15*(4
*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^2 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^3)*x^7 +
1/2*((5*B*a^2*b^4 + 2*A*a*b^5)*d^3 + 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e
+ 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^2 + (2*B*a^5*b + 5*A*a^4*b^2)*e^3)*x^6
+ 1/5*(5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3 + 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)
*d^2*e + 9*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^2 + (B*a^6 + 6*A*a^5*b)*e^3)*x^5
+ 1/4*(A*a^6*e^3 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3 + 9*(2*B*a^5*b + 5*A*
a^4*b^2)*d^2*e + 3*(B*a^6 + 6*A*a^5*b)*d*e^2)*x^4 + (A*a^6*d*e^2 + (2*B*a^
5*b + 5*A*a^4*b^2)*d^3 + (B*a^6 + 6*A*a^5*b)*d^2*e)*x^3 + 1/2*(3*A*a^6*d^2
*e + (B*a^6 + 6*A*a^5*b)*d^3)*x^2

```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 802 vs. $2(155) = 310$.

Time = 0.06 (sec) , antiderivative size = 802, normalized size of antiderivative = 5.04

$$\begin{aligned}
 \int (a + bx)^6 (A + Bx)(d + ex)^3 dx = & Aa^6 d^3 x + \frac{Bb^6 e^3 x^{11}}{11} \\
 & + x^{10} \left(\frac{Ab^6 e^3}{10} + \frac{3Bab^5 e^3}{5} + \frac{3Bb^6 d e^2}{10} \right) + x^9 \\
 & \cdot \left(\frac{2Aab^5 e^3}{3} + \frac{Ab^6 d e^2}{3} + \frac{5Ba^2 b^4 e^3}{3} + 2Bab^5 d e^2 \right. \\
 & \quad \left. + \frac{Bb^6 d^2 e}{3} \right) + x^8 \\
 & \cdot \left(\frac{15Aa^2 b^4 e^3}{8} + \frac{9Aab^5 d e^2}{4} + \frac{3Ab^6 d^2 e}{8} + \frac{5Ba^3 b^3 e^3}{2} \right. \\
 & \quad \left. + \frac{45Ba^2 b^4 d e^2}{8} + \frac{9Bab^5 d^2 e}{4} + \frac{Bb^6 d^3}{8} \right) \\
 & + x^7 \cdot \left(\frac{20Aa^3 b^3 e^3}{7} + \frac{45Aa^2 b^4 d e^2}{7} + \frac{18Aab^5 d^2 e}{7} \right. \\
 & \quad \left. + \frac{Ab^6 d^3}{7} + \frac{15Ba^4 b^2 e^3}{7} + \frac{60Ba^3 b^3 d e^2}{7} \right. \\
 & \quad \left. + \frac{45Ba^2 b^4 d^2 e}{7} + \frac{6Bab^5 d^3}{7} \right) + x^6 \\
 & \cdot \left(\frac{5Aa^4 b^2 e^3}{2} + 10Aa^3 b^3 d e^2 + \frac{15Aa^2 b^4 d^2 e}{2} + Aab^5 d^3 \right. \\
 & \quad \left. + Ba^5 b e^3 + \frac{15Ba^4 b^2 d e^2}{2} + 10Ba^3 b^3 d^2 e + \frac{5Ba^2 b^4 d^3}{2} \right) \\
 & + x^5 \\
 & \cdot \left(\frac{6Aa^5 b e^3}{5} + 9Aa^4 b^2 d e^2 + 12Aa^3 b^3 d^2 e + 3Aa^2 b^4 d^3 \right. \\
 & \quad \left. + \frac{Ba^6 e^3}{5} + \frac{18Ba^5 b d e^2}{5} + 9Ba^4 b^2 d^2 e + 4Ba^3 b^3 d^3 \right) \\
 & + x^4 \left(\frac{Aa^6 e^3}{4} + \frac{9Aa^5 b d e^2}{2} + \frac{45Aa^4 b^2 d^2 e}{4} \right. \\
 & \quad \left. + 5Aa^3 b^3 d^3 + \frac{3Ba^6 d e^2}{4} + \frac{9Ba^5 b d^2 e}{2} + \frac{15Ba^4 b^2 d^3}{4} \right) \\
 & + x^3 (Aa^6 d e^2 + 6Aa^5 b d^2 e + 5Aa^4 b^2 d^3 + Ba^6 d^2 e \\
 & \quad + 2Ba^5 b d^3) + x^2 \cdot \left(\frac{3Aa^6 d^2 e}{2} + 3Aa^5 b d^3 + \frac{Ba^6 d^3}{2} \right)
 \end{aligned}$$

input

```
integrate((b*x+a)**6*(B*x+A)*(e*x+d)**3,x)
```

output

```

A*a**6*d**3*x + B*b**6*e**3*x**11/11 + x**10*(A*b**6*e**3/10 + 3*B*a*b**5*
e**3/5 + 3*B*b**6*d*e**2/10) + x**9*(2*A*a*b**5*e**3/3 + A*b**6*d*e**2/3 +
5*B*a**2*b**4*e**3/3 + 2*B*a*b**5*d*e**2 + B*b**6*d**2*e/3) + x**8*(15*A*
a**2*b**4*e**3/8 + 9*A*a*b**5*d*e**2/4 + 3*A*b**6*d**2*e/8 + 5*B*a**3*b**3
*e**3/2 + 45*B*a**2*b**4*d*e**2/8 + 9*B*a*b**5*d**2*e/4 + B*b**6*d**3/8) +
x**7*(20*A*a**3*b**3*e**3/7 + 45*A*a**2*b**4*d*e**2/7 + 18*A*a*b**5*d**2*
e/7 + A*b**6*d**3/7 + 15*B*a**4*b**2*e**3/7 + 60*B*a**3*b**3*d*e**2/7 + 45
*B*a**2*b**4*d**2*e/7 + 6*B*a*b**5*d**3/7) + x**6*(5*A*a**4*b**2*e**3/2 +
10*A*a**3*b**3*d*e**2 + 15*A*a**2*b**4*d**2*e/2 + A*a*b**5*d**3 + B*a**5*b
*e**3 + 15*B*a**4*b**2*d*e**2/2 + 10*B*a**3*b**3*d**2*e + 5*B*a**2*b**4*d*
*3/2) + x**5*(6*A*a**5*b*e**3/5 + 9*A*a**4*b**2*d*e**2 + 12*A*a**3*b**3*d*
*2*e + 3*A*a**2*b**4*d**3 + B*a**6*e**3/5 + 18*B*a**5*b*d*e**2/5 + 9*B*a**
4*b**2*d**2*e + 4*B*a**3*b**3*d**3) + x**4*(A*a**6*e**3/4 + 9*A*a**5*b*d*e
**2/2 + 45*A*a**4*b**2*d**2*e/4 + 5*A*a**3*b**3*d**3 + 3*B*a**6*d*e**2/4 +
9*B*a**5*b*d**2*e/2 + 15*B*a**4*b**2*d**3/4) + x**3*(A*a**6*d*e**2 + 6*A*
a**5*b*d**2*e + 5*A*a**4*b**2*d**3 + B*a**6*d**2*e + 2*B*a**5*b*d**3) + x
*2*(3*A*a**6*d**2*e/2 + 3*A*a**5*b*d**3 + B*a**6*d**3/2)

```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 643 vs. $2(149) = 298$.

Time = 0.05 (sec) , antiderivative size = 643, normalized size of antiderivative = 4.04

$$\int (a + bx)^6 (A + Bx)(d + ex)^3 dx$$

$$= \frac{1}{11} Bb^6 e^3 x^{11} + Aa^6 d^3 x + \frac{1}{10} (3 Bb^6 d e^2 + (6 Bab^5 + Ab^6) e^3) x^{10}$$

$$+ \frac{1}{3} (Bb^6 d^2 e + (6 Bab^5 + Ab^6) d e^2 + (5 Ba^2 b^4 + 2 Aab^5) e^3) x^9$$

$$+ \frac{1}{8} (Bb^6 d^3 + 3 (6 Bab^5 + Ab^6) d^2 e + 9 (5 Ba^2 b^4 + 2 Aab^5) d e^2 + 5 (4 Ba^3 b^3 + 3 Aa^2 b^4) e^3) x^8$$

$$+ \frac{1}{7} ((6 Bab^5 + Ab^6) d^3 + 9 (5 Ba^2 b^4 + 2 Aab^5) d^2 e + 15 (4 Ba^3 b^3 + 3 Aa^2 b^4) d e^2 + 5 (3 Ba^4 b^2 + 4 Aa^3 b^3) e^3) x^7$$

$$+ \frac{1}{2} ((5 Ba^2 b^4 + 2 Aab^5) d^3 + 5 (4 Ba^3 b^3 + 3 Aa^2 b^4) d^2 e + 5 (3 Ba^4 b^2 + 4 Aa^3 b^3) d e^2 + (2 Ba^5 b + 5 Aa^4 b^2) e^3) x^6$$

$$+ \frac{1}{5} (5 (4 Ba^3 b^3 + 3 Aa^2 b^4) d^3 + 15 (3 Ba^4 b^2 + 4 Aa^3 b^3) d^2 e + 9 (2 Ba^5 b + 5 Aa^4 b^2) d e^2 + (Ba^6 + 6 Aa^5 b) e^3) x^5$$

$$+ \frac{1}{4} (Aa^6 e^3 + 5 (3 Ba^4 b^2 + 4 Aa^3 b^3) d^3 + 9 (2 Ba^5 b + 5 Aa^4 b^2) d^2 e + 3 (Ba^6 + 6 Aa^5 b) d e^2) x^4$$

$$+ (Aa^6 d e^2 + (2 Ba^5 b + 5 Aa^4 b^2) d^3 + (Ba^6 + 6 Aa^5 b) d^2 e) x^3$$

$$+ \frac{1}{2} (3 Aa^6 d^2 e + (Ba^6 + 6 Aa^5 b) d^3) x^2$$

input

```
integrate((b*x+a)^6*(B*x+A)*(e*x+d)^3,x, algorithm="maxima")
```

output

```
1/11*B*b^6*e^3*x^11 + A*a^6*d^3*x + 1/10*(3*B*b^6*d*e^2 + (6*B*a*b^5 + A*b^6)*e^3)*x^10 + 1/3*(B*b^6*d^2*e + (6*B*a*b^5 + A*b^6)*d*e^2 + (5*B*a^2*b^4 + 2*A*a*b^5)*e^3)*x^9 + 1/8*(B*b^6*d^3 + 3*(6*B*a*b^5 + A*b^6)*d^2*e + 9*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^2 + 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^3)*x^8 + 1/7*((6*B*a*b^5 + A*b^6)*d^3 + 9*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e + 15*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^2 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^3)*x^7 + 1/2*((5*B*a^2*b^4 + 2*A*a*b^5)*d^3 + 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^2 + (2*B*a^5*b + 5*A*a^4*b^2)*e^3)*x^6 + 1/5*(5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3 + 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e + 9*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^2 + (B*a^6 + 6*A*a^5*b)*e^3)*x^5 + 1/4*(A*a^6*e^3 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3 + 9*(2*B*a^5*b + 5*A*a^4*b^2)*d^2*e + 3*(B*a^6 + 6*A*a^5*b)*d*e^2)*x^4 + (A*a^6*d*e^2 + (2*B*a^5*b + 5*A*a^4*b^2)*d^3 + (B*a^6 + 6*A*a^5*b)*d^2*e)*x^3 + 1/2*(3*A*a^6*d^2*e + (B*a^6 + 6*A*a^5*b)*d^3)*x^2
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 782 vs. $2(149) = 298$.

Time = 0.12 (sec) , antiderivative size = 782, normalized size of antiderivative = 4.92

$$\int (a + bx)^6 (A + Bx)(d + ex)^3 dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)*(e*x+d)^3,x, algorithm="giac")`

output

```
1/11*B*b^6*e^3*x^11 + 3/10*B*b^6*d*e^2*x^10 + 3/5*B*a*b^5*e^3*x^10 + 1/10*
A*b^6*e^3*x^10 + 1/3*B*b^6*d^2*e*x^9 + 2*B*a*b^5*d*e^2*x^9 + 1/3*A*b^6*d*e
^2*x^9 + 5/3*B*a^2*b^4*e^3*x^9 + 2/3*A*a*b^5*e^3*x^9 + 1/8*B*b^6*d^3*x^8 +
9/4*B*a*b^5*d^2*e*x^8 + 3/8*A*b^6*d^2*e*x^8 + 45/8*B*a^2*b^4*d*e^2*x^8 +
9/4*A*a*b^5*d*e^2*x^8 + 5/2*B*a^3*b^3*e^3*x^8 + 15/8*A*a^2*b^4*e^3*x^8 + 6
/7*B*a*b^5*d^3*x^7 + 1/7*A*b^6*d^3*x^7 + 45/7*B*a^2*b^4*d^2*e*x^7 + 18/7*A
*a*b^5*d^2*e*x^7 + 60/7*B*a^3*b^3*d*e^2*x^7 + 45/7*A*a^2*b^4*d*e^2*x^7 + 1
5/7*B*a^4*b^2*e^3*x^7 + 20/7*A*a^3*b^3*e^3*x^7 + 5/2*B*a^2*b^4*d^3*x^6 + A
*a*b^5*d^3*x^6 + 10*B*a^3*b^3*d^2*e*x^6 + 15/2*A*a^2*b^4*d^2*e*x^6 + 15/2*
B*a^4*b^2*d*e^2*x^6 + 10*A*a^3*b^3*d*e^2*x^6 + B*a^5*b*e^3*x^6 + 5/2*A*a^4
*b^2*e^3*x^6 + 4*B*a^3*b^3*d^3*x^5 + 3*A*a^2*b^4*d^3*x^5 + 9*B*a^4*b^2*d^2
*e*x^5 + 12*A*a^3*b^3*d^2*e*x^5 + 18/5*B*a^5*b*d*e^2*x^5 + 9*A*a^4*b^2*d*e
^2*x^5 + 1/5*B*a^6*e^3*x^5 + 6/5*A*a^5*b*e^3*x^5 + 15/4*B*a^4*b^2*d^3*x^4
+ 5*A*a^3*b^3*d^3*x^4 + 9/2*B*a^5*b*d^2*e*x^4 + 45/4*A*a^4*b^2*d^2*e*x^4 +
3/4*B*a^6*d*e^2*x^4 + 9/2*A*a^5*b*d*e^2*x^4 + 1/4*A*a^6*e^3*x^4 + 2*B*a^5
*b*d^3*x^3 + 5*A*a^4*b^2*d^3*x^3 + B*a^6*d^2*e*x^3 + 6*A*a^5*b*d^2*e*x^3 +
A*a^6*d*e^2*x^3 + 1/2*B*a^6*d^3*x^2 + 3*A*a^5*b*d^3*x^2 + 3/2*A*a^6*d^2*e
*x^2 + A*a^6*d^3*x
```

Mupad [B] (verification not implemented)

Time = 1.00 (sec) , antiderivative size = 649, normalized size of antiderivative = 4.08

$$\begin{aligned}
\int (a + bx)^6 (A + Bx)(d + ex)^3 dx = & x^3 (B a^6 d^2 e + A a^6 d e^2 + 2 B a^5 b d^3 + 6 A a^5 b d^2 e \\
& + 5 A a^4 b^2 d^3) \\
& + x^9 \left(\frac{5 B a^2 b^4 e^3}{3} + 2 B a b^5 d e^2 + \frac{2 A a b^5 e^3}{3} \right. \\
& \left. + \frac{B b^6 d^2 e}{3} + \frac{A b^6 d e^2}{3} \right) + x^4 \left(\frac{3 B a^6 d e^2}{4} \right. \\
& \left. + \frac{A a^6 e^3}{4} + \frac{9 B a^5 b d^2 e}{2} + \frac{9 A a^5 b d e^2}{2} \right. \\
& \left. + \frac{15 B a^4 b^2 d^3}{4} + \frac{45 A a^4 b^2 d^2 e}{4} + 5 A a^3 b^3 d^3 \right) \\
& + x^8 \left(\frac{5 B a^3 b^3 e^3}{2} + \frac{45 B a^2 b^4 d e^2}{8} + \frac{15 A a^2 b^4 e^3}{8} \right. \\
& \left. + \frac{9 B a b^5 d^2 e}{4} + \frac{9 A a b^5 d e^2}{4} + \frac{B b^6 d^3}{8} \right. \\
& \left. + \frac{3 A b^6 d^2 e}{8} \right) + x^6 \left(B a^5 b e^3 + \frac{15 B a^4 b^2 d e^2}{2} \right. \\
& \left. + \frac{5 A a^4 b^2 e^3}{2} + 10 B a^3 b^3 d^2 e + 10 A a^3 b^3 d e^2 \right. \\
& \left. + \frac{5 B a^2 b^4 d^3}{2} + \frac{15 A a^2 b^4 d^2 e}{2} + A a b^5 d^3 \right) \\
& + x^5 \left(\frac{B a^6 e^3}{5} + \frac{18 B a^5 b d e^2}{5} + \frac{6 A a^5 b e^3}{5} \right. \\
& \left. + 9 B a^4 b^2 d^2 e + 9 A a^4 b^2 d e^2 + 4 B a^3 b^3 d^3 \right. \\
& \left. + 12 A a^3 b^3 d^2 e + 3 A a^2 b^4 d^3 \right) \\
& + x^7 \left(\frac{15 B a^4 b^2 e^3}{7} + \frac{60 B a^3 b^3 d e^2}{7} + \frac{20 A a^3 b^3 e^3}{7} \right. \\
& \left. + \frac{45 B a^2 b^4 d^2 e}{7} + \frac{45 A a^2 b^4 d e^2}{7} + \frac{6 B a b^5 d^3}{7} \right. \\
& \left. + \frac{18 A a b^5 d^2 e}{7} + \frac{A b^6 d^3}{7} \right) \\
& + \frac{a^5 d^2 x^2 (3 A a e + 6 A b d + B a d)}{2} \\
& + \frac{b^5 e^2 x^{10} (A b e + 6 B a e + 3 B b d)}{10} \\
& + A a^6 d^3 x + \frac{B b^6 e^3 x^{11}}{11}
\end{aligned}$$

input `int((A + B*x)*(a + b*x)^6*(d + e*x)^3,x)`

output

$$\begin{aligned} & x^3*(2*B*a^5*b*d^3 + A*a^6*d*e^2 + B*a^6*d^2*e + 5*A*a^4*b^2*d^3 + 6*A*a^5 \\ & *b*d^2*e) + x^9*((2*A*a*b^5*e^3)/3 + (A*b^6*d*e^2)/3 + (B*b^6*d^2*e)/3 + (\\ & 5*B*a^2*b^4*e^3)/3 + 2*B*a*b^5*d*e^2) + x^4*((A*a^6*e^3)/4 + (3*B*a^6*d*e^ \\ & 2)/4 + 5*A*a^3*b^3*d^3 + (15*B*a^4*b^2*d^3)/4 + (45*A*a^4*b^2*d^2*e)/4 + (\\ & 9*A*a^5*b*d*e^2)/2 + (9*B*a^5*b*d^2*e)/2) + x^8*((B*b^6*d^3)/8 + (3*A*b^6* \\ & d^2*e)/8 + (15*A*a^2*b^4*e^3)/8 + (5*B*a^3*b^3*e^3)/2 + (45*B*a^2*b^4*d*e^ \\ & 2)/8 + (9*A*a*b^5*d*e^2)/4 + (9*B*a*b^5*d^2*e)/4) + x^6*(A*a*b^5*d^3 + B*a \\ & ^5*b*e^3 + (5*A*a^4*b^2*e^3)/2 + (5*B*a^2*b^4*d^3)/2 + (15*A*a^2*b^4*d^2*e \\ &)/2 + 10*A*a^3*b^3*d*e^2 + 10*B*a^3*b^3*d^2*e + (15*B*a^4*b^2*d*e^2)/2) + \\ & x^5*((B*a^6*e^3)/5 + (6*A*a^5*b*e^3)/5 + 3*A*a^2*b^4*d^3 + 4*B*a^3*b^3*d^3 \\ & + 12*A*a^3*b^3*d^2*e + 9*A*a^4*b^2*d*e^2 + 9*B*a^4*b^2*d^2*e + (18*B*a^5* \\ & b*d*e^2)/5) + x^7*((A*b^6*d^3)/7 + (6*B*a*b^5*d^3)/7 + (20*A*a^3*b^3*e^3)/ \\ & 7 + (15*B*a^4*b^2*e^3)/7 + (45*A*a^2*b^4*d*e^2)/7 + (45*B*a^2*b^4*d^2*e)/7 \\ & + (60*B*a^3*b^3*d*e^2)/7 + (18*A*a*b^5*d^2*e)/7) + (a^5*d^2*x^2*(3*A*a*e \\ & + 6*A*b*d + B*a*d))/2 + (b^5*e^2*x^10*(A*b*e + 6*B*a*e + 3*B*b*d))/10 + A* \\ & a^6*d^3*x + (B*b^6*e^3*x^11)/11 \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 421, normalized size of antiderivative = 2.65

$$\int (a + bx)^6 (A + Bx)(d + ex)^3 dx$$

$$= \frac{x(120b^7e^3x^{10} + 924ab^6e^3x^9 + 396b^7de^2x^9 + 3080a^2b^5e^3x^8 + 3080ab^6de^2x^8 + 440b^7d^2ex^8 + 5775a^3b^4e^3x^7 + 3080a^2b^5de^2x^7 + 180ab^6d^2ex^7 + 180a^2b^5d^2e^2x^7 + 120ab^6d^3ex^6 + 120a^2b^5d^3e^2x^6 + 60ab^6d^4ex^6 + 60a^2b^5d^4e^2x^6 + 30ab^6d^5ex^5 + 30a^2b^5d^5e^2x^5 + 15ab^6d^6ex^5 + 15a^2b^5d^6e^2x^5 + 5ab^6d^7ex^4 + 5a^2b^5d^7e^2x^4 + b^6d^8ex^4 + a^2b^5d^8e^2x^4 + 5ab^6d^9ex^3 + 5a^2b^5d^9e^2x^3 + b^6d^{10}ex^3 + a^2b^5d^{10}e^2x^3 + 5ab^6d^{11}ex^2 + 5a^2b^5d^{11}e^2x^2 + b^6d^{12}ex^2 + a^2b^5d^{12}e^2x^2 + 5ab^6d^{13}ex + 5a^2b^5d^{13}e^2x + b^6d^{14}ex + a^2b^5d^{14}e^2x + b^6d^{15}e^2x + a^2b^5d^{15}e^2x + b^6d^{16}e^2x + a^2b^5d^{16}e^2x + b^6d^{17}e^2x + a^2b^5d^{17}e^2x + b^6d^{18}e^2x + a^2b^5d^{18}e^2x + b^6d^{19}e^2x + a^2b^5d^{19}e^2x + b^6d^{20}e^2x + a^2b^5d^{20}e^2x + b^6d^{21}e^2x + a^2b^5d^{21}e^2x + b^6d^{22}e^2x + a^2b^5d^{22}e^2x + b^6d^{23}e^2x + a^2b^5d^{23}e^2x + b^6d^{24}e^2x + a^2b^5d^{24}e^2x + b^6d^{25}e^2x + a^2b^5d^{25}e^2x + b^6d^{26}e^2x + a^2b^5d^{26}e^2x + b^6d^{27}e^2x + a^2b^5d^{27}e^2x + b^6d^{28}e^2x + a^2b^5d^{28}e^2x + b^6d^{29}e^2x + a^2b^5d^{29}e^2x + b^6d^{30}e^2x + a^2b^5d^{30}e^2x + b^6d^{31}e^2x + a^2b^5d^{31}e^2x + b^6d^{32}e^2x + a^2b^5d^{32}e^2x + b^6d^{33}e^2x + a^2b^5d^{33}e^2x + b^6d^{34}e^2x + a^2b^5d^{34}e^2x + b^6d^{35}e^2x + a^2b^5d^{35}e^2x + b^6d^{36}e^2x + a^2b^5d^{36}e^2x + b^6d^{37}e^2x + a^2b^5d^{37}e^2x + b^6d^{38}e^2x + a^2b^5d^{38}e^2x + b^6d^{39}e^2x + a^2b^5d^{39}e^2x + b^6d^{40}e^2x + a^2b^5d^{40}e^2x + b^6d^{41}e^2x + a^2b^5d^{41}e^2x + b^6d^{42}e^2x + a^2b^5d^{42}e^2x + b^6d^{43}e^2x + a^2b^5d^{43}e^2x + b^6d^{44}e^2x + a^2b^5d^{44}e^2x + b^6d^{45}e^2x + a^2b^5d^{45}e^2x + b^6d^{46}e^2x + a^2b^5d^{46}e^2x + b^6d^{47}e^2x + a^2b^5d^{47}e^2x + b^6d^{48}e^2x + a^2b^5d^{48}e^2x + b^6d^{49}e^2x + a^2b^5d^{49}e^2x + b^6d^{50}e^2x + a^2b^5d^{50}e^2x + b^6d^{51}e^2x + a^2b^5d^{51}e^2x + b^6d^{52}e^2x + a^2b^5d^{52}e^2x + b^6d^{53}e^2x + a^2b^5d^{53}e^2x + b^6d^{54}e^2x + a^2b^5d^{54}e^2x + b^6d^{55}e^2x + a^2b^5d^{55}e^2x + b^6d^{56}e^2x + a^2b^5d^{56}e^2x + b^6d^{57}e^2x + a^2b^5d^{57}e^2x + b^6d^{58}e^2x + a^2b^5d^{58}e^2x + b^6d^{59}e^2x + a^2b^5d^{59}e^2x + b^6d^{60}e^2x + a^2b^5d^{60}e^2x + b^6d^{61}e^2x + a^2b^5d^{61}e^2x + b^6d^{62}e^2x + a^2b^5d^{62}e^2x + b^6d^{63}e^2x + a^2b^5d^{63}e^2x + b^6d^{64}e^2x + a^2b^5d^{64}e^2x + b^6d^{65}e^2x + a^2b^5d^{65}e^2x + b^6d^{66}e^2x + a^2b^5d^{66}e^2x + b^6d^{67}e^2x + a^2b^5d^{67}e^2x + b^6d^{68}e^2x + a^2b^5d^{68}e^2x + b^6d^{69}e^2x + a^2b^5d^{69}e^2x + b^6d^{70}e^2x + a^2b^5d^{70}e^2x + b^6d^{71}e^2x + a^2b^5d^{71}e^2x + b^6d^{72}e^2x + a^2b^5d^{72}e^2x + b^6d^{73}e^2x + a^2b^5d^{73}e^2x + b^6d^{74}e^2x + a^2b^5d^{74}e^2x + b^6d^{75}e^2x + a^2b^5d^{75}e^2x + b^6d^{76}e^2x + a^2b^5d^{76}e^2x + b^6d^{77}e^2x + a^2b^5d^{77}e^2x + b^6d^{78}e^2x + a^2b^5d^{78}e^2x + b^6d^{79}e^2x + a^2b^5d^{79}e^2x + b^6d^{80}e^2x + a^2b^5d^{80}e^2x + b^6d^{81}e^2x + a^2b^5d^{81}e^2x + b^6d^{82}e^2x + a^2b^5d^{82}e^2x + b^6d^{83}e^2x + a^2b^5d^{83}e^2x + b^6d^{84}e^2x + a^2b^5d^{84}e^2x + b^6d^{85}e^2x + a^2b^5d^{85}e^2x + b^6d^{86}e^2x + a^2b^5d^{86}e^2x + b^6d^{87}e^2x + a^2b^5d^{87}e^2x + b^6d^{88}e^2x + a^2b^5d^{88}e^2x + b^6d^{89}e^2x + a^2b^5d^{89}e^2x + b^6d^{90}e^2x + a^2b^5d^{90}e^2x + b^6d^{91}e^2x + a^2b^5d^{91}e^2x + b^6d^{92}e^2x + a^2b^5d^{92}e^2x + b^6d^{93}e^2x + a^2b^5d^{93}e^2x + b^6d^{94}e^2x + a^2b^5d^{94}e^2x + b^6d^{95}e^2x + a^2b^5d^{95}e^2x + b^6d^{96}e^2x + a^2b^5d^{96}e^2x + b^6d^{97}e^2x + a^2b^5d^{97}e^2x + b^6d^{98}e^2x + a^2b^5d^{98}e^2x + b^6d^{99}e^2x + a^2b^5d^{99}e^2x + b^6d^{100}e^2x + a^2b^5d^{100}e^2x + b^6d^{101}e^2x + a^2b^5d^{101}e^2x + b^6d^{102}e^2x + a^2b^5d^{102}e^2x + b^6d^{103}e^2x + a^2b^5d^{103}e^2x + b^6d^{104}e^2x + a^2b^5d^{104}e^2x + b^6d^{105}e^2x + a^2b^5d^{105}e^2x + b^6d^{106}e^2x + a^2b^5d^{106}e^2x + b^6d^{107}e^2x + a^2b^5d^{107}e^2x + b^6d^{108}e^2x + a^2b^5d^{108}e^2x + b^6d^{109}e^2x + a^2b^5d^{109}e^2x + b^6d^{110}e^2x + a^2b^5d^{110}e^2x + b^6d^{111}e^2x + a^2b^5d^{111}e^2x + b^6d^{112}e^2x + a^2b^5d^{112}e^2x + b^6d^{113}e^2x + a^2b^5d^{113}e^2x + b^6d^{114}e^2x + a^2b^5d^{114}e^2x + b^6d^{115}e^2x + a^2b^5d^{115}e^2x + b^6d^{116}e^2x + a^2b^5d^{116}e^2x + b^6d^{117}e^2x + a^2b^5d^{117}e^2x + b^6d^{118}e^2x + a^2b^5d^{118}e^2x + b^6d^{119}e^2x + a^2b^5d^{119}e^2x + b^6d^{120}e^2x + a^2b^5d^{120}e^2x + b^6d^{121}e^2x + a^2b^5d^{121}e^2x + b^6d^{122}e^2x + a^2b^5d^{122}e^2x + b^6d^{123}e^2x + a^2b^5d^{123}e^2x + b^6d^{124}e^2x + a^2b^5d^{124}e^2x + b^6d^{125}e^2x + a^2b^5d^{125}e^2x + b^6d^{126}e^2x + a^2b^5d^{126}e^2x + b^6d^{127}e^2x + a^2b^5d^{127}e^2x + b^6d^{128}e^2x + a^2b^5d^{128}e^2x + b^6d^{129}e^2x + a^2b^5d^{129}e^2x + b^6d^{130}e^2x + a^2b^5d^{130}e^2x + b^6d^{131}e^2x + a^2b^5d^{131}e^2x + b^6d^{132}e^2x + a^2b^5d^{132}e^2x + b^6d^{133}e^2x + a^2b^5d^{133}e^2x + b^6d^{134}e^2x + a^2b^5d^{134}e^2x + b^6d^{135}e^2x + a^2b^5d^{135}e^2x + b^6d^{136}e^2x + a^2b^5d^{136}e^2x + b^6d^{137}e^2x + a^2b^5d^{137}e^2x + b^6d^{138}e^2x + a^2b^5d^{138}e^2x + b^6d^{139}e^2x + a^2b^5d^{139}e^2x + b^6d^{140}e^2x + a^2b^5d^{140}e^2x + b^6d^{141}e^2x + a^2b^5d^{141}e^2x + b^6d^{142}e^2x + a^2b^5d^{142}e^2x + b^6d^{143}e^2x + a^2b^5d^{143}e^2x + b^6d^{144}e^2x + a^2b^5d^{144}e^2x + b^6d^{145}e^2x + a^2b^5d^{145}e^2x + b^6d^{146}e^2x + a^2b^5d^{146}e^2x + b^6d^{147}e^2x + a^2b^5d^{147}e^2x + b^6d^{148}e^2x + a^2b^5d^{148}e^2x + b^6d^{149}e^2x + a^2b^5d^{149}e^2x + b^6d^{150}e^2x + a^2b^5d^{150}e^2x + b^6d^{151}e^2x + a^2b^5d^{151}e^2x + b^6d^{152}e^2x + a^2b^5d^{152}e^2x + b^6d^{153}e^2x + a^2b^5d^{153}e^2x + b^6d^{154}e^2x + a^2b^5d^{154}e^2x + b^6d^{155}e^2x + a^2b^5d^{155}e^2x + b^6d^{156}e^2x + a^2b^5d^{156}e^2x + b^6d^{157}e^2x + a^2b^5d^{157}e^2x + b^6d^{158}e^2x + a^2b^5d^{158}e^2x + b^6d^{159}e^2x + a^2b^5d^{159}e^2x + b^6d^{160}e^2x + a^2b^5d^{160}e^2x + b^6d^{161}e^2x + a^2b^5d^{161}e^2x + b^6d^{162}e^2x + a^2b^5d^{162}e^2x + b^6d^{163}e^2x + a^2b^5d^{163}e^2x + b^6d^{164}e^2x + a^2b^5d^{164}e^2x + b^6d^{165}e^2x + a^2b^5d^{165}e^2x + b^6d^{166}e^2x + a^2b^5d^{166}e^2x + b^6d^{167}e^2x + a^2b^5d^{167}e^2x + b^6d^{168}e^2x + a^2b^5d^{168}e^2x + b^6d^{169}e^2x + a^2b^5d^{169}e^2x + b^6d^{170}e^2x + a^2b^5d^{170}e^2x + b^6d^{171}e^2x + a^2b^5d^{171}e^2x + b^6d^{172}e^2x + a^2b^5d^{172}e^2x + b^6d^{173}e^2x + a^2b^5d^{173}e^2x + b^6d^{174}e^2x + a^2b^5d^{174}e^2x + b^6d^{175}e^2x + a^2b^5d^{175}e^2x + b^6d^{176}e^2x + a^2b^5d^{176}e^2x + b^6d^{177}e^2x + a^2b^5d^{177}e^2x + b^6d^{178}e^2x + a^2b^5d^{178}e^2x + b^6d^{179}e^2x + a^2b^5d^{179}e^2x + b^6d^{180}e^2x + a^2b^5d^{180}e^2x + b^6d^{181}e^2x + a^2b^5d^{181}e^2x + b^6d^{182}e^2x + a^2b^5d^{182}e^2x + b^6d^{183}e^2x + a^2b^5d^{183}e^2x + b^6d^{184}e^2x + a^2b^5d^{184}e^2x + b^6d^{185}e^2x + a^2b^5d^{185}e^2x + b^6d^{186}e^2x + a^2b^5d^{186}e^2x + b^6d^{187}e^2x + a^2b^5d^{187}e^2x + b^6d^{188}e^2x + a^2b^5d^{188}e^2x + b^6d^{189}e^2x + a^2b^5d^{189}e^2x + b^6d^{190}e^2x + a^2b^5d^{190}e^2x + b^6d^{191}e^2x + a^2b^5d^{191}e^2x + b^6d^{192}e^2x + a^2b^5d^{192}e^2x + b^6d^{193}e^2x + a^2b^5d^{193}e^2x + b^6d^{194}e^2x + a^2b^5d^{194}e^2x + b^6d^{195}e^2x + a^2b^5d^{195}e^2x + b^6d^{196}e^2x + a^2b^5d^{196}e^2x + b^6d^{197}e^2x + a^2b^5d^{197}e^2x + b^6d^{198}e^2x + a^2b^5d^{198}e^2x + b^6d^{199}e^2x + a^2b^5d^{199}e^2x + b^6d^{200}e^2x + a^2b^5d^{200}e^2x + b^6d^{201}e^2x + a^2b^5d^{201}e^2x + b^6d^{202}e^2x + a^2b^5d^{202}e^2x + b^6d^{203}e^2x + a^2b^5d^{203}e^2x + b^6d^{204}e^2x + a^2b^5d^{204}e^2x + b^6d^{205}e^2x + a^2b^5d^{205}e^2x + b^6d^{206}e^2x + a^2b^5d^{206}e^2x + b^6d^{207}e^2x + a^2b^5d^{207}e^2x + b^6d^{208}e^2x + a^2b^5d^{208}e^2x + b^6d^{209}e^2x + a^2b^5d^{209}e^2x + b^6d^{210}e^2x + a^2b^5d^{210}e^2x + b^6d^{211}e^2x + a^2b^5d^{211}e^2x + b^6d^{212}e^2x + a^2b^5d^{212}e^2x + b^6d^{213}e^2x + a^2b^5d^{213}e^2x + b^6d^{214}e^2x + a^2b^5d^{214}e^2x + b^6d^{215}e^2x + a^2b^5d^{215}e^2x + b^6d^{216}e^2x + a^2b^5d^{216}e^2x + b^6d^{217}e^2x + a^2b^5d^{217}e^2x + b^6d^{218}e^2x + a^2b^5d^{218}e^2x + b^6d^{219}e^2x + a^2b^5d^{219}e^2x + b^6d^{220}e^2x + a^2b^5d^{220}e^2x + b^6d^{221}e^2x + a^2b^5d^{221}e^2x + b^6d^{222}e^2x + a^2b^5d^{222}e^2x + b^6d^{223}e^2x + a^2b^5d^{223}e^2x + b^6d^{224}e^2x + a^2b^5d^{224}e^2x + b^6d^{225}e^2x + a^2b^5d^{225}e^2x + b^6d^{226}e^2x + a^2b^5d^{226}e^2x + b^6d^{227}e^2x + a^2b^5d^{227}e^2x + b^6d^{228}e^2x + a^2b^5d^{228}e^2x + b^6d^{229}e^2x + a^2b^5d^{229}e^2x + b^6d^{230}e^2x + a^2b^5d^{230}e^2x + b^6d^{231}e^2x + a^2b^5d^{231}e^2x + b^6d^{232}e^2x + a^2b^5d^{232}e^2x + b^6d^{233}e^2x + a^2b^5d^{233}e^2x + b^6d^{234}e^2x + a^2b^5d^{234}e^2x + b^6d^{235}e^2x + a^2b^5d^{235}e^2x + b^6d^{236}e^2x + a^2b^5d^{236}e^2x + b^6d^{237}e^2x + a^2b^5d^{237}e^2x + b^6d^{238}e^2x + a^2b^5d^{238}e^2x + b^6d^{239}e^2x + a^2b^5d^{239}e^2x + b^6d^{240}e^2x + a^2b^5d^{240}e^2x + b^6d^{241}e^2x + a^2b^5d^{241}e^2x + b^6d^{242}e^2x + a^2b^5d^{242}e^2x + b^6d^{243}e^2x + a^2b^5d^{243}e^2x + b^6d^{244}e^2x + a^2b^5d^{244}e^2x + b^6d^{245}e^2x + a^2b^5d^{245}e^2x + b^6d^{246}e^2x + a^2b^5d^{246}e^2x + b^6d^{247}e^2x + a^2b^5d^{247}e^2x + b^6d^{248}e^2x + a^2b^5d^{248}e^2x + b^6d^{249}e^2x + a^2b^5d^{249}e^2x + b^6d^{250}e^2x + a^2b^5d^{250}e^2x + b^6d^{251}e^2x + a^2b^5d^{251}e^2x + b^6d^{252}e^2x + a^2b^5d^{252}e^2x + b^6d^{253}e^2x + a^2b^5d^{253}e^2x + b^6d^{254}e^2x + a^2b^5d^{254}e^2x + b^6d^{255}e^2x + a^2b^5d^{255}e^2x + b^6d^{256}e^2x + a^2b^5d^{256}e^2x + b^6d^{257}e^2x + a^2b^5d^{257}e^2x + b^6d^{258}e^2x + a^2b^5d^{258}e^2x + b^6d^{259}e^2x + a^2b^5d^{259}e^2x + b^6d^{260}e^2x + a^2b^5d^{260}e^2x + b^6d^{261}e^2x + a^2b^5d^{261}e^2x + b^6d^{262}e^2x + a^2b^5d^{262}e^2x + b^6d^{263}e^2x + a^2b^5d^{263}e^2x + b^6d^{264}e^2x + a^2b^5d^{264}e^2x + b^6d^{265}e^2x + a^2b^5d^{265}e^2x + b^6d^{266}e^2x + a^2b^5d^{266}e^2x + b^6d^{267}e^2x + a^2b^5d^{267}e^2x + b^6d^{268}e^2x + a^2b^5d^{268}e^2x + b^6d^{269}e^2x + a^2b^5d^{269}e^2x + b^6d^{270}e^2x + a^2b^5d^{270}e^2x + b^6d^{271}e^2x + a^2b^5d^{271}e^2x + b^6d^{272}e^2x + a^2b^5d^{272}e^2x + b^6d^{273}e^2x + a^2b^5d^{273}e^2x + b^6d^{274}e^2x + a^2b^5d^{274}e^2x + b^6d^{275}e^2x + a^2b^5d^{275}e^2x + b^6d^{276}e^2x + a^2b^5d^{276}e^2x + b^6d^{277}e^2x + a^2b^5d^{277}e^2x + b^6d^{278}e^2x + a^2b^5d^{278}e^2x + b^6d^{279}e^2x + a^2b^5d^{279}e^2x + b^6d^{280}e^2x + a^2b^5d^{280}e^2x + b^6d^{281}e^2x + a^2b^5d^{281}e^2x + b^6d^{282}e^2x + a^2b^5d^{282}e^2x + b^6d^{283}e^2x + a^2b^5d^{283}e^2x + b^6d^{284}e^2x + a^2b^5d^{284}e^2x + b^6d^{285}e^2x + a^2b^5d^{285}e^2x + b^6d^{286}e^2x + a^2b^5d^{286}e^2x + b^6d^{287}e^2x + a^2b^5d^{287}e^2x + b^6d^{288}e^2x + a^2b^5d^{288}e^2x + b^6d^{289}e^2x + a^2b^5d^{289}e^2x + b^6d^{290}e^2x + a^2b^5d^{290}e^2x + b^6d^{291}e^2x + a^2b^5d^{291}e^2x + b^6d^{292}e^2x + a^2b^5d^{292}e^2x + b^6d^{293}e^2x + a^2b^5d^{293}e^2x + b^6d^{294}e^2x + a^2b^5d^{294}e^2x + b^6d^{295}e^2x + a^2b^5d^{295}e^2x + b^6d^{296}e^2x + a^2b^5d^{296}e^2x + b^6d^{297}e^2x + a^2b^5d^{297}e^2x + b^6d^{298}e^2x + a^2b^5d^{298}e^2x + b^6d^{299}e^2x + a^2b^5d^{299}e^2x + b^6d^{300}e^2x + a^2b^5d^{300}e^2x + b^6d^{301}e^2x + a^2b^5d^{301}e^2x + b^6d^{302}e^2x + a^2b^5d^{302}e^2x + b^6d^{303}e^2x + a^2b^5d^{303}e^2x + b^6d^{304}e^2x + a^2b^5d^{304}e^2x + b^6d^{305}e^2x + a^2b^5d^{305}e^2x + b^6d^{306}e^2x + a^2b^5d^{306}e^2x + b^6d^{307}e^2x + a^2b^5d^{307}e^2x + b^6d^{308}e^2x + a^2b^5d^{308}e^2x + b^6d^{309}e^2x + a^2b^5d^{309}e^2x + b^6d^{310}e^2x + a^2b^5d^{310}e^2x + b^6d^{311}e^2x + a^2b^5d^{311}e^2x + b^6d^{312}e^2x + a^2b^5d^{312}e^2x + b^6d^{313}e^2x + a^2b^5d^{313}e^2x + b^6d^{314}e^2x + a^2b^5d^{314}e^2x + b^6d^{315}e^2x + a^2b^5d^{315}e^2x + b^6d^{316}e^2x + a^2b^5d^{316}e^2x + b^6d^{317}e^2x + a^2b^5d^{317}e^2x + b^6d^{318}e^2x + a^2b^5d^{318}e^2x + b^6d^{319}e^2x + a^2b^5d^{319}e^2x + b^6d^{320}e^2x + a^2b^5d^{320}e^2x + b^6d^{321}e^2x + a^2b^5d^{321}e^2x + b^6d^{322}e^2x + a^2b^5d^{322}e^2x + b^6d^{323}e^2x + a^2b^5d^{323}e^2x + b^6d^{324}e^2x + a^2b^5d^{324}e^2x + b^6d^{325}e^2x + a^2b^5d^{325}e^2x + b^6d^{326}e^2x + a^2b^5d^{326}e^2x + b^6d^{327}e^2x + a^2b^5d^{327}e^2x + b^6d^{328}e^2x + a^2b^5d^{328}e^2x + b^6d^{329}e^2x + a^2b^5d^{329}e^2x + b^6d^{330}e^2x + a^2b^5d^{330}e^2x + b^6d^{331}e^2x + a^2b^5d^{331}e^2x + b^6d^{332}e^2x + a^2b^5d^{332}e^2x + b^6d^{333}e^2x + a^2b^5d^{333}e^2x + b^6d^{334}e^2x + a^2b^5d^{334}e^2x + b^6d^{335}e^2x + a^2b^5d^{335}e^2x + b^6d^{336}e^2x + a^2b^5d^{336}e^2x + b^6d^{337}e^2x + a^2b^5d^{337}e^2x + b^6d^{338}e^2x + a^2b^5d^{338}e^2x + b^6d^{339}e^2x + a^2b^5d^{339}e^2x + b^6d^{340}e^2x + a^2b^5d^{340}e^2x + b^6d^{341}e^2x + a^2b^5d^{341}e^2x + b^6d^{342}e^2x + a^2b^5d^{342}e^2x + b^6d^{343}e^2x + a^2b^5d^{343}e^2x + b^6d^{344}e^2x + a^2b^5d^{344}e^2x + b^6d^{345}e^2x + a^2b^5d^{345}e^2x + b^6d^{346}e^2x + a^2b^5d^{346}e^2x + b^6d^{347}e^2x + a^2b^5d^{347}e^2x + b^6d^{348}e^2x + a^2b^5d^{348}e^2x + b^6d^{349}e^2x + a^2b^5d^{349}e^2x + b^6d^{350}e^2x + a^2b^5d^{350}e^2x + b^6d^{351}e^2x + a^2b^5d^{351}e^2x + b^6d^{352}e^2x + a^2b^5d^{352}e^2x + b^6d^{353}e^2x + a^2b^5d^{353}e^2x + b^6d^{354}e^2x + a^2b^5d^{354}e^2x + b^6d^{355}e^2x + a^2b^5d^{355}e^2x + b^6d^{356}e^2x + a^2b^5d^{356}e^2x + b^6d^{357}e^2x + a^2b^5d^{357}e^2x + b^6d^{358}e^2x + a^2b^5d^{358}e^2x + b^6d^{359}e^2x + a^2b^5d^{359}e^2x + b^6d^{360}e^2x + a^2b^5d^{360}e^2x + b^6d^{361}e^2x + a^2b^5d^{361}e^2x + b^6d^{362}e^2x + a^2b^5d^{362}e^2x + b^6d^{363}e^2x$$

output

```
(x*(1320*a**7*d**3 + 1980*a**7*d**2*e*x + 1320*a**7*d*e**2*x**2 + 330*a**7*
e**3*x**3 + 4620*a**6*b*d**3*x + 9240*a**6*b*d**2*e*x**2 + 6930*a**6*b*d*
e**2*x**3 + 1848*a**6*b*e**3*x**4 + 9240*a**5*b**2*d**3*x**2 + 20790*a**5*
b**2*d**2*e*x**3 + 16632*a**5*b**2*d*e**2*x**4 + 4620*a**5*b**2*e**3*x**5
+ 11550*a**4*b**3*d**3*x**3 + 27720*a**4*b**3*d**2*e*x**4 + 23100*a**4*b**
3*d*e**2*x**5 + 6600*a**4*b**3*e**3*x**6 + 9240*a**3*b**4*d**3*x**4 + 2310
0*a**3*b**4*d**2*e*x**5 + 19800*a**3*b**4*d*e**2*x**6 + 5775*a**3*b**4*e**
3*x**7 + 4620*a**2*b**5*d**3*x**5 + 11880*a**2*b**5*d**2*e*x**6 + 10395*a**
2*b**5*d*e**2*x**7 + 3080*a**2*b**5*e**3*x**8 + 1320*a*b**6*d**3*x**6 + 3
465*a*b**6*d**2*e*x**7 + 3080*a*b**6*d*e**2*x**8 + 924*a*b**6*e**3*x**9 +
165*b**7*d**3*x**7 + 440*b**7*d**2*e*x**8 + 396*b**7*d*e**2*x**9 + 120*b**
7*e**3*x**10))/1320
```


3.47 $\int (a + bx)^6 (A + Bx)(d + ex)^2 dx$

Optimal result	476
Mathematica [B] (verified)	477
Rubi [A] (verified)	477
Maple [B] (verified)	479
Fricas [B] (verification not implemented)	480
Sympy [B] (verification not implemented)	481
Maxima [B] (verification not implemented)	482
Giac [B] (verification not implemented)	483
Mupad [B] (verification not implemented)	486
Reduce [B] (verification not implemented)	487

Optimal result

Integrand size = 20, antiderivative size = 118

$$\int (a + bx)^6 (A + Bx)(d + ex)^2 dx = \frac{(Ab - aB)(bd - ae)^2(a + bx)^7}{7b^4} + \frac{(bd - ae)(bBd + 2Abe - 3aBe)(a + bx)^8}{8b^4} + \frac{e(2bBd + Abe - 3aBe)(a + bx)^9}{9b^4} + \frac{Be^2(a + bx)^{10}}{10b^4}$$

output

```
1/7*(A*b-B*a)*(-a*e+b*d)^2*(b*x+a)^7/b^4+1/8*(-a*e+b*d)*(2*A*b*e-3*B*a*e+B
*b*d)*(b*x+a)^8/b^4+1/9*e*(A*b*e-3*B*a*e+2*B*b*d)*(b*x+a)^9/b^4+1/10*B*e^2
*(b*x+a)^10/b^4
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 386 vs. $2(118) = 236$.

Time = 0.14 (sec) , antiderivative size = 386, normalized size of antiderivative = 3.27

$$\int (a + bx)^6 (A + Bx)(d + ex)^2 dx$$

$$= \frac{x(210a^6(4A(3d^2 + 3dex + e^2x^2) + Bx(6d^2 + 8dex + 3e^2x^2)) + 252a^5bx(5A(6d^2 + 8dex + 3e^2x^2) + 2Bx(10d^2 + 15dex + 6e^2x^2)) + 630a^4b^2x^2(2A(10d^2 + 15dex + 6e^2x^2) + Bx(15d^2 + 24dex + 10e^2x^2)) + 120a^3b^3x^3(7A(15d^2 + 24dex + 10e^2x^2) + 4Bx(21d^2 + 35dex + 15e^2x^2)) + 45a^2b^4x^4(8A(21d^2 + 35dex + 15e^2x^2) + 5Bx(28d^2 + 48dex + 21e^2x^2)) + 30ab^5x^5(3A(28d^2 + 48dex + 21e^2x^2) + 2Bx(36d^2 + 63dex + 28e^2x^2)) + b^6x^6(10A(36d^2 + 63dex + 28e^2x^2) + 7Bx(45d^2 + 80dex + 36e^2x^2)))}{2520}$$

input `Integrate[(a + b*x)^6*(A + B*x)*(d + e*x)^2,x]`

output

```
(x*(210*a^6*(4*A*(3*d^2 + 3*d*e*x + e^2*x^2) + B*x*(6*d^2 + 8*d*e*x + 3*e^2*x^2)) + 252*a^5*b*x*(5*A*(6*d^2 + 8*d*e*x + 3*e^2*x^2) + 2*B*x*(10*d^2 + 15*d*e*x + 6*e^2*x^2)) + 630*a^4*b^2*x^2*(2*A*(10*d^2 + 15*d*e*x + 6*e^2*x^2) + B*x*(15*d^2 + 24*d*e*x + 10*e^2*x^2)) + 120*a^3*b^3*x^3*(7*A*(15*d^2 + 24*d*e*x + 10*e^2*x^2) + 4*B*x*(21*d^2 + 35*d*e*x + 15*e^2*x^2)) + 45*a^2*b^4*x^4*(8*A*(21*d^2 + 35*d*e*x + 15*e^2*x^2) + 5*B*x*(28*d^2 + 48*d*e*x + 21*e^2*x^2)) + 30*a*b^5*x^5*(3*A*(28*d^2 + 48*d*e*x + 21*e^2*x^2) + 2*B*x*(36*d^2 + 63*d*e*x + 28*e^2*x^2)) + b^6*x^6*(10*A*(36*d^2 + 63*d*e*x + 28*e^2*x^2) + 7*B*x*(45*d^2 + 80*d*e*x + 36*e^2*x^2)))/2520
```

Rubi [A] (verified)

Time = 0.55 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^6 (A + Bx)(d + ex)^2 dx$$

↓ 86

$$\int \left(\frac{e(a + bx)^8(-3aBe + Abe + 2bBd)}{b^3} + \frac{(a + bx)^7(bd - ae)(-3aBe + 2Abe + bBd)}{b^3} + \frac{(a + bx)^6(Ab - aB)(bd - ae)}{b^3} \right) dx$$

$$\begin{array}{c} \downarrow \text{2009} \\ \frac{e(a+bx)^9(-3aBe + Abe + 2bBd)}{9b^4} + \frac{(a+bx)^8(bd - ae)(-3aBe + 2Abe + bBd)}{8b^4} + \\ \frac{(a+bx)^7(Ab - aB)(bd - ae)^2}{7b^4} + \frac{Be^2(a+bx)^{10}}{10b^4} \end{array}$$

input `Int[(a + b*x)^6*(A + B*x)*(d + e*x)^2,x]`

output `((A*b - a*B)*(b*d - a*e)^2*(a + b*x)^7)/(7*b^4) + ((b*d - a*e)*(b*B*d + 2*A*b*e - 3*a*B*e)*(a + b*x)^8)/(8*b^4) + (e*(2*b*B*d + A*b*e - 3*a*B*e)*(a + b*x)^9)/(9*b^4) + (B*e^2*(a + b*x)^10)/(10*b^4)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;`
`FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 468 vs. $2(110) = 220$.

Time = 0.18 (sec) , antiderivative size = 469, normalized size of antiderivative = 3.97

method	result
default	$\frac{b^6 B e^2 x^{10}}{10} + \frac{((b^6 A + 6a b^5 B)e^2 + 2b^6 B d e)x^9}{9} + \frac{((6a b^5 A + 15a^2 b^4 B)e^2 + 2(b^6 A + 6a b^5 B)d e + b^6 B d^2)x^8}{8} + \frac{((15a^2 b^4 A + 20a b^3 B)e^2 + 2(6a^2 b^3 A + 15a b^2 B)d e + b^6 B d^2)x^7}{7} + \frac{((15a^2 b^4 A + 20a b^3 B)e^2 + 2(6a^2 b^3 A + 15a b^2 B)d e + b^6 B d^2)x^6}{6} + \frac{((15a^2 b^4 A + 20a b^3 B)e^2 + 2(6a^2 b^3 A + 15a b^2 B)d e + b^6 B d^2)x^5}{5} + \frac{((15a^2 b^4 A + 20a b^3 B)e^2 + 2(6a^2 b^3 A + 15a b^2 B)d e + b^6 B d^2)x^4}{4} + \frac{((15a^2 b^4 A + 20a b^3 B)e^2 + 2(6a^2 b^3 A + 15a b^2 B)d e + b^6 B d^2)x^3}{3} + \frac{((15a^2 b^4 A + 20a b^3 B)e^2 + 2(6a^2 b^3 A + 15a b^2 B)d e + b^6 B d^2)x^2}{2} + \frac{((15a^2 b^4 A + 20a b^3 B)e^2 + 2(6a^2 b^3 A + 15a b^2 B)d e + b^6 B d^2)x}{1} + \frac{((15a^2 b^4 A + 20a b^3 B)e^2 + 2(6a^2 b^3 A + 15a b^2 B)d e + b^6 B d^2)}{0}$
norman	$\frac{b^6 B e^2 x^{10}}{10} + (\frac{1}{9} A b^6 e^2 + \frac{2}{3} B a b^5 e^2 + \frac{2}{9} b^6 B d e) x^9 + (\frac{3}{4} A a b^5 e^2 + \frac{1}{4} A b^6 d e + \frac{15}{8} B a^2 b^4 e^2 + \frac{3}{2} B a b^5 d e + \frac{15}{8} B a^2 b^4 d^2) x^8 + \frac{x(252b^6 B e^2 x^9 + 280A b^6 e^2 x^8 + 1680B a b^5 e^2 x^8 + 560B b^6 d e x^8 + 1890A a b^5 e^2 x^7 + 630A b^6 d e x^7 + 4725B a^2 b^4 e^2 x^7 + 3780B a b^5 d e x^7 + 1575B a^2 b^4 d^2 x^7 + 1296B a b^5 d^2 x^7 + 864B a^2 b^4 d^2 x^7 + 648B a b^5 d^2 x^7 + 432B a^2 b^4 d^2 x^7 + 324B a b^5 d^2 x^7 + 216B a^2 b^4 d^2 x^7 + 162B a b^5 d^2 x^7 + 108B a^2 b^4 d^2 x^7 + 81B a b^5 d^2 x^7 + 54B a^2 b^4 d^2 x^7 + 40.5B a b^5 d^2 x^7 + 27B a^2 b^4 d^2 x^7 + 20.25B a b^5 d^2 x^7 + 13.5B a^2 b^4 d^2 x^7 + 10.125B a b^5 d^2 x^7 + 6.75B a^2 b^4 d^2 x^7 + 5.0625B a b^5 d^2 x^7 + 3.375B a^2 b^4 d^2 x^7 + 2.53125B a b^5 d^2 x^7 + 1.6875B a^2 b^4 d^2 x^7 + 1.265625B a b^5 d^2 x^7 + 0.84375B a^2 b^4 d^2 x^7 + 0.6328125B a b^5 d^2 x^7 + 0.421875B a^2 b^4 d^2 x^7 + 0.31640625B a b^5 d^2 x^7 + 0.2109375B a^2 b^4 d^2 x^7 + 0.158203125B a b^5 d^2 x^7 + 0.11865234375B a^2 b^4 d^2 x^7 + 0.0890796875B a b^5 d^2 x^7 + 0.0668095703125B a^2 b^4 d^2 x^7 + 0.0501071875B a b^5 d^2 x^7 + 0.03758046875B a^2 b^4 d^2 x^7 + 0.0281853515625B a b^5 d^2 x^7 + 0.021139015625B a^2 b^4 d^2 x^7 + 0.015854271875B a b^5 d^2 x^7 + 0.011890703125B a^2 b^4 d^2 x^7 + 0.00891796875B a b^5 d^2 x^7 + 0.006688515625B a^2 b^4 d^2 x^7 + 0.00501640625B a b^5 d^2 x^7 + 0.0037548046875B a^2 b^4 d^2 x^7 + 0.002816103515625B a b^5 d^2 x^7 + 0.002112078125B a^2 b^4 d^2 x^7 + 0.001584059375B a b^5 d^2 x^7 + 0.00118804453125B a^2 b^4 d^2 x^7 + 0.000891033203125B a b^5 d^2 x^7 + 0.0006680248046875B a^2 b^4 d^2 x^7 + 0.00050101853125B a b^5 d^2 x^7 + 0.0003750141875B a^2 b^4 d^2 x^7 + 0.000281260640625B a b^5 d^2 x^7 + 0.00021124548046875B a^2 b^4 d^2 x^7 + 0.0001584341015625B a b^5 d^2 x^7 + 0.000118825578125B a^2 b^4 d^2 x^7 + 8.9104171875e-05B a b^5 d^2 x^7 + 6.68031296875e-05B a^2 b^4 d^2 x^7 + 5.0102346875e-05B a b^5 d^2 x^7 + 3.7501759375e-05B a^2 b^4 d^2 x^7 + 2.81263696875e-05B a b^5 d^2 x^7 + 2.1124883046875e-05B a^2 b^4 d^2 x^7 + 1.58436623125e-05B a b^5 d^2 x^7 + 1.188274671875e-05B a^2 b^4 d^2 x^7 + 8.91105903125e-06B a b^5 d^2 x^7 + 6.68079421875e-06B a^2 b^4 d^2 x^7 + 5.010595640625e-06B a b^5 d^2 x^7 + 3.75044671875e-06B a^2 b^4 d^2 x^7 + 2.81267259375e-06B a b^5 d^2 x^7 + 2.1124990625e-06B a^2 b^4 d^2 x^7 + 1.5843796875e-06B a b^5 d^2 x^7 + 1.1882846875e-06B a^2 b^4 d^2 x^7 + 8.911153125e-07B a b^5 d^2 x^7 + 6.6808871875e-07B a^2 b^4 d^2 x^7 + 5.01068046875e-07B a b^5 d^2 x^7 + 3.750571875e-07B a^2 b^4 d^2 x^7 + 2.812763046875e-07B a b^5 d^2 x^7 + 2.112584375e-07B a^2 b^4 d^2 x^7 + 1.5844640625e-07B a b^5 d^2 x^7 + 1.1883703125e-07B a^2 b^4 d^2 x^7 + 8.9112471875e-08B a b^5 d^2 x^7 + 6.68097846875e-08B a^2 b^4 d^2 x^7 + 5.01076640625e-08B a b^5 d^2 x^7 + 3.7506571875e-08B a^2 b^4 d^2 x^7 + 2.81284875e-08B a b^5 d^2 x^7 + 2.1126690625e-08B a^2 b^4 d^2 x^7 + 1.58454846875e-08B a b^5 d^2 x^7 + 1.18845571875e-08B a^2 b^4 d^2 x^7 + 8.9113411875e-09B a b^5 d^2 x^7 + 6.6810646875e-09B a^2 b^4 d^2 x^7 + 5.01085246875e-09B a b^5 d^2 x^7 + 3.750743046875e-09B a^2 b^4 d^2 x^7 + 2.812934375e-09B a b^5 d^2 x^7 + 2.1127540625e-09B a^2 b^4 d^2 x^7 + 1.584632875e-09B a b^5 d^2 x^7 + 1.1885410625e-09B a^2 b^4 d^2 x^7 + 8.9114351875e-10B a b^5 d^2 x^7 + 6.6811506875e-10B a^2 b^4 d^2 x^7 + 5.01093846875e-10B a b^5 d^2 x^7 + 3.75082875e-10B a^2 b^4 d^2 x^7 + 2.8130196875e-10B a b^5 d^2 x^7 + 2.1128390625e-10B a^2 b^4 d^2 x^7 + 1.584717275e-10B a b^5 d^2 x^7 + 1.18862646875e-10B a^2 b^4 d^2 x^7 + 8.9115291875e-11B a b^5 d^2 x^7 + 6.6812366875e-11B a^2 b^4 d^2 x^7 + 5.01102446875e-11B a b^5 d^2 x^7 + 3.7509146875e-11B a^2 b^4 d^2 x^7 + 2.813105375e-11B a b^5 d^2 x^7 + 2.1129240625e-11B a^2 b^4 d^2 x^7 + 1.584801675e-11B a b^5 d^2 x^7 + 1.188711875e-11B a^2 b^4 d^2 x^7 + 8.9116231875e-12B a b^5 d^2 x^7 + 6.6813226875e-12B a^2 b^4 d^2 x^7 + 5.01111046875e-12B a b^5 d^2 x^7 + 3.7510006875e-12B a^2 b^4 d^2 x^7 + 2.8131906875e-12B a b^5 d^2 x^7 + 2.1130090625e-12B a^2 b^4 d^2 x^7 + 1.584886075e-12B a b^5 d^2 x^7 + 1.188797275e-12B a^2 b^4 d^2 x^7 + 8.9117171875e-13B a b^5 d^2 x^7 + 6.6814086875e-13B a^2 b^4 d^2 x^7 + 5.01119646875e-13B a b^5 d^2 x^7 + 3.7510866875e-13B a^2 b^4 d^2 x^7 + 2.813276375e-13B a b^5 d^2 x^7 + 2.1130940625e-13B a^2 b^4 d^2 x^7 + 1.584970475e-13B a b^5 d^2 x^7 + 1.188882675e-13B a^2 b^4 d^2 x^7 + 8.9118111875e-14B a b^5 d^2 x^7 + 6.6814946875e-14B a^2 b^4 d^2 x^7 + 5.01128246875e-14B a b^5 d^2 x^7 + 3.7511726875e-14B a^2 b^4 d^2 x^7 + 2.81336206875e-14B a b^5 d^2 x^7 + 2.1131790625e-14B a^2 b^4 d^2 x^7 + 1.585054875e-14B a b^5 d^2 x^7 + 1.188968075e-14B a^2 b^4 d^2 x^7 + 8.9119051875e-15B a b^5 d^2 x^7 + 6.6815806875e-15B a^2 b^4 d^2 x^7 + 5.01136846875e-15B a b^5 d^2 x^7 + 3.7512586875e-15B a^2 b^4 d^2 x^7 + 2.8134476875e-15B a b^5 d^2 x^7 + 2.1132640625e-15B a^2 b^4 d^2 x^7 + 1.585139275e-15B a b^5 d^2 x^7 + 1.18905346875e-15B a^2 b^4 d^2 x^7 + 8.9120001875e-16B a b^5 d^2 x^7 + 6.6816666875e-16B a^2 b^4 d^2 x^7 + 5.01145446875e-16B a b^5 d^2 x^7 + 3.7513446875e-16B a^2 b^4 d^2 x^7 + 2.813533375e-16B a b^5 d^2 x^7 + 2.1133490625e-16B a^2 b^4 d^2 x^7 + 1.585223675e-16B a b^5 d^2 x^7 + 1.189138875e-16B a^2 b^4 d^2 x^7 + 8.9120941875e-17B a b^5 d^2 x^7 + 6.6817526875e-17B a^2 b^4 d^2 x^7 + 5.01154046875e-17B a b^5 d^2 x^7 + 3.7514306875e-17B a^2 b^4 d^2 x^7 + 2.81361906875e-17B a b^5 d^2 x^7 + 2.1134340625e-17B a^2 b^4 d^2 x^7 + 1.585308075e-17B a b^5 d^2 x^7 + 1.189224275e-17B a^2 b^4 d^2 x^7 + 8.9121881875e-18B a b^5 d^2 x^7 + 6.6818386875e-18B a^2 b^4 d^2 x^7 + 5.01162646875e-18B a b^5 d^2 x^7 + 3.7515166875e-18B a^2 b^4 d^2 x^7 + 2.8137046875e-18B a b^5 d^2 x^7 + 2.1135190625e-18B a^2 b^4 d^2 x^7 + 1.585392475e-18B a b^5 d^2 x^7 + 1.189309675e-18B a^2 b^4 d^2 x^7 + 8.9122821875e-19B a b^5 d^2 x^7 + 6.6819246875e-19B a^2 b^4 d^2 x^7 + 5.01171246875e-19B a b^5 d^2 x^7 + 3.7516026875e-19B a^2 b^4 d^2 x^7 + 2.813790375e-19B a b^5 d^2 x^7 + 2.1136040625e-19B a^2 b^4 d^2 x^7 + 1.585476875e-19B a b^5 d^2 x^7 + 1.189395075e-19B a^2 b^4 d^2 x^7 + 8.9123761875e-20B a b^5 d^2 x^7 + 6.6820106875e-20B a^2 b^4 d^2 x^7 + 5.01179846875e-20B a b^5 d^2 x^7 + 3.7516886875e-20B a^2 b^4 d^2 x^7 + 2.81387606875e-20B a b^5 d^2 x^7 + 2.1136890625e-20B a^2 b^4 d^2 x^7 + 1.585561275e-20B a b^5 d^2 x^7 + 1.18948046875e-20B a^2 b^4 d^2 x^7 + 8.9124701875e-21B a b^5 d^2 x^7 + 6.6820966875e-21B a^2 b^4 d^2 x^7 + 5.01188446875e-21B a b^5 d^2 x^7 + 3.7517746875e-21B a^2 b^4 d^2 x^7 + 2.8139616875e-21B a b^5 d^2 x^7 + 2.1137740625e-21B a^2 b^4 d^2 x^7 + 1.585645675e-21B a b^5 d^2 x^7 + 1.189565875e-21B a^2 b^4 d^2 x^7 + 8.9125641875e-22B a b^5 d^2 x^7 + 6.6821826875e-22B a^2 b^4 d^2 x^7 + 5.01197046875e-22B a b^5 d^2 x^7 + 3.7518606875e-22B a^2 b^4 d^2 x^7 + 2.814047375e-22B a b^5 d^2 x^7 + 2.1138590625e-22B a^2 b^4 d^2 x^7 + 1.585730075e-22B a b^5 d^2 x^7 + 1.189651275e-22B a^2 b^4 d^2 x^7 + 8.9126581875e-23B a b^5 d^2 x^7 + 6.6822686875e-23B a^2 b^4 d^2 x^7 + 5.01205646875e-23B a b^5 d^2 x^7 + 3.7519466875e-23B a^2 b^4 d^2 x^7 + 2.81413306875e-23B a b^5 d^2 x^7 + 2.1139440625e-23B a^2 b^4 d^2 x^7 + 1.585814475e-23B a b^5 d^2 x^7 + 1.189736675e-23B a^2 b^4 d^2 x^7 + 8.9127521875e-24B a b^5 d^2 x^7 + 6.6823546875e-24B a^2 b^4 d^2 x^7 + 5.01214246875e-24B a b^5 d^2 x^7 + 3.7520326875e-24B a^2 b^4 d^2 x^7 + 2.8142186875e-24B a b^5 d^2 x^7 + 2.1140290625e-24B a^2 b^4 d^2 x^7 + 1.585898875e-24B a b^5 d^2 x^7 + 1.189822075e-24B a^2 b^4 d^2 x^7 + 8.9128461875e-25B a b^5 d^2 x^7 + 6.6824406875e-25B a^2 b^4 d^2 x^7 + 5.01222846875e-25B a b^5 d^2 x^7 + 3.7521186875e-25B a^2 b^4 d^2 x^7 + 2.814304375e-25B a b^5 d^2 x^7 + 2.1141140625e-25B a^2 b^4 d^2 x^7 + 1.585983275e-25B a b^5 d^2 x^7 + 1.18990746875e-25B a^2 b^4 d^2 x^7 + 8.9129401875e-26B a b^5 d^2 x^7 + 6.6825266875e-26B a^2 b^4 d^2 x^7 + 5.01231446875e-26B a b^5 d^2 x^7 + 3.7522046875e-26B a^2 b^4 d^2 x^7 + 2.81439006875e-26B a b^5 d^2 x^7 + 2.1141990625e-26B a^2 b^4 d^2 x^7 + 1.586067675e-26B a b^5 d^2 x^7 + 1.189992875e-26B a^2 b^4 d^2 x^7 + 8.9130341875e-27B a b^5 d^2 x^7 + 6.6826126875e-27B a^2 b^4 d^2 x^7 + 5.01240046875e-27B a b^5 d^2 x^7 + 3.7522906875e-27B a^2 b^4 d^2 x^7 + 2.8144756875e-27B a b^5 d^2 x^7 + 2.1142840625e-27B a^2 b^4 d^2 x^7 + 1.586152075e-27B a b^5 d^2 x^7 + 1.190078275e-27B a^2 b^4 d^2 x^7 + 8.9131281875e-28B a b^5 d^2 x^7 + 6.6826986875e-28B a^2 b^4 d^2 x^7 + 5.01248646875e-28B a b^5 d^2 x^7 + 3.7523766875e-28B a^2 b^4 d^2 x^7 + 2.814561375e-28B a b^5 d^2 x^7 + 2.1143690625e-28B a^2 b^4 d^2 x^7 + 1.586236475e-28B a b^5 d^2 x^7 + 1.190163675e-28B a^2 b^4 d^2 x^7 + 8.9132221875e-29B a b^5 d^2 x^7 + 6.6827846875e-29B a^2 b^4 d^2 x^7 + 5.01257246875e-29B a b^5 d^2 x^7 + 3.7524626875e-29B a^2 b^4 d^2 x^7 + 2.81464706875e-29B a b^5 d^2 x^7 + 2.1144540625e-29B a^2 b^4 d^2 x^7 + 1.586320875e-29B a b^5 d^2 x^7 + 1.190249075e-29B a^2 b^4 d^2 x^7 + 8.9133161875e-30B a b^5 d^2 x^7 + 6.6828706875e-30B a^2 b^4 d^2 x^7 + 5.01265846875e-30B a b^5 d^2 x^7 + 3.7525486875e-30B a^2 b^4 d^2 x^7 + 2.8147326875e-30B a b^5 d^2 x^7 + 2.1145390625e-30B a^2 b^4 d^2 x^7 + 1.586405275e-30B a b^5 d^2 x^7 + 1.19033446875e-30B a^2 b^4 d^2 x^7 + 8.9134101875e-31B a b^5 d^2 x^7 + 6.6829566875e-31B a^2 b^4 d^2 x^7 + 5.01274446875e-31B a b^5 d^2 x^7 + 3.7526346875e-31B a^2 b^4 d^2 x^7 + 2.814818375e-31B a b^5 d^2 x^7 + 2.1146240625e-31B a^2 b^4 d^2 x^7 + 1.586489675e-31B a b^5 d^2 x^7 + 1.190419875e-31B a^2 b^4 d^2 x^7 + 8.9135041875e-32B a b^5 d^2 x^7 + 6.6830426875e-32B a^2 b^4 d^2 x^7 + 5.01283046875e-32B a b^5 d^2 x^7 + 3.7527206875e-32B a^2 b^4 d^2 x^7 + 2.81490406875e-32B a b^5 d^2 x^7 + 2.1147090625e-32B a^2 b^4 d^2 x^7 + 1.586574075e-32B a b^5 d^2 x^7 + 1.190505275e-32B a^2 b^4 d^2 x^7 + 8.9135981875e-33B a b^5 d^2 x^7 + 6.6831286875e-33B a^2 b^4 d^2 x^7 + 5.01291646875e-33B a b^5 d^2 x^7 + 3.7528066875e-33B a^2 b^4 d^2 x^7 + 2.8149896875e-33B a b^5 d^2 x^7 + 2.1147940625e-33B a^2 b^4 d^2 x^7 + 1.586658475e-33B a b^5 d^2 x^7 + 1.190590675e-33B a^2 b^4 d^2 x^7 + 8.9136921875e-34B a b^5 d^2 x^7 + 6.6832146875e-34B a^2 b^4 d^2 x^7 + 5.01300246875e-34B a b^5 d^2 x^7 + 3.7528926875e-34B a^2 b^4 d^2 x^7 + 2.815075375e-34B a b^5 d^2 x^7 + 2.1148790625e-34B a^2 b^4 d^2 x^7 + 1.586742875e-34B a b^5 d^2 x^7 + 1.190676075e-34B a^2 b^4 d^2 x^7 + 8.9137861875e-35B a b^5 d^2 x^7 + 6.6833006875e-35B a^2 b^4 d^2 x^7 + 5.01308846875e-35B a b^5 d^2 x^7 + 3.7529786875e-35B a^2 b^4 d^2 x^7 + 2.81516106875e-35B a b^5 d^2 x^7 + 2.1149640625e-35B a^2 b^4 d^2 x^7 + 1.586827275e-35B a b^5 d^2 x^7 + 1.19076146875e-35B a^2 b^4 d^2 x^7 + 8.9138801875e-36B a b^5 d^2 x^7 + 6.6833866875e-36B a^2 b^4 d^2 x^7 + 5.01317446875e-36B a b^5 d^2 x^7 + 3.7530646875e-36B a^2 b^4 d^2 x^7 + 2.8152466875e-36B a b^5 d^2 x^7 + 2.1150490625e-36B a^2 b^4 d^2 x^7 + 1.586911675e-36B a b^5 d^2 x^7 + 1.190846875e-36B a^2 b^4 d^2 x^7 + 8.9139741875e-37B a b^5 d^2 x^7 + 6.6834726875e-37B a^2 b^4 d^2 x^7 + 5.01326046875e-37B a b^5 d^2 x^7 + 3.7531506875e-37B a^2 b^4 d^2 x^7 + 2.815332375e-37B a b^5 d^2 x^7 + 2.1151340625e-37B a^2 b^4 d^2 x^7 + 1.587006075e-37B a b^5 d^2 x^7 + 1.190932275e-37B a^2 b^4 d^2 x^7 + 8.9140681875e-38B a b^5 d^2 x^7 + 6.6835586875e-38B a^2 b^4 d^2 x^7 + 5.01334646875e-38B a b^5 d^2 x^7 + 3.7532366875e-38B a^2 b^4 d^2 x^7 + 2.81541806875e-38B a b^5 d^2 x^7 + 2.1152190625e-38B a^2 b^4 d^2 x^7 + 1.587090475e-38B a b^5 d^2 x^7 + 1.191017675e-38B a^2 b^4 d^2 x^7 + 8.9141621875e-39B a b^5 d^2 x^7 + 6.6836446875e-39B a^2 b^4 d^2 x^7 + 5.01343246875e-39B a b^5 d^2 x^7 + 3.7533226875e-39B a^2 b^4 d^2 x^7 + 2.8155036875e-39B a b^5 d^2 x^7 + 2.1153040625e-39B a^2 b^4 d^2 x^7 + 1.587174875e-39B a b^5 d^2 x^7 + 1.191103075e-39B a^2 b^4 d^2 x^7 + 8.9142561875e-40B a b^5 d^2 x^7 + 6.6837306875e-40B a^2 b^4 d^2 x^7 + 5.01351846875e-40B a b^5 d^2 x^7 + 3.7534086875e-40B a^2 b^4 d^2 x^7 + 2.815589375e-40B a b^5 d^2 x^7 + 2.1153890625e-40B a^2 b^4 d^2 x^7 + 1.587259275e-40B a b^5 d^2 x^7 + 1.19118846875e-40B a^2 b^4 d^2 x^7 + 8.9143501875e-41B a b^5 d^2 x^7 + 6.6838166875e-41B a^2 b^4 d^2 x^7 + 5.01360446875e-41B a b^5 d^2 x^7 + 3.7534946875e-41B a^2 b^4 d^2 x^7 + 2.81567506875e-41B a b^5 d^2 x^7 + 2.1154740625e-41B a^2 b^4 d^2 x^7 + 1.587343675e-41B a b^5 d^2 x^7 + 1.191273875e-41B a^2 b^4 d^2 x^7 + 8.9144441875e-42B a b^5 d^2 x^7 + 6.6839026875e-42B a^2 b^4 d^2 x^7 + 5.01369046875e-42B a b^5 d^2 x^7 + 3.7535806875e-42B a^2 b^4 d^2 x^7 + 2.8157606875e-42B a b^5 d^2 x^7 + 2.1155590625e-42B a^2 b^4 d^2 x^7 + 1.587428075e-42B a b^5 d^2 x^7 + 1.191359275e-42B a^2 b^4 d^2 x^7 + 8.9145381875e-43B a b^5 d^2 x^7 + 6.6839886875e-43B a^2 b^4 d^2 x^7 + 5.01377646875e-43B a b^5 d^2 x^7 + 3.7536666875e-43B a^2 b^4 d^2 x^7 + 2.815846375e-43B a b^5 d^2 x^7 + 2.1156440625e-43B a^2 b^4 d^2 x^7 + 1.587512475e-43B a b^5 d^2 x^7 + 1.191444675e-43B a^2 b^4 d^2 x^7 + 8.9146321875e-44B a b^5 d^2 x^7 + 6.6840746875e-44B a^2 b^4 d^2 x^7 + 5.01386246875e-44B a b^5 d^2 x^7 + 3.7537526875e-44B a^2 b^4 d^2 x^7 + 2.81593206875e-44B a b^5 d^2 x^7 + 2.1157290625e-44B a^2 b^4 d^2 x^7 + 1.587596875e-44B a b^5 d^2 x^7 + 1.191530075e-44B a^2 b^4 d^2 x^7 + 8.9147261875e-45B a b^5 d^2 x^7 + 6.6841606875e-45B a^2 b^4 d^2 x^7 + 5.01394846875e-45B a b^5 d^2 x^7 + 3.7538386875e-45B a^2 b^4 d^2 x^7 + 2.8160176875e-45B a b^5 d^2 x^7 + 2.1158140625e-45B a^2 b^4 d^2 x^7 + 1.587681275e-45B a b^5 d^2 x^7 + 1.19161546875e-45B a^2 b^4 d^2 x^7 + 8.9148201$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 476 vs. $2(110) = 220$.

Time = 0.07 (sec) , antiderivative size = 476, normalized size of antiderivative = 4.03

$$\int (a + bx)^6 (A + Bx)(d + ex)^2 dx$$

$$= \frac{1}{10} Bb^6 e^2 x^{10} + Aa^6 d^2 x + \frac{1}{9} (2 Bb^6 de + (6 Bab^5 + Ab^6) e^2) x^9$$

$$+ \frac{1}{8} (Bb^6 d^2 + 2 (6 Bab^5 + Ab^6) de + 3 (5 Ba^2 b^4 + 2 Aab^5) e^2) x^8$$

$$+ \frac{1}{7} ((6 Bab^5 + Ab^6) d^2 + 6 (5 Ba^2 b^4 + 2 Aab^5) de + 5 (4 Ba^3 b^3 + 3 Aa^2 b^4) e^2) x^7$$

$$+ \frac{1}{6} (3 (5 Ba^2 b^4 + 2 Aab^5) d^2 + 10 (4 Ba^3 b^3 + 3 Aa^2 b^4) de + 5 (3 Ba^4 b^2 + 4 Aa^3 b^3) e^2) x^6$$

$$+ \frac{1}{5} (5 (4 Ba^3 b^3 + 3 Aa^2 b^4) d^2 + 10 (3 Ba^4 b^2 + 4 Aa^3 b^3) de + 3 (2 Ba^5 b + 5 Aa^4 b^2) e^2) x^5$$

$$+ \frac{1}{4} (5 (3 Ba^4 b^2 + 4 Aa^3 b^3) d^2 + 6 (2 Ba^5 b + 5 Aa^4 b^2) de + (Ba^6 + 6 Aa^5 b) e^2) x^4$$

$$+ \frac{1}{3} (Aa^6 e^2 + 3 (2 Ba^5 b + 5 Aa^4 b^2) d^2 + 2 (Ba^6 + 6 Aa^5 b) de) x^3$$

$$+ \frac{1}{2} (2 Aa^6 de + (Ba^6 + 6 Aa^5 b) d^2) x^2$$

input `integrate((b*x+a)^6*(B*x+A)*(e*x+d)^2,x, algorithm="fricas")`

output `1/10*B*b^6*e^2*x^10 + A*a^6*d^2*x + 1/9*(2*B*b^6*d*e + (6*B*a*b^5 + A*b^6)*e^2)*x^9 + 1/8*(B*b^6*d^2 + 2*(6*B*a*b^5 + A*b^6)*d*e + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*e^2)*x^8 + 1/7*((6*B*a*b^5 + A*b^6)*d^2 + 6*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e + 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^2)*x^7 + 1/6*(3*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2 + 10*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^2)*x^6 + 1/5*(5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2 + 10*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e + 3*(2*B*a^5*b + 5*A*a^4*b^2)*e^2)*x^5 + 1/4*(5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2 + 6*(2*B*a^5*b + 5*A*a^4*b^2)*d*e + (B*a^6 + 6*A*a^5*b)*e^2)*x^4 + 1/3*(A*a^6*e^2 + 3*(2*B*a^5*b + 5*A*a^4*b^2)*d^2 + 2*(B*a^6 + 6*A*a^5*b)*d*e)*x^3 + 1/2*(2*A*a^6*d*e + (B*a^6 + 6*A*a^5*b)*d^2)*x^2`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 568 vs. $2(116) = 232$.

Time = 0.05 (sec) , antiderivative size = 568, normalized size of antiderivative = 4.81

$$\begin{aligned}
 \int (a + bx)^6 (A + Bx)(d + ex)^2 dx = & Aa^6 d^2 x + \frac{Bb^6 e^2 x^{10}}{10} \\
 & + x^9 \left(\frac{Ab^6 e^2}{9} + \frac{2Bab^5 e^2}{3} + \frac{2Bb^6 de}{9} \right) + x^8 \\
 & \cdot \left(\frac{3Aab^5 e^2}{4} + \frac{Ab^6 de}{4} + \frac{15Ba^2 b^4 e^2}{8} + \frac{3Bab^5 de}{2} \right. \\
 & \quad \left. + \frac{Bb^6 d^2}{8} \right) + x^7 \cdot \left(\frac{15Aa^2 b^4 e^2}{7} + \frac{12Aab^5 de}{7} \right. \\
 & \quad \left. + \frac{Ab^6 d^2}{7} + \frac{20Ba^3 b^3 e^2}{7} + \frac{30Ba^2 b^4 de}{7} + \frac{6Bab^5 d^2}{7} \right) \\
 & + x^6 \cdot \left(\frac{10Aa^3 b^3 e^2}{3} + 5Aa^2 b^4 de + Aab^5 d^2 \right. \\
 & \quad \left. + \frac{5Ba^4 b^2 e^2}{2} + \frac{20Ba^3 b^3 de}{3} + \frac{5Ba^2 b^4 d^2}{2} \right) + x^5 \\
 & \cdot \left(3Aa^4 b^2 e^2 + 8Aa^3 b^3 de + 3Aa^2 b^4 d^2 + \frac{6Ba^5 b e^2}{5} \right. \\
 & \quad \left. + 6Ba^4 b^2 de + 4Ba^3 b^3 d^2 \right) + x^4 \\
 & \cdot \left(\frac{3Aa^5 b e^2}{2} + \frac{15Aa^4 b^2 de}{2} + 5Aa^3 b^3 d^2 + \frac{Ba^6 e^2}{4} \right. \\
 & \quad \left. + 3Ba^5 b de + \frac{15Ba^4 b^2 d^2}{4} \right) \\
 & + x^3 \left(\frac{Aa^6 e^2}{3} + 4Aa^5 b de + 5Aa^4 b^2 d^2 + \frac{2Ba^6 de}{3} \right. \\
 & \quad \left. + 2Ba^5 b d^2 \right) + x^2 \left(Aa^6 de + 3Aa^5 b d^2 + \frac{Ba^6 d^2}{2} \right)
 \end{aligned}$$

input `integrate((b*x+a)**6*(B*x+A)*(e*x+d)**2,x)`

output

```
A*a**6*d**2*x + B*b**6*e**2*x**10/10 + x**9*(A*b**6*e**2/9 + 2*B*a*b**5*e*
*2/3 + 2*B*b**6*d*e/9) + x**8*(3*A*a*b**5*e**2/4 + A*b**6*d*e/4 + 15*B*a**
2*b**4*e**2/8 + 3*B*a*b**5*d*e/2 + B*b**6*d**2/8) + x**7*(15*A*a**2*b**4*e
**2/7 + 12*A*a*b**5*d*e/7 + A*b**6*d**2/7 + 20*B*a**3*b**3*e**2/7 + 30*B*a
**2*b**4*d*e/7 + 6*B*a*b**5*d**2/7) + x**6*(10*A*a**3*b**3*e**2/3 + 5*A*a*
*2*b**4*d*e + A*a*b**5*d**2 + 5*B*a**4*b**2*e**2/2 + 20*B*a**3*b**3*d*e/3
+ 5*B*a**2*b**4*d**2/2) + x**5*(3*A*a**4*b**2*e**2 + 8*A*a**3*b**3*d*e + 3
*A*a**2*b**4*d**2 + 6*B*a**5*b*e**2/5 + 6*B*a**4*b**2*d*e + 4*B*a**3*b**3*
d**2) + x**4*(3*A*a**5*b*e**2/2 + 15*A*a**4*b**2*d*e/2 + 5*A*a**3*b**3*d**
2 + B*a**6*e**2/4 + 3*B*a**5*b*d*e + 15*B*a**4*b**2*d**2/4) + x**3*(A*a**6
*e**2/3 + 4*A*a**5*b*d*e + 5*A*a**4*b**2*d**2 + 2*B*a**6*d*e/3 + 2*B*a**5*
b*d**2) + x**2*(A*a**6*d*e + 3*A*a**5*b*d**2 + B*a**6*d**2/2)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 476 vs. $2(110) = 220$.

Time = 0.04 (sec) , antiderivative size = 476, normalized size of antiderivative = 4.03

$$\begin{aligned}
 & \int (a + bx)^6 (A + Bx)(d + ex)^2 dx \\
 &= \frac{1}{10} Bb^6 e^2 x^{10} + Aa^6 d^2 x + \frac{1}{9} (2 Bb^6 de + (6 Bab^5 + Ab^6) e^2) x^9 \\
 &+ \frac{1}{8} (Bb^6 d^2 + 2 (6 Bab^5 + Ab^6) de + 3 (5 Ba^2 b^4 + 2 Aab^5) e^2) x^8 \\
 &+ \frac{1}{7} ((6 Bab^5 + Ab^6) d^2 + 6 (5 Ba^2 b^4 + 2 Aab^5) de + 5 (4 Ba^3 b^3 + 3 Aa^2 b^4) e^2) x^7 \\
 &+ \frac{1}{6} (3 (5 Ba^2 b^4 + 2 Aab^5) d^2 + 10 (4 Ba^3 b^3 + 3 Aa^2 b^4) de + 5 (3 Ba^4 b^2 + 4 Aa^3 b^3) e^2) x^6 \\
 &+ \frac{1}{5} (5 (4 Ba^3 b^3 + 3 Aa^2 b^4) d^2 + 10 (3 Ba^4 b^2 + 4 Aa^3 b^3) de + 3 (2 Ba^5 b + 5 Aa^4 b^2) e^2) x^5 \\
 &+ \frac{1}{4} (5 (3 Ba^4 b^2 + 4 Aa^3 b^3) d^2 + 6 (2 Ba^5 b + 5 Aa^4 b^2) de + (Ba^6 + 6 Aa^5 b) e^2) x^4 \\
 &+ \frac{1}{3} (Aa^6 e^2 + 3 (2 Ba^5 b + 5 Aa^4 b^2) d^2 + 2 (Ba^6 + 6 Aa^5 b) de) x^3 \\
 &+ \frac{1}{2} (2 Aa^6 de + (Ba^6 + 6 Aa^5 b) d^2) x^2
 \end{aligned}$$

input

```
integrate((b*x+a)^6*(B*x+A)*(e*x+d)^2,x, algorithm="maxima")
```

output

```

1/10*B*b^6*e^2*x^10 + A*a^6*d^2*x + 1/9*(2*B*b^6*d*e + (6*B*a*b^5 + A*b^6)
*e^2)*x^9 + 1/8*(B*b^6*d^2 + 2*(6*B*a*b^5 + A*b^6)*d*e + 3*(5*B*a^2*b^4 +
2*A*a*b^5)*e^2)*x^8 + 1/7*((6*B*a*b^5 + A*b^6)*d^2 + 6*(5*B*a^2*b^4 + 2*A*
a*b^5)*d*e + 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^2)*x^7 + 1/6*(3*(5*B*a^2*b^4
+ 2*A*a*b^5)*d^2 + 10*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e + 5*(3*B*a^4*b^2 + 4
*A*a^3*b^3)*e^2)*x^6 + 1/5*(5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2 + 10*(3*B*a^
4*b^2 + 4*A*a^3*b^3)*d*e + 3*(2*B*a^5*b + 5*A*a^4*b^2)*e^2)*x^5 + 1/4*(5*(
3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2 + 6*(2*B*a^5*b + 5*A*a^4*b^2)*d*e + (B*a^6
+ 6*A*a^5*b)*e^2)*x^4 + 1/3*(A*a^6*e^2 + 3*(2*B*a^5*b + 5*A*a^4*b^2)*d^2 +
2*(B*a^6 + 6*A*a^5*b)*d*e)*x^3 + 1/2*(2*A*a^6*d*e + (B*a^6 + 6*A*a^5*b)*d
^2)*x^2

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 552 vs. $2(110) = 220$.

Time = 0.12 (sec) , antiderivative size = 552, normalized size of antiderivative = 4.68

$$\begin{aligned}
 \int (a + bx)^6 (A + Bx)(d + ex)^2 dx = & \frac{1}{10} Bb^6 e^2 x^{10} + \frac{2}{9} Bb^6 dex^9 + \frac{2}{3} Bab^5 e^2 x^9 \\
 & + \frac{1}{9} Ab^6 e^2 x^9 + \frac{1}{8} Bb^6 d^2 x^8 + \frac{3}{2} Bab^5 dex^8 \\
 & + \frac{1}{4} Ab^6 dex^8 + \frac{15}{8} Ba^2 b^4 e^2 x^8 + \frac{3}{4} Aab^5 e^2 x^8 \\
 & + \frac{6}{7} Bab^5 d^2 x^7 + \frac{1}{7} Ab^6 d^2 x^7 + \frac{30}{7} Ba^2 b^4 dex^7 \\
 & + \frac{12}{7} Aab^5 dex^7 + \frac{20}{7} Ba^3 b^3 e^2 x^7 + \frac{15}{7} Aa^2 b^4 e^2 x^7 \\
 & + \frac{5}{2} Ba^2 b^4 d^2 x^6 + Aab^5 d^2 x^6 + \frac{20}{3} Ba^3 b^3 dex^6 \\
 & + 5 Aa^2 b^4 dex^6 + \frac{5}{2} Ba^4 b^2 e^2 x^6 + \frac{10}{3} Aa^3 b^3 e^2 x^6 \\
 & + 4 Ba^3 b^3 d^2 x^5 + 3 Aa^2 b^4 d^2 x^5 + 6 Ba^4 b^2 dex^5 \\
 & + 8 Aa^3 b^3 dex^5 + \frac{6}{5} Ba^5 b e^2 x^5 + 3 Aa^4 b^2 e^2 x^5 \\
 & + \frac{15}{4} Ba^4 b^2 d^2 x^4 + 5 Aa^3 b^3 d^2 x^4 + 3 Ba^5 b dex^4 \\
 & + \frac{15}{2} Aa^4 b^2 dex^4 + \frac{1}{4} Ba^6 e^2 x^4 + \frac{3}{2} Aa^5 b e^2 x^4 \\
 & + 2 Ba^5 b d^2 x^3 + 5 Aa^4 b^2 d^2 x^3 + \frac{2}{3} Ba^6 dex^3 \\
 & + 4 Aa^5 b dex^3 + \frac{1}{3} Aa^6 e^2 x^3 + \frac{1}{2} Ba^6 d^2 x^2 \\
 & + 3 Aa^5 b d^2 x^2 + Aa^6 dex^2 + Aa^6 d^2 x
 \end{aligned}$$

input `integrate((b*x+a)^6*(B*x+A)*(e*x+d)^2,x, algorithm="giac")`

output

$$\begin{aligned}
& 1/10*B*b^6*e^2*x^10 + 2/9*B*b^6*d*e*x^9 + 2/3*B*a*b^5*e^2*x^9 + 1/9*A*b^6* \\
& e^2*x^9 + 1/8*B*b^6*d^2*x^8 + 3/2*B*a*b^5*d*e*x^8 + 1/4*A*b^6*d*e*x^8 + 15 \\
& /8*B*a^2*b^4*e^2*x^8 + 3/4*A*a*b^5*e^2*x^8 + 6/7*B*a*b^5*d^2*x^7 + 1/7*A*b \\
& ^6*d^2*x^7 + 30/7*B*a^2*b^4*d*e*x^7 + 12/7*A*a*b^5*d*e*x^7 + 20/7*B*a^3*b^ \\
& 3*e^2*x^7 + 15/7*A*a^2*b^4*e^2*x^7 + 5/2*B*a^2*b^4*d^2*x^6 + A*a*b^5*d^2*x \\
& ^6 + 20/3*B*a^3*b^3*d*e*x^6 + 5*A*a^2*b^4*d*e*x^6 + 5/2*B*a^4*b^2*e^2*x^6 \\
& + 10/3*A*a^3*b^3*e^2*x^6 + 4*B*a^3*b^3*d^2*x^5 + 3*A*a^2*b^4*d^2*x^5 + 6*B \\
& *a^4*b^2*d*e*x^5 + 8*A*a^3*b^3*d*e*x^5 + 6/5*B*a^5*b*e^2*x^5 + 3*A*a^4*b^2 \\
& *e^2*x^5 + 15/4*B*a^4*b^2*d^2*x^4 + 5*A*a^3*b^3*d^2*x^4 + 3*B*a^5*b*d*e*x^ \\
& 4 + 15/2*A*a^4*b^2*d*e*x^4 + 1/4*B*a^6*e^2*x^4 + 3/2*A*a^5*b*e^2*x^4 + 2*B \\
& *a^5*b*d^2*x^3 + 5*A*a^4*b^2*d^2*x^3 + 2/3*B*a^6*d*e*x^3 + 4*A*a^5*b*d*e*x \\
& ^3 + 1/3*A*a^6*e^2*x^3 + 1/2*B*a^6*d^2*x^2 + 3*A*a^5*b*d^2*x^2 + A*a^6*d*e \\
& *x^2 + A*a^6*d^2*x
\end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 457, normalized size of antiderivative = 3.87

$$\begin{aligned}
\int (a + bx)^6 (A + Bx)(d + ex)^2 dx = & x^4 \left(\frac{B a^6 e^2}{4} + 3 B a^5 b d e + \frac{3 A a^5 b e^2}{2} \right. \\
& \left. + \frac{15 B a^4 b^2 d^2}{4} + \frac{15 A a^4 b^2 d e}{2} + 5 A a^3 b^3 d^2 \right) \\
& + x^7 \left(\frac{20 B a^3 b^3 e^2}{7} + \frac{30 B a^2 b^4 d e}{7} + \frac{15 A a^2 b^4 e^2}{7} \right. \\
& \left. + \frac{6 B a b^5 d^2}{7} + \frac{12 A a b^5 d e}{7} + \frac{A b^6 d^2}{7} \right) \\
& + x^5 \left(\frac{6 B a^5 b e^2}{5} + 6 B a^4 b^2 d e + 3 A a^4 b^2 e^2 \right. \\
& \left. + 4 B a^3 b^3 d^2 + 8 A a^3 b^3 d e + 3 A a^2 b^4 d^2 \right) \\
& + x^6 \left(\frac{5 B a^4 b^2 e^2}{2} + \frac{20 B a^3 b^3 d e}{3} + \frac{10 A a^3 b^3 e^2}{3} \right. \\
& \left. + \frac{5 B a^2 b^4 d^2}{2} + 5 A a^2 b^4 d e + A a b^5 d^2 \right) \\
& + x^3 \left(\frac{2 B a^6 d e}{3} + \frac{A a^6 e^2}{3} + 2 B a^5 b d^2 + 4 A a^5 b d e \right. \\
& \left. + 5 A a^4 b^2 d^2 \right) + x^8 \left(\frac{15 B a^2 b^4 e^2}{8} + \frac{3 B a b^5 d e}{2} \right. \\
& \left. + \frac{3 A a b^5 e^2}{4} + \frac{B b^6 d^2}{8} + \frac{A b^6 d e}{4} \right) \\
& + A a^6 d^2 x + \frac{a^5 d x^2 (2 A a e + 6 A b d + B a d)}{2} \\
& + \frac{b^5 e x^9 (A b e + 6 B a e + 2 B b d)}{9} + \frac{B b^6 e^2 x^{10}}{10}
\end{aligned}$$

input

```
int((A + B*x)*(a + b*x)^6*(d + e*x)^2,x)
```

output

```
x^4*((B*a^6*e^2)/4 + (3*A*a^5*b*e^2)/2 + 5*A*a^3*b^3*d^2 + (15*B*a^4*b^2*d^2)/4 + 3*B*a^5*b*d*e + (15*A*a^4*b^2*d*e)/2) + x^7*((A*b^6*d^2)/7 + (6*B*a*b^5*d^2)/7 + (15*A*a^2*b^4*e^2)/7 + (20*B*a^3*b^3*e^2)/7 + (12*A*a*b^5*d*e)/7 + (30*B*a^2*b^4*d*e)/7) + x^5*((6*B*a^5*b*e^2)/5 + 3*A*a^2*b^4*d^2 + 3*A*a^4*b^2*e^2 + 4*B*a^3*b^3*d^2 + 8*A*a^3*b^3*d*e + 6*B*a^4*b^2*d*e) + x^6*(A*a*b^5*d^2 + (10*A*a^3*b^3*e^2)/3 + (5*B*a^2*b^4*d^2)/2 + (5*B*a^4*b^2*e^2)/2 + 5*A*a^2*b^4*d*e + (20*B*a^3*b^3*d*e)/3) + x^3*((A*a^6*e^2)/3 + (2*B*a^6*d*e)/3 + 2*B*a^5*b*d^2 + 5*A*a^4*b^2*d^2 + 4*A*a^5*b*d*e) + x^8*((B*b^6*d^2)/8 + (A*b^6*d*e)/4 + (3*A*a*b^5*e^2)/4 + (15*B*a^2*b^4*e^2)/8 + (3*B*a*b^5*d*e)/2) + A*a^6*d^2*x + (a^5*d*x^2*(2*A*a*e + 6*A*b*d + B*a*d))/2 + (b^5*e*x^9*(A*b*e + 6*B*a*e + 2*B*b*d))/9 + (B*b^6*e^2*x^10)/10
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 295, normalized size of antiderivative = 2.50

$$\int (a + bx)^6 (A + Bx)(d + ex)^2 dx$$

$$= \frac{x(36b^7e^2x^9 + 280ab^6e^2x^8 + 80b^7dex^8 + 945a^2b^5e^2x^7 + 630ab^6dex^7 + 45b^7d^2x^7 + 1800a^3b^4e^2x^6 + 2160a^2b^5dex^6 + 1260ab^6d^2x^5 + 360a^2b^5d^2x^5 + 2160a^2b^5d^2ex^6 + 945a^2b^5e^2x^7 + 360ab^6d^2x^6 + 630ab^6dex^7 + 280ab^6e^2x^8 + 45b^7d^2x^7 + 80b^7dex^8 + 36b^7e^2x^9)}{360}$$

input

```
int((b*x+a)^6*(B*x+A)*(e*x+d)^2,x)
```

output

```
(x*(360*a**7*d**2 + 360*a**7*d*e*x + 120*a**7*e**2*x**2 + 1260*a**6*b*d**2*x + 1680*a**6*b*d*e*x**2 + 630*a**6*b*e**2*x**3 + 2520*a**5*b**2*d**2*x**2 + 3780*a**5*b**2*d*e*x**3 + 1512*a**5*b**2*e**2*x**4 + 3150*a**4*b**3*d**2*x**3 + 5040*a**4*b**3*d*e*x**4 + 2100*a**4*b**3*e**2*x**5 + 2520*a**3*b**4*d**2*x**4 + 4200*a**3*b**4*d*e*x**5 + 1800*a**3*b**4*e**2*x**6 + 1260*a**2*b**5*d**2*x**5 + 2160*a**2*b**5*d*e*x**6 + 945*a**2*b**5*e**2*x**7 + 360*a*b**6*d**2*x**6 + 630*a*b**6*d*e*x**7 + 280*a*b**6*e**2*x**8 + 45*b**7*d**2*x**7 + 80*b**7*d*e*x**8 + 36*b**7*e**2*x**9))/360
```

3.48 $\int (a + bx)^6 (A + Bx)(d + ex) dx$

Optimal result	488
Mathematica [B] (verified)	488
Rubi [A] (verified)	489
Maple [B] (verified)	490
Fricas [B] (verification not implemented)	491
Sympy [B] (verification not implemented)	491
Maxima [B] (verification not implemented)	493
Giac [B] (verification not implemented)	493
Mupad [B] (verification not implemented)	495
Reduce [B] (verification not implemented)	495

Optimal result

Integrand size = 18, antiderivative size = 75

$$\int (a + bx)^6 (A + Bx)(d + ex) dx = \frac{(Ab - aB)(bd - ae)(a + bx)^7}{7b^3} + \frac{(bBd + Abe - 2aBe)(a + bx)^8}{8b^3} + \frac{Be(a + bx)^9}{9b^3}$$

output

```
1/7*(A*b-B*a)*(-a*e+b*d)*(b*x+a)^7/b^3+1/8*(A*b*e-2*B*a*e+B*b*d)*(b*x+a)^8/b^3+1/9*B*e*(b*x+a)^9/b^3
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 231 vs. 2(75) = 150.

Time = 0.09 (sec) , antiderivative size = 231, normalized size of antiderivative = 3.08

$$\int (a + bx)^6 (A + Bx)(d + ex) dx = \frac{1}{504}x(84a^6(3A(2d + ex) + Bx(3d + 2ex)) + 126a^4b^2x^2(5A(4d + 3ex) + 3Bx(5d + 4ex)) + 252a^5bx(Bx(4d + 3ex) + A(6d + 4ex)) + 168a^3b^3x^3(3A(5d + 4ex) + 2Bx(6d + 5ex)) + 36a^2b^4x^4(7A(6d + 5ex) + 5Bx(7d + 6ex)) + 18ab^5x^5(4A(7d + 6ex) + 3Bx(8d + 7ex)) + b^6x^6(9A(8d + 7ex) + 7Bx(9d + 8ex)))$$

input `Integrate[(a + b*x)^6*(A + B*x)*(d + e*x), x]`

output $(x*(84*a^6*(3*A*(2*d + e*x) + B*x*(3*d + 2*e*x)) + 126*a^4*b^2*x^2*(5*A*(4*d + 3*e*x) + 3*B*x*(5*d + 4*e*x)) + 252*a^5*b*x*(B*x*(4*d + 3*e*x) + A*(6*d + 4*e*x)) + 168*a^3*b^3*x^3*(3*A*(5*d + 4*e*x) + 2*B*x*(6*d + 5*e*x)) + 36*a^2*b^4*x^4*(7*A*(6*d + 5*e*x) + 5*B*x*(7*d + 6*e*x)) + 18*a*b^5*x^5*(4*A*(7*d + 6*e*x) + 3*B*x*(8*d + 7*e*x)) + b^6*x^6*(9*A*(8*d + 7*e*x) + 7*B*x*(9*d + 8*e*x)))/504$

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^6 (A + Bx)(d + ex) dx$$

$$\downarrow 86$$

$$\int \left(\frac{(a + bx)^7 (-2aBe + Abe + bBd)}{b^2} + \frac{(a + bx)^6 (Ab - aB)(bd - ae)}{b^2} + \frac{Be(a + bx)^8}{b^2} \right) dx$$

$$\downarrow 2009$$

$$\frac{(a + bx)^8 (-2aBe + Abe + bBd)}{8b^3} + \frac{(a + bx)^7 (Ab - aB)(bd - ae)}{7b^3} + \frac{Be(a + bx)^9}{9b^3}$$

input `Int[(a + b*x)^6*(A + B*x)*(d + e*x), x]`

output $((A*b - a*B)*(b*d - a*e)*(a + b*x)^7)/(7*b^3) + ((b*B*d + A*b*e - 2*a*B*e)*(a + b*x)^8)/(8*b^3) + (B*e*(a + b*x)^9)/(9*b^3)$

Definitions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 278 vs. $2(69) = 138$.

Time = 0.15 (sec) , antiderivative size = 279, normalized size of antiderivative = 3.72

method	result
norman	$\frac{b^6 B e x^9}{9} + \left(\frac{1}{8} A b^6 e + \frac{3}{4} B a b^5 e + \frac{1}{8} b^6 B d\right) x^8 + \left(\frac{6}{7} A a b^5 e + \frac{1}{7} A b^6 d + \frac{15}{7} B a^2 b^4 e + \frac{6}{7} B a b^5 d\right) x^7$
default	$\frac{b^6 B e x^9}{9} + \frac{((b^6 A + 6 a b^5 B) e + b^6 B d) x^8}{8} + \frac{((6 a b^5 A + 15 a^2 b^4 B) e + (b^6 A + 6 a b^5 B) d) x^7}{7} + \frac{((15 a^2 b^4 A + 20 a^3 b^3 B) e + (6 a b^5 A + 6 a^2 b^4 B) d) x^6}{6}$
orering	$x(56 b^6 B e x^8 + 63 A b^6 e x^7 + 378 B a b^5 e x^7 + 63 B b^6 d x^7 + 432 A a b^5 e x^6 + 72 A b^6 d x^6 + 1080 B a^2 b^4 e x^6 + 432 B a b^5 d x^6 + 1260 A a^2 b^4 e x^5 + 1260 A a b^5 d x^5 + 1260 B a^2 b^4 d x^5 + 1260 B a b^5 d x^5 + 1260 A a^2 b^4 e x^4 + 1260 A a b^5 d x^4 + 1260 B a^2 b^4 d x^4 + 1260 B a b^5 d x^4 + 1260 A a^2 b^4 e x^3 + 1260 A a b^5 d x^3 + 1260 B a^2 b^4 d x^3 + 1260 B a b^5 d x^3 + 1260 A a^2 b^4 e x^2 + 1260 A a b^5 d x^2 + 1260 B a^2 b^4 d x^2 + 1260 B a b^5 d x^2 + 1260 A a^2 b^4 e x + 1260 A a b^5 d x + 1260 B a^2 b^4 d x + 1260 B a b^5 d x + 1260 A a^2 b^4 e + 1260 A a b^5 d + 1260 B a^2 b^4 d + 1260 B a b^5 d)$
gosper	$\frac{1}{3} x^3 B a^6 e + \frac{1}{2} x^2 a^6 A e + \frac{15}{4} x^4 A a^4 b^2 e + 5 x^4 A a^3 b^3 d + \frac{1}{2} x^2 B a^6 d + \frac{1}{9} b^6 B e x^9 + a^6 A d x + \frac{10}{3} x^7$
risch	$\frac{1}{3} x^3 B a^6 e + \frac{1}{2} x^2 a^6 A e + \frac{15}{4} x^4 A a^4 b^2 e + 5 x^4 A a^3 b^3 d + \frac{1}{2} x^2 B a^6 d + \frac{1}{9} b^6 B e x^9 + a^6 A d x + \frac{10}{3} x^7$
parallelrisch	$\frac{1}{3} x^3 B a^6 e + \frac{1}{2} x^2 a^6 A e + \frac{15}{4} x^4 A a^4 b^2 e + 5 x^4 A a^3 b^3 d + \frac{1}{2} x^2 B a^6 d + \frac{1}{9} b^6 B e x^9 + a^6 A d x + \frac{10}{3} x^7$

input

```
int((b*x+a)^6*(B*x+A)*(e*x+d),x,method=_RETURNVERBOSE)
```

output

```
1/9*b^6*B*e*x^9+(1/8*A*b^6*e+3/4*B*a*b^5*e+1/8*b^6*B*d)*x^8+(6/7*A*a*b^5*e+1/7*A*b^6*d+15/7*B*a^2*b^4*e+6/7*B*a*b^5*d)*x^7+(5/2*A*a^2*b^4*e+A*a*b^5*d+10/3*B*a^3*b^3*e+5/2*B*a^2*b^4*d)*x^6+(4*A*a^3*b^3*e+3*A*a^2*b^4*d+3*B*a^4*b^2*e+4*B*a^3*b^3*d)*x^5+(15/4*A*a^4*b^2*e+5*A*a^3*b^3*d+3/2*B*a^5*b*e+15/4*B*a^4*b^2*d)*x^4+(2*A*a^5*b*e+5*A*a^4*b^2*d+1/3*B*a^6*e+2*B*a^5*b*d)*x^3+(1/2*a^6*A*e+3*A*a^5*b*d+1/2*B*a^6*d)*x^2+a^6*A*d*x
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 297 vs. $2(69) = 138$.

Time = 0.07 (sec) , antiderivative size = 297, normalized size of antiderivative = 3.96

$$\begin{aligned} & \int (a + bx)^6 (A + Bx)(d + ex) dx \\ &= \frac{1}{9} Bb^6 ex^9 + Aa^6 dx + \frac{1}{8} (Bb^6 d + (6 Bab^5 + Ab^6)e)x^8 \\ & \quad + \frac{1}{7} ((6 Bab^5 + Ab^6)d + 3(5 Ba^2 b^4 + 2 Aab^5)e)x^7 \\ & \quad + \frac{1}{6} (3(5 Ba^2 b^4 + 2 Aab^5)d + 5(4 Ba^3 b^3 + 3 Aa^2 b^4)e)x^6 \\ & \quad + ((4 Ba^3 b^3 + 3 Aa^2 b^4)d + (3 Ba^4 b^2 + 4 Aa^3 b^3)e)x^5 \\ & \quad + \frac{1}{4} (5(3 Ba^4 b^2 + 4 Aa^3 b^3)d + 3(2 Ba^5 b + 5 Aa^4 b^2)e)x^4 \\ & \quad + \frac{1}{3} (3(2 Ba^5 b + 5 Aa^4 b^2)d + (Ba^6 + 6 Aa^5 b)e)x^3 + \frac{1}{2} (Aa^6 e + (Ba^6 + 6 Aa^5 b)d)x^2 \end{aligned}$$

input `integrate((b*x+a)^6*(B*x+A)*(e*x+d),x, algorithm="fricas")`

output

```
1/9*B*b^6*e*x^9 + A*a^6*d*x + 1/8*(B*b^6*d + (6*B*a*b^5 + A*b^6)*e)*x^8 +
1/7*((6*B*a*b^5 + A*b^6)*d + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*e)*x^7 + 1/6*(3*(
5*B*a^2*b^4 + 2*A*a*b^5)*d + 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e)*x^6 + ((4*B*
a^3*b^3 + 3*A*a^2*b^4)*d + (3*B*a^4*b^2 + 4*A*a^3*b^3)*e)*x^5 + 1/4*(5*(3*
B*a^4*b^2 + 4*A*a^3*b^3)*d + 3*(2*B*a^5*b + 5*A*a^4*b^2)*e)*x^4 + 1/3*(3*(
2*B*a^5*b + 5*A*a^4*b^2)*d + (B*a^6 + 6*A*a^5*b)*e)*x^3 + 1/2*(A*a^6*e + (
B*a^6 + 6*A*a^5*b)*d)*x^2
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 333 vs. $2(71) = 142$.

Time = 0.04 (sec) , antiderivative size = 333, normalized size of antiderivative = 4.44

$$\int (a + bx)^6 (A + Bx)(d + ex) dx = Aa^6 dx + \frac{Bb^6 ex^9}{9} + x^8 \left(\frac{Ab^6 e}{8} + \frac{3Bab^5 e}{4} + \frac{Bb^6 d}{8} \right) + x^7 \cdot \left(\frac{6Aab^5 e}{7} + \frac{Ab^6 d}{7} + \frac{15Ba^2 b^4 e}{7} + \frac{6Bab^5 d}{7} \right) + x^6 \cdot \left(\frac{5Aa^2 b^4 e}{2} + Aab^5 d + \frac{10Ba^3 b^3 e}{3} + \frac{5Ba^2 b^4 d}{2} \right) + x^5 \cdot (4Aa^3 b^3 e + 3Aa^2 b^4 d + 3Ba^4 b^2 e + 4Ba^3 b^3 d) + x^4 \cdot \left(\frac{15Aa^4 b^2 e}{4} + 5Aa^3 b^3 d + \frac{3Ba^5 b e}{2} + \frac{15Ba^4 b^2 d}{4} \right) + x^3 \cdot \left(2Aa^5 b e + 5Aa^4 b^2 d + \frac{Ba^6 e}{3} + 2Ba^5 b d \right) + x^2 \left(\frac{Aa^6 e}{2} + 3Aa^5 b d + \frac{Ba^6 d}{2} \right)$$

input `integrate((b*x+a)**6*(B*x+A)*(e*x+d),x)`

output `A*a**6*d*x + B*b**6*e*x**9/9 + x**8*(A*b**6*e/8 + 3*B*a*b**5*e/4 + B*b**6*d/8) + x**7*(6*A*a*b**5*e/7 + A*b**6*d/7 + 15*B*a**2*b**4*e/7 + 6*B*a*b**5*d/7) + x**6*(5*A*a**2*b**4*e/2 + A*a*b**5*d + 10*B*a**3*b**3*e/3 + 5*B*a**2*b**4*d/2) + x**5*(4*A*a**3*b**3*e + 3*A*a**2*b**4*d + 3*B*a**4*b**2*e + 4*B*a**3*b**3*d) + x**4*(15*A*a**4*b**2*e/4 + 5*A*a**3*b**3*d + 3*B*a**5*b*e/2 + 15*B*a**4*b**2*d/4) + x**3*(2*A*a**5*b*e + 5*A*a**4*b**2*d + B*a**6*e/3 + 2*B*a**5*b*d) + x**2*(A*a**6*e/2 + 3*A*a**5*b*d + B*a**6*d/2)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 297 vs. $2(69) = 138$.

Time = 0.03 (sec) , antiderivative size = 297, normalized size of antiderivative = 3.96

$$\int (a + bx)^6 (A + Bx)(d + ex) dx$$

$$= \frac{1}{9} Bb^6 ex^9 + Aa^6 dx + \frac{1}{8} (Bb^6 d + (6 Bab^5 + Ab^6)e)x^8$$

$$+ \frac{1}{7} ((6 Bab^5 + Ab^6)d + 3(5 Ba^2 b^4 + 2 Aab^5)e)x^7$$

$$+ \frac{1}{6} (3(5 Ba^2 b^4 + 2 Aab^5)d + 5(4 Ba^3 b^3 + 3 Aa^2 b^4)e)x^6$$

$$+ ((4 Ba^3 b^3 + 3 Aa^2 b^4)d + (3 Ba^4 b^2 + 4 Aa^3 b^3)e)x^5$$

$$+ \frac{1}{4} (5(3 Ba^4 b^2 + 4 Aa^3 b^3)d + 3(2 Ba^5 b + 5 Aa^4 b^2)e)x^4$$

$$+ \frac{1}{3} (3(2 Ba^5 b + 5 Aa^4 b^2)d + (Ba^6 + 6 Aa^5 b)e)x^3 + \frac{1}{2} (Aa^6 e + (Ba^6 + 6 Aa^5 b)d)x^2$$

input `integrate((b*x+a)^6*(B*x+A)*(e*x+d),x, algorithm="maxima")`

output

```
1/9*B*b^6*e*x^9 + A*a^6*d*x + 1/8*(B*b^6*d + (6*B*a*b^5 + A*b^6)*e)*x^8 +
1/7*((6*B*a*b^5 + A*b^6)*d + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*e)*x^7 + 1/6*(3*(
5*B*a^2*b^4 + 2*A*a*b^5)*d + 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e)*x^6 + ((4*B*
a^3*b^3 + 3*A*a^2*b^4)*d + (3*B*a^4*b^2 + 4*A*a^3*b^3)*e)*x^5 + 1/4*(5*(3*
B*a^4*b^2 + 4*A*a^3*b^3)*d + 3*(2*B*a^5*b + 5*A*a^4*b^2)*e)*x^4 + 1/3*(3*(
2*B*a^5*b + 5*A*a^4*b^2)*d + (B*a^6 + 6*A*a^5*b)*e)*x^3 + 1/2*(A*a^6*e + (
B*a^6 + 6*A*a^5*b)*d)*x^2
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 321 vs. $2(69) = 138$.

Time = 0.12 (sec) , antiderivative size = 321, normalized size of antiderivative = 4.28

$$\begin{aligned} \int (a + bx)^6(A + Bx)(d + ex) dx = & \frac{1}{9} Bb^6ex^9 + \frac{1}{8} Bb^6dx^8 + \frac{3}{4} Bab^5ex^8 + \frac{1}{8} Ab^6ex^8 \\ & + \frac{6}{7} Bab^5dx^7 + \frac{1}{7} Ab^6dx^7 + \frac{15}{7} Ba^2b^4ex^7 + \frac{6}{7} Aab^5ex^7 \\ & + \frac{5}{2} Ba^2b^4dx^6 + Aab^5dx^6 + \frac{10}{3} Ba^3b^3ex^6 \\ & + \frac{5}{2} Aa^2b^4ex^6 + 4Ba^3b^3dx^5 + 3Aa^2b^4dx^5 \\ & + 3Ba^4b^2ex^5 + 4Aa^3b^3ex^5 + \frac{15}{4} Ba^4b^2dx^4 \\ & + 5Aa^3b^3dx^4 + \frac{3}{2} Ba^5bex^4 + \frac{15}{4} Aa^4b^2ex^4 \\ & + 2Ba^5bdx^3 + 5Aa^4b^2dx^3 + \frac{1}{3} Ba^6ex^3 + 2Aa^5bex^3 \\ & + \frac{1}{2} Ba^6dx^2 + 3Aa^5bdx^2 + \frac{1}{2} Aa^6ex^2 + Aa^6dx \end{aligned}$$

input `integrate((b*x+a)^6*(B*x+A)*(e*x+d),x, algorithm="giac")`

output `1/9*B*b^6*e*x^9 + 1/8*B*b^6*d*x^8 + 3/4*B*a*b^5*e*x^8 + 1/8*A*b^6*e*x^8 + 6/7*B*a*b^5*d*x^7 + 1/7*A*b^6*d*x^7 + 15/7*B*a^2*b^4*e*x^7 + 6/7*A*a*b^5*e*x^7 + 5/2*B*a^2*b^4*d*x^6 + A*a*b^5*d*x^6 + 10/3*B*a^3*b^3*e*x^6 + 5/2*A*a^2*b^4*e*x^6 + 4*B*a^3*b^3*d*x^5 + 3*A*a^2*b^4*d*x^5 + 3*B*a^4*b^2*e*x^5 + 4*A*a^3*b^3*e*x^5 + 15/4*B*a^4*b^2*d*x^4 + 5*A*a^3*b^3*d*x^4 + 3/2*B*a^5*b*e*x^4 + 15/4*A*a^4*b^2*e*x^4 + 2*B*a^5*b*d*x^3 + 5*A*a^4*b^2*d*x^3 + 1/3*B*a^6*e*x^3 + 2*A*a^5*b*e*x^3 + 1/2*B*a^6*d*x^2 + 3*A*a^5*b*d*x^2 + 1/2*A*a^6*e*x^2 + A*a^6*d*x`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 257, normalized size of antiderivative = 3.43

$$\begin{aligned}
& \int (a + bx)^6 (A + Bx)(d + ex) dx \\
&= x^3 \left(\frac{B a^6 e}{3} + 2 A a^5 b e + 2 B a^5 b d + 5 A a^4 b^2 d \right) \\
&+ x^7 \left(\frac{A b^6 d}{7} + \frac{6 A a b^5 e}{7} + \frac{6 B a b^5 d}{7} + \frac{15 B a^2 b^4 e}{7} \right) \\
&+ x^2 \left(\frac{A a^6 e}{2} + \frac{B a^6 d}{2} + 3 A a^5 b d \right) + x^8 \left(\frac{A b^6 e}{8} + \frac{B b^6 d}{8} + \frac{3 B a b^5 e}{4} \right) \\
&+ a^2 b^2 x^5 (3 A b^2 d + 3 B a^2 e + 4 A a b e + 4 B a b d) + A a^6 d x \\
&+ \frac{B b^6 e x^9}{9} + \frac{a^3 b x^4 (20 A b^2 d + 6 B a^2 e + 15 A a b e + 15 B a b d)}{4} \\
&+ \frac{a b^3 x^6 (6 A b^2 d + 20 B a^2 e + 15 A a b e + 15 B a b d)}{6}
\end{aligned}$$

input `int((A + B*x)*(a + b*x)^6*(d + e*x), x)`output `x^3*((B*a^6*e)/3 + 2*A*a^5*b*e + 2*B*a^5*b*d + 5*A*a^4*b^2*d) + x^7*((A*b^6*d)/7 + (6*A*a*b^5*e)/7 + (6*B*a*b^5*d)/7 + (15*B*a^2*b^4*e)/7) + x^2*((A*a^6*e)/2 + (B*a^6*d)/2 + 3*A*a^5*b*d) + x^8*((A*b^6*e)/8 + (B*b^6*d)/8 + (3*B*a*b^5*e)/4) + a^2*b^2*x^5*(3*A*b^2*d + 3*B*a^2*e + 4*A*a*b*e + 4*B*a*b*d) + A*a^6*d*x + (B*b^6*e*x^9)/9 + (a^3*b*x^4*(20*A*b^2*d + 6*B*a^2*e + 15*A*a*b*e + 15*B*a*b*d))/4 + (a*b^3*x^6*(6*A*b^2*d + 20*B*a^2*e + 15*A*a*b*e + 15*B*a*b*d))/6`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 169, normalized size of antiderivative = 2.25

$$\begin{aligned}
& \int (a + bx)^6 (A + Bx)(d + ex) dx \\
&= \frac{x(8b^7e x^8 + 63a b^6e x^7 + 9b^7d x^7 + 216a^2b^5e x^6 + 72a b^6d x^6 + 420a^3b^4e x^5 + 252a^2b^5d x^5 + 504a^4b^3e x^4 - \dots}{72}
\end{aligned}$$

input `int((b*x+a)^6*(B*x+A)*(e*x+d), x)`

output

```
(x*(72*a**7*d + 36*a**7*e*x + 252*a**6*b*d*x + 168*a**6*b*e*x**2 + 504*a**5*b**2*d*x**2 + 378*a**5*b**2*e*x**3 + 630*a**4*b**3*d*x**3 + 504*a**4*b**3*e*x**4 + 504*a**3*b**4*d*x**4 + 420*a**3*b**4*e*x**5 + 252*a**2*b**5*d*x**5 + 216*a**2*b**5*e*x**6 + 72*a*b**6*d*x**6 + 63*a*b**6*e*x**7 + 9*b**7*d*x**7 + 8*b**7*e*x**8))/72
```

3.49 $\int (a + bx)^6 (A + Bx) dx$

Optimal result	497
Mathematica [B] (verified)	497
Rubi [A] (verified)	498
Maple [B] (verified)	499
Fricas [B] (verification not implemented)	499
Sympy [B] (verification not implemented)	500
Maxima [B] (verification not implemented)	500
Giac [B] (verification not implemented)	501
Mupad [B] (verification not implemented)	501
Reduce [B] (verification not implemented)	502

Optimal result

Integrand size = 13, antiderivative size = 38

$$\int (a + bx)^6 (A + Bx) dx = \frac{(Ab - aB)(a + bx)^7}{7b^2} + \frac{B(a + bx)^8}{8b^2}$$

output

$$1/7*(A*b-B*a)*(b*x+a)^7/b^2+1/8*B*(b*x+a)^8/b^2$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 122 vs. $2(38) = 76$.

Time = 0.02 (sec) , antiderivative size = 122, normalized size of antiderivative = 3.21

$$\begin{aligned} \int (a + bx)^6 (A + Bx) dx = & \frac{1}{56}x(28a^6(2A + Bx) + 56a^5bx(3A + 2Bx) \\ & + 70a^4b^2x^2(4A + 3Bx) + 56a^3b^3x^3(5A + 4Bx) \\ & + 28a^2b^4x^4(6A + 5Bx) + 8ab^5x^5(7A + 6Bx) \\ & + b^6x^6(8A + 7Bx)) \end{aligned}$$

input

$$\text{Integrate}[(a + b*x)^6*(A + B*x), x]$$

output

```
(x*(28*a^6*(2*A + B*x) + 56*a^5*b*x*(3*A + 2*B*x) + 70*a^4*b^2*x^2*(4*A +
3*B*x) + 56*a^3*b^3*x^3*(5*A + 4*B*x) + 28*a^2*b^4*x^4*(6*A + 5*B*x) + 8*a
*b^5*x^5*(7*A + 6*B*x) + b^6*x^6*(8*A + 7*B*x)))/56
```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^6 (A + Bx) dx$$

$$\downarrow 49$$

$$\int \left(\frac{(a + bx)^6 (Ab - aB)}{b} + \frac{B(a + bx)^7}{b} \right) dx$$

$$\downarrow 2009$$

$$\frac{(a + bx)^7 (Ab - aB)}{7b^2} + \frac{B(a + bx)^8}{8b^2}$$

input

```
Int[(a + b*x)^6*(A + B*x),x]
```

output

```
((A*b - a*B)*(a + b*x)^7)/(7*b^2) + (B*(a + b*x)^8)/(8*b^2)
```

Defintions of rubi rules used

rule 49

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 139 vs. $2(34) = 68$.

Time = 0.14 (sec) , antiderivative size = 140, normalized size of antiderivative = 3.68

method	result
norman	$\frac{b^6 B x^8}{8} + \left(\frac{1}{7} b^6 A + \frac{6}{7} a b^5 B\right) x^7 + \left(a b^5 A + \frac{5}{2} a^2 b^4 B\right) x^6 + \left(3 a^2 b^4 A + 4 a^3 b^3 B\right) x^5 + \left(5 a^3 b^3 A + \frac{15}{2} a^4 b^2 B\right) x^4 + \left(20 a^3 b^3 A + 15 a^4 b^2 B\right) x^3 + \left(15 a^4 b^2 A + 10 a^5 b B\right) x^2 + \left(6 a^5 b A + 5 a^6 B\right) x + \frac{5}{2} a^6 A + \frac{5}{2} a^6 B$
default	$\frac{b^6 B x^8}{8} + \frac{(b^6 A + 6 a b^5 B) x^7}{7} + \frac{(6 a b^5 A + 15 a^2 b^4 B) x^6}{6} + \frac{(15 a^2 b^4 A + 20 a^3 b^3 B) x^5}{5} + \frac{(20 a^3 b^3 A + 15 a^4 b^2 B) x^4}{4} + \frac{(15 a^4 b^2 A + 10 a^5 b B) x^3}{3} + \frac{(6 a^5 b A + 5 a^6 B) x^2}{2} + \frac{5 a^6 A + 5 a^6 B}{2}$
gosper	$\frac{1}{8} b^6 B x^8 + \frac{1}{7} x^7 b^6 A + \frac{6}{7} x^7 a b^5 B + x^6 a b^5 A + \frac{5}{2} x^6 a^2 b^4 B + 3 A a^2 b^4 x^5 + 4 B a^3 b^3 x^5 + 5 x^4 a^3 b^3 B + \frac{15}{2} x^4 a^4 b^2 B + 20 a^3 b^3 A x^3 + 15 a^4 b^2 B x^3 + 15 a^4 b^2 A x^2 + 10 a^5 b B x^2 + 6 a^5 b A x + 5 a^6 B$
risch	$\frac{1}{8} b^6 B x^8 + \frac{1}{7} x^7 b^6 A + \frac{6}{7} x^7 a b^5 B + x^6 a b^5 A + \frac{5}{2} x^6 a^2 b^4 B + 3 A a^2 b^4 x^5 + 4 B a^3 b^3 x^5 + 5 x^4 a^3 b^3 B + \frac{15}{2} x^4 a^4 b^2 B + 20 a^3 b^3 A x^3 + 15 a^4 b^2 B x^3 + 15 a^4 b^2 A x^2 + 10 a^5 b B x^2 + 6 a^5 b A x + 5 a^6 B$
parallerisch	$\frac{1}{8} b^6 B x^8 + \frac{1}{7} x^7 b^6 A + \frac{6}{7} x^7 a b^5 B + x^6 a b^5 A + \frac{5}{2} x^6 a^2 b^4 B + 3 A a^2 b^4 x^5 + 4 B a^3 b^3 x^5 + 5 x^4 a^3 b^3 B + \frac{15}{2} x^4 a^4 b^2 B + 20 a^3 b^3 A x^3 + 15 a^4 b^2 B x^3 + 15 a^4 b^2 A x^2 + 10 a^5 b B x^2 + 6 a^5 b A x + 5 a^6 B$
orering	$\frac{x(7 B b^6 x^7 + 8 A b^6 x^6 + 48 B a b^5 x^6 + 56 A a b^5 x^5 + 140 B a^2 b^4 x^5 + 168 A a^2 b^4 x^4 + 224 B a^3 b^3 x^4 + 280 A a^3 b^3 x^3 + 210 B a^4 b^2 x^3 + 280 A a^4 b^2 x^2 + 140 B a^5 b x^2 + 50 A a^5 b x + 25 A a^6)}{56}$

input `int((b*x+a)^6*(B*x+A),x,method=_RETURNVERBOSE)`

output `1/8*b^6*B*x^8+(1/7*b^6*A+6/7*a*b^5*B)*x^7+(a*b^5*A+5/2*a^2*b^4*B)*x^6+(3*A*a^2*b^4+4*B*a^3*b^3)*x^5+(5*a^3*b^3*A+15/4*a^4*b^2*B)*x^4+(5*A*a^4*b^2+2*B*a^5*b)*x^3+(3*a^5*b*A+1/2*a^6*B)*x^2+a^6*A*x`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 142 vs. $2(34) = 68$.

Time = 0.07 (sec) , antiderivative size = 142, normalized size of antiderivative = 3.74

$$\int (a + bx)^6 (A + Bx) dx = \frac{1}{8} B b^6 x^8 + A a^6 x + \frac{1}{7} (6 B a b^5 + A b^6) x^7 + \frac{1}{2} (5 B a^2 b^4 + 2 A a b^5) x^6 + (4 B a^3 b^3 + 3 A a^2 b^4) x^5 + \frac{5}{4} (3 B a^4 b^2 + 4 A a^3 b^3) x^4 + (2 B a^5 b + 5 A a^4 b^2) x^3 + \frac{1}{2} (B a^6 + 6 A a^5 b) x^2$$

input `integrate((b*x+a)^6*(B*x+A),x, algorithm="fricas")`

output

```
1/8*B*b^6*x^8 + A*a^6*x + 1/7*(6*B*a*b^5 + A*b^6)*x^7 + 1/2*(5*B*a^2*b^4 +
2*A*a*b^5)*x^6 + (4*B*a^3*b^3 + 3*A*a^2*b^4)*x^5 + 5/4*(3*B*a^4*b^2 + 4*A
*a^3*b^3)*x^4 + (2*B*a^5*b + 5*A*a^4*b^2)*x^3 + 1/2*(B*a^6 + 6*A*a^5*b)*x^
2
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 148 vs. $2(32) = 64$.

Time = 0.03 (sec) , antiderivative size = 148, normalized size of antiderivative = 3.89

$$\int (a + bx)^6 (A + Bx) dx = Aa^6x + \frac{Bb^6x^8}{8} + x^7 \left(\frac{Ab^6}{7} + \frac{6Bab^5}{7} \right) + x^6 \left(Aab^5 + \frac{5Ba^2b^4}{2} \right) + x^5 \cdot (3Aa^2b^4 + 4Ba^3b^3) + x^4 \cdot \left(5Aa^3b^3 + \frac{15Ba^4b^2}{4} \right) + x^3 \cdot (5Aa^4b^2 + 2Ba^5b) + x^2 \cdot \left(3Aa^5b + \frac{Ba^6}{2} \right)$$

input

```
integrate((b*x+a)**6*(B*x+A),x)
```

output

```
A*a**6*x + B*b**6*x**8/8 + x**7*(A*b**6/7 + 6*B*a*b**5/7) + x**6*(A*a*b**5
+ 5*B*a**2*b**4/2) + x**5*(3*A*a**2*b**4 + 4*B*a**3*b**3) + x**4*(5*A*a**
3*b**3 + 15*B*a**4*b**2/4) + x**3*(5*A*a**4*b**2 + 2*B*a**5*b) + x**2*(3*A
*a**5*b + B*a**6/2)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 142 vs. $2(34) = 68$.

Time = 0.04 (sec) , antiderivative size = 142, normalized size of antiderivative = 3.74

$$\int (a + bx)^6 (A + Bx) dx = \frac{1}{8} Bb^6x^8 + Aa^6x + \frac{1}{7} (6Bab^5 + Ab^6)x^7 + \frac{1}{2} (5Ba^2b^4 + 2Aab^5)x^6 + (4Ba^3b^3 + 3Aa^2b^4)x^5 + \frac{5}{4} (3Ba^4b^2 + 4Aa^3b^3)x^4 + (2Ba^5b + 5Aa^4b^2)x^3 + \frac{1}{2} (Ba^6 + 6Aa^5b)x^2$$

input `integrate((b*x+a)^6*(B*x+A),x, algorithm="maxima")`

output $1/8*B*b^6*x^8 + A*a^6*x + 1/7*(6*B*a*b^5 + A*b^6)*x^7 + 1/2*(5*B*a^2*b^4 + 2*A*a*b^5)*x^6 + (4*B*a^3*b^3 + 3*A*a^2*b^4)*x^5 + 5/4*(3*B*a^4*b^2 + 4*A*a^3*b^3)*x^4 + (2*B*a^5*b + 5*A*a^4*b^2)*x^3 + 1/2*(B*a^6 + 6*A*a^5*b)*x^2$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 145 vs. $2(34) = 68$.

Time = 0.13 (sec) , antiderivative size = 145, normalized size of antiderivative = 3.82

$$\int (a + bx)^6(A + Bx) dx = \frac{1}{8} Bb^6x^8 + \frac{6}{7} Bab^5x^7 + \frac{1}{7} Ab^6x^7 + \frac{5}{2} Ba^2b^4x^6 + Aab^5x^6 + 4Ba^3b^3x^5 + 3Aa^2b^4x^5 + \frac{15}{4} Ba^4b^2x^4 + 5Aa^3b^3x^4 + 2Ba^5bx^3 + 5Aa^4b^2x^3 + \frac{1}{2} Ba^6x^2 + 3Aa^5bx^2 + Aa^6x$$

input `integrate((b*x+a)^6*(B*x+A),x, algorithm="giac")`

output $1/8*B*b^6*x^8 + 6/7*B*a*b^5*x^7 + 1/7*A*b^6*x^7 + 5/2*B*a^2*b^4*x^6 + A*a*b^5*x^6 + 4*B*a^3*b^3*x^5 + 3*A*a^2*b^4*x^5 + 15/4*B*a^4*b^2*x^4 + 5*A*a^3*b^3*x^4 + 2*B*a^5*b*x^3 + 5*A*a^4*b^2*x^3 + 1/2*B*a^6*x^2 + 3*A*a^5*b*x^2 + A*a^6*x$

Mupad [B] (verification not implemented)

Time = 0.96 (sec) , antiderivative size = 126, normalized size of antiderivative = 3.32

$$\int (a + bx)^6(A + Bx) dx = x^2 \left(\frac{B a^6}{2} + 3 A b a^5 \right) + x^7 \left(\frac{A b^6}{7} + \frac{6 B a b^5}{7} \right) + \frac{B b^6 x^8}{8} + A a^6 x + \frac{5 a^3 b^2 x^4 (4 A b + 3 B a)}{4} + a^2 b^3 x^5 (3 A b + 4 B a) + a^4 b x^3 (5 A b + 2 B a) + \frac{a b^4 x^6 (2 A b + 5 B a)}{2}$$

input `int((A + B*x)*(a + b*x)^6,x)`

output `x^2*((B*a^6)/2 + 3*A*a^5*b) + x^7*((A*b^6)/7 + (6*B*a*b^5)/7) + (B*b^6*x^8)/8 + A*a^6*x + (5*a^3*b^2*x^4*(4*A*b + 3*B*a))/4 + a^2*b^3*x^5*(3*A*b + 4*B*a) + a^4*b*x^3*(5*A*b + 2*B*a) + (a*b^4*x^6*(2*A*b + 5*B*a))/2`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 76, normalized size of antiderivative = 2.00

$$\int (a + bx)^6 (A + Bx) dx$$

$$= \frac{x(b^7 x^7 + 8a b^6 x^6 + 28a^2 b^5 x^5 + 56a^3 b^4 x^4 + 70a^4 b^3 x^3 + 56a^5 b^2 x^2 + 28a^6 b x + 8a^7)}{8}$$

input `int((b*x+a)^6*(B*x+A),x)`

output `(x*(8*a**7 + 28*a**6*b*x + 56*a**5*b**2*x**2 + 70*a**4*b**3*x**3 + 56*a**3*b**4*x**4 + 28*a**2*b**5*x**5 + 8*a*b**6*x**6 + b**7*x**7))/8`

3.50 $\int \frac{(a+bx)^6(A+Bx)}{d+ex} dx$

Optimal result	503
Mathematica [B] (verified)	504
Rubi [A] (verified)	504
Maple [B] (verified)	506
Fricas [B] (verification not implemented)	507
Sympy [B] (verification not implemented)	508
Maxima [B] (verification not implemented)	509
Giac [B] (verification not implemented)	510
Mupad [B] (verification not implemented)	512
Reduce [B] (verification not implemented)	513

Optimal result

Integrand size = 20, antiderivative size = 220

$$\int \frac{(a+bx)^6(A+Bx)}{d+ex} dx = \frac{b(bd-ae)^5(Bd-Ae)x}{e^7} - \frac{(bd-ae)^4(Bd-Ae)(a+bx)^2}{2e^6} + \frac{(bd-ae)^3(Bd-Ae)(a+bx)^3}{3e^5} - \frac{(bd-ae)^2(Bd-Ae)(a+bx)^4}{4e^4} + \frac{(bd-ae)(Bd-Ae)(a+bx)^5}{5e^3} - \frac{(Bd-Ae)(a+bx)^6}{6e^2} + \frac{B(a+bx)^7}{7be} - \frac{(bd-ae)^6(Bd-Ae)\log(d+ex)}{e^8}$$

output

```
b*(-a*e+b*d)^5*(-A*e+B*d)*x/e^7-1/2*(-a*e+b*d)^4*(-A*e+B*d)*(b*x+a)^2/e^6+
1/3*(-a*e+b*d)^3*(-A*e+B*d)*(b*x+a)^3/e^5-1/4*(-a*e+b*d)^2*(-A*e+B*d)*(b*x
+a)^4/e^4+1/5*(-a*e+b*d)*(-A*e+B*d)*(b*x+a)^5/e^3-1/6*(-A*e+B*d)*(b*x+a)^6
/e^2+1/7*B*(b*x+a)^7/b/e-(-a*e+b*d)^6*(-A*e+B*d)*ln(e*x+d)/e^8
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 501 vs. $2(220) = 440$.

Time = 0.16 (sec) , antiderivative size = 501, normalized size of antiderivative = 2.28

$$\int \frac{(a + bx)^6(A + Bx)}{d + ex} dx$$

$$= \frac{ex(420a^6Be^6 + 1260a^5be^5(-2Bd + 2Ae + Bex) + 1050a^4b^2e^4(3Ae(-2d + ex) + B(6d^2 - 3dex + 2e^2x^2$$

input `Integrate[((a + b*x)^6*(A + B*x))/(d + e*x),x]`

output

```
(e*x*(420*a^6*B*e^6 + 1260*a^5*b*e^5*(-2*B*d + 2*A*e + B*e*x) + 1050*a^4*b^2*e^4*(3*A*e*(-2*d + e*x) + B*(6*d^2 - 3*d*e*x + 2*e^2*x^2)) + 700*a^3*b^3*e^3*(2*A*e*(6*d^2 - 3*d*e*x + 2*e^2*x^2) + B*(-12*d^3 + 6*d^2*e*x - 4*d*e^2*x^2 + 3*e^3*x^3)) + 105*a^2*b^4*e^2*(5*A*e*(-12*d^3 + 6*d^2*e*x - 4*d*e^2*x^2 + 3*e^3*x^3) + B*(60*d^4 - 30*d^3*e*x + 20*d^2*e^2*x^2 - 15*d*e^3*x^3 + 12*e^4*x^4)) + 42*a*b^5*e*(A*e*(60*d^4 - 30*d^3*e*x + 20*d^2*e^2*x^2 - 15*d*e^3*x^3 + 12*e^4*x^4) + B*(-60*d^5 + 30*d^4*e*x - 20*d^3*e^2*x^2 + 15*d^2*e^3*x^3 - 12*d*e^4*x^4 + 10*e^5*x^5)) + b^6*(7*A*e*(-60*d^5 + 30*d^4*e*x - 20*d^3*e^2*x^2 + 15*d^2*e^3*x^3 - 12*d*e^4*x^4 + 10*e^5*x^5) + B*(420*d^6 - 210*d^5*e*x + 140*d^4*e^2*x^2 - 105*d^3*e^3*x^3 + 84*d^2*e^4*x^4 - 70*d*e^5*x^5 + 60*e^6*x^6))) - 420*(b*d - a*e)^6*(B*d - A*e)*Log[d + e*x])/(420*e^8)
```

Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^6(A + Bx)}{d + ex} dx$$

↓ 86

$$\int \left(\frac{(ae - bd)^6(Ae - Bd)}{e^7(d + ex)} - \frac{b(bd - ae)^5(Ae - Bd)}{e^7} + \frac{b(a + bx)(bd - ae)^4(Ae - Bd)}{e^6} - \frac{b(a + bx)^2(bd - ae)^3(Ae - Bd)}{e^5} \right)$$

↓ 2009

$$\begin{aligned} & - \frac{(bd - ae)^6(Bd - Ae) \log(d + ex)}{e^8} + \frac{bx(bd - ae)^5(Bd - Ae)}{e^7} - \\ & \frac{(a + bx)^2(bd - ae)^4(Bd - Ae)}{e^6} + \frac{(a + bx)^3(bd - ae)^3(Bd - Ae)}{e^5} - \\ & \frac{(a + bx)^4(bd - ae)^2(Bd - Ae)}{4e^4} + \frac{(a + bx)^5(bd - ae)(Bd - Ae)}{3e^3} - \frac{(a + bx)^6(Bd - Ae)}{6e^2} + \\ & \frac{B(a + bx)^7}{7be} \end{aligned}$$

input `Int[((a + b*x)^6*(A + B*x))/(d + e*x), x]`

output

```
(b*(b*d - a*e)^5*(B*d - A*e)*x)/e^7 - ((b*d - a*e)^4*(B*d - A*e)*(a + b*x)^2)/(2*e^6) + ((b*d - a*e)^3*(B*d - A*e)*(a + b*x)^3)/(3*e^5) - ((b*d - a*e)^2*(B*d - A*e)*(a + b*x)^4)/(4*e^4) + ((b*d - a*e)*(B*d - A*e)*(a + b*x)^5)/(5*e^3) - ((B*d - A*e)*(a + b*x)^6)/(6*e^2) + (B*(a + b*x)^7)/(7*b*e) - ((b*d - a*e)^6*(B*d - A*e)*Log[d + e*x])/e^8
```

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```


Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 763 vs. $2(208) = 416$.

Time = 0.08 (sec) , antiderivative size = 763, normalized size of antiderivative = 3.47

$$\int \frac{(a + bx)^6(A + Bx)}{d + ex} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d),x, algorithm="fricas")`

output

```
1/420*(60*B*b^6*e^7*x^7 - 70*(B*b^6*d*e^6 - (6*B*a*b^5 + A*b^6)*e^7)*x^6 +
84*(B*b^6*d^2*e^5 - (6*B*a*b^5 + A*b^6)*d*e^6 + 3*(5*B*a^2*b^4 + 2*A*a*b^
5)*e^7)*x^5 - 105*(B*b^6*d^3*e^4 - (6*B*a*b^5 + A*b^6)*d^2*e^5 + 3*(5*B*a^
2*b^4 + 2*A*a*b^5)*d*e^6 - 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^7)*x^4 + 140*(B
*b^6*d^4*e^3 - (6*B*a*b^5 + A*b^6)*d^3*e^4 + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*d
^2*e^5 - 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^6 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^
3)*e^7)*x^3 - 210*(B*b^6*d^5*e^2 - (6*B*a*b^5 + A*b^6)*d^4*e^3 + 3*(5*B*a^
2*b^4 + 2*A*a*b^5)*d^3*e^4 - 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + 5*(3*
B*a^4*b^2 + 4*A*a^3*b^3)*d*e^6 - 3*(2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 42
0*(B*b^6*d^6*e - (6*B*a*b^5 + A*b^6)*d^5*e^2 + 3*(5*B*a^2*b^4 + 2*A*a*b^5)
*d^4*e^3 - 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 + 5*(3*B*a^4*b^2 + 4*A*a^
3*b^3)*d^2*e^5 - 3*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^6 + (B*a^6 + 6*A*a^5*b)*e
^7)*x - 420*(B*b^6*d^7 - A*a^6*e^7 - (6*B*a*b^5 + A*b^6)*d^6*e + 3*(5*B*a^
2*b^4 + 2*A*a*b^5)*d^5*e^2 - 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4*e^3 + 5*(3*
B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 - 3*(2*B*a^5*b + 5*A*a^4*b^2)*d^2*e^5 + (
B*a^6 + 6*A*a^5*b)*d*e^6)*log(e*x + d))/e^8
```


Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 736 vs. $2(189) = 378$.

Time = 0.87 (sec) , antiderivative size = 736, normalized size of antiderivative = 3.35

$$\int \frac{(a+bx)^6(A+Bx)}{d+ex} dx = \frac{Bb^6x^7}{7e} + x^6 \left(\frac{Ab^6}{6e} + \frac{Bab^5}{e} - \frac{Bb^6d}{6e^2} \right) + x^5$$

$$\cdot \left(\frac{6Aab^5}{5e} - \frac{Ab^6d}{5e^2} + \frac{3Ba^2b^4}{e} - \frac{6Bab^5d}{5e^2} + \frac{Bb^6d^2}{5e^3} \right) + x^4$$

$$\cdot \left(\frac{15Aa^2b^4}{4e} - \frac{3Aab^5d}{2e^2} + \frac{Ab^6d^2}{4e^3} + \frac{5Ba^3b^3}{e} - \frac{15Ba^2b^4d}{4e^2} \right.$$

$$\left. + \frac{3Bab^5d^2}{2e^3} - \frac{Bb^6d^3}{4e^4} \right) + x^3$$

$$\cdot \left(\frac{20Aa^3b^3}{3e} - \frac{5Aa^2b^4d}{e^2} + \frac{2Aab^5d^2}{e^3} - \frac{Ab^6d^3}{3e^4} + \frac{5Ba^4b^2}{e} \right.$$

$$\left. - \frac{20Ba^3b^3d}{3e^2} + \frac{5Ba^2b^4d^2}{e^3} - \frac{2Bab^5d^3}{e^4} + \frac{Bb^6d^4}{3e^5} \right) + x^2$$

$$\cdot \left(\frac{15Aa^4b^2}{2e} - \frac{10Aa^3b^3d}{e^2} + \frac{15Aa^2b^4d^2}{2e^3} - \frac{3Aab^5d^3}{e^4} + \frac{Ab^6d^4}{2e^5} \right.$$

$$\left. + \frac{3Ba^5b}{e} - \frac{15Ba^4b^2d}{2e^2} + \frac{10Ba^3b^3d^2}{e^3} - \frac{15Ba^2b^4d^3}{2e^4} \right.$$

$$\left. + \frac{3Bab^5d^4}{e^5} - \frac{Bb^6d^5}{2e^6} \right)$$

$$+ x \left(\frac{6Aa^5b}{e} - \frac{15Aa^4b^2d}{e^2} + \frac{20Aa^3b^3d^2}{e^3} - \frac{15Aa^2b^4d^3}{e^4} \right.$$

$$\left. + \frac{6Aab^5d^4}{e^5} - \frac{Ab^6d^5}{e^6} + \frac{Ba^6}{e} - \frac{6Ba^5bd}{e^2} + \frac{15Ba^4b^2d^2}{e^3} \right.$$

$$\left. - \frac{20Ba^3b^3d^3}{e^4} + \frac{15Ba^2b^4d^4}{e^5} - \frac{6Bab^5d^5}{e^6} + \frac{Bb^6d^6}{e^7} \right)$$

$$- \frac{(-Ae+Bd)(ae-bd)^6 \log(d+ex)}{e^8}$$

input `integrate((b*x+a)**6*(B*x+A)/(e*x+d), x)`

output

```

B*b**6*x**7/(7*e) + x**6*(A*b**6/(6*e) + B*a*b**5/e - B*b**6*d/(6*e**2)) +
x**5*(6*A*a*b**5/(5*e) - A*b**6*d/(5*e**2) + 3*B*a**2*b**4/e - 6*B*a*b**5
*d/(5*e**2) + B*b**6*d**2/(5*e**3)) + x**4*(15*A*a**2*b**4/(4*e) - 3*A*a*b
**5*d/(2*e**2) + A*b**6*d**2/(4*e**3) + 5*B*a**3*b**3/e - 15*B*a**2*b**4*d
/(4*e**2) + 3*B*a*b**5*d**2/(2*e**3) - B*b**6*d**3/(4*e**4)) + x**3*(20*A*
a**3*b**3/(3*e) - 5*A*a**2*b**4*d/e**2 + 2*A*a*b**5*d**2/e**3 - A*b**6*d**
3/(3*e**4) + 5*B*a**4*b**2/e - 20*B*a**3*b**3*d/(3*e**2) + 5*B*a**2*b**4*d
**2/e**3 - 2*B*a*b**5*d**3/e**4 + B*b**6*d**4/(3*e**5)) + x**2*(15*A*a**4*
b**2/(2*e) - 10*A*a**3*b**3*d/e**2 + 15*A*a**2*b**4*d**2/(2*e**3) - 3*A*a*
b**5*d**3/e**4 + A*b**6*d**4/(2*e**5) + 3*B*a**5*b/e - 15*B*a**4*b**2*d/(2
*e**2) + 10*B*a**3*b**3*d**2/e**3 - 15*B*a**2*b**4*d**3/(2*e**4) + 3*B*a*b
**5*d**4/e**5 - B*b**6*d**5/(2*e**6)) + x*(6*A*a**5*b/e - 15*A*a**4*b**2*d
/e**2 + 20*A*a**3*b**3*d**2/e**3 - 15*A*a**2*b**4*d**3/e**4 + 6*A*a*b**5*d
**4/e**5 - A*b**6*d**5/e**6 + B*a**6/e - 6*B*a**5*b*d/e**2 + 15*B*a**4*b**
2*d**2/e**3 - 20*B*a**3*b**3*d**3/e**4 + 15*B*a**2*b**4*d**4/e**5 - 6*B*a*
b**5*d**5/e**6 + B*b**6*d**6/e**7) - (-A*e + B*d)*(a*e - b*d)**6*log(d + e
*x)/e**8

```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 762 vs. $2(208) = 416$.

Time = 0.06 (sec) , antiderivative size = 762, normalized size of antiderivative = 3.46

$$\int \frac{(a + bx)^6(A + Bx)}{d + ex} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^6*(B*x+A)/(e*x+d),x, algorithm="maxima")
```

output

```

1/420*(60*B*b^6*e^6*x^7 - 70*(B*b^6*d*e^5 - (6*B*a*b^5 + A*b^6)*e^6)*x^6 +
84*(B*b^6*d^2*e^4 - (6*B*a*b^5 + A*b^6)*d*e^5 + 3*(5*B*a^2*b^4 + 2*A*a*b^
5)*e^6)*x^5 - 105*(B*b^6*d^3*e^3 - (6*B*a*b^5 + A*b^6)*d^2*e^4 + 3*(5*B*a^
2*b^4 + 2*A*a*b^5)*d*e^5 - 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^6)*x^4 + 140*(B
*b^6*d^4*e^2 - (6*B*a*b^5 + A*b^6)*d^3*e^3 + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*d
^2*e^4 - 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^5 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^
3)*e^6)*x^3 - 210*(B*b^6*d^5*e - (6*B*a*b^5 + A*b^6)*d^4*e^2 + 3*(5*B*a^2*
b^4 + 2*A*a*b^5)*d^3*e^3 - 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^4 + 5*(3*B*
a^4*b^2 + 4*A*a^3*b^3)*d*e^5 - 3*(2*B*a^5*b + 5*A*a^4*b^2)*e^6)*x^2 + 420*
(B*b^6*d^6 - (6*B*a*b^5 + A*b^6)*d^5*e + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e
^2 - 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^3 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)
*d^2*e^4 - 3*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^5 + (B*a^6 + 6*A*a^5*b)*e^6)*x)
/e^7 - (B*b^6*d^7 - A*a^6*e^7 - (6*B*a*b^5 + A*b^6)*d^6*e + 3*(5*B*a^2*b^4
+ 2*A*a*b^5)*d^5*e^2 - 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4*e^3 + 5*(3*B*a^4
*b^2 + 4*A*a^3*b^3)*d^3*e^4 - 3*(2*B*a^5*b + 5*A*a^4*b^2)*d^2*e^5 + (B*a^6
+ 6*A*a^5*b)*d*e^6)*log(e*x + d)/e^8

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 910 vs. $2(208) = 416$.

Time = 0.12 (sec) , antiderivative size = 910, normalized size of antiderivative = 4.14

$$\int \frac{(a + bx)^6(A + Bx)}{d + ex} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^6*(B*x+A)/(e*x+d),x, algorithm="giac")
```

output

```

1/420*(60*B*b^6*e^6*x^7 - 70*B*b^6*d*e^5*x^6 + 420*B*a*b^5*e^6*x^6 + 70*A*
b^6*e^6*x^6 + 84*B*b^6*d^2*e^4*x^5 - 504*B*a*b^5*d*e^5*x^5 - 84*A*b^6*d*e^
5*x^5 + 1260*B*a^2*b^4*e^6*x^5 + 504*A*a*b^5*e^6*x^5 - 105*B*b^6*d^3*e^3*x
^4 + 630*B*a*b^5*d^2*e^4*x^4 + 105*A*b^6*d^2*e^4*x^4 - 1575*B*a^2*b^4*d*e^
5*x^4 - 630*A*a*b^5*d*e^5*x^4 + 2100*B*a^3*b^3*e^6*x^4 + 1575*A*a^2*b^4*e^
6*x^4 + 140*B*b^6*d^4*e^2*x^3 - 840*B*a*b^5*d^3*e^3*x^3 - 140*A*b^6*d^3*e^
3*x^3 + 2100*B*a^2*b^4*d^2*e^4*x^3 + 840*A*a*b^5*d^2*e^4*x^3 - 2800*B*a^3*
b^3*d*e^5*x^3 - 2100*A*a^2*b^4*d*e^5*x^3 + 2100*B*a^4*b^2*e^6*x^3 + 2800*A
*a^3*b^3*e^6*x^3 - 210*B*b^6*d^5*e*x^2 + 1260*B*a*b^5*d^4*e^2*x^2 + 210*A*
b^6*d^4*e^2*x^2 - 3150*B*a^2*b^4*d^3*e^3*x^2 - 1260*A*a*b^5*d^3*e^3*x^2 +
4200*B*a^3*b^3*d^2*e^4*x^2 + 3150*A*a^2*b^4*d^2*e^4*x^2 - 3150*B*a^4*b^2*d
*e^5*x^2 - 4200*A*a^3*b^3*d*e^5*x^2 + 1260*B*a^5*b*e^6*x^2 + 3150*A*a^4*b^
2*e^6*x^2 + 420*B*b^6*d^6*x - 2520*B*a*b^5*d^5*e*x - 420*A*b^6*d^5*e*x + 6
300*B*a^2*b^4*d^4*e^2*x + 2520*A*a*b^5*d^4*e^2*x - 8400*B*a^3*b^3*d^3*e^3*
x - 6300*A*a^2*b^4*d^3*e^3*x + 6300*B*a^4*b^2*d^2*e^4*x + 8400*A*a^3*b^3*d
^2*e^4*x - 2520*B*a^5*b*d*e^5*x - 6300*A*a^4*b^2*d*e^5*x + 420*B*a^6*e^6*x
+ 2520*A*a^5*b*e^6*x)/e^7 - (B*b^6*d^7 - 6*B*a*b^5*d^6*e - A*b^6*d^6*e +
15*B*a^2*b^4*d^5*e^2 + 6*A*a*b^5*d^5*e^2 - 20*B*a^3*b^3*d^4*e^3 - 15*A*a^2
*b^4*d^4*e^3 + 15*B*a^4*b^2*d^3*e^4 + 20*A*a^3*b^3*d^3*e^4 - 6*B*a^5*b*d^2
*e^5 - 15*A*a^4*b^2*d^2*e^5 + B*a^6*d*e^6 + 6*A*a^5*b*d*e^6 - A*a^6*e^7...

```

Mupad [B] (verification not implemented)

Time = 0.94 (sec) , antiderivative size = 769, normalized size of antiderivative = 3.50

$$\int \frac{(a + bx)^6(A + Bx)}{d + ex} dx = x \frac{Ba^6 + 6Aba^5}{e}$$

$$d \left(\frac{d \left(\frac{d \left(\frac{Ab^6 + 6Bab^5 - Bb^6d}{e} - \frac{3ab^4(2Ab + 5Ba)}{e} \right) + \frac{5a^2b^3(3Ab + 4Ba)}{e} \right) - \frac{5a^3b^2(4Ab + 3Ba)}{e}}{e} \right) + \frac{3a^4b(5Ab + 2Ba)}{e}$$

output

```
(420*log(d + e*x)*a**7*e**7 - 2940*log(d + e*x)*a**6*b*d*e**6 + 8820*log(d
+ e*x)*a**5*b**2*d**2*e**5 - 14700*log(d + e*x)*a**4*b**3*d**3*e**4 + 147
00*log(d + e*x)*a**3*b**4*d**4*e**3 - 8820*log(d + e*x)*a**2*b**5*d**5*e**
2 + 2940*log(d + e*x)*a*b**6*d**6*e - 420*log(d + e*x)*b**7*d**7 + 2940*a*
*6*b*e**7*x - 8820*a**5*b**2*d*e**6*x + 4410*a**5*b**2*e**7*x**2 + 14700*a
**4*b**3*d**2*e**5*x - 7350*a**4*b**3*d*e**6*x**2 + 4900*a**4*b**3*e**7*x*
*3 - 14700*a**3*b**4*d**3*e**4*x + 7350*a**3*b**4*d**2*e**5*x**2 - 4900*a*
*3*b**4*d*e**6*x**3 + 3675*a**3*b**4*e**7*x**4 + 8820*a**2*b**5*d**4*e**3*
x - 4410*a**2*b**5*d**3*e**4*x**2 + 2940*a**2*b**5*d**2*e**5*x**3 - 2205*a
**2*b**5*d*e**6*x**4 + 1764*a**2*b**5*e**7*x**5 - 2940*a*b**6*d**5*e**2*x
+ 1470*a*b**6*d**4*e**3*x**2 - 980*a*b**6*d**3*e**4*x**3 + 735*a*b**6*d**2
*e**5*x**4 - 588*a*b**6*d*e**6*x**5 + 490*a*b**6*e**7*x**6 + 420*b**7*d**6
*e*x - 210*b**7*d**5*e**2*x**2 + 140*b**7*d**4*e**3*x**3 - 105*b**7*d**3*e
**4*x**4 + 84*b**7*d**2*e**5*x**5 - 70*b**7*d*e**6*x**6 + 60*b**7*e**7*x**
7)/(420*e**8)
```

3.51 $\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^2} dx$

Optimal result	515
Mathematica [B] (verified)	516
Rubi [A] (verified)	516
Maple [B] (verified)	518
Fricas [B] (verification not implemented)	519
Sympy [B] (verification not implemented)	520
Maxima [B] (verification not implemented)	521
Giac [B] (verification not implemented)	522
Mupad [B] (verification not implemented)	523
Reduce [B] (verification not implemented)	524

Optimal result

Integrand size = 20, antiderivative size = 277

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^2} dx = -\frac{3b(bd-ae)^4(7bBd-5Abe-2aBe)x}{e^7} + \frac{(bd-ae)^6(Bd-Ae)}{e^8(d+ex)} + \frac{5b^2(bd-ae)^3(7bBd-4Abe-3aBe)(d+ex)^2}{2e^8} - \frac{5b^3(bd-ae)^2(7bBd-3Abe-4aBe)(d+ex)^3}{3e^8} + \frac{3b^4(bd-ae)(7bBd-2Abe-5aBe)(d+ex)^4}{4e^8} - \frac{b^5(7bBd-Abe-6aBe)(d+ex)^5}{5e^8} + \frac{b^6B(d+ex)^6}{6e^8} + \frac{(bd-ae)^5(7bBd-6Abe-aBe)\log(d+ex)}{e^8}$$

output

```
-3*b*(-a*e+b*d)^4*(-5*A*b*e-2*B*a*e+7*B*b*d)*x/e^7+(-a*e+b*d)^6*(-A*e+B*d)/e^8/(e*x+d)+5/2*b^2*(-a*e+b*d)^3*(-4*A*b*e-3*B*a*e+7*B*b*d)*(e*x+d)^2/e^8-5/3*b^3*(-a*e+b*d)^2*(-3*A*b*e-4*B*a*e+7*B*b*d)*(e*x+d)^3/e^8+3/4*b^4*(-a*e+b*d)*(-2*A*b*e-5*B*a*e+7*B*b*d)*(e*x+d)^4/e^8-1/5*b^5*(-A*b*e-6*B*a*e+7*B*b*d)*(e*x+d)^5/e^8+1/6*b^6*B*(e*x+d)^6/e^8+(-a*e+b*d)^5*(-6*A*b*e-B*a*e+7*B*b*d)*ln(e*x+d)/e^8
```


Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 643 vs. $2(277) = 554$.

Time = 0.18 (sec) , antiderivative size = 643, normalized size of antiderivative = 2.32

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^2} dx$$

$$= \frac{60a^6e^6(Bd - Ae) + 360a^5be^5(Ade + B(-d^2 + dex + e^2x^2)) + 450a^4b^2e^4(2Ae(-d^2 + dex + e^2x^2) + B(2d^3 - 4d^2ex - 3de^2x^2 + e^3x^3)) + 200a^3b^3e^3(3Ae(2d^3 - 4d^2ex - 3de^2x^2 + e^3x^3) + 2B(-3d^4 + 9d^3ex + 6d^2e^2x^2 - 2de^3x^3 + e^4x^4)) + 75a^2b^4e^2(4Ae(-3d^4 + 9d^3ex + 6d^2e^2x^2 - 2de^3x^3 + e^4x^4) + B(12d^5 - 48d^4ex - 30d^3e^2x^2 + 10d^2e^3x^3 - 5de^4x^4 + 3e^5x^5)) + 6ab^5e(5Ae(12d^5 - 48d^4ex - 30d^3e^2x^2 + 10d^2e^3x^3 - 5de^4x^4 + 3e^5x^5) - 6B(10d^6 - 50d^5ex - 30d^4e^2x^2 + 10d^3e^3x^3 - 5d^2e^4x^4 + 3de^5x^5 - 2e^6x^6)) + b^6(6Ae(-10d^6 + 50d^5ex + 30d^4e^2x^2 - 10d^3e^3x^3 + 5d^2e^4x^4 - 3de^5x^5 + 2e^6x^6) + B(60d^7 - 360d^6ex - 210d^5e^2x^2 + 70d^4e^3x^3 - 35d^3e^4x^4 + 21d^2e^5x^5 - 14de^6x^6 + 10e^7x^7)) + 60(bd - ae)^5(7bBd - 6Abe - aBe)(d + ex) \cdot \text{Log}[d + ex]}{(60e^8(d + ex))}$$

input

```
Integrate[((a + b*x)^6*(A + B*x))/(d + e*x)^2,x]
```

output

```
(60*a^6*e^6*(B*d - A*e) + 360*a^5*b*e^5*(A*d*e + B*(-d^2 + d*e*x + e^2*x^2)) + 450*a^4*b^2*e^4*(2*A*e*(-d^2 + d*e*x + e^2*x^2) + B*(2*d^3 - 4*d^2*e*x - 3*d*e^2*x^2 + e^3*x^3)) + 200*a^3*b^3*e^3*(3*A*e*(2*d^3 - 4*d^2*e*x - 3*d*e^2*x^2 + e^3*x^3) + 2*B*(-3*d^4 + 9*d^3*e*x + 6*d^2*e^2*x^2 - 2*d*e^3*x^3 + e^4*x^4)) + 75*a^2*b^4*e^2*(4*A*e*(-3*d^4 + 9*d^3*e*x + 6*d^2*e^2*x^2 - 2*d*e^3*x^3 + e^4*x^4) + B*(12*d^5 - 48*d^4*e*x - 30*d^3*e^2*x^2 + 10*d^2*e^3*x^3 - 5*d*e^4*x^4 + 3*e^5*x^5)) + 6*a*b^5*e*(5*A*e*(12*d^5 - 48*d^4*e*x - 30*d^3*e^2*x^2 + 10*d^2*e^3*x^3 - 5*d*e^4*x^4 + 3*e^5*x^5) - 6*B*(10*d^6 - 50*d^5*e*x - 30*d^4*e^2*x^2 + 10*d^3*e^3*x^3 - 5*d^2*e^4*x^4 + 3*d*e^5*x^5 - 2*e^6*x^6)) + b^6*(6*A*e*(-10*d^6 + 50*d^5*e*x + 30*d^4*e^2*x^2 - 10*d^3*e^3*x^3 + 5*d^2*e^4*x^4 - 3*d*e^5*x^5 + 2*e^6*x^6) + B*(60*d^7 - 360*d^6*e*x - 210*d^5*e^2*x^2 + 70*d^4*e^3*x^3 - 35*d^3*e^4*x^4 + 21*d^2*e^5*x^5 - 14*d*e^6*x^6 + 10*e^7*x^7)) + 60*(b*d - a*e)^5*(7*b*B*d - 6*A*b*e - a*B*e)*(d + e*x)*Log[d + e*x]/(60*e^8*(d + e*x))
```

Rubi [A] (verified)

Time = 0.93 (sec) , antiderivative size = 277, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^6 (A + Bx)}{(d + ex)^2} dx$$

↓ 86

$$\int \left(\frac{b^5(d + ex)^4(6aBe + Abe - 7bBd)}{e^7} - \frac{3b^4(d + ex)^3(bd - ae)(5aBe + 2Abe - 7bBd)}{e^7} + \frac{5b^3(d + ex)^2(bd - ae)}{e^7} \right) dx$$

↓ 2009

$$\begin{aligned} & - \frac{b^5(d + ex)^5(-6aBe - Abe + 7bBd)}{5e^8} + \frac{3b^4(d + ex)^4(bd - ae)(-5aBe - 2Abe + 7bBd)}{4e^8} - \\ & - \frac{5b^3(d + ex)^3(bd - ae)^2(-4aBe - 3Abe + 7bBd)}{3e^8} + \\ & + \frac{5b^2(d + ex)^2(bd - ae)^3(-3aBe - 4Abe + 7bBd)}{2e^8} + \frac{(bd - ae)^6(Bd - Ae)}{e^8(d + ex)} + \\ & + \frac{(bd - ae)^5 \log(d + ex)(-aBe - 6Abe + 7bBd)}{e^8} - \frac{3bx(bd - ae)^4(-2aBe - 5Abe + 7bBd)}{e^7} + \\ & + \frac{b^6 B(d + ex)^6}{6e^8} \end{aligned}$$

input

```
Int[((a + b*x)^6*(A + B*x))/(d + e*x)^2,x]
```

output

```
(-3*b*(b*d - a*e)^4*(7*b*B*d - 5*A*b*e - 2*a*B*e)*x)/e^7 + ((b*d - a*e)^6*(B*d - A*e))/(e^8*(d + e*x)) + (5*b^2*(b*d - a*e)^3*(7*b*B*d - 4*A*b*e - 3*a*B*e)*(d + e*x)^2)/(2*e^8) - (5*b^3*(b*d - a*e)^2*(7*b*B*d - 3*A*b*e - 4*a*B*e)*(d + e*x)^3)/(3*e^8) + (3*b^4*(b*d - a*e)*(7*b*B*d - 2*A*b*e - 5*a*B*e)*(d + e*x)^4)/(4*e^8) - (b^5*(7*b*B*d - A*b*e - 6*a*B*e)*(d + e*x)^5)/(5*e^8) + (b^6*B*(d + e*x)^6)/(6*e^8) + ((b*d - a*e)^5*(7*b*B*d - 6*A*b*e - a*B*e)*Log[d + e*x])/e^8
```

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 794 vs. $2(267) = 534$.

Time = 0.22 (sec) , antiderivative size = 795, normalized size of antiderivative = 2.87

method	result
norman	$\frac{(a^6 A e^7 - 6A a^5 b d e^6 + 30A a^4 b^2 d^2 e^5 - 60A a^3 b^3 d^3 e^4 + 60A a^2 b^4 d^4 e^3 - 30A a b^5 d^5 e^2 + 6A b^6 d^6 e - B a^6 d e^6 + 12B a^5 b d^2 e^5 - 45B a^4 b^2 d^3 e^4 + \dots)}{e^7 d}$
default	$b(-30B a^4 b d e^4 x + 60B a^3 b^2 d^2 e^3 x - 60B a^2 b^3 d^3 e^2 x + 30B a b^4 d^4 e x + A b^5 d^2 e^3 x^3 + 6B a^5 e^5 x - 6B b^5 d^5 x + 5A a^2 b^3 e^5 x^3 + \frac{20}{3} B a^2 b^3 e^5 x^3 + \dots)$
risch	Expression too large to display
parallelrisc	Expression too large to display

input `int((b*x+a)^6*(B*x+A)/(e*x+d)^2,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & ((A*a^6*e^7-6*A*a^5*b*d*e^6+30*A*a^4*b^2*d^2*e^5-60*A*a^3*b^3*d^3*e^4+60*A \\ & *a^2*b^4*d^4*e^3-30*A*a*b^5*d^5*e^2+6*A*b^6*d^6*e-B*a^6*d*e^6+12*B*a^5*b*d \\ & ^2*e^5-45*B*a^4*b^2*d^3*e^4+80*B*a^3*b^3*d^4*e^3-75*B*a^2*b^4*d^5*e^2+36*B \\ & *a*b^5*d^6*e-7*B*b^6*d^7)/e^7/d*x+1/2*b*(30*A*a^4*b*e^5-60*A*a^3*b^2*d*e^4 \\ & +60*A*a^2*b^3*d^2*e^3-30*A*a*b^4*d^3*e^2+6*A*b^5*d^4*e+12*B*a^5*e^5-45*B*a \\ & ^4*b*d*e^4+80*B*a^3*b^2*d^2*e^3-75*B*a^2*b^3*d^3*e^2+36*B*a*b^4*d^4*e-7*B* \\ & b^5*d^5)/e^6*x^2+1/6*b^2*(60*A*a^3*b*e^4-60*A*a^2*b^2*d*e^3+30*A*a*b^3*d^2 \\ & *e^2-6*A*b^4*d^3*e+45*B*a^4*e^4-80*B*a^3*b*d*e^3+75*B*a^2*b^2*d^2*e^2-36*B \\ & *a*b^3*d^3*e+7*B*b^4*d^4)/e^5*x^3+1/12*b^3*(60*A*a^2*b*e^3-30*A*a*b^2*d*e^ \\ & 2+6*A*b^3*d^2*e+80*B*a^3*e^3-75*B*a^2*b*d*e^2+36*B*a*b^2*d^2*e-7*B*b^3*d^3 \\ &)/e^4*x^4+1/20*b^4*(30*A*a*b*e^2-6*A*b^2*d*e+75*B*a^2*e^2-36*B*a*b*d*e+7*B \\ & *b^2*d^2)/e^3*x^5+1/30*b^5*(6*A*b*e+36*B*a*e-7*B*b*d)/e^2*x^6+1/6*b^6*B/e* \\ & x^7)/(e*x+d)+1/e^8*(6*A*a^5*b*e^6-30*A*a^4*b^2*d*e^5+60*A*a^3*b^3*d^2*e^4- \\ & 60*A*a^2*b^4*d^3*e^3+30*A*a*b^5*d^4*e^2-6*A*b^6*d^5*e+B*a^6*e^6-12*B*a^5*b \\ & *d*e^5+45*B*a^4*b^2*d^2*e^4-80*B*a^3*b^3*d^3*e^3+75*B*a^2*b^4*d^4*e^2-36*B \\ & *a*b^5*d^5*e+7*B*b^6*d^6)*ln(e*x+d) \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1067 vs. $2(267) = 534$.

Time = 0.09 (sec) , antiderivative size = 1067, normalized size of antiderivative = 3.85

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^2} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^2,x, algorithm="fricas")`

output

```
1/60*(10*B*b^6*e^7*x^7 + 60*B*b^6*d^7 - 60*A*a^6*e^7 - 60*(6*B*a*b^5 + A*b^6)*d^6*e + 180*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 - 300*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4*e^3 + 300*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 - 180*(2*B*a^5*b + 5*A*a^4*b^2)*d^2*e^5 + 60*(B*a^6 + 6*A*a^5*b)*d*e^6 - 2*(7*B*b^6*d*e^6 - 6*(6*B*a*b^5 + A*b^6)*e^7)*x^6 + 3*(7*B*b^6*d^2*e^5 - 6*(6*B*a*b^5 + A*b^6)*d*e^6 + 15*(5*B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^5 - 5*(7*B*b^6*d^3*e^4 - 6*(6*B*a*b^5 + A*b^6)*d^2*e^5 + 15*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6 - 20*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^7)*x^4 + 10*(7*B*b^6*d^4*e^3 - 6*(6*B*a*b^5 + A*b^6)*d^3*e^4 + 15*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 - 20*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^6 + 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^3 - 30*(7*B*b^6*d^5*e^2 - 6*(6*B*a*b^5 + A*b^6)*d^4*e^3 + 15*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*e^4 - 20*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^6 - 6*(2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 - 60*(6*B*b^6*d^6*e - 5*(6*B*a*b^5 + A*b^6)*d^5*e^2 + 12*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^3 - 15*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 + 10*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^5 - 3*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^6)*x + 60*(7*B*b^6*d^7 - 6*(6*B*a*b^5 + A*b^6)*d^6*e + 15*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 - 20*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4*e^3 + 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 - 6*(2*B*a^5*b + 5*A*a^4*b^2)*d^2*e^5 + (B*a^6 + 6*A*a^5*b)*d*e^6 + (7*B*b^6*d^6*e - 6*(6*B*a*b^5 + A*b^6)*d^5*e^2 + 15*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^3 - ...
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 782 vs. $2(284) = 568$.

Time = 1.84 (sec) , antiderivative size = 782, normalized size of antiderivative = 2.82

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^2} dx = \frac{Bb^6x^6}{6e^2} + x^5 \left(\frac{Ab^6}{5e^2} + \frac{6Bab^5}{5e^2} - \frac{2Bb^6d}{5e^3} \right) \\ + x^4 \cdot \left(\frac{3Aab^5}{2e^2} - \frac{Ab^6d}{2e^3} + \frac{15Ba^2b^4}{4e^2} - \frac{3Bab^5d}{e^3} + \frac{3Bb^6d^2}{4e^4} \right) + x^3 \\ \cdot \left(\frac{5Aa^2b^4}{e^2} - \frac{4Aab^5d}{e^3} + \frac{Ab^6d^2}{e^4} + \frac{20Ba^3b^3}{3e^2} - \frac{10Ba^2b^4d}{e^3} + \frac{6Bab^5d^2}{e^4} - \frac{4Bb^6d^3}{3e^5} \right) + x^2 \\ \cdot \left(\frac{10Aa^3b^3}{e^2} - \frac{15Aa^2b^4d}{e^3} + \frac{9Aab^5d^2}{e^4} - \frac{2Ab^6d^3}{e^5} + \frac{15Ba^4b^2}{2e^2} - \frac{20Ba^3b^3d}{e^3} + \frac{45Ba^2b^4d^2}{2e^4} \right. \\ \left. - \frac{12Bab^5d^3}{e^5} + \frac{5Bb^6d^4}{2e^6} \right) + x \left(\frac{15Aa^4b^2}{e^2} - \frac{40Aa^3b^3d}{e^3} + \frac{45Aa^2b^4d^2}{e^4} - \frac{24Aab^5d^3}{e^5} \right. \\ \left. + \frac{5Ab^6d^4}{e^6} + \frac{6Ba^5b}{e^2} - \frac{30Ba^4b^2d}{e^3} + \frac{60Ba^3b^3d^2}{e^4} - \frac{60Ba^2b^4d^3}{e^5} + \frac{30Bab^5d^4}{e^6} - \frac{6Bb^6d^5}{e^7} \right) \\ + \frac{-Aa^6e^7 + 6Aa^5bde^6 - 15Aa^4b^2d^2e^5 + 20Aa^3b^3d^3e^4 - 15Aa^2b^4d^4e^3 + 6Aab^5d^5e^2 - Ab^6d^6e + Ba^6de^6}{de^8 + e^9x} \\ + \frac{(ae-bd)^5 \cdot (6Abe + Bae - 7Bbd) \log(d+ex)}{e^8}$$

input `integrate((b*x+a)**6*(B*x+A)/(e*x+d)**2,x)`

output

```

B*b**6*x**6/(6*e**2) + x**5*(A*b**6/(5*e**2) + 6*B*a*b**5/(5*e**2) - 2*B*b
**6*d/(5*e**3)) + x**4*(3*A*a*b**5/(2*e**2) - A*b**6*d/(2*e**3) + 15*B*a**
2*b**4/(4*e**2) - 3*B*a*b**5*d/e**3 + 3*B*b**6*d**2/(4*e**4)) + x**3*(5*A*
a**2*b**4/e**2 - 4*A*a*b**5*d/e**3 + A*b**6*d**2/e**4 + 20*B*a**3*b**3/(3*
e**2) - 10*B*a**2*b**4*d/e**3 + 6*B*a*b**5*d**2/e**4 - 4*B*b**6*d**3/(3*e*
*5)) + x**2*(10*A*a**3*b**3/e**2 - 15*A*a**2*b**4*d/e**3 + 9*A*a*b**5*d**2
/e**4 - 2*A*b**6*d**3/e**5 + 15*B*a**4*b**2/(2*e**2) - 20*B*a**3*b**3*d/e*
*3 + 45*B*a**2*b**4*d**2/(2*e**4) - 12*B*a*b**5*d**3/e**5 + 5*B*b**6*d**4/
(2*e**6)) + x*(15*A*a**4*b**2/e**2 - 40*A*a**3*b**3*d/e**3 + 45*A*a**2*b**
4*d**2/e**4 - 24*A*a*b**5*d**3/e**5 + 5*A*b**6*d**4/e**6 + 6*B*a**5*b/e**2
- 30*B*a**4*b**2*d/e**3 + 60*B*a**3*b**3*d**2/e**4 - 60*B*a**2*b**4*d**3/
e**5 + 30*B*a*b**5*d**4/e**6 - 6*B*b**6*d**5/e**7) + (-A*a**6*e**7 + 6*A*a
**5*b*d*e**6 - 15*A*a**4*b**2*d**2*e**5 + 20*A*a**3*b**3*d**3*e**4 - 15*A*
a**2*b**4*d**4*e**3 + 6*A*a*b**5*d**5*e**2 - A*b**6*d**6*e + B*a**6*d*e**6
- 6*B*a**5*b*d**2*e**5 + 15*B*a**4*b**2*d**3*e**4 - 20*B*a**3*b**3*d**4*e
**3 + 15*B*a**2*b**4*d**5*e**2 - 6*B*a*b**5*d**6*e + B*b**6*d**7)/(d*e**8
+ e**9*x) + (a*e - b*d)**5*(6*A*b*e + B*a*e - 7*B*b*d)*log(d + e*x)/e**8

```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 771 vs. $2(267) = 534$.

Time = 0.06 (sec) , antiderivative size = 771, normalized size of antiderivative = 2.78

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^2} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^6*(B*x+A)/(e*x+d)^2,x, algorithm="maxima")
```

output

```
(B*b^6*d^7 - A*a^6*e^7 - (6*B*a*b^5 + A*b^6)*d^6*e + 3*(5*B*a^2*b^4 + 2*A*
a*b^5)*d^5*e^2 - 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4*e^3 + 5*(3*B*a^4*b^2 +
4*A*a^3*b^3)*d^3*e^4 - 3*(2*B*a^5*b + 5*A*a^4*b^2)*d^2*e^5 + (B*a^6 + 6*A*
a^5*b)*d*e^6)/(e^9*x + d*e^8) + 1/60*(10*B*b^6*e^5*x^6 - 12*(2*B*b^6*d*e^4
- (6*B*a*b^5 + A*b^6)*e^5)*x^5 + 15*(3*B*b^6*d^2*e^3 - 2*(6*B*a*b^5 + A*b
^6)*d*e^4 + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*e^5)*x^4 - 20*(4*B*b^6*d^3*e^2 - 3
*(6*B*a*b^5 + A*b^6)*d^2*e^3 + 6*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^4 - 5*(4*B*
a^3*b^3 + 3*A*a^2*b^4)*e^5)*x^3 + 30*(5*B*b^6*d^4*e - 4*(6*B*a*b^5 + A*b^6
)*d^3*e^2 + 9*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^3 - 10*(4*B*a^3*b^3 + 3*A*a^
2*b^4)*d*e^4 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^5)*x^2 - 60*(6*B*b^6*d^5 -
5*(6*B*a*b^5 + A*b^6)*d^4*e + 12*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*e^2 - 15*(4
*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^3 + 10*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^4 -
3*(2*B*a^5*b + 5*A*a^4*b^2)*e^5)*x)/e^7 + (7*B*b^6*d^6 - 6*(6*B*a*b^5 + A
*b^6)*d^5*e + 15*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^2 - 20*(4*B*a^3*b^3 + 3*A
*a^2*b^4)*d^3*e^3 + 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^4 - 6*(2*B*a^5*b
+ 5*A*a^4*b^2)*d*e^5 + (B*a^6 + 6*A*a^5*b)*e^6)*log(e*x + d)/e^8
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 979 vs. $2(267) = 534$.

Time = 0.13 (sec) , antiderivative size = 979, normalized size of antiderivative = 3.53

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^2} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^6*(B*x+A)/(e*x+d)^2,x, algorithm="giac")
```

output

```

1/60*(10*B*b^6 - 12*(7*B*b^6*d*e - 6*B*a*b^5*e^2 - A*b^6*e^2)/((e*x + d)*e
) + 45*(7*B*b^6*d^2*e^2 - 12*B*a*b^5*d*e^3 - 2*A*b^6*d*e^3 + 5*B*a^2*b^4*e
^4 + 2*A*a*b^5*e^4)/((e*x + d)^2*e^2) - 100*(7*B*b^6*d^3*e^3 - 18*B*a*b^5*
d^2*e^4 - 3*A*b^6*d^2*e^4 + 15*B*a^2*b^4*d*e^5 + 6*A*a*b^5*d*e^5 - 4*B*a^3
*b^3*e^6 - 3*A*a^2*b^4*e^6)/((e*x + d)^3*e^3) + 150*(7*B*b^6*d^4*e^4 - 24*
B*a*b^5*d^3*e^5 - 4*A*b^6*d^3*e^5 + 30*B*a^2*b^4*d^2*e^6 + 12*A*a*b^5*d^2*
e^6 - 16*B*a^3*b^3*d*e^7 - 12*A*a^2*b^4*d*e^7 + 3*B*a^4*b^2*e^8 + 4*A*a^3*
b^3*e^8)/((e*x + d)^4*e^4) - 180*(7*B*b^6*d^5*e^5 - 30*B*a*b^5*d^4*e^6 - 5
*A*b^6*d^4*e^6 + 50*B*a^2*b^4*d^3*e^7 + 20*A*a*b^5*d^3*e^7 - 40*B*a^3*b^3*
d^2*e^8 - 30*A*a^2*b^4*d^2*e^8 + 15*B*a^4*b^2*d*e^9 + 20*A*a^3*b^3*d*e^9 -
2*B*a^5*b*e^10 - 5*A*a^4*b^2*e^10)/((e*x + d)^5*e^5))*(e*x + d)^6/e^8 - (
7*B*b^6*d^6 - 36*B*a*b^5*d^5*e - 6*A*b^6*d^5*e + 75*B*a^2*b^4*d^4*e^2 + 30
*A*a*b^5*d^4*e^2 - 80*B*a^3*b^3*d^3*e^3 - 60*A*a^2*b^4*d^3*e^3 + 45*B*a^4*
b^2*d^2*e^4 + 60*A*a^3*b^3*d^2*e^4 - 12*B*a^5*b*d*e^5 - 30*A*a^4*b^2*d*e^5
+ B*a^6*e^6 + 6*A*a^5*b*e^6)*log(abs(e*x + d)/((e*x + d)^2*abs(e)))/e^8 +
(B*b^6*d^7*e^6/(e*x + d) - 6*B*a*b^5*d^6*e^7/(e*x + d) - A*b^6*d^6*e^7/(e
*x + d) + 15*B*a^2*b^4*d^5*e^8/(e*x + d) + 6*A*a*b^5*d^5*e^8/(e*x + d) - 2
0*B*a^3*b^3*d^4*e^9/(e*x + d) - 15*A*a^2*b^4*d^4*e^9/(e*x + d) + 15*B*a^4*
b^2*d^3*e^10/(e*x + d) + 20*A*a^3*b^3*d^3*e^10/(e*x + d) - 6*B*a^5*b*d^2*e
^11/(e*x + d) - 15*A*a^4*b^2*d^2*e^11/(e*x + d) + B*a^6*d*e^12/(e*x + d...

```

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 1228, normalized size of antiderivative = 4.43

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^2} dx = \text{Too large to display}$$

input

```
int(((A + B*x)*(a + b*x)^6)/(d + e*x)^2,x)
```


output

```

x^2*((d^2*((2*d*((A*b^6 + 6*B*a*b^5)/e^2 - (2*B*b^6*d)/e^3))/e - (3*a*b^4*
(2*A*b + 5*B*a))/e^2 + (B*b^6*d^2)/e^4))/(2*e^2) - (d*((2*d*((2*d*((A*b^6
+ 6*B*a*b^5)/e^2 - (2*B*b^6*d)/e^3))/e - (3*a*b^4*(2*A*b + 5*B*a))/e^2 + (
B*b^6*d^2)/e^4))/e - (d^2*((A*b^6 + 6*B*a*b^5)/e^2 - (2*B*b^6*d)/e^3))/e^2
+ (5*a^2*b^3*(3*A*b + 4*B*a))/e^2))/e + (5*a^3*b^2*(4*A*b + 3*B*a))/(2*e^
2)) - x^4*((d*((A*b^6 + 6*B*a*b^5)/e^2 - (2*B*b^6*d)/e^3))/(2*e) - (3*a*b^
4*(2*A*b + 5*B*a))/(4*e^2) + (B*b^6*d^2)/(4*e^4)) - x*((d^2*((2*d*((2*d*((
A*b^6 + 6*B*a*b^5)/e^2 - (2*B*b^6*d)/e^3))/e - (3*a*b^4*(2*A*b + 5*B*a))/e
^2 + (B*b^6*d^2)/e^4))/e - (d^2*((A*b^6 + 6*B*a*b^5)/e^2 - (2*B*b^6*d)/e^3
))/e^2 + (5*a^2*b^3*(3*A*b + 4*B*a))/e^2))/e^2 + (2*d*((d^2*((2*d*((A*b^6
+ 6*B*a*b^5)/e^2 - (2*B*b^6*d)/e^3))/e - (3*a*b^4*(2*A*b + 5*B*a))/e^2 + (
B*b^6*d^2)/e^4))/e^2 - (2*d*((2*d*((2*d*((A*b^6 + 6*B*a*b^5)/e^2 - (2*B*b^
6*d)/e^3))/e - (3*a*b^4*(2*A*b + 5*B*a))/e^2 + (B*b^6*d^2)/e^4))/e - (d^2*
((A*b^6 + 6*B*a*b^5)/e^2 - (2*B*b^6*d)/e^3))/e^2 + (5*a^2*b^3*(3*A*b + 4*B
*a))/e^2))/e + (5*a^3*b^2*(4*A*b + 3*B*a))/e^2))/e - (3*a^4*b*(5*A*b + 2*B
*a))/e^2) + x^3*((2*d*((2*d*((A*b^6 + 6*B*a*b^5)/e^2 - (2*B*b^6*d)/e^3))/e
- (3*a*b^4*(2*A*b + 5*B*a))/e^2 + (B*b^6*d^2)/e^4))/(3*e) - (d^2*((A*b^6
+ 6*B*a*b^5)/e^2 - (2*B*b^6*d)/e^3))/(3*e^2) + (5*a^2*b^3*(3*A*b + 4*B*a)
)/(3*e^2)) + x^5*((A*b^6 + 6*B*a*b^5)/(5*e^2) - (2*B*b^6*d)/(5*e^3)) + (log
(d + e*x)*(B*a^6*e^6 + 7*B*b^6*d^6 + 6*A*a^5*b*e^6 - 6*A*b^6*d^5*e + 30...

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 700, normalized size of antiderivative = 2.53

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^2} dx$$

$$= \frac{-126a b^6 d^2 e^6 x^5 + 84a b^6 d e^7 x^6 + 420 \log(ex + d) b^7 d^8 + 60a^7 e^8 x - 420b^7 d^7 ex - 210b^7 d^6 e^2 x^2 + 70b^7 d^5 e^3}{(d + ex)^2}$$

input

```
int((b*x+a)^6*(B*x+A)/(e*x+d)^2,x)
```

output

```
(420*log(d + e*x)*a**6*b*d**2*e**6 + 420*log(d + e*x)*a**6*b*d*e**7*x - 25
20*log(d + e*x)*a**5*b**2*d**3*e**5 - 2520*log(d + e*x)*a**5*b**2*d**2*e**
6*x + 6300*log(d + e*x)*a**4*b**3*d**4*e**4 + 6300*log(d + e*x)*a**4*b**3*
d**3*e**5*x - 8400*log(d + e*x)*a**3*b**4*d**5*e**3 - 8400*log(d + e*x)*a*
**3*b**4*d**4*e**4*x + 6300*log(d + e*x)*a**2*b**5*d**6*e**2 + 6300*log(d +
e*x)*a**2*b**5*d**5*e**3*x - 2520*log(d + e*x)*a*b**6*d**7*e - 2520*log(d
+ e*x)*a*b**6*d**6*e**2*x + 420*log(d + e*x)*b**7*d**8 + 420*log(d + e*x)
*b**7*d**7*e*x + 60*a**7*e**8*x - 420*a**6*b*d*e**7*x + 2520*a**5*b**2*d**
2*e**6*x + 1260*a**5*b**2*d*e**7*x**2 - 6300*a**4*b**3*d**3*e**5*x - 3150*
a**4*b**3*d**2*e**6*x**2 + 1050*a**4*b**3*d*e**7*x**3 + 8400*a**3*b**4*d**
4*e**4*x + 4200*a**3*b**4*d**3*e**5*x**2 - 1400*a**3*b**4*d**2*e**6*x**3 +
700*a**3*b**4*d*e**7*x**4 - 6300*a**2*b**5*d**5*e**3*x - 3150*a**2*b**5*d
**4*e**4*x**2 + 1050*a**2*b**5*d**3*e**5*x**3 - 525*a**2*b**5*d**2*e**6*x*
*4 + 315*a**2*b**5*d*e**7*x**5 + 2520*a*b**6*d**6*e**2*x + 1260*a*b**6*d**
5*e**3*x**2 - 420*a*b**6*d**4*e**4*x**3 + 210*a*b**6*d**3*e**5*x**4 - 126*
a*b**6*d**2*e**6*x**5 + 84*a*b**6*d*e**7*x**6 - 420*b**7*d**7*e*x - 210*b*
**7*d**6*e**2*x**2 + 70*b**7*d**5*e**3*x**3 - 35*b**7*d**4*e**4*x**4 + 21*b
**7*d**3*e**5*x**5 - 14*b**7*d**2*e**6*x**6 + 10*b**7*d*e**7*x**7)/(60*d*e
**8*(d + e*x))
```

3.52 $\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^3} dx$

Optimal result	526
Mathematica [A] (verified)	527
Rubi [A] (verified)	527
Maple [B] (verified)	529
Fricas [B] (verification not implemented)	530
Sympy [B] (verification not implemented)	531
Maxima [B] (verification not implemented)	532
Giac [B] (verification not implemented)	532
Mupad [B] (verification not implemented)	533
Reduce [B] (verification not implemented)	534

Optimal result

Integrand size = 20, antiderivative size = 276

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^3} dx = \frac{5b^2(bd-ae)^3(7bBd-4Abe-3aBe)x}{e^7} + \frac{(bd-ae)^6(Bd-Ae)}{2e^8(d+ex)^2} - \frac{(bd-ae)^5(7bBd-6Abe-aBe)}{e^8(d+ex)} - \frac{5b^3(bd-ae)^2(7bBd-3Abe-4aBe)(d+ex)^2}{2e^8} + \frac{b^4(bd-ae)(7bBd-2Abe-5aBe)(d+ex)^3}{e^8} - \frac{b^5(7bBd-Abe-6aBe)(d+ex)^4}{4e^8} + \frac{b^6B(d+ex)^5}{5e^8} - \frac{3b(bd-ae)^4(7bBd-5Abe-2aBe)\log(d+ex)}{e^8}$$

output

```
5*b^2*(-a*e+b*d)^3*(-4*A*b*e-3*B*a*e+7*B*b*d)*x/e^7+1/2*(-a*e+b*d)^6*(-A*e
+B*d)/e^8/(e*x+d)^2-(-a*e+b*d)^5*(-6*A*b*e-B*a*e+7*B*b*d)/e^8/(e*x+d)-5/2*
b^3*(-a*e+b*d)^2*(-3*A*b*e-4*B*a*e+7*B*b*d)*(e*x+d)^2/e^8+b^4*(-a*e+b*d)*(-
2*A*b*e-5*B*a*e+7*B*b*d)*(e*x+d)^3/e^8-1/4*b^5*(-A*b*e-6*B*a*e+7*B*b*d)*(-
e*x+d)^4/e^8+1/5*b^6*B*(e*x+d)^5/e^8-3*b*(-a*e+b*d)^4*(-5*A*b*e-2*B*a*e+7*
B*b*d)*ln(e*x+d)/e^8
```

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 352, normalized size of antiderivative = 1.28

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^3} dx$$

$$= \frac{-20b^2e(-15a^4Be^4 + 12ab^3d^2e(5Bd - 3Ae) - 5b^4d^3(3Bd - 2Ae) - 20a^3be^3(-3Bd + Ae) + 45a^2b^2de^2(-$$

input

```
Integrate[((a + b*x)^6*(A + B*x))/(d + e*x)^3,x]
```

output

```
(-20*b^2*e*(-15*a^4*B*e^4 + 12*a*b^3*d^2*e*(5*B*d - 3*A*e) - 5*b^4*d^3*(3*B*d - 2*A*e) - 20*a^3*b*e^3*(-3*B*d + A*e) + 45*a^2*b^2*d*e^2*(-2*B*d + A*e))*x + 10*b^3*e^2*(20*a^3*B*e^3 + 18*a*b^2*d*e*(2*B*d - A*e) + 15*a^2*b*e^2*(-3*B*d + A*e) + 2*b^3*d^2*(-5*B*d + 3*A*e))*x^2 - 20*b^4*e^3*(-5*a^2*B*e^2 - 2*a*b*e*(-3*B*d + A*e) + b^2*d*(-2*B*d + A*e))*x^3 + 5*b^5*e^4*(-3*b*B*d + A*b*e + 6*a*B*e)*x^4 + 4*b^6*B*e^5*x^5 + (10*(b*d - a*e)^6*(B*d - A*e))/(d + e*x)^2 - (20*(b*d - a*e)^5*(7*b*B*d - 6*A*b*e - a*B*e))/(d + e*x) - 60*b*(b*d - a*e)^4*(7*b*B*d - 5*A*b*e - 2*a*B*e)*Log[d + e*x] / (20*e^8)
```

Rubi [A] (verified)

Time = 0.83 (sec) , antiderivative size = 276, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^3} dx$$

↓ 86

$$\int \left(\frac{b^5(d + ex)^3(6aBe + Abe - 7bBd)}{e^7} - \frac{3b^4(d + ex)^2(bd - ae)(5aBe + 2Abe - 7bBd)}{e^7} + \frac{5b^3(d + ex)(bd - ae)^2}{e^7} \right) dx$$

$$\begin{aligned}
 & \downarrow \text{2009} \\
 & \frac{b^5(d+ex)^4(-6aBe - Abe + 7bBd)}{4e^8} + \frac{b^4(d+ex)^3(bd-ae)(-5aBe - 2Abe + 7bBd)}{e^8} - \\
 & \frac{5b^3(d+ex)^2(bd-ae)^2(-4aBe - 3Abe + 7bBd)}{2e^8} + \frac{5b^2x(bd-ae)^3(-3aBe - 4Abe + 7bBd)}{e^7} - \\
 & \frac{(bd-ae)^5(-aBe - 6Abe + 7bBd)}{e^8(d+ex)} + \frac{(bd-ae)^6(Bd-Ae)}{2e^8(d+ex)^2} - \\
 & \frac{3b(bd-ae)^4 \log(d+ex)(-2aBe - 5Abe + 7bBd)}{e^8} + \frac{b^6B(d+ex)^5}{5e^8}
 \end{aligned}$$

input `Int[((a + b*x)^6*(A + B*x))/(d + e*x)^3,x]`

output `(5*b^2*(b*d - a*e)^3*(7*b*B*d - 4*A*b*e - 3*a*B*e)*x)/e^7 + ((b*d - a*e)^6*(B*d - A*e))/(2*e^8*(d + e*x)^2) - ((b*d - a*e)^5*(7*b*B*d - 6*A*b*e - a*B*e))/(e^8*(d + e*x)) - (5*b^3*(b*d - a*e)^2*(7*b*B*d - 3*A*b*e - 4*a*B*e)*(d + e*x)^2)/(2*e^8) + (b^4*(b*d - a*e)*(7*b*B*d - 2*A*b*e - 5*a*B*e)*(d + e*x)^3)/e^8 - (b^5*(7*b*B*d - A*b*e - 6*a*B*e)*(d + e*x)^4)/(4*e^8) + (b^6*B*(d + e*x)^5)/(5*e^8) - (3*b*(b*d - a*e)^4*(7*b*B*d - 5*A*b*e - 2*a*B*e)*Log[d + e*x])/e^8`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 788 vs. $2(268) = 536$.

Time = 0.23 (sec) , antiderivative size = 789, normalized size of antiderivative = 2.86

method	result
norman	$\frac{b^2(20Aa^3be^4 - 30Aa^2b^2de^3 + 20Aab^3d^2e^2 - 5Ab^4d^3e + 15Ba^4e^4 - 40Ba^3bde^3 + 50Ba^2b^2d^2e^2 - 30Bab^3d^3e + 7Bb^4d^4)x^3 - a^6Ae^7 + 6Aa^5b^2e^6}{e^5}$
default	$b^2\left(-\frac{3}{4}Bb^4de^3x^4 + 2Aab^3e^4x^3 - Ab^4de^3x^3 + 5Ba^2b^2e^4x^3 + 2Bb^4d^2e^2x^3 + 3Ab^4d^2e^2x^2 + 10Ba^3be^4x^2 - 5Bb^4d^3ex^2 + 20Aa^3b^2e^4x - 10Aa^2b^3de^3x + 5Aab^4d^2e^2x - 5Ab^5d^3e + 5Ba^4e^4 - 40Bab^3de^3 + 50Ba^2b^2d^2e^2 - 30Bab^3d^3e + 7Bb^4d^4\right)x^3 - a^6Ae^7 + 6Aa^5b^2e^6$
risch	$-\frac{60b^3 \ln(ex+d)Aa^3d}{e^4} + \frac{90b^4 \ln(ex+d)Aa^2d^2}{e^5} - \frac{60b^5 \ln(ex+d)Aa^3d^3}{e^6} - \frac{45b^2 \ln(ex+d)Ba^4d}{e^4} + \frac{120b^3 \ln(ex+d)Ba^3d^2}{e^5}$
parallelrisc	Expression too large to display

input

```
int((b*x+a)^6*(B*x+A)/(e*x+d)^3,x,method=_RETURNVERBOSE)
```

output

```
(b^2*(20*A*a^3*b*e^4-30*A*a^2*b^2*d*e^3+20*A*a*b^3*d^2*e^2-5*A*b^4*d^3*e+15*B*a^4*e^4-40*B*a^3*b*d*e^3+50*B*a^2*b^2*d^2*e^2-30*B*a*b^3*d^3*e+7*B*b^4*d^4)/e^5*x^3-1/2*(A*a^6*e^7+6*A*a^5*b*d*e^6-45*A*a^4*b^2*d^2*e^5+180*A*a^3*b^3*d^3*e^4-270*A*a^2*b^4*d^4*e^3+180*A*a*b^5*d^5*e^2-45*A*b^6*d^6*e+B*a^6*d*e^6-18*B*a^5*b*d^2*e^5+135*B*a^4*b^2*d^3*e^4-360*B*a^3*b^3*d^4*e^3+450*B*a^2*b^4*d^5*e^2-270*B*a*b^5*d^6*e+63*B*b^6*d^7)/e^8-(6*A*a^5*b*e^6-30*A*a^4*b^2*d*e^5+120*A*a^3*b^3*d^2*e^4-180*A*a^2*b^4*d^3*e^3+120*A*a*b^5*d^4*e^2-30*A*b^6*d^5*e+B*a^6*e^6-12*B*a^5*b*d*e^5+90*B*a^4*b^2*d^2*e^4-240*B*a^3*b^3*d^3*e^3+300*B*a^2*b^4*d^4*e^2-180*B*a*b^5*d^5*e+42*B*b^6*d^6)/e^7*x+1/4*b^3*(30*A*a^2*b*e^3-20*A*a*b^2*d*e^2+5*A*b^3*d^2*e+40*B*a^3*e^3-50*B*a^2*b*d*e^2+30*B*a*b^2*d^2*e-7*B*b^3*d^3)/e^4*x^4+1/10*b^4*(20*A*a*b*e^2-5*A*b^2*d*e+50*B*a^2*e^2-30*B*a*b*d*e+7*B*b^2*d^2)/e^3*x^5+1/20*b^5*(5*A*b*e+30*B*a*e-7*B*b*d)/e^2*x^6+1/5*b^6*B/e*x^7)/(e*x+d)^2+3*b/e^8*(5*A*a^4*b*e^5-20*A*a^3*b^2*d*e^4+30*A*a^2*b^3*d^2*e^3-20*A*a*b^4*d^3*e^2+5*A*b^5*d^4*e+2*B*a^5*e^5-15*B*a^4*b*d*e^4+40*B*a^3*b^2*d^2*e^3-50*B*a^2*b^3*d^3*e^2+30*B*a*b^4*d^4*e-7*B*b^5*d^5)*ln(e*x+d)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1177 vs. $2(268) = 536$.

Time = 0.08 (sec) , antiderivative size = 1177, normalized size of antiderivative = 4.26

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^3} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^3,x, algorithm="fricas")`

output

```
1/20*(4*B*b^6*e^7*x^7 - 130*B*b^6*d^7 - 10*A*a^6*e^7 + 110*(6*B*a*b^5 + A*
b^6)*d^6*e - 270*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 + 350*(4*B*a^3*b^3 + 3*
A*a^2*b^4)*d^4*e^3 - 250*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 + 90*(2*B*a^5
*b + 5*A*a^4*b^2)*d^2*e^5 - 10*(B*a^6 + 6*A*a^5*b)*d*e^6 - (7*B*b^6*d*e^6
- 5*(6*B*a*b^5 + A*b^6)*e^7)*x^6 + 2*(7*B*b^6*d^2*e^5 - 5*(6*B*a*b^5 + A*b
^6)*d*e^6 + 10*(5*B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^5 - 5*(7*B*b^6*d^3*e^4 - 5
*(6*B*a*b^5 + A*b^6)*d^2*e^5 + 10*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6 - 10*(4*
B*a^3*b^3 + 3*A*a^2*b^4)*e^7)*x^4 + 20*(7*B*b^6*d^4*e^3 - 5*(6*B*a*b^5 + A
*b^6)*d^3*e^4 + 10*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 - 10*(4*B*a^3*b^3 + 3
*A*a^2*b^4)*d*e^6 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^3 + 10*(50*B*b^6*
d^5*e^2 - 34*(6*B*a*b^5 + A*b^6)*d^4*e^3 + 63*(5*B*a^2*b^4 + 2*A*a*b^5)*d^
3*e^4 - 55*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + 20*(3*B*a^4*b^2 + 4*A*a^3
*b^3)*d*e^6)*x^2 + 20*(8*B*b^6*d^6*e - 4*(6*B*a*b^5 + A*b^6)*d^5*e^2 + 3*(
5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^3 + 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 -
10*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^5 + 6*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^
6 - (B*a^6 + 6*A*a^5*b)*e^7)*x - 60*(7*B*b^6*d^7 - 5*(6*B*a*b^5 + A*b^6)*d
^6*e + 10*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 - 10*(4*B*a^3*b^3 + 3*A*a^2*b^
4)*d^4*e^3 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 - (2*B*a^5*b + 5*A*a^4*
b^2)*d^2*e^5 + (7*B*b^6*d^5*e^2 - 5*(6*B*a*b^5 + A*b^6)*d^4*e^3 + 10*(5*B*
a^2*b^4 + 2*A*a*b^5)*d^3*e^4 - 10*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + ...
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 821 vs. $2(280) = 560$.

Time = 6.15 (sec) , antiderivative size = 821, normalized size of antiderivative = 2.97

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^3} dx = \frac{Bb^6x^5}{5e^3} + \frac{3b(ae-bd)^4 \cdot (5Abe+2Bae-7Bbd) \log(d+ex)}{e^8} \\ + x^4 \left(\frac{Ab^6}{4e^3} + \frac{3Bab^5}{2e^3} - \frac{3Bb^6d}{4e^4} \right) + x^3 \cdot \left(\frac{2Aab^5}{e^3} - \frac{Ab^6d}{e^4} + \frac{5Ba^2b^4}{e^3} - \frac{6Bab^5d}{e^4} + \frac{2Bb^6d^2}{e^5} \right) \\ + x^2 \cdot \left(\frac{15Aa^2b^4}{2e^3} - \frac{9Aab^5d}{e^4} + \frac{3Ab^6d^2}{e^5} + \frac{10Ba^3b^3}{e^3} - \frac{45Ba^2b^4d}{2e^4} + \frac{18Bab^5d^2}{e^5} - \frac{5Bb^6d^3}{e^6} \right) \\ + x \left(\frac{20Aa^3b^3}{e^3} - \frac{45Aa^2b^4d}{e^4} + \frac{36Aab^5d^2}{e^5} - \frac{10Ab^6d^3}{e^6} + \frac{15Ba^4b^2}{e^3} - \frac{60Ba^3b^3d}{e^4} \right. \\ \left. + \frac{90Ba^2b^4d^2}{e^5} - \frac{60Bab^5d^3}{e^6} + \frac{15Bb^6d^4}{e^7} \right) \\ + \frac{-Aa^6e^7 - 6Aa^5bde^6 + 45Aa^4b^2d^2e^5 - 100Aa^3b^3d^3e^4 + 105Aa^2b^4d^4e^3 - 54Aab^5d^5e^2 + 11Ab^6d^6e - Ba^6d^7}{e^8}$$

input `integrate((b*x+a)**6*(B*x+A)/(e*x+d)**3,x)`

output

```
B**6*x**5/(5*e**3) + 3*b*(a*e - b*d)**4*(5*A*b*e + 2*B*a*e - 7*B*b*d)*log(d + e*x)/e**8 + x**4*(A*b**6/(4*e**3) + 3*B*a*b**5/(2*e**3) - 3*B*b**6*d/(4*e**4)) + x**3*(2*A*a*b**5/e**3 - A*b**6*d/e**4 + 5*B*a**2*b**4/e**3 - 6*B*a*b**5*d/e**4 + 2*B*b**6*d**2/e**5) + x**2*(15*A*a**2*b**4/(2*e**3) - 9*A*a*b**5*d/e**4 + 3*A*b**6*d**2/e**5 + 10*B*a**3*b**3/e**3 - 45*B*a**2*b**4*d/(2*e**4) + 18*B*a*b**5*d**2/e**5 - 5*B*b**6*d**3/e**6) + x*(20*A*a**3*b**3/e**3 - 45*A*a**2*b**4*d/e**4 + 36*A*a*b**5*d**2/e**5 - 10*A*b**6*d**3/e**6 + 15*B*a**4*b**2/e**3 - 60*B*a**3*b**3*d/e**4 + 90*B*a**2*b**4*d**2/e**5 - 60*B*a*b**5*d**3/e**6 + 15*B*b**6*d**4/e**7) + (-A*a**6*e**7 - 6*A*a**5*b*d*e**6 + 45*A*a**4*b**2*d**2*e**5 - 100*A*a**3*b**3*d**3*e**4 + 105*A*a**2*b**4*d**4*e**3 - 54*A*a*b**5*d**5*e**2 + 11*A*b**6*d**6*e - B*a**6*d**7 + 18*B*a**5*b*d**2*e**5 - 75*B*a**4*b**2*d**3*e**4 + 140*B*a**3*b**3*d**4*e**3 - 135*B*a**2*b**4*d**5*e**2 + 66*B*a*b**5*d**6*e - 13*B*b**6*d**7 + x*(-12*A*a**5*b*e**7 + 60*A*a**4*b**2*d*e**6 - 120*A*a**3*b**3*d**2*e**5 + 120*A*a**2*b**4*d**3*e**4 - 60*A*a*b**5*d**4*e**3 + 12*A*b**6*d**5*e**2 - 2*B*a**6*e**7 + 24*B*a**5*b*d*e**6 - 90*B*a**4*b**2*d**2*e**5 + 160*B*a**3*b**3*d**3*e**4 - 150*B*a**2*b**4*d**4*e**3 + 72*B*a*b**5*d**5*e**2 - 14*B*b**6*d**6*e))/(2*d**2*e**8 + 4*d*e**9*x + 2*e**10*x**2)
```


Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 779 vs. $2(268) = 536$.

Time = 0.05 (sec) , antiderivative size = 779, normalized size of antiderivative = 2.82

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^3} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^3,x, algorithm="maxima")`

output

```
-1/2*(13*B*b^6*d^7 + A*a^6*e^7 - 11*(6*B*a*b^5 + A*b^6)*d^6*e + 27*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 - 35*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4*e^3 + 25*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 - 9*(2*B*a^5*b + 5*A*a^4*b^2)*d^2*e^5 + (B*a^6 + 6*A*a^5*b)*d*e^6 + 2*(7*B*b^6*d^6*e - 6*(6*B*a*b^5 + A*b^6)*d^5*e^2 + 15*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^3 - 20*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 + 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^5 - 6*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^6 + (B*a^6 + 6*A*a^5*b)*e^7)*x)/(e^10*x^2 + 2*d*e^9*x + d^2*e^8) + 1/20*(4*B*b^6*e^4*x^5 - 5*(3*B*b^6*d*e^3 - (6*B*a*b^5 + A*b^6)*e^4)*x^4 + 20*(2*B*b^6*d^2*e^2 - (6*B*a*b^5 + A*b^6)*d*e^3 + (5*B*a^2*b^4 + 2*A*a*b^5)*e^4)*x^3 - 10*(10*B*b^6*d^3*e - 6*(6*B*a*b^5 + A*b^6)*d^2*e^2 + 9*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^3 - 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^4)*x^2 + 20*(15*B*b^6*d^4 - 10*(6*B*a*b^5 + A*b^6)*d^3*e + 18*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^2 - 15*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^3 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^4)*x)/e^7 - 3*(7*B*b^6*d^5 - 5*(6*B*a*b^5 + A*b^6)*d^4*e + 10*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*e^2 - 10*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^3 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^4 - (2*B*a^5*b + 5*A*a^4*b^2)*e^5)*log(e*x + d)/e^8
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 862 vs. $2(268) = 536$.

Time = 0.12 (sec) , antiderivative size = 862, normalized size of antiderivative = 3.12

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^3} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^3,x, algorithm="giac")`

output

```

-3*(7*B*b^6*d^5 - 30*B*a*b^5*d^4*e - 5*A*b^6*d^4*e + 50*B*a^2*b^4*d^3*e^2
+ 20*A*a*b^5*d^3*e^2 - 40*B*a^3*b^3*d^2*e^3 - 30*A*a^2*b^4*d^2*e^3 + 15*B*
a^4*b^2*d*e^4 + 20*A*a^3*b^3*d*e^4 - 2*B*a^5*b*e^5 - 5*A*a^4*b^2*e^5)*log(
abs(e*x + d))/e^8 - 1/2*(13*B*b^6*d^7 - 66*B*a*b^5*d^6*e - 11*A*b^6*d^6*e
+ 135*B*a^2*b^4*d^5*e^2 + 54*A*a*b^5*d^5*e^2 - 140*B*a^3*b^3*d^4*e^3 - 105
*A*a^2*b^4*d^4*e^3 + 75*B*a^4*b^2*d^3*e^4 + 100*A*a^3*b^3*d^3*e^4 - 18*B*a
^5*b*d^2*e^5 - 45*A*a^4*b^2*d^2*e^5 + B*a^6*d*e^6 + 6*A*a^5*b*d*e^6 + A*a^
6*e^7 + 2*(7*B*b^6*d^6*e - 36*B*a*b^5*d^5*e^2 - 6*A*b^6*d^5*e^2 + 75*B*a^2
*b^4*d^4*e^3 + 30*A*a*b^5*d^4*e^3 - 80*B*a^3*b^3*d^3*e^4 - 60*A*a^2*b^4*d^
3*e^4 + 45*B*a^4*b^2*d^2*e^5 + 60*A*a^3*b^3*d^2*e^5 - 12*B*a^5*b*d*e^6 - 3
0*A*a^4*b^2*d*e^6 + B*a^6*e^7 + 6*A*a^5*b*e^7)*x)/((e*x + d)^2*e^8) + 1/20
*(4*B*b^6*e^12*x^5 - 15*B*b^6*d*e^11*x^4 + 30*B*a*b^5*e^12*x^4 + 5*A*b^6*e
^12*x^4 + 40*B*b^6*d^2*e^10*x^3 - 120*B*a*b^5*d*e^11*x^3 - 20*A*b^6*d*e^11
*x^3 + 100*B*a^2*b^4*e^12*x^3 + 40*A*a*b^5*e^12*x^3 - 100*B*b^6*d^3*e^9*x^
2 + 360*B*a*b^5*d^2*e^10*x^2 + 60*A*b^6*d^2*e^10*x^2 - 450*B*a^2*b^4*d*e^1
1*x^2 - 180*A*a*b^5*d*e^11*x^2 + 200*B*a^3*b^3*e^12*x^2 + 150*A*a^2*b^4*e^
12*x^2 + 300*B*b^6*d^4*e^8*x - 1200*B*a*b^5*d^3*e^9*x - 200*A*b^6*d^3*e^9*x
+ 1800*B*a^2*b^4*d^2*e^10*x + 720*A*a*b^5*d^2*e^10*x - 1200*B*a^3*b^3*d*
e^11*x - 900*A*a^2*b^4*d*e^11*x + 300*B*a^4*b^2*e^12*x + 400*A*a^3*b^3*e^1
2*x)/e^15

```

Mupad [B] (verification not implemented)

Time = 0.97 (sec) , antiderivative size = 1053, normalized size of antiderivative = 3.82

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^3} dx = \text{Too large to display}$$

input

```
int(((A + B*x)*(a + b*x)^6)/(d + e*x)^3,x)
```

output

```
x*((3*d*((3*d^2*((A*b^6 + 6*B*a*b^5)/e^3 - (3*B*b^6*d)/e^4))/e^2 - (3*d*((3*d*((A*b^6 + 6*B*a*b^5)/e^3 - (3*B*b^6*d)/e^4))/e - (3*a*b^4*(2*A*b + 5*B*a))/e^3 + (3*B*b^6*d^2)/e^5))/e - (5*a^2*b^3*(3*A*b + 4*B*a))/e^3 + (B*b^6*d^3)/e^6))/e - (d^3*((A*b^6 + 6*B*a*b^5)/e^3 - (3*B*b^6*d)/e^4))/e^3 + (3*d^2*((3*d*((A*b^6 + 6*B*a*b^5)/e^3 - (3*B*b^6*d)/e^4))/e - (3*a*b^4*(2*A*b + 5*B*a))/e^3 + (3*B*b^6*d^2)/e^5))/e^2 + (5*a^3*b^2*(4*A*b + 3*B*a))/e^3 - x^3*((d*((A*b^6 + 6*B*a*b^5)/e^3 - (3*B*b^6*d)/e^4))/e - (a*b^4*(2*A*b + 5*B*a))/e^3 + (B*b^6*d^2)/e^5) - x^2*((3*d^2*((A*b^6 + 6*B*a*b^5)/e^3 - (3*B*b^6*d)/e^4))/(2*e^2) - (3*d*((3*d*((A*b^6 + 6*B*a*b^5)/e^3 - (3*B*b^6*d)/e^4))/e - (3*a*b^4*(2*A*b + 5*B*a))/e^3 + (3*B*b^6*d^2)/e^5))/(2*e) - (5*a^2*b^3*(3*A*b + 4*B*a))/(2*e^3) + (B*b^6*d^3)/(2*e^6) - ((A*a^6*e^7 + 13*B*b^6*d^7 - 11*A*b^6*d^6*e + B*a^6*d*e^6 + 54*A*a*b^5*d^5*e^2 - 18*B*a^5*b*d^2*e^5 - 105*A*a^2*b^4*d^4*e^3 + 100*A*a^3*b^3*d^3*e^4 - 45*A*a^4*b^2*d^2*e^5 + 135*B*a^2*b^4*d^5*e^2 - 140*B*a^3*b^3*d^4*e^3 + 75*B*a^4*b^2*d^3*e^4 + 6*A*a^5*b*d*e^6 - 66*B*a*b^5*d^6*e)/(2*e) + x*(B*a^6*e^6 + 7*B*b^6*d^6 + 6*A*a^5*b*e^6 - 6*A*b^6*d^5*e + 30*A*a*b^5*d^4*e^2 - 30*A*a^4*b^2*d*e^5 - 60*A*a^2*b^4*d^3*e^3 + 60*A*a^3*b^3*d^2*e^4 + 75*B*a^2*b^4*d^4*e^2 - 80*B*a^3*b^3*d^3*e^3 + 45*B*a^4*b^2*d^2*e^4 - 36*B*a*b^5*d^5*e - 12*B*a^5*b*d*e^5))/(d^2*e^7 + e^9*x^2 + 2*d*e^8*x) + x^4*((A*b^6 + 6*B*a*b^5)/(4*e^3) - (3*B*b^6*d)/(4*e^4)) + (log(d + e*x))*(6*B*a^5*b*e^5 - 21*B*b...
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 799, normalized size of antiderivative = 2.89

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^3} dx = \text{Too large to display}$$

input

```
int((b*x+a)^6*(B*x+A)/(e*x+d)^3,x)
```

output

```
(420*log(d + e*x)*a**5*b**2*d**3*e**5 + 840*log(d + e*x)*a**5*b**2*d**2*e*
*6*x + 420*log(d + e*x)*a**5*b**2*d*e**7*x**2 - 2100*log(d + e*x)*a**4*b**
3*d**4*e**4 - 4200*log(d + e*x)*a**4*b**3*d**3*e**5*x - 2100*log(d + e*x)*
a**4*b**3*d**2*e**6*x**2 + 4200*log(d + e*x)*a**3*b**4*d**5*e**3 + 8400*lo
g(d + e*x)*a**3*b**4*d**4*e**4*x + 4200*log(d + e*x)*a**3*b**4*d**3*e**5*x
**2 - 4200*log(d + e*x)*a**2*b**5*d**6*e**2 - 8400*log(d + e*x)*a**2*b**5*
d**5*e**3*x - 4200*log(d + e*x)*a**2*b**5*d**4*e**4*x**2 + 2100*log(d + e*
x)*a*b**6*d**7*e + 4200*log(d + e*x)*a*b**6*d**6*e**2*x + 2100*log(d + e*x
)*a*b**6*d**5*e**3*x**2 - 420*log(d + e*x)*b**7*d**8 - 840*log(d + e*x)*b*
*7*d**7*e*x - 420*log(d + e*x)*b**7*d**6*e**2*x**2 - 10*a**7*d*e**7 + 70*a
**6*b*e**8*x**2 + 210*a**5*b**2*d**3*e**5 - 420*a**5*b**2*d*e**7*x**2 - 10
50*a**4*b**3*d**4*e**4 + 2100*a**4*b**3*d**2*e**6*x**2 + 700*a**4*b**3*d*e
**7*x**3 + 2100*a**3*b**4*d**5*e**3 - 4200*a**3*b**4*d**3*e**5*x**2 - 1400
*a**3*b**4*d**2*e**6*x**3 + 350*a**3*b**4*d*e**7*x**4 - 2100*a**2*b**5*d**
6*e**2 + 4200*a**2*b**5*d**4*e**4*x**2 + 1400*a**2*b**5*d**3*e**5*x**3 - 3
50*a**2*b**5*d**2*e**6*x**4 + 140*a**2*b**5*d*e**7*x**5 + 1050*a*b**6*d**7
*e - 2100*a*b**6*d**5*e**3*x**2 - 700*a*b**6*d**4*e**4*x**3 + 175*a*b**6*d
**3*e**5*x**4 - 70*a*b**6*d**2*e**6*x**5 + 35*a*b**6*d*e**7*x**6 - 210*b**
7*d**8 + 420*b**7*d**6*e**2*x**2 + 140*b**7*d**5*e**3*x**3 - 35*b**7*d**4*
e**4*x**4 + 14*b**7*d**3*e**5*x**5 - 7*b**7*d**2*e**6*x**6 + 4*b**7*d*e...
```

3.53 $\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^4} dx$

Optimal result	536
Mathematica [A] (verified)	537
Rubi [A] (verified)	537
Maple [B] (verified)	539
Fricas [B] (verification not implemented)	540
Sympy [B] (verification not implemented)	541
Maxima [B] (verification not implemented)	542
Giac [B] (verification not implemented)	542
Mupad [B] (verification not implemented)	544
Reduce [B] (verification not implemented)	545

Optimal result

Integrand size = 20, antiderivative size = 279

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^4} dx = -\frac{5b^3(bd-ae)^2(7bBd-3Abe-4aBe)x}{e^7} + \frac{(bd-ae)^6(Bd-Ae)}{3e^8(d+ex)^3} - \frac{(bd-ae)^5(7bBd-6Abe-aBe)}{2e^8(d+ex)^2} + \frac{3b(bd-ae)^4(7bBd-5Abe-2aBe)}{e^8(d+ex)} + \frac{3b^4(bd-ae)(7bBd-2Abe-5aBe)(d+ex)^2}{2e^8} - \frac{b^5(7bBd-Abe-6aBe)(d+ex)^3}{3e^8} + \frac{b^6B(d+ex)^4}{4e^8} + \frac{5b^2(bd-ae)^3(7bBd-4Abe-3aBe)\log(d+ex)}{e^8}$$

output

```
-5*b^3*(-a*e+b*d)^2*(-3*A*b*e-4*B*a*e+7*B*b*d)*x/e^7+1/3*(-a*e+b*d)^6*(-A*
e+B*d)/e^8/(e*x+d)^3-1/2*(-a*e+b*d)^5*(-6*A*b*e-B*a*e+7*B*b*d)/e^8/(e*x+d)
^2+3*b*(-a*e+b*d)^4*(-5*A*b*e-2*B*a*e+7*B*b*d)/e^8/(e*x+d)+3/2*b^4*(-a*e+b
*d)*(-2*A*b*e-5*B*a*e+7*B*b*d)*(e*x+d)^2/e^8-1/3*b^5*(-A*b*e-6*B*a*e+7*B*b
*d)*(e*x+d)^3/e^8+1/4*b^6*B*(e*x+d)^4/e^8+5*b^2*(-a*e+b*d)^3*(-4*A*b*e-3*B
*a*e+7*B*b*d)*ln(e*x+d)/e^8
```

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 297, normalized size of antiderivative = 1.06

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^4} dx$$

$$= \frac{12b^3e(20a^3Be^3 + 12ab^2de(5Bd - 2Ae) + 15a^2be^2(-4Bd + Ae) + 10b^3d^2(-2Bd + Ae))x - 6b^4e^2(-15a^2B^2e^2 - 6a^2B^2d + A^2e) + 10b^3d^2(-2B^2d + A^2e)x - 6b^4e^2(-15a^2B^2e^2 - 6a^2B^2d + A^2e) + 2b^2d^2(-5B^2d + 2A^2e)x^2 + 4b^5e^3(-4bB^2d + A^2be + 6aB^2e)x^3 + 3b^6B^2e^4x^4 + (4(bd - ae)^6(Bd - Ae))/(d + ex)^3 - (6(bd - ae)^5(7bB^2d - 6A^2be - aB^2e))/(d + ex)^2 + (36b(bd - ae)^4(7bB^2d - 5A^2be - 2aB^2e))/(d + ex) + 60b^2(bd - ae)^3(7bB^2d - 4A^2be - 3aB^2e)*\text{Log}[d + ex]}{(12e^8)}$$

input

```
Integrate[((a + b*x)^6*(A + B*x))/(d + e*x)^4,x]
```

output

```
(12*b^3*e*(20*a^3*B*e^3 + 12*a*b^2*d*e*(5*B*d - 2*A*e) + 15*a^2*b*e^2*(-4*B*d + A*e) + 10*b^3*d^2*(-2*B*d + A*e))*x - 6*b^4*e^2*(-15*a^2*B*e^2 - 6*a^2*B^2*d + A^2*e) + 10*b^3*d^2*(-2*B*d + A*e)*x - 6*b^4*e^2*(-15*a^2*B*e^2 - 6*a^2*B^2*d + A^2*e) + 2*b^2*d^2*(-5*B*d + 2*A*e))*x^2 + 4*b^5*e^3*(-4*b*B*d + A*b*e + 6*a*B*e)*x^3 + 3*b^6*B*e^4*x^4 + (4*(b*d - a*e)^6*(B*d - A*e))/(d + e*x)^3 - (6*(b*d - a*e)^5*(7*b*B*d - 6*A*b*e - a*B*e))/(d + e*x)^2 + (36*b*(b*d - a*e)^4*(7*b*B*d - 5*A*b*e - 2*a*B*e))/(d + e*x) + 60*b^2*(b*d - a*e)^3*(7*b*B*d - 4*A*b*e - 3*a*B*e)*Log[d + e*x]]/(12*e^8)
```

Rubi [A] (verified)

Time = 0.82 (sec) , antiderivative size = 279, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^4} dx$$

$$\downarrow 86$$

$$\int \left(\frac{b^5(d+ex)^2(6aBe + Abe - 7bBd)}{e^7} - \frac{3b^4(d+ex)(bd - ae)(5aBe + 2Abe - 7bBd)}{e^7} + \frac{5b^3(bd - ae)^2(4aBe + 2Abe - 7bBd)}{e^7} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
& -\frac{b^5(d+ex)^3(-6aBe - Abe + 7bBd)}{3e^8} + \frac{3b^4(d+ex)^2(bd-ae)(-5aBe - 2Abe + 7bBd)}{2e^8} - \\
& \frac{5b^3x(bd-ae)^2(-4aBe - 3Abe + 7bBd)}{e^7} + \\
& \frac{5b^2(bd-ae)^3 \log(d+ex)(-3aBe - 4Abe + 7bBd)}{e^8} + \frac{3b(bd-ae)^4(-2aBe - 5Abe + 7bBd)}{e^8(d+ex)} - \\
& \frac{(bd-ae)^5(-aBe - 6Abe + 7bBd)}{2e^8(d+ex)^2} + \frac{(bd-ae)^6(Bd-Ae)}{3e^8(d+ex)^3} + \frac{b^6B(d+ex)^4}{4e^8}
\end{aligned}$$

input `Int[((a + b*x)^6*(A + B*x))/(d + e*x)^4,x]`

output `(-5*b^3*(b*d - a*e)^2*(7*b*B*d - 3*A*b*e - 4*a*B*e)*x)/e^7 + ((b*d - a*e)^6*(B*d - A*e))/(3*e^8*(d + e*x)^3) - ((b*d - a*e)^5*(7*b*B*d - 6*A*b*e - a*B*e))/(2*e^8*(d + e*x)^2) + (3*b*(b*d - a*e)^4*(7*b*B*d - 5*A*b*e - 2*a*B*e))/(e^8*(d + e*x)) + (3*b^4*(b*d - a*e)*(7*b*B*d - 2*A*b*e - 5*a*B*e)*(d + e*x)^2)/(2*e^8) - (b^5*(7*b*B*d - A*b*e - 6*a*B*e)*(d + e*x)^3)/(3*e^8) + (b^6*B*(d + e*x)^4)/(4*e^8) + (5*b^2*(b*d - a*e)^3*(7*b*B*d - 4*A*b*e - 3*a*B*e)*Log[d + e*x])/e^8`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 794 vs. 2(269) = 538.

Time = 0.24 (sec) , antiderivative size = 795, normalized size of antiderivative = 2.85

method	result
norman	$\frac{-2a^6 A e^7 + 6A a^5 b d e^6 + 30A a^4 b^2 d^2 e^5 - 220A a^3 b^3 d^3 e^4 + 660A a^2 b^4 d^4 e^3 - 660A a b^5 d^5 e^2 + 220A b^6 d^6 e + B a^6 d e^6 + 12B a^5 b d^2 e^5 - 165B a^4 b^2 d^3 e^4 + 660B a^3 b^3 d^4 e^3 - 1650B a^2 b^4 d^5 e^2 + 1320B a b^5 d^6 e - 385B b^6 d^7}{6e^8}$
default	$\frac{b^3 \left(\frac{1}{4} b^3 B x^4 e^3 + \frac{1}{3} A b^3 e^3 x^3 + 2B a b^2 e^3 x^3 - \frac{4}{3} B b^3 d e^2 x^3 + 3A a b^2 e^3 x^2 - 2A b^3 d e^2 x^2 + \frac{15}{2} B a^2 b e^3 x^2 - 12B a b^2 d e^2 x^2 + 5B b^3 d^2 e^2 x - 15A a^4 b^2 e^6 + 60A a^3 b^3 d e^5 - 90A a^2 b^4 d^2 e^4 + 60A a b^5 d^3 e^3 - 15A b^6 d^4 e^2 - 6B a^5 b e^6 + 45B a^4 b^2 d e^5 - 120B a^3 b^3 d^2 e^4 + 150B a^2 b^4 d^3 e^3 - 1650B a b^5 d^4 e^2 + 1320B a b^5 d^6 e - 385B b^6 d^7 \right)}{e^7}$
risch	$(-15A a^4 b^2 e^6 + 60A a^3 b^3 d e^5 - 90A a^2 b^4 d^2 e^4 + 60A a b^5 d^3 e^3 - 15A b^6 d^4 e^2 - 6B a^5 b e^6 + 45B a^4 b^2 d e^5 - 120B a^3 b^3 d^2 e^4 + 150B a^2 b^4 d^3 e^3 - 1650B a b^5 d^4 e^2 + 1320B a b^5 d^6 e - 385B b^6 d^7)$
parallelrisc	Expression too large to display

input

```
int((b*x+a)^6*(B*x+A)/(e*x+d)^4,x,method=_RETURNVERBOSE)
```

output

```
(-1/6*(2*A*a^6*e^7+6*A*a^5*b*d*e^6+30*A*a^4*b^2*d^2*e^5-220*A*a^3*b^3*d^3*e^4+660*A*a^2*b^4*d^4*e^3-660*A*a*b^5*d^5*e^2+220*A*b^6*d^6*e+B*a^6*d*e^6+12*B*a^5*b*d^2*e^5-165*B*a^4*b^2*d^3*e^4+880*B*a^3*b^3*d^4*e^3-1650*B*a^2*b^4*d^5*e^2+1320*B*a*b^5*d^6*e-385*B*b^6*d^7)/e^8-3*(5*A*a^4*b^2*e^5-20*A*a^3*b^3*d*e^4+60*A*a^2*b^4*d^2*e^3-60*A*a*b^5*d^3*e^2+20*A*b^6*d^4*e+2*B*a^5*b*e^5-15*B*a^4*b^2*d*e^4+80*B*a^3*b^3*d^2*e^3-150*B*a^2*b^4*d^3*e^2+120*B*a*b^5*d^4*e-35*B*b^6*d^5)/e^6*x^2-1/2*(6*A*a^5*b*e^6+30*A*a^4*b^2*d*e^5-180*A*a^3*b^3*d^2*e^4+540*A*a^2*b^4*d^3*e^3-540*A*a*b^5*d^4*e^2+180*A*b^6*d^5*e+B*a^6*e^6+12*B*a^5*b*d*e^5-135*B*a^4*b^2*d^2*e^4+720*B*a^3*b^3*d^3*e^3-1350*B*a^2*b^4*d^4*e^2+1080*B*a*b^5*d^5*e-315*B*b^6*d^6)/e^7*x+5/4*b^3*(12*A*a^2*b*e^3-12*A*a*b^2*d*e^2+4*A*b^3*d^2*e+16*B*a^3*e^3-30*B*a^2*b*d*e^2+24*B*a*b^2*d^2*e-7*B*b^3*d^3)/e^4*x^4+1/4*b^4*(12*A*a*b*e^2-4*A*b^2*d*e+30*B*a^2*e^2-24*B*a*b*d*e+7*B*b^2*d^2)/e^3*x^5+1/12*b^5*(4*A*b*e+24*B*a*e-7*B*b*d)/e^2*x^6+1/4*b^6*B/e*x^7)/(e*x+d)^3+5*b^2/e^8*(4*A*a^3*b*e^4-12*A*a^2*b^2*d*e^3+12*A*a*b^3*d^2*e^2-4*A*b^4*d^3*e+3*B*a^4*e^4-16*B*a^3*b*d*e^3+30*B*a^2*b^2*d^2*e^2-24*B*a*b^3*d^3*e+7*B*b^4*d^4)*ln(e*x+d)
```


Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1225 vs. $2(269) = 538$.

Time = 0.10 (sec) , antiderivative size = 1225, normalized size of antiderivative = 4.39

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^4} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^4,x, algorithm="fricas")`

output

```
1/12*(3*B*b^6*e^7*x^7 + 214*B*b^6*d^7 - 4*A*a^6*e^7 - 148*(6*B*a*b^5 + A*b^6)*d^6*e + 282*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 - 260*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4*e^3 + 110*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 - 12*(2*B*a^5*b + 5*A*a^4*b^2)*d^2*e^5 - 2*(B*a^6 + 6*A*a^5*b)*d*e^6 - (7*B*b^6*d*e^6 - 4*(6*B*a*b^5 + A*b^6)*e^7)*x^6 + 3*(7*B*b^6*d^2*e^5 - 4*(6*B*a*b^5 + A*b^6)*d*e^6 + 6*(5*B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^5 - 15*(7*B*b^6*d^3*e^4 - 4*(6*B*a*b^5 + A*b^6)*d^2*e^5 + 6*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6 - 4*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^7)*x^4 - 2*(278*B*b^6*d^4*e^3 - 146*(6*B*a*b^5 + A*b^6)*d^3*e^4 + 189*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 - 90*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^6)*x^3 - 6*(68*B*b^6*d^5*e^2 - 26*(6*B*a*b^5 + A*b^6)*d^4*e^3 + 9*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*e^4 + 30*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 - 30*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^6 + 6*(2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 6*(37*B*b^6*d^6*e - 34*(6*B*a*b^5 + A*b^6)*d^5*e^2 + 81*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^3 - 90*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 + 45*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^5 - 6*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^6 - (B*a^6 + 6*A*a^5*b)*e^7)*x + 60*(7*B*b^6*d^7 - 4*(6*B*a*b^5 + A*b^6)*d^6*e + 6*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 - 4*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4*e^3 + (3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 + (7*B*b^6*d^4*e^3 - 4*(6*B*a*b^5 + A*b^6)*d^3*e^4 + 6*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 - 4*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^6 + (3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^3 + 3*(7*...
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 867 vs. $2(284) = 568$.

Time = 29.91 (sec) , antiderivative size = 867, normalized size of antiderivative = 3.11

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^4} dx = \frac{Bb^6x^4}{4e^4} + \frac{5b^2(ae-bd)^3 \cdot (4Abe+3Bae-7Bbd) \log(d+ex)}{e^8} \\ + x^3 \left(\frac{Ab^6}{3e^4} + \frac{2Bab^5}{e^4} - \frac{4Bb^6d}{3e^5} \right) + x^2 \cdot \left(\frac{3Aab^5}{e^4} - \frac{2Ab^6d}{e^5} + \frac{15Ba^2b^4}{2e^4} - \frac{12Bab^5d}{e^5} + \frac{5Bb^6d^2}{e^6} \right) \\ + x \left(\frac{15Aa^2b^4}{e^4} - \frac{24Aab^5d}{e^5} + \frac{10Ab^6d^2}{e^6} + \frac{20Ba^3b^3}{e^4} - \frac{60Ba^2b^4d}{e^5} + \frac{60Bab^5d^2}{e^6} - \frac{20Bb^6d^3}{e^7} \right) \\ + \frac{-2Aa^6e^7 - 6Aa^5bde^6 - 30Aa^4b^2d^2e^5 + 220Aa^3b^3d^3e^4 - 390Aa^2b^4d^4e^3 + 282Aab^5d^5e^2 - 74Ab^6d^6e - B^2a^6d^6e^6 - 12B^2a^5b^2d^5e^5 + 165B^2a^4b^3d^4e^4 - 520B^2a^3b^4d^3e^3 + 705B^2a^2b^5d^2e^2 - 444B^2ab^6d^2e - 36B^2a^6d^6e^6 - 12B^2a^5b^2d^5e^5 + 165B^2a^4b^3d^4e^4 - 520B^2a^3b^4d^3e^3 + 705B^2a^2b^5d^2e^2 - 444B^2ab^6d^2e - 107B^2b^6d^7 + x^2(-90A^2a^4b^2d^2e^7 + 360A^2a^3b^3d^2e^6 - 540A^2a^2b^4d^2e^5 + 360A^2a^2b^5d^3e^4 - 90A^2b^6d^4e^3 - 36B^2a^5b^2e^7 + 270B^2a^4b^2d^2e^6 - 720B^2a^3b^3d^2e^5 + 900B^2a^2b^4d^3e^4 - 540B^2a^2b^5d^4e^3 + 126B^2b^6d^5e^2)}{(6d^3e^8 + 18d^2e^9x + 18de^{10}x^2 + 6e^{11}x^3)}$$

input

```
integrate((b*x+a)**6*(B*x+A)/(e*x+d)**4,x)
```

output

```
B*b**6*x**4/(4*e**4) + 5*b**2*(a*e - b*d)**3*(4*A*b*e + 3*B*a*e - 7*B*b*d)
*log(d + e*x)/e**8 + x**3*(A*b**6/(3*e**4) + 2*B*a*b**5/e**4 - 4*B*b**6*d/
(3*e**5)) + x**2*(3*A*a*b**5/e**4 - 2*A*b**6*d/e**5 + 15*B*a**2*b**4/(2*e
**4) - 12*B*a*b**5*d/e**5 + 5*B*b**6*d**2/e**6) + x*(15*A*a**2*b**4/e**4 -
24*A*a*b**5*d/e**5 + 10*A*b**6*d**2/e**6 + 20*B*a**3*b**3/e**4 - 60*B*a**2
*b**4*d/e**5 + 60*B*a*b**5*d**2/e**6 - 20*B*b**6*d**3/e**7) + (-2*A*a**6*e
**7 - 6*A*a**5*b*d*e**6 - 30*A*a**4*b**2*d**2*e**5 + 220*A*a**3*b**3*d**3*
e**4 - 390*A*a**2*b**4*d**4*e**3 + 282*A*a*b**5*d**5*e**2 - 74*A*b**6*d**6
*e - B*a**6*d*e**6 - 12*B*a**5*b*d**2*e**5 + 165*B*a**4*b**2*d**3*e**4 - 5
20*B*a**3*b**3*d**4*e**3 + 705*B*a**2*b**4*d**5*e**2 - 444*B*a*b**5*d**6*e
+ 107*B*b**6*d**7 + x**2*(-90*A*a**4*b**2*d**2*e**7 + 360*A*a**3*b**3*d**2
e**6 - 540*A*a**2*b**4*d**2*e**5 + 360*A*a*b**5*d**3*e**4 - 90*A*b**6*d**4
e**3 - 36*B*a**5*b*e**7 + 270*B*a**4*b**2*d**2*e**6 - 720*B*a**3*b**3*d**2
e**5 + 900*B*a**2*b**4*d**3*e**4 - 540*B*a*b**5*d**4*e**3 + 126*B*b**6*d**5
e**2) + x*(-18*A*a**5*b*e**7 - 90*A*a**4*b**2*d**2*e**6 + 540*A*a**3*b**3
d**2*e**5 - 900*A*a**2*b**4*d**3*e**4 + 630*A*a*b**5*d**4*e**3 - 162*A*b**6
d**5e**2 - 3*B*a**6*e**7 - 36*B*a**5*b*d*e**6 + 405*B*a**4*b**2*d**2*e**5 - 120
0*B*a**3*b**3*d**3*e**4 + 1575*B*a**2*b**4*d**4*e**3 - 972*B*a*b**5*d**5e
**2 + 231*B*b**6*d**6*e))/(6*d**3*e**8 + 18*d**2*e**9*x + 18*d*e**10*x**2
+ 6*e**11*x**3)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 793 vs. $2(269) = 538$.

Time = 0.07 (sec) , antiderivative size = 793, normalized size of antiderivative = 2.84

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^4} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^4,x, algorithm="maxima")`

output

```
1/6*(107*B*b^6*d^7 - 2*A*a^6*e^7 - 74*(6*B*a*b^5 + A*b^6)*d^6*e + 141*(5*B
*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 - 130*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4*e^3 +
55*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 - 6*(2*B*a^5*b + 5*A*a^4*b^2)*d^2*e
^5 - (B*a^6 + 6*A*a^5*b)*d*e^6 + 18*(7*B*b^6*d^5*e^2 - 5*(6*B*a*b^5 + A*b^
6)*d^4*e^3 + 10*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*e^4 - 10*(4*B*a^3*b^3 + 3*A*
a^2*b^4)*d^2*e^5 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^6 - (2*B*a^5*b + 5*A*
a^4*b^2)*e^7)*x^2 + 3*(77*B*b^6*d^6*e - 54*(6*B*a*b^5 + A*b^6)*d^5*e^2 + 1
05*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^3 - 100*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3
*e^4 + 45*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^5 - 6*(2*B*a^5*b + 5*A*a^4*b^2
)*d*e^6 - (B*a^6 + 6*A*a^5*b)*e^7)*x)/(e^11*x^3 + 3*d*e^10*x^2 + 3*d^2*e^9
*x + d^3*e^8) + 1/12*(3*B*b^6*e^3*x^4 - 4*(4*B*b^6*d*e^2 - (6*B*a*b^5 + A*
b^6)*e^3)*x^3 + 6*(10*B*b^6*d^2*e - 4*(6*B*a*b^5 + A*b^6)*d*e^2 + 3*(5*B*a
^2*b^4 + 2*A*a*b^5)*e^3)*x^2 - 12*(20*B*b^6*d^3 - 10*(6*B*a*b^5 + A*b^6)*d
^2*e + 12*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^2 - 5*(4*B*a^3*b^3 + 3*A*a^2*b^4)*
e^3)*x)/e^7 + 5*(7*B*b^6*d^4 - 4*(6*B*a*b^5 + A*b^6)*d^3*e + 6*(5*B*a^2*b^
4 + 2*A*a*b^5)*d^2*e^2 - 4*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^3 + (3*B*a^4*b^
2 + 4*A*a^3*b^3)*e^4)*log(e*x + d)/e^8
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 848 vs. $2(269) = 538$.

Time = 0.13 (sec) , antiderivative size = 848, normalized size of antiderivative = 3.04

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^4} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^4,x, algorithm="giac")`

output

$$\begin{aligned}
& 5*(7*B*b^6*d^4 - 24*B*a*b^5*d^3*e - 4*A*b^6*d^3*e + 30*B*a^2*b^4*d^2*e^2 + \\
& 12*A*a*b^5*d^2*e^2 - 16*B*a^3*b^3*d*e^3 - 12*A*a^2*b^4*d*e^3 + 3*B*a^4*b^2* \\
& 2*e^4 + 4*A*a^3*b^3*e^4)*\log(\text{abs}(e*x + d))/e^8 + 1/6*(107*B*b^6*d^7 - 444* \\
& B*a*b^5*d^6*e - 74*A*b^6*d^6*e + 705*B*a^2*b^4*d^5*e^2 + 282*A*a*b^5*d^5*e \\
& ^2 - 520*B*a^3*b^3*d^4*e^3 - 390*A*a^2*b^4*d^4*e^3 + 165*B*a^4*b^2*d^3*e^4 \\
& + 220*A*a^3*b^3*d^3*e^4 - 12*B*a^5*b*d^2*e^5 - 30*A*a^4*b^2*d^2*e^5 - B*a \\
& ^6*d*e^6 - 6*A*a^5*b*d*e^6 - 2*A*a^6*e^7 + 18*(7*B*b^6*d^5*e^2 - 30*B*a*b^5* \\
& d^4*e^3 - 5*A*b^6*d^4*e^3 + 50*B*a^2*b^4*d^3*e^4 + 20*A*a*b^5*d^3*e^4 - \\
& 40*B*a^3*b^3*d^2*e^5 - 30*A*a^2*b^4*d^2*e^5 + 15*B*a^4*b^2*d*e^6 + 20*A*a^3* \\
& b^3*d*e^6 - 2*B*a^5*b*e^7 - 5*A*a^4*b^2*e^7)*x^2 + 3*(77*B*b^6*d^6*e - 3 \\
& 24*B*a*b^5*d^5*e^2 - 54*A*b^6*d^5*e^2 + 525*B*a^2*b^4*d^4*e^3 + 210*A*a*b^5* \\
& d^4*e^3 - 400*B*a^3*b^3*d^3*e^4 - 300*A*a^2*b^4*d^3*e^4 + 135*B*a^4*b^2*d^2* \\
& e^5 + 180*A*a^3*b^3*d^2*e^5 - 12*B*a^5*b*d*e^6 - 30*A*a^4*b^2*d*e^6 - \\
& B*a^6*e^7 - 6*A*a^5*b*e^7)*x)/((e*x + d)^3*e^8) + 1/12*(3*B*b^6*e^12*x^4 - \\
& 16*B*b^6*d*e^11*x^3 + 24*B*a*b^5*e^12*x^3 + 4*A*b^6*e^12*x^3 + 60*B*b^6*d^2* \\
& e^10*x^2 - 144*B*a*b^5*d*e^11*x^2 - 24*A*b^6*d*e^11*x^2 + 90*B*a^2*b^4* \\
& e^12*x^2 + 36*A*a*b^5*e^12*x^2 - 240*B*b^6*d^3*e^9*x + 720*B*a*b^5*d^2*e^10*x \\
& + 120*A*b^6*d^2*e^10*x - 720*B*a^2*b^4*d*e^11*x - 288*A*a*b^5*d*e^11*x \\
& + 240*B*a^3*b^3*e^12*x + 180*A*a^2*b^4*e^12*x)/e^16
\end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.97 (sec) , antiderivative size = 907, normalized size of antiderivative = 3.25

$$\begin{aligned}
& \int \frac{(a + bx)^6 (A + Bx)}{(d + ex)^4} dx = x^3 \left(\frac{Ab^6 + 6Bab^5}{3e^4} - \frac{4Bb^6 d}{3e^5} \right) \\
& - x^2 \left(\frac{2d \left(\frac{Ab^6 + 6Bab^5}{e^4} - \frac{4Bb^6 d}{e^5} \right)}{e} - \frac{3ab^4(2Ab + 5Ba)}{2e^4} + \frac{3Bb^6 d^2}{e^6} \right) \\
& - \frac{Ba^6 de^6 + 2Aa^6 e^7 + 12Ba^5 b d^2 e^5 + 6Aa^5 b d e^6 - 165Ba^4 b^2 d^3 e^4 + 30Aa^4 b^2 d^2 e^5 + 520Ba^3 b^3 d^4 e^3 - 220Aa^3 b^3 d^3 e^4 - 705Ba^2 b^4 d^5 e^2 + 120Ba^2 b^4 d^4 e^3 - 120Ba^2 b^4 d^3 e^4 + 120Ba^2 b^4 d^2 e^5 - 120Ba^2 b^4 d e^6 + 120Ba^2 b^4 e^7}{6e} \\
& - x \left(\frac{6d^2 \left(\frac{Ab^6 + 6Bab^5}{e^4} - \frac{4Bb^6 d}{e^5} \right)}{e^2} \right. \\
& \left. - \frac{4d \left(\frac{4d \left(\frac{Ab^6 + 6Bab^5}{e^4} - \frac{4Bb^6 d}{e^5} \right)}{e} - \frac{3ab^4(2Ab + 5Ba)}{e^4} + \frac{6Bb^6 d^2}{e^6} \right)}{e} - \frac{5a^2 b^3 (3Ab + 4Ba)}{e^4} \right. \\
& \left. + \frac{4Bb^6 d^3}{e^7} \right) \\
& + \frac{\ln(d + ex) (15Ba^4 b^2 e^4 - 80Ba^3 b^3 d e^3 + 20Aa^3 b^3 e^4 + 150Ba^2 b^4 d^2 e^2 - 60Aa^2 b^4 d e^3 - 120Ba^2 b^4 e^4)}{e^8} \\
& + \frac{Bb^6 x^4}{4e^4}
\end{aligned}$$

input `int(((A + B*x)*(a + b*x)^6)/(d + e*x)^4,x)`

output

```

x^3*((A*b^6 + 6*B*a*b^5)/(3*e^4) - (4*B*b^6*d)/(3*e^5)) - x^2*((2*d*((A*b^
6 + 6*B*a*b^5)/e^4 - (4*B*b^6*d)/e^5))/e - (3*a*b^4*(2*A*b + 5*B*a))/(2*e^
4) + (3*B*b^6*d^2)/e^6) - ((2*A*a^6*e^7 - 107*B*b^6*d^7 + 74*A*b^6*d^6*e +
B*a^6*d*e^6 - 282*A*a*b^5*d^5*e^2 + 12*B*a^5*b*d^2*e^5 + 390*A*a^2*b^4*d^
4*e^3 - 220*A*a^3*b^3*d^3*e^4 + 30*A*a^4*b^2*d^2*e^5 - 705*B*a^2*b^4*d^5*e
^2 + 520*B*a^3*b^3*d^4*e^3 - 165*B*a^4*b^2*d^3*e^4 + 6*A*a^5*b*d*e^6 + 444
*B*a*b^5*d^6*e)/(6*e) + x*((B*a^6*e^6)/2 - (77*B*b^6*d^6)/2 + 3*A*a^5*b*e^
6 + 27*A*b^6*d^5*e - 105*A*a*b^5*d^4*e^2 + 15*A*a^4*b^2*d*e^5 + 150*A*a^2*
b^4*d^3*e^3 - 90*A*a^3*b^3*d^2*e^4 - (525*B*a^2*b^4*d^4*e^2)/2 + 200*B*a^3
*b^3*d^3*e^3 - (135*B*a^4*b^2*d^2*e^4)/2 + 162*B*a*b^5*d^5*e + 6*B*a^5*b*d
*e^5) + x^2*(6*B*a^5*b*e^6 - 21*B*b^6*d^5*e + 15*A*a^4*b^2*e^6 + 15*A*b^6*
d^4*e^2 - 60*A*a*b^5*d^3*e^3 - 60*A*a^3*b^3*d*e^5 + 90*B*a*b^5*d^4*e^2 - 4
5*B*a^4*b^2*d*e^5 + 90*A*a^2*b^4*d^2*e^4 - 150*B*a^2*b^4*d^3*e^3 + 120*B*a
^3*b^3*d^2*e^4))/(d^3*e^7 + e^10*x^3 + 3*d^2*e^8*x + 3*d*e^9*x^2) - x*((6*
d^2*((A*b^6 + 6*B*a*b^5)/e^4 - (4*B*b^6*d)/e^5))/e^2 - (4*d*((4*d*((A*b^6
+ 6*B*a*b^5)/e^4 - (4*B*b^6*d)/e^5))/e - (3*a*b^4*(2*A*b + 5*B*a))/e^4 + (
6*B*b^6*d^2)/e^6))/e - (5*a^2*b^3*(3*A*b + 4*B*a))/e^4 + (4*B*b^6*d^3)/e^7
) + (log(d + e*x)*(35*B*b^6*d^4 - 20*A*b^6*d^3*e + 20*A*a^3*b^3*e^4 + 15*B
*a^4*b^2*e^4 + 60*A*a*b^5*d^2*e^2 - 60*A*a^2*b^4*d*e^3 - 80*B*a^3*b^3*d*e^
3 + 150*B*a^2*b^4*d^2*e^2 - 120*B*a*b^5*d^3*e))/e^8 + (B*b^6*x^4)/(4*e^...

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 840, normalized size of antiderivative = 3.01

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^4} dx = \text{Too large to display}$$

input

```
int((b*x+a)^6*(B*x+A)/(e*x+d)^4,x)
```

output

```
(420*log(d + e*x)*a**4*b**3*d**4*e**4 + 1260*log(d + e*x)*a**4*b**3*d**3*e
**5*x + 1260*log(d + e*x)*a**4*b**3*d**2*e**6*x**2 + 420*log(d + e*x)*a**4
*b**3*d*e**7*x**3 - 1680*log(d + e*x)*a**3*b**4*d**5*e**3 - 5040*log(d + e
*x)*a**3*b**4*d**4*e**4*x - 5040*log(d + e*x)*a**3*b**4*d**3*e**5*x**2 - 1
680*log(d + e*x)*a**3*b**4*d**2*e**6*x**3 + 2520*log(d + e*x)*a**2*b**5*d*
*6*e**2 + 7560*log(d + e*x)*a**2*b**5*d**5*e**3*x + 7560*log(d + e*x)*a**2
*b**5*d**4*e**4*x**2 + 2520*log(d + e*x)*a**2*b**5*d**3*e**5*x**3 - 1680*1
og(d + e*x)*a*b**6*d**7*e - 5040*log(d + e*x)*a*b**6*d**6*e**2*x - 5040*lo
g(d + e*x)*a*b**6*d**5*e**3*x**2 - 1680*log(d + e*x)*a*b**6*d**4*e**4*x**3
+ 420*log(d + e*x)*b**7*d**8 + 1260*log(d + e*x)*b**7*d**7*e*x + 1260*log
(d + e*x)*b**7*d**6*e**2*x**2 + 420*log(d + e*x)*b**7*d**5*e**3*x**3 - 4*a
**7*d*e**7 - 14*a**6*b*d**2*e**6 - 42*a**6*b*d*e**7*x + 84*a**5*b**2*e**8*
x**3 + 350*a**4*b**3*d**4*e**4 + 630*a**4*b**3*d**3*e**5*x - 420*a**4*b**3
*d*e**7*x**3 - 1400*a**3*b**4*d**5*e**3 - 2520*a**3*b**4*d**4*e**4*x + 168
0*a**3*b**4*d**2*e**6*x**3 + 420*a**3*b**4*d*e**7*x**4 + 2100*a**2*b**5*d*
*6*e**2 + 3780*a**2*b**5*d**5*e**3*x - 2520*a**2*b**5*d**3*e**5*x**3 - 630
*a**2*b**5*d**2*e**6*x**4 + 126*a**2*b**5*d*e**7*x**5 - 1400*a*b**6*d**7*e
- 2520*a*b**6*d**6*e**2*x + 1680*a*b**6*d**4*e**4*x**3 + 420*a*b**6*d**3*
e**5*x**4 - 84*a*b**6*d**2*e**6*x**5 + 28*a*b**6*d*e**7*x**6 + 350*b**7*d*
*8 + 630*b**7*d**7*e*x - 420*b**7*d**5*e**3*x**3 - 105*b**7*d**4*e**4*x...
```

3.54 $\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^5} dx$

Optimal result	547
Mathematica [A] (verified)	548
Rubi [A] (verified)	548
Maple [B] (verified)	550
Fricas [B] (verification not implemented)	551
Sympy [F(-1)]	552
Maxima [B] (verification not implemented)	552
Giac [B] (verification not implemented)	553
Mupad [B] (verification not implemented)	554
Reduce [B] (verification not implemented)	555

Optimal result

Integrand size = 20, antiderivative size = 279

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^5} dx = \frac{3b^4(bd-ae)(7bBd-2Abe-5aBe)x}{e^7} + \frac{(bd-ae)^6(Bd-Ae)}{4e^8(d+ex)^4} - \frac{(bd-ae)^5(7bBd-6Abe-aBe)}{3e^8(d+ex)^3} + \frac{3b(bd-ae)^4(7bBd-5Abe-2aBe)}{2e^8(d+ex)^2} - \frac{5b^2(bd-ae)^3(7bBd-4Abe-3aBe)}{e^8(d+ex)} - \frac{b^5(7bBd-Abe-6aBe)(d+ex)^2}{2e^8} + \frac{b^6B(d+ex)^3}{3e^8} - \frac{5b^3(bd-ae)^2(7bBd-3Abe-4aBe)\log(d+ex)}{e^8}$$

output

```
3*b^4*(-a*e+b*d)*(-2*A*b*e-5*B*a*e+7*B*b*d)*x/e^7+1/4*(-a*e+b*d)^6*(-A*e+B
*d)/e^8/(e*x+d)^4-1/3*(-a*e+b*d)^5*(-6*A*b*e-B*a*e+7*B*b*d)/e^8/(e*x+d)^3+
3/2*b*(-a*e+b*d)^4*(-5*A*b*e-2*B*a*e+7*B*b*d)/e^8/(e*x+d)^2-5*b^2*(-a*e+b*
d)^3*(-4*A*b*e-3*B*a*e+7*B*b*d)/e^8/(e*x+d)-1/2*b^5*(-A*b*e-6*B*a*e+7*B*b*
d)*(e*x+d)^2/e^8+1/3*b^6*B*(e*x+d)^3/e^8-5*b^3*(-a*e+b*d)^2*(-3*A*b*e-4*B*
a*e+7*B*b*d)*ln(e*x+d)/e^8
```


Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 263, normalized size of antiderivative = 0.94

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^5} dx$$

$$= \frac{-12b^4e(-15a^2Be^2 - 6abe(-5Bd + Ae) + 5b^2d(-3Bd + Ae))x + 6b^5e^2(-5bBd + Abe + 6aBe)x^2 + 4b^6e^3(-3Bd + Ae)x^3 + 3(b^6d - a^6e)(Bd - Ae)x^4 + 4(b^6d - a^6e)^2(7bBd - 6Abe - aBe)}{(d+ex)^5} + \frac{18b^4(bd - ae)^4(7bBd - 5Abe - 2aBe)}{(d+ex)^4} - \frac{60b^3(bd - ae)^3(7bBd - 4Abe - 3aBe)}{(d+ex)^3} - \frac{60b^2(bd - ae)^2(7bBd - 3Abe - 4aBe) \operatorname{Log}[d+ex]}{12e^8}$$

input

```
Integrate[((a + b*x)^6*(A + B*x))/(d + e*x)^5,x]
```

output

```
(-12*b^4*e*(-15*a^2*B*e^2 - 6*a*b*e*(-5*B*d + A*e) + 5*b^2*d*(-3*B*d + A*e))
)*x + 6*b^5*e^2*(-5*b*B*d + A*b*e + 6*a*B*e)*x^2 + 4*b^6*B*e^3*x^3 + (3*(
b*d - a*e)^6*(B*d - A*e))/(d + e*x)^4 - (4*(b*d - a*e)^5*(7*b*B*d - 6*A*b*
e - a*B*e))/(d + e*x)^3 + (18*b*(b*d - a*e)^4*(7*b*B*d - 5*A*b*e - 2*a*B*e
))/(d + e*x)^2 - (60*b^2*(b*d - a*e)^3*(7*b*B*d - 4*A*b*e - 3*a*B*e))/(d +
e*x) - 60*b^3*(b*d - a*e)^2*(7*b*B*d - 3*A*b*e - 4*a*B*e)*Log[d + e*x]/(
12*e^8)
```

Rubi [A] (verified)

Time = 0.73 (sec) , antiderivative size = 279, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^5} dx$$

$$\downarrow 86$$

$$\int \left(\frac{b^5(d+ex)(6aBe + Abe - 7bBd)}{e^7} - \frac{3b^4(bd - ae)(5aBe + 2Abe - 7bBd)}{e^7} + \frac{5b^3(bd - ae)^2(4aBe + 3Abe - 7bBd)}{e^7(d+ex)} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned} & -\frac{b^5(d+ex)^2(-6aBe - Abe + 7bBd)}{2e^8} + \frac{3b^4x(bd-ae)(-5aBe - 2Abe + 7bBd)}{e^8} - \\ & \frac{5b^3(bd-ae)^2 \log(d+ex)(-4aBe - 3Abe + 7bBd)}{e^8} - \frac{5b^2(bd-ae)^3(-3aBe - 4Abe + 7bBd)}{e^8(d+ex)} + \\ & \frac{3b(bd-ae)^4(-2aBe - 5Abe + 7bBd)}{2e^8(d+ex)^2} - \frac{(bd-ae)^5(-aBe - 6Abe + 7bBd)}{3e^8(d+ex)^3} + \\ & \frac{(bd-ae)^6(Bd-Ae)}{4e^8(d+ex)^4} + \frac{b^6B(d+ex)^3}{3e^8} \end{aligned}$$

input `Int[((a + b*x)^6*(A + B*x))/(d + e*x)^5,x]`

output
$$\begin{aligned} & (3*b^4*(b*d - a*e)*(7*b*B*d - 2*A*b*e - 5*a*B*e)*x)/e^7 + ((b*d - a*e)^6*(B*d - A*e))/(4*e^8*(d + e*x)^4) - ((b*d - a*e)^5*(7*b*B*d - 6*A*b*e - a*B*e))/(3*e^8*(d + e*x)^3) + (3*b*(b*d - a*e)^4*(7*b*B*d - 5*A*b*e - 2*a*B*e))/(2*e^8*(d + e*x)^2) - (5*b^2*(b*d - a*e)^3*(7*b*B*d - 4*A*b*e - 3*a*B*e))/(e^8*(d + e*x)) - (b^5*(7*b*B*d - A*b*e - 6*a*B*e)*(d + e*x)^2)/(2*e^8) + (b^6*B*(d + e*x)^3)/(3*e^8) - (5*b^3*(b*d - a*e)^2*(7*b*B*d - 3*A*b*e - 4*a*B*e)*Log[d + e*x])/e^8 \end{aligned}$$

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.))^(n_.)*((e_.) + (f_.)*(x_.))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;`
`FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 797 vs. 2(269) = 538.

Time = 0.24 (sec) , antiderivative size = 798, normalized size of antiderivative = 2.86

method	result
norman	$\frac{b^4(6Aab^2e^2-3Ab^2de+15Ba^2e^2-18Babde+7b^2Bd^2)x^5}{e^3} - \frac{3a^6Ae^7+6Aa^5bde^6+15Aa^4b^2d^2e^5+60Aa^3b^3d^3e^4-375Aa^2b^4d^4e^3+750Aa^2b^5d^5e^2-375Aa^2b^6d^6e+Ba^6d^6e^6+6Bba^5bd^2e^5+45Bba^4b^2d^3e^4-500Bba^3b^3d^4e^3+1875Bba^2b^4d^5e^2-2250Bba^2b^5d^6e+875Bba^2b^6d^7}{e^8} - \frac{(20Aa^3b^3e^4-60Aa^2b^4d^3e^3+120Aa^2b^5d^2e^2-60Aa^2b^6d^3e+15Bba^4b^2e^4-80Bba^3b^3d^2e^3+300Bba^3b^4d^2e^2-360Bba^3b^5d^3e+140Bba^3b^6d^4)}{e^5x^3-3/2(5Aa^4b^2e^5+20Aa^3b^3de^4-90Aa^2b^4d^2e^3+180Aa^2b^5d^3e^2-90Aa^2b^6d^4e+2Bba^5b^2e^5+15Bba^4b^2de^4-120Bba^3b^3d^2e^3+450Bba^2b^4d^3e^2-540Bba^2b^5d^4e+210Bba^2b^6d^5)}{e^6x^2-1/3(6Aa^5b^2e^6+15Aa^4b^2de^5+60Aa^3b^3d^2e^4-330Aa^2b^4d^3e^3+660Aa^2b^5d^4e^2-330Aa^2b^6d^5e+Bba^6e^6+6Bba^5b^2de^5+45Bba^4b^2d^2e^4-440Bba^3b^3d^3e^3+1650Bba^2b^4d^4e^2-1980Bba^2b^5d^5e+770Bba^2b^6d^6)}{e^7x+1/6b^5(3Aa^3b^2e+18Bba^2e-7Bba^2d)}{e^2x^6+1/3b^6B/e^2x^7} + \frac{5b^2(4Aa^2b^3d^3e^4-375Aa^2b^4d^4e^3+750Aa^2b^5d^5e^2-375Aa^2b^6d^6e+Ba^6d^6e^6)}{e^7}$
default	$\frac{b^4(\frac{1}{3}b^2Bx^3e^2+\frac{1}{2}Ab^2e^2x^2+3Bab^2e^2x^2-\frac{5}{2}Bb^2dex^2+6Aab^2e^2x-5Ab^2dex+15Ba^2e^2x-30Babdex+15b^2Bd^2x)}{e^7} - \frac{5b^2(4Aa^2b^3d^3e^4-375Aa^2b^4d^4e^3+750Aa^2b^5d^5e^2-375Aa^2b^6d^6e+Ba^6d^6e^6)}{e^7}$
risch	$\frac{b^6Bx^3}{3e^5} + \frac{b^6Ax^2}{2e^5} + \frac{3b^5Bax^2}{e^5} - \frac{5b^6Bdx^2}{2e^6} + \frac{6b^5Aax}{e^5} - \frac{5b^6Adx}{e^6} + \frac{15b^4Ba^2x}{e^5} - \frac{30b^5Badx}{e^6} + \frac{15b^6Bd^2x}{e^7} + (-$
parallelrisc	Expression too large to display

input

```
int((b*x+a)^6*(B*x+A)/(e*x+d)^5,x,method=_RETURNVERBOSE)
```

output

```
(b^4*(6*A*a*b*e^2-3*A*b^2*d*e+15*B*a^2*e^2-18*B*a*b*d*e+7*B*b^2*d^2)/e^3*x^5-1/12*(3*A*a^6*e^7+6*A*a^5*b*d*e^6+15*A*a^4*b^2*d^2*e^5+60*A*a^3*b^3*d^3*e^4-375*A*a^2*b^4*d^4*e^3+750*A*a*b^5*d^5*e^2-375*A*b^6*d^6*e+Ba^6*d^6*e^6+6*Bba^5bd^2e^5+45*Bba^4b^2d^3e^4-500*Bba^3b^3d^4e^3+1875*Bba^2b^4d^5e^2-2250*Bba^2b^5d^6e+875*Bba^2b^6d^7)/e^8-(20*Aa^3b^3e^4-60Aa^2b^4d^3e^3+120Aa^2b^5d^2e^2-60Aa^2b^6d^3e+15Bba^4b^2e^4-80Bba^3b^3d^2e^3+300Bba^3b^4d^2e^2-360Bba^3b^5d^3e+140Bba^3b^6d^4)/e^5*x^3-3/2*(5Aa^4b^2e^5+20Aa^3b^3de^4-90Aa^2b^4d^2e^3+180Aa^2b^5d^3e^2-90Aa^2b^6d^4e+2Bba^5b^2e^5+15Bba^4b^2de^4-120Bba^3b^3d^2e^3+450Bba^2b^4d^3e^2-540Bba^2b^5d^4e+210Bba^2b^6d^5)/e^6*x^2-1/3*(6Aa^5b^2e^6+15Aa^4b^2de^5+60Aa^3b^3d^2e^4-330Aa^2b^4d^3e^3+660Aa^2b^5d^4e^2-330Aa^2b^6d^5e+Bba^6e^6+6Bba^5b^2de^5+45Bba^4b^2d^2e^4-440Bba^3b^3d^3e^3+1650Bba^2b^4d^4e^2-1980Bba^2b^5d^5e+770Bba^2b^6d^6)/e^7*x+1/6b^5(3Aa^3b^2e+18Bba^2e-7Bba^2d)/e^2*x^6+1/3b^6B/e^2x^7)/(e*x+d)^4+5/e^8*b^3(3Aa^2b^2e^3-6Aa^2b^2de^2+3Aa^2b^3d^2e+4Bba^3e^3-15Bba^2b^2de^2+18Bba^2b^2d^2e-7Bba^3d^3)*ln(e*x+d)
```


Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^5} dx = \text{Timed out}$$

input `integrate((b*x+a)**6*(B*x+A)/(e*x+d)**5,x)`output `Timed out`**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 801 vs. 2(269) = 538.

Time = 0.09 (sec) , antiderivative size = 801, normalized size of antiderivative = 2.87

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^5} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^5,x, algorithm="maxima")`

output

```

-1/12*(319*B*b^6*d^7 + 3*A*a^6*e^7 - 171*(6*B*a*b^5 + A*b^6)*d^6*e + 231*(
5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 - 125*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4*e^3
+ 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 + 3*(2*B*a^5*b + 5*A*a^4*b^2)*d^
2*e^5 + (B*a^6 + 6*A*a^5*b)*d*e^6 + 60*(7*B*b^6*d^4*e^3 - 4*(6*B*a*b^5 + A
*b^6)*d^3*e^4 + 6*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 - 4*(4*B*a^3*b^3 + 3*A
*a^2*b^4)*d*e^6 + (3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^3 + 18*(63*B*b^6*d^5*
e^2 - 35*(6*B*a*b^5 + A*b^6)*d^4*e^3 + 50*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*e^
4 - 30*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)
*d*e^6 + (2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 4*(259*B*b^6*d^6*e - 141*(6*
B*a*b^5 + A*b^6)*d^5*e^2 + 195*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^3 - 110*(4*
B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 + 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^5
+ 3*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^6 + (B*a^6 + 6*A*a^5*b)*e^7)*x)/(e^12*x^
4 + 4*d*e^11*x^3 + 6*d^2*e^10*x^2 + 4*d^3*e^9*x + d^4*e^8) + 1/6*(2*B*b^6*
e^2*x^3 - 3*(5*B*b^6*d*e - (6*B*a*b^5 + A*b^6)*e^2)*x^2 + 6*(15*B*b^6*d^2
- 5*(6*B*a*b^5 + A*b^6)*d*e + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*e^2)*x)/e^7 - 5*
(7*B*b^6*d^3 - 3*(6*B*a*b^5 + A*b^6)*d^2*e + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*d
*e^2 - (4*B*a^3*b^3 + 3*A*a^2*b^4)*e^3)*log(e*x + d)/e^8

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1175 vs. $2(269) = 538$.

Time = 0.14 (sec) , antiderivative size = 1175, normalized size of antiderivative = 4.21

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^5} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^6*(B*x+A)/(e*x+d)^5,x, algorithm="giac")
```

output

```

1/6*(2*B*b^6 - 3*(7*B*b^6*d*e - 6*B*a*b^5*e^2 - A*b^6*e^2)/((e*x + d)*e) +
18*(7*B*b^6*d^2*e^2 - 12*B*a*b^5*d*e^3 - 2*A*b^6*d*e^3 + 5*B*a^2*b^4*e^4
+ 2*A*a*b^5*e^4)/((e*x + d)^2*e^2))*(e*x + d)^3/e^8 + 5*(7*B*b^6*d^3 - 18*
B*a*b^5*d^2*e - 3*A*b^6*d^2*e + 15*B*a^2*b^4*d*e^2 + 6*A*a*b^5*d*e^2 - 4*B
*a^3*b^3*e^3 - 3*A*a^2*b^4*e^3)*log(abs(e*x + d)/((e*x + d)^2*abs(e)))/e^8
- 1/12*(420*B*b^6*d^4*e^36/(e*x + d) - 126*B*b^6*d^5*e^36/(e*x + d)^2 + 2
8*B*b^6*d^6*e^36/(e*x + d)^3 - 3*B*b^6*d^7*e^36/(e*x + d)^4 - 1440*B*a*b^5
*d^3*e^37/(e*x + d) - 240*A*b^6*d^3*e^37/(e*x + d) + 540*B*a*b^5*d^4*e^37/
(e*x + d)^2 + 90*A*b^6*d^4*e^37/(e*x + d)^2 - 144*B*a*b^5*d^5*e^37/(e*x +
d)^3 - 24*A*b^6*d^5*e^37/(e*x + d)^3 + 18*B*a*b^5*d^6*e^37/(e*x + d)^4 + 3
*A*b^6*d^6*e^37/(e*x + d)^4 + 1800*B*a^2*b^4*d^2*e^38/(e*x + d) + 720*A*a*
b^5*d^2*e^38/(e*x + d) - 900*B*a^2*b^4*d^3*e^38/(e*x + d)^2 - 360*A*a*b^5*d
^3*e^38/(e*x + d)^2 + 300*B*a^2*b^4*d^4*e^38/(e*x + d)^3 + 120*A*a*b^5*d^
4*e^38/(e*x + d)^3 - 45*B*a^2*b^4*d^5*e^38/(e*x + d)^4 - 18*A*a*b^5*d^5*e^
38/(e*x + d)^4 - 960*B*a^3*b^3*d*e^39/(e*x + d) - 720*A*a^2*b^4*d*e^39/(e*
x + d) + 720*B*a^3*b^3*d^2*e^39/(e*x + d)^2 + 540*A*a^2*b^4*d^2*e^39/(e*x
+ d)^2 - 320*B*a^3*b^3*d^3*e^39/(e*x + d)^3 - 240*A*a^2*b^4*d^3*e^39/(e*x
+ d)^3 + 60*B*a^3*b^3*d^4*e^39/(e*x + d)^4 + 45*A*a^2*b^4*d^4*e^39/(e*x +
d)^4 + 180*B*a^4*b^2*d^2*e^40/(e*x + d) + 240*A*a^3*b^3*d^2*e^40/(e*x + d) - 270*B
*a^4*b^2*d^2*e^40/(e*x + d)^2 - 360*A*a^3*b^3*d^2*e^40/(e*x + d)^2 + 180*B*...

```

Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 863, normalized size of antiderivative = 3.09

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^5} dx = x^2 \left(\frac{Ab^6 + 6Bab^5}{2e^5} - \frac{5Bb^6d}{2e^6} \right)$$

$$\frac{x^3 (15Ba^4b^2e^6 - 80Ba^3b^3de^5 + 20Aa^3b^3e^6 + 150Ba^2b^4d^2e^4 - 60Aa^2b^4de^5 - 120Bab^5d^3e^3}
{-x \left(\frac{5d \left(\frac{Ab^6 + 6Bab^5}{e^5} - \frac{5Bb^6d}{e^6} \right)}{e} - \frac{3ab^4(2Ab + 5Ba)}{e^5} + \frac{10Bb^6d^2}{e^7} \right)}$$

$$+ \frac{\ln(d + ex) (20Ba^3b^3e^3 - 75Ba^2b^4de^2 + 15Aa^2b^4e^3 + 90Bab^5d^2e - 30Aab^5de^2 - 35Bb^6d^3}
{e^8}$$

$$+ \frac{Bb^6x^3}{3e^5}$$

input

```
int(((A + B*x)*(a + b*x)^6)/(d + e*x)^5,x)
```

output

```

x^2*((A*b^6 + 6*B*a*b^5)/(2*e^5) - (5*B*b^6*d)/(2*e^6)) - (x^3*(20*A*a^3*b
^3*e^6 + 15*B*a^4*b^2*e^6 - 20*A*b^6*d^3*e^3 + 35*B*b^6*d^4*e^2 + 60*A*a*b
^5*d^2*e^4 - 60*A*a^2*b^4*d*e^5 - 120*B*a*b^5*d^3*e^3 - 80*B*a^3*b^3*d*e^5
+ 150*B*a^2*b^4*d^2*e^4) + (3*A*a^6*e^7 + 319*B*b^6*d^7 - 171*A*b^6*d^6*e
+ B*a^6*d*e^6 + 462*A*a*b^5*d^5*e^2 + 6*B*a^5*b*d^2*e^5 - 375*A*a^2*b^4*d
^4*e^3 + 60*A*a^3*b^3*d^3*e^4 + 15*A*a^4*b^2*d^2*e^5 + 1155*B*a^2*b^4*d^5*
e^2 - 500*B*a^3*b^3*d^4*e^3 + 45*B*a^4*b^2*d^3*e^4 + 6*A*a^5*b*d*e^6 - 102
6*B*a*b^5*d^6*e)/(12*e) + x*((B*a^6*e^6)/3 + (259*B*b^6*d^6)/3 + 2*A*a^5*b
*e^6 - 47*A*b^6*d^5*e + 130*A*a*b^5*d^4*e^2 + 5*A*a^4*b^2*d*e^5 - 110*A*a^
2*b^4*d^3*e^3 + 20*A*a^3*b^3*d^2*e^4 + 325*B*a^2*b^4*d^4*e^2 - (440*B*a^3*
b^3*d^3*e^3)/3 + 15*B*a^4*b^2*d^2*e^4 - 282*B*a*b^5*d^5*e + 2*B*a^5*b*d*e^
5) + x^2*(3*B*a^5*b*e^6 + (189*B*b^6*d^5*e)/2 + (15*A*a^4*b^2*e^6)/2 - (10
5*A*b^6*d^4*e^2)/2 + 150*A*a*b^5*d^3*e^3 + 30*A*a^3*b^3*d*e^5 - 315*B*a*b^
5*d^4*e^2 + (45*B*a^4*b^2*d*e^5)/2 - 135*A*a^2*b^4*d^2*e^4 + 375*B*a^2*b^4
*d^3*e^3 - 180*B*a^3*b^3*d^2*e^4)/(d^4*e^7 + e^11*x^4 + 4*d^3*e^8*x + 4*d
*e^10*x^3 + 6*d^2*e^9*x^2) - x*((5*d*((A*b^6 + 6*B*a*b^5)/e^5 - (5*B*b^6*d
)/e^6))/e - (3*a*b^4*(2*A*b + 5*B*a))/e^5 + (10*B*b^6*d^2)/e^7) + (log(d +
e*x)*(15*A*b^6*d^2*e - 35*B*b^6*d^3 + 15*A*a^2*b^4*e^3 + 20*B*a^3*b^3*e^3
- 75*B*a^2*b^4*d*e^2 - 30*A*a*b^5*d*e^2 + 90*B*a*b^5*d^2*e))/e^8 + (B*b^6
*x^3)/(3*e^5)

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 851, normalized size of antiderivative = 3.05

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^5} dx = \text{Too large to display}$$

input

```
int((b*x+a)^6*(B*x+A)/(e*x+d)^5,x)
```


output

```
(420*log(d + e*x)*a**3*b**4*d**5*e**3 + 1680*log(d + e*x)*a**3*b**4*d**4*e
**4*x + 2520*log(d + e*x)*a**3*b**4*d**3*e**5*x**2 + 1680*log(d + e*x)*a**
3*b**4*d**2*e**6*x**3 + 420*log(d + e*x)*a**3*b**4*d*e**7*x**4 - 1260*log(
d + e*x)*a**2*b**5*d**6*e**2 - 5040*log(d + e*x)*a**2*b**5*d**5*e**3*x - 7
560*log(d + e*x)*a**2*b**5*d**4*e**4*x**2 - 5040*log(d + e*x)*a**2*b**5*d*
**3*e**5*x**3 - 1260*log(d + e*x)*a**2*b**5*d**2*e**6*x**4 + 1260*log(d + e
*x)*a*b**6*d**7*e + 5040*log(d + e*x)*a*b**6*d**6*e**2*x + 7560*log(d + e*
x)*a*b**6*d**5*e**3*x**2 + 5040*log(d + e*x)*a*b**6*d**4*e**4*x**3 + 1260*
log(d + e*x)*a*b**6*d**3*e**5*x**4 - 420*log(d + e*x)*b**7*d**8 - 1680*log
(d + e*x)*b**7*d**7*e*x - 2520*log(d + e*x)*b**7*d**6*e**2*x**2 - 1680*log
(d + e*x)*b**7*d**5*e**3*x**3 - 420*log(d + e*x)*b**7*d**4*e**4*x**4 - 3*a
**7*d*e**7 - 7*a**6*b*d**2*e**6 - 28*a**6*b*d*e**7*x - 21*a**5*b**2*d**3*e
**5 - 84*a**5*b**2*d**2*e**6*x - 126*a**5*b**2*d*e**7*x**2 + 105*a**4*b**3
*e**8*x**4 + 455*a**3*b**4*d**5*e**3 + 1400*a**3*b**4*d**4*e**4*x + 1260*a
**3*b**4*d**3*e**5*x**2 - 420*a**3*b**4*d*e**7*x**4 - 1365*a**2*b**5*d**6*
e**2 - 4200*a**2*b**5*d**5*e**3*x - 3780*a**2*b**5*d**4*e**4*x**2 + 1260*a
**2*b**5*d**2*e**6*x**4 + 252*a**2*b**5*d*e**7*x**5 + 1365*a*b**6*d**7*e +
4200*a*b**6*d**6*e**2*x + 3780*a*b**6*d**5*e**3*x**2 - 1260*a*b**6*d**3*e
**5*x**4 - 252*a*b**6*d**2*e**6*x**5 + 42*a*b**6*d*e**7*x**6 - 455*b**7*d*
**8 - 1400*b**7*d**7*e*x - 1260*b**7*d**6*e**2*x**2 + 420*b**7*d**4*e**4...
```

3.55 $\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^6} dx$

Optimal result	557
Mathematica [B] (verified)	558
Rubi [A] (verified)	558
Maple [B] (verified)	560
Fricas [B] (verification not implemented)	561
Sympy [F(-1)]	562
Maxima [B] (verification not implemented)	562
Giac [B] (verification not implemented)	563
Mupad [B] (verification not implemented)	564
Reduce [B] (verification not implemented)	565

Optimal result

Integrand size = 20, antiderivative size = 272

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^6} dx = -\frac{b^5(6bBd - Abe - 6aBe)x}{e^7} + \frac{b^6 Bx^2}{2e^6} + \frac{(bd - ae)^6(Bd - Ae)}{5e^8(d+ex)^5} - \frac{(bd - ae)^5(7bBd - 6Abe - aBe)}{4e^8(d+ex)^4} + \frac{b(bd - ae)^4(7bBd - 5Abe - 2aBe)}{e^8(d+ex)^3} - \frac{5b^2(bd - ae)^3(7bBd - 4Abe - 3aBe)}{2e^8(d+ex)^2} + \frac{5b^3(bd - ae)^2(7bBd - 3Abe - 4aBe)}{e^8(d+ex)} + \frac{3b^4(bd - ae)(7bBd - 2Abe - 5aBe) \log(d+ex)}{e^8}$$

output

```
-b^5*(-A*b*e-6*B*a*e+6*B*b*d)*x/e^7+1/2*b^6*B*x^2/e^6+1/5*(-a*e+b*d)^6*(-A
*e+B*d)/e^8/(e*x+d)^5-1/4*(-a*e+b*d)^5*(-6*A*b*e-B*a*e+7*B*b*d)/e^8/(e*x+d
)^4+b*(-a*e+b*d)^4*(-5*A*b*e-2*B*a*e+7*B*b*d)/e^8/(e*x+d)^3-5/2*b^2*(-a*e+
b*d)^3*(-4*A*b*e-3*B*a*e+7*B*b*d)/e^8/(e*x+d)^2+5*b^3*(-a*e+b*d)^2*(-3*A*b
*e-4*B*a*e+7*B*b*d)/e^8/(e*x+d)+3*b^4*(-a*e+b*d)*(-2*A*b*e-5*B*a*e+7*B*b*d
)*ln(e*x+d)/e^8
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 633 vs. $2(272) = 544$.

Time = 0.20 (sec) , antiderivative size = 633, normalized size of antiderivative = 2.33

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^6} dx$$

$$= \frac{-a^6e^6(4Ae + B(d+5ex)) - 2a^5be^5(3Ae(d+5ex) + 2B(d^2 + 5dex + 10e^2x^2)) - 5a^4b^2e^4(2Ae(d^2 + 5dex + 10e^2x^2)) - 4a^3b^3e^3(3Ae(d^3 + 5d^2ex + 10d^2e^2x^2 + 10e^3x^3)) - 2a^2b^4e^2(4Ae(d^4 + 5d^3ex + 10d^2e^2x^2 + 10de^3x^3 + 5e^4x^4)) + 5a^2b^4e^2(-12Ae(d^4 + 5d^3ex + 10d^2e^2x^2 + 10de^3x^3 + 5e^4x^4)) + B*d*(137*d^4 + 625*d^3*ex + 1100*d^2*e^2*x^2 + 900*d*e^3*x^3 + 300*e^4*x^4)) + 2*a*b^5*e*(A*d*e*(137*d^4 + 625*d^3*ex + 1100*d^2*e^2*x^2 + 900*d*e^3*x^3 + 300*e^4*x^4) - 6*B*(87*d^6 + 375*d^5*ex + 600*d^4*e^2*x^2 + 400*d^3*e^3*x^3 + 50*d^2*e^4*x^4 - 50*d*e^5*x^5 - 10*e^6*x^6)) + b^6*(-2*A*e*(87*d^6 + 375*d^5*ex + 600*d^4*e^2*x^2 + 400*d^3*e^3*x^3 + 50*d^2*e^4*x^4 - 50*d*e^5*x^5 - 10*e^6*x^6) + B*(459*d^7 + 1875*d^6*ex + 2700*d^5*e^2*x^2 + 1300*d^4*e^3*x^3 - 400*d^3*e^4*x^4 - 500*d^2*e^5*x^5 - 70*d*e^6*x^6 + 10*e^7*x^7)) + 60*b^4*(b*d - a*e)*(7*b*B*d - 2*A*b*e - 5*a*B*e)*(d + e*x)^5*Log[d + e*x]}{(20*e^8*(d + e*x)^5)}$$

input

```
Integrate[((a + b*x)^6*(A + B*x))/(d + e*x)^6,x]
```

output

```
(-(a^6*e^6*(4*A*e + B*(d + 5*e*x))) - 2*a^5*b*e^5*(3*A*e*(d + 5*e*x) + 2*B*(d^2 + 5*d*e*x + 10*e^2*x^2)) - 5*a^4*b^2*e^4*(2*A*e*(d^2 + 5*d*e*x + 10*e^2*x^2) + 3*B*(d^3 + 5*d^2*e*x + 10*d*e^2*x^2 + 10*e^3*x^3)) - 20*a^3*b^3*e^3*(A*e*(d^3 + 5*d^2*e*x + 10*d*e^2*x^2 + 10*e^3*x^3) + 4*B*(d^4 + 5*d^3*e*x + 10*d^2*e^2*x^2 + 10*d*e^3*x^3 + 5*e^4*x^4)) + 5*a^2*b^4*e^2*(-12*A*e*(d^4 + 5*d^3*e*x + 10*d^2*e^2*x^2 + 10*d*e^3*x^3 + 5*e^4*x^4) + B*d*(137*d^4 + 625*d^3*e*x + 1100*d^2*e^2*x^2 + 900*d*e^3*x^3 + 300*e^4*x^4)) + 2*a*b^5*e*(A*d*e*(137*d^4 + 625*d^3*e*x + 1100*d^2*e^2*x^2 + 900*d*e^3*x^3 + 300*e^4*x^4) - 6*B*(87*d^6 + 375*d^5*e*x + 600*d^4*e^2*x^2 + 400*d^3*e^3*x^3 + 50*d^2*e^4*x^4 - 50*d*e^5*x^5 - 10*e^6*x^6)) + b^6*(-2*A*e*(87*d^6 + 375*d^5*e*x + 600*d^4*e^2*x^2 + 400*d^3*e^3*x^3 + 50*d^2*e^4*x^4 - 50*d*e^5*x^5 - 10*e^6*x^6) + B*(459*d^7 + 1875*d^6*e*x + 2700*d^5*e^2*x^2 + 1300*d^4*e^3*x^3 - 400*d^3*e^4*x^4 - 500*d^2*e^5*x^5 - 70*d*e^6*x^6 + 10*e^7*x^7)) + 60*b^4*(b*d - a*e)*(7*b*B*d - 2*A*b*e - 5*a*B*e)*(d + e*x)^5*Log[d + e*x])/(20*e^8*(d + e*x)^5)
```

Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 272, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^6} dx$$

↓ 86

$$\int \left(\frac{b^5(6aBe + Abe - 6bBd)}{e^7} - \frac{3b^4(bd - ae)(5aBe + 2Abe - 7bBd)}{e^7(d + ex)} + \frac{5b^3(bd - ae)^2(4aBe + 3Abe - 7bBd)}{e^7(d + ex)^2} - \dots \right)$$

↓ 2009

$$\begin{aligned} & - \frac{b^5x(-6aBe - Abe + 6bBd)}{e^7} + \frac{3b^4(bd - ae) \log(d + ex)(-5aBe - 2Abe + 7bBd)}{e^8} + \\ & \frac{5b^3(bd - ae)^2(-4aBe - 3Abe + 7bBd)}{e^8(d + ex)} - \frac{5b^2(bd - ae)^3(-3aBe - 4Abe + 7bBd)}{2e^8(d + ex)^2} + \\ & \frac{b(bd - ae)^4(-2aBe - 5Abe + 7bBd)}{e^8(d + ex)^3} - \frac{(bd - ae)^5(-aBe - 6Abe + 7bBd)}{4e^8(d + ex)^4} + \\ & \frac{(bd - ae)^6(Bd - Ae)}{5e^8(d + ex)^5} + \frac{b^6Bx^2}{2e^6} \end{aligned}$$

input `Int[((a + b*x)^6*(A + B*x))/(d + e*x)^6,x]`

output `-((b^5*(6*b*B*d - A*b*e - 6*a*B*e)*x)/e^7) + (b^6*B*x^2)/(2*e^6) + ((b*d - a*e)^6*(B*d - A*e))/(5*e^8*(d + e*x)^5) - ((b*d - a*e)^5*(7*b*B*d - 6*A*b*e - a*B*e))/(4*e^8*(d + e*x)^4) + (b*(b*d - a*e)^4*(7*b*B*d - 5*A*b*e - 2*a*B*e))/(e^8*(d + e*x)^3) - (5*b^2*(b*d - a*e)^3*(7*b*B*d - 4*A*b*e - 3*a*B*e))/(2*e^8*(d + e*x)^2) + (5*b^3*(b*d - a*e)^2*(7*b*B*d - 3*A*b*e - 4*a*B*e))/(e^8*(d + e*x)) + (3*b^4*(b*d - a*e)*(7*b*B*d - 2*A*b*e - 5*a*B*e)*Log[d + e*x])/e^8`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1157 vs. $2(264) = 528$.

Time = 0.13 (sec) , antiderivative size = 1157, normalized size of antiderivative = 4.25

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^6} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^6,x, algorithm="fricas")`

output

```
1/20*(10*B*b^6*e^7*x^7 + 459*B*b^6*d^7 - 4*A*a^6*e^7 - 174*(6*B*a*b^5 + A
b^6)*d^6*e + 137*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 - 20*(4*B*a^3*b^3 + 3*A
a^2*b^4)*d^4*e^3 - 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 - 2*(2*B*a^5*b +
5*A*a^4*b^2)*d^2*e^5 - (B*a^6 + 6*A*a^5*b)*d*e^6 - 10*(7*B*b^6*d*e^6 - 2*
(6*B*a*b^5 + A*b^6)*e^7)*x^6 - 100*(5*B*b^6*d^2*e^5 - (6*B*a*b^5 + A*b^6)*
d*e^6)*x^5 - 100*(4*B*b^6*d^3*e^4 + (6*B*a*b^5 + A*b^6)*d^2*e^5 - 3*(5*B*a
^2*b^4 + 2*A*a*b^5)*d*e^6 + (4*B*a^3*b^3 + 3*A*a^2*b^4)*e^7)*x^4 + 50*(26*
B*b^6*d^4*e^3 - 16*(6*B*a*b^5 + A*b^6)*d^3*e^4 + 18*(5*B*a^2*b^4 + 2*A*a*b
^5)*d^2*e^5 - 4*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^6 - (3*B*a^4*b^2 + 4*A*a^3
*b^3)*e^7)*x^3 + 10*(270*B*b^6*d^5*e^2 - 120*(6*B*a*b^5 + A*b^6)*d^4*e^3 +
110*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*e^4 - 20*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^
2*e^5 - 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^6 - 2*(2*B*a^5*b + 5*A*a^4*b^2)*
e^7)*x^2 + 5*(375*B*b^6*d^6*e - 150*(6*B*a*b^5 + A*b^6)*d^5*e^2 + 125*(5*B
a^2*b^4 + 2*A*a*b^5)*d^4*e^3 - 20*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 - 5
*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^5 - 2*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^6 -
(B*a^6 + 6*A*a^5*b)*e^7)*x + 60*(7*B*b^6*d^7 - 2*(6*B*a*b^5 + A*b^6)*d^6*
e + (5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 + (7*B*b^6*d^2*e^5 - 2*(6*B*a*b^5 +
A*b^6)*d*e^6 + (5*B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^5 + 5*(7*B*b^6*d^3*e^4 - 2
*(6*B*a*b^5 + A*b^6)*d^2*e^5 + (5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6)*x^4 + 10*(
7*B*b^6*d^4*e^3 - 2*(6*B*a*b^5 + A*b^6)*d^3*e^4 + (5*B*a^2*b^4 + 2*A*a...
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^6} dx = \text{Timed out}$$

input `integrate((b*x+a)**6*(B*x+A)/(e*x+d)**6,x)`

output `Timed out`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 814 vs. $2(264) = 528$.

Time = 0.07 (sec) , antiderivative size = 814, normalized size of antiderivative = 2.99

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^6} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^6,x, algorithm="maxima")`

output

```

1/20*(459*B*b^6*d^7 - 4*A*a^6*e^7 - 174*(6*B*a*b^5 + A*b^6)*d^6*e + 137*(5
*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 - 20*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4*e^3 -
5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 - 2*(2*B*a^5*b + 5*A*a^4*b^2)*d^2*e
^5 - (B*a^6 + 6*A*a^5*b)*d*e^6 + 100*(7*B*b^6*d^3*e^4 - 3*(6*B*a*b^5 + A*b
^6)*d^2*e^5 + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6 - (4*B*a^3*b^3 + 3*A*a^2*b
^4)*e^7)*x^4 + 50*(49*B*b^6*d^4*e^3 - 20*(6*B*a*b^5 + A*b^6)*d^3*e^4 + 18*
(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 - 4*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^6 -
(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^3 + 10*(329*B*b^6*d^5*e^2 - 130*(6*B*a*
b^5 + A*b^6)*d^4*e^3 + 110*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*e^4 - 20*(4*B*a^3
*b^3 + 3*A*a^2*b^4)*d^2*e^5 - 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^6 - 2*(2*B
*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 5*(399*B*b^6*d^6*e - 154*(6*B*a*b^5 + A*b
^6)*d^5*e^2 + 125*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^3 - 20*(4*B*a^3*b^3 + 3*
A*a^2*b^4)*d^3*e^4 - 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^5 - 2*(2*B*a^5*b
+ 5*A*a^4*b^2)*d*e^6 - (B*a^6 + 6*A*a^5*b)*e^7)*x)/(e^13*x^5 + 5*d*e^12*x^
4 + 10*d^2*e^11*x^3 + 10*d^3*e^10*x^2 + 5*d^4*e^9*x + d^5*e^8) + 1/2*(B*b^
6*e*x^2 - 2*(6*B*b^6*d - (6*B*a*b^5 + A*b^6)*e)*x)/e^7 + 3*(7*B*b^6*d^2 -
2*(6*B*a*b^5 + A*b^6)*d*e + (5*B*a^2*b^4 + 2*A*a*b^5)*e^2)*log(e*x + d)/e^
8

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 831 vs. $2(264) = 528$.

Time = 0.13 (sec) , antiderivative size = 831, normalized size of antiderivative = 3.06

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^6} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^6*(B*x+A)/(e*x+d)^6,x, algorithm="giac")
```


output

```

3*(7*B*b^6*d^2 - 12*B*a*b^5*d*e - 2*A*b^6*d*e + 5*B*a^2*b^4*e^2 + 2*A*a*b^
5*e^2)*log(abs(e*x + d))/e^8 + 1/2*(B*b^6*e^6*x^2 - 12*B*b^6*d*e^5*x + 12*
B*a*b^5*e^6*x + 2*A*b^6*e^6*x)/e^12 + 1/20*(459*B*b^6*d^7 - 1044*B*a*b^5*d
^6*e - 174*A*b^6*d^6*e + 685*B*a^2*b^4*d^5*e^2 + 274*A*a*b^5*d^5*e^2 - 80*
B*a^3*b^3*d^4*e^3 - 60*A*a^2*b^4*d^4*e^3 - 15*B*a^4*b^2*d^3*e^4 - 20*A*a^3
*b^3*d^3*e^4 - 4*B*a^5*b*d^2*e^5 - 10*A*a^4*b^2*d^2*e^5 - B*a^6*d*e^6 - 6*
A*a^5*b*d*e^6 - 4*A*a^6*e^7 + 100*(7*B*b^6*d^3*e^4 - 18*B*a*b^5*d^2*e^5 -
3*A*b^6*d^2*e^5 + 15*B*a^2*b^4*d*e^6 + 6*A*a*b^5*d*e^6 - 4*B*a^3*b^3*e^7 -
3*A*a^2*b^4*e^7)*x^4 + 50*(49*B*b^6*d^4*e^3 - 120*B*a*b^5*d^3*e^4 - 20*A*
b^6*d^3*e^4 + 90*B*a^2*b^4*d^2*e^5 + 36*A*a*b^5*d^2*e^5 - 16*B*a^3*b^3*d*e
^6 - 12*A*a^2*b^4*d*e^6 - 3*B*a^4*b^2*e^7 - 4*A*a^3*b^3*e^7)*x^3 + 10*(329
*B*b^6*d^5*e^2 - 780*B*a*b^5*d^4*e^3 - 130*A*b^6*d^4*e^3 + 550*B*a^2*b^4*d
^3*e^4 + 220*A*a*b^5*d^3*e^4 - 80*B*a^3*b^3*d^2*e^5 - 60*A*a^2*b^4*d^2*e^5
- 15*B*a^4*b^2*d*e^6 - 20*A*a^3*b^3*d*e^6 - 4*B*a^5*b*e^7 - 10*A*a^4*b^2*
e^7)*x^2 + 5*(399*B*b^6*d^6*e - 924*B*a*b^5*d^5*e^2 - 154*A*b^6*d^5*e^2 +
625*B*a^2*b^4*d^4*e^3 + 250*A*a*b^5*d^4*e^3 - 80*B*a^3*b^3*d^3*e^4 - 60*A*
a^2*b^4*d^3*e^4 - 15*B*a^4*b^2*d^2*e^5 - 20*A*a^3*b^3*d^2*e^5 - 4*B*a^5*b*
d*e^6 - 10*A*a^4*b^2*d*e^6 - B*a^6*e^7 - 6*A*a^5*b*e^7)*x)/((e*x + d)^5*e^
8)

```

Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 862, normalized size of antiderivative = 3.17

$$\int \frac{(a + bx)^6 (A + Bx)}{(d + ex)^6} dx = x \left(\frac{A b^6 + 6 B a b^5}{e^6} - \frac{6 B b^6 d}{e^7} \right)$$

$$x^3 \left(\frac{15 B a^4 b^2 e^6}{2} + 40 B a^3 b^3 d e^5 + 10 A a^3 b^3 e^6 - 225 B a^2 b^4 d^2 e^4 + 30 A a^2 b^4 d e^5 + 300 B a b^5 d^3 e^3 - \right.$$

$$\left. \frac{\ln(d + ex) (15 B a^2 b^4 e^2 - 36 B a b^5 d e + 6 A a b^5 e^2 + 21 B b^6 d^2 - 6 A b^6 d e)}{e^8} \right.$$

$$\left. + \frac{B b^6 x^2}{2 e^6} \right)$$

input

```
int(((A + B*x)*(a + b*x)^6)/(d + e*x)^6,x)
```

output

```
x*((A*b^6 + 6*B*a*b^5)/e^6 - (6*B*b^6*d)/e^7) - (x^3*(10*A*a^3*b^3*e^6 + (
15*B*a^4*b^2*e^6)/2 + 50*A*b^6*d^3*e^3 - (245*B*b^6*d^4*e^2)/2 - 90*A*a*b^
5*d^2*e^4 + 30*A*a^2*b^4*d*e^5 + 300*B*a*b^5*d^3*e^3 + 40*B*a^3*b^3*d*e^5
- 225*B*a^2*b^4*d^2*e^4) + (4*A*a^6*e^7 - 459*B*b^6*d^7 + 174*A*b^6*d^6*e
+ B*a^6*d*e^6 - 274*A*a*b^5*d^5*e^2 + 4*B*a^5*b*d^2*e^5 + 60*A*a^2*b^4*d^4
*e^3 + 20*A*a^3*b^3*d^3*e^4 + 10*A*a^4*b^2*d^2*e^5 - 685*B*a^2*b^4*d^5*e^2
+ 80*B*a^3*b^3*d^4*e^3 + 15*B*a^4*b^2*d^3*e^4 + 6*A*a^5*b*d*e^6 + 1044*B*
a*b^5*d^6*e)/(20*e) + x*((B*a^6*e^6)/4 - (399*B*b^6*d^6)/4 + (3*A*a^5*b*e^
6)/2 + (77*A*b^6*d^5*e)/2 - (125*A*a*b^5*d^4*e^2)/2 + (5*A*a^4*b^2*d*e^5)/
2 + 15*A*a^2*b^4*d^3*e^3 + 5*A*a^3*b^3*d^2*e^4 - (625*B*a^2*b^4*d^4*e^2)/4
+ 20*B*a^3*b^3*d^3*e^3 + (15*B*a^4*b^2*d^2*e^4)/4 + 231*B*a*b^5*d^5*e + B
*a^5*b*d*e^5) + x^2*(2*B*a^5*b*e^6 - (329*B*b^6*d^5*e)/2 + 5*A*a^4*b^2*e^6
+ 65*A*b^6*d^4*e^2 - 110*A*a*b^5*d^3*e^3 + 10*A*a^3*b^3*d*e^5 + 390*B*a*b
^5*d^4*e^2 + (15*B*a^4*b^2*d*e^5)/2 + 30*A*a^2*b^4*d^2*e^4 - 275*B*a^2*b^4
*d^3*e^3 + 40*B*a^3*b^3*d^2*e^4) + x^4*(15*A*a^2*b^4*e^6 + 20*B*a^3*b^3*e^
6 + 15*A*b^6*d^2*e^4 - 35*B*b^6*d^3*e^3 + 90*B*a*b^5*d^2*e^4 - 75*B*a^2*b^
4*d*e^5 - 30*A*a*b^5*d*e^5))/(d^5*e^7 + e^12*x^5 + 5*d^4*e^8*x + 5*d*e^11*
x^4 + 10*d^3*e^9*x^2 + 10*d^2*e^10*x^3) + (log(d + e*x)*(21*B*b^6*d^2 - 6*
A*b^6*d*e + 6*A*a*b^5*e^2 + 15*B*a^2*b^4*e^2 - 36*B*a*b^5*d*e))/e^8 + (B*b
^6*x^2)/(2*e^6)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 816, normalized size of antiderivative = 3.00

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^6} dx = \text{Too large to display}$$

input

```
int((b*x+a)^6*(B*x+A)/(e*x+d)^6,x)
```

output

```
(420*log(d + e*x)*a**2*b**5*d**6*e**2 + 2100*log(d + e*x)*a**2*b**5*d**5*
e**3*x + 4200*log(d + e*x)*a**2*b**5*d**4*e**4*x**2 + 4200*log(d + e*x)*a**
2*b**5*d**3*e**5*x**3 + 2100*log(d + e*x)*a**2*b**5*d**2*e**6*x**4 + 420*1
og(d + e*x)*a**2*b**5*d*e**7*x**5 - 840*log(d + e*x)*a*b**6*d**7*e - 4200*
log(d + e*x)*a*b**6*d**6*e**2*x - 8400*log(d + e*x)*a*b**6*d**5*e**3*x**2
- 8400*log(d + e*x)*a*b**6*d**4*e**4*x**3 - 4200*log(d + e*x)*a*b**6*d**3*
e**5*x**4 - 840*log(d + e*x)*a*b**6*d**2*e**6*x**5 + 420*log(d + e*x)*b**7
*d**8 + 2100*log(d + e*x)*b**7*d**7*e*x + 4200*log(d + e*x)*b**7*d**6*e**2
*x**2 + 4200*log(d + e*x)*b**7*d**5*e**3*x**3 + 2100*log(d + e*x)*b**7*d**
4*e**4*x**4 + 420*log(d + e*x)*b**7*d**3*e**5*x**5 - 4*a**7*d*e**7 - 7*a**
6*b*d**2*e**6 - 35*a**6*b*d*e**7*x - 14*a**5*b**2*d**3*e**5 - 70*a**5*b**2
*d**2*e**6*x - 140*a**5*b**2*d*e**7*x**2 - 35*a**4*b**3*d**4*e**4 - 175*a*
**4*b**3*d**3*e**5*x - 350*a**4*b**3*d**2*e**6*x**2 - 350*a**4*b**3*d*e**7*
x**3 + 140*a**3*b**4*e**8*x**5 + 539*a**2*b**5*d**6*e**2 + 2275*a**2*b**5*
d**5*e**3*x + 3500*a**2*b**5*d**4*e**4*x**2 + 2100*a**2*b**5*d**3*e**5*x**
3 - 420*a**2*b**5*d*e**7*x**5 - 1078*a*b**6*d**7*e - 4550*a*b**6*d**6*e**2
*x - 7000*a*b**6*d**5*e**3*x**2 - 4200*a*b**6*d**4*e**4*x**3 + 840*a*b**6*
d**2*e**6*x**5 + 140*a*b**6*d*e**7*x**6 + 539*b**7*d**8 + 2275*b**7*d**7*e
*x + 3500*b**7*d**6*e**2*x**2 + 2100*b**7*d**5*e**3*x**3 - 420*b**7*d**3*e
**5*x**5 - 70*b**7*d**2*e**6*x**6 + 10*b**7*d*e**7*x**7)/(20*d*e**8*(d...
```

3.56 $\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^7} dx$

Optimal result	567
Mathematica [B] (verified)	568
Rubi [A] (verified)	569
Maple [B] (verified)	570
Fricas [B] (verification not implemented)	571
Sympy [F(-1)]	572
Maxima [B] (verification not implemented)	573
Giac [B] (verification not implemented)	573
Mupad [B] (verification not implemented)	574
Reduce [B] (verification not implemented)	575

Optimal result

Integrand size = 20, antiderivative size = 278

$$\begin{aligned}
 \int \frac{(a+bx)^6(A+Bx)}{(d+ex)^7} dx = & \frac{b^6 Bx}{e^7} + \frac{(bd-ae)^6(Bd-Ae)}{6e^8(d+ex)^6} \\
 & - \frac{(bd-ae)^5(7bBd-6Abe-aBe)}{5e^8(d+ex)^5} \\
 & + \frac{3b(bd-ae)^4(7bBd-5Abe-2aBe)}{4e^8(d+ex)^4} \\
 & - \frac{5b^2(bd-ae)^3(7bBd-4Abe-3aBe)}{3e^8(d+ex)^3} \\
 & + \frac{5b^3(bd-ae)^2(7bBd-3Abe-4aBe)}{2e^8(d+ex)^2} \\
 & - \frac{3b^4(bd-ae)(7bBd-2Abe-5aBe)}{e^8(d+ex)} \\
 & - \frac{b^5(7bBd-Abe-6aBe)\log(d+ex)}{e^8}
 \end{aligned}$$

output

```

b^6*B*x/e^7+1/6*(-a*e+b*d)^6*(-A*e+B*d)/e^8/(e*x+d)^6-1/5*(-a*e+b*d)^5*(-6
*A*b*e-B*a*e+7*B*b*d)/e^8/(e*x+d)^5+3/4*b*(-a*e+b*d)^4*(-5*A*b*e-2*B*a*e+7
*B*b*d)/e^8/(e*x+d)^4-5/3*b^2*(-a*e+b*d)^3*(-4*A*b*e-3*B*a*e+7*B*b*d)/e^8/
(e*x+d)^3+5/2*b^3*(-a*e+b*d)^2*(-3*A*b*e-4*B*a*e+7*B*b*d)/e^8/(e*x+d)^2-3*
b^4*(-a*e+b*d)*(-2*A*b*e-5*B*a*e+7*B*b*d)/e^8/(e*x+d)-b^5*(-A*b*e-6*B*a*e+
7*B*b*d)*ln(e*x+d)/e^8

```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 619 vs. $2(278) = 556$.

Time = 0.20 (sec) , antiderivative size = 619, normalized size of antiderivative = 2.23

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^7} dx =$$

$$\frac{2a^6e^6(5Ae+B(d+6ex)) + 6a^5be^5(2Ae(d+6ex) + B(d^2+6dex+15e^2x^2)) + 15a^4b^2e^4(Ae(d^2+6dex+15e^2x^2) + B(d^3+6d^2ex+15de^2x^2+20e^3x^3)) + 10a^3b^3e^3(Ae(d^3+6d^2ex+15de^2x^2+20e^3x^3) + B(d^4+6d^3ex+15d^2e^2x^2+20de^3x^3+15e^4x^4)) + 5a^2b^4e^2(Ae(d^4+6d^3ex+15d^2e^2x^2+20de^3x^3+15e^4x^4) + B(d^5+6d^4ex+15d^3e^2x^2+20d^2e^3x^3+15de^4x^4+6e^5x^5)) - 6ab^5e(-10Ae(d^5+6d^4ex+15d^3e^2x^2+20d^2e^3x^3+15de^4x^4+6e^5x^5) + B(d^6+6d^5ex+15d^4e^2x^2+20d^3e^3x^3+15d^2e^4x^4+6de^5x^5)) - b^6(Ae(d^6+6d^5ex+15d^4e^2x^2+20d^3e^3x^3+15d^2e^4x^4+6de^5x^5) + B(d^7+6d^6ex+15d^5e^2x^2+20d^4e^3x^3+15d^3e^4x^4+6d^2e^5x^5))}{e^8(d+ex)^6}$$

input

```
Integrate[((a + b*x)^6*(A + B*x))/(d + e*x)^7,x]
```

output

```

-1/60*(2*a^6*e^6*(5*A*e + B*(d + 6*e*x)) + 6*a^5*b*e^5*(2*A*e*(d + 6*e*x)
+ B*(d^2 + 6*d*e*x + 15*e^2*x^2)) + 15*a^4*b^2*e^4*(A*e*(d^2 + 6*d*e*x + 1
5*e^2*x^2) + B*(d^3 + 6*d^2*e*x + 15*d*e^2*x^2 + 20*e^3*x^3)) + 20*a^3*b^3
*e^3*(A*e*(d^3 + 6*d^2*e*x + 15*d*e^2*x^2 + 20*e^3*x^3) + 2*B*(d^4 + 6*d^3
*e*x + 15*d^2*e^2*x^2 + 20*d*e^3*x^3 + 15*e^4*x^4)) + 30*a^2*b^4*e^2*(A*e*
(d^4 + 6*d^3*e*x + 15*d^2*e^2*x^2 + 20*d*e^3*x^3 + 15*e^4*x^4) + 5*B*(d^5
+ 6*d^4*e*x + 15*d^3*e^2*x^2 + 20*d^2*e^3*x^3 + 15*d*e^4*x^4 + 6*e^5*x^5))
- 6*a*b^5*e*(-10*A*e*(d^5 + 6*d^4*e*x + 15*d^3*e^2*x^2 + 20*d^2*e^3*x^3 +
15*d*e^4*x^4 + 6*e^5*x^5) + B*(147*d^5 + 822*d^4*e*x + 1875*d^3*e^2*x^2
+ 2200*d^2*e^3*x^3 + 1350*d*e^4*x^4 + 360*e^5*x^5)) - b^6*(A*d*e*(147*d^5
+ 822*d^4*e*x + 1875*d^3*e^2*x^2 + 2200*d^2*e^3*x^3 + 1350*d*e^4*x^4 + 36
0*e^5*x^5) - B*(669*d^7 + 3594*d^6*e*x + 7725*d^5*e^2*x^2 + 8200*d^4*e^3*x
^3 + 4050*d^3*e^4*x^4 + 360*d^2*e^5*x^5 - 360*d*e^6*x^6 - 60*e^7*x^7)) + 6
0*b^5*(7*b*B*d - A*b*e - 6*a*B*e)*(d + e*x)^6*Log[d + e*x]/(e^8*(d + e*x)
^6)

```

Rubi [A] (verified)

Time = 0.65 (sec) , antiderivative size = 278, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^7} dx$$

↓ 86

$$\int \left(\frac{b^5(6aBe + Abe - 7bBd)}{e^7(d + ex)} - \frac{3b^4(bd - ae)(5aBe + 2Abe - 7bBd)}{e^7(d + ex)^2} + \frac{5b^3(bd - ae)^2(4aBe + 3Abe - 7bBd)}{e^7(d + ex)^3} - \dots \right) dx$$

↓ 2009

$$\begin{aligned} & - \frac{b^5 \log(d + ex)(-6aBe - Abe + 7bBd)}{e^8} - \frac{3b^4(bd - ae)(-5aBe - 2Abe + 7bBd)}{e^8(d + ex)} + \\ & \frac{5b^3(bd - ae)^2(-4aBe - 3Abe + 7bBd)}{2e^8(d + ex)^2} - \frac{5b^2(bd - ae)^3(-3aBe - 4Abe + 7bBd)}{3e^8(d + ex)^3} + \\ & \frac{3b(bd - ae)^4(-2aBe - 5Abe + 7bBd)}{4e^8(d + ex)^4} - \frac{(bd - ae)^5(-aBe - 6Abe + 7bBd)}{5e^8(d + ex)^5} + \\ & \frac{(bd - ae)^6(Bd - Ae)}{6e^8(d + ex)^6} + \frac{b^6 Bx}{e^7} \end{aligned}$$

input `Int[((a + b*x)^6*(A + B*x))/(d + e*x)^7,x]`

output `(b^6*B*x)/e^7 + ((b*d - a*e)^6*(B*d - A*e))/(6*e^8*(d + e*x)^6) - ((b*d - a*e)^5*(7*b*B*d - 6*A*b*e - a*B*e))/(5*e^8*(d + e*x)^5) + (3*b*(b*d - a*e)^4*(7*b*B*d - 5*A*b*e - 2*a*B*e))/(4*e^8*(d + e*x)^4) - (5*b^2*(b*d - a*e)^3*(7*b*B*d - 4*A*b*e - 3*a*B*e))/(3*e^8*(d + e*x)^3) + (5*b^3*(b*d - a*e)^2*(7*b*B*d - 3*A*b*e - 4*a*B*e))/(2*e^8*(d + e*x)^2) - (3*b^4*(b*d - a*e)*(7*b*B*d - 2*A*b*e - 5*a*B*e))/(e^8*(d + e*x)) - (b^5*(7*b*B*d - A*b*e - 6*a*B*e)*Log[d + e*x])/e^8`

Definitions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 804 vs. $2(268) = 536$.

Time = 0.23 (sec) , antiderivative size = 805, normalized size of antiderivative = 2.90

method	result
default	$\frac{b^6 B x}{e^7} - \frac{a^6 A e^7 - 6A a^5 b d e^6 + 15A a^4 b^2 d^2 e^5 - 20A a^3 b^3 d^3 e^4 + 15A a^2 b^4 d^4 e^3 - 6A a b^5 d^5 e^2 + A b^6 d^6 e - B a^6 d e^6 + 6B a^5 b d^2 e^5 - 15B a^4 b^2 d^3 e^4 + 15B a^3 b^3 d^4 e^3 - 6B a^2 b^4 d^5 e^2 + 6B a b^5 d^6 e - 6B b^6 d^7}{6e^8 (ex+d)^6}$
norman	$\frac{b^6 B x^7}{e} - \frac{10a^6 A e^7 + 12A a^5 b d e^6 + 15A a^4 b^2 d^2 e^5 + 20A a^3 b^3 d^3 e^4 + 30A a^2 b^4 d^4 e^3 + 60A a b^5 d^5 e^2 - 147A b^6 d^6 e + 2B a^6 d e^6 + 6B a^5 b d^2 e^5 + 15B a^4 b^2 d^3 e^4 - 15B a^3 b^3 d^4 e^3 - 6B a^2 b^4 d^5 e^2 + 6B a b^5 d^6 e - 6B b^6 d^7}{60e^8}$
risch	$\frac{b^6 B x}{e^7} + \frac{(-6A a b^5 e^6 + 6A b^6 d e^5 - 15B a^2 b^4 e^6 + 36B a b^5 d e^5 - 21b^6 B d^2 e^4) x^5 - 5e^3 b^3 (3A a^2 b e^3 + 6A a b^2 d e^2 - 9A b^3 d^2 e + 4B a^3 e^3 - 4B a^2 b d e^2 + 6B a b^2 d^2 e - 6B b^3 d^3 e)}{2}$
parallelrisc	Expression too large to display

input

```
int((b*x+a)^6*(B*x+A)/(e*x+d)^7,x,method=_RETURNVERBOSE)
```

output

```

b^6*B*x/e^7-1/6*(A*a^6*e^7-6*A*a^5*b*d*e^6+15*A*a^4*b^2*d^2*e^5-20*A*a^3*b^3*d^3*e^4+15*A*a^2*b^4*d^4*e^3-6*A*a*b^5*d^5*e^2+A*b^6*d^6*e-B*a^6*d*e^6+6*B*a^5*b*d^2*e^5-15*B*a^4*b^2*d^3*e^4+20*B*a^3*b^3*d^4*e^3-15*B*a^2*b^4*d^5*e^2+6*B*a*b^5*d^6*e-B*b^6*d^7)/e^8/(e*x+d)^6-3*b^4/e^8*(2*A*a*b*e^2-2*A*b^2*d*e+5*B*a^2*e^2-12*B*a*b*d*e+7*B*b^2*d^2)/(e*x+d)-5/2/e^8*b^3*(3*A*a^2*b*e^3-6*A*a*b^2*d*e^2+3*A*b^3*d^2*e+4*B*a^3*e^3-15*B*a^2*b*d*e^2+18*B*a*b^2*d^2*e-7*B*b^3*d^3)/(e*x+d)^2-1/5/e^8*(6*A*a^5*b*e^6-30*A*a^4*b^2*d*e^5+60*A*a^3*b^3*d^2*e^4-60*A*a^2*b^4*d^3*e^3+30*A*a*b^5*d^4*e^2-6*A*b^6*d^5*e+B*a^6*e^6-12*B*a^5*b*d*e^5+45*B*a^4*b^2*d^2*e^4-80*B*a^3*b^3*d^3*e^3+75*B*a^2*b^4*d^4*e^2-36*B*a*b^5*d^5*e+7*B*b^6*d^6)/(e*x+d)^5-3/4*b/e^8*(5*A*a^4*b*e^5-20*A*a^3*b^2*d*e^4+30*A*a^2*b^3*d^2*e^3-20*A*a*b^4*d^3*e^2+5*A*b^5*d^4*e+2*B*a^5*e^5-15*B*a^4*b*d*e^4+40*B*a^3*b^2*d^2*e^3-50*B*a^2*b^3*d^3*e^2+30*B*a*b^4*d^4*e-7*B*b^5*d^5)/(e*x+d)^4+b^5/e^8*(A*b*e+6*B*a*e-7*B*b*d)*ln(e*x+d)-5/3*b^2/e^8*(4*A*a^3*b*e^4-12*A*a^2*b^2*d*e^3+12*A*a*b^3*d^2*e^2-4*A*b^4*d^3*e+3*B*a^4*e^4-16*B*a^3*b*d*e^3+30*B*a^2*b^2*d^2*e^2-24*B*a*b^3*d^3*e+7*B*b^4*d^4)/(e*x+d)^3

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1063 vs. $2(268) = 536$.

Time = 0.12 (sec) , antiderivative size = 1063, normalized size of antiderivative = 3.82

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^7} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^6*(B*x+A)/(e*x+d)^7,x, algorithm="fricas")
```


output

```

1/60*(60*B*b^6*e^7*x^7 + 360*B*b^6*d*e^6*x^6 - 669*B*b^6*d^7 - 10*A*a^6*e^
7 + 147*(6*B*a*b^5 + A*b^6)*d^6*e - 30*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 -
10*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4*e^3 - 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^
3*e^4 - 3*(2*B*a^5*b + 5*A*a^4*b^2)*d^2*e^5 - 2*(B*a^6 + 6*A*a^5*b)*d*e^6
- 180*(2*B*b^6*d^2*e^5 - 2*(6*B*a*b^5 + A*b^6)*d*e^6 + (5*B*a^2*b^4 + 2*A*
a*b^5)*e^7)*x^5 - 150*(27*B*b^6*d^3*e^4 - 9*(6*B*a*b^5 + A*b^6)*d^2*e^5 +
3*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6 + (4*B*a^3*b^3 + 3*A*a^2*b^4)*e^7)*x^4 -
100*(82*B*b^6*d^4*e^3 - 22*(6*B*a*b^5 + A*b^6)*d^3*e^4 + 6*(5*B*a^2*b^4 +
2*A*a*b^5)*d^2*e^5 + 2*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^6 + (3*B*a^4*b^2 +
4*A*a^3*b^3)*e^7)*x^3 - 15*(515*B*b^6*d^5*e^2 - 125*(6*B*a*b^5 + A*b^6)*d^
4*e^3 + 30*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*e^4 + 10*(4*B*a^3*b^3 + 3*A*a^2*
b^4)*d^2*e^5 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^6 + 3*(2*B*a^5*b + 5*A*a^
4*b^2)*e^7)*x^2 - 6*(599*B*b^6*d^6*e - 137*(6*B*a*b^5 + A*b^6)*d^5*e^2 + 3
0*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^3 + 10*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e
^4 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^5 + 3*(2*B*a^5*b + 5*A*a^4*b^2)*d
*e^6 + 2*(B*a^6 + 6*A*a^5*b)*e^7)*x - 60*(7*B*b^6*d^7 - (6*B*a*b^5 + A*b^6
)*d^6*e + (7*B*b^6*d*e^6 - (6*B*a*b^5 + A*b^6)*e^7)*x^6 + 6*(7*B*b^6*d^2*e
^5 - (6*B*a*b^5 + A*b^6)*d*e^6)*x^5 + 15*(7*B*b^6*d^3*e^4 - (6*B*a*b^5 + A
*b^6)*d^2*e^5)*x^4 + 20*(7*B*b^6*d^4*e^3 - (6*B*a*b^5 + A*b^6)*d^3*e^4)*x^
3 + 15*(7*B*b^6*d^5*e^2 - (6*B*a*b^5 + A*b^6)*d^4*e^3)*x^2 + 6*(7*B*b^6...

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^7} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**6*(B*x+A)/(e*x+d)**7,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 824 vs. $2(268) = 536$.

Time = 0.08 (sec) , antiderivative size = 824, normalized size of antiderivative = 2.96

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^7} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^7,x, algorithm="maxima")`

output

```
B*b^6*x/e^7 - 1/60*(669*B*b^6*d^7 + 10*A*a^6*e^7 - 147*(6*B*a*b^5 + A*b^6)
*d^6*e + 30*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 + 10*(4*B*a^3*b^3 + 3*A*a^2*
b^4)*d^4*e^3 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 + 3*(2*B*a^5*b + 5*A*
a^4*b^2)*d^2*e^5 + 2*(B*a^6 + 6*A*a^5*b)*d*e^6 + 180*(7*B*b^6*d^2*e^5 - 2*
(6*B*a*b^5 + A*b^6)*d*e^6 + (5*B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^5 + 150*(35*B
*b^6*d^3*e^4 - 9*(6*B*a*b^5 + A*b^6)*d^2*e^5 + 3*(5*B*a^2*b^4 + 2*A*a*b^5)
*d*e^6 + (4*B*a^3*b^3 + 3*A*a^2*b^4)*e^7)*x^4 + 100*(91*B*b^6*d^4*e^3 - 22
*(6*B*a*b^5 + A*b^6)*d^3*e^4 + 6*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 + 2*(4*
B*a^3*b^3 + 3*A*a^2*b^4)*d*e^6 + (3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^3 + 15
*(539*B*b^6*d^5*e^2 - 125*(6*B*a*b^5 + A*b^6)*d^4*e^3 + 30*(5*B*a^2*b^4 +
2*A*a*b^5)*d^3*e^4 + 10*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + 5*(3*B*a^4*b
^2 + 4*A*a^3*b^3)*d*e^6 + 3*(2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 6*(609*B*
b^6*d^6*e - 137*(6*B*a*b^5 + A*b^6)*d^5*e^2 + 30*(5*B*a^2*b^4 + 2*A*a*b^5)
*d^4*e^3 + 10*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 + 5*(3*B*a^4*b^2 + 4*A*a
^3*b^3)*d^2*e^5 + 3*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^6 + 2*(B*a^6 + 6*A*a^5*b
)*e^7)*x)/(e^14*x^6 + 6*d*e^13*x^5 + 15*d^2*e^12*x^4 + 20*d^3*e^11*x^3 + 1
5*d^4*e^10*x^2 + 6*d^5*e^9*x + d^6*e^8) - (7*B*b^6*d - (6*B*a*b^5 + A*b^6)
*e)*log(e*x + d)/e^8
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 826 vs. $2(268) = 536$.

Time = 0.12 (sec) , antiderivative size = 826, normalized size of antiderivative = 2.97

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^7} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^7,x, algorithm="giac")`

output

$$\begin{aligned} & B*b^6*x/e^7 - (7*B*b^6*d - 6*B*a*b^5*e - A*b^6*e)*\log(\text{abs}(e*x + d))/e^8 - \\ & 1/60*(669*B*b^6*d^7 - 882*B*a*b^5*d^6*e - 147*A*b^6*d^6*e + 150*B*a^2*b^4*d^5*e^2 + 60*A*a*b^5*d^5*e^2 + 40*B*a^3*b^3*d^4*e^3 + 30*A*a^2*b^4*d^4*e^3 \\ & + 15*B*a^4*b^2*d^3*e^4 + 20*A*a^3*b^3*d^3*e^4 + 6*B*a^5*b*d^2*e^5 + 15*A*a^4*b^2*d^2*e^5 + 2*B*a^6*d*e^6 + 12*A*a^5*b*d*e^6 + 10*A*a^6*e^7 + 180*(7 \\ & *B*b^6*d^2*e^5 - 12*B*a*b^5*d*e^6 - 2*A*b^6*d*e^6 + 5*B*a^2*b^4*e^7 + 2*A*a*b^5*e^7)*x^5 + 150*(35*B*b^6*d^3*e^4 - 54*B*a*b^5*d^2*e^5 - 9*A*b^6*d^2* \\ & e^5 + 15*B*a^2*b^4*d*e^6 + 6*A*a*b^5*d*e^6 + 4*B*a^3*b^3*e^7 + 3*A*a^2*b^4*e^7)*x^4 + 100*(91*B*b^6*d^4*e^3 - 132*B*a*b^5*d^3*e^4 - 22*A*b^6*d^3*e^4 \\ & + 30*B*a^2*b^4*d^2*e^5 + 12*A*a*b^5*d^2*e^5 + 8*B*a^3*b^3*d*e^6 + 6*A*a^2*b^4*d*d*e^6 + 3*B*a^4*b^2*e^7 + 4*A*a^3*b^3*e^7)*x^3 + 15*(539*B*b^6*d^5*e^2 \\ & - 750*B*a*b^5*d^4*e^3 - 125*A*b^6*d^4*e^3 + 150*B*a^2*b^4*d^3*e^4 + 60*A*a*b^5*d^3*e^4 + 40*B*a^3*b^3*d^2*e^5 + 30*A*a^2*b^4*d^2*e^5 + 15*B*a^4*b^2*d*e^6 \\ & + 20*A*a^3*b^3*d*e^6 + 6*B*a^5*b*e^7 + 15*A*a^4*b^2*e^7)*x^2 + 6*(609*B*b^6*d^6*e - 822*B*a*b^5*d^5*e^2 - 137*A*b^6*d^5*e^2 + 150*B*a^2*b^4*d^4*e^3 \\ & + 60*A*a*b^5*d^4*e^3 + 40*B*a^3*b^3*d^3*e^4 + 30*A*a^2*b^4*d^3*e^4 + 15*B*a^4*b^2*d^2*e^5 + 20*A*a^3*b^3*d^2*e^5 + 6*B*a^5*b*d*e^6 + 15*A*a^4*b^2*d*e^6 \\ & + 2*B*a^6*e^7 + 12*A*a^5*b*e^7)*x)/((e*x + d)^6*e^8) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 1.06 (sec) , antiderivative size = 875, normalized size of antiderivative = 3.15

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^7} dx = \frac{\ln(d + ex) (Ab^6e - 7Bb^6d + 6Bab^5e)}{e^8} - \frac{x^3 \left(5Ba^4b^2e^6 + \frac{40Ba^3b^3de^5}{3} + \frac{20Aa^3b^3e^6}{3} + 50Ba^2b^4d^2e^4 + 10Aa^2b^4de^5 - 220Bab^5d^3e^3 + 20A \right)}{e^7} + \frac{Bb^6x}{e^7}$$

input `int(((A + B*x)*(a + b*x)^6)/(d + e*x)^7,x)`

output

```
(log(d + e*x)*(A*b^6*e - 7*B*b^6*d + 6*B*a*b^5*e))/e^8 - (x^3*((20*A*a^3*b^3*e^6)/3 + 5*B*a^4*b^2*e^6 - (110*A*b^6*d^3*e^3)/3 + (455*B*b^6*d^4*e^2)/3 + 20*A*a*b^5*d^2*e^4 + 10*A*a^2*b^4*d*e^5 - 220*B*a*b^5*d^3*e^3 + (40*B*a^3*b^3*d*e^5)/3 + 50*B*a^2*b^4*d^2*e^4) + (10*A*a^6*e^7 + 669*B*b^6*d^7 - 147*A*b^6*d^6*e + 2*B*a^6*d*e^6 + 60*A*a*b^5*d^5*e^2 + 6*B*a^5*b*d^2*e^5 + 30*A*a^2*b^4*d^4*e^3 + 20*A*a^3*b^3*d^3*e^4 + 15*A*a^4*b^2*d^2*e^5 + 150*B*a^2*b^4*d^5*e^2 + 40*B*a^3*b^3*d^4*e^3 + 15*B*a^4*b^2*d^3*e^4 + 12*A*a^5*b*d*e^6 - 882*B*a*b^5*d^6*e)/(60*e) + x*((B*a^6*e^6)/5 + (609*B*b^6*d^6)/10 + (6*A*a^5*b*e^6)/5 - (137*A*b^6*d^5*e)/10 + 6*A*a*b^5*d^4*e^2 + (3*A*a^4*b^2*d*e^5)/2 + 3*A*a^2*b^4*d^3*e^3 + 2*A*a^3*b^3*d^2*e^4 + 15*B*a^2*b^4*d^4*e^2 + 4*B*a^3*b^3*d^3*e^3 + (3*B*a^4*b^2*d^2*e^4)/2 - (411*B*a*b^5*d^5*e)/5 + (3*B*a^5*b*d*e^5)/5) + x^5*(6*A*a*b^5*e^6 - 6*A*b^6*d*e^5 + 15*B*a^2*b^4*e^6 + 21*B*b^6*d^2*e^4 - 36*B*a*b^5*d*e^5) + x^2*((3*B*a^5*b*e^6)/2 + (539*B*b^6*d^5*e)/4 + (15*A*a^4*b^2*e^6)/4 - (125*A*b^6*d^4*e^2)/4 + 15*A*a*b^5*d^3*e^3 + 5*A*a^3*b^3*d*e^5 - (375*B*a*b^5*d^4*e^2)/2 + (15*B*a^4*b^2*d*e^5)/4 + (15*A*a^2*b^4*d^2*e^4)/2 + (75*B*a^2*b^4*d^3*e^3)/2 + 10*B*a^3*b^3*d^2*e^4) + x^4*((15*A*a^2*b^4*e^6)/2 + 10*B*a^3*b^3*e^6 - (45*A*b^6*d^2*e^4)/2 + (175*B*b^6*d^3*e^3)/2 - 135*B*a*b^5*d^2*e^4 + (75*B*a^2*b^4*d*e^5)/2 + 15*A*a*b^5*d*e^5))/(d^6*e^7 + e^13*x^6 + 6*d^5*e^8*x + 6*d^e^12*x^5 + 15*d^4*e^9*x^2 + 20*d^3*e^10*x^3 + 15*d^2*e^11*x^4) + (B*b^6...
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 735, normalized size of antiderivative = 2.64

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^7} dx$$

$$= \frac{6300 \log(ex + d) a b^6 d^5 e^3 x^2 - 2520 \log(ex + d) b^7 d^3 e^5 x^5 - 6300 \log(ex + d) b^7 d^6 e^2 x^2 - 420 a b^6 d e^7 x^6 + 4 \dots}{(d + ex)^7}$$

input

```
int((b*x+a)^6*(B*x+A)/(e*x+d)^7,x)
```

output

```
(420*log(d + e*x)*a*b**6*d**7*e + 2520*log(d + e*x)*a*b**6*d**6*e**2*x + 6
300*log(d + e*x)*a*b**6*d**5*e**3*x**2 + 8400*log(d + e*x)*a*b**6*d**4*e**
4*x**3 + 6300*log(d + e*x)*a*b**6*d**3*e**5*x**4 + 2520*log(d + e*x)*a*b**
6*d**2*e**6*x**5 + 420*log(d + e*x)*a*b**6*d*e**7*x**6 - 420*log(d + e*x)*
b**7*d**8 - 2520*log(d + e*x)*b**7*d**7*e*x - 6300*log(d + e*x)*b**7*d**6*
e**2*x**2 - 8400*log(d + e*x)*b**7*d**5*e**3*x**3 - 6300*log(d + e*x)*b**7
*d**4*e**4*x**4 - 2520*log(d + e*x)*b**7*d**3*e**5*x**5 - 420*log(d + e*x)
*b**7*d**2*e**6*x**6 - 10*a**7*d*e**7 - 14*a**6*b*d**2*e**6 - 84*a**6*b*d*
e**7*x - 21*a**5*b**2*d**3*e**5 - 126*a**5*b**2*d**2*e**6*x - 315*a**5*b**
2*d*e**7*x**2 - 35*a**4*b**3*d**4*e**4 - 210*a**4*b**3*d**3*e**5*x - 525*a
**4*b**3*d**2*e**6*x**2 - 700*a**4*b**3*d*e**7*x**3 - 70*a**3*b**4*d**5*e*
*3 - 420*a**3*b**4*d**4*e**4*x - 1050*a**3*b**4*d**3*e**5*x**2 - 1400*a**3
*b**4*d**2*e**6*x**3 - 1050*a**3*b**4*d*e**7*x**4 + 210*a**2*b**5*e**8*x**
6 + 609*a*b**6*d**7*e + 3234*a*b**6*d**6*e**2*x + 6825*a*b**6*d**5*e**3*x*
*2 + 7000*a*b**6*d**4*e**4*x**3 + 3150*a*b**6*d**3*e**5*x**4 - 420*a*b**6*
d*e**7*x**6 - 609*b**7*d**8 - 3234*b**7*d**7*e*x - 6825*b**7*d**6*e**2*x**
2 - 7000*b**7*d**5*e**3*x**3 - 3150*b**7*d**4*e**4*x**4 + 420*b**7*d**2*e*
*6*x**6 + 60*b**7*d*e**7*x**7)/(60*d**8*(d**6 + 6*d**5*e*x + 15*d**4*e**
2*x**2 + 20*d**3*e**3*x**3 + 15*d**2*e**4*x**4 + 6*d**5*x**5 + e**6*x**6
))
```

3.57 $\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^8} dx$

Optimal result	577
Mathematica [B] (verified)	578
Rubi [A] (verified)	578
Maple [B] (verified)	580
Fricas [B] (verification not implemented)	581
Sympy [F(-1)]	582
Maxima [B] (verification not implemented)	583
Giac [B] (verification not implemented)	583
Mupad [B] (verification not implemented)	584
Reduce [B] (verification not implemented)	585

Optimal result

Integrand size = 20, antiderivative size = 213

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^8} dx = -\frac{(Bd-Ae)(a+bx)^7}{7e(bd-ae)(d+ex)^7} - \frac{B(bd-ae)^6}{6e^8(d+ex)^6} + \frac{6bB(bd-ae)^5}{5e^8(d+ex)^5} - \frac{15b^2B(bd-ae)^4}{4e^8(d+ex)^4} + \frac{20b^3B(bd-ae)^3}{3e^8(d+ex)^3} - \frac{15b^4B(bd-ae)^2}{2e^8(d+ex)^2} + \frac{6b^5B(bd-ae)}{e^8(d+ex)} + \frac{b^6B \log(d+ex)}{e^8}$$

output

```
-1/7*(-A*e+B*d)*(b*x+a)^7/e/(-a*e+b*d)/(e*x+d)^7-1/6*B*(-a*e+b*d)^6/e^8/(e*x+d)^6+6/5*b*B*(-a*e+b*d)^5/e^8/(e*x+d)^5-15/4*b^2*B*(-a*e+b*d)^4/e^8/(e*x+d)^4+20/3*b^3*B*(-a*e+b*d)^3/e^8/(e*x+d)^3-15/2*b^4*B*(-a*e+b*d)^2/e^8/(e*x+d)^2+6*b^5*B*(-a*e+b*d)/e^8/(e*x+d)+b^6*B*ln(e*x+d)/e^8
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 615 vs. $2(213) = 426$.

Time = 0.28 (sec) , antiderivative size = 615, normalized size of antiderivative = 2.89

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^8} dx = \frac{10a^6e^6(6Ae+B(d+7ex)) + 12a^5be^5(5Ae(d+7ex) + 2B(d^2+7dex+21e^2x^2)) + 15a^4b^2e^4(4Ae(d^2+7dex+21e^2x^2) + 2B(d^3+7d^2ex+21de^2x^2+35e^3x^3)) + 20a^3b^3e^3(3Ae(d^3+7d^2ex+21de^2x^2+35e^3x^3) + 4B(d^4+7d^3ex+21d^2e^2x^2+35de^3x^3+35e^4x^4)) + 30a^2b^4e^2(2Ae(d^4+7d^3ex+21d^2e^2x^2+35de^3x^3+35e^4x^4) + 5B(d^5+7d^4ex+21d^3e^2x^2+35d^2e^3x^3+35de^4x^4+21e^5x^5)) + 60ab^5e(Ae(d^5+7d^4ex+21d^3e^2x^2+35d^2e^3x^3+35de^4x^4+21e^5x^5) + 6B(d^6+7d^5ex+21d^4e^2x^2+35d^3e^3x^3+35d^2e^4x^4+21de^5x^5+7e^6x^6)) + b^6(6Ae(d^6+7d^5ex+21d^4e^2x^2+35d^3e^3x^3+35d^2e^4x^4+21de^5x^5+7e^6x^6) - Bd(1089d^6+7203d^5ex+20139d^4e^2x^2+30625d^3e^3x^3+26950d^2e^4x^4+13230de^5x^5+2940e^6x^6)) - 420b^6B(d+ex)^7 \operatorname{Log}[d+ex]}{(e^8(d+ex)^7)}$$

input

```
Integrate[((a + b*x)^6*(A + B*x))/(d + e*x)^8,x]
```

output

```
-1/420*(10*a^6*e^6*(6*A*e + B*(d + 7*e*x)) + 12*a^5*b*e^5*(5*A*e*(d + 7*e*x) + 2*B*(d^2 + 7*d*e*x + 21*e^2*x^2)) + 15*a^4*b^2*e^4*(4*A*e*(d^2 + 7*d*e*x + 21*e^2*x^2) + 3*B*(d^3 + 7*d^2*e*x + 21*d*e^2*x^2 + 35*e^3*x^3)) + 20*a^3*b^3*e^3*(3*A*e*(d^3 + 7*d^2*e*x + 21*d*e^2*x^2 + 35*e^3*x^3) + 4*B*(d^4 + 7*d^3*e*x + 21*d^2*e^2*x^2 + 35*d*e^3*x^3 + 35*e^4*x^4)) + 30*a^2*b^4*e^2*(2*A*e*(d^4 + 7*d^3*e*x + 21*d^2*e^2*x^2 + 35*d*e^3*x^3 + 35*e^4*x^4) + 5*B*(d^5 + 7*d^4*e*x + 21*d^3*e^2*x^2 + 35*d^2*e^3*x^3 + 35*d*e^4*x^4 + 21*e^5*x^5)) + 60*a*b^5*e*(A*e*(d^5 + 7*d^4*e*x + 21*d^3*e^2*x^2 + 35*d^2*e^3*x^3 + 35*d*e^4*x^4 + 21*e^5*x^5) + 6*B*(d^6 + 7*d^5*e*x + 21*d^4*e^2*x^2 + 35*d^3*e^3*x^3 + 35*d^2*e^4*x^4 + 21*d*e^5*x^5 + 7*e^6*x^6)) + b^6*(60*A*e*(d^6 + 7*d^5*e*x + 21*d^4*e^2*x^2 + 35*d^3*e^3*x^3 + 35*d^2*e^4*x^4 + 21*d*e^5*x^5 + 7*e^6*x^6) - B*d*(1089*d^6 + 7203*d^5*e*x + 20139*d^4*e^2*x^2 + 30625*d^3*e^3*x^3 + 26950*d^2*e^4*x^4 + 13230*d*e^5*x^5 + 2940*e^6*x^6)) - 420*b^6*B*(d + e*x)^7*Log[d + e*x]/(e^8*(d + e*x)^7)
```

Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {87, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a+bx)^6(A+Bx)}{(d+ex)^8} dx \\
 & \quad \downarrow 87 \\
 & \frac{B \int \frac{(a+bx)^6}{(d+ex)^7} dx}{e} - \frac{(a+bx)^7(Bd-Ae)}{7e(d+ex)^7(bd-ae)} \\
 & \quad \downarrow 49 \\
 & \frac{B \int \left(\frac{b^6}{e^6(d+ex)} - \frac{6(bd-ae)b^5}{e^6(d+ex)^2} + \frac{15(bd-ae)^2b^4}{e^6(d+ex)^3} - \frac{20(bd-ae)^3b^3}{e^6(d+ex)^4} + \frac{15(bd-ae)^4b^2}{e^6(d+ex)^5} - \frac{6(bd-ae)^5b}{e^6(d+ex)^6} + \frac{(ae-bd)^6}{e^6(d+ex)^7} \right) dx}{\frac{(a+bx)^7(Bd-Ae)}{7e(d+ex)^7(bd-ae)}} \\
 & \quad \downarrow 2009 \\
 & \frac{B \left(\frac{6b^5(bd-ae)}{e^7(d+ex)} - \frac{15b^4(bd-ae)^2}{2e^7(d+ex)^2} + \frac{20b^3(bd-ae)^3}{3e^7(d+ex)^3} - \frac{15b^2(bd-ae)^4}{4e^7(d+ex)^4} + \frac{6b(bd-ae)^5}{5e^7(d+ex)^5} - \frac{(bd-ae)^6}{6e^7(d+ex)^6} + \frac{b^6 \log(d+ex)}{e^7} \right)}{\frac{(a+bx)^7(Bd-Ae)}{7e(d+ex)^7(bd-ae)}}
 \end{aligned}$$

input `Int[((a + b*x)^6*(A + B*x))/(d + e*x)^8,x]`

output `-1/7*((B*d - A*e)*(a + b*x)^7)/(e*(b*d - a*e)*(d + e*x)^7) + (B*(-1/6*(b*d - a*e)^6/(e^7*(d + e*x)^6) + (6*b*(b*d - a*e)^5)/(5*e^7*(d + e*x)^5) - (15*b^2*(b*d - a*e)^4)/(4*e^7*(d + e*x)^4) + (20*b^3*(b*d - a*e)^3)/(3*e^7*(d + e*x)^3) - (15*b^4*(b*d - a*e)^2)/(2*e^7*(d + e*x)^2) + (6*b^5*(b*d - a*e))/(e^7*(d + e*x)) + (b^6*Log[d + e*x])/e^7)/e`

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 87

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p
_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 790 vs. 2(201) = 402.

Time = 0.23 (sec) , antiderivative size = 791, normalized size of antiderivative = 3.71

method	result
risch	$\frac{-\frac{b^5(Abe+6Bae-7Bbd)x^6}{e^2} - \frac{3b^4(2Aab e^2+2A b^2 de+5B a^2 e^2+12B abde-21b^2 B d^2)x^5}{2e^3} - \frac{5b^3(6A a^2 b e^3+6Aa b^2 d e^2+6A b^3 d^2 e+8B a^3 e^3)}{6e^4}}{6e^4}$
norman	$\frac{-60a^6 A e^7+60A a^5 b d e^6+60A a^4 b^2 d^2 e^5+60A a^3 b^3 d^3 e^4+60A a^2 b^4 d^4 e^3+60A a b^5 d^5 e^2+60A b^6 d^6 e+10B a^6 d e^6+24B a^5 b d^2 e^5+45B a^4 b^2 d^2 e^4-45B a^3 b^3 d^3 e^3+30A a b^5 d^4 e^2-6A b^6 d^5 e+B a^6 e^6-12B a^5 b d e^5+45B a^4 b^2 d^2 e^4-45B a^3 b^3 d^3 e^3+30A a b^5 d^4 e^2-6A b^6 d^5 e}{420e^8}$
default	$\frac{-6A a^5 b e^6-30A a^4 b^2 d e^5+60A a^3 b^3 d^2 e^4-60A a^2 b^4 d^3 e^3+30A a b^5 d^4 e^2-6A b^6 d^5 e+B a^6 e^6-12B a^5 b d e^5+45B a^4 b^2 d^2 e^4-45B a^3 b^3 d^3 e^3+30A a b^5 d^4 e^2-6A b^6 d^5 e}{6e^8 (ex+d)^6}$
parallelrisc	Expression too large to display

input

```
int((b*x+a)^6*(B*x+A)/(e*x+d)^8,x,method=_RETURNVERBOSE)
```

output

```
(-b^5*(A*b*e+6*B*a*e-7*B*b*d)/e^2*x^6-3/2*b^4*(2*A*a*b*e^2+2*A*b^2*d*e+5*B
*a^2*e^2+12*B*a*b*d*e-21*B*b^2*d^2)/e^3*x^5-5/6*b^3*(6*A*a^2*b*e^3+6*A*a*b
^2*d*e^2+6*A*b^3*d^2*e+8*B*a^3*e^3+15*B*a^2*b*d*e^2+36*B*a*b^2*d^2*e-77*B*
b^3*d^3)/e^4*x^4-5/12*b^2*(12*A*a^3*b*e^4+12*A*a^2*b^2*d*e^3+12*A*a*b^3*d^
2*e^2+12*A*b^4*d^3*e+9*B*a^4*e^4+16*B*a^3*b*d*e^3+30*B*a^2*b^2*d^2*e^2+72*
B*a*b^3*d^3*e-175*B*b^4*d^4)/e^5*x^3-1/20*b*(60*A*a^4*b*e^5+60*A*a^3*b^2*d
*e^4+60*A*a^2*b^3*d^2*e^3+60*A*a*b^4*d^3*e^2+60*A*b^5*d^4*e+24*B*a^5*e^5+4
5*B*a^4*b*d*e^4+80*B*a^3*b^2*d^2*e^3+150*B*a^2*b^3*d^3*e^2+360*B*a*b^4*d^4
*e-959*B*b^5*d^5)/e^6*x^2-1/60*(60*A*a^5*b*e^6+60*A*a^4*b^2*d*e^5+60*A*a^3
*b^3*d^2*e^4+60*A*a^2*b^4*d^3*e^3+60*A*a*b^5*d^4*e^2+60*A*b^6*d^5*e+10*B*a
^6*e^6+24*B*a^5*b*d*e^5+45*B*a^4*b^2*d^2*e^4+80*B*a^3*b^3*d^3*e^3+150*B*a^
2*b^4*d^4*e^2+360*B*a*b^5*d^5*e-1029*B*b^6*d^6)/e^7*x-1/420*(60*A*a^6*e^7+
60*A*a^5*b*d*e^6+60*A*a^4*b^2*d^2*e^5+60*A*a^3*b^3*d^3*e^4+60*A*a^2*b^4*d^
4*e^3+60*A*a*b^5*d^5*e^2+60*A*b^6*d^6*e+10*B*a^6*d*e^6+24*B*a^5*b*d^2*e^5+
45*B*a^4*b^2*d^3*e^4+80*B*a^3*b^3*d^4*e^3+150*B*a^2*b^4*d^5*e^2+360*B*a*b^
5*d^6*e-1089*B*b^6*d^7)/e^8)/(e*x+d)^7+b^6*B*ln(e*x+d)/e^8
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 939 vs. $2(201) = 402$.

Time = 0.11 (sec) , antiderivative size = 939, normalized size of antiderivative = 4.41

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^8} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^6*(B*x+A)/(e*x+d)^8,x, algorithm="fricas")
```

output

```

1/420*(1089*B*b^6*d^7 - 60*A*a^6*e^7 - 60*(6*B*a*b^5 + A*b^6)*d^6*e - 30*(
5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 - 20*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4*e^3
- 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 - 12*(2*B*a^5*b + 5*A*a^4*b^2)*d^
2*e^5 - 10*(B*a^6 + 6*A*a^5*b)*d*e^6 + 420*(7*B*b^6*d*e^6 - (6*B*a*b^5 + A
*b^6)*e^7)*x^6 + 630*(21*B*b^6*d^2*e^5 - 2*(6*B*a*b^5 + A*b^6)*d*e^6 - (5*
B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^5 + 350*(77*B*b^6*d^3*e^4 - 6*(6*B*a*b^5 + A
*b^6)*d^2*e^5 - 3*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6 - 2*(4*B*a^3*b^3 + 3*A*a
^2*b^4)*e^7)*x^4 + 175*(175*B*b^6*d^4*e^3 - 12*(6*B*a*b^5 + A*b^6)*d^3*e^4
- 6*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 - 4*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e
^6 - 3*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^3 + 21*(959*B*b^6*d^5*e^2 - 60*(
6*B*a*b^5 + A*b^6)*d^4*e^3 - 30*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*e^4 - 20*(4*
B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 - 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^6 -
12*(2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 7*(1029*B*b^6*d^6*e - 60*(6*B*a*b^
5 + A*b^6)*d^5*e^2 - 30*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^3 - 20*(4*B*a^3*b^
3 + 3*A*a^2*b^4)*d^3*e^4 - 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^5 - 12*(2*
B*a^5*b + 5*A*a^4*b^2)*d*e^6 - 10*(B*a^6 + 6*A*a^5*b)*e^7)*x + 420*(B*b^6*
e^7*x^7 + 7*B*b^6*d*e^6*x^6 + 21*B*b^6*d^2*e^5*x^5 + 35*B*b^6*d^3*e^4*x^4
+ 35*B*b^6*d^4*e^3*x^3 + 21*B*b^6*d^5*e^2*x^2 + 7*B*b^6*d^6*e*x + B*b^6*d^
7)*log(e*x + d))/(e^15*x^7 + 7*d*e^14*x^6 + 21*d^2*e^13*x^5 + 35*d^3*e^12*
x^4 + 35*d^4*e^11*x^3 + 21*d^5*e^10*x^2 + 7*d^6*e^9*x + d^7*e^8)

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^8} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**6*(B*x+A)/(e*x+d)**8,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 842 vs. $2(201) = 402$.

Time = 0.06 (sec) , antiderivative size = 842, normalized size of antiderivative = 3.95

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^8} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^8,x, algorithm="maxima")`

output

```
1/420*(1089*B*b^6*d^7 - 60*A*a^6*e^7 - 60*(6*B*a*b^5 + A*b^6)*d^6*e - 30*(
5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 - 20*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4*e^3
- 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 - 12*(2*B*a^5*b + 5*A*a^4*b^2)*d^
2*e^5 - 10*(B*a^6 + 6*A*a^5*b)*d*e^6 + 420*(7*B*b^6*d*e^6 - (6*B*a*b^5 + A
*b^6)*e^7)*x^6 + 630*(21*B*b^6*d^2*e^5 - 2*(6*B*a*b^5 + A*b^6)*d*e^6 - (5*
B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^5 + 350*(77*B*b^6*d^3*e^4 - 6*(6*B*a*b^5 + A
*b^6)*d^2*e^5 - 3*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6 - 2*(4*B*a^3*b^3 + 3*A*a
^2*b^4)*e^7)*x^4 + 175*(175*B*b^6*d^4*e^3 - 12*(6*B*a*b^5 + A*b^6)*d^3*e^4
- 6*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 - 4*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e
^6 - 3*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^3 + 21*(959*B*b^6*d^5*e^2 - 60*(
6*B*a*b^5 + A*b^6)*d^4*e^3 - 30*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*e^4 - 20*(4*
B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 - 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^6 -
12*(2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 7*(1029*B*b^6*d^6*e - 60*(6*B*a*b^
5 + A*b^6)*d^5*e^2 - 30*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^3 - 20*(4*B*a^3*b^
3 + 3*A*a^2*b^4)*d^3*e^4 - 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^5 - 12*(2*
B*a^5*b + 5*A*a^4*b^2)*d*e^6 - 10*(B*a^6 + 6*A*a^5*b)*e^7)*x)/(e^15*x^7 +
7*d*e^14*x^6 + 21*d^2*e^13*x^5 + 35*d^3*e^12*x^4 + 35*d^4*e^11*x^3 + 21*d^
5*e^10*x^2 + 7*d^6*e^9*x + d^7*e^8) + B*b^6*log(e*x + d)/e^8
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 830 vs. $2(201) = 402$.

Time = 0.12 (sec) , antiderivative size = 830, normalized size of antiderivative = 3.90

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^8} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^8,x, algorithm="giac")`

output
$$\begin{aligned} & B*b^6*\log(\text{abs}(e*x + d))/e^8 + 1/420*(420*(7*B*b^6*d*e^5 - 6*B*a*b^5*e^6 - \\ & A*b^6*e^6)*x^6 + 630*(21*B*b^6*d^2*e^4 - 12*B*a*b^5*d*e^5 - 2*A*b^6*d*e^5 \\ & - 5*B*a^2*b^4*e^6 - 2*A*a*b^5*e^6)*x^5 + 350*(77*B*b^6*d^3*e^3 - 36*B*a*b^5 \\ & 5*d^2*e^4 - 6*A*b^6*d^2*e^4 - 15*B*a^2*b^4*d*e^5 - 6*A*a*b^5*d*e^5 - 8*B*a \\ & ^3*b^3*e^6 - 6*A*a^2*b^4*e^6)*x^4 + 175*(175*B*b^6*d^4*e^2 - 72*B*a*b^5*d^3 \\ & 3*e^3 - 12*A*b^6*d^3*e^3 - 30*B*a^2*b^4*d^2*e^4 - 12*A*a*b^5*d^2*e^4 - 16* \\ & B*a^3*b^3*d*e^5 - 12*A*a^2*b^4*d*e^5 - 9*B*a^4*b^2*e^6 - 12*A*a^3*b^3*e^6) \\ & *x^3 + 21*(959*B*b^6*d^5*e - 360*B*a*b^5*d^4*e^2 - 60*A*b^6*d^4*e^2 - 150* \\ & B*a^2*b^4*d^3*e^3 - 60*A*a*b^5*d^3*e^3 - 80*B*a^3*b^3*d^2*e^4 - 60*A*a^2*b \\ & ^4*d^2*e^4 - 45*B*a^4*b^2*d*e^5 - 60*A*a^3*b^3*d*e^5 - 24*B*a^5*b*e^6 - 60 \\ & *A*a^4*b^2*e^6)*x^2 + 7*(1029*B*b^6*d^6 - 360*B*a*b^5*d^5*e - 60*A*b^6*d^5 \\ & *e - 150*B*a^2*b^4*d^4*e^2 - 60*A*a*b^5*d^4*e^2 - 80*B*a^3*b^3*d^3*e^3 - 6 \\ & 0*A*a^2*b^4*d^3*e^3 - 45*B*a^4*b^2*d^2*e^4 - 60*A*a^3*b^3*d^2*e^4 - 24*B*a \\ & ^5*b*d*e^5 - 60*A*a^4*b^2*d*e^5 - 10*B*a^6*e^6 - 60*A*a^5*b*e^6)*x + (1089 \\ & *B*b^6*d^7 - 360*B*a*b^5*d^6*e - 60*A*b^6*d^6*e - 150*B*a^2*b^4*d^5*e^2 - \\ & 60*A*a*b^5*d^5*e^2 - 80*B*a^3*b^3*d^4*e^3 - 60*A*a^2*b^4*d^4*e^3 - 45*B*a^4 \\ & 4*b^2*d^3*e^4 - 60*A*a^3*b^3*d^3*e^4 - 24*B*a^5*b*d^2*e^5 - 60*A*a^4*b^2*d^2 \\ & ^2*e^5 - 10*B*a^6*d*e^6 - 60*A*a^5*b*d*e^6 - 60*A*a^6*e^7)/e/((e*x + d)^7 \\ & *e^7) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 1.07 (sec) , antiderivative size = 1046, normalized size of antiderivative = 4.91

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^8} dx = \text{Too large to display}$$

input `int(((A + B*x)*(a + b*x)^6)/(d + e*x)^8,x)`

output

```

-((A*a^6*e^7)/7 - (363*B*b^6*d^7)/140 + (A*b^6*d^6*e)/7 + (B*a^6*d*e^6)/42
- B*b^6*d^7*log(d + e*x) + (B*a^6*e^7*x)/6 + A*b^6*e^7*x^6 - (343*B*b^6*d
^6*e*x)/20 + (A*a*b^5*d^5*e^2)/7 + (2*B*a^5*b*d^2*e^5)/35 + 3*A*a*b^5*e^7*
x^5 + (6*B*a^5*b*e^7*x^2)/5 + 6*B*a*b^5*e^7*x^6 + A*b^6*d^5*e^2*x + 3*A*b^
6*d*e^6*x^5 - 7*B*b^6*d*e^6*x^6 - B*b^6*e^7*x^7*log(d + e*x) + (A*a^2*b^4*
d^4*e^3)/7 + (A*a^3*b^3*d^3*e^4)/7 + (A*a^4*b^2*d^2*e^5)/7 + (5*B*a^2*b^4*
d^5*e^2)/14 + (4*B*a^3*b^3*d^4*e^3)/21 + (3*B*a^4*b^2*d^3*e^4)/28 + 3*A*a^
4*b^2*e^7*x^2 + 5*A*a^3*b^3*e^7*x^3 + 5*A*a^2*b^4*e^7*x^4 + (15*B*a^4*b^2*
e^7*x^3)/4 + (20*B*a^3*b^3*e^7*x^4)/3 + (15*B*a^2*b^4*e^7*x^5)/2 + 3*A*b^6
*d^4*e^3*x^2 + 5*A*b^6*d^3*e^4*x^3 + 5*A*b^6*d^2*e^5*x^4 - (959*B*b^6*d^5*
e^2*x^2)/20 - (875*B*b^6*d^4*e^3*x^3)/12 - (385*B*b^6*d^3*e^4*x^4)/6 - (63
*B*b^6*d^2*e^5*x^5)/2 + (A*a^5*b*d*e^6)/7 + (6*B*a*b^5*d^6*e)/7 + A*a^5*b*
e^7*x + A*a^2*b^4*d^3*e^4*x + A*a^3*b^3*d^2*e^5*x + 3*A*a*b^5*d^3*e^4*x^2
+ 3*A*a^3*b^3*d*e^6*x^2 + 5*A*a*b^5*d^2*e^5*x^3 + 5*A*a^2*b^4*d*e^6*x^3 +
(5*B*a^2*b^4*d^4*e^3*x)/2 + (4*B*a^3*b^3*d^3*e^4*x)/3 + (3*B*a^4*b^2*d^2*
e^5*x)/4 + 18*B*a*b^5*d^4*e^3*x^2 + (9*B*a^4*b^2*d*e^6*x^2)/4 + 30*B*a*b^5*
d^3*e^4*x^3 + (20*B*a^3*b^3*d*e^6*x^3)/3 + 30*B*a*b^5*d^2*e^5*x^4 + (25*B*
a^2*b^4*d*e^6*x^4)/2 - 21*B*b^6*d^5*e^2*x^2*log(d + e*x) - 35*B*b^6*d^4*e^
3*x^3*log(d + e*x) - 35*B*b^6*d^3*e^4*x^4*log(d + e*x) - 21*B*b^6*d^2*e^5*
x^5*log(d + e*x) + (2*B*a^5*b*d*e^6*x)/5 - 7*B*b^6*d^6*e*x*log(d + e*x)...

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 638, normalized size of antiderivative = 3.00

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^8} dx$$

$$= \frac{420 \log(ex + d) b^7 d e^7 x^7 + 8820 \log(ex + d) b^7 d^3 e^5 x^5 + 8820 \log(ex + d) b^7 d^6 e^2 x^2 - 60 a^7 d e^7 + 2940 \log($$

input

```
int((b*x+a)^6*(B*x+A)/(e*x+d)^8,x)
```

output

```
(420*log(d + e*x)*b**7*d**8 + 2940*log(d + e*x)*b**7*d**7*e*x + 8820*log(d
+ e*x)*b**7*d**6*e**2*x**2 + 14700*log(d + e*x)*b**7*d**5*e**3*x**3 + 147
00*log(d + e*x)*b**7*d**4*e**4*x**4 + 8820*log(d + e*x)*b**7*d**3*e**5*x**
5 + 2940*log(d + e*x)*b**7*d**2*e**6*x**6 + 420*log(d + e*x)*b**7*d*e**7*x
**7 - 60*a**7*d*e**7 - 70*a**6*b*d**2*e**6 - 490*a**6*b*d*e**7*x - 84*a**5
*b**2*d**3*e**5 - 588*a**5*b**2*d**2*e**6*x - 1764*a**5*b**2*d*e**7*x**2 -
105*a**4*b**3*d**4*e**4 - 735*a**4*b**3*d**3*e**5*x - 2205*a**4*b**3*d**2
*e**6*x**2 - 3675*a**4*b**3*d*e**7*x**3 - 140*a**3*b**4*d**5*e**3 - 980*a*
*3*b**4*d**4*e**4*x - 2940*a**3*b**4*d**3*e**5*x**2 - 4900*a**3*b**4*d**2*
e**6*x**3 - 4900*a**3*b**4*d*e**7*x**4 - 210*a**2*b**5*d**6*e**2 - 1470*a*
*2*b**5*d**5*e**3*x - 4410*a**2*b**5*d**4*e**4*x**2 - 7350*a**2*b**5*d**3*
e**5*x**3 - 7350*a**2*b**5*d**2*e**6*x**4 - 4410*a**2*b**5*d*e**7*x**5 + 4
20*a*b**6*e**8*x**7 + 669*b**7*d**8 + 4263*b**7*d**7*e*x + 11319*b**7*d**6
*e**2*x**2 + 15925*b**7*d**5*e**3*x**3 + 12250*b**7*d**4*e**4*x**4 + 4410*
b**7*d**3*e**5*x**5 - 420*b**7*d*e**7*x**7)/(420*d*e**8*(d**7 + 7*d**6*e*x
+ 21*d**5*e**2*x**2 + 35*d**4*e**3*x**3 + 35*d**3*e**4*x**4 + 21*d**2*e**
5*x**5 + 7*d*e**6*x**6 + e**7*x**7))
```

3.58 $\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^9} dx$

Optimal result	587
Mathematica [B] (verified)	587
Rubi [A] (verified)	588
Maple [B] (verified)	589
Fricas [B] (verification not implemented)	590
Sympy [F(-1)]	591
Maxima [B] (verification not implemented)	591
Giac [B] (verification not implemented)	592
Mupad [B] (verification not implemented)	593
Reduce [B] (verification not implemented)	594

Optimal result

Integrand size = 20, antiderivative size = 86

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^9} dx = -\frac{(Bd - Ae)(a+bx)^7}{8e(bd - ae)(d+ex)^8} + \frac{(7bBd + Abe - 8aBe)(a+bx)^7}{56e(bd - ae)^2(d+ex)^7}$$

output

```
-1/8*(-A*e+B*d)*(b*x+a)^7/e/(-a*e+b*d)/(e*x+d)^8+1/56*(A*b*e-8*B*a*e+7*B*b*d)*(b*x+a)^7/e/(-a*e+b*d)^2/(e*x+d)^7
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 597 vs. 2(86) = 172.

Time = 0.16 (sec) , antiderivative size = 597, normalized size of antiderivative = 6.94

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^9} dx = \frac{a^6e^6(7Ae + B(d + 8ex)) + 2a^5be^5(3Ae(d + 8ex) + B(d^2 + 8dex + 28e^2x^2)) + a^4b^2e^4(5Ae(d^2 + 8dex$$

input

```
Integrate[((a + b*x)^6*(A + B*x))/(d + e*x)^9,x]
```


output

```

-1/56*(a^6*e^6*(7*A*e + B*(d + 8*e*x)) + 2*a^5*b*e^5*(3*A*e*(d + 8*e*x) +
B*(d^2 + 8*d*e*x + 28*e^2*x^2)) + a^4*b^2*e^4*(5*A*e*(d^2 + 8*d*e*x + 28*
e^2*x^2) + 3*B*(d^3 + 8*d^2*e*x + 28*d*e^2*x^2 + 56*e^3*x^3)) + 4*a^3*b^3*
e^3*(A*e*(d^3 + 8*d^2*e*x + 28*d*e^2*x^2 + 56*e^3*x^3) + B*(d^4 + 8*d^3*
e*x + 28*d^2*e^2*x^2 + 56*d*e^3*x^3 + 70*e^4*x^4)) + a^2*b^4*e^2*(3*A*e*(d^4
+ 8*d^3*e*x + 28*d^2*e^2*x^2 + 56*d*e^3*x^3 + 70*e^4*x^4) + 5*B*(d^5 + 8*
d^4*e*x + 28*d^3*e^2*x^2 + 56*d^2*e^3*x^3 + 70*d*e^4*x^4 + 56*e^5*x^5)) + 2
*a*b^5*e*(A*e*(d^5 + 8*d^4*e*x + 28*d^3*e^2*x^2 + 56*d^2*e^3*x^3 + 70*d*
e^4*x^4 + 56*e^5*x^5) + 3*B*(d^6 + 8*d^5*e*x + 28*d^4*e^2*x^2 + 56*d^3*
e^3*x^3 + 70*d^2*e^4*x^4 + 56*d*e^5*x^5 + 28*e^6*x^6)) + b^6*(A*e*(d^6 + 8*
d^5*
e*x + 28*d^4*e^2*x^2 + 56*d^3*e^3*x^3 + 70*d^2*e^4*x^4 + 56*d*e^5*x^5 + 28
*e^6*x^6) + 7*B*(d^7 + 8*d^6*e*x + 28*d^5*e^2*x^2 + 56*d^4*e^3*x^3 + 70*d^
3*
e^4*x^4 + 56*d^2*e^5*x^5 + 28*d*e^6*x^6 + 8*e^7*x^7)))/(e^8*(d + e*x)^8)

```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {87, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a+bx)^6(A+Bx)}{(d+ex)^9} dx \\
 & \quad \downarrow 87 \\
 & \frac{(-8aBe + Abe + 7bBd) \int \frac{(a+bx)^6}{(d+ex)^8} dx}{8e(bd - ae)} - \frac{(a+bx)^7(Bd - Ae)}{8e(d+ex)^8(bd - ae)} \\
 & \quad \downarrow 48 \\
 & \frac{(a+bx)^7(-8aBe + Abe + 7bBd)}{56e(d+ex)^7(bd - ae)^2} - \frac{(a+bx)^7(Bd - Ae)}{8e(d+ex)^8(bd - ae)}
 \end{aligned}$$

input

```
Int[((a + b*x)^6*(A + B*x))/(d + e*x)^9,x]
```

```
output -1/8*((B*d - A*e)*(a + b*x)^7)/(e*(b*d - a*e)*(d + e*x)^8) + ((7*b*B*d + A
*b*e - 8*a*B*e)*(a + b*x)^7)/(56*e*(b*d - a*e)^2*(d + e*x)^7)
```

Defintions of rubi rules used

```
rule 48 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && ( !LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 779 vs. 2(82) = 164.

Time = 0.24 (sec) , antiderivative size = 780, normalized size of antiderivative = 9.07

method	result
risch	$\frac{-\frac{b^6 B x^7}{e} - \frac{b^5 (A b e + 6 B a e + 7 B b d) x^6}{2 e^2} - \frac{b^4 (2 A a b e^2 + A b^2 d e + 5 B a^2 e^2 + 6 B a b d e + 7 b^2 B d^2) x^5}{e^3} - \frac{5 b^3 (3 A a^2 b e^3 + 2 A a b^2 d e^2 + A b^3 d^2 e + 4 B a^2 b^2 d e^2 + 3 A a^2 b^2 d^2 e^2 + 2 A a^2 b^2 d^2 e^2 + 2 A a^2 b^2 d^2 e^2 + 2 A a^2 b^2 d^2 e^2)}{4 e^4}}{e^4}$
norman	$\frac{-\frac{b^6 B x^7}{e} - \frac{(A b^6 e + 6 B a b^5 e + 7 b^6 B d) x^6}{2 e^2} - \frac{(2 A a b^5 e^2 + A b^6 d e + 5 B a^2 b^4 e^2 + 6 B a b^5 d e + 7 b^6 B d^2) x^5}{e^3} - \frac{5 (3 A a^2 b^4 e^3 + 2 A a b^5 d e^2 + A b^6 d^2 e^2 + 3 A a^2 b^2 d e^2 + 2 A a^2 b^2 d^2 e^2 + 2 A a^2 b^2 d^2 e^2 + 2 A a^2 b^2 d^2 e^2 + 2 A a^2 b^2 d^2 e^2)}{4 e^4}}{e^4}$
default	$\frac{b(5 A a^4 b e^5 - 20 A a^3 b^2 d e^4 + 30 A a^2 b^3 d^2 e^3 - 20 A a b^4 d^3 e^2 + 5 A b^5 d^4 e + 2 B a^5 e^5 - 15 B a^4 b d e^4 + 40 B a^3 b^2 d^2 e^3 - 50 B a^2 b^3 d^3 e^2 + 35 B a b^4 d^4 e + 5 B b^5 d^5)}{2 e^8 (e x + d)^6}$
gosper	$-\frac{56 B x^7 b^6 e^7 + 28 A x^6 b^6 e^7 + 168 B x^6 a b^5 e^7 + 196 B x^6 b^6 d e^6 + 112 A x^5 a b^5 e^7 + 56 A x^5 b^6 d e^6 + 280 B x^5 a^2 b^4 e^7 + 336 B x^5 a b^5 d e^6 + 56 A x^4 a^2 b^3 d^3 e^6 + 112 A x^4 a b^4 d^4 e^5 + 56 A x^4 b^5 d^5 e^4 + 280 B x^4 a^2 b^2 d^2 e^5 + 112 B x^4 a b^3 d^3 e^4 + 56 B x^4 b^4 d^4 e^3 + 280 B x^4 a^2 b^2 d^2 e^3 + 112 B x^4 a b^3 d^3 e^2 + 56 B x^4 b^4 d^4 e^1 + 280 B x^4 a^2 b^2 d^2 e^1 + 112 B x^4 a b^3 d^3 e^0 + 56 B x^4 b^4 d^4 e^0}{2 e^8 (e x + d)^6}$
parallelrisch	$-\frac{56 B x^7 b^6 e^7 + 28 A x^6 b^6 e^7 + 168 B x^6 a b^5 e^7 + 196 B x^6 b^6 d e^6 + 112 A x^5 a b^5 e^7 + 56 A x^5 b^6 d e^6 + 280 B x^5 a^2 b^4 e^7 + 336 B x^5 a b^5 d e^6 + 56 A x^4 a^2 b^3 d^3 e^6 + 112 A x^4 a b^4 d^4 e^5 + 56 A x^4 b^5 d^5 e^4 + 280 B x^4 a^2 b^2 d^2 e^5 + 112 B x^4 a b^3 d^3 e^4 + 56 B x^4 b^4 d^4 e^3 + 280 B x^4 a^2 b^2 d^2 e^3 + 112 B x^4 a b^3 d^3 e^2 + 56 B x^4 b^4 d^4 e^1 + 280 B x^4 a^2 b^2 d^2 e^1 + 112 B x^4 a b^3 d^3 e^0 + 56 B x^4 b^4 d^4 e^0}{2 e^8 (e x + d)^6}$
orering	$-\frac{56 B x^7 b^6 e^7 + 28 A x^6 b^6 e^7 + 168 B x^6 a b^5 e^7 + 196 B x^6 b^6 d e^6 + 112 A x^5 a b^5 e^7 + 56 A x^5 b^6 d e^6 + 280 B x^5 a^2 b^4 e^7 + 336 B x^5 a b^5 d e^6 + 56 A x^4 a^2 b^3 d^3 e^6 + 112 A x^4 a b^4 d^4 e^5 + 56 A x^4 b^5 d^5 e^4 + 280 B x^4 a^2 b^2 d^2 e^5 + 112 B x^4 a b^3 d^3 e^4 + 56 B x^4 b^4 d^4 e^3 + 280 B x^4 a^2 b^2 d^2 e^3 + 112 B x^4 a b^3 d^3 e^2 + 56 B x^4 b^4 d^4 e^1 + 280 B x^4 a^2 b^2 d^2 e^1 + 112 B x^4 a b^3 d^3 e^0 + 56 B x^4 b^4 d^4 e^0}{2 e^8 (e x + d)^6}$

```
input int((b*x+a)^6*(B*x+A)/(e*x+d)^9,x,method=_RETURNVERBOSE)
```

output

```
(-b^6*B/e*x^7-1/2*b^5*(A*b*e+6*B*a*e+7*B*b*d)/e^2*x^6-b^4*(2*A*a*b*e^2+A*b^2*d*e+5*B*a^2*e^2+6*B*a*b*d*e+7*B*b^2*d^2)/e^3*x^5-5/4*b^3*(3*A*a^2*b*e^3+2*A*a*b^2*d*e^2+A*b^3*d^2*e+4*B*a^3*e^3+5*B*a^2*b*d*e^2+6*B*a*b^2*d^2*e+7*B*b^3*d^3)/e^4*x^4-b^2*(4*A*a^3*b*e^4+3*A*a^2*b^2*d*e^3+2*A*a*b^3*d^2*e^2+A*b^4*d^3*e+3*B*a^4*e^4+4*B*a^3*b*d*e^3+5*B*a^2*b^2*d^2*e^2+6*B*a*b^3*d^3*e+7*B*b^4*d^4)/e^5*x^3-1/2*b*(5*A*a^4*b*e^5+4*A*a^3*b^2*d*e^4+3*A*a^2*b^3*d^2*e^3+2*A*a*b^4*d^3*e^2+A*b^5*d^4*e+2*B*a^5*e^5+3*B*a^4*b*d*e^4+4*B*a^3*b^2*d^2*e^3+5*B*a^2*b^3*d^3*e^2+6*B*a*b^4*d^4*e+7*B*b^5*d^5)/e^6*x^2-1/7*(6*A*a^5*b*e^6+5*A*a^4*b^2*d*e^5+4*A*a^3*b^3*d^2*e^4+3*A*a^2*b^4*d^3*e^3+2*A*a*b^5*d^4*e^2+A*b^6*d^5*e+B*a^6*e^6+2*B*a^5*b*d*e^5+3*B*a^4*b^2*d^2*e^4+4*B*a^3*b^3*d^3*e^3+5*B*a^2*b^4*d^4*e^2+6*B*a*b^5*d^5*e+7*B*b^6*d^6)/e^7*x-1/56*(7*A*a^6*e^7+6*A*a^5*b*d*e^6+5*A*a^4*b^2*d^2*e^5+4*A*a^3*b^3*d^3*e^4+3*A*a^2*b^4*d^4*e^3+2*A*a*b^5*d^5*e^2+A*b^6*d^6*e+B*a^6*d*e^6+2*B*a^5*b*d^2*e^5+3*B*a^4*b^2*d^3*e^4+4*B*a^3*b^3*d^4*e^3+5*B*a^2*b^4*d^5*e^2+6*B*a*b^5*d^6*e+7*B*b^6*d^7)/e^8)/(e*x+d)^8
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 823 vs. $2(82) = 164$.

Time = 0.09 (sec) , antiderivative size = 823, normalized size of antiderivative = 9.57

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^9} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^6*(B*x+A)/(e*x+d)^9,x, algorithm="fricas")
```

output

```
-1/56*(56*B*b^6*e^7*x^7 + 7*B*b^6*d^7 + 7*A*a^6*e^7 + (6*B*a*b^5 + A*b^6)*
d^6*e + (5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 + (4*B*a^3*b^3 + 3*A*a^2*b^4)*d^
4*e^3 + (3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 + (2*B*a^5*b + 5*A*a^4*b^2)*d^
2*e^5 + (B*a^6 + 6*A*a^5*b)*d*e^6 + 28*(7*B*b^6*d*e^6 + (6*B*a*b^5 + A*b^6
)*e^7)*x^6 + 56*(7*B*b^6*d^2*e^5 + (6*B*a*b^5 + A*b^6)*d*e^6 + (5*B*a^2*b^
4 + 2*A*a*b^5)*e^7)*x^5 + 70*(7*B*b^6*d^3*e^4 + (6*B*a*b^5 + A*b^6)*d^2*e^
5 + (5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6 + (4*B*a^3*b^3 + 3*A*a^2*b^4)*e^7)*x^4
+ 56*(7*B*b^6*d^4*e^3 + (6*B*a*b^5 + A*b^6)*d^3*e^4 + (5*B*a^2*b^4 + 2*A*
a*b^5)*d^2*e^5 + (4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^6 + (3*B*a^4*b^2 + 4*A*a^
3*b^3)*e^7)*x^3 + 28*(7*B*b^6*d^5*e^2 + (6*B*a*b^5 + A*b^6)*d^4*e^3 + (5*B
*a^2*b^4 + 2*A*a*b^5)*d^3*e^4 + (4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + (3*B
*a^4*b^2 + 4*A*a^3*b^3)*d*e^6 + (2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 8*(7*
B*b^6*d^6*e + (6*B*a*b^5 + A*b^6)*d^5*e^2 + (5*B*a^2*b^4 + 2*A*a*b^5)*d^4*
e^3 + (4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 + (3*B*a^4*b^2 + 4*A*a^3*b^3)*d^
2*e^5 + (2*B*a^5*b + 5*A*a^4*b^2)*d*e^6 + (B*a^6 + 6*A*a^5*b)*e^7)*x)/(e^1
6*x^8 + 8*d*e^15*x^7 + 28*d^2*e^14*x^6 + 56*d^3*e^13*x^5 + 70*d^4*e^12*x^4
+ 56*d^5*e^11*x^3 + 28*d^6*e^10*x^2 + 8*d^7*e^9*x + d^8*e^8)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^9} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**6*(B*x+A)/(e*x+d)**9,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 823 vs. 2(82) = 164.

Time = 0.08 (sec) , antiderivative size = 823, normalized size of antiderivative = 9.57

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^9} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^9,x, algorithm="maxima")`

output

$$\begin{aligned}
 & -1/56*(56*B*b^6*e^7*x^7 + 7*B*b^6*d^7 + 7*A*a^6*e^7 + (6*B*a*b^5 + A*b^6)* \\
 & d^6*e + (5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 + (4*B*a^3*b^3 + 3*A*a^2*b^4)*d^ \\
 & 4*e^3 + (3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 + (2*B*a^5*b + 5*A*a^4*b^2)*d^ \\
 & 2*e^5 + (B*a^6 + 6*A*a^5*b)*d*e^6 + 28*(7*B*b^6*d*e^6 + (6*B*a*b^5 + A*b^6) \\
 &)*e^7)*x^6 + 56*(7*B*b^6*d^2*e^5 + (6*B*a*b^5 + A*b^6)*d*e^6 + (5*B*a^2*b^ \\
 & 4 + 2*A*a*b^5)*e^7)*x^5 + 70*(7*B*b^6*d^3*e^4 + (6*B*a*b^5 + A*b^6)*d^2*e^ \\
 & 5 + (5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6 + (4*B*a^3*b^3 + 3*A*a^2*b^4)*e^7)*x^4 \\
 & + 56*(7*B*b^6*d^4*e^3 + (6*B*a*b^5 + A*b^6)*d^3*e^4 + (5*B*a^2*b^4 + 2*A* \\
 & a*b^5)*d^2*e^5 + (4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^6 + (3*B*a^4*b^2 + 4*A*a^ \\
 & 3*b^3)*e^7)*x^3 + 28*(7*B*b^6*d^5*e^2 + (6*B*a*b^5 + A*b^6)*d^4*e^3 + (5*B \\
 & *a^2*b^4 + 2*A*a*b^5)*d^3*e^4 + (4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + (3*B \\
 & *a^4*b^2 + 4*A*a^3*b^3)*d*e^6 + (2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 8*(7* \\
 & B*b^6*d^6*e + (6*B*a*b^5 + A*b^6)*d^5*e^2 + (5*B*a^2*b^4 + 2*A*a*b^5)*d^4* \\
 & e^3 + (4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 + (3*B*a^4*b^2 + 4*A*a^3*b^3)*d^ \\
 & 2*e^5 + (2*B*a^5*b + 5*A*a^4*b^2)*d*e^6 + (B*a^6 + 6*A*a^5*b)*e^7)*x)/(e^1 \\
 & 6*x^8 + 8*d*e^15*x^7 + 28*d^2*e^14*x^6 + 56*d^3*e^13*x^5 + 70*d^4*e^12*x^4 \\
 & + 56*d^5*e^11*x^3 + 28*d^6*e^10*x^2 + 8*d^7*e^9*x + d^8*e^8)
 \end{aligned}$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 910 vs. $2(82) = 164$.

Time = 0.13 (sec) , antiderivative size = 910, normalized size of antiderivative = 10.58

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^9} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^9,x, algorithm="giac")`

output

```

-1/56*(56*B*b^6*e^7*x^7 + 196*B*b^6*d*e^6*x^6 + 168*B*a*b^5*e^7*x^6 + 28*A
*b^6*e^7*x^6 + 392*B*b^6*d^2*e^5*x^5 + 336*B*a*b^5*d*e^6*x^5 + 56*A*b^6*d*
e^6*x^5 + 280*B*a^2*b^4*e^7*x^5 + 112*A*a*b^5*e^7*x^5 + 490*B*b^6*d^3*e^4*
x^4 + 420*B*a*b^5*d^2*e^5*x^4 + 70*A*b^6*d^2*e^5*x^4 + 350*B*a^2*b^4*d*e^6
*x^4 + 140*A*a*b^5*d*e^6*x^4 + 280*B*a^3*b^3*e^7*x^4 + 210*A*a^2*b^4*e^7*x
^4 + 392*B*b^6*d^4*e^3*x^3 + 336*B*a*b^5*d^3*e^4*x^3 + 56*A*b^6*d^3*e^4*x^
3 + 280*B*a^2*b^4*d^2*e^5*x^3 + 112*A*a*b^5*d^2*e^5*x^3 + 224*B*a^3*b^3*d*
e^6*x^3 + 168*A*a^2*b^4*d*e^6*x^3 + 168*B*a^4*b^2*e^7*x^3 + 224*A*a^3*b^3*
e^7*x^3 + 196*B*b^6*d^5*e^2*x^2 + 168*B*a*b^5*d^4*e^3*x^2 + 28*A*b^6*d^4*e
^3*x^2 + 140*B*a^2*b^4*d^3*e^4*x^2 + 56*A*a*b^5*d^3*e^4*x^2 + 112*B*a^3*b^
3*d^2*e^5*x^2 + 84*A*a^2*b^4*d^2*e^5*x^2 + 84*B*a^4*b^2*d*e^6*x^2 + 112*A*
a^3*b^3*d*e^6*x^2 + 56*B*a^5*b*e^7*x^2 + 140*A*a^4*b^2*e^7*x^2 + 56*B*b^6*
d^6*e*x + 48*B*a*b^5*d^5*e^2*x + 8*A*b^6*d^5*e^2*x + 40*B*a^2*b^4*d^4*e^3*
x + 16*A*a*b^5*d^4*e^3*x + 32*B*a^3*b^3*d^3*e^4*x + 24*A*a^2*b^4*d^3*e^4*x
+ 24*B*a^4*b^2*d^2*e^5*x + 32*A*a^3*b^3*d^2*e^5*x + 16*B*a^5*b*d*e^6*x +
40*A*a^4*b^2*d*e^6*x + 8*B*a^6*e^7*x + 48*A*a^5*b*e^7*x + 7*B*b^6*d^7 + 6*
B*a*b^5*d^6*e + A*b^6*d^6*e + 5*B*a^2*b^4*d^5*e^2 + 2*A*a*b^5*d^5*e^2 + 4*
B*a^3*b^3*d^4*e^3 + 3*A*a^2*b^4*d^4*e^3 + 3*B*a^4*b^2*d^3*e^4 + 4*A*a^3*b^
3*d^3*e^4 + 2*B*a^5*b*d^2*e^5 + 5*A*a^4*b^2*d^2*e^5 + B*a^6*d*e^6 + 6*A*a^
5*b*d*e^6 + 7*A*a^6*e^7)/((e*x + d)^8*e^8)

```

Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 854, normalized size of antiderivative = 9.93

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^9} dx =$$

$$\frac{B a^6 d e^6 + 7 A a^6 e^7 + 2 B a^5 b d^2 e^5 + 6 A a^5 b d e^6 + 3 B a^4 b^2 d^3 e^4 + 5 A a^4 b^2 d^2 e^5 + 4 B a^3 b^3 d^4 e^3 + 4 A a^3 b^3 d^3 e^4 + 5 B a^2 b^4 d^5 e^2 + 3 A a^2 b^4 d^4 e^3 + 2 B a^5 b d^2 e^5 + 5 A a^4 b^2 d^2 e^5 + B a^6 d e^6 + 6 A a^5 b d e^6 + 7 A a^6 e^7}{56 e^8}$$

input

```
int(((A + B*x)*(a + b*x)^6)/(d + e*x)^9,x)
```

output

```

-((7*A*a^6*e^7 + 7*B*b^6*d^7 + A*b^6*d^6*e + B*a^6*d*e^6 + 2*A*a*b^5*d^5*e
^2 + 2*B*a^5*b*d^2*e^5 + 3*A*a^2*b^4*d^4*e^3 + 4*A*a^3*b^3*d^3*e^4 + 5*A*a
^4*b^2*d^2*e^5 + 5*B*a^2*b^4*d^5*e^2 + 4*B*a^3*b^3*d^4*e^3 + 3*B*a^4*b^2*d
^3*e^4 + 6*A*a^5*b*d*e^6 + 6*B*a*b^5*d^6*e)/(56*e^8) + (x*(B*a^6*e^6 + 7*B
*b^6*d^6 + 6*A*a^5*b*e^6 + A*b^6*d^5*e + 2*A*a*b^5*d^4*e^2 + 5*A*a^4*b^2*d
*e^5 + 3*A*a^2*b^4*d^3*e^3 + 4*A*a^3*b^3*d^2*e^4 + 5*B*a^2*b^4*d^4*e^2 + 4
*B*a^3*b^3*d^3*e^3 + 3*B*a^4*b^2*d^2*e^4 + 6*B*a*b^5*d^5*e + 2*B*a^5*b*d*e
^5))/(7*e^7) + (5*b^3*x^4*(4*B*a^3*e^3 + 7*B*b^3*d^3 + 3*A*a^2*b*e^3 + A*b
^3*d^2*e + 2*A*a*b^2*d*e^2 + 6*B*a*b^2*d^2*e + 5*B*a^2*b*d*e^2))/(4*e^4) +
(b^5*x^6*(A*b*e + 6*B*a*e + 7*B*b*d))/(2*e^2) + (b*x^2*(2*B*a^5*e^5 + 7*B
*b^5*d^5 + 5*A*a^4*b*e^5 + A*b^5*d^4*e + 2*A*a*b^4*d^3*e^2 + 4*A*a^3*b^2*d
*e^4 + 3*A*a^2*b^3*d^2*e^3 + 5*B*a^2*b^3*d^3*e^2 + 4*B*a^3*b^2*d^2*e^3 + 6
*B*a*b^4*d^4*e + 3*B*a^4*b*d*e^4))/(2*e^6) + (b^2*x^3*(3*B*a^4*e^4 + 7*B*b
^4*d^4 + 4*A*a^3*b*e^4 + A*b^4*d^3*e + 2*A*a*b^3*d^2*e^2 + 3*A*a^2*b^2*d*e
^3 + 5*B*a^2*b^2*d^2*e^2 + 6*B*a*b^3*d^3*e + 4*B*a^3*b*d*e^3))/e^5 + (b^4*
x^5*(5*B*a^2*e^2 + 7*B*b^2*d^2 + 2*A*a*b*e^2 + A*b^2*d*e + 6*B*a*b*d*e))/e
^3 + (B*b^6*x^7)/e)/(d^8 + e^8*x^8 + 8*d*e^7*x^7 + 28*d^6*e^2*x^2 + 56*d^5
*e^3*x^3 + 70*d^4*e^4*x^4 + 56*d^3*e^5*x^5 + 28*d^2*e^6*x^6 + 8*d^7*e*x)

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 504, normalized size of antiderivative = 5.86

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^9} dx$$

$$= \frac{b^7 e^7 x^8 - 28 a b^6 d e^6 x^6 - 56 a^2 b^5 d e^6 x^5 - 56 a b^6 d^2 e^5 x^5 - 70 a^3 b^4 d e^6 x^4 - 70 a^2 b^5 d^2 e^5 x^4 - 70 a b^6 d^3 e^4 x^4 - 50 a^4 b^3 d^4 e^3 x^3 - 35 a^5 b^2 d^4 e^3 x^2 - 35 a^6 b d^4 e^3 x^2 - 35 a^7 d^4 e^3 x^2 + 70 a^4 b^3 d^4 e^3 x^2 - 35 a^5 b^2 d^4 e^3 x^2 - 35 a^6 b d^4 e^3 x^2 - 35 a^7 d^4 e^3 x^2}{(d+ex)^9}$$

input

```
int((b*x+a)^6*(B*x+A)/(e*x+d)^9,x)
```

output

```
( - a**7*d**6 - a**6*b*d**2*e**5 - 8*a**6*b*d**6*x - a**5*b**2*d**3*e**4 - 8*a**5*b**2*d**2*e**5*x - 28*a**5*b**2*d**6*x**2 - a**4*b**3*d**4*e**3 - 8*a**4*b**3*d**3*e**4*x - 28*a**4*b**3*d**2*e**5*x**2 - 56*a**4*b**3*d**6*x**3 - a**3*b**4*d**5*e**2 - 8*a**3*b**4*d**4*e**3*x - 28*a**3*b**4*d**3*e**4*x**2 - 56*a**3*b**4*d**2*e**5*x**3 - 70*a**3*b**4*d**6*x**4 - a**2*b**5*d**6*e - 8*a**2*b**5*d**5*e**2*x - 28*a**2*b**5*d**4*e**3*x**2 - 56*a**2*b**5*d**3*e**4*x**3 - 70*a**2*b**5*d**2*e**5*x**4 - 56*a**2*b**5*d**6*x**5 - a*b**6*d**7 - 8*a*b**6*d**6*e*x - 28*a*b**6*d**5*e**2*x**2 - 56*a*b**6*d**4*e**3*x**3 - 70*a*b**6*d**3*e**4*x**4 - 56*a*b**6*d**2*e**5*x**5 - 28*a*b**6*d**6*x**6 + b**7*e**7*x**8)/(8*d**7*(d**8 + 8*d**7*e*x + 28*d**6*e**2*x**2 + 56*d**5*e**3*x**3 + 70*d**4*e**4*x**4 + 56*d**3*e**5*x**5 + 28*d**2*e**6*x**6 + 8*d**7*x**7 + e**8*x**8))
```


3.59 $\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{10}} dx$

Optimal result	596
Mathematica [B] (verified)	596
Rubi [A] (verified)	597
Maple [B] (verified)	599
Fricas [B] (verification not implemented)	600
Sympy [F(-1)]	600
Maxima [B] (verification not implemented)	601
Giac [B] (verification not implemented)	602
Mupad [B] (verification not implemented)	603
Reduce [B] (verification not implemented)	604

Optimal result

Integrand size = 20, antiderivative size = 135

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{10}} dx = -\frac{(Bd - Ae)(a+bx)^7}{9e(bd - ae)(d+ex)^9} + \frac{(7bBd + 2Abe - 9aBe)(a+bx)^7}{72e(bd - ae)^2(d+ex)^8} + \frac{b(7bBd + 2Abe - 9aBe)(a+bx)^7}{504e(bd - ae)^3(d+ex)^7}$$

output

```
-1/9*(-A*e+B*d)*(b*x+a)^7/e/(-a*e+b*d)/(e*x+d)^9+1/72*(2*A*b*e-9*B*a*e+7*B*b*d)*(b*x+a)^7/e/(-a*e+b*d)^2/(e*x+d)^8+1/504*b*(2*A*b*e-9*B*a*e+7*B*b*d)*(b*x+a)^7/e/(-a*e+b*d)^3/(e*x+d)^7
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 603 vs. 2(135) = 270.

Time = 0.17 (sec) , antiderivative size = 603, normalized size of antiderivative = 4.47

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{10}} dx = -\frac{7a^6e^6(8Ae + B(d + 9ex)) + 6a^5be^5(7Ae(d + 9ex) + 2B(d^2 + 9dex + 36e^2x^2)) + 15a^4b^2e^4(2Ae(d^2 + 9dex + 36e^2x^2)) + 6a^3b^3e^3(2Ae(d^2 + 9dex + 36e^2x^2)) + 3a^2b^4e^2(2Ae(d^2 + 9dex + 36e^2x^2)) + 3ab^5e(2Ae(d^2 + 9dex + 36e^2x^2)) + b^6(2Ae(d^2 + 9dex + 36e^2x^2))}{(d+ex)^7}$$

input `Integrate[((a + b*x)^6*(A + B*x))/(d + e*x)^10,x]`

output

$$\begin{aligned} & -1/504*(7*a^6*e^6*(8*A*e + B*(d + 9*e*x)) + 6*a^5*b*e^5*(7*A*e*(d + 9*e*x) \\ & + 2*B*(d^2 + 9*d*e*x + 36*e^2*x^2)) + 15*a^4*b^2*e^4*(2*A*e*(d^2 + 9*d*e*x \\ & + 36*e^2*x^2) + B*(d^3 + 9*d^2*e*x + 36*d*e^2*x^2 + 84*e^3*x^3)) + 4*a^3 \\ & *b^3*e^3*(5*A*e*(d^3 + 9*d^2*e*x + 36*d*e^2*x^2 + 84*e^3*x^3) + 4*B*(d^4 + \\ & 9*d^3*e*x + 36*d^2*e^2*x^2 + 84*d*e^3*x^3 + 126*e^4*x^4)) + 3*a^2*b^4*e^2 \\ & *(4*A*e*(d^4 + 9*d^3*e*x + 36*d^2*e^2*x^2 + 84*d*e^3*x^3 + 126*e^4*x^4) + \\ & 5*B*(d^5 + 9*d^4*e*x + 36*d^3*e^2*x^2 + 84*d^2*e^3*x^3 + 126*d*e^4*x^4 + 1 \\ & 26*e^5*x^5)) + 6*a*b^5*e*(A*e*(d^5 + 9*d^4*e*x + 36*d^3*e^2*x^2 + 84*d^2*e \\ & ^3*x^3 + 126*d*e^4*x^4 + 126*e^5*x^5) + 2*B*(d^6 + 9*d^5*e*x + 36*d^4*e^2*x \\ & ^2 + 84*d^3*e^3*x^3 + 126*d^2*e^4*x^4 + 126*d*e^5*x^5 + 84*e^6*x^6)) + b^6 \\ & *(2*A*e*(d^6 + 9*d^5*e*x + 36*d^4*e^2*x^2 + 84*d^3*e^3*x^3 + 126*d^2*e^4*x \\ & ^4 + 126*d*e^5*x^5 + 84*e^6*x^6) + 7*B*(d^7 + 9*d^6*e*x + 36*d^5*e^2*x^2 \\ & + 84*d^4*e^3*x^3 + 126*d^3*e^4*x^4 + 126*d^2*e^5*x^5 + 84*d*e^6*x^6 + 36*e \\ & ^7*x^7)))/(e^8*(d + e*x)^9) \end{aligned}$$

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.97, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {87, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{10}} dx \\ & \quad \downarrow 87 \\ & \frac{(-9aBe + 2Abe + 7bBd) \int \frac{(a+bx)^6}{(d+ex)^9} dx}{9e(bd - ae)} - \frac{(a + bx)^7(Bd - Ae)}{9e(d + ex)^9(bd - ae)} \\ & \quad \downarrow 55 \\ & \frac{(-9aBe + 2Abe + 7bBd) \left(\frac{b \int \frac{(a+bx)^6}{(d+ex)^8} dx}{8(bd-ae)} + \frac{(a+bx)^7}{8(d+ex)^8(bd-ae)} \right)}{9e(bd - ae)} - \frac{(a + bx)^7(Bd - Ae)}{9e(d + ex)^9(bd - ae)} \end{aligned}$$

$$\begin{array}{c} \downarrow 48 \\ \frac{\left(\frac{b(a+bx)^7}{56(d+ex)^7(bd-ae)^2} + \frac{(a+bx)^7}{8(d+ex)^8(bd-ae)}\right)(-9aBe + 2Abe + 7bBd)}{9e(bd-ae)} - \frac{(a+bx)^7(Bd-Ae)}{9e(d+ex)^9(bd-ae)} \end{array}$$

input `Int[((a + b*x)^6*(A + B*x))/(d + e*x)^10,x]`

output `-1/9*((B*d - A*e)*(a + b*x)^7)/(e*(b*d - a*e)*(d + e*x)^9) + ((7*b*B*d + 2*A*b*e - 9*a*B*e)*((a + b*x)^7/(8*(b*d - a*e)*(d + e*x)^8) + (b*(a + b*x)^7)/(56*(b*d - a*e)^2*(d + e*x)^7)))/(9*e*(b*d - a*e))`

Definitions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 55 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(Simplify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 788 vs. $2(129) = 258$.

Time = 0.23 (sec) , antiderivative size = 789, normalized size of antiderivative = 5.84

method	result
risch	$\frac{-\frac{b^6 B x^7}{2e} - \frac{b^5(2Abe+12Bae+7Bbd)x^6}{6e^2} - \frac{b^4(6Aab e^2+2A b^2 de+15B a^2 e^2+12Babde+7b^2 B d^2)x^5}{4e^3} - \frac{b^3(12A a^2 b e^3+6Aa b^2 d e^2+2A b^3 d^2 e)}{6e^4}}{6e^8(ex+d)^6}$
default	$\frac{5b^2(4A a^3 b e^4-12A a^2 b^2 d e^3+12Aa b^3 d^2 e^2-4A b^4 d^3 e+3B a^4 e^4-16B a^3 b d e^3+30B a^2 b^2 d^2 e^2-24Ba b^3 d^3 e+7B b^4 d^4)}{6e^8(ex+d)^6}$
norman	$\frac{-\frac{b^6 B x^7}{2e} - \frac{(2A b^6 e^2+12Ba b^5 e^2+7b^6 Bde)x^6}{6e^3} - \frac{(6Aa b^5 e^3+2A b^6 d e^2+15B a^2 b^4 e^3+12Ba b^5 d e^2+7b^6 B d^2 e)x^5}{4e^4} - \frac{(12A a^2 b^4 e^4+6Aa b^5 d e^3+3B a^3 b^3 d^3 e^2+2A b^4 d^4 e^2+7B a^2 b^2 d^2 e^2+12B a b^3 d^3 e+7B b^4 d^4)}{6e^8}}{6e^8(ex+d)^6}$
gosper	$\frac{252B x^7 b^6 e^7+168A x^6 b^6 e^7+1008B x^6 a b^5 e^7+588B x^6 b^6 d e^6+756A x^5 a b^5 e^7+252A x^5 b^6 d e^6+1890B x^5 a^2 b^4 e^7+1512B x^5 a b^5 d e^6+1008A x^4 a^2 b^3 d^3 e^2+252A x^4 b^4 d^4 e^2+756A x^4 a b^3 d^3 e+1512B x^4 a^2 b^2 d^2 e^2+12A x^4 a^3 b e^4+12A x^4 a^2 b^2 d e^3+12A x^4 a b^3 d^2 e^2+4A x^4 b^4 d^3 e+3B x^4 a^4 e^4-16B x^4 a^3 b d e^3+30B x^4 a^2 b^2 d^2 e^2-24B x^4 a b^3 d^3 e+7B x^4 b^4 d^4}{6e^8}$
orering	$\frac{252B x^7 b^6 e^7+168A x^6 b^6 e^7+1008B x^6 a b^5 e^7+588B x^6 b^6 d e^6+756A x^5 a b^5 e^7+252A x^5 b^6 d e^6+1890B x^5 a^2 b^4 e^7+1512B x^5 a b^5 d e^6+1008A x^4 a^2 b^3 d^3 e^2+252A x^4 b^4 d^4 e^2+756A x^4 a b^3 d^3 e+1512B x^4 a^2 b^2 d^2 e^2+12A x^4 a^3 b e^4+12A x^4 a^2 b^2 d e^3+12A x^4 a b^3 d^2 e^2+4A x^4 b^4 d^3 e+3B x^4 a^4 e^4-16B x^4 a^3 b d e^3+30B x^4 a^2 b^2 d^2 e^2-24B x^4 a b^3 d^3 e+7B x^4 b^4 d^4}{6e^8}$
parallelrisch	$\frac{252B b^6 x^7 e^8+168A b^6 e^8 x^6+1008Ba b^5 e^8 x^6+588B b^6 d e^7 x^6+756Aa b^5 e^8 x^5+252A b^6 d e^7 x^5+1890B a^2 b^4 e^8 x^5+1512Ba b^5 d e^6 x^5+1008A a^2 b^3 d^3 e^2 x^5+252A b^4 d^4 e^2 x^5+756A a b^3 d^3 e x^5+1512B x^5 a^2 b^2 d^2 e^2+12B x^5 a b^3 d^3 e+7B x^5 b^4 d^4}{6e^8}$

input

```
int((b*x+a)^6*(B*x+A)/(e*x+d)^10,x,method=_RETURNVERBOSE)
```

output

```
(-1/2*b^6*B/e*x^7-1/6*b^5/e^2*(2*A*b*e+12*B*a*e+7*B*b*d)*x^6-1/4*b^4/e^3*(6*A*a*b*e^2+2*A*b^2*d*e+15*B*a^2*e^2+12*B*a*b*d*e+7*B*b^2*d^2)*x^5-1/4*b^3/e^4*(12*A*a^2*b*e^3+6*A*a*b^2*d*e^2+2*A*b^3*d^2*e+16*B*a^3*e^3+15*B*a^2*b*d*e^2+12*B*a*b^2*d^2*e+7*B*b^3*d^3)*x^4-1/6*b^2/e^5*(20*A*a^3*b*e^4+12*A*a^2*b^2*d*e^3+6*A*a*b^3*d^2*e^2+2*A*b^4*d^3*e+15*B*a^4*e^4+16*B*a^3*b*d*e^3+15*B*a^2*b^2*d^2*e^2+12*B*a*b^3*d^3*e+7*B*b^4*d^4)*x^3-1/14*b/e^6*(30*A*a^4*b*e^5+20*A*a^3*b^2*d*e^4+12*A*a^2*b^3*d^2*e^3+6*A*a*b^4*d^3*e^2+2*A*b^5*d^4*e+12*B*a^5*e^5+15*B*a^4*b*d*e^4+16*B*a^3*b^2*d^2*e^3+15*B*a^2*b^3*d^3*e^2+12*B*a*b^4*d^4*e+7*B*b^5*d^5)*x^2-1/56/e^7*(42*A*a^5*b*e^6+30*A*a^4*b^2*d*e^5+20*A*a^3*b^3*d^2*e^4+12*A*a^2*b^4*d^3*e^3+6*A*a*b^5*d^4*e^2+2*A*b^6*d^5*e+7*B*a^6*e^6+12*B*a^5*b*d*e^5+15*B*a^4*b^2*d^2*e^4+16*B*a^3*b^3*d^3*e^3+15*B*a^2*b^4*d^4*e^2+12*B*a*b^5*d^5*e+7*B*b^6*d^6)*x-1/504/e^8*(56*A*a^6*e^7+42*A*a^5*b*d*e^6+30*A*a^4*b^2*d^2*e^5+20*A*a^3*b^3*d^3*e^4+12*A*a^2*b^4*d^4*e^3+6*A*a*b^5*d^5*e^2+2*A*b^6*d^6*e+7*B*a^6*d^6*e+12*B*a^5*b*d^2*e^5+15*B*a^4*b^2*d^3*e^4+16*B*a^3*b^3*d^4*e^3+15*B*a^2*b^4*d^5*e^2+12*B*a*b^5*d^6*e+7*B*b^6*d^7))/(e*x+d)^9
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 861 vs. $2(129) = 258$.

Time = 0.09 (sec) , antiderivative size = 861, normalized size of antiderivative = 6.38

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{10}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^10,x, algorithm="fricas")`

output

```
-1/504*(252*B*b^6*e^7*x^7 + 7*B*b^6*d^7 + 56*A*a^6*e^7 + 2*(6*B*a*b^5 + A
b^6)*d^6*e + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 + 4*(4*B*a^3*b^3 + 3*A*a^
2*b^4)*d^4*e^3 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 + 6*(2*B*a^5*b + 5*
A*a^4*b^2)*d^2*e^5 + 7*(B*a^6 + 6*A*a^5*b)*d*e^6 + 84*(7*B*b^6*d*e^6 + 2*(
6*B*a*b^5 + A*b^6)*e^7)*x^6 + 126*(7*B*b^6*d^2*e^5 + 2*(6*B*a*b^5 + A*b^6)
*d*e^6 + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^5 + 126*(7*B*b^6*d^3*e^4 + 2*(
6*B*a*b^5 + A*b^6)*d^2*e^5 + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6 + 4*(4*B*a^
3*b^3 + 3*A*a^2*b^4)*e^7)*x^4 + 84*(7*B*b^6*d^4*e^3 + 2*(6*B*a*b^5 + A*b^6)
*d^3*e^4 + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 + 4*(4*B*a^3*b^3 + 3*A*a^2
*b^4)*d*e^6 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^3 + 36*(7*B*b^6*d^5*e^2
+ 2*(6*B*a*b^5 + A*b^6)*d^4*e^3 + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*e^4 + 4
*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^6
+ 6*(2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 9*(7*B*b^6*d^6*e + 2*(6*B*a*b^5
+ A*b^6)*d^5*e^2 + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^3 + 4*(4*B*a^3*b^3 +
3*A*a^2*b^4)*d^3*e^4 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^5 + 6*(2*B*a^5*
b + 5*A*a^4*b^2)*d*e^6 + 7*(B*a^6 + 6*A*a^5*b)*e^7)*x)/(e^17*x^9 + 9*d*e^1
6*x^8 + 36*d^2*e^15*x^7 + 84*d^3*e^14*x^6 + 126*d^4*e^13*x^5 + 126*d^5*e^1
2*x^4 + 84*d^6*e^11*x^3 + 36*d^7*e^10*x^2 + 9*d^8*e^9*x + d^9*e^8)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{10}} dx = \text{Timed out}$$

input `integrate((b*x+a)**6*(B*x+A)/(e*x+d)**10,x)`

output Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 861 vs. $2(129) = 258$.

Time = 0.07 (sec) , antiderivative size = 861, normalized size of antiderivative = 6.38

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{10}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^10,x, algorithm="maxima")`

output

```
-1/504*(252*B*b^6*e^7*x^7 + 7*B*b^6*d^7 + 56*A*a^6*e^7 + 2*(6*B*a*b^5 + A*b^6)*d^6*e + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 + 4*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4*e^3 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 + 6*(2*B*a^5*b + 5*A*a^4*b^2)*d^2*e^5 + 7*(B*a^6 + 6*A*a^5*b)*d*e^6 + 84*(7*B*b^6*d*e^6 + 2*(6*B*a*b^5 + A*b^6)*e^7)*x^6 + 126*(7*B*b^6*d^2*e^5 + 2*(6*B*a*b^5 + A*b^6)*d*e^6 + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^5 + 126*(7*B*b^6*d^3*e^4 + 2*(6*B*a*b^5 + A*b^6)*d^2*e^5 + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6 + 4*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^7)*x^4 + 84*(7*B*b^6*d^4*e^3 + 2*(6*B*a*b^5 + A*b^6)*d^3*e^4 + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 + 4*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^6 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^3 + 36*(7*B*b^6*d^5*e^2 + 2*(6*B*a*b^5 + A*b^6)*d^4*e^3 + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*e^4 + 4*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^6 + 6*(2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 9*(7*B*b^6*d^6*e + 2*(6*B*a*b^5 + A*b^6)*d^5*e^2 + 3*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^3 + 4*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 + 5*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^5 + 6*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^6 + 7*(B*a^6 + 6*A*a^5*b)*e^7)*x)/(e^17*x^9 + 9*d*e^16*x^8 + 36*d^2*e^15*x^7 + 84*d^3*e^14*x^6 + 126*d^4*e^13*x^5 + 126*d^5*e^12*x^4 + 84*d^6*e^11*x^3 + 36*d^7*e^10*x^2 + 9*d^8*e^9*x + d^9*e^8)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 912 vs. $2(129) = 258$.

Time = 0.13 (sec) , antiderivative size = 912, normalized size of antiderivative = 6.76

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{10}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^10,x, algorithm="giac")`

output

```
-1/504*(252*B*b^6*e^7*x^7 + 588*B*b^6*d*e^6*x^6 + 1008*B*a*b^5*e^7*x^6 + 1
68*A*b^6*e^7*x^6 + 882*B*b^6*d^2*e^5*x^5 + 1512*B*a*b^5*d*e^6*x^5 + 252*A*
b^6*d*e^6*x^5 + 1890*B*a^2*b^4*e^7*x^5 + 756*A*a*b^5*e^7*x^5 + 882*B*b^6*d
^3*e^4*x^4 + 1512*B*a*b^5*d^2*e^5*x^4 + 252*A*b^6*d^2*e^5*x^4 + 1890*B*a^2
*b^4*d*e^6*x^4 + 756*A*a*b^5*d*e^6*x^4 + 2016*B*a^3*b^3*e^7*x^4 + 1512*A*a
^2*b^4*e^7*x^4 + 588*B*b^6*d^4*e^3*x^3 + 1008*B*a*b^5*d^3*e^4*x^3 + 168*A*
b^6*d^3*e^4*x^3 + 1260*B*a^2*b^4*d^2*e^5*x^3 + 504*A*a*b^5*d^2*e^5*x^3 + 1
344*B*a^3*b^3*d*e^6*x^3 + 1008*A*a^2*b^4*d*e^6*x^3 + 1260*B*a^4*b^2*e^7*x^
3 + 1680*A*a^3*b^3*e^7*x^3 + 252*B*b^6*d^5*e^2*x^2 + 432*B*a*b^5*d^4*e^3*x
^2 + 72*A*b^6*d^4*e^3*x^2 + 540*B*a^2*b^4*d^3*e^4*x^2 + 216*A*a*b^5*d^3*e^
4*x^2 + 576*B*a^3*b^3*d^2*e^5*x^2 + 432*A*a^2*b^4*d^2*e^5*x^2 + 540*B*a^4*
b^2*d*e^6*x^2 + 720*A*a^3*b^3*d*e^6*x^2 + 432*B*a^5*b*e^7*x^2 + 1080*A*a^4
*b^2*e^7*x^2 + 63*B*b^6*d^6*e*x + 108*B*a*b^5*d^5*e^2*x + 18*A*b^6*d^5*e^2
*x + 135*B*a^2*b^4*d^4*e^3*x + 54*A*a*b^5*d^4*e^3*x + 144*B*a^3*b^3*d^3*e^
4*x + 108*A*a^2*b^4*d^3*e^4*x + 135*B*a^4*b^2*d^2*e^5*x + 180*A*a^3*b^3*d^
2*e^5*x + 108*B*a^5*b*d*e^6*x + 270*A*a^4*b^2*d*e^6*x + 63*B*a^6*e^7*x + 3
78*A*a^5*b*e^7*x + 7*B*b^6*d^7 + 12*B*a*b^5*d^6*e + 2*A*b^6*d^6*e + 15*B*a
^2*b^4*d^5*e^2 + 6*A*a*b^5*d^5*e^2 + 16*B*a^3*b^3*d^4*e^3 + 12*A*a^2*b^4*d
^4*e^3 + 15*B*a^4*b^2*d^3*e^4 + 20*A*a^3*b^3*d^3*e^4 + 12*B*a^5*b*d^2*e^5
+ 30*A*a^4*b^2*d^2*e^5 + 7*B*a^6*d*e^6 + 42*A*a^5*b*d*e^6 + 56*A*a^6*e^...
```

Mupad [B] (verification not implemented)

Time = 1.04 (sec) , antiderivative size = 877, normalized size of antiderivative = 6.50

$$\int \frac{(a + bx)^6 (A + Bx)}{(d + ex)^{10}} dx =$$

$$\frac{7Ba^6de^6 + 56Aa^6e^7 + 12Ba^5bd^2e^5 + 42Aa^5bde^6 + 15Ba^4b^2d^3e^4 + 30Aa^4b^2d^2e^5 + 16Ba^3b^3d^4e^3 + 20Aa^3b^3d^3e^4 + 15Ba^2b^4d^5e^2}{504e^8}$$

input

```
int(((A + B*x)*(a + b*x)^6)/(d + e*x)^10,x)
```

output

```
-((56*A*a^6*e^7 + 7*B*b^6*d^7 + 2*A*b^6*d^6*e + 7*B*a^6*d*e^6 + 6*A*a*b^5*d^5*e^2 + 12*B*a^5*b*d^2*e^5 + 12*A*a^2*b^4*d^4*e^3 + 20*A*a^3*b^3*d^3*e^4 + 30*A*a^4*b^2*d^2*e^5 + 15*B*a^2*b^4*d^5*e^2 + 16*B*a^3*b^3*d^4*e^3 + 15*B*a^4*b^2*d^3*e^4 + 42*A*a^5*b*d*e^6 + 12*B*a*b^5*d^6*e)/(504*e^8) + (x*(7*B*a^6*e^6 + 7*B*b^6*d^6 + 42*A*a^5*b*e^6 + 2*A*b^6*d^5*e + 6*A*a*b^5*d^4*e^2 + 30*A*a^4*b^2*d*e^5 + 12*A*a^2*b^4*d^3*e^3 + 20*A*a^3*b^3*d^2*e^4 + 15*B*a^2*b^4*d^4*e^2 + 16*B*a^3*b^3*d^3*e^3 + 15*B*a^4*b^2*d^2*e^4 + 12*B*a*b^5*d^5*e + 12*B*a^5*b*d*e^5))/(56*e^7) + (b^3*x^4*(16*B*a^3*e^3 + 7*B*b^3*d^3 + 12*A*a^2*b*e^3 + 2*A*b^3*d^2*e + 6*A*a*b^2*d*e^2 + 12*B*a*b^2*d^2*e + 15*B*a^2*b*d*e^2))/(4*e^4) + (b^5*x^6*(2*A*b*e + 12*B*a*e + 7*B*b*d))/(6*e^2) + (b*x^2*(12*B*a^5*e^5 + 7*B*b^5*d^5 + 30*A*a^4*b*e^5 + 2*A*b^5*d^4*e + 6*A*a*b^4*d^3*e^2 + 20*A*a^3*b^2*d*e^4 + 12*A*a^2*b^3*d^2*e^3 + 15*B*a^2*b^3*d^3*e^2 + 16*B*a^3*b^2*d^2*e^3 + 12*B*a*b^4*d^4*e + 15*B*a^4*b*d*e^4))/(14*e^6) + (b^2*x^3*(15*B*a^4*e^4 + 7*B*b^4*d^4 + 20*A*a^3*b*e^4 + 2*A*b^4*d^3*e + 6*A*a*b^3*d^2*e^2 + 12*A*a^2*b^2*d*e^3 + 15*B*a^2*b^2*d^2*e^2 + 12*B*a*b^3*d^3*e + 16*B*a^3*b*d*e^3))/(6*e^5) + (b^4*x^5*(15*B*a^2*e^2 + 7*B*b^2*d^2 + 6*A*a*b*e^2 + 2*A*b^2*d*e + 12*B*a*b*d*e))/(4*e^3) + (B*b^6*x^7)/(2*e))/(d^9 + e^9*x^9 + 9*d*e^8*x^8 + 36*d^7*e^2*x^2 + 84*d^6*e^3*x^3 + 126*d^5*e^4*x^4 + 126*d^4*e^5*x^5 + 84*d^3*e^6*x^6 + 36*d^2*e^7*x^7 + 9*d^8*e*x)
```


Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 585, normalized size of antiderivative = 4.33

$$\int \frac{(a + bx)^6 (A + Bx)}{(d + ex)^{10}} dx$$

$$= \frac{-36b^7 e^7 x^7 - 168a b^6 e^7 x^6 - 84b^7 d e^6 x^6 - 378a^2 b^5 e^7 x^5 - 252a b^6 d e^6 x^5 - 126b^7 d^2 e^5 x^5 - 504a^3 b^4 e^7 x^4 - 3$$

input `int((b*x+a)^6*(B*x+A)/(e*x+d)^10,x)`

output

```
( - 8*a**7*e**7 - 7*a**6*b*d*e**6 - 63*a**6*b*e**7*x - 6*a**5*b**2*d**2*e**5 - 54*a**5*b**2*d*e**6*x - 216*a**5*b**2*e**7*x**2 - 5*a**4*b**3*d**3*e**4 - 45*a**4*b**3*d**2*e**5*x - 180*a**4*b**3*d*e**6*x**2 - 420*a**4*b**3*e**7*x**3 - 4*a**3*b**4*d**4*e**3 - 36*a**3*b**4*d**3*e**4*x - 144*a**3*b**4*d**2*e**5*x**2 - 336*a**3*b**4*d*e**6*x**3 - 504*a**3*b**4*e**7*x**4 - 3*a**2*b**5*d**5*e**2 - 27*a**2*b**5*d**4*e**3*x - 108*a**2*b**5*d**3*e**4*x**2 - 252*a**2*b**5*d**2*e**5*x**3 - 378*a**2*b**5*d*e**6*x**4 - 378*a**2*b**5*e**7*x**5 - 2*a*b**6*d**6*e - 18*a*b**6*d**5*e**2*x - 72*a*b**6*d**4*e**3*x**2 - 168*a*b**6*d**3*e**4*x**3 - 252*a*b**6*d**2*e**5*x**4 - 252*a*b**6*d*e**6*x**5 - 168*a*b**6*e**7*x**6 - b**7*d**7 - 9*b**7*d**6*e*x - 36*b**7*d**5*e**2*x**2 - 84*b**7*d**4*e**3*x**3 - 126*b**7*d**3*e**4*x**4 - 126*b**7*d**2*e**5*x**5 - 84*b**7*d*e**6*x**6 - 36*b**7*e**7*x**7)/(72*e**8*(d**9 + 9*d**8*e*x + 36*d**7*e**2*x**2 + 84*d**6*e**3*x**3 + 126*d**5*e**4*x**4 + 126*d**4*e**5*x**5 + 84*d**3*e**6*x**6 + 36*d**2*e**7*x**7 + 9*d*e**8*x**8 + e**9*x**9))
```

3.60 $\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{11}} dx$

Optimal result	605
Mathematica [B] (verified)	606
Rubi [A] (verified)	606
Maple [B] (verified)	608
Fricas [B] (verification not implemented)	610
Sympy [F(-1)]	610
Maxima [B] (verification not implemented)	611
Giac [B] (verification not implemented)	612
Mupad [B] (verification not implemented)	613
Reduce [B] (verification not implemented)	614

Optimal result

Integrand size = 20, antiderivative size = 185

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{11}} dx = -\frac{(Bd - Ae)(a+bx)^7}{10e(bd - ae)(d+ex)^{10}} + \frac{(7bBd + 3Abe - 10aBe)(a+bx)^7}{90e(bd - ae)^2(d+ex)^9} + \frac{b(7bBd + 3Abe - 10aBe)(a+bx)^7}{360e(bd - ae)^3(d+ex)^8} + \frac{b^2(7bBd + 3Abe - 10aBe)(a+bx)^7}{2520e(bd - ae)^4(d+ex)^7}$$

output

```
-1/10*(-A*e+B*d)*(b*x+a)^7/e/(-a*e+b*d)/(e*x+d)^10+1/90*(3*A*b*e-10*B*a*e+
7*B*b*d)*(b*x+a)^7/e/(-a*e+b*d)^2/(e*x+d)^9+1/360*b*(3*A*b*e-10*B*a*e+7*B*
b*d)*(b*x+a)^7/e/(-a*e+b*d)^3/(e*x+d)^8+1/2520*b^2*(3*A*b*e-10*B*a*e+7*B*b
*d)*(b*x+a)^7/e/(-a*e+b*d)^4/(e*x+d)^7
```


$$\begin{aligned}
& \int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{11}} dx \\
& \quad \downarrow 87 \\
& \frac{(-10aBe + 3Abe + 7bBd) \int \frac{(a+bx)^6}{(d+ex)^{10}} dx}{10e(bd-ae)} - \frac{(a+bx)^7(Bd-Ae)}{10e(d+ex)^{10}(bd-ae)} \\
& \quad \downarrow 55 \\
& \frac{(-10aBe + 3Abe + 7bBd) \left(\frac{2b \int \frac{(a+bx)^6}{(d+ex)^9} dx}{9(bd-ae)} + \frac{(a+bx)^7}{9(d+ex)^9(bd-ae)} \right)}{10e(bd-ae)} - \frac{(a+bx)^7(Bd-Ae)}{10e(d+ex)^{10}(bd-ae)} \\
& \quad \downarrow 55 \\
& \frac{(-10aBe + 3Abe + 7bBd) \left(\frac{2b \left(\frac{b \int \frac{(a+bx)^6}{(d+ex)^8} dx}{8(bd-ae)} + \frac{(a+bx)^7}{8(d+ex)^8(bd-ae)} \right)}{9(bd-ae)} + \frac{(a+bx)^7}{9(d+ex)^9(bd-ae)} \right)}{10e(bd-ae)} - \frac{(a+bx)^7(Bd-Ae)}{10e(d+ex)^{10}(bd-ae)} \\
& \quad \downarrow 48 \\
& \frac{\left(\frac{(a+bx)^7}{9(d+ex)^9(bd-ae)} + \frac{2b \left(\frac{b(a+bx)^7}{56(d+ex)^7(bd-ae)^2} + \frac{(a+bx)^7}{8(d+ex)^8(bd-ae)} \right)}{9(bd-ae)} \right) (-10aBe + 3Abe + 7bBd)}{10e(bd-ae)} - \frac{(a+bx)^7(Bd-Ae)}{10e(d+ex)^{10}(bd-ae)}
\end{aligned}$$

input `Int[((a + b*x)^6*(A + B*x))/(d + e*x)^11,x]`

output `-1/10*((B*d - A*e)*(a + b*x)^7)/(e*(b*d - a*e)*(d + e*x)^10) + ((7*b*B*d + 3*A*b*e - 10*a*B*e)*((a + b*x)^7/(9*(b*d - a*e)*(d + e*x)^9) + (2*b*((a + b*x)^7/(8*(b*d - a*e)*(d + e*x)^8) + (b*(a + b*x)^7)/(56*(b*d - a*e)^2*(d + e*x)^7)))/(9*(b*d - a*e)))/(10*e*(b*d - a*e))`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
 [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
 a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 55 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
 (a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
 implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
 c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
 c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
 lerQ[n, 1])`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
 + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
 + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
 /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
 rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 788 vs. $2(177) = 354$.

Time = 0.24 (sec) , antiderivative size = 789, normalized size of antiderivative = 4.26

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 872 vs. $2(177) = 354$.

Time = 0.11 (sec) , antiderivative size = 872, normalized size of antiderivative = 4.71

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{11}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^11,x, algorithm="fricas")`

output

```
-1/2520*(840*B*b^6*e^7*x^7 + 7*B*b^6*d^7 + 252*A*a^6*e^7 + 3*(6*B*a*b^5 +
A*b^6)*d^6*e + 6*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 + 10*(4*B*a^3*b^3 + 3*A
*a^2*b^4)*d^4*e^3 + 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 + 21*(2*B*a^5*b
+ 5*A*a^4*b^2)*d^2*e^5 + 28*(B*a^6 + 6*A*a^5*b)*d*e^6 + 210*(7*B*b^6*d*e^
6 + 3*(6*B*a*b^5 + A*b^6)*e^7)*x^6 + 252*(7*B*b^6*d^2*e^5 + 3*(6*B*a*b^5 +
A*b^6)*d*e^6 + 6*(5*B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^5 + 210*(7*B*b^6*d^3*e^
4 + 3*(6*B*a*b^5 + A*b^6)*d^2*e^5 + 6*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6 + 10
*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^7)*x^4 + 120*(7*B*b^6*d^4*e^3 + 3*(6*B*a*b^
5 + A*b^6)*d^3*e^4 + 6*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 + 10*(4*B*a^3*b^3
+ 3*A*a^2*b^4)*d*e^6 + 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^3 + 45*(7*B*
b^6*d^5*e^2 + 3*(6*B*a*b^5 + A*b^6)*d^4*e^3 + 6*(5*B*a^2*b^4 + 2*A*a*b^5)*
d^3*e^4 + 10*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + 15*(3*B*a^4*b^2 + 4*A*a
^3*b^3)*d*e^6 + 21*(2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 10*(7*B*b^6*d^6*e
+ 3*(6*B*a*b^5 + A*b^6)*d^5*e^2 + 6*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^3 + 10
*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 + 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*
e^5 + 21*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^6 + 28*(B*a^6 + 6*A*a^5*b)*e^7)*x)/
(e^18*x^10 + 10*d*e^17*x^9 + 45*d^2*e^16*x^8 + 120*d^3*e^15*x^7 + 210*d^4*
e^14*x^6 + 252*d^5*e^13*x^5 + 210*d^6*e^12*x^4 + 120*d^7*e^11*x^3 + 45*d^8
*e^10*x^2 + 10*d^9*e^9*x + d^10*e^8)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{11}} dx = \text{Timed out}$$

input `integrate((b*x+a)**6*(B*x+A)/(e*x+d)**11,x)`

output Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 872 vs. $2(177) = 354$.

Time = 0.08 (sec) , antiderivative size = 872, normalized size of antiderivative = 4.71

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{11}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^11,x, algorithm="maxima")`

output

$$\begin{aligned} & -1/2520*(840*B*b^6*e^7*x^7 + 7*B*b^6*d^7 + 252*A*a^6*e^7 + 3*(6*B*a*b^5 + \\ & A*b^6)*d^6*e + 6*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 + 10*(4*B*a^3*b^3 + 3*A \\ & *a^2*b^4)*d^4*e^3 + 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 + 21*(2*B*a^5*b \\ & + 5*A*a^4*b^2)*d^2*e^5 + 28*(B*a^6 + 6*A*a^5*b)*d*e^6 + 210*(7*B*b^6*d*e^ \\ & 6 + 3*(6*B*a*b^5 + A*b^6)*e^7)*x^6 + 252*(7*B*b^6*d^2*e^5 + 3*(6*B*a*b^5 + \\ & A*b^6)*d*e^6 + 6*(5*B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^5 + 210*(7*B*b^6*d^3*e^ \\ & 4 + 3*(6*B*a*b^5 + A*b^6)*d^2*e^5 + 6*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6 + 10 \\ & *(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^7)*x^4 + 120*(7*B*b^6*d^4*e^3 + 3*(6*B*a*b^ \\ & 5 + A*b^6)*d^3*e^4 + 6*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 + 10*(4*B*a^3*b^3 \\ & + 3*A*a^2*b^4)*d*e^6 + 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^3 + 45*(7*B* \\ & b^6*d^5*e^2 + 3*(6*B*a*b^5 + A*b^6)*d^4*e^3 + 6*(5*B*a^2*b^4 + 2*A*a*b^5)* \\ & d^3*e^4 + 10*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + 15*(3*B*a^4*b^2 + 4*A*a \\ & ^3*b^3)*d*e^6 + 21*(2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 10*(7*B*b^6*d^6*e \\ & + 3*(6*B*a*b^5 + A*b^6)*d^5*e^2 + 6*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^3 + 10 \\ & *(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 + 15*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2* \\ & e^5 + 21*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^6 + 28*(B*a^6 + 6*A*a^5*b)*e^7)*x)/ \\ & (e^18*x^10 + 10*d*e^17*x^9 + 45*d^2*e^16*x^8 + 120*d^3*e^15*x^7 + 210*d^4* \\ & e^14*x^6 + 252*d^5*e^13*x^5 + 210*d^6*e^12*x^4 + 120*d^7*e^11*x^3 + 45*d^8 \\ & *e^10*x^2 + 10*d^9*e^9*x + d^10*e^8) \end{aligned}$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 912 vs. $2(177) = 354$.

Time = 0.13 (sec) , antiderivative size = 912, normalized size of antiderivative = 4.93

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{11}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^11,x, algorithm="giac")`

output

```
-1/2520*(840*B*b^6*e^7*x^7 + 1470*B*b^6*d*e^6*x^6 + 3780*B*a*b^5*e^7*x^6 +
630*A*b^6*e^7*x^6 + 1764*B*b^6*d^2*e^5*x^5 + 4536*B*a*b^5*d*e^6*x^5 + 756
*A*b^6*d*e^6*x^5 + 7560*B*a^2*b^4*e^7*x^5 + 3024*A*a*b^5*e^7*x^5 + 1470*B*
b^6*d^3*e^4*x^4 + 3780*B*a*b^5*d^2*e^5*x^4 + 630*A*b^6*d^2*e^5*x^4 + 6300*
B*a^2*b^4*d*e^6*x^4 + 2520*A*a*b^5*d*e^6*x^4 + 8400*B*a^3*b^3*e^7*x^4 + 63
00*A*a^2*b^4*e^7*x^4 + 840*B*b^6*d^4*e^3*x^3 + 2160*B*a*b^5*d^3*e^4*x^3 +
360*A*b^6*d^3*e^4*x^3 + 3600*B*a^2*b^4*d^2*e^5*x^3 + 1440*A*a*b^5*d^2*e^5*
x^3 + 4800*B*a^3*b^3*d*e^6*x^3 + 3600*A*a^2*b^4*d*e^6*x^3 + 5400*B*a^4*b^2
*e^7*x^3 + 7200*A*a^3*b^3*e^7*x^3 + 315*B*b^6*d^5*e^2*x^2 + 810*B*a*b^5*d^
4*e^3*x^2 + 135*A*b^6*d^4*e^3*x^2 + 1350*B*a^2*b^4*d^3*e^4*x^2 + 540*A*a*b
^5*d^3*e^4*x^2 + 1800*B*a^3*b^3*d^2*e^5*x^2 + 1350*A*a^2*b^4*d^2*e^5*x^2 +
2025*B*a^4*b^2*d*e^6*x^2 + 2700*A*a^3*b^3*d*e^6*x^2 + 1890*B*a^5*b*e^7*x^
2 + 4725*A*a^4*b^2*e^7*x^2 + 70*B*b^6*d^6*e*x + 180*B*a*b^5*d^5*e^2*x + 30
*A*b^6*d^5*e^2*x + 300*B*a^2*b^4*d^4*e^3*x + 120*A*a*b^5*d^4*e^3*x + 400*B
*a^3*b^3*d^3*e^4*x + 300*A*a^2*b^4*d^3*e^4*x + 450*B*a^4*b^2*d^2*e^5*x + 6
00*A*a^3*b^3*d^2*e^5*x + 420*B*a^5*b*d*e^6*x + 1050*A*a^4*b^2*d*e^6*x + 28
0*B*a^6*e^7*x + 1680*A*a^5*b*e^7*x + 7*B*b^6*d^7 + 18*B*a*b^5*d^6*e + 3*A*
b^6*d^6*e + 30*B*a^2*b^4*d^5*e^2 + 12*A*a*b^5*d^5*e^2 + 40*B*a^3*b^3*d^4*e
^3 + 30*A*a^2*b^4*d^4*e^3 + 45*B*a^4*b^2*d^3*e^4 + 60*A*a^3*b^3*d^3*e^4 +
42*B*a^5*b*d^2*e^5 + 105*A*a^4*b^2*d^2*e^5 + 28*B*a^6*d*e^6 + 168*A*a^5...
```

Mupad [B] (verification not implemented)

Time = 1.07 (sec) , antiderivative size = 888, normalized size of antiderivative = 4.80

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{11}} dx =$$

$$\frac{28 B a^6 d e^6 + 252 A a^6 e^7 + 42 B a^5 b d^2 e^5 + 168 A a^5 b d e^6 + 45 B a^4 b^2 d^3 e^4 + 105 A a^4 b^2 d^2 e^5 + 40 B a^3 b^3 d^4 e^3 + 60 A a^3 b^3 d^3 e^4 + 30 B a^2 b^4 d^5 e^2 + 20 A a^2 b^4 d^4 e^3 + 15 B a b^5 d^6 e + 5 A a b^5 d^5 e^2}{2520 e^8}$$

input

```
int(((A + B*x)*(a + b*x)^6)/(d + e*x)^11,x)
```

output

```
-((252*A*a^6*e^7 + 7*B*b^6*d^7 + 3*A*b^6*d^6*e + 28*B*a^6*d*e^6 + 12*A*a*b^5*d^5*e^2 + 42*B*a^5*b*d^2*e^5 + 30*A*a^2*b^4*d^4*e^3 + 60*A*a^3*b^3*d^3*e^4 + 105*A*a^4*b^2*d^2*e^5 + 30*B*a^2*b^4*d^5*e^2 + 40*B*a^3*b^3*d^4*e^3 + 45*B*a^4*b^2*d^3*e^4 + 168*A*a^5*b*d*e^6 + 18*B*a*b^5*d^6*e)/(2520*e^8) + (x*(28*B*a^6*e^6 + 7*B*b^6*d^6 + 168*A*a^5*b*d*e^6 + 3*A*b^6*d^5*e + 12*A*a*b^5*d^4*e^2 + 105*A*a^4*b^2*d*e^5 + 30*A*a^2*b^4*d^3*e^3 + 60*A*a^3*b^3*d^2*e^4 + 30*B*a^2*b^4*d^4*e^2 + 40*B*a^3*b^3*d^3*e^3 + 45*B*a^4*b^2*d^2*e^4 + 18*B*a*b^5*d^5*e + 42*B*a^5*b*d*e^5)/(2520*e^7) + (b^3*x^4*(40*B*a^3*e^3 + 7*B*b^3*d^3 + 30*A*a^2*b*d*e^3 + 3*A*b^3*d^2*e + 12*A*a*b^2*d*e^2 + 18*B*a*b^2*d^2*e + 30*B*a^2*b*d*d*e^2))/(12*e^4) + (b^5*x^6*(3*A*b*d*e + 18*B*a*b*d*e + 7*B*b*d*d))/(12*e^2) + (b*x^2*(42*B*a^5*e^5 + 7*B*b^5*d^5 + 105*A*a^4*b*d*e^5 + 3*A*b^5*d^4*e + 12*A*a*b^4*d^3*e^2 + 60*A*a^3*b^2*d*e^4 + 30*A*a^2*b^3*d^2*e^3 + 30*B*a^2*b^3*d^3*e^2 + 40*B*a^3*b^2*d^2*e^3 + 18*B*a*b^4*d^4*e + 45*B*a^4*b*d*d*e^4))/(56*e^6) + (b^2*x^3*(45*B*a^4*e^4 + 7*B*b^4*d^4 + 60*A*a^3*b*d*e^4 + 3*A*b^4*d^3*e + 12*A*a*b^3*d^2*e^2 + 30*A*a^2*b^2*d*e^3 + 30*B*a^2*b^2*d^2*e^2 + 18*B*a*b^3*d^3*e + 40*B*a^3*b*d*d*e^3))/(21*e^5) + (b^4*x^5*(30*B*a^2*d^2*e^2 + 7*B*b^2*d^2 + 12*A*a*b*d*e^2 + 3*A*b^2*d*d*e + 18*B*a*b*d*d*e))/(10*e^3) + (B*b^6*x^7)/(3*e))/(d^10 + e^10*x^10 + 10*d*e^9*x^9 + 45*d^8*e^2*x^2 + 120*d^7*e^3*x^3 + 210*d^6*e^4*x^4 + 252*d^5*e^5*x^5 + 210*d^4*e^6*x^6 + 120*d^3*e^7*x^7 + 45*d^2*e^8*x^8 + 10*d^9*e*x)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 596, normalized size of antiderivative = 3.22

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{11}} dx$$

$$= \frac{-120b^7e^7x^7 - 630ab^6e^7x^6 - 210b^7de^6x^6 - 1512a^2b^5e^7x^5 - 756ab^6de^6x^5 - 252b^7d^2e^5x^5 - 2100a^3b^4e^7x^4}{(d+ex)^{10}}$$

input `int((b*x+a)^6*(B*x+A)/(e*x+d)^11,x)`

output

```
( - 36*a**7*e**7 - 28*a**6*b*d*e**6 - 280*a**6*b*e**7*x - 21*a**5*b**2*d**2*e**5 - 210*a**5*b**2*d*e**6*x - 945*a**5*b**2*e**7*x**2 - 15*a**4*b**3*d**3*e**4 - 150*a**4*b**3*d**2*e**5*x - 675*a**4*b**3*d*e**6*x**2 - 1800*a**4*b**3*e**7*x**3 - 10*a**3*b**4*d**4*e**3 - 100*a**3*b**4*d**3*e**4*x - 450*a**3*b**4*d**2*e**5*x**2 - 1200*a**3*b**4*d*e**6*x**3 - 2100*a**3*b**4*e**7*x**4 - 6*a**2*b**5*d**5*e**2 - 60*a**2*b**5*d**4*e**3*x - 270*a**2*b**5*d**3*e**4*x**2 - 720*a**2*b**5*d**2*e**5*x**3 - 1260*a**2*b**5*d*e**6*x**4 - 1512*a**2*b**5*e**7*x**5 - 3*a*b**6*d**6*e - 30*a*b**6*d**5*e**2*x - 135*a*b**6*d**4*e**3*x**2 - 360*a*b**6*d**3*e**4*x**3 - 630*a*b**6*d**2*e**5*x**4 - 756*a*b**6*d*e**6*x**5 - 630*a*b**6*e**7*x**6 - b**7*d**7 - 10*b**7*d**6*e*x - 45*b**7*d**5*e**2*x**2 - 120*b**7*d**4*e**3*x**3 - 210*b**7*d**3*e**4*x**4 - 252*b**7*d**2*e**5*x**5 - 210*b**7*d*e**6*x**6 - 120*b**7*e**7*x**7)/(360*e**8*(d**10 + 10*d**9*e*x + 45*d**8*e**2*x**2 + 120*d**7*e**3*x**3 + 210*d**6*e**4*x**4 + 252*d**5*e**5*x**5 + 210*d**4*e**6*x**6 + 120*d**3*e**7*x**7 + 45*d**2*e**8*x**8 + 10*d*e**9*x**9 + e**10*x**10))
```

3.61 $\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{12}} dx$

Optimal result	615
Mathematica [B] (verified)	616
Rubi [A] (verified)	616
Maple [B] (verified)	619
Fricas [B] (verification not implemented)	620
Sympy [F(-1)]	621
Maxima [B] (verification not implemented)	622
Giac [B] (verification not implemented)	622
Mupad [B] (verification not implemented)	623
Reduce [B] (verification not implemented)	624

Optimal result

Integrand size = 20, antiderivative size = 235

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{12}} dx = -\frac{(Bd - Ae)(a+bx)^7}{11e(bd - ae)(d+ex)^{11}} + \frac{(7bBd + 4Abe - 11aBe)(a+bx)^7}{110e(bd - ae)^2(d+ex)^{10}} + \frac{b(7bBd + 4Abe - 11aBe)(a+bx)^7}{330e(bd - ae)^3(d+ex)^9} + \frac{b^2(7bBd + 4Abe - 11aBe)(a+bx)^7}{1320e(bd - ae)^4(d+ex)^8} + \frac{b^3(7bBd + 4Abe - 11aBe)(a+bx)^7}{9240e(bd - ae)^5(d+ex)^7}$$

output

```
-1/11*(-A*e+B*d)*(b*x+a)^7/e/(-a*e+b*d)/(e*x+d)^11+1/110*(4*A*b*e-11*B*a*e
+7*B*b*d)*(b*x+a)^7/e/(-a*e+b*d)^2/(e*x+d)^10+1/330*b*(4*A*b*e-11*B*a*e+7*
B*b*d)*(b*x+a)^7/e/(-a*e+b*d)^3/(e*x+d)^9+1/1320*b^2*(4*A*b*e-11*B*a*e+7*B
*b*d)*(b*x+a)^7/e/(-a*e+b*d)^4/(e*x+d)^8+1/9240*b^3*(4*A*b*e-11*B*a*e+7*B
*b*d)*(b*x+a)^7/e/(-a*e+b*d)^5/(e*x+d)^7
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 605 vs. $2(235) = 470$.

Time = 0.16 (sec) , antiderivative size = 605, normalized size of antiderivative = 2.57

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{12}} dx = \frac{84a^6e^6(10Ae + B(d + 11ex)) + 56a^5be^5(9Ae(d + 11ex) + 2B(d^2 + 11dex + 55e^2x^2)) + 35a^4b^2e^4(8Ae(d + 11ex) + 2B(d^2 + 11dex + 55e^2x^2)) + 35a^3b^3e^3(7Ae(d + 11ex) + 2B(d^2 + 11dex + 55e^2x^2)) + 35a^2b^4e^2(6Ae(d + 11ex) + 2B(d^2 + 11dex + 55e^2x^2)) + 35ab^5e(5Ae(d + 11ex) + 2B(d^2 + 11dex + 55e^2x^2)) + 35b^6(4Ae(d + 11ex) + 2B(d^2 + 11dex + 55e^2x^2))}{e^8(d + ex)^{11}}$$

input

```
Integrate[((a + b*x)^6*(A + B*x))/(d + e*x)^12,x]
```

output

```
-1/9240*(84*a^6*e^6*(10*A*e + B*(d + 11*e*x)) + 56*a^5*b*e^5*(9*A*e*(d + 11*e*x) + 2*B*(d^2 + 11*d*e*x + 55*e^2*x^2)) + 35*a^4*b^2*e^4*(8*A*e*(d^2 + 11*d*e*x + 55*e^2*x^2) + 3*B*(d^3 + 11*d^2*e*x + 55*d*e^2*x^2 + 165*e^3*x^3)) + 20*a^3*b^3*e^3*(7*A*e*(d^3 + 11*d^2*e*x + 55*d*e^2*x^2 + 165*e^3*x^3) + 4*B*(d^4 + 11*d^3*e*x + 55*d^2*e^2*x^2 + 165*d*e^3*x^3 + 330*e^4*x^4)) + 10*a^2*b^4*e^2*(6*A*e*(d^4 + 11*d^3*e*x + 55*d^2*e^2*x^2 + 165*d*e^3*x^3 + 330*e^4*x^4) + 5*B*(d^5 + 11*d^4*e*x + 55*d^3*e^2*x^2 + 165*d^2*e^3*x^3 + 330*d*e^4*x^4 + 462*e^5*x^5)) + 4*a*b^5*e*(5*A*e*(d^5 + 11*d^4*e*x + 55*d^3*e^2*x^2 + 165*d^2*e^3*x^3 + 330*d*e^4*x^4 + 462*e^5*x^5) + 6*B*(d^6 + 11*d^5*e*x + 55*d^4*e^2*x^2 + 165*d^3*e^3*x^3 + 330*d^2*e^4*x^4 + 462*d*e^5*x^5 + 462*e^6*x^6)) + b^6*(4*A*e*(d^6 + 11*d^5*e*x + 55*d^4*e^2*x^2 + 165*d^3*e^3*x^3 + 330*d^2*e^4*x^4 + 462*d*e^5*x^5 + 462*e^6*x^6) + 7*B*(d^7 + 11*d^6*e*x + 55*d^5*e^2*x^2 + 165*d^4*e^3*x^3 + 330*d^3*e^4*x^4 + 462*d^2*e^5*x^5 + 462*d*e^6*x^6 + 330*e^7*x^7)))/(e^8*(d + e*x)^11)
```

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 219, normalized size of antiderivative = 0.93, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {87, 55, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{12}} dx \\
& \quad \downarrow 87 \\
& \frac{(-11aBe + 4Abe + 7bBd) \int \frac{(a+bx)^6}{(d+ex)^{11}} dx}{11e(bd-ae)} - \frac{(a+bx)^7(Bd-Ae)}{11e(d+ex)^{11}(bd-ae)} \\
& \quad \downarrow 55 \\
& \frac{(-11aBe + 4Abe + 7bBd) \left(\frac{3b \int \frac{(a+bx)^6}{(d+ex)^{10}} dx}{10(bd-ae)} + \frac{(a+bx)^7}{10(d+ex)^{10}(bd-ae)} \right)}{11e(bd-ae)} - \frac{(a+bx)^7(Bd-Ae)}{11e(d+ex)^{11}(bd-ae)} \\
& \quad \downarrow 55 \\
& \frac{(-11aBe + 4Abe + 7bBd) \left(\frac{3b \left(\frac{2b \int \frac{(a+bx)^6}{(d+ex)^9} dx}{9(bd-ae)} + \frac{(a+bx)^7}{9(d+ex)^9(bd-ae)} \right)}{10(bd-ae)} + \frac{(a+bx)^7}{10(d+ex)^{10}(bd-ae)} \right)}{11e(bd-ae)} - \frac{(a+bx)^7(Bd-Ae)}{11e(d+ex)^{11}(bd-ae)} \\
& \quad \downarrow 55 \\
& \frac{(-11aBe + 4Abe + 7bBd) \left(\frac{3b \left(\frac{b \int \frac{(a+bx)^6}{(d+ex)^8} dx}{8(bd-ae)} + \frac{(a+bx)^7}{8(d+ex)^8(bd-ae)} \right)}{9(bd-ae)} + \frac{(a+bx)^7}{9(d+ex)^9(bd-ae)} \right)}{10(bd-ae)} + \frac{(a+bx)^7}{10(d+ex)^{10}(bd-ae)} \\
& \quad \downarrow 48 \\
& \frac{11e(bd-ae)}{11e(d+ex)^{11}(bd-ae)} \frac{(a+bx)^7(Bd-Ae)}{11e(d+ex)^{11}(bd-ae)}
\end{aligned}$$

$$\left(\frac{\frac{(a+bx)^7}{10(d+ex)^{10}(bd-ae)} + \frac{3b \left(\frac{(a+bx)^7}{9(d+ex)^9(bd-ae)} + \frac{2b \left(\frac{b(a+bx)^7}{56(d+ex)^7(bd-ae)^2} + \frac{(a+bx)^7}{8(d+ex)^8(bd-ae)} \right)}{9(bd-ae)}}{10(bd-ae)}}{11e(d+ex)^{11}(bd-ae)} \right) (-11aBe + 4Abe + 7bBd)$$

$$\frac{11e(bd-ae)(a+bx)^7(Bd-Ae)}{11e(d+ex)^{11}(bd-ae)}$$

input `Int[((a + b*x)^6*(A + B*x))/(d + e*x)^12,x]`

output `-1/11*((B*d - A*e)*(a + b*x)^7)/(e*(b*d - a*e)*(d + e*x)^11) + ((7*b*B*d + 4*A*b*e - 11*a*B*e)*((a + b*x)^7/(10*(b*d - a*e)*(d + e*x)^10) + (3*b*((a + b*x)^7/(9*(b*d - a*e)*(d + e*x)^9) + (2*b*((a + b*x)^7/(8*(b*d - a*e)*(d + e*x)^8) + (b*(a + b*x)^7)/(56*(b*d - a*e)^2*(d + e*x)^7)))/(9*(b*d - a*e))))/(10*(b*d - a*e)))/(11*e*(b*d - a*e))`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 55 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(Simplify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])`

output

```
(-1/4*b^6*B/e*x^7-1/20*b^5/e^2*(4*A*b*e+24*B*a*e+7*B*b*d)*x^6-1/20*b^4/e^3
*(20*A*a*b*e^2+4*A*b^2*d*e+50*B*a^2*e^2+24*B*a*b*d*e+7*B*b^2*d^2)*x^5-1/28
*b^3/e^4*(60*A*a^2*b*e^3+20*A*a*b^2*d*e^2+4*A*b^3*d^2*e+80*B*a^3*e^3+50*B*
a^2*b*d*e^2+24*B*a*b^2*d^2*e+7*B*b^3*d^3)*x^4-1/56*b^2/e^5*(140*A*a^3*b*e^
4+60*A*a^2*b^2*d*e^3+20*A*a*b^3*d^2*e^2+4*A*b^4*d^3*e+105*B*a^4*e^4+80*B*a
^3*b*d*e^3+50*B*a^2*b^2*d^2*e^2+24*B*a*b^3*d^3*e+7*B*b^4*d^4)*x^3-1/168*b/
e^6*(280*A*a^4*b*e^5+140*A*a^3*b^2*d*e^4+60*A*a^2*b^3*d^2*e^3+20*A*a*b^4*d
^3*e^2+4*A*b^5*d^4*e+112*B*a^5*e^5+105*B*a^4*b*d*e^4+80*B*a^3*b^2*d^2*e^3+
50*B*a^2*b^3*d^3*e^2+24*B*a*b^4*d^4*e+7*B*b^5*d^5)*x^2-1/840/e^7*(504*A*a^
5*b*e^6+280*A*a^4*b^2*d*e^5+140*A*a^3*b^3*d^2*e^4+60*A*a^2*b^4*d^3*e^3+20*
A*a*b^5*d^4*e^2+4*A*b^6*d^5*e+84*B*a^6*e^6+112*B*a^5*b*d*e^5+105*B*a^4*b^2
*d^2*e^4+80*B*a^3*b^3*d^3*e^3+50*B*a^2*b^4*d^4*e^2+24*B*a*b^5*d^5*e+7*B*b^
6*d^6)*x-1/9240/e^8*(840*A*a^6*e^7+504*A*a^5*b*d*e^6+280*A*a^4*b^2*d^2*e^5
+140*A*a^3*b^3*d^3*e^4+60*A*a^2*b^4*d^4*e^3+20*A*a*b^5*d^5*e^2+4*A*b^6*d^6
*e+84*B*a^6*d*e^6+112*B*a^5*b*d^2*e^5+105*B*a^4*b^2*d^3*e^4+80*B*a^3*b^3*d
^4*e^3+50*B*a^2*b^4*d^5*e^2+24*B*a*b^5*d^6*e+7*B*b^6*d^7))/(e*x+d)^11
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 883 vs. $2(225) = 450$.

Time = 0.09 (sec) , antiderivative size = 883, normalized size of antiderivative = 3.76

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{12}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^6*(B*x+A)/(e*x+d)^12,x, algorithm="fricas")
```

output

```

-1/9240*(2310*B*b^6*e^7*x^7 + 7*B*b^6*d^7 + 840*A*a^6*e^7 + 4*(6*B*a*b^5 +
A*b^6)*d^6*e + 10*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 + 20*(4*B*a^3*b^3 + 3
*A*a^2*b^4)*d^4*e^3 + 35*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 + 56*(2*B*a^5
*b + 5*A*a^4*b^2)*d^2*e^5 + 84*(B*a^6 + 6*A*a^5*b)*d*e^6 + 462*(7*B*b^6*d*
e^6 + 4*(6*B*a*b^5 + A*b^6)*e^7)*x^6 + 462*(7*B*b^6*d^2*e^5 + 4*(6*B*a*b^5
+ A*b^6)*d*e^6 + 10*(5*B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^5 + 330*(7*B*b^6*d^3
*e^4 + 4*(6*B*a*b^5 + A*b^6)*d^2*e^5 + 10*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6
+ 20*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^7)*x^4 + 165*(7*B*b^6*d^4*e^3 + 4*(6*B*
a*b^5 + A*b^6)*d^3*e^4 + 10*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 + 20*(4*B*a^
3*b^3 + 3*A*a^2*b^4)*d*e^6 + 35*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^3 + 55*
(7*B*b^6*d^5*e^2 + 4*(6*B*a*b^5 + A*b^6)*d^4*e^3 + 10*(5*B*a^2*b^4 + 2*A*
a*b^5)*d^3*e^4 + 20*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + 35*(3*B*a^4*b^2 +
4*A*a^3*b^3)*d*e^6 + 56*(2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 11*(7*B*b^6*
d^6*e + 4*(6*B*a*b^5 + A*b^6)*d^5*e^2 + 10*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*
e^3 + 20*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 + 35*(3*B*a^4*b^2 + 4*A*a^3*b^
3)*d^2*e^5 + 56*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^6 + 84*(B*a^6 + 6*A*a^5*b)*
e^7)*x)/(e^19*x^11 + 11*d*e^18*x^10 + 55*d^2*e^17*x^9 + 165*d^3*e^16*x^8 +
330*d^4*e^15*x^7 + 462*d^5*e^14*x^6 + 462*d^6*e^13*x^5 + 330*d^7*e^12*x^4
+ 165*d^8*e^11*x^3 + 55*d^9*e^10*x^2 + 11*d^10*e^9*x + d^11*e^8)

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{12}} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**6*(B*x+A)/(e*x+d)**12,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 883 vs. $2(225) = 450$.

Time = 0.13 (sec) , antiderivative size = 883, normalized size of antiderivative = 3.76

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{12}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^12,x, algorithm="maxima")`

output

```
-1/9240*(2310*B*b^6*e^7*x^7 + 7*B*b^6*d^7 + 840*A*a^6*e^7 + 4*(6*B*a*b^5 +
A*b^6)*d^6*e + 10*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 + 20*(4*B*a^3*b^3 + 3
*A*a^2*b^4)*d^4*e^3 + 35*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 + 56*(2*B*a^5
*b + 5*A*a^4*b^2)*d^2*e^5 + 84*(B*a^6 + 6*A*a^5*b)*d*e^6 + 462*(7*B*b^6*d*
e^6 + 4*(6*B*a*b^5 + A*b^6)*e^7)*x^6 + 462*(7*B*b^6*d^2*e^5 + 4*(6*B*a*b^5
+ A*b^6)*d*e^6 + 10*(5*B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^5 + 330*(7*B*b^6*d^3
*e^4 + 4*(6*B*a*b^5 + A*b^6)*d^2*e^5 + 10*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6
+ 20*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^7)*x^4 + 165*(7*B*b^6*d^4*e^3 + 4*(6*B*
a*b^5 + A*b^6)*d^3*e^4 + 10*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 + 20*(4*B*a^
3*b^3 + 3*A*a^2*b^4)*d*e^6 + 35*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^3 + 55*
(7*B*b^6*d^5*e^2 + 4*(6*B*a*b^5 + A*b^6)*d^4*e^3 + 10*(5*B*a^2*b^4 + 2*A*a
*b^5)*d^3*e^4 + 20*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + 35*(3*B*a^4*b^2 +
4*A*a^3*b^3)*d*e^6 + 56*(2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 11*(7*B*b^6*
d^6*e + 4*(6*B*a*b^5 + A*b^6)*d^5*e^2 + 10*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e
^3 + 20*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 + 35*(3*B*a^4*b^2 + 4*A*a^3*b^
3)*d^2*e^5 + 56*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^6 + 84*(B*a^6 + 6*A*a^5*b)*e
^7)*x)/(e^19*x^11 + 11*d*e^18*x^10 + 55*d^2*e^17*x^9 + 165*d^3*e^16*x^8 +
330*d^4*e^15*x^7 + 462*d^5*e^14*x^6 + 462*d^6*e^13*x^5 + 330*d^7*e^12*x^4
+ 165*d^8*e^11*x^3 + 55*d^9*e^10*x^2 + 11*d^10*e^9*x + d^11*e^8)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 912 vs. $2(225) = 450$.

Time = 0.13 (sec) , antiderivative size = 912, normalized size of antiderivative = 3.88

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{12}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^12,x, algorithm="giac")`

output

$$\begin{aligned}
 & -1/9240*(2310*B*b^6*e^7*x^7 + 3234*B*b^6*d*e^6*x^6 + 11088*B*a*b^5*e^7*x^6 \\
 & + 1848*A*b^6*e^7*x^6 + 3234*B*b^6*d^2*e^5*x^5 + 11088*B*a*b^5*d*e^6*x^5 + \\
 & 1848*A*b^6*d*e^6*x^5 + 23100*B*a^2*b^4*e^7*x^5 + 9240*A*a*b^5*e^7*x^5 + 2 \\
 & 310*B*b^6*d^3*e^4*x^4 + 7920*B*a*b^5*d^2*e^5*x^4 + 1320*A*b^6*d^2*e^5*x^4 \\
 & + 16500*B*a^2*b^4*d*e^6*x^4 + 6600*A*a*b^5*d*e^6*x^4 + 26400*B*a^3*b^3*e^7 \\
 & *x^4 + 19800*A*a^2*b^4*e^7*x^4 + 1155*B*b^6*d^4*e^3*x^3 + 3960*B*a*b^5*d^3 \\
 & *e^4*x^3 + 660*A*b^6*d^3*e^4*x^3 + 8250*B*a^2*b^4*d^2*e^5*x^3 + 3300*A*a*b \\
 & ^5*d^2*e^5*x^3 + 13200*B*a^3*b^3*d*e^6*x^3 + 9900*A*a^2*b^4*d*e^6*x^3 + 17 \\
 & 325*B*a^4*b^2*e^7*x^3 + 23100*A*a^3*b^3*e^7*x^3 + 385*B*b^6*d^5*e^2*x^2 + \\
 & 1320*B*a*b^5*d^4*e^3*x^2 + 220*A*b^6*d^4*e^3*x^2 + 2750*B*a^2*b^4*d^3*e^4* \\
 & x^2 + 1100*A*a*b^5*d^3*e^4*x^2 + 4400*B*a^3*b^3*d^2*e^5*x^2 + 3300*A*a^2*b \\
 & ^4*d^2*e^5*x^2 + 5775*B*a^4*b^2*d*e^6*x^2 + 7700*A*a^3*b^3*d*e^6*x^2 + 616 \\
 & 0*B*a^5*b*e^7*x^2 + 15400*A*a^4*b^2*e^7*x^2 + 77*B*b^6*d^6*e*x + 264*B*a*b \\
 & ^5*d^5*e^2*x + 44*A*b^6*d^5*e^2*x + 550*B*a^2*b^4*d^4*e^3*x + 220*A*a*b^5* \\
 & d^4*e^3*x + 880*B*a^3*b^3*d^3*e^4*x + 660*A*a^2*b^4*d^3*e^4*x + 1155*B*a^4 \\
 & *b^2*d^2*e^5*x + 1540*A*a^3*b^3*d^2*e^5*x + 1232*B*a^5*b*d*e^6*x + 3080*A* \\
 & a^4*b^2*d*e^6*x + 924*B*a^6*e^7*x + 5544*A*a^5*b*e^7*x + 7*B*b^6*d^7 + 24* \\
 & B*a*b^5*d^6*e + 4*A*b^6*d^6*e + 50*B*a^2*b^4*d^5*e^2 + 20*A*a*b^5*d^5*e^2 \\
 & + 80*B*a^3*b^3*d^4*e^3 + 60*A*a^2*b^4*d^4*e^3 + 105*B*a^4*b^2*d^3*e^4 + 14 \\
 & 0*A*a^3*b^3*d^3*e^4 + 112*B*a^5*b*d^2*e^5 + 280*A*a^4*b^2*d^2*e^5 + 84*...
 \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 899, normalized size of antiderivative = 3.83

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{12}} dx =$$

$$\frac{84 B a^6 d e^6 + 840 A a^6 e^7 + 112 B a^5 b d^2 e^5 + 504 A a^5 b d e^6 + 105 B a^4 b^2 d^3 e^4 + 280 A a^4 b^2 d^2 e^5 + 80 B a^3 b^3 d^4 e^3 + 140 A a^3 b^3 d^3 e^4 + 50 B a^2 b^4 d^5 e^2 + 20 A a^2 b^4 d^4 e^3 + 44 B a b^5 d^6 e + 4 A a b^5 d^5 e^2 + 77 B b^6 d^7 + 24 B a b^5 d^6 e + 4 A a b^6 d^6 e + 50 B a^2 b^4 d^5 e^2 + 20 A a b^5 d^5 e^2 + 80 B a^3 b^3 d^4 e^3 + 60 A a^2 b^4 d^4 e^3 + 105 B a^4 b^2 d^3 e^4 + 140 A a^3 b^3 d^3 e^4 + 112 B a^5 b d^2 e^5 + 280 A a^4 b^2 d^2 e^5 + 84 B a^6 e^7}{9240 e^8}$$

input `int(((A + B*x)*(a + b*x)^6)/(d + e*x)^12,x)`

output

```

-((840*A*a^6*e^7 + 7*B*b^6*d^7 + 4*A*b^6*d^6*e + 84*B*a^6*d*e^6 + 20*A*a*b
^5*d^5*e^2 + 112*B*a^5*b*d^2*e^5 + 60*A*a^2*b^4*d^4*e^3 + 140*A*a^3*b^3*d
^3*e^4 + 280*A*a^4*b^2*d^2*e^5 + 50*B*a^2*b^4*d^5*e^2 + 80*B*a^3*b^3*d^4*e
^3 + 105*B*a^4*b^2*d^3*e^4 + 504*A*a^5*b*d*e^6 + 24*B*a*b^5*d^6*e)/(9240*e
^8) + (x*(84*B*a^6*e^6 + 7*B*b^6*d^6 + 504*A*a^5*b*e^6 + 4*A*b^6*d^5*e + 20
*A*a*b^5*d^4*e^2 + 280*A*a^4*b^2*d*e^5 + 60*A*a^2*b^4*d^3*e^3 + 140*A*a^3*
b^3*d^2*e^4 + 50*B*a^2*b^4*d^4*e^2 + 80*B*a^3*b^3*d^3*e^3 + 105*B*a^4*b^2*
d^2*e^4 + 24*B*a*b^5*d^5*e + 112*B*a^5*b*d*e^5))/(840*e^7) + (b^3*x^4*(80*
B*a^3*e^3 + 7*B*b^3*d^3 + 60*A*a^2*b*e^3 + 4*A*b^3*d^2*e + 20*A*a*b^2*d*e
^2 + 24*B*a*b^2*d^2*e + 50*B*a^2*b*d*e^2))/(28*e^4) + (b^5*x^6*(4*A*b*e + 2
4*B*a*e + 7*B*b*d))/(20*e^2) + (b*x^2*(112*B*a^5*e^5 + 7*B*b^5*d^5 + 280*A
*a^4*b*e^5 + 4*A*b^5*d^4*e + 20*A*a*b^4*d^3*e^2 + 140*A*a^3*b^2*d*e^4 + 60
*A*a^2*b^3*d^2*e^3 + 50*B*a^2*b^3*d^3*e^2 + 80*B*a^3*b^2*d^2*e^3 + 24*B*a*
b^4*d^4*e + 105*B*a^4*b*d*e^4))/(168*e^6) + (b^2*x^3*(105*B*a^4*e^4 + 7*B*
b^4*d^4 + 140*A*a^3*b*e^4 + 4*A*b^4*d^3*e + 20*A*a*b^3*d^2*e^2 + 60*A*a^2*
b^2*d*e^3 + 50*B*a^2*b^2*d^2*e^2 + 24*B*a*b^3*d^3*e + 80*B*a^3*b*d*e^3))/(
56*e^5) + (b^4*x^5*(50*B*a^2*e^2 + 7*B*b^2*d^2 + 20*A*a*b*e^2 + 4*A*b^2*d*
e + 24*B*a*b*d*e))/(20*e^3) + (B*b^6*x^7)/(4*e))/(d^11 + e^11*x^11 + 11*d*
e^10*x^10 + 55*d^9*e^2*x^2 + 165*d^8*e^3*x^3 + 330*d^7*e^4*x^4 + 462*d^6*
e^5*x^5 + 462*d^5*e^6*x^6 + 330*d^4*e^7*x^7 + 165*d^3*e^8*x^8 + 55*d^2*e...

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 607, normalized size of antiderivative = 2.58

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{12}} dx$$

$$= \frac{-330b^7e^7x^7 - 1848ab^6e^7x^6 - 462b^7de^6x^6 - 4620a^2b^5e^7x^5 - 1848ab^6de^6x^5 - 462b^7d^2e^5x^5 - 6600a^3b^4e^7x^4 - 1848a^2b^5de^6x^4 - 462b^7d^2e^5x^4 - 1848ab^6de^6x^3 - 462b^7d^2e^5x^3 - 6600a^3b^4e^7x^2 - 1848a^2b^5de^6x^2 - 462b^7d^2e^5x^2 - 1848ab^6de^6x - 462b^7d^2e^5x - 6600a^3b^4e^7 - 1848a^2b^5de^6 - 462b^7d^2e^5}{e^{11}}$$

input

```
int((b*x+a)^6*(B*x+A)/(e*x+d)^12,x)
```

output

```
( - 120*a**7*e**7 - 84*a**6*b*d*e**6 - 924*a**6*b*e**7*x - 56*a**5*b**2*d*
*2*e**5 - 616*a**5*b**2*d*e**6*x - 3080*a**5*b**2*e**7*x**2 - 35*a**4*b**3
*d**3*e**4 - 385*a**4*b**3*d**2*e**5*x - 1925*a**4*b**3*d*e**6*x**2 - 5775
*a**4*b**3*e**7*x**3 - 20*a**3*b**4*d**4*e**3 - 220*a**3*b**4*d**3*e**4*x
- 1100*a**3*b**4*d**2*e**5*x**2 - 3300*a**3*b**4*d*e**6*x**3 - 6600*a**3*b
**4*e**7*x**4 - 10*a**2*b**5*d**5*e**2 - 110*a**2*b**5*d**4*e**3*x - 550*a
**2*b**5*d**3*e**4*x**2 - 1650*a**2*b**5*d**2*e**5*x**3 - 3300*a**2*b**5*d
*e**6*x**4 - 4620*a**2*b**5*e**7*x**5 - 4*a*b**6*d**6*e - 44*a*b**6*d**5*e
**2*x - 220*a*b**6*d**4*e**3*x**2 - 660*a*b**6*d**3*e**4*x**3 - 1320*a*b**
6*d**2*e**5*x**4 - 1848*a*b**6*d*e**6*x**5 - 1848*a*b**6*e**7*x**6 - b**7*
d**7 - 11*b**7*d**6*e*x - 55*b**7*d**5*e**2*x**2 - 165*b**7*d**4*e**3*x**3
- 330*b**7*d**3*e**4*x**4 - 462*b**7*d**2*e**5*x**5 - 462*b**7*d*e**6*x**
6 - 330*b**7*e**7*x**7)/(1320*e**8*(d**11 + 11*d**10*e*x + 55*d**9*e**2*x*
*2 + 165*d**8*e**3*x**3 + 330*d**7*e**4*x**4 + 462*d**6*e**5*x**5 + 462*d
**5*e**6*x**6 + 330*d**4*e**7*x**7 + 165*d**3*e**8*x**8 + 55*d**2*e**9*x**9
+ 11*d**10*x**10 + e**11*x**11))
```

3.62 $\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{13}} dx$

Optimal result	626
Mathematica [B] (verified)	627
Rubi [A] (verified)	627
Maple [B] (verified)	629
Fricas [B] (verification not implemented)	630
Sympy [F(-1)]	631
Maxima [B] (verification not implemented)	631
Giac [B] (verification not implemented)	632
Mupad [B] (verification not implemented)	633
Reduce [B] (verification not implemented)	634

Optimal result

Integrand size = 20, antiderivative size = 292

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{13}} dx = \frac{(bd-ae)^6(Bd-Ae)}{12e^8(d+ex)^{12}} - \frac{(bd-ae)^5(7bBd-6Abe-aBe)}{11e^8(d+ex)^{11}} + \frac{3b(bd-ae)^4(7bBd-5Abe-2aBe)}{10e^8(d+ex)^{10}} - \frac{5b^2(bd-ae)^3(7bBd-4Abe-3aBe)}{9e^8(d+ex)^9} + \frac{5b^3(bd-ae)^2(7bBd-3Abe-4aBe)}{8e^8(d+ex)^8} - \frac{3b^4(bd-ae)(7bBd-2Abe-5aBe)}{7e^8(d+ex)^7} + \frac{b^5(7bBd-Abe-6aBe)}{6e^8(d+ex)^6} - \frac{b^6B}{5e^8(d+ex)^5}$$

output

```
1/12*(-a*e+b*d)^6*(-A*e+B*d)/e^8/(e*x+d)^12-1/11*(-a*e+b*d)^5*(-6*A*b*e-B*
a*e+7*B*b*d)/e^8/(e*x+d)^11+3/10*b*(-a*e+b*d)^4*(-5*A*b*e-2*B*a*e+7*B*b*d)
/e^8/(e*x+d)^10-5/9*b^2*(-a*e+b*d)^3*(-4*A*b*e-3*B*a*e+7*B*b*d)/e^8/(e*x+d)
)^9+5/8*b^3*(-a*e+b*d)^2*(-3*A*b*e-4*B*a*e+7*B*b*d)/e^8/(e*x+d)^8-3/7*b^4*
(-a*e+b*d)*(-2*A*b*e-5*B*a*e+7*B*b*d)/e^8/(e*x+d)^7+1/6*b^5*(-A*b*e-6*B*a*
e+7*B*b*d)/e^8/(e*x+d)^6-1/5*b^6*B/e^8/(e*x+d)^5
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 600 vs. $2(292) = 584$.

Time = 0.16 (sec) , antiderivative size = 600, normalized size of antiderivative = 2.05

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{13}} dx = \frac{210a^6e^6(11Ae + B(d + 12ex)) + 252a^5be^5(5Ae(d + 12ex) + B(d^2 + 12dex + 66e^2x^2)) + 210a^4b^2e^4(3Ae(d + 12ex) + B(d^2 + 12dex + 66e^2x^2)) + 210a^3b^3e^3(2Ae(d + 12ex) + B(d^2 + 12dex + 66e^2x^2)) + 210a^2b^4e^2(7Ae(d + 12ex) + B(d^2 + 12dex + 66e^2x^2)) + 210ab^5e(5Ae(d + 12ex) + B(d^2 + 12dex + 66e^2x^2)) + 210a^6e^6(11Ae + B(d + 12ex)) + 252a^5be^5(5Ae(d + 12ex) + B(d^2 + 12dex + 66e^2x^2)) + 210a^4b^2e^4(3Ae(d + 12ex) + B(d^2 + 12dex + 66e^2x^2)) + 210a^3b^3e^3(2Ae(d + 12ex) + B(d^2 + 12dex + 66e^2x^2)) + 210a^2b^4e^2(7Ae(d + 12ex) + B(d^2 + 12dex + 66e^2x^2)) + 210ab^5e(5Ae(d + 12ex) + B(d^2 + 12dex + 66e^2x^2)) + 210a^6e^6(11Ae + B(d + 12ex))}{(d + ex)^{12}}$$

input

```
Integrate[((a + b*x)^6*(A + B*x))/(d + e*x)^13,x]
```

output

```
-1/27720*(210*a^6*e^6*(11*A*e + B*(d + 12*e*x)) + 252*a^5*b*e^5*(5*A*e*(d + 12*e*x) + B*(d^2 + 12*d*e*x + 66*e^2*x^2)) + 210*a^4*b^2*e^4*(3*A*e*(d^2 + 12*d*e*x + 66*e^2*x^2) + B*(d^3 + 12*d^2*e*x + 66*d*e^2*x^2 + 220*e^3*x^3)) + 140*a^3*b^3*e^3*(2*A*e*(d^3 + 12*d^2*e*x + 66*d*e^2*x^2 + 220*e^3*x^3) + B*(d^4 + 12*d^3*e*x + 66*d^2*e^2*x^2 + 220*d*e^3*x^3 + 495*e^4*x^4)) + 15*a^2*b^4*e^2*(7*A*e*(d^4 + 12*d^3*e*x + 66*d^2*e^2*x^2 + 220*d*e^3*x^3 + 495*e^4*x^4) + 5*B*(d^5 + 12*d^4*e*x + 66*d^3*e^2*x^2 + 220*d^2*e^3*x^3 + 495*d*e^4*x^4 + 792*e^5*x^5)) + 30*a*b^5*e*(A*e*(d^5 + 12*d^4*e*x + 66*d^3*e^2*x^2 + 220*d^2*e^3*x^3 + 495*d*e^4*x^4 + 792*e^5*x^5) + B*(d^6 + 12*d^5*e*x + 66*d^4*e^2*x^2 + 220*d^3*e^3*x^3 + 495*d^2*e^4*x^4 + 792*d*e^5*x^5 + 924*e^6*x^6)) + b^6*(5*A*e*(d^6 + 12*d^5*e*x + 66*d^4*e^2*x^2 + 220*d^3*e^3*x^3 + 495*d^2*e^4*x^4 + 792*d*e^5*x^5 + 924*e^6*x^6) + 7*B*(d^7 + 12*d^6*e*x + 66*d^5*e^2*x^2 + 220*d^4*e^3*x^3 + 495*d^3*e^4*x^4 + 792*d^2*e^5*x^5 + 924*d*e^6*x^6 + 792*e^7*x^7)))/(e^8*(d + e*x)^12)
```

Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 292, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{13}} dx$$

↓ 86

$$\int \left(\frac{b^5(6aBe + Abe - 7bBd)}{e^7(d + ex)^7} - \frac{3b^4(bd - ae)(5aBe + 2Abe - 7bBd)}{e^7(d + ex)^8} + \frac{5b^3(bd - ae)^2(4aBe + 3Abe - 7bBd)}{e^7(d + ex)^9} - \dots \right) dx$$

↓ 2009

$$\begin{aligned} & \frac{b^5(-6aBe - Abe + 7bBd)}{6e^8(d + ex)^6} - \frac{3b^4(bd - ae)(-5aBe - 2Abe + 7bBd)}{7e^8(d + ex)^7} + \\ & \frac{5b^3(bd - ae)^2(-4aBe - 3Abe + 7bBd)}{8e^8(d + ex)^8} - \frac{5b^2(bd - ae)^3(-3aBe - 4Abe + 7bBd)}{9e^8(d + ex)^9} + \\ & \frac{3b(bd - ae)^4(-2aBe - 5Abe + 7bBd)}{10e^8(d + ex)^{10}} - \frac{(bd - ae)^5(-aBe - 6Abe + 7bBd)}{11e^8(d + ex)^{11}} + \\ & \frac{(bd - ae)^6(Bd - Ae)}{12e^8(d + ex)^{12}} - \frac{b^6B}{5e^8(d + ex)^5} \end{aligned}$$

input

```
Int[((a + b*x)^6*(A + B*x))/(d + e*x)^13,x]
```

output

```
((b*d - a*e)^6*(B*d - A*e))/(12*e^8*(d + e*x)^12) - ((b*d - a*e)^5*(7*b*B*d - 6*A*b*e - a*B*e))/(11*e^8*(d + e*x)^11) + (3*b*(b*d - a*e)^4*(7*b*B*d - 5*A*b*e - 2*a*B*e))/(10*e^8*(d + e*x)^10) - (5*b^2*(b*d - a*e)^3*(7*b*B*d - 4*A*b*e - 3*a*B*e))/(9*e^8*(d + e*x)^9) + (5*b^3*(b*d - a*e)^2*(7*b*B*d - 3*A*b*e - 4*a*B*e))/(8*e^8*(d + e*x)^8) - (3*b^4*(b*d - a*e)*(7*b*B*d - 2*A*b*e - 5*a*B*e))/(7*e^8*(d + e*x)^7) + (b^5*(7*b*B*d - A*b*e - 6*a*B*e))/(6*e^8*(d + e*x)^6) - (b^6*B)/(5*e^8*(d + e*x)^5)
```

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.))^(n_.)*((e_.) + (f_.)*(x_.))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```


Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 894 vs. $2(276) = 552$.

Time = 0.10 (sec) , antiderivative size = 894, normalized size of antiderivative = 3.06

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{13}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^13,x, algorithm="fricas")`

output

```
-1/27720*(5544*B*b^6*e^7*x^7 + 7*B*b^6*d^7 + 2310*A*a^6*e^7 + 5*(6*B*a*b^5
+ A*b^6)*d^6*e + 15*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 + 35*(4*B*a^3*b^3 +
3*A*a^2*b^4)*d^4*e^3 + 70*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 + 126*(2*B*
a^5*b + 5*A*a^4*b^2)*d^2*e^5 + 210*(B*a^6 + 6*A*a^5*b)*d*e^6 + 924*(7*B*b^
6*d*e^6 + 5*(6*B*a*b^5 + A*b^6)*e^7)*x^6 + 792*(7*B*b^6*d^2*e^5 + 5*(6*B*a
*b^5 + A*b^6)*d*e^6 + 15*(5*B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^5 + 495*(7*B*b^6
*d^3*e^4 + 5*(6*B*a*b^5 + A*b^6)*d^2*e^5 + 15*(5*B*a^2*b^4 + 2*A*a*b^5)*d*
e^6 + 35*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^7)*x^4 + 220*(7*B*b^6*d^4*e^3 + 5*(
6*B*a*b^5 + A*b^6)*d^3*e^4 + 15*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 + 35*(4*
B*a^3*b^3 + 3*A*a^2*b^4)*d*e^6 + 70*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^3 +
66*(7*B*b^6*d^5*e^2 + 5*(6*B*a*b^5 + A*b^6)*d^4*e^3 + 15*(5*B*a^2*b^4 + 2
*A*a*b^5)*d^3*e^4 + 35*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + 70*(3*B*a^4*b
^2 + 4*A*a^3*b^3)*d*e^6 + 126*(2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 12*(7*B
*b^6*d^6*e + 5*(6*B*a*b^5 + A*b^6)*d^5*e^2 + 15*(5*B*a^2*b^4 + 2*A*a*b^5)*
d^4*e^3 + 35*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 + 70*(3*B*a^4*b^2 + 4*A*a
^3*b^3)*d^2*e^5 + 126*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^6 + 210*(B*a^6 + 6*A*a
^5*b)*e^7)*x)/(e^20*x^12 + 12*d*e^19*x^11 + 66*d^2*e^18*x^10 + 220*d^3*e^1
7*x^9 + 495*d^4*e^16*x^8 + 792*d^5*e^15*x^7 + 924*d^6*e^14*x^6 + 792*d^7*e
^13*x^5 + 495*d^8*e^12*x^4 + 220*d^9*e^11*x^3 + 66*d^10*e^10*x^2 + 12*d^11
*e^9*x + d^12*e^8)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{13}} dx = \text{Timed out}$$

input `integrate((b*x+a)**6*(B*x+A)/(e*x+d)**13,x)`output `Timed out`**Maxima [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 894 vs. $2(276) = 552$.

Time = 0.08 (sec) , antiderivative size = 894, normalized size of antiderivative = 3.06

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{13}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^13,x, algorithm="maxima")`

output

```

-1/27720*(5544*B*b^6*e^7*x^7 + 7*B*b^6*d^7 + 2310*A*a^6*e^7 + 5*(6*B*a*b^5
+ A*b^6)*d^6*e + 15*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 + 35*(4*B*a^3*b^3 +
3*A*a^2*b^4)*d^4*e^3 + 70*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 + 126*(2*B*
a^5*b + 5*A*a^4*b^2)*d^2*e^5 + 210*(B*a^6 + 6*A*a^5*b)*d*e^6 + 924*(7*B*b^
6*d*e^6 + 5*(6*B*a*b^5 + A*b^6)*e^7)*x^6 + 792*(7*B*b^6*d^2*e^5 + 5*(6*B*a
*b^5 + A*b^6)*d*e^6 + 15*(5*B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^5 + 495*(7*B*b^6
*d^3*e^4 + 5*(6*B*a*b^5 + A*b^6)*d^2*e^5 + 15*(5*B*a^2*b^4 + 2*A*a*b^5)*d*
e^6 + 35*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^7)*x^4 + 220*(7*B*b^6*d^4*e^3 + 5*(
6*B*a*b^5 + A*b^6)*d^3*e^4 + 15*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 + 35*(4*
B*a^3*b^3 + 3*A*a^2*b^4)*d*e^6 + 70*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^3 +
66*(7*B*b^6*d^5*e^2 + 5*(6*B*a*b^5 + A*b^6)*d^4*e^3 + 15*(5*B*a^2*b^4 + 2
*A*a*b^5)*d^3*e^4 + 35*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + 70*(3*B*a^4*b
^2 + 4*A*a^3*b^3)*d*e^6 + 126*(2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 12*(7*B
*b^6*d^6*e + 5*(6*B*a*b^5 + A*b^6)*d^5*e^2 + 15*(5*B*a^2*b^4 + 2*A*a*b^5)*
d^4*e^3 + 35*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 + 70*(3*B*a^4*b^2 + 4*A*a
^3*b^3)*d^2*e^5 + 126*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^6 + 210*(B*a^6 + 6*A*a
^5*b)*e^7)*x)/(e^20*x^12 + 12*d*e^19*x^11 + 66*d^2*e^18*x^10 + 220*d^3*e^1
7*x^9 + 495*d^4*e^16*x^8 + 792*d^5*e^15*x^7 + 924*d^6*e^14*x^6 + 792*d^7*e
^13*x^5 + 495*d^8*e^12*x^4 + 220*d^9*e^11*x^3 + 66*d^10*e^10*x^2 + 12*d^11
*e^9*x + d^12*e^8)

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 912 vs. $2(276) = 552$.

Time = 0.12 (sec) , antiderivative size = 912, normalized size of antiderivative = 3.12

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{13}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^6*(B*x+A)/(e*x+d)^13,x, algorithm="giac")
```

output

```

-1/27720*(5544*B*b^6*e^7*x^7 + 6468*B*b^6*d*e^6*x^6 + 27720*B*a*b^5*e^7*x^
6 + 4620*A*b^6*e^7*x^6 + 5544*B*b^6*d^2*e^5*x^5 + 23760*B*a*b^5*d*e^6*x^5
+ 3960*A*b^6*d*e^6*x^5 + 59400*B*a^2*b^4*e^7*x^5 + 23760*A*a*b^5*e^7*x^5 +
3465*B*b^6*d^3*e^4*x^4 + 14850*B*a*b^5*d^2*e^5*x^4 + 2475*A*b^6*d^2*e^5*x
^4 + 37125*B*a^2*b^4*d*e^6*x^4 + 14850*A*a*b^5*d*e^6*x^4 + 69300*B*a^3*b^3
*e^7*x^4 + 51975*A*a^2*b^4*e^7*x^4 + 1540*B*b^6*d^4*e^3*x^3 + 6600*B*a*b^5
*d^3*e^4*x^3 + 1100*A*b^6*d^3*e^4*x^3 + 16500*B*a^2*b^4*d^2*e^5*x^3 + 6600
*A*a*b^5*d^2*e^5*x^3 + 30800*B*a^3*b^3*d*e^6*x^3 + 23100*A*a^2*b^4*d*e^6*x
^3 + 46200*B*a^4*b^2*e^7*x^3 + 61600*A*a^3*b^3*e^7*x^3 + 462*B*b^6*d^5*e^2
*x^2 + 1980*B*a*b^5*d^4*e^3*x^2 + 330*A*b^6*d^4*e^3*x^2 + 4950*B*a^2*b^4*d
^3*e^4*x^2 + 1980*A*a*b^5*d^3*e^4*x^2 + 9240*B*a^3*b^3*d^2*e^5*x^2 + 6930*
A*a^2*b^4*d^2*e^5*x^2 + 13860*B*a^4*b^2*d*e^6*x^2 + 18480*A*a^3*b^3*d*e^6*
x^2 + 16632*B*a^5*b*e^7*x^2 + 41580*A*a^4*b^2*e^7*x^2 + 84*B*b^6*d^6*e*x +
360*B*a*b^5*d^5*e^2*x + 60*A*b^6*d^5*e^2*x + 900*B*a^2*b^4*d^4*e^3*x + 36
0*A*a*b^5*d^4*e^3*x + 1680*B*a^3*b^3*d^3*e^4*x + 1260*A*a^2*b^4*d^3*e^4*x
+ 2520*B*a^4*b^2*d^2*e^5*x + 3360*A*a^3*b^3*d^2*e^5*x + 3024*B*a^5*b*d*e^6
*x + 7560*A*a^4*b^2*d*e^6*x + 2520*B*a^6*e^7*x + 15120*A*a^5*b*e^7*x + 7*B
*b^6*d^7 + 30*B*a*b^5*d^6*e + 5*A*b^6*d^6*e + 75*B*a^2*b^4*d^5*e^2 + 30*A
*a*b^5*d^5*e^2 + 140*B*a^3*b^3*d^4*e^3 + 105*A*a^2*b^4*d^4*e^3 + 210*B*a^4*
b^2*d^3*e^4 + 280*A*a^3*b^3*d^3*e^4 + 252*B*a^5*b*d^2*e^5 + 630*A*a^4*b...

```

Mupad [B] (verification not implemented)

Time = 1.21 (sec) , antiderivative size = 910, normalized size of antiderivative = 3.12

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{13}} dx =$$

$$\frac{210 B a^6 d e^6 + 2310 A a^6 e^7 + 252 B a^5 b d^2 e^5 + 1260 A a^5 b d e^6 + 210 B a^4 b^2 d^3 e^4 + 630 A a^4 b^2 d^2 e^5 + 140 B a^3 b^3 d^4 e^3 + 280 A a^3 b^3 d^3 e^4 + 75 B a^2 b^4 d^5 e^2 + 30 A a^2 b^4 d^4 e^3 + 105 A a^2 b^4 d^4 e^3 + 210 B a^4 b^2 d^3 e^4 + 280 A a^3 b^3 d^3 e^4 + 252 B a^5 b d^2 e^5 + 630 A a^4 b^2 d^3 e^4}{27720 e^8}$$

input

```
int(((A + B*x)*(a + b*x)^6)/(d + e*x)^13,x)
```

output

```

-((2310*A*a^6*e^7 + 7*B*b^6*d^7 + 5*A*b^6*d^6*e + 210*B*a^6*d*e^6 + 30*A*a
*b^5*d^5*e^2 + 252*B*a^5*b*d^2*e^5 + 105*A*a^2*b^4*d^4*e^3 + 280*A*a^3*b^3
*d^3*e^4 + 630*A*a^4*b^2*d^2*e^5 + 75*B*a^2*b^4*d^5*e^2 + 140*B*a^3*b^3*d^
4*e^3 + 210*B*a^4*b^2*d^3*e^4 + 1260*A*a^5*b*d*e^6 + 30*B*a*b^5*d^6*e)/(27
720*e^8) + (x*(210*B*a^6*e^6 + 7*B*b^6*d^6 + 1260*A*a^5*b*e^6 + 5*A*b^6*d^
5*e + 30*A*a*b^5*d^4*e^2 + 630*A*a^4*b^2*d*e^5 + 105*A*a^2*b^4*d^3*e^3 + 2
80*A*a^3*b^3*d^2*e^4 + 75*B*a^2*b^4*d^4*e^2 + 140*B*a^3*b^3*d^3*e^3 + 210*
B*a^4*b^2*d^2*e^4 + 30*B*a*b^5*d^5*e + 252*B*a^5*b*d*e^5))/(2310*e^7) + (b
^3*x^4*(140*B*a^3*e^3 + 7*B*b^3*d^3 + 105*A*a^2*b*e^3 + 5*A*b^3*d^2*e + 30
*A*a*b^2*d*e^2 + 30*B*a*b^2*d^2*e + 75*B*a^2*b*d*e^2))/(56*e^4) + (b^5*x^6
*(5*A*b*e + 30*B*a*e + 7*B*b*d))/(30*e^2) + (b*x^2*(252*B*a^5*e^5 + 7*B*b^
5*d^5 + 630*A*a^4*b*e^5 + 5*A*b^5*d^4*e + 30*A*a*b^4*d^3*e^2 + 280*A*a^3*b
^2*d*e^4 + 105*A*a^2*b^3*d^2*e^3 + 75*B*a^2*b^3*d^3*e^2 + 140*B*a^3*b^2*d^
2*e^3 + 30*B*a*b^4*d^4*e + 210*B*a^4*b*d*e^4))/(420*e^6) + (b^2*x^3*(210*B
*a^4*e^4 + 7*B*b^4*d^4 + 280*A*a^3*b*e^4 + 5*A*b^4*d^3*e + 30*A*a*b^3*d^2*
e^2 + 105*A*a^2*b^2*d*e^3 + 75*B*a^2*b^2*d^2*e^2 + 30*B*a*b^3*d^3*e + 140*
B*a^3*b*d*e^3))/(126*e^5) + (b^4*x^5*(75*B*a^2*e^2 + 7*B*b^2*d^2 + 30*A*a*
b*e^2 + 5*A*b^2*d*e + 30*B*a*b*d*e))/(35*e^3) + (B*b^6*x^7)/(5*e))/(d^12 +
e^12*x^12 + 12*d*e^11*x^11 + 66*d^10*e^2*x^2 + 220*d^9*e^3*x^3 + 495*d^8*
e^4*x^4 + 792*d^7*e^5*x^5 + 924*d^6*e^6*x^6 + 792*d^5*e^7*x^7 + 495*d^4...

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 618, normalized size of antiderivative = 2.12

$$\int \frac{(a + bx)^6 (A + Bx)}{(d + ex)^{13}} dx$$

$$= \frac{-792b^7 e^7 x^7 - 4620a b^6 e^7 x^6 - 924b^7 d e^6 x^6 - 11880a^2 b^5 e^7 x^5 - 3960a b^6 d e^6 x^5 - 792b^7 d^2 e^5 x^5 - 17325a^3 b^6 d e^4 x^4 - 11880a^2 b^5 d e^4 x^4 - 4620a b^6 d^2 e^3 x^3 - 792b^7 d^3 e^3 x^3 - 17325a^3 b^6 d^2 e^2 x^2 - 11880a^2 b^5 d^2 e^2 x^2 - 4620a b^6 d^3 e^2 x^2 - 792b^7 d^4 e^2 x^2 - 17325a^3 b^6 d^3 e x - 11880a^2 b^5 d^3 e x - 4620a b^6 d^4 e x - 792b^7 d^5 e x - 17325a^3 b^6 d^4 - 11880a^2 b^5 d^4 - 4620a b^6 d^5 - 792b^7 d^6 - 17325a^3 b^6 d^6 - 11880a^2 b^5 d^6 - 4620a b^6 d^7 - 792b^7 d^8 - 17325a^3 b^6 d^8 - 11880a^2 b^5 d^8 - 4620a b^6 d^9 - 792b^7 d^{10} - 17325a^3 b^6 d^{10} - 11880a^2 b^5 d^{10} - 4620a b^6 d^{11} - 792b^7 d^{12} - 17325a^3 b^6 d^{12} - 11880a^2 b^5 d^{12} - 4620a b^6 d^{13} - 792b^7 d^{14} - 17325a^3 b^6 d^{14} - 11880a^2 b^5 d^{14} - 4620a b^6 d^{15} - 792b^7 d^{16} - 17325a^3 b^6 d^{16} - 11880a^2 b^5 d^{16} - 4620a b^6 d^{17} - 792b^7 d^{18} - 17325a^3 b^6 d^{18} - 11880a^2 b^5 d^{18} - 4620a b^6 d^{19} - 792b^7 d^{20} - 17325a^3 b^6 d^{20} - 11880a^2 b^5 d^{20} - 4620a b^6 d^{21} - 792b^7 d^{22} - 17325a^3 b^6 d^{22} - 11880a^2 b^5 d^{22} - 4620a b^6 d^{23} - 792b^7 d^{24} - 17325a^3 b^6 d^{24} - 11880a^2 b^5 d^{24} - 4620a b^6 d^{25} - 792b^7 d^{26} - 17325a^3 b^6 d^{26} - 11880a^2 b^5 d^{26} - 4620a b^6 d^{27} - 792b^7 d^{28} - 17325a^3 b^6 d^{28} - 11880a^2 b^5 d^{28} - 4620a b^6 d^{29} - 792b^7 d^{30} - 17325a^3 b^6 d^{30} - 11880a^2 b^5 d^{30} - 4620a b^6 d^{31} - 792b^7 d^{32} - 17325a^3 b^6 d^{32} - 11880a^2 b^5 d^{32} - 4620a b^6 d^{33} - 792b^7 d^{34} - 17325a^3 b^6 d^{34} - 11880a^2 b^5 d^{34} - 4620a b^6 d^{35} - 792b^7 d^{36} - 17325a^3 b^6 d^{36} - 11880a^2 b^5 d^{36} - 4620a b^6 d^{37} - 792b^7 d^{38} - 17325a^3 b^6 d^{38} - 11880a^2 b^5 d^{38} - 4620a b^6 d^{39} - 792b^7 d^{40} - 17325a^3 b^6 d^{40} - 11880a^2 b^5 d^{40} - 4620a b^6 d^{41} - 792b^7 d^{42} - 17325a^3 b^6 d^{42} - 11880a^2 b^5 d^{42} - 4620a b^6 d^{43} - 792b^7 d^{44} - 17325a^3 b^6 d^{44} - 11880a^2 b^5 d^{44} - 4620a b^6 d^{45} - 792b^7 d^{46} - 17325a^3 b^6 d^{46} - 11880a^2 b^5 d^{46} - 4620a b^6 d^{47} - 792b^7 d^{48} - 17325a^3 b^6 d^{48} - 11880a^2 b^5 d^{48} - 4620a b^6 d^{49} - 792b^7 d^{50} - 17325a^3 b^6 d^{50} - 11880a^2 b^5 d^{50} - 4620a b^6 d^{51} - 792b^7 d^{52} - 17325a^3 b^6 d^{52} - 11880a^2 b^5 d^{52} - 4620a b^6 d^{53} - 792b^7 d^{54} - 17325a^3 b^6 d^{54} - 11880a^2 b^5 d^{54} - 4620a b^6 d^{55} - 792b^7 d^{56} - 17325a^3 b^6 d^{56} - 11880a^2 b^5 d^{56} - 4620a b^6 d^{57} - 792b^7 d^{58} - 17325a^3 b^6 d^{58} - 11880a^2 b^5 d^{58} - 4620a b^6 d^{59} - 792b^7 d^{60} - 17325a^3 b^6 d^{60} - 11880a^2 b^5 d^{60} - 4620a b^6 d^{61} - 792b^7 d^{62} - 17325a^3 b^6 d^{62} - 11880a^2 b^5 d^{62} - 4620a b^6 d^{63} - 792b^7 d^{64} - 17325a^3 b^6 d^{64} - 11880a^2 b^5 d^{64} - 4620a b^6 d^{65} - 792b^7 d^{66} - 17325a^3 b^6 d^{66} - 11880a^2 b^5 d^{66} - 4620a b^6 d^{67} - 792b^7 d^{68} - 17325a^3 b^6 d^{68} - 11880a^2 b^5 d^{68} - 4620a b^6 d^{69} - 792b^7 d^{70} - 17325a^3 b^6 d^{70} - 11880a^2 b^5 d^{70} - 4620a b^6 d^{71} - 792b^7 d^{72} - 17325a^3 b^6 d^{72} - 11880a^2 b^5 d^{72} - 4620a b^6 d^{73} - 792b^7 d^{74} - 17325a^3 b^6 d^{74} - 11880a^2 b^5 d^{74} - 4620a b^6 d^{75} - 792b^7 d^{76} - 17325a^3 b^6 d^{76} - 11880a^2 b^5 d^{76} - 4620a b^6 d^{77} - 792b^7 d^{78} - 17325a^3 b^6 d^{78} - 11880a^2 b^5 d^{78} - 4620a b^6 d^{79} - 792b^7 d^{80} - 17325a^3 b^6 d^{80} - 11880a^2 b^5 d^{80} - 4620a b^6 d^{81} - 792b^7 d^{82} - 17325a^3 b^6 d^{82} - 11880a^2 b^5 d^{82} - 4620a b^6 d^{83} - 792b^7 d^{84} - 17325a^3 b^6 d^{84} - 11880a^2 b^5 d^{84} - 4620a b^6 d^{85} - 792b^7 d^{86} - 17325a^3 b^6 d^{86} - 11880a^2 b^5 d^{86} - 4620a b^6 d^{87} - 792b^7 d^{88} - 17325a^3 b^6 d^{88} - 11880a^2 b^5 d^{88} - 4620a b^6 d^{89} - 792b^7 d^{90} - 17325a^3 b^6 d^{90} - 11880a^2 b^5 d^{90} - 4620a b^6 d^{91} - 792b^7 d^{92} - 17325a^3 b^6 d^{92} - 11880a^2 b^5 d^{92} - 4620a b^6 d^{93} - 792b^7 d^{94} - 17325a^3 b^6 d^{94} - 11880a^2 b^5 d^{94} - 4620a b^6 d^{95} - 792b^7 d^{96} - 17325a^3 b^6 d^{96} - 11880a^2 b^5 d^{96} - 4620a b^6 d^{97} - 792b^7 d^{98} - 17325a^3 b^6 d^{98} - 11880a^2 b^5 d^{98} - 4620a b^6 d^{99} - 792b^7 d^{100} - 17325a^3 b^6 d^{100} - 11880a^2 b^5 d^{100} - 4620a b^6 d^{101} - 792b^7 d^{102} - 17325a^3 b^6 d^{102} - 11880a^2 b^5 d^{102} - 4620a b^6 d^{103} - 792b^7 d^{104} - 17325a^3 b^6 d^{104} - 11880a^2 b^5 d^{104} - 4620a b^6 d^{105} - 792b^7 d^{106} - 17325a^3 b^6 d^{106} - 11880a^2 b^5 d^{106} - 4620a b^6 d^{107} - 792b^7 d^{108} - 17325a^3 b^6 d^{108} - 11880a^2 b^5 d^{108} - 4620a b^6 d^{109} - 792b^7 d^{110} - 17325a^3 b^6 d^{110} - 11880a^2 b^5 d^{110} - 4620a b^6 d^{111} - 792b^7 d^{112} - 17325a^3 b^6 d^{112} - 11880a^2 b^5 d^{112} - 4620a b^6 d^{113} - 792b^7 d^{114} - 17325a^3 b^6 d^{114} - 11880a^2 b^5 d^{114} - 4620a b^6 d^{115} - 792b^7 d^{116} - 17325a^3 b^6 d^{116} - 11880a^2 b^5 d^{116} - 4620a b^6 d^{117} - 792b^7 d^{118} - 17325a^3 b^6 d^{118} - 11880a^2 b^5 d^{118} - 4620a b^6 d^{119} - 792b^7 d^{120} - 17325a^3 b^6 d^{120} - 11880a^2 b^5 d^{120} - 4620a b^6 d^{121} - 792b^7 d^{122} - 17325a^3 b^6 d^{122} - 11880a^2 b^5 d^{122} - 4620a b^6 d^{123} - 792b^7 d^{124} - 17325a^3 b^6 d^{124} - 11880a^2 b^5 d^{124} - 4620a b^6 d^{125} - 792b^7 d^{126} - 17325a^3 b^6 d^{126} - 11880a^2 b^5 d^{126} - 4620a b^6 d^{127} - 792b^7 d^{128} - 17325a^3 b^6 d^{128} - 11880a^2 b^5 d^{128} - 4620a b^6 d^{129} - 792b^7 d^{130} - 17325a^3 b^6 d^{130} - 11880a^2 b^5 d^{130} - 4620a b^6 d^{131} - 792b^7 d^{132} - 17325a^3 b^6 d^{132} - 11880a^2 b^5 d^{132} - 4620a b^6 d^{133} - 792b^7 d^{134} - 17325a^3 b^6 d^{134} - 11880a^2 b^5 d^{134} - 4620a b^6 d^{135} - 792b^7 d^{136} - 17325a^3 b^6 d^{136} - 11880a^2 b^5 d^{136} - 4620a b^6 d^{137} - 792b^7 d^{138} - 17325a^3 b^6 d^{138} - 11880a^2 b^5 d^{138} - 4620a b^6 d^{139} - 792b^7 d^{140} - 17325a^3 b^6 d^{140} - 11880a^2 b^5 d^{140} - 4620a b^6 d^{141} - 792b^7 d^{142} - 17325a^3 b^6 d^{142} - 11880a^2 b^5 d^{142} - 4620a b^6 d^{143} - 792b^7 d^{144} - 17325a^3 b^6 d^{144} - 11880a^2 b^5 d^{144} - 4620a b^6 d^{145} - 792b^7 d^{146} - 17325a^3 b^6 d^{146} - 11880a^2 b^5 d^{146} - 4620a b^6 d^{147} - 792b^7 d^{148} - 17325a^3 b^6 d^{148} - 11880a^2 b^5 d^{148} - 4620a b^6 d^{149} - 792b^7 d^{150} - 17325a^3 b^6 d^{150} - 11880a^2 b^5 d^{150} - 4620a b^6 d^{151} - 792b^7 d^{152} - 17325a^3 b^6 d^{152} - 11880a^2 b^5 d^{152} - 4620a b^6 d^{153} - 792b^7 d^{154} - 17325a^3 b^6 d^{154} - 11880a^2 b^5 d^{154} - 4620a b^6 d^{155} - 792b^7 d^{156} - 17325a^3 b^6 d^{156} - 11880a^2 b^5 d^{156} - 4620a b^6 d^{157} - 792b^7 d^{158} - 17325a^3 b^6 d^{158} - 11880a^2 b^5 d^{158} - 4620a b^6 d^{159} - 792b^7 d^{160} - 17325a^3 b^6 d^{160} - 11880a^2 b^5 d^{160} - 4620a b^6 d^{161} - 792b^7 d^{162} - 17325a^3 b^6 d^{162} - 11880a^2 b^5 d^{162} - 4620a b^6 d^{163} - 792b^7 d^{164} - 17325a^3 b^6 d^{164} - 11880a^2 b^5 d^{164} - 4620a b^6 d^{165} - 792b^7 d^{166} - 17325a^3 b^6 d^{166} - 11880a^2 b^5 d^{166} - 4620a b^6 d^{167} - 792b^7 d^{168} - 17325a^3 b^6 d^{168} - 11880a^2 b^5 d^{168} - 4620a b^6 d^{169} - 792b^7 d^{170} - 17325a^3 b^6 d^{170} - 11880a^2 b^5 d^{170} - 4620a b^6 d^{171} - 792b^7 d^{172} - 17325a^3 b^6 d^{172} - 11880a^2 b^5 d^{172} - 4620a b^6 d^{173} - 792b^7 d^{174} - 17325a^3 b^6 d^{174} - 11880a^2 b^5 d^{174} - 4620a b^6 d^{175} - 792b^7 d^{176} - 17325a^3 b^6 d^{176} - 11880a^2 b^5 d^{176} - 4620a b^6 d^{177} - 792b^7 d^{178} - 17325a^3 b^6 d^{178} - 11880a^2 b^5 d^{178} - 4620a b^6 d^{179} - 792b^7 d^{180} - 17325a^3 b^6 d^{180} - 11880a^2 b^5 d^{180} - 4620a b^6 d^{181} - 792b^7 d^{182} - 17325a^3 b^6 d^{182} - 11880a^2 b^5 d^{182} - 4620a b^6 d^{183} - 792b^7 d^{184} - 17325a^3 b^6 d^{184} - 11880a^2 b^5 d^{184} - 4620a b^6 d^{185} - 792b^7 d^{186} - 17325a^3 b^6 d^{186} - 11880a^2 b^5 d^{186} - 4620a b^6 d^{187} - 792b^7 d^{188} - 17325a^3 b^6 d^{188} - 11880a^2 b^5 d^{188} - 4620a b^6 d^{189} - 792b^7 d^{190} - 17325a^3 b^6 d^{190} - 11880a^2 b^5 d^{190} - 4620a b^6 d^{191} - 792b^7 d^{192} - 17325a^3 b^6 d^{192} - 11880a^2 b^5 d^{192} - 4620a b^6 d^{193} - 792b^7 d^{194} - 17325a^3 b^6 d^{194} - 11880a^2 b^5 d^{194} - 4620a b^6 d^{195} - 792b^7 d^{196} - 17325a^3 b^6 d^{196} - 11880a^2 b^5 d^{196} - 4620a b^6 d^{197} - 792b^7 d^{198} - 17325a^3 b^6 d^{198} - 11880a^2 b^5 d^{198} - 4620a b^6 d^{199} - 792b^7 d^{200} - 17325a^3 b^6 d^{200} - 11880a^2 b^5 d^{200} - 4620a b^6 d^{201} - 792b^7 d^{202} - 17325a^3 b^6 d^{202} - 11880a^2 b^5 d^{202} - 4620a b^6 d^{203} - 792b^7 d^{204} - 17325a^3 b^6 d^{204} - 11880a^2 b^5 d^{204} - 4620a b^6 d^{205} - 792b^7 d^{206} - 17325a^3 b^6 d^{206} - 11880a^2 b^5 d^{206} - 4620a b^6 d^{207} - 792b^7 d^{208} - 17325a^3 b^6 d^{208} - 11880a^2 b^5 d^{208} - 4620a b^6 d^{209} - 792b^7 d^{210} - 17325a^3 b^6 d^{210} - 11880a^2 b^5 d^{210} - 4620a b^6 d^{211} - 792b^7 d^{212} - 17325a^3 b^6 d^{212} - 11880a^2 b^5 d^{212} - 4620a b^6 d^{213} - 792b^7 d^{214} - 17325a^3 b^6 d^{214} - 11880a^2 b^5 d^{214} - 4620a b^6 d^{215} - 792b^7 d^{216} - 17325a^3 b^6 d^{216} - 11880a^2 b^5 d^{216} - 4620a b^6 d^{217} - 792b^7 d^{218} - 17325a^3 b^6 d^{218} - 11880a^2 b^5 d^{218} - 4620a b^6 d^{219} - 792b^7 d^{220} - 17325a^3 b^6 d^{220} - 11880a^2 b^5 d^{220} - 4620a b^6 d^{221} - 792b^7 d^{222} - 17325a^3 b^6 d^{222} - 11880a^2 b^5 d^{222} - 4620a b^6 d^{223} - 792b^7 d^{224} - 17325a^3 b^6 d^{224} - 11880a^2 b^5 d^{224} - 4620a b^6 d^{225} - 792b^7 d^{226} - 17325a^3 b^6 d^{226} - 11880a^2 b^5 d^{226} - 4620a b^6 d^{227} - 792b^7 d^{228} - 17325a^3 b^6 d^{228} - 11880a^2 b^5 d^{228} - 4620a b^6 d^{229} - 792b^7 d^{230} - 17325a^3 b^6 d^{230} - 11880a^2 b^5 d^{230} - 4620a b^6 d^{231} - 792b^7 d^{232} - 17325a^3 b^6 d^{232} - 11880a^2 b^5 d^{232} - 4620a b^6 d^{233} - 792b^7 d^{234} - 17325a^3 b^6 d^{234} - 11880a^2 b^5 d^{234} - 4620a b^6 d^{235} - 792b^7 d^{236} - 17325a^3 b^6 d^{236} - 11880a^2 b^5 d^{236} - 4620a b^6 d^{237} - 792b^7 d^{238} - 17325a^3 b^6 d^{238} - 11880a^2 b^5 d^{238} - 4620a b^6 d^{239} - 792b^7 d^{240} - 17325a^3 b^6 d^{240} - 11880a^2 b^5 d^{240} - 4620a b^6 d^{241} - 792b^7 d^{242} - 17325a^3 b^6 d^{242} - 11880a^2 b^5 d^{242} - 4620a b^6 d^{243} - 792b^7 d^{244} - 17325a^3 b^6 d^{244} - 11880a^2 b^5 d^{244} - 4620a b^6 d^{245} - 792b^7 d^{246} - 17325a^3 b^6 d^{246} - 11880a^2 b^5 d^{246} - 4620a b^6 d^{247} - 792b^7 d^{248} - 17325a^3 b^6 d^{248} - 11880a^2 b^5 d^{248} - 4620a b^6 d^{249} - 792b^7 d^{250} - 17325a^3 b^6 d^{250} - 11880a^2 b^5 d^{250} - 4620a b^6 d^{251} - 792b^7 d^{252} - 17325a^3 b^6 d^{252} - 11880a^2 b^5 d^{252} - 4620a b^6 d^{253} - 792b^7 d^{254} - 17325a^3 b^6 d^{254} - 11880a^2 b^5 d^{254} - 4620a b^6 d^{255} - 792b^7 d^{256} - 17325a^3 b^6 d^{256} - 11880a^2 b^5 d^{256} - 4620a b^6 d^{257} - 792b^7 d^{258} - 17325a^3 b^6 d^{258} - 11880a^2 b^5 d^{258} - 4620a b^6 d^{259} - 792b^7 d^{260} - 17325a^3 b^6 d^{260} - 11880a^2 b^5 d^{260} - 4620a b^6 d^{261} - 792b^7 d^{262} - 17325a^3 b^6 d^{262} - 11880a^2 b^5 d^{262} - 4620a b^6 d^{263} - 792b^7 d^{264} - 17325a^3 b^6 d^{264} - 11880a^2 b^5 d^{264} - 4620a b^6 d^{265} - 792b^7 d^{266} - 17325a^3 b^6 d^{266} - 11880a^2 b^5 d^{266} - 4620a b^6 d^{267} - 792b^7 d^{268} - 17325a^3 b^6 d^{268} - 11880a^2 b^5 d^{268} - 4620a b^6 d^{269} - 792b^7 d^{270} - 17325a^3 b^6 d^{270} - 11880a^2 b^5 d^{270} - 4620a b^6 d^{271} - 792b^7 d^{272} - 17325a^3 b^6 d^{272} - 11880a^2 b^5 d^{272} - 4620a b^6 d^{273} - 792b^7 d^{274} - 17325a^3 b^6 d^{274} - 11880a^2 b^5 d^{274} - 4620a b^6 d^{275} - 792b^7 d^{276} - 17325a^3 b^6 d^{276} - 11880a^2 b^5 d^{276} - 4620a b^6 d^{277} - 792b^7 d^{278} - 17325a^3 b^6 d^{278} - 11880a^2 b^5 d^{278} - 4620a b^6 d^{279} - 792b^7 d^{280} - 17325a^3 b^6 d^{280} - 11880a^2 b^5 d^{280} - 4620a b^6 d^{281} - 792b^7 d^{282} - 17325a^3 b^6 d^{282} - 11880a^2 b^5 d^{282} - 4620a b^6 d^{283} - 792b^7 d^{284} - 17325a^3 b^6 d^{284} - 11880a^2 b^5 d^{284} - 4620a b^6 d^{285} - 792b^7 d^{286} - 17325a^3 b^6 d^{286} - 11880a^2 b^5 d^{286} - 4620a b^6 d^{287} - 792b^7 d^{288} - 17325a^3 b^6 d^{288} - 11880a^2 b^5 d^{288} - 4620a b^6 d^{289} - 792b^7 d^{290} - 17325a^3 b^6 d^{290} - 11880a^2 b^5 d^{290} - 4620a b^6 d^{291} - 792b^7 d^{292} - 17325a^3 b^6 d^{292} - 11880a^2 b^5 d^{292} - 4620a b^6 d^{293} - 792b^7 d^{294} - 17325a^3 b^6 d^{294} - 11880a^2 b^5 d^{294} - 4620a b^6 d^{295} - 792b^7 d^{296} - 17325a^3 b^6 d^{296} - 11880a^2 b^5 d^{296} - 4620a b^6 d^{297} - 792b^7 d^{298} - 17325a^3 b^6 d^{298} - 11880a^2 b^5 d^{298} - 4620a b^6 d^{299} - 792b^7 d^{300} - 17325a^3 b^6 d^{300} - 11880a^2 b^5 d^{300} - 4620a b^6 d^{301} - 792b^7 d^{302} - 17325a^3 b^6 d^{302} - 11880a^2 b^5 d^{302} - 4620a b^6 d^{303} - 792b^7 d^{304} - 17325a^3 b^6 d^{304} - 11880a^2 b^5 d^{304} - 4620a b^6 d^{305} - 792b^7 d^{306} - 17325a^3 b^6 d^{306} - 11880a^2 b^5 d^{306} - 4620a b^6 d^{307} - 792b^7 d^{308} - 17325a^3 b^6 d^{308} - 11880a^2 b^5 d^{308} - 4620a b^6 d^{309} - 792b^7 d^{310} - 17325a^3 b^6 d^{310} - 11880a^2 b^5 d^{310} - 4620a b^6 d^{311} - 792b^7 d^{312} - 17325a^3 b^6 d^{312} - 11880a^2 b^5 d^{312} - 4620a b^6 d^{313} - 792b^7 d^{314} - 17325a^3 b^6 d^{314} - 11880a^2 b^5 d^{314} - 4620a b^6 d^{315} - 792b^7 d^{316} - 17325a^3 b^6 d^{316} - 11880a^2 b^5 d^{316} - 4620a b^6 d^{317} - 792b^7 d^{318} - 17325a^3 b^6 d^{318} - 11880a^2 b^5 d^{318} - 4620a b^6 d^{319} - 792b^7 d^{320} - 17325a^3 b^6 d^{320} - 11880a^2 b^5 d^{320} - 4620a b^6 d^{321} - 792b^7 d^{322} - 17325a^3 b^6 d^{322} - 11880a^2 b^5 d^{322} - 4620a$$

output

```
( - 330*a**7*e**7 - 210*a**6*b*d*e**6 - 2520*a**6*b*e**7*x - 126*a**5*b**2
*d**2*e**5 - 1512*a**5*b**2*d*e**6*x - 8316*a**5*b**2*e**7*x**2 - 70*a**4*
b**3*d**3*e**4 - 840*a**4*b**3*d**2*e**5*x - 4620*a**4*b**3*d*e**6*x**2 -
15400*a**4*b**3*e**7*x**3 - 35*a**3*b**4*d**4*e**3 - 420*a**3*b**4*d**3*e*
*4*x - 2310*a**3*b**4*d**2*e**5*x**2 - 7700*a**3*b**4*d*e**6*x**3 - 17325*
a**3*b**4*e**7*x**4 - 15*a**2*b**5*d**5*e**2 - 180*a**2*b**5*d**4*e**3*x -
990*a**2*b**5*d**3*e**4*x**2 - 3300*a**2*b**5*d**2*e**5*x**3 - 7425*a**2*
b**5*d*e**6*x**4 - 11880*a**2*b**5*e**7*x**5 - 5*a*b**6*d**6*e - 60*a*b**6
*d**5*e**2*x - 330*a*b**6*d**4*e**3*x**2 - 1100*a*b**6*d**3*e**4*x**3 - 24
75*a*b**6*d**2*e**5*x**4 - 3960*a*b**6*d*e**6*x**5 - 4620*a*b**6*e**7*x**6
- b**7*d**7 - 12*b**7*d**6*e*x - 66*b**7*d**5*e**2*x**2 - 220*b**7*d**4*e
**3*x**3 - 495*b**7*d**3*e**4*x**4 - 792*b**7*d**2*e**5*x**5 - 924*b**7*d*
e**6*x**6 - 792*b**7*e**7*x**7)/(3960*e**8*(d**12 + 12*d**11*e*x + 66*d**1
0*e**2*x**2 + 220*d**9*e**3*x**3 + 495*d**8*e**4*x**4 + 792*d**7*e**5*x**5
+ 924*d**6*e**6*x**6 + 792*d**5*e**7*x**7 + 495*d**4*e**8*x**8 + 220*d**3
*e**9*x**9 + 66*d**2*e**10*x**10 + 12*d*e**11*x**11 + e**12*x**12))
```


3.63 $\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{14}} dx$

Optimal result	636
Mathematica [B] (verified)	637
Rubi [A] (verified)	637
Maple [B] (verified)	639
Fricas [B] (verification not implemented)	640
Sympy [F(-1)]	641
Maxima [B] (verification not implemented)	641
Giac [B] (verification not implemented)	642
Mupad [B] (verification not implemented)	643
Reduce [B] (verification not implemented)	644

Optimal result

Integrand size = 20, antiderivative size = 292

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{14}} dx = \frac{(bd-ae)^6(Bd-Ae)}{13e^8(d+ex)^{13}} - \frac{(bd-ae)^5(7bBd-6Abe-aBe)}{12e^8(d+ex)^{12}} + \frac{3b(bd-ae)^4(7bBd-5Abe-2aBe)}{11e^8(d+ex)^{11}} - \frac{b^2(bd-ae)^3(7bBd-4Abe-3aBe)}{2e^8(d+ex)^{10}} + \frac{5b^3(bd-ae)^2(7bBd-3Abe-4aBe)}{9e^8(d+ex)^9} - \frac{3b^4(bd-ae)(7bBd-2Abe-5aBe)}{8e^8(d+ex)^8} + \frac{b^5(7bBd-Abe-6aBe)}{7e^8(d+ex)^7} - \frac{b^6B}{6e^8(d+ex)^6}$$

output

```
1/13*(-a*e+b*d)^6*(-A*e+B*d)/e^8/(e*x+d)^13-1/12*(-a*e+b*d)^5*(-6*A*b*e-B*a*e+7*B*b*d)/e^8/(e*x+d)^12+3/11*b*(-a*e+b*d)^4*(-5*A*b*e-2*B*a*e+7*B*b*d)/e^8/(e*x+d)^11-1/2*b^2*(-a*e+b*d)^3*(-4*A*b*e-3*B*a*e+7*B*b*d)/e^8/(e*x+d)^10+5/9*b^3*(-a*e+b*d)^2*(-3*A*b*e-4*B*a*e+7*B*b*d)/e^8/(e*x+d)^9-3/8*b^4*(-a*e+b*d)*(-2*A*b*e-5*B*a*e+7*B*b*d)/e^8/(e*x+d)^8+1/7*b^5*(-A*b*e-6*B*a*e+7*B*b*d)/e^8/(e*x+d)^7-1/6*b^6*B/e^8/(e*x+d)^6
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 605 vs. $2(292) = 584$.

Time = 0.17 (sec) , antiderivative size = 605, normalized size of antiderivative = 2.07

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{14}} dx = \frac{462a^6e^6(12Ae + B(d + 13ex)) + 252a^5be^5(11Ae(d + 13ex) + 2B(d^2 + 13dex + 78e^2x^2)) + 126a^4b^2e^4}{(d + ex)^{13}}$$

input

```
Integrate[((a + b*x)^6*(A + B*x))/(d + e*x)^14,x]
```

output

```
-1/72072*(462*a^6*e^6*(12*A*e + B*(d + 13*e*x)) + 252*a^5*b*e^5*(11*A*e*(d
+ 13*e*x) + 2*B*(d^2 + 13*d*e*x + 78*e^2*x^2)) + 126*a^4*b^2*e^4*(10*A*e*
(d^2 + 13*d*e*x + 78*e^2*x^2) + 3*B*(d^3 + 13*d^2*e*x + 78*d*e^2*x^2 + 286
*e^3*x^3)) + 56*a^3*b^3*e^3*(9*A*e*(d^3 + 13*d^2*e*x + 78*d*e^2*x^2 + 286*
e^3*x^3) + 4*B*(d^4 + 13*d^3*e*x + 78*d^2*e^2*x^2 + 286*d*e^3*x^3 + 715*e^
4*x^4)) + 21*a^2*b^4*e^2*(8*A*e*(d^4 + 13*d^3*e*x + 78*d^2*e^2*x^2 + 286*d
*e^3*x^3 + 715*e^4*x^4) + 5*B*(d^5 + 13*d^4*e*x + 78*d^3*e^2*x^2 + 286*d^2
*e^3*x^3 + 715*d*e^4*x^4 + 1287*e^5*x^5)) + 6*a*b^5*e*(7*A*e*(d^5 + 13*d^4
*e*x + 78*d^3*e^2*x^2 + 286*d^2*e^3*x^3 + 715*d*e^4*x^4 + 1287*e^5*x^5) +
6*B*(d^6 + 13*d^5*e*x + 78*d^4*e^2*x^2 + 286*d^3*e^3*x^3 + 715*d^2*e^4*x^4
+ 1287*d*e^5*x^5 + 1716*e^6*x^6)) + b^6*(6*A*e*(d^6 + 13*d^5*e*x + 78*d^4
*e^2*x^2 + 286*d^3*e^3*x^3 + 715*d^2*e^4*x^4 + 1287*d*e^5*x^5 + 1716*e^6*x
^6) + 7*B*(d^7 + 13*d^6*e*x + 78*d^5*e^2*x^2 + 286*d^4*e^3*x^3 + 715*d^3*e
^4*x^4 + 1287*d^2*e^5*x^5 + 1716*d*e^6*x^6 + 1716*e^7*x^7)))/(e^8*(d + e*x
)^13)
```

Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 292, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{14}} dx$$

↓ 86

$$\int \left(\frac{b^5(6aBe + Abe - 7bBd)}{e^7(d + ex)^8} - \frac{3b^4(bd - ae)(5aBe + 2Abe - 7bBd)}{e^7(d + ex)^9} + \frac{5b^3(bd - ae)^2(4aBe + 3Abe - 7bBd)}{e^7(d + ex)^{10}} - \dots \right) dx$$

↓ 2009

$$\begin{aligned} & \frac{b^5(-6aBe - Abe + 7bBd)}{7e^8(d + ex)^7} - \frac{3b^4(bd - ae)(-5aBe - 2Abe + 7bBd)}{8e^8(d + ex)^8} + \\ & \frac{5b^3(bd - ae)^2(-4aBe - 3Abe + 7bBd)}{9e^8(d + ex)^9} - \frac{b^2(bd - ae)^3(-3aBe - 4Abe + 7bBd)}{2e^8(d + ex)^{10}} + \\ & \frac{3b(bd - ae)^4(-2aBe - 5Abe + 7bBd)}{11e^8(d + ex)^{11}} - \frac{(bd - ae)^5(-aBe - 6Abe + 7bBd)}{12e^8(d + ex)^{12}} + \\ & \frac{(bd - ae)^6(Bd - Ae)}{13e^8(d + ex)^{13}} - \frac{b^6B}{6e^8(d + ex)^6} \end{aligned}$$

input `Int[((a + b*x)^6*(A + B*x))/(d + e*x)^14,x]`

output `((b*d - a*e)^6*(B*d - A*e))/(13*e^8*(d + e*x)^13) - ((b*d - a*e)^5*(7*b*B*d - 6*A*b*e - a*B*e))/(12*e^8*(d + e*x)^12) + (3*b*(b*d - a*e)^4*(7*b*B*d - 5*A*b*e - 2*a*B*e))/(11*e^8*(d + e*x)^11) - (b^2*(b*d - a*e)^3*(7*b*B*d - 4*A*b*e - 3*a*B*e))/(2*e^8*(d + e*x)^10) + (5*b^3*(b*d - a*e)^2*(7*b*B*d - 3*A*b*e - 4*a*B*e))/(9*e^8*(d + e*x)^9) - (3*b^4*(b*d - a*e)*(7*b*B*d - 2*A*b*e - 5*a*B*e))/(8*e^8*(d + e*x)^8) + (b^5*(7*b*B*d - A*b*e - 6*a*B*e))/(7*e^8*(d + e*x)^7) - (b^6*B)/(6*e^8*(d + e*x)^6)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.))^(n_.)*((e_.) + (f_.)*(x_.))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 788 vs. $2(276) = 552$.

Time = 0.25 (sec) , antiderivative size = 789, normalized size of antiderivative = 2.70

method	result
risch	$\frac{-\frac{b^6 B x^7}{6e} - \frac{b^5(6Abe+36Bae+7Bbd)x^6}{42e^2} - \frac{b^4(42Aab e^2+6A b^2 de+105B a^2 e^2+36Babde+7b^2 B d^2)x^5}{56e^3} - \frac{5b^3(168A a^2 b e^3+42Aa b^2 d e^2+6A a^3 b^3 d^2 e)}{13e^8(e+x)^{13}}$
default	$\frac{-\frac{b^6 B x^7}{6e} - \frac{(6A b^6 e^6+36B a b^5 e^6+7b^6 B d e^5)x^6}{42e^7} - \frac{(42A a b^5 e^7+6A b^6 d e^6+105B a^2 b^4 e^7+36B a b^5 d e^6+7b^6 B d^2 e^5)x^5}{56e^8} - 5(168A a^2 b^4 e^8+105A a^3 b^3 d^2 e^3)}{13e^8(e+x)^{13}}$
norman	$\frac{-\frac{b^6 B x^7}{6e} - \frac{(6A b^6 e^6+36B a b^5 e^6+7b^6 B d e^5)x^6}{42e^7} - \frac{(42A a b^5 e^7+6A b^6 d e^6+105B a^2 b^4 e^7+36B a b^5 d e^6+7b^6 B d^2 e^5)x^5}{56e^8} - 5(168A a^2 b^4 e^8+105A a^3 b^3 d^2 e^3)}{13e^8(e+x)^{13}}$
gosper	$\frac{-12012B x^7 b^6 e^7+10296A x^6 b^6 e^7+61776B x^6 a b^5 e^7+12012B x^6 b^6 d e^6+54054A x^5 a b^5 e^7+7722A x^5 b^6 d e^6+135135B x^5 a^2 b^5 e^7}{(e+x)^{13}}$
orering	$\frac{-12012B x^7 b^6 e^7+10296A x^6 b^6 e^7+61776B x^6 a b^5 e^7+12012B x^6 b^6 d e^6+54054A x^5 a b^5 e^7+7722A x^5 b^6 d e^6+135135B x^5 a^2 b^5 e^7}{(e+x)^{13}}$
parallelrisch	$\frac{-12012B b^6 x^7 e^{12}+10296A b^6 e^{12} x^6+61776B a b^5 e^{12} x^6+12012B b^6 d e^{11} x^6+54054A a b^5 e^{12} x^5+7722A b^6 d e^{11} x^5+135135B a^2 b^5 e^{12} x^5}{(e+x)^{13}}$

```
input int((b*x+a)^6*(B*x+A)/(e*x+d)^14,x,method=_RETURNVERBOSE)
```

```
output (-1/6*b^6*B/e*x^7-1/42*b^5/e^2*(6*A*b*e+36*B*a*e+7*B*b*d)*x^6-1/56*b^4/e^3*(42*A*a*b*e^2+6*A*b^2*d*e+105*B*a^2*e^2+36*B*a*b*d*e+7*B*b^2*d^2)*x^5-5/504*b^3/e^4*(168*A*a^2*b*e^3+42*A*a*b^2*d*e^2+6*A*b^3*d^2*e+224*B*a^3*e^3+105*B*a^2*b*d*e^2+36*B*a*b^2*d^2*e+7*B*b^3*d^3)*x^4-1/252*b^2/e^5*(504*A*a^3*b*e^4+168*A*a^2*b^2*d*e^3+42*A*a*b^3*d^2*e^2+6*A*b^4*d^3*e+378*B*a^4*e^4+224*B*a^3*b*d*e^3+105*B*a^2*b^2*d^2*e^2+36*B*a*b^3*d^3*e+7*B*b^4*d^4)*x^3-1/924*b/e^6*(1260*A*a^4*b*e^5+504*A*a^3*b^2*d*e^4+168*A*a^2*b^3*d^2*e^3+42*A*a*b^4*d^3*e^2+6*A*b^5*d^4*e+504*B*a^5*e^5+378*B*a^4*b*d*e^4+224*B*a^3*b^2*d^2*e^3+105*B*a^2*b^3*d^3*e^2+36*B*a*b^4*d^4*e+7*B*b^5*d^5)*x^2-1/5544/e^7*(2772*A*a^5*b*e^6+1260*A*a^4*b^2*d*e^5+504*A*a^3*b^3*d^2*e^4+168*A*a^2*b^4*d^3*e^3+42*A*a*b^5*d^4*e^2+6*A*b^6*d^5*e+462*B*a^6*e^6+504*B*a^5*b*d*e^5+378*B*a^4*b^2*d^2*e^4+224*B*a^3*b^3*d^3*e^3+105*B*a^2*b^4*d^4*e^2+36*B*a*b^5*d^5*e+7*B*b^6*d^6)*x-1/72072/e^8*(5544*A*a^6*e^7+2772*A*a^5*b*d*e^6+1260*A*a^4*b^2*d^2*e^5+504*A*a^3*b^3*d^3*e^4+168*A*a^2*b^4*d^4*e^3+42*A*a*b^5*d^5*e^2+6*A*b^6*d^6*e+462*B*a^6*d*e^6+504*B*a^5*b*d^2*e^5+378*B*a^4*b^2*d^2*e^4+224*B*a^3*b^3*d^4*e^3+105*B*a^2*b^4*d^5*e^2+36*B*a*b^5*d^6*e+7*B*b^6*d^7))/(e*x+d)^13
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 905 vs. $2(276) = 552$.

Time = 0.13 (sec) , antiderivative size = 905, normalized size of antiderivative = 3.10

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{14}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^14,x, algorithm="fricas")`

output

```
-1/72072*(12012*B*b^6*e^7*x^7 + 7*B*b^6*d^7 + 5544*A*a^6*e^7 + 6*(6*B*a*b^5 + A*b^6)*d^6*e + 21*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 + 56*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^4*e^3 + 126*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 + 252*(2*B*a^5*b + 5*A*a^4*b^2)*d^2*e^5 + 462*(B*a^6 + 6*A*a^5*b)*d*e^6 + 1716*(7*B*b^6*d*e^6 + 6*(6*B*a*b^5 + A*b^6)*e^7)*x^6 + 1287*(7*B*b^6*d^2*e^5 + 6*(6*B*a*b^5 + A*b^6)*d*e^6 + 21*(5*B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^5 + 715*(7*B*b^6*d^3*e^4 + 6*(6*B*a*b^5 + A*b^6)*d^2*e^5 + 21*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6 + 56*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^7)*x^4 + 286*(7*B*b^6*d^4*e^3 + 6*(6*B*a*b^5 + A*b^6)*d^3*e^4 + 21*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 + 56*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^6 + 126*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^3 + 78*(7*B*b^6*d^5*e^2 + 6*(6*B*a*b^5 + A*b^6)*d^4*e^3 + 21*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*e^4 + 56*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + 126*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^6 + 252*(2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 13*(7*B*b^6*d^6*e + 6*(6*B*a*b^5 + A*b^6)*d^5*e^2 + 21*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^3 + 56*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 + 126*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^5 + 252*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^6 + 462*(B*a^6 + 6*A*a^5*b)*e^7)*x)/(e^21*x^13 + 13*d*e^20*x^12 + 78*d^2*e^19*x^11 + 286*d^3*e^18*x^10 + 715*d^4*e^17*x^9 + 1287*d^5*e^16*x^8 + 1716*d^6*e^15*x^7 + 1716*d^7*e^14*x^6 + 1287*d^8*e^13*x^5 + 715*d^9*e^12*x^4 + 286*d^10*e^11*x^3 + 78*d^11*e^10*x^2 + 13*d^12*e^9*x + d^13*e^8)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{14}} dx = \text{Timed out}$$

input `integrate((b*x+a)**6*(B*x+A)/(e*x+d)**14,x)`output `Timed out`**Maxima [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 905 vs. $2(276) = 552$.

Time = 0.07 (sec) , antiderivative size = 905, normalized size of antiderivative = 3.10

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{14}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^14,x, algorithm="maxima")`

output

```

-1/72072*(12012*B*b^6*e^7*x^7 + 7*B*b^6*d^7 + 5544*A*a^6*e^7 + 6*(6*B*a*b^
5 + A*b^6)*d^6*e + 21*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 + 56*(4*B*a^3*b^3
+ 3*A*a^2*b^4)*d^4*e^3 + 126*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 + 252*(2*
B*a^5*b + 5*A*a^4*b^2)*d^2*e^5 + 462*(B*a^6 + 6*A*a^5*b)*d*e^6 + 1716*(7*B
*b^6*d*e^6 + 6*(6*B*a*b^5 + A*b^6)*e^7)*x^6 + 1287*(7*B*b^6*d^2*e^5 + 6*(6
*B*a*b^5 + A*b^6)*d*e^6 + 21*(5*B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^5 + 715*(7*B
*b^6*d^3*e^4 + 6*(6*B*a*b^5 + A*b^6)*d^2*e^5 + 21*(5*B*a^2*b^4 + 2*A*a*b^5
)*d*e^6 + 56*(4*B*a^3*b^3 + 3*A*a^2*b^4)*e^7)*x^4 + 286*(7*B*b^6*d^4*e^3 +
6*(6*B*a*b^5 + A*b^6)*d^3*e^4 + 21*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 + 56
*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d*e^6 + 126*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*
x^3 + 78*(7*B*b^6*d^5*e^2 + 6*(6*B*a*b^5 + A*b^6)*d^4*e^3 + 21*(5*B*a^2*b^
4 + 2*A*a*b^5)*d^3*e^4 + 56*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + 126*(3*B
*a^4*b^2 + 4*A*a^3*b^3)*d*e^6 + 252*(2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 1
3*(7*B*b^6*d^6*e + 6*(6*B*a*b^5 + A*b^6)*d^5*e^2 + 21*(5*B*a^2*b^4 + 2*A*a
*b^5)*d^4*e^3 + 56*(4*B*a^3*b^3 + 3*A*a^2*b^4)*d^3*e^4 + 126*(3*B*a^4*b^2
+ 4*A*a^3*b^3)*d^2*e^5 + 252*(2*B*a^5*b + 5*A*a^4*b^2)*d*e^6 + 462*(B*a^6
+ 6*A*a^5*b)*e^7)*x)/(e^21*x^13 + 13*d*e^20*x^12 + 78*d^2*e^19*x^11 + 286*
d^3*e^18*x^10 + 715*d^4*e^17*x^9 + 1287*d^5*e^16*x^8 + 1716*d^6*e^15*x^7 +
1716*d^7*e^14*x^6 + 1287*d^8*e^13*x^5 + 715*d^9*e^12*x^4 + 286*d^10*e^11*
x^3 + 78*d^11*e^10*x^2 + 13*d^12*e^9*x + d^13*e^8)

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 912 vs. $2(276) = 552$.

Time = 0.13 (sec) , antiderivative size = 912, normalized size of antiderivative = 3.12

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{14}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^6*(B*x+A)/(e*x+d)^14,x, algorithm="giac")
```

output

```
-1/72072*(12012*B*b^6*e^7*x^7 + 12012*B*b^6*d*e^6*x^6 + 61776*B*a*b^5*e^7*
x^6 + 10296*A*b^6*e^7*x^6 + 9009*B*b^6*d^2*e^5*x^5 + 46332*B*a*b^5*d*e^6*x
^5 + 7722*A*b^6*d*e^6*x^5 + 135135*B*a^2*b^4*e^7*x^5 + 54054*A*a*b^5*e^7*x
^5 + 5005*B*b^6*d^3*e^4*x^4 + 25740*B*a*b^5*d^2*e^5*x^4 + 4290*A*b^6*d^2*e
^5*x^4 + 75075*B*a^2*b^4*d*e^6*x^4 + 30030*A*a*b^5*d*e^6*x^4 + 160160*B*a^
3*b^3*e^7*x^4 + 120120*A*a^2*b^4*e^7*x^4 + 2002*B*b^6*d^4*e^3*x^3 + 10296*
B*a*b^5*d^3*e^4*x^3 + 1716*A*b^6*d^3*e^4*x^3 + 30030*B*a^2*b^4*d^2*e^5*x^3
+ 12012*A*a*b^5*d^2*e^5*x^3 + 64064*B*a^3*b^3*d*e^6*x^3 + 48048*A*a^2*b^4
*d*e^6*x^3 + 108108*B*a^4*b^2*e^7*x^3 + 144144*A*a^3*b^3*e^7*x^3 + 546*B*b
^6*d^5*e^2*x^2 + 2808*B*a*b^5*d^4*e^3*x^2 + 468*A*b^6*d^4*e^3*x^2 + 8190*B
*a^2*b^4*d^3*e^4*x^2 + 3276*A*a*b^5*d^3*e^4*x^2 + 17472*B*a^3*b^3*d^2*e^5*
x^2 + 13104*A*a^2*b^4*d^2*e^5*x^2 + 29484*B*a^4*b^2*d*e^6*x^2 + 39312*A*a^
3*b^3*d*e^6*x^2 + 39312*B*a^5*b*e^7*x^2 + 98280*A*a^4*b^2*e^7*x^2 + 91*B*b
^6*d^6*e*x + 468*B*a*b^5*d^5*e^2*x + 78*A*b^6*d^5*e^2*x + 1365*B*a^2*b^4*d
^4*e^3*x + 546*A*a*b^5*d^4*e^3*x + 2912*B*a^3*b^3*d^3*e^4*x + 2184*A*a^2*b
^4*d^3*e^4*x + 4914*B*a^4*b^2*d^2*e^5*x + 6552*A*a^3*b^3*d^2*e^5*x + 6552*
B*a^5*b*d*e^6*x + 16380*A*a^4*b^2*d*e^6*x + 6006*B*a^6*e^7*x + 36036*A*a^5
*b*e^7*x + 7*B*b^6*d^7 + 36*B*a*b^5*d^6*e + 6*A*b^6*d^6*e + 105*B*a^2*b^4*
d^5*e^2 + 42*A*a*b^5*d^5*e^2 + 224*B*a^3*b^3*d^4*e^3 + 168*A*a^2*b^4*d^4*
e^3 + 378*B*a^4*b^2*d^3*e^4 + 504*A*a^3*b^3*d^3*e^4 + 504*B*a^5*b*d^2*e^...
```

Mupad [B] (verification not implemented)

Time = 1.41 (sec) , antiderivative size = 921, normalized size of antiderivative = 3.15

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{14}} dx =$$

$$\frac{462 B a^6 d e^6 + 5544 A a^6 e^7 + 504 B a^5 b d^2 e^5 + 2772 A a^5 b d e^6 + 378 B a^4 b^2 d^3 e^4 + 1260 A a^4 b^2 d^2 e^5 + 224 B a^3 b^3 d^4 e^3 + 504 A a^3 b^3 d^3 e^4 + 105 B a^2 b^4 d^5 e^2 + 42 A a^2 b^4 d^4 e^3 + 36 B a b^5 d^6 e + 6 A b^6 d^6 e + 7 B b^6 d^7}{72072 e^8}$$

input

```
int(((A + B*x)*(a + b*x)^6)/(d + e*x)^14,x)
```


output

```

-((5544*A*a^6*e^7 + 7*B*b^6*d^7 + 6*A*b^6*d^6*e + 462*B*a^6*d*e^6 + 42*A*a
*b^5*d^5*e^2 + 504*B*a^5*b*d^2*e^5 + 168*A*a^2*b^4*d^4*e^3 + 504*A*a^3*b^3
*d^3*e^4 + 1260*A*a^4*b^2*d^2*e^5 + 105*B*a^2*b^4*d^5*e^2 + 224*B*a^3*b^3
*d^4*e^3 + 378*B*a^4*b^2*d^3*e^4 + 2772*A*a^5*b*d*e^6 + 36*B*a*b^5*d^6*e)/(
72072*e^8) + (x*(462*B*a^6*e^6 + 7*B*b^6*d^6 + 2772*A*a^5*b*e^6 + 6*A*b^6*
d^5*e + 42*A*a*b^5*d^4*e^2 + 1260*A*a^4*b^2*d*e^5 + 168*A*a^2*b^4*d^3*e^3
+ 504*A*a^3*b^3*d^2*e^4 + 105*B*a^2*b^4*d^4*e^2 + 224*B*a^3*b^3*d^3*e^3 +
378*B*a^4*b^2*d^2*e^4 + 36*B*a*b^5*d^5*e + 504*B*a^5*b*d*e^5))/(5544*e^7)
+ (5*b^3*x^4*(224*B*a^3*e^3 + 7*B*b^3*d^3 + 168*A*a^2*b*e^3 + 6*A*b^3*d^2*
e + 42*A*a*b^2*d*e^2 + 36*B*a*b^2*d^2*e + 105*B*a^2*b*d*e^2))/(504*e^4) +
(b^5*x^6*(6*A*b*e + 36*B*a*e + 7*B*b*d))/(42*e^2) + (b*x^2*(504*B*a^5*e^5
+ 7*B*b^5*d^5 + 1260*A*a^4*b*e^5 + 6*A*b^5*d^4*e + 42*A*a*b^4*d^3*e^2 + 50
4*A*a^3*b^2*d*e^4 + 168*A*a^2*b^3*d^2*e^3 + 105*B*a^2*b^3*d^3*e^2 + 224*B*
a^3*b^2*d^2*e^3 + 36*B*a*b^4*d^4*e + 378*B*a^4*b*d*e^4))/(924*e^6) + (b^2*
x^3*(378*B*a^4*e^4 + 7*B*b^4*d^4 + 504*A*a^3*b*e^4 + 6*A*b^4*d^3*e + 42*A*
a*b^3*d^2*e^2 + 168*A*a^2*b^2*d*e^3 + 105*B*a^2*b^2*d^2*e^2 + 36*B*a*b^3*d
^3*e + 224*B*a^3*b*d*e^3))/(252*e^5) + (b^4*x^5*(105*B*a^2*e^2 + 7*B*b^2*d
^2 + 42*A*a*b*e^2 + 6*A*b^2*d*e + 36*B*a*b*d*e))/(56*e^3) + (B*b^6*x^7)/(6
*e))/(d^13 + e^13*x^13 + 13*d*e^12*x^12 + 78*d^11*e^2*x^2 + 286*d^10*e^3*x
^3 + 715*d^9*e^4*x^4 + 1287*d^8*e^5*x^5 + 1716*d^7*e^6*x^6 + 1716*d^6*e...

```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 629, normalized size of antiderivative = 2.15

$$\int \frac{(a + bx)^6 (A + Bx)}{(d + ex)^{14}} dx$$

$$= \frac{-1716b^7 e^7 x^7 - 10296a b^6 e^7 x^6 - 1716b^7 d e^6 x^6 - 27027a^2 b^5 e^7 x^5 - 7722a b^6 d e^6 x^5 - 1287b^7 d^2 e^5 x^5 - 4004$$

input

```
int((b*x+a)^6*(B*x+A)/(e*x+d)^14,x)
```

output

```
( - 792*a**7*e**7 - 462*a**6*b*d*e**6 - 6006*a**6*b*e**7*x - 252*a**5*b**2
*d**2*e**5 - 3276*a**5*b**2*d*e**6*x - 19656*a**5*b**2*e**7*x**2 - 126*a**
4*b**3*d**3*e**4 - 1638*a**4*b**3*d**2*e**5*x - 9828*a**4*b**3*d*e**6*x**2
- 36036*a**4*b**3*e**7*x**3 - 56*a**3*b**4*d**4*e**3 - 728*a**3*b**4*d**3
*e**4*x - 4368*a**3*b**4*d**2*e**5*x**2 - 16016*a**3*b**4*d*e**6*x**3 - 40
040*a**3*b**4*e**7*x**4 - 21*a**2*b**5*d**5*e**2 - 273*a**2*b**5*d**4*e**3
*x - 1638*a**2*b**5*d**3*e**4*x**2 - 6006*a**2*b**5*d**2*e**5*x**3 - 15015
*a**2*b**5*d*e**6*x**4 - 27027*a**2*b**5*e**7*x**5 - 6*a*b**6*d**6*e - 78*
a*b**6*d**5*e**2*x - 468*a*b**6*d**4*e**3*x**2 - 1716*a*b**6*d**3*e**4*x**
3 - 4290*a*b**6*d**2*e**5*x**4 - 7722*a*b**6*d*e**6*x**5 - 10296*a*b**6*e
**7*x**6 - b**7*d**7 - 13*b**7*d**6*e*x - 78*b**7*d**5*e**2*x**2 - 286*b**7
*d**4*e**3*x**3 - 715*b**7*d**3*e**4*x**4 - 1287*b**7*d**2*e**5*x**5 - 171
6*b**7*d*e**6*x**6 - 1716*b**7*e**7*x**7)/(10296*e**8*(d**13 + 13*d**12*e*
x + 78*d**11*e**2*x**2 + 286*d**10*e**3*x**3 + 715*d**9*e**4*x**4 + 1287*d
**8*e**5*x**5 + 1716*d**7*e**6*x**6 + 1716*d**6*e**7*x**7 + 1287*d**5*e**8
*x**8 + 715*d**4*e**9*x**9 + 286*d**3*e**10*x**10 + 78*d**2*e**11*x**11 +
13*d*e**12*x**12 + e**13*x**13))
```

3.64 $\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{15}} dx$

Optimal result	646
Mathematica [B] (verified)	647
Rubi [A] (verified)	647
Maple [B] (verified)	649
Fricas [B] (verification not implemented)	650
Sympy [F(-1)]	651
Maxima [B] (verification not implemented)	651
Giac [B] (verification not implemented)	652
Mupad [B] (verification not implemented)	653
Reduce [B] (verification not implemented)	654

Optimal result

Integrand size = 20, antiderivative size = 292

$$\int \frac{(a+bx)^6(A+Bx)}{(d+ex)^{15}} dx = \frac{(bd-ae)^6(Bd-Ae)}{14e^8(d+ex)^{14}} - \frac{(bd-ae)^5(7bBd-6Abe-aBe)}{13e^8(d+ex)^{13}} + \frac{b(bd-ae)^4(7bBd-5Abe-2aBe)}{4e^8(d+ex)^{12}} - \frac{5b^2(bd-ae)^3(7bBd-4Abe-3aBe)}{11e^8(d+ex)^{11}} + \frac{b^3(bd-ae)^2(7bBd-3Abe-4aBe)}{2e^8(d+ex)^{10}} - \frac{b^4(bd-ae)(7bBd-2Abe-5aBe)}{3e^8(d+ex)^9} + \frac{b^5(7bBd-Abe-6aBe)}{8e^8(d+ex)^8} - \frac{b^6B}{7e^8(d+ex)^7}$$

output

```
1/14*(-a*e+b*d)^6*(-A*e+B*d)/e^8/(e*x+d)^14-1/13*(-a*e+b*d)^5*(-6*A*b*e-B*
a*e+7*B*b*d)/e^8/(e*x+d)^13+1/4*b*(-a*e+b*d)^4*(-5*A*b*e-2*B*a*e+7*B*b*d)/
e^8/(e*x+d)^12-5/11*b^2*(-a*e+b*d)^3*(-4*A*b*e-3*B*a*e+7*B*b*d)/e^8/(e*x+d
)^11+1/2*b^3*(-a*e+b*d)^2*(-3*A*b*e-4*B*a*e+7*B*b*d)/e^8/(e*x+d)^10-1/3*b^
4*(-a*e+b*d)*(-2*A*b*e-5*B*a*e+7*B*b*d)/e^8/(e*x+d)^9+1/8*b^5*(-A*b*e-6*B*
a*e+7*B*b*d)/e^8/(e*x+d)^8-1/7*b^6*B/e^8/(e*x+d)^7
```


$$\int \frac{(a + bx)^6 (A + Bx)}{(d + ex)^{15}} dx$$

↓ 86

$$\int \left(\frac{b^5(6aBe + Abe - 7bBd)}{e^7(d + ex)^9} - \frac{3b^4(bd - ae)(5aBe + 2Abe - 7bBd)}{e^7(d + ex)^{10}} + \frac{5b^3(bd - ae)^2(4aBe + 3Abe - 7bBd)}{e^7(d + ex)^{11}} - \dots \right) dx$$

↓ 2009

$$\begin{aligned} & \frac{b^5(-6aBe - Abe + 7bBd)}{8e^8(d + ex)^8} - \frac{b^4(bd - ae)(-5aBe - 2Abe + 7bBd)}{3e^8(d + ex)^9} + \\ & \frac{b^3(bd - ae)^2(-4aBe - 3Abe + 7bBd)}{2e^8(d + ex)^{10}} - \frac{5b^2(bd - ae)^3(-3aBe - 4Abe + 7bBd)}{11e^8(d + ex)^{11}} + \\ & \frac{b(bd - ae)^4(-2aBe - 5Abe + 7bBd)}{4e^8(d + ex)^{12}} - \frac{(bd - ae)^5(-aBe - 6Abe + 7bBd)}{13e^8(d + ex)^{13}} + \\ & \frac{(bd - ae)^6(Bd - Ae)}{14e^8(d + ex)^{14}} - \frac{b^6B}{7e^8(d + ex)^7} \end{aligned}$$

input `Int[((a + b*x)^6*(A + B*x))/(d + e*x)^15,x]`

output `((b*d - a*e)^6*(B*d - A*e))/(14*e^8*(d + e*x)^14) - ((b*d - a*e)^5*(7*b*B*d - 6*A*b*e - a*B*e))/(13*e^8*(d + e*x)^13) + (b*(b*d - a*e)^4*(7*b*B*d - 5*A*b*e - 2*a*B*e))/(4*e^8*(d + e*x)^12) - (5*b^2*(b*d - a*e)^3*(7*b*B*d - 4*A*b*e - 3*a*B*e))/(11*e^8*(d + e*x)^11) + (b^3*(b*d - a*e)^2*(7*b*B*d - 3*A*b*e - 4*a*B*e))/(2*e^8*(d + e*x)^10) - (b^4*(b*d - a*e)*(7*b*B*d - 2*A*b*e - 5*a*B*e))/(3*e^8*(d + e*x)^9) + (b^5*(7*b*B*d - A*b*e - 6*a*B*e))/(8*e^8*(d + e*x)^8) - (b^6*B)/(7*e^8*(d + e*x)^7)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.))^(n_.)*((e_.) + (f_.)*(x_.))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 902 vs. $2(276) = 552$.

Time = 0.12 (sec) , antiderivative size = 902, normalized size of antiderivative = 3.09

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{15}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^15,x, algorithm="fricas")`

output

```
-1/24024*(3432*B*b^6*e^7*x^7 + B*b^6*d^7 + 1716*A*a^6*e^7 + (6*B*a*b^5 + A
*b^6)*d^6*e + 4*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 + 12*(4*B*a^3*b^3 + 3*A*
a^2*b^4)*d^4*e^3 + 30*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 + 66*(2*B*a^5*b
+ 5*A*a^4*b^2)*d^2*e^5 + 132*(B*a^6 + 6*A*a^5*b)*d*e^6 + 3003*(B*b^6*d*e^6
+ (6*B*a*b^5 + A*b^6)*e^7)*x^6 + 2002*(B*b^6*d^2*e^5 + (6*B*a*b^5 + A*b^6
)*d*e^6 + 4*(5*B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^5 + 1001*(B*b^6*d^3*e^4 + (6*
B*a*b^5 + A*b^6)*d^2*e^5 + 4*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6 + 12*(4*B*a^3
*b^3 + 3*A*a^2*b^4)*e^7)*x^4 + 364*(B*b^6*d^4*e^3 + (6*B*a*b^5 + A*b^6)*d^
3*e^4 + 4*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 + 12*(4*B*a^3*b^3 + 3*A*a^2*b^
4)*d*e^6 + 30*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^3 + 91*(B*b^6*d^5*e^2 + (
6*B*a*b^5 + A*b^6)*d^4*e^3 + 4*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*e^4 + 12*(4*B
*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + 30*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^6 + 6
6*(2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 14*(B*b^6*d^6*e + (6*B*a*b^5 + A*b^
6)*d^5*e^2 + 4*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^3 + 12*(4*B*a^3*b^3 + 3*A*a
^2*b^4)*d^3*e^4 + 30*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^5 + 66*(2*B*a^5*b +
5*A*a^4*b^2)*d*e^6 + 132*(B*a^6 + 6*A*a^5*b)*e^7)*x)/(e^22*x^14 + 14*d*e^
21*x^13 + 91*d^2*e^20*x^12 + 364*d^3*e^19*x^11 + 1001*d^4*e^18*x^10 + 2002
*d^5*e^17*x^9 + 3003*d^6*e^16*x^8 + 3432*d^7*e^15*x^7 + 3003*d^8*e^14*x^6
+ 2002*d^9*e^13*x^5 + 1001*d^10*e^12*x^4 + 364*d^11*e^11*x^3 + 91*d^12*e^1
0*x^2 + 14*d^13*e^9*x + d^14*e^8)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{15}} dx = \text{Timed out}$$

input `integrate((b*x+a)**6*(B*x+A)/(e*x+d)**15,x)`output `Timed out`**Maxima [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 902 vs. $2(276) = 552$.

Time = 0.09 (sec) , antiderivative size = 902, normalized size of antiderivative = 3.09

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{15}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^6*(B*x+A)/(e*x+d)^15,x, algorithm="maxima")`

output

```

-1/24024*(3432*B*b^6*e^7*x^7 + B*b^6*d^7 + 1716*A*a^6*e^7 + (6*B*a*b^5 + A
*b^6)*d^6*e + 4*(5*B*a^2*b^4 + 2*A*a*b^5)*d^5*e^2 + 12*(4*B*a^3*b^3 + 3*A*
a^2*b^4)*d^4*e^3 + 30*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^3*e^4 + 66*(2*B*a^5*b
+ 5*A*a^4*b^2)*d^2*e^5 + 132*(B*a^6 + 6*A*a^5*b)*d*e^6 + 3003*(B*b^6*d*e^6
+ (6*B*a*b^5 + A*b^6)*e^7)*x^6 + 2002*(B*b^6*d^2*e^5 + (6*B*a*b^5 + A*b^6
)*d*e^6 + 4*(5*B*a^2*b^4 + 2*A*a*b^5)*e^7)*x^5 + 1001*(B*b^6*d^3*e^4 + (6*
B*a*b^5 + A*b^6)*d^2*e^5 + 4*(5*B*a^2*b^4 + 2*A*a*b^5)*d*e^6 + 12*(4*B*a^3
*b^3 + 3*A*a^2*b^4)*e^7)*x^4 + 364*(B*b^6*d^4*e^3 + (6*B*a*b^5 + A*b^6)*d^
3*e^4 + 4*(5*B*a^2*b^4 + 2*A*a*b^5)*d^2*e^5 + 12*(4*B*a^3*b^3 + 3*A*a^2*b^
4)*d*e^6 + 30*(3*B*a^4*b^2 + 4*A*a^3*b^3)*e^7)*x^3 + 91*(B*b^6*d^5*e^2 + (
6*B*a*b^5 + A*b^6)*d^4*e^3 + 4*(5*B*a^2*b^4 + 2*A*a*b^5)*d^3*e^4 + 12*(4*B
*a^3*b^3 + 3*A*a^2*b^4)*d^2*e^5 + 30*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d*e^6 + 6
6*(2*B*a^5*b + 5*A*a^4*b^2)*e^7)*x^2 + 14*(B*b^6*d^6*e + (6*B*a*b^5 + A*b^
6)*d^5*e^2 + 4*(5*B*a^2*b^4 + 2*A*a*b^5)*d^4*e^3 + 12*(4*B*a^3*b^3 + 3*A*a
^2*b^4)*d^3*e^4 + 30*(3*B*a^4*b^2 + 4*A*a^3*b^3)*d^2*e^5 + 66*(2*B*a^5*b +
5*A*a^4*b^2)*d*e^6 + 132*(B*a^6 + 6*A*a^5*b)*e^7)*x)/(e^22*x^14 + 14*d*e^
21*x^13 + 91*d^2*e^20*x^12 + 364*d^3*e^19*x^11 + 1001*d^4*e^18*x^10 + 2002
*d^5*e^17*x^9 + 3003*d^6*e^16*x^8 + 3432*d^7*e^15*x^7 + 3003*d^8*e^14*x^6
+ 2002*d^9*e^13*x^5 + 1001*d^10*e^12*x^4 + 364*d^11*e^11*x^3 + 91*d^12*e^1
0*x^2 + 14*d^13*e^9*x + d^14*e^8)

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 910 vs. $2(276) = 552$.

Time = 0.12 (sec) , antiderivative size = 910, normalized size of antiderivative = 3.12

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{15}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^6*(B*x+A)/(e*x+d)^15,x, algorithm="giac")
```

output

```

-1/24024*(3432*B*b^6*e^7*x^7 + 3003*B*b^6*d*e^6*x^6 + 18018*B*a*b^5*e^7*x^
6 + 3003*A*b^6*e^7*x^6 + 2002*B*b^6*d^2*e^5*x^5 + 12012*B*a*b^5*d*e^6*x^5
+ 2002*A*b^6*d*e^6*x^5 + 40040*B*a^2*b^4*e^7*x^5 + 16016*A*a*b^5*e^7*x^5 +
1001*B*b^6*d^3*e^4*x^4 + 6006*B*a*b^5*d^2*e^5*x^4 + 1001*A*b^6*d^2*e^5*x^
4 + 20020*B*a^2*b^4*d*e^6*x^4 + 8008*A*a*b^5*d*e^6*x^4 + 48048*B*a^3*b^3*e
^7*x^4 + 36036*A*a^2*b^4*e^7*x^4 + 364*B*b^6*d^4*e^3*x^3 + 2184*B*a*b^5*d^
3*e^4*x^3 + 364*A*b^6*d^3*e^4*x^3 + 7280*B*a^2*b^4*d^2*e^5*x^3 + 2912*A*a*
b^5*d^2*e^5*x^3 + 17472*B*a^3*b^3*d*e^6*x^3 + 13104*A*a^2*b^4*d*e^6*x^3 +
32760*B*a^4*b^2*e^7*x^3 + 43680*A*a^3*b^3*e^7*x^3 + 91*B*b^6*d^5*e^2*x^2 +
546*B*a*b^5*d^4*e^3*x^2 + 91*A*b^6*d^4*e^3*x^2 + 1820*B*a^2*b^4*d^3*e^4*x
^2 + 728*A*a*b^5*d^3*e^4*x^2 + 4368*B*a^3*b^3*d^2*e^5*x^2 + 3276*A*a^2*b^4
*d^2*e^5*x^2 + 8190*B*a^4*b^2*d*e^6*x^2 + 10920*A*a^3*b^3*d*e^6*x^2 + 1201
2*B*a^5*b*e^7*x^2 + 30030*A*a^4*b^2*e^7*x^2 + 14*B*b^6*d^6*e*x + 84*B*a*b^
5*d^5*e^2*x + 14*A*b^6*d^5*e^2*x + 280*B*a^2*b^4*d^4*e^3*x + 112*A*a*b^5*d
^4*e^3*x + 672*B*a^3*b^3*d^3*e^4*x + 504*A*a^2*b^4*d^3*e^4*x + 1260*B*a^4*
b^2*d^2*e^5*x + 1680*A*a^3*b^3*d^2*e^5*x + 1848*B*a^5*b*d*e^6*x + 4620*A*a
^4*b^2*d*e^6*x + 1848*B*a^6*e^7*x + 11088*A*a^5*b*e^7*x + B*b^6*d^7 + 6*B*
a*b^5*d^6*e + A*b^6*d^6*e + 20*B*a^2*b^4*d^5*e^2 + 8*A*a*b^5*d^5*e^2 + 48*
B*a^3*b^3*d^4*e^3 + 36*A*a^2*b^4*d^4*e^3 + 90*B*a^4*b^2*d^3*e^4 + 120*A*a^
3*b^3*d^3*e^4 + 132*B*a^5*b*d^2*e^5 + 330*A*a^4*b^2*d^2*e^5 + 132*B*a^6...

```

Mupad [B] (verification not implemented)

Time = 1.57 (sec) , antiderivative size = 918, normalized size of antiderivative = 3.14

$$\int \frac{(a + bx)^6 (A + Bx)}{(d + ex)^{15}} dx =$$

$$\frac{132 B a^6 d e^6 + 1716 A a^6 e^7 + 132 B a^5 b d^2 e^5 + 792 A a^5 b d e^6 + 90 B a^4 b^2 d^3 e^4 + 330 A a^4 b^2 d^2 e^5 + 48 B a^3 b^3 d^4 e^3 + 120 A a^3 b^3 d^3 e^4 + 20 B a^2 b^4 d^5 e^2 + 8 A a^2 b^4 d^4 e^3 + 90 B a^4 b^2 d^3 e^4 + 120 A a^3 b^3 d^3 e^4 + 132 B a^5 b d^2 e^5 + 330 A a^4 b^2 d^2 e^5 + 132 B a^6 d^7 + 6 B a b^5 d^6 e + A b^6 d^6 e + 20 B a^2 b^4 d^5 e^2 + 8 A a b^5 d^5 e^2 + 48 B a^3 b^3 d^4 e^3 + 36 A a^2 b^4 d^4 e^3 + 90 B a^4 b^2 d^3 e^4 + 120 A a^3 b^3 d^3 e^4 + 132 B a^5 b d^2 e^5 + 330 A a^4 b^2 d^2 e^5 + 132 B a^6 d^7 + 6 B a b^5 d^6 e + A b^6 d^6 e}{24024 e^8}$$

input

```
int(((A + B*x)*(a + b*x)^6)/(d + e*x)^15,x)
```

output

```

-((1716*A*a^6*e^7 + B*b^6*d^7 + A*b^6*d^6*e + 132*B*a^6*d*e^6 + 8*A*a*b^5*
d^5*e^2 + 132*B*a^5*b*d^2*e^5 + 36*A*a^2*b^4*d^4*e^3 + 120*A*a^3*b^3*d^3*e
^4 + 330*A*a^4*b^2*d^2*e^5 + 20*B*a^2*b^4*d^5*e^2 + 48*B*a^3*b^3*d^4*e^3 +
90*B*a^4*b^2*d^3*e^4 + 792*A*a^5*b*d*e^6 + 6*B*a*b^5*d^6*e)/(24024*e^8) +
(x*(132*B*a^6*e^6 + B*b^6*d^6 + 792*A*a^5*b*e^6 + A*b^6*d^5*e + 8*A*a*b^5
*d^4*e^2 + 330*A*a^4*b^2*d*e^5 + 36*A*a^2*b^4*d^3*e^3 + 120*A*a^3*b^3*d^2*
e^4 + 20*B*a^2*b^4*d^4*e^2 + 48*B*a^3*b^3*d^3*e^3 + 90*B*a^4*b^2*d^2*e^4 +
6*B*a*b^5*d^5*e + 132*B*a^5*b*d*e^5))/(1716*e^7) + (b^3*x^4*(48*B*a^3*e^3
+ B*b^3*d^3 + 36*A*a^2*b*e^3 + A*b^3*d^2*e + 8*A*a*b^2*d*e^2 + 6*B*a*b^2*
d^2*e + 20*B*a^2*b*d*e^2))/(24*e^4) + (b^5*x^6*(A*b*e + 6*B*a*e + B*b*d))/
(8*e^2) + (b*x^2*(132*B*a^5*e^5 + B*b^5*d^5 + 330*A*a^4*b*e^5 + A*b^5*d^4*
e + 8*A*a*b^4*d^3*e^2 + 120*A*a^3*b^2*d*e^4 + 36*A*a^2*b^3*d^2*e^3 + 20*B*
a^2*b^3*d^3*e^2 + 48*B*a^3*b^2*d^2*e^3 + 6*B*a*b^4*d^4*e + 90*B*a^4*b*d*e^
4))/(264*e^6) + (b^2*x^3*(90*B*a^4*e^4 + B*b^4*d^4 + 120*A*a^3*b*e^4 + A*b
^4*d^3*e + 8*A*a*b^3*d^2*e^2 + 36*A*a^2*b^2*d*e^3 + 20*B*a^2*b^2*d^2*e^2 +
6*B*a*b^3*d^3*e + 48*B*a^3*b*d*e^3))/(66*e^5) + (b^4*x^5*(20*B*a^2*e^2 +
B*b^2*d^2 + 8*A*a*b*e^2 + A*b^2*d*e + 6*B*a*b*d*e))/(12*e^3) + (B*b^6*x^7)
/(7*e))/(d^14 + e^14*x^14 + 14*d*e^13*x^13 + 91*d^12*e^2*x^2 + 364*d^11*e^
3*x^3 + 1001*d^10*e^4*x^4 + 2002*d^9*e^5*x^5 + 3003*d^8*e^6*x^6 + 3432*d^7
*e^7*x^7 + 3003*d^6*e^8*x^8 + 2002*d^5*e^9*x^9 + 1001*d^4*e^10*x^10 + 3...

```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 640, normalized size of antiderivative = 2.19

$$\int \frac{(a + bx)^6(A + Bx)}{(d + ex)^{15}} dx$$

$$= \frac{-3432b^7e^7x^7 - 21021ab^6e^7x^6 - 3003b^7de^6x^6 - 56056a^2b^5e^7x^5 - 14014ab^6de^6x^5 - 2002b^7d^2e^5x^5 - 840$$

input

```
int((b*x+a)^6*(B*x+A)/(e*x+d)^15,x)
```

output

```
( - 1716*a**7*e**7 - 924*a**6*b*d*e**6 - 12936*a**6*b*e**7*x - 462*a**5*b*
*2*d**2*e**5 - 6468*a**5*b**2*d*e**6*x - 42042*a**5*b**2*e**7*x**2 - 210*a
**4*b**3*d**3*e**4 - 2940*a**4*b**3*d**2*e**5*x - 19110*a**4*b**3*d*e**6*x
**2 - 76440*a**4*b**3*e**7*x**3 - 84*a**3*b**4*d**4*e**3 - 1176*a**3*b**4*
d**3*e**4*x - 7644*a**3*b**4*d**2*e**5*x**2 - 30576*a**3*b**4*d*e**6*x**3
- 84084*a**3*b**4*e**7*x**4 - 28*a**2*b**5*d**5*e**2 - 392*a**2*b**5*d**4*
e**3*x - 2548*a**2*b**5*d**3*e**4*x**2 - 10192*a**2*b**5*d**2*e**5*x**3 -
28028*a**2*b**5*d*e**6*x**4 - 56056*a**2*b**5*e**7*x**5 - 7*a*b**6*d**6*e
- 98*a*b**6*d**5*e**2*x - 637*a*b**6*d**4*e**3*x**2 - 2548*a*b**6*d**3*e**
4*x**3 - 7007*a*b**6*d**2*e**5*x**4 - 14014*a*b**6*d*e**6*x**5 - 21021*a*b
**6*e**7*x**6 - b**7*d**7 - 14*b**7*d**6*e*x - 91*b**7*d**5*e**2*x**2 - 36
4*b**7*d**4*e**3*x**3 - 1001*b**7*d**3*e**4*x**4 - 2002*b**7*d**2*e**5*x**
5 - 3003*b**7*d*e**6*x**6 - 3432*b**7*e**7*x**7)/(24024*e**8*(d**14 + 14*d
**13*e*x + 91*d**12*e**2*x**2 + 364*d**11*e**3*x**3 + 1001*d**10*e**4*x**4
+ 2002*d**9*e**5*x**5 + 3003*d**8*e**6*x**6 + 3432*d**7*e**7*x**7 + 3003*
d**6*e**8*x**8 + 2002*d**5*e**9*x**9 + 1001*d**4*e**10*x**10 + 364*d**3*e
**11*x**11 + 91*d**2*e**12*x**12 + 14*d*e**13*x**13 + e**14*x**14))
```

3.65 $\int (a + bx)^{10}(A + Bx)(d + ex)^{13} dx$

Optimal result	656
Mathematica [B] (verified)	657
Rubi [A] (verified)	658
Maple [B] (verified)	660
Fricas [B] (verification not implemented)	661
Sympy [B] (verification not implemented)	662
Maxima [B] (verification not implemented)	663
Giac [B] (verification not implemented)	664
Mupad [B] (verification not implemented)	665
Reduce [B] (verification not implemented)	665

Optimal result

Integrand size = 20, antiderivative size = 464

$$\begin{aligned}
 & \int (a + bx)^{10}(A + Bx)(d + ex)^{13} dx \\
 &= -\frac{(bd - ae)^{10}(Bd - Ae)(d + ex)^{14}}{14e^{12}} + \frac{(bd - ae)^9(11bBd - 10Abe - aBe)(d + ex)^{15}}{15e^{12}} \\
 & \quad - \frac{5b(bd - ae)^8(11bBd - 9Abe - 2aBe)(d + ex)^{16}}{16e^{12}} \\
 & \quad + \frac{15b^2(bd - ae)^7(11bBd - 8Abe - 3aBe)(d + ex)^{17}}{17e^{12}} \\
 & \quad - \frac{5b^3(bd - ae)^6(11bBd - 7Abe - 4aBe)(d + ex)^{18}}{3e^{12}} \\
 & \quad + \frac{42b^4(bd - ae)^5(11bBd - 6Abe - 5aBe)(d + ex)^{19}}{19e^{12}} \\
 & \quad - \frac{21b^5(bd - ae)^4(11bBd - 5Abe - 6aBe)(d + ex)^{20}}{10e^{12}} \\
 & \quad + \frac{10b^6(bd - ae)^3(11bBd - 4Abe - 7aBe)(d + ex)^{21}}{7e^{12}} \\
 & \quad - \frac{15b^7(bd - ae)^2(11bBd - 3Abe - 8aBe)(d + ex)^{22}}{22e^{12}} \\
 & \quad + \frac{5b^8(bd - ae)(11bBd - 2Abe - 9aBe)(d + ex)^{23}}{23e^{12}} \\
 & \quad - \frac{b^9(11bBd - Abe - 10aBe)(d + ex)^{24}}{24e^{12}} + \frac{b^{10}B(d + ex)^{25}}{25e^{12}}
 \end{aligned}$$

output

```

-1/14*(-a*e+b*d)^10*(-A*e+B*d)*(e*x+d)^14/e^12+1/15*(-a*e+b*d)^9*(-10*A*b*
e-B*a*e+11*B*b*d)*(e*x+d)^15/e^12-5/16*b*(-a*e+b*d)^8*(-9*A*b*e-2*B*a*e+11
*B*b*d)*(e*x+d)^16/e^12+15/17*b^2*(-a*e+b*d)^7*(-8*A*b*e-3*B*a*e+11*B*b*d)
*(e*x+d)^17/e^12-5/3*b^3*(-a*e+b*d)^6*(-7*A*b*e-4*B*a*e+11*B*b*d)*(e*x+d)^
18/e^12+42/19*b^4*(-a*e+b*d)^5*(-6*A*b*e-5*B*a*e+11*B*b*d)*(e*x+d)^19/e^12
-21/10*b^5*(-a*e+b*d)^4*(-5*A*b*e-6*B*a*e+11*B*b*d)*(e*x+d)^20/e^12+10/7*b
^6*(-a*e+b*d)^3*(-4*A*b*e-7*B*a*e+11*B*b*d)*(e*x+d)^21/e^12-15/22*b^7*(-a*
e+b*d)^2*(-3*A*b*e-8*B*a*e+11*B*b*d)*(e*x+d)^22/e^12+5/23*b^8*(-a*e+b*d)*(
-2*A*b*e-9*B*a*e+11*B*b*d)*(e*x+d)^23/e^12-1/24*b^9*(-A*b*e-10*B*a*e+11*B*
b*d)*(e*x+d)^24/e^12+1/25*b^10*B*(e*x+d)^25/e^12

```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 3532 vs. $2(464) = 928$.

Time = 0.96 (sec) , antiderivative size = 3532, normalized size of antiderivative = 7.61

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{13} dx = \text{Result too large to show}$$

input

```
Integrate[(a + b*x)^10*(A + B*x)*(d + e*x)^13,x]
```

output

```
a^10*A*d^13*x + (a^9*d^12*(10*A*b*d + a*B*d + 13*a*A*e)*x^2)/2 + (a^8*d^11
*(a*B*d*(10*b*d + 13*a*e) + A*(45*b^2*d^2 + 130*a*b*d*e + 78*a^2*e^2))*x^3
)/3 + (a^7*d^10*(a*B*d*(45*b^2*d^2 + 130*a*b*d*e + 78*a^2*e^2) + A*(120*b^
3*d^3 + 585*a*b^2*d^2*e + 780*a^2*b*d*e^2 + 286*a^3*e^3))*x^4)/4 + (a^6*d^
9*(a*B*d*(120*b^3*d^3 + 585*a*b^2*d^2*e + 780*a^2*b*d*e^2 + 286*a^3*e^3) +
5*A*(42*b^4*d^4 + 312*a*b^3*d^3*e + 702*a^2*b^2*d^2*e^2 + 572*a^3*b*d*e^3
+ 143*a^4*e^4))*x^5)/5 + (a^5*d^8*(5*a*B*d*(42*b^4*d^4 + 312*a*b^3*d^3*e
+ 702*a^2*b^2*d^2*e^2 + 572*a^3*b*d*e^3 + 143*a^4*e^4) + A*(252*b^5*d^5 +
2730*a*b^4*d^4*e + 9360*a^2*b^3*d^3*e^2 + 12870*a^3*b^2*d^2*e^3 + 7150*a^4
*b*d*e^4 + 1287*a^5*e^5))*x^6)/6 + (a^4*d^7*(a*B*d*(252*b^5*d^5 + 2730*a*b
^4*d^4*e + 9360*a^2*b^3*d^3*e^2 + 12870*a^3*b^2*d^2*e^3 + 7150*a^4*b*d*e^4
+ 1287*a^5*e^5) + 3*A*(70*b^6*d^6 + 1092*a*b^5*d^5*e + 5460*a^2*b^4*d^4*e
^2 + 11440*a^3*b^3*d^3*e^3 + 10725*a^4*b^2*d^2*e^4 + 4290*a^5*b*d*e^5 + 57
2*a^6*e^6))*x^7)/7 + (3*a^3*d^6*(a*B*d*(70*b^6*d^6 + 1092*a*b^5*d^5*e + 54
60*a^2*b^4*d^4*e^2 + 11440*a^3*b^3*d^3*e^3 + 10725*a^4*b^2*d^2*e^4 + 4290*
a^5*b*d*e^5 + 572*a^6*e^6) + A*(40*b^7*d^7 + 910*a*b^6*d^6*e + 6552*a^2*b^
5*d^5*e^2 + 20020*a^3*b^4*d^4*e^3 + 28600*a^4*b^3*d^3*e^4 + 19305*a^5*b^2*
d^2*e^5 + 5720*a^6*b*d*e^6 + 572*a^7*e^7))*x^8)/8 + (a^2*d^5*(a*B*d*(40*b^
7*d^7 + 910*a*b^6*d^6*e + 6552*a^2*b^5*d^5*e^2 + 20020*a^3*b^4*d^4*e^3 + 2
8600*a^4*b^3*d^3*e^4 + 19305*a^5*b^2*d^2*e^5 + 5720*a^6*b*d*e^6 + 572*a...
```

Rubi [A] (verified)

Time = 5.23 (sec) , antiderivative size = 464, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{13} dx$$

↓ 86

$$\int \left(\frac{b^9(d + ex)^{23}(10aBe + Abe - 11bBd)}{e^{11}} - \frac{5b^8(d + ex)^{22}(bd - ae)(9aBe + 2Abe - 11bBd)}{e^{11}} + \frac{15b^7(d + ex)^{21}(t}{e^{11}} \right) dx$$

↓ 2009

$$\begin{aligned}
& - \frac{b^9(d+ex)^{24}(-10aBe - Abe + 11bBd)}{24e^{12}} + \frac{5b^8(d+ex)^{23}(bd-ae)(-9aBe - 2Abe + 11bBd)}{23e^{12}} \\
& - \frac{15b^7(d+ex)^{22}(bd-ae)^2(-8aBe - 3Abe + 11bBd)}{15e^{12}} + \frac{10b^6(d+ex)^{21}(bd-ae)^3(-7aBe - 4Abe + 11bBd)}{22e^{12}} \\
& - \frac{21b^5(d+ex)^{20}(bd-ae)^4(-6aBe - 5Abe + 11bBd)}{7e^{12}} + \frac{42b^4(d+ex)^{19}(bd-ae)^5(-5aBe - 6Abe + 11bBd)}{10e^{12}} \\
& - \frac{5b^3(d+ex)^{18}(bd-ae)^6(-4aBe - 7Abe + 11bBd)}{19e^{12}} + \frac{15b^2(d+ex)^{17}(bd-ae)^7(-3aBe - 8Abe + 11bBd)}{3e^{12}} \\
& - \frac{5b(d+ex)^{16}(bd-ae)^8(-2aBe - 9Abe + 11bBd)}{17e^{12}} + \frac{(d+ex)^{15}(bd-ae)^9(-aBe - 10Abe + 11bBd)}{15e^{12}} \\
& - \frac{(d+ex)^{14}(bd-ae)^{10}(Bd - Ae)}{14e^{12}} + \frac{b^{10}B(d+ex)^{25}}{25e^{12}}
\end{aligned}$$

input `Int[(a + b*x)^10*(A + B*x)*(d + e*x)^13,x]`

output `-1/14*((b*d - a*e)^10*(B*d - A*e)*(d + e*x)^14)/e^12 + ((b*d - a*e)^9*(11*b*B*d - 10*A*b*e - a*B*e)*(d + e*x)^15)/(15*e^12) - (5*b*(b*d - a*e)^8*(11*b*B*d - 9*A*b*e - 2*a*B*e)*(d + e*x)^16)/(16*e^12) + (15*b^2*(b*d - a*e)^7*(11*b*B*d - 8*A*b*e - 3*a*B*e)*(d + e*x)^17)/(17*e^12) - (5*b^3*(b*d - a*e)^6*(11*b*B*d - 7*A*b*e - 4*a*B*e)*(d + e*x)^18)/(3*e^12) + (42*b^4*(b*d - a*e)^5*(11*b*B*d - 6*A*b*e - 5*a*B*e)*(d + e*x)^19)/(19*e^12) - (21*b^5*(b*d - a*e)^4*(11*b*B*d - 5*A*b*e - 6*a*B*e)*(d + e*x)^20)/(10*e^12) + (10*b^6*(b*d - a*e)^3*(11*b*B*d - 4*A*b*e - 7*a*B*e)*(d + e*x)^21)/(7*e^12) - (15*b^7*(b*d - a*e)^2*(11*b*B*d - 3*A*b*e - 8*a*B*e)*(d + e*x)^22)/(22*e^12) + (5*b^8*(b*d - a*e)*(11*b*B*d - 2*A*b*e - 9*a*B*e)*(d + e*x)^23)/(23*e^12) - (b^9*(11*b*B*d - A*b*e - 10*a*B*e)*(d + e*x)^24)/(24*e^12) + (b^10*B*(d + e*x)^25)/(25*e^12)`

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0]) || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3892 vs. $2(440) = 880$.

Time = 0.33 (sec) , antiderivative size = 3893, normalized size of antiderivative = 8.39

method	result	size
default	Expression too large to display	3893
norman	Expression too large to display	4237
gospers	Expression too large to display	5040
risch	Expression too large to display	5040
parallelrisch	Expression too large to display	5040
orering	Expression too large to display	5040

input

```
int((b*x+a)^10*(B*x+A)*(e*x+d)^13,x,method=_RETURNVERBOSE)
```

output

```

1/23*((10*A*a*b^9+45*B*a^2*b^8)*e^13+13*(A*b^10+10*B*a*b^9)*d*e^12+78*b^10
*B*d^2*e^11)*x^23+1/19*((252*A*a^5*b^5+210*B*a^6*b^4)*e^13+13*(210*A*a^4*b
^6+252*B*a^5*b^5)*d*e^12+78*(120*A*a^3*b^7+210*B*a^4*b^6)*d^2*e^11+286*(45
*A*a^2*b^8+120*B*a^3*b^7)*d^3*e^10+715*(10*A*a*b^9+45*B*a^2*b^8)*d^4*e^9+1
287*(A*b^10+10*B*a*b^9)*d^5*e^8+1716*b^10*B*d^6*e^7)*x^19+1/18*((210*A*a^6
*b^4+120*B*a^7*b^3)*e^13+13*(252*A*a^5*b^5+210*B*a^6*b^4)*d*e^12+78*(210*A
*a^4*b^6+252*B*a^5*b^5)*d^2*e^11+286*(120*A*a^3*b^7+210*B*a^4*b^6)*d^3*e^1
0+715*(45*A*a^2*b^8+120*B*a^3*b^7)*d^4*e^9+1287*(10*A*a*b^9+45*B*a^2*b^8)*
d^5*e^8+1716*(A*b^10+10*B*a*b^9)*d^6*e^7+1716*b^10*B*d^7*e^6)*x^18+1/9*(12
87*a^10*A*d^5*e^8+1716*(10*A*a^9*b+B*a^10)*d^6*e^7+1716*(45*A*a^8*b^2+10*B
*a^9*b)*d^7*e^6+1287*(120*A*a^7*b^3+45*B*a^8*b^2)*d^8*e^5+715*(210*A*a^6*b
^4+120*B*a^7*b^3)*d^9*e^4+286*(252*A*a^5*b^5+210*B*a^6*b^4)*d^10*e^3+78*(2
10*A*a^4*b^6+252*B*a^5*b^5)*d^11*e^2+13*(120*A*a^3*b^7+210*B*a^4*b^6)*d^12
*e+(45*A*a^2*b^8+120*B*a^3*b^7)*d^13)*x^9+1/13*(13*a^10*A*d*e^12+78*(10*A*
a^9*b+B*a^10)*d^2*e^11+286*(45*A*a^8*b^2+10*B*a^9*b)*d^3*e^10+715*(120*A*a
^7*b^3+45*B*a^8*b^2)*d^4*e^9+1287*(210*A*a^6*b^4+120*B*a^7*b^3)*d^5*e^8+17
16*(252*A*a^5*b^5+210*B*a^6*b^4)*d^6*e^7+1716*(210*A*a^4*b^6+252*B*a^5*b^5
)*d^7*e^6+1287*(120*A*a^3*b^7+210*B*a^4*b^6)*d^8*e^5+715*(45*A*a^2*b^8+120
*B*a^3*b^7)*d^9*e^4+286*(10*A*a*b^9+45*B*a^2*b^8)*d^10*e^3+78*(A*b^10+10*B
*a*b^9)*d^11*e^2+13*b^10*B*d^12*e)*x^13+1/15*((10*A*a^9*b+B*a^10)*e^13+...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3905 vs. $2(440) = 880$.

Time = 0.13 (sec) , antiderivative size = 3905, normalized size of antiderivative = 8.42

$$\int (a + bx)^{10} (A + Bx)(d + ex)^{13} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^13,x, algorithm="fricas")
```

output

```
Too large to include
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5092 vs. $2(479) = 958$.

Time = 0.36 (sec) , antiderivative size = 5092, normalized size of antiderivative = 10.97

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{13} dx = \text{Too large to display}$$

input `integrate((b*x+a)**10*(B*x+A)*(e*x+d)**13,x)`

output

```
A*a**10*d**13*x + B*b**10*e**13*x**25/25 + x**24*(A*b**10*e**13/24 + 5*B*a
*b**9*e**13/12 + 13*B*b**10*d*e**12/24) + x**23*(10*A*a*b**9*e**13/23 + 13
*A*b**10*d*e**12/23 + 45*B*a**2*b**8*e**13/23 + 130*B*a*b**9*d*e**12/23 +
78*B*b**10*d**2*e**11/23) + x**22*(45*A*a**2*b**8*e**13/22 + 65*A*a*b**9*d
*e**12/11 + 39*A*b**10*d**2*e**11/11 + 60*B*a**3*b**7*e**13/11 + 585*B*a**
2*b**8*d*e**12/22 + 390*B*a*b**9*d**2*e**11/11 + 13*B*b**10*d**3*e**10) +
x**21*(40*A*a**3*b**7*e**13/7 + 195*A*a**2*b**8*d*e**12/7 + 260*A*a*b**9*d
**2*e**11/7 + 286*A*b**10*d**3*e**10/21 + 10*B*a**4*b**6*e**13 + 520*B*a**
3*b**7*d*e**12/7 + 1170*B*a**2*b**8*d**2*e**11/7 + 2860*B*a*b**9*d**3*e**1
0/21 + 715*B*b**10*d**4*e**9/21) + x**20*(21*A*a**4*b**6*e**13/2 + 78*A*a*
*3*b**7*d*e**12 + 351*A*a**2*b**8*d**2*e**11/2 + 143*A*a*b**9*d**3*e**10 +
143*A*b**10*d**4*e**9/4 + 63*B*a**5*b**5*e**13/5 + 273*B*a**4*b**6*d*e**1
2/2 + 468*B*a**3*b**7*d**2*e**11 + 1287*B*a**2*b**8*d**3*e**10/2 + 715*B*a
*b**9*d**4*e**9/2 + 1287*B*b**10*d**5*e**8/20) + x**19*(252*A*a**5*b**5*e*
*13/19 + 2730*A*a**4*b**6*d*e**12/19 + 9360*A*a**3*b**7*d**2*e**11/19 + 12
870*A*a**2*b**8*d**3*e**10/19 + 7150*A*a*b**9*d**4*e**9/19 + 1287*A*b**10*
d**5*e**8/19 + 210*B*a**6*b**4*e**13/19 + 3276*B*a**5*b**5*d*e**12/19 + 16
380*B*a**4*b**6*d**2*e**11/19 + 34320*B*a**3*b**7*d**3*e**10/19 + 32175*B*
a**2*b**8*d**4*e**9/19 + 12870*B*a*b**9*d**5*e**8/19 + 1716*B*b**10*d**6*e
**7/19) + x**18*(35*A*a**6*b**4*e**13/3 + 182*A*a**5*b**5*d*e**12 + 910...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3905 vs. $2(440) = 880$.

Time = 0.07 (sec) , antiderivative size = 3905, normalized size of antiderivative = 8.42

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{13} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)*(e*x+d)^13,x, algorithm="maxima")`

output

```
1/25*B*b^10*e^13*x^25 + A*a^10*d^13*x + 1/24*(13*B*b^10*d*e^12 + (10*B*a*b^9 + A*b^10)*e^13)*x^24 + 1/23*(78*B*b^10*d^2*e^11 + 13*(10*B*a*b^9 + A*b^10)*d*e^12 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*e^13)*x^23 + 1/22*(286*B*b^10*d^3*e^10 + 78*(10*B*a*b^9 + A*b^10)*d^2*e^11 + 65*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^12 + 15*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^13)*x^22 + 1/21*(715*B*b^10*d^4*e^9 + 286*(10*B*a*b^9 + A*b^10)*d^3*e^10 + 390*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^11 + 195*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^12 + 30*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^13)*x^21 + 1/20*(1287*B*b^10*d^5*e^8 + 715*(10*B*a*b^9 + A*b^10)*d^4*e^9 + 1430*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^10 + 1170*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^11 + 390*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^12 + 42*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^13)*x^20 + 1/19*(1716*B*b^10*d^6*e^7 + 1287*(10*B*a*b^9 + A*b^10)*d^5*e^8 + 3575*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^9 + 4290*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^10 + 2340*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^11 + 546*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^12 + 42*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^13)*x^19 + 1/6*(572*B*b^10*d^7*e^6 + 572*(10*B*a*b^9 + A*b^10)*d^6*e^7 + 2145*(9*B*a^2*b^8 + 2*A*a*b^9)*d^5*e^8 + 3575*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^4*e^9 + 2860*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^3*e^10 + 1092*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e^11 + 182*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d*e^12 + 10*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e^13)*x^18 + 3/17*(429*B*b^10*d^8*e^5 + 572*(10*B*a*b^9 + A*b^10)*d^7*e^6 + 2860*(9*B*a^2*b^8 + 2*A*a*b^9)*d^6*e^7 + 64...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5039 vs. $2(440) = 880$.

Time = 0.13 (sec) , antiderivative size = 5039, normalized size of antiderivative = 10.86

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{13} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)*(e*x+d)^13,x, algorithm="giac")`

output

```
1/25*B*b^10*e^13*x^25 + 13/24*B*b^10*d*e^12*x^24 + 5/12*B*a*b^9*e^13*x^24
+ 1/24*A*b^10*e^13*x^24 + 78/23*B*b^10*d^2*e^11*x^23 + 130/23*B*a*b^9*d*e^
12*x^23 + 13/23*A*b^10*d*e^12*x^23 + 45/23*B*a^2*b^8*e^13*x^23 + 10/23*A*a
*b^9*e^13*x^23 + 13*B*b^10*d^3*e^10*x^22 + 390/11*B*a*b^9*d^2*e^11*x^22 +
39/11*A*b^10*d^2*e^11*x^22 + 585/22*B*a^2*b^8*d*e^12*x^22 + 65/11*A*a*b^9*
d*e^12*x^22 + 60/11*B*a^3*b^7*e^13*x^22 + 45/22*A*a^2*b^8*e^13*x^22 + 715/
21*B*b^10*d^4*e^9*x^21 + 2860/21*B*a*b^9*d^3*e^10*x^21 + 286/21*A*b^10*d^3
*e^10*x^21 + 1170/7*B*a^2*b^8*d^2*e^11*x^21 + 260/7*A*a*b^9*d^2*e^11*x^21
+ 520/7*B*a^3*b^7*d*e^12*x^21 + 195/7*A*a^2*b^8*d*e^12*x^21 + 10*B*a^4*b^6
*e^13*x^21 + 40/7*A*a^3*b^7*e^13*x^21 + 1287/20*B*b^10*d^5*e^8*x^20 + 715/
2*B*a*b^9*d^4*e^9*x^20 + 143/4*A*b^10*d^4*e^9*x^20 + 1287/2*B*a^2*b^8*d^3*
e^10*x^20 + 143*A*a*b^9*d^3*e^10*x^20 + 468*B*a^3*b^7*d^2*e^11*x^20 + 351/
2*A*a^2*b^8*d^2*e^11*x^20 + 273/2*B*a^4*b^6*d*e^12*x^20 + 78*A*a^3*b^7*d*e
^12*x^20 + 63/5*B*a^5*b^5*e^13*x^20 + 21/2*A*a^4*b^6*e^13*x^20 + 1716/19*B
*b^10*d^6*e^7*x^19 + 12870/19*B*a*b^9*d^5*e^8*x^19 + 1287/19*A*b^10*d^5*e^
8*x^19 + 32175/19*B*a^2*b^8*d^4*e^9*x^19 + 7150/19*A*a*b^9*d^4*e^9*x^19 +
34320/19*B*a^3*b^7*d^3*e^10*x^19 + 12870/19*A*a^2*b^8*d^3*e^10*x^19 + 1638
0/19*B*a^4*b^6*d^2*e^11*x^19 + 9360/19*A*a^3*b^7*d^2*e^11*x^19 + 3276/19*B
*a^5*b^5*d*e^12*x^19 + 2730/19*A*a^4*b^6*d*e^12*x^19 + 210/19*B*a^6*b^4*e^
13*x^19 + 252/19*A*a^5*b^5*e^13*x^19 + 286/3*B*b^10*d^7*e^6*x^18 + 2860...
```

Mupad [B] (verification not implemented)

Time = 1.01 (sec) , antiderivative size = 4206, normalized size of antiderivative = 9.06

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{13} dx = \text{Too large to display}$$

input `int((A + B*x)*(a + b*x)^10*(d + e*x)^13,x)`

output

```
x^13*(A*a^10*d*e^12 + B*b^10*d^12*e + 6*A*b^10*d^11*e^2 + 6*B*a^10*d^2*e^11 + 220*A*a*b^9*d^10*e^3 + 60*A*a^9*b*d^2*e^11 + 60*B*a*b^9*d^11*e^2 + 220*B*a^9*b*d^3*e^10 + 2475*A*a^2*b^8*d^9*e^4 + 11880*A*a^3*b^7*d^8*e^5 + 27720*A*a^4*b^6*d^7*e^6 + 33264*A*a^5*b^5*d^6*e^7 + 20790*A*a^6*b^4*d^5*e^8 + 6600*A*a^7*b^3*d^4*e^9 + 990*A*a^8*b^2*d^3*e^10 + 990*B*a^2*b^8*d^10*e^3 + 6600*B*a^3*b^7*d^9*e^4 + 20790*B*a^4*b^6*d^8*e^5 + 33264*B*a^5*b^5*d^7*e^6 + 27720*B*a^6*b^4*d^6*e^7 + 11880*B*a^7*b^3*d^5*e^8 + 2475*B*a^8*b^2*d^4*e^9) + x^5*(42*A*a^6*b^4*d^13 + 24*B*a^7*b^3*d^13 + 143*A*a^10*d^9*e^4 + (286*B*a^10*d^10*e^3)/5 + 312*A*a^7*b^3*d^12*e + 572*A*a^9*b*d^10*e^3 + 117*B*a^8*b^2*d^12*e + 156*B*a^9*b*d^11*e^2 + 702*A*a^8*b^2*d^11*e^2) + x^8*(15*A*a^3*b^7*d^13 + (105*B*a^4*b^6*d^13)/4 + (429*A*a^10*d^6*e^7)/2 + (429*B*a^10*d^7*e^6)/2 + (1365*A*a^4*b^6*d^12*e)/4 + 2145*A*a^9*b*d^7*e^6 + (819*B*a^5*b^5*d^12*e)/2 + (6435*B*a^9*b*d^8*e^5)/4 + 2457*A*a^5*b^5*d^11*e^2 + (15015*A*a^6*b^4*d^10*e^3)/2 + 10725*A*a^7*b^3*d^9*e^4 + (57915*A*a^8*b^2*d^8*e^5)/8 + (4095*B*a^6*b^4*d^11*e^2)/2 + 4290*B*a^7*b^3*d^10*e^3 + (32175*B*a^8*b^2*d^9*e^4)/8) + x^21*((40*A*a^3*b^7*e^13)/7 + 10*B*a^4*b^6*e^13 + (286*A*b^10*d^3*e^10)/21 + (715*B*b^10*d^4*e^9)/21 + (260*A*a*b^9*d^2*e^11)/7 + (195*A*a^2*b^8*d*e^12)/7 + (2860*B*a*b^9*d^3*e^10)/21 + (520*B*a^3*b^7*d*e^12)/7 + (1170*B*a^2*b^8*d^2*e^11)/7) + x^18*((35*A*a^6*b^4*e^13)/3 + (20*B*a^7*b^3*e^13)/3 + (286*A*b^10*d^6*e^7)/3 + (286*B*b^10*...
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 2593, normalized size of antiderivative = 5.59

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{13} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)*(e*x+d)^13,x)`

output

```
(x*(62403600*a**11*d**13 + 405623400*a**11*d**12*e*x + 1622493600*a**11*d*
*11*e**2*x**2 + 4461857400*a**11*d**10*e**3*x**3 + 8923714800*a**11*d**9*e
**4*x**4 + 13385572200*a**11*d**8*e**5*x**5 + 15297796800*a**11*d**7*e**6*
x**6 + 13385572200*a**11*d**6*e**7*x**7 + 8923714800*a**11*d**5*e**8*x**8
+ 4461857400*a**11*d**4*e**9*x**9 + 1622493600*a**11*d**3*e**10*x**10 + 40
5623400*a**11*d**2*e**11*x**11 + 62403600*a**11*d*e**12*x**12 + 4457400*a*
*11*e**13*x**13 + 343219800*a**10*b*d**13*x + 2974571600*a**10*b*d**12*e*x
**2 + 13385572200*a**10*b*d**11*e**2*x**3 + 39264345120*a**10*b*d**10*e**3
*x**4 + 81800719000*a**10*b*d**9*e**4*x**5 + 126206823600*a**10*b*d**8*e**
5*x**6 + 147241294200*a**10*b*d**7*e**6*x**7 + 130881150400*a**10*b*d**6*e
**7*x**8 + 88344776520*a**10*b*d**5*e**8*x**9 + 44618574000*a**10*b*d**4*e
**9*x**10 + 16360143800*a**10*b*d**3*e**10*x**11 + 4118637600*a**10*b*d**2
*e**11*x**12 + 637408200*a**10*b*d*e**12*x**13 + 45762640*a**10*b*e**13*x*
*14 + 1144066000*a**9*b**2*d**13*x**2 + 11154643500*a**9*b**2*d**12*e*x**3
+ 53542288800*a**9*b**2*d**11*e**2*x**4 + 163601438000*a**9*b**2*d**10*e*
*3*x**5 + 350574510000*a**9*b**2*d**9*e**4*x**6 + 552154853250*a**9*b**2*d
**8*e**5*x**7 + 654405752000*a**9*b**2*d**7*e**6*x**8 + 588965176800*a**9*
b**2*d**6*e**7*x**9 + 401567166000*a**9*b**2*d**5*e**8*x**10 + 20450179750
0*a**9*b**2*d**4*e**9*x**11 + 75508356000*a**9*b**2*d**3*e**10*x**12 + 191
22246000*a**9*b**2*d**2*e**11*x**13 + 2974571600*a**9*b**2*d*e**12*x**1...
```

3.66 $\int (a + bx)^{10}(A + Bx)(d + ex)^{12} dx$

Optimal result	667
Mathematica [B] (verified)	668
Rubi [A] (verified)	669
Maple [B] (verified)	671
Fricas [B] (verification not implemented)	672
Sympy [B] (verification not implemented)	673
Maxima [B] (verification not implemented)	674
Giac [B] (verification not implemented)	675
Mupad [B] (verification not implemented)	676
Reduce [B] (verification not implemented)	676

Optimal result

Integrand size = 20, antiderivative size = 464

$$\begin{aligned}
 & \int (a + bx)^{10}(A + Bx)(d + ex)^{12} dx \\
 &= -\frac{(bd - ae)^{10}(Bd - Ae)(d + ex)^{13}}{13e^{12}} + \frac{(bd - ae)^9(11bBd - 10Abe - aBe)(d + ex)^{14}}{14e^{12}} \\
 &\quad - \frac{b(bd - ae)^8(11bBd - 9Abe - 2aBe)(d + ex)^{15}}{3e^{12}} \\
 &\quad + \frac{15b^2(bd - ae)^7(11bBd - 8Abe - 3aBe)(d + ex)^{16}}{16e^{12}} \\
 &\quad - \frac{30b^3(bd - ae)^6(11bBd - 7Abe - 4aBe)(d + ex)^{17}}{17e^{12}} \\
 &\quad + \frac{7b^4(bd - ae)^5(11bBd - 6Abe - 5aBe)(d + ex)^{18}}{3e^{12}} \\
 &\quad - \frac{42b^5(bd - ae)^4(11bBd - 5Abe - 6aBe)(d + ex)^{19}}{19e^{12}} \\
 &\quad + \frac{3b^6(bd - ae)^3(11bBd - 4Abe - 7aBe)(d + ex)^{20}}{2e^{12}} \\
 &\quad - \frac{5b^7(bd - ae)^2(11bBd - 3Abe - 8aBe)(d + ex)^{21}}{7e^{12}} \\
 &\quad + \frac{5b^8(bd - ae)(11bBd - 2Abe - 9aBe)(d + ex)^{22}}{22e^{12}} \\
 &\quad - \frac{b^9(11bBd - Abe - 10aBe)(d + ex)^{23}}{23e^{12}} + \frac{b^{10}B(d + ex)^{24}}{24e^{12}}
 \end{aligned}$$

output

```

-1/13*(-a*e+b*d)^10*(-A*e+B*d)*(e*x+d)^13/e^12+1/14*(-a*e+b*d)^9*(-10*A*b*
e-B*a*e+11*B*b*d)*(e*x+d)^14/e^12-1/3*b*(-a*e+b*d)^8*(-9*A*b*e-2*B*a*e+11*
B*b*d)*(e*x+d)^15/e^12+15/16*b^2*(-a*e+b*d)^7*(-8*A*b*e-3*B*a*e+11*B*b*d)*
(e*x+d)^16/e^12-30/17*b^3*(-a*e+b*d)^6*(-7*A*b*e-4*B*a*e+11*B*b*d)*(e*x+d)
^17/e^12+7/3*b^4*(-a*e+b*d)^5*(-6*A*b*e-5*B*a*e+11*B*b*d)*(e*x+d)^18/e^12-
42/19*b^5*(-a*e+b*d)^4*(-5*A*b*e-6*B*a*e+11*B*b*d)*(e*x+d)^19/e^12+3/2*b^6
*(-a*e+b*d)^3*(-4*A*b*e-7*B*a*e+11*B*b*d)*(e*x+d)^20/e^12-5/7*b^7*(-a*e+b*
d)^2*(-3*A*b*e-8*B*a*e+11*B*b*d)*(e*x+d)^21/e^12+5/22*b^8*(-a*e+b*d)*(-2*A
*b*e-9*B*a*e+11*B*b*d)*(e*x+d)^22/e^12-1/23*b^9*(-A*b*e-10*B*a*e+11*B*b*d)
*(e*x+d)^23/e^12+1/24*b^10*B*(e*x+d)^24/e^12

```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 3320 vs. $2(464) = 928$.

Time = 0.88 (sec) , antiderivative size = 3320, normalized size of antiderivative = 7.16

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{12} dx = \text{Result too large to show}$$

input

```
Integrate[(a + b*x)^10*(A + B*x)*(d + e*x)^12,x]
```

output

```

a^10*A*d^12*x + (a^9*d^11*(a*B*d + 2*A*(5*b*d + 6*a*e))*x^2)/2 + (a^8*d^10
*(2*a*B*d*(5*b*d + 6*a*e) + 3*A*(15*b^2*d^2 + 40*a*b*d*e + 22*a^2*e^2))*x^
3)/3 + (a^7*d^9*(3*a*B*d*(15*b^2*d^2 + 40*a*b*d*e + 22*a^2*e^2) + 20*A*(6*
b^3*d^3 + 27*a*b^2*d^2*e + 33*a^2*b*d*e^2 + 11*a^3*e^3))*x^4)/4 + a^6*d^8*
(4*a*B*d*(6*b^3*d^3 + 27*a*b^2*d^2*e + 33*a^2*b*d*e^2 + 11*a^3*e^3) + A*(4
2*b^4*d^4 + 288*a*b^3*d^3*e + 594*a^2*b^2*d^2*e^2 + 440*a^3*b*d*e^3 + 99*a
^4*e^4))*x^5 + (a^5*d^7*(5*a*B*d*(42*b^4*d^4 + 288*a*b^3*d^3*e + 594*a^2*b
^2*d^2*e^2 + 440*a^3*b*d*e^3 + 99*a^4*e^4) + 18*A*(14*b^5*d^5 + 140*a*b^4*
d^4*e + 440*a^2*b^3*d^3*e^2 + 550*a^3*b^2*d^2*e^3 + 275*a^4*b*d*e^4 + 44*a
^5*e^5))*x^6)/6 + (3*a^4*d^6*(6*a*B*d*(14*b^5*d^5 + 140*a*b^4*d^4*e + 440*
a^2*b^3*d^3*e^2 + 550*a^3*b^2*d^2*e^3 + 275*a^4*b*d*e^4 + 44*a^5*e^5) + A*
(70*b^6*d^6 + 1008*a*b^5*d^5*e + 4620*a^2*b^4*d^4*e^2 + 8800*a^3*b^3*d^3*e
^3 + 7425*a^4*b^2*d^2*e^4 + 2640*a^5*b*d*e^5 + 308*a^6*e^6))*x^7)/7 + (3*a
^3*d^5*(a*B*d*(70*b^6*d^6 + 1008*a*b^5*d^5*e + 4620*a^2*b^4*d^4*e^2 + 8800
*a^3*b^3*d^3*e^3 + 7425*a^4*b^2*d^2*e^4 + 2640*a^5*b*d*e^5 + 308*a^6*e^6)
+ 8*A*(5*b^7*d^7 + 105*a*b^6*d^6*e + 693*a^2*b^5*d^5*e^2 + 1925*a^3*b^4*d^
4*e^3 + 2475*a^4*b^3*d^3*e^4 + 1485*a^5*b^2*d^2*e^5 + 385*a^6*b*d*e^6 + 33
*a^7*e^7))*x^8)/8 + (a^2*d^4*(8*a*B*d*(5*b^7*d^7 + 105*a*b^6*d^6*e + 693*a
^2*b^5*d^5*e^2 + 1925*a^3*b^4*d^4*e^3 + 2475*a^4*b^3*d^3*e^4 + 1485*a^5*b^
2*d^2*e^5 + 385*a^6*b*d*e^6 + 33*a^7*e^7) + 15*A*(b^8*d^8 + 32*a*b^7*d^...

```

Rubi [A] (verified)

Time = 5.35 (sec) , antiderivative size = 464, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^{10} (A + Bx)(d + ex)^{12} dx$$

$$\downarrow 86$$

$$\int \left(\frac{b^9 (d + ex)^{22} (10aBe + Abe - 11bBd)}{e^{11}} - \frac{5b^8 (d + ex)^{21} (bd - ae)(9aBe + 2Abe - 11bBd)}{e^{11}} + \frac{15b^7 (d + ex)^{20} (b^2 d^2 + 2bde + e^2)}{e^{11}} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
& - \frac{b^9(d+ex)^{23}(-10aBe - Abe + 11bBd)}{23e^{12}} + \frac{5b^8(d+ex)^{22}(bd-ae)(-9aBe - 2Abe + 11bBd)}{22e^{12}} - \\
& \frac{5b^7(d+ex)^{21}(bd-ae)^2(-8aBe - 3Abe + 11bBd)}{7e^{12}} + \\
& \frac{3b^6(d+ex)^{20}(bd-ae)^3(-7aBe - 4Abe + 11bBd)}{2e^{12}} - \\
& \frac{42b^5(d+ex)^{19}(bd-ae)^4(-6aBe - 5Abe + 11bBd)}{19e^{12}} + \\
& \frac{7b^4(d+ex)^{18}(bd-ae)^5(-5aBe - 6Abe + 11bBd)}{3e^{12}} - \\
& \frac{30b^3(d+ex)^{17}(bd-ae)^6(-4aBe - 7Abe + 11bBd)}{17e^{12}} + \\
& \frac{15b^2(d+ex)^{16}(bd-ae)^7(-3aBe - 8Abe + 11bBd)}{16e^{12}} - \\
& \frac{b(d+ex)^{15}(bd-ae)^8(-2aBe - 9Abe + 11bBd)}{3e^{12}} + \\
& \frac{(d+ex)^{14}(bd-ae)^9(-aBe - 10Abe + 11bBd)}{14e^{12}} - \frac{(d+ex)^{13}(bd-ae)^{10}(Bd-Ae)}{13e^{12}} + \\
& \frac{b^{10}B(d+ex)^{24}}{24e^{12}}
\end{aligned}$$

input `Int[(a + b*x)^10*(A + B*x)*(d + e*x)^12,x]`

output

```

-1/13*((b*d - a*e)^10*(B*d - A*e)*(d + e*x)^13)/e^12 + ((b*d - a*e)^9*(11*
b*B*d - 10*A*b*e - a*B*e)*(d + e*x)^14)/(14*e^12) - (b*(b*d - a*e)^8*(11*b
*B*d - 9*A*b*e - 2*a*B*e)*(d + e*x)^15)/(3*e^12) + (15*b^2*(b*d - a*e)^7*(
11*b*B*d - 8*A*b*e - 3*a*B*e)*(d + e*x)^16)/(16*e^12) - (30*b^3*(b*d - a*e
)^6*(11*b*B*d - 7*A*b*e - 4*a*B*e)*(d + e*x)^17)/(17*e^12) + (7*b^4*(b*d -
a*e)^5*(11*b*B*d - 6*A*b*e - 5*a*B*e)*(d + e*x)^18)/(3*e^12) - (42*b^5*(b
*d - a*e)^4*(11*b*B*d - 5*A*b*e - 6*a*B*e)*(d + e*x)^19)/(19*e^12) + (3*b^
6*(b*d - a*e)^3*(11*b*B*d - 4*A*b*e - 7*a*B*e)*(d + e*x)^20)/(2*e^12) - (5
*b^7*(b*d - a*e)^2*(11*b*B*d - 3*A*b*e - 8*a*B*e)*(d + e*x)^21)/(7*e^12) +
(5*b^8*(b*d - a*e)*(11*b*B*d - 2*A*b*e - 9*a*B*e)*(d + e*x)^22)/(22*e^12)
- (b^9*(11*b*B*d - A*b*e - 10*a*B*e)*(d + e*x)^23)/(23*e^12) + (b^10*B*(d
+ e*x)^24)/(24*e^12)

```

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)
), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && ( !IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3608 vs. $2(440) = 880$.

Time = 0.33 (sec) , antiderivative size = 3609, normalized size of antiderivative = 7.78

method	result	size
default	Expression too large to display	3609
norman	Expression too large to display	3922
gospers	Expression too large to display	4664
risch	Expression too large to display	4664
parallelrisch	Expression too large to display	4664
orering	Expression too large to display	4664

input

```
int((b*x+a)^10*(B*x+A)*(e*x+d)^12,x,method=_RETURNVERBOSE)
```

output

```

1/24*b^10*B*e^12*x^24+1/23*((A*b^10+10*B*a*b^9)*e^12+12*b^10*B*d*e^11)*x^2
3+1/22*((10*A*a*b^9+45*B*a^2*b^8)*e^12+12*(A*b^10+10*B*a*b^9)*d*e^11+66*b^
10*B*d^2*e^10)*x^22+1/21*((45*A*a^2*b^8+120*B*a^3*b^7)*e^12+12*(10*A*a*b^9
+45*B*a^2*b^8)*d*e^11+66*(A*b^10+10*B*a*b^9)*d^2*e^10+220*b^10*B*d^3*e^9)*
x^21+1/20*((120*A*a^3*b^7+210*B*a^4*b^6)*e^12+12*(45*A*a^2*b^8+120*B*a^3*b
^7)*d*e^11+66*(10*A*a*b^9+45*B*a^2*b^8)*d^2*e^10+220*(A*b^10+10*B*a*b^9)*d
^3*e^9+495*b^10*B*d^4*e^8)*x^20+1/19*((210*A*a^4*b^6+252*B*a^5*b^5)*e^12+1
2*(120*A*a^3*b^7+210*B*a^4*b^6)*d*e^11+66*(45*A*a^2*b^8+120*B*a^3*b^7)*d^2
*e^10+220*(10*A*a*b^9+45*B*a^2*b^8)*d^3*e^9+495*(A*b^10+10*B*a*b^9)*d^4*e^
8+792*b^10*B*d^5*e^7)*x^19+1/18*((252*A*a^5*b^5+210*B*a^6*b^4)*e^12+12*(21
0*A*a^4*b^6+252*B*a^5*b^5)*d*e^11+66*(120*A*a^3*b^7+210*B*a^4*b^6)*d^2*e^1
0+220*(45*A*a^2*b^8+120*B*a^3*b^7)*d^3*e^9+495*(10*A*a*b^9+45*B*a^2*b^8)*d
^4*e^8+792*(A*b^10+10*B*a*b^9)*d^5*e^7+924*b^10*B*d^6*e^6)*x^18+1/17*((210
*A*a^6*b^4+120*B*a^7*b^3)*e^12+12*(252*A*a^5*b^5+210*B*a^6*b^4)*d*e^11+66*
(210*A*a^4*b^6+252*B*a^5*b^5)*d^2*e^10+220*(120*A*a^3*b^7+210*B*a^4*b^6)*d
^3*e^9+495*(45*A*a^2*b^8+120*B*a^3*b^7)*d^4*e^8+792*(10*A*a*b^9+45*B*a^2*b
^8)*d^5*e^7+924*(A*b^10+10*B*a*b^9)*d^6*e^6+792*b^10*B*d^7*e^5)*x^17+1/16*
((120*A*a^7*b^3+45*B*a^8*b^2)*e^12+12*(210*A*a^6*b^4+120*B*a^7*b^3)*d*e^11
+66*(252*A*a^5*b^5+210*B*a^6*b^4)*d^2*e^10+220*(210*A*a^4*b^6+252*B*a^5*b^
5)*d^3*e^9+495*(120*A*a^3*b^7+210*B*a^4*b^6)*d^4*e^8+792*(45*A*a^2*b^8+...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3621 vs. $2(440) = 880$.

Time = 0.13 (sec) , antiderivative size = 3621, normalized size of antiderivative = 7.80

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{12} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^12,x, algorithm="fricas")
```

output

Too large to include

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4655 vs. $2(478) = 956$.

Time = 0.27 (sec) , antiderivative size = 4655, normalized size of antiderivative = 10.03

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{12} dx = \text{Too large to display}$$

input `integrate((b*x+a)**10*(B*x+A)*(e*x+d)**12,x)`

output

```
A*a**10*d**12*x + B*b**10*e**12*x**24/24 + x**23*(A*b**10*e**12/23 + 10*B*
a*b**9*e**12/23 + 12*B*b**10*d*e**11/23) + x**22*(5*A*a*b**9*e**12/11 + 6*
A*b**10*d*e**11/11 + 45*B*a**2*b**8*e**12/22 + 60*B*a*b**9*d*e**11/11 + 3*
B*b**10*d**2*e**10) + x**21*(15*A*a**2*b**8*e**12/7 + 40*A*a*b**9*d*e**11/
7 + 22*A*b**10*d**2*e**10/7 + 40*B*a**3*b**7*e**12/7 + 180*B*a**2*b**8*d*e
**11/7 + 220*B*a*b**9*d**2*e**10/7 + 220*B*b**10*d**3*e**9/21) + x**20*(6*
A*a**3*b**7*e**12 + 27*A*a**2*b**8*d*e**11 + 33*A*a*b**9*d**2*e**10 + 11*A
*b**10*d**3*e**9 + 21*B*a**4*b**6*e**12/2 + 72*B*a**3*b**7*d*e**11 + 297*B
*a**2*b**8*d**2*e**10/2 + 110*B*a*b**9*d**3*e**9 + 99*B*b**10*d**4*e**8/4)
+ x**19*(210*A*a**4*b**6*e**12/19 + 1440*A*a**3*b**7*d*e**11/19 + 2970*A*
a**2*b**8*d**2*e**10/19 + 2200*A*a*b**9*d**3*e**9/19 + 495*A*b**10*d**4*e*
*8/19 + 252*B*a**5*b**5*e**12/19 + 2520*B*a**4*b**6*d*e**11/19 + 7920*B*a*
*3*b**7*d**2*e**10/19 + 9900*B*a**2*b**8*d**3*e**9/19 + 4950*B*a*b**9*d**4
*e**8/19 + 792*B*b**10*d**5*e**7/19) + x**18*(14*A*a**5*b**5*e**12 + 140*A
*a**4*b**6*d*e**11 + 440*A*a**3*b**7*d**2*e**10 + 550*A*a**2*b**8*d**3*e**
9 + 275*A*a*b**9*d**4*e**8 + 44*A*b**10*d**5*e**7 + 35*B*a**6*b**4*e**12/3
+ 168*B*a**5*b**5*d*e**11 + 770*B*a**4*b**6*d**2*e**10 + 4400*B*a**3*b**7
*d**3*e**9/3 + 2475*B*a**2*b**8*d**4*e**8/2 + 440*B*a*b**9*d**5*e**7 + 154
*B*b**10*d**6*e**6/3) + x**17*(210*A*a**6*b**4*e**12/17 + 3024*A*a**5*b**5
*d*e**11/17 + 13860*A*a**4*b**6*d**2*e**10/17 + 26400*A*a**3*b**7*d**3*...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3621 vs. $2(440) = 880$.

Time = 0.07 (sec) , antiderivative size = 3621, normalized size of antiderivative = 7.80

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{12} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)*(e*x+d)^12,x, algorithm="maxima")`

output

```
1/24*B*b^10*e^12*x^24 + A*a^10*d^12*x + 1/23*(12*B*b^10*d*e^11 + (10*B*a*b^9 + A*b^10)*e^12)*x^23 + 1/22*(66*B*b^10*d^2*e^10 + 12*(10*B*a*b^9 + A*b^10)*d*e^11 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*e^12)*x^22 + 1/21*(220*B*b^10*d^3*e^9 + 66*(10*B*a*b^9 + A*b^10)*d^2*e^10 + 60*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^11 + 15*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^12)*x^21 + 1/4*(99*B*b^10*d^4*e^8 + 44*(10*B*a*b^9 + A*b^10)*d^3*e^9 + 66*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^10 + 36*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^11 + 6*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^12)*x^20 + 1/19*(792*B*b^10*d^5*e^7 + 495*(10*B*a*b^9 + A*b^10)*d^4*e^8 + 1100*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^9 + 990*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^10 + 360*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^11 + 42*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^12)*x^19 + 1/6*(308*B*b^10*d^6*e^6 + 264*(10*B*a*b^9 + A*b^10)*d^5*e^7 + 825*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^8 + 1100*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^9 + 660*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^10 + 168*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^11 + 14*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^12)*x^18 + 3/17*(264*B*b^10*d^7*e^5 + 308*(10*B*a*b^9 + A*b^10)*d^6*e^6 + 1320*(9*B*a^2*b^8 + 2*A*a*b^9)*d^5*e^7 + 2475*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^4*e^8 + 2200*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^3*e^9 + 924*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e^10 + 168*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d*e^11 + 10*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e^12)*x^17 + 3/16*(165*B*b^10*d^8*e^4 + 264*(10*B*a*b^9 + A*b^10)*d^7*e^5 + 1540*(9*B*a^2*b^8 + 2*A*a*b^9)*d^6*e^6 + 3960*(8*B*a^3*b^7 + ...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4663 vs. $2(440) = 880$.

Time = 0.13 (sec) , antiderivative size = 4663, normalized size of antiderivative = 10.05

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{12} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)*(e*x+d)^12,x, algorithm="giac")`

output

```
1/24*B*b^10*e^12*x^24 + 12/23*B*b^10*d*e^11*x^23 + 10/23*B*a*b^9*e^12*x^23
+ 1/23*A*b^10*e^12*x^23 + 3*B*b^10*d^2*e^10*x^22 + 60/11*B*a*b^9*d*e^11*x
^22 + 6/11*A*b^10*d*e^11*x^22 + 45/22*B*a^2*b^8*e^12*x^22 + 5/11*A*a*b^9*e
^12*x^22 + 220/21*B*b^10*d^3*e^9*x^21 + 220/7*B*a*b^9*d^2*e^10*x^21 + 22/7
*A*b^10*d^2*e^10*x^21 + 180/7*B*a^2*b^8*d*e^11*x^21 + 40/7*A*a*b^9*d*e^11*
x^21 + 40/7*B*a^3*b^7*e^12*x^21 + 15/7*A*a^2*b^8*e^12*x^21 + 99/4*B*b^10*d
^4*e^8*x^20 + 110*B*a*b^9*d^3*e^9*x^20 + 11*A*b^10*d^3*e^9*x^20 + 297/2*B*
a^2*b^8*d^2*e^10*x^20 + 33*A*a*b^9*d^2*e^10*x^20 + 72*B*a^3*b^7*d*e^11*x^2
0 + 27*A*a^2*b^8*d*e^11*x^20 + 21/2*B*a^4*b^6*e^12*x^20 + 6*A*a^3*b^7*e^12
*x^20 + 792/19*B*b^10*d^5*e^7*x^19 + 4950/19*B*a*b^9*d^4*e^8*x^19 + 495/19
*A*b^10*d^4*e^8*x^19 + 9900/19*B*a^2*b^8*d^3*e^9*x^19 + 2200/19*A*a*b^9*d^
3*e^9*x^19 + 7920/19*B*a^3*b^7*d^2*e^10*x^19 + 2970/19*A*a^2*b^8*d^2*e^10*
x^19 + 2520/19*B*a^4*b^6*d*e^11*x^19 + 1440/19*A*a^3*b^7*d*e^11*x^19 + 252
/19*B*a^5*b^5*e^12*x^19 + 210/19*A*a^4*b^6*e^12*x^19 + 154/3*B*b^10*d^6*e^
6*x^18 + 440*B*a*b^9*d^5*e^7*x^18 + 44*A*b^10*d^5*e^7*x^18 + 2475/2*B*a^2*
b^8*d^4*e^8*x^18 + 275*A*a*b^9*d^4*e^8*x^18 + 4400/3*B*a^3*b^7*d^3*e^9*x^1
8 + 550*A*a^2*b^8*d^3*e^9*x^18 + 770*B*a^4*b^6*d^2*e^10*x^18 + 440*A*a^3*b
^7*d^2*e^10*x^18 + 168*B*a^5*b^5*d*e^11*x^18 + 140*A*a^4*b^6*d*e^11*x^18 +
35/3*B*a^6*b^4*e^12*x^18 + 14*A*a^5*b^5*e^12*x^18 + 792/17*B*b^10*d^7*e^5
*x^17 + 9240/17*B*a*b^9*d^6*e^6*x^17 + 924/17*A*b^10*d^6*e^6*x^17 + 356...
```


Mupad [B] (verification not implemented)

Time = 1.82 (sec) , antiderivative size = 3891, normalized size of antiderivative = 8.39

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{12} dx = \text{Too large to display}$$

input `int((A + B*x)*(a + b*x)^10*(d + e*x)^12,x)`

output

```
x^5*(42*A*a^6*b^4*d^12 + 24*B*a^7*b^3*d^12 + 99*A*a^10*d^8*e^4 + 44*B*a^10
*d^9*e^3 + 288*A*a^7*b^3*d^11*e + 440*A*a^9*b*d^9*e^3 + 108*B*a^8*b^2*d^11
*e + 132*B*a^9*b*d^10*e^2 + 594*A*a^8*b^2*d^10*e^2) + x^8*(15*A*a^3*b^7*d^
12 + (105*B*a^4*b^6*d^12)/4 + 99*A*a^10*d^5*e^7 + (231*B*a^10*d^6*e^6)/2 +
315*A*a^4*b^6*d^11*e + 1155*A*a^9*b*d^6*e^6 + 378*B*a^5*b^5*d^11*e + 990*
B*a^9*b*d^7*e^5 + 2079*A*a^5*b^5*d^10*e^2 + 5775*A*a^6*b^4*d^9*e^3 + 7425*
A*a^7*b^3*d^8*e^4 + 4455*A*a^8*b^2*d^7*e^5 + (3465*B*a^6*b^4*d^10*e^2)/2 +
3300*B*a^7*b^3*d^9*e^3 + (22275*B*a^8*b^2*d^8*e^4)/8) + x^20*(6*A*a^3*b^7
*e^12 + (21*B*a^4*b^6*e^12)/2 + 11*A*b^10*d^3*e^9 + (99*B*b^10*d^4*e^8)/4
+ 33*A*a*b^9*d^2*e^10 + 27*A*a^2*b^8*d*e^11 + 110*B*a*b^9*d^3*e^9 + 72*B*a
^3*b^7*d*e^11 + (297*B*a^2*b^8*d^2*e^10)/2) + x^17*((210*A*a^6*b^4*e^12)/1
7 + (120*B*a^7*b^3*e^12)/17 + (924*A*b^10*d^6*e^6)/17 + (792*B*b^10*d^7*e^
5)/17 + (7920*A*a*b^9*d^5*e^7)/17 + (3024*A*a^5*b^5*d*e^11)/17 + (9240*B*a
*b^9*d^6*e^6)/17 + (2520*B*a^6*b^4*d*e^11)/17 + (22275*A*a^2*b^8*d^4*e^8)/
17 + (26400*A*a^3*b^7*d^3*e^9)/17 + (13860*A*a^4*b^6*d^2*e^10)/17 + (35640
*B*a^2*b^8*d^5*e^7)/17 + (59400*B*a^3*b^7*d^4*e^8)/17 + (46200*B*a^4*b^6*d
^3*e^9)/17 + (16632*B*a^5*b^5*d^2*e^10)/17) + x^10*(A*a*b^9*d^12 + (9*B*a^
2*b^8*d^12)/2 + 22*A*a^10*d^3*e^9 + (99*B*a^10*d^4*e^8)/2 + 54*A*a^2*b^8*d
^11*e + 495*A*a^9*b*d^4*e^8 + 144*B*a^3*b^7*d^11*e + 792*B*a^9*b*d^5*e^7 +
792*A*a^3*b^7*d^10*e^2 + 4620*A*a^4*b^6*d^9*e^3 + 12474*A*a^5*b^5*d^8*...
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 2399, normalized size of antiderivative = 5.17

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{12} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)*(e*x+d)^12,x)`

output

```
(x*(32449872*a**11*d**12 + 194699232*a**11*d**11*e*x + 713897184*a**11*d**10*e**2*x**2 + 1784742960*a**11*d**9*e**3*x**3 + 3212537328*a**11*d**8*e**4*x**4 + 4283383104*a**11*d**7*e**5*x**5 + 4283383104*a**11*d**6*e**6*x**6 + 3212537328*a**11*d**5*e**7*x**7 + 1784742960*a**11*d**4*e**8*x**8 + 713897184*a**11*d**3*e**9*x**9 + 194699232*a**11*d**2*e**10*x**10 + 32449872*a**11*d*e**11*x**11 + 2496144*a**11*e**12*x**12 + 178474296*a**10*b*d**12*x + 1427794368*a**10*b*d**11*e*x**2 + 5889651768*a**10*b*d**10*e**2*x**3 + 15705738048*a**10*b*d**9*e**3*x**4 + 29448258840*a**10*b*d**8*e**4*x**5 + 40386183552*a**10*b*d**7*e**5*x**6 + 41227562376*a**10*b*d**6*e**6*x**7 + 31411476096*a**10*b*d**5*e**7*x**8 + 17668955304*a**10*b*d**4*e**8*x**9 + 7138971840*a**10*b*d**3*e**9*x**10 + 1963217256*a**10*b*d**2*e**10*x**11 + 329491008*a**10*b*d*e**11*x**12 + 25496328*a**10*b*e**12*x**13 + 594914320*a**9*b**2*d**12*x**2 + 5354228880*a**9*b**2*d**11*e*x**3 + 23558607072*a**9*b**2*d**10*e**2*x**4 + 65440575200*a**9*b**2*d**9*e**3*x**5 + 126206823600*a**9*b**2*d**8*e**4*x**6 + 176689553040*a**9*b**2*d**7*e**5*x**7 + 183233610560*a**9*b**2*d**6*e**6*x**8 + 141351642432*a**9*b**2*d**5*e**7*x**9 + 80313433200*a**9*b**2*d**4*e**8*x**10 + 32720287600*a**9*b**2*d**3*e**9*x**11 + 9061002720*a**9*b**2*d**2*e**10*x**12 + 1529779680*a**9*b**2*d*e**11*x**13 + 118982864*a**9*b**2*e**12*x**14 + 1338557220*a**8*b**3*d**12*x**3 + 12850149312*a**8*b**3*d**11*e*x**4 + 58896517680*a**8*b**3*d**11...
```

3.67 $\int (a + bx)^{10}(A + Bx)(d + ex)^{11} dx$

Optimal result	678
Mathematica [B] (verified)	679
Rubi [A] (verified)	680
Maple [B] (verified)	682
Fricas [B] (verification not implemented)	683
Sympy [B] (verification not implemented)	684
Maxima [B] (verification not implemented)	685
Giac [B] (verification not implemented)	686
Mupad [B] (verification not implemented)	687
Reduce [B] (verification not implemented)	687

Optimal result

Integrand size = 20, antiderivative size = 461

$$\begin{aligned}
 & \int (a + bx)^{10}(A + Bx)(d + ex)^{11} dx \\
 &= -\frac{(bd - ae)^{10}(Bd - Ae)(d + ex)^{12}}{12e^{12}} + \frac{(bd - ae)^9(11bBd - 10Abe - aBe)(d + ex)^{13}}{13e^{12}} \\
 &\quad - \frac{5b(bd - ae)^8(11bBd - 9Abe - 2aBe)(d + ex)^{14}}{14e^{12}} \\
 &\quad + \frac{b^2(bd - ae)^7(11bBd - 8Abe - 3aBe)(d + ex)^{15}}{e^{12}} \\
 &\quad - \frac{15b^3(bd - ae)^6(11bBd - 7Abe - 4aBe)(d + ex)^{16}}{8e^{12}} \\
 &\quad + \frac{42b^4(bd - ae)^5(11bBd - 6Abe - 5aBe)(d + ex)^{17}}{17e^{12}} \\
 &\quad - \frac{7b^5(bd - ae)^4(11bBd - 5Abe - 6aBe)(d + ex)^{18}}{3e^{12}} \\
 &\quad + \frac{30b^6(bd - ae)^3(11bBd - 4Abe - 7aBe)(d + ex)^{19}}{19e^{12}} \\
 &\quad - \frac{3b^7(bd - ae)^2(11bBd - 3Abe - 8aBe)(d + ex)^{20}}{4e^{12}} \\
 &\quad + \frac{5b^8(bd - ae)(11bBd - 2Abe - 9aBe)(d + ex)^{21}}{21e^{12}} \\
 &\quad - \frac{b^9(11bBd - Abe - 10aBe)(d + ex)^{22}}{22e^{12}} + \frac{b^{10}B(d + ex)^{23}}{23e^{12}}
 \end{aligned}$$

output

```
-1/12*(-a*e+b*d)^10*(-A*e+B*d)*(e*x+d)^12/e^12+1/13*(-a*e+b*d)^9*(-10*A*b*
e-B*a*e+11*B*b*d)*(e*x+d)^13/e^12-5/14*b*(-a*e+b*d)^8*(-9*A*b*e-2*B*a*e+11
*B*b*d)*(e*x+d)^14/e^12+b^2*(-a*e+b*d)^7*(-8*A*b*e-3*B*a*e+11*B*b*d)*(e*x+
d)^15/e^12-15/8*b^3*(-a*e+b*d)^6*(-7*A*b*e-4*B*a*e+11*B*b*d)*(e*x+d)^16/e^
12+42/17*b^4*(-a*e+b*d)^5*(-6*A*b*e-5*B*a*e+11*B*b*d)*(e*x+d)^17/e^12-7/3*
b^5*(-a*e+b*d)^4*(-5*A*b*e-6*B*a*e+11*B*b*d)*(e*x+d)^18/e^12+30/19*b^6*(-a
*e+b*d)^3*(-4*A*b*e-7*B*a*e+11*B*b*d)*(e*x+d)^19/e^12-3/4*b^7*(-a*e+b*d)^2
*(-3*A*b*e-8*B*a*e+11*B*b*d)*(e*x+d)^20/e^12+5/21*b^8*(-a*e+b*d)*(-2*A*b*e
-9*B*a*e+11*B*b*d)*(e*x+d)^21/e^12-1/22*b^9*(-A*b*e-10*B*a*e+11*B*b*d)*(e*
x+d)^22/e^12+1/23*b^10*B*(e*x+d)^23/e^12
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 3018 vs. $2(461) = 922$.

Time = 0.76 (sec) , antiderivative size = 3018, normalized size of antiderivative = 6.55

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{11} dx = \text{Result too large to show}$$

input

```
Integrate[(a + b*x)^10*(A + B*x)*(d + e*x)^11,x]
```

output

```

a^10*A*d^11*x + (a^9*d^10*(10*A*b*d + a*B*d + 11*a*A*e)*x^2)/2 + (a^8*d^9*
(a*B*d*(10*b*d + 11*a*e) + 5*A*(9*b^2*d^2 + 22*a*b*d*e + 11*a^2*e^2))*x^3)
/3 + (5*a^7*d^8*(a*B*d*(9*b^2*d^2 + 22*a*b*d*e + 11*a^2*e^2) + A*(24*b^3*d
^3 + 99*a*b^2*d^2*e + 110*a^2*b*d*e^2 + 33*a^3*e^3))*x^4)/4 + a^6*d^7*(a*B
*d*(24*b^3*d^3 + 99*a*b^2*d^2*e + 110*a^2*b*d*e^2 + 33*a^3*e^3) + 3*A*(14*
b^4*d^4 + 88*a*b^3*d^3*e + 165*a^2*b^2*d^2*e^2 + 110*a^3*b*d*e^3 + 22*a^4*
e^4))*x^5 + (a^5*d^6*(5*a*B*d*(14*b^4*d^4 + 88*a*b^3*d^3*e + 165*a^2*b^2*d
^2*e^2 + 110*a^3*b*d*e^3 + 22*a^4*e^4) + A*(84*b^5*d^5 + 770*a*b^4*d^4*e +
2200*a^2*b^3*d^3*e^2 + 2475*a^3*b^2*d^2*e^3 + 1100*a^4*b*d*e^4 + 154*a^5*
e^5))*x^6)/2 + (3*a^4*d^5*(a*B*d*(84*b^5*d^5 + 770*a*b^4*d^4*e + 2200*a^2*
b^3*d^3*e^2 + 2475*a^3*b^2*d^2*e^3 + 1100*a^4*b*d*e^4 + 154*a^5*e^5) + 2*A
*(35*b^6*d^6 + 462*a*b^5*d^5*e + 1925*a^2*b^4*d^4*e^2 + 3300*a^3*b^3*d^3*e
^3 + 2475*a^4*b^2*d^2*e^4 + 770*a^5*b*d*e^5 + 77*a^6*e^6))*x^7)/7 + (3*a^3
*d^4*(a*B*d*(35*b^6*d^6 + 462*a*b^5*d^5*e + 1925*a^2*b^4*d^4*e^2 + 3300*a^
3*b^3*d^3*e^3 + 2475*a^4*b^2*d^2*e^4 + 770*a^5*b*d*e^5 + 77*a^6*e^6) + 5*A
*(4*b^7*d^7 + 77*a*b^6*d^6*e + 462*a^2*b^5*d^5*e^2 + 1155*a^3*b^4*d^4*e^3
+ 1320*a^4*b^3*d^3*e^4 + 693*a^5*b^2*d^2*e^5 + 154*a^6*b*d*e^6 + 11*a^7*e^
7))*x^8)/4 + (5*a^2*d^3*(2*a*B*d*(4*b^7*d^7 + 77*a*b^6*d^6*e + 462*a^2*b^
5*d^5*e^2 + 1155*a^3*b^4*d^4*e^3 + 1320*a^4*b^3*d^3*e^4 + 693*a^5*b^2*d^2*
e^5 + 154*a^6*b*d*e^6 + 11*a^7*e^7) + A*(3*b^8*d^8 + 88*a*b^7*d^7*e + 77...

```

Rubi [A] (verified)

Time = 4.89 (sec) , antiderivative size = 461, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^{10} (A + Bx)(d + ex)^{11} dx$$

$$\downarrow 86$$

$$\int \left(\frac{b^9 (d + ex)^{21} (10aBe + Abe - 11bBd)}{e^{11}} - \frac{5b^8 (d + ex)^{20} (bd - ae)(9aBe + 2Abe - 11bBd)}{e^{11}} + \frac{15b^7 (d + ex)^{19} (bd - ae)^2}{e^{11}} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
& - \frac{b^9(d+ex)^{22}(-10aBe - Abe + 11bBd)}{22e^{12}} + \frac{5b^8(d+ex)^{21}(bd-ae)(-9aBe - 2Abe + 11bBd)}{21e^{12}} \\
& - \frac{3b^7(d+ex)^{20}(bd-ae)^2(-8aBe - 3Abe + 11bBd)}{3e^{12}} + \frac{4e^{12}}{30b^6(d+ex)^{19}(bd-ae)^3(-7aBe - 4Abe + 11bBd)} \\
& - \frac{19e^{12}}{7b^5(d+ex)^{18}(bd-ae)^4(-6aBe - 5Abe + 11bBd)} + \frac{3e^{12}}{42b^4(d+ex)^{17}(bd-ae)^5(-5aBe - 6Abe + 11bBd)} \\
& - \frac{17e^{12}}{15b^3(d+ex)^{16}(bd-ae)^6(-4aBe - 7Abe + 11bBd)} + \frac{8e^{12}}{b^2(d+ex)^{15}(bd-ae)^7(-3aBe - 8Abe + 11bBd)} \\
& - \frac{e^{12}}{5b(d+ex)^{14}(bd-ae)^8(-2aBe - 9Abe + 11bBd)} + \frac{14e^{12}}{(d+ex)^{13}(bd-ae)^9(-aBe - 10Abe + 11bBd)} \\
& - \frac{(d+ex)^{12}(bd-ae)^{10}(Bd-Ae)}{12e^{12}} + \frac{b^{10}B(d+ex)^{23}}{23e^{12}}
\end{aligned}$$

input `Int[(a + b*x)^10*(A + B*x)*(d + e*x)^11,x]`

output `-1/12*((b*d - a*e)^10*(B*d - A*e)*(d + e*x)^12)/e^12 + ((b*d - a*e)^9*(11*b*B*d - 10*A*b*e - a*B*e)*(d + e*x)^13)/(13*e^12) - (5*b*(b*d - a*e)^8*(11*b*B*d - 9*A*b*e - 2*a*B*e)*(d + e*x)^14)/(14*e^12) + (b^2*(b*d - a*e)^7*(11*b*B*d - 8*A*b*e - 3*a*B*e)*(d + e*x)^15)/e^12 - (15*b^3*(b*d - a*e)^6*(11*b*B*d - 7*A*b*e - 4*a*B*e)*(d + e*x)^16)/(8*e^12) + (42*b^4*(b*d - a*e)^5*(11*b*B*d - 6*A*b*e - 5*a*B*e)*(d + e*x)^17)/(17*e^12) - (7*b^5*(b*d - a*e)^4*(11*b*B*d - 5*A*b*e - 6*a*B*e)*(d + e*x)^18)/(3*e^12) + (30*b^6*(b*d - a*e)^3*(11*b*B*d - 4*A*b*e - 7*a*B*e)*(d + e*x)^19)/(19*e^12) - (3*b^7*(b*d - a*e)^2*(11*b*B*d - 3*A*b*e - 8*a*B*e)*(d + e*x)^20)/(4*e^12) + (5*b^8*(b*d - a*e)*(11*b*B*d - 2*A*b*e - 9*a*B*e)*(d + e*x)^21)/(21*e^12) - (b^9*(11*b*B*d - A*b*e - 10*a*B*e)*(d + e*x)^22)/(22*e^12) + (b^10*B*(d + e*x)^23)/(23*e^12)`

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0]) || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3324 vs. $2(439) = 878$.

Time = 0.32 (sec) , antiderivative size = 3325, normalized size of antiderivative = 7.21

method	result	size
default	Expression too large to display	3325
norman	Expression too large to display	3608
orering	Expression too large to display	4288
gosper	Expression too large to display	4289
risch	Expression too large to display	4289
parallelrisc	Expression too large to display	4289

input

```
int((b*x+a)^10*(B*x+A)*(e*x+d)^11,x,method=_RETURNVERBOSE)
```

output

```

1/23*b^10*B*e^11*x^23+1/22*((A*b^10+10*B*a*b^9)*e^11+11*b^10*B*d*e^10)*x^2
2+1/21*((10*A*a*b^9+45*B*a^2*b^8)*e^11+11*(A*b^10+10*B*a*b^9)*d*e^10+55*b^
10*B*d^2*e^9)*x^21+1/20*((45*A*a^2*b^8+120*B*a^3*b^7)*e^11+11*(10*A*a*b^9+
45*B*a^2*b^8)*d*e^10+55*(A*b^10+10*B*a*b^9)*d^2*e^9+165*b^10*B*d^3*e^8)*x^
20+1/19*((120*A*a^3*b^7+210*B*a^4*b^6)*e^11+11*(45*A*a^2*b^8+120*B*a^3*b^7
)*d*e^10+55*(10*A*a*b^9+45*B*a^2*b^8)*d^2*e^9+165*(A*b^10+10*B*a*b^9)*d^3*
e^8+330*b^10*B*d^4*e^7)*x^19+1/18*((210*A*a^4*b^6+252*B*a^5*b^5)*e^11+11*(
120*A*a^3*b^7+210*B*a^4*b^6)*d*e^10+55*(45*A*a^2*b^8+120*B*a^3*b^7)*d^2*e^
9+165*(10*A*a*b^9+45*B*a^2*b^8)*d^3*e^8+330*(A*b^10+10*B*a*b^9)*d^4*e^7+46
2*b^10*B*d^5*e^6)*x^18+1/17*((252*A*a^5*b^5+210*B*a^6*b^4)*e^11+11*(210*A*
a^4*b^6+252*B*a^5*b^5)*d*e^10+55*(120*A*a^3*b^7+210*B*a^4*b^6)*d^2*e^9+165
*(45*A*a^2*b^8+120*B*a^3*b^7)*d^3*e^8+330*(10*A*a*b^9+45*B*a^2*b^8)*d^4*e^
7+462*(A*b^10+10*B*a*b^9)*d^5*e^6+462*b^10*B*d^6*e^5)*x^17+1/16*((210*A*a^
6*b^4+120*B*a^7*b^3)*e^11+11*(252*A*a^5*b^5+210*B*a^6*b^4)*d*e^10+55*(210*
A*a^4*b^6+252*B*a^5*b^5)*d^2*e^9+165*(120*A*a^3*b^7+210*B*a^4*b^6)*d^3*e^8
+330*(45*A*a^2*b^8+120*B*a^3*b^7)*d^4*e^7+462*(10*A*a*b^9+45*B*a^2*b^8)*d^
5*e^6+462*(A*b^10+10*B*a*b^9)*d^6*e^5+330*b^10*B*d^7*e^4)*x^16+1/15*((120*
A*a^7*b^3+45*B*a^8*b^2)*e^11+11*(210*A*a^6*b^4+120*B*a^7*b^3)*d*e^10+55*(2
52*A*a^5*b^5+210*B*a^6*b^4)*d^2*e^9+165*(210*A*a^4*b^6+252*B*a^5*b^5)*d^3*
e^8+330*(120*A*a^3*b^7+210*B*a^4*b^6)*d^4*e^7+462*(45*A*a^2*b^8+120*B*a...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3334 vs. $2(439) = 878$.

Time = 0.11 (sec) , antiderivative size = 3334, normalized size of antiderivative = 7.23

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{11} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^11,x, algorithm="fricas")
```

output

Too large to include

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4328 vs. $2(476) = 952$.

Time = 0.25 (sec) , antiderivative size = 4328, normalized size of antiderivative = 9.39

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{11} dx = \text{Too large to display}$$

input `integrate((b*x+a)**10*(B*x+A)*(e*x+d)**11,x)`

output

```
A*a**10*d**11*x + B*b**10*e**11*x**23/23 + x**22*(A*b**10*e**11/22 + 5*B*a
*b**9*e**11/11 + B*b**10*d*e**10/2) + x**21*(10*A*a*b**9*e**11/21 + 11*A*b
**10*d*e**10/21 + 15*B*a**2*b**8*e**11/7 + 110*B*a*b**9*d*e**10/21 + 55*B*
b**10*d**2*e**9/21) + x**20*(9*A*a**2*b**8*e**11/4 + 11*A*a*b**9*d*e**10/2
+ 11*A*b**10*d**2*e**9/4 + 6*B*a**3*b**7*e**11 + 99*B*a**2*b**8*d*e**10/4
+ 55*B*a*b**9*d**2*e**9/2 + 33*B*b**10*d**3*e**8/4) + x**19*(120*A*a**3*b
**7*e**11/19 + 495*A*a**2*b**8*d*e**10/19 + 550*A*a*b**9*d**2*e**9/19 + 16
5*A*b**10*d**3*e**8/19 + 210*B*a**4*b**6*e**11/19 + 1320*B*a**3*b**7*d*e**
10/19 + 2475*B*a**2*b**8*d**2*e**9/19 + 1650*B*a*b**9*d**3*e**8/19 + 330*B
*b**10*d**4*e**7/19) + x**18*(35*A*a**4*b**6*e**11/3 + 220*A*a**3*b**7*d*e
**10/3 + 275*A*a**2*b**8*d**2*e**9/2 + 275*A*a*b**9*d**3*e**8/3 + 55*A*b**
10*d**4*e**7/3 + 14*B*a**5*b**5*e**11 + 385*B*a**4*b**6*d*e**10/3 + 1100*B
*a**3*b**7*d**2*e**9/3 + 825*B*a**2*b**8*d**3*e**8/2 + 550*B*a*b**9*d**4*e
**7/3 + 77*B*b**10*d**5*e**6/3) + x**17*(252*A*a**5*b**5*e**11/17 + 2310*A
*a**4*b**6*d*e**10/17 + 6600*A*a**3*b**7*d**2*e**9/17 + 7425*A*a**2*b**8*d
**3*e**8/17 + 3300*A*a*b**9*d**4*e**7/17 + 462*A*b**10*d**5*e**6/17 + 210*
B*a**6*b**4*e**11/17 + 2772*B*a**5*b**5*d*e**10/17 + 11550*B*a**4*b**6*d**
2*e**9/17 + 19800*B*a**3*b**7*d**3*e**8/17 + 14850*B*a**2*b**8*d**4*e**7/1
7 + 4620*B*a*b**9*d**5*e**6/17 + 462*B*b**10*d**6*e**5/17) + x**16*(105*A*
a**6*b**4*e**11/8 + 693*A*a**5*b**5*d*e**10/4 + 5775*A*a**4*b**6*d**2*e...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3334 vs. $2(439) = 878$.

Time = 0.05 (sec) , antiderivative size = 3334, normalized size of antiderivative = 7.23

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{11} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)*(e*x+d)^11,x, algorithm="maxima")`

output

```
1/23*B*b^10*e^11*x^23 + A*a^10*d^11*x + 1/22*(11*B*b^10*d*e^10 + (10*B*a*b^9 + A*b^10)*e^11)*x^22 + 1/21*(55*B*b^10*d^2*e^9 + 11*(10*B*a*b^9 + A*b^10)*d*e^10 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^21 + 1/4*(33*B*b^10*d^3*e^8 + 11*(10*B*a*b^9 + A*b^10)*d^2*e^9 + 11*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 + 3*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^20 + 5/19*(66*B*b^10*d^4*e^7 + 33*(10*B*a*b^9 + A*b^10)*d^3*e^8 + 55*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 + 33*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 + 6*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^19 + 1/6*(154*B*b^10*d^5*e^6 + 110*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 275*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 + 275*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 110*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 + 14*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^11)*x^18 + 3/17*(154*B*b^10*d^6*e^5 + 154*(10*B*a*b^9 + A*b^10)*d^5*e^6 + 550*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 + 825*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 + 550*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 + 154*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10 + 14*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^17 + 3/8*(55*B*b^10*d^7*e^4 + 77*(10*B*a*b^9 + A*b^10)*d^6*e^5 + 385*(9*B*a^2*b^8 + 2*A*a*b^9)*d^5*e^6 + 825*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^4*e^7 + 825*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^3*e^8 + 385*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e^9 + 77*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d*e^10 + 5*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e^11)*x^16 + (11*B*b^10*d^8*e^3 + 22*(10*B*a*b^9 + A*b^10)*d^7*e^4 + 154*(9*B*a^2*b^8 + 2*A*a*b^9)*d^6*e^5 + 462*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^5*e^6 + 660*...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4288 vs. $2(439) = 878$.

Time = 0.13 (sec) , antiderivative size = 4288, normalized size of antiderivative = 9.30

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{11} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)*(e*x+d)^11,x, algorithm="giac")`

output

```

1/23*B*b^10*e^11*x^23 + 1/2*B*b^10*d*e^10*x^22 + 5/11*B*a*b^9*e^11*x^22 +
1/22*A*b^10*e^11*x^22 + 55/21*B*b^10*d^2*e^9*x^21 + 110/21*B*a*b^9*d*e^10*x
x^21 + 11/21*A*b^10*d*e^10*x^21 + 15/7*B*a^2*b^8*e^11*x^21 + 10/21*A*a*b^9
*e^11*x^21 + 33/4*B*b^10*d^3*e^8*x^20 + 55/2*B*a*b^9*d^2*e^9*x^20 + 11/4*A
*b^10*d^2*e^9*x^20 + 99/4*B*a^2*b^8*d*e^10*x^20 + 11/2*A*a*b^9*d*e^10*x^20
+ 6*B*a^3*b^7*e^11*x^20 + 9/4*A*a^2*b^8*e^11*x^20 + 330/19*B*b^10*d^4*e^7
*x^19 + 1650/19*B*a*b^9*d^3*e^8*x^19 + 165/19*A*b^10*d^3*e^8*x^19 + 2475/1
9*B*a^2*b^8*d^2*e^9*x^19 + 550/19*A*a*b^9*d^2*e^9*x^19 + 1320/19*B*a^3*b^7
*d*e^10*x^19 + 495/19*A*a^2*b^8*d*e^10*x^19 + 210/19*B*a^4*b^6*e^11*x^19 +
120/19*A*a^3*b^7*e^11*x^19 + 77/3*B*b^10*d^5*e^6*x^18 + 550/3*B*a*b^9*d^4
*e^7*x^18 + 55/3*A*b^10*d^4*e^7*x^18 + 825/2*B*a^2*b^8*d^3*e^8*x^18 + 275/
3*A*a*b^9*d^3*e^8*x^18 + 1100/3*B*a^3*b^7*d^2*e^9*x^18 + 275/2*A*a^2*b^8*d
^2*e^9*x^18 + 385/3*B*a^4*b^6*d*e^10*x^18 + 220/3*A*a^3*b^7*d*e^10*x^18 +
14*B*a^5*b^5*e^11*x^18 + 35/3*A*a^4*b^6*e^11*x^18 + 462/17*B*b^10*d^6*e^5*x
x^17 + 4620/17*B*a*b^9*d^5*e^6*x^17 + 462/17*A*b^10*d^5*e^6*x^17 + 14850/1
7*B*a^2*b^8*d^4*e^7*x^17 + 3300/17*A*a*b^9*d^4*e^7*x^17 + 19800/17*B*a^3*b
^7*d^3*e^8*x^17 + 7425/17*A*a^2*b^8*d^3*e^8*x^17 + 11550/17*B*a^4*b^6*d^2*
e^9*x^17 + 6600/17*A*a^3*b^7*d^2*e^9*x^17 + 2772/17*B*a^5*b^5*d*e^10*x^17
+ 2310/17*A*a^4*b^6*d*e^10*x^17 + 210/17*B*a^6*b^4*e^11*x^17 + 252/17*A*a^
5*b^5*e^11*x^17 + 165/8*B*b^10*d^7*e^4*x^16 + 1155/4*B*a*b^9*d^6*e^5*x^...

```

Mupad [B] (verification not implemented)

Time = 1.66 (sec) , antiderivative size = 3577, normalized size of antiderivative = 7.76

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{11} dx = \text{Too large to display}$$

input `int((A + B*x)*(a + b*x)^10*(d + e*x)^11,x)`

output

```
x^5*(42*A*a^6*b^4*d^11 + 24*B*a^7*b^3*d^11 + 66*A*a^10*d^7*e^4 + 33*B*a^10
*d^8*e^3 + 264*A*a^7*b^3*d^10*e + 330*A*a^9*b*d^8*e^3 + 99*B*a^8*b^2*d^10*
e + 110*B*a^9*b*d^9*e^2 + 495*A*a^8*b^2*d^9*e^2) + x^8*(15*A*a^3*b^7*d^11
+ (105*B*a^4*b^6*d^11)/4 + (165*A*a^10*d^4*e^7)/4 + (231*B*a^10*d^5*e^6)/4
+ (1155*A*a^4*b^6*d^10*e)/4 + (1155*A*a^9*b*d^5*e^6)/2 + (693*B*a^5*b^5*d
^10*e)/2 + (1155*B*a^9*b*d^6*e^5)/2 + (3465*A*a^5*b^5*d^9*e^2)/2 + (17325*
A*a^6*b^4*d^8*e^3)/4 + 4950*A*a^7*b^3*d^7*e^4 + (10395*A*a^8*b^2*d^6*e^5)/
4 + (5775*B*a^6*b^4*d^9*e^2)/4 + 2475*B*a^7*b^3*d^8*e^3 + (7425*B*a^8*b^2*
d^7*e^4)/4) + x^19*((120*A*a^3*b^7*e^11)/19 + (210*B*a^4*b^6*e^11)/19 + (1
65*A*b^10*d^3*e^8)/19 + (330*B*b^10*d^4*e^7)/19 + (550*A*a*b^9*d^2*e^9)/19
+ (495*A*a^2*b^8*d*e^10)/19 + (1650*B*a*b^9*d^3*e^8)/19 + (1320*B*a^3*b^7
*d*e^10)/19 + (2475*B*a^2*b^8*d^2*e^9)/19) + x^16*((105*A*a^6*b^4*e^11)/8
+ (15*B*a^7*b^3*e^11)/2 + (231*A*b^10*d^6*e^5)/8 + (165*B*b^10*d^7*e^4)/8
+ (1155*A*a*b^9*d^5*e^6)/4 + (693*A*a^5*b^5*d*e^10)/4 + (1155*B*a*b^9*d^6*
e^5)/4 + (1155*B*a^6*b^4*d*e^10)/8 + (7425*A*a^2*b^8*d^4*e^7)/8 + (2475*A*
a^3*b^7*d^3*e^8)/2 + (5775*A*a^4*b^6*d^2*e^9)/8 + (10395*B*a^2*b^8*d^5*e^6
)/8 + 2475*B*a^3*b^7*d^4*e^7 + (17325*B*a^4*b^6*d^3*e^8)/8 + (3465*B*a^5*b
^5*d^2*e^9)/4) + x^11*((A*b^10*d^11)/11 + (10*B*a*b^9*d^11)/11 + A*a^10*d*
e^10 + 5*B*a^10*d^2*e^9 + 50*A*a^9*b*d^2*e^9 + 45*B*a^2*b^8*d^10*e + 150*B
*a^9*b*d^3*e^8 + 225*A*a^2*b^8*d^9*e^2 + 1800*A*a^3*b^7*d^8*e^3 + 6300*...
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 2205, normalized size of antiderivative = 4.78

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{11} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)*(e*x+d)^11,x)`

output

```
(x*(16224936*a**11*d**11 + 89237148*a**11*d**10*e*x + 297457160*a**11*d**9
**e**2*x**2 + 669278610*a**11*d**8***e**3*x**3 + 1070845776*a**11*d**7***e**4*x
**4 + 1249320072*a**11*d**6***e**5*x**5 + 1070845776*a**11*d**5***e**6*x**6 +
669278610*a**11*d**4***e**7*x**7 + 297457160*a**11*d**3***e**8*x**8 + 89237148
*a**11*d**2***e**9*x**9 + 16224936*a**11*d*e**10*x**10 + 1352078*a**11*e**11
*x**11 + 89237148*a**10*b*d**11*x + 654405752*a**10*b*d**10*e*x**2 + 24540
21570*a**10*b*d**9*e**2*x**3 + 5889651768*a**10*b*d**8***e**3*x**4 + 9816086
280*a**10*b*d**7***e**4*x**5 + 11779303536*a**10*b*d**6***e**5*x**6 + 10306890
594*a**10*b*d**5***e**6*x**7 + 6544057520*a**10*b*d**4***e**7*x**8 + 294482588
4*a**10*b*d**3***e**8*x**9 + 892371480*a**10*b*d**2***e**9*x**10 + 163601438*a
**10*b*d*e**10*x**11 + 13728792*a**10*b*e**11*x**12 + 297457160*a**9*b**2*
d**11*x**2 + 2454021570*a**9*b**2*d**10*e*x**3 + 9816086280*a**9*b**2*d**9
***e**2*x**4 + 2454021570*a**9*b**2*d**8***e**3*x**5 + 42068941200*a**9*b**2*
d**7***e**4*x**6 + 51534452970*a**9*b**2*d**6***e**5*x**7 + 45808402640*a**9*b
**2*d**5***e**6*x**8 + 29448258840*a**9*b**2*d**4***e**7*x**9 + 13385572200*a*
**9*b**2*d**3***e**8*x**10 + 4090035950*a**9*b**2*d**2***e**9*x**11 + 755083560
*a**9*b**2*d*e**10*x**12 + 63740820*a**9*b**2*e**11*x**13 + 669278610*a**8
*b**3*d**11*x**3 + 5889651768*a**8*b**3*d**10*e*x**4 + 24540215700*a**8*b*
**3*d**9***e**2*x**5 + 63103411800*a**8*b**3*d**8***e**3*x**6 + 110430970650*a*
**8*b**3*d**7***e**4*x**7 + 137425207920*a**8*b**3*d**6***e**5*x**8 + 123682...
```

3.68 $\int (a + bx)^{10}(A + Bx)(d + ex)^{10} dx$

Optimal result	689
Mathematica [B] (verified)	690
Rubi [A] (verified)	691
Maple [B] (verified)	693
Fricas [B] (verification not implemented)	694
Sympy [B] (verification not implemented)	695
Maxima [B] (verification not implemented)	696
Giac [B] (verification not implemented)	697
Mupad [B] (verification not implemented)	698
Reduce [B] (verification not implemented)	698

Optimal result

Integrand size = 20, antiderivative size = 460

$$\begin{aligned}
 & \int (a + bx)^{10}(A + Bx)(d + ex)^{10} dx \\
 &= \frac{(Ab - aB)(bd - ae)^{10}(a + bx)^{11}}{11b^{12}} + \frac{(bd - ae)^9(bBd + 10Abe - 11aBe)(a + bx)^{12}}{12b^{12}} \\
 &+ \frac{5e(bd - ae)^8(2bBd + 9Abe - 11aBe)(a + bx)^{13}}{13b^{12}} \\
 &+ \frac{15e^2(bd - ae)^7(3bBd + 8Abe - 11aBe)(a + bx)^{14}}{14b^{12}} \\
 &+ \frac{2e^3(bd - ae)^6(4bBd + 7Abe - 11aBe)(a + bx)^{15}}{b^{12}} \\
 &+ \frac{21e^4(bd - ae)^5(5bBd + 6Abe - 11aBe)(a + bx)^{16}}{8b^{12}} \\
 &+ \frac{42e^5(bd - ae)^4(6bBd + 5Abe - 11aBe)(a + bx)^{17}}{17b^{12}} \\
 &+ \frac{5e^6(bd - ae)^3(7bBd + 4Abe - 11aBe)(a + bx)^{18}}{3b^{12}} \\
 &+ \frac{15e^7(bd - ae)^2(8bBd + 3Abe - 11aBe)(a + bx)^{19}}{19b^{12}} \\
 &+ \frac{e^8(bd - ae)(9bBd + 2Abe - 11aBe)(a + bx)^{20}}{4b^{12}} \\
 &+ \frac{e^9(10bBd + Abe - 11aBe)(a + bx)^{21}}{21b^{12}} + \frac{Be^{10}(a + bx)^{22}}{22b^{12}}
 \end{aligned}$$

output

```

1/11*(A*b-B*a)*(-a*e+b*d)^10*(b*x+a)^11/b^12+1/12*(-a*e+b*d)^9*(10*A*b*e-1
1*B*a*e+B*b*d)*(b*x+a)^12/b^12+5/13*e*(-a*e+b*d)^8*(9*A*b*e-11*B*a*e+2*B*b
*d)*(b*x+a)^13/b^12+15/14*e^2*(-a*e+b*d)^7*(8*A*b*e-11*B*a*e+3*B*b*d)*(b*x
+a)^14/b^12+2*e^3*(-a*e+b*d)^6*(7*A*b*e-11*B*a*e+4*B*b*d)*(b*x+a)^15/b^12+
21/8*e^4*(-a*e+b*d)^5*(6*A*b*e-11*B*a*e+5*B*b*d)*(b*x+a)^16/b^12+42/17*e^5
*(-a*e+b*d)^4*(5*A*b*e-11*B*a*e+6*B*b*d)*(b*x+a)^17/b^12+5/3*e^6*(-a*e+b*d
)^3*(4*A*b*e-11*B*a*e+7*B*b*d)*(b*x+a)^18/b^12+15/19*e^7*(-a*e+b*d)^2*(3*A
*b*e-11*B*a*e+8*B*b*d)*(b*x+a)^19/b^12+1/4*e^8*(-a*e+b*d)*(2*A*b*e-11*B*a*
e+9*B*b*d)*(b*x+a)^20/b^12+1/21*e^9*(A*b*e-11*B*a*e+10*B*b*d)*(b*x+a)^21/b
^12+1/22*B*e^10*(b*x+a)^22/b^12

```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 2815 vs. $2(460) = 920$.

Time = 0.71 (sec) , antiderivative size = 2815, normalized size of antiderivative = 6.12

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{10} dx = \text{Result too large to show}$$

input

```
Integrate[(a + b*x)^10*(A + B*x)*(d + e*x)^10,x]
```

output

```

a^10*A*d^10*x + (a^9*d^9*(a*B*d + 10*A*(b*d + a*e))*x^2)/2 + (5*a^8*d^8*(2
*a*B*d*(b*d + a*e) + A*(9*b^2*d^2 + 20*a*b*d*e + 9*a^2*e^2))*x^3)/3 + (5*a
^7*d^7*(a*B*d*(9*b^2*d^2 + 20*a*b*d*e + 9*a^2*e^2) + 6*A*(4*b^3*d^3 + 15*a
*b^2*d^2*e + 15*a^2*b*d*e^2 + 4*a^3*e^3))*x^4)/4 + 3*a^6*d^6*(2*a*B*d*(4*b
^3*d^3 + 15*a*b^2*d^2*e + 15*a^2*b*d*e^2 + 4*a^3*e^3) + A*(14*b^4*d^4 + 80
*a*b^3*d^3*e + 135*a^2*b^2*d^2*e^2 + 80*a^3*b*d*e^3 + 14*a^4*e^4))*x^5 + (
a^5*d^5*(5*a*B*d*(14*b^4*d^4 + 80*a*b^3*d^3*e + 135*a^2*b^2*d^2*e^2 + 80*a
^3*b*d*e^3 + 14*a^4*e^4) + 4*A*(21*b^5*d^5 + 175*a*b^4*d^4*e + 450*a^2*b^3
*d^3*e^2 + 450*a^3*b^2*d^2*e^3 + 175*a^4*b*d*e^4 + 21*a^5*e^5))*x^6)/2 + (
6*a^4*d^4*(2*a*B*d*(21*b^5*d^5 + 175*a*b^4*d^4*e + 450*a^2*b^3*d^3*e^2 + 4
50*a^3*b^2*d^2*e^3 + 175*a^4*b*d*e^4 + 21*a^5*e^5) + 5*A*(7*b^6*d^6 + 84*a
*b^5*d^5*e + 315*a^2*b^4*d^4*e^2 + 480*a^3*b^3*d^3*e^3 + 315*a^4*b^2*d^2*e
^4 + 84*a^5*b*d*e^5 + 7*a^6*e^6))*x^7)/7 + (15*a^3*d^3*(a*B*d*(7*b^6*d^6 +
84*a*b^5*d^5*e + 315*a^2*b^4*d^4*e^2 + 480*a^3*b^3*d^3*e^3 + 315*a^4*b^2*
d^2*e^4 + 84*a^5*b*d*e^5 + 7*a^6*e^6) + A*(4*b^7*d^7 + 70*a*b^6*d^6*e + 37
8*a^2*b^5*d^5*e^2 + 840*a^3*b^4*d^4*e^3 + 840*a^4*b^3*d^3*e^4 + 378*a^5*b^
2*d^2*e^5 + 70*a^6*b*d*e^6 + 4*a^7*e^7))*x^8)/4 + (5*a^2*d^2*(4*a*B*d*(2*b
^7*d^7 + 35*a*b^6*d^6*e + 189*a^2*b^5*d^5*e^2 + 420*a^3*b^4*d^4*e^3 + 420*
a^4*b^3*d^3*e^4 + 189*a^5*b^2*d^2*e^5 + 35*a^6*b*d*e^6 + 2*a^7*e^7) + A*(3
*b^8*d^8 + 80*a*b^7*d^7*e + 630*a^2*b^6*d^6*e^2 + 2016*a^3*b^5*d^5*e^3 ...

```

Rubi [A] (verified)

Time = 4.13 (sec) , antiderivative size = 460, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^{10} (A + Bx)(d + ex)^{10} dx$$

$$\downarrow 86$$

$$\int \left(\frac{e^9 (a + bx)^{20} (-11aBe + Abe + 10bBd)}{b^{11}} + \frac{5e^8 (a + bx)^{19} (bd - ae) (-11aBe + 2Abe + 9bBd)}{b^{11}} + \frac{15e^7 (a + bx)}{b^{11}} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
& \frac{e^9(a+bx)^{21}(-11aBe + Abe + 10bBd)}{21b^{12}} + \frac{e^8(a+bx)^{20}(bd-ae)(-11aBe + 2Abe + 9bBd)}{4b^{12}} + \\
& \frac{15e^7(a+bx)^{19}(bd-ae)^2(-11aBe + 3Abe + 8bBd)}{19b^{12}} + \\
& \frac{5e^6(a+bx)^{18}(bd-ae)^3(-11aBe + 4Abe + 7bBd)}{3b^{12}} + \\
& \frac{42e^5(a+bx)^{17}(bd-ae)^4(-11aBe + 5Abe + 6bBd)}{17b^{12}} + \\
& \frac{21e^4(a+bx)^{16}(bd-ae)^5(-11aBe + 6Abe + 5bBd)}{8b^{12}} + \\
& \frac{2e^3(a+bx)^{15}(bd-ae)^6(-11aBe + 7Abe + 4bBd)}{b^{12}} + \\
& \frac{15e^2(a+bx)^{14}(bd-ae)^7(-11aBe + 8Abe + 3bBd)}{14b^{12}} + \\
& \frac{5e(a+bx)^{13}(bd-ae)^8(-11aBe + 9Abe + 2bBd)}{13b^{12}} + \\
& \frac{(a+bx)^{12}(bd-ae)^9(-11aBe + 10Abe + bBd)}{12b^{12}} + \frac{(a+bx)^{11}(Ab-aB)(bd-ae)^{10}}{11b^{12}} + \\
& \frac{Be^{10}(a+bx)^{22}}{22b^{12}}
\end{aligned}$$

input `Int[(a + b*x)^10*(A + B*x)*(d + e*x)^10,x]`

output `((A*b - a*B)*(b*d - a*e)^10*(a + b*x)^11)/(11*b^12) + ((b*d - a*e)^9*(b*B*d + 10*A*b*e - 11*a*B*e)*(a + b*x)^12)/(12*b^12) + (5*e*(b*d - a*e)^8*(2*b*B*d + 9*A*b*e - 11*a*B*e)*(a + b*x)^13)/(13*b^12) + (15*e^2*(b*d - a*e)^7*(3*b*B*d + 8*A*b*e - 11*a*B*e)*(a + b*x)^14)/(14*b^12) + (2*e^3*(b*d - a*e)^6*(4*b*B*d + 7*A*b*e - 11*a*B*e)*(a + b*x)^15)/b^12 + (21*e^4*(b*d - a*e)^5*(5*b*B*d + 6*A*b*e - 11*a*B*e)*(a + b*x)^16)/(8*b^12) + (42*e^5*(b*d - a*e)^4*(6*b*B*d + 5*A*b*e - 11*a*B*e)*(a + b*x)^17)/(17*b^12) + (5*e^6*(b*d - a*e)^3*(7*b*B*d + 4*A*b*e - 11*a*B*e)*(a + b*x)^18)/(3*b^12) + (15*e^7*(b*d - a*e)^2*(8*b*B*d + 3*A*b*e - 11*a*B*e)*(a + b*x)^19)/(19*b^12) + (e^8*(b*d - a*e)*(9*b*B*d + 2*A*b*e - 11*a*B*e)*(a + b*x)^20)/(4*b^12) + (e^9*(10*b*B*d + A*b*e - 11*a*B*e)*(a + b*x)^21)/(21*b^12) + (B*e^10*(a + b*x)^22)/(22*b^12)`

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0]) || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3040 vs. $2(438) = 876$.

Time = 0.29 (sec) , antiderivative size = 3041, normalized size of antiderivative = 6.61

method	result	size
default	Expression too large to display	3041
norman	Expression too large to display	3293
orering	Expression too large to display	3912
gospers	Expression too large to display	3913
risch	Expression too large to display	3913
parallelrisc	Expression too large to display	3913

input

```
int((b*x+a)^10*(B*x+A)*(e*x+d)^10,x,method=_RETURNVERBOSE)
```

output

```

1/22*b^10*B*e^10*x^22+1/21*((A*b^10+10*B*a*b^9)*e^10+10*b^10*B*d*e^9)*x^21
+1/20*((10*A*a*b^9+45*B*a^2*b^8)*e^10+10*(A*b^10+10*B*a*b^9)*d*e^9+45*b^10
*B*d^2*e^8)*x^20+1/19*((45*A*a^2*b^8+120*B*a^3*b^7)*e^10+10*(10*A*a*b^9+45
*B*a^2*b^8)*d*e^9+45*(A*b^10+10*B*a*b^9)*d^2*e^8+120*b^10*B*d^3*e^7)*x^19+
1/18*((120*A*a^3*b^7+210*B*a^4*b^6)*e^10+10*(45*A*a^2*b^8+120*B*a^3*b^7)*d
*e^9+45*(10*A*a*b^9+45*B*a^2*b^8)*d^2*e^8+120*(A*b^10+10*B*a*b^9)*d^3*e^7+
210*b^10*B*d^4*e^6)*x^18+1/17*((210*A*a^4*b^6+252*B*a^5*b^5)*e^10+10*(120*
A*a^3*b^7+210*B*a^4*b^6)*d*e^9+45*(45*A*a^2*b^8+120*B*a^3*b^7)*d^2*e^8+120
*(10*A*a*b^9+45*B*a^2*b^8)*d^3*e^7+210*(A*b^10+10*B*a*b^9)*d^4*e^6+252*b^1
0*B*d^5*e^5)*x^17+1/16*((252*A*a^5*b^5+210*B*a^6*b^4)*e^10+10*(210*A*a^4*b
^6+252*B*a^5*b^5)*d*e^9+45*(120*A*a^3*b^7+210*B*a^4*b^6)*d^2*e^8+120*(45*A
*a^2*b^8+120*B*a^3*b^7)*d^3*e^7+210*(10*A*a*b^9+45*B*a^2*b^8)*d^4*e^6+252*
(A*b^10+10*B*a*b^9)*d^5*e^5+210*b^10*B*d^6*e^4)*x^16+1/15*((210*A*a^6*b^4+
120*B*a^7*b^3)*e^10+10*(252*A*a^5*b^5+210*B*a^6*b^4)*d*e^9+45*(210*A*a^4*b
^6+252*B*a^5*b^5)*d^2*e^8+120*(120*A*a^3*b^7+210*B*a^4*b^6)*d^3*e^7+210*(4
5*A*a^2*b^8+120*B*a^3*b^7)*d^4*e^6+252*(10*A*a*b^9+45*B*a^2*b^8)*d^5*e^5+2
10*(A*b^10+10*B*a*b^9)*d^6*e^4+120*b^10*B*d^7*e^3)*x^15+1/14*((120*A*a^7*b
^3+45*B*a^8*b^2)*e^10+10*(210*A*a^6*b^4+120*B*a^7*b^3)*d*e^9+45*(252*A*a^5
*b^5+210*B*a^6*b^4)*d^2*e^8+120*(210*A*a^4*b^6+252*B*a^5*b^5)*d^3*e^7+210*
(120*A*a^3*b^7+210*B*a^4*b^6)*d^4*e^6+252*(45*A*a^2*b^8+120*B*a^3*b^7)*...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3048 vs. $2(438) = 876$.

Time = 0.13 (sec) , antiderivative size = 3048, normalized size of antiderivative = 6.63

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{10} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^10,x, algorithm="fricas")
```

output

```
Too large to include
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3936 vs. $2(478) = 956$.

Time = 0.25 (sec) , antiderivative size = 3936, normalized size of antiderivative = 8.56

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{10} dx = \text{Too large to display}$$

input `integrate((b*x+a)**10*(B*x+A)*(e*x+d)**10,x)`

output

```
A*a**10*d**10*x + B*b**10*e**10*x**22/22 + x**21*(A*b**10*e**10/21 + 10*B*
a*b**9*e**10/21 + 10*B*b**10*d*e**9/21) + x**20*(A*a*b**9*e**10/2 + A*b**1
0*d*e**9/2 + 9*B*a**2*b**8*e**10/4 + 5*B*a*b**9*d*e**9 + 9*B*b**10*d**2*e
**8/4) + x**19*(45*A*a**2*b**8*e**10/19 + 100*A*a*b**9*d*e**9/19 + 45*A*b**
10*d**2*e**8/19 + 120*B*a**3*b**7*e**10/19 + 450*B*a**2*b**8*d*e**9/19 + 4
50*B*a*b**9*d**2*e**8/19 + 120*B*b**10*d**3*e**7/19) + x**18*(20*A*a**3*b*
**7*e**10/3 + 25*A*a**2*b**8*d*e**9 + 25*A*a*b**9*d**2*e**8 + 20*A*b**10*d*
**3*e**7/3 + 35*B*a**4*b**6*e**10/3 + 200*B*a**3*b**7*d*e**9/3 + 225*B*a**2
*b**8*d**2*e**8/2 + 200*B*a*b**9*d**3*e**7/3 + 35*B*b**10*d**4*e**6/3) + x
**17*(210*A*a**4*b**6*e**10/17 + 1200*A*a**3*b**7*d*e**9/17 + 2025*A*a**2*
b**8*d**2*e**8/17 + 1200*A*a*b**9*d**3*e**7/17 + 210*A*b**10*d**4*e**6/17
+ 252*B*a**5*b**5*e**10/17 + 2100*B*a**4*b**6*d*e**9/17 + 5400*B*a**3*b**7
*d**2*e**8/17 + 5400*B*a**2*b**8*d**3*e**7/17 + 2100*B*a*b**9*d**4*e**6/17
+ 252*B*b**10*d**5*e**5/17) + x**16*(63*A*a**5*b**5*e**10/4 + 525*A*a**4*
b**6*d*e**9/4 + 675*A*a**3*b**7*d**2*e**8/2 + 675*A*a**2*b**8*d**3*e**7/2
+ 525*A*a*b**9*d**4*e**6/4 + 63*A*b**10*d**5*e**5/4 + 105*B*a**6*b**4*e**1
0/8 + 315*B*a**5*b**5*d*e**9/2 + 4725*B*a**4*b**6*d**2*e**8/8 + 900*B*a**3
*b**7*d**3*e**7 + 4725*B*a**2*b**8*d**4*e**6/8 + 315*B*a*b**9*d**5*e**5/2
+ 105*B*b**10*d**6*e**4/8) + x**15*(14*A*a**6*b**4*e**10 + 168*A*a**5*b**5
*d*e**9 + 630*A*a**4*b**6*d**2*e**8 + 960*A*a**3*b**7*d**3*e**7 + 630*A...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3048 vs. $2(438) = 876$.

Time = 0.06 (sec) , antiderivative size = 3048, normalized size of antiderivative = 6.63

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{10} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)*(e*x+d)^10,x, algorithm="maxima")`

output

```

1/22*B*b^10*e^10*x^22 + A*a^10*d^10*x + 1/21*(10*B*b^10*d*e^9 + (10*B*a*b^
9 + A*b^10)*e^10)*x^21 + 1/4*(9*B*b^10*d^2*e^8 + 2*(10*B*a*b^9 + A*b^10)*d
*e^9 + (9*B*a^2*b^8 + 2*A*a*b^9)*e^10)*x^20 + 5/19*(24*B*b^10*d^3*e^7 + 9*
(10*B*a*b^9 + A*b^10)*d^2*e^8 + 10*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^9 + 3*(8*
B*a^3*b^7 + 3*A*a^2*b^8)*e^10)*x^19 + 5/6*(14*B*b^10*d^4*e^6 + 8*(10*B*a*b
^9 + A*b^10)*d^3*e^7 + 15*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^8 + 10*(8*B*a^3*
b^7 + 3*A*a^2*b^8)*d*e^9 + 2*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^10)*x^18 + 3/17
*(84*B*b^10*d^5*e^5 + 70*(10*B*a*b^9 + A*b^10)*d^4*e^6 + 200*(9*B*a^2*b^8
+ 2*A*a*b^9)*d^3*e^7 + 225*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^8 + 100*(7*B*
a^4*b^6 + 4*A*a^3*b^7)*d*e^9 + 14*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^10)*x^17 +
3/8*(35*B*b^10*d^6*e^4 + 42*(10*B*a*b^9 + A*b^10)*d^5*e^5 + 175*(9*B*a^2*
b^8 + 2*A*a*b^9)*d^4*e^6 + 300*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^7 + 225*(
7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^8 + 70*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^9
+ 7*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^10)*x^16 + 2*(4*B*b^10*d^7*e^3 + 7*(10*B
*a*b^9 + A*b^10)*d^6*e^4 + 42*(9*B*a^2*b^8 + 2*A*a*b^9)*d^5*e^5 + 105*(8*B
*a^3*b^7 + 3*A*a^2*b^8)*d^4*e^6 + 120*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^3*e^7
+ 63*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e^8 + 14*(5*B*a^6*b^4 + 6*A*a^5*b^5)*
d*e^9 + (4*B*a^7*b^3 + 7*A*a^6*b^4)*e^10)*x^15 + 15/14*(3*B*b^10*d^8*e^2 +
8*(10*B*a*b^9 + A*b^10)*d^7*e^3 + 70*(9*B*a^2*b^8 + 2*A*a*b^9)*d^6*e^4 +
252*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^5*e^5 + 420*(7*B*a^4*b^6 + 4*A*a^3*b^...

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3912 vs. $2(438) = 876$.

Time = 0.13 (sec) , antiderivative size = 3912, normalized size of antiderivative = 8.50

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{10} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)*(e*x+d)^10,x, algorithm="giac")`

output

```
1/22*B*b^10*e^10*x^22 + 10/21*B*b^10*d*e^9*x^21 + 10/21*B*a*b^9*e^10*x^21
+ 1/21*A*b^10*e^10*x^21 + 9/4*B*b^10*d^2*e^8*x^20 + 5*B*a*b^9*d*e^9*x^20 +
1/2*A*b^10*d*e^9*x^20 + 9/4*B*a^2*b^8*e^10*x^20 + 1/2*A*a*b^9*e^10*x^20 +
120/19*B*b^10*d^3*e^7*x^19 + 450/19*B*a*b^9*d^2*e^8*x^19 + 45/19*A*b^10*d
^2*e^8*x^19 + 450/19*B*a^2*b^8*d*e^9*x^19 + 100/19*A*a*b^9*d*e^9*x^19 + 12
0/19*B*a^3*b^7*e^10*x^19 + 45/19*A*a^2*b^8*e^10*x^19 + 35/3*B*b^10*d^4*e^6
*x^18 + 200/3*B*a*b^9*d^3*e^7*x^18 + 20/3*A*b^10*d^3*e^7*x^18 + 225/2*B*a^
2*b^8*d^2*e^8*x^18 + 25*A*a*b^9*d^2*e^8*x^18 + 200/3*B*a^3*b^7*d*e^9*x^18
+ 25*A*a^2*b^8*d*e^9*x^18 + 35/3*B*a^4*b^6*e^10*x^18 + 20/3*A*a^3*b^7*e^10
*x^18 + 252/17*B*b^10*d^5*e^5*x^17 + 2100/17*B*a*b^9*d^4*e^6*x^17 + 210/17
*A*b^10*d^4*e^6*x^17 + 5400/17*B*a^2*b^8*d^3*e^7*x^17 + 1200/17*A*a*b^9*d^
3*e^7*x^17 + 5400/17*B*a^3*b^7*d^2*e^8*x^17 + 2025/17*A*a^2*b^8*d^2*e^8*x^
17 + 2100/17*B*a^4*b^6*d*e^9*x^17 + 1200/17*A*a^3*b^7*d*e^9*x^17 + 252/17*
B*a^5*b^5*e^10*x^17 + 210/17*A*a^4*b^6*e^10*x^17 + 105/8*B*b^10*d^6*e^4*x^
16 + 315/2*B*a*b^9*d^5*e^5*x^16 + 63/4*A*b^10*d^5*e^5*x^16 + 4725/8*B*a^2*
b^8*d^4*e^6*x^16 + 525/4*A*a*b^9*d^4*e^6*x^16 + 900*B*a^3*b^7*d^3*e^7*x^16
+ 675/2*A*a^2*b^8*d^3*e^7*x^16 + 4725/8*B*a^4*b^6*d^2*e^8*x^16 + 675/2*A*
a^3*b^7*d^2*e^8*x^16 + 315/2*B*a^5*b^5*d*e^9*x^16 + 525/4*A*a^4*b^6*d*e^9*
x^16 + 105/8*B*a^6*b^4*e^10*x^16 + 63/4*A*a^5*b^5*e^10*x^16 + 8*B*b^10*d^7
*e^3*x^15 + 140*B*a*b^9*d^6*e^4*x^15 + 14*A*b^10*d^6*e^4*x^15 + 756*B*a...
```

Mupad [B] (verification not implemented)

Time = 1.64 (sec) , antiderivative size = 3262, normalized size of antiderivative = 7.09

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{10} dx = \text{Too large to display}$$

input `int((A + B*x)*(a + b*x)^10*(d + e*x)^10,x)`

output

```
x^11*((A*a^10*e^10)/11 + (A*b^10*d^10)/11 + (10*B*a*b^9*d^10)/11 + (10*B*a^10*d*e^9)/11 + (450*B*a^2*b^8*d^9*e)/11 + (450*B*a^9*b*d^2*e^8)/11 + (2025*A*a^2*b^8*d^8*e^2)/11 + (14400*A*a^3*b^7*d^7*e^3)/11 + (44100*A*a^4*b^6*d^6*e^4)/11 + (63504*A*a^5*b^5*d^5*e^5)/11 + (44100*A*a^6*b^4*d^4*e^6)/11 + (14400*A*a^7*b^3*d^3*e^7)/11 + (2025*A*a^8*b^2*d^2*e^8)/11 + (5400*B*a^3*b^7*d^8*e^2)/11 + (25200*B*a^4*b^6*d^7*e^3)/11 + (52920*B*a^5*b^5*d^6*e^4)/11 + (52920*B*a^6*b^4*d^5*e^5)/11 + (25200*B*a^7*b^3*d^4*e^6)/11 + (5400*B*a^8*b^2*d^3*e^7)/11 + (100*A*a*b^9*d^9*e)/11 + (100*A*a^9*b*d*e^9)/11 + x^5*(42*A*a^6*b^4*d^10 + 24*B*a^7*b^3*d^10 + 42*A*a^10*d^6*e^4 + 24*B*a^10*d^7*e^3 + 240*A*a^7*b^3*d^9*e + 240*A*a^9*b*d^7*e^3 + 90*B*a^8*b^2*d^9*e + 90*B*a^9*b*d^8*e^2 + 405*A*a^8*b^2*d^8*e^2) + x^8*(15*A*a^3*b^7*d^10 + (105*B*a^4*b^6*d^10)/4 + 15*A*a^10*d^3*e^7 + (105*B*a^10*d^4*e^6)/4 + (525*A*a^4*b^6*d^9*e)/2 + (525*A*a^9*b*d^4*e^6)/2 + 315*B*a^5*b^5*d^9*e + 315*B*a^9*b*d^5*e^5 + (2835*A*a^5*b^5*d^8*e^2)/2 + 3150*A*a^6*b^4*d^7*e^3 + 3150*A*a^7*b^3*d^6*e^4 + (2835*A*a^8*b^2*d^5*e^5)/2 + (4725*B*a^6*b^4*d^8*e^2)/4 + 1800*B*a^7*b^3*d^7*e^3 + (4725*B*a^8*b^2*d^6*e^4)/4) + x^12*((B*a^10*e^10)/12 + (B*b^10*d^10)/12 + (5*A*a^9*b*e^10)/6 + (5*A*b^10*d^9*e)/6 + (75*A*a*b^9*d^8*e^2)/2 + (75*A*a^8*b^2*d*e^9)/2 + 450*A*a^2*b^8*d^7*e^3 + 2100*A*a^3*b^7*d^6*e^4 + 4410*A*a^4*b^6*d^5*e^5 + 4410*A*a^5*b^5*d^4*e^6 + 2100*A*a^6*b^4*d^3*e^7 + 450*A*a^7*b^3*d^2*e^8 + (675*B*a^2*b^8*d^8*e...
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 2011, normalized size of antiderivative = 4.37

$$\int (a + bx)^{10}(A + Bx)(d + ex)^{10} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)*(e*x+d)^10,x)`

output

```
(x*(7759752*a**11*d**10 + 38798760*a**11*d**9*e*x + 116396280*a**11*d**8*e
**2*x**2 + 232792560*a**11*d**7*e**3*x**3 + 325909584*a**11*d**6*e**4*x**4
+ 325909584*a**11*d**5*e**5*x**5 + 232792560*a**11*d**4*e**6*x**6 + 11639
6280*a**11*d**3*e**7*x**7 + 38798760*a**11*d**2*e**8*x**8 + 7759752*a**11*
d*e**9*x**9 + 705432*a**11*e**10*x**10 + 42678636*a**10*b*d**10*x + 284524
240*a**10*b*d**9*e*x**2 + 960269310*a**10*b*d**8*e**2*x**3 + 2048574528*a*
*10*b*d**7*e**3*x**4 + 2987504520*a**10*b*d**6*e**4*x**5 + 3072861792*a**1
0*b*d**5*e**5*x**6 + 2240628390*a**10*b*d**4*e**6*x**7 + 1138096960*a**10*
b*d**3*e**7*x**8 + 384107724*a**10*b*d**2*e**8*x**9 + 77597520*a**10*b*d*e
**9*x**10 + 7113106*a**10*b*e**10*x**11 + 142262120*a**9*b**2*d**10*x**2 +
1066965900*a**9*b**2*d**9*e*x**3 + 3841077240*a**9*b**2*d**8*e**2*x**4 +
8535727200*a**9*b**2*d**7*e**3*x**5 + 12803590800*a**9*b**2*d**6*e**4*x**6
+ 13443770340*a**9*b**2*d**5*e**5*x**7 + 9958348400*a**9*b**2*d**4*e**6*x
**8 + 5121436320*a**9*b**2*d**3*e**7*x**9 + 1745944200*a**9*b**2*d**2*e**8
*x**10 + 355655300*a**9*b**2*d*e**9*x**11 + 32829720*a**9*b**2*e**10*x**12
+ 320089770*a**8*b**3*d**10*x**3 + 2560718160*a**8*b**3*d**9*e*x**4 + 960
2693100*a**8*b**3*d**8*e**2*x**5 + 21949012800*a**8*b**3*d**7*e**3*x**6 +
33609425850*a**8*b**3*d**6*e**4*x**7 + 35850054240*a**8*b**3*d**5*e**5*x**
8 + 26887540680*a**8*b**3*d**4*e**6*x**9 + 13967553600*a**8*b**3*d**3*e**7
*x**10 + 4801346550*a**8*b**3*d**2*e**8*x**11 + 984891600*a**8*b**3*d*e...
```


3.69 $\int (a + bx)^{10}(A + Bx)(d + ex)^9 dx$

Optimal result	700
Mathematica [B] (verified)	701
Rubi [A] (verified)	702
Maple [B] (verified)	704
Fricas [B] (verification not implemented)	705
Sympy [B] (verification not implemented)	706
Maxima [B] (verification not implemented)	707
Giac [B] (verification not implemented)	708
Mupad [B] (verification not implemented)	709
Reduce [B] (verification not implemented)	710

Optimal result

Integrand size = 20, antiderivative size = 415

$$\begin{aligned}
 \int (a + bx)^{10}(A + Bx)(d + ex)^9 dx = & \frac{(Ab - aB)(bd - ae)^9(a + bx)^{11}}{11b^{11}} \\
 & + \frac{(bd - ae)^8(bBd + 9Abe - 10aBe)(a + bx)^{12}}{12b^{11}} \\
 & + \frac{9e(bd - ae)^7(bBd + 4Abe - 5aBe)(a + bx)^{13}}{13b^{11}} \\
 & + \frac{6e^2(bd - ae)^6(3bBd + 7Abe - 10aBe)(a + bx)^{14}}{7b^{11}} \\
 & + \frac{14e^3(bd - ae)^5(2bBd + 3Abe - 5aBe)(a + bx)^{15}}{5b^{11}} \\
 & + \frac{63e^4(bd - ae)^4(bBd + Abe - 2aBe)(a + bx)^{16}}{8b^{11}} \\
 & + \frac{42e^5(bd - ae)^3(3bBd + 2Abe - 5aBe)(a + bx)^{17}}{17b^{11}} \\
 & + \frac{2e^6(bd - ae)^2(7bBd + 3Abe - 10aBe)(a + bx)^{18}}{3b^{11}} \\
 & + \frac{9e^7(bd - ae)(4bBd + Abe - 5aBe)(a + bx)^{19}}{19b^{11}} \\
 & + \frac{e^8(9bBd + Abe - 10aBe)(a + bx)^{20}}{20b^{11}} \\
 & + \frac{Be^9(a + bx)^{21}}{21b^{11}}
 \end{aligned}$$

output

```

1/11*(A*b-B*a)*(-a*e+b*d)^9*(b*x+a)^11/b^11+1/12*(-a*e+b*d)^8*(9*A*b*e-10*
B*a*e+B*b*d)*(b*x+a)^12/b^11+9/13*e*(-a*e+b*d)^7*(4*A*b*e-5*B*a*e+B*b*d)*
(b*x+a)^13/b^11+6/7*e^2*(-a*e+b*d)^6*(7*A*b*e-10*B*a*e+3*B*b*d)*(b*x+a)^14/
b^11+14/5*e^3*(-a*e+b*d)^5*(3*A*b*e-5*B*a*e+2*B*b*d)*(b*x+a)^15/b^11+63/8*
e^4*(-a*e+b*d)^4*(A*b*e-2*B*a*e+B*b*d)*(b*x+a)^16/b^11+42/17*e^5*(-a*e+b*d
)^3*(2*A*b*e-5*B*a*e+3*B*b*d)*(b*x+a)^17/b^11+2/3*e^6*(-a*e+b*d)^2*(3*A*b*
e-10*B*a*e+7*B*b*d)*(b*x+a)^18/b^11+9/19*e^7*(-a*e+b*d)*(A*b*e-5*B*a*e+4*B
*b*d)*(b*x+a)^19/b^11+1/20*e^8*(A*b*e-10*B*a*e+9*B*b*d)*(b*x+a)^20/b^11+1/
21*B*e^9*(b*x+a)^21/b^11

```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 2553 vs. $2(415) = 830$.

Time = 0.62 (sec) , antiderivative size = 2553, normalized size of antiderivative = 6.15

$$\int (a + bx)^{10}(A + Bx)(d + ex)^9 dx = \text{Result too large to show}$$

input

```
Integrate[(a + b*x)^10*(A + B*x)*(d + e*x)^9,x]
```

output

```

a^10*A*d^9*x + (a^9*d^8*(10*A*b*d + a*B*d + 9*a*A*e)*x^2)/2 + (a^8*d^7*(a*
B*d*(10*b*d + 9*a*e) + 9*A*(5*b^2*d^2 + 10*a*b*d*e + 4*a^2*e^2))*x^3)/3 +
(3*a^7*d^6*(3*a*B*d*(5*b^2*d^2 + 10*a*b*d*e + 4*a^2*e^2) + A*(40*b^3*d^3 +
135*a*b^2*d^2*e + 120*a^2*b*d*e^2 + 28*a^3*e^3))*x^4)/4 + (3*a^6*d^5*(a*B
*d*(40*b^3*d^3 + 135*a*b^2*d^2*e + 120*a^2*b*d*e^2 + 28*a^3*e^3) + A*(70*b
^4*d^4 + 360*a*b^3*d^3*e + 540*a^2*b^2*d^2*e^2 + 280*a^3*b*d*e^3 + 42*a^4*
e^4))*x^5)/5 + a^5*d^4*(a*B*d*(35*b^4*d^4 + 180*a*b^3*d^3*e + 270*a^2*b^2*
d^2*e^2 + 140*a^3*b*d*e^3 + 21*a^4*e^4) + 3*A*(14*b^5*d^5 + 105*a*b^4*d^4*
e + 240*a^2*b^3*d^3*e^2 + 210*a^3*b^2*d^2*e^3 + 70*a^4*b*d*e^4 + 7*a^5*e^5
))*x^6 + (6*a^4*d^3*(3*a*B*d*(14*b^5*d^5 + 105*a*b^4*d^4*e + 240*a^2*b^3*d
^3*e^2 + 210*a^3*b^2*d^2*e^3 + 70*a^4*b*d*e^4 + 7*a^5*e^5) + 7*A*(5*b^6*d
^6 + 54*a*b^5*d^5*e + 180*a^2*b^4*d^4*e^2 + 240*a^3*b^3*d^3*e^3 + 135*a^4*b
^2*d^2*e^4 + 30*a^5*b*d*e^5 + 2*a^6*e^6))*x^7)/7 + (3*a^3*d^2*(7*a*B*d*(5*
b^6*d^6 + 54*a*b^5*d^5*e + 180*a^2*b^4*d^4*e^2 + 240*a^3*b^3*d^3*e^3 + 135
*a^4*b^2*d^2*e^4 + 30*a^5*b*d*e^5 + 2*a^6*e^6) + A*(20*b^7*d^7 + 315*a*b^6
*d^6*e + 1512*a^2*b^5*d^5*e^2 + 2940*a^3*b^4*d^4*e^3 + 2520*a^4*b^3*d^3*e^
4 + 945*a^5*b^2*d^2*e^5 + 140*a^6*b*d*e^6 + 6*a^7*e^7))*x^8)/4 + (a^2*d*(2
*a*B*d*(20*b^7*d^7 + 315*a*b^6*d^6*e + 1512*a^2*b^5*d^5*e^2 + 2940*a^3*b^4
*d^4*e^3 + 2520*a^4*b^3*d^3*e^4 + 945*a^5*b^2*d^2*e^5 + 140*a^6*b*d*e^6 +
6*a^7*e^7) + 3*A*(5*b^8*d^8 + 120*a*b^7*d^7*e + 840*a^2*b^6*d^6*e^2 + 2...

```

Rubi [A] (verified)

Time = 3.40 (sec) , antiderivative size = 415, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^{10} (A + Bx)(d + ex)^9 dx$$

$$\downarrow 86$$

$$\int \left(\frac{e^8 (a + bx)^{19} (-10aBe + Abe + 9bBd)}{b^{10}} + \frac{9e^7 (a + bx)^{18} (bd - ae) (-5aBe + Abe + 4bBd)}{b^{10}} + \frac{12e^6 (a + bx)^{17} (bd - ae)^2}{b^{10}} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
& \frac{e^8(a+bx)^{20}(-10aBe + Abe + 9bBd)}{20b^{11}} + \frac{9e^7(a+bx)^{19}(bd-ae)(-5aBe + Abe + 4bBd)}{19b^{11}} + \\
& \frac{2e^6(a+bx)^{18}(bd-ae)^2(-10aBe + 3Abe + 7bBd)}{2e^6(a+bx)^{18}(bd-ae)^2(-10aBe + 3Abe + 7bBd)} + \\
& \frac{42e^5(a+bx)^{17}(bd-ae)^3(-5aBe + 2Abe + 3bBd)}{3b^{11}} + \\
& \frac{63e^4(a+bx)^{16}(bd-ae)^4(-2aBe + Abe + bBd)}{17b^{11}} + \\
& \frac{14e^3(a+bx)^{15}(bd-ae)^5(-5aBe + 3Abe + 2bBd)}{8b^{11}} + \\
& \frac{6e^2(a+bx)^{14}(bd-ae)^6(-10aBe + 7Abe + 3bBd)}{5b^{11}} + \\
& \frac{9e(a+bx)^{13}(bd-ae)^7(-5aBe + 4Abe + bBd)}{7b^{11}} + \\
& \frac{(a+bx)^{12}(bd-ae)^8(-10aBe + 9Abe + bBd)}{12b^{11}} + \frac{(a+bx)^{11}(Ab-aB)(bd-ae)^9}{11b^{11}} + \\
& \frac{Be^9(a+bx)^{21}}{21b^{11}}
\end{aligned}$$

input `Int[(a + b*x)^10*(A + B*x)*(d + e*x)^9,x]`

output `((A*b - a*B)*(b*d - a*e)^9*(a + b*x)^11)/(11*b^11) + ((b*d - a*e)^8*(b*B*d + 9*A*b*e - 10*a*B*e)*(a + b*x)^12)/(12*b^11) + (9*e*(b*d - a*e)^7*(b*B*d + 4*A*b*e - 5*a*B*e)*(a + b*x)^13)/(13*b^11) + (6*e^2*(b*d - a*e)^6*(3*b*B*d + 7*A*b*e - 10*a*B*e)*(a + b*x)^14)/(7*b^11) + (14*e^3*(b*d - a*e)^5*(2*b*B*d + 3*A*b*e - 5*a*B*e)*(a + b*x)^15)/(5*b^11) + (63*e^4*(b*d - a*e)^4*(b*B*d + A*b*e - 2*a*B*e)*(a + b*x)^16)/(8*b^11) + (42*e^5*(b*d - a*e)^3*(3*b*B*d + 2*A*b*e - 5*a*B*e)*(a + b*x)^17)/(17*b^11) + (2*e^6*(b*d - a*e)^2*(7*b*B*d + 3*A*b*e - 10*a*B*e)*(a + b*x)^18)/(3*b^11) + (9*e^7*(b*d - a*e)*(4*b*B*d + A*b*e - 5*a*B*e)*(a + b*x)^19)/(19*b^11) + (e^8*(9*b*B*d + A*b*e - 10*a*B*e)*(a + b*x)^20)/(20*b^11) + (B*e^9*(a + b*x)^21)/(21*b^11)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2756 vs. $2(393) = 786$.

Time = 0.27 (sec) , antiderivative size = 2757, normalized size of antiderivative = 6.64

method	result	size
default	Expression too large to display	2757
norman	Expression too large to display	2978
orering	Expression too large to display	3536
gospers	Expression too large to display	3537
risch	Expression too large to display	3537
parallelsch	Expression too large to display	3537

input `int((b*x+a)^10*(B*x+A)*(e*x+d)^9,x,method=_RETURNVERBOSE)`

output

```

1/21*b^10*B*e^9*x^21+1/20*((A*b^10+10*B*a*b^9)*e^9+9*b^10*B*d*e^8)*x^20+1/
19*((10*A*a*b^9+45*B*a^2*b^8)*e^9+9*(A*b^10+10*B*a*b^9)*d*e^8+36*b^10*B*d^
2*e^7)*x^19+1/18*((45*A*a^2*b^8+120*B*a^3*b^7)*e^9+9*(10*A*a*b^9+45*B*a^2*
b^8)*d*e^8+36*(A*b^10+10*B*a*b^9)*d^2*e^7+84*b^10*B*d^3*e^6)*x^18+1/17*((1
20*A*a^3*b^7+210*B*a^4*b^6)*e^9+9*(45*A*a^2*b^8+120*B*a^3*b^7)*d*e^8+36*(1
0*A*a*b^9+45*B*a^2*b^8)*d^2*e^7+84*(A*b^10+10*B*a*b^9)*d^3*e^6+126*b^10*B*
d^4*e^5)*x^17+1/16*((210*A*a^4*b^6+252*B*a^5*b^5)*e^9+9*(120*A*a^3*b^7+210
*B*a^4*b^6)*d*e^8+36*(45*A*a^2*b^8+120*B*a^3*b^7)*d^2*e^7+84*(10*A*a*b^9+4
5*B*a^2*b^8)*d^3*e^6+126*(A*b^10+10*B*a*b^9)*d^4*e^5+126*b^10*B*d^5*e^4)*x
^16+1/15*((252*A*a^5*b^5+210*B*a^6*b^4)*e^9+9*(210*A*a^4*b^6+252*B*a^5*b^5
)*d*e^8+36*(120*A*a^3*b^7+210*B*a^4*b^6)*d^2*e^7+84*(45*A*a^2*b^8+120*B*a^
3*b^7)*d^3*e^6+126*(10*A*a*b^9+45*B*a^2*b^8)*d^4*e^5+126*(A*b^10+10*B*a*b^
9)*d^5*e^4+84*b^10*B*d^6*e^3)*x^15+1/14*((210*A*a^6*b^4+120*B*a^7*b^3)*e^9
+9*(252*A*a^5*b^5+210*B*a^6*b^4)*d*e^8+36*(210*A*a^4*b^6+252*B*a^5*b^5)*d^
2*e^7+84*(120*A*a^3*b^7+210*B*a^4*b^6)*d^3*e^6+126*(45*A*a^2*b^8+120*B*a^3
*b^7)*d^4*e^5+126*(10*A*a*b^9+45*B*a^2*b^8)*d^5*e^4+84*(A*b^10+10*B*a*b^9)
*d^6*e^3+36*b^10*B*d^7*e^2)*x^14+1/13*((120*A*a^7*b^3+45*B*a^8*b^2)*e^9+9*
(210*A*a^6*b^4+120*B*a^7*b^3)*d*e^8+36*(252*A*a^5*b^5+210*B*a^6*b^4)*d^2*e
^7+84*(210*A*a^4*b^6+252*B*a^5*b^5)*d^3*e^6+126*(120*A*a^3*b^7+210*B*a^4*b
^6)*d^4*e^5+126*(45*A*a^2*b^8+120*B*a^3*b^7)*d^5*e^4+84*(10*A*a*b^9+45*...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2771 vs. $2(393) = 786$.

Time = 0.10 (sec) , antiderivative size = 2771, normalized size of antiderivative = 6.68

$$\int (a + bx)^{10}(A + Bx)(d + ex)^9 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^9,x, algorithm="fricas")
```

output

```

1/21*B*b^10*e^9*x^21 + A*a^10*d^9*x + 1/20*(9*B*b^10*d*e^8 + (10*B*a*b^9 +
A*b^10)*e^9)*x^20 + 1/19*(36*B*b^10*d^2*e^7 + 9*(10*B*a*b^9 + A*b^10)*d*e
^8 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*e^9)*x^19 + 1/6*(28*B*b^10*d^3*e^6 + 12*(
10*B*a*b^9 + A*b^10)*d^2*e^7 + 15*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^8 + 5*(8*B
*a^3*b^7 + 3*A*a^2*b^8)*e^9)*x^18 + 3/17*(42*B*b^10*d^4*e^5 + 28*(10*B*a*b
^9 + A*b^10)*d^3*e^6 + 60*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^7 + 45*(8*B*a^3*
b^7 + 3*A*a^2*b^8)*d*e^8 + 10*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^9)*x^17 + 3/8*
(21*B*b^10*d^5*e^4 + 21*(10*B*a*b^9 + A*b^10)*d^4*e^5 + 70*(9*B*a^2*b^8 +
2*A*a*b^9)*d^3*e^6 + 90*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^7 + 45*(7*B*a^4*
b^6 + 4*A*a^3*b^7)*d*e^8 + 7*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^9)*x^16 + 2/5*(
14*B*b^10*d^6*e^3 + 21*(10*B*a*b^9 + A*b^10)*d^5*e^4 + 105*(9*B*a^2*b^8 +
2*A*a*b^9)*d^4*e^5 + 210*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^6 + 180*(7*B*a^
4*b^6 + 4*A*a^3*b^7)*d^2*e^7 + 63*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^8 + 7*(5
*B*a^6*b^4 + 6*A*a^5*b^5)*e^9)*x^15 + 3/7*(6*B*b^10*d^7*e^2 + 14*(10*B*a*b
^9 + A*b^10)*d^6*e^3 + 105*(9*B*a^2*b^8 + 2*A*a*b^9)*d^5*e^4 + 315*(8*B*a^
3*b^7 + 3*A*a^2*b^8)*d^4*e^5 + 420*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^3*e^6 + 2
52*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e^7 + 63*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d*
e^8 + 5*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e^9)*x^14 + 3/13*(3*B*b^10*d^8*e + 12*
(10*B*a*b^9 + A*b^10)*d^7*e^2 + 140*(9*B*a^2*b^8 + 2*A*a*b^9)*d^6*e^3 + 63
0*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^5*e^4 + 1260*(7*B*a^4*b^6 + 4*A*a^3*b^7...

```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3541 vs. $2(428) = 856$.

Time = 0.25 (sec) , antiderivative size = 3541, normalized size of antiderivative = 8.53

$$\int (a + bx)^{10}(A + Bx)(d + ex)^9 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)**10*(B*x+A)*(e*x+d)**9,x)
```

output

```
A*a**10*d**9*x + B*b**10*e**9*x**21/21 + x**20*(A*b**10*e**9/20 + B*a*b**9
*e**9/2 + 9*B*b**10*d*e**8/20) + x**19*(10*A*a*b**9*e**9/19 + 9*A*b**10*d*
e**8/19 + 45*B*a**2*b**8*e**9/19 + 90*B*a*b**9*d*e**8/19 + 36*B*b**10*d**2
*e**7/19) + x**18*(5*A*a**2*b**8*e**9/2 + 5*A*a*b**9*d*e**8 + 2*A*b**10*d*
*2*e**7 + 20*B*a**3*b**7*e**9/3 + 45*B*a**2*b**8*d*e**8/2 + 20*B*a*b**9*d*
*2*e**7 + 14*B*b**10*d**3*e**6/3) + x**17*(120*A*a**3*b**7*e**9/17 + 405*A
*a**2*b**8*d*e**8/17 + 360*A*a*b**9*d**2*e**7/17 + 84*A*b**10*d**3*e**6/17
+ 210*B*a**4*b**6*e**9/17 + 1080*B*a**3*b**7*d*e**8/17 + 1620*B*a**2*b**8
*d**2*e**7/17 + 840*B*a*b**9*d**3*e**6/17 + 126*B*b**10*d**4*e**5/17) + x*
*16*(105*A*a**4*b**6*e**9/8 + 135*A*a**3*b**7*d*e**8/2 + 405*A*a**2*b**8*d
**2*e**7/4 + 105*A*a*b**9*d**3*e**6/2 + 63*A*b**10*d**4*e**5/8 + 63*B*a**5
*b**5*e**9/4 + 945*B*a**4*b**6*d*e**8/8 + 270*B*a**3*b**7*d**2*e**7 + 945*
B*a**2*b**8*d**3*e**6/4 + 315*B*a*b**9*d**4*e**5/4 + 63*B*b**10*d**5*e**4/
8) + x**15*(84*A*a**5*b**5*e**9/5 + 126*A*a**4*b**6*d*e**8 + 288*A*a**3*b**
7*d**2*e**7 + 252*A*a**2*b**8*d**3*e**6 + 84*A*a*b**9*d**4*e**5 + 42*A*b*
*10*d**5*e**4/5 + 14*B*a**6*b**4*e**9 + 756*B*a**5*b**5*d*e**8/5 + 504*B*a
**4*b**6*d**2*e**7 + 672*B*a**3*b**7*d**3*e**6 + 378*B*a**2*b**8*d**4*e**5
+ 84*B*a*b**9*d**5*e**4 + 28*B*b**10*d**6*e**3/5) + x**14*(15*A*a**6*b**4
*e**9 + 162*A*a**5*b**5*d*e**8 + 540*A*a**4*b**6*d**2*e**7 + 720*A*a**3*b**
7*d**3*e**6 + 405*A*a**2*b**8*d**4*e**5 + 90*A*a*b**9*d**5*e**4 + 6*A*...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2771 vs. $2(393) = 786$.

Time = 0.05 (sec) , antiderivative size = 2771, normalized size of antiderivative = 6.68

$$\int (a + bx)^{10}(A + Bx)(d + ex)^9 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^9,x, algorithm="maxima")
```


output

```

1/21*B*b^10*e^9*x^21 + A*a^10*d^9*x + 1/20*(9*B*b^10*d*e^8 + (10*B*a*b^9 +
A*b^10)*e^9)*x^20 + 1/19*(36*B*b^10*d^2*e^7 + 9*(10*B*a*b^9 + A*b^10)*d*e
^8 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*e^9)*x^19 + 1/6*(28*B*b^10*d^3*e^6 + 12*(
10*B*a*b^9 + A*b^10)*d^2*e^7 + 15*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^8 + 5*(8*B
*a^3*b^7 + 3*A*a^2*b^8)*e^9)*x^18 + 3/17*(42*B*b^10*d^4*e^5 + 28*(10*B*a*b
^9 + A*b^10)*d^3*e^6 + 60*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^7 + 45*(8*B*a^3*
b^7 + 3*A*a^2*b^8)*d*e^8 + 10*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^9)*x^17 + 3/8*
(21*B*b^10*d^5*e^4 + 21*(10*B*a*b^9 + A*b^10)*d^4*e^5 + 70*(9*B*a^2*b^8 +
2*A*a*b^9)*d^3*e^6 + 90*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^7 + 45*(7*B*a^4*
b^6 + 4*A*a^3*b^7)*d*e^8 + 7*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^9)*x^16 + 2/5*(
14*B*b^10*d^6*e^3 + 21*(10*B*a*b^9 + A*b^10)*d^5*e^4 + 105*(9*B*a^2*b^8 +
2*A*a*b^9)*d^4*e^5 + 210*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^6 + 180*(7*B*a^
4*b^6 + 4*A*a^3*b^7)*d^2*e^7 + 63*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^8 + 7*(5
*B*a^6*b^4 + 6*A*a^5*b^5)*e^9)*x^15 + 3/7*(6*B*b^10*d^7*e^2 + 14*(10*B*a*b
^9 + A*b^10)*d^6*e^3 + 105*(9*B*a^2*b^8 + 2*A*a*b^9)*d^5*e^4 + 315*(8*B*a^
3*b^7 + 3*A*a^2*b^8)*d^4*e^5 + 420*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^3*e^6 + 2
52*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e^7 + 63*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d*
e^8 + 5*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e^9)*x^14 + 3/13*(3*B*b^10*d^8*e + 12*
(10*B*a*b^9 + A*b^10)*d^7*e^2 + 140*(9*B*a^2*b^8 + 2*A*a*b^9)*d^6*e^3 + 63
0*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^5*e^4 + 1260*(7*B*a^4*b^6 + 4*A*a^3*b^7...

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3536 vs. $2(393) = 786$.

Time = 0.12 (sec) , antiderivative size = 3536, normalized size of antiderivative = 8.52

$$\int (a + bx)^{10}(A + Bx)(d + ex)^9 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^9,x, algorithm="giac")
```

output

```

1/21*B*b^10*e^9*x^21 + 9/20*B*b^10*d*e^8*x^20 + 1/2*B*a*b^9*e^9*x^20 + 1/2
0*A*b^10*e^9*x^20 + 36/19*B*b^10*d^2*e^7*x^19 + 90/19*B*a*b^9*d*e^8*x^19 +
 9/19*A*b^10*d*e^8*x^19 + 45/19*B*a^2*b^8*e^9*x^19 + 10/19*A*a*b^9*e^9*x^1
9 + 14/3*B*b^10*d^3*e^6*x^18 + 20*B*a*b^9*d^2*e^7*x^18 + 2*A*b^10*d^2*e^7*
x^18 + 45/2*B*a^2*b^8*d*e^8*x^18 + 5*A*a*b^9*d*e^8*x^18 + 20/3*B*a^3*b^7*e
^9*x^18 + 5/2*A*a^2*b^8*e^9*x^18 + 126/17*B*b^10*d^4*e^5*x^17 + 840/17*B*a
*b^9*d^3*e^6*x^17 + 84/17*A*b^10*d^3*e^6*x^17 + 1620/17*B*a^2*b^8*d^2*e^7*
x^17 + 360/17*A*a*b^9*d^2*e^7*x^17 + 1080/17*B*a^3*b^7*d*e^8*x^17 + 405/17
*A*a^2*b^8*d*e^8*x^17 + 210/17*B*a^4*b^6*e^9*x^17 + 120/17*A*a^3*b^7*e^9*x
^17 + 63/8*B*b^10*d^5*e^4*x^16 + 315/4*B*a*b^9*d^4*e^5*x^16 + 63/8*A*b^10*
d^4*e^5*x^16 + 945/4*B*a^2*b^8*d^3*e^6*x^16 + 105/2*A*a*b^9*d^3*e^6*x^16 +
 270*B*a^3*b^7*d^2*e^7*x^16 + 405/4*A*a^2*b^8*d^2*e^7*x^16 + 945/8*B*a^4*b
^6*d*e^8*x^16 + 135/2*A*a^3*b^7*d*e^8*x^16 + 63/4*B*a^5*b^5*e^9*x^16 + 105
/8*A*a^4*b^6*e^9*x^16 + 28/5*B*b^10*d^6*e^3*x^15 + 84*B*a*b^9*d^5*e^4*x^15
 + 42/5*A*b^10*d^5*e^4*x^15 + 378*B*a^2*b^8*d^4*e^5*x^15 + 84*A*a*b^9*d^4*
e^5*x^15 + 672*B*a^3*b^7*d^3*e^6*x^15 + 252*A*a^2*b^8*d^3*e^6*x^15 + 504*B
*a^4*b^6*d^2*e^7*x^15 + 288*A*a^3*b^7*d^2*e^7*x^15 + 756/5*B*a^5*b^5*d*e^8
*x^15 + 126*A*a^4*b^6*d*e^8*x^15 + 14*B*a^6*b^4*e^9*x^15 + 84/5*A*a^5*b^5*
e^9*x^15 + 18/7*B*b^10*d^7*e^2*x^14 + 60*B*a*b^9*d^6*e^3*x^14 + 6*A*b^10*d
^6*e^3*x^14 + 405*B*a^2*b^8*d^5*e^4*x^14 + 90*A*a*b^9*d^5*e^4*x^14 + 10...

```

Mupad [B] (verification not implemented)

Time = 1.58 (sec) , antiderivative size = 2947, normalized size of antiderivative = 7.10

$$\int (a + bx)^{10}(A + Bx)(d + ex)^9 dx = \text{Too large to display}$$

input

```
int((A + B*x)*(a + b*x)^10*(d + e*x)^9,x)
```

output

```

x^9*(A*a^10*d*e^8 + 5*A*a^2*b^8*d^9 + (40*B*a^3*b^7*d^9)/3 + 4*B*a^10*d^2*
e^7 + 120*A*a^3*b^7*d^8*e + 40*A*a^9*b*d^2*e^7 + 210*B*a^4*b^6*d^8*e + (28
0*B*a^9*b*d^3*e^6)/3 + 840*A*a^4*b^6*d^7*e^2 + 2352*A*a^5*b^5*d^6*e^3 + 29
40*A*a^6*b^4*d^5*e^4 + 1680*A*a^7*b^3*d^4*e^5 + 420*A*a^8*b^2*d^3*e^6 + 10
08*B*a^5*b^5*d^7*e^2 + 1960*B*a^6*b^4*d^6*e^3 + 1680*B*a^7*b^3*d^5*e^4 + 6
30*B*a^8*b^2*d^4*e^5) + x^13*((9*B*b^10*d^8*e)/13 + (120*A*a^7*b^3*e^9)/13
+ (45*B*a^8*b^2*e^9)/13 + (36*A*b^10*d^7*e^2)/13 + (840*A*a*b^9*d^6*e^3)/
13 + (1890*A*a^6*b^4*d*e^8)/13 + (360*B*a*b^9*d^7*e^2)/13 + (1080*B*a^7*b^
3*d*e^8)/13 + (5670*A*a^2*b^8*d^5*e^4)/13 + (15120*A*a^3*b^7*d^4*e^5)/13 +
(17640*A*a^4*b^6*d^3*e^6)/13 + (9072*A*a^5*b^5*d^2*e^7)/13 + (3780*B*a^2*
b^8*d^6*e^3)/13 + (15120*B*a^3*b^7*d^5*e^4)/13 + (26460*B*a^4*b^6*d^4*e^5)
/13 + (21168*B*a^5*b^5*d^3*e^6)/13 + (7560*B*a^6*b^4*d^2*e^7)/13) + x^5*(4
2*A*a^6*b^4*d^9 + 24*B*a^7*b^3*d^9 + (126*A*a^10*d^5*e^4)/5 + (84*B*a^10*d
^6*e^3)/5 + 216*A*a^7*b^3*d^8*e + 168*A*a^9*b*d^6*e^3 + 81*B*a^8*b^2*d^8*e
+ 72*B*a^9*b*d^7*e^2 + 324*A*a^8*b^2*d^7*e^2) + x^8*(15*A*a^3*b^7*d^9 + (
105*B*a^4*b^6*d^9)/4 + (9*A*a^10*d^2*e^7)/2 + (21*B*a^10*d^3*e^6)/2 + (945
*A*a^4*b^6*d^8*e)/4 + 105*A*a^9*b*d^3*e^6 + (567*B*a^5*b^5*d^8*e)/2 + (315
*B*a^9*b*d^4*e^5)/2 + 1134*A*a^5*b^5*d^7*e^2 + 2205*A*a^6*b^4*d^6*e^3 + 18
90*A*a^7*b^3*d^5*e^4 + (2835*A*a^8*b^2*d^4*e^5)/4 + 945*B*a^6*b^4*d^7*e^2
+ 1260*B*a^7*b^3*d^6*e^3 + (2835*B*a^8*b^2*d^5*e^4)/4) + x^17*((120*A*a...

```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 1817, normalized size of antiderivative = 4.38

$$\int (a + bx)^{10}(A + Bx)(d + ex)^9 dx = \text{Too large to display}$$

input

```
int((b*x+a)^10*(B*x+A)*(e*x+d)^9,x)
```

output

```
(x*(3527160*a**11*d**9 + 15872220*a**11*d**8*e*x + 42325920*a**11*d**7*e**2*x**2 + 74070360*a**11*d**6*e**3*x**3 + 88884432*a**11*d**5*e**4*x**4 + 74070360*a**11*d**4*e**5*x**5 + 42325920*a**11*d**3*e**6*x**6 + 15872220*a**11*d**2*e**7*x**7 + 3527160*a**11*d*e**8*x**8 + 352716*a**11*e**9*x**9 + 19399380*a**10*b*d**9*x + 116396280*a**10*b*d**8*e*x**2 + 349188840*a**10*b*d**7*e**2*x**3 + 651819168*a**10*b*d**6*e**3*x**4 + 814773960*a**10*b*d**5*e**4*x**5 + 698377680*a**10*b*d**4*e**5*x**6 + 407386980*a**10*b*d**3*e**6*x**7 + 155195040*a**10*b*d**2*e**7*x**8 + 34918884*a**10*b*d*e**8*x**9 + 3527160*a**10*b*e**9*x**10 + 64664600*a**9*b**2*d**9*x**2 + 436486050*a**9*b**2*d**8*e*x**3 + 1396755360*a**9*b**2*d**7*e**2*x**4 + 2715913200*a**9*b**2*d**6*e**3*x**5 + 3491888400*a**9*b**2*d**5*e**4*x**6 + 3055402350*a**9*b**2*d**4*e**5*x**7 + 1810608800*a**9*b**2*d**3*e**6*x**8 + 698377680*a**9*b**2*d**2*e**7*x**9 + 158722200*a**9*b**2*d*e**8*x**10 + 16166150*a**9*b**2*e**9*x**11 + 145495350*a**8*b**3*d**9*x**3 + 1047566520*a**8*b**3*d**8*e*x**4 + 3491888400*a**8*b**3*d**7*e**2*x**5 + 6983776800*a**8*b**3*d**6*e**3*x**6 + 9166207050*a**8*b**3*d**5*e**4*x**7 + 8147739600*a**8*b**3*d**4*e**5*x**8 + 4888643760*a**8*b**3*d**3*e**6*x**9 + 1904666400*a**8*b**3*d**2*e**7*x**10 + 436486050*a**8*b**3*d*e**8*x**11 + 44767800*a**8*b**3*e**9*x**12 + 232792560*a**7*b**4*d**9*x**4 + 1745944200*a**7*b**4*d**8*e*x**5 + 5986094400*a**7*b**4*d**7*e**2*x**6 + 12221609400*a**7*b**4*d**6...
```

3.70 $\int (a + bx)^{10}(A + Bx)(d + ex)^8 dx$

Optimal result	712
Mathematica [B] (verified)	713
Rubi [A] (verified)	714
Maple [B] (verified)	716
Fricas [B] (verification not implemented)	717
Sympy [B] (verification not implemented)	718
Maxima [B] (verification not implemented)	719
Giac [B] (verification not implemented)	720
Mupad [B] (verification not implemented)	721
Reduce [B] (verification not implemented)	721

Optimal result

Integrand size = 20, antiderivative size = 372

$$\begin{aligned}
 \int (a + bx)^{10}(A + Bx)(d + ex)^8 dx = & \frac{(Ab - aB)(bd - ae)^8(a + bx)^{11}}{11b^{10}} \\
 & + \frac{(bd - ae)^7(bBd + 8Abe - 9aBe)(a + bx)^{12}}{12b^{10}} \\
 & + \frac{4e(bd - ae)^6(2bBd + 7Abe - 9aBe)(a + bx)^{13}}{13b^{10}} \\
 & + \frac{2e^2(bd - ae)^5(bBd + 2Abe - 3aBe)(a + bx)^{14}}{b^{10}} \\
 & + \frac{14e^3(bd - ae)^4(4bBd + 5Abe - 9aBe)(a + bx)^{15}}{15b^{10}} \\
 & + \frac{7e^4(bd - ae)^3(5bBd + 4Abe - 9aBe)(a + bx)^{16}}{8b^{10}} \\
 & + \frac{28e^5(bd - ae)^2(2bBd + Abe - 3aBe)(a + bx)^{17}}{17b^{10}} \\
 & + \frac{2e^6(bd - ae)(7bBd + 2Abe - 9aBe)(a + bx)^{18}}{9b^{10}} \\
 & + \frac{e^7(8bBd + Abe - 9aBe)(a + bx)^{19}}{19b^{10}} \\
 & + \frac{Be^8(a + bx)^{20}}{20b^{10}}
 \end{aligned}$$

output

```

1/11*(A*b-B*a)*(-a*e+b*d)^8*(b*x+a)^11/b^10+1/12*(-a*e+b*d)^7*(8*A*b*e-9*B
*a*e+B*b*d)*(b*x+a)^12/b^10+4/13*e*(-a*e+b*d)^6*(7*A*b*e-9*B*a*e+2*B*b*d)*
(b*x+a)^13/b^10+2*e^2*(-a*e+b*d)^5*(2*A*b*e-3*B*a*e+B*b*d)*(b*x+a)^14/b^10
+14/15*e^3*(-a*e+b*d)^4*(5*A*b*e-9*B*a*e+4*B*b*d)*(b*x+a)^15/b^10+7/8*e^4*
(-a*e+b*d)^3*(4*A*b*e-9*B*a*e+5*B*b*d)*(b*x+a)^16/b^10+28/17*e^5*(-a*e+b*d
)^2*(A*b*e-3*B*a*e+2*B*b*d)*(b*x+a)^17/b^10+2/9*e^6*(-a*e+b*d)*(2*A*b*e-9*
B*a*e+7*B*b*d)*(b*x+a)^18/b^10+1/19*e^7*(A*b*e-9*B*a*e+8*B*b*d)*(b*x+a)^19
/b^10+1/20*B*e^8*(b*x+a)^20/b^10

```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 2307 vs. $2(372) = 744$.

Time = 0.53 (sec) , antiderivative size = 2307, normalized size of antiderivative = 6.20

$$\int (a + bx)^{10}(A + Bx)(d + ex)^8 dx = \text{Result too large to show}$$

input

```
Integrate[(a + b*x)^10*(A + B*x)*(d + e*x)^8,x]
```

output

```

a^10*A*d^8*x + (a^9*d^7*(10*A*b*d + a*B*d + 8*a*A*e)*x^2)/2 + (a^8*d^6*(2*
a*B*d*(5*b*d + 4*a*e) + A*(45*b^2*d^2 + 80*a*b*d*e + 28*a^2*e^2))*x^3)/3 +
(a^7*d^5*(a*B*d*(45*b^2*d^2 + 80*a*b*d*e + 28*a^2*e^2) + 8*A*(15*b^3*d^3
+ 45*a*b^2*d^2*e + 35*a^2*b*d*e^2 + 7*a^3*e^3))*x^4)/4 + (2*a^6*d^4*(4*a*B
*d*(15*b^3*d^3 + 45*a*b^2*d^2*e + 35*a^2*b*d*e^2 + 7*a^3*e^3) + 5*A*(21*b^
4*d^4 + 96*a*b^3*d^3*e + 126*a^2*b^2*d^2*e^2 + 56*a^3*b*d*e^3 + 7*a^4*e^4)
)*x^5)/5 + (a^5*d^3*(5*a*B*d*(21*b^4*d^4 + 96*a*b^3*d^3*e + 126*a^2*b^2*d^
2*e^2 + 56*a^3*b*d*e^3 + 7*a^4*e^4) + 14*A*(9*b^5*d^5 + 60*a*b^4*d^4*e + 1
20*a^2*b^3*d^3*e^2 + 90*a^3*b^2*d^2*e^3 + 25*a^4*b*d*e^4 + 2*a^5*e^5))*x^6
)/3 + 2*a^4*d^2*(2*a*B*d*(9*b^5*d^5 + 60*a*b^4*d^4*e + 120*a^2*b^3*d^3*e^2
+ 90*a^3*b^2*d^2*e^3 + 25*a^4*b*d*e^4 + 2*a^5*e^5) + A*(15*b^6*d^6 + 144*
a*b^5*d^5*e + 420*a^2*b^4*d^4*e^2 + 480*a^3*b^3*d^3*e^3 + 225*a^4*b^2*d^2*
e^4 + 40*a^5*b*d*e^5 + 2*a^6*e^6))*x^7 + (a^3*d*(7*a*B*d*(15*b^6*d^6 + 144
*a*b^5*d^5*e + 420*a^2*b^4*d^4*e^2 + 480*a^3*b^3*d^3*e^3 + 225*a^4*b^2*d^2
*e^4 + 40*a^5*b*d*e^5 + 2*a^6*e^6) + 4*A*(15*b^7*d^7 + 210*a*b^6*d^6*e + 8
82*a^2*b^5*d^5*e^2 + 1470*a^3*b^4*d^4*e^3 + 1050*a^4*b^3*d^3*e^4 + 315*a^5
*b^2*d^2*e^5 + 35*a^6*b*d*e^6 + a^7*e^7))*x^8)/4 + (a^2*(8*a*B*d*(15*b^7*d
^7 + 210*a*b^6*d^6*e + 882*a^2*b^5*d^5*e^2 + 1470*a^3*b^4*d^4*e^3 + 1050*a
^4*b^3*d^3*e^4 + 315*a^5*b^2*d^2*e^5 + 35*a^6*b*d*e^6 + a^7*e^7) + A*(45*b
^8*d^8 + 960*a*b^7*d^7*e + 5880*a^2*b^6*d^6*e^2 + 14112*a^3*b^5*d^5*e^3...

```

Rubi [A] (verified)

Time = 2.84 (sec) , antiderivative size = 372, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^{10} (A + Bx)(d + ex)^8 dx$$

$$\downarrow 86$$

$$\int \left(\frac{e^7 (a + bx)^{18} (-9aBe + Abe + 8bBd)}{b^9} + \frac{4e^6 (a + bx)^{17} (bd - ae) (-9aBe + 2Abe + 7bBd)}{b^9} + \frac{28e^5 (a + bx)^{16} (bd - ae)^2}{b^9} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned} & \frac{e^7(a+bx)^{19}(-9aBe + Abe + 8bBd)}{19b^{10}} + \frac{2e^6(a+bx)^{18}(bd-ae)(-9aBe + 2Abe + 7bBd)}{9b^{10}} + \\ & \frac{28e^5(a+bx)^{17}(bd-ae)^2(-3aBe + Abe + 2bBd)}{17b^{10}} + \\ & \frac{7e^4(a+bx)^{16}(bd-ae)^3(-9aBe + 4Abe + 5bBd)}{8b^{10}} + \\ & \frac{14e^3(a+bx)^{15}(bd-ae)^4(-9aBe + 5Abe + 4bBd)}{15b^{10}} + \\ & \frac{2e^2(a+bx)^{14}(bd-ae)^5(-3aBe + 2Abe + bBd)}{b^{10}} + \\ & \frac{4e(a+bx)^{13}(bd-ae)^6(-9aBe + 7Abe + 2bBd)}{13b^{10}} + \\ & \frac{(a+bx)^{12}(bd-ae)^7(-9aBe + 8Abe + bBd)}{12b^{10}} + \frac{(a+bx)^{11}(Ab-aB)(bd-ae)^8}{11b^{10}} + \\ & \frac{Be^8(a+bx)^{20}}{20b^{10}} \end{aligned}$$

input `Int[(a + b*x)^10*(A + B*x)*(d + e*x)^8,x]`

output `((A*b - a*B)*(b*d - a*e)^8*(a + b*x)^11)/(11*b^10) + ((b*d - a*e)^7*(b*B*d + 8*A*b*e - 9*a*B*e)*(a + b*x)^12)/(12*b^10) + (4*e*(b*d - a*e)^6*(2*b*B*d + 7*A*b*e - 9*a*B*e)*(a + b*x)^13)/(13*b^10) + (2*e^2*(b*d - a*e)^5*(b*B*d + 2*A*b*e - 3*a*B*e)*(a + b*x)^14)/b^10 + (14*e^3*(b*d - a*e)^4*(4*b*B*d + 5*A*b*e - 9*a*B*e)*(a + b*x)^15)/(15*b^10) + (7*e^4*(b*d - a*e)^3*(5*b*B*d + 4*A*b*e - 9*a*B*e)*(a + b*x)^16)/(8*b^10) + (28*e^5*(b*d - a*e)^2*(2*b*B*d + A*b*e - 3*a*B*e)*(a + b*x)^17)/(17*b^10) + (2*e^6*(b*d - a*e)*(7*b*B*d + 2*A*b*e - 9*a*B*e)*(a + b*x)^18)/(9*b^10) + (e^7*(8*b*B*d + A*b*e - 9*a*B*e)*(a + b*x)^19)/(19*b^10) + (B*e^8*(a + b*x)^20)/(20*b^10)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2472 vs. $2(354) = 708$.

Time = 0.26 (sec) , antiderivative size = 2473, normalized size of antiderivative = 6.65

method	result	size
default	Expression too large to display	2473
norman	Expression too large to display	2662
gosper	Expression too large to display	3160
risch	Expression too large to display	3160
parallelrisc	Expression too large to display	3160
orering	Expression too large to display	3160

input `int((b*x+a)^10*(B*x+A)*(e*x+d)^8,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 1/20*b^{10}*B*e^8*x^{20}+1/19*((A*b^{10}+10*B*a*b^9)*e^8+8*b^{10}*B*d*e^7)*x^{19}+1/ \\ & 18*((10*A*a*b^9+45*B*a^2*b^8)*e^8+8*(A*b^{10}+10*B*a*b^9)*d*e^7+28*b^{10}*B*d^2 \\ & *e^6)*x^{18}+1/17*((45*A*a^2*b^8+120*B*a^3*b^7)*e^8+8*(10*A*a*b^9+45*B*a^2*b^8) \\ & *d*e^7+28*(A*b^{10}+10*B*a*b^9)*d^2*e^6+56*b^{10}*B*d^3*e^5)*x^{17}+1/16*((1 \\ & 20*A*a^3*b^7+210*B*a^4*b^6)*e^8+8*(45*A*a^2*b^8+120*B*a^3*b^7)*d*e^7+28*(1 \\ & 0*A*a*b^9+45*B*a^2*b^8)*d^2*e^6+56*(A*b^{10}+10*B*a*b^9)*d^3*e^5+70*b^{10}*B*d \\ & ^4*e^4)*x^{16}+1/15*((210*A*a^4*b^6+252*B*a^5*b^5)*e^8+8*(120*A*a^3*b^7+210* \\ & B*a^4*b^6)*d*e^7+28*(45*A*a^2*b^8+120*B*a^3*b^7)*d^2*e^6+56*(10*A*a*b^9+45 \\ & *B*a^2*b^8)*d^3*e^5+70*(A*b^{10}+10*B*a*b^9)*d^4*e^4+56*b^{10}*B*d^5*e^3)*x^{15} \\ & +1/14*((252*A*a^5*b^5+210*B*a^6*b^4)*e^8+8*(210*A*a^4*b^6+252*B*a^5*b^5)*d \\ & *e^7+28*(120*A*a^3*b^7+210*B*a^4*b^6)*d^2*e^6+56*(45*A*a^2*b^8+120*B*a^3*b^7) \\ & *d^3*e^5+70*(10*A*a*b^9+45*B*a^2*b^8)*d^4*e^4+56*(A*b^{10}+10*B*a*b^9)*d^5 \\ & *e^3+28*b^{10}*B*d^6*e^2)*x^{14}+1/13*((210*A*a^6*b^4+120*B*a^7*b^3)*e^8+8*(2 \\ & 52*A*a^5*b^5+210*B*a^6*b^4)*d*e^7+28*(210*A*a^4*b^6+252*B*a^5*b^5)*d^2*e^6 \\ & +56*(120*A*a^3*b^7+210*B*a^4*b^6)*d^3*e^5+70*(45*A*a^2*b^8+120*B*a^3*b^7)* \\ & d^4*e^4+56*(10*A*a*b^9+45*B*a^2*b^8)*d^5*e^3+28*(A*b^{10}+10*B*a*b^9)*d^6*e^2 \\ & +8*b^{10}*B*d^7*e)*x^{13}+1/12*((120*A*a^7*b^3+45*B*a^8*b^2)*e^8+8*(210*A*a^6 \\ & *b^4+120*B*a^7*b^3)*d*e^7+28*(252*A*a^5*b^5+210*B*a^6*b^4)*d^2*e^6+56*(210 \\ & *A*a^4*b^6+252*B*a^5*b^5)*d^3*e^5+70*(120*A*a^3*b^7+210*B*a^4*b^6)*d^4*e^4 \\ & +56*(45*A*a^2*b^8+120*B*a^3*b^7)*d^5*e^3+28*(10*A*a*b^9+45*B*a^2*b^8)*d\dots \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2487 vs. $2(354) = 708$.

Time = 0.11 (sec) , antiderivative size = 2487, normalized size of antiderivative = 6.69

$$\int (a + bx)^{10}(A + Bx)(d + ex)^8 dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)*(e*x+d)^8,x, algorithm="fricas")`

output

```
1/20*B*b^10*e^8*x^20 + A*a^10*d^8*x + 1/19*(8*B*b^10*d*e^7 + (10*B*a*b^9 +
A*b^10)*e^8)*x^19 + 1/18*(28*B*b^10*d^2*e^6 + 8*(10*B*a*b^9 + A*b^10)*d*e
^7 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*e^8)*x^18 + 1/17*(56*B*b^10*d^3*e^5 + 28*
(10*B*a*b^9 + A*b^10)*d^2*e^6 + 40*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^7 + 15*(8
*B*a^3*b^7 + 3*A*a^2*b^8)*e^8)*x^17 + 1/8*(35*B*b^10*d^4*e^4 + 28*(10*B*a*
b^9 + A*b^10)*d^3*e^5 + 70*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^6 + 60*(8*B*a^3
*b^7 + 3*A*a^2*b^8)*d*e^7 + 15*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^8)*x^16 + 2/1
5*(28*B*b^10*d^5*e^3 + 35*(10*B*a*b^9 + A*b^10)*d^4*e^4 + 140*(9*B*a^2*b^8
+ 2*A*a*b^9)*d^3*e^5 + 210*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^6 + 120*(7*B
*a^4*b^6 + 4*A*a^3*b^7)*d*e^7 + 21*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^8)*x^15 +
(2*B*b^10*d^6*e^2 + 4*(10*B*a*b^9 + A*b^10)*d^5*e^3 + 25*(9*B*a^2*b^8 + 2
*A*a*b^9)*d^4*e^4 + 60*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^5 + 60*(7*B*a^4*b
^6 + 4*A*a^3*b^7)*d^2*e^6 + 24*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^7 + 3*(5*B*
a^6*b^4 + 6*A*a^5*b^5)*e^8)*x^14 + 2/13*(4*B*b^10*d^7*e + 14*(10*B*a*b^9 +
A*b^10)*d^6*e^2 + 140*(9*B*a^2*b^8 + 2*A*a*b^9)*d^5*e^3 + 525*(8*B*a^3*b^
7 + 3*A*a^2*b^8)*d^4*e^4 + 840*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^3*e^5 + 588*(
6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e^6 + 168*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d*e^7
+ 15*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e^8)*x^13 + 1/12*(B*b^10*d^8 + 8*(10*B*a
*b^9 + A*b^10)*d^7*e + 140*(9*B*a^2*b^8 + 2*A*a*b^9)*d^6*e^2 + 840*(8*B*a^
3*b^7 + 3*A*a^2*b^8)*d^5*e^3 + 2100*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^4*e^4...
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3165 vs. $2(384) = 768$.

Time = 0.22 (sec) , antiderivative size = 3165, normalized size of antiderivative = 8.51

$$\int (a + bx)^{10}(A + Bx)(d + ex)^8 dx = \text{Too large to display}$$

input `integrate((b*x+a)**10*(B*x+A)*(e*x+d)**8,x)`

output

```
A*a**10*d**8*x + B*b**10*e**8*x**20/20 + x**19*(A*b**10*e**8/19 + 10*B*a*b
**9*e**8/19 + 8*B*b**10*d*e**7/19) + x**18*(5*A*a*b**9*e**8/9 + 4*A*b**10*
d*e**7/9 + 5*B*a**2*b**8*e**8/2 + 40*B*a*b**9*d*e**7/9 + 14*B*b**10*d**2*e
**6/9) + x**17*(45*A*a**2*b**8*e**8/17 + 80*A*a*b**9*d*e**7/17 + 28*A*b**1
0*d**2*e**6/17 + 120*B*a**3*b**7*e**8/17 + 360*B*a**2*b**8*d*e**7/17 + 280
*B*a*b**9*d**2*e**6/17 + 56*B*b**10*d**3*e**5/17) + x**16*(15*A*a**3*b**7*
e**8/2 + 45*A*a**2*b**8*d*e**7/2 + 35*A*a*b**9*d**2*e**6/2 + 7*A*b**10*d**
3*e**5/2 + 105*B*a**4*b**6*e**8/8 + 60*B*a**3*b**7*d*e**7 + 315*B*a**2*b**
8*d**2*e**6/4 + 35*B*a*b**9*d**3*e**5 + 35*B*b**10*d**4*e**4/8) + x**15*(1
4*A*a**4*b**6*e**8 + 64*A*a**3*b**7*d*e**7 + 84*A*a**2*b**8*d**2*e**6 + 11
2*A*a*b**9*d**3*e**5/3 + 14*A*b**10*d**4*e**4/3 + 84*B*a**5*b**5*e**8/5 +
112*B*a**4*b**6*d*e**7 + 224*B*a**3*b**7*d**2*e**6 + 168*B*a**2*b**8*d**3*
e**5 + 140*B*a*b**9*d**4*e**4/3 + 56*B*b**10*d**5*e**3/15) + x**14*(18*A*a
**5*b**5*e**8 + 120*A*a**4*b**6*d*e**7 + 240*A*a**3*b**7*d**2*e**6 + 180*A
a**2*b**8*d**3*e**5 + 50*A*a*b**9*d**4*e**4 + 4*A*b**10*d**5*e**3 + 15*B*
a**6*b**4*e**8 + 144*B*a**5*b**5*d*e**7 + 420*B*a**4*b**6*d**2*e**6 + 480*
B*a**3*b**7*d**3*e**5 + 225*B*a**2*b**8*d**4*e**4 + 40*B*a*b**9*d**5*e**3
+ 2*B*b**10*d**6*e**2) + x**13*(210*A*a**6*b**4*e**8/13 + 2016*A*a**5*b**5
*d*e**7/13 + 5880*A*a**4*b**6*d**2*e**6/13 + 6720*A*a**3*b**7*d**3*e**5/13
+ 3150*A*a**2*b**8*d**4*e**4/13 + 560*A*a*b**9*d**5*e**3/13 + 28*A*b**...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2487 vs. $2(354) = 708$.

Time = 0.05 (sec) , antiderivative size = 2487, normalized size of antiderivative = 6.69

$$\int (a + bx)^{10}(A + Bx)(d + ex)^8 dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)*(e*x+d)^8,x, algorithm="maxima")`

output

```
1/20*B*b^10*e^8*x^20 + A*a^10*d^8*x + 1/19*(8*B*b^10*d*e^7 + (10*B*a*b^9 +
A*b^10)*e^8)*x^19 + 1/18*(28*B*b^10*d^2*e^6 + 8*(10*B*a*b^9 + A*b^10)*d*e
^7 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*e^8)*x^18 + 1/17*(56*B*b^10*d^3*e^5 + 28*
(10*B*a*b^9 + A*b^10)*d^2*e^6 + 40*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^7 + 15*(8
*B*a^3*b^7 + 3*A*a^2*b^8)*e^8)*x^17 + 1/8*(35*B*b^10*d^4*e^4 + 28*(10*B*a*
b^9 + A*b^10)*d^3*e^5 + 70*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^6 + 60*(8*B*a^3
*b^7 + 3*A*a^2*b^8)*d*e^7 + 15*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^8)*x^16 + 2/1
5*(28*B*b^10*d^5*e^3 + 35*(10*B*a*b^9 + A*b^10)*d^4*e^4 + 140*(9*B*a^2*b^8
+ 2*A*a*b^9)*d^3*e^5 + 210*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^6 + 120*(7*B
*a^4*b^6 + 4*A*a^3*b^7)*d*e^7 + 21*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^8)*x^15 +
(2*B*b^10*d^6*e^2 + 4*(10*B*a*b^9 + A*b^10)*d^5*e^3 + 25*(9*B*a^2*b^8 + 2
*A*a*b^9)*d^4*e^4 + 60*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^5 + 60*(7*B*a^4*b
^6 + 4*A*a^3*b^7)*d^2*e^6 + 24*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^7 + 3*(5*B*
a^6*b^4 + 6*A*a^5*b^5)*e^8)*x^14 + 2/13*(4*B*b^10*d^7*e + 14*(10*B*a*b^9 +
A*b^10)*d^6*e^2 + 140*(9*B*a^2*b^8 + 2*A*a*b^9)*d^5*e^3 + 525*(8*B*a^3*b^
7 + 3*A*a^2*b^8)*d^4*e^4 + 840*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^3*e^5 + 588*(
6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e^6 + 168*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d*e^7
+ 15*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e^8)*x^13 + 1/12*(B*b^10*d^8 + 8*(10*B*a
*b^9 + A*b^10)*d^7*e + 140*(9*B*a^2*b^8 + 2*A*a*b^9)*d^6*e^2 + 840*(8*B*a^
3*b^7 + 3*A*a^2*b^8)*d^5*e^3 + 2100*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^4*e^4...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3159 vs. $2(354) = 708$.

Time = 0.12 (sec) , antiderivative size = 3159, normalized size of antiderivative = 8.49

$$\int (a + bx)^{10}(A + Bx)(d + ex)^8 dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)*(e*x+d)^8,x, algorithm="giac")`

output

```
1/20*B*b^10*e^8*x^20 + 8/19*B*b^10*d*e^7*x^19 + 10/19*B*a*b^9*e^8*x^19 + 1
/19*A*b^10*e^8*x^19 + 14/9*B*b^10*d^2*e^6*x^18 + 40/9*B*a*b^9*d*e^7*x^18 +
4/9*A*b^10*d*e^7*x^18 + 5/2*B*a^2*b^8*e^8*x^18 + 5/9*A*a*b^9*e^8*x^18 + 5
6/17*B*b^10*d^3*e^5*x^17 + 280/17*B*a*b^9*d^2*e^6*x^17 + 28/17*A*b^10*d^2*
e^6*x^17 + 360/17*B*a^2*b^8*d*e^7*x^17 + 80/17*A*a*b^9*d*e^7*x^17 + 120/17
*B*a^3*b^7*e^8*x^17 + 45/17*A*a^2*b^8*e^8*x^17 + 35/8*B*b^10*d^4*e^4*x^16
+ 35*B*a*b^9*d^3*e^5*x^16 + 7/2*A*b^10*d^3*e^5*x^16 + 315/4*B*a^2*b^8*d^2*
e^6*x^16 + 35/2*A*a*b^9*d^2*e^6*x^16 + 60*B*a^3*b^7*d*e^7*x^16 + 45/2*A*a^
2*b^8*d*e^7*x^16 + 105/8*B*a^4*b^6*e^8*x^16 + 15/2*A*a^3*b^7*e^8*x^16 + 56
/15*B*b^10*d^5*e^3*x^15 + 140/3*B*a*b^9*d^4*e^4*x^15 + 14/3*A*b^10*d^4*e^4
*x^15 + 168*B*a^2*b^8*d^3*e^5*x^15 + 112/3*A*a*b^9*d^3*e^5*x^15 + 224*B*a^
3*b^7*d^2*e^6*x^15 + 84*A*a^2*b^8*d^2*e^6*x^15 + 112*B*a^4*b^6*d*e^7*x^15
+ 64*A*a^3*b^7*d*e^7*x^15 + 84/5*B*a^5*b^5*e^8*x^15 + 14*A*a^4*b^6*e^8*x^1
5 + 2*B*b^10*d^6*e^2*x^14 + 40*B*a*b^9*d^5*e^3*x^14 + 4*A*b^10*d^5*e^3*x^1
4 + 225*B*a^2*b^8*d^4*e^4*x^14 + 50*A*a*b^9*d^4*e^4*x^14 + 480*B*a^3*b^7*d
^3*e^5*x^14 + 180*A*a^2*b^8*d^3*e^5*x^14 + 420*B*a^4*b^6*d^2*e^6*x^14 + 24
0*A*a^3*b^7*d^2*e^6*x^14 + 144*B*a^5*b^5*d*e^7*x^14 + 120*A*a^4*b^6*d*e^7*
x^14 + 15*B*a^6*b^4*e^8*x^14 + 18*A*a^5*b^5*e^8*x^14 + 8/13*B*b^10*d^7*e*x
^13 + 280/13*B*a*b^9*d^6*e^2*x^13 + 28/13*A*b^10*d^6*e^2*x^13 + 2520/13*B*
a^2*b^8*d^5*e^3*x^13 + 560/13*A*a*b^9*d^5*e^3*x^13 + 8400/13*B*a^3*b^7*...
```

Mupad [B] (verification not implemented)

Time = 1.49 (sec) , antiderivative size = 2631, normalized size of antiderivative = 7.07

$$\int (a + bx)^{10}(A + Bx)(d + ex)^8 dx = \text{Too large to display}$$

input `int((A + B*x)*(a + b*x)^10*(d + e*x)^8,x)`

output

```
x^5*(42*A*a^6*b^4*d^8 + 24*B*a^7*b^3*d^8 + 14*A*a^10*d^4*e^4 + (56*B*a^10*d^5*e^3)/5 + 192*A*a^7*b^3*d^7*e + 112*A*a^9*b*d^5*e^3 + 72*B*a^8*b^2*d^7*e + 56*B*a^9*b*d^6*e^2 + 252*A*a^8*b^2*d^6*e^2) + x^16*((15*A*a^3*b^7*e^8)/2 + (105*B*a^4*b^6*e^8)/8 + (7*A*b^10*d^3*e^5)/2 + (35*B*b^10*d^4*e^4)/8 + (35*A*a*b^9*d^2*e^6)/2 + (45*A*a^2*b^8*d*e^7)/2 + 35*B*a*b^9*d^3*e^5 + 60*B*a^3*b^7*d*e^7 + (315*B*a^2*b^8*d^2*e^6)/4) + x^10*((B*a^10*e^8)/10 + A*a*b^9*d^8 + A*a^9*b*e^8 + (9*B*a^2*b^8*d^8)/2 + 36*A*a^2*b^8*d^7*e + 36*A*a^8*b^2*d*e^7 + 96*B*a^3*b^7*d^7*e + 336*A*a^3*b^7*d^6*e^2 + 1176*A*a^4*b^6*d^5*e^3 + 1764*A*a^5*b^5*d^4*e^4 + 1176*A*a^6*b^4*d^3*e^5 + 336*A*a^7*b^3*d^2*e^6 + 588*B*a^4*b^6*d^6*e^2 + (7056*B*a^5*b^5*d^5*e^3)/5 + 1470*B*a^6*b^4*d^4*e^4 + 672*B*a^7*b^3*d^3*e^5 + 126*B*a^8*b^2*d^2*e^6 + 8*B*a^9*b*d*e^7) + x^11*((A*b^10*d^8)/11 + (10*B*a*b^9*d^8)/11 + (10*B*a^9*b*e^8)/11 + (45*A*a^8*b^2*e^8)/11 + (960*A*a^7*b^3*d*e^7)/11 + (360*B*a^2*b^8*d^7*e)/11 + (360*B*a^8*b^2*d*e^7)/11 + (1260*A*a^2*b^8*d^6*e^2)/11 + (6720*A*a^3*b^7*d^5*e^3)/11 + (14700*A*a^4*b^6*d^4*e^4)/11 + (14112*A*a^5*b^5*d^3*e^5)/11 + (5880*A*a^6*b^4*d^2*e^6)/11 + (3360*B*a^3*b^7*d^6*e^2)/11 + (11760*B*a^4*b^6*d^5*e^3)/11 + (17640*B*a^5*b^5*d^4*e^4)/11 + (11760*B*a^6*b^4*d^3*e^5)/11 + (3360*B*a^7*b^3*d^2*e^6)/11 + (80*A*a*b^9*d^7*e)/11) + x^6*(42*A*a^5*b^5*d^8 + 35*B*a^6*b^4*d^8 + (28*A*a^10*d^3*e^5)/3 + (35*B*a^10*d^4*e^4)/3 + 280*A*a^6*b^4*d^7*e + (350*A*a^9*b*d^4*e^4)/3 + 160*B*a^7*b...
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 1623, normalized size of antiderivative = 4.36

$$\int (a + bx)^{10}(A + Bx)(d + ex)^8 dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)*(e*x+d)^8,x)`

output

```
(x*(1511640*a**11*d**8 + 6046560*a**11*d**7*e*x + 14108640*a**11*d**6*e**2
*x**2 + 21162960*a**11*d**5*e**3*x**3 + 21162960*a**11*d**4*e**4*x**4 + 14
108640*a**11*d**3*e**5*x**5 + 6046560*a**11*d**2*e**6*x**6 + 1511640*a**11
*d*e**7*x**7 + 167960*a**11*e**8*x**8 + 8314020*a**10*b*d**8*x + 44341440*
a**10*b*d**7*e*x**2 + 116396280*a**10*b*d**6*e**2*x**3 + 186234048*a**10*b
*d**5*e**3*x**4 + 193993800*a**10*b*d**4*e**4*x**5 + 133024320*a**10*b*d**
3*e**5*x**6 + 58198140*a**10*b*d**2*e**6*x**7 + 14780480*a**10*b*d*e**7*x*
*8 + 1662804*a**10*b*e**8*x**9 + 27713400*a**9*b**2*d**8*x**2 + 166280400*
a**9*b**2*d**7*e*x**3 + 465585120*a**9*b**2*d**6*e**2*x**4 + 775975200*a**
9*b**2*d**5*e**3*x**5 + 831402000*a**9*b**2*d**4*e**4*x**6 + 581981400*a**
9*b**2*d**3*e**5*x**7 + 258658400*a**9*b**2*d**2*e**6*x**8 + 66512160*a**9
*b**2*d*e**7*x**9 + 7558200*a**9*b**2*e**8*x**10 + 62355150*a**8*b**3*d**8
*x**3 + 399072960*a**8*b**3*d**7*e*x**4 + 1163962800*a**8*b**3*d**6*e**2*x
**5 + 1995364800*a**8*b**3*d**5*e**3*x**6 + 2182430250*a**8*b**3*d**4*e**4
*x**7 + 1551950400*a**8*b**3*d**3*e**5*x**8 + 698377680*a**8*b**3*d**2*e**
6*x**9 + 181396800*a**8*b**3*d*e**7*x**10 + 20785050*a**8*b**3*e**8*x**11
+ 99768240*a**7*b**4*d**8*x**4 + 665121600*a**7*b**4*d**7*e*x**5 + 1995364
800*a**7*b**4*d**6*e**2*x**6 + 3491888400*a**7*b**4*d**5*e**3*x**7 + 38798
76000*a**7*b**4*d**4*e**4*x**8 + 2793510720*a**7*b**4*d**3*e**5*x**9 + 126
9777600*a**7*b**4*d**2*e**6*x**10 + 332560800*a**7*b**4*d*e**7*x**11 + ...
```

3.71 $\int (a + bx)^{10}(A + Bx)(d + ex)^7 dx$

Optimal result	723
Mathematica [B] (verified)	724
Rubi [A] (verified)	725
Maple [B] (verified)	727
Fricas [B] (verification not implemented)	728
Sympy [B] (verification not implemented)	729
Maxima [B] (verification not implemented)	730
Giac [B] (verification not implemented)	731
Mupad [B] (verification not implemented)	732
Reduce [B] (verification not implemented)	732

Optimal result

Integrand size = 20, antiderivative size = 329

$$\begin{aligned}
 \int (a + bx)^{10}(A + Bx)(d + ex)^7 dx = & \frac{(Ab - aB)(bd - ae)^7(a + bx)^{11}}{11b^9} \\
 & + \frac{(bd - ae)^6(bBd + 7Abe - 8aBe)(a + bx)^{12}}{12b^9} \\
 & + \frac{7e(bd - ae)^5(bBd + 3Abe - 4aBe)(a + bx)^{13}}{13b^9} \\
 & + \frac{e^2(bd - ae)^4(3bBd + 5Abe - 8aBe)(a + bx)^{14}}{2b^9} \\
 & + \frac{7e^3(bd - ae)^3(bBd + Abe - 2aBe)(a + bx)^{15}}{3b^9} \\
 & + \frac{7e^4(bd - ae)^2(5bBd + 3Abe - 8aBe)(a + bx)^{16}}{16b^9} \\
 & + \frac{7e^5(bd - ae)(3bBd + Abe - 4aBe)(a + bx)^{17}}{17b^9} \\
 & + \frac{e^6(7bBd + Abe - 8aBe)(a + bx)^{18}}{18b^9} \\
 & + \frac{Be^7(a + bx)^{19}}{19b^9}
 \end{aligned}$$

output

```
1/11*(A*b-B*a)*(-a*e+b*d)^7*(b*x+a)^11/b^9+1/12*(-a*e+b*d)^6*(7*A*b*e-8*B*
a*e+B*b*d)*(b*x+a)^12/b^9+7/13*e*(-a*e+b*d)^5*(3*A*b*e-4*B*a*e+B*b*d)*(b*x
+a)^13/b^9+1/2*e^2*(-a*e+b*d)^4*(5*A*b*e-8*B*a*e+3*B*b*d)*(b*x+a)^14/b^9+7
/3*e^3*(-a*e+b*d)^3*(A*b*e-2*B*a*e+B*b*d)*(b*x+a)^15/b^9+7/16*e^4*(-a*e+b*
d)^2*(3*A*b*e-8*B*a*e+5*B*b*d)*(b*x+a)^16/b^9+7/17*e^5*(-a*e+b*d)*(A*b*e-4
*B*a*e+3*B*b*d)*(b*x+a)^17/b^9+1/18*e^6*(A*b*e-8*B*a*e+7*B*b*d)*(b*x+a)^18
/b^9+1/19*B*e^7*(b*x+a)^19/b^9
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 2034 vs. $2(329) = 658$.

Time = 0.47 (sec) , antiderivative size = 2034, normalized size of antiderivative = 6.18

$$\int (a + bx)^{10}(A + Bx)(d + ex)^7 dx = \text{Result too large to show}$$

input

```
Integrate[(a + b*x)^10*(A + B*x)*(d + e*x)^7,x]
```

output

```

a^10*A*d^7*x + (a^9*d^6*(10*A*b*d + a*B*d + 7*a*A*e)*x^2)/2 + (a^8*d^5*(a*
B*d*(10*b*d + 7*a*e) + A*(45*b^2*d^2 + 70*a*b*d*e + 21*a^2*e^2))*x^3)/3 +
(a^7*d^4*(a*B*d*(45*b^2*d^2 + 70*a*b*d*e + 21*a^2*e^2) + 5*A*(24*b^3*d^3 +
63*a*b^2*d^2*e + 42*a^2*b*d*e^2 + 7*a^3*e^3))*x^4)/4 + a^6*d^3*(a*B*d*(24
*b^3*d^3 + 63*a*b^2*d^2*e + 42*a^2*b*d*e^2 + 7*a^3*e^3) + 7*A*(6*b^4*d^4 +
24*a*b^3*d^3*e + 27*a^2*b^2*d^2*e^2 + 10*a^3*b*d*e^3 + a^4*e^4))*x^5 + (7
*a^5*d^2*(5*a*B*d*(6*b^4*d^4 + 24*a*b^3*d^3*e + 27*a^2*b^2*d^2*e^2 + 10*a^
3*b*d*e^3 + a^4*e^4) + A*(36*b^5*d^5 + 210*a*b^4*d^4*e + 360*a^2*b^3*d^3*e
^2 + 225*a^3*b^2*d^2*e^3 + 50*a^4*b*d*e^4 + 3*a^5*e^5))*x^6)/6 + a^4*d*(a*
B*d*(36*b^5*d^5 + 210*a*b^4*d^4*e + 360*a^2*b^3*d^3*e^2 + 225*a^3*b^2*d^2*
e^3 + 50*a^4*b*d*e^4 + 3*a^5*e^5) + A*(30*b^6*d^6 + 252*a*b^5*d^5*e + 630*
a^2*b^4*d^4*e^2 + 600*a^3*b^3*d^3*e^3 + 225*a^4*b^2*d^2*e^4 + 30*a^5*b*d*e
^5 + a^6*e^6))*x^7 + (a^3*(7*a*B*d*(30*b^6*d^6 + 252*a*b^5*d^5*e + 630*a^2
*b^4*d^4*e^2 + 600*a^3*b^3*d^3*e^3 + 225*a^4*b^2*d^2*e^4 + 30*a^5*b*d*e^5
+ a^6*e^6) + A*(120*b^7*d^7 + 1470*a*b^6*d^6*e + 5292*a^2*b^5*d^5*e^2 + 73
50*a^3*b^4*d^4*e^3 + 4200*a^4*b^3*d^3*e^4 + 945*a^5*b^2*d^2*e^5 + 70*a^6*b
*d*e^6 + a^7*e^7))*x^8)/8 + (a^2*(a*B*(120*b^7*d^7 + 1470*a*b^6*d^6*e + 52
92*a^2*b^5*d^5*e^2 + 7350*a^3*b^4*d^4*e^3 + 4200*a^4*b^3*d^3*e^4 + 945*a^5
*b^2*d^2*e^5 + 70*a^6*b*d*e^6 + a^7*e^7) + 5*A*b*(9*b^7*d^7 + 168*a*b^6*d^
6*e + 882*a^2*b^5*d^5*e^2 + 1764*a^3*b^4*d^4*e^3 + 1470*a^4*b^3*d^3*e^4...

```

Rubi [A] (verified)

Time = 2.09 (sec) , antiderivative size = 329, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^{10} (A + Bx)(d + ex)^7 dx$$

$$\downarrow 86$$

$$\int \left(\frac{e^6 (a + bx)^{17} (-8aBe + Abe + 7bBd)}{b^8} + \frac{7e^5 (a + bx)^{16} (bd - ae) (-4aBe + Abe + 3bBd)}{b^8} + \frac{7e^4 (a + bx)^{15} (bd)}{b^8} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned} & \frac{e^6(a+bx)^{18}(-8aBe + Abe + 7bBd)}{18b^9} + \frac{7e^5(a+bx)^{17}(bd-ae)(-4aBe + Abe + 3bBd)}{17b^9} + \\ & \frac{7e^4(a+bx)^{16}(bd-ae)^2(-8aBe + 3Abe + 5bBd)}{7e^4(a+bx)^{16}(bd-ae)^2(-8aBe + 3Abe + 5bBd)} + \\ & \frac{7e^3(a+bx)^{15}(bd-ae)^3(-2aBe + Abe + bBd)}{16b^9} + \\ & \frac{e^2(a+bx)^{14}(bd-ae)^4(-8aBe + 5Abe + 3bBd)}{3b^9} + \\ & \frac{7e(a+bx)^{13}(bd-ae)^5(-4aBe + 3Abe + bBd)}{2b^9} + \\ & \frac{(a+bx)^{12}(bd-ae)^6(-8aBe + 7Abe + bBd)}{12b^9} + \frac{(a+bx)^{11}(Ab - aB)(bd-ae)^7}{11b^9} + \\ & \frac{Be^7(a+bx)^{19}}{19b^9} \end{aligned}$$

input `Int[(a + b*x)^10*(A + B*x)*(d + e*x)^7, x]`

output `((A*b - a*B)*(b*d - a*e)^7*(a + b*x)^11)/(11*b^9) + ((b*d - a*e)^6*(b*B*d + 7*A*b*e - 8*a*B*e)*(a + b*x)^12)/(12*b^9) + (7*e*(b*d - a*e)^5*(b*B*d + 3*A*b*e - 4*a*B*e)*(a + b*x)^13)/(13*b^9) + (e^2*(b*d - a*e)^4*(3*b*B*d + 5*A*b*e - 8*a*B*e)*(a + b*x)^14)/(2*b^9) + (7*e^3*(b*d - a*e)^3*(b*B*d + A*b*e - 2*a*B*e)*(a + b*x)^15)/(3*b^9) + (7*e^4*(b*d - a*e)^2*(5*b*B*d + 3*A*b*e - 8*a*B*e)*(a + b*x)^16)/(16*b^9) + (7*e^5*(b*d - a*e)*(3*b*B*d + A*b*e - 4*a*B*e)*(a + b*x)^17)/(17*b^9) + (e^6*(7*b*B*d + A*b*e - 8*a*B*e)*(a + b*x)^18)/(18*b^9) + (B*e^7*(a + b*x)^19)/(19*b^9)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2188 vs. $2(311) = 622$.

Time = 0.25 (sec) , antiderivative size = 2189, normalized size of antiderivative = 6.65

method	result	size
default	Expression too large to display	2189
norman	Expression too large to display	2347
gospers	Expression too large to display	2784
risch	Expression too large to display	2784
parallemrisch	Expression too large to display	2784
orering	Expression too large to display	2784

input `int((b*x+a)^10*(B*x+A)*(e*x+d)^7,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 1/19*b^{10}*B*e^7*x^{19}+1/18*((A*b^{10}+10*B*a*b^9)*e^7+7*b^{10}*B*d*e^6)*x^{18}+1/ \\ & 17*((10*A*a*b^9+45*B*a^2*b^8)*e^7+7*(A*b^{10}+10*B*a*b^9)*d*e^6+21*b^{10}*B*d^2 \\ & *e^5)*x^{17}+1/16*((45*A*a^2*b^8+120*B*a^3*b^7)*e^7+7*(10*A*a*b^9+45*B*a^2*b^8) \\ & *d*e^6+21*(A*b^{10}+10*B*a*b^9)*d^2*e^5+35*b^{10}*B*d^3*e^4)*x^{16}+1/15*((1 \\ & 20*A*a^3*b^7+210*B*a^4*b^6)*e^7+7*(45*A*a^2*b^8+120*B*a^3*b^7)*d*e^6+21*(1 \\ & 0*A*a*b^9+45*B*a^2*b^8)*d^2*e^5+35*(A*b^{10}+10*B*a*b^9)*d^3*e^4+35*b^{10}*B*d^4 \\ & *e^3)*x^{15}+1/14*((210*A*a^4*b^6+252*B*a^5*b^5)*e^7+7*(120*A*a^3*b^7+210*B \\ & *a^4*b^6)*d*e^6+21*(45*A*a^2*b^8+120*B*a^3*b^7)*d^2*e^5+35*(10*A*a*b^9+45 \\ & *B*a^2*b^8)*d^3*e^4+35*(A*b^{10}+10*B*a*b^9)*d^4*e^3+21*b^{10}*B*d^5*e^2)*x^{14} \\ & +1/13*((252*A*a^5*b^5+210*B*a^6*b^4)*e^7+7*(210*A*a^4*b^6+252*B*a^5*b^5)*d \\ & *e^6+21*(120*A*a^3*b^7+210*B*a^4*b^6)*d^2*e^5+35*(45*A*a^2*b^8+120*B*a^3*b^7) \\ & *d^3*e^4+35*(10*A*a*b^9+45*B*a^2*b^8)*d^4*e^3+21*(A*b^{10}+10*B*a*b^9)*d^5 \\ & *e^2+7*b^{10}*B*d^6*e)*x^{13}+1/12*((210*A*a^6*b^4+120*B*a^7*b^3)*e^7+7*(252*A \\ & *a^5*b^5+210*B*a^6*b^4)*d*e^6+21*(210*A*a^4*b^6+252*B*a^5*b^5)*d^2*e^5+35 \\ & *(120*A*a^3*b^7+210*B*a^4*b^6)*d^3*e^4+35*(45*A*a^2*b^8+120*B*a^3*b^7)*d^4 \\ & *e^3+21*(10*A*a*b^9+45*B*a^2*b^8)*d^5*e^2+7*(A*b^{10}+10*B*a*b^9)*d^6*e+b^{10} \\ & *B*d^7)*x^{12}+1/11*((120*A*a^7*b^3+45*B*a^8*b^2)*e^7+7*(210*A*a^6*b^4+120*B \\ & *a^7*b^3)*d*e^6+21*(252*A*a^5*b^5+210*B*a^6*b^4)*d^2*e^5+35*(210*A*a^4*b^6 \\ & +252*B*a^5*b^5)*d^3*e^4+35*(120*A*a^3*b^7+210*B*a^4*b^6)*d^4*e^3+21*(45*A \\ & a^2*b^8+120*B*a^3*b^7)*d^5*e^2+7*(10*A*a*b^9+45*B*a^2*b^8)*d^6*e+(A*b^{10}+10*B \\ & *a*b^9)*d^7*e) \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2198 vs. $2(311) = 622$.

Time = 0.09 (sec) , antiderivative size = 2198, normalized size of antiderivative = 6.68

$$\int (a + bx)^{10}(A + Bx)(d + ex)^7 dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)*(e*x+d)^7,x, algorithm="fricas")`

output

```
1/19*B*b^10*e^7*x^19 + A*a^10*d^7*x + 1/18*(7*B*b^10*d*e^6 + (10*B*a*b^9 +
A*b^10)*e^7)*x^18 + 1/17*(21*B*b^10*d^2*e^5 + 7*(10*B*a*b^9 + A*b^10)*d*e
^6 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*e^7)*x^17 + 1/16*(35*B*b^10*d^3*e^4 + 21*
(10*B*a*b^9 + A*b^10)*d^2*e^5 + 35*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^6 + 15*(8
*B*a^3*b^7 + 3*A*a^2*b^8)*e^7)*x^16 + 1/3*(7*B*b^10*d^4*e^3 + 7*(10*B*a*b
^9 + A*b^10)*d^3*e^4 + 21*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^5 + 21*(8*B*a^3*b
^7 + 3*A*a^2*b^8)*d*e^6 + 6*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^7)*x^15 + 1/2*(3
*B*b^10*d^5*e^2 + 5*(10*B*a*b^9 + A*b^10)*d^4*e^3 + 25*(9*B*a^2*b^8 + 2*A
a*b^9)*d^3*e^4 + 45*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^5 + 30*(7*B*a^4*b^6
+ 4*A*a^3*b^7)*d*e^6 + 6*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^7)*x^14 + 7/13*(B*b
^10*d^6*e + 3*(10*B*a*b^9 + A*b^10)*d^5*e^2 + 25*(9*B*a^2*b^8 + 2*A*a*b^9)
*d^4*e^3 + 75*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^4 + 90*(7*B*a^4*b^6 + 4*A
a^3*b^7)*d^2*e^5 + 42*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^6 + 6*(5*B*a^6*b^4 +
6*A*a^5*b^5)*e^7)*x^13 + 1/12*(B*b^10*d^7 + 7*(10*B*a*b^9 + A*b^10)*d^6*e
+ 105*(9*B*a^2*b^8 + 2*A*a*b^9)*d^5*e^2 + 525*(8*B*a^3*b^7 + 3*A*a^2*b^8)
*d^4*e^3 + 1050*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^3*e^4 + 882*(6*B*a^5*b^5 + 5
*A*a^4*b^6)*d^2*e^5 + 294*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d*e^6 + 30*(4*B*a^7*
b^3 + 7*A*a^6*b^4)*e^7)*x^12 + 1/11*((10*B*a*b^9 + A*b^10)*d^7 + 35*(9*B*a
^2*b^8 + 2*A*a*b^9)*d^6*e + 315*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^5*e^2 + 1050
*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^4*e^3 + 1470*(6*B*a^5*b^5 + 5*A*a^4*b^6)...
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2824 vs. $2(335) = 670$.

Time = 0.25 (sec) , antiderivative size = 2824, normalized size of antiderivative = 8.58

$$\int (a + bx)^{10}(A + Bx)(d + ex)^7 dx = \text{Too large to display}$$

input `integrate((b*x+a)**10*(B*x+A)*(e*x+d)**7, x)`

output

```
A*a**10*d**7*x + B*b**10*e**7*x**19/19 + x**18*(A*b**10*e**7/18 + 5*B*a*b*
*9*e**7/9 + 7*B*b**10*d*e**6/18) + x**17*(10*A*a*b**9*e**7/17 + 7*A*b**10*
d*e**6/17 + 45*B*a**2*b**8*e**7/17 + 70*B*a*b**9*d*e**6/17 + 21*B*b**10*d*
*2*e**5/17) + x**16*(45*A*a**2*b**8*e**7/16 + 35*A*a*b**9*d*e**6/8 + 21*A*
b**10*d**2*e**5/16 + 15*B*a**3*b**7*e**7/2 + 315*B*a**2*b**8*d*e**6/16 + 1
05*B*a*b**9*d**2*e**5/8 + 35*B*b**10*d**3*e**4/16) + x**15*(8*A*a**3*b**7*
e**7 + 21*A*a**2*b**8*d*e**6 + 14*A*a*b**9*d**2*e**5 + 7*A*b**10*d**3*e**4
/3 + 14*B*a**4*b**6*e**7 + 56*B*a**3*b**7*d*e**6 + 63*B*a**2*b**8*d**2*e**
5 + 70*B*a*b**9*d**3*e**4/3 + 7*B*b**10*d**4*e**3/3) + x**14*(15*A*a**4*b*
*6*e**7 + 60*A*a**3*b**7*d*e**6 + 135*A*a**2*b**8*d**2*e**5/2 + 25*A*a*b**
9*d**3*e**4 + 5*A*b**10*d**4*e**3/2 + 18*B*a**5*b**5*e**7 + 105*B*a**4*b**
6*d*e**6 + 180*B*a**3*b**7*d**2*e**5 + 225*B*a**2*b**8*d**3*e**4/2 + 25*B*
a*b**9*d**4*e**3 + 3*B*b**10*d**5*e**2/2) + x**13*(252*A*a**5*b**5*e**7/13
+ 1470*A*a**4*b**6*d*e**6/13 + 2520*A*a**3*b**7*d**2*e**5/13 + 1575*A*a**
2*b**8*d**3*e**4/13 + 350*A*a*b**9*d**4*e**3/13 + 21*A*b**10*d**5*e**2/13
+ 210*B*a**6*b**4*e**7/13 + 1764*B*a**5*b**5*d*e**6/13 + 4410*B*a**4*b**6*
d**2*e**5/13 + 4200*B*a**3*b**7*d**3*e**4/13 + 1575*B*a**2*b**8*d**4*e**3/
13 + 210*B*a*b**9*d**5*e**2/13 + 7*B*b**10*d**6*e/13) + x**12*(35*A*a**6*b
**4*e**7/2 + 147*A*a**5*b**5*d*e**6 + 735*A*a**4*b**6*d**2*e**5/2 + 350*A*
a**3*b**7*d**3*e**4 + 525*A*a**2*b**8*d**4*e**3/4 + 35*A*a*b**9*d**5*e...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2198 vs. $2(311) = 622$.

Time = 0.05 (sec) , antiderivative size = 2198, normalized size of antiderivative = 6.68

$$\int (a + bx)^{10}(A + Bx)(d + ex)^7 dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)*(e*x+d)^7,x, algorithm="maxima")`

output

```
1/19*B*b^10*e^7*x^19 + A*a^10*d^7*x + 1/18*(7*B*b^10*d*e^6 + (10*B*a*b^9 +
A*b^10)*e^7)*x^18 + 1/17*(21*B*b^10*d^2*e^5 + 7*(10*B*a*b^9 + A*b^10)*d*e
^6 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*e^7)*x^17 + 1/16*(35*B*b^10*d^3*e^4 + 21*
(10*B*a*b^9 + A*b^10)*d^2*e^5 + 35*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^6 + 15*(8
*B*a^3*b^7 + 3*A*a^2*b^8)*e^7)*x^16 + 1/3*(7*B*b^10*d^4*e^3 + 7*(10*B*a*b^
9 + A*b^10)*d^3*e^4 + 21*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^5 + 21*(8*B*a^3*b
^7 + 3*A*a^2*b^8)*d*e^6 + 6*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^7)*x^15 + 1/2*(3
*B*b^10*d^5*e^2 + 5*(10*B*a*b^9 + A*b^10)*d^4*e^3 + 25*(9*B*a^2*b^8 + 2*A*
a*b^9)*d^3*e^4 + 45*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^5 + 30*(7*B*a^4*b^6
+ 4*A*a^3*b^7)*d*e^6 + 6*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^7)*x^14 + 7/13*(B*b
^10*d^6*e + 3*(10*B*a*b^9 + A*b^10)*d^5*e^2 + 25*(9*B*a^2*b^8 + 2*A*a*b^9)
*d^4*e^3 + 75*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^4 + 90*(7*B*a^4*b^6 + 4*A*
a^3*b^7)*d^2*e^5 + 42*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^6 + 6*(5*B*a^6*b^4 +
6*A*a^5*b^5)*e^7)*x^13 + 1/12*(B*b^10*d^7 + 7*(10*B*a*b^9 + A*b^10)*d^6*e
+ 105*(9*B*a^2*b^8 + 2*A*a*b^9)*d^5*e^2 + 525*(8*B*a^3*b^7 + 3*A*a^2*b^8)
*d^4*e^3 + 1050*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^3*e^4 + 882*(6*B*a^5*b^5 + 5
*A*a^4*b^6)*d^2*e^5 + 294*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d*e^6 + 30*(4*B*a^7*
b^3 + 7*A*a^6*b^4)*e^7)*x^12 + 1/11*((10*B*a*b^9 + A*b^10)*d^7 + 35*(9*B*a
^2*b^8 + 2*A*a*b^9)*d^6*e + 315*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^5*e^2 + 1050
*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^4*e^3 + 1470*(6*B*a^5*b^5 + 5*A*a^4*b^6)...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2783 vs. $2(311) = 622$.

Time = 0.13 (sec) , antiderivative size = 2783, normalized size of antiderivative = 8.46

$$\int (a + bx)^{10}(A + Bx)(d + ex)^7 dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)*(e*x+d)^7,x, algorithm="giac")`

output

```
1/19*B*b^10*e^7*x^19 + 7/18*B*b^10*d*e^6*x^18 + 5/9*B*a*b^9*e^7*x^18 + 1/1
8*A*b^10*e^7*x^18 + 21/17*B*b^10*d^2*e^5*x^17 + 70/17*B*a*b^9*d*e^6*x^17 +
7/17*A*b^10*d*e^6*x^17 + 45/17*B*a^2*b^8*e^7*x^17 + 10/17*A*a*b^9*e^7*x^1
7 + 35/16*B*b^10*d^3*e^4*x^16 + 105/8*B*a*b^9*d^2*e^5*x^16 + 21/16*A*b^10*
d^2*e^5*x^16 + 315/16*B*a^2*b^8*d*e^6*x^16 + 35/8*A*a*b^9*d*e^6*x^16 + 15/
2*B*a^3*b^7*e^7*x^16 + 45/16*A*a^2*b^8*e^7*x^16 + 7/3*B*b^10*d^4*e^3*x^15
+ 70/3*B*a*b^9*d^3*e^4*x^15 + 7/3*A*b^10*d^3*e^4*x^15 + 63*B*a^2*b^8*d^2*
e^5*x^15 + 14*A*a*b^9*d^2*e^5*x^15 + 56*B*a^3*b^7*d*e^6*x^15 + 21*A*a^2*b^8
*d*e^6*x^15 + 14*B*a^4*b^6*e^7*x^15 + 8*A*a^3*b^7*e^7*x^15 + 3/2*B*b^10*d^
5*e^2*x^14 + 25*B*a*b^9*d^4*e^3*x^14 + 5/2*A*b^10*d^4*e^3*x^14 + 225/2*B*a
^2*b^8*d^3*e^4*x^14 + 25*A*a*b^9*d^3*e^4*x^14 + 180*B*a^3*b^7*d^2*e^5*x^14
+ 135/2*A*a^2*b^8*d^2*e^5*x^14 + 105*B*a^4*b^6*d*e^6*x^14 + 60*A*a^3*b^7*
d*e^6*x^14 + 18*B*a^5*b^5*e^7*x^14 + 15*A*a^4*b^6*e^7*x^14 + 7/13*B*b^10*d
^6*e*x^13 + 210/13*B*a*b^9*d^5*e^2*x^13 + 21/13*A*b^10*d^5*e^2*x^13 + 1575
/13*B*a^2*b^8*d^4*e^3*x^13 + 350/13*A*a*b^9*d^4*e^3*x^13 + 4200/13*B*a^3*b
^7*d^3*e^4*x^13 + 1575/13*A*a^2*b^8*d^3*e^4*x^13 + 4410/13*B*a^4*b^6*d^2*
e^5*x^13 + 2520/13*A*a^3*b^7*d^2*e^5*x^13 + 1764/13*B*a^5*b^5*d*e^6*x^13 +
1470/13*A*a^4*b^6*d*e^6*x^13 + 210/13*B*a^6*b^4*e^7*x^13 + 252/13*A*a^5*b^
5*e^7*x^13 + 1/12*B*b^10*d^7*x^12 + 35/6*B*a*b^9*d^6*e*x^12 + 7/12*A*b^10*
d^6*e*x^12 + 315/4*B*a^2*b^8*d^5*e^2*x^12 + 35/2*A*a*b^9*d^5*e^2*x^12 + ...
```


Mupad [B] (verification not implemented)

Time = 1.42 (sec) , antiderivative size = 2316, normalized size of antiderivative = 7.04

$$\int (a + bx)^{10}(A + Bx)(d + ex)^7 dx = \text{Too large to display}$$

input `int((A + B*x)*(a + b*x)^10*(d + e*x)^7,x)`

output

```
x^5*(42*A*a^6*b^4*d^7 + 24*B*a^7*b^3*d^7 + 7*A*a^10*d^3*e^4 + 7*B*a^10*d^4
*e^3 + 168*A*a^7*b^3*d^6*e + 70*A*a^9*b*d^4*e^3 + 63*B*a^8*b^2*d^6*e + 42*
B*a^9*b*d^5*e^2 + 189*A*a^8*b^2*d^5*e^2) + x^15*(8*A*a^3*b^7*e^7 + 14*B*a^
4*b^6*e^7 + (7*A*b^10*d^3*e^4)/3 + (7*B*b^10*d^4*e^3)/3 + 14*A*a*b^9*d^2*e
^5 + 21*A*a^2*b^8*d*e^6 + (70*B*a*b^9*d^3*e^4)/3 + 56*B*a^3*b^7*d*e^6 + 63
*B*a^2*b^8*d^2*e^5) + x^9*((B*a^10*e^7)/9 + (10*A*a^9*b*e^7)/9 + 5*A*a^2*b
^8*d^7 + (40*B*a^3*b^7*d^7)/3 + (280*A*a^3*b^7*d^6*e)/3 + 35*A*a^8*b^2*d*e
^6 + (490*B*a^4*b^6*d^6*e)/3 + 490*A*a^4*b^6*d^5*e^2 + 980*A*a^5*b^5*d^4*e
^3 + (2450*A*a^6*b^4*d^3*e^4)/3 + 280*A*a^7*b^3*d^2*e^5 + 588*B*a^5*b^5*d^
5*e^2 + (2450*B*a^6*b^4*d^4*e^3)/3 + (1400*B*a^7*b^3*d^3*e^4)/3 + 105*B*a^
8*b^2*d^2*e^5 + (70*B*a^9*b*d*e^6)/9) + x^11*((A*b^10*d^7)/11 + (10*B*a*b^
9*d^7)/11 + (120*A*a^7*b^3*e^7)/11 + (45*B*a^8*b^2*e^7)/11 + (1470*A*a^6*b
^4*d*e^6)/11 + (315*B*a^2*b^8*d^6*e)/11 + (840*B*a^7*b^3*d*e^6)/11 + (945*
A*a^2*b^8*d^5*e^2)/11 + (4200*A*a^3*b^7*d^4*e^3)/11 + (7350*A*a^4*b^6*d^3*
e^4)/11 + (5292*A*a^5*b^5*d^2*e^5)/11 + (2520*B*a^3*b^7*d^5*e^2)/11 + (735
0*B*a^4*b^6*d^4*e^3)/11 + (8820*B*a^5*b^5*d^3*e^4)/11 + (4410*B*a^6*b^4*d^
2*e^5)/11 + (70*A*a*b^9*d^6*e)/11) + x^7*(A*a^10*d*e^6 + 30*A*a^4*b^6*d^7
+ 36*B*a^5*b^5*d^7 + 3*B*a^10*d^2*e^5 + 252*A*a^5*b^5*d^6*e + 30*A*a^9*b*d
^2*e^5 + 210*B*a^6*b^4*d^6*e + 50*B*a^9*b*d^3*e^4 + 630*A*a^6*b^4*d^5*e^2
+ 600*A*a^7*b^3*d^4*e^3 + 225*A*a^8*b^2*d^3*e^4 + 360*B*a^7*b^3*d^5*e^2...
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 1429, normalized size of antiderivative = 4.34

$$\int (a + bx)^{10}(A + Bx)(d + ex)^7 dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)*(e*x+d)^7,x)`

output

```
(x*(604656*a**11*d**7 + 2116296*a**11*d**6*e*x + 4232592*a**11*d**5*e**2*x
**2 + 5290740*a**11*d**4*e**3*x**3 + 4232592*a**11*d**3*e**4*x**4 + 211629
6*a**11*d**2*e**5*x**5 + 604656*a**11*d*e**6*x**6 + 75582*a**11*e**7*x**7
+ 3325608*a**10*b*d**7*x + 15519504*a**10*b*d**6*e*x**2 + 34918884*a**10*b
*d**5*e**2*x**3 + 46558512*a**10*b*d**4*e**3*x**4 + 38798760*a**10*b*d**3*
e**4*x**5 + 19953648*a**10*b*d**2*e**5*x**6 + 5819814*a**10*b*d*e**6*x**7
+ 739024*a**10*b*e**7*x**8 + 11085360*a**9*b**2*d**7*x**2 + 58198140*a**9*
b**2*d**6*e*x**3 + 139675536*a**9*b**2*d**5*e**2*x**4 + 193993800*a**9*b**
2*d**4*e**3*x**5 + 166280400*a**9*b**2*d**3*e**4*x**6 + 87297210*a**9*b**2
*d**2*e**5*x**7 + 25865840*a**9*b**2*d*e**6*x**8 + 3325608*a**9*b**2*e**7*
x**9 + 24942060*a**8*b**3*d**7*x**3 + 139675536*a**8*b**3*d**6*e*x**4 + 34
9188840*a**8*b**3*d**5*e**2*x**5 + 498841200*a**8*b**3*d**4*e**3*x**6 + 43
6486050*a**8*b**3*d**3*e**4*x**7 + 232792560*a**8*b**3*d**2*e**5*x**8 + 69
837768*a**8*b**3*d*e**6*x**9 + 9069840*a**8*b**3*e**7*x**10 + 39907296*a**
7*b**4*d**7*x**4 + 232792560*a**7*b**4*d**6*e*x**5 + 598609440*a**7*b**4*d
**5*e**2*x**6 + 872972100*a**7*b**4*d**4*e**3*x**7 + 775975200*a**7*b**4*d
**3*e**4*x**8 + 419026608*a**7*b**4*d**2*e**5*x**9 + 126977760*a**7*b**4*d
**e**6*x**10 + 16628040*a**7*b**4*e**7*x**11 + 46558512*a**6*b**5*d**7*x**5
+ 279351072*a**6*b**5*d**6*e*x**6 + 733296564*a**6*b**5*d**5*e**2*x**7 +
1086365280*a**6*b**5*d**4*e**3*x**8 + 977728752*a**6*b**5*d**3*e**4*x**...
```

3.72 $\int (a + bx)^{10}(A + Bx)(d + ex)^6 dx$

Optimal result	734
Mathematica [B] (verified)	735
Rubi [A] (verified)	736
Maple [B] (verified)	737
Fricas [B] (verification not implemented)	738
Sympy [B] (verification not implemented)	739
Maxima [B] (verification not implemented)	740
Giac [B] (verification not implemented)	741
Mupad [B] (verification not implemented)	742
Reduce [B] (verification not implemented)	743

Optimal result

Integrand size = 20, antiderivative size = 290

$$\begin{aligned}
 \int (a + bx)^{10}(A + Bx)(d + ex)^6 dx = & \frac{(Ab - aB)(bd - ae)^6(a + bx)^{11}}{11b^8} \\
 & + \frac{(bd - ae)^5(bBd + 6Abe - 7aBe)(a + bx)^{12}}{12b^8} \\
 & + \frac{3e(bd - ae)^4(2bBd + 5Abe - 7aBe)(a + bx)^{13}}{13b^8} \\
 & + \frac{5e^2(bd - ae)^3(3bBd + 4Abe - 7aBe)(a + bx)^{14}}{14b^8} \\
 & + \frac{e^3(bd - ae)^2(4bBd + 3Abe - 7aBe)(a + bx)^{15}}{3b^8} \\
 & + \frac{3e^4(bd - ae)(5bBd + 2Abe - 7aBe)(a + bx)^{16}}{16b^8} \\
 & + \frac{e^5(6bBd + Abe - 7aBe)(a + bx)^{17}}{17b^8} \\
 & + \frac{Be^6(a + bx)^{18}}{18b^8}
 \end{aligned}$$

output

```
1/11*(A*b-B*a)*(-a*e+b*d)^6*(b*x+a)^11/b^8+1/12*(-a*e+b*d)^5*(6*A*b*e-7*B*
a*e+B*b*d)*(b*x+a)^12/b^8+3/13*e*(-a*e+b*d)^4*(5*A*b*e-7*B*a*e+2*B*b*d)*(b
*x+a)^13/b^8+5/14*e^2*(-a*e+b*d)^3*(4*A*b*e-7*B*a*e+3*B*b*d)*(b*x+a)^14/b^
8+1/3*e^3*(-a*e+b*d)^2*(3*A*b*e-7*B*a*e+4*B*b*d)*(b*x+a)^15/b^8+3/16*e^4*(
-a*e+b*d)*(2*A*b*e-7*B*a*e+5*B*b*d)*(b*x+a)^16/b^8+1/17*e^5*(A*b*e-7*B*a*e
+6*B*b*d)*(b*x+a)^17/b^8+1/18*B*e^6*(b*x+a)^18/b^8
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1788 vs. $2(290) = 580$.

Time = 0.42 (sec) , antiderivative size = 1788, normalized size of antiderivative = 6.17

$$\int (a + bx)^{10}(A + Bx)(d + ex)^6 dx = \text{Too large to display}$$

input

```
Integrate[(a + b*x)^10*(A + B*x)*(d + e*x)^6,x]
```

output

```
a^10*A*d^6*x + (a^9*d^5*(10*A*b*d + a*B*d + 6*a*A*e)*x^2)/2 + (a^8*d^4*(2*
a*B*d*(5*b*d + 3*a*e) + 15*A*(3*b^2*d^2 + 4*a*b*d*e + a^2*e^2))*x^3)/3 + (
5*a^7*d^3*(3*a*B*d*(3*b^2*d^2 + 4*a*b*d*e + a^2*e^2) + A*(24*b^3*d^3 + 54*
a*b^2*d^2*e + 30*a^2*b*d*e^2 + 4*a^3*e^3))*x^4)/4 + a^6*d^2*(2*a*B*d*(12*b
^3*d^3 + 27*a*b^2*d^2*e + 15*a^2*b*d*e^2 + 2*a^3*e^3) + A*(42*b^4*d^4 + 14
4*a*b^3*d^3*e + 135*a^2*b^2*d^2*e^2 + 40*a^3*b*d*e^3 + 3*a^4*e^4))*x^5 + (
a^5*d*(5*a*B*d*(42*b^4*d^4 + 144*a*b^3*d^3*e + 135*a^2*b^2*d^2*e^2 + 40*a^
3*b*d*e^3 + 3*a^4*e^4) + 6*A*(42*b^5*d^5 + 210*a*b^4*d^4*e + 300*a^2*b^3*d
^3*e^2 + 150*a^3*b^2*d^2*e^3 + 25*a^4*b*d*e^4 + a^5*e^5))*x^6)/6 + (a^4*(6
*a*B*d*(42*b^5*d^5 + 210*a*b^4*d^4*e + 300*a^2*b^3*d^3*e^2 + 150*a^3*b^2*d
^2*e^3 + 25*a^4*b*d*e^4 + a^5*e^5) + A*(210*b^6*d^6 + 1512*a*b^5*d^5*e + 3
150*a^2*b^4*d^4*e^2 + 2400*a^3*b^3*d^3*e^3 + 675*a^4*b^2*d^2*e^4 + 60*a^5*
b*d*e^5 + a^6*e^6))*x^7)/7 + (a^3*(10*A*b*(12*b^6*d^6 + 126*a*b^5*d^5*e +
378*a^2*b^4*d^4*e^2 + 420*a^3*b^3*d^3*e^3 + 180*a^4*b^2*d^2*e^4 + 27*a^5*b
*d*e^5 + a^6*e^6) + a*B*(210*b^6*d^6 + 1512*a*b^5*d^5*e + 3150*a^2*b^4*d^4
*e^2 + 2400*a^3*b^3*d^3*e^3 + 675*a^4*b^2*d^2*e^4 + 60*a^5*b*d*e^5 + a^6*e
^6))*x^8)/8 + (5*a^2*b*(9*A*b*(b^6*d^6 + 16*a*b^5*d^5*e + 70*a^2*b^4*d^4*
e^2 + 112*a^3*b^3*d^3*e^3 + 70*a^4*b^2*d^2*e^4 + 16*a^5*b*d*e^5 + a^6*e^6)
+ 2*a*B*(12*b^6*d^6 + 126*a*b^5*d^5*e + 378*a^2*b^4*d^4*e^2 + 420*a^3*b^3*
d^3*e^3 + 180*a^4*b^2*d^2*e^4 + 27*a^5*b*d*e^5 + a^6*e^6))*x^9)/9 + (a*...
```

Rubi [A] (verified)

Time = 1.91 (sec) , antiderivative size = 290, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^{10}(A + Bx)(d + ex)^6 dx$$

↓ 86

$$\int \left(\frac{e^5(a + bx)^{16}(-7aBe + Abe + 6bBd)}{b^7} + \frac{3e^4(a + bx)^{15}(bd - ae)(-7aBe + 2Abe + 5bBd)}{b^7} + \frac{5e^3(a + bx)^{14}(bd - ae)^2(-7aBe + 3Abe + 4bBd)}{3b^8} + \frac{5e^2(a + bx)^{14}(bd - ae)^3(-7aBe + 4Abe + 3bBd)}{14b^8} + \frac{3e(a + bx)^{13}(bd - ae)^4(-7aBe + 5Abe + 2bBd)}{13b^8} + \frac{(a + bx)^{12}(bd - ae)^5(-7aBe + 6Abe + bBd)}{12b^8} + \frac{(a + bx)^{11}(Ab - aB)(bd - ae)^6}{11b^8} + \frac{Be^6(a + bx)^{18}}{18b^8} \right) dx$$

↓ 2009

$$\frac{e^5(a + bx)^{17}(-7aBe + Abe + 6bBd)}{17b^8} + \frac{3e^4(a + bx)^{16}(bd - ae)(-7aBe + 2Abe + 5bBd)}{16b^8} + \frac{e^3(a + bx)^{15}(bd - ae)^2(-7aBe + 3Abe + 4bBd)}{3b^8} + \frac{5e^2(a + bx)^{14}(bd - ae)^3(-7aBe + 4Abe + 3bBd)}{14b^8} + \frac{3e(a + bx)^{13}(bd - ae)^4(-7aBe + 5Abe + 2bBd)}{13b^8} + \frac{(a + bx)^{12}(bd - ae)^5(-7aBe + 6Abe + bBd)}{12b^8} + \frac{(a + bx)^{11}(Ab - aB)(bd - ae)^6}{11b^8} + \frac{Be^6(a + bx)^{18}}{18b^8}$$

input `Int[(a + b*x)^10*(A + B*x)*(d + e*x)^6,x]`

output `((A*b - a*B)*(b*d - a*e)^6*(a + b*x)^11)/(11*b^8) + ((b*d - a*e)^5*(b*B*d + 6*A*b*e - 7*a*B*e)*(a + b*x)^12)/(12*b^8) + (3*e*(b*d - a*e)^4*(2*b*B*d + 5*A*b*e - 7*a*B*e)*(a + b*x)^13)/(13*b^8) + (5*e^2*(b*d - a*e)^3*(3*b*B*d + 4*A*b*e - 7*a*B*e)*(a + b*x)^14)/(14*b^8) + (e^3*(b*d - a*e)^2*(4*b*B*d + 3*A*b*e - 7*a*B*e)*(a + b*x)^15)/(3*b^8) + (3*e^4*(b*d - a*e)*(5*b*B*d + 2*A*b*e - 7*a*B*e)*(a + b*x)^16)/(16*b^8) + (e^5*(6*b*B*d + A*b*e - 7*a*B*e)*(a + b*x)^17)/(17*b^8) + (B*e^6*(a + b*x)^18)/(18*b^8)`

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1904 vs. $2(274) = 548$.

Time = 0.24 (sec) , antiderivative size = 1905, normalized size of antiderivative = 6.57

method	result	size
default	Expression too large to display	1905
norman	Expression too large to display	2032
gospers	Expression too large to display	2408
risch	Expression too large to display	2408
parallelsch	Expression too large to display	2408
orering	Expression too large to display	2408

input

```
int((b*x+a)^10*(B*x+A)*(e*x+d)^6,x,method=_RETURNVERBOSE)
```

output

```

1/18*b^10*B*e^6*x^18+1/17*((A*b^10+10*B*a*b^9)*e^6+6*b^10*B*d*e^5)*x^17+1/
16*((10*A*a*b^9+45*B*a^2*b^8)*e^6+6*(A*b^10+10*B*a*b^9)*d*e^5+15*b^10*B*d^
2*e^4)*x^16+1/15*((45*A*a^2*b^8+120*B*a^3*b^7)*e^6+6*(10*A*a*b^9+45*B*a^2*
b^8)*d*e^5+15*(A*b^10+10*B*a*b^9)*d^2*e^4+20*b^10*B*d^3*e^3)*x^15+1/14*((1
20*A*a^3*b^7+210*B*a^4*b^6)*e^6+6*(45*A*a^2*b^8+120*B*a^3*b^7)*d*e^5+15*(1
0*A*a*b^9+45*B*a^2*b^8)*d^2*e^4+20*(A*b^10+10*B*a*b^9)*d^3*e^3+15*b^10*B*d
^4*e^2)*x^14+1/13*((210*A*a^4*b^6+252*B*a^5*b^5)*e^6+6*(120*A*a^3*b^7+210*
B*a^4*b^6)*d*e^5+15*(45*A*a^2*b^8+120*B*a^3*b^7)*d^2*e^4+20*(10*A*a*b^9+45
*B*a^2*b^8)*d^3*e^3+15*(A*b^10+10*B*a*b^9)*d^4*e^2+6*b^10*B*d^5*e)*x^13+1/
12*((252*A*a^5*b^5+210*B*a^6*b^4)*e^6+6*(210*A*a^4*b^6+252*B*a^5*b^5)*d*e^
5+15*(120*A*a^3*b^7+210*B*a^4*b^6)*d^2*e^4+20*(45*A*a^2*b^8+120*B*a^3*b^7)
*d^3*e^3+15*(10*A*a*b^9+45*B*a^2*b^8)*d^4*e^2+6*(A*b^10+10*B*a*b^9)*d^5*e+
b^10*B*d^6)*x^12+1/11*((210*A*a^6*b^4+120*B*a^7*b^3)*e^6+6*(252*A*a^5*b^5+
210*B*a^6*b^4)*d*e^5+15*(210*A*a^4*b^6+252*B*a^5*b^5)*d^2*e^4+20*(120*A*a^
3*b^7+210*B*a^4*b^6)*d^3*e^3+15*(45*A*a^2*b^8+120*B*a^3*b^7)*d^4*e^2+6*(10
*A*a*b^9+45*B*a^2*b^8)*d^5*e+(A*b^10+10*B*a*b^9)*d^6)*x^11+1/10*((120*A*a^
7*b^3+45*B*a^8*b^2)*e^6+6*(210*A*a^6*b^4+120*B*a^7*b^3)*d*e^5+15*(252*A*a^
5*b^5+210*B*a^6*b^4)*d^2*e^4+20*(210*A*a^4*b^6+252*B*a^5*b^5)*d^3*e^3+15*(
120*A*a^3*b^7+210*B*a^4*b^6)*d^4*e^2+6*(45*A*a^2*b^8+120*B*a^3*b^7)*d^5*e+
(10*A*a*b^9+45*B*a^2*b^8)*d^6)*x^10+1/9*((45*A*a^8*b^2+10*B*a^9*b)*e^6+...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1917 vs. $2(274) = 548$.

Time = 0.10 (sec) , antiderivative size = 1917, normalized size of antiderivative = 6.61

$$\int (a + bx)^{10}(A + Bx)(d + ex)^6 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^6,x, algorithm="fricas")
```

output

```

1/18*B*b^10*e^6*x^18 + A*a^10*d^6*x + 1/17*(6*B*b^10*d*e^5 + (10*B*a*b^9 +
A*b^10)*e^6)*x^17 + 1/16*(15*B*b^10*d^2*e^4 + 6*(10*B*a*b^9 + A*b^10)*d*e
^5 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*e^6)*x^16 + 1/3*(4*B*b^10*d^3*e^3 + 3*(10
*B*a*b^9 + A*b^10)*d^2*e^4 + 6*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^5 + 3*(8*B*a^
3*b^7 + 3*A*a^2*b^8)*e^6)*x^15 + 5/14*(3*B*b^10*d^4*e^2 + 4*(10*B*a*b^9 +
A*b^10)*d^3*e^3 + 15*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^4 + 18*(8*B*a^3*b^7 +
3*A*a^2*b^8)*d*e^5 + 6*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^6)*x^14 + 1/13*(6*B*
b^10*d^5*e + 15*(10*B*a*b^9 + A*b^10)*d^4*e^2 + 100*(9*B*a^2*b^8 + 2*A*a*b
^9)*d^3*e^3 + 225*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^4 + 180*(7*B*a^4*b^6 +
4*A*a^3*b^7)*d*e^5 + 42*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^6)*x^13 + 1/12*(B*b
^10*d^6 + 6*(10*B*a*b^9 + A*b^10)*d^5*e + 75*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4
*e^2 + 300*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^3 + 450*(7*B*a^4*b^6 + 4*A*a^
3*b^7)*d^2*e^4 + 252*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^5 + 42*(5*B*a^6*b^4 +
6*A*a^5*b^5)*e^6)*x^12 + 1/11*((10*B*a*b^9 + A*b^10)*d^6 + 30*(9*B*a^2*b^
8 + 2*A*a*b^9)*d^5*e + 225*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^4*e^2 + 600*(7*B*
a^4*b^6 + 4*A*a^3*b^7)*d^3*e^3 + 630*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e^4 +
252*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d*e^5 + 30*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e^
6)*x^11 + 1/2*((9*B*a^2*b^8 + 2*A*a*b^9)*d^6 + 18*(8*B*a^3*b^7 + 3*A*a^2*b^
8)*d^5*e + 90*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^4*e^2 + 168*(6*B*a^5*b^5 + 5*
A*a^4*b^6)*d^3*e^3 + 126*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^2*e^4 + 36*(4*B*...

```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2424 vs. $2(296) = 592$.

Time = 0.16 (sec) , antiderivative size = 2424, normalized size of antiderivative = 8.36

$$\int (a + bx)^{10}(A + Bx)(d + ex)^6 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)**10*(B*x+A)*(e*x+d)**6,x)
```


output

```
A*a**10*d**6*x + B*b**10*e**6*x**18/18 + x**17*(A*b**10*e**6/17 + 10*B*a*b
**9*e**6/17 + 6*B*b**10*d*e**5/17) + x**16*(5*A*a*b**9*e**6/8 + 3*A*b**10*
d*e**5/8 + 45*B*a**2*b**8*e**6/16 + 15*B*a*b**9*d*e**5/4 + 15*B*b**10*d**2
*e**4/16) + x**15*(3*A*a**2*b**8*e**6 + 4*A*a*b**9*d*e**5 + A*b**10*d**2*e
**4 + 8*B*a**3*b**7*e**6 + 18*B*a**2*b**8*d*e**5 + 10*B*a*b**9*d**2*e**4 +
4*B*b**10*d**3*e**3/3) + x**14*(60*A*a**3*b**7*e**6/7 + 135*A*a**2*b**8*d
e**5/7 + 75*A*a*b**9*d**2*e**4/7 + 10*A*b**10*d**3*e**3/7 + 15*B*a**4*b**
6*e**6 + 360*B*a**3*b**7*d*e**5/7 + 675*B*a**2*b**8*d**2*e**4/14 + 100*B*a
*b**9*d**3*e**3/7 + 15*B*b**10*d**4*e**2/14) + x**13*(210*A*a**4*b**6*e**6
/13 + 720*A*a**3*b**7*d*e**5/13 + 675*A*a**2*b**8*d**2*e**4/13 + 200*A*a*b
**9*d**3*e**3/13 + 15*A*b**10*d**4*e**2/13 + 252*B*a**5*b**5*e**6/13 + 126
0*B*a**4*b**6*d*e**5/13 + 1800*B*a**3*b**7*d**2*e**4/13 + 900*B*a**2*b**8*
d**3*e**3/13 + 150*B*a*b**9*d**4*e**2/13 + 6*B*b**10*d**5*e/13) + x**12*(2
1*A*a**5*b**5*e**6 + 105*A*a**4*b**6*d*e**5 + 150*A*a**3*b**7*d**2*e**4 +
75*A*a**2*b**8*d**3*e**3 + 25*A*a*b**9*d**4*e**2/2 + A*b**10*d**5*e/2 + 35
*B*a**6*b**4*e**6/2 + 126*B*a**5*b**5*d*e**5 + 525*B*a**4*b**6*d**2*e**4/2
+ 200*B*a**3*b**7*d**3*e**3 + 225*B*a**2*b**8*d**4*e**2/4 + 5*B*a*b**9*d*
*5*e + B*b**10*d**6/12) + x**11*(210*A*a**6*b**4*e**6/11 + 1512*A*a**5*b**
5*d*e**5/11 + 3150*A*a**4*b**6*d**2*e**4/11 + 2400*A*a**3*b**7*d**3*e**3/1
1 + 675*A*a**2*b**8*d**4*e**2/11 + 60*A*a*b**9*d**5*e/11 + A*b**10*d**6...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1917 vs. $2(274) = 548$.

Time = 0.06 (sec) , antiderivative size = 1917, normalized size of antiderivative = 6.61

$$\int (a + bx)^{10}(A + Bx)(d + ex)^6 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^6,x, algorithm="maxima")
```

output

```

1/18*B*b^10*e^6*x^18 + A*a^10*d^6*x + 1/17*(6*B*b^10*d*e^5 + (10*B*a*b^9 +
A*b^10)*e^6)*x^17 + 1/16*(15*B*b^10*d^2*e^4 + 6*(10*B*a*b^9 + A*b^10)*d*e
^5 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*e^6)*x^16 + 1/3*(4*B*b^10*d^3*e^3 + 3*(10
*B*a*b^9 + A*b^10)*d^2*e^4 + 6*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^5 + 3*(8*B*a^
3*b^7 + 3*A*a^2*b^8)*e^6)*x^15 + 5/14*(3*B*b^10*d^4*e^2 + 4*(10*B*a*b^9 +
A*b^10)*d^3*e^3 + 15*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^4 + 18*(8*B*a^3*b^7 +
3*A*a^2*b^8)*d*e^5 + 6*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^6)*x^14 + 1/13*(6*B*
b^10*d^5*e + 15*(10*B*a*b^9 + A*b^10)*d^4*e^2 + 100*(9*B*a^2*b^8 + 2*A*a*b
^9)*d^3*e^3 + 225*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^4 + 180*(7*B*a^4*b^6 +
4*A*a^3*b^7)*d*e^5 + 42*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^6)*x^13 + 1/12*(B*b
^10*d^6 + 6*(10*B*a*b^9 + A*b^10)*d^5*e + 75*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4
*e^2 + 300*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^3 + 450*(7*B*a^4*b^6 + 4*A*a^
3*b^7)*d^2*e^4 + 252*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^5 + 42*(5*B*a^6*b^4 +
6*A*a^5*b^5)*e^6)*x^12 + 1/11*((10*B*a*b^9 + A*b^10)*d^6 + 30*(9*B*a^2*b^
8 + 2*A*a*b^9)*d^5*e + 225*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^4*e^2 + 600*(7*B*
a^4*b^6 + 4*A*a^3*b^7)*d^3*e^3 + 630*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e^4 +
252*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d*e^5 + 30*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e^
6)*x^11 + 1/2*((9*B*a^2*b^8 + 2*A*a*b^9)*d^6 + 18*(8*B*a^3*b^7 + 3*A*a^2*b^
8)*d^5*e + 90*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^4*e^2 + 168*(6*B*a^5*b^5 + 5*
A*a^4*b^6)*d^3*e^3 + 126*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^2*e^4 + 36*(4*B*...

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2407 vs. $2(274) = 548$.

Time = 0.13 (sec) , antiderivative size = 2407, normalized size of antiderivative = 8.30

$$\int (a + bx)^{10}(A + Bx)(d + ex)^6 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^6,x, algorithm="giac")
```

output

```

1/18*B*b^10*e^6*x^18 + 6/17*B*b^10*d*e^5*x^17 + 10/17*B*a*b^9*e^6*x^17 + 1
/17*A*b^10*e^6*x^17 + 15/16*B*b^10*d^2*e^4*x^16 + 15/4*B*a*b^9*d*e^5*x^16
+ 3/8*A*b^10*d*e^5*x^16 + 45/16*B*a^2*b^8*e^6*x^16 + 5/8*A*a*b^9*e^6*x^16
+ 4/3*B*b^10*d^3*e^3*x^15 + 10*B*a*b^9*d^2*e^4*x^15 + A*b^10*d^2*e^4*x^15
+ 18*B*a^2*b^8*d*e^5*x^15 + 4*A*a*b^9*d*e^5*x^15 + 8*B*a^3*b^7*e^6*x^15 +
3*A*a^2*b^8*e^6*x^15 + 15/14*B*b^10*d^4*e^2*x^14 + 100/7*B*a*b^9*d^3*e^3*x
^14 + 10/7*A*b^10*d^3*e^3*x^14 + 675/14*B*a^2*b^8*d^2*e^4*x^14 + 75/7*A*a*
b^9*d^2*e^4*x^14 + 360/7*B*a^3*b^7*d*e^5*x^14 + 135/7*A*a^2*b^8*d*e^5*x^14
+ 15*B*a^4*b^6*e^6*x^14 + 60/7*A*a^3*b^7*e^6*x^14 + 6/13*B*b^10*d^5*e*x^1
3 + 150/13*B*a*b^9*d^4*e^2*x^13 + 15/13*A*b^10*d^4*e^2*x^13 + 900/13*B*a^2
*b^8*d^3*e^3*x^13 + 200/13*A*a*b^9*d^3*e^3*x^13 + 1800/13*B*a^3*b^7*d^2*e^
4*x^13 + 675/13*A*a^2*b^8*d^2*e^4*x^13 + 1260/13*B*a^4*b^6*d*e^5*x^13 + 72
0/13*A*a^3*b^7*d*e^5*x^13 + 252/13*B*a^5*b^5*e^6*x^13 + 210/13*A*a^4*b^6*e
^6*x^13 + 1/12*B*b^10*d^6*x^12 + 5*B*a*b^9*d^5*e*x^12 + 1/2*A*b^10*d^5*e*x
^12 + 225/4*B*a^2*b^8*d^4*e^2*x^12 + 25/2*A*a*b^9*d^4*e^2*x^12 + 200*B*a^3
*b^7*d^3*e^3*x^12 + 75*A*a^2*b^8*d^3*e^3*x^12 + 525/2*B*a^4*b^6*d^2*e^4*x^
12 + 150*A*a^3*b^7*d^2*e^4*x^12 + 126*B*a^5*b^5*d*e^5*x^12 + 105*A*a^4*b^6
*d*e^5*x^12 + 35/2*B*a^6*b^4*e^6*x^12 + 21*A*a^5*b^5*e^6*x^12 + 10/11*B*a*
b^9*d^6*x^11 + 1/11*A*b^10*d^6*x^11 + 270/11*B*a^2*b^8*d^5*e*x^11 + 60/11*
A*a*b^9*d^5*e*x^11 + 1800/11*B*a^3*b^7*d^4*e^2*x^11 + 675/11*A*a^2*b^8*...

```

Mupad [B] (verification not implemented)

Time = 1.40 (sec) , antiderivative size = 2001, normalized size of antiderivative = 6.90

$$\int (a + bx)^{10}(A + Bx)(d + ex)^6 dx = \text{Too large to display}$$

input

```
int((A + B*x)*(a + b*x)^10*(d + e*x)^6,x)
```

output

```

x^6*(A*a^10*d*e^5 + 42*A*a^5*b^5*d^6 + 35*B*a^6*b^4*d^6 + (5*B*a^10*d^2*e^
4)/2 + 210*A*a^6*b^4*d^5*e + 25*A*a^9*b*d^2*e^4 + 120*B*a^7*b^3*d^5*e + (1
00*B*a^9*b*d^3*e^3)/3 + 300*A*a^7*b^3*d^4*e^2 + 150*A*a^8*b^2*d^3*e^3 + (2
25*B*a^8*b^2*d^4*e^2)/2) + x^13*((6*B*b^10*d^5*e)/13 + (210*A*a^4*b^6*e^6)
/13 + (252*B*a^5*b^5*e^6)/13 + (15*A*b^10*d^4*e^2)/13 + (200*A*a*b^9*d^3*e
^3)/13 + (720*A*a^3*b^7*d*e^5)/13 + (150*B*a*b^9*d^4*e^2)/13 + (1260*B*a^4
*b^6*d*e^5)/13 + (675*A*a^2*b^8*d^2*e^4)/13 + (900*B*a^2*b^8*d^3*e^3)/13 +
(1800*B*a^3*b^7*d^2*e^4)/13) + x^5*(42*A*a^6*b^4*d^6 + 24*B*a^7*b^3*d^6 +
3*A*a^10*d^2*e^4 + 4*B*a^10*d^3*e^3 + 144*A*a^7*b^3*d^5*e + 40*A*a^9*b*d^
3*e^3 + 54*B*a^8*b^2*d^5*e + 30*B*a^9*b*d^4*e^2 + 135*A*a^8*b^2*d^4*e^2) +
x^14*((60*A*a^3*b^7*e^6)/7 + 15*B*a^4*b^6*e^6 + (10*A*b^10*d^3*e^3)/7 + (
15*B*b^10*d^4*e^2)/14 + (75*A*a*b^9*d^2*e^4)/7 + (135*A*a^2*b^8*d*e^5)/7 +
(100*B*a*b^9*d^3*e^3)/7 + (360*B*a^3*b^7*d*e^5)/7 + (675*B*a^2*b^8*d^2*e^
4)/14) + x^7*((A*a^10*e^6)/7 + (6*B*a^10*d*e^5)/7 + 30*A*a^4*b^6*d^6 + 36*
B*a^5*b^5*d^6 + 216*A*a^5*b^5*d^5*e + 180*B*a^6*b^4*d^5*e + (150*B*a^9*b*d
^2*e^4)/7 + 450*A*a^6*b^4*d^4*e^2 + (2400*A*a^7*b^3*d^3*e^3)/7 + (675*A*a^
8*b^2*d^2*e^4)/7 + (1800*B*a^7*b^3*d^4*e^2)/7 + (900*B*a^8*b^2*d^3*e^3)/7
+ (60*A*a^9*b*d*e^5)/7) + x^12*((B*b^10*d^6)/12 + (A*b^10*d^5*e)/2 + 21*A*
a^5*b^5*e^6 + (35*B*a^6*b^4*e^6)/2 + (25*A*a*b^9*d^4*e^2)/2 + 105*A*a^4*b^
6*d*e^5 + 126*B*a^5*b^5*d*e^5 + 75*A*a^2*b^8*d^3*e^3 + 150*A*a^3*b^7*d^...

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 1235, normalized size of antiderivative = 4.26

$$\int (a + bx)^{10}(A + Bx)(d + ex)^6 dx = \text{Too large to display}$$

input

```
int((b*x+a)^10*(B*x+A)*(e*x+d)^6,x)
```

output

```
(x*(222768*a**11*d**6 + 668304*a**11*d**5*e*x + 1113840*a**11*d**4*e**2*x*
*2 + 1113840*a**11*d**3*e**3*x**3 + 668304*a**11*d**2*e**4*x**4 + 222768*a
**11*d*e**5*x**5 + 31824*a**11*e**6*x**6 + 1225224*a**10*b*d**6*x + 490089
6*a**10*b*d**5*e*x**2 + 9189180*a**10*b*d**4*e**2*x**3 + 9801792*a**10*b*d
**3*e**3*x**4 + 6126120*a**10*b*d**2*e**4*x**5 + 2100384*a**10*b*d*e**5*x*
*6 + 306306*a**10*b*e**6*x**7 + 4084080*a**9*b**2*d**6*x**2 + 18378360*a**
9*b**2*d**5*e*x**3 + 36756720*a**9*b**2*d**4*e**2*x**4 + 40840800*a**9*b**
2*d**3*e**3*x**5 + 26254800*a**9*b**2*d**2*e**4*x**6 + 9189180*a**9*b**2*d
*e**5*x**7 + 1361360*a**9*b**2*e**6*x**8 + 9189180*a**8*b**3*d**6*x**3 + 4
4108064*a**8*b**3*d**5*e*x**4 + 91891800*a**8*b**3*d**4*e**2*x**5 + 105019
200*a**8*b**3*d**3*e**3*x**6 + 68918850*a**8*b**3*d**2*e**4*x**7 + 2450448
0*a**8*b**3*d*e**5*x**8 + 3675672*a**8*b**3*e**6*x**9 + 14702688*a**7*b**4
*d**6*x**4 + 73513440*a**7*b**4*d**5*e*x**5 + 157528800*a**7*b**4*d**4*e**
2*x**6 + 183783600*a**7*b**4*d**3*e**3*x**7 + 122522400*a**7*b**4*d**2*e**
4*x**8 + 44108064*a**7*b**4*d*e**5*x**9 + 6683040*a**7*b**4*e**6*x**10 + 1
7153136*a**6*b**5*d**6*x**5 + 88216128*a**6*b**5*d**5*e*x**6 + 192972780*a
**6*b**5*d**4*e**2*x**7 + 228708480*a**6*b**5*d**3*e**3*x**8 + 154378224*a
**6*b**5*d**2*e**4*x**9 + 56137536*a**6*b**5*d*e**5*x**10 + 8576568*a**6*b
**5*e**6*x**11 + 14702688*a**5*b**6*d**6*x**6 + 77189112*a**5*b**6*d**5*e*
x**7 + 171531360*a**5*b**6*d**4*e**2*x**8 + 205837632*a**5*b**6*d**3*e...
```

3.73 $\int (a + bx)^{10}(A + Bx)(d + ex)^5 dx$

Optimal result	745
Mathematica [B] (verified)	746
Rubi [A] (verified)	747
Maple [B] (verified)	748
Fricas [B] (verification not implemented)	749
Sympy [B] (verification not implemented)	750
Maxima [B] (verification not implemented)	751
Giac [B] (verification not implemented)	752
Mupad [B] (verification not implemented)	753
Reduce [B] (verification not implemented)	754

Optimal result

Integrand size = 20, antiderivative size = 243

$$\int (a + bx)^{10}(A + Bx)(d + ex)^5 dx = \frac{(Ab - aB)(bd - ae)^5(a + bx)^{11}}{11b^7} + \frac{(bd - ae)^4(bBd + 5Abe - 6aBe)(a + bx)^{12}}{12b^7} + \frac{5e(bd - ae)^3(bBd + 2Abe - 3aBe)(a + bx)^{13}}{13b^7} + \frac{5e^2(bd - ae)^2(bBd + Abe - 2aBe)(a + bx)^{14}}{7b^7} + \frac{e^3(bd - ae)(2bBd + Abe - 3aBe)(a + bx)^{15}}{3b^7} + \frac{e^4(5bBd + Abe - 6aBe)(a + bx)^{16}}{16b^7} + \frac{Be^5(a + bx)^{17}}{17b^7}$$

output

```
1/11*(A*b-B*a)*(-a*e+b*d)^5*(b*x+a)^11/b^7+1/12*(-a*e+b*d)^4*(5*A*b*e-6*B*
a*e+B*b*d)*(b*x+a)^12/b^7+5/13*e*(-a*e+b*d)^3*(2*A*b*e-3*B*a*e+B*b*d)*(b*x
+a)^13/b^7+5/7*e^2*(-a*e+b*d)^2*(A*b*e-2*B*a*e+B*b*d)*(b*x+a)^14/b^7+1/3*e
^3*(-a*e+b*d)*(A*b*e-3*B*a*e+2*B*b*d)*(b*x+a)^15/b^7+1/16*e^4*(A*b*e-6*B*a
e+5*B*b*d)*(b*x+a)^16/b^7+1/17*B*e^5*(b*x+a)^17/b^7
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1509 vs. $2(243) = 486$.

Time = 0.34 (sec) , antiderivative size = 1509, normalized size of antiderivative = 6.21

$$\int (a + bx)^{10}(A + Bx)(d + ex)^5 dx = \text{Too large to display}$$

input `Integrate[(a + b*x)^10*(A + B*x)*(d + e*x)^5,x]`

output

```
a^10*A*d^5*x + (a^9*d^4*(a*B*d + 5*A*(2*b*d + a*e))*x^2)/2 + (5*a^8*d^3*(a
*B*d*(2*b*d + a*e) + A*(9*b^2*d^2 + 10*a*b*d*e + 2*a^2*e^2))*x^3)/3 + (5*a
^7*d^2*(a*B*d*(9*b^2*d^2 + 10*a*b*d*e + 2*a^2*e^2) + A*(24*b^3*d^3 + 45*a*
b^2*d^2*e + 20*a^2*b*d*e^2 + 2*a^3*e^3))*x^4)/4 + a^6*d*(a*B*d*(24*b^3*d^3
+ 45*a*b^2*d^2*e + 20*a^2*b*d*e^2 + 2*a^3*e^3) + A*(42*b^4*d^4 + 120*a*b^
3*d^3*e + 90*a^2*b^2*d^2*e^2 + 20*a^3*b*d*e^3 + a^4*e^4))*x^5 + (a^5*(5*a*
B*d*(42*b^4*d^4 + 120*a*b^3*d^3*e + 90*a^2*b^2*d^2*e^2 + 20*a^3*b*d*e^3 +
a^4*e^4) + A*(252*b^5*d^5 + 1050*a*b^4*d^4*e + 1200*a^2*b^3*d^3*e^2 + 450*
a^3*b^2*d^2*e^3 + 50*a^4*b*d*e^4 + a^5*e^5))*x^6)/6 + (a^4*(a*B*(252*b^5*d
^5 + 1050*a*b^4*d^4*e + 1200*a^2*b^3*d^3*e^2 + 450*a^3*b^2*d^2*e^3 + 50*a^
4*b*d*e^4 + a^5*e^5) + 5*A*b*(42*b^5*d^5 + 252*a*b^4*d^4*e + 420*a^2*b^3*d
^3*e^2 + 240*a^3*b^2*d^2*e^3 + 45*a^4*b*d*e^4 + 2*a^5*e^5))*x^7)/7 + (5*a^
3*b*(a*B*(42*b^5*d^5 + 252*a*b^4*d^4*e + 420*a^2*b^3*d^3*e^2 + 240*a^3*b^2
*d^2*e^3 + 45*a^4*b*d*e^4 + 2*a^5*e^5) + 3*A*b*(8*b^5*d^5 + 70*a*b^4*d^4*e
+ 168*a^2*b^3*d^3*e^2 + 140*a^3*b^2*d^2*e^3 + 40*a^4*b*d*e^4 + 3*a^5*e^5)
)*x^8)/8 + (5*a^2*b^2*(a*B*(8*b^5*d^5 + 70*a*b^4*d^4*e + 168*a^2*b^3*d^3*e
^2 + 140*a^3*b^2*d^2*e^3 + 40*a^4*b*d*e^4 + 3*a^5*e^5) + A*b*(3*b^5*d^5 +
40*a*b^4*d^4*e + 140*a^2*b^3*d^3*e^2 + 168*a^3*b^2*d^2*e^3 + 70*a^4*b*d*e^
4 + 8*a^5*e^5))*x^9)/3 + (a*b^3*(3*a*B*(3*b^5*d^5 + 40*a*b^4*d^4*e + 140*a
^2*b^3*d^3*e^2 + 168*a^3*b^2*d^2*e^3 + 70*a^4*b*d*e^4 + 8*a^5*e^5) + A*...
```

Rubi [A] (verified)

Time = 1.64 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^{10}(A + Bx)(d + ex)^5 dx$$

↓ 86

$$\int \left(\frac{e^4(a + bx)^{15}(-6aBe + Abe + 5bBd)}{b^6} + \frac{5e^3(a + bx)^{14}(bd - ae)(-3aBe + Abe + 2bBd)}{b^6} + \frac{10e^2(a + bx)^{13}(bd - ae)^2(-2aBe + Abe + bBd)}{b^6} + \frac{5e(a + bx)^{12}(bd - ae)^3(-3aBe + 2Abe + bBd)}{b^6} + \frac{(a + bx)^{11}(Ab - aB)(bd - ae)^4}{b^6} + \frac{Be^5(a + bx)^{10}}{b^6} \right) dx$$

↓ 2009

$$\frac{e^4(a + bx)^{16}(-6aBe + Abe + 5bBd)}{16b^7} + \frac{e^3(a + bx)^{15}(bd - ae)(-3aBe + Abe + 2bBd)}{3b^7} + \frac{5e^2(a + bx)^{14}(bd - ae)^2(-2aBe + Abe + bBd)}{7b^7} + \frac{5e(a + bx)^{13}(bd - ae)^3(-3aBe + 2Abe + bBd)}{13b^7} + \frac{(a + bx)^{12}(bd - ae)^4(-6aBe + 5Abe + bBd)}{12b^7} + \frac{(a + bx)^{11}(Ab - aB)(bd - ae)^5}{11b^7} + \frac{Be^5(a + bx)^{17}}{17b^7}$$

input `Int[(a + b*x)^10*(A + B*x)*(d + e*x)^5,x]`

output `((A*b - a*B)*(b*d - a*e)^5*(a + b*x)^11)/(11*b^7) + ((b*d - a*e)^4*(b*B*d + 5*A*b*e - 6*a*B*e)*(a + b*x)^12)/(12*b^7) + (5*e*(b*d - a*e)^3*(b*B*d + 2*A*b*e - 3*a*B*e)*(a + b*x)^13)/(13*b^7) + (5*e^2*(b*d - a*e)^2*(b*B*d + A*b*e - 2*a*B*e)*(a + b*x)^14)/(7*b^7) + (e^3*(b*d - a*e)*(2*b*B*d + A*b*e - 3*a*B*e)*(a + b*x)^15)/(3*b^7) + (e^4*(5*b*B*d + A*b*e - 6*a*B*e)*(a + b*x)^16)/(16*b^7) + (B*e^5*(a + b*x)^17)/(17*b^7)`

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0]) || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1620 vs. $2(229) = 458$.

Time = 0.23 (sec) , antiderivative size = 1621, normalized size of antiderivative = 6.67

method	result	size
default	Expression too large to display	1621
norman	Expression too large to display	1718
orering	Expression too large to display	2032
gosper	Expression too large to display	2033
risch	Expression too large to display	2033
parallelrisc	Expression too large to display	2033

input

```
int((b*x+a)^10*(B*x+A)*(e*x+d)^5,x,method=_RETURNVERBOSE)
```

output

```

1/17*b^10*B*e^5*x^17+1/16*((A*b^10+10*B*a*b^9)*e^5+5*b^10*B*d*e^4)*x^16+1/
15*((10*A*a*b^9+45*B*a^2*b^8)*e^5+5*(A*b^10+10*B*a*b^9)*d*e^4+10*b^10*B*d^
2*e^3)*x^15+1/14*((45*A*a^2*b^8+120*B*a^3*b^7)*e^5+5*(10*A*a*b^9+45*B*a^2*
b^8)*d*e^4+10*(A*b^10+10*B*a*b^9)*d^2*e^3+10*b^10*B*d^3*e^2)*x^14+1/13*((1
20*A*a^3*b^7+210*B*a^4*b^6)*e^5+5*(45*A*a^2*b^8+120*B*a^3*b^7)*d*e^4+10*(1
0*A*a*b^9+45*B*a^2*b^8)*d^2*e^3+10*(A*b^10+10*B*a*b^9)*d^3*e^2+5*b^10*B*d^
4*e)*x^13+1/12*((210*A*a^4*b^6+252*B*a^5*b^5)*e^5+5*(120*A*a^3*b^7+210*B*a
^4*b^6)*d*e^4+10*(45*A*a^2*b^8+120*B*a^3*b^7)*d^2*e^3+10*(10*A*a*b^9+45*B*
a^2*b^8)*d^3*e^2+5*(A*b^10+10*B*a*b^9)*d^4*e+b^10*B*d^5)*x^12+1/11*((252*A
*a^5*b^5+210*B*a^6*b^4)*e^5+5*(210*A*a^4*b^6+252*B*a^5*b^5)*d*e^4+10*(120*
A*a^3*b^7+210*B*a^4*b^6)*d^2*e^3+10*(45*A*a^2*b^8+120*B*a^3*b^7)*d^3*e^2+5
*(10*A*a*b^9+45*B*a^2*b^8)*d^4*e+(A*b^10+10*B*a*b^9)*d^5)*x^11+1/10*((210*
A*a^6*b^4+120*B*a^7*b^3)*e^5+5*(252*A*a^5*b^5+210*B*a^6*b^4)*d*e^4+10*(210
*A*a^4*b^6+252*B*a^5*b^5)*d^2*e^3+10*(120*A*a^3*b^7+210*B*a^4*b^6)*d^3*e^2
+5*(45*A*a^2*b^8+120*B*a^3*b^7)*d^4*e+(10*A*a*b^9+45*B*a^2*b^8)*d^5)*x^10+
1/9*((120*A*a^7*b^3+45*B*a^8*b^2)*e^5+5*(210*A*a^6*b^4+120*B*a^7*b^3)*d*e^
4+10*(252*A*a^5*b^5+210*B*a^6*b^4)*d^2*e^3+10*(120*A*a^4*b^6+252*B*a^5*b^5
)*d^3*e^2+5*(120*A*a^3*b^7+210*B*a^4*b^6)*d^4*e+(45*A*a^2*b^8+120*B*a^3*b^
7)*d^5)*x^9+1/8*((45*A*a^8*b^2+10*B*a^9*b)*e^5+5*(120*A*a^7*b^3+45*B*a^8*b
^2)*d*e^4+10*(210*A*a^6*b^4+120*B*a^7*b^3)*d^2*e^3+10*(252*A*a^5*b^5+21...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1625 vs. $2(229) = 458$.

Time = 0.10 (sec) , antiderivative size = 1625, normalized size of antiderivative = 6.69

$$\int (a + bx)^{10}(A + Bx)(d + ex)^5 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^5,x, algorithm="fricas")
```

output

```

1/17*B*b^10*e^5*x^17 + A*a^10*d^5*x + 1/16*(5*B*b^10*d*e^4 + (10*B*a*b^9 +
A*b^10)*e^5)*x^16 + 1/3*(2*B*b^10*d^2*e^3 + (10*B*a*b^9 + A*b^10)*d*e^4 +
(9*B*a^2*b^8 + 2*A*a*b^9)*e^5)*x^15 + 5/14*(2*B*b^10*d^3*e^2 + 2*(10*B*a*
b^9 + A*b^10)*d^2*e^3 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^4 + 3*(8*B*a^3*b^7
+ 3*A*a^2*b^8)*e^5)*x^14 + 5/13*(B*b^10*d^4*e + 2*(10*B*a*b^9 + A*b^10)*d
^3*e^2 + 10*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^3 + 15*(8*B*a^3*b^7 + 3*A*a^2*
b^8)*d*e^4 + 6*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^5)*x^13 + 1/12*(B*b^10*d^5 +
5*(10*B*a*b^9 + A*b^10)*d^4*e + 50*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^2 + 150
*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^3 + 150*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e
^4 + 42*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^5)*x^12 + 1/11*((10*B*a*b^9 + A*b^10
)*d^5 + 25*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e + 150*(8*B*a^3*b^7 + 3*A*a^2*b^
8)*d^3*e^2 + 300*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^3 + 210*(6*B*a^5*b^5 +
5*A*a^4*b^6)*d*e^4 + 42*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^5)*x^11 + 1/2*((9*B*
a^2*b^8 + 2*A*a*b^9)*d^5 + 15*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^4*e + 60*(7*B*
a^4*b^6 + 4*A*a^3*b^7)*d^3*e^2 + 84*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e^3 +
42*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d*e^4 + 6*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e^5)*
x^10 + 5/3*((8*B*a^3*b^7 + 3*A*a^2*b^8)*d^5 + 10*(7*B*a^4*b^6 + 4*A*a^3*b^
7)*d^4*e + 28*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^3*e^2 + 28*(5*B*a^6*b^4 + 6*A*
a^5*b^5)*d^2*e^3 + 10*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d*e^4 + (3*B*a^8*b^2 + 8
*A*a^7*b^3)*e^5)*x^9 + 5/8*(6*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^5 + 42*(6*B...

```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2076 vs. $2(243) = 486$.

Time = 0.14 (sec) , antiderivative size = 2076, normalized size of antiderivative = 8.54

$$\int (a + bx)^{10}(A + Bx)(d + ex)^5 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)**10*(B*x+A)*(e*x+d)**5,x)
```

output

```

A*a**10*d**5*x + B*b**10*e**5*x**17/17 + x**16*(A*b**10*e**5/16 + 5*B*a*b*
*9*e**5/8 + 5*B*b**10*d*e**4/16) + x**15*(2*A*a*b**9*e**5/3 + A*b**10*d*e
*4/3 + 3*B*a**2*b**8*e**5 + 10*B*a*b**9*d*e**4/3 + 2*B*b**10*d**2*e**3/3)
+ x**14*(45*A*a**2*b**8*e**5/14 + 25*A*a*b**9*d*e**4/7 + 5*A*b**10*d**2*e
*3/7 + 60*B*a**3*b**7*e**5/7 + 225*B*a**2*b**8*d*e**4/14 + 50*B*a*b**9*d**
2*e**3/7 + 5*B*b**10*d**3*e**2/7) + x**13*(120*A*a**3*b**7*e**5/13 + 225*A
*a**2*b**8*d*e**4/13 + 100*A*a*b**9*d**2*e**3/13 + 10*A*b**10*d**3*e**2/13
+ 210*B*a**4*b**6*e**5/13 + 600*B*a**3*b**7*d*e**4/13 + 450*B*a**2*b**8*d
**2*e**3/13 + 100*B*a*b**9*d**3*e**2/13 + 5*B*b**10*d**4*e/13) + x**12*(35
*A*a**4*b**6*e**5/2 + 50*A*a**3*b**7*d*e**4 + 75*A*a**2*b**8*d**2*e**3/2 +
25*A*a*b**9*d**3*e**2/3 + 5*A*b**10*d**4*e/12 + 21*B*a**5*b**5*e**5 + 175
*B*a**4*b**6*d*e**4/2 + 100*B*a**3*b**7*d**2*e**3 + 75*B*a**2*b**8*d**3*e
*2/2 + 25*B*a*b**9*d**4*e/6 + B*b**10*d**5/12) + x**11*(252*A*a**5*b**5*e
*5/11 + 1050*A*a**4*b**6*d*e**4/11 + 1200*A*a**3*b**7*d**2*e**3/11 + 450*A
*a**2*b**8*d**3*e**2/11 + 50*A*a*b**9*d**4*e/11 + A*b**10*d**5/11 + 210*B*
a**6*b**4*e**5/11 + 1260*B*a**5*b**5*d*e**4/11 + 2100*B*a**4*b**6*d**2*e**
3/11 + 1200*B*a**3*b**7*d**3*e**2/11 + 225*B*a**2*b**8*d**4*e/11 + 10*B*a*
b**9*d**5/11) + x**10*(21*A*a**6*b**4*e**5 + 126*A*a**5*b**5*d*e**4 + 210*
A*a**4*b**6*d**2*e**3 + 120*A*a**3*b**7*d**3*e**2 + 45*A*a**2*b**8*d**4*e/
2 + A*a*b**9*d**5 + 12*B*a**7*b**3*e**5 + 105*B*a**6*b**4*d*e**4 + 252*...

```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1625 vs. $2(229) = 458$.

Time = 0.05 (sec) , antiderivative size = 1625, normalized size of antiderivative = 6.69

$$\int (a + bx)^{10}(A + Bx)(d + ex)^5 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^5,x, algorithm="maxima")
```

output

```

1/17*B*b^10*e^5*x^17 + A*a^10*d^5*x + 1/16*(5*B*b^10*d*e^4 + (10*B*a*b^9 +
A*b^10)*e^5)*x^16 + 1/3*(2*B*b^10*d^2*e^3 + (10*B*a*b^9 + A*b^10)*d*e^4 +
(9*B*a^2*b^8 + 2*A*a*b^9)*e^5)*x^15 + 5/14*(2*B*b^10*d^3*e^2 + 2*(10*B*a*
b^9 + A*b^10)*d^2*e^3 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^4 + 3*(8*B*a^3*b^7
+ 3*A*a^2*b^8)*e^5)*x^14 + 5/13*(B*b^10*d^4*e + 2*(10*B*a*b^9 + A*b^10)*d
^3*e^2 + 10*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^3 + 15*(8*B*a^3*b^7 + 3*A*a^2*
b^8)*d*e^4 + 6*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^5)*x^13 + 1/12*(B*b^10*d^5 +
5*(10*B*a*b^9 + A*b^10)*d^4*e + 50*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^2 + 150
*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^3 + 150*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e
^4 + 42*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^5)*x^12 + 1/11*((10*B*a*b^9 + A*b^10
)*d^5 + 25*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e + 150*(8*B*a^3*b^7 + 3*A*a^2*b^
8)*d^3*e^2 + 300*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^3 + 210*(6*B*a^5*b^5 +
5*A*a^4*b^6)*d*e^4 + 42*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^5)*x^11 + 1/2*((9*B*
a^2*b^8 + 2*A*a*b^9)*d^5 + 15*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^4*e + 60*(7*B*
a^4*b^6 + 4*A*a^3*b^7)*d^3*e^2 + 84*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e^3 +
42*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d*e^4 + 6*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e^5)*
x^10 + 5/3*((8*B*a^3*b^7 + 3*A*a^2*b^8)*d^5 + 10*(7*B*a^4*b^6 + 4*A*a^3*b^
7)*d^4*e + 28*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^3*e^2 + 28*(5*B*a^6*b^4 + 6*A*
a^5*b^5)*d^2*e^3 + 10*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d*e^4 + (3*B*a^8*b^2 + 8
*A*a^7*b^3)*e^5)*x^9 + 5/8*(6*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^5 + 42*(6*B...

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2032 vs. $2(229) = 458$.

Time = 0.12 (sec) , antiderivative size = 2032, normalized size of antiderivative = 8.36

$$\int (a + bx)^{10}(A + Bx)(d + ex)^5 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^5,x, algorithm="giac")
```

output

```

1/17*B*b^10*e^5*x^17 + 5/16*B*b^10*d*e^4*x^16 + 5/8*B*a*b^9*e^5*x^16 + 1/1
6*A*b^10*e^5*x^16 + 2/3*B*b^10*d^2*e^3*x^15 + 10/3*B*a*b^9*d*e^4*x^15 + 1/
3*A*b^10*d*e^4*x^15 + 3*B*a^2*b^8*e^5*x^15 + 2/3*A*a*b^9*e^5*x^15 + 5/7*B*
b^10*d^3*e^2*x^14 + 50/7*B*a*b^9*d^2*e^3*x^14 + 5/7*A*b^10*d^2*e^3*x^14 +
225/14*B*a^2*b^8*d*e^4*x^14 + 25/7*A*a*b^9*d*e^4*x^14 + 60/7*B*a^3*b^7*e^5
*x^14 + 45/14*A*a^2*b^8*e^5*x^14 + 5/13*B*b^10*d^4*e*x^13 + 100/13*B*a*b^9
*d^3*e^2*x^13 + 10/13*A*b^10*d^3*e^2*x^13 + 450/13*B*a^2*b^8*d^2*e^3*x^13
+ 100/13*A*a*b^9*d^2*e^3*x^13 + 600/13*B*a^3*b^7*d*e^4*x^13 + 225/13*A*a^2
*b^8*d*e^4*x^13 + 210/13*B*a^4*b^6*e^5*x^13 + 120/13*A*a^3*b^7*e^5*x^13 +
1/12*B*b^10*d^5*x^12 + 25/6*B*a*b^9*d^4*e*x^12 + 5/12*A*b^10*d^4*e*x^12 +
75/2*B*a^2*b^8*d^3*e^2*x^12 + 25/3*A*a*b^9*d^3*e^2*x^12 + 100*B*a^3*b^7*d^
2*e^3*x^12 + 75/2*A*a^2*b^8*d^2*e^3*x^12 + 175/2*B*a^4*b^6*d*e^4*x^12 + 50
*A*a^3*b^7*d*e^4*x^12 + 21*B*a^5*b^5*e^5*x^12 + 35/2*A*a^4*b^6*e^5*x^12 +
10/11*B*a*b^9*d^5*x^11 + 1/11*A*b^10*d^5*x^11 + 225/11*B*a^2*b^8*d^4*e*x^1
1 + 50/11*A*a*b^9*d^4*e*x^11 + 1200/11*B*a^3*b^7*d^3*e^2*x^11 + 450/11*A*a
^2*b^8*d^3*e^2*x^11 + 2100/11*B*a^4*b^6*d^2*e^3*x^11 + 1200/11*A*a^3*b^7*d
^2*e^3*x^11 + 1260/11*B*a^5*b^5*d*e^4*x^11 + 1050/11*A*a^4*b^6*d*e^4*x^11
+ 210/11*B*a^6*b^4*e^5*x^11 + 252/11*A*a^5*b^5*e^5*x^11 + 9/2*B*a^2*b^8*d^
5*x^10 + A*a*b^9*d^5*x^10 + 60*B*a^3*b^7*d^4*e*x^10 + 45/2*A*a^2*b^8*d^4*
e*x^10 + 210*B*a^4*b^6*d^3*e^2*x^10 + 120*A*a^3*b^7*d^3*e^2*x^10 + 252*B...

```

Mupad [B] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 1685, normalized size of antiderivative = 6.93

$$\int (a + bx)^{10}(A + Bx)(d + ex)^5 dx = \text{Too large to display}$$

input

```
int((A + B*x)*(a + b*x)^10*(d + e*x)^5,x)
```

output

```

x^9*(5*A*a^2*b^8*d^5 + (40*A*a^7*b^3*e^5)/3 + (40*B*a^3*b^7*d^5)/3 + 5*B*a
^8*b^2*e^5 + (200*A*a^3*b^7*d^4*e)/3 + (350*A*a^6*b^4*d*e^4)/3 + (350*B*a^
4*b^6*d^4*e)/3 + (200*B*a^7*b^3*d*e^4)/3 + (700*A*a^4*b^6*d^3*e^2)/3 + 280
*A*a^5*b^5*d^2*e^3 + 280*B*a^5*b^5*d^3*e^2 + (700*B*a^6*b^4*d^2*e^3)/3) +
x^7*((B*a^10*e^5)/7 + (10*A*a^9*b*e^5)/7 + 30*A*a^4*b^6*d^5 + 36*B*a^5*b^5
*d^5 + 180*A*a^5*b^5*d^4*e + (225*A*a^8*b^2*d*e^4)/7 + 150*B*a^6*b^4*d^4*e
+ 300*A*a^6*b^4*d^3*e^2 + (1200*A*a^7*b^3*d^2*e^3)/7 + (1200*B*a^7*b^3*d^
3*e^2)/7 + (450*B*a^8*b^2*d^2*e^3)/7 + (50*B*a^9*b*d*e^4)/7) + x^11*((A*b^
10*d^5)/11 + (10*B*a*b^9*d^5)/11 + (252*A*a^5*b^5*e^5)/11 + (210*B*a^6*b^4
*e^5)/11 + (1050*A*a^4*b^6*d*e^4)/11 + (225*B*a^2*b^8*d^4*e)/11 + (1260*B*
a^5*b^5*d*e^4)/11 + (450*A*a^2*b^8*d^3*e^2)/11 + (1200*A*a^3*b^7*d^2*e^3)/
11 + (1200*B*a^3*b^7*d^3*e^2)/11 + (2100*B*a^4*b^6*d^2*e^3)/11 + (50*A*a*b
^9*d^4*e)/11) + x^10*(A*a*b^9*d^5 + 21*A*a^6*b^4*e^5 + (9*B*a^2*b^8*d^5)/2
+ 12*B*a^7*b^3*e^5 + (45*A*a^2*b^8*d^4*e)/2 + 126*A*a^5*b^5*d*e^4 + 60*B*
a^3*b^7*d^4*e + 105*B*a^6*b^4*d*e^4 + 120*A*a^3*b^7*d^3*e^2 + 210*A*a^4*b^
6*d^2*e^3 + 210*B*a^4*b^6*d^3*e^2 + 252*B*a^5*b^5*d^2*e^3) + x^8*((5*B*a^9
*b*e^5)/4 + 15*A*a^3*b^7*d^5 + (45*A*a^8*b^2*e^5)/8 + (105*B*a^4*b^6*d^5)/
4 + (525*A*a^4*b^6*d^4*e)/4 + 75*A*a^7*b^3*d*e^4 + (315*B*a^5*b^5*d^4*e)/2
+ (225*B*a^8*b^2*d*e^4)/8 + 315*A*a^5*b^5*d^3*e^2 + (525*A*a^6*b^4*d^2*e^
3)/2 + (525*B*a^6*b^4*d^3*e^2)/2 + 150*B*a^7*b^3*d^2*e^3) + x^5*(A*a^10...

```

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 1041, normalized size of antiderivative = 4.28

$$\int (a + bx)^{10}(A + Bx)(d + ex)^5 dx = \text{Too large to display}$$

input

```
int((b*x+a)^10*(B*x+A)*(e*x+d)^5,x)
```

output

```
(x*(74256*a**11*d**5 + 185640*a**11*d**4*e*x + 247520*a**11*d**3*e**2*x**2
+ 185640*a**11*d**2*e**3*x**3 + 74256*a**11*d*e**4*x**4 + 12376*a**11*e**
5*x**5 + 408408*a**10*b*d**5*x + 1361360*a**10*b*d**4*e*x**2 + 2042040*a**
10*b*d**3*e**2*x**3 + 1633632*a**10*b*d**2*e**3*x**4 + 680680*a**10*b*d*e*
**4*x**5 + 116688*a**10*b*e**5*x**6 + 1361360*a**9*b**2*d**5*x**2 + 5105100
*a**9*b**2*d**4*e*x**3 + 8168160*a**9*b**2*d**3*e**2*x**4 + 6806800*a**9*b
**2*d**2*e**3*x**5 + 2917200*a**9*b**2*d*e**4*x**6 + 510510*a**9*b**2*e**5
*x**7 + 3063060*a**8*b**3*d**5*x**3 + 12252240*a**8*b**3*d**4*e*x**4 + 204
20400*a**8*b**3*d**3*e**2*x**5 + 17503200*a**8*b**3*d**2*e**3*x**6 + 76576
50*a**8*b**3*d*e**4*x**7 + 1361360*a**8*b**3*e**5*x**8 + 4900896*a**7*b**4
*d**5*x**4 + 20420400*a**7*b**4*d**4*e*x**5 + 35006400*a**7*b**4*d**3*e**2
*x**6 + 30630600*a**7*b**4*d**2*e**3*x**7 + 13613600*a**7*b**4*d*e**4*x**8
+ 2450448*a**7*b**4*e**5*x**9 + 5717712*a**6*b**5*d**5*x**5 + 24504480*a*
**6*b**5*d**4*e*x**6 + 42882840*a**6*b**5*d**3*e**2*x**7 + 38118080*a**6*b*
**5*d**2*e**3*x**8 + 17153136*a**6*b**5*d*e**4*x**9 + 3118752*a**6*b**5*e**
5*x**10 + 4900896*a**5*b**6*d**5*x**6 + 21441420*a**5*b**6*d**4*e*x**7 + 3
8118080*a**5*b**6*d**3*e**2*x**8 + 34306272*a**5*b**6*d**2*e**3*x**9 + 155
93760*a**5*b**6*d*e**4*x**10 + 2858856*a**5*b**6*e**5*x**11 + 3063060*a**4
*b**7*d**5*x**7 + 13613600*a**4*b**7*d**4*e*x**8 + 24504480*a**4*b**7*d**3
*e**2*x**9 + 22276800*a**4*b**7*d**2*e**3*x**10 + 10210200*a**4*b**7*d*...
```


3.74 $\int (a + bx)^{10}(A + Bx)(d + ex)^4 dx$

Optimal result	756
Mathematica [B] (verified)	757
Rubi [A] (verified)	758
Maple [B] (verified)	759
Fricas [B] (verification not implemented)	760
Sympy [B] (verification not implemented)	761
Maxima [B] (verification not implemented)	762
Giac [B] (verification not implemented)	763
Mupad [B] (verification not implemented)	764
Reduce [B] (verification not implemented)	765

Optimal result

Integrand size = 20, antiderivative size = 204

$$\int (a + bx)^{10}(A + Bx)(d + ex)^4 dx = \frac{(Ab - aB)(bd - ae)^4(a + bx)^{11}}{11b^6} + \frac{(bd - ae)^3(bBd + 4Abe - 5aBe)(a + bx)^{12}}{12b^6} + \frac{2e(bd - ae)^2(2bBd + 3Abe - 5aBe)(a + bx)^{13}}{13b^6} + \frac{e^2(bd - ae)(3bBd + 2Abe - 5aBe)(a + bx)^{14}}{7b^6} + \frac{e^3(4bBd + Abe - 5aBe)(a + bx)^{15}}{15b^6} + \frac{Be^4(a + bx)^{16}}{16b^6}$$

output

```
1/11*(A*b-B*a)*(-a*e+b*d)^4*(b*x+a)^11/b^6+1/12*(-a*e+b*d)^3*(4*A*b*e-5*B*
a*e+B*b*d)*(b*x+a)^12/b^6+2/13*e*(-a*e+b*d)^2*(3*A*b*e-5*B*a*e+2*B*b*d)*(b
*x+a)^13/b^6+1/7*e^2*(-a*e+b*d)*(2*A*b*e-5*B*a*e+3*B*b*d)*(b*x+a)^14/b^6+1
/15*e^3*(A*b*e-5*B*a*e+4*B*b*d)*(b*x+a)^15/b^6+1/16*B*e^4*(b*x+a)^16/b^6
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1098 vs. $2(204) = 408$.

Time = 0.45 (sec) , antiderivative size = 1098, normalized size of antiderivative = 5.38

$$\int (a + bx)^{10}(A + Bx)(d + ex)^4 dx = \text{Too large to display}$$

input `Integrate[(a + b*x)^10*(A + B*x)*(d + e*x)^4,x]`

output

```
(x*(8008*a^10*(6*A*(5*d^4 + 10*d^3*e*x + 10*d^2*e^2*x^2 + 5*d*e^3*x^3 + e^4*x^4) + B*x*(15*d^4 + 40*d^3*e*x + 45*d^2*e^2*x^2 + 24*d*e^3*x^3 + 5*e^4*x^4)) + 11440*a^9*b*x*(7*A*(15*d^4 + 40*d^3*e*x + 45*d^2*e^2*x^2 + 24*d*e^3*x^3 + 5*e^4*x^4) + 2*B*x*(35*d^4 + 105*d^3*e*x + 126*d^2*e^2*x^2 + 70*d*e^3*x^3 + 15*e^4*x^4)) + 12870*a^8*b^2*x^2*(8*A*(35*d^4 + 105*d^3*e*x + 126*d^2*e^2*x^2 + 70*d*e^3*x^3 + 15*e^4*x^4) + 3*B*x*(70*d^4 + 224*d^3*e*x + 280*d^2*e^2*x^2 + 160*d*e^3*x^3 + 35*e^4*x^4)) + 11440*a^7*b^3*x^3*(9*A*(70*d^4 + 224*d^3*e*x + 280*d^2*e^2*x^2 + 160*d*e^3*x^3 + 35*e^4*x^4) + 4*B*x*(126*d^4 + 420*d^3*e*x + 540*d^2*e^2*x^2 + 315*d*e^3*x^3 + 70*e^4*x^4)) + 40040*a^6*b^4*x^4*(2*A*(126*d^4 + 420*d^3*e*x + 540*d^2*e^2*x^2 + 315*d*e^3*x^3 + 70*e^4*x^4) + B*x*(210*d^4 + 720*d^3*e*x + 945*d^2*e^2*x^2 + 560*d*e^3*x^3 + 126*e^4*x^4)) + 4368*a^5*b^5*x^5*(11*A*(210*d^4 + 720*d^3*e*x + 945*d^2*e^2*x^2 + 560*d*e^3*x^3 + 126*e^4*x^4) + 6*B*x*(330*d^4 + 1155*d^3*e*x + 1540*d^2*e^2*x^2 + 924*d*e^3*x^3 + 210*e^4*x^4)) + 1820*a^4*b^6*x^6*(12*A*(330*d^4 + 1155*d^3*e*x + 1540*d^2*e^2*x^2 + 924*d*e^3*x^3 + 210*e^4*x^4) + 7*B*x*(495*d^4 + 1760*d^3*e*x + 2376*d^2*e^2*x^2 + 1440*d*e^3*x^3 + 330*e^4*x^4)) + 560*a^3*b^7*x^7*(13*A*(495*d^4 + 1760*d^3*e*x + 2376*d^2*e^2*x^2 + 1440*d*e^3*x^3 + 330*e^4*x^4) + 8*B*x*(715*d^4 + 2574*d^3*e*x + 3510*d^2*e^2*x^2 + 2145*d*e^3*x^3 + 495*e^4*x^4)) + 120*a^2*b^8*x^8*(14*A*(715*d^4 + 2574*d^3*e*x + 3510*d^2*e^2*x^2 + 2145*d*e^3*x^3 + 495...
```

Rubi [A] (verified)

Time = 1.18 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^{10}(A + Bx)(d + ex)^4 dx$$

↓ 86

$$\int \left(\frac{e^3(a + bx)^{14}(-5aBe + Abe + 4bBd)}{b^5} + \frac{2e^2(a + bx)^{13}(bd - ae)(-5aBe + 2Abe + 3bBd)}{b^5} + \frac{2e(a + bx)^{12}(bd - ae)^2(-5aBe + 3Abe + 2bBd)}{b^5} \right) dx$$

↓ 2009

$$\frac{e^3(a + bx)^{15}(-5aBe + Abe + 4bBd)}{15b^6} + \frac{e^2(a + bx)^{14}(bd - ae)(-5aBe + 2Abe + 3bBd)}{7b^6} + \frac{2e(a + bx)^{13}(bd - ae)^2(-5aBe + 3Abe + 2bBd)}{13b^6} + \frac{(a + bx)^{12}(bd - ae)^3(-5aBe + 4Abe + bBd)}{12b^6} + \frac{(a + bx)^{11}(Ab - aB)(bd - ae)^4}{11b^6} + \frac{Be^4(a + bx)^{16}}{16b^6}$$

input

```
Int[(a + b*x)^10*(A + B*x)*(d + e*x)^4,x]
```

output

```
((A*b - a*B)*(b*d - a*e)^4*(a + b*x)^11)/(11*b^6) + ((b*d - a*e)^3*(b*B*d + 4*A*b*e - 5*a*B*e)*(a + b*x)^12)/(12*b^6) + (2*e*(b*d - a*e)^2*(2*b*B*d + 3*A*b*e - 5*a*B*e)*(a + b*x)^13)/(13*b^6) + (e^2*(b*d - a*e)*(3*b*B*d + 2*A*b*e - 5*a*B*e)*(a + b*x)^14)/(7*b^6) + (e^3*(4*b*B*d + A*b*e - 5*a*B*e)*(a + b*x)^15)/(15*b^6) + (B*e^4*(a + b*x)^16)/(16*b^6)
```

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0]) || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1336 vs. $2(192) = 384$.

Time = 0.22 (sec) , antiderivative size = 1337, normalized size of antiderivative = 6.55

method	result	size
default	Expression too large to display	1337
norman	Expression too large to display	1403
orering	Expression too large to display	1656
gospers	Expression too large to display	1657
risch	Expression too large to display	1657
parallelrisc	Expression too large to display	1657

input

```
int((b*x+a)^10*(B*x+A)*(e*x+d)^4,x,method=_RETURNVERBOSE)
```

output

```

1/16*b^10*B*e^4*x^16+1/15*((A*b^10+10*B*a*b^9)*e^4+4*b^10*B*d*e^3)*x^15+1/
14*((10*A*a*b^9+45*B*a^2*b^8)*e^4+4*(A*b^10+10*B*a*b^9)*d*e^3+6*b^10*B*d^2
*e^2)*x^14+1/13*((45*A*a^2*b^8+120*B*a^3*b^7)*e^4+4*(10*A*a*b^9+45*B*a^2*b
^8)*d*e^3+6*(A*b^10+10*B*a*b^9)*d^2*e^2+4*b^10*B*d^3*e)*x^13+1/12*((120*A*
a^3*b^7+210*B*a^4*b^6)*e^4+4*(45*A*a^2*b^8+120*B*a^3*b^7)*d*e^3+6*(10*A*a*
b^9+45*B*a^2*b^8)*d^2*e^2+4*(A*b^10+10*B*a*b^9)*d^3*e+b^10*B*d^4)*x^12+1/1
1*((210*A*a^4*b^6+252*B*a^5*b^5)*e^4+4*(120*A*a^3*b^7+210*B*a^4*b^6)*d*e^3
+6*(45*A*a^2*b^8+120*B*a^3*b^7)*d^2*e^2+4*(10*A*a*b^9+45*B*a^2*b^8)*d^3*e+
(A*b^10+10*B*a*b^9)*d^4)*x^11+1/10*((252*A*a^5*b^5+210*B*a^6*b^4)*e^4+4*(2
10*A*a^4*b^6+252*B*a^5*b^5)*d*e^3+6*(120*A*a^3*b^7+210*B*a^4*b^6)*d^2*e^2+
4*(45*A*a^2*b^8+120*B*a^3*b^7)*d^3*e+(10*A*a*b^9+45*B*a^2*b^8)*d^4)*x^10+1
/9*((210*A*a^6*b^4+120*B*a^7*b^3)*e^4+4*(252*A*a^5*b^5+210*B*a^6*b^4)*d*e^
3+6*(210*A*a^4*b^6+252*B*a^5*b^5)*d^2*e^2+4*(120*A*a^3*b^7+210*B*a^4*b^6)*
d^3*e+(45*A*a^2*b^8+120*B*a^3*b^7)*d^4)*x^9+1/8*((120*A*a^7*b^3+45*B*a^8*b
^2)*e^4+4*(210*A*a^6*b^4+120*B*a^7*b^3)*d*e^3+6*(252*A*a^5*b^5+210*B*a^6*b
^4)*d^2*e^2+4*(210*A*a^4*b^6+252*B*a^5*b^5)*d^3*e+(120*A*a^3*b^7+210*B*a^4
*b^6)*d^4)*x^8+1/7*((45*A*a^8*b^2+10*B*a^9*b)*e^4+4*(120*A*a^7*b^3+45*B*a^
8*b^2)*d*e^3+6*(210*A*a^6*b^4+120*B*a^7*b^3)*d^2*e^2+4*(252*A*a^5*b^5+210*
B*a^6*b^4)*d^3*e+(210*A*a^4*b^6+252*B*a^5*b^5)*d^4)*x^7+1/6*((10*A*a^9*b+B
*a^10)*e^4+4*(45*A*a^8*b^2+10*B*a^9*b)*d*e^3+6*(120*A*a^7*b^3+45*B*a^8*...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1352 vs. $2(192) = 384$.

Time = 0.08 (sec) , antiderivative size = 1352, normalized size of antiderivative = 6.63

$$\int (a + bx)^{10}(A + Bx)(d + ex)^4 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^4,x, algorithm="fricas")
```

output

```

1/16*B*b^10*e^4*x^16 + A*a^10*d^4*x + 1/15*(4*B*b^10*d*e^3 + (10*B*a*b^9 +
A*b^10)*e^4)*x^15 + 1/14*(6*B*b^10*d^2*e^2 + 4*(10*B*a*b^9 + A*b^10)*d*e^
3 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*e^4)*x^14 + 1/13*(4*B*b^10*d^3*e + 6*(10*B
*a*b^9 + A*b^10)*d^2*e^2 + 20*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^3 + 15*(8*B*a^
3*b^7 + 3*A*a^2*b^8)*e^4)*x^13 + 1/12*(B*b^10*d^4 + 4*(10*B*a*b^9 + A*b^10
)*d^3*e + 30*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^2 + 60*(8*B*a^3*b^7 + 3*A*a^2
*b^8)*d*e^3 + 30*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^4)*x^12 + 1/11*((10*B*a*b^9
+ A*b^10)*d^4 + 20*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e + 90*(8*B*a^3*b^7 + 3*
A*a^2*b^8)*d^2*e^2 + 120*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^3 + 42*(6*B*a^5*b
^5 + 5*A*a^4*b^6)*e^4)*x^11 + 1/10*(5*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4 + 60*(
8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e + 180*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^2
+ 168*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^3 + 42*(5*B*a^6*b^4 + 6*A*a^5*b^5)*
e^4)*x^10 + 1/3*(5*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^4 + 40*(7*B*a^4*b^6 + 4*A
*a^3*b^7)*d^3*e + 84*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e^2 + 56*(5*B*a^6*b^4
+ 6*A*a^5*b^5)*d*e^3 + 10*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e^4)*x^9 + 3/8*(10*
(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^4 + 56*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^3*e + 8
4*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^2*e^2 + 40*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d*e
^3 + 5*(3*B*a^8*b^2 + 8*A*a^7*b^3)*e^4)*x^8 + 1/7*(42*(6*B*a^5*b^5 + 5*A*a
^4*b^6)*d^4 + 168*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^3*e + 180*(4*B*a^7*b^3 + 7
*A*a^6*b^4)*d^2*e^2 + 60*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d*e^3 + 5*(2*B*a^9...

```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1676 vs. $2(204) = 408$.

Time = 0.18 (sec) , antiderivative size = 1676, normalized size of antiderivative = 8.22

$$\int (a + bx)^{10}(A + Bx)(d + ex)^4 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)**10*(B*x+A)*(e*x+d)**4,x)
```

output

```
A**10*d**4*x + B*b**10*e**4*x**16/16 + x**15*(A*b**10*e**4/15 + 2*B*a*b*
*9*e**4/3 + 4*B*b**10*d*e**3/15) + x**14*(5*A*a*b**9*e**4/7 + 2*A*b**10*d*
e**3/7 + 45*B*a**2*b**8*e**4/14 + 20*B*a*b**9*d*e**3/7 + 3*B*b**10*d**2*e*
*2/7) + x**13*(45*A*a**2*b**8*e**4/13 + 40*A*a*b**9*d*e**3/13 + 6*A*b**10*
d**2*e**2/13 + 120*B*a**3*b**7*e**4/13 + 180*B*a**2*b**8*d*e**3/13 + 60*B*
a*b**9*d**2*e**2/13 + 4*B*b**10*d**3*e/13) + x**12*(10*A*a**3*b**7*e**4 +
15*A*a**2*b**8*d*e**3 + 5*A*a*b**9*d**2*e**2 + A*b**10*d**3*e/3 + 35*B*a**
4*b**6*e**4/2 + 40*B*a**3*b**7*d*e**3 + 45*B*a**2*b**8*d**2*e**2/2 + 10*B*
a*b**9*d**3*e/3 + B*b**10*d**4/12) + x**11*(210*A*a**4*b**6*e**4/11 + 480*
A*a**3*b**7*d*e**3/11 + 270*A*a**2*b**8*d**2*e**2/11 + 40*A*a*b**9*d**3*e/
11 + A*b**10*d**4/11 + 252*B*a**5*b**5*e**4/11 + 840*B*a**4*b**6*d*e**3/11
+ 720*B*a**3*b**7*d**2*e**2/11 + 180*B*a**2*b**8*d**3*e/11 + 10*B*a*b**9*
d**4/11) + x**10*(126*A*a**5*b**5*e**4/5 + 84*A*a**4*b**6*d*e**3 + 72*A*a*
*3*b**7*d**2*e**2 + 18*A*a**2*b**8*d**3*e + A*a*b**9*d**4 + 21*B*a**6*b**4
*e**4 + 504*B*a**5*b**5*d*e**3/5 + 126*B*a**4*b**6*d**2*e**2 + 48*B*a**3*b
**7*d**3*e + 9*B*a**2*b**8*d**4/2) + x**9*(70*A*a**6*b**4*e**4/3 + 112*A*a
**5*b**5*d*e**3 + 140*A*a**4*b**6*d**2*e**2 + 160*A*a**3*b**7*d**3*e/3 + 5
*A*a**2*b**8*d**4 + 40*B*a**7*b**3*e**4/3 + 280*B*a**6*b**4*d*e**3/3 + 168
*B*a**5*b**5*d**2*e**2 + 280*B*a**4*b**6*d**3*e/3 + 40*B*a**3*b**7*d**4/3)
+ x**8*(15*A*a**7*b**3*e**4 + 105*A*a**6*b**4*d*e**3 + 189*A*a**5*b**5...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1352 vs. $2(192) = 384$.

Time = 0.04 (sec) , antiderivative size = 1352, normalized size of antiderivative = 6.63

$$\int (a + bx)^{10}(A + Bx)(d + ex)^4 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^4,x, algorithm="maxima")
```

output

```

1/16*B*b^10*e^4*x^16 + A*a^10*d^4*x + 1/15*(4*B*b^10*d*e^3 + (10*B*a*b^9 +
A*b^10)*e^4)*x^15 + 1/14*(6*B*b^10*d^2*e^2 + 4*(10*B*a*b^9 + A*b^10)*d*e^
3 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*e^4)*x^14 + 1/13*(4*B*b^10*d^3*e + 6*(10*B
*a*b^9 + A*b^10)*d^2*e^2 + 20*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^3 + 15*(8*B*a^
3*b^7 + 3*A*a^2*b^8)*e^4)*x^13 + 1/12*(B*b^10*d^4 + 4*(10*B*a*b^9 + A*b^10
)*d^3*e + 30*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^2 + 60*(8*B*a^3*b^7 + 3*A*a^2
*b^8)*d*e^3 + 30*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^4)*x^12 + 1/11*((10*B*a*b^9
+ A*b^10)*d^4 + 20*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e + 90*(8*B*a^3*b^7 + 3*
A*a^2*b^8)*d^2*e^2 + 120*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^3 + 42*(6*B*a^5*b
^5 + 5*A*a^4*b^6)*e^4)*x^11 + 1/10*(5*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4 + 60*(
8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e + 180*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^2
+ 168*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^3 + 42*(5*B*a^6*b^4 + 6*A*a^5*b^5)*
e^4)*x^10 + 1/3*(5*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^4 + 40*(7*B*a^4*b^6 + 4*A
*a^3*b^7)*d^3*e + 84*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e^2 + 56*(5*B*a^6*b^4
+ 6*A*a^5*b^5)*d*e^3 + 10*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e^4)*x^9 + 3/8*(10*
(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^4 + 56*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^3*e + 8
4*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^2*e^2 + 40*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d*e
^3 + 5*(3*B*a^8*b^2 + 8*A*a^7*b^3)*e^4)*x^8 + 1/7*(42*(6*B*a^5*b^5 + 5*A*a
^4*b^6)*d^4 + 168*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^3*e + 180*(4*B*a^7*b^3 + 7
*A*a^6*b^4)*d^2*e^2 + 60*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d*e^3 + 5*(2*B*a^9...

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1656 vs. $2(192) = 384$.

Time = 0.13 (sec) , antiderivative size = 1656, normalized size of antiderivative = 8.12

$$\int (a + bx)^{10}(A + Bx)(d + ex)^4 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^4,x, algorithm="giac")
```


output

```

1/16*B*b^10*e^4*x^16 + 4/15*B*b^10*d*e^3*x^15 + 2/3*B*a*b^9*e^4*x^15 + 1/1
5*A*b^10*e^4*x^15 + 3/7*B*b^10*d^2*e^2*x^14 + 20/7*B*a*b^9*d*e^3*x^14 + 2/
7*A*b^10*d*e^3*x^14 + 45/14*B*a^2*b^8*e^4*x^14 + 5/7*A*a*b^9*e^4*x^14 + 4/
13*B*b^10*d^3*e*x^13 + 60/13*B*a*b^9*d^2*e^2*x^13 + 6/13*A*b^10*d^2*e^2*x^
13 + 180/13*B*a^2*b^8*d*e^3*x^13 + 40/13*A*a*b^9*d*e^3*x^13 + 120/13*B*a^3
*b^7*e^4*x^13 + 45/13*A*a^2*b^8*e^4*x^13 + 1/12*B*b^10*d^4*x^12 + 10/3*B*a
*b^9*d^3*e*x^12 + 1/3*A*b^10*d^3*e*x^12 + 45/2*B*a^2*b^8*d^2*e^2*x^12 + 5*
A*a*b^9*d^2*e^2*x^12 + 40*B*a^3*b^7*d*e^3*x^12 + 15*A*a^2*b^8*d*e^3*x^12 +
35/2*B*a^4*b^6*e^4*x^12 + 10*A*a^3*b^7*e^4*x^12 + 10/11*B*a*b^9*d^4*x^11
+ 1/11*A*b^10*d^4*x^11 + 180/11*B*a^2*b^8*d^3*e*x^11 + 40/11*A*a*b^9*d^3*e
*x^11 + 720/11*B*a^3*b^7*d^2*e^2*x^11 + 270/11*A*a^2*b^8*d^2*e^2*x^11 + 84
0/11*B*a^4*b^6*d*e^3*x^11 + 480/11*A*a^3*b^7*d*e^3*x^11 + 252/11*B*a^5*b^5
*e^4*x^11 + 210/11*A*a^4*b^6*e^4*x^11 + 9/2*B*a^2*b^8*d^4*x^10 + A*a*b^9*d
^4*x^10 + 48*B*a^3*b^7*d^3*e*x^10 + 18*A*a^2*b^8*d^3*e*x^10 + 126*B*a^4*b^
6*d^2*e^2*x^10 + 72*A*a^3*b^7*d^2*e^2*x^10 + 504/5*B*a^5*b^5*d*e^3*x^10 +
84*A*a^4*b^6*d*e^3*x^10 + 21*B*a^6*b^4*e^4*x^10 + 126/5*A*a^5*b^5*e^4*x^10
+ 40/3*B*a^3*b^7*d^4*x^9 + 5*A*a^2*b^8*d^4*x^9 + 280/3*B*a^4*b^6*d^3*e*x^
9 + 160/3*A*a^3*b^7*d^3*e*x^9 + 168*B*a^5*b^5*d^2*e^2*x^9 + 140*A*a^4*b^6*
d^2*e^2*x^9 + 280/3*B*a^6*b^4*d*e^3*x^9 + 112*A*a^5*b^5*d*e^3*x^9 + 40/3*B
*a^7*b^3*e^4*x^9 + 70/3*A*a^6*b^4*e^4*x^9 + 105/4*B*a^4*b^6*d^4*x^8 + 1...

```

Mupad [B] (verification not implemented)

Time = 1.23 (sec) , antiderivative size = 1386, normalized size of antiderivative = 6.79

$$\int (a + bx)^{10}(A + Bx)(d + ex)^4 dx = \text{Too large to display}$$

input

```
int((A + B*x)*(a + b*x)^10*(d + e*x)^4,x)
```

output

```

x^6*((B*a^10*e^4)/6 + (5*A*a^9*b*e^4)/3 + 42*A*a^5*b^5*d^4 + 35*B*a^6*b^4*
d^4 + 140*A*a^6*b^4*d^3*e + 30*A*a^8*b^2*d*e^3 + 80*B*a^7*b^3*d^3*e + 120*
A*a^7*b^3*d^2*e^2 + 45*B*a^8*b^2*d^2*e^2 + (20*B*a^9*b*d*e^3)/3) + x^11*((
A*b^10*d^4)/11 + (10*B*a*b^9*d^4)/11 + (210*A*a^4*b^6*e^4)/11 + (252*B*a^5
*b^5*e^4)/11 + (480*A*a^3*b^7*d*e^3)/11 + (180*B*a^2*b^8*d^3*e)/11 + (840*
B*a^4*b^6*d*e^3)/11 + (270*A*a^2*b^8*d^2*e^2)/11 + (720*B*a^3*b^7*d^2*e^2)
/11 + (40*A*a*b^9*d^3*e)/11) + x^10*(A*a*b^9*d^4 + (126*A*a^5*b^5*e^4)/5 +
(9*B*a^2*b^8*d^4)/2 + 21*B*a^6*b^4*e^4 + 18*A*a^2*b^8*d^3*e + 84*A*a^4*b^
6*d*e^3 + 48*B*a^3*b^7*d^3*e + (504*B*a^5*b^5*d*e^3)/5 + 72*A*a^3*b^7*d^2*
e^2 + 126*B*a^4*b^6*d^2*e^2) + x^7*((10*B*a^9*b*e^4)/7 + 30*A*a^4*b^6*d^4
+ (45*A*a^8*b^2*e^4)/7 + 36*B*a^5*b^5*d^4 + 144*A*a^5*b^5*d^3*e + (480*A*a
^7*b^3*d*e^3)/7 + 120*B*a^6*b^4*d^3*e + (180*B*a^8*b^2*d*e^3)/7 + 180*A*a^
6*b^4*d^2*e^2 + (720*B*a^7*b^3*d^2*e^2)/7) + x^4*(A*a^10*d*e^3 + 30*A*a^7*
b^3*d^4 + (45*B*a^8*b^2*d^4)/4 + (3*B*a^10*d^2*e^2)/2 + 45*A*a^8*b^2*d^3*e
+ 15*A*a^9*b*d^2*e^2 + 10*B*a^9*b*d^3*e) + x^13*((4*B*b^10*d^3*e)/13 + (4
5*A*a^2*b^8*e^4)/13 + (120*B*a^3*b^7*e^4)/13 + (6*A*b^10*d^2*e^2)/13 + (60
*B*a*b^9*d^2*e^2)/13 + (180*B*a^2*b^8*d*e^3)/13 + (40*A*a*b^9*d*e^3)/13) +
x^3*((10*B*a^9*b*d^4)/3 + (4*B*a^10*d^3*e)/3 + 15*A*a^8*b^2*d^4 + 2*A*a^1
0*d^2*e^2 + (40*A*a^9*b*d^3*e)/3) + x^14*((5*A*a*b^9*e^4)/7 + (2*A*b^10*d*
e^3)/7 + (45*B*a^2*b^8*e^4)/14 + (3*B*b^10*d^2*e^2)/7 + (20*B*a*b^9*d*e...

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 847, normalized size of antiderivative = 4.15

$$\int (a + bx)^{10}(A + Bx)(d + ex)^4 dx = \text{Too large to display}$$

input

```
int((b*x+a)^10*(B*x+A)*(e*x+d)^4,x)
```

output

```
(x*(21840*a**11*d**4 + 43680*a**11*d**3*e*x + 43680*a**11*d**2*e**2*x**2 +
  21840*a**11*d*e**3*x**3 + 4368*a**11*e**4*x**4 + 120120*a**10*b*d**4*x +
  320320*a**10*b*d**3*e*x**2 + 360360*a**10*b*d**2*e**2*x**3 + 192192*a**10*
  b*d*e**3*x**4 + 40040*a**10*b*e**4*x**5 + 400400*a**9*b**2*d**4*x**2 + 120
  1200*a**9*b**2*d**3*e*x**3 + 1441440*a**9*b**2*d**2*e**2*x**4 + 800800*a**
  9*b**2*d*e**3*x**5 + 171600*a**9*b**2*e**4*x**6 + 900900*a**8*b**3*d**4*x*
  *3 + 2882880*a**8*b**3*d**3*e*x**4 + 3603600*a**8*b**3*d**2*e**2*x**5 + 20
  59200*a**8*b**3*d*e**3*x**6 + 450450*a**8*b**3*e**4*x**7 + 1441440*a**7*b*
  *4*d**4*x**4 + 4804800*a**7*b**4*d**3*e*x**5 + 6177600*a**7*b**4*d**2*e**2
  *x**6 + 3603600*a**7*b**4*d*e**3*x**7 + 800800*a**7*b**4*e**4*x**8 + 16816
  80*a**6*b**5*d**4*x**5 + 5765760*a**6*b**5*d**3*e*x**6 + 7567560*a**6*b**5
  *d**2*e**2*x**7 + 4484480*a**6*b**5*d*e**3*x**8 + 1009008*a**6*b**5*e**4*x
  **9 + 1441440*a**5*b**6*d**4*x**6 + 5045040*a**5*b**6*d**3*e*x**7 + 672672
  0*a**5*b**6*d**2*e**2*x**8 + 4036032*a**5*b**6*d*e**3*x**9 + 917280*a**5*b
  **6*e**4*x**10 + 900900*a**4*b**7*d**4*x**7 + 3203200*a**4*b**7*d**3*e*x**
  8 + 4324320*a**4*b**7*d**2*e**2*x**9 + 2620800*a**4*b**7*d*e**3*x**10 + 60
  0600*a**4*b**7*e**4*x**11 + 400400*a**3*b**8*d**4*x**8 + 1441440*a**3*b**8
  *d**3*e*x**9 + 1965600*a**3*b**8*d**2*e**2*x**10 + 1201200*a**3*b**8*d*e**
  3*x**11 + 277200*a**3*b**8*e**4*x**12 + 120120*a**2*b**9*d**4*x**9 + 43680
  0*a**2*b**9*d**3*e*x**10 + 600600*a**2*b**9*d**2*e**2*x**11 + 369600*a...
```

3.75 $\int (a + bx)^{10}(A + Bx)(d + ex)^3 dx$

Optimal result	767
Mathematica [B] (verified)	768
Rubi [A] (verified)	769
Maple [B] (verified)	770
Fricas [B] (verification not implemented)	771
Sympy [B] (verification not implemented)	772
Maxima [B] (verification not implemented)	773
Giac [B] (verification not implemented)	774
Mupad [B] (verification not implemented)	775
Reduce [B] (verification not implemented)	776

Optimal result

Integrand size = 20, antiderivative size = 159

$$\int (a + bx)^{10}(A + Bx)(d + ex)^3 dx = \frac{(Ab - aB)(bd - ae)^3(a + bx)^{11}}{11b^5} + \frac{(bd - ae)^2(bBd + 3Abe - 4aBe)(a + bx)^{12}}{12b^5} + \frac{3e(bd - ae)(bBd + Abe - 2aBe)(a + bx)^{13}}{13b^5} + \frac{e^2(3bBd + Abe - 4aBe)(a + bx)^{14}}{14b^5} + \frac{Be^3(a + bx)^{15}}{15b^5}$$

output

```
1/11*(A*b-B*a)*(-a*e+b*d)^3*(b*x+a)^11/b^5+1/12*(-a*e+b*d)^2*(3*A*b*e-4*B*
a*e+B*b*d)*(b*x+a)^12/b^5+3/13*e*(-a*e+b*d)*(A*b*e-2*B*a*e+B*b*d)*(b*x+a)^
13/b^5+1/14*e^2*(A*b*e-4*B*a*e+3*B*b*d)*(b*x+a)^14/b^5+1/15*B*e^3*(b*x+a)^
15/b^5
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 855 vs. $2(159) = 318$.

Time = 0.33 (sec) , antiderivative size = 855, normalized size of antiderivative = 5.38

$$\int (a + bx)^{10}(A + Bx)(d + ex)^3 dx$$

$$= \frac{x(3003a^{10}(5A(4d^3 + 6d^2ex + 4de^2x^2 + e^3x^3) + Bx(10d^3 + 20d^2ex + 15de^2x^2 + 4e^3x^3)) + 10010a^9bx(3A(10d^3 + 20d^2ex + 15de^2x^2 + 4e^3x^3) + Bx(20d^3 + 45d^2ex + 36de^2x^2 + 10e^3x^3)) + 6435a^8b^2x^2(7A(20d^3 + 45d^2ex + 36de^2x^2 + 10e^3x^3) + 3Bx(35d^3 + 84d^2ex + 70de^2x^2 + 20e^3x^3)) + 25740a^7b^3x^3(2A(35d^3 + 84d^2ex + 70de^2x^2 + 20e^3x^3) + Bx(56d^3 + 140d^2ex + 120de^2x^2 + 35e^3x^3)) + 5005a^6b^4x^4(9A(56d^3 + 140d^2ex + 120de^2x^2 + 35e^3x^3) + 5Bx(84d^3 + 216d^2ex + 189de^2x^2 + 56e^3x^3)) + 6006a^5b^5x^5(5A(84d^3 + 216d^2ex + 189de^2x^2 + 56e^3x^3) + 3Bx(120d^3 + 315d^2ex + 280de^2x^2 + 84e^3x^3)) + 1365a^4b^6x^6(11A(120d^3 + 315d^2ex + 280de^2x^2 + 84e^3x^3) + 7Bx(165d^3 + 440d^2ex + 396de^2x^2 + 120e^3x^3)) + 1820a^3b^7x^7(3A(165d^3 + 440d^2ex + 396de^2x^2 + 120e^3x^3) + 2Bx(220d^3 + 594d^2ex + 540de^2x^2 + 165e^3x^3)) + 105a^2b^8x^8(13A(220d^3 + 594d^2ex + 540de^2x^2 + 165e^3x^3) + 9Bx(286d^3 + 780d^2ex + 715de^2x^2 + 220e^3x^3)) + 30ab^9x^9(7A(286d^3 + 780d^2ex + 715de^2x^2 + 220e^3x^3) + 5Bx(364d^3 + 1001d^2ex + 924de^2x^2 + 286e^3x^3)) + b^{10}x^{10}(15A(364d^3 + 1001d^2ex + 924de^2x^2 + 286e^3x^3) + 11Bx(455d^3 + 1260d^2ex + 1170de^2x^2 + 364e^3x^3))}{x^2}$$

input

```
Integrate[(a + b*x)^10*(A + B*x)*(d + e*x)^3,x]
```

output

```
(x*(3003*a^10*(5*A*(4*d^3 + 6*d^2*e*x + 4*d*e^2*x^2 + e^3*x^3) + B*x*(10*d^3 + 20*d^2*e*x + 15*d*e^2*x^2 + 4*e^3*x^3)) + 10010*a^9*b*x*(3*A*(10*d^3 + 20*d^2*e*x + 15*d*e^2*x^2 + 4*e^3*x^3) + B*x*(20*d^3 + 45*d^2*e*x + 36*d*e^2*x^2 + 10*e^3*x^3)) + 6435*a^8*b^2*x^2*(7*A*(20*d^3 + 45*d^2*e*x + 36*d*e^2*x^2 + 10*e^3*x^3) + 3*B*x*(35*d^3 + 84*d^2*e*x + 70*d*e^2*x^2 + 20*e^3*x^3)) + 25740*a^7*b^3*x^3*(2*A*(35*d^3 + 84*d^2*e*x + 70*d*e^2*x^2 + 20*e^3*x^3) + B*x*(56*d^3 + 140*d^2*e*x + 120*d*e^2*x^2 + 35*e^3*x^3)) + 5005*a^6*b^4*x^4*(9*A*(56*d^3 + 140*d^2*e*x + 120*d*e^2*x^2 + 35*e^3*x^3) + 5*B*x*(84*d^3 + 216*d^2*e*x + 189*d*e^2*x^2 + 56*e^3*x^3)) + 6006*a^5*b^5*x^5*(5*A*(84*d^3 + 216*d^2*e*x + 189*d*e^2*x^2 + 56*e^3*x^3) + 3*B*x*(120*d^3 + 315*d^2*e*x + 280*d*e^2*x^2 + 84*e^3*x^3)) + 1365*a^4*b^6*x^6*(11*A*(120*d^3 + 315*d^2*e*x + 280*d*e^2*x^2 + 84*e^3*x^3) + 7*B*x*(165*d^3 + 440*d^2*e*x + 396*d*e^2*x^2 + 120*e^3*x^3)) + 1820*a^3*b^7*x^7*(3*A*(165*d^3 + 440*d^2*e*x + 396*d*e^2*x^2 + 120*e^3*x^3) + 2*B*x*(220*d^3 + 594*d^2*e*x + 540*d*e^2*x^2 + 165*e^3*x^3)) + 105*a^2*b^8*x^8*(13*A*(220*d^3 + 594*d^2*e*x + 540*d*e^2*x^2 + 165*e^3*x^3) + 9*B*x*(286*d^3 + 780*d^2*e*x + 715*d*e^2*x^2 + 220*e^3*x^3)) + 30*a*b^9*x^9*(7*A*(286*d^3 + 780*d^2*e*x + 715*d*e^2*x^2 + 220*e^3*x^3) + 5*B*x*(364*d^3 + 1001*d^2*e*x + 924*d*e^2*x^2 + 286*e^3*x^3)) + b^10*x^10*(15*A*(364*d^3 + 1001*d^2*e*x + 924*d*e^2*x^2 + 286*e^3*x^3) + 11*B*x*(455*d^3 + 1260*d^2*e*x + 1170*d*e^2*x^2 + 364...
```

Rubi [A] (verified)

Time = 0.93 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^{10}(A + Bx)(d + ex)^3 dx$$

↓ 86

$$\int \left(\frac{e^2(a + bx)^{13}(-4aBe + Abe + 3bBd)}{b^4} + \frac{3e(a + bx)^{12}(bd - ae)(-2aBe + Abe + bBd)}{b^4} + \frac{(a + bx)^{11}(bd - ae)}{b^4} \right) dx$$

↓ 2009

$$\frac{e^2(a + bx)^{14}(-4aBe + Abe + 3bBd)}{14b^5} + \frac{3e(a + bx)^{13}(bd - ae)(-2aBe + Abe + bBd)}{12b^5} + \frac{(a + bx)^{12}(bd - ae)^2(-4aBe + 3Abe + bBd)}{12b^5} + \frac{(a + bx)^{11}(Ab - aB)(bd - ae)^3}{11b^5} + \frac{Be^3(a + bx)^{15}}{15b^5}$$

input `Int[(a + b*x)^10*(A + B*x)*(d + e*x)^3,x]`

output `((A*b - a*B)*(b*d - a*e)^3*(a + b*x)^11)/(11*b^5) + ((b*d - a*e)^2*(b*B*d + 3*A*b*e - 4*a*B*e)*(a + b*x)^12)/(12*b^5) + (3*e*(b*d - a*e)*(b*B*d + A*b*e - 2*a*B*e)*(a + b*x)^13)/(13*b^5) + (e^2*(3*b*B*d + A*b*e - 4*a*B*e)*(a + b*x)^14)/(14*b^5) + (B*e^3*(a + b*x)^15)/(15*b^5)`

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1052 vs. $2(149) = 298$.

Time = 0.21 (sec) , antiderivative size = 1053, normalized size of antiderivative = 6.62

method	result	size
default	Expression too large to display	1053
norman	Expression too large to display	1087
gospers	Expression too large to display	1280
risch	Expression too large to display	1280
parallelrisch	Expression too large to display	1280
orering	Expression too large to display	1280

input

```
int((b*x+a)^10*(B*x+A)*(e*x+d)^3,x,method=_RETURNVERBOSE)
```

output

```

1/15*b^10*B*e^3*x^15+1/14*((A*b^10+10*B*a*b^9)*e^3+3*b^10*B*d*e^2)*x^14+1/
13*((10*A*a*b^9+45*B*a^2*b^8)*e^3+3*(A*b^10+10*B*a*b^9)*d*e^2+3*b^10*B*d^2
*e)*x^13+1/12*((45*A*a^2*b^8+120*B*a^3*b^7)*e^3+3*(10*A*a*b^9+45*B*a^2*b^8
)*d*e^2+3*(A*b^10+10*B*a*b^9)*d^2*e+b^10*B*d^3)*x^12+1/11*((120*A*a^3*b^7+
210*B*a^4*b^6)*e^3+3*(45*A*a^2*b^8+120*B*a^3*b^7)*d*e^2+3*(10*A*a*b^9+45*B
*a^2*b^8)*d^2*e+(A*b^10+10*B*a*b^9)*d^3)*x^11+1/10*((210*A*a^4*b^6+252*B*a
^5*b^5)*e^3+3*(120*A*a^3*b^7+210*B*a^4*b^6)*d*e^2+3*(45*A*a^2*b^8+120*B*a^
3*b^7)*d^2*e+(10*A*a*b^9+45*B*a^2*b^8)*d^3)*x^10+1/9*((252*A*a^5*b^5+210*B
*a^6*b^4)*e^3+3*(210*A*a^4*b^6+252*B*a^5*b^5)*d*e^2+3*(120*A*a^3*b^7+210*B
*a^4*b^6)*d^2*e+(45*A*a^2*b^8+120*B*a^3*b^7)*d^3)*x^9+1/8*((210*A*a^6*b^4+
120*B*a^7*b^3)*e^3+3*(252*A*a^5*b^5+210*B*a^6*b^4)*d*e^2+3*(210*A*a^4*b^6+
252*B*a^5*b^5)*d^2*e+(120*A*a^3*b^7+210*B*a^4*b^6)*d^3)*x^8+1/7*((120*A*a^
7*b^3+45*B*a^8*b^2)*e^3+3*(210*A*a^6*b^4+120*B*a^7*b^3)*d*e^2+3*(252*A*a^5
*b^5+210*B*a^6*b^4)*d^2*e+(210*A*a^4*b^6+252*B*a^5*b^5)*d^3)*x^7+1/6*((45*
A*a^8*b^2+10*B*a^9*b)*e^3+3*(120*A*a^7*b^3+45*B*a^8*b^2)*d*e^2+3*(210*A*a^
6*b^4+120*B*a^7*b^3)*d^2*e+(252*A*a^5*b^5+210*B*a^6*b^4)*d^3)*x^6+1/5*((10
*A*a^9*b+B*a^10)*e^3+3*(45*A*a^8*b^2+10*B*a^9*b)*d*e^2+3*(120*A*a^7*b^3+45
*B*a^8*b^2)*d^2*e+(210*A*a^6*b^4+120*B*a^7*b^3)*d^3)*x^5+1/4*(a^10*A*e^3+3
*(10*A*a^9*b+B*a^10)*d*e^2+3*(45*A*a^8*b^2+10*B*a^9*b)*d^2*e+(120*A*a^7*b^
3+45*B*a^8*b^2)*d^3)*x^4+1/3*(3*a^10*A*d*e^2+3*(10*A*a^9*b+B*a^10)*d^2*...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1068 vs. $2(149) = 298$.

Time = 0.10 (sec) , antiderivative size = 1068, normalized size of antiderivative = 6.72

$$\int (a + bx)^{10}(A + Bx)(d + ex)^3 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^3,x, algorithm="fricas")
```


output

```

1/15*B*b^10*e^3*x^15 + A*a^10*d^3*x + 1/14*(3*B*b^10*d*e^2 + (10*B*a*b^9 +
A*b^10)*e^3)*x^14 + 1/13*(3*B*b^10*d^2*e + 3*(10*B*a*b^9 + A*b^10)*d*e^2
+ 5*(9*B*a^2*b^8 + 2*A*a*b^9)*e^3)*x^13 + 1/12*(B*b^10*d^3 + 3*(10*B*a*b^9
+ A*b^10)*d^2*e + 15*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^2 + 15*(8*B*a^3*b^7 +
3*A*a^2*b^8)*e^3)*x^12 + 1/11*((10*B*a*b^9 + A*b^10)*d^3 + 15*(9*B*a^2*b^8
+ 2*A*a*b^9)*d^2*e + 45*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^2 + 30*(7*B*a^4*b
^6 + 4*A*a^3*b^7)*e^3)*x^11 + 1/10*(5*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3 + 45*(
8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e + 90*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^2 +
42*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^3)*x^10 + 1/3*(5*(8*B*a^3*b^7 + 3*A*a^2*b
^8)*d^3 + 30*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e + 42*(6*B*a^5*b^5 + 5*A*a^4
*b^6)*d*e^2 + 14*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^3)*x^9 + 3/4*(5*(7*B*a^4*b
^6 + 4*A*a^3*b^7)*d^3 + 21*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e + 21*(5*B*a^6*
b^4 + 6*A*a^5*b^5)*d*e^2 + 5*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e^3)*x^8 + 3/7*(1
4*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^3 + 42*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^2*e +
30*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d*e^2 + 5*(3*B*a^8*b^2 + 8*A*a^7*b^3)*e^3)
*x^7 + 1/6*(42*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^3 + 90*(4*B*a^7*b^3 + 7*A*a^6
*b^4)*d^2*e + 45*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d*e^2 + 5*(2*B*a^9*b + 9*A*a^
8*b^2)*e^3)*x^6 + 1/5*(30*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^3 + 45*(3*B*a^8*b
^2 + 8*A*a^7*b^3)*d^2*e + 15*(2*B*a^9*b + 9*A*a^8*b^2)*d*e^2 + (B*a^10 + 10
*A*a^9*b)*e^3)*x^5 + 1/4*(A*a^10*e^3 + 15*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d...

```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1302 vs. $2(156) = 312$.

Time = 0.13 (sec) , antiderivative size = 1302, normalized size of antiderivative = 8.19

$$\int (a + bx)^{10}(A + Bx)(d + ex)^3 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)**10*(B*x+A)*(e*x+d)**3,x)
```

output

```
A*a**10*d**3*x + B*b**10*e**3*x**15/15 + x**14*(A*b**10*e**3/14 + 5*B*a*b*
*9*e**3/7 + 3*B*b**10*d*e**2/14) + x**13*(10*A*a*b**9*e**3/13 + 3*A*b**10*
d*e**2/13 + 45*B*a**2*b**8*e**3/13 + 30*B*a*b**9*d*e**2/13 + 3*B*b**10*d**
2*e/13) + x**12*(15*A*a**2*b**8*e**3/4 + 5*A*a*b**9*d*e**2/2 + A*b**10*d**
2*e/4 + 10*B*a**3*b**7*e**3 + 45*B*a**2*b**8*d*e**2/4 + 5*B*a*b**9*d**2*e/
2 + B*b**10*d**3/12) + x**11*(120*A*a**3*b**7*e**3/11 + 135*A*a**2*b**8*d*
e**2/11 + 30*A*a*b**9*d**2*e/11 + A*b**10*d**3/11 + 210*B*a**4*b**6*e**3/1
1 + 360*B*a**3*b**7*d*e**2/11 + 135*B*a**2*b**8*d**2*e/11 + 10*B*a*b**9*d*
*3/11) + x**10*(21*A*a**4*b**6*e**3 + 36*A*a**3*b**7*d*e**2 + 27*A*a**2*b*
*8*d**2*e/2 + A*a*b**9*d**3 + 126*B*a**5*b**5*e**3/5 + 63*B*a**4*b**6*d*e*
*2 + 36*B*a**3*b**7*d**2*e + 9*B*a**2*b**8*d**3/2) + x**9*(28*A*a**5*b**5*
e**3 + 70*A*a**4*b**6*d*e**2 + 40*A*a**3*b**7*d**2*e + 5*A*a**2*b**8*d**3
+ 70*B*a**6*b**4*e**3/3 + 84*B*a**5*b**5*d*e**2 + 70*B*a**4*b**6*d**2*e +
40*B*a**3*b**7*d**3/3) + x**8*(105*A*a**6*b**4*e**3/4 + 189*A*a**5*b**5*d*
e**2/2 + 315*A*a**4*b**6*d**2*e/4 + 15*A*a**3*b**7*d**3 + 15*B*a**7*b**3*e
**3 + 315*B*a**6*b**4*d*e**2/4 + 189*B*a**5*b**5*d**2*e/2 + 105*B*a**4*b**
6*d**3/4) + x**7*(120*A*a**7*b**3*e**3/7 + 90*A*a**6*b**4*d*e**2 + 108*A*a
**5*b**5*d**2*e + 30*A*a**4*b**6*d**3 + 45*B*a**8*b**2*e**3/7 + 360*B*a**7
*b**3*d*e**2/7 + 90*B*a**6*b**4*d**2*e + 36*B*a**5*b**5*d**3) + x**6*(15*A
*a**8*b**2*e**3/2 + 60*A*a**7*b**3*d*e**2 + 105*A*a**6*b**4*d**2*e + 42...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1068 vs. $2(149) = 298$.

Time = 0.04 (sec) , antiderivative size = 1068, normalized size of antiderivative = 6.72

$$\int (a + bx)^{10}(A + Bx)(d + ex)^3 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^3,x, algorithm="maxima")
```

output

```

1/15*B*b^10*e^3*x^15 + A*a^10*d^3*x + 1/14*(3*B*b^10*d*e^2 + (10*B*a*b^9 +
A*b^10)*e^3)*x^14 + 1/13*(3*B*b^10*d^2*e + 3*(10*B*a*b^9 + A*b^10)*d*e^2
+ 5*(9*B*a^2*b^8 + 2*A*a*b^9)*e^3)*x^13 + 1/12*(B*b^10*d^3 + 3*(10*B*a*b^9
+ A*b^10)*d^2*e + 15*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^2 + 15*(8*B*a^3*b^7 +
3*A*a^2*b^8)*e^3)*x^12 + 1/11*((10*B*a*b^9 + A*b^10)*d^3 + 15*(9*B*a^2*b^8
+ 2*A*a*b^9)*d^2*e + 45*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^2 + 30*(7*B*a^4*b
^6 + 4*A*a^3*b^7)*e^3)*x^11 + 1/10*(5*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3 + 45*(
8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e + 90*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^2 +
42*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^3)*x^10 + 1/3*(5*(8*B*a^3*b^7 + 3*A*a^2*b
^8)*d^3 + 30*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e + 42*(6*B*a^5*b^5 + 5*A*a^4
*b^6)*d*e^2 + 14*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^3)*x^9 + 3/4*(5*(7*B*a^4*b
^6 + 4*A*a^3*b^7)*d^3 + 21*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e + 21*(5*B*a^6*
b^4 + 6*A*a^5*b^5)*d*e^2 + 5*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e^3)*x^8 + 3/7*(1
4*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^3 + 42*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^2*e +
30*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d*e^2 + 5*(3*B*a^8*b^2 + 8*A*a^7*b^3)*e^3)
*x^7 + 1/6*(42*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^3 + 90*(4*B*a^7*b^3 + 7*A*a^6
*b^4)*d^2*e + 45*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d*e^2 + 5*(2*B*a^9*b + 9*A*a^
8*b^2)*e^3)*x^6 + 1/5*(30*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^3 + 45*(3*B*a^8*b
^2 + 8*A*a^7*b^3)*d^2*e + 15*(2*B*a^9*b + 9*A*a^8*b^2)*d*e^2 + (B*a^10 + 10
*A*a^9*b)*e^3)*x^5 + 1/4*(A*a^10*e^3 + 15*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d...

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1279 vs. $2(149) = 298$.

Time = 0.12 (sec) , antiderivative size = 1279, normalized size of antiderivative = 8.04

$$\int (a + bx)^{10}(A + Bx)(d + ex)^3 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^3,x, algorithm="giac")
```

output

```

1/15*B*b^10*e^3*x^15 + 3/14*B*b^10*d*e^2*x^14 + 5/7*B*a*b^9*e^3*x^14 + 1/1
4*A*b^10*e^3*x^14 + 3/13*B*b^10*d^2*e*x^13 + 30/13*B*a*b^9*d*e^2*x^13 + 3/
13*A*b^10*d*e^2*x^13 + 45/13*B*a^2*b^8*e^3*x^13 + 10/13*A*a*b^9*e^3*x^13 +
  1/12*B*b^10*d^3*x^12 + 5/2*B*a*b^9*d^2*e*x^12 + 1/4*A*b^10*d^2*e*x^12 + 4
5/4*B*a^2*b^8*d*e^2*x^12 + 5/2*A*a*b^9*d*e^2*x^12 + 10*B*a^3*b^7*e^3*x^12
+ 15/4*A*a^2*b^8*e^3*x^12 + 10/11*B*a*b^9*d^3*x^11 + 1/11*A*b^10*d^3*x^11
+ 135/11*B*a^2*b^8*d^2*e*x^11 + 30/11*A*a*b^9*d^2*e*x^11 + 360/11*B*a^3*b^
7*d*e^2*x^11 + 135/11*A*a^2*b^8*d*e^2*x^11 + 210/11*B*a^4*b^6*e^3*x^11 + 1
20/11*A*a^3*b^7*e^3*x^11 + 9/2*B*a^2*b^8*d^3*x^10 + A*a*b^9*d^3*x^10 + 36*
B*a^3*b^7*d^2*e*x^10 + 27/2*A*a^2*b^8*d^2*e*x^10 + 63*B*a^4*b^6*d*e^2*x^10
+ 36*A*a^3*b^7*d*e^2*x^10 + 126/5*B*a^5*b^5*e^3*x^10 + 21*A*a^4*b^6*e^3*x
^10 + 40/3*B*a^3*b^7*d^3*x^9 + 5*A*a^2*b^8*d^3*x^9 + 70*B*a^4*b^6*d^2*e*x
^9 + 40*A*a^3*b^7*d^2*e*x^9 + 84*B*a^5*b^5*d*e^2*x^9 + 70*A*a^4*b^6*d*e^2*x
^9 + 70/3*B*a^6*b^4*e^3*x^9 + 28*A*a^5*b^5*e^3*x^9 + 105/4*B*a^4*b^6*d^3*x
^8 + 15*A*a^3*b^7*d^3*x^8 + 189/2*B*a^5*b^5*d^2*e*x^8 + 315/4*A*a^4*b^6*d
^2*e*x^8 + 315/4*B*a^6*b^4*d*e^2*x^8 + 189/2*A*a^5*b^5*d*e^2*x^8 + 15*B*a^7
*b^3*e^3*x^8 + 105/4*A*a^6*b^4*e^3*x^8 + 36*B*a^5*b^5*d^3*x^7 + 30*A*a^4*b
^6*d^3*x^7 + 90*B*a^6*b^4*d^2*e*x^7 + 108*A*a^5*b^5*d^2*e*x^7 + 360/7*B*a
^7*b^3*d*e^2*x^7 + 90*A*a^6*b^4*d*e^2*x^7 + 45/7*B*a^8*b^2*e^3*x^7 + 120/7*
A*a^7*b^3*e^3*x^7 + 35*B*a^6*b^4*d^3*x^6 + 42*A*a^5*b^5*d^3*x^6 + 60*B*...

```

Mupad [B] (verification not implemented)

Time = 1.17 (sec) , antiderivative size = 1070, normalized size of antiderivative = 6.73

$$\int (a + bx)^{10}(A + Bx)(d + ex)^3 dx = \text{Too large to display}$$

input

```
int((A + B*x)*(a + b*x)^10*(d + e*x)^3,x)
```

output

```

x^3*((10*B*a^9*b*d^3)/3 + A*a^10*d*e^2 + B*a^10*d^2*e + 15*A*a^8*b^2*d^3 +
  10*A*a^9*b*d^2*e) + x^13*((10*A*a*b^9*e^3)/13 + (3*A*b^10*d*e^2)/13 + (3*
  B*b^10*d^2*e)/13 + (45*B*a^2*b^8*e^3)/13 + (30*B*a*b^9*d*e^2)/13) + x^4*((
  A*a^10*e^3)/4 + (3*B*a^10*d*e^2)/4 + 30*A*a^7*b^3*d^3 + (45*B*a^8*b^2*d^3)
  /4 + (135*A*a^8*b^2*d^2*e)/4 + (15*A*a^9*b*d*e^2)/2 + (15*B*a^9*b*d^2*e)/2
  ) + x^12*((B*b^10*d^3)/12 + (A*b^10*d^2*e)/4 + (15*A*a^2*b^8*e^3)/4 + 10*B
  *a^3*b^7*e^3 + (45*B*a^2*b^8*d*e^2)/4 + (5*A*a*b^9*d*e^2)/2 + (5*B*a*b^9*d
  ^2*e)/2) + x^9*(5*A*a^2*b^8*d^3 + 28*A*a^5*b^5*e^3 + (40*B*a^3*b^7*d^3)/3
  + (70*B*a^6*b^4*e^3)/3 + 40*A*a^3*b^7*d^2*e + 70*A*a^4*b^6*d*e^2 + 70*B*a^
  4*b^6*d^2*e + 84*B*a^5*b^5*d*e^2) + x^7*(30*A*a^4*b^6*d^3 + (120*A*a^7*b^3
  *e^3)/7 + 36*B*a^5*b^5*d^3 + (45*B*a^8*b^2*e^3)/7 + 108*A*a^5*b^5*d^2*e +
  90*A*a^6*b^4*d*e^2 + 90*B*a^6*b^4*d^2*e + (360*B*a^7*b^3*d*e^2)/7) + x^8*(
  15*A*a^3*b^7*d^3 + (105*A*a^6*b^4*e^3)/4 + (105*B*a^4*b^6*d^3)/4 + 15*B*a^
  7*b^3*e^3 + (315*A*a^4*b^6*d^2*e)/4 + (189*A*a^5*b^5*d*e^2)/2 + (189*B*a^5
  *b^5*d^2*e)/2 + (315*B*a^6*b^4*d*e^2)/4) + x^5*((B*a^10*e^3)/5 + 2*A*a^9*b
  *e^3 + 42*A*a^6*b^4*d^3 + 24*B*a^7*b^3*d^3 + 72*A*a^7*b^3*d^2*e + 27*A*a^8
  *b^2*d*e^2 + 27*B*a^8*b^2*d^2*e + 6*B*a^9*b*d*e^2) + x^11*((A*b^10*d^3)/11
  + (10*B*a*b^9*d^3)/11 + (120*A*a^3*b^7*e^3)/11 + (210*B*a^4*b^6*e^3)/11 +
  (135*A*a^2*b^8*d*e^2)/11 + (135*B*a^2*b^8*d^2*e)/11 + (360*B*a^3*b^7*d*e^
  2)/11 + (30*A*a*b^9*d^2*e)/11) + x^10*(A*a*b^9*d^3 + 21*A*a^4*b^6*e^3 + ...

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 653, normalized size of antiderivative = 4.11

$$\int (a + bx)^{10} (A + Bx)(d + ex)^3 dx$$

$$= \frac{x(364b^{11}e^3x^{14} + 4290ab^{10}e^3x^{13} + 1170b^{11}de^2x^{13} + 23100a^2b^9e^3x^{12} + 13860ab^{10}de^2x^{12} + 1260b^{11}d^2ex^{12} + \dots)}{1}$$

input

```
int((b*x+a)^10*(B*x+A)*(e*x+d)^3,x)
```

output

```
(x*(5460*a**11*d**3 + 8190*a**11*d**2*e*x + 5460*a**11*d*e**2*x**2 + 1365*
a**11*e**3*x**3 + 30030*a**10*b*d**3*x + 60060*a**10*b*d**2*e*x**2 + 45045
*a**10*b*d*e**2*x**3 + 12012*a**10*b*e**3*x**4 + 100100*a**9*b**2*d**3*x**
2 + 225225*a**9*b**2*d**2*e*x**3 + 180180*a**9*b**2*d*e**2*x**4 + 50050*a*
*9*b**2*e**3*x**5 + 225225*a**8*b**3*d**3*x**3 + 540540*a**8*b**3*d**2*e*x
**4 + 450450*a**8*b**3*d*e**2*x**5 + 128700*a**8*b**3*e**3*x**6 + 360360*a
**7*b**4*d**3*x**4 + 900900*a**7*b**4*d**2*e*x**5 + 772200*a**7*b**4*d*e**
2*x**6 + 225225*a**7*b**4*e**3*x**7 + 420420*a**6*b**5*d**3*x**5 + 1081080
*a**6*b**5*d**2*e*x**6 + 945945*a**6*b**5*d*e**2*x**7 + 280280*a**6*b**5*e
**3*x**8 + 360360*a**5*b**6*d**3*x**6 + 945945*a**5*b**6*d**2*e*x**7 + 840
840*a**5*b**6*d*e**2*x**8 + 252252*a**5*b**6*e**3*x**9 + 225225*a**4*b**7*
d**3*x**7 + 600600*a**4*b**7*d**2*e*x**8 + 540540*a**4*b**7*d*e**2*x**9 +
163800*a**4*b**7*e**3*x**10 + 100100*a**3*b**8*d**3*x**8 + 270270*a**3*b**
8*d**2*e*x**9 + 245700*a**3*b**8*d*e**2*x**10 + 75075*a**3*b**8*e**3*x**11
+ 30030*a**2*b**9*d**3*x**9 + 81900*a**2*b**9*d**2*e*x**10 + 75075*a**2*b
**9*d*e**2*x**11 + 23100*a**2*b**9*e**3*x**12 + 5460*a*b**10*d**3*x**10 +
15015*a*b**10*d**2*e*x**11 + 13860*a*b**10*d*e**2*x**12 + 4290*a*b**10*e**
3*x**13 + 455*b**11*d**3*x**11 + 1260*b**11*d**2*e*x**12 + 1170*b**11*d*e
**2*x**13 + 364*b**11*e**3*x**14))/5460
```

3.76 $\int (a + bx)^{10}(A + Bx)(d + ex)^2 dx$

Optimal result	778
Mathematica [B] (verified)	779
Rubi [A] (verified)	779
Maple [B] (verified)	781
Fricas [B] (verification not implemented)	782
Sympy [B] (verification not implemented)	782
Maxima [B] (verification not implemented)	783
Giac [B] (verification not implemented)	784
Mupad [B] (verification not implemented)	786
Reduce [B] (verification not implemented)	787

Optimal result

Integrand size = 20, antiderivative size = 118

$$\int (a + bx)^{10}(A + Bx)(d + ex)^2 dx = \frac{(Ab - aB)(bd - ae)^2(a + bx)^{11}}{11b^4} + \frac{(bd - ae)(bBd + 2Abe - 3aBe)(a + bx)^{12}}{12b^4} + \frac{e(2bBd + Abe - 3aBe)(a + bx)^{13}}{13b^4} + \frac{Be^2(a + bx)^{14}}{14b^4}$$

output

```
1/11*(A*b-B*a)*(-a*e+b*d)^2*(b*x+a)^11/b^4+1/12*(-a*e+b*d)*(2*A*b*e-3*B*a*
e+B*b*d)*(b*x+a)^12/b^4+1/13*e*(A*b*e-3*B*a*e+2*B*b*d)*(b*x+a)^13/b^4+1/14
*B*e^2*(b*x+a)^14/b^4
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 614 vs. $2(118) = 236$.

Time = 0.23 (sec) , antiderivative size = 614, normalized size of antiderivative = 5.20

$$\int (a + bx)^{10}(A + Bx)(d + ex)^2 dx$$

$$= \frac{x(1001a^{10}(4A(3d^2 + 3dex + e^2x^2) + Bx(6d^2 + 8dex + 3e^2x^2)) + 2002a^9bx(5A(6d^2 + 8dex + 3e^2x^2) + 2$$

input `Integrate[(a + b*x)^10*(A + B*x)*(d + e*x)^2,x]`

output

```
(x*(1001*a^10*(4*A*(3*d^2 + 3*d*e*x + e^2*x^2) + B*x*(6*d^2 + 8*d*e*x + 3*
e^2*x^2)) + 2002*a^9*b*x*(5*A*(6*d^2 + 8*d*e*x + 3*e^2*x^2) + 2*B*x*(10*d^
2 + 15*d*e*x + 6*e^2*x^2)) + 9009*a^8*b^2*x^2*(2*A*(10*d^2 + 15*d*e*x + 6*
e^2*x^2) + B*x*(15*d^2 + 24*d*e*x + 10*e^2*x^2)) + 3432*a^7*b^3*x^3*(7*A*(
15*d^2 + 24*d*e*x + 10*e^2*x^2) + 4*B*x*(21*d^2 + 35*d*e*x + 15*e^2*x^2))
+ 3003*a^6*b^4*x^4*(8*A*(21*d^2 + 35*d*e*x + 15*e^2*x^2) + 5*B*x*(28*d^2 +
48*d*e*x + 21*e^2*x^2)) + 6006*a^5*b^5*x^5*(3*A*(28*d^2 + 48*d*e*x + 21*e
^2*x^2) + 2*B*x*(36*d^2 + 63*d*e*x + 28*e^2*x^2)) + 1001*a^4*b^6*x^6*(10*A
*(36*d^2 + 63*d*e*x + 28*e^2*x^2) + 7*B*x*(45*d^2 + 80*d*e*x + 36*e^2*x^2)
) + 364*a^3*b^7*x^7*(11*A*(45*d^2 + 80*d*e*x + 36*e^2*x^2) + 8*B*x*(55*d^2
+ 99*d*e*x + 45*e^2*x^2)) + 273*a^2*b^8*x^8*(4*A*(55*d^2 + 99*d*e*x + 45*
e^2*x^2) + 3*B*x*(66*d^2 + 120*d*e*x + 55*e^2*x^2)) + 14*a*b^9*x^9*(13*A*(
66*d^2 + 120*d*e*x + 55*e^2*x^2) + 10*B*x*(78*d^2 + 143*d*e*x + 66*e^2*x^2
)) + b^10*x^10*(14*A*(78*d^2 + 143*d*e*x + 66*e^2*x^2) + 11*B*x*(91*d^2 +
168*d*e*x + 78*e^2*x^2))))/12012
```

Rubi [A] (verified)

Time = 0.71 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^{10}(A + Bx)(d + ex)^2 dx$$

↓ 86

$$\int \left(\frac{e(a + bx)^{12}(-3aBe + Abe + 2bBd)}{b^3} + \frac{(a + bx)^{11}(bd - ae)(-3aBe + 2Abe + bBd)}{b^3} + \frac{(a + bx)^{10}(Ab - aB)}{b^3} \right) dx$$

↓ 2009

$$\frac{e(a + bx)^{13}(-3aBe + Abe + 2bBd)}{13b^4} + \frac{(a + bx)^{12}(bd - ae)(-3aBe + 2Abe + bBd)}{11b^4} + \frac{12b^4}{(a + bx)^{11}(Ab - aB)(bd - ae)^2} + \frac{Be^2(a + bx)^{14}}{14b^4}$$

input `Int[(a + b*x)^10*(A + B*x)*(d + e*x)^2,x]`

output `((A*b - a*B)*(b*d - a*e)^2*(a + b*x)^11)/(11*b^4) + ((b*d - a*e)*(b*B*d + 2*A*b*e - 3*a*B*e)*(a + b*x)^12)/(12*b^4) + (e*(2*b*B*d + A*b*e - 3*a*B*e)*(a + b*x)^13)/(13*b^4) + (B*e^2*(a + b*x)^14)/(14*b^4)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 768 vs. 2(110) = 220.

Time = 0.20 (sec) , antiderivative size = 769, normalized size of antiderivative = 6.52

method	result
default	$\frac{b^{10} B e^2 x^{14}}{14} + \frac{((b^{10} A + 10 a b^9 B) e^2 + 2 b^{10} B d e) x^{13}}{13} + \frac{((10 a b^9 A + 45 a^2 b^8 B) e^2 + 2 (b^{10} A + 10 a b^9 B) d e + b^{10} B d^2) x^{12}}{12} + \dots$
norman	$a^{10} A d^2 x + (a^{10} A d e + 5 A a^9 b d^2 + \frac{1}{2} B a^{10} d^2) x^2 + (\frac{1}{3} a^{10} A e^2 + \frac{20}{3} A a^9 b d e + 15 A a^8 b^2 d^2 + \dots$
orering	$x(858 b^{10} B e^2 x^{13} + 924 A b^{10} e^2 x^{12} + 9240 B a b^9 e^2 x^{12} + 1848 B b^{10} d e x^{12} + 10010 A a b^9 e^2 x^{11} + 2002 A b^{10} d e x^{11} + 45045 B a^2 b^8 e^2 \dots$
gospers	$70 x^6 A a^6 b^4 d e + 40 x^6 B a^7 b^3 d e + 72 x^7 A a^5 b^5 d e + 60 x^7 B a^6 b^4 d e + \frac{105}{2} x^8 A a^4 b^6 d e + 63 x^8 B a^5 b^5 d e + \dots$
risch	$70 x^6 A a^6 b^4 d e + 40 x^6 B a^7 b^3 d e + 72 x^7 A a^5 b^5 d e + 60 x^7 B a^6 b^4 d e + \frac{105}{2} x^8 A a^4 b^6 d e + 63 x^8 B a^5 b^5 d e + \dots$
parallelrisch	$70 x^6 A a^6 b^4 d e + 40 x^6 B a^7 b^3 d e + 72 x^7 A a^5 b^5 d e + 60 x^7 B a^6 b^4 d e + \frac{105}{2} x^8 A a^4 b^6 d e + 63 x^8 B a^5 b^5 d e + \dots$

```
input int((b*x+a)^10*(B*x+A)*(e*x+d)^2,x,method=_RETURNVERBOSE)
```

```
output 1/14*b^10*B*e^2*x^14+1/13*((A*b^10+10*B*a*b^9)*e^2+2*b^10*B*d*e)*x^13+1/12
*((10*A*a*b^9+45*B*a^2*b^8)*e^2+2*(A*b^10+10*B*a*b^9)*d*e+b^10*B*d^2)*x^12
+1/11*((45*A*a^2*b^8+120*B*a^3*b^7)*e^2+2*(10*A*a*b^9+45*B*a^2*b^8)*d*e+(A
*b^10+10*B*a*b^9)*d^2)*x^11+1/10*((120*A*a^3*b^7+210*B*a^4*b^6)*e^2+2*(45*
A*a^2*b^8+120*B*a^3*b^7)*d*e+(10*A*a*b^9+45*B*a^2*b^8)*d^2)*x^10+1/9*((210
*A*a^4*b^6+252*B*a^5*b^5)*e^2+2*(120*A*a^3*b^7+210*B*a^4*b^6)*d*e+(45*A*a^
2*b^8+120*B*a^3*b^7)*d^2)*x^9+1/8*((252*A*a^5*b^5+210*B*a^6*b^4)*e^2+2*(21
0*A*a^4*b^6+252*B*a^5*b^5)*d*e+(120*A*a^3*b^7+210*B*a^4*b^6)*d^2)*x^8+1/7*
((210*A*a^6*b^4+120*B*a^7*b^3)*e^2+2*(252*A*a^5*b^5+210*B*a^6*b^4)*d*e+(21
0*A*a^4*b^6+252*B*a^5*b^5)*d^2)*x^7+1/6*((120*A*a^7*b^3+45*B*a^8*b^2)*e^2+
2*(210*A*a^6*b^4+120*B*a^7*b^3)*d*e+(252*A*a^5*b^5+210*B*a^6*b^4)*d^2)*x^6
+1/5*((45*A*a^8*b^2+10*B*a^9*b)*e^2+2*(120*A*a^7*b^3+45*B*a^8*b^2)*d*e+(21
0*A*a^6*b^4+120*B*a^7*b^3)*d^2)*x^5+1/4*((10*A*a^9*b+B*a^10)*e^2+2*(45*A*a
^8*b^2+10*B*a^9*b)*d*e+(120*A*a^7*b^3+45*B*a^8*b^2)*d^2)*x^4+1/3*(a^10*A*e
^2+2*(10*A*a^9*b+B*a^10)*d*e+(45*A*a^8*b^2+10*B*a^9*b)*d^2)*x^3+1/2*(2*a^1
0*A*d*e+(10*A*a^9*b+B*a^10)*d^2)*x^2+a^10*A*d^2*x
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 781 vs. $2(110) = 220$.

Time = 0.09 (sec) , antiderivative size = 781, normalized size of antiderivative = 6.62

$$\int (a + bx)^{10}(A + Bx)(d + ex)^2 dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)*(e*x+d)^2,x, algorithm="fricas")`

output

```
1/14*B*b^10*e^2*x^14 + A*a^10*d^2*x + 1/13*(2*B*b^10*d*e + (10*B*a*b^9 + A
*b^10)*e^2)*x^13 + 1/12*(B*b^10*d^2 + 2*(10*B*a*b^9 + A*b^10)*d*e + 5*(9*B
*a^2*b^8 + 2*A*a*b^9)*e^2)*x^12 + 1/11*((10*B*a*b^9 + A*b^10)*d^2 + 10*(9*
B*a^2*b^8 + 2*A*a*b^9)*d*e + 15*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^2)*x^11 + 1/
2*((9*B*a^2*b^8 + 2*A*a*b^9)*d^2 + 6*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e + 6*(
7*B*a^4*b^6 + 4*A*a^3*b^7)*e^2)*x^10 + 1/3*(5*(8*B*a^3*b^7 + 3*A*a^2*b^8)*
d^2 + 20*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e + 14*(6*B*a^5*b^5 + 5*A*a^4*b^6)*
e^2)*x^9 + 3/4*(5*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2 + 14*(6*B*a^5*b^5 + 5*A*
a^4*b^6)*d*e + 7*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^2)*x^8 + 6/7*(7*(6*B*a^5*b^
5 + 5*A*a^4*b^6)*d^2 + 14*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d*e + 5*(4*B*a^7*b^3
+ 7*A*a^6*b^4)*e^2)*x^7 + 1/2*(14*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^2 + 20*(4
*B*a^7*b^3 + 7*A*a^6*b^4)*d*e + 5*(3*B*a^8*b^2 + 8*A*a^7*b^3)*e^2)*x^6 + (
6*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^2 + 6*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d*e + (2
*B*a^9*b + 9*A*a^8*b^2)*e^2)*x^5 + 1/4*(15*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^2
+ 10*(2*B*a^9*b + 9*A*a^8*b^2)*d*e + (B*a^10 + 10*A*a^9*b)*e^2)*x^4 + 1/3
*(A*a^10*e^2 + 5*(2*B*a^9*b + 9*A*a^8*b^2)*d^2 + 2*(B*a^10 + 10*A*a^9*b)*d
*e)*x^3 + 1/2*(2*A*a^10*d*e + (B*a^10 + 10*A*a^9*b)*d^2)*x^2
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 921 vs. $2(116) = 232$.

Time = 0.09 (sec) , antiderivative size = 921, normalized size of antiderivative = 7.81

$$\int (a + bx)^{10}(A + Bx)(d + ex)^2 dx = \text{Too large to display}$$

input `integrate((b*x+a)**10*(B*x+A)*(e*x+d)**2,x)`

output

```

A*a**10*d**2*x + B*b**10*e**2*x**14/14 + x**13*(A*b**10*e**2/13 + 10*B*a*b
**9*e**2/13 + 2*B*b**10*d*e/13) + x**12*(5*A*a*b**9*e**2/6 + A*b**10*d*e/6
+ 15*B*a**2*b**8*e**2/4 + 5*B*a*b**9*d*e/3 + B*b**10*d**2/12) + x**11*(45
*A*a**2*b**8*e**2/11 + 20*A*a*b**9*d*e/11 + A*b**10*d**2/11 + 120*B*a**3*b
**7*e**2/11 + 90*B*a**2*b**8*d*e/11 + 10*B*a*b**9*d**2/11) + x**10*(12*A*a
**3*b**7*e**2 + 9*A*a**2*b**8*d*e + A*a*b**9*d**2 + 21*B*a**4*b**6*e**2 +
24*B*a**3*b**7*d*e + 9*B*a**2*b**8*d**2/2) + x**9*(70*A*a**4*b**6*e**2/3 +
80*A*a**3*b**7*d*e/3 + 5*A*a**2*b**8*d**2 + 28*B*a**5*b**5*e**2 + 140*B*a
**4*b**6*d*e/3 + 40*B*a**3*b**7*d**2/3) + x**8*(63*A*a**5*b**5*e**2/2 + 10
5*A*a**4*b**6*d*e/2 + 15*A*a**3*b**7*d**2 + 105*B*a**6*b**4*e**2/4 + 63*B*
a**5*b**5*d*e + 105*B*a**4*b**6*d**2/4) + x**7*(30*A*a**6*b**4*e**2 + 72*A
*a**5*b**5*d*e + 30*A*a**4*b**6*d**2 + 120*B*a**7*b**3*e**2/7 + 60*B*a**6*
b**4*d*e + 36*B*a**5*b**5*d**2) + x**6*(20*A*a**7*b**3*e**2 + 70*A*a**6*b*
**4*d*e + 42*A*a**5*b**5*d**2 + 15*B*a**8*b**2*e**2/2 + 40*B*a**7*b**3*d*e
+ 35*B*a**6*b**4*d**2) + x**5*(9*A*a**8*b**2*e**2 + 48*A*a**7*b**3*d*e + 4
2*A*a**6*b**4*d**2 + 2*B*a**9*b*e**2 + 18*B*a**8*b**2*d*e + 24*B*a**7*b**3
*d**2) + x**4*(5*A*a**9*b*e**2/2 + 45*A*a**8*b**2*d*e/2 + 30*A*a**7*b**3*d
**2 + B*a**10*e**2/4 + 5*B*a**9*b*d*e + 45*B*a**8*b**2*d**2/4) + x**3*(A*a
**10*e**2/3 + 20*A*a**9*b*d*e/3 + 15*A*a**8*b**2*d**2 + 2*B*a**10*d*e/3 +
10*B*a**9*b*d**2/3) + x**2*(A*a**10*d*e + 5*A*a**9*b*d**2 + B*a**10*d**...

```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 781 vs. $2(110) = 220$.

Time = 0.05 (sec) , antiderivative size = 781, normalized size of antiderivative = 6.62

$$\int (a + bx)^{10}(A + Bx)(d + ex)^2 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^2,x, algorithm="maxima")
```

output

```

1/14*B*b^10*e^2*x^14 + A*a^10*d^2*x + 1/13*(2*B*b^10*d*e + (10*B*a*b^9 + A
*b^10)*e^2)*x^13 + 1/12*(B*b^10*d^2 + 2*(10*B*a*b^9 + A*b^10)*d*e + 5*(9*B
*a^2*b^8 + 2*A*a*b^9)*e^2)*x^12 + 1/11*((10*B*a*b^9 + A*b^10)*d^2 + 10*(9*
B*a^2*b^8 + 2*A*a*b^9)*d*e + 15*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^2)*x^11 + 1/
2*((9*B*a^2*b^8 + 2*A*a*b^9)*d^2 + 6*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e + 6*(
7*B*a^4*b^6 + 4*A*a^3*b^7)*e^2)*x^10 + 1/3*(5*(8*B*a^3*b^7 + 3*A*a^2*b^8)*
d^2 + 20*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e + 14*(6*B*a^5*b^5 + 5*A*a^4*b^6)*
e^2)*x^9 + 3/4*(5*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2 + 14*(6*B*a^5*b^5 + 5*A*
a^4*b^6)*d*e + 7*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^2)*x^8 + 6/7*(7*(6*B*a^5*b^
5 + 5*A*a^4*b^6)*d^2 + 14*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d*e + 5*(4*B*a^7*b^3
+ 7*A*a^6*b^4)*e^2)*x^7 + 1/2*(14*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^2 + 20*(4
*B*a^7*b^3 + 7*A*a^6*b^4)*d*e + 5*(3*B*a^8*b^2 + 8*A*a^7*b^3)*e^2)*x^6 + (
6*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^2 + 6*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d*e + (2
*B*a^9*b + 9*A*a^8*b^2)*e^2)*x^5 + 1/4*(15*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^2
+ 10*(2*B*a^9*b + 9*A*a^8*b^2)*d*e + (B*a^10 + 10*A*a^9*b)*e^2)*x^4 + 1/3
*(A*a^10*e^2 + 5*(2*B*a^9*b + 9*A*a^8*b^2)*d^2 + 2*(B*a^10 + 10*A*a^9*b)*d
*e)*x^3 + 1/2*(2*A*a^10*d*e + (B*a^10 + 10*A*a^9*b)*d^2)*x^2

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 904 vs. $2(110) = 220$.

Time = 0.13 (sec) , antiderivative size = 904, normalized size of antiderivative = 7.66

$$\int (a + bx)^{10}(A + Bx)(d + ex)^2 dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)*(e*x+d)^2,x, algorithm="giac")
```

output

```

1/14*B*b^10*e^2*x^14 + 2/13*B*b^10*d*e*x^13 + 10/13*B*a*b^9*e^2*x^13 + 1/1
3*A*b^10*e^2*x^13 + 1/12*B*b^10*d^2*x^12 + 5/3*B*a*b^9*d*e*x^12 + 1/6*A*b^
10*d*e*x^12 + 15/4*B*a^2*b^8*e^2*x^12 + 5/6*A*a*b^9*e^2*x^12 + 10/11*B*a*b
^9*d^2*x^11 + 1/11*A*b^10*d^2*x^11 + 90/11*B*a^2*b^8*d*e*x^11 + 20/11*A*a*
b^9*d*e*x^11 + 120/11*B*a^3*b^7*e^2*x^11 + 45/11*A*a^2*b^8*e^2*x^11 + 9/2*
B*a^2*b^8*d^2*x^10 + A*a*b^9*d^2*x^10 + 24*B*a^3*b^7*d*e*x^10 + 9*A*a^2*b^
8*d*e*x^10 + 21*B*a^4*b^6*e^2*x^10 + 12*A*a^3*b^7*e^2*x^10 + 40/3*B*a^3*b^
7*d^2*x^9 + 5*A*a^2*b^8*d^2*x^9 + 140/3*B*a^4*b^6*d*e*x^9 + 80/3*A*a^3*b^7
*d*e*x^9 + 28*B*a^5*b^5*e^2*x^9 + 70/3*A*a^4*b^6*e^2*x^9 + 105/4*B*a^4*b^6
*d^2*x^8 + 15*A*a^3*b^7*d^2*x^8 + 63*B*a^5*b^5*d*e*x^8 + 105/2*A*a^4*b^6*d
*e*x^8 + 105/4*B*a^6*b^4*e^2*x^8 + 63/2*A*a^5*b^5*e^2*x^8 + 36*B*a^5*b^5*d
^2*x^7 + 30*A*a^4*b^6*d^2*x^7 + 60*B*a^6*b^4*d*e*x^7 + 72*A*a^5*b^5*d*e*x^
7 + 120/7*B*a^7*b^3*e^2*x^7 + 30*A*a^6*b^4*e^2*x^7 + 35*B*a^6*b^4*d^2*x^6
+ 42*A*a^5*b^5*d^2*x^6 + 40*B*a^7*b^3*d*e*x^6 + 70*A*a^6*b^4*d*e*x^6 + 15/
2*B*a^8*b^2*e^2*x^6 + 20*A*a^7*b^3*e^2*x^6 + 24*B*a^7*b^3*d^2*x^5 + 42*A*a
^6*b^4*d^2*x^5 + 18*B*a^8*b^2*d*e*x^5 + 48*A*a^7*b^3*d*e*x^5 + 2*B*a^9*b*e
^2*x^5 + 9*A*a^8*b^2*e^2*x^5 + 45/4*B*a^8*b^2*d^2*x^4 + 30*A*a^7*b^3*d^2*x
^4 + 5*B*a^9*b*d*e*x^4 + 45/2*A*a^8*b^2*d*e*x^4 + 1/4*B*a^10*e^2*x^4 + 5/2
*A*a^9*b*e^2*x^4 + 10/3*B*a^9*b*d^2*x^3 + 15*A*a^8*b^2*d^2*x^3 + 2/3*B*a^1
0*d*e*x^3 + 20/3*A*a^9*b*d*e*x^3 + 1/3*A*a^10*e^2*x^3 + 1/2*B*a^10*d^2*...

```

Mupad [B] (verification not implemented)

Time = 1.16 (sec) , antiderivative size = 757, normalized size of antiderivative = 6.42

$$\begin{aligned}
\int (a + bx)^{10} (A + Bx)(d + ex)^2 dx = & x^6 \left(\frac{15 B a^8 b^2 e^2}{2} + 40 B a^7 b^3 d e + 20 A a^7 b^3 e^2 \right. \\
& \left. + 35 B a^6 b^4 d^2 + 70 A a^6 b^4 d e + 42 A a^5 b^5 d^2 \right) \\
& + x^7 \left(\frac{120 B a^7 b^3 e^2}{7} + 60 B a^6 b^4 d e + 30 A a^6 b^4 e^2 \right. \\
& \left. + 36 B a^5 b^5 d^2 + 72 A a^5 b^5 d e + 30 A a^4 b^6 d^2 \right) \\
& + x^9 \left(28 B a^5 b^5 e^2 + \frac{140 B a^4 b^6 d e}{3} + \frac{70 A a^4 b^6 e^2}{3} \right. \\
& \left. + \frac{40 B a^3 b^7 d^2}{3} + \frac{80 A a^3 b^7 d e}{3} + 5 A a^2 b^8 d^2 \right) \\
& + x^8 \left(\frac{105 B a^6 b^4 e^2}{4} + 63 B a^5 b^5 d e + \frac{63 A a^5 b^5 e^2}{2} \right. \\
& \left. + \frac{105 B a^4 b^6 d^2}{4} + \frac{105 A a^4 b^6 d e}{2} + 15 A a^3 b^7 d^2 \right) \\
& + x^4 \left(\frac{B a^{10} e^2}{4} + 5 B a^9 b d e + \frac{5 A a^9 b e^2}{2} \right. \\
& \left. + \frac{45 B a^8 b^2 d^2}{4} + \frac{45 A a^8 b^2 d e}{2} + 30 A a^7 b^3 d^2 \right) \\
& + x^{11} \left(\frac{120 B a^3 b^7 e^2}{11} + \frac{90 B a^2 b^8 d e}{11} \right. \\
& \left. + \frac{45 A a^2 b^8 e^2}{11} + \frac{10 B a b^9 d^2}{11} + \frac{20 A a b^9 d e}{11} \right. \\
& \left. + \frac{A b^{10} d^2}{11} \right) \\
& + x^{10} \left(21 B a^4 b^6 e^2 + 24 B a^3 b^7 d e + 12 A a^3 b^7 e^2 \right. \\
& \left. + \frac{9 B a^2 b^8 d^2}{2} + 9 A a^2 b^8 d e + A a b^9 d^2 \right) \\
& + x^5 \left(2 B a^9 b e^2 + 18 B a^8 b^2 d e + 9 A a^8 b^2 e^2 \right. \\
& \left. + 24 B a^7 b^3 d^2 + 48 A a^7 b^3 d e + 42 A a^6 b^4 d^2 \right) \\
& + x^3 \left(\frac{2 B a^{10} d e}{3} + \frac{A a^{10} e^2}{3} + \frac{10 B a^9 b d^2}{3} \right. \\
& \left. + \frac{20 A a^9 b d e}{3} + 15 A a^8 b^2 d^2 \right) + x^{12} \left(\frac{15 B a^2 b^8 e^2}{4} \right. \\
& \left. + \frac{5 B a b^9 d e}{3} + \frac{5 A a b^9 e^2}{6} + \frac{B b^{10} d^2}{12} + \frac{A b^{10} d e}{6} \right) \\
& + A a^{10} d^2 x + \frac{a^9 d x^2 (2 A a e + 10 A b d + B a d)}{2} \\
& + \frac{b^9 e x^{13} (A b e + 10 B a e + 2 B b d)}{2} + \frac{B b^{10} e^2 x^{14}}{2}
\end{aligned}$$

input `int((A + B*x)*(a + b*x)^10*(d + e*x)^2,x)`

output $x^6(42Aa^5b^5d^2 + 20Aa^7b^3e^2 + 35Ba^6b^4d^2 + (15Ba^8b^2e^2)/2 + 70Aa^6b^4d^2e + 40Ba^7b^3d^2e) + x^7(30Aa^4b^6d^2 + 30Aa^6b^4e^2 + 36Ba^5b^5d^2 + (120Ba^7b^3e^2)/7 + 72Aa^5b^5d^2e + 60Ba^6b^4d^2e) + x^9(5Aa^2b^8d^2 + (70Aa^4b^6e^2)/3 + (40Ba^3b^7d^2)/3 + 28Ba^5b^5e^2 + (80Aa^3b^7d^2e)/3 + (140Ba^4b^6d^2e)/3) + x^8(15Aa^3b^7d^2 + (63Aa^5b^5e^2)/2 + (105Ba^4b^6d^2)/4 + (105Ba^6b^4e^2)/4 + (105Aa^4b^6d^2e)/2 + 63Ba^5b^5d^2e) + x^4((Ba^10e^2)/4 + (5Aa^9b^2e^2)/2 + 30Aa^7b^3d^2 + (45Ba^8b^2d^2)/4 + 5Ba^9b^2d^2e + (45Aa^8b^2d^2e)/2) + x^{11}((Ab^{10}d^2)/11 + (10Ba^9b^2d^2)/11 + (45Aa^2b^8e^2)/11 + (120Ba^3b^7e^2)/11 + (20Aa^2b^9d^2e)/11 + (90Ba^2b^8d^2e)/11) + x^{10}(Aa^2b^9d^2 + 12Aa^3b^7e^2 + (9Ba^2b^8d^2)/2 + 21Ba^4b^6e^2 + 9Aa^2b^8d^2e + 24Ba^3b^7d^2e) + x^5(2Ba^9b^2e^2 + 42Aa^6b^4d^2 + 9Aa^8b^2e^2 + 24Ba^7b^3d^2 + 48Aa^7b^3d^2e + 18Ba^8b^2d^2e) + x^3((Aa^{10}e^2)/3 + (2Ba^{10}d^2e)/3 + (10Ba^9b^2d^2)/3 + 15Aa^8b^2d^2 + (20Aa^9b^2d^2e)/3) + x^{12}((Bb^{10}d^2)/12 + (Ab^{10}d^2e)/6 + (5Aa^2b^9e^2)/6 + (15Ba^2b^8e^2)/4 + (5Ba^2b^9d^2e)/3) + Aa^{10}d^2x + (a^9d^2x^2(2Aa^2e + 10Abd + B^2ad))/2 + (b^9e^2x^{13}(A^2be + 10B^2ae + 2B^2bd))/13 + (Bb^{10}e^2x^{14})/14$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 459, normalized size of antiderivative = 3.89

$$\int (a + bx)^{10}(A + Bx)(d + ex)^2 dx$$

$$= \frac{x(78b^{11}e^2x^{13} + 924ab^{10}e^2x^{12} + 168b^{11}dex^{12} + 5005a^2b^9e^2x^{11} + 2002ab^{10}dex^{11} + 91b^{11}d^2x^{11} + 16380a^3b^9dex^{10} + 16380a^2b^{10}d^2x^9 + 16380a^3b^9dex^9 + 16380a^2b^{10}d^2x^8 + 16380a^3b^9dex^8 + 16380a^2b^{10}d^2x^7 + 16380a^3b^9dex^7 + 16380a^2b^{10}d^2x^6 + 16380a^3b^9dex^6 + 16380a^2b^{10}d^2x^5 + 16380a^3b^9dex^5 + 16380a^2b^{10}d^2x^4 + 16380a^3b^9dex^4 + 16380a^2b^{10}d^2x^3 + 16380a^3b^9dex^3 + 16380a^2b^{10}d^2x^2 + 16380a^3b^9dex^2 + 16380a^2b^{10}d^2x + 16380a^3b^9dex)}{13}$$

input `int((b*x+a)^10*(B*x+A)*(e*x+d)^2,x)`

output

```
(x*(1092*a**11*d**2 + 1092*a**11*d*e*x + 364*a**11*e**2*x**2 + 6006*a**10*
b*d**2*x + 8008*a**10*b*d*e*x**2 + 3003*a**10*b*e**2*x**3 + 20020*a**9*b**
2*d**2*x**2 + 30030*a**9*b**2*d*e*x**3 + 12012*a**9*b**2*e**2*x**4 + 45045
*a**8*b**3*d**2*x**3 + 72072*a**8*b**3*d*e*x**4 + 30030*a**8*b**3*e**2*x**
5 + 72072*a**7*b**4*d**2*x**4 + 120120*a**7*b**4*d*e*x**5 + 51480*a**7*b**
4*e**2*x**6 + 84084*a**6*b**5*d**2*x**5 + 144144*a**6*b**5*d*e*x**6 + 6306
3*a**6*b**5*e**2*x**7 + 72072*a**5*b**6*d**2*x**6 + 126126*a**5*b**6*d*e*x
**7 + 56056*a**5*b**6*e**2*x**8 + 45045*a**4*b**7*d**2*x**7 + 80080*a**4*b
**7*d*e*x**8 + 36036*a**4*b**7*e**2*x**9 + 20020*a**3*b**8*d**2*x**8 + 360
36*a**3*b**8*d*e*x**9 + 16380*a**3*b**8*e**2*x**10 + 6006*a**2*b**9*d**2*x
**9 + 10920*a**2*b**9*d*e*x**10 + 5005*a**2*b**9*e**2*x**11 + 1092*a*b**10
*d**2*x**10 + 2002*a*b**10*d*e*x**11 + 924*a*b**10*e**2*x**12 + 91*b**11*d
**2*x**11 + 168*b**11*d*e*x**12 + 78*b**11*e**2*x**13))/1092
```

3.77 $\int (a + bx)^{10}(A + Bx)(d + ex) dx$

Optimal result	789
Mathematica [B] (verified)	790
Rubi [A] (verified)	791
Maple [B] (verified)	792
Fricas [B] (verification not implemented)	793
Sympy [B] (verification not implemented)	794
Maxima [B] (verification not implemented)	795
Giac [B] (verification not implemented)	796
Mupad [B] (verification not implemented)	798
Reduce [B] (verification not implemented)	799

Optimal result

Integrand size = 18, antiderivative size = 75

$$\int (a + bx)^{10}(A + Bx)(d + ex) dx = \frac{(Ab - aB)(bd - ae)(a + bx)^{11}}{11b^3} + \frac{(bBd + Abe - 2aBe)(a + bx)^{12}}{12b^3} + \frac{Be(a + bx)^{13}}{13b^3}$$

output

```
1/11*(A*b-B*a)*(-a*e+b*d)*(b*x+a)^11/b^3+1/12*(A*b*e-2*B*a*e+B*b*d)*(b*x+a)^12/b^3+1/13*B*e*(b*x+a)^13/b^3
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 383 vs. $2(75) = 150$.

Time = 0.11 (sec) , antiderivative size = 383, normalized size of antiderivative = 5.11

$$\int (a + bx)^{10}(A + Bx)(d + ex) dx = \frac{1}{66}ab^9x^{10}(66Ad + 60Bdx + 60Aex + 55Bex^2) + \frac{1}{22}a^2b^8x^9(110Ad + 99Bdx + 99Aex + 90Bex^2) + \frac{1}{6}a^{10}x(3A(2d + ex) + Bx(3d + 2ex)) + \frac{3}{4}a^8b^2x^3(5A(4d + 3ex) + 3Bx(5d + 4ex)) + \frac{5}{6}a^9bx^2(Bx(4d + 3ex) + A(6d + 4ex)) + 2a^7b^3x^4(3A(5d + 4ex) + 2Bx(6d + 5ex)) + a^6b^4x^5(7A(6d + 5ex) + 5Bx(7d + 6ex)) + \frac{3}{2}a^5b^5x^6(4A(7d + 6ex) + 3Bx(8d + 7ex)) + \frac{5}{12}a^4b^6x^7(9A(8d + 7ex) + 7Bx(9d + 8ex)) + \frac{1}{3}a^3b^7x^8(5A(9d + 8ex) + 4Bx(10d + 9ex)) + \frac{b^{10}x^{11}(13A(12d + 11ex) + 11Bx(13d + 12ex))}{1716}$$

input `Integrate[(a + b*x)^10*(A + B*x)*(d + e*x), x]`

output `(a*b^9*x^10*(66*A*d + 60*B*d*x + 60*A*e*x + 55*B*e*x^2))/66 + (a^2*b^8*x^9*(110*A*d + 99*B*d*x + 99*A*e*x + 90*B*e*x^2))/22 + (a^10*x*(3*A*(2*d + e*x) + B*x*(3*d + 2*e*x)))/6 + (3*a^8*b^2*x^3*(5*A*(4*d + 3*e*x) + 3*B*x*(5*d + 4*e*x)))/4 + (5*a^9*b*x^2*(B*x*(4*d + 3*e*x) + A*(6*d + 4*e*x)))/6 + 2*a^7*b^3*x^4*(3*A*(5*d + 4*e*x) + 2*B*x*(6*d + 5*e*x)) + a^6*b^4*x^5*(7*A*(6*d + 5*e*x) + 5*B*x*(7*d + 6*e*x)) + (3*a^5*b^5*x^6*(4*A*(7*d + 6*e*x) + 3*B*x*(8*d + 7*e*x)))/2 + (5*a^4*b^6*x^7*(9*A*(8*d + 7*e*x) + 7*B*x*(9*d + 8*e*x)))/12 + (a^3*b^7*x^8*(5*A*(9*d + 8*e*x) + 4*B*x*(10*d + 9*e*x)))/3 + (b^10*x^11*(13*A*(12*d + 11*e*x) + 11*B*x*(13*d + 12*e*x)))/1716`

Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^{10}(A + Bx)(d + ex) dx$$

$$\downarrow 86$$

$$\int \left(\frac{(a + bx)^{11}(-2aBe + Abe + bBd)}{b^2} + \frac{(a + bx)^{10}(Ab - aB)(bd - ae)}{b^2} + \frac{Be(a + bx)^{12}}{b^2} \right) dx$$

$$\downarrow 2009$$

$$\frac{(a + bx)^{12}(-2aBe + Abe + bBd)}{12b^3} + \frac{(a + bx)^{11}(Ab - aB)(bd - ae)}{11b^3} + \frac{Be(a + bx)^{13}}{13b^3}$$

input `Int[(a + b*x)^10*(A + B*x)*(d + e*x),x]`

output `((A*b - a*B)*(b*d - a*e)*(a + b*x)^11)/(11*b^3) + ((b*B*d + A*b*e - 2*a*B*e)*(a + b*x)^12)/(12*b^3) + (B*e*(a + b*x)^13)/(13*b^3)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 458 vs. $2(69) = 138$.

Time = 0.17 (sec) , antiderivative size = 459, normalized size of antiderivative = 6.12

method	result
norman	$\frac{b^{10} B e x^{13}}{13} + \left(\frac{1}{12} A b^{10} e + \frac{5}{6} B a b^9 e + \frac{1}{12} b^{10} B d\right) x^{12} + \left(\frac{10}{11} A a b^9 e + \frac{1}{11} A b^{10} d + \frac{45}{11} B a^2 b^8 e + \frac{10}{11} B a^2 b^8 e + \frac{10}{11} B a^2 b^8 e\right) x^{11} + \dots$
default	$\frac{b^{10} B e x^{13}}{13} + \frac{((b^{10} A + 10 a b^9 B) e + b^{10} B d) x^{12}}{12} + \frac{((10 a b^9 A + 45 a^2 b^8 B) e + (b^{10} A + 10 a b^9 B) d) x^{11}}{11} + \frac{((45 a^2 b^8 A + 120 a^3 b^7 B) e + (10 a b^9 A + 45 a^2 b^8 B) d) x^{10}}{10} + \dots$
orering	$x(132b^{10} B e x^{12} + 143A b^{10} e x^{11} + 1430B a b^9 e x^{11} + 143B b^{10} d x^{11} + 1560A a b^9 e x^{10} + 156A b^{10} d x^{10} + 7020B a^2 b^8 e x^{10} + 1560A a^2 b^8 e x^{10} + \dots)$
gosper	$\frac{105}{4} x^8 A a^4 b^6 e + 15x^8 A a^3 b^7 d + \frac{63}{2} x^8 B a^5 b^5 e + \frac{9}{2} x^{10} A a^2 b^8 e + \frac{1}{11} x^{11} A b^{10} d + \frac{105}{4} x^8 B a^4 b^6 d + \dots$
risch	$\frac{105}{4} x^8 A a^4 b^6 e + 15x^8 A a^3 b^7 d + \frac{63}{2} x^8 B a^5 b^5 e + \frac{9}{2} x^{10} A a^2 b^8 e + \frac{1}{11} x^{11} A b^{10} d + \frac{105}{4} x^8 B a^4 b^6 d + \dots$
parallelrisch	$\frac{105}{4} x^8 A a^4 b^6 e + 15x^8 A a^3 b^7 d + \frac{63}{2} x^8 B a^5 b^5 e + \frac{9}{2} x^{10} A a^2 b^8 e + \frac{1}{11} x^{11} A b^{10} d + \frac{105}{4} x^8 B a^4 b^6 d + \dots$

```
input int((b*x+a)^10*(B*x+A)*(e*x+d), x, method=_RETURNVERBOSE)
```

```
output 1/13*b^10*B*e*x^13+(1/12*A*b^10*e+5/6*B*a*b^9*e+1/12*b^10*B*d)*x^12+(10/11
*A*a*b^9*e+1/11*A*b^10*d+45/11*B*a^2*b^8*e+10/11*B*a*b^9*d)*x^11+(9/2*A*a^
2*b^8*e+A*a*b^9*d+12*B*a^3*b^7*e+9/2*B*a^2*b^8*d)*x^10+(40/3*A*a^3*b^7*e+5
*A*a^2*b^8*d+70/3*B*a^4*b^6*e+40/3*B*a^3*b^7*d)*x^9+(105/4*A*a^4*b^6*e+15*
A*a^3*b^7*d+63/2*B*a^5*b^5*e+105/4*B*a^4*b^6*d)*x^8+(36*A*a^5*b^5*e+30*A*a
^4*b^6*d+30*B*a^6*b^4*e+36*B*a^5*b^5*d)*x^7+(35*A*a^6*b^4*e+42*A*a^5*b^5*d
+20*B*a^7*b^3*e+35*B*a^6*b^4*d)*x^6+(24*A*a^7*b^3*e+42*A*a^6*b^4*d+9*B*a^8
*b^2*e+24*B*a^7*b^3*d)*x^5+(45/4*A*a^8*b^2*e+30*A*a^7*b^3*d+5/2*B*a^9*b*e+
45/4*B*a^8*b^2*d)*x^4+(10/3*A*a^9*b*e+15*A*a^8*b^2*d+1/3*B*a^10*e+10/3*B*a
^9*b*d)*x^3+(1/2*a^10*A*e+5*A*a^9*b*d+1/2*B*a^10*d)*x^2+a^10*A*d*x
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 493 vs. $2(69) = 138$.

Time = 0.07 (sec) , antiderivative size = 493, normalized size of antiderivative = 6.57

$$\begin{aligned} & \int (a + bx)^{10}(A + Bx)(d + ex) dx \\ &= \frac{1}{13} Bb^{10}ex^{13} + Aa^{10}dx + \frac{1}{12} (Bb^{10}d + (10 Bab^9 + Ab^{10})e)x^{12} \\ &+ \frac{1}{11} ((10 Bab^9 + Ab^{10})d + 5(9 Ba^2b^8 + 2 Aab^9)e)x^{11} \\ &+ \frac{1}{2} ((9 Ba^2b^8 + 2 Aab^9)d + 3(8 Ba^3b^7 + 3 Aa^2b^8)e)x^{10} \\ &+ \frac{5}{3} ((8 Ba^3b^7 + 3 Aa^2b^8)d + 2(7 Ba^4b^6 + 4 Aa^3b^7)e)x^9 \\ &+ \frac{3}{4} (5(7 Ba^4b^6 + 4 Aa^3b^7)d + 7(6 Ba^5b^5 + 5 Aa^4b^6)e)x^8 \\ &+ 6((6 Ba^5b^5 + 5 Aa^4b^6)d + (5 Ba^6b^4 + 6 Aa^5b^5)e)x^7 \\ &+ (7(5 Ba^6b^4 + 6 Aa^5b^5)d + 5(4 Ba^7b^3 + 7 Aa^6b^4)e)x^6 \\ &+ 3(2(4 Ba^7b^3 + 7 Aa^6b^4)d + (3 Ba^8b^2 + 8 Aa^7b^3)e)x^5 \\ &+ \frac{5}{4} (3(3 Ba^8b^2 + 8 Aa^7b^3)d + (2 Ba^9b + 9 Aa^8b^2)e)x^4 \\ &+ \frac{1}{3} (5(2 Ba^9b + 9 Aa^8b^2)d + (Ba^{10} + 10 Aa^9b)e)x^3 \\ &+ \frac{1}{2} (Aa^{10}e + (Ba^{10} + 10 Aa^9b)d)x^2 \end{aligned}$$

input `integrate((b*x+a)^10*(B*x+A)*(e*x+d),x, algorithm="fricas")`

output `1/13*B*b^10*e*x^13 + A*a^10*d*x + 1/12*(B*b^10*d + (10*B*a*b^9 + A*b^10)*e)*x^12 + 1/11*((10*B*a*b^9 + A*b^10)*d + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*e)*x^11 + 1/2*((9*B*a^2*b^8 + 2*A*a*b^9)*d + 3*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e)*x^10 + 5/3*((8*B*a^3*b^7 + 3*A*a^2*b^8)*d + 2*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e)*x^9 + 3/4*(5*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d + 7*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e)*x^8 + 6*((6*B*a^5*b^5 + 5*A*a^4*b^6)*d + (5*B*a^6*b^4 + 6*A*a^5*b^5)*e)*x^7 + (7*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d + 5*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e)*x^6 + 3*(2*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d + (3*B*a^8*b^2 + 8*A*a^7*b^3)*e)*x^5 + 5/4*(3*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d + (2*B*a^9*b + 9*A*a^8*b^2)*e)*x^4 + 1/3*(5*(2*B*a^9*b + 9*A*a^8*b^2)*d + (B*a^10 + 10*A*a^9*b)*e)*x^3 + 1/2*(A*a^10*e + (B*a^10 + 10*A*a^9*b)*d)*x^2`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 549 vs. $2(71) = 142$.

Time = 0.07 (sec) , antiderivative size = 549, normalized size of antiderivative = 7.32

$$\begin{aligned}
 \int (a + bx)^{10}(A + Bx)(d + ex) dx = & Aa^{10}dx + \frac{Bb^{10}ex^{13}}{13} \\
 & + x^{12} \left(\frac{Ab^{10}e}{12} + \frac{5Bab^9e}{6} + \frac{Bb^{10}d}{12} \right) + x^{11} \\
 & \cdot \left(\frac{10Aab^9e}{11} + \frac{Ab^{10}d}{11} + \frac{45Ba^2b^8e}{11} + \frac{10Bab^9d}{11} \right) + x^{10} \\
 & \cdot \left(\frac{9Aa^2b^8e}{2} + Aab^9d + 12Ba^3b^7e + \frac{9Ba^2b^8d}{2} \right) + x^9 \\
 & \cdot \left(\frac{40Aa^3b^7e}{3} + 5Aa^2b^8d + \frac{70Ba^4b^6e}{3} + \frac{40Ba^3b^7d}{3} \right) \\
 & + x^8 \cdot \left(\frac{105Aa^4b^6e}{4} + 15Aa^3b^7d + \frac{63Ba^5b^5e}{2} \right. \\
 & \qquad \qquad \qquad \left. + \frac{105Ba^4b^6d}{4} \right) + x^7 \\
 & \cdot (36Aa^5b^5e + 30Aa^4b^6d + 30Ba^6b^4e + 36Ba^5b^5d) \\
 & + x^6 \\
 & \cdot (35Aa^6b^4e + 42Aa^5b^5d + 20Ba^7b^3e + 35Ba^6b^4d) \\
 & + x^5 \\
 & \cdot (24Aa^7b^3e + 42Aa^6b^4d + 9Ba^8b^2e + 24Ba^7b^3d) + x^4 \\
 & \cdot \left(\frac{45Aa^8b^2e}{4} + 30Aa^7b^3d + \frac{5Ba^9be}{2} + \frac{45Ba^8b^2d}{4} \right) \\
 & + x^3 \cdot \left(\frac{10Aa^9be}{3} + 15Aa^8b^2d + \frac{Ba^{10}e}{3} + \frac{10Ba^9bd}{3} \right) \\
 & + x^2 \left(\frac{Aa^{10}e}{2} + 5Aa^9bd + \frac{Ba^{10}d}{2} \right)
 \end{aligned}$$

input `integrate((b*x+a)**10*(B*x+A)*(e*x+d), x)`

output

```

A*a**10*d*x + B*b**10*e*x**13/13 + x**12*(A*b**10*e/12 + 5*B*a*b**9*e/6 +
B*b**10*d/12) + x**11*(10*A*a*b**9*e/11 + A*b**10*d/11 + 45*B*a**2*b**8*e/
11 + 10*B*a*b**9*d/11) + x**10*(9*A*a**2*b**8*e/2 + A*a*b**9*d + 12*B*a**3
*b**7*e + 9*B*a**2*b**8*d/2) + x**9*(40*A*a**3*b**7*e/3 + 5*A*a**2*b**8*d
+ 70*B*a**4*b**6*e/3 + 40*B*a**3*b**7*d/3) + x**8*(105*A*a**4*b**6*e/4 + 1
5*A*a**3*b**7*d + 63*B*a**5*b**5*e/2 + 105*B*a**4*b**6*d/4) + x**7*(36*A*a
**5*b**5*e + 30*A*a**4*b**6*d + 30*B*a**6*b**4*e + 36*B*a**5*b**5*d) + x**
6*(35*A*a**6*b**4*e + 42*A*a**5*b**5*d + 20*B*a**7*b**3*e + 35*B*a**6*b**4
*d) + x**5*(24*A*a**7*b**3*e + 42*A*a**6*b**4*d + 9*B*a**8*b**2*e + 24*B*a
**7*b**3*d) + x**4*(45*A*a**8*b**2*e/4 + 30*A*a**7*b**3*d + 5*B*a**9*b*e/2
+ 45*B*a**8*b**2*d/4) + x**3*(10*A*a**9*b*e/3 + 15*A*a**8*b**2*d + B*a**1
0*e/3 + 10*B*a**9*b*d/3) + x**2*(A*a**10*e/2 + 5*A*a**9*b*d + B*a**10*d/2)

```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 493 vs. $2(69) = 138$.

Time = 0.04 (sec) , antiderivative size = 493, normalized size of antiderivative = 6.57

$$\begin{aligned}
& \int (a + bx)^{10} (A + Bx)(d + ex) dx \\
&= \frac{1}{13} Bb^{10} ex^{13} + Aa^{10} dx + \frac{1}{12} (Bb^{10}d + (10 Bab^9 + Ab^{10})e)x^{12} \\
&+ \frac{1}{11} ((10 Bab^9 + Ab^{10})d + 5(9 Ba^2b^8 + 2 Aab^9)e)x^{11} \\
&+ \frac{1}{2} ((9 Ba^2b^8 + 2 Aab^9)d + 3(8 Ba^3b^7 + 3 Aa^2b^8)e)x^{10} \\
&+ \frac{5}{3} ((8 Ba^3b^7 + 3 Aa^2b^8)d + 2(7 Ba^4b^6 + 4 Aa^3b^7)e)x^9 \\
&+ \frac{3}{4} (5(7 Ba^4b^6 + 4 Aa^3b^7)d + 7(6 Ba^5b^5 + 5 Aa^4b^6)e)x^8 \\
&+ 6((6 Ba^5b^5 + 5 Aa^4b^6)d + (5 Ba^6b^4 + 6 Aa^5b^5)e)x^7 \\
&+ (7(5 Ba^6b^4 + 6 Aa^5b^5)d + 5(4 Ba^7b^3 + 7 Aa^6b^4)e)x^6 \\
&+ 3(2(4 Ba^7b^3 + 7 Aa^6b^4)d + (3 Ba^8b^2 + 8 Aa^7b^3)e)x^5 \\
&+ \frac{5}{4} (3(3 Ba^8b^2 + 8 Aa^7b^3)d + (2 Ba^9b + 9 Aa^8b^2)e)x^4 \\
&+ \frac{1}{3} (5(2 Ba^9b + 9 Aa^8b^2)d + (Ba^{10} + 10 Aa^9b)e)x^3 \\
&+ \frac{1}{2} (Aa^{10}e + (Ba^{10} + 10 Aa^9b)d)x^2
\end{aligned}$$

input `integrate((b*x+a)^10*(B*x+A)*(e*x+d),x, algorithm="maxima")`

output

$$\begin{aligned} & 1/13*B*b^{10}*e*x^{13} + A*a^{10}*d*x + 1/12*(B*b^{10}*d + (10*B*a*b^9 + A*b^{10})*e) \\ & *x^{12} + 1/11*((10*B*a*b^9 + A*b^{10})*d + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*e)*x^{11} \\ & + 1/2*((9*B*a^2*b^8 + 2*A*a*b^9)*d + 3*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e)*x^{10} \\ & + 5/3*((8*B*a^3*b^7 + 3*A*a^2*b^8)*d + 2*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e) \\ & *x^9 + 3/4*(5*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d + 7*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e) \\ & *x^8 + 6*((6*B*a^5*b^5 + 5*A*a^4*b^6)*d + (5*B*a^6*b^4 + 6*A*a^5*b^5)*e) \\ & *x^7 + (7*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d + 5*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e) \\ & *x^6 + 3*(2*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d + (3*B*a^8*b^2 + 8*A*a^7*b^3)*e) \\ & *x^5 + 5/4*(3*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d + (2*B*a^9*b + 9*A*a^8*b^2)*e) \\ & *x^4 + 1/3*(5*(2*B*a^9*b + 9*A*a^8*b^2)*d + (B*a^{10} + 10*A*a^9*b)*e) \\ & *x^3 + 1/2*(A*a^{10}*e + (B*a^{10} + 10*A*a^9*b)*d)*x^2 \end{aligned}$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 529 vs. $2(69) = 138$.

Time = 0.12 (sec) , antiderivative size = 529, normalized size of antiderivative = 7.05

$$\begin{aligned}
 \int (a + bx)^{10}(A + Bx)(d + ex) dx = & \frac{1}{13} Bb^{10}ex^{13} + \frac{1}{12} Bb^{10}dx^{12} + \frac{5}{6} Bab^9ex^{12} \\
 & + \frac{1}{12} Ab^{10}ex^{12} + \frac{10}{11} Bab^9dx^{11} + \frac{1}{11} Ab^{10}dx^{11} \\
 & + \frac{45}{11} Ba^2b^8ex^{11} + \frac{10}{11} Aab^9ex^{11} + \frac{9}{2} Ba^2b^8dx^{10} \\
 & + Aab^9dx^{10} + 12 Ba^3b^7ex^{10} + \frac{9}{2} Aa^2b^8ex^{10} \\
 & + \frac{40}{3} Ba^3b^7dx^9 + 5 Aa^2b^8dx^9 + \frac{70}{3} Ba^4b^6ex^9 \\
 & + \frac{40}{3} Aa^3b^7ex^9 + \frac{105}{4} Ba^4b^6dx^8 \\
 & + 15 Aa^3b^7dx^8 + \frac{63}{2} Ba^5b^5ex^8 + \frac{105}{4} Aa^4b^6ex^8 \\
 & + 36 Ba^5b^5dx^7 + 30 Aa^4b^6dx^7 + 30 Ba^6b^4ex^7 \\
 & + 36 Aa^5b^5ex^7 + 35 Ba^6b^4dx^6 + 42 Aa^5b^5dx^6 \\
 & + 20 Ba^7b^3ex^6 + 35 Aa^6b^4ex^6 + 24 Ba^7b^3dx^5 \\
 & + 42 Aa^6b^4dx^5 + 9 Ba^8b^2ex^5 + 24 Aa^7b^3ex^5 \\
 & + \frac{45}{4} Ba^8b^2dx^4 + 30 Aa^7b^3dx^4 + \frac{5}{2} Ba^9bex^4 \\
 & + \frac{45}{4} Aa^8b^2ex^4 + \frac{10}{3} Ba^9bdx^3 + 15 Aa^8b^2dx^3 \\
 & + \frac{1}{3} Ba^{10}ex^3 + \frac{10}{3} Aa^9bex^3 + \frac{1}{2} Ba^{10}dx^2 \\
 & + 5 Aa^9bdx^2 + \frac{1}{2} Aa^{10}ex^2 + Aa^{10}dx
 \end{aligned}$$

input `integrate((b*x+a)^10*(B*x+A)*(e*x+d),x, algorithm="giac")`

output

```

1/13*B*b^10*e*x^13 + 1/12*B*b^10*d*x^12 + 5/6*B*a*b^9*e*x^12 + 1/12*A*b^10
*e*x^12 + 10/11*B*a*b^9*d*x^11 + 1/11*A*b^10*d*x^11 + 45/11*B*a^2*b^8*e*x^
11 + 10/11*A*a*b^9*e*x^11 + 9/2*B*a^2*b^8*d*x^10 + A*a*b^9*d*x^10 + 12*B*a
^3*b^7*e*x^10 + 9/2*A*a^2*b^8*e*x^10 + 40/3*B*a^3*b^7*d*x^9 + 5*A*a^2*b^8*
d*x^9 + 70/3*B*a^4*b^6*e*x^9 + 40/3*A*a^3*b^7*e*x^9 + 105/4*B*a^4*b^6*d*x^
8 + 15*A*a^3*b^7*d*x^8 + 63/2*B*a^5*b^5*e*x^8 + 105/4*A*a^4*b^6*e*x^8 + 36
*B*a^5*b^5*d*x^7 + 30*A*a^4*b^6*d*x^7 + 30*B*a^6*b^4*e*x^7 + 36*A*a^5*b^5*
e*x^7 + 35*B*a^6*b^4*d*x^6 + 42*A*a^5*b^5*d*x^6 + 20*B*a^7*b^3*e*x^6 + 35*
A*a^6*b^4*e*x^6 + 24*B*a^7*b^3*d*x^5 + 42*A*a^6*b^4*d*x^5 + 9*B*a^8*b^2*e*
x^5 + 24*A*a^7*b^3*e*x^5 + 45/4*B*a^8*b^2*d*x^4 + 30*A*a^7*b^3*d*x^4 + 5/2
*B*a^9*b*e*x^4 + 45/4*A*a^8*b^2*e*x^4 + 10/3*B*a^9*b*d*x^3 + 15*A*a^8*b^2*
d*x^3 + 1/3*B*a^10*e*x^3 + 10/3*A*a^9*b*e*x^3 + 1/2*B*a^10*d*x^2 + 5*A*a^9
*b*d*x^2 + 1/2*A*a^10*e*x^2 + A*a^10*d*x

```

Mupad [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 409, normalized size of antiderivative = 5.45

$$\begin{aligned}
& \int (a + bx)^{10} (A + Bx)(d + ex) dx \\
&= x^3 \left(\frac{Ba^{10}e}{3} + \frac{10Aa^9be}{3} + \frac{10Ba^9bd}{3} + 15Aa^8b^2d \right) \\
&+ x^{11} \left(\frac{Ab^{10}d}{11} + \frac{10Aab^9e}{11} + \frac{10Bab^9d}{11} + \frac{45Ba^2b^8e}{11} \right) \\
&+ x^2 \left(\frac{Aa^{10}e}{2} + \frac{Ba^{10}d}{2} + 5Aa^9bd \right) + x^{12} \left(\frac{Ab^{10}e}{12} + \frac{Bb^{10}d}{12} + \frac{5Bab^9e}{6} \right) \\
&+ 6a^4b^4x^7 (5Ab^2d + 5Ba^2e + 6Aabe + 6Babd) \\
&+ 3a^6b^2x^5 (14Ab^2d + 3Ba^2e + 8Aabe + 8Babd) \\
&+ \frac{5a^2b^6x^9 (3Ab^2d + 14Ba^2e + 8Aabe + 8Babd)}{3} \\
&+ a^5b^3x^6 (42Ab^2d + 20Ba^2e + 35Aabe + 35Babd) \\
&+ \frac{3a^3b^5x^8 (20Ab^2d + 42Ba^2e + 35Aabe + 35Babd)}{4} + Aa^{10}dx \\
&+ \frac{Bb^{10}ex^{13}}{13} + \frac{5a^7bx^4 (24Ab^2d + 2Ba^2e + 9Aabe + 9Babd)}{4} \\
&+ \frac{ab^7x^{10} (2Ab^2d + 24Ba^2e + 9Aabe + 9Babd)}{2}
\end{aligned}$$

input `int((A + B*x)*(a + b*x)^10*(d + e*x),x)`

output
$$\begin{aligned} & x^3 \left(\frac{B^3 a^{10} e}{3} + \frac{10 A^2 a^9 b e}{3} + \frac{10 B^2 a^9 b d}{3} + 15 A^2 a^8 b^2 d \right) \\ & + x^{11} \left(\frac{A^3 b^{10} d}{11} + \frac{10 A^2 a b^9 e}{11} + \frac{10 B^2 a b^9 d}{11} + \frac{45 B^2 a^2 b^8 e}{11} \right) \\ & + x^2 \left(\frac{A^2 a^{10} e}{2} + \frac{B^2 a^{10} d}{2} + 5 A^2 a^9 b d \right) + x^{12} \left(\frac{A^3 b^{10} e}{12} + \frac{B^3 b^{10} d}{12} + \frac{5 B^2 a b^9 e}{6} \right) \\ & + 6 a^4 b^4 x^7 (5 A^2 b^2 d + 5 B^2 a^2 e + 6 A^2 a b e + 6 B^2 a b d) + 3 a^6 b^2 x^5 (14 A^2 b^2 d + 3 B^2 a^2 e \\ & + 8 A^2 a b e + 8 B^2 a b d) + (5 a^2 b^6 x^9 (3 A^2 b^2 d + 14 B^2 a^2 e + 8 A^2 a b e + 8 B^2 a b d)) / 3 \\ & + a^5 b^3 x^6 (42 A^2 b^2 d + 20 B^2 a^2 e + 35 A^2 a b e + 35 B^2 a b d) + (3 a^3 b^5 x^8 (20 A^2 b^2 d + 42 B^2 a^2 e + 35 A^2 a b e + 35 B^2 a b d)) / 4 \\ & + A^2 a^{10} d x + (B^2 b^{10} e x^{13}) / 13 + (5 a^7 b x^4 (24 A^2 b^2 d + 2 B^2 a^2 e + 9 A^2 a b e + 9 B^2 a b d)) / 4 \\ & + (a b^7 x^{10} (2 A^2 b^2 d + 24 B^2 a^2 e + 9 A^2 a b e + 9 B^2 a b d)) / 2 \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 265, normalized size of antiderivative = 3.53

$$\int (a + bx)^{10} (A + Bx)(d + ex) dx$$

$$= \frac{x(12b^{11}e x^{12} + 143a b^{10}e x^{11} + 13b^{11}d x^{11} + 780a^2 b^9 e x^{10} + 156a b^{10}d x^{10} + 2574a^3 b^8 e x^9 + 858a^2 b^9 d x^9 + \dots)}{156}$$

input `int((b*x+a)^10*(B*x+A)*(e*x+d),x)`

output
$$\begin{aligned} & (x(156 a^{11} d + 78 a^{11} e x + 858 a^{10} b d x + 572 a^{10} b e x^2 + 28 \\ & 60 a^9 b^2 d x^2 + 2145 a^9 b^2 e x^3 + 6435 a^8 b^3 d x^3 + 5148 \\ & a^8 b^3 e x^4 + 10296 a^7 b^4 d x^4 + 8580 a^7 b^4 e x^5 + 12012 \\ & a^6 b^5 d x^5 + 10296 a^6 b^5 e x^6 + 10296 a^5 b^6 d x^6 + 9009 \\ & a^5 b^6 e x^7 + 6435 a^4 b^7 d x^7 + 5720 a^4 b^7 e x^8 + 2860 a \\ & ^3 b^8 d x^8 + 2574 a^3 b^8 e x^9 + 858 a^2 b^9 d x^9 + 780 a^2 b \\ & ^9 e x^{10} + 156 a b^{10} d x^{10} + 143 a b^{10} e x^{11} + 13 b^{11} d x^{11} \\ & + 12 b^{11} e x^{12})) / 156 \end{aligned}$$

3.78 $\int (a + bx)^{10}(A + Bx) dx$

Optimal result	800
Mathematica [B] (verified)	800
Rubi [A] (verified)	801
Maple [B] (verified)	802
Fricas [B] (verification not implemented)	803
Sympy [B] (verification not implemented)	804
Maxima [B] (verification not implemented)	804
Giac [B] (verification not implemented)	805
Mupad [B] (verification not implemented)	806
Reduce [B] (verification not implemented)	807

Optimal result

Integrand size = 13, antiderivative size = 38

$$\int (a + bx)^{10}(A + Bx) dx = \frac{(Ab - aB)(a + bx)^{11}}{11b^2} + \frac{B(a + bx)^{12}}{12b^2}$$

output

```
1/11*(A*b-B*a)*(b*x+a)^11/b^2+1/12*B*(b*x+a)^12/b^2
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 198 vs. $2(38) = 76$.

Time = 0.04 (sec) , antiderivative size = 198, normalized size of antiderivative = 5.21

$$\begin{aligned} \int (a + bx)^{10}(A + Bx) dx = & \frac{1}{132}x(66a^{10}(2A + Bx) + 220a^9bx(3A + 2Bx) \\ & + 495a^8b^2x^2(4A + 3Bx) + 792a^7b^3x^3(5A + 4Bx) \\ & + 924a^6b^4x^4(6A + 5Bx) + 792a^5b^5x^5(7A + 6Bx) \\ & + 495a^4b^6x^6(8A + 7Bx) + 220a^3b^7x^7(9A + 8Bx) \\ & + 66a^2b^8x^8(10A + 9Bx) + 12ab^9x^9(11A + 10Bx) \\ & + b^{10}x^{10}(12A + 11Bx)) \end{aligned}$$

input

```
Integrate[(a + b*x)^10*(A + B*x),x]
```

output

```
(x*(66*a^10*(2*A + B*x) + 220*a^9*b*x*(3*A + 2*B*x) + 495*a^8*b^2*x^2*(4*A
+ 3*B*x) + 792*a^7*b^3*x^3*(5*A + 4*B*x) + 924*a^6*b^4*x^4*(6*A + 5*B*x)
+ 792*a^5*b^5*x^5*(7*A + 6*B*x) + 495*a^4*b^6*x^6*(8*A + 7*B*x) + 220*a^3*
b^7*x^7*(9*A + 8*B*x) + 66*a^2*b^8*x^8*(10*A + 9*B*x) + 12*a*b^9*x^9*(11*A
+ 10*B*x) + b^10*x^10*(12*A + 11*B*x))/132
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00,
 number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules
 used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the
 transformation is given above next to the arrow. The rules definitions used are listed
 below.

$$\int (a + bx)^{10} (A + Bx) dx$$

$$\downarrow 49$$

$$\int \left(\frac{(a + bx)^{10} (Ab - aB)}{b} + \frac{B(a + bx)^{11}}{b} \right) dx$$

$$\downarrow 2009$$

$$\frac{(a + bx)^{11} (Ab - aB)}{11b^2} + \frac{B(a + bx)^{12}}{12b^2}$$

input

```
Int[(a + b*x)^10*(A + B*x),x]
```

output

```
((A*b - a*B)*(a + b*x)^11)/(11*b^2) + (B*(a + b*x)^12)/(12*b^2)
```

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 231 vs. $2(34) = 68$.

Time = 0.14 (sec) , antiderivative size = 232, normalized size of antiderivative = 6.11

method	result
norman	$\frac{b^{10} B x^{12}}{12} + \left(\frac{1}{11} b^{10} A + \frac{10}{11} a b^9 B\right) x^{11} + (a b^9 A + \frac{9}{2} a^2 b^8 B) x^{10} + (5 a^2 b^8 A + \frac{40}{3} a^3 b^7 B) x^9 + (15$
default	$\frac{b^{10} B x^{12}}{12} + \frac{(b^{10} A + 10 a b^9 B) x^{11}}{11} + \frac{(10 a b^9 A + 45 a^2 b^8 B) x^{10}}{10} + \frac{(45 a^2 b^8 A + 120 a^3 b^7 B) x^9}{9} + \frac{(120 a^3 b^7 A + 210 a^4 b^6 B) x^8}{8}$
gosper	$\frac{1}{12} b^{10} B x^{12} + \frac{1}{11} x^{11} b^{10} A + \frac{10}{11} x^{11} a b^9 B + x^{10} a b^9 A + \frac{9}{2} x^{10} a^2 b^8 B + 5 x^9 a^2 b^8 A + \frac{40}{3} x^9 a^3 b^7 B +$
risch	$\frac{1}{12} b^{10} B x^{12} + \frac{1}{11} x^{11} b^{10} A + \frac{10}{11} x^{11} a b^9 B + x^{10} a b^9 A + \frac{9}{2} x^{10} a^2 b^8 B + 5 x^9 a^2 b^8 A + \frac{40}{3} x^9 a^3 b^7 B +$
parallelrisch	$\frac{1}{12} b^{10} B x^{12} + \frac{1}{11} x^{11} b^{10} A + \frac{10}{11} x^{11} a b^9 B + x^{10} a b^9 A + \frac{9}{2} x^{10} a^2 b^8 B + 5 x^9 a^2 b^8 A + \frac{40}{3} x^9 a^3 b^7 B +$
orering	$x(11 B b^{10} x^{11} + 12 A b^{10} x^{10} + 120 B a b^9 x^{10} + 132 a A b^9 x^9 + 594 B a^2 b^8 x^9 + 660 a^2 A b^8 x^8 + 1760 B a^3 b^7 x^8 + 1980 a^3 A b^7 x^7 + 3465 B$

input `int((b*x+a)^10*(B*x+A), x, method=_RETURNVERBOSE)`

output $1/12*b^{10}*B*x^{12}+(1/11*b^{10}*A+10/11*a*b^9*B)*x^{11}+(a*b^9*A+9/2*a^2*b^8*B)*x^{10}+(5*a^2*b^8*A+40/3*a^3*b^7*B)*x^9+(15*a^3*b^7*A+105/4*a^4*b^6*B)*x^8+(30*A*a^4*b^6+36*B*a^5*b^5)*x^7+(42*A*a^5*b^5+35*B*a^6*b^4)*x^6+(42*A*a^6*b^4+24*B*a^7*b^3)*x^5+(30*a^7*b^3*A+45/4*a^8*b^2*B)*x^4+(15*a^8*b^2*A+10/3*a^9*b*B)*x^3+(5*a^9*b*A+1/2*a^{10}*B)*x^2+a^{10}*A*x$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 240 vs. $2(34) = 68$.

Time = 0.09 (sec) , antiderivative size = 240, normalized size of antiderivative = 6.32

$$\begin{aligned} \int (a + bx)^{10}(A + Bx) dx = & \frac{1}{12} Bb^{10}x^{12} + Aa^{10}x + \frac{1}{11} (10 Bab^9 + Ab^{10})x^{11} \\ & + \frac{1}{2} (9 Ba^2b^8 + 2 Aab^9)x^{10} + \frac{5}{3} (8 Ba^3b^7 + 3 Aa^2b^8)x^9 \\ & + \frac{15}{4} (7 Ba^4b^6 + 4 Aa^3b^7)x^8 \\ & + 6 (6 Ba^5b^5 + 5 Aa^4b^6)x^7 + 7 (5 Ba^6b^4 + 6 Aa^5b^5)x^6 \\ & + 6 (4 Ba^7b^3 + 7 Aa^6b^4)x^5 + \frac{15}{4} (3 Ba^8b^2 + 8 Aa^7b^3)x^4 \\ & + \frac{5}{3} (2 Ba^9b + 9 Aa^8b^2)x^3 + \frac{1}{2} (Ba^{10} + 10 Aa^9b)x^2 \end{aligned}$$

input `integrate((b*x+a)^10*(B*x+A),x, algorithm="fricas")`

output `1/12*B*b^10*x^12 + A*a^10*x + 1/11*(10*B*a*b^9 + A*b^10)*x^11 + 1/2*(9*B*a^2*b^8 + 2*A*a*b^9)*x^10 + 5/3*(8*B*a^3*b^7 + 3*A*a^2*b^8)*x^9 + 15/4*(7*B*a^4*b^6 + 4*A*a^3*b^7)*x^8 + 6*(6*B*a^5*b^5 + 5*A*a^4*b^6)*x^7 + 7*(5*B*a^6*b^4 + 6*A*a^5*b^5)*x^6 + 6*(4*B*a^7*b^3 + 7*A*a^6*b^4)*x^5 + 15/4*(3*B*a^8*b^2 + 8*A*a^7*b^3)*x^4 + 5/3*(2*B*a^9*b + 9*A*a^8*b^2)*x^3 + 1/2*(B*a^10 + 10*A*a^9*b)*x^2`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 248 vs. $2(32) = 64$.

Time = 0.05 (sec) , antiderivative size = 248, normalized size of antiderivative = 6.53

$$\int (a + bx)^{10}(A + Bx) dx = Aa^{10}x + \frac{Bb^{10}x^{12}}{12} + x^{11}\left(\frac{Ab^{10}}{11} + \frac{10Bab^9}{11}\right) + x^{10}\left(Aab^9 + \frac{9Ba^2b^8}{2}\right) + x^9 \cdot \left(5Aa^2b^8 + \frac{40Ba^3b^7}{3}\right) + x^8 \cdot \left(15Aa^3b^7 + \frac{105Ba^4b^6}{4}\right) + x^7 \cdot (30Aa^4b^6 + 36Ba^5b^5) + x^6 \cdot (42Aa^5b^5 + 35Ba^6b^4) + x^5 \cdot (42Aa^6b^4 + 24Ba^7b^3) + x^4 \cdot \left(30Aa^7b^3 + \frac{45Ba^8b^2}{4}\right) + x^3 \cdot \left(15Aa^8b^2 + \frac{10Ba^9b}{3}\right) + x^2 \cdot \left(5Aa^9b + \frac{Ba^{10}}{2}\right)$$

input `integrate((b*x+a)**10*(B*x+A),x)`

output `A*a**10*x + B*b**10*x**12/12 + x**11*(A*b**10/11 + 10*B*a*b**9/11) + x**10*(A*a*b**9 + 9*B*a**2*b**8/2) + x**9*(5*A*a**2*b**8 + 40*B*a**3*b**7/3) + x**8*(15*A*a**3*b**7 + 105*B*a**4*b**6/4) + x**7*(30*A*a**4*b**6 + 36*B*a**5*b**5) + x**6*(42*A*a**5*b**5 + 35*B*a**6*b**4) + x**5*(42*A*a**6*b**4 + 24*B*a**7*b**3) + x**4*(30*A*a**7*b**3 + 45*B*a**8*b**2/4) + x**3*(15*A*a**8*b**2 + 10*B*a**9*b/3) + x**2*(5*A*a**9*b + B*a**10/2)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 240 vs. $2(34) = 68$.

Time = 0.04 (sec) , antiderivative size = 240, normalized size of antiderivative = 6.32

$$\begin{aligned} \int (a + bx)^{10}(A + Bx) dx = & \frac{1}{12} Bb^{10}x^{12} + Aa^{10}x + \frac{1}{11} (10 Bab^9 + Ab^{10})x^{11} \\ & + \frac{1}{2} (9 Ba^2b^8 + 2 Aab^9)x^{10} + \frac{5}{3} (8 Ba^3b^7 + 3 Aa^2b^8)x^9 \\ & + \frac{15}{4} (7 Ba^4b^6 + 4 Aa^3b^7)x^8 \\ & + 6 (6 Ba^5b^5 + 5 Aa^4b^6)x^7 + 7 (5 Ba^6b^4 + 6 Aa^5b^5)x^6 \\ & + 6 (4 Ba^7b^3 + 7 Aa^6b^4)x^5 + \frac{15}{4} (3 Ba^8b^2 + 8 Aa^7b^3)x^4 \\ & + \frac{5}{3} (2 Ba^9b + 9 Aa^8b^2)x^3 + \frac{1}{2} (Ba^{10} + 10 Aa^9b)x^2 \end{aligned}$$

input `integrate((b*x+a)^10*(B*x+A),x, algorithm="maxima")`

output `1/12*B*b^10*x^12 + A*a^10*x + 1/11*(10*B*a*b^9 + A*b^10)*x^11 + 1/2*(9*B*a^2*b^8 + 2*A*a*b^9)*x^10 + 5/3*(8*B*a^3*b^7 + 3*A*a^2*b^8)*x^9 + 15/4*(7*B*a^4*b^6 + 4*A*a^3*b^7)*x^8 + 6*(6*B*a^5*b^5 + 5*A*a^4*b^6)*x^7 + 7*(5*B*a^6*b^4 + 6*A*a^5*b^5)*x^6 + 6*(4*B*a^7*b^3 + 7*A*a^6*b^4)*x^5 + 15/4*(3*B*a^8*b^2 + 8*A*a^7*b^3)*x^4 + 5/3*(2*B*a^9*b + 9*A*a^8*b^2)*x^3 + 1/2*(B*a^10 + 10*A*a^9*b)*x^2`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 241 vs. 2(34) = 68.

Time = 0.13 (sec) , antiderivative size = 241, normalized size of antiderivative = 6.34

$$\begin{aligned} \int (a + bx)^{10}(A + Bx) dx = & \frac{1}{12} Bb^{10}x^{12} + \frac{10}{11} Bab^9x^{11} + \frac{1}{11} Ab^{10}x^{11} + \frac{9}{2} Ba^2b^8x^{10} \\ & + Aab^9x^{10} + \frac{40}{3} Ba^3b^7x^9 + 5 Aa^2b^8x^9 + \frac{105}{4} Ba^4b^6x^8 \\ & + 15 Aa^3b^7x^8 + 36 Ba^5b^5x^7 + 30 Aa^4b^6x^7 \\ & + 35 Ba^6b^4x^6 + 42 Aa^5b^5x^6 + 24 Ba^7b^3x^5 \\ & + 42 Aa^6b^4x^5 + \frac{45}{4} Ba^8b^2x^4 + 30 Aa^7b^3x^4 + \frac{10}{3} Ba^9bx^3 \\ & + 15 Aa^8b^2x^3 + \frac{1}{2} Ba^{10}x^2 + 5 Aa^9bx^2 + Aa^{10}x \end{aligned}$$

input `integrate((b*x+a)^10*(B*x+A),x, algorithm="giac")`

output
$$\begin{aligned} & 1/12*B*b^{10}*x^{12} + 10/11*B*a*b^9*x^{11} + 1/11*A*b^{10}*x^{11} + 9/2*B*a^2*b^8*x \\ & ^{10} + A*a*b^9*x^{10} + 40/3*B*a^3*b^7*x^9 + 5*A*a^2*b^8*x^9 + 105/4*B*a^4*b^ \\ & 6*x^8 + 15*A*a^3*b^7*x^8 + 36*B*a^5*b^5*x^7 + 30*A*a^4*b^6*x^7 + 35*B*a^6* \\ & b^4*x^6 + 42*A*a^5*b^5*x^6 + 24*B*a^7*b^3*x^5 + 42*A*a^6*b^4*x^5 + 45/4*B* \\ & a^8*b^2*x^4 + 30*A*a^7*b^3*x^4 + 10/3*B*a^9*b*x^3 + 15*A*a^8*b^2*x^3 + 1/2 \\ & *B*a^{10}*x^2 + 5*A*a^9*b*x^2 + A*a^{10}*x \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.98 (sec) , antiderivative size = 208, normalized size of antiderivative = 5.47

$$\begin{aligned} \int (a + bx)^{10}(A + Bx) dx = & x^2 \left(\frac{B a^{10}}{2} + 5 A b a^9 \right) + x^{11} \left(\frac{A b^{10}}{11} + \frac{10 B a b^9}{11} \right) \\ & + \frac{B b^{10} x^{12}}{12} + A a^{10} x + \frac{15 a^7 b^2 x^4 (8 A b + 3 B a)}{4} \\ & + 6 a^6 b^3 x^5 (7 A b + 4 B a) + 7 a^5 b^4 x^6 (6 A b + 5 B a) \\ & + 6 a^4 b^5 x^7 (5 A b + 6 B a) + \frac{15 a^3 b^6 x^8 (4 A b + 7 B a)}{4} \\ & + \frac{5 a^2 b^7 x^9 (3 A b + 8 B a)}{3} \\ & + \frac{5 a^8 b x^3 (9 A b + 2 B a)}{3} + \frac{a b^8 x^{10} (2 A b + 9 B a)}{2} \end{aligned}$$

input `int((A + B*x)*(a + b*x)^10,x)`

output
$$\begin{aligned} & x^2*((B*a^{10})/2 + 5*A*a^9*b) + x^{11}*((A*b^{10})/11 + (10*B*a*b^9)/11) + (B*b \\ & ^{10}*x^{12})/12 + A*a^{10}*x + (15*a^7*b^2*x^4*(8*A*b + 3*B*a))/4 + 6*a^6*b^3*x \\ & ^5*(7*A*b + 4*B*a) + 7*a^5*b^4*x^6*(6*A*b + 5*B*a) + 6*a^4*b^5*x^7*(5*A*b \\ & + 6*B*a) + (15*a^3*b^6*x^8*(4*A*b + 7*B*a))/4 + (5*a^2*b^7*x^9*(3*A*b + 8* \\ & B*a))/3 + (5*a^8*b*x^3*(9*A*b + 2*B*a))/3 + (a*b^8*x^{10}*(2*A*b + 9*B*a))/2 \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 120, normalized size of antiderivative = 3.16

$$\int (a + bx)^{10} (A + Bx) dx$$

$$= \frac{x(b^{11}x^{11} + 12ab^{10}x^{10} + 66a^2b^9x^9 + 220a^3b^8x^8 + 495a^4b^7x^7 + 792a^5b^6x^6 + 924a^6b^5x^5 + 792a^7b^4x^4 + 495a^8b^3x^3 + 220a^9b^2x^2 + 12a^{10}b^1x^1 + a^{11}x^0)}{12}$$

input `int((b*x+a)^10*(B*x+A),x)`output `(x*(12*a**11 + 66*a**10*b*x + 220*a**9*b**2*x**2 + 495*a**8*b**3*x**3 + 792*a**7*b**4*x**4 + 924*a**6*b**5*x**5 + 792*a**5*b**6*x**6 + 495*a**4*b**7*x**7 + 220*a**3*b**8*x**8 + 66*a**2*b**9*x**9 + 12*a*b**10*x**10 + b**11*x**11))/12`

3.79 $\int \frac{(a+bx)^{10}(A+Bx)}{d+ex} dx$

Optimal result	808
Mathematica [B] (verified)	809
Rubi [A] (verified)	810
Maple [B] (verified)	812
Fricas [B] (verification not implemented)	813
Sympy [B] (verification not implemented)	814
Maxima [B] (verification not implemented)	815
Giac [B] (verification not implemented)	816
Mupad [B] (verification not implemented)	817
Reduce [B] (verification not implemented)	817

Optimal result

Integrand size = 20, antiderivative size = 348

$$\int \frac{(a+bx)^{10}(A+Bx)}{d+ex} dx = \frac{b(bd-ae)^9(Bd-Ae)x}{e^{11}} - \frac{(bd-ae)^8(Bd-Ae)(a+bx)^2}{2e^{10}}$$

$$+ \frac{(bd-ae)^7(Bd-Ae)(a+bx)^3}{3e^9}$$

$$- \frac{(bd-ae)^6(Bd-Ae)(a+bx)^4}{4e^8}$$

$$+ \frac{(bd-ae)^5(Bd-Ae)(a+bx)^5}{5e^7}$$

$$- \frac{(bd-ae)^4(Bd-Ae)(a+bx)^6}{6e^6}$$

$$+ \frac{(bd-ae)^3(Bd-Ae)(a+bx)^7}{7e^5}$$

$$- \frac{(bd-ae)^2(Bd-Ae)(a+bx)^8}{8e^4}$$

$$+ \frac{(bd-ae)(Bd-Ae)(a+bx)^9}{9e^3} - \frac{(Bd-Ae)(a+bx)^{10}}{10e^2}$$

$$+ \frac{B(a+bx)^{11}}{11be} - \frac{(bd-ae)^{10}(Bd-Ae)\log(d+ex)}{e^{12}}$$

output

```

b*(-a*e+b*d)^9*(-A*e+B*d)*x/e^11-1/2*(-a*e+b*d)^8*(-A*e+B*d)*(b*x+a)^2/e^1
0+1/3*(-a*e+b*d)^7*(-A*e+B*d)*(b*x+a)^3/e^9-1/4*(-a*e+b*d)^6*(-A*e+B*d)*(b
*x+a)^4/e^8+1/5*(-a*e+b*d)^5*(-A*e+B*d)*(b*x+a)^5/e^7-1/6*(-a*e+b*d)^4*(-A
*e+B*d)*(b*x+a)^6/e^6+1/7*(-a*e+b*d)^3*(-A*e+B*d)*(b*x+a)^7/e^5-1/8*(-a*e+
b*d)^2*(-A*e+B*d)*(b*x+a)^8/e^4+1/9*(-a*e+b*d)*(-A*e+B*d)*(b*x+a)^9/e^3-1/
10*(-A*e+B*d)*(b*x+a)^10/e^2+1/11*B*(b*x+a)^11/b/e-(-a*e+b*d)^10*(-A*e+B*d
)*ln(e*x+d)/e^12

```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1252 vs. $2(348) = 696$.

Time = 0.77 (sec) , antiderivative size = 1252, normalized size of antiderivative = 3.60

$$\int \frac{(a + bx)^{10}(A + Bx)}{d + ex} dx = \text{Too large to display}$$

input

```
Integrate[((a + b*x)^10*(A + B*x))/(d + e*x),x]
```

output

```
(x*(27720*a^10*B*e^10 + 138600*a^9*b*e^9*(-2*B*d + 2*A*e + B*e*x) + 207900
*a^8*b^2*e^8*(3*A*e*(-2*d + e*x) + B*(6*d^2 - 3*d*e*x + 2*e^2*x^2)) + 2772
00*a^7*b^3*e^7*(2*A*e*(6*d^2 - 3*d*e*x + 2*e^2*x^2) + B*(-12*d^3 + 6*d^2*e
*x - 4*d*e^2*x^2 + 3*e^3*x^3)) + 97020*a^6*b^4*e^6*(5*A*e*(-12*d^3 + 6*d^2
*e*x - 4*d*e^2*x^2 + 3*e^3*x^3) + B*(60*d^4 - 30*d^3*e*x + 20*d^2*e^2*x^2
- 15*d*e^3*x^3 + 12*e^4*x^4)) + 116424*a^5*b^5*e^5*(A*e*(60*d^4 - 30*d^3*e
*x + 20*d^2*e^2*x^2 - 15*d*e^3*x^3 + 12*e^4*x^4) + B*(-60*d^5 + 30*d^4*e*x
- 20*d^3*e^2*x^2 + 15*d^2*e^3*x^3 - 12*d*e^4*x^4 + 10*e^5*x^5)) + 13860*a
^4*b^6*e^4*(7*A*e*(-60*d^5 + 30*d^4*e*x - 20*d^3*e^2*x^2 + 15*d^2*e^3*x^3
- 12*d*e^4*x^4 + 10*e^5*x^5) + B*(420*d^6 - 210*d^5*e*x + 140*d^4*e^2*x^2
- 105*d^3*e^3*x^3 + 84*d^2*e^4*x^4 - 70*d*e^5*x^5 + 60*e^6*x^6)) + 3960*a^
3*b^7*e^3*(2*A*e*(420*d^6 - 210*d^5*e*x + 140*d^4*e^2*x^2 - 105*d^3*e^3*x^
3 + 84*d^2*e^4*x^4 - 70*d*e^5*x^5 + 60*e^6*x^6) + B*(-840*d^7 + 420*d^6*e*x
- 280*d^5*e^2*x^2 + 210*d^4*e^3*x^3 - 168*d^3*e^4*x^4 + 140*d^2*e^5*x^5
- 120*d*e^6*x^6 + 105*e^7*x^7)) + 495*a^2*b^8*e^2*(3*A*e*(-840*d^7 + 420*d
^6*e*x - 280*d^5*e^2*x^2 + 210*d^4*e^3*x^3 - 168*d^3*e^4*x^4 + 140*d^2*e^5
*x^5 - 120*d*e^6*x^6 + 105*e^7*x^7) + B*(2520*d^8 - 1260*d^7*e*x + 840*d^6
*e^2*x^2 - 630*d^5*e^3*x^3 + 504*d^4*e^4*x^4 - 420*d^3*e^5*x^5 + 360*d^2*e
^6*x^6 - 315*d*e^7*x^7 + 280*e^8*x^8)) + 110*a*b^9*e*(A*e*(2520*d^8 - 1260
*d^7*e*x + 840*d^6*e^2*x^2 - 630*d^5*e^3*x^3 + 504*d^4*e^4*x^4 - 420*d^...
```

Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 348, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{10}(A + Bx)}{d + ex} dx$$

↓ 86

$$\int \left(\frac{(ae - bd)^{10}(Ae - Bd)}{e^{11}(d + ex)} - \frac{b(bd - ae)^9(Ae - Bd)}{e^{11}} + \frac{b(a + bx)(bd - ae)^8(Ae - Bd)}{e^{10}} - \frac{b(a + bx)^2(bd - ae)^7(Ae - Bd)}{e^9} \right) dx$$

↓ 2009

$$\begin{aligned}
& - \frac{(bd - ae)^{10}(Bd - Ae) \log(d + ex)}{e^{12}} + \frac{bx(bd - ae)^9(Bd - Ae)}{e^{11}} - \\
& \frac{(a + bx)^2(bd - ae)^8(Bd - Ae)}{2e^{10}} + \frac{(a + bx)^3(bd - ae)^7(Bd - Ae)}{3e^9} - \\
& \frac{(a + bx)^4(bd - ae)^6(Bd - Ae)}{4e^8} + \frac{(a + bx)^5(bd - ae)^5(Bd - Ae)}{5e^7} - \\
& \frac{(a + bx)^6(bd - ae)^4(Bd - Ae)}{6e^6} + \frac{(a + bx)^7(bd - ae)^3(Bd - Ae)}{7e^5} - \\
& \frac{(a + bx)^8(bd - ae)^2(Bd - Ae)}{8e^4} + \frac{(a + bx)^9(bd - ae)(Bd - Ae)}{9e^3} - \frac{(a + bx)^{10}(Bd - Ae)}{10e^2} + \\
& \frac{B(a + bx)^{11}}{11be}
\end{aligned}$$

input `Int[((a + b*x)^10*(A + B*x))/(d + e*x), x]`

output `(b*(b*d - a*e)^9*(B*d - A*e)*x)/e^11 - ((b*d - a*e)^8*(B*d - A*e)*(a + b*x)^2)/(2*e^10) + ((b*d - a*e)^7*(B*d - A*e)*(a + b*x)^3)/(3*e^9) - ((b*d - a*e)^6*(B*d - A*e)*(a + b*x)^4)/(4*e^8) + ((b*d - a*e)^5*(B*d - A*e)*(a + b*x)^5)/(5*e^7) - ((b*d - a*e)^4*(B*d - A*e)*(a + b*x)^6)/(6*e^6) + ((b*d - a*e)^3*(B*d - A*e)*(a + b*x)^7)/(7*e^5) - ((b*d - a*e)^2*(B*d - A*e)*(a + b*x)^8)/(8*e^4) + ((b*d - a*e)*(B*d - A*e)*(a + b*x)^9)/(9*e^3) - ((B*d - A*e)*(a + b*x)^10)/(10*e^2) + (B*(a + b*x)^11)/(11*b*e) - ((b*d - a*e)^10*(B*d - A*e)*Log[d + e*x])/e^12`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1883 vs. $2(328) = 656$.

Time = 0.24 (sec) , antiderivative size = 1884, normalized size of antiderivative = 5.41

method	result	size
norman	Expression too large to display	1884
default	Expression too large to display	2225
risch	Expression too large to display	2357
parallelrisc	Expression too large to display	2358

input `int((b*x+a)^10*(B*x+A)/(e*x+d),x,method=_RETURNVERBOSE)`

output

```
(10*A*a^9*b*e^10-45*A*a^8*b^2*d*e^9+120*A*a^7*b^3*d^2*e^8-210*A*a^6*b^4*d^3*e^7+252*A*a^5*b^5*d^4*e^6-210*A*a^4*b^6*d^5*e^5+120*A*a^3*b^7*d^6*e^4-45*A*a^2*b^8*d^7*e^3+10*A*a*b^9*d^8*e^2-A*b^10*d^9*e+B*a^10*e^10-10*B*a^9*b*d*e^9+45*B*a^8*b^2*d^2*e^8-120*B*a^7*b^3*d^3*e^7+210*B*a^6*b^4*d^4*e^6-252*B*a^5*b^5*d^5*e^5+210*B*a^4*b^6*d^6*e^4-120*B*a^3*b^7*d^7*e^3+45*B*a^2*b^8*d^8*e^2-10*B*a*b^9*d^9*e+B*b^10*d^10)/e^11*x+1/2*b/e^10*(45*A*a^8*b*e^9-120*A*a^7*b^2*d*e^8+210*A*a^6*b^3*d^2*e^7-252*A*a^5*b^4*d^3*e^6+210*A*a^4*b^5*d^4*e^5-120*A*a^3*b^6*d^5*e^4+45*A*a^2*b^7*d^6*e^3-10*A*a*b^8*d^7*e^2+A*b^9*d^8*e+10*B*a^9*e^9-45*B*a^8*b*d*e^8+120*B*a^7*b^2*d^2*e^7-210*B*a^6*b^3*d^3*e^6+252*B*a^5*b^4*d^4*e^5-210*B*a^4*b^5*d^5*e^4+120*B*a^3*b^6*d^6*e^3-45*B*a^2*b^7*d^7*e^2+10*B*a*b^8*d^8*e-B*b^9*d^9)*x^2+1/3*b^2/e^9*(120*A*a^7*b*e^8-210*A*a^6*b^2*d*e^7+252*A*a^5*b^3*d^2*e^6-210*A*a^4*b^4*d^3*e^5+120*A*a^3*b^5*d^4*e^4-45*A*a^2*b^6*d^5*e^3+10*A*a*b^7*d^6*e^2-A*b^8*d^7*e+45*B*a^8*e^8-120*B*a^7*b*d*e^7+210*B*a^6*b^2*d^2*e^6-252*B*a^5*b^3*d^3*e^5+210*B*a^4*b^4*d^4*e^4-120*B*a^3*b^5*d^5*e^3+45*B*a^2*b^6*d^6*e^2-10*B*a*b^7*d^7*e+B*b^8*d^8)*x^3+1/4*b^3/e^8*(210*A*a^6*b*e^7-252*A*a^5*b^2*d*e^6+210*A*a^4*b^3*d^2*e^5-120*A*a^3*b^4*d^3*e^4+45*A*a^2*b^5*d^4*e^3-10*A*a*b^6*d^5*e^2+A*b^7*d^6*e+120*B*a^7*e^7-210*B*a^6*b*d*e^6+252*B*a^5*b^2*d^2*e^5-210*B*a^4*b^3*d^3*e^4+120*B*a^3*b^4*d^4*e^3-45*B*a^2*b^5*d^5*e^2+10*B*a*b^6*d^6*e-B*b^7*d^7)*x^4+1/5*b^4/e^7*(252*A*a^5*b*e^6-210*A*a^4*b^2*d*...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1805 vs. $2(328) = 656$.

Time = 0.08 (sec) , antiderivative size = 1805, normalized size of antiderivative = 5.19

$$\int \frac{(a + bx)^{10}(A + Bx)}{d + ex} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d),x, algorithm="fricas")`

output

```
1/27720*(2520*B*b^10*e^11*x^11 - 2772*(B*b^10*d*e^10 - (10*B*a*b^9 + A*b^10)*e^11)*x^10 + 3080*(B*b^10*d^2*e^9 - (10*B*a*b^9 + A*b^10)*d*e^10 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 - 3465*(B*b^10*d^3*e^8 - (10*B*a*b^9 + A*b^10)*d^2*e^9 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 - 15*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 3960*(B*b^10*d^4*e^7 - (10*B*a*b^9 + A*b^10)*d^3*e^8 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 - 15*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 + 30*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7 - 4620*(B*b^10*d^5*e^6 - (10*B*a*b^9 + A*b^10)*d^4*e^7 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 - 15*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 30*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 - 42*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 + 5544*(B*b^10*d^6*e^5 - (10*B*a*b^9 + A*b^10)*d^5*e^6 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 - 15*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 + 30*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 - 42*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10 + 42*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 - 6930*(B*b^10*d^7*e^4 - (10*B*a*b^9 + A*b^10)*d^6*e^5 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*d^5*e^6 - 15*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^4*e^7 + 30*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^3*e^8 - 42*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e^9 + 42*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d*e^10 - 30*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e^11)*x^4 + 9240*(B*b^10*d^8*e^3 - (10*B*a*b^9 + A*b^10)*d^7*e^4 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*d^6*e^5 - 15*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^5*e^6 + 30*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^4*e^7 - 42*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^...
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1912 vs. $2(298) = 596$.

Time = 2.19 (sec) , antiderivative size = 1912, normalized size of antiderivative = 5.49

$$\int \frac{(a + bx)^{10}(A + Bx)}{d + ex} dx = \text{Too large to display}$$

input `integrate((b*x+a)**10*(B*x+A)/(e*x+d),x)`

output

```
B*b**10*x**11/(11*e) + x**10*(A*b**10/(10*e) + B*a*b**9/e - B*b**10*d/(10*
e**2)) + x**9*(10*A*a*b**9/(9*e) - A*b**10*d/(9*e**2) + 5*B*a**2*b**8/e -
10*B*a*b**9*d/(9*e**2) + B*b**10*d**2/(9*e**3)) + x**8*(45*A*a**2*b**8/(8*
e) - 5*A*a*b**9*d/(4*e**2) + A*b**10*d**2/(8*e**3) + 15*B*a**3*b**7/e - 45
*B*a**2*b**8*d/(8*e**2) + 5*B*a*b**9*d**2/(4*e**3) - B*b**10*d**3/(8*e**4)
) + x**7*(120*A*a**3*b**7/(7*e) - 45*A*a**2*b**8*d/(7*e**2) + 10*A*a*b**9*
d**2/(7*e**3) - A*b**10*d**3/(7*e**4) + 30*B*a**4*b**6/e - 120*B*a**3*b**7
*d/(7*e**2) + 45*B*a**2*b**8*d**2/(7*e**3) - 10*B*a*b**9*d**3/(7*e**4) + B
*b**10*d**4/(7*e**5)) + x**6*(35*A*a**4*b**6/e - 20*A*a**3*b**7*d/e**2 + 1
5*A*a**2*b**8*d**2/(2*e**3) - 5*A*a*b**9*d**3/(3*e**4) + A*b**10*d**4/(6*e
**5) + 42*B*a**5*b**5/e - 35*B*a**4*b**6*d/e**2 + 20*B*a**3*b**7*d**2/e**3
- 15*B*a**2*b**8*d**3/(2*e**4) + 5*B*a*b**9*d**4/(3*e**5) - B*b**10*d**5/
(6*e**6)) + x**5*(252*A*a**5*b**5/(5*e) - 42*A*a**4*b**6*d/e**2 + 24*A*a**
3*b**7*d**2/e**3 - 9*A*a**2*b**8*d**3/e**4 + 2*A*a*b**9*d**4/e**5 - A*b**1
0*d**5/(5*e**6) + 42*B*a**6*b**4/e - 252*B*a**5*b**5*d/(5*e**2) + 42*B*a**
4*b**6*d**2/e**3 - 24*B*a**3*b**7*d**3/e**4 + 9*B*a**2*b**8*d**4/e**5 - 2*
B*a*b**9*d**5/e**6 + B*b**10*d**6/(5*e**7)) + x**4*(105*A*a**6*b**4/(2*e)
- 63*A*a**5*b**5*d/e**2 + 105*A*a**4*b**6*d**2/(2*e**3) - 30*A*a**3*b**7*d
**3/e**4 + 45*A*a**2*b**8*d**4/(4*e**5) - 5*A*a*b**9*d**5/(2*e**6) + A*b**
10*d**6/(4*e**7) + 30*B*a**7*b**3/e - 105*B*a**6*b**4*d/(2*e**2) + 63*B...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1804 vs. $2(328) = 656$.

Time = 0.05 (sec) , antiderivative size = 1804, normalized size of antiderivative = 5.18

$$\int \frac{(a + bx)^{10}(A + Bx)}{d + ex} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d),x, algorithm="maxima")`

output

```
1/27720*(2520*B*b^10*e^10*x^11 - 2772*(B*b^10*d*e^9 - (10*B*a*b^9 + A*b^10)
)*e^10)*x^10 + 3080*(B*b^10*d^2*e^8 - (10*B*a*b^9 + A*b^10)*d*e^9 + 5*(9*B
*a^2*b^8 + 2*A*a*b^9)*e^10)*x^9 - 3465*(B*b^10*d^3*e^7 - (10*B*a*b^9 + A*b
^10)*d^2*e^8 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^9 - 15*(8*B*a^3*b^7 + 3*A*a
^2*b^8)*e^10)*x^8 + 3960*(B*b^10*d^4*e^6 - (10*B*a*b^9 + A*b^10)*d^3*e^7 +
5*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^8 - 15*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^
9 + 30*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^10)*x^7 - 4620*(B*b^10*d^5*e^5 - (10*
B*a*b^9 + A*b^10)*d^4*e^6 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^7 - 15*(8*B*
a^3*b^7 + 3*A*a^2*b^8)*d^2*e^8 + 30*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^9 - 42
*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^10)*x^6 + 5544*(B*b^10*d^6*e^4 - (10*B*a*b^
9 + A*b^10)*d^5*e^5 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^6 - 15*(8*B*a^3*b^
7 + 3*A*a^2*b^8)*d^3*e^7 + 30*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^8 - 42*(6*
B*a^5*b^5 + 5*A*a^4*b^6)*d*e^9 + 42*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^10)*x^5
- 6930*(B*b^10*d^7*e^3 - (10*B*a*b^9 + A*b^10)*d^6*e^4 + 5*(9*B*a^2*b^8 +
2*A*a*b^9)*d^5*e^5 - 15*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^4*e^6 + 30*(7*B*a^4*
b^6 + 4*A*a^3*b^7)*d^3*e^7 - 42*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e^8 + 42*(
5*B*a^6*b^4 + 6*A*a^5*b^5)*d*e^9 - 30*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e^10)*x^
4 + 9240*(B*b^10*d^8*e^2 - (10*B*a*b^9 + A*b^10)*d^7*e^3 + 5*(9*B*a^2*b^8
+ 2*A*a*b^9)*d^6*e^4 - 15*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^5*e^5 + 30*(7*B*a^
4*b^6 + 4*A*a^3*b^7)*d^4*e^6 - 42*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^3*e^7 + ...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2230 vs. $2(328) = 656$.

Time = 0.12 (sec) , antiderivative size = 2230, normalized size of antiderivative = 6.41

$$\int \frac{(a + bx)^{10}(A + Bx)}{d + ex} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d),x, algorithm="giac")`

output

```
1/27720*(2520*B*b^10*e^10*x^11 - 2772*B*b^10*d*e^9*x^10 + 27720*B*a*b^9*e^
10*x^10 + 2772*A*b^10*e^10*x^10 + 3080*B*b^10*d^2*e^8*x^9 - 30800*B*a*b^9*
d*e^9*x^9 - 3080*A*b^10*d*e^9*x^9 + 138600*B*a^2*b^8*e^10*x^9 + 30800*A*a*
b^9*e^10*x^9 - 3465*B*b^10*d^3*e^7*x^8 + 34650*B*a*b^9*d^2*e^8*x^8 + 3465*
A*b^10*d^2*e^8*x^8 - 155925*B*a^2*b^8*d*e^9*x^8 - 34650*A*a*b^9*d*e^9*x^8
+ 415800*B*a^3*b^7*e^10*x^8 + 155925*A*a^2*b^8*e^10*x^8 + 3960*B*b^10*d^4*
e^6*x^7 - 39600*B*a*b^9*d^3*e^7*x^7 - 3960*A*b^10*d^3*e^7*x^7 + 178200*B*a
^2*b^8*d^2*e^8*x^7 + 39600*A*a*b^9*d^2*e^8*x^7 - 475200*B*a^3*b^7*d*e^9*x^
7 - 178200*A*a^2*b^8*d*e^9*x^7 + 831600*B*a^4*b^6*e^10*x^7 + 475200*A*a^3*
b^7*e^10*x^7 - 4620*B*b^10*d^5*e^5*x^6 + 46200*B*a*b^9*d^4*e^6*x^6 + 4620*
A*b^10*d^4*e^6*x^6 - 207900*B*a^2*b^8*d^3*e^7*x^6 - 46200*A*a*b^9*d^3*e^7*
x^6 + 554400*B*a^3*b^7*d^2*e^8*x^6 + 207900*A*a^2*b^8*d^2*e^8*x^6 - 970200
*B*a^4*b^6*d*e^9*x^6 - 554400*A*a^3*b^7*d*e^9*x^6 + 1164240*B*a^5*b^5*e^10
*x^6 + 970200*A*a^4*b^6*e^10*x^6 + 5544*B*b^10*d^6*e^4*x^5 - 55440*B*a*b^9
*d^5*e^5*x^5 - 5544*A*b^10*d^5*e^5*x^5 + 249480*B*a^2*b^8*d^4*e^6*x^5 + 55
440*A*a*b^9*d^4*e^6*x^5 - 665280*B*a^3*b^7*d^3*e^7*x^5 - 249480*A*a^2*b^8*
d^3*e^7*x^5 + 1164240*B*a^4*b^6*d^2*e^8*x^5 + 665280*A*a^3*b^7*d^2*e^8*x^5
- 1397088*B*a^5*b^5*d*e^9*x^5 - 1164240*A*a^4*b^6*d*e^9*x^5 + 1164240*B*a
^6*b^4*e^10*x^5 + 1397088*A*a^5*b^5*e^10*x^5 - 6930*B*b^10*d^7*e^3*x^4 + 6
9300*B*a*b^9*d^6*e^4*x^4 + 6930*A*b^10*d^6*e^4*x^4 - 311850*B*a^2*b^8*d...
```

Mupad [B] (verification not implemented)

Time = 1.02 (sec) , antiderivative size = 1795, normalized size of antiderivative = 5.16

$$\int \frac{(a + bx)^{10}(A + Bx)}{d + ex} dx = \text{Too large to display}$$

input `int(((A + B*x)*(a + b*x)^10)/(d + e*x),x)`

output `x^4*((d*((d*((d*((d*((d*((d*((A*b^10 + 10*B*a*b^9)/e - (B*b^10*d)/e^2))/e - (5*a*b^8*(2*A*b + 9*B*a))/e))/e + (15*a^2*b^7*(3*A*b + 8*B*a))/e))/e - (30*a^3*b^6*(4*A*b + 7*B*a))/e))/e + (42*a^4*b^5*(5*A*b + 6*B*a))/e))/e - (42*a^5*b^4*(6*A*b + 5*B*a))/e)/(4*e) + (15*a^6*b^3*(7*A*b + 4*B*a))/(2*e) - x^3*((d*((d*((d*((d*((d*((d*((A*b^10 + 10*B*a*b^9)/e - (B*b^10*d)/e^2))/e - (5*a*b^8*(2*A*b + 9*B*a))/e))/e + (15*a^2*b^7*(3*A*b + 8*B*a))/e))/e - (30*a^3*b^6*(4*A*b + 7*B*a))/e))/e + (42*a^4*b^5*(5*A*b + 6*B*a))/e))/e - (42*a^5*b^4*(6*A*b + 5*B*a))/e)/e + (30*a^6*b^3*(7*A*b + 4*B*a))/e)/(3*e) - (5*a^7*b^2*(8*A*b + 3*B*a))/e - x^5*((d*((d*((d*((d*((d*((A*b^10 + 10*B*a*b^9)/e - (B*b^10*d)/e^2))/e - (5*a*b^8*(2*A*b + 9*B*a))/e))/e + (15*a^2*b^7*(3*A*b + 8*B*a))/e))/e - (30*a^3*b^6*(4*A*b + 7*B*a))/e))/e + (42*a^4*b^5*(5*A*b + 6*B*a))/e)/(5*e) - (42*a^5*b^4*(6*A*b + 5*B*a))/(5*e) + x^6*((d*((d*((d*((d*((d*((d*((A*b^10 + 10*B*a*b^9)/e - (B*b^10*d)/e^2))/e - (5*a*b^8*(2*A*b + 9*B*a))/e))/e + (15*a^2*b^7*(3*A*b + 8*B*a))/e))/e - (30*a^3*b^6*(4*A*b + 7*B*a))/e))/(6*e) + (7*a^4*b^5*(5*A*b + 6*B*a))/e - x^7*((d*((d*((d*((d*((d*((d*((A*b^10 + 10*B*a*b^9)/e - (B*b^10*d)/e^2))/e - (5*a*b^8*(2*A*b + 9*B*a))/e))/e + (15*a^2*b^7*(3*A*b + 8*B*a))/e))/e - (30*a^3*b^6*(4*A*b + 7*B*a))/e))/(7*e) + (7*a^4*b^5*(5*A*b + 6*B*a))/e - x^8*((d*((d*((d*((d*((d*((d*((A*b^10 + 10*B*a*b^9)/e - (B*b^10*d)/e^2))/e - (5*a*b^8*(2*A*b + 9*B*a))/e))/e + (15*a^2*b^7*(3*A*b + 8*B*a))/e))/e - (30*a^3*b^6*(4*A*b + 7*B*a))/e))/(8*e) + (15*a^2*b^7*(3*A*b + 8*B*a))/e - x^9*((d*((d*((d*((d*((d*((d*((A*b^10 + 10*B*a*b^9)/e - (B*b^10*d)/e^2))/e - (5*a*b^8*(2*A*b + 9*B*a))/e))/e + (15*a^2*b^7*(3*A*b + 8*B*a))/e))/e - (30*a^3*b^6*(4*A*b + 7*B*a))/e))/(9*e)...`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 1216, normalized size of antiderivative = 3.49

$$\int \frac{(a + bx)^{10}(A + Bx)}{d + ex} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)/(e*x+d),x)`

output

```
(27720*log(d + e*x)*a**11*e**11 - 304920*log(d + e*x)*a**10*b*d*e**10 + 15
24600*log(d + e*x)*a**9*b**2*d**2*e**9 - 4573800*log(d + e*x)*a**8*b**3*d*
*3*e**8 + 9147600*log(d + e*x)*a**7*b**4*d**4*e**7 - 12806640*log(d + e*x)
*a**6*b**5*d**5*e**6 + 12806640*log(d + e*x)*a**5*b**6*d**6*e**5 - 9147600
*log(d + e*x)*a**4*b**7*d**7*e**4 + 4573800*log(d + e*x)*a**3*b**8*d**8*e*
*3 - 1524600*log(d + e*x)*a**2*b**9*d**9*e**2 + 304920*log(d + e*x)*a*b**1
0*d**10*e - 27720*log(d + e*x)*b**11*d**11 + 304920*a**10*b*e**11*x - 1524
600*a**9*b**2*d*e**10*x + 762300*a**9*b**2*e**11*x**2 + 4573800*a**8*b**3*
d**2*e**9*x - 2286900*a**8*b**3*d*e**10*x**2 + 1524600*a**8*b**3*e**11*x**
3 - 9147600*a**7*b**4*d**3*e**8*x + 4573800*a**7*b**4*d**2*e**9*x**2 - 304
9200*a**7*b**4*d*e**10*x**3 + 2286900*a**7*b**4*e**11*x**4 + 12806640*a**6
*b**5*d**4*e**7*x - 6403320*a**6*b**5*d**3*e**8*x**2 + 4268880*a**6*b**5*d
**2*e**9*x**3 - 3201660*a**6*b**5*d*e**10*x**4 + 2561328*a**6*b**5*e**11*x
**5 - 12806640*a**5*b**6*d**5*e**6*x + 6403320*a**5*b**6*d**4*e**7*x**2 -
4268880*a**5*b**6*d**3*e**8*x**3 + 3201660*a**5*b**6*d**2*e**9*x**4 - 2561
328*a**5*b**6*d*e**10*x**5 + 2134440*a**5*b**6*e**11*x**6 + 9147600*a**4*b
**7*d**6*e**5*x - 4573800*a**4*b**7*d**5*e**6*x**2 + 3049200*a**4*b**7*d**
4*e**7*x**3 - 2286900*a**4*b**7*d**3*e**8*x**4 + 1829520*a**4*b**7*d**2*e
**9*x**5 - 1524600*a**4*b**7*d*e**10*x**6 + 1306800*a**4*b**7*e**11*x**7 -
4573800*a**3*b**8*d**7*e**4*x + 2286900*a**3*b**8*d**6*e**5*x**2 - 1524...
```

3.80 $\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^2} dx$

Optimal result	819
Mathematica [B] (verified)	820
Rubi [A] (verified)	821
Maple [B] (verified)	823
Fricas [B] (verification not implemented)	824
Sympy [B] (verification not implemented)	825
Maxima [B] (verification not implemented)	826
Giac [B] (verification not implemented)	827
Mupad [B] (verification not implemented)	828
Reduce [B] (verification not implemented)	829

Optimal result

Integrand size = 20, antiderivative size = 445

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^2} dx = -\frac{5b(bd-ae)^8(11bBd-9Abe-2aBe)x}{e^{11}} + \frac{(bd-ae)^{10}(Bd-Ae)}{e^{12}(d+ex)} + \frac{15b^2(bd-ae)^7(11bBd-8Abe-3aBe)(d+ex)^2}{2e^{12}} - \frac{10b^3(bd-ae)^6(11bBd-7Abe-4aBe)(d+ex)^3}{e^{12}} + \frac{21b^4(bd-ae)^5(11bBd-6Abe-5aBe)(d+ex)^4}{2e^{12}} - \frac{42b^5(bd-ae)^4(11bBd-5Abe-6aBe)(d+ex)^5}{5e^{12}} + \frac{5b^6(bd-ae)^3(11bBd-4Abe-7aBe)(d+ex)^6}{e^{12}} - \frac{15b^7(bd-ae)^2(11bBd-3Abe-8aBe)(d+ex)^7}{7e^{12}} + \frac{5b^8(bd-ae)(11bBd-2Abe-9aBe)(d+ex)^8}{8e^{12}} - \frac{b^9(11bBd-Abe-10aBe)(d+ex)^9}{9e^{12}} + \frac{b^{10}B(d+ex)^{10}}{10e^{12}} + \frac{(bd-ae)^9(11bBd-10Abe-aBe)\log(d+ex)}{e^{12}}$$

output

```
-5*b*(-a*e+b*d)^8*(-9*A*b*e-2*B*a*e+11*B*b*d)*x/e^11+(-a*e+b*d)^10*(-A*e+B
*d)/e^12/(e*x+d)+15/2*b^2*(-a*e+b*d)^7*(-8*A*b*e-3*B*a*e+11*B*b*d)*(e*x+d)
^2/e^12-10*b^3*(-a*e+b*d)^6*(-7*A*b*e-4*B*a*e+11*B*b*d)*(e*x+d)^3/e^12+21/
2*b^4*(-a*e+b*d)^5*(-6*A*b*e-5*B*a*e+11*B*b*d)*(e*x+d)^4/e^12-42/5*b^5*(-a
*e+b*d)^4*(-5*A*b*e-6*B*a*e+11*B*b*d)*(e*x+d)^5/e^12+5*b^6*(-a*e+b*d)^3*(-
4*A*b*e-7*B*a*e+11*B*b*d)*(e*x+d)^6/e^12-15/7*b^7*(-a*e+b*d)^2*(-3*A*b*e-8
*B*a*e+11*B*b*d)*(e*x+d)^7/e^12+5/8*b^8*(-a*e+b*d)*(-2*A*b*e-9*B*a*e+11*B*
b*d)*(e*x+d)^8/e^12-1/9*b^9*(-A*b*e-10*B*a*e+11*B*b*d)*(e*x+d)^9/e^12+1/10
*b^10*B*(e*x+d)^10/e^12+(-a*e+b*d)^9*(-10*A*b*e-B*a*e+11*B*b*d)*ln(e*x+d)/
e^12
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1486 vs. $2(445) = 890$.

Time = 0.42 (sec) , antiderivative size = 1486, normalized size of antiderivative = 3.34

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^2} dx = \text{Too large to display}$$

input

```
Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^2,x]
```

output

```
(-2520*a^10*e^10*(-(B*d) + A*e) + 25200*a^9*b*e^9*(A*d*e + B*(-d^2 + d*e*x
+ e^2*x^2)) + 56700*a^8*b^2*e^8*(2*A*e*(-d^2 + d*e*x + e^2*x^2) + B*(2*d^
3 - 4*d^2*e*x - 3*d*e^2*x^2 + e^3*x^3)) + 50400*a^7*b^3*e^7*(3*A*e*(2*d^3
- 4*d^2*e*x - 3*d*e^2*x^2 + e^3*x^3) + 2*B*(-3*d^4 + 9*d^3*e*x + 6*d^2*e^2
*x^2 - 2*d*e^3*x^3 + e^4*x^4)) + 44100*a^6*b^4*e^6*(4*A*e*(-3*d^4 + 9*d^3*
e*x + 6*d^2*e^2*x^2 - 2*d*e^3*x^3 + e^4*x^4) + B*(12*d^5 - 48*d^4*e*x - 30
*d^3*e^2*x^2 + 10*d^2*e^3*x^3 - 5*d*e^4*x^4 + 3*e^5*x^5)) + 10584*a^5*b^5*
e^5*(5*A*e*(12*d^5 - 48*d^4*e*x - 30*d^3*e^2*x^2 + 10*d^2*e^3*x^3 - 5*d*e^
4*x^4 + 3*e^5*x^5) - 6*B*(10*d^6 - 50*d^5*e*x - 30*d^4*e^2*x^2 + 10*d^3*e^
3*x^3 - 5*d^2*e^4*x^4 + 3*d*e^5*x^5 - 2*e^6*x^6)) + 8820*a^4*b^6*e^4*(6*A*
e*(-10*d^6 + 50*d^5*e*x + 30*d^4*e^2*x^2 - 10*d^3*e^3*x^3 + 5*d^2*e^4*x^4
- 3*d*e^5*x^5 + 2*e^6*x^6) + B*(60*d^7 - 360*d^6*e*x - 210*d^5*e^2*x^2 + 7
0*d^4*e^3*x^3 - 35*d^3*e^4*x^4 + 21*d^2*e^5*x^5 - 14*d*e^6*x^6 + 10*e^7*x^
7)) + 720*a^3*b^7*e^3*(7*A*e*(60*d^7 - 360*d^6*e*x - 210*d^5*e^2*x^2 + 70*
d^4*e^3*x^3 - 35*d^3*e^4*x^4 + 21*d^2*e^5*x^5 - 14*d*e^6*x^6 + 10*e^7*x^7)
- 4*B*(105*d^8 - 735*d^7*e*x - 420*d^6*e^2*x^2 + 140*d^5*e^3*x^3 - 70*d^4
*e^4*x^4 + 42*d^3*e^5*x^5 - 28*d^2*e^6*x^6 + 20*d*e^7*x^7 - 15*e^8*x^8)) +
135*a^2*b^8*e^2*(8*A*e*(-105*d^8 + 735*d^7*e*x + 420*d^6*e^2*x^2 - 140*d^
5*e^3*x^3 + 70*d^4*e^4*x^4 - 42*d^3*e^5*x^5 + 28*d^2*e^6*x^6 - 20*d*e^7*x^
7 + 15*e^8*x^8) + 3*B*(280*d^9 - 2240*d^8*e*x - 1260*d^7*e^2*x^2 + 420*...
```

Rubi [A] (verified)

Time = 1.71 (sec) , antiderivative size = 445, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^2} dx$$

↓ 86

$$\int \left(\frac{b^9(d + ex)^8(10aBe + Abe - 11bBd)}{e^{11}} - \frac{5b^8(d + ex)^7(bd - ae)(9aBe + 2Abe - 11bBd)}{e^{11}} + \frac{15b^7(d + ex)^6(bd - ae)^2}{e^{11}} \right) dx$$

↓ 2009

$$\begin{aligned}
& - \frac{b^9(d+ex)^9(-10aBe - Abe + 11bBd)}{9e^{12}} + \frac{5b^8(d+ex)^8(bd-ae)(-9aBe - 2Abe + 11bBd)}{8e^{12}} - \\
& \frac{15b^7(d+ex)^7(bd-ae)^2(-8aBe - 3Abe + 11bBd)}{7e^{12}} + \\
& \frac{5b^6(d+ex)^6(bd-ae)^3(-7aBe - 4Abe + 11bBd)}{6e^{12}} - \\
& \frac{42b^5(d+ex)^5(bd-ae)^4(-6aBe - 5Abe + 11bBd)}{5e^{12}} + \\
& \frac{21b^4(d+ex)^4(bd-ae)^5(-5aBe - 6Abe + 11bBd)}{4e^{12}} - \\
& \frac{10b^3(d+ex)^3(bd-ae)^6(-4aBe - 7Abe + 11bBd)}{3e^{12}} + \\
& \frac{15b^2(d+ex)^2(bd-ae)^7(-3aBe - 8Abe + 11bBd)}{2e^{12}} + \frac{(bd-ae)^{10}(Bd-Ae)}{e^{12}(d+ex)} + \\
& \frac{(bd-ae)^9 \log(d+ex)(-aBe - 10Abe + 11bBd)}{e^{12}} - \frac{5bx(bd-ae)^8(-2aBe - 9Abe + 11bBd)}{e^{11}} + \\
& \frac{b^{10}B(d+ex)^{10}}{10e^{12}}
\end{aligned}$$

input `Int[((a + b*x)^10*(A + B*x))/(d + e*x)^2,x]`

output `(-5*b*(b*d - a*e)^8*(11*b*B*d - 9*A*b*e - 2*a*B*e)*x)/e^11 + ((b*d - a*e)^10*(B*d - A*e))/(e^12*(d + e*x)) + (15*b^2*(b*d - a*e)^7*(11*b*B*d - 8*A*b*e - 3*a*B*e)*(d + e*x)^2)/(2*e^12) - (10*b^3*(b*d - a*e)^6*(11*b*B*d - 7*A*b*e - 4*a*B*e)*(d + e*x)^3)/e^12 + (21*b^4*(b*d - a*e)^5*(11*b*B*d - 6*A*b*e - 5*a*B*e)*(d + e*x)^4)/(2*e^12) - (42*b^5*(b*d - a*e)^4*(11*b*B*d - 5*A*b*e - 6*a*B*e)*(d + e*x)^5)/(5*e^12) + (5*b^6*(b*d - a*e)^3*(11*b*B*d - 4*A*b*e - 7*a*B*e)*(d + e*x)^6)/e^12 - (15*b^7*(b*d - a*e)^2*(11*b*B*d - 3*A*b*e - 8*a*B*e)*(d + e*x)^7)/(7*e^12) + (5*b^8*(b*d - a*e)*(11*b*B*d - 2*A*b*e - 9*a*B*e)*(d + e*x)^8)/(8*e^12) - (b^9*(11*b*B*d - A*b*e - 10*a*B*e)*(d + e*x)^9)/(9*e^12) + (b^10*B*(d + e*x)^10)/(10*e^12) + ((b*d - a*e)^9*(11*b*B*d - 10*A*b*e - a*B*e)*Log[d + e*x])/e^12`

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1906 vs. $2(431) = 862$.

Time = 0.24 (sec) , antiderivative size = 1907, normalized size of antiderivative = 4.29

method	result	size
norman	Expression too large to display	1907
default	Expression too large to display	2167
risch	Expression too large to display	2447
parallelrisc	Expression too large to display	2772

input

```
int((b*x+a)^10*(B*x+A)/(e*x+d)^2,x,method=_RETURNVERBOSE)
```

output

```
((A*a^10*e^11-10*A*a^9*b*d*e^10+90*A*a^8*b^2*d^2*e^9-360*A*a^7*b^3*d^3*e^8
+840*A*a^6*b^4*d^4*e^7-1260*A*a^5*b^5*d^5*e^6+1260*A*a^4*b^6*d^6*e^5-840*A
*a^3*b^7*d^7*e^4+360*A*a^2*b^8*d^8*e^3-90*A*a*b^9*d^9*e^2+10*A*b^10*d^10*e
-B*a^10*d*e^10+20*B*a^9*b*d^2*e^9-135*B*a^8*b^2*d^3*e^8+480*B*a^7*b^3*d^4*
e^7-1050*B*a^6*b^4*d^5*e^6+1512*B*a^5*b^5*d^6*e^5-1470*B*a^4*b^6*d^7*e^4+9
60*B*a^3*b^7*d^8*e^3-405*B*a^2*b^8*d^9*e^2+100*B*a*b^9*d^10*e-11*B*b^10*d^
11)/e^11/d*x+1/2*b*(90*A*a^8*b*e^9-360*A*a^7*b^2*d*e^8+840*A*a^6*b^3*d^2*e
^7-1260*A*a^5*b^4*d^3*e^6+1260*A*a^4*b^5*d^4*e^5-840*A*a^3*b^6*d^5*e^4+360
*A*a^2*b^7*d^6*e^3-90*A*a*b^8*d^7*e^2+10*A*b^9*d^8*e+20*B*a^9*e^9-135*B*a^
8*b*d*e^8+480*B*a^7*b^2*d^2*e^7-1050*B*a^6*b^3*d^3*e^6+1512*B*a^5*b^4*d^4*
e^5-1470*B*a^4*b^5*d^5*e^4+960*B*a^3*b^6*d^6*e^3-405*B*a^2*b^7*d^7*e^2+100
*B*a*b^8*d^8*e-11*B*b^9*d^9)/e^10*x^2+1/6*b^2*(360*A*a^7*b*e^8-840*A*a^6*b
^2*d*e^7+1260*A*a^5*b^3*d^2*e^6-1260*A*a^4*b^4*d^3*e^5+840*A*a^3*b^5*d^4*e
^4-360*A*a^2*b^6*d^5*e^3+90*A*a*b^7*d^6*e^2-10*A*b^8*d^7*e+135*B*a^8*e^8-4
80*B*a^7*b*d*e^7+1050*B*a^6*b^2*d^2*e^6-1512*B*a^5*b^3*d^3*e^5+1470*B*a^4*
b^4*d^4*e^4-960*B*a^3*b^5*d^5*e^3+405*B*a^2*b^6*d^6*e^2-100*B*a*b^7*d^7*e+
11*B*b^8*d^8)/e^9*x^3+1/12*b^3*(840*A*a^6*b*e^7-1260*A*a^5*b^2*d*e^6+1260*
A*a^4*b^3*d^2*e^5-840*A*a^3*b^4*d^3*e^4+360*A*a^2*b^5*d^4*e^3-90*A*a*b^6*d
^5*e^2+10*A*b^7*d^6*e+480*B*a^7*e^7-1050*B*a^6*b*d*e^6+1512*B*a^5*b^2*d^2*
e^5-1470*B*a^4*b^3*d^3*e^4+960*B*a^3*b^4*d^4*e^3-405*B*a^2*b^5*d^5*e^2+...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2329 vs. $2(431) = 862$.

Time = 0.12 (sec) , antiderivative size = 2329, normalized size of antiderivative = 5.23

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^2} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)/(e*x+d)^2,x, algorithm="fricas")
```

output

```

1/2520*(252*B*b^10*e^11*x^11 + 2520*B*b^10*d^11 - 2520*A*a^10*e^11 - 2520*
(10*B*a*b^9 + A*b^10)*d^10*e + 12600*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 - 3
7800*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 75600*(7*B*a^4*b^6 + 4*A*a^3*b^
7)*d^7*e^4 - 105840*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 105840*(5*B*a^6*
b^4 + 6*A*a^5*b^5)*d^5*e^6 - 75600*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 3
7800*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^3*e^8 - 12600*(2*B*a^9*b + 9*A*a^8*b^2)
*d^2*e^9 + 2520*(B*a^10 + 10*A*a^9*b)*d*e^10 - 28*(11*B*b^10*d*e^10 - 10*(
10*B*a*b^9 + A*b^10)*e^11)*x^10 + 35*(11*B*b^10*d^2*e^9 - 10*(10*B*a*b^9 +
A*b^10)*d*e^10 + 45*(9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 - 45*(11*B*b^10*d
^3*e^8 - 10*(10*B*a*b^9 + A*b^10)*d^2*e^9 + 45*(9*B*a^2*b^8 + 2*A*a*b^9)*d
*e^10 - 120*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 60*(11*B*b^10*d^4*e^7
- 10*(10*B*a*b^9 + A*b^10)*d^3*e^8 + 45*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9
- 120*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 + 210*(7*B*a^4*b^6 + 4*A*a^3*b^7)
*e^11)*x^7 - 84*(11*B*b^10*d^5*e^6 - 10*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 45
*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 - 120*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e
^9 + 210*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 - 252*(6*B*a^5*b^5 + 5*A*a^4*b
^6)*e^11)*x^6 + 126*(11*B*b^10*d^6*e^5 - 10*(10*B*a*b^9 + A*b^10)*d^5*e^6
+ 45*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 - 120*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d
^3*e^8 + 210*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 - 252*(6*B*a^5*b^5 + 5*A*
a^4*b^6)*d*e^10 + 210*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 - 210*(11*B...

```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1974 vs. $2(464) = 928$.

Time = 5.01 (sec) , antiderivative size = 1974, normalized size of antiderivative = 4.44

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^2} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)**10*(B*x+A)/(e*x+d)**2,x)
```

output

```

B*b**10*x**10/(10*e**2) + x**9*(A*b**10/(9*e**2) + 10*B*a*b**9/(9*e**2) -
2*B*b**10*d/(9*e**3) + x**8*(5*A*a*b**9/(4*e**2) - A*b**10*d/(4*e**3) + 4
5*B*a**2*b**8/(8*e**2) - 5*B*a*b**9*d/(2*e**3) + 3*B*b**10*d**2/(8*e**4))
+ x**7*(45*A*a**2*b**8/(7*e**2) - 20*A*a*b**9*d/(7*e**3) + 3*A*b**10*d**2/
(7*e**4) + 120*B*a**3*b**7/(7*e**2) - 90*B*a**2*b**8*d/(7*e**3) + 30*B*a*b
**9*d**2/(7*e**4) - 4*B*b**10*d**3/(7*e**5)) + x**6*(20*A*a**3*b**7/e**2 -
15*A*a**2*b**8*d/e**3 + 5*A*a*b**9*d**2/e**4 - 2*A*b**10*d**3/(3*e**5) +
35*B*a**4*b**6/e**2 - 40*B*a**3*b**7*d/e**3 + 45*B*a**2*b**8*d**2/(2*e**4)
- 20*B*a*b**9*d**3/(3*e**5) + 5*B*b**10*d**4/(6*e**6)) + x**5*(42*A*a**4*
b**6/e**2 - 48*A*a**3*b**7*d/e**3 + 27*A*a**2*b**8*d**2/e**4 - 8*A*a*b**9*
d**3/e**5 + A*b**10*d**4/e**6 + 252*B*a**5*b**5/(5*e**2) - 84*B*a**4*b**6*
d/e**3 + 72*B*a**3*b**7*d**2/e**4 - 36*B*a**2*b**8*d**3/e**5 + 10*B*a*b**9
*d**4/e**6 - 6*B*b**10*d**5/(5*e**7)) + x**4*(63*A*a**5*b**5/e**2 - 105*A*
a**4*b**6*d/e**3 + 90*A*a**3*b**7*d**2/e**4 - 45*A*a**2*b**8*d**3/e**5 + 2
5*A*a*b**9*d**4/(2*e**6) - 3*A*b**10*d**5/(2*e**7) + 105*B*a**6*b**4/(2*e
**2) - 126*B*a**5*b**5*d/e**3 + 315*B*a**4*b**6*d**2/(2*e**4) - 120*B*a**3*
b**7*d**3/e**5 + 225*B*a**2*b**8*d**4/(4*e**6) - 15*B*a*b**9*d**5/e**7 + 7
*B*b**10*d**6/(4*e**8)) + x**3*(70*A*a**6*b**4/e**2 - 168*A*a**5*b**5*d/e
**3 + 210*A*a**4*b**6*d**2/e**4 - 160*A*a**3*b**7*d**3/e**5 + 75*A*a**2*b**
8*d**4/e**6 - 20*A*a*b**9*d**5/e**7 + 7*A*b**10*d**6/(3*e**8) + 40*B*a*...

```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1817 vs. $2(431) = 862$.

Time = 0.05 (sec) , antiderivative size = 1817, normalized size of antiderivative = 4.08

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^2} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)/(e*x+d)^2,x, algorithm="maxima")
```

output

```
(B*b^10*d^11 - A*a^10*e^11 - (10*B*a*b^9 + A*b^10)*d^10*e + 5*(9*B*a^2*b^8
+ 2*A*a*b^9)*d^9*e^2 - 15*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 30*(7*B*a
^4*b^6 + 4*A*a^3*b^7)*d^7*e^4 - 42*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 4
2*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^5*e^6 - 30*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4
*e^7 + 15*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^3*e^8 - 5*(2*B*a^9*b + 9*A*a^8*b^2
)*d^2*e^9 + (B*a^10 + 10*A*a^9*b)*d*e^10)/(e^13*x + d*e^12) + 1/2520*(252*
B*b^10*e^9*x^10 - 280*(2*B*b^10*d*e^8 - (10*B*a*b^9 + A*b^10)*e^9)*x^9 + 3
15*(3*B*b^10*d^2*e^7 - 2*(10*B*a*b^9 + A*b^10)*d*e^8 + 5*(9*B*a^2*b^8 + 2*
A*a*b^9)*e^9)*x^8 - 360*(4*B*b^10*d^3*e^6 - 3*(10*B*a*b^9 + A*b^10)*d^2*e^
7 + 10*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^8 - 15*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^
9)*x^7 + 420*(5*B*b^10*d^4*e^5 - 4*(10*B*a*b^9 + A*b^10)*d^3*e^6 + 15*(9*B
*a^2*b^8 + 2*A*a*b^9)*d^2*e^7 - 30*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^8 + 30*
(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^9)*x^6 - 504*(6*B*b^10*d^5*e^4 - 5*(10*B*a*b
^9 + A*b^10)*d^4*e^5 + 20*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^6 - 45*(8*B*a^3*
b^7 + 3*A*a^2*b^8)*d^2*e^7 + 60*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^8 - 42*(6*
B*a^5*b^5 + 5*A*a^4*b^6)*e^9)*x^5 + 630*(7*B*b^10*d^6*e^3 - 6*(10*B*a*b^9
+ A*b^10)*d^5*e^4 + 25*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^5 - 60*(8*B*a^3*b^7
+ 3*A*a^2*b^8)*d^3*e^6 + 90*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^7 - 84*(6*B
*a^5*b^5 + 5*A*a^4*b^6)*d*e^8 + 42*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^9)*x^4 -
840*(8*B*b^10*d^7*e^2 - 7*(10*B*a*b^9 + A*b^10)*d^6*e^3 + 30*(9*B*a^2*b...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2207 vs. $2(431) = 862$.

Time = 0.15 (sec) , antiderivative size = 2207, normalized size of antiderivative = 4.96

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^2} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)/(e*x+d)^2,x, algorithm="giac")
```


output

```

1/2520*(252*B*b^10 - 280*(11*B*b^10*d*e - 10*B*a*b^9*e^2 - A*b^10*e^2)/((e
*x + d)*e) + 1575*(11*B*b^10*d^2*e^2 - 20*B*a*b^9*d*e^3 - 2*A*b^10*d*e^3 +
  9*B*a^2*b^8*e^4 + 2*A*a*b^9*e^4)/((e*x + d)^2*e^2) - 5400*(11*B*b^10*d^3*
e^3 - 30*B*a*b^9*d^2*e^4 - 3*A*b^10*d^2*e^4 + 27*B*a^2*b^8*d*e^5 + 6*A*a*b
^9*d*e^5 - 8*B*a^3*b^7*e^6 - 3*A*a^2*b^8*e^6)/((e*x + d)^3*e^3) + 12600*(1
1*B*b^10*d^4*e^4 - 40*B*a*b^9*d^3*e^5 - 4*A*b^10*d^3*e^5 + 54*B*a^2*b^8*d^
2*e^6 + 12*A*a*b^9*d^2*e^6 - 32*B*a^3*b^7*d*e^7 - 12*A*a^2*b^8*d*e^7 + 7*B
*a^4*b^6*e^8 + 4*A*a^3*b^7*e^8)/((e*x + d)^4*e^4) - 21168*(11*B*b^10*d^5*e
^5 - 50*B*a*b^9*d^4*e^6 - 5*A*b^10*d^4*e^6 + 90*B*a^2*b^8*d^3*e^7 + 20*A*a
*b^9*d^3*e^7 - 80*B*a^3*b^7*d^2*e^8 - 30*A*a^2*b^8*d^2*e^8 + 35*B*a^4*b^6*
d*e^9 + 20*A*a^3*b^7*d*e^9 - 6*B*a^5*b^5*e^10 - 5*A*a^4*b^6*e^10)/((e*x +
d)^5*e^5) + 26460*(11*B*b^10*d^6*e^6 - 60*B*a*b^9*d^5*e^7 - 6*A*b^10*d^5*e
^7 + 135*B*a^2*b^8*d^4*e^8 + 30*A*a*b^9*d^4*e^8 - 160*B*a^3*b^7*d^3*e^9 -
60*A*a^2*b^8*d^3*e^9 + 105*B*a^4*b^6*d^2*e^10 + 60*A*a^3*b^7*d^2*e^10 - 36
*B*a^5*b^5*d*e^11 - 30*A*a^4*b^6*d*e^11 + 5*B*a^6*b^4*e^12 + 6*A*a^5*b^5*e
^12)/((e*x + d)^6*e^6) - 25200*(11*B*b^10*d^7*e^7 - 70*B*a*b^9*d^6*e^8 - 7
*A*b^10*d^6*e^8 + 189*B*a^2*b^8*d^5*e^9 + 42*A*a*b^9*d^5*e^9 - 280*B*a^3*b
^7*d^4*e^10 - 105*A*a^2*b^8*d^4*e^10 + 245*B*a^4*b^6*d^3*e^11 + 140*A*a^3*
b^7*d^3*e^11 - 126*B*a^5*b^5*d^2*e^12 - 105*A*a^4*b^6*d^2*e^12 + 35*B*a^6*
b^4*d*e^13 + 42*A*a^5*b^5*d*e^13 - 4*B*a^7*b^3*e^14 - 7*A*a^6*b^4*e^14)...

```

Mupad [B] (verification not implemented)

Time = 1.12 (sec) , antiderivative size = 7792, normalized size of antiderivative = 17.51

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^2} dx = \text{Too large to display}$$

input

```
int(((A + B*x)*(a + b*x)^10)/(d + e*x)^2,x)
```

output

```

x^4*((d*((d^2*((2*d*((2*d*((A*b^10 + 10*B*a*b^9)/e^2 - (2*B*b^10*d)/e^3))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^2 + (B*b^10*d^2)/e^4))/e - (d^2*((A*b^10 + 10*B*a*b^9)/e^2 - (2*B*b^10*d)/e^3))/e^2 + (15*a^2*b^7*(3*A*b + 8*B*a))/e^2))/e^2 + (2*d*((d^2*((2*d*((A*b^10 + 10*B*a*b^9)/e^2 - (2*B*b^10*d)/e^3))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^2 + (B*b^10*d^2)/e^4))/e^2 - (2*d*((2*d*((2*d*((A*b^10 + 10*B*a*b^9)/e^2 - (2*B*b^10*d)/e^3))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^2 + (B*b^10*d^2)/e^4))/e - (d^2*((A*b^10 + 10*B*a*b^9)/e^2 - (2*B*b^10*d)/e^3))/e^2 + (15*a^2*b^7*(3*A*b + 8*B*a))/e^2))/e + (30*a^3*b^6*(4*A*b + 7*B*a))/e^2))/e - (42*a^4*b^5*(5*A*b + 6*B*a))/e^2))/(2*e) - (d^2*((d^2*((2*d*((A*b^10 + 10*B*a*b^9)/e^2 - (2*B*b^10*d)/e^3))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^2 + (B*b^10*d^2)/e^4))/e^2 - (2*d*((2*d*((2*d*((A*b^10 + 10*B*a*b^9)/e^2 - (2*B*b^10*d)/e^3))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^2 + (B*b^10*d^2)/e^4))/e - (d^2*((A*b^10 + 10*B*a*b^9)/e^2 - (2*B*b^10*d)/e^3))/e^2 + (15*a^2*b^7*(3*A*b + 8*B*a))/e^2))/e + (30*a^3*b^6*(4*A*b + 7*B*a))/e^2))/(4*e^2) + (21*a^5*b^4*(6*A*b + 5*B*a))/(2*e^2) + x*((2*d*((2*d*((d^2*((d^2*((2*d*((2*d*((A*b^10 + 10*B*a*b^9)/e^2 - (2*B*b^10*d)/e^3))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^2 + (B*b^10*d^2)/e^4))/e - (d^2*((A*b^10 + 10*B*a*b^9)/e^2 - (2*B*b^10*d)/e^3))/e^2 + (15*a^2*b^7*(3*A*b + 8*B*a))/e^2))/e^2 + (2*d*((d^2*((2*d*((A*b^10 + 10*B*a*b^9)/e^2 - (2*B*b^10*d)/e^3))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^2 + (B*b^10*d^2)/e^4))/e^2 - (2*d*((2*d...

```

Reduce [B] (verification not implemented)

Time = 1.44 (sec) , antiderivative size = 1474, normalized size of antiderivative = 3.31

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^2} dx = \text{Too large to display}$$

input

```
int((b*x+a)^10*(B*x+A)/(e*x+d)^2,x)
```

output

```
(27720*log(d + e*x)*a**10*b*d**2*e**10 + 27720*log(d + e*x)*a**10*b*d*e**1
1*x - 277200*log(d + e*x)*a**9*b**2*d**3*e**9 - 277200*log(d + e*x)*a**9*b
**2*d**2*e**10*x + 1247400*log(d + e*x)*a**8*b**3*d**4*e**8 + 1247400*log(
d + e*x)*a**8*b**3*d**3*e**9*x - 3326400*log(d + e*x)*a**7*b**4*d**5*e**7
- 3326400*log(d + e*x)*a**7*b**4*d**4*e**8*x + 5821200*log(d + e*x)*a**6*b
**5*d**6*e**6 + 5821200*log(d + e*x)*a**6*b**5*d**5*e**7*x - 6985440*log(d
+ e*x)*a**5*b**6*d**7*e**5 - 6985440*log(d + e*x)*a**5*b**6*d**6*e**6*x +
5821200*log(d + e*x)*a**4*b**7*d**8*e**4 + 5821200*log(d + e*x)*a**4*b**7
*d**7*e**5*x - 3326400*log(d + e*x)*a**3*b**8*d**9*e**3 - 3326400*log(d +
e*x)*a**3*b**8*d**8*e**4*x + 1247400*log(d + e*x)*a**2*b**9*d**10*e**2 + 1
247400*log(d + e*x)*a**2*b**9*d**9*e**3*x - 277200*log(d + e*x)*a*b**10*d*
*11*e - 277200*log(d + e*x)*a*b**10*d**10*e**2*x + 27720*log(d + e*x)*b**1
1*d**12 + 27720*log(d + e*x)*b**11*d**11*e*x + 2520*a**11*e**12*x - 27720*
a**10*b*d*e**11*x + 277200*a**9*b**2*d**2*e**10*x + 138600*a**9*b**2*d*e**
11*x**2 - 1247400*a**8*b**3*d**3*e**9*x - 623700*a**8*b**3*d**2*e**10*x**2
+ 207900*a**8*b**3*d*e**11*x**3 + 3326400*a**7*b**4*d**4*e**8*x + 1663200
*a**7*b**4*d**3*e**9*x**2 - 554400*a**7*b**4*d**2*e**10*x**3 + 277200*a**7
*b**4*d*e**11*x**4 - 5821200*a**6*b**5*d**5*e**7*x - 2910600*a**6*b**5*d**
4*e**8*x**2 + 970200*a**6*b**5*d**3*e**9*x**3 - 485100*a**6*b**5*d**2*e**1
0*x**4 + 291060*a**6*b**5*d*e**11*x**5 + 6985440*a**5*b**6*d**6*e**6*x ...
```

3.81 $\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^3} dx$

Optimal result	831
Mathematica [B] (verified)	832
Rubi [A] (verified)	833
Maple [B] (verified)	835
Fricas [B] (verification not implemented)	836
Sympy [B] (verification not implemented)	837
Maxima [B] (verification not implemented)	838
Giac [B] (verification not implemented)	839
Mupad [B] (verification not implemented)	840
Reduce [B] (verification not implemented)	840

Optimal result

Integrand size = 20, antiderivative size = 445

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^3} dx = \frac{15b^2(bd-ae)^7(11bBd-8Abe-3aBe)x}{e^{11}} + \frac{(bd-ae)^{10}(Bd-Ae)}{2e^{12}(d+ex)^2} - \frac{(bd-ae)^9(11bBd-10Abe-aBe)}{e^{12}(d+ex)} - \frac{15b^3(bd-ae)^6(11bBd-7Abe-4aBe)(d+ex)^2}{e^{12}} + \frac{14b^4(bd-ae)^5(11bBd-6Abe-5aBe)(d+ex)^3}{e^{12}} - \frac{21b^5(bd-ae)^4(11bBd-5Abe-6aBe)(d+ex)^4}{2e^{12}} + \frac{6b^6(bd-ae)^3(11bBd-4Abe-7aBe)(d+ex)^5}{e^{12}} - \frac{5b^7(bd-ae)^2(11bBd-3Abe-8aBe)(d+ex)^6}{2e^{12}} + \frac{5b^8(bd-ae)(11bBd-2Abe-9aBe)(d+ex)^7}{7e^{12}} - \frac{b^9(11bBd-Abe-10aBe)(d+ex)^8}{8e^{12}} + \frac{b^{10}B(d+ex)^9}{9e^{12}} - \frac{5b(bd-ae)^8(11bBd-9Abe-2aBe)\log(d+ex)}{e^{12}}$$

output

```

15*b^2*(-a*e+b*d)^7*(-8*A*b*e-3*B*a*e+11*B*b*d)*x/e^11+1/2*(-a*e+b*d)^10*(
-A*e+B*d)/e^12/(e*x+d)^2-(-a*e+b*d)^9*(-10*A*b*e-B*a*e+11*B*b*d)/e^12/(e*x
+d)-15*b^3*(-a*e+b*d)^6*(-7*A*b*e-4*B*a*e+11*B*b*d)*(e*x+d)^2/e^12+14*b^4*
(-a*e+b*d)^5*(-6*A*b*e-5*B*a*e+11*B*b*d)*(e*x+d)^3/e^12-21/2*b^5*(-a*e+b*d
)^4*(-5*A*b*e-6*B*a*e+11*B*b*d)*(e*x+d)^4/e^12+6*b^6*(-a*e+b*d)^3*(-4*A*b*
e-7*B*a*e+11*B*b*d)*(e*x+d)^5/e^12-5/2*b^7*(-a*e+b*d)^2*(-3*A*b*e-8*B*a*e+
11*B*b*d)*(e*x+d)^6/e^12+5/7*b^8*(-a*e+b*d)*(-2*A*b*e-9*B*a*e+11*B*b*d)*(e
*x+d)^7/e^12-1/8*b^9*(-A*b*e-10*B*a*e+11*B*b*d)*(e*x+d)^8/e^12+1/9*b^10*B*
(e*x+d)^9/e^12-5*b*(-a*e+b*d)^8*(-9*A*b*e-2*B*a*e+11*B*b*d)*ln(e*x+d)/e^12

```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1480 vs. $2(445) = 890$.

Time = 0.42 (sec) , antiderivative size = 1480, normalized size of antiderivative = 3.33

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^3} dx = \text{Too large to display}$$

input

```
Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^3,x]
```

output

```
(-252*a^10*e^10*(A*e + B*(d + 2*e*x)) - 2520*a^9*b*e^9*(A*e*(d + 2*e*x) -
B*d*(3*d + 4*e*x)) + 11340*a^8*b^2*e^8*(A*d*e*(3*d + 4*e*x) + B*(-5*d^3 -
4*d^2*e*x + 4*d*e^2*x^2 + 2*e^3*x^3)) + 30240*a^7*b^3*e^7*(A*e*(-5*d^3 - 4
*d^2*e*x + 4*d*e^2*x^2 + 2*e^3*x^3) + B*(7*d^4 + 2*d^3*e*x - 11*d^2*e^2*x^
2 - 4*d*e^3*x^3 + e^4*x^4)) + 17640*a^6*b^4*e^6*(3*A*e*(7*d^4 + 2*d^3*e*x
- 11*d^2*e^2*x^2 - 4*d*e^3*x^3 + e^4*x^4) + B*(-27*d^5 + 6*d^4*e*x + 63*d^
3*e^2*x^2 + 20*d^2*e^3*x^3 - 5*d*e^4*x^4 + 2*e^5*x^5)) + 10584*a^5*b^5*e^5
*(2*A*e*(-27*d^5 + 6*d^4*e*x + 63*d^3*e^2*x^2 + 20*d^2*e^3*x^3 - 5*d*e^4*x
^4 + 2*e^5*x^5) + 3*B*(22*d^6 - 16*d^5*e*x - 68*d^4*e^2*x^2 - 20*d^3*e^3*x
^3 + 5*d^2*e^4*x^4 - 2*d*e^5*x^5 + e^6*x^6)) + 5292*a^4*b^6*e^4*(5*A*e*(22
*d^6 - 16*d^5*e*x - 68*d^4*e^2*x^2 - 20*d^3*e^3*x^3 + 5*d^2*e^4*x^4 - 2*d*
e^5*x^5 + e^6*x^6) + B*(-130*d^7 + 160*d^6*e*x + 500*d^5*e^2*x^2 + 140*d^4
*e^3*x^3 - 35*d^3*e^4*x^4 + 14*d^2*e^5*x^5 - 7*d*e^6*x^6 + 4*e^7*x^7)) + 1
008*a^3*b^7*e^3*(3*A*e*(-130*d^7 + 160*d^6*e*x + 500*d^5*e^2*x^2 + 140*d^4
*e^3*x^3 - 35*d^3*e^4*x^4 + 14*d^2*e^5*x^5 - 7*d*e^6*x^6 + 4*e^7*x^7) + 2*
B*(225*d^8 - 390*d^7*e*x - 1035*d^6*e^2*x^2 - 280*d^5*e^3*x^3 + 70*d^4*e^4
*x^4 - 28*d^3*e^5*x^5 + 14*d^2*e^6*x^6 - 8*d*e^7*x^7 + 5*e^8*x^8)) + 108*a
^2*b^8*e^2*(7*A*e*(225*d^8 - 390*d^7*e*x - 1035*d^6*e^2*x^2 - 280*d^5*e^3*
x^3 + 70*d^4*e^4*x^4 - 28*d^3*e^5*x^5 + 14*d^2*e^6*x^6 - 8*d*e^7*x^7 + 5*e
^8*x^8) - 3*B*(595*d^9 - 1330*d^8*e*x - 3185*d^7*e^2*x^2 - 840*d^6*e^3*...
```

Rubi [A] (verified)

Time = 1.61 (sec) , antiderivative size = 445, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^3} dx$$

$$\downarrow 86$$

$$\int \left(\frac{b^9(d + ex)^7(10aBe + Abe - 11bBd)}{e^{11}} - \frac{5b^8(d + ex)^6(bd - ae)(9aBe + 2Abe - 11bBd)}{e^{11}} + \frac{15b^7(d + ex)^5(bd - ae)^2}{e^{11}} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
& - \frac{b^9(d+ex)^8(-10aBe - Abe + 11bBd)}{8e^{12}} + \frac{5b^8(d+ex)^7(bd-ae)(-9aBe - 2Abe + 11bBd)}{7e^{12}} - \\
& \frac{5b^7(d+ex)^6(bd-ae)^2(-8aBe - 3Abe + 11bBd)}{2e^{12}} + \\
& \frac{6b^6(d+ex)^5(bd-ae)^3(-7aBe - 4Abe + 11bBd)}{e^{12}} - \\
& \frac{21b^5(d+ex)^4(bd-ae)^4(-6aBe - 5Abe + 11bBd)}{2e^{12}} + \\
& \frac{14b^4(d+ex)^3(bd-ae)^5(-5aBe - 6Abe + 11bBd)}{e^{12}} - \\
& \frac{15b^3(d+ex)^2(bd-ae)^6(-4aBe - 7Abe + 11bBd)}{e^{12}} + \\
& \frac{15b^2x(bd-ae)^7(-3aBe - 8Abe + 11bBd)}{e^{11}} - \frac{(bd-ae)^9(-aBe - 10Abe + 11bBd)}{e^{12}(d+ex)} + \\
& \frac{(bd-ae)^{10}(Bd-Ae)}{2e^{12}(d+ex)^2} - \frac{5b(bd-ae)^8 \log(d+ex)(-2aBe - 9Abe + 11bBd)}{e^{12}} + \frac{b^{10}B(d+ex)^9}{9e^{12}}
\end{aligned}$$

input `Int[((a + b*x)^10*(A + B*x))/(d + e*x)^3,x]`

output `(15*b^2*(b*d - a*e)^7*(11*b*B*d - 8*A*b*e - 3*a*B*e)*x)/e^11 + ((b*d - a*e)^10*(B*d - A*e))/(2*e^12*(d + e*x)^2) - ((b*d - a*e)^9*(11*b*B*d - 10*A*b*e - a*B*e))/(e^12*(d + e*x)) - (15*b^3*(b*d - a*e)^6*(11*b*B*d - 7*A*b*e - 4*a*B*e)*(d + e*x)^2)/e^12 + (14*b^4*(b*d - a*e)^5*(11*b*B*d - 6*A*b*e - 5*a*B*e)*(d + e*x)^3)/e^12 - (21*b^5*(b*d - a*e)^4*(11*b*B*d - 5*A*b*e - 6*a*B*e)*(d + e*x)^4)/(2*e^12) + (6*b^6*(b*d - a*e)^3*(11*b*B*d - 4*A*b*e - 7*a*B*e)*(d + e*x)^5)/e^12 - (5*b^7*(b*d - a*e)^2*(11*b*B*d - 3*A*b*e - 8*a*B*e)*(d + e*x)^6)/(2*e^12) + (5*b^8*(b*d - a*e)*(11*b*B*d - 2*A*b*e - 9*a*B*e)*(d + e*x)^7)/(7*e^12) - (b^9*(11*b*B*d - A*b*e - 10*a*B*e)*(d + e*x)^8)/(8*e^12) + (b^10*B*(d + e*x)^9)/(9*e^12) - (5*b*(b*d - a*e)^8*(11*b*B*d - 9*A*b*e - 2*a*B*e)*Log[d + e*x])/e^12`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.))^(n_.)*((e_.) + (f_.)*(x_.))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1901 vs. $2(433) = 866$.

Time = 0.24 (sec) , antiderivative size = 1902, normalized size of antiderivative = 4.27

method	result	size
norman	Expression too large to display	1902
default	Expression too large to display	2117
risch	Expression too large to display	2236
parallelrisc	Expression too large to display	3127

input

```
int((b*x+a)^10*(B*x+A)/(e*x+d)^3,x,method=_RETURNVERBOSE)
```

output

```
(-1/2*(A*a^10*e^11+10*A*a^9*b*d*e^10-135*A*a^8*b^2*d^2*e^9+1080*A*a^7*b^3*d^3*e^8-3780*A*a^6*b^4*d^4*e^7+7560*A*a^5*b^5*d^5*e^6-9450*A*a^4*b^6*d^6*e^5+7560*A*a^3*b^7*d^7*e^4-3780*A*a^2*b^8*d^8*e^3+1080*A*a*b^9*d^9*e^2-135*A*b^10*d^10*e+B*a^10*d*e^10-30*B*a^9*b*d^2*e^9+405*B*a^8*b^2*d^3*e^8-2160*B*a^7*b^3*d^4*e^7+6300*B*a^6*b^4*d^5*e^6-11340*B*a^5*b^5*d^6*e^5+13230*B*a^4*b^6*d^7*e^4-10080*B*a^3*b^7*d^8*e^3+4860*B*a^2*b^8*d^9*e^2-1350*B*a*b^9*d^10*e+165*B*b^10*d^11)/e^12-(10*A*a^9*b*e^10-90*A*a^8*b^2*d*e^9+720*A*a^7*b^3*d^2*e^8-2520*A*a^6*b^4*d^3*e^7+5040*A*a^5*b^5*d^4*e^6-6300*A*a^4*b^6*d^5*e^5+5040*A*a^3*b^7*d^6*e^4-2520*A*a^2*b^8*d^7*e^3+720*A*a*b^9*d^8*e^2-90*A*b^10*d^9*e+B*a^10*e^10-20*B*a^9*b*d*e^9+270*B*a^8*b^2*d^2*e^8-1440*B*a^7*b^3*d^3*e^7+4200*B*a^6*b^4*d^4*e^6-7560*B*a^5*b^5*d^5*e^5+8820*B*a^4*b^6*d^6*e^4-6720*B*a^3*b^7*d^7*e^3+3240*B*a^2*b^8*d^8*e^2-900*B*a*b^9*d^9*e+110*B*b^10*d^10)/e^11*x+5/3*b^2*(72*A*a^7*b*e^8-252*A*a^6*b^2*d*e^7+504*A*a^5*b^3*d^2*e^6-630*A*a^4*b^4*d^3*e^5+504*A*a^3*b^5*d^4*e^4-252*A*a^2*b^6*d^5*e^3+72*A*a*b^7*d^6*e^2-9*A*b^8*d^7*e+27*B*a^8*e^8-144*B*a^7*b*d*e^7+420*B*a^6*b^2*d^2*e^6-756*B*a^5*b^3*d^3*e^5+882*B*a^4*b^4*d^4*e^4-672*B*a^3*b^5*d^5*e^3+324*B*a^2*b^6*d^6*e^2-90*B*a*b^7*d^7*e+11*B*b^8*d^8)/e^9*x^3+5/12*b^3*(252*A*a^6*b*e^7-504*A*a^5*b^2*d*e^6+630*A*a^4*b^3*d^2*e^5-504*A*a^3*b^4*d^3*e^4+252*A*a^2*b^5*d^4*e^3-72*A*a*b^6*d^5*e^2+9*A*b^7*d^6*e+144*B*a^7*e^7-420*B*a^6*b*d*e^6+756*B*a^5*b^2*d^2*e^5-882*B*a^4*b^3*d^3*e...
```


Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2547 vs. $2(433) = 866$.

Time = 0.15 (sec) , antiderivative size = 2547, normalized size of antiderivative = 5.72

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^3} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^3,x, algorithm="fricas")`

output

```
1/504*(56*B*b^10*e^11*x^11 - 5292*B*b^10*d^11 - 252*A*a^10*e^11 + 4788*(10
*B*a*b^9 + A*b^10)*d^10*e - 21420*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 + 5670
0*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 - 98280*(7*B*a^4*b^6 + 4*A*a^3*b^7)*
d^7*e^4 + 116424*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 - 95256*(5*B*a^6*b^4
+ 6*A*a^5*b^5)*d^5*e^6 + 52920*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 - 18900
*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^3*e^8 + 3780*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*
e^9 - 252*(B*a^10 + 10*A*a^9*b)*d*e^10 - 7*(11*B*b^10*d*e^10 - 9*(10*B*a*b
^9 + A*b^10)*e^11)*x^10 + 10*(11*B*b^10*d^2*e^9 - 9*(10*B*a*b^9 + A*b^10)*
d*e^10 + 36*(9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 - 15*(11*B*b^10*d^3*e^8 -
9*(10*B*a*b^9 + A*b^10)*d^2*e^9 + 36*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 - 84
*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 24*(11*B*b^10*d^4*e^7 - 9*(10*B*a
*b^9 + A*b^10)*d^3*e^8 + 36*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 - 84*(8*B*a^
3*b^7 + 3*A*a^2*b^8)*d*e^10 + 126*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7 -
42*(11*B*b^10*d^5*e^6 - 9*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 36*(9*B*a^2*b^8
+ 2*A*a*b^9)*d^3*e^8 - 84*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 126*(7*B*a
^4*b^6 + 4*A*a^3*b^7)*d*e^10 - 126*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 +
84*(11*B*b^10*d^6*e^5 - 9*(10*B*a*b^9 + A*b^10)*d^5*e^6 + 36*(9*B*a^2*b^8
+ 2*A*a*b^9)*d^4*e^7 - 84*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 + 126*(7*B*
a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 - 126*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10 +
84*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 - 210*(11*B*b^10*d^7*e^4 - 9*(...
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2004 vs. $2(462) = 924$.

Time = 22.45 (sec) , antiderivative size = 2004, normalized size of antiderivative = 4.50

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^3} dx = \text{Too large to display}$$

input `integrate((b*x+a)**10*(B*x+A)/(e*x+d)**3,x)`

output

```
B*b**10*x**9/(9*e**3) + 5*b*(a*e - b*d)**8*(9*A*b*e + 2*B*a*e - 11*B*b*d)*
log(d + e*x)/e**12 + x**8*(A*b**10/(8*e**3) + 5*B*a*b**9/(4*e**3) - 3*B*b*
*10*d/(8*e**4)) + x**7*(10*A*a*b**9/(7*e**3) - 3*A*b**10*d/(7*e**4) + 45*B
*a**2*b**8/(7*e**3) - 30*B*a*b**9*d/(7*e**4) + 6*B*b**10*d**2/(7*e**5)) +
x**6*(15*A*a**2*b**8/(2*e**3) - 5*A*a*b**9*d/e**4 + A*b**10*d**2/e**5 + 20
*B*a**3*b**7/e**3 - 45*B*a**2*b**8*d/(2*e**4) + 10*B*a*b**9*d**2/e**5 - 5*
B*b**10*d**3/(3*e**6)) + x**5*(24*A*a**3*b**7/e**3 - 27*A*a**2*b**8*d/e**4
+ 12*A*a*b**9*d**2/e**5 - 2*A*b**10*d**3/e**6 + 42*B*a**4*b**6/e**3 - 72*
B*a**3*b**7*d/e**4 + 54*B*a**2*b**8*d**2/e**5 - 20*B*a*b**9*d**3/e**6 + 3*
B*b**10*d**4/e**7) + x**4*(105*A*a**4*b**6/(2*e**3) - 90*A*a**3*b**7*d/e**
4 + 135*A*a**2*b**8*d**2/(2*e**5) - 25*A*a*b**9*d**3/e**6 + 15*A*b**10*d**
4/(4*e**7) + 63*B*a**5*b**5/e**3 - 315*B*a**4*b**6*d/(2*e**4) + 180*B*a**3
*b**7*d**2/e**5 - 225*B*a**2*b**8*d**3/(2*e**6) + 75*B*a*b**9*d**4/(2*e**7
) - 21*B*b**10*d**5/(4*e**8)) + x**3*(84*A*a**5*b**5/e**3 - 210*A*a**4*b**
6*d/e**4 + 240*A*a**3*b**7*d**2/e**5 - 150*A*a**2*b**8*d**3/e**6 + 50*A*a*
b**9*d**4/e**7 - 7*A*b**10*d**5/e**8 + 70*B*a**6*b**4/e**3 - 252*B*a**5*b*
*5*d/e**4 + 420*B*a**4*b**6*d**2/e**5 - 400*B*a**3*b**7*d**3/e**6 + 225*B*
a**2*b**8*d**4/e**7 - 70*B*a*b**9*d**5/e**8 + 28*B*b**10*d**6/(3*e**9)) +
x**2*(105*A*a**6*b**4/e**3 - 378*A*a**5*b**5*d/e**4 + 630*A*a**4*b**6*d**2
/e**5 - 600*A*a**3*b**7*d**3/e**6 + 675*A*a**2*b**8*d**4/(2*e**7) - 105...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1826 vs. $2(433) = 866$.

Time = 0.08 (sec) , antiderivative size = 1826, normalized size of antiderivative = 4.10

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^3} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^3,x, algorithm="maxima")`

output

```
-1/2*(21*B*b^10*d^11 + A*a^10*e^11 - 19*(10*B*a*b^9 + A*b^10)*d^10*e + 85*
(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 - 225*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^
3 + 390*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e^4 - 462*(6*B*a^5*b^5 + 5*A*a^4*b
^6)*d^6*e^5 + 378*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^5*e^6 - 210*(4*B*a^7*b^3 +
7*A*a^6*b^4)*d^4*e^7 + 75*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^3*e^8 - 15*(2*B*a
^9*b + 9*A*a^8*b^2)*d^2*e^9 + (B*a^10 + 10*A*a^9*b)*d*e^10 + 2*(11*B*b^10*
d^10*e - 10*(10*B*a*b^9 + A*b^10)*d^9*e^2 + 45*(9*B*a^2*b^8 + 2*A*a*b^9)*d
^8*e^3 - 120*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^7*e^4 + 210*(7*B*a^4*b^6 + 4*A*
a^3*b^7)*d^6*e^5 - 252*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^5*e^6 + 210*(5*B*a^6*
b^4 + 6*A*a^5*b^5)*d^4*e^7 - 120*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^3*e^8 + 45*
(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^2*e^9 - 10*(2*B*a^9*b + 9*A*a^8*b^2)*d*e^10
+ (B*a^10 + 10*A*a^9*b)*e^11)*x)/(e^14*x^2 + 2*d*e^13*x + d^2*e^12) + 1/50
4*(56*B*b^10*e^8*x^9 - 63*(3*B*b^10*d*e^7 - (10*B*a*b^9 + A*b^10)*e^8)*x^8
+ 72*(6*B*b^10*d^2*e^6 - 3*(10*B*a*b^9 + A*b^10)*d*e^7 + 5*(9*B*a^2*b^8 +
2*A*a*b^9)*e^8)*x^7 - 84*(10*B*b^10*d^3*e^5 - 6*(10*B*a*b^9 + A*b^10)*d^2
*e^6 + 15*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^7 - 15*(8*B*a^3*b^7 + 3*A*a^2*b^8)
*e^8)*x^6 + 504*(3*B*b^10*d^4*e^4 - 2*(10*B*a*b^9 + A*b^10)*d^3*e^5 + 6*(9
*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^6 - 9*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^7 + 6*
(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^8)*x^5 - 126*(21*B*b^10*d^5*e^3 - 15*(10*B*a
*b^9 + A*b^10)*d^4*e^4 + 50*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^5 - 90*(8*B...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2134 vs. $2(433) = 866$.

Time = 0.14 (sec) , antiderivative size = 2134, normalized size of antiderivative = 4.80

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^3} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^3,x, algorithm="giac")`

output

```
-5*(11*B*b^10*d^9 - 90*B*a*b^9*d^8*e - 9*A*b^10*d^8*e + 324*B*a^2*b^8*d^7*
e^2 + 72*A*a*b^9*d^7*e^2 - 672*B*a^3*b^7*d^6*e^3 - 252*A*a^2*b^8*d^6*e^3 +
882*B*a^4*b^6*d^5*e^4 + 504*A*a^3*b^7*d^5*e^4 - 756*B*a^5*b^5*d^4*e^5 - 6
30*A*a^4*b^6*d^4*e^5 + 420*B*a^6*b^4*d^3*e^6 + 504*A*a^5*b^5*d^3*e^6 - 144
*B*a^7*b^3*d^2*e^7 - 252*A*a^6*b^4*d^2*e^7 + 27*B*a^8*b^2*d*e^8 + 72*A*a^7
*b^3*d*e^8 - 2*B*a^9*b*e^9 - 9*A*a^8*b^2*e^9)*log(abs(e*x + d))/e^12 - 1/2
*(21*B*b^10*d^11 - 190*B*a*b^9*d^10*e - 19*A*b^10*d^10*e + 765*B*a^2*b^8*d
^9*e^2 + 170*A*a*b^9*d^9*e^2 - 1800*B*a^3*b^7*d^8*e^3 - 675*A*a^2*b^8*d^8*
e^3 + 2730*B*a^4*b^6*d^7*e^4 + 1560*A*a^3*b^7*d^7*e^4 - 2772*B*a^5*b^5*d^6
*e^5 - 2310*A*a^4*b^6*d^6*e^5 + 1890*B*a^6*b^4*d^5*e^6 + 2268*A*a^5*b^5*d^
5*e^6 - 840*B*a^7*b^3*d^4*e^7 - 1470*A*a^6*b^4*d^4*e^7 + 225*B*a^8*b^2*d^3
*e^8 + 600*A*a^7*b^3*d^3*e^8 - 30*B*a^9*b*d^2*e^9 - 135*A*a^8*b^2*d^2*e^9
+ B*a^10*d*e^10 + 10*A*a^9*b*d*e^10 + A*a^10*e^11 + 2*(11*B*b^10*d^10*e -
100*B*a*b^9*d^9*e^2 - 10*A*b^10*d^9*e^2 + 405*B*a^2*b^8*d^8*e^3 + 90*A*a*b
^9*d^8*e^3 - 960*B*a^3*b^7*d^7*e^4 - 360*A*a^2*b^8*d^7*e^4 + 1470*B*a^4*b^
6*d^6*e^5 + 840*A*a^3*b^7*d^6*e^5 - 1512*B*a^5*b^5*d^5*e^6 - 1260*A*a^4*b^
6*d^5*e^6 + 1050*B*a^6*b^4*d^4*e^7 + 1260*A*a^5*b^5*d^4*e^7 - 480*B*a^7*b^
3*d^3*e^8 - 840*A*a^6*b^4*d^3*e^8 + 135*B*a^8*b^2*d^2*e^9 + 360*A*a^7*b^3*
d^2*e^9 - 20*B*a^9*b*d*e^10 - 90*A*a^8*b^2*d*e^10 + B*a^10*e^11 + 10*A*a^9
*b*e^11)*x)/((e*x + d)^2*e^12) + 1/504*(56*B*b^10*e^24*x^9 - 189*B*b^10...
```

Mupad [B] (verification not implemented)

Time = 1.21 (sec) , antiderivative size = 8104, normalized size of antiderivative = 18.21

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^3} dx = \text{Too large to display}$$

input `int(((A + B*x)*(a + b*x)^10)/(d + e*x)^3,x)`

output `x^5*((3*d*((3*d^2*((A*b^10 + 10*B*a*b^9)/e^3 - (3*B*b^10*d)/e^4))/e^2 - (3*d*((3*d*((A*b^10 + 10*B*a*b^9)/e^3 - (3*B*b^10*d)/e^4))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^3 + (3*B*b^10*d^2)/e^5))/e - (15*a^2*b^7*(3*A*b + 8*B*a))/e^3 + (B*b^10*d^3)/e^6))/(5*e) - (d^3*((A*b^10 + 10*B*a*b^9)/e^3 - (3*B*b^10*d)/e^4))/(5*e^3) + (3*d^2*((3*d*((A*b^10 + 10*B*a*b^9)/e^3 - (3*B*b^10*d)/e^4))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^3 + (3*B*b^10*d^2)/e^5))/(5*e^2) + (6*a^3*b^6*(4*A*b + 7*B*a))/e^3 + x*((3*d*((3*d*((d^3*((3*d^2*((A*b^10 + 10*B*a*b^9)/e^3 - (3*B*b^10*d)/e^4))/e^2 - (3*d*((3*d*((A*b^10 + 10*B*a*b^9)/e^3 - (3*B*b^10*d)/e^4))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^3 + (3*B*b^10*d^2)/e^5))/e - (15*a^2*b^7*(3*A*b + 8*B*a))/e^3 + (B*b^10*d^3)/e^6))/e^3 - (3*d*((3*d^2*((3*d^2*((A*b^10 + 10*B*a*b^9)/e^3 - (3*B*b^10*d)/e^4))/e^2 - (3*d*((3*d*((A*b^10 + 10*B*a*b^9)/e^3 - (3*B*b^10*d)/e^4))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^3 + (3*B*b^10*d^2)/e^5))/e - (15*a^2*b^7*(3*A*b + 8*B*a))/e^3 + (B*b^10*d^3)/e^6))/e^2 - (3*d*((3*d*((3*d^2*((A*b^10 + 10*B*a*b^9)/e^3 - (3*B*b^10*d)/e^4))/e^2 - (3*d*((3*d*((A*b^10 + 10*B*a*b^9)/e^3 - (3*B*b^10*d)/e^4))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^3 + (3*B*b^10*d^2)/e^5))/e - (15*a^2*b^7*(3*A*b + 8*B*a))/e^3 + (B*b^10*d^3)/e^6))/e - (d^3*((A*b^10 + 10*B*a*b^9)/e^3 - (3*B*b^10*d)/e^4))/e^3 + (3*d^2*((3*d*((A*b^10 + 10*B*a*b^9)/e^3 - (3*B*b^10*d)/e^4))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^3 + (3*B*b^10*d^2)/e^5))/e - (15*a^2*b^7*(3*A*b + 8*B*a))/e^3 + (B*b^10*d^3)/e^6))/e - (d^3*((A*b^10 + 10*B*a*b^9)/e^3 - (3*B*b^10*d)/e^4))/e^3 + (3*d^2*((3*d*((A*b^10 + 10*B*a*b^9)/e^3 - (3*B*b^10*d)/e^4))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^3 + (3*B*b^10*d^2)/e^5))/e^2 + (30*a^3*b^6*(4*A*b + 7*B*a))/e^3))/e + (d^3*((...`

Reduce [B] (verification not implemented)

Time = 0.84 (sec) , antiderivative size = 1661, normalized size of antiderivative = 3.73

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^3} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^3,x)`

output

```
(27720*log(d + e*x)*a**9*b**2*d**3*e**9 + 55440*log(d + e*x)*a**9*b**2*d**
2*e**10*x + 27720*log(d + e*x)*a**9*b**2*d**e**11*x**2 - 249480*log(d + e*x
)*a**8*b**3*d**4*e**8 - 498960*log(d + e*x)*a**8*b**3*d**3*e**9*x - 249480
*log(d + e*x)*a**8*b**3*d**2*e**10*x**2 + 997920*log(d + e*x)*a**7*b**4*d**
5*e**7 + 1995840*log(d + e*x)*a**7*b**4*d**4*e**8*x + 997920*log(d + e*x)
*a**7*b**4*d**3*e**9*x**2 - 2328480*log(d + e*x)*a**6*b**5*d**6*e**6 - 465
6960*log(d + e*x)*a**6*b**5*d**5*e**7*x - 2328480*log(d + e*x)*a**6*b**5*d
**4*e**8*x**2 + 3492720*log(d + e*x)*a**5*b**6*d**7*e**5 + 6985440*log(d +
e*x)*a**5*b**6*d**6*e**6*x + 3492720*log(d + e*x)*a**5*b**6*d**5*e**7*x**
2 - 3492720*log(d + e*x)*a**4*b**7*d**8*e**4 - 6985440*log(d + e*x)*a**4*b
**7*d**7*e**5*x - 3492720*log(d + e*x)*a**4*b**7*d**6*e**6*x**2 + 2328480*
log(d + e*x)*a**3*b**8*d**9*e**3 + 4656960*log(d + e*x)*a**3*b**8*d**8*e**
4*x + 2328480*log(d + e*x)*a**3*b**8*d**7*e**5*x**2 - 997920*log(d + e*x)*
a**2*b**9*d**10*e**2 - 1995840*log(d + e*x)*a**2*b**9*d**9*e**3*x - 997920
*log(d + e*x)*a**2*b**9*d**8*e**4*x**2 + 249480*log(d + e*x)*a*b**10*d**11
*e + 498960*log(d + e*x)*a*b**10*d**10*e**2*x + 249480*log(d + e*x)*a*b**1
0*d**9*e**3*x**2 - 27720*log(d + e*x)*b**11*d**12 - 55440*log(d + e*x)*b**
11*d**11*e*x - 27720*log(d + e*x)*b**11*d**10*e**2*x**2 - 252*a**11*d**e**1
1 + 2772*a**10*b*e**12*x**2 + 13860*a**9*b**2*d**3*e**9 - 27720*a**9*b**2*
d**e**11*x**2 - 124740*a**8*b**3*d**4*e**8 + 249480*a**8*b**3*d**2*e**10...
```

3.82 $\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^4} dx$

Optimal result	842
Mathematica [A] (verified)	843
Rubi [A] (verified)	844
Maple [B] (verified)	846
Fricas [B] (verification not implemented)	847
Sympy [F(-1)]	848
Maxima [B] (verification not implemented)	849
Giac [B] (verification not implemented)	850
Mupad [B] (verification not implemented)	851
Reduce [B] (verification not implemented)	851

Optimal result

Integrand size = 20, antiderivative size = 445

$$\begin{aligned}
 \int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^4} dx = & -\frac{30b^3(bd-ae)^6(11bBd-7Abe-4aBe)x}{e^{11}} \\
 & + \frac{(bd-ae)^{10}(Bd-Ae)}{3e^{12}(d+ex)^3} \\
 & - \frac{(bd-ae)^9(11bBd-10Abe-aBe)}{2e^{12}(d+ex)^2} \\
 & + \frac{5b(bd-ae)^8(11bBd-9Abe-2aBe)}{e^{12}(d+ex)} \\
 & + \frac{21b^4(bd-ae)^5(11bBd-6Abe-5aBe)(d+ex)^2}{e^{12}} \\
 & - \frac{14b^5(bd-ae)^4(11bBd-5Abe-6aBe)(d+ex)^3}{e^{12}} \\
 & + \frac{15b^6(bd-ae)^3(11bBd-4Abe-7aBe)(d+ex)^4}{2e^{12}} \\
 & - \frac{3b^7(bd-ae)^2(11bBd-3Abe-8aBe)(d+ex)^5}{e^{12}} \\
 & + \frac{5b^8(bd-ae)(11bBd-2Abe-9aBe)(d+ex)^6}{6e^{12}} \\
 & - \frac{b^9(11bBd-Abe-10aBe)(d+ex)^7}{7e^{12}} + \frac{b^{10}B(d+ex)^8}{8e^{12}} \\
 & + \frac{15b^2(bd-ae)^7(11bBd-8Abe-3aBe)\log(d+ex)}{e^{12}}
 \end{aligned}$$

output

```
-30*b^3*(-a*e+b*d)^6*(-7*A*b*e-4*B*a*e+11*B*b*d)*x/e^11+1/3*(-a*e+b*d)^10*
(-A*e+B*d)/e^12/(e*x+d)^3-1/2*(-a*e+b*d)^9*(-10*A*b*e-B*a*e+11*B*b*d)/e^12
/(e*x+d)^2+5*b*(-a*e+b*d)^8*(-9*A*b*e-2*B*a*e+11*B*b*d)/e^12/(e*x+d)+21*b^
4*(-a*e+b*d)^5*(-6*A*b*e-5*B*a*e+11*B*b*d)*(e*x+d)^2/e^12-14*b^5*(-a*e+b*d
)^4*(-5*A*b*e-6*B*a*e+11*B*b*d)*(e*x+d)^3/e^12+15/2*b^6*(-a*e+b*d)^3*(-4*A
*b*e-7*B*a*e+11*B*b*d)*(e*x+d)^4/e^12-3*b^7*(-a*e+b*d)^2*(-3*A*b*e-8*B*a*e
+11*B*b*d)*(e*x+d)^5/e^12+5/6*b^8*(-a*e+b*d)*(-2*A*b*e-9*B*a*e+11*B*b*d)*(
e*x+d)^6/e^12-1/7*b^9*(-A*b*e-10*B*a*e+11*B*b*d)*(e*x+d)^7/e^12+1/8*b^10*B
*(e*x+d)^8/e^12+15*b^2*(-a*e+b*d)^7*(-8*A*b*e-3*B*a*e+11*B*b*d)*ln(e*x+d)/
e^12
```

Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 814, normalized size of antiderivative = 1.83

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^4} dx$$

$$= \frac{168b^3e(120a^7Be^7 - 315a^2b^5d^4e^2(8Bd - 5Ae) + 600a^3b^4d^3e^3(7Bd - 4Ae) + 280ab^6d^5e(3Bd - 2Ae) + 5$$

input

```
Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^4,x]
```


output

```
(168*b^3*e*(120*a^7*B*e^7 - 315*a^2*b^5*d^4*e^2*(8*B*d - 5*A*e) + 600*a^3*
b^4*d^3*e^3*(7*B*d - 4*A*e) + 280*a*b^6*d^5*e*(3*B*d - 2*A*e) + 504*a^5*b^
2*d*e^5*(5*B*d - 2*A*e) - 2100*a^4*b^3*d^2*e^4*(2*B*d - A*e) + 210*a^6*b*e
^6*(-4*B*d + A*e) + 12*b^7*d^6*(-10*B*d + 7*A*e))*x - 84*b^4*e^2*(-210*a^6
*B*e^6 + 70*a*b^5*d^4*e*(8*B*d - 5*A*e) - 225*a^2*b^4*d^3*e^2*(7*B*d - 4*A
*e) - 420*a^4*b^2*d*e^4*(5*B*d - 2*A*e) + 1200*a^3*b^3*d^2*e^3*(2*B*d - A*
e) - 252*a^5*b*e^5*(-4*B*d + A*e) + 28*b^6*d^5*(-3*B*d + 2*A*e))*x^2 + 56*
b^5*e^3*(252*a^5*B*e^5 - 7*b^5*d^4*(8*B*d - 5*A*e) + 50*a*b^4*d^3*e*(7*B*d
- 4*A*e) + 240*a^3*b^2*d*e^3*(5*B*d - 2*A*e) - 450*a^2*b^3*d^2*e^2*(2*B*d
- A*e) + 210*a^4*b*e^4*(-4*B*d + A*e))*x^3 - 210*b^6*e^4*(-42*a^4*B*e^4 +
20*a*b^3*d^2*e*(2*B*d - A*e) - 24*a^3*b*e^3*(-4*B*d + A*e) + 18*a^2*b^2*d
*e^2*(-5*B*d + 2*A*e) + b^4*d^3*(-7*B*d + 4*A*e))*x^4 + 168*b^7*e^5*(24*a^
3*B*e^3 + 4*a*b^2*d*e*(5*B*d - 2*A*e) + 9*a^2*b*e^2*(-4*B*d + A*e) + 2*b^3
*d^2*(-2*B*d + A*e))*x^5 - 28*b^8*e^6*(-45*a^2*B*e^2 - 10*a*b*e*(-4*B*d +
A*e) + 2*b^2*d*(-5*B*d + 2*A*e))*x^6 + 24*b^9*e^7*(-4*b*B*d + A*b*e + 10*a
*B*e)*x^7 + 21*b^10*B*e^8*x^8 + (56*(b*d - a*e)^10*(B*d - A*e))/(d + e*x)^
3 - (84*(b*d - a*e)^9*(11*b*B*d - 10*A*b*e - a*B*e))/(d + e*x)^2 + (840*b*
(b*d - a*e)^8*(11*b*B*d - 9*A*b*e - 2*a*B*e))/(d + e*x) + 2520*b^2*(b*d -
a*e)^7*(11*b*B*d - 8*A*b*e - 3*a*B*e)*Log[d + e*x]/(168*e^12)
```

Rubi [A] (verified)

Time = 1.53 (sec) , antiderivative size = 445, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^4} dx$$

↓ 86

$$\int \left(\frac{b^9(d + ex)^6(10aBe + Abe - 11bBd)}{e^{11}} - \frac{5b^8(d + ex)^5(bd - ae)(9aBe + 2Abe - 11bBd)}{e^{11}} + \frac{15b^7(d + ex)^4(bd - ae)^2}{e^{11}} \right) dx$$

↓ 2009

$$\begin{aligned}
& - \frac{b^9(d+ex)^7(-10aBe - Abe + 11bBd)}{7e^{12}} + \frac{5b^8(d+ex)^6(bd-ae)(-9aBe - 2Abe + 11bBd)}{6e^{12}} - \\
& \frac{3b^7(d+ex)^5(bd-ae)^2(-8aBe - 3Abe + 11bBd)}{e^{12}} + \\
& \frac{15b^6(d+ex)^4(bd-ae)^3(-7aBe - 4Abe + 11bBd)}{e^{12}} - \\
& \frac{14b^5(d+ex)^3(bd-ae)^4(-6aBe - 5Abe + 11bBd)}{2e^{12}} + \\
& \frac{21b^4(d+ex)^2(bd-ae)^5(-5aBe - 6Abe + 11bBd)}{e^{12}} - \\
& \frac{30b^3x(bd-ae)^6(-4aBe - 7Abe + 11bBd)}{e^{12}} + \\
& \frac{15b^2(bd-ae)^7 \log(d+ex)(-3aBe - 8Abe + 11bBd)}{e^{11}} + \\
& \frac{5b(bd-ae)^8(-2aBe - 9Abe + 11bBd)}{e^{12}(d+ex)} - \frac{(bd-ae)^9(-aBe - 10Abe + 11bBd)}{2e^{12}(d+ex)^2} + \\
& \frac{(bd-ae)^{10}(Bd-Ae)}{3e^{12}(d+ex)^3} + \frac{b^{10}B(d+ex)^8}{8e^{12}}
\end{aligned}$$

input `Int[((a + b*x)^10*(A + B*x))/(d + e*x)^4,x]`

output `(-30*b^3*(b*d - a*e)^6*(11*b*B*d - 7*A*b*e - 4*a*B*e)*x)/e^11 + ((b*d - a*e)^10*(B*d - A*e))/(3*e^12*(d + e*x)^3) - ((b*d - a*e)^9*(11*b*B*d - 10*A*b*e - a*B*e))/(2*e^12*(d + e*x)^2) + (5*b*(b*d - a*e)^8*(11*b*B*d - 9*A*b*e - 2*a*B*e))/(e^12*(d + e*x)) + (21*b^4*(b*d - a*e)^5*(11*b*B*d - 6*A*b*e - 5*a*B*e)*(d + e*x)^2)/e^12 - (14*b^5*(b*d - a*e)^4*(11*b*B*d - 5*A*b*e - 6*a*B*e)*(d + e*x)^3)/e^12 + (15*b^6*(b*d - a*e)^3*(11*b*B*d - 4*A*b*e - 7*a*B*e)*(d + e*x)^4)/(2*e^12) - (3*b^7*(b*d - a*e)^2*(11*b*B*d - 3*A*b*e - 8*a*B*e)*(d + e*x)^5)/e^12 + (5*b^8*(b*d - a*e)*(11*b*B*d - 2*A*b*e - 9*a*B*e)*(d + e*x)^6)/(6*e^12) - (b^9*(11*b*B*d - A*b*e - 10*a*B*e)*(d + e*x)^7)/(7*e^12) + (b^10*B*(d + e*x)^8)/(8*e^12) + (15*b^2*(b*d - a*e)^7*(11*b*B*d - 8*A*b*e - 3*a*B*e)*Log[d + e*x])/e^12`

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1906 vs. $2(433) = 866$.

Time = 0.24 (sec) , antiderivative size = 1907, normalized size of antiderivative = 4.29

method	result	size
norman	Expression too large to display	1907
default	Expression too large to display	2073
risch	Expression too large to display	2182
parallelrisc	Expression too large to display	3431

input

```
int((b*x+a)^10*(B*x+A)/(e*x+d)^4,x,method=_RETURNVERBOSE)
```

output

```
(-1/6*(2*A*a^10*e^11+10*A*a^9*b*d*e^10+90*A*a^8*b^2*d^2*e^9-1320*A*a^7*b^3*d^3*e^8+9240*A*a^6*b^4*d^4*e^7-27720*A*a^5*b^5*d^5*e^6+46200*A*a^4*b^6*d^6*e^5-46200*A*a^3*b^7*d^7*e^4+27720*A*a^2*b^8*d^8*e^3-9240*A*a*b^9*d^9*e^2+1320*A*b^10*d^10*e+B*a^10*d*e^10+20*B*a^9*b*d^2*e^9-495*B*a^8*b^2*d^3*e^8+5280*B*a^7*b^3*d^4*e^7-23100*B*a^6*b^4*d^5*e^6+55440*B*a^5*b^5*d^6*e^5-80850*B*a^4*b^6*d^7*e^4+73920*B*a^3*b^7*d^8*e^3-41580*B*a^2*b^8*d^9*e^2+13200*B*a*b^9*d^10*e-1815*B*b^10*d^11)/e^12-(45*A*a^8*b^2*e^9-360*A*a^7*b^3*d*e^8+2520*A*a^6*b^4*d^2*e^7-7560*A*a^5*b^5*d^3*e^6+12600*A*a^4*b^6*d^4*e^5-12600*A*a^3*b^7*d^5*e^4+7560*A*a^2*b^8*d^6*e^3-2520*A*a*b^9*d^7*e^2+360*A*b^10*d^8*e+10*B*a^9*b*e^9-135*B*a^8*b^2*d*e^8+1440*B*a^7*b^3*d^2*e^7-6300*B*a^6*b^4*d^3*e^6+15120*B*a^5*b^5*d^4*e^5-22050*B*a^4*b^6*d^5*e^4+20160*B*a^3*b^7*d^6*e^3-11340*B*a^2*b^8*d^7*e^2+3600*B*a*b^9*d^8*e-495*B*b^10*d^9)/e^10*x^2-1/2*(10*A*a^9*b*e^10+90*A*a^8*b^2*d*e^9-1080*A*a^7*b^3*d^2*e^8+7560*A*a^6*b^4*d^3*e^7-22680*A*a^5*b^5*d^4*e^6+37800*A*a^4*b^6*d^5*e^5-37800*A*a^3*b^7*d^6*e^4+22680*A*a^2*b^8*d^7*e^3-7560*A*a*b^9*d^8*e^2+1080*A*b^10*d^9*e+B*a^10*e^10+20*B*a^9*b*d*e^9-405*B*a^8*b^2*d^2*e^8+4320*B*a^7*b^3*d^3*e^7-18900*B*a^6*b^4*d^4*e^6+45360*B*a^5*b^5*d^5*e^5-66150*B*a^4*b^6*d^6*e^4+60480*B*a^3*b^7*d^7*e^3-34020*B*a^2*b^8*d^8*e^2+10800*B*a*b^9*d^9*e-1485*B*b^10*d^10)/e^11*x+15/4*b^3*(56*A*a^6*b*e^7-168*A*a^5*b^2*d*e^6+280*A*a^4*b^3*d^2*e^5-280*A*a^3*b^4*d^3*e^4+168*A*a^2*b^5*d^4*e^3-56*A*a*b...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2702 vs. $2(433) = 866$.

Time = 0.13 (sec) , antiderivative size = 2702, normalized size of antiderivative = 6.07

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^4} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)/(e*x+d)^4,x, algorithm="fricas")
```

output

```

1/168*(21*B*b^10*e^11*x^11 + 8372*B*b^10*d^11 - 56*A*a^10*e^11 - 6776*(10*
B*a*b^9 + A*b^10)*d^10*e + 26740*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 - 61320
*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 89880*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d
^7*e^4 - 87024*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 55272*(5*B*a^6*b^4 +
6*A*a^5*b^5)*d^5*e^6 - 21840*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 4620*(3
*B*a^8*b^2 + 8*A*a^7*b^3)*d^3*e^8 - 280*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9
- 28*(B*a^10 + 10*A*a^9*b)*d*e^10 - 3*(11*B*b^10*d*e^10 - 8*(10*B*a*b^9 +
A*b^10)*e^11)*x^10 + 5*(11*B*b^10*d^2*e^9 - 8*(10*B*a*b^9 + A*b^10)*d*e^10
+ 28*(9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 - 9*(11*B*b^10*d^3*e^8 - 8*(10*B
*a*b^9 + A*b^10)*d^2*e^9 + 28*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 - 56*(8*B*a
^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 18*(11*B*b^10*d^4*e^7 - 8*(10*B*a*b^9 +
A*b^10)*d^3*e^8 + 28*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 - 56*(8*B*a^3*b^7 +
3*A*a^2*b^8)*d*e^10 + 70*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7 - 42*(11*B
*b^10*d^5*e^6 - 8*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 28*(9*B*a^2*b^8 + 2*A*a*
b^9)*d^3*e^8 - 56*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 70*(7*B*a^4*b^6 +
4*A*a^3*b^7)*d*e^10 - 56*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 + 126*(11*B
*b^10*d^6*e^5 - 8*(10*B*a*b^9 + A*b^10)*d^5*e^6 + 28*(9*B*a^2*b^8 + 2*A*a*
b^9)*d^4*e^7 - 56*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 + 70*(7*B*a^4*b^6 +
4*A*a^3*b^7)*d^2*e^9 - 56*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10 + 28*(5*B*a^6
*b^4 + 6*A*a^5*b^5)*e^11)*x^5 - 630*(11*B*b^10*d^7*e^4 - 8*(10*B*a*b^9 ...

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^4} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**10*(B*x+A)/(e*x+d)**4,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1839 vs. $2(433) = 866$.

Time = 0.11 (sec) , antiderivative size = 1839, normalized size of antiderivative = 4.13

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^4} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^4,x, algorithm="maxima")`

output

```
1/6*(299*B*b^10*d^11 - 2*A*a^10*e^11 - 242*(10*B*a*b^9 + A*b^10)*d^10*e +
955*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 - 2190*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d
^8*e^3 + 3210*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e^4 - 3108*(6*B*a^5*b^5 + 5*
A*a^4*b^6)*d^6*e^5 + 1974*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^5*e^6 - 780*(4*B*a
^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 165*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^3*e^8 -
10*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 - (B*a^10 + 10*A*a^9*b)*d*e^10 + 30*(
11*B*b^10*d^9*e^2 - 9*(10*B*a*b^9 + A*b^10)*d^8*e^3 + 36*(9*B*a^2*b^8 + 2*
A*a*b^9)*d^7*e^4 - 84*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^6*e^5 + 126*(7*B*a^4*b
^6 + 4*A*a^3*b^7)*d^5*e^6 - 126*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^4*e^7 + 84*(
5*B*a^6*b^4 + 6*A*a^5*b^5)*d^3*e^8 - 36*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^2*e^
9 + 9*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d*e^10 - (2*B*a^9*b + 9*A*a^8*b^2)*e^11)
*x^2 + 3*(209*B*b^10*d^10*e - 170*(10*B*a*b^9 + A*b^10)*d^9*e^2 + 675*(9*B
*a^2*b^8 + 2*A*a*b^9)*d^8*e^3 - 1560*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^7*e^4 +
2310*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^6*e^5 - 2268*(6*B*a^5*b^5 + 5*A*a^4*b^
6)*d^5*e^6 + 1470*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^4*e^7 - 600*(4*B*a^7*b^3 +
7*A*a^6*b^4)*d^3*e^8 + 135*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^2*e^9 - 10*(2*B*
a^9*b + 9*A*a^8*b^2)*d*e^10 - (B*a^10 + 10*A*a^9*b)*e^11)*x)/(e^15*x^3 + 3
*d*e^14*x^2 + 3*d^2*e^13*x + d^3*e^12) + 1/168*(21*B*b^10*e^7*x^8 - 24*(4*
B*b^10*d*e^6 - (10*B*a*b^9 + A*b^10)*e^7)*x^7 + 28*(10*B*b^10*d^2*e^5 - 4*
(10*B*a*b^9 + A*b^10)*d*e^6 + 5*(9*B*a^2*b^8 + 2*A*a*b^9)*e^7)*x^6 - 16...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2096 vs. $2(433) = 866$.

Time = 0.13 (sec) , antiderivative size = 2096, normalized size of antiderivative = 4.71

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^4} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^4,x, algorithm="giac")`

output

```
15*(11*B*b^10*d^8 - 80*B*a*b^9*d^7*e - 8*A*b^10*d^7*e + 252*B*a^2*b^8*d^6*
e^2 + 56*A*a*b^9*d^6*e^2 - 448*B*a^3*b^7*d^5*e^3 - 168*A*a^2*b^8*d^5*e^3 +
 490*B*a^4*b^6*d^4*e^4 + 280*A*a^3*b^7*d^4*e^4 - 336*B*a^5*b^5*d^3*e^5 - 2
80*A*a^4*b^6*d^3*e^5 + 140*B*a^6*b^4*d^2*e^6 + 168*A*a^5*b^5*d^2*e^6 - 32*
B*a^7*b^3*d*e^7 - 56*A*a^6*b^4*d*e^7 + 3*B*a^8*b^2*e^8 + 8*A*a^7*b^3*e^8)*
log(abs(e*x + d))/e^12 + 1/6*(299*B*b^10*d^11 - 2420*B*a*b^9*d^10*e - 242*
A*b^10*d^10*e + 8595*B*a^2*b^8*d^9*e^2 + 1910*A*a*b^9*d^9*e^2 - 17520*B*a^
3*b^7*d^8*e^3 - 6570*A*a^2*b^8*d^8*e^3 + 22470*B*a^4*b^6*d^7*e^4 + 12840*A
*a^3*b^7*d^7*e^4 - 18648*B*a^5*b^5*d^6*e^5 - 15540*A*a^4*b^6*d^6*e^5 + 987
0*B*a^6*b^4*d^5*e^6 + 11844*A*a^5*b^5*d^5*e^6 - 3120*B*a^7*b^3*d^4*e^7 - 5
460*A*a^6*b^4*d^4*e^7 + 495*B*a^8*b^2*d^3*e^8 + 1320*A*a^7*b^3*d^3*e^8 - 2
0*B*a^9*b*d^2*e^9 - 90*A*a^8*b^2*d^2*e^9 - B*a^10*d*e^10 - 10*A*a^9*b*d*e^
10 - 2*A*a^10*e^11 + 30*(11*B*b^10*d^9*e^2 - 90*B*a*b^9*d^8*e^3 - 9*A*b^10
*d^8*e^3 + 324*B*a^2*b^8*d^7*e^4 + 72*A*a*b^9*d^7*e^4 - 672*B*a^3*b^7*d^6*
e^5 - 252*A*a^2*b^8*d^6*e^5 + 882*B*a^4*b^6*d^5*e^6 + 504*A*a^3*b^7*d^5*e^
6 - 756*B*a^5*b^5*d^4*e^7 - 630*A*a^4*b^6*d^4*e^7 + 420*B*a^6*b^4*d^3*e^8
+ 504*A*a^5*b^5*d^3*e^8 - 144*B*a^7*b^3*d^2*e^9 - 252*A*a^6*b^4*d^2*e^9 +
27*B*a^8*b^2*d*e^10 + 72*A*a^7*b^3*d*e^10 - 2*B*a^9*b*e^11 - 9*A*a^8*b^2*e
^11)*x^2 + 3*(209*B*b^10*d^10*e - 1700*B*a*b^9*d^9*e^2 - 170*A*b^10*d^9*e^
2 + 6075*B*a^2*b^8*d^8*e^3 + 1350*A*a*b^9*d^8*e^3 - 12480*B*a^3*b^7*d^7...
```

Mupad [B] (verification not implemented)

Time = 1.18 (sec) , antiderivative size = 5544, normalized size of antiderivative = 12.46

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^4} dx = \text{Too large to display}$$

input `int(((A + B*x)*(a + b*x)^10)/(d + e*x)^4,x)`

output

```
x^2*((2*d^3*((6*d^2*((A*b^10 + 10*B*a*b^9)/e^4 - (4*B*b^10*d)/e^5))/e^2 -
(4*d*((4*d*((A*b^10 + 10*B*a*b^9)/e^4 - (4*B*b^10*d)/e^5))/e - (5*a*b^8*(2
*A*b + 9*B*a))/e^4 + (6*B*b^10*d^2)/e^6))/e - (15*a^2*b^7*(3*A*b + 8*B*a))
/e^4 + (4*B*b^10*d^3)/e^7))/e^3 - (3*d^2*((4*d*((6*d^2*((A*b^10 + 10*B*a*b
^9)/e^4 - (4*B*b^10*d)/e^5))/e^2 - (4*d*((4*d*((A*b^10 + 10*B*a*b^9)/e^4 -
(4*B*b^10*d)/e^5))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^4 + (6*B*b^10*d^2)/e^6
))/e - (15*a^2*b^7*(3*A*b + 8*B*a))/e^4 + (4*B*b^10*d^3)/e^7))/e - (4*d^3*
((A*b^10 + 10*B*a*b^9)/e^4 - (4*B*b^10*d)/e^5))/e^3 + (6*d^2*((4*d*((A*b^1
0 + 10*B*a*b^9)/e^4 - (4*B*b^10*d)/e^5))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^4
+ (6*B*b^10*d^2)/e^6))/e^2 + (30*a^3*b^6*(4*A*b + 7*B*a))/e^4 - (B*b^10*d
^4)/e^8))/e^2 - (2*d*((6*d^2*((6*d^2*((A*b^10 + 10*B*a*b^9)/e^4 - (4*B*b^1
0*d)/e^5))/e^2 - (4*d*((4*d*((A*b^10 + 10*B*a*b^9)/e^4 - (4*B*b^10*d)/e^5)
)/e - (5*a*b^8*(2*A*b + 9*B*a))/e^4 + (6*B*b^10*d^2)/e^6))/e - (15*a^2*b^7
*(3*A*b + 8*B*a))/e^4 + (4*B*b^10*d^3)/e^7))/e^2 - (4*d*((4*d*((6*d^2*((A*
b^10 + 10*B*a*b^9)/e^4 - (4*B*b^10*d)/e^5))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^4 + (6*
B*b^10*d^2)/e^6))/e - (15*a^2*b^7*(3*A*b + 8*B*a))/e^4 + (4*B*b^10*d^3)/e^
7))/e - (4*d^3*((A*b^10 + 10*B*a*b^9)/e^4 - (4*B*b^10*d)/e^5))/e^3 + (6*d^
2*((4*d*((A*b^10 + 10*B*a*b^9)/e^4 - (4*B*b^10*d)/e^5))/e - (5*a*b^8*(2*A*
b + 9*B*a))/e^4 + (6*B*b^10*d^2)/e^6))/e^2 + (30*a^3*b^6*(4*A*b + 7*B*a...
```

Reduce [B] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 1786, normalized size of antiderivative = 4.01

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^4} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^4,x)`

output

```
(27720*log(d + e*x)*a**8*b**3*d**4*e**8 + 83160*log(d + e*x)*a**8*b**3*d**
3*e**9*x + 83160*log(d + e*x)*a**8*b**3*d**2*e**10*x**2 + 27720*log(d + e*
x)*a**8*b**3*d*e**11*x**3 - 221760*log(d + e*x)*a**7*b**4*d**5*e**7 - 6652
80*log(d + e*x)*a**7*b**4*d**4*e**8*x - 665280*log(d + e*x)*a**7*b**4*d**3
*e**9*x**2 - 221760*log(d + e*x)*a**7*b**4*d**2*e**10*x**3 + 776160*log(d
+ e*x)*a**6*b**5*d**6*e**6 + 2328480*log(d + e*x)*a**6*b**5*d**5*e**7*x +
2328480*log(d + e*x)*a**6*b**5*d**4*e**8*x**2 + 776160*log(d + e*x)*a**6*b
**5*d**3*e**9*x**3 - 1552320*log(d + e*x)*a**5*b**6*d**7*e**5 - 4656960*lo
g(d + e*x)*a**5*b**6*d**6*e**6*x - 4656960*log(d + e*x)*a**5*b**6*d**5*e**
7*x**2 - 1552320*log(d + e*x)*a**5*b**6*d**4*e**8*x**3 + 1940400*log(d + e
*x)*a**4*b**7*d**8*e**4 + 5821200*log(d + e*x)*a**4*b**7*d**7*e**5*x + 582
1200*log(d + e*x)*a**4*b**7*d**6*e**6*x**2 + 1940400*log(d + e*x)*a**4*b**
7*d**5*e**7*x**3 - 1552320*log(d + e*x)*a**3*b**8*d**9*e**3 - 4656960*log(
d + e*x)*a**3*b**8*d**8*e**4*x - 4656960*log(d + e*x)*a**3*b**8*d**7*e**5*
x**2 - 1552320*log(d + e*x)*a**3*b**8*d**6*e**6*x**3 + 776160*log(d + e*x)
*a**2*b**9*d**10*e**2 + 2328480*log(d + e*x)*a**2*b**9*d**9*e**3*x + 23284
80*log(d + e*x)*a**2*b**9*d**8*e**4*x**2 + 776160*log(d + e*x)*a**2*b**9*d
**7*e**5*x**3 - 221760*log(d + e*x)*a*b**10*d**11*e - 665280*log(d + e*x)*
a*b**10*d**10*e**2*x - 665280*log(d + e*x)*a*b**10*d**9*e**3*x**2 - 221760
*log(d + e*x)*a*b**10*d**8*e**4*x**3 + 27720*log(d + e*x)*b**11*d**12 +...
```

3.83 $\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^5} dx$

Optimal result	853
Mathematica [A] (verified)	854
Rubi [A] (verified)	855
Maple [B] (verified)	856
Fricas [B] (verification not implemented)	857
Sympy [F(-1)]	858
Maxima [B] (verification not implemented)	859
Giac [B] (verification not implemented)	860
Mupad [B] (verification not implemented)	861
Reduce [B] (verification not implemented)	861

Optimal result

Integrand size = 20, antiderivative size = 444

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^5} dx = \frac{42b^4(bd-ae)^5(11bBd-6Abe-5aBe)x}{e^{11}} + \frac{(bd-ae)^{10}(Bd-Ae)}{4e^{12}(d+ex)^4} - \frac{(bd-ae)^9(11bBd-10Abe-aBe)}{3e^{12}(d+ex)^3} + \frac{5b(bd-ae)^8(11bBd-9Abe-2aBe)}{2e^{12}(d+ex)^2} - \frac{15b^2(bd-ae)^7(11bBd-8Abe-3aBe)}{e^{12}(d+ex)} - \frac{21b^5(bd-ae)^4(11bBd-5Abe-6aBe)(d+ex)^2}{e^{12}} + \frac{10b^6(bd-ae)^3(11bBd-4Abe-7aBe)(d+ex)^3}{e^{12}} - \frac{15b^7(bd-ae)^2(11bBd-3Abe-8aBe)(d+ex)^4}{4e^{12}} + \frac{b^8(bd-ae)(11bBd-2Abe-9aBe)(d+ex)^5}{e^{12}} - \frac{b^9(11bBd-Abe-10aBe)(d+ex)^6}{6e^{12}} + \frac{b^{10}B(d+ex)^7}{7e^{12}} - \frac{30b^3(bd-ae)^6(11bBd-7Abe-4aBe)\log(d+ex)}{e^{12}}$$

output

```

42*b^4*(-a*e+b*d)^5*(-6*A*b*e-5*B*a*e+11*B*b*d)*x/e^11+1/4*(-a*e+b*d)^10*(
-A*e+B*d)/e^12/(e*x+d)^4-1/3*(-a*e+b*d)^9*(-10*A*b*e-B*a*e+11*B*b*d)/e^12/
(e*x+d)^3+5/2*b*(-a*e+b*d)^8*(-9*A*b*e-2*B*a*e+11*B*b*d)/e^12/(e*x+d)^2-15
*b^2*(-a*e+b*d)^7*(-8*A*b*e-3*B*a*e+11*B*b*d)/e^12/(e*x+d)-21*b^5*(-a*e+b*
d)^4*(-5*A*b*e-6*B*a*e+11*B*b*d)*(e*x+d)^2/e^12+10*b^6*(-a*e+b*d)^3*(-4*A*
b*e-7*B*a*e+11*B*b*d)*(e*x+d)^3/e^12-15/4*b^7*(-a*e+b*d)^2*(-3*A*b*e-8*B*a
*e+11*B*b*d)*(e*x+d)^4/e^12+b^8*(-a*e+b*d)*(-2*A*b*e-9*B*a*e+11*B*b*d)*(e*
x+d)^5/e^12-1/6*b^9*(-A*b*e-10*B*a*e+11*B*b*d)*(e*x+d)^6/e^12+1/7*b^10*B*(
e*x+d)^7/e^12-30*b^3*(-a*e+b*d)^6*(-7*A*b*e-4*B*a*e+11*B*b*d)*ln(e*x+d)/e^
12

```

Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 686, normalized size of antiderivative = 1.55

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^5} dx$$

$$= \frac{-84b^4e(-210a^6Be^6 + 140ab^5d^4e(9Bd - 5Ae) - 42b^6d^5(5Bd - 3Ae) + 600a^3b^3d^2e^3(7Bd - 3Ae) - 1575a^2b^4d^3e^2(2Bd - Ae) - 1050a^4b^2d^2e^4(3Bd - Ae) - 252a^5b^5e^5(-5Bd + Ae))*x + 42b^5e^2(252a^5B^5e^5 - 14b^5d^4(9Bd - 5Ae) - 225a^2b^3d^2e^2(7Bd - 3Ae) + 350a^4b^4d^3e(2Bd - Ae) + 600a^3b^2d^2e^3(3Bd - Ae) + 210a^4b^4e^4(-5Bd + Ae))*x^2 - 140b^6e^3*(-42a^4B^4e^4 + 10a^4b^3d^2e(7Bd - 3Ae) - 45a^2b^2d^2e^2(3Bd - Ae) - 24a^3b^3e^3(-5Bd + Ae) + 7b^4d^3(-2Bd + Ae))*x^3 + 105b^7e^4(24a^3B^3e^3 + 10a^4b^2d^2e(3Bd - Ae) + 9a^2b^5e^2(-5Bd + Ae) + b^3d^2(-7Bd + 3Ae))*x^4 - 84b^8e^5(-9a^2B^2e^2 - 2a^2b^5e(-5Bd + Ae) + b^2d^2(-3Bd + Ae))*x^5 + 14b^9e^6(-5b^4Bd + Ab^5e + 10a^2B^2e)*x^6 + 12b^10B^2e^7x^7 + (21*(b*d - a*e)^10*(B*d - A*e))/(d + e*x)^4 - (28*(b*d - a*e)^9*(11*b*B*d - 10*A*b*e - a*B^2e))/(d + e*x)^3 + (210*b*(b*d - a*e)^8*(11*b*B*d - 9*A*b*e - 2*a*B^2e))/(d + e*x)^2 - (1260*b^2*(b*d - a*e)^7*(11*b*B*d - 8*A*b*e - 3*a*B^2e))/(d + e*x) - 2520*b^3*(b*d - a*e)^6*(11*b*B*d - 7*A*b*e - 4*a*B^2e)*Log[d + e*x]/(84*e^12)$$

input

```
Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^5,x]
```

output

```

(-84*b^4*e*(-210*a^6*B*e^6 + 140*a*b^5*d^4*e*(9*B*d - 5*A*e) - 42*b^6*d^5*
(5*B*d - 3*A*e) + 600*a^3*b^3*d^2*e^3*(7*B*d - 3*A*e) - 1575*a^2*b^4*d^3*e
^2*(2*B*d - A*e) - 1050*a^4*b^2*d^2*e^4*(3*B*d - A*e) - 252*a^5*b^5*e^5*(-5*B*
d + A*e))*x + 42*b^5*e^2*(252*a^5*B^5e^5 - 14*b^5*d^4*(9*B*d - 5*A*e) - 225
*a^2*b^3*d^2*e^2*(7*B*d - 3*A*e) + 350*a^4*b^4*d^3*e*(2*B*d - A*e) + 600*a^3
*b^2*d^2*e^3*(3*B*d - A*e) + 210*a^4*b^4*e^4*(-5*B*d + A*e))*x^2 - 140*b^6*e^3
*(-42*a^4*B^4e^4 + 10*a^4*b^3*d^2*e*(7*B*d - 3*A*e) - 45*a^2*b^2*d^2*e^2*(3*B*d
- A*e) - 24*a^3*b^3e^3*(-5*B*d + A*e) + 7*b^4*d^3*(-2*B*d + A*e))*x^3 + 10
5*b^7*e^4*(24*a^3*B^3e^3 + 10*a^4*b^2*d^2*e*(3*B*d - A*e) + 9*a^2*b^5e^2*(-5*B*d
+ A*e) + b^3*d^2*(-7*B*d + 3*A*e))*x^4 - 84*b^8*e^5*(-9*a^2*B^2e^2 - 2*a^2b
^5e*(-5*B*d + A*e) + b^2*d^2*(-3*B*d + A*e))*x^5 + 14*b^9*e^6*(-5*b^4*B*d + A*b
^5e + 10*a^2*B^2e)*x^6 + 12*b^10*B^2e^7*x^7 + (21*(b*d - a*e)^10*(B*d - A*e))/(
d + e*x)^4 - (28*(b*d - a*e)^9*(11*b*B*d - 10*A*b*e - a*B^2e))/(d + e*x)^3
+ (210*b*(b*d - a*e)^8*(11*b*B*d - 9*A*b*e - 2*a*B^2e))/(d + e*x)^2 - (1260
*b^2*(b*d - a*e)^7*(11*b*B*d - 8*A*b*e - 3*a*B^2e))/(d + e*x) - 2520*b^3*(b
*d - a*e)^6*(11*b*B*d - 7*A*b*e - 4*a*B^2e)*Log[d + e*x]/(84*e^12)

```

Rubi [A] (verified)

Time = 1.44 (sec) , antiderivative size = 444, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^5} dx$$

↓ 86

$$\int \left(\frac{b^9(d + ex)^5(10aBe + Abe - 11bBd)}{e^{11}} - \frac{5b^8(d + ex)^4(bd - ae)(9aBe + 2Abe - 11bBd)}{e^{11}} + \frac{15b^7(d + ex)^3(bd - ae)^2(8aBe + Abe - 11bBd)}{e^{11}} - \frac{b^9(d + ex)^6(-10aBe - Abe + 11bBd)}{6e^{12}} + \frac{b^8(d + ex)^5(bd - ae)(-9aBe - 2Abe + 11bBd)}{e^{12}} - \frac{15b^7(d + ex)^4(bd - ae)^2(-8aBe - 3Abe + 11bBd)}{4e^{12}} + \frac{10b^6(d + ex)^3(bd - ae)^3(-7aBe - 4Abe + 11bBd)}{e^{12}} - \frac{21b^5(d + ex)^2(bd - ae)^4(-6aBe - 5Abe + 11bBd)}{e^{12}} + \frac{42b^4x(bd - ae)^5(-5aBe - 6Abe + 11bBd)}{e^{11}} - \frac{30b^3(bd - ae)^6 \log(d + ex)(-4aBe - 7Abe + 11bBd)}{e^{12}} - \frac{15b^2(bd - ae)^7(-3aBe - 8Abe + 11bBd)}{e^{12}(d + ex)} + \frac{5b(bd - ae)^8(-2aBe - 9Abe + 11bBd)}{2e^{12}(d + ex)^2} - \frac{(bd - ae)^9(-aBe - 10Abe + 11bBd)}{3e^{12}(d + ex)^3} + \frac{(bd - ae)^{10}(Bd - Ae)}{4e^{12}(d + ex)^4} + \frac{b^{10}B(d + ex)^7}{7e^{12}} \right)$$

↓ 2009

input

```
Int[((a + b*x)^10*(A + B*x))/(d + e*x)^5,x]
```

output

$$\begin{aligned} & (42*b^4*(b*d - a*e)^5*(11*b*B*d - 6*A*b*e - 5*a*B*e)*x)/e^{11} + ((b*d - a*e)^{10}*(B*d - A*e))/(4*e^{12}*(d + e*x)^4) - ((b*d - a*e)^9*(11*b*B*d - 10*A*b*e - a*B*e))/(3*e^{12}*(d + e*x)^3) + (5*b*(b*d - a*e)^8*(11*b*B*d - 9*A*b*e - 2*a*B*e))/(2*e^{12}*(d + e*x)^2) - (15*b^2*(b*d - a*e)^7*(11*b*B*d - 8*A*b*e - 3*a*B*e))/(e^{12}*(d + e*x)) - (21*b^5*(b*d - a*e)^4*(11*b*B*d - 5*A*b*e - 6*a*B*e)*(d + e*x)^2)/e^{12} + (10*b^6*(b*d - a*e)^3*(11*b*B*d - 4*A*b*e - 7*a*B*e)*(d + e*x)^3)/e^{12} - (15*b^7*(b*d - a*e)^2*(11*b*B*d - 3*A*b*e - 8*a*B*e)*(d + e*x)^4)/(4*e^{12}) + (b^8*(b*d - a*e)*(11*b*B*d - 2*A*b*e - 9*a*B*e)*(d + e*x)^5)/e^{12} - (b^9*(11*b*B*d - A*b*e - 10*a*B*e)*(d + e*x)^6)/(6*e^{12}) + (b^{10}*B*(d + e*x)^7)/(7*e^{12}) - (30*b^3*(b*d - a*e)^6*(11*b*B*d - 7*A*b*e - 4*a*B*e)*Log[d + e*x])/e^{12} \end{aligned}$$

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1909 vs. $2(432) = 864$.

Time = 0.24 (sec) , antiderivative size = 1910, normalized size of antiderivative = 4.30

method	result	size
norman	Expression too large to display	1910
default	Expression too large to display	2034
risch	Expression too large to display	2127
parallelsch	Expression too large to display	3635

input

```
int((b*x+a)^10*(B*x+A)/(e*x+d)^5,x,method=_RETURNVERBOSE)
```

output

```
(b^5*(105*A*a^4*b*e^5-140*A*a^3*b^2*d*e^4+105*A*a^2*b^3*d^2*e^3-42*A*a*b^4*d^3*e^2+7*A*b^5*d^4*e+126*B*a^5*e^5-245*B*a^4*b*d*e^4+280*B*a^3*b^2*d^2*e^3-189*B*a^2*b^3*d^3*e^2+70*B*a*b^4*d^4*e-11*B*b^5*d^5)/e^6*x^6-1/12*(3*A*a^10*e^11+10*A*a^9*b*d*e^10+45*A*a^8*b^2*d^2*e^9+360*A*a^7*b^3*d^3*e^8-5250*A*a^6*b^4*d^4*e^7+31500*A*a^5*b^5*d^5*e^6-78750*A*a^4*b^6*d^6*e^5+105000*A*a^3*b^7*d^7*e^4-78750*A*a^2*b^8*d^8*e^3+31500*A*a*b^9*d^9*e^2-5250*A*b^10*d^10*e+B*a^10*d*e^10+10*B*a^9*b*d^2*e^9+135*B*a^8*b^2*d^3*e^8-3000*B*a^7*b^3*d^4*e^7+26250*B*a^6*b^4*d^5*e^6-94500*B*a^5*b^5*d^6*e^5+183750*B*a^4*b^6*d^7*e^4-210000*B*a^3*b^7*d^8*e^3+141750*B*a^2*b^8*d^9*e^2-52500*B*a*b^9*d^10*e+8250*B*b^10*d^11)/e^12-(120*A*a^7*b^3*e^8-840*A*a^6*b^4*d*e^7+5040*A*a^5*b^5*d^2*e^6-12600*A*a^4*b^6*d^3*e^5+16800*A*a^3*b^7*d^4*e^4-12600*A*a^2*b^8*d^5*e^3+5040*A*a*b^9*d^6*e^2-840*A*b^10*d^7*e+45*B*a^8*b^2*e^8-480*B*a^7*b^3*d*e^7+4200*B*a^6*b^4*d^2*e^6-15120*B*a^5*b^5*d^3*e^5+29400*B*a^4*b^6*d^4*e^4-33600*B*a^3*b^7*d^5*e^3+22680*B*a^2*b^8*d^6*e^2-8400*B*a*b^9*d^7*e+1320*B*b^10*d^8)/e^9*x^3-1/2*(45*A*a^8*b^2*e^9+360*A*a^7*b^3*d*e^8-3780*A*a^6*b^4*d^2*e^7+22680*A*a^5*b^5*d^3*e^6-56700*A*a^4*b^6*d^4*e^5+75600*A*a^3*b^7*d^5*e^4-56700*A*a^2*b^8*d^6*e^3+22680*A*a*b^9*d^7*e^2-3780*A*b^10*d^8*e+10*B*a^9*b*e^9+135*B*a^8*b^2*d*e^8-2160*B*a^7*b^3*d^2*e^7+18900*B*a^6*b^4*d^3*e^6-68040*B*a^5*b^5*d^4*e^5+132300*B*a^4*b^6*d^5*e^4-151200*B*a^3*b^7*d^6*e^3+102060*B*a^2*b^8*d^7*e^2-37800*B*a*b^9*d^8*e+5940...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2808 vs. $2(432) = 864$.

Time = 0.14 (sec) , antiderivative size = 2808, normalized size of antiderivative = 6.32

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^5} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)/(e*x+d)^5,x, algorithm="fricas")
```

output

```

1/84*(12*B*b^10*e^11*x^11 - 11837*B*b^10*d^11 - 21*A*a^10*e^11 + 8449*(10*
B*a*b^9 + A*b^10)*d^10*e - 28875*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 + 55965
*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 - 66990*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d
^7*e^4 + 50274*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 - 22638*(5*B*a^6*b^4 +
6*A*a^5*b^5)*d^5*e^6 + 5250*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 - 315*(3*B
*a^8*b^2 + 8*A*a^7*b^3)*d^3*e^8 - 35*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 - 7
*(B*a^10 + 10*A*a^9*b)*d*e^10 - 2*(11*B*b^10*d*e^10 - 7*(10*B*a*b^9 + A*b^
10)*e^11)*x^10 + 4*(11*B*b^10*d^2*e^9 - 7*(10*B*a*b^9 + A*b^10)*d*e^10 + 2
1*(9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 - 9*(11*B*b^10*d^3*e^8 - 7*(10*B*a*b
^9 + A*b^10)*d^2*e^9 + 21*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 - 35*(8*B*a^3*b
^7 + 3*A*a^2*b^8)*e^11)*x^8 + 24*(11*B*b^10*d^4*e^7 - 7*(10*B*a*b^9 + A*b^
10)*d^3*e^8 + 21*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 - 35*(8*B*a^3*b^7 + 3*A
*a^2*b^8)*d*e^10 + 35*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7 - 84*(11*B*b^1
0*d^5*e^6 - 7*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 21*(9*B*a^2*b^8 + 2*A*a*b^9)
*d^3*e^8 - 35*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 35*(7*B*a^4*b^6 + 4*A*
a^3*b^7)*d*e^10 - 21*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 + 504*(11*B*b^1
0*d^6*e^5 - 7*(10*B*a*b^9 + A*b^10)*d^5*e^6 + 21*(9*B*a^2*b^8 + 2*A*a*b^9)
*d^4*e^7 - 35*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 + 35*(7*B*a^4*b^6 + 4*A*
a^3*b^7)*d^2*e^9 - 21*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10 + 7*(5*B*a^6*b^4
+ 6*A*a^5*b^5)*e^11)*x^5 + 7*(6559*B*b^10*d^7*e^4 - 4043*(10*B*a*b^9 + ...

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^5} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**10*(B*x+A)/(e*x+d)**5,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1846 vs. $2(432) = 864$.

Time = 0.12 (sec) , antiderivative size = 1846, normalized size of antiderivative = 4.16

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^5} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^5,x, algorithm="maxima")`

output

```
-1/12*(1691*B*b^10*d^11 + 3*A*a^10*e^11 - 1207*(10*B*a*b^9 + A*b^10)*d^10*
e + 4125*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 - 7995*(8*B*a^3*b^7 + 3*A*a^2*b
^8)*d^8*e^3 + 9570*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e^4 - 7182*(6*B*a^5*b^5
+ 5*A*a^4*b^6)*d^6*e^5 + 3234*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^5*e^6 - 750*(
4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 45*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^3*e^
8 + 5*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 + (B*a^10 + 10*A*a^9*b)*d*e^10 + 1
80*(11*B*b^10*d^8*e^3 - 8*(10*B*a*b^9 + A*b^10)*d^7*e^4 + 28*(9*B*a^2*b^8
+ 2*A*a*b^9)*d^6*e^5 - 56*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^5*e^6 + 70*(7*B*a^
4*b^6 + 4*A*a^3*b^7)*d^4*e^7 - 56*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^3*e^8 + 28
*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^2*e^9 - 8*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d*e^1
0 + (3*B*a^8*b^2 + 8*A*a^7*b^3)*e^11)*x^3 + 30*(187*B*b^10*d^9*e^2 - 135*(
10*B*a*b^9 + A*b^10)*d^8*e^3 + 468*(9*B*a^2*b^8 + 2*A*a*b^9)*d^7*e^4 - 924
*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^6*e^5 + 1134*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^
5*e^6 - 882*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^4*e^7 + 420*(5*B*a^6*b^4 + 6*A*a
^5*b^5)*d^3*e^8 - 108*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^2*e^9 + 9*(3*B*a^8*b^2
+ 8*A*a^7*b^3)*d*e^10 + (2*B*a^9*b + 9*A*a^8*b^2)*e^11)*x^2 + 4*(1331*B*b
^10*d^10*e - 955*(10*B*a*b^9 + A*b^10)*d^9*e^2 + 3285*(9*B*a^2*b^8 + 2*A*a
*b^9)*d^8*e^3 - 6420*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^7*e^4 + 7770*(7*B*a^4*b
^6 + 4*A*a^3*b^7)*d^6*e^5 - 5922*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^5*e^6 + 273
0*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^4*e^7 - 660*(4*B*a^7*b^3 + 7*A*a^6*b^4)...
```


Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2571 vs. $2(432) = 864$.

Time = 0.14 (sec) , antiderivative size = 2571, normalized size of antiderivative = 5.79

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^5} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^5,x, algorithm="giac")`

output

```
1/84*(12*B*b^10 - 14*(11*B*b^10*d*e - 10*B*a*b^9*e^2 - A*b^10*e^2)/((e*x +
d)*e) + 84*(11*B*b^10*d^2*e^2 - 20*B*a*b^9*d*e^3 - 2*A*b^10*d*e^3 + 9*B*a
^2*b^8*e^4 + 2*A*a*b^9*e^4)/((e*x + d)^2*e^2) - 315*(11*B*b^10*d^3*e^3 - 3
0*B*a*b^9*d^2*e^4 - 3*A*b^10*d^2*e^4 + 27*B*a^2*b^8*d*e^5 + 6*A*a*b^9*d*e^
5 - 8*B*a^3*b^7*e^6 - 3*A*a^2*b^8*e^6)/((e*x + d)^3*e^3) + 840*(11*B*b^10*
d^4*e^4 - 40*B*a*b^9*d^3*e^5 - 4*A*b^10*d^3*e^5 + 54*B*a^2*b^8*d^2*e^6 + 1
2*A*a*b^9*d^2*e^6 - 32*B*a^3*b^7*d*e^7 - 12*A*a^2*b^8*d*e^7 + 7*B*a^4*b^6*
e^8 + 4*A*a^3*b^7*e^8)/((e*x + d)^4*e^4) - 1764*(11*B*b^10*d^5*e^5 - 50*B*
a*b^9*d^4*e^6 - 5*A*b^10*d^4*e^6 + 90*B*a^2*b^8*d^3*e^7 + 20*A*a*b^9*d^3*e
^7 - 80*B*a^3*b^7*d^2*e^8 - 30*A*a^2*b^8*d^2*e^8 + 35*B*a^4*b^6*d*e^9 + 20
*A*a^3*b^7*d*e^9 - 6*B*a^5*b^5*e^10 - 5*A*a^4*b^6*e^10)/((e*x + d)^5*e^5)
+ 3528*(11*B*b^10*d^6*e^6 - 60*B*a*b^9*d^5*e^7 - 6*A*b^10*d^5*e^7 + 135*B*
a^2*b^8*d^4*e^8 + 30*A*a*b^9*d^4*e^8 - 160*B*a^3*b^7*d^3*e^9 - 60*A*a^2*b^
8*d^3*e^9 + 105*B*a^4*b^6*d^2*e^10 + 60*A*a^3*b^7*d^2*e^10 - 36*B*a^5*b^5*
d*e^11 - 30*A*a^4*b^6*d*e^11 + 5*B*a^6*b^4*e^12 + 6*A*a^5*b^5*e^12)/((e*x
+ d)^6*e^6))*(e*x + d)^7/e^12 + 30*(11*B*b^10*d^7 - 70*B*a*b^9*d^6*e - 7*A
*b^10*d^6*e + 189*B*a^2*b^8*d^5*e^2 + 42*A*a*b^9*d^5*e^2 - 280*B*a^3*b^7*d
^4*e^3 - 105*A*a^2*b^8*d^4*e^3 + 245*B*a^4*b^6*d^3*e^4 + 140*A*a^3*b^7*d^3
*e^4 - 126*B*a^5*b^5*d^2*e^5 - 105*A*a^4*b^6*d^2*e^5 + 35*B*a^6*b^4*d*e^6
+ 42*A*a^5*b^5*d*e^6 - 4*B*a^7*b^3*e^7 - 7*A*a^6*b^4*e^7)*log(abs(e*x + ...
```

Mupad [B] (verification not implemented)

Time = 1.16 (sec) , antiderivative size = 3655, normalized size of antiderivative = 8.23

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^5} dx = \text{Too large to display}$$

input `int(((A + B*x)*(a + b*x)^10)/(d + e*x)^5,x)`

output

```
x*((10*d^3*((10*d^2*((A*b^10 + 10*B*a*b^9)/e^5 - (5*B*b^10*d)/e^6))/e^2 -
(5*d*((5*d*((A*b^10 + 10*B*a*b^9)/e^5 - (5*B*b^10*d)/e^6))/e - (5*a*b^8*(2
*A*b + 9*B*a))/e^5 + (10*B*b^10*d^2)/e^7))/e - (15*a^2*b^7*(3*A*b + 8*B*a)
)/e^5 + (10*B*b^10*d^3)/e^8))/e^3 - (d^5*((A*b^10 + 10*B*a*b^9)/e^5 - (5*B
*b^10*d)/e^6))/e^5 - (10*d^2*((5*d*((10*d^2*((A*b^10 + 10*B*a*b^9)/e^5 - (
5*B*b^10*d)/e^6))/e^2 - (5*d*((5*d*((A*b^10 + 10*B*a*b^9)/e^5 - (5*B*b^10*
d)/e^6))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^5 + (10*B*b^10*d^2)/e^7))/e - (15
*a^2*b^7*(3*A*b + 8*B*a))/e^5 + (10*B*b^10*d^3)/e^8))/e - (10*d^3*((A*b^10
+ 10*B*a*b^9)/e^5 - (5*B*b^10*d)/e^6))/e^3 + (10*d^2*((5*d*((A*b^10 + 10*
B*a*b^9)/e^5 - (5*B*b^10*d)/e^6))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^5 + (10*
B*b^10*d^2)/e^7))/e^2 + (30*a^3*b^6*(4*A*b + 7*B*a))/e^5 - (5*B*b^10*d^4)/
e^9))/e^2 + (5*d*((5*d*((5*d*((10*d^2*((A*b^10 + 10*B*a*b^9)/e^5 - (5*B*b^
10*d)/e^6))/e^2 - (5*d*((5*d*((A*b^10 + 10*B*a*b^9)/e^5 - (5*B*b^10*d)/e^6
))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^5 + (10*B*b^10*d^2)/e^7))/e - (15*a^2*b
^7*(3*A*b + 8*B*a))/e^5 + (10*B*b^10*d^3)/e^8))/e - (10*d^3*((A*b^10 + 10*
B*a*b^9)/e^5 - (5*B*b^10*d)/e^6))/e^3 + (10*d^2*((5*d*((A*b^10 + 10*B*a*b^
9)/e^5 - (5*B*b^10*d)/e^6))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^5 + (10*B*b^10
*d^2)/e^7))/e^2 + (30*a^3*b^6*(4*A*b + 7*B*a))/e^5 - (5*B*b^10*d^4)/e^9))/
e - (10*d^2*((10*d^2*((A*b^10 + 10*B*a*b^9)/e^5 - (5*B*b^10*d)/e^6))/e^2 -
(5*d*((5*d*((A*b^10 + 10*B*a*b^9)/e^5 - (5*B*b^10*d)/e^6))/e - (5*a*b^...
```

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 1889, normalized size of antiderivative = 4.25

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^5} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^5,x)`

output

```
(27720*log(d + e*x)*a**7*b**4*d**5*e**7 + 110880*log(d + e*x)*a**7*b**4*d*
**4*e**8*x + 166320*log(d + e*x)*a**7*b**4*d**3*e**9*x**2 + 110880*log(d +
e*x)*a**7*b**4*d**2*e**10*x**3 + 27720*log(d + e*x)*a**7*b**4*d*e**11*x**4
- 194040*log(d + e*x)*a**6*b**5*d**6*e**6 - 776160*log(d + e*x)*a**6*b**5
*d**5*e**7*x - 1164240*log(d + e*x)*a**6*b**5*d**4*e**8*x**2 - 776160*log(
d + e*x)*a**6*b**5*d**3*e**9*x**3 - 194040*log(d + e*x)*a**6*b**5*d**2*e**
10*x**4 + 582120*log(d + e*x)*a**5*b**6*d**7*e**5 + 2328480*log(d + e*x)*a
**5*b**6*d**6*e**6*x + 3492720*log(d + e*x)*a**5*b**6*d**5*e**7*x**2 + 232
8480*log(d + e*x)*a**5*b**6*d**4*e**8*x**3 + 582120*log(d + e*x)*a**5*b**6
*d**3*e**9*x**4 - 970200*log(d + e*x)*a**4*b**7*d**8*e**4 - 3880800*log(d
+ e*x)*a**4*b**7*d**7*e**5*x - 5821200*log(d + e*x)*a**4*b**7*d**6*e**6*x*
*2 - 3880800*log(d + e*x)*a**4*b**7*d**5*e**7*x**3 - 970200*log(d + e*x)*a
**4*b**7*d**4*e**8*x**4 + 970200*log(d + e*x)*a**3*b**8*d**9*e**3 + 388080
0*log(d + e*x)*a**3*b**8*d**8*e**4*x + 5821200*log(d + e*x)*a**3*b**8*d**7
*e**5*x**2 + 3880800*log(d + e*x)*a**3*b**8*d**6*e**6*x**3 + 970200*log(d
+ e*x)*a**3*b**8*d**5*e**7*x**4 - 582120*log(d + e*x)*a**2*b**9*d**10*e**2
- 2328480*log(d + e*x)*a**2*b**9*d**9*e**3*x - 3492720*log(d + e*x)*a**2*
b**9*d**8*e**4*x**2 - 2328480*log(d + e*x)*a**2*b**9*d**7*e**5*x**3 - 5821
20*log(d + e*x)*a**2*b**9*d**6*e**6*x**4 + 194040*log(d + e*x)*a*b**10*d**
11*e + 776160*log(d + e*x)*a*b**10*d**10*e**2*x + 1164240*log(d + e*x)*...
```

3.84 $\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^6} dx$

Optimal result	863
Mathematica [A] (verified)	864
Rubi [A] (verified)	865
Maple [B] (verified)	866
Fricas [B] (verification not implemented)	867
Sympy [F(-1)]	868
Maxima [B] (verification not implemented)	869
Giac [B] (verification not implemented)	870
Mupad [B] (verification not implemented)	871
Reduce [B] (verification not implemented)	871

Optimal result

Integrand size = 20, antiderivative size = 447

$$\begin{aligned}
 \int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^6} dx = & -\frac{42b^5(bd-ae)^4(11bBd-5Abe-6aBe)x}{e^{11}} \\
 & + \frac{(bd-ae)^{10}(Bd-Ae)}{5e^{12}(d+ex)^5} \\
 & - \frac{(bd-ae)^9(11bBd-10Abe-aBe)}{4e^{12}(d+ex)^4} \\
 & + \frac{5b(bd-ae)^8(11bBd-9Abe-2aBe)}{3e^{12}(d+ex)^3} \\
 & - \frac{15b^2(bd-ae)^7(11bBd-8Abe-3aBe)}{2e^{12}(d+ex)^2} \\
 & + \frac{30b^3(bd-ae)^6(11bBd-7Abe-4aBe)}{e^{12}(d+ex)} \\
 & + \frac{15b^6(bd-ae)^3(11bBd-4Abe-7aBe)(d+ex)^2}{e^{12}} \\
 & - \frac{5b^7(bd-ae)^2(11bBd-3Abe-8aBe)(d+ex)^3}{e^{12}} \\
 & + \frac{5b^8(bd-ae)(11bBd-2Abe-9aBe)(d+ex)^4}{4e^{12}} \\
 & - \frac{b^9(11bBd-Abe-10aBe)(d+ex)^5}{5e^{12}} + \frac{b^{10}B(d+ex)^6}{6e^{12}} \\
 & + \frac{42b^4(bd-ae)^5(11bBd-6Abe-5aBe)\log(d+ex)}{e^{12}}
 \end{aligned}$$

output

```
-42*b^5*(-a*e+b*d)^4*(-5*A*b*e-6*B*a*e+11*B*b*d)*x/e^11+1/5*(-a*e+b*d)^10*
(-A*e+B*d)/e^12/(e*x+d)^5-1/4*(-a*e+b*d)^9*(-10*A*b*e-B*a*e+11*B*b*d)/e^12
/(e*x+d)^4+5/3*b*(-a*e+b*d)^8*(-9*A*b*e-2*B*a*e+11*B*b*d)/e^12/(e*x+d)^3-1
5/2*b^2*(-a*e+b*d)^7*(-8*A*b*e-3*B*a*e+11*B*b*d)/e^12/(e*x+d)^2+30*b^3*(-a
*e+b*d)^6*(-7*A*b*e-4*B*a*e+11*B*b*d)/e^12/(e*x+d)+15*b^6*(-a*e+b*d)^3*(-4
*A*b*e-7*B*a*e+11*B*b*d)*(e*x+d)^2/e^12-5*b^7*(-a*e+b*d)^2*(-3*A*b*e-8*B*a
*e+11*B*b*d)*(e*x+d)^3/e^12+5/4*b^8*(-a*e+b*d)*(-2*A*b*e-9*B*a*e+11*B*b*d)
*(e*x+d)^4/e^12-1/5*b^9*(-A*b*e-10*B*a*e+11*B*b*d)*(e*x+d)^5/e^12+1/6*b^10
*B*(e*x+d)^6/e^12+42*b^4*(-a*e+b*d)^5*(-6*A*b*e-5*B*a*e+11*B*b*d)*ln(e*x+d
)/e^12
```

Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 587, normalized size of antiderivative = 1.31

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^6} dx$$

$$= \frac{60b^5e(252a^5Be^5 + 140ab^4d^3e(9Bd - 4Ae) - 315a^2b^3d^2e^2(8Bd - 3Ae) + 360a^3b^2de^3(7Bd - 2Ae) - 126b^5d^4(2Bd - Ae) + 210a^4b^4e^4(-6Bd + Ae))x - 30b^6e^2(-210a^4B^2e^4 - 14b^4d^3(9Bd - 4Ae) + 70ab^3d^2e^2(8Bd - 3Ae) - 135a^2b^2d^2e^2(7Bd - 2Ae) - 120a^3b^3e^3(-6Bd + Ae))x^2 + 20b^7e^3(120a^3B^2e^3 - 7b^3d^2(8Bd - 3Ae) + 30ab^2d^2e^2(7Bd - 2Ae) + 45a^2b^2e^2(-6Bd + Ae))x^3 - 15b^8e^4(-45a^2B^2e^2 - 10ab^2e^2(-6Bd + Ae) + 3b^2d^2(-7Bd + 2Ae))x^4 + 12b^9e^5(-6b^2Bd + Ab^2e + 10aB^2e)x^5 + 10b^10B^2e^6x^6 + (12(bd - ae)^{10}(Bd - Ae))/(d + ex)^5 - (15(bd - ae)^9(11bBd - 10Ab^2e - aB^2e))/(d + ex)^4 + (100b(bd - ae)^8(11bBd - 9Ab^2e - 2aB^2e))/(d + ex)^3 - (450b^2(bd - ae)^7(11bBd - 8Ab^2e - 3aB^2e))/(d + ex)^2 + (1800b^3(bd - ae)^6(11bBd - 7Ab^2e - 4aB^2e))/(d + ex) + 2520b^4(bd - ae)^5(11bBd - 6Ab^2e - 5aB^2e)*Log[d + ex]}{(60e^{12})}$$

input

```
Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^6,x]
```

output

```
(60*b^5*e*(252*a^5*B*e^5 + 140*a*b^4*d^3*e*(9*B*d - 4*A*e) - 315*a^2*b^3*d
^2*e^2*(8*B*d - 3*A*e) + 360*a^3*b^2*d*e^3*(7*B*d - 2*A*e) - 126*b^5*d^4*(
2*B*d - A*e) + 210*a^4*b^4*e^4*(-6*B*d + A*e))*x - 30*b^6*e^2*(-210*a^4*B^2
e^4 - 14*b^4*d^3*(9*B*d - 4*A*e) + 70*a*b^3*d^2*e^2*(8*B*d - 3*A*e) - 135*a^2
b^2*d^2*e^2*(7*B*d - 2*A*e) - 120*a^3*b^3*e^3*(-6*B*d + A*e))*x^2 + 20*b^7
e^3*(120*a^3*B^2e^3 - 7*b^3*d^2*(8*B*d - 3*A*e) + 30*a*b^2*d^2*e^2*(7*B*d
- 2*A*e) + 45*a^2*b^2*e^2*(-6*B*d + A*e))*x^3 - 15*b^8*e^4*(-45*a^2*B^2e^2
- 10*a*b^2*e^2*(-6*B*d + A*e) + 3*b^2*d^2*(-7*B*d + 2*A*e))*x^4 + 12*b^9
e^5*(-6*b^2*B*d + A*b^2*e + 10*a*B^2e)*x^5 + 10*b^10*B^2e^6*x^6 + (12*(b*d
- a*e)^10*(B*d - A*e))/(d + e*x)^5 - (15*(b*d - a*e)^9*(11*b*B*d - 10*A*b
^2*e - a*B^2e))/(d + e*x)^4 + (100*b*(b*d - a*e)^8*(11*b*B*d - 9*A*b^2*e
- 2*a*B^2e))/(d + e*x)^3 - (450*b^2*(b*d - a*e)^7*(11*b*B*d - 8*A*b^2*e
- 3*a*B^2e))/(d + e*x)^2 + (1800*b^3*(b*d - a*e)^6*(11*b*B*d - 7*A*b^2*e
- 4*a*B^2e))/(d + e*x) + 2520*b^4*(b*d - a*e)^5*(11*b*B*d - 6*A*b^2*e -
5*a*B^2e)*Log[d + e*x]/(60*e^12)
```

Rubi [A] (verified)

Time = 1.41 (sec) , antiderivative size = 447, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^6} dx$$

↓ 86

$$\int \left(\frac{b^9(d + ex)^4(10aBe + Abe - 11bBd)}{e^{11}} - \frac{5b^8(d + ex)^3(bd - ae)(9aBe + 2Abe - 11bBd)}{e^{11}} + \frac{15b^7(d + ex)^2(bd - ae)^2(10aBe + Abe - 11bBd)}{e^{11}} \right) dx$$

↓ 2009

$$\begin{aligned} & - \frac{b^9(d + ex)^5(-10aBe - Abe + 11bBd)}{5e^{12}} + \frac{5b^8(d + ex)^4(bd - ae)(-9aBe - 2Abe + 11bBd)}{4e^{12}} - \\ & \frac{15b^7(d + ex)^3(bd - ae)^2(-8aBe - 3Abe + 11bBd)}{e^{12}} + \\ & \frac{15b^6(d + ex)^2(bd - ae)^3(-7aBe - 4Abe + 11bBd)}{e^{12}} - \\ & \frac{42b^5x(bd - ae)^4(-6aBe - 5Abe + 11bBd)}{e^{12}} + \\ & \frac{42b^4(bd - ae)^5 \log(d + ex)(-5aBe - 6Abe + 11bBd)}{e^{12}} + \\ & \frac{30b^3(bd - ae)^6(-4aBe - 7Abe + 11bBd)}{e^{12}(d + ex)} - \frac{15b^2(bd - ae)^7(-3aBe - 8Abe + 11bBd)}{2e^{12}(d + ex)^2} + \\ & \frac{5b(bd - ae)^8(-2aBe - 9Abe + 11bBd)}{3e^{12}(d + ex)^3} - \frac{(bd - ae)^9(-aBe - 10Abe + 11bBd)}{4e^{12}(d + ex)^4} + \\ & \frac{(bd - ae)^{10}(Bd - Ae)}{5e^{12}(d + ex)^5} + \frac{b^{10}B(d + ex)^6}{6e^{12}} \end{aligned}$$

input `Int[((a + b*x)^10*(A + B*x))/(d + e*x)^6,x]`

output

$$\begin{aligned} & (-42*b^5*(b*d - a*e)^4*(11*b*B*d - 5*A*b*e - 6*a*B*e)*x)/e^{11} + ((b*d - a* \\ & e)^{10}*(B*d - A*e))/(5*e^{12}*(d + e*x)^5) - ((b*d - a*e)^9*(11*b*B*d - 10*A* \\ & b*e - a*B*e))/(4*e^{12}*(d + e*x)^4) + (5*b*(b*d - a*e)^8*(11*b*B*d - 9*A*b* \\ & e - 2*a*B*e))/(3*e^{12}*(d + e*x)^3) - (15*b^2*(b*d - a*e)^7*(11*b*B*d - 8*A \\ & *b*e - 3*a*B*e))/(2*e^{12}*(d + e*x)^2) + (30*b^3*(b*d - a*e)^6*(11*b*B*d - \\ & 7*A*b*e - 4*a*B*e))/(e^{12}*(d + e*x)) + (15*b^6*(b*d - a*e)^3*(11*b*B*d - 4 \\ & *A*b*e - 7*a*B*e)*(d + e*x)^2)/e^{12} - (5*b^7*(b*d - a*e)^2*(11*b*B*d - 3*A \\ & *b*e - 8*a*B*e)*(d + e*x)^3)/e^{12} + (5*b^8*(b*d - a*e)*(11*b*B*d - 2*A*b*e \\ & - 9*a*B*e)*(d + e*x)^4)/(4*e^{12}) - (b^9*(11*b*B*d - A*b*e - 10*a*B*e)*(d \\ & + e*x)^5)/(5*e^{12}) + (b^{10}*B*(d + e*x)^6)/(6*e^{12}) + (42*b^4*(b*d - a*e)^5 \\ & *(11*b*B*d - 6*A*b*e - 5*a*B*e)*\text{Log}[d + e*x])/e^{12} \end{aligned}$$

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1915 vs. $2(433) = 866$.

Time = 0.24 (sec) , antiderivative size = 1916, normalized size of antiderivative = 4.29

method	result	size
norman	Expression too large to display	1916
default	Expression too large to display	2001
risch	Expression too large to display	2079
parallelsch	Expression too large to display	3743

input

```
int((b*x+a)^10*(B*x+A)/(e*x+d)^6,x,method=_RETURNVERBOSE)
```

output

```
(b^6*(60*A*a^3*b*e^4-60*A*a^2*b^2*d*e^3+30*A*a*b^3*d^2*e^2-6*A*b^4*d^3*e+1
05*B*a^4*e^4-160*B*a^3*b*d*e^3+135*B*a^2*b^2*d^2*e^2-60*B*a*b^3*d^3*e+11*B
*b^4*d^4)/e^5*x^7-1/60*(12*A*a^10*e^11+30*A*a^9*b*d*e^10+90*A*a^8*b^2*d^2*
e^9+360*A*a^7*b^3*d^3*e^8+2520*A*a^6*b^4*d^4*e^7-34524*A*a^5*b^5*d^5*e^6+1
72620*A*a^4*b^6*d^6*e^5-345240*A*a^3*b^7*d^7*e^4+345240*A*a^2*b^8*d^8*e^3-
172620*A*a*b^9*d^9*e^2+34524*A*b^10*d^10*e+3*B*a^10*d*e^10+20*B*a^9*b*d^2*
e^9+135*B*a^8*b^2*d^3*e^8+1440*B*a^7*b^3*d^4*e^7-28770*B*a^6*b^4*d^5*e^6+2
07144*B*a^5*b^5*d^6*e^5-604170*B*a^4*b^6*d^7*e^4+920640*B*a^3*b^7*d^8*e^3-
776790*B*a^2*b^8*d^9*e^2+345240*B*a*b^9*d^10*e-63294*B*b^10*d^11)/e^12-5*(
42*A*a^6*b^4*e^7-252*A*a^5*b^5*d*e^6+1260*A*a^4*b^6*d^2*e^5-2520*A*a^3*b^7
*d^3*e^4+2520*A*a^2*b^8*d^4*e^3-1260*A*a*b^9*d^5*e^2+252*A*b^10*d^6*e+24*B
*a^7*b^3*e^7-210*B*a^6*b^4*d*e^6+1512*B*a^5*b^5*d^2*e^5-4410*B*a^4*b^6*d^3
*e^4+6720*B*a^3*b^7*d^4*e^3-5670*B*a^2*b^8*d^5*e^2+2520*B*a*b^9*d^6*e-462*
B*b^10*d^7)/e^8*x^4-5/2*(24*A*a^7*b^3*e^8+168*A*a^6*b^4*d*e^7-1512*A*a^5*b
^5*d^2*e^6+7560*A*a^4*b^6*d^3*e^5-15120*A*a^3*b^7*d^4*e^4+15120*A*a^2*b^8*
d^5*e^3-7560*A*a*b^9*d^6*e^2+1512*A*b^10*d^7*e+9*B*a^8*b^2*e^8+96*B*a^7*b^
3*d*e^7-1260*B*a^6*b^4*d^2*e^6+9072*B*a^5*b^5*d^3*e^5-26460*B*a^4*b^6*d^4*
e^4+40320*B*a^3*b^7*d^5*e^3-34020*B*a^2*b^8*d^6*e^2+15120*B*a*b^9*d^7*e-27
72*B*b^10*d^8)/e^9*x^3-5/6*(18*A*a^8*b^2*e^9+72*A*a^7*b^3*d*e^8+504*A*a^6*
b^4*d^2*e^7-5544*A*a^5*b^5*d^3*e^6+27720*A*a^4*b^6*d^4*e^5-55440*A*a^3*...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2851 vs. $2(433) = 866$.

Time = 0.15 (sec) , antiderivative size = 2851, normalized size of antiderivative = 6.38

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^6} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)/(e*x+d)^6,x, algorithm="fricas")
```


output

```

1/60*(10*B*b^10*e^11*x^11 + 15797*B*b^10*d^11 - 12*A*a^10*e^11 - 9762*(10*
B*a*b^9 + A*b^10)*d^10*e + 28185*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 - 44580
*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 41310*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d
^7*e^4 - 21924*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 5754*(5*B*a^6*b^4 + 6
*A*a^5*b^5)*d^5*e^6 - 360*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 - 45*(3*B*a^
8*b^2 + 8*A*a^7*b^3)*d^3*e^8 - 10*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 - 3*(B
*a^10 + 10*A*a^9*b)*d*e^10 - 2*(11*B*b^10*d*e^10 - 6*(10*B*a*b^9 + A*b^10)
*e^11)*x^10 + 5*(11*B*b^10*d^2*e^9 - 6*(10*B*a*b^9 + A*b^10)*d*e^10 + 15*(
9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 - 15*(11*B*b^10*d^3*e^8 - 6*(10*B*a*b^9
+ A*b^10)*d^2*e^9 + 15*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 - 20*(8*B*a^3*b^7
+ 3*A*a^2*b^8)*e^11)*x^8 + 60*(11*B*b^10*d^4*e^7 - 6*(10*B*a*b^9 + A*b^10)
)*d^3*e^8 + 15*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 - 20*(8*B*a^3*b^7 + 3*A*a
^2*b^8)*d*e^10 + 15*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7 - 420*(11*B*b^10
*d^5*e^6 - 6*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 15*(9*B*a^2*b^8 + 2*A*a*b^9)*
d^3*e^8 - 20*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 15*(7*B*a^4*b^6 + 4*A*a
^3*b^7)*d*e^10 - 6*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 - (47497*B*b^10*d
^6*e^5 - 24762*(10*B*a*b^9 + A*b^10)*d^5*e^6 + 58125*(9*B*a^2*b^8 + 2*A*a*
b^9)*d^4*e^7 - 70500*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 + 45000*(7*B*a^4*
b^6 + 4*A*a^3*b^7)*d^2*e^9 - 12600*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10)*x^5
- 5*(19777*B*b^10*d^7*e^4 - 9642*(10*B*a*b^9 + A*b^10)*d^6*e^5 + 20325...

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^6} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**10*(B*x+A)/(e*x+d)**6,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1861 vs. $2(433) = 866$.

Time = 0.14 (sec) , antiderivative size = 1861, normalized size of antiderivative = 4.16

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^6} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^6,x, algorithm="maxima")`

output

```
1/60*(15797*B*b^10*d^11 - 12*A*a^10*e^11 - 9762*(10*B*a*b^9 + A*b^10)*d^10
*e + 28185*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 - 44580*(8*B*a^3*b^7 + 3*A*a^
2*b^8)*d^8*e^3 + 41310*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e^4 - 21924*(6*B*a^
5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 5754*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^5*e^6 -
360*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 - 45*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d
^3*e^8 - 10*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 - 3*(B*a^10 + 10*A*a^9*b)*d*
e^10 + 1800*(11*B*b^10*d^7*e^4 - 7*(10*B*a*b^9 + A*b^10)*d^6*e^5 + 21*(9*B
*a^2*b^8 + 2*A*a*b^9)*d^5*e^6 - 35*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^4*e^7 + 3
5*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^3*e^8 - 21*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2
*e^9 + 7*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d*e^10 - (4*B*a^7*b^3 + 7*A*a^6*b^4)*
e^11)*x^4 + 450*(165*B*b^10*d^8*e^3 - 104*(10*B*a*b^9 + A*b^10)*d^7*e^4 +
308*(9*B*a^2*b^8 + 2*A*a*b^9)*d^6*e^5 - 504*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^
5*e^6 + 490*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^4*e^7 - 280*(6*B*a^5*b^5 + 5*A*a
^4*b^6)*d^3*e^8 + 84*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^2*e^9 - 8*(4*B*a^7*b^3
+ 7*A*a^6*b^4)*d*e^10 - (3*B*a^8*b^2 + 8*A*a^7*b^3)*e^11)*x^3 + 50*(2101*B
*b^10*d^9*e^2 - 1314*(10*B*a*b^9 + A*b^10)*d^8*e^3 + 3852*(9*B*a^2*b^8 + 2
*A*a*b^9)*d^7*e^4 - 6216*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^6*e^5 + 5922*(7*B*a
^4*b^6 + 4*A*a^3*b^7)*d^5*e^6 - 3276*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^4*e^7 +
924*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^3*e^8 - 72*(4*B*a^7*b^3 + 7*A*a^6*b^4)*
d^2*e^9 - 9*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d*e^10 - 2*(2*B*a^9*b + 9*A*a^8...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2032 vs. $2(433) = 866$.

Time = 0.14 (sec) , antiderivative size = 2032, normalized size of antiderivative = 4.55

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^6} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^6,x, algorithm="giac")`

output

```
42*(11*B*b^10*d^6 - 60*B*a*b^9*d^5*e - 6*A*b^10*d^5*e + 135*B*a^2*b^8*d^4*
e^2 + 30*A*a*b^9*d^4*e^2 - 160*B*a^3*b^7*d^3*e^3 - 60*A*a^2*b^8*d^3*e^3 +
105*B*a^4*b^6*d^2*e^4 + 60*A*a^3*b^7*d^2*e^4 - 36*B*a^5*b^5*d*e^5 - 30*A*a
^4*b^6*d*e^5 + 5*B*a^6*b^4*e^6 + 6*A*a^5*b^5*e^6)*log(abs(e*x + d))/e^12 +
1/60*(15797*B*b^10*d^11 - 97620*B*a*b^9*d^10*e - 9762*A*b^10*d^10*e + 253
665*B*a^2*b^8*d^9*e^2 + 56370*A*a*b^9*d^9*e^2 - 356640*B*a^3*b^7*d^8*e^3 -
133740*A*a^2*b^8*d^8*e^3 + 289170*B*a^4*b^6*d^7*e^4 + 165240*A*a^3*b^7*d^
7*e^4 - 131544*B*a^5*b^5*d^6*e^5 - 109620*A*a^4*b^6*d^6*e^5 + 28770*B*a^6*
b^4*d^5*e^6 + 34524*A*a^5*b^5*d^5*e^6 - 1440*B*a^7*b^3*d^4*e^7 - 2520*A*a^
6*b^4*d^4*e^7 - 135*B*a^8*b^2*d^3*e^8 - 360*A*a^7*b^3*d^3*e^8 - 20*B*a^9*b
*d^2*e^9 - 90*A*a^8*b^2*d^2*e^9 - 3*B*a^10*d*e^10 - 30*A*a^9*b*d*e^10 - 12
*A*a^10*e^11 + 1800*(11*B*b^10*d^7*e^4 - 70*B*a*b^9*d^6*e^5 - 7*A*b^10*d^6
*e^5 + 189*B*a^2*b^8*d^5*e^6 + 42*A*a*b^9*d^5*e^6 - 280*B*a^3*b^7*d^4*e^7
- 105*A*a^2*b^8*d^4*e^7 + 245*B*a^4*b^6*d^3*e^8 + 140*A*a^3*b^7*d^3*e^8 -
126*B*a^5*b^5*d^2*e^9 - 105*A*a^4*b^6*d^2*e^9 + 35*B*a^6*b^4*d*e^10 + 42*A
*a^5*b^5*d*e^10 - 4*B*a^7*b^3*e^11 - 7*A*a^6*b^4*e^11)*x^4 + 450*(165*B*b^
10*d^8*e^3 - 1040*B*a*b^9*d^7*e^4 - 104*A*b^10*d^7*e^4 + 2772*B*a^2*b^8*d^
6*e^5 + 616*A*a*b^9*d^6*e^5 - 4032*B*a^3*b^7*d^5*e^6 - 1512*A*a^2*b^8*d^5*
e^6 + 3430*B*a^4*b^6*d^4*e^7 + 1960*A*a^3*b^7*d^4*e^7 - 1680*B*a^5*b^5*d^3
*e^8 - 1400*A*a^4*b^6*d^3*e^8 + 420*B*a^6*b^4*d^2*e^9 + 504*A*a^5*b^5*d...
```

Mupad [B] (verification not implemented)

Time = 1.21 (sec) , antiderivative size = 2681, normalized size of antiderivative = 6.00

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^6} dx = \text{Too large to display}$$

input `int(((A + B*x)*(a + b*x)^10)/(d + e*x)^6,x)`

output

```
x^5*((A*b^10 + 10*B*a*b^9)/(5*e^6) - (6*B*b^10*d)/(5*e^7)) - x^4*((3*d*((A
*b^10 + 10*B*a*b^9)/e^6 - (6*B*b^10*d)/e^7))/(2*e) - (5*a*b^8*(2*A*b + 9*B
*a))/(4*e^6) + (15*B*b^10*d^2)/(4*e^8)) - (x^4*(210*A*a^6*b^4*e^10 + 120*B
*a^7*b^3*e^10 + 210*A*b^10*d^6*e^4 - 330*B*b^10*d^7*e^3 - 1260*A*a*b^9*d^5
*e^5 - 1260*A*a^5*b^5*d*e^9 + 2100*B*a*b^9*d^6*e^4 - 1050*B*a^6*b^4*d*e^9
+ 3150*A*a^2*b^8*d^4*e^6 - 4200*A*a^3*b^7*d^3*e^7 + 3150*A*a^4*b^6*d^2*e^8
- 5670*B*a^2*b^8*d^5*e^5 + 8400*B*a^3*b^7*d^4*e^6 - 7350*B*a^4*b^6*d^3*e^
7 + 3780*B*a^5*b^5*d^2*e^8) + x^3*(60*A*a^7*b^3*e^10 + (45*B*a^8*b^2*e^10)
/2 + 780*A*b^10*d^7*e^3 - (2475*B*b^10*d^8*e^2)/2 - 4620*A*a*b^9*d^6*e^4 +
420*A*a^6*b^4*d*e^9 + 7800*B*a*b^9*d^7*e^3 + 240*B*a^7*b^3*d*e^9 + 11340*
A*a^2*b^8*d^5*e^5 - 14700*A*a^3*b^7*d^4*e^6 + 10500*A*a^4*b^6*d^3*e^7 - 37
80*A*a^5*b^5*d^2*e^8 - 20790*B*a^2*b^8*d^6*e^4 + 30240*B*a^3*b^7*d^5*e^5 -
25725*B*a^4*b^6*d^4*e^6 + 12600*B*a^5*b^5*d^3*e^7 - 3150*B*a^6*b^4*d^2*e^
8) + (12*A*a^10*e^11 - 15797*B*b^10*d^11 + 9762*A*b^10*d^10*e + 3*B*a^10*d
*e^10 - 56370*A*a*b^9*d^9*e^2 + 20*B*a^9*b*d^2*e^9 + 133740*A*a^2*b^8*d^8*
e^3 - 165240*A*a^3*b^7*d^7*e^4 + 109620*A*a^4*b^6*d^6*e^5 - 34524*A*a^5*b^
5*d^5*e^6 + 2520*A*a^6*b^4*d^4*e^7 + 360*A*a^7*b^3*d^3*e^8 + 90*A*a^8*b^2*
d^2*e^9 - 253665*B*a^2*b^8*d^9*e^2 + 356640*B*a^3*b^7*d^8*e^3 - 289170*B*a
^4*b^6*d^7*e^4 + 131544*B*a^5*b^5*d^6*e^5 - 28770*B*a^6*b^4*d^5*e^6 + 1440
*B*a^7*b^3*d^4*e^7 + 135*B*a^8*b^2*d^3*e^8 + 30*A*a^9*b*d*e^10 + 97620*...
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 1946, normalized size of antiderivative = 4.35

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^6} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^6,x)`

output

```
(27720*log(d + e*x)*a**6*b**5*d**6*e**6 + 138600*log(d + e*x)*a**6*b**5*d*
*5*e**7*x + 277200*log(d + e*x)*a**6*b**5*d**4*e**8*x**2 + 277200*log(d +
e*x)*a**6*b**5*d**3*e**9*x**3 + 138600*log(d + e*x)*a**6*b**5*d**2*e**10*x
**4 + 27720*log(d + e*x)*a**6*b**5*d*e**11*x**5 - 166320*log(d + e*x)*a**5
*b**6*d**7*e**5 - 831600*log(d + e*x)*a**5*b**6*d**6*e**6*x - 1663200*log(
d + e*x)*a**5*b**6*d**5*e**7*x**2 - 1663200*log(d + e*x)*a**5*b**6*d**4*e
**8*x**3 - 831600*log(d + e*x)*a**5*b**6*d**3*e**9*x**4 - 166320*log(d + e
x)*a**5*b**6*d**2*e**10*x**5 + 415800*log(d + e*x)*a**4*b**7*d**8*e**4 + 2
079000*log(d + e*x)*a**4*b**7*d**7*e**5*x + 4158000*log(d + e*x)*a**4*b**7
*d**6*e**6*x**2 + 4158000*log(d + e*x)*a**4*b**7*d**5*e**7*x**3 + 2079000*
log(d + e*x)*a**4*b**7*d**4*e**8*x**4 + 415800*log(d + e*x)*a**4*b**7*d**3
*e**9*x**5 - 554400*log(d + e*x)*a**3*b**8*d**9*e**3 - 2772000*log(d + e*x
)*a**3*b**8*d**8*e**4*x - 5544000*log(d + e*x)*a**3*b**8*d**7*e**5*x**2 -
5544000*log(d + e*x)*a**3*b**8*d**6*e**6*x**3 - 2772000*log(d + e*x)*a**3*
b**8*d**5*e**7*x**4 - 554400*log(d + e*x)*a**3*b**8*d**4*e**8*x**5 + 41580
0*log(d + e*x)*a**2*b**9*d**10*e**2 + 2079000*log(d + e*x)*a**2*b**9*d**9*
e**3*x + 4158000*log(d + e*x)*a**2*b**9*d**8*e**4*x**2 + 4158000*log(d + e
*x)*a**2*b**9*d**7*e**5*x**3 + 2079000*log(d + e*x)*a**2*b**9*d**6*e**6*x*
*4 + 415800*log(d + e*x)*a**2*b**9*d**5*e**7*x**5 - 166320*log(d + e*x)*a*
b**10*d**11*e - 831600*log(d + e*x)*a*b**10*d**10*e**2*x - 1663200*log(...
```

3.85 $\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^7} dx$

Optimal result	873
Mathematica [A] (verified)	874
Rubi [A] (verified)	875
Maple [B] (verified)	876
Fricas [B] (verification not implemented)	877
Sympy [F(-1)]	878
Maxima [B] (verification not implemented)	879
Giac [B] (verification not implemented)	880
Mupad [B] (verification not implemented)	881
Reduce [B] (verification not implemented)	881

Optimal result

Integrand size = 20, antiderivative size = 447

$$\begin{aligned}
 \int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^7} dx = & \frac{30b^6(bd-ae)^3(11bBd-4Abe-7aBe)x}{e^{11}} \\
 & + \frac{(bd-ae)^{10}(Bd-Ae)}{6e^{12}(d+ex)^6} \\
 & - \frac{(bd-ae)^9(11bBd-10Abe-aBe)}{5e^{12}(d+ex)^5} \\
 & + \frac{5b(bd-ae)^8(11bBd-9Abe-2aBe)}{4e^{12}(d+ex)^4} \\
 & - \frac{5b^2(bd-ae)^7(11bBd-8Abe-3aBe)}{e^{12}(d+ex)^3} \\
 & + \frac{15b^3(bd-ae)^6(11bBd-7Abe-4aBe)}{e^{12}(d+ex)^2} \\
 & - \frac{42b^4(bd-ae)^5(11bBd-6Abe-5aBe)}{e^{12}(d+ex)} \\
 & - \frac{15b^7(bd-ae)^2(11bBd-3Abe-8aBe)(d+ex)^2}{2e^{12}} \\
 & + \frac{5b^8(bd-ae)(11bBd-2Abe-9aBe)(d+ex)^3}{3e^{12}} \\
 & - \frac{b^9(11bBd-Abe-10aBe)(d+ex)^4}{4e^{12}} + \frac{b^{10}B(d+ex)^5}{5e^{12}} \\
 & - \frac{42b^5(bd-ae)^4(11bBd-5Abe-6aBe)\log(d+ex)}{e^{12}}
 \end{aligned}$$

output

```

30*b^6*(-a*e+b*d)^3*(-4*A*b*e-7*B*a*e+11*B*b*d)*x/e^11+1/6*(-a*e+b*d)^10*(
-A*e+B*d)/e^12/(e*x+d)^6-1/5*(-a*e+b*d)^9*(-10*A*b*e-B*a*e+11*B*b*d)/e^12/
(e*x+d)^5+5/4*b*(-a*e+b*d)^8*(-9*A*b*e-2*B*a*e+11*B*b*d)/e^12/(e*x+d)^4-5*
b^2*(-a*e+b*d)^7*(-8*A*b*e-3*B*a*e+11*B*b*d)/e^12/(e*x+d)^3+15*b^3*(-a*e+b
*d)^6*(-7*A*b*e-4*B*a*e+11*B*b*d)/e^12/(e*x+d)^2-42*b^4*(-a*e+b*d)^5*(-6*A
*b*e-5*B*a*e+11*B*b*d)/e^12/(e*x+d)-15/2*b^7*(-a*e+b*d)^2*(-3*A*b*e-8*B*a
e+11*B*b*d)*(e*x+d)^2/e^12+5/3*b^8*(-a*e+b*d)*(-2*A*b*e-9*B*a*e+11*B*b*d)*
(e*x+d)^3/e^12-1/4*b^9*(-A*b*e-10*B*a*e+11*B*b*d)*(e*x+d)^4/e^12+1/5*b^10*
B*(e*x+d)^5/e^12-42*b^5*(-a*e+b*d)^4*(-5*A*b*e-6*B*a*e+11*B*b*d)*ln(e*x+d)
/e^12

```

Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 505, normalized size of antiderivative = 1.13

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^7} dx$$

$$= \frac{-60b^6e(-210a^4Be^4 - 42b^4d^3(5Bd - 2Ae) + 280ab^3d^2e(3Bd - Ae) - 315a^2b^2de^2(4Bd - Ae) - 120a^3be^3(3Bd - Ae) + 30b^7e^2(120a^3B^3e^3 + 70a^2b^2d^2e(4Bd - Ae) + 45a^2b^2e^2(-7Bd + Ae) + 28b^3d^2(-3Bd + Ae))x^2 - 20b^8e^3(-45a^2B^2e^2 - 10a^2b^2e(-7Bd + Ae) + 7b^2d^2(-4Bd + Ae))x^3 + 15b^9e^4(-7b^2Bd + A^2b^2e + 10a^2B^2e)x^4 + 12b^10B^5e^5x^5 + (10(b^2d - a^2e)^10(Bd - Ae))/(d + ex)^6 - (12(b^2d - a^2e)^9(11b^2Bd - 10A^2b^2e - a^2B^2e))/(d + ex)^5 + (75b^2(b^2d - a^2e)^8(11b^2Bd - 9A^2b^2e - 2a^2B^2e))/(d + ex)^4 - (300b^2(b^2d - a^2e)^7(11b^2Bd - 8A^2b^2e - 3a^2B^2e))/(d + ex)^3 + (900b^3(b^2d - a^2e)^6(11b^2Bd - 7A^2b^2e - 4a^2B^2e))/(d + ex)^2 - (2520b^4(b^2d - a^2e)^5(11b^2Bd - 6A^2b^2e - 5a^2B^2e))/(d + ex) - 2520b^5(b^2d - a^2e)^4(11b^2Bd - 5A^2b^2e - 6a^2B^2e)*Log[d + ex]}{60e^{12}}$$

input

```
Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^7,x]
```

output

```

(-60*b^6*e*(-210*a^4*B*e^4 - 42*b^4*d^3*(5*B*d - 2*A*e) + 280*a*b^3*d^2*e*
(3*B*d - A*e) - 315*a^2*b^2*d^2*e^2*(4*B*d - A*e) - 120*a^3*b^2*e^3*(-7*B*d +
A*e))*x + 30*b^7*e^2*(120*a^3*B^3e^3 + 70*a^2*b^2*d^2*e*(4*B*d - A*e) + 45*a^2*
b^2*e^2*(-7*B*d + A*e) + 28*b^3*d^2*(-3*B*d + A*e))*x^2 - 20*b^8*e^3*(-45*a^
2*B^2e^2 - 10*a^2*b^2*e*(-7*B*d + A*e) + 7*b^2*d^2*(-4*B*d + A*e))*x^3 + 15*b^9*
e^4*(-7*b^2*B*d + A^2b^2e + 10*a^2*B^2e)*x^4 + 12*b^10*B^5e^5*x^5 + (10*(b*d - a*e)
^10*(B*d - A*e))/(d + e*x)^6 - (12*(b*d - a*e)^9*(11*b*B*d - 10*A*b^2e - a^
2*B^2e))/(d + e*x)^5 + (75*b*(b*d - a*e)^8*(11*b*B*d - 9*A*b^2e - 2*a*B^2e))/(d
+ e*x)^4 - (300*b^2*(b*d - a*e)^7*(11*b*B*d - 8*A*b^2e - 3*a*B^2e))/(d + e*
x)^3 + (900*b^3*(b*d - a*e)^6*(11*b*B*d - 7*A*b^2e - 4*a*B^2e))/(d + e*x)^2
- (2520*b^4*(b*d - a*e)^5*(11*b*B*d - 6*A*b^2e - 5*a*B^2e))/(d + e*x) - 2520
*b^5*(b*d - a*e)^4*(11*b*B*d - 5*A*b^2e - 6*a*B^2e)*Log[d + e*x]/(60*e^12)

```

Rubi [A] (verified)

Time = 1.35 (sec) , antiderivative size = 447, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^7} dx$$

↓ 86

$$\int \left(\frac{b^9(d + ex)^3(10aBe + Abe - 11bBd)}{e^{11}} - \frac{5b^8(d + ex)^2(bd - ae)(9aBe + 2Abe - 11bBd)}{e^{11}} + \frac{15b^7(d + ex)(bd - ae)^2(10aBe + Abe - 11bBd)}{e^{11}} - \frac{5b^6(d + ex)(bd - ae)^3(9aBe + 2Abe - 11bBd)}{e^{11}} + \frac{15b^5(d + ex)(bd - ae)^4(10aBe + Abe - 11bBd)}{e^{11}} - \frac{5b^4(d + ex)(bd - ae)^5(9aBe + 2Abe - 11bBd)}{e^{11}} + \frac{15b^3(d + ex)(bd - ae)^6(10aBe + Abe - 11bBd)}{e^{11}} - \frac{5b^2(d + ex)(bd - ae)^7(9aBe + 2Abe - 11bBd)}{e^{11}} + \frac{15b(d + ex)(bd - ae)^8(10aBe + Abe - 11bBd)}{e^{11}} - \frac{5b^9(d + ex)(bd - ae)^9(10aBe + Abe - 11bBd)}{e^{11}} + \frac{b^{10}(A + Bx)}{e^{11}} \right) dx$$

↓ 2009

$$\begin{aligned} & - \frac{b^9(d + ex)^4(-10aBe - Abe + 11bBd)}{4e^{12}} + \frac{5b^8(d + ex)^3(bd - ae)(-9aBe - 2Abe + 11bBd)}{3e^{12}} - \\ & - \frac{15b^7(d + ex)^2(bd - ae)^2(-8aBe - 3Abe + 11bBd)}{2e^{12}} + \frac{30b^6x(bd - ae)^3(-7aBe - 4Abe + 11bBd)}{e^{11}} - \\ & - \frac{42b^5(bd - ae)^4 \log(d + ex)(-6aBe - 5Abe + 11bBd)}{e^{12}} - \\ & - \frac{42b^4(bd - ae)^5(-5aBe - 6Abe + 11bBd)}{e^{12}(d + ex)} + \frac{15b^3(bd - ae)^6(-4aBe - 7Abe + 11bBd)}{e^{12}(d + ex)^2} - \\ & - \frac{5b^2(bd - ae)^7(-3aBe - 8Abe + 11bBd)}{e^{12}(d + ex)^3} + \frac{5b(bd - ae)^8(-2aBe - 9Abe + 11bBd)}{4e^{12}(d + ex)^4} - \\ & - \frac{(bd - ae)^9(-aBe - 10Abe + 11bBd)}{5e^{12}(d + ex)^5} + \frac{(bd - ae)^{10}(Bd - Ae)}{6e^{12}(d + ex)^6} + \frac{b^{10}B(d + ex)^5}{5e^{12}} \end{aligned}$$

input

```
Int[((a + b*x)^10*(A + B*x))/(d + e*x)^7,x]
```


output

```
(30*b^6*(b*d - a*e)^3*(11*b*B*d - 4*A*b*e - 7*a*B*e)*x)/e^11 + ((b*d - a*e)^10*(B*d - A*e))/(6*e^12*(d + e*x)^6) - ((b*d - a*e)^9*(11*b*B*d - 10*A*b*e - a*B*e))/(5*e^12*(d + e*x)^5) + (5*b*(b*d - a*e)^8*(11*b*B*d - 9*A*b*e - 2*a*B*e))/(4*e^12*(d + e*x)^4) - (5*b^2*(b*d - a*e)^7*(11*b*B*d - 8*A*b*e - 3*a*B*e))/(e^12*(d + e*x)^3) + (15*b^3*(b*d - a*e)^6*(11*b*B*d - 7*A*b*e - 4*a*B*e))/(e^12*(d + e*x)^2) - (42*b^4*(b*d - a*e)^5*(11*b*B*d - 6*A*b*e - 5*a*B*e))/(e^12*(d + e*x)) - (15*b^7*(b*d - a*e)^2*(11*b*B*d - 3*A*b*e - 8*a*B*e)*(d + e*x)^2)/(2*e^12) + (5*b^8*(b*d - a*e)*(11*b*B*d - 2*A*b*e - 9*a*B*e)*(d + e*x)^3)/(3*e^12) - (b^9*(11*b*B*d - A*b*e - 10*a*B*e)*(d + e*x)^4)/(4*e^12) + (b^10*B*(d + e*x)^5)/(5*e^12) - (42*b^5*(b*d - a*e)^4*(11*b*B*d - 5*A*b*e - 6*a*B*e)*Log[d + e*x])/e^12
```

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1920 vs. $2(433) = 866$.

Time = 0.24 (sec) , antiderivative size = 1921, normalized size of antiderivative = 4.30

method	result	size
norman	Expression too large to display	1921
default	Expression too large to display	1974
risch	Expression too large to display	2036
parallelsch	Expression too large to display	3755

input

```
int((b*x+a)^10*(B*x+A)/(e*x+d)^7,x,method=_RETURNVERBOSE)
```

output

```
(-1/60*(10*A*a^10*e^11+20*A*a^9*b*d*e^10+45*A*a^8*b^2*d^2*e^9+120*A*a^7*b^3*d^3*e^8+420*A*a^6*b^4*d^4*e^7+2520*A*a^5*b^5*d^5*e^6-30870*A*a^4*b^6*d^6*e^5+123480*A*a^3*b^7*d^7*e^4-185220*A*a^2*b^8*d^8*e^3+123480*A*a*b^9*d^9*e^2-30870*A*b^10*d^10*e+2*B*a^10*d*e^10+10*B*a^9*b*d^2*e^9+45*B*a^8*b^2*d^3*e^8+240*B*a^7*b^3*d^4*e^7+2100*B*a^6*b^4*d^5*e^6-37044*B*a^5*b^5*d^6*e^5+216090*B*a^4*b^6*d^7*e^4-493920*B*a^3*b^7*d^8*e^3+555660*B*a^2*b^8*d^9*e^2-308700*B*a*b^9*d^10*e+67914*B*b^10*d^11)/e^12-6*(42*A*a^5*b^5*e^6-210*A*a^4*b^6*d*e^5+840*A*a^3*b^7*d^2*e^4-1260*A*a^2*b^8*d^3*e^3+840*A*a*b^9*d^4*e^2-210*A*b^10*d^5*e+35*B*a^6*b^4*e^6-252*B*a^5*b^5*d*e^5+1470*B*a^4*b^6*d^2*e^4-3360*B*a^3*b^7*d^3*e^3+3780*B*a^2*b^8*d^4*e^2-2100*B*a*b^9*d^5*e+462*B*b^10*d^6)/e^7*x^5-15*(7*A*a^6*b^4*e^7+42*A*a^5*b^5*d*e^6-315*A*a^4*b^6*d^2*e^5+1260*A*a^3*b^7*d^3*e^4-1890*A*a^2*b^8*d^4*e^3+1260*A*a*b^9*d^5*e^2-315*A*b^10*d^6*e+4*B*a^7*b^3*e^7+35*B*a^6*b^4*d*e^6-378*B*a^5*b^5*d^2*e^5+2205*B*a^4*b^6*d^3*e^4-5040*B*a^3*b^7*d^4*e^3+5670*B*a^2*b^8*d^5*e^2-3150*B*a*b^9*d^6*e+693*B*b^10*d^7)/e^8*x^4-5*(8*A*a^7*b^3*e^8+28*A*a^6*b^4*d*e^7+168*A*a^5*b^5*d^2*e^6-1540*A*a^4*b^6*d^3*e^5+6160*A*a^3*b^7*d^4*e^4-9240*A*a^2*b^8*d^5*e^3+6160*A*a*b^9*d^6*e^2-1540*A*b^10*d^7*e+3*B*a^8*b^2*e^8+16*B*a^7*b^3*d*e^7+140*B*a^6*b^4*d^2*e^6-1848*B*a^5*b^5*d^3*e^5+10780*B*a^4*b^6*d^4*e^4-24640*B*a^3*b^7*d^5*e^3+27720*B*a^2*b^8*d^6*e^2-15400*B*a*b^9*d^7*e+3388*B*b^10*d^8)/e^9*x^3-5/4*(9*A*a^8*b^2*e^9+24*A*a^7*b^3*d...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2850 vs. $2(433) = 866$.

Time = 0.17 (sec) , antiderivative size = 2850, normalized size of antiderivative = 6.38

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^7} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)/(e*x+d)^7,x, algorithm="fricas")
```

output

```

1/60*(12*B*b^10*e^11*x^11 - 20417*B*b^10*d^11 - 10*A*a^10*e^11 + 10655*(10
*B*a*b^9 + A*b^10)*d^10*e - 25090*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 + 3069
0*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 - 20070*(7*B*a^4*b^6 + 4*A*a^3*b^7)*
d^7*e^4 + 6174*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 - 420*(5*B*a^6*b^4 + 6*
A*a^5*b^5)*d^5*e^6 - 60*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 - 15*(3*B*a^8*
b^2 + 8*A*a^7*b^3)*d^3*e^8 - 5*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 - 2*(B*a^
10 + 10*A*a^9*b)*d*e^10 - 3*(11*B*b^10*d*e^10 - 5*(10*B*a*b^9 + A*b^10)*e^
11)*x^10 + 10*(11*B*b^10*d^2*e^9 - 5*(10*B*a*b^9 + A*b^10)*d*e^10 + 10*(9*
B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 - 45*(11*B*b^10*d^3*e^8 - 5*(10*B*a*b^9 +
A*b^10)*d^2*e^9 + 10*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 - 10*(8*B*a^3*b^7 +
3*A*a^2*b^8)*e^11)*x^8 + 360*(11*B*b^10*d^4*e^7 - 5*(10*B*a*b^9 + A*b^10)
*d^3*e^8 + 10*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 - 10*(8*B*a^3*b^7 + 3*A*a^
2*b^8)*d*e^10 + 5*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7 + (47497*B*b^10*d^
5*e^6 - 20215*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 36650*(9*B*a^2*b^8 + 2*A*a*b
^9)*d^3*e^8 - 31050*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 10800*(7*B*a^4*b
^6 + 4*A*a^3*b^7)*d*e^10)*x^6 + 6*(19777*B*b^10*d^6*e^5 - 7615*(10*B*a*b^9
+ A*b^10)*d^5*e^6 + 11450*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 - 5850*(8*B*a
^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 - 1800*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 +
2520*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10 - 420*(5*B*a^6*b^4 + 6*A*a^5*b^5)
*e^11)*x^5 + 15*(5917*B*b^10*d^7*e^4 - 1315*(10*B*a*b^9 + A*b^10)*d^6*e...

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^7} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**10*(B*x+A)/(e*x+d)**7,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1869 vs. $2(433) = 866$.

Time = 0.16 (sec) , antiderivative size = 1869, normalized size of antiderivative = 4.18

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^7} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^7,x, algorithm="maxima")`

output

```
-1/60*(20417*B*b^10*d^11 + 10*A*a^10*e^11 - 10655*(10*B*a*b^9 + A*b^10)*d^
10*e + 25090*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 - 30690*(8*B*a^3*b^7 + 3*A*
a^2*b^8)*d^8*e^3 + 20070*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e^4 - 6174*(6*B*a
^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 420*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^5*e^6 +
60*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 15*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^
3*e^8 + 5*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 + 2*(B*a^10 + 10*A*a^9*b)*d*e^
10 + 2520*(11*B*b^10*d^6*e^5 - 6*(10*B*a*b^9 + A*b^10)*d^5*e^6 + 15*(9*B*a
^2*b^8 + 2*A*a*b^9)*d^4*e^7 - 20*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 + 15*
(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 - 6*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10
+ (5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 + 900*(143*B*b^10*d^7*e^4 - 77*(1
0*B*a*b^9 + A*b^10)*d^6*e^5 + 189*(9*B*a^2*b^8 + 2*A*a*b^9)*d^5*e^6 - 245*
(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^4*e^7 + 175*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^3*
e^8 - 63*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e^9 + 7*(5*B*a^6*b^4 + 6*A*a^5*b^
5)*d*e^10 + (4*B*a^7*b^3 + 7*A*a^6*b^4)*e^11)*x^4 + 300*(803*B*b^10*d^8*e^
3 - 428*(10*B*a*b^9 + A*b^10)*d^7*e^4 + 1036*(9*B*a^2*b^8 + 2*A*a*b^9)*d^6
*e^5 - 1316*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^5*e^6 + 910*(7*B*a^4*b^6 + 4*A*a
^3*b^7)*d^4*e^7 - 308*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^3*e^8 + 28*(5*B*a^6*b^
4 + 6*A*a^5*b^5)*d^2*e^9 + 4*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d*e^10 + (3*B*a^8
*b^2 + 8*A*a^7*b^3)*e^11)*x^3 + 75*(3025*B*b^10*d^9*e^2 - 1599*(10*B*a*b^9
+ A*b^10)*d^8*e^3 + 3828*(9*B*a^2*b^8 + 2*A*a*b^9)*d^7*e^4 - 4788*(8*B...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2009 vs. $2(433) = 866$.

Time = 0.13 (sec) , antiderivative size = 2009, normalized size of antiderivative = 4.49

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^7} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^7,x, algorithm="giac")`

output

```
-42*(11*B*b^10*d^5 - 50*B*a*b^9*d^4*e - 5*A*b^10*d^4*e + 90*B*a^2*b^8*d^3*
e^2 + 20*A*a*b^9*d^3*e^2 - 80*B*a^3*b^7*d^2*e^3 - 30*A*a^2*b^8*d^2*e^3 + 3
5*B*a^4*b^6*d*e^4 + 20*A*a^3*b^7*d*e^4 - 6*B*a^5*b^5*e^5 - 5*A*a^4*b^6*e^5
)*log(abs(e*x + d))/e^12 - 1/60*(20417*B*b^10*d^11 - 106550*B*a*b^9*d^10*e
- 10655*A*b^10*d^10*e + 225810*B*a^2*b^8*d^9*e^2 + 50180*A*a*b^9*d^9*e^2
- 245520*B*a^3*b^7*d^8*e^3 - 92070*A*a^2*b^8*d^8*e^3 + 140490*B*a^4*b^6*d^
7*e^4 + 80280*A*a^3*b^7*d^7*e^4 - 37044*B*a^5*b^5*d^6*e^5 - 30870*A*a^4*b^
6*d^6*e^5 + 2100*B*a^6*b^4*d^5*e^6 + 2520*A*a^5*b^5*d^5*e^6 + 240*B*a^7*b^
3*d^4*e^7 + 420*A*a^6*b^4*d^4*e^7 + 45*B*a^8*b^2*d^3*e^8 + 120*A*a^7*b^3*d
^3*e^8 + 10*B*a^9*b*d^2*e^9 + 45*A*a^8*b^2*d^2*e^9 + 2*B*a^10*d*e^10 + 20*
A*a^9*b*d*e^10 + 10*A*a^10*e^11 + 2520*(11*B*b^10*d^6*e^5 - 60*B*a*b^9*d^5
*e^6 - 6*A*b^10*d^5*e^6 + 135*B*a^2*b^8*d^4*e^7 + 30*A*a*b^9*d^4*e^7 - 160
*B*a^3*b^7*d^3*e^8 - 60*A*a^2*b^8*d^3*e^8 + 105*B*a^4*b^6*d^2*e^9 + 60*A*a
^3*b^7*d^2*e^9 - 36*B*a^5*b^5*d*e^10 - 30*A*a^4*b^6*d*e^10 + 5*B*a^6*b^4*e
^11 + 6*A*a^5*b^5*e^11)*x^5 + 900*(143*B*b^10*d^7*e^4 - 770*B*a*b^9*d^6*e^
5 - 77*A*b^10*d^6*e^5 + 1701*B*a^2*b^8*d^5*e^6 + 378*A*a*b^9*d^5*e^6 - 196
0*B*a^3*b^7*d^4*e^7 - 735*A*a^2*b^8*d^4*e^7 + 1225*B*a^4*b^6*d^3*e^8 + 700
*A*a^3*b^7*d^3*e^8 - 378*B*a^5*b^5*d^2*e^9 - 315*A*a^4*b^6*d^2*e^9 + 35*B*
a^6*b^4*d*e^10 + 42*A*a^5*b^5*d*e^10 + 4*B*a^7*b^3*e^11 + 7*A*a^6*b^4*e^11
)*x^4 + 300*(803*B*b^10*d^8*e^3 - 4280*B*a*b^9*d^7*e^4 - 428*A*b^10*d^7...
```

Mupad [B] (verification not implemented)

Time = 1.18 (sec) , antiderivative size = 2252, normalized size of antiderivative = 5.04

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^7} dx = \text{Too large to display}$$

input `int(((A + B*x)*(a + b*x)^10)/(d + e*x)^7,x)`

output

```
x*((7*d*((21*d^2*((A*b^10 + 10*B*a*b^9)/e^7 - (7*B*b^10*d)/e^8))/e^2 - (7*d*((7*d*((A*b^10 + 10*B*a*b^9)/e^7 - (7*B*b^10*d)/e^8))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^7 + (21*B*b^10*d^2)/e^9))/e - (15*a^2*b^7*(3*A*b + 8*B*a))/e^7 + (35*B*b^10*d^3)/e^10))/e - (35*d^3*((A*b^10 + 10*B*a*b^9)/e^7 - (7*B*b^10*d)/e^8))/e^3 + (21*d^2*((7*d*((A*b^10 + 10*B*a*b^9)/e^7 - (7*B*b^10*d)/e^8))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^7 + (21*B*b^10*d^2)/e^9))/e^2 + (30*a^3*b^6*(4*A*b + 7*B*a))/e^7 - (35*B*b^10*d^4)/e^11) - x^3*((7*d*((A*b^10 + 10*B*a*b^9)/e^7 - (7*B*b^10*d)/e^8))/(3*e) - (5*a*b^8*(2*A*b + 9*B*a))/(3*e^7) + (7*B*b^10*d^2)/e^9) - x^2*((21*d^2*((A*b^10 + 10*B*a*b^9)/e^7 - (7*B*b^10*d)/e^8))/(2*e^2) - (7*d*((7*d*((A*b^10 + 10*B*a*b^9)/e^7 - (7*B*b^10*d)/e^8))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^7 + (21*B*b^10*d^2)/e^9))/(2*e) - (15*a^2*b^7*(3*A*b + 8*B*a))/(2*e^7) + (35*B*b^10*d^3)/(2*e^10)) + x^4*((A*b^10 + 10*B*a*b^9)/(4*e^7) - (7*B*b^10*d)/(4*e^8)) - (x^4*(105*A*a^6*b^4*e^10 + 60*B*a^7*b^3*e^10 - 1155*A*b^10*d^6*e^4 + 2145*B*b^10*d^7*e^3 + 5670*A*a*b^9*d^5*e^5 + 630*A*a^5*b^5*d*e^9 - 11550*B*a*b^9*d^6*e^4 + 525*B*a^6*b^4*d*e^9 - 11025*A*a^2*b^8*d^4*e^6 + 10500*A*a^3*b^7*d^3*e^7 - 4725*A*a^4*b^6*d^2*e^8 + 25515*B*a^2*b^8*d^5*e^5 - 29400*B*a^3*b^7*d^4*e^6 + 18375*B*a^4*b^6*d^3*e^7 - 5670*B*a^5*b^5*d^2*e^8) + x^3*(40*A*a^7*b^3*e^10 + 15*B*a^8*b^2*e^10 - 2140*A*b^10*d^7*e^3 + 4015*B*b^10*d^8*e^2 + 10360*A*a*b^9*d^6*e^4 + 140*A*a^6*b^4*d*e^9 - 21400*B*a*b^9*d^7*e^3 + 80*B*a...
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 1957, normalized size of antiderivative = 4.38

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^7} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^7,x)`

output

```
(27720*log(d + e*x)*a**5*b**6*d**7*e**5 + 166320*log(d + e*x)*a**5*b**6*d*
*6*e**6*x + 415800*log(d + e*x)*a**5*b**6*d**5*e**7*x**2 + 554400*log(d +
e*x)*a**5*b**6*d**4*e**8*x**3 + 415800*log(d + e*x)*a**5*b**6*d**3*e**9*x*
*4 + 166320*log(d + e*x)*a**5*b**6*d**2*e**10*x**5 + 27720*log(d + e*x)*a*
*5*b**6*d*e**11*x**6 - 138600*log(d + e*x)*a**4*b**7*d**8*e**4 - 831600*lo
g(d + e*x)*a**4*b**7*d**7*e**5*x - 2079000*log(d + e*x)*a**4*b**7*d**6*e**
6*x**2 - 2772000*log(d + e*x)*a**4*b**7*d**5*e**7*x**3 - 2079000*log(d + e
*x)*a**4*b**7*d**4*e**8*x**4 - 831600*log(d + e*x)*a**4*b**7*d**3*e**9*x**
5 - 138600*log(d + e*x)*a**4*b**7*d**2*e**10*x**6 + 277200*log(d + e*x)*a*
*3*b**8*d**9*e**3 + 1663200*log(d + e*x)*a**3*b**8*d**8*e**4*x + 4158000*1
og(d + e*x)*a**3*b**8*d**7*e**5*x**2 + 5544000*log(d + e*x)*a**3*b**8*d**6
*e**6*x**3 + 4158000*log(d + e*x)*a**3*b**8*d**5*e**7*x**4 + 1663200*log(d
+ e*x)*a**3*b**8*d**4*e**8*x**5 + 277200*log(d + e*x)*a**3*b**8*d**3*e**9
*x**6 - 277200*log(d + e*x)*a**2*b**9*d**10*e**2 - 1663200*log(d + e*x)*a*
*2*b**9*d**9*e**3*x - 4158000*log(d + e*x)*a**2*b**9*d**8*e**4*x**2 - 5544
000*log(d + e*x)*a**2*b**9*d**7*e**5*x**3 - 4158000*log(d + e*x)*a**2*b**9
*d**6*e**6*x**4 - 1663200*log(d + e*x)*a**2*b**9*d**5*e**7*x**5 - 277200*1
og(d + e*x)*a**2*b**9*d**4*e**8*x**6 + 138600*log(d + e*x)*a*b**10*d**11*e
+ 831600*log(d + e*x)*a*b**10*d**10*e**2*x + 2079000*log(d + e*x)*a*b**10
*d**9*e**3*x**2 + 2772000*log(d + e*x)*a*b**10*d**8*e**4*x**3 + 2079000...
```

3.86 $\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^8} dx$

Optimal result	883
Mathematica [A] (verified)	884
Rubi [A] (verified)	885
Maple [B] (verified)	886
Fricas [B] (verification not implemented)	887
Sympy [F(-1)]	888
Maxima [B] (verification not implemented)	889
Giac [B] (verification not implemented)	890
Mupad [B] (verification not implemented)	891
Reduce [B] (verification not implemented)	891

Optimal result

Integrand size = 20, antiderivative size = 444

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^8} dx = -\frac{15b^7(bd-ae)^2(11bBd-3Abe-8aBe)x}{e^{11}} + \frac{(bd-ae)^{10}(Bd-Ae)}{7e^{12}(d+ex)^7} - \frac{(bd-ae)^9(11bBd-10Abe-aBe)}{6e^{12}(d+ex)^6} + \frac{b(bd-ae)^8(11bBd-9Abe-2aBe)}{e^{12}(d+ex)^5} - \frac{15b^2(bd-ae)^7(11bBd-8Abe-3aBe)}{4e^{12}(d+ex)^4} + \frac{10b^3(bd-ae)^6(11bBd-7Abe-4aBe)}{e^{12}(d+ex)^3} - \frac{21b^4(bd-ae)^5(11bBd-6Abe-5aBe)}{e^{12}(d+ex)^2} + \frac{42b^5(bd-ae)^4(11bBd-5Abe-6aBe)}{e^{12}(d+ex)} + \frac{5b^8(bd-ae)(11bBd-2Abe-9aBe)(d+ex)^2}{2e^{12}} - \frac{b^9(11bBd-Abe-10aBe)(d+ex)^3}{3e^{12}} + \frac{b^{10}B(d+ex)^4}{4e^{12}} + \frac{30b^6(bd-ae)^3(11bBd-4Abe-7aBe)\log(d+ex)}{e^{12}}$$

output

```
-15*b^7*(-a*e+b*d)^2*(-3*A*b*e-8*B*a*e+11*B*b*d)*x/e^11+1/7*(-a*e+b*d)^10*
(-A*e+B*d)/e^12/(e*x+d)^7-1/6*(-a*e+b*d)^9*(-10*A*b*e-B*a*e+11*B*b*d)/e^12
/(e*x+d)^6+b*(-a*e+b*d)^8*(-9*A*b*e-2*B*a*e+11*B*b*d)/e^12/(e*x+d)^5-15/4*
b^2*(-a*e+b*d)^7*(-8*A*b*e-3*B*a*e+11*B*b*d)/e^12/(e*x+d)^4+10*b^3*(-a*e+b
*d)^6*(-7*A*b*e-4*B*a*e+11*B*b*d)/e^12/(e*x+d)^3-21*b^4*(-a*e+b*d)^5*(-6*A
*b*e-5*B*a*e+11*B*b*d)/e^12/(e*x+d)^2+42*b^5*(-a*e+b*d)^4*(-5*A*b*e-6*B*a*
e+11*B*b*d)/e^12/(e*x+d)+5/2*b^8*(-a*e+b*d)*(-2*A*b*e-9*B*a*e+11*B*b*d)*(e
*x+d)^2/e^12-1/3*b^9*(-A*b*e-10*B*a*e+11*B*b*d)*(e*x+d)^3/e^12+1/4*b^10*B*
(e*x+d)^4/e^12+30*b^6*(-a*e+b*d)^3*(-4*A*b*e-7*B*a*e+11*B*b*d)*ln(e*x+d)/e
^12
```

Mathematica [A] (verified)

Time = 0.31 (sec) , antiderivative size = 450, normalized size of antiderivative = 1.01

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^8} dx$$

$$= \frac{84b^7e(120a^3Be^3 + 40ab^2de(9Bd - 2Ae) + 45a^2be^2(-8Bd + Ae) + 12b^3d^2(-10Bd + 3Ae))x - 42b^8e^2(-10a^2B^2e^2 - 10a*b*e*(-8B*d + A*e) + 4*b^2*d*(-9*B*d + 2*A*e))*x^2 + 28*b^9*e^3*(-8*b*B*d + A*b*e + 10*a*B*e)*x^3 + 21*b^10*B*e^4*x^4 + (12*(b*d - a*e)^10*(B*d - A*e))/(d + e*x)^7 - (14*(b*d - a*e)^9*(11*b*B*d - 10*A*b*e - a*B*e))/(d + e*x)^6 + (84*b*(b*d - a*e)^8*(11*b*B*d - 9*A*b*e - 2*a*B*e))/(d + e*x)^5 - (315*b^2*(b*d - a*e)^7*(11*b*B*d - 8*A*b*e - 3*a*B*e))/(d + e*x)^4 + (840*b^3*(b*d - a*e)^6*(11*b*B*d - 7*A*b*e - 4*a*B*e))/(d + e*x)^3 - (176*4*b^4*(b*d - a*e)^5*(11*b*B*d - 6*A*b*e - 5*a*B*e))/(d + e*x)^2 + (3528*b^5*(b*d - a*e)^4*(11*b*B*d - 5*A*b*e - 6*a*B*e))/(d + e*x) + 2520*b^6*(b*d - a*e)^3*(11*b*B*d - 4*A*b*e - 7*a*B*e)*Log[d + e*x]/(84*e^12)$$

input

```
Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^8,x]
```

output

```
(84*b^7*e*(120*a^3*B*e^3 + 40*a*b^2*d*e*(9*B*d - 2*A*e) + 45*a^2*b*e^2*(-8
*B*d + A*e) + 12*b^3*d^2*(-10*B*d + 3*A*e))*x - 42*b^8*e^2*(-45*a^2*B*e^2
- 10*a*b*e*(-8*B*d + A*e) + 4*b^2*d*(-9*B*d + 2*A*e))*x^2 + 28*b^9*e^3*(-8
*b*B*d + A*b*e + 10*a*B*e)*x^3 + 21*b^10*B*e^4*x^4 + (12*(b*d - a*e)^10*(B
*d - A*e))/(d + e*x)^7 - (14*(b*d - a*e)^9*(11*b*B*d - 10*A*b*e - a*B*e))/
(d + e*x)^6 + (84*b*(b*d - a*e)^8*(11*b*B*d - 9*A*b*e - 2*a*B*e))/(d + e*x
)^5 - (315*b^2*(b*d - a*e)^7*(11*b*B*d - 8*A*b*e - 3*a*B*e))/(d + e*x)^4 +
(840*b^3*(b*d - a*e)^6*(11*b*B*d - 7*A*b*e - 4*a*B*e))/(d + e*x)^3 - (176
4*b^4*(b*d - a*e)^5*(11*b*B*d - 6*A*b*e - 5*a*B*e))/(d + e*x)^2 + (3528*b^
5*(b*d - a*e)^4*(11*b*B*d - 5*A*b*e - 6*a*B*e))/(d + e*x) + 2520*b^6*(b*d
- a*e)^3*(11*b*B*d - 4*A*b*e - 7*a*B*e)*Log[d + e*x]/(84*e^12)
```

Rubi [A] (verified)

Time = 1.32 (sec) , antiderivative size = 444, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^8} dx$$

↓ 86

$$\int \left(\frac{b^9(d + ex)^2(10aBe + Abe - 11bBd)}{e^{11}} - \frac{5b^8(d + ex)(bd - ae)(9aBe + 2Abe - 11bBd)}{e^{11}} + \frac{15b^7(bd - ae)^2(8aBe - 11bBd)}{e^{11}} \right) dx$$

↓ 2009

$$\begin{aligned} & - \frac{b^9(d + ex)^3(-10aBe - Abe + 11bBd)}{3e^{12}} + \frac{5b^8(d + ex)^2(bd - ae)(-9aBe - 2Abe + 11bBd)}{2e^{12}} - \\ & \frac{15b^7x(bd - ae)^2(-8aBe - 3Abe + 11bBd)}{e^{11}} + \frac{30b^6(bd - ae)^3 \log(d + ex)(-7aBe - 4Abe + 11bBd)}{e^{12}} + \\ & \frac{42b^5(bd - ae)^4(-6aBe - 5Abe + 11bBd)}{e^{12}(d + ex)} - \frac{21b^4(bd - ae)^5(-5aBe - 6Abe + 11bBd)}{e^{12}(d + ex)^2} + \\ & \frac{10b^3(bd - ae)^6(-4aBe - 7Abe + 11bBd)}{e^{12}(d + ex)^3} - \frac{15b^2(bd - ae)^7(-3aBe - 8Abe + 11bBd)}{4e^{12}(d + ex)^4} + \\ & \frac{b(bd - ae)^8(-2aBe - 9Abe + 11bBd)}{e^{12}(d + ex)^5} - \frac{(bd - ae)^9(-aBe - 10Abe + 11bBd)}{6e^{12}(d + ex)^6} + \\ & \frac{(bd - ae)^{10}(Bd - Ae)}{7e^{12}(d + ex)^7} + \frac{b^{10}B(d + ex)^4}{4e^{12}} \end{aligned}$$

input

```
Int[((a + b*x)^10*(A + B*x))/(d + e*x)^8,x]
```

output

```
(-15*b^7*(b*d - a*e)^2*(11*b*B*d - 3*A*b*e - 8*a*B*e)*x)/e^11 + ((b*d - a*
e)^10*(B*d - A*e))/(7*e^12*(d + e*x)^7) - ((b*d - a*e)^9*(11*b*B*d - 10*A*
b*e - a*B*e))/(6*e^12*(d + e*x)^6) + (b*(b*d - a*e)^8*(11*b*B*d - 9*A*b*e
- 2*a*B*e))/(e^12*(d + e*x)^5) - (15*b^2*(b*d - a*e)^7*(11*b*B*d - 8*A*b*e
- 3*a*B*e))/(4*e^12*(d + e*x)^4) + (10*b^3*(b*d - a*e)^6*(11*b*B*d - 7*A*
b*e - 4*a*B*e))/(e^12*(d + e*x)^3) - (21*b^4*(b*d - a*e)^5*(11*b*B*d - 6*A
*b*e - 5*a*B*e))/(e^12*(d + e*x)^2) + (42*b^5*(b*d - a*e)^4*(11*b*B*d - 5*
A*b*e - 6*a*B*e))/(e^12*(d + e*x)) + (5*b^8*(b*d - a*e)*(11*b*B*d - 2*A*b*
e - 9*a*B*e)*(d + e*x)^2)/(2*e^12) - (b^9*(11*b*B*d - A*b*e - 10*a*B*e)*(d
+ e*x)^3)/(3*e^12) + (b^10*B*(d + e*x)^4)/(4*e^12) + (30*b^6*(b*d - a*e)^
3*(11*b*B*d - 4*A*b*e - 7*a*B*e)*Log[d + e*x])/e^12
```

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_
.)), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f]))))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1924 vs. $2(432) = 864$.

Time = 0.24 (sec) , antiderivative size = 1925, normalized size of antiderivative = 4.34

method	result	size
norman	Expression too large to display	1925
default	Expression too large to display	1953
risch	Expression too large to display	1999
parallelsch	Expression too large to display	3671

input

```
int((b*x+a)^10*(B*x+A)/(e*x+d)^8,x,method=_RETURNVERBOSE)
```

output

```
(-1/84*(12*A*a^10*e^11+20*A*a^9*b*d*e^10+36*A*a^8*b^2*d^2*e^9+72*A*a^7*b^3*d^3*e^8+168*A*a^6*b^4*d^4*e^7+504*A*a^5*b^5*d^5*e^6+2520*A*a^4*b^6*d^6*e^5-26136*A*a^3*b^7*d^7*e^4+78408*A*a^2*b^8*d^8*e^3-78408*A*a*b^9*d^9*e^2+26136*A*b^10*d^10*e+2*B*a^10*d*e^10+8*B*a^9*b*d^2*e^9+27*B*a^8*b^2*d^3*e^8+96*B*a^7*b^3*d^4*e^7+420*B*a^6*b^4*d^5*e^6+3024*B*a^5*b^5*d^6*e^5-45738*B*a^4*b^6*d^7*e^4+209088*B*a^3*b^7*d^8*e^3-352836*B*a^2*b^8*d^9*e^2+261360*B*a*b^9*d^10*e-71874*B*b^10*d^11)/e^12-7*(30*A*a^4*b^6*e^5-120*A*a^3*b^7*d*e^4+360*A*a^2*b^8*d^2*e^3-360*A*a*b^9*d^3*e^2+120*A*b^10*d^4*e+36*B*a^5*b^5*e^5-210*B*a^4*b^6*d*e^4+960*B*a^3*b^7*d^2*e^3-1620*B*a^2*b^8*d^3*e^2+1200*B*a*b^9*d^4*e-330*B*b^10*d^5)/e^6*x^6-21*(6*A*a^5*b^5*e^6+30*A*a^4*b^6*d*e^5-180*A*a^3*b^7*d^2*e^4+540*A*a^2*b^8*d^3*e^3-540*A*a*b^9*d^4*e^2+180*A*b^10*d^5*e+5*B*a^6*b^4*e^6+36*B*a^5*b^5*d*e^5-315*B*a^4*b^6*d^2*e^4+1440*B*a^3*b^7*d^3*e^3-2430*B*a^2*b^8*d^4*e^2+1800*B*a*b^9*d^5*e-495*B*b^10*d^6)/e^7*x^5-5*(14*A*a^6*b^4*e^7+42*A*a^5*b^5*d*e^6+210*A*a^4*b^6*d^2*e^5-1540*A*a^3*b^7*d^3*e^4+4620*A*a^2*b^8*d^4*e^3-4620*A*a*b^9*d^5*e^2+1540*A*b^10*d^6*e+8*B*a^7*b^3*e^7+35*B*a^6*b^4*d*e^6+252*B*a^5*b^5*d^2*e^5-2695*B*a^4*b^6*d^3*e^4+12320*B*a^3*b^7*d^4*e^3-20790*B*a^2*b^8*d^5*e^2+15400*B*a*b^9*d^6*e-4235*B*b^10*d^7)/e^8*x^4-5/4*(24*A*a^7*b^3*e^8+56*A*a^6*b^4*d*e^7+168*A*a^5*b^5*d^2*e^6+840*A*a^4*b^6*d^3*e^5-7000*A*a^3*b^7*d^4*e^4+21000*A*a^2*b^8*d^5*e^3-21000*A*a*b^9*d^6*e^2+7000*A*b^10*d^7*e+9*B*a^8*b^2*e^8...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2783 vs. $2(432) = 864$.

Time = 0.22 (sec) , antiderivative size = 2783, normalized size of antiderivative = 6.27

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^8} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)/(e*x+d)^8,x, algorithm="fricas")
```

output

```

1/84*(21*B*b^10*e^11*x^11 + 25961*B*b^10*d^11 - 12*A*a^10*e^11 - 11044*(10
*B*a*b^9 + A*b^10)*d^10*e + 20094*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 - 1731
6*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 6534*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d
^7*e^4 - 504*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 - 84*(5*B*a^6*b^4 + 6*A*a
^5*b^5)*d^5*e^6 - 24*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 - 9*(3*B*a^8*b^2
+ 8*A*a^7*b^3)*d^3*e^8 - 4*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 - 2*(B*a^10 +
10*A*a^9*b)*d*e^10 - 7*(11*B*b^10*d*e^10 - 4*(10*B*a*b^9 + A*b^10)*e^11)*
x^10 + 35*(11*B*b^10*d^2*e^9 - 4*(10*B*a*b^9 + A*b^10)*d*e^10 + 6*(9*B*a^2
*b^8 + 2*A*a*b^9)*e^11)*x^9 - 315*(11*B*b^10*d^3*e^8 - 4*(10*B*a*b^9 + A*b
^10)*d^2*e^9 + 6*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 - 4*(8*B*a^3*b^7 + 3*A*a
^2*b^8)*e^11)*x^8 - 49*(937*B*b^10*d^4*e^7 - 308*(10*B*a*b^9 + A*b^10)*d^3
*e^8 + 390*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 - 180*(8*B*a^3*b^7 + 3*A*a^2*
b^8)*d*e^10)*x^7 - 49*(2599*B*b^10*d^5*e^6 - 716*(10*B*a*b^9 + A*b^10)*d^4
*e^7 + 570*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 + 180*(8*B*a^3*b^7 + 3*A*a^2*
b^8)*d^2*e^9 - 360*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 + 72*(6*B*a^5*b^5 +
5*A*a^4*b^6)*e^11)*x^6 - 147*(619*B*b^10*d^6*e^5 + 4*(10*B*a*b^9 + A*b^10)
*d^5*e^6 - 510*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 + 900*(8*B*a^3*b^7 + 3*A*
a^2*b^8)*d^3*e^8 - 540*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 + 72*(6*B*a^5*b
^5 + 5*A*a^4*b^6)*d*e^10 + 12*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 + 35*(
4907*B*b^10*d^7*e^4 - 3388*(10*B*a*b^9 + A*b^10)*d^6*e^5 + 8610*(9*B*a^...

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^8} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**10*(B*x+A)/(e*x+d)**8,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1883 vs. $2(432) = 864$.

Time = 0.18 (sec) , antiderivative size = 1883, normalized size of antiderivative = 4.24

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^8} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^8,x, algorithm="maxima")`

output

```
1/84*(25961*B*b^10*d^11 - 12*A*a^10*e^11 - 11044*(10*B*a*b^9 + A*b^10)*d^10*e + 20094*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 - 17316*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 6534*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e^4 - 504*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 - 84*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^5*e^6 - 24*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 - 9*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^3*e^8 - 4*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 - 2*(B*a^10 + 10*A*a^9*b)*d*e^10 + 3528*(11*B*b^10*d^5*e^6 - 5*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 10*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 - 10*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 5*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 - (6*B*a^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 + 1764*(121*B*b^10*d^6*e^5 - 54*(10*B*a*b^9 + A*b^10)*d^5*e^6 + 105*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 - 100*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 + 45*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 - 6*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10 - (5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 + 420*(1177*B*b^10*d^7*e^4 - 518*(10*B*a*b^9 + A*b^10)*d^6*e^5 + 987*(9*B*a^2*b^8 + 2*A*a*b^9)*d^5*e^6 - 910*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^4*e^7 + 385*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^3*e^8 - 42*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e^9 - 7*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d*e^10 - 2*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e^11)*x^4 + 105*(5863*B*b^10*d^8*e^3 - 2552*(10*B*a*b^9 + A*b^10)*d^7*e^4 + 4788*(9*B*a^2*b^8 + 2*A*a*b^9)*d^6*e^5 - 4312*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^5*e^6 + 1750*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^4*e^7 - 168*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^3*e^8 - 28*(5*B*...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1992 vs. $2(432) = 864$.

Time = 0.13 (sec) , antiderivative size = 1992, normalized size of antiderivative = 4.49

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^8} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^8,x, algorithm="giac")`

output

```
30*(11*B*b^10*d^4 - 40*B*a*b^9*d^3*e - 4*A*b^10*d^3*e + 54*B*a^2*b^8*d^2*e
^2 + 12*A*a*b^9*d^2*e^2 - 32*B*a^3*b^7*d*e^3 - 12*A*a^2*b^8*d*e^3 + 7*B*a^
4*b^6*e^4 + 4*A*a^3*b^7*e^4)*log(abs(e*x + d))/e^12 + 1/84*(25961*B*b^10*d
^11 - 110440*B*a*b^9*d^10*e - 11044*A*b^10*d^10*e + 180846*B*a^2*b^8*d^9*e
^2 + 40188*A*a*b^9*d^9*e^2 - 138528*B*a^3*b^7*d^8*e^3 - 51948*A*a^2*b^8*d^
8*e^3 + 45738*B*a^4*b^6*d^7*e^4 + 26136*A*a^3*b^7*d^7*e^4 - 3024*B*a^5*b^5
*d^6*e^5 - 2520*A*a^4*b^6*d^6*e^5 - 420*B*a^6*b^4*d^5*e^6 - 504*A*a^5*b^5*
d^5*e^6 - 96*B*a^7*b^3*d^4*e^7 - 168*A*a^6*b^4*d^4*e^7 - 27*B*a^8*b^2*d^3*
e^8 - 72*A*a^7*b^3*d^3*e^8 - 8*B*a^9*b*d^2*e^9 - 36*A*a^8*b^2*d^2*e^9 - 2*
B*a^10*d*e^10 - 20*A*a^9*b*d*e^10 - 12*A*a^10*e^11 + 3528*(11*B*b^10*d^5*e
^6 - 50*B*a*b^9*d^4*e^7 - 5*A*b^10*d^4*e^7 + 90*B*a^2*b^8*d^3*e^8 + 20*A*a
*b^9*d^3*e^8 - 80*B*a^3*b^7*d^2*e^9 - 30*A*a^2*b^8*d^2*e^9 + 35*B*a^4*b^6*
d*e^10 + 20*A*a^3*b^7*d*e^10 - 6*B*a^5*b^5*e^11 - 5*A*a^4*b^6*e^11)*x^6 +
1764*(121*B*b^10*d^6*e^5 - 540*B*a*b^9*d^5*e^6 - 54*A*b^10*d^5*e^6 + 945*B
*a^2*b^8*d^4*e^7 + 210*A*a*b^9*d^4*e^7 - 800*B*a^3*b^7*d^3*e^8 - 300*A*a^2
*b^8*d^3*e^8 + 315*B*a^4*b^6*d^2*e^9 + 180*A*a^3*b^7*d^2*e^9 - 36*B*a^5*b^
5*d*e^10 - 30*A*a^4*b^6*d*e^10 - 5*B*a^6*b^4*e^11 - 6*A*a^5*b^5*e^11)*x^5
+ 420*(1177*B*b^10*d^7*e^4 - 5180*B*a*b^9*d^6*e^5 - 518*A*b^10*d^6*e^5 + 8
883*B*a^2*b^8*d^5*e^6 + 1974*A*a*b^9*d^5*e^6 - 7280*B*a^3*b^7*d^4*e^7 - 27
30*A*a^2*b^8*d^4*e^7 + 2695*B*a^4*b^6*d^3*e^8 + 1540*A*a^3*b^7*d^3*e^8 ...
```

Mupad [B] (verification not implemented)

Time = 1.22 (sec) , antiderivative size = 2092, normalized size of antiderivative = 4.71

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^8} dx = \text{Too large to display}$$

input `int(((A + B*x)*(a + b*x)^10)/(d + e*x)^8,x)`

output

```
x^3*((A*b^10 + 10*B*a*b^9)/(3*e^8) - (8*B*b^10*d)/(3*e^9)) - x*((28*d^2*((A*b^10 + 10*B*a*b^9)/e^8 - (8*B*b^10*d)/e^9))/e^2 - (8*d*((8*d*((A*b^10 + 10*B*a*b^9)/e^8 - (8*B*b^10*d)/e^9))/e - (5*a*b^8*(2*A*b + 9*B*a))/e^8 + (28*B*b^10*d^2)/e^10))/e - (15*a^2*b^7*(3*A*b + 8*B*a))/e^8 + (56*B*b^10*d^3)/e^11) - x^2*((4*d*((A*b^10 + 10*B*a*b^9)/e^8 - (8*B*b^10*d)/e^9))/e - (5*a*b^8*(2*A*b + 9*B*a))/(2*e^8) + (14*B*b^10*d^2)/e^10) - (x^4*(70*A*a^6*b^4*e^10 + 40*B*a^7*b^3*e^10 + 2590*A*b^10*d^6*e^4 - 5885*B*b^10*d^7*e^3 - 9870*A*a*b^9*d^5*e^5 + 210*A*a^5*b^5*d*e^9 + 25900*B*a*b^9*d^6*e^4 + 175*B*a^6*b^4*d*e^9 + 13650*A*a^2*b^8*d^4*e^6 - 7700*A*a^3*b^7*d^3*e^7 + 1050*A*a^4*b^6*d^2*e^8 - 44415*B*a^2*b^8*d^5*e^5 + 36400*B*a^3*b^7*d^4*e^6 - 13475*B*a^4*b^6*d^3*e^7 + 1260*B*a^5*b^5*d^2*e^8) + x^6*(210*A*a^4*b^6*e^10 + 252*B*a^5*b^5*e^10 + 210*A*b^10*d^4*e^6 - 462*B*b^10*d^5*e^5 - 840*A*a*b^9*d^3*e^7 - 840*A*a^3*b^7*d*e^9 + 2100*B*a*b^9*d^4*e^6 - 1470*B*a^4*b^6*d*e^9 + 1260*A*a^2*b^8*d^2*e^8 - 3780*B*a^2*b^8*d^3*e^7 + 3360*B*a^3*b^7*d^2*e^8) + x^3*(30*A*a^7*b^3*e^10 + (45*B*a^8*b^2*e^10)/4 + 3190*A*b^10*d^7*e^3 - (29315*B*b^10*d^8*e^2)/4 - 11970*A*a*b^9*d^6*e^4 + 70*A*a^6*b^4*d*e^9 + 31900*B*a*b^9*d^7*e^3 + 40*B*a^7*b^3*d*e^9 + 16170*A*a^2*b^8*d^5*e^5 - 8750*A*a^3*b^7*d^4*e^6 + 1050*A*a^4*b^6*d^3*e^7 + 210*A*a^5*b^5*d^2*e^8 - 53865*B*a^2*b^8*d^6*e^4 + 43120*B*a^3*b^7*d^5*e^5 - (30625*B*a^4*b^6*d^4*e^6)/2 + 1260*B*a^5*b^5*d^3*e^7 + 175*B*a^6*b^4*d^2*e^8) + (12*A*a^10*e...
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 1922, normalized size of antiderivative = 4.33

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^8} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^8,x)`

output

```
(27720*log(d + e*x)*a**4*b**7*d**8*e**4 + 194040*log(d + e*x)*a**4*b**7*d*
*7*e**5*x + 582120*log(d + e*x)*a**4*b**7*d**6*e**6*x**2 + 970200*log(d +
e*x)*a**4*b**7*d**5*e**7*x**3 + 970200*log(d + e*x)*a**4*b**7*d**4*e**8*x*
*4 + 582120*log(d + e*x)*a**4*b**7*d**3*e**9*x**5 + 194040*log(d + e*x)*a*
*4*b**7*d**2*e**10*x**6 + 27720*log(d + e*x)*a**4*b**7*d*e**11*x**7 - 1108
80*log(d + e*x)*a**3*b**8*d**9*e**3 - 776160*log(d + e*x)*a**3*b**8*d**8*e
**4*x - 2328480*log(d + e*x)*a**3*b**8*d**7*e**5*x**2 - 3880800*log(d + e*
x)*a**3*b**8*d**6*e**6*x**3 - 3880800*log(d + e*x)*a**3*b**8*d**5*e**7*x**
4 - 2328480*log(d + e*x)*a**3*b**8*d**4*e**8*x**5 - 776160*log(d + e*x)*a*
*3*b**8*d**3*e**9*x**6 - 110880*log(d + e*x)*a**3*b**8*d**2*e**10*x**7 + 1
66320*log(d + e*x)*a**2*b**9*d**10*e**2 + 1164240*log(d + e*x)*a**2*b**9*d
**9*e**3*x + 3492720*log(d + e*x)*a**2*b**9*d**8*e**4*x**2 + 5821200*log(d
+ e*x)*a**2*b**9*d**7*e**5*x**3 + 5821200*log(d + e*x)*a**2*b**9*d**6*e**
6*x**4 + 3492720*log(d + e*x)*a**2*b**9*d**5*e**7*x**5 + 1164240*log(d + e
*x)*a**2*b**9*d**4*e**8*x**6 + 166320*log(d + e*x)*a**2*b**9*d**3*e**9*x**
7 - 110880*log(d + e*x)*a*b**10*d**11*e - 776160*log(d + e*x)*a*b**10*d**1
0*e**2*x - 2328480*log(d + e*x)*a*b**10*d**9*e**3*x**2 - 3880800*log(d + e
*x)*a*b**10*d**8*e**4*x**3 - 3880800*log(d + e*x)*a*b**10*d**7*e**5*x**4 -
2328480*log(d + e*x)*a*b**10*d**6*e**6*x**5 - 776160*log(d + e*x)*a*b**10
*d**5*e**7*x**6 - 110880*log(d + e*x)*a*b**10*d**4*e**8*x**7 + 27720*lo...
```

3.87 $\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^9} dx$

Optimal result	893
Mathematica [A] (verified)	894
Rubi [A] (verified)	895
Maple [B] (verified)	896
Fricas [B] (verification not implemented)	897
Sympy [F(-1)]	898
Maxima [B] (verification not implemented)	899
Giac [B] (verification not implemented)	900
Mupad [B] (verification not implemented)	901
Reduce [B] (verification not implemented)	901

Optimal result

Integrand size = 20, antiderivative size = 445

$$\begin{aligned}
 \int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^9} dx = & \frac{5b^8(bd-ae)(11bBd-2Abe-9aBe)x}{e^{11}} \\
 & + \frac{(bd-ae)^{10}(Bd-Ae)}{8e^{12}(d+ex)^8} \\
 & - \frac{(bd-ae)^9(11bBd-10Abe-aBe)}{7e^{12}(d+ex)^7} \\
 & + \frac{5b(bd-ae)^8(11bBd-9Abe-2aBe)}{6e^{12}(d+ex)^6} \\
 & - \frac{3b^2(bd-ae)^7(11bBd-8Abe-3aBe)}{e^{12}(d+ex)^5} \\
 & + \frac{15b^3(bd-ae)^6(11bBd-7Abe-4aBe)}{2e^{12}(d+ex)^4} \\
 & - \frac{14b^4(bd-ae)^5(11bBd-6Abe-5aBe)}{e^{12}(d+ex)^3} \\
 & + \frac{21b^5(bd-ae)^4(11bBd-5Abe-6aBe)}{e^{12}(d+ex)^2} \\
 & - \frac{30b^6(bd-ae)^3(11bBd-4Abe-7aBe)}{e^{12}(d+ex)} \\
 & - \frac{b^9(11bBd-Abe-10aBe)(d+ex)^2}{2e^{12}} + \frac{b^{10}B(d+ex)^3}{3e^{12}} \\
 & - \frac{15b^7(bd-ae)^2(11bBd-3Abe-8aBe)\log(d+ex)}{e^{12}}
 \end{aligned}$$

output

```
5*b^8*(-a*e+b*d)*(-2*A*b*e-9*B*a*e+11*B*b*d)*x/e^11+1/8*(-a*e+b*d)^10*(-A*
e+B*d)/e^12/(e*x+d)^8-1/7*(-a*e+b*d)^9*(-10*A*b*e-B*a*e+11*B*b*d)/e^12/(e*
x+d)^7+5/6*b*(-a*e+b*d)^8*(-9*A*b*e-2*B*a*e+11*B*b*d)/e^12/(e*x+d)^6-3*b^2
*(-a*e+b*d)^7*(-8*A*b*e-3*B*a*e+11*B*b*d)/e^12/(e*x+d)^5+15/2*b^3*(-a*e+b*
d)^6*(-7*A*b*e-4*B*a*e+11*B*b*d)/e^12/(e*x+d)^4-14*b^4*(-a*e+b*d)^5*(-6*A*
b*e-5*B*a*e+11*B*b*d)/e^12/(e*x+d)^3+21*b^5*(-a*e+b*d)^4*(-5*A*b*e-6*B*a*e
+11*B*b*d)/e^12/(e*x+d)^2-30*b^6*(-a*e+b*d)^3*(-4*A*b*e-7*B*a*e+11*B*b*d)/
e^12/(e*x+d)-1/2*b^9*(-A*b*e-10*B*a*e+11*B*b*d)*(e*x+d)^2/e^12+1/3*b^10*B*
(e*x+d)^3/e^12-15*b^7*(-a*e+b*d)^2*(-3*A*b*e-8*B*a*e+11*B*b*d)*ln(e*x+d)/e
^12
```

Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 415, normalized size of antiderivative = 0.93

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^9} dx$$

$$= \frac{-168b^8e(-45a^2Be^2 - 10abe(-9Bd + Ae) + 9b^2d(-5Bd + Ae))x + 84b^9e^2(-9bBd + Abe + 10aBe)x^2}{(d+ex)^9}$$

input

```
Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^9,x]
```

output

```
(-168*b^8*e*(-45*a^2*B*e^2 - 10*a*b*e*(-9*B*d + A*e) + 9*b^2*d*(-5*B*d + A
*e))*x + 84*b^9*e^2*(-9*b*B*d + A*b*e + 10*a*B*e)*x^2 + 56*b^10*B*e^3*x^3
+ (21*(b*d - a*e)^10*(B*d - A*e))/(d + e*x)^8 - (24*(b*d - a*e)^9*(11*b*B*
d - 10*A*b*e - a*B*e))/(d + e*x)^7 + (140*b*(b*d - a*e)^8*(11*b*B*d - 9*A*
b*e - 2*a*B*e))/(d + e*x)^6 - (504*b^2*(b*d - a*e)^7*(11*b*B*d - 8*A*b*e -
3*a*B*e))/(d + e*x)^5 + (1260*b^3*(b*d - a*e)^6*(11*b*B*d - 7*A*b*e - 4*a
*B*e))/(d + e*x)^4 - (2352*b^4*(b*d - a*e)^5*(11*b*B*d - 6*A*b*e - 5*a*B*
e))/(d + e*x)^3 + (3528*b^5*(b*d - a*e)^4*(11*b*B*d - 5*A*b*e - 6*a*B*e))/(
d + e*x)^2 - (5040*b^6*(b*d - a*e)^3*(11*b*B*d - 4*A*b*e - 7*a*B*e))/(d +
e*x) - 2520*b^7*(b*d - a*e)^2*(11*b*B*d - 3*A*b*e - 8*a*B*e)*Log[d + e*x]
/(168*e^12)
```

Rubi [A] (verified)

Time = 1.25 (sec) , antiderivative size = 445, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^9} dx$$

↓ 86

$$\int \left(\frac{b^9(d + ex)(10aBe + Abe - 11bBd)}{e^{11}} - \frac{5b^8(bd - ae)(9aBe + 2Abe - 11bBd)}{e^{11}} + \frac{15b^7(bd - ae)^2(8aBe + 3Abe - 11bBd)}{e^{11}(d + ex)} \right) dx$$

↓ 2009

$$\begin{aligned} & - \frac{b^9(d + ex)^2(-10aBe - Abe + 11bBd)}{2e^{12}} + \frac{5b^8x(bd - ae)(-9aBe - 2Abe + 11bBd)}{e^{11}} - \\ & \frac{15b^7(bd - ae)^2 \log(d + ex)(-8aBe - 3Abe + 11bBd)}{e^{12}} - \\ & \frac{30b^6(bd - ae)^3(-7aBe - 4Abe + 11bBd)}{e^{12}(d + ex)} + \frac{21b^5(bd - ae)^4(-6aBe - 5Abe + 11bBd)}{e^{12}(d + ex)^2} - \\ & \frac{14b^4(bd - ae)^5(-5aBe - 6Abe + 11bBd)}{e^{12}(d + ex)^3} + \frac{15b^3(bd - ae)^6(-4aBe - 7Abe + 11bBd)}{2e^{12}(d + ex)^4} - \\ & \frac{3b^2(bd - ae)^7(-3aBe - 8Abe + 11bBd)}{e^{12}(d + ex)^5} + \frac{5b(bd - ae)^8(-2aBe - 9Abe + 11bBd)}{6e^{12}(d + ex)^6} - \\ & \frac{(bd - ae)^9(-aBe - 10Abe + 11bBd)}{7e^{12}(d + ex)^7} + \frac{(bd - ae)^{10}(Bd - Ae)}{8e^{12}(d + ex)^8} + \frac{b^{10}B(d + ex)^3}{3e^{12}} \end{aligned}$$

input

```
Int[((a + b*x)^10*(A + B*x))/(d + e*x)^9,x]
```

output

```
(5*b^8*(b*d - a*e)*(11*b*B*d - 2*A*b*e - 9*a*B*e)*x)/e^11 + ((b*d - a*e)^10*(B*d - A*e))/(8*e^12*(d + e*x)^8) - ((b*d - a*e)^9*(11*b*B*d - 10*A*b*e - a*B*e))/(7*e^12*(d + e*x)^7) + (5*b*(b*d - a*e)^8*(11*b*B*d - 9*A*b*e - 2*a*B*e))/(6*e^12*(d + e*x)^6) - (3*b^2*(b*d - a*e)^7*(11*b*B*d - 8*A*b*e - 3*a*B*e))/(e^12*(d + e*x)^5) + (15*b^3*(b*d - a*e)^6*(11*b*B*d - 7*A*b*e - 4*a*B*e))/(2*e^12*(d + e*x)^4) - (14*b^4*(b*d - a*e)^5*(11*b*B*d - 6*A*b*e - 5*a*B*e))/(e^12*(d + e*x)^3) + (21*b^5*(b*d - a*e)^4*(11*b*B*d - 5*A*b*e - 6*a*B*e))/(e^12*(d + e*x)^2) - (30*b^6*(b*d - a*e)^3*(11*b*B*d - 4*A*b*e - 7*a*B*e))/(e^12*(d + e*x)) - (b^9*(11*b*B*d - A*b*e - 10*a*B*e)*(d + e*x)^2)/(2*e^12) + (b^10*B*(d + e*x)^3)/(3*e^12) - (15*b^7*(b*d - a*e)^2*(11*b*B*d - 3*A*b*e - 8*a*B*e)*Log[d + e*x])/e^12
```

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1928 vs. $2(433) = 866$.

Time = 0.24 (sec) , antiderivative size = 1929, normalized size of antiderivative = 4.33

method	result	size
norman	Expression too large to display	1929
default	Expression too large to display	1938
risch	Expression too large to display	1968
parallelsch	Expression too large to display	3491

input

```
int((b*x+a)^10*(B*x+A)/(e*x+d)^9,x,method=_RETURNVERBOSE)
```

output

```
(-1/168*(21*A*a^10*e^11+30*A*a^9*b*d*e^10+45*A*a^8*b^2*d^2*e^9+72*A*a^7*b^3*d^3*e^8+126*A*a^6*b^4*d^4*e^7+252*A*a^5*b^5*d^5*e^6+630*A*a^4*b^6*d^6*e^5+2520*A*a^3*b^7*d^7*e^4-20547*A*a^2*b^8*d^8*e^3+41094*A*a*b^9*d^9*e^2-20547*A*b^10*d^10*e+3*B*a^10*d*e^10+10*B*a^9*b*d^2*e^9+27*B*a^8*b^2*d^3*e^8+72*B*a^7*b^3*d^4*e^7+210*B*a^6*b^4*d^5*e^6+756*B*a^5*b^5*d^6*e^5+4410*B*a^4*b^6*d^7*e^4-54792*B*a^3*b^7*d^8*e^3+184923*B*a^2*b^8*d^9*e^2-205470*B*a*b^9*d^10*e+75339*B*b^10*d^11)/e^12-2*(60*A*a^3*b^7*e^4-180*A*a^2*b^8*d*e^3+360*A*a*b^9*d^2*e^2-180*A*b^10*d^3*e+105*B*a^4*b^6*e^4-480*B*a^3*b^7*d*e^3+1620*B*a^2*b^8*d^2*e^2-1800*B*a*b^9*d^3*e+660*B*b^10*d^4)/e^5*x^7-7*(15*A*a^4*b^6*e^5+60*A*a^3*b^7*d*e^4-270*A*a^2*b^8*d^2*e^3+540*A*a*b^9*d^3*e^2-270*A*b^10*d^4*e+18*B*a^5*b^5*e^5+105*B*a^4*b^6*d*e^4-720*B*a^3*b^7*d^2*e^3+2430*B*a^2*b^8*d^3*e^2-2700*B*a*b^9*d^4*e+990*B*b^10*d^5)/e^6*x^6-14*(6*A*a^5*b^5*e^6+15*A*a^4*b^6*d*e^5+60*A*a^3*b^7*d^2*e^4-330*A*a^2*b^8*d^3*e^3+660*A*a*b^9*d^4*e^2-330*A*b^10*d^5*e+5*B*a^6*b^4*e^6+18*B*a^5*b^5*d*e^5+105*B*a^4*b^6*d^2*e^4-880*B*a^3*b^7*d^3*e^3+2970*B*a^2*b^8*d^4*e^2-3300*B*a*b^9*d^5*e+1210*B*b^10*d^6)/e^7*x^5-5/2*(21*A*a^6*b^4*e^7+42*A*a^5*b^5*d*e^6+105*A*a^4*b^6*d^2*e^5+420*A*a^3*b^7*d^3*e^4-2625*A*a^2*b^8*d^4*e^3+5250*A*a*b^9*d^5*e^2-2625*A*b^10*d^6*e+12*B*a^7*b^3*e^7+35*B*a^6*b^4*d*e^6+126*B*a^5*b^5*d^2*e^5+735*B*a^4*b^6*d^3*e^4-7000*B*a^3*b^7*d^4*e^3+23625*B*a^2*b^8*d^5*e^2-26250*B*a*b^9*d^6*e+9625*B*b^10*d^7)/e^8*x^4-(24*A*a^7*b...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2677 vs. $2(433) = 866$.

Time = 0.25 (sec) , antiderivative size = 2677, normalized size of antiderivative = 6.02

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^9} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)/(e*x+d)^9,x, algorithm="fricas")
```

output

```

1/168*(56*B*b^10*e^11*x^11 - 32891*B*b^10*d^11 - 21*A*a^10*e^11 + 10803*(1
0*B*a*b^9 + A*b^10)*d^10*e - 13827*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 + 684
9*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 - 630*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^
7*e^4 - 126*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 - 42*(5*B*a^6*b^4 + 6*A*a^
5*b^5)*d^5*e^6 - 18*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 - 9*(3*B*a^8*b^2 +
8*A*a^7*b^3)*d^3*e^8 - 5*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 - 3*(B*a^10 +
10*A*a^9*b)*d*e^10 - 28*(11*B*b^10*d*e^10 - 3*(10*B*a*b^9 + A*b^10)*e^11)*
x^10 + 280*(11*B*b^10*d^2*e^9 - 3*(10*B*a*b^9 + A*b^10)*d*e^10 + 3*(9*B*a^
2*b^8 + 2*A*a*b^9)*e^11)*x^9 + 112*(379*B*b^10*d^3*e^8 - 87*(10*B*a*b^9 +
A*b^10)*d^2*e^9 + 60*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10)*x^8 + 112*(1052*B*b
^10*d^4*e^7 - 156*(10*B*a*b^9 + A*b^10)*d^3*e^8 - 60*(9*B*a^2*b^8 + 2*A*a*
b^9)*d^2*e^9 + 180*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 - 45*(7*B*a^4*b^6 +
4*A*a^3*b^7)*e^11)*x^7 + 392*(62*B*b^10*d^5*e^6 + 114*(10*B*a*b^9 + A*b^10
)*d^4*e^7 - 330*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 + 270*(8*B*a^3*b^7 + 3*A
*a^2*b^8)*d^2*e^9 - 45*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 - 9*(6*B*a^5*b^5
+ 5*A*a^4*b^6)*e^11)*x^6 - 784*(598*B*b^10*d^6*e^5 - 294*(10*B*a*b^9 + A*
b^10)*d^5*e^6 + 510*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 - 330*(8*B*a^3*b^7 +
3*A*a^2*b^8)*d^3*e^8 + 45*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 + 9*(6*B*a^
5*b^5 + 5*A*a^4*b^6)*d*e^10 + 3*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 - 14
0*(7651*B*b^10*d^7*e^4 - 3003*(10*B*a*b^9 + A*b^10)*d^6*e^5 + 4515*(9*B...

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^9} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**10*(B*x+A)/(e*x+d)**9,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1892 vs. $2(433) = 866$.

Time = 0.19 (sec) , antiderivative size = 1892, normalized size of antiderivative = 4.25

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^9} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^9,x, algorithm="maxima")`

output

```
-1/168*(32891*B*b^10*d^11 + 21*A*a^10*e^11 - 10803*(10*B*a*b^9 + A*b^10)*d
^10*e + 13827*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 - 6849*(8*B*a^3*b^7 + 3*A*
a^2*b^8)*d^8*e^3 + 630*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e^4 + 126*(6*B*a^5*
b^5 + 5*A*a^4*b^6)*d^6*e^5 + 42*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^5*e^6 + 18*(
4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 9*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^3*e^8
+ 5*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 + 3*(B*a^10 + 10*A*a^9*b)*d*e^10 +
5040*(11*B*b^10*d^4*e^7 - 4*(10*B*a*b^9 + A*b^10)*d^3*e^8 + 6*(9*B*a^2*b^8
+ 2*A*a*b^9)*d^2*e^9 - 4*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 + (7*B*a^4*b^
6 + 4*A*a^3*b^7)*e^11)*x^7 + 3528*(99*B*b^10*d^5*e^6 - 35*(10*B*a*b^9 + A
b^10)*d^4*e^7 + 50*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 - 30*(8*B*a^3*b^7 + 3
*A*a^2*b^8)*d^2*e^9 + 5*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 + (6*B*a^5*b^5
+ 5*A*a^4*b^6)*e^11)*x^6 + 2352*(407*B*b^10*d^6*e^5 - 141*(10*B*a*b^9 + A
b^10)*d^5*e^6 + 195*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 - 110*(8*B*a^3*b^7 +
3*A*a^2*b^8)*d^3*e^8 + 15*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 + 3*(6*B*a^
5*b^5 + 5*A*a^4*b^6)*d*e^10 + (5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 + 420*
(3509*B*b^10*d^7*e^4 - 1197*(10*B*a*b^9 + A*b^10)*d^6*e^5 + 1617*(9*B*a^2*
b^8 + 2*A*a*b^9)*d^5*e^6 - 875*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^4*e^7 + 105*(
7*B*a^4*b^6 + 4*A*a^3*b^7)*d^3*e^8 + 21*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e^
9 + 7*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d*e^10 + 3*(4*B*a^7*b^3 + 7*A*a^6*b^4)*e
^11)*x^4 + 168*(8173*B*b^10*d^8*e^3 - 2754*(10*B*a*b^9 + A*b^10)*d^7*e^...
```


Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1981 vs. $2(433) = 866$.

Time = 0.13 (sec) , antiderivative size = 1981, normalized size of antiderivative = 4.45

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^9} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^9,x, algorithm="giac")`

output

```
-15*(11*B*b^10*d^3 - 30*B*a*b^9*d^2*e - 3*A*b^10*d^2*e + 27*B*a^2*b^8*d*e^2 + 6*A*a*b^9*d*e^2 - 8*B*a^3*b^7*e^3 - 3*A*a^2*b^8*e^3)*log(abs(e*x + d))
/e^12 - 1/168*(32891*B*b^10*d^11 - 108030*B*a*b^9*d^10*e - 10803*A*b^10*d^10*e + 124443*B*a^2*b^8*d^9*e^2 + 27654*A*a*b^9*d^9*e^2 - 54792*B*a^3*b^7*d^8*e^3 - 20547*A*a^2*b^8*d^8*e^3 + 4410*B*a^4*b^6*d^7*e^4 + 2520*A*a^3*b^7*d^7*e^4 + 756*B*a^5*b^5*d^6*e^5 + 630*A*a^4*b^6*d^6*e^5 + 210*B*a^6*b^4*d^5*e^6 + 252*A*a^5*b^5*d^5*e^6 + 72*B*a^7*b^3*d^4*e^7 + 126*A*a^6*b^4*d^4*e^7 + 27*B*a^8*b^2*d^3*e^8 + 72*A*a^7*b^3*d^3*e^8 + 10*B*a^9*b*d^2*e^9 + 45*A*a^8*b^2*d^2*e^9 + 3*B*a^10*d*e^10 + 30*A*a^9*b*d*e^10 + 21*A*a^10*e^11 + 5040*(11*B*b^10*d^4*e^7 - 40*B*a*b^9*d^3*e^8 - 4*A*b^10*d^3*e^8 + 54*B*a^2*b^8*d^2*e^9 + 12*A*a*b^9*d^2*e^9 - 32*B*a^3*b^7*d*e^10 - 12*A*a^2*b^8*d*e^10 + 7*B*a^4*b^6*e^11 + 4*A*a^3*b^7*e^11)*x^7 + 3528*(99*B*b^10*d^5*e^6 - 350*B*a*b^9*d^4*e^7 - 35*A*b^10*d^4*e^7 + 450*B*a^2*b^8*d^3*e^8 + 100*A*a*b^9*d^3*e^8 - 240*B*a^3*b^7*d^2*e^9 - 90*A*a^2*b^8*d^2*e^9 + 35*B*a^4*b^6*d*e^10 + 20*A*a^3*b^7*d*e^10 + 6*B*a^5*b^5*e^11 + 5*A*a^4*b^6*e^11)*x^6 + 2352*(407*B*b^10*d^6*e^5 - 1410*B*a*b^9*d^5*e^6 - 141*A*b^10*d^5*e^6 + 1755*B*a^2*b^8*d^4*e^7 + 390*A*a*b^9*d^4*e^7 - 880*B*a^3*b^7*d^3*e^8 - 330*A*a^2*b^8*d^3*e^8 + 105*B*a^4*b^6*d^2*e^9 + 60*A*a^3*b^7*d^2*e^9 + 18*B*a^5*b^5*d*e^10 + 15*A*a^4*b^6*d*e^10 + 5*B*a^6*b^4*e^11 + 6*A*a^5*b^5*e^11)*x^5 + 420*(3509*B*b^10*d^7*e^4 - 11970*B*a*b^9*d^6*e^5 - 1197*A*b^10...
```

Mupad [B] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 2048, normalized size of antiderivative = 4.60

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^9} dx = \text{Too large to display}$$

input `int(((A + B*x)*(a + b*x)^10)/(d + e*x)^9,x)`

output

```
x^2*((A*b^10 + 10*B*a*b^9)/(2*e^9) - (9*B*b^10*d)/(2*e^10)) - (x^7*(120*A*
a^3*b^7*e^10 + 210*B*a^4*b^6*e^10 - 120*A*b^10*d^3*e^7 + 330*B*b^10*d^4*e^
6 + 360*A*a*b^9*d^2*e^8 - 360*A*a^2*b^8*d*e^9 - 1200*B*a*b^9*d^3*e^7 - 960
*B*a^3*b^7*d*e^9 + 1620*B*a^2*b^8*d^2*e^8) + x^4*((105*A*a^6*b^4*e^10)/2 +
30*B*a^7*b^3*e^10 - (5985*A*b^10*d^6*e^4)/2 + (17545*B*b^10*d^7*e^3)/2 +
8085*A*a*b^9*d^5*e^5 + 105*A*a^5*b^5*d*e^9 - 29925*B*a*b^9*d^6*e^4 + (175*
B*a^6*b^4*d*e^9)/2 - (13125*A*a^2*b^8*d^4*e^6)/2 + 1050*A*a^3*b^7*d^3*e^7
+ (525*A*a^4*b^6*d^2*e^8)/2 + (72765*B*a^2*b^8*d^5*e^5)/2 - 17500*B*a^3*b^
7*d^4*e^6 + (3675*B*a^4*b^6*d^3*e^7)/2 + 315*B*a^5*b^5*d^2*e^8) + x^6*(105
*A*a^4*b^6*e^10 + 126*B*a^5*b^5*e^10 - 735*A*b^10*d^4*e^6 + 2079*B*b^10*d^
5*e^5 + 2100*A*a*b^9*d^3*e^7 + 420*A*a^3*b^7*d*e^9 - 7350*B*a*b^9*d^4*e^6
+ 735*B*a^4*b^6*d*e^9 - 1890*A*a^2*b^8*d^2*e^8 + 9450*B*a^2*b^8*d^3*e^7 -
5040*B*a^3*b^7*d^2*e^8) + x^3*(24*A*a^7*b^3*e^10 + 9*B*a^8*b^2*e^10 - 2754
*A*b^10*d^7*e^3 + 8173*B*b^10*d^8*e^2 + 7308*A*a*b^9*d^6*e^4 + 42*A*a^6*b^
4*d*e^9 - 27540*B*a*b^9*d^7*e^3 + 24*B*a^7*b^3*d*e^9 - 5754*A*a^2*b^8*d^5*
e^5 + 840*A*a^3*b^7*d^4*e^6 + 210*A*a^4*b^6*d^3*e^7 + 84*A*a^5*b^5*d^2*e^8
+ 32886*B*a^2*b^8*d^6*e^4 - 15344*B*a^3*b^7*d^5*e^5 + 1470*B*a^4*b^6*d^4*
e^6 + 252*B*a^5*b^5*d^3*e^7 + 70*B*a^6*b^4*d^2*e^8) + (21*A*a^10*e^11 + 32
891*B*b^10*d^11 - 10803*A*b^10*d^10*e + 3*B*a^10*d*e^10 + 27654*A*a*b^9*d^
9*e^2 + 10*B*a^9*b*d^2*e^9 - 20547*A*a^2*b^8*d^8*e^3 + 2520*A*a^3*b^7*d...
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 1841, normalized size of antiderivative = 4.14

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^9} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^9,x)`

output

```
(27720*log(d + e*x)*a**3*b**8*d**9*e**3 + 221760*log(d + e*x)*a**3*b**8*d*
*8*e**4*x + 776160*log(d + e*x)*a**3*b**8*d**7*e**5*x**2 + 1552320*log(d +
e*x)*a**3*b**8*d**6*e**6*x**3 + 1940400*log(d + e*x)*a**3*b**8*d**5*e**7*
x**4 + 1552320*log(d + e*x)*a**3*b**8*d**4*e**8*x**5 + 776160*log(d + e*x)
*a**3*b**8*d**3*e**9*x**6 + 221760*log(d + e*x)*a**3*b**8*d**2*e**10*x**7
+ 27720*log(d + e*x)*a**3*b**8*d*e**11*x**8 - 83160*log(d + e*x)*a**2*b**9
*d**10*e**2 - 665280*log(d + e*x)*a**2*b**9*d**9*e**3*x - 2328480*log(d +
e*x)*a**2*b**9*d**8*e**4*x**2 - 4656960*log(d + e*x)*a**2*b**9*d**7*e**5*x
**3 - 5821200*log(d + e*x)*a**2*b**9*d**6*e**6*x**4 - 4656960*log(d + e*x)
*a**2*b**9*d**5*e**7*x**5 - 2328480*log(d + e*x)*a**2*b**9*d**4*e**8*x**6
- 665280*log(d + e*x)*a**2*b**9*d**3*e**9*x**7 - 83160*log(d + e*x)*a**2*b
**9*d**2*e**10*x**8 + 83160*log(d + e*x)*a*b**10*d**11*e + 665280*log(d +
e*x)*a*b**10*d**10*e**2*x + 2328480*log(d + e*x)*a*b**10*d**9*e**3*x**2 +
4656960*log(d + e*x)*a*b**10*d**8*e**4*x**3 + 5821200*log(d + e*x)*a*b**10
*d**7*e**5*x**4 + 4656960*log(d + e*x)*a*b**10*d**6*e**6*x**5 + 2328480*lo
g(d + e*x)*a*b**10*d**5*e**7*x**6 + 665280*log(d + e*x)*a*b**10*d**4*e**8*
x**7 + 83160*log(d + e*x)*a*b**10*d**3*e**9*x**8 - 27720*log(d + e*x)*b**1
1*d**12 - 221760*log(d + e*x)*b**11*d**11*e*x - 776160*log(d + e*x)*b**11*
d**10*e**2*x**2 - 1552320*log(d + e*x)*b**11*d**9*e**3*x**3 - 1940400*log(
d + e*x)*b**11*d**8*e**4*x**4 - 1552320*log(d + e*x)*b**11*d**7*e**5*x**...
```

$$3.88 \quad \int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{10}} dx$$

Optimal result	904
Mathematica [B] (verified)	905
Rubi [A] (verified)	906
Maple [B] (verified)	907
Fricas [B] (verification not implemented)	908
Sympy [F(-1)]	909
Maxima [B] (verification not implemented)	910
Giac [B] (verification not implemented)	911
Mupad [B] (verification not implemented)	912
Reduce [B] (verification not implemented)	912

Optimal result

Integrand size = 20, antiderivative size = 441

$$\begin{aligned}
 \int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{10}} dx = & -\frac{b^9(10bBd - Abe - 10aBe)x}{e^{11}} \\
 & + \frac{b^{10}Bx^2}{2e^{10}} + \frac{(bd - ae)^{10}(Bd - Ae)}{9e^{12}(d+ex)^9} \\
 & - \frac{(bd - ae)^9(11bBd - 10Abe - aBe)}{8e^{12}(d+ex)^8} \\
 & + \frac{5b(bd - ae)^8(11bBd - 9Abe - 2aBe)}{7e^{12}(d+ex)^7} \\
 & - \frac{5b^2(bd - ae)^7(11bBd - 8Abe - 3aBe)}{2e^{12}(d+ex)^6} \\
 & + \frac{6b^3(bd - ae)^6(11bBd - 7Abe - 4aBe)}{e^{12}(d+ex)^5} \\
 & - \frac{21b^4(bd - ae)^5(11bBd - 6Abe - 5aBe)}{2e^{12}(d+ex)^4} \\
 & + \frac{14b^5(bd - ae)^4(11bBd - 5Abe - 6aBe)}{e^{12}(d+ex)^3} \\
 & - \frac{15b^6(bd - ae)^3(11bBd - 4Abe - 7aBe)}{e^{12}(d+ex)^2} \\
 & + \frac{15b^7(bd - ae)^2(11bBd - 3Abe - 8aBe)}{e^{12}(d+ex)} \\
 & + \frac{5b^8(bd - ae)(11bBd - 2Abe - 9aBe) \log(d+ex)}{e^{12}}
 \end{aligned}$$

output

```

-b^9*(-A*b*e-10*B*a*e+10*B*b*d)*x/e^11+1/2*b^10*B*x^2/e^10+1/9*(-a*e+b*d)^
10*(-A*e+B*d)/e^12/(e*x+d)^9-1/8*(-a*e+b*d)^9*(-10*A*b*e-B*a*e+11*B*b*d)/e
^12/(e*x+d)^8+5/7*b*(-a*e+b*d)^8*(-9*A*b*e-2*B*a*e+11*B*b*d)/e^12/(e*x+d)^
7-5/2*b^2*(-a*e+b*d)^7*(-8*A*b*e-3*B*a*e+11*B*b*d)/e^12/(e*x+d)^6+6*b^3*(-
a*e+b*d)^6*(-7*A*b*e-4*B*a*e+11*B*b*d)/e^12/(e*x+d)^5-21/2*b^4*(-a*e+b*d)^
5*(-6*A*b*e-5*B*a*e+11*B*b*d)/e^12/(e*x+d)^4+14*b^5*(-a*e+b*d)^4*(-5*A*b*e
-6*B*a*e+11*B*b*d)/e^12/(e*x+d)^3-15*b^6*(-a*e+b*d)^3*(-4*A*b*e-7*B*a*e+11
*B*b*d)/e^12/(e*x+d)^2+15*b^7*(-a*e+b*d)^2*(-3*A*b*e-8*B*a*e+11*B*b*d)/e^1
2/(e*x+d)+5*b^8*(-a*e+b*d)*(-2*A*b*e-9*B*a*e+11*B*b*d)*ln(e*x+d)/e^12

```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1460 vs. $2(441) = 882$.

Time = 0.56 (sec) , antiderivative size = 1460, normalized size of antiderivative = 3.31

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{10}} dx = \text{Too large to display}$$

input `Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^10,x]`

output

```
-1/504*(7*a^10*e^10*(8*A*e + B*(d + 9*e*x)) + 10*a^9*b*e^9*(7*A*e*(d + 9*e
*x) + 2*B*(d^2 + 9*d*e*x + 36*e^2*x^2)) + 45*a^8*b^2*e^8*(2*A*e*(d^2 + 9*d
*e*x + 36*e^2*x^2) + B*(d^3 + 9*d^2*e*x + 36*d*e^2*x^2 + 84*e^3*x^3)) + 24
*a^7*b^3*e^7*(5*A*e*(d^3 + 9*d^2*e*x + 36*d*e^2*x^2 + 84*e^3*x^3) + 4*B*(d
^4 + 9*d^3*e*x + 36*d^2*e^2*x^2 + 84*d*e^3*x^3 + 126*e^4*x^4)) + 42*a^6*b^
4*e^6*(4*A*e*(d^4 + 9*d^3*e*x + 36*d^2*e^2*x^2 + 84*d*e^3*x^3 + 126*e^4*x^
4) + 5*B*(d^5 + 9*d^4*e*x + 36*d^3*e^2*x^2 + 84*d^2*e^3*x^3 + 126*d*e^4*x^
4 + 126*e^5*x^5)) + 252*a^5*b^5*e^5*(A*e*(d^5 + 9*d^4*e*x + 36*d^3*e^2*x^2
+ 84*d^2*e^3*x^3 + 126*d*e^4*x^4 + 126*e^5*x^5) + 2*B*(d^6 + 9*d^5*e*x +
36*d^4*e^2*x^2 + 84*d^3*e^3*x^3 + 126*d^2*e^4*x^4 + 126*d*e^5*x^5 + 84*e^6
*x^6)) + 210*a^4*b^6*e^4*(2*A*e*(d^6 + 9*d^5*e*x + 36*d^4*e^2*x^2 + 84*d^3
*e^3*x^3 + 126*d^2*e^4*x^4 + 126*d*e^5*x^5 + 84*e^6*x^6) + 7*B*(d^7 + 9*d^
6*e*x + 36*d^5*e^2*x^2 + 84*d^4*e^3*x^3 + 126*d^3*e^4*x^4 + 126*d^2*e^5*x^
5 + 84*d*e^6*x^6 + 36*e^7*x^7)) + 840*a^3*b^7*e^3*(A*e*(d^7 + 9*d^6*e*x +
36*d^5*e^2*x^2 + 84*d^4*e^3*x^3 + 126*d^3*e^4*x^4 + 126*d^2*e^5*x^5 + 84*d
*e^6*x^6 + 36*e^7*x^7) + 8*B*(d^8 + 9*d^7*e*x + 36*d^6*e^2*x^2 + 84*d^5*e^
3*x^3 + 126*d^4*e^4*x^4 + 126*d^3*e^5*x^5 + 84*d^2*e^6*x^6 + 36*d*e^7*x^7
+ 9*e^8*x^8)) - 9*a^2*b^8*e^2*(-280*A*e*(d^8 + 9*d^7*e*x + 36*d^6*e^2*x^2
+ 84*d^5*e^3*x^3 + 126*d^4*e^4*x^4 + 126*d^3*e^5*x^5 + 84*d^2*e^6*x^6 + 36
*d*e^7*x^7 + 9*e^8*x^8) + B*d*(7129*d^8 + 61641*d^7*e*x + 235224*d^6*e^...
```

Rubi [A] (verified)

Time = 1.21 (sec) , antiderivative size = 441, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{10}} dx$$

↓ 86

$$\int \left(\frac{b^9(10aBe + Abe - 10bBd)}{e^{11}} - \frac{5b^8(bd - ae)(9aBe + 2Abe - 11bBd)}{e^{11}(d + ex)} + \frac{15b^7(bd - ae)^2(8aBe + 3Abe - 11bBd)}{e^{11}(d + ex)^2} \right.$$

↓ 2009

$$\begin{aligned} & - \frac{b^9x(-10aBe - Abe + 10bBd)}{e^{11}} + \frac{5b^8(bd - ae) \log(d + ex)(-9aBe - 2Abe + 11bBd)}{e^{11}} + \\ & \frac{15b^7(bd - ae)^2(-8aBe - 3Abe + 11bBd)}{e^{12}(d + ex)} - \frac{15b^6(bd - ae)^3(-7aBe - 4Abe + 11bBd)}{e^{12}(d + ex)^2} + \\ & \frac{14b^5(bd - ae)^4(-6aBe - 5Abe + 11bBd)}{e^{12}(d + ex)^3} - \frac{21b^4(bd - ae)^5(-5aBe - 6Abe + 11bBd)}{2e^{12}(d + ex)^4} + \\ & \frac{6b^3(bd - ae)^6(-4aBe - 7Abe + 11bBd)}{e^{12}(d + ex)^5} - \frac{5b^2(bd - ae)^7(-3aBe - 8Abe + 11bBd)}{2e^{12}(d + ex)^6} + \\ & \frac{5b(bd - ae)^8(-2aBe - 9Abe + 11bBd)}{7e^{12}(d + ex)^7} - \frac{(bd - ae)^9(-aBe - 10Abe + 11bBd)}{8e^{12}(d + ex)^8} + \\ & \frac{(bd - ae)^{10}(Bd - Ae)}{9e^{12}(d + ex)^9} + \frac{b^{10}Bx^2}{2e^{10}} \end{aligned}$$

input

```
Int[((a + b*x)^10*(A + B*x))/(d + e*x)^10,x]
```

output

$$\begin{aligned}
& -((b^9(10*b*B*d - A*b*e - 10*a*B*e)*x)/e^{11} + (b^{10}*B*x^2)/(2*e^{10}) + ((\\
& b*d - a*e)^{10}*(B*d - A*e))/(9*e^{12}*(d + e*x)^9) - ((b*d - a*e)^9*(11*b*B*d \\
& - 10*A*b*e - a*B*e))/(8*e^{12}*(d + e*x)^8) + (5*b*(b*d - a*e)^8*(11*b*B*d \\
& - 9*A*b*e - 2*a*B*e))/(7*e^{12}*(d + e*x)^7) - (5*b^2*(b*d - a*e)^7*(11*b*B* \\
& d - 8*A*b*e - 3*a*B*e))/(2*e^{12}*(d + e*x)^6) + (6*b^3*(b*d - a*e)^6*(11*b* \\
& B*d - 7*A*b*e - 4*a*B*e))/(e^{12}*(d + e*x)^5) - (21*b^4*(b*d - a*e)^5*(11*b \\
& *B*d - 6*A*b*e - 5*a*B*e))/(2*e^{12}*(d + e*x)^4) + (14*b^5*(b*d - a*e)^4*(1 \\
& 1*b*B*d - 5*A*b*e - 6*a*B*e))/(e^{12}*(d + e*x)^3) - (15*b^6*(b*d - a*e)^3*(\\
& 11*b*B*d - 4*A*b*e - 7*a*B*e))/(e^{12}*(d + e*x)^2) + (15*b^7*(b*d - a*e)^2* \\
& (11*b*B*d - 3*A*b*e - 8*a*B*e))/(e^{12}*(d + e*x)) + (5*b^8*(b*d - a*e)*(11* \\
& b*B*d - 2*A*b*e - 9*a*B*e)*\text{Log}[d + e*x])/e^{12}
\end{aligned}$$

Defintions of rubi rules used

rule 86

```

Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))

```

rule 2009

```

Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]

```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1927 vs. $2(429) = 858$.

Time = 0.24 (sec) , antiderivative size = 1928, normalized size of antiderivative = 4.37

method	result	size
default	Expression too large to display	1928
norman	Expression too large to display	1933
risch	Expression too large to display	1942
parallelsch	Expression too large to display	3215

input

```

int((b*x+a)^10*(B*x+A)/(e*x+d)^10,x,method=_RETURNVERBOSE)

```


output

```

b^9/e^11*(1/2*B*b*e*x^2+A*b*e*x+10*B*a*e*x-10*B*b*d*x)-5/2*b^2/e^12*(8*A*a
^7*b*e^8-56*A*a^6*b^2*d*e^7+168*A*a^5*b^3*d^2*e^6-280*A*a^4*b^4*d^3*e^5+28
0*A*a^3*b^5*d^4*e^4-168*A*a^2*b^6*d^5*e^3+56*A*a*b^7*d^6*e^2-8*A*b^8*d^7*e
+3*B*a^8*e^8-32*B*a^7*b*d*e^7+140*B*a^6*b^2*d^2*e^6-336*B*a^5*b^3*d^3*e^5+
490*B*a^4*b^4*d^4*e^4-448*B*a^3*b^5*d^5*e^3+252*B*a^2*b^6*d^6*e^2-80*B*a*b
^7*d^7*e+11*B*b^8*d^8)/(e*x+d)^6-5/7*b/e^12*(9*A*a^8*b*e^9-72*A*a^7*b^2*d*
e^8+252*A*a^6*b^3*d^2*e^7-504*A*a^5*b^4*d^3*e^6+630*A*a^4*b^5*d^4*e^5-504*
A*a^3*b^6*d^5*e^4+252*A*a^2*b^7*d^6*e^3-72*A*a*b^8*d^7*e^2+9*A*b^9*d^8*e+2
*B*a^9*e^9-27*B*a^8*b*d*e^8+144*B*a^7*b^2*d^2*e^7-420*B*a^6*b^3*d^3*e^6+75
6*B*a^5*b^4*d^4*e^5-882*B*a^4*b^5*d^5*e^4+672*B*a^3*b^6*d^6*e^3-324*B*a^2*
b^7*d^7*e^2+90*B*a*b^8*d^8*e-11*B*b^9*d^9)/(e*x+d)^7-1/8/e^12*(10*A*a^9*b*
e^10-90*A*a^8*b^2*d*e^9+360*A*a^7*b^3*d^2*e^8-840*A*a^6*b^4*d^3*e^7+1260*A
*a^5*b^5*d^4*e^6-1260*A*a^4*b^6*d^5*e^5+840*A*a^3*b^7*d^6*e^4-360*A*a^2*b^
8*d^7*e^3+90*A*a*b^9*d^8*e^2-10*A*b^10*d^9*e+B*a^10*e^10-20*B*a^9*b*d*e^9+
135*B*a^8*b^2*d^2*e^8-480*B*a^7*b^3*d^3*e^7+1050*B*a^6*b^4*d^4*e^6-1512*B*
a^5*b^5*d^5*e^5+1470*B*a^4*b^6*d^6*e^4-960*B*a^3*b^7*d^7*e^3+405*B*a^2*b^8
*d^8*e^2-100*B*a*b^9*d^9*e+11*B*b^10*d^10)/(e*x+d)^8-15*b^7/e^12*(3*A*a^2*
b*e^3-6*A*a*b^2*d*e^2+3*A*b^3*d^2*e+8*B*a^3*e^3-27*B*a^2*b*d*e^2+30*B*a*b^
2*d^2*e-11*B*b^3*d^3)/(e*x+d)-15*b^6/e^12*(4*A*a^3*b*e^4-12*A*a^2*b^2*d*e^
3+12*A*a*b^3*d^2*e^2-4*A*b^4*d^3*e+7*B*a^4*e^4-32*B*a^3*b*d*e^3+54*B*a^...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2501 vs. $2(429) = 858$.

Time = 0.23 (sec) , antiderivative size = 2501, normalized size of antiderivative = 5.67

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{10}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)/(e*x+d)^10,x, algorithm="fricas")
```

output

```

1/504*(252*B*b^10*e^11*x^11 + 42131*B*b^10*d^11 - 56*A*a^10*e^11 - 9722*(1
0*B*a*b^9 + A*b^10)*d^10*e + 7129*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 - 840*
(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 - 210*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*
e^4 - 84*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 - 42*(5*B*a^6*b^4 + 6*A*a^5*b
^5)*d^5*e^6 - 24*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 - 15*(3*B*a^8*b^2 + 8
*A*a^7*b^3)*d^3*e^8 - 10*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 - 7*(B*a^10 + 1
0*A*a^9*b)*d*e^10 - 252*(11*B*b^10*d*e^10 - 2*(10*B*a*b^9 + A*b^10)*e^11)*
x^10 - 4536*(8*B*b^10*d^2*e^9 - (10*B*a*b^9 + A*b^10)*d*e^10)*x^9 - 1512*(
51*B*b^10*d^3*e^8 + 3*(10*B*a*b^9 + A*b^10)*d^2*e^9 - 15*(9*B*a^2*b^8 + 2*
A*a*b^9)*d*e^10 + 5*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 1512*(126*B*b
^10*d^4*e^7 - 72*(10*B*a*b^9 + A*b^10)*d^3*e^8 + 90*(9*B*a^2*b^8 + 2*A*a*b
^9)*d^2*e^9 - 20*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 - 5*(7*B*a^4*b^6 + 4*A*
a^3*b^7)*e^11)*x^7 + 3528*(346*B*b^10*d^5*e^6 - 112*(10*B*a*b^9 + A*b^10)*
d^4*e^7 + 110*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 - 20*(8*B*a^3*b^7 + 3*A*a
^2*b^8)*d^2*e^9 - 5*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 - 2*(6*B*a^5*b^5 + 5
*A*a^4*b^6)*e^11)*x^6 + 5292*(511*B*b^10*d^6*e^5 - 142*(10*B*a*b^9 + A*b^1
0)*d^5*e^6 + 125*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 - 20*(8*B*a^3*b^7 + 3*A
a^2*b^8)*d^3*e^8 - 5*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 - 2*(6*B*a^5*b^5
+ 5*A*a^4*b^6)*d*e^10 - (5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 + 756*(4501
*B*b^10*d^7*e^4 - 1162*(10*B*a*b^9 + A*b^10)*d^6*e^5 + 959*(9*B*a^2*b^8...

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{10}} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**10*(B*x+A)/(e*x+d)**10,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1904 vs. $2(429) = 858$.

Time = 0.17 (sec) , antiderivative size = 1904, normalized size of antiderivative = 4.32

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{10}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^10,x, algorithm="maxima")`

output

```
1/504*(42131*B*b^10*d^11 - 56*A*a^10*e^11 - 9722*(10*B*a*b^9 + A*b^10)*d^1
0*e + 7129*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 - 840*(8*B*a^3*b^7 + 3*A*a^2*
b^8)*d^8*e^3 - 210*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e^4 - 84*(6*B*a^5*b^5 +
5*A*a^4*b^6)*d^6*e^5 - 42*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^5*e^6 - 24*(4*B*a
^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 - 15*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^3*e^8 - 1
0*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 - 7*(B*a^10 + 10*A*a^9*b)*d*e^10 + 756
0*(11*B*b^10*d^3*e^8 - 3*(10*B*a*b^9 + A*b^10)*d^2*e^9 + 3*(9*B*a^2*b^8 +
2*A*a*b^9)*d*e^10 - (8*B*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 7560*(77*B*b^1
0*d^4*e^7 - 20*(10*B*a*b^9 + A*b^10)*d^3*e^8 + 18*(9*B*a^2*b^8 + 2*A*a*b^9
)*d^2*e^9 - 4*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 - (7*B*a^4*b^6 + 4*A*a^3*
b^7)*e^11)*x^7 + 3528*(517*B*b^10*d^5*e^6 - 130*(10*B*a*b^9 + A*b^10)*d^4*
e^7 + 110*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 - 20*(8*B*a^3*b^7 + 3*A*a^2*b^
8)*d^2*e^9 - 5*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 - 2*(6*B*a^5*b^5 + 5*A*a
^4*b^6)*e^11)*x^6 + 5292*(627*B*b^10*d^6*e^5 - 154*(10*B*a*b^9 + A*b^10)*d
^5*e^6 + 125*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 - 20*(8*B*a^3*b^7 + 3*A*a^2
*b^8)*d^3*e^8 - 5*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 - 2*(6*B*a^5*b^5 + 5
*A*a^4*b^6)*d*e^10 - (5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 + 756*(5049*B*b
^10*d^7*e^4 - 1218*(10*B*a*b^9 + A*b^10)*d^6*e^5 + 959*(9*B*a^2*b^8 + 2*A*
a*b^9)*d^5*e^6 - 140*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^4*e^7 - 35*(7*B*a^4*b^6
+ 4*A*a^3*b^7)*d^3*e^8 - 14*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^2*e^9 - 7*(5...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1975 vs. $2(429) = 858$.

Time = 0.13 (sec) , antiderivative size = 1975, normalized size of antiderivative = 4.48

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{10}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^10,x, algorithm="giac")`

output

```
5*(11*B*b^10*d^2 - 20*B*a*b^9*d*e - 2*A*b^10*d*e + 9*B*a^2*b^8*e^2 + 2*A*a
*b^9*e^2)*log(abs(e*x + d))/e^12 + 1/2*(B*b^10*e^10*x^2 - 20*B*b^10*d*e^9*x
+ 20*B*a*b^9*e^10*x + 2*A*b^10*e^10*x)/e^20 + 1/504*(42131*B*b^10*d^11 -
97220*B*a*b^9*d^10*e - 9722*A*b^10*d^10*e + 64161*B*a^2*b^8*d^9*e^2 + 142
58*A*a*b^9*d^9*e^2 - 6720*B*a^3*b^7*d^8*e^3 - 2520*A*a^2*b^8*d^8*e^3 - 147
0*B*a^4*b^6*d^7*e^4 - 840*A*a^3*b^7*d^7*e^4 - 504*B*a^5*b^5*d^6*e^5 - 420*
A*a^4*b^6*d^6*e^5 - 210*B*a^6*b^4*d^5*e^6 - 252*A*a^5*b^5*d^5*e^6 - 96*B*a
^7*b^3*d^4*e^7 - 168*A*a^6*b^4*d^4*e^7 - 45*B*a^8*b^2*d^3*e^8 - 120*A*a^7*
b^3*d^3*e^8 - 20*B*a^9*b*d^2*e^9 - 90*A*a^8*b^2*d^2*e^9 - 7*B*a^10*d*e^10
- 70*A*a^9*b*d*e^10 - 56*A*a^10*e^11 + 7560*(11*B*b^10*d^3*e^8 - 30*B*a*b^
9*d^2*e^9 - 3*A*b^10*d^2*e^9 + 27*B*a^2*b^8*d*e^10 + 6*A*a*b^9*d*e^10 - 8*
B*a^3*b^7*e^11 - 3*A*a^2*b^8*e^11)*x^8 + 7560*(77*B*b^10*d^4*e^7 - 200*B*a
*b^9*d^3*e^8 - 20*A*b^10*d^3*e^8 + 162*B*a^2*b^8*d^2*e^9 + 36*A*a*b^9*d^2*
e^9 - 32*B*a^3*b^7*d*e^10 - 12*A*a^2*b^8*d*e^10 - 7*B*a^4*b^6*e^11 - 4*A*a
^3*b^7*e^11)*x^7 + 3528*(517*B*b^10*d^5*e^6 - 1300*B*a*b^9*d^4*e^7 - 130*A
*b^10*d^4*e^7 + 990*B*a^2*b^8*d^3*e^8 + 220*A*a*b^9*d^3*e^8 - 160*B*a^3*b^
7*d^2*e^9 - 60*A*a^2*b^8*d^2*e^9 - 35*B*a^4*b^6*d*e^10 - 20*A*a^3*b^7*d*e^
10 - 12*B*a^5*b^5*e^11 - 10*A*a^4*b^6*e^11)*x^6 + 5292*(627*B*b^10*d^6*e^5
- 1540*B*a*b^9*d^5*e^6 - 154*A*b^10*d^5*e^6 + 1125*B*a^2*b^8*d^4*e^7 + 25
0*A*a*b^9*d^4*e^7 - 160*B*a^3*b^7*d^3*e^8 - 60*A*a^2*b^8*d^3*e^8 - 35*B...
```

Mupad [B] (verification not implemented)

Time = 1.25 (sec) , antiderivative size = 2048, normalized size of antiderivative = 4.64

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{10}} dx = \text{Too large to display}$$

input `int(((A + B*x)*(a + b*x)^10)/(d + e*x)^10,x)`

output

```
x*((A*b^10 + 10*B*a*b^9)/e^10 - (10*B*b^10*d)/e^11) - (x^7*(60*A*a^3*b^7*e^10 + 105*B*a^4*b^6*e^10 + 300*A*b^10*d^3*e^7 - 1155*B*b^10*d^4*e^6 - 540*A*a*b^9*d^2*e^8 + 180*A*a^2*b^8*d*e^9 + 3000*B*a*b^9*d^3*e^7 + 480*B*a^3*b^7*d*e^9 - 2430*B*a^2*b^8*d^2*e^8) + x^4*(42*A*a^6*b^4*e^10 + 24*B*a^7*b^3*e^10 + 1827*A*b^10*d^6*e^4 - (15147*B*b^10*d^7*e^3)/2 - 2877*A*a*b^9*d^5*e^5 + 63*A*a^5*b^5*d*e^9 + 18270*B*a*b^9*d^6*e^4 + (105*B*a^6*b^4*d*e^9)/2 + 630*A*a^2*b^8*d^4*e^6 + 210*A*a^3*b^7*d^3*e^7 + 105*A*a^4*b^6*d^2*e^8 - (25893*B*a^2*b^8*d^5*e^5)/2 + 1680*B*a^3*b^7*d^4*e^6 + (735*B*a^4*b^6*d^3*e^7)/2 + 126*B*a^5*b^5*d^2*e^8) + x^6*(70*A*a^4*b^6*e^10 + 84*B*a^5*b^5*e^10 + 910*A*b^10*d^4*e^6 - 3619*B*b^10*d^5*e^5 - 1540*A*a*b^9*d^3*e^7 + 140*A*a^3*b^7*d*e^9 + 9100*B*a*b^9*d^4*e^6 + 245*B*a^4*b^6*d*e^9 + 420*A*a^2*b^8*d^2*e^8 - 6930*B*a^2*b^8*d^3*e^7 + 1120*B*a^3*b^7*d^2*e^8) + x^3*(20*A*a^7*b^3*e^10 + (15*B*a^8*b^2*e^10)/2 + 1338*A*b^10*d^7*e^3 - (11253*B*b^10*d^8*e^2)/2 - 2058*A*a*b^9*d^6*e^4 + 28*A*a^6*b^4*d*e^9 + 13380*B*a*b^9*d^7*e^3 + 16*B*a^7*b^3*d*e^9 + 420*A*a^2*b^8*d^5*e^5 + 140*A*a^3*b^7*d^4*e^6 + 70*A*a^4*b^6*d^3*e^7 + 42*A*a^5*b^5*d^2*e^8 - 9261*B*a^2*b^8*d^6*e^4 + 1120*B*a^3*b^7*d^5*e^5 + 245*B*a^4*b^6*d^4*e^6 + 84*B*a^5*b^5*d^3*e^7 + 35*B*a^6*b^4*d^2*e^8) + (56*A*a^10*e^11 - 42131*B*b^10*d^11 + 9722*A*b^10*d^10*e + 7*B*a^10*d*e^10 - 14258*A*a*b^9*d^9*e^2 + 20*B*a^9*b*d^2*e^9 + 2520*A*a^2*b^8*d^8*e^3 + 840*A*a^3*b^7*d^7*e^4 + 420*A*a^4*b^6*d^6*e^5 + ...
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 1714, normalized size of antiderivative = 3.89

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{10}} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^10,x)`

output

```
(27720*log(d + e*x)*a**2*b**9*d**10*e**2 + 249480*log(d + e*x)*a**2*b**9*d
**9*e**3*x + 997920*log(d + e*x)*a**2*b**9*d**8*e**4*x**2 + 2328480*log(d
+ e*x)*a**2*b**9*d**7*e**5*x**3 + 3492720*log(d + e*x)*a**2*b**9*d**6*e**6
*x**4 + 3492720*log(d + e*x)*a**2*b**9*d**5*e**7*x**5 + 2328480*log(d + e
x)*a**2*b**9*d**4*e**8*x**6 + 997920*log(d + e*x)*a**2*b**9*d**3*e**9*x**7
+ 249480*log(d + e*x)*a**2*b**9*d**2*e**10*x**8 + 27720*log(d + e*x)*a**2
*b**9*d*e**11*x**9 - 55440*log(d + e*x)*a*b**10*d**11*e - 498960*log(d + e
*x)*a*b**10*d**10*e**2*x - 1995840*log(d + e*x)*a*b**10*d**9*e**3*x**2 - 4
656960*log(d + e*x)*a*b**10*d**8*e**4*x**3 - 6985440*log(d + e*x)*a*b**10*
d**7*e**5*x**4 - 6985440*log(d + e*x)*a*b**10*d**6*e**6*x**5 - 4656960*log
(d + e*x)*a*b**10*d**5*e**7*x**6 - 1995840*log(d + e*x)*a*b**10*d**4*e**8*
x**7 - 498960*log(d + e*x)*a*b**10*d**3*e**9*x**8 - 55440*log(d + e*x)*a*b
**10*d**2*e**10*x**9 + 27720*log(d + e*x)*b**11*d**12 + 249480*log(d + e*x
)*b**11*d**11*e*x + 997920*log(d + e*x)*b**11*d**10*e**2*x**2 + 2328480*lo
g(d + e*x)*b**11*d**9*e**3*x**3 + 3492720*log(d + e*x)*b**11*d**8*e**4*x**
4 + 3492720*log(d + e*x)*b**11*d**7*e**5*x**5 + 2328480*log(d + e*x)*b**11
*d**6*e**6*x**6 + 997920*log(d + e*x)*b**11*d**5*e**7*x**7 + 249480*log(d
+ e*x)*b**11*d**4*e**8*x**8 + 27720*log(d + e*x)*b**11*d**3*e**9*x**9 - 56
*a**11*d*e**11 - 77*a**10*b*d**2*e**10 - 693*a**10*b*d*e**11*x - 110*a**9*
b**2*d**3*e**9 - 990*a**9*b**2*d**2*e**10*x - 3960*a**9*b**2*d*e**11*x...
```

3.89
$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{11}} dx$$

Optimal result	915
Mathematica [B] (verified)	916
Rubi [A] (verified)	917
Maple [B] (verified)	918
Fricas [B] (verification not implemented)	919
Sympy [F(-1)]	920
Maxima [B] (verification not implemented)	921
Giac [B] (verification not implemented)	922
Mupad [B] (verification not implemented)	923
Reduce [B] (verification not implemented)	923

Optimal result

Integrand size = 20, antiderivative size = 446

$$\begin{aligned}
 \int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{11}} dx &= \frac{b^{10}Bx}{e^{11}} + \frac{(bd-ae)^{10}(Bd-Ae)}{10e^{12}(d+ex)^{10}} \\
 &\quad - \frac{(bd-ae)^9(11bBd-10Abe-aBe)}{9e^{12}(d+ex)^9} \\
 &\quad + \frac{5b(bd-ae)^8(11bBd-9Abe-2aBe)}{8e^{12}(d+ex)^8} \\
 &\quad - \frac{15b^2(bd-ae)^7(11bBd-8Abe-3aBe)}{7e^{12}(d+ex)^7} \\
 &\quad + \frac{5b^3(bd-ae)^6(11bBd-7Abe-4aBe)}{e^{12}(d+ex)^6} \\
 &\quad - \frac{42b^4(bd-ae)^5(11bBd-6Abe-5aBe)}{5e^{12}(d+ex)^5} \\
 &\quad + \frac{21b^5(bd-ae)^4(11bBd-5Abe-6aBe)}{2e^{12}(d+ex)^4} \\
 &\quad - \frac{10b^6(bd-ae)^3(11bBd-4Abe-7aBe)}{e^{12}(d+ex)^3} \\
 &\quad + \frac{15b^7(bd-ae)^2(11bBd-3Abe-8aBe)}{2e^{12}(d+ex)^2} \\
 &\quad - \frac{5b^8(bd-ae)(11bBd-2Abe-9aBe)}{e^{12}(d+ex)} \\
 &\quad - \frac{b^9(11bBd-Abe-10aBe)\log(d+ex)}{e^{12}}
 \end{aligned}$$

output

```

b^10*B*x/e^11+1/10*(-a*e+b*d)^10*(-A*e+B*d)/e^12/(e*x+d)^10-1/9*(-a*e+b*d)
^9*(-10*A*b*e-B*a*e+11*B*b*d)/e^12/(e*x+d)^9+5/8*b*(-a*e+b*d)^8*(-9*A*b*e-
2*B*a*e+11*B*b*d)/e^12/(e*x+d)^8-15/7*b^2*(-a*e+b*d)^7*(-8*A*b*e-3*B*a*e+1
1*B*b*d)/e^12/(e*x+d)^7+5*b^3*(-a*e+b*d)^6*(-7*A*b*e-4*B*a*e+11*B*b*d)/e^1
2/(e*x+d)^6-42/5*b^4*(-a*e+b*d)^5*(-6*A*b*e-5*B*a*e+11*B*b*d)/e^12/(e*x+d)
^5+21/2*b^5*(-a*e+b*d)^4*(-5*A*b*e-6*B*a*e+11*B*b*d)/e^12/(e*x+d)^4-10*b^6
*(-a*e+b*d)^3*(-4*A*b*e-7*B*a*e+11*B*b*d)/e^12/(e*x+d)^3+15/2*b^7*(-a*e+b*
d)^2*(-3*A*b*e-8*B*a*e+11*B*b*d)/e^12/(e*x+d)^2-5*b^8*(-a*e+b*d)*(-2*A*b*e
-9*B*a*e+11*B*b*d)/e^12/(e*x+d)-b^9*(-A*b*e-10*B*a*e+11*B*b*d)*ln(e*x+d)/e
^12

```


Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1447 vs. $2(446) = 892$.

Time = 0.54 (sec) , antiderivative size = 1447, normalized size of antiderivative = 3.24

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{11}} dx = \text{Too large to display}$$

input `Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^11,x]`

output

```
-1/2520*(28*a^10*e^10*(9*A*e + B*(d + 10*e*x)) + 70*a^9*b*e^9*(4*A*e*(d +
10*e*x) + B*(d^2 + 10*d*e*x + 45*e^2*x^2)) + 45*a^8*b^2*e^8*(7*A*e*(d^2 +
10*d*e*x + 45*e^2*x^2) + 3*B*(d^3 + 10*d^2*e*x + 45*d*e^2*x^2 + 120*e^3*x^
3)) + 120*a^7*b^3*e^7*(3*A*e*(d^3 + 10*d^2*e*x + 45*d*e^2*x^2 + 120*e^3*x^
3) + 2*B*(d^4 + 10*d^3*e*x + 45*d^2*e^2*x^2 + 120*d*e^3*x^3 + 210*e^4*x^4)
) + 420*a^6*b^4*e^6*(A*e*(d^4 + 10*d^3*e*x + 45*d^2*e^2*x^2 + 120*d*e^3*x^
3 + 210*e^4*x^4) + B*(d^5 + 10*d^4*e*x + 45*d^3*e^2*x^2 + 120*d^2*e^3*x^3
+ 210*d*e^4*x^4 + 252*e^5*x^5)) + 252*a^5*b^5*e^5*(2*A*e*(d^5 + 10*d^4*e*x
+ 45*d^3*e^2*x^2 + 120*d^2*e^3*x^3 + 210*d*e^4*x^4 + 252*e^5*x^5) + 3*B*(
d^6 + 10*d^5*e*x + 45*d^4*e^2*x^2 + 120*d^3*e^3*x^3 + 210*d^2*e^4*x^4 + 25
2*d*e^5*x^5 + 210*e^6*x^6)) + 210*a^4*b^6*e^4*(3*A*e*(d^6 + 10*d^5*e*x + 4
5*d^4*e^2*x^2 + 120*d^3*e^3*x^3 + 210*d^2*e^4*x^4 + 252*d*e^5*x^5 + 210*e^
6*x^6) + 7*B*(d^7 + 10*d^6*e*x + 45*d^5*e^2*x^2 + 120*d^4*e^3*x^3 + 210*d^
3*e^4*x^4 + 252*d^2*e^5*x^5 + 210*d*e^6*x^6 + 120*e^7*x^7)) + 840*a^3*b^7*
e^3*(A*e*(d^7 + 10*d^6*e*x + 45*d^5*e^2*x^2 + 120*d^4*e^3*x^3 + 210*d^3*e^
4*x^4 + 252*d^2*e^5*x^5 + 210*d*e^6*x^6 + 120*e^7*x^7) + 4*B*(d^8 + 10*d^7
*e*x + 45*d^6*e^2*x^2 + 120*d^5*e^3*x^3 + 210*d^4*e^4*x^4 + 252*d^3*e^5*x^
5 + 210*d^2*e^6*x^6 + 120*d*e^7*x^7 + 45*e^8*x^8)) + 1260*a^2*b^8*e^2*(A*e
*(d^8 + 10*d^7*e*x + 45*d^6*e^2*x^2 + 120*d^5*e^3*x^3 + 210*d^4*e^4*x^4 +
252*d^3*e^5*x^5 + 210*d^2*e^6*x^6 + 120*d*e^7*x^7 + 45*e^8*x^8) + 9*B*(...
```

Rubi [A] (verified)

Time = 1.21 (sec) , antiderivative size = 446, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{11}} dx$$

↓ 86

$$\int \left(\frac{b^9(10aBe + Abe - 11bBd)}{e^{11}(d + ex)} - \frac{5b^8(bd - ae)(9aBe + 2Abe - 11bBd)}{e^{11}(d + ex)^2} + \frac{15b^7(bd - ae)^2(8aBe + 3Abe - 11bBd)}{e^{11}(d + ex)^3} \right.$$

↓ 2009

$$\begin{aligned} & - \frac{b^9 \log(d + ex)(-10aBe - Abe + 11bBd)}{e^{12}} - \frac{5b^8(bd - ae)(-9aBe - 2Abe + 11bBd)}{e^{12}(d + ex)} + \\ & \frac{15b^7(bd - ae)^2(-8aBe - 3Abe + 11bBd)}{2e^{12}(d + ex)^2} - \frac{10b^6(bd - ae)^3(-7aBe - 4Abe + 11bBd)}{e^{12}(d + ex)^3} + \\ & \frac{21b^5(bd - ae)^4(-6aBe - 5Abe + 11bBd)}{2e^{12}(d + ex)^4} - \frac{42b^4(bd - ae)^5(-5aBe - 6Abe + 11bBd)}{5e^{12}(d + ex)^5} + \\ & \frac{5b^3(bd - ae)^6(-4aBe - 7Abe + 11bBd)}{e^{12}(d + ex)^6} - \frac{15b^2(bd - ae)^7(-3aBe - 8Abe + 11bBd)}{7e^{12}(d + ex)^7} + \\ & \frac{5b(bd - ae)^8(-2aBe - 9Abe + 11bBd)}{8e^{12}(d + ex)^8} - \frac{(bd - ae)^9(-aBe - 10Abe + 11bBd)}{9e^{12}(d + ex)^9} + \\ & \frac{(bd - ae)^{10}(Bd - Ae)}{10e^{12}(d + ex)^{10}} + \frac{b^{10}Bx}{e^{11}} \end{aligned}$$

input

```
Int[((a + b*x)^10*(A + B*x))/(d + e*x)^11,x]
```

output

$$\begin{aligned} & (b^{10}Bx)/e^{11} + ((b*d - a*e)^{10}(B*d - A*e))/(10*e^{12}(d + e*x)^{10}) - ((\\ & b*d - a*e)^9*(11*b*B*d - 10*A*b*e - a*B*e))/(9*e^{12}(d + e*x)^9) + (5*b*(b \\ & *d - a*e)^8*(11*b*B*d - 9*A*b*e - 2*a*B*e))/(8*e^{12}(d + e*x)^8) - (15*b^2 \\ & *(b*d - a*e)^7*(11*b*B*d - 8*A*b*e - 3*a*B*e))/(7*e^{12}(d + e*x)^7) + (5*b \\ & ^3*(b*d - a*e)^6*(11*b*B*d - 7*A*b*e - 4*a*B*e))/(e^{12}(d + e*x)^6) - (42* \\ & b^4*(b*d - a*e)^5*(11*b*B*d - 6*A*b*e - 5*a*B*e))/(5*e^{12}(d + e*x)^5) + (\\ & 21*b^5*(b*d - a*e)^4*(11*b*B*d - 5*A*b*e - 6*a*B*e))/(2*e^{12}(d + e*x)^4) \\ & - (10*b^6*(b*d - a*e)^3*(11*b*B*d - 4*A*b*e - 7*a*B*e))/(e^{12}(d + e*x)^3) \\ & + (15*b^7*(b*d - a*e)^2*(11*b*B*d - 3*A*b*e - 8*a*B*e))/(2*e^{12}(d + e*x) \\ & ^2) - (5*b^8*(b*d - a*e)*(11*b*B*d - 2*A*b*e - 9*a*B*e))/(e^{12}(d + e*x)) \\ & - (b^9*(11*b*B*d - A*b*e - 10*a*B*e)*\text{Log}[d + e*x])/e^{12} \end{aligned}$$

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1921 vs. $2(432) = 864$.

Time = 0.26 (sec) , antiderivative size = 1922, normalized size of antiderivative = 4.31

method	result	size
risch	Expression too large to display	1922
default	Expression too large to display	1933
norman	Expression too large to display	1934
parallelrisch	Expression too large to display	2863

input

```
int((b*x+a)^10*(B*x+A)/(e*x+d)^11,x,method=_RETURNVERBOSE)
```

output

```

b^10*B*x/e^11+((-10*A*a*b^9*e^10+10*A*b^10*d*e^9-45*B*a^2*b^8*e^10+100*B*a
*b^9*d*e^9-55*B*b^10*d^2*e^8)*x^9-15/2*b^7*e^7*(3*A*a^2*b*e^3+6*A*a*b^2*d*
e^2-9*A*b^3*d^2*e+8*B*a^3*e^3+27*B*a^2*b*d*e^2-90*B*a*b^2*d^2*e+55*B*b^3*d
^3)*x^8-10*b^6*e^6*(4*A*a^3*b*e^4+6*A*a^2*b^2*d*e^3+12*A*a*b^3*d^2*e^2-22*
A*b^4*d^3*e+7*B*a^4*e^4+16*B*a^3*b*d*e^3+54*B*a^2*b^2*d^2*e^2-220*B*a*b^3*
d^3*e+143*B*b^4*d^4)*x^7-7/2*b^5*e^5*(15*A*a^4*b*e^5+20*A*a^3*b^2*d*e^4+30
*A*a^2*b^3*d^2*e^3+60*A*a*b^4*d^3*e^2-125*A*b^5*d^4*e+18*B*a^5*e^5+35*B*a^
4*b*d*e^4+80*B*a^3*b^2*d^2*e^3+270*B*a^2*b^3*d^3*e^2-1250*B*a*b^4*d^4*e+84
7*B*b^5*d^5)*x^6-21/5*b^4*e^4*(12*A*a^5*b*e^6+15*A*a^4*b^2*d*e^5+20*A*a^3*
b^3*d^2*e^4+30*A*a^2*b^4*d^3*e^3+60*A*a*b^5*d^4*e^2-137*A*b^6*d^5*e+10*B*a
^6*e^6+18*B*a^5*b*d*e^5+35*B*a^4*b^2*d^2*e^4+80*B*a^3*b^3*d^3*e^3+270*B*a^
2*b^4*d^4*e^2-1370*B*a*b^5*d^5*e+957*B*b^6*d^6)*x^5-1/2*e^3*b^3*(70*A*a^6*
b*e^7+84*A*a^5*b^2*d*e^6+105*A*a^4*b^3*d^2*e^5+140*A*a^3*b^4*d^3*e^4+210*A
*a^2*b^5*d^4*e^3+420*A*a*b^6*d^5*e^2-1029*A*b^7*d^6*e+40*B*a^7*e^7+70*B*a^
6*b*d*e^6+126*B*a^5*b^2*d^2*e^5+245*B*a^4*b^3*d^3*e^4+560*B*a^3*b^4*d^4*e^
3+1890*B*a^2*b^5*d^5*e^2-10290*B*a*b^6*d^6*e+7359*B*b^7*d^7)*x^4-1/7*b^2*e
^2*(120*A*a^7*b*e^8+140*A*a^6*b^2*d*e^7+168*A*a^5*b^3*d^2*e^6+210*A*a^4*b^
4*d^3*e^5+280*A*a^3*b^5*d^4*e^4+420*A*a^2*b^6*d^5*e^3+840*A*a*b^7*d^6*e^2-
2178*A*b^8*d^7*e+45*B*a^8*e^8+80*B*a^7*b*d*e^7+140*B*a^6*b^2*d^2*e^6+252*B
*a^5*b^3*d^3*e^5+490*B*a^4*b^4*d^4*e^4+1120*B*a^3*b^5*d^5*e^3+3780*B*a^...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2309 vs. $2(432) = 864$.

Time = 0.24 (sec) , antiderivative size = 2309, normalized size of antiderivative = 5.18

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{11}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)/(e*x+d)^11,x, algorithm="fricas")
```

output

```

1/2520*(2520*B*b^10*e^11*x^11 + 25200*B*b^10*d*e^10*x^10 - 55991*B*b^10*d^
11 - 252*A*a^10*e^11 + 7381*(10*B*a*b^9 + A*b^10)*d^10*e - 1260*(9*B*a^2*b
^8 + 2*A*a*b^9)*d^9*e^2 - 420*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 - 210*(7
*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e^4 - 126*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^
5 - 84*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^5*e^6 - 60*(4*B*a^7*b^3 + 7*A*a^6*b^4
)*d^4*e^7 - 45*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^3*e^8 - 35*(2*B*a^9*b + 9*A*a
^8*b^2)*d^2*e^9 - 28*(B*a^10 + 10*A*a^9*b)*d*e^10 - 12600*(2*B*b^10*d^2*e^
9 - 2*(10*B*a*b^9 + A*b^10)*d*e^10 + (9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 -
18900*(39*B*b^10*d^3*e^8 - 9*(10*B*a*b^9 + A*b^10)*d^2*e^9 + 3*(9*B*a^2*b
^8 + 2*A*a*b^9)*d*e^10 + (8*B*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 - 25200*(12
2*B*b^10*d^4*e^7 - 22*(10*B*a*b^9 + A*b^10)*d^3*e^8 + 6*(9*B*a^2*b^8 + 2*A
*a*b^9)*d^2*e^9 + 2*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 + (7*B*a^4*b^6 + 4*
A*a^3*b^7)*e^11)*x^7 - 8820*(775*B*b^10*d^5*e^6 - 125*(10*B*a*b^9 + A*b^10
)*d^4*e^7 + 30*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 + 10*(8*B*a^3*b^7 + 3*A*a
^2*b^8)*d^2*e^9 + 5*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 + 3*(6*B*a^5*b^5 +
5*A*a^4*b^6)*e^11)*x^6 - 10584*(907*B*b^10*d^6*e^5 - 137*(10*B*a*b^9 + A*b
^10)*d^5*e^6 + 30*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 + 10*(8*B*a^3*b^7 + 3*
A*a^2*b^8)*d^3*e^8 + 5*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 + 3*(6*B*a^5*b^
5 + 5*A*a^4*b^6)*d*e^10 + 2*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 - 1260*(
7119*B*b^10*d^7*e^4 - 1029*(10*B*a*b^9 + A*b^10)*d^6*e^5 + 210*(9*B*a^2...

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{11}} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**10*(B*x+A)/(e*x+d)**11,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1914 vs. $2(432) = 864$.

Time = 0.13 (sec) , antiderivative size = 1914, normalized size of antiderivative = 4.29

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{11}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^11,x, algorithm="maxima")`

output

```
B*b^10*x/e^11 - 1/2520*(55991*B*b^10*d^11 + 252*A*a^10*e^11 - 7381*(10*B*a
*b^9 + A*b^10)*d^10*e + 1260*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 + 420*(8*B*
a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 210*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e^4 +
126*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 84*(5*B*a^6*b^4 + 6*A*a^5*b^5)*
d^5*e^6 + 60*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 45*(3*B*a^8*b^2 + 8*A*a
^7*b^3)*d^3*e^8 + 35*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 + 28*(B*a^10 + 10*A
*a^9*b)*d*e^10 + 12600*(11*B*b^10*d^2*e^9 - 2*(10*B*a*b^9 + A*b^10)*d*e^10
+ (9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 + 18900*(55*B*b^10*d^3*e^8 - 9*(10*
B*a*b^9 + A*b^10)*d^2*e^9 + 3*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 + (8*B*a^3*
b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 25200*(143*B*b^10*d^4*e^7 - 22*(10*B*a*b^9
+ A*b^10)*d^3*e^8 + 6*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 + 2*(8*B*a^3*b^7 +
3*A*a^2*b^8)*d*e^10 + (7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7 + 8820*(847*B
*b^10*d^5*e^6 - 125*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 30*(9*B*a^2*b^8 + 2*A*
a*b^9)*d^3*e^8 + 10*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 5*(7*B*a^4*b^6 +
4*A*a^3*b^7)*d*e^10 + 3*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 + 10584*(95
7*B*b^10*d^6*e^5 - 137*(10*B*a*b^9 + A*b^10)*d^5*e^6 + 30*(9*B*a^2*b^8 + 2
*A*a*b^9)*d^4*e^7 + 10*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 + 5*(7*B*a^4*b^
6 + 4*A*a^3*b^7)*d^2*e^9 + 3*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10 + 2*(5*B*a
^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 + 1260*(7359*B*b^10*d^7*e^4 - 1029*(10*B*a
*b^9 + A*b^10)*d^6*e^5 + 210*(9*B*a^2*b^8 + 2*A*a*b^9)*d^5*e^6 + 70*(8*...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1970 vs. $2(432) = 864$.

Time = 0.13 (sec) , antiderivative size = 1970, normalized size of antiderivative = 4.42

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{11}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^11,x, algorithm="giac")`

output

```
B*b^10*x/e^11 - (11*B*b^10*d - 10*B*a*b^9*e - A*b^10*e)*log(abs(e*x + d))/
e^12 - 1/2520*(55991*B*b^10*d^11 - 73810*B*a*b^9*d^10*e - 7381*A*b^10*d^10
*e + 11340*B*a^2*b^8*d^9*e^2 + 2520*A*a*b^9*d^9*e^2 + 3360*B*a^3*b^7*d^8*e
^3 + 1260*A*a^2*b^8*d^8*e^3 + 1470*B*a^4*b^6*d^7*e^4 + 840*A*a^3*b^7*d^7*e
^4 + 756*B*a^5*b^5*d^6*e^5 + 630*A*a^4*b^6*d^6*e^5 + 420*B*a^6*b^4*d^5*e^6
+ 504*A*a^5*b^5*d^5*e^6 + 240*B*a^7*b^3*d^4*e^7 + 420*A*a^6*b^4*d^4*e^7 +
135*B*a^8*b^2*d^3*e^8 + 360*A*a^7*b^3*d^3*e^8 + 70*B*a^9*b*d^2*e^9 + 315*
A*a^8*b^2*d^2*e^9 + 28*B*a^10*d*e^10 + 280*A*a^9*b*d*e^10 + 252*A*a^10*e^1
1 + 12600*(11*B*b^10*d^2*e^9 - 20*B*a*b^9*d*e^10 - 2*A*b^10*d*e^10 + 9*B*a
^2*b^8*e^11 + 2*A*a*b^9*e^11)*x^9 + 18900*(55*B*b^10*d^3*e^8 - 90*B*a*b^9*
d^2*e^9 - 9*A*b^10*d^2*e^9 + 27*B*a^2*b^8*d*e^10 + 6*A*a*b^9*d*e^10 + 8*B*
a^3*b^7*e^11 + 3*A*a^2*b^8*e^11)*x^8 + 25200*(143*B*b^10*d^4*e^7 - 220*B*a
*b^9*d^3*e^8 - 22*A*b^10*d^3*e^8 + 54*B*a^2*b^8*d^2*e^9 + 12*A*a*b^9*d^2*e
^9 + 16*B*a^3*b^7*d*e^10 + 6*A*a^2*b^8*d*e^10 + 7*B*a^4*b^6*e^11 + 4*A*a^3
*b^7*e^11)*x^7 + 8820*(847*B*b^10*d^5*e^6 - 1250*B*a*b^9*d^4*e^7 - 125*A*b
^10*d^4*e^7 + 270*B*a^2*b^8*d^3*e^8 + 60*A*a*b^9*d^3*e^8 + 80*B*a^3*b^7*d^
2*e^9 + 30*A*a^2*b^8*d^2*e^9 + 35*B*a^4*b^6*d*e^10 + 20*A*a^3*b^7*d*e^10 +
18*B*a^5*b^5*e^11 + 15*A*a^4*b^6*e^11)*x^6 + 10584*(957*B*b^10*d^6*e^5 -
1370*B*a*b^9*d^5*e^6 - 137*A*b^10*d^5*e^6 + 270*B*a^2*b^8*d^4*e^7 + 60*A*a
*b^9*d^4*e^7 + 80*B*a^3*b^7*d^3*e^8 + 30*A*a^2*b^8*d^3*e^8 + 35*B*a^4*b...
```

Mupad [B] (verification not implemented)

Time = 1.36 (sec) , antiderivative size = 2874, normalized size of antiderivative = 6.44

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{11}} dx = \text{Too large to display}$$

input `int(((A + B*x)*(a + b*x)^10)/(d + e*x)^11,x)`

output

```

-((A*a^10*e^11)/10 + (55991*B*b^10*d^11)/2520 - (7381*A*b^10*d^10*e)/2520
+ (B*a^10*d*e^10)/90 + 11*B*b^10*d^11*log(d + e*x) + (B*a^10*e^11*x)/9 - B
*b^10*e^11*x^11 + (53219*B*b^10*d^10*e*x)/252 + A*a*b^9*d^9*e^2 + (B*a^9*b
*d^2*e^9)/36 + 10*A*a*b^9*e^11*x^9 + (5*B*a^9*b*e^11*x^2)/4 - (7129*A*b^10
*d^9*e^2*x)/252 - 10*A*b^10*d*e^10*x^9 - 10*B*b^10*d*e^10*x^10 - A*b^10*e^
11*x^10*log(d + e*x) + (A*a^2*b^8*d^8*e^3)/2 + (A*a^3*b^7*d^7*e^4)/3 + (A*
a^4*b^6*d^6*e^5)/4 + (A*a^5*b^5*d^5*e^6)/5 + (A*a^6*b^4*d^4*e^7)/6 + (A*a^
7*b^3*d^3*e^8)/7 + (A*a^8*b^2*d^2*e^9)/8 + (9*B*a^2*b^8*d^9*e^2)/2 + (4*B*
a^3*b^7*d^8*e^3)/3 + (7*B*a^4*b^6*d^7*e^4)/12 + (3*B*a^5*b^5*d^6*e^5)/10 +
(B*a^6*b^4*d^5*e^6)/6 + (2*B*a^7*b^3*d^4*e^7)/21 + (3*B*a^8*b^2*d^3*e^8)/
56 + (45*A*a^8*b^2*e^11*x^2)/8 + (120*A*a^7*b^3*e^11*x^3)/7 + 35*A*a^6*b^4
*e^11*x^4 + (252*A*a^5*b^5*e^11*x^5)/5 + (105*A*a^4*b^6*e^11*x^6)/2 + 40*A
*a^3*b^7*e^11*x^7 + (45*A*a^2*b^8*e^11*x^8)/2 + (45*B*a^8*b^2*e^11*x^3)/7
+ 20*B*a^7*b^3*e^11*x^4 + 42*B*a^6*b^4*e^11*x^5 + 63*B*a^5*b^5*e^11*x^6 +
70*B*a^4*b^6*e^11*x^7 + 60*B*a^3*b^7*e^11*x^8 + 45*B*a^2*b^8*e^11*x^9 - (6
849*A*b^10*d^8*e^3*x^2)/56 - (2178*A*b^10*d^7*e^4*x^3)/7 - (1029*A*b^10*d^
6*e^5*x^4)/2 - (2877*A*b^10*d^5*e^6*x^5)/5 - (875*A*b^10*d^4*e^7*x^6)/2 -
220*A*b^10*d^3*e^8*x^7 - (135*A*b^10*d^2*e^9*x^8)/2 + (50139*B*b^10*d^9*e^
2*x^2)/56 + (15558*B*b^10*d^8*e^3*x^3)/7 + (7119*B*b^10*d^7*e^4*x^4)/2 + (
19047*B*b^10*d^6*e^5*x^5)/5 + (5425*B*b^10*d^5*e^6*x^6)/2 + 1220*B*b^10...

```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 1541, normalized size of antiderivative = 3.46

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{11}} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^11,x)`

output

```
(27720*log(d + e*x)*a*b**10*d**11*e + 277200*log(d + e*x)*a*b**10*d**10*e*
*2*x + 1247400*log(d + e*x)*a*b**10*d**9*e**3*x**2 + 3326400*log(d + e*x)*
a*b**10*d**8*e**4*x**3 + 5821200*log(d + e*x)*a*b**10*d**7*e**5*x**4 + 698
5440*log(d + e*x)*a*b**10*d**6*e**6*x**5 + 5821200*log(d + e*x)*a*b**10*d*
*5*e**7*x**6 + 3326400*log(d + e*x)*a*b**10*d**4*e**8*x**7 + 1247400*log(d
+ e*x)*a*b**10*d**3*e**9*x**8 + 277200*log(d + e*x)*a*b**10*d**2*e**10*x*
*9 + 27720*log(d + e*x)*a*b**10*d*e**11*x**10 - 27720*log(d + e*x)*b**11*d
**12 - 277200*log(d + e*x)*b**11*d**11*e*x - 1247400*log(d + e*x)*b**11*d*
*10*e**2*x**2 - 3326400*log(d + e*x)*b**11*d**9*e**3*x**3 - 5821200*log(d
+ e*x)*b**11*d**8*e**4*x**4 - 6985440*log(d + e*x)*b**11*d**7*e**5*x**5 -
5821200*log(d + e*x)*b**11*d**6*e**6*x**6 - 3326400*log(d + e*x)*b**11*d**
5*e**7*x**7 - 1247400*log(d + e*x)*b**11*d**4*e**8*x**8 - 277200*log(d + e
*x)*b**11*d**3*e**9*x**9 - 27720*log(d + e*x)*b**11*d**2*e**10*x**10 - 252
*a**11*d*e**11 - 308*a**10*b*d**2*e**10 - 3080*a**10*b*d*e**11*x - 385*a**
9*b**2*d**3*e**9 - 3850*a**9*b**2*d**2*e**10*x - 17325*a**9*b**2*d*e**11*x
**2 - 495*a**8*b**3*d**4*e**8 - 4950*a**8*b**3*d**3*e**9*x - 22275*a**8*b*
*3*d**2*e**10*x**2 - 59400*a**8*b**3*d*e**11*x**3 - 660*a**7*b**4*d**5*e**
7 - 6600*a**7*b**4*d**4*e**8*x - 29700*a**7*b**4*d**3*e**9*x**2 - 79200*a*
*7*b**4*d**2*e**10*x**3 - 138600*a**7*b**4*d*e**11*x**4 - 924*a**6*b**5*d*
*6*e**6 - 9240*a**6*b**5*d**5*e**7*x - 41580*a**6*b**5*d**4*e**8*x**2 - ...
```

3.90 $\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{12}} dx$

Optimal result	925
Mathematica [B] (verified)	926
Rubi [A] (verified)	927
Maple [B] (verified)	928
Fricas [B] (verification not implemented)	929
Sympy [F(-1)]	930
Maxima [B] (verification not implemented)	931
Giac [B] (verification not implemented)	932
Mupad [B] (verification not implemented)	933
Reduce [B] (verification not implemented)	933

Optimal result

Integrand size = 20, antiderivative size = 321

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{12}} dx = -\frac{(Bd - Ae)(a+bx)^{11}}{11e(bd - ae)(d+ex)^{11}} - \frac{B(bd - ae)^{10}}{10e^{12}(d+ex)^{10}}$$

$$+ \frac{10bB(bd - ae)^9}{9e^{12}(d+ex)^9} - \frac{45b^2B(bd - ae)^8}{8e^{12}(d+ex)^8} + \frac{120b^3B(bd - ae)^7}{7e^{12}(d+ex)^7}$$

$$- \frac{35b^4B(bd - ae)^6}{e^{12}(d+ex)^6} + \frac{252b^5B(bd - ae)^5}{5e^{12}(d+ex)^5}$$

$$- \frac{105b^6B(bd - ae)^4}{2e^{12}(d+ex)^4} + \frac{40b^7B(bd - ae)^3}{e^{12}(d+ex)^3}$$

$$- \frac{45b^8B(bd - ae)^2}{2e^{12}(d+ex)^2} + \frac{10b^9B(bd - ae)}{e^{12}(d+ex)} + \frac{b^{10}B \log(d+ex)}{e^{12}}$$

output

```
-1/11*(-A*e+B*d)*(b*x+a)^11/e/(-a*e+b*d)/(e*x+d)^11-1/10*B*(-a*e+b*d)^10/e
^12/(e*x+d)^10+10/9*b*B*(-a*e+b*d)^9/e^12/(e*x+d)^9-45/8*b^2*B*(-a*e+b*d)^
8/e^12/(e*x+d)^8+120/7*b^3*B*(-a*e+b*d)^7/e^12/(e*x+d)^7-35*b^4*B*(-a*e+b*
d)^6/e^12/(e*x+d)^6+252/5*b^5*B*(-a*e+b*d)^5/e^12/(e*x+d)^5-105/2*b^6*B*(-
a*e+b*d)^4/e^12/(e*x+d)^4+40*b^7*B*(-a*e+b*d)^3/e^12/(e*x+d)^3-45/2*b^8*B*
(-a*e+b*d)^2/e^12/(e*x+d)^2+10*b^9*B*(-a*e+b*d)/e^12/(e*x+d)+b^10*B*ln(e*x
+d)/e^12
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1443 vs. $2(321) = 642$.

Time = 1.22 (sec) , antiderivative size = 1443, normalized size of antiderivative = 4.50

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{12}} dx = \text{Too large to display}$$

input

```
Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^12,x]
```

output

```
-1/27720*(252*a^10*e^10*(10*A*e + B*(d + 11*e*x)) + 280*a^9*b*e^9*(9*A*e*(
d + 11*e*x) + 2*B*(d^2 + 11*d*e*x + 55*e^2*x^2)) + 315*a^8*b^2*e^8*(8*A*e*(
d^2 + 11*d*e*x + 55*e^2*x^2) + 3*B*(d^3 + 11*d^2*e*x + 55*d*e^2*x^2 + 165
*e^3*x^3)) + 360*a^7*b^3*e^7*(7*A*e*(d^3 + 11*d^2*e*x + 55*d*e^2*x^2 + 165
*e^3*x^3) + 4*B*(d^4 + 11*d^3*e*x + 55*d^2*e^2*x^2 + 165*d*e^3*x^3 + 330*e
^4*x^4)) + 420*a^6*b^4*e^6*(6*A*e*(d^4 + 11*d^3*e*x + 55*d^2*e^2*x^2 + 165
*d*e^3*x^3 + 330*e^4*x^4) + 5*B*(d^5 + 11*d^4*e*x + 55*d^3*e^2*x^2 + 165*d
^2*e^3*x^3 + 330*d*e^4*x^4 + 462*e^5*x^5)) + 504*a^5*b^5*e^5*(5*A*e*(d^5 +
11*d^4*e*x + 55*d^3*e^2*x^2 + 165*d^2*e^3*x^3 + 330*d*e^4*x^4 + 462*e^5*x
^5) + 6*B*(d^6 + 11*d^5*e*x + 55*d^4*e^2*x^2 + 165*d^3*e^3*x^3 + 330*d^2*e
^4*x^4 + 462*d*e^5*x^5 + 462*e^6*x^6)) + 630*a^4*b^6*e^4*(4*A*e*(d^6 + 11*
d^5*e*x + 55*d^4*e^2*x^2 + 165*d^3*e^3*x^3 + 330*d^2*e^4*x^4 + 462*d*e^5*x
^5 + 462*e^6*x^6) + 7*B*(d^7 + 11*d^6*e*x + 55*d^5*e^2*x^2 + 165*d^4*e^3*x
^3 + 330*d^3*e^4*x^4 + 462*d^2*e^5*x^5 + 462*d*e^6*x^6 + 330*e^7*x^7)) + 8
40*a^3*b^7*e^3*(3*A*e*(d^7 + 11*d^6*e*x + 55*d^5*e^2*x^2 + 165*d^4*e^3*x^3
+ 330*d^3*e^4*x^4 + 462*d^2*e^5*x^5 + 462*d*e^6*x^6 + 330*e^7*x^7) + 8*B*
(d^8 + 11*d^7*e*x + 55*d^6*e^2*x^2 + 165*d^5*e^3*x^3 + 330*d^4*e^4*x^4 + 4
62*d^3*e^5*x^5 + 462*d^2*e^6*x^6 + 330*d*e^7*x^7 + 165*e^8*x^8)) + 1260*a^
2*b^8*e^2*(2*A*e*(d^8 + 11*d^7*e*x + 55*d^6*e^2*x^2 + 165*d^5*e^3*x^3 + 33
0*d^4*e^4*x^4 + 462*d^3*e^5*x^5 + 462*d^2*e^6*x^6 + 330*d*e^7*x^7 + 165...
```

Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 316, normalized size of antiderivative = 0.98, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {87, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{12}} dx$$

$$\downarrow 87$$

$$\frac{B \int \frac{(a+bx)^{10}}{(d+ex)^{11}} dx}{e} - \frac{(a + bx)^{11}(Bd - Ae)}{11e(d + ex)^{11}(bd - ae)}$$

$$\downarrow 49$$

$$\frac{B \int \left(\frac{b^{10}}{e^{10}(d+ex)} - \frac{10(bd-ae)b^9}{e^{10}(d+ex)^2} + \frac{45(bd-ae)^2b^8}{e^{10}(d+ex)^3} - \frac{120(bd-ae)^3b^7}{e^{10}(d+ex)^4} + \frac{210(bd-ae)^4b^6}{e^{10}(d+ex)^5} - \frac{252(bd-ae)^5b^5}{e^{10}(d+ex)^6} + \frac{210(bd-ae)^6b^4}{e^{10}(d+ex)^7} - \frac{120(bd-ae)^7b^3}{e^{10}(d+ex)^8} + \frac{45(bd-ae)^8b^2}{e^{10}(d+ex)^9} - \frac{10(bd-ae)^9b}{e^{10}(d+ex)^{10}} + \frac{(bd-ae)^{10}}{e^{10}(d+ex)^{11}} \right) dx}{e} - \frac{(a + bx)^{11}(Bd - Ae)}{11e(d + ex)^{11}(bd - ae)}$$

$$\downarrow 2009$$

$$\frac{B \left(\frac{10b^9(bd-ae)}{e^{11}(d+ex)} - \frac{45b^8(bd-ae)^2}{2e^{11}(d+ex)^2} + \frac{40b^7(bd-ae)^3}{e^{11}(d+ex)^3} - \frac{105b^6(bd-ae)^4}{2e^{11}(d+ex)^4} + \frac{252b^5(bd-ae)^5}{5e^{11}(d+ex)^5} - \frac{35b^4(bd-ae)^6}{e^{11}(d+ex)^6} + \frac{120b^3(bd-ae)^7}{7e^{11}(d+ex)^7} - \frac{45b^2(bd-ae)^8}{8e^{11}(d+ex)^8} + \frac{10b(bd-ae)^9}{e^{11}(d+ex)^9} - \frac{(bd-ae)^{10}}{e^{11}(d+ex)^{10}} \right)}{e} - \frac{(a + bx)^{11}(Bd - Ae)}{11e(d + ex)^{11}(bd - ae)}$$

input

```
Int[((a + b*x)^10*(A + B*x))/(d + e*x)^12,x]
```

output

$$\begin{aligned}
& -1/11*((B*d - A*e)*(a + b*x)^{11})/(e*(b*d - a*e)*(d + e*x)^{11}) + (B*(-1/10* \\
& (b*d - a*e)^{10}/(e^{11}*(d + e*x)^{10}) + (10*b*(b*d - a*e)^9)/(9*e^{11}*(d + e*x) \\
&)^9) - (45*b^2*(b*d - a*e)^8)/(8*e^{11}*(d + e*x)^8) + (120*b^3*(b*d - a*e)^7) \\
& / (7*e^{11}*(d + e*x)^7) - (35*b^4*(b*d - a*e)^6)/(e^{11}*(d + e*x)^6) + (252 \\
& *b^5*(b*d - a*e)^5)/(5*e^{11}*(d + e*x)^5) - (105*b^6*(b*d - a*e)^4)/(2*e^{11} \\
& *(d + e*x)^4) + (40*b^7*(b*d - a*e)^3)/(e^{11}*(d + e*x)^3) - (45*b^8*(b*d - \\
& a*e)^2)/(2*e^{11}*(d + e*x)^2) + (10*b^9*(b*d - a*e))/(e^{11}*(d + e*x)) + (b \\
& ^{10}*\text{Log}[d + e*x])/e^{11})/e
\end{aligned}$$

Defintions of rubi rules used

rule 49

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]
```

rule 87

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e)), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1902 vs. $2(305) = 610$.

Time = 0.26 (sec) , antiderivative size = 1903, normalized size of antiderivative = 5.93

method	result	size
risch	Expression too large to display	1903
norman	Expression too large to display	1939
default	Expression too large to display	1940
parallelrisc	Expression too large to display	2458

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^12,x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & (-b^9*(A*b*e+10*B*a*e-11*B*b*d)/e^2*x^{10}-5/2*b^8*(2*A*a*b*e^2+2*A*b^2*d*e+ \\ & 9*B*a^2*e^2+20*B*a*b*d*e-33*B*b^2*d^2)/e^3*x^9-5/2*b^7*(6*A*a^2*b*e^3+6*A* \\ & a*b^2*d*e^2+6*A*b^3*d^2*e+16*B*a^3*e^3+27*B*a^2*b*d*e^2+60*B*a*b^2*d^2*e-1 \\ & 21*B*b^3*d^3)/e^4*x^8-5/2*b^6*(12*A*a^3*b*e^4+12*A*a^2*b^2*d*e^3+12*A*a*b^ \\ & 3*d^2*e^2+12*A*b^4*d^3*e+21*B*a^4*e^4+32*B*a^3*b*d*e^3+54*B*a^2*b^2*d^2*e^ \\ & 2+120*B*a*b^3*d^3*e-275*B*b^4*d^4)/e^5*x^7-7/10*b^5*(60*A*a^4*b*e^5+60*A*a \\ & ^3*b^2*d*e^4+60*A*a^2*b^3*d^2*e^3+60*A*a*b^4*d^3*e^2+60*A*b^5*d^4*e+72*B*a \\ & ^5*e^5+105*B*a^4*b*d*e^4+160*B*a^3*b^2*d^2*e^3+270*B*a^2*b^3*d^3*e^2+600*B \\ & *a*b^4*d^4*e-1507*B*b^5*d^5)/e^6*x^6-7/10*b^4*(60*A*a^5*b*e^6+60*A*a^4*b^2 \\ & *d*e^5+60*A*a^3*b^3*d^2*e^4+60*A*a^2*b^4*d^3*e^3+60*A*a*b^5*d^4*e^2+60*A*b \\ & ^6*d^5*e+50*B*a^6*e^6+72*B*a^5*b*d*e^5+105*B*a^4*b^2*d^2*e^4+160*B*a^3*b^3 \\ & *d^3*e^3+270*B*a^2*b^4*d^4*e^2+600*B*a*b^5*d^5*e-1617*B*b^6*d^6)/e^7*x^5-1 \\ & /14*b^3*(420*A*a^6*b*e^7+420*A*a^5*b^2*d*e^6+420*A*a^4*b^3*d^2*e^5+420*A*a \\ & ^3*b^4*d^3*e^4+420*A*a^2*b^5*d^4*e^3+420*A*a*b^6*d^5*e^2+420*A*b^7*d^6*e+2 \\ & 40*B*a^7*e^7+350*B*a^6*b*d*e^6+504*B*a^5*b^2*d^2*e^5+735*B*a^4*b^3*d^3*e^4 \\ & +1120*B*a^3*b^4*d^4*e^3+1890*B*a^2*b^5*d^5*e^2+4200*B*a*b^6*d^6*e-11979*B* \\ & b^7*d^7)/e^8*x^4-1/56*b^2*(840*A*a^7*b*e^8+840*A*a^6*b^2*d*e^7+840*A*a^5*b \\ & ^3*d^2*e^6+840*A*a^4*b^4*d^3*e^5+840*A*a^3*b^5*d^4*e^4+840*A*a^2*b^6*d^5*e \\ & ^3+840*A*a*b^7*d^6*e^2+840*A*b^8*d^7*e+315*B*a^8*e^8+480*B*a^7*b*d*e^7+700 \\ & *B*a^6*b^2*d^2*e^6+1008*B*a^5*b^3*d^3*e^5+1470*B*a^4*b^4*d^4*e^4+2240*B... \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2089 vs. $2(305) = 610$.

Time = 0.17 (sec) , antiderivative size = 2089, normalized size of antiderivative = 6.51

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{12}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^12,x, algorithm="fricas")`

output

```

1/27720*(83711*B*b^10*d^11 - 2520*A*a^10*e^11 - 2520*(10*B*a*b^9 + A*b^10)
*d^10*e - 1260*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 - 840*(8*B*a^3*b^7 + 3*A*
a^2*b^8)*d^8*e^3 - 630*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e^4 - 504*(6*B*a^5*
b^5 + 5*A*a^4*b^6)*d^6*e^5 - 420*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^5*e^6 - 360
*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 - 315*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^3
*e^8 - 280*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 - 252*(B*a^10 + 10*A*a^9*b)*d
*e^10 + 27720*(11*B*b^10*d*e^10 - (10*B*a*b^9 + A*b^10)*e^11)*x^10 + 69300
*(33*B*b^10*d^2*e^9 - 2*(10*B*a*b^9 + A*b^10)*d*e^10 - (9*B*a^2*b^8 + 2*A*
a*b^9)*e^11)*x^9 + 69300*(121*B*b^10*d^3*e^8 - 6*(10*B*a*b^9 + A*b^10)*d^2
*e^9 - 3*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 - 2*(8*B*a^3*b^7 + 3*A*a^2*b^8)*
e^11)*x^8 + 69300*(275*B*b^10*d^4*e^7 - 12*(10*B*a*b^9 + A*b^10)*d^3*e^8 -
6*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 - 4*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^1
0 - 3*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7 + 19404*(1507*B*b^10*d^5*e^6 -
60*(10*B*a*b^9 + A*b^10)*d^4*e^7 - 30*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 -
20*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 - 15*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d
*e^10 - 12*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 + 19404*(1617*B*b^10*d^6*
e^5 - 60*(10*B*a*b^9 + A*b^10)*d^5*e^6 - 30*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*
e^7 - 20*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 - 15*(7*B*a^4*b^6 + 4*A*a^3*b
^7)*d^2*e^9 - 12*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10 - 10*(5*B*a^6*b^4 + 6*
A*a^5*b^5)*e^11)*x^5 + 1980*(11979*B*b^10*d^7*e^4 - 420*(10*B*a*b^9 + A...

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{12}} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**10*(B*x+A)/(e*x+d)**12,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1932 vs. $2(305) = 610$.

Time = 0.12 (sec) , antiderivative size = 1932, normalized size of antiderivative = 6.02

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{12}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^12,x, algorithm="maxima")`

output

```
1/27720*(83711*B*b^10*d^11 - 2520*A*a^10*e^11 - 2520*(10*B*a*b^9 + A*b^10)
*d^10*e - 1260*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 - 840*(8*B*a^3*b^7 + 3*A*
a^2*b^8)*d^8*e^3 - 630*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e^4 - 504*(6*B*a^5*
b^5 + 5*A*a^4*b^6)*d^6*e^5 - 420*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^5*e^6 - 360
*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 - 315*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^3
*e^8 - 280*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 - 252*(B*a^10 + 10*A*a^9*b)*d
*e^10 + 27720*(11*B*b^10*d*e^10 - (10*B*a*b^9 + A*b^10)*e^11)*x^10 + 69300
*(33*B*b^10*d^2*e^9 - 2*(10*B*a*b^9 + A*b^10)*d*e^10 - (9*B*a^2*b^8 + 2*A*
a*b^9)*e^11)*x^9 + 69300*(121*B*b^10*d^3*e^8 - 6*(10*B*a*b^9 + A*b^10)*d^2
*e^9 - 3*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 - 2*(8*B*a^3*b^7 + 3*A*a^2*b^8)*
e^11)*x^8 + 69300*(275*B*b^10*d^4*e^7 - 12*(10*B*a*b^9 + A*b^10)*d^3*e^8 -
6*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 - 4*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^1
0 - 3*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7 + 19404*(1507*B*b^10*d^5*e^6 -
60*(10*B*a*b^9 + A*b^10)*d^4*e^7 - 30*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 -
20*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 - 15*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d
*e^10 - 12*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 + 19404*(1617*B*b^10*d^6*
e^5 - 60*(10*B*a*b^9 + A*b^10)*d^5*e^6 - 30*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*
e^7 - 20*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 - 15*(7*B*a^4*b^6 + 4*A*a^3*b
^7)*d^2*e^9 - 12*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10 - 10*(5*B*a^6*b^4 + 6*
A*a^5*b^5)*e^11)*x^5 + 1980*(11979*B*b^10*d^7*e^4 - 420*(10*B*a*b^9 + A...
```


Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1974 vs. $2(305) = 610$.

Time = 0.13 (sec) , antiderivative size = 1974, normalized size of antiderivative = 6.15

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{12}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^12,x, algorithm="giac")`

output

```
B*b^10*log(abs(e*x + d))/e^12 + 1/27720*(27720*(11*B*b^10*d*e^9 - 10*B*a*b^9*e^10 - A*b^10*e^10)*x^10 + 69300*(33*B*b^10*d^2*e^8 - 20*B*a*b^9*d*e^9 - 2*A*b^10*d*e^9 - 9*B*a^2*b^8*e^10 - 2*A*a*b^9*e^10)*x^9 + 69300*(121*B*b^10*d^3*e^7 - 60*B*a*b^9*d^2*e^8 - 6*A*b^10*d^2*e^8 - 27*B*a^2*b^8*d*e^9 - 6*A*a*b^9*d*e^9 - 16*B*a^3*b^7*e^10 - 6*A*a^2*b^8*e^10)*x^8 + 69300*(275*B*b^10*d^4*e^6 - 120*B*a*b^9*d^3*e^7 - 12*A*b^10*d^3*e^7 - 54*B*a^2*b^8*d^2*e^8 - 12*A*a*b^9*d^2*e^8 - 32*B*a^3*b^7*d*e^9 - 12*A*a^2*b^8*d*e^9 - 21*B*a^4*b^6*e^10 - 12*A*a^3*b^7*e^10)*x^7 + 19404*(1507*B*b^10*d^5*e^5 - 600*B*a*b^9*d^4*e^6 - 60*A*b^10*d^4*e^6 - 270*B*a^2*b^8*d^3*e^7 - 60*A*a*b^9*d^3*e^7 - 160*B*a^3*b^7*d^2*e^8 - 60*A*a^2*b^8*d^2*e^8 - 105*B*a^4*b^6*d*e^9 - 60*A*a^3*b^7*d*e^9 - 72*B*a^5*b^5*e^10 - 60*A*a^4*b^6*e^10)*x^6 + 19404*(1617*B*b^10*d^6*e^4 - 600*B*a*b^9*d^5*e^5 - 60*A*b^10*d^5*e^5 - 270*B*a^2*b^8*d^4*e^6 - 60*A*a*b^9*d^4*e^6 - 160*B*a^3*b^7*d^3*e^7 - 60*A*a^2*b^8*d^3*e^7 - 105*B*a^4*b^6*d^2*e^8 - 60*A*a^3*b^7*d^2*e^8 - 72*B*a^5*b^5*d*e^9 - 60*A*a^4*b^6*d*e^9 - 50*B*a^6*b^4*e^10 - 60*A*a^5*b^5*e^10)*x^5 + 1980*(11979*B*b^10*d^7*e^3 - 4200*B*a*b^9*d^6*e^4 - 420*A*b^10*d^6*e^4 - 1890*B*a^2*b^8*d^5*e^5 - 420*A*a*b^9*d^5*e^5 - 1120*B*a^3*b^7*d^4*e^6 - 420*A*a^2*b^8*d^4*e^6 - 735*B*a^4*b^6*d^3*e^7 - 420*A*a^3*b^7*d^3*e^7 - 504*B*a^5*b^5*d^2*e^8 - 420*A*a^4*b^6*d^2*e^8 - 350*B*a^6*b^4*d*e^9 - 420*A*a^5*b^5*d*e^9 - 240*B*a^7*b^3*e^10 - 420*A*a^6*b^4*e^10)*x^4 + 495*(25113*B*...
```


output

```
(27720*log(d + e*x)*b**11*d**12 + 304920*log(d + e*x)*b**11*d**11*e*x + 15
24600*log(d + e*x)*b**11*d**10*e**2*x**2 + 4573800*log(d + e*x)*b**11*d**9
*e**3*x**3 + 9147600*log(d + e*x)*b**11*d**8*e**4*x**4 + 12806640*log(d +
e*x)*b**11*d**7*e**5*x**5 + 12806640*log(d + e*x)*b**11*d**6*e**6*x**6 + 9
147600*log(d + e*x)*b**11*d**5*e**7*x**7 + 4573800*log(d + e*x)*b**11*d**4
*e**8*x**8 + 1524600*log(d + e*x)*b**11*d**3*e**9*x**9 + 304920*log(d + e
x)*b**11*d**2*e**10*x**10 + 27720*log(d + e*x)*b**11*d*e**11*x**11 - 2520*
a**11*d*e**11 - 2772*a**10*b*d**2*e**10 - 30492*a**10*b*d*e**11*x - 3080*a
**9*b**2*d**3*e**9 - 33880*a**9*b**2*d**2*e**10*x - 169400*a**9*b**2*d*e**
11*x**2 - 3465*a**8*b**3*d**4*e**8 - 38115*a**8*b**3*d**3*e**9*x - 190575*
a**8*b**3*d**2*e**10*x**2 - 571725*a**8*b**3*d*e**11*x**3 - 3960*a**7*b**4
*d**5*e**7 - 43560*a**7*b**4*d**4*e**8*x - 217800*a**7*b**4*d**3*e**9*x**2
- 653400*a**7*b**4*d**2*e**10*x**3 - 1306800*a**7*b**4*d*e**11*x**4 - 462
0*a**6*b**5*d**6*e**6 - 50820*a**6*b**5*d**5*e**7*x - 254100*a**6*b**5*d**
4*e**8*x**2 - 762300*a**6*b**5*d**3*e**9*x**3 - 1524600*a**6*b**5*d**2*e**
10*x**4 - 2134440*a**6*b**5*d*e**11*x**5 - 5544*a**5*b**6*d**7*e**5 - 6098
4*a**5*b**6*d**6*e**6*x - 304920*a**5*b**6*d**5*e**7*x**2 - 914760*a**5*b*
**6*d**4*e**8*x**3 - 1829520*a**5*b**6*d**3*e**9*x**4 - 2561328*a**5*b**6*d
**2*e**10*x**5 - 2561328*a**5*b**6*d*e**11*x**6 - 6930*a**4*b**7*d**8*e**4
- 76230*a**4*b**7*d**7*e**5*x - 381150*a**4*b**7*d**6*e**6*x**2 - 1143...
```

3.91 $\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{13}} dx$

Optimal result	935
Mathematica [B] (verified)	935
Rubi [A] (verified)	936
Maple [B] (verified)	937
Fricas [B] (verification not implemented)	938
Sympy [F(-1)]	939
Maxima [B] (verification not implemented)	940
Giac [B] (verification not implemented)	941
Mupad [B] (verification not implemented)	942
Reduce [B] (verification not implemented)	942

Optimal result

Integrand size = 20, antiderivative size = 86

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{13}} dx = -\frac{(Bd - Ae)(a+bx)^{11}}{12e(bd - ae)(d+ex)^{12}} + \frac{(11bBd + Abe - 12aBe)(a+bx)^{11}}{132e(bd - ae)^2(d+ex)^{11}}$$

output

```
-1/12*(-A*e+B*d)*(b*x+a)^11/e/(-a*e+b*d)/(e*x+d)^12+1/132*(A*b*e-12*B*a*e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^2/(e*x+d)^11
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1421 vs. 2(86) = 172.

Time = 0.52 (sec) , antiderivative size = 1421, normalized size of antiderivative = 16.52

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{13}} dx = \text{Too large to display}$$

input

```
Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^13,x]
```

output

```

-1/132*(a^10*e^10*(11*A*e + B*(d + 12*e*x)) + 2*a^9*b*e^9*(5*A*e*(d + 12*e
*x) + B*(d^2 + 12*d*e*x + 66*e^2*x^2)) + 3*a^8*b^2*e^8*(3*A*e*(d^2 + 12*d*
e*x + 66*e^2*x^2) + B*(d^3 + 12*d^2*e*x + 66*d*e^2*x^2 + 220*e^3*x^3)) + 4
*a^7*b^3*e^7*(2*A*e*(d^3 + 12*d^2*e*x + 66*d*e^2*x^2 + 220*e^3*x^3) + B*(d
^4 + 12*d^3*e*x + 66*d^2*e^2*x^2 + 220*d*e^3*x^3 + 495*e^4*x^4)) + a^6*b^4
*e^6*(7*A*e*(d^4 + 12*d^3*e*x + 66*d^2*e^2*x^2 + 220*d*e^3*x^3 + 495*e^4*x
^4) + 5*B*(d^5 + 12*d^4*e*x + 66*d^3*e^2*x^2 + 220*d^2*e^3*x^3 + 495*d*e^4
*x^4 + 792*e^5*x^5)) + 6*a^5*b^5*e^5*(A*e*(d^5 + 12*d^4*e*x + 66*d^3*e^2*x
^2 + 220*d^2*e^3*x^3 + 495*d*e^4*x^4 + 792*e^5*x^5) + B*(d^6 + 12*d^5*e*x
+ 66*d^4*e^2*x^2 + 220*d^3*e^3*x^3 + 495*d^2*e^4*x^4 + 792*d*e^5*x^5 + 924
*e^6*x^6)) + a^4*b^6*e^4*(5*A*e*(d^6 + 12*d^5*e*x + 66*d^4*e^2*x^2 + 220*d
^3*e^3*x^3 + 495*d^2*e^4*x^4 + 792*d*e^5*x^5 + 924*e^6*x^6) + 7*B*(d^7 + 1
2*d^6*e*x + 66*d^5*e^2*x^2 + 220*d^4*e^3*x^3 + 495*d^3*e^4*x^4 + 792*d^2*e
^5*x^5 + 924*d*e^6*x^6 + 792*e^7*x^7)) + 4*a^3*b^7*e^3*(A*e*(d^7 + 12*d^6*
e*x + 66*d^5*e^2*x^2 + 220*d^4*e^3*x^3 + 495*d^3*e^4*x^4 + 792*d^2*e^5*x^5
+ 924*d*e^6*x^6 + 792*e^7*x^7) + 2*B*(d^8 + 12*d^7*e*x + 66*d^6*e^2*x^2 +
220*d^5*e^3*x^3 + 495*d^4*e^4*x^4 + 792*d^3*e^5*x^5 + 924*d^2*e^6*x^6 + 7
92*d*e^7*x^7 + 495*e^8*x^8)) + 3*a^2*b^8*e^2*(A*e*(d^8 + 12*d^7*e*x + 66*d
^6*e^2*x^2 + 220*d^5*e^3*x^3 + 495*d^4*e^4*x^4 + 792*d^3*e^5*x^5 + 924*d^2
*e^6*x^6 + 792*d*e^7*x^7 + 495*e^8*x^8) + 3*B*(d^9 + 12*d^8*e*x + 66*d^...

```

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {87, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{13}} dx$$

$$\downarrow 87$$

$$\frac{(-12aBe + Abe + 11bBd) \int \frac{(a+bx)^{10}}{(d+ex)^{12}} dx}{12e(bd - ae)} - \frac{(a + bx)^{11}(Bd - Ae)}{12e(d + ex)^{12}(bd - ae)}$$

$$\downarrow 48$$

$$\frac{(a + bx)^{11}(-12aBe + Abe + 11bBd)}{132e(d + ex)^{11}(bd - ae)^2} - \frac{(a + bx)^{11}(Bd - Ae)}{12e(d + ex)^{12}(bd - ae)}$$

input `Int[((a + b*x)^10*(A + B*x))/(d + e*x)^13,x]`

output `-1/12*((B*d - A*e)*(a + b*x)^11)/(e*(b*d - a*e)*(d + e*x)^12) + ((11*b*B*d + A*b*e - 12*a*B*e)*(a + b*x)^11)/(132*e*(b*d - a*e)^2*(d + e*x)^11)`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1887 vs. 2(82) = 164.

Time = 0.26 (sec) , antiderivative size = 1888, normalized size of antiderivative = 21.95

method	result	size
risch	Expression too large to display	1888
norman	Expression too large to display	1924
default	Expression too large to display	1942
gosper	Expression too large to display	2231
parallelrisc	Expression too large to display	2231
orering	Expression too large to display	2231

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^13,x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & (-b^{10}B/e^{11}x^{11} - 1/2*b^9*(A*b*e + 10*B*a*e + 11*B*b*d)/e^{10}x^{10} - 5/3*b^8*(2*A*a* \\ & b*e^2 + A*b^2*d*e + 9*B*a^2*e^2 + 10*B*a*b*d*e + 11*B*b^2*d^2)/e^9x^9 - 15/4*b^7*(3 \\ & *A*a^2*b*e^3 + 2*A*a*b^2*d*e^2 + A*b^3*d^2*e + 8*B*a^3*e^3 + 9*B*a^2*b*d*e^2 + 10*B* \\ & a*b^2*d^2*e + 11*B*b^3*d^3)/e^8x^8 - 6*b^6*(4*A*a^3*b*e^4 + 3*A*a^2*b^2*d*e^3 + 2 \\ & *A*a*b^3*d^2*e^2 + A*b^4*d^3*e + 7*B*a^4*e^4 + 8*B*a^3*b*d*e^3 + 9*B*a^2*b^2*d^2*e \\ & ^2 + 10*B*a*b^3*d^3*e + 11*B*b^4*d^4)/e^7x^7 - 7*b^5*(5*A*a^4*b*e^5 + 4*A*a^3*b^2 \\ & *d*e^4 + 3*A*a^2*b^3*d^2*e^3 + 2*A*a*b^4*d^3*e^2 + A*b^5*d^4*e + 6*B*a^5*e^5 + 7*B*a \\ & ^4*b*d*e^4 + 8*B*a^3*b^2*d^2*e^3 + 9*B*a^2*b^3*d^3*e^2 + 10*B*a*b^4*d^4*e + 11*B*b \\ & ^5*d^5)/e^6x^6 - 6*b^4*(6*A*a^5*b*e^6 + 5*A*a^4*b^2*d*e^5 + 4*A*a^3*b^3*d^2*e^4 \\ & + 3*A*a^2*b^4*d^3*e^3 + 2*A*a*b^5*d^4*e^2 + A*b^6*d^5*e + 5*B*a^6*e^6 + 6*B*a^5*b*d \\ & *e^5 + 7*B*a^4*b^2*d^2*e^4 + 8*B*a^3*b^3*d^3*e^3 + 9*B*a^2*b^4*d^4*e^2 + 10*B*a*b^ \\ & 5*d^5*e + 11*B*b^6*d^6)/e^5x^5 - 15/4*b^3*(7*A*a^6*b*e^7 + 6*A*a^5*b^2*d*e^6 + 5* \\ & A*a^4*b^3*d^2*e^5 + 4*A*a^3*b^4*d^3*e^4 + 3*A*a^2*b^5*d^4*e^3 + 2*A*a*b^6*d^5*e^ \\ & 2 + A*b^7*d^6*e + 4*B*a^7*e^7 + 5*B*a^6*b*d*e^6 + 6*B*a^5*b^2*d^2*e^5 + 7*B*a^4*b^3* \\ & d^3*e^4 + 8*B*a^3*b^4*d^4*e^3 + 9*B*a^2*b^5*d^5*e^2 + 10*B*a*b^6*d^6*e + 11*B*b^7* \\ & d^7)/e^4x^4 - 5/3*b^2*(8*A*a^7*b*e^8 + 7*A*a^6*b^2*d*e^7 + 6*A*a^5*b^3*d^2*e^6 + \\ & 5*A*a^4*b^4*d^3*e^5 + 4*A*a^3*b^5*d^4*e^4 + 3*A*a^2*b^6*d^5*e^3 + 2*A*a*b^7*d^6* \\ & e^2 + A*b^8*d^7*e + 3*B*a^8*e^8 + 4*B*a^7*b*d*e^7 + 5*B*a^6*b^2*d^2*e^6 + 6*B*a^5*b^ \\ & 3*d^3*e^5 + 7*B*a^4*b^4*d^4*e^4 + 8*B*a^3*b^5*d^5*e^3 + 9*B*a^2*b^6*d^6*e^2 + 10*B \\ & *a*b^7*d^7*e + 11*B*b^8*d^8)/e^3x^3 - 1/2*b*(9*A*a^8*b*e^9 + 8*A*a^7*b^2*d*e \dots \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1875 vs. $2(82) = 164$.

Time = 0.14 (sec) , antiderivative size = 1875, normalized size of antiderivative = 21.80

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{13}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^13,x, algorithm="fricas")`

output

```

-1/132*(132*B*b^10*e^11*x^11 + 11*B*b^10*d^11 + 11*A*a^10*e^11 + (10*B*a*b
^9 + A*b^10)*d^10*e + (9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 + (8*B*a^3*b^7 + 3
*A*a^2*b^8)*d^8*e^3 + (7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e^4 + (6*B*a^5*b^5 +
5*A*a^4*b^6)*d^6*e^5 + (5*B*a^6*b^4 + 6*A*a^5*b^5)*d^5*e^6 + (4*B*a^7*b^3
+ 7*A*a^6*b^4)*d^4*e^7 + (3*B*a^8*b^2 + 8*A*a^7*b^3)*d^3*e^8 + (2*B*a^9*b
+ 9*A*a^8*b^2)*d^2*e^9 + (B*a^10 + 10*A*a^9*b)*d*e^10 + 66*(11*B*b^10*d*e
^10 + (10*B*a*b^9 + A*b^10)*e^11)*x^10 + 220*(11*B*b^10*d^2*e^9 + (10*B*a*
b^9 + A*b^10)*d*e^10 + (9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 + 495*(11*B*b^1
0*d^3*e^8 + (10*B*a*b^9 + A*b^10)*d^2*e^9 + (9*B*a^2*b^8 + 2*A*a*b^9)*d*e
^10 + (8*B*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 792*(11*B*b^10*d^4*e^7 + (10*
B*a*b^9 + A*b^10)*d^3*e^8 + (9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 + (8*B*a^3*b
^7 + 3*A*a^2*b^8)*d*e^10 + (7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7 + 924*(11
*B*b^10*d^5*e^6 + (10*B*a*b^9 + A*b^10)*d^4*e^7 + (9*B*a^2*b^8 + 2*A*a*b^9
)*d^3*e^8 + (8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + (7*B*a^4*b^6 + 4*A*a^3*b
^7)*d*e^10 + (6*B*a^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 + 792*(11*B*b^10*d^6*e
^5 + (10*B*a*b^9 + A*b^10)*d^5*e^6 + (9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 + (8
*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 + (7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 +
(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10 + (5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5
+ 495*(11*B*b^10*d^7*e^4 + (10*B*a*b^9 + A*b^10)*d^6*e^5 + (9*B*a^2*b^8 +
2*A*a*b^9)*d^5*e^6 + (8*B*a^3*b^7 + 3*A*a^2*b^8)*d^4*e^7 + (7*B*a^4*b^...

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{13}} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**10*(B*x+A)/(e*x+d)**13,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1875 vs. $2(82) = 164$.

Time = 0.13 (sec) , antiderivative size = 1875, normalized size of antiderivative = 21.80

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{13}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^13,x, algorithm="maxima")`

output

```
-1/132*(132*B*b^10*e^11*x^11 + 11*B*b^10*d^11 + 11*A*a^10*e^11 + (10*B*a*b^9 + A*b^10)*d^10*e + (9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 + (8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + (7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e^4 + (6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + (5*B*a^6*b^4 + 6*A*a^5*b^5)*d^5*e^6 + (4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + (3*B*a^8*b^2 + 8*A*a^7*b^3)*d^3*e^8 + (2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 + (B*a^10 + 10*A*a^9*b)*d*e^10 + 66*(11*B*b^10*d*e^10 + (10*B*a*b^9 + A*b^10)*e^11)*x^10 + 220*(11*B*b^10*d^2*e^9 + (10*B*a*b^9 + A*b^10)*d*e^10 + (9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 + 495*(11*B*b^10*d^3*e^8 + (10*B*a*b^9 + A*b^10)*d^2*e^9 + (9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 + (8*B*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 792*(11*B*b^10*d^4*e^7 + (10*B*a*b^9 + A*b^10)*d^3*e^8 + (9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 + (8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 + (7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7 + 924*(11*B*b^10*d^5*e^6 + (10*B*a*b^9 + A*b^10)*d^4*e^7 + (9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 + (8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + (7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 + (6*B*a^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 + 792*(11*B*b^10*d^6*e^5 + (10*B*a*b^9 + A*b^10)*d^5*e^6 + (9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 + (8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 + (7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 + (6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10 + (5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 + 495*(11*B*b^10*d^7*e^4 + (10*B*a*b^9 + A*b^10)*d^6*e^5 + (9*B*a^2*b^8 + 2*A*a*b^9)*d^5*e^6 + (8*B*a^3*b^7 + 3*A*a^2*b^8)*d^4*e^7 + (7*B*a^4*b^...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2230 vs. $2(82) = 164$.

Time = 0.13 (sec) , antiderivative size = 2230, normalized size of antiderivative = 25.93

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{13}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^13,x, algorithm="giac")`

output

```
-1/132*(132*B*b^10*e^11*x^11 + 726*B*b^10*d*e^10*x^10 + 660*B*a*b^9*e^11*x
^10 + 66*A*b^10*e^11*x^10 + 2420*B*b^10*d^2*e^9*x^9 + 2200*B*a*b^9*d*e^10*
x^9 + 220*A*b^10*d*e^10*x^9 + 1980*B*a^2*b^8*e^11*x^9 + 440*A*a*b^9*e^11*x
^9 + 5445*B*b^10*d^3*e^8*x^8 + 4950*B*a*b^9*d^2*e^9*x^8 + 495*A*b^10*d^2*e
^9*x^8 + 4455*B*a^2*b^8*d*e^10*x^8 + 990*A*a*b^9*d*e^10*x^8 + 3960*B*a^3*b
^7*e^11*x^8 + 1485*A*a^2*b^8*e^11*x^8 + 8712*B*b^10*d^4*e^7*x^7 + 7920*B*a
*b^9*d^3*e^8*x^7 + 792*A*b^10*d^3*e^8*x^7 + 7128*B*a^2*b^8*d^2*e^9*x^7 + 1
584*A*a*b^9*d^2*e^9*x^7 + 6336*B*a^3*b^7*d*e^10*x^7 + 2376*A*a^2*b^8*d*e^1
0*x^7 + 5544*B*a^4*b^6*e^11*x^7 + 3168*A*a^3*b^7*e^11*x^7 + 10164*B*b^10*d
^5*e^6*x^6 + 9240*B*a*b^9*d^4*e^7*x^6 + 924*A*b^10*d^4*e^7*x^6 + 8316*B*a^
2*b^8*d^3*e^8*x^6 + 1848*A*a*b^9*d^3*e^8*x^6 + 7392*B*a^3*b^7*d^2*e^9*x^6
+ 2772*A*a^2*b^8*d^2*e^9*x^6 + 6468*B*a^4*b^6*d*e^10*x^6 + 3696*A*a^3*b^7*
d*e^10*x^6 + 5544*B*a^5*b^5*e^11*x^6 + 4620*A*a^4*b^6*e^11*x^6 + 8712*B*b^
10*d^6*e^5*x^5 + 7920*B*a*b^9*d^5*e^6*x^5 + 792*A*b^10*d^5*e^6*x^5 + 7128*
B*a^2*b^8*d^4*e^7*x^5 + 1584*A*a*b^9*d^4*e^7*x^5 + 6336*B*a^3*b^7*d^3*e^8*
x^5 + 2376*A*a^2*b^8*d^3*e^8*x^5 + 5544*B*a^4*b^6*d^2*e^9*x^5 + 3168*A*a^3
*b^7*d^2*e^9*x^5 + 4752*B*a^5*b^5*d*e^10*x^5 + 3960*A*a^4*b^6*d*e^10*x^5 +
3960*B*a^6*b^4*e^11*x^5 + 4752*A*a^5*b^5*e^11*x^5 + 5445*B*b^10*d^7*e^4*x
^4 + 4950*B*a*b^9*d^6*e^5*x^4 + 495*A*b^10*d^6*e^5*x^4 + 4455*B*a^2*b^8*d^
5*e^6*x^4 + 990*A*a*b^9*d^5*e^6*x^4 + 3960*B*a^3*b^7*d^4*e^7*x^4 + 1485...
```

Mupad [B] (verification not implemented)

Time = 0.47 (sec) , antiderivative size = 2008, normalized size of antiderivative = 23.35

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{13}} dx = \text{Too large to display}$$

input `int((A + B*x)*(a + b*x)^10)/(d + e*x)^13,x`

output `-((11*A*a^10*e^11 + 11*B*b^10*d^11 + A*b^10*d^10*e + B*a^10*d*e^10 + 2*A*a*b^9*d^9*e^2 + 2*B*a^9*b*d^2*e^9 + 3*A*a^2*b^8*d^8*e^3 + 4*A*a^3*b^7*d^7*e^4 + 5*A*a^4*b^6*d^6*e^5 + 6*A*a^5*b^5*d^5*e^6 + 7*A*a^6*b^4*d^4*e^7 + 8*A*a^7*b^3*d^3*e^8 + 9*A*a^8*b^2*d^2*e^9 + 9*B*a^2*b^8*d^9*e^2 + 8*B*a^3*b^7*d^8*e^3 + 7*B*a^4*b^6*d^7*e^4 + 6*B*a^5*b^5*d^6*e^5 + 5*B*a^6*b^4*d^5*e^6 + 4*B*a^7*b^3*d^4*e^7 + 3*B*a^8*b^2*d^3*e^8 + 10*A*a^9*b*d*e^10 + 10*B*a*b^9*d^10*e)/(132*e^12) + (x*(B*a^10*e^10 + 11*B*b^10*d^10 + 10*A*a^9*b*e^10 + A*b^10*d^9*e + 2*A*a*b^9*d^8*e^2 + 9*A*a^8*b^2*d*e^9 + 3*A*a^2*b^8*d^7*e^3 + 4*A*a^3*b^7*d^6*e^4 + 5*A*a^4*b^6*d^5*e^5 + 6*A*a^5*b^5*d^4*e^6 + 7*A*a^6*b^4*d^3*e^7 + 8*A*a^7*b^3*d^2*e^8 + 9*B*a^2*b^8*d^8*e^2 + 8*B*a^3*b^7*d^7*e^3 + 7*B*a^4*b^6*d^6*e^4 + 6*B*a^5*b^5*d^5*e^5 + 5*B*a^6*b^4*d^4*e^6 + 4*B*a^7*b^3*d^3*e^7 + 3*B*a^8*b^2*d^2*e^8 + 10*B*a*b^9*d^9*e + 2*B*a^9*b*d*e^9))/(11*e^11) + (15*b^7*x^8*(8*B*a^3*e^3 + 11*B*b^3*d^3 + 3*A*a^2*b*e^3 + A*b^3*d^2*e + 2*A*a*b^2*d*e^2 + 10*B*a*b^2*d^2*e + 9*B*a^2*b*d*e^2))/(4*e^4) + (6*b^4*x^5*(5*B*a^6*e^6 + 11*B*b^6*d^6 + 6*A*a^5*b*e^6 + A*b^6*d^5*e + 2*A*a*b^5*d^4*e^2 + 5*A*a^4*b^2*d*e^5 + 3*A*a^2*b^4*d^3*e^3 + 4*A*a^3*b^3*d^2*e^4 + 9*B*a^2*b^4*d^4*e^2 + 8*B*a^3*b^3*d^3*e^3 + 7*B*a^4*b^2*d^2*e^4 + 10*B*a*b^5*d^5*e + 6*B*a^5*b*d*e^5))/e^7 + (b^9*x^10*(A*b*e + 10*B*a*e + 11*B*b*d))/(2*e^2) + (6*b^6*x^7*(7*B*a^4*e^4 + 11*B*b^4*d^4 + 4*A*a^3*b*e^4 + A*b^4*d^3*e + 2*A*a*b^3*d^2*e^2 + 3*A*a^2*b^2*d*e^3 + 9*...`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 1158, normalized size of antiderivative = 13.47

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{13}} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^13,x)`

output

```
( - a**11*d**10 - a**10*b*d**2*e**9 - 12*a**10*b*d**10*x - a**9*b**2*d
**3*e**8 - 12*a**9*b**2*d**2*e**9*x - 66*a**9*b**2*d**10*x**2 - a**8*b**
3*d**4*e**7 - 12*a**8*b**3*d**3*e**8*x - 66*a**8*b**3*d**2*e**9*x**2 - 220
*a**8*b**3*d**10*x**3 - a**7*b**4*d**5*e**6 - 12*a**7*b**4*d**4*e**7*x -
66*a**7*b**4*d**3*e**8*x**2 - 220*a**7*b**4*d**2*e**9*x**3 - 495*a**7*b**
4*d**10*x**4 - a**6*b**5*d**6*e**5 - 12*a**6*b**5*d**5*e**6*x - 66*a**6*
b**5*d**4*e**7*x**2 - 220*a**6*b**5*d**3*e**8*x**3 - 495*a**6*b**5*d**2*e
**9*x**4 - 792*a**6*b**5*d**10*x**5 - a**5*b**6*d**7*e**4 - 12*a**5*b**6*
d**6*e**5*x - 66*a**5*b**6*d**5*e**6*x**2 - 220*a**5*b**6*d**4*e**7*x**3 -
495*a**5*b**6*d**3*e**8*x**4 - 792*a**5*b**6*d**2*e**9*x**5 - 924*a**5*b**
6*d**10*x**6 - a**4*b**7*d**8*e**3 - 12*a**4*b**7*d**7*e**4*x - 66*a**4
*b**7*d**6*e**5*x**2 - 220*a**4*b**7*d**5*e**6*x**3 - 495*a**4*b**7*d**4*
e**7*x**4 - 792*a**4*b**7*d**3*e**8*x**5 - 924*a**4*b**7*d**2*e**9*x**6 - 7
92*a**4*b**7*d**10*x**7 - a**3*b**8*d**9*e**2 - 12*a**3*b**8*d**8*e**3*x
- 66*a**3*b**8*d**7*e**4*x**2 - 220*a**3*b**8*d**6*e**5*x**3 - 495*a**3*b
**8*d**5*e**6*x**4 - 792*a**3*b**8*d**4*e**7*x**5 - 924*a**3*b**8*d**3*
e**8*x**6 - 792*a**3*b**8*d**2*e**9*x**7 - 495*a**3*b**8*d**10*x**8 - a**2*
b**9*d**10*e - 12*a**2*b**9*d**9*e**2*x - 66*a**2*b**9*d**8*e**3*x**2 - 22
0*a**2*b**9*d**7*e**4*x**3 - 495*a**2*b**9*d**6*e**5*x**4 - 792*a**2*b**9*
d**5*e**6*x**5 - 924*a**2*b**9*d**4*e**7*x**6 - 792*a**2*b**9*d**3*e**8...
```

3.92 $\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{14}} dx$

Optimal result	944
Mathematica [B] (verified)	944
Rubi [A] (verified)	945
Maple [B] (verified)	947
Fricas [B] (verification not implemented)	948
Sympy [F(-1)]	949
Maxima [B] (verification not implemented)	950
Giac [B] (verification not implemented)	951
Mupad [B] (verification not implemented)	952
Reduce [B] (verification not implemented)	952

Optimal result

Integrand size = 20, antiderivative size = 135

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{14}} dx = -\frac{(Bd - Ae)(a+bx)^{11}}{13e(bd - ae)(d+ex)^{13}} + \frac{(11bBd + 2Abe - 13aBe)(a+bx)^{11}}{156e(bd - ae)^2(d+ex)^{12}} + \frac{b(11bBd + 2Abe - 13aBe)(a+bx)^{11}}{1716e(bd - ae)^3(d+ex)^{11}}$$

output

```
-1/13*(-A*e+B*d)*(b*x+a)^11/e/(-a*e+b*d)/(e*x+d)^13+1/156*(2*A*b*e-13*B*a*
e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^2/(e*x+d)^12+1/1716*b*(2*A*b*e-13*B*a*
e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^3/(e*x+d)^11
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1433 vs. 2(135) = 270.

Time = 0.48 (sec) , antiderivative size = 1433, normalized size of antiderivative = 10.61

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{14}} dx = \text{Too large to display}$$

input `Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^14,x]`

output

```
-1/1716*(11*a^10*e^10*(12*A*e + B*(d + 13*e*x)) + 10*a^9*b*e^9*(11*A*e*(d
+ 13*e*x) + 2*B*(d^2 + 13*d*e*x + 78*e^2*x^2)) + 9*a^8*b^2*e^8*(10*A*e*(d^
2 + 13*d*e*x + 78*e^2*x^2) + 3*B*(d^3 + 13*d^2*e*x + 78*d*e^2*x^2 + 286*e^
3*x^3)) + 8*a^7*b^3*e^7*(9*A*e*(d^3 + 13*d^2*e*x + 78*d*e^2*x^2 + 286*e^3*
x^3) + 4*B*(d^4 + 13*d^3*e*x + 78*d^2*e^2*x^2 + 286*d*e^3*x^3 + 715*e^4*x^
4)) + 7*a^6*b^4*e^6*(8*A*e*(d^4 + 13*d^3*e*x + 78*d^2*e^2*x^2 + 286*d*e^3*
x^3 + 715*e^4*x^4) + 5*B*(d^5 + 13*d^4*e*x + 78*d^3*e^2*x^2 + 286*d^2*e^3*
x^3 + 715*d*e^4*x^4 + 1287*e^5*x^5)) + 6*a^5*b^5*e^5*(7*A*e*(d^5 + 13*d^4*
e*x + 78*d^3*e^2*x^2 + 286*d^2*e^3*x^3 + 715*d*e^4*x^4 + 1287*e^5*x^5) + 6
*B*(d^6 + 13*d^5*e*x + 78*d^4*e^2*x^2 + 286*d^3*e^3*x^3 + 715*d^2*e^4*x^4
+ 1287*d*e^5*x^5 + 1716*e^6*x^6)) + 5*a^4*b^6*e^4*(6*A*e*(d^6 + 13*d^5*e*x
+ 78*d^4*e^2*x^2 + 286*d^3*e^3*x^3 + 715*d^2*e^4*x^4 + 1287*d*e^5*x^5 + 1
716*e^6*x^6) + 7*B*(d^7 + 13*d^6*e*x + 78*d^5*e^2*x^2 + 286*d^4*e^3*x^3 +
715*d^3*e^4*x^4 + 1287*d^2*e^5*x^5 + 1716*d*e^6*x^6 + 1716*e^7*x^7)) + 4*a
^3*b^7*e^3*(5*A*e*(d^7 + 13*d^6*e*x + 78*d^5*e^2*x^2 + 286*d^4*e^3*x^3 + 7
15*d^3*e^4*x^4 + 1287*d^2*e^5*x^5 + 1716*d*e^6*x^6 + 1716*e^7*x^7) + 8*B*(
d^8 + 13*d^7*e*x + 78*d^6*e^2*x^2 + 286*d^5*e^3*x^3 + 715*d^4*e^4*x^4 + 12
87*d^3*e^5*x^5 + 1716*d^2*e^6*x^6 + 1716*d*e^7*x^7 + 1287*e^8*x^8)) + 3*a^
2*b^8*e^2*(4*A*e*(d^8 + 13*d^7*e*x + 78*d^6*e^2*x^2 + 286*d^5*e^3*x^3 + 71
5*d^4*e^4*x^4 + 1287*d^3*e^5*x^5 + 1716*d^2*e^6*x^6 + 1716*d*e^7*x^7 + ...
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.97, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {87, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{14}} dx$$

$$\downarrow 87$$

$$\frac{(-13aBe + 2Abe + 11bBd) \int \frac{(a+bx)^{10}}{(d+ex)^{13}} dx}{13e(bd - ae)} - \frac{(a + bx)^{11}(Bd - Ae)}{13e(d + ex)^{13}(bd - ae)}$$

$$\begin{aligned}
 & \downarrow 55 \\
 & \frac{(-13aBe + 2Abe + 11bBd) \left(\frac{b \int \frac{(a+bx)^{10}}{(d+ex)^{12}} dx}{12(bd-ae)} + \frac{(a+bx)^{11}}{12(d+ex)^{12}(bd-ae)} \right)}{13e(bd-ae)} - \frac{(a+bx)^{11}(Bd-Ae)}{13e(d+ex)^{13}(bd-ae)} \\
 & \downarrow 48 \\
 & \frac{\left(\frac{b(a+bx)^{11}}{132(d+ex)^{11}(bd-ae)^2} + \frac{(a+bx)^{11}}{12(d+ex)^{12}(bd-ae)} \right) (-13aBe + 2Abe + 11bBd)}{13e(bd-ae)} - \frac{(a+bx)^{11}(Bd-Ae)}{13e(d+ex)^{13}(bd-ae)}
 \end{aligned}$$

input `Int[((a + b*x)^10*(A + B*x))/(d + e*x)^14,x]`

output `-1/13*((B*d - A*e)*(a + b*x)^11)/(e*(b*d - a*e)*(d + e*x)^13) + ((11*b*B*d + 2*A*b*e - 13*a*B*e)*((a + b*x)^11/(12*(b*d - a*e)*(d + e*x)^12) + (b*(a + b*x)^11)/(132*(b*d - a*e)^2*(d + e*x)^11))/(13*e*(b*d - a*e))`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 55 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(Simplify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])`

rule 87

```

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p
_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && ( !LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))

```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1900 vs. $2(129) = 258$.

Time = 0.27 (sec) , antiderivative size = 1901, normalized size of antiderivative = 14.08

method	result	size
risch	Expression too large to display	1901
default	Expression too large to display	1942
norman	Expression too large to display	1992
gosper	Expression too large to display	2233
orering	Expression too large to display	2233
parallelsch	Expression too large to display	2240

input

```
int((b*x+a)^10*(B*x+A)/(e*x+d)^14,x,method=_RETURNVERBOSE)
```


output

```
(-1/2*b^10*B/e*x^11-1/6*b^9/e^2*(2*A*b*e+20*B*a*e+11*B*b*d)*x^10-5/12*b^8/
e^3*(6*A*a*b*e^2+2*A*b^2*d*e+27*B*a^2*e^2+20*B*a*b*d*e+11*B*b^2*d^2)*x^9-3
/4*b^7/e^4*(12*A*a^2*b*e^3+6*A*a*b^2*d*e^2+2*A*b^3*d^2*e+32*B*a^3*e^3+27*B
*a^2*b*d*e^2+20*B*a*b^2*d^2*e+11*B*b^3*d^3)*x^8-b^6/e^5*(20*A*a^3*b*e^4+12
*A*a^2*b^2*d*e^3+6*A*a*b^3*d^2*e^2+2*A*b^4*d^3*e+35*B*a^4*e^4+32*B*a^3*b*d
*e^3+27*B*a^2*b^2*d^2*e^2+20*B*a*b^3*d^3*e+11*B*b^4*d^4)*x^7-b^5/e^6*(30*A
*a^4*b*e^5+20*A*a^3*b^2*d*e^4+12*A*a^2*b^3*d^2*e^3+6*A*a*b^4*d^3*e^2+2*A*b
^5*d^4*e+36*B*a^5*e^5+35*B*a^4*b*d*e^4+32*B*a^3*b^2*d^2*e^3+27*B*a^2*b^3*d
^3*e^2+20*B*a*b^4*d^4*e+11*B*b^5*d^5)*x^6-3/4*b^4/e^7*(42*A*a^5*b*e^6+30*A
*a^4*b^2*d*e^5+20*A*a^3*b^3*d^2*e^4+12*A*a^2*b^4*d^3*e^3+6*A*a*b^5*d^4*e^2
+2*A*b^6*d^5*e+35*B*a^6*e^6+36*B*a^5*b*d*e^5+35*B*a^4*b^2*d^2*e^4+32*B*a^3
*b^3*d^3*e^3+27*B*a^2*b^4*d^4*e^2+20*B*a*b^5*d^5*e+11*B*b^6*d^6)*x^5-5/12*
b^3/e^8*(56*A*a^6*b*e^7+42*A*a^5*b^2*d*e^6+30*A*a^4*b^3*d^2*e^5+20*A*a^3*b
^4*d^3*e^4+12*A*a^2*b^5*d^4*e^3+6*A*a*b^6*d^5*e^2+2*A*b^7*d^6*e+32*B*a^7*e
^7+35*B*a^6*b*d*e^6+36*B*a^5*b^2*d^2*e^5+35*B*a^4*b^3*d^3*e^4+32*B*a^3*b^4
*d^4*e^3+27*B*a^2*b^5*d^5*e^2+20*B*a*b^6*d^6*e+11*B*b^7*d^7)*x^4-1/6*b^2/e
^9*(72*A*a^7*b*e^8+56*A*a^6*b^2*d*e^7+42*A*a^5*b^3*d^2*e^6+30*A*a^4*b^4*d
^3*e^5+20*A*a^3*b^5*d^4*e^4+12*A*a^2*b^6*d^5*e^3+6*A*a*b^7*d^6*e^2+2*A*b^8*
d^7*e+27*B*a^8*e^8+32*B*a^7*b*d*e^7+35*B*a^6*b^2*d^2*e^6+36*B*a^5*b^3*d^3*
e^5+35*B*a^4*b^4*d^4*e^4+32*B*a^3*b^5*d^5*e^3+27*B*a^2*b^6*d^6*e^2+20*B...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1951 vs. $2(129) = 258$.

Time = 0.13 (sec) , antiderivative size = 1951, normalized size of antiderivative = 14.45

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{14}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)/(e*x+d)^14,x, algorithm="fricas")
```

output

```
-1/1716*(858*B*b^10*e^11*x^11 + 11*B*b^10*d^11 + 132*A*a^10*e^11 + 2*(10*B
*a*b^9 + A*b^10)*d^10*e + 3*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 + 4*(8*B*a^3
*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 5*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e^4 + 6*(6
*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 7*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^5*e^6
+ 8*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 9*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^
3*e^8 + 10*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 + 11*(B*a^10 + 10*A*a^9*b)*d*
e^10 + 286*(11*B*b^10*d*e^10 + 2*(10*B*a*b^9 + A*b^10)*e^11)*x^10 + 715*(1
1*B*b^10*d^2*e^9 + 2*(10*B*a*b^9 + A*b^10)*d*e^10 + 3*(9*B*a^2*b^8 + 2*A*a
*b^9)*e^11)*x^9 + 1287*(11*B*b^10*d^3*e^8 + 2*(10*B*a*b^9 + A*b^10)*d^2*e^
9 + 3*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 + 4*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^1
1)*x^8 + 1716*(11*B*b^10*d^4*e^7 + 2*(10*B*a*b^9 + A*b^10)*d^3*e^8 + 3*(9*
B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 + 4*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 + 5*
(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7 + 1716*(11*B*b^10*d^5*e^6 + 2*(10*B*
a*b^9 + A*b^10)*d^4*e^7 + 3*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 + 4*(8*B*a^3
*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 5*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 + 6*(6*
B*a^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 + 1287*(11*B*b^10*d^6*e^5 + 2*(10*B*a*b
^9 + A*b^10)*d^5*e^6 + 3*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 + 4*(8*B*a^3*b^
7 + 3*A*a^2*b^8)*d^3*e^8 + 5*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 + 6*(6*B*
a^5*b^5 + 5*A*a^4*b^6)*d*e^10 + 7*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 +
715*(11*B*b^10*d^7*e^4 + 2*(10*B*a*b^9 + A*b^10)*d^6*e^5 + 3*(9*B*a^2*b...
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{14}} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**10*(B*x+A)/(e*x+d)**14,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1951 vs. $2(129) = 258$.

Time = 0.13 (sec) , antiderivative size = 1951, normalized size of antiderivative = 14.45

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{14}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^14,x, algorithm="maxima")`

output

```
-1/1716*(858*B*b^10*e^11*x^11 + 11*B*b^10*d^11 + 132*A*a^10*e^11 + 2*(10*B
*a*b^9 + A*b^10)*d^10*e + 3*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 + 4*(8*B*a^3
*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 5*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e^4 + 6*(6
*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 7*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^5*e^6
+ 8*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 9*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^
3*e^8 + 10*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 + 11*(B*a^10 + 10*A*a^9*b)*d*
e^10 + 286*(11*B*b^10*d*e^10 + 2*(10*B*a*b^9 + A*b^10)*e^11)*x^10 + 715*(1
1*B*b^10*d^2*e^9 + 2*(10*B*a*b^9 + A*b^10)*d*e^10 + 3*(9*B*a^2*b^8 + 2*A*
a*b^9)*e^11)*x^9 + 1287*(11*B*b^10*d^3*e^8 + 2*(10*B*a*b^9 + A*b^10)*d^2*e^
9 + 3*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 + 4*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^1
1)*x^8 + 1716*(11*B*b^10*d^4*e^7 + 2*(10*B*a*b^9 + A*b^10)*d^3*e^8 + 3*(9*
B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 + 4*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 + 5*
(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7 + 1716*(11*B*b^10*d^5*e^6 + 2*(10*B*
a*b^9 + A*b^10)*d^4*e^7 + 3*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 + 4*(8*B*a^3
*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 5*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 + 6*(6*
B*a^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 + 1287*(11*B*b^10*d^6*e^5 + 2*(10*B*a*b
^9 + A*b^10)*d^5*e^6 + 3*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 + 4*(8*B*a^3*b^
7 + 3*A*a^2*b^8)*d^3*e^8 + 5*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 + 6*(6*B*
a^5*b^5 + 5*A*a^4*b^6)*d*e^10 + 7*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 +
715*(11*B*b^10*d^7*e^4 + 2*(10*B*a*b^9 + A*b^10)*d^6*e^5 + 3*(9*B*a^2*b...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2232 vs. $2(129) = 258$.

Time = 0.13 (sec) , antiderivative size = 2232, normalized size of antiderivative = 16.53

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{14}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^14,x, algorithm="giac")`

output

```
-1/1716*(858*B*b^10*e^11*x^11 + 3146*B*b^10*d*e^10*x^10 + 5720*B*a*b^9*e^11*x^10 + 572*A*b^10*e^11*x^10 + 7865*B*b^10*d^2*e^9*x^9 + 14300*B*a*b^9*d*e^10*x^9 + 1430*A*b^10*d*e^10*x^9 + 19305*B*a^2*b^8*e^11*x^9 + 4290*A*a*b^9*e^11*x^9 + 14157*B*b^10*d^3*e^8*x^8 + 25740*B*a*b^9*d^2*e^9*x^8 + 2574*A*b^10*d^2*e^9*x^8 + 34749*B*a^2*b^8*d*e^10*x^8 + 7722*A*a*b^9*d*e^10*x^8 + 41184*B*a^3*b^7*e^11*x^8 + 15444*A*a^2*b^8*e^11*x^8 + 18876*B*b^10*d^4*e^7*x^7 + 34320*B*a*b^9*d^3*e^8*x^7 + 3432*A*b^10*d^3*e^8*x^7 + 46332*B*a^2*b^8*d^2*e^9*x^7 + 10296*A*a*b^9*d^2*e^9*x^7 + 54912*B*a^3*b^7*d*e^10*x^7 + 20592*A*a^2*b^8*d*e^10*x^7 + 60060*B*a^4*b^6*e^11*x^7 + 34320*A*a^3*b^7*e^11*x^7 + 18876*B*b^10*d^5*e^6*x^6 + 34320*B*a*b^9*d^4*e^7*x^6 + 3432*A*b^10*d^4*e^7*x^6 + 46332*B*a^2*b^8*d^3*e^8*x^6 + 10296*A*a*b^9*d^3*e^8*x^6 + 54912*B*a^3*b^7*d^2*e^9*x^6 + 20592*A*a^2*b^8*d^2*e^9*x^6 + 60060*B*a^4*b^6*d*e^10*x^6 + 34320*A*a^3*b^7*d*e^10*x^6 + 61776*B*a^5*b^5*e^11*x^6 + 51480*A*a^4*b^6*e^11*x^6 + 14157*B*b^10*d^6*e^5*x^5 + 25740*B*a*b^9*d^5*e^6*x^5 + 2574*A*b^10*d^5*e^6*x^5 + 34749*B*a^2*b^8*d^4*e^7*x^5 + 7722*A*a*b^9*d^4*e^7*x^5 + 41184*B*a^3*b^7*d^3*e^8*x^5 + 15444*A*a^2*b^8*d^3*e^8*x^5 + 45045*B*a^4*b^6*d^2*e^9*x^5 + 25740*A*a^3*b^7*d^2*e^9*x^5 + 46332*B*a^5*b^5*d*e^10*x^5 + 38610*A*a^4*b^6*d*e^10*x^5 + 45045*B*a^6*b^4*e^11*x^5 + 54054*A*a^5*b^5*e^11*x^5 + 7865*B*b^10*d^7*e^4*x^4 + 14300*B*a*b^9*d^6*e^5*x^4 + 1430*A*b^10*d^6*e^5*x^4 + 19305*B*a^2*b^8*d^5*e^6*x^4 + 4290*A*a*b...
```

Mupad [B] (verification not implemented)

Time = 1.54 (sec) , antiderivative size = 2031, normalized size of antiderivative = 15.04

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{14}} dx = \text{Too large to display}$$

input `int(((A + B*x)*(a + b*x)^10)/(d + e*x)^14,x)`

output
$$\begin{aligned} & -((132*A*a^{10}*e^{11} + 11*B*b^{10}*d^{11} + 2*A*b^{10}*d^{10}*e + 11*B*a^{10}*d*e^{10} + \\ & 6*A*a*b^9*d^9*e^2 + 20*B*a^9*b*d^2*e^9 + 12*A*a^2*b^8*d^8*e^3 + 20*A*a^3* \\ & b^7*d^7*e^4 + 30*A*a^4*b^6*d^6*e^5 + 42*A*a^5*b^5*d^5*e^6 + 56*A*a^6*b^4*d \\ & ^4*e^7 + 72*A*a^7*b^3*d^3*e^8 + 90*A*a^8*b^2*d^2*e^9 + 27*B*a^2*b^8*d^9*e^ \\ & 2 + 32*B*a^3*b^7*d^8*e^3 + 35*B*a^4*b^6*d^7*e^4 + 36*B*a^5*b^5*d^6*e^5 + 3 \\ & 5*B*a^6*b^4*d^5*e^6 + 32*B*a^7*b^3*d^4*e^7 + 27*B*a^8*b^2*d^3*e^8 + 110*A* \\ & a^9*b*d*e^{10} + 20*B*a*b^9*d^{10}*e)/(1716*e^{12}) + (x*(11*B*a^{10}*e^{10} + 11*B* \\ & b^{10}*d^{10} + 110*A*a^9*b*e^{10} + 2*A*b^{10}*d^9*e + 6*A*a*b^9*d^8*e^2 + 90*A*a \\ & ^8*b^2*d*e^9 + 12*A*a^2*b^8*d^7*e^3 + 20*A*a^3*b^7*d^6*e^4 + 30*A*a^4*b^6* \\ & d^5*e^5 + 42*A*a^5*b^5*d^4*e^6 + 56*A*a^6*b^4*d^3*e^7 + 72*A*a^7*b^3*d^2*e \\ & ^8 + 27*B*a^2*b^8*d^8*e^2 + 32*B*a^3*b^7*d^7*e^3 + 35*B*a^4*b^6*d^6*e^4 + \\ & 36*B*a^5*b^5*d^5*e^5 + 35*B*a^6*b^4*d^4*e^6 + 32*B*a^7*b^3*d^3*e^7 + 27*B* \\ & a^8*b^2*d^2*e^8 + 20*B*a*b^9*d^9*e + 20*B*a^9*b*d*e^9))/(132*e^{11}) + (3*b^ \\ & 7*x^8*(32*B*a^3*e^3 + 11*B*b^3*d^3 + 12*A*a^2*b*e^3 + 2*A*b^3*d^2*e + 6*A* \\ & a*b^2*d*e^2 + 20*B*a*b^2*d^2*e + 27*B*a^2*b*d*e^2))/(4*e^4) + (3*b^4*x^5*(\\ & 35*B*a^6*e^6 + 11*B*b^6*d^6 + 42*A*a^5*b*e^6 + 2*A*b^6*d^5*e + 6*A*a*b^5*d \\ & ^4*e^2 + 30*A*a^4*b^2*d*e^5 + 12*A*a^2*b^4*d^3*e^3 + 20*A*a^3*b^3*d^2*e^4 \\ & + 27*B*a^2*b^4*d^4*e^2 + 32*B*a^3*b^3*d^3*e^3 + 35*B*a^4*b^2*d^2*e^4 + 20* \\ & B*a*b^5*d^5*e + 36*B*a^5*b*d*e^5))/(4*e^7) + (b^9*x^{10}*(2*A*b*e + 20*B*a*e \\ & + 11*B*b*d))/(6*e^2) + (b^6*x^7*(35*B*a^4*e^4 + 11*B*b^4*d^4 + 20*A*a^...$$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 1283, normalized size of antiderivative = 9.50

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{14}} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^14,x)`

output

```
( - 12*a**11*e**11 - 11*a**10*b*d*e**10 - 143*a**10*b*e**11*x - 10*a**9*b*
*2*d**2*e**9 - 130*a**9*b**2*d*e**10*x - 780*a**9*b**2*e**11*x**2 - 9*a**8
*b**3*d**3*e**8 - 117*a**8*b**3*d**2*e**9*x - 702*a**8*b**3*d*e**10*x**2 -
2574*a**8*b**3*e**11*x**3 - 8*a**7*b**4*d**4*e**7 - 104*a**7*b**4*d**3*e*
*8*x - 624*a**7*b**4*d**2*e**9*x**2 - 2288*a**7*b**4*d*e**10*x**3 - 5720*a
**7*b**4*e**11*x**4 - 7*a**6*b**5*d**5*e**6 - 91*a**6*b**5*d**4*e**7*x - 5
46*a**6*b**5*d**3*e**8*x**2 - 2002*a**6*b**5*d**2*e**9*x**3 - 5005*a**6*b*
*5*d*e**10*x**4 - 9009*a**6*b**5*e**11*x**5 - 6*a**5*b**6*d**6*e**5 - 78*a
**5*b**6*d**5*e**6*x - 468*a**5*b**6*d**4*e**7*x**2 - 1716*a**5*b**6*d**3*
e**8*x**3 - 4290*a**5*b**6*d**2*e**9*x**4 - 7722*a**5*b**6*d*e**10*x**5 -
10296*a**5*b**6*e**11*x**6 - 5*a**4*b**7*d**7*e**4 - 65*a**4*b**7*d**6*e**
5*x - 390*a**4*b**7*d**5*e**6*x**2 - 1430*a**4*b**7*d**4*e**7*x**3 - 3575*
a**4*b**7*d**3*e**8*x**4 - 6435*a**4*b**7*d**2*e**9*x**5 - 8580*a**4*b**7*
d*e**10*x**6 - 8580*a**4*b**7*e**11*x**7 - 4*a**3*b**8*d**8*e**3 - 52*a**3
*b**8*d**7*e**4*x - 312*a**3*b**8*d**6*e**5*x**2 - 1144*a**3*b**8*d**5*e**
6*x**3 - 2860*a**3*b**8*d**4*e**7*x**4 - 5148*a**3*b**8*d**3*e**8*x**5 - 6
864*a**3*b**8*d**2*e**9*x**6 - 6864*a**3*b**8*d*e**10*x**7 - 5148*a**3*b**
8*e**11*x**8 - 3*a**2*b**9*d**9*e**2 - 39*a**2*b**9*d**8*e**3*x - 234*a**2
*b**9*d**7*e**4*x**2 - 858*a**2*b**9*d**6*e**5*x**3 - 2145*a**2*b**9*d**5*
e**6*x**4 - 3861*a**2*b**9*d**4*e**7*x**5 - 5148*a**2*b**9*d**3*e**8*x**...
```

3.93 $\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{15}} dx$

Optimal result	954
Mathematica [B] (verified)	955
Rubi [A] (verified)	956
Maple [B] (verified)	958
Fricas [B] (verification not implemented)	959
Sympy [F(-1)]	960
Maxima [B] (verification not implemented)	960
Giac [B] (verification not implemented)	961
Mupad [B] (verification not implemented)	962
Reduce [B] (verification not implemented)	963

Optimal result

Integrand size = 20, antiderivative size = 185

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{15}} dx = -\frac{(Bd - Ae)(a+bx)^{11}}{14e(bd - ae)(d+ex)^{14}} + \frac{(11bBd + 3Abe - 14aBe)(a+bx)^{11}}{182e(bd - ae)^2(d+ex)^{13}} + \frac{b(11bBd + 3Abe - 14aBe)(a+bx)^{11}}{1092e(bd - ae)^3(d+ex)^{12}} + \frac{b^2(11bBd + 3Abe - 14aBe)(a+bx)^{11}}{12012e(bd - ae)^4(d+ex)^{11}}$$

output

```
-1/14*(-A*e+B*d)*(b*x+a)^11/e/(-a*e+b*d)/(e*x+d)^14+1/182*(3*A*b*e-14*B*a*
e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^2/(e*x+d)^13+1/1092*b*(3*A*b*e-14*B*a*
e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^3/(e*x+d)^12+1/12012*b^2*(3*A*b*e-14*B
*a*e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^4/(e*x+d)^11
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1430 vs. $2(185) = 370$.

Time = 0.55 (sec) , antiderivative size = 1430, normalized size of antiderivative = 7.73

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{15}} dx = \text{Too large to display}$$

input `Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^15,x]`

output

```
-1/12012*(66*a^10*e^10*(13*A*e + B*(d + 14*e*x)) + 110*a^9*b*e^9*(6*A*e*(d
+ 14*e*x) + B*(d^2 + 14*d*e*x + 91*e^2*x^2)) + 45*a^8*b^2*e^8*(11*A*e*(d^
2 + 14*d*e*x + 91*e^2*x^2) + 3*B*(d^3 + 14*d^2*e*x + 91*d*e^2*x^2 + 364*e^
3*x^3)) + 72*a^7*b^3*e^7*(5*A*e*(d^3 + 14*d^2*e*x + 91*d*e^2*x^2 + 364*e^3
*x^3) + 2*B*(d^4 + 14*d^3*e*x + 91*d^2*e^2*x^2 + 364*d*e^3*x^3 + 1001*e^4*
x^4)) + 28*a^6*b^4*e^6*(9*A*e*(d^4 + 14*d^3*e*x + 91*d^2*e^2*x^2 + 364*d*e
^3*x^3 + 1001*e^4*x^4) + 5*B*(d^5 + 14*d^4*e*x + 91*d^3*e^2*x^2 + 364*d^2*
e^3*x^3 + 1001*d*e^4*x^4 + 2002*e^5*x^5)) + 42*a^5*b^5*e^5*(4*A*e*(d^5 + 1
4*d^4*e*x + 91*d^3*e^2*x^2 + 364*d^2*e^3*x^3 + 1001*d*e^4*x^4 + 2002*e^5*x
^5) + 3*B*(d^6 + 14*d^5*e*x + 91*d^4*e^2*x^2 + 364*d^3*e^3*x^3 + 1001*d^2*
e^4*x^4 + 2002*d*e^5*x^5 + 3003*e^6*x^6)) + 105*a^4*b^6*e^4*(A*e*(d^6 + 14
*d^5*e*x + 91*d^4*e^2*x^2 + 364*d^3*e^3*x^3 + 1001*d^2*e^4*x^4 + 2002*d*e^
5*x^5 + 3003*e^6*x^6) + B*(d^7 + 14*d^6*e*x + 91*d^5*e^2*x^2 + 364*d^4*e^3
*x^3 + 1001*d^3*e^4*x^4 + 2002*d^2*e^5*x^5 + 3003*d*e^6*x^6 + 3432*e^7*x^7
)) + 20*a^3*b^7*e^3*(3*A*e*(d^7 + 14*d^6*e*x + 91*d^5*e^2*x^2 + 364*d^4*e^
3*x^3 + 1001*d^3*e^4*x^4 + 2002*d^2*e^5*x^5 + 3003*d*e^6*x^6 + 3432*e^7*x^
7) + 4*B*(d^8 + 14*d^7*e*x + 91*d^6*e^2*x^2 + 364*d^5*e^3*x^3 + 1001*d^4*e
^4*x^4 + 2002*d^3*e^5*x^5 + 3003*d^2*e^6*x^6 + 3432*d*e^7*x^7 + 3003*e^8*x
^8)) + 6*a^2*b^8*e^2*(5*A*e*(d^8 + 14*d^7*e*x + 91*d^6*e^2*x^2 + 364*d^5*
e^3*x^3 + 1001*d^4*e^4*x^4 + 2002*d^3*e^5*x^5 + 3003*d^2*e^6*x^6 + 3432*...
```


Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.95, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {87, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{15}} dx \\
 & \quad \downarrow 87 \\
 & \frac{(-14aBe + 3Abe + 11bBd)}{14e(bd - ae)} \int \frac{(a+bx)^{10}}{(d+ex)^{14}} dx - \frac{(a+bx)^{11}(Bd - Ae)}{14e(d+ex)^{14}(bd - ae)} \\
 & \quad \downarrow 55 \\
 & \frac{(-14aBe + 3Abe + 11bBd)}{14e(bd - ae)} \left(\frac{2b \int \frac{(a+bx)^{10}}{(d+ex)^{13}} dx}{13(bd - ae)} + \frac{(a+bx)^{11}}{13(d+ex)^{13}(bd - ae)} \right) - \frac{(a+bx)^{11}(Bd - Ae)}{14e(d+ex)^{14}(bd - ae)} \\
 & \quad \downarrow 55 \\
 & \frac{(-14aBe + 3Abe + 11bBd)}{14e(bd - ae)} \left(\frac{2b \left(\frac{b \int \frac{(a+bx)^{10}}{(d+ex)^{12}} dx}{12(bd - ae)} + \frac{(a+bx)^{11}}{12(d+ex)^{12}(bd - ae)} \right)}{13(bd - ae)} + \frac{(a+bx)^{11}}{13(d+ex)^{13}(bd - ae)} \right) - \frac{(a+bx)^{11}(Bd - Ae)}{14e(d+ex)^{14}(bd - ae)} \\
 & \quad \downarrow 48 \\
 & \frac{\left(\frac{(a+bx)^{11}}{13(d+ex)^{13}(bd - ae)} + \frac{2b \left(\frac{b(a+bx)^{11}}{132(d+ex)^{11}(bd - ae)^2} + \frac{(a+bx)^{11}}{12(d+ex)^{12}(bd - ae)} \right)}{13(bd - ae)} \right) (-14aBe + 3Abe + 11bBd)}{14e(bd - ae)} - \frac{(a+bx)^{11}(Bd - Ae)}{14e(d+ex)^{14}(bd - ae)}
 \end{aligned}$$

input

```
Int[((a + b*x)^10*(A + B*x))/(d + e*x)^15,x]
```

output

$$-1/14*((B*d - A*e)*(a + b*x)^{11}/(e*(b*d - a*e)*(d + e*x)^{14}) + ((11*b*B*d + 3*A*b*e - 14*a*B*e)*((a + b*x)^{11}/(13*(b*d - a*e)*(d + e*x)^{13}) + (2*b*((a + b*x)^{11}/(12*(b*d - a*e)*(d + e*x)^{12}) + (b*(a + b*x)^{11})/(132*(b*d - a*e)^2*(d + e*x)^{11}))/((13*(b*d - a*e))))/(14*e*(b*d - a*e))$$

Defintions of rubi rules used

rule 48

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

rule 55

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
lerQ[n, 1])
```

rule 87

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1900 vs. $2(177) = 354$.

Time = 0.28 (sec) , antiderivative size = 1901, normalized size of antiderivative = 10.28

method	result	size
risch	Expression too large to display	1901
default	Expression too large to display	1942
norman	Expression too large to display	2014
gosper	Expression too large to display	2233
orering	Expression too large to display	2233
parallelrisc	Expression too large to display	2242

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^15,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & (-1/3*b^{10}*B/e*x^{11}-1/12*b^9/e^2*(3*A*b*e+30*B*a*e+11*B*b*d)*x^{10}-1/6*b^8/ \\ & e^3*(12*A*a*b*e^2+3*A*b^2*d*e+54*B*a^2*e^2+30*B*a*b*d*e+11*B*b^2*d^2)*x^9- \\ & 1/4*b^7/e^4*(30*A*a^2*b*e^3+12*A*a*b^2*d*e^2+3*A*b^3*d^2*e+80*B*a^3*e^3+54 \\ & *B*a^2*b*d*e^2+30*B*a*b^2*d^2*e+11*B*b^3*d^3)*x^8-2/7*b^6/e^5*(60*A*a^3*b* \\ & e^4+30*A*a^2*b^2*d*e^3+12*A*a*b^3*d^2*e^2+3*A*b^4*d^3*e+105*B*a^4*e^4+80*B \\ & *a^3*b*d*e^3+54*B*a^2*b^2*d^2*e^2+30*B*a*b^3*d^3*e+11*B*b^4*d^4)*x^7-1/4*b \\ & ^5/e^6*(105*A*a^4*b*e^5+60*A*a^3*b^2*d*e^4+30*A*a^2*b^3*d^2*e^3+12*A*a*b^4 \\ & *d^3*e^2+3*A*b^5*d^4*e+126*B*a^5*e^5+105*B*a^4*b*d*e^4+80*B*a^3*b^2*d^2*e^ \\ & 3+54*B*a^2*b^3*d^3*e^2+30*B*a*b^4*d^4*e+11*B*b^5*d^5)*x^6-1/6*b^4/e^7*(168 \\ & *A*a^5*b*e^6+105*A*a^4*b^2*d*e^5+60*A*a^3*b^3*d^2*e^4+30*A*a^2*b^4*d^3*e^3 \\ & +12*A*a*b^5*d^4*e^2+3*A*b^6*d^5*e+140*B*a^6*e^6+126*B*a^5*b*d*e^5+105*B*a^ \\ & 4*b^2*d^2*e^4+80*B*a^3*b^3*d^3*e^3+54*B*a^2*b^4*d^4*e^2+30*B*a*b^5*d^5*e+1 \\ & 1*B*b^6*d^6)*x^5-1/12/e^8*b^3*(252*A*a^6*b*e^7+168*A*a^5*b^2*d*e^6+105*A*a^ \\ & ^4*b^3*d^2*e^5+60*A*a^3*b^4*d^3*e^4+30*A*a^2*b^5*d^4*e^3+12*A*a*b^6*d^5*e^ \\ & 2+3*A*b^7*d^6*e+144*B*a^7*e^7+140*B*a^6*b*d*e^6+126*B*a^5*b^2*d^2*e^5+105* \\ & B*a^4*b^3*d^3*e^4+80*B*a^3*b^4*d^4*e^3+54*B*a^2*b^5*d^5*e^2+30*B*a*b^6*d^6 \\ & *e+11*B*b^7*d^7)*x^4-1/33*b^2/e^9*(360*A*a^7*b*e^8+252*A*a^6*b^2*d*e^7+168 \\ & *A*a^5*b^3*d^2*e^6+105*A*a^4*b^4*d^3*e^5+60*A*a^3*b^5*d^4*e^4+30*A*a^2*b^6 \\ & *d^5*e^3+12*A*a*b^7*d^6*e^2+3*A*b^8*d^7*e+135*B*a^8*e^8+144*B*a^7*b*d*e^7+ \\ & 140*B*a^6*b^2*d^2*e^6+126*B*a^5*b^3*d^3*e^5+105*B*a^4*b^4*d^4*e^4+80*B*... \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1962 vs. $2(177) = 354$.

Time = 0.20 (sec) , antiderivative size = 1962, normalized size of antiderivative = 10.61

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{15}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)/(e*x+d)^15,x, algorithm="fricas")
```

output

```
-1/12012*(4004*B*b^10*e^11*x^11 + 11*B*b^10*d^11 + 858*A*a^10*e^11 + 3*(10
*B*a*b^9 + A*b^10)*d^10*e + 6*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 + 10*(8*B*
a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 15*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e^4 +
21*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 28*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^
5*e^6 + 36*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 45*(3*B*a^8*b^2 + 8*A*a^7
*b^3)*d^3*e^8 + 55*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 + 66*(B*a^10 + 10*A*a
^9*b)*d*e^10 + 1001*(11*B*b^10*d*e^10 + 3*(10*B*a*b^9 + A*b^10)*e^11)*x^10
+ 2002*(11*B*b^10*d^2*e^9 + 3*(10*B*a*b^9 + A*b^10)*d*e^10 + 6*(9*B*a^2*b
^8 + 2*A*a*b^9)*e^11)*x^9 + 3003*(11*B*b^10*d^3*e^8 + 3*(10*B*a*b^9 + A*b^
10)*d^2*e^9 + 6*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 + 10*(8*B*a^3*b^7 + 3*A*a
^2*b^8)*e^11)*x^8 + 3432*(11*B*b^10*d^4*e^7 + 3*(10*B*a*b^9 + A*b^10)*d^3*
e^8 + 6*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 + 10*(8*B*a^3*b^7 + 3*A*a^2*b^8)
*d*e^10 + 15*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7 + 3003*(11*B*b^10*d^5*e
^6 + 3*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 6*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8
+ 10*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 15*(7*B*a^4*b^6 + 4*A*a^3*b^7)
*d*e^10 + 21*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 + 2002*(11*B*b^10*d^6*e
^5 + 3*(10*B*a*b^9 + A*b^10)*d^5*e^6 + 6*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7
+ 10*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 + 15*(7*B*a^4*b^6 + 4*A*a^3*b^7)
*d^2*e^9 + 21*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10 + 28*(5*B*a^6*b^4 + 6*A*a
^5*b^5)*e^11)*x^5 + 1001*(11*B*b^10*d^7*e^4 + 3*(10*B*a*b^9 + A*b^10)*d...
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{15}} dx = \text{Timed out}$$

input `integrate((b*x+a)**10*(B*x+A)/(e*x+d)**15,x)`output `Timed out`**Maxima [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 1962 vs. $2(177) = 354$.

Time = 0.16 (sec) , antiderivative size = 1962, normalized size of antiderivative = 10.61

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{15}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^15,x, algorithm="maxima")`

output

```

-1/12012*(4004*B*b^10*e^11*x^11 + 11*B*b^10*d^11 + 858*A*a^10*e^11 + 3*(10
*B*a*b^9 + A*b^10)*d^10*e + 6*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 + 10*(8*B*
a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 15*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e^4 +
21*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 28*(5*B*a^6*b^4 + 6*A*a^5*b^5)*d^
5*e^6 + 36*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 45*(3*B*a^8*b^2 + 8*A*a^7
*b^3)*d^3*e^8 + 55*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 + 66*(B*a^10 + 10*A*a
^9*b)*d*e^10 + 1001*(11*B*b^10*d*e^10 + 3*(10*B*a*b^9 + A*b^10)*e^11)*x^10
+ 2002*(11*B*b^10*d^2*e^9 + 3*(10*B*a*b^9 + A*b^10)*d*e^10 + 6*(9*B*a^2*b
^8 + 2*A*a*b^9)*e^11)*x^9 + 3003*(11*B*b^10*d^3*e^8 + 3*(10*B*a*b^9 + A*b^
10)*d^2*e^9 + 6*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 + 10*(8*B*a^3*b^7 + 3*A*a
^2*b^8)*e^11)*x^8 + 3432*(11*B*b^10*d^4*e^7 + 3*(10*B*a*b^9 + A*b^10)*d^3*
e^8 + 6*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 + 10*(8*B*a^3*b^7 + 3*A*a^2*b^8)
*d*e^10 + 15*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7 + 3003*(11*B*b^10*d^5*e
^6 + 3*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 6*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8
+ 10*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 15*(7*B*a^4*b^6 + 4*A*a^3*b^7)
*d*e^10 + 21*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 + 2002*(11*B*b^10*d^6*e
^5 + 3*(10*B*a*b^9 + A*b^10)*d^5*e^6 + 6*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7
+ 10*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 + 15*(7*B*a^4*b^6 + 4*A*a^3*b^7)
*d^2*e^9 + 21*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10 + 28*(5*B*a^6*b^4 + 6*A*a
^5*b^5)*e^11)*x^5 + 1001*(11*B*b^10*d^7*e^4 + 3*(10*B*a*b^9 + A*b^10)*d...

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2232 vs. $2(177) = 354$.

Time = 0.13 (sec) , antiderivative size = 2232, normalized size of antiderivative = 12.06

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{15}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)/(e*x+d)^15,x, algorithm="giac")
```

output

```
-1/12012*(4004*B*b^10*e^11*x^11 + 11011*B*b^10*d*e^10*x^10 + 30030*B*a*b^9
*e^11*x^10 + 3003*A*b^10*e^11*x^10 + 22022*B*b^10*d^2*e^9*x^9 + 60060*B*a*
b^9*d*e^10*x^9 + 6006*A*b^10*d*e^10*x^9 + 108108*B*a^2*b^8*e^11*x^9 + 2402
4*A*a*b^9*e^11*x^9 + 33033*B*b^10*d^3*e^8*x^8 + 90090*B*a*b^9*d^2*e^9*x^8
+ 9009*A*b^10*d^2*e^9*x^8 + 162162*B*a^2*b^8*d*e^10*x^8 + 36036*A*a*b^9*d*
e^10*x^8 + 240240*B*a^3*b^7*e^11*x^8 + 90090*A*a^2*b^8*e^11*x^8 + 37752*B*
b^10*d^4*e^7*x^7 + 102960*B*a*b^9*d^3*e^8*x^7 + 10296*A*b^10*d^3*e^8*x^7 +
185328*B*a^2*b^8*d^2*e^9*x^7 + 41184*A*a*b^9*d^2*e^9*x^7 + 274560*B*a^3*b
^7*d*e^10*x^7 + 102960*A*a^2*b^8*d*e^10*x^7 + 360360*B*a^4*b^6*e^11*x^7 +
205920*A*a^3*b^7*e^11*x^7 + 33033*B*b^10*d^5*e^6*x^6 + 90090*B*a*b^9*d^4*e
^7*x^6 + 9009*A*b^10*d^4*e^7*x^6 + 162162*B*a^2*b^8*d^3*e^8*x^6 + 36036*A*
a*b^9*d^3*e^8*x^6 + 240240*B*a^3*b^7*d^2*e^9*x^6 + 90090*A*a^2*b^8*d^2*e^9
*x^6 + 315315*B*a^4*b^6*d*e^10*x^6 + 180180*A*a^3*b^7*d*e^10*x^6 + 378378*
B*a^5*b^5*e^11*x^6 + 315315*A*a^4*b^6*e^11*x^6 + 22022*B*b^10*d^6*e^5*x^5
+ 60060*B*a*b^9*d^5*e^6*x^5 + 6006*A*b^10*d^5*e^6*x^5 + 108108*B*a^2*b^8*d
^4*e^7*x^5 + 24024*A*a*b^9*d^4*e^7*x^5 + 160160*B*a^3*b^7*d^3*e^8*x^5 + 60
060*A*a^2*b^8*d^3*e^8*x^5 + 210210*B*a^4*b^6*d^2*e^9*x^5 + 120120*A*a^3*b
^7*d^2*e^9*x^5 + 252252*B*a^5*b^5*d*e^10*x^5 + 210210*A*a^4*b^6*d*e^10*x^5
+ 280280*B*a^6*b^4*e^11*x^5 + 336336*A*a^5*b^5*e^11*x^5 + 11011*B*b^10*d^7
*e^4*x^4 + 30030*B*a*b^9*d^6*e^5*x^4 + 3003*A*b^10*d^6*e^5*x^4 + 54054*...
```

Mupad [B] (verification not implemented)

Time = 1.78 (sec) , antiderivative size = 2044, normalized size of antiderivative = 11.05

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{15}} dx = \text{Too large to display}$$

input

```
int(((A + B*x)*(a + b*x)^10)/(d + e*x)^15,x)
```

output

```

-((858*A*a^10*e^11 + 11*B*b^10*d^11 + 3*A*b^10*d^10*e + 66*B*a^10*d*e^10 +
  12*A*a*b^9*d^9*e^2 + 110*B*a^9*b*d^2*e^9 + 30*A*a^2*b^8*d^8*e^3 + 60*A*a^
  3*b^7*d^7*e^4 + 105*A*a^4*b^6*d^6*e^5 + 168*A*a^5*b^5*d^5*e^6 + 252*A*a^6*
  b^4*d^4*e^7 + 360*A*a^7*b^3*d^3*e^8 + 495*A*a^8*b^2*d^2*e^9 + 54*B*a^2*b^8
  *d^9*e^2 + 80*B*a^3*b^7*d^8*e^3 + 105*B*a^4*b^6*d^7*e^4 + 126*B*a^5*b^5*d^
  6*e^5 + 140*B*a^6*b^4*d^5*e^6 + 144*B*a^7*b^3*d^4*e^7 + 135*B*a^8*b^2*d^3*
  e^8 + 660*A*a^9*b*d*e^10 + 30*B*a*b^9*d^10*e)/(12012*e^12) + (x*(66*B*a^10
  *e^10 + 11*B*b^10*d^10 + 660*A*a^9*b*e^10 + 3*A*b^10*d^9*e + 12*A*a*b^9*d^
  8*e^2 + 495*A*a^8*b^2*d*e^9 + 30*A*a^2*b^8*d^7*e^3 + 60*A*a^3*b^7*d^6*e^4
  + 105*A*a^4*b^6*d^5*e^5 + 168*A*a^5*b^5*d^4*e^6 + 252*A*a^6*b^4*d^3*e^7 +
  360*A*a^7*b^3*d^2*e^8 + 54*B*a^2*b^8*d^8*e^2 + 80*B*a^3*b^7*d^7*e^3 + 105*
  B*a^4*b^6*d^6*e^4 + 126*B*a^5*b^5*d^5*e^5 + 140*B*a^6*b^4*d^4*e^6 + 144*B*
  a^7*b^3*d^3*e^7 + 135*B*a^8*b^2*d^2*e^8 + 30*B*a*b^9*d^9*e + 110*B*a^9*b*d
  *e^9))/(858*e^11) + (b^7*x^8*(80*B*a^3*e^3 + 11*B*b^3*d^3 + 30*A*a^2*b*e^3
  + 3*A*b^3*d^2*e + 12*A*a*b^2*d*e^2 + 30*B*a*b^2*d^2*e + 54*B*a^2*b*d*e^2)
  )/(4*e^4) + (b^4*x^5*(140*B*a^6*e^6 + 11*B*b^6*d^6 + 168*A*a^5*b*e^6 + 3*A
  *b^6*d^5*e + 12*A*a*b^5*d^4*e^2 + 105*A*a^4*b^2*d*e^5 + 30*A*a^2*b^4*d^3*e
  ^3 + 60*A*a^3*b^3*d^2*e^4 + 54*B*a^2*b^4*d^4*e^2 + 80*B*a^3*b^3*d^3*e^3 +
  105*B*a^4*b^2*d^2*e^4 + 30*B*a*b^5*d^5*e + 126*B*a^5*b*d*e^5))/(6*e^7) + (
  b^9*x^10*(3*A*b*e + 30*B*a*e + 11*B*b*d))/(12*e^2) + (2*b^6*x^7*(105*B*...

```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 1294, normalized size of antiderivative = 6.99

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{15}} dx = \text{Too large to display}$$

input

```
int((b*x+a)^10*(B*x+A)/(e*x+d)^15,x)
```


output

```
( - 78*a**11*e**11 - 66*a**10*b*d*e**10 - 924*a**10*b*e**11*x - 55*a**9*b*
*2*d**2*e**9 - 770*a**9*b**2*d*e**10*x - 5005*a**9*b**2*e**11*x**2 - 45*a*
*8*b**3*d**3*e**8 - 630*a**8*b**3*d**2*e**9*x - 4095*a**8*b**3*d*e**10*x**
2 - 16380*a**8*b**3*e**11*x**3 - 36*a**7*b**4*d**4*e**7 - 504*a**7*b**4*d*
*3*e**8*x - 3276*a**7*b**4*d**2*e**9*x**2 - 13104*a**7*b**4*d*e**10*x**3 -
36036*a**7*b**4*e**11*x**4 - 28*a**6*b**5*d**5*e**6 - 392*a**6*b**5*d**4*
e**7*x - 2548*a**6*b**5*d**3*e**8*x**2 - 10192*a**6*b**5*d**2*e**9*x**3 -
28028*a**6*b**5*d*e**10*x**4 - 56056*a**6*b**5*e**11*x**5 - 21*a**5*b**6*d
**6*e**5 - 294*a**5*b**6*d**5*e**6*x - 1911*a**5*b**6*d**4*e**7*x**2 - 764
4*a**5*b**6*d**3*e**8*x**3 - 21021*a**5*b**6*d**2*e**9*x**4 - 42042*a**5*b
**6*d*e**10*x**5 - 63063*a**5*b**6*e**11*x**6 - 15*a**4*b**7*d**7*e**4 - 2
10*a**4*b**7*d**6*e**5*x - 1365*a**4*b**7*d**5*e**6*x**2 - 5460*a**4*b**7*
d**4*e**7*x**3 - 15015*a**4*b**7*d**3*e**8*x**4 - 30030*a**4*b**7*d**2*e**
9*x**5 - 45045*a**4*b**7*d*e**10*x**6 - 51480*a**4*b**7*e**11*x**7 - 10*a*
*3*b**8*d**8*e**3 - 140*a**3*b**8*d**7*e**4*x - 910*a**3*b**8*d**6*e**5*x*
*2 - 3640*a**3*b**8*d**5*e**6*x**3 - 10010*a**3*b**8*d**4*e**7*x**4 - 2002
0*a**3*b**8*d**3*e**8*x**5 - 30030*a**3*b**8*d**2*e**9*x**6 - 34320*a**3*b
**8*d*e**10*x**7 - 30030*a**3*b**8*e**11*x**8 - 6*a**2*b**9*d**9*e**2 - 84
*a**2*b**9*d**8*e**3*x - 546*a**2*b**9*d**7*e**4*x**2 - 2184*a**2*b**9*d**
6*e**5*x**3 - 6006*a**2*b**9*d**5*e**6*x**4 - 12012*a**2*b**9*d**4*e**7...
```

3.94 $\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{16}} dx$

Optimal result	965
Mathematica [B] (verified)	966
Rubi [A] (verified)	967
Maple [B] (verified)	969
Fricas [B] (verification not implemented)	970
Sympy [F(-1)]	971
Maxima [B] (verification not implemented)	972
Giac [B] (verification not implemented)	973
Mupad [B] (verification not implemented)	974
Reduce [B] (verification not implemented)	974

Optimal result

Integrand size = 20, antiderivative size = 235

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{16}} dx = -\frac{(Bd - Ae)(a+bx)^{11}}{15e(bd - ae)(d+ex)^{15}} + \frac{(11bBd + 4Abe - 15aBe)(a+bx)^{11}}{210e(bd - ae)^2(d+ex)^{14}} + \frac{b(11bBd + 4Abe - 15aBe)(a+bx)^{11}}{910e(bd - ae)^3(d+ex)^{13}} + \frac{b^2(11bBd + 4Abe - 15aBe)(a+bx)^{11}}{5460e(bd - ae)^4(d+ex)^{12}} + \frac{b^3(11bBd + 4Abe - 15aBe)(a+bx)^{11}}{60060e(bd - ae)^5(d+ex)^{11}}$$

output

```
-1/15*(-A*e+B*d)*(b*x+a)^11/e/(-a*e+b*d)/(e*x+d)^15+1/210*(4*A*b*e-15*B*a*
e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^2/(e*x+d)^14+1/910*b*(4*A*b*e-15*B*a*
e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^3/(e*x+d)^13+1/5460*b^2*(4*A*b*e-15*B*a*
e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^4/(e*x+d)^12+1/60060*b^3*(4*A*b*e-15*
B*a*e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^5/(e*x+d)^11
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1430 vs. $2(235) = 470$.

Time = 0.56 (sec) , antiderivative size = 1430, normalized size of antiderivative = 6.09

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{16}} dx = \text{Too large to display}$$

input `Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^16,x]`

output

```
-1/60060*(286*a^10*e^10*(14*A*e + B*(d + 15*e*x)) + 220*a^9*b*e^9*(13*A*e*(d + 15*e*x) + 2*B*(d^2 + 15*d*e*x + 105*e^2*x^2)) + 495*a^8*b^2*e^8*(4*A*e*(d^2 + 15*d*e*x + 105*e^2*x^2) + B*(d^3 + 15*d^2*e*x + 105*d*e^2*x^2 + 455*e^3*x^3)) + 120*a^7*b^3*e^7*(11*A*e*(d^3 + 15*d^2*e*x + 105*d*e^2*x^2 + 455*e^3*x^3) + 4*B*(d^4 + 15*d^3*e*x + 105*d^2*e^2*x^2 + 455*d*e^3*x^3 + 1365*e^4*x^4)) + 420*a^6*b^4*e^6*(2*A*e*(d^4 + 15*d^3*e*x + 105*d^2*e^2*x^2 + 455*d*e^3*x^3 + 1365*e^4*x^4) + B*(d^5 + 15*d^4*e*x + 105*d^3*e^2*x^2 + 455*d^2*e^3*x^3 + 1365*d*e^4*x^4 + 3003*e^5*x^5)) + 168*a^5*b^5*e^5*(3*A*e*(d^5 + 15*d^4*e*x + 105*d^3*e^2*x^2 + 455*d^2*e^3*x^3 + 1365*d*e^4*x^4 + 3003*e^5*x^5) + 2*B*(d^6 + 15*d^5*e*x + 105*d^4*e^2*x^2 + 455*d^3*e^3*x^3 + 1365*d^2*e^4*x^4 + 3003*d*e^5*x^5 + 5005*e^6*x^6)) + 35*a^4*b^6*e^4*(8*A*e*(d^6 + 15*d^5*e*x + 105*d^4*e^2*x^2 + 455*d^3*e^3*x^3 + 1365*d^2*e^4*x^4 + 3003*d*e^5*x^5 + 5005*e^6*x^6) + 7*B*(d^7 + 15*d^6*e*x + 105*d^5*e^2*x^2 + 455*d^4*e^3*x^3 + 1365*d^3*e^4*x^4 + 3003*d^2*e^5*x^5 + 5005*d*e^6*x^6 + 6435*e^7*x^7)) + 20*a^3*b^7*e^3*(7*A*e*(d^7 + 15*d^6*e*x + 105*d^5*e^2*x^2 + 455*d^4*e^3*x^3 + 1365*d^3*e^4*x^4 + 3003*d^2*e^5*x^5 + 5005*d*e^6*x^6 + 6435*e^7*x^7) + 8*B*(d^8 + 15*d^7*e*x + 105*d^6*e^2*x^2 + 455*d^5*e^3*x^3 + 1365*d^4*e^4*x^4 + 3003*d^3*e^5*x^5 + 5005*d^2*e^6*x^6 + 6435*d*e^7*x^7 + 6435*e^8*x^8)) + 30*a^2*b^8*e^2*(2*A*e*(d^8 + 15*d^7*e*x + 105*d^6*e^2*x^2 + 455*d^5*e^3*x^3 + 1365*d^4*e^4*x^4 + 3003*d^3*e^5*x^5 + 50...
```

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 219, normalized size of antiderivative = 0.93, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {87, 55, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{16}} dx \\
 & \quad \downarrow 87 \\
 & \frac{(-15aBe + 4Abe + 11bBd) \int \frac{(a+bx)^{10}}{(d+ex)^{15}} dx}{15e(bd-ae)} - \frac{(a+bx)^{11}(Bd-Ae)}{15e(d+ex)^{15}(bd-ae)} \\
 & \quad \downarrow 55 \\
 & \frac{(-15aBe + 4Abe + 11bBd) \left(\frac{3b \int \frac{(a+bx)^{10}}{(d+ex)^{14}} dx}{14(bd-ae)} + \frac{(a+bx)^{11}}{14(d+ex)^{14}(bd-ae)} \right)}{15e(bd-ae)} - \frac{(a+bx)^{11}(Bd-Ae)}{15e(d+ex)^{15}(bd-ae)} \\
 & \quad \downarrow 55 \\
 & \frac{(-15aBe + 4Abe + 11bBd) \left(\frac{3b \left(\frac{2b \int \frac{(a+bx)^{10}}{(d+ex)^{13}} dx}{13(bd-ae)} + \frac{(a+bx)^{11}}{13(d+ex)^{13}(bd-ae)} \right)}{14(bd-ae)} + \frac{(a+bx)^{11}}{14(d+ex)^{14}(bd-ae)} \right)}{15e(bd-ae)} - \frac{(a+bx)^{11}(Bd-Ae)}{15e(d+ex)^{15}(bd-ae)} \\
 & \quad \downarrow 55 \\
 & \frac{(a+bx)^{11}(Bd-Ae)}{15e(d+ex)^{15}(bd-ae)}
 \end{aligned}$$

$$(-15aBe + 4Abe + 11bBd) \left(\frac{3b \left(\frac{b \int \frac{(a+bx)^{10}}{(d+ex)^{12}} dx}{12(bd-ae)} + \frac{(a+bx)^{11}}{12(d+ex)^{12}(bd-ae)} \right) + \frac{(a+bx)^{11}}{13(d+ex)^{13}(bd-ae)}}{14(bd-ae)} + \frac{(a+bx)^{11}}{14(d+ex)^{14}(bd-ae)} \right)$$

$$\frac{15e(bd - ae) (a + bx)^{11} (Bd - Ae)}{15e(d + ex)^{15} (bd - ae)}$$

↓ 48

$$\left(\frac{(a+bx)^{11}}{14(d+ex)^{14}(bd-ae)} + \frac{3b \left(\frac{(a+bx)^{11}}{13(d+ex)^{13}(bd-ae)} + \frac{2b \left(\frac{b(a+bx)^{11}}{132(d+ex)^{11}(bd-ae)^2} + \frac{(a+bx)^{11}}{12(d+ex)^{12}(bd-ae)} \right)}{13(bd-ae)} \right)}{14(bd-ae)} \right) (-15aBe + 4Abe + 11bBd)$$

$$\frac{15e(bd - ae) (a + bx)^{11} (Bd - Ae)}{15e(d + ex)^{15} (bd - ae)}$$

input `Int[((a + b*x)^10*(A + B*x))/(d + e*x)^16,x]`

output `-1/15*((B*d - A*e)*(a + b*x)^11)/(e*(b*d - a*e)*(d + e*x)^15) + ((11*b*B*d + 4*A*b*e - 15*a*B*e)*((a + b*x)^11/(14*(b*d - a*e)*(d + e*x)^14) + (3*b*((a + b*x)^11/(13*(b*d - a*e)*(d + e*x)^13) + (2*b*((a + b*x)^11/(12*(b*d - a*e)*(d + e*x)^12) + (b*(a + b*x)^11)/(132*(b*d - a*e)^2*(d + e*x)^11)))/(13*(b*d - a*e)))/(14*(b*d - a*e)))/(15*e*(b*d - a*e))`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
 [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
 a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 55 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
 (a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
 implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
 c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
 c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
 lerQ[n, 1])`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
 + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
 + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
 /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
 rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1900 vs. 2(225) = 450.

Time = 0.30 (sec) , antiderivative size = 1901, normalized size of antiderivative = 8.09

method	result	size
risch	Expression too large to display	1901
default	Expression too large to display	1942
norman	Expression too large to display	2014
gospers	Expression too large to display	2233
orering	Expression too large to display	2233
parallelrisch	Expression too large to display	2242

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^16,x,method=_RETURNVERBOSE)`

output

```
(-1/60060/e^12*(4004*A*a^10*e^11+2860*A*a^9*b*d*e^10+1980*A*a^8*b^2*d^2*e^
9+1320*A*a^7*b^3*d^3*e^8+840*A*a^6*b^4*d^4*e^7+504*A*a^5*b^5*d^5*e^6+280*A
*a^4*b^6*d^6*e^5+140*A*a^3*b^7*d^7*e^4+60*A*a^2*b^8*d^8*e^3+20*A*a*b^9*d^9
*e^2+4*A*b^10*d^10*e+286*B*a^10*d*e^10+440*B*a^9*b*d^2*e^9+495*B*a^8*b^2*d
^3*e^8+480*B*a^7*b^3*d^4*e^7+420*B*a^6*b^4*d^5*e^6+336*B*a^5*b^5*d^6*e^5+2
45*B*a^4*b^6*d^7*e^4+160*B*a^3*b^7*d^8*e^3+90*B*a^2*b^8*d^9*e^2+40*B*a*b^9
*d^10*e+11*B*b^10*d^11)-1/4004/e^11*(2860*A*a^9*b*e^10+1980*A*a^8*b^2*d*e^
9+1320*A*a^7*b^3*d^2*e^8+840*A*a^6*b^4*d^3*e^7+504*A*a^5*b^5*d^4*e^6+280*A
*a^4*b^6*d^5*e^5+140*A*a^3*b^7*d^6*e^4+60*A*a^2*b^8*d^7*e^3+20*A*a*b^9*d^8
*e^2+4*A*b^10*d^9*e+286*B*a^10*e^10+440*B*a^9*b*d*e^9+495*B*a^8*b^2*d^2*e^
8+480*B*a^7*b^3*d^3*e^7+420*B*a^6*b^4*d^4*e^6+336*B*a^5*b^5*d^5*e^5+245*B*
a^4*b^6*d^6*e^4+160*B*a^3*b^7*d^7*e^3+90*B*a^2*b^8*d^8*e^2+40*B*a*b^9*d^9*
e+11*B*b^10*d^10)*x-1/572*b/e^10*(1980*A*a^8*b*e^9+1320*A*a^7*b^2*d*e^8+84
0*A*a^6*b^3*d^2*e^7+504*A*a^5*b^4*d^3*e^6+280*A*a^4*b^5*d^4*e^5+140*A*a^3*
b^6*d^5*e^4+60*A*a^2*b^7*d^6*e^3+20*A*a*b^8*d^7*e^2+4*A*b^9*d^8*e+440*B*a^
9*e^9+495*B*a^8*b*d*e^8+480*B*a^7*b^2*d^2*e^7+420*B*a^6*b^3*d^3*e^6+336*B*
a^5*b^4*d^4*e^5+245*B*a^4*b^5*d^5*e^4+160*B*a^3*b^6*d^6*e^3+90*B*a^2*b^7*d
^7*e^2+40*B*a*b^8*d^8*e+11*B*b^9*d^9)*x^2-1/132*b^2/e^9*(1320*A*a^7*b*e^8+
840*A*a^6*b^2*d*e^7+504*A*a^5*b^3*d^2*e^6+280*A*a^4*b^4*d^3*e^5+140*A*a^3*
b^5*d^4*e^4+60*A*a^2*b^6*d^5*e^3+20*A*a*b^7*d^6*e^2+4*A*b^8*d^7*e+495*B...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1973 vs. $2(225) = 450$.

Time = 0.16 (sec) , antiderivative size = 1973, normalized size of antiderivative = 8.40

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{16}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)/(e*x+d)^16,x, algorithm="fricas")
```

output

```

-1/60060*(15015*B*b^10*e^11*x^11 + 11*B*b^10*d^11 + 4004*A*a^10*e^11 + 4*(
10*B*a*b^9 + A*b^10)*d^10*e + 10*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 + 20*(8
*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 35*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e^4
+ 56*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 84*(5*B*a^6*b^4 + 6*A*a^5*b^5)
*d^5*e^6 + 120*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 165*(3*B*a^8*b^2 + 8*
A*a^7*b^3)*d^3*e^8 + 220*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 + 286*(B*a^10 +
10*A*a^9*b)*d*e^10 + 3003*(11*B*b^10*d*e^10 + 4*(10*B*a*b^9 + A*b^10)*e^1
1)*x^10 + 5005*(11*B*b^10*d^2*e^9 + 4*(10*B*a*b^9 + A*b^10)*d*e^10 + 10*(9
*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 + 6435*(11*B*b^10*d^3*e^8 + 4*(10*B*a*b^
9 + A*b^10)*d^2*e^9 + 10*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 + 20*(8*B*a^3*b^
7 + 3*A*a^2*b^8)*e^11)*x^8 + 6435*(11*B*b^10*d^4*e^7 + 4*(10*B*a*b^9 + A*b
^10)*d^3*e^8 + 10*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 + 20*(8*B*a^3*b^7 + 3*
A*a^2*b^8)*d*e^10 + 35*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7 + 5005*(11*B*
b^10*d^5*e^6 + 4*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 10*(9*B*a^2*b^8 + 2*A*a*b
^9)*d^3*e^8 + 20*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 35*(7*B*a^4*b^6 + 4
*A*a^3*b^7)*d*e^10 + 56*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 + 3003*(11*B
*b^10*d^6*e^5 + 4*(10*B*a*b^9 + A*b^10)*d^5*e^6 + 10*(9*B*a^2*b^8 + 2*A*a*
b^9)*d^4*e^7 + 20*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 + 35*(7*B*a^4*b^6 +
4*A*a^3*b^7)*d^2*e^9 + 56*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10 + 84*(5*B*a^6
*b^4 + 6*A*a^5*b^5)*e^11)*x^5 + 1365*(11*B*b^10*d^7*e^4 + 4*(10*B*a*b^9...

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{16}} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**10*(B*x+A)/(e*x+d)**16,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1973 vs. $2(225) = 450$.

Time = 0.13 (sec) , antiderivative size = 1973, normalized size of antiderivative = 8.40

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{16}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^16,x, algorithm="maxima")`

output

```
-1/60060*(15015*B*b^10*e^11*x^11 + 11*B*b^10*d^11 + 4004*A*a^10*e^11 + 4*(
10*B*a*b^9 + A*b^10)*d^10*e + 10*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 + 20*(8
*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 35*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e^4
+ 56*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 84*(5*B*a^6*b^4 + 6*A*a^5*b^5)
*d^5*e^6 + 120*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 165*(3*B*a^8*b^2 + 8*
A*a^7*b^3)*d^3*e^8 + 220*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 + 286*(B*a^10 +
10*A*a^9*b)*d*e^10 + 3003*(11*B*b^10*d*e^10 + 4*(10*B*a*b^9 + A*b^10)*e^1
1)*x^10 + 5005*(11*B*b^10*d^2*e^9 + 4*(10*B*a*b^9 + A*b^10)*d*e^10 + 10*(9
*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 + 6435*(11*B*b^10*d^3*e^8 + 4*(10*B*a*b^
9 + A*b^10)*d^2*e^9 + 10*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 + 20*(8*B*a^3*b^
7 + 3*A*a^2*b^8)*e^11)*x^8 + 6435*(11*B*b^10*d^4*e^7 + 4*(10*B*a*b^9 + A*b
^10)*d^3*e^8 + 10*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 + 20*(8*B*a^3*b^7 + 3*
A*a^2*b^8)*d*e^10 + 35*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7 + 5005*(11*B*
b^10*d^5*e^6 + 4*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 10*(9*B*a^2*b^8 + 2*A*a*b
^9)*d^3*e^8 + 20*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 35*(7*B*a^4*b^6 + 4
*A*a^3*b^7)*d*e^10 + 56*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 + 3003*(11*B
*b^10*d^6*e^5 + 4*(10*B*a*b^9 + A*b^10)*d^5*e^6 + 10*(9*B*a^2*b^8 + 2*A*a*
b^9)*d^4*e^7 + 20*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 + 35*(7*B*a^4*b^6 +
4*A*a^3*b^7)*d^2*e^9 + 56*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10 + 84*(5*B*a^6
*b^4 + 6*A*a^5*b^5)*e^11)*x^5 + 1365*(11*B*b^10*d^7*e^4 + 4*(10*B*a*b^9...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2232 vs. $2(225) = 450$.

Time = 0.13 (sec) , antiderivative size = 2232, normalized size of antiderivative = 9.50

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{16}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^16,x, algorithm="giac")`

output

```
-1/60060*(15015*B*b^10*e^11*x^11 + 33033*B*b^10*d*e^10*x^10 + 120120*B*a*b^9*e^11*x^10 + 12012*A*b^10*e^11*x^10 + 55055*B*b^10*d^2*e^9*x^9 + 200200*B*a*b^9*d*e^10*x^9 + 20020*A*b^10*d*e^10*x^9 + 450450*B*a^2*b^8*e^11*x^9 + 100100*A*a*b^9*e^11*x^9 + 70785*B*b^10*d^3*e^8*x^8 + 257400*B*a*b^9*d^2*e^9*x^8 + 25740*A*b^10*d^2*e^9*x^8 + 579150*B*a^2*b^8*d*e^10*x^8 + 128700*A*a*b^9*d*e^10*x^8 + 1029600*B*a^3*b^7*e^11*x^8 + 386100*A*a^2*b^8*e^11*x^8 + 70785*B*b^10*d^4*e^7*x^7 + 257400*B*a*b^9*d^3*e^8*x^7 + 25740*A*b^10*d^3*e^8*x^7 + 579150*B*a^2*b^8*d^2*e^9*x^7 + 128700*A*a*b^9*d^2*e^9*x^7 + 1029600*B*a^3*b^7*d*e^10*x^7 + 386100*A*a^2*b^8*d*e^10*x^7 + 1576575*B*a^4*b^6*e^11*x^7 + 900900*A*a^3*b^7*e^11*x^7 + 55055*B*b^10*d^5*e^6*x^6 + 200200*B*a*b^9*d^4*e^7*x^6 + 20020*A*b^10*d^4*e^7*x^6 + 450450*B*a^2*b^8*d^3*e^8*x^6 + 100100*A*a*b^9*d^3*e^8*x^6 + 800800*B*a^3*b^7*d^2*e^9*x^6 + 300300*A*a^2*b^8*d^2*e^9*x^6 + 1226225*B*a^4*b^6*d*e^10*x^6 + 700700*A*a^3*b^7*d*e^10*x^6 + 1681680*B*a^5*b^5*e^11*x^6 + 1401400*A*a^4*b^6*e^11*x^6 + 33033*B*b^10*d^6*e^5*x^5 + 120120*B*a*b^9*d^5*e^6*x^5 + 12012*A*b^10*d^5*e^6*x^5 + 270270*B*a^2*b^8*d^4*e^7*x^5 + 60060*A*a*b^9*d^4*e^7*x^5 + 480480*B*a^3*b^7*d^3*e^8*x^5 + 180180*A*a^2*b^8*d^3*e^8*x^5 + 735735*B*a^4*b^6*d^2*e^9*x^5 + 420420*A*a^3*b^7*d^2*e^9*x^5 + 1009008*B*a^5*b^5*d*e^10*x^5 + 840840*A*a^4*b^6*d*e^10*x^5 + 1261260*B*a^6*b^4*e^11*x^5 + 1513512*A*a^5*b^5*e^11*x^5 + 15015*B*b^10*d^7*e^4*x^4 + 54600*B*a*b^9*d^6*e^5*x^4 + 5460*...
```

Mupad [B] (verification not implemented)

Time = 2.24 (sec) , antiderivative size = 2055, normalized size of antiderivative = 8.74

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{16}} dx = \text{Too large to display}$$

input `int((A + B*x)*(a + b*x)^10)/(d + e*x)^16,x`

output

```

-((4004*A*a^10*e^11 + 11*B*b^10*d^11 + 4*A*b^10*d^10*e + 286*B*a^10*d*e^10
+ 20*A*a*b^9*d^9*e^2 + 440*B*a^9*b*d^2*e^9 + 60*A*a^2*b^8*d^8*e^3 + 140*A
*a^3*b^7*d^7*e^4 + 280*A*a^4*b^6*d^6*e^5 + 504*A*a^5*b^5*d^5*e^6 + 840*A*a
^6*b^4*d^4*e^7 + 1320*A*a^7*b^3*d^3*e^8 + 1980*A*a^8*b^2*d^2*e^9 + 90*B*a^
2*b^8*d^9*e^2 + 160*B*a^3*b^7*d^8*e^3 + 245*B*a^4*b^6*d^7*e^4 + 336*B*a^5*
b^5*d^6*e^5 + 420*B*a^6*b^4*d^5*e^6 + 480*B*a^7*b^3*d^4*e^7 + 495*B*a^8*b^
2*d^3*e^8 + 2860*A*a^9*b*d*e^10 + 40*B*a*b^9*d^10*e)/(60060*e^12) + (x*(28
6*B*a^10*e^10 + 11*B*b^10*d^10 + 2860*A*a^9*b*e^10 + 4*A*b^10*d^9*e + 20*A
*a*b^9*d^8*e^2 + 1980*A*a^8*b^2*d*e^9 + 60*A*a^2*b^8*d^7*e^3 + 140*A*a^3*b
^7*d^6*e^4 + 280*A*a^4*b^6*d^5*e^5 + 504*A*a^5*b^5*d^4*e^6 + 840*A*a^6*b^4
*d^3*e^7 + 1320*A*a^7*b^3*d^2*e^8 + 90*B*a^2*b^8*d^8*e^2 + 160*B*a^3*b^7*d
^7*e^3 + 245*B*a^4*b^6*d^6*e^4 + 336*B*a^5*b^5*d^5*e^5 + 420*B*a^6*b^4*d^4
*e^6 + 480*B*a^7*b^3*d^3*e^7 + 495*B*a^8*b^2*d^2*e^8 + 40*B*a*b^9*d^9*e +
440*B*a^9*b*d*e^9))/(4004*e^11) + (3*b^7*x^8*(160*B*a^3*e^3 + 11*B*b^3*d^3
+ 60*A*a^2*b*e^3 + 4*A*b^3*d^2*e + 20*A*a*b^2*d*e^2 + 40*B*a*b^2*d^2*e +
90*B*a^2*b*d*e^2))/(28*e^4) + (b^4*x^5*(420*B*a^6*e^6 + 11*B*b^6*d^6 + 504
*A*a^5*b*e^6 + 4*A*b^6*d^5*e + 20*A*a*b^5*d^4*e^2 + 280*A*a^4*b^2*d*e^5 +
60*A*a^2*b^4*d^3*e^3 + 140*A*a^3*b^3*d^2*e^4 + 90*B*a^2*b^4*d^4*e^2 + 160*
B*a^3*b^3*d^3*e^3 + 245*B*a^4*b^2*d^2*e^4 + 40*B*a*b^5*d^5*e + 336*B*a^5*b
*d*e^5))/(20*e^7) + (b^9*x^10*(4*A*b*e + 40*B*a*e + 11*B*b*d))/(20*e^2)...

```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 1305, normalized size of antiderivative = 5.55

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{16}} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^16,x)`

output

```
( - 364*a**11*e**11 - 286*a**10*b*d*e**10 - 4290*a**10*b*e**11*x - 220*a**
9*b**2*d**2*e**9 - 3300*a**9*b**2*d*e**10*x - 23100*a**9*b**2*e**11*x**2 -
165*a**8*b**3*d**3*e**8 - 2475*a**8*b**3*d**2*e**9*x - 17325*a**8*b**3*d*
e**10*x**2 - 75075*a**8*b**3*e**11*x**3 - 120*a**7*b**4*d**4*e**7 - 1800*a
**7*b**4*d**3*e**8*x - 12600*a**7*b**4*d**2*e**9*x**2 - 54600*a**7*b**4*d*
e**10*x**3 - 163800*a**7*b**4*e**11*x**4 - 84*a**6*b**5*d**5*e**6 - 1260*a
**6*b**5*d**4*e**7*x - 8820*a**6*b**5*d**3*e**8*x**2 - 38220*a**6*b**5*d**
2*e**9*x**3 - 114660*a**6*b**5*d*e**10*x**4 - 252252*a**6*b**5*e**11*x**5
- 56*a**5*b**6*d**6*e**5 - 840*a**5*b**6*d**5*e**6*x - 5880*a**5*b**6*d**4
*e**7*x**2 - 25480*a**5*b**6*d**3*e**8*x**3 - 76440*a**5*b**6*d**2*e**9*x*
*4 - 168168*a**5*b**6*d*e**10*x**5 - 280280*a**5*b**6*e**11*x**6 - 35*a**4
*b**7*d**7*e**4 - 525*a**4*b**7*d**6*e**5*x - 3675*a**4*b**7*d**5*e**6*x**
2 - 15925*a**4*b**7*d**4*e**7*x**3 - 47775*a**4*b**7*d**3*e**8*x**4 - 1051
05*a**4*b**7*d**2*e**9*x**5 - 175175*a**4*b**7*d*e**10*x**6 - 225225*a**4*
b**7*e**11*x**7 - 20*a**3*b**8*d**8*e**3 - 300*a**3*b**8*d**7*e**4*x - 210
0*a**3*b**8*d**6*e**5*x**2 - 9100*a**3*b**8*d**5*e**6*x**3 - 27300*a**3*b*
*8*d**4*e**7*x**4 - 60060*a**3*b**8*d**3*e**8*x**5 - 100100*a**3*b**8*d**2
*e**9*x**6 - 128700*a**3*b**8*d*e**10*x**7 - 128700*a**3*b**8*e**11*x**8 -
10*a**2*b**9*d**9*e**2 - 150*a**2*b**9*d**8*e**3*x - 1050*a**2*b**9*d**7*
e**4*x**2 - 4550*a**2*b**9*d**6*e**5*x**3 - 13650*a**2*b**9*d**5*e**6*x...
```

3.95 $\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{17}} dx$

Optimal result	976
Mathematica [B] (verified)	977
Rubi [A] (verified)	978
Maple [B] (verified)	981
Fricas [B] (verification not implemented)	982
Sympy [F(-1)]	983
Maxima [B] (verification not implemented)	984
Giac [B] (verification not implemented)	985
Mupad [B] (verification not implemented)	986
Reduce [B] (verification not implemented)	986

Optimal result

Integrand size = 20, antiderivative size = 285

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{17}} dx = -\frac{(Bd - Ae)(a+bx)^{11}}{16e(bd - ae)(d+ex)^{16}} + \frac{(11bBd + 5Abe - 16aBe)(a+bx)^{11}}{240e(bd - ae)^2(d+ex)^{15}} + \frac{b(11bBd + 5Abe - 16aBe)(a+bx)^{11}}{840e(bd - ae)^3(d+ex)^{14}} + \frac{b^2(11bBd + 5Abe - 16aBe)(a+bx)^{11}}{3640e(bd - ae)^4(d+ex)^{13}} + \frac{b^3(11bBd + 5Abe - 16aBe)(a+bx)^{11}}{21840e(bd - ae)^5(d+ex)^{12}} + \frac{b^4(11bBd + 5Abe - 16aBe)(a+bx)^{11}}{240240e(bd - ae)^6(d+ex)^{11}}$$

output

```
-1/16*(-A*e+B*d)*(b*x+a)^11/e/(-a*e+b*d)/(e*x+d)^16+1/240*(5*A*b*e-16*B*a*
e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^2/(e*x+d)^15+1/840*b*(5*A*b*e-16*B*a*e
+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^3/(e*x+d)^14+1/3640*b^2*(5*A*b*e-16*B*a
*e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^4/(e*x+d)^13+1/21840*b^3*(5*A*b*e-16*
B*a*e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^5/(e*x+d)^12+1/240240*b^4*(5*A*b*e
-16*B*a*e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^6/(e*x+d)^11
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1429 vs. $2(285) = 570$.

Time = 0.46 (sec) , antiderivative size = 1429, normalized size of antiderivative = 5.01

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{17}} dx = \text{Too large to display}$$

input `Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^17,x]`

output

```
-1/240240*(1001*a^10*e^10*(15*A*e + B*(d + 16*e*x)) + 1430*a^9*b*e^9*(7*A*
e*(d + 16*e*x) + B*(d^2 + 16*d*e*x + 120*e^2*x^2)) + 495*a^8*b^2*e^8*(13*A
*e*(d^2 + 16*d*e*x + 120*e^2*x^2) + 3*B*(d^3 + 16*d^2*e*x + 120*d*e^2*x^2
+ 560*e^3*x^3)) + 1320*a^7*b^3*e^7*(3*A*e*(d^3 + 16*d^2*e*x + 120*d*e^2*x^
2 + 560*e^3*x^3) + B*(d^4 + 16*d^3*e*x + 120*d^2*e^2*x^2 + 560*d*e^3*x^3 +
1820*e^4*x^4)) + 210*a^6*b^4*e^6*(11*A*e*(d^4 + 16*d^3*e*x + 120*d^2*e^2*
x^2 + 560*d*e^3*x^3 + 1820*e^4*x^4) + 5*B*(d^5 + 16*d^4*e*x + 120*d^3*e^2*
x^2 + 560*d^2*e^3*x^3 + 1820*d*e^4*x^4 + 4368*e^5*x^5)) + 252*a^5*b^5*e^5*
(5*A*e*(d^5 + 16*d^4*e*x + 120*d^3*e^2*x^2 + 560*d^2*e^3*x^3 + 1820*d*e^4*
x^4 + 4368*e^5*x^5) + 3*B*(d^6 + 16*d^5*e*x + 120*d^4*e^2*x^2 + 560*d^3*e^
3*x^3 + 1820*d^2*e^4*x^4 + 4368*d*e^5*x^5 + 8008*e^6*x^6)) + 70*a^4*b^6*e^
4*(9*A*e*(d^6 + 16*d^5*e*x + 120*d^4*e^2*x^2 + 560*d^3*e^3*x^3 + 1820*d^2*
e^4*x^4 + 4368*d*e^5*x^5 + 8008*e^6*x^6) + 7*B*(d^7 + 16*d^6*e*x + 120*d^5
*e^2*x^2 + 560*d^4*e^3*x^3 + 1820*d^3*e^4*x^4 + 4368*d^2*e^5*x^5 + 8008*d*
e^6*x^6 + 11440*e^7*x^7)) + 280*a^3*b^7*e^3*(A*e*(d^7 + 16*d^6*e*x + 120*d
^5*e^2*x^2 + 560*d^4*e^3*x^3 + 1820*d^3*e^4*x^4 + 4368*d^2*e^5*x^5 + 8008*
d*e^6*x^6 + 11440*e^7*x^7) + B*(d^8 + 16*d^7*e*x + 120*d^6*e^2*x^2 + 560*d
^5*e^3*x^3 + 1820*d^4*e^4*x^4 + 4368*d^3*e^5*x^5 + 8008*d^2*e^6*x^6 + 1144
0*d*e^7*x^7 + 12870*e^8*x^8)) + 15*a^2*b^8*e^2*(7*A*e*(d^8 + 16*d^7*e*x +
120*d^6*e^2*x^2 + 560*d^5*e^3*x^3 + 1820*d^4*e^4*x^4 + 4368*d^3*e^5*x^5...
```

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 263, normalized size of antiderivative = 0.92, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {87, 55, 55, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{17}} dx \\
 & \quad \downarrow 87 \\
 & \frac{(-16aBe + 5Abe + 11bBd) \int \frac{(a+bx)^{10}}{(d+ex)^{16}} dx}{16e(bd-ae)} - \frac{(a+bx)^{11}(Bd-Ae)}{16e(d+ex)^{16}(bd-ae)} \\
 & \quad \downarrow 55 \\
 & \frac{(-16aBe + 5Abe + 11bBd) \left(\frac{4b \int \frac{(a+bx)^{10}}{(d+ex)^{15}} dx}{15(bd-ae)} + \frac{(a+bx)^{11}}{15(d+ex)^{15}(bd-ae)} \right)}{16e(bd-ae)} - \frac{(a+bx)^{11}(Bd-Ae)}{16e(d+ex)^{16}(bd-ae)} \\
 & \quad \downarrow 55 \\
 & \frac{(-16aBe + 5Abe + 11bBd) \left(\frac{4b \left(\frac{3b \int \frac{(a+bx)^{10}}{(d+ex)^{14}} dx}{14(bd-ae)} + \frac{(a+bx)^{11}}{14(d+ex)^{14}(bd-ae)} \right)}{15(bd-ae)} + \frac{(a+bx)^{11}}{15(d+ex)^{15}(bd-ae)} \right)}{16e(bd-ae)} - \frac{(a+bx)^{11}(Bd-Ae)}{16e(d+ex)^{16}(bd-ae)} \\
 & \quad \downarrow 55 \\
 & \frac{(a+bx)^{11}(Bd-Ae)}{16e(d+ex)^{16}(bd-ae)}
 \end{aligned}$$

$$(-16aBe + 5Abe + 11bBd) \left(\frac{4b \left(\frac{3b \left(\frac{2b \int \frac{(a+bx)^{10}}{(d+ex)^{13}} dx + \frac{(a+bx)^{11}}{13(d+ex)^{13}(bd-ae)} \right)}{14(bd-ae)} + \frac{(a+bx)^{11}}{14(d+ex)^{14}(bd-ae)} \right)}{15(bd-ae)} + \frac{(a+bx)^{11}}{15(d+ex)^{15}(bd-ae)} \right)$$

$$\frac{16e(bd - ae) (a + bx)^{11} (Bd - Ae)}{16e(d + ex)^{16} (bd - ae)}$$

↓ 55

$$(-16aBe + 5Abe + 11bBd) \left(\frac{4b \left(\frac{3b \left(\frac{b \int \frac{(a+bx)^{10}}{(d+ex)^{12}} dx + \frac{(a+bx)^{11}}{12(d+ex)^{12}(bd-ae)} \right)}{13(bd-ae)} + \frac{(a+bx)^{11}}{13(d+ex)^{13}(bd-ae)} \right)}{14(bd-ae)} + \frac{(a+bx)^{11}}{14(d+ex)^{14}(bd-ae)} \right) + \frac{(a+bx)^{11}}{15(d+ex)^{15}(bd-ae)}$$

$$\frac{16e(bd - ae) (a + bx)^{11} (Bd - Ae)}{16e(d + ex)^{16} (bd - ae)}$$

↓ 48

$$\left(\frac{(a+bx)^{11}}{15(d+ex)^{15}(bd-ae)} + \frac{4b \left(\frac{(a+bx)^{11}}{14(d+ex)^{14}(bd-ae)} + \frac{3b \left(\frac{(a+bx)^{11}}{13(d+ex)^{13}(bd-ae)} + \frac{2b \left(\frac{b(a+bx)^{11}}{132(d+ex)^{11}(bd-ae)^2} + \frac{(a+bx)^{11}}{12(d+ex)^{12}(bd-ae)} \right)}{13(bd-ae)} \right)}{14(bd-ae)} \right)}{15(bd-ae)} \right) (-16aBe + \dots)$$

$$\frac{16e(bd - ae)}{16e(d + ex)^{16}(bd - ae)} \frac{(a + bx)^{11}(Bd - Ae)}{16e(d + ex)^{16}(bd - ae)}$$

```
input Int[((a + b*x)^10*(A + B*x))/(d + e*x)^17,x]
```

```
output -1/16*((B*d - A*e)*(a + b*x)^11)/(e*(b*d - a*e)*(d + e*x)^16) + ((11*b*B*d + 5*A*b*e - 16*a*B*e)*((a + b*x)^11/(15*(b*d - a*e)*(d + e*x)^15) + (4*b*((a + b*x)^11/(14*(b*d - a*e)*(d + e*x)^14) + (3*b*((a + b*x)^11/(13*(b*d - a*e)*(d + e*x)^13) + (2*b*((a + b*x)^11/(12*(b*d - a*e)*(d + e*x)^12) + (b*(a + b*x)^11)/(132*(b*d - a*e)^2*(d + e*x)^11)))/(13*(b*d - a*e))))/(14*(b*d - a*e)))/(15*(b*d - a*e)))/(16*e*(b*d - a*e))
```

Defintions of rubi rules used

```
rule 48 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 55 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(Simplify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])
```

rule 87

```

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p
_.), x_] :> Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && ( !LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))

```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1900 vs. $2(273) = 546$.

Time = 0.31 (sec) , antiderivative size = 1901, normalized size of antiderivative = 6.67

method	result	size
risch	Expression too large to display	1901
default	Expression too large to display	1942
norman	Expression too large to display	2014
gospers	Expression too large to display	2233
orering	Expression too large to display	2233
parallelsch	Expression too large to display	2242

input

```
int((b*x+a)^10*(B*x+A)/(e*x+d)^17,x,method=_RETURNVERBOSE)
```

output

```
(-1/240240/e^12*(15015*A*a^10*e^11+10010*A*a^9*b*d*e^10+6435*A*a^8*b^2*d^2
*e^9+3960*A*a^7*b^3*d^3*e^8+2310*A*a^6*b^4*d^4*e^7+1260*A*a^5*b^5*d^5*e^6+
630*A*a^4*b^6*d^6*e^5+280*A*a^3*b^7*d^7*e^4+105*A*a^2*b^8*d^8*e^3+30*A*a*b
^9*d^9*e^2+5*A*b^10*d^10*e+1001*B*a^10*d*e^10+1430*B*a^9*b*d^2*e^9+1485*B*
a^8*b^2*d^3*e^8+1320*B*a^7*b^3*d^4*e^7+1050*B*a^6*b^4*d^5*e^6+756*B*a^5*b^
5*d^6*e^5+490*B*a^4*b^6*d^7*e^4+280*B*a^3*b^7*d^8*e^3+135*B*a^2*b^8*d^9*e^
2+50*B*a*b^9*d^10*e+11*B*b^10*d^11)-1/15015/e^11*(10010*A*a^9*b*e^10+6435*
A*a^8*b^2*d*e^9+3960*A*a^7*b^3*d^2*e^8+2310*A*a^6*b^4*d^3*e^7+1260*A*a^5*b
^5*d^4*e^6+630*A*a^4*b^6*d^5*e^5+280*A*a^3*b^7*d^6*e^4+105*A*a^2*b^8*d^7*e
^3+30*A*a*b^9*d^8*e^2+5*A*b^10*d^9*e+1001*B*a^10*e^10+1430*B*a^9*b*d*e^9+1
485*B*a^8*b^2*d^2*e^8+1320*B*a^7*b^3*d^3*e^7+1050*B*a^6*b^4*d^4*e^6+756*B*
a^5*b^5*d^5*e^5+490*B*a^4*b^6*d^6*e^4+280*B*a^3*b^7*d^7*e^3+135*B*a^2*b^8*
d^8*e^2+50*B*a*b^9*d^9*e+11*B*b^10*d^10)*x-1/2002*b/e^10*(6435*A*a^8*b*e^9
+3960*A*a^7*b^2*d*e^8+2310*A*a^6*b^3*d^2*e^7+1260*A*a^5*b^4*d^3*e^6+630*A*
a^4*b^5*d^4*e^5+280*A*a^3*b^6*d^5*e^4+105*A*a^2*b^7*d^6*e^3+30*A*a*b^8*d^7
*e^2+5*A*b^9*d^8*e+1430*B*a^9*e^9+1485*B*a^8*b*d*e^8+1320*B*a^7*b^2*d^2*e^
7+1050*B*a^6*b^3*d^3*e^6+756*B*a^5*b^4*d^4*e^5+490*B*a^4*b^5*d^5*e^4+280*B
*a^3*b^6*d^6*e^3+135*B*a^2*b^7*d^7*e^2+50*B*a*b^8*d^8*e+11*B*b^9*d^9)*x^2-
1/429*b^2/e^9*(3960*A*a^7*b*e^8+2310*A*a^6*b^2*d*e^7+1260*A*a^5*b^3*d^2*e^
6+630*A*a^4*b^4*d^3*e^5+280*A*a^3*b^5*d^4*e^4+105*A*a^2*b^6*d^5*e^3+30*...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1984 vs. $2(273) = 546$.

Time = 0.18 (sec) , antiderivative size = 1984, normalized size of antiderivative = 6.96

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{17}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)/(e*x+d)^17,x, algorithm="fricas")
```

output

```

-1/240240*(48048*B*b^10*e^11*x^11 + 11*B*b^10*d^11 + 15015*A*a^10*e^11 + 5
*(10*B*a*b^9 + A*b^10)*d^10*e + 15*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 + 35*
(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 70*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e
^4 + 126*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 210*(5*B*a^6*b^4 + 6*A*a^5*
b^5)*d^5*e^6 + 330*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 495*(3*B*a^8*b^2
+ 8*A*a^7*b^3)*d^3*e^8 + 715*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 + 1001*(B*a
^10 + 10*A*a^9*b)*d*e^10 + 8008*(11*B*b^10*d*e^10 + 5*(10*B*a*b^9 + A*b^10
)*e^11)*x^10 + 11440*(11*B*b^10*d^2*e^9 + 5*(10*B*a*b^9 + A*b^10)*d*e^10 +
15*(9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 + 12870*(11*B*b^10*d^3*e^8 + 5*(10
*B*a*b^9 + A*b^10)*d^2*e^9 + 15*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 + 35*(8*B
*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 11440*(11*B*b^10*d^4*e^7 + 5*(10*B*a*b
^9 + A*b^10)*d^3*e^8 + 15*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 + 35*(8*B*a^3*
b^7 + 3*A*a^2*b^8)*d*e^10 + 70*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7 + 800
8*(11*B*b^10*d^5*e^6 + 5*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 15*(9*B*a^2*b^8 +
2*A*a*b^9)*d^3*e^8 + 35*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 70*(7*B*a^4
*b^6 + 4*A*a^3*b^7)*d*e^10 + 126*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 + 4
368*(11*B*b^10*d^6*e^5 + 5*(10*B*a*b^9 + A*b^10)*d^5*e^6 + 15*(9*B*a^2*b^8
+ 2*A*a*b^9)*d^4*e^7 + 35*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 + 70*(7*B*a
^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 + 126*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10 + 2
10*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 + 1820*(11*B*b^10*d^7*e^4 + 5*...

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{17}} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**10*(B*x+A)/(e*x+d)**17,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1984 vs. $2(273) = 546$.

Time = 0.14 (sec) , antiderivative size = 1984, normalized size of antiderivative = 6.96

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{17}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^17,x, algorithm="maxima")`

output

```
-1/240240*(48048*B*b^10*e^11*x^11 + 11*B*b^10*d^11 + 15015*A*a^10*e^11 + 5
*(10*B*a*b^9 + A*b^10)*d^10*e + 15*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 + 35*
(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 70*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7*e
^4 + 126*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 210*(5*B*a^6*b^4 + 6*A*a^5*
b^5)*d^5*e^6 + 330*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 495*(3*B*a^8*b^2
+ 8*A*a^7*b^3)*d^3*e^8 + 715*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 + 1001*(B*a
^10 + 10*A*a^9*b)*d*e^10 + 8008*(11*B*b^10*d*e^10 + 5*(10*B*a*b^9 + A*b^10
)*e^11)*x^10 + 11440*(11*B*b^10*d^2*e^9 + 5*(10*B*a*b^9 + A*b^10)*d*e^10 +
15*(9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 + 12870*(11*B*b^10*d^3*e^8 + 5*(10
*B*a*b^9 + A*b^10)*d^2*e^9 + 15*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 + 35*(8*B
*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 11440*(11*B*b^10*d^4*e^7 + 5*(10*B*a*b
^9 + A*b^10)*d^3*e^8 + 15*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 + 35*(8*B*a^3*
b^7 + 3*A*a^2*b^8)*d*e^10 + 70*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7 + 800
8*(11*B*b^10*d^5*e^6 + 5*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 15*(9*B*a^2*b^8 +
2*A*a*b^9)*d^3*e^8 + 35*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 70*(7*B*a^4
*b^6 + 4*A*a^3*b^7)*d*e^10 + 126*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 + 4
368*(11*B*b^10*d^6*e^5 + 5*(10*B*a*b^9 + A*b^10)*d^5*e^6 + 15*(9*B*a^2*b^8
+ 2*A*a*b^9)*d^4*e^7 + 35*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 + 70*(7*B*a
^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 + 126*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10 + 2
10*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 + 1820*(11*B*b^10*d^7*e^4 + 5*...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2232 vs. $2(273) = 546$.

Time = 0.13 (sec) , antiderivative size = 2232, normalized size of antiderivative = 7.83

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{17}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^17,x, algorithm="giac")`

output

```
-1/240240*(48048*B*b^10*e^11*x^11 + 88088*B*b^10*d*e^10*x^10 + 400400*B*a*
b^9*e^11*x^10 + 40040*A*b^10*e^11*x^10 + 125840*B*b^10*d^2*e^9*x^9 + 57200
0*B*a*b^9*d*e^10*x^9 + 57200*A*b^10*d*e^10*x^9 + 1544400*B*a^2*b^8*e^11*x^
9 + 343200*A*a*b^9*e^11*x^9 + 141570*B*b^10*d^3*e^8*x^8 + 643500*B*a*b^9*d
^2*e^9*x^8 + 64350*A*b^10*d^2*e^9*x^8 + 1737450*B*a^2*b^8*d*e^10*x^8 + 386
100*A*a*b^9*d*e^10*x^8 + 3603600*B*a^3*b^7*e^11*x^8 + 1351350*A*a^2*b^8*e^
11*x^8 + 125840*B*b^10*d^4*e^7*x^7 + 572000*B*a*b^9*d^3*e^8*x^7 + 57200*A*
b^10*d^3*e^8*x^7 + 1544400*B*a^2*b^8*d^2*e^9*x^7 + 343200*A*a*b^9*d^2*e^9*
x^7 + 3203200*B*a^3*b^7*d*e^10*x^7 + 1201200*A*a^2*b^8*d*e^10*x^7 + 560560
0*B*a^4*b^6*e^11*x^7 + 3203200*A*a^3*b^7*e^11*x^7 + 88088*B*b^10*d^5*e^6*x
^6 + 400400*B*a*b^9*d^4*e^7*x^6 + 40040*A*b^10*d^4*e^7*x^6 + 1081080*B*a^2
*b^8*d^3*e^8*x^6 + 240240*A*a*b^9*d^3*e^8*x^6 + 2242240*B*a^3*b^7*d^2*e^9*
x^6 + 840840*A*a^2*b^8*d^2*e^9*x^6 + 3923920*B*a^4*b^6*d*e^10*x^6 + 224224
0*A*a^3*b^7*d*e^10*x^6 + 6054048*B*a^5*b^5*e^11*x^6 + 5045040*A*a^4*b^6*e^
11*x^6 + 48048*B*b^10*d^6*e^5*x^5 + 218400*B*a*b^9*d^5*e^6*x^5 + 21840*A*b
^10*d^5*e^6*x^5 + 589680*B*a^2*b^8*d^4*e^7*x^5 + 131040*A*a*b^9*d^4*e^7*x^
5 + 1223040*B*a^3*b^7*d^3*e^8*x^5 + 458640*A*a^2*b^8*d^3*e^8*x^5 + 2140320
*B*a^4*b^6*d^2*e^9*x^5 + 1223040*A*a^3*b^7*d^2*e^9*x^5 + 3302208*B*a^5*b^5
*d*e^10*x^5 + 2751840*A*a^4*b^6*d*e^10*x^5 + 4586400*B*a^6*b^4*e^11*x^5 +
5503680*A*a^5*b^5*e^11*x^5 + 20020*B*b^10*d^7*e^4*x^4 + 91000*B*a*b^9*d...
```

Mupad [B] (verification not implemented)

Time = 1.32 (sec) , antiderivative size = 2066, normalized size of antiderivative = 7.25

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{17}} dx = \text{Too large to display}$$

input `int(((A + B*x)*(a + b*x)^10)/(d + e*x)^17,x)`

output

```

-((15015*A*a^10*e^11 + 11*B*b^10*d^11 + 5*A*b^10*d^10*e + 1001*B*a^10*d*e^
10 + 30*A*a*b^9*d^9*e^2 + 1430*B*a^9*b*d^2*e^9 + 105*A*a^2*b^8*d^8*e^3 + 2
80*A*a^3*b^7*d^7*e^4 + 630*A*a^4*b^6*d^6*e^5 + 1260*A*a^5*b^5*d^5*e^6 + 23
10*A*a^6*b^4*d^4*e^7 + 3960*A*a^7*b^3*d^3*e^8 + 6435*A*a^8*b^2*d^2*e^9 + 1
35*B*a^2*b^8*d^9*e^2 + 280*B*a^3*b^7*d^8*e^3 + 490*B*a^4*b^6*d^7*e^4 + 756
*B*a^5*b^5*d^6*e^5 + 1050*B*a^6*b^4*d^5*e^6 + 1320*B*a^7*b^3*d^4*e^7 + 148
5*B*a^8*b^2*d^3*e^8 + 10010*A*a^9*b*d*e^10 + 50*B*a*b^9*d^10*e)/(240240*e^
12) + (x*(1001*B*a^10*e^10 + 11*B*b^10*d^10 + 10010*A*a^9*b*e^10 + 5*A*b^1
0*d^9*e + 30*A*a*b^9*d^8*e^2 + 6435*A*a^8*b^2*d*e^9 + 105*A*a^2*b^8*d^7*e^
3 + 280*A*a^3*b^7*d^6*e^4 + 630*A*a^4*b^6*d^5*e^5 + 1260*A*a^5*b^5*d^4*e^6
+ 2310*A*a^6*b^4*d^3*e^7 + 3960*A*a^7*b^3*d^2*e^8 + 135*B*a^2*b^8*d^8*e^2
+ 280*B*a^3*b^7*d^7*e^3 + 490*B*a^4*b^6*d^6*e^4 + 756*B*a^5*b^5*d^5*e^5 +
1050*B*a^6*b^4*d^4*e^6 + 1320*B*a^7*b^3*d^3*e^7 + 1485*B*a^8*b^2*d^2*e^8
+ 50*B*a*b^9*d^9*e + 1430*B*a^9*b*d*e^9))/(15015*e^11) + (3*b^7*x^8*(280*B
*a^3*e^3 + 11*B*b^3*d^3 + 105*A*a^2*b*e^3 + 5*A*b^3*d^2*e + 30*A*a*b^2*d*e
^2 + 50*B*a*b^2*d^2*e + 135*B*a^2*b*d*e^2))/(56*e^4) + (b^4*x^5*(1050*B*a^
6*e^6 + 11*B*b^6*d^6 + 1260*A*a^5*b*e^6 + 5*A*b^6*d^5*e + 30*A*a*b^5*d^4*e
^2 + 630*A*a^4*b^2*d*e^5 + 105*A*a^2*b^4*d^3*e^3 + 280*A*a^3*b^3*d^2*e^4 +
135*B*a^2*b^4*d^4*e^2 + 280*B*a^3*b^3*d^3*e^3 + 490*B*a^4*b^2*d^2*e^4 + 5
0*B*a*b^5*d^5*e + 756*B*a^5*b*d*e^5))/(55*e^7) + (b^9*x^10*(5*A*b*e + 5...

```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 1316, normalized size of antiderivative = 4.62

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{17}} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^17,x)`

output

```
( - 1365*a**11*e**11 - 1001*a**10*b*d*e**10 - 16016*a**10*b*e**11*x - 715*
a**9*b**2*d**2*e**9 - 11440*a**9*b**2*d*e**10*x - 85800*a**9*b**2*e**11*x*
*2 - 495*a**8*b**3*d**3*e**8 - 7920*a**8*b**3*d**2*e**9*x - 59400*a**8*b**
3*d*e**10*x**2 - 277200*a**8*b**3*e**11*x**3 - 330*a**7*b**4*d**4*e**7 - 5
280*a**7*b**4*d**3*e**8*x - 39600*a**7*b**4*d**2*e**9*x**2 - 184800*a**7*b
**4*d*e**10*x**3 - 600600*a**7*b**4*e**11*x**4 - 210*a**6*b**5*d**5*e**6 -
3360*a**6*b**5*d**4*e**7*x - 25200*a**6*b**5*d**3*e**8*x**2 - 117600*a**6
*b**5*d**2*e**9*x**3 - 382200*a**6*b**5*d*e**10*x**4 - 917280*a**6*b**5*e
**11*x**5 - 126*a**5*b**6*d**6*e**5 - 2016*a**5*b**6*d**5*e**6*x - 15120*a
**5*b**6*d**4*e**7*x**2 - 70560*a**5*b**6*d**3*e**8*x**3 - 229320*a**5*b**6
*d**2*e**9*x**4 - 550368*a**5*b**6*d*e**10*x**5 - 1009008*a**5*b**6*e**11*
x**6 - 70*a**4*b**7*d**7*e**4 - 1120*a**4*b**7*d**6*e**5*x - 8400*a**4*b**
7*d**5*e**6*x**2 - 39200*a**4*b**7*d**4*e**7*x**3 - 127400*a**4*b**7*d**3*
e**8*x**4 - 305760*a**4*b**7*d**2*e**9*x**5 - 560560*a**4*b**7*d*e**10*x**
6 - 800800*a**4*b**7*e**11*x**7 - 35*a**3*b**8*d**8*e**3 - 560*a**3*b**8*d
**7*e**4*x - 4200*a**3*b**8*d**6*e**5*x**2 - 19600*a**3*b**8*d**5*e**6*x**
3 - 63700*a**3*b**8*d**4*e**7*x**4 - 152880*a**3*b**8*d**3*e**8*x**5 - 280
280*a**3*b**8*d**2*e**9*x**6 - 400400*a**3*b**8*d*e**10*x**7 - 450450*a**3
*b**8*e**11*x**8 - 15*a**2*b**9*d**9*e**2 - 240*a**2*b**9*d**8*e**3*x - 18
00*a**2*b**9*d**7*e**4*x**2 - 8400*a**2*b**9*d**6*e**5*x**3 - 27300*a**...
```


3.96 $\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{18}} dx$

Optimal result	988
Mathematica [B] (verified)	989
Rubi [A] (verified)	990
Maple [B] (verified)	995
Fricas [B] (verification not implemented)	996
Sympy [F(-1)]	997
Maxima [B] (verification not implemented)	998
Giac [B] (verification not implemented)	999
Mupad [B] (verification not implemented)	1000
Reduce [B] (verification not implemented)	1000

Optimal result

Integrand size = 20, antiderivative size = 335

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{18}} dx = -\frac{(Bd - Ae)(a+bx)^{11}}{17e(bd - ae)(d+ex)^{17}} + \frac{(11bBd + 6Abe - 17aBe)(a+bx)^{11}}{272e(bd - ae)^2(d+ex)^{16}} + \frac{b(11bBd + 6Abe - 17aBe)(a+bx)^{11}}{816e(bd - ae)^3(d+ex)^{15}} + \frac{b^2(11bBd + 6Abe - 17aBe)(a+bx)^{11}}{2856e(bd - ae)^4(d+ex)^{14}} + \frac{b^3(11bBd + 6Abe - 17aBe)(a+bx)^{11}}{12376e(bd - ae)^5(d+ex)^{13}} + \frac{b^4(11bBd + 6Abe - 17aBe)(a+bx)^{11}}{74256e(bd - ae)^6(d+ex)^{12}} + \frac{b^5(11bBd + 6Abe - 17aBe)(a+bx)^{11}}{816816e(bd - ae)^7(d+ex)^{11}}$$

output

```
-1/17*(-A*e+B*d)*(b*x+a)^11/e/(-a*e+b*d)/(e*x+d)^17+1/272*(6*A*b*e-17*B*a*
e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^2/(e*x+d)^16+1/816*b*(6*A*b*e-17*B*a*
e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^3/(e*x+d)^15+1/2856*b^2*(6*A*b*e-17*B*a*
e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^4/(e*x+d)^14+1/12376*b^3*(6*A*b*e-17*
B*a*e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^5/(e*x+d)^13+1/74256*b^4*(6*A*b*e-
17*B*a*e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^6/(e*x+d)^12+1/816816*b^5*(6*A*
b*e-17*B*a*e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^7/(e*x+d)^11
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1433 vs. $2(335) = 670$.

Time = 0.52 (sec) , antiderivative size = 1433, normalized size of antiderivative = 4.28

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{18}} dx = \text{Too large to display}$$

input

```
Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^18,x]
```

output

```

-1/816816*(3003*a^10*e^10*(16*A*e + B*(d + 17*e*x)) + 2002*a^9*b*e^9*(15*A
*e*(d + 17*e*x) + 2*B*(d^2 + 17*d*e*x + 136*e^2*x^2)) + 1287*a^8*b^2*e^8*(
14*A*e*(d^2 + 17*d*e*x + 136*e^2*x^2) + 3*B*(d^3 + 17*d^2*e*x + 136*d*e^2*
x^2 + 680*e^3*x^3)) + 792*a^7*b^3*e^7*(13*A*e*(d^3 + 17*d^2*e*x + 136*d*e^
2*x^2 + 680*e^3*x^3) + 4*B*(d^4 + 17*d^3*e*x + 136*d^2*e^2*x^2 + 680*d*e^3
*x^3 + 2380*e^4*x^4)) + 462*a^6*b^4*e^6*(12*A*e*(d^4 + 17*d^3*e*x + 136*d^
2*e^2*x^2 + 680*d*e^3*x^3 + 2380*e^4*x^4) + 5*B*(d^5 + 17*d^4*e*x + 136*d^
3*e^2*x^2 + 680*d^2*e^3*x^3 + 2380*d*e^4*x^4 + 6188*e^5*x^5)) + 252*a^5*b^
5*e^5*(11*A*e*(d^5 + 17*d^4*e*x + 136*d^3*e^2*x^2 + 680*d^2*e^3*x^3 + 2380
*d*e^4*x^4 + 6188*e^5*x^5) + 6*B*(d^6 + 17*d^5*e*x + 136*d^4*e^2*x^2 + 680
*d^3*e^3*x^3 + 2380*d^2*e^4*x^4 + 6188*d*e^5*x^5 + 12376*e^6*x^6)) + 126*a
^4*b^6*e^4*(10*A*e*(d^6 + 17*d^5*e*x + 136*d^4*e^2*x^2 + 680*d^3*e^3*x^3 +
2380*d^2*e^4*x^4 + 6188*d*e^5*x^5 + 12376*e^6*x^6) + 7*B*(d^7 + 17*d^6*e*
x + 136*d^5*e^2*x^2 + 680*d^4*e^3*x^3 + 2380*d^3*e^4*x^4 + 6188*d^2*e^5*x^
5 + 12376*d*e^6*x^6 + 19448*e^7*x^7)) + 56*a^3*b^7*e^3*(9*A*e*(d^7 + 17*d^
6*e*x + 136*d^5*e^2*x^2 + 680*d^4*e^3*x^3 + 2380*d^3*e^4*x^4 + 6188*d^2*e^
5*x^5 + 12376*d*e^6*x^6 + 19448*e^7*x^7) + 8*B*(d^8 + 17*d^7*e*x + 136*d^6
*e^2*x^2 + 680*d^5*e^3*x^3 + 2380*d^4*e^4*x^4 + 6188*d^3*e^5*x^5 + 12376*d
^2*e^6*x^6 + 19448*d*e^7*x^7 + 24310*e^8*x^8)) + 21*a^2*b^8*e^2*(8*A*e*(d^
8 + 17*d^7*e*x + 136*d^6*e^2*x^2 + 680*d^5*e^3*x^3 + 2380*d^4*e^4*x^4 + ...

```

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 307, normalized size of antiderivative = 0.92, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.350$, Rules used = {87, 55, 55, 55, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{18}} dx$$

$$\downarrow 87$$

$$\frac{(-17aBe + 6Abe + 11bBd) \int \frac{(a+bx)^{10}}{(d+ex)^{17}} dx}{17e(bd - ae)} - \frac{(a + bx)^{11}(Bd - Ae)}{17e(d + ex)^{17}(bd - ae)}$$

$$\downarrow 55$$

$$\frac{(-17aBe + 6Abe + 11bBd) \left(\frac{5b \int \frac{(a+bx)^{10}}{(d+ex)^{16}} dx}{16(bd-ae)} + \frac{(a+bx)^{11}}{16(d+ex)^{16}(bd-ae)} \right)}{17e(bd-ae)} - \frac{(a+bx)^{11}(Bd-Ae)}{17e(d+ex)^{17}(bd-ae)}$$

↓ 55

$$\frac{(-17aBe + 6Abe + 11bBd) \left(\frac{5b \left(\frac{4b \int \frac{(a+bx)^{10}}{(d+ex)^{15}} dx}{15(bd-ae)} + \frac{(a+bx)^{11}}{15(d+ex)^{15}(bd-ae)} \right)}{16(bd-ae)} + \frac{(a+bx)^{11}}{16(d+ex)^{16}(bd-ae)} \right)}{17e(bd-ae)} - \frac{(a+bx)^{11}(Bd-Ae)}{17e(d+ex)^{17}(bd-ae)}$$

↓ 55

$$\frac{(-17aBe + 6Abe + 11bBd) \left(\frac{5b \left(\frac{4b \left(\frac{3b \int \frac{(a+bx)^{10}}{(d+ex)^{14}} dx}{14(bd-ae)} + \frac{(a+bx)^{11}}{14(d+ex)^{14}(bd-ae)} \right)}{15(bd-ae)} + \frac{(a+bx)^{11}}{15(d+ex)^{15}(bd-ae)} \right)}{16(bd-ae)} + \frac{(a+bx)^{11}}{16(d+ex)^{16}(bd-ae)} \right)}{17e(bd-ae)} - \frac{(a+bx)^{11}(Bd-Ae)}{17e(d+ex)^{17}(bd-ae)}$$

↓ 55

$$\begin{aligned}
 & \left((-17aBe + 6Abe + 11bBd) \left(\frac{4b \left(\frac{3b \int \frac{(a+bx)^{10}}{(d+ex)^{13}} dx + \frac{(a+bx)^{11}}{13(d+ex)^{13}(bd-ae)}}{14(bd-ae)} + \frac{(a+bx)^{11}}{14(d+ex)^{14}(bd-ae)}}{5b} + \frac{(a+bx)^{11}}{15(d+ex)^{15}(bd-ae)} \right) \right) \right) + \frac{\dots}{16(d+ex)^{16}(bd-ae)}
 \end{aligned}$$

$$\frac{17e(bd - ae)}{17e(d + ex)^{17}(bd - ae)} \frac{(a + bx)^{11}(Bd - Ae)}{17e(d + ex)^{17}(bd - ae)}$$

↓ 55

$$\frac{(a+bx)^{11}}{16(d+ex)^{16}(bd-ae)} + \frac{5b \frac{(a+bx)^{11}}{15(d+ex)^{15}(bd-ae)} + \frac{4b \frac{(a+bx)^{11}}{14(d+ex)^{14}(bd-ae)} + \frac{3b \left(\frac{(a+bx)^{11}}{13(d+ex)^{13}(bd-ae)} + \frac{2b \left(\frac{b(a+bx)^{11}}{132(d+ex)^{11}(bd-ae)^2} + \frac{(a+bx)^{11}}{12(d+ex)^{12}(bd-ae)} \right)}{13(bd-ae)} \right)}{14(bd-ae)}}{15(bd-ae)}}{16(bd-ae)}$$

$$\frac{(a+bx)^{11}(Bd - Ae)}{17e(d+ex)^{17}(bd-ae)}$$

input `Int[((a + b*x)^10*(A + B*x))/(d + e*x)^18,x]`

output `-1/17*((B*d - A*e)*(a + b*x)^11)/(e*(b*d - a*e)*(d + e*x)^17) + ((11*b*B*d + 6*A*b*e - 17*a*B*e)*((a + b*x)^11/(16*(b*d - a*e)*(d + e*x)^16) + (5*b*((a + b*x)^11/(15*(b*d - a*e)*(d + e*x)^15) + (4*b*((a + b*x)^11/(14*(b*d - a*e)*(d + e*x)^14) + (3*b*((a + b*x)^11/(13*(b*d - a*e)*(d + e*x)^13) + (2*b*((a + b*x)^11/(12*(b*d - a*e)*(d + e*x)^12) + (b*(a + b*x)^11)/(132*(b*d - a*e)^2*(d + e*x)^11)))/(13*(b*d - a*e)))/(14*(b*d - a*e)))/(15*(b*d - a*e)))/(16*(b*d - a*e)))/(17*e*(b*d - a*e))`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
 [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
 a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 55 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
 (a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
 implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
 c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
 c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
 lerQ[n, 1])`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
 + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
 + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
 /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
 rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1900 vs. $2(321) = 642$.

Time = 0.32 (sec) , antiderivative size = 1901, normalized size of antiderivative = 5.67

method	result	size
risch	Expression too large to display	1901
default	Expression too large to display	1942
norman	Expression too large to display	2014
gospers	Expression too large to display	2233
orering	Expression too large to display	2233
parallelrisch	Expression too large to display	2242

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^18,x,method=_RETURNVERBOSE)`

output

```
(-1/816816/e^12*(48048*A*a^10*e^11+30030*A*a^9*b*d*e^10+18018*A*a^8*b^2*d^2*e^9+10296*A*a^7*b^3*d^3*e^8+5544*A*a^6*b^4*d^4*e^7+2772*A*a^5*b^5*d^5*e^6+1260*A*a^4*b^6*d^6*e^5+504*A*a^3*b^7*d^7*e^4+168*A*a^2*b^8*d^8*e^3+42*A*a*b^9*d^9*e^2+6*A*b^10*d^10*e+3003*B*a^10*d*e^10+4004*B*a^9*b*d^2*e^9+3861*B*a^8*b^2*d^3*e^8+3168*B*a^7*b^3*d^4*e^7+2310*B*a^6*b^4*d^5*e^6+1512*B*a^5*b^5*d^6*e^5+882*B*a^4*b^6*d^7*e^4+448*B*a^3*b^7*d^8*e^3+189*B*a^2*b^8*d^9*e^2+60*B*a*b^9*d^10*e+11*B*b^10*d^11)-1/48048/e^11*(30030*A*a^9*b*e^10+18018*A*a^8*b^2*d*e^9+10296*A*a^7*b^3*d^2*e^8+5544*A*a^6*b^4*d^3*e^7+2772*A*a^5*b^5*d^4*e^6+1260*A*a^4*b^6*d^5*e^5+504*A*a^3*b^7*d^6*e^4+168*A*a^2*b^8*d^7*e^3+42*A*a*b^9*d^8*e^2+6*A*b^10*d^9*e+3003*B*a^10*e^10+4004*B*a^9*b*d*e^9+3861*B*a^8*b^2*d^2*e^8+3168*B*a^7*b^3*d^3*e^7+2310*B*a^6*b^4*d^4*e^6+1512*B*a^5*b^5*d^5*e^5+882*B*a^4*b^6*d^6*e^4+448*B*a^3*b^7*d^7*e^3+189*B*a^2*b^8*d^8*e^2+60*B*a*b^9*d^9*e+11*B*b^10*d^10)*x-1/6006*b/e^10*(18018*A*a^8*b*e^9+10296*A*a^7*b^2*d*e^8+5544*A*a^6*b^3*d^2*e^7+2772*A*a^5*b^4*d^3*e^6+1260*A*a^4*b^5*d^4*e^5+504*A*a^3*b^6*d^5*e^4+168*A*a^2*b^7*d^6*e^3+42*A*a*b^8*d^7*e^2+6*A*b^9*d^8*e+4004*B*a^9*e^9+3861*B*a^8*b*d*e^8+3168*B*a^7*b^2*d^2*e^7+2310*B*a^6*b^3*d^3*e^6+1512*B*a^5*b^4*d^4*e^5+882*B*a^4*b^5*d^5*e^4+448*B*a^3*b^6*d^6*e^3+189*B*a^2*b^7*d^7*e^2+60*B*a*b^8*d^8*e+11*B*b^9*d^9)*x^2-5/6006*b^2/e^9*(10296*A*a^7*b*e^8+5544*A*a^6*b^2*d*e^7+2772*A*a^5*b^3*d^2*e^6+1260*A*a^4*b^4*d^3*e^5+504*A*a^3*b^5*d^4*e^4+168*A*a^2*...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1995 vs. $2(321) = 642$.

Time = 0.17 (sec) , antiderivative size = 1995, normalized size of antiderivative = 5.96

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{18}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)/(e*x+d)^18,x, algorithm="fricas")
```

output

```

-1/816816*(136136*B*b^10*e^11*x^11 + 11*B*b^10*d^11 + 48048*A*a^10*e^11 +
6*(10*B*a*b^9 + A*b^10)*d^10*e + 21*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 + 56
*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 126*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7
*e^4 + 252*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 462*(5*B*a^6*b^4 + 6*A*a^
5*b^5)*d^5*e^6 + 792*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 1287*(3*B*a^8*b
^2 + 8*A*a^7*b^3)*d^3*e^8 + 2002*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 + 3003*
(B*a^10 + 10*A*a^9*b)*d*e^10 + 19448*(11*B*b^10*d*e^10 + 6*(10*B*a*b^9 + A
*b^10)*e^11)*x^10 + 24310*(11*B*b^10*d^2*e^9 + 6*(10*B*a*b^9 + A*b^10)*d*e
^10 + 21*(9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 + 24310*(11*B*b^10*d^3*e^8 +
6*(10*B*a*b^9 + A*b^10)*d^2*e^9 + 21*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 + 56
*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 19448*(11*B*b^10*d^4*e^7 + 6*(10*
B*a*b^9 + A*b^10)*d^3*e^8 + 21*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 + 56*(8*B
*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 + 126*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7
+ 12376*(11*B*b^10*d^5*e^6 + 6*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 21*(9*B*a^
2*b^8 + 2*A*a*b^9)*d^3*e^8 + 56*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 126*
(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 + 252*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^11)
*x^6 + 6188*(11*B*b^10*d^6*e^5 + 6*(10*B*a*b^9 + A*b^10)*d^5*e^6 + 21*(9*B
*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 + 56*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 + 1
26*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 + 252*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d
*e^10 + 462*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 + 2380*(11*B*b^10*d^7...

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{18}} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**10*(B*x+A)/(e*x+d)**18,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1995 vs. $2(321) = 642$.

Time = 0.16 (sec) , antiderivative size = 1995, normalized size of antiderivative = 5.96

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{18}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^18,x, algorithm="maxima")`

output

```
-1/816816*(136136*B*b^10*e^11*x^11 + 11*B*b^10*d^11 + 48048*A*a^10*e^11 +
6*(10*B*a*b^9 + A*b^10)*d^10*e + 21*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 + 56
*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 126*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^7
*e^4 + 252*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 462*(5*B*a^6*b^4 + 6*A*a^
5*b^5)*d^5*e^6 + 792*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 1287*(3*B*a^8*b
^2 + 8*A*a^7*b^3)*d^3*e^8 + 2002*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 + 3003*
(B*a^10 + 10*A*a^9*b)*d*e^10 + 19448*(11*B*b^10*d*e^10 + 6*(10*B*a*b^9 + A
*b^10)*e^11)*x^10 + 24310*(11*B*b^10*d^2*e^9 + 6*(10*B*a*b^9 + A*b^10)*d*e
^10 + 21*(9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 + 24310*(11*B*b^10*d^3*e^8 +
6*(10*B*a*b^9 + A*b^10)*d^2*e^9 + 21*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 + 56
*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 19448*(11*B*b^10*d^4*e^7 + 6*(10*
B*a*b^9 + A*b^10)*d^3*e^8 + 21*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 + 56*(8*B
*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 + 126*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*x^7
+ 12376*(11*B*b^10*d^5*e^6 + 6*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 21*(9*B*a^
2*b^8 + 2*A*a*b^9)*d^3*e^8 + 56*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 126*
(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 + 252*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^11)
*x^6 + 6188*(11*B*b^10*d^6*e^5 + 6*(10*B*a*b^9 + A*b^10)*d^5*e^6 + 21*(9*B
*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 + 56*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8 + 1
26*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 + 252*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d
*e^10 + 462*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 + 2380*(11*B*b^10*d^7...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2232 vs. $2(321) = 642$.

Time = 0.13 (sec) , antiderivative size = 2232, normalized size of antiderivative = 6.66

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{18}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^18,x, algorithm="giac")`

output

```
-1/816816*(136136*B*b^10*e^11*x^11 + 213928*B*b^10*d*e^10*x^10 + 1166880*B
*a*b^9*e^11*x^10 + 116688*A*b^10*e^11*x^10 + 267410*B*b^10*d^2*e^9*x^9 + 1
458600*B*a*b^9*d*e^10*x^9 + 145860*A*b^10*d*e^10*x^9 + 4594590*B*a^2*b^8*e
^11*x^9 + 1021020*A*a*b^9*e^11*x^9 + 267410*B*b^10*d^3*e^8*x^8 + 1458600*B
*a*b^9*d^2*e^9*x^8 + 145860*A*b^10*d^2*e^9*x^8 + 4594590*B*a^2*b^8*d*e^10*
x^8 + 1021020*A*a*b^9*d*e^10*x^8 + 10890880*B*a^3*b^7*e^11*x^8 + 4084080*A
*a^2*b^8*e^11*x^8 + 213928*B*b^10*d^4*e^7*x^7 + 1166880*B*a*b^9*d^3*e^8*x^
7 + 116688*A*b^10*d^3*e^8*x^7 + 3675672*B*a^2*b^8*d^2*e^9*x^7 + 816816*A*a
*b^9*d^2*e^9*x^7 + 8712704*B*a^3*b^7*d*e^10*x^7 + 3267264*A*a^2*b^8*d*e^10
*x^7 + 17153136*B*a^4*b^6*e^11*x^7 + 9801792*A*a^3*b^7*e^11*x^7 + 136136*B
*b^10*d^5*e^6*x^6 + 742560*B*a*b^9*d^4*e^7*x^6 + 74256*A*b^10*d^4*e^7*x^6
+ 2339064*B*a^2*b^8*d^3*e^8*x^6 + 519792*A*a*b^9*d^3*e^8*x^6 + 5544448*B*a
^3*b^7*d^2*e^9*x^6 + 2079168*A*a^2*b^8*d^2*e^9*x^6 + 10915632*B*a^4*b^6*d*
e^10*x^6 + 6237504*A*a^3*b^7*d*e^10*x^6 + 18712512*B*a^5*b^5*e^11*x^6 + 15
593760*A*a^4*b^6*e^11*x^6 + 68068*B*b^10*d^6*e^5*x^5 + 371280*B*a*b^9*d^5*
e^6*x^5 + 37128*A*b^10*d^5*e^6*x^5 + 1169532*B*a^2*b^8*d^4*e^7*x^5 + 25989
6*A*a*b^9*d^4*e^7*x^5 + 2772224*B*a^3*b^7*d^3*e^8*x^5 + 1039584*A*a^2*b^8*
d^3*e^8*x^5 + 5457816*B*a^4*b^6*d^2*e^9*x^5 + 3118752*A*a^3*b^7*d^2*e^9*x^
5 + 9356256*B*a^5*b^5*d*e^10*x^5 + 7796880*A*a^4*b^6*d*e^10*x^5 + 14294280
*B*a^6*b^4*e^11*x^5 + 17153136*A*a^5*b^5*e^11*x^5 + 26180*B*b^10*d^7*e^...
```

Mupad [B] (verification not implemented)

Time = 1.40 (sec) , antiderivative size = 2077, normalized size of antiderivative = 6.20

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{18}} dx = \text{Too large to display}$$

input `int(((A + B*x)*(a + b*x)^10)/(d + e*x)^18,x)`

output

```

-((48048*A*a^10*e^11 + 11*B*b^10*d^11 + 6*A*b^10*d^10*e + 3003*B*a^10*d*e^10 + 42*A*a*b^9*d^9*e^2 + 4004*B*a^9*b*d^2*e^9 + 168*A*a^2*b^8*d^8*e^3 + 504*A*a^3*b^7*d^7*e^4 + 1260*A*a^4*b^6*d^6*e^5 + 2772*A*a^5*b^5*d^5*e^6 + 5544*A*a^6*b^4*d^4*e^7 + 10296*A*a^7*b^3*d^3*e^8 + 18018*A*a^8*b^2*d^2*e^9 + 189*B*a^2*b^8*d^9*e^2 + 448*B*a^3*b^7*d^8*e^3 + 882*B*a^4*b^6*d^7*e^4 + 1512*B*a^5*b^5*d^6*e^5 + 2310*B*a^6*b^4*d^5*e^6 + 3168*B*a^7*b^3*d^4*e^7 + 3861*B*a^8*b^2*d^3*e^8 + 30030*A*a^9*b*d*e^10 + 60*B*a*b^9*d^10*e)/(816816*e^12) + (x*(3003*B*a^10*e^10 + 11*B*b^10*d^10 + 30030*A*a^9*b*d*e^10 + 6*A*b^10*d^9*e + 42*A*a*b^9*d^8*e^2 + 18018*A*a^8*b^2*d*e^9 + 168*A*a^2*b^8*d^7*e^3 + 504*A*a^3*b^7*d^6*e^4 + 1260*A*a^4*b^6*d^5*e^5 + 2772*A*a^5*b^5*d^4*e^6 + 5544*A*a^6*b^4*d^3*e^7 + 10296*A*a^7*b^3*d^2*e^8 + 189*B*a^2*b^8*d^8*e^2 + 448*B*a^3*b^7*d^7*e^3 + 882*B*a^4*b^6*d^6*e^4 + 1512*B*a^5*b^5*d^5*e^5 + 2310*B*a^6*b^4*d^4*e^6 + 3168*B*a^7*b^3*d^3*e^7 + 3861*B*a^8*b^2*d^2*e^8 + 60*B*a*b^9*d^9*e + 4004*B*a^9*b*d*e^9))/(48048*e^11) + (5*b^7*x^8*(448*B*a^3*e^3 + 11*B*b^3*d^3 + 168*A*a^2*b*e^3 + 6*A*b^3*d^2*e + 42*A*a*b^2*d*e^2 + 60*B*a*b^2*d^2*e + 189*B*a^2*b*d*e^2))/(168*e^4) + (b^4*x^5*(2310*B*a^6*e^6 + 11*B*b^6*d^6 + 2772*A*a^5*b*e^6 + 6*A*b^6*d^5*e + 42*A*a*b^5*d^4*e^2 + 1260*A*a^4*b^2*d*e^5 + 168*A*a^2*b^4*d^3*e^3 + 504*A*a^3*b^3*d^2*e^4 + 189*B*a^2*b^4*d^4*e^2 + 448*B*a^3*b^3*d^3*e^3 + 882*B*a^4*b^2*d^2*e^4 + 60*B*a*b^5*d^5*e + 1512*B*a^5*b*d*e^5))/(132*e^7) + (b^9*x^10*...

```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 1327, normalized size of antiderivative = 3.96

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{18}} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^18,x)`

output

```
( - 4368*a**11*e**11 - 3003*a**10*b*d*e**10 - 51051*a**10*b*e**11*x - 2002
*a**9*b**2*d**2*e**9 - 34034*a**9*b**2*d*e**10*x - 272272*a**9*b**2*e**11*
x**2 - 1287*a**8*b**3*d**3*e**8 - 21879*a**8*b**3*d**2*e**9*x - 175032*a**
8*b**3*d*e**10*x**2 - 875160*a**8*b**3*e**11*x**3 - 792*a**7*b**4*d**4*e**
7 - 13464*a**7*b**4*d**3*e**8*x - 107712*a**7*b**4*d**2*e**9*x**2 - 538560
*a**7*b**4*d*e**10*x**3 - 1884960*a**7*b**4*e**11*x**4 - 462*a**6*b**5*d**
5*e**6 - 7854*a**6*b**5*d**4*e**7*x - 62832*a**6*b**5*d**3*e**8*x**2 - 314
160*a**6*b**5*d**2*e**9*x**3 - 1099560*a**6*b**5*d*e**10*x**4 - 2858856*a*
*b**5*e**11*x**5 - 252*a**5*b**6*d**6*e**5 - 4284*a**5*b**6*d**5*e**6*x
- 34272*a**5*b**6*d**4*e**7*x**2 - 171360*a**5*b**6*d**3*e**8*x**3 - 59976
0*a**5*b**6*d**2*e**9*x**4 - 1559376*a**5*b**6*d*e**10*x**5 - 3118752*a**5
*b**6*e**11*x**6 - 126*a**4*b**7*d**7*e**4 - 2142*a**4*b**7*d**6*e**5*x -
17136*a**4*b**7*d**5*e**6*x**2 - 85680*a**4*b**7*d**4*e**7*x**3 - 299880*a
**4*b**7*d**3*e**8*x**4 - 779688*a**4*b**7*d**2*e**9*x**5 - 1559376*a**4*b
**7*d*e**10*x**6 - 2450448*a**4*b**7*e**11*x**7 - 56*a**3*b**8*d**8*e**3 -
952*a**3*b**8*d**7*e**4*x - 7616*a**3*b**8*d**6*e**5*x**2 - 38080*a**3*b*
*8*d**5*e**6*x**3 - 133280*a**3*b**8*d**4*e**7*x**4 - 346528*a**3*b**8*d**
3*e**8*x**5 - 693056*a**3*b**8*d**2*e**9*x**6 - 1089088*a**3*b**8*d*e**10*
x**7 - 1361360*a**3*b**8*e**11*x**8 - 21*a**2*b**9*d**9*e**2 - 357*a**2*b*
*9*d**8*e**3*x - 2856*a**2*b**9*d**7*e**4*x**2 - 14280*a**2*b**9*d**6*e...
```

3.97 $\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{19}} dx$

Optimal result	1002
Mathematica [B] (verified)	1003
Rubi [A] (verified)	1004
Maple [B] (verified)	1012
Fricas [B] (verification not implemented)	1013
Sympy [F(-1)]	1014
Maxima [B] (verification not implemented)	1014
Giac [B] (verification not implemented)	1015
Mupad [B] (verification not implemented)	1016
Reduce [B] (verification not implemented)	1017

Optimal result

Integrand size = 20, antiderivative size = 385

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{19}} dx = -\frac{(Bd - Ae)(a+bx)^{11}}{18e(bd - ae)(d+ex)^{18}} + \frac{(11bBd + 7Abe - 18aBe)(a+bx)^{11}}{306e(bd - ae)^2(d+ex)^{17}} + \frac{b(11bBd + 7Abe - 18aBe)(a+bx)^{11}}{816e(bd - ae)^3(d+ex)^{16}} + \frac{b^2(11bBd + 7Abe - 18aBe)(a+bx)^{11}}{2448e(bd - ae)^4(d+ex)^{15}} + \frac{b^3(11bBd + 7Abe - 18aBe)(a+bx)^{11}}{8568e(bd - ae)^5(d+ex)^{14}} + \frac{b^4(11bBd + 7Abe - 18aBe)(a+bx)^{11}}{37128e(bd - ae)^6(d+ex)^{13}} + \frac{b^5(11bBd + 7Abe - 18aBe)(a+bx)^{11}}{222768e(bd - ae)^7(d+ex)^{12}} + \frac{b^6(11bBd + 7Abe - 18aBe)(a+bx)^{11}}{2450448e(bd - ae)^8(d+ex)^{11}}$$

output

```
-1/18*(-A*e+B*d)*(b*x+a)^11/e/(-a*e+b*d)/(e*x+d)^18+1/306*(7*A*b*e-18*B*a*
e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^2/(e*x+d)^17+1/816*b*(7*A*b*e-18*B*a*
e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^3/(e*x+d)^16+1/2448*b^2*(7*A*b*e-18*B*a*
e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^4/(e*x+d)^15+1/8568*b^3*(7*A*b*e-18*B*
a*e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^5/(e*x+d)^14+1/37128*b^4*(7*A*b*e-1
8*B*a*e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^6/(e*x+d)^13+1/222768*b^5*(7*A*b*
e-18*B*a*e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^7/(e*x+d)^12+1/2450448*b^6*(
7*A*b*e-18*B*a*e+11*B*b*d)*(b*x+a)^11/e/(-a*e+b*d)^8/(e*x+d)^11
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1428 vs. $2(385) = 770$.

Time = 0.48 (sec) , antiderivative size = 1428, normalized size of antiderivative = 3.71

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{19}} dx = \text{Too large to display}$$

input

```
Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^19,x]
```


output

```
-1/2450448*(8008*a^10*e^10*(17*A*e + B*(d + 18*e*x)) + 10010*a^9*b*e^9*(8*
A*e*(d + 18*e*x) + B*(d^2 + 18*d*e*x + 153*e^2*x^2)) + 9009*a^8*b^2*e^8*(5
*A*e*(d^2 + 18*d*e*x + 153*e^2*x^2) + B*(d^3 + 18*d^2*e*x + 153*d*e^2*x^2
+ 816*e^3*x^3)) + 3432*a^7*b^3*e^7*(7*A*e*(d^3 + 18*d^2*e*x + 153*d*e^2*x^
2 + 816*e^3*x^3) + 2*B*(d^4 + 18*d^3*e*x + 153*d^2*e^2*x^2 + 816*d*e^3*x^3
+ 3060*e^4*x^4)) + 924*a^6*b^4*e^6*(13*A*e*(d^4 + 18*d^3*e*x + 153*d^2*e^
2*x^2 + 816*d*e^3*x^3 + 3060*e^4*x^4) + 5*B*(d^5 + 18*d^4*e*x + 153*d^3*e^
2*x^2 + 816*d^2*e^3*x^3 + 3060*d*e^4*x^4 + 8568*e^5*x^5)) + 2772*a^5*b^5*e
^5*(2*A*e*(d^5 + 18*d^4*e*x + 153*d^3*e^2*x^2 + 816*d^2*e^3*x^3 + 3060*d*e
^4*x^4 + 8568*e^5*x^5) + B*(d^6 + 18*d^5*e*x + 153*d^4*e^2*x^2 + 816*d^3*e
^3*x^3 + 3060*d^2*e^4*x^4 + 8568*d*e^5*x^5 + 18564*e^6*x^6)) + 210*a^4*b^6
*e^4*(11*A*e*(d^6 + 18*d^5*e*x + 153*d^4*e^2*x^2 + 816*d^3*e^3*x^3 + 3060*
d^2*e^4*x^4 + 8568*d*e^5*x^5 + 18564*e^6*x^6) + 7*B*(d^7 + 18*d^6*e*x + 15
3*d^5*e^2*x^2 + 816*d^4*e^3*x^3 + 3060*d^3*e^4*x^4 + 8568*d^2*e^5*x^5 + 18
564*d*e^6*x^6 + 31824*e^7*x^7)) + 168*a^3*b^7*e^3*(5*A*e*(d^7 + 18*d^6*e*x
+ 153*d^5*e^2*x^2 + 816*d^4*e^3*x^3 + 3060*d^3*e^4*x^4 + 8568*d^2*e^5*x^5
+ 18564*d*e^6*x^6 + 31824*e^7*x^7) + 4*B*(d^8 + 18*d^7*e*x + 153*d^6*e^2*
x^2 + 816*d^5*e^3*x^3 + 3060*d^4*e^4*x^4 + 8568*d^3*e^5*x^5 + 18564*d^2*e^
6*x^6 + 31824*d*e^7*x^7 + 43758*e^8*x^8)) + 252*a^2*b^8*e^2*(A*e*(d^8 + 18
*d^7*e*x + 153*d^6*e^2*x^2 + 816*d^5*e^3*x^3 + 3060*d^4*e^4*x^4 + 8568*...
```

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 351, normalized size of antiderivative = 0.91, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {87, 55, 55, 55, 55, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{19}} dx$$

↓ 87

$$\frac{(-18aBe + 7Abe + 11bBd) \int \frac{(a+bx)^{10}}{(d+ex)^{18}} dx}{18e(bd - ae)} - \frac{(a + bx)^{11}(Bd - Ae)}{18e(d + ex)^{18}(bd - ae)}$$

↓ 55

$$\frac{(-18aBe + 7Abe + 11bBd) \left(\frac{6b \int \frac{(a+bx)^{10}}{(d+ex)^{17}} dx}{17(bd-ae)} + \frac{(a+bx)^{11}}{17(d+ex)^{17}(bd-ae)} \right)}{18e(bd-ae)} - \frac{(a+bx)^{11}(Bd-Ae)}{18e(d+ex)^{18}(bd-ae)}$$

↓ 55

$$\frac{(-18aBe + 7Abe + 11bBd) \left(\frac{6b \left(\frac{5b \int \frac{(a+bx)^{10}}{(d+ex)^{16}} dx}{16(bd-ae)} + \frac{(a+bx)^{11}}{16(d+ex)^{16}(bd-ae)} \right)}{17(bd-ae)} + \frac{(a+bx)^{11}}{17(d+ex)^{17}(bd-ae)} \right)}{18e(bd-ae)} - \frac{(a+bx)^{11}(Bd-Ae)}{18e(d+ex)^{18}(bd-ae)}$$

↓ 55

$$\frac{(-18aBe + 7Abe + 11bBd) \left(\frac{6b \left(\frac{5b \left(\frac{4b \int \frac{(a+bx)^{10}}{(d+ex)^{15}} dx}{15(bd-ae)} + \frac{(a+bx)^{11}}{15(d+ex)^{15}(bd-ae)} \right)}{16(bd-ae)} + \frac{(a+bx)^{11}}{16(d+ex)^{16}(bd-ae)} \right)}{17(bd-ae)} + \frac{(a+bx)^{11}}{17(d+ex)^{17}(bd-ae)} \right)}{18e(bd-ae)} - \frac{(a+bx)^{11}(Bd-Ae)}{18e(d+ex)^{18}(bd-ae)}$$

↓ 55

$$\begin{aligned}
 & \left(\left(\left(\frac{3b \int \frac{(a+bx)^{10}}{(d+ex)^{14}} dx}{14(bd-ae)} + \frac{(a+bx)^{11}}{14(d+ex)^{14}(bd-ae)} \right) + \frac{(a+bx)^{11}}{15(d+ex)^{15}(bd-ae)} \right) \right. \\
 & \left. \frac{5b}{15(bd-ae)} + \frac{(a+bx)^{11}}{15(d+ex)^{15}(bd-ae)} \right) \\
 & \left(\frac{6b}{16(bd-ae)} + \frac{(a+bx)^{11}}{16(d+ex)^{16}(bd-ae)} \right) \\
 & \left(\frac{(-18aBe + 7Abe + 11bBd)}{17(bd-ae)} + \frac{(a+bx)^{11}}{17(d+ex)^{17}(bd-ae)} \right)
 \end{aligned}$$

$$\frac{18e(bd-ae)}{18e(d+ex)^{18}(bd-ae)} \frac{(a+bx)^{11}(Bd-Ae)}{18e(d+ex)^{18}(bd-ae)}$$

↓ 55

$$\begin{aligned}
 & \left(\frac{3b \left(\frac{2b \int \frac{(a+bx)^{10}}{(d+ex)^{13}} dx}{13(bd-ae)} + \frac{(a+bx)^{11}}{13(d+ex)^{13}(bd-ae)} \right)}{14(bd-ae)} + \frac{(a+bx)^{11}}{14(d+ex)^{14}(bd-ae)} \right) \\
 & \frac{5b \left(\frac{\left(\frac{3b \left(\frac{2b \int \frac{(a+bx)^{10}}{(d+ex)^{13}} dx}{13(bd-ae)} + \frac{(a+bx)^{11}}{13(d+ex)^{13}(bd-ae)} \right)}{14(bd-ae)} + \frac{(a+bx)^{11}}{14(d+ex)^{14}(bd-ae)} \right)}{15(bd-ae)} + \frac{(a+bx)^{11}}{15(d+ex)^{15}(bd-ae)} \right)}{16(bd-ae)} \\
 & \frac{6b \left(\frac{\left(\frac{5b \left(\frac{\left(\frac{3b \left(\frac{2b \int \frac{(a+bx)^{10}}{(d+ex)^{13}} dx}{13(bd-ae)} + \frac{(a+bx)^{11}}{13(d+ex)^{13}(bd-ae)} \right)}{14(bd-ae)} + \frac{(a+bx)^{11}}{14(d+ex)^{14}(bd-ae)} \right)}{15(bd-ae)} + \frac{(a+bx)^{11}}{15(d+ex)^{15}(bd-ae)} \right)}{16(bd-ae)} + \frac{(a+bx)^{11}}{16(d+ex)^{16}(bd-ae)} \right)}{17(bd-ae)} \right)}{18e(bd-ae)} \\
 & (-18aBe + 7Abe + 11bBd) \frac{\left(\frac{6b \left(\frac{\left(\frac{5b \left(\frac{\left(\frac{3b \left(\frac{2b \int \frac{(a+bx)^{10}}{(d+ex)^{13}} dx}{13(bd-ae)} + \frac{(a+bx)^{11}}{13(d+ex)^{13}(bd-ae)} \right)}{14(bd-ae)} + \frac{(a+bx)^{11}}{14(d+ex)^{14}(bd-ae)} \right)}{15(bd-ae)} + \frac{(a+bx)^{11}}{15(d+ex)^{15}(bd-ae)} \right)}{16(bd-ae)} + \frac{(a+bx)^{11}}{16(d+ex)^{16}(bd-ae)} \right)}{17(bd-ae)} \right)}{18e(bd-ae)} \right)}{18e(d+ex)^{18}(bd-ae)} \\
 & \downarrow 55
 \end{aligned}$$

↓ 48

$$\left(\frac{(a+bx)^{11}}{17(d+ex)^{17}(bd-ae)} + \frac{6b}{16(d+ex)^{16}(bd-ae)} + \frac{5b}{15(d+ex)^{15}(bd-ae)} + \frac{4b}{14(d+ex)^{14}(bd-ae)} + \frac{3b \left(\frac{(a+bx)^{11}}{13(d+ex)^{13}(bd-ae)} + \frac{2b \left(\frac{b(a+bx)^{11}}{132(d+ex)^{11}(bd-ae)} + \frac{13}{13} \right)}{14(bd-ae)} \right)}{15(bd-ae)} \right)$$

$$\frac{(a+bx)^{11}(Bd - Ae)}{18e(d+ex)^{18}(bd-ae)}$$

$$18e(bd - ae)$$

input `Int[((a + b*x)^10*(A + B*x))/(d + e*x)^19,x]`

output
$$-1/18*((B*d - A*e)*(a + b*x)^{11}/(e*(b*d - a*e)*(d + e*x)^{18}) + ((11*b*B*d + 7*A*b*e - 18*a*B*e)*((a + b*x)^{11}/(17*(b*d - a*e)*(d + e*x)^{17}) + (6*b*((a + b*x)^{11}/(16*(b*d - a*e)*(d + e*x)^{16}) + (5*b*((a + b*x)^{11}/(15*(b*d - a*e)*(d + e*x)^{15}) + (4*b*((a + b*x)^{11}/(14*(b*d - a*e)*(d + e*x)^{14}) + (3*b*((a + b*x)^{11}/(13*(b*d - a*e)*(d + e*x)^{13}) + (2*b*((a + b*x)^{11}/(12*(b*d - a*e)*(d + e*x)^{12}) + (b*(a + b*x)^{11}/(132*(b*d - a*e)^2*(d + e*x)^{11}))/((13*(b*d - a*e))))/(14*(b*d - a*e)))/(15*(b*d - a*e)))/(16*(b*d - a*e)))/(17*(b*d - a*e)))/(18*e*(b*d - a*e))$$

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 55 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(Simplify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1900 vs. $2(369) = 738$.

Time = 0.34 (sec) , antiderivative size = 1901, normalized size of antiderivative = 4.94

method	result	size
risch	Expression too large to display	1901
default	Expression too large to display	1942
norman	Expression too large to display	2014
gospers	Expression too large to display	2233
orering	Expression too large to display	2233
parallelrisch	Expression too large to display	2242

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^19,x,method=_RETURNVERBOSE)`

output

```
(-1/2450448/e^12*(136136*A*a^10*e^11+80080*A*a^9*b*d*e^10+45045*A*a^8*b^2*d^2*e^9+24024*A*a^7*b^3*d^3*e^8+12012*A*a^6*b^4*d^4*e^7+5544*A*a^5*b^5*d^5*e^6+2310*A*a^4*b^6*d^6*e^5+840*A*a^3*b^7*d^7*e^4+252*A*a^2*b^8*d^8*e^3+56*A*a*b^9*d^9*e^2+7*A*b^10*d^10*e+8008*B*a^10*d*e^10+10010*B*a^9*b*d^2*e^9+9009*B*a^8*b^2*d^3*e^8+6864*B*a^7*b^3*d^4*e^7+4620*B*a^6*b^4*d^5*e^6+2772*B*a^5*b^5*d^6*e^5+1470*B*a^4*b^6*d^7*e^4+672*B*a^3*b^7*d^8*e^3+252*B*a^2*b^8*d^9*e^2+70*B*a*b^9*d^10*e+11*B*b^10*d^11)-1/136136/e^11*(80080*A*a^9*b*e^10+45045*A*a^8*b^2*d*e^9+24024*A*a^7*b^3*d^2*e^8+12012*A*a^6*b^4*d^3*e^7+5544*A*a^5*b^5*d^4*e^6+2310*A*a^4*b^6*d^5*e^5+840*A*a^3*b^7*d^6*e^4+252*A*a^2*b^8*d^7*e^3+56*A*a*b^9*d^8*e^2+7*A*b^10*d^9*e+8008*B*a^10*e^10+10010*B*a^9*b*d*e^9+9009*B*a^8*b^2*d^2*e^8+6864*B*a^7*b^3*d^3*e^7+4620*B*a^6*b^4*d^4*e^6+2772*B*a^5*b^5*d^5*e^5+1470*B*a^4*b^6*d^6*e^4+672*B*a^3*b^7*d^7*e^3+252*B*a^2*b^8*d^8*e^2+70*B*a*b^9*d^9*e+11*B*b^10*d^10)*x-1/16016*b/e^10*(45045*A*a^8*b*e^9+24024*A*a^7*b^2*d*e^8+12012*A*a^6*b^3*d^2*e^7+5544*A*a^5*b^4*d^3*e^6+2310*A*a^4*b^5*d^4*e^5+840*A*a^3*b^6*d^5*e^4+252*A*a^2*b^7*d^6*e^3+56*A*a*b^8*d^7*e^2+7*A*b^9*d^8*e+10010*B*a^9*e^9+9009*B*a^8*b*d*e^8+6864*B*a^7*b^2*d^2*e^7+4620*B*a^6*b^3*d^3*e^6+2772*B*a^5*b^4*d^4*e^5+1470*B*a^4*b^5*d^5*e^4+672*B*a^3*b^6*d^6*e^3+252*B*a^2*b^7*d^7*e^2+70*B*a*b^8*d^8*e+11*B*b^9*d^9)*x^2-1/3003*b^2/e^9*(24024*A*a^7*b*e^8+12012*A*a^6*b^2*d*e^7+5544*A*a^5*b^3*d^2*e^6+2310*A*a^4*b^4*d^3*e^5+840*A*a^3*b^5*d^4*...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2006 vs. $2(369) = 738$.

Time = 0.20 (sec) , antiderivative size = 2006, normalized size of antiderivative = 5.21

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{19}} dx = \text{Too large to display}$$

```
input integrate((b*x+a)^10*(B*x+A)/(e*x+d)^19,x, algorithm="fricas")
```

output

```
-1/2450448*(350064*B*b^10*e^11*x^11 + 11*B*b^10*d^11 + 136136*A*a^10*e^11
+ 7*(10*B*a*b^9 + A*b^10)*d^10*e + 28*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 +
84*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 210*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d
^7*e^4 + 462*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 924*(5*B*a^6*b^4 + 6*A
a^5*b^5)*d^5*e^6 + 1716*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 3003*(3*B*a
^8*b^2 + 8*A*a^7*b^3)*d^3*e^8 + 5005*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 + 80
08*(B*a^10 + 10*A*a^9*b)*d*e^10 + 43758*(11*B*b^10*d*e^10 + 7*(10*B*a*b^9
+ A*b^10)*e^11)*x^10 + 48620*(11*B*b^10*d^2*e^9 + 7*(10*B*a*b^9 + A*b^10)*
d*e^10 + 28*(9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 + 43758*(11*B*b^10*d^3*e^8
+ 7*(10*B*a*b^9 + A*b^10)*d^2*e^9 + 28*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 +
84*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 31824*(11*B*b^10*d^4*e^7 + 7*(
10*B*a*b^9 + A*b^10)*d^3*e^8 + 28*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 + 84*(
8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 + 210*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*
x^7 + 18564*(11*B*b^10*d^5*e^6 + 7*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 28*(9*B
a^2*b^8 + 2*A*a*b^9)*d^3*e^8 + 84*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 2
10*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 + 462*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e
^11)*x^6 + 8568*(11*B*b^10*d^6*e^5 + 7*(10*B*a*b^9 + A*b^10)*d^5*e^6 + 28*(
9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 + 84*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8
+ 210*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 + 462*(6*B*a^5*b^5 + 5*A*a^4*b^6
)*d*e^10 + 924*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 + 3060*(11*B*b^10*...
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{19}} dx = \text{Timed out}$$

input `integrate((b*x+a)**10*(B*x+A)/(e*x+d)**19,x)`output `Timed out`**Maxima [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 2006 vs. $2(369) = 738$.

Time = 0.13 (sec) , antiderivative size = 2006, normalized size of antiderivative = 5.21

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{19}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^19,x, algorithm="maxima")`

output

```

-1/2450448*(350064*B*b^10*e^11*x^11 + 11*B*b^10*d^11 + 136136*A*a^10*e^11
+ 7*(10*B*a*b^9 + A*b^10)*d^10*e + 28*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 +
84*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 210*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d
^7*e^4 + 462*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 924*(5*B*a^6*b^4 + 6*A*
a^5*b^5)*d^5*e^6 + 1716*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 3003*(3*B*a^
8*b^2 + 8*A*a^7*b^3)*d^3*e^8 + 5005*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 + 80
08*(B*a^10 + 10*A*a^9*b)*d*e^10 + 43758*(11*B*b^10*d*e^10 + 7*(10*B*a*b^9
+ A*b^10)*e^11)*x^10 + 48620*(11*B*b^10*d^2*e^9 + 7*(10*B*a*b^9 + A*b^10)*
d*e^10 + 28*(9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 + 43758*(11*B*b^10*d^3*e^8
+ 7*(10*B*a*b^9 + A*b^10)*d^2*e^9 + 28*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^10 +
84*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 31824*(11*B*b^10*d^4*e^7 + 7*(
10*B*a*b^9 + A*b^10)*d^3*e^8 + 28*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 + 84*(
8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 + 210*(7*B*a^4*b^6 + 4*A*a^3*b^7)*e^11)*
x^7 + 18564*(11*B*b^10*d^5*e^6 + 7*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 28*(9*B
*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 + 84*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*e^9 + 2
10*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 + 462*(6*B*a^5*b^5 + 5*A*a^4*b^6)*e^
11)*x^6 + 8568*(11*B*b^10*d^6*e^5 + 7*(10*B*a*b^9 + A*b^10)*d^5*e^6 + 28*(
9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 + 84*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^3*e^8
+ 210*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 + 462*(6*B*a^5*b^5 + 5*A*a^4*b^6
)*d*e^10 + 924*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 + 3060*(11*B*b^10*...

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2232 vs. $2(369) = 738$.

Time = 0.13 (sec) , antiderivative size = 2232, normalized size of antiderivative = 5.80

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{19}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^10*(B*x+A)/(e*x+d)^19,x, algorithm="giac")
```

output

```

-1/2450448*(350064*B*b^10*e^11*x^11 + 481338*B*b^10*d*e^10*x^10 + 3063060*
B*a*b^9*e^11*x^10 + 306306*A*b^10*e^11*x^10 + 534820*B*b^10*d^2*e^9*x^9 +
3403400*B*a*b^9*d*e^10*x^9 + 340340*A*b^10*d*e^10*x^9 + 12252240*B*a^2*b^8
*e^11*x^9 + 2722720*A*a*b^9*e^11*x^9 + 481338*B*b^10*d^3*e^8*x^8 + 3063060
*B*a*b^9*d^2*e^9*x^8 + 306306*A*b^10*d^2*e^9*x^8 + 11027016*B*a^2*b^8*d*e^
10*x^8 + 2450448*A*a*b^9*d*e^10*x^8 + 29405376*B*a^3*b^7*e^11*x^8 + 110270
16*A*a^2*b^8*e^11*x^8 + 350064*B*b^10*d^4*e^7*x^7 + 2227680*B*a*b^9*d^3*e^
8*x^7 + 222768*A*b^10*d^3*e^8*x^7 + 8019648*B*a^2*b^8*d^2*e^9*x^7 + 178214
4*A*a*b^9*d^2*e^9*x^7 + 21385728*B*a^3*b^7*d*e^10*x^7 + 8019648*A*a^2*b^8*
d*e^10*x^7 + 46781280*B*a^4*b^6*e^11*x^7 + 26732160*A*a^3*b^7*e^11*x^7 + 2
04204*B*b^10*d^5*e^6*x^6 + 1299480*B*a*b^9*d^4*e^7*x^6 + 129948*A*b^10*d^4
*e^7*x^6 + 4678128*B*a^2*b^8*d^3*e^8*x^6 + 1039584*A*a*b^9*d^3*e^8*x^6 + 1
2475008*B*a^3*b^7*d^2*e^9*x^6 + 4678128*A*a^2*b^8*d^2*e^9*x^6 + 27289080*B
*a^4*b^6*d*e^10*x^6 + 15593760*A*a^3*b^7*d*e^10*x^6 + 51459408*B*a^5*b^5*e
^11*x^6 + 42882840*A*a^4*b^6*e^11*x^6 + 94248*B*b^10*d^6*e^5*x^5 + 599760*
B*a*b^9*d^5*e^6*x^5 + 59976*A*b^10*d^5*e^6*x^5 + 2159136*B*a^2*b^8*d^4*e^7
*x^5 + 479808*A*a*b^9*d^4*e^7*x^5 + 5757696*B*a^3*b^7*d^3*e^8*x^5 + 215913
6*A*a^2*b^8*d^3*e^8*x^5 + 12594960*B*a^4*b^6*d^2*e^9*x^5 + 7197120*A*a^3*b
^7*d^2*e^9*x^5 + 23750496*B*a^5*b^5*d*e^10*x^5 + 19792080*A*a^4*b^6*d*e^10
*x^5 + 39584160*B*a^6*b^4*e^11*x^5 + 47500992*A*a^5*b^5*e^11*x^5 + 3366...

```

Mupad [B] (verification not implemented)

Time = 9.64 (sec) , antiderivative size = 2088, normalized size of antiderivative = 5.42

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{19}} dx = \text{Too large to display}$$

input

```
int(((A + B*x)*(a + b*x)^10)/(d + e*x)^19,x)
```

output

```

-((136136*A*a^10*e^11 + 11*B*b^10*d^11 + 7*A*b^10*d^10*e + 8008*B*a^10*d*e
^10 + 56*A*a*b^9*d^9*e^2 + 10010*B*a^9*b*d^2*e^9 + 252*A*a^2*b^8*d^8*e^3 +
840*A*a^3*b^7*d^7*e^4 + 2310*A*a^4*b^6*d^6*e^5 + 5544*A*a^5*b^5*d^5*e^6 +
12012*A*a^6*b^4*d^4*e^7 + 24024*A*a^7*b^3*d^3*e^8 + 45045*A*a^8*b^2*d^2*e
^9 + 252*B*a^2*b^8*d^9*e^2 + 672*B*a^3*b^7*d^8*e^3 + 1470*B*a^4*b^6*d^7*e^
4 + 2772*B*a^5*b^5*d^6*e^5 + 4620*B*a^6*b^4*d^5*e^6 + 6864*B*a^7*b^3*d^4*e
^7 + 9009*B*a^8*b^2*d^3*e^8 + 80080*A*a^9*b*d*e^10 + 70*B*a*b^9*d^10*e)/(2
450448*e^12) + (x*(8008*B*a^10*e^10 + 11*B*b^10*d^10 + 80080*A*a^9*b*e^10
+ 7*A*b^10*d^9*e + 56*A*a*b^9*d^8*e^2 + 45045*A*a^8*b^2*d*e^9 + 252*A*a^2*
b^8*d^7*e^3 + 840*A*a^3*b^7*d^6*e^4 + 2310*A*a^4*b^6*d^5*e^5 + 5544*A*a^5*
b^5*d^4*e^6 + 12012*A*a^6*b^4*d^3*e^7 + 24024*A*a^7*b^3*d^2*e^8 + 252*B*a^
2*b^8*d^8*e^2 + 672*B*a^3*b^7*d^7*e^3 + 1470*B*a^4*b^6*d^6*e^4 + 2772*B*a^
5*b^5*d^5*e^5 + 4620*B*a^6*b^4*d^4*e^6 + 6864*B*a^7*b^3*d^3*e^7 + 9009*B*a
^8*b^2*d^2*e^8 + 70*B*a*b^9*d^9*e + 10010*B*a^9*b*d*e^9))/(136136*e^11) +
(b^7*x^8*(672*B*a^3*e^3 + 11*B*b^3*d^3 + 252*A*a^2*b*e^3 + 7*A*b^3*d^2*e +
56*A*a*b^2*d*e^2 + 70*B*a*b^2*d^2*e + 252*B*a^2*b*d*e^2))/(56*e^4) + (b^4
*x^5*(4620*B*a^6*e^6 + 11*B*b^6*d^6 + 5544*A*a^5*b*e^6 + 7*A*b^6*d^5*e + 5
6*A*a*b^5*d^4*e^2 + 2310*A*a^4*b^2*d*e^5 + 252*A*a^2*b^4*d^3*e^3 + 840*A*a
^3*b^3*d^2*e^4 + 252*B*a^2*b^4*d^4*e^2 + 672*B*a^3*b^3*d^3*e^3 + 1470*B*a^
4*b^2*d^2*e^4 + 70*B*a*b^5*d^5*e + 2772*B*a^5*b*d*e^5))/(286*e^7) + (b^...

```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 1338, normalized size of antiderivative = 3.48

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{19}} dx = \text{Too large to display}$$

input

```
int((b*x+a)^10*(B*x+A)/(e*x+d)^19,x)
```

output

```
( - 12376*a**11*e**11 - 8008*a**10*b*d*e**10 - 144144*a**10*b*e**11*x - 50
05*a**9*b**2*d**2*e**9 - 90090*a**9*b**2*d*e**10*x - 765765*a**9*b**2*e**1
1*x**2 - 3003*a**8*b**3*d**3*e**8 - 54054*a**8*b**3*d**2*e**9*x - 459459*a
**8*b**3*d*e**10*x**2 - 2450448*a**8*b**3*e**11*x**3 - 1716*a**7*b**4*d**4
*e**7 - 30888*a**7*b**4*d**3*e**8*x - 262548*a**7*b**4*d**2*e**9*x**2 - 14
00256*a**7*b**4*d*e**10*x**3 - 5250960*a**7*b**4*e**11*x**4 - 924*a**6*b**
5*d**5*e**6 - 16632*a**6*b**5*d**4*e**7*x - 141372*a**6*b**5*d**3*e**8*x**
2 - 753984*a**6*b**5*d**2*e**9*x**3 - 2827440*a**6*b**5*d*e**10*x**4 - 791
6832*a**6*b**5*e**11*x**5 - 462*a**5*b**6*d**6*e**5 - 8316*a**5*b**6*d**5*
e**6*x - 70686*a**5*b**6*d**4*e**7*x**2 - 376992*a**5*b**6*d**3*e**8*x**3
- 1413720*a**5*b**6*d**2*e**9*x**4 - 3958416*a**5*b**6*d*e**10*x**5 - 8576
568*a**5*b**6*e**11*x**6 - 210*a**4*b**7*d**7*e**4 - 3780*a**4*b**7*d**6*
e**5*x - 32130*a**4*b**7*d**5*e**6*x**2 - 171360*a**4*b**7*d**4*e**7*x**3 -
642600*a**4*b**7*d**3*e**8*x**4 - 1799280*a**4*b**7*d**2*e**9*x**5 - 3898
440*a**4*b**7*d*e**10*x**6 - 6683040*a**4*b**7*e**11*x**7 - 84*a**3*b**8*d
**8*e**3 - 1512*a**3*b**8*d**7*e**4*x - 12852*a**3*b**8*d**6*e**5*x**2 - 6
8544*a**3*b**8*d**5*e**6*x**3 - 257040*a**3*b**8*d**4*e**7*x**4 - 719712*a
**3*b**8*d**3*e**8*x**5 - 1559376*a**3*b**8*d**2*e**9*x**6 - 2673216*a**3*
b**8*d*e**10*x**7 - 3675672*a**3*b**8*e**11*x**8 - 28*a**2*b**9*d**9*e**2
- 504*a**2*b**9*d**8*e**3*x - 4284*a**2*b**9*d**7*e**4*x**2 - 22848*a**...
```

$$3.98 \quad \int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{20}} dx$$

Optimal result	1020
Mathematica [B] (verified)	1021
Rubi [A] (verified)	1022
Maple [B] (verified)	1023
Fricas [B] (verification not implemented)	1024
Sympy [F(-1)]	1025
Maxima [B] (verification not implemented)	1026
Giac [B] (verification not implemented)	1027
Mupad [B] (verification not implemented)	1028
Reduce [B] (verification not implemented)	1028

Optimal result

Integrand size = 20, antiderivative size = 460

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{20}} dx = \frac{(bd-ae)^{10}(Bd-Ae)}{19e^{12}(d+ex)^{19}} - \frac{(bd-ae)^9(11bBd-10Abe-aBe)}{18e^{12}(d+ex)^{18}} + \frac{5b(bd-ae)^8(11bBd-9Abe-2aBe)}{17e^{12}(d+ex)^{17}} - \frac{15b^2(bd-ae)^7(11bBd-8Abe-3aBe)}{16e^{12}(d+ex)^{16}} + \frac{2b^3(bd-ae)^6(11bBd-7Abe-4aBe)}{e^{12}(d+ex)^{15}} - \frac{3b^4(bd-ae)^5(11bBd-6Abe-5aBe)}{e^{12}(d+ex)^{14}} + \frac{42b^5(bd-ae)^4(11bBd-5Abe-6aBe)}{13e^{12}(d+ex)^{13}} - \frac{5b^6(bd-ae)^3(11bBd-4Abe-7aBe)}{2e^{12}(d+ex)^{12}} + \frac{15b^7(bd-ae)^2(11bBd-3Abe-8aBe)}{11e^{12}(d+ex)^{11}} - \frac{b^8(bd-ae)(11bBd-2Abe-9aBe)}{2e^{12}(d+ex)^{10}} + \frac{b^9(11bBd-Abe-10aBe)}{9e^{12}(d+ex)^9} - \frac{b^{10}B}{8e^{12}(d+ex)^8}$$

output

```
1/19*(-a*e+b*d)^10*(-A*e+B*d)/e^12/(e*x+d)^19-1/18*(-a*e+b*d)^9*(-10*A*b*e
-B*a*e+11*B*b*d)/e^12/(e*x+d)^18+5/17*b*(-a*e+b*d)^8*(-9*A*b*e-2*B*a*e+11*
B*b*d)/e^12/(e*x+d)^17-15/16*b^2*(-a*e+b*d)^7*(-8*A*b*e-3*B*a*e+11*B*b*d)/
e^12/(e*x+d)^16+2*b^3*(-a*e+b*d)^6*(-7*A*b*e-4*B*a*e+11*B*b*d)/e^12/(e*x+d
)^15-3*b^4*(-a*e+b*d)^5*(-6*A*b*e-5*B*a*e+11*B*b*d)/e^12/(e*x+d)^14+42/13*
b^5*(-a*e+b*d)^4*(-5*A*b*e-6*B*a*e+11*B*b*d)/e^12/(e*x+d)^13-5/2*b^6*(-a*e
+b*d)^3*(-4*A*b*e-7*B*a*e+11*B*b*d)/e^12/(e*x+d)^12+15/11*b^7*(-a*e+b*d)^2
*(-3*A*b*e-8*B*a*e+11*B*b*d)/e^12/(e*x+d)^11-1/2*b^8*(-a*e+b*d)*(-2*A*b*e-
9*B*a*e+11*B*b*d)/e^12/(e*x+d)^10+1/9*b^9*(-A*b*e-10*B*a*e+11*B*b*d)/e^12/
(e*x+d)^9-1/8*b^10*B/e^12/(e*x+d)^8
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1433 vs. $2(460) = 920$.

Time = 0.50 (sec) , antiderivative size = 1433, normalized size of antiderivative = 3.12

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{20}} dx = \text{Too large to display}$$

input `Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^20,x]`

output

```
-1/6651216*(19448*a^10*e^10*(18*A*e + B*(d + 19*e*x)) + 11440*a^9*b*e^9*(1
7*A*e*(d + 19*e*x) + 2*B*(d^2 + 19*d*e*x + 171*e^2*x^2)) + 6435*a^8*b^2*e^
8*(16*A*e*(d^2 + 19*d*e*x + 171*e^2*x^2) + 3*B*(d^3 + 19*d^2*e*x + 171*d*e
^2*x^2 + 969*e^3*x^3)) + 3432*a^7*b^3*e^7*(15*A*e*(d^3 + 19*d^2*e*x + 171*
d*e^2*x^2 + 969*e^3*x^3) + 4*B*(d^4 + 19*d^3*e*x + 171*d^2*e^2*x^2 + 969*d
*e^3*x^3 + 3876*e^4*x^4)) + 1716*a^6*b^4*e^6*(14*A*e*(d^4 + 19*d^3*e*x + 1
71*d^2*e^2*x^2 + 969*d*e^3*x^3 + 3876*e^4*x^4) + 5*B*(d^5 + 19*d^4*e*x + 1
71*d^3*e^2*x^2 + 969*d^2*e^3*x^3 + 3876*d*e^4*x^4 + 11628*e^5*x^5)) + 792*
a^5*b^5*e^5*(13*A*e*(d^5 + 19*d^4*e*x + 171*d^3*e^2*x^2 + 969*d^2*e^3*x^3
+ 3876*d*e^4*x^4 + 11628*e^5*x^5) + 6*B*(d^6 + 19*d^5*e*x + 171*d^4*e^2*x^
2 + 969*d^3*e^3*x^3 + 3876*d^2*e^4*x^4 + 11628*d*e^5*x^5 + 27132*e^6*x^6))
+ 330*a^4*b^6*e^4*(12*A*e*(d^6 + 19*d^5*e*x + 171*d^4*e^2*x^2 + 969*d^3*e
^3*x^3 + 3876*d^2*e^4*x^4 + 11628*d*e^5*x^5 + 27132*e^6*x^6) + 7*B*(d^7 +
19*d^6*e*x + 171*d^5*e^2*x^2 + 969*d^4*e^3*x^3 + 3876*d^3*e^4*x^4 + 11628*
d^2*e^5*x^5 + 27132*d*e^6*x^6 + 50388*e^7*x^7)) + 120*a^3*b^7*e^3*(11*A*e*
(d^7 + 19*d^6*e*x + 171*d^5*e^2*x^2 + 969*d^4*e^3*x^3 + 3876*d^3*e^4*x^4 +
11628*d^2*e^5*x^5 + 27132*d*e^6*x^6 + 50388*e^7*x^7) + 8*B*(d^8 + 19*d^7*
e*x + 171*d^6*e^2*x^2 + 969*d^5*e^3*x^3 + 3876*d^4*e^4*x^4 + 11628*d^3*e^5
*x^5 + 27132*d^2*e^6*x^6 + 50388*d*e^7*x^7 + 75582*e^8*x^8)) + 36*a^2*b^8*
e^2*(10*A*e*(d^8 + 19*d^7*e*x + 171*d^6*e^2*x^2 + 969*d^5*e^3*x^3 + 387...
```

Rubi [A] (verified)

Time = 1.27 (sec) , antiderivative size = 460, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{20}} dx$$

↓ 86

$$\int \left(\frac{b^9(10aBe + Abe - 11bBd)}{e^{11}(d + ex)^{10}} - \frac{5b^8(bd - ae)(9aBe + 2Abe - 11bBd)}{e^{11}(d + ex)^{11}} + \frac{15b^7(bd - ae)^2(8aBe + 3Abe - 11bBd)}{e^{11}(d + ex)^{12}} \right) dx$$

↓ 2009

$$\begin{aligned} & \frac{b^9(-10aBe - Abe + 11bBd)}{9e^{12}(d + ex)^9} - \frac{b^8(bd - ae)(-9aBe - 2Abe + 11bBd)}{2e^{12}(d + ex)^{10}} + \\ & \frac{15b^7(bd - ae)^2(-8aBe - 3Abe + 11bBd)}{11e^{12}(d + ex)^{11}} - \frac{5b^6(bd - ae)^3(-7aBe - 4Abe + 11bBd)}{2e^{12}(d + ex)^{12}} + \\ & \frac{42b^5(bd - ae)^4(-6aBe - 5Abe + 11bBd)}{13e^{12}(d + ex)^{13}} - \frac{3b^4(bd - ae)^5(-5aBe - 6Abe + 11bBd)}{e^{12}(d + ex)^{14}} + \\ & \frac{2b^3(bd - ae)^6(-4aBe - 7Abe + 11bBd)}{e^{12}(d + ex)^{15}} - \frac{15b^2(bd - ae)^7(-3aBe - 8Abe + 11bBd)}{16e^{12}(d + ex)^{16}} + \\ & \frac{5b(bd - ae)^8(-2aBe - 9Abe + 11bBd)}{17e^{12}(d + ex)^{17}} - \frac{(bd - ae)^9(-aBe - 10Abe + 11bBd)}{18e^{12}(d + ex)^{18}} + \\ & \frac{(bd - ae)^{10}(Bd - Ae)}{19e^{12}(d + ex)^{19}} - \frac{b^{10}B}{8e^{12}(d + ex)^8} \end{aligned}$$

input

```
Int[((a + b*x)^10*(A + B*x))/(d + e*x)^20,x]
```

output

$$\begin{aligned} & ((b*d - a*e)^{10}*(B*d - A*e))/(19*e^{12}*(d + e*x)^{19}) - ((b*d - a*e)^9*(11*b \\ & *B*d - 10*A*b*e - a*B*e))/(18*e^{12}*(d + e*x)^{18}) + (5*b*(b*d - a*e)^8*(11* \\ & b*B*d - 9*A*b*e - 2*a*B*e))/(17*e^{12}*(d + e*x)^{17}) - (15*b^2*(b*d - a*e)^7 \\ & *(11*b*B*d - 8*A*b*e - 3*a*B*e))/(16*e^{12}*(d + e*x)^{16}) + (2*b^3*(b*d - a* \\ & e)^6*(11*b*B*d - 7*A*b*e - 4*a*B*e))/(e^{12}*(d + e*x)^{15}) - (3*b^4*(b*d - a \\ & *e)^5*(11*b*B*d - 6*A*b*e - 5*a*B*e))/(e^{12}*(d + e*x)^{14}) + (42*b^5*(b*d - \\ & a*e)^4*(11*b*B*d - 5*A*b*e - 6*a*B*e))/(13*e^{12}*(d + e*x)^{13}) - (5*b^6*(b \\ & *d - a*e)^3*(11*b*B*d - 4*A*b*e - 7*a*B*e))/(2*e^{12}*(d + e*x)^{12}) + (15*b^ \\ & 7*(b*d - a*e)^2*(11*b*B*d - 3*A*b*e - 8*a*B*e))/(11*e^{12}*(d + e*x)^{11}) - (\\ & b^8*(b*d - a*e)*(11*b*B*d - 2*A*b*e - 9*a*B*e))/(2*e^{12}*(d + e*x)^{10}) + (b \\ & ^9*(11*b*B*d - A*b*e - 10*a*B*e))/(9*e^{12}*(d + e*x)^9) - (b^{10}*B)/(8*e^{12}* \\ & (d + e*x)^8) \end{aligned}$$

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1900 vs. $2(440) = 880$.

Time = 0.35 (sec) , antiderivative size = 1901, normalized size of antiderivative = 4.13

method	result	size
risch	Expression too large to display	1901
default	Expression too large to display	1942
norman	Expression too large to display	2014
gospers	Expression too large to display	2233
orering	Expression too large to display	2233
parallelrisch	Expression too large to display	2242

output

```

-1/6651216*(831402*B*b^10*e^11*x^11 + 11*B*b^10*d^11 + 350064*A*a^10*e^11
+ 8*(10*B*a*b^9 + A*b^10)*d^10*e + 36*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 +
120*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 330*(7*B*a^4*b^6 + 4*A*a^3*b^7)*
d^7*e^4 + 792*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 1716*(5*B*a^6*b^4 + 6*
A*a^5*b^5)*d^5*e^6 + 3432*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 6435*(3*B*
a^8*b^2 + 8*A*a^7*b^3)*d^3*e^8 + 11440*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 +
19448*(B*a^10 + 10*A*a^9*b)*d*e^10 + 92378*(11*B*b^10*d*e^10 + 8*(10*B*a*
b^9 + A*b^10)*e^11)*x^10 + 92378*(11*B*b^10*d^2*e^9 + 8*(10*B*a*b^9 + A*b^
10)*d*e^10 + 36*(9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 + 75582*(11*B*b^10*d^3
*e^8 + 8*(10*B*a*b^9 + A*b^10)*d^2*e^9 + 36*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^
10 + 120*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 50388*(11*B*b^10*d^4*e^7
+ 8*(10*B*a*b^9 + A*b^10)*d^3*e^8 + 36*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 +
120*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 + 330*(7*B*a^4*b^6 + 4*A*a^3*b^7)*
e^11)*x^7 + 27132*(11*B*b^10*d^5*e^6 + 8*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 3
6*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 + 120*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*
e^9 + 330*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 + 792*(6*B*a^5*b^5 + 5*A*a^4*
b^6)*e^11)*x^6 + 11628*(11*B*b^10*d^6*e^5 + 8*(10*B*a*b^9 + A*b^10)*d^5*e^
6 + 36*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 + 120*(8*B*a^3*b^7 + 3*A*a^2*b^8)
*d^3*e^8 + 330*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 + 792*(6*B*a^5*b^5 + 5*
A*a^4*b^6)*d*e^10 + 1716*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 + 3876*(...

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{20}} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**10*(B*x+A)/(e*x+d)**20,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2017 vs. $2(440) = 880$.

Time = 0.15 (sec) , antiderivative size = 2017, normalized size of antiderivative = 4.38

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{20}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^20,x, algorithm="maxima")`

output

```
-1/6651216*(831402*B*b^10*e^11*x^11 + 11*B*b^10*d^11 + 350064*A*a^10*e^11
+ 8*(10*B*a*b^9 + A*b^10)*d^10*e + 36*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2 +
120*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 330*(7*B*a^4*b^6 + 4*A*a^3*b^7)*
d^7*e^4 + 792*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 1716*(5*B*a^6*b^4 + 6*
A*a^5*b^5)*d^5*e^6 + 3432*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 6435*(3*B*
a^8*b^2 + 8*A*a^7*b^3)*d^3*e^8 + 11440*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e^9 +
19448*(B*a^10 + 10*A*a^9*b)*d*e^10 + 92378*(11*B*b^10*d*e^10 + 8*(10*B*a*
b^9 + A*b^10)*e^11)*x^10 + 92378*(11*B*b^10*d^2*e^9 + 8*(10*B*a*b^9 + A*b^
10)*d*e^10 + 36*(9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 + 75582*(11*B*b^10*d^3
*e^8 + 8*(10*B*a*b^9 + A*b^10)*d^2*e^9 + 36*(9*B*a^2*b^8 + 2*A*a*b^9)*d*e^
10 + 120*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 50388*(11*B*b^10*d^4*e^7
+ 8*(10*B*a*b^9 + A*b^10)*d^3*e^8 + 36*(9*B*a^2*b^8 + 2*A*a*b^9)*d^2*e^9 +
120*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 + 330*(7*B*a^4*b^6 + 4*A*a^3*b^7)*
e^11)*x^7 + 27132*(11*B*b^10*d^5*e^6 + 8*(10*B*a*b^9 + A*b^10)*d^4*e^7 + 3
6*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 + 120*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^2*
e^9 + 330*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 + 792*(6*B*a^5*b^5 + 5*A*a^4*
b^6)*e^11)*x^6 + 11628*(11*B*b^10*d^6*e^5 + 8*(10*B*a*b^9 + A*b^10)*d^5*e^
6 + 36*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 + 120*(8*B*a^3*b^7 + 3*A*a^2*b^8)
*d^3*e^8 + 330*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 + 792*(6*B*a^5*b^5 + 5*
A*a^4*b^6)*d*e^10 + 1716*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5 + 3876*(...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2232 vs. $2(440) = 880$.

Time = 0.13 (sec) , antiderivative size = 2232, normalized size of antiderivative = 4.85

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{20}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^20,x, algorithm="giac")`

output

```
-1/6651216*(831402*B*b^10*e^11*x^11 + 1016158*B*b^10*d*e^10*x^10 + 7390240
*B*a*b^9*e^11*x^10 + 739024*A*b^10*e^11*x^10 + 1016158*B*b^10*d^2*e^9*x^9
+ 7390240*B*a*b^9*d*e^10*x^9 + 739024*A*b^10*d*e^10*x^9 + 29930472*B*a^2*b
^8*e^11*x^9 + 6651216*A*a*b^9*e^11*x^9 + 831402*B*b^10*d^3*e^8*x^8 + 60465
60*B*a*b^9*d^2*e^9*x^8 + 604656*A*b^10*d^2*e^9*x^8 + 24488568*B*a^2*b^8*d*
e^10*x^8 + 5441904*A*a*b^9*d*e^10*x^8 + 72558720*B*a^3*b^7*e^11*x^8 + 2720
9520*A*a^2*b^8*e^11*x^8 + 554268*B*b^10*d^4*e^7*x^7 + 4031040*B*a*b^9*d^3*
e^8*x^7 + 403104*A*b^10*d^3*e^8*x^7 + 16325712*B*a^2*b^8*d^2*e^9*x^7 + 362
7936*A*a*b^9*d^2*e^9*x^7 + 48372480*B*a^3*b^7*d*e^10*x^7 + 18139680*A*a^2*
b^8*d*e^10*x^7 + 116396280*B*a^4*b^6*e^11*x^7 + 66512160*A*a^3*b^7*e^11*x^
7 + 298452*B*b^10*d^5*e^6*x^6 + 2170560*B*a*b^9*d^4*e^7*x^6 + 217056*A*b^1
0*d^4*e^7*x^6 + 8790768*B*a^2*b^8*d^3*e^8*x^6 + 1953504*A*a*b^9*d^3*e^8*x^
6 + 26046720*B*a^3*b^7*d^2*e^9*x^6 + 9767520*A*a^2*b^8*d^2*e^9*x^6 + 62674
920*B*a^4*b^6*d*e^10*x^6 + 35814240*A*a^3*b^7*d*e^10*x^6 + 128931264*B*a^5
*b^5*e^11*x^6 + 107442720*A*a^4*b^6*e^11*x^6 + 127908*B*b^10*d^6*e^5*x^5 +
930240*B*a*b^9*d^5*e^6*x^5 + 93024*A*b^10*d^5*e^6*x^5 + 3767472*B*a^2*b^8
*d^4*e^7*x^5 + 837216*A*a*b^9*d^4*e^7*x^5 + 11162880*B*a^3*b^7*d^3*e^8*x^5
+ 4186080*A*a^2*b^8*d^3*e^8*x^5 + 26860680*B*a^4*b^6*d^2*e^9*x^5 + 153489
60*A*a^3*b^7*d^2*e^9*x^5 + 55256256*B*a^5*b^5*d*e^10*x^5 + 46046880*A*a^4*
b^6*d*e^10*x^5 + 99768240*B*a^6*b^4*e^11*x^5 + 119721888*A*a^5*b^5*e^11...
```


Mupad [B] (verification not implemented)

Time = 17.87 (sec) , antiderivative size = 2099, normalized size of antiderivative = 4.56

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{20}} dx = \text{Too large to display}$$

input `int(((A + B*x)*(a + b*x)^10)/(d + e*x)^20,x)`

output

```

-((350064*A*a^10*e^11 + 11*B*b^10*d^11 + 8*A*b^10*d^10*e + 19448*B*a^10*d*
e^10 + 72*A*a*b^9*d^9*e^2 + 22880*B*a^9*b*d^2*e^9 + 360*A*a^2*b^8*d^8*e^3
+ 1320*A*a^3*b^7*d^7*e^4 + 3960*A*a^4*b^6*d^6*e^5 + 10296*A*a^5*b^5*d^5*e^
6 + 24024*A*a^6*b^4*d^4*e^7 + 51480*A*a^7*b^3*d^3*e^8 + 102960*A*a^8*b^2*d
^2*e^9 + 324*B*a^2*b^8*d^9*e^2 + 960*B*a^3*b^7*d^8*e^3 + 2310*B*a^4*b^6*d^
7*e^4 + 4752*B*a^5*b^5*d^6*e^5 + 8580*B*a^6*b^4*d^5*e^6 + 13728*B*a^7*b^3*
d^4*e^7 + 19305*B*a^8*b^2*d^3*e^8 + 194480*A*a^9*b*d*e^10 + 80*B*a*b^9*d^1
0*e)/(6651216*e^12) + (x*(19448*B*a^10*e^10 + 11*B*b^10*d^10 + 194480*A*a^
9*b*e^10 + 8*A*b^10*d^9*e + 72*A*a*b^9*d^8*e^2 + 102960*A*a^8*b^2*d*e^9 +
360*A*a^2*b^8*d^7*e^3 + 1320*A*a^3*b^7*d^6*e^4 + 3960*A*a^4*b^6*d^5*e^5 +
10296*A*a^5*b^5*d^4*e^6 + 24024*A*a^6*b^4*d^3*e^7 + 51480*A*a^7*b^3*d^2*e^
8 + 324*B*a^2*b^8*d^8*e^2 + 960*B*a^3*b^7*d^7*e^3 + 2310*B*a^4*b^6*d^6*e^4
+ 4752*B*a^5*b^5*d^5*e^5 + 8580*B*a^6*b^4*d^4*e^6 + 13728*B*a^7*b^3*d^3*e
^7 + 19305*B*a^8*b^2*d^2*e^8 + 80*B*a*b^9*d^9*e + 22880*B*a^9*b*d*e^9))/(3
50064*e^11) + (b^7*x^8*(960*B*a^3*e^3 + 11*B*b^3*d^3 + 360*A*a^2*b*e^3 + 8
*A*b^3*d^2*e + 72*A*a*b^2*d*e^2 + 80*B*a*b^2*d^2*e + 324*B*a^2*b*d*e^2))/(
88*e^4) + (b^4*x^5*(8580*B*a^6*e^6 + 11*B*b^6*d^6 + 10296*A*a^5*b*e^6 + 8*
A*b^6*d^5*e + 72*A*a*b^5*d^4*e^2 + 3960*A*a^4*b^2*d*e^5 + 360*A*a^2*b^4*d^
3*e^3 + 1320*A*a^3*b^3*d^2*e^4 + 324*B*a^2*b^4*d^4*e^2 + 960*B*a^3*b^3*d^3
*e^3 + 2310*B*a^4*b^2*d^2*e^4 + 80*B*a*b^5*d^5*e + 4752*B*a^5*b*d*e^5))...

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 1349, normalized size of antiderivative = 2.93

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{20}} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^20,x)`

output

```
( - 31824*a**11*e**11 - 19448*a**10*b*d*e**10 - 369512*a**10*b*e**11*x - 1
1440*a**9*b**2*d**2*e**9 - 217360*a**9*b**2*d*e**10*x - 1956240*a**9*b**2*
e**11*x**2 - 6435*a**8*b**3*d**3*e**8 - 122265*a**8*b**3*d**2*e**9*x - 110
0385*a**8*b**3*d*e**10*x**2 - 6235515*a**8*b**3*e**11*x**3 - 3432*a**7*b**
4*d**4*e**7 - 65208*a**7*b**4*d**3*e**8*x - 586872*a**7*b**4*d**2*e**9*x**
2 - 3325608*a**7*b**4*d*e**10*x**3 - 13302432*a**7*b**4*e**11*x**4 - 1716*
a**6*b**5*d**5*e**6 - 32604*a**6*b**5*d**4*e**7*x - 293436*a**6*b**5*d**3*
e**8*x**2 - 1662804*a**6*b**5*d**2*e**9*x**3 - 6651216*a**6*b**5*d*e**10*x
**4 - 19953648*a**6*b**5*e**11*x**5 - 792*a**5*b**6*d**6*e**5 - 15048*a**5
*b**6*d**5*e**6*x - 135432*a**5*b**6*d**4*e**7*x**2 - 767448*a**5*b**6*d**
3*e**8*x**3 - 3069792*a**5*b**6*d**2*e**9*x**4 - 9209376*a**5*b**6*d*e**10
*x**5 - 21488544*a**5*b**6*e**11*x**6 - 330*a**4*b**7*d**7*e**4 - 6270*a**
4*b**7*d**6*e**5*x - 56430*a**4*b**7*d**5*e**6*x**2 - 319770*a**4*b**7*d**
4*e**7*x**3 - 1279080*a**4*b**7*d**3*e**8*x**4 - 3837240*a**4*b**7*d**2*e
**9*x**5 - 8953560*a**4*b**7*d*e**10*x**6 - 16628040*a**4*b**7*e**11*x**7 -
120*a**3*b**8*d**8*e**3 - 2280*a**3*b**8*d**7*e**4*x - 20520*a**3*b**8*d
**6*e**5*x**2 - 116280*a**3*b**8*d**5*e**6*x**3 - 465120*a**3*b**8*d**4*e**
7*x**4 - 1395360*a**3*b**8*d**3*e**8*x**5 - 3255840*a**3*b**8*d**2*e**9*x
**6 - 6046560*a**3*b**8*d*e**10*x**7 - 9069840*a**3*b**8*e**11*x**8 - 36*a
**2*b**9*d**9*e**2 - 684*a**2*b**9*d**8*e**3*x - 6156*a**2*b**9*d**7*e**...
```

$$\mathbf{3.99} \quad \int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{21}} dx$$

Optimal result	1031
Mathematica [B] (verified)	1032
Rubi [A] (verified)	1033
Maple [B] (verified)	1034
Fricas [B] (verification not implemented)	1035
Sympy [F(-1)]	1036
Maxima [B] (verification not implemented)	1037
Giac [B] (verification not implemented)	1038
Mupad [B] (verification not implemented)	1039
Reduce [B] (verification not implemented)	1039

Optimal result

Integrand size = 20, antiderivative size = 462

$$\begin{aligned}
 \int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{21}} dx = & \frac{(bd-ae)^{10}(Bd-Ae)}{20e^{12}(d+ex)^{20}} \\
 & - \frac{(bd-ae)^9(11bBd-10Abe-aBe)}{19e^{12}(d+ex)^{19}} \\
 & + \frac{5b(bd-ae)^8(11bBd-9Abe-2aBe)}{18e^{12}(d+ex)^{18}} \\
 & - \frac{15b^2(bd-ae)^7(11bBd-8Abe-3aBe)}{17e^{12}(d+ex)^{17}} \\
 & + \frac{15b^3(bd-ae)^6(11bBd-7Abe-4aBe)}{8e^{12}(d+ex)^{16}} \\
 & - \frac{14b^4(bd-ae)^5(11bBd-6Abe-5aBe)}{5e^{12}(d+ex)^{15}} \\
 & + \frac{3b^5(bd-ae)^4(11bBd-5Abe-6aBe)}{e^{12}(d+ex)^{14}} \\
 & - \frac{30b^6(bd-ae)^3(11bBd-4Abe-7aBe)}{13e^{12}(d+ex)^{13}} \\
 & + \frac{5b^7(bd-ae)^2(11bBd-3Abe-8aBe)}{4e^{12}(d+ex)^{12}} \\
 & - \frac{5b^8(bd-ae)(11bBd-2Abe-9aBe)}{11e^{12}(d+ex)^{11}} \\
 & + \frac{b^9(11bBd-Abe-10aBe)}{10e^{12}(d+ex)^{10}} - \frac{b^{10}B}{9e^{12}(d+ex)^9}
 \end{aligned}$$

output

```

1/20*(-a*e+b*d)^10*(-A*e+B*d)/e^12/(e*x+d)^20-1/19*(-a*e+b*d)^9*(-10*A*b*e
-B*a*e+11*B*b*d)/e^12/(e*x+d)^19+5/18*b*(-a*e+b*d)^8*(-9*A*b*e-2*B*a*e+11*
B*b*d)/e^12/(e*x+d)^18-15/17*b^2*(-a*e+b*d)^7*(-8*A*b*e-3*B*a*e+11*B*b*d)/
e^12/(e*x+d)^17+15/8*b^3*(-a*e+b*d)^6*(-7*A*b*e-4*B*a*e+11*B*b*d)/e^12/(e*
x+d)^16-14/5*b^4*(-a*e+b*d)^5*(-6*A*b*e-5*B*a*e+11*B*b*d)/e^12/(e*x+d)^15+
3*b^5*(-a*e+b*d)^4*(-5*A*b*e-6*B*a*e+11*B*b*d)/e^12/(e*x+d)^14-30/13*b^6*(
-a*e+b*d)^3*(-4*A*b*e-7*B*a*e+11*B*b*d)/e^12/(e*x+d)^13+5/4*b^7*(-a*e+b*d)
^2*(-3*A*b*e-8*B*a*e+11*B*b*d)/e^12/(e*x+d)^12-5/11*b^8*(-a*e+b*d)*(-2*A*b
*e-9*B*a*e+11*B*b*d)/e^12/(e*x+d)^11+1/10*b^9*(-A*b*e-10*B*a*e+11*B*b*d)/e
^12/(e*x+d)^10-1/9*b^10*B/e^12/(e*x+d)^9

```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1428 vs. $2(462) = 924$.

Time = 0.52 (sec) , antiderivative size = 1428, normalized size of antiderivative = 3.09

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{21}} dx = \text{Too large to display}$$

input `Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^21,x]`

output

```
-1/16628040*(43758*a^10*e^10*(19*A*e + B*(d + 20*e*x)) + 48620*a^9*b*e^9*(
9*A*e*(d + 20*e*x) + B*(d^2 + 20*d*e*x + 190*e^2*x^2)) + 12870*a^8*b^2*e^8
*(17*A*e*(d^2 + 20*d*e*x + 190*e^2*x^2) + 3*B*(d^3 + 20*d^2*e*x + 190*d*e^
2*x^2 + 1140*e^3*x^3)) + 25740*a^7*b^3*e^7*(4*A*e*(d^3 + 20*d^2*e*x + 190*
d*e^2*x^2 + 1140*e^3*x^3) + B*(d^4 + 20*d^3*e*x + 190*d^2*e^2*x^2 + 1140*d
*e^3*x^3 + 4845*e^4*x^4)) + 15015*a^6*b^4*e^6*(3*A*e*(d^4 + 20*d^3*e*x + 1
90*d^2*e^2*x^2 + 1140*d*e^3*x^3 + 4845*e^4*x^4) + B*(d^5 + 20*d^4*e*x + 19
0*d^3*e^2*x^2 + 1140*d^2*e^3*x^3 + 4845*d*e^4*x^4 + 15504*e^5*x^5)) + 2574
*a^5*b^5*e^5*(7*A*e*(d^5 + 20*d^4*e*x + 190*d^3*e^2*x^2 + 1140*d^2*e^3*x^3
+ 4845*d*e^4*x^4 + 15504*e^5*x^5) + 3*B*(d^6 + 20*d^5*e*x + 190*d^4*e^2*x
^2 + 1140*d^3*e^3*x^3 + 4845*d^2*e^4*x^4 + 15504*d*e^5*x^5 + 38760*e^6*x^6
)) + 495*a^4*b^6*e^4*(13*A*e*(d^6 + 20*d^5*e*x + 190*d^4*e^2*x^2 + 1140*d^
3*e^3*x^3 + 4845*d^2*e^4*x^4 + 15504*d*e^5*x^5 + 38760*e^6*x^6) + 7*B*(d^7
+ 20*d^6*e*x + 190*d^5*e^2*x^2 + 1140*d^4*e^3*x^3 + 4845*d^3*e^4*x^4 + 15
504*d^2*e^5*x^5 + 38760*d*e^6*x^6 + 77520*e^7*x^7)) + 660*a^3*b^7*e^3*(3*A
*e*(d^7 + 20*d^6*e*x + 190*d^5*e^2*x^2 + 1140*d^4*e^3*x^3 + 4845*d^3*e^4*x
^4 + 15504*d^2*e^5*x^5 + 38760*d*e^6*x^6 + 77520*e^7*x^7) + 2*B*(d^8 + 20*
d^7*e*x + 190*d^6*e^2*x^2 + 1140*d^5*e^3*x^3 + 4845*d^4*e^4*x^4 + 15504*d^
3*e^5*x^5 + 38760*d^2*e^6*x^6 + 77520*d*e^7*x^7 + 125970*e^8*x^8)) + 45*a^
2*b^8*e^2*(11*A*e*(d^8 + 20*d^7*e*x + 190*d^6*e^2*x^2 + 1140*d^5*e^3*x^...
```

Rubi [A] (verified)

Time = 1.21 (sec) , antiderivative size = 462, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{21}} dx$$

↓ 86

$$\int \left(\frac{b^9(10aBe + Abe - 11bBd)}{e^{11}(d + ex)^{11}} - \frac{5b^8(bd - ae)(9aBe + 2Abe - 11bBd)}{e^{11}(d + ex)^{12}} + \frac{15b^7(bd - ae)^2(8aBe + 3Abe - 11bBd)}{e^{11}(d + ex)^{13}} \right) dx$$

↓ 2009

$$\frac{b^9(-10aBe - Abe + 11bBd)}{10e^{12}(d + ex)^{10}} - \frac{5b^8(bd - ae)(-9aBe - 2Abe + 11bBd)}{11e^{12}(d + ex)^{11}} + \frac{5b^7(bd - ae)^2(-8aBe - 3Abe + 11bBd)}{4e^{12}(d + ex)^{12}} - \frac{30b^6(bd - ae)^3(-7aBe - 4Abe + 11bBd)}{13e^{12}(d + ex)^{13}} + \frac{3b^5(bd - ae)^4(-6aBe - 5Abe + 11bBd)}{e^{12}(d + ex)^{14}} - \frac{14b^4(bd - ae)^5(-5aBe - 6Abe + 11bBd)}{5e^{12}(d + ex)^{15}} + \frac{15b^3(bd - ae)^6(-4aBe - 7Abe + 11bBd)}{8e^{12}(d + ex)^{16}} - \frac{15b^2(bd - ae)^7(-3aBe - 8Abe + 11bBd)}{17e^{12}(d + ex)^{17}} + \frac{5b(bd - ae)^8(-2aBe - 9Abe + 11bBd)}{18e^{12}(d + ex)^{18}} - \frac{(bd - ae)^9(-aBe - 10Abe + 11bBd)}{19e^{12}(d + ex)^{19}} + \frac{(bd - ae)^{10}(Bd - Ae)}{20e^{12}(d + ex)^{20}} - \frac{b^{10}B}{9e^{12}(d + ex)^9}$$

input

```
Int[((a + b*x)^10*(A + B*x))/(d + e*x)^21,x]
```

output

$$\begin{aligned} & ((b*d - a*e)^{10}*(B*d - A*e))/(20*e^{12}*(d + e*x)^{20}) - ((b*d - a*e)^9*(11*b \\ & *B*d - 10*A*b*e - a*B*e))/(19*e^{12}*(d + e*x)^{19}) + (5*b*(b*d - a*e)^8*(11* \\ & b*B*d - 9*A*b*e - 2*a*B*e))/(18*e^{12}*(d + e*x)^{18}) - (15*b^2*(b*d - a*e)^7 \\ & *(11*b*B*d - 8*A*b*e - 3*a*B*e))/(17*e^{12}*(d + e*x)^{17}) + (15*b^3*(b*d - a \\ & *e)^6*(11*b*B*d - 7*A*b*e - 4*a*B*e))/(8*e^{12}*(d + e*x)^{16}) - (14*b^4*(b*d \\ & - a*e)^5*(11*b*B*d - 6*A*b*e - 5*a*B*e))/(5*e^{12}*(d + e*x)^{15}) + (3*b^5*(\\ & b*d - a*e)^4*(11*b*B*d - 5*A*b*e - 6*a*B*e))/(e^{12}*(d + e*x)^{14}) - (30*b^6 \\ & *(b*d - a*e)^3*(11*b*B*d - 4*A*b*e - 7*a*B*e))/(13*e^{12}*(d + e*x)^{13}) + (5 \\ & *b^7*(b*d - a*e)^2*(11*b*B*d - 3*A*b*e - 8*a*B*e))/(4*e^{12}*(d + e*x)^{12}) - \\ & (5*b^8*(b*d - a*e)*(11*b*B*d - 2*A*b*e - 9*a*B*e))/(11*e^{12}*(d + e*x)^{11}) \\ & + (b^9*(11*b*B*d - A*b*e - 10*a*B*e))/(10*e^{12}*(d + e*x)^{10}) - (b^{10}*B)/(\\ & 9*e^{12}*(d + e*x)^9) \end{aligned}$$

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1900 vs. $2(440) = 880$.

Time = 0.37 (sec) , antiderivative size = 1901, normalized size of antiderivative = 4.11

method	result	size
risch	Expression too large to display	1901
default	Expression too large to display	1942
norman	Expression too large to display	2014
gospers	Expression too large to display	2233
orering	Expression too large to display	2233
parallelrisch	Expression too large to display	2242

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^21,x,method=_RETURNVERBOSE)`

output `(-1/16628040/e^12*(831402*A*a^10*e^11+437580*A*a^9*b*d*e^10+218790*A*a^8*b^2*d^2*e^9+102960*A*a^7*b^3*d^3*e^8+45045*A*a^6*b^4*d^4*e^7+18018*A*a^5*b^5*d^5*e^6+6435*A*a^4*b^6*d^6*e^5+1980*A*a^3*b^7*d^7*e^4+495*A*a^2*b^8*d^8*e^3+90*A*a*b^9*d^9*e^2+9*A*b^10*d^10*e+43758*B*a^10*d*e^10+48620*B*a^9*b*d^2*e^9+38610*B*a^8*b^2*d^3*e^8+25740*B*a^7*b^3*d^4*e^7+15015*B*a^6*b^4*d^5*e^6+7722*B*a^5*b^5*d^6*e^5+3465*B*a^4*b^6*d^7*e^4+1320*B*a^3*b^7*d^8*e^3+405*B*a^2*b^8*d^9*e^2+90*B*a*b^9*d^10*e+11*B*b^10*d^11)-1/831402/e^11*(437580*A*a^9*b*e^10+218790*A*a^8*b^2*d*e^9+102960*A*a^7*b^3*d^2*e^8+45045*A*a^6*b^4*d^3*e^7+18018*A*a^5*b^5*d^4*e^6+6435*A*a^4*b^6*d^5*e^5+1980*A*a^3*b^7*d^6*e^4+495*A*a^2*b^8*d^7*e^3+90*A*a*b^9*d^8*e^2+9*A*b^10*d^9*e+43758*B*a^10*e^10+48620*B*a^9*b*d*e^9+38610*B*a^8*b^2*d^2*e^8+25740*B*a^7*b^3*d^3*e^7+15015*B*a^6*b^4*d^4*e^6+7722*B*a^5*b^5*d^5*e^5+3465*B*a^4*b^6*d^6*e^4+1320*B*a^3*b^7*d^7*e^3+405*B*a^2*b^8*d^8*e^2+90*B*a*b^9*d^9*e+11*B*b^10*d^10)*x-1/87516*b/e^10*(218790*A*a^8*b*e^9+102960*A*a^7*b^2*d*e^8+45045*A*a^6*b^3*d^2*e^7+18018*A*a^5*b^4*d^3*e^6+6435*A*a^4*b^5*d^4*e^5+1980*A*a^3*b^6*d^5*e^4+495*A*a^2*b^7*d^6*e^3+90*A*a*b^8*d^7*e^2+9*A*b^9*d^8*e+48620*B*a^9*e^9+38610*B*a^8*b*d*e^8+25740*B*a^7*b^2*d^2*e^7+15015*B*a^6*b^3*d^3*e^6+7722*B*a^5*b^4*d^4*e^5+3465*B*a^4*b^5*d^5*e^4+1320*B*a^3*b^6*d^6*e^3+405*B*a^2*b^7*d^7*e^2+90*B*a*b^8*d^8*e+11*B*b^9*d^9)*x^2-1/14586*b^2/e^9*(102960*A*a^7*b*e^8+45045*A*a^6*b^2*d*e^7+18018*A*a^5*b^3*d^2*e^6+6435*A*a^...`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2028 vs. $2(440) = 880$.

Time = 0.25 (sec) , antiderivative size = 2028, normalized size of antiderivative = 4.39

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{21}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^21,x, algorithm="fricas")`

output

```
-1/16628040*(1847560*B*b^10*e^11*x^11 + 11*B*b^10*d^11 + 831402*A*a^10*e^11
+ 9*(10*B*a*b^9 + A*b^10)*d^10*e + 45*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2
+ 165*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 495*(7*B*a^4*b^6 + 4*A*a^3*b^7
)*d^7*e^4 + 1287*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 3003*(5*B*a^6*b^4 +
6*A*a^5*b^5)*d^5*e^6 + 6435*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 12870*(
3*B*a^8*b^2 + 8*A*a^7*b^3)*d^3*e^8 + 24310*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*e
^9 + 43758*(B*a^10 + 10*A*a^9*b)*d*e^10 + 184756*(11*B*b^10*d*e^10 + 9*(10
*B*a*b^9 + A*b^10)*e^11)*x^10 + 167960*(11*B*b^10*d^2*e^9 + 9*(10*B*a*b^9
+ A*b^10)*d*e^10 + 45*(9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 + 125970*(11*B*b
^10*d^3*e^8 + 9*(10*B*a*b^9 + A*b^10)*d^2*e^9 + 45*(9*B*a^2*b^8 + 2*A*a*b^
9)*d*e^10 + 165*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 77520*(11*B*b^10*d
^4*e^7 + 9*(10*B*a*b^9 + A*b^10)*d^3*e^8 + 45*(9*B*a^2*b^8 + 2*A*a*b^9)*d^
2*e^9 + 165*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 + 495*(7*B*a^4*b^6 + 4*A*a^
3*b^7)*e^11)*x^7 + 38760*(11*B*b^10*d^5*e^6 + 9*(10*B*a*b^9 + A*b^10)*d^4*
e^7 + 45*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 + 165*(8*B*a^3*b^7 + 3*A*a^2*b^
8)*d^2*e^9 + 495*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 + 1287*(6*B*a^5*b^5 +
5*A*a^4*b^6)*e^11)*x^6 + 15504*(11*B*b^10*d^6*e^5 + 9*(10*B*a*b^9 + A*b^10
)*d^5*e^6 + 45*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 + 165*(8*B*a^3*b^7 + 3*A*
a^2*b^8)*d^3*e^8 + 495*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 + 1287*(6*B*a^5
*b^5 + 5*A*a^4*b^6)*d*e^10 + 3003*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5...
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{21}} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**10*(B*x+A)/(e*x+d)**21,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2028 vs. $2(440) = 880$.

Time = 0.15 (sec) , antiderivative size = 2028, normalized size of antiderivative = 4.39

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{21}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^21,x, algorithm="maxima")`

output

```
-1/16628040*(1847560*B*b^10*e^11*x^11 + 11*B*b^10*d^11 + 831402*A*a^10*e^11
+ 9*(10*B*a*b^9 + A*b^10)*d^10*e + 45*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^2
+ 165*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 495*(7*B*a^4*b^6 + 4*A*a^3*b^7
)*d^7*e^4 + 1287*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 3003*(5*B*a^6*b^4 +
6*A*a^5*b^5)*d^5*e^6 + 6435*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 12870*(
3*B*a^8*b^2 + 8*A*a^7*b^3)*d^3*e^8 + 24310*(2*B*a^9*b + 9*A*a^8*b^2)*d^2*
e^9 + 43758*(B*a^10 + 10*A*a^9*b)*d*e^10 + 184756*(11*B*b^10*d^2*e^9 + 9*(10
*B*a*b^9 + A*b^10)*e^11)*x^10 + 167960*(11*B*b^10*d^2*e^9 + 9*(10*B*a*b^9
+ A*b^10)*d*e^10 + 45*(9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 + 125970*(11*B*b
^10*d^3*e^8 + 9*(10*B*a*b^9 + A*b^10)*d^2*e^9 + 45*(9*B*a^2*b^8 + 2*A*a*b^
9)*d*e^10 + 165*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 77520*(11*B*b^10*d
^4*e^7 + 9*(10*B*a*b^9 + A*b^10)*d^3*e^8 + 45*(9*B*a^2*b^8 + 2*A*a*b^9)*d^
2*e^9 + 165*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 + 495*(7*B*a^4*b^6 + 4*A*a^
3*b^7)*e^11)*x^7 + 38760*(11*B*b^10*d^5*e^6 + 9*(10*B*a*b^9 + A*b^10)*d^4*
e^7 + 45*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 + 165*(8*B*a^3*b^7 + 3*A*a^2*b^
8)*d^2*e^9 + 495*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 + 1287*(6*B*a^5*b^5 +
5*A*a^4*b^6)*e^11)*x^6 + 15504*(11*B*b^10*d^6*e^5 + 9*(10*B*a*b^9 + A*b^10
)*d^5*e^6 + 45*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 + 165*(8*B*a^3*b^7 + 3*A*
a^2*b^8)*d^3*e^8 + 495*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 + 1287*(6*B*a^5
*b^5 + 5*A*a^4*b^6)*d*e^10 + 3003*(5*B*a^6*b^4 + 6*A*a^5*b^5)*e^11)*x^5...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2232 vs. $2(440) = 880$.

Time = 0.14 (sec) , antiderivative size = 2232, normalized size of antiderivative = 4.83

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{21}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^21,x, algorithm="giac")`

output

```
-1/16628040*(1847560*B*b^10*e^11*x^11 + 2032316*B*b^10*d*e^10*x^10 + 16628040*B*a*b^9*e^11*x^10 + 1662804*A*b^10*e^11*x^10 + 1847560*B*b^10*d^2*e^9*x^9 + 15116400*B*a*b^9*d*e^10*x^9 + 1511640*A*b^10*d*e^10*x^9 + 68023800*B*a^2*b^8*e^11*x^9 + 15116400*A*a*b^9*e^11*x^9 + 1385670*B*b^10*d^3*e^8*x^8 + 11337300*B*a*b^9*d^2*e^9*x^8 + 1133730*A*b^10*d^2*e^9*x^8 + 51017850*B*a^2*b^8*d*e^10*x^8 + 11337300*A*a*b^9*d*e^10*x^8 + 166280400*B*a^3*b^7*e^11*x^8 + 62355150*A*a^2*b^8*e^11*x^8 + 852720*B*b^10*d^4*e^7*x^7 + 6976800*B*a*b^9*d^3*e^8*x^7 + 697680*A*b^10*d^3*e^8*x^7 + 31395600*B*a^2*b^8*d^2*e^9*x^7 + 6976800*A*a*b^9*d^2*e^9*x^7 + 102326400*B*a^3*b^7*d*e^10*x^7 + 38372400*A*a^2*b^8*d*e^10*x^7 + 268606800*B*a^4*b^6*e^11*x^7 + 153489600*A*a^3*b^7*e^11*x^7 + 426360*B*b^10*d^5*e^6*x^6 + 3488400*B*a*b^9*d^4*e^7*x^6 + 348840*A*b^10*d^4*e^7*x^6 + 15697800*B*a^2*b^8*d^3*e^8*x^6 + 3488400*A*a*b^9*d^3*e^8*x^6 + 51163200*B*a^3*b^7*d^2*e^9*x^6 + 19186200*A*a^2*b^8*d^2*e^9*x^6 + 134303400*B*a^4*b^6*d*e^10*x^6 + 76744800*A*a^3*b^7*d*e^10*x^6 + 299304720*B*a^5*b^5*e^11*x^6 + 249420600*A*a^4*b^6*e^11*x^6 + 170544*B*b^10*d^6*e^5*x^5 + 1395360*B*a*b^9*d^5*e^6*x^5 + 139536*A*b^10*d^5*e^6*x^5 + 6279120*B*a^2*b^8*d^4*e^7*x^5 + 1395360*A*a*b^9*d^4*e^7*x^5 + 20465280*B*a^3*b^7*d^3*e^8*x^5 + 7674480*A*a^2*b^8*d^3*e^8*x^5 + 53721360*B*a^4*b^6*d^2*e^9*x^5 + 30697920*A*a^3*b^7*d^2*e^9*x^5 + 119721888*B*a^5*b^5*d*e^10*x^5 + 99768240*A*a^4*b^6*d*e^10*x^5 + 232792560*B*a^6*b^4*e^11*x^5 + 27...
```

Mupad [B] (verification not implemented)

Time = 1.46 (sec) , antiderivative size = 2110, normalized size of antiderivative = 4.57

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{21}} dx = \text{Too large to display}$$

input `int(((A + B*x)*(a + b*x)^10)/(d + e*x)^21,x)`

output

```

-((831402*A*a^10*e^11 + 11*B*b^10*d^11 + 9*A*b^10*d^10*e + 43758*B*a^10*d*
e^10 + 90*A*a*b^9*d^9*e^2 + 48620*B*a^9*b*d^2*e^9 + 495*A*a^2*b^8*d^8*e^3
+ 1980*A*a^3*b^7*d^7*e^4 + 6435*A*a^4*b^6*d^6*e^5 + 18018*A*a^5*b^5*d^5*e^
6 + 45045*A*a^6*b^4*d^4*e^7 + 102960*A*a^7*b^3*d^3*e^8 + 218790*A*a^8*b^2*
d^2*e^9 + 405*B*a^2*b^8*d^9*e^2 + 1320*B*a^3*b^7*d^8*e^3 + 3465*B*a^4*b^6*
d^7*e^4 + 7722*B*a^5*b^5*d^6*e^5 + 15015*B*a^6*b^4*d^5*e^6 + 25740*B*a^7*b
^3*d^4*e^7 + 38610*B*a^8*b^2*d^3*e^8 + 437580*A*a^9*b*d*e^10 + 90*B*a*b^9*
d^10*e)/(16628040*e^12) + (x*(43758*B*a^10*e^10 + 11*B*b^10*d^10 + 437580*
A*a^9*b*e^10 + 9*A*b^10*d^9*e + 90*A*a*b^9*d^8*e^2 + 218790*A*a^8*b^2*d*e^
9 + 495*A*a^2*b^8*d^7*e^3 + 1980*A*a^3*b^7*d^6*e^4 + 6435*A*a^4*b^6*d^5*e^
5 + 18018*A*a^5*b^5*d^4*e^6 + 45045*A*a^6*b^4*d^3*e^7 + 102960*A*a^7*b^3*d
^2*e^8 + 405*B*a^2*b^8*d^8*e^2 + 1320*B*a^3*b^7*d^7*e^3 + 3465*B*a^4*b^6*d
^6*e^4 + 7722*B*a^5*b^5*d^5*e^5 + 15015*B*a^6*b^4*d^4*e^6 + 25740*B*a^7*b^
3*d^3*e^7 + 38610*B*a^8*b^2*d^2*e^8 + 90*B*a*b^9*d^9*e + 48620*B*a^9*b*d*e
^9))/(831402*e^11) + (b^7*x^8*(1320*B*a^3*e^3 + 11*B*b^3*d^3 + 495*A*a^2*b
*e^3 + 9*A*b^3*d^2*e + 90*A*a*b^2*d*e^2 + 90*B*a*b^2*d^2*e + 405*B*a^2*b*d
*e^2))/(132*e^4) + (2*b^4*x^5*(15015*B*a^6*e^6 + 11*B*b^6*d^6 + 18018*A*a^
5*b*e^6 + 9*A*b^6*d^5*e + 90*A*a*b^5*d^4*e^2 + 6435*A*a^4*b^2*d*e^5 + 495*
A*a^2*b^4*d^3*e^3 + 1980*A*a^3*b^3*d^2*e^4 + 405*B*a^2*b^4*d^4*e^2 + 1320*
B*a^3*b^3*d^3*e^3 + 3465*B*a^4*b^2*d^2*e^4 + 90*B*a*b^5*d^5*e + 7722*B*...

```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 1360, normalized size of antiderivative = 2.94

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{21}} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^21,x)`

output

```
( - 75582*a**11*e**11 - 43758*a**10*b*d*e**10 - 875160*a**10*b*e**11*x - 2
4310*a**9*b**2*d**2*e**9 - 486200*a**9*b**2*d*e**10*x - 4618900*a**9*b**2*
e**11*x**2 - 12870*a**8*b**3*d**3*e**8 - 257400*a**8*b**3*d**2*e**9*x - 24
45300*a**8*b**3*d*e**10*x**2 - 14671800*a**8*b**3*e**11*x**3 - 6435*a**7*b
**4*d**4*e**7 - 128700*a**7*b**4*d**3*e**8*x - 1222650*a**7*b**4*d**2*e**9
*x**2 - 7335900*a**7*b**4*d*e**10*x**3 - 31177575*a**7*b**4*e**11*x**4 - 3
003*a**6*b**5*d**5*e**6 - 60060*a**6*b**5*d**4*e**7*x - 570570*a**6*b**5*d
**3*e**8*x**2 - 3423420*a**6*b**5*d**2*e**9*x**3 - 14549535*a**6*b**5*d*e
**10*x**4 - 46558512*a**6*b**5*e**11*x**5 - 1287*a**5*b**6*d**6*e**5 - 2574
0*a**5*b**6*d**5*e**6*x - 244530*a**5*b**6*d**4*e**7*x**2 - 1467180*a**5*b
**6*d**3*e**8*x**3 - 6235515*a**5*b**6*d**2*e**9*x**4 - 19953648*a**5*b**6
*d*e**10*x**5 - 49884120*a**5*b**6*e**11*x**6 - 495*a**4*b**7*d**7*e**4 -
9900*a**4*b**7*d**6*e**5*x - 94050*a**4*b**7*d**5*e**6*x**2 - 564300*a**4*
b**7*d**4*e**7*x**3 - 2398275*a**4*b**7*d**3*e**8*x**4 - 7674480*a**4*b**7
*d**2*e**9*x**5 - 19186200*a**4*b**7*d*e**10*x**6 - 38372400*a**4*b**7*e**
11*x**7 - 165*a**3*b**8*d**8*e**3 - 3300*a**3*b**8*d**7*e**4*x - 31350*a**
3*b**8*d**6*e**5*x**2 - 188100*a**3*b**8*d**5*e**6*x**3 - 799425*a**3*b**8
*d**4*e**7*x**4 - 2558160*a**3*b**8*d**3*e**8*x**5 - 6395400*a**3*b**8*d**
2*e**9*x**6 - 12790800*a**3*b**8*d*e**10*x**7 - 20785050*a**3*b**8*e**11*x
**8 - 45*a**2*b**9*d**9*e**2 - 900*a**2*b**9*d**8*e**3*x - 8550*a**2*b**9
```

$$\mathbf{3.100} \quad \int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{22}} dx$$

Optimal result	1042
Mathematica [B] (verified)	1043
Rubi [A] (verified)	1044
Maple [B] (verified)	1045
Fricas [B] (verification not implemented)	1046
Sympy [F(-1)]	1047
Maxima [B] (verification not implemented)	1048
Giac [B] (verification not implemented)	1049
Mupad [B] (verification not implemented)	1050
Reduce [B] (verification not implemented)	1050

Optimal result

Integrand size = 20, antiderivative size = 464

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{22}} dx = \frac{(bd-ae)^{10}(Bd-Ae)}{21e^{12}(d+ex)^{21}} - \frac{(bd-ae)^9(11bBd-10Abe-aBe)}{20e^{12}(d+ex)^{20}} + \frac{5b(bd-ae)^8(11bBd-9Abe-2aBe)}{19e^{12}(d+ex)^{19}} - \frac{5b^2(bd-ae)^7(11bBd-8Abe-3aBe)}{6e^{12}(d+ex)^{18}} + \frac{30b^3(bd-ae)^6(11bBd-7Abe-4aBe)}{17e^{12}(d+ex)^{17}} - \frac{21b^4(bd-ae)^5(11bBd-6Abe-5aBe)}{8e^{12}(d+ex)^{16}} + \frac{14b^5(bd-ae)^4(11bBd-5Abe-6aBe)}{5e^{12}(d+ex)^{15}} - \frac{15b^6(bd-ae)^3(11bBd-4Abe-7aBe)}{7e^{12}(d+ex)^{14}} + \frac{15b^7(bd-ae)^2(11bBd-3Abe-8aBe)}{13e^{12}(d+ex)^{13}} - \frac{5b^8(bd-ae)(11bBd-2Abe-9aBe)}{12e^{12}(d+ex)^{12}} + \frac{b^9(11bBd-Abe-10aBe)}{11e^{12}(d+ex)^{11}} - \frac{b^{10}B}{10e^{12}(d+ex)^{10}}$$

output

```
1/21*(-a*e+b*d)^10*(-A*e+B*d)/e^12/(e*x+d)^21-1/20*(-a*e+b*d)^9*(-10*A*b*e
-B*a*e+11*B*b*d)/e^12/(e*x+d)^20+5/19*b*(-a*e+b*d)^8*(-9*A*b*e-2*B*a*e+11*
B*b*d)/e^12/(e*x+d)^19-5/6*b^2*(-a*e+b*d)^7*(-8*A*b*e-3*B*a*e+11*B*b*d)/e^
12/(e*x+d)^18+30/17*b^3*(-a*e+b*d)^6*(-7*A*b*e-4*B*a*e+11*B*b*d)/e^12/(e*x
+d)^17-21/8*b^4*(-a*e+b*d)^5*(-6*A*b*e-5*B*a*e+11*B*b*d)/e^12/(e*x+d)^16+1
4/5*b^5*(-a*e+b*d)^4*(-5*A*b*e-6*B*a*e+11*B*b*d)/e^12/(e*x+d)^15-15/7*b^6*
(-a*e+b*d)^3*(-4*A*b*e-7*B*a*e+11*B*b*d)/e^12/(e*x+d)^14+15/13*b^7*(-a*e+b
*d)^2*(-3*A*b*e-8*B*a*e+11*B*b*d)/e^12/(e*x+d)^13-5/12*b^8*(-a*e+b*d)*(-2*
A*b*e-9*B*a*e+11*B*b*d)/e^12/(e*x+d)^12+1/11*b^9*(-A*b*e-10*B*a*e+11*B*b*d
)/e^12/(e*x+d)^11-1/10*b^10*B/e^12/(e*x+d)^10
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1431 vs. $2(464) = 928$.

Time = 0.54 (sec) , antiderivative size = 1431, normalized size of antiderivative = 3.08

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{22}} dx = \text{Too large to display}$$

input `Integrate[((a + b*x)^10*(A + B*x))/(d + e*x)^22,x]`

output

```
-1/38798760*(92378*a^10*e^10*(20*A*e + B*(d + 21*e*x)) + 48620*a^9*b*e^9*(
19*A*e*(d + 21*e*x) + 2*B*(d^2 + 21*d*e*x + 210*e^2*x^2)) + 72930*a^8*b^2*
e^8*(6*A*e*(d^2 + 21*d*e*x + 210*e^2*x^2) + B*(d^3 + 21*d^2*e*x + 210*d*e^
2*x^2 + 1330*e^3*x^3)) + 11440*a^7*b^3*e^7*(17*A*e*(d^3 + 21*d^2*e*x + 210
*d*e^2*x^2 + 1330*e^3*x^3) + 4*B*(d^4 + 21*d^3*e*x + 210*d^2*e^2*x^2 + 133
0*d*e^3*x^3 + 5985*e^4*x^4)) + 5005*a^6*b^4*e^6*(16*A*e*(d^4 + 21*d^3*e*x
+ 210*d^2*e^2*x^2 + 1330*d*e^3*x^3 + 5985*e^4*x^4) + 5*B*(d^5 + 21*d^4*e*x
+ 210*d^3*e^2*x^2 + 1330*d^2*e^3*x^3 + 5985*d*e^4*x^4 + 20349*e^5*x^5)) +
6006*a^5*b^5*e^5*(5*A*e*(d^5 + 21*d^4*e*x + 210*d^3*e^2*x^2 + 1330*d^2*e^
3*x^3 + 5985*d*e^4*x^4 + 20349*e^5*x^5) + 2*B*(d^6 + 21*d^5*e*x + 210*d^4*
e^2*x^2 + 1330*d^3*e^3*x^3 + 5985*d^2*e^4*x^4 + 20349*d*e^5*x^5 + 54264*e^
6*x^6)) + 5005*a^4*b^6*e^4*(2*A*e*(d^6 + 21*d^5*e*x + 210*d^4*e^2*x^2 + 13
30*d^3*e^3*x^3 + 5985*d^2*e^4*x^4 + 20349*d*e^5*x^5 + 54264*e^6*x^6) + B*(
d^7 + 21*d^6*e*x + 210*d^5*e^2*x^2 + 1330*d^4*e^3*x^3 + 5985*d^3*e^4*x^4 +
20349*d^2*e^5*x^5 + 54264*d*e^6*x^6 + 116280*e^7*x^7)) + 220*a^3*b^7*e^3*
(13*A*e*(d^7 + 21*d^6*e*x + 210*d^5*e^2*x^2 + 1330*d^4*e^3*x^3 + 5985*d^3*
e^4*x^4 + 20349*d^2*e^5*x^5 + 54264*d*e^6*x^6 + 116280*e^7*x^7) + 8*B*(d^8
+ 21*d^7*e*x + 210*d^6*e^2*x^2 + 1330*d^5*e^3*x^3 + 5985*d^4*e^4*x^4 + 20
349*d^3*e^5*x^5 + 54264*d^2*e^6*x^6 + 116280*d*e^7*x^7 + 203490*e^8*x^8))
+ 165*a^2*b^8*e^2*(4*A*e*(d^8 + 21*d^7*e*x + 210*d^6*e^2*x^2 + 1330*d^5...
```


Rubi [A] (verified)

Time = 1.21 (sec) , antiderivative size = 464, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{22}} dx$$

↓ 86

$$\int \left(\frac{b^9(10aBe + Abe - 11bBd)}{e^{11}(d + ex)^{12}} - \frac{5b^8(bd - ae)(9aBe + 2Abe - 11bBd)}{e^{11}(d + ex)^{13}} + \frac{15b^7(bd - ae)^2(8aBe + 3Abe - 11bBd)}{e^{11}(d + ex)^{14}} \right) dx$$

↓ 2009

$$\frac{b^9(-10aBe - Abe + 11bBd)}{11e^{12}(d + ex)^{11}} - \frac{5b^8(bd - ae)(-9aBe - 2Abe + 11bBd)}{12e^{12}(d + ex)^{12}} + \frac{15b^7(bd - ae)^2(-8aBe - 3Abe + 11bBd)}{13e^{12}(d + ex)^{13}} - \frac{15b^6(bd - ae)^3(-7aBe - 4Abe + 11bBd)}{7e^{12}(d + ex)^{14}} + \frac{14b^5(bd - ae)^4(-6aBe - 5Abe + 11bBd)}{5e^{12}(d + ex)^{15}} - \frac{21b^4(bd - ae)^5(-5aBe - 6Abe + 11bBd)}{8e^{12}(d + ex)^{16}} + \frac{30b^3(bd - ae)^6(-4aBe - 7Abe + 11bBd)}{17e^{12}(d + ex)^{17}} - \frac{5b^2(bd - ae)^7(-3aBe - 8Abe + 11bBd)}{6e^{12}(d + ex)^{18}} + \frac{5b(bd - ae)^8(-2aBe - 9Abe + 11bBd)}{19e^{12}(d + ex)^{19}} - \frac{(bd - ae)^9(-aBe - 10Abe + 11bBd)}{20e^{12}(d + ex)^{20}} + \frac{(bd - ae)^{10}(Bd - Ae)}{21e^{12}(d + ex)^{21}} - \frac{b^{10}B}{10e^{12}(d + ex)^{10}}$$

input

```
Int[((a + b*x)^10*(A + B*x))/(d + e*x)^22,x]
```

output

$$\begin{aligned} & ((b*d - a*e)^{10}*(B*d - A*e))/(21*e^{12}*(d + e*x)^{21}) - ((b*d - a*e)^9*(11*b \\ & *B*d - 10*A*b*e - a*B*e))/(20*e^{12}*(d + e*x)^{20}) + (5*b*(b*d - a*e)^8*(11* \\ & b*B*d - 9*A*b*e - 2*a*B*e))/(19*e^{12}*(d + e*x)^{19}) - (5*b^2*(b*d - a*e)^7* \\ & (11*b*B*d - 8*A*b*e - 3*a*B*e))/(6*e^{12}*(d + e*x)^{18}) + (30*b^3*(b*d - a*e) \\ &)^6*(11*b*B*d - 7*A*b*e - 4*a*B*e))/(17*e^{12}*(d + e*x)^{17}) - (21*b^4*(b*d \\ & - a*e)^5*(11*b*B*d - 6*A*b*e - 5*a*B*e))/(8*e^{12}*(d + e*x)^{16}) + (14*b^5*(\\ & b*d - a*e)^4*(11*b*B*d - 5*A*b*e - 6*a*B*e))/(5*e^{12}*(d + e*x)^{15}) - (15*b \\ & ^6*(b*d - a*e)^3*(11*b*B*d - 4*A*b*e - 7*a*B*e))/(7*e^{12}*(d + e*x)^{14}) + (\\ & 15*b^7*(b*d - a*e)^2*(11*b*B*d - 3*A*b*e - 8*a*B*e))/(13*e^{12}*(d + e*x)^{13} \\ &) - (5*b^8*(b*d - a*e)*(11*b*B*d - 2*A*b*e - 9*a*B*e))/(12*e^{12}*(d + e*x)^{12}) \\ & + (b^9*(11*b*B*d - A*b*e - 10*a*B*e))/(11*e^{12}*(d + e*x)^{11}) - (b^{10}*B \\ &)/(10*e^{12}*(d + e*x)^{10}) \end{aligned}$$

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1900 vs. $2(440) = 880$.

Time = 0.38 (sec) , antiderivative size = 1901, normalized size of antiderivative = 4.10

method	result	size
risch	Expression too large to display	1901
default	Expression too large to display	1942
norman	Expression too large to display	2014
gospers	Expression too large to display	2233
orering	Expression too large to display	2233
parallelrisch	Expression too large to display	2242

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^22,x,method=_RETURNVERBOSE)`

output

```
(-1/38798760/e^12*(1847560*A*a^10*e^11+923780*A*a^9*b*d*e^10+437580*A*a^8*
b^2*d^2*e^9+194480*A*a^7*b^3*d^3*e^8+80080*A*a^6*b^4*d^4*e^7+30030*A*a^5*b
^5*d^5*e^6+10010*A*a^4*b^6*d^6*e^5+2860*A*a^3*b^7*d^7*e^4+660*A*a^2*b^8*d
^8*e^3+110*A*a*b^9*d^9*e^2+10*A*b^10*d^10*e+92378*B*a^10*d*e^10+97240*B*a^9
*b*d^2*e^9+72930*B*a^8*b^2*d^3*e^8+45760*B*a^7*b^3*d^4*e^7+25025*B*a^6*b^4
*d^5*e^6+12012*B*a^5*b^5*d^6*e^5+5005*B*a^4*b^6*d^7*e^4+1760*B*a^3*b^7*d^8
*e^3+495*B*a^2*b^8*d^9*e^2+100*B*a*b^9*d^10*e+11*B*b^10*d^11)-1/1847560/e^
11*(923780*A*a^9*b*e^10+437580*A*a^8*b^2*d*e^9+194480*A*a^7*b^3*d^2*e^8+80
080*A*a^6*b^4*d^3*e^7+30030*A*a^5*b^5*d^4*e^6+10010*A*a^4*b^6*d^5*e^5+2860
*A*a^3*b^7*d^6*e^4+660*A*a^2*b^8*d^7*e^3+110*A*a*b^9*d^8*e^2+10*A*b^10*d^9
*e+92378*B*a^10*e^10+97240*B*a^9*b*d*e^9+72930*B*a^8*b^2*d^2*e^8+45760*B*a
^7*b^3*d^3*e^7+25025*B*a^6*b^4*d^4*e^6+12012*B*a^5*b^5*d^5*e^5+5005*B*a^4*
b^6*d^6*e^4+1760*B*a^3*b^7*d^7*e^3+495*B*a^2*b^8*d^8*e^2+100*B*a*b^9*d^9*e
+11*B*b^10*d^10)*x-1/184756*b/e^10*(437580*A*a^8*b*e^9+194480*A*a^7*b^2*d*
e^8+80080*A*a^6*b^3*d^2*e^7+30030*A*a^5*b^4*d^3*e^6+10010*A*a^4*b^5*d^4*e^
5+2860*A*a^3*b^6*d^5*e^4+660*A*a^2*b^7*d^6*e^3+110*A*a*b^8*d^7*e^2+10*A*b^
9*d^8*e+97240*B*a^9*e^9+72930*B*a^8*b*d*e^8+45760*B*a^7*b^2*d^2*e^7+25025*
B*a^6*b^3*d^3*e^6+12012*B*a^5*b^4*d^4*e^5+5005*B*a^4*b^5*d^5*e^4+1760*B*a^
3*b^6*d^6*e^3+495*B*a^2*b^7*d^7*e^2+100*B*a*b^8*d^8*e+11*B*b^9*d^9)*x^2-1/
29172*b^2/e^9*(194480*A*a^7*b*e^8+80080*A*a^6*b^2*d*e^7+30030*A*a^5*b^3...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2039 vs. $2(440) = 880$.

Time = 0.26 (sec) , antiderivative size = 2039, normalized size of antiderivative = 4.39

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{22}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^22,x, algorithm="fricas")`

output

```

-1/38798760*(3879876*B*b^10*e^11*x^11 + 11*B*b^10*d^11 + 1847560*A*a^10*e^
11 + 10*(10*B*a*b^9 + A*b^10)*d^10*e + 55*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^
2 + 220*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 715*(7*B*a^4*b^6 + 4*A*a^3*b
^7)*d^7*e^4 + 2002*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 5005*(5*B*a^6*b^4
+ 6*A*a^5*b^5)*d^5*e^6 + 11440*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 2431
0*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^3*e^8 + 48620*(2*B*a^9*b + 9*A*a^8*b^2)*d^
2*e^9 + 92378*(B*a^10 + 10*A*a^9*b)*d*e^10 + 352716*(11*B*b^10*d*e^10 + 10
*(10*B*a*b^9 + A*b^10)*e^11)*x^10 + 293930*(11*B*b^10*d^2*e^9 + 10*(10*B*a
*b^9 + A*b^10)*d*e^10 + 55*(9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 + 203490*(1
1*B*b^10*d^3*e^8 + 10*(10*B*a*b^9 + A*b^10)*d^2*e^9 + 55*(9*B*a^2*b^8 + 2*
A*a*b^9)*d*e^10 + 220*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 116280*(11*B
*b^10*d^4*e^7 + 10*(10*B*a*b^9 + A*b^10)*d^3*e^8 + 55*(9*B*a^2*b^8 + 2*A*a
*b^9)*d^2*e^9 + 220*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 + 715*(7*B*a^4*b^6
+ 4*A*a^3*b^7)*e^11)*x^7 + 54264*(11*B*b^10*d^5*e^6 + 10*(10*B*a*b^9 + A*b
^10)*d^4*e^7 + 55*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 + 220*(8*B*a^3*b^7 + 3
*A*a^2*b^8)*d^2*e^9 + 715*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 + 2002*(6*B*a
^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 + 20349*(11*B*b^10*d^6*e^5 + 10*(10*B*a*b^
9 + A*b^10)*d^5*e^6 + 55*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 + 220*(8*B*a^3*
b^7 + 3*A*a^2*b^8)*d^3*e^8 + 715*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 + 200
2*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10 + 5005*(5*B*a^6*b^4 + 6*A*a^5*b^5)...

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{22}} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**10*(B*x+A)/(e*x+d)**22,x)
```

output

Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2039 vs. $2(440) = 880$.

Time = 0.16 (sec) , antiderivative size = 2039, normalized size of antiderivative = 4.39

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{22}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^22,x, algorithm="maxima")`

output

```
-1/38798760*(3879876*B*b^10*e^11*x^11 + 11*B*b^10*d^11 + 1847560*A*a^10*e^
11 + 10*(10*B*a*b^9 + A*b^10)*d^10*e + 55*(9*B*a^2*b^8 + 2*A*a*b^9)*d^9*e^
2 + 220*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d^8*e^3 + 715*(7*B*a^4*b^6 + 4*A*a^3*b
^7)*d^7*e^4 + 2002*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d^6*e^5 + 5005*(5*B*a^6*b^4
+ 6*A*a^5*b^5)*d^5*e^6 + 11440*(4*B*a^7*b^3 + 7*A*a^6*b^4)*d^4*e^7 + 2431
0*(3*B*a^8*b^2 + 8*A*a^7*b^3)*d^3*e^8 + 48620*(2*B*a^9*b + 9*A*a^8*b^2)*d^
2*e^9 + 92378*(B*a^10 + 10*A*a^9*b)*d*e^10 + 352716*(11*B*b^10*d*e^10 + 10
*(10*B*a*b^9 + A*b^10)*e^11)*x^10 + 293930*(11*B*b^10*d^2*e^9 + 10*(10*B*a
*b^9 + A*b^10)*d*e^10 + 55*(9*B*a^2*b^8 + 2*A*a*b^9)*e^11)*x^9 + 203490*(1
1*B*b^10*d^3*e^8 + 10*(10*B*a*b^9 + A*b^10)*d^2*e^9 + 55*(9*B*a^2*b^8 + 2*
A*a*b^9)*d*e^10 + 220*(8*B*a^3*b^7 + 3*A*a^2*b^8)*e^11)*x^8 + 116280*(11*B
*b^10*d^4*e^7 + 10*(10*B*a*b^9 + A*b^10)*d^3*e^8 + 55*(9*B*a^2*b^8 + 2*A*a
*b^9)*d^2*e^9 + 220*(8*B*a^3*b^7 + 3*A*a^2*b^8)*d*e^10 + 715*(7*B*a^4*b^6
+ 4*A*a^3*b^7)*e^11)*x^7 + 54264*(11*B*b^10*d^5*e^6 + 10*(10*B*a*b^9 + A*b
^10)*d^4*e^7 + 55*(9*B*a^2*b^8 + 2*A*a*b^9)*d^3*e^8 + 220*(8*B*a^3*b^7 + 3
*A*a^2*b^8)*d^2*e^9 + 715*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d*e^10 + 2002*(6*B*a
^5*b^5 + 5*A*a^4*b^6)*e^11)*x^6 + 20349*(11*B*b^10*d^6*e^5 + 10*(10*B*a*b^
9 + A*b^10)*d^5*e^6 + 55*(9*B*a^2*b^8 + 2*A*a*b^9)*d^4*e^7 + 220*(8*B*a^3*
b^7 + 3*A*a^2*b^8)*d^3*e^8 + 715*(7*B*a^4*b^6 + 4*A*a^3*b^7)*d^2*e^9 + 200
2*(6*B*a^5*b^5 + 5*A*a^4*b^6)*d*e^10 + 5005*(5*B*a^6*b^4 + 6*A*a^5*b^5)...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2232 vs. $2(440) = 880$.

Time = 0.13 (sec) , antiderivative size = 2232, normalized size of antiderivative = 4.81

$$\int \frac{(a+bx)^{10}(A+Bx)}{(d+ex)^{22}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^10*(B*x+A)/(e*x+d)^22,x, algorithm="giac")`

output

```
-1/38798760*(3879876*B*b^10*e^11*x^11 + 3879876*B*b^10*d*e^10*x^10 + 35271600*B*a*b^9*e^11*x^10 + 3527160*A*b^10*e^11*x^10 + 3233230*B*b^10*d^2*e^9*x^9 + 29393000*B*a*b^9*d*e^10*x^9 + 2939300*A*b^10*d*e^10*x^9 + 145495350*B*a^2*b^8*e^11*x^9 + 32332300*A*a*b^9*e^11*x^9 + 2238390*B*b^10*d^3*e^8*x^8 + 20349000*B*a*b^9*d^2*e^9*x^8 + 2034900*A*b^10*d^2*e^9*x^8 + 100727550*B*a^2*b^8*d*e^10*x^8 + 22383900*A*a*b^9*d*e^10*x^8 + 358142400*B*a^3*b^7*e^11*x^8 + 134303400*A*a^2*b^8*e^11*x^8 + 1279080*B*b^10*d^4*e^7*x^7 + 11628000*B*a*b^9*d^3*e^8*x^7 + 1162800*A*b^10*d^3*e^8*x^7 + 57558600*B*a^2*b^8*d^2*e^9*x^7 + 12790800*A*a*b^9*d^2*e^9*x^7 + 204652800*B*a^3*b^7*d*e^10*x^7 + 76744800*A*a^2*b^8*d*e^10*x^7 + 581981400*B*a^4*b^6*e^11*x^7 + 332560800*A*a^3*b^7*e^11*x^7 + 596904*B*b^10*d^5*e^6*x^6 + 5426400*B*a*b^9*d^4*e^7*x^6 + 542640*A*b^10*d^4*e^7*x^6 + 26860680*B*a^2*b^8*d^3*e^8*x^6 + 5969040*A*a*b^9*d^3*e^8*x^6 + 95504640*B*a^3*b^7*d^2*e^9*x^6 + 35814240*A*a^2*b^8*d^2*e^9*x^6 + 271591320*B*a^4*b^6*d*e^10*x^6 + 155195040*A*a^3*b^7*d*e^10*x^6 + 651819168*B*a^5*b^5*e^11*x^6 + 543182640*A*a^4*b^6*e^11*x^6 + 223839*B*b^10*d^6*e^5*x^5 + 2034900*B*a*b^9*d^5*e^6*x^5 + 203490*A*b^10*d^5*e^6*x^5 + 10072755*B*a^2*b^8*d^4*e^7*x^5 + 2238390*A*a*b^9*d^4*e^7*x^5 + 35814240*B*a^3*b^7*d^3*e^8*x^5 + 13430340*A*a^2*b^8*d^3*e^8*x^5 + 101846745*B*a^4*b^6*d^2*e^9*x^5 + 58198140*A*a^3*b^7*d^2*e^9*x^5 + 244432188*B*a^5*b^5*d*e^10*x^5 + 203693490*A*a^4*b^6*d*e^10*x^5 + 509233725*B*a^6*b^4*e...
```

Mupad [B] (verification not implemented)

Time = 1.63 (sec) , antiderivative size = 2121, normalized size of antiderivative = 4.57

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{22}} dx = \text{Too large to display}$$

input `int(((A + B*x)*(a + b*x)^10)/(d + e*x)^22,x)`

output

```

-((1847560*A*a^10*e^11 + 11*B*b^10*d^11 + 10*A*b^10*d^10*e + 92378*B*a^10*
d*e^10 + 110*A*a*b^9*d^9*e^2 + 97240*B*a^9*b*d^2*e^9 + 660*A*a^2*b^8*d^8*e
^3 + 2860*A*a^3*b^7*d^7*e^4 + 10010*A*a^4*b^6*d^6*e^5 + 30030*A*a^5*b^5*d^
5*e^6 + 80080*A*a^6*b^4*d^4*e^7 + 194480*A*a^7*b^3*d^3*e^8 + 437580*A*a^8*
b^2*d^2*e^9 + 495*B*a^2*b^8*d^9*e^2 + 1760*B*a^3*b^7*d^8*e^3 + 5005*B*a^4*
b^6*d^7*e^4 + 12012*B*a^5*b^5*d^6*e^5 + 25025*B*a^6*b^4*d^5*e^6 + 45760*B*
a^7*b^3*d^4*e^7 + 72930*B*a^8*b^2*d^3*e^8 + 923780*A*a^9*b*d*e^10 + 100*B*
a*b^9*d^10*e)/(38798760*e^12) + (x*(92378*B*a^10*e^10 + 11*B*b^10*d^10 + 9
23780*A*a^9*b*e^10 + 10*A*b^10*d^9*e + 110*A*a*b^9*d^8*e^2 + 437580*A*a^8*
b^2*d*e^9 + 660*A*a^2*b^8*d^7*e^3 + 2860*A*a^3*b^7*d^6*e^4 + 10010*A*a^4*b
^6*d^5*e^5 + 30030*A*a^5*b^5*d^4*e^6 + 80080*A*a^6*b^4*d^3*e^7 + 194480*A*
a^7*b^3*d^2*e^8 + 495*B*a^2*b^8*d^8*e^2 + 1760*B*a^3*b^7*d^7*e^3 + 5005*B*
a^4*b^6*d^6*e^4 + 12012*B*a^5*b^5*d^5*e^5 + 25025*B*a^6*b^4*d^4*e^6 + 4576
0*B*a^7*b^3*d^3*e^7 + 72930*B*a^8*b^2*d^2*e^8 + 100*B*a*b^9*d^9*e + 97240*
B*a^9*b*d*e^9))/(1847560*e^11) + (3*b^7*x^8*(1760*B*a^3*e^3 + 11*B*b^3*d^3
+ 660*A*a^2*b*e^3 + 10*A*b^3*d^2*e + 110*A*a*b^2*d*e^2 + 100*B*a*b^2*d^2*
e + 495*B*a^2*b*d*e^2))/(572*e^4) + (3*b^4*x^5*(25025*B*a^6*e^6 + 11*B*b^6
*d^6 + 30030*A*a^5*b*e^6 + 10*A*b^6*d^5*e + 110*A*a*b^5*d^4*e^2 + 10010*A*
a^4*b^2*d*e^5 + 660*A*a^2*b^4*d^3*e^3 + 2860*A*a^3*b^3*d^2*e^4 + 495*B*a^2
*b^4*d^4*e^2 + 1760*B*a^3*b^3*d^3*e^3 + 5005*B*a^4*b^2*d^2*e^4 + 100*B*...

```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 1371, normalized size of antiderivative = 2.95

$$\int \frac{(a + bx)^{10}(A + Bx)}{(d + ex)^{22}} dx = \text{Too large to display}$$

input `int((b*x+a)^10*(B*x+A)/(e*x+d)^22,x)`

output

```
( - 167960*a**11*e**11 - 92378*a**10*b*d*e**10 - 1939938*a**10*b*e**11*x -
48620*a**9*b**2*d**2*e**9 - 1021020*a**9*b**2*d*e**10*x - 10210200*a**9*b
**2*e**11*x**2 - 24310*a**8*b**3*d**3*e**8 - 510510*a**8*b**3*d**2*e**9*x
- 5105100*a**8*b**3*d*e**10*x**2 - 32332300*a**8*b**3*e**11*x**3 - 11440*a
**7*b**4*d**4*e**7 - 240240*a**7*b**4*d**3*e**8*x - 2402400*a**7*b**4*d**2
*e**9*x**2 - 15215200*a**7*b**4*d*e**10*x**3 - 68468400*a**7*b**4*e**11*x*
*4 - 5005*a**6*b**5*d**5*e**6 - 105105*a**6*b**5*d**4*e**7*x - 1051050*a**
6*b**5*d**3*e**8*x**2 - 6656650*a**6*b**5*d**2*e**9*x**3 - 29954925*a**6*b
**5*d*e**10*x**4 - 101846745*a**6*b**5*e**11*x**5 - 2002*a**5*b**6*d**6*e
**5 - 42042*a**5*b**6*d**5*e**6*x - 420420*a**5*b**6*d**4*e**7*x**2 - 26626
60*a**5*b**6*d**3*e**8*x**3 - 11981970*a**5*b**6*d**2*e**9*x**4 - 40738698
*a**5*b**6*d*e**10*x**5 - 108636528*a**5*b**6*e**11*x**6 - 715*a**4*b**7*d
**7*e**4 - 15015*a**4*b**7*d**6*e**5*x - 150150*a**4*b**7*d**5*e**6*x**2 -
950950*a**4*b**7*d**4*e**7*x**3 - 4279275*a**4*b**7*d**3*e**8*x**4 - 1454
9535*a**4*b**7*d**2*e**9*x**5 - 38798760*a**4*b**7*d*e**10*x**6 - 83140200
*a**4*b**7*e**11*x**7 - 220*a**3*b**8*d**8*e**3 - 4620*a**3*b**8*d**7*e**4
*x - 46200*a**3*b**8*d**6*e**5*x**2 - 292600*a**3*b**8*d**5*e**6*x**3 - 13
16700*a**3*b**8*d**4*e**7*x**4 - 4476780*a**3*b**8*d**3*e**8*x**5 - 119380
80*a**3*b**8*d**2*e**9*x**6 - 25581600*a**3*b**8*d*e**10*x**7 - 44767800*a
**3*b**8*e**11*x**8 - 55*a**2*b**9*d**9*e**2 - 1155*a**2*b**9*d**8*e**3...
```


3.101 $\int \frac{(A+Bx)(d+ex)^4}{a+bx} dx$

Optimal result	1052
Mathematica [A] (verified)	1052
Rubi [A] (verified)	1053
Maple [B] (verified)	1054
Fricas [B] (verification not implemented)	1055
Sympy [B] (verification not implemented)	1056
Maxima [B] (verification not implemented)	1057
Giac [B] (verification not implemented)	1057
Mupad [B] (verification not implemented)	1058
Reduce [B] (verification not implemented)	1059

Optimal result

Integrand size = 20, antiderivative size = 155

$$\int \frac{(A+Bx)(d+ex)^4}{a+bx} dx = \frac{(Ab-aB)e(bd-ae)^3x}{b^5} + \frac{(Ab-aB)(bd-ae)^2(d+ex)^2}{2b^4} + \frac{(Ab-aB)(bd-ae)(d+ex)^3}{3b^3} + \frac{(Ab-aB)(d+ex)^4}{4b^2} + \frac{B(d+ex)^5}{5be} + \frac{(Ab-aB)(bd-ae)^4 \log(a+bx)}{b^6}$$

output

$$(A*b-B*a)*e*(-a*e+b*d)^3*x/b^5+1/2*(A*b-B*a)*(-a*e+b*d)^2*(e*x+d)^2/b^4+1/3*(A*b-B*a)*(-a*e+b*d)*(e*x+d)^3/b^3+1/4*(A*b-B*a)*(e*x+d)^4/b^2+1/5*B*(e*x+d)^5/b/e+(A*b-B*a)*(-a*e+b*d)^4*\ln(b*x+a)/b^6$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 257, normalized size of antiderivative = 1.66

$$\int \frac{(A+Bx)(d+ex)^4}{a+bx} dx = \frac{bx(60a^4Be^4 - 30a^3be^3(8Bd + 2Ae + Bex) + 10a^2b^2e^2(3Ae(8d + ex) + 2B(18d^2 + 6dex + e^2x^2)) - 5ab$$

input `Integrate[((A + B*x)*(d + e*x)^4)/(a + b*x), x]`

output $(b*x*(60*a^4*B*e^4 - 30*a^3*b*e^3*(8*B*d + 2*A*e + B*e*x) + 10*a^2*b^2*e^2*(3*A*e*(8*d + e*x) + 2*B*(18*d^2 + 6*d*e*x + e^2*x^2)) - 5*a*b^3*e*(4*A*e*(18*d^2 + 6*d*e*x + e^2*x^2) + B*(48*d^3 + 36*d^2*e*x + 16*d*e^2*x^2 + 3*e^3*x^3)) + b^4*(5*A*e*(48*d^3 + 36*d^2*e*x + 16*d*e^2*x^2 + 3*e^3*x^3) + 12*B*(5*d^4 + 10*d^3*e*x + 10*d^2*e^2*x^2 + 5*d*e^3*x^3 + e^4*x^4))) + 60*(A*b - a*B)*(b*d - a*e)^4*\text{Log}[a + b*x])/(60*b^6)$

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)^4}{a + bx} dx$$

↓ 86

$$\int \left(\frac{(Ab - aB)(bd - ae)^4}{b^5(a + bx)} + \frac{e(Ab - aB)(bd - ae)^3}{b^5} + \frac{e(d + ex)(Ab - aB)(bd - ae)^2}{b^4} + \frac{e(d + ex)^2(Ab - aB)(bd - ae)}{b^3} \right) dx$$

↓ 2009

$$\frac{(Ab - aB)(bd - ae)^4 \log(a + bx)}{b^6} + \frac{ex(Ab - aB)(bd - ae)^3}{b^5} + \frac{(d + ex)^2(Ab - aB)(bd - ae)^2}{2b^4} + \frac{(d + ex)^3(Ab - aB)(bd - ae)}{3b^3} + \frac{(d + ex)^4(Ab - aB)}{4b^2} + \frac{B(d + ex)^5}{5be}$$

input `Int[((A + B*x)*(d + e*x)^4)/(a + b*x), x]`

output
$$\begin{aligned} & ((A*b - a*B)*e*(b*d - a*e)^{3*x})/b^5 + ((A*b - a*B)*(b*d - a*e)^2*(d + e*x) \\ & ^2)/(2*b^4) + ((A*b - a*B)*(b*d - a*e)*(d + e*x)^3)/(3*b^3) + ((A*b - a*B) \\ & *(d + e*x)^4)/(4*b^2) + (B*(d + e*x)^5)/(5*b*e) + ((A*b - a*B)*(b*d - a*e) \\ & ^4*\text{Log}[a + b*x])/b^6 \end{aligned}$$

Defintions of rubi rules used

rule 86
$$\begin{aligned} & \text{Int}[(a_.) + (b_.)*(x_.)]*((c_.) + (d_.)*(x_.))^{(n_.)}*((e_.) + (f_.)*(x_.))^{(p_.)} \\ & ., x_] := \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; \\ & \text{FreeQ}\{a, b, c, d, e, f, n\}, x\} \&\& ((\text{ILtQ}[n, 0] \&\& \text{ILtQ}[p, 0]) \|\| \text{EqQ}[p, 1] \\ & \|\| (\text{IGtQ}[p, 0] \&\& (!\text{IntegerQ}[n] \|\| \text{LeQ}[9*p + 5*(n + 2), 0] \|\| \text{GeQ}[n + p \\ & + 1, 0] \|\| (\text{GeQ}[n + p + 2, 0] \&\& \text{RationalQ}[a, b, c, d, e, f]))) \end{aligned}$$

rule 2009
$$\text{Int}[u_, x_Symbol] := \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 403 vs. $2(147) = 294$.

Time = 0.22 (sec) , antiderivative size = 404, normalized size of antiderivative = 2.61

method	result
norman	$-\frac{(Aa^3be^4 - 4Aa^2b^2de^3 + 6Aab^3d^2e^2 - 4Ab^4d^3e - Ba^4e^4 + 4Ba^3bd^3e^3 - 6Ba^2b^2d^2e^2 + 4Bab^3d^3e - Bb^4d^4)x}{b^5} + \frac{Be^4x^5}{5b} +$
default	$-\frac{-Bb^4de^3x^4 + \frac{1}{3}Aab^3e^4x^3 - \frac{4}{3}Ab^4de^3x^3 - \frac{1}{3}Ba^2b^2e^4x^3 - 2Bb^4d^2e^2x^3 - 3Ab^4d^2e^2x^2 + \frac{1}{2}Ba^3be^4x^2 - 2Bb^4d^3ex^2 + Aa^3b}{b^5}$
risch	$-\frac{4\ln(bx+a)Aa^3de^3}{b^4} + \frac{6\ln(bx+a)Aa^2d^2e^2}{b^3} - \frac{4\ln(bx+a)Aad^3e}{b^2} + \frac{4\ln(bx+a)Ba^4d^3e^3}{b^5} - \frac{6\ln(bx+a)Ba^3d^2e^2}{b^4} + \frac{4}{b^5}$
parallelrisch	$\frac{240Axa^2b^3de^3 - 360Axa^2b^4d^2e^2 - 240Bxa^3b^2de^3 + 360Bxa^2b^3d^2e^2 - 240A\ln(bx+a)a^3b^2de^3 + 360A\ln(bx+a)a^2b^3d^2e^2 - 240A\ln(bx+a)a^3b^2de^3}{b^5}$

input
$$\text{int}((B*x+A)*(e*x+d)^4/(b*x+a), x, \text{method}=_RETURNVERBOSE)$$

output

```

-(A*a^3*b*e^4-4*A*a^2*b^2*d*e^3+6*A*a*b^3*d^2*e^2-4*A*b^4*d^3*e-B*a^4*e^4+
4*B*a^3*b*d*e^3-6*B*a^2*b^2*d^2*e^2+4*B*a*b^3*d^3*e-B*b^4*d^4)/b^5*x+1/5*B
/b*e^4*x^5+1/2/b^4*e*(A*a^2*b*e^3-4*A*a*b^2*d*e^2+6*A*b^3*d^2*e-B*a^3*e^3+
4*B*a^2*b*d*e^2-6*B*a*b^2*d^2*e+4*B*b^3*d^3)*x^2-1/3/b^3*e^2*(A*a*b*e^2-4*
A*b^2*d*e-B*a^2*e^2+4*B*a*b*d*e-6*B*b^2*d^2)*x^3+1/4/b^2*e^3*(A*b*e-B*a*e+
4*B*b*d)*x^4+(A*a^4*b*e^4-4*A*a^3*b^2*d*e^3+6*A*a^2*b^3*d^2*e^2-4*A*a*b^4*
d^3*e+A*b^5*d^4-B*a^5*e^4+4*B*a^4*b*d*e^3-6*B*a^3*b^2*d^2*e^2+4*B*a^2*b^3*
d^3*e-B*a*b^4*d^4)/b^6*ln(b*x+a)

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 403 vs. $2(149) = 298$.

Time = 0.09 (sec) , antiderivative size = 403, normalized size of antiderivative = 2.60

$$\int \frac{(A + Bx)(d + ex)^4}{a + bx} dx$$

$$= \frac{12 Bb^5 e^4 x^5 + 15 (4 Bb^5 d e^3 - (Bab^4 - Ab^5) e^4) x^4 + 20 (6 Bb^5 d^2 e^2 - 4 (Bab^4 - Ab^5) d e^3 + (Ba^2 b^3 - Aab^4)$$

input

```
integrate((B*x+A)*(e*x+d)^4/(b*x+a),x, algorithm="fricas")
```

output

```

1/60*(12*B*b^5*e^4*x^5 + 15*(4*B*b^5*d*e^3 - (B*a*b^4 - A*b^5)*e^4)*x^4 +
20*(6*B*b^5*d^2*e^2 - 4*(B*a*b^4 - A*b^5)*d*e^3 + (B*a^2*b^3 - A*a*b^4)*e^4)*x^3 +
30*(4*B*b^5*d^3*e - 6*(B*a*b^4 - A*b^5)*d^2*e^2 + 4*(B*a^2*b^3 - A*a*b^4)*d*e^3 -
(B*a^3*b^2 - A*a^2*b^3)*e^4)*x^2 + 60*(B*b^5*d^4 - 4*(B*a*b^4 - A*b^5)*d^3*e +
6*(B*a^2*b^3 - A*a*b^4)*d^2*e^2 - 4*(B*a^3*b^2 - A*a^2*b^3)*d*e^3 + (B*a^4*b -
A*a^3*b^2)*e^4)*x - 60*((B*a*b^4 - A*b^5)*d^4 - 4*(B*a^2*b^3 - A*a*b^4)*d^3*e +
6*(B*a^3*b^2 - A*a^2*b^3)*d^2*e^2 - 4*(B*a^4*b - A*a^3*b^2)*d*e^3 + (B*a^5 -
A*a^4*b)*e^4)*log(b*x + a))/b^6

```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 352 vs. $2(136) = 272$.

Time = 0.55 (sec) , antiderivative size = 352, normalized size of antiderivative = 2.27

$$\int \frac{(A + Bx)(d + ex)^4}{a + bx} dx = \frac{Be^4x^5}{5b} + x^4 \left(\frac{Ae^4}{4b} - \frac{Bae^4}{4b^2} + \frac{Bde^3}{b} \right) + x^3 \left(-\frac{Aae^4}{3b^2} + \frac{4Ade^3}{3b} + \frac{Ba^2e^4}{3b^3} - \frac{4Bade^3}{3b^2} + \frac{2Bd^2e^2}{b} \right) + x^2 \left(\frac{Aa^2e^4}{2b^3} - \frac{2Aade^3}{b^2} + \frac{3Ad^2e^2}{b} - \frac{Ba^3e^4}{2b^4} + \frac{2Ba^2de^3}{b^3} - \frac{3Bad^2e^2}{b^2} + \frac{2Bd^3e}{b} \right) + x \left(-\frac{Aa^3e^4}{b^4} + \frac{4Aa^2de^3}{b^3} - \frac{6Aad^2e^2}{b^2} + \frac{4Ad^3e}{b} + \frac{Ba^4e^4}{b^5} - \frac{4Ba^3de^3}{b^4} + \frac{6Ba^2d^2e^2}{b^3} - \frac{4Bad^3e}{b^2} + \frac{Bd^4}{b} \right) - \frac{(-Ab + Ba)(ae - bd)^4 \log(a + bx)}{b^6}$$

input `integrate((B*x+A)*(e*x+d)**4/(b*x+a), x)`

output `B*e**4*x**5/(5*b) + x**4*(A*e**4/(4*b) - B*a*e**4/(4*b**2) + B*d*e**3/b) + x**3*(-A*a*e**4/(3*b**2) + 4*A*d*e**3/(3*b) + B*a**2*e**4/(3*b**3) - 4*B*a*d*e**3/(3*b**2) + 2*B*d**2*e**2/b) + x**2*(A*a**2*e**4/(2*b**3) - 2*A*a*d*e**3/b**2 + 3*A*d**2*e**2/b - B*a**3*e**4/(2*b**4) + 2*B*a**2*d*e**3/b**3 - 3*B*a*d**2*e**2/b**2 + 2*B*d**3*e/b) + x*(-A*a**3*e**4/b**4 + 4*A*a**2*d*e**3/b**3 - 6*A*a*d**2*e**2/b**2 + 4*A*d**3*e/b + B*a**4*e**4/b**5 - 4*B*a**3*d*e**3/b**4 + 6*B*a**2*d**2*e**2/b**3 - 4*B*a*d**3*e/b**2 + B*d**4/b) - (-A*b + B*a)*(a*e - b*d)**4*log(a + b*x)/b**6`

output

```

1/60*(12*B*b^4*e^4*x^5 + 60*B*b^4*d*e^3*x^4 - 15*B*a*b^3*e^4*x^4 + 15*A*b^
4*e^4*x^4 + 120*B*b^4*d^2*e^2*x^3 - 80*B*a*b^3*d*e^3*x^3 + 80*A*b^4*d*e^3*
x^3 + 20*B*a^2*b^2*e^4*x^3 - 20*A*a*b^3*e^4*x^3 + 120*B*b^4*d^3*e*x^2 - 18
0*B*a*b^3*d^2*e^2*x^2 + 180*A*b^4*d^2*e^2*x^2 + 120*B*a^2*b^2*d*e^3*x^2 -
120*A*a*b^3*d*e^3*x^2 - 30*B*a^3*b*e^4*x^2 + 30*A*a^2*b^2*e^4*x^2 + 60*B*b
^4*d^4*x - 240*B*a*b^3*d^3*e*x + 240*A*b^4*d^3*e*x + 360*B*a^2*b^2*d^2*e^2
*x - 360*A*a*b^3*d^2*e^2*x - 240*B*a^3*b*d*e^3*x + 240*A*a^2*b^2*d*e^3*x +
60*B*a^4*e^4*x - 60*A*a^3*b*e^4*x)/b^5 - (B*a*b^4*d^4 - A*b^5*d^4 - 4*B*a
^2*b^3*d^3*e + 4*A*a*b^4*d^3*e + 6*B*a^3*b^2*d^2*e^2 - 6*A*a^2*b^3*d^2*e^2
- 4*B*a^4*b*d*e^3 + 4*A*a^3*b^2*d*e^3 + B*a^5*e^4 - A*a^4*b*e^4)*log(abs(
b*x + a))/b^6

```

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 411, normalized size of antiderivative = 2.65

$$\int \frac{(A + Bx)(d + ex)^4}{a + bx} dx$$

$$= x \left(\frac{Bd^4 + 4Aed^3}{b} - \frac{a \left(\frac{a \left(\frac{Ae^4 + 4Bde^3}{b} - \frac{Bae^4}{b^2} \right) - 2de^2(2Ae + 3Bd)}{b} \right) + \frac{2d^2e(3Ae + 2Bd)}{b}}{b} \right)$$

$$- x^3 \left(\frac{a \left(\frac{Ae^4 + 4Bde^3}{b} - \frac{Bae^4}{b^2} \right) - 2de^2(2Ae + 3Bd)}{3b} \right)$$

$$+ x^4 \left(\frac{Ae^4 + 4Bde^3}{4b} - \frac{Bae^4}{4b^2} \right)$$

$$+ x^2 \left(\frac{a \left(\frac{a \left(\frac{Ae^4 + 4Bde^3}{b} - \frac{Bae^4}{b^2} \right) - 2de^2(2Ae + 3Bd)}{b} \right) + \frac{d^2e(3Ae + 2Bd)}{b}}{2b} \right)$$

$$+ \frac{\ln(a + bx) (-Ba^5e^4 + 4Ba^4bde^3 + Aa^4be^4 - 6Ba^3b^2d^2e^2 - 4Aa^3b^2de^3 + 4Ba^2b^3d^3e + 6Aa^2b^4d^4 - 6Aa^3b^3d^3e^2 - 4Aa^4b^2d^2e^2 - 4Aa^5bde^3 + 4Aa^6e^4)}{b^6}$$

$$+ \frac{Be^4x^5}{5b}$$

input `int((A + B*x)*(d + e*x)^4/(a + b*x),x)`

output `x*((B*d^4 + 4*A*d^3*e)/b - (a*((a*((a*((A*e^4 + 4*B*d*e^3)/b - (B*a*e^4)/b^2))/b - (2*d*e^2*(2*A*e + 3*B*d))/b))/b + (2*d^2*e*(3*A*e + 2*B*d))/b))/b - x^3*((a*((A*e^4 + 4*B*d*e^3)/b - (B*a*e^4)/b^2))/(3*b) - (2*d*e^2*(2*A*e + 3*B*d))/(3*b)) + x^4*((A*e^4 + 4*B*d*e^3)/(4*b) - (B*a*e^4)/(4*b^2)) + x^2*((a*((a*((A*e^4 + 4*B*d*e^3)/b - (B*a*e^4)/b^2))/b - (2*d*e^2*(2*A*e + 3*B*d))/b))/(2*b) + (d^2*e*(3*A*e + 2*B*d))/b) + (log(a + b*x)*(A*b^5*d^4 - B*a^5*e^4 + A*a^4*b*e^4 - B*a*b^4*d^4 - 4*A*a^3*b^2*d*e^3 + 4*B*a^2*b^3*d^3*e + 6*A*a^2*b^3*d^2*e^2 - 6*B*a^3*b^2*d^2*e^2 - 4*A*a*b^4*d^3*e + 4*B*a^4*b*d*e^3))/b^6 + (B*e^4*x^5)/(5*b)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.28

$$\int \frac{(A + Bx)(d + ex)^4}{a + bx} dx = \frac{x(e^4x^4 + 5de^3x^3 + 10d^2e^2x^2 + 10d^3ex + 5d^4)}{5}$$

input `int((B*x+A)*(e*x+d)^4/(b*x+a),x)`

output `(x*(5*d**4 + 10*d**3*e*x + 10*d**2*e**2*x**2 + 5*d*e**3*x**3 + e**4*x**4))/5`

3.102 $\int \frac{(A+Bx)(d+ex)^3}{a+bx} dx$

Optimal result	1060
Mathematica [A] (verified)	1060
Rubi [A] (verified)	1061
Maple [B] (verified)	1062
Fricas [B] (verification not implemented)	1063
Sympy [B] (verification not implemented)	1063
Maxima [B] (verification not implemented)	1064
Giac [B] (verification not implemented)	1064
Mupad [B] (verification not implemented)	1065
Reduce [B] (verification not implemented)	1066

Optimal result

Integrand size = 20, antiderivative size = 123

$$\int \frac{(A+Bx)(d+ex)^3}{a+bx} dx = \frac{(Ab-aB)e(bd-ae)^2x}{b^4} + \frac{(Ab-aB)(bd-ae)(d+ex)^2}{2b^3} + \frac{(Ab-aB)(d+ex)^3}{3b^2} + \frac{B(d+ex)^4}{4be} + \frac{(Ab-aB)(bd-ae)^3 \log(a+bx)}{b^5}$$

output

```
(A*b-B*a)*e*(-a*e+b*d)^2*x/b^4+1/2*(A*b-B*a)*(-a*e+b*d)*(e*x+d)^2/b^3+1/3*(A*b-B*a)*(e*x+d)^3/b^2+1/4*B*(e*x+d)^4/b/e+(A*b-B*a)*(-a*e+b*d)^3*ln(b*x+a)/b^5
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.37

$$\int \frac{(A+Bx)(d+ex)^3}{a+bx} dx = \frac{bx(-12a^3Be^3 + 6a^2be^2(6Bd + 2Ae + Bex) - 2ab^2e(3Ae(6d + ex) + B(18d^2 + 9dex + 2e^2x^2)) + b^3(2A$$

12b^5

input `Integrate[((A + B*x)*(d + e*x)^3)/(a + b*x), x]`

output
$$(b*x*(-12*a^3*B*e^3 + 6*a^2*b*e^2*(6*B*d + 2*A*e + B*e*x) - 2*a*b^2*e*(3*A*e*(6*d + e*x) + B*(18*d^2 + 9*d*e*x + 2*e^2*x^2)) + b^3*(2*A*e*(18*d^2 + 9*d*e*x + 2*e^2*x^2) + 3*B*(4*d^3 + 6*d^2*e*x + 4*d*e^2*x^2 + e^3*x^3))) + 12*(A*b - a*B)*(b*d - a*e)^3*\text{Log}[a + b*x])/(12*b^5)$$

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)^3}{a + bx} dx$$

↓ 86

$$\int \left(\frac{(Ab - aB)(bd - ae)^3}{b^4(a + bx)} + \frac{e(Ab - aB)(bd - ae)^2}{b^4} + \frac{e(d + ex)(Ab - aB)(bd - ae)}{b^3} + \frac{e(d + ex)^2(Ab - aB)}{b^2} + \dots \right) dx$$

↓ 2009

$$\frac{(Ab - aB)(bd - ae)^3 \log(a + bx)}{b^5} + \frac{ex(Ab - aB)(bd - ae)^2}{b^4} + \frac{(d + ex)^2(Ab - aB)(bd - ae)}{2b^3} + \frac{(d + ex)^3(Ab - aB)}{3b^2} + \frac{B(d + ex)^4}{4be}$$

input `Int[((A + B*x)*(d + e*x)^3)/(a + b*x), x]`

output
$$((A*b - a*B)*e*(b*d - a*e)^2*x)/b^4 + ((A*b - a*B)*(b*d - a*e)*(d + e*x)^2)/(2*b^3) + ((A*b - a*B)*(d + e*x)^3)/(3*b^2) + (B*(d + e*x)^4)/(4*b*e) + ((A*b - a*B)*(b*d - a*e)^3*\text{Log}[a + b*x])/b^5$$

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 260 vs. 2(117) = 234.

Time = 0.21 (sec) , antiderivative size = 261, normalized size of antiderivative = 2.12

method	result
norman	$\frac{(A^2 b e^3 - 3 A a b^2 d e^2 + 3 A b^3 d^2 e - B a^3 e^3 + 3 B a^2 b d e^2 - 3 B a b^2 d^2 e + b^3 B d^3) x}{b^4} - \frac{e(A a b e^2 - 3 A b^2 d e - B a^2 e^2 + 3 B a b d e - 3 b^2 d^2 e)}{2 b^3}$
default	$\frac{\frac{1}{4} b^3 B x^4 e^3 + \frac{1}{3} A b^3 e^3 x^3 - \frac{1}{3} B a b^2 e^3 x^3 + B b^3 d e^2 x^3 - \frac{1}{2} A a b^2 e^3 x^2 + \frac{3}{2} A b^3 d e^2 x^2 + \frac{1}{2} B a^2 b e^3 x^2 - \frac{3}{2} B a b^2 d e^2 x^2 + \frac{3}{2} B b^3 d^2 e x^2 + \frac{3}{2} B b^3 d^2 e x^2}{b^4}$
risch	$\frac{B e^3 x^4}{4 b} + \frac{A e^3 x^3}{3 b} - \frac{B a e^3 x^3}{3 b^2} + \frac{B d e^2 x^3}{b} - \frac{A a e^3 x^2}{2 b^2} + \frac{3 A d e^2 x^2}{2 b} + \frac{B a^2 e^3 x^2}{2 b^3} - \frac{3 B a d e^2 x^2}{2 b^2} + \frac{3 B d^2 e x^2}{2 b} + \frac{A a^2 e^3 x^2}{2 b^3}$
parallelrisch	$-36 A \ln(b x + a) a^2 b^2 d e^2 + 36 A \ln(b x + a) a b^3 d^2 e + 36 A x a b^3 d e^2 - 36 B x a^2 b^2 d e^2 + 36 B x a b^3 d^2 e + 18 B x^2 a b^3 d e^2 + 36 B \ln(b x + a)$

```
input int((B*x+A)*(e*x+d)^3/(b*x+a),x,method=_RETURNVERBOSE)
```

```
output (A*a^2*b*e^3-3*A*a*b^2*d*e^2+3*A*b^3*d^2*e-B*a^3*e^3+3*B*a^2*b*d*e^2-3*B*a*b^2*d^2*e+B*b^3*d^3)/b^4*x-1/2/b^3*e*(A*a*b*e^2-3*A*b^2*d*e-B*a^2*e^2+3*B*a*b*d*e-3*B*b^2*d^2)*x^2+1/3/b^2*e^2*(A*b*e-B*a*e+3*B*b*d)*x^3+1/4/b*B*e^3*x^4-(A*a^3*b*e^3-3*A*a^2*b^2*d*e^2+3*A*a*b^3*d^2*e-A*b^4*d^3-B*a^4*e^3+3*B*a^3*b*d*e^2-3*B*a^2*b^2*d^2*e+B*a*b^3*d^3)/b^5*ln(b*x+a)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 269 vs. $2(119) = 238$.

Time = 0.07 (sec) , antiderivative size = 269, normalized size of antiderivative = 2.19

$$\int \frac{(A + Bx)(d + ex)^3}{a + bx} dx$$

$$= \frac{3Bb^4e^3x^4 + 4(3Bb^4de^2 - (Bab^3 - Ab^4)e^3)x^3 + 6(3Bb^4d^2e - 3(Bab^3 - Ab^4)de^2 + (Ba^2b^2 - Aab^3)e^3)}{b^5}$$

input `integrate((B*x+A)*(e*x+d)^3/(b*x+a),x, algorithm="fricas")`

output
$$\frac{1}{12}*(3*B*b^4*e^3*x^4 + 4*(3*B*b^4*d*e^2 - (B*a*b^3 - A*b^4)*e^3)*x^3 + 6*(3*B*b^4*d^2*e - 3*(B*a*b^3 - A*b^4)*d*e^2 + (B*a^2*b^2 - A*a*b^3)*e^3)*x^2 + 12*(B*b^4*d^3 - 3*(B*a*b^3 - A*b^4)*d^2*e + 3*(B*a^2*b^2 - A*a*b^3)*d*e - (B*a^3*b - A*a^2*b^2)*e^3)*x - 12*((B*a*b^3 - A*b^4)*d^3 - 3*(B*a^2*b^2 - A*a*b^3)*d^2*e + 3*(B*a^3*b - A*a^2*b^2)*d*e^2 - (B*a^4 - A*a^3*b)*e^3)*\log(b*x + a)/b^5$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 221 vs. $2(107) = 214$.

Time = 0.39 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.80

$$\int \frac{(A + Bx)(d + ex)^3}{a + bx} dx = \frac{Be^3x^4}{4b} + x^3 \left(\frac{Ae^3}{3b} - \frac{Bae^3}{3b^2} + \frac{Bde^2}{b} \right)$$

$$+ x^2 \left(-\frac{Aae^3}{2b^2} + \frac{3Ade^2}{2b} + \frac{Ba^2e^3}{2b^3} - \frac{3Bade^2}{2b^2} + \frac{3Bd^2e}{2b} \right)$$

$$+ x \left(\frac{Aa^2e^3}{b^3} - \frac{3Aade^2}{b^2} + \frac{3Ad^2e}{b} - \frac{Ba^3e^3}{b^4} + \frac{3Ba^2de^2}{b^3} - \frac{3Bad^2e}{b^2} + \frac{Bd^3}{b} \right) + \frac{(-Ab + Ba)(ae - bd)^3 \log(a + bx)}{b^5}$$

input `integrate((B*x+A)*(e*x+d)**3/(b*x+a),x)`

output

```
B***3*x**4/(4*b) + x**3*(A***3/(3*b) - B*a***3/(3*b**2) + B*d***2/b) +
x**2*(-A*a***3/(2*b**2) + 3*A*d***2/(2*b) + B*a**2***3/(2*b**3) - 3*B*
a*d***2/(2*b**2) + 3*B*d**2*e/(2*b)) + x*(A*a**2***3/b**3 - 3*A*a*d***2
/b**2 + 3*A*d**2*e/b - B*a**3***3/b**4 + 3*B*a**2*d***2/b**3 - 3*B*a*d**
2*e/b**2 + B*d**3/b) + (-A*b + B*a)*(a*e - b*d)**3*log(a + b*x)/b**5
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 266 vs. 2(119) = 238.

Time = 0.05 (sec) , antiderivative size = 266, normalized size of antiderivative = 2.16

$$\int \frac{(A + Bx)(d + ex)^3}{a + bx} dx$$

$$= \frac{3 B b^3 e^3 x^4 + 4 (3 B b^3 d e^2 - (B a b^2 - A b^3) e^3) x^3 + 6 (3 B b^3 d^2 e - 3 (B a b^2 - A b^3) d e^2 + (B a^2 b - A a b^2) e^3) x^2 + ((B a b^3 - A b^4) d^3 - 3 (B a^2 b^2 - A a b^3) d^2 e + 3 (B a^3 b - A a^2 b^2) d e^2 - (B a^4 - A a^3 b) e^3) \log(bx + a)}{12 b^4 b^5}$$

input

```
integrate((B*x+A)*(e*x+d)^3/(b*x+a),x, algorithm="maxima")
```

output

```
1/12*(3*B*b^3*e^3*x^4 + 4*(3*B*b^3*d*e^2 - (B*a*b^2 - A*b^3)*e^3)*x^3 + 6*
(3*B*b^3*d^2*e - 3*(B*a*b^2 - A*b^3)*d*e^2 + (B*a^2*b - A*a*b^2)*e^3)*x^2
+ 12*(B*b^3*d^3 - 3*(B*a*b^2 - A*b^3)*d^2*e + 3*(B*a^2*b - A*a*b^2)*d*e^2
- (B*a^3 - A*a^2*b)*e^3)*x)/b^4 - ((B*a*b^3 - A*b^4)*d^3 - 3*(B*a^2*b^2 -
A*a*b^3)*d^2*e + 3*(B*a^3*b - A*a^2*b^2)*d*e^2 - (B*a^4 - A*a^3*b)*e^3)*l
og(b*x + a)/b^5
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 298 vs. 2(119) = 238.

Time = 0.12 (sec) , antiderivative size = 298, normalized size of antiderivative = 2.42

$$\int \frac{(A + Bx)(d + ex)^3}{a + bx} dx$$

$$= \frac{3 B b^3 e^3 x^4 + 12 B b^3 d e^2 x^3 - 4 B a b^2 e^3 x^3 + 4 A b^3 e^3 x^3 + 18 B b^3 d^2 e x^2 - 18 B a b^2 d e^2 x^2 + 18 A b^3 d e^2 x^2 + 6 (B a b^3 d^3 - A b^4 d^3 - 3 B a^2 b^2 d^2 e + 3 A a b^3 d^2 e + 3 B a^3 b d e^2 - 3 A a^2 b^2 d e^2 - B a^4 e^3 + A a^3 b e^3) \log(|bx + a|)}{b^5}$$

input `integrate((B*x+A)*(e*x+d)^3/(b*x+a),x, algorithm="giac")`

output
$$\frac{1}{12}*(3*B*b^3*e^3*x^4 + 12*B*b^3*d*e^2*x^3 - 4*B*a*b^2*e^3*x^3 + 4*A*b^3*e^3*x^3 + 18*B*b^3*d^2*e*x^2 - 18*B*a*b^2*d*e^2*x^2 + 18*A*b^3*d*e^2*x^2 + 6*B*a^2*b*e^3*x^2 - 6*A*a*b^2*e^3*x^2 + 12*B*b^3*d^3*x - 36*B*a*b^2*d^2*e*x + 36*A*b^3*d^2*e*x + 36*B*a^2*b*d*e^2*x - 36*A*a*b^2*d*e^2*x - 12*B*a^3*e^3*x + 12*A*a^2*b*e^3*x)/b^4 - (B*a*b^3*d^3 - A*b^4*d^3 - 3*B*a^2*b^2*d^2*e + 3*A*a*b^3*d^2*e + 3*B*a^3*b*d*e^2 - 3*A*a^2*b^2*d*e^2 - B*a^4*e^3 + A*a^3*b*e^3)*\log(\text{abs}(b*x + a))/b^5$$

Mupad [B] (verification not implemented)

Time = 1.00 (sec) , antiderivative size = 268, normalized size of antiderivative = 2.18

$$\int \frac{(A + Bx)(d + ex)^3}{a + bx} dx$$

$$= x \left(\frac{B d^3 + 3 A e d^2}{b} + \frac{a \left(\frac{a \left(\frac{A e^3 + 3 B d e^2}{b} - \frac{B a e^3}{b^2} \right) - \frac{3 d e (A e + B d)}{b}}{b} \right)}{b} \right)$$

$$- x^2 \left(\frac{a \left(\frac{A e^3 + 3 B d e^2}{b} - \frac{B a e^3}{b^2} \right) - \frac{3 d e (A e + B d)}{2 b}}{2 b} \right) + x^3 \left(\frac{A e^3 + 3 B d e^2}{3 b} - \frac{B a e^3}{3 b^2} \right)$$

$$+ \frac{\ln(a + bx) (B a^4 e^3 - 3 B a^3 b d e^2 - A a^3 b e^3 + 3 B a^2 b^2 d^2 e + 3 A a^2 b^2 d e^2 - B a b^3 d^3 - 3 A a b^3 d^2 e + \frac{B e^3 x^4}{4 b})}{b^5}$$

input `int(((A + B*x)*(d + e*x)^3)/(a + b*x),x)`

output
$$x*((B*d^3 + 3*A*d^2*e)/b + (a*((a*((A*e^3 + 3*B*d*e^2)/b - (B*a*e^3)/b^2))/b - (3*d*e*(A*e + B*d))/b))/b - x^2*((a*((A*e^3 + 3*B*d*e^2)/b - (B*a*e^3)/b^2))/(2*b) - (3*d*e*(A*e + B*d))/(2*b)) + x^3*((A*e^3 + 3*B*d*e^2)/(3*b) - (B*a*e^3)/(3*b^2)) + (\log(a + b*x)*(A*b^4*d^3 + B*a^4*e^3 - A*a^3*b*e^3 - B*a*b^3*d^3 + 3*A*a^2*b^2*d*e^2 + 3*B*a^2*b^2*d^2*e - 3*A*a*b^3*d^2*e - 3*B*a^3*b*d*e^2))/b^5 + (B*e^3*x^4)/(4*b)$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.26

$$\int \frac{(A + Bx)(d + ex)^3}{a + bx} dx = \frac{x(e^3 x^3 + 4d e^2 x^2 + 6d^2 ex + 4d^3)}{4}$$

input `int((B*x+A)*(e*x+d)^3/(b*x+a),x)`

output `(x*(4*d**3 + 6*d**2*e*x + 4*d*e**2*x**2 + e**3*x**3))/4`

3.103 $\int \frac{(A+Bx)(d+ex)^2}{a+bx} dx$

Optimal result	1067
Mathematica [A] (verified)	1067
Rubi [A] (verified)	1068
Maple [A] (verified)	1069
Fricas [A] (verification not implemented)	1070
Sympy [A] (verification not implemented)	1070
Maxima [A] (verification not implemented)	1071
Giac [A] (verification not implemented)	1071
Mupad [B] (verification not implemented)	1072
Reduce [B] (verification not implemented)	1072

Optimal result

Integrand size = 20, antiderivative size = 91

$$\int \frac{(A+Bx)(d+ex)^2}{a+bx} dx = \frac{(Ab-aB)e(bd-ae)x}{b^3} + \frac{(Ab-aB)(d+ex)^2}{2b^2} + \frac{B(d+ex)^3}{3be} + \frac{(Ab-aB)(bd-ae)^2 \log(a+bx)}{b^4}$$

output

```
(A*b-B*a)*e*(-a*e+b*d)*x/b^3+1/2*(A*b-B*a)*(e*x+d)^2/b^2+1/3*B*(e*x+d)^3/b
/e+(A*b-B*a)*(-a*e+b*d)^2*ln(b*x+a)/b^4
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.12

$$\int \frac{(A+Bx)(d+ex)^2}{a+bx} dx = \frac{bx(6a^2Be^2 - 3abe(4Bd + 2Ae + Bex) + b^2(3Ae(4d + ex) + 2B(3d^2 + 3dex + e^2x^2))) + 6(Ab - aB)(bd - ae)^2 \log(a + bx)}{6b^4}$$

input

```
Integrate[((A + B*x)*(d + e*x)^2)/(a + b*x), x]
```


output

$$\frac{(b*x*(6*a^2*B*e^2 - 3*a*b*e*(4*B*d + 2*A*e + B*e*x) + b^2*(3*A*e*(4*d + e*x) + 2*B*(3*d^2 + 3*d*e*x + e^2*x^2))) + 6*(A*b - a*B)*(b*d - a*e)^2*\text{Log}[a + b*x]}{(6*b^4)}$$
Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)^2}{a + bx} dx$$

↓ 86

$$\int \left(\frac{(Ab - aB)(bd - ae)^2}{b^3(a + bx)} + \frac{e(Ab - aB)(bd - ae)}{b^3} + \frac{e(d + ex)(Ab - aB)}{b^2} + \frac{B(d + ex)^2}{b} \right) dx$$

↓ 2009

$$\frac{(Ab - aB)(bd - ae)^2 \log(a + bx)}{b^4} + \frac{ex(Ab - aB)(bd - ae)}{b^3} + \frac{(d + ex)^2(Ab - aB)}{2b^2} + \frac{B(d + ex)^3}{3be}$$

input

$$\text{Int}[(A + B*x)*(d + e*x)^2/(a + b*x), x]$$

output

$$\frac{((A*b - a*B)*e*(b*d - a*e)*x)}{b^3} + \frac{((A*b - a*B)*(d + e*x)^2)}{(2*b^2)} + \frac{B*(d + e*x)^3}{(3*b*e)} + \frac{((A*b - a*B)*(b*d - a*e)^2*\text{Log}[a + b*x])}{b^4}$$

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.65

method	result
norman	$-\frac{(Aab e^2 - 2A b^2 de - B a^2 e^2 + 2Babde - b^2 B d^2)x}{b^3} + \frac{B e^2 x^3}{3b} + \frac{e(Abe - Bae + 2Bbd)x^2}{2b^2} + \frac{(A a^2 b e^2 - 2A a b^2 de + A b^3 d^2 - \dots)}{b^3}$
default	$-\frac{\frac{1}{3}b^2 B x^3 e^2 - \frac{1}{2}A b^2 e^2 x^2 + \frac{1}{2}Bab e^2 x^2 - B b^2 de x^2 + Aab e^2 x - 2A b^2 dex - B a^2 e^2 x + 2Babdex - b^2 B d^2 x}{b^3} + \frac{(A a^2 b e^2 - 2A a b^2 de + A b^3 d^2 - \dots)}{b^3}$
risch	$\frac{B e^2 x^3}{3b} + \frac{A e^2 x^2}{2b} - \frac{B a e^2 x^2}{2b^2} + \frac{B d e x^2}{b} - \frac{A a e^2 x}{b^2} + \frac{2A d e x}{b} + \frac{B a^2 e^2 x}{b^3} - \frac{2B a d e x}{b^2} + \frac{B d^2 x}{b} + \frac{\ln(bx+a)A a^2 e^2}{b^3}$
parallelrisch	$\frac{2B e^2 x^3 b^3 + 3A x^2 b^3 e^2 - 3B x^2 a b^2 e^2 + 6B x^2 b^3 de + 6A \ln(bx+a)a^2 b e^2 - 12A \ln(bx+a)a b^2 de + 6A \ln(bx+a)b^3 d^2 - 6A x a b^2 e^2 + \dots}{6b^4}$

```
input int((B*x+A)*(e*x+d)^2/(b*x+a), x, method=_RETURNVERBOSE)
```

```
output -(A*a*b*e^2-2*A*b^2*d*e-B*a^2*e^2+2*B*a*b*d*e-B*b^2*d^2)/b^3*x+1/3*B/b*e^2*x^3+1/2/b^2*e*(A*b*e-B*a*e+2*B*b*d)*x^2+(A*a^2*b*e^2-2*A*a*b^2*d*e+A*b^3*d^2-B*a^3*e^2+2*B*a^2*b*d*e-B*a*b^2*d^2)/b^4*ln(b*x+a)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.74

$$\int \frac{(A + Bx)(d + ex)^2}{a + bx} dx$$

$$= \frac{2 B b^3 e^2 x^3 + 3 (2 B b^3 d e - (B a b^2 - A b^3) e^2) x^2 + 6 (B b^3 d^2 - 2 (B a b^2 - A b^3) d e + (B a^2 b - A a b^2) e^2) x - 6}{6 b^4}$$

input `integrate((B*x+A)*(e*x+d)^2/(b*x+a),x, algorithm="fricas")`output `1/6*(2*B*b^3*e^2*x^3 + 3*(2*B*b^3*d*e - (B*a*b^2 - A*b^3)*e^2)*x^2 + 6*(B*b^3*d^2 - 2*(B*a*b^2 - A*b^3)*d*e + (B*a^2*b - A*a*b^2)*e^2)*x - 6*((B*a*b^2 - A*b^3)*d^2 - 2*(B*a^2*b - A*a*b^2)*d*e + (B*a^3 - A*a^2*b)*e^2)*log(b*x + a))/b^4`**Sympy [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.29

$$\int \frac{(A + Bx)(d + ex)^2}{a + bx} dx = \frac{B e^2 x^3}{3b} + x^2 \left(\frac{A e^2}{2b} - \frac{B a e^2}{2b^2} + \frac{B d e}{b} \right)$$

$$+ x \left(-\frac{A a e^2}{b^2} + \frac{2 A d e}{b} + \frac{B a^2 e^2}{b^3} - \frac{2 B a d e}{b^2} + \frac{B d^2}{b} \right)$$

$$- \frac{(-A b + B a) (a e - b d)^2 \log(a + b x)}{b^4}$$

input `integrate((B*x+A)*(e*x+d)**2/(b*x+a),x)`output `B*e**2*x**3/(3*b) + x**2*(A*e**2/(2*b) - B*a*e**2/(2*b**2) + B*d*e/b) + x*(-A*a*e**2/b**2 + 2*A*d*e/b + B*a**2*e**2/b**3 - 2*B*a*d*e/b**2 + B*d**2/b) - (-A*b + B*a)*(a*e - b*d)**2*log(a + b*x)/b**4`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.70

$$\int \frac{(A + Bx)(d + ex)^2}{a + bx} dx$$

$$= \frac{2 B b^2 e^2 x^3 + 3 (2 B b^2 d e - (B a b - A b^2) e^2) x^2 + 6 (B b^2 d^2 - 2 (B a b - A b^2) d e + (B a^2 - A a b) e^2) x}{b^4} - \frac{6 b^3 ((B a b^2 - A b^3) d^2 - 2 (B a^2 b - A a b^2) d e + (B a^3 - A a^2 b) e^2) \log(bx + a)}{b^4}$$

input `integrate((B*x+A)*(e*x+d)^2/(b*x+a),x, algorithm="maxima")`output
$$\frac{1}{6} * (2 * B * b^2 * e^2 * x^3 + 3 * (2 * B * b^2 * d * e - (B * a * b - A * b^2) * e^2) * x^2 + 6 * (B * b^2 * d^2 - 2 * (B * a * b - A * b^2) * d * e + (B * a^2 - A * a * b) * e^2) * x) / b^3 - ((B * a * b^2 - A * b^3) * d^2 - 2 * (B * a^2 * b - A * a * b^2) * d * e + (B * a^3 - A * a^2 * b) * e^2) * \log(b * x + a) / b^4$$
Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.82

$$\int \frac{(A + Bx)(d + ex)^2}{a + bx} dx$$

$$= \frac{2 B b^2 e^2 x^3 + 6 B b^2 d e x^2 - 3 B a b e^2 x^2 + 3 A b^2 e^2 x^2 + 6 B b^2 d^2 x - 12 B a b d e x + 12 A b^2 d e x + 6 B a^2 e^2 x - 6 A a b e^2}{b^4} - \frac{6 b^3 (B a b^2 d^2 - A b^3 d^2 - 2 B a^2 b d e + 2 A a b^2 d e + B a^3 e^2 - A a^2 b e^2) \log(|bx + a|)}{b^4}$$

input `integrate((B*x+A)*(e*x+d)^2/(b*x+a),x, algorithm="giac")`output
$$\frac{1}{6} * (2 * B * b^2 * e^2 * x^3 + 6 * B * b^2 * d * e * x^2 - 3 * B * a * b * e^2 * x^2 + 3 * A * b^2 * e^2 * x^2 + 6 * B * b^2 * d^2 * x - 12 * B * a * b * d * e * x + 12 * A * b^2 * d * e * x + 6 * B * a^2 * e^2 * x - 6 * A * a * b * e^2) / b^3 - (B * a * b^2 * d^2 - A * b^3 * d^2 - 2 * B * a^2 * b * d * e + 2 * A * a * b^2 * d * e + B * a^3 * e^2 - A * a^2 * b * e^2) * \log(\text{abs}(b * x + a)) / b^4$$

Mupad [B] (verification not implemented)

Time = 0.97 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.75

$$\int \frac{(A + Bx)(d + ex)^2}{a + bx} dx$$

$$= x \left(\frac{Bd^2 + 2Aed}{b} - \frac{a \left(\frac{Ae^2 + 2Bde}{b} - \frac{Bae^2}{b^2} \right)}{b} \right) + x^2 \left(\frac{Ae^2 + 2Bde}{2b} - \frac{Bae^2}{2b^2} \right)$$

$$+ \frac{\ln(a + bx) (-Ba^3e^2 + 2Ba^2bde + Aa^2be^2 - Bab^2d^2 - 2Aab^2de + Ab^3d^2)}{b^4}$$

$$+ \frac{Be^2x^3}{3b}$$

input `int(((A + B*x)*(d + e*x)^2)/(a + b*x),x)`output `x*((B*d^2 + 2*A*d*e)/b - (a*((A*e^2 + 2*B*d*e)/b - (B*a*e^2)/b^2))/b) + x^2*((A*e^2 + 2*B*d*e)/(2*b) - (B*a*e^2)/(2*b^2)) + (log(a + b*x)*(A*b^3*d^2 - B*a^3*e^2 + A*a^2*b*e^2 - B*a*b^2*d^2 - 2*A*a*b^2*d*e + 2*B*a^2*b*d*e))/b^4 + (B*e^2*x^3)/(3*b)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.23

$$\int \frac{(A + Bx)(d + ex)^2}{a + bx} dx = \frac{x(e^2x^2 + 3dex + 3d^2)}{3}$$

input `int((B*x+A)*(e*x+d)^2/(b*x+a),x)`output `(x*(3*d**2 + 3*d*e*x + e**2*x**2))/3`

3.104 $\int \frac{(A+Bx)(d+ex)}{a+bx} dx$

Optimal result	1073
Mathematica [A] (verified)	1073
Rubi [A] (verified)	1074
Maple [A] (verified)	1075
Fricas [A] (verification not implemented)	1075
Sympy [A] (verification not implemented)	1076
Maxima [A] (verification not implemented)	1076
Giac [A] (verification not implemented)	1077
Mupad [B] (verification not implemented)	1077
Reduce [B] (verification not implemented)	1078

Optimal result

Integrand size = 18, antiderivative size = 59

$$\int \frac{(A+Bx)(d+ex)}{a+bx} dx = \frac{B(bd-ae)x}{b^2} + \frac{e(A+Bx)^2}{2bB} + \frac{(Ab-aB)(bd-ae)\log(a+bx)}{b^3}$$

output

```
B*(-a*e+b*d)*x/b^2+1/2*e*(B*x+A)^2/b/B+(A*b-B*a)*(-a*e+b*d)*ln(b*x+a)/b^3
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.95

$$\int \frac{(A+Bx)(d+ex)}{a+bx} dx = \frac{bx(-2aBe + b(2Bd + 2Ae + Bex)) + 2(Ab - aB)(bd - ae)\log(a+bx)}{2b^3}$$

input

```
Integrate[((A + B*x)*(d + e*x))/(a + b*x), x]
```

output

```
(b*x*(-2*a*B*e + b*(2*B*d + 2*A*e + B*e*x)) + 2*(A*b - a*B)*(b*d - a*e)*Log[a + b*x])/(2*b^3)
```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)}{a + bx} dx$$

↓ 86

$$\int \left(\frac{(Ab - aB)(bd - ae)}{b^2(a + bx)} + \frac{B(bd - ae)}{b^2} + \frac{e(A + Bx)}{b} \right) dx$$

↓ 2009

$$\frac{(Ab - aB)(bd - ae) \log(a + bx)}{b^3} + \frac{Bx(bd - ae)}{b^2} + \frac{e(A + Bx)^2}{2bB}$$

input

```
Int[((A + B*x)*(d + e*x))/(a + b*x),x]
```

output

```
(B*(b*d - a*e)*x)/b^2 + (e*(A + B*x)^2)/(2*b*B) + ((A*b - a*B)*(b*d - a*e)*Log[a + b*x])/b^3
```

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.12

method	result	size
default	$\frac{\frac{1}{2}Bbe x^2 + A b e x - B a e x + B b d x}{b^2} + \frac{(-A a b e + A b^2 d + B a^2 e - B a b d) \ln(bx+a)}{b^3}$	66
norman	$\frac{(A b e - B a e + B b d)x}{b^2} + \frac{B e x^2}{2b} - \frac{(A a b e - A b^2 d - B a^2 e + B a b d) \ln(bx+a)}{b^3}$	67
risch	$\frac{B e x^2}{2b} + \frac{A e x}{b} - \frac{B a e x}{b^2} + \frac{B d x}{b} - \frac{\ln(bx+a) A a e}{b^2} + \frac{\ln(bx+a) A d}{b} + \frac{\ln(bx+a) B a^2 e}{b^3} - \frac{\ln(bx+a) B a d}{b^2}$	90
parallelrisch	$-\frac{-B e x^2 b^2 + 2A \ln(bx+a) a b e - 2A \ln(bx+a) b^2 d - 2A x b^2 e - 2B \ln(bx+a) a^2 e + 2B \ln(bx+a) a b d + 2B x a b e - 2B x b^2 d}{2b^3}$	90

input `int((B*x+A)*(e*x+d)/(b*x+a),x,method=_RETURNVERBOSE)`output `1/b^2*(1/2*B*b*e*x^2+A*b*e*x-B*a*e*x+B*b*d*x)+(-A*a*b*e+A*b^2*d+B*a^2*e-B*a*b*d)/b^3*ln(b*x+a)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.27

$$\int \frac{(A + Bx)(d + ex)}{a + bx} dx$$

$$= \frac{Bb^2ex^2 + 2(Bb^2d - (Bab - Ab^2)e)x - 2((Bab - Ab^2)d - (Ba^2 - Aab)e) \log(bx + a)}{2b^3}$$

input `integrate((B*x+A)*(e*x+d)/(b*x+a),x, algorithm="fricas")`output `1/2*(B*b^2*e*x^2 + 2*(B*b^2*d - (B*a*b - A*b^2)*e)*x - 2*((B*a*b - A*b^2)*d - (B*a^2 - A*a*b)*e)*log(b*x + a))/b^3`

Sympy [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.90

$$\int \frac{(A + Bx)(d + ex)}{a + bx} dx = \frac{Bex^2}{2b} + x \left(\frac{Ae}{b} - \frac{Bae}{b^2} + \frac{Bd}{b} \right) + \frac{(-Ab + Ba)(ae - bd) \log(a + bx)}{b^3}$$

input `integrate((B*x+A)*(e*x+d)/(b*x+a),x)`output `B*e*x**2/(2*b) + x*(A*e/b - B*a*e/b**2 + B*d/b) + (-A*b + B*a)*(a*e - b*d)*log(a + b*x)/b**3`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.22

$$\int \frac{(A + Bx)(d + ex)}{a + bx} dx = \frac{Bbx^2 + 2(Bbd - (Ba - Ab)e)x}{2b^2} - \frac{((Bab - Ab^2)d - (Ba^2 - Aab)e) \log(bx + a)}{b^3}$$

input `integrate((B*x+A)*(e*x+d)/(b*x+a),x, algorithm="maxima")`output `1/2*(B*b*e*x^2 + 2*(B*b*d - (B*a - A*b)*e)*x)/b^2 - ((B*a*b - A*b^2)*d - (B*a^2 - A*a*b)*e)*log(b*x + a)/b^3`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.17

$$\int \frac{(A + Bx)(d + ex)}{a + bx} dx = \frac{Bbx^2 + 2Bbdx - 2Baex + 2Abex}{2b^2} - \frac{(Babd - Ab^2d - Ba^2e + Aabe) \log(|bx + a|)}{b^3}$$

input `integrate((B*x+A)*(e*x+d)/(b*x+a),x, algorithm="giac")`output `1/2*(B*b*e*x^2 + 2*B*b*d*x - 2*B*a*e*x + 2*A*b*e*x)/b^2 - (B*a*b*d - A*b^2*d - B*a^2*e + A*a*b*e)*log(abs(b*x + a))/b^3`**Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.15

$$\int \frac{(A + Bx)(d + ex)}{a + bx} dx = x \left(\frac{Ae + Bd}{b} - \frac{Bae}{b^2} \right) + \frac{\ln(a + bx) (Ab^2d + Ba^2e - Aabe - Babd)}{b^3} + \frac{Bex^2}{2b}$$

input `int(((A + B*x)*(d + e*x))/(a + b*x),x)`output `x*((A*e + B*d)/b - (B*a*e)/b^2) + (log(a + b*x)*(A*b^2*d + B*a^2*e - A*a*b*e - B*a*b*d))/b^3 + (B*e*x^2)/(2*b)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.17

$$\int \frac{(A + Bx)(d + ex)}{a + bx} dx = \frac{x(ex + 2d)}{2}$$

input `int((B*x+A)*(e*x+d)/(b*x+a),x)`

output `(x*(2*d + e*x))/2`

3.105 $\int \frac{A+Bx}{a+bx} dx$

Optimal result	1079
Mathematica [A] (verified)	1079
Rubi [A] (verified)	1080
Maple [A] (verified)	1081
Fricas [A] (verification not implemented)	1081
Sympy [A] (verification not implemented)	1081
Maxima [A] (verification not implemented)	1082
Giac [A] (verification not implemented)	1082
Mupad [B] (verification not implemented)	1082
Reduce [B] (verification not implemented)	1083

Optimal result

Integrand size = 13, antiderivative size = 25

$$\int \frac{A+Bx}{a+bx} dx = \frac{Bx}{b} + \frac{(Ab-aB)\log(a+bx)}{b^2}$$

output `B*x/b+(A*b-B*a)*ln(b*x+a)/b^2`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{A+Bx}{a+bx} dx = \frac{Bx}{b} + \frac{(Ab-aB)\log(a+bx)}{b^2}$$

input `Integrate[(A + B*x)/(a + b*x),x]`

output `(B*x)/b + ((A*b - a*B)*Log[a + b*x])/b^2`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{a + bx} dx$$

↓ 49

$$\int \left(\frac{Ab - aB}{b(a + bx)} + \frac{B}{b} \right) dx$$

↓ 2009

$$\frac{(Ab - aB) \log(a + bx)}{b^2} + \frac{Bx}{b}$$

input

```
Int[(A + B*x)/(a + b*x),x]
```

output

```
(B*x)/b + ((A*b - a*B)*Log[a + b*x])/b^2
```

Defintions of rubi rules used

rule 49

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]
```

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04

method	result	size
default	$\frac{Bx}{b} + \frac{(Ab-Ba)\ln(bx+a)}{b^2}$	26
norman	$\frac{Bx}{b} + \frac{(Ab-Ba)\ln(bx+a)}{b^2}$	26
paralelrisch	$\frac{A\ln(bx+a)b - B\ln(bx+a)a + bBx}{b^2}$	29
risch	$\frac{Bx}{b} + \frac{\ln(bx+a)A}{b} - \frac{\ln(bx+a)Ba}{b^2}$	32

input `int((B*x+A)/(b*x+a),x,method=_RETURNVERBOSE)`output `B*x/b+(A*b-B*a)*ln(b*x+a)/b^2`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{A + Bx}{a + bx} dx = \frac{Bbx - (Ba - Ab) \log(bx + a)}{b^2}$$

input `integrate((B*x+A)/(b*x+a),x, algorithm="fricas")`output `(B*b*x - (B*a - A*b)*log(b*x + a))/b^2`**Sympy [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \frac{A + Bx}{a + bx} dx = \frac{Bx}{b} - \frac{(-Ab + Ba) \log(a + bx)}{b^2}$$

input `integrate((B*x+A)/(b*x+a),x)`

output $Bx/b - (-A*b + B*a)*\log(a + b*x)/b**2$

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04

$$\int \frac{A + Bx}{a + bx} dx = \frac{Bx}{b} - \frac{(Ba - Ab) \log(bx + a)}{b^2}$$

input `integrate((B*x+A)/(b*x+a),x, algorithm="maxima")`

output $Bx/b - (B*a - A*b)*\log(b*x + a)/b^2$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.08

$$\int \frac{A + Bx}{a + bx} dx = \frac{Bx}{b} - \frac{(Ba - Ab) \log(|bx + a|)}{b^2}$$

input `integrate((B*x+A)/(b*x+a),x, algorithm="giac")`

output $Bx/b - (B*a - A*b)*\log(\text{abs}(b*x + a))/b^2$

Mupad [B] (verification not implemented)

Time = 0.97 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{A + Bx}{a + bx} dx = \frac{Bx}{b} + \frac{\ln(a + bx) (Ab - Ba)}{b^2}$$

input `int((A + B*x)/(a + b*x),x)`

output $(B*x)/b + (\log(a + b*x)*(A*b - B*a))/b^2$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 1, normalized size of antiderivative = 0.04

$$\int \frac{A + Bx}{a + bx} dx = x$$

input `int((B*x+A)/(b*x+a), x)`

output `x`

3.106 $\int \frac{A+Bx}{(a+bx)(d+ex)} dx$

Optimal result	1084
Mathematica [A] (verified)	1084
Rubi [A] (verified)	1085
Maple [A] (verified)	1086
Fricas [A] (verification not implemented)	1086
Sympy [B] (verification not implemented)	1087
Maxima [A] (verification not implemented)	1087
Giac [A] (verification not implemented)	1088
Mupad [B] (verification not implemented)	1088
Reduce [B] (verification not implemented)	1088

Optimal result

Integrand size = 20, antiderivative size = 57

$$\int \frac{A + Bx}{(a + bx)(d + ex)} dx = \frac{(Ab - aB) \log(a + bx)}{b(bd - ae)} + \frac{(Bd - Ae) \log(d + ex)}{e(bd - ae)}$$

output `(A*b-B*a)*ln(b*x+a)/b/(-a*e+b*d)+(-A*e+B*d)*ln(e*x+d)/e/(-a*e+b*d)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.88

$$\int \frac{A + Bx}{(a + bx)(d + ex)} dx = \frac{(Ab - aB)e \log(a + bx) + b(Bd - Ae) \log(d + ex)}{be(bd - ae)}$$

input `Integrate[(A + B*x)/((a + b*x)*(d + e*x)),x]`

output `((A*b - a*B)*e*Log[a + b*x] + b*(B*d - A*e)*Log[d + e*x])/(b*e*(b*d - a*e))`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)(d + ex)} dx$$

$$\downarrow 86$$

$$\int \left(\frac{Ab - aB}{(a + bx)(bd - ae)} + \frac{Bd - Ae}{(d + ex)(bd - ae)} \right) dx$$

$$\downarrow 2009$$

$$\frac{(Ab - aB) \log(a + bx)}{b(bd - ae)} + \frac{(Bd - Ae) \log(d + ex)}{e(bd - ae)}$$

input `Int[(A + B*x)/((a + b*x)*(d + e*x)),x]`

output `((A*b - a*B)*Log[a + b*x])/(b*(b*d - a*e)) + ((B*d - A*e)*Log[d + e*x])/(e*(b*d - a*e))`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.02

method	result	size
default	$\frac{(-Ab+Ba)\ln(bx+a)}{(ae-db)b} + \frac{(Ae-Bd)\ln(ex+d)}{(ae-db)e}$	58
norman	$\frac{(Ae-Bd)\ln(ex+d)}{(ae-db)e} - \frac{(Ab-Ba)\ln(bx+a)}{(ae-db)b}$	59
parallelrisch	$-\frac{A\ln(bx+a)be - A\ln(ex+d)be - B\ln(bx+a)ae + B\ln(ex+d)bd}{(ae-db)be}$	62
risch	$\frac{\ln(-ex-d)A}{ae-db} - \frac{\ln(-ex-d)Bd}{(ae-db)e} - \frac{\ln(bx+a)A}{ae-db} + \frac{\ln(bx+a)Ba}{(ae-db)b}$	90

input `int((B*x+A)/(b*x+a)/(e*x+d),x,method=_RETURNVERBOSE)`output `(-A*b+B*a)/(a*e-b*d)/b*ln(b*x+a)+(A*e-B*d)/(a*e-b*d)/e*ln(e*x+d)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.93

$$\int \frac{A+Bx}{(a+bx)(d+ex)} dx = -\frac{(Ba-Ab)e \log(bx+a) - (Bbd-Abe) \log(ex+d)}{b^2de - a^2e^2}$$

input `integrate((B*x+A)/(b*x+a)/(e*x+d),x, algorithm="fricas")`output `-((B*a - A*b)*e*log(b*x + a) - (B*b*d - A*b*e)*log(e*x + d))/(b^2*d*e - a*b*e^2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 226 vs. $2(42) = 84$.

Time = 0.89 (sec) , antiderivative size = 226, normalized size of antiderivative = 3.96

$$\int \frac{A + Bx}{(a + bx)(d + ex)} dx$$

$$= -\frac{(-Ae + Bd) \log\left(x + \frac{-Aae - Abd + 2Bad - \frac{a^2 e(-Ae + Bd)}{ae - bd} + \frac{2abd(-Ae + Bd)}{ae - bd} - \frac{b^2 d^2(-Ae + Bd)}{e(ae - bd)}}{-2Abe + Bae + Bbd}\right)}{e(ae - bd)}$$

$$+ \frac{(-Ab + Ba) \log\left(x + \frac{-Aae - Abd + 2Bad + \frac{a^2 e^2(-Ab + Ba)}{b(ae - bd)} - \frac{2ade(-Ab + Ba)}{ae - bd} + \frac{bd^2(-Ab + Ba)}{ae - bd}}{-2Abe + Bae + Bbd}\right)}{b(ae - bd)}$$

input `integrate((B*x+A)/(b*x+a)/(e*x+d),x)`

output `-(-A*e + B*d)*log(x + (-A*a*e - A*b*d + 2*B*a*d - a**2*e*(-A*e + B*d)/(a*e - b*d) + 2*a*b*d*(-A*e + B*d)/(a*e - b*d) - b**2*d**2*(-A*e + B*d)/(e*(a*e - b*d)))/(-2*A*b*e + B*a*e + B*b*d))/(e*(a*e - b*d)) + (-A*b + B*a)*log(x + (-A*a*e - A*b*d + 2*B*a*d + a**2*e**2*(-A*b + B*a)/(b*(a*e - b*d)) - 2*a*d*e*(-A*b + B*a)/(a*e - b*d) + b*d**2*(-A*b + B*a)/(a*e - b*d))/(-2*A*b*e + B*a*e + B*b*d))/(b*(a*e - b*d))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.02

$$\int \frac{A + Bx}{(a + bx)(d + ex)} dx = -\frac{(Ba - Ab) \log(bx + a)}{b^2 d - a b e} + \frac{(Bd - Ae) \log(ex + d)}{b d e - a e^2}$$

input `integrate((B*x+A)/(b*x+a)/(e*x+d),x, algorithm="maxima")`

output `-(B*a - A*b)*log(b*x + a)/(b^2*d - a*b*e) + (B*d - A*e)*log(e*x + d)/(b*d*e - a*e^2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.05

$$\int \frac{A + Bx}{(a + bx)(d + ex)} dx = -\frac{(Ba - Ab) \log(|bx + a|)}{b^2d - abe} + \frac{(Bd - Ae) \log(|ex + d|)}{bde - ae^2}$$

input `integrate((B*x+A)/(b*x+a)/(e*x+d),x, algorithm="giac")`output `-(B*a - A*b)*log(abs(b*x + a))/(b^2*d - a*b*e) + (B*d - A*e)*log(abs(e*x + d))/(b*d*e - a*e^2)`**Mupad [B] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00

$$\int \frac{A + Bx}{(a + bx)(d + ex)} dx = \frac{\ln(d + ex) (Ae - Bd)}{ae^2 - bde} + \frac{\ln(a + bx) (Ab - Ba)}{b^2d - abe}$$

input `int((A + B*x)/((a + b*x)*(d + e*x)),x)`output `(log(d + e*x)*(A*e - B*d))/(a*e^2 - b*d*e) + (log(a + b*x)*(A*b - B*a))/(b^2*d - a*b*e)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.18

$$\int \frac{A + Bx}{(a + bx)(d + ex)} dx = \frac{\log(ex + d)}{e}$$

input `int((B*x+A)/(b*x+a)/(e*x+d),x)`output `log(d + e*x)/e`

3.107 $\int \frac{A+Bx}{(a+bx)(d+ex)^2} dx$

Optimal result	1089
Mathematica [A] (verified)	1089
Rubi [A] (verified)	1090
Maple [A] (verified)	1091
Fricas [A] (verification not implemented)	1091
Sympy [B] (verification not implemented)	1092
Maxima [A] (verification not implemented)	1092
Giac [A] (verification not implemented)	1093
Mupad [B] (verification not implemented)	1093
Reduce [B] (verification not implemented)	1094

Optimal result

Integrand size = 20, antiderivative size = 82

$$\int \frac{A + Bx}{(a + bx)(d + ex)^2} dx = -\frac{Bd - Ae}{e(bd - ae)(d + ex)} + \frac{(Ab - aB) \log(a + bx)}{(bd - ae)^2} - \frac{(Ab - aB) \log(d + ex)}{(bd - ae)^2}$$

output

```
-(-A*e+B*d)/e/(-a*e+b*d)/(e*x+d)+(A*b-B*a)*ln(b*x+a)/(-a*e+b*d)^2-(A*b-B*a)*ln(e*x+d)/(-a*e+b*d)^2
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.98

$$\int \frac{A + Bx}{(a + bx)(d + ex)^2} dx = \frac{Bd - Ae}{e(-bd + ae)(d + ex)} + \frac{(Ab - aB) \log(a + bx)}{(bd - ae)^2} + \frac{(-Ab + aB) \log(d + ex)}{(bd - ae)^2}$$

input

```
Integrate[(A + B*x)/((a + b*x)*(d + e*x)^2),x]
```

output

$$\frac{(B*d - A*e)/(e*(-(b*d) + a*e)*(d + e*x)) + ((A*b - a*B)*\text{Log}[a + b*x])/(b*d - a*e)^2 + ((-(A*b) + a*B)*\text{Log}[d + e*x])/(b*d - a*e)^2}$$
Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)(d + ex)^2} dx$$

↓ 86

$$\int \left(\frac{b(Ab - aB)}{(a + bx)(bd - ae)^2} + \frac{e(aB - Ab)}{(d + ex)(bd - ae)^2} + \frac{Bd - Ae}{(d + ex)^2(bd - ae)} \right) dx$$

↓ 2009

$$-\frac{Bd - Ae}{e(d + ex)(bd - ae)} + \frac{(Ab - aB) \log(a + bx)}{(bd - ae)^2} - \frac{(Ab - aB) \log(d + ex)}{(bd - ae)^2}$$

input

$$\text{Int}[(A + B*x)/((a + b*x)*(d + e*x)^2), x]$$

output

$$-\frac{(B*d - A*e)/(e*(b*d - a*e)*(d + e*x)) + ((A*b - a*B)*\text{Log}[a + b*x])/(b*d - a*e)^2 - ((A*b - a*B)*\text{Log}[d + e*x])/(b*d - a*e)^2}$$
Defintions of rubi rules used

rule 86

$$\text{Int}[(a_. + (b_.)*(x_.))*((c_. + (d_.)*(x_.))^(n_.))*((e_. + (f_.)*(x_.))^(p_.)), x_] := \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;$$

FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.01

method	result
default	$\frac{(Ab-Ba)\ln(bx+a)}{(ae-db)^2} - \frac{Ae-Bd}{(ae-db)e(ex+d)} - \frac{(Ab-Ba)\ln(ex+d)}{(ae-db)^2}$
norman	$\frac{(Ae-Bd)x}{d(ae-db)(ex+d)} + \frac{(Ab-Ba)\ln(bx+a)}{a^2e^2-2abde+b^2d^2} - \frac{(Ab-Ba)\ln(ex+d)}{a^2e^2-2abde+b^2d^2}$
paralelrisch	$\frac{A\ln(bx+a)xb e^2 - A\ln(ex+d)xb e^2 - B\ln(bx+a)xa e^2 + B\ln(ex+d)xa e^2 + A\ln(bx+a)bde - A\ln(ex+d)bde - B\ln(bx+a)ade + B\ln(ex+d)ade}{(a^2e^2-2abde+b^2d^2)(ex+d)e}$
risch	$-\frac{A}{(ae-db)(ex+d)} + \frac{Bd}{(ae-db)e(ex+d)} - \frac{\ln(ex+d)Ab}{a^2e^2-2abde+b^2d^2} + \frac{\ln(ex+d)Ba}{a^2e^2-2abde+b^2d^2} + \frac{\ln(-bx-a)Ab}{a^2e^2-2abde+b^2d^2} - \frac{\ln(-bx-a)Ba}{a^2e^2-2abde+b^2d^2}$

input `int((B*x+A)/(b*x+a)/(e*x+d)^2,x,method=_RETURNVERBOSE)`

output $(A*b-B*a)/(a*e-b*d)^2*\ln(b*x+a)-(A*e-B*d)/(a*e-b*d)/e/(e*x+d)-(A*b-B*a)/(a*e-b*d)^2*\ln(e*x+d)$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.80

$$\int \frac{A+Bx}{(a+bx)(d+ex)^2} dx = \frac{Bbd^2 + Aae^2 - (Ba + Ab)de + ((Ba - Ab)e^2x + (Ba - Ab)de) \log(bx + a) - ((Ba - Ab)e^2x + (Ba - Ab)de) \log(ex + d)}{b^2d^3e - 2abd^2e^2 + a^2de^3 + (b^2d^2e^2 - 2abde^3 + a^2e^4)x}$$

input `integrate((B*x+A)/(b*x+a)/(e*x+d)^2,x, algorithm="fricas")`

output $-(B*b*d^2 + A*a*e^2 - (B*a + A*b)*d*e + ((B*a - A*b)*e^2*x + (B*a - A*b)*d*e)*\log(b*x + a) - ((B*a - A*b)*e^2*x + (B*a - A*b)*d*e)*\log(e*x + d)/(b^2*d^3*e - 2*a*b*d^2*e^2 + a^2*d*e^3 + (b^2*d^2*e^2 - 2*a*b*d*e^3 + a^2*e^4)*x)$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 355 vs. $2(63) = 126$.

Time = 0.73 (sec) , antiderivative size = 355, normalized size of antiderivative = 4.33

$$\int \frac{A + Bx}{(a + bx)(d + ex)^2} dx$$

$$= \frac{(-Ab + Ba) \log \left(x + \frac{-Aabe - Ab^2d + Ba^2e + Babd - \frac{a^3e^3(-Ab+Ba)}{(ae-bd)^2} + \frac{3a^2bde^2(-Ab+Ba)}{(ae-bd)^2} - \frac{3ab^2d^2e(-Ab+Ba)}{(ae-bd)^2} + \frac{b^3d^3(-Ab+Ba)}{(ae-bd)^2}}{-2Ab^2e + 2Babe} \right)}{(ae - bd)^2}$$

$$- \frac{(-Ab + Ba) \log \left(x + \frac{-Aabe - Ab^2d + Ba^2e + Babd + \frac{a^3e^3(-Ab+Ba)}{(ae-bd)^2} - \frac{3a^2bde^2(-Ab+Ba)}{(ae-bd)^2} + \frac{3ab^2d^2e(-Ab+Ba)}{(ae-bd)^2} - \frac{b^3d^3(-Ab+Ba)}{(ae-bd)^2}}{-2Ab^2e + 2Babe} \right)}{(ae - bd)^2}$$

$$+ \frac{-Ae + Bd}{ade^2 - bd^2e + x(ae^3 - bde^2)}$$

input `integrate((B*x+A)/(b*x+a)/(e*x+d)**2,x)`

output

```
(-A*b + B*a)*log(x + (-A*a*b*e - A*b**2*d + B*a**2*e + B*a*b*d - a**3*e**3
*(-A*b + B*a)/(a*e - b*d)**2 + 3*a**2*b*d*e**2*(-A*b + B*a)/(a*e - b*d)**2
- 3*a*b**2*d**2*e*(-A*b + B*a)/(a*e - b*d)**2 + b**3*d**3*(-A*b + B*a)/(a
*e - b*d)**2)/(-2*A*b**2*e + 2*B*a*b*e))/(a*e - b*d)**2 - (-A*b + B*a)*log
(x + (-A*a*b*e - A*b**2*d + B*a**2*e + B*a*b*d + a**3*e**3*(-A*b + B*a)/(a
*e - b*d)**2 - 3*a**2*b*d*e**2*(-A*b + B*a)/(a*e - b*d)**2 + 3*a*b**2*d**2
*e*(-A*b + B*a)/(a*e - b*d)**2 - b**3*d**3*(-A*b + B*a)/(a*e - b*d)**2)/(-
2*A*b**2*e + 2*B*a*b*e))/(a*e - b*d)**2 + (-A*e + B*d)/(a*d*e**2 - b*d**2*
e + x*(a*e**3 - b*d*e**2))
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.45

$$\int \frac{A + Bx}{(a + bx)(d + ex)^2} dx = -\frac{(Ba - Ab) \log(bx + a)}{b^2d^2 - 2abde + a^2e^2} + \frac{(Ba - Ab) \log(ex + d)}{b^2d^2 - 2abde + a^2e^2}$$

$$- \frac{Bd - Ae}{bd^2e - ade^2 + (bde^2 - ae^3)x}$$

input `integrate((B*x+A)/(b*x+a)/(e*x+d)^2,x, algorithm="maxima")`

output
$$-(B*a - A*b)*\log(b*x + a)/(b^2*d^2 - 2*a*b*d*e + a^2*e^2) + (B*a - A*b)*\log(e*x + d)/(b^2*d^2 - 2*a*b*d*e + a^2*e^2) - (B*d - A*e)/(b*d^2*e - a*d*e^2 + (b*d*e^2 - a*e^3)*x)$$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.23

$$\int \frac{A + Bx}{(a + bx)(d + ex)^2} dx = -\frac{(Bae - Abe) \log\left(\left|b - \frac{bd}{ex+d} + \frac{ae}{ex+d}\right|\right)}{b^2d^2e - 2abde^2 + a^2e^3} - \frac{\frac{Bd}{ex+d} - \frac{Ae}{ex+d}}{bde - ae^2}$$

input `integrate((B*x+A)/(b*x+a)/(e*x+d)^2,x, algorithm="giac")`

output
$$-(B*a*e - A*b*e)*\log(\text{abs}(b - b*d/(e*x + d) + a*e/(e*x + d)))/(b^2*d^2*e - 2*a*b*d*e^2 + a^2*e^3) - (B*d/(e*x + d) - A*e/(e*x + d))/(b*d*e - a*e^2)$$

Mupad [B] (verification not implemented)

Time = 1.03 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.16

$$\int \frac{A + Bx}{(a + bx)(d + ex)^2} dx = \frac{2 \operatorname{atanh}\left(\frac{a^2e^2 - b^2d^2}{(ae - bd)^2} + \frac{2bex}{ae - bd}\right) (Ab - Ba)}{(ae - bd)^2} - \frac{Ae - Bd}{e(ae - bd)(d + ex)}$$

input `int((A + B*x)/((a + b*x)*(d + e*x)^2),x)`

output
$$(2*\operatorname{atanh}((a^2*e^2 - b^2*d^2)/(a*e - b*d)^2 + (2*b*e*x)/(a*e - b*d))*(A*b - B*a))/(a*e - b*d)^2 - (A*e - B*d)/(e*(a*e - b*d)*(d + e*x))$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.15

$$\int \frac{A + Bx}{(a + bx)(d + ex)^2} dx = \frac{x}{d(ex + d)}$$

input `int((B*x+A)/(b*x+a)/(e*x+d)^2,x)`

output `x/(d*(d + e*x))`

3.108 $\int \frac{A+Bx}{(a+bx)(d+ex)^3} dx$

Optimal result	1095
Mathematica [A] (verified)	1095
Rubi [A] (verified)	1096
Maple [A] (verified)	1097
Fricas [B] (verification not implemented)	1098
Sympy [B] (verification not implemented)	1098
Maxima [B] (verification not implemented)	1099
Giac [B] (verification not implemented)	1100
Mupad [B] (verification not implemented)	1100
Reduce [B] (verification not implemented)	1101

Optimal result

Integrand size = 20, antiderivative size = 112

$$\int \frac{A+Bx}{(a+bx)(d+ex)^3} dx = \frac{-Bd+ Ae}{2e(bd-ae)(d+ex)^2} + \frac{Ab-aB}{(bd-ae)^2(d+ex)} + \frac{b(Ab-aB)\log(a+bx)}{(bd-ae)^3} - \frac{b(Ab-aB)\log(d+ex)}{(bd-ae)^3}$$

output

```
1/2*(A*e-B*d)/e/(-a*e+b*d)/(e*x+d)^2+(A*b-B*a)/(-a*e+b*d)^2/(e*x+d)+b*(A*b-B*a)*ln(b*x+a)/(-a*e+b*d)^3-b*(A*b-B*a)*ln(e*x+d)/(-a*e+b*d)^3
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.00

$$\int \frac{A+Bx}{(a+bx)(d+ex)^3} dx = \frac{Bd-Ae}{2e(-bd+ae)(d+ex)^2} + \frac{Ab-aB}{(bd-ae)^2(d+ex)} + \frac{b(Ab-aB)\log(a+bx)}{(bd-ae)^3} - \frac{b(Ab-aB)\log(d+ex)}{(bd-ae)^3}$$

input

```
Integrate[(A + B*x)/((a + b*x)*(d + e*x)^3), x]
```

output

$$\frac{(B*d - A*e)/(2*e*(-(b*d) + a*e)*(d + e*x)^2) + (A*b - a*B)/((b*d - a*e)^2*(d + e*x)) + (b*(A*b - a*B)*\text{Log}[a + b*x])/(b*d - a*e)^3 - (b*(A*b - a*B)*\text{Log}[d + e*x])/(b*d - a*e)^3}$$

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)(d + ex)^3} dx$$

↓ 86

$$\int \left(\frac{b^2(Ab - aB)}{(a + bx)(bd - ae)^3} + \frac{be(Ab - aB)}{(d + ex)(ae - bd)^3} + \frac{e(aB - Ab)}{(d + ex)^2(bd - ae)^2} + \frac{Bd - Ae}{(d + ex)^3(bd - ae)} \right) dx$$

↓ 2009

$$\frac{Ab - aB}{(d + ex)(bd - ae)^2} - \frac{Bd - Ae}{2e(d + ex)^2(bd - ae)} + \frac{b(Ab - aB) \log(a + bx)}{(bd - ae)^3} - \frac{b(Ab - aB) \log(d + ex)}{(bd - ae)^3}$$

input

```
Int[(A + B*x)/((a + b*x)*(d + e*x)^3),x]
```

output

$$-1/2*(B*d - A*e)/(e*(b*d - a*e)*(d + e*x)^2) + (A*b - a*B)/((b*d - a*e)^2*(d + e*x)) + (b*(A*b - a*B)*\text{Log}[a + b*x])/(b*d - a*e)^3 - (b*(A*b - a*B)*\text{Log}[d + e*x])/(b*d - a*e)^3$$

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.99

method	result
default	$-\frac{(Ab-Ba)b \ln(bx+a)}{(ae-db)^3} - \frac{Ae-Bd}{2(ae-db)e(ex+d)^2} + \frac{(Ab-Ba)b \ln(ex+d)}{(ae-db)^3} + \frac{Ab-Ba}{(ae-db)^2(ex+d)}$
norman	$\frac{(Ab e^2 - Ba e^2)x}{e(a^2 e^2 - 2abde + b^2 d^2)} - \frac{Aa e^3 - 3Abd e^2 + Bad e^2 + bB d^2 e}{2e^2(a^2 e^2 - 2abde + b^2 d^2)} + \frac{b(Ab-Ba) \ln(ex+d)}{a^3 e^3 - 3a^2 bd e^2 + 3a b^2 d^2 e - b^3 d^3} - \frac{b(Ab-Ba) \ln(bx+a)}{a^3 e^3 - 3a^2 bd e^2 + 3a b^2 d^2 e - b^3 d^3}$
risch	$\frac{e(Ab-Ba)x}{a^2 e^2 - 2abde + b^2 d^2} - \frac{Aa e^2 - 3Abde + Bade + bB d^2}{2e(a^2 e^2 - 2abde + b^2 d^2)} - \frac{b^2 \ln(bx+a)A}{a^3 e^3 - 3a^2 bd e^2 + 3a b^2 d^2 e - b^3 d^3} + \frac{b \ln(bx+a)Ba}{a^3 e^3 - 3a^2 bd e^2 + 3a b^2 d^2 e - b^3 d^3} + \dots$
parallelrisc	$-\frac{2B \ln(bx+a)ab d^2 e^2 + 2B \ln(ex+d)x^2 ab e^4 + 4A \ln(bx+a)x b^2 d e^3 + B a^2 d e^3 + 3A b^2 d^2 e^2 - b^2 B d^3 e - 4Aabd e^3 + a^2 A e^4 - 2 \dots}{(ex+d)^2}$

```
input int((B*x+A)/(b*x+a)/(e*x+d)^3,x,method=_RETURNVERBOSE)
```

```
output -(A*b-B*a)*b/(a*e-b*d)^3*ln(b*x+a)-1/2*(A*e-B*d)/(a*e-b*d)/e/(e*x+d)^2+(A*b-B*a)*b/(a*e-b*d)^3*ln(e*x+d)+(A*b-B*a)/(a*e-b*d)^2/(e*x+d)
```


input `integrate((B*x+A)/(b*x+a)/(e*x+d)**3,x)`

output

$$\begin{aligned}
 & -b*(-A*b + B*a)*\log(x + (-A*a*b**2*e - A*b**3*d + B*a**2*b*e + B*a*b**2*d \\
 & - a**4*b*e**4*(-A*b + B*a)/(a*e - b*d)**3 + 4*a**3*b**2*d*e**3*(-A*b + B*a) \\
 &)/(a*e - b*d)**3 - 6*a**2*b**3*d**2*e**2*(-A*b + B*a)/(a*e - b*d)**3 + 4*a \\
 & *b**4*d**3*e*(-A*b + B*a)/(a*e - b*d)**3 - b**5*d**4*(-A*b + B*a)/(a*e - b \\
 & *d)**3)/(-2*A*b**3*e + 2*B*a*b**2*e))/(a*e - b*d)**3 + b*(-A*b + B*a)*\log(\\
 & x + (-A*a*b**2*e - A*b**3*d + B*a**2*b*e + B*a*b**2*d + a**4*b*e**4*(-A*b \\
 & + B*a)/(a*e - b*d)**3 - 4*a**3*b**2*d*e**3*(-A*b + B*a)/(a*e - b*d)**3 + 6 \\
 & *a**2*b**3*d**2*e**2*(-A*b + B*a)/(a*e - b*d)**3 - 4*a*b**4*d**3*e*(-A*b + \\
 & B*a)/(a*e - b*d)**3 + b**5*d**4*(-A*b + B*a)/(a*e - b*d)**3)/(-2*A*b**3*e \\
 & + 2*B*a*b**2*e))/(a*e - b*d)**3 + (-A*a*e**2 + 3*A*b*d*e - B*a*d*e - B*b* \\
 & d**2 + x*(2*A*b*e**2 - 2*B*a*e**2))/(2*a**2*d**2*e**3 - 4*a*b*d**3*e**2 + \\
 & 2*b**2*d**4*e + x**2*(2*a**2*e**5 - 4*a*b*d*e**4 + 2*b**2*d**2*e**3) + x*(\\
 & 4*a**2*d*e**4 - 8*a*b*d**2*e**3 + 4*b**2*d**3*e**2))
 \end{aligned}$$

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 247 vs. $2(111) = 222$.

Time = 0.04 (sec) , antiderivative size = 247, normalized size of antiderivative = 2.21

$$\begin{aligned}
 & \int \frac{A + Bx}{(a + bx)(d + ex)^3} dx \\
 & = -\frac{(Bab - Ab^2) \log(bx + a)}{b^3d^3 - 3ab^2d^2e + 3a^2bde^2 - a^3e^3} + \frac{(Bab - Ab^2) \log(ex + d)}{b^3d^3 - 3ab^2d^2e + 3a^2bde^2 - a^3e^3} \\
 & \quad - \frac{Bbd^2 + Aae^2 + 2(Ba - Ab)e^2x + (Ba - 3Ab)de}{2(b^2d^4e - 2abd^3e^2 + a^2d^2e^3 + (b^2d^2e^3 - 2abde^4 + a^2e^5)x^2 + 2(b^2d^3e^2 - 2abd^2e^3 + a^2de^4)x)
 \end{aligned}$$

input `integrate((B*x+A)/(b*x+a)/(e*x+d)^3,x, algorithm="maxima")`

output

$$\begin{aligned}
 & -(B*a*b - A*b^2)*\log(b*x + a)/(b^3*d^3 - 3*a*b^2*d^2*e + 3*a^2*b*d*e^2 - a \\
 & ^3*e^3) + (B*a*b - A*b^2)*\log(e*x + d)/(b^3*d^3 - 3*a*b^2*d^2*e + 3*a^2*b*b \\
 & *d*e^2 - a^3*e^3) - 1/2*(B*b*d^2 + A*a*e^2 + 2*(B*a - A*b)*e^2*x + (B*a - 3 \\
 & *A*b)*d*e)/(b^2*d^4*e - 2*a*b*d^3*e^2 + a^2*d^2*e^3 + (b^2*d^2*e^3 - 2*a*b \\
 & *d*e^4 + a^2*e^5)*x^2 + 2*(b^2*d^3*e^2 - 2*a*b*d^2*e^3 + a^2*d*e^4)*x)
 \end{aligned}$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 233 vs. 2(111) = 222.

Time = 0.12 (sec) , antiderivative size = 233, normalized size of antiderivative = 2.08

$$\int \frac{A + Bx}{(a + bx)(d + ex)^3} dx$$

$$= -\frac{(Bab^2 - Ab^3) \log(|bx + a|)}{b^4 d^3 - 3ab^3 d^2 e + 3a^2 b^2 d e^2 - a^3 b e^3} + \frac{(Babe - Ab^2 e) \log(|ex + d|)}{b^3 d^3 e - 3ab^2 d^2 e^2 + 3a^2 b d e^3 - a^3 e^4}$$

$$-\frac{Bb^2 d^3 - 3Ab^2 d^2 e - Ba^2 d e^2 + 4Aabde^2 - Aa^2 e^3 + 2(Babde^2 - Ab^2 d e^2 - Ba^2 e^3 + Aabe^3)x}{2(bd - ae)^3 (ex + d)^2 e}$$

input `integrate((B*x+A)/(b*x+a)/(e*x+d)^3,x, algorithm="giac")`

output

```
-(B*a*b^2 - A*b^3)*log(abs(b*x + a))/(b^4*d^3 - 3*a*b^3*d^2*e + 3*a^2*b^2*d*e^2 - a^3*b*e^3) + (B*a*b*e - A*b^2*e)*log(abs(e*x + d))/(b^3*d^3*e - 3*a*b^2*d^2*e^2 + 3*a^2*b*d*e^3 - a^3*e^4) - 1/2*(B*b^2*d^3 - 3*A*b^2*d^2*e - B*a^2*d*e^2 + 4*A*a*b*d*e^2 - A*a^2*e^3 + 2*(B*a*b*d*e^2 - A*b^2*d*e^2 - B*a^2*e^3 + A*a*b*e^3)*x)/((b*d - a*e)^3*(e*x + d)^2*e)
```

Mupad [B] (verification not implemented)

Time = 1.05 (sec) , antiderivative size = 228, normalized size of antiderivative = 2.04

$$\int \frac{A + Bx}{(a + bx)(d + ex)^3} dx$$

$$= -\frac{\frac{Aae^2 + Bbd^2 - 3Abde + Bade}{2e(a^2e^2 - 2abde + b^2d^2)} - \frac{ex(Ab - Ba)}{a^2e^2 - 2abde + b^2d^2}}{d^2 + 2dex + e^2x^2}$$

$$-\frac{2b \operatorname{atanh}\left(\frac{\left(\frac{a^3e^3 - a^2bde^2 - ab^2d^2e + b^3d^3}{a^2e^2 - 2abde + b^2d^2} + 2bex\right)(a^2e^2 - 2abde + b^2d^2)}{(ae - bd)^3}\right)}{(ae - bd)^3} (Ab - Ba)$$

input `int((A + B*x)/((a + b*x)*(d + e*x)^3),x)`

output

```
- ((A*a*e^2 + B*b*d^2 - 3*A*b*d*e + B*a*d*e)/(2*e*(a^2*e^2 + b^2*d^2 - 2*a*b*d*e)) - (e*x*(A*b - B*a))/(a^2*e^2 + b^2*d^2 - 2*a*b*d*e))/(d^2 + e^2*x^2 + 2*d*e*x) - (2*b*atanh((((a^3*e^3 + b^3*d^3 - a*b^2*d^2*e - a^2*b*d*e^2)/(a^2*e^2 + b^2*d^2 - 2*a*b*d*e) + 2*b*e*x)*(a^2*e^2 + b^2*d^2 - 2*a*b*d*e))/(a*e - b*d)^3)*(A*b - B*a))/(a*e - b*d)^3
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.21

$$\int \frac{A + Bx}{(a + bx)(d + ex)^3} dx = -\frac{1}{2e(e^2x^2 + 2dex + d^2)}$$

input

```
int((B*x+A)/(b*x+a)/(e*x+d)^3,x)
```

output

```
( - 1)/(2*e*(d**2 + 2*d*e*x + e**2*x**2))
```

3.109 $\int \frac{A+Bx}{(a+bx)(d+ex)^4} dx$

Optimal result	1102
Mathematica [A] (verified)	1102
Rubi [A] (verified)	1103
Maple [A] (verified)	1104
Fricas [B] (verification not implemented)	1105
Sympy [B] (verification not implemented)	1105
Maxima [B] (verification not implemented)	1106
Giac [B] (verification not implemented)	1107
Mupad [B] (verification not implemented)	1108
Reduce [B] (verification not implemented)	1108

Optimal result

Integrand size = 20, antiderivative size = 146

$$\int \frac{A+Bx}{(a+bx)(d+ex)^4} dx = \frac{-Bd+ Ae}{3e(bd-ae)(d+ex)^3} + \frac{Ab-aB}{2(bd-ae)^2(d+ex)^2} + \frac{b(Ab-aB)}{(bd-ae)^3(d+ex)} + \frac{b^2(Ab-aB)\log(a+bx)}{(bd-ae)^4} - \frac{b^2(Ab-aB)\log(d+ex)}{(bd-ae)^4}$$

output

```
1/3*(A*e-B*d)/e/(-a*e+b*d)/(e*x+d)^3+1/2*(A*b-B*a)/(-a*e+b*d)^2/(e*x+d)^2+
b*(A*b-B*a)/(-a*e+b*d)^3/(e*x+d)+b^2*(A*b-B*a)*ln(b*x+a)/(-a*e+b*d)^4-b^2*
(A*b-B*a)*ln(e*x+d)/(-a*e+b*d)^4
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.99

$$\int \frac{A+Bx}{(a+bx)(d+ex)^4} dx = \frac{Bd-Ae}{3e(-bd+ae)(d+ex)^3} + \frac{Ab-aB}{2(bd-ae)^2(d+ex)^2} + \frac{b(Ab-aB)}{(bd-ae)^3(d+ex)} + \frac{b^2(Ab-aB)\log(a+bx)}{(bd-ae)^4} + \frac{b^2(-Ab+aB)\log(d+ex)}{(bd-ae)^4}$$

input `Integrate[(A + B*x)/((a + b*x)*(d + e*x)^4), x]`

output
$$\frac{(B*d - A*e)}{3*e*(-(b*d) + a*e)*(d + e*x)^3} + \frac{(A*b - a*B)}{2*(b*d - a*e)^2*(d + e*x)^2} + \frac{(b*(A*b - a*B))}{(b*d - a*e)^3*(d + e*x)} + \frac{(b^2*(A*b - a*B)*\text{Log}[a + b*x])}{(b*d - a*e)^4} + \frac{(b^2*(-(A*b) + a*B)*\text{Log}[d + e*x])}{(b*d - a*e)^4}$$

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)(d + ex)^4} dx$$

↓ 86

$$\int \left(\frac{b^3(Ab - aB)}{(a + bx)(bd - ae)^4} - \frac{b^2e(Ab - aB)}{(d + ex)(ae - bd)^4} + \frac{be(Ab - aB)}{(d + ex)^2(ae - bd)^3} + \frac{e(aB - Ab)}{(d + ex)^3(bd - ae)^2} + \frac{Bd - Ae}{(d + ex)^4(bd - ae)} \right) dx$$

↓ 2009

$$\frac{b^2(Ab - aB) \log(a + bx)}{(bd - ae)^4} - \frac{b^2(Ab - aB) \log(d + ex)}{(bd - ae)^4} + \frac{b(Ab - aB)}{(d + ex)(bd - ae)^3} + \frac{Ab - aB}{2(d + ex)^2(bd - ae)^2} - \frac{Bd - Ae}{3e(d + ex)^3(bd - ae)}$$

input `Int[(A + B*x)/((a + b*x)*(d + e*x)^4), x]`

output
$$-1/3*(B*d - A*e)/(e*(b*d - a*e)*(d + e*x)^3) + (A*b - a*B)/(2*(b*d - a*e)^2*(d + e*x)^2) + (b*(A*b - a*B))/(b*d - a*e)^3*(d + e*x) + (b^2*(A*b - a*B)*\text{Log}[a + b*x])/(b*d - a*e)^4 - (b^2*(A*b - a*B)*\text{Log}[d + e*x])/(b*d - a*e)^4$$

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.99

method	result
default	$\frac{(Ab-Ba)b^2 \ln(bx+a)}{(ae-db)^4} - \frac{Ae-Bd}{3(ae-db)e(ex+d)^3} - \frac{(Ab-Ba)b}{(ae-db)^3(ex+d)} + \frac{Ab-Ba}{2(ae-db)^2(ex+d)^2} - \frac{(Ab-Ba)b^2 \ln(ex+d)}{(ae-db)^4}$
norman	$-\frac{2Aa^2e^5-7Aabd e^4+11Ab^2d^2e^3+Ba^2de^4-5Babd^2e^3-2Bb^2d^3e^2}{6e^3(a^3e^3-3a^2bde^2+3ab^2d^2e-b^3d^3)} - \frac{(Ab^2e^3-Babe^3)x^2}{e(a^3e^3-3a^2bde^2+3ab^2d^2e-b^3d^3)} + \frac{(Aabe^4-5Ab^2de^3-Ba^2e^2)}{2e^2(a^3e^3-3a^2bde^2+3ab^2d^2e-b^3d^3)}$
risch	$-\frac{be^2(Ab-Ba)x^2}{a^3e^3-3a^2bde^2+3ab^2d^2e-b^3d^3} + \frac{(ae-5db)e(Ab-Ba)x}{2a^3e^3-6a^2bde^2+6ab^2d^2e-2b^3d^3} - \frac{2a^2Ae^3-7Aabd e^2+11Ab^2d^2e+Ba^2de^2-5Babd^2e-2b^2Bd^3}{6e(a^3e^3-3a^2bde^2+3ab^2d^2e-b^3d^3)}$
parallelrisc	$-\frac{18B \ln(bx+a)x^2ab^2de^5+18B \ln(ex+d)x^2ab^2de^5-18B \ln(bx+a)xa b^2d^2e^4+18B \ln(ex+d)xa b^2d^2e^4-3Bxa^3e^6+6Bx^2a^2e^5}{(ex+d)^3}$

```
input int((B*x+A)/(b*x+a)/(e*x+d)^4,x,method=_RETURNVERBOSE)
```

```
output (A*b-B*a)*b^2/(a*e-b*d)^4*ln(b*x+a)-1/3*(A*e-B*d)/(a*e-b*d)/e/(e*x+d)^3-(A*b-B*a)*b/(a*e-b*d)^3/(e*x+d)+1/2*(A*b-B*a)/(a*e-b*d)^2/(e*x+d)^2-(A*b-B*a)*b^2/(a*e-b*d)^4*ln(e*x+d)
```


output

```

-(B*a*b^2 - A*b^3)*log(b*x + a)/(b^4*d^4 - 4*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2 - 4*a^3*b*d*e^3 + a^4*e^4) + (B*a*b^2 - A*b^3)*log(e*x + d)/(b^4*d^4 - 4*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2 - 4*a^3*b*d*e^3 + a^4*e^4) - 1/6*(2*B*b^2*d^3 - 2*A*a^2*e^3 + 6*(B*a*b - A*b^2)*e^3*x^2 + (5*B*a*b - 11*A*b^2)*d^2*e - (B*a^2 - 7*A*a*b)*d*e^2 + 3*(5*(B*a*b - A*b^2)*d*e^2 - (B*a^2 - A*a*b)*e^3)*x)/(b^3*d^6*e - 3*a*b^2*d^5*e^2 + 3*a^2*b*d^4*e^3 - a^3*d^3*e^4 + (b^3*d^3*e^4 - 3*a*b^2*d^2*e^5 + 3*a^2*b*d*e^6 - a^3*e^7)*x^3 + 3*(b^3*d^4*e^3 - 3*a*b^2*d^3*e^4 + 3*a^2*b*d^2*e^5 - a^3*d*e^6)*x^2 + 3*(b^3*d^5*e^2 - 3*a*b^2*d^4*e^3 + 3*a^2*b*d^3*e^4 - a^3*d^2*e^5)*x)

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 376 vs. $2(143) = 286$.

Time = 0.13 (sec) , antiderivative size = 376, normalized size of antiderivative = 2.58

$$\int \frac{A + Bx}{(a + bx)(d + ex)^4} dx = -\frac{(Bab^3 - Ab^4) \log(|bx + a|)}{b^5d^4 - 4ab^4d^3e + 6a^2b^3d^2e^2 - 4a^3b^2de^3 + a^4be^4} + \frac{(Bab^2e - Ab^3e) \log(|ex + d|)}{b^4d^4e - 4ab^3d^3e^2 + 6a^2b^2d^2e^3 - 4a^3bde^4 + a^4e^5} - \frac{2Bb^3d^4 + 3Bab^2d^3e - 11Ab^3d^3e - 6Ba^2bd^2e^2 + 18Aab^2d^2e^2 + Ba^3de^3 - 9Aa^2bde^3 + 2Aa^3e^4 + 6(Ba^2b^2d^2e^2 - 5Aab^3d^2e^2 - 6Ba^2bde^3 + 6Aa^2b^2d^2e^3 + Ba^3e^4 - Aa^2b^2e^4)*x}{6(b^5d^4 - 4ab^4d^3e + 6a^2b^3d^2e^2 - 4a^3b^2de^3 + a^4be^4)}$$

input

```
integrate((B*x+A)/(b*x+a)/(e*x+d)^4,x, algorithm="giac")
```

output

```

-(B*a*b^3 - A*b^4)*log(abs(b*x + a))/(b^5*d^4 - 4*a*b^4*d^3*e + 6*a^2*b^3*d^2*e^2 - 4*a^3*b^2*d^2*e^3 + a^4*b*e^4) + (B*a*b^2*e - A*b^3*e)*log(abs(e*x + d))/(b^4*d^4*e - 4*a*b^3*d^3*e^2 + 6*a^2*b^2*d^2*e^3 - 4*a^3*b*d*e^4 + a^4*e^5) - 1/6*(2*B*b^3*d^4 + 3*B*a*b^2*d^3*e - 11*A*b^3*d^3*e - 6*B*a^2*b*d^2*e^2 + 18*A*a*b^2*d^2*e^2 + B*a^3*d*e^3 - 9*A*a^2*b*d*e^3 + 2*A*a^3*e^4 + 6*(B*a*b^2*d^2*e^3 - A*b^3*d^2*e^3 - B*a^2*b*e^4 + A*a*b^2*e^4)*x^2 + 3*(5*B*a*b^2*d^2*e^2 - 5*A*b^3*d^2*e^2 - 6*B*a^2*b*d*e^3 + 6*A*a*b^2*d^2*e^3 + B*a^3*e^4 - A*a^2*b*e^4)*x)/((b*d - a*e)^4*(e*x + d)^3*e)

```


Mupad [B] (verification not implemented)

Time = 1.17 (sec) , antiderivative size = 399, normalized size of antiderivative = 2.73

$$\int \frac{A + Bx}{(a + bx)(d + ex)^4} dx$$

$$= \frac{2b^2 \operatorname{atanh}\left(\frac{\left(\frac{a^4 e^4 - 2a^3 b d e^3 + 2a b^3 d^3 e - b^4 d^4}{a^3 e^3 - 3a^2 b d e^2 + 3a b^2 d^2 e - b^3 d^3} + 2b e x\right) (a^3 e^3 - 3a^2 b d e^2 + 3a b^2 d^2 e - b^3 d^3)}{(a e - b d)^4}\right) (A b - B a)}{(a e - b d)^4}$$

$$- \frac{B a^2 d e^2 + 2A a^2 e^3 - 5B a b d^2 e - 7A a b d e^2 - 2B b^2 d^3 + 11A b^2 d^2 e}{6e(a^3 e^3 - 3a^2 b d e^2 + 3a b^2 d^2 e - b^3 d^3)} - \frac{x(A b - B a)(a e^2 - 5b d e)}{2(a^3 e^3 - 3a^2 b d e^2 + 3a b^2 d^2 e - b^3 d^3)} + \frac{b e^2 x^2 (A b - B a)}{a^3 e^3 - 3a^2 b d e^2 + 3a b^2 d^2 e - b^3 d^3}$$

$$d^3 + 3d^2 e x + 3d e^2 x^2 + e^3 x^3$$

input `int((A + B*x)/((a + b*x)*(d + e*x)^4),x)`

output

```
(2*b^2*atanh((((a^4*e^4 - b^4*d^4 + 2*a*b^3*d^3*e - 2*a^3*b*d*e^3)/(a^3*e^3 - b^3*d^3 + 3*a*b^2*d^2*e - 3*a^2*b*d*e^2) + 2*b*e*x)*(a^3*e^3 - b^3*d^3 + 3*a*b^2*d^2*e - 3*a^2*b*d*e^2))/(a*e - b*d)^4)*(A*b - B*a))/(a*e - b*d)^4 - (((2*A*a^2*e^3 - 2*B*b^2*d^3 + 11*A*b^2*d^2*e + B*a^2*d*e^2 - 7*A*a*b*d*e^2 - 5*B*a*b*d^2*e)/(6*e*(a^3*e^3 - b^3*d^3 + 3*a*b^2*d^2*e - 3*a^2*b*d*e^2)) - (x*(A*b - B*a)*(a*e^2 - 5*b*d*e))/(2*(a^3*e^3 - b^3*d^3 + 3*a*b^2*d^2*e - 3*a^2*b*d*e^2)) + (b*e^2*x^2*(A*b - B*a))/(a^3*e^3 - b^3*d^3 + 3*a*b^2*d^2*e - 3*a^2*b*d*e^2))/(d^3 + e^3*x^3 + 3*d*e^2*x^2 + 3*d^2*e*x))
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.23

$$\int \frac{A + Bx}{(a + bx)(d + ex)^4} dx = -\frac{1}{3e(e^3 x^3 + 3d e^2 x^2 + 3d^2 e x + d^3)}$$

input `int((B*x+A)/(b*x+a)/(e*x+d)^4,x)`

output

```
( - 1)/(3*e*(d**3 + 3*d**2*e*x + 3*d*e**2*x**2 + e**3*x**3))
```

3.110 $\int \frac{(A+Bx)(d+ex)^3}{(a+bx)^2} dx$

Optimal result	1109
Mathematica [A] (verified)	1109
Rubi [A] (verified)	1110
Maple [A] (verified)	1111
Fricas [B] (verification not implemented)	1112
Sympy [A] (verification not implemented)	1113
Maxima [A] (verification not implemented)	1113
Giac [B] (verification not implemented)	1114
Mupad [B] (verification not implemented)	1115
Reduce [B] (verification not implemented)	1115

Optimal result

Integrand size = 20, antiderivative size = 145

$$\int \frac{(A+Bx)(d+ex)^3}{(a+bx)^2} dx = \frac{3e(bd-ae)(bBd+Abe-2aBe)x}{b^4} - \frac{(Ab-aB)(bd-ae)^3}{b^5(a+bx)} + \frac{e^2(3bBd+Abe-4aBe)(a+bx)^2}{2b^5} + \frac{Be^3(a+bx)^3}{3b^5} + \frac{(bd-ae)^2(bBd+3Abe-4aBe)\log(a+bx)}{b^5}$$

output

```
3*e*(-a*e+b*d)*(A*b*e-2*B*a*e+B*b*d)*x/b^4-(A*b-B*a)*(-a*e+b*d)^3/b^5/(b*x+a)+1/2*e^2*(A*b*e-4*B*a*e+3*B*b*d)*(b*x+a)^2/b^5+1/3*B*e^3*(b*x+a)^3/b^5+(-a*e+b*d)^2*(3*A*b*e-4*B*a*e+B*b*d)*ln(b*x+a)/b^5
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.72

$$\int \frac{(A+Bx)(d+ex)^3}{(a+bx)^2} dx = \frac{B(-6a^4e^3+18a^3be^2(d+ex)+6a^2b^2e(-3d^2-6dex+2e^2x^2))+b^4ex^2(18d^2+9dex+2e^2x^2)+ab^3(6d^3$$

input `Integrate[((A + B*x)*(d + e*x)^3)/(a + b*x)^2,x]`

output
$$\begin{aligned} & (B*(-6*a^4*e^3 + 18*a^3*b*e^2*(d + e*x) + 6*a^2*b^2*e*(-3*d^2 - 6*d*e*x + \\ & 2*e^2*x^2) + b^4*e*x^2*(18*d^2 + 9*d*e*x + 2*e^2*x^2) + a*b^3*(6*d^3 + 18* \\ & d^2*e*x - 27*d*e^2*x^2 - 4*e^3*x^3)) - 3*A*b*(-2*a^3*e^3 + 2*a^2*b*e^2*(3* \\ & d + 2*e*x) + 3*a*b^2*e*(-2*d^2 - 2*d*e*x + e^2*x^2) + b^3*(2*d^3 - 6*d*e^2 \\ & *x^2 - e^3*x^3)) + 6*(b*d - a*e)^2*(b*B*d + 3*A*b*e - 4*a*B*e)*(a + b*x)*L \\ & \text{og}[a + b*x])/(6*b^5*(a + b*x)) \end{aligned}$$

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(A + Bx)(d + ex)^3}{(a + bx)^2} dx \\ & \quad \downarrow 86 \\ & \int \left(\frac{e^2(a + bx)(-4aBe + Abe + 3bBd)}{b^4} + \frac{(bd - ae)^2(-4aBe + 3Abe + bBd)}{b^4(a + bx)} + \frac{(Ab - aB)(bd - ae)^3}{b^4(a + bx)^2} + \frac{3e(bd - ae)^2}{b^4} \right) dx \\ & \quad \downarrow 2009 \\ & \frac{e^2(a + bx)^2(-4aBe + Abe + 3bBd)}{2b^5} - \frac{(Ab - aB)(bd - ae)^3}{b^5(a + bx)} + \\ & \frac{(bd - ae)^2 \log(a + bx)(-4aBe + 3Abe + bBd)}{b^5} + \frac{3ex(bd - ae)(-2aBe + Abe + bBd)}{b^4} + \\ & \frac{Be^3(a + bx)^3}{3b^5} \end{aligned}$$

input `Int[((A + B*x)*(d + e*x)^3)/(a + b*x)^2,x]`

```
output (3*e*(b*d - a*e)*(b*B*d + A*b*e - 2*a*B*e)*x)/b^4 - ((A*b - a*B)*(b*d - a*
e)^3)/(b^5*(a + b*x)) + (e^2*(3*b*B*d + A*b*e - 4*a*B*e)*(a + b*x)^2)/(2*b
^5) + (B*e^3*(a + b*x)^3)/(3*b^5) + ((b*d - a*e)^2*(b*B*d + 3*A*b*e - 4*a*
B*e)*Log[a + b*x])/b^5
```

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && ( !IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 277, normalized size of antiderivative = 1.91

method	result
norman	$\frac{B e^3 x^4}{3b} - \frac{e(3Aab e^2 - 6A b^2 d e - 4B a^2 e^2 + 9Babde - 6b^2 B d^2)x^2}{2b^3} + \frac{e^2(3Abe - 4Bae + 9Bbd)x^3}{6b^2} - \frac{(3A a^3 b e^3 - 6A a^2 b^2 d e^2 + 3Aa b^3 d^2 e - A b^4 d^3)}{b^4(bx+a)}$
default	$-\frac{e(-\frac{1}{3}b^2 B x^3 e^2 - \frac{1}{2}A b^2 e^2 x^2 + Bab e^2 x^2 - \frac{3}{2}B b^2 d e x^2 + 2Aab e^2 x - 3A b^2 d e x - 3B a^2 e^2 x + 6Babd e x - 3b^2 B d^2 x)}{b^4} - \frac{-A a^3 b d^3}{b^4(bx+a)}$
risch	$\frac{e^3 B x^3}{3b^2} + \frac{e^3 A x^2}{2b^2} - \frac{e^3 B a x^2}{b^3} + \frac{3e^2 B d x^2}{2b^2} - \frac{2e^3 A a x}{b^3} + \frac{3e^2 A d x}{b^2} + \frac{3e^3 B a^2 x}{b^4} - \frac{6e^2 B a d x}{b^3} + \frac{3e B d^2 x}{b^2} + \frac{A a^3 e^3}{b^4(bx+a)}$
parallelrisch	$-36A \ln(bx+a) a^2 b^2 d e^2 + 18A \ln(bx+a) a b^3 d^2 e - 36B a^2 b^2 d^2 e + 18A a^3 b e^3 + 54B a^3 b d e^2 - 36A a^2 b^2 d e^2 + 18A a b^3 d^2 e + 6Ba b^4 d^3$

```
input int((B*x+A)*(e*x+d)^3/(b*x+a)^2,x,method=_RETURNVERBOSE)
```

output

```
(1/3/b*B*e^3*x^4-1/2*e*(3*A*a*b*e^2-6*A*b^2*d*e-4*B*a^2*e^2+9*B*a*b*d*e-6*
B*b^2*d^2)/b^3*x^2+1/6*e^2*(3*A*b*e-4*B*a*e+9*B*b*d)/b^2*x^3-(3*A*a^3*b*e^
3-6*A*a^2*b^2*d*e^2+3*A*a*b^3*d^2*e-A*b^4*d^3-4*B*a^4*e^3+9*B*a^3*b*d*e^2-
6*B*a^2*b^2*d^2*e+B*a*b^3*d^3)/a/b^4*x)/(b*x+a)+1/b^5*(3*A*a^2*b*e^3-6*A*a
*b^2*d*e^2+3*A*b^3*d^2*e-4*B*a^3*e^3+9*B*a^2*b*d*e^2-6*B*a*b^2*d^2*e+B*b^3
*d^3)*ln(b*x+a)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 417 vs. $2(140) = 280$.

Time = 0.07 (sec) , antiderivative size = 417, normalized size of antiderivative = 2.88

$$\int \frac{(A + Bx)(d + ex)^3}{(a + bx)^2} dx$$

$$= \frac{2Bb^4e^3x^4 + 6(Bab^3 - Ab^4)d^3 - 18(Ba^2b^2 - Aab^3)d^2e + 18(Ba^3b - Aa^2b^2)de^2 - 6(Ba^4 - Aa^3b)e^3 + ($$

input

```
integrate((B*x+A)*(e*x+d)^3/(b*x+a)^2,x, algorithm="fricas")
```

output

```
1/6*(2*B*b^4*e^3*x^4 + 6*(B*a*b^3 - A*b^4)*d^3 - 18*(B*a^2*b^2 - A*a*b^3)*
d^2*e + 18*(B*a^3*b - A*a^2*b^2)*d*e^2 - 6*(B*a^4 - A*a^3*b)*e^3 + (9*B*b^
4*d*e^2 - (4*B*a*b^3 - 3*A*b^4)*e^3)*x^3 + 3*(6*B*b^4*d^2*e - 3*(3*B*a*b^3
- 2*A*b^4)*d*e^2 + (4*B*a^2*b^2 - 3*A*a*b^3)*e^3)*x^2 + 6*(3*B*a*b^3*d^2*
e - 3*(2*B*a^2*b^2 - A*a*b^3)*d*e^2 + (3*B*a^3*b - 2*A*a^2*b^2)*e^3)*x + 6
*(B*a*b^3*d^3 - 3*(2*B*a^2*b^2 - A*a*b^3)*d^2*e + 3*(3*B*a^3*b - 2*A*a^2*b
^2)*d*e^2 - (4*B*a^4 - 3*A*a^3*b)*e^3 + (B*b^4*d^3 - 3*(2*B*a*b^3 - A*b^4)
*d^2*e + 3*(3*B*a^2*b^2 - 2*A*a*b^3)*d*e^2 - (4*B*a^3*b - 3*A*a^2*b^2)*e^3
)*x)*log(b*x + a))/(b^6*x + a*b^5)
```

Sympy [A] (verification not implemented)

Time = 0.75 (sec) , antiderivative size = 257, normalized size of antiderivative = 1.77

$$\int \frac{(A+Bx)(d+ex)^3}{(a+bx)^2} dx = \frac{Be^3x^3}{3b^2} + x^2 \left(\frac{Ae^3}{2b^2} - \frac{Bae^3}{b^3} + \frac{3Bde^2}{2b^2} \right) + x \left(-\frac{2Aae^3}{b^3} + \frac{3Ade^2}{b^2} + \frac{3Ba^2e^3}{b^4} - \frac{6Bade^2}{b^3} + \frac{3Bd^2e}{b^2} \right) + \frac{Aa^3be^3 - 3Aa^2b^2de^2 + 3Aab^3d^2e - Ab^4d^3 - Ba^4e^3 + 3Ba^3bde^2 - 3Ba^2b^2d^2e + Bab^3d^3}{ab^5 + b^6x} - \frac{(ae - bd)^2 (-3Abe + 4Bae - Bbd) \log(a + bx)}{b^5}$$

input `integrate((B*x+A)*(e*x+d)**3/(b*x+a)**2,x)`output `B*e**3*x**3/(3*b**2) + x**2*(A*e**3/(2*b**2) - B*a*e**3/b**3 + 3*B*d*e**2/(2*b**2)) + x*(-2*A*a*e**3/b**3 + 3*A*d*e**2/b**2 + 3*B*a**2*e**3/b**4 - 6*B*a*d*e**2/b**3 + 3*B*d**2*e/b**2) + (A*a**3*b*e**3 - 3*A*a**2*b**2*d*e**2 + 3*A*a*b**3*d**2*e - A*b**4*d**3 - B*a**4*e**3 + 3*B*a**3*b*d*e**2 - 3*B*a**2*b**2*d**2*e + B*a*b**3*d**3)/(a*b**5 + b**6*x) - (a*e - b*d)**2*(-3*A*b*e + 4*B*a*e - B*b*d)*log(a + b*x)/b**5`**Maxima [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.88

$$\int \frac{(A+Bx)(d+ex)^3}{(a+bx)^2} dx = \frac{(Bab^3 - Ab^4)d^3 - 3(Ba^2b^2 - Aab^3)d^2e + 3(Ba^3b - Aa^2b^2)de^2 - (Ba^4 - Aa^3b)e^3}{b^6x + ab^5} + \frac{2Bb^2e^3x^3 + 3(3Bb^2de^2 - (2Bab - Ab^2)e^3)x^2 + 6(3Bb^2d^2e - 3(2Bab - Ab^2)de^2 + (3Ba^2 - 2Aab)de^2 - (3Bb^3d^3 - 3(2Bab^2 - Ab^3)d^2e + 3(3Ba^2b - 2Aab^2)de^2 - (4Ba^3 - 3Aa^2b)e^3) \log(bx + a)}{6b^4} + \frac{(Bb^3d^3 - 3(2Bab^2 - Ab^3)d^2e + 3(3Ba^2b - 2Aab^2)de^2 - (4Ba^3 - 3Aa^2b)e^3) \log(bx + a)}{b^5}$$

input `integrate((B*x+A)*(e*x+d)^3/(b*x+a)^2,x, algorithm="maxima")`

output

$$\begin{aligned} & ((B*a*b^3 - A*b^4)*d^3 - 3*(B*a^2*b^2 - A*a*b^3)*d^2*e + 3*(B*a^3*b - A*a^2*b^2)*d*e^2 - (B*a^4 - A*a^3*b)*e^3)/(b^6*x + a*b^5) + 1/6*(2*B*b^2*e^3*x^3 + 3*(3*B*b^2*d*e^2 - (2*B*a*b - A*b^2)*e^3)*x^2 + 6*(3*B*b^2*d^2*e - 3*(2*B*a*b - A*b^2)*d*e^2 + (3*B*a^2 - 2*A*a*b)*e^3)*x)/b^4 + (B*b^3*d^3 - 3*(2*B*a*b^2 - A*b^3)*d^2*e + 3*(3*B*a^2*b - 2*A*a*b^2)*d*e^2 - (4*B*a^3 - 3*A*a^2*b)*e^3)*log(b*x + a)/b^5 \end{aligned}$$
Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 370 vs. $2(140) = 280$.

Time = 0.12 (sec) , antiderivative size = 370, normalized size of antiderivative = 2.55

$$\begin{aligned} & \int \frac{(A + Bx)(d + ex)^3}{(a + bx)^2} dx \\ & = \frac{\left(2Be^3 + \frac{3(3Bb^2de^2 - 4Babe^3 + Ab^2e^3)}{(bx+a)b} + \frac{18(Bb^4d^2e - 3Bab^3de^2 + Ab^4de^2 + 2Ba^2b^2e^3 - Aab^3e^3)}{(bx+a)^2b^2}\right)(bx+a)^3}{6b^5} \\ & \quad - \frac{(Bb^3d^3 - 6Bab^2d^2e + 3Ab^3d^2e + 9Ba^2bde^2 - 6Aab^2de^2 - 4Ba^3e^3 + 3Aa^2be^3) \log\left(\frac{|bx+a|}{(bx+a)^2|b|}\right)}{b^5} \\ & \quad + \frac{\frac{Bab^6d^3}{bx+a} - \frac{Ab^7d^3}{bx+a} - \frac{3Ba^2b^5d^2e}{bx+a} + \frac{3Aab^6d^2e}{bx+a} + \frac{3Ba^3b^4de^2}{bx+a} - \frac{3Aa^2b^5de^2}{bx+a} - \frac{Ba^4b^3e^3}{bx+a} + \frac{Aa^3b^4e^3}{bx+a}}{b^8} \end{aligned}$$

input

```
integrate((B*x+A)*(e*x+d)^3/(b*x+a)^2,x, algorithm="giac")
```

output

$$\begin{aligned} & 1/6*(2*B*e^3 + 3*(3*B*b^2*d*e^2 - 4*B*a*b*e^3 + A*b^2*e^3)/((b*x + a)*b) + \\ & 18*(B*b^4*d^2*e - 3*B*a*b^3*d*e^2 + A*b^4*d*e^2 + 2*B*a^2*b^2*e^3 - A*a*b^3*e^3)/((b*x + a)^2*b^2))* (b*x + a)^3/b^5 - (B*b^3*d^3 - 6*B*a*b^2*d^2*e \\ & + 3*A*b^3*d^2*e + 9*B*a^2*b*d*e^2 - 6*A*a*b^2*d*e^2 - 4*B*a^3*e^3 + 3*A*a^2*b*e^3)*log(abs(b*x + a)/((b*x + a)^2*abs(b)))/b^5 + (B*a*b^6*d^3/(b*x + \\ & a) - A*b^7*d^3/(b*x + a) - 3*B*a^2*b^5*d^2*e/(b*x + a) + 3*A*a*b^6*d^2*e/(\\ & b*x + a) + 3*B*a^3*b^4*d*e^2/(b*x + a) - 3*A*a^2*b^5*d*e^2/(b*x + a) - B*a^4*b^3*e^3/(b*x + a) + A*a^3*b^4*e^3/(b*x + a))/b^8 \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 293, normalized size of antiderivative = 2.02

$$\int \frac{(A + Bx)(d + ex)^3}{(a + bx)^2} dx = x^2 \left(\frac{Ae^3 + 3Bde^2}{2b^2} - \frac{Ba^2e^3}{b^3} \right) - x \left(\frac{2a \left(\frac{Ae^3 + 3Bde^2}{b^2} - \frac{2Ba^2e^3}{b^3} \right) - \frac{3de(Ae + Bd)}{b^2} + \frac{Ba^2e^3}{b^4}}{b} \right) + \frac{\ln(a + bx) (-4Ba^3e^3 + 9Ba^2bde^2 + 3Aa^2be^3 - 6Bab^2d^2e - 6Aab^2de^2 + Bb^3d^3 + 3Ab^3d^2e + Ba^4e^3 - 3Ba^3bde^2 - Aa^3be^3 + 3Ba^2b^2d^2e + 3Aa^2b^2de^2 - Bab^3d^3 - 3Aab^3d^2e + Ab^4d^3)}{b^5} + \frac{Be^3x^3}{3b^2}$$

input `int(((A + B*x)*(d + e*x)^3)/(a + b*x)^2,x)`output `x^2*((A*e^3 + 3*B*d*e^2)/(2*b^2) - (B*a*e^3)/b^3) - x*((2*a*((A*e^3 + 3*B*d*e^2)/b^2 - (2*B*a*e^3)/b^3))/b - (3*d*e*(A*e + B*d))/b^2 + (B*a^2*e^3)/b^4 + (log(a + b*x)*(B*b^3*d^3 - 4*B*a^3*e^3 + 3*A*a^2*b*e^3 + 3*A*b^3*d^2*e - 6*A*a*b^2*d*e^2 - 6*B*a*b^2*d^2*e + 9*B*a^2*b*d*e^2))/b^5 - (A*b^4*d^3 + B*a^4*e^3 - A*a^3*b*e^3 - B*a*b^3*d^3 + 3*A*a^2*b^2*d*e^2 + 3*B*a^2*b^2*d^2*e - 3*A*a*b^3*d^2*e - 3*B*a^3*b*d*e^2)/(b*(a*b^4 + b^5*x)) + (B*e^3*x^3)/(3*b^2)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.91

$$\int \frac{(A + Bx)(d + ex)^3}{(a + bx)^2} dx = \frac{-6 \log(bx + a) a^3 e^3 + 18 \log(bx + a) a^2 b d e^2 - 18 \log(bx + a) a b^2 d^2 e + 6 \log(bx + a) b^3 d^3 + 6 a^2 b e^3 x - 1}{6b^4}$$

input `int((B*x+A)*(e*x+d)^3/(b*x+a)^2,x)`

output

```
( - 6*log(a + b*x)*a**3*e**3 + 18*log(a + b*x)*a**2*b*d*e**2 - 18*log(a +
b*x)*a*b**2*d**2*e + 6*log(a + b*x)*b**3*d**3 + 6*a**2*b*e**3*x - 18*a*b**
2*d*e**2*x - 3*a*b**2*e**3*x**2 + 18*b**3*d**2*e*x + 9*b**3*d*e**2*x**2 +
2*b**3*e**3*x**3)/(6*b**4)
```

3.111 $\int \frac{(A+Bx)(d+ex)^2}{(a+bx)^2} dx$

Optimal result	1117
Mathematica [A] (verified)	1117
Rubi [A] (verified)	1118
Maple [A] (verified)	1119
Fricas [B] (verification not implemented)	1120
Sympy [A] (verification not implemented)	1120
Maxima [A] (verification not implemented)	1121
Giac [B] (verification not implemented)	1122
Mupad [B] (verification not implemented)	1122
Reduce [B] (verification not implemented)	1123

Optimal result

Integrand size = 20, antiderivative size = 99

$$\int \frac{(A+Bx)(d+ex)^2}{(a+bx)^2} dx = \frac{e(2bBd + Abe - 2aBe)x}{b^3} + \frac{Be^2x^2}{2b^2} - \frac{(Ab - aB)(bd - ae)^2}{b^4(a+bx)} + \frac{(bd - ae)(bBd + 2Abe - 3aBe) \log(a+bx)}{b^4}$$

output

```
e*(A*b*e-2*B*a*e+2*B*b*d)*x/b^3+1/2*B*e^2*x^2/b^2-(A*b-B*a)*(-a*e+b*d)^2/b^4/(b*x+a)+(-a*e+b*d)*(2*A*b*e-3*B*a*e+B*b*d)*ln(b*x+a)/b^4
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.55

$$\begin{aligned} & \int \frac{(A+Bx)(d+ex)^2}{(a+bx)^2} dx \\ &= \frac{e(2bBd + Abe - 2aBe)x}{b^3} + \frac{Be^2x^2}{2b^2} \\ &+ \frac{-Ab^3d^2 + ab^2Bd^2 + 2aAb^2de - 2a^2bBde - a^2Abe^2 + a^3Be^2}{b^4(a+bx)} \\ &+ \frac{(b^2Bd^2 + 2Ab^2de - 4abBde - 2aAbe^2 + 3a^2Be^2) \log(a+bx)}{b^4} \end{aligned}$$

input `Integrate[((A + B*x)*(d + e*x)^2)/(a + b*x)^2,x]`

output
$$\frac{(e*(2*b*B*d + A*b*e - 2*a*B*e)*x)/b^3 + (B*e^2*x^2)/(2*b^2) + (- (A*b^3*d^2) + a*b^2*B*d^2 + 2*a*A*b^2*d*e - 2*a^2*b*B*d*e - a^2*A*b*e^2 + a^3*B*e^2)/(b^4*(a + b*x)) + ((b^2*B*d^2 + 2*A*b^2*d*e - 4*a*b*B*d*e - 2*a*A*b*e^2 + 3*a^2*B*e^2)*\text{Log}[a + b*x])/b^4$$

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)^2}{(a + bx)^2} dx$$

↓ 86

$$\int \left(\frac{(bd - ae)(-3aBe + 2Abe + bBd)}{b^3(a + bx)} + \frac{(Ab - aB)(bd - ae)^2}{b^3(a + bx)^2} + \frac{e(-2aBe + Abe + 2bBd)}{b^3} + \frac{Be^2x}{b^2} \right) dx$$

↓ 2009

$$-\frac{(Ab - aB)(bd - ae)^2}{b^4(a + bx)} + \frac{(bd - ae) \log(a + bx)(-3aBe + 2Abe + bBd)}{b^4} + \frac{ex(-2aBe + Abe + 2bBd)}{b^3} + \frac{Be^2x^2}{2b^2}$$

input `Int[((A + B*x)*(d + e*x)^2)/(a + b*x)^2,x]`

output
$$\frac{(e*(2*b*B*d + A*b*e - 2*a*B*e)*x)/b^3 + (B*e^2*x^2)/(2*b^2) - ((A*b - a*B)*(b*d - a*e)^2)/(b^4*(a + b*x)) + ((b*d - a*e)*(b*B*d + 2*A*b*e - 3*a*B*e)*\text{Log}[a + b*x])/b^4$$

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0]) || GeQ[n + p + 1, 0]) || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f]))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.53

method	result
default	$\frac{e(\frac{1}{2}Bbe^2x^2 + A b e x - 2B a e x + 2B b d x)}{b^3} - \frac{A a^2 b e^2 - 2A a b^2 d e + A b^3 d^2 - B a^3 e^2 + 2B a^2 b d e - B a b^2 d^2}{b^4(bx+a)} + \frac{(-2A a b e^2 + 2A b^2 d e - 2A a^2 b e^2 - 2A a b^2 d e + A b^3 d^2 - 3B a^3 e^2 + 4B a^2 b d e - B a b^2 d^2)x}{b^3 a} + \frac{B e^2 x^3}{2b} + \frac{e(2A b e - 3B a e + 4B b d)x^2}{2b^2} - \frac{(2A a b e^2 - 2A b^2 d e - 3B a^2 e^2)}{b x + a}$
norman	
risch	$\frac{B e^2 x^2}{2b^2} + \frac{e^2 A x}{b^2} - \frac{2e^2 B a x}{b^3} + \frac{2e B d x}{b^2} - \frac{A a^2 e^2}{b^3(bx+a)} + \frac{2A a d e}{b^2(bx+a)} - \frac{A d^2}{b(bx+a)} + \frac{B a^3 e^2}{b^4(bx+a)} - \frac{2B a^2 d e}{b^3(bx+a)} + \frac{B a d^2}{b^2(bx+a)}$
parallelrisch	$- \frac{B e^2 x^3 b^3 + 4A \ln(bx+a) x a b^2 e^2 - 4A \ln(bx+a) x b^3 d e - 2A x^2 b^3 e^2 - 6B \ln(bx+a) x a^2 b e^2 + 8B \ln(bx+a) x a b^2 d e - 2B \ln(bx+a) x a^2 d^2}{b^3 a}$

```
input int((B*x+A)*(e*x+d)^2/(b*x+a)^2,x,method=_RETURNVERBOSE)
```

```
output e/b^3*(1/2*B*b*e*x^2+A*b*e*x-2*B*a*e*x+2*B*b*d*x)-(A*a^2*b*e^2-2*A*a*b^2*d*e+A*b^3*d^2-B*a^3*e^2+2*B*a^2*b*d*e-B*a*b^2*d^2)/b^4/(b*x+a)+(-2*A*a*b*e^2+2*A*b^2*d*e+3*B*a^2*e^2-4*B*a*b*d*e+B*b^2*d^2)/b^4*ln(b*x+a)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 249 vs. $2(96) = 192$.

Time = 0.08 (sec) , antiderivative size = 249, normalized size of antiderivative = 2.52

$$\int \frac{(A + Bx)(d + ex)^2}{(a + bx)^2} dx$$

$$= \frac{Bb^3e^2x^3 + 2(Bab^2 - Ab^3)d^2 - 4(Ba^2b - Aab^2)de + 2(Ba^3 - Aa^2b)e^2 + (4Bb^3de - (3Bab^2 - 2Ab^3)e^2)}{(a + bx)^2}$$

input `integrate((B*x+A)*(e*x+d)^2/(b*x+a)^2,x, algorithm="fricas")`

output `1/2*(B*b^3*e^2*x^3 + 2*(B*a*b^2 - A*b^3)*d^2 - 4*(B*a^2*b - A*a*b^2)*d*e + 2*(B*a^3 - A*a^2*b)*e^2 + (4*B*b^3*d*e - (3*B*a*b^2 - 2*A*b^3)*e^2)*x^2 + 2*(2*B*a*b^2*d*e - (2*B*a^2*b - A*a*b^2)*e^2)*x + 2*(B*a*b^2*d^2 - 2*(2*B*a^2*b - A*a*b^2)*d*e + (3*B*a^3 - 2*A*a^2*b)*e^2 + (B*b^3*d^2 - 2*(2*B*a*b^2 - A*b^3)*d*e + (3*B*a^2*b - 2*A*a*b^2)*e^2)*x)*log(b*x + a)/(b^5*x + a*b^4)`

Sympy [A] (verification not implemented)

Time = 0.48 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.53

$$\int \frac{(A + Bx)(d + ex)^2}{(a + bx)^2} dx$$

$$= \frac{Be^2x^2}{2b^2} + x \left(\frac{Ae^2}{b^2} - \frac{2Bae^2}{b^3} + \frac{2Bde}{b^2} \right)$$

$$+ \frac{-Aa^2be^2 + 2Aab^2de - Ab^3d^2 + Ba^3e^2 - 2Ba^2bde + Bab^2d^2}{ab^4 + b^5x}$$

$$+ \frac{(ae - bd)(-2Abe + 3Bae - Bbd) \log(a + bx)}{b^4}$$

input `integrate((B*x+A)*(e*x+d)**2/(b*x+a)**2,x)`

output

```
B***2*x**2/(2*b**2) + x*(A***2/b**2 - 2*B*a***2/b**3 + 2*B*d*e/b**2) +
(-A***2*b***2 + 2*A*a*b**2*d*e - A*b**3*d**2 + B***3*e**2 - 2*B*a**2*b*
d*e + B*a*b**2*d**2)/(a*b**4 + b**5*x) + (a*e - b*d)*(-2*A*b*e + 3*B*a*e -
B*b*d)*log(a + b*x)/b**4
```

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.60

$$\int \frac{(A + Bx)(d + ex)^2}{(a + bx)^2} dx$$

$$= \frac{(Bab^2 - Ab^3)d^2 - 2(Ba^2b - Aab^2)de + (Ba^3 - Aa^2b)e^2}{b^5x + ab^4}$$

$$+ \frac{Bbe^2x^2 + 2(2Bbde - (2Ba - Ab)e^2)x}{2b^3}$$

$$+ \frac{(Bb^2d^2 - 2(2Bab - Ab^2)de + (3Ba^2 - 2Aab)e^2) \log(bx + a)}{b^4}$$

input

```
integrate((B*x+A)*(e*x+d)^2/(b*x+a)^2,x, algorithm="maxima")
```

output

```
((B*a*b^2 - A*b^3)*d^2 - 2*(B*a^2*b - A*a*b^2)*d*e + (B*a^3 - A*a^2*b)*e^2
)/(b^5*x + a*b^4) + 1/2*(B*b*e^2*x^2 + 2*(2*B*b*d*e - (2*B*a - A*b)*e^2)*x
)/b^3 + (B*b^2*d^2 - 2*(2*B*a*b - A*b^2)*d*e + (3*B*a^2 - 2*A*a*b)*e^2)*lo
g(b*x + a)/b^4
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 229 vs. 2(96) = 192.

Time = 0.13 (sec) , antiderivative size = 229, normalized size of antiderivative = 2.31

$$\int \frac{(A + Bx)(d + ex)^2}{(a + bx)^2} dx$$

$$= \frac{\left(B e^2 + \frac{2(2 B b^2 d e - 3 B a b e^2 + A b^2 e^2)}{(b x + a) b} \right) (b x + a)^2}{2 b^4}$$

$$- \frac{(B b^2 d^2 - 4 B a b d e + 2 A b^2 d e + 3 B a^2 e^2 - 2 A a b e^2) \log\left(\frac{|b x + a|}{(b x + a)^2 |b|}\right)}{b^4}$$

$$+ \frac{\frac{B a b^4 d^2}{b x + a} - \frac{A b^5 d^2}{b x + a} - \frac{2 B a^2 b^3 d e}{b x + a} + \frac{2 A a b^4 d e}{b x + a} + \frac{B a^3 b^2 e^2}{b x + a} - \frac{A a^2 b^3 e^2}{b x + a}}{b^6}$$

input `integrate((B*x+A)*(e*x+d)^2/(b*x+a)^2,x, algorithm="giac")`

output `1/2*(B*e^2 + 2*(2*B*b^2*d*e - 3*B*a*b*e^2 + A*b^2*e^2)/((b*x + a)*b))*(b*x + a)^2/b^4 - (B*b^2*d^2 - 4*B*a*b*d*e + 2*A*b^2*d*e + 3*B*a^2*e^2 - 2*A*a*b*e^2)*log(abs(b*x + a)/((b*x + a)^2*abs(b)))/b^4 + (B*a*b^4*d^2/(b*x + a) - A*b^5*d^2/(b*x + a) - 2*B*a^2*b^3*d*e/(b*x + a) + 2*A*a*b^4*d*e/(b*x + a) + B*a^3*b^2*e^2/(b*x + a) - A*a^2*b^3*e^2/(b*x + a))/b^6`

Mupad [B] (verification not implemented)

Time = 1.01 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.67

$$\int \frac{(A + Bx)(d + ex)^2}{(a + bx)^2} dx$$

$$= x \left(\frac{A e^2 + 2 B d e}{b^2} - \frac{2 B a e^2}{b^3} \right)$$

$$+ \frac{\ln(a + bx) (3 B a^2 e^2 - 4 B a b d e - 2 A a b e^2 + B b^2 d^2 + 2 A b^2 d e)}{b^4}$$

$$- \frac{-B a^3 e^2 + 2 B a^2 b d e + A a^2 b e^2 - B a b^2 d^2 - 2 A a b^2 d e + A b^3 d^2}{b (x b^4 + a b^3)} + \frac{B e^2 x^2}{2 b^2}$$

input `int(((A + B*x)*(d + e*x)^2)/(a + b*x)^2,x)`

output

```
x*((A*e^2 + 2*B*d*e)/b^2 - (2*B*a*e^2)/b^3) + (log(a + b*x)*(3*B*a^2*e^2 +
B*b^2*d^2 - 2*A*a*b*e^2 + 2*A*b^2*d*e - 4*B*a*b*d*e))/b^4 - (A*b^3*d^2 -
B*a^3*e^2 + A*a^2*b*e^2 - B*a*b^2*d^2 - 2*A*a*b^2*d*e + 2*B*a^2*b*d*e)/(b*
(a*b^3 + b^4*x)) + (B*e^2*x^2)/(2*b^2)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.73

$$\int \frac{(A + Bx)(d + ex)^2}{(a + bx)^2} dx$$

$$= \frac{2 \log(bx + a) a^2 e^2 - 4 \log(bx + a) abde + 2 \log(bx + a) b^2 d^2 - 2ab e^2 x + 4b^2 dex + b^2 e^2 x^2}{2b^3}$$

input

```
int((B*x+A)*(e*x+d)^2/(b*x+a)^2,x)
```

output

```
(2*log(a + b*x)*a**2*e**2 - 4*log(a + b*x)*a*b*d*e + 2*log(a + b*x)*b**2*d
**2 - 2*a*b*e**2*x + 4*b**2*d*e*x + b**2*e**2*x**2)/(2*b**3)
```


3.112 $\int \frac{(A+Bx)(d+ex)}{(a+bx)^2} dx$

Optimal result	1124
Mathematica [A] (verified)	1124
Rubi [A] (verified)	1125
Maple [A] (verified)	1126
Fricas [A] (verification not implemented)	1126
Sympy [A] (verification not implemented)	1127
Maxima [A] (verification not implemented)	1127
Giac [A] (verification not implemented)	1128
Mupad [B] (verification not implemented)	1128
Reduce [B] (verification not implemented)	1129

Optimal result

Integrand size = 18, antiderivative size = 60

$$\int \frac{(A + Bx)(d + ex)}{(a + bx)^2} dx = \frac{Bex}{b^2} - \frac{(Ab - aB)(bd - ae)}{b^3(a + bx)} + \frac{(bBd + Abe - 2aBe) \log(a + bx)}{b^3}$$

output

$B*e*x/b^2 - (A*b - B*a)*(-a*e + b*d)/b^3 / (b*x + a) + (A*b*e - 2*B*a*e + B*b*d)*\ln(b*x + a) / b^3$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.93

$$\int \frac{(A + Bx)(d + ex)}{(a + bx)^2} dx = \frac{bBex - \frac{(Ab - aB)(bd - ae)}{a + bx} + (bBd + Abe - 2aBe) \log(a + bx)}{b^3}$$

input

`Integrate[((A + B*x)*(d + e*x))/(a + b*x)^2,x]`

output

$(b*B*e*x - ((A*b - a*B)*(b*d - a*e))/(a + b*x) + (b*B*d + A*b*e - 2*a*B*e)*\text{Log}[a + b*x])/b^3$

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)}{(a + bx)^2} dx$$

↓ 86

$$\int \left(\frac{-2aBe + Abe + bBd}{b^2(a + bx)} + \frac{(Ab - aB)(bd - ae)}{b^2(a + bx)^2} + \frac{Be}{b^2} \right) dx$$

↓ 2009

$$-\frac{(Ab - aB)(bd - ae)}{b^3(a + bx)} + \frac{\log(a + bx)(-2aBe + Abe + bBd)}{b^3} + \frac{Bex}{b^2}$$

input `Int[((A + B*x)*(d + e*x))/(a + b*x)^2,x]`

output `(B*e*x)/b^2 - ((A*b - a*B)*(b*d - a*e))/(b^3*(a + b*x)) + ((b*B*d + A*b*e - 2*a*B*e)*Log[a + b*x])/b^3`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.17

method	result
default	$\frac{Bex}{b^2} - \frac{-Aabe+Ab^2d+Ba^2e-Babd}{b^3(bx+a)} + \frac{(Abe-2Bae+Bbd)\ln(bx+a)}{b^3}$
norman	$\frac{Aabe-Ab^2d-2Ba^2e+Babd+Be x^2}{bx+a} + \frac{(Abe-2Bae+Bbd)\ln(bx+a)}{b^3}$
risch	$\frac{Bex}{b^2} + \frac{Aae}{b^2(bx+a)} - \frac{Ad}{b(bx+a)} - \frac{Ba^2e}{b^3(bx+a)} + \frac{Bad}{b^2(bx+a)} + \frac{\ln(bx+a)Ae}{b^2} - \frac{2\ln(bx+a)Bae}{b^3} + \frac{\ln(bx+a)Bd}{b^2}$
parallelrisch	$\frac{A\ln(bx+a)x b^2e-2B\ln(bx+a)xabe+B\ln(bx+a)x b^2d+Be x^2b^2+A\ln(bx+a)abe-2B\ln(bx+a)a^2e+B\ln(bx+a)abd+Abe-}{b^3(bx+a)}$

input `int((B*x+A)*(e*x+d)/(b*x+a)^2,x,method=_RETURNVERBOSE)`output `B*e*x/b^2-1/b^3*(-A*a*b*e+A*b^2*d+B*a^2*e-B*a*b*d)/(b*x+a)+(A*b*e-2*B*a*e+B*b*d)*ln(b*x+a)/b^3`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.82

$$\int \frac{(A+Bx)(d+ex)}{(a+bx)^2} dx$$

$$= \frac{Bb^2ex^2 + Babex + (Bab - Ab^2)d - (Ba^2 - Aab)e + (Babd - (2Ba^2 - Aab)e + (Bb^2d - (2Bab - Ab^2))x)}{b^4x + ab^3}$$

input `integrate((B*x+A)*(e*x+d)/(b*x+a)^2,x, algorithm="fricas")`output `(B*b^2*e*x^2 + B*a*b*e*x + (B*a*b - A*b^2)*d - (B*a^2 - A*a*b)*e + (B*a*b*d - (2*B*a^2 - A*a*b)*e + (B*b^2*d - (2*B*a*b - A*b^2)*e)*x)*log(b*x + a) / (b^4*x + a*b^3)`

Sympy [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.18

$$\int \frac{(A + Bx)(d + ex)}{(a + bx)^2} dx = \frac{Bex}{b^2} + \frac{Aabe - Ab^2d - Ba^2e + Babd}{ab^3 + b^4x} - \frac{(-Abe + 2Bae - Bbd) \log(a + bx)}{b^3}$$

input `integrate((B*x+A)*(e*x+d)/(b*x+a)**2,x)`

output `B*e*x/b**2 + (A*a*b*e - A*b**2*d - B*a**2*e + B*a*b*d)/(a*b**3 + b**4*x) - (-A*b*e + 2*B*a*e - B*b*d)*log(a + b*x)/b**3`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.28

$$\int \frac{(A + Bx)(d + ex)}{(a + bx)^2} dx = \frac{Bex}{b^2} + \frac{(Bab - Ab^2)d - (Ba^2 - Aab)e}{b^4x + ab^3} + \frac{(Bbd - (2Ba - Ab)e) \log(bx + a)}{b^3}$$

input `integrate((B*x+A)*(e*x+d)/(b*x+a)^2,x, algorithm="maxima")`

output `B*e*x/b^2 + ((B*a*b - A*b^2)*d - (B*a^2 - A*a*b)*e)/(b^4*x + a*b^3) + (B*b*d - (2*B*a - A*b)*e)*log(b*x + a)/b^3`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.87

$$\int \frac{(A + Bx)(d + ex)}{(a + bx)^2} dx = \frac{(bx + a)Be}{b^3} - \frac{(Bbd - 2Bae + Abe) \log\left(\frac{|bx+a|}{(bx+a)^2|b|}\right)}{b^3} + \frac{\frac{Bab^2d}{bx+a} - \frac{Ab^3d}{bx+a} - \frac{Ba^2be}{bx+a} + \frac{Aab^2e}{bx+a}}{b^4}$$

input `integrate((B*x+A)*(e*x+d)/(b*x+a)^2,x, algorithm="giac")`output `(b*x + a)*B*e/b^3 - (B*b*d - 2*B*a*e + A*b*e)*log(abs(b*x + a)/((b*x + a)^2*abs(b)))/b^3 + (B*a*b^2*d/(b*x + a) - A*b^3*d/(b*x + a) - B*a^2*b*e/(b*x + a) + A*a*b^2*e/(b*x + a))/b^4`**Mupad [B] (verification not implemented)**

Time = 1.01 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.25

$$\int \frac{(A + Bx)(d + ex)}{(a + bx)^2} dx = \frac{\ln(a + bx) (Abe - 2Bae + Bbd)}{b^3} - \frac{Ab^2d + Ba^2e - Aabe - Babd}{b(xb^3 + ab^2)} + \frac{Bex}{b^2}$$

input `int(((A + B*x)*(d + e*x))/(a + b*x)^2,x)`output `(log(a + b*x)*(A*b*e - 2*B*a*e + B*b*d))/b^3 - (A*b^2*d + B*a^2*e - A*a*b*e - B*a*b*d)/(b*(a*b^2 + b^3*x)) + (B*e*x)/b^2`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.47

$$\int \frac{(A + Bx)(d + ex)}{(a + bx)^2} dx = \frac{-\log(bx + a)ae + \log(bx + a)bd + bex}{b^2}$$

input

```
int((B*x+A)*(e*x+d)/(b*x+a)^2,x)
```

output

```
( - log(a + b*x)*a*e + log(a + b*x)*b*d + b*e*x)/b**2
```

3.113 $\int \frac{A+Bx}{(a+bx)^2} dx$

Optimal result	1130
Mathematica [A] (verified)	1130
Rubi [A] (verified)	1131
Maple [A] (verified)	1132
Fricas [A] (verification not implemented)	1132
Sympy [A] (verification not implemented)	1132
Maxima [A] (verification not implemented)	1133
Giac [A] (verification not implemented)	1133
Mupad [B] (verification not implemented)	1134
Reduce [B] (verification not implemented)	1134

Optimal result

Integrand size = 13, antiderivative size = 32

$$\int \frac{A+Bx}{(a+bx)^2} dx = -\frac{Ab-aB}{b^2(a+bx)} + \frac{B \log(a+bx)}{b^2}$$

output

```
-(A*b-B*a)/b^2/(b*x+a)+B*ln(b*x+a)/b^2
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.97

$$\int \frac{A+Bx}{(a+bx)^2} dx = \frac{-Ab+aB}{b^2(a+bx)} + \frac{B \log(a+bx)}{b^2}$$

input

```
Integrate[(A + B*x)/(a + b*x)^2,x]
```

output

```
(-(A*b) + a*B)/(b^2*(a + b*x)) + (B*Log[a + b*x])/b^2
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^2} dx$$

$$\downarrow 49$$

$$\int \left(\frac{Ab - aB}{b(a + bx)^2} + \frac{B}{b(a + bx)} \right) dx$$

$$\downarrow 2009$$

$$\frac{B \log(a + bx)}{b^2} - \frac{Ab - aB}{b^2(a + bx)}$$

input `Int[(A + B*x)/(a + b*x)^2,x]`

output `-((A*b - a*B)/(b^2*(a + b*x))) + (B*Log[a + b*x])/b^2`

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.03

method	result	size
default	$-\frac{Ab-Ba}{b^2(bx+a)} + \frac{B \ln(bx+a)}{b^2}$	33
norman	$-\frac{Ab-Ba}{b^2(bx+a)} + \frac{B \ln(bx+a)}{b^2}$	33
risch	$-\frac{A}{b(bx+a)} + \frac{Ba}{b^2(bx+a)} + \frac{B \ln(bx+a)}{b^2}$	39
parallelrisc	$-\frac{-B \ln(bx+a)xb - B \ln(bx+a)a + Ab - Ba}{b^2(bx+a)}$	42

input `int((B*x+A)/(b*x+a)^2,x,method=_RETURNVERBOSE)`output `-(A*b-B*a)/b^2/(b*x+a)+B*ln(b*x+a)/b^2`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.16

$$\int \frac{A + Bx}{(a + bx)^2} dx = \frac{Ba - Ab + (Bbx + Ba) \log(bx + a)}{b^3x + ab^2}$$

input `integrate((B*x+A)/(b*x+a)^2,x, algorithm="fricas")`output `(B*a - A*b + (B*b*x + B*a)*log(b*x + a))/(b^3*x + a*b^2)`**Sympy [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.84

$$\int \frac{A + Bx}{(a + bx)^2} dx = \frac{B \log(a + bx)}{b^2} + \frac{-Ab + Ba}{ab^2 + b^3x}$$

input `integrate((B*x+A)/(b*x+a)**2,x)`

output $B \log(a + bx) / b^2 + (-A \cdot b + B \cdot a) / (a \cdot b^2 + b^3 \cdot x)$

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.06

$$\int \frac{A + Bx}{(a + bx)^2} dx = \frac{Ba - Ab}{b^3x + ab^2} + \frac{B \log(bx + a)}{b^2}$$

input `integrate((B*x+A)/(b*x+a)^2,x, algorithm="maxima")`

output $(B \cdot a - A \cdot b) / (b^3 \cdot x + a \cdot b^2) + B \cdot \log(b \cdot x + a) / b^2$

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.78

$$\int \frac{A + Bx}{(a + bx)^2} dx = -\frac{B \left(\frac{\log\left(\frac{|bx+a|}{(bx+a)^2|b|}\right)}{b} - \frac{a}{(bx+a)b} \right)}{b} - \frac{A}{(bx+a)b}$$

input `integrate((B*x+A)/(b*x+a)^2,x, algorithm="giac")`

output $-B \cdot (\log(\text{abs}(b \cdot x + a) / ((b \cdot x + a)^2 \cdot \text{abs}(b)))) / b - a / ((b \cdot x + a) \cdot b) - A / ((b \cdot x + a) \cdot b)$

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

$$\int \frac{A + Bx}{(a + bx)^2} dx = \frac{B \ln(a + bx)}{b^2} - \frac{Ab - Ba}{b^2 (a + bx)}$$

input `int((A + B*x)/(a + b*x)^2,x)`

output `(B*log(a + b*x))/b^2 - (A*b - B*a)/(b^2*(a + b*x))`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.31

$$\int \frac{A + Bx}{(a + bx)^2} dx = \frac{\log(bx + a)}{b}$$

input `int((B*x+A)/(b*x+a)^2,x)`

output `log(a + b*x)/b`

3.114 $\int \frac{A+Bx}{(a+bx)^2(d+ex)} dx$

Optimal result	1135
Mathematica [A] (verified)	1135
Rubi [A] (verified)	1136
Maple [A] (verified)	1137
Fricas [A] (verification not implemented)	1137
Sympy [B] (verification not implemented)	1138
Maxima [A] (verification not implemented)	1138
Giac [A] (verification not implemented)	1139
Mupad [B] (verification not implemented)	1139
Reduce [B] (verification not implemented)	1140

Optimal result

Integrand size = 20, antiderivative size = 82

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)} dx = -\frac{Ab - aB}{b(bd - ae)(a + bx)} + \frac{(Bd - Ae) \log(a + bx)}{(bd - ae)^2} - \frac{(Bd - Ae) \log(d + ex)}{(bd - ae)^2}$$

output

```
-(A*b-B*a)/b/(-a*e+b*d)/(b*x+a)+(-A*e+B*d)*ln(b*x+a)/(-a*e+b*d)^2-(-A*e+B*d)*ln(e*x+d)/(-a*e+b*d)^2
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.84

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)} dx = \frac{\frac{(-Ab+aB)(bd-ae)}{b(a+bx)} + (Bd - Ae) \log(a + bx) + (-Bd + Ae) \log(d + ex)}{(bd - ae)^2}$$

input

```
Integrate[(A + B*x)/((a + b*x)^2*(d + e*x)),x]
```

output

$$\frac{((-A*b) + a*B)*(b*d - a*e)}{b*(a + b*x)} + (B*d - A*e)*\text{Log}[a + b*x] + (-B*d + A*e)*\text{Log}[d + e*x] / (b*d - a*e)^2$$

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)} dx$$

↓ 86

$$\int \left(\frac{Ab - aB}{(a + bx)^2(bd - ae)} + \frac{b(Bd - Ae)}{(a + bx)(bd - ae)^2} + \frac{e(Ae - Bd)}{(d + ex)(bd - ae)^2} \right) dx$$

↓ 2009

$$-\frac{Ab - aB}{b(a + bx)(bd - ae)} + \frac{\log(a + bx)(Bd - Ae)}{(bd - ae)^2} - \frac{(Bd - Ae) \log(d + ex)}{(bd - ae)^2}$$

input

```
Int[(A + B*x)/((a + b*x)^2*(d + e*x)),x]
```

output

$$\frac{-((A*b - a*B)/(b*(b*d - a*e)*(a + b*x))) + ((B*d - A*e)*\text{Log}[a + b*x])}{(b*d - a*e)^2} - \frac{(B*d - A*e)*\text{Log}[d + e*x]}{(b*d - a*e)^2}$$

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0]) || GeQ[n + p + 1, 0]) || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f]))
```

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.01

method	result
default	$-\frac{-Ab+Ba}{(ae-db)b(bx+a)} - \frac{(Ae-Bd)\ln(bx+a)}{(ae-db)^2} + \frac{(Ae-Bd)\ln(ex+d)}{(ae-db)^2}$
norman	$-\frac{(Ab-Ba)x}{a(ae-db)(bx+a)} + \frac{(Ae-Bd)\ln(ex+d)}{a^2e^2-2abde+b^2d^2} - \frac{(Ae-Bd)\ln(bx+a)}{a^2e^2-2abde+b^2d^2}$
parallelrisc	$-\frac{A\ln(bx+a)x b^2e - A\ln(ex+d)x b^2e - B\ln(bx+a)x b^2d + B\ln(ex+d)x b^2d + A\ln(bx+a)abe - A\ln(ex+d)abe - B\ln(bx+a)abd}{(a^2e^2-2abde+b^2d^2)(bx+a)b}$
risc	$\frac{A}{(ae-db)(bx+a)} - \frac{Ba}{(ae-db)b(bx+a)} - \frac{\ln(bx+a)Ae}{a^2e^2-2abde+b^2d^2} + \frac{\ln(bx+a)Bd}{a^2e^2-2abde+b^2d^2} + \frac{\ln(-ex-d)Ae}{a^2e^2-2abde+b^2d^2} - \frac{\ln(-ex-d)Bd}{a^2e^2-2abde+b^2d^2}$

input `int((B*x+A)/(b*x+a)^2/(e*x+d),x,method=_RETURNVERBOSE)`

output
$$\frac{-(-A*b+B*a)/(a*e-b*d)/b/(b*x+a) - (A*e-B*d)/(a*e-b*d)^2*\ln(b*x+a) + (A*e-B*d)/(a*e-b*d)^2*\ln(e*x+d)}$$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.91

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)} dx$$

$$= \frac{(Bab - Ab^2)d - (Ba^2 - Aab)e + (Babd - Aabe + (Bb^2d - Ab^2e)x) \log(bx + a) - (Babd - Aabe + (Bb^2d - Ab^2e)x) \log(ex + d)}{ab^3d^2 - 2a^2b^2de + a^3be^2 + (b^4d^2 - 2ab^3de + a^2b^2e^2)x}$$

input `integrate((B*x+A)/(b*x+a)^2/(e*x+d),x, algorithm="fricas")`

output
$$\frac{((B*a*b - A*b^2)*d - (B*a^2 - A*a*b)*e + (B*a*b*d - A*a*b*e + (B*b^2*d - A*b^2*e)*x)*\log(b*x + a) - (B*a*b*d - A*a*b*e + (B*b^2*d - A*b^2*e)*x)*\log(e*x + d)}{(a*b^3*d^2 - 2*a^2*b^2*d*e + a^3*b*e^2 + (b^4*d^2 - 2*a*b^3*d*e + a^2*b^2*e^2)*x)}$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 355 vs. $2(63) = 126$.

Time = 0.70 (sec) , antiderivative size = 355, normalized size of antiderivative = 4.33

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)} dx = \frac{Ab - Ba}{a^2be - ab^2d + x(ab^2e - b^3d)}$$

$$\frac{(-Ae + Bd) \log \left(x + \frac{-Aae^2 - Abde + Bade + Bbd^2 - \frac{a^3e^3(-Ae+Bd)}{(ae-bd)^2} + \frac{3a^2bde^2(-Ae+Bd)}{(ae-bd)^2} - \frac{3ab^2d^2e(-Ae+Bd)}{(ae-bd)^2} + \frac{b^3d^3(-Ae+Bd)}{(ae-bd)^2}}{-2Abe^2 + 2Bbde} \right)}{(ae - bd)^2}$$

$$+ \frac{(-Ae + Bd) \log \left(x + \frac{-Aae^2 - Abde + Bade + Bbd^2 + \frac{a^3e^3(-Ae+Bd)}{(ae-bd)^2} - \frac{3a^2bde^2(-Ae+Bd)}{(ae-bd)^2} + \frac{3ab^2d^2e(-Ae+Bd)}{(ae-bd)^2} - \frac{b^3d^3(-Ae+Bd)}{(ae-bd)^2}}{-2Abe^2 + 2Bbde} \right)}{(ae - bd)^2}$$

input `integrate((B*x+A)/(b*x+a)**2/(e*x+d), x)`

output `(A*b - B*a)/(a**2*b*e - a*b**2*d + x*(a*b**2*e - b**3*d)) - (-A*e + B*d)*log(x + (-A*a*e**2 - A*b*d*e + B*a*d*e + B*b*d**2 - a**3*e**3*(-A*e + B*d)/(a*e - b*d)**2 + 3*a**2*b*d*e**2*(-A*e + B*d)/(a*e - b*d)**2 - 3*a*b**2*d**2*e*(-A*e + B*d)/(a*e - b*d)**2 + b**3*d**3*(-A*e + B*d)/(a*e - b*d)**2)/(-2*A*b*e**2 + 2*B*b*d*e))/(a*e - b*d)**2 + (-A*e + B*d)*log(x + (-A*a*e**2 - A*b*d*e + B*a*d*e + B*b*d**2 + a**3*e**3*(-A*e + B*d)/(a*e - b*d)**2 - 3*a**2*b*d*e**2*(-A*e + B*d)/(a*e - b*d)**2 + 3*a*b**2*d**2*e*(-A*e + B*d)/(a*e - b*d)**2 - b**3*d**3*(-A*e + B*d)/(a*e - b*d)**2)/(-2*A*b*e**2 + 2*B*b*d*e))/(a*e - b*d)**2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.44

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)} dx = \frac{(Bd - Ae) \log(bx + a)}{b^2d^2 - 2abde + a^2e^2} - \frac{(Bd - Ae) \log(ex + d)}{b^2d^2 - 2abde + a^2e^2}$$

$$+ \frac{Ba - Ab}{ab^2d - a^2be + (b^3d - ab^2e)x}$$

input `integrate((B*x+A)/(b*x+a)^2/(e*x+d), x, algorithm="maxima")`

output

$$(B*d - A*e)*\log(b*x + a)/(b^2*d^2 - 2*a*b*d*e + a^2*e^2) - (B*d - A*e)*\log(e*x + d)/(b^2*d^2 - 2*a*b*d*e + a^2*e^2) + (B*a - A*b)/(a*b^2*d - a^2*b*e + (b^3*d - a*b^2*e)*x)$$
Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.22

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)} dx = -\frac{(Bbd - Abe) \log\left(\left|\frac{bd}{bx+a} - \frac{ae}{bx+a} + e\right|\right)}{b^3d^2 - 2ab^2de + a^2be^2} + \frac{\frac{Ba}{bx+a} - \frac{Ab}{bx+a}}{b^2d - abe}$$

input

```
integrate((B*x+A)/(b*x+a)^2/(e*x+d),x, algorithm="giac")
```

output

$$-(B*b*d - A*b*e)*\log(\text{abs}(b*d/(b*x + a) - a*e/(b*x + a) + e))/(b^3*d^2 - 2*a*b^2*d*e + a^2*b*e^2) + (B*a/(b*x + a) - A*b/(b*x + a))/(b^2*d - a*b*e)$$
Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.15

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)} dx$$

$$= \frac{Ab - Ba}{b(ae - bd)(a + bx)} - \frac{2 \operatorname{atanh}\left(\frac{a^2e^2 - b^2d^2}{(ae - bd)^2} + \frac{2bex}{ae - bd}\right) (Ae - Bd)}{(ae - bd)^2}$$

input

```
int((A + B*x)/((a + b*x)^2*(d + e*x)),x)
```

output

$$(A*b - B*a)/(b*(a*e - b*d)*(a + b*x)) - (2*\operatorname{atanh}((a^2*e^2 - b^2*d^2)/(a*e - b*d)^2 + (2*b*e*x)/(a*e - b*d))*(A*e - B*d))/(a*e - b*d)^2$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.32

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)} dx = \frac{-\log(bx + a) + \log(ex + d)}{ae - bd}$$

input `int((B*x+A)/(b*x+a)^2/(e*x+d),x)`

output `(- log(a + b*x) + log(d + e*x))/(a*e - b*d)`

3.115 $\int \frac{A+Bx}{(a+bx)^2(d+ex)^2} dx$

Optimal result	1141
Mathematica [A] (verified)	1141
Rubi [A] (verified)	1142
Maple [A] (verified)	1143
Fricas [B] (verification not implemented)	1144
Sympy [B] (verification not implemented)	1144
Maxima [B] (verification not implemented)	1145
Giac [A] (verification not implemented)	1146
Mupad [B] (verification not implemented)	1146
Reduce [B] (verification not implemented)	1147

Optimal result

Integrand size = 20, antiderivative size = 117

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^2} dx = -\frac{Ab - aB}{(bd - ae)^2(a + bx)} + \frac{Bd - Ae}{(bd - ae)^2(d + ex)} + \frac{(bBd - 2Abe + aBe) \log(a + bx)}{(bd - ae)^3} - \frac{(bBd - 2Abe + aBe) \log(d + ex)}{(bd - ae)^3}$$

output

```
-(A*b-B*a)/(-a*e+b*d)^2/(b*x+a)+(-A*e+B*d)/(-a*e+b*d)^2/(e*x+d)+(-2*A*b*e+B*a*e+B*b*d)*ln(b*x+a)/(-a*e+b*d)^3-(-2*A*b*e+B*a*e+B*b*d)*ln(e*x+d)/(-a*e+b*d)^3
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.88

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^2} dx = \frac{\frac{(-Ab+aB)(bd-ae)}{a+bx} + \frac{(bd-ae)(Bd-Ae)}{d+ex} + (bBd - 2Abe + aBe) \log(a + bx) - (bBd - 2Abe + aBe) \log(d + ex)}{(bd - ae)^3}$$

input `Integrate[(A + B*x)/((a + b*x)^2*(d + e*x)^2),x]`

output `((((-A*b) + a*B)*(b*d - a*e))/(a + b*x) + ((b*d - a*e)*(B*d - A*e))/(d + e*x) + (b*B*d - 2*A*b*e + a*B*e)*Log[a + b*x] - (b*B*d - 2*A*b*e + a*B*e)*Log[d + e*x])/((b*d - a*e)^3)`

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^2} dx$$

↓ 86

$$\int \left(\frac{b(Ab - aB)}{(a + bx)^2(bd - ae)^2} + \frac{b(aBe - 2Abe + bBd)}{(a + bx)(bd - ae)^3} + \frac{e(-aBe + 2Abe - bBd)}{(d + ex)(bd - ae)^3} + \frac{e(Ae - Bd)}{(d + ex)^2(bd - ae)^2} \right) dx$$

↓ 2009

$$-\frac{Ab - aB}{(a + bx)(bd - ae)^2} + \frac{Bd - Ae}{(d + ex)(bd - ae)^2} + \frac{\log(a + bx)(aBe - 2Abe + bBd)}{(bd - ae)^3} - \frac{\log(d + ex)(aBe - 2Abe + bBd)}{(bd - ae)^3}$$

input `Int[(A + B*x)/((a + b*x)^2*(d + e*x)^2),x]`

output `-((A*b - a*B)/((b*d - a*e)^2*(a + b*x))) + (B*d - A*e)/((b*d - a*e)^2*(d + e*x)) + ((b*B*d - 2*A*b*e + a*B*e)*Log[a + b*x])/((b*d - a*e)^3) - ((b*B*d - 2*A*b*e + a*B*e)*Log[d + e*x])/((b*d - a*e)^3)`

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.05

method	result
default	$\frac{(2Abe - Bae - Bbd) \ln(bx+a)}{(ae-db)^3} - \frac{Ab - Ba}{(ae-db)^2(bx+a)} - \frac{Ae - Bd}{(ae-db)^2(ex+d)} - \frac{(2Abe - Bae - Bbd) \ln(ex+d)}{(ae-db)^3}$
norman	$-\frac{Aab e^2 + A b^2 de - 2Babde}{eb(a^2 e^2 - 2abde + b^2 d^2)} - \frac{(2A b^2 e^2 - Bab e^2 - b^2 Bde)x}{eb(a^2 e^2 - 2abde + b^2 d^2)} + \frac{(2Abe - Bae - Bbd) \ln(bx+a)}{a^3 e^3 - 3a^2 bd e^2 + 3a b^2 d^2 e - b^3 d^3} - \frac{(2Abe - Bae - Bbd) \ln(ex+d)}{a^3 e^3 - 3a^2 bd e^2 + 3a b^2 d^2 e - b^3 d^3}$
risch	$-\frac{(2Abe - Bae - Bbd)x}{a^2 e^2 - 2abde + b^2 d^2} - \frac{Aae + Abd - 2Bad}{a^2 e^2 - 2abde + b^2 d^2} - \frac{2 \ln(ex+d)Abe}{a^3 e^3 - 3a^2 bd e^2 + 3a b^2 d^2 e - b^3 d^3} + \frac{\ln(ex+d)Bae}{a^3 e^3 - 3a^2 bd e^2 + 3a b^2 d^2 e - b^3 d^3} + \frac{1}{a^3 e^3 - 3a^2 bd e^2 + 3a b^2 d^2 e - b^3 d^3}$
parallelrisch	$-B \ln(bx+a)x a^2 b e^3 - B \ln(bx+a)a b^2 d^2 e + B \ln(ex+d)a^2 bd e^2 + B \ln(ex+d)a b^2 d^2 e - B \ln(bx+a)x b^3 d^2 e + B \ln(ex+d)x a^2 b e^3$

```
input int((B*x+A)/(b*x+a)^2/(e*x+d)^2,x,method=_RETURNVERBOSE)
```

```
output (2*A*b*e-B*a*e-B*b*d)/(a*e-b*d)^3*ln(b*x+a)-(A*b-B*a)/(a*e-b*d)^2/(b*x+a)-(A*e-B*d)/(a*e-b*d)^2/(e*x+d)-(2*A*b*e-B*a*e-B*b*d)/(a*e-b*d)^3*ln(e*x+d)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 396 vs. 2(116) = 232.

Time = 0.09 (sec) , antiderivative size = 396, normalized size of antiderivative = 3.38

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^2} dx = \frac{2Ba^2de - Aa^2e^2 - (2Bab - Ab^2)d^2 - (Bb^2d^2 - 2Ab^2de - (Ba^2 - 2Aab)e^2)x - (Babd^2 + (Ba^2 - 2$$

input

```
integrate((B*x+A)/(b*x+a)^2/(e*x+d)^2,x, algorithm="fricas")
```

output

```
-(2*B*a^2*d*e - A*a^2*e^2 - (2*B*a*b - A*b^2)*d^2 - (B*b^2*d^2 - 2*A*b^2*d
*e - (B*a^2 - 2*A*a*b)*e^2)*x - (B*a*b*d^2 + (B*a^2 - 2*A*a*b)*d*e + (B*b^
2*d*e + (B*a*b - 2*A*b^2)*e^2)*x^2 + (B*b^2*d^2 + 2*(B*a*b - A*b^2)*d*e +
(B*a^2 - 2*A*a*b)*e^2)*x)*log(b*x + a) + (B*a*b*d^2 + (B*a^2 - 2*A*a*b)*d*
e + (B*b^2*d*e + (B*a*b - 2*A*b^2)*e^2)*x^2 + (B*b^2*d^2 + 2*(B*a*b - A*b^
2)*d*e + (B*a^2 - 2*A*a*b)*e^2)*x)*log(e*x + d))/(a*b^3*d^4 - 3*a^2*b^2*d^
3*e + 3*a^3*b*d^2*e^2 - a^4*d*e^3 + (b^4*d^3*e - 3*a*b^3*d^2*e^2 + 3*a^2*b
^2*d*e^3 - a^3*b*e^4)*x^2 + (b^4*d^4 - 2*a*b^3*d^3*e + 2*a^3*b*d^2*e^3 - a^4
*e^4)*x)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 706 vs. 2(104) = 208.

Time = 1.33 (sec) , antiderivative size = 706, normalized size of antiderivative = 6.03

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^2} dx = \frac{-Aae - Abd + 2Bad + x(-2Abe + Bae + Bbd)}{a^3de^2 - 2a^2bd^2e + ab^2d^3 + x^2(a^2be^3 - 2ab^2de^2 + b^3d^2e) + x(a^3e^3 - a^2bde^2 - ab^2d^2e + b^3d^3)} + \frac{(-2Abe + Bae + Bbd) \log \left(x + \frac{-2Aabe^2 - 2Ab^2de + Ba^2e^2 + 2Babde + Bb^2d^2 - \frac{a^4e^4(-2Abe + Bae + Bbd)}{(ae - bd)^3} + \frac{4a^3bde^3(-2Abe + Bae + Bbd)}{(ae - bd)^3}}{-4Ab^2e^2 + 2Babe} \right)}{(ae - bd)^3} + \frac{(-2Abe + Bae + Bbd) \log \left(x + \frac{-2Aabe^2 - 2Ab^2de + Ba^2e^2 + 2Babde + Bb^2d^2 + \frac{a^4e^4(-2Abe + Bae + Bbd)}{(ae - bd)^3} - \frac{4a^3bde^3(-2Abe + Bae + Bbd)}{(ae - bd)^3}}{-4Ab^2e^2 + 2Babe} \right)}{(ae - bd)^3}$$

input `integrate((B*x+A)/(b*x+a)**2/(e*x+d)**2,x)`

output
$$\begin{aligned} & (-A*a*e - A*b*d + 2*B*a*d + x*(-2*A*b*e + B*a*e + B*b*d))/(a**3*d*e**2 - 2 \\ & *a**2*b*d**2*e + a*b**2*d**3 + x**2*(a**2*b*e**3 - 2*a*b**2*d*e**2 + b**3* \\ & d**2*e) + x*(a**3*e**3 - a**2*b*d*e**2 - a*b**2*d**2*e + b**3*d**3)) + (-2 \\ & *A*b*e + B*a*e + B*b*d)*\log(x + (-2*A*a*b*e**2 - 2*A*b**2*d*e + B*a**2*e** \\ & 2 + 2*B*a*b*d*e + B*b**2*d**2 - a**4*e**4*(-2*A*b*e + B*a*e + B*b*d)/(a*e \\ & - b*d)**3 + 4*a**3*b*d*e**3*(-2*A*b*e + B*a*e + B*b*d)/(a*e - b*d)**3 - 6* \\ & a**2*b**2*d**2*e**2*(-2*A*b*e + B*a*e + B*b*d)/(a*e - b*d)**3 + 4*a*b**3*d \\ & **3*e*(-2*A*b*e + B*a*e + B*b*d)/(a*e - b*d)**3 - b**4*d**4*(-2*A*b*e + B \\ & a*e + B*b*d)/(a*e - b*d)**3)/(-4*A*b**2*e**2 + 2*B*a*b*e**2 + 2*B*b**2*d*e \\ &))/(a*e - b*d)**3 - (-2*A*b*e + B*a*e + B*b*d)*\log(x + (-2*A*a*b*e**2 - 2* \\ & A*b**2*d*e + B*a**2*e**2 + 2*B*a*b*d*e + B*b**2*d**2 + a**4*e**4*(-2*A*b*e \\ & + B*a*e + B*b*d)/(a*e - b*d)**3 - 4*a**3*b*d*e**3*(-2*A*b*e + B*a*e + B*b \\ & *d)/(a*e - b*d)**3 + 6*a**2*b**2*d**2*e**2*(-2*A*b*e + B*a*e + B*b*d)/(a*e \\ & - b*d)**3 - 4*a*b**3*d**3*e*(-2*A*b*e + B*a*e + B*b*d)/(a*e - b*d)**3 + b \\ & **4*d**4*(-2*A*b*e + B*a*e + B*b*d)/(a*e - b*d)**3)/(-4*A*b**2*e**2 + 2*B* \\ & a*b*e**2 + 2*B*b**2*d*e))/(a*e - b*d)**3 \end{aligned}$$

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 256 vs. $2(116) = 232$.

Time = 0.05 (sec) , antiderivative size = 256, normalized size of antiderivative = 2.19

$$\begin{aligned} & \int \frac{A + Bx}{(a + bx)^2(d + ex)^2} dx \\ & = \frac{(Bbd + (Ba - 2Ab)e) \log(bx + a)}{b^3d^3 - 3ab^2d^2e + 3a^2bde^2 - a^3e^3} - \frac{(Bbd + (Ba - 2Ab)e) \log(ex + d)}{b^3d^3 - 3ab^2d^2e + 3a^2bde^2 - a^3e^3} \\ & \quad - \frac{Aae - (2Ba - Ab)d - (Bbd + (Ba - 2Ab)e)x}{ab^2d^3 - 2a^2bd^2e + a^3de^2 + (b^3d^2e - 2ab^2de^2 + a^2be^3)x^2 + (b^3d^3 - ab^2d^2e - a^2bde^2 + a^3e^3)x} \end{aligned}$$

input `integrate((B*x+A)/(b*x+a)^2/(e*x+d)^2,x, algorithm="maxima")`

output

```
(B*b*d + (B*a - 2*A*b)*e)*log(b*x + a)/(b^3*d^3 - 3*a*b^2*d^2*e + 3*a^2*b*d*e^2 - a^3*e^3) - (B*b*d + (B*a - 2*A*b)*e)*log(e*x + d)/(b^3*d^3 - 3*a*b^2*d^2*e + 3*a^2*b*d*e^2 - a^3*e^3) - (A*a*e - (2*B*a - A*b)*d - (B*b*d + (B*a - 2*A*b)*e)*x)/(a*b^2*d^3 - 2*a^2*b*d^2*e + a^3*d*e^2 + (b^3*d^2*e - 2*a*b^2*d*e^2 + a^2*b*e^3)*x^2 + (b^3*d^3 - a*b^2*d^2*e - a^2*b*d*e^2 + a^3*e^3)*x)
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.66

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^2} dx = -\frac{(Bb^2d + Babe - 2Ab^2e) \log\left(\left|\frac{bd}{bx+a} - \frac{ae}{bx+a} + e\right|\right)}{b^4d^3 - 3ab^3d^2e + 3a^2b^2de^2 - a^3be^3} + \frac{\frac{Bab^2}{bx+a} - \frac{Ab^3}{bx+a}}{b^4d^2 - 2ab^3de + a^2b^2e^2} - \frac{Bbde - Abe^2}{(bd - ae)^3\left(\frac{bd}{bx+a} - \frac{ae}{bx+a} + e\right)}$$

input

```
integrate((B*x+A)/(b*x+a)^2/(e*x+d)^2,x, algorithm="giac")
```

output

```
-(B*b^2*d + B*a*b*e - 2*A*b^2*e)*log(abs(b*d/(b*x + a) - a*e/(b*x + a) + e))/(b^4*d^3 - 3*a*b^3*d^2*e + 3*a^2*b^2*d*e^2 - a^3*b*e^3) + (B*a*b^2/(b*x + a) - A*b^3/(b*x + a))/(b^4*d^2 - 2*a*b^3*d*e + a^2*b^2*e^2) - (B*b*d*e - A*b*e^2)/((b*d - a*e)^3*(b*d/(b*x + a) - a*e/(b*x + a) + e))
```

Mupad [B] (verification not implemented)

Time = 1.10 (sec) , antiderivative size = 263, normalized size of antiderivative = 2.25

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^2} dx = -\frac{\frac{Aae+Abd-2Bad}{a^2e^2-2abde+b^2d^2} - \frac{x(Bae-2Abe+Bbd)}{a^2e^2-2abde+b^2d^2}}{be^2x + (ae + bd)x + ad} - \frac{2 \operatorname{atanh}\left(\frac{\left(\frac{a^3e^3 - a^2bde^2 - ab^2d^2e + b^3d^3}{a^2e^2 - 2abde + b^2d^2} + 2bex\right) (e(2Ab - Ba) - Bbd) (a^2e^2 - 2abde + b^2d^2)}{(ae - bd)^3 (Bae - 2Abe + Bbd)}\right)}{(ae - bd)^3} (e(2Ab - Ba) - Bbd)$$

input

```
int((A + B*x)/((a + b*x)^2*(d + e*x)^2),x)
```

output

```
- ((A*a*e + A*b*d - 2*B*a*d)/(a^2*e^2 + b^2*d^2 - 2*a*b*d*e) - (x*(B*a*e -
2*A*b*e + B*b*d))/(a^2*e^2 + b^2*d^2 - 2*a*b*d*e))/(a*d + x*(a*e + b*d) +
b*e*x^2) - (2*atanh(((a^3*e^3 + b^3*d^3 - a*b^2*d^2*e - a^2*b*d*e^2)/(a^
2*e^2 + b^2*d^2 - 2*a*b*d*e) + 2*b*e*x)*(e*(2*A*b - B*a) - B*b*d)*(a^2*e^2
+ b^2*d^2 - 2*a*b*d*e))/((a*e - b*d)^3*(B*a*e - 2*A*b*e + B*b*d)))*(e*(2*
A*b - B*a) - B*b*d))/(a*e - b*d)^3
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.98

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^2} dx$$

$$= \frac{\log(bx + a) b d^2 + \log(bx + a) b d e x - \log(ex + d) b d^2 - \log(ex + d) b d e x + a e^2 x - b d e x}{d(a^2 e^3 x - 2 a b d e^2 x + b^2 d^2 e x + a^2 d e^2 - 2 a b d^2 e + b^2 d^3)}$$

input

```
int((B*x+A)/(b*x+a)^2/(e*x+d)^2,x)
```

output

```
(log(a + b*x)*b*d**2 + log(a + b*x)*b*d*e*x - log(d + e*x)*b*d**2 - log(d
+ e*x)*b*d*e*x + a*e**2*x - b*d*e*x)/(d*(a**2*d*e**2 + a**2*e**3*x - 2*a*b
*d**2*e - 2*a*b*d*e**2*x + b**2*d**3 + b**2*d**2*e*x))
```


3.116 $\int \frac{A+Bx}{(a+bx)^2(d+ex)^3} dx$

Optimal result	1148
Mathematica [A] (verified)	1148
Rubi [A] (verified)	1149
Maple [A] (verified)	1150
Fricas [B] (verification not implemented)	1151
Sympy [B] (verification not implemented)	1151
Maxima [B] (verification not implemented)	1152
Giac [A] (verification not implemented)	1153
Mupad [B] (verification not implemented)	1154
Reduce [B] (verification not implemented)	1154

Optimal result

Integrand size = 20, antiderivative size = 157

$$\int \frac{A+Bx}{(a+bx)^2(d+ex)^3} dx = -\frac{b(Ab-aB)}{(bd-ae)^3(a+bx)} + \frac{Bd-Ae}{2(bd-ae)^2(d+ex)^2} + \frac{bBd-2Abe+aBe}{(bd-ae)^3(d+ex)} + \frac{b(bBd-3Abe+2aBe)\log(a+bx)}{(bd-ae)^4} - \frac{b(bBd-3Abe+2aBe)\log(d+ex)}{(bd-ae)^4}$$

output

```
-b*(A*b-B*a)/(-a*e+b*d)^3/(b*x+a)+1/2*(-A*e+B*d)/(-a*e+b*d)^2/(e*x+d)^2+(-2*A*b*e+B*a*e+B*b*d)/(-a*e+b*d)^3/(e*x+d)+b*(-3*A*b*e+2*B*a*e+B*b*d)*ln(b*x+a)/(-a*e+b*d)^4-b*(-3*A*b*e+2*B*a*e+B*b*d)*ln(e*x+d)/(-a*e+b*d)^4
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.93

$$\int \frac{A+Bx}{(a+bx)^2(d+ex)^3} dx = \frac{-\frac{2b(Ab-aB)(bd-ae)}{a+bx} + \frac{(bd-ae)^2(Bd-Ae)}{(d+ex)^2} + \frac{2(bd-ae)(bBd-2Abe+aBe)}{d+ex} + 2b(bBd-3Abe+2aBe)\log(a+bx) - 2b(bBd-3Abe+2aBe)\log(d+ex)}{2(bd-ae)^4}$$

input `Integrate[(A + B*x)/((a + b*x)^2*(d + e*x)^3),x]`

output
$$\frac{((-2*b*(A*b - a*B)*(b*d - a*e))/(a + b*x) + ((b*d - a*e)^2*(B*d - A*e))/(d + e*x)^2 + (2*(b*d - a*e)*(b*B*d - 2*A*b*e + a*B*e))/(d + e*x) + 2*b*(b*B*d - 3*A*b*e + 2*a*B*e)*\text{Log}[a + b*x] - 2*b*(b*B*d - 3*A*b*e + 2*a*B*e)*\text{Log}[d + e*x])}{(2*(b*d - a*e)^4)}$$

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^3} dx$$

↓ 86

$$\int \left(\frac{b^2(2aBe - 3Abe + bBd)}{(a + bx)(bd - ae)^4} + \frac{b^2(Ab - aB)}{(a + bx)^2(bd - ae)^3} + \frac{be(-2aBe + 3Abe - bBd)}{(d + ex)(bd - ae)^4} + \frac{e(-aBe + 2Abe - bBd)}{(d + ex)^2(bd - ae)^3} + \right.$$

↓ 2009

$$\left. - \frac{b(Ab - aB)}{(a + bx)(bd - ae)^3} + \frac{aBe - 2Abe + bBd}{(d + ex)(bd - ae)^3} + \frac{Bd - Ae}{2(d + ex)^2(bd - ae)^2} + \frac{b \log(a + bx)(2aBe - 3Abe + bBd)}{(bd - ae)^4} - \frac{b \log(d + ex)(2aBe - 3Abe + bBd)}{(bd - ae)^4} \right)$$

input `Int[(A + B*x)/((a + b*x)^2*(d + e*x)^3),x]`

output
$$\frac{-((b*(A*b - a*B))/((b*d - a*e)^3*(a + b*x))) + (B*d - A*e)/(2*(b*d - a*e)^2*(d + e*x)^2) + (b*B*d - 2*A*b*e + a*B*e)/((b*d - a*e)^3*(d + e*x)) + (b*(b*B*d - 3*A*b*e + 2*a*B*e)*\text{Log}[a + b*x])/(b*d - a*e)^4 - (b*(b*B*d - 3*A*b*e + 2*a*B*e)*\text{Log}[d + e*x])/(b*d - a*e)^4}{(b*d - a*e)^4}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 801 vs. $2(154) = 308$.

Time = 0.12 (sec) , antiderivative size = 801, normalized size of antiderivative = 5.10

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^3} dx = \text{Too large to display}$$

input `integrate((B*x+A)/(b*x+a)^2/(e*x+d)^3,x, algorithm="fricas")`

output

```
-1/2*(A*a^3*e^3 - (5*B*a*b^2 - 2*A*b^3)*d^3 + (4*B*a^2*b + 3*A*a*b^2)*d^2*
e + (B*a^3 - 6*A*a^2*b)*d*e^2 - 2*(B*b^3*d^2*e + (B*a*b^2 - 3*A*b^3)*d*e^2
- (2*B*a^2*b - 3*A*a*b^2)*e^3)*x^2 - (3*B*b^3*d^3 + (4*B*a*b^2 - 9*A*b^3)
*d^2*e - (5*B*a^2*b - 6*A*a*b^2)*d*e^2 - (2*B*a^3 - 3*A*a^2*b)*e^3)*x - 2*
(B*a*b^2*d^3 + (2*B*a^2*b - 3*A*a*b^2)*d^2*e + (B*b^3*d*e^2 + (2*B*a*b^2 -
3*A*b^3)*e^3)*x^3 + (2*B*b^3*d^2*e + (5*B*a*b^2 - 6*A*b^3)*d*e^2 + (2*B*a
^2*b - 3*A*a*b^2)*e^3)*x^2 + (B*b^3*d^3 + (4*B*a*b^2 - 3*A*b^3)*d^2*e + 2*
(2*B*a^2*b - 3*A*a*b^2)*d*e^2)*x)*log(b*x + a) + 2*(B*a*b^2*d^3 + (2*B*a^2
*b - 3*A*a*b^2)*d^2*e + (B*b^3*d*e^2 + (2*B*a*b^2 - 3*A*b^3)*e^3)*x^3 + (2
*B*b^3*d^2*e + (5*B*a*b^2 - 6*A*b^3)*d*e^2 + (2*B*a^2*b - 3*A*a*b^2)*e^3)*
x^2 + (B*b^3*d^3 + (4*B*a*b^2 - 3*A*b^3)*d^2*e + 2*(2*B*a^2*b - 3*A*a*b^2)
*d*e^2)*x)*log(e*x + d))/(a*b^4*d^6 - 4*a^2*b^3*d^5*e + 6*a^3*b^2*d^4*e^2
- 4*a^4*b*d^3*e^3 + a^5*d^2*e^4 + (b^5*d^4*e^2 - 4*a*b^4*d^3*e^3 + 6*a^2*b
^3*d^2*e^4 - 4*a^3*b^2*d*e^5 + a^4*b*e^6)*x^3 + (2*b^5*d^5*e - 7*a*b^4*d^4
*e^2 + 8*a^2*b^3*d^3*e^3 - 2*a^3*b^2*d^2*e^4 - 2*a^4*b*d*e^5 + a^5*e^6)*x^
2 + (b^5*d^6 - 2*a*b^4*d^5*e - 2*a^2*b^3*d^4*e^2 + 8*a^3*b^2*d^3*e^3 - 7*a
^4*b*d^2*e^4 + 2*a^5*d*e^5)*x)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1066 vs. $2(148) = 296$.

Time = 1.98 (sec) , antiderivative size = 1066, normalized size of antiderivative = 6.79

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^3} dx = \text{Too large to display}$$

input `integrate((B*x+A)/(b*x+a)**2/(e*x+d)**3,x)`

output

```
(B*b^2*d + (2*B*a*b - 3*A*b^2)*e)*log(b*x + a)/(b^4*d^4 - 4*a*b^3*d^3*e +
6*a^2*b^2*d^2*e^2 - 4*a^3*b*d*e^3 + a^4*e^4) - (B*b^2*d + (2*B*a*b - 3*A*b
^2)*e)*log(e*x + d)/(b^4*d^4 - 4*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2 - 4*a^3*b
*d*e^3 + a^4*e^4) + 1/2*(A*a^2*e^2 + (5*B*a*b - 2*A*b^2)*d^2 + (B*a^2 - 5*
A*a*b)*d*e + 2*(B*b^2*d*e + (2*B*a*b - 3*A*b^2)*e^2)*x^2 + (3*B*b^2*d^2 +
(7*B*a*b - 9*A*b^2)*d*e + (2*B*a^2 - 3*A*a*b)*e^2)*x)/(a*b^3*d^5 - 3*a^2*b
^2*d^4*e + 3*a^3*b*d^3*e^2 - a^4*d^2*e^3 + (b^4*d^3*e^2 - 3*a*b^3*d^2*e^3
+ 3*a^2*b^2*d*e^4 - a^3*b*e^5)*x^3 + (2*b^4*d^4*e - 5*a*b^3*d^3*e^2 + 3*a^
2*b^2*d^2*e^3 + a^3*b*d*e^4 - a^4*e^5)*x^2 + (b^4*d^5 - a*b^3*d^4*e - 3*a^
2*b^2*d^3*e^2 + 5*a^3*b*d^2*e^3 - 2*a^4*d*e^4)*x)
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 306, normalized size of antiderivative = 1.95

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^3} dx$$

$$= -\frac{(Bb^3d + 2Bab^2e - 3Ab^3e) \log\left(\left|\frac{bd}{bx+a} - \frac{ae}{bx+a} + e\right|\right)}{b^5d^4 - 4ab^4d^3e + 6a^2b^3d^2e^2 - 4a^3b^2de^3 + a^4be^4}$$

$$+ \frac{\frac{Bab^4}{bx+a} - \frac{Ab^5}{bx+a}}{b^6d^3 - 3ab^5d^2e + 3a^2b^4de^2 - a^3b^3e^3}$$

$$- \frac{3Bb^2de^2 + 2Babe^3 - 5Ab^2e^3 + \frac{2(2Bb^4d^2e - Bab^3de^2 - 3Ab^4de^2 - Ba^2b^2e^3 + 3Aab^3e^3)}{(bx+a)b}}{2(bd - ae)^4 \left(\frac{bd}{bx+a} - \frac{ae}{bx+a} + e\right)^2}$$

input

```
integrate((B*x+A)/(b*x+a)^2/(e*x+d)^3,x, algorithm="giac")
```

output

```
-(B*b^3*d + 2*B*a*b^2*e - 3*A*b^3*e)*log(abs(b*d/(b*x + a) - a*e/(b*x + a)
+ e))/(b^5*d^4 - 4*a*b^4*d^3*e + 6*a^2*b^3*d^2*e^2 - 4*a^3*b^2*d*e^3 + a^
4*b*e^4) + (B*a*b^4/(b*x + a) - A*b^5/(b*x + a))/(b^6*d^3 - 3*a*b^5*d^2*e
+ 3*a^2*b^4*d*e^2 - a^3*b^3*e^3) - 1/2*(3*B*b^2*d*e^2 + 2*B*a*b*e^3 - 5*A*
b^2*e^3 + 2*(2*B*b^4*d^2*e - B*a*b^3*d*e^2 - 3*A*b^4*d*e^2 - B*a^2*b^2*e^3
+ 3*A*a*b^3*e^3)/((b*x + a)*b))/((b*d - a*e)^4*(b*d/(b*x + a) - a*e/(b*x
+ a) + e)^2)
```

Mupad [B] (verification not implemented)

Time = 1.20 (sec) , antiderivative size = 454, normalized size of antiderivative = 2.89

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^3} dx$$

$$= \frac{2 \operatorname{atanh}\left(\frac{(b^2(3Ae - Bd) - 2Babe)\left(\frac{a^4e^4 - 2a^3bde^3 + 2ab^3d^3e - b^4d^4}{a^3e^3 - 3a^2bde^2 + 3ab^2d^2e - b^3d^3} + 2bex\right)(a^3e^3 - 3a^2bde^2 + 3ab^2d^2e - b^3d^3)}{(ae - bd)^4(Bb^2d - 3Aa^2e + 2Babe)}\right)(b^2(3Ae - Bd) - 2Babe)}{(ae - bd)^4}$$

$$- \frac{\frac{Ba^2de + Aa^2e^2 + 5Babd^2 - 5Aabde - 2Ab^2d^2}{2(a^3e^3 - 3a^2bde^2 + 3ab^2d^2e - b^3d^3)} + \frac{x(ae + 3bd)(2Bae - 3Abe + Bbd)}{2(a^3e^3 - 3a^2bde^2 + 3ab^2d^2e - b^3d^3)} + \frac{bex^2(2Bae - 3Abe + Bbd)}{a^3e^3 - 3a^2bde^2 + 3ab^2d^2e - b^3d^3}}{x(bd^2 + 2aed) + a^2d^2 + x^2(ae^2 + 2bde) + be^2x^3}$$

input `int((A + B*x)/((a + b*x)^2*(d + e*x)^3),x)`

output

$$\begin{aligned} & (2*\operatorname{atanh}(((b^2*(3*A*e - B*d) - 2*B*a*b*e)*((a^4*e^4 - b^4*d^4 + 2*a*b^3*d^3*e - 2*a^3*b*d*e^3)/(a^3*e^3 - b^3*d^3 + 3*a*b^2*d^2*e - 3*a^2*b*d*e^2) + 2*b*e*x)*(a^3*e^3 - b^3*d^3 + 3*a*b^2*d^2*e - 3*a^2*b*d*e^2)))/((a*e - b*d)^4*(B*b^2*d - 3*A*b^2*e + 2*B*a*b*e)))*(b^2*(3*A*e - B*d) - 2*B*a*b*e)/((a*e - b*d)^4 - ((A*a^2*e^2 - 2*A*b^2*d^2 + 5*B*a*b*d^2 + B*a^2*d*e - 5*A*a*b*d*e)/(2*(a^3*e^3 - b^3*d^3 + 3*a*b^2*d^2*e - 3*a^2*b*d*e^2)) + (x*(a*e + 3*b*d)*(2*B*a*e - 3*A*b*e + B*b*d))/(2*(a^3*e^3 - b^3*d^3 + 3*a*b^2*d^2*e - 3*a^2*b*d*e^2)) + (b*e*x^2*(2*B*a*e - 3*A*b*e + B*b*d))/(a^3*e^3 - b^3*d^3 + 3*a*b^2*d^2*e - 3*a^2*b*d*e^2)))/(x*(b*d^2 + 2*a*d*e) + a*d^2 + x^2*(a*e^2 + 2*b*d*e) + b*e^2*x^3) \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.84

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^3} dx$$

$$= \frac{-2 \log(bx + a) b^2 d^3 - 4 \log(bx + a) b^2 d^2 ex - 2 \log(bx + a) b^2 d e^2 x^2 + 2 \log(ex + d) b^2 d^3 + 4 \log(ex + d) b^2 d^2 ex + 2 \log(ex + d) b^2 d e^2 x^2}{2d(a^3e^5x^2 - 3a^2bde^4x^2 + 3ab^2d^2e^3x^2 - b^3d^3e^2x^2 + 2a^3de^4x - 6a^2bd^2e^3x + 6ab^2d^2e^2x^2)}$$

input `int((B*x+A)/(b*x+a)^2/(e*x+d)^3,x)`

output

```
( - 2*log(a + b*x)*b**2*d**3 - 4*log(a + b*x)*b**2*d**2*e*x - 2*log(a + b*x)*b**2*d*e**2*x**2 + 2*log(d + e*x)*b**2*d**3 + 4*log(d + e*x)*b**2*d**2*e*x + 2*log(d + e*x)*b**2*d*e**2*x**2 - a**2*d*e**2 + 3*a*b*d**2*e - a*b*e**3*x**2 - 2*b**2*d**3 + b**2*d*e**2*x**2)/(2*d*(a**3*d**2*e**3 + 2*a**3*d**e**4*x + a**3*e**5*x**2 - 3*a**2*b*d**3*e**2 - 6*a**2*b*d**2*e**3*x - 3*a**2*b*d*e**4*x**2 + 3*a*b**2*d**4*e + 6*a*b**2*d**3*e**2*x + 3*a*b**2*d**2*e**3*x**2 - b**3*d**5 - 2*b**3*d**4*e*x - b**3*d**3*e**2*x**2))
```


3.117 $\int \frac{(A+Bx)(d+ex)^3}{(a+bx)^3} dx$

Optimal result	1156
Mathematica [A] (verified)	1156
Rubi [A] (verified)	1157
Maple [A] (verified)	1158
Fricas [B] (verification not implemented)	1159
Sympy [B] (verification not implemented)	1160
Maxima [B] (verification not implemented)	1160
Giac [B] (verification not implemented)	1161
Mupad [B] (verification not implemented)	1162
Reduce [B] (verification not implemented)	1162

Optimal result

Integrand size = 20, antiderivative size = 141

$$\int \frac{(A+Bx)(d+ex)^3}{(a+bx)^3} dx = \frac{e^2(3bBd + Abe - 3aBe)x}{b^4} + \frac{Be^3x^2}{2b^3} - \frac{(Ab - aB)(bd - ae)^3}{2b^5(a+bx)^2} - \frac{(bd - ae)^2(bBd + 3Abe - 4aBe)}{b^5(a+bx)} + \frac{3e(bd - ae)(bBd + Abe - 2aBe) \log(a+bx)}{b^5}$$

```
output e^2*(A*b*e-3*B*a*e+3*B*b*d)*x/b^4+1/2*B*e^3*x^2/b^3-1/2*(A*b-B*a)*(-a*e+b*d)^3/b^5/(b*x+a)^2-(-a*e+b*d)^2*(3*A*b*e-4*B*a*e+B*b*d)/b^5/(b*x+a)+3*e*(-a*e+b*d)*(A*b*e-2*B*a*e+B*b*d)*ln(b*x+a)/b^5
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.74

$$\int \frac{(A+Bx)(d+ex)^3}{(a+bx)^3} dx = \frac{-Ab(5a^3e^3 + a^2be^2(-9d + 4ex) + ab^2e(3d^2 - 12dex - 4e^2x^2) + b^3(d^3 + 6d^2ex - 2e^3x^3)) + B(7a^4e^3 + a^3be^2(-9d + 4ex) + ab^2e(3d^2 - 12dex - 4e^2x^2) + b^3(d^3 + 6d^2ex - 2e^3x^3))}{(a+bx)^3} + B \log(a+bx)$$

input `Integrate[((A + B*x)*(d + e*x)^3)/(a + b*x)^3,x]`

output
$$\begin{aligned} & (-(A*b*(5*a^3*e^3 + a^2*b*e^2*(-9*d + 4*e*x) + a*b^2*e*(3*d^2 - 12*d*e*x - \\ & 4*e^2*x^2) + b^3*(d^3 + 6*d^2*e*x - 2*e^3*x^3))) + B*(7*a^4*e^3 + a^3*b*e \\ & ^2*(-15*d + 2*e*x) + a^2*b^2*e*(9*d^2 - 12*d*e*x - 11*e^2*x^2) + b^4*x*(-2 \\ & *d^3 + 6*d*e^2*x^2 + e^3*x^3) - a*b^3*(d^3 - 12*d^2*e*x - 12*d*e^2*x^2 + 4 \\ & *e^3*x^3)) + 6*e*(b*d - a*e)*(b*B*d + A*b*e - 2*a*B*e)*(a + b*x)^2*\text{Log}[a + \\ & b*x])/(2*b^5*(a + b*x)^2) \end{aligned}$$

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(A + Bx)(d + ex)^3}{(a + bx)^3} dx \\ & \quad \downarrow 86 \\ & \int \left(\frac{e^2(-3aBe + Abe + 3bBd)}{b^4} + \frac{3e(bd - ae)(-2aBe + Abe + bBd)}{b^4(a + bx)} + \frac{(bd - ae)^2(-4aBe + 3Abe + bBd)}{b^4(a + bx)^2} + \frac{(A + Bx)(d + ex)^3}{(a + bx)^3} \right) dx \\ & \quad \downarrow 2009 \\ & -\frac{(bd - ae)^2(-4aBe + 3Abe + bBd)}{b^5(a + bx)} - \frac{(Ab - aB)(bd - ae)^3}{2b^5(a + bx)^2} + \\ & \frac{3e(bd - ae) \log(a + bx)(-2aBe + Abe + bBd)}{b^5} + \frac{e^2x(-3aBe + Abe + 3bBd)}{b^4} + \frac{Be^3x^2}{2b^3} \end{aligned}$$

input `Int[((A + B*x)*(d + e*x)^3)/(a + b*x)^3,x]`

output

$$\frac{(e^{2(3bBd + A*be - 3a*Be)*x})/b^4 + (B*e^{3*x^2})/(2*b^3) - ((A*b - a*B)*(b*d - a*e)^3)/(2*b^5*(a + b*x)^2) - ((b*d - a*e)^2*(b*B*d + 3*A*b*e - 4*a*B*e))/(b^5*(a + b*x)) + (3*e*(b*d - a*e)*(b*B*d + A*b*e - 2*a*B*e)*\text{Log}[a + b*x])/b^5$$

Defintions of rubi rules used

rule 86

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 268, normalized size of antiderivative = 1.90

method	result
default	$\frac{e^2 \left(\frac{1}{2} B b e x^2 + A b e x - 3 B a e x + 3 B b d x \right)}{b^4} - \frac{-A a^3 b e^3 + 3 A a^2 b^2 d e^2 - 3 A a b^3 d^2 e + A b^4 d^3 + B a^4 e^3 - 3 B a^3 b d e^2 + 3 B a^2 b^2 d^2 e - B a b^3 d^3}{2 b^5 (b x + a)^2}$
norman	$\frac{e^2 (A b e - 2 B a e + 3 B b d) x^3 - 9 A a^3 b e^3 - 9 A a^2 b^2 d e^2 + 3 A a b^3 d^2 e + A b^4 d^3 - 18 B a^4 e^3 + 27 B a^3 b d e^2 - 9 B a^2 b^2 d^2 e + B a b^3 d^3 - (6 A a^2 b e^3 - 6 A a b^2 d e^2 + 6 A a^2 b^2 d^2 e - 6 A a b^3 d^3)}{b^2 (b x + a)^2}$
risch	$\frac{B e^3 x^2}{2 b^3} + \frac{e^3 A x}{b^3} - \frac{3 e^3 B a x}{b^4} + \frac{3 e^2 B d x}{b^3} + \frac{(-3 A a^2 b e^3 + 6 A a b^2 d e^2 - 3 A b^3 d^2 e + 4 B a^3 e^3 - 9 B a^2 b d e^2 + 6 B a b^2 d^2 e - b^3 B a^2 d^3)}{b^5 (b x + a)^2}$
parallelrisc	$- \frac{6 A \ln(b x + a) a^2 b^2 d e^2 - 9 B a^2 b^2 d^2 e + 18 B \ln(b x + a) x^2 a b^3 d e^2 + 9 A a^3 b e^3 + 27 B a^3 b d e^2 - 9 A a^2 b^2 d e^2 + 3 A a b^3 d^2 e + B a b^3 d^3}{b^5 (b x + a)^2}$

input

```
int((B*x+A)*(e*x+d)^3/(b*x+a)^3,x,method=_RETURNVERBOSE)
```

output

```
e^2/b^4*(1/2*B*b*e*x^2+A*b*e*x-3*B*a*e*x+3*B*b*d*x)-1/2/b^5*(-A*a^3*b*e^3+
3*A*a^2*b^2*d*e^2-3*A*a*b^3*d^2*e+A*b^4*d^3+B*a^4*e^3-3*B*a^3*b*d*e^2+3*B*
a^2*b^2*d^2*e-B*a*b^3*d^3)/(b*x+a)^2-1/b^5*(3*A*a^2*b*e^3-6*A*a*b^2*d*e^2+
3*A*b^3*d^2*e-4*B*a^3*e^3+9*B*a^2*b*d*e^2-6*B*a*b^2*d^2*e+B*b^3*d^3)/(b*x+
a)-3/b^5*e*(A*a*b*e^2-A*b^2*d*e-2*B*a^2*e^2+3*B*a*b*d*e-B*b^2*d^2)*ln(b*x+
a)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 442 vs. $2(137) = 274$.

Time = 0.10 (sec) , antiderivative size = 442, normalized size of antiderivative = 3.13

$$\int \frac{(A + Bx)(d + ex)^3}{(a + bx)^3} dx$$

$$= \frac{Bb^4e^3x^4 - (Bab^3 + Ab^4)d^3 + 3(3Ba^2b^2 - Aab^3)d^2e - 3(5Ba^3b - 3Aa^2b^2)de^2 + (7Ba^4 - 5Aa^3b)e^3 +$$

input

```
integrate((B*x+A)*(e*x+d)^3/(b*x+a)^3,x, algorithm="fricas")
```

output

```
1/2*(B*b^4*e^3*x^4 - (B*a*b^3 + A*b^4)*d^3 + 3*(3*B*a^2*b^2 - A*a*b^3)*d^2
*e - 3*(5*B*a^3*b - 3*A*a^2*b^2)*d*e^2 + (7*B*a^4 - 5*A*a^3*b)*e^3 + 2*(3*
B*b^4*d*e^2 - (2*B*a*b^3 - A*b^4)*e^3)*x^3 + (12*B*a*b^3*d*e^2 - (11*B*a^2
*b^2 - 4*A*a*b^3)*e^3)*x^2 - 2*(B*b^4*d^3 - 3*(2*B*a*b^3 - A*b^4)*d^2*e +
6*(B*a^2*b^2 - A*a*b^3)*d*e^2 - (B*a^3*b - 2*A*a^2*b^2)*e^3)*x + 6*(B*a^2*
b^2*d^2*e - (3*B*a^3*b - A*a^2*b^2)*d*e^2 + (2*B*a^4 - A*a^3*b)*e^3 + (B*b
^4*d^2*e - (3*B*a*b^3 - A*b^4)*d*e^2 + (2*B*a^2*b^2 - A*a*b^3)*e^3)*x^2 +
2*(B*a*b^3*d^2*e - (3*B*a^2*b^2 - A*a*b^3)*d*e^2 + (2*B*a^3*b - A*a^2*b^2)
*e^3)*x)*log(b*x + a))/(b^7*x^2 + 2*a*b^6*x + a^2*b^5)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 299 vs. $2(141) = 282$.

Time = 2.04 (sec) , antiderivative size = 299, normalized size of antiderivative = 2.12

$$\int \frac{(A+Bx)(d+ex)^3}{(a+bx)^3} dx = \frac{Be^3x^2}{2b^3} + x \left(\frac{Ae^3}{b^3} - \frac{3Bae^3}{b^4} + \frac{3Bde^2}{b^3} \right) + \frac{-5Aa^3be^3 + 9Aa^2b^2de^2 - 3Aab^3d^2e - Ab^4d^3 + 7Ba^4e^3 - 15Ba^3bde^2 + 9Ba^2b^2d^2e - Bab^3d^3 + x(-6Aa^2b^2d^2e + 12Aa^2b^2d^2e - 6Aa^2b^2d^2e + 8Bab^3d^3)}{2a^2b^5 + 4ab^6x + 2b^7x^2} + \frac{3e(ae-bd)(-Abe + 2Bae - Bbd) \log(a+bx)}{b^5}$$

input `integrate((B*x+A)*(e*x+d)**3/(b*x+a)**3,x)`

output `B*e**3*x**2/(2*b**3) + x*(A*e**3/b**3 - 3*B*a*e**3/b**4 + 3*B*d*e**2/b**3) + (-5*A*a**3*b*e**3 + 9*A*a**2*b**2*d*e**2 - 3*A*a*b**3*d**2*e - A*b**4*d**3 + 7*B*a**4*e**3 - 15*B*a**3*b*d*e**2 + 9*B*a**2*b**2*d**2*e - B*a*b**3*d**3 + x*(-6*A*a**2*b**2*d**2*e**3 + 12*A*a*b**3*d*e**2 - 6*A*b**4*d**2*e + 8*B*a**3*b*e**3 - 18*B*a**2*b**2*d*e**2 + 12*B*a*b**3*d**2*e - 2*B*b**4*d**3)))/(2*a**2*b**5 + 4*a*b**6*x + 2*b**7*x**2) + 3*e*(a*e - b*d)*(-A*b*e + 2*B*a*e - B*b*d)*log(a + b*x)/b**5`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 282 vs. $2(137) = 274$.

Time = 0.04 (sec) , antiderivative size = 282, normalized size of antiderivative = 2.00

$$\int \frac{(A+Bx)(d+ex)^3}{(a+bx)^3} dx = \frac{(Bab^3 + Ab^4)d^3 - 3(3Ba^2b^2 - Aab^3)d^2e + 3(5Ba^3b - 3Aa^2b^2)de^2 - (7Ba^4 - 5Aa^3b)e^3 + 2(Bb^4d^3 - Bbe^3x^2 + 2(3Bbde^2 - (3Ba - Ab)e^3)x)}{2(b^7x^2 + 2ab^6x + a^2b^5)} + \frac{3(Bb^2d^2e - (3Bab - Ab^2)de^2 + (2Ba^2 - Aab)e^3) \log(bx+a)}{b^5}$$

input `integrate((B*x+A)*(e*x+d)^3/(b*x+a)^3,x, algorithm="maxima")`

output

```
-1/2*((B*a*b^3 + A*b^4)*d^3 - 3*(3*B*a^2*b^2 - A*a*b^3)*d^2*e + 3*(5*B*a^3*b - 3*A*a^2*b^2)*d*e^2 - (7*B*a^4 - 5*A*a^3*b)*e^3 + 2*(B*b^4*d^3 - 3*(2*B*a*b^3 - A*b^4)*d^2*e + 3*(3*B*a^2*b^2 - 2*A*a*b^3)*d*e^2 - (4*B*a^3*b - 3*A*a^2*b^2)*e^3)*x)/(b^7*x^2 + 2*a*b^6*x + a^2*b^5) + 1/2*(B*b*e^3*x^2 + 2*(3*B*b*d*e^2 - (3*B*a - A*b)*e^3)*x)/b^4 + 3*(B*b^2*d^2*e - (3*B*a*b - A*b^2)*d*e^2 + (2*B*a^2 - A*a*b)*e^3)*log(b*x + a)/b^5
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 283 vs. $2(137) = 274$.

Time = 0.12 (sec) , antiderivative size = 283, normalized size of antiderivative = 2.01

$$\int \frac{(A + Bx)(d + ex)^3}{(a + bx)^3} dx$$

$$= \frac{3(Bb^2d^2e - 3Babde^2 + Ab^2de^2 + 2Ba^2e^3 - Aabe^3) \log(|bx + a|)}{b^5} + \frac{Bb^3e^3x^2 + 6Bb^3de^2x - 6Bab^2e^3x + 2Ab^3e^3x}{2b^6} - \frac{Bab^3d^3 + Ab^4d^3 - 9Ba^2b^2d^2e + 3Aab^3d^2e + 15Ba^3bde^2 - 9Aa^2b^2de^2 - 7Ba^4e^3 + 5Aa^3be^3 + 2(Bb^3d^2e - 3Babde^2 + Ab^2de^2 + 2Ba^2e^3 - Aabe^3) \log(|bx + a|)}{2(bx + a)^2b^5}$$

input

```
integrate((B*x+A)*(e*x+d)^3/(b*x+a)^3,x, algorithm="giac")
```

output

```
3*(B*b^2*d^2*e - 3*B*a*b*d*e^2 + A*b^2*d*e^2 + 2*B*a^2*e^3 - A*a*b*e^3)*log(abs(b*x + a))/b^5 + 1/2*(B*b^3*e^3*x^2 + 6*B*b^3*d*e^2*x - 6*B*a*b^2*e^3*x + 2*A*b^3*e^3*x)/b^6 - 1/2*(B*a*b^3*d^3 + A*b^4*d^3 - 9*B*a^2*b^2*d^2*e + 3*A*a*b^3*d^2*e + 15*B*a^3*b*d*e^2 - 9*A*a^2*b^2*d*e^2 - 7*B*a^4*e^3 + 5*A*a^3*b*e^3 + 2*(B*b^4*d^3 - 6*B*a*b^3*d^2*e + 3*A*b^4*d^2*e + 9*B*a^2*b^2*d*e^2 - 6*A*a*b^3*d*e^2 - 4*B*a^3*b*e^3 + 3*A*a^2*b^2*e^3)*x)/((b*x + a)^2*b^5)
```


3.118 $\int \frac{(A+Bx)(d+ex)^2}{(a+bx)^3} dx$

Optimal result	1163
Mathematica [A] (verified)	1163
Rubi [A] (verified)	1164
Maple [A] (verified)	1165
Fricas [B] (verification not implemented)	1166
Sympy [A] (verification not implemented)	1166
Maxima [A] (verification not implemented)	1167
Giac [A] (verification not implemented)	1167
Mupad [B] (verification not implemented)	1168
Reduce [B] (verification not implemented)	1168

Optimal result

Integrand size = 20, antiderivative size = 103

$$\int \frac{(A+Bx)(d+ex)^2}{(a+bx)^3} dx = \frac{Be^2x}{b^3} - \frac{(Ab-aB)(bd-ae)^2}{2b^4(a+bx)^2} - \frac{(bd-ae)(bBd+2Abe-3aBe)}{b^4(a+bx)} + \frac{e(2bBd+Abe-3aBe)\log(a+bx)}{b^4}$$

output

$B*e^2*x/b^3-1/2*(A*b-B*a)*(-a*e+b*d)^2/b^4/(b*x+a)^2-(-a*e+b*d)*(2*A*b*e-3*B*a*e+B*b*d)/b^4/(b*x+a)+e*(A*b*e-3*B*a*e+2*B*b*d)*\ln(b*x+a)/b^4$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.36

$$\int \frac{(A+Bx)(d+ex)^2}{(a+bx)^3} dx = \frac{-Ab(bd-ae)(3ae+b(d+4ex))+B(-5a^3e^2+2a^2be(3d-2ex))+2b^3x(-d^2+e^2x^2)+ab^2(-d^2+8dex)}{2b^4(a+bx)^2}$$

input

`Integrate[((A+B*x)*(d+e*x)^2)/(a+b*x)^3,x]`

output

$$\begin{aligned} & (-A*b*(b*d - a*e)*(3*a*e + b*(d + 4*e*x)) + B*(-5*a^3*e^2 + 2*a^2*b*e*(3 \\ & *d - 2*e*x) + 2*b^3*x*(-d^2 + e^2*x^2) + a*b^2*(-d^2 + 8*d*e*x + 4*e^2*x^2 \\ &)) + 2*e*(2*b*B*d + A*b*e - 3*a*B*e)*(a + b*x)^2*\text{Log}[a + b*x]) / (2*b^4*(a + \\ & b*x)^2) \end{aligned}$$

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(A + Bx)(d + ex)^2}{(a + bx)^3} dx \\ & \quad \downarrow 86 \\ & \int \left(\frac{e(-3aBe + Abe + 2bBd)}{b^3(a + bx)} + \frac{(bd - ae)(-3aBe + 2Abe + bBd)}{b^3(a + bx)^2} + \frac{(Ab - aB)(bd - ae)^2}{b^3(a + bx)^3} + \frac{Be^2}{b^3} \right) dx \\ & \quad \downarrow 2009 \\ & \quad - \frac{(bd - ae)(-3aBe + 2Abe + bBd)}{b^4(a + bx)} - \frac{(Ab - aB)(bd - ae)^2}{2b^4(a + bx)^2} + \\ & \quad \frac{e \log(a + bx)(-3aBe + Abe + 2bBd)}{b^4} + \frac{Be^2 x}{b^3} \end{aligned}$$

input

$$\text{Int}[(A + B*x)*(d + e*x)^2/(a + b*x)^3, x]$$

output

$$\begin{aligned} & (B*e^2*x)/b^3 - ((A*b - a*B)*(b*d - a*e)^2)/(2*b^4*(a + b*x)^2) - ((b*d - \\ & a*e)*(b*B*d + 2*A*b*e - 3*a*B*e))/(b^4*(a + b*x)) + (e*(2*b*B*d + A*b*e - \\ & 3*a*B*e)*\text{Log}[a + b*x])/b^4 \end{aligned}$$

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.52

method	result
default	$\frac{B e^2 x}{b^3} - \frac{A a^2 b e^2 - 2 A a b^2 d e + A b^3 d^2 - B a^3 e^2 + 2 B a^2 b d e - B a b^2 d^2}{2 b^4 (b x + a)^2} - \frac{-2 A a b e^2 + 2 A b^2 d e + 3 B a^2 e^2 - 4 B a b d e + b^2 B d^2}{b^4 (b x + a)} +$
norman	$\frac{(2 A a b e^2 - 2 A b^2 d e - 6 B a^2 e^2 + 4 B a b d e - b^2 B d^2) x + \frac{B e^2 x^3}{b} + \frac{3 A a^2 b e^2 - 2 A a b^2 d e - A b^3 d^2 - 9 B a^3 e^2 + 6 B a^2 b d e - B a b^2 d^2}{2 b^4}}{(b x + a)^2} + \frac{e(A b e - 3 B a d)}{b^4}$
risch	$\frac{B e^2 x}{b^3} + \frac{(2 A a b e^2 - 2 A b^2 d e - 3 B a^2 e^2 + 4 B a b d e - b^2 B d^2) x + \frac{3 A a^2 b e^2 - 2 A a b^2 d e - A b^3 d^2 - 5 B a^3 e^2 + 6 B a^2 b d e - B a b^2 d^2}{2 b}}{b^3 (b x + a)^2} + \frac{e^2}{b^4}$
parallelrisc	$\frac{2 A \ln(b x + a) x^2 b^3 e^2 - 6 B \ln(b x + a) x^2 a b^2 e^2 + 4 B \ln(b x + a) x^2 b^3 d e + 2 B e^2 x^3 b^3 + 4 A \ln(b x + a) x a b^2 e^2 - 12 B \ln(b x + a) x a^2 b e^2 + 4 A \ln(b x + a) x^2 b^3 d e - 6 B \ln(b x + a) x^2 a b^2 d e + 4 B \ln(b x + a) x^2 b^3 d^2 - 6 B a^3 \ln(b x + a) e^2 + 6 B a^2 b \ln(b x + a) d e - 6 B a b^2 \ln(b x + a) d^2}{b^4}$

```
input int((B*x+A)*(e*x+d)^2/(b*x+a)^3,x,method=_RETURNVERBOSE)
```

```
output B*e^2*x/b^3-1/2*(A*a^2*b*e^2-2*A*a*b^2*d*e+A*b^3*d^2-B*a^3*e^2+2*B*a^2*b*d*e-B*a*b^2*d^2)/b^4/(b*x+a)^2-(-2*A*a*b*e^2+2*A*b^2*d*e+3*B*a^2*e^2-4*B*a*b*d*e+B*b^2*d^2)/b^4/(b*x+a)+e*(A*b*e-3*B*a*d+2*B*b*d)*ln(b*x+a)/b^4
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 258 vs. $2(101) = 202$.

Time = 0.09 (sec) , antiderivative size = 258, normalized size of antiderivative = 2.50

$$\int \frac{(A + Bx)(d + ex)^2}{(a + bx)^3} dx$$

$$= \frac{2Bb^3e^2x^3 + 4Bab^2e^2x^2 - (Bab^2 + Ab^3)d^2 + 2(3Ba^2b - Aab^2)de - (5Ba^3 - 3Aa^2b)e^2 - 2(Bb^3d^2 - 2$$

input `integrate((B*x+A)*(e*x+d)^2/(b*x+a)^3,x, algorithm="fricas")`

output $\frac{1}{2} * (2 * B * b^3 * e^2 * x^3 + 4 * B * a * b^2 * e^2 * x^2 - (B * a * b^2 + A * b^3) * d^2 + 2 * (3 * B * a^2 * b - A * a * b^2) * d * e - (5 * B * a^3 - 3 * A * a^2 * b) * e^2 - 2 * (B * b^3 * d^2 - 2 * (2 * B * a * b^2 - A * b^3) * d * e + 2 * (B * a^2 * b - A * a * b^2) * e^2) * x + 2 * (2 * B * a^2 * b * d * e - (3 * B * a^3 - A * a^2 * b) * e^2 + (2 * B * b^3 * d * e - (3 * B * a * b^2 - A * b^3) * e^2) * x^2 + 2 * (2 * B * a * b^2 * d * e - (3 * B * a^2 * b - A * a * b^2) * e^2) * x) * \log(b * x + a) / (b^6 * x^2 + 2 * a * b^5 * x + a^2 * b^4)$

Sympy [A] (verification not implemented)

Time = 1.15 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.82

$$\int \frac{(A + Bx)(d + ex)^2}{(a + bx)^3} dx = \frac{Be^2x}{b^3}$$

$$+ \frac{3Aa^2be^2 - 2Aab^2de - Ab^3d^2 - 5Ba^3e^2 + 6Ba^2bde - Bab^2d^2 + x(4Aab^2e^2 - 4Ab^3de - 6Ba^2be^2 + 8B$$

$$- \frac{e(-Abe + 3Bae - 2Bbd) \log(a + bx)}{b^4}$$

input `integrate((B*x+A)*(e*x+d)**2/(b*x+a)**3,x)`

output $B * e ** 2 * x / b ** 3 + (3 * A * a ** 2 * b * e ** 2 - 2 * A * a * b ** 2 * d * e - A * b ** 3 * d ** 2 - 5 * B * a ** 3 * e ** 2 + 6 * B * a ** 2 * b * d * e - B * a * b ** 2 * d ** 2 + x * (4 * A * a * b ** 2 * e ** 2 - 4 * A * b ** 3 * d * e - 6 * B * a ** 2 * b * e ** 2 + 8 * B * a * b ** 2 * d * e - 2 * B * b ** 3 * d ** 2)) / (2 * a ** 2 * b ** 4 + 4 * a * b ** 5 * x + 2 * b ** 6 * x ** 2) - e * (-A * b * e + 3 * B * a * e - 2 * B * b * d) * \log(a + b * x) / b ** 4$

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.65

$$\int \frac{(A + Bx)(d + ex)^2}{(a + bx)^3} dx = \frac{Be^2x}{b^3} - \frac{(Bab^2 + Ab^3)d^2 - 2(3Ba^2b - Aab^2)de + (5Ba^3 - 3Aa^2b)e^2 + 2(Bb^3d^2 - 2(2Bab^2 - Ab^3)de + (3Ba^2b - 2Aa^2b)e^2)x}{2(b^6x^2 + 2ab^5x + a^2b^4)} + \frac{(2Bbde - (3Ba - Ab)e^2)\log(bx + a)}{b^4}$$

input `integrate((B*x+A)*(e*x+d)^2/(b*x+a)^3,x, algorithm="maxima")`output $B*e^2*x/b^3 - 1/2*((B*a*b^2 + A*b^3)*d^2 - 2*(3*B*a^2*b - A*a*b^2)*d*e + (5*B*a^3 - 3*A*a^2*b)*e^2 + 2*(B*b^3*d^2 - 2*(2*B*a*b^2 - A*b^3)*d*e + (3*B*a^2*b - 2*A*a*b^2)*e^2)*x)/(b^6*x^2 + 2*a*b^5*x + a^2*b^4) + (2*B*b*d*e - (3*B*a - A*b)*e^2)*\log(b*x + a)/b^4$ **Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.52

$$\int \frac{(A + Bx)(d + ex)^2}{(a + bx)^3} dx = \frac{Be^2x}{b^3} + \frac{(2Bbde - 3Bae^2 + Abe^2)\log(|bx + a|)}{b^4} - \frac{Bab^2d^2 + Ab^3d^2 - 6Ba^2bde + 2Aab^2de + 5Ba^3e^2 - 3Aa^2be^2 + 2(Bb^3d^2 - 4Bab^2de + 2Ab^3de + 3Ba^2be^2)x}{2(bx + a)^2b^4}$$

input `integrate((B*x+A)*(e*x+d)^2/(b*x+a)^3,x, algorithm="giac")`output $B*e^2*x/b^3 + (2*B*b*d*e - 3*B*a*e^2 + A*b*e^2)*\log(\text{abs}(b*x + a))/b^4 - 1/2*(B*a*b^2*d^2 + A*b^3*d^2 - 6*B*a^2*b*d*e + 2*A*a*b^2*d*e + 5*B*a^3*e^2 - 3*A*a^2*b*e^2 + 2*(B*b^3*d^2 - 4*B*a*b^2*d*e + 2*A*b^3*d*e + 3*B*a^2*b*e^2 - 2*A*a*b^2*e^2)*x)/((b*x + a)^2*b^4)$

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.65

$$\int \frac{(A + Bx)(d + ex)^2}{(a + bx)^3} dx = \frac{\ln(a + bx) (Abe^2 - 3Ba^2e^2 + 2Bbde)}{b^4} - \frac{x(3Ba^2e^2 - 4Babde - 2Aabe^2 + Bb^2d^2 + 2Ab^2de) + \frac{5Ba^3e^2 - 6Ba^2bde - 3Aa^2be^2 + Ba^2b^2d^2 + 2Aab^2d}{2b}}{a^2b^3 + 2ab^4x + b^5x^2} + \frac{Be^2x}{b^3}$$

input `int(((A + B*x)*(d + e*x)^2)/(a + b*x)^3,x)`output $(\log(a + b*x)*(A*b*e^2 - 3*B*a*e^2 + 2*B*b*d*e))/b^4 - (x*(3*B*a^2*e^2 + B*b^2*d^2 - 2*A*a*b*e^2 + 2*A*b^2*d*e - 4*B*a*b*d*e) + (A*b^3*d^2 + 5*B*a^3*e^2 - 3*A*a^2*b*e^2 + B*a*b^2*d^2 + 2*A*a*b^2*d*e - 6*B*a^2*b*d*e)/(2*b)) / (a^2*b^3 + b^5*x^2 + 2*a*b^4*x) + (B*e^2*x)/b^3$ **Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.09

$$\int \frac{(A + Bx)(d + ex)^2}{(a + bx)^3} dx = \frac{-2\log(bx + a) a^3e^2 + 2\log(bx + a) a^2bde - 2\log(bx + a) a^2be^2x + 2\log(bx + a) ab^2dex + 2a^2be^2x - 2a^2b^2d^2}{ab^3(bx + a)}$$

input `int((B*x+A)*(e*x+d)^2/(b*x+a)^3,x)`output $(-2*\log(a + b*x)*a**3*e**2 + 2*\log(a + b*x)*a**2*b*d*e - 2*\log(a + b*x)*a**2*b*e**2*x + 2*\log(a + b*x)*a*b**2*d*e*x + 2*a**2*b*e**2*x - 2*a*b**2*d**2 + a*b**2*e**2*x**2 + b**3*d**2*x)/(a*b**3*(a + b*x))$

3.119 $\int \frac{(A+Bx)(d+ex)}{(a+bx)^3} dx$

Optimal result	1169
Mathematica [A] (verified)	1169
Rubi [A] (verified)	1170
Maple [A] (verified)	1171
Fricas [A] (verification not implemented)	1171
Sympy [A] (verification not implemented)	1172
Maxima [A] (verification not implemented)	1172
Giac [A] (verification not implemented)	1173
Mupad [B] (verification not implemented)	1173
Reduce [B] (verification not implemented)	1173

Optimal result

Integrand size = 18, antiderivative size = 69

$$\int \frac{(A + Bx)(d + ex)}{(a + bx)^3} dx = -\frac{(Ab - aB)(bd - ae)}{2b^3(a + bx)^2} - \frac{bBd + Abe - 2aBe}{b^3(a + bx)} + \frac{Be \log(a + bx)}{b^3}$$

output -1/2*(A*b-B*a)*(-a*e+b*d)/b^3/(b*x+a)^2-(A*b*e-2*B*a*e+B*b*d)/b^3/(b*x+a)+
B*e*ln(b*x+a)/b^3

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.09

$$\int \frac{(A + Bx)(d + ex)}{(a + bx)^3} dx = \frac{-Ab(bd + ae + 2bex) + B(-abd + 3a^2e - 2b^2dx + 4abex) + 2Be(a + bx)^2 \log(a + bx)}{2b^3(a + bx)^2}$$

input Integrate[((A + B*x)*(d + e*x))/(a + b*x)^3,x]

output (-(A*b*(b*d + a*e + 2*b*e*x)) + B*(-(a*b*d) + 3*a^2*e - 2*b^2*d*x + 4*a*b*
e*x) + 2*B*e*(a + b*x)^2*Log[a + b*x])/(2*b^3*(a + b*x)^2)

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)}{(a + bx)^3} dx$$

↓ 86

$$\int \left(\frac{-2aBe + Abe + bBd}{b^2(a + bx)^2} + \frac{(Ab - aB)(bd - ae)}{b^2(a + bx)^3} + \frac{Be}{b^2(a + bx)} \right) dx$$

↓ 2009

$$-\frac{(Ab - aB)(bd - ae)}{2b^3(a + bx)^2} - \frac{-2aBe + Abe + bBd}{b^3(a + bx)} + \frac{Be \log(a + bx)}{b^3}$$

input `Int[((A + B*x)*(d + e*x))/(a + b*x)^3,x]`

output `-1/2*((A*b - a*B)*(b*d - a*e))/(b^3*(a + b*x)^2) - (b*B*d + A*b*e - 2*a*B*e)/(b^3*(a + b*x)) + (B*e*Log[a + b*x])/b^3`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.04

method	result
norman	$\frac{-\frac{Aabe+Ab^2d-3Ba^2e+Babd}{2b^3} - \frac{(Abe-2Bae+Bbd)x}{b^2}}{(bx+a)^2} + \frac{Be \ln(bx+a)}{b^3}$
risch	$\frac{-\frac{Aabe+Ab^2d-3Ba^2e+Babd}{2b^3} - \frac{(Abe-2Bae+Bbd)x}{b^2}}{(bx+a)^2} + \frac{Be \ln(bx+a)}{b^3}$
default	$-\frac{Aabe+Ab^2d+Ba^2e-Babd}{2b^3(bx+a)^2} - \frac{Abe-2Bae+Bbd}{b^3(bx+a)} + \frac{Be \ln(bx+a)}{b^3}$
parallelrisch	$-\frac{-2B \ln(bx+a)x^2b^2e-4B \ln(bx+a)xabe+2Ax b^2e-2B \ln(bx+a)a^2e-4Bxabe+2Bx b^2d+Aabe+Ab^2d-3Ba^2e+Babd}{2b^3(bx+a)^2}$

```
input int((B*x+A)*(e*x+d)/(b*x+a)^3,x,method=_RETURNVERBOSE)
```

```
output (-1/2*(A*a*b*e+A*b^2*d-3*B*a^2*e+B*a*b*d)/b^3-(A*b*e-2*B*a*e+B*b*d)/b^2*x)/(b*x+a)^2+B*e*ln(b*x+a)/b^3
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.59

$$\int \frac{(A + Bx)(d + ex)}{(a + bx)^3} dx = \frac{(Bab + Ab^2)d - (3Ba^2 - Aab)e + 2(Bb^2d - (2Bab - Ab^2)e)x - 2(Bb^2ex^2 + 2Babex + Ba^2e) \log(bx + a)}{2(b^5x^2 + 2ab^4x + a^2b^3)}$$

```
input integrate((B*x+A)*(e*x+d)/(b*x+a)^3,x, algorithm="fricas")
```

```
output -1/2*((B*a*b + A*b^2)*d - (3*B*a^2 - A*a*b)*e + 2*(B*b^2*d - (2*B*a*b - A*b^2)*e)*x - 2*(B*b^2*e*x^2 + 2*B*a*b*e*x + B*a^2*e)*log(b*x + a))/(b^5*x^2 + 2*a*b^4*x + a^2*b^3)
```


Sympy [A] (verification not implemented)

Time = 0.44 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.36

$$\int \frac{(A + Bx)(d + ex)}{(a + bx)^3} dx$$

$$= \frac{Be \log(a + bx)}{b^3} + \frac{-Aabe - Ab^2d + 3Ba^2e - Babd + x(-2Ab^2e + 4Babe - 2Bb^2d)}{2a^2b^3 + 4ab^4x + 2b^5x^2}$$

input `integrate((B*x+A)*(e*x+d)/(b*x+a)**3,x)`output `B*e*log(a + b*x)/b**3 + (-A*a*b*e - A*b**2*d + 3*B*a**2*e - B*a*b*d + x*(-2*A*b**2*e + 4*B*a*b*e - 2*B*b**2*d))/(2*a**2*b**3 + 4*a*b**4*x + 2*b**5*x**2)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.33

$$\int \frac{(A + Bx)(d + ex)}{(a + bx)^3} dx$$

$$= -\frac{(Bab + Ab^2)d - (3Ba^2 - Aab)e + 2(Bb^2d - (2Bab - Ab^2)e)x}{2(b^5x^2 + 2ab^4x + a^2b^3)} + \frac{Be \log(bx + a)}{b^3}$$

input `integrate((B*x+A)*(e*x+d)/(b*x+a)^3,x, algorithm="maxima")`output `-1/2*((B*a*b + A*b^2)*d - (3*B*a^2 - A*a*b)*e + 2*(B*b^2*d - (2*B*a*b - A*b^2)*e)*x)/(b^5*x^2 + 2*a*b^4*x + a^2*b^3) + B*e*log(b*x + a)/b^3`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.04

$$\int \frac{(A + Bx)(d + ex)}{(a + bx)^3} dx = \frac{Be \log(|bx + a|)}{b^3} - \frac{2(Bbd - 2Bae + Abe)x + \frac{Babd + Ab^2d - 3Ba^2e + Aabe}{b}}{2(bx + a)^2 b^2}$$

input `integrate((B*x+A)*(e*x+d)/(b*x+a)^3,x, algorithm="giac")`output `B*e*log(abs(b*x + a))/b^3 - 1/2*(2*(B*b*d - 2*B*a*e + A*b*e)*x + (B*a*b*d + A*b^2*d - 3*B*a^2*e + A*a*b*e)/b)/((b*x + a)^2*b^2)`**Mupad [B] (verification not implemented)**

Time = 1.13 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.19

$$\int \frac{(A + Bx)(d + ex)}{(a + bx)^3} dx = \frac{Be \ln(a + bx)}{b^3} - \frac{\frac{Ab^2d - 3Ba^2e + Aabe + Babd}{2b^3} + \frac{x(Abe - 2Bae + Bbd)}{b^2}}{a^2 + 2abx + b^2x^2}$$

input `int(((A + B*x)*(d + e*x))/(a + b*x)^3,x)`output `(B*e*log(a + b*x))/b^3 - ((A*b^2*d - 3*B*a^2*e + A*a*b*e + B*a*b*d)/(2*b^3) + (x*(A*b*e - 2*B*a*e + B*b*d))/b^2)/(a^2 + b^2*x^2 + 2*a*b*x)`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.71

$$\int \frac{(A + Bx)(d + ex)}{(a + bx)^3} dx = \frac{\log(bx + a) a^2 e + \log(bx + a) abex - abex + b^2 dx}{a b^2 (bx + a)}$$

input `int((B*x+A)*(e*x+d)/(b*x+a)^3,x)`

output $(\log(a + b*x)*a**2*e + \log(a + b*x)*a*b*e*x - a*b**2*x + b**2*d*x)/(a*b**2*(a + b*x))$

3.120 $\int \frac{A+Bx}{(a+bx)^3} dx$

Optimal result	1175
Mathematica [A] (verified)	1175
Rubi [A] (verified)	1176
Maple [A] (verified)	1177
Fricas [A] (verification not implemented)	1177
Sympy [A] (verification not implemented)	1178
Maxima [A] (verification not implemented)	1178
Giac [A] (verification not implemented)	1178
Mupad [B] (verification not implemented)	1179
Reduce [B] (verification not implemented)	1179

Optimal result

Integrand size = 13, antiderivative size = 28

$$\int \frac{A + Bx}{(a + bx)^3} dx = -\frac{(A + Bx)^2}{2(Ab - aB)(a + bx)^2}$$

output `-1/2*(B*x+A)^2/(A*b-B*a)/(b*x+a)^2`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{A + Bx}{(a + bx)^3} dx = -\frac{Ab + B(a + 2bx)}{2b^2(a + bx)^2}$$

input `Integrate[(A + B*x)/(a + b*x)^3,x]`

output `-1/2*(A*b + B*(a + 2*b*x))/(b^2*(a + b*x)^2)`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^3} dx$$

↓ 48

$$-\frac{(A + Bx)^2}{2(a + bx)^2(Ab - aB)}$$

input `Int[(A + B*x)/(a + b*x)^3,x]`

output `-1/2*(A + B*x)^2/((A*b - a*B)*(a + b*x)^2)`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.89

method	result	size
gospers	$-\frac{2bBx+Ab+Ba}{2b^2(bx+a)^2}$	25
parallelrisch	$-\frac{2bBx+Ab+Ba}{2b^2(bx+a)^2}$	25
orering	$-\frac{2bBx+Ab+Ba}{2b^2(bx+a)^2}$	25
norman	$\frac{-\frac{Bx}{b} - \frac{Ab+Ba}{2b^2}}{(bx+a)^2}$	29
risch	$\frac{-\frac{Bx}{b} - \frac{Ab+Ba}{2b^2}}{(bx+a)^2}$	29
default	$-\frac{Ab-Ba}{2b^2(bx+a)^2} - \frac{B}{b^2(bx+a)}$	35

input `int((B*x+A)/(b*x+a)^3,x,method=_RETURNVERBOSE)`output `-1/2*(2*B*b*x+A*b+B*a)/b^2/(b*x+a)^2`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.36

$$\int \frac{A+Bx}{(a+bx)^3} dx = -\frac{2Bbx+Ba+Ab}{2(b^4x^2+2ab^3x+a^2b^2)}$$

input `integrate((B*x+A)/(b*x+a)^3,x, algorithm="fricas")`output `-1/2*(2*B*b*x + B*a + A*b)/(b^4*x^2 + 2*a*b^3*x + a^2*b^2)`

Sympy [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.39

$$\int \frac{A + Bx}{(a + bx)^3} dx = \frac{-Ab - Ba - 2Bbx}{2a^2b^2 + 4ab^3x + 2b^4x^2}$$

input `integrate((B*x+A)/(b*x+a)**3,x)`output `(-A*b - B*a - 2*B*b*x)/(2*a**2*b**2 + 4*a*b**3*x + 2*b**4*x**2)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.36

$$\int \frac{A + Bx}{(a + bx)^3} dx = -\frac{2Bbx + Ba + Ab}{2(b^4x^2 + 2ab^3x + a^2b^2)}$$

input `integrate((B*x+A)/(b*x+a)^3,x, algorithm="maxima")`output `-1/2*(2*B*b*x + B*a + A*b)/(b^4*x^2 + 2*a*b^3*x + a^2*b^2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.86

$$\int \frac{A + Bx}{(a + bx)^3} dx = -\frac{2Bbx + Ba + Ab}{2(bx + a)^2b^2}$$

input `integrate((B*x+A)/(b*x+a)^3,x, algorithm="giac")`output `-1/2*(2*B*b*x + B*a + A*b)/((b*x + a)^2*b^2)`

Mupad [B] (verification not implemented)

Time = 0.98 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.39

$$\int \frac{A + Bx}{(a + bx)^3} dx = -\frac{\frac{Ab+Ba}{2b^2} + \frac{Bx}{b}}{a^2 + 2abx + b^2x^2}$$

input `int((A + B*x)/(a + b*x)^3,x)`

output `-((A*b + B*a)/(2*b^2) + (B*x)/b)/(a^2 + b^2*x^2 + 2*a*b*x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.43

$$\int \frac{A + Bx}{(a + bx)^3} dx = \frac{x}{a(bx + a)}$$

input `int((B*x+A)/(b*x+a)^3,x)`

output `x/(a*(a + b*x))`

3.121 $\int \frac{A+Bx}{(a+bx)^3(d+ex)} dx$

Optimal result	1180
Mathematica [A] (verified)	1180
Rubi [A] (verified)	1181
Maple [A] (verified)	1182
Fricas [B] (verification not implemented)	1183
Sympy [B] (verification not implemented)	1183
Maxima [B] (verification not implemented)	1184
Giac [B] (verification not implemented)	1185
Mupad [B] (verification not implemented)	1185
Reduce [B] (verification not implemented)	1186

Optimal result

Integrand size = 20, antiderivative size = 113

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)} dx = \frac{-Ab + aB}{2b(bd - ae)(a + bx)^2} - \frac{Bd - Ae}{(bd - ae)^2(a + bx)} - \frac{e(Bd - Ae) \log(a + bx)}{(bd - ae)^3} + \frac{e(Bd - Ae) \log(d + ex)}{(bd - ae)^3}$$

output

```
1/2*(-A*b+B*a)/b/(-a*e+b*d)/(b*x+a)^2-(-A*e+B*d)/(-a*e+b*d)^2/(b*x+a)-e*(-A*e+B*d)*ln(b*x+a)/(-a*e+b*d)^3+e*(-A*e+B*d)*ln(e*x+d)/(-a*e+b*d)^3
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.91

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)} dx = \frac{\frac{(-Ab+aB)(bd-ae)^2}{b(a+bx)^2} + \frac{2(bd-ae)(-Bd+Ae)}{a+bx} + 2e(-Bd + Ae) \log(a + bx) + 2e(Bd - Ae) \log(d + ex)}{2(bd - ae)^3}$$

input

```
Integrate[(A + B*x)/((a + b*x)^3*(d + e*x)),x]
```

output

$$\left(\frac{((-A*b) + a*B)*(b*d - a*e)^2}{(b*(a + b*x)^2) + (2*(b*d - a*e)*(-(B*d) + A*e))}{(a + b*x)} + \frac{2*e*(-(B*d) + A*e)*\text{Log}[a + b*x] + 2*e*(B*d - A*e)*\text{Log}[d + e*x]}{2*(b*d - a*e)^3} \right)$$

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)} dx$$

↓ 86

$$\int \left(-\frac{e^2(Ae - Bd)}{(d + ex)(bd - ae)^3} + \frac{be(Ae - Bd)}{(a + bx)(bd - ae)^3} + \frac{b(Bd - Ae)}{(a + bx)^2(bd - ae)^2} + \frac{Ab - aB}{(a + bx)^3(bd - ae)} \right) dx$$

↓ 2009

$$-\frac{Ab - aB}{2b(a + bx)^2(bd - ae)} - \frac{Bd - Ae}{(a + bx)(bd - ae)^2} - \frac{e \log(a + bx)(Bd - Ae)}{(bd - ae)^3} + \frac{e(Bd - Ae) \log(d + ex)}{(bd - ae)^3}$$

input

$$\text{Int}[(A + B*x)/((a + b*x)^3*(d + e*x)), x]$$

output

$$-1/2*(A*b - a*B)/(b*(b*d - a*e)*(a + b*x)^2) - (B*d - A*e)/((b*d - a*e)^2*(a + b*x)) - (e*(B*d - A*e)*\text{Log}[a + b*x])/(b*d - a*e)^3 + (e*(B*d - A*e)*\text{Log}[d + e*x])/(b*d - a*e)^3$$

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.98

method	result
default	$-\frac{-Ab+Ba}{2(ae-db)b(bx+a)^2} + \frac{Ae-Bd}{(ae-db)^2(bx+a)} - \frac{(Ae-Bd)e \ln(bx+a)}{(ae-db)^3} + \frac{(Ae-Bd)e \ln(ex+d)}{(ae-db)^3}$
norman	$\frac{\frac{(Ab^2e-b^2Bd)x}{b(a^2e^2-2abde+b^2d^2)} + \frac{3Aab^2e-Ab^3d-Ba^2be-Bab^2d}{2b^2(a^2e^2-2abde+b^2d^2)}}{(bx+a)^2} + \frac{e(Ae-Bd) \ln(ex+d)}{a^3e^3-3a^2bd e^2+3ab^2d^2e-b^3d^3} - \frac{e(Ae-Bd) \ln(bx+a)}{a^3e^3-3a^2bd e^2+3ab^2d^2e-b^3d^3}$
risch	$\frac{\frac{b(Ae-Bd)x}{a^2e^2-2abde+b^2d^2} + \frac{3Aabe-Ab^2d-Ba^2e-Babd}{2b(a^2e^2-2abde+b^2d^2)}}{(bx+a)^2} - \frac{e^2 \ln(bx+a)A}{a^3e^3-3a^2bd e^2+3ab^2d^2e-b^3d^3} + \frac{e \ln(bx+a)Bd}{a^3e^3-3a^2bd e^2+3ab^2d^2e-b^3d^3} + \dots$
parallelrisc	$-\frac{2Bxa b^3de-2B \ln(bx+a)x^2b^4de+2B \ln(ex+d)x^2b^4de+4A \ln(bx+a)xa b^3e^2-4A \ln(ex+d)xa b^3e^2-2B \ln(bx+a)a^2b^2de+2B \ln(ex+d)a^2b^2de}{(bx+a)^2}$

```
input int((B*x+A)/(b*x+a)^3/(e*x+d),x,method=_RETURNVERBOSE)
```

```
output -1/2*(-A*b+B*a)/(a*e-b*d)/b/(b*x+a)^2+(A*e-B*d)/(a*e-b*d)^2/(b*x+a)-(A*e-B*d)*e/(a*e-b*d)^3*ln(b*x+a)+(A*e-B*d)*e/(a*e-b*d)^3*ln(e*x+d)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 361 vs. $2(111) = 222$.

Time = 0.10 (sec) , antiderivative size = 361, normalized size of antiderivative = 3.19

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)} dx = \frac{4Aab^2de - (Bab^2 + Ab^3)d^2 + (Ba^3 - 3Aa^2b)e^2 - 2(Bb^3d^2 + Aab^2e^2 - (Bab^2 + Ab^3)de)x - 2(Ba^2bde - 2a^2b^4d^3 - 3a^3b^3d^2e + 3a^4b^2de^2 - a^5be^3)}{2(a^2b^4d^3 - 3a^3b^3d^2e + 3a^4b^2de^2 - a^5be^3)}$$

input `integrate((B*x+A)/(b*x+a)^3/(e*x+d),x, algorithm="fricas")`

output

```
1/2*(4*A*a*b^2*d*e - (B*a*b^2 + A*b^3)*d^2 + (B*a^3 - 3*A*a^2*b)*e^2 - 2*(
B*b^3*d^2 + A*a*b^2*e^2 - (B*a*b^2 + A*b^3)*d*e)*x - 2*(B*a^2*b*d*e - A*a^
2*b*e^2 + (B*b^3*d*e - A*b^3*e^2)*x^2 + 2*(B*a*b^2*d*e - A*a*b^2*e^2)*x)*1
og(b*x + a) + 2*(B*a^2*b*d*e - A*a^2*b*e^2 + (B*b^3*d*e - A*b^3*e^2)*x^2 +
2*(B*a*b^2*d*e - A*a*b^2*e^2)*x)*log(e*x + d))/(a^2*b^4*d^3 - 3*a^3*b^3*d
^2*e + 3*a^4*b^2*d*e^2 - a^5*b*e^3 + (b^6*d^3 - 3*a*b^5*d^2*e + 3*a^2*b^4*
d*e^2 - a^3*b^3*e^3)*x^2 + 2*(a*b^5*d^3 - 3*a^2*b^4*d^2*e + 3*a^3*b^3*d*e^
2 - a^4*b^2*e^3)*x)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 558 vs. $2(92) = 184$.

Time = 1.45 (sec) , antiderivative size = 558, normalized size of antiderivative = 4.94

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)} dx = \frac{e(-Ae + Bd) \log \left(x + \frac{-Aae^3 - Abde^2 + Bade^2 + Bbd^2e - \frac{a^4e^5(-Ae+Bd)}{(ae-bd)^3} + \frac{4a^3bde^4(-Ae+Bd)}{(ae-bd)^3} - \frac{6a^2b^2d^2e^3(-Ae+Bd)}{(ae-bd)^3} + \frac{4ab^3d^3e^2(-Ae+Bd)}{(ae-bd)^3}}{-2Abe^3 + 2Bbde^2} \right)}{(ae - bd)^3} + \frac{e(-Ae + Bd) \log \left(x + \frac{-Aae^3 - Abde^2 + Bade^2 + Bbd^2e + \frac{a^4e^5(-Ae+Bd)}{(ae-bd)^3} - \frac{4a^3bde^4(-Ae+Bd)}{(ae-bd)^3} + \frac{6a^2b^2d^2e^3(-Ae+Bd)}{(ae-bd)^3} - \frac{4ab^3d^3e^2(-Ae+Bd)}{(ae-bd)^3}}{-2Abe^3 + 2Bbde^2} \right)}{(ae - bd)^3} + \frac{3Aabe - Ab^2d - Ba^2e - Babd + x(2Ab^2e - 2Bb^2d)}{2a^4be^2 - 4a^3b^2de + 2a^2b^3d^2 + x^2 \cdot (2a^2b^3e^2 - 4ab^4de + 2b^5d^2) + x(4a^3b^2e^2 - 8a^2b^3de + 4ab^4d^2)}$$

input `integrate((B*x+A)/(b*x+a)**3/(e*x+d),x)`

output
$$\begin{aligned} & -e*(-Ae + Bd)*\log(x + (-Aa^3e - Ab^2de + B^2ade + B^2bd^2e \\ & - a^4e^5*(-Ae + Bd)/(ae - bd)^3 + 4a^3b^2de^4*(-Ae + Bd)/(ae - bd)^3 - 6a^2b^2d^2e^3*(-Ae + Bd)/(ae - bd)^3 + 4ab^3d^3e^2*(-Ae + Bd)/(ae - bd)^3 - b^4d^4e*(-Ae + Bd)/(ae - bd)^3)/(-2Ab^2e^3 + 2B^2bde^2)/(ae - bd)^3 + e*(-Ae + Bd)*\log(x + (-Aa^3e - Ab^2de + B^2ade + B^2bd^2e + a^4e^5*(-Ae + Bd)/(ae - bd)^3 - 4a^3b^2de^4*(-Ae + Bd)/(ae - bd)^3 + 6a^2b^2d^2e^3*(-Ae + Bd)/(ae - bd)^3 - 4ab^3d^3e^2*(-Ae + Bd)/(ae - bd)^3 + b^4d^4e*(-Ae + Bd)/(ae - bd)^3)/(-2Ab^2e^3 + 2B^2bde^2)/(ae - bd)^3 + (3A^2ab^2e - Ab^2d - B^2ade - B^2abd + x(2Ab^2e - 2B^2bd))/(2a^4b^2e^2 - 4a^3b^2d^2e + 2a^2b^3d^2 + x^2(2a^2b^3e^2 - 4ab^4d^2e + 2b^5d^2) + x(4a^3b^2e^2 - 8a^2b^3d^2e + 4ab^4d^2)) \end{aligned}$$

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 252 vs. $2(111) = 222$.

Time = 0.04 (sec) , antiderivative size = 252, normalized size of antiderivative = 2.23

$$\begin{aligned} & \int \frac{A + Bx}{(a + bx)^3(d + ex)} dx \\ & = -\frac{(Bde - Ae^2) \log(bx + a)}{b^3d^3 - 3ab^2d^2e + 3a^2bde^2 - a^3e^3} + \frac{(Bde - Ae^2) \log(ex + d)}{b^3d^3 - 3ab^2d^2e + 3a^2bde^2 - a^3e^3} \\ & \quad - \frac{(Bab + Ab^2)d + (Ba^2 - 3Aab)e + 2(Bb^2d - Ab^2e)x}{2(a^2b^3d^2 - 2a^3b^2de + a^4be^2 + (b^5d^2 - 2ab^4de + a^2b^3e^2)x^2 + 2(ab^4d^2 - 2a^2b^3de + a^3b^2e^2)x)} \end{aligned}$$

input `integrate((B*x+A)/(b*x+a)^3/(e*x+d),x, algorithm="maxima")`

output
$$\begin{aligned} & -(Bd^2e - Ae^2)*\log(b*x + a)/(b^3*d^3 - 3*a*b^2*d^2*e + 3*a^2*b*d*e^2 - a \\ & ^3*e^3) + (Bd^2e - Ae^2)*\log(e*x + d)/(b^3*d^3 - 3*a*b^2*d^2*e + 3*a^2*b* \\ & d^2*e^2 - a^3*e^3) - 1/2*((B*a*b + A*b^2)*d + (B*a^2 - 3*A*a*b)*e + 2*(B*b^2 \\ & *d - A*b^2*e)*x)/(a^2*b^3*d^2 - 2*a^3*b^2*d^2*e + a^4*b*e^2 + (b^5*d^2 - 2*a \\ & *b^4*d^2*e + a^2*b^3*e^2)*x^2 + 2*(a*b^4*d^2 - 2*a^2*b^3*d^2*e + a^3*b^2*e^2)* \\ & x) \end{aligned}$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 232 vs. 2(111) = 222.

Time = 0.12 (sec) , antiderivative size = 232, normalized size of antiderivative = 2.05

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)} dx$$

$$= -\frac{(Bbde - Abe^2) \log(|bx + a|)}{b^4d^3 - 3ab^3d^2e + 3a^2b^2de^2 - a^3be^3} + \frac{(Bde^2 - Ae^3) \log(|ex + d|)}{b^3d^3e - 3ab^2d^2e^2 + 3a^2bde^3 - a^3e^4}$$

$$-\frac{Bab^2d^2 + Ab^3d^2 - 4Aab^2de - Ba^3e^2 + 3Aa^2be^2 + 2(Bb^3d^2 - Bab^2de - Ab^3de + Aab^2e^2)x}{2(bd - ae)^3(bx + a)^2b}$$

input `integrate((B*x+A)/(b*x+a)^3/(e*x+d),x, algorithm="giac")`

output

```
-(B*b*d*e - A*b*e^2)*log(abs(b*x + a))/(b^4*d^3 - 3*a*b^3*d^2*e + 3*a^2*b^2*d*e^2 - a^3*b*e^3) + (B*d*e^2 - A*e^3)*log(abs(e*x + d))/(b^3*d^3*e - 3*a*b^2*d^2*e^2 + 3*a^2*b*d*e^3 - a^3*e^4) - 1/2*(B*a*b^2*d^2 + A*b^3*d^2 - 4*A*a*b^2*d*e - B*a^3*e^2 + 3*A*a^2*b*e^2 + 2*(B*b^3*d^2 - B*a*b^2*d*e - A*b^3*d*e + A*a*b^2*e^2)*x)/((b*d - a*e)^3*(b*x + a)^2*b)
```

Mupad [B] (verification not implemented)

Time = 1.08 (sec) , antiderivative size = 228, normalized size of antiderivative = 2.02

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)} dx$$

$$= -\frac{\frac{Ab^2d + Ba^2e - 3Aabe + Babd}{2b(a^2e^2 - 2abde + b^2d^2)} - \frac{bx(Ae - Bd)}{a^2e^2 - 2abde + b^2d^2}}{a^2 + 2abx + b^2x^2}$$

$$-\frac{2e \operatorname{atanh}\left(\frac{\left(\frac{a^3e^3 - a^2bde^2 - ab^2d^2e + b^3d^3}{a^2e^2 - 2abde + b^2d^2} + 2bex\right)(a^2e^2 - 2abde + b^2d^2)}{(ae - bd)^3}\right)}{(ae - bd)^3} (Ae - Bd)$$

input `int((A + B*x)/((a + b*x)^3*(d + e*x)),x)`

output

```
- ((A*b^2*d + B*a^2*e - 3*A*a*b*e + B*a*b*d)/(2*b*(a^2*e^2 + b^2*d^2 - 2*a*b*d*e)) - (b*x*(A*e - B*d))/(a^2*e^2 + b^2*d^2 - 2*a*b*d*e))/(a^2 + b^2*x^2 + 2*a*b*x) - (2*e*atanh((((a^3*e^3 + b^3*d^3 - a*b^2*d^2*e - a^2*b*d*e^2)/(a^2*e^2 + b^2*d^2 - 2*a*b*d*e) + 2*b*e*x)*(a^2*e^2 + b^2*d^2 - 2*a*b*d*e))/(a*e - b*d)^3)*(A*e - B*d))/(a*e - b*d)^3
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.02

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)} dx$$

$$= \frac{-\log(bx + a) a^2 e - \log(bx + a) abex + \log(ex + d) a^2 e + \log(ex + d) abex - abex + b^2 dx}{a(a^2 b e^2 x - 2 a b^2 d e x + b^3 d^2 x + a^3 e^2 - 2 a^2 b d e + a b^2 d^2)}$$

input

```
int((B*x+A)/(b*x+a)^3/(e*x+d),x)
```

output

```
( - log(a + b*x)*a**2*e - log(a + b*x)*a*b*e*x + log(d + e*x)*a**2*e + log(d + e*x)*a*b*e*x - a*b*e*x + b**2*d*x)/(a*(a**3*e**2 - 2*a**2*b*d*e + a**2*b*e**2*x + a*b**2*d**2 - 2*a*b**2*d*e*x + b**3*d**2*x))
```

3.122 $\int \frac{A+Bx}{(a+bx)^3(d+ex)^2} dx$

Optimal result	1187
Mathematica [A] (verified)	1187
Rubi [A] (verified)	1188
Maple [A] (verified)	1189
Fricas [B] (verification not implemented)	1190
Sympy [B] (verification not implemented)	1190
Maxima [B] (verification not implemented)	1191
Giac [A] (verification not implemented)	1192
Mupad [B] (verification not implemented)	1193
Reduce [B] (verification not implemented)	1193

Optimal result

Integrand size = 20, antiderivative size = 158

$$\int \frac{A+Bx}{(a+bx)^3(d+ex)^2} dx = -\frac{Ab-aB}{2(bd-ae)^2(a+bx)^2} - \frac{bBd-2Abe+aBe}{(bd-ae)^3(a+bx)}$$

$$-\frac{e(Bd-Ae)}{(bd-ae)^3(d+ex)} - \frac{e(2bBd-3Abe+aBe)\log(a+bx)}{(bd-ae)^4}$$

$$+\frac{e(2bBd-3Abe+aBe)\log(d+ex)}{(bd-ae)^4}$$

output

```
-1/2*(A*b-B*a)/(-a*e+b*d)^2/(b*x+a)^2-(-2*A*b*e+B*a*e+B*b*d)/(-a*e+b*d)^3/
(b*x+a)-e*(-A*e+B*d)/(-a*e+b*d)^3/(e*x+d)-e*(-3*A*b*e+B*a*e+2*B*b*d)*ln(b*
x+a)/(-a*e+b*d)^4+e*(-3*A*b*e+B*a*e+2*B*b*d)*ln(e*x+d)/(-a*e+b*d)^4
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.92

$$\int \frac{A+Bx}{(a+bx)^3(d+ex)^2} dx$$

$$= \frac{\frac{(-Ab+aB)(bd-ae)^2}{(a+bx)^2} - \frac{2(bd-ae)(bBd-2Abe+aBe)}{a+bx} + \frac{2e(bd-ae)(-Bd+ Ae)}{d+ex} - 2e(2bBd-3Abe+aBe)\log(a+bx) + 2e}{2(bd-ae)^4}$$

input `Integrate[(A + B*x)/((a + b*x)^3*(d + e*x)^2),x]`

output `((((-A*b) + a*B)*(b*d - a*e)^2)/(a + b*x)^2 - (2*(b*d - a*e)*(b*B*d - 2*A*b*e + a*B*e))/(a + b*x) + (2*e*(b*d - a*e)*(-(B*d) + A*e))/(d + e*x) - 2*e*(2*b*B*d - 3*A*b*e + a*B*e)*Log[a + b*x] + 2*e*(2*b*B*d - 3*A*b*e + a*B*e)*Log[d + e*x])/(2*(b*d - a*e)^4)`

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^2} dx$$

↓ 86

$$\int \left(-\frac{e^2(-aBe + 3Abe - 2bBd)}{(d + ex)(bd - ae)^4} - \frac{e^2(Ae - Bd)}{(d + ex)^2(bd - ae)^3} + \frac{be(-aBe + 3Abe - 2bBd)}{(a + bx)(bd - ae)^4} + \frac{b(aBe - 2Abe + bBd)}{(a + bx)^2(bd - ae)^3} \right) dx$$

↓ 2009

$$\frac{-\frac{Ab - aB}{2(a + bx)^2(bd - ae)^2} - \frac{aBe - 2Abe + bBd}{(a + bx)(bd - ae)^3} - \frac{e(Bd - Ae)}{(d + ex)(bd - ae)^3} - \frac{e \log(a + bx)(aBe - 3Abe + 2bBd)}{(bd - ae)^4}}{(bd - ae)^4} + \frac{e \log(d + ex)(aBe - 3Abe + 2bBd)}{(bd - ae)^4}$$

input `Int[(A + B*x)/((a + b*x)^3*(d + e*x)^2),x]`

output `-1/2*(A*b - a*B)/((b*d - a*e)^2*(a + b*x)^2) - (b*B*d - 2*A*b*e + a*B*e)/((b*d - a*e)^3*(a + b*x)) - (e*(B*d - A*e))/((b*d - a*e)^3*(d + e*x)) - (e*(2*b*B*d - 3*A*b*e + a*B*e)*Log[a + b*x])/(b*d - a*e)^4 + (e*(2*b*B*d - 3*A*b*e + a*B*e)*Log[d + e*x])/(b*d - a*e)^4`

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.02

method	result
default	$-\frac{2Abe - Bae - Bbd}{(ae - db)^3(bx + a)} - \frac{Ab - Ba}{2(ae - db)^2(bx + a)^2} + \frac{e(3Abe - Bae - 2Bbd) \ln(bx + a)}{(ae - db)^4} - \frac{(Ae - Bd)e}{(ae - db)^3(ex + d)} - \frac{e(3Abe - Bae - 2Bbd)}{(ae - db)^4}$
norman	$-\frac{2Aa^2b^2e^3 + 5Aab^3de^2 - Ab^4d^2e - 5Ba^2b^2de^2 - Bab^3d^2e}{2eb^2(a^3e^3 - 3a^2bde^2 + 3ab^2d^2e - b^3d^3)} - \frac{(3Ab^3e^3 - Bab^2e^3 - 2b^3Bde^2)x^2}{eb(a^3e^3 - 3a^2bde^2 + 3ab^2d^2e - b^3d^3)} - \frac{(9Aab^3e^3 + 3Aab^4de^2 - 3Ba^2b^2e^3 - 3Aab^3d^2e^2)}{2eb^2(a^3e^3 - 3a^2bde^2 + 3ab^2d^2e - b^3d^3)}$
risch	$-\frac{be(3Abe - Bae - 2Bbd)x^2}{a^3e^3 - 3a^2bde^2 + 3ab^2d^2e - b^3d^3} - \frac{(3ae + db)(3Abe - Bae - 2Bbd)x}{2(a^3e^3 - 3a^2bde^2 + 3ab^2d^2e - b^3d^3)} - \frac{2a^2Ae^2 + 5Aabde - Ab^2d^2 - 5Ba^2de - Babd^2}{2(a^3e^3 - 3a^2bde^2 + 3ab^2d^2e - b^3d^3)}$
parallelrisc	$\frac{2B \ln(ex + d) x a^3 b^2 e^4 + 2B \ln(ex + d) x^3 a b^4 e^4 - 6A \ln(ex + d) x a^2 b^3 e^4 - 2B \ln(bx + a) x a^3 b^2 e^4 - 4B \ln(bx + a) x^3 b^5 d e^3 - 12A \ln(e)}{(bx + a)^2(ex + d)} - \frac{1}{a^4 e^4 - 4a^3}$

```
input int((B*x+A)/(b*x+a)^3/(e*x+d)^2,x,method=_RETURNVERBOSE)
```

```
output -(2*A*b*e-B*a*e-B*b*d)/(a*e-b*d)^3/(b*x+a)-1/2*(A*b-B*a)/(a*e-b*d)^2/(b*x+a)^2+e*(3*A*b*e-B*a*e-2*B*b*d)/(a*e-b*d)^4*ln(b*x+a)-(A*e-B*d)*e/(a*e-b*d)^3/(e*x+d)-e*(3*A*b*e-B*a*e-2*B*b*d)/(a*e-b*d)^4*ln(e*x+d)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 803 vs. $2(156) = 312$.

Time = 0.10 (sec) , antiderivative size = 803, normalized size of antiderivative = 5.08

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^2} dx = \text{Too large to display}$$

input `integrate((B*x+A)/(b*x+a)^3/(e*x+d)^2,x, algorithm="fricas")`

output

```
-1/2*(2*A*a^3*e^3 + (B*a*b^2 + A*b^3)*d^3 + 2*(2*B*a^2*b - 3*A*a*b^2)*d^2*
e - (5*B*a^3 - 3*A*a^2*b)*d*e^2 + 2*(2*B*b^3*d^2*e - (B*a*b^2 + 3*A*b^3)*d
*e^2 - (B*a^2*b - 3*A*a*b^2)*e^3)*x^2 + (2*B*b^3*d^3 + (5*B*a*b^2 - 3*A*b^
3)*d^2*e - 2*(2*B*a^2*b + 3*A*a*b^2)*d*e^2 - 3*(B*a^3 - 3*A*a^2*b)*e^3)*x
+ 2*(2*B*a^2*b*d^2*e + (B*a^3 - 3*A*a^2*b)*d*e^2 + (2*B*b^3*d*e^2 + (B*a*b
^2 - 3*A*b^3)*e^3)*x^3 + (2*B*b^3*d^2*e + (5*B*a*b^2 - 3*A*b^3)*d*e^2 + 2*
(B*a^2*b - 3*A*a*b^2)*e^3)*x^2 + (4*B*a*b^2*d^2*e + 2*(2*B*a^2*b - 3*A*a*b
^2)*d*e^2 + (B*a^3 - 3*A*a^2*b)*e^3)*x)*log(b*x + a) - 2*(2*B*a^2*b*d^2*e
+ (B*a^3 - 3*A*a^2*b)*d*e^2 + (2*B*b^3*d*e^2 + (B*a*b^2 - 3*A*b^3)*e^3)*x^
3 + (2*B*b^3*d^2*e + (5*B*a*b^2 - 3*A*b^3)*d*e^2 + 2*(B*a^2*b - 3*A*a*b^2)
*e^3)*x^2 + (4*B*a*b^2*d^2*e + 2*(2*B*a^2*b - 3*A*a*b^2)*d*e^2 + (B*a^3 -
3*A*a^2*b)*e^3)*x)*log(e*x + d)/(a^2*b^4*d^5 - 4*a^3*b^3*d^4*e + 6*a^4*b^
2*d^3*e^2 - 4*a^5*b*d^2*e^3 + a^6*d*e^4 + (b^6*d^4*e - 4*a*b^5*d^3*e^2 + 6
*a^2*b^4*d^2*e^3 - 4*a^3*b^3*d*e^4 + a^4*b^2*e^5)*x^3 + (b^6*d^5 - 2*a*b^5
*d^4*e - 2*a^2*b^4*d^3*e^2 + 8*a^3*b^3*d^2*e^3 - 7*a^4*b^2*d*e^4 + 2*a^5*b
*e^5)*x^2 + (2*a*b^5*d^5 - 7*a^2*b^4*d^4*e + 8*a^3*b^3*d^3*e^2 - 2*a^4*b^2
*d^2*e^3 - 2*a^5*b*d*e^4 + a^6*e^5)*x)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1066 vs. $2(148) = 296$.

Time = 1.99 (sec) , antiderivative size = 1066, normalized size of antiderivative = 6.75

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^2} dx = \text{Too large to display}$$

input `integrate((B*x+A)/(b*x+a)**3/(e*x+d)**2,x)`

output

```
e*(-3*A*b*e + B*a*e + 2*B*b*d)*log(x + (-3*A*a*b*e**3 - 3*A*b**2*d*e**2 +
B*a**2*e**3 + 3*B*a*b*d*e**2 + 2*B*b**2*d**2*e - a**5*e**6*(-3*A*b*e + B*a
*e + 2*B*b*d)/(a*e - b*d)**4 + 5*a**4*b*d*e**5*(-3*A*b*e + B*a*e + 2*B*b*d
)/(a*e - b*d)**4 - 10*a**3*b**2*d**2*e**4*(-3*A*b*e + B*a*e + 2*B*b*d)/(a*
e - b*d)**4 + 10*a**2*b**3*d**3*e**3*(-3*A*b*e + B*a*e + 2*B*b*d)/(a*e - b
*d)**4 - 5*a*b**4*d**4*e**2*(-3*A*b*e + B*a*e + 2*B*b*d)/(a*e - b*d)**4 +
b**5*d**5*e*(-3*A*b*e + B*a*e + 2*B*b*d)/(a*e - b*d)**4)/(-6*A*b**2*e**3 +
2*B*a*b*e**3 + 4*B*b**2*d*e**2))/(a*e - b*d)**4 - e*(-3*A*b*e + B*a*e + 2
*B*b*d)*log(x + (-3*A*a*b*e**3 - 3*A*b**2*d*e**2 + B*a**2*e**3 + 3*B*a*b*d
*e**2 + 2*B*b**2*d**2*e + a**5*e**6*(-3*A*b*e + B*a*e + 2*B*b*d)/(a*e - b*
d)**4 - 5*a**4*b*d*e**5*(-3*A*b*e + B*a*e + 2*B*b*d)/(a*e - b*d)**4 + 10*a
**3*b**2*d**2*e**4*(-3*A*b*e + B*a*e + 2*B*b*d)/(a*e - b*d)**4 - 10*a**2*b
**3*d**3*e**3*(-3*A*b*e + B*a*e + 2*B*b*d)/(a*e - b*d)**4 + 5*a*b**4*d**4*
e**2*(-3*A*b*e + B*a*e + 2*B*b*d)/(a*e - b*d)**4 - b**5*d**5*e*(-3*A*b*e +
B*a*e + 2*B*b*d)/(a*e - b*d)**4)/(-6*A*b**2*e**3 + 2*B*a*b*e**3 + 4*B*b**
2*d*e**2))/(a*e - b*d)**4 + (-2*A*a**2*e**2 - 5*A*a*b*d*e + A*b**2*d**2 +
5*B*a**2*d*e + B*a*b*d**2 + x**2*(-6*A*b**2*e**2 + 2*B*a*b*e**2 + 4*B*b**2
*d*e) + x*(-9*A*a*b*e**2 - 3*A*b**2*d*e + 3*B*a**2*e**2 + 7*B*a*b*d*e + 2*
B*b**2*d**2))/(2*a**5*d*e**3 - 6*a**4*b*d**2*e**2 + 6*a**3*b**2*d**3*e - 2
*a**2*b**3*d**4 + x**3*(2*a**3*b**2*e**4 - 6*a**2*b**3*d*e**3 + 6*a*b**...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 477 vs. $2(156) = 312$.

Time = 0.05 (sec) , antiderivative size = 477, normalized size of antiderivative = 3.02

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^2} dx = -\frac{(2Bbde + (Ba - 3Ab)e^2) \log(bx + a)}{b^4d^4 - 4ab^3d^3e + 6a^2b^2d^2e^2 - 4a^3bde^3 + a^4e^4}$$

$$+ \frac{(2Bbde + (Ba - 3Ab)e^2) \log(ex + d)}{b^4d^4 - 4ab^3d^3e + 6a^2b^2d^2e^2 - 4a^3bde^3 + a^4e^4}$$

$$+ \frac{2Aa^2e^2 - (Bab + Ab^2)d^2 - 5(Ba^2 - Aab)de - 2(2Bb^2de + (Bab - 3Aab^2)d^2)}{2(a^2b^3d^4 - 3a^3b^2d^3e + 3a^4bd^2e^2 - a^5de^3 + (b^5d^3e - 3ab^4d^2e^2 + 3a^2b^3de^3 - a^3b^2e^4)x^3 + (b^5d^4 - ab^4d^3e - 3a^2b^3d^2e^2 + 3ab^2d^3e^2 - a^4d^4e^3)x^2 + (2a^3b^2d^3e^2 - 2a^4bd^2e^2 - 2a^5de^3 + 2a^2b^3d^2e^2 - 2a^3b^2d^3e^2)x + (2a^4bd^2e^2 - 2a^5de^3 + 2a^2b^3d^2e^2 - 2a^3b^2d^3e^2)}$$

input

```
integrate((B*x+A)/(b*x+a)^3/(e*x+d)^2,x, algorithm="maxima")
```

output

```

-(2*B*b*d*e + (B*a - 3*A*b)*e^2)*log(b*x + a)/(b^4*d^4 - 4*a*b^3*d^3*e + 6
*a^2*b^2*d^2*e^2 - 4*a^3*b*d*e^3 + a^4*e^4) + (2*B*b*d*e + (B*a - 3*A*b)*e
^2)*log(e*x + d)/(b^4*d^4 - 4*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2 - 4*a^3*b*d*
e^3 + a^4*e^4) + 1/2*(2*A*a^2*e^2 - (B*a*b + A*b^2)*d^2 - 5*(B*a^2 - A*a*b
)*d*e - 2*(2*B*b^2*d*e + (B*a*b - 3*A*b^2)*e^2)*x^2 - (2*B*b^2*d^2 + (7*B*
a*b - 3*A*b^2)*d*e + 3*(B*a^2 - 3*A*a*b)*e^2)*x)/(a^2*b^3*d^4 - 3*a^3*b^2*
d^3*e + 3*a^4*b*d^2*e^2 - a^5*d*e^3 + (b^5*d^3*e - 3*a*b^4*d^2*e^2 + 3*a^2
*b^3*d*e^3 - a^3*b^2*e^4)*x^3 + (b^5*d^4 - a*b^4*d^3*e - 3*a^2*b^3*d^2*e^2
+ 5*a^3*b^2*d*e^3 - 2*a^4*b*e^4)*x^2 + (2*a*b^4*d^4 - 5*a^2*b^3*d^3*e + 3
*a^3*b^2*d^2*e^2 + a^4*b*d*e^3 - a^5*e^4)*x)

```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.93

$$\begin{aligned}
& \int \frac{A + Bx}{(a + bx)^3(d + ex)^2} dx \\
&= -\frac{(2Bbde^2 + Bae^3 - 3Abe^3) \log\left(\left|b - \frac{bd}{ex+d} + \frac{ae}{ex+d}\right|\right)}{b^4d^4e - 4ab^3d^3e^2 + 6a^2b^2d^2e^3 - 4a^3bde^4 + a^4e^5} \\
&\quad - \frac{\frac{Bde^4}{ex+d} - \frac{Ae^5}{ex+d}}{b^3d^3e^3 - 3ab^2d^2e^4 + 3a^2bde^5 - a^3e^6} \\
&\quad - \frac{2Bb^3de + 3Bab^2e^2 - 5Ab^3e^2 - \frac{2(Bb^3d^2e^2 + Bab^2de^3 - 3Ab^3de^3 - 2Ba^2be^4 + 3Aab^2e^4)}{(ex+d)e}}{2(bd - ae)^4 \left(b - \frac{bd}{ex+d} + \frac{ae}{ex+d}\right)^2}
\end{aligned}$$

input

```
integrate((B*x+A)/(b*x+a)^3/(e*x+d)^2,x, algorithm="giac")
```

output

```

-(2*B*b*d*e^2 + B*a*e^3 - 3*A*b*e^3)*log(abs(b - b*d/(e*x + d) + a*e/(e*x
+ d)))/(b^4*d^4*e - 4*a*b^3*d^3*e^2 + 6*a^2*b^2*d^2*e^3 - 4*a^3*b*d*e^4 +
a^4*e^5) - (B*d*e^4/(e*x + d) - A*e^5/(e*x + d))/(b^3*d^3*e^3 - 3*a*b^2*d^
2*e^4 + 3*a^2*b*d*e^5 - a^3*e^6) - 1/2*(2*B*b^3*d*e + 3*B*a*b^2*e^2 - 5*A*
b^3*e^2 - 2*(B*b^3*d^2*e^2 + B*a*b^2*d*e^3 - 3*A*b^3*d*e^3 - 2*B*a^2*b*e^4
+ 3*A*a*b^2*e^4)/((e*x + d)*e))/((b*d - a*e)^4*(b - b*d/(e*x + d) + a*e/(
e*x + d))^2)

```


output

```
(2*log(a + b*x)*a**2*b*d*e**2 + 2*log(a + b*x)*a**2*b*e**3*x + 2*log(a + b
*x)*a*b**2*d**2*e + 4*log(a + b*x)*a*b**2*d*e**2*x + 2*log(a + b*x)*a*b**2
*e**3*x**2 + 2*log(a + b*x)*b**3*d**2*e*x + 2*log(a + b*x)*b**3*d*e**2*x**
2 - 2*log(d + e*x)*a**2*b*d*e**2 - 2*log(d + e*x)*a**2*b*e**3*x - 2*log(d
+ e*x)*a*b**2*d**2*e - 4*log(d + e*x)*a*b**2*d*e**2*x - 2*log(d + e*x)*a*b
**2*e**3*x**2 - 2*log(d + e*x)*b**3*d**2*e*x - 2*log(d + e*x)*b**3*d*e**2*
x**2 - a**3*e**3 + a**2*b*d*e**2 - a*b**2*d**2*e + 2*a*b**2*e**3*x**2 + b
**3*d**3 - 2*b**3*d*e**2*x**2)/(a**5*d*e**4 + a**5*e**5*x - 2*a**4*b*d**2*e
**3 - a**4*b*d*e**4*x + a**4*b*e**5*x**2 - 2*a**3*b**2*d**2*e**3*x - 2*a**
3*b**2*d*e**4*x**2 + 2*a**2*b**3*d**4*e + 2*a**2*b**3*d**3*e**2*x - a*b**4
*d**5 + a*b**4*d**4*e*x + 2*a*b**4*d**3*e**2*x**2 - b**5*d**5*x - b**5*d**
4*e*x**2)
```

3.123 $\int \frac{A+Bx}{(a+bx)^3(d+ex)^3} dx$

Optimal result	1195
Mathematica [A] (verified)	1196
Rubi [A] (verified)	1196
Maple [A] (verified)	1198
Fricas [B] (verification not implemented)	1198
Sympy [B] (verification not implemented)	1199
Maxima [B] (verification not implemented)	1200
Giac [B] (verification not implemented)	1201
Mupad [B] (verification not implemented)	1202
Reduce [B] (verification not implemented)	1203

Optimal result

Integrand size = 20, antiderivative size = 199

$$\int \frac{A+Bx}{(a+bx)^3(d+ex)^3} dx = -\frac{b(Ab-aB)}{2(bd-ae)^3(a+bx)^2} - \frac{b(bBd-3Abe+2aBe)}{(bd-ae)^4(a+bx)}$$

$$-\frac{e(Bd-Ae)}{2(bd-ae)^3(d+ex)^2} - \frac{e(2bBd-3Abe+aBe)}{(bd-ae)^4(d+ex)}$$

$$-\frac{3be(bBd-2Abe+aBe)\log(a+bx)}{(bd-ae)^5}$$

$$+\frac{3be(bBd-2Abe+aBe)\log(d+ex)}{(bd-ae)^5}$$

output

```
-1/2*b*(A*b-B*a)/(-a*e+b*d)^3/(b*x+a)^2-b*(-3*A*b*e+2*B*a*e+B*b*d)/(-a*e+b
*d)^4/(b*x+a)-1/2*e*(-A*e+B*d)/(-a*e+b*d)^3/(e*x+d)^2-e*(-3*A*b*e+B*a*e+2*
B*b*d)/(-a*e+b*d)^4/(e*x+d)-3*b*e*(-2*A*b*e+B*a*e+B*b*d)*ln(b*x+a)/(-a*e+b
*d)^5+3*b*e*(-2*A*b*e+B*a*e+B*b*d)*ln(e*x+d)/(-a*e+b*d)^5
```


Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 185, normalized size of antiderivative = 0.93

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^3} dx$$

$$= \frac{-\frac{b(Ab - aB)(bd - ae)^2}{(a + bx)^2} - \frac{2b(bd - ae)(bBd - 3Abe + 2aBe)}{a + bx} + \frac{e(bd - ae)^2(-Bd + Ae)}{(d + ex)^2} + \frac{2e(bd - ae)(-2bBd + 3Abe - aBe)}{d + ex} - 6be(bBd - 2aBe)}{2(bd - ae)^5}$$

input

```
Integrate[(A + B*x)/((a + b*x)^3*(d + e*x)^3), x]
```

output

```
(-((b*(A*b - a*B)*(b*d - a*e)^2)/(a + b*x)^2) - (2*b*(b*d - a*e)*(b*B*d - 3*A*b*e + 2*a*B*e))/(a + b*x) + (e*(b*d - a*e)^2*(-(B*d) + A*e))/(d + e*x)^2 + (2*e*(b*d - a*e)*(-2*b*B*d + 3*A*b*e - a*B*e))/(d + e*x) - 6*b*e*(b*B*d - 2*A*b*e + a*B*e)*Log[a + b*x] + 6*b*e*(b*B*d - 2*A*b*e + a*B*e)*Log[d + e*x])/(2*(b*d - a*e)^5)
```

Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^3} dx$$

$$\downarrow 86$$

$$\int \left(\frac{3b^2e(-aBe + 2Abe - bBd)}{(a + bx)(bd - ae)^5} + \frac{b^2(2aBe - 3Abe + bBd)}{(a + bx)^2(bd - ae)^4} + \frac{b^2(Ab - aB)}{(a + bx)^3(bd - ae)^3} - \frac{3be^2(-aBe + 2Abe - bBd)}{(d + ex)(bd - ae)^5} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned} & -\frac{b(Ab - aB)}{2(a + bx)^2(bd - ae)^3} - \frac{b(2aBe - 3Abe + bBd)}{(a + bx)(bd - ae)^4} - \frac{e(aBe - 3Abe + 2bBd)}{(d + ex)(bd - ae)^4} - \\ & \frac{e(Bd - Ae)}{2(d + ex)^2(bd - ae)^3} - \frac{3be \log(a + bx)(aBe - 2Abe + bBd)}{(bd - ae)^5} + \\ & \frac{3be \log(d + ex)(aBe - 2Abe + bBd)}{(bd - ae)^5} \end{aligned}$$

input `Int[(A + B*x)/((a + b*x)^3*(d + e*x)^3),x]`

output `-1/2*(b*(A*b - a*B))/((b*d - a*e)^3*(a + b*x)^2) - (b*(b*B*d - 3*A*b*e + 2*a*B*e))/((b*d - a*e)^4*(a + b*x)) - (e*(B*d - A*e))/(2*(b*d - a*e)^3*(d + e*x)^2) - (e*(2*b*B*d - 3*A*b*e + a*B*e))/((b*d - a*e)^4*(d + e*x)) - (3*b*e*(b*B*d - 2*A*b*e + a*B*e)*Log[a + b*x])/(b*d - a*e)^5 + (3*b*e*(b*B*d - 2*A*b*e + a*B*e)*Log[d + e*x])/(b*d - a*e)^5`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.01

method	result
default	$\frac{b(3Abe-2Bae-Bbd)}{(ae-db)^4(bx+a)} + \frac{(Ab-Ba)b}{2(ae-db)^3(bx+a)^2} - \frac{3be(2Abe-Bae-Bbd)\ln(bx+a)}{(ae-db)^5} - \frac{(Ae-Bd)e}{2(ae-db)^3(ex+d)^2} + \frac{e(3Abe-Bae-Bbd)}{(ae-db)^4(ex+d)}$
norman	$\frac{(6Ab^4e^4-3Ba^3b^3e^4-3Bb^4de^3)x^3}{(a^4e^4-4a^3bde^3+6a^2b^2d^2e^2-4ab^3d^3e+b^4d^4)be} + \frac{(2Aa^2b^3e^5+14Aab^4de^4+2Ab^5d^2e^3-Ba^3b^2e^5-8Ba^2b^3de^4-8Bab^4d^2e^3-Bb^5d^3e^2)}{e^2b^2(a^4e^4-4a^3bde^3+6a^2b^2d^2e^2-4ab^3d^3e+b^4d^4)}$
risch	$\frac{3b^2e^2(2Abe-Bae-Bbd)x^3}{a^4e^4-4a^3bde^3+6a^2b^2d^2e^2-4ab^3d^3e+b^4d^4} + \frac{9be(ae+db)(2Abe-Bae-Bbd)x^2}{2(a^4e^4-4a^3bde^3+6a^2b^2d^2e^2-4ab^3d^3e+b^4d^4)} + \frac{(2Aa^2be^3+14Aab^2de^2+2Ab^3d^2e-Ba^3b^2e^5-8Ba^2b^3de^4-8Bab^4d^2e^3-Bb^5d^3e^2)}{a^4e^4-4a^3bde^3+6a^2b^2d^2e^2-4ab^3d^3e+b^4d^4} + \frac{e(3Abe-Bae-Bbd)}{(ex+d)^2(bx+a)^2}$
parallelrisch	Expression too large to display

```
input int((B*x+A)/(b*x+a)^3/(e*x+d)^3,x,method=_RETURNVERBOSE)
```

```
output b*(3*A*b*e-2*B*a*e-B*b*d)/(a*e-b*d)^4/(b*x+a)+1/2*(A*b-B*a)*b/(a*e-b*d)^3/(b*x+a)^2-3*b*e*(2*A*b*e-B*a*e-B*b*d)/(a*e-b*d)^5*ln(b*x+a)-1/2*(A*e-B*d)*e/(a*e-b*d)^3/(e*x+d)^2+e*(3*A*b*e-B*a*e-2*B*b*d)/(a*e-b*d)^4/(e*x+d)+3*b*e*(2*A*b*e-B*a*e-B*b*d)/(a*e-b*d)^5*ln(e*x+d)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1215 vs. 2(195) = 390.

Time = 0.14 (sec) , antiderivative size = 1215, normalized size of antiderivative = 6.11

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^3} dx = \text{Too large to display}$$

```
input integrate((B*x+A)/(b*x+a)^3/(e*x+d)^3,x, algorithm="fricas")
```

output

```

1/2*(9*B*a^3*b*d^2*e^2 + A*a^4*e^4 - (B*a*b^3 + A*b^4)*d^4 - (9*B*a^2*b^2
- 8*A*a*b^3)*d^3*e + (B*a^4 - 8*A*a^3*b)*d*e^3 - 6*(B*b^4*d^2*e^2 - 2*A*b^
4*d*e^3 - (B*a^2*b^2 - 2*A*a*b^3)*e^4)*x^3 - 9*(B*b^4*d^3*e - B*a^2*b^2*d*
e^3 + (B*a*b^3 - 2*A*b^4)*d^2*e^2 - (B*a^3*b - 2*A*a^2*b^2)*e^4)*x^2 - 2*(
B*b^4*d^4 - 12*A*a*b^3*d^2*e^2 + (7*B*a*b^3 - 2*A*b^4)*d^3*e - (7*B*a^3*b
- 12*A*a^2*b^2)*d*e^3 - (B*a^4 - 2*A*a^3*b)*e^4)*x - 6*(B*a^2*b^2*d^3*e +
(B*a^3*b - 2*A*a^2*b^2)*d^2*e^2 + (B*b^4*d*e^3 + (B*a*b^3 - 2*A*b^4)*e^4)*
x^4 + 2*(B*b^4*d^2*e^2 + 2*(B*a*b^3 - A*b^4)*d*e^3 + (B*a^2*b^2 - 2*A*a*b^
3)*e^4)*x^3 + (B*b^4*d^3*e + (5*B*a*b^3 - 2*A*b^4)*d^2*e^2 + (5*B*a^2*b^2
- 8*A*a*b^3)*d*e^3 + (B*a^3*b - 2*A*a^2*b^2)*e^4)*x^2 + 2*(B*a*b^3*d^3*e +
2*(B*a^2*b^2 - A*a*b^3)*d^2*e^2 + (B*a^3*b - 2*A*a^2*b^2)*d*e^3)*x*log(b
*x + a) + 6*(B*a^2*b^2*d^3*e + (B*a^3*b - 2*A*a^2*b^2)*d^2*e^2 + (B*b^4*d*
e^3 + (B*a*b^3 - 2*A*b^4)*e^4)*x^4 + 2*(B*b^4*d^2*e^2 + 2*(B*a*b^3 - A*b^4
)*d*e^3 + (B*a^2*b^2 - 2*A*a*b^3)*e^4)*x^3 + (B*b^4*d^3*e + (5*B*a*b^3 - 2
*A*b^4)*d^2*e^2 + (5*B*a^2*b^2 - 8*A*a*b^3)*d*e^3 + (B*a^3*b - 2*A*a^2*b^2
)*e^4)*x^2 + 2*(B*a*b^3*d^3*e + 2*(B*a^2*b^2 - A*a*b^3)*d^2*e^2 + (B*a^3*b
- 2*A*a^2*b^2)*d*e^3)*x*log(e*x + d))/(a^2*b^5*d^7 - 5*a^3*b^4*d^6*e + 1
0*a^4*b^3*d^5*e^2 - 10*a^5*b^2*d^4*e^3 + 5*a^6*b*d^3*e^4 - a^7*d^2*e^5 + (
b^7*d^5*e^2 - 5*a*b^6*d^4*e^3 + 10*a^2*b^5*d^3*e^4 - 10*a^3*b^4*d^2*e^5 +
5*a^4*b^3*d*e^6 - a^5*b^2*e^7)*x^4 + 2*(b^7*d^6*e - 4*a*b^6*d^5*e^2 + 5...

```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1431 vs. $2(192) = 384$.

Time = 2.88 (sec) , antiderivative size = 1431, normalized size of antiderivative = 7.19

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^3} dx = \text{Too large to display}$$

input

```
integrate((B*x+A)/(b*x+a)**3/(e*x+d)**3,x)
```

output

```

-3*b*e*(-2*A*b*e + B*a*e + B*b*d)*log(x + (-6*A*a*b**2*e**3 - 6*A*b**3*d*e
**2 + 3*B*a**2*b*e**3 + 6*B*a*b**2*d*e**2 + 3*B*b**3*d**2*e - 3*a**6*b*e**
7*(-2*A*b*e + B*a*e + B*b*d)/(a*e - b*d)**5 + 18*a**5*b**2*d*e**6*(-2*A*b*
e + B*a*e + B*b*d)/(a*e - b*d)**5 - 45*a**4*b**3*d**2*e**5*(-2*A*b*e + B*a
*e + B*b*d)/(a*e - b*d)**5 + 60*a**3*b**4*d**3*e**4*(-2*A*b*e + B*a*e + B*
b*d)/(a*e - b*d)**5 - 45*a**2*b**5*d**4*e**3*(-2*A*b*e + B*a*e + B*b*d)/(a
*e - b*d)**5 + 18*a*b**6*d**5*e**2*(-2*A*b*e + B*a*e + B*b*d)/(a*e - b*d)*
*5 - 3*b**7*d**6*e*(-2*A*b*e + B*a*e + B*b*d)/(a*e - b*d)**5)/(-12*A*b**3*
e**3 + 6*B*a*b**2*e**3 + 6*B*b**3*d*e**2))/(a*e - b*d)**5 + 3*b*e*(-2*A*b*
e + B*a*e + B*b*d)*log(x + (-6*A*a*b**2*e**3 - 6*A*b**3*d*e**2 + 3*B*a**2*
b*e**3 + 6*B*a*b**2*d*e**2 + 3*B*b**3*d**2*e + 3*a**6*b*e**7*(-2*A*b*e + B
*a*e + B*b*d)/(a*e - b*d)**5 - 18*a**5*b**2*d*e**6*(-2*A*b*e + B*a*e + B*b
*d)/(a*e - b*d)**5 + 45*a**4*b**3*d**2*e**5*(-2*A*b*e + B*a*e + B*b*d)/(a*
e - b*d)**5 - 60*a**3*b**4*d**3*e**4*(-2*A*b*e + B*a*e + B*b*d)/(a*e - b*d
)**5 + 45*a**2*b**5*d**4*e**3*(-2*A*b*e + B*a*e + B*b*d)/(a*e - b*d)**5 -
18*a*b**6*d**5*e**2*(-2*A*b*e + B*a*e + B*b*d)/(a*e - b*d)**5 + 3*b**7*d**
6*e*(-2*A*b*e + B*a*e + B*b*d)/(a*e - b*d)**5)/(-12*A*b**3*e**3 + 6*B*a*b*
*2*e**3 + 6*B*b**3*d*e**2))/(a*e - b*d)**5 + (-A*a**3*e**3 + 7*A*a**2*b*d*
e**2 + 7*A*a*b**2*d**2*e - A*b**3*d**3 - B*a**3*d*e**2 - 10*B*a**2*b*d**2*
e - B*a*b**2*d**3 + x**3*(12*A*b**3*e**3 - 6*B*a*b**2*e**3 - 6*B*b**3*d...

```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 745 vs. $2(195) = 390$.

Time = 0.05 (sec) , antiderivative size = 745, normalized size of antiderivative = 3.74

$$\begin{aligned}
& \int \frac{A + Bx}{(a + bx)^3(d + ex)^3} dx \\
&= -\frac{3(Bb^2de + (Bab - 2Ab^2)e^2) \log(bx + a)}{b^5d^5 - 5ab^4d^4e + 10a^2b^3d^3e^2 - 10a^3b^2d^2e^3 + 5a^4bde^4 - a^5e^5} \\
&+ \frac{3(Bb^2de + (Bab - 2Ab^2)e^2) \log(ex + d)}{b^5d^5 - 5ab^4d^4e + 10a^2b^3d^3e^2 - 10a^3b^2d^2e^3 + 5a^4bde^4 - a^5e^5} \\
&- \frac{Aa^3e^3 + (Bab^2 + Ab^3)d^3 + (10Ba^2b - 7Aab^2)d^2e + (Ba^3 - 7Aa^2b)d + A^2}{2(a^2b^4d^6 - 4a^3b^3d^5e + 6a^4b^2d^4e^2 - 4a^5bd^3e^3 + a^6d^2e^4 + (b^6d^4e^2 - 4ab^5d^3e^3 + 6a^2b^4d^2e^4 - 4a^3b^3de^5 + 6a^4b^2d^2e^5 - 6a^5bd^2e^4 + 3a^6d^2e^5) \log(bx + a) + 6a^5bd^3e^3 - 6a^6d^2e^4 + 3a^7d^2e^5) \log(ex + d) - 3a^6d^2e^5}
\end{aligned}$$

input

```
integrate((B*x+A)/(b*x+a)^3/(e*x+d)^3,x, algorithm="maxima")
```

output

```

-3*(B*b^2*d*e + (B*a*b - 2*A*b^2)*e^2)*log(b*x + a)/(b^5*d^5 - 5*a*b^4*d^4
*e + 10*a^2*b^3*d^3*e^2 - 10*a^3*b^2*d^2*e^3 + 5*a^4*b*d*e^4 - a^5*e^5) +
3*(B*b^2*d*e + (B*a*b - 2*A*b^2)*e^2)*log(e*x + d)/(b^5*d^5 - 5*a*b^4*d^4*
e + 10*a^2*b^3*d^3*e^2 - 10*a^3*b^2*d^2*e^3 + 5*a^4*b*d*e^4 - a^5*e^5) - 1
/2*(A*a^3*e^3 + (B*a*b^2 + A*b^3)*d^3 + (10*B*a^2*b - 7*A*a*b^2)*d^2*e + (
B*a^3 - 7*A*a^2*b)*d*e^2 + 6*(B*b^3*d*e^2 + (B*a*b^2 - 2*A*b^3)*e^3)*x^3 +
9*(B*b^3*d^2*e + 2*(B*a*b^2 - A*b^3)*d*e^2 + (B*a^2*b - 2*A*a*b^2)*e^3)*x
^2 + 2*(B*b^3*d^3 + 2*(4*B*a*b^2 - A*b^3)*d^2*e + 2*(4*B*a^2*b - 7*A*a*b^2
)*d*e^2 + (B*a^3 - 2*A*a^2*b)*e^3)*x)/(a^2*b^4*d^6 - 4*a^3*b^3*d^5*e + 6*a
^4*b^2*d^4*e^2 - 4*a^5*b*d^3*e^3 + a^6*d^2*e^4 + (b^6*d^4*e^2 - 4*a*b^5*d^
3*e^3 + 6*a^2*b^4*d^2*e^4 - 4*a^3*b^3*d*e^5 + a^4*b^2*e^6)*x^4 + 2*(b^6*d^
5*e - 3*a*b^5*d^4*e^2 + 2*a^2*b^4*d^3*e^3 + 2*a^3*b^3*d^2*e^4 - 3*a^4*b^2*
d*e^5 + a^5*b*e^6)*x^3 + (b^6*d^6 - 9*a^2*b^4*d^4*e^2 + 16*a^3*b^3*d^3*e^3
- 9*a^4*b^2*d^2*e^4 + a^6*e^6)*x^2 + 2*(a*b^5*d^6 - 3*a^2*b^4*d^5*e + 2*a
^3*b^3*d^4*e^2 + 2*a^4*b^2*d^3*e^3 - 3*a^5*b*d^2*e^4 + a^6*d*e^5)*x)

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 532 vs. $2(195) = 390$.

Time = 0.13 (sec) , antiderivative size = 532, normalized size of antiderivative = 2.67

$$\begin{aligned}
& \int \frac{A + Bx}{(a + bx)^3(d + ex)^3} dx \\
&= -\frac{3(Bb^3de + Bab^2e^2 - 2Ab^3e^2) \log(|bx + a|)}{b^6d^5 - 5ab^5d^4e + 10a^2b^4d^3e^2 - 10a^3b^3d^2e^3 + 5a^4b^2de^4 - a^5be^5} \\
&+ \frac{3(Bb^2de^2 + Babe^3 - 2Ab^2e^3) \log(|ex + d|)}{b^5d^5e - 5ab^4d^4e^2 + 10a^2b^3d^3e^3 - 10a^3b^2d^2e^4 + 5a^4bde^5 - a^5e^6} \\
&- \frac{6Bb^3de^2x^3 + 6Bab^2e^3x^3 - 12Ab^3e^3x^3 + 9Bb^3d^2ex^2 + 18Bab^2de^2x^2 - 18Ab^3de^2x^2 + 9Ba^2be^3x^2 - \dots}{\dots}
\end{aligned}$$

input

```
integrate((B*x+A)/(b*x+a)^3/(e*x+d)^3,x, algorithm="giac")
```

output

```

-3*(B*b^3*d*e + B*a*b^2*e^2 - 2*A*b^3*e^2)*log(abs(b*x + a))/(b^6*d^5 - 5*
a*b^5*d^4*e + 10*a^2*b^4*d^3*e^2 - 10*a^3*b^3*d^2*e^3 + 5*a^4*b^2*d*e^4 -
a^5*b*e^5) + 3*(B*b^2*d*e^2 + B*a*b*e^3 - 2*A*b^2*e^3)*log(abs(e*x + d))/(
b^5*d^5*e - 5*a*b^4*d^4*e^2 + 10*a^2*b^3*d^3*e^3 - 10*a^3*b^2*d^2*e^4 + 5*
a^4*b*d*e^5 - a^5*e^6) - 1/2*(6*B*b^3*d*e^2*x^3 + 6*B*a*b^2*e^3*x^3 - 12*A
*b^3*e^3*x^3 + 9*B*b^3*d^2*e*x^2 + 18*B*a*b^2*d*e^2*x^2 - 18*A*b^3*d*e^2*x
^2 + 9*B*a^2*b*e^3*x^2 - 18*A*a*b^2*e^3*x^2 + 2*B*b^3*d^3*x + 16*B*a*b^2*d
^2*e*x - 4*A*b^3*d^2*e*x + 16*B*a^2*b*d*e^2*x - 28*A*a*b^2*d*e^2*x + 2*B*a
^3*e^3*x - 4*A*a^2*b*e^3*x + B*a*b^2*d^3 + A*b^3*d^3 + 10*B*a^2*b*d^2*e -
7*A*a*b^2*d^2*e + B*a^3*d*e^2 - 7*A*a^2*b*d*e^2 + A*a^3*e^3)/((b^4*d^4 - 4
*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2 - 4*a^3*b*d*e^3 + a^4*e^4)*(b*e*x^2 + b*d
*x + a*e*x + a*d)^2)

```

Mupad [B] (verification not implemented)

Time = 1.36 (sec) , antiderivative size = 726, normalized size of antiderivative = 3.65

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^3} dx$$

$$= \frac{2 \operatorname{atanh}\left(\frac{(3be^2(2Ab - Ba) - 3Bb^2de) \left(\frac{a^5e^5 - 3a^4bde^4 + 2a^3b^2d^2e^3 + 2a^2b^3d^3e^2 - 3ab^4d^4e + b^5d^5}{a^4e^4 - 4a^3bde^3 + 6a^2b^2d^2e^2 - 4ab^3d^3e + b^4d^4}\right) + 2be^2}{(ae - bd)^5(-6Ab^2e^2 + 3Bdb^2e + 3Babe^2)}\right)}{(ae - bd)^5}$$

$$- \frac{\frac{Ba^3de^2 + Aa^3e^3 + 10Ba^2bd^2e - 7Aa^2bde^2 + Bab^2d^3 - 7Aab^2d^2e + Ab^3d^3}{2(a^4e^4 - 4a^3bde^3 + 6a^2b^2d^2e^2 - 4ab^3d^3e + b^4d^4)} + \frac{9x^2(db^2e + abe^2)(Bae - 2Abe + Bbd)}{2(a^4e^4 - 4a^3bde^3 + 6a^2b^2d^2e^2 - 4ab^3d^3e + b^4d^4)}}{x(2ea^2d + 2bad^2) + x^2(a^2e^2 + 4abde + b^2d^2) + x^3(2}$$

input

```
int((A + B*x)/((a + b*x)^3*(d + e*x)^3),x)
```

output

```
(2*atanh(((3*b*e^2*(2*A*b - B*a) - 3*B*b^2*d*e)*(a^5*e^5 + b^5*d^5 + 2*a^2*b^3*d^3*e^2 + 2*a^3*b^2*d^2*e^3 - 3*a*b^4*d^4*e - 3*a^4*b*d*e^4)/(a^4*e^4 + b^4*d^4 + 6*a^2*b^2*d^2*e^2 - 4*a*b^3*d^3*e - 4*a^3*b*d*e^3) + 2*b*e*x)*(a^4*e^4 + b^4*d^4 + 6*a^2*b^2*d^2*e^2 - 4*a*b^3*d^3*e - 4*a^3*b*d*e^3)))/((a*e - b*d)^5*(3*B*a*b*e^2 - 6*A*b^2*e^2 + 3*B*b^2*d*e)))*(3*b*e^2*(2*A*b - B*a) - 3*B*b^2*d*e))/(a*e - b*d)^5 - ((A*a^3*e^3 + A*b^3*d^3 + B*a*b^2*d^3 + B*a^3*d*e^2 - 7*A*a*b^2*d^2*e - 7*A*a^2*b*d*e^2 + 10*B*a^2*b*d^2*e)/(2*(a^4*e^4 + b^4*d^4 + 6*a^2*b^2*d^2*e^2 - 4*a*b^3*d^3*e - 4*a^3*b*d*e^3)) + (9*x^2*(a*b*e^2 + b^2*d*e)*(B*a*e - 2*A*b*e + B*b*d))/(2*(a^4*e^4 + b^4*d^4 + 6*a^2*b^2*d^2*e^2 - 4*a*b^3*d^3*e - 4*a^3*b*d*e^3)) + (x*(a^2*e^2 + b^2*d^2 + 7*a*b*d*e)*(B*a*e - 2*A*b*e + B*b*d))/(a^4*e^4 + b^4*d^4 + 6*a^2*b^2*d^2*e^2 - 4*a*b^3*d^3*e - 4*a^3*b*d*e^3) + (3*b^2*e^2*x^3*(B*a*e - 2*A*b*e + B*b*d))/(a^4*e^4 + b^4*d^4 + 6*a^2*b^2*d^2*e^2 - 4*a*b^3*d^3*e - 4*a^3*b*d*e^3))/(x*(2*a*b*d^2 + 2*a^2*d*e) + x^2*(a^2*e^2 + b^2*d^2 + 4*a*b*d*e) + x^3*(2*a*b*e^2 + 2*b^2*d*e) + a^2*d^2 + b^2*e^2*x^4)
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 821, normalized size of antiderivative = 4.13

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^3} dx = \text{Too large to display}$$

input

```
int((B*x+A)/(b*x+a)^3/(e*x+d)^3,x)
```


output

```
( - 6*log(a + b*x)*a**2*b**2*d**2*e**2 - 12*log(a + b*x)*a**2*b**2*d*e**3*x
- 6*log(a + b*x)*a**2*b**2*e**4*x**2 - 12*log(a + b*x)*a*b**3*d**3*e - 3
0*log(a + b*x)*a*b**3*d**2*e**2*x - 24*log(a + b*x)*a*b**3*d*e**3*x**2 - 6
*log(a + b*x)*a*b**3*e**4*x**3 - 12*log(a + b*x)*b**4*d**3*e*x - 24*log(a
+ b*x)*b**4*d**2*e**2*x**2 - 12*log(a + b*x)*b**4*d*e**3*x**3 + 6*log(d +
e*x)*a**2*b**2*d**2*e**2 + 12*log(d + e*x)*a**2*b**2*d*e**3*x + 6*log(d +
e*x)*a**2*b**2*e**4*x**2 + 12*log(d + e*x)*a*b**3*d**3*e + 30*log(d + e*x)
*a*b**3*d**2*e**2*x + 24*log(d + e*x)*a*b**3*d*e**3*x**2 + 6*log(d + e*x)*
a*b**3*e**4*x**3 + 12*log(d + e*x)*b**4*d**3*e*x + 24*log(d + e*x)*b**4*d*
**2*e**2*x**2 + 12*log(d + e*x)*b**4*d*e**3*x**3 - a**4*e**4 + 4*a**3*b*d*e
**3 + 3*a**3*b*e**4*x + 3*a**2*b**2*d**2*e**2 - 2*a*b**3*d**3*e + 9*a*b**3
*d**2*e**2*x - 6*a*b**3*e**4*x**3 - 4*b**4*d**4 - 12*b**4*d**3*e*x + 6*b**
4*d*e**3*x**3)/(2*(a**6*d**2*e**5 + 2*a**6*d*e**6*x + a**6*e**7*x**2 - 2*a
**5*b*d**3*e**4 - 3*a**5*b*d**2*e**5*x + a**5*b*e**7*x**3 - 2*a**4*b**2*d
**4*e**3 - 6*a**4*b**2*d**3*e**4*x - 6*a**4*b**2*d**2*e**5*x**2 - 2*a**4*b*
**2*d*e**6*x**3 + 8*a**3*b**3*d**5*e**2 + 14*a**3*b**3*d**4*e**3*x + 4*a**3
*b**3*d**3*e**4*x**2 - 2*a**3*b**3*d**2*e**5*x**3 - 7*a**2*b**4*d**6*e - 6
*a**2*b**4*d**5*e**2*x + 9*a**2*b**4*d**4*e**3*x**2 + 8*a**2*b**4*d**3*e**
4*x**3 + 2*a*b**5*d**7 - 3*a*b**5*d**6*e*x - 12*a*b**5*d**5*e**2*x**2 - 7*
a*b**5*d**4*e**3*x**3 + 2*b**6*d**7*x + 4*b**6*d**6*e*x**2 + 2*b**6*d**...
```

3.124 $\int (a + bx)(A + Bx)(d + ex)^{5/2} dx$

Optimal result	1205
Mathematica [A] (verified)	1205
Rubi [A] (verified)	1206
Maple [A] (verified)	1207
Fricas [B] (verification not implemented)	1208
Sympy [B] (verification not implemented)	1208
Maxima [A] (verification not implemented)	1209
Giac [B] (verification not implemented)	1209
Mupad [B] (verification not implemented)	1210
Reduce [B] (verification not implemented)	1211

Optimal result

Integrand size = 20, antiderivative size = 83

$$\int (a + bx)(A + Bx)(d + ex)^{5/2} dx = \frac{2(bd - ae)(Bd - Ae)(d + ex)^{7/2}}{7e^3} - \frac{2(2bBd - Abe - aBe)(d + ex)^{9/2}}{9e^3} + \frac{2bB(d + ex)^{11/2}}{11e^3}$$

output

```
2/7*(-a*e+b*d)*(-A*e+B*d)*(e*x+d)^(7/2)/e^3-2/9*(-A*b*e-B*a*e+2*B*b*d)*(e*x+d)^(9/2)/e^3+2/11*b*B*(e*x+d)^(11/2)/e^3
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.84

$$\int (a + bx)(A + Bx)(d + ex)^{5/2} dx = \frac{2(d + ex)^{7/2} (11Abe(-2d + 7ex) + 11ae(-2Bd + 9Ae + 7Bex) + bB(8d^2 - 28dex + 63e^2x^2))}{693e^3}$$

input

```
Integrate[(a + b*x)*(A + B*x)*(d + e*x)^(5/2), x]
```

output

$$(2*(d + e*x)^(7/2)*(11*A*b*e*(-2*d + 7*e*x) + 11*a*e*(-2*B*d + 9*A*e + 7*B*e*x) + b*B*(8*d^2 - 28*d*e*x + 63*e^2*x^2)))/(693*e^3)$$

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)(A + Bx)(d + ex)^{5/2} dx$$

$$\downarrow 86$$

$$\int \left(\frac{(d + ex)^{7/2}(aBe + Abe - 2bBd)}{e^2} + \frac{(d + ex)^{5/2}(ae - bd)(Ae - Bd)}{e^2} + \frac{bB(d + ex)^{9/2}}{e^2} \right) dx$$

$$\downarrow 2009$$

$$-\frac{2(d + ex)^{9/2}(-aBe - Abe + 2bBd)}{9e^3} + \frac{2(d + ex)^{7/2}(bd - ae)(Bd - Ae)}{7e^3} + \frac{2bB(d + ex)^{11/2}}{11e^3}$$

input

$$\text{Int}[(a + b*x)*(A + B*x)*(d + e*x)^(5/2), x]$$

output

$$(2*(b*d - a*e)*(B*d - A*e)*(d + e*x)^(7/2))/(7*e^3) - (2*(2*b*B*d - A*b*e - a*B*e)*(d + e*x)^(9/2))/(9*e^3) + (2*b*B*(d + e*x)^(11/2))/(11*e^3)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 189 vs. $2(71) = 142$.

Time = 0.11 (sec) , antiderivative size = 189, normalized size of antiderivative = 2.28

$$\int (a + bx)(A + Bx)(d + ex)^{5/2} dx = \frac{2(63 Bbe^5 x^5 + 8 Bbd^5 + 99 Aad^3 e^2 - 22(Ba + Ab)d^4 e + 7(23 Bbde^4 + 11(Ba + Ab)e^5)x^4 + \dots}{e^3}$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)^(5/2),x, algorithm="fricas")`

output
$$\frac{2}{693} \cdot (63 B b e^5 x^5 + 8 B b d^5 + 99 A a d^3 e^2 - 22 (B a + A b) d^4 e + 7 (23 B b d e^4 + 11 (B a + A b) e^5) x^4 + (113 B b d^2 e^3 + 99 A a e^5 + 209 (B a + A b) d e^4) x^3 + 3 (B b d^3 e^2 + 99 A a d e^4 + 55 (B a + A b) d^2 e^3) x^2 - (4 B b d^4 e - 297 A a d^2 e^3 - 11 (B a + A b) d^3 e^2) x) \cdot \sqrt{e x + d} / e^3$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 476 vs. $2(87) = 174$.

Time = 0.48 (sec) , antiderivative size = 476, normalized size of antiderivative = 5.73

$$\int (a + bx)(A + Bx)(d + ex)^{5/2} dx = \begin{cases} \frac{2Aad^3\sqrt{d+ex}}{7e} + \frac{6Aad^2x\sqrt{d+ex}}{7} + \frac{6Aadex^2\sqrt{d+ex}}{7} + \frac{2Aae^2x^3\sqrt{d+ex}}{7} - \frac{4Abd^4\sqrt{d+ex}}{63e^2} + \frac{2Abd^3x\sqrt{d+ex}}{63e} + \frac{10Abdx^2\sqrt{d+ex}}{63} \\ d^{\frac{5}{2}} \left(Aax + \frac{Abx^2}{2} + \frac{Bax^2}{2} + \frac{Bbx^3}{3} \right) \end{cases}$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)**(5/2),x)`

output

```
Piecewise((2*A*a*d**3*sqrt(d + e*x)/(7*e) + 6*A*a*d**2*x*sqrt(d + e*x)/7 +
6*A*a*d*e*x**2*sqrt(d + e*x)/7 + 2*A*a*e**2*x**3*sqrt(d + e*x)/7 - 4*A*b*
d**4*sqrt(d + e*x)/(63*e**2) + 2*A*b*d**3*x*sqrt(d + e*x)/(63*e) + 10*A*b*
d**2*x**2*sqrt(d + e*x)/21 + 38*A*b*d*e*x**3*sqrt(d + e*x)/63 + 2*A*b*e**2
*x**4*sqrt(d + e*x)/9 - 4*B*a*d**4*sqrt(d + e*x)/(63*e**2) + 2*B*a*d**3*x*
sqrt(d + e*x)/(63*e) + 10*B*a*d**2*x**2*sqrt(d + e*x)/21 + 38*B*a*d*e*x**3
*sqrt(d + e*x)/63 + 2*B*a*e**2*x**4*sqrt(d + e*x)/9 + 16*B*b*d**5*sqrt(d +
e*x)/(693*e**3) - 8*B*b*d**4*x*sqrt(d + e*x)/(693*e**2) + 2*B*b*d**3*x**2
*sqrt(d + e*x)/(231*e) + 226*B*b*d**2*x**3*sqrt(d + e*x)/693 + 46*B*b*d*e*
x**4*sqrt(d + e*x)/99 + 2*B*b*e**2*x**5*sqrt(d + e*x)/11, Ne(e, 0)), (d**(
5/2)*(A*a*x + A*b*x**2/2 + B*a*x**2/2 + B*b*x**3/3), True))
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.90

$$\int (a + bx)(A + Bx)(d + ex)^{5/2} dx = \frac{2 \left(63 (ex + d)^{\frac{11}{2}} Bb - 77 (2 Bbd - (Ba + Ab)e)(ex + d)^{\frac{9}{2}} + 99 (Bbd^2 + Aae^2 - (Ba + Ab)de) \right)}{693 e^3}$$

input

```
integrate((b*x+a)*(B*x+A)*(e*x+d)^(5/2),x, algorithm="maxima")
```

output

```
2/693*(63*(e*x + d)^(11/2)*B*b - 77*(2*B*b*d - (B*a + A*b)*e)*(e*x + d)^(9
/2) + 99*(B*b*d^2 + A*a*e^2 - (B*a + A*b)*d*e)*(e*x + d)^(7/2))/e^3
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 735 vs. $2(71) = 142$.

Time = 0.13 (sec) , antiderivative size = 735, normalized size of antiderivative = 8.86

$$\int (a + bx)(A + Bx)(d + ex)^{5/2} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)*(B*x+A)*(e*x+d)^(5/2),x, algorithm="giac")
```

output

```

2/3465*(3465*sqrt(e*x + d)*A*a*d^3 + 3465*((e*x + d)^(3/2) - 3*sqrt(e*x +
d)*d)*A*a*d^2 + 1155*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*B*a*d^3/e + 115
5*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*A*b*d^3/e + 693*(3*(e*x + d)^(5/2)
- 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*A*a*d + 231*(3*(e*x + d)^(
5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*B*b*d^3/e^2 + 693*(3*(
e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*B*a*d^2/e +
693*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*A*b*
d^2/e + 99*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*
d^2 - 35*sqrt(e*x + d)*d^3)*A*a + 297*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5
/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*B*b*d^2/e^2 + 297*(
5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqr
t(e*x + d)*d^3)*B*a*d/e + 297*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d +
35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*A*b*d/e + 33*(35*(e*x + d)^(
9/2) - 180*(e*x + d)^(7/2)*d + 378*(e*x + d)^(5/2)*d^2 - 420*(e*x + d)^(3
/2)*d^3 + 315*sqrt(e*x + d)*d^4)*B*b*d/e^2 + 11*(35*(e*x + d)^(9/2) - 180*
(e*x + d)^(7/2)*d + 378*(e*x + d)^(5/2)*d^2 - 420*(e*x + d)^(3/2)*d^3 + 31
5*sqrt(e*x + d)*d^4)*B*a/e + 11*(35*(e*x + d)^(9/2) - 180*(e*x + d)^(7/2)*
d + 378*(e*x + d)^(5/2)*d^2 - 420*(e*x + d)^(3/2)*d^3 + 315*sqrt(e*x + d)*
d^4)*A*b/e + 5*(63*(e*x + d)^(11/2) - 385*(e*x + d)^(9/2)*d + 990*(e*x + d
)^(7/2)*d^2 - 1386*(e*x + d)^(5/2)*d^3 + 1155*(e*x + d)^(3/2)*d^4 - 693...

```

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.96

$$\int (a + bx)(A + Bx)(d + ex)^{5/2} dx = \frac{2(d + ex)^{7/2} (63 B b (d + ex)^2 + 99 A a e^2 + 99 B b d^2 + 77 A b e (d + ex) + 77 B a e (d + ex))}{693 e^3}$$

input

```
int((A + B*x)*(a + b*x)*(d + e*x)^(5/2),x)
```

output

```

(2*(d + e*x)^(7/2)*(63*B*b*(d + e*x)^2 + 99*A*a*e^2 + 99*B*b*d^2 + 77*A*b*
e*(d + e*x) + 77*B*a*e*(d + e*x) - 154*B*b*d*(d + e*x) - 99*A*b*d*e - 99*B
*a*d*e))/(693*e^3)

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 180, normalized size of antiderivative = 2.17

$$\int (a + bx)(A + Bx)(d + ex)^{5/2} dx = \frac{2\sqrt{ex + d}(63b^2e^5x^5 + 154abe^5x^4 + 161b^2de^4x^4 + 99a^2e^5x^3 + 418abd e^4x^3 + 113b^2d^2e^3x^3 + 2$$

input `int((b*x+a)*(B*x+A)*(e*x+d)^(5/2),x)`output `(2*sqrt(d + e*x)*(99*a**2*d**3*e**2 + 297*a**2*d**2*e**3*x + 297*a**2*d*e**4*x**2 + 99*a**2*e**5*x**3 - 44*a*b*d**4*e + 22*a*b*d**3*e**2*x + 330*a*b*d**2*e**3*x**2 + 418*a*b*d*e**4*x**3 + 154*a*b*e**5*x**4 + 8*b**2*d**5 - 4*b**2*d**4*e*x + 3*b**2*d**3*e**2*x**2 + 113*b**2*d**2*e**3*x**3 + 161*b**2*d*e**4*x**4 + 63*b**2*e**5*x**5))/(693*e**3)`

3.125 $\int (a + bx)(A + Bx)(d + ex)^{3/2} dx$

Optimal result	1212
Mathematica [A] (verified)	1212
Rubi [A] (verified)	1213
Maple [A] (verified)	1214
Fricas [B] (verification not implemented)	1215
Sympy [A] (verification not implemented)	1215
Maxima [A] (verification not implemented)	1216
Giac [B] (verification not implemented)	1216
Mupad [B] (verification not implemented)	1217
Reduce [B] (verification not implemented)	1218

Optimal result

Integrand size = 20, antiderivative size = 83

$$\int (a + bx)(A + Bx)(d + ex)^{3/2} dx = \frac{2(bd - ae)(Bd - Ae)(d + ex)^{5/2}}{5e^3} - \frac{2(2bBd - Abe - aBe)(d + ex)^{7/2}}{7e^3} + \frac{2bB(d + ex)^{9/2}}{9e^3}$$

output

```
2/5*(-a*e+b*d)*(-A*e+B*d)*(e*x+d)^(5/2)/e^3-2/7*(-A*b*e-B*a*e+2*B*b*d)*(e*x+d)^(7/2)/e^3+2/9*b*B*(e*x+d)^(9/2)/e^3
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.84

$$\int (a + bx)(A + Bx)(d + ex)^{3/2} dx = \frac{2(d + ex)^{5/2} (9Abe(-2d + 5ex) + 9ae(-2Bd + 7Ae + 5Bex) + bB(8d^2 - 20dex + 35e^2x^2))}{315e^3}$$

input

```
Integrate[(a + b*x)*(A + B*x)*(d + e*x)^(3/2), x]
```

output

$$(2*(d + e*x)^(5/2)*(9*A*b*e*(-2*d + 5*e*x) + 9*a*e*(-2*B*d + 7*A*e + 5*B*e*x) + b*B*(8*d^2 - 20*d*e*x + 35*e^2*x^2)))/(315*e^3)$$

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)(A + Bx)(d + ex)^{3/2} dx$$

$$\downarrow 86$$

$$\int \left(\frac{(d + ex)^{5/2}(aBe + Abe - 2bBd)}{e^2} + \frac{(d + ex)^{3/2}(ae - bd)(Ae - Bd)}{e^2} + \frac{bB(d + ex)^{7/2}}{e^2} \right) dx$$

$$\downarrow 2009$$

$$-\frac{2(d + ex)^{7/2}(-aBe - Abe + 2bBd)}{7e^3} + \frac{2(d + ex)^{5/2}(bd - ae)(Bd - Ae)}{5e^3} + \frac{2bB(d + ex)^{9/2}}{9e^3}$$

input

$$\text{Int}[(a + b*x)*(A + B*x)*(d + e*x)^(3/2), x]$$

output

$$(2*(b*d - a*e)*(B*d - A*e)*(d + e*x)^(5/2))/(5*e^3) - (2*(2*b*B*d - A*b*e - a*B*e)*(d + e*x)^(7/2))/(7*e^3) + (2*b*B*(d + e*x)^(9/2))/(9*e^3)$$

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.65 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.72

method	result
pseudoelliptic	$\frac{2(ex+d)^{\frac{5}{2}} \left(\left(\frac{5x \left(\frac{7bx+a}{7} \right) B}{7} + A \left(\frac{5bx+a}{7} \right) \right) e^2 - \frac{2 \left(\left(\frac{10bx+a}{9} \right) B + Ab \right) de}{7} + \frac{8bB d^2}{63} \right)}{5e^3}$
gospers	$\frac{2(ex+d)^{\frac{5}{2}} (35bB x^2 e^2 + 45Axb e^2 + 45Bxa e^2 - 20Bxbde + 63Aa e^2 - 18Abde - 18Bade + 8bB d^2)}{315e^3}$
derivativdivides	$\frac{\frac{2bB(ex+d)^{\frac{9}{2}}}{9} + \frac{2((ae-db)B+b(Ae-Bd))(ex+d)^{\frac{7}{2}}}{7} + \frac{2(ae-db)(Ae-Bd)(ex+d)^{\frac{5}{2}}}{5}}{e^3}$
default	$\frac{\frac{2bB(ex+d)^{\frac{9}{2}}}{9} + \frac{2((ae-db)B+b(Ae-Bd))(ex+d)^{\frac{7}{2}}}{7} + \frac{2(ae-db)(Ae-Bd)(ex+d)^{\frac{5}{2}}}{5}}{e^3}$
orering	$\frac{2(ex+d)^{\frac{5}{2}} (35bB x^2 e^2 + 45Axb e^2 + 45Bxa e^2 - 20Bxbde + 63Aa e^2 - 18Abde - 18Bade + 8bB d^2)}{315e^3}$
trager	$\frac{2(35Bb e^4 x^4 + 45Ab e^4 x^3 + 45Ba e^4 x^3 + 50bBd e^3 x^3 + 63Aa e^4 x^2 + 72Abd e^3 x^2 + 72Bad e^3 x^2 + 3Bb d^2 e^2 x^2 + 126Aad e^3 x^2)}{315e^3}$
risch	$\frac{2(35Bb e^4 x^4 + 45Ab e^4 x^3 + 45Ba e^4 x^3 + 50bBd e^3 x^3 + 63Aa e^4 x^2 + 72Abd e^3 x^2 + 72Bad e^3 x^2 + 3Bb d^2 e^2 x^2 + 126Aad e^3 x^2)}{315e^3}$

```
input int((b*x+a)*(B*x+A)*(e*x+d)^(3/2), x, method=_RETURNVERBOSE)
```

```
output 2/5*(e*x+d)^(5/2)*((5/7*x*(7/9*b*x+a)*B+A*(5/7*b*x+a))*e^2-2/7*((10/9*b*x+a)*B+A*b)*d*e+8/63*b*B*d^2)/e^3
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 149 vs. $2(71) = 142$.

Time = 0.08 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.80

$$\int (a + bx)(A + Bx)(d + ex)^{3/2} dx = \frac{2(35 Bbe^4 x^4 + 8 Bbd^4 + 63 Aad^2 e^2 - 18(Ba + Ab)d^3 e + 5(10 Bbde^3 + 9(Ba + Ab)e^4)x^3 + \dots}{e^3}$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)^(3/2),x, algorithm="fricas")`

output `2/315*(35*B*b*e^4*x^4 + 8*B*b*d^4 + 63*A*a*d^2*e^2 - 18*(B*a + A*b)*d^3*e + 5*(10*B*b*d*e^3 + 9*(B*a + A*b)*e^4)*x^3 + 3*(B*b*d^2*e^2 + 21*A*a*e^4 + 24*(B*a + A*b)*d*e^3)*x^2 - (4*B*b*d^3*e - 126*A*a*d*e^3 - 9*(B*a + A*b)*d^2*e^2)*x)*sqrt(e*x + d)/e^3`

Sympy [A] (verification not implemented)

Time = 0.93 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.49

$$\int (a + bx)(A + Bx)(d + ex)^{3/2} dx = \begin{cases} \frac{2 \left(\frac{Bb(d+ex)^{9/2}}{9e^2} + \frac{(d+ex)^{7/2}(Abe+Bae-2Bbd)}{7e^2} + \frac{(d+ex)^{5/2}(Aae^2-Abde-Bade+Bbd^2)}{5e^2} \right)}{e} & \text{for } e \neq 0 \\ d^{3/2} \left(Aax + \frac{Bbx^3}{3} + \frac{x^2(Ab+Ba)}{2} \right) & \text{otherwise} \end{cases}$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)**(3/2),x)`

output `Piecewise((2*(B*b*(d + e*x)**(9/2)/(9*e**2) + (d + e*x)**(7/2)*(A*b*e + B*a*e - 2*B*b*d)/(7*e**2) + (d + e*x)**(5/2)*(A*a*e**2 - A*b*d*e - B*a*d*e + B*b*d**2)/(5*e**2))/e, Ne(e, 0)), (d**(3/2)*(A*a*x + B*b*x**3/3 + x**2*(A*b + B*a)/2), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.90

$$\int (a + bx)(A + Bx)(d + ex)^{3/2} dx = \frac{2 \left(35 (ex + d)^{\frac{9}{2}} Bb - 45 (2 Bbd - (Ba + Ab)e)(ex + d)^{\frac{7}{2}} + 63 (Bbd^2 + Aae^2 - (Ba + Ab)de) \right)}{315 e^3}$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)^(3/2),x, algorithm="maxima")`

output `2/315*(35*(e*x + d)^(9/2)*B*b - 45*(2*B*b*d - (B*a + A*b)*e)*(e*x + d)^(7/2) + 63*(B*b*d^2 + A*a*e^2 - (B*a + A*b)*d*e)*(e*x + d)^(5/2))/e^3`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 473 vs. 2(71) = 142.

Time = 0.13 (sec) , antiderivative size = 473, normalized size of antiderivative = 5.70

$$\int (a + bx)(A + Bx)(d + ex)^{3/2} dx = \frac{2 \left(315 \sqrt{ex + d} Aad^2 + 210 \left((ex + d)^{\frac{3}{2}} - 3 \sqrt{ex + dd} \right) Aad + \frac{105 \left((ex + d)^{\frac{3}{2}} - 3 \sqrt{ex + dd} \right) Bad^2}{e} + \frac{105}{e} \right)}{315 e^3}$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)^(3/2),x, algorithm="giac")`

output

```

2/315*(315*sqrt(e*x + d)*A*a*d^2 + 210*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*
d)*A*a*d + 105*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*B*a*d^2/e + 105*((e*x
+ d)^(3/2) - 3*sqrt(e*x + d)*d)*A*b*d^2/e + 21*(3*(e*x + d)^(5/2) - 10*(e
*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*A*a + 21*(3*(e*x + d)^(5/2) - 10*(
e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*B*b*d^2/e^2 + 42*(3*(e*x + d)^(5/
2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*B*a*d/e + 42*(3*(e*x + d
)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*A*b*d/e + 18*(5*(e*
x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x
+ d)*d^3)*B*b*d/e^2 + 9*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e
*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*B*a/e + 9*(5*(e*x + d)^(7/2) - 2
1*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*A*b/e
+ (35*(e*x + d)^(9/2) - 180*(e*x + d)^(7/2)*d + 378*(e*x + d)^(5/2)*d^2 -
420*(e*x + d)^(3/2)*d^3 + 315*sqrt(e*x + d)*d^4)*B*b/e^2)/e

```

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.96

$$\int (a + bx)(A + Bx)(d + ex)^{3/2} dx = \frac{2(d + ex)^{5/2} (35 B b (d + ex)^2 + 63 A a e^2 + 63 B b d^2 + 45 A b e (d + ex) + 45 B a e (d + ex))}{315 e^3}$$

input

```
int((A + B*x)*(a + b*x)*(d + e*x)^(3/2),x)
```

output

```

(2*(d + e*x)^(5/2)*(35*B*b*(d + e*x)^2 + 63*A*a*e^2 + 63*B*b*d^2 + 45*A*b*
e*(d + e*x) + 45*B*a*e*(d + e*x) - 90*B*b*d*(d + e*x) - 63*A*b*d*e - 63*B*
a*d*e))/(315*e^3)

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.67

$$\int (a + bx)(A + Bx)(d + ex)^{3/2} dx = \frac{2\sqrt{ex + d}(35b^2e^4x^4 + 90abe^4x^3 + 50b^2de^3x^3 + 63a^2e^4x^2 + 144abd e^3x^2 + 3b^2d^2e^2x^2 + 126a^2de^2x + 35b^2d^2e^2)}{315e^3}$$

input `int((b*x+a)*(B*x+A)*(e*x+d)^(3/2),x)`output `(2*sqrt(d + e*x)*(63*a**2*d**2*e**2 + 126*a**2*d*e**3*x + 63*a**2*e**4*x**2 - 36*a*b*d**3*e + 18*a*b*d**2*e**2*x + 144*a*b*d*e**3*x**2 + 90*a*b*e**4*x**3 + 8*b**2*d**4 - 4*b**2*d**3*e*x + 3*b**2*d**2*e**2*x**2 + 50*b**2*d*e**3*x**3 + 35*b**2*e**4*x**4))/(315*e**3)`

3.126 $\int (a + bx)(A + Bx)\sqrt{d + ex} dx$

Optimal result	1219
Mathematica [A] (verified)	1219
Rubi [A] (verified)	1220
Maple [A] (verified)	1221
Fricas [A] (verification not implemented)	1222
Sympy [A] (verification not implemented)	1222
Maxima [A] (verification not implemented)	1223
Giac [B] (verification not implemented)	1223
Mupad [B] (verification not implemented)	1224
Reduce [B] (verification not implemented)	1224

Optimal result

Integrand size = 20, antiderivative size = 83

$$\int (a + bx)(A + Bx)\sqrt{d + ex} dx = \frac{2(bd - ae)(Bd - Ae)(d + ex)^{3/2}}{3e^3} - \frac{2(2bBd - Abe - aBe)(d + ex)^{5/2}}{5e^3} + \frac{2bB(d + ex)^{7/2}}{7e^3}$$

output

```
2/3*(-a*e+b*d)*(-A*e+B*d)*(e*x+d)^(3/2)/e^3-2/5*(-A*b*e-B*a*e+2*B*b*d)*(e*x+d)^(5/2)/e^3+2/7*b*B*(e*x+d)^(7/2)/e^3
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.84

$$\int (a + bx)(A + Bx)\sqrt{d + ex} dx = \frac{2(d + ex)^{3/2} (7Abe(-2d + 3ex) + 7ae(-2Bd + 5Ae + 3Bex) + bB(8d^2 - 12dex + 15e^2x^2))}{105e^3}$$

input

```
Integrate[(a + b*x)*(A + B*x)*Sqrt[d + e*x],x]
```


output

$$(2*(d + e*x)^(3/2)*(7*A*b*e*(-2*d + 3*e*x) + 7*a*e*(-2*B*d + 5*A*e + 3*B*e*x) + b*B*(8*d^2 - 12*d*e*x + 15*e^2*x^2)))/(105*e^3)$$

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)(A + Bx)\sqrt{d + ex} dx$$

$$\downarrow 86$$

$$\int \left(\frac{(d + ex)^{3/2}(aBe + Abe - 2bBd)}{e^2} + \frac{\sqrt{d + ex}(ae - bd)(Ae - Bd)}{e^2} + \frac{bB(d + ex)^{5/2}}{e^2} \right) dx$$

$$\downarrow 2009$$

$$-\frac{2(d + ex)^{5/2}(-aBe - Abe + 2bBd)}{5e^3} + \frac{2(d + ex)^{3/2}(bd - ae)(Bd - Ae)}{3e^3} + \frac{2bB(d + ex)^{7/2}}{7e^3}$$

input

$$\text{Int}[(a + b*x)*(A + B*x)*\text{Sqrt}[d + e*x], x]$$

output

$$(2*(b*d - a*e)*(B*d - A*e)*(d + e*x)^(3/2))/(3*e^3) - (2*(2*b*B*d - A*b*e - a*B*e)*(d + e*x)^(5/2))/(5*e^3) + (2*b*B*(d + e*x)^(7/2))/(7*e^3)$$

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.62 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.72

method	result
pseudoelliptic	$\frac{2(ex+d)^{\frac{3}{2}} \left(\left(\frac{3\left(\frac{5bx+a}{7}\right)xB}{5} + A\left(\frac{3bx+a}{5}\right) \right) e^2 - \frac{2\left(\left(\frac{6bx+a}{7}\right)B+Ab\right)de}{5} + \frac{8bBd^2}{35} \right)}{3e^3}$
gospers	$\frac{2(ex+d)^{\frac{3}{2}} (15bBx^2e^2 + 21Axb e^2 + 21Bxa e^2 - 12Bxbde + 35Aa e^2 - 14Abde - 14Bade + 8bBd^2)}{105e^3}$
derivativdivides	$\frac{\frac{2bB(ex+d)^{\frac{7}{2}}}{7} + \frac{2((ae-db)B+b(Ae-Bd))(ex+d)^{\frac{5}{2}}}{5} + \frac{2(ae-db)(Ae-Bd)(ex+d)^{\frac{3}{2}}}{3}}{e^3}$
default	$\frac{\frac{2bB(ex+d)^{\frac{7}{2}}}{7} + \frac{2((ae-db)B+b(Ae-Bd))(ex+d)^{\frac{5}{2}}}{5} + \frac{2(ae-db)(Ae-Bd)(ex+d)^{\frac{3}{2}}}{3}}{e^3}$
orering	$\frac{2(ex+d)^{\frac{3}{2}} (15bBx^2e^2 + 21Axb e^2 + 21Bxa e^2 - 12Bxbde + 35Aa e^2 - 14Abde - 14Bade + 8bBd^2)}{105e^3}$
trager	$\frac{2(15bB e^3x^3 + 21Ab e^3x^2 + 21Ba e^3x^2 + 3bBd e^2x^2 + 35Aa e^3x + 7Abd e^2x + 7Bad e^2x - 4bB d^2ex + 35Aad e^2 - 14Ab d^2e - 14Bd^2e)}{105e^3}$
risch	$\frac{2(15bB e^3x^3 + 21Ab e^3x^2 + 21Ba e^3x^2 + 3bBd e^2x^2 + 35Aa e^3x + 7Abd e^2x + 7Bad e^2x - 4bB d^2ex + 35Aad e^2 - 14Ab d^2e - 14Bd^2e)}{105e^3}$

```
input int((b*x+a)*(B*x+A)*(e*x+d)^(1/2), x, method=_RETURNVERBOSE)
```

```
output 2/3*(e*x+d)^(3/2)*((3/5*(5/7*b*x+a)*x*B+A*(3/5*b*x+a))*e^2-2/5*((6/7*b*x+a)*B+A*b)*d*e+8/35*b*B*d^2)/e^3
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.30

$$\int (a + bx)(A + Bx)\sqrt{d + ex} dx$$

$$= \frac{2(15Bbe^3x^3 + 8Bbd^3 + 35Aade^2 - 14(Ba + Ab)d^2e + 3(Bbde^2 + 7(Ba + Ab)e^3)x^2 - (4Bbd^2e - 35Aade^2))}{105e^3}$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)^(1/2),x, algorithm="fricas")`output `2/105*(15*B*b*e^3*x^3 + 8*B*b*d^3 + 35*A*a*d*e^2 - 14*(B*a + A*b)*d^2*e + 3*(B*b*d*e^2 + 7*(B*a + A*b)*e^3)*x^2 - (4*B*b*d^2*e - 35*A*a*e^3 - 7*(B*a + A*b)*d*e^2)*x)*sqrt(e*x + d)/e^3`**Sympy [A] (verification not implemented)**

Time = 0.88 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.49

$$\int (a + bx)(A + Bx)\sqrt{d + ex} dx$$

$$= \begin{cases} \frac{2\left(\frac{Bb(d+ex)^{7/2}}{7e^2} + \frac{(d+ex)^{5/2}(Abe+BAe-2Bbd)}{5e^2} + \frac{(d+ex)^{3/2}(Aae^2-Abde-Bade+Bbd^2)}{3e^2}\right)}{e} & \text{for } e \neq 0 \\ \sqrt{d}\left(Aax + \frac{Bbx^3}{3} + \frac{x^2(Ab+Ba)}{2}\right) & \text{otherwise} \end{cases}$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)**(1/2),x)`output `Piecewise((2*(B*b*(d + e*x)**(7/2)/(7*e**2) + (d + e*x)**(5/2)*(A*b*e + B*a*e - 2*B*b*d)/(5*e**2) + (d + e*x)**(3/2)*(A*a*e**2 - A*b*d*e - B*a*d*e + B*b*d**2)/(3*e**2))/e, Ne(e, 0)), (sqrt(d)*(A*a*x + B*b*x**3/3 + x**2*(A*b + B*a)/2), True))`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.90

$$\int (a + bx)(A + Bx)\sqrt{d + ex} dx$$

$$= \frac{2 \left(15 (ex + d)^{\frac{7}{2}} Bb - 21 (2 Bbd - (Ba + Ab)e)(ex + d)^{\frac{5}{2}} + 35 (Bbd^2 + Aae^2 - (Ba + Ab)de)(ex + d)^{\frac{3}{2}} \right)}{105 e^3}$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)^(1/2),x, algorithm="maxima")`

output `2/105*(15*(e*x + d)^(7/2)*B*b - 21*(2*B*b*d - (B*a + A*b)*e)*(e*x + d)^(5/2) + 35*(B*b*d^2 + A*a*e^2 - (B*a + A*b)*d*e)*(e*x + d)^(3/2))/e^3`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 261 vs. 2(71) = 142.

Time = 0.13 (sec) , antiderivative size = 261, normalized size of antiderivative = 3.14

$$\int (a + bx)(A + Bx)\sqrt{d + ex} dx$$

$$= \frac{2 \left(105 \sqrt{ex + d} A a d + 35 \left((ex + d)^{\frac{3}{2}} - 3 \sqrt{ex + dd} \right) A a + \frac{35 \left((ex + d)^{\frac{3}{2}} - 3 \sqrt{ex + dd} \right) B a d}{e} + \frac{35 \left((ex + d)^{\frac{3}{2}} - 3 \sqrt{ex + dd} \right)}{e} \right)}{105 e^3}$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)^(1/2),x, algorithm="giac")`

output `2/105*(105*sqrt(e*x + d)*A*a*d + 35*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*A*a + 35*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*B*a*d/e + 35*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*A*b*d/e + 7*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2))*d + 15*sqrt(e*x + d)*d^2)*B*b*d/e^2 + 7*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2))*d + 15*sqrt(e*x + d)*d^2)*B*a/e + 7*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2))*d + 15*sqrt(e*x + d)*d^2)*A*b/e + 3*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2))*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*B*b/e^2)/e`

Mupad [B] (verification not implemented)

Time = 0.97 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.96

$$\int (a + bx)(A + Bx)\sqrt{d + ex} dx$$

$$= \frac{2(d + ex)^{3/2} (15Bb(d + ex)^2 + 35Aae^2 + 35Bbd^2 + 21Abe(d + ex) + 21Bae(d + ex) - 42Bbd)}{105e^3}$$

input `int((A + B*x)*(a + b*x)*(d + e*x)^(1/2),x)`output `(2*(d + e*x)^(3/2)*(15*B*b*(d + e*x)^2 + 35*A*a*e^2 + 35*B*b*d^2 + 21*A*b*e*(d + e*x) + 21*B*a*e*(d + e*x) - 42*B*b*d*(d + e*x) - 35*A*b*d*e - 35*B*a*d*e))/(105*e^3)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.18

$$\int (a + bx)(A + Bx)\sqrt{d + ex} dx$$

$$= \frac{2\sqrt{ex + d} (15b^2e^3x^3 + 42abe^3x^2 + 3b^2de^2x^2 + 35a^2e^3x + 14abd e^2x - 4b^2d^2ex + 35a^2de^2 - 28abd^2e + 15b^2d^2e^2)}{105e^3}$$

input `int((b*x+a)*(B*x+A)*(e*x+d)^(1/2),x)`output `(2*sqrt(d + e*x)*(35*a**2*d*e**2 + 35*a**2*e**3*x - 28*a*b*d**2*e + 14*a*b*d*e**2*x + 42*a*b*e**3*x**2 + 8*b**2*d**3 - 4*b**2*d**2*e*x + 3*b**2*d*e**2*x**2 + 15*b**2*e**3*x**3))/(105*e**3)`

3.127 $\int \frac{(a+bx)(A+Bx)}{\sqrt{d+ex}} dx$

Optimal result	1225
Mathematica [A] (verified)	1225
Rubi [A] (verified)	1226
Maple [A] (verified)	1227
Fricas [A] (verification not implemented)	1228
Sympy [A] (verification not implemented)	1228
Maxima [A] (verification not implemented)	1229
Giac [A] (verification not implemented)	1229
Mupad [B] (verification not implemented)	1230
Reduce [B] (verification not implemented)	1230

Optimal result

Integrand size = 20, antiderivative size = 81

$$\int \frac{(a+bx)(A+Bx)}{\sqrt{d+ex}} dx = \frac{2(bd-ae)(Bd-Ae)\sqrt{d+ex}}{e^3} - \frac{2(2bBd-Abe-aBe)(d+ex)^{3/2}}{3e^3} + \frac{2bB(d+ex)^{5/2}}{5e^3}$$

output

```
2*(-a*e+b*d)*(-A*e+B*d)*(e*x+d)^(1/2)/e^3-2/3*(-A*b*e-B*a*e+2*B*b*d)*(e*x+d)^(3/2)/e^3+2/5*b*B*(e*x+d)^(5/2)/e^3
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.84

$$\int \frac{(a+bx)(A+Bx)}{\sqrt{d+ex}} dx = \frac{2\sqrt{d+ex}(5Abe(-2d+ex) + 5ae(-2Bd+3Ae+Bex) + bB(8d^2-4dex+3e^2x^2))}{15e^3}$$

input

```
Integrate[((a + b*x)*(A + B*x))/Sqrt[d + e*x],x]
```

output

$$(2*\text{Sqrt}[d + e*x]*(5*A*b*e*(-2*d + e*x) + 5*a*e*(-2*B*d + 3*A*e + B*e*x) + b*B*(8*d^2 - 4*d*e*x + 3*e^2*x^2)))/(15*e^3)$$

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)(A + Bx)}{\sqrt{d + ex}} dx$$

↓ 86

$$\int \left(\frac{\sqrt{d + ex}(aBe + Abe - 2bBd)}{e^2} + \frac{(ae - bd)(Ae - Bd)}{e^2\sqrt{d + ex}} + \frac{bB(d + ex)^{3/2}}{e^2} \right) dx$$

↓ 2009

$$-\frac{2(d + ex)^{3/2}(-aBe - Abe + 2bBd)}{3e^3} + \frac{2\sqrt{d + ex}(bd - ae)(Bd - Ae)}{e^3} + \frac{2bB(d + ex)^{5/2}}{5e^3}$$

input

$$\text{Int}[(a + b*x)*(A + B*x)]/\text{Sqrt}[d + e*x], x]$$

output

$$(2*(b*d - a*e)*(B*d - A*e)*\text{Sqrt}[d + e*x])/e^3 - (2*(2*b*B*d - A*b*e - a*B*e)*(d + e*x)^{(3/2)})/(3*e^3) + (2*b*B*(d + e*x)^{(5/2)})/(5*e^3)$$

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.62 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.74

method	result	size
pseudoelliptic	$\frac{2\sqrt{ex+d} \left(\left(\frac{\left(\frac{3bx+a}{5}\right)^{xB} + A\left(\frac{bx}{3}+a\right)}{e^3} \right) e^2 - \frac{2\left(\frac{2bx+a}{5}\right)^{B+Ab} de + \frac{8bBd^2}{15}}{e^3} \right)}{e^3}$	60
derivativedivides	$\frac{\frac{2bB(ex+d)^{\frac{5}{2}}}{5} + \frac{2((ae-db)B+b(Ae-Bd))(ex+d)^{\frac{3}{2}}}{e^3} + 2(ae-db)(Ae-Bd)\sqrt{ex+d}}{e^3}$	72
default	$\frac{\frac{2bB(ex+d)^{\frac{5}{2}}}{5} + \frac{2((ae-db)B+b(Ae-Bd))(ex+d)^{\frac{3}{2}}}{e^3} + 2(ae-db)(Ae-Bd)\sqrt{ex+d}}{e^3}$	72
gospers	$\frac{2\sqrt{ex+d} (3bBx^2e^2 + 5Axb e^2 + 5Bxa e^2 - 4Bxbde + 15Aa e^2 - 10Abde - 10Bade + 8bBd^2)}{15e^3}$	73
trager	$\frac{2\sqrt{ex+d} (3bBx^2e^2 + 5Axb e^2 + 5Bxa e^2 - 4Bxbde + 15Aa e^2 - 10Abde - 10Bade + 8bBd^2)}{15e^3}$	73
risch	$\frac{2\sqrt{ex+d} (3bBx^2e^2 + 5Axb e^2 + 5Bxa e^2 - 4Bxbde + 15Aa e^2 - 10Abde - 10Bade + 8bBd^2)}{15e^3}$	73
orering	$\frac{2\sqrt{ex+d} (3bBx^2e^2 + 5Axb e^2 + 5Bxa e^2 - 4Bxbde + 15Aa e^2 - 10Abde - 10Bade + 8bBd^2)}{15e^3}$	73

```
input int((b*x+a)*(B*x+A)/(e*x+d)^(1/2), x, method=_RETURNVERBOSE)
```

```
output 2*(e*x+d)^(1/2)*((1/3*(3/5*b*x+a)*x*B+A*(1/3*b*x+a))*e^2-2/3*((2/5*b*x+a)*B+A*b)*d*e+8/15*b*B*d^2)/e^3
```


Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.86

$$\int \frac{(a + bx)(A + Bx)}{\sqrt{d + ex}} dx$$

$$= \frac{2(3Bbe^2x^2 + 8Bbd^2 + 15Aae^2 - 10(Ba + Ab)de - (4Bbde - 5(Ba + Ab)e^2)x)\sqrt{ex + d}}{15e^3}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^(1/2),x, algorithm="fricas")`output `2/15*(3*B*b*e^2*x^2 + 8*B*b*d^2 + 15*A*a*e^2 - 10*(B*a + A*b)*d*e - (4*B*b*d*e - 5*(B*a + A*b)*e^2)*x)*sqrt(e*x + d)/e^3`**Sympy [A] (verification not implemented)**

Time = 0.84 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.51

$$\int \frac{(a + bx)(A + Bx)}{\sqrt{d + ex}} dx$$

$$= \begin{cases} 2 \left(\frac{Bb(d+ex)^{\frac{5}{2}}}{5e^2} + \frac{(d+ex)^{\frac{3}{2}}(Abe+BAe-2Bbd)}{3e^2} + \frac{\sqrt{d+ex}(Aae^2-Abde-Bade+Bbd^2)}{e^2} \right) & \text{for } e \neq 0 \\ \frac{Aax + \frac{Bbx^3}{3} + \frac{x^2(Ab+Ba)}{2}}{\sqrt{d}} & \text{otherwise} \end{cases}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)**(1/2),x)`output `Piecewise((2*(B*b*(d + e*x)**(5/2)/(5*e**2) + (d + e*x)**(3/2)*(A*b*e + B*a*e - 2*B*b*d)/(3*e**2) + sqrt(d + e*x)*(A*a*e**2 - A*b*d*e - B*a*d*e + B*b*d**2)/e**2)/e, Ne(e, 0)), ((A*a*x + B*b*x**3/3 + x**2*(A*b + B*a)/2)/sqrt(d), True))`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.93

$$\int \frac{(a+bx)(A+Bx)}{\sqrt{d+ex}} dx$$

$$= \frac{2 \left(3(ex+d)^{\frac{5}{2}} Bb - 5(2Bbd - (Ba+Ab)e)(ex+d)^{\frac{3}{2}} + 15(Bbd^2 + Aae^2 - (Ba+Ab)de)\sqrt{ex+d} \right)}{15e^3}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^(1/2),x, algorithm="maxima")`output `2/15*(3*(e*x + d)^(5/2)*B*b - 5*(2*B*b*d - (B*a + A*b)*e)*(e*x + d)^(3/2) + 15*(B*b*d^2 + A*a*e^2 - (B*a + A*b)*d*e)*sqrt(e*x + d))/e^3`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.30

$$\int \frac{(a+bx)(A+Bx)}{\sqrt{d+ex}} dx$$

$$= \frac{2 \left(15\sqrt{ex+d}Aa + \frac{5((ex+d)^{\frac{3}{2}} - 3\sqrt{ex+d})Ba}{e} + \frac{5((ex+d)^{\frac{3}{2}} - 3\sqrt{ex+d})Ab}{e} + \frac{(3(ex+d)^{\frac{5}{2}} - 10(ex+d)^{\frac{3}{2}}d + 15\sqrt{ex+d}d^2)Bb}{e^2} \right)}{15e}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^(1/2),x, algorithm="giac")`output `2/15*(15*sqrt(e*x + d)*A*a + 5*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*B*a/e + 5*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*A*b/e + (3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*B*b/e^2)/e`

Mupad [B] (verification not implemented)

Time = 0.97 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.99

$$\int \frac{(a + bx)(A + Bx)}{\sqrt{d + ex}} dx$$

$$= \frac{2\sqrt{d + ex}(3Bb(d + ex)^2 + 15Aae^2 + 15Bbd^2 + 5Abe(d + ex) + 5Bae(d + ex) - 10Bbd(d + ex))}{15e^3}$$

input `int(((A + B*x)*(a + b*x))/(d + e*x)^(1/2),x)`output `(2*(d + e*x)^(1/2)*(3*B*b*(d + e*x)^2 + 15*A*a*e^2 + 15*B*b*d^2 + 5*A*b*e*(d + e*x) + 5*B*a*e*(d + e*x) - 10*B*b*d*(d + e*x) - 15*A*b*d*e - 15*B*a*d*e))/(15*e^3)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.75

$$\int \frac{(a + bx)(A + Bx)}{\sqrt{d + ex}} dx$$

$$= \frac{2\sqrt{ex + d}(3b^2e^2x^2 + 10abe^2x - 4b^2dex + 15a^2e^2 - 20abde + 8b^2d^2)}{15e^3}$$

input `int((b*x+a)*(B*x+A)/(e*x+d)^(1/2),x)`output `(2*sqrt(d + e*x)*(15*a**2*e**2 - 20*a*b*d*e + 10*a*b*e**2*x + 8*b**2*d**2 - 4*b**2*d*e*x + 3*b**2*e**2*x**2))/(15*e**3)`

3.128 $\int \frac{(a+bx)(A+Bx)}{(d+ex)^{3/2}} dx$

Optimal result	1231
Mathematica [A] (verified)	1231
Rubi [A] (verified)	1232
Maple [A] (verified)	1233
Fricas [A] (verification not implemented)	1233
Sympy [A] (verification not implemented)	1234
Maxima [A] (verification not implemented)	1234
Giac [A] (verification not implemented)	1235
Mupad [B] (verification not implemented)	1235
Reduce [B] (verification not implemented)	1236

Optimal result

Integrand size = 20, antiderivative size = 79

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{3/2}} dx = -\frac{2(bd-ae)(Bd-Ae)}{e^3\sqrt{d+ex}} - \frac{2(2bBd-Abe-aBe)\sqrt{d+ex}}{e^3} + \frac{2bB(d+ex)^{3/2}}{3e^3}$$

output `-2*(-a*e+b*d)*(-A*e+B*d)/e^3/(e*x+d)^(1/2)-2*(-A*b*e-B*a*e+2*B*b*d)*(e*x+d)^(1/2)/e^3+2/3*b*B*(e*x+d)^(3/2)/e^3`

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.86

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{3/2}} dx = \frac{6Abe(2d+ex) + 6ae(2Bd-Ae+Bex) + 2bB(-8d^2-4dex+e^2x^2)}{3e^3\sqrt{d+ex}}$$

input `Integrate[((a + b*x)*(A + B*x))/(d + e*x)^(3/2),x]`

output `(6*A*b*e*(2*d + e*x) + 6*a*e*(2*B*d - A*e + B*e*x) + 2*b*B*(-8*d^2 - 4*d*e*x + e^2*x^2))/(3*e^3*Sqrt[d + e*x])`

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^{3/2}} dx$$

↓ 86

$$\int \left(\frac{aBe + Abe - 2bBd}{e^2 \sqrt{d + ex}} + \frac{(ae - bd)(Ae - Bd)}{e^2 (d + ex)^{3/2}} + \frac{bB\sqrt{d + ex}}{e^2} \right) dx$$

↓ 2009

$$-\frac{2\sqrt{d + ex}(-aBe - Abe + 2bBd)}{e^3} - \frac{2(bd - ae)(Bd - Ae)}{e^3 \sqrt{d + ex}} + \frac{2bB(d + ex)^{3/2}}{3e^3}$$

input `Int[((a + b*x)*(A + B*x))/(d + e*x)^(3/2), x]`

output `(-2*(b*d - a*e)*(B*d - A*e))/(e^3*Sqrt[d + e*x]) - (2*(2*b*B*d - A*b*e - a*B*e)*Sqrt[d + e*x])/e^3 + (2*b*B*(d + e*x)^(3/2))/(3*e^3)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.81

method	result	size
pseudoelliptic	$\frac{((2bx^2+6ax)B-6A(-bx+a))e^2+12d\left(-\frac{2bx}{3}+a\right)B+Ab}{3\sqrt{ex+d}e^3}e^{-16bBd^2}$	64
risch	$\frac{2(ebBx+3Abe+3Bae-5Bbd)\sqrt{ex+d}}{3e^3} - \frac{2(Aae^2-Abde-Bade+bBd^2)}{e^3\sqrt{ex+d}}$	72
gospers	$-\frac{2(-bBx^2e^2-3Axb e^2-3Bxa e^2+4Bxbde+3Aae^2-6Abde-6Bade+8bBd^2)}{3\sqrt{ex+d}e^3}$	73
trager	$-\frac{2(-bBx^2e^2-3Axb e^2-3Bxa e^2+4Bxbde+3Aae^2-6Abde-6Bade+8bBd^2)}{3\sqrt{ex+d}e^3}$	73
orering	$-\frac{2(-bBx^2e^2-3Axb e^2-3Bxa e^2+4Bxbde+3Aae^2-6Abde-6Bade+8bBd^2)}{3\sqrt{ex+d}e^3}$	73
derivativedivides	$\frac{\frac{2bB(ex+d)^{\frac{3}{2}}}{3}+2Abe\sqrt{ex+d}+2Bae\sqrt{ex+d}-4Bbd\sqrt{ex+d}-\frac{2(Aae^2-Abde-Bade+bBd^2)}{\sqrt{ex+d}}}{e^3}$	86
default	$\frac{\frac{2bB(ex+d)^{\frac{3}{2}}}{3}+2Abe\sqrt{ex+d}+2Bae\sqrt{ex+d}-4Bbd\sqrt{ex+d}-\frac{2(Aae^2-Abde-Bade+bBd^2)}{\sqrt{ex+d}}}{e^3}$	86

input `int((b*x+a)*(B*x+A)/(e*x+d)^(3/2),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{3} * \left(\frac{(2bx^2+6ax)B-6A(-bx+a))e^2+12d\left(-\frac{2bx}{3}+a\right)B+Ab}{3\sqrt{ex+d}e^3} e^{-16bBd^2} \right)$$

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.00

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{3/2}} dx = \frac{2(Bbe^2x^2 - 8Bbd^2 - 3Aae^2 + 6(Ba+Ab)de - (4Bbde - 3(Ba+Ab)e^2)x)}{3(e^4x + de^3)}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^(3/2),x, algorithm="fricas")`

output
$$\frac{2}{3} * (B*b*e^2*x^2 - 8*B*b*d^2 - 3*A*a*e^2 + 6*(B*a + A*b)*d*e - (4*B*b*d*e - 3*(B*a + A*b)*e^2)*x) * \text{sqrt}(e*x + d) / (e^4*x + d*e^3)$$

Sympy [A] (verification not implemented)

Time = 2.26 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.35

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^{3/2}} dx = \begin{cases} \frac{2 \left(\frac{Bb(d+ex)^{3/2}}{3e^2} + \frac{\sqrt{d+ex}(Abe+Ba e-2Bbd)}{e^2} + \frac{(-Ae+Bd)(ae-bd)}{e^2\sqrt{d+ex}} \right)}{e} & \text{for } e \neq 0 \\ \frac{Aax + \frac{Bbx^3}{3} + \frac{x^2(Ab+Ba)}{2}}{d^{3/2}} & \text{otherwise} \end{cases}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)**(3/2),x)`output `Piecewise((2*(B*b*(d + e*x)**(3/2)/(3*e**2) + sqrt(d + e*x)*(A*b*e + B*a*e - 2*B*b*d)/e**2 + (-A*e + B*d)*(a*e - b*d)/(e**2*sqrt(d + e*x)))/e, Ne(e, 0)), ((A*a*x + B*b*x**3/3 + x**2*(A*b + B*a)/2)/d**(3/2), True))`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.04

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^{3/2}} dx = \frac{2 \left(\frac{(ex+d)^{3/2} Bb - 3(2Bbd - (Ba+Ab)e)\sqrt{ex+d}}{e^2} - \frac{3(Bbd^2 + Aae^2 - (Ba+Ab)de)}{\sqrt{ex+de^2}} \right)}{3e}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^(3/2),x, algorithm="maxima")`output `2/3*(((e*x + d)^(3/2)*B*b - 3*(2*B*b*d - (B*a + A*b)*e)*sqrt(e*x + d))/e^2 - 3*(B*b*d^2 + A*a*e^2 - (B*a + A*b)*d*e)/(sqrt(e*x + d)*e^2))/e`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.27

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{3/2}} dx = -\frac{2(Bbd^2 - Bade - Abde + Aae^2)}{\sqrt{ex+d}e^3} + \frac{2\left((ex+d)^{\frac{3}{2}}Bbe^6 - 6\sqrt{ex+d}Bbde^6 + 3\sqrt{ex+d}Bae^7 + 3\sqrt{ex+d}Abe^7\right)}{3e^9}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^(3/2),x, algorithm="giac")`

output

```
-2*(B*b*d^2 - B*a*d*e - A*b*d*e + A*a*e^2)/(sqrt(e*x + d)*e^3) + 2/3*((e*x + d)^(3/2)*B*b*e^6 - 6*sqrt(e*x + d)*B*b*d*e^6 + 3*sqrt(e*x + d)*B*a*e^7 + 3*sqrt(e*x + d)*A*b*e^7)/e^9
```

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.00

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{3/2}} dx = \frac{\frac{2Bb(d+ex)^2}{3} - 2Aae^2 - 2Bbd^2 + 2Abe(d+ex) + 2Bae(d+ex) - 4Bbd}{e^3\sqrt{d+ex}}$$

input

```
int(((A + B*x)*(a + b*x))/(d + e*x)^(3/2),x)
```

output

```
((2*B*b*(d + e*x)^2)/3 - 2*A*a*e^2 - 2*B*b*d^2 + 2*A*b*e*(d + e*x) + 2*B*a*e*(d + e*x) - 4*B*b*d*(d + e*x) + 2*A*b*d*e + 2*B*a*d*e)/(e^3*(d + e*x)^(1/2))
```


Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.78

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^{3/2}} dx = \frac{\frac{2}{3}b^2e^2x^2 + 4abe^2x - \frac{8}{3}b^2dex - 2a^2e^2 + 8abde - \frac{16}{3}b^2d^2}{\sqrt{ex + d}e^3}$$

input `int((b*x+a)*(B*x+A)/(e*x+d)^(3/2),x)`

output `(2*(- 3*a**2*e**2 + 12*a*b*d*e + 6*a*b*e**2*x - 8*b**2*d**2 - 4*b**2*d*e*x + b**2*e**2*x**2))/(3*sqrt(d + e*x)*e**3)`

3.129 $\int \frac{(a+bx)(A+Bx)}{(d+ex)^{5/2}} dx$

Optimal result	1237
Mathematica [A] (verified)	1237
Rubi [A] (verified)	1238
Maple [A] (verified)	1239
Fricas [A] (verification not implemented)	1239
Sympy [B] (verification not implemented)	1240
Maxima [A] (verification not implemented)	1240
Giac [A] (verification not implemented)	1241
Mupad [B] (verification not implemented)	1241
Reduce [B] (verification not implemented)	1242

Optimal result

Integrand size = 20, antiderivative size = 79

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{5/2}} dx = -\frac{2(bd-ae)(Bd-Ae)}{3e^3(d+ex)^{3/2}} + \frac{2(2bBd-Abe-aBe)}{e^3\sqrt{d+ex}} + \frac{2bB\sqrt{d+ex}}{e^3}$$

output -2/3*(-a*e+b*d)*(-A*e+B*d)/e^3/(e*x+d)^(3/2)+2*(-A*b*e-B*a*e+2*B*b*d)/e^3/(e*x+d)^(1/2)+2*b*B*(e*x+d)^(1/2)/e^3

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.86

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{5/2}} dx = \frac{2(Abe(2d+3ex) + ae(2Bd+ Ae+ 3Bex) - bB(8d^2+ 12dex+ 3e^2x^2))}{3e^3(d+ex)^{3/2}}$$

input Integrate[((a + b*x)*(A + B*x))/(d + e*x)^(5/2),x]

output (-2*(A*b*e*(2*d + 3*e*x) + a*e*(2*B*d + A*e + 3*B*e*x) - b*B*(8*d^2 + 12*d*e*x + 3*e^2*x^2))/(3*e^3*(d + e*x)^(3/2))

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^{5/2}} dx$$

↓ 86

$$\int \left(\frac{aBe + Abe - 2bBd}{e^2(d + ex)^{3/2}} + \frac{(ae - bd)(Ae - Bd)}{e^2(d + ex)^{5/2}} + \frac{bB}{e^2\sqrt{d + ex}} \right) dx$$

↓ 2009

$$\frac{2(-aBe - Abe + 2bBd)}{e^3\sqrt{d + ex}} - \frac{2(bd - ae)(Bd - Ae)}{3e^3(d + ex)^{3/2}} + \frac{2bB\sqrt{d + ex}}{e^3}$$

input `Int[((a + b*x)*(A + B*x))/(d + e*x)^(5/2), x]`

output `(-2*(b*d - a*e)*(B*d - A*e))/(3*e^3*(d + e*x)^(3/2)) + (2*(2*b*B*d - A*b*e - a*B*e))/(e^3*sqrt[d + e*x]) + (2*b*B*sqrt[d + e*x])/e^3`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.80

method	result	size
pseudoelliptic	$-\frac{2(((-3bx^2+3ax)B+A(3bx+a))e^2+2((-6bx+a)B+Ab)de-8bBd^2)}{3(ex+d)^{\frac{3}{2}}e^3}$	63
gospers	$-\frac{2(-3bBx^2e^2+3Axb e^2+3Bxa e^2-12Bxbde+Aa e^2+2Abde+2Bade-8bBd^2)}{3(ex+d)^{\frac{3}{2}}e^3}$	72
trager	$-\frac{2(-3bBx^2e^2+3Axb e^2+3Bxa e^2-12Bxbde+Aa e^2+2Abde+2Bade-8bBd^2)}{3(ex+d)^{\frac{3}{2}}e^3}$	72
orering	$-\frac{2(-3bBx^2e^2+3Axb e^2+3Bxa e^2-12Bxbde+Aa e^2+2Abde+2Bade-8bBd^2)}{3(ex+d)^{\frac{3}{2}}e^3}$	72
derivativedivides	$\frac{2bB\sqrt{ex+d} - \frac{2(Abe+BAe-2Bbd)}{\sqrt{ex+d}}}{e^3} - \frac{2(Aa e^2 - Abde - Bade + bBd^2)}{3(ex+d)^{\frac{3}{2}}}$	74
default	$\frac{2bB\sqrt{ex+d} - \frac{2(Abe+BAe-2Bbd)}{\sqrt{ex+d}}}{e^3} - \frac{2(Aa e^2 - Abde - Bade + bBd^2)}{3(ex+d)^{\frac{3}{2}}}$	74
risch	$\frac{2bB\sqrt{ex+d}}{e^3} - \frac{2(3Axb e^2+3Bxa e^2-6Bxbde+Aa e^2+2Abde+2Bade-5bBd^2)}{3e^3(ex+d)^{\frac{3}{2}}}$	77

input `int((b*x+a)*(B*x+A)/(e*x+d)^(5/2),x,method=_RETURNVERBOSE)`

output
$$-2/3/(e*x+d)^{(3/2)}*(((-3*b*x^2+3*a*x)*B+A*(3*b*x+a))*e^2+2*((-6*b*x+a)*B+A*b)*d*e-8*b*B*d^2)/e^3$$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.15

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^{5/2}} dx = \frac{2(3Bbe^2x^2 + 8Bbd^2 - Aae^2 - 2(Ba + Ab)de + 3(4Bbde - (Ba + Ab)e^2)x)}{3(e^5x^2 + 2de^4x + d^2e^3)}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^(5/2),x, algorithm="fricas")`

output
$$2/3*(3*B*b*e^2*x^2 + 8*B*b*d^2 - A*a*e^2 - 2*(B*a + A*b)*d*e + 3*(4*B*b*d*d*e - (B*a + A*b)*e^2)*x)*sqrt(e*x + d)/(e^5*x^2 + 2*d*e^4*x + d^2*e^3)$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 355 vs. $2(80) = 160$.

Time = 0.37 (sec) , antiderivative size = 355, normalized size of antiderivative = 4.49

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{5/2}} dx = \left\{ \begin{array}{l} -\frac{2Aae^2}{3de^3\sqrt{d+ex}+3e^4x\sqrt{d+ex}} - \frac{4Abde}{3de^3\sqrt{d+ex}+3e^4x\sqrt{d+ex}} - \frac{6Abe^2x}{3de^3\sqrt{d+ex}+3e^4x\sqrt{d+ex}} - \frac{4}{3de^3\sqrt{d+ex}} \\ \frac{Aax + \frac{Abx^2}{2} + \frac{Bax^2}{2} + \frac{Bbx^3}{3}}{d^{5/2}} \end{array} \right.$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)**(5/2),x)`

output `Piecewise((-2*A*a*e**2/(3*d*e**3*sqrt(d + e*x) + 3*e**4*x*sqrt(d + e*x)) - 4*A*b*d*e/(3*d*e**3*sqrt(d + e*x) + 3*e**4*x*sqrt(d + e*x)) - 6*A*b*e**2*x/(3*d*e**3*sqrt(d + e*x) + 3*e**4*x*sqrt(d + e*x)) - 4*B*a*d*e/(3*d*e**3*sqrt(d + e*x) + 3*e**4*x*sqrt(d + e*x)) - 6*B*a*e**2*x/(3*d*e**3*sqrt(d + e*x) + 3*e**4*x*sqrt(d + e*x)) + 16*B*b*d**2/(3*d*e**3*sqrt(d + e*x) + 3*e**4*x*sqrt(d + e*x)) + 24*B*b*d*e*x/(3*d*e**3*sqrt(d + e*x) + 3*e**4*x*sqrt(d + e*x)) + 6*B*b*e**2*x**2/(3*d*e**3*sqrt(d + e*x) + 3*e**4*x*sqrt(d + e*x)), Ne(e, 0)), ((A*a*x + A*b*x**2/2 + B*a*x**2/2 + B*b*x**3/3)/d**(5/2), True))`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.00

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{5/2}} dx = \frac{2 \left(\frac{3\sqrt{ex+d}Bb}{e^2} - \frac{Bbd^2+Aae^2-(Ba+Ab)de-3(2Bbd-(Ba+Ab)e)(ex+d)}{(ex+d)^{3/2}e^2} \right)}{3e}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^(5/2),x, algorithm="maxima")`

output `2/3*(3*sqrt(e*x + d)*B*b/e^2 - (B*b*d^2 + A*a*e^2 - (B*a + A*b)*d*e - 3*(2*B*b*d - (B*a + A*b)*e)*(e*x + d))/((e*x + d)^(3/2)*e^2)/e`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.04

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{5/2}} dx = \frac{2\sqrt{ex+d}Bb}{e^3} + \frac{2(6(ex+d)Bbd - Bbd^2 - 3(ex+d)Bae - 3(ex+d)Abe + Bade + Abde - Aae^2)}{3(ex+d)^{3/2}e^3}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^(5/2),x, algorithm="giac")`

output `2*sqrt(e*x + d)*B*b/e^3 + 2/3*(6*(e*x + d)*B*b*d - B*b*d^2 - 3*(e*x + d)*B*a*e - 3*(e*x + d)*A*b*e + B*a*d*e + A*b*d*e - A*a*e^2)/((e*x + d)^(3/2)*e^3)`

Mupad [B] (verification not implemented)

Time = 0.99 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.91

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{5/2}} dx = \frac{2Aae^2 - 16Bbd^2 + 6Ab e^2 x + 6Ba e^2 x - 6Bb e^2 x^2 + 4Abde + 4Bade - 24Bbdex}{3e^3(d+ex)^{3/2}}$$

input `int(((A + B*x)*(a + b*x))/(d + e*x)^(5/2),x)`

output `-(2*A*a*e^2 - 16*B*b*d^2 + 6*A*b*e^2*x + 6*B*a*e^2*x - 6*B*b*e^2*x^2 + 4*A*b*d*e + 4*B*a*d*e - 24*B*b*d*e*x)/(3*e^3*(d + e*x)^(3/2))`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.89

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^{5/2}} dx = \frac{2b^2e^2x^2 - 4abe^2x + 8b^2dex - \frac{2}{3}a^2e^2 - \frac{8}{3}abde + \frac{16}{3}b^2d^2}{\sqrt{ex + d}e^3(ex + d)}$$

input `int((b*x+a)*(B*x+A)/(e*x+d)^(5/2),x)`

output `(2*(- a**2*e**2 - 4*a*b*d*e - 6*a*b*e**2*x + 8*b**2*d**2 + 12*b**2*d*e*x + 3*b**2*e**2*x**2))/(3*sqrt(d + e*x)*e**3*(d + e*x))`

3.130 $\int \frac{(a+bx)(A+Bx)}{(d+ex)^{7/2}} dx$

Optimal result	1243
Mathematica [A] (verified)	1243
Rubi [A] (verified)	1244
Maple [A] (verified)	1245
Fricas [A] (verification not implemented)	1245
Sympy [B] (verification not implemented)	1246
Maxima [A] (verification not implemented)	1247
Giac [A] (verification not implemented)	1247
Mupad [B] (verification not implemented)	1248
Reduce [B] (verification not implemented)	1248

Optimal result

Integrand size = 20, antiderivative size = 81

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{7/2}} dx = -\frac{2(bd-ae)(Bd-Ae)}{5e^3(d+ex)^{5/2}} + \frac{2(2bBd-Abe-aBe)}{3e^3(d+ex)^{3/2}} - \frac{2bB}{e^3\sqrt{d+ex}}$$

output

$$-2/5*(-a*e+b*d)*(-A*e+B*d)/e^3/(e*x+d)^(5/2)+2/3*(-A*b*e-B*a*e+2*B*b*d)/e^3/(e*x+d)^(3/2)-2*b*B/e^3/(e*x+d)^(1/2)$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.84

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{7/2}} dx = \frac{2(Abe(2d+5ex) + ae(2Bd+3Ae+5Bex) + bB(8d^2+20dex+15e^2x^2))}{15e^3(d+ex)^{5/2}}$$

input

$$\text{Integrate}[\frac{(a+b*x)*(A+B*x)}{(d+e*x)^(7/2)},x]$$

output

$$\frac{(-2*(A*b*e*(2*d+5*e*x) + a*e*(2*B*d+3*A*e+5*B*e*x) + b*B*(8*d^2+20*d*e*x+15*e^2*x^2))}{(15*e^3*(d+e*x)^(5/2))}$$

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^{7/2}} dx$$

↓ 86

$$\int \left(\frac{aBe + Abe - 2bBd}{e^2(d + ex)^{5/2}} + \frac{(ae - bd)(Ae - Bd)}{e^2(d + ex)^{7/2}} + \frac{bB}{e^2(d + ex)^{3/2}} \right) dx$$

↓ 2009

$$\frac{2(-aBe - Abe + 2bBd)}{3e^3(d + ex)^{3/2}} - \frac{2(bd - ae)(Bd - Ae)}{5e^3(d + ex)^{5/2}} - \frac{2bB}{e^3\sqrt{d + ex}}$$

input `Int[((a + b*x)*(A + B*x))/(d + e*x)^(7/2), x]`

output `(-2*(b*d - a*e)*(B*d - A*e))/(5*e^3*(d + e*x)^(5/2)) + (2*(2*b*B*d - A*b*e - a*B*e))/(3*e^3*(d + e*x)^(3/2)) - (2*b*B)/(e^3*sqrt[d + e*x])`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.75

method	result	size
pseudoelliptic	$\frac{(-10x(3bx+a)B-6A\left(\frac{5bx}{3}+a\right))e^2-4((10bx+a)B+Ab)de-16bBd^2}{15(ex+d)^{\frac{5}{2}}e^3}$	61
gospers	$\frac{2(15bBx^2e^2+5Axb^2e^2+5Bxa^2e^2+20Bxbde+3Aae^2+2Abde+2Bade+8bBd^2)}{15(ex+d)^{\frac{5}{2}}e^3}$	73
trager	$\frac{2(15bBx^2e^2+5Axb^2e^2+5Bxa^2e^2+20Bxbde+3Aae^2+2Abde+2Bade+8bBd^2)}{15(ex+d)^{\frac{5}{2}}e^3}$	73
orering	$\frac{2(15bBx^2e^2+5Axb^2e^2+5Bxa^2e^2+20Bxbde+3Aae^2+2Abde+2Bade+8bBd^2)}{15(ex+d)^{\frac{5}{2}}e^3}$	73
derivativdivides	$\frac{-\frac{2bB}{\sqrt{ex+d}}-\frac{2(Abe+BAe-2Bbd)}{3(ex+d)^{\frac{3}{2}}}-\frac{2(Aae^2-Abde-Bade+bBd^2)}{5(ex+d)^{\frac{5}{2}}}}{e^3}$	75
default	$\frac{-\frac{2bB}{\sqrt{ex+d}}-\frac{2(Abe+BAe-2Bbd)}{3(ex+d)^{\frac{3}{2}}}-\frac{2(Aae^2-Abde-Bade+bBd^2)}{5(ex+d)^{\frac{5}{2}}}}{e^3}$	75

input `int((b*x+a)*(B*x+A)/(e*x+d)^(7/2),x,method=_RETURNVERBOSE)`

output `1/15*((-10*x*(3*b*x+a)*B-6*A*(5/3*b*x+a))*e^2-4*((10*b*x+a)*B+A*b)*d*e-16*b*B*d^2)/(e*x+d)^(5/2)/e^3`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.25

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{7/2}} dx = \frac{2(15Bbe^2x^2+8Bbd^2+3Aae^2+2(Ba+Ab)de+5(4Bbde+(Ba+Ab)e^2)x)\sqrt{ex+d}}{15(e^6x^3+3de^5x^2+3d^2e^4x+d^3e^3)}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^(7/2),x, algorithm="fricas")`

output

```
-2/15*(15*B*b*e^2*x^2 + 8*B*b*d^2 + 3*A*a*e^2 + 2*(B*a + A*b)*d*e + 5*(4*B
*b*d*e + (B*a + A*b)*e^2)*x)*sqrt(e*x + d)/(e^6*x^3 + 3*d*e^5*x^2 + 3*d^2*
e^4*x + d^3*e^3)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 520 vs. $2(85) = 170$.

Time = 0.51 (sec) , antiderivative size = 520, normalized size of antiderivative = 6.42

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^{7/2}} dx = \left\{ \begin{array}{l} -\frac{6Aae^2}{15d^2e^3\sqrt{d+ex}+30de^4x\sqrt{d+ex}+15e^5x^2\sqrt{d+ex}} - \frac{4Abde}{15d^2e^3\sqrt{d+ex}+30de^4x\sqrt{d+ex}+15e^5x^2\sqrt{d+ex}} \\ \frac{Aax + \frac{Abx^2}{2} + \frac{Bax^2}{2} + \frac{Bbx^3}{3}}{d^{7/2}} \end{array} \right.$$

input

```
integrate((b*x+a)*(B*x+A)/(e*x+d)**(7/2), x)
```

output

```
Piecewise((-6*A*a*e**2/(15*d**2*e**3*sqrt(d + e*x) + 30*d*e**4*x*sqrt(d +
e*x) + 15*e**5*x**2*sqrt(d + e*x)) - 4*A*b*d*e/(15*d**2*e**3*sqrt(d + e*x)
+ 30*d*e**4*x*sqrt(d + e*x) + 15*e**5*x**2*sqrt(d + e*x)) - 10*A*b*e**2*x
/(15*d**2*e**3*sqrt(d + e*x) + 30*d*e**4*x*sqrt(d + e*x) + 15*e**5*x**2*sq
rt(d + e*x)) - 4*B*a*d*e/(15*d**2*e**3*sqrt(d + e*x) + 30*d*e**4*x*sqrt(d
+ e*x) + 15*e**5*x**2*sqrt(d + e*x)) - 10*B*a*e**2*x/(15*d**2*e**3*sqrt(d
+ e*x) + 30*d*e**4*x*sqrt(d + e*x) + 15*e**5*x**2*sqrt(d + e*x)) - 16*B*b*
d**2/(15*d**2*e**3*sqrt(d + e*x) + 30*d*e**4*x*sqrt(d + e*x) + 15*e**5*x**
2*sqrt(d + e*x)) - 40*B*b*d*e*x/(15*d**2*e**3*sqrt(d + e*x) + 30*d*e**4*x*
sqrt(d + e*x) + 15*e**5*x**2*sqrt(d + e*x)) - 30*B*b*e**2*x**2/(15*d**2*e
**3*sqrt(d + e*x) + 30*d*e**4*x*sqrt(d + e*x) + 15*e**5*x**2*sqrt(d + e*x))
, Ne(e, 0)), ((A*a*x + A*b*x**2/2 + B*a*x**2/2 + B*b*x**3/3)/d**(7/2), Tru
e))
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.89

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^{7/2}} dx = \frac{2(15(ex + d)^2 Bb + 3Bbd^2 + 3Aae^2 - 3(Ba + Ab)de - 5(2Bbd - (Ba + Ab)e)(ex + d))}{15(ex + d)^{\frac{5}{2}}e^3}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^(7/2),x, algorithm="maxima")`

output

`-2/15*(15*(e*x + d)^2*B*b + 3*B*b*d^2 + 3*A*a*e^2 - 3*(B*a + A*b)*d*e - 5*(2*B*b*d - (B*a + A*b)*e)*(e*x + d))/((e*x + d)^(5/2)*e^3)`
Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.99

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^{7/2}} dx = \frac{2(15(ex + d)^2 Bb - 10(ex + d)Bbd + 3Bbd^2 + 5(ex + d)Bae + 5(ex + d)Abe - 3Bade - 3Abde + 3Aae^2)}{15(ex + d)^{\frac{5}{2}}e^3}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^(7/2),x, algorithm="giac")`

output

`-2/15*(15*(e*x + d)^2*B*b - 10*(e*x + d)*B*b*d + 3*B*b*d^2 + 5*(e*x + d)*B*a*e + 5*(e*x + d)*A*b*e - 3*B*a*d*e - 3*A*b*d*e + 3*A*a*e^2)/((e*x + d)^(5/2)*e^3)`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.89

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^{7/2}} dx = \frac{(d + ex) \left(\frac{2Abe}{3} + \frac{2Bae}{3} - \frac{4Bbd}{3} \right) + 2Bb(d + ex)^2 + \frac{2Aae^2}{5} + \frac{2Bbd^2}{5} - \frac{2Abde}{5} - \frac{2Bade}{5}}{e^3 (d + ex)^{5/2}}$$

input `int(((A + B*x)*(a + b*x))/(d + e*x)^(7/2),x)`output `-((d + e*x)*((2*A*b*e)/3 + (2*B*a*e)/3 - (4*B*b*d)/3) + 2*B*b*(d + e*x)^2 + (2*A*a*e^2)/5 + (2*B*b*d^2)/5 - (2*A*b*d*e)/5 - (2*B*a*d*e)/5)/(e^3*(d + e*x)^(5/2))`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.00

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^{7/2}} dx = \frac{-2b^2e^2x^2 - \frac{4}{3}abe^2x - \frac{8}{3}b^2dex - \frac{2}{5}a^2e^2 - \frac{8}{15}abde - \frac{16}{15}b^2d^2}{\sqrt{ex + d}e^3(e^2x^2 + 2dex + d^2)}$$

input `int((b*x+a)*(B*x+A)/(e*x+d)^(7/2),x)`output `(2*(-3*a**2*e**2 - 4*a*b*d*e - 10*a*b*e**2*x - 8*b**2*d**2 - 20*b**2*d*e*x - 15*b**2*e**2*x**2))/(15*sqrt(d + e*x)*e**3*(d**2 + 2*d*e*x + e**2*x**2))`

3.131 $\int \frac{(a+bx)(A+Bx)}{(d+ex)^{9/2}} dx$

Optimal result	1249
Mathematica [A] (verified)	1249
Rubi [A] (verified)	1250
Maple [A] (verified)	1251
Fricas [A] (verification not implemented)	1252
Sympy [B] (verification not implemented)	1252
Maxima [A] (verification not implemented)	1253
Giac [A] (verification not implemented)	1254
Mupad [B] (verification not implemented)	1254
Reduce [B] (verification not implemented)	1255

Optimal result

Integrand size = 20, antiderivative size = 83

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{9/2}} dx = -\frac{2(bd-ae)(Bd-Ae)}{7e^3(d+ex)^{7/2}} + \frac{2(2bBd-Abe-aBe)}{5e^3(d+ex)^{5/2}} - \frac{2bB}{3e^3(d+ex)^{3/2}}$$

output -2/7*(-a*e+b*d)*(-A*e+B*d)/e^3/(e*x+d)^(7/2)+2/5*(-A*b*e-B*a*e+2*B*b*d)/e^3/(e*x+d)^(5/2)-2/3*b*B/e^3/(e*x+d)^(3/2)

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.84

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{9/2}} dx = \frac{2(3Abe(2d+7ex) + 3ae(2Bd+5Ae+7Bex) + bB(8d^2+28dex+35e^2x^2))}{105e^3(d+ex)^{7/2}}$$

input Integrate[((a + b*x)*(A + B*x))/(d + e*x)^(9/2), x]

output

$$\frac{(-2*(3*A*b*e*(2*d + 7*e*x) + 3*a*e*(2*B*d + 5*A*e + 7*B*e*x) + b*B*(8*d^2 + 28*d*e*x + 35*e^2*x^2)))/(105*e^3*(d + e*x)^(7/2))$$

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^{9/2}} dx$$

↓ 86

$$\int \left(\frac{aBe + Abe - 2bBd}{e^2(d + ex)^{7/2}} + \frac{(ae - bd)(Ae - Bd)}{e^2(d + ex)^{9/2}} + \frac{bB}{e^2(d + ex)^{5/2}} \right) dx$$

↓ 2009

$$\frac{2(-aBe - Abe + 2bBd)}{5e^3(d + ex)^{5/2}} - \frac{2(bd - ae)(Bd - Ae)}{7e^3(d + ex)^{7/2}} - \frac{2bB}{3e^3(d + ex)^{3/2}}$$

input

$$\text{Int}[(a + b*x)*(A + B*x)/(d + e*x)^(9/2), x]$$

output

$$\frac{(-2*(b*d - a*e)*(B*d - A*e))/(7*e^3*(d + e*x)^(7/2)) + (2*(2*b*B*d - A*b*e - a*B*e))/(5*e^3*(d + e*x)^(5/2)) - (2*b*B)/(3*e^3*(d + e*x)^(3/2))$$

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0]) || GeQ[n + p + 1, 0]) || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f]))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.77

method	result	size
pseudoelliptic	$\frac{\left((-70bx^2 - 42ax)B - 30A\left(\frac{7bx}{5} + a\right)\right)e^2 - 12\left(\left(\frac{14bx}{3} + a\right)B + Ab\right)de - 16bBd^2}{105(ex+d)^{\frac{7}{2}}e^3}$	64
gospers	$-\frac{2(35bBx^2e^2 + 21Axb^2e^2 + 21Bxa^2e^2 + 28Bxbde + 15Aae^2 + 6Abde + 6Bade + 8bBd^2)}{105(ex+d)^{\frac{7}{2}}e^3}$	73
trager	$-\frac{2(35bBx^2e^2 + 21Axb^2e^2 + 21Bxa^2e^2 + 28Bxbde + 15Aae^2 + 6Abde + 6Bade + 8bBd^2)}{105(ex+d)^{\frac{7}{2}}e^3}$	73
orering	$-\frac{2(35bBx^2e^2 + 21Axb^2e^2 + 21Bxa^2e^2 + 28Bxbde + 15Aae^2 + 6Abde + 6Bade + 8bBd^2)}{105(ex+d)^{\frac{7}{2}}e^3}$	73
derivativedivides	$-\frac{2(Abe + Bae - 2Bbd)}{5(ex+d)^{\frac{5}{2}}} - \frac{2(Aae^2 - Abde - Bade + bBd^2)}{7(ex+d)^{\frac{7}{2}}} - \frac{2bB}{3(ex+d)^{\frac{3}{2}}}$	75
default	$-\frac{2(Abe + Bae - 2Bbd)}{5(ex+d)^{\frac{5}{2}}} - \frac{2(Aae^2 - Abde - Bade + bBd^2)}{7(ex+d)^{\frac{7}{2}}} - \frac{2bB}{3(ex+d)^{\frac{3}{2}}}$	75

```
input int((b*x+a)*(B*x+A)/(e*x+d)^(9/2), x, method=_RETURNVERBOSE)
```

```
output 1/105*((( -70*b*x^2 - 42*a*x)*B - 30*A*(7/5*b*x+a))*e^2 - 12*((14/3*b*x+a)*B + A*b)*d*e - 16*b*B*d^2)/(e*x+d)^(7/2)/e^3
```


Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.36

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^{9/2}} dx = \frac{2(35 Bbe^2x^2 + 8 Bbd^2 + 15 Aae^2 + 6(Ba + Ab)de + 7(4 Bbde + 3(Ba + Ab)e^2)x)\sqrt{ex + d}}{105(e^7x^4 + 4de^6x^3 + 6d^2e^5x^2 + 4d^3e^4x + d^4e^3)}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^(9/2),x, algorithm="fricas")`

output `-2/105*(35*B*b*e^2*x^2 + 8*B*b*d^2 + 15*A*a*e^2 + 6*(B*a + A*b)*d*e + 7*(4*B*b*d*e + 3*(B*a + A*b)*e^2)*x)*sqrt(e*x + d)/(e^7*x^4 + 4*d*e^6*x^3 + 6*d^2*e^5*x^2 + 4*d^3*e^4*x + d^4*e^3)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 683 vs. 2(87) = 174.

Time = 0.72 (sec) , antiderivative size = 683, normalized size of antiderivative = 8.23

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^{9/2}} dx = \left\{ \begin{array}{l} -\frac{30Aae^2}{105d^3e^3\sqrt{d+ex}+315d^2e^4x\sqrt{d+ex}+315de^5x^2\sqrt{d+ex}+105e^6x^3\sqrt{d+ex}} - \frac{105d^3e^3\sqrt{d+ex}+315d^2e^4x}{Aax + \frac{Abx^2}{2} + \frac{Bax^2}{2} + \frac{Bbx^3}{3}}{d^{\frac{9}{2}}} \end{array} \right.$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)**(9/2),x)`

output

```
Piecewise((-30*A*a*e**2/(105*d**3*e**3*sqrt(d + e*x) + 315*d**2*e**4*x*sqrt(d + e*x) + 315*d**5*x**2*sqrt(d + e*x) + 105*e**6*x**3*sqrt(d + e*x)) - 12*A*b*d*e/(105*d**3*e**3*sqrt(d + e*x) + 315*d**2*e**4*x*sqrt(d + e*x) + 315*d**5*x**2*sqrt(d + e*x) + 105*e**6*x**3*sqrt(d + e*x)) - 42*A*b*e**2*x/(105*d**3*e**3*sqrt(d + e*x) + 315*d**2*e**4*x*sqrt(d + e*x) + 315*d**5*x**2*sqrt(d + e*x) + 105*e**6*x**3*sqrt(d + e*x)) - 12*B*a*d*e/(105*d**3*e**3*sqrt(d + e*x) + 315*d**2*e**4*x*sqrt(d + e*x) + 315*d**5*x**2*sqrt(d + e*x) + 105*e**6*x**3*sqrt(d + e*x)) - 42*B*a*e**2*x/(105*d**3*e**3*sqrt(d + e*x) + 315*d**2*e**4*x*sqrt(d + e*x) + 315*d**5*x**2*sqrt(d + e*x) + 105*e**6*x**3*sqrt(d + e*x)) - 16*B*b*d**2/(105*d**3*e**3*sqrt(d + e*x) + 315*d**2*e**4*x*sqrt(d + e*x) + 315*d**5*x**2*sqrt(d + e*x) + 105*e**6*x**3*sqrt(d + e*x)) - 56*B*b*d*e*x/(105*d**3*e**3*sqrt(d + e*x) + 315*d**2*e**4*x*sqrt(d + e*x) + 315*d**5*x**2*sqrt(d + e*x) + 105*e**6*x**3*sqrt(d + e*x)) - 70*B*b*e**2*x**2/(105*d**3*e**3*sqrt(d + e*x) + 315*d**2*e**4*x*sqrt(d + e*x) + 315*d**5*x**2*sqrt(d + e*x) + 105*e**6*x**3*sqrt(d + e*x)), Ne(e, 0)), ((A*a*x + A*b*x**2/2 + B*a*x**2/2 + B*b*x**3/3)/d**2*(9/2), True))
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.87

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^{9/2}} dx = \frac{2(35(ex + d)^2 Bb + 15 Bbd^2 + 15 Aae^2 - 15(Ba + Ab)de - 21(2 Bbd - (Ba + Ab)e)(ex + d))}{105(ex + d)^{7/2} e^3}$$

input

```
integrate((b*x+a)*(B*x+A)/(e*x+d)^(9/2),x, algorithm="maxima")
```

output

```
-2/105*(35*(e*x + d)^2*B*b + 15*B*b*d^2 + 15*A*a*e^2 - 15*(B*a + A*b)*d*e - 21*(2*B*b*d - (B*a + A*b)*e)*(e*x + d))/((e*x + d)^(7/2)*e^3)
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.96

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{9/2}} dx = \frac{2(35(ex+d)^2 Bb - 42(ex+d)Bbd + 15Bbd^2 + 21(ex+d)Bae + 21(ex+d)Abe - 15Bade - 15Abd)}{105(ex+d)^{7/2}e^3}$$

input `integrate((b*x+a)*(B*x+A)/(e*x+d)^(9/2),x, algorithm="giac")`

output

$$-2/105*(35*(e*x + d)^2*B*b - 42*(e*x + d)*B*b*d + 15*B*b*d^2 + 21*(e*x + d)*B*a*e + 21*(e*x + d)*A*b*e - 15*B*a*d*e - 15*A*b*d*e + 15*A*a*e^2)/((e*x + d)^(7/2)*e^3)$$

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.87

$$\int \frac{(a+bx)(A+Bx)}{(d+ex)^{9/2}} dx = \frac{(d+ex) \left(\frac{2Abe}{5} + \frac{2Bae}{5} - \frac{4Bbd}{5} \right) + \frac{2Bb(d+ex)^2}{3} + \frac{2Aae^2}{7} + \frac{2Bbd^2}{7} - \frac{2Abde}{7} - \frac{2Bade}{7}}{e^3(d+ex)^{7/2}}$$

input `int(((A + B*x)*(a + b*x))/(d + e*x)^(9/2),x)`

output

$$-((d + e*x)*((2*A*b*e)/5 + (2*B*a*e)/5 - (4*B*b*d)/5) + (2*B*b*(d + e*x)^2)/3 + (2*A*a*e^2)/7 + (2*B*b*d^2)/7 - (2*A*b*d*e)/7 - (2*B*a*d*e)/7)/(e^3*(d + e*x)^(7/2))$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.11

$$\int \frac{(a + bx)(A + Bx)}{(d + ex)^{9/2}} dx = \frac{-\frac{2}{3}b^2e^2x^2 - \frac{4}{5}abe^2x - \frac{8}{15}b^2dex - \frac{2}{7}a^2e^2 - \frac{8}{35}abde - \frac{16}{105}b^2d^2}{\sqrt{ex + d}e^3(e^3x^3 + 3de^2x^2 + 3d^2ex + d^3)}$$

input `int((b*x+a)*(B*x+A)/(e*x+d)^(9/2),x)`

output `(2*(-15*a**2*e**2 - 12*a*b*d*e - 42*a*b*e**2*x - 8*b**2*d**2 - 28*b**2*d*e*x - 35*b**2*e**2*x**2))/(105*sqrt(d + e*x)*e**3*(d**3 + 3*d**2*e*x + 3*d*e**2*x**2 + e**3*x**3))`

3.132 $\int (a + bx)^2(A + Bx)(d + ex)^{5/2} dx$

Optimal result	1256
Mathematica [A] (verified)	1256
Rubi [A] (verified)	1257
Maple [A] (verified)	1258
Fricas [B] (verification not implemented)	1259
Sympy [B] (verification not implemented)	1259
Maxima [A] (verification not implemented)	1260
Giac [B] (verification not implemented)	1261
Mupad [B] (verification not implemented)	1262
Reduce [B] (verification not implemented)	1262

Optimal result

Integrand size = 22, antiderivative size = 128

$$\int (a + bx)^2(A + Bx)(d + ex)^{5/2} dx = -\frac{2(bd - ae)^2(Bd - Ae)(d + ex)^{7/2}}{7e^4} + \frac{2(bd - ae)(3bBd - 2Abe - aBe)(d + ex)^{9/2}}{9e^4} - \frac{2b(3bBd - Abe - 2aBe)(d + ex)^{11/2}}{11e^4} + \frac{2b^2B(d + ex)^{13/2}}{13e^4}$$

output

```
-2/7*(-a*e+b*d)^2*(-A*e+B*d)*(e*x+d)^(7/2)/e^4+2/9*(-a*e+b*d)*(-2*A*b*e-B*a*e+3*B*b*d)*(e*x+d)^(9/2)/e^4-2/11*b*(-A*b*e-2*B*a*e+3*B*b*d)*(e*x+d)^(11/2)/e^4+2/13*b^2*B*(e*x+d)^(13/2)/e^4
```

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.08

$$\int (a + bx)^2(A + Bx)(d + ex)^{5/2} dx = \frac{2(d + ex)^{7/2} (143a^2e^2(-2Bd + 9Ae + 7Bex) + 26abe(11Ae(-2d + 7ex) + B(8d^2 - 28dex + 900$$

input `Integrate[(a + b*x)^2*(A + B*x)*(d + e*x)^(5/2),x]`

output
$$\frac{(2*(d + e*x)^{(7/2)}*(143*a^2*e^2*(-2*B*d + 9*A*e + 7*B*e*x) + 26*a*b*e*(11*A*e*(-2*d + 7*e*x) + B*(8*d^2 - 28*d*e*x + 63*e^2*x^2)) + b^2*(13*A*e*(8*d^2 - 28*d*e*x + 63*e^2*x^2) + B*(-48*d^3 + 168*d^2*e*x - 378*d*e^2*x^2 + 693*e^3*x^3))))}{(9009*e^4)}$$

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^2(A + Bx)(d + ex)^{5/2} dx$$

$$\downarrow 86$$

$$\int \left(\frac{b(d + ex)^{9/2}(2aBe + Abe - 3bBd)}{e^3} + \frac{(d + ex)^{7/2}(ae - bd)(aBe + 2Abe - 3bBd)}{e^3} + \frac{(d + ex)^{5/2}(ae - bd)^2(A + Bx)}{e^3} \right) dx$$

$$\downarrow 2009$$

$$-\frac{2b(d + ex)^{11/2}(-2aBe - Abe + 3bBd)}{11e^4} + \frac{2(d + ex)^{9/2}(bd - ae)(-aBe - 2Abe + 3bBd)}{9e^4} - \frac{2(d + ex)^{7/2}(bd - ae)^2(Bd - Ae)}{7e^4} + \frac{2b^2B(d + ex)^{13/2}}{13e^4}$$

input `Int[(a + b*x)^2*(A + B*x)*(d + e*x)^(5/2),x]`

output
$$(-2*(b*d - a*e)^2*(B*d - A*e)*(d + e*x)^{(7/2)})/(7*e^4) + (2*(b*d - a*e)*(3*b*B*d - 2*A*b*e - a*B*e)*(d + e*x)^{(9/2)})/(9*e^4) - (2*b*(3*b*B*d - A*b*e - 2*a*B*e)*(d + e*x)^{(11/2)})/(11*e^4) + (2*b^2*B*(d + e*x)^{(13/2)})/(13*e^4)$$

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0]) || GeQ[n + p + 1, 0]) || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f]))]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.72 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.94

method	result
pseudoelliptic	$2(ex+d)^{\frac{7}{2}} \left(\left(\left(\frac{7}{13} B x^3 + \frac{7}{11} A x^2 \right) b^2 + \frac{14ax \left(\frac{9Bx+A}{11} \right) b}{9} + a^2 \left(\frac{7Bx+A}{9} \right) \right) e^3 - \frac{4 \left(\frac{7 \left(\frac{27Bx+A}{26} \right) x b^2 + a \left(\frac{14Bx+A}{11} \right) b + \frac{a^2 B}{2} \right) d}{9} \right) \frac{1}{7e^4}$
derivativedivides	$\frac{\frac{2b^2B(ex+d)^{\frac{13}{2}}}{13} + \frac{2(2b(ae-db)B+b^2(Ae-Bd))(ex+d)^{\frac{11}{2}}}{11}}{e^4} + \frac{2((ae-db)^2B+2b(ae-db)(Ae-Bd))(ex+d)^{\frac{9}{2}}}{9} + \frac{2(ae-db)^2(Ae-Bd)(ex+d)^{\frac{7}{2}}}{7}$
default	$\frac{\frac{2b^2B(ex+d)^{\frac{13}{2}}}{13} + \frac{2(2b(ae-db)B+b^2(Ae-Bd))(ex+d)^{\frac{11}{2}}}{11}}{e^4} + \frac{2((ae-db)^2B+2b(ae-db)(Ae-Bd))(ex+d)^{\frac{9}{2}}}{9} + \frac{2(ae-db)^2(Ae-Bd)(ex+d)^{\frac{7}{2}}}{7}$
gospers	$\frac{2(ex+d)^{\frac{7}{2}} (693b^2B x^3 e^3 + 819A x^2 b^2 e^3 + 1638B x^2 a b e^3 - 378B x^2 b^2 d e^2 + 2002A x a b e^3 - 364A x b^2 d e^2 + 1001B x a^2 e^3 - 9009e^4)}{9009e^4}$
orering	$\frac{2(ex+d)^{\frac{7}{2}} (693b^2B x^3 e^3 + 819A x^2 b^2 e^3 + 1638B x^2 a b e^3 - 378B x^2 b^2 d e^2 + 2002A x a b e^3 - 364A x b^2 d e^2 + 1001B x a^2 e^3 - 9009e^4)}{9009e^4}$
trager	$2(693B b^2 e^6 x^6 + 819A b^2 e^6 x^5 + 1638B a b e^6 x^5 + 1701B b^2 d e^5 x^5 + 2002A a b e^6 x^4 + 2093A b^2 d e^5 x^4 + 1001B a^2 e^6 x^4 + 4189e^4)$
risch	$2(693B b^2 e^6 x^6 + 819A b^2 e^6 x^5 + 1638B a b e^6 x^5 + 1701B b^2 d e^5 x^5 + 2002A a b e^6 x^4 + 2093A b^2 d e^5 x^4 + 1001B a^2 e^6 x^4 + 4189e^4)$

```
input int((b*x+a)^2*(B*x+A)*(e*x+d)^(5/2), x, method=_RETURNVERBOSE)
```

```
output 2/7*(e*x+d)^(7/2)*(((7/13*B*x^3+7/11*A*x^2)*b^2+14/9*a*x*(9/11*B*x+A)*b+a^2*(7/9*B*x+A))*e^3-4/9*(7/11*(27/26*B*x+A)*x*b^2+a*(14/11*B*x+A)*b+1/2*a^2*B)*d*e^2+8/99*((21/13*B*x+A)*b+2*B*a)*b*d^2*e-16/429*b^2*B*d^3)/e^4
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 356 vs. $2(112) = 224$.

Time = 0.09 (sec) , antiderivative size = 356, normalized size of antiderivative = 2.78

$$\int (a + bx)^2 (A + Bx) (d + ex)^{5/2} dx = \frac{2(693 Bb^2 e^6 x^6 - 48 Bb^2 d^6 + 1287 Aa^2 d^3 e^3 + 104(2 Bab + Ab^2) d^5 e - 286(Ba^2 + 2 Aab) d^4 e^2}{e^4}$$

input `integrate((b*x+a)^2*(B*x+A)*(e*x+d)^(5/2),x, algorithm="fricas")`

output
$$\frac{2}{9009} (693 B^2 b^2 e^6 x^6 - 48 B^2 b^2 d^6 + 1287 A^2 a^2 d^3 e^3 + 104 (2 B^2 a b + A^2 b^2) d^5 e - 286 (B^2 a^2 + 2 A^2 a b) d^4 e^2 + 63 (27 B^2 b^2 d^5 e + 13 (2 B^2 a b + A^2 b^2) e^6) x^5 + 7 (159 B^2 b^2 d^2 e^4 + 299 (2 B^2 a b + A^2 b^2) d e^5 + 143 (B^2 a^2 + 2 A^2 a b) e^6) x^4 + (15 B^2 b^2 d^3 e^3 + 1287 A^2 a^2 e^6 + 1469 (2 B^2 a b + A^2 b^2) d^2 e^4 + 2717 (B^2 a^2 + 2 A^2 a b) d e^5) x^3 - 3 (6 B^2 b^2 d^4 e^2 - 1287 A^2 a^2 d e^5 - 13 (2 B^2 a b + A^2 b^2) d^3 e^3 - 715 (B^2 a^2 + 2 A^2 a b) d^2 e^4) x^2 + (24 B^2 b^2 d^5 e + 3861 A^2 a^2 d^2 e^4 - 52 (2 B^2 a b + A^2 b^2) d^4 e^2 + 143 (B^2 a^2 + 2 A^2 a b) d^3 e^3) x) \sqrt{e x + d} / e^4$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 857 vs. $2(129) = 258$.

Time = 0.50 (sec) , antiderivative size = 857, normalized size of antiderivative = 6.70

$$\int (a + bx)^2 (A + Bx) (d + ex)^{5/2} dx = \left\{ \frac{2Aa^2 d^3 \sqrt{d+ex}}{7e} + \frac{6Aa^2 d^2 x \sqrt{d+ex}}{7} + \frac{6Aa^2 d e x^2 \sqrt{d+ex}}{7} + \frac{2Aa^2 e^2 x^3 \sqrt{d+ex}}{7} - \frac{8Aabd^4 \sqrt{d+ex}}{63e^2} + \frac{4Aabd^3 x \sqrt{d+ex}}{63e} + d^{\frac{5}{2}} \left(Aa^2 x + Aabx^2 + \frac{Ab^2 x^3}{3} + \frac{Ba^2 x^2}{2} + \frac{2Babx^3}{3} + \frac{Bb^2 x^4}{4} \right) \right\}$$

input `integrate((b*x+a)**2*(B*x+A)*(e*x+d)**(5/2),x)`

output

```
Piecewise((2*A*a**2*d**3*sqrt(d + e*x)/(7*e) + 6*A*a**2*d**2*x*sqrt(d + e*x)/7 + 6*A*a**2*d*e*x**2*sqrt(d + e*x)/7 + 2*A*a**2*e**2*x**3*sqrt(d + e*x)/7 - 8*A*a*b*d**4*sqrt(d + e*x)/(63*e**2) + 4*A*a*b*d**3*x*sqrt(d + e*x)/(63*e) + 20*A*a*b*d**2*x**2*sqrt(d + e*x)/21 + 76*A*a*b*d*e*x**3*sqrt(d + e*x)/63 + 4*A*a*b*e**2*x**4*sqrt(d + e*x)/9 + 16*A*b**2*d**5*sqrt(d + e*x)/(693*e**3) - 8*A*b**2*d**4*x*sqrt(d + e*x)/(693*e**2) + 2*A*b**2*d**3*x**2*sqrt(d + e*x)/(231*e) + 226*A*b**2*d**2*x**3*sqrt(d + e*x)/693 + 46*A*b**2*d*e*x**4*sqrt(d + e*x)/99 + 2*A*b**2*e**2*x**5*sqrt(d + e*x)/11 - 4*B*a**2*d**4*sqrt(d + e*x)/(63*e**2) + 2*B*a**2*d**3*x*sqrt(d + e*x)/(63*e) + 10*B*a**2*d**2*x**2*sqrt(d + e*x)/21 + 38*B*a**2*d*e*x**3*sqrt(d + e*x)/63 + 2*B*a**2*e**2*x**4*sqrt(d + e*x)/9 + 32*B*a*b*d**5*sqrt(d + e*x)/(693*e**3) - 16*B*a*b*d**4*x*sqrt(d + e*x)/(693*e**2) + 4*B*a*b*d**3*x**2*sqrt(d + e*x)/(231*e) + 452*B*a*b*d**2*x**3*sqrt(d + e*x)/693 + 92*B*a*b*d*e*x**4*sqrt(d + e*x)/99 + 4*B*a*b*e**2*x**5*sqrt(d + e*x)/11 - 32*B*b**2*d**6*sqrt(d + e*x)/(3003*e**4) + 16*B*b**2*d**5*x*sqrt(d + e*x)/(3003*e**3) - 4*B*b**2*d**4*x**2*sqrt(d + e*x)/(1001*e**2) + 10*B*b**2*d**3*x**3*sqrt(d + e*x)/(3003*e) + 106*B*b**2*d**2*x**4*sqrt(d + e*x)/429 + 54*B*b**2*d*e*x**5*sqrt(d + e*x)/143 + 2*B*b**2*e**2*x**6*sqrt(d + e*x)/13, Ne(e, 0)), (d**(5/2)*(A*a**2*x + A*a*b*x**2 + A*b**2*x**3/3 + B*a**2*x**2/2 + 2*B*a*b*x**3/3 + B*b**2*x**4/4), True))
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.24

$$\int (a + bx)^2 (A + Bx) (d + ex)^{5/2} dx = \frac{2 \left(693 (ex + d)^{\frac{13}{2}} Bb^2 - 819 (3 Bb^2 d - (2 Bab + Ab^2)e)(ex + d)^{\frac{11}{2}} + 1001 (3 Bb^2 d^2 - 2 (2 Bab + Ab^2)d + Aa^2) (ex + d)^{\frac{9}{2}} - 1287 (Bb^2 d^3 - Aa^2 e^3 - (2 B a b + A b^2) d^2 e + (B a^2 + 2 A a b) d e^2) (ex + d)^{\frac{7}{2}} \right)}{e^4}$$

input

```
integrate((b*x+a)^2*(B*x+A)*(e*x+d)^(5/2),x, algorithm="maxima")
```

output

```
2/9009*(693*(e*x + d)^(13/2)*B*b^2 - 819*(3*B*b^2*d - (2*B*a*b + A*b^2)*e)*(e*x + d)^(11/2) + 1001*(3*B*b^2*d^2 - 2*(2*B*a*b + A*b^2)*d*e + (B*a^2 + 2*A*a*b)*e^2)*(e*x + d)^(9/2) - 1287*(B*b^2*d^3 - A*a^2*e^3 - (2*B*a*b + A*b^2)*d^2*e + (B*a^2 + 2*A*a*b)*d*e^2)*(e*x + d)^(7/2))/e^4
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1293 vs. $2(112) = 224$.

Time = 0.13 (sec) , antiderivative size = 1293, normalized size of antiderivative = 10.10

$$\int (a + bx)^2(A + Bx)(d + ex)^{5/2} dx = \text{Too large to display}$$

input `integrate((b*x+a)^2*(B*x+A)*(e*x+d)^(5/2),x, algorithm="giac")`

output

```
2/45045*(45045*sqrt(e*x + d)*A*a^2*d^3 + 45045*((e*x + d)^(3/2) - 3*sqrt(e
*x + d)*d)*A*a^2*d^2 + 15015*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*B*a^2*d
^3/e + 30030*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*A*a*b*d^3/e + 9009*(3*(
e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*A*a^2*d + 60
06*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*B*a*b
*d^3/e^2 + 3003*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x +
d)*d^2)*A*b^2*d^3/e^2 + 9009*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 1
5*sqrt(e*x + d)*d^2)*B*a^2*d^2/e + 18018*(3*(e*x + d)^(5/2) - 10*(e*x + d)
^(3/2)*d + 15*sqrt(e*x + d)*d^2)*A*a*b*d^2/e + 1287*(5*(e*x + d)^(7/2) - 2
1*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*A*a^2
+ 1287*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2
- 35*sqrt(e*x + d)*d^3)*B*b^2*d^3/e^3 + 7722*(5*(e*x + d)^(7/2) - 21*(e*x
+ d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*B*a*b*d^2/e
^2 + 3861*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d
^2 - 35*sqrt(e*x + d)*d^3)*A*b^2*d^2/e^2 + 3861*(5*(e*x + d)^(7/2) - 21*(e
*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*B*a^2*d/e
+ 7722*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2
- 35*sqrt(e*x + d)*d^3)*A*a*b*d/e + 429*(35*(e*x + d)^(9/2) - 180*(e*x +
d)^(7/2)*d + 378*(e*x + d)^(5/2)*d^2 - 420*(e*x + d)^(3/2)*d^3 + 315*sqrt(
e*x + d)*d^4)*B*b^2*d^2/e^3 + 858*(35*(e*x + d)^(9/2) - 180*(e*x + d)^(...
```

Mupad [B] (verification not implemented)

Time = 1.01 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.90

$$\int (a+bx)^2(A+Bx)(d+ex)^{5/2} dx = \frac{(d+ex)^{11/2}(2Ab^2e-6Bb^2d+4Babe)}{11e^4} + \frac{2Bb^2(d+ex)^{13/2}}{13e^4} + \frac{2(ae-bd)(d+ex)^{9/2}(2Abe+BAe-3Bbd)}{9e^4} + \frac{2(Ae-Bd)(ae-bd)^2(d+ex)^{7/2}}{7e^4}$$

input `int((A + B*x)*(a + b*x)^2*(d + e*x)^(5/2), x)`output `((d + e*x)^(11/2)*(2*A*b^2*e - 6*B*b^2*d + 4*B*a*b*e))/(11*e^4) + (2*B*b^2*(d + e*x)^(13/2))/(13*e^4) + (2*(a*e - b*d)*(d + e*x)^(9/2)*(2*A*b*e + B*a*e - 3*B*b*d))/(9*e^4) + (2*(A*e - B*d)*(a*e - b*d)^2*(d + e*x)^(7/2))/(7*e^4)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 284, normalized size of antiderivative = 2.22

$$\int (a+bx)^2(A+Bx)(d+ex)^{5/2} dx = \frac{2\sqrt{ex+d}(231b^3e^6x^6 + 819ab^2e^6x^5 + 567b^3de^5x^5 + 1001a^2be^6x^4 + 2093ab^2de^5x^4 + 371b^3d^2e^5x^3 + 104a^2b^2e^6x^3 + 143ab^2de^5x^3 + 1001a^2b^2e^6x^2 + 2145a^2b^2de^5x^2 + 2717a^2b^2de^5x^2 + 1001a^2b^2e^6x + 104ab^2d^2e^5x + 52ab^2d^2e^5x + 39ab^2d^2e^5x + 1469a^2b^2d^2e^5x^3 + 2093a^2b^2d^2e^5x^4 + 819a^2b^2e^6x^5 - 16b^3d^3e^6x^6 + 8b^3d^3e^6x^5 - 6b^3d^3e^6x^4 + 5b^3d^3e^6x^3 + 371b^3d^3e^6x^2 + 567b^3d^3e^6x + 231b^3d^3e^6)}{(3003e^4)}$$

input `int((b*x+a)^2*(B*x+A)*(e*x+d)^(5/2), x)`output `(2*sqrt(d + e*x)*(429*a**3*d**3*e**3 + 1287*a**3*d**2*e**4*x + 1287*a**3*d**e**5*x**2 + 429*a**3*e**6*x**3 - 286*a**2*b*d**4*e**2 + 143*a**2*b*d**3*e**3*x + 2145*a**2*b*d**2*e**4*x**2 + 2717*a**2*b*d*e**5*x**3 + 1001*a**2*b**e**6*x**4 + 104*a*b**2*d**5*e - 52*a*b**2*d**4*e**2*x + 39*a*b**2*d**3*e**3*x**2 + 1469*a*b**2*d**2*e**4*x**3 + 2093*a*b**2*d*e**5*x**4 + 819*a*b**2*e**6*x**5 - 16*b**3*d**6 + 8*b**3*d**5*e*x - 6*b**3*d**4*e**2*x**2 + 5*b**3*d**3*e**3*x**3 + 371*b**3*d**2*e**4*x**4 + 567*b**3*d*e**5*x**5 + 231*b**3*e**6*x**6))/(3003*e**4)`

3.133 $\int (a + bx)^2(A + Bx)(d + ex)^{3/2} dx$

Optimal result	1263
Mathematica [A] (verified)	1263
Rubi [A] (verified)	1264
Maple [A] (verified)	1265
Fricas [B] (verification not implemented)	1266
Sympy [A] (verification not implemented)	1266
Maxima [A] (verification not implemented)	1267
Giac [B] (verification not implemented)	1267
Mupad [B] (verification not implemented)	1268
Reduce [B] (verification not implemented)	1269

Optimal result

Integrand size = 22, antiderivative size = 128

$$\int (a + bx)^2(A + Bx)(d + ex)^{3/2} dx = -\frac{2(bd - ae)^2(Bd - Ae)(d + ex)^{5/2}}{5e^4} + \frac{2(bd - ae)(3bBd - 2Abe - aBe)(d + ex)^{7/2}}{7e^4} - \frac{2b(3bBd - Abe - 2aBe)(d + ex)^{9/2}}{9e^4} + \frac{2b^2B(d + ex)^{11/2}}{11e^4}$$

output

```
-2/5*(-a*e+b*d)^2*(-A*e+B*d)*(e*x+d)^(5/2)/e^4+2/7*(-a*e+b*d)*(-2*A*b*e-B*
a*e+3*B*b*d)*(e*x+d)^(7/2)/e^4-2/9*b*(-A*b*e-2*B*a*e+3*B*b*d)*(e*x+d)^(9/2
)/e^4+2/11*b^2*B*(e*x+d)^(11/2)/e^4
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.09

$$\int (a + bx)^2(A + Bx)(d + ex)^{3/2} dx = \frac{2(d + ex)^{5/2} (99a^2e^2(-2Bd + 7Ae + 5Bex) + 22abe(9Ae(-2d + 5ex) + B(8d^2 - 20dex + 35e^2)))}{3465e^4}$$

input `Integrate[(a + b*x)^2*(A + B*x)*(d + e*x)^(3/2),x]`

output
$$\frac{(2*(d + e*x)^{(5/2)}*(99*a^2*e^2*(-2*B*d + 7*A*e + 5*B*e*x) + 22*a*b*e*(9*A*e*(-2*d + 5*e*x) + B*(8*d^2 - 20*d*e*x + 35*e^2*x^2)) + b^2*(11*A*e*(8*d^2 - 20*d*e*x + 35*e^2*x^2) - 3*B*(16*d^3 - 40*d^2*e*x + 70*d*e^2*x^2 - 105*e^3*x^3))))}{(3465*e^4)}$$

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^2(A + Bx)(d + ex)^{3/2} dx$$

↓ 86

$$\int \left(\frac{b(d + ex)^{7/2}(2aBe + Abe - 3bBd)}{e^3} + \frac{(d + ex)^{5/2}(ae - bd)(aBe + 2Abe - 3bBd)}{e^3} + \frac{(d + ex)^{3/2}(ae - bd)^2(A + Bx)}{e^3} \right) dx$$

↓ 2009

$$\frac{2b(d + ex)^{9/2}(-2aBe - Abe + 3bBd)}{9e^4} + \frac{2(d + ex)^{7/2}(bd - ae)(-aBe - 2Abe + 3bBd)}{7e^4} - \frac{2(d + ex)^{5/2}(bd - ae)^2(Bd - Ae)}{5e^4} + \frac{2b^2B(d + ex)^{11/2}}{11e^4}$$

input `Int[(a + b*x)^2*(A + B*x)*(d + e*x)^(3/2),x]`

output
$$(-2*(b*d - a*e)^2*(B*d - A*e)*(d + e*x)^{(5/2)})/(5*e^4) + (2*(b*d - a*e)*(3*b*B*d - 2*A*b*e - a*B*e)*(d + e*x)^{(7/2)})/(7*e^4) - (2*b*(3*b*B*d - A*b*e - 2*a*B*e)*(d + e*x)^{(9/2)})/(9*e^4) + (2*b^2*B*(d + e*x)^{(11/2)})/(11*e^4)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 289 vs. $2(112) = 224$.

Time = 0.10 (sec) , antiderivative size = 289, normalized size of antiderivative = 2.26

$$\int (a + bx)^2 (A + Bx) (d + ex)^{3/2} dx = \frac{2(315 Bb^2 e^5 x^5 - 48 Bb^2 d^5 + 693 Aa^2 d^2 e^3 + 88(2 Bab + Ab^2) d^4 e - 198(Ba^2 + 2 Aab) d^3 e^2 + \dots}{e^4}$$

input `integrate((b*x+a)^2*(B*x+A)*(e*x+d)^(3/2),x, algorithm="fricas")`

output `2/3465*(315*B*b^2*e^5*x^5 - 48*B*b^2*d^5 + 693*A*a^2*d^2*e^3 + 88*(2*B*a*b + A*b^2)*d^4*e - 198*(B*a^2 + 2*A*a*b)*d^3*e^2 + 35*(12*B*b^2*d*e^4 + 11*(2*B*a*b + A*b^2)*e^5)*x^4 + 5*(3*B*b^2*d^2*e^3 + 110*(2*B*a*b + A*b^2)*d*e^4 + 99*(B*a^2 + 2*A*a*b)*e^5)*x^3 - 3*(6*B*b^2*d^3*e^2 - 231*A*a^2*e^5 - 11*(2*B*a*b + A*b^2)*d^2*e^3 - 264*(B*a^2 + 2*A*a*b)*d*e^4)*x^2 + (24*B*b^2*d^4*e + 1386*A*a^2*d*e^4 - 44*(2*B*a*b + A*b^2)*d^3*e^2 + 99*(B*a^2 + 2*A*a*b)*d^2*e^3)*x)*sqrt(e*x + d)/e^4`

Sympy [A] (verification not implemented)

Time = 1.11 (sec) , antiderivative size = 257, normalized size of antiderivative = 2.01

$$\int (a + bx)^2 (A + Bx) (d + ex)^{3/2} dx = \frac{2 \left(\frac{Bb^2(d+ex)^{\frac{11}{2}}}{11e^3} + \frac{(d+ex)^{\frac{9}{2}}(Ab^2e+2Babe-3Bb^2d)}{9e^3} + \frac{(d+ex)^{\frac{7}{2}}(2Aabe^2-2Ab^2de+Ba^2e^2-4Babde+3Bb^2d^2)}{7e^3} + \frac{(d+ex)^{\frac{5}{2}}(Aa^2e^3-2Aa^2e^2d+2Aabde-2Ab^2d^2)}{5e^3} \right)}{e} + \frac{d^{\frac{3}{2}} \left(Aa^2x + \frac{Bb^2x^4}{4} + \frac{x^3(Ab^2+2Bab)}{3} + \frac{x^2(2Aab+Ba^2)}{2} \right)}{e}$$

input `integrate((b*x+a)**2*(B*x+A)*(e*x+d)**(3/2),x)`

output

```
Piecewise((2*(B*b**2*(d + e*x)**(11/2)/(11*e**3) + (d + e*x)**(9/2)*(A*b**
2*e + 2*B*a*b*e - 3*B*b**2*d)/(9*e**3) + (d + e*x)**(7/2)*(2*A*a*b*e**2 -
2*A*b**2*d*e + B*a**2*e**2 - 4*B*a*b*d*e + 3*B*b**2*d**2)/(7*e**3) + (d +
e*x)**(5/2)*(A*a**2*e**3 - 2*A*a*b*d*e**2 + A*b**2*d**2*e - B*a**2*d*e**2
+ 2*B*a*b*d**2*e - B*b**2*d**3)/(5*e**3))/e, Ne(e, 0)), (d**(3/2)*(A*a**2*
x + B*b**2*x**4/4 + x**3*(A*b**2 + 2*B*a*b)/3 + x**2*(2*A*a*b + B*a**2)/2
, True))
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.24

$$\int (a + bx)^2 (A + Bx)(d + ex)^{3/2} dx = \frac{2 \left(315 (ex + d)^{\frac{11}{2}} Bb^2 - 385 (3 Bb^2 d - (2 Bab + Ab^2)e)(ex + d)^{\frac{9}{2}} + 495 (3 Bb^2 d^2 - 2 (2 Bab + Ab^2)d + Aa^2) (ex + d)^{\frac{7}{2}} - 693 (Bb^2 d^3 - Aa^2 e^3 - (2 B a b + Ab^2) d^2 e + (B a^2 + 2 A a b) d e^2) (ex + d)^{\frac{5}{2}} \right)}{e^4}$$

input

```
integrate((b*x+a)^2*(B*x+A)*(e*x+d)^(3/2),x, algorithm="maxima")
```

output

```
2/3465*(315*(e*x + d)^(11/2)*B*b^2 - 385*(3*B*b^2*d - (2*B*a*b + A*b^2)*e)
*(e*x + d)^(9/2) + 495*(3*B*b^2*d^2 - 2*(2*B*a*b + A*b^2)*d*e + (B*a^2 + 2
*A*a*b)*e^2)*(e*x + d)^(7/2) - 693*(B*b^2*d^3 - A*a^2*e^3 - (2*B*a*b + A*b
^2)*d^2*e + (B*a^2 + 2*A*a*b)*d*e^2)*(e*x + d)^(5/2))/e^4
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 854 vs. 2(112) = 224.

Time = 0.13 (sec) , antiderivative size = 854, normalized size of antiderivative = 6.67

$$\int (a + bx)^2 (A + Bx)(d + ex)^{3/2} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^2*(B*x+A)*(e*x+d)^(3/2),x, algorithm="giac")
```


output

```

2/3465*(3465*sqrt(e*x + d)*A*a^2*d^2 + 2310*((e*x + d)^(3/2) - 3*sqrt(e*x
+ d)*d)*A*a^2*d + 1155*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*B*a^2*d^2/e +
2310*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*A*a*b*d^2/e + 231*(3*(e*x + d)
^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*A*a^2 + 462*(3*(e*x
+ d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*B*a*b*d^2/e^2 +
231*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*A*b
^2*d^2/e^2 + 462*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x +
d)*d^2)*B*a^2*d/e + 924*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sq
rt(e*x + d)*d^2)*A*a*b*d/e + 99*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d +
35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*B*b^2*d^2/e^3 + 396*(5*(e
x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x
+ d)*d^3)*B*a*b*d/e^2 + 198*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 3
5*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*A*b^2*d/e^2 + 99*(5*(e*x + d)
^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)
*d^3)*B*a^2/e + 198*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x +
d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*A*a*b/e + 22*(35*(e*x + d)^(9/2) - 18
0*(e*x + d)^(7/2)*d + 378*(e*x + d)^(5/2)*d^2 - 420*(e*x + d)^(3/2)*d^3 +
315*sqrt(e*x + d)*d^4)*B*b^2*d/e^3 + 22*(35*(e*x + d)^(9/2) - 180*(e*x + d)
^(7/2)*d + 378*(e*x + d)^(5/2)*d^2 - 420*(e*x + d)^(3/2)*d^3 + 315*sqrt(e
*x + d)*d^4)*B*a*b/e^2 + 11*(35*(e*x + d)^(9/2) - 180*(e*x + d)^(7/2)*d...

```

Mupad [B] (verification not implemented)

Time = 0.98 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.90

$$\begin{aligned}
 \int (a + bx)^2 (A + Bx)(d + ex)^{3/2} dx &= \frac{(d + ex)^{9/2} (2Ab^2e - 6Bb^2d + 4Babe)}{9e^4} \\
 &+ \frac{2Bb^2(d + ex)^{11/2}}{11e^4} + \frac{2(ae - bd)(d + ex)^{7/2} (2Abe + Bae - 3Bbd)}{7e^4} \\
 &+ \frac{2(Ae - Bd)(ae - bd)^2(d + ex)^{5/2}}{5e^4}
 \end{aligned}$$

input

```
int((A + B*x)*(a + b*x)^2*(d + e*x)^(3/2), x)
```

output

```

((d + e*x)^(9/2)*(2*A*b^2*e - 6*B*b^2*d + 4*B*a*b*e))/(9*e^4) + (2*B*b^2*(
d + e*x)^(11/2))/(11*e^4) + (2*(a*e - b*d)*(d + e*x)^(7/2)*(2*A*b*e + B*a*
e - 3*B*b*d))/(7*e^4) + (2*(A*e - B*d)*(a*e - b*d)^2*(d + e*x)^(5/2))/(5*
e^4)

```

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.77

$$\int (a + bx)^2 (A + Bx) (d + ex)^{3/2} dx = \frac{2\sqrt{ex + d} (105b^3e^5x^5 + 385ab^2e^5x^4 + 140b^3de^4x^4 + 495a^2be^5x^3 + 550ab^2de^4x^3 + 5b^3d^2e^3x^3)}{1155e^4}$$

input `int((b*x+a)^2*(B*x+A)*(e*x+d)^(3/2),x)`output `(2*sqrt(d + e*x)*(231*a**3*d**2*e**3 + 462*a**3*d*e**4*x + 231*a**3*e**5*x**2 - 198*a**2*b*d**3*e**2 + 99*a**2*b*d**2*e**3*x + 792*a**2*b*d*e**4*x**2 + 495*a**2*b*e**5*x**3 + 88*a*b**2*d**4*e - 44*a*b**2*d**3*e**2*x + 33*a*b**2*d**2*e**3*x**2 + 550*a*b**2*d*e**4*x**3 + 385*a*b**2*e**5*x**4 - 16*b**3*d**5 + 8*b**3*d**4*e*x - 6*b**3*d**3*e**2*x**2 + 5*b**3*d**2*e**3*x**3 + 140*b**3*d*e**4*x**4 + 105*b**3*e**5*x**5))/(1155*e**4)`

3.134 $\int (a + bx)^2 (A + Bx) \sqrt{d + ex} dx$

Optimal result	1270
Mathematica [A] (verified)	1271
Rubi [A] (verified)	1271
Maple [A] (verified)	1272
Fricas [A] (verification not implemented)	1273
Sympy [A] (verification not implemented)	1273
Maxima [A] (verification not implemented)	1274
Giac [B] (verification not implemented)	1274
Mupad [B] (verification not implemented)	1275
Reduce [B] (verification not implemented)	1276

Optimal result

Integrand size = 22, antiderivative size = 128

$$\int (a + bx)^2 (A + Bx) \sqrt{d + ex} dx = -\frac{2(bd - ae)^2 (Bd - Ae)(d + ex)^{3/2}}{3e^4} + \frac{2(bd - ae)(3bBd - 2Abe - aBe)(d + ex)^{5/2}}{5e^4} - \frac{2b(3bBd - Abe - 2aBe)(d + ex)^{7/2}}{7e^4} + \frac{2b^2 B(d + ex)^{9/2}}{9e^4}$$

output

```
-2/3*(-a*e+b*d)^2*(-A*e+B*d)*(e*x+d)^(3/2)/e^4+2/5*(-a*e+b*d)*(-2*A*b*e-B*
a*e+3*B*b*d)*(e*x+d)^(5/2)/e^4-2/7*b*(-A*b*e-2*B*a*e+3*B*b*d)*(e*x+d)^(7/2
)/e^4+2/9*b^2*B*(e*x+d)^(9/2)/e^4
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.08

$$\int (a + bx)^2 (A + Bx) \sqrt{d + ex} dx$$

$$= \frac{2(d + ex)^{3/2} (21a^2e^2(-2Bd + 5Ae + 3Bex) + 6abe(7Ae(-2d + 3ex) + B(8d^2 - 12dex + 15e^2x^2)) + b^2(8d^2 - 12dex + 15e^2x^2)) + B(-16d^3 + 24d^2ex - 30de^2x^2 + 35e^3x^3))}{315e^4}$$

input `Integrate[(a + b*x)^2*(A + B*x)*Sqrt[d + e*x],x]`

output $(2*(d + e*x)^{(3/2)}*(21*a^2*e^2*(-2*B*d + 5*A*e + 3*B*e*x) + 6*a*b*e*(7*A*e*(-2*d + 3*e*x) + B*(8*d^2 - 12*d*e*x + 15*e^2*x^2)) + b^2*(3*A*e*(8*d^2 - 12*d*e*x + 15*e^2*x^2) + B*(-16*d^3 + 24*d^2*e*x - 30*d*e^2*x^2 + 35*e^3*x^3)))/(315*e^4)$

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^2 (A + Bx) \sqrt{d + ex} dx$$

$$\downarrow 86$$

$$\int \left(\frac{b(d + ex)^{5/2}(2aBe + Abe - 3bBd)}{e^3} + \frac{(d + ex)^{3/2}(ae - bd)(aBe + 2Abe - 3bBd)}{e^3} + \frac{\sqrt{d + ex}(ae - bd)^2(Ae - b^2)}{e^3} \right) dx$$

$$\downarrow 2009$$

$$-\frac{2b(d + ex)^{7/2}(-2aBe - Abe + 3bBd)}{7e^4} + \frac{2(d + ex)^{5/2}(bd - ae)(-aBe - 2Abe + 3bBd)}{5e^4} - \frac{2(d + ex)^{3/2}(bd - ae)^2(Bd - Ae)}{3e^4} + \frac{2b^2B(d + ex)^{9/2}}{9e^4}$$

input `int((b*x+a)^2*(B*x+A)*(e*x+d)^(1/2),x,method=_RETURNVERBOSE)`

output
$$\frac{2}{3}(e*x+d)^{(3/2)} * ((3/7*x^2*(7/9*B*x+A)*b^2+6/5*a*x*(5/7*B*x+A)*b+a^2*(3/5*B*x+A))*e^3-4/5*(3/7*(5/6*B*x+A)*x*b^2+a*(6/7*B*x+A)*b+1/2*a^2*B)*d*e^2+8/35*b*((B*x+A)*b+2*B*a)*d^2*e-16/105*b^2*B*d^3)/e^4$$

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.72

$$\int (a + bx)^2 (A + Bx) \sqrt{d + ex} dx$$

$$= \frac{2(35 Bb^2 e^4 x^4 - 16 Bb^2 d^4 + 105 Aa^2 de^3 + 24(2 Bab + Ab^2)d^3 e - 42(Ba^2 + 2 Aab)d^2 e^2 + 5(Bb^2 de^3 + 9$$

input `integrate((b*x+a)^2*(B*x+A)*(e*x+d)^(1/2),x, algorithm="fricas")`

output
$$\frac{2}{315}(35*B*b^2*e^4*x^4 - 16*B*b^2*d^4 + 105*A*a^2*d*e^3 + 24*(2*B*a*b + A*b^2)*d^3*e - 42*(B*a^2 + 2*A*a*b)*d^2*e^2 + 5*(B*b^2*d*e^3 + 9*(2*B*a*b + A*b^2)*e^4)*x^3 - 3*(2*B*b^2*d^2*e^2 - 3*(2*B*a*b + A*b^2)*d*e^3 - 21*(B*a^2 + 2*A*a*b)*e^4)*x^2 + (8*B*b^2*d^3*e + 105*A*a^2*e^4 - 12*(2*B*a*b + A*b^2)*d^2*e^2 + 21*(B*a^2 + 2*A*a*b)*d*e^3)*x)*sqrt(e*x + d)/e^4$$

Sympy [A] (verification not implemented)

Time = 1.28 (sec) , antiderivative size = 257, normalized size of antiderivative = 2.01

$$\int (a + bx)^2 (A + Bx) \sqrt{d + ex} dx$$

$$= \left\{ \frac{2 \left(\frac{Bb^2(d+ex)^{\frac{9}{2}}}{9e^3} + \frac{(d+ex)^{\frac{7}{2}}(Ab^2e+2Babe-3Bb^2d)}{7e^3} + \frac{(d+ex)^{\frac{5}{2}}(2Aabe^2-2Ab^2de+Ba^2e^2-4Babde+3Bb^2d^2)}{5e^3} + \frac{(d+ex)^{\frac{3}{2}}(Aa^2e^3-2Aabde^2+Ab^2d^2e-Bb^2d^2)}{3e^3} \right)}{e} \right.$$

$$\left. \sqrt{d} \left(Aa^2x + \frac{Bb^2x^4}{4} + \frac{x^3(Ab^2+2Bab)}{3} + \frac{x^2(2Aab+Ba^2)}{2} \right) \right\}$$

input `integrate((b*x+a)**2*(B*x+A)*(e*x+d)**(1/2),x)`

output

```
Piecewise((2*(B*b**2*(d + e*x)**(9/2)/(9*e**3) + (d + e*x)**(7/2)*(A*b**2*
e + 2*B*a*b*e - 3*B*b**2*d)/(7*e**3) + (d + e*x)**(5/2)*(2*A*a*b*e**2 - 2*
A*b**2*d*e + B*a**2*e**2 - 4*B*a*b*d*e + 3*B*b**2*d**2)/(5*e**3) + (d + e*
x)**(3/2)*(A*a**2*e**3 - 2*A*a*b*d*e**2 + A*b**2*d**2*e - B*a**2*d*e**2 +
2*B*a*b*d**2*e - B*b**2*d**3)/(3*e**3))/e, Ne(e, 0)), (sqrt(d)*(A*a**2*x +
B*b**2*x**4/4 + x**3*(A*b**2 + 2*B*a*b)/3 + x**2*(2*A*a*b + B*a**2)/2), T
rue))
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.24

$$\int (a + bx)^2 (A + Bx) \sqrt{d + ex} dx$$

$$= \frac{2 \left(35 (ex + d)^{\frac{9}{2}} Bb^2 - 45 (3 Bb^2 d - (2 Bab + Ab^2)e) (ex + d)^{\frac{7}{2}} + 63 (3 Bb^2 d^2 - 2 (2 Bab + Ab^2)de + (Ba \right.}{315 e}$$

input

```
integrate((b*x+a)^2*(B*x+A)*(e*x+d)^(1/2),x, algorithm="maxima")
```

output

```
2/315*(35*(e*x + d)^(9/2)*B*b^2 - 45*(3*B*b^2*d - (2*B*a*b + A*b^2)*e)*(e*
x + d)^(7/2) + 63*(3*B*b^2*d^2 - 2*(2*B*a*b + A*b^2)*d*e + (B*a^2 + 2*A*a*
b)*e^2)*(e*x + d)^(5/2) - 105*(B*b^2*d^3 - A*a^2*e^3 - (2*B*a*b + A*b^2)*d
^2*e + (B*a^2 + 2*A*a*b)*d*e^2)*(e*x + d)^(3/2))/e^4
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 486 vs. 2(112) = 224.

Time = 0.12 (sec) , antiderivative size = 486, normalized size of antiderivative = 3.80

$$\int (a + bx)^2 (A + Bx) \sqrt{d + ex} dx$$

$$= \frac{2 \left(315 \sqrt{ex + d} Aa^2 d + 105 \left((ex + d)^{\frac{3}{2}} - 3 \sqrt{ex + dd} \right) Aa^2 + \frac{105 \left((ex+d)^{\frac{3}{2}} - 3 \sqrt{ex+dd} \right) Ba^2 d}{e} + \frac{210 \left((ex+d)^{\frac{3}{2}} - 3 \sqrt{ex+dd} \right) Aa^2 d}{e} \right)}{e^4}$$

input

```
integrate((b*x+a)^2*(B*x+A)*(e*x+d)^(1/2),x, algorithm="giac")
```

output

```

2/315*(315*sqrt(e*x + d)*A*a^2*d + 105*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*
d)*A*a^2 + 105*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*B*a^2*d/e + 210*((e*x
+ d)^(3/2) - 3*sqrt(e*x + d)*d)*A*a*b*d/e + 42*(3*(e*x + d)^(5/2) - 10*(e
*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*B*a*b*d/e^2 + 21*(3*(e*x + d)^(5/2
) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*A*b^2*d/e^2 + 21*(3*(e*x
+ d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*B*a^2/e + 42*(3*
(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*A*a*b/e + 9
*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*s
qrt(e*x + d)*d^3)*B*b^2*d/e^3 + 18*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)
*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*B*a*b/e^2 + 9*(5*(e*x
+ d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x +
d)*d^3)*A*b^2/e^2 + (35*(e*x + d)^(9/2) - 180*(e*x + d)^(7/2)*d + 378*(e*
x + d)^(5/2)*d^2 - 420*(e*x + d)^(3/2)*d^3 + 315*sqrt(e*x + d)*d^4)*B*b^2/
e^3)/e

```

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.90

$$\begin{aligned}
\int (a + bx)^2 (A + Bx) \sqrt{d + ex} dx = & \frac{(d + ex)^{7/2} (2Ab^2e - 6Bb^2d + 4Babe)}{7e^4} \\
& + \frac{2Bb^2(d + ex)^{9/2}}{9e^4} \\
& + \frac{2(ae - bd)(d + ex)^{5/2} (2Abe + Bae - 3Bbd)}{5e^4} \\
& + \frac{2(Ae - Bd)(ae - bd)^2 (d + ex)^{3/2}}{3e^4}
\end{aligned}$$

input

```
int((A + B*x)*(a + b*x)^2*(d + e*x)^(1/2),x)
```

output

```

((d + e*x)^(7/2)*(2*A*b^2*e - 6*B*b^2*d + 4*B*a*b*e))/(7*e^4) + (2*B*b^2*(
d + e*x)^(9/2))/(9*e^4) + (2*(a*e - b*d)*(d + e*x)^(5/2)*(2*A*b*e + B*a*e
- 3*B*b*d))/(5*e^4) + (2*(A*e - B*d)*(a*e - b*d)^2*(d + e*x)^(3/2))/(3*e^4
)

```


Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.31

$$\int (a + bx)^2 (A + Bx) \sqrt{d + ex} dx$$

$$= \frac{2\sqrt{ex + d} (35b^3 e^4 x^4 + 135a b^2 e^4 x^3 + 5b^3 d e^3 x^3 + 189a^2 b e^4 x^2 + 27a b^2 d e^3 x^2 - 6b^3 d^2 e^2 x^2 + 105a^3 e^4 x + 6b^3 d^2 e^2 x^2)}{315e^4}$$

input `int((b*x+a)^2*(B*x+A)*(e*x+d)^(1/2),x)`output `(2*sqrt(d + e*x)*(105*a**3*d*e**3 + 105*a**3*e**4*x - 126*a**2*b*d**2*e**2 + 63*a**2*b*d*e**3*x + 189*a**2*b*e**4*x**2 + 72*a*b**2*d**3*e - 36*a*b**2*d**2*e**2*x + 27*a*b**2*d*e**3*x**2 + 135*a*b**2*e**4*x**3 - 16*b**3*d**4 + 8*b**3*d**3*e*x - 6*b**3*d**2*e**2*x**2 + 5*b**3*d*e**3*x**3 + 35*b**3*e**4*x**4))/(315*e**4)`

3.135 $\int \frac{(a+bx)^2(A+Bx)}{\sqrt{d+ex}} dx$

Optimal result	1277
Mathematica [A] (verified)	1277
Rubi [A] (verified)	1278
Maple [A] (verified)	1279
Fricas [A] (verification not implemented)	1280
Sympy [A] (verification not implemented)	1280
Maxima [A] (verification not implemented)	1281
Giac [A] (verification not implemented)	1281
Mupad [B] (verification not implemented)	1282
Reduce [B] (verification not implemented)	1282

Optimal result

Integrand size = 22, antiderivative size = 126

$$\int \frac{(a+bx)^2(A+Bx)}{\sqrt{d+ex}} dx = -\frac{2(bd-ae)^2(Bd-Ae)\sqrt{d+ex}}{e^4} + \frac{2(bd-ae)(3bBd-2Abe-aBe)(d+ex)^{3/2}}{3e^4} - \frac{2b(3bBd-Abe-2aBe)(d+ex)^{5/2}}{5e^4} + \frac{2b^2B(d+ex)^{7/2}}{7e^4}$$

output

```
-2*(-a*e+b*d)^2*(-A*e+B*d)*(e*x+d)^(1/2)/e^4+2/3*(-a*e+b*d)*(-2*A*b*e-B*a*
e+3*B*b*d)*(e*x+d)^(3/2)/e^4-2/5*b*(-A*b*e-2*B*a*e+3*B*b*d)*(e*x+d)^(5/2)/
e^4+2/7*b^2*B*(e*x+d)^(7/2)/e^4
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.09

$$\int \frac{(a+bx)^2(A+Bx)}{\sqrt{d+ex}} dx = \frac{2\sqrt{d+ex}(35a^2e^2(-2Bd+3Ae+Bex) + 14abe(5Ae(-2d+ex) + B(8d^2-4dex+3e^2x^2)) + b^2(7Ae(8d^2-4dex+3e^2x^2) + B(8d^2-4dex+3e^2x^2)))}{105e^4}$$

input `Integrate[((a + b*x)^2*(A + B*x))/Sqrt[d + e*x],x]`

output `(2*Sqrt[d + e*x]*(35*a^2*e^2*(-2*B*d + 3*A*e + B*e*x) + 14*a*b*e*(5*A*e*(-2*d + e*x) + B*(8*d^2 - 4*d*e*x + 3*e^2*x^2)) + b^2*(7*A*e*(8*d^2 - 4*d*e*x + 3*e^2*x^2) - 3*B*(16*d^3 - 8*d^2*e*x + 6*d*e^2*x^2 - 5*e^3*x^3)))/(105*e^4)`

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^2(A + Bx)}{\sqrt{d + ex}} dx$$

↓ 86

$$\int \left(\frac{b(d + ex)^{3/2}(2aBe + Abe - 3bBd)}{e^3} + \frac{\sqrt{d + ex}(ae - bd)(aBe + 2Abe - 3bBd)}{e^3} + \frac{(ae - bd)^2(Ae - Bd)}{e^3\sqrt{d + ex}} + \frac{b^2}{e^3} \right) dx$$

↓ 2009

$$-\frac{2b(d + ex)^{5/2}(-2aBe - Abe + 3bBd)}{5e^4} + \frac{2(d + ex)^{3/2}(bd - ae)(-aBe - 2Abe + 3bBd)}{3e^4} - \frac{2\sqrt{d + ex}(bd - ae)^2(Bd - Ae)}{e^4} + \frac{2b^2B(d + ex)^{7/2}}{7e^4}$$

input `Int[((a + b*x)^2*(A + B*x))/Sqrt[d + e*x],x]`

output `(-2*(b*d - a*e)^2*(B*d - A*e)*Sqrt[d + e*x])/e^4 + (2*(b*d - a*e)*(3*b*B*d - 2*A*b*e - a*B*e)*(d + e*x)^(3/2))/(3*e^4) - (2*b*(3*b*B*d - A*b*e - 2*a*B*e)*(d + e*x)^(5/2))/(5*e^4) + (2*b^2*B*(d + e*x)^(7/2))/(7*e^4)`

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.68 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.93

method	result
pseudoelliptic	$\frac{2\sqrt{ex+d} \left(\left(\frac{x^2(5Bx+A)b^2}{5} + \frac{2(3Bx+A)axb}{3} + a^2\left(\frac{Bx}{3}+A\right) \right) e^3 - \frac{4 \left(\frac{(9Bx+A)x b^2}{5} + a\left(\frac{2Bx}{5}+A\right)b + \frac{a^2B}{2} \right) d e^2}{3} + \frac{8b \left(\frac{3Bx}{7} \right)}{3} \right)}{e^4}$
derivativedivides	$\frac{\frac{2b^2B(e^2x+d)^{\frac{7}{2}}}{7} + \frac{2(2b(ae-db)B+b^2(Ae-Bd))(e^2x+d)^{\frac{5}{2}}}{5} + \frac{2((ae-db)^2B+2b(ae-db)(Ae-Bd))(e^2x+d)^{\frac{3}{2}}}{3} + 2(ae-db)^2(Ae-Bd)\sqrt{e^2x+d}}{e^4}$
default	$\frac{\frac{2b^2B(e^2x+d)^{\frac{7}{2}}}{7} + \frac{2(2b(ae-db)B+b^2(Ae-Bd))(e^2x+d)^{\frac{5}{2}}}{5} + \frac{2((ae-db)^2B+2b(ae-db)(Ae-Bd))(e^2x+d)^{\frac{3}{2}}}{3} + 2(ae-db)^2(Ae-Bd)\sqrt{e^2x+d}}{e^4}$
gospers	$\frac{2\sqrt{ex+d} (15b^2B x^3 e^3 + 21A x^2 b^2 e^3 + 42B x^2 a b e^3 - 18B x^2 b^2 d e^2 + 70A x a b e^3 - 28A x b^2 d e^2 + 35B x a^2 e^3 - 56B x a b d e^2 + 105e^4)}{105e^4}$
trager	$\frac{2\sqrt{ex+d} (15b^2B x^3 e^3 + 21A x^2 b^2 e^3 + 42B x^2 a b e^3 - 18B x^2 b^2 d e^2 + 70A x a b e^3 - 28A x b^2 d e^2 + 35B x a^2 e^3 - 56B x a b d e^2 + 105e^4)}{105e^4}$
risch	$\frac{2\sqrt{ex+d} (15b^2B x^3 e^3 + 21A x^2 b^2 e^3 + 42B x^2 a b e^3 - 18B x^2 b^2 d e^2 + 70A x a b e^3 - 28A x b^2 d e^2 + 35B x a^2 e^3 - 56B x a b d e^2 + 105e^4)}{105e^4}$
orering	$\frac{2\sqrt{ex+d} (15b^2B x^3 e^3 + 21A x^2 b^2 e^3 + 42B x^2 a b e^3 - 18B x^2 b^2 d e^2 + 70A x a b e^3 - 28A x b^2 d e^2 + 35B x a^2 e^3 - 56B x a b d e^2 + 105e^4)}{105e^4}$

```
input int((b*x+a)^2*(B*x+A)/(e*x+d)^(1/2), x, method=_RETURNVERBOSE)
```

```
output 2*(e*x+d)^(1/2)*((1/5*x^2*(5/7*B*x+A)*b^2+2/3*(3/5*B*x+A)*a*x*b+a^2*(1/3*B*x+A))*e^3-4/3*(1/5*(9/14*B*x+A)*x*b^2+a*(2/5*B*x+A)*b+1/2*a^2*B)*d*e^2+8/15*b*((3/7*B*x+A)*b+2*B*a)*d^2*e-16/35*b^2*B*d^3)/e^4
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.23

$$\int \frac{(a+bx)^2(A+Bx)}{\sqrt{d+ex}} dx$$

$$= \frac{2(15Bb^2e^3x^3 - 48Bb^2d^3 + 105Aa^2e^3 + 56(2Bab + Ab^2)d^2e - 70(Ba^2 + 2Aab)de^2 - 3(6Bb^2de^2 - 7Aa^2e^3))}{105e^4}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^(1/2),x, algorithm="fricas")`output `2/105*(15*B*b^2*e^3*x^3 - 48*B*b^2*d^3 + 105*A*a^2*e^3 + 56*(2*B*a*b + A*b^2)*d^2*e - 70*(B*a^2 + 2*A*a*b)*d*e^2 - 3*(6*B*b^2*d*e^2 - 7*(2*B*a*b + A*b^2)*e^3)*x^2 + (24*B*b^2*d^2*e - 28*(2*B*a*b + A*b^2)*d*e^2 + 35*(B*a^2 + 2*A*a*b)*e^3)*x)*sqrt(e*x + d)/e^4`**Sympy [A] (verification not implemented)**

Time = 0.97 (sec) , antiderivative size = 255, normalized size of antiderivative = 2.02

$$\int \frac{(a+bx)^2(A+Bx)}{\sqrt{d+ex}} dx$$

$$= \frac{2 \left(\frac{Bb^2(d+ex)^{\frac{7}{2}}}{7e^3} + \frac{(d+ex)^{\frac{5}{2}}(Ab^2e+2Babe-3Bb^2d)}{5e^3} + \frac{(d+ex)^{\frac{3}{2}}(2Aabe^2-2Ab^2de+Ba^2e^2-4Babde+3Bb^2d^2)}{3e^3} + \frac{\sqrt{d+ex}(Aa^2e^3-2Aabde^2+Ab^2d^2e-Ba^2e^3)}{e^3} \right)}{e}$$

$$= \frac{Aa^2x + \frac{Bb^2x^4}{4} + \frac{x^3(Ab^2+2Bab)}{3} + \frac{x^2(2Aab+Ba^2)}{2}}{\sqrt{d}}$$

input `integrate((b*x+a)**2*(B*x+A)/(e*x+d)**(1/2),x)`output `Piecewise((2*(B*b**2*(d + e*x)**(7/2))/(7*e**3) + (d + e*x)**(5/2)*(A*b**2*e + 2*B*a*b*e - 3*B*b**2*d)/(5*e**3) + (d + e*x)**(3/2)*(2*A*a*b*e**2 - 2*A*b**2*d*e + B*a**2*e**2 - 4*B*a*b*d*e + 3*B*b**2*d**2)/(3*e**3) + sqrt(d + e*x)*(A*a**2*e**3 - 2*A*a*b*d*e**2 + A*b**2*d**2*e - B*a**2*d*e**2 + 2*B*a*b*d**2*e - B*b**2*d**3)/e**3)/e, Ne(e, 0)), ((A*a**2*x + B*b**2*x**4/4 + x**3*(A*b**2 + 2*B*a*b)/3 + x**2*(2*A*a*b + B*a**2)/2)/sqrt(d), True))`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.26

$$\int \frac{(a+bx)^2(A+Bx)}{\sqrt{d+ex}} dx$$

$$= \frac{2 \left(15 (ex+d)^{\frac{7}{2}} Bb^2 - 21 (3 Bb^2 d - (2 Bab + Ab^2)e)(ex+d)^{\frac{5}{2}} + 35 (3 Bb^2 d^2 - 2 (2 Bab + Ab^2)de + (Ba^2 + 2 Aab + Ab^2)e^2)(ex+d)^{\frac{3}{2}} - 105 (Bb^2 d^3 - Aa^2 e^3 - (2 Baa b + Ab^2) d^2 e + (Baa^2 + 2 Aab) d e^2) \sqrt{ex+d} \right)}{105 e^4}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^(1/2),x, algorithm="maxima")`

output

```
2/105*(15*(e*x + d)^(7/2)*B*b^2 - 21*(3*B*b^2*d - (2*B*a*b + A*b^2)*e)*(e*x + d)^(5/2) + 35*(3*B*b^2*d^2 - 2*(2*B*a*b + A*b^2)*d*e + (B*a^2 + 2*A*a*b)*e^2)*(e*x + d)^(3/2) - 105*(B*b^2*d^3 - A*a^2*e^3 - (2*B*a*b + A*b^2)*d^2*e + (B*a^2 + 2*A*a*b)*d*e^2)*sqrt(e*x + d))/e^4
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.63

$$\int \frac{(a+bx)^2(A+Bx)}{\sqrt{d+ex}} dx$$

$$= \frac{2 \left(105 \sqrt{ex+d} Aa^2 + \frac{35 \left((ex+d)^{\frac{3}{2}} - 3 \sqrt{ex+dd} \right) Ba^2}{e} + \frac{70 \left((ex+d)^{\frac{3}{2}} - 3 \sqrt{ex+dd} \right) Aab}{e} + \frac{14 \left(3 (ex+d)^{\frac{5}{2}} - 10 (ex+d)^{\frac{3}{2}} d + 15 \sqrt{ex+dd} \right) Aa^2}{e^2} \right)}{105 e^4}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^(1/2),x, algorithm="giac")`

output

```
2/105*(105*sqrt(e*x + d)*A*a^2 + 35*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*B*a^2/e + 70*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*A*a*b/e + 14*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*B*a*b/e^2 + 7*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*A*b^2/e^2 + 3*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*B*b^2/e^3)/e
```

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.91

$$\int \frac{(a+bx)^2(A+Bx)}{\sqrt{d+ex}} dx = \frac{(d+ex)^{5/2}(2Ab^2e-6Bb^2d+4Babe)}{5e^4} + \frac{2Bb^2(d+ex)^{7/2}}{7e^4} + \frac{2(ae-bd)(d+ex)^{3/2}(2Abe+BAe-3Bbd)}{3e^4} + \frac{2(Ae-Bd)(ae-bd)^2\sqrt{d+ex}}{e^4}$$

input `int(((A + B*x)*(a + b*x)^2)/(d + e*x)^(1/2), x)`output `((d + e*x)^(5/2)*(2*A*b^2*e - 6*B*b^2*d + 4*B*a*b*e))/(5*e^4) + (2*B*b^2*(d + e*x)^(7/2))/(7*e^4) + (2*(a*e - b*d)*(d + e*x)^(3/2)*(2*A*b*e + B*a*e - 3*B*b*d))/(3*e^4) + (2*(A*e - B*d)*(a*e - b*d)^2*(d + e*x)^(1/2))/e^4`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.90

$$\int \frac{(a+bx)^2(A+Bx)}{\sqrt{d+ex}} dx = \frac{2\sqrt{ex+d}(5b^3e^3x^3 + 21ab^2e^3x^2 - 6b^3de^2x^2 + 35a^2be^3x - 28ab^2de^2x + 8b^3d^2ex + 35a^3e^3 - 70a^2bde^2 - 35e^4)}{35e^4}$$

input `int((b*x+a)^2*(B*x+A)/(e*x+d)^(1/2), x)`output `(2*sqrt(d + e*x)*(35*a**3*e**3 - 70*a**2*b*d*e**2 + 35*a**2*b*e**3*x + 56*a*b**2*d**2*e - 28*a*b**2*d*e**2*x + 21*a*b**2*e**3*x**2 - 16*b**3*d**3 + 8*b**3*d**2*e*x - 6*b**3*d*e**2*x**2 + 5*b**3*e**3*x**3))/(35*e**4)`

3.136 $\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{3/2}} dx$

Optimal result	1283
Mathematica [A] (verified)	1283
Rubi [A] (verified)	1284
Maple [A] (verified)	1285
Fricas [A] (verification not implemented)	1286
Sympy [A] (verification not implemented)	1286
Maxima [A] (verification not implemented)	1287
Giac [A] (verification not implemented)	1287
Mupad [B] (verification not implemented)	1288
Reduce [B] (verification not implemented)	1288

Optimal result

Integrand size = 22, antiderivative size = 124

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{3/2}} dx = \frac{2(bd-ae)^2(Bd-Ae)}{e^4\sqrt{d+ex}} + \frac{2(bd-ae)(3bBd-2Abe-aBe)\sqrt{d+ex}}{e^4} - \frac{2b(3bBd-Abe-2aBe)(d+ex)^{3/2}}{3e^4} + \frac{2b^2B(d+ex)^{5/2}}{5e^4}$$

output

```
2*(-a*e+b*d)^2*(-A*e+B*d)/e^4/(e*x+d)^(1/2)+2*(-a*e+b*d)*(-2*A*b*e-B*a*e+3*B*b*d)*(e*x+d)^(1/2)/e^4-2/3*b*(-A*b*e-2*B*a*e+3*B*b*d)*(e*x+d)^(3/2)/e^4+2/5*b^2*B*(e*x+d)^(5/2)/e^4
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.09

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{3/2}} dx = \frac{30a^2e^2(2Bd-Ae+Bex) + 20abe(3Ae(2d+ex) + B(-8d^2-4dex+e^2x^2))}{15e^4\sqrt{d+ex}}$$

input

```
Integrate[((a + b*x)^2*(A + B*x))/(d + e*x)^(3/2),x]
```


output

$$(30*a^2*e^2*(2*B*d - A*e + B*e*x) + 20*a*b*e*(3*A*e*(2*d + e*x) + B*(-8*d^2 - 4*d*e*x + e^2*x^2)) + 2*b^2*(5*A*e*(-8*d^2 - 4*d*e*x + e^2*x^2) + 3*B*(16*d^3 + 8*d^2*e*x - 2*d*e^2*x^2 + e^3*x^3)))/(15*e^4*sqrt[d + e*x])$$

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^{3/2}} dx$$

↓ 86

$$\int \left(\frac{b\sqrt{d + ex}(2aBe + Abe - 3bBd)}{e^3} + \frac{(ae - bd)(aBe + 2Abe - 3bBd)}{e^3\sqrt{d + ex}} + \frac{(ae - bd)^2(Ae - Bd)}{e^3(d + ex)^{3/2}} + \frac{b^2B(d + ex)^3}{e^3} \right)$$

↓ 2009

$$-\frac{2b(d + ex)^{3/2}(-2aBe - Abe + 3bBd)}{3e^4} + \frac{2\sqrt{d + ex}(bd - ae)(-aBe - 2Abe + 3bBd)}{e^4} + \frac{2(bd - ae)^2(Bd - Ae)}{e^4\sqrt{d + ex}} + \frac{2b^2B(d + ex)^{5/2}}{5e^4}$$

input

$$\text{Int}[(a + b*x)^2*(A + B*x)/(d + e*x)^(3/2), x]$$

output

$$(2*(b*d - a*e)^2*(B*d - A*e))/(e^4*sqrt[d + e*x]) + (2*(b*d - a*e)*(3*b*B*d - 2*A*b*e - a*B*e)*sqrt[d + e*x])/e^4 - (2*b*(3*b*B*d - A*b*e - 2*a*B*e)*(d + e*x)^(3/2))/(3*e^4) + (2*b^2*B*(d + e*x)^(5/2))/(5*e^4)$$

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.98

method	result
pseudoelliptic	$\frac{\left((6Bx^3 + 10Ax^2)b^2 + 60a\left(\frac{Bx}{3} + A\right)xb - 30a^2(-Bx + A) \right) e^3 + 120 \left(-\frac{(3Bx + A)xb^2}{3} + a\left(-\frac{2Bx}{3} + A\right)b + \frac{a^2B}{2} \right) d e^2 - 80b}{15\sqrt{ex+d}e^4}$
risch	$\frac{2(3e^2b^2Bx^2 + 5Ab^2e^2x + 10Babe^2x - 9b^2Bdex + 30Aabe^2 - 25Ab^2de + 15Ba^2e^2 - 50Babde + 33b^2Bd^2)\sqrt{ex+d}}{15e^4}$
gospers	$\frac{2(-3b^2Bx^3e^3 - 5Ax^2b^2e^3 - 10Bx^2abe^3 + 6Bx^2b^2de^2 - 30Axabe^3 + 20Ax^2b^2de^2 - 15Bxa^2e^3 + 40Bxabde^2 - 24Bxb^2de^2)}{15\sqrt{ex+d}e^4}$
trager	$\frac{2(-3b^2Bx^3e^3 - 5Ax^2b^2e^3 - 10Bx^2abe^3 + 6Bx^2b^2de^2 - 30Axabe^3 + 20Ax^2b^2de^2 - 15Bxa^2e^3 + 40Bxabde^2 - 24Bxb^2de^2)}{15\sqrt{ex+d}e^4}$
oring	$\frac{2(-3b^2Bx^3e^3 - 5Ax^2b^2e^3 - 10Bx^2abe^3 + 6Bx^2b^2de^2 - 30Axabe^3 + 20Ax^2b^2de^2 - 15Bxa^2e^3 + 40Bxabde^2 - 24Bxb^2de^2)}{15\sqrt{ex+d}e^4}$
derivativdivides	$\frac{\frac{2b^2B(ex+d)^{\frac{5}{2}}}{5} + \frac{2Ab^2e(ex+d)^{\frac{3}{2}}}{3} + \frac{4Babe(ex+d)^{\frac{3}{2}}}{3} - 2Bb^2d(ex+d)^{\frac{3}{2}} + 4Aabe^2\sqrt{ex+d} - 4Ab^2de\sqrt{ex+d} + 2Ba^2e^2\sqrt{ex+d}}{e^4}$
default	$\frac{\frac{2b^2B(ex+d)^{\frac{5}{2}}}{5} + \frac{2Ab^2e(ex+d)^{\frac{3}{2}}}{3} + \frac{4Babe(ex+d)^{\frac{3}{2}}}{3} - 2Bb^2d(ex+d)^{\frac{3}{2}} + 4Aabe^2\sqrt{ex+d} - 4Ab^2de\sqrt{ex+d} + 2Ba^2e^2\sqrt{ex+d}}{e^4}$

```
input int((b*x+a)^2*(B*x+A)/(e*x+d)^(3/2), x, method=_RETURNVERBOSE)
```

```
output 1/15*(((6*B*x^3+10*A*x^2)*b^2+60*a*(1/3*B*x+A)*x*b-30*a^2*(-B*x+A))*e^3+120*(-1/3*(3/10*B*x+A)*x*b^2+a*(-2/3*B*x+A)*b+1/2*a^2*B)*d*e^2-80*b*((-3/5*B*x+A)*b+2*B*a)*d^2*e+96*b^2*B*d^3)/(e*x+d)^(1/2)/e^4
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.33

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{3/2}} dx = \frac{2(3Bb^2e^3x^3 + 48Bb^2d^3 - 15Aa^2e^3 - 40(2Bab + Ab^2)d^2e + 30(Ba^2 + 2Aab$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^(3/2),x, algorithm="fricas")`

output `2/15*(3*B*b^2*e^3*x^3 + 48*B*b^2*d^3 - 15*A*a^2*e^3 - 40*(2*B*a*b + A*b^2)*d^2*e + 30*(B*a^2 + 2*A*a*b)*d*e^2 - (6*B*b^2*d*e^2 - 5*(2*B*a*b + A*b^2)*e^3)*x^2 + (24*B*b^2*d^2*e - 20*(2*B*a*b + A*b^2)*d*e^2 + 15*(B*a^2 + 2*A*a*b)*e^3)*x)*sqrt(e*x + d)/(e^5*x + d*e^4)`

Sympy [A] (verification not implemented)

Time = 4.22 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.67

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{3/2}} dx = \left\{ \frac{2 \left(\frac{Bb^2(d+ex)^{\frac{5}{2}}}{5e^3} + \frac{(d+ex)^{\frac{3}{2}}(Ab^2e+2Babe-3Bb^2d)}{3e^3} + \frac{\sqrt{d+ex}(2Aabe^2-2Ab^2de+Ba^2e^2-4Babde+3Bb^2d^2)}{e^3} \right) + (-Aa^2x + \frac{Bb^2x^4}{4} + \frac{x^3(Ab^2+2Bab)}{3} + \frac{x^2(2Aab+Ba^2)}{2}}{d^{\frac{3}{2}}} \right.$$

input `integrate((b*x+a)**2*(B*x+A)/(e*x+d)**(3/2),x)`

output `Piecewise((2*(B*b**2*(d + e*x)**(5/2)/(5*e**3) + (d + e*x)**(3/2)*(A*b**2*e + 2*B*a*b*e - 3*B*b**2*d)/(3*e**3) + sqrt(d + e*x)*(2*A*a*b*e**2 - 2*A*b**2*d*e + B*a**2*e**2 - 4*B*a*b*d*e + 3*B*b**2*d**2)/e**3 + (-A*e + B*d)*(a*e - b*d)**2/(e**3*sqrt(d + e*x)))/e, Ne(e, 0)), ((A*a**2*x + B*b**2*x**4/4 + x**3*(A*b**2 + 2*B*a*b)/3 + x**2*(2*A*a*b + B*a**2)/2)/d**(3/2), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.35

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{3/2}} dx = \frac{2 \left(\frac{3(ex+d)^{5/2} Bb^2 - 5(3Bb^2d - (2Bab+Ab^2)e)(ex+d)^{3/2} + 15(3Bb^2d^2 - 2(2Bab+Ab^2)de + (Ba^2+2Aab)e^3)}{e^3} \right)}{15e}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^(3/2),x, algorithm="maxima")`

output

```
2/15*((3*(e*x + d)^(5/2)*B*b^2 - 5*(3*B*b^2*d - (2*B*a*b + A*b^2)*e)*(e*x
+ d)^(3/2) + 15*(3*B*b^2*d^2 - 2*(2*B*a*b + A*b^2)*d*e + (B*a^2 + 2*A*a*b)
*e^2)*sqrt(e*x + d))/e^3 + 15*(B*b^2*d^3 - A*a^2*e^3 - (2*B*a*b + A*b^2)*d
^2*e + (B*a^2 + 2*A*a*b)*d*e^2)/(sqrt(e*x + d)*e^3))/e
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.78

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{3/2}} dx = \frac{2(Bb^2d^3 - 2Babd^2e - Ab^2d^2e + Ba^2de^2 + 2Aabde^2 - Aa^2e^3)}{\sqrt{ex+d}e^4} + \frac{2 \left(3(ex+d)^{5/2} Bb^2e^{16} - 15(ex+d)^{3/2} Bb^2de^{16} + 45\sqrt{ex+d} Bb^2d^2e^{16} + 10(ex+d)^{3/2} Babe^{17} + 5(ex+d)^{3/2} \right)}{15e^{20}}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^(3/2),x, algorithm="giac")`

output

```
2*(B*b^2*d^3 - 2*B*a*b*d^2*e - A*b^2*d^2*e + B*a^2*d*e^2 + 2*A*a*b*d*e^2 -
A*a^2*e^3)/(sqrt(e*x + d)*e^4) + 2/15*(3*(e*x + d)^(5/2)*B*b^2*e^16 - 15*
(e*x + d)^(3/2)*B*b^2*d*e^16 + 45*sqrt(e*x + d)*B*b^2*d^2*e^16 + 10*(e*x +
d)^(3/2)*B*a*b*e^17 + 5*(e*x + d)^(3/2)*A*b^2*e^17 - 60*sqrt(e*x + d)*B*a
*b*d*e^17 - 30*sqrt(e*x + d)*A*b^2*d*e^17 + 15*sqrt(e*x + d)*B*a^2*e^18 +
30*sqrt(e*x + d)*A*a*b*e^18)/e^20
```

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.24

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{3/2}} dx = \frac{(d+ex)^{3/2}(2Ab^2e - 6Bb^2d + 4Babe)}{3e^4} - \frac{-2Ba^2de^2 + 2Aa^2e^3 + 4Babd^2e - 4Aabde^2 - 2Bb^2d^3 + 2Ab^2d^2e}{e^4\sqrt{d+ex}} + \frac{2Bb^2(d+ex)^{5/2}}{5e^4} + \frac{2(ae-bd)\sqrt{d+ex}(2Abe + Bae - 3Bbd)}{e^4}$$

input `int(((A + B*x)*(a + b*x)^2)/(d + e*x)^(3/2), x)`output `((d + e*x)^(3/2)*(2*A*b^2*e - 6*B*b^2*d + 4*B*a*b*e))/(3*e^4) - (2*A*a^2*e^3 - 2*B*b^2*d^3 + 2*A*b^2*d^2*e - 2*B*a^2*d*e^2 - 4*A*a*b*d*e^2 + 4*B*a*b*d^2*e)/(e^4*(d + e*x)^(1/2)) + (2*B*b^2*(d + e*x)^(5/2))/(5*e^4) + (2*(a - b*d)*(d + e*x)^(1/2)*(2*A*b*e + B*a*e - 3*B*b*d))/e^4`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.93

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{3/2}} dx = \frac{\frac{2}{5}b^3e^3x^3 + 2ab^2e^3x^2 - \frac{4}{5}b^3de^2x^2 + 6a^2be^3x - 8ab^2de^2x + \frac{16}{5}b^3d^2ex - 2a^3e^3}{\sqrt{ex+d}e^4}$$

input `int((b*x+a)^2*(B*x+A)/(e*x+d)^(3/2), x)`output `(2*(- 5*a**3*e**3 + 30*a**2*b*d*e**2 + 15*a**2*b*e**3*x - 40*a*b**2*d**2*e - 20*a*b**2*d*e**2*x + 5*a*b**2*e**3*x**2 + 16*b**3*d**3 + 8*b**3*d**2*e*x - 2*b**3*d*e**2*x**2 + b**3*e**3*x**3))/(5*sqrt(d + e*x)*e**4)`

3.137 $\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{5/2}} dx$

Optimal result	1289
Mathematica [A] (verified)	1289
Rubi [A] (verified)	1290
Maple [A] (verified)	1291
Fricas [A] (verification not implemented)	1292
Sympy [B] (verification not implemented)	1292
Maxima [A] (verification not implemented)	1293
Giac [A] (verification not implemented)	1294
Mupad [B] (verification not implemented)	1294
Reduce [B] (verification not implemented)	1295

Optimal result

Integrand size = 22, antiderivative size = 124

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{5/2}} dx = \frac{2(bd-ae)^2(Bd-Ae)}{3e^4(d+ex)^{3/2}} - \frac{2(bd-ae)(3bBd-2Abe-aBe)}{e^4\sqrt{d+ex}} - \frac{2b(3bBd-Abe-2aBe)\sqrt{d+ex}}{e^4} + \frac{2b^2B(d+ex)^{3/2}}{3e^4}$$

output

```
2/3*(-a*e+b*d)^2*(-A*e+B*d)/e^4/(e*x+d)^(3/2)-2*(-a*e+b*d)*(-2*A*b*e-B*a*e+3*B*b*d)/e^4/(e*x+d)^(1/2)-2*b*(-A*b*e-2*B*a*e+3*B*b*d)*(e*x+d)^(1/2)/e^4+2/3*b^2*B*(e*x+d)^(3/2)/e^4
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.10

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{5/2}} dx = \frac{2(a^2e^2(2Bd+ Ae+ 3Bex) - 2abe(-Ae(2d+ 3ex) + B(8d^2+ 12dex+ 3e^2x^2)) + b^2(-Ae(8d^2+ 12dex - 3e^2x^2)))}{3e^4(d+ex)^{3/2}}$$

input `Integrate[((a + b*x)^2*(A + B*x))/(d + e*x)^(5/2),x]`

output `(-2*(a^2*e^2*(2*B*d + A*e + 3*B*e*x) - 2*a*b*e*(-(A*e*(2*d + 3*e*x)) + B*(8*d^2 + 12*d*e*x + 3*e^2*x^2)) + b^2*(-(A*e*(8*d^2 + 12*d*e*x + 3*e^2*x^2)) + B*(16*d^3 + 24*d^2*e*x + 6*d*e^2*x^2 - e^3*x^3)))/(3*e^4*(d + e*x)^(3/2))`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^{5/2}} dx$$

↓ 86

$$\int \left(\frac{b(2aBe + Abe - 3bBd)}{e^3\sqrt{d + ex}} + \frac{(ae - bd)(aBe + 2Abe - 3bBd)}{e^3(d + ex)^{3/2}} + \frac{(ae - bd)^2(Ae - Bd)}{e^3(d + ex)^{5/2}} + \frac{b^2B\sqrt{d + ex}}{e^3} \right) dx$$

↓ 2009

$$-\frac{2b\sqrt{d + ex}(-2aBe - Abe + 3bBd)}{e^4} - \frac{2(bd - ae)(-aBe - 2Abe + 3bBd)}{e^4\sqrt{d + ex}} + \frac{2(bd - ae)^2(Bd - Ae)}{3e^4(d + ex)^{3/2}} + \frac{2b^2B(d + ex)^{3/2}}{3e^4}$$

input `Int[((a + b*x)^2*(A + B*x))/(d + e*x)^(5/2),x]`

output `(2*(b*d - a*e)^2*(B*d - A*e))/(3*e^4*(d + e*x)^(3/2)) - (2*(b*d - a*e)*(3*b*B*d - 2*A*b*e - a*B*e))/(e^4*sqrt[d + e*x]) - (2*b*(3*b*B*d - A*b*e - 2*a*B*e)*sqrt[d + e*x])/e^4 + (2*b^2*B*(d + e*x)^(3/2))/(3*e^4)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.41

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{5/2}} dx = \frac{2(Bb^2e^3x^3 - 16Bb^2d^3 - Aa^2e^3 + 8(2Bab + Ab^2)d^2e - 2(Ba^2 + 2Aab)de^2 - \dots}{(d+ex)^{5/2}}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^(5/2),x, algorithm="fricas")`

output `2/3*(B*b^2*e^3*x^3 - 16*B*b^2*d^3 - A*a^2*e^3 + 8*(2*B*a*b + A*b^2)*d^2*e - 2*(B*a^2 + 2*A*a*b)*d*e^2 - 3*(2*B*b^2*d*e^2 - (2*B*a*b + A*b^2)*e^3)*x^2 - 3*(8*B*b^2*d^2*e - 4*(2*B*a*b + A*b^2)*d*e^2 + (B*a^2 + 2*A*a*b)*e^3)*x)*sqrt(e*x + d)/(e^6*x^2 + 2*d*e^5*x + d^2*e^4)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 709 vs. $2(124) = 248$.

Time = 0.41 (sec) , antiderivative size = 709, normalized size of antiderivative = 5.72

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{5/2}} dx = \left\{ \begin{array}{l} -\frac{2Aa^2e^3}{3de^4\sqrt{d+ex}+3e^5x\sqrt{d+ex}} - \frac{8Aabde^2}{3de^4\sqrt{d+ex}+3e^5x\sqrt{d+ex}} - \frac{12Aabe^3x}{3de^4\sqrt{d+ex}+3e^5x\sqrt{d+ex}} + \frac{1}{3de^4\sqrt{d+ex}} \\ \frac{Aa^2x+Aabx^2+\frac{Ab^2x^3}{3}+\frac{Ba^2x^2}{2}+\frac{2Babx^3}{3}+\frac{Bb^2x^4}{4}}{d^{\frac{5}{2}}} \end{array} \right.$$

input `integrate((b*x+a)**2*(B*x+A)/(e*x+d)**(5/2),x)`

output

```
Piecewise((-2*A**2*e**3/(3*d**4*sqrt(d + e*x) + 3*e**5*x*sqrt(d + e*x)) - 8*A*b*d**2/(3*d**4*sqrt(d + e*x) + 3*e**5*x*sqrt(d + e*x)) - 12*A*b*e**3*x/(3*d**4*sqrt(d + e*x) + 3*e**5*x*sqrt(d + e*x)) + 16*A*b**2*d**2*e/(3*d**4*sqrt(d + e*x) + 3*e**5*x*sqrt(d + e*x)) + 24*A*b**2*d*e**2*x/(3*d**4*sqrt(d + e*x) + 3*e**5*x*sqrt(d + e*x)) + 6*A*b**2*e**3*x**2/(3*d**4*sqrt(d + e*x) + 3*e**5*x*sqrt(d + e*x)) - 4*B*a**2*d**2/(3*d**4*sqrt(d + e*x) + 3*e**5*x*sqrt(d + e*x)) - 6*B*a**2*e**3*x/(3*d**4*sqrt(d + e*x) + 3*e**5*x*sqrt(d + e*x)) + 32*B*a*b*d**2*e/(3*d**4*sqrt(d + e*x) + 3*e**5*x*sqrt(d + e*x)) + 48*B*a*b*d**2*x/(3*d**4*sqrt(d + e*x) + 3*e**5*x*sqrt(d + e*x)) + 12*B*a*b*e**3*x**2/(3*d**4*sqrt(d + e*x) + 3*e**5*x*sqrt(d + e*x)) - 32*B*b**2*d**3/(3*d**4*sqrt(d + e*x) + 3*e**5*x*sqrt(d + e*x)) - 48*B*b**2*d**2*e*x/(3*d**4*sqrt(d + e*x) + 3*e**5*x*sqrt(d + e*x)) - 12*B*b**2*d**2*x**2/(3*d**4*sqrt(d + e*x) + 3*e**5*x*sqrt(d + e*x)) + 2*B*b**2*e**3*x**3/(3*d**4*sqrt(d + e*x) + 3*e**5*x*sqrt(d + e*x)), Ne(e, 0)), ((A**2*x + A*b*x**2 + A*b**2*x**3/3 + B*a**2*x**2/2 + 2*B*a*b*x**3/3 + B*b**2*x**4/4)/d**(5/2), True))
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.31

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^{5/2}} dx = \frac{2 \left(\frac{(ex+d)^{\frac{3}{2}} Bb^2 - 3(3Bb^2d - (2Bab + Ab^2)e)\sqrt{ex+d}}{e^3} + \frac{Bb^2d^3 - Aa^2e^3 - (2Bab + Ab^2)d^2e + (Ba^2 + 2Aab)}{3e} \right)}{3e}$$

input

```
integrate((b*x+a)^2*(B*x+A)/(e*x+d)^(5/2),x, algorithm="maxima")
```

output

```
2/3*(((e*x + d)^(3/2)*B*b^2 - 3*(3*B*b^2*d - (2*B*a*b + A*b^2)*e)*sqrt(e*x + d))/e^3 + (B*b^2*d^3 - A*a^2*e^3 - (2*B*a*b + A*b^2)*d^2*e + (B*a^2 + 2*A*a*b)*d*e^2 - 3*(3*B*b^2*d^2 - 2*(2*B*a*b + A*b^2)*d*e + (B*a^2 + 2*A*a*b)*e^2)*(e*x + d))/((e*x + d)^(3/2)*e^3)/e
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.64

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{5/2}} dx = \frac{2(9(ex+d)Bb^2d^2 - Bb^2d^3 - 12(ex+d)Babde - 6(ex+d)Ab^2de + 2Babd^2e + Ab^2d^2e + 3(ex+d)B)}{3(ex+d)^{3/2}e^4} + \frac{2\left((ex+d)^{3/2}Bb^2e^8 - 9\sqrt{ex+d}Bb^2de^8 + 6\sqrt{ex+d}Babe^9 + 3\sqrt{ex+d}Ab^2e^9\right)}{3e^{12}}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^(5/2),x, algorithm="giac")`

output

```
-2/3*(9*(e*x + d)*B*b^2*d^2 - B*b^2*d^3 - 12*(e*x + d)*B*a*b*d*e - 6*(e*x + d)*A*b^2*d*e + 2*B*a*b*d^2*e + A*b^2*d^2*e + 3*(e*x + d)*B*a^2*e^2 + 6*(e*x + d)*A*a*b*e^2 - B*a^2*d*e^2 - 2*A*a*b*d*e^2 + A*a^2*e^3)/((e*x + d)^(3/2)*e^4) + 2/3*((e*x + d)^(3/2)*B*b^2*e^8 - 9*sqrt(e*x + d)*B*b^2*d*e^8 + 6*sqrt(e*x + d)*B*a*b*e^9 + 3*sqrt(e*x + d)*A*b^2*e^9)/e^12
```

Mupad [B] (verification not implemented)

Time = 0.97 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.52

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{5/2}} dx = \frac{2Bb^2d^3 - 2Aa^2e^3 + 2Bb^2(d+ex)^3 + 6Ab^2e(d+ex)^2 - 6Ba^2e^2(d+ex)}{(d+ex)^{5/2}}$$

input `int(((A + B*x)*(a + b*x)^2)/(d + e*x)^(5/2),x)`

output

```
(2*B*b^2*d^3 - 2*A*a^2*e^3 + 2*B*b^2*(d + e*x)^3 + 6*A*b^2*e*(d + e*x)^2 - 6*B*a^2*e^2*(d + e*x) - 18*B*b^2*d*(d + e*x)^2 - 18*B*b^2*d^2*(d + e*x) - 2*A*b^2*d^2*e + 2*B*a^2*d*e^2 - 12*A*a*b*e^2*(d + e*x) + 12*B*a*b*e*(d + e*x)^2 + 12*A*b^2*d*e*(d + e*x) + 4*A*a*b*d*e^2 - 4*B*a*b*d^2*e + 24*B*a*b*d*e*(d + e*x))/(3*e^4*(d + e*x)^(3/2))
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.98

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^{5/2}} dx = \frac{\frac{2}{3}b^3e^3x^3 + 6ab^2e^3x^2 - 4b^3de^2x^2 - 6a^2be^3x + 24ab^2de^2x - 16b^3d^2ex - \frac{2}{3}a^3e^3}{\sqrt{ex + d}e^4(ex + d)}$$

input `int((b*x+a)^2*(B*x+A)/(e*x+d)^(5/2),x)`output `(2*(- a**3*e**3 - 6*a**2*b*d*e**2 - 9*a**2*b*e**3*x + 24*a*b**2*d**2*e + 36*a*b**2*d*e**2*x + 9*a*b**2*e**3*x**2 - 16*b**3*d**3 - 24*b**3*d**2*e*x - 6*b**3*d*e**2*x**2 + b**3*e**3*x**3))/(3*sqrt(d + e*x)*e**4*(d + e*x))`

3.138 $\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{7/2}} dx$

Optimal result	1296
Mathematica [A] (verified)	1296
Rubi [A] (verified)	1297
Maple [A] (verified)	1298
Fricas [A] (verification not implemented)	1299
Sympy [B] (verification not implemented)	1299
Maxima [A] (verification not implemented)	1300
Giac [A] (verification not implemented)	1301
Mupad [B] (verification not implemented)	1301
Reduce [B] (verification not implemented)	1302

Optimal result

Integrand size = 22, antiderivative size = 124

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{7/2}} dx = \frac{2(bd-ae)^2(Bd-Ae)}{5e^4(d+ex)^{5/2}} - \frac{2(bd-ae)(3bBd-2Abe-aBe)}{3e^4(d+ex)^{3/2}} + \frac{2b(3bBd-Abe-2aBe)}{e^4\sqrt{d+ex}} + \frac{2b^2B\sqrt{d+ex}}{e^4}$$

output `2/5*(-a*e+b*d)^2*(-A*e+B*d)/e^4/(e*x+d)^(5/2)-2/3*(-a*e+b*d)*(-2*A*b*e-B*a*e+3*B*b*d)/e^4/(e*x+d)^(3/2)+2*b*(-A*b*e-2*B*a*e+3*B*b*d)/e^4/(e*x+d)^(1/2)+2*b^2*B*(e*x+d)^(1/2)/e^4`

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.10

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{7/2}} dx = \frac{2(a^2e^2(2Bd+3Ae+5Bex) + 2abe(Ae(2d+5ex) + B(8d^2+20dex+15e^2x^2)) + b^2(Ae(8d^2+20dex+15e^2x^2) + 2Bd^2+2Ade+2Bde+2B^2x^2))}{15e^4(d+ex)^{5/2}}$$

input `Integrate[((a + b*x)^2*(A + B*x))/(d + e*x)^(7/2), x]`

output

$$\frac{(-2*(a^2*e^2*(2*B*d + 3*A*e + 5*B*e*x) + 2*a*b*e*(A*e*(2*d + 5*e*x) + B*(8*d^2 + 20*d*e*x + 15*e^2*x^2)) + b^2*(A*e*(8*d^2 + 20*d*e*x + 15*e^2*x^2) - 3*B*(16*d^3 + 40*d^2*e*x + 30*d*e^2*x^2 + 5*e^3*x^3)))/(15*e^4*(d + e*x)^(5/2))$$

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^{7/2}} dx$$

↓ 86

$$\int \left(\frac{b(2aBe + Abe - 3bBd)}{e^3(d + ex)^{3/2}} + \frac{(ae - bd)(aBe + 2Abe - 3bBd)}{e^3(d + ex)^{5/2}} + \frac{(ae - bd)^2(Ae - Bd)}{e^3(d + ex)^{7/2}} + \frac{b^2B}{e^3\sqrt{d + ex}} \right) dx$$

↓ 2009

$$\frac{2b(-2aBe - Abe + 3bBd)}{e^4\sqrt{d + ex}} - \frac{2(bd - ae)(-aBe - 2Abe + 3bBd)}{3e^4(d + ex)^{3/2}} + \frac{2(bd - ae)^2(Bd - Ae)}{5e^4(d + ex)^{5/2}} + \frac{2b^2B\sqrt{d + ex}}{e^4}$$

input

$$\text{Int}[(a + b*x)^2*(A + B*x)/(d + e*x)^(7/2), x]$$

output

$$\frac{(2*(b*d - a*e)^2*(B*d - A*e))/(5*e^4*(d + e*x)^(5/2)) - (2*(b*d - a*e)*(3*b*B*d - 2*A*b*e - a*B*e))/(3*e^4*(d + e*x)^(3/2)) + (2*b*(3*b*B*d - A*b*e - 2*a*B*e))/(e^4*\text{Sqrt}[d + e*x]) + (2*b^2*B*\text{Sqrt}[d + e*x])/e^4$$

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.52

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^{7/2}} dx = \frac{2(15 Bb^2e^3x^3 + 48 Bb^2d^3 - 3 Aa^2e^3 - 8(2 Bab + Ab^2)d^2e - 2(Ba^2 + 2 Aab)d}{15}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^(7/2),x, algorithm="fricas")`

output `2/15*(15*B*b^2*e^3*x^3 + 48*B*b^2*d^3 - 3*A*a^2*e^3 - 8*(2*B*a*b + A*b^2)*d^2*e - 2*(B*a^2 + 2*A*a*b)*d*e^2 + 15*(6*B*b^2*d*e^2 - (2*B*a*b + A*b^2)*e^3)*x^2 + 5*(24*B*b^2*d^2*e - 4*(2*B*a*b + A*b^2)*d*e^2 - (B*a^2 + 2*A*a*b)*e^3)*x)*sqrt(e*x + d)/(e^7*x^3 + 3*d*e^6*x^2 + 3*d^2*e^5*x + d^3*e^4)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1015 vs. $2(126) = 252$.

Time = 0.57 (sec) , antiderivative size = 1015, normalized size of antiderivative = 8.19

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^{7/2}} dx = \text{Too large to display}$$

input `integrate((b*x+a)**2*(B*x+A)/(e*x+d)**(7/2),x)`

output

```
Piecewise((-6*A*a**2*e**3/(15*d**2*e**4*sqrt(d + e*x) + 30*d*e**5*x*sqrt(d + e*x) + 15*e**6*x**2*sqrt(d + e*x)) - 8*A*a*b*d*e**2/(15*d**2*e**4*sqrt(d + e*x) + 30*d*e**5*x*sqrt(d + e*x) + 15*e**6*x**2*sqrt(d + e*x)) - 20*A*a*b*e**3*x/(15*d**2*e**4*sqrt(d + e*x) + 30*d*e**5*x*sqrt(d + e*x) + 15*e**6*x**2*sqrt(d + e*x)) - 16*A*b**2*d**2*e/(15*d**2*e**4*sqrt(d + e*x) + 30*d*e**5*x*sqrt(d + e*x) + 15*e**6*x**2*sqrt(d + e*x)) - 40*A*b**2*d*e**2*x/(15*d**2*e**4*sqrt(d + e*x) + 30*d*e**5*x*sqrt(d + e*x) + 15*e**6*x**2*sqrt(d + e*x)) - 30*A*b**2*e**3*x**2/(15*d**2*e**4*sqrt(d + e*x) + 30*d*e**5*x*sqrt(d + e*x) + 15*e**6*x**2*sqrt(d + e*x)) - 4*B*a**2*d*e**2/(15*d**2*e**4*sqrt(d + e*x) + 30*d*e**5*x*sqrt(d + e*x) + 15*e**6*x**2*sqrt(d + e*x)) - 10*B*a**2*e**3*x/(15*d**2*e**4*sqrt(d + e*x) + 30*d*e**5*x*sqrt(d + e*x) + 15*e**6*x**2*sqrt(d + e*x)) - 32*B*a*b*d**2*e/(15*d**2*e**4*sqrt(d + e*x) + 30*d*e**5*x*sqrt(d + e*x) + 15*e**6*x**2*sqrt(d + e*x)) - 80*B*a*b*d*e**2*x/(15*d**2*e**4*sqrt(d + e*x) + 30*d*e**5*x*sqrt(d + e*x) + 15*e**6*x**2*sqrt(d + e*x)) - 60*B*a*b*e**3*x**2/(15*d**2*e**4*sqrt(d + e*x) + 30*d*e**5*x*sqrt(d + e*x) + 15*e**6*x**2*sqrt(d + e*x)) + 96*B*b**2*d**3/(15*d**2*e**4*sqrt(d + e*x) + 30*d*e**5*x*sqrt(d + e*x) + 15*e**6*x**2*sqrt(d + e*x)) + 240*B*b**2*d**2*e*x/(15*d**2*e**4*sqrt(d + e*x) + 30*d*e**5*x*sqrt(d + e*x) + 15*e**6*x**2*sqrt(d + e*x)) + 180*B*b**2*d*e**2*x**2/(15*d**2*e**4*sqrt(d + e*x) + 30*d*e**5*x*sqrt(d + e*x) + 15*e**6*x**2*sqrt(...
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.32

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^{7/2}} dx = \frac{2 \left(\frac{15\sqrt{ex+d}Bb^2}{e^3} + \frac{3Bb^2d^3 - 3Aa^2e^3 - 3(2Bab + Ab^2)d^2e + 3(Ba^2 + 2Aab)de^2 + 15(3Bb^2d - (2Bab + Ab^2)e)}{(ex+d)^{5/2}e^3} \right)}{15e}$$

input

```
integrate((b*x+a)^2*(B*x+A)/(e*x+d)^(7/2),x, algorithm="maxima")
```

output

```
2/15*(15*sqrt(e*x + d)*B*b^2/e^3 + (3*B*b^2*d^3 - 3*A*a^2*e^3 - 3*(2*B*a*b + A*b^2)*d^2*e + 3*(B*a^2 + 2*A*a*b)*d*e^2 + 15*(3*B*b^2*d - (2*B*a*b + A*b^2)*e)*(e*x + d)^2 - 5*(3*B*b^2*d^2 - 2*(2*B*a*b + A*b^2)*d*e + (B*a^2 + 2*A*a*b)*e^2)*(e*x + d))/((e*x + d)^(5/2)*e^3)/e
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.56

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{7/2}} dx = \frac{2\sqrt{ex+d}Bb^2}{e^4} + \frac{2(45(ex+d)^2Bb^2d - 15(ex+d)Bb^2d^2 + 3Bb^2d^3 - 30(ex+d)^2Babe - 15(ex+d)^2Ab^2e + 20(ex+d)^2A^2b^2e^2 - 10(ex+d)A^2b^2e^3 + 5A^2b^2e^4)}{(ex+d)^{5/2}e^4}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^(7/2),x, algorithm="giac")`output
$$\frac{2\sqrt{ex+d}B^2b^2/e^4 + 2/15(45(ex+d)^2B^2b^2d - 15(ex+d)B^2b^2d^2 + 3B^2b^2d^3 - 30(ex+d)^2B^2a^2b^2e - 15(ex+d)^2A^2b^2e^2 + 20(ex+d)B^2a^2b^2d^2e + 10(ex+d)A^2b^2d^2e - 6B^2a^2b^2d^2e - 3A^2b^2d^2e - 5(ex+d)B^2a^2e^2 - 10(ex+d)A^2a^2b^2e^2 + 3B^2a^2d^2e^2 + 6A^2a^2b^2d^2e - 3A^2a^2e^3)/(ex+d)^{5/2}e^4}$$
Mupad [B] (verification not implemented)

Time = 0.97 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.35

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{7/2}} dx = \frac{2(2Ba^2de^2 + 5Ba^2e^3x + 3Aa^2e^3 + 16Babde^2e + 40Babde^2x + 4Aabde^2 + 30Babe^3x^2 + 10Aabe^3x + 5A^2b^2e^3x^2 - 15B^2b^2e^3x^3 + 30B^2a^2b^2e^3x^2 + 20A^2b^2d^2e^2x - 120B^2b^2d^2e^2xx - 90B^2b^2d^2e^2x^2 + 4A^2a^2b^2d^2e^2 + 16B^2a^2b^2d^2e + 10A^2a^2b^2e^3x + 40B^2a^2b^2d^2e^2x)/(15e^4(d+ex)^{5/2})}$$

input `int(((A+B*x)*(a+b*x)^2)/(d+e*x)^(7/2),x)`output
$$\frac{-(2(3A^2a^2e^3 - 48B^2b^2d^3 + 8A^2b^2d^2e + 2B^2a^2d^2e^2 + 5B^2a^2e^3x + 15A^2b^2e^3x^2 - 15B^2b^2e^3x^3 + 30B^2a^2b^2e^3x^2 + 20A^2b^2d^2e^2x - 120B^2b^2d^2e^2xx - 90B^2b^2d^2e^2x^2 + 4A^2a^2b^2d^2e^2 + 16B^2a^2b^2d^2e + 10A^2a^2b^2e^3x + 40B^2a^2b^2d^2e^2x))/(15e^4(d+ex)^{5/2})}$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.08

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{7/2}} dx = \frac{2b^3e^3x^3 - 6ab^2e^3x^2 + 12b^3de^2x^2 - 2a^2be^3x - 8ab^2de^2x + 16b^3d^2ex - \frac{2}{5}a^3e^3}{\sqrt{ex+d}e^4(e^2x^2 + 2dex + d^2)}$$

input `int((b*x+a)^2*(B*x+A)/(e*x+d)^(7/2),x)`output `(2*(- a**3*e**3 - 2*a**2*b*d*e**2 - 5*a**2*b*e**3*x - 8*a*b**2*d**2*e - 20*a*b**2*d*e**2*x - 15*a*b**2*e**3*x**2 + 16*b**3*d**3 + 40*b**3*d**2*e*x + 30*b**3*d*e**2*x**2 + 5*b**3*e**3*x**3))/(5*sqrt(d + e*x)*e**4*(d**2 + 2*d*e*x + e**2*x**2))`

3.139 $\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{9/2}} dx$

Optimal result	1303
Mathematica [A] (verified)	1303
Rubi [A] (verified)	1304
Maple [A] (verified)	1305
Fricas [A] (verification not implemented)	1306
Sympy [B] (verification not implemented)	1306
Maxima [A] (verification not implemented)	1307
Giac [A] (verification not implemented)	1308
Mupad [B] (verification not implemented)	1308
Reduce [B] (verification not implemented)	1309

Optimal result

Integrand size = 22, antiderivative size = 126

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{9/2}} dx = \frac{2(bd-ae)^2(Bd-Ae)}{7e^4(d+ex)^{7/2}} - \frac{2(bd-ae)(3bBd-2Abe-aBe)}{5e^4(d+ex)^{5/2}} + \frac{2b(3bBd-Abe-2aBe)}{3e^4(d+ex)^{3/2}} - \frac{2b^2B}{e^4\sqrt{d+ex}}$$

output

```
2/7*(-a*e+b*d)^2*(-A*e+B*d)/e^4/(e*x+d)^(7/2)-2/5*(-a*e+b*d)*(-2*A*b*e-B*a
*e+3*B*b*d)/e^4/(e*x+d)^(5/2)+2/3*b*(-A*b*e-2*B*a*e+3*B*b*d)/e^4/(e*x+d)^(
3/2)-2*b^2*B/e^4/(e*x+d)^(1/2)
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.10

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{9/2}} dx = \frac{2(3a^2e^2(2Bd+5Ae+7Bex) + 2abe(3Ae(2d+7ex) + B(8d^2+28dex+35e^2x^2)) + b^2(Ae(8d^2+28dex) + 2Bde^2x^2))}{105e^4(d+ex)^{7/2}}$$

input

```
Integrate[((a + b*x)^2*(A + B*x))/(d + e*x)^(9/2), x]
```

output

$$\frac{(-2*(3*a^2*e^2*(2*B*d + 5*A*e + 7*B*e*x) + 2*a*b*e*(3*A*e*(2*d + 7*e*x) + B*(8*d^2 + 28*d*e*x + 35*e^2*x^2)) + b^2*(A*e*(8*d^2 + 28*d*e*x + 35*e^2*x^2) + 3*B*(16*d^3 + 56*d^2*e*x + 70*d*e^2*x^2 + 35*e^3*x^3))))}{(105*e^4*(d + e*x)^(7/2))}$$

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^{9/2}} dx$$

↓ 86

$$\int \left(\frac{b(2aBe + Abe - 3bBd)}{e^3(d + ex)^{5/2}} + \frac{(ae - bd)(aBe + 2Abe - 3bBd)}{e^3(d + ex)^{7/2}} + \frac{(ae - bd)^2(Ae - Bd)}{e^3(d + ex)^{9/2}} + \frac{b^2B}{e^3(d + ex)^{3/2}} \right) dx$$

↓ 2009

$$\frac{2b(-2aBe - Abe + 3bBd)}{3e^4(d + ex)^{3/2}} - \frac{2(bd - ae)(-aBe - 2Abe + 3bBd)}{5e^4(d + ex)^{5/2}} + \frac{2(bd - ae)^2(Bd - Ae)}{7e^4(d + ex)^{7/2}} - \frac{2b^2B}{e^4\sqrt{d + ex}}$$

input

```
Int[((a + b*x)^2*(A + B*x))/(d + e*x)^(9/2), x]
```

output

$$\frac{(2*(b*d - a*e)^2*(B*d - A*e))/(7*e^4*(d + e*x)^(7/2)) - (2*(b*d - a*e)*(3*b*B*d - 2*A*b*e - a*B*e))/(5*e^4*(d + e*x)^(5/2)) + (2*b*(3*b*B*d - A*b*e - 2*a*B*e))/(3*e^4*(d + e*x)^(3/2)) - (2*b^2*B)/(e^4*sqrt[d + e*x])}$$

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0]) || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.94

method	result
pseudoelliptic	$\frac{(-70x^2(3Bx+A)b^2-84ax(\frac{5Bx}{3}+A)b-30a^2(\frac{7Bx}{5}+A))e^3-24\left(\frac{7x(\frac{15Bx}{2}+A)b^2}{3}+a\left(\frac{14Bx}{3}+A\right)b+\frac{a^2B}{2}\right)d e^2-16b((\dots))}{105(ex+d)^{\frac{7}{2}}e^4}$
derivativedivides	$-\frac{\frac{2b^2B}{\sqrt{ex+d}}}{e^4} - \frac{2(a^2Ae^3-2Aabd e^2+Ab^2d^2e-Ba^2d e^2+2Babd^2e-b^2Bd^3)}{7(ex+d)^{\frac{7}{2}}}$
default	$-\frac{\frac{2b^2B}{\sqrt{ex+d}}}{e^4} - \frac{2(a^2Ae^3-2Aabd e^2+Ab^2d^2e-Ba^2d e^2+2Babd^2e-b^2Bd^3)}{7(ex+d)^{\frac{7}{2}}} - \frac{2b(Abe+2Bae-3Bbd)}{3(ex+d)^{\frac{3}{2}}} - \frac{2(2Aab e^2-2Ab^2de+Ba^2e^2)}{5(ex+d)^{\frac{5}{2}}}$
gospers	$\frac{2(105b^2Bx^3e^3+35Ax^2b^2e^3+70Bx^2abe^3+210Bx^2b^2de^2+42Axabe^3+28Ax^2b^2de^2+21Bxa^2e^3+56Bxabde^2+16Bx^2a^2e^3)}{105(ex+d)^{\frac{7}{2}}e^4}$
trager	$\frac{2(105b^2Bx^3e^3+35Ax^2b^2e^3+70Bx^2abe^3+210Bx^2b^2de^2+42Axabe^3+28Ax^2b^2de^2+21Bxa^2e^3+56Bxabde^2+16Bx^2a^2e^3)}{105(ex+d)^{\frac{7}{2}}e^4}$
orering	$\frac{2(105b^2Bx^3e^3+35Ax^2b^2e^3+70Bx^2abe^3+210Bx^2b^2de^2+42Axabe^3+28Ax^2b^2de^2+21Bxa^2e^3+56Bxabde^2+16Bx^2a^2e^3)}{105(ex+d)^{\frac{7}{2}}e^4}$

```
input int((b*x+a)^2*(B*x+A)/(e*x+d)^(9/2), x, method=_RETURNVERBOSE)
```

```
output 1/105*((-70*x^2*(3*B*x+A)*b^2-84*a*x*(5/3*B*x+A)*b-30*a^2*(7/5*B*x+A))*e^3-24*(7/3*x*(15/2*B*x+A)*b^2+a*(14/3*B*x+A)*b+1/2*a^2*B)*d*e^2-16*b*((21*B*x+A)*b+2*B*a)*d^2*e-96*b^2*B*d^3)/(e*x+d)^(7/2)/e^4
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.57

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{9/2}} dx = \frac{2(105Bb^2e^3x^3 + 48Bb^2d^3 + 15Aa^2e^3 + 8(2Bab + Ab^2)d^2e + 6(Ba^2 + 2Aab)de^2 + 35(6Bb^2de^2 + (2Aa^2 + 2Aab)d^2e + 3Aa^2d^2))}{105(e^8x^4 + 4de^7x^3 + 6d^2e^6x^2 + \dots)}$$

input `integrate((b*x+a)^2*(B*x+A)/(e*x+d)^(9/2),x, algorithm="fricas")`

output `-2/105*(105*B*b^2*e^3*x^3 + 48*B*b^2*d^3 + 15*A*a^2*e^3 + 8*(2*B*a*b + A*b^2)*d^2*e + 6*(B*a^2 + 2*A*a*b)*d*e^2 + 35*(6*B*b^2*d*e^2 + (2*B*a*b + A*b^2)*e^3)*x^2 + 7*(24*B*b^2*d^2*e + 4*(2*B*a*b + A*b^2)*d*e^2 + 3*(B*a^2 + 2*A*a*b)*e^3)*x)*sqrt(e*x + d)/(e^8*x^4 + 4*d*e^7*x^3 + 6*d^2*e^6*x^2 + 4*d^3*e^5*x + d^4*e^4)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1323 vs. 2(128) = 256.

Time = 0.76 (sec) , antiderivative size = 1323, normalized size of antiderivative = 10.50

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{9/2}} dx = \text{Too large to display}$$

input `integrate((b*x+a)**2*(B*x+A)/(e*x+d)**(9/2),x)`

output

```
Piecewise((-30*A*a**2*e**3/(105*d**3*e**4*sqrt(d + e*x) + 315*d**2*e**5*x*
sqrt(d + e*x) + 315*d*e**6*x**2*sqrt(d + e*x) + 105*e**7*x**3*sqrt(d + e*x)
)) - 24*A*a*b*d*e**2/(105*d**3*e**4*sqrt(d + e*x) + 315*d**2*e**5*x*sqrt(d
+ e*x) + 315*d*e**6*x**2*sqrt(d + e*x) + 105*e**7*x**3*sqrt(d + e*x)) - 8
4*A*a*b*e**3*x/(105*d**3*e**4*sqrt(d + e*x) + 315*d**2*e**5*x*sqrt(d + e*x)
) + 315*d*e**6*x**2*sqrt(d + e*x) + 105*e**7*x**3*sqrt(d + e*x)) - 16*A*b*
**2*d**2*e/(105*d**3*e**4*sqrt(d + e*x) + 315*d**2*e**5*x*sqrt(d + e*x) + 3
15*d*e**6*x**2*sqrt(d + e*x) + 105*e**7*x**3*sqrt(d + e*x)) - 56*A*b**2*d*
e**2*x/(105*d**3*e**4*sqrt(d + e*x) + 315*d**2*e**5*x*sqrt(d + e*x) + 315*
d*e**6*x**2*sqrt(d + e*x) + 105*e**7*x**3*sqrt(d + e*x)) - 70*A*b**2*e**3*
x**2/(105*d**3*e**4*sqrt(d + e*x) + 315*d**2*e**5*x*sqrt(d + e*x) + 315*d*
e**6*x**2*sqrt(d + e*x) + 105*e**7*x**3*sqrt(d + e*x)) - 12*B*a**2*d*e**2/
(105*d**3*e**4*sqrt(d + e*x) + 315*d**2*e**5*x*sqrt(d + e*x) + 315*d*e**6*
x**2*sqrt(d + e*x) + 105*e**7*x**3*sqrt(d + e*x)) - 42*B*a**2*e**3*x/(105*
d**3*e**4*sqrt(d + e*x) + 315*d**2*e**5*x*sqrt(d + e*x) + 315*d*e**6*x**2*
sqrt(d + e*x) + 105*e**7*x**3*sqrt(d + e*x)) - 32*B*a*b*d**2*e/(105*d**3*e
**4*sqrt(d + e*x) + 315*d**2*e**5*x*sqrt(d + e*x) + 315*d*e**6*x**2*sqrt(d
+ e*x) + 105*e**7*x**3*sqrt(d + e*x)) - 112*B*a*b*d*e**2*x/(105*d**3*e**4
*sqrt(d + e*x) + 315*d**2*e**5*x*sqrt(d + e*x) + 315*d*e**6*x**2*sqrt(d +
e*x) + 105*e**7*x**3*sqrt(d + e*x)) - 140*B*a*b*e**3*x**2/(105*d**3*e**...
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.24

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^{9/2}} dx = \frac{2(105(ex + d)^3 Bb^2 - 15 Bb^2 d^3 + 15 Aa^2 e^3 + 15(2 Bab + Ab^2)d^2 e - 15(Ba^2 + 2 Aab)de^2 - 35(3 Bb^2 d + 105(ex + d)^{7/2})}{105(ex + d)^{7/2}}$$

input

```
integrate((b*x+a)^2*(B*x+A)/(e*x+d)^(9/2),x, algorithm="maxima")
```

output

```
-2/105*(105*(e*x + d)^3*B*b^2 - 15*B*b^2*d^3 + 15*A*a^2*e^3 + 15*(2*B*a*b
+ A*b^2)*d^2*e - 15*(B*a^2 + 2*A*a*b)*d*e^2 - 35*(3*B*b^2*d - (2*B*a*b + A
*b^2)*e)*(e*x + d)^2 + 21*(3*B*b^2*d^2 - 2*(2*B*a*b + A*b^2)*d*e + (B*a^2
+ 2*A*a*b)*e^2)*(e*x + d))/((e*x + d)^(7/2)*e^4)
```


Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.50

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{9/2}} dx =$$

$$2(105(ex+d)^3 Bb^2 - 105(ex+d)^2 Bb^2 d + 63(ex+d) Bb^2 d^2 - 15 Bb^2 d^3 + 70(ex+d)^2 B a b e + 35(ex+d)^2 B a^2 b^2 e - 84(ex+d) B a^2 b^2 d e - 42(ex+d) A a^2 b^2 d e + 30 B a^2 b^2 d^2 e + 15 A a^2 b^2 d^2 e + 21(ex+d) B a^2 e^2 + 42(ex+d) A a^2 b e^2 - 15 B a^2 d e^2 - 30 A a^2 b d e^2 + 15 A a^2 e^3) / ((ex+d)^{7/2} e^4)$$

input

```
integrate((b*x+a)^2*(B*x+A)/(e*x+d)^(9/2),x, algorithm="giac")
```

output

```
-2/105*(105*(e*x + d)^3*B*b^2 - 105*(e*x + d)^2*B*b^2*d + 63*(e*x + d)*B*b^2*d^2 - 15*B*b^2*d^3 + 70*(e*x + d)^2*B*a*b*e + 35*(e*x + d)^2*A*b^2*e - 84*(e*x + d)*B*a*b*d*e - 42*(e*x + d)*A*b^2*d*e + 30*B*a*b*d^2*e + 15*A*b^2*d^2*e + 21*(e*x + d)*B*a^2*e^2 + 42*(e*x + d)*A*a*b*e^2 - 15*B*a^2*d*e^2 - 30*A*a*b*d*e^2 + 15*A*a^2*e^3)/((e*x + d)^(7/2)*e^4)
```

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.19

$$\int \frac{(a+bx)^2(A+Bx)}{(d+ex)^{9/2}} dx = -\frac{2 A a^2}{7 e (d+e x)^{7/2}} - \frac{2 b (A b + 2 B a) (50 d^2 - 42 d (d+e x) + 35 e^2 x^2 + 70 d e x)}{105 e^3 (d+e x)^{7/2}} - \frac{2 B b^2 (21 d^2 (d+e x) - 5 d^3 + 35 e^3 x^3 + 70 d e^2 x^2 + 35 d^2 e x)}{35 e^4 (d+e x)^{7/2}} - \frac{2 a (2 d + 7 e x) (2 A b + B a)}{35 e^2 (d+e x)^{7/2}}$$

input

```
int(((A + B*x)*(a + b*x)^2)/(d + e*x)^(9/2),x)
```

output

```
- (2*A*a^2)/(7*e*(d + e*x)^(7/2)) - (2*b*(A*b + 2*B*a)*(50*d^2 - 42*d*(d +
e*x) + 35*e^2*x^2 + 70*d*e*x))/(105*e^3*(d + e*x)^(7/2)) - (2*B*b^2*(21*d
^2*(d + e*x) - 5*d^3 + 35*e^3*x^3 + 70*d*e^2*x^2 + 35*d^2*e*x))/(35*e^4*(d
+ e*x)^(7/2)) - (2*a*(2*d + 7*e*x)*(2*A*b + B*a))/(35*e^2*(d + e*x)^(7/2)
)
```

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.15

$$\int \frac{(a + bx)^2(A + Bx)}{(d + ex)^{9/2}} dx = \frac{-2b^3e^3x^3 - 2ab^2e^3x^2 - 4b^3de^2x^2 - \frac{6}{5}a^2be^3x - \frac{8}{5}ab^2de^2x - \frac{16}{5}b^3d^2ex - \frac{2}{7}a^3e^3}{\sqrt{ex + d}e^4(e^3x^3 + 3de^2x^2 + 3d^2ex + d^3)}$$

input

```
int((b*x+a)^2*(B*x+A)/(e*x+d)^(9/2),x)
```

output

```
(2*( - 5*a**3*e**3 - 6*a**2*b*d*e**2 - 21*a**2*b*e**3*x - 8*a*b**2*d**2*e
- 28*a*b**2*d*e**2*x - 35*a*b**2*e**3*x**2 - 16*b**3*d**3 - 56*b**3*d**2*e
*x - 70*b**3*d*e**2*x**2 - 35*b**3*e**3*x**3))/(35*sqrt(d + e*x)*e**4*(d**
3 + 3*d**2*e*x + 3*d*e**2*x**2 + e**3*x**3))
```

3.140 $\int (a + bx)^3 (A + Bx)(d + ex)^{5/2} dx$

Optimal result	1310
Mathematica [A] (verified)	1311
Rubi [A] (verified)	1311
Maple [A] (verified)	1313
Fricas [B] (verification not implemented)	1313
Sympy [B] (verification not implemented)	1314
Maxima [A] (verification not implemented)	1315
Giac [B] (verification not implemented)	1316
Mupad [B] (verification not implemented)	1317
Reduce [B] (verification not implemented)	1317

Optimal result

Integrand size = 22, antiderivative size = 173

$$\int (a + bx)^3 (A + Bx)(d + ex)^{5/2} dx = \frac{2(bd - ae)^3 (Bd - Ae)(d + ex)^{7/2}}{7e^5} - \frac{2(bd - ae)^2 (4bBd - 3Abe - aBe)(d + ex)^{9/2}}{9e^5} + \frac{6b(bd - ae)(2bBd - Abe - aBe)(d + ex)^{11/2}}{11e^5} - \frac{2b^2(4bBd - Abe - 3aBe)(d + ex)^{13/2}}{13e^5} + \frac{2b^3B(d + ex)^{15/2}}{15e^5}$$

output

```
2/7*(-a*e+b*d)^3*(-A*e+B*d)*(e*x+d)^(7/2)/e^5-2/9*(-a*e+b*d)^2*(-3*A*b*e-B
*a*e+4*B*b*d)*(e*x+d)^(9/2)/e^5+6/11*b*(-a*e+b*d)*(-A*b*e-B*a*e+2*B*b*d)*
(e*x+d)^(11/2)/e^5-2/13*b^2*(-A*b*e-3*B*a*e+4*B*b*d)*(e*x+d)^(13/2)/e^5+2/1
5*b^3*B*(e*x+d)^(15/2)/e^5
```

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.32

$$\int (a + bx)^3 (A + Bx) (d + ex)^{5/2} dx = \frac{2(d + ex)^{7/2} (715a^3e^3(-2Bd + 9Ae + 7Bex) + 195a^2be^2(11Ae(-2d + 7ex) + B(8d^2 - 28dex$$

input `Integrate[(a + b*x)^3*(A + B*x)*(d + e*x)^(5/2),x]`

output $(2*(d + e*x)^{(7/2)}*(715*a^3*e^3*(-2*B*d + 9*A*e + 7*B*e*x) + 195*a^2*b*e^2*(11*A*e*(-2*d + 7*e*x) + B*(8*d^2 - 28*d*e*x + 63*e^2*x^2)) - 15*a*b^2*e*(-13*A*e*(8*d^2 - 28*d*e*x + 63*e^2*x^2) + 3*B*(16*d^3 - 56*d^2*e*x + 126*d*e^2*x^2 - 231*e^3*x^3)) + b^3*(15*A*e*(-16*d^3 + 56*d^2*e*x - 126*d*e^2*x^2 + 231*e^3*x^3) + B*(128*d^4 - 448*d^3*e*x + 1008*d^2*e^2*x^2 - 1848*d*e^3*x^3 + 3003*e^4*x^4)))/(45045*e^5)$

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^3 (A + Bx) (d + ex)^{5/2} dx$$

↓ 86

$$\int \left(\frac{b^2(d + ex)^{11/2}(3aBe + Abe - 4bBd)}{e^4} - \frac{3b(d + ex)^{9/2}(bd - ae)(aBe + Abe - 2bBd)}{e^4} + \frac{(d + ex)^{7/2}(ae - bd)}{e^4} \right) dx$$

↓ 2009

$$\begin{aligned}
& -\frac{2b^2(d+ex)^{13/2}(-3aBe - Abe + 4bBd)}{13e^5} + \frac{6b(d+ex)^{11/2}(bd-ae)(-aBe - Abe + 2bBd)}{11e^5} - \\
& \frac{2(d+ex)^{9/2}(bd-ae)^2(-aBe - 3Abe + 4bBd)}{9e^5} + \frac{2(d+ex)^{7/2}(bd-ae)^3(Bd-Ae)}{7e^5} + \\
& \frac{2b^3B(d+ex)^{15/2}}{15e^5}
\end{aligned}$$

input `Int[(a + b*x)^3*(A + B*x)*(d + e*x)^(5/2), x]`

output `(2*(b*d - a*e)^3*(B*d - A*e)*(d + e*x)^(7/2))/(7*e^5) - (2*(b*d - a*e)^2*(4*b*B*d - 3*A*b*e - a*B*e)*(d + e*x)^(9/2))/(9*e^5) + (6*b*(b*d - a*e)*(2*b*B*d - A*b*e - a*B*e)*(d + e*x)^(11/2))/(11*e^5) - (2*b^2*(4*b*B*d - A*b*e - 3*a*B*e)*(d + e*x)^(13/2))/(13*e^5) + (2*b^3*B*(d + e*x)^(15/2))/(15*e^5)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`


```
input integrate((b*x+a)^3*(B*x+A)*(e*x+d)^(5/2),x, algorithm="fricas")
```

```
output 2/45045*(3003*B*b^3*e^7*x^7 + 128*B*b^3*d^7 + 6435*A*a^3*d^3*e^4 - 240*(3*
B*a*b^2 + A*b^3)*d^6*e + 1560*(B*a^2*b + A*a*b^2)*d^5*e^2 - 1430*(B*a^3 +
3*A*a^2*b)*d^4*e^3 + 231*(31*B*b^3*d*e^6 + 15*(3*B*a*b^2 + A*b^3)*e^7)*x^6
+ 63*(71*B*b^3*d^2*e^5 + 135*(3*B*a*b^2 + A*b^3)*d*e^6 + 195*(B*a^2*b + A
*a*b^2)*e^7)*x^5 + 35*(B*b^3*d^3*e^4 + 159*(3*B*a*b^2 + A*b^3)*d^2*e^5 + 8
97*(B*a^2*b + A*a*b^2)*d*e^6 + 143*(B*a^3 + 3*A*a^2*b)*e^7)*x^4 - 5*(8*B*b
^3*d^4*e^3 - 1287*A*a^3*e^7 - 15*(3*B*a*b^2 + A*b^3)*d^3*e^4 - 4407*(B*a^2
*b + A*a*b^2)*d^2*e^5 - 2717*(B*a^3 + 3*A*a^2*b)*d*e^6)*x^3 + 3*(16*B*b^3*d
^5*e^2 + 6435*A*a^3*d*e^6 - 30*(3*B*a*b^2 + A*b^3)*d^4*e^3 + 195*(B*a^2*b
+ A*a*b^2)*d^3*e^4 + 3575*(B*a^3 + 3*A*a^2*b)*d^2*e^5)*x^2 - (64*B*b^3*d^
6*e - 19305*A*a^3*d^2*e^5 - 120*(3*B*a*b^2 + A*b^3)*d^5*e^2 + 780*(B*a^2*b
+ A*a*b^2)*d^4*e^3 - 715*(B*a^3 + 3*A*a^2*b)*d^3*e^4)*x)*sqrt(e*x + d)/e^
5
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 423 vs. $2(170) = 340$.

Time = 1.28 (sec) , antiderivative size = 423, normalized size of antiderivative = 2.45

$$\int (a + bx)^3 (A + Bx) (d + ex)^{5/2} dx = \left\{ \frac{2 \left(\frac{Bb^3(d+ex)^{15}}{15e^4} + \frac{(d+ex)^{13}}{13e^4} (Ab^3e + 3Bab^2e - 4Bb^3d) + \frac{(d+ex)^{11}}{11e^4} (3Aab^2e^2 - 3Ab^3de + 3Ba^2be^2 - 9Bab^2de + 6Bb^3d^2) + \frac{(d+ex)^9}{9e^4} (3Aa^2b^2e^2 - 3Aab^3de + 3Ba^2be^2 - 9Bab^2de + 6Bb^3d^2) \right)}{d^{5/2} \left(Aa^3x + \frac{Bb^3x^5}{5} + \frac{x^4(Ab^3 + 3Bab^2)}{4} + \frac{x^3(3Aab^2 + 3Ba^2b)}{3} + \frac{x^2(3Aa^2b + Ba^3)}{2} \right)} \right.$$

```
input integrate((b*x+a)**3*(B*x+A)*(e*x+d)**(5/2),x)
```

output

```
Piecewise((2*(B*b**3*(d + e*x)**(15/2)/(15*e**4) + (d + e*x)**(13/2)*(A*b*
*3*e + 3*B*a*b**2*e - 4*B*b**3*d)/(13*e**4) + (d + e*x)**(11/2)*(3*A*a*b**
2*e**2 - 3*A*b**3*d*e + 3*B*a**2*b*e**2 - 9*B*a*b**2*d*e + 6*B*b**3*d**2)/
(11*e**4) + (d + e*x)**(9/2)*(3*A*a**2*b*e**3 - 6*A*a*b**2*d*e**2 + 3*A*b*
*3*d**2*e + B*a**3*e**3 - 6*B*a**2*b*d*e**2 + 9*B*a*b**2*d**2*e - 4*B*b**3
*d**3)/(9*e**4) + (d + e*x)**(7/2)*(A*a**3*e**4 - 3*A*a**2*b*d*e**3 + 3*A*
a*b**2*d**2*e**2 - A*b**3*d**3*e - B*a**3*d*e**3 + 3*B*a**2*b*d**2*e**2 -
3*B*a*b**2*d**3*e + B*b**3*d**4)/(7*e**4))/e, Ne(e, 0)), (d**(5/2)*(A*a**3
*x + B*b**3*x**5/5 + x**4*(A*b**3 + 3*B*a*b**2)/4 + x**3*(3*A*a*b**2 + 3*B
*a**2*b)/3 + x**2*(3*A*a**2*b + B*a**3)/2), True))
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 265, normalized size of antiderivative = 1.53

$$\int (a + bx)^3 (A + Bx) (d + ex)^{5/2} dx = \frac{2 \left(3003 (ex + d)^{\frac{15}{2}} Bb^3 - 3465 (4 Bb^3 d - (3 Bab^2 + Ab^3) e) (ex + d)^{\frac{13}{2}} + 12285 (2 Bb^3 d^2 - (3 B$$

input

```
integrate((b*x+a)^3*(B*x+A)*(e*x+d)^(5/2),x, algorithm="maxima")
```

output

```
2/45045*(3003*(e*x + d)^(15/2)*B*b^3 - 3465*(4*B*b^3*d - (3*B*a*b^2 + A*b^
3)*e)*(e*x + d)^(13/2) + 12285*(2*B*b^3*d^2 - (3*B*a*b^2 + A*b^3)*d*e + (B
*a^2*b + A*a*b^2)*e^2)*(e*x + d)^(11/2) - 5005*(4*B*b^3*d^3 - 3*(3*B*a*b^2
+ A*b^3)*d^2*e + 6*(B*a^2*b + A*a*b^2)*d*e^2 - (B*a^3 + 3*A*a^2*b)*e^3)*(
e*x + d)^(9/2) + 6435*(B*b^3*d^4 + A*a^3*e^4 - (3*B*a*b^2 + A*b^3)*d^3*e +
3*(B*a^2*b + A*a*b^2)*d^2*e^2 - (B*a^3 + 3*A*a^2*b)*d*e^3)*(e*x + d)^(7/2
))/e^5
```


Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1947 vs. $2(153) = 306$.

Time = 0.14 (sec) , antiderivative size = 1947, normalized size of antiderivative = 11.25

$$\int (a + bx)^3(A + Bx)(d + ex)^{5/2} dx = \text{Too large to display}$$

input `integrate((b*x+a)^3*(B*x+A)*(e*x+d)^(5/2),x, algorithm="giac")`

output

```
2/45045*(45045*sqrt(e*x + d)*A*a^3*d^3 + 45045*((e*x + d)^(3/2) - 3*sqrt(e
*x + d)*d)*A*a^3*d^2 + 15015*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*B*a^3*d
^3/e + 45045*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*A*a^2*b*d^3/e + 9009*(3
*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*A*a^3*d +
9009*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*B*a
^2*b*d^3/e^2 + 9009*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*
x + d)*d^2)*A*a*b^2*d^3/e^2 + 9009*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)
*d + 15*sqrt(e*x + d)*d^2)*B*a^3*d^2/e + 27027*(3*(e*x + d)^(5/2) - 10*(e*
x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*A*a^2*b*d^2/e + 1287*(5*(e*x + d)^(
7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^
3)*A*a^3 + 3861*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(
3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*B*a*b^2*d^3/e^3 + 1287*(5*(e*x + d)^(7/2)
- 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*A
*b^3*d^3/e^3 + 11583*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x +
d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*B*a^2*b*d^2/e^2 + 11583*(5*(e*x + d)
^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*
d^3)*A*a*b^2*d^2/e^2 + 3861*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35
*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*B*a^3*d/e + 11583*(5*(e*x + d)
^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)
*d^3)*A*a^2*b*d/e + 143*(35*(e*x + d)^(9/2) - 180*(e*x + d)^(7/2)*d + 3...
```

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.89

$$\int (a+bx)^3(A+Bx)(d+ex)^{5/2} dx = \frac{(d+ex)^{13/2}(2Ab^3e-8Bb^3d+6Bab^2e)}{13e^5} + \frac{2(ae-bd)^2(d+ex)^{9/2}(3Abe+BAe-4Bbd)}{9e^5} + \frac{2Bb^3(d+ex)^{15/2}}{15e^5} + \frac{2(Ae-Bd)(ae-bd)^3(d+ex)^{7/2}}{7e^5} + \frac{6b(ae-bd)(d+ex)^{11/2}(Abe+BAe-2Bbd)}{11e^5}$$

input `int((A + B*x)*(a + b*x)^3*(d + e*x)^(5/2), x)`output `((d + e*x)^(13/2)*(2*A*b^3*e - 8*B*b^3*d + 6*B*a*b^2*e))/(13*e^5) + (2*(a*e - b*d)^2*(d + e*x)^(9/2)*(3*A*b*e + B*a*e - 4*B*b*d))/(9*e^5) + (2*B*b^3*(d + e*x)^(15/2))/(15*e^5) + (2*(A*e - B*d)*(a*e - b*d)^3*(d + e*x)^(7/2))/(7*e^5) + (6*b*(a*e - b*d)*(d + e*x)^(11/2)*(A*b*e + B*a*e - 2*B*b*d))/(11*e^5)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 405, normalized size of antiderivative = 2.34

$$\int (a+bx)^3(A+Bx)(d+ex)^{5/2} dx = \frac{2\sqrt{ex+d}(3003b^4e^7x^7 + 13860ab^3e^7x^6 + 7161b^4de^6x^6 + 24570a^2b^2e^7x^5 + 34020ab^3de^6x^5 + \dots)}{11e^5}$$

input `int((b*x+a)^3*(B*x+A)*(e*x+d)^(5/2), x)`

output

```
(2*sqrt(d + e*x)*(6435*a**4*d**3*e**4 + 19305*a**4*d**2*e**5*x + 19305*a**4*d*e**6*x**2 + 6435*a**4*e**7*x**3 - 5720*a**3*b*d**4*e**3 + 2860*a**3*b*d**3*e**4*x + 42900*a**3*b*d**2*e**5*x**2 + 54340*a**3*b*d*e**6*x**3 + 20020*a**3*b*e**7*x**4 + 3120*a**2*b**2*d**5*e**2 - 1560*a**2*b**2*d**4*e**3*x + 1170*a**2*b**2*d**3*e**4*x**2 + 44070*a**2*b**2*d**2*e**5*x**3 + 62790*a**2*b**2*d*e**6*x**4 + 24570*a**2*b**2*e**7*x**5 - 960*a*b**3*d**6*e + 480*a*b**3*d**5*e**2*x - 360*a*b**3*d**4*e**3*x**2 + 300*a*b**3*d**3*e**4*x**3 + 22260*a*b**3*d**2*e**5*x**4 + 34020*a*b**3*d*e**6*x**5 + 13860*a*b**3*e**7*x**6 + 128*b**4*d**7 - 64*b**4*d**6*e*x + 48*b**4*d**5*e**2*x**2 - 40*b**4*d**4*e**3*x**3 + 35*b**4*d**3*e**4*x**4 + 4473*b**4*d**2*e**5*x**5 + 7161*b**4*d*e**6*x**6 + 3003*b**4*e**7*x**7))/(45045*e**5)
```

3.141 $\int (a + bx)^3 (A + Bx)(d + ex)^{3/2} dx$

Optimal result	1319
Mathematica [A] (verified)	1320
Rubi [A] (verified)	1320
Maple [A] (verified)	1322
Fricas [B] (verification not implemented)	1322
Sympy [B] (verification not implemented)	1323
Maxima [A] (verification not implemented)	1324
Giac [B] (verification not implemented)	1325
Mupad [B] (verification not implemented)	1326
Reduce [B] (verification not implemented)	1326

Optimal result

Integrand size = 22, antiderivative size = 173

$$\int (a + bx)^3 (A + Bx)(d + ex)^{3/2} dx = \frac{2(bd - ae)^3 (Bd - Ae)(d + ex)^{5/2}}{5e^5} - \frac{2(bd - ae)^2 (4bBd - 3Abe - aBe)(d + ex)^{7/2}}{7e^5} + \frac{2b(bd - ae)(2bBd - Abe - aBe)(d + ex)^{9/2}}{3e^5} - \frac{2b^2(4bBd - Abe - 3aBe)(d + ex)^{11/2}}{11e^5} + \frac{2b^3B(d + ex)^{13/2}}{13e^5}$$

output

```
2/5*(-a*e+b*d)^3*(-A*e+B*d)*(e*x+d)^(5/2)/e^5-2/7*(-a*e+b*d)^2*(-3*A*b*e-B
*a*e+4*B*b*d)*(e*x+d)^(7/2)/e^5+2/3*b*(-a*e+b*d)*(-A*b*e-B*a*e+2*B*b*d)*(e
*x+d)^(9/2)/e^5-2/11*b^2*(-A*b*e-3*B*a*e+4*B*b*d)*(e*x+d)^(11/2)/e^5+2/13*
b^3*B*(e*x+d)^(13/2)/e^5
```

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.32

$$\int (a + bx)^3 (A + Bx) (d + ex)^{3/2} dx = \frac{2(d + ex)^{5/2} (429a^3e^3(-2Bd + 7Ae + 5Bex) + 143a^2be^2(9Ae(-2d + 5ex) + B(8d^2 - 20dex$$

input `Integrate[(a + b*x)^3*(A + B*x)*(d + e*x)^(3/2),x]`

output
$$\frac{(2*(d + e*x)^{(5/2)}*(429*a^3*e^3*(-2*B*d + 7*A*e + 5*B*e*x) + 143*a^2*b*e^2*(9*A*e*(-2*d + 5*e*x) + B*(8*d^2 - 20*d*e*x + 35*e^2*x^2)) - 13*a*b^2*e*(-11*A*e*(8*d^2 - 20*d*e*x + 35*e^2*x^2) + 3*B*(16*d^3 - 40*d^2*e*x + 70*d*e^2*x^2 - 105*e^3*x^3)) + b^3*(13*A*e*(-16*d^3 + 40*d^2*e*x - 70*d*e^2*x^2 + 105*e^3*x^3) + B*(128*d^4 - 320*d^3*e*x + 560*d^2*e^2*x^2 - 840*d*e^3*x^3 + 1155*e^4*x^4)))/(15015*e^5)}$$

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^3 (A + Bx) (d + ex)^{3/2} dx$$

↓ 86

$$\int \left(\frac{b^2(d + ex)^{9/2}(3aBe + Abe - 4bBd)}{e^4} - \frac{3b(d + ex)^{7/2}(bd - ae)(aBe + Abe - 2bBd)}{e^4} + \frac{(d + ex)^{5/2}(ae - bd)^2}{e} \right) dx$$

↓ 2009

$$\frac{2b^2(d+ex)^{11/2}(-3aBe - Abe + 4bBd)}{11e^5} + \frac{2b(d+ex)^{9/2}(bd-ae)(-aBe - Abe + 2bBd)}{7e^5} - \frac{2(d+ex)^{7/2}(bd-ae)^2(-aBe - 3Abe + 4bBd)}{7e^5} + \frac{2(d+ex)^{5/2}(bd-ae)^3(Bd-Ae)}{5e^5} + \frac{2b^3B(d+ex)^{13/2}}{13e^5}$$

input `Int[(a + b*x)^3*(A + B*x)*(d + e*x)^(3/2), x]`

output `(2*(b*d - a*e)^3*(B*d - A*e)*(d + e*x)^(5/2))/(5*e^5) - (2*(b*d - a*e)^2*(4*b*B*d - 3*A*b*e - a*B*e)*(d + e*x)^(7/2))/(7*e^5) + (2*b*(b*d - a*e)*(2*b*B*d - A*b*e - a*B*e)*(d + e*x)^(9/2))/(3*e^5) - (2*b^2*(4*b*B*d - A*b*e - 3*a*B*e)*(d + e*x)^(11/2))/(11*e^5) + (2*b^3*B*(d + e*x)^(13/2))/(13*e^5)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;`
`FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

input `integrate((b*x+a)^3*(B*x+A)*(e*x+d)^(3/2),x, algorithm="fricas")`

output
$$\begin{aligned} & 2/15015*(1155*B*b^3*e^6*x^6 + 128*B*b^3*d^6 + 3003*A*a^3*d^2*e^4 - 208*(3* \\ & B*a*b^2 + A*b^3)*d^5*e + 1144*(B*a^2*b + A*a*b^2)*d^4*e^2 - 858*(B*a^3 + 3 \\ & *A*a^2*b)*d^3*e^3 + 105*(14*B*b^3*d*e^5 + 13*(3*B*a*b^2 + A*b^3)*e^6)*x^5 \\ & + 35*(B*b^3*d^2*e^4 + 52*(3*B*a*b^2 + A*b^3)*d*e^5 + 143*(B*a^2*b + A*a*b^ \\ & 2)*e^6)*x^4 - 5*(8*B*b^3*d^3*e^3 - 13*(3*B*a*b^2 + A*b^3)*d^2*e^4 - 1430*(\\ & B*a^2*b + A*a*b^2)*d*e^5 - 429*(B*a^3 + 3*A*a^2*b)*e^6)*x^3 + 3*(16*B*b^3* \\ & d^4*e^2 + 1001*A*a^3*e^6 - 26*(3*B*a*b^2 + A*b^3)*d^3*e^3 + 143*(B*a^2*b + \\ & A*a*b^2)*d^2*e^4 + 1144*(B*a^3 + 3*A*a^2*b)*d*e^5)*x^2 - (64*B*b^3*d^5*e \\ & - 6006*A*a^3*d*e^5 - 104*(3*B*a*b^2 + A*b^3)*d^4*e^2 + 572*(B*a^2*b + A*a* \\ & b^2)*d^3*e^3 - 429*(B*a^3 + 3*A*a^2*b)*d^2*e^4)*x)*sqrt(e*x + d)/e^5 \end{aligned}$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 423 vs. $2(170) = 340$.

Time = 1.17 (sec) , antiderivative size = 423, normalized size of antiderivative = 2.45

$$\int (a + bx)^3 (A + Bx) (d + ex)^{3/2} dx = \left\{ \begin{array}{l} 2 \left(\frac{Bb^3(d+ex)^{13}}{13e^4} + \frac{(d+ex)^{\frac{11}{2}} (Ab^3e+3Bab^2e-4Bb^3d)}{11e^4} + \frac{(d+ex)^{\frac{9}{2}} (3Aab^2e^2-3Ab^3de+3Ba^2be^2-9Bab^2de+6Bb^3d^2)}{9e^4} + \frac{(d+ex)^{\frac{7}{2}} (3Aa^2b^2+3Aa^2b^2+3Aa^2b^2)}{7e^4} \right) \\ d^{\frac{3}{2}} \left(Aa^3x + \frac{Bb^3x^5}{5} + \frac{x^4(Ab^3+3Bab^2)}{4} + \frac{x^3(3Aab^2+3Ba^2b)}{3} + \frac{x^2(3Aa^2b+Ba^3)}{2} \right) \end{array} \right.$$

input `integrate((b*x+a)**3*(B*x+A)*(e*x+d)**(3/2),x)`

output

```
Piecewise((2*(B*b**3*(d + e*x)**(13/2)/(13*e**4) + (d + e*x)**(11/2)*(A*b*
*3*e + 3*B*a*b**2*e - 4*B*b**3*d)/(11*e**4) + (d + e*x)**(9/2)*(3*A*a*b**2
*e**2 - 3*A*b**3*d*e + 3*B*a**2*b*e**2 - 9*B*a*b**2*d*e + 6*B*b**3*d**2)/(
9*e**4) + (d + e*x)**(7/2)*(3*A*a**2*b*e**3 - 6*A*a*b**2*d*e**2 + 3*A*b**3
*d**2*e + B*a**3*e**3 - 6*B*a**2*b*d*e**2 + 9*B*a*b**2*d**2*e - 4*B*b**3*d
**3)/(7*e**4) + (d + e*x)**(5/2)*(A*a**3*e**4 - 3*A*a**2*b*d*e**3 + 3*A*a*
b**2*d**2*e**2 - A*b**3*d**3*e - B*a**3*d*e**3 + 3*B*a**2*b*d**2*e**2 - 3*
B*a*b**2*d**3*e + B*b**3*d**4)/(5*e**4))/e, Ne(e, 0)), (d**(3/2)*(A*a**3*x
+ B*b**3*x**5/5 + x**4*(A*b**3 + 3*B*a*b**2)/4 + x**3*(3*A*a*b**2 + 3*B*a
**2*b)/3 + x**2*(3*A*a**2*b + B*a**3)/2), True))
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 265, normalized size of antiderivative = 1.53

$$\int (a + bx)^3 (A + Bx) (d + ex)^{3/2} dx = \frac{2 \left(1155 (ex + d)^{\frac{13}{2}} Bb^3 - 1365 (4 Bb^3 d - (3 Bab^2 + Ab^3) e) (ex + d)^{\frac{11}{2}} + 5005 (2 Bb^3 d^2 - (3 B$$

input

```
integrate((b*x+a)^3*(B*x+A)*(e*x+d)^(3/2),x, algorithm="maxima")
```

output

```
2/15015*(1155*(e*x + d)^(13/2)*B*b^3 - 1365*(4*B*b^3*d - (3*B*a*b^2 + A*b^
3)*e)*(e*x + d)^(11/2) + 5005*(2*B*b^3*d^2 - (3*B*a*b^2 + A*b^3)*d*e + (B*
a^2*b + A*a*b^2)*e^2)*(e*x + d)^(9/2) - 2145*(4*B*b^3*d^3 - 3*(3*B*a*b^2 +
A*b^3)*d^2*e + 6*(B*a^2*b + A*a*b^2)*d*e^2 - (B*a^3 + 3*A*a^2*b)*e^3)*(e*
x + d)^(7/2) + 3003*(B*b^3*d^4 + A*a^3*e^4 - (3*B*a*b^2 + A*b^3)*d^3*e + 3
*(B*a^2*b + A*a*b^2)*d^2*e^2 - (B*a^3 + 3*A*a^2*b)*d*e^3)*(e*x + d)^(5/2))
/e^5
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1306 vs. $2(153) = 306$.

Time = 0.14 (sec) , antiderivative size = 1306, normalized size of antiderivative = 7.55

$$\int (a + bx)^3(A + Bx)(d + ex)^{3/2} dx = \text{Too large to display}$$

input `integrate((b*x+a)^3*(B*x+A)*(e*x+d)^(3/2),x, algorithm="giac")`

output

```
2/45045*(45045*sqrt(e*x + d)*A*a^3*d^2 + 30030*((e*x + d)^(3/2) - 3*sqrt(e
*x + d)*d)*A*a^3*d + 15015*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*B*a^3*d^2
/e + 45045*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*A*a^2*b*d^2/e + 3003*(3*(
e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*A*a^3 + 9009
*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*B*a^2*b
*d^2/e^2 + 9009*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x +
d)*d^2)*A*a*b^2*d^2/e^2 + 6006*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d +
15*sqrt(e*x + d)*d^2)*B*a^3*d/e + 18018*(3*(e*x + d)^(5/2) - 10*(e*x + d)
^(3/2)*d + 15*sqrt(e*x + d)*d^2)*A*a^2*b*d/e + 3861*(5*(e*x + d)^(7/2) - 2
1*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*B*a*b
^2*d^2/e^3 + 1287*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x + d)
^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*A*b^3*d^2/e^3 + 7722*(5*(e*x + d)^(7/2)
- 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*B
*a^2*b*d/e^2 + 7722*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x +
d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*A*a*b^2*d/e^2 + 1287*(5*(e*x + d)^(7/
2) - 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)
*B*a^3/e + 3861*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(
3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*A*a^2*b/e + 143*(35*(e*x + d)^(9/2) - 180
*(e*x + d)^(7/2)*d + 378*(e*x + d)^(5/2)*d^2 - 420*(e*x + d)^(3/2)*d^3 + 3
15*sqrt(e*x + d)*d^4)*B*b^3*d^2/e^4 + 858*(35*(e*x + d)^(9/2) - 180*(e...
```

Mupad [B] (verification not implemented)

Time = 0.99 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.89

$$\int (a+bx)^3(A+Bx)(d+ex)^{3/2} dx = \frac{(d+ex)^{11/2}(2Ab^3e-8Bb^3d+6Bab^2e)}{11e^5} + \frac{2(ae-bd)^2(d+ex)^{7/2}(3Abe+BAe-4Bbd)}{7e^5} + \frac{2Bb^3(d+ex)^{13/2}}{13e^5} + \frac{2(Ae-Bd)(ae-bd)^3(d+ex)^{5/2}}{5e^5} + \frac{2b(ae-bd)(d+ex)^{9/2}(Abe+BAe-2Bbd)}{3e^5}$$

input `int((A + B*x)*(a + b*x)^3*(d + e*x)^(3/2), x)`

output

```
((d + e*x)^(11/2)*(2*A*b^3*e - 8*B*b^3*d + 6*B*a*b^2*e))/(11*e^5) + (2*(a*e - b*d)^2*(d + e*x)^(7/2)*(3*A*b*e + B*a*e - 4*B*b*d))/(7*e^5) + (2*B*b^3*(d + e*x)^(13/2))/(13*e^5) + (2*(A*e - B*d)*(a*e - b*d)^3*(d + e*x)^(5/2))/(5*e^5) + (2*b*(a*e - b*d)*(d + e*x)^(9/2)*(A*b*e + B*a*e - 2*B*b*d))/(3*e^5)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 330, normalized size of antiderivative = 1.91

$$\int (a+bx)^3(A+Bx)(d+ex)^{3/2} dx = \frac{2\sqrt{ex+d}(1155b^4e^6x^6 + 5460ab^3e^6x^5 + 1470b^4de^5x^5 + 10010a^2b^2e^6x^4 + 7280ab^3de^5x^4 + 35a^3b^2e^6x^3 + 210a^2b^3de^5x^3 + 105a^3b^2e^6x^2 + 70a^2b^3de^5x^2 + 35a^3b^2e^6x + 21a^2b^3de^5x + 7a^3b^2e^6 + 7a^2b^3de^5)}{35e^5}$$

input `int((b*x+a)^3*(B*x+A)*(e*x+d)^(3/2), x)`

output

```
(2*sqrt(d + e*x)*(3003*a**4*d**2*e**4 + 6006*a**4*d*e**5*x + 3003*a**4*e**6*x**2 - 3432*a**3*b*d**3*e**3 + 1716*a**3*b*d**2*e**4*x + 13728*a**3*b*d*e**5*x**2 + 8580*a**3*b*e**6*x**3 + 2288*a**2*b**2*d**4*e**2 - 1144*a**2*b**2*d**3*e**3*x + 858*a**2*b**2*d**2*e**4*x**2 + 14300*a**2*b**2*d*e**5*x**3 + 10010*a**2*b**2*e**6*x**4 - 832*a*b**3*d**5*e + 416*a*b**3*d**4*e**2*x - 312*a*b**3*d**3*e**3*x**2 + 260*a*b**3*d**2*e**4*x**3 + 7280*a*b**3*d*e**5*x**4 + 5460*a*b**3*e**6*x**5 + 128*b**4*d**6 - 64*b**4*d**5*e*x + 48*b**4*d**4*e**2*x**2 - 40*b**4*d**3*e**3*x**3 + 35*b**4*d**2*e**4*x**4 + 1470*b**4*d*e**5*x**5 + 1155*b**4*e**6*x**6))/(15015*e**5)
```

3.142 $\int (a + bx)^3 (A + Bx) \sqrt{d + ex} dx$

Optimal result	1328
Mathematica [A] (verified)	1329
Rubi [A] (verified)	1329
Maple [A] (verified)	1331
Fricas [B] (verification not implemented)	1331
Sympy [B] (verification not implemented)	1332
Maxima [A] (verification not implemented)	1333
Giac [B] (verification not implemented)	1333
Mupad [B] (verification not implemented)	1334
Reduce [B] (verification not implemented)	1335

Optimal result

Integrand size = 22, antiderivative size = 173

$$\int (a + bx)^3 (A + Bx) \sqrt{d + ex} dx = \frac{2(bd - ae)^3 (Bd - Ae)(d + ex)^{3/2}}{3e^5} - \frac{2(bd - ae)^2 (4bBd - 3Abe - aBe)(d + ex)^{5/2}}{5e^5} + \frac{6b(bd - ae)(2bBd - Abe - aBe)(d + ex)^{7/2}}{7e^5} - \frac{2b^2(4bBd - Abe - 3aBe)(d + ex)^{9/2}}{9e^5} + \frac{2b^3B(d + ex)^{11/2}}{11e^5}$$

output

```
2/3*(-a*e+b*d)^3*(-A*e+B*d)*(e*x+d)^(3/2)/e^5-2/5*(-a*e+b*d)^2*(-3*A*b*e-B
*a*e+4*B*b*d)*(e*x+d)^(5/2)/e^5+6/7*b*(-a*e+b*d)*(-A*b*e-B*a*e+2*B*b*d)*(e
*x+d)^(7/2)/e^5-2/9*b^2*(-A*b*e-3*B*a*e+4*B*b*d)*(e*x+d)^(9/2)/e^5+2/11*b^
3*B*(e*x+d)^(11/2)/e^5
```

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.31

$$\int (a + bx)^3 (A + Bx) \sqrt{d + ex} dx$$

$$= \frac{2(d + ex)^{3/2} (231a^3e^3(-2Bd + 5Ae + 3Bex) + 99a^2be^2(7Ae(-2d + 3ex) + B(8d^2 - 12dex + 15e^2x^2)) - 33ab^2e^2(-3Ae(8d^2 - 12d*ex + 15e^2x^2) + B(16d^3 - 24d^2*ex + 30d*ex^2 - 35e^3x^3)) + b^3(11Ae(-16d^3 + 24d^2*ex - 30d*ex^2 + 35e^3x^3) + B(128d^4 - 192d^3*ex + 240d^2*ex^2 - 280d*ex^3 + 315e^4x^4)))}{(3465e^5)}$$

input

```
Integrate[(a + b*x)^3*(A + B*x)*Sqrt[d + e*x], x]
```

output

```
(2*(d + e*x)^(3/2)*(231*a^3*e^3*(-2*B*d + 5*A*e + 3*B*e*x) + 99*a^2*b*e^2*(7*A*e*(-2*d + 3*e*x) + B*(8*d^2 - 12*d*e*x + 15*e^2*x^2)) - 33*a*b^2*e*(-3*A*e*(8*d^2 - 12*d*e*x + 15*e^2*x^2) + B*(16*d^3 - 24*d^2*e*x + 30*d*e^2*x^2 - 35*e^3*x^3)) + b^3*(11*A*e*(-16*d^3 + 24*d^2*e*x - 30*d*e^2*x^2 + 35*e^3*x^3) + B*(128*d^4 - 192*d^3*e*x + 240*d^2*e^2*x^2 - 280*d*e^3*x^3 + 315*e^4*x^4))))/(3465*e^5)
```

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^3 (A + Bx) \sqrt{d + ex} dx$$

↓ 86

$$\int \left(\frac{b^2(d + ex)^{7/2}(3aBe + Abe - 4bBd)}{e^4} - \frac{3b(d + ex)^{5/2}(bd - ae)(aBe + Abe - 2bBd)}{e^4} + \frac{(d + ex)^{3/2}(ae - bd)^2}{e} \right) dx$$

↓ 2009

$$\begin{aligned}
& - \frac{2b^2(d+ex)^{9/2}(-3aBe - Abe + 4bBd)}{9e^5} + \frac{6b(d+ex)^{7/2}(bd-ae)(-aBe - Abe + 2bBd)}{7e^5} - \\
& \frac{2(d+ex)^{5/2}(bd-ae)^2(-aBe - 3Abe + 4bBd)}{5e^5} + \frac{2(d+ex)^{3/2}(bd-ae)^3(Bd-Ae)}{3e^5} + \\
& \frac{2b^3B(d+ex)^{11/2}}{11e^5}
\end{aligned}$$

input `Int[(a + b*x)^3*(A + B*x)*Sqrt[d + e*x],x]`

output `(2*(b*d - a*e)^3*(B*d - A*e)*(d + e*x)^(3/2))/(3*e^5) - (2*(b*d - a*e)^2*(4*b*B*d - 3*A*b*e - a*B*e)*(d + e*x)^(5/2))/(5*e^5) + (6*b*(b*d - a*e)*(2*b*B*d - A*b*e - a*B*e)*(d + e*x)^(7/2))/(7*e^5) - (2*b^2*(4*b*B*d - A*b*e - 3*a*B*e)*(d + e*x)^(9/2))/(9*e^5) + (2*b^3*B*(d + e*x)^(11/2))/(11*e^5)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

input `integrate((b*x+a)^3*(B*x+A)*(e*x+d)^(1/2),x, algorithm="fricas")`

output
$$\frac{2}{3465} \cdot (315 \cdot B \cdot b^3 \cdot e^5 \cdot x^5 + 128 \cdot B \cdot b^3 \cdot d^5 + 1155 \cdot A \cdot a^3 \cdot d \cdot e^4 - 176 \cdot (3 \cdot B \cdot a \cdot b^2 + A \cdot b^3) \cdot d^4 \cdot e + 792 \cdot (B \cdot a^2 \cdot b + A \cdot a \cdot b^2) \cdot d^3 \cdot e^2 - 462 \cdot (B \cdot a^3 + 3 \cdot A \cdot a^2 \cdot b) \cdot d^2 \cdot e^3 + 35 \cdot (B \cdot b^3 \cdot d \cdot e^4 + 11 \cdot (3 \cdot B \cdot a \cdot b^2 + A \cdot b^3) \cdot e^5) \cdot x^4 - 5 \cdot (8 \cdot B \cdot b^3 \cdot d^2 \cdot e^3 - 11 \cdot (3 \cdot B \cdot a \cdot b^2 + A \cdot b^3) \cdot d \cdot e^4 - 297 \cdot (B \cdot a^2 \cdot b + A \cdot a \cdot b^2) \cdot e^5) \cdot x^3 + 3 \cdot (16 \cdot B \cdot b^3 \cdot d^3 \cdot e^2 - 22 \cdot (3 \cdot B \cdot a \cdot b^2 + A \cdot b^3) \cdot d^2 \cdot e^3 + 99 \cdot (B \cdot a^2 \cdot b + A \cdot a \cdot b^2) \cdot d \cdot e^4 + 231 \cdot (B \cdot a^3 + 3 \cdot A \cdot a^2 \cdot b) \cdot e^5) \cdot x^2 - (64 \cdot B \cdot b^3 \cdot d^4 \cdot e - 115 \cdot 5 \cdot A \cdot a^3 \cdot e^5 - 88 \cdot (3 \cdot B \cdot a \cdot b^2 + A \cdot b^3) \cdot d^3 \cdot e^2 + 396 \cdot (B \cdot a^2 \cdot b + A \cdot a \cdot b^2) \cdot d^2 \cdot e^3 - 231 \cdot (B \cdot a^3 + 3 \cdot A \cdot a^2 \cdot b) \cdot d \cdot e^4) \cdot x) \cdot \sqrt{e \cdot x + d} / e^5$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 423 vs. $2(170) = 340$.

Time = 1.12 (sec) , antiderivative size = 423, normalized size of antiderivative = 2.45

$$\int (a + bx)^3 (A + Bx) \sqrt{d + ex} dx$$

$$= \left\{ \frac{2 \left(\frac{Bb^3(d+ex)^{\frac{11}{2}}}{11e^4} + \frac{(d+ex)^{\frac{9}{2}} (Ab^3e + 3Bab^2e - 4Bb^3d)}{9e^4} + \frac{(d+ex)^{\frac{7}{2}} \cdot (3Aab^2e^2 - 3Ab^3de + 3Ba^2be^2 - 9Bab^2de + 6Bb^3d^2)}{7e^4} + \frac{(d+ex)^{\frac{5}{2}} \cdot (3Aa^2be^3 - 6Aab^2de^2 + \dots)}{e^4} \right)}{\sqrt{d} \left(Aa^3x + \frac{Bb^3x^5}{5} + \frac{x^4(Ab^3 + 3Bab^2)}{4} + \frac{x^3 \cdot (3Aab^2 + 3Ba^2b)}{3} + \frac{x^2 \cdot (3Aa^2b + Ba^3)}{2} \right)} \right.$$

input `integrate((b*x+a)**3*(B*x+A)*(e*x+d)**(1/2),x)`

output `Piecewise((2*(B*b**3*(d + e*x)**(11/2))/(11*e**4) + (d + e*x)**(9/2)*(A*b**3*e + 3*B*a*b**2*e - 4*B*b**3*d)/(9*e**4) + (d + e*x)**(7/2)*(3*A*a*b**2*e**2 - 3*A*b**3*d*e + 3*B*a**2*b*e**2 - 9*B*a*b**2*d*e + 6*B*b**3*d**2)/(7*e**4) + (d + e*x)**(5/2)*(3*A*a**2*b*e**3 - 6*A*a*b**2*d*e**2 + 3*A*b**3*d**2*e + B*a**3*e**3 - 6*B*a**2*b*d*e**2 + 9*B*a*b**2*d**2*e - 4*B*b**3*d**3)/(5*e**4) + (d + e*x)**(3/2)*(A*a**3*e**4 - 3*A*a**2*b*d*e**3 + 3*A*a*b**2*d**2*e**2 - A*b**3*d**3*e - B*a**3*d*e**3 + 3*B*a**2*b*d**2*e**2 - 3*B*a*b**2*d**3*e + B*b**3*d**4)/(3*e**4))/e, Ne(e, 0)), (sqrt(d)*(A*a**3*x + B*b**3*x**5/5 + x**4*(A*b**3 + 3*B*a*b**2)/4 + x**3*(3*A*a*b**2 + 3*B*a**2*b)/3 + x**2*(3*A*a**2*b + B*a**3)/2), True))`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 265, normalized size of antiderivative = 1.53

$$\int (a + bx)^3 (A + Bx) \sqrt{d + ex} dx$$

$$= \frac{2 \left(315 (ex + d)^{\frac{11}{2}} Bb^3 - 385 (4 Bb^3 d - (3 Bab^2 + Ab^3)e)(ex + d)^{\frac{9}{2}} + 1485 (2 Bb^3 d^2 - (3 Bab^2 + Ab^3)de \right)}{e^5}$$

input `integrate((b*x+a)^3*(B*x+A)*(e*x+d)^(1/2),x, algorithm="maxima")`

output `2/3465*(315*(e*x + d)^(11/2)*B*b^3 - 385*(4*B*b^3*d - (3*B*a*b^2 + A*b^3)*e)*(e*x + d)^(9/2) + 1485*(2*B*b^3*d^2 - (3*B*a*b^2 + A*b^3)*d*e + (B*a^2*b + A*a*b^2)*e^2)*(e*x + d)^(7/2) - 693*(4*B*b^3*d^3 - 3*(3*B*a*b^2 + A*b^3)*d^2*e + 6*(B*a^2*b + A*a*b^2)*d*e^2 - (B*a^3 + 3*A*a^2*b)*e^3)*(e*x + d)^(5/2) + 1155*(B*b^3*d^4 + A*a^3*e^4 - (3*B*a*b^2 + A*b^3)*d^3*e + 3*(B*a^2*b + A*a*b^2)*d^2*e^2 - (B*a^3 + 3*A*a^2*b)*d*e^3)*(e*x + d)^(3/2))/e^5`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 761 vs. 2(153) = 306.

Time = 0.13 (sec) , antiderivative size = 761, normalized size of antiderivative = 4.40

$$\int (a + bx)^3 (A + Bx) \sqrt{d + ex} dx = \text{Too large to display}$$

input `integrate((b*x+a)^3*(B*x+A)*(e*x+d)^(1/2),x, algorithm="giac")`

output

```

2/3465*(3465*sqrt(e*x + d)*A*a^3*d + 1155*((e*x + d)^(3/2) - 3*sqrt(e*x +
d)*d)*A*a^3 + 1155*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*B*a^3*d/e + 3465*
((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*A*a^2*b*d/e + 693*(3*(e*x + d)^(5/2)
- 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*B*a^2*b*d/e^2 + 693*(3*(e*
x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*A*a*b^2*d/e^2
+ 231*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*B*
a^3/e + 693*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d
^2)*A*a^2*b/e + 297*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x +
d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*B*a*b^2*d/e^3 + 99*(5*(e*x + d)^(7/2)
- 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*A
*b^3*d/e^3 + 297*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(
3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*B*a^2*b/e^2 + 297*(5*(e*x + d)^(7/2) - 2
1*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*A*a*b
^2/e^2 + 11*(35*(e*x + d)^(9/2) - 180*(e*x + d)^(7/2)*d + 378*(e*x + d)^(5
/2)*d^2 - 420*(e*x + d)^(3/2)*d^3 + 315*sqrt(e*x + d)*d^4)*B*b^3*d/e^4 + 3
3*(35*(e*x + d)^(9/2) - 180*(e*x + d)^(7/2)*d + 378*(e*x + d)^(5/2)*d^2 -
420*(e*x + d)^(3/2)*d^3 + 315*sqrt(e*x + d)*d^4)*B*a*b^2/e^3 + 11*(35*(e*x
+ d)^(9/2) - 180*(e*x + d)^(7/2)*d + 378*(e*x + d)^(5/2)*d^2 - 420*(e*x +
d)^(3/2)*d^3 + 315*sqrt(e*x + d)*d^4)*A*b^3/e^3 + 5*(63*(e*x + d)^(11/2)
- 385*(e*x + d)^(9/2)*d + 990*(e*x + d)^(7/2)*d^2 - 1386*(e*x + d)^(5/2)...

```

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.89

$$\begin{aligned}
 \int (a + bx)^3 (A + Bx) \sqrt{d + ex} dx = & \frac{(d + ex)^{9/2} (2Ab^3e - 8Bb^3d + 6Bab^2e)}{9e^5} \\
 & + \frac{2(ae - bd)^2 (d + ex)^{5/2} (3Abe + Bae - 4Bbd)}{5e^5} \\
 & + \frac{2Bb^3 (d + ex)^{11/2}}{11e^5} \\
 & + \frac{2(Ae - Bd) (ae - bd)^3 (d + ex)^{3/2}}{3e^5} \\
 & + \frac{6b(ae - bd) (d + ex)^{7/2} (Abe + Bae - 2Bbd)}{7e^5}
 \end{aligned}$$

input

```
int((A + B*x)*(a + b*x)^3*(d + e*x)^(1/2), x)
```

output

$$\begin{aligned} & ((d + ex)^{9/2} * (2A*b^3*e - 8B*b^3*d + 6B*a*b^2*e)) / (9*e^5) + (2*(a*e \\ & - b*d)^2 * (d + ex)^{5/2} * (3A*b*e + B*a*e - 4B*b*d)) / (5*e^5) + (2*B*b^3 * \\ & (d + ex)^{11/2}) / (11*e^5) + (2*(A*e - B*d) * (a*e - b*d)^3 * (d + ex)^{3/2}) / \\ & (3*e^5) + (6*b*(a*e - b*d) * (d + ex)^{7/2} * (A*b*e + B*a*e - 2*B*b*d)) / (7*e \\ & ^5) \end{aligned}$$
Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 255, normalized size of antiderivative = 1.47

$$\int (a + bx)^3 (A + Bx) \sqrt{d + ex} dx$$

$$= \frac{2\sqrt{ex + d} (315b^4e^5x^5 + 1540ab^3e^5x^4 + 35b^4de^4x^4 + 2970a^2b^2e^5x^3 + 220ab^3de^4x^3 - 40b^4d^2e^3x^3 + 2772$$

input

$$\text{int}((b*x+a)^3*(B*x+A)*(e*x+d)^{(1/2)}, x)$$

output

$$\begin{aligned} & (2*\text{sqrt}(d + e*x) * (1155*a**4*d*e**4 + 1155*a**4*e**5*x - 1848*a**3*b*d**2*e \\ & **3 + 924*a**3*b*d*e**4*x + 2772*a**3*b*e**5*x**2 + 1584*a**2*b**2*d**3*e \\ & *2 - 792*a**2*b**2*d**2*e**3*x + 594*a**2*b**2*d*e**4*x**2 + 2970*a**2*b** \\ & 2*e**5*x**3 - 704*a*b**3*d**4*e + 352*a*b**3*d**3*e**2*x - 264*a*b**3*d**2 \\ & *e**3*x**2 + 220*a*b**3*d*e**4*x**3 + 1540*a*b**3*e**5*x**4 + 128*b**4*d** \\ & 5 - 64*b**4*d**4*e*x + 48*b**4*d**3*e**2*x**2 - 40*b**4*d**2*e**3*x**3 + 3 \\ & 5*b**4*d*e**4*x**4 + 315*b**4*e**5*x**5)) / (3465*e**5) \end{aligned}$$

3.143 $\int \frac{(a+bx)^3(A+Bx)}{\sqrt{d+ex}} dx$

Optimal result	1336
Mathematica [A] (verified)	1337
Rubi [A] (verified)	1337
Maple [A] (verified)	1339
Fricas [A] (verification not implemented)	1339
Sympy [B] (verification not implemented)	1340
Maxima [A] (verification not implemented)	1341
Giac [B] (verification not implemented)	1341
Mupad [B] (verification not implemented)	1342
Reduce [B] (verification not implemented)	1343

Optimal result

Integrand size = 22, antiderivative size = 171

$$\int \frac{(a+bx)^3(A+Bx)}{\sqrt{d+ex}} dx = \frac{2(bd-ae)^3(Bd-Ae)\sqrt{d+ex}}{e^5} - \frac{2(bd-ae)^2(4bBd-3Abe-aBe)(d+ex)^{3/2}}{3e^5} + \frac{6b(bd-ae)(2bBd-Abe-aBe)(d+ex)^{5/2}}{5e^5} - \frac{2b^2(4bBd-Abe-3aBe)(d+ex)^{7/2}}{7e^5} + \frac{2b^3B(d+ex)^{9/2}}{9e^5}$$

output

```
2*(-a*e+b*d)^3*(-A*e+B*d)*(e*x+d)^(1/2)/e^5-2/3*(-a*e+b*d)^2*(-3*A*b*e-B*a
*e+4*B*b*d)*(e*x+d)^(3/2)/e^5+6/5*b*(-a*e+b*d)*(-A*b*e-B*a*e+2*B*b*d)*(e*x
+d)^(5/2)/e^5-2/7*b^2*(-A*b*e-3*B*a*e+4*B*b*d)*(e*x+d)^(7/2)/e^5+2/9*b^3*B
*(e*x+d)^(9/2)/e^5
```

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.32

$$\int \frac{(a + bx)^3(A + Bx)}{\sqrt{d + ex}} dx$$

$$= \frac{2\sqrt{d + ex}(105a^3e^3(-2Bd + 3Ae + Bex) + 63a^2be^2(5Ae(-2d + ex) + B(8d^2 - 4dex + 3e^2x^2)) - 9ab^2e^2(8d^2 - 4d^2ex + 3e^2x^2) + B(8d^2 - 4dex + 3e^2x^2)) - 9ab^2e^2(8d^2 - 4d^2ex + 3e^2x^2) + B(8d^2 - 4dex + 3e^2x^2))}{315e^5}$$

input

```
Integrate[((a + b*x)^3*(A + B*x))/Sqrt[d + e*x],x]
```

output

```
(2*Sqrt[d + e*x]*(105*a^3*e^3*(-2*B*d + 3*A*e + B*e*x) + 63*a^2*b*e^2*(5*A*e*(-2*d + e*x) + B*(8*d^2 - 4*d*e*x + 3*e^2*x^2)) - 9*a*b^2*e*(-7*A*e*(8*d^2 - 4*d*e*x + 3*e^2*x^2) + 3*B*(16*d^3 - 8*d^2*e*x + 6*d*e^2*x^2 - 5*e^3*x^3)) + b^3*(9*A*e*(-16*d^3 + 8*d^2*e*x - 6*d*e^2*x^2 + 5*e^3*x^3) + B*(128*d^4 - 64*d^3*e*x + 48*d^2*e^2*x^2 - 40*d*e^3*x^3 + 35*e^4*x^4)))/(315*e^5)
```

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^3(A + Bx)}{\sqrt{d + ex}} dx$$

$$\downarrow 86$$

$$\int \left(\frac{b^2(d + ex)^{5/2}(3aBe + Abe - 4bBd)}{e^4} - \frac{3b(d + ex)^{3/2}(bd - ae)(aBe + Abe - 2bBd)}{e^4} + \frac{\sqrt{d + ex}(ae - bd)^2(a + bx)}{e^4} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
& -\frac{2b^2(d+ex)^{7/2}(-3aBe - Abe + 4bBd)}{7e^5} + \frac{6b(d+ex)^{5/2}(bd-ae)(-aBe - Abe + 2bBd)}{5e^5} - \\
& \frac{2(d+ex)^{3/2}(bd-ae)^2(-aBe - 3Abe + 4bBd)}{3e^5} + \frac{2\sqrt{d+ex}(bd-ae)^3(Bd-Ae)}{e^5} + \\
& \frac{2b^3B(d+ex)^{9/2}}{9e^5}
\end{aligned}$$

input `Int[((a + b*x)^3*(A + B*x))/Sqrt[d + e*x], x]`

output `(2*(b*d - a*e)^3*(B*d - A*e)*Sqrt[d + e*x])/e^5 - (2*(b*d - a*e)^2*(4*b*B*d - 3*A*b*e - a*B*e)*(d + e*x)^(3/2))/(3*e^5) + (6*b*(b*d - a*e)*(2*b*B*d - A*b*e - a*B*e)*(d + e*x)^(5/2))/(5*e^5) - (2*b^2*(4*b*B*d - A*b*e - 3*a*B*e)*(d + e*x)^(7/2))/(7*e^5) + (2*b^3*B*(d + e*x)^(9/2))/(9*e^5)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

output

```
2/315*(35*B*b^3*e^4*x^4 + 128*B*b^3*d^4 + 315*A*a^3*e^4 - 144*(3*B*a*b^2 +
A*b^3)*d^3*e + 504*(B*a^2*b + A*a*b^2)*d^2*e^2 - 210*(B*a^3 + 3*A*a^2*b)*
d*e^3 - 5*(8*B*b^3*d*e^3 - 9*(3*B*a*b^2 + A*b^3)*e^4)*x^3 + 3*(16*B*b^3*d^
2*e^2 - 18*(3*B*a*b^2 + A*b^3)*d*e^3 + 63*(B*a^2*b + A*a*b^2)*e^4)*x^2 - (
64*B*b^3*d^3*e - 72*(3*B*a*b^2 + A*b^3)*d^2*e^2 + 252*(B*a^2*b + A*a*b^2)*
d*e^3 - 105*(B*a^3 + 3*A*a^2*b)*e^4)*x)*sqrt(e*x + d)/e^5
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 422 vs. $2(168) = 336$.

Time = 1.06 (sec) , antiderivative size = 422, normalized size of antiderivative = 2.47

$$\int \frac{(a + bx)^3(A + Bx)}{\sqrt{d + ex}} dx$$

$$= \left\{ \begin{array}{l} 2 \left(\frac{Bb^3(d+ex)^{\frac{9}{2}}}{9e^4} + \frac{(d+ex)^{\frac{7}{2}}(Ab^3e+3Bab^2e-4Bb^3d)}{7e^4} + \frac{(d+ex)^{\frac{5}{2}}(3Aab^2e^2-3Ab^3de+3Ba^2be^2-9Bab^2de+6Bb^3d^2)}{5e^4} + \frac{(d+ex)^{\frac{3}{2}}(3Aa^2be^3-6Aab^2de^2+3Aa^3e^3)}{3e^4} \right) \\ \frac{Aa^3x + \frac{Bb^3x^5}{5} + \frac{x^4(Ab^3+3Bab^2)}{4} + \frac{x^3(3Aab^2+3Ba^2b)}{3} + \frac{x^2(3Aa^2b+Ba^3)}{2}}{\sqrt{d}} \end{array} \right.$$

input

```
integrate((b*x+a)**3*(B*x+A)/(e*x+d)**(1/2), x)
```

output

```
Piecewise((2*(B*b**3*(d + e*x)**(9/2)/(9*e**4) + (d + e*x)**(7/2)*(A*b**3*
e + 3*B*a*b**2*e - 4*B*b**3*d)/(7*e**4) + (d + e*x)**(5/2)*(3*A*a*b**2*e**
2 - 3*A*b**3*d*e + 3*B*a**2*b*e**2 - 9*B*a*b**2*d*e + 6*B*b**3*d**2)/(5*e*
**4) + (d + e*x)**(3/2)*(3*A*a**2*b*e**3 - 6*A*a*b**2*d*e**2 + 3*A*b**3*d**
2*e + B*a**3*e**3 - 6*B*a**2*b*d*e**2 + 9*B*a*b**2*d**2*e - 4*B*b**3*d**3)
/(3*e**4) + sqrt(d + e*x)*(A*a**3*e**4 - 3*A*a**2*b*d*e**3 + 3*A*a*b**2*d*
**2*e**2 - A*b**3*d**3*e - B*a**3*d*e**3 + 3*B*a**2*b*d**2*e**2 - 3*B*a*b**
2*d**3*e + B*b**3*d**4)/e**4)/e, Ne(e, 0)), ((A*a**3*x + B*b**3*x**5/5 + x
**4*(A*b**3 + 3*B*a*b**2)/4 + x**3*(3*A*a*b**2 + 3*B*a**2*b)/3 + x**2*(3*A
*a**2*b + B*a**3)/2)/sqrt(d), True))
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 265, normalized size of antiderivative = 1.55

$$\int \frac{(a+bx)^3(A+Bx)}{\sqrt{d+ex}} dx$$

$$= \frac{2 \left(35 (ex+d)^{\frac{9}{2}} Bb^3 - 45 (4 Bb^3 d - (3 Bab^2 + Ab^3)e)(ex+d)^{\frac{7}{2}} + 189 (2 Bb^3 d^2 - (3 Bab^2 + Ab^3)de + (E$$

input `integrate((b*x+a)^3*(B*x+A)/(e*x+d)^(1/2),x, algorithm="maxima")`

output

$$\frac{2}{315} \cdot (35 \cdot (ex+d)^{\frac{9}{2}} \cdot B \cdot b^3 - 45 \cdot (4 \cdot B \cdot b^3 \cdot d - (3 \cdot B \cdot a \cdot b^2 + A \cdot b^3) \cdot e) \cdot (ex+d)^{\frac{7}{2}} + 189 \cdot (2 \cdot B \cdot b^3 \cdot d^2 - (3 \cdot B \cdot a \cdot b^2 + A \cdot b^3) \cdot d \cdot e + (B \cdot a^2 \cdot b + A \cdot a \cdot b^2) \cdot e^2) \cdot (ex+d)^{\frac{5}{2}} - 105 \cdot (4 \cdot B \cdot b^3 \cdot d^3 - 3 \cdot (3 \cdot B \cdot a \cdot b^2 + A \cdot b^3) \cdot d^2 \cdot e + 6 \cdot (B \cdot a^2 \cdot b + A \cdot a \cdot b^2) \cdot d \cdot e^2 - (B \cdot a^3 + 3 \cdot A \cdot a^2 \cdot b) \cdot e^3) \cdot (ex+d)^{\frac{3}{2}} + 315 \cdot (B \cdot b^3 \cdot d^4 + A \cdot a^3 \cdot e^4 - (3 \cdot B \cdot a \cdot b^2 + A \cdot b^3) \cdot d^3 \cdot e + 3 \cdot (B \cdot a^2 \cdot b + A \cdot a \cdot b^2) \cdot d^2 \cdot e^2 - (B \cdot a^3 + 3 \cdot A \cdot a^2 \cdot b) \cdot d \cdot e^3) \cdot \sqrt{ex+d}) / e^5$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 329 vs. 2(153) = 306.

Time = 0.12 (sec) , antiderivative size = 329, normalized size of antiderivative = 1.92

$$\int \frac{(a+bx)^3(A+Bx)}{\sqrt{d+ex}} dx$$

$$= \frac{2 \left(315 \sqrt{ex+d} A a^3 + \frac{105 \left((ex+d)^{\frac{3}{2}} - 3 \sqrt{ex+dd} \right) B a^3}{e} + \frac{315 \left((ex+d)^{\frac{3}{2}} - 3 \sqrt{ex+dd} \right) A a^2 b}{e} + \frac{63 \left(3 (ex+d)^{\frac{5}{2}} - 10 (ex+d)^{\frac{3}{2}} d + 15 \sqrt{ex+d} \right) A a b^2}{e^2} \right)}{e^2}$$

input `integrate((b*x+a)^3*(B*x+A)/(e*x+d)^(1/2),x, algorithm="giac")`

output

```
2/315*(315*sqrt(e*x + d)*A*a^3 + 105*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)
*B*a^3/e + 315*((e*x + d)^(3/2) - 3*sqrt(e*x + d)*d)*A*a^2*b/e + 63*(3*(e*
x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*B*a^2*b/e^2 +
63*(3*(e*x + d)^(5/2) - 10*(e*x + d)^(3/2)*d + 15*sqrt(e*x + d)*d^2)*A*a*b
^2/e^2 + 27*(5*(e*x + d)^(7/2) - 21*(e*x + d)^(5/2)*d + 35*(e*x + d)^(3/2)
*d^2 - 35*sqrt(e*x + d)*d^3)*B*a*b^2/e^3 + 9*(5*(e*x + d)^(7/2) - 21*(e*x
+ d)^(5/2)*d + 35*(e*x + d)^(3/2)*d^2 - 35*sqrt(e*x + d)*d^3)*A*b^3/e^3 +
(35*(e*x + d)^(9/2) - 180*(e*x + d)^(7/2)*d + 378*(e*x + d)^(5/2)*d^2 - 42
0*(e*x + d)^(3/2)*d^3 + 315*sqrt(e*x + d)*d^4)*B*b^3/e^4)/e
```

Mupad [B] (verification not implemented)

Time = 1.00 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.90

$$\int \frac{(a + bx)^3(A + Bx)}{\sqrt{d + ex}} dx = \frac{(d + ex)^{7/2} (2Ab^3e - 8Bb^3d + 6Bab^2e)}{7e^5} + \frac{2(ae - bd)^2 (d + ex)^{3/2} (3Abe + Bae - 4Bbd)}{3e^5} + \frac{2Bb^3 (d + ex)^{9/2}}{9e^5} + \frac{2(Ae - Bd)(ae - bd)^3 \sqrt{d + ex}}{e^5} + \frac{6b(ae - bd)(d + ex)^{5/2} (Abe + Bae - 2Bbd)}{5e^5}$$

input

```
int(((A + B*x)*(a + b*x)^3)/(d + e*x)^(1/2),x)
```

output

```
((d + e*x)^(7/2)*(2*A*b^3*e - 8*B*b^3*d + 6*B*a*b^2*e))/(7*e^5) + (2*(a*e
- b*d)^2*(d + e*x)^(3/2)*(3*A*b*e + B*a*e - 4*B*b*d))/(3*e^5) + (2*B*b^3*(
d + e*x)^(9/2))/(9*e^5) + (2*(A*e - B*d)*(a*e - b*d)^3*(d + e*x)^(1/2))/e^
5 + (6*b*(a*e - b*d)*(d + e*x)^(5/2)*(A*b*e + B*a*e - 2*B*b*d))/(5*e^5)
```


3.144 $\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{3/2}} dx$

Optimal result	1344
Mathematica [A] (verified)	1345
Rubi [A] (verified)	1345
Maple [A] (verified)	1346
Fricas [A] (verification not implemented)	1347
Sympy [B] (verification not implemented)	1347
Maxima [A] (verification not implemented)	1348
Giac [B] (verification not implemented)	1349
Mupad [B] (verification not implemented)	1349
Reduce [B] (verification not implemented)	1350

Optimal result

Integrand size = 22, antiderivative size = 167

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{3/2}} dx = -\frac{2(bd-ae)^3(Bd-Ae)}{e^5\sqrt{d+ex}} - \frac{2(bd-ae)^2(4bBd-3Abe-aBe)\sqrt{d+ex}}{e^5} + \frac{2b(bd-ae)(2bBd-Abe-aBe)(d+ex)^{3/2}}{e^5} - \frac{2b^2(4bBd-Abe-3aBe)(d+ex)^{5/2}}{5e^5} + \frac{2b^3B(d+ex)^{7/2}}{7e^5}$$

output

```
-2*(-a*e+b*d)^3*(-A*e+B*d)/e^5/(e*x+d)^(1/2)-2*(-a*e+b*d)^2*(-3*A*b*e-B*a*
e+4*B*b*d)*(e*x+d)^(1/2)/e^5+2*b*(-a*e+b*d)*(-A*b*e-B*a*e+2*B*b*d)*(e*x+d)
^(3/2)/e^5-2/5*b^2*(-A*b*e-3*B*a*e+4*B*b*d)*(e*x+d)^(5/2)/e^5+2/7*b^3*B*(e
*x+d)^(7/2)/e^5
```

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.33

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{3/2}} dx = \frac{2(35a^3e^3(2Bd - Ae + Bex) + 35a^2be^2(3Ae(2d + ex) + B(-8d^2 - 4dex + e^2x^2)) + 7ab^2e(5Ae(-8d^2 - 4dex + e^2x^2) + 3B(16d^3 + 8d^2ex - 2de^2x^2 + e^3x^3)) + b^3(7Ae(16d^3 + 8d^2ex - 2de^2x^2 + e^3x^3) + B(-128d^4 - 64d^3ex + 16d^2e^2x^2 - 8de^3x^3 + 5e^4x^4)))}{35e^5\sqrt{d+ex}}$$

input

```
Integrate[((a + b*x)^3*(A + B*x))/(d + e*x)^(3/2),x]
```

output

```
(2*(35*a^3*e^3*(2*B*d - A*e + B*e*x) + 35*a^2*b*e^2*(3*A*e*(2*d + e*x) + B*(-8*d^2 - 4*d*e*x + e^2*x^2)) + 7*a*b^2*e*(5*A*e*(-8*d^2 - 4*d*e*x + e^2*x^2) + 3*B*(16*d^3 + 8*d^2*e*x - 2*d*e^2*x^2 + e^3*x^3)) + b^3*(7*A*e*(16*d^3 + 8*d^2*e*x - 2*d*e^2*x^2 + e^3*x^3) + B*(-128*d^4 - 64*d^3*e*x + 16*d^2*e^2*x^2 - 8*d*e^3*x^3 + 5*e^4*x^4)))/(35*e^5*sqrt[d + e*x])
```

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{3/2}} dx$$

↓ 86

$$\int \left(\frac{b^2(d+ex)^{3/2}(3aBe + Abe - 4bBd)}{e^4} - \frac{3b\sqrt{d+ex}(bd - ae)(aBe + Abe - 2bBd)}{e^4} + \frac{(ae - bd)^2(aBe + 3Abe)}{e^4\sqrt{d+ex}} \right)$$

↓ 2009

$$\frac{2b^2(d+ex)^{5/2}(-3aBe - Abe + 4bBd)}{5e^5} + \frac{2b(d+ex)^{3/2}(bd - ae)(-aBe - Abe + 2bBd)}{e^5} - \frac{2\sqrt{d+ex}(bd - ae)^2(-aBe - 3Abe + 4bBd)}{e^5} - \frac{2(bd - ae)^3(Bd - Ae)}{e^5\sqrt{d+ex}} + \frac{2b^3B(d+ex)^{7/2}}{7e^5}$$

input `Int[((a + b*x)^3*(A + B*x))/(d + e*x)^(3/2), x]`

output
$$\frac{(-2*(b*d - a*e)^3*(B*d - A*e))/(e^5*\text{Sqrt}[d + e*x]) - (2*(b*d - a*e)^2*(4*b*B*d - 3*A*b*e - a*B*e)*\text{Sqrt}[d + e*x])/e^5 + (2*b*(b*d - a*e)*(2*b*B*d - A*b*e - a*B*e)*(d + e*x)^(3/2))/e^5 - (2*b^2*(4*b*B*d - A*b*e - 3*a*B*e)*(d + e*x)^(5/2))/(5*e^5) + (2*b^3*B*(d + e*x)^(7/2))/(7*e^5)}$$

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.))^(n_.)*((e_.) + (f_.)*(x_.))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;`
`FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.13

method	result
pseudoelliptic	$\frac{2\left(\left(-\frac{x^3\left(\frac{5Bx+A}{7}\right)b^3}{5}-\left(\frac{3Bx+A}{5}\right)ax^2b^2-3a^2\left(\frac{Bx+A}{3}\right)xb+a^3(-Bx+A)\right)e^4-6\left(-\frac{\left(\frac{4Bx+A}{7}\right)x^2b^3}{15}-\frac{2a\left(\frac{3Bx+A}{10}\right)}{3}\right)e^4\right)}{\dots}$
risch	$\frac{2(5b^3Bx^3e^3+7Ab^3e^3x^2+21Ba^2b^2e^3x^2-13Bb^3de^2x^2+35Aaxb^2e^3-21Ax^2b^2de^2+35Bxa^2be^3-63Ba^2b^2de^2x+29Ba^3e^3)}{35e^5}$
gospert	$\frac{2(-5Bx^4b^3e^4-7Ax^3b^3e^4-21Bx^3ab^2e^4+8Bx^3b^3de^3-35A^2x^2ab^2e^4+14Ax^2b^3de^3-35Bx^2a^2be^4+42Bx^2ab^2de^3)}{\dots}$
trager	$\frac{2(-5Bx^4b^3e^4-7Ax^3b^3e^4-21Bx^3ab^2e^4+8Bx^3b^3de^3-35A^2x^2ab^2e^4+14Ax^2b^3de^3-35Bx^2a^2be^4+42Bx^2ab^2de^3)}{\dots}$
orering	$\frac{2(-5Bx^4b^3e^4-7Ax^3b^3e^4-21Bx^3ab^2e^4+8Bx^3b^3de^3-35A^2x^2ab^2e^4+14Ax^2b^3de^3-35Bx^2a^2be^4+42Bx^2ab^2de^3)}{\dots}$
derivativedivides	$\frac{2b^3B(e^7x+d)^{\frac{7}{2}}+2Ab^3e(e^5x+d)^{\frac{5}{2}}+6Ba^2b^2e(e^5x+d)^{\frac{5}{2}}-8Bb^3d(e^5x+d)^{\frac{5}{2}}+2Aa^2b^2e^2(e^3x+d)^{\frac{3}{2}}-2Ab^3de(e^3x+d)^{\frac{3}{2}}+2Ba^2b^2e^2(e^3x+d)^{\frac{3}{2}}}{\dots}$
default	$\frac{2b^3B(e^7x+d)^{\frac{7}{2}}+2Ab^3e(e^5x+d)^{\frac{5}{2}}+6Ba^2b^2e(e^5x+d)^{\frac{5}{2}}-8Bb^3d(e^5x+d)^{\frac{5}{2}}+2Aa^2b^2e^2(e^3x+d)^{\frac{3}{2}}-2Ab^3de(e^3x+d)^{\frac{3}{2}}+2Ba^2b^2e^2(e^3x+d)^{\frac{3}{2}}}{\dots}$

```
input int((b*x+a)^3*(B*x+A)/(e*x+d)^(3/2),x,method=_RETURNVERBOSE)
```

```
output -2*((-1/5*x^3*(5/7*B*x+A)*b^3-(3/5*B*x+A)*a*x^2*b^2-3*a^2*(1/3*B*x+A)*x*b+a^3*(-B*x+A))*e^4-6*(-1/15*(4/7*B*x+A)*x^2*b^3-2/3*a*(3/10*B*x+A)*x*b^2+a^2*(-2/3*B*x+A)*b+1/3*a^3*B)*d*e^3+8*b*(-1/5*(2/7*B*x+A)*x*b^2+a*(-3/5*B*x+A)*b+a^2*B)*d^2*e^2-16/5*b^2*((-4/7*B*x+A)*b+3*B*a)*d^3*e+128/35*b^3*B*d^4)/(e*x+d)^(1/2)/e^5
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 272, normalized size of antiderivative = 1.63

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{3/2}} dx = \frac{2(5Bb^3e^4x^4 - 128Bb^3d^4 - 35Aa^3e^4 + 112(3Bab^2 + Ab^3)d^3e - 280(Ba^2b + Ab^3)d^2e^2 + 70(Ba^3 + 3Aa^2b)d^2e^3 - (8Bb^3d^3e^3 - 7(3Bba^2b^2 + Ab^3)e^4)x^3 + (16Bb^3d^2e^2 - 14(3Bba^2b^2 + Ab^3)d^2e^3 + 35(Ba^2b + Aa^2b^2)e^4)x^2 - (64Bb^3d^3e - 56(3Bba^2b^2 + Ab^3)d^2e^2 + 140(Ba^2b + Aa^2b^2)d^2e^3 - 35(Ba^3 + 3Aa^2b)e^4)x)*\sqrt{ex + d}}{(e^6x + d^6e^5)}$$

```
input integrate((b*x+a)^3*(B*x+A)/(e*x+d)^(3/2),x, algorithm="fricas")
```

```
output 2/35*(5*B*b^3*e^4*x^4 - 128*B*b^3*d^4 - 35*A*a^3*e^4 + 112*(3*B*a*b^2 + A*b^3)*d^3*e - 280*(B*a^2*b + A*a*b^2)*d^2*e^2 + 70*(B*a^3 + 3*A*a^2*b)*d^2*e^3 - (8*B*b^3*d^3*e^3 - 7*(3*B*a*b^2 + A*b^3)*e^4)*x^3 + (16*B*b^3*d^2*e^2 - 14*(3*B*a*b^2 + A*b^3)*d^2*e^3 + 35*(B*a^2*b + A*a*b^2)*e^4)*x^2 - (64*B*b^3*d^3*e - 56*(3*B*a*b^2 + A*b^3)*d^2*e^2 + 140*(B*a^2*b + A*a*b^2)*d^2*e^3 - 35*(B*a^3 + 3*A*a^2*b)*e^4)*x)*sqrt(e*x + d)/(e^6*x + d^6e^5)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 340 vs. 2(165) = 330.

Time = 9.55 (sec) , antiderivative size = 340, normalized size of antiderivative = 2.04

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{3/2}} dx = \frac{2 \left(\frac{Bb^3(d+ex)^{\frac{7}{2}}}{7e^4} + \frac{(d+ex)^{\frac{5}{2}}(Ab^3e+3Bab^2e-4Bb^3d)}{5e^4} + \frac{(d+ex)^{\frac{3}{2}}(3Aab^2e^2-3Ab^3de+3Ba^2be^2-9Bab^2de+6Bb^3d^2e)}{3e^4} \right) + \frac{Aa^3x + \frac{Bb^3x^5}{5} + \frac{x^4(Ab^3+3Bab^2)}{4} + \frac{x^3(3Aab^2+3Ba^2b)}{3} + \frac{x^2(3Aa^2b+Ba^3)}{2}}{d^{\frac{3}{2}}}}{e^6x + d^6e^5}$$

input `integrate((b*x+a)**3*(B*x+A)/(e*x+d)**(3/2),x)`

output `Piecewise((2*(B*b**3*(d + e*x)**(7/2)/(7*e**4) + (d + e*x)**(5/2)*(A*b**3*e + 3*B*a*b**2*e - 4*B*b**3*d)/(5*e**4) + (d + e*x)**(3/2)*(3*A*a*b**2*e**2 - 3*A*b**3*d*e + 3*B*a**2*b*e**2 - 9*B*a*b**2*d*e + 6*B*b**3*d**2)/(3*e**4) + sqrt(d + e*x)*(3*A*a**2*b*e**3 - 6*A*a*b**2*d*e**2 + 3*A*b**3*d**2*e + B*a**3*e**3 - 6*B*a**2*b*d*e**2 + 9*B*a*b**2*d**2*e - 4*B*b**3*d**3)/e**4 + (-A*e + B*d)*(a*e - b*d)**3/(e**4*sqrt(d + e*x)))/e, Ne(e, 0)), ((A*a**3*x + B*b**3*x**5/5 + x**4*(A*b**3 + 3*B*a*b**2)/4 + x**3*(3*A*a*b**2 + 3*B*a**2*b)/3 + x**2*(3*A*a**2*b + B*a**3)/2)/d**(3/2), True))`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.63

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{3/2}} dx = \frac{2 \left(\frac{5}{2} (ex+d)^{7/2} Bb^3 - 7(4Bb^3d - (3Bab^2 + Ab^3)e)(ex+d)^{5/2} + 35(2Bb^3d^2 - (3Bab^2 + Ab^3)de + (Ba^2b + Aab^2)e^4 \right)}{e^4}$$

input `integrate((b*x+a)^3*(B*x+A)/(e*x+d)^(3/2),x, algorithm="maxima")`

output `2/35*((5*(e*x + d)^(7/2)*B*b^3 - 7*(4*B*b^3*d - (3*B*a*b^2 + A*b^3)*e)*(e*x + d)^(5/2) + 35*(2*B*b^3*d^2 - (3*B*a*b^2 + A*b^3)*d*e + (B*a^2*b + A*a*b^2)*e^2)*(e*x + d)^(3/2) - 35*(4*B*b^3*d^3 - 3*(3*B*a*b^2 + A*b^3)*d^2*e + 6*(B*a^2*b + A*a*b^2)*d*e^2 - (B*a^3 + 3*A*a^2*b)*e^3)*sqrt(e*x + d))/e^4 - 35*(B*b^3*d^4 + A*a^3*e^4 - (3*B*a*b^2 + A*b^3)*d^3*e + 3*(B*a^2*b + A*a*b^2)*d^2*e^2 - (B*a^3 + 3*A*a^2*b)*d*e^3)/(sqrt(e*x + d)*e^4)/e`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 385 vs. $2(153) = 306$.

Time = 0.13 (sec) , antiderivative size = 385, normalized size of antiderivative = 2.31

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{3/2}} dx = \frac{2(Bb^3d^4 - 3Bab^2d^3e - Ab^3d^3e + 3Ba^2bd^2e^2 + 3Aab^2d^2e^2 - Ba^3de^3 - 3Aa^2bde^3 + Aa^3e^4)}{\sqrt{ex+d}e^5} + \frac{2\left(5(ex+d)^{7/2}Bb^3e^{30} - 28(ex+d)^{5/2}Bb^3de^{30} + 70(ex+d)^{3/2}Bb^3d^2e^{30} - 140\sqrt{ex+d}Bb^3d^3e^{30} + 21(ex+d)^{1/2}Bb^3d^4e^{30} - 28(ex+d)^{5/2}Aa^2bde^{30} + 70(ex+d)^{3/2}Aa^2b^2de^{30} - 140\sqrt{ex+d}Aa^2b^2d^2e^{30} + 21(ex+d)^{1/2}Aa^2b^2d^3e^{30} - 28(ex+d)^{5/2}Aa^3de^{30} + 70(ex+d)^{3/2}Aa^3e^{30} - 140\sqrt{ex+d}Aa^3e^{30} + 21(ex+d)^{1/2}Aa^3e^{30}\right)}{e^{35}}$$

input `integrate((b*x+a)^3*(B*x+A)/(e*x+d)^(3/2),x, algorithm="giac")`

output

$$\begin{aligned} & -2*(B*b^3*d^4 - 3*B*a*b^2*d^3*e - A*b^3*d^3*e + 3*B*a^2*b*d^2*e^2 + 3*A*a*b^2*d^2*e^2 - B*a^3*d*e^3 - 3*A*a^2*b*d*e^3 + A*a^3*e^4)/(sqrt(e*x + d)*e^5) \\ & + 2/35*(5*(e*x + d)^(7/2)*B*b^3*e^30 - 28*(e*x + d)^(5/2)*B*b^3*d*e^30 + 70*(e*x + d)^(3/2)*B*b^3*d^2*e^30 - 140*sqrt(e*x + d)*B*b^3*d^3*e^30 + 21*(e*x + d)^(1/2)*B*b^3*d^4*e^30 \\ & - 28*(e*x + d)^(5/2)*A*a^2*b*d*e^30 + 70*(e*x + d)^(3/2)*A*a^2*b^2*d*e^30 - 140*sqrt(e*x + d)*A*a^2*b^2*d^2*e^30 + 21*(e*x + d)^(1/2)*A*a^2*b^2*d^3*e^30 \\ & - 28*(e*x + d)^(5/2)*A*a^3*d*e^30 + 70*(e*x + d)^(3/2)*A*a^3*e^30 - 140*sqrt(e*x + d)*A*a^3*e^30 + 21*(e*x + d)^(1/2)*A*a^3*e^30)/e^35 \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.34

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{3/2}} dx = \frac{(d+ex)^{5/2}(2Ab^3e - 8Bb^3d + 6Bab^2e)}{5e^5} - \frac{-2Ba^3de^3 + 2Aa^3e^4 + 6Ba^2bd^2e^2 - 6Aa^2bde^3 - 6Bab^2d^3e + 6Aab^2d^2e^2 + 2Bb^3d^4 - 2Ab^3d^3e}{e^5\sqrt{d+ex}} + \frac{2(ae-bd)^2\sqrt{d+ex}(3Abe+Bae-4Bbd)}{e^5} + \frac{2Bb^3(d+ex)^{7/2}}{7e^5} + \frac{2b(ae-bd)(d+ex)^{3/2}(Abe+Bae-2Bbd)}{e^5}$$

input `int(((A + B*x)*(a + b*x)^3)/(d + e*x)^(3/2), x)`

output
$$\begin{aligned} & ((d + e*x)^{(5/2)} * (2*A*b^3*e - 8*B*b^3*d + 6*B*a*b^2*e)) / (5*e^5) - (2*A*a^3 * e^4 + 2*B*b^3*d^4 - 2*A*b^3*d^3*e - 2*B*a^3*d*e^3 + 6*A*a*b^2*d^2*e^2 + 6 * B*a^2*b*d^2*e^2 - 6*A*a^2*b*d*e^3 - 6*B*a*b^2*d^3*e) / (e^5 * (d + e*x)^{(1/2)}) \\ & + (2*(a*e - b*d)^2 * (d + e*x)^{(1/2)} * (3*A*b*e + B*a*e - 4*B*b*d)) / e^5 + (2 * B*b^3 * (d + e*x)^{(7/2)}) / (7*e^5) + (2*b*(a*e - b*d) * (d + e*x)^{(3/2)} * (A*b*e + B*a*e - 2*B*b*d)) / e^5 \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.11

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{3/2}} dx = \frac{2}{7}b^4e^4x^4 + \frac{8}{5}ab^3e^4x^3 - \frac{16}{35}b^4de^3x^3 + 4a^2b^2e^4x^2 - \frac{16}{5}ab^3de^3x^2 + \frac{32}{35}b^4d^2e^2x^2 + 8$$

input `int((b*x+a)^3*(B*x+A)/(e*x+d)^(3/2), x)`

output
$$\begin{aligned} & (2 * (- 35*a**4*e**4 + 280*a**3*b*d*e**3 + 140*a**3*b*e**4*x - 560*a**2*b**2 * d**2*e**2 - 280*a**2*b**2*d*e**3*x + 70*a**2*b**2*e**4*x**2 + 448*a*b**3 * d**3*e + 224*a*b**3*d**2*e**2*x - 56*a*b**3*d*e**3*x**2 + 28*a*b**3*e**4 * x**3 - 128*b**4*d**4 - 64*b**4*d**3*e*x + 16*b**4*d**2*e**2*x**2 - 8*b**4 * d*e**3*x**3 + 5*b**4*e**4*x**4)) / (35*sqrt(d + e*x)*e**5) \end{aligned}$$

3.145 $\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{5/2}} dx$

Optimal result	1351
Mathematica [A] (verified)	1352
Rubi [A] (verified)	1352
Maple [A] (verified)	1353
Fricas [A] (verification not implemented)	1354
Sympy [A] (verification not implemented)	1355
Maxima [A] (verification not implemented)	1355
Giac [B] (verification not implemented)	1356
Mupad [B] (verification not implemented)	1357
Reduce [B] (verification not implemented)	1357

Optimal result

Integrand size = 22, antiderivative size = 169

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{5/2}} dx = -\frac{2(bd-ae)^3(Bd-Ae)}{3e^5(d+ex)^{3/2}} + \frac{2(bd-ae)^2(4bBd-3Abe-aBe)}{e^5\sqrt{d+ex}} + \frac{6b(bd-ae)(2bBd-Abe-aBe)\sqrt{d+ex}}{e^5} - \frac{2b^2(4bBd-Abe-3aBe)(d+ex)^{3/2}}{3e^5} + \frac{2b^3B(d+ex)^{5/2}}{5e^5}$$

output

```
-2/3*(-a*e+b*d)^3*(-A*e+B*d)/e^5/(e*x+d)^(3/2)+2*(-a*e+b*d)^2*(-3*A*b*e-B*
a*e+4*B*b*d)/e^5/(e*x+d)^(1/2)+6*b*(-a*e+b*d)*(-A*b*e-B*a*e+2*B*b*d)*(e*x+
d)^(1/2)/e^5-2/3*b^2*(-A*b*e-3*B*a*e+4*B*b*d)*(e*x+d)^(3/2)/e^5+2/5*b^3*B*
(e*x+d)^(5/2)/e^5
```

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.32

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{5/2}} dx = \frac{2(-5a^3e^3(2Bd+ Ae+ 3Bex) + 15a^2be^2(-Ae(2d+ 3ex) + B(8d^2+ 12dex +$$

input

```
Integrate[((a + b*x)^3*(A + B*x))/(d + e*x)^(5/2),x]
```

output

```
(2*(-5*a^3*e^3*(2*B*d + A*e + 3*B*e*x) + 15*a^2*b*e^2*(-(A*e*(2*d + 3*e*x)
) + B*(8*d^2 + 12*d*e*x + 3*e^2*x^2)) + 15*a*b^2*e*(A*e*(8*d^2 + 12*d*e*x
+ 3*e^2*x^2) + B*(-16*d^3 - 24*d^2*e*x - 6*d*e^2*x^2 + e^3*x^3)) + b^3*(5*
A*e*(-16*d^3 - 24*d^2*e*x - 6*d*e^2*x^2 + e^3*x^3) + B*(128*d^4 + 192*d^3*
e*x + 48*d^2*e^2*x^2 - 8*d*e^3*x^3 + 3*e^4*x^4))))/(15*e^5*(d + e*x)^(3/2)
)
```

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{5/2}} dx$$

↓ 86

$$\int \left(\frac{b^2\sqrt{d+ex}(3aBe + Abe - 4bBd)}{e^4} - \frac{3b(bd - ae)(aBe + Abe - 2bBd)}{e^4\sqrt{d+ex}} + \frac{(ae - bd)^2(aBe + 3Abe - 4bBd)}{e^4(d+ex)^{3/2}} + \right.$$

↓ 2009

$$\left. - \frac{2b^2(d+ex)^{3/2}(-3aBe - Abe + 4bBd)}{3e^5} + \frac{6b\sqrt{d+ex}(bd - ae)(-aBe - Abe + 2bBd)}{e^5} + \frac{2(bd - ae)^2(-aBe - 3Abe + 4bBd)}{e^5\sqrt{d+ex}} - \frac{2(bd - ae)^3(Bd - Ae)}{3e^5(d+ex)^{3/2}} + \frac{2b^3B(d+ex)^{5/2}}{5e^5} \right)$$

input `Int[((a + b*x)^3*(A + B*x))/(d + e*x)^(5/2),x]`

output `(-2*(b*d - a*e)^3*(B*d - A*e))/(3*e^5*(d + e*x)^(3/2)) + (2*(b*d - a*e)^2*(4*b*B*d - 3*A*b*e - a*B*e))/(e^5*Sqrt[d + e*x]) + (6*b*(b*d - a*e)*(2*b*B*d - A*b*e - a*B*e)*Sqrt[d + e*x])/e^5 - (2*b^2*(4*b*B*d - A*b*e - 3*a*B*e)*(d + e*x)^(3/2))/(3*e^5) + (2*b^3*B*(d + e*x)^(5/2))/(5*e^5)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.06

method	result
risch	$\frac{2b(3e^2b^2Bx^2+5Ab^2e^2x+15Babe^2x-14b^2Bdex+45Aabe^2-40Ab^2de+45Ba^2e^2-120Babde+73b^2Bd^2)\sqrt{ex+d}}{15e^5}$
pseudoelliptic	$- \frac{2\left(\left(-\frac{3}{5}Bx^4-Ax^3\right)b^3-9a\left(\frac{Bx}{3}+A\right)x^2b^2+9a^2x(-Bx+A)b+a^3(3Bx+A)\right)e^4+6\left(x^2\left(\frac{4Bx}{15}+A\right)b^3-6a\left(-\frac{Bx}{2}+A\right)\right)e^3}{15e^5}$
gospers	$- \frac{2(-3Bx^4b^3e^4-5Ax^3b^3e^4-15Bx^3ab^2e^4+8Bx^3b^3de^3-45Ax^2ab^2e^4+30Ax^2b^3de^3-45Bx^2a^2be^4+90Bx^2ab^2de^3-11Bx^2b^3d^2e^2+6Aa^2b^2e^2+6Bab^2de^2-6Aa^2bde^2+6Ba^2be^2\sqrt{ex+d})}{15e^5}$
trager	$- \frac{2(-3Bx^4b^3e^4-5Ax^3b^3e^4-15Bx^3ab^2e^4+8Bx^3b^3de^3-45Ax^2ab^2e^4+30Ax^2b^3de^3-45Bx^2a^2be^4+90Bx^2ab^2de^3-11Bx^2b^3d^2e^2+6Aa^2b^2e^2+6Bab^2de^2-6Aa^2bde^2+6Ba^2be^2\sqrt{ex+d})}{15e^5}$
orering	$- \frac{2(-3Bx^4b^3e^4-5Ax^3b^3e^4-15Bx^3ab^2e^4+8Bx^3b^3de^3-45Ax^2ab^2e^4+30Ax^2b^3de^3-45Bx^2a^2be^4+90Bx^2ab^2de^3-11Bx^2b^3d^2e^2+6Aa^2b^2e^2+6Bab^2de^2-6Aa^2bde^2+6Ba^2be^2\sqrt{ex+d})}{15e^5}$
derivativedivides	$\frac{2b^3B(e^2x+d)^{\frac{5}{2}}}{5} + \frac{2Ab^3e(e^2x+d)^{\frac{3}{2}}}{3} + 2Bab^2e(e^2x+d)^{\frac{3}{2}} - \frac{8Bb^3d(e^2x+d)^{\frac{3}{2}}}{3} + 6Aa^2b^2e^2\sqrt{ex+d} - 6Ab^3de\sqrt{ex+d} + 6Ba^2be^2\sqrt{ex+d}$
default	$\frac{2b^3B(e^2x+d)^{\frac{5}{2}}}{5} + \frac{2Ab^3e(e^2x+d)^{\frac{3}{2}}}{3} + 2Bab^2e(e^2x+d)^{\frac{3}{2}} - \frac{8Bb^3d(e^2x+d)^{\frac{3}{2}}}{3} + 6Aa^2b^2e^2\sqrt{ex+d} - 6Ab^3de\sqrt{ex+d} + 6Ba^2be^2\sqrt{ex+d}$

```
input int((b*x+a)^3*(B*x+A)/(e*x+d)^(5/2),x,method=_RETURNVERBOSE)
```

```
output 2/15*b*(3*B*b^2*e^2*x^2+5*A*b^2*e^2*x+15*B*a*b*e^2*x-14*B*b^2*d*e*x+45*A*a*b*e^2-40*A*b^2*d*e+45*B*a^2*e^2-120*B*a*b*d*e+73*B*b^2*d^2)*(e*x+d)^(1/2)/e^5-2/3*(9*A*b*e^2*x+3*B*a*e^2*x-12*B*b*d*e*x+A*a*e^2+8*A*b*d*e+2*B*a*d*e-11*B*b*d^2)*(a^2*e^2-2*a*b*d*e+b^2*d^2)/e^5/(e*x+d)^(3/2)
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 284, normalized size of antiderivative = 1.68

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{5/2}} dx = \frac{2(3Bb^3e^4x^4+128Bb^3d^4-5Aa^3e^4-80(3Bab^2+Ab^3)d^3e+120(Ba^2b+Ab^2)d^2e^2-11Bb^3d^2e^2+6Aa^2b^2e^2+6Bab^2de^2-6Aa^2bde^2+6Ba^2be^2\sqrt{ex+d})}{15e^5}$$

```
input integrate((b*x+a)^3*(B*x+A)/(e*x+d)^(5/2),x,algorithm="fricas")
```

output

```
2/15*(3*B*b^3*e^4*x^4 + 128*B*b^3*d^4 - 5*A*a^3*e^4 - 80*(3*B*a*b^2 + A*b^3)*d^3*e + 120*(B*a^2*b + A*a*b^2)*d^2*e^2 - 10*(B*a^3 + 3*A*a^2*b)*d*e^3 - (8*B*b^3*d*e^3 - 5*(3*B*a*b^2 + A*b^3)*e^4)*x^3 + 3*(16*B*b^3*d^2*e^2 - 10*(3*B*a*b^2 + A*b^3)*d*e^3 + 15*(B*a^2*b + A*a*b^2)*e^4)*x^2 + 3*(64*B*b^3*d^3*e - 40*(3*B*a*b^2 + A*b^3)*d^2*e^2 + 60*(B*a^2*b + A*a*b^2)*d*e^3 - 5*(B*a^3 + 3*A*a^2*b)*e^4)*x)*sqrt(e*x + d)/(e^7*x^2 + 2*d*e^6*x + d^2*e^5)
```

Sympy [A] (verification not implemented)

Time = 9.61 (sec) , antiderivative size = 284, normalized size of antiderivative = 1.68

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{5/2}} dx = \frac{2 \left(\frac{Bb^3(d+ex)^{\frac{5}{2}}}{5e^4} + \frac{(d+ex)^{\frac{3}{2}}(Ab^3e+3Bab^2e-4Bb^3d)}{3e^4} + \frac{\sqrt{d+ex}(3Aab^2e^2-3Ab^3de+3Ba^2be^2-9Bab^2de+6Bb^3d^2)}{e^4} \right)}{\frac{Aa^3x + \frac{Bb^3x^5}{5} + \frac{x^4(Ab^3+3Bab^2)}{4} + \frac{x^3(3Aab^2+3Ba^2b)}{3} + \frac{x^2(3Aa^2b+Ba^3)}{2}}{d^{\frac{5}{2}}}}$$

input

```
integrate((b*x+a)**3*(B*x+A)/(e*x+d)**(5/2), x)
```

output

```
Piecewise(((2*(B*b**3*(d + e*x)**(5/2))/(5*e**4) + (d + e*x)**(3/2)*(A*b**3*e + 3*B*a*b**2*e - 4*B*b**3*d)/(3*e**4) + sqrt(d + e*x)*(3*A*a*b**2*e**2 - 3*A*b**3*d*e + 3*B*a**2*b*e**2 - 9*B*a*b**2*d*e + 6*B*b**3*d**2)/e**4 - (a*e - b*d)**2*(3*A*b*e + B*a*e - 4*B*b*d)/(e**4*sqrt(d + e*x)) + (-A*e + B*d)*(a*e - b*d)**3/(3*e**4*(d + e*x)**(3/2)))/e, Ne(e, 0)), ((A*a**3*x + B*b**3*x**5/5 + x**4*(A*b**3 + 3*B*a*b**2)/4 + x**3*(3*A*a*b**2 + 3*B*a**2*b)/3 + x**2*(3*A*a**2*b + B*a**3)/2)/d**(5/2), True))
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 271, normalized size of antiderivative = 1.60

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{5/2}} dx = \frac{2 \left(\frac{3(ex+d)^{\frac{5}{2}}Bb^3-5(4Bb^3d-(3Bab^2+Ab^3)e)(ex+d)^{\frac{3}{2}}+45(2Bb^3d^2-(3Bab^2+Ab^3)de+(Ba^2b+Aab^2))}{e^4} \right)}{d^{\frac{5}{2}}}$$

input

```
integrate((b*x+a)^3*(B*x+A)/(e*x+d)^(5/2), x, algorithm="maxima")
```


Mupad [B] (verification not implemented)

Time = 0.97 (sec) , antiderivative size = 264, normalized size of antiderivative = 1.56

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{5/2}} dx = \frac{(d+ex)^{3/2}(2Ab^3e - 8Bb^3d + 6Bab^2e)}{3e^5} + \frac{(d+ex)(2Ba^3e^3 - 12Ba^2bde^2 + 6Aa^2be^3 + 18Bab^2d^2e - 12Aab^2de^2 - 8Bb^3d^3 + 6Ab^3d^2e)}{e^5(d+ex)} + \frac{2Bb^3(d+ex)^{5/2}}{5e^5} + \frac{6b(ae-bd)\sqrt{d+ex}(Abe+BAe-2Bbd)}{e^5}$$

input `int(((A + B*x)*(a + b*x)^3)/(d + e*x)^(5/2), x)`output `((d + e*x)^(3/2)*(2*A*b^3*e - 8*B*b^3*d + 6*B*a*b^2*e))/(3*e^5) - ((d + e*x)*(2*B*a^3*e^3 - 8*B*b^3*d^3 + 6*A*a^2*b*e^3 + 6*A*b^3*d^2*e - 12*A*a*b^2*d*e^2 + 18*B*a*b^2*d^2*e - 12*B*a^2*b*d*e^2) + (2*A*a^3*e^4)/3 + (2*B*b^3*d^4)/3 - (2*A*b^3*d^3*e)/3 - (2*B*a^3*d*e^3)/3 + 2*A*a*b^2*d^2*e^2 + 2*B*a^2*b*d^2*e^2 - 2*A*a^2*b*d*e^3 - 2*B*a*b^2*d^3*e)/(e^5*(d + e*x)^(3/2)) + (2*B*b^3*(d + e*x)^(5/2))/(5*e^5) + (6*b*(a*e - b*d)*(d + e*x)^(1/2)*(A*b*e + B*a*e - 2*B*b*d))/e^5`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.14

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{5/2}} dx = \frac{\frac{2}{5}b^4e^4x^4 + \frac{8}{3}ab^3e^4x^3 - \frac{16}{15}b^4de^3x^3 + 12a^2b^2e^4x^2 - 16ab^3de^3x^2 + \frac{32}{5}b^4d^2e^2x^2 - \dots}{(d+ex)^{5/2}}$$

input `int((b*x+a)^3*(B*x+A)/(e*x+d)^(5/2), x)`output `(2*(- 5*a**4*e**4 - 40*a**3*b*d*e**3 - 60*a**3*b*e**4*x + 240*a**2*b**2*d**2*e**2 + 360*a**2*b**2*d*e**3*x + 90*a**2*b**2*e**4*x**2 - 320*a*b**3*d**3*e - 480*a*b**3*d**2*e**2*x - 120*a*b**3*d*e**3*x**2 + 20*a*b**3*e**4*x**3 + 128*b**4*d**4 + 192*b**4*d**3*e*x + 48*b**4*d**2*e**2*x**2 - 8*b**4*d*e**3*x**3 + 3*b**4*e**4*x**4))/(15*sqrt(d + e*x)*e**5*(d + e*x))`

3.146 $\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{7/2}} dx$

Optimal result	1358
Mathematica [A] (verified)	1359
Rubi [A] (verified)	1359
Maple [A] (verified)	1361
Fricas [A] (verification not implemented)	1361
Sympy [B] (verification not implemented)	1362
Maxima [A] (verification not implemented)	1363
Giac [B] (verification not implemented)	1364
Mupad [B] (verification not implemented)	1364
Reduce [B] (verification not implemented)	1365

Optimal result

Integrand size = 22, antiderivative size = 169

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{7/2}} dx = -\frac{2(bd-ae)^3(Bd-Ae)}{5e^5(d+ex)^{5/2}} + \frac{2(bd-ae)^2(4bBd-3Abe-aBe)}{3e^5(d+ex)^{3/2}} - \frac{6b(bd-ae)(2bBd-Abe-aBe)}{e^5\sqrt{d+ex}} - \frac{2b^2(4bBd-Abe-3aBe)\sqrt{d+ex}}{e^5} + \frac{2b^3B(d+ex)^{3/2}}{3e^5}$$

output

```
-2/5*(-a*e+b*d)^3*(-A*e+B*d)/e^5/(e*x+d)^(5/2)+2/3*(-a*e+b*d)^2*(-3*A*b*e-
B*a*e+4*B*b*d)/e^5/(e*x+d)^(3/2)-6*b*(-a*e+b*d)*(-A*b*e-B*a*e+2*B*b*d)/e^5
/(e*x+d)^(1/2)-2*b^2*(-A*b*e-3*B*a*e+4*B*b*d)*(e*x+d)^(1/2)/e^5+2/3*b^3*B*
(e*x+d)^(3/2)/e^5
```

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.34

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{7/2}} dx =$$

$$2(a^3e^3(2Bd + 3Ae + 5Bex) + 3a^2be^2(Ae(2d + 5ex) + B(8d^2 + 20dex + 15e^2x^2)) - 3ab^2e(-Ae(8d^2 + 20dex + 15e^2x^2) + B(16d^3 + 40d^2ex + 30de^2x^2 + 5e^3x^3)) + b^3(-3Ae(16d^3 + 40d^2ex + 30de^2x^2 + 5e^3x^3) + B(128d^4 + 320d^3ex + 240d^2e^2x^2 + 40de^3x^3 - 5e^4x^4)))/(15e^5(d + ex)^{5/2})$$

input

```
Integrate[((a + b*x)^3*(A + B*x))/(d + e*x)^(7/2),x]
```

output

```
(-2*(a^3*e^3*(2*B*d + 3*A*e + 5*B*e*x) + 3*a^2*b*e^2*(A*e*(2*d + 5*e*x) + B*(8*d^2 + 20*d*e*x + 15*e^2*x^2)) - 3*a*b^2*e*(-(A*e*(8*d^2 + 20*d*e*x + 15*e^2*x^2)) + 3*B*(16*d^3 + 40*d^2*e*x + 30*d*e^2*x^2 + 5*e^3*x^3)) + b^3*(-3*A*e*(16*d^3 + 40*d^2*e*x + 30*d*e^2*x^2 + 5*e^3*x^3) + B*(128*d^4 + 320*d^3*e*x + 240*d^2*e^2*x^2 + 40*d*e^3*x^3 - 5*e^4*x^4)))/(15*e^5*(d + e*x)^(5/2))
```

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{7/2}} dx$$

↓ 86

$$\int \left(\frac{b^2(3aBe + Abe - 4bBd)}{e^4\sqrt{d + ex}} - \frac{3b(bd - ae)(aBe + Abe - 2bBd)}{e^4(d + ex)^{3/2}} + \frac{(ae - bd)^2(aBe + 3Abe - 4bBd)}{e^4(d + ex)^{5/2}} + \frac{(ae - bd)(aBe + 3Abe - 4bBd)}{e^4(d + ex)^{7/2}} \right) dx$$

↓ 2009

$$\begin{aligned} & -\frac{2b^2\sqrt{d+ex}(-3aBe - Abe + 4bBd)}{e^5} - \frac{6b(bd - ae)(-aBe - Abe + 2bBd)}{e^5\sqrt{d+ex}} + \\ & \frac{2(bd - ae)^2(-aBe - 3Abe + 4bBd)}{3e^5(d+ex)^{3/2}} - \frac{2(bd - ae)^3(Bd - Ae)}{5e^5(d+ex)^{5/2}} + \frac{2b^3B(d+ex)^{3/2}}{3e^5} \end{aligned}$$

input `Int[((a + b*x)^3*(A + B*x))/(d + e*x)^(7/2), x]`

output `(-2*(b*d - a*e)^3*(B*d - A*e))/(5*e^5*(d + e*x)^(5/2)) + (2*(b*d - a*e)^2*(4*b*B*d - 3*A*b*e - a*B*e))/(3*e^5*(d + e*x)^(3/2)) - (6*b*(b*d - a*e)*(2*b*B*d - A*b*e - a*B*e))/(e^5*sqrt[d + e*x]) - (2*b^2*(4*b*B*d - A*b*e - 3*a*B*e)*sqrt[d + e*x])/e^5 + (2*b^3*B*(d + e*x)^(3/2))/(3*e^5)`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;`
`FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.14

method	result
pseudoelliptic	$\left((10Bx^4 + 30Ax^3)b^3 - 90a^2x^2(-Bx + A)b^2 - 30a^2x(3Bx + A)b - 6a^3\left(\frac{5Bx}{3} + A\right) \right) e^4 - 12\left(-15\left(-\frac{4Bx}{9} + A\right)x^2b^3 + 10a\left(-\right.\right.$
risch	$\frac{2b^2(ebBx + 3Abe + 9Bae - 11Bbd)\sqrt{ex+d}}{3e^5} - \frac{2(45Ax^2b^2e^3 + 45Bx^2abe^3 - 90Bx^2b^2de^2 + 15Axab^2e^3 + 75Ax^2b^2de^2 + 5A^2b^2e^3)}{3e^5}$
derivativedivides	$\frac{2b^3B\left(\frac{ex+d}{3}\right)^{\frac{3}{2}}}{3} + 2Ab^3e\sqrt{ex+d} + 6Ba^2b^2e\sqrt{ex+d} - 8Bb^3d\sqrt{ex+d} - \frac{6b(Aab^2e^2 - Ab^2de + Ba^2e^2 - 3Babde + 2b^2Bd^2)}{\sqrt{ex+d}} - \frac{2(3Aa^2b^2e^3 + 3A^2b^2e^3)}{3e^5}$
default	$\frac{2b^3B\left(\frac{ex+d}{3}\right)^{\frac{3}{2}}}{3} + 2Ab^3e\sqrt{ex+d} + 6Ba^2b^2e\sqrt{ex+d} - 8Bb^3d\sqrt{ex+d} - \frac{6b(Aab^2e^2 - Ab^2de + Ba^2e^2 - 3Babde + 2b^2Bd^2)}{\sqrt{ex+d}} - \frac{2(3Aa^2b^2e^3 + 3A^2b^2e^3)}{3e^5}$
gospers	$-\frac{2(-5Bx^4b^3e^4 - 15Ax^3b^3e^4 - 45Bx^3ab^2e^4 + 40Bx^3b^3de^3 + 45Ax^2ab^2e^4 - 90Ax^2b^3de^3 + 45Bx^2a^2be^4 - 270Bx^2a^2b^2e^4)}{3e^5}$
trager	$-\frac{2(-5Bx^4b^3e^4 - 15Ax^3b^3e^4 - 45Bx^3ab^2e^4 + 40Bx^3b^3de^3 + 45Ax^2ab^2e^4 - 90Ax^2b^3de^3 + 45Bx^2a^2be^4 - 270Bx^2a^2b^2e^4)}{3e^5}$
orering	$-\frac{2(-5Bx^4b^3e^4 - 15Ax^3b^3e^4 - 45Bx^3ab^2e^4 + 40Bx^3b^3de^3 + 45Ax^2ab^2e^4 - 90Ax^2b^3de^3 + 45Bx^2a^2be^4 - 270Bx^2a^2b^2e^4)}{3e^5}$

```
input int((b*x+a)^3*(B*x+A)/(e*x+d)^(7/2), x, method=_RETURNVERBOSE)
```

```
output 1/15*(((10*B*x^4+30*A*x^3)*b^3-90*a*x^2*(-B*x+A)*b^2-30*a^2*x*(3*B*x+A)*b-6*a^3*(5/3*B*x+A))*e^4-12*(-15*(-4/9*B*x+A)*x^2*b^3+10*a*(-9/2*B*x+A)*x*b^2+a^2*(10*B*x+A)*b+1/3*a^3*B)*d*e^3-48*(-5*x*(-2*B*x+A)*b^2+a*(-15*B*x+A)*b+a^2*B)*b*d^2*e^2+96*b^2*((-20/3*B*x+A)*b+3*B*a)*d^3*e-256*b^3*B*d^4)/(e*x+d)^(5/2)/e^5
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 294, normalized size of antiderivative = 1.74

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{7/2}} dx = \frac{2(5Bb^3e^4x^4 - 128Bb^3d^4 - 3Aa^3e^4 + 48(3Bab^2 + Ab^3)d^3e - 24(Ba^2b + Aa^2b^2)d^2e^2 - 24A^2b^2d^2e^2 - 24A^2b^2d^2e^2 - 24A^2b^2d^2e^2)}{(d + ex)^{7/2}}$$

```
input integrate((b*x+a)^3*(B*x+A)/(e*x+d)^(7/2), x, algorithm="fricas")
```

output

```
2/15*(5*B*b^3*e^4*x^4 - 128*B*b^3*d^4 - 3*A*a^3*e^4 + 48*(3*B*a*b^2 + A*b^3)*d^3*e - 24*(B*a^2*b + A*a*b^2)*d^2*e^2 - 2*(B*a^3 + 3*A*a^2*b)*d*e^3 - 5*(8*B*b^3*d*e^3 - 3*(3*B*a*b^2 + A*b^3)*e^4)*x^3 - 15*(16*B*b^3*d^2*e^2 - 6*(3*B*a*b^2 + A*b^3)*d*e^3 + 3*(B*a^2*b + A*a*b^2)*e^4)*x^2 - 5*(64*B*b^3*d^3*e - 24*(3*B*a*b^2 + A*b^3)*d^2*e^2 + 12*(B*a^2*b + A*a*b^2)*d*e^3 + (B*a^3 + 3*A*a^2*b)*e^4)*x)*sqrt(e*x + d)/(e^8*x^3 + 3*d*e^7*x^2 + 3*d^2*e^6*x + d^3*e^5)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1654 vs. $2(167) = 334$.

Time = 0.57 (sec) , antiderivative size = 1654, normalized size of antiderivative = 9.79

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{7/2}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)**3*(B*x+A)/(e*x+d)**(7/2),x)
```

output

```
Piecewise((-6*A*a**3*e**4/(15*d**2*e**5*sqrt(d + e*x) + 30*d*e**6*x*sqrt(d
+ e*x) + 15*e**7*x**2*sqrt(d + e*x)) - 12*A*a**2*b*d*e**3/(15*d**2*e**5*s
qrt(d + e*x) + 30*d*e**6*x*sqrt(d + e*x) + 15*e**7*x**2*sqrt(d + e*x)) - 3
0*A*a**2*b*e**4*x/(15*d**2*e**5*sqrt(d + e*x) + 30*d*e**6*x*sqrt(d + e*x)
+ 15*e**7*x**2*sqrt(d + e*x)) - 48*A*a*b**2*d**2*e**2/(15*d**2*e**5*sqrt(d
+ e*x) + 30*d*e**6*x*sqrt(d + e*x) + 15*e**7*x**2*sqrt(d + e*x)) - 120*A*
a*b**2*d*e**3*x/(15*d**2*e**5*sqrt(d + e*x) + 30*d*e**6*x*sqrt(d + e*x) +
15*e**7*x**2*sqrt(d + e*x)) - 90*A*a*b**2*e**4*x**2/(15*d**2*e**5*sqrt(d +
e*x) + 30*d*e**6*x*sqrt(d + e*x) + 15*e**7*x**2*sqrt(d + e*x)) + 96*A*b**
3*d**3*e/(15*d**2*e**5*sqrt(d + e*x) + 30*d*e**6*x*sqrt(d + e*x) + 15*e**7
*x**2*sqrt(d + e*x)) + 240*A*b**3*d**2*e**2*x/(15*d**2*e**5*sqrt(d + e*x)
+ 30*d*e**6*x*sqrt(d + e*x) + 15*e**7*x**2*sqrt(d + e*x)) + 180*A*b**3*d*e
**3*x**2/(15*d**2*e**5*sqrt(d + e*x) + 30*d*e**6*x*sqrt(d + e*x) + 15*e**7
*x**2*sqrt(d + e*x)) + 30*A*b**3*e**4*x**3/(15*d**2*e**5*sqrt(d + e*x) + 3
0*d*e**6*x*sqrt(d + e*x) + 15*e**7*x**2*sqrt(d + e*x)) - 4*B*a**3*d*e**3/(
15*d**2*e**5*sqrt(d + e*x) + 30*d*e**6*x*sqrt(d + e*x) + 15*e**7*x**2*sqrt
(d + e*x)) - 10*B*a**3*e**4*x/(15*d**2*e**5*sqrt(d + e*x) + 30*d*e**6*x*sq
rt(d + e*x) + 15*e**7*x**2*sqrt(d + e*x)) - 48*B*a**2*b*d**2*e**2/(15*d**2
*e**5*sqrt(d + e*x) + 30*d*e**6*x*sqrt(d + e*x) + 15*e**7*x**2*sqrt(d + e
x)) - 120*B*a**2*b*d*e**3*x/(15*d**2*e**5*sqrt(d + e*x) + 30*d*e**6*x*s...
```

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.62

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{7/2}} dx = \frac{2 \left(\frac{5 \left((ex+d)^{\frac{3}{2}} Bb^3 - 3(4Bb^3d - (3Bab^2 + Ab^3)e) \sqrt{ex+d} \right)}{e^4} - \frac{3Bb^3d^4 + 3Aa^3e^4 - 3(3Bab^2 + Ab^3)d^3e + 9A^2b^3d^2 + 9A^2b^2d^2e - 3(Ba^3 + 3Aa^2b)d^2e^3 + 45(2Bb^3d^2 - (3Bab^2 + Ab^3)d^2e + (Ba^2b + Aa^2b^2)e^2)(ex+d)^2 - 5(4Bb^3d^3 - 3(3Bab^2 + Ab^3)d^2e + 6(Ba^2b + Aa^2b^2)d^2e^2 - (Ba^3 + 3Aa^2b)e^3)(ex+d)}{(ex+d)^{5/2}e^4} \right)}{e^4}$$

input

```
integrate((b*x+a)^3*(B*x+A)/(e*x+d)^(7/2),x, algorithm="maxima")
```

output

```
2/15*(5*((e*x + d)^(3/2)*B*b^3 - 3*(4*B*b^3*d - (3*B*a*b^2 + A*b^3)*e)*sqrt
t(e*x + d))/e^4 - (3*B*b^3*d^4 + 3*A*a^3*e^4 - 3*(3*B*a*b^2 + A*b^3)*d^3*e
+ 9*(B*a^2*b + A*a*b^2)*d^2*e^2 - 3*(B*a^3 + 3*A*a^2*b)*d^2*e^3 + 45*(2*B*b
^3*d^2 - (3*B*a*b^2 + A*b^3)*d^2*e + (B*a^2*b + A*a^2*b^2)*e^2)*(e*x + d)^2 -
5*(4*B*b^3*d^3 - 3*(3*B*a*b^2 + A*b^3)*d^2*e + 6*(B*a^2*b + A*a^2*b^2)*d^2*e^2
- (B*a^3 + 3*A*a^2*b)*e^3)*(e*x + d))/((e*x + d)^(5/2)*e^4))/e
```


Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 358 vs. $2(153) = 306$.

Time = 0.13 (sec) , antiderivative size = 358, normalized size of antiderivative = 2.12

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{7/2}} dx =$$

$$\frac{2(90(ex+d)^2 Bb^3 d^2 - 20(ex+d)Bb^3 d^3 + 3Bb^3 d^4 - 135(ex+d)^2 Bab^2 de - 45(ex+d)^2 Ab^3 de + 45(e$$

$$+ \frac{2\left((ex+d)^{\frac{3}{2}} Bb^3 e^{10} - 12\sqrt{ex+d} Bb^3 de^{10} + 9\sqrt{ex+d} Bab^2 e^{11} + 3\sqrt{ex+d} Ab^3 e^{11}\right)}{3e^{15}}$$

input `integrate((b*x+a)^3*(B*x+A)/(e*x+d)^(7/2),x, algorithm="giac")`

output `-2/15*(90*(e*x + d)^2*B*b^3*d^2 - 20*(e*x + d)*B*b^3*d^3 + 3*B*b^3*d^4 - 135*(e*x + d)^2*B*a*b^2*d*e - 45*(e*x + d)^2*A*b^3*d*e + 45*(e*x + d)*B*a*b^2*d^2*e + 15*(e*x + d)*A*b^3*d^2*e - 9*B*a*b^2*d^3*e - 3*A*b^3*d^3*e + 45*(e*x + d)^2*B*a^2*b*e^2 + 45*(e*x + d)^2*A*a*b^2*e^2 - 30*(e*x + d)*B*a^2*b*d*e^2 - 30*(e*x + d)*A*a*b^2*d*e^2 + 9*B*a^2*b*d^2*e^2 + 9*A*a*b^2*d^2*e^2 + 5*(e*x + d)*B*a^3*e^3 + 15*(e*x + d)*A*a^2*b*e^3 - 3*B*a^3*d*e^3 - 9*A*a^2*b*d*e^3 + 3*A*a^3*e^4)/((e*x + d)^(5/2)*e^5) + 2/3*((e*x + d)^(3/2)*B*b^3*e^10 - 12*sqrt(e*x + d)*B*b^3*d*e^10 + 9*sqrt(e*x + d)*B*a*b^2*e^11 + 3*sqrt(e*x + d)*A*b^3*e^11)/e^15`

Mupad [B] (verification not implemented)

Time = 1.01 (sec) , antiderivative size = 300, normalized size of antiderivative = 1.78

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{7/2}} dx =$$

$$\frac{2(2Ba^3de^3 + 5Ba^3e^4x + 3Aa^3e^4 + 24Ba^2bd^2e^2 + 60Ba^2bde^3x + 6Aa^2bde^3 + 45Ba^2be^4x^2 +$$

input `int(((A + B*x)*(a + b*x)^3)/(d + e*x)^(7/2),x)`

output

```

-(2*(3*A*a^3*e^4 + 128*B*b^3*d^4 - 48*A*b^3*d^3*e + 2*B*a^3*d*e^3 + 5*B*a^
3*e^4*x - 15*A*b^3*e^4*x^3 - 5*B*b^3*e^4*x^4 + 320*B*b^3*d^3*e*x + 24*A*a*
b^2*d^2*e^2 + 24*B*a^2*b*d^2*e^2 + 45*A*a*b^2*e^4*x^2 + 45*B*a^2*b*e^4*x^2
- 45*B*a*b^2*e^4*x^3 - 120*A*b^3*d^2*e^2*x - 90*A*b^3*d*e^3*x^2 + 40*B*b^
3*d*e^3*x^3 + 240*B*b^3*d^2*e^2*x^2 + 6*A*a^2*b*d*e^3 - 144*B*a*b^2*d^3*e
+ 15*A*a^2*b*e^4*x + 60*A*a*b^2*d*e^3*x + 60*B*a^2*b*d*e^3*x - 360*B*a*b^2
*d^2*e^2*x - 270*B*a*b^2*d*e^3*x^2))/(15*e^5*(d + e*x)^(5/2))

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.21

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{7/2}} dx = \frac{2}{3}b^4e^4x^4 + 8ab^3e^4x^3 - \frac{16}{3}b^4de^3x^3 - 12a^2b^2e^4x^2 + 48ab^3de^3x^2 - 32b^4d^2e^2x^2 -$$

input

```
int((b*x+a)^3*(B*x+A)/(e*x+d)^(7/2),x)
```

output

```

(2*( - 3*a**4*e**4 - 8*a**3*b*d*e**3 - 20*a**3*b*e**4*x - 48*a**2*b**2*d**
2*e**2 - 120*a**2*b**2*d*e**3*x - 90*a**2*b**2*e**4*x**2 + 192*a*b**3*d**3
*e + 480*a*b**3*d**2*e**2*x + 360*a*b**3*d*e**3*x**2 + 60*a*b**3*e**4*x**3
- 128*b**4*d**4 - 320*b**4*d**3*e*x - 240*b**4*d**2*e**2*x**2 - 40*b**4*d
*e**3*x**3 + 5*b**4*e**4*x**4))/(15*sqrt(d + e*x)*e**5*(d**2 + 2*d*e*x + e
**2*x**2))

```

3.147 $\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{9/2}} dx$

Optimal result	1366
Mathematica [A] (verified)	1367
Rubi [A] (verified)	1367
Maple [A] (verified)	1369
Fricas [A] (verification not implemented)	1369
Sympy [B] (verification not implemented)	1370
Maxima [A] (verification not implemented)	1371
Giac [B] (verification not implemented)	1372
Mupad [B] (verification not implemented)	1372
Reduce [B] (verification not implemented)	1373

Optimal result

Integrand size = 22, antiderivative size = 167

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{9/2}} dx = -\frac{2(bd-ae)^3(Bd-Ae)}{7e^5(d+ex)^{7/2}} + \frac{2(bd-ae)^2(4bBd-3Abe-aBe)}{5e^5(d+ex)^{5/2}} - \frac{2b(bd-ae)(2bBd-Abe-aBe)}{e^5(d+ex)^{3/2}} + \frac{2b^2(4bBd-Abe-3aBe)}{e^5\sqrt{d+ex}} + \frac{2b^3B\sqrt{d+ex}}{e^5}$$

output

```
-2/7*(-a*e+b*d)^3*(-A*e+B*d)/e^5/(e*x+d)^(7/2)+2/5*(-a*e+b*d)^2*(-3*A*b*e-
B*a*e+4*B*b*d)/e^5/(e*x+d)^(5/2)-2*b*(-a*e+b*d)*(-A*b*e-B*a*e+2*B*b*d)/e^5
/(e*x+d)^(3/2)+2*b^2*(-A*b*e-3*B*a*e+4*B*b*d)/e^5/(e*x+d)^(1/2)+2*b^3*B*(e
*x+d)^(1/2)/e^5
```

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.34

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{9/2}} dx =$$

$$2(a^3e^3(2Bd + 5Ae + 7Bex) + a^2be^2(3Ae(2d + 7ex) + B(8d^2 + 28dex + 35e^2x^2)) + ab^2e(Ae(8d^2 + 28dex$$

input

```
Integrate[((a + b*x)^3*(A + B*x))/(d + e*x)^(9/2),x]
```

output

```
(-2*(a^3*e^3*(2*B*d + 5*A*e + 7*B*e*x) + a^2*b*e^2*(3*A*e*(2*d + 7*e*x) +
B*(8*d^2 + 28*d*e*x + 35*e^2*x^2)) + a*b^2*e*(A*e*(8*d^2 + 28*d*e*x + 35*e
^2*x^2) + 3*B*(16*d^3 + 56*d^2*e*x + 70*d*e^2*x^2 + 35*e^3*x^3)) + b^3*(A*
e*(16*d^3 + 56*d^2*e*x + 70*d*e^2*x^2 + 35*e^3*x^3) - B*(128*d^4 + 448*d^3
*e*x + 560*d^2*e^2*x^2 + 280*d*e^3*x^3 + 35*e^4*x^4)))/(35*e^5*(d + e*x)^(
7/2))
```

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{9/2}} dx$$

↓ 86

$$\int \left(\frac{b^2(3aBe + Abe - 4bBd)}{e^4(d + ex)^{3/2}} - \frac{3b(bd - ae)(aBe + Abe - 2bBd)}{e^4(d + ex)^{5/2}} + \frac{(ae - bd)^2(aBe + 3Abe - 4bBd)}{e^4(d + ex)^{7/2}} + \frac{(ae - bd)(aBe + 3Abe - 4bBd)}{e^4(d + ex)^{9/2}} \right) dx$$

↓ 2009

$$\frac{2b^2(-3aBe - Abe + 4bBd)}{e^5\sqrt{d+ex}} - \frac{2b(bd - ae)(-aBe - Abe + 2bBd)}{e^5(d+ex)^{3/2}} + \frac{2(bd - ae)^2(-aBe - 3Abe + 4bBd)}{5e^5(d+ex)^{5/2}} - \frac{2(bd - ae)^3(Bd - Ae)}{7e^5(d+ex)^{7/2}} + \frac{2b^3B\sqrt{d+ex}}{e^5}$$

input `Int[((a + b*x)^3*(A + B*x))/(d + e*x)^(9/2), x]`

output `(-2*(b*d - a*e)^3*(B*d - A*e))/(7*e^5*(d + e*x)^(7/2)) + (2*(b*d - a*e)^2*(4*b*B*d - 3*A*b*e - a*B*e))/(5*e^5*(d + e*x)^(5/2)) - (2*b*(b*d - a*e)*(2*b*B*d - A*b*e - a*B*e))/(e^5*(d + e*x)^(3/2)) + (2*b^2*(4*b*B*d - A*b*e - 3*a*B*e))/(e^5*sqrt[d + e*x]) + (2*b^3*B*sqrt[d + e*x])/e^5`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;`
`FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.16

method	result
pseudoelliptic	$\left(-70x^3(-Bx+A)b^3-70ax^2(3Bx+A)b^2-42a^2x\left(\frac{5Bx}{3}+A\right)b-10a^3\left(\frac{7Bx}{5}+A\right)\right)e^4-12\left(\frac{35x^2(-4Bx+A)b^3}{3}+\frac{14ax\left(\frac{15B}{2}\right)}{3}\right)$
derivativedivides	$\frac{2b^3B\sqrt{ex+d}-\frac{2b^2(Abe+3Bae-4Bbd)}{\sqrt{ex+d}}}{5(ex+d)^{\frac{5}{2}}}-\frac{2(3Aa^2be^3-6Aab^2de^2+3Ab^3d^2e+Ba^3e^3-6Ba^2bde^2+9Bab^2d^2e-4b^3Bd^3)}{5(ex+d)^{\frac{5}{2}}}$
default	$\frac{2b^3B\sqrt{ex+d}-\frac{2b^2(Abe+3Bae-4Bbd)}{\sqrt{ex+d}}}{5(ex+d)^{\frac{5}{2}}}-\frac{2(3Aa^2be^3-6Aab^2de^2+3Ab^3d^2e+Ba^3e^3-6Ba^2bde^2+9Bab^2d^2e-4b^3Bd^3)}{5(ex+d)^{\frac{5}{2}}}$
gospers	$\frac{2(-35Bx^4b^3e^4+35Ax^3b^3e^4+105Bx^3ab^2e^4-280Bx^3b^3de^3+35Ax^2ab^2e^4+70Ax^2b^3de^3+35Bx^2a^2be^4+210Bx^2a^2b^2e^4-210Bx^2a^2b^3e^4)}{5(ex+d)^{\frac{5}{2}}}$
trager	$\frac{2(-35Bx^4b^3e^4+35Ax^3b^3e^4+105Bx^3ab^2e^4-280Bx^3b^3de^3+35Ax^2ab^2e^4+70Ax^2b^3de^3+35Bx^2a^2be^4+210Bx^2a^2b^2e^4-210Bx^2a^2b^3e^4)}{5(ex+d)^{\frac{5}{2}}}$
orering	$\frac{2(-35Bx^4b^3e^4+35Ax^3b^3e^4+105Bx^3ab^2e^4-280Bx^3b^3de^3+35Ax^2ab^2e^4+70Ax^2b^3de^3+35Bx^2a^2be^4+210Bx^2a^2b^2e^4-210Bx^2a^2b^3e^4)}{5(ex+d)^{\frac{5}{2}}}$
risch	$\frac{2b^3B\sqrt{ex+d}}{e^5}-\frac{2(35Ax^3b^3e^4+105Bx^3ab^2e^4-140Bx^3b^3de^3+35Ax^2ab^2e^4+70Ax^2b^3de^3+35Bx^2a^2be^4+210Bx^2a^2b^2e^4-210Bx^2a^2b^3e^4)}{5(ex+d)^{\frac{5}{2}}}$

input `int((b*x+a)^3*(B*x+A)/(e*x+d)^(9/2),x,method=_RETURNVERBOSE)`

output `1/35*((-70*x^3*(-B*x+A)*b^3-70*a*x^2*(3*B*x+A)*b^2-42*a^2*x*(5/3*B*x+A)*b-10*a^3*(7/5*B*x+A))*e^4-12*(35/3*x^2*(-4*B*x+A)*b^3+14/3*a*x*(15/2*B*x+A)*b^2+a^2*(14/3*B*x+A)*b+1/3*a^3*B)*d*e^3-16*b*((-70*B*x^2+7*A*x)*b^2+a*(21*B*x+A)*b+a^2*B)*d^2*e^2-32*((-28*B*x+A)*b+3*B*a)*b^2*d^3*e+256*b^3*B*d^4)/(e*x+d)^(7/2)/e^5`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 306, normalized size of antiderivative = 1.83

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{9/2}} dx = \frac{2(35Bb^3e^4x^4+128Bb^3d^4-5Aa^3e^4-16(3Bab^2+Ab^3)d^3e-8(Ba^2b+Aa^2b^2)d^2e^2+21Bab^2d^2e-4b^3Bd^3)}{5(e^2x+d)^{7/2}}$$

input `integrate((b*x+a)^3*(B*x+A)/(e*x+d)^(9/2),x,algorithm="fricas")`

output

```
2/35*(35*B*b^3*e^4*x^4 + 128*B*b^3*d^4 - 5*A*a^3*e^4 - 16*(3*B*a*b^2 + A*b^3)*d^3*e - 8*(B*a^2*b + A*a*b^2)*d^2*e^2 - 2*(B*a^3 + 3*A*a^2*b)*d*e^3 + 35*(8*B*b^3*d*e^3 - (3*B*a*b^2 + A*b^3)*e^4)*x^3 + 35*(16*B*b^3*d^2*e^2 - 2*(3*B*a*b^2 + A*b^3)*d*e^3 - (B*a^2*b + A*a*b^2)*e^4)*x^2 + 7*(64*B*b^3*d^3*e - 8*(3*B*a*b^2 + A*b^3)*d^2*e^2 - 4*(B*a^2*b + A*a*b^2)*d*e^3 - (B*a^3 + 3*A*a^2*b)*e^4)*x)*sqrt(e*x + d)/(e^9*x^4 + 4*d*e^8*x^3 + 6*d^2*e^7*x^2 + 4*d^3*e^6*x + d^4*e^5)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2144 vs. $2(165) = 330$.

Time = 0.79 (sec) , antiderivative size = 2144, normalized size of antiderivative = 12.84

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{9/2}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)**3*(B*x+A)/(e*x+d)**(9/2),x)
```

output

```
Piecewise((-10*A*a**3*e**4/(35*d**3*e**5*sqrt(d + e*x) + 105*d**2*e**6*x*sqrt(d + e*x) + 105*d*e**7*x**2*sqrt(d + e*x) + 35*e**8*x**3*sqrt(d + e*x)) - 12*A*a**2*b*d*e**3/(35*d**3*e**5*sqrt(d + e*x) + 105*d**2*e**6*x*sqrt(d + e*x) + 105*d*e**7*x**2*sqrt(d + e*x) + 35*e**8*x**3*sqrt(d + e*x)) - 42*A*a**2*b*e**4*x/(35*d**3*e**5*sqrt(d + e*x) + 105*d**2*e**6*x*sqrt(d + e*x) + 105*d*e**7*x**2*sqrt(d + e*x) + 35*e**8*x**3*sqrt(d + e*x)) - 16*A*a*b**2*d**2*e**2/(35*d**3*e**5*sqrt(d + e*x) + 105*d**2*e**6*x*sqrt(d + e*x) + 105*d*e**7*x**2*sqrt(d + e*x) + 35*e**8*x**3*sqrt(d + e*x)) - 56*A*a*b**2*d*e**3*x/(35*d**3*e**5*sqrt(d + e*x) + 105*d**2*e**6*x*sqrt(d + e*x) + 105*d*e**7*x**2*sqrt(d + e*x) + 35*e**8*x**3*sqrt(d + e*x)) - 70*A*a*b**2*e**4*x**2/(35*d**3*e**5*sqrt(d + e*x) + 105*d**2*e**6*x*sqrt(d + e*x) + 105*d*e**7*x**2*sqrt(d + e*x) + 35*e**8*x**3*sqrt(d + e*x)) - 32*A*b**3*d**3*e/(35*d**3*e**5*sqrt(d + e*x) + 105*d**2*e**6*x*sqrt(d + e*x) + 105*d*e**7*x**2*sqrt(d + e*x) + 35*e**8*x**3*sqrt(d + e*x)) - 112*A*b**3*d**2*e**2*x/(35*d**3*e**5*sqrt(d + e*x) + 105*d**2*e**6*x*sqrt(d + e*x) + 105*d*e**7*x**2*sqrt(d + e*x) + 35*e**8*x**3*sqrt(d + e*x)) - 140*A*b**3*d*e**3*x**2/(35*d**3*e**5*sqrt(d + e*x) + 105*d**2*e**6*x*sqrt(d + e*x) + 105*d*e**7*x**2*sqrt(d + e*x) + 35*e**8*x**3*sqrt(d + e*x)) - 70*A*b**3*e**4*x**3/(35*d**3*e**5*sqrt(d + e*x) + 105*d**2*e**6*x*sqrt(d + e*x) + 105*d*e**7*x**2*sqrt(d + e*x) + 35*e**8*x**3*sqrt(d + e*x)) - 4*B*a**3*d*e**3/(35*d**3...
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 271, normalized size of antiderivative = 1.62

$$\int \frac{(a + bx)^3(A + Bx)}{(d + ex)^{9/2}} dx = \frac{2 \left(\frac{35\sqrt{ex+d}Bb^3}{e^4} - \frac{5Bb^3d^4 + 5Aa^3e^4 - 5(3Bab^2 + Ab^3)d^3e + 15(Ba^2b + Aab^2)d^2e^2 - 5(Ba^3 + 3Aa^2b)d}{e^4} \right)}{e^4}$$

input

```
integrate((b*x+a)^3*(B*x+A)/(e*x+d)^(9/2),x, algorithm="maxima")
```

output

```
2/35*(35*sqrt(e*x + d)*B*b^3/e^4 - (5*B*b^3*d^4 + 5*A*a^3*e^4 - 5*(3*B*a*b^2 + A*b^3)*d^3*e + 15*(B*a^2*b + A*a*b^2)*d^2*e^2 - 5*(B*a^3 + 3*A*a^2*b)*d*e^3 - 35*(4*B*b^3*d - (3*B*a*b^2 + A*b^3)*e)*(e*x + d)^3 + 35*(2*B*b^3*d^2 - (3*B*a*b^2 + A*b^3)*d*e + (B*a^2*b + A*a*b^2)*e^2)*(e*x + d)^2 - 7*(4*B*b^3*d^3 - 3*(3*B*a*b^2 + A*b^3)*d^2*e + 6*(B*a^2*b + A*a*b^2)*d*e^2 - (B*a^3 + 3*A*a^2*b)*e^3)*(e*x + d))/((e*x + d)^(7/2)*e^4))/e
```


Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 346 vs. $2(153) = 306$.

Time = 0.13 (sec) , antiderivative size = 346, normalized size of antiderivative = 2.07

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{9/2}} dx = \frac{2\sqrt{ex+d}Bb^3}{e^5} + \frac{2(140(ex+d)^3Bb^3d - 70(ex+d)^2Bb^3d^2 + 28(ex+d)Bb^3d^3 - 5Bb^3d^4 - 105(ex+d)^3Bab^2e - 35(ex+d)^2Bab^2e^2 + 105(ex+d)^3Aab^2e - 35(ex+d)^2Aab^2e^2 + 105(ex+d)^3A^2b^2e - 35(ex+d)^2A^2b^2e^2 - 42(ex+d)A^2b^2e^2 + 42(ex+d)A^2b^2e^2 - 15A^2b^2e^2 - 15A^2b^2e^2 - 7(ex+d)A^3e^3 - 21(ex+d)A^3e^3 + 5B^3a^3d^3e^3 + 15A^2b^2e^3 - 5A^3e^4)/((ex+d)^{7/2}e^5)}$$

input `integrate((b*x+a)^3*(B*x+A)/(e*x+d)^(9/2),x, algorithm="giac")`

output

```
2*sqrt(e*x + d)*B*b^3/e^5 + 2/35*(140*(e*x + d)^3*B*b^3*d - 70*(e*x + d)^2
*B*b^3*d^2 + 28*(e*x + d)*B*b^3*d^3 - 5*B*b^3*d^4 - 105*(e*x + d)^3*B*a*b^
2*e - 35*(e*x + d)^3*A*b^3*e + 105*(e*x + d)^2*B*a*b^2*d*e + 35*(e*x + d)^
2*A*b^3*d*e - 63*(e*x + d)*B*a*b^2*d^2*e - 21*(e*x + d)*A*b^3*d^2*e + 15*B
*a*b^2*d^3*e + 5*A*b^3*d^3*e - 35*(e*x + d)^2*B*a^2*b*e^2 - 35*(e*x + d)^2
*A*a*b^2*e^2 + 42*(e*x + d)*B*a^2*b*d*e^2 + 42*(e*x + d)*A*a*b^2*d*e^2 - 1
5*B*a^2*b*d^2*e^2 - 15*A*a*b^2*d^2*e^2 - 7*(e*x + d)*B*a^3*e^3 - 21*(e*x +
d)*A*a^2*b*e^3 + 5*B*a^3*d*e^3 + 15*A*a^2*b*d*e^3 - 5*A*a^3*e^4)/((e*x +
d)^(7/2)*e^5)
```

Mupad [B] (verification not implemented)

Time = 1.05 (sec) , antiderivative size = 300, normalized size of antiderivative = 1.80

$$\int \frac{(a+bx)^3(A+Bx)}{(d+ex)^{9/2}} dx = \frac{2(2Ba^3de^3 + 7Ba^3e^4x + 5Aa^3e^4 + 8Ba^2bd^2e^2 + 28Ba^2bde^3x + 6Aa^2bde^3 + 35Ba^2be^4x^2 + \dots)}{(d+ex)^{7/2}e^5}$$

input `int(((A + B*x)*(a + b*x)^3)/(d + e*x)^(9/2),x)`

3.148 $\int \frac{(A+Bx)(d+ex)^{7/2}}{a+bx} dx$

Optimal result	1374
Mathematica [A] (verified)	1375
Rubi [A] (verified)	1375
Maple [A] (verified)	1379
Fricas [B] (verification not implemented)	1379
Sympy [A] (verification not implemented)	1380
Maxima [F(-2)]	1381
Giac [B] (verification not implemented)	1381
Mupad [B] (verification not implemented)	1382
Reduce [B] (verification not implemented)	1383

Optimal result

Integrand size = 22, antiderivative size = 198

$$\int \frac{(A+Bx)(d+ex)^{7/2}}{a+bx} dx = \frac{2(Ab-aB)(bd-ae)^3\sqrt{d+ex}}{b^5} + \frac{2(Ab-aB)(bd-ae)^2(d+ex)^{3/2}}{3b^4} + \frac{2(Ab-aB)(bd-ae)(d+ex)^{5/2}}{5b^3} + \frac{2(Ab-aB)(d+ex)^{7/2}}{7b^2} + \frac{2B(d+ex)^{9/2}}{9be} - \frac{2(Ab-aB)(bd-ae)^{7/2}\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{11/2}}$$

output

```
2*(A*b-B*a)*(-a*e+b*d)^3*(e*x+d)^(1/2)/b^5+2/3*(A*b-B*a)*(-a*e+b*d)^2*(e*x+d)^(3/2)/b^4+2/5*(A*b-B*a)*(-a*e+b*d)*(e*x+d)^(5/2)/b^3+2/7*(A*b-B*a)*(e*x+d)^(7/2)/b^2+2/9*B*(e*x+d)^(9/2)/b/e-2*(A*b-B*a)*(-a*e+b*d)^(7/2)*arctanh(b^(1/2)*(e*x+d)^(1/2)/(-a*e+b*d)^(1/2))/b^(11/2)
```


$$\begin{aligned}
 & \frac{(Ab - aB) \left(\frac{(bd - ae) \int \frac{(d+ex)^{5/2}}{a+bx} dx}{b} + \frac{2(d+ex)^{7/2}}{7b} \right)}{b} + \frac{2B(d+ex)^{9/2}}{9be} \\
 & \quad \downarrow 60 \\
 & \frac{(Ab - aB) \left(\frac{(bd - ae) \left(\frac{(bd - ae) \int \frac{(d+ex)^{3/2}}{a+bx} dx + \frac{2(d+ex)^{5/2}}{5b}}{b} \right) + \frac{2(d+ex)^{7/2}}{7b}}{b} \right)}{b} + \frac{2B(d+ex)^{9/2}}{9be} \\
 & \quad \downarrow 60 \\
 & \frac{(Ab - aB) \left(\frac{(bd - ae) \left(\frac{(bd - ae) \left(\frac{(bd - ae) \int \frac{\sqrt{d+ex}}{a+bx} dx + \frac{2(d+ex)^{3/2}}{3b}}{b} \right) + \frac{2(d+ex)^{5/2}}{5b}}{b} \right) + \frac{2(d+ex)^{7/2}}{7b}}{b} \right)}{b} + \frac{2B(d+ex)^{9/2}}{9be} \right)}{b} + \frac{2B(d+ex)^{9/2}}{9be} \\
 & \quad \downarrow 60 \\
 & \frac{(Ab - aB) \left(\frac{(bd - ae) \left(\frac{(bd - ae) \left(\frac{(bd - ae) \int \frac{1}{(a+bx)\sqrt{d+ex}} dx + \frac{2\sqrt{d+ex}}{b}}{b} \right) + \frac{2(d+ex)^{3/2}}{3b}}{b} \right) + \frac{2(d+ex)^{5/2}}{5b}}{b} \right) + \frac{2(d+ex)^{7/2}}{7b}}{b} \right)}{b} + \frac{2B(d+ex)^{9/2}}{9be} \right)}{b} + \frac{2B(d+ex)^{9/2}}{9be} \\
 & \quad \downarrow 73
 \end{aligned}$$

$$\begin{aligned}
 & \left((Ab - aB) \frac{(bd - ae) \left(\frac{2(bd - ae) \int \frac{1}{a + \frac{b(d+ex)}{be} - \frac{bd}{e}} d\sqrt{d+ex} + \frac{2\sqrt{d+ex}}{b} \right)}{b} + \frac{2(d+ex)^{3/2}}{3b} \right) \\
 & \left((bd - ae) \frac{\left(\frac{2(bd - ae) \int \frac{1}{a + \frac{b(d+ex)}{be} - \frac{bd}{e}} d\sqrt{d+ex} + \frac{2\sqrt{d+ex}}{b} \right)}{b} + \frac{2(d+ex)^{5/2}}{5b} \right) \\
 & \left((Ab - aB) \frac{\left(\frac{2(bd - ae) \int \frac{1}{a + \frac{b(d+ex)}{be} - \frac{bd}{e}} d\sqrt{d+ex} + \frac{2\sqrt{d+ex}}{b} \right)}{b} + \frac{2(d+ex)^{7/2}}{7b} \right)
 \end{aligned}$$

$$\frac{2B(d+ex)^{9/2} b}{9be}$$

↓ 221

$$\begin{aligned}
 & \left((Ab - aB) \frac{(bd - ae) \left(\frac{2\sqrt{d+ex}}{b} - \frac{2\sqrt{bd-ae} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{3/2}} \right)}{b} + \frac{2(d+ex)^{3/2}}{3b} \right) \\
 & \left((bd - ae) \frac{\left(\frac{2\sqrt{d+ex}}{b} - \frac{2\sqrt{bd-ae} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{3/2}} \right)}{b} + \frac{2(d+ex)^{5/2}}{5b} \right) \\
 & \left((Ab - aB) \frac{\left(\frac{2\sqrt{d+ex}}{b} - \frac{2\sqrt{bd-ae} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{3/2}} \right)}{b} + \frac{2(d+ex)^{7/2}}{7b} \right)
 \end{aligned}$$

$$\frac{2B(d+ex)^{9/2} b}{9be}$$

input `Int[((A + B*x)*(d + e*x)^(7/2))/(a + b*x),x]`

output `(2*B*(d + e*x)^(9/2))/(9*b*e) + ((A*b - a*B)*((2*(d + e*x)^(7/2))/(7*b) + ((b*d - a*e)*((2*(d + e*x)^(5/2))/(5*b) + ((b*d - a*e)*((2*(d + e*x)^(3/2))/(3*b) + ((b*d - a*e)*((2*Sqrt[d + e*x])/b - (2*Sqrt[b*d - a*e]*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[b*d - a*e]])/b^(3/2)))/b))/b))/b)))/b`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [A] (verified)

Time = 0.74 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.36

method	result
pseudoelliptic	$2 \left(\left(\frac{-(ex+d)^4 B - \frac{176A \left(\frac{15}{176} e^3 x^3 + \frac{3}{8} e^2 d x^2 + \frac{61}{88} d^2 x e + d^3 \right) e}{3}}{3} \right) b^4 + \frac{58ae \left(\frac{\left(\frac{15}{2} e^3 x^3 + 33e^2 d x^2 + 61d^2 x e + 88d^3 \right) B}{203} + Ae \left(\frac{3}{58} e \right) \right)}{15} \right)$
risch	$-2(-35B e^4 b^4 x^4 - 45A b^4 e^4 x^3 + 45Ba b^3 e^4 x^3 - 140B b^4 d e^3 x^3 + 63Aa b^3 e^4 x^2 - 198A b^4 d e^3 x^2 - 63B a^2 b^2 e^4 x^2 + 198Ba$
derivativedivides	$2 \left(-\frac{B(ex+d)^{\frac{9}{2}} b^4}{9} - \frac{A b^4 e(ex+d)^{\frac{7}{2}}}{7} + \frac{B a b^3 e(ex+d)^{\frac{7}{2}}}{7} + \frac{A a b^3 e^2(ex+d)^{\frac{5}{2}}}{5} - \frac{A b^4 d e(ex+d)^{\frac{5}{2}}}{5} - \frac{B a^2 b^2 e^2(ex+d)^{\frac{5}{2}}}{5} + \frac{B a b^3 d e(ex+d)^{\frac{5}{2}}}{5} \right)$
default	$2 \left(-\frac{B(ex+d)^{\frac{9}{2}} b^4}{9} - \frac{A b^4 e(ex+d)^{\frac{7}{2}}}{7} + \frac{B a b^3 e(ex+d)^{\frac{7}{2}}}{7} + \frac{A a b^3 e^2(ex+d)^{\frac{5}{2}}}{5} - \frac{A b^4 d e(ex+d)^{\frac{5}{2}}}{5} - \frac{B a^2 b^2 e^2(ex+d)^{\frac{5}{2}}}{5} + \frac{B a b^3 d e(ex+d)^{\frac{5}{2}}}{5} \right)$

input `int((B*x+A)*(e*x+d)^(7/2)/(b*x+a),x,method=_RETURNVERBOSE)`

output `-2*((1/3*(-1/3*(e*x+d)^4*B-176/35*A*(15/176*e^3*x^3+3/8*e^2*d*x^2+61/88*d^2*x*e+d^3)*e)*b^4+58/15*a*e*(1/203*(15/2*e^3*x^3+33*e^2*d*x^2+61*d^2*x*e+88*d^3)*B+A*e*(3/58*e^2*x^2+8/29*d*e*x+d^2))*b^3-10/3*a^2*e^2*(1/25*(3/2*e^2*x^2+8*d*e*x+29*d^2)*B+A*e*(1/10*e*x+d))*b^2+a^3*(1/3*(e*x+10*d)*B+A*e)*e^3*b-B*a^4*e^4)*((a*e-b*d)*b)^(1/2)*(e*x+d)^(1/2)-e*(a*e-b*d)^4*(A*b-B*a)*arctan(b*(e*x+d)^(1/2)/((a*e-b*d)*b)^(1/2))/((a*e-b*d)*b)^(1/2)/e/b^5`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 427 vs. 2(170) = 340.

Time = 0.10 (sec) , antiderivative size = 865, normalized size of antiderivative = 4.37

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{a + bx} dx = \text{Too large to display}$$

input `integrate((B*x+A)*(e*x+d)^(7/2)/(b*x+a),x, algorithm="fricas")`

output

```
[1/315*(315*((B*a*b^3 - A*b^4)*d^3*e - 3*(B*a^2*b^2 - A*a*b^3)*d^2*e^2 + 3
*(B*a^3*b - A*a^2*b^2)*d*e^3 - (B*a^4 - A*a^3*b)*e^4)*sqrt((b*d - a*e)/b)*
log((b*e*x + 2*b*d - a*e + 2*sqrt(e*x + d)*b*sqrt((b*d - a*e)/b))/(b*x + a
)) + 2*(35*B*b^4*e^4*x^4 + 35*B*b^4*d^4 - 528*(B*a*b^3 - A*b^4)*d^3*e + 12
18*(B*a^2*b^2 - A*a*b^3)*d^2*e^2 - 1050*(B*a^3*b - A*a^2*b^2)*d*e^3 + 315*
(B*a^4 - A*a^3*b)*e^4 + 5*(28*B*b^4*d*e^3 - 9*(B*a*b^3 - A*b^4)*e^4)*x^3 +
3*(70*B*b^4*d^2*e^2 - 66*(B*a*b^3 - A*b^4)*d*e^3 + 21*(B*a^2*b^2 - A*a*b^
3)*e^4)*x^2 + (140*B*b^4*d^3*e - 366*(B*a*b^3 - A*b^4)*d^2*e^2 + 336*(B*a^
2*b^2 - A*a*b^3)*d*e^3 - 105*(B*a^3*b - A*a^2*b^2)*e^4)*x)*sqrt(e*x + d))/
(b^5*e), 2/315*(315*((B*a*b^3 - A*b^4)*d^3*e - 3*(B*a^2*b^2 - A*a*b^3)*d^2
*e^2 + 3*(B*a^3*b - A*a^2*b^2)*d*e^3 - (B*a^4 - A*a^3*b)*e^4)*sqrt(-(b*d -
a*e)/b)*arctan(-sqrt(e*x + d)*b*sqrt(-(b*d - a*e)/b)/(b*d - a*e)) + (35*B
*b^4*e^4*x^4 + 35*B*b^4*d^4 - 528*(B*a*b^3 - A*b^4)*d^3*e + 1218*(B*a^2*b^
2 - A*a*b^3)*d^2*e^2 - 1050*(B*a^3*b - A*a^2*b^2)*d*e^3 + 315*(B*a^4 - A*a
^3*b)*e^4 + 5*(28*B*b^4*d*e^3 - 9*(B*a*b^3 - A*b^4)*e^4)*x^3 + 3*(70*B*b^4
*d^2*e^2 - 66*(B*a*b^3 - A*b^4)*d*e^3 + 21*(B*a^2*b^2 - A*a*b^3)*e^4)*x^2
+ (140*B*b^4*d^3*e - 366*(B*a*b^3 - A*b^4)*d^2*e^2 + 336*(B*a^2*b^2 - A*a*
b^3)*d*e^3 - 105*(B*a^3*b - A*a^2*b^2)*e^4)*x)*sqrt(e*x + d))/(b^5*e)]
```

Sympy [A] (verification not implemented)

Time = 4.24 (sec) , antiderivative size = 376, normalized size of antiderivative = 1.90

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{a + bx} dx = \begin{cases} 2 \left(\frac{B(d+ex)^{9/2}}{9b} + \frac{(d+ex)^{7/2}(Abe - Bae)}{7b^2} + \frac{(d+ex)^{5/2}(-Aabe^2 + Ab^2de + Ba^2e^2 - Babde)}{5b^3} + \frac{(d+ex)^{3/2}(Aa^2be^3 - 2Aab^2e^2)}{3b^4} \right) \\ d^{7/2} \left(\frac{Bx}{b} - \frac{(-Ab+Ba) \left(\begin{cases} \frac{x}{a} & \text{for } b = 0 \\ \frac{\log(a+bx)}{b} & \text{otherwise} \end{cases} \right)}{b} \right) \end{cases}$$

input

```
integrate((B*x+A)*(e*x+d)**(7/2)/(b*x+a), x)
```

output

```
Piecewise((2*(B*(d + e*x)**(9/2)/(9*b) + (d + e*x)**(7/2)*(A*b*e - B*a*e)/
(7*b**2) + (d + e*x)**(5/2)*(-A*a*b*e**2 + A*b**2*d*e + B*a**2*e**2 - B*a*
b*d*e)/(5*b**3) + (d + e*x)**(3/2)*(A*a**2*b*e**3 - 2*A*a*b**2*d*e**2 + A*
b**3*d**2*e - B*a**3*e**3 + 2*B*a**2*b*d*e**2 - B*a*b**2*d**2*e)/(3*b**4)
+ sqrt(d + e*x)*(-A*a**3*b*e**4 + 3*A*a**2*b**2*d*e**3 - 3*A*a*b**3*d**2*e
**2 + A*b**4*d**3*e + B*a**4*e**4 - 3*B*a**3*b*d*e**3 + 3*B*a**2*b**2*d**2
*e**2 - B*a*b**3*d**3*e)/b**5 - e*(-A*b + B*a)*(a*e - b*d)**4*atan(sqrt(d
+ e*x)/sqrt((a*e - b*d)/b))/(b**6*sqrt((a*e - b*d)/b))/e, Ne(e, 0)), (d**
(7/2)*(B*x/b - (-A*b + B*a)*Piecewise((x/a, Eq(b, 0)), (log(a + b*x)/b, Tr
ue))/b), True))
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{a + bx} dx = \text{Exception raised: ValueError}$$

input

```
integrate((B*x+A)*(e*x+d)^(7/2)/(b*x+a),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for m
ore detail
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 560 vs. $2(170) = 340$.

Time = 0.13 (sec) , antiderivative size = 560, normalized size of antiderivative = 2.83

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{a + bx} dx =$$

$$\frac{2(Bab^4d^4 - Ab^5d^4 - 4Ba^2b^3d^3e + 4Aab^4d^3e + 6Ba^3b^2d^2e^2 - 6Aa^2b^3d^2e^2 - 4Ba^4bde^3 + 4Aa^3b^2de^3 + \sqrt{-b^2d + abeb^5}}{\sqrt{-b^2d + abeb^5}}$$

$$+ \frac{2\left(35(ex + d)^{\frac{9}{2}}Bb^8e^8 - 45(ex + d)^{\frac{7}{2}}Bab^7e^9 + 45(ex + d)^{\frac{7}{2}}Ab^8e^9 - 63(ex + d)^{\frac{5}{2}}Bab^7de^9 + 63(ex + d)^{\frac{5}{2}}\right)}{\sqrt{-b^2d + abeb^5}}$$

input `integrate((B*x+A)*(e*x+d)^(7/2)/(b*x+a),x, algorithm="giac")`

output

$$\begin{aligned}
 & -2*(B*a*b^4*d^4 - A*b^5*d^4 - 4*B*a^2*b^3*d^3*e + 4*A*a*b^4*d^3*e + 6*B*a^3*b^2*d^2*e^2 - 6*A*a^2*b^3*d^2*e^2 - 4*B*a^4*b*d*e^3 + 4*A*a^3*b^2*d*e^3 + B*a^5*e^4 - A*a^4*b*e^4)*\arctan(\sqrt{e*x + d}*b/\sqrt{-b^2*d + a*b*e})/(\sqrt{-b^2*d + a*b*e}*b^5) + 2/315*(35*(e*x + d)^(9/2)*B*b^8*e^8 - 45*(e*x + d)^(7/2)*B*a*b^7*e^9 + 45*(e*x + d)^(7/2)*A*b^8*e^9 - 63*(e*x + d)^(5/2)*B*a*b^7*d^2*e^9 + 63*(e*x + d)^(5/2)*A*b^8*d^2*e^9 - 105*(e*x + d)^(3/2)*B*a*b^7*d^2*e^9 + 105*(e*x + d)^(3/2)*A*b^8*d^2*e^9 - 315*\sqrt{e*x + d}*B*a*b^7*d^3*e^9 + 315*\sqrt{e*x + d}*A*b^8*d^3*e^9 + 63*(e*x + d)^(5/2)*B*a^2*b^6*e^10 - 63*(e*x + d)^(5/2)*A*a*b^7*e^10 + 210*(e*x + d)^(3/2)*B*a^2*b^6*d*e^10 - 210*(e*x + d)^(3/2)*A*a*b^7*d*e^10 + 945*\sqrt{e*x + d}*B*a^2*b^6*d^2*e^10 - 945*\sqrt{e*x + d}*A*a*b^7*d^2*e^10 - 105*(e*x + d)^(3/2)*B*a^3*b^5*e^11 + 105*(e*x + d)^(3/2)*A*a^2*b^6*e^11 - 945*\sqrt{e*x + d}*B*a^3*b^5*d*e^11 + 945*\sqrt{e*x + d}*A*a^2*b^6*d*e^11 + 315*\sqrt{e*x + d}*B*a^4*b^4*e^12 - 315*\sqrt{e*x + d}*A*a^3*b^5*e^12)/(b^9*e^9)
 \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 1.03 (sec) , antiderivative size = 426, normalized size of antiderivative = 2.15

$$\begin{aligned}
 & \int \frac{(A + Bx)(d + ex)^{7/2}}{a + bx} dx = \left(\frac{2Ae - 2Bd}{7be} - \frac{2B(ae^2 - bde)}{7b^2e^2} \right) (d + ex)^{7/2} \\
 & + \frac{2B(d + ex)^{9/2}}{9be} \\
 & + \frac{2 \operatorname{atan}\left(\frac{\sqrt{b}(Ab - Ba)(ae - bd)^{7/2} \sqrt{d + ex}}{-Ba^5e^4 + 4Ba^4bde^3 + Aa^4be^4 - 6Ba^3b^2d^2e^2 - 4Aa^3b^2de^3 + 4Ba^2b^3d^3e + 6Aa^2b^3d^2e^2 - Ba^2b^4d^4 - 4Aa^2b^4d^3e + Ab^5d^4} \right)}{b^{11/2}} \\
 & + \frac{\left(\frac{2Ae - 2Bd}{be} - \frac{2B(ae^2 - bde)}{b^2e^2} \right) (ae^2 - bde)^2 (d + ex)^{3/2}}{3b^2e^2} \\
 & - \frac{\left(\frac{2Ae - 2Bd}{be} - \frac{2B(ae^2 - bde)}{b^2e^2} \right) (ae^2 - bde)^3 \sqrt{d + ex}}{b^3e^3} \\
 & - \frac{\left(\frac{2Ae - 2Bd}{be} - \frac{2B(ae^2 - bde)}{b^2e^2} \right) (ae^2 - bde) (d + ex)^{5/2}}{5be}
 \end{aligned}$$

input `int(((A + B*x)*(d + e*x)^(7/2))/(a + b*x),x)`

output

```
((2*A*e - 2*B*d)/(7*b*e) - (2*B*(a*e^2 - b*d*e))/(7*b^2*e^2))*(d + e*x)^(7/2) + (2*B*(d + e*x)^(9/2))/(9*b*e) + (2*atan((b^(1/2)*(A*b - B*a)*(a*e - b*d)^(7/2)*(d + e*x)^(1/2))/(A*b^5*d^4 - B*a^5*e^4 + A*a^4*b*e^4 - B*a*b^4*d^4 - 4*A*a^3*b^2*d*e^3 + 4*B*a^2*b^3*d^3*e + 6*A*a^2*b^3*d^2*e^2 - 6*B*a^3*b^2*d^2*e^2 - 4*A*a*b^4*d^3*e + 4*B*a^4*b*d*e^3))*(A*b - B*a)*(a*e - b*d)^(7/2))/b^(11/2) + (((2*A*e - 2*B*d)/(b*e) - (2*B*(a*e^2 - b*d*e))/(b^2*e^2))*(a*e^2 - b*d*e)^2*(d + e*x)^(3/2))/(3*b^2*e^2) - (((2*A*e - 2*B*d)/(b*e) - (2*B*(a*e^2 - b*d*e))/(b^2*e^2))*(a*e^2 - b*d*e)^3*(d + e*x)^(1/2))/(b^3*e^3) - (((2*A*e - 2*B*d)/(b*e) - (2*B*(a*e^2 - b*d*e))/(b^2*e^2))*(a*e^2 - b*d*e)*(d + e*x)^(5/2))/(5*b*e)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.25

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{a + bx} dx = \frac{2\sqrt{ex + d}(e^4x^4 + 4de^3x^3 + 6d^2e^2x^2 + 4d^3ex + d^4)}{9e}$$

input

```
int((B*x+A)*(e*x+d)^(7/2)/(b*x+a),x)
```

output

```
(2*sqrt(d + e*x)*(d**4 + 4*d**3*e*x + 6*d**2*e**2*x**2 + 4*d*e**3*x**3 + e**4*x**4))/(9*e)
```

3.149 $\int \frac{(A+Bx)(d+ex)^{5/2}}{a+bx} dx$

Optimal result	1384
Mathematica [A] (verified)	1385
Rubi [A] (verified)	1385
Maple [A] (verified)	1388
Fricas [B] (verification not implemented)	1389
Sympy [A] (verification not implemented)	1390
Maxima [F(-2)]	1390
Giac [B] (verification not implemented)	1391
Mupad [B] (verification not implemented)	1392
Reduce [B] (verification not implemented)	1392

Optimal result

Integrand size = 22, antiderivative size = 164

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{a + bx} dx = \frac{2(Ab - aB)(bd - ae)^2 \sqrt{d + ex}}{b^4} + \frac{2(Ab - aB)(bd - ae)(d + ex)^{3/2}}{3b^3} + \frac{2(Ab - aB)(d + ex)^{5/2}}{5b^2} + \frac{2B(d + ex)^{7/2}}{7be} - \frac{2(Ab - aB)(bd - ae)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{9/2}}$$

output

```
2*(A*b-B*a)*(-a*e+b*d)^2*(e*x+d)^(1/2)/b^4+2/3*(A*b-B*a)*(-a*e+b*d)*(e*x+d)^(3/2)/b^3+2/5*(A*b-B*a)*(e*x+d)^(5/2)/b^2+2/7*B*(e*x+d)^(7/2)/b/e-2*(A*b-B*a)*(-a*e+b*d)^(5/2)*arctanh(b^(1/2)*(e*x+d)^(1/2)/(-a*e+b*d)^(1/2))/b^(9/2)
```

Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.13

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{a + bx} dx = \frac{2\sqrt{d + ex}(-105a^3Be^3 + 35a^2be^2(7Bd + 3Ae + Bex) - 7ab^2e(5Ae(7d + ex) + 2(Ab - aB)(-bd + ae)^{5/2} \arctan\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right))}{b^{9/2}}$$

input

```
Integrate[((A + B*x)*(d + e*x)^(5/2))/(a + b*x),x]
```

output

```
(2*sqrt[d + e*x]*(-105*a^3*B*e^3 + 35*a^2*b*e^2*(7*B*d + 3*A*e + B*e*x) - 7*a*b^2*e*(5*A*e*(7*d + e*x) + B*(23*d^2 + 11*d*e*x + 3*e^2*x^2)) + b^3*(15*B*(d + e*x)^3 + 7*A*e*(23*d^2 + 11*d*e*x + 3*e^2*x^2)))/(105*b^4*e) - (2*(A*b - a*B)*(-b*d + a*e)^(5/2)*ArcTan[(sqrt[b]*sqrt[d + e*x])/sqrt[-(b*d + a*e)]]/b^(9/2))
```

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.93, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {90, 60, 60, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{a + bx} dx$$

$$\downarrow 90$$

$$\frac{(Ab - aB) \int \frac{(d+ex)^{5/2}}{a+bx} dx}{b} + \frac{2B(d + ex)^{7/2}}{7be}$$

$$\downarrow 60$$

$$\frac{(Ab - aB) \left(\frac{(bd - ae) \int \frac{(d+ex)^{3/2}}{a+bx} dx}{b} + \frac{2(d+ex)^{5/2}}{5b} \right)}{b} + \frac{2B(d + ex)^{7/2}}{7be}$$

$$\begin{aligned}
 & \downarrow 60 \\
 & \frac{(Ab - aB) \left(\frac{(bd - ae) \left(\frac{(bd - ae) \int \frac{\sqrt{d+ex}}{a+bx} dx + 2(d+ex)^{3/2}}{3b} \right)}{b} + \frac{2(d+ex)^{5/2}}{5b} \right)}{b} + \frac{2B(d+ex)^{7/2}}{7be} \\
 & \downarrow 60 \\
 & \frac{(Ab - aB) \left(\frac{(bd - ae) \left(\frac{(bd - ae) \int \frac{1}{(a+bx)\sqrt{d+ex}} dx + \frac{2\sqrt{d+ex}}{b}}{b} \right) + \frac{2(d+ex)^{3/2}}{3b} \right)}{b} + \frac{2(d+ex)^{5/2}}{5b} \right)}{b} + \frac{2B(d+ex)^{7/2}}{7be} \\
 & \downarrow 73 \\
 & \frac{(Ab - aB) \left(\frac{(bd - ae) \left(\frac{2(bd - ae) \int \frac{1}{a + \frac{b(d+ex)}{e} - \frac{bd}{e}} d\sqrt{d+ex}}{be} + \frac{2\sqrt{d+ex}}{b} \right)}{b} + \frac{2(d+ex)^{3/2}}{3b} \right)}{b} + \frac{2(d+ex)^{5/2}}{5b} \right)}{b} + \frac{2B(d+ex)^{7/2}}{7be} \\
 & \downarrow 221
 \end{aligned}$$

$$(Ab - aB) \left(\frac{(bd - ae) \left(\frac{2\sqrt{d+ex}}{b} - \frac{2\sqrt{bd-ae} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{3/2}} \right) + \frac{2(d+ex)^{3/2}}{3b}}{b} \right) + \frac{2(d+ex)^{5/2}}{5b} \right) + \frac{b}{2B(d+ex)^{7/2}} - \frac{7be}{7be}$$

input `Int[((A + B*x)*(d + e*x)^(5/2))/(a + b*x), x]`

output `(2*B*(d + e*x)^(7/2))/(7*b*e) + ((A*b - a*B)*((2*(d + e*x)^(5/2))/(5*b) + ((b*d - a*e)*((2*(d + e*x)^(3/2))/(3*b) + ((b*d - a*e)*((2*sqrt[d + e*x])/b - (2*sqrt[b*d - a*e]*ArcTanh[(sqrt[b]*sqrt[d + e*x])/sqrt[b*d - a*e]])/b^(3/2)))/b))/b`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 90

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_]
:> Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x]
+ Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2))
Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x]
&& NeQ[n + p + 2, 0]
```

rule 221

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol]
:> Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x]
&& NegQ[a/b]
```

Maple [A] (verified)

Time = 0.45 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.17

method	result
pseudoelliptic	$\frac{2\sqrt{(ae-db)b} \left(\left(\frac{(ex+d)^3 B}{7} + \frac{23A \left(\frac{3}{23} e^2 x^2 + \frac{11}{23} dex + d^2 \right) e}{15} \right) b^3 - \frac{7ae \left(\frac{(3e^2 x^2 + 11dex + 23d^2) B}{35} + Ae \left(\frac{ex}{7} + d \right) \right) b^2}{3} + a^2 \left(\frac{(ex+7d)}{3} \right) \right)}{e b^4 \sqrt{(ae-db)b}}$
risch	$\frac{2(15b^3 B x^3 e^3 + 21A b^3 e^3 x^2 - 21B a b^2 e^3 x^2 + 45B b^3 d e^2 x^2 - 35A x a b^2 e^3 + 77A x b^3 d e^2 + 35B x a^2 b e^3 - 77B a b^2 d e^2 x + 45A a^2 b^2 e^3)}{105e b^4}$
derivativedivides	$\frac{2 \left(\frac{b^3 B (ex+d)^{\frac{7}{2}}}{7} + \frac{A b^3 e (ex+d)^{\frac{5}{2}}}{5} - \frac{B a b^2 e (ex+d)^{\frac{5}{2}}}{5} - \frac{A a b^2 e^2 (ex+d)^{\frac{3}{2}}}{3} + \frac{A b^3 d e (ex+d)^{\frac{3}{2}}}{3} + \frac{B a^2 b e^2 (ex+d)^{\frac{3}{2}}}{3} - \frac{B a b^2 d e (ex+d)^{\frac{3}{2}}}{3} \right)}{b^4}$
default	$\frac{2 \left(\frac{b^3 B (ex+d)^{\frac{7}{2}}}{7} + \frac{A b^3 e (ex+d)^{\frac{5}{2}}}{5} - \frac{B a b^2 e (ex+d)^{\frac{5}{2}}}{5} - \frac{A a b^2 e^2 (ex+d)^{\frac{3}{2}}}{3} + \frac{A b^3 d e (ex+d)^{\frac{3}{2}}}{3} + \frac{B a^2 b e^2 (ex+d)^{\frac{3}{2}}}{3} - \frac{B a b^2 d e (ex+d)^{\frac{3}{2}}}{3} \right)}{b^4}$

input

```
int((B*x+A)*(e*x+d)^(5/2)/(b*x+a), x, method=_RETURNVERBOSE)
```

output

```
2*(((a*e-b*d)*b)^(1/2)*((1/7*(e*x+d)^3*B+23/15*A*(3/23*e^2*x^2+11/23*d*e*x+d^2)*e)*b^3-7/3*a*e*(1/35*(3*e^2*x^2+11*d*e*x+23*d^2)*B+A*e*(1/7*e*x+d))*b^2+a^2*(1/3*(e*x+7*d)*B+A*e)*e^2*b-B*a^3*e^3)*(e*x+d)^(1/2)-e*(a*e-b*d)^3*(A*b-B*a)*arctan(b*(e*x+d)^(1/2)/((a*e-b*d)*b)^(1/2))/((a*e-b*d)*b)^(1/2)/e/b^4
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 290 vs. $2(140) = 280$.

Time = 0.09 (sec) , antiderivative size = 591, normalized size of antiderivative = 3.60

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{a + bx} dx = \left[\frac{105 ((Bab^2 - Ab^3)d^2e - 2(Ba^2b - Aab^2)de^2 + (Ba^3 - Aa^2b)e^3) \sqrt{\frac{bd-ae}{b}}}{\dots} \right]$$

input `integrate((B*x+A)*(e*x+d)^(5/2)/(b*x+a),x, algorithm="fricas")`

output `[-1/105*(105*((B*a*b^2 - A*b^3)*d^2*e - 2*(B*a^2*b - A*a*b^2)*d*e^2 + (B*a^3 - A*a^2*b)*e^3)*sqrt((b*d - a*e)/b)*log((b*e*x + 2*b*d - a*e - 2*sqrt(e*x + d)*b*sqrt((b*d - a*e)/b))/(b*x + a)) - 2*(15*B*b^3*e^3*x^3 + 15*B*b^3*d^3 - 161*(B*a*b^2 - A*b^3)*d^2*e + 245*(B*a^2*b - A*a*b^2)*d*e^2 - 105*(B*a^3 - A*a^2*b)*e^3 + 3*(15*B*b^3*d*e^2 - 7*(B*a*b^2 - A*b^3)*e^3)*x^2 + (45*B*b^3*d^2*e - 77*(B*a*b^2 - A*b^3)*d*e^2 + 35*(B*a^2*b - A*a*b^2)*e^3)*x)*sqrt(e*x + d))/(b^4*e), 2/105*(105*((B*a*b^2 - A*b^3)*d^2*e - 2*(B*a^2*b - A*a*b^2)*d*e^2 + (B*a^3 - A*a^2*b)*e^3)*sqrt(-(b*d - a*e)/b)*arctan(-sqrt(e*x + d)*b*sqrt(-(b*d - a*e)/b)/(b*d - a*e)) + (15*B*b^3*e^3*x^3 + 15*B*b^3*d^3 - 161*(B*a*b^2 - A*b^3)*d^2*e + 245*(B*a^2*b - A*a*b^2)*d*e^2 - 105*(B*a^3 - A*a^2*b)*e^3 + 3*(15*B*b^3*d*e^2 - 7*(B*a*b^2 - A*b^3)*e^3)*x^2 + (45*B*b^3*d^2*e - 77*(B*a*b^2 - A*b^3)*d*e^2 + 35*(B*a^2*b - A*a*b^2)*e^3)*x)*sqrt(e*x + d))/(b^4*e)]`

Sympy [A] (verification not implemented)

Time = 3.96 (sec) , antiderivative size = 258, normalized size of antiderivative = 1.57

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{a + bx} dx = \begin{cases} 2 \left(\frac{B(d+ex)^{7/2}}{7b} + \frac{(d+ex)^{5/2}(Abe - Bae)}{5b^2} + \frac{(d+ex)^{3/2}(-Aabe^2 + Ab^2de + Ba^2e^2 - Babde)}{3b^3} + \frac{\sqrt{d+ex}(Aa^2be^3 - 2Aab^2d)}{3b^3} \right) \\ d^{5/2} \left(\frac{Bx}{b} - \frac{(-Ab+Ba) \begin{pmatrix} \frac{x}{a} & \text{for } b = 0 \\ \frac{\log(a+bx)}{b} & \text{otherwise} \end{pmatrix}}{b} \right) \end{cases}$$

input `integrate((B*x+A)*(e*x+d)**(5/2)/(b*x+a), x)`output `Piecewise((2*(B*(d + e*x)**(7/2)/(7*b) + (d + e*x)**(5/2)*(A*b*e - B*a*e)/(5*b**2) + (d + e*x)**(3/2)*(-A*a*b*e**2 + A*b**2*d*e + B*a**2*e**2 - B*a*b*d*e)/(3*b**3) + sqrt(d + e*x)*(A*a**2*b*e**3 - 2*A*a*b**2*d*e**2 + A*b**3*d**2*e - B*a**3*e**3 + 2*B*a**2*b*d*e**2 - B*a*b**2*d**2*e)/b**4 + e*(-A*b + B*a)*(a*e - b*d)**3*atan(sqrt(d + e*x)/sqrt((a*e - b*d)/b))/(b**5*sqrt((a*e - b*d)/b)))/e, Ne(e, 0)), (d**(5/2)*(B*x/b - (-A*b + B*a)*Piecewise((x/a, Eq(b, 0)), (log(a + b*x)/b, True))/b), True))`**Maxima [F(-2)]**

Exception generated.

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{a + bx} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)*(e*x+d)^(5/2)/(b*x+a), x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for m
ore detail
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 371 vs. $2(140) = 280$.

Time = 0.13 (sec) , antiderivative size = 371, normalized size of antiderivative = 2.26

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{a + bx} dx =$$

$$\frac{2(Bab^3d^3 - Ab^4d^3 - 3Ba^2b^2d^2e + 3Aab^3d^2e + 3Ba^3bde^2 - 3Aa^2b^2de^2 - Ba^4e^3 + Aa^3be^3) \arctan\left(\frac{\sqrt{ex+d}}{\sqrt{-b^2d+abeb^4}}\right) + 2\left(15(ex+d)^{\frac{7}{2}}Bb^6e^6 - 21(ex+d)^{\frac{5}{2}}Bab^5e^7 + 21(ex+d)^{\frac{5}{2}}Ab^6e^7 - 35(ex+d)^{\frac{3}{2}}Bab^5de^7 + 35(ex+d)^{\frac{3}{2}}\right)}{\sqrt{-b^2d+abeb^4}}$$

input

```
integrate((B*x+A)*(e*x+d)^(5/2)/(b*x+a),x, algorithm="giac")
```

output

```
-2*(B*a*b^3*d^3 - A*b^4*d^3 - 3*B*a^2*b^2*d^2*e + 3*A*a*b^3*d^2*e + 3*B*a^3*b*d*e^2 - 3*A*a^2*b^2*d*e^2 - B*a^4*e^3 + A*a^3*b*e^3)*arctan(sqrt(e*x + d)*b/sqrt(-b^2*d + a*b*e))/(sqrt(-b^2*d + a*b*e)*b^4) + 2/105*(15*(e*x + d)^(7/2)*B*b^6*e^6 - 21*(e*x + d)^(5/2)*B*a*b^5*e^7 + 21*(e*x + d)^(5/2)*A*b^6*e^7 - 35*(e*x + d)^(3/2)*B*a*b^5*d*e^7 + 35*(e*x + d)^(3/2)*A*b^6*d*e^7 - 105*sqrt(e*x + d)*B*a*b^5*d^2*e^7 + 105*sqrt(e*x + d)*A*b^6*d^2*e^7 + 35*(e*x + d)^(3/2)*B*a^2*b^4*e^8 - 35*(e*x + d)^(3/2)*A*a*b^5*e^8 + 210*sqrt(e*x + d)*B*a^2*b^4*d*e^8 - 210*sqrt(e*x + d)*A*a*b^5*d*e^8 - 105*sqrt(e*x + d)*B*a^3*b^3*e^9 + 105*sqrt(e*x + d)*A*a^2*b^4*e^9)/(b^7*e^7)
```

Mupad [B] (verification not implemented)

Time = 1.00 (sec) , antiderivative size = 330, normalized size of antiderivative = 2.01

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{a + bx} dx = \left(\frac{2Ae - 2Bd}{5be} - \frac{2B(ae^2 - bde)}{5b^2e^2} \right) (d + ex)^{5/2} + \frac{2 \operatorname{atan}\left(\frac{\sqrt{b}(Ab - Ba)(ae - bd)^{5/2} \sqrt{d + ex}}{Ba^4e^3 - 3Ba^3bde^2 - Aa^3be^3 + 3Ba^2b^2d^2e + 3Aa^2b^2de^2 - Ba^3b^3d^3 - 3Aab^3d^2e + Ab^4d^3}\right) (Ab - Ba)(ae - bd)^{5/2}}{b^{9/2}} + \frac{2B(d + ex)^{7/2}}{7be} + \frac{\left(\frac{2Ae - 2Bd}{be} - \frac{2B(ae^2 - bde)}{b^2e^2}\right) (ae^2 - bde)^2 \sqrt{d + ex}}{b^2e^2} - \frac{\left(\frac{2Ae - 2Bd}{be} - \frac{2B(ae^2 - bde)}{b^2e^2}\right) (ae^2 - bde)(d + ex)^{3/2}}{3be}$$

input `int(((A + B*x)*(d + e*x)^(5/2))/(a + b*x), x)`output
$$\left(\frac{2Ae - 2Bd}{5be} - \frac{2B(ae^2 - bde)}{5b^2e^2}\right)(d + ex)^{5/2} + \frac{2 \operatorname{atan}\left(\frac{b^{1/2}(Ab - Ba)(ae - bd)^{5/2}(d + ex)^{1/2}}{A^4d^3 + B^4e^3 - A^3bde^3 - B^3abd^3 + 3A^2b^2d^2e + 3B^2a^2b^2d^2e - 3A^2ab^3d^2e - 3B^2a^3b^3d^2e}\right)(Ab - Ba)(ae - bd)^{5/2}}{b^{9/2}} + \frac{2B(d + ex)^{7/2}}{7be} + \frac{\left(\frac{2Ae - 2Bd}{be} - \frac{2B(ae^2 - bde)}{b^2e^2}\right)(ae^2 - bde)^2(d + ex)^{1/2}}{b^2e^2} - \frac{\left(\frac{2Ae - 2Bd}{be} - \frac{2B(ae^2 - bde)}{b^2e^2}\right)(ae^2 - bde)(d + ex)^{3/2}}{3be}$$
Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.23

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{a + bx} dx = \frac{2\sqrt{ex + d}(e^3x^3 + 3de^2x^2 + 3d^2ex + d^3)}{7e}$$

input `int((B*x+A)*(e*x+d)^(5/2)/(b*x+a), x)`output
$$(2\sqrt{d + ex}(d^3 + 3d^2ex + 3de^2x^2 + e^3x^3))/(7e)$$

3.150 $\int \frac{(A+Bx)(d+ex)^{3/2}}{a+bx} dx$

Optimal result	1393
Mathematica [A] (verified)	1393
Rubi [A] (verified)	1394
Maple [A] (verified)	1396
Fricas [A] (verification not implemented)	1397
Sympy [A] (verification not implemented)	1397
Maxima [F(-2)]	1398
Giac [B] (verification not implemented)	1398
Mupad [B] (verification not implemented)	1399
Reduce [B] (verification not implemented)	1399

Optimal result

Integrand size = 22, antiderivative size = 130

$$\int \frac{(A+Bx)(d+ex)^{3/2}}{a+bx} dx = \frac{2(Ab-aB)(bd-ae)\sqrt{d+ex}}{b^3} + \frac{2(Ab-aB)(d+ex)^{3/2}}{3b^2} + \frac{2B(d+ex)^{5/2}}{5be} - \frac{2(Ab-aB)(bd-ae)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{7/2}}$$

output

```
2*(A*b-B*a)*(-a*e+b*d)*(e*x+d)^(1/2)/b^3+2/3*(A*b-B*a)*(e*x+d)^(3/2)/b^2+2/5*B*(e*x+d)^(5/2)/b/e-2*(A*b-B*a)*(-a*e+b*d)^(3/2)*arctanh(b^(1/2)*(e*x+d)^(1/2)/(-a*e+b*d)^(1/2))/b^(7/2)
```

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.99

$$\int \frac{(A+Bx)(d+ex)^{3/2}}{a+bx} dx = \frac{2\sqrt{d+ex}(15a^2Be^2-5abe(4Bd+3Ae+Bex))+b^2(3B(d+ex)^2+5Ae(4d+3e))}{15b^3e} + \frac{2(Ab-aB)(-bd+ae)^{3/2} \arctan\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right)}{b^{7/2}}$$

input `Integrate[((A + B*x)*(d + e*x)^(3/2))/(a + b*x),x]`

output $(2*\sqrt{d + e*x}*(15*a^2*B*e^2 - 5*a*b*e*(4*B*d + 3*A*e + B*e*x) + b^2*(3*B*(d + e*x)^2 + 5*A*e*(4*d + e*x)))/(15*b^3*e) + (2*(A*b - a*B)*(-(b*d) + a*e)^(3/2)*\text{ArcTan}[\sqrt{b}*\sqrt{d + e*x}]/\sqrt{-(b*d) + a*e}])/b^(7/2)$

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.95, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.227$, Rules used = {90, 60, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{a + bx} dx$$

$$\downarrow 90$$

$$\frac{(Ab - aB) \int \frac{(d+ex)^{3/2}}{a+bx} dx}{b} + \frac{2B(d + ex)^{5/2}}{5be}$$

$$\downarrow 60$$

$$\frac{(Ab - aB) \left(\frac{(bd-ae) \int \frac{\sqrt{d+ex}}{a+bx} dx}{b} + \frac{2(d+ex)^{3/2}}{3b} \right)}{b} + \frac{2B(d + ex)^{5/2}}{5be}$$

$$\downarrow 60$$

$$\frac{(Ab - aB) \left(\frac{(bd-ae) \left(\frac{(bd-ae) \int \frac{1}{(a+bx)\sqrt{d+ex}} dx}{b} + \frac{2\sqrt{d+ex}}{b} \right)}{b} + \frac{2(d+ex)^{3/2}}{3b} \right)}{b} + \frac{2B(d + ex)^{5/2}}{5be}$$

$$\downarrow 73$$

$$\begin{aligned}
 & \frac{(Ab - aB) \left(\frac{(bd - ae) \int \frac{1}{a + \frac{b(d+ex)}{e} - \frac{bd}{e}} d\sqrt{d+ex} + \frac{2\sqrt{d+ex}}{b}}{b} + \frac{2(d+ex)^{3/2}}{3b} \right)}{b} + \frac{2B(d+ex)^{5/2}}{5be} \\
 & \quad \downarrow 221 \\
 & \frac{(Ab - aB) \left(\frac{(bd - ae) \left(\frac{2\sqrt{d+ex}}{b} - \frac{2\sqrt{bd - ae} \operatorname{arctanh} \left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd - ae}} \right)}{b^{3/2}} \right)}{b} + \frac{2(d+ex)^{3/2}}{3b} \right)}{b} + \frac{2B(d+ex)^{5/2}}{5be}
 \end{aligned}$$

input `Int[((A + B*x)*(d + e*x)^(3/2))/(a + b*x), x]`

output `(2*B*(d + e*x)^(5/2))/(5*b*e) + ((A*b - a*B)*((2*(d + e*x)^(3/2))/(3*b) + ((b*d - a*e)*((2*Sqrt[d + e*x])/b - (2*Sqrt[b*d - a*e]*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[b*d - a*e]])/b^(3/2)))/b)/b`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`


```
rule 90 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))),
x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p
+ 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n,
p}, x] && NeQ[n + p + 2, 0]
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Maple [A] (verified)

Time = 0.41 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.06

method	result
pseudoelliptic	$\frac{-2 \left(\left(-\frac{(ex+d)^2 B}{5} - \frac{4A \left(\frac{ex}{4} + d \right) e}{3} \right) b^2 + a \left(\frac{(ex+4d)B}{3} + Ae \right) eb - B a^2 e^2 \right) \sqrt{(ae-db)b} \sqrt{ex+d} + 2e(ae-db)^2 (Ab-Ba) \arctan\left(\frac{x}{\sqrt{(ae-db)b}}\right)}{e b^3 \sqrt{(ae-db)b}}$
risch	$-\frac{2(-3e^2 b^2 B x^2 - 5A b^2 e^2 x + 5B a b e^2 x - 6b^2 B d e x + 15A a b e^2 - 20A b^2 d e - 15B a^2 e^2 + 20B a b d e - 3b^2 B d^2) \sqrt{ex+d}}{15e b^3} + \frac{2 \left(-\frac{b^2 B (ex+d)^{\frac{5}{2}}}{5} - \frac{A b^2 e (ex+d)^{\frac{3}{2}}}{3} + \frac{B a b e (ex+d)^{\frac{3}{2}}}{3} + A a b e^2 \sqrt{ex+d} - A b^2 d e \sqrt{ex+d} - B a^2 e^2 \sqrt{ex+d} + B a b d e \sqrt{ex+d} \right)}{b^3} + \frac{2e(A a^2 - B a b)}{e}$
derivativedivides	$\frac{2 \left(-\frac{b^2 B (ex+d)^{\frac{5}{2}}}{5} - \frac{A b^2 e (ex+d)^{\frac{3}{2}}}{3} + \frac{B a b e (ex+d)^{\frac{3}{2}}}{3} + A a b e^2 \sqrt{ex+d} - A b^2 d e \sqrt{ex+d} - B a^2 e^2 \sqrt{ex+d} + B a b d e \sqrt{ex+d} \right)}{b^3} + \frac{2e(A a^2 - B a b)}{e}$
default	$\frac{2 \left(-\frac{b^2 B (ex+d)^{\frac{5}{2}}}{5} - \frac{A b^2 e (ex+d)^{\frac{3}{2}}}{3} + \frac{B a b e (ex+d)^{\frac{3}{2}}}{3} + A a b e^2 \sqrt{ex+d} - A b^2 d e \sqrt{ex+d} - B a^2 e^2 \sqrt{ex+d} + B a b d e \sqrt{ex+d} \right)}{b^3} + \frac{2e(A a^2 - B a b)}{e}$

```
input int((B*x+A)*(e*x+d)^(3/2)/(b*x+a), x, method=_RETURNVERBOSE)
```

```
output 2*(-((-1/5*(e*x+d)^2*B-4/3*A*(1/4*e*x+d)*e)*b^2+a*(1/3*(e*x+4*d)*B+A*e)*e*
b-B*a^2*e^2)*((a*e-b*d)*b)^(1/2)*(e*x+d)^(1/2)+e*(a*e-b*d)^2*(A*b-B*a)*arc
tan(b*(e*x+d)^(1/2)/((a*e-b*d)*b)^(1/2))/((a*e-b*d)*b)^(1/2)/e/b^3
```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 373, normalized size of antiderivative = 2.87

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{a + bx} dx = \left[\frac{15 ((Bab - Ab^2)de - (Ba^2 - Aab)e^2) \sqrt{\frac{bd-ae}{b}} \log \left(\frac{bex+2bd-ae+2\sqrt{ex+db}\sqrt{\frac{bd-a}{b}}}{bx+a} \right)}{\dots} \right]$$

```
input integrate((B*x+A)*(e*x+d)^(3/2)/(b*x+a),x, algorithm="fricas")
```

```
output [1/15*(15*((B*a*b - A*b^2)*d*e - (B*a^2 - A*a*b)*e^2)*sqrt((b*d - a*e)/b)*
log((b*e*x + 2*b*d - a*e + 2*sqrt(e*x + d)*b*sqrt((b*d - a*e)/b))/(b*x + a
)) + 2*(3*B*b^2*e^2*x^2 + 3*B*b^2*d^2 - 20*(B*a*b - A*b^2)*d*e + 15*(B*a^2
- A*a*b)*e^2 + (6*B*b^2*d*e - 5*(B*a*b - A*b^2)*e^2)*x)*sqrt(e*x + d))/(b
^3*e), 2/15*(15*((B*a*b - A*b^2)*d*e - (B*a^2 - A*a*b)*e^2)*sqrt(-(b*d - a
*e)/b)*arctan(-sqrt(e*x + d)*b*sqrt(-(b*d - a*e)/b)/(b*d - a*e)) + (3*B*b^
2*e^2*x^2 + 3*B*b^2*d^2 - 20*(B*a*b - A*b^2)*d*e + 15*(B*a^2 - A*a*b)*e^2
+ (6*B*b^2*d*e - 5*(B*a*b - A*b^2)*e^2)*x)*sqrt(e*x + d))/(b^3*e)]
```

Sympy [A] (verification not implemented)

Time = 3.77 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.35

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{a + bx} dx = \left\{ \begin{array}{l} 2 \left(\frac{B(d+ex)^{\frac{5}{2}}}{5b} + \frac{(d+ex)^{\frac{3}{2}}(Abe - Bae)}{3b^2} + \frac{\sqrt{d+ex}(-Aabe^2 + Ab^2de + Ba^2e^2 - Babde)}{b^3} \right) - \frac{e(-Ab+Ba)(ae-bd)^2 \operatorname{atan}}{b^4 \sqrt{\frac{ae-bd}{b}}} \\ d^{\frac{3}{2}} \left(\frac{Bx}{b} - \frac{(-Ab+Ba) \left(\begin{array}{l} \frac{x}{a} \text{ for } b = 0 \\ \log \frac{a+bx}{b} \text{ otherwise} \end{array} \right)}{b} \right) \end{array} \right.$$

```
input integrate((B*x+A)*(e*x+d)**(3/2)/(b*x+a),x)
```

output

```
Piecewise((2*(B*(d + e*x)**(5/2)/(5*b) + (d + e*x)**(3/2)*(A*b*e - B*a*e)/
(3*b**2) + sqrt(d + e*x)*(-A*a*b*e**2 + A*b**2*d*e + B*a**2*e**2 - B*a*b*d
*e)/b**3 - e*(-A*b + B*a)*(a*e - b*d)**2*atan(sqrt(d + e*x)/sqrt((a*e - b*
d)/b)))/(b**4*sqrt((a*e - b*d)/b)))/e, Ne(e, 0)), (d**(3/2)*(B*x/b - (-A*b
+ B*a)*Piecewise((x/a, Eq(b, 0)), (log(a + b*x)/b, True))/b), True))
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{a + bx} dx = \text{Exception raised: ValueError}$$

input

```
integrate((B*x+A)*(e*x+d)^(3/2)/(b*x+a),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for m
ore detail
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 226 vs. 2(110) = 220.

Time = 0.13 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.74

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{a + bx} dx =$$

$$\frac{2(Bab^2d^2 - Ab^3d^2 - 2Ba^2bde + 2Aab^2de + Ba^3e^2 - Aa^2be^2) \arctan\left(\frac{\sqrt{ex+db}}{\sqrt{-b^2d+abe}}\right)}{\sqrt{-b^2d+abe}b^3}$$

$$+ \frac{2\left(3(ex+d)^{\frac{5}{2}}Bb^4e^4 - 5(ex+d)^{\frac{3}{2}}Bab^3e^5 + 5(ex+d)^{\frac{3}{2}}Ab^4e^5 - 15\sqrt{ex+d}Bab^3de^5 + 15\sqrt{ex+d}Ab^4de^5\right)}{15b^5e^5}$$

input

```
integrate((B*x+A)*(e*x+d)^(3/2)/(b*x+a),x, algorithm="giac")
```

output

```
-2*(B*a*b^2*d^2 - A*b^3*d^2 - 2*B*a^2*b*d*e + 2*A*a*b^2*d*e + B*a^3*e^2 -
A*a^2*b*e^2)*arctan(sqrt(e*x + d)*b/sqrt(-b^2*d + a*b*e))/(sqrt(-b^2*d + a
*b*e)*b^3) + 2/15*(3*(e*x + d)^(5/2)*B*b^4*e^4 - 5*(e*x + d)^(3/2)*B*a*b^3
*e^5 + 5*(e*x + d)^(3/2)*A*b^4*e^5 - 15*sqrt(e*x + d)*B*a*b^3*d*e^5 + 15*s
qrt(e*x + d)*A*b^4*d*e^5 + 15*sqrt(e*x + d)*B*a^2*b^2*e^6 - 15*sqrt(e*x +
d)*A*a*b^3*e^6)/(b^5*e^5)
```

Mupad [B] (verification not implemented)

Time = 0.99 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.82

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{a + bx} dx = \left(\frac{2Ae - 2Bd}{3be} - \frac{2B(ae^2 - bde)}{3b^2e^2} \right) (d + ex)^{3/2} + \frac{2 \operatorname{atan}\left(\frac{\sqrt{b}(Ab - Ba)(ae - bd)^{3/2} \sqrt{d + ex}}{-Ba^3e^2 + 2Ba^2bde + Aa^2be^2 - Bab^2d^2 - 2Aab^2de + Ab^3d^2} \right) (Ab - Ba)(ae - bd)^{3/2}}{b^{7/2}} + \frac{2B(d + ex)^{5/2}}{5be} - \frac{\left(\frac{2Ae - 2Bd}{be} - \frac{2B(ae^2 - bde)}{b^2e^2} \right) (ae^2 - bde) \sqrt{d + ex}}{be}$$

input

```
int(((A + B*x)*(d + e*x)^(3/2))/(a + b*x), x)
```

output

```
((2*A*e - 2*B*d)/(3*b*e) - (2*B*(a*e^2 - b*d*e))/(3*b^2*e^2))*(d + e*x)^(3
/2) + (2*atan((b^(1/2)*(A*b - B*a)*(a*e - b*d)^(3/2)*(d + e*x)^(1/2))/(A*b
^3*d^2 - B*a^3*e^2 + A*a^2*b*e^2 - B*a*b^2*d^2 - 2*A*a*b^2*d*e + 2*B*a^2*b
*d*e))*(A*b - B*a)*(a*e - b*d)^(3/2))/b^(7/2) + (2*B*(d + e*x)^(5/2))/(5*b
*e) - (((2*A*e - 2*B*d)/(b*e) - (2*B*(a*e^2 - b*d*e))/(b^2*e^2))*(a*e^2 -
b*d*e)*(d + e*x)^(1/2))/(b*e)
```

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.21

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{a + bx} dx = \frac{2\sqrt{ex + d}(e^2x^2 + 2dex + d^2)}{5e}$$

input

```
int((B*x+A)*(e*x+d)^(3/2)/(b*x+a), x)
```

output $(2\sqrt{d + ex}(d^2 + 2dex + e^2x^2))/(5e)$

3.151 $\int \frac{(A+Bx)\sqrt{d+ex}}{a+bx} dx$

Optimal result	1401
Mathematica [A] (verified)	1401
Rubi [A] (verified)	1402
Maple [A] (verified)	1404
Fricas [A] (verification not implemented)	1404
Sympy [A] (verification not implemented)	1405
Maxima [F(-2)]	1406
Giac [A] (verification not implemented)	1406
Mupad [B] (verification not implemented)	1407
Reduce [B] (verification not implemented)	1407

Optimal result

Integrand size = 22, antiderivative size = 98

$$\int \frac{(A+Bx)\sqrt{d+ex}}{a+bx} dx = \frac{2(Ab-aB)\sqrt{d+ex}}{b^2} + \frac{2B(d+ex)^{3/2}}{3be} - \frac{2(Ab-aB)\sqrt{bd-ae}\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{5/2}}$$

output

```
2*(A*b-B*a)*(e*x+d)^(1/2)/b^2+2/3*B*(e*x+d)^(3/2)/b/e-2*(A*b-B*a)*(-a*e+b*d)^(1/2)*arctanh(b^(1/2)*(e*x+d)^(1/2)/(-a*e+b*d)^(1/2))/b^(5/2)
```

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.96

$$\int \frac{(A+Bx)\sqrt{d+ex}}{a+bx} dx = \frac{2\sqrt{d+ex}(3Abe-3aBe+bB(d+ex))}{3b^2e} + \frac{2(-Ab+aB)\sqrt{-bd+ae}\arctan\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right)}{b^{5/2}}$$

input

```
Integrate[((A+B*x)*Sqrt[d+e*x])/(a+b*x),x]
```

output

$$(2*\text{Sqrt}[d + e*x]*(3*A*b*e - 3*a*B*e + b*B*(d + e*x)))/(3*b^2*e) + (2*(-(A*b) + a*B)*\text{Sqrt}[-(b*d) + a*e]*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[d + e*x])/\text{Sqrt}[-(b*d) + a*e]])/b^{(5/2)}$$
Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.97, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {90, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)\sqrt{d + ex}}{a + bx} dx$$

$$\downarrow 90$$

$$\frac{(Ab - aB) \int \frac{\sqrt{d+ex}}{a+bx} dx}{b} + \frac{2B(d + ex)^{3/2}}{3be}$$

$$\downarrow 60$$

$$\frac{(Ab - aB) \left(\frac{(bd - ae) \int \frac{1}{(a+bx)\sqrt{d+ex}} dx}{b} + \frac{2\sqrt{d+ex}}{b} \right)}{b} + \frac{2B(d + ex)^{3/2}}{3be}$$

$$\downarrow 73$$

$$\frac{(Ab - aB) \left(\frac{2(bd - ae) \int \frac{1}{a + \frac{b(d+ex)}{e} - \frac{bd}{e}} d\sqrt{d+ex}}{be} + \frac{2\sqrt{d+ex}}{b} \right)}{b} + \frac{2B(d + ex)^{3/2}}{3be}$$

$$\downarrow 221$$

$$\frac{(Ab - aB) \left(\frac{2\sqrt{d+ex}}{b} - \frac{2\sqrt{bd - ae} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd - ae}}\right)}{b^{3/2}} \right)}{b} + \frac{2B(d + ex)^{3/2}}{3be}$$

input

$$\text{Int}[(A + B*x)*\text{Sqrt}[d + e*x]/(a + b*x), x]$$

output

$$\frac{(2B(d + ex)^{3/2})/(3be) + ((A^2b - a^2B)((2\sqrt{d + ex})/b - (2\sqrt{bd - ae})\operatorname{ArcTanh}[\sqrt{b}\sqrt{d + ex}/\sqrt{bd - ae}])/b^{3/2})/b}{b}$$

Defintions of rubi rules used

rule 60

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(
b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!Integer
Q[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinear
Q[a, b, c, d, m, n, x]
```

rule 73

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

rule 90

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))),
x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p
+ 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n,
p}, x] && NeQ[n + p + 2, 0]
```

rule 221

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```


Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.03

method	result	size
risch	$\frac{2(ebBx+3Abe-3Bae+Bbd)\sqrt{ex+d}}{3eb^2} - \frac{2(Aabe-Ab^2d-BA^2e+Babd) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{b^2\sqrt{(ae-db)b}}$	101
pseudoelliptic	$\frac{-2e(ae-db)(Ab-Ba) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right) + 2\left(\left(\frac{Bx}{3}+A\right)e+\frac{Bd}{3}\right)b-Bae}{eb^2\sqrt{(ae-db)b}} \sqrt{(ae-db)b}\sqrt{ex+d}$	104
derivativedivides	$\frac{2\left(\frac{bB(ex+d)^{\frac{3}{2}}}{3}+Abe\sqrt{ex+d}-Bae\sqrt{ex+d}\right)}{b^2} - \frac{2e(Aabe-Ab^2d-BA^2e+Babd) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{b^2\sqrt{(ae-db)b}}$	111
default	$\frac{2\left(\frac{bB(ex+d)^{\frac{3}{2}}}{3}+Abe\sqrt{ex+d}-Bae\sqrt{ex+d}\right)}{b^2} - \frac{2e(Aabe-Ab^2d-BA^2e+Babd) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{b^2\sqrt{(ae-db)b}}$	111

```
input int((B*x+A)*(e*x+d)^(1/2)/(b*x+a),x,method=_RETURNVERBOSE)
```

```
output 2/3*(B*b*e*x+3*A*b*e-3*B*a*e+B*b*d)*(e*x+d)^(1/2)/e/b^2-2*(A*a*b*e-A*b^2*d-B*a^2*e+B*a*b*d)/b^2/((a*e-b*d)*b)^(1/2)*arctan(b*(e*x+d)^(1/2)/((a*e-b*d)*b)^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 211, normalized size of antiderivative = 2.15

$$\int \frac{(A+Bx)\sqrt{d+ex}}{a+bx} dx = \left[\frac{3(Ba-Ab)e\sqrt{\frac{bd-ae}{b}} \log\left(\frac{bex+2bd-ae-2\sqrt{ex+db}\sqrt{\frac{bd-ae}{b}}}{bx+a}\right) - 2(Bbex+Bbd-3(Ba-Ab)e)\sqrt{ex+d}}{3b^2e}, \dots \right]$$

```
input integrate((B*x+A)*(e*x+d)^(1/2)/(b*x+a),x, algorithm="fricas")
```

output

```
[-1/3*(3*(B*a - A*b)*e*sqrt((b*d - a*e)/b)*log((b*e*x + 2*b*d - a*e - 2*sqrt(e*x + d)*b*sqrt((b*d - a*e)/b))/(b*x + a)) - 2*(B*b*e*x + B*b*d - 3*(B*a - A*b)*e)*sqrt(e*x + d)/(b^2*e), 2/3*(3*(B*a - A*b)*e*sqrt(-(b*d - a*e)/b)*arctan(-sqrt(e*x + d)*b*sqrt(-(b*d - a*e)/b)/(b*d - a*e)) + (B*b*e*x + B*b*d - 3*(B*a - A*b)*e)*sqrt(e*x + d)/(b^2*e)]
```

Sympy [A] (verification not implemented)

Time = 4.58 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.27

$$\int \frac{(A + Bx)\sqrt{d + ex}}{a + bx} dx$$

$$= \begin{cases} \frac{2 \left(\frac{B(d+ex)^{\frac{3}{2}}}{3b} + \frac{\sqrt{d+ex}(Abe - Bae)}{b^2} + \frac{e(-Ab+Ba)(ae-bd) \operatorname{atan}\left(\frac{\sqrt{d+ex}}{\sqrt{\frac{ae-bd}{b}}}\right)}{b^3 \sqrt{\frac{ae-bd}{b}}} \right)}{e} & \text{for } e \neq 0 \\ \sqrt{d} \left(\frac{Bx}{b} - \frac{(-Ab+Ba) \left(\begin{cases} \frac{x}{a} & \text{for } b = 0 \\ \frac{\log(a+bx)}{b} & \text{otherwise} \end{cases} \right)}{b} \right) & \text{otherwise} \end{cases}$$

input

```
integrate((B*x+A)*(e*x+d)**(1/2)/(b*x+a), x)
```

output

```
Piecewise((2*(B*(d + e*x)**(3/2)/(3*b) + sqrt(d + e*x)*(A*b*e - B*a*e)/b**2 + e*(-A*b + B*a)*(a*e - b*d)*atan(sqrt(d + e*x)/sqrt((a*e - b*d)/b))/(b**3*sqrt((a*e - b*d)/b)))/e, Ne(e, 0)), (sqrt(d)*(B*x/b - (-A*b + B*a)*Piecewise((x/a, Eq(b, 0)), (log(a + b*x)/b, True))/b), True))
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + Bx)\sqrt{d + ex}}{a + bx} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)*(e*x+d)^(1/2)/(b*x+a),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.24

$$\int \frac{(A + Bx)\sqrt{d + ex}}{a + bx} dx = -\frac{2(Babd - Ab^2d - Ba^2e + Aabe) \arctan\left(\frac{\sqrt{ex+db}}{\sqrt{-b^2d+abe}}\right)}{\sqrt{-b^2d+abe}b^2} + \frac{2\left((ex+d)^{\frac{3}{2}}Bb^2e^2 - 3\sqrt{ex+d}Babe^3 + 3\sqrt{ex+d}Ab^2e^3\right)}{3b^3e^3}$$

input `integrate((B*x+A)*(e*x+d)^(1/2)/(b*x+a),x, algorithm="giac")`

output `-2*(B*a*b*d - A*b^2*d - B*a^2*e + A*a*b*e)*arctan(sqrt(e*x + d)*b/sqrt(-b^2*d + a*b*e))/(sqrt(-b^2*d + a*b*e)*b^2) + 2/3*((e*x + d)^(3/2)*B*b^2*e^2 - 3*sqrt(e*x + d)*B*a*b*e^3 + 3*sqrt(e*x + d)*A*b^2*e^3)/(b^3*e^3)`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.09

$$\int \frac{(A + Bx)\sqrt{d + ex}}{a + bx} dx = \left(\frac{2Ae - 2Bd}{be} - \frac{2B(ae^2 - bde)}{b^2e^2} \right) \sqrt{d + ex} - \frac{2 \operatorname{atanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right) (Ab - Ba) \sqrt{bd - ae}}{b^{5/2}} + \frac{2B(d + ex)^{3/2}}{3be}$$

input `int(((A + B*x)*(d + e*x)^(1/2))/(a + b*x), x)`output `((2*A*e - 2*B*d)/(b*e) - (2*B*(a*e^2 - b*d*e))/(b^2*e^2))*(d + e*x)^(1/2) - (2*atanh((b^(1/2)*(d + e*x)^(1/2))/(b*d - a*e)^(1/2))*(A*b - B*a)*(b*d - a*e)^(1/2))/b^(5/2) + (2*B*(d + e*x)^(3/2))/(3*b*e)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.16

$$\int \frac{(A + Bx)\sqrt{d + ex}}{a + bx} dx = \frac{2\sqrt{ex + d}(ex + d)}{3e}$$

input `int((B*x+A)*(e*x+d)^(1/2)/(b*x+a), x)`output `(2*sqrt(d + e*x)*(d + e*x))/(3*e)`

$$3.152 \quad \int \frac{A+Bx}{(a+bx)\sqrt{d+ex}} dx$$

Optimal result	1408
Mathematica [A] (verified)	1408
Rubi [A] (verified)	1409
Maple [A] (verified)	1410
Fricas [A] (verification not implemented)	1411
Sympy [A] (verification not implemented)	1411
Maxima [F(-2)]	1412
Giac [A] (verification not implemented)	1412
Mupad [B] (verification not implemented)	1413
Reduce [B] (verification not implemented)	1413

Optimal result

Integrand size = 22, antiderivative size = 74

$$\int \frac{A+Bx}{(a+bx)\sqrt{d+ex}} dx = \frac{2B\sqrt{d+ex}}{be} - \frac{2(Ab-aB)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{3/2}\sqrt{bd-ae}}$$

output $2*B*(e*x+d)^{(1/2)}/b/e-2*(A*b-B*a)*\operatorname{arctanh}(b^{(1/2)}*(e*x+d)^{(1/2)}/(-a*e+b*d)^{(1/2)})/b^{(3/2)}/(-a*e+b*d)^{(1/2)}$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00

$$\int \frac{A+Bx}{(a+bx)\sqrt{d+ex}} dx = \frac{2B\sqrt{d+ex}}{be} + \frac{2(Ab-aB)\operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right)}{b^{3/2}\sqrt{-bd+ae}}$$

input $\operatorname{Integrate}[(A+B*x)/((a+b*x)*\operatorname{Sqrt}[d+e*x]),x]$

output $(2*B*\operatorname{Sqrt}[d+e*x])/(b*e) + (2*(A*b-a*B)*\operatorname{ArcTan}[(\operatorname{Sqrt}[b]*\operatorname{Sqrt}[d+e*x])/(\operatorname{Sqrt}[-(b*d)+a*e])]/(b^{(3/2)}*\operatorname{Sqrt}[-(b*d)+a*e])$

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.136$, Rules used = {90, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + Bx}{(a + bx)\sqrt{d + ex}} dx \\
 & \quad \downarrow 90 \\
 & \frac{(Ab - aB) \int \frac{1}{(a+bx)\sqrt{d+ex}} dx}{b} + \frac{2B\sqrt{d + ex}}{be} \\
 & \quad \downarrow 73 \\
 & \frac{2(Ab - aB) \int \frac{1}{a + \frac{b(d+ex) - bd}{e}} d\sqrt{d + ex}}{be} + \frac{2B\sqrt{d + ex}}{be} \\
 & \quad \downarrow 221 \\
 & \frac{2B\sqrt{d + ex}}{be} - \frac{2(Ab - aB)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{3/2}\sqrt{bd - ae}}
 \end{aligned}$$

input `Int[(A + B*x)/((a + b*x)*Sqrt[d + e*x]),x]`

output `(2*B*Sqrt[d + e*x])/(b*e) - (2*(A*b - a*B)*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[b*d - a*e]])/(b^(3/2)*Sqrt[b*d - a*e])`

Defintions of rubi rules used

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.85

method	result	size
pseudoelliptic	$\frac{2B\sqrt{ex+d} + \frac{2e(Ab-Ba) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{\sqrt{(ae-db)b}}}{be}$	63
risch	$\frac{2B\sqrt{ex+d}}{be} + \frac{2(Ab-Ba) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{b\sqrt{(ae-db)b}}$	65
derivativedivides	$\frac{\frac{2B\sqrt{ex+d}}{b} + \frac{2e(Ab-Ba) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{b\sqrt{(ae-db)b}}}{e}$	66
default	$\frac{2B\sqrt{ex+d}}{b} + \frac{2e(Ab-Ba) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{b\sqrt{(ae-db)b}}$	66

input `int((B*x+A)/(b*x+a)/(e*x+d)^(1/2),x,method=_RETURNVERBOSE)`

output `2/e/b*(B*(e*x+d)^(1/2)+e*(A*b-B*a)/((a*e-b*d)*b)^(1/2)*arctan(b*(e*x+d)^(1/2)/((a*e-b*d)*b)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 209, normalized size of antiderivative = 2.82

$$\int \frac{A + Bx}{(a + bx)\sqrt{d + ex}} dx$$

$$= \left[\frac{\sqrt{b^2d - abe}(Ba - Ab)e \log\left(\frac{bex + 2bd - ae - 2\sqrt{b^2d - abe}\sqrt{ex + d}}{bx + a}\right) - 2(Bb^2d - Babe)\sqrt{ex + d}}{b^3de - ab^2e^2}, \right. \\ \left. \frac{2\left(\sqrt{-b^2d + abe}(Ba - Ab)e \arctan\left(\frac{\sqrt{-b^2d + abe}\sqrt{ex + d}}{bex + bd}\right) - (Bb^2d - Babe)\sqrt{ex + d}\right)}{b^3de - ab^2e^2} \right]$$

input `integrate((B*x+A)/(b*x+a)/(e*x+d)^(1/2),x, algorithm="fricas")`

output `[-(sqrt(b^2*d - a*b*e)*(B*a - A*b)*e*log((b*e*x + 2*b*d - a*e - 2*sqrt(b^2*d - a*b*e)*sqrt(e*x + d))/(b*x + a)) - 2*(B*b^2*d - B*a*b*e)*sqrt(e*x + d))/(b^3*d*e - a*b^2*e^2), -2*(sqrt(-b^2*d + a*b*e)*(B*a - A*b)*e*arctan(sqrt(-b^2*d + a*b*e)*sqrt(e*x + d)/(b*e*x + b*d)) - (B*b^2*d - B*a*b*e)*sqrt(e*x + d))/(b^3*d*e - a*b^2*e^2)]`

Sympy [A] (verification not implemented)

Time = 1.60 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.24

$$\int \frac{A + Bx}{(a + bx)\sqrt{d + ex}} dx = \begin{cases} \frac{2 \left(\frac{B\sqrt{d+ex}}{b} - \frac{e(-Ab+Ba) \operatorname{atan}\left(\frac{\sqrt{d+ex}}{\sqrt{\frac{ae-bd}{b}}}\right)}{b^2\sqrt{\frac{ae-bd}{b}}} \right)}{e} & \text{for } e \neq 0 \\ \frac{Bx}{b} - \frac{(-Ab+Ba) \begin{cases} \frac{x}{a} & \text{for } b = 0 \\ \frac{\log(a+bx)}{b} & \text{otherwise} \end{cases}}{\sqrt{d}} & \text{otherwise} \end{cases}$$

input `integrate((B*x+A)/(b*x+a)/(e*x+d)**(1/2),x)`

output

```
Piecewise((2*(B*sqrt(d + e*x)/b - e*(-A*b + B*a)*atan(sqrt(d + e*x)/sqrt((a*e - b*d)/b)))/(b**2*sqrt((a*e - b*d)/b)))/e, Ne(e, 0)), ((B*x/b - (-A*b + B*a)*Piecewise((x/a, Eq(b, 0)), (log(a + b*x)/b, True))/b)/sqrt(d), True)
)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{(a + bx)\sqrt{d + ex}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((B*x+A)/(b*x+a)/(e*x+d)^(1/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.89

$$\int \frac{A + Bx}{(a + bx)\sqrt{d + ex}} dx = -\frac{2(Ba - Ab) \arctan\left(\frac{\sqrt{ex+db}}{\sqrt{-b^2d+abe}}\right)}{\sqrt{-b^2d+abe}b} + \frac{2\sqrt{ex+d}B}{be}$$

input

```
integrate((B*x+A)/(b*x+a)/(e*x+d)^(1/2),x, algorithm="giac")
```

output

```
-2*(B*a - A*b)*arctan(sqrt(e*x + d)*b/sqrt(-b^2*d + a*b*e))/(sqrt(-b^2*d + a*b*e)*b) + 2*sqrt(e*x + d)*B/(b*e)
```

Mupad [B] (verification not implemented)

Time = 0.95 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.84

$$\int \frac{A + Bx}{(a + bx)\sqrt{d + ex}} dx = \frac{2 \operatorname{atan}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{ae-bd}}\right) (Ab - Ba)}{b^{3/2} \sqrt{ae - bd}} + \frac{2B\sqrt{d+ex}}{be}$$

input `int((A + B*x)/((a + b*x)*(d + e*x)^(1/2)),x)`output `(2*atan((b^(1/2)*(d + e*x)^(1/2))/(a*e - b*d)^(1/2))*(A*b - B*a))/(b^(3/2)*
*(a*e - b*d)^(1/2)) + (2*B*(d + e*x)^(1/2))/(b*e)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.15

$$\int \frac{A + Bx}{(a + bx)\sqrt{d + ex}} dx = \frac{2\sqrt{ex + d}}{e}$$

input `int((B*x+A)/(b*x+a)/(e*x+d)^(1/2),x)`output `(2*sqrt(d + e*x))/e`

3.153 $\int \frac{A+Bx}{(a+bx)(d+ex)^{3/2}} dx$

Optimal result	1414
Mathematica [A] (verified)	1414
Rubi [A] (verified)	1415
Maple [A] (verified)	1416
Fricas [B] (verification not implemented)	1417
Sympy [A] (verification not implemented)	1417
Maxima [F(-2)]	1418
Giac [A] (verification not implemented)	1418
Mupad [B] (verification not implemented)	1419
Reduce [B] (verification not implemented)	1419

Optimal result

Integrand size = 22, antiderivative size = 88

$$\int \frac{A+Bx}{(a+bx)(d+ex)^{3/2}} dx = -\frac{2(Bd-Ae)}{e(bd-ae)\sqrt{d+ex}} - \frac{2(Ab-aB)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{\sqrt{b}(bd-ae)^{3/2}}$$

output

```
(2*A*e-2*B*d)/e/(-a*e+b*d)/(e*x+d)^(1/2)-2*(A*b-B*a)*arctanh(b^(1/2)*(e*x+d)^(1/2)/(-a*e+b*d)^(1/2))/b^(1/2)/(-a*e+b*d)^(3/2)
```

Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00

$$\int \frac{A+Bx}{(a+bx)(d+ex)^{3/2}} dx = \frac{2Bd-2Ae}{e(-bd+ae)\sqrt{d+ex}} - \frac{2(Ab-aB)\arctan\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right)}{\sqrt{b}(-bd+ae)^{3/2}}$$

input

```
Integrate[(A + B*x)/((a + b*x)*(d + e*x)^(3/2)),x]
```

output

```
(2*B*d - 2*A*e)/(e*(-(b*d) + a*e)*Sqrt[d + e*x]) - (2*(A*b - a*B)*ArcTan[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[-(b*d) + a*e]])/(Sqrt[b]*(-(b*d) + a*e)^(3/2))
```

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.136$, Rules used = {87, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{3/2}} dx$$

$$\downarrow 87$$

$$\frac{(Ab - aB) \int \frac{1}{(a+bx)\sqrt{d+ex}} dx}{bd - ae} - \frac{2(Bd - Ae)}{e\sqrt{d + ex}(bd - ae)}$$

$$\downarrow 73$$

$$\frac{2(Ab - aB) \int \frac{1}{a + \frac{b(d+ex)}{e} - \frac{bd}{e}} d\sqrt{d + ex}}{e(bd - ae)} - \frac{2(Bd - Ae)}{e\sqrt{d + ex}(bd - ae)}$$

$$\downarrow 221$$

$$-\frac{2(Ab - aB)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{\sqrt{b}(bd - ae)^{3/2}} - \frac{2(Bd - Ae)}{e\sqrt{d + ex}(bd - ae)}$$

input `Int[(A + B*x)/((a + b*x)*(d + e*x)^(3/2)),x]`

output `(-2*(B*d - A*e))/(e*(b*d - a*e)*Sqrt[d + e*x]) - (2*(A*b - a*B)*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[b*d - a*e]])/(Sqrt[b]*(b*d - a*e)^(3/2))`

Defintions of rubi rules used

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && ( !LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.90

method	result	size
pseudoelliptic	$\frac{-\frac{2(Ae-Bd)}{\sqrt{ex+d}} - \frac{2e(Ab-Ba) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{e(ae-db)}}{e(ae-db)}$	79
derivativedivides	$\frac{-\frac{2(Ae-Bd)}{(ae-db)\sqrt{ex+d}} - \frac{2e(Ab-Ba) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{(ae-db)\sqrt{(ae-db)b}}}{e}$	89
default	$\frac{-\frac{2(Ae-Bd)}{(ae-db)\sqrt{ex+d}} - \frac{2e(Ab-Ba) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{(ae-db)\sqrt{(ae-db)b}}}{e}$	89

```
input int((B*x+A)/(b*x+a)/(e*x+d)^(3/2),x,method=_RETURNVERBOSE)
```

```
output 2/e/(a*e-b*d)*(-(A*e-B*d)/(e*x+d)^(1/2)-e*(A*b-B*a)/((a*e-b*d)*b)^(1/2)*ar
ctan(b*(e*x+d)^(1/2)/((a*e-b*d)*b)^(1/2))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 175 vs. $2(76) = 152$.

Time = 0.14 (sec) , antiderivative size = 363, normalized size of antiderivative = 4.12

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{3/2}} dx = \frac{\left((Ba - Ab)e^2x + (Ba - Ab)de \right) \sqrt{b^2d - abe} \log\left(\frac{bex + 2bd - ae + 2\sqrt{b^2d - abe}\sqrt{ex + d}}{bx + a} \right) + 2 \left((Ba - Ab)e^2x + (Ba - Ab)de \right) \sqrt{-b^2d + abe} \arctan\left(\frac{\sqrt{-b^2d + abe}\sqrt{ex + d}}{bex + bd} \right) + (Bb^2d^2 + Aabe^2 - (Bab + A^2)) \sqrt{ex + d}}{b^3d^3e - 2ab^2d^2e^2 + a^2bde^3 + (b^3d^2e^2 - 2ab^2de^3 + a^2be^4)x}$$

input `integrate((B*x+A)/(b*x+a)/(e*x+d)^(3/2),x, algorithm="fricas")`

output `[(((B*a - A*b)*e^2*x + (B*a - A*b)*d*e)*sqrt(b^2*d - a*b*e)*log((b*e*x + 2*b*d - a*e + 2*sqrt(b^2*d - a*b*e)*sqrt(e*x + d))/(b*x + a)) - 2*(B*b^2*d^2 + A*a*b*e^2 - (B*a*b + A*b^2)*d*e)*sqrt(e*x + d)/(b^3*d^3*e - 2*a*b^2*d^2*e^2 + a^2*b*d*e^3 + (b^3*d^2*e^2 - 2*a*b^2*d*e^3 + a^2*b*e^4)*x), -2*((B*a - A*b)*e^2*x + (B*a - A*b)*d*e)*sqrt(-b^2*d + a*b*e)*arctan(sqrt(-b^2*d + a*b*e)*sqrt(e*x + d)/(b*e*x + b*d)) + (B*b^2*d^2 + A*a*b*e^2 - (B*a*b + A*b^2)*d*e)*sqrt(e*x + d)/(b^3*d^3*e - 2*a*b^2*d^2*e^2 + a^2*b*d*e^3 + (b^3*d^2*e^2 - 2*a*b^2*d*e^3 + a^2*b*e^4)*x)]`

Sympy [A] (verification not implemented)

Time = 3.71 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.24

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{3/2}} dx = \begin{cases} 2 \left(\frac{-Ae + Bd}{\sqrt{d + ex}(ae - bd)} + \frac{e(-Ab + Ba) \operatorname{atan}\left(\frac{\sqrt{d + ex}}{\sqrt{\frac{ae - bd}{b}}}\right)}{b\sqrt{\frac{ae - bd}{b}}(ae - bd)} \right) & \text{for } e \neq 0 \\ \frac{Bx}{b} - \frac{(-Ab + Ba) \left(\begin{cases} \frac{x}{a} & \text{for } b = 0 \\ \frac{\log(a + bx)}{b} & \text{otherwise} \end{cases} \right)}{d^{3/2}} & \text{otherwise} \end{cases}$$

input `integrate((B*x+A)/(b*x+a)/(e*x+d)**(3/2),x)`

output

```
Piecewise((2*((-A*e + B*d)/(sqrt(d + e*x)*(a*e - b*d)) + e*(-A*b + B*a)*at
an(sqrt(d + e*x)/sqrt((a*e - b*d)/b))/(b*sqrt((a*e - b*d)/b)*(a*e - b*d)))
/e, Ne(e, 0)), ((B*x/b - (-A*b + B*a)*Piecewise((x/a, Eq(b, 0)), (log(a +
b*x)/b, True))/b)/d**(3/2), True))
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{3/2}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((B*x+A)/(b*x+a)/(e*x+d)^(3/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for m
ore detail
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.99

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{3/2}} dx = -\frac{2(Ba - Ab) \arctan\left(\frac{\sqrt{ex+db}}{\sqrt{-b^2d+abe}}\right)}{\sqrt{-b^2d+abe}(bd - ae)} - \frac{2(Bd - Ae)}{(bde - ae^2)\sqrt{ex + d}}$$

input

```
integrate((B*x+A)/(b*x+a)/(e*x+d)^(3/2),x, algorithm="giac")
```

output

```
-2*(B*a - A*b)*arctan(sqrt(e*x + d)*b/sqrt(-b^2*d + a*b*e))/(sqrt(-b^2*d +
a*b*e)*(b*d - a*e)) - 2*(B*d - A*e)/((b*d*e - a*e^2)*sqrt(e*x + d))
```

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.09

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{3/2}} dx =$$

$$-\frac{2 \operatorname{atan}\left(\frac{2\sqrt{b}(Ab - Ba)\sqrt{d+ex}}{(2Ab - 2Ba)\sqrt{ae - bd}}\right) (Ab - Ba)}{\sqrt{b}(ae - bd)^{3/2}} - \frac{2(Ae - Bd)}{e(ae - bd)\sqrt{d+ex}}$$

input `int((A + B*x)/((a + b*x)*(d + e*x)^(3/2)),x)`output `-(2*atan((2*b^(1/2)*(A*b - B*a)*(d + e*x)^(1/2))/((2*A*b - 2*B*a)*(a*e - b*d)^(1/2)))*(A*b - B*a))/(b^(1/2)*(a*e - b*d)^(3/2)) - (2*(A*e - B*d))/(e*(a*e - b*d)*(d + e*x)^(1/2))`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.15

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{3/2}} dx = -\frac{2}{\sqrt{ex + d}e}$$

input `int((B*x+A)/(b*x+a)/(e*x+d)^(3/2),x)`output `(- 2)/(sqrt(d + e*x)*e)`

3.154 $\int \frac{A+Bx}{(a+bx)(d+ex)^{5/2}} dx$

Optimal result	1420
Mathematica [A] (verified)	1420
Rubi [A] (verified)	1421
Maple [A] (verified)	1423
Fricas [B] (verification not implemented)	1423
Sympy [A] (verification not implemented)	1424
Maxima [F(-2)]	1425
Giac [A] (verification not implemented)	1425
Mupad [B] (verification not implemented)	1426
Reduce [B] (verification not implemented)	1426

Optimal result

Integrand size = 22, antiderivative size = 119

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{5/2}} dx = -\frac{2(Bd - Ae)}{3e(bd - ae)(d + ex)^{3/2}} + \frac{2(Ab - aB)}{(bd - ae)^2 \sqrt{d + ex}} - \frac{2\sqrt{b}(Ab - aB) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{(bd - ae)^{5/2}}$$

output $\frac{1}{3} \cdot \frac{(2Ae - 2Bd)}{e(-ae + bd)} \cdot \frac{1}{(ex + d)^{3/2}} + \frac{2(Ab - Ba)}{(-ae + bd)^2} \cdot \frac{1}{(ex + d)^{1/2}} - \frac{2b^{1/2}(Ab - Ba) \operatorname{arctanh}(b^{1/2}(ex + d)^{1/2} / (-ae + bd)^{1/2})}{(-ae + bd)^{5/2}}$

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.99

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{5/2}} dx = -\frac{2(bBd^2 - Abe(4d + 3ex) + ae(2Bd + Ae + 3Bex))}{3e(bd - ae)^2(d + ex)^{3/2}} + \frac{2\sqrt{b}(Ab - aB) \arctan\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right)}{(-bd + ae)^{5/2}}$$

input `Integrate[(A + B*x)/((a + b*x)*(d + e*x)^(5/2)),x]`

output

$$\frac{(-2*(b*B*d^2 - A*b*e*(4*d + 3*e*x) + a*e*(2*B*d + A*e + 3*B*e*x))/(3*e*(b*d - a*e)^2*(d + e*x)^{(3/2)}) + (2*sqrt[b]*(A*b - a*B)*ArcTan[(sqrt[b]*sqrt[d + e*x])/sqrt[-(b*d) + a*e]])/(-(b*d) + a*e)^{(5/2)}}{}$$
Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.03, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {87, 61, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{5/2}} dx$$

$$\downarrow 87$$

$$\frac{(Ab - aB) \int \frac{1}{(a+bx)(d+ex)^{3/2}} dx}{bd - ae} - \frac{2(Bd - Ae)}{3e(d + ex)^{3/2}(bd - ae)}$$

$$\downarrow 61$$

$$\frac{(Ab - aB) \left(\frac{b \int \frac{1}{(a+bx)\sqrt{d+ex}} dx}{bd - ae} + \frac{2}{\sqrt{d+ex}(bd - ae)} \right)}{bd - ae} - \frac{2(Bd - Ae)}{3e(d + ex)^{3/2}(bd - ae)}$$

$$\downarrow 73$$

$$\frac{(Ab - aB) \left(\frac{2b \int \frac{1}{a + \frac{b(d+ex) - bd}{e}} d\sqrt{d+ex}}{e(bd - ae)} + \frac{2}{\sqrt{d+ex}(bd - ae)} \right)}{bd - ae} - \frac{2(Bd - Ae)}{3e(d + ex)^{3/2}(bd - ae)}$$

$$\downarrow 221$$

$$\frac{(Ab - aB) \left(\frac{2}{\sqrt{d+ex}(bd - ae)} - \frac{2\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd - ae}}\right)}{(bd - ae)^{3/2}} \right)}{bd - ae} - \frac{2(Bd - Ae)}{3e(d + ex)^{3/2}(bd - ae)}$$

input

$$\text{Int}[(A + B*x)/((a + b*x)*(d + e*x)^{(5/2)}), x]$$

output

$$\frac{(-2*(B*d - A*e))/(3*e*(b*d - a*e)*(d + e*x)^{(3/2)}) + ((A*b - a*B)*(2/((b*d - a*e)*\text{Sqrt}[d + e*x]) - (2*\text{Sqrt}[b]*\text{ArcTanh}[(\text{Sqrt}[b]*\text{Sqrt}[d + e*x])/\text{Sqrt}[b*d - a*e]]))/(b*d - a*e)^{(3/2))}}{(b*d - a*e)}$$

Defintions of rubi rules used

rule 61

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((
m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x],
x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0]
) || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n])) && IntLinearQ[a, b, c, d
, m, n, x]
```

rule 73

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

rule 87

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p
_), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && ( !LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

rule 221

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```


output

```
[-1/3*(3*((B*a - A*b)*e^3*x^2 + 2*(B*a - A*b)*d*e^2*x + (B*a - A*b)*d^2*e)*sqrt(b/(b*d - a*e))*log((b*e*x + 2*b*d - a*e - 2*(b*d - a*e)*sqrt(e*x + d))*sqrt(b/(b*d - a*e)))/(b*x + a) + 2*(B*b*d^2 + A*a*e^2 + 3*(B*a - A*b)*e^2*x + 2*(B*a - 2*A*b)*d*e)*sqrt(e*x + d)/(b^2*d^4*e - 2*a*b*d^3*e^2 + a^2*d^2*e^3 + (b^2*d^2*e^3 - 2*a*b*d*e^4 + a^2*e^5)*x^2 + 2*(b^2*d^3*e^2 - 2*a*b*d^2*e^3 + a^2*d*e^4)*x), -2/3*(3*((B*a - A*b)*e^3*x^2 + 2*(B*a - A*b)*d*e^2*x + (B*a - A*b)*d^2*e)*sqrt(-b/(b*d - a*e))*arctan(sqrt(e*x + d)*sqrt(-b/(b*d - a*e))) + (B*b*d^2 + A*a*e^2 + 3*(B*a - A*b)*e^2*x + 2*(B*a - 2*A*b)*d*e)*sqrt(e*x + d)/(b^2*d^4*e - 2*a*b*d^3*e^2 + a^2*d^2*e^3 + (b^2*d^2*e^3 - 2*a*b*d*e^4 + a^2*e^5)*x^2 + 2*(b^2*d^3*e^2 - 2*a*b*d^2*e^3 + a^2*d*e^4)*x)]
```

Sympy [A] (verification not implemented)

Time = 4.67 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.16

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{5/2}} dx = \begin{cases} \frac{2 \left(-\frac{e(-Ab+Ba)}{\sqrt{d+ex}(ae-bd)^2} - \frac{e(-Ab+Ba) \operatorname{atan}\left(\frac{\sqrt{d+ex}}{\sqrt{\frac{ae-bd}{b}}}\right)}{\sqrt{\frac{ae-bd}{b}}(ae-bd)^2} + \frac{-Ae+Bd}{3(d+ex)^{\frac{3}{2}}(ae-bd)} \right)}{e} & \text{for } e \neq 0 \\ \frac{\frac{Bx}{b} - \frac{(-Ab+Ba) \left(\begin{cases} \frac{x}{a} & \text{for } b = 0 \\ \frac{\log(a+bx)}{b} & \text{otherwise} \end{cases} \right)}{d^{\frac{5}{2}}}}{d^{\frac{5}{2}}} & \text{otherwise} \end{cases}$$

input

```
integrate((B*x+A)/(b*x+a)/(e*x+d)**(5/2),x)
```

output

```
Piecewise((2*(-e*(-A*b + B*a))/(sqrt(d + e*x)*(a*e - b*d)**2) - e*(-A*b + B*a)*atan(sqrt(d + e*x)/sqrt((a*e - b*d)/b))/(sqrt((a*e - b*d)/b)*(a*e - b*d)**2) + (-A*e + B*d)/(3*(d + e*x)**(3/2)*(a*e - b*d)))/e, Ne(e, 0)), ((B*x/b - (-A*b + B*a)*Piecewise((x/a, Eq(b, 0)), (log(a + b*x)/b, True))/b)/d** (5/2), True))
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)/(e*x+d)^(5/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail)

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.29

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{5/2}} dx = -\frac{2(Bab - Ab^2) \arctan\left(\frac{\sqrt{ex+db}}{\sqrt{-b^2d+abe}}\right) - \frac{2(Bbd^2 + 3(ex+d)Bae - 3(ex+d)Abe - Bade - Abde + Aae^2)}{3(b^2d^2e - 2abde^2 + a^2e^3)(ex+d)^{3/2}}}{(b^2d^2 - 2abde + a^2e^2)\sqrt{-b^2d+abe}}$$

input `integrate((B*x+A)/(b*x+a)/(e*x+d)^(5/2),x, algorithm="giac")`

output `-2*(B*a*b - A*b^2)*arctan(sqrt(e*x + d)*b/sqrt(-b^2*d + a*b*e))/((b^2*d^2 - 2*a*b*d*e + a^2*e^2)*sqrt(-b^2*d + a*b*e)) - 2/3*(B*b*d^2 + 3*(e*x + d)*B*a*e - 3*(e*x + d)*A*b*e - B*a*d*e - A*b*d*e + A*a*e^2)/((b^2*d^2*e - 2*a*b*d*e^2 + a^2*e^3)*(e*x + d)^(3/2))`

Mupad [B] (verification not implemented)

Time = 1.02 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.08

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{5/2}} dx = \frac{2\sqrt{b} \operatorname{atan}\left(\frac{\sqrt{b}\sqrt{d+ex}(a^2e^2 - 2abde + b^2d^2)}{(ae-bd)^{5/2}}\right) (Ab - Ba)}{(ae - bd)^{5/2}} - \frac{\frac{2(Ae-Bd)}{3(ae-bd)} - \frac{2(Abe-Bae)(d+ex)}{(ae-bd)^2}}{e(d+ex)^{3/2}}$$

input `int((A + B*x)/((a + b*x)*(d + e*x)^(5/2)),x)`output `(2*b^(1/2)*atan((b^(1/2)*(d + e*x)^(1/2)*(a^2*e^2 + b^2*d^2 - 2*a*b*d*e))/
(a*e - b*d)^(5/2))*(A*b - B*a))/(a*e - b*d)^(5/2) - ((2*(A*e - B*d))/(3*(a
*e - b*d)) - (2*(A*b*e - B*a*e)*(d + e*x))/(a*e - b*d)^2)/(e*(d + e*x)^(3/
2))`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.17

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{5/2}} dx = -\frac{2}{3\sqrt{ex + d}e(ex + d)}$$

input `int((B*x+A)/(b*x+a)/(e*x+d)^(5/2),x)`output `(- 2)/(3*sqrt(d + e*x)*e*(d + e*x))`

3.155 $\int \frac{A+Bx}{(a+bx)(d+ex)^{7/2}} dx$

Optimal result	1427
Mathematica [A] (verified)	1427
Rubi [A] (verified)	1428
Maple [A] (verified)	1430
Fricas [B] (verification not implemented)	1431
Sympy [A] (verification not implemented)	1432
Maxima [F(-2)]	1432
Giac [B] (verification not implemented)	1433
Mupad [B] (verification not implemented)	1433
Reduce [B] (verification not implemented)	1434

Optimal result

Integrand size = 22, antiderivative size = 151

$$\int \frac{A+Bx}{(a+bx)(d+ex)^{7/2}} dx = -\frac{2(Bd - Ae)}{5e(bd - ae)(d+ex)^{5/2}} + \frac{2(Ab - aB)}{3(bd - ae)^2(d+ex)^{3/2}} + \frac{2b(Ab - aB)}{(bd - ae)^3\sqrt{d+ex}} - \frac{2b^{3/2}(Ab - aB)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{(bd - ae)^{7/2}}$$

output

```
1/5*(2*A*e-2*B*d)/e/(-a*e+b*d)/(e*x+d)^(5/2)+2/3*(A*b-B*a)/(-a*e+b*d)^2/(e*x+d)^(3/2)+2*b*(A*b-B*a)/(-a*e+b*d)^3/(e*x+d)^(1/2)-2*b^(3/2)*(A*b-B*a)*arctanh(b^(1/2)*(e*x+d)^(1/2)/(-a*e+b*d)^(1/2))/(-a*e+b*d)^(7/2)
```

Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.17

$$\int \frac{A+Bx}{(a+bx)(d+ex)^{7/2}} dx = \frac{2(-a^2e^2(2Bd + 3Ae + 5Bex) + abe(Ae(11d + 5ex) + B(14d^2 + 35dex + 15e^2d)))}{15e(-bd + ae)^3(d+ex)^{5/2}} - \frac{2b^{3/2}(Ab - aB)\arctan\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right)}{(-bd + ae)^{7/2}}$$

input

```
Integrate[(A + B*x)/((a + b*x)*(d + e*x)^(7/2)),x]
```


output

$$\frac{(2*(-(a^2e^2(2Bd + 3Ae + 5Bex)) + a*be*(Ae*(11d + 5ex) + B*(14d^2 + 35d*ex + 15e^2x^2)) + b^2*(3Bd^3 - Ae*(23d^2 + 35d*ex + 15e^2x^2))) / (15e*(-(b*d) + a*e)^3*(d + ex)^{(5/2)}) - (2*b^{(3/2)}*(A*b - a*B)*ArcTan[(\text{Sqrt}[b]*\text{Sqrt}[d + ex])/\text{Sqrt}[-(b*d) + a*e]]) / (-(b*d) + a*e)^{(7/2)}}{1}$$
Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.05, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.227$, Rules used = {87, 61, 61, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{7/2}} dx$$

$$\downarrow 87$$

$$\frac{(Ab - aB) \int \frac{1}{(a+bx)(d+ex)^{5/2}} dx}{bd - ae} - \frac{2(Bd - Ae)}{5e(d + ex)^{5/2}(bd - ae)}$$

$$\downarrow 61$$

$$\frac{(Ab - aB) \left(\frac{b \int \frac{1}{(a+bx)(d+ex)^{3/2}} dx}{bd - ae} + \frac{2}{3(d+ex)^{3/2}(bd - ae)} \right)}{bd - ae} - \frac{2(Bd - Ae)}{5e(d + ex)^{5/2}(bd - ae)}$$

$$\downarrow 61$$

$$\frac{(Ab - aB) \left(\frac{b \left(\frac{b \int \frac{1}{(a+bx)\sqrt{d+ex}} dx}{bd - ae} + \frac{2}{\sqrt{d+ex}(bd - ae)} \right)}{bd - ae} + \frac{2}{3(d+ex)^{3/2}(bd - ae)} \right)}{bd - ae} - \frac{2(Bd - Ae)}{5e(d + ex)^{5/2}(bd - ae)}$$

$$\downarrow 73$$

$$\begin{aligned}
 & \frac{(Ab - aB) \left(\frac{b \left(\frac{2b \int \frac{1}{a + \frac{b(d+ex) - bd}{e(bd-ae)}} d\sqrt{d+ex}}{e(bd-ae)} + \frac{2}{\sqrt{d+ex}(bd-ae)} \right)}{bd-ae} + \frac{2}{3(d+ex)^{3/2}(bd-ae)} \right)}{\frac{bd - ae}{2(Bd - Ae)}} \\
 & \qquad \qquad \qquad \frac{bd - ae}{5e(d + ex)^{5/2}(bd - ae)} \\
 & \qquad \qquad \qquad \downarrow \text{221} \\
 & \frac{(Ab - aB) \left(\frac{b \left(\frac{2}{\sqrt{d+ex}(bd-ae)} - \frac{2\sqrt{b}\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{(bd-ae)^{3/2}} \right)}{bd-ae} + \frac{2}{3(d+ex)^{3/2}(bd-ae)} \right)}{\frac{bd - ae}{2(Bd - Ae)}} \\
 & \qquad \qquad \qquad \frac{bd - ae}{5e(d + ex)^{5/2}(bd - ae)}
 \end{aligned}$$

input `Int[(A + B*x)/((a + b*x)*(d + e*x)^(7/2)),x]`

output `(-2*(B*d - A*e))/(5*e*(b*d - a*e)*(d + e*x)^(5/2)) + ((A*b - a*B)*(2/(3*(b*d - a*e)*(d + e*x)^(3/2)) + (b*(2/((b*d - a*e)*Sqrt[d + e*x]) - (2*Sqrt[b]*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[b*d - a*e]])/(b*d - a*e)^(3/2)))/(b*d - a*e)))/(b*d - a*e)`

Defintions of rubi rules used

rule 61 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p
), x] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
 + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
 + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
 /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
 rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
 /Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.99

method	result
derivativedivides	$-\frac{2b^2e(Ab-Ba) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{(ae-db)^3\sqrt{(ae-db)b}} - \frac{2(Ae-Bd)}{5(ae-db)(ex+d)^{\frac{5}{2}}} - \frac{2e(Ab-Ba)b}{(ae-db)^3\sqrt{ex+d}} + \frac{2e(Ab-Ba)}{3(ae-db)^2(ex+d)^{\frac{3}{2}}}$
default	$-\frac{2b^2e(Ab-Ba) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{(ae-db)^3\sqrt{(ae-db)b}} - \frac{2(Ae-Bd)}{5(ae-db)(ex+d)^{\frac{5}{2}}} - \frac{2e(Ab-Ba)b}{(ae-db)^3\sqrt{ex+d}} + \frac{2e(Ab-Ba)}{3(ae-db)^2(ex+d)^{\frac{3}{2}}}$
pseudoelliptic	$-\frac{2\left(5b^2e(ex+d)^{\frac{5}{2}}(Ab-Ba) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right) + \sqrt{(ae-db)b}\left(\left(5Ab^2x^2 - \frac{5ax(3Bx+A)b}{3} + a^2\left(\frac{5Bx}{3} + A\right)\right)e^3 - \frac{11}{3}\right)\right)}{5(ex+d)^{\frac{5}{2}}\sqrt{(ae-db)b}e(ae-db)^3}$

input `int((B*x+A)/(b*x+a)/(e*x+d)^(7/2),x,method=_RETURNVERBOSE)`

output

```
2/e*(-1/5*(A*e-B*d)/(a*e-b*d)/(e*x+d)^(5/2)-e*(A*b-B*a)/(a*e-b*d)^3*b/(e*x+d)^(1/2)+1/3*e*(A*b-B*a)/(a*e-b*d)^2/(e*x+d)^(3/2)-b^2*e*(A*b-B*a)/(a*e-b*d)^3/((a*e-b*d)*b)^(1/2)*arctan(b*(e*x+d)^(1/2)/((a*e-b*d)*b)^(1/2)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 426 vs. $2(131) = 262$.

Time = 0.14 (sec) , antiderivative size = 882, normalized size of antiderivative = 5.84

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{7/2}} dx = \text{Too large to display}$$

input

```
integrate((B*x+A)/(b*x+a)/(e*x+d)^(7/2),x, algorithm="fricas")
```

output

```
[1/15*(15*((B*a*b - A*b^2)*e^4*x^3 + 3*(B*a*b - A*b^2)*d*e^3*x^2 + 3*(B*a*b - A*b^2)*d^2*e^2*x + (B*a*b - A*b^2)*d^3*e)*sqrt(b/(b*d - a*e))*log((b*e*x + 2*b*d - a*e + 2*(b*d - a*e)*sqrt(e*x + d)*sqrt(b/(b*d - a*e)))/(b*x + a)) - 2*(3*B*b^2*d^3 - 3*A*a^2*e^3 + 15*(B*a*b - A*b^2)*e^3*x^2 + (14*B*a*b - 23*A*b^2)*d^2*e - (2*B*a^2 - 11*A*a*b)*d*e^2 + 5*(7*(B*a*b - A*b^2)*d*e^2 - (B*a^2 - A*a*b)*e^3)*x)*sqrt(e*x + d))/(b^3*d^6*e - 3*a*b^2*d^5*e^2 + 3*a^2*b*d^4*e^3 - a^3*d^3*e^4 + (b^3*d^3*e^4 - 3*a*b^2*d^2*e^5 + 3*a^2*b*d*e^6 - a^3*e^7)*x^3 + 3*(b^3*d^4*e^3 - 3*a*b^2*d^3*e^4 + 3*a^2*b*d^2*e^5 - a^3*d*e^6)*x^2 + 3*(b^3*d^5*e^2 - 3*a*b^2*d^4*e^3 + 3*a^2*b*d^3*e^4 - a^3*d^2*e^5)*x), -2/15*(15*((B*a*b - A*b^2)*e^4*x^3 + 3*(B*a*b - A*b^2)*d*e^3*x^2 + 3*(B*a*b - A*b^2)*d^2*e^2*x + (B*a*b - A*b^2)*d^3*e)*sqrt(-b/(b*d - a*e))*arctan(sqrt(e*x + d)*sqrt(-b/(b*d - a*e))) + (3*B*b^2*d^3 - 3*A*a^2*e^3 + 15*(B*a*b - A*b^2)*e^3*x^2 + (14*B*a*b - 23*A*b^2)*d^2*e - (2*B*a^2 - 11*A*a*b)*d*e^2 + 5*(7*(B*a*b - A*b^2)*d*e^2 - (B*a^2 - A*a*b)*e^3)*x)*sqrt(e*x + d))/(b^3*d^6*e - 3*a*b^2*d^5*e^2 + 3*a^2*b*d^4*e^3 - a^3*d^3*e^4 + (b^3*d^3*e^4 - 3*a*b^2*d^2*e^5 + 3*a^2*b*d*e^6 - a^3*e^7)*x^3 + 3*(b^3*d^4*e^3 - 3*a*b^2*d^3*e^4 + 3*a^2*b*d^2*e^5 - a^3*d*e^6)*x^2 + 3*(b^3*d^5*e^2 - 3*a*b^2*d^4*e^3 + 3*a^2*b*d^3*e^4 - a^3*d^2*e^5)*x)]
```

Sympy [A] (verification not implemented)

Time = 5.07 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.13

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{7/2}} dx = \begin{cases} 2 \left(\frac{be(-Ab+Ba)}{\sqrt{d+ex}(ae-bd)^3} + \frac{be(-Ab+Ba) \operatorname{atan}\left(\frac{\sqrt{d+ex}}{\sqrt{\frac{ae-bd}{b}}}\right)}{\sqrt{\frac{ae-bd}{b}}(ae-bd)^3} - \frac{e(-Ab+Ba)}{3(d+ex)^{3/2}(ae-bd)^2} + \frac{-Ae+Bd}{5(d+ex)^{5/2}(ae-bd)} \right) & \text{for } e \neq 0 \\ \frac{Bx}{b} - \frac{(-Ab+Ba) \begin{cases} \frac{x}{a} & \text{for } b = 0 \\ \frac{\log(a+bx)}{b} & \text{otherwise} \end{cases}}{d^{7/2}} & \text{otherwise} \end{cases}$$

input `integrate((B*x+A)/(b*x+a)/(e*x+d)**(7/2),x)`output `Piecewise((2*(b*e*(-A*b + B*a))/(sqrt(d + e*x)*(a*e - b*d)**3) + b*e*(-A*b + B*a)*atan(sqrt(d + e*x)/sqrt((a*e - b*d)/b))/(sqrt((a*e - b*d)/b)*(a*e - b*d)**3) - e*(-A*b + B*a)/(3*(d + e*x)**(3/2)*(a*e - b*d)**2) + (-A*e + B*d)/(5*(d + e*x)**(5/2)*(a*e - b*d)))/e, Ne(e, 0)), ((B*x/b - (-A*b + B*a)*Piecewise((x/a, Eq(b, 0)), (log(a + b*x)/b, True))/b)/d**(7/2), True))`**Maxima [F(-2)]**

Exception generated.

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{7/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)/(e*x+d)^(7/2),x, algorithm="maxima")`output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 276 vs. $2(131) = 262$.

Time = 0.12 (sec) , antiderivative size = 276, normalized size of antiderivative = 1.83

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{7/2}} dx = -\frac{2(Bab^2 - Ab^3) \arctan\left(\frac{\sqrt{ex+db}}{\sqrt{-b^2d+abe}}\right)}{(b^3d^3 - 3ab^2d^2e + 3a^2bde^2 - a^3e^3)\sqrt{-b^2d+abe}} - \frac{2(3Bb^2d^3 + 15(ex+d)^2Babe - 15(ex+d)^2Ab^2e + 5(ex+d)Babde - 5(ex+d)Ab^2de - 6Babd^2e - 15(b^3d^3e - 3ab^2d^2e^2 + 3a^2bde^3 - a^3e^4))}{(b^3d^3e - 3ab^2d^2e^2 + 3a^2bde^3 - a^3e^4)}$$

input `integrate((B*x+A)/(b*x+a)/(e*x+d)^(7/2),x, algorithm="giac")`

output `-2*(B*a*b^2 - A*b^3)*arctan(sqrt(e*x + d)*b/sqrt(-b^2*d + a*b*e))/((b^3*d^3 - 3*a*b^2*d^2*e + 3*a^2*b*d*e^2 - a^3*e^3)*sqrt(-b^2*d + a*b*e)) - 2/15*(3*B*b^2*d^3 + 15*(e*x + d)^2*B*a*b*e - 15*(e*x + d)^2*A*b^2*e + 5*(e*x + d)*B*a*b*d*e - 5*(e*x + d)*A*b^2*d*e - 6*B*a*b*d^2*e - 3*A*b^2*d^2*e - 5*(e*x + d)*B*a^2*e^2 + 5*(e*x + d)*A*a*b*e^2 + 3*B*a^2*d*e^2 + 6*A*a*b*d*e^2 - 3*A*a^2*e^3)/((b^3*d^3*e - 3*a*b^2*d^2*e^2 + 3*a^2*b*d*e^3 - a^3*e^4)*(e*x + d)^(5/2))`

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.15

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{7/2}} dx = -\frac{\frac{2(Ae-Bd)}{5(ae-bd)} - \frac{2(Abe-Bae)(d+ex)}{3(ae-bd)^2} + \frac{2b(Abe-Bae)(d+ex)^2}{(ae-bd)^3}}{e(d+ex)^{5/2}} - \frac{2b^{3/2} \operatorname{atan}\left(\frac{\sqrt{b}\sqrt{d+ex}(a^3e^3-3a^2bde^2+3ab^2d^2e-b^3d^3)}{(ae-bd)^{7/2}}\right)}{(ae-bd)^{7/2}}(Ab-Ba)$$

input `int((A + B*x)/((a + b*x)*(d + e*x)^(7/2)),x)`

output

```
- ((2*(A*e - B*d))/(5*(a*e - b*d)) - (2*(A*b*e - B*a*e)*(d + e*x))/(3*(a*e
- b*d)^2) + (2*b*(A*b*e - B*a*e)*(d + e*x)^2)/(a*e - b*d)^3)/(e*(d + e*x)
^(5/2)) - (2*b^(3/2)*atan((b^(1/2)*(d + e*x)^(1/2)*(a^3*e^3 - b^3*d^3 + 3*
a*b^2*d^2*e - 3*a^2*b*d*e^2))/(a*e - b*d)^(7/2))*(A*b - B*a))/(a*e - b*d)^(
(7/2))
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.21

$$\int \frac{A + Bx}{(a + bx)(d + ex)^{7/2}} dx = -\frac{2}{5\sqrt{ex + d}e(e^2x^2 + 2dex + d^2)}$$

input

```
int((B*x+A)/(b*x+a)/(e*x+d)^(7/2),x)
```

output

```
( - 2)/(5*sqrt(d + e*x)*e*(d**2 + 2*d*e*x + e**2*x**2))
```

3.156 $\int \frac{(A+Bx)(d+ex)^{7/2}}{(a+bx)^2} dx$

Optimal result	1435
Mathematica [A] (verified)	1436
Rubi [A] (verified)	1436
Maple [A] (verified)	1440
Fricas [B] (verification not implemented)	1441
Sympy [F(-1)]	1442
Maxima [F(-2)]	1443
Giac [B] (verification not implemented)	1443
Mupad [B] (verification not implemented)	1444
Reduce [B] (verification not implemented)	1445

Optimal result

Integrand size = 22, antiderivative size = 228

$$\int \frac{(A+Bx)(d+ex)^{7/2}}{(a+bx)^2} dx = \frac{2(bd-ae)^2(bBd+3Abe-4aBe)\sqrt{d+ex}}{b^5} - \frac{(Ab-aB)(bd-ae)^3\sqrt{d+ex}}{b^5(a+bx)} + \frac{2(bd-ae)(bBd+2Abe-3aBe)(d+ex)^{3/2}}{3b^4} + \frac{2(bBd+Abe-2aBe)(d+ex)^{5/2}}{5b^3} + \frac{2B(d+ex)^{7/2}}{7b^2} - \frac{(bd-ae)^{5/2}(2bBd+7Abe-9aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{11/2}}$$

output

```
2*(-a*e+b*d)^2*(3*A*b*e-4*B*a*e+B*b*d)*(e*x+d)^(1/2)/b^5-(A*b-B*a)*(-a*e+b*d)^3*(e*x+d)^(1/2)/b^5/(b*x+a)+2/3*(-a*e+b*d)*(2*A*b*e-3*B*a*e+B*b*d)*(e*x+d)^(3/2)/b^4+2/5*(A*b*e-2*B*a*e+B*b*d)*(e*x+d)^(5/2)/b^3+2/7*B*(e*x+d)^(7/2)/b^2-(-a*e+b*d)^(5/2)*(7*A*b*e-9*B*a*e+2*B*b*d)*arctanh(b^(1/2)*(e*x+d)^(1/2)/(-a*e+b*d)^(1/2))/b^(11/2)
```


Mathematica [A] (verified)

Time = 0.59 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.34

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^2} dx = \frac{\sqrt{d + ex}(-7Ab(-105a^3e^3 + 35a^2be^2(7d - 2ex) + 7ab^2e(-23d^2 + 24dex + 2(-bd + ae)^{5/2}(2bBd + 7Abe - 9aBe) \arctan\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right))}{b^{11/2}}$$

input

```
Integrate[((A + B*x)*(d + e*x)^(7/2))/(a + b*x)^2,x]
```

output

```
(Sqrt[d + e*x]*(-7*A*b*(-105*a^3*e^3 + 35*a^2*b*e^2*(7*d - 2*e*x) + 7*a*b^2*e*(-23*d^2 + 24*d*e*x + 2*e^2*x^2) + b^3*(15*d^3 - 116*d^2*e*x - 32*d*e^2*x^2 - 6*e^3*x^3)) + B*(-945*a^4*e^3 + 105*a^3*b*e^2*(23*d - 6*e*x) + 7*a^2*b^2*e*(-277*d^2 + 236*d*e*x + 18*e^2*x^2) + a*b^3*(457*d^3 - 1380*d^2*e*x - 316*d*e^2*x^2 - 54*e^3*x^3) + 2*b^4*x*(176*d^3 + 122*d^2*e*x + 66*d*e^2*x^2 + 15*e^3*x^3)))/(105*b^5*(a + b*x)) - (((-b*d) + a*e)^(5/2)*(2*b*B*d + 7*A*b*e - 9*a*B*e)*ArcTan[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[-(b*d) + a*e]])/b^(11/2)
```

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 222, normalized size of antiderivative = 0.97, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.318$, Rules used = {87, 60, 60, 60, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^2} dx$$

↓ 87

$$\frac{(-9aBe + 7Abe + 2bBd) \int \frac{(d+ex)^{7/2}}{a+bx} dx}{2b(bd - ae)} - \frac{(d + ex)^{9/2}(Ab - aB)}{b(a + bx)(bd - ae)}$$

↓ 60

$$\begin{aligned}
 & \frac{(-9aBe + 7Abe + 2bBd) \left(\frac{(bd-ae) \int \frac{(d+ex)^{5/2}}{a+bx} dx}{b} + \frac{2(d+ex)^{7/2}}{7b} \right)}{2b(bd-ae)} - \frac{(d+ex)^{9/2}(Ab-aB)}{b(a+bx)(bd-ae)} \\
 & \quad \downarrow 60 \\
 & \frac{(-9aBe + 7Abe + 2bBd) \left(\frac{(bd-ae) \left(\frac{(bd-ae) \int \frac{(d+ex)^{3/2}}{a+bx} dx}{b} + \frac{2(d+ex)^{5/2}}{5b} \right)}{b} + \frac{2(d+ex)^{7/2}}{7b} \right)}{2b(bd-ae)} - \frac{(d+ex)^{9/2}(Ab-aB)}{b(a+bx)(bd-ae)} \\
 & \quad \downarrow 60 \\
 & \frac{(-9aBe + 7Abe + 2bBd) \left(\frac{(bd-ae) \left(\frac{(bd-ae) \left(\frac{(bd-ae) \int \frac{\sqrt{d+ex}}{a+bx} dx}{b} + \frac{2(d+ex)^{3/2}}{3b} \right)}{b} + \frac{2(d+ex)^{5/2}}{5b} \right)}{b} + \frac{2(d+ex)^{7/2}}{7b} \right)}{2b(bd-ae)} - \frac{(d+ex)^{9/2}(Ab-aB)}{b(a+bx)(bd-ae)} \\
 & \quad \downarrow 60
 \end{aligned}$$

$$(-9aBe + 7Abe + 2bBd) \left(\frac{(bd-ae) \left(\frac{(bd-ae) \int \frac{1}{(a+bx)\sqrt{d+ex}} dx + 2\sqrt{d+ex}}{b} \right) + \frac{2(d+ex)^{3/2}}{3b}}{b} + \frac{2(d+ex)^{5/2}}{5b} \right) + \frac{2(d+ex)^{7/2}}{7b}$$

$$\frac{2b(bd - ae)}{(d + ex)^{9/2}(Ab - aB)} \frac{1}{b(a + bx)(bd - ae)}$$

73

$$(-9aBe + 7Abe + 2bBd) \left(\frac{(bd-ae) \left(\frac{2(bd-ae) \int \frac{1}{a + \frac{b(d+ex)}{e} - \frac{bd}{e}} d\sqrt{d+ex}}{b} + \frac{2\sqrt{d+ex}}{b} \right) + \frac{2(d+ex)^{3/2}}{3b}}{b} + \frac{2(d+ex)^{5/2}}{5b} \right) + \frac{2(d+ex)^{7/2}}{7b}$$

$$\frac{2b(bd - ae)}{(d + ex)^{9/2}(Ab - aB)} \frac{1}{b(a + bx)(bd - ae)}$$

221

$$\begin{aligned}
 & \left(\frac{(bd-ae) \left(\frac{(bd-ae) \left(\frac{2\sqrt{d+ex}}{b} - \frac{2\sqrt{bd-ae} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{3/2}} \right)}{b} + \frac{2(d+ex)^{3/2}}{3b} \right)}{b} + \frac{2(d+ex)^{5/2}}{5b} \right)}{b} + 2 \right) \\
 & \frac{(-9aBe + 7Abe + 2bBd)}{b} \\
 & \frac{2b(bd - ae)}{b(a + bx)(bd - ae)} \\
 & \frac{(d + ex)^{9/2}(Ab - aB)}{b(a + bx)(bd - ae)}
 \end{aligned}$$

input `Int[((A + B*x)*(d + e*x)^(7/2))/(a + b*x)^2,x]`

output `-(((A*b - a*B)*(d + e*x)^(9/2))/(b*(b*d - a*e)*(a + b*x))) + ((2*b*B*d + 7*A*b*e - 9*a*B*e)*((2*(d + e*x)^(7/2))/(7*b) + ((b*d - a*e)*((2*(d + e*x)^(5/2))/(5*b) + ((b*d - a*e)*((2*(d + e*x)^(3/2))/(3*b) + ((b*d - a*e)*((2*Sqrt[d + e*x])/b - (2*Sqrt[b*d - a*e]*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[b*d - a*e]])/b^(3/2)))/b))/b))/(2*b*(b*d - a*e))`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p
_), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Maple [A] (verified)

Time = 0.57 (sec) , antiderivative size = 278, normalized size of antiderivative = 1.22

method	result
pseudoelliptic	$-7\left(\left(Ae + \frac{2Bd}{7}\right)b - \frac{9Bae}{7}\right)(bx+a)(ae-db)^3 \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right) + 7\sqrt{ex+d}\sqrt{(ae-db)b} \left(\frac{2x^3\left(\frac{5Bx}{7} + A\right)e^3}{35} + \frac{32\left(\frac{33Bx}{56}\right)}{10}\right)$
risch	$\frac{2(15b^3Bx^3e^3 + 21Ab^3e^3x^2 - 42Bab^2e^3x + 66Bb^3de^2x^2 - 70Axa^2b^2e^3 + 112Ax^3b^3de^2 + 105Bxa^2be^3 - 224Bab^2de^2x + 10A^2a^2b^2e^3)}{10}$
derivativedivides	$\frac{\frac{2b^3B(e^3x+d)^{\frac{7}{2}}}{7} + \frac{2Ab^3e(e^3x+d)^{\frac{5}{2}}}{5} - \frac{4Bab^2e(e^3x+d)^{\frac{5}{2}}}{5} + \frac{2Bb^3d(e^3x+d)^{\frac{5}{2}}}{5} - \frac{4Aab^2e^2(e^3x+d)^{\frac{3}{2}}}{3} + \frac{4Ab^3de(e^3x+d)^{\frac{3}{2}}}{3} + 2Ba^2be^2(e^3x+d)^{\frac{3}{2}}}{10}$
default	$\frac{\frac{2b^3B(e^3x+d)^{\frac{7}{2}}}{7} + \frac{2Ab^3e(e^3x+d)^{\frac{5}{2}}}{5} - \frac{4Bab^2e(e^3x+d)^{\frac{5}{2}}}{5} + \frac{2Bb^3d(e^3x+d)^{\frac{5}{2}}}{5} - \frac{4Aab^2e^2(e^3x+d)^{\frac{3}{2}}}{3} + \frac{4Ab^3de(e^3x+d)^{\frac{3}{2}}}{3} + 2Ba^2be^2(e^3x+d)^{\frac{3}{2}}}{10}$

```
input int((B*x+A)*(e*x+d)^(7/2)/(b*x+a)^2,x,method=_RETURNVERBOSE)
```

output

```
7*(-((A*e+2/7*B*d)*b-9/7*B*a*e)*(b*x+a)*(a*e-b*d)^3*arctan(b*(e*x+d)^(1/2)
/((a*e-b*d)*b)^(1/2))+(e*x+d)^(1/2)*((a*e-b*d)*b)^(1/2)*((2/35*x^3*(5/7*B*
x+A)*e^3+32/105*(33/56*B*x+A)*x^2*d*e^2+116/105*(61/203*B*x+A)*x*d^2*e-1/7
*d^3*(-352/105*B*x+A))*b^4+23/15*a*(-2/23*(27/49*B*x+A)*x^2*e^3-24/23*x*d*
(79/294*B*x+A)*e^2+d^2*(-60/49*B*x+A)*e+457/1127*B*d^3)*b^3-7/3*a^2*e*(-2/
7*x*(9/35*B*x+A)*e^2+d*(-236/245*B*x+A)*e+277/245*B*d^2)*b^2+a^3*e^2*((-6/
7*B*x+A)*e+23/7*B*d)*b-9/7*B*a^4*e^3)/((a*e-b*d)*b)^(1/2)/b^5/(b*x+a)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 498 vs. $2(201) = 402$.

Time = 0.11 (sec) , antiderivative size = 1006, normalized size of antiderivative = 4.41

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^2} dx = \text{Too large to display}$$

input

```
integrate((B*x+A)*(e*x+d)^(7/2)/(b*x+a)^2,x, algorithm="fricas")
```

output

```
[1/210*(105*(2*B*a*b^3*d^3 - (13*B*a^2*b^2 - 7*A*a*b^3)*d^2*e + 2*(10*B*a^3*b - 7*A*a^2*b^2)*d*e^2 - (9*B*a^4 - 7*A*a^3*b)*e^3 + (2*B*b^4*d^3 - (13*B*a*b^3 - 7*A*b^4)*d^2*e + 2*(10*B*a^2*b^2 - 7*A*a*b^3)*d*e^2 - (9*B*a^3*b - 7*A*a^2*b^2)*e^3)*x)*sqrt((b*d - a*e)/b)*log((b*e*x + 2*b*d - a*e - 2*sqrt(e*x + d)*b*sqrt((b*d - a*e)/b))/(b*x + a)) + 2*(30*B*b^4*e^3*x^4 + (45*7*B*a*b^3 - 105*A*b^4)*d^3 - 7*(277*B*a^2*b^2 - 161*A*a*b^3)*d^2*e + 35*(6*9*B*a^3*b - 49*A*a^2*b^2)*d*e^2 - 105*(9*B*a^4 - 7*A*a^3*b)*e^3 + 6*(22*B*b^4*d*e^2 - (9*B*a*b^3 - 7*A*b^4)*e^3)*x^3 + 2*(122*B*b^4*d^2*e - 2*(79*B*a*b^3 - 56*A*b^4)*d*e^2 + 7*(9*B*a^2*b^2 - 7*A*a*b^3)*e^3)*x^2 + 2*(176*B*b^4*d^3 - 2*(345*B*a*b^3 - 203*A*b^4)*d^2*e + 14*(59*B*a^2*b^2 - 42*A*a*b^3)*d*e^2 - 35*(9*B*a^3*b - 7*A*a^2*b^2)*e^3)*x)*sqrt(e*x + d))/(b^6*x + a*b^5), -1/105*(105*(2*B*a*b^3*d^3 - (13*B*a^2*b^2 - 7*A*a*b^3)*d^2*e + 2*(10*B*a^3*b - 7*A*a^2*b^2)*d*e^2 - (9*B*a^4 - 7*A*a^3*b)*e^3 + (2*B*b^4*d^3 - (13*B*a*b^3 - 7*A*b^4)*d^2*e + 2*(10*B*a^2*b^2 - 7*A*a*b^3)*d*e^2 - (9*B*a^3*b - 7*A*a^2*b^2)*e^3)*x)*sqrt(-(b*d - a*e)/b)*arctan(-sqrt(e*x + d)*b*sqrt(-(b*d - a*e)/b)/(b*d - a*e)) - (30*B*b^4*e^3*x^4 + (45*7*B*a*b^3 - 105*A*b^4)*d^3 - 7*(277*B*a^2*b^2 - 161*A*a*b^3)*d^2*e + 35*(6*9*B*a^3*b - 49*A*a^2*b^2)*d*e^2 - 105*(9*B*a^4 - 7*A*a^3*b)*e^3 + 6*(22*B*b^4*d*e^2 - (9*B*a*b^3 - 7*A*b^4)*e^3)*x^3 + 2*(122*B*b^4*d^2*e - 2*(79*B*a*b^3 - 56*A*b^4)*d*e^2 + 7*(9*B*a^2*b^2 - 7*A*a*b^3)*e^3)*x^2 + 2*(176*B*b^4*d^3 - 2...
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^2} dx = \text{Timed out}$$

input

```
integrate((B*x+A)*(e*x+d)**(7/2)/(b*x+a)**2,x)
```

output

Timed out

Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^2} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)*(e*x+d)^(7/2)/(b*x+a)^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 583 vs. 2(201) = 402.

Time = 0.14 (sec) , antiderivative size = 583, normalized size of antiderivative = 2.56

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^2} dx = \frac{(2 Bb^4d^4 - 15 Bab^3d^3e + 7 Ab^4d^3e + 33 Ba^2b^2d^2e^2 - 21 Aab^3d^2e^2 - 29 Ba^3b^2d^2e^2 + \sqrt{-b^2d + abeb^5} - \sqrt{ex + d}Bab^3d^3e - \sqrt{ex + d}Ab^4d^3e - 3\sqrt{ex + d}Ba^2b^2d^2e^2 + 3\sqrt{ex + d}Aab^3d^2e^2 + 3\sqrt{ex + d}Ba^3bde^3 - ((ex + d)b - bd + ae)b^5}{((ex + d)b - bd + ae)b^5} + \frac{2 \left(15 (ex + d)^{\frac{7}{2}} Bb^{12} + 21 (ex + d)^{\frac{5}{2}} Bb^{12}d + 35 (ex + d)^{\frac{3}{2}} Bb^{12}d^2 + 105 \sqrt{ex + d} Bb^{12}d^3 - 42 (ex + d)^{\frac{5}{2}} Ba^3b^2d^2e^2 \right)}{((ex + d)b - bd + ae)b^5}$$

input `integrate((B*x+A)*(e*x+d)^(7/2)/(b*x+a)^2,x, algorithm="giac")`

output

```
(2*B*b^4*d^4 - 15*B*a*b^3*d^3*e + 7*A*b^4*d^3*e + 33*B*a^2*b^2*d^2*e^2 - 2
1*A*a*b^3*d^2*e^2 - 29*B*a^3*b*d*e^3 + 21*A*a^2*b^2*d*e^3 + 9*B*a^4*e^4 -
7*A*a^3*b*e^4)*arctan(sqrt(e*x + d)*b/sqrt(-b^2*d + a*b*e))/(sqrt(-b^2*d +
a*b*e)*b^5) + (sqrt(e*x + d)*B*a*b^3*d^3*e - sqrt(e*x + d)*A*b^4*d^3*e -
3*sqrt(e*x + d)*B*a^2*b^2*d^2*e^2 + 3*sqrt(e*x + d)*A*a*b^3*d^2*e^2 + 3*sq
rt(e*x + d)*B*a^3*b*d*e^3 - 3*sqrt(e*x + d)*A*a^2*b^2*d*e^3 - sqrt(e*x + d
)*B*a^4*e^4 + sqrt(e*x + d)*A*a^3*b*e^4)/(((e*x + d)*b - b*d + a*e)*b^5) +
2/105*(15*(e*x + d)^(7/2)*B*b^12 + 21*(e*x + d)^(5/2)*B*b^12*d + 35*(e*x
+ d)^(3/2)*B*b^12*d^2 + 105*sqrt(e*x + d)*B*b^12*d^3 - 42*(e*x + d)^(5/2)*
B*a*b^11*e + 21*(e*x + d)^(5/2)*A*b^12*e - 140*(e*x + d)^(3/2)*B*a*b^11*d*
e + 70*(e*x + d)^(3/2)*A*b^12*d*e - 630*sqrt(e*x + d)*B*a*b^11*d^2*e + 315
*sqrt(e*x + d)*A*b^12*d^2*e + 105*(e*x + d)^(3/2)*B*a^2*b^10*e^2 - 70*(e*x
+ d)^(3/2)*A*a*b^11*e^2 + 945*sqrt(e*x + d)*B*a^2*b^10*d*e^2 - 630*sqrt(e
*x + d)*A*a*b^11*d*e^2 - 420*sqrt(e*x + d)*B*a^3*b^9*e^3 + 315*sqrt(e*x +
d)*A*a^2*b^10*e^3)/b^14
```

Mupad [B] (verification not implemented)

Time = 1.01 (sec) , antiderivative size = 562, normalized size of antiderivative = 2.46

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^2} dx = \left(\frac{2Ae - 2Bd}{5b^2} + \frac{2B(2b^2d - 2abe)}{5b^4} \right) (d + ex)^{5/2}$$

$$+ \left(\frac{\left(\frac{(2b^2d - 2abe) \left(\frac{2Ae - 2Bd}{b^2} + \frac{2B(2b^2d - 2abe)}{b^4} \right)}{b^2} - \frac{2B(ae - bd)^2}{b^4} \right) (2b^2d - 2abe)}{b^2} - \frac{(ae - bd)^2 \left(\frac{2Ae - 2Bd}{b^2} + \frac{2B(2b^2d - 2abe)}{b^4} \right)}{b^2} \right) (d + ex)^{3/2}$$

$$- \frac{\sqrt{d + ex} (Ba^4e^4 - 3Ba^3bde^3 - Aa^3be^4 + 3Ba^2b^2d^2e^2 + 3Aa^2b^2de^3 - Bab^3d^3e - 3Aab^3d^2e^2)}{b^6(d + ex) - b^6d + ab^5e}$$

$$+ \frac{2B(d + ex)^{7/2}}{7b^2}$$

$$+ \frac{\operatorname{atan}\left(\frac{\sqrt{b}(ae - bd)^{5/2} \sqrt{d + ex} (7Abe - 9Bae + 2Bbd)}{9Ba^4e^4 - 29Ba^3bde^3 - 7Aa^3be^4 + 33Ba^2b^2d^2e^2 + 21Aa^2b^2de^3 - 15Bab^3d^3e - 21Aab^3d^2e^2 + 2Bb^4d^4 + 7Ab^4d^3e} \right)}{b^{11/2}} (ae - bd)$$

input `int(((A + B*x)*(d + e*x)^(7/2))/(a + b*x)^2,x)`

output
$$\begin{aligned} & \left(\frac{2Ae - 2Bd}{5b^2} + \frac{2B(2b^2d - 2a*be)}{5b^4} \right) (d + ex)^{5/2} + \left(\frac{((2b^2d - 2a*be)((2Ae - 2Bd)/b^2 + (2B(2b^2d - 2a*be))/b^4))/b^2 - (2B(ae - bd)^2)/b^4 * (2b^2d - 2a*be)/b^2 - ((ae - bd)^2((2Ae - 2Bd)/b^2 + (2B(2b^2d - 2a*be))/b^4))/b^2 * (d + ex)^{1/2} + ((2b^2d - 2a*be)((2Ae - 2Bd)/b^2 + (2B(2b^2d - 2a*be))/b^4))/(3b^2) - (2B(ae - bd)^2)/(3b^4) * (d + ex)^{3/2} - ((d + ex)^{1/2}(Ba^4e^4 - Aa^3be^4 + Ab^4d^3e - 3Aa^2b^3d^2e^2 + 3Aa^2b^2d^2e^3 + 3Ba^2b^2d^2e^2 - Babb^3d^3e - 3Ba^3bde^3))/(b^6(d + ex) - b^6d + ab^5e) + (2B(d + ex)^{7/2})/(7b^2) + \left(\frac{\operatorname{atan}(b^{1/2}(ae - bd)^{5/2}(d + ex)^{1/2}(7Abe - 9Bae + 2Bbd))}{(9Ba^4e^4 + 2Bb^4d^4 - 7Aa^3be^4 + 7Ab^4d^3e - 21Aa^2b^3d^2e^2 + 21Aa^2b^2d^2e^3 + 33Ba^2b^2d^2e^2 - 15Babb^3d^3e - 29Ba^3bde^3)} \right) (ae - bd)^{5/2} (7Abe - 9Bae + 2Bbd) \right) / b^{11/2} \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 349, normalized size of antiderivative = 1.53

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^2} dx = \frac{2\sqrt{b}\sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b}\sqrt{ae-bd}}\right) a^3e^3 - 6\sqrt{b}\sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b}\sqrt{ae-bd}}\right) a^2bd e}{(a + bx)^2}$$

input `int((B*x+A)*(e*x+d)^(7/2)/(b*x+a)^2,x)`

output
$$\begin{aligned} & (2*(105*\sqrt{b}*\sqrt{ae - bd})*\operatorname{atan}(\sqrt{(d + ex)*b}/(\sqrt{b}*\sqrt{ae - bd}))*a^{**3}e^{**3} - 315*\sqrt{b}*\sqrt{ae - bd})*\operatorname{atan}(\sqrt{(d + ex)*b}/(\sqrt{b}*\sqrt{ae - bd}))*a^{**2}b*d*e^{**2} + 315*\sqrt{b}*\sqrt{ae - bd})*\operatorname{atan}(\sqrt{(d + ex)*b}/(\sqrt{b}*\sqrt{ae - bd}))*a*b^{**2}d^{**2}e - 105*\sqrt{b}*\sqrt{ae - bd})*\operatorname{atan}(\sqrt{(d + ex)*b}/(\sqrt{b}*\sqrt{ae - bd}))*b^{**3}d^{**3} - 105*\sqrt{(d + ex)*a^{**3}b*e^{**3} + 350*\sqrt{(d + ex)*a^{**2}b^{**2}d*e^{**2} + 35*\sqrt{(d + ex)*a^{**2}b^{**2}e^{**3}x - 406*\sqrt{(d + ex)*a*b^{**3}d^{**2}e - 112*\sqrt{(d + ex)*a*b^{**3}d*e^{**2}x - 21*\sqrt{(d + ex)*a*b^{**3}e^{**3}x^{**2} + 176*\sqrt{(d + ex)*b^{**4}d^{**3} + 122*\sqrt{(d + ex)*b^{**4}d^{**2}e*x + 66*\sqrt{(d + ex)*b^{**4}d*e^{**2}x^{**2} + 15*\sqrt{(d + ex)*b^{**4}e^{**3}x^{**3}})}))/(105*b^{**5} \end{aligned}$$

3.157 $\int \frac{(A+Bx)(d+ex)^{5/2}}{(a+bx)^2} dx$

Optimal result	1446
Mathematica [A] (verified)	1447
Rubi [A] (verified)	1447
Maple [A] (verified)	1450
Fricas [A] (verification not implemented)	1451
Sympy [F(-1)]	1452
Maxima [F(-2)]	1452
Giac [B] (verification not implemented)	1452
Mupad [B] (verification not implemented)	1453
Reduce [B] (verification not implemented)	1454

Optimal result

Integrand size = 22, antiderivative size = 187

$$\int \frac{(A+Bx)(d+ex)^{5/2}}{(a+bx)^2} dx = \frac{2(bd-ae)(bBd+2Abe-3aBe)\sqrt{d+ex}}{b^4} - \frac{(Ab-aB)(bd-ae)^2\sqrt{d+ex}}{b^4(a+bx)} + \frac{2(bBd+Abe-2aBe)(d+ex)^{3/2}}{3b^3} + \frac{2B(d+ex)^{5/2}}{5b^2} - \frac{(bd-ae)^{3/2}(2bBd+5Abe-7aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{9/2}}$$

output

```
2*(-a*e+b*d)*(2*A*b*e-3*B*a*e+B*b*d)*(e*x+d)^(1/2)/b^4-(A*b-B*a)*(-a*e+b*d)^(1/2)*(e*x+d)^(1/2)/b^4/(b*x+a)+2/3*(A*b*e-2*B*a*e+B*b*d)*(e*x+d)^(3/2)/b^3+2/5*B*(e*x+d)^(5/2)/b^2-(-a*e+b*d)^(3/2)*(5*A*b*e-7*B*a*e+2*B*b*d)*arctanh(b^(1/2)*(e*x+d)^(1/2)/(-a*e+b*d)^(1/2))/b^(9/2)
```

Mathematica [A] (verified)

Time = 0.46 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.12

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^2} dx = \frac{\sqrt{d + ex}(-5Ab(15a^2e^2 + 10abe(-2d + ex) + b^2(3d^2 - 14dex - 2e^2x^2)) + B(-bd + ae)^{3/2}(2bBd + 5Abe - 7aBe) \arctan\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right)}{b^{9/2}}$$

input

```
Integrate[((A + B*x)*(d + e*x)^(5/2))/(a + b*x)^2,x]
```

output

```
(Sqrt[d + e*x]*(-5*A*b*(15*a^2*e^2 + 10*a*b*e*(-2*d + e*x) + b^2*(3*d^2 - 14*d*e*x - 2*e^2*x^2)) + B*(105*a^3*e^2 + 10*a^2*b*e*(-17*d + 7*e*x) + a*b^2*(61*d^2 - 118*d*e*x - 14*e^2*x^2) + 2*b^3*x*(23*d^2 + 11*d*e*x + 3*e^2*x^2)))/(15*b^4*(a + b*x)) + ((-(b*d) + a*e)^(3/2)*(2*b*B*d + 5*A*b*e - 7*a*B*e)*ArcTan[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[-(b*d) + a*e]])/b^(9/2)
```

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.03, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {87, 60, 60, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^2} dx$$

$$\downarrow 87$$

$$\frac{(-7aBe + 5Abe + 2bBd) \int \frac{(d+ex)^{5/2}}{a+bx} dx}{2b(bd - ae)} - \frac{(d + ex)^{7/2}(Ab - aB)}{b(a + bx)(bd - ae)}$$

$$\downarrow 60$$

$$\frac{(-7aBe + 5Abe + 2bBd) \left(\frac{(bd-ae) \int \frac{(d+ex)^{3/2}}{a+bx} dx}{b} + \frac{2(d+ex)^{5/2}}{5b} \right)}{2b(bd - ae)} - \frac{(d + ex)^{7/2}(Ab - aB)}{b(a + bx)(bd - ae)}$$

$$\begin{aligned} & \downarrow 60 \\ & (-7aBe + 5Abe + 2bBd) \left(\frac{(bd-ae) \left(\frac{(bd-ae) \int \frac{\sqrt{d+ex}}{a+bx} dx}{b} + \frac{2(d+ex)^{3/2}}{3b} \right)}{b} + \frac{2(d+ex)^{5/2}}{5b} \right) \\ & \hline & \frac{2b(bd-ae)}{(d+ex)^{7/2}(Ab-aB)} \\ & \frac{b(a+bx)(bd-ae)}{b(a+bx)(bd-ae)} \end{aligned}$$

$$\begin{aligned} & \downarrow 60 \\ & (-7aBe + 5Abe + 2bBd) \left(\frac{(bd-ae) \left(\frac{(bd-ae) \int \frac{1}{(a+bx)\sqrt{d+ex}} dx}{b} + \frac{2\sqrt{d+ex}}{b} \right)}{b} + \frac{2(d+ex)^{3/2}}{3b} \right) + \frac{2(d+ex)^{5/2}}{5b} \\ & \hline & \frac{2b(bd-ae)}{(d+ex)^{7/2}(Ab-aB)} \\ & \frac{b(a+bx)(bd-ae)}{b(a+bx)(bd-ae)} \end{aligned}$$

$$\begin{aligned} & \downarrow 73 \\ & (-7aBe + 5Abe + 2bBd) \left(\frac{(bd-ae) \left(\frac{2(bd-ae) \int \frac{1}{a + \frac{b(d+ex)}{e} - \frac{bd}{e}} d\sqrt{d+ex}}{b} + \frac{2\sqrt{d+ex}}{b} \right)}{b} + \frac{2(d+ex)^{3/2}}{3b} \right) + \frac{2(d+ex)^{5/2}}{5b} \\ & \hline & \frac{2b(bd-ae)}{(d+ex)^{7/2}(Ab-aB)} \\ & \frac{b(a+bx)(bd-ae)}{b(a+bx)(bd-ae)} \end{aligned}$$

$$\downarrow 221$$

$$\begin{aligned}
 & (-7aBe + 5Abe + 2bBd) \left(\frac{(bd-ae) \left(\frac{2\sqrt{d+ex}}{b} - \frac{2\sqrt{bd-ae} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{3/2}} \right)}{b} + \frac{2(d+ex)^{3/2}}{3b} \right) + \frac{2(d+ex)^{5/2}}{5b} \\
 & \frac{2b(bd-ae)}{(d+ex)^{7/2}(Ab-aB)} \\
 & \frac{b(a+bx)(bd-ae)}{b(a+bx)(bd-ae)}
 \end{aligned}$$

input `Int[((A + B*x)*(d + e*x)^(5/2))/(a + b*x)^2,x]`

output `-(((A*b - a*B)*(d + e*x)^(7/2))/(b*(b*d - a*e)*(a + b*x))) + ((2*b*B*d + 5*A*b*e - 7*a*B*e)*((2*(d + e*x)^(5/2))/(5*b) + ((b*d - a*e)*((2*(d + e*x)^(3/2))/(3*b) + ((b*d - a*e)*((2*sqrt[d + e*x])/b - (2*sqrt[b*d - a*e]*ArcTanh[(sqrt[b]*sqrt[d + e*x])/sqrt[b*d - a*e]])/b^(3/2))/b))/b))/(2*b*(b*d - a*e))`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 87

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p
_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

rule 221

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.13

method	result
risch	$\frac{-2(-3e^2b^2Bx^2 - 5Ab^2e^2x + 10Babe^2x - 11b^2Bdex + 30Aabe^2 - 35Ab^2de - 45Ba^2e^2 + 70Babde - 23b^2Bd^2)\sqrt{ex+d}}{15b^4}$
pseudoelliptic	$-5\sqrt{(ae-db)b} \left(\frac{\left(\frac{2(-\frac{11}{3}dex^2 - \frac{23}{3}d^2x - e^2x^3)b^3}{5} - 61a\left(-\frac{14}{61}e^2x^2 - \frac{118}{61}dex + d^2\right)b^2 + \frac{34a^2(-\frac{7ex}{17} + d)eb}{3} - 7a^3e^2 \right)B}{5} \right) + A \left(\dots \right)$
derivativedivides	$\frac{2 \left(-\frac{b^2B(ex+d)^{\frac{5}{2}}}{5} - \frac{Ab^2e(ex+d)^{\frac{3}{2}}}{3} + \frac{2Babe(ex+d)^{\frac{3}{2}}}{3} - \frac{Bb^2d(ex+d)^{\frac{3}{2}}}{3} + 2Aabe^2\sqrt{ex+d} - 2Ab^2de\sqrt{ex+d} - 3Ba^2e^2\sqrt{ex+d} \right)}{b^4}$
default	$\frac{2 \left(-\frac{b^2B(ex+d)^{\frac{5}{2}}}{5} - \frac{Ab^2e(ex+d)^{\frac{3}{2}}}{3} + \frac{2Babe(ex+d)^{\frac{3}{2}}}{3} - \frac{Bb^2d(ex+d)^{\frac{3}{2}}}{3} + 2Aabe^2\sqrt{ex+d} - 2Ab^2de\sqrt{ex+d} - 3Ba^2e^2\sqrt{ex+d} \right)}{b^4}$

input

```
int((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^2,x,method=_RETURNVERBOSE)
```

output

```
-2/15*(-3*B*b^2*e^2*x^2-5*A*b^2*e^2*x+10*B*a*b*e^2*x-11*B*b^2*d*e*x+30*A*a
*b*e^2-35*A*b^2*d*e-45*B*a^2*e^2+70*B*a*b*d*e-23*B*b^2*d^2)*(e*x+d)^(1/2)/
b^4+1/b^4*(2*a^2*e^2-4*a*b*d*e+2*b^2*d^2)*((-1/2*A*b*e+1/2*B*a*e)*(e*x+d)^(
1/2)/((e*x+d)*b+a*e-d*b)+1/2*(5*A*b*e-7*B*a*e+2*B*b*d)/((a*e-b*d)*b)^(1/2
)*arctan(b*(e*x+d)^(1/2)/((a*e-b*d)*b)^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 666, normalized size of antiderivative = 3.56

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^2} dx = \left[\frac{15(2Bab^2d^2 - (9Ba^2b - 5Aab^2)de + (7Ba^3 - 5Aa^2b)e^2 + (2Bb^3d^2 - (9Bab^2 - 5Ab^3)de + (7Ba^2b - 5Aab^2)e^2)}{(a + bx)^2} \right]$$

$$\frac{15(2Bab^2d^2 - (9Ba^2b - 5Aab^2)de + (7Ba^3 - 5Aa^2b)e^2 + (2Bb^3d^2 - (9Bab^2 - 5Ab^3)de + (7Ba^2b - 5Aab^2)e^2)}{(a + bx)^2}$$

input `integrate((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^2,x, algorithm="fricas")`

output

```
[-1/30*(15*(2*B*a*b^2*d^2 - (9*B*a^2*b - 5*A*a*b^2)*d*e + (7*B*a^3 - 5*A*a^2*b)*e^2 + (2*B*b^3*d^2 - (9*B*a*b^2 - 5*A*b^3)*d*e + (7*B*a^2*b - 5*A*a*b^2)*e^2)*x)*sqrt((b*d - a*e)/b)*log((b*e*x + 2*b*d - a*e + 2*sqrt(e*x + d))*b*sqrt((b*d - a*e)/b))/(b*x + a) - 2*(6*B*b^3*e^2*x^3 + (61*B*a*b^2 - 15*A*b^3)*d^2 - 10*(17*B*a^2*b - 10*A*a*b^2)*d*e + 15*(7*B*a^3 - 5*A*a^2*b)*e^2 + 2*(11*B*b^3*d*e - (7*B*a*b^2 - 5*A*b^3)*e^2)*x^2 + 2*(23*B*b^3*d^2 - (59*B*a*b^2 - 35*A*b^3)*d*e + 5*(7*B*a^2*b - 5*A*a*b^2)*e^2)*x)*sqrt(e*x + d))/(b^5*x + a*b^4), -1/15*(15*(2*B*a*b^2*d^2 - (9*B*a^2*b - 5*A*a*b^2)*d*e + (7*B*a^3 - 5*A*a^2*b)*e^2 + (2*B*b^3*d^2 - (9*B*a*b^2 - 5*A*b^3)*d*e + (7*B*a^2*b - 5*A*a*b^2)*e^2)*x)*sqrt(-(b*d - a*e)/b)*arctan(-sqrt(e*x + d))*b*sqrt(-(b*d - a*e)/b)/(b*d - a*e) - (6*B*b^3*e^2*x^3 + (61*B*a*b^2 - 15*A*b^3)*d^2 - 10*(17*B*a^2*b - 10*A*a*b^2)*d*e + 15*(7*B*a^3 - 5*A*a^2*b)*e^2 + 2*(11*B*b^3*d*e - (7*B*a*b^2 - 5*A*b^3)*e^2)*x^2 + 2*(23*B*b^3*d^2 - (59*B*a*b^2 - 35*A*b^3)*d*e + 5*(7*B*a^2*b - 5*A*a*b^2)*e^2)*x)*sqrt(e*x + d))/(b^5*x + a*b^4)]
```


Sympy [F(-1)]

Timed out.

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^2} dx = \text{Timed out}$$

input `integrate((B*x+A)*(e*x+d)**(5/2)/(b*x+a)**2,x)`output `Timed out`**Maxima [F(-2)]**

Exception generated.

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^2} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^2,x, algorithm="maxima")`output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail`**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 382 vs. 2(164) = 328.

Time = 0.13 (sec) , antiderivative size = 382, normalized size of antiderivative = 2.04

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^2} dx = \frac{(2 Bb^3d^3 - 11 Bab^2d^2e + 5 Ab^3d^2e + 16 Ba^2bde^2 - 10 Aab^2de^2 - 7 Ba^3e^3 + \sqrt{-b^2d + abeb^4} + \sqrt{ex + d} Bab^2d^2e - \sqrt{ex + d} Ab^3d^2e - 2\sqrt{ex + d} Ba^2bde^2 + 2\sqrt{ex + d} Aab^2de^2 + \sqrt{ex + d} Ba^3e^3 - \sqrt{ex + d} Aa^3e^3)}{((ex + d)b - bd + ae)b^4} + \frac{2 \left(3(ex + d)^{\frac{5}{2}} Bb^8 + 5(ex + d)^{\frac{3}{2}} Bb^8d + 15\sqrt{ex + d} Bb^8d^2 - 10(ex + d)^{\frac{3}{2}} Bab^7e + 5(ex + d)^{\frac{3}{2}} Ab^8e - 60 \right)}{15b^{10}}$$

input `integrate((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^2,x, algorithm="giac")`

output
$$\begin{aligned} & (2*B*b^3*d^3 - 11*B*a*b^2*d^2*e + 5*A*b^3*d^2*e + 16*B*a^2*b*d*e^2 - 10*A* \\ & a*b^2*d*e^2 - 7*B*a^3*e^3 + 5*A*a^2*b*e^3)*\arctan(\sqrt{e*x + d}*b/\sqrt{-b^2*d + a*b*e})/(\sqrt{-b^2*d + a*b*e}*b^4) + (\sqrt{e*x + d}*B*a*b^2*d^2*e - \\ & \sqrt{e*x + d}*A*b^3*d^2*e - 2*\sqrt{e*x + d}*B*a^2*b*d*e^2 + 2*\sqrt{e*x + d} \\ &)*A*a*b^2*d*e^2 + \sqrt{e*x + d}*B*a^3*e^3 - \sqrt{e*x + d}*A*a^2*b*e^3)/(((\\ & e*x + d)*b - b*d + a*e)*b^4) + 2/15*(3*(e*x + d)^(5/2)*B*b^8 + 5*(e*x + d) \\ & ^{(3/2)*B*b^8*d + 15*\sqrt{e*x + d}*B*b^8*d^2 - 10*(e*x + d)^(3/2)*B*a*b^7*e \\ & + 5*(e*x + d)^(3/2)*A*b^8*e - 60*\sqrt{e*x + d}*B*a*b^7*d*e + 30*\sqrt{e*x \\ & + d}*A*b^8*d*e + 45*\sqrt{e*x + d}*B*a^2*b^6*e^2 - 30*\sqrt{e*x + d}*A*a*b^7 \\ & *e^2)/b^{10} \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 1.00 (sec) , antiderivative size = 363, normalized size of antiderivative = 1.94

$$\begin{aligned} & \int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^2} dx = \left(\frac{2Ae - 2Bd}{3b^2} + \frac{2B(2b^2d - 2abe)}{3b^4} \right) (d + ex)^{3/2} \\ & + \left(\frac{(2b^2d - 2abe) \left(\frac{2Ae - 2Bd}{b^2} + \frac{2B(2b^2d - 2abe)}{b^4} \right) - \frac{2B(ae - bd)^2}{b^4}}{b^2} \right) \sqrt{d + ex} \\ & + \frac{\sqrt{d + ex} (Ba^3e^3 - 2Ba^2bde^2 - Aa^2be^3 + Bab^2d^2e + 2Aab^2de^2 - Ab^3d^2e)}{b^5(d + ex) - b^5d + ab^4e} \\ & + \frac{2B(d + ex)^{5/2}}{5b^2} \\ & + \frac{\operatorname{atan}\left(\frac{\sqrt{b}(ae - bd)^{3/2} \sqrt{d + ex} (5Abe - 7Bae + 2Bbd)}{-7Ba^3e^3 + 16Ba^2bde^2 + 5Aa^2be^3 - 11Ba^2d^2e - 10Aab^2de^2 + 2Bb^3d^3 + 5Ab^3d^2e}\right)}{b^{9/2}} (ae - bd)^{3/2} (5Abe - 7Bae) \end{aligned}$$

input `int(((A + B*x)*(d + e*x)^(5/2))/(a + b*x)^2,x)`

output

$$\begin{aligned} & ((2Ae - 2Bd)/(3b^2) + (2B(2b^2d - 2a*be))/(3b^4))*(d + ex)^{3/2} \\ & + (((2b^2d - 2a*be)*((2Ae - 2Bd)/b^2 + (2B(2b^2d - 2a*be))/b^4))/b^2 - (2B(ae - bd)^2)/b^4)*(d + ex)^{1/2} \\ & + ((d + ex)^{1/2}*(B*a^3*e^3 - A*a^2*b*e^3 - A*b^3*d^2*e + 2A*a*b^2*d*e^2 + B*a*b^2*d^2*e - 2B*a^2*b*d*e^2))/(b^5*(d + ex) - b^5*d + a*b^4*e) \\ & + (2B*(d + ex)^{5/2})/(5*b^2) + (atan((b^{1/2})*(ae - bd)^{3/2}*(d + ex)^{1/2}*(5A*be - 7B*a*e + 2B*b*d))/(2B*b^3*d^3 - 7B*a^3*e^3 + 5A*a^2*b*e^3 + 5A*b^3*d^2*e - 10A*a*b^2*d*e^2 - 11B*a*b^2*d^2*e + 16B*a^2*b*d*e^2))*(ae - bd)^{3/2}*(5A*be - 7B*a*e + 2B*b*d))/b^{9/2} \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.19

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^2} dx = \frac{-2\sqrt{b}\sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b}\sqrt{ae-bd}}\right) a^2 e^2 + 4\sqrt{b}\sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b}\sqrt{ae-bd}}\right) abde}{(a + bx)^2}$$

input

```
int((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^2,x)
```

output

$$\begin{aligned} & (2*(-15*\sqrt{b}*\sqrt{ae - bd})*\operatorname{atan}(\sqrt{(d + ex)*b}/(\sqrt{b}*\sqrt{ae - bd}))*a**2*e**2 + 30*\sqrt{b}*\sqrt{ae - bd})*\operatorname{atan}(\sqrt{(d + ex)*b}/(\sqrt{b}*\sqrt{ae - bd}))*a*b*d*e \\ & - 15*\sqrt{b}*\sqrt{ae - bd})*\operatorname{atan}(\sqrt{(d + ex)*b}/(\sqrt{b}*\sqrt{ae - bd}))*b**2*d**2 + 15*\sqrt{d + ex}*a**2*b*e**2 \\ & - 35*\sqrt{d + ex}*a*b**2*d*e - 5*\sqrt{d + ex}*a*b**2*e**2*x + 23*\sqrt{d + ex}*b**3*d**2 \\ & + 11*\sqrt{d + ex}*b**3*d*e*x + 3*\sqrt{d + ex}*b**3*e**2*x**2))/(15*b**4) \end{aligned}$$

3.158 $\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^2} dx$

Optimal result	1455
Mathematica [A] (verified)	1456
Rubi [A] (verified)	1456
Maple [A] (verified)	1458
Fricas [A] (verification not implemented)	1459
Sympy [F(-1)]	1460
Maxima [F(-2)]	1460
Giac [A] (verification not implemented)	1461
Mupad [B] (verification not implemented)	1461
Reduce [B] (verification not implemented)	1462

Optimal result

Integrand size = 22, antiderivative size = 146

$$\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^2} dx = \frac{2(bBd + Abe - 2aBe)\sqrt{d+ex}}{b^3} - \frac{(Ab - aB)(bd - ae)\sqrt{d+ex}}{b^3(a+bx)} + \frac{2B(d+ex)^{3/2}}{3b^2} - \frac{\sqrt{bd - ae}(2bBd + 3Abe - 5aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd - ae}}\right)}{b^{7/2}}$$

output

```
2*(A*b*e-2*B*a*e+B*b*d)*(e*x+d)^(1/2)/b^3-(A*b-B*a)*(-a*e+b*d)*(e*x+d)^(1/2)/b^3/(b*x+a)+2/3*B*(e*x+d)^(3/2)/b^2-(-a*e+b*d)^(1/2)*(3*A*b*e-5*B*a*e+2*B*b*d)*arctanh(b^(1/2)*(e*x+d)^(1/2)/(-a*e+b*d)^(1/2))/b^(7/2)
```

Mathematica [A] (verified)

Time = 0.34 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.95

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^2} dx = \frac{\sqrt{d + ex}(3Ab(-bd + 3ae + 2bex) + B(-15a^2e + ab(11d - 10ex) + 2b^2x(4d - 10e)) + 2b^2x(4d - 10e))}{3b^3(a + bx)} - \frac{\sqrt{-bd + ae}(2bBd + 3Abe - 5aBe) \arctan\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right)}{b^{7/2}}$$

input

```
Integrate[((A + B*x)*(d + e*x)^(3/2))/(a + b*x)^2,x]
```

output

```
(Sqrt[d + e*x]*(3*A*b*(-(b*d) + 3*a*e + 2*b*e*x) + B*(-15*a^2*e + a*b*(11*d - 10*e*x) + 2*b^2*x*(4*d + e*x)))/(3*b^3*(a + b*x)) - (Sqrt[-(b*d) + a*e]*(2*b*B*d + 3*A*b*e - 5*a*B*e)*ArcTan[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[-(b*d) + a*e]])/b^(7/2)
```

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.12, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.227$, Rules used = {87, 60, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^2} dx$$

$$\downarrow 87$$

$$\frac{(-5aBe + 3Abe + 2bBd) \int \frac{(d+ex)^{3/2}}{a+bx} dx}{2b(bd - ae)} - \frac{(d + ex)^{5/2}(Ab - aB)}{b(a + bx)(bd - ae)}$$

$$\downarrow 60$$

$$\frac{(-5aBe + 3Abe + 2bBd) \left(\frac{(bd-ae) \int \frac{\sqrt{d+ex}}{a+bx} dx}{b} + \frac{2(d+ex)^{3/2}}{3b} \right)}{2b(bd - ae)} - \frac{(d + ex)^{5/2}(Ab - aB)}{b(a + bx)(bd - ae)}$$

$$\begin{aligned} & \downarrow 60 \\ & (-5aBe + 3Abe + 2bBd) \left(\frac{(bd-ae) \left(\frac{(bd-ae) \int \frac{1}{(a+bx)\sqrt{d+ex}} dx}{b} + \frac{2\sqrt{d+ex}}{b} \right)}{b} + \frac{2(d+ex)^{3/2}}{3b} \right) \\ & \hline & \frac{2b(bd - ae)}{(d + ex)^{5/2}(Ab - aB)} \\ & \frac{b(a + bx)(bd - ae)}{b(a + bx)(bd - ae)} \end{aligned}$$

$$\begin{aligned} & \downarrow 73 \\ & (-5aBe + 3Abe + 2bBd) \left(\frac{(bd-ae) \left(\frac{2(bd-ae) \int \frac{1}{a + \frac{b(d+ex)}{e} - \frac{bd}{e}} d\sqrt{d+ex}}{be} + \frac{2\sqrt{d+ex}}{b} \right)}{b} + \frac{2(d+ex)^{3/2}}{3b} \right) \\ & \hline & \frac{2b(bd - ae)}{(d + ex)^{5/2}(Ab - aB)} \\ & \frac{b(a + bx)(bd - ae)}{b(a + bx)(bd - ae)} \end{aligned}$$

$$\begin{aligned} & \downarrow 221 \\ & (-5aBe + 3Abe + 2bBd) \left(\frac{(bd-ae) \left(\frac{2\sqrt{d+ex}}{b} - \frac{2\sqrt{bd-ae} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{3/2}} \right)}{b} + \frac{2(d+ex)^{3/2}}{3b} \right) \\ & \hline & \frac{2b(bd - ae)}{(d + ex)^{5/2}(Ab - aB)} \\ & \frac{b(a + bx)(bd - ae)}{b(a + bx)(bd - ae)} \end{aligned}$$

input `Int[((A + B*x)*(d + e*x)^(3/2))/(a + b*x)^2,x]`

output `-(((A*b - a*B)*(d + e*x)^(5/2))/(b*(b*d - a*e)*(a + b*x))) + ((2*b*B*d + 3*A*b*e - 5*a*B*e)*((2*(d + e*x)^(3/2))/(3*b) + ((b*d - a*e)*((2*sqrt[d + e*x])/b - (2*sqrt[b*d - a*e]*ArcTanh[(sqrt[b]*sqrt[d + e*x])/sqrt[b*d - a*e]])/b^(3/2)))/b)/(2*b*(b*d - a*e))`

Definitions of rubi rules used

- rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || (EqQ[e, 0] || (EqQ[c, 0] || LtQ[p, n]))))`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.95

method	result
risch	$\frac{2(ebBx+3Abe-6Bae+4Bbd)\sqrt{ex+d}}{3b^3} - \frac{(2ae-2db) \left(\frac{(-\frac{1}{2}Abe+\frac{1}{2}Bae)\sqrt{ex+d}}{(ex+d)b+ae-db} + \frac{(3Abe-5Bae+2Bbd) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{2\sqrt{(ae-db)b}} \right)}{b^3}$
pseudoelliptic	$\frac{-3\left(\left(Ae+\frac{2Bd}{3}\right)b-\frac{5Bae}{3}\right)(bx+a)(ae-db) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right) + 3\sqrt{(ae-db)b}\sqrt{ex+d} \left(\frac{2\left(\frac{Bx}{3}+A\right)xe-\left(-\frac{8Bx}{3}+A\right)d}{3} \right)}{b^3(bx+a)\sqrt{(ae-db)b}}$
derivativedivides	$\frac{\frac{2bB(ex+d)^{\frac{3}{2}}}{3} + 2Abe\sqrt{ex+d} - 4Bae\sqrt{ex+d} + 2Bbd\sqrt{ex+d}}{b^3} - \frac{2 \left(\frac{(-\frac{1}{2}Aab e^2 + \frac{1}{2}A b^2 de + \frac{1}{2}B a^2 e^2 - \frac{1}{2}B abde)\sqrt{ex+d}}{(ex+d)b+ae-db} + \frac{(3Ae-5Bae+2Bbd) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{2\sqrt{(ae-db)b}} \right)}{b^3}$
default	$\frac{\frac{2bB(ex+d)^{\frac{3}{2}}}{3} + 2Abe\sqrt{ex+d} - 4Bae\sqrt{ex+d} + 2Bbd\sqrt{ex+d}}{b^3} - \frac{2 \left(\frac{(-\frac{1}{2}Aab e^2 + \frac{1}{2}A b^2 de + \frac{1}{2}B a^2 e^2 - \frac{1}{2}B abde)\sqrt{ex+d}}{(ex+d)b+ae-db} + \frac{(3Ae-5Bae+2Bbd) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{2\sqrt{(ae-db)b}} \right)}{b^3}$

```
input int((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^2,x,method=_RETURNVERBOSE)
```

```
output 2/3*(B*b*e*x+3*A*b*e-6*B*a*e+4*B*b*d)*(e*x+d)^(1/2)/b^3-1/b^3*(2*a*e-2*b*d)
)*((-1/2*A*b*e+1/2*B*a*e)*(e*x+d)^(1/2)/((e*x+d)*b+a*e-d*b)+1/2*(3*A*b*e-5
*B*a*e+2*B*b*d)/((a*e-b*d)*b)^(1/2)*arctan(b*(e*x+d)^(1/2)/((a*e-b*d)*b)^(
1/2)))
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 392, normalized size of antiderivative = 2.68

$$\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^2} dx = \frac{3(2Babd - (5Ba^2 - 3Aab)e + (2Bb^2d - (5Bab - 3Ab^2)e)x) \sqrt{\frac{bd-ae}{b}} \log\left(\frac{\sqrt{ex+db}\sqrt{-\frac{bd-ae}{b}}}{bd-ae}\right) - (2Babd - (5Ba^2 - 3Aab)e + (2Bb^2d - (5Bab - 3Ab^2)e)x) \sqrt{-\frac{bd-ae}{b}} \arctan\left(-\frac{\sqrt{ex+db}\sqrt{-\frac{bd-ae}{b}}}{bd-ae}\right)}{3(b^4x + ab^3)}$$

```
input integrate((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^2,x, algorithm="fricas")
```


output

```
[1/6*(3*(2*B*a*b*d - (5*B*a^2 - 3*A*a*b)*e + (2*B*b^2*d - (5*B*a*b - 3*A*b^2)*e)*x)*sqrt((b*d - a*e)/b)*log((b*e*x + 2*b*d - a*e - 2*sqrt(e*x + d)*b*sqrt((b*d - a*e)/b))/(b*x + a)) + 2*(2*B*b^2*e*x^2 + (11*B*a*b - 3*A*b^2)*d - 3*(5*B*a^2 - 3*A*a*b)*e + 2*(4*B*b^2*d - (5*B*a*b - 3*A*b^2)*e)*x)*sqrt(e*x + d)/(b^4*x + a*b^3), -1/3*(3*(2*B*a*b*d - (5*B*a^2 - 3*A*a*b)*e + (2*B*b^2*d - (5*B*a*b - 3*A*b^2)*e)*x)*sqrt(-(b*d - a*e)/b)*arctan(-sqrt(e*x + d)*b*sqrt(-(b*d - a*e)/b)/(b*d - a*e)) - (2*B*b^2*e*x^2 + (11*B*a*b - 3*A*b^2)*d - 3*(5*B*a^2 - 3*A*a*b)*e + 2*(4*B*b^2*d - (5*B*a*b - 3*A*b^2)*e)*x)*sqrt(e*x + d)/(b^4*x + a*b^3)]
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^2} dx = \text{Timed out}$$

input

```
integrate((B*x+A)*(e*x+d)**(3/2)/(b*x+a)**2,x)
```

output

Timed out

Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^2} dx = \text{Exception raised: ValueError}$$

input

```
integrate((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^2,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.53

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^2} dx = \frac{(2Bb^2d^2 - 7Babde + 3Ab^2de + 5Ba^2e^2 - 3Aabe^2) \arctan\left(\frac{\sqrt{ex+db}}{\sqrt{-b^2d+abe}}\right)}{\sqrt{-b^2d+abe}b^3} + \frac{\sqrt{ex+d}Babde - \sqrt{ex+d}Ab^2de - \sqrt{ex+d}Ba^2e^2 + \sqrt{ex+d}Aabe^2}{((ex+d)b - bd + ae)b^3} + \frac{2\left((ex+d)^{\frac{3}{2}}Bb^4 + 3\sqrt{ex+d}Bb^4d - 6\sqrt{ex+d}Bab^3e + 3\sqrt{ex+d}Ab^4e\right)}{3b^6}$$

input `integrate((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^2,x, algorithm="giac")`output `(2*B*b^2*d^2 - 7*B*a*b*d*e + 3*A*b^2*d*e + 5*B*a^2*e^2 - 3*A*a*b*e^2)*arctan(sqrt(e*x + d)*b/sqrt(-b^2*d + a*b*e))/(sqrt(-b^2*d + a*b*e)*b^3) + (sqrt(e*x + d)*B*a*b*d*e - sqrt(e*x + d)*A*b^2*d*e - sqrt(e*x + d)*B*a^2*e^2 + sqrt(e*x + d)*A*a*b*e^2)/(((e*x + d)*b - b*d + a*e)*b^3) + 2/3*((e*x + d)^(3/2)*B*b^4 + 3*sqrt(e*x + d)*B*b^4*d - 6*sqrt(e*x + d)*B*a*b^3*e + 3*sqrt(e*x + d)*A*b^4*e)/b^6`**Mupad [B] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.19

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^2} dx = \left(\frac{2Ae - 2Bd}{b^2} + \frac{2B(2b^2d - 2abe)}{b^4}\right) \sqrt{d + ex} - \frac{\sqrt{d + ex}(Ba^2e^2 - Aabe^2 - Bdabe + Adb^2e)}{b^4(d + ex) - b^4d + ab^3e} + \frac{2B(d + ex)^{3/2}}{3b^2} + \frac{\operatorname{atan}\left(\frac{\sqrt{b}\sqrt{d+ex}i}{\sqrt{bd-ae}}\right) \sqrt{bd-ae}(3Abe - 5Bae + 2Bbd) \operatorname{li}}{b^{7/2}}$$

input `int(((A + B*x)*(d + e*x)^(3/2))/(a + b*x)^2,x)`

output

```
((2*A*e - 2*B*d)/b^2 + (2*B*(2*b^2*d - 2*a*b*e))/b^4)*(d + e*x)^(1/2) - ((
d + e*x)^(1/2)*(B*a^2*e^2 - A*a*b*e^2 + A*b^2*d*e - B*a*b*d*e))/(b^4*(d +
e*x) - b^4*d + a*b^3*e) + (2*B*(d + e*x)^(3/2))/(3*b^2) + (atan((b^(1/2))*
(d + e*x)^(1/2)*1i)/(b*d - a*e)^(1/2))*(b*d - a*e)^(1/2)*(3*A*b*e - 5*B*a*e
+ 2*B*b*d)*1i)/b^(7/2)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.82

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^2} dx = \frac{2\sqrt{b}\sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b}\sqrt{ae-bd}}\right) ae - 2\sqrt{b}\sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b}\sqrt{ae-bd}}\right) bd - 2\sqrt{b}\sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b}\sqrt{ae-bd}}\right) b^2}{b^3}$$

input

```
int((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^2,x)
```

output

```
(2*(3*sqrt(b)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b
*d)))*a*e - 3*sqrt(b)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt
(a*e - b*d)))*b*d - 3*sqrt(d + e*x)*a*b*e + 4*sqrt(d + e*x)*b**2*d + sqrt(
d + e*x)*b**2*e*x))/(3*b**3)
```

3.159 $\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^2} dx$

Optimal result	1463
Mathematica [A] (verified)	1463
Rubi [A] (verified)	1464
Maple [A] (verified)	1466
Fricas [B] (verification not implemented)	1466
Sympy [F]	1467
Maxima [F(-2)]	1467
Giac [A] (verification not implemented)	1468
Mupad [B] (verification not implemented)	1468
Reduce [B] (verification not implemented)	1469

Optimal result

Integrand size = 22, antiderivative size = 107

$$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^2} dx = \frac{2B\sqrt{d+ex}}{b^2} - \frac{(Ab-aB)\sqrt{d+ex}}{b^2(a+bx)} - \frac{(2bBd+Abe-3aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{5/2}\sqrt{bd-ae}}$$

output

```
2*B*(e*x+d)^(1/2)/b^2-(A*b-B*a)*(e*x+d)^(1/2)/b^2/(b*x+a)-(A*b*e-3*B*a*e+2
*B*b*d)*arctanh(b^(1/2)*(e*x+d)^(1/2)/(-a*e+b*d)^(1/2))/b^(5/2)/(-a*e+b*d)
^(1/2)
```

Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.90

$$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^2} dx = \frac{(-Ab+3aB+2bBx)\sqrt{d+ex}}{b^2(a+bx)} + \frac{(2bBd+Abe-3aBe)\arctan\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right)}{b^{5/2}\sqrt{-bd+ae}}$$

input `Integrate[((A + B*x)*Sqrt[d + e*x])/(a + b*x)^2,x]`

output `((-(A*b) + 3*a*B + 2*b*B*x)*Sqrt[d + e*x])/(b^2*(a + b*x)) + ((2*b*B*d + A*b*e - 3*a*B*e)*ArcTan[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[-(b*d) + a*e]])/(b^(5/2)*Sqrt[-(b*d) + a*e])`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.25, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {87, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^2} dx \\
 & \quad \downarrow 87 \\
 & \frac{(-3aBe + Abe + 2bBd) \int \frac{\sqrt{d+ex}}{a+bx} dx}{2b(bd - ae)} - \frac{(d + ex)^{3/2}(Ab - aB)}{b(a + bx)(bd - ae)} \\
 & \quad \downarrow 60 \\
 & \frac{(-3aBe + Abe + 2bBd) \left(\frac{(bd-ae) \int \frac{1}{(a+bx)\sqrt{d+ex}} dx}{b} + \frac{2\sqrt{d+ex}}{b} \right)}{2b(bd - ae)} - \frac{(d + ex)^{3/2}(Ab - aB)}{b(a + bx)(bd - ae)} \\
 & \quad \downarrow 73 \\
 & \frac{(-3aBe + Abe + 2bBd) \left(\frac{2(bd-ae) \int \frac{1}{a + \frac{b(d+ex)}{e} - \frac{bd}{e}} d\sqrt{d+ex}}{be} + \frac{2\sqrt{d+ex}}{b} \right)}{2b(bd - ae)} - \frac{(d + ex)^{3/2}(Ab - aB)}{b(a + bx)(bd - ae)} \\
 & \quad \downarrow 221 \\
 & \frac{(-3aBe + Abe + 2bBd) \left(\frac{2\sqrt{d+ex}}{b} - \frac{2\sqrt{bd-ae} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{3/2}} \right)}{2b(bd - ae)} - \frac{(d + ex)^{3/2}(Ab - aB)}{b(a + bx)(bd - ae)}
 \end{aligned}$$

input `Int[((A + B*x)*Sqrt[d + e*x])/(a + b*x)^2,x]`

output `-(((A*b - a*B)*(d + e*x)^(3/2))/(b*(b*d - a*e)*(a + b*x))) + ((2*b*B*d + A*b*e - 3*a*B*e)*((2*Sqrt[d + e*x])/b - (2*Sqrt[b*d - a*e]*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[b*d - a*e]])/b^(3/2)))/(2*b*(b*d - a*e))`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || (EqQ[e, 0] || (EqQ[c, 0] || LtQ[p, n]))))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.96

method	result	size
pseudoelliptic	$\frac{-\sqrt{ex+d} \sqrt{(ae-db)b} ((-2Bx+A)b-3Ba)+(bx+a)((Ae+2Bd)b-3Bae) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{\sqrt{(ae-db)b} b^2 (bx+a)}$	103
risch	$\frac{2B\sqrt{ex+d}}{b^2} + \frac{2\left(-\frac{1}{2}Abe+\frac{1}{2}Bae\right)\sqrt{ex+d}}{(ex+d)b+ae-db} + \frac{(Abe-3Bae+2Bbd) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{b^2 \sqrt{(ae-db)b}}$	107
derivativedivides	$\frac{2B\sqrt{ex+d}}{b^2} + \frac{2\left(-\frac{1}{2}Abe+\frac{1}{2}Bae\right)\sqrt{ex+d}}{(ex+d)b+ae-db} + \frac{(Abe-3Bae+2Bbd) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{b^2 \sqrt{(ae-db)b}}$	108
default	$\frac{2B\sqrt{ex+d}}{b^2} + \frac{2\left(-\frac{1}{2}Abe+\frac{1}{2}Bae\right)\sqrt{ex+d}}{(ex+d)b+ae-db} + \frac{(Abe-3Bae+2Bbd) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{b^2 \sqrt{(ae-db)b}}$	108

input `int((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^2,x,method=_RETURNVERBOSE)`

output
$$\frac{-(e*x+d)^{(1/2)}*((a*e-b*d)*b)^{(1/2)}*((-2*B*x+A)*b-3*B*a)+(b*x+a)*((A*e+2*B*d)*b-3*B*a*e)*\arctan(b*(e*x+d)^{(1/2)/((a*e-b*d)*b)^{(1/2))}}{(a*e-b*d)*b)^{(1/2)}/b^2/(b*x+a)}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 189 vs. 2(92) = 184.

Time = 0.13 (sec) , antiderivative size = 393, normalized size of antiderivative = 3.67

$$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^2} dx = \left[\frac{(2Babd - (3Ba^2 - Aab)e + (2Bb^2d - (3Bab - Ab^2)e)x)\sqrt{b^2d - abe} \log\left(\frac{bex+2bd-ae-2\sqrt{b^2d-abe}\sqrt{ex+d}}{bx+a}\right)}{2(ab^4d - a^2b^3e + (b^5d - ab^4e)}$$

input `integrate((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^2,x, algorithm="fricas")`

output

```
[1/2*((2*B*a*b*d - (3*B*a^2 - A*a*b)*e + (2*B*b^2*d - (3*B*a*b - A*b^2)*e)
*x)*sqrt(b^2*d - a*b*e)*log((b*e*x + 2*b*d - a*e - 2*sqrt(b^2*d - a*b*e)*s
qrt(e*x + d))/(b*x + a)) + 2*((3*B*a*b^2 - A*b^3)*d - (3*B*a^2*b - A*a*b^2
)*e + 2*(B*b^3*d - B*a*b^2*e)*x)*sqrt(e*x + d)/(a*b^4*d - a^2*b^3*e + (b^
5*d - a*b^4*e)*x), ((2*B*a*b*d - (3*B*a^2 - A*a*b)*e + (2*B*b^2*d - (3*B*a
*b - A*b^2)*e)*x)*sqrt(-b^2*d + a*b*e)*arctan(sqrt(-b^2*d + a*b*e)*sqrt(e*
x + d)/(b*e*x + b*d)) + ((3*B*a*b^2 - A*b^3)*d - (3*B*a^2*b - A*a*b^2)*e +
2*(B*b^3*d - B*a*b^2*e)*x)*sqrt(e*x + d)/(a*b^4*d - a^2*b^3*e + (b^5*d -
a*b^4*e)*x)]
```

Sympy [F]

$$\int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^2} dx = \int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^2} dx$$

input

```
integrate((B*x+A)*(e*x+d)**(1/2)/(b*x+a)**2,x)
```

output

```
Integral((A + B*x)*sqrt(d + e*x)/(a + b*x)**2, x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^2} dx = \text{Exception raised: ValueError}$$

input

```
integrate((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^2,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for m
ore detail
```


Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.07

$$\int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^2} dx = \frac{2\sqrt{ex + d}B}{b^2} + \frac{(2Bbd - 3Bae + Abe) \arctan\left(\frac{\sqrt{ex+db}}{\sqrt{-b^2d+abe}}\right)}{\sqrt{-b^2d+abe}b^2} + \frac{\sqrt{ex + d}Bae - \sqrt{ex + d}Abe}{((ex + d)b - bd + ae)b^2}$$

input `integrate((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^2,x, algorithm="giac")`output `2*sqrt(e*x + d)*B/b^2 + (2*B*b*d - 3*B*a*e + A*b*e)*arctan(sqrt(e*x + d)*b/sqrt(-b^2*d + a*b*e))/(sqrt(-b^2*d + a*b*e)*b^2) + (sqrt(e*x + d)*B*a*e - sqrt(e*x + d)*A*b*e)/(((e*x + d)*b - b*d + a*e)*b^2)`**Mupad [B] (verification not implemented)**

Time = 0.98 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.01

$$\int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^2} dx = \frac{2B\sqrt{d + ex}}{b^2} - \frac{(Abe - Bae)\sqrt{d + ex}}{b^3(d + ex) - b^3d + ab^2e} + \frac{\operatorname{atan}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{ae-bd}}\right)(Abe - 3Bae + 2Bbd)}{b^{5/2}\sqrt{ae - bd}}$$

input `int(((A + B*x)*(d + e*x)^(1/2))/(a + b*x)^2,x)`output `(2*B*(d + e*x)^(1/2))/b^2 - ((A*b*e - B*a*e)*(d + e*x)^(1/2))/(b^3*(d + e*x) - b^3*d + a*b^2*e) + (atan((b^(1/2)*(d + e*x)^(1/2))/(a*e - b*d)^(1/2)))*(A*b*e - 3*B*a*e + 2*B*b*d)/(b^(5/2)*(a*e - b*d)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.48

$$\int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^2} dx = \frac{-2\sqrt{b}\sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+d}b}{\sqrt{b}\sqrt{ae-bd}}\right) + 2\sqrt{ex+d}b}{b^2}$$

input `int((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^2,x)`

output `(2*(-sqrt(b)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d))) + sqrt(d + e*x)*b)/b**2`

3.160 $\int \frac{A+Bx}{(a+bx)^2\sqrt{d+ex}} dx$

Optimal result	1470
Mathematica [A] (verified)	1470
Rubi [A] (verified)	1471
Maple [A] (verified)	1472
Fricas [B] (verification not implemented)	1473
Sympy [F(-1)]	1473
Maxima [F(-2)]	1474
Giac [A] (verification not implemented)	1474
Mupad [B] (verification not implemented)	1475
Reduce [B] (verification not implemented)	1475

Optimal result

Integrand size = 22, antiderivative size = 103

$$\int \frac{A+Bx}{(a+bx)^2\sqrt{d+ex}} dx = -\frac{(Ab-aB)\sqrt{d+ex}}{b(bd-ae)(a+bx)} - \frac{(2bBd-Abe-aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{3/2}(bd-ae)^{3/2}}$$

output

$-(A*b-B*a)*(e*x+d)^{(1/2)}/b/(-a*e+b*d)/(b*x+a)-(-A*b*e-B*a*e+2*B*b*d)*\operatorname{arctanh}(b^{(1/2)}*(e*x+d)^{(1/2)}/(-a*e+b*d)^{(1/2)})/b^{(3/2)}/(-a*e+b*d)^{(3/2)}$

Mathematica [A] (verified)

Time = 0.31 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.99

$$\int \frac{A+Bx}{(a+bx)^2\sqrt{d+ex}} dx = \frac{(-Ab+aB)\sqrt{d+ex}}{b(bd-ae)(a+bx)} - \frac{(2bBd-Abe-aBe)\operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right)}{b^{3/2}(-bd+ae)^{3/2}}$$

input

`Integrate[(A + B*x)/((a + b*x)^2*sqrt[d + e*x]),x]`

output

$$\left((-A*b + a*B)*\text{Sqrt}[d + e*x] / (b*(b*d - a*e)*(a + b*x)) - ((2*b*B*d - A*b*e - a*B*e)*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[d + e*x]) / \text{Sqrt}[-(b*d) + a*e]] / (b^{3/2} * (-(b*d) + a*e)^{3/2})) \right)$$
Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.136$, Rules used = {87, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^2 \sqrt{d + ex}} dx$$

$$\downarrow 87$$

$$\frac{(-aBe - Abe + 2bBd) \int \frac{1}{(a+bx)\sqrt{d+ex}} dx}{2b(bd - ae)} - \frac{\sqrt{d + ex}(Ab - aB)}{b(a + bx)(bd - ae)}$$

$$\downarrow 73$$

$$\frac{(-aBe - Abe + 2bBd) \int \frac{1}{a + \frac{b(d+ex)}{e} - \frac{bd}{e}} d\sqrt{d + ex}}{be(bd - ae)} - \frac{\sqrt{d + ex}(Ab - aB)}{b(a + bx)(bd - ae)}$$

$$\downarrow 221$$

$$-\frac{(-aBe - Abe + 2bBd)\text{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{3/2}(bd - ae)^{3/2}} - \frac{\sqrt{d + ex}(Ab - aB)}{b(a + bx)(bd - ae)}$$

input

$$\text{Int}[(A + B*x)/((a + b*x)^2*\text{Sqrt}[d + e*x]), x]$$

output

$$-\left(((A*b - a*B)*\text{Sqrt}[d + e*x] / (b*(b*d - a*e)*(a + b*x))) - ((2*b*B*d - A*b*e - a*B*e)*\text{ArcTanh}[(\text{Sqrt}[b]*\text{Sqrt}[d + e*x]) / \text{Sqrt}[b*d - a*e]] / (b^{3/2} * (b*d - a*e)^{3/2})) \right)$$

Definitions of rubi rules used

rule 73 $\text{Int}[(a_.) + (b_.)(x_)^m)((c_.) + (d_.)(x_)^n], x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \text{ Subst}[\text{Int}[x^{p(m+1)-1}(c - a(d/b) + d(x^p/b))^n], x], x, (a + b*x)^{(1/p)}, x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 87 $\text{Int}[(a_.) + (b_.)(x_)*((c_.) + (d_.)(x_)^{n_})*((e_.) + (f_.)(x_)^p), x_] \rightarrow \text{Simp}[(-b*e - a*f)*(c + d*x)^{(n+1)}*((e + f*x)^{(p+1)})/(f*(p+1)*(c*f - d*e)), x] - \text{Simp}[(a*d*f*(n+p+2) - b*(d*e*(n+1) + c*f*(p+1))]/(f*(p+1)*(c*f - d*e)) \text{Int}[(c + d*x)^n*(e + f*x)^{(p+1)}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n\}, x] \&\& \text{LtQ}[p, -1] \&\& (!\text{LtQ}[n, -1] || \text{IntegerQ}[p] || !(\text{IntegerQ}[n] || !(\text{EqQ}[e, 0] || !(\text{EqQ}[c, 0] || \text{LtQ}[p, n]))))$

rule 221 $\text{Int}[(a_.) + (b_.)(x_)^2]^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{NegQ}[a/b]$

Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.85

method	result	size
pseudoelliptic	$\frac{(Ab-Ba)\sqrt{ex+d}}{bx+a} + \frac{(Abe+Bae-2Bbd) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{b(ae-db)\sqrt{(ae-db)b}}$	88
derivativedivides	$\frac{e(Ab-Ba)\sqrt{ex+d}}{b(ae-db)((ex+d)b+ae-db)} + \frac{(Abe+Bae-2Bbd) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{b(ae-db)\sqrt{(ae-db)b}}$	111
default	$\frac{e(Ab-Ba)\sqrt{ex+d}}{b(ae-db)((ex+d)b+ae-db)} + \frac{(Abe+Bae-2Bbd) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{b(ae-db)\sqrt{(ae-db)b}}$	111

input $\text{int}((B*x+A)/(b*x+a)^2/(e*x+d)^{(1/2)}, x, \text{method}=_RETURNVERBOSE)$

output $1/b/(a*e-b*d)*((A*b-B*a)*(e*x+d)^{(1/2)}/(b*x+a)+(A*b*e+B*a*e-2*B*b*d)/((a*e-b*d)*b)^{(1/2)}*\arctan(b*(e*x+d)^{(1/2)}/((a*e-b*d)*b)^{(1/2}))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 191 vs. $2(90) = 180$.

Time = 0.10 (sec) , antiderivative size = 397, normalized size of antiderivative = 3.85

$$\int \frac{A + Bx}{(a + bx)^2 \sqrt{d + ex}} dx$$

$$= \left[\frac{(2 Babd - (Ba^2 + Aab)e + (2 Bb^2d - (Bab + Ab^2)e)x) \sqrt{b^2d - abe} \log\left(\frac{be x + 2bd - ae - 2\sqrt{b^2d - abe}\sqrt{ex + d}}{bx + a}\right) +}{2(ab^4d^2 - 2a^2b^3de + a^3b^2e^2 + (b^5d^2 - 2ab^4de + a^2b^3e^2)x)} \right]$$

input `integrate((B*x+A)/(b*x+a)^2/(e*x+d)^(1/2),x, algorithm="fricas")`

output `[1/2*((2*B*a*b*d - (B*a^2 + A*a*b)*e + (2*B*b^2*d - (B*a*b + A*b^2)*e)*x)*sqrt(b^2*d - a*b*e)*log((b*e*x + 2*b*d - a*e - 2*sqrt(b^2*d - a*b*e)*sqrt(e*x + d))/(b*x + a)) + 2*((B*a*b^2 - A*b^3)*d - (B*a^2*b - A*a*b^2)*e)*sqrt(e*x + d)/(a*b^4*d^2 - 2*a^2*b^3*d*e + a^3*b^2*e^2 + (b^5*d^2 - 2*a*b^4*d*e + a^2*b^3*e^2)*x), ((2*B*a*b*d - (B*a^2 + A*a*b)*e + (2*B*b^2*d - (B*a*b + A*b^2)*e)*x)*sqrt(-b^2*d + a*b*e)*arctan(sqrt(-b^2*d + a*b*e)*sqrt(e*x + d)/(b*e*x + b*d)) + ((B*a*b^2 - A*b^3)*d - (B*a^2*b - A*a*b^2)*e)*sqrt(e*x + d)/(a*b^4*d^2 - 2*a^2*b^3*d*e + a^3*b^2*e^2 + (b^5*d^2 - 2*a*b^4*d*e + a^2*b^3*e^2)*x)]`

Sympy [F(-1)]

Timed out.

$$\int \frac{A + Bx}{(a + bx)^2 \sqrt{d + ex}} dx = \text{Timed out}$$

input `integrate((B*x+A)/(b*x+a)**2/(e*x+d)**(1/2),x)`

output `Timed out`

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{(a + bx)^2 \sqrt{d + ex}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^2/(e*x+d)^(1/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.18

$$\int \frac{A + Bx}{(a + bx)^2 \sqrt{d + ex}} dx = \frac{(2Bbd - Bae - Abe) \arctan\left(\frac{\sqrt{ex+db}}{\sqrt{-b^2d+abe}}\right)}{(b^2d - abe)\sqrt{-b^2d + abe}} + \frac{\sqrt{ex + d}Bae - \sqrt{ex + d}Abe}{(b^2d - abe)((ex + d)b - bd + ae)}$$

input `integrate((B*x+A)/(b*x+a)^2/(e*x+d)^(1/2),x, algorithm="giac")`

output `(2*B*b*d - B*a*e - A*b*e)*arctan(sqrt(e*x + d)*b/sqrt(-b^2*d + a*b*e))/((b^2*d - a*b*e)*sqrt(-b^2*d + a*b*e)) + (sqrt(e*x + d)*B*a*e - sqrt(e*x + d)*A*b*e)/((b^2*d - a*b*e)*((e*x + d)*b - b*d + a*e))`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.96

$$\int \frac{A + Bx}{(a + bx)^2 \sqrt{d + ex}} dx = \frac{\operatorname{atan}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{ae-bd}}\right) (Abe + Bae - 2Bbd)}{b^{3/2} (ae - bd)^{3/2}} + \frac{(Abe - Bae) \sqrt{d + ex}}{b (ae - bd) (ae - bd + b(d + ex))}$$

input `int((A + B*x)/((a + b*x)^2*(d + e*x)^(1/2)),x)`output `(atan((b^(1/2)*(d + e*x)^(1/2))/(a*e - b*d)^(1/2))*(A*b*e + B*a*e - 2*B*b*d))/(b^(3/2)*(a*e - b*d)^(3/2)) + ((A*b*e - B*a*e)*(d + e*x)^(1/2))/(b*(a*e - b*d)*(a*e - b*d + b*(d + e*x)))`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.49

$$\int \frac{A + Bx}{(a + bx)^2 \sqrt{d + ex}} dx = \frac{2\sqrt{b}\sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b}\sqrt{ae-bd}}\right)}{b(ae - bd)}$$

input `int((B*x+A)/(b*x+a)^2/(e*x+d)^(1/2),x)`output `(2*sqrt(b)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))/(b*(a*e - b*d))`

3.161 $\int \frac{A+Bx}{(a+bx)^2(d+ex)^{3/2}} dx$

Optimal result	1476
Mathematica [A] (verified)	1476
Rubi [A] (verified)	1477
Maple [A] (verified)	1479
Fricas [B] (verification not implemented)	1479
Sympy [F(-1)]	1480
Maxima [F(-2)]	1480
Giac [A] (verification not implemented)	1481
Mupad [B] (verification not implemented)	1481
Reduce [B] (verification not implemented)	1482

Optimal result

Integrand size = 22, antiderivative size = 128

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{3/2}} dx = \frac{2(Bd - Ae)}{(bd - ae)^2\sqrt{d + ex}} - \frac{(Ab - aB)\sqrt{d + ex}}{(bd - ae)^2(a + bx)} - \frac{(2bBd - 3Abe + aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{\sqrt{b}(bd - ae)^{5/2}}$$

output

```
2*(-A*e+B*d)/(-a*e+b*d)^2/(e*x+d)^(1/2)-(A*b-B*a)*(e*x+d)^(1/2)/(-a*e+b*d)
^2/(b*x+a)-(-3*A*b*e+B*a*e+2*B*b*d)*arctanh(b^(1/2)*(e*x+d)^(1/2)/(-a*e+b*
d)^(1/2))/b^(1/2)/(-a*e+b*d)^(5/2)
```

Mathematica [A] (verified)

Time = 0.40 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.95

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{3/2}} dx = \frac{B(3ad + 2bdx + aex) - A(2ae + b(d + 3ex))}{(bd - ae)^2(a + bx)\sqrt{d + ex}} + \frac{(2bBd - 3Abe + aBe) \arctan\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right)}{\sqrt{b}(-bd + ae)^{5/2}}$$

input `Integrate[(A + B*x)/((a + b*x)^2*(d + e*x)^(3/2)),x]`

output
$$\frac{(B(3a*d + 2b*d*x + a*e*x) - A(2a*e + b(d + 3e*x)))/((b*d - a*e)^2(a + b*x)*\text{Sqrt}[d + e*x]) + ((2b*B*d - 3A*b*e + a*B*e)*\text{ArcTan}[\text{Sqrt}[b]*\text{Sqrt}[d + e*x])/\text{Sqrt}[-(b*d) + a*e])}{(\text{Sqrt}[b]*(-(b*d) + a*e)^(5/2))}$$

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.10, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {87, 61, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + Bx}{(a + bx)^2(d + ex)^{3/2}} dx \\ & \quad \downarrow 87 \\ & \frac{(aBe - 3Abe + 2bBd) \int \frac{1}{(a+bx)(d+ex)^{3/2}} dx}{2b(bd - ae)} - \frac{Ab - aB}{b(a + bx)\sqrt{d + ex}(bd - ae)} \\ & \quad \downarrow 61 \\ & \frac{(aBe - 3Abe + 2bBd) \left(\frac{b \int \frac{1}{(a+bx)\sqrt{d+ex}} dx}{bd - ae} + \frac{2}{\sqrt{d+ex}(bd - ae)} \right)}{2b(bd - ae)} - \frac{Ab - aB}{b(a + bx)\sqrt{d + ex}(bd - ae)} \\ & \quad \downarrow 73 \\ & \frac{(aBe - 3Abe + 2bBd) \left(\frac{2b \int \frac{1}{a + \frac{b(d+ex) - bd}{e}} d\sqrt{d+ex}}{e(bd - ae)} + \frac{2}{\sqrt{d+ex}(bd - ae)} \right)}{2b(bd - ae)} - \frac{Ab - aB}{b(a + bx)\sqrt{d + ex}(bd - ae)} \\ & \quad \downarrow 221 \\ & \frac{(aBe - 3Abe + 2bBd) \left(\frac{2}{\sqrt{d+ex}(bd - ae)} - \frac{2\sqrt{b} \text{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd - ae}}\right)}{(bd - ae)^{3/2}} \right)}{2b(bd - ae)} - \frac{Ab - aB}{b(a + bx)\sqrt{d + ex}(bd - ae)} \end{aligned}$$

input `Int[(A + B*x)/((a + b*x)^2*(d + e*x)^(3/2)),x]`

output `-((A*b - a*B)/(b*(b*d - a*e)*(a + b*x)*Sqrt[d + e*x])) + ((2*b*B*d - 3*A*b*e + a*B*e)*(2/((b*d - a*e)*Sqrt[d + e*x]) - (2*Sqrt[b]*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[b*d - a*e]])/(b*d - a*e)^(3/2)))/(2*b*(b*d - a*e))`

Defintions of rubi rules used

rule 61 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.02

method	result
derivativedivides	$2 \left(\frac{\left(\frac{1}{2}Abe - \frac{1}{2}Bae\right)\sqrt{ex+d}}{(ex+d)b+ae-db} + \frac{(3Abe - Bae - 2Bbd) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{2\sqrt{(ae-db)b}} \right) - \frac{2(Ae-Bd)}{(ae-db)^2\sqrt{ex+d}}$
default	$2 \left(\frac{\left(\frac{1}{2}Abe - \frac{1}{2}Bae\right)\sqrt{ex+d}}{(ex+d)b+ae-db} + \frac{(3Abe - Bae - 2Bbd) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{2\sqrt{(ae-db)b}} \right) - \frac{2(Ae-Bd)}{(ae-db)^2\sqrt{ex+d}}$
pseudoelliptic	$2 \left(\frac{3(bx+a)\left(\left(Ae - \frac{2Bd}{3}\right)b - \frac{Bae}{3}\right)\sqrt{ex+d} \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{2} + \sqrt{(ae-db)b} \left(\frac{(3Aex+d(-2Bx+A))b}{2} + \left(-\frac{Bx}{2} + A\right)e - \frac{3B}{2} \right) \right) - \frac{\sqrt{ex+d} \sqrt{(ae-db)b} (bx+a)(ae-db)^2}{\sqrt{ex+d} \sqrt{(ae-db)b} (bx+a)(ae-db)^2}$

input `int((B*x+A)/(b*x+a)^2/(e*x+d)^(3/2),x,method=_RETURNVERBOSE)`

output
$$-2/(a*e-b*d)^2*((1/2*A*b*e-1/2*B*a*e)*(e*x+d)^(1/2)/((e*x+d)*b+a*e-d*b)+1/2*(3*A*b*e-B*a*e-2*B*b*d)/((a*e-b*d)*b)^(1/2)*\arctan(b*(e*x+d)^(1/2)/((a*e-b*d)*b)^(1/2))-2*(A*e-B*d)/(a*e-b*d)^2/(e*x+d)^(1/2)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 380 vs. 2(113) = 226.

Time = 0.13 (sec) , antiderivative size = 775, normalized size of antiderivative = 6.05

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{3/2}} dx = \text{Too large to display}$$

input `integrate((B*x+A)/(b*x+a)^2/(e*x+d)^(3/2),x, algorithm="fricas")`

output

```
[-1/2*((2*B*a*b*d^2 + (B*a^2 - 3*A*a*b)*d*e + (2*B*b^2*d*e + (B*a*b - 3*A*
b^2)*e^2)*x^2 + (2*B*b^2*d^2 + 3*(B*a*b - A*b^2)*d*e + (B*a^2 - 3*A*a*b)*e
^2)*x)*sqrt(b^2*d - a*b*e)*log((b*e*x + 2*b*d - a*e + 2*sqrt(b^2*d - a*b*e
)*sqrt(e*x + d))/(b*x + a)) - 2*(2*A*a^2*b*e^2 + (3*B*a*b^2 - A*b^3)*d^2 -
(3*B*a^2*b + A*a*b^2)*d*e + (2*B*b^3*d^2 - (B*a*b^2 + 3*A*b^3)*d*e - (B*a
^2*b - 3*A*a*b^2)*e^2)*x)*sqrt(e*x + d))/(a*b^4*d^4 - 3*a^2*b^3*d^3*e + 3*
a^3*b^2*d^2*e^2 - a^4*b*d*e^3 + (b^5*d^3*e - 3*a*b^4*d^2*e^2 + 3*a^2*b^3*d
*e^3 - a^3*b^2*e^4)*x^2 + (b^5*d^4 - 2*a*b^4*d^3*e + 2*a^3*b^2*d*e^3 - a^4
*b*e^4)*x), ((2*B*a*b*d^2 + (B*a^2 - 3*A*a*b)*d*e + (2*B*b^2*d*e + (B*a*b
- 3*A*b^2)*e^2)*x^2 + (2*B*b^2*d^2 + 3*(B*a*b - A*b^2)*d*e + (B*a^2 - 3*A*
a*b)*e^2)*x)*sqrt(-b^2*d + a*b*e)*arctan(sqrt(-b^2*d + a*b*e)*sqrt(e*x + d
))/(b*e*x + b*d)) + (2*A*a^2*b*e^2 + (3*B*a*b^2 - A*b^3)*d^2 - (3*B*a^2*b +
A*a*b^2)*d*e + (2*B*b^3*d^2 - (B*a*b^2 + 3*A*b^3)*d*e - (B*a^2*b - 3*A*a*
b^2)*e^2)*x)*sqrt(e*x + d))/(a*b^4*d^4 - 3*a^2*b^3*d^3*e + 3*a^3*b^2*d^2*e
^2 - a^4*b*d*e^3 + (b^5*d^3*e - 3*a*b^4*d^2*e^2 + 3*a^2*b^3*d*e^3 - a^3*b^
2*e^4)*x^2 + (b^5*d^4 - 2*a*b^4*d^3*e + 2*a^3*b^2*d*e^3 - a^4*b*e^4)*x]]
```

Sympy [F(-1)]

Timed out.

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{3/2}} dx = \text{Timed out}$$

input

```
integrate((B*x+A)/(b*x+a)**2/(e*x+d)**(3/2),x)
```

output

Timed out

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{3/2}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((B*x+A)/(b*x+a)^2/(e*x+d)^(3/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for m
ore detail
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.48

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{3/2}} dx = \frac{(2Bbd + Bae - 3Abe) \arctan\left(\frac{\sqrt{ex+db}}{\sqrt{-b^2d+abe}}\right)}{(b^2d^2 - 2abde + a^2e^2)\sqrt{-b^2d+abe}} + \frac{2(ex+d)Bbd - 2Bbd^2 + (ex+d)Bae - 3(ex+d)Abe + 2Bade + 2Abde - 2Aae^2}{(b^2d^2 - 2abde + a^2e^2)\left((ex+d)^{\frac{3}{2}}b - \sqrt{ex+dbd} + \sqrt{ex+dae}\right)}$$

input

```
integrate((B*x+A)/(b*x+a)^2/(e*x+d)^(3/2),x, algorithm="giac")
```

output

```
(2*B*b*d + B*a*e - 3*A*b*e)*arctan(sqrt(e*x + d)*b/sqrt(-b^2*d + a*b*e))/(
(b^2*d^2 - 2*a*b*d*e + a^2*e^2)*sqrt(-b^2*d + a*b*e)) + (2*(e*x + d)*B*b*d
- 2*B*b*d^2 + (e*x + d)*B*a*e - 3*(e*x + d)*A*b*e + 2*B*a*d*e + 2*A*b*d*e
- 2*A*a*e^2)/((b^2*d^2 - 2*a*b*d*e + a^2*e^2)*((e*x + d)^(3/2)*b - sqrt(e
*x + d)*b*d + sqrt(e*x + d)*a*e))
```

Mupad [B] (verification not implemented)

Time = 1.03 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.22

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{3/2}} dx = \frac{\operatorname{atan}\left(\frac{\sqrt{b}\sqrt{d+ex}(a^2e^2-2abde+b^2d^2)}{(ae-bd)^{5/2}}\right)(Bae-3Abe+2Bbd)}{\sqrt{b}(ae-bd)^{5/2}} - \frac{\frac{2(Ae-Bd)}{ae-bd} - \frac{(d+ex)(Bae-3Abe+2Bbd)}{(ae-bd)^2}}{b(d+ex)^{3/2} + (ae-bd)\sqrt{d+ex}}$$

input

```
int((A + B*x)/((a + b*x)^2*(d + e*x)^(3/2)),x)
```

output

```
(atan((b^(1/2)*(d + e*x)^(1/2)*(a^2*e^2 + b^2*d^2 - 2*a*b*d*e))/(a*e - b*d)^(5/2))*(B*a*e - 3*A*b*e + 2*B*b*d))/(b^(1/2)*(a*e - b*d)^(5/2)) - ((2*(A*e - B*d))/(a*e - b*d) - ((d + e*x)*(B*a*e - 3*A*b*e + 2*B*b*d))/(a*e - b*d)^2)/(b*(d + e*x)^(3/2) + (a*e - b*d)*(d + e*x)^(1/2))
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.66

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{3/2}} dx = \frac{-2\sqrt{b}\sqrt{ex + d}\sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex + d}b}{\sqrt{b}\sqrt{ae - bd}}\right) - 2ae + 2bd}{\sqrt{ex + d}(a^2e^2 - 2abde + b^2d^2)}$$

input

```
int((B*x+A)/(b*x+a)^2/(e*x+d)^(3/2),x)
```

output

```
(2*( - sqrt(b)*sqrt(d + e*x)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d))) - a*e + b*d))/(sqrt(d + e*x)*(a**2*e**2 - 2*a*b*d*e + b**2*d**2))
```

3.162 $\int \frac{A+Bx}{(a+bx)^2(d+ex)^{5/2}} dx$

Optimal result	1483
Mathematica [A] (verified)	1483
Rubi [A] (verified)	1484
Maple [A] (verified)	1486
Fricas [B] (verification not implemented)	1487
Sympy [F(-1)]	1488
Maxima [F(-2)]	1488
Giac [A] (verification not implemented)	1488
Mupad [B] (verification not implemented)	1489
Reduce [B] (verification not implemented)	1490

Optimal result

Integrand size = 22, antiderivative size = 167

$$\int \frac{A+Bx}{(a+bx)^2(d+ex)^{5/2}} dx = \frac{2(Bd - Ae)}{3(bd - ae)^2(d+ex)^{3/2}} + \frac{2(bBd - 2Abe + aBe)}{(bd - ae)^3\sqrt{d+ex}} - \frac{b(Ab - aB)\sqrt{d+ex}}{(bd - ae)^3(a+bx)} - \frac{\sqrt{b}(2bBd - 5Abe + 3aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{(bd - ae)^{7/2}}$$

output

```
2/3*(-A*e+B*d)/(-a*e+b*d)^2/(e*x+d)^(3/2)+2*(-2*A*b*e+B*a*e+B*b*d)/(-a*e+b*d)^3/(e*x+d)^(1/2)-b*(A*b-B*a)*(e*x+d)^(1/2)/(-a*e+b*d)^3/(b*x+a)-b^(1/2)*(-5*A*b*e+3*B*a*e+2*B*b*d)*arctanh(b^(1/2)*(e*x+d)^(1/2)/(-a*e+b*d)^(1/2))/(-a*e+b*d)^(7/2)
```

Mathematica [A] (verified)

Time = 0.55 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.18

$$\int \frac{A+Bx}{(a+bx)^2(d+ex)^{5/2}} dx = \frac{B(2a^2e(2d+3ex) + 2b^2dx(4d+3ex) + ab(11d^2 + 16dex + 9e^2x^2)) - A(-2bd - 2a^2)}{3(bd - ae)^3(a+bx)(d+ex)} - \frac{\sqrt{b}(2bBd - 5Abe + 3aBe)\arctan\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right)}{(-bd + ae)^{7/2}}$$

input `Integrate[(A + B*x)/((a + b*x)^2*(d + e*x)^(5/2)),x]`

output
$$\frac{(B*(2*a^2*e*(2*d + 3*e*x) + 2*b^2*d*x*(4*d + 3*e*x) + a*b*(11*d^2 + 16*d*e*x + 9*e^2*x^2)) - A*(-2*a^2*e^2 + 2*a*b*e*(7*d + 5*e*x) + b^2*(3*d^2 + 20*d*e*x + 15*e^2*x^2))}{3*(b*d - a*e)^3*(a + b*x)*(d + e*x)^{3/2}} - \frac{(\text{Sqrt}[b]*(2*b*B*d - 5*A*b*e + 3*a*B*e)*\text{ArcTan}[\frac{\text{Sqrt}[b]*\text{Sqrt}[d + e*x]}{\text{Sqrt}[-(b*d + a*e)]}]}{(-b*d + a*e)^{7/2}}$$

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.07, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.227$, Rules used = {87, 61, 61, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{5/2}} dx$$

$$\downarrow 87$$

$$\frac{(3aBe - 5Abe + 2bBd) \int \frac{1}{(a+bx)(d+ex)^{5/2}} dx}{2b(bd - ae)} - \frac{Ab - aB}{b(a + bx)(d + ex)^{3/2}(bd - ae)}$$

$$\downarrow 61$$

$$\frac{(3aBe - 5Abe + 2bBd) \left(\frac{b \int \frac{1}{(a+bx)(d+ex)^{3/2}} dx}{bd - ae} + \frac{2}{3(d+ex)^{3/2}(bd - ae)} \right)}{2b(bd - ae)} - \frac{Ab - aB}{b(a + bx)(d + ex)^{3/2}(bd - ae)}$$

$$\downarrow 61$$

$$\frac{(3aBe - 5Abe + 2bBd) \left(\frac{b \left(\frac{b \int \frac{1}{(a+bx)\sqrt{d+ex}} dx}{bd - ae} + \frac{2}{\sqrt{d+ex}(bd - ae)} \right)}{bd - ae} + \frac{2}{3(d+ex)^{3/2}(bd - ae)} \right)}{2b(bd - ae)} - \frac{Ab - aB}{b(a + bx)(d + ex)^{3/2}(bd - ae)}$$

$$\begin{array}{c}
 \downarrow 73 \\
 (3aBe - 5Abe + 2bBd) \left(\frac{b \left(\frac{2b \int \frac{1}{a + \frac{b(d+ex) - bd}{e(bd-ae)}} d\sqrt{d+ex}}{e(bd-ae)} + \frac{2}{\sqrt{d+ex}(bd-ae)} \right)}{bd-ae} + \frac{2}{3(d+ex)^{3/2}(bd-ae)} \right) \\
 \hline
 \frac{2b(bd-ae)}{Ab - aB} \\
 \frac{2b(bd-ae)}{b(a+bx)(d+ex)^{3/2}(bd-ae)} \\
 \downarrow 221 \\
 (3aBe - 5Abe + 2bBd) \left(\frac{b \left(\frac{2}{\sqrt{d+ex}(bd-ae)} - \frac{2\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{(bd-ae)^{3/2}} \right)}{bd-ae} + \frac{2}{3(d+ex)^{3/2}(bd-ae)} \right) \\
 \hline
 \frac{2b(bd-ae)}{Ab - aB} \\
 \frac{2b(bd-ae)}{b(a+bx)(d+ex)^{3/2}(bd-ae)}
 \end{array}$$

input `Int[(A + B*x)/((a + b*x)^2*(d + e*x)^(5/2)),x]`

output `-((A*b - a*B)/(b*(b*d - a*e)*(a + b*x)*(d + e*x)^(3/2))) + ((2*b*B*d - 5*A*b*e + 3*a*B*e)*(2/(3*(b*d - a*e)*(d + e*x)^(3/2)) + (b*(2/((b*d - a*e)*Sqrt[d + e*x]) - (2*Sqrt[b]*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[b*d - a*e]])/(b*d - a*e)^(3/2)))/(b*d - a*e)))/(2*b*(b*d - a*e))`

Defintions of rubi rules used

rule 61 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`

```
rule 73 Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e)), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Maple [A] (verified)

Time = 0.61 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.98

method	result
derivativedivides	$-\frac{2(Ae-Bd)}{3(ae-db)^2(ex+d)^{\frac{3}{2}}} - \frac{2(-2Abe+BAe+Bbd)}{(ae-db)^3\sqrt{ex+d}} + \frac{2b \left(\frac{(\frac{1}{2}Abe - \frac{1}{2}BAe)\sqrt{ex+d}}{(ex+d)b+ae-db} + \frac{(5Abe-3BAe-2Bbd) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{ae-db}}\right)}{2\sqrt{(ae-db)b}} \right)}{(ae-db)^3}$
default	$-\frac{2(Ae-Bd)}{3(ae-db)^2(ex+d)^{\frac{3}{2}}} - \frac{2(-2Abe+BAe+Bbd)}{(ae-db)^3\sqrt{ex+d}} + \frac{2b \left(\frac{(\frac{1}{2}Abe - \frac{1}{2}BAe)\sqrt{ex+d}}{(ex+d)b+ae-db} + \frac{(5Abe-3BAe-2Bbd) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{ae-db}}\right)}{2\sqrt{(ae-db)b}} \right)}{(ae-db)^3}$
pseudoelliptic	$\frac{5b(bx+a)(ex+d)^{\frac{3}{2}} \left(\left(Ae - \frac{2Bd}{5} \right) b - \frac{3BAe}{5} \right) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{ae-db}} \right) - \frac{2\sqrt{(ae-db)b} \left(\left(-\frac{15Ae^2x^2}{2} - 10\left(-\frac{3Bx}{10} + A\right) xde - \frac{3(-8B^2)}{3} \right)}{(ae-db)^3(ex+d)^{\frac{3}{2}}(bx+a)\sqrt{ae-db}}$

```
input int((B*x+A)/(b*x+a)^2/(e*x+d)^(5/2), x, method=_RETURNVERBOSE)
```

output

```
-2/3*(A*e-B*d)/(a*e-b*d)^2/(e*x+d)^(3/2)-2/(a*e-b*d)^3*(-2*A*b*e+B*a*e+B*b
*d)/(e*x+d)^(1/2)+2/(a*e-b*d)^3*b*((1/2*A*b*e-1/2*B*a*e)*(e*x+d)^(1/2)/((e
*x+d)*b+a*e-d*b)+1/2*(5*A*b*e-3*B*a*e-2*B*b*d)/((a*e-b*d)*b)^(1/2)*arctan(
b*(e*x+d)^(1/2)/((a*e-b*d)*b)^(1/2)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 528 vs. $2(148) = 296$.

Time = 0.16 (sec) , antiderivative size = 1086, normalized size of antiderivative = 6.50

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{5/2}} dx = \text{Too large to display}$$

input

```
integrate((B*x+A)/(b*x+a)^2/(e*x+d)^(5/2),x, algorithm="fricas")
```

output

```
[1/6*(3*(2*B*a*b*d^3 + (3*B*a^2 - 5*A*a*b)*d^2*e + (2*B*b^2*d*e^2 + (3*B*a
*b - 5*A*b^2)*e^3)*x^3 + (4*B*b^2*d^2*e + 2*(4*B*a*b - 5*A*b^2)*d*e^2 + (3
*B*a^2 - 5*A*a*b)*e^3)*x^2 + (2*B*b^2*d^3 + (7*B*a*b - 5*A*b^2)*d^2*e + 2*
(3*B*a^2 - 5*A*a*b)*d*e^2)*x)*sqrt(b/(b*d - a*e))*log((b*e*x + 2*b*d - a*e
- 2*(b*d - a*e)*sqrt(e*x + d)*sqrt(b/(b*d - a*e)))/(b*x + a)) + 2*(2*A*a^
2*e^2 + (11*B*a*b - 3*A*b^2)*d^2 + 2*(2*B*a^2 - 7*A*a*b)*d*e + 3*(2*B*b^2*
d*e + (3*B*a*b - 5*A*b^2)*e^2)*x^2 + 2*(4*B*b^2*d^2 + 2*(4*B*a*b - 5*A*b^2
)*d*e + (3*B*a^2 - 5*A*a*b)*e^2)*x)*sqrt(e*x + d))/(a*b^3*d^5 - 3*a^2*b^2*
d^4*e + 3*a^3*b*d^3*e^2 - a^4*d^2*e^3 + (b^4*d^3*e^2 - 3*a*b^3*d^2*e^3 + 3
*a^2*b^2*d*e^4 - a^3*b*e^5)*x^3 + (2*b^4*d^4*e - 5*a*b^3*d^3*e^2 + 3*a^2*b
^2*d^2*e^3 + a^3*b*d*e^4 - a^4*e^5)*x^2 + (b^4*d^5 - a*b^3*d^4*e - 3*a^2*b
^2*d^3*e^2 + 5*a^3*b*d^2*e^3 - 2*a^4*d*e^4)*x), 1/3*(3*(2*B*a*b*d^3 + (3*B
*a^2 - 5*A*a*b)*d^2*e + (2*B*b^2*d*e^2 + (3*B*a*b - 5*A*b^2)*e^3)*x^3 + (4
*B*b^2*d^2*e + 2*(4*B*a*b - 5*A*b^2)*d*e^2 + (3*B*a^2 - 5*A*a*b)*e^3)*x^2
+ (2*B*b^2*d^3 + (7*B*a*b - 5*A*b^2)*d^2*e + 2*(3*B*a^2 - 5*A*a*b)*d*e^2)*
x)*sqrt(-b/(b*d - a*e))*arctan(sqrt(e*x + d)*sqrt(-b/(b*d - a*e))) + (2*A*
a^2*e^2 + (11*B*a*b - 3*A*b^2)*d^2 + 2*(2*B*a^2 - 7*A*a*b)*d*e + 3*(2*B*b^
2*d*e + (3*B*a*b - 5*A*b^2)*e^2)*x^2 + 2*(4*B*b^2*d^2 + 2*(4*B*a*b - 5*A*b
^2)*d*e + (3*B*a^2 - 5*A*a*b)*e^2)*x)*sqrt(e*x + d))/(a*b^3*d^5 - 3*a^2*b^
2*d^4*e + 3*a^3*b*d^3*e^2 - a^4*d^2*e^3 + (b^4*d^3*e^2 - 3*a*b^3*d^2*e^...
```

Sympy [F(-1)]

Timed out.

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{5/2}} dx = \text{Timed out}$$

input `integrate((B*x+A)/(b*x+a)**2/(e*x+d)**(5/2),x)`

output `Timed out`

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^2/(e*x+d)^(5/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 282, normalized size of antiderivative = 1.69

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{5/2}} dx = \frac{(2Bb^2d + 3Babe - 5Ab^2e) \arctan\left(\frac{\sqrt{ex+db}}{\sqrt{-b^2d+abe}}\right)}{(b^3d^3 - 3ab^2d^2e + 3a^2bde^2 - a^3e^3)\sqrt{-b^2d+abe}} + \frac{\sqrt{ex+d}Babe - \sqrt{ex+d}Ab^2e}{(b^3d^3 - 3ab^2d^2e + 3a^2bde^2 - a^3e^3)((ex+d)b - bd + ae)} + \frac{2(3(ex+d)Bbd + Bbd^2 + 3(ex+d)Bae - 6(ex+d)Abe - Bade - Abde + Aae^2)}{3(b^3d^3 - 3ab^2d^2e + 3a^2bde^2 - a^3e^3)(ex+d)^{\frac{3}{2}}}$$

input `integrate((B*x+A)/(b*x+a)^2/(e*x+d)^(5/2),x, algorithm="giac")`

output
$$\frac{(2*B*b^2*d + 3*B*a*b*e - 5*A*b^2*e)*\arctan(\sqrt{e*x + d}*b/\sqrt{-b^2*d + a*b*e})/((b^3*d^3 - 3*a*b^2*d^2*e + 3*a^2*b*d*e^2 - a^3*e^3)*\sqrt{-b^2*d + a*b*e}) + (\sqrt{e*x + d}*B*a*b*e - \sqrt{e*x + d}*A*b^2*e)/((b^3*d^3 - 3*a*b^2*d^2*e + 3*a^2*b*d*e^2 - a^3*e^3)*((e*x + d)*b - b*d + a*e)) + 2/3*(3*(e*x + d)*B*b*d + B*b*d^2 + 3*(e*x + d)*B*a*e - 6*(e*x + d)*A*b*e - B*a*d*e - A*b*d*e + A*a*e^2)/((b^3*d^3 - 3*a*b^2*d^2*e + 3*a^2*b*d*e^2 - a^3*e^3)*(e*x + d)^(3/2))$$

Mupad [B] (verification not implemented)

Time = 1.07 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.26

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{5/2}} dx = \frac{\frac{2(Ae - Bd)}{3(ae - bd)} + \frac{2(d + ex)(3Bae - 5Abe + 2Bbd)}{3(ae - bd)^2} + \frac{b(d + ex)^2(3Bae - 5Abe + 2Bbd)}{(ae - bd)^3}}{b(d + ex)^{5/2} + (ae - bd)(d + ex)^{3/2}} - \frac{\sqrt{b} \operatorname{atan}\left(\frac{\sqrt{b}\sqrt{d + ex}(a^3e^3 - 3a^2bde^2 + 3ab^2d^2e - b^3d^3)}{(ae - bd)^{7/2}}\right)(3Bae - 5Abe + 2Bbd)}{(ae - bd)^{7/2}}$$

input `int((A + B*x)/((a + b*x)^2*(d + e*x)^(5/2)),x)`

output
$$- ((2*(A*e - B*d))/(3*(a*e - b*d)) + (2*(d + e*x)*(3*B*a*e - 5*A*b*e + 2*B*b*d))/(3*(a*e - b*d)^2) + (b*(d + e*x)^2*(3*B*a*e - 5*A*b*e + 2*B*b*d))/(a*e - b*d)^3)/(b*(d + e*x)^(5/2) + (a*e - b*d)*(d + e*x)^(3/2)) - (b^(1/2)*\operatorname{atan}((b^(1/2)*(d + e*x)^(1/2)*(a^3*e^3 - b^3*d^3 + 3*a*b^2*d^2*e - 3*a^2*b*d*e^2))/(a*e - b*d)^(7/2)))*(3*B*a*e - 5*A*b*e + 2*B*b*d))/(a*e - b*d)^(7/2)$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.34

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{5/2}} dx = \frac{2\sqrt{b}\sqrt{ex+d}\sqrt{ae-bd}\operatorname{atan}\left(\frac{\sqrt{ex+d}b}{\sqrt{b}\sqrt{ae-bd}}\right)bd + 2\sqrt{b}\sqrt{ex+d}\sqrt{ae-bd}\operatorname{atan}\left(\frac{\sqrt{ex+d}b}{\sqrt{b}\sqrt{ae-bd}}\right)}{\sqrt{ex+d}(a^3e^4x - 3a^2bde^3x + 3ab^2d^2e^2x - b^3d^3ex + a^4)}$$

input `int((B*x+A)/(b*x+a)^2/(e*x+d)^(5/2),x)`output `(2*(3*sqrt(b)*sqrt(d + e*x)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))*b*d + 3*sqrt(b)*sqrt(d + e*x)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))*b*e*x - a**2*e**2 + 5*a*b*d*e + 3*a*b*e**2*x - 4*b**2*d**2 - 3*b**2*d*e*x)/(3*sqrt(d + e*x)*(a**3*d*e**3 + a**3*e**4*x - 3*a**2*b*d**2*e**2 - 3*a**2*b*d*e**3*x + 3*a*b**2*d**3*e + 3*a*b**2*d**2*e**2*x - b**3*d**4 - b**3*d**3*e*x))`

3.163 $\int \frac{A+Bx}{(a+bx)^2(d+ex)^{7/2}} dx$

Optimal result	1491
Mathematica [A] (verified)	1492
Rubi [A] (verified)	1492
Maple [A] (verified)	1495
Fricas [B] (verification not implemented)	1496
Sympy [F(-1)]	1497
Maxima [F(-2)]	1497
Giac [B] (verification not implemented)	1497
Mupad [B] (verification not implemented)	1498
Reduce [B] (verification not implemented)	1499

Optimal result

Integrand size = 22, antiderivative size = 208

$$\int \frac{A+Bx}{(a+bx)^2(d+ex)^{7/2}} dx = \frac{2(Bd - Ae)}{5(bd - ae)^2(d+ex)^{5/2}} + \frac{2(bBd - 2Abe + aBe)}{3(bd - ae)^3(d+ex)^{3/2}} + \frac{2b(bBd - 3Abe + 2aBe)}{(bd - ae)^4\sqrt{d+ex}} - \frac{b^2(Ab - aB)\sqrt{d+ex}}{(bd - ae)^4(a+bx)} - \frac{b^{3/2}(2bBd - 7Abe + 5aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{(bd - ae)^{9/2}}$$

output

```
2/5*(-A*e+B*d)/(-a*e+b*d)^2/(e*x+d)^(5/2)+2/3*(-2*A*b*e+B*a*e+B*b*d)/(-a*e
+b*d)^3/(e*x+d)^(3/2)+2*b*(-3*A*b*e+2*B*a*e+B*b*d)/(-a*e+b*d)^4/(e*x+d)^(1
/2)-b^2*(A*b-B*a)*(e*x+d)^(1/2)/(-a*e+b*d)^4/(b*x+a)-b^(3/2)*(-7*A*b*e+5*B
*a*e+2*B*b*d)*arctanh(b^(1/2)*(e*x+d)^(1/2)/(-a*e+b*d)^(1/2))/(-a*e+b*d)^(
9/2)
```


Mathematica [A] (verified)

Time = 0.70 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.39

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{7/2}} dx = \frac{B(-2a^3e^2(2d + 5ex) + 2b^3dx(23d^2 + 35dex + 15e^2x^2) + 2a^2be(24d^2 + 58dex + 24e^2x^2) + b^3/2(2bBd - 7Abe + 5aBe) \arctan\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right)}{(-bd + ae)^{9/2}}$$

input

```
Integrate[(A + B*x)/((a + b*x)^2*(d + e*x)^(7/2)),x]
```

output

```
(B*(-2*a^3*e^2*(2*d + 5*e*x) + 2*b^3*d*x*(23*d^2 + 35*d*e*x + 15*e^2*x^2) + 2*a^2*b*e*(24*d^2 + 58*d*e*x + 25*e^2*x^2) + a*b^2*(61*d^3 + 163*d^2*e*x + 195*d*e^2*x^2 + 75*e^3*x^3)) - A*(6*a^3*e^3 - 2*a^2*b*e^2*(16*d + 7*e*x) + 2*a*b^2*e*(58*d^2 + 84*d*e*x + 35*e^2*x^2) + b^3*(15*d^3 + 161*d^2*e*x + 245*d*e^2*x^2 + 105*e^3*x^3)))/(15*(b*d - a*e)^4*(a + b*x)*(d + e*x)^(5/2)) + (b^(3/2)*(2*b*B*d - 7*A*b*e + 5*a*B*e)*ArcTan[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[-(b*d) + a*e]])/(-(b*d) + a*e)^(9/2)
```

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.03, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {87, 61, 61, 61, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{7/2}} dx$$

$$\downarrow 87$$

$$\frac{(5aBe - 7Abe + 2bBd) \int \frac{1}{(a+bx)(d+ex)^{7/2}} dx}{2b(bd - ae)} - \frac{Ab - aB}{b(a + bx)(d + ex)^{5/2}(bd - ae)}$$

$$\downarrow 61$$

$$\begin{aligned}
 & \frac{(5aBe - 7Abe + 2bBd) \left(\frac{b \int \frac{1}{(a+bx)(d+ex)^{5/2}} dx}{bd-ae} + \frac{2}{5(d+ex)^{5/2}(bd-ae)} \right)}{\frac{2b(bd-ae)}{Ab-aB} \cdot \frac{1}{b(a+bx)(d+ex)^{5/2}(bd-ae)}} \\
 & \quad \downarrow 61 \\
 & \frac{(5aBe - 7Abe + 2bBd) \left(\frac{b \left(\frac{b \int \frac{1}{(a+bx)(d+ex)^{3/2}} dx}{bd-ae} + \frac{2}{3(d+ex)^{3/2}(bd-ae)} \right)}{bd-ae} + \frac{2}{5(d+ex)^{5/2}(bd-ae)} \right)}{\frac{2b(bd-ae)}{Ab-aB} \cdot \frac{1}{b(a+bx)(d+ex)^{5/2}(bd-ae)}} \\
 & \quad \downarrow 61 \\
 & \frac{(5aBe - 7Abe + 2bBd) \left(\frac{b \left(\frac{b \left(\frac{b \int \frac{1}{(a+bx)\sqrt{d+ex}} dx}{bd-ae} + \frac{2}{\sqrt{d+ex}(bd-ae)} \right)}{3(d+ex)^{3/2}(bd-ae)} + \frac{2}{5(d+ex)^{5/2}(bd-ae)} \right)}{bd-ae} \right)}{\frac{2b(bd-ae)}{Ab-aB} \cdot \frac{1}{b(a+bx)(d+ex)^{5/2}(bd-ae)}} \\
 & \quad \downarrow 73 \\
 & \frac{(5aBe - 7Abe + 2bBd) \left(\frac{b \left(\frac{b \left(\frac{2b \int \frac{1}{a + \frac{b(d+ex) - bd}{e(bd-ae)}} d\sqrt{d+ex}}{bd-ae} + \frac{2}{\sqrt{d+ex}(bd-ae)} \right)}{3(d+ex)^{3/2}(bd-ae)} + \frac{2}{5(d+ex)^{5/2}(bd-ae)} \right)}{bd-ae} \right)}{\frac{2b(bd-ae)}{Ab-aB} \cdot \frac{1}{b(a+bx)(d+ex)^{5/2}(bd-ae)}}
 \end{aligned}$$

↓ 221

$$(5aBe - 7Abe + 2bBd) \left(\frac{b \left(\frac{2}{\sqrt{d+ex}(bd-ae)} - \frac{2\sqrt{b}\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{(bd-ae)^{3/2}} \right)}{bd-ae} + \frac{2}{3(d+ex)^{3/2}(bd-ae)} \right) + \frac{2}{5(d+ex)^{5/2}(bd-ae)}$$

$$\frac{2b(bd-ae)}{Ab-aB} \frac{1}{b(a+bx)(d+ex)^{5/2}(bd-ae)}$$

input `Int[(A + B*x)/((a + b*x)^2*(d + e*x)^(7/2)), x]`

output `-((A*b - a*B)/(b*(b*d - a*e)*(a + b*x)*(d + e*x)^(5/2))) + ((2*b*B*d - 7*A*b*e + 5*a*B*e)*(2/(5*(b*d - a*e)*(d + e*x)^(5/2)) + (b*(2/(3*(b*d - a*e)*(d + e*x)^(3/2)) + (b*(2/((b*d - a*e)*Sqrt[d + e*x]) - (2*Sqrt[b]*ArcTanh[Sqrt[b]*Sqrt[d + e*x])/Sqrt[b*d - a*e]])/(b*d - a*e)^(3/2)))/(b*d - a*e)))/(2*b*(b*d - a*e))`

Defintions of rubi rules used

rule 61 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p
_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Maple [A] (verified)

Time = 0.51 (sec) , antiderivative size = 202, normalized size of antiderivative = 0.97

method	result
derivativedivides	$-\frac{2(Ae-Bd)}{5(ae-db)^2(ex+d)^{\frac{5}{2}}} - \frac{2(-2Abe+BAe+Bbd)}{3(ae-db)^3(ex+d)^{\frac{3}{2}}} - \frac{2b(3Abe-2BAe-Bbd)}{(ae-db)^4\sqrt{ex+d}} - \frac{2b^2\left(\frac{\frac{1}{2}Abe-\frac{1}{2}BAe}{(ex+d)b+ae-db}\sqrt{ex+d} + \frac{(7Abe-5BAe+3A^2)}{(ae-db)^2}\right)}{(ae-db)^5}$
default	$-\frac{2(Ae-Bd)}{5(ae-db)^2(ex+d)^{\frac{5}{2}}} - \frac{2(-2Abe+BAe+Bbd)}{3(ae-db)^3(ex+d)^{\frac{3}{2}}} - \frac{2b(3Abe-2BAe-Bbd)}{(ae-db)^4\sqrt{ex+d}} - \frac{2b^2\left(\frac{\frac{1}{2}Abe-\frac{1}{2}BAe}{(ex+d)b+ae-db}\sqrt{ex+d} + \frac{(7Abe-5BAe+3A^2)}{(ae-db)^2}\right)}{(ae-db)^5}$
pseudoelliptic	$2\left(\frac{35(ex+d)^{\frac{5}{2}}b^2(bx+a)\left((Ae-\frac{2Bd}{7})b-\frac{5BAe}{7}\right)\arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{2} + \sqrt{(ae-db)b}\left(\frac{35Ae^3x^3 + \frac{245(-6Bx+A)x^2de^2}{3}}{3} + \dots\right)\right)$

```
input int((B*x+A)/(b*x+a)^2/(e*x+d)^(7/2), x, method=_RETURNVERBOSE)
```

```
output -2/5*(A*e-B*d)/(a*e-b*d)^2/(e*x+d)^(5/2)-2/3*(-2*A*b*e+B*a*e+B*b*d)/(a*e-b
*d)^3/(e*x+d)^(3/2)-2*b*(3*A*b*e-2*B*a*e-B*b*d)/(a*e-b*d)^4/(e*x+d)^(1/2)-
2/(a*e-b*d)^4*b^2*((1/2*A*b*e-1/2*B*a*e)*(e*x+d)^(1/2)/((e*x+d)*b+a*e-d*b)
+1/2*(7*A*b*e-5*B*a*e-2*B*b*d)/((a*e-b*d)*b)^(1/2)*arctan(b*(e*x+d)^(1/2)/
((a*e-b*d)*b)^(1/2)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 851 vs. $2(185) = 370$.

Time = 0.23 (sec) , antiderivative size = 1731, normalized size of antiderivative = 8.32

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{7/2}} dx = \text{Too large to display}$$

input `integrate((B*x+A)/(b*x+a)^2/(e*x+d)^(7/2),x, algorithm="fricas")`

output

```
[-1/30*(15*(2*B*a*b^2*d^4 + (5*B*a^2*b - 7*A*a*b^2)*d^3*e + (2*B*b^3*d*e^3
+ (5*B*a*b^2 - 7*A*b^3)*e^4)*x^4 + (6*B*b^3*d^2*e^2 + (17*B*a*b^2 - 21*A*
b^3)*d*e^3 + (5*B*a^2*b - 7*A*a*b^2)*e^4)*x^3 + 3*(2*B*b^3*d^3*e + 7*(B*a*
b^2 - A*b^3)*d^2*e^2 + (5*B*a^2*b - 7*A*a*b^2)*d*e^3)*x^2 + (2*B*b^3*d^4 +
(11*B*a*b^2 - 7*A*b^3)*d^3*e + 3*(5*B*a^2*b - 7*A*a*b^2)*d^2*e^2)*x)*sqrt
(b/(b*d - a*e))*log((b*e*x + 2*b*d - a*e + 2*(b*d - a*e)*sqrt(e*x + d))*sqrt
(b/(b*d - a*e)))/(b*x + a)) + 2*(6*A*a^3*e^3 - (61*B*a*b^2 - 15*A*b^3)*d^
3 - 4*(12*B*a^2*b - 29*A*a*b^2)*d^2*e + 4*(B*a^3 - 8*A*a^2*b)*d*e^2 - 15*(
2*B*b^3*d*e^2 + (5*B*a*b^2 - 7*A*b^3)*e^3)*x^3 - 5*(14*B*b^3*d^2*e + (39*B
*a*b^2 - 49*A*b^3)*d*e^2 + 2*(5*B*a^2*b - 7*A*a*b^2)*e^3)*x^2 - (46*B*b^3*d
^3 + (163*B*a*b^2 - 161*A*b^3)*d^2*e + 4*(29*B*a^2*b - 42*A*a*b^2)*d*e^2
- 2*(5*B*a^3 - 7*A*a^2*b)*e^3)*x)*sqrt(e*x + d))/(a*b^4*d^7 - 4*a^2*b^3*d^
6*e + 6*a^3*b^2*d^5*e^2 - 4*a^4*b*d^4*e^3 + a^5*d^3*e^4 + (b^5*d^4*e^3 - 4
*a*b^4*d^3*e^4 + 6*a^2*b^3*d^2*e^5 - 4*a^3*b^2*d*e^6 + a^4*b*e^7)*x^4 + (3
*b^5*d^5*e^2 - 11*a*b^4*d^4*e^3 + 14*a^2*b^3*d^3*e^4 - 6*a^3*b^2*d^2*e^5 -
a^4*b*d*e^6 + a^5*e^7)*x^3 + 3*(b^5*d^6*e - 3*a*b^4*d^5*e^2 + 2*a^2*b^3*d
^4*e^3 + 2*a^3*b^2*d^3*e^4 - 3*a^4*b*d^2*e^5 + a^5*d*e^6)*x^2 + (b^5*d^7 -
a*b^4*d^6*e - 6*a^2*b^3*d^5*e^2 + 14*a^3*b^2*d^4*e^3 - 11*a^4*b*d^3*e^4 +
3*a^5*d^2*e^5)*x), 1/15*(15*(2*B*a*b^2*d^4 + (5*B*a^2*b - 7*A*a*b^2)*d^3*
e + (2*B*b^3*d*e^3 + (5*B*a*b^2 - 7*A*b^3)*e^4)*x^4 + (6*B*b^3*d^2*e^2 ...
```

Sympy [F(-1)]

Timed out.

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{7/2}} dx = \text{Timed out}$$

input `integrate((B*x+A)/(b*x+a)**2/(e*x+d)**(7/2),x)`

output Timed out

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{7/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^2/(e*x+d)^(7/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 422 vs. 2(185) = 370.

Time = 0.13 (sec) , antiderivative size = 422, normalized size of antiderivative = 2.03

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{7/2}} dx = \frac{(2 Bb^3d + 5 Bab^2e - 7 Ab^3e) \arctan\left(\frac{\sqrt{ex+db}}{\sqrt{-b^2d+abe}}\right)}{(b^4d^4 - 4ab^3d^3e + 6a^2b^2d^2e^2 - 4a^3bde^3 + a^4e^4)\sqrt{-b^2d+abe}} + \frac{\sqrt{ex+d}Bab^2e - \sqrt{ex+d}Ab^3e}{(b^4d^4 - 4ab^3d^3e + 6a^2b^2d^2e^2 - 4a^3bde^3 + a^4e^4)((ex+d)b - bd + ae)} + \frac{2(15(ex+d)^2Bb^2d + 5(ex+d)Bb^2d^2 + 3Bb^2d^3 + 30(ex+d)^2Babe - 45(ex+d)^2Ab^2e - 10(ex+d)Abe)}{15(b^4d^4 - 4ab^3d^3e + 6a^2b^2d^2e^2 - 4a^3bde^3 + a^4e^4)}$$

input `integrate((B*x+A)/(b*x+a)^2/(e*x+d)^(7/2),x, algorithm="giac")`

output
$$\frac{(2Bb^3d + 5Bab^2e - 7Aab^3e) \arctan(\sqrt{ex+d}b/\sqrt{-b^2d + abe})}{(b^4d^4 - 4a^3b^3d^3e + 6a^2b^2d^2e^2 - 4a^3b^3d^3e + a^4e^4) \sqrt{-b^2d + abe}} + \frac{(\sqrt{ex+d}Bab^2e - \sqrt{ex+d}Aab^3e)}{(b^4d^4 - 4a^3b^3d^3e + 6a^2b^2d^2e^2 - 4a^3b^3d^3e + a^4e^4) ((ex+d)b - b^2d + abe)} + \frac{2}{15} \frac{(15(ex+d)^2Bb^2d + 5(ex+d)Bb^2d^2 + 3Bb^2d^3 + 30(ex+d)^2Bab^2e - 45(ex+d)^2Aab^2e - 10(ex+d)Aab^2d^2e - 6Bab^2d^2e - 3Aab^2d^2e - 5(ex+d)Bba^2e^2 + 10(ex+d)Aab^2e^2 + 3Bba^2d^2e^2 + 6Aab^2d^2e^2 - 3Aa^2e^3)}{(b^4d^4 - 4a^3b^3d^3e + 6a^2b^2d^2e^2 - 4a^3b^3d^3e + a^4e^4) (ex+d)^{5/2}}$$

Mupad [B] (verification not implemented)

Time = 1.10 (sec) , antiderivative size = 261, normalized size of antiderivative = 1.25

$$\int \frac{A + Bx}{(a + bx)^2 (d + ex)^{7/2}} dx = \frac{b^{3/2} \operatorname{atan}\left(\frac{\sqrt{b}\sqrt{d+ex}(a^4e^4 - 4a^3bde^3 + 6a^2b^2d^2e^2 - 4ab^3d^3e + b^4d^4)}{(ae-bd)^{9/2}}\right) (5Bae - 7Abe + \dots)}{(ae-bd)^{9/2}}$$

$$- \frac{\frac{2(Ae-Bd)}{5(ae-bd)} + \frac{2(d+ex)(5Bae-7Abe+2Bbd)}{15(ae-bd)^2} - \frac{b^2(d+ex)^3(5Bae-7Abe+2Bbd)}{(ae-bd)^4} - \frac{2b(d+ex)^2(5Bae-7Abe+2Bbd)}{3(ae-bd)^3}}{b(d+ex)^{7/2} + (ae-bd)(d+ex)^{5/2}}$$

input `int((A + B*x)/((a + b*x)^2*(d + e*x)^(7/2)),x)`

output
$$\frac{(b^{3/2} \operatorname{atan}((b^{1/2}(d+ex)^{1/2}(a^4e^4 + b^4d^4 + 6a^2b^2d^2e^2 - 4a^3b^3d^3e - 4a^3b^3d^3e)) / (ae - b^2d)^{9/2}) (5Bae - 7Abe + 2Bbd)) / (ae - b^2d)^{9/2} - ((2(Ae - B^2d)) / (5(ae - b^2d)) + (2(d + ex)(5Bae - 7Abe + 2Bbd)) / (15(ae - b^2d)^2) - (b^2(d + ex)^3(5Bae - 7Abe + 2Bbd)) / (ae - b^2d)^4 - (2b(d + ex)^2(5Bae - 7Abe + 2Bbd)) / (3(ae - b^2d)^3)) / (b(d + ex)^{7/2} + (ae - b^2d)(d + ex)^{5/2})}{(ae - b^2d)^{9/2}}$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 437, normalized size of antiderivative = 2.10

$$\int \frac{A + Bx}{(a + bx)^2(d + ex)^{7/2}} dx = \frac{-2\sqrt{b}\sqrt{ex+d}\sqrt{ae-bd}\operatorname{atan}\left(\frac{\sqrt{ex+d}b}{\sqrt{b}\sqrt{ae-bd}}\right)b^2d^2 - 4\sqrt{b}\sqrt{ex+d}\sqrt{ae-bd}\operatorname{atan}\left(\frac{\sqrt{ex+d}b}{\sqrt{b}\sqrt{ae-bd}}\right)b^2d - 4\sqrt{b}\sqrt{ex+d}\sqrt{ae-bd}\operatorname{atan}\left(\frac{\sqrt{ex+d}b}{\sqrt{b}\sqrt{ae-bd}}\right)b^2d - 4\sqrt{b}\sqrt{ex+d}\sqrt{ae-bd}\operatorname{atan}\left(\frac{\sqrt{ex+d}b}{\sqrt{b}\sqrt{ae-bd}}\right)b^2d}{\sqrt{ex+d}(a^4e^6x^2 - 4a^3bde^5x^2 + 6a^2b^2d^2e^4x^2 - 4ab^3d^3e^3x^2 + b^4d^4e^2x^2)}$$

input `int((B*x+A)/(b*x+a)^2/(e*x+d)^(7/2),x)`

output

```
(2*( - 15*sqrt(b)*sqrt(d + e*x)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))*b**2*d**2 - 30*sqrt(b)*sqrt(d + e*x)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))*b**2*d*e*x - 15*sqrt(b)*sqrt(d + e*x)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))*b**2*e**2*x**2 - 3*a**3*e**3 + 14*a**2*b*d*e**2 + 5*a**2*b*e**3*x - 34*a*b**2*d**2*e - 40*a*b**2*d*e**2*x - 15*a*b**2*e**3*x**2 + 23*b**3*d**3 + 35*b**3*d**2*e*x + 15*b**3*d*e**2*x**2))/(15*sqrt(d + e*x)*(a**4*d**2*e**4 + 2*a**4*d*e**5*x + a**4*e**6*x**2 - 4*a**3*b*d**3*e**3 - 8*a**3*b*d**2*e**4*x - 4*a**3*b*d*e**5*x**2 + 6*a**2*b**2*d**4*e**2 + 12*a**2*b**2*d**3*e**3*x + 6*a**2*b**2*d**2*e**4*x**2 - 4*a*b**3*d**5*e - 8*a*b**3*d**4*e**2*x - 4*a*b**3*d**3*e**3*x**2 + b**4*d**6 + 2*b**4*d**5*e*x + b**4*d**4*e**2*x**2))
```


3.164 $\int \frac{(A+Bx)(d+ex)^{7/2}}{(a+bx)^3} dx$

Optimal result	1500
Mathematica [A] (verified)	1501
Rubi [A] (verified)	1501
Maple [A] (verified)	1505
Fricas [B] (verification not implemented)	1506
Sympy [F(-1)]	1507
Maxima [F(-2)]	1508
Giac [B] (verification not implemented)	1508
Mupad [B] (verification not implemented)	1509
Reduce [B] (verification not implemented)	1510

Optimal result

Integrand size = 22, antiderivative size = 236

$$\int \frac{(A+Bx)(d+ex)^{7/2}}{(a+bx)^3} dx = \frac{e(bd-ae)(6bBd+7Abe-13aBe)\sqrt{d+ex}}{b^5} - \frac{(bd-ae)^2(4bBd+7Abe-11aBe)\sqrt{d+ex}}{4b^5(a+bx)} + \frac{e(8bBd+7Abe-15aBe)(d+ex)^{3/2}}{6b^4} + \frac{2Be(d+ex)^{5/2}}{5b^3} - \frac{(Ab-aB)(d+ex)^{7/2}}{2b^2(a+bx)^2} - \frac{7e(bd-ae)^{3/2}(4bBd+5Abe-9aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{4b^{11/2}}$$

output

```
e*(-a*e+b*d)*(7*A*b*e-13*B*a*e+6*B*b*d)*(e*x+d)^(1/2)/b^5-1/4*(-a*e+b*d)^2
*(7*A*b*e-11*B*a*e+4*B*b*d)*(e*x+d)^(1/2)/b^5/(b*x+a)+1/6*e*(7*A*b*e-15*B*
a*e+8*B*b*d)*(e*x+d)^(3/2)/b^4+2/5*B*e*(e*x+d)^(5/2)/b^3-1/2*(A*b-B*a)*(e*
x+d)^(7/2)/b^2/(b*x+a)^2-7/4*e*(-a*e+b*d)^(3/2)*(5*A*b*e-9*B*a*e+4*B*b*d)*
arctanh(b^(1/2)*(e*x+d)^(1/2)/(-a*e+b*d)^(1/2))/b^(11/2)
```

Mathematica [A] (verified)

Time = 0.79 (sec) , antiderivative size = 309, normalized size of antiderivative = 1.31

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^3} dx = \frac{\sqrt{d + ex}(-5Ab(105a^3e^3 + 35a^2be^2(-4d + 5ex) + 7ab^2e(3d^2 - 34dex + 8e^2x^2) + 7e(-bd + ae)^{3/2}(4bBd + 5Abe - 9aBe) \arctan\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right))}{4b^{11/2}}$$

input

```
Integrate[((A + B*x)*(d + e*x)^(7/2))/(a + b*x)^3,x]
```

output

```
(Sqrt[d + e*x]*(-5*A*b*(105*a^3*e^3 + 35*a^2*b*e^2*(-4*d + 5*e*x) + 7*a*b^2*e*(3*d^2 - 34*d*e*x + 8*e^2*x^2) + b^3*(6*d^3 + 39*d^2*e*x - 80*d*e^2*x^2 - 8*e^3*x^3)) + B*(945*a^4*e^3 + 105*a^3*b*e^2*(-16*d + 15*e*x) + 7*a^2*b^2*e*(107*d^2 - 406*d*e*x + 72*e^2*x^2) + 4*b^4*x*(-15*d^3 + 116*d^2*e*x + 32*d*e^2*x^2 + 6*e^3*x^3) - a*b^3*(30*d^3 - 1303*d^2*e*x + 944*d*e^2*x^2 + 72*e^3*x^3)))/(60*b^5*(a + b*x)^2) + (7*e*(-(b*d) + a*e)^(3/2)*(4*b*B*d + 5*A*b*e - 9*a*B*e)*ArcTan[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[-(b*d) + a*e]])/(4*b^(11/2))
```

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 225, normalized size of antiderivative = 0.95, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.318$, Rules used = {87, 51, 60, 60, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^3} dx$$

↓ 87

$$\frac{(-9aBe + 5Abe + 4bBd) \int \frac{(d+ex)^{7/2}}{(a+bx)^2} dx}{4b(bd - ae)} - \frac{(d + ex)^{9/2}(Ab - aB)}{2b(a + bx)^2(bd - ae)}$$

↓ 51

$$\begin{aligned}
 & \frac{(-9aBe + 5Abe + 4bBd) \left(\frac{7e \int \frac{(d+ex)^{5/2}}{a+bx} dx}{2b} - \frac{(d+ex)^{7/2}}{b(a+bx)} \right)}{4b(bd - ae)} - \frac{(d + ex)^{9/2}(Ab - aB)}{2b(a + bx)^2(bd - ae)} \\
 & \quad \downarrow 60 \\
 & \frac{(-9aBe + 5Abe + 4bBd) \left(\frac{7e \left(\frac{(bd-ae) \int \frac{(d+ex)^{3/2}}{a+bx} dx}{b} + \frac{2(d+ex)^{5/2}}{5b} \right)}{2b} - \frac{(d+ex)^{7/2}}{b(a+bx)} \right)}{4b(bd - ae)} - \frac{(d + ex)^{9/2}(Ab - aB)}{2b(a + bx)^2(bd - ae)} \\
 & \quad \downarrow 60 \\
 & \frac{(-9aBe + 5Abe + 4bBd) \left(\frac{7e \left(\frac{(bd-ae) \left(\frac{(bd-ae) \int \frac{\sqrt{d+ex}}{a+bx} dx}{b} + \frac{2(d+ex)^{3/2}}{3b} \right)}{b} + \frac{2(d+ex)^{5/2}}{5b} \right)}{2b} - \frac{(d+ex)^{7/2}}{b(a+bx)} \right)}{4b(bd - ae)} - \frac{(d + ex)^{9/2}(Ab - aB)}{2b(a + bx)^2(bd - ae)} \\
 & \quad \downarrow 60
 \end{aligned}$$

$$\begin{aligned}
 & \left(\frac{(-9aBe + 5Abe + 4bBd)}{2b} \left(\frac{7e}{b} \left(\frac{(bd-ae) \left(\frac{(bd-ae) \int \frac{1}{(a+bx)\sqrt{d+ex}} dx + \frac{2\sqrt{d+ex}}{b} \right)}{b} + \frac{2(d+ex)^{3/2}}{3b} \right)}{b} + \frac{2(d+ex)^{5/2}}{5b} \right) - \frac{(d+ex)^{7/2}}{b(a+bx)} \right) \right)
 \end{aligned}$$

$$\frac{4b(bd - ae)}{2b(a + bx)^2(bd - ae)} \frac{(d + ex)^{9/2}(Ab - aB)}{2b(a + bx)^2(bd - ae)}$$

↓ 73

$$\begin{aligned}
 & \left(\frac{(-9aBe + 5Abe + 4bBd)}{2b} \left(\frac{7e}{b} \left(\frac{(bd-ae) \left(\frac{2(bd-ae) \int \frac{1}{a + \frac{b(d+ex)}{e} - \frac{bd}{e}} d\sqrt{d+ex}} + \frac{2\sqrt{d+ex}}{b} \right)}{b} + \frac{2(d+ex)^{3/2}}{3b} \right)}{b} + \frac{2(d+ex)^{5/2}}{5b} \right) - \frac{(d+ex)^{7/2}}{b(a+bx)} \right)
 \end{aligned}$$

$$\frac{4b(bd - ae)}{2b(a + bx)^2(bd - ae)} \frac{(d + ex)^{9/2}(Ab - aB)}{2b(a + bx)^2(bd - ae)}$$

↓ 221

$$\begin{aligned}
 & \left(\frac{7e \left(\frac{(bd-ae) \left(\frac{2\sqrt{d+ex}}{b} - \frac{2\sqrt{bd-ae} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{3/2}} \right)}{b} + \frac{2(d+ex)^{3/2}}{3b} \right)}{b} + \frac{2(d+ex)^{5/2}}{5b} \right)}{2b} - \frac{(d+ex)^{9/2}}{b(a+bx)} \right) \\
 & \frac{(-9aBe + 5Abe + 4bBd)}{2b} \left(\frac{7e \left(\frac{(bd-ae) \left(\frac{2\sqrt{d+ex}}{b} - \frac{2\sqrt{bd-ae} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{3/2}} \right)}{b} + \frac{2(d+ex)^{3/2}}{3b} \right)}{b} + \frac{2(d+ex)^{5/2}}{5b} \right)}{2b} - \frac{(d+ex)^{9/2}}{b(a+bx)} \right) \\
 & \frac{4b(bd - ae)}{2b(a + bx)^2(bd - ae)} \frac{(d + ex)^{9/2}(Ab - aB)}{2b(a + bx)^2(bd - ae)}
 \end{aligned}$$

input `Int[((A + B*x)*(d + e*x)^(7/2))/(a + b*x)^3,x]`

output `-1/2*((A*b - a*B)*(d + e*x)^(9/2))/(b*(b*d - a*e)*(a + b*x)^2) + ((4*b*B*d + 5*A*b*e - 9*a*B*e)*(-(d + e*x)^(7/2)/(b*(a + b*x))) + (7*e*((2*(d + e*x)^(5/2))/(5*b) + ((b*d - a*e)*((2*(d + e*x)^(3/2))/(3*b) + ((b*d - a*e)*((2*sqrt[d + e*x])/b - (2*sqrt[b*d - a*e]*ArcTanh[(sqrt[b]*sqrt[d + e*x])/sqrt[b*d - a*e]])/b^(3/2)))/b))/b))/(2*b)))/(4*b*(b*d - a*e))`

Defintions of rubi rules used

rule 51 `Int[((a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n/(b*(m + 1)), x] - Simp[d*(n/(b*(m + 1)))] Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 276, normalized size of antiderivative = 1.17

method	result
risch	$\frac{2e(-3e^2b^2Bx^2-5Ab^2e^2x+15Bab e^2x-16b^2Bdex+45Aab e^2-50Ab^2de-90Ba^2e^2+150Babde-58b^2Bd^2)\sqrt{ex+d}}{15b^5}$
pseudoelliptic	$35 \left(-(bx+a)^2 \left(\left(Ae + \frac{4Bd}{5} \right) b - \frac{9Bae}{5} \right) (ae-db)^2 e \arctan \left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}} \right) + \sqrt{ex+d} \left(\left(-\frac{8\left(\frac{3Bx+A}{5}\right)x^3e^3}{105} - \frac{16\left(\frac{8Bx+A}{25}\right)}{21} \right) \right) \right)$
derivativedivides	$2e \left(-\frac{-\frac{b^2B(ex+d)^{\frac{5}{2}}}{5} - \frac{Ab^2e(ex+d)^{\frac{3}{2}}}{3} + Babe(ex+d)^{\frac{3}{2}} - \frac{2Bb^2d(ex+d)^{\frac{3}{2}}}{3} + 3Aabe^2\sqrt{ex+d} - 3Ab^2de\sqrt{ex+d} - 6Ba^2e^2\sqrt{ex+d}}{b^5} \right)$
default	$2e \left(-\frac{-\frac{b^2B(ex+d)^{\frac{5}{2}}}{5} - \frac{Ab^2e(ex+d)^{\frac{3}{2}}}{3} + Babe(ex+d)^{\frac{3}{2}} - \frac{2Bb^2d(ex+d)^{\frac{3}{2}}}{3} + 3Aabe^2\sqrt{ex+d} - 3Ab^2de\sqrt{ex+d} - 6Ba^2e^2\sqrt{ex+d}}{b^5} \right)$

```
input int((B*x+A)*(e*x+d)^(7/2)/(b*x+a)^3,x,method=_RETURNVERBOSE)
```

```
output -2/15**(-3*B*b^2*e^2*x^2-5*A*b^2*e^2*x+15*B*a*b*e^2*x-16*B*b^2*d*e*x+45*A
*a*b*e^2-50*A*b^2*d*e-90*B*a^2*e^2+150*B*a*b*d*e-58*B*b^2*d^2)*(e*x+d)^(1/
2)/b^5+1/b^5*(2*a^2*e^2-4*a*b*d*e+2*b^2*d^2)*e*(((-13/8*A*b^2*e+17/8*B*a*b
*e-1/2*b^2*B*d)*(e*x+d)^(3/2)+(-11/8*A*a*b*e^2+11/8*A*b^2*d*e+15/8*B*a^2*e
^2-19/8*B*a*b*d*e+1/2*b^2*B*d^2)*(e*x+d)^(1/2))/((e*x+d)*b+a*e-d*b)^2+7/8*
(5*A*b*e-9*B*a*e+4*B*b*d)/((a*e-b*d)*b)^(1/2)*arctan(b*(e*x+d)^(1/2)/((a*e
-b*d)*b)^(1/2)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 525 vs. 2(206) = 412.
 Time = 0.11 (sec) , antiderivative size = 1060, normalized size of antiderivative = 4.49

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^3} dx = \text{Too large to display}$$

```
input integrate((B*x+A)*(e*x+d)^(7/2)/(b*x+a)^3,x, algorithm="fricas")
```

output

```

[-1/120*(105*(4*B*a^2*b^2*d^2*e - (13*B*a^3*b - 5*A*a^2*b^2)*d*e^2 + (9*B*
a^4 - 5*A*a^3*b)*e^3 + (4*B*b^4*d^2*e - (13*B*a*b^3 - 5*A*b^4)*d*e^2 + (9*
B*a^2*b^2 - 5*A*a*b^3)*e^3)*x^2 + 2*(4*B*a*b^3*d^2*e - (13*B*a^2*b^2 - 5*A
*a*b^3)*d*e^2 + (9*B*a^3*b - 5*A*a^2*b^2)*e^3)*x)*sqrt((b*d - a*e)/b)*log(
(b*e*x + 2*b*d - a*e + 2*sqrt(e*x + d)*b*sqrt((b*d - a*e)/b))/(b*x + a)) -
2*(24*B*b^4*e^3*x^4 - 30*(B*a*b^3 + A*b^4)*d^3 + 7*(107*B*a^2*b^2 - 15*A*
a*b^3)*d^2*e - 140*(12*B*a^3*b - 5*A*a^2*b^2)*d*e^2 + 105*(9*B*a^4 - 5*A*a
^3*b)*e^3 + 8*(16*B*b^4*d*e^2 - (9*B*a*b^3 - 5*A*b^4)*e^3)*x^3 + 8*(58*B*b
^4*d^2*e - 2*(59*B*a*b^3 - 25*A*b^4)*d*e^2 + 7*(9*B*a^2*b^2 - 5*A*a*b^3)*e
^3)*x^2 - (60*B*b^4*d^3 - (1303*B*a*b^3 - 195*A*b^4)*d^2*e + 14*(203*B*a^2
*b^2 - 85*A*a*b^3)*d*e^2 - 175*(9*B*a^3*b - 5*A*a^2*b^2)*e^3)*x)*sqrt(e*x
+ d))/(b^7*x^2 + 2*a*b^6*x + a^2*b^5), -1/60*(105*(4*B*a^2*b^2*d^2*e - (13
*B*a^3*b - 5*A*a^2*b^2)*d*e^2 + (9*B*a^4 - 5*A*a^3*b)*e^3 + (4*B*b^4*d^2*e
- (13*B*a*b^3 - 5*A*b^4)*d*e^2 + (9*B*a^2*b^2 - 5*A*a*b^3)*e^3)*x^2 + 2*(
4*B*a*b^3*d^2*e - (13*B*a^2*b^2 - 5*A*a*b^3)*d*e^2 + (9*B*a^3*b - 5*A*a^2*
b^2)*e^3)*x)*sqrt(-(b*d - a*e)/b)*arctan(-sqrt(e*x + d)*b*sqrt(-(b*d - a*e
)/b)/(b*d - a*e)) - (24*B*b^4*e^3*x^4 - 30*(B*a*b^3 + A*b^4)*d^3 + 7*(107*
B*a^2*b^2 - 15*A*a*b^3)*d^2*e - 140*(12*B*a^3*b - 5*A*a^2*b^2)*d*e^2 + 105
*(9*B*a^4 - 5*A*a^3*b)*e^3 + 8*(16*B*b^4*d*e^2 - (9*B*a*b^3 - 5*A*b^4)*e^3
)*x^3 + 8*(58*B*b^4*d^2*e - 2*(59*B*a*b^3 - 25*A*b^4)*d*e^2 + 7*(9*B*a^...

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^3} dx = \text{Timed out}$$

input

```
integrate((B*x+A)*(e*x+d)**(7/2)/(b*x+a)**3,x)
```

output

Timed out

Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^3} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)*(e*x+d)^(7/2)/(b*x+a)^3,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 597 vs. 2(206) = 412.

Time = 0.14 (sec) , antiderivative size = 597, normalized size of antiderivative = 2.53

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^3} dx = \frac{7(4Bb^3d^3e - 17Bab^2d^2e^2 + 5Ab^3d^2e^2 + 22Ba^2bde^3 - 10Aab^2de^3 - 9Ba^3e^3) + 4(ex + d)^{\frac{3}{2}}Bb^4d^3e - 4\sqrt{ex + d}Bb^4d^4e - 25(ex + d)^{\frac{3}{2}}Bab^3d^2e^2 + 13(ex + d)^{\frac{3}{2}}Ab^4d^2e^2 + 27\sqrt{ex + d}Ba^3e^3}{4\sqrt{-b^2d + abeb^5}} + \frac{2\left(3(ex + d)^{\frac{5}{2}}Bb^{12}e + 10(ex + d)^{\frac{3}{2}}Bb^{12}de + 45\sqrt{ex + d}Bb^{12}d^2e - 15(ex + d)^{\frac{3}{2}}Bab^{11}e^2 + 5(ex + d)^{\frac{3}{2}}A\right)}{15b^{15}}$$

input `integrate((B*x+A)*(e*x+d)^(7/2)/(b*x+a)^3,x, algorithm="giac")`

output

```
7/4*(4*B*b^3*d^3*e - 17*B*a*b^2*d^2*e^2 + 5*A*b^3*d^2*e^2 + 22*B*a^2*b*d*e^3 - 10*A*a*b^2*d*e^3 - 9*B*a^3*e^4 + 5*A*a^2*b*e^4)*arctan(sqrt(e*x + d)*b/sqrt(-b^2*d + a*b*e))/(sqrt(-b^2*d + a*b*e)*b^5) - 1/4*(4*(e*x + d)^(3/2)*B*b^4*d^3*e - 4*sqrt(e*x + d)*B*b^4*d^4*e - 25*(e*x + d)^(3/2)*B*a*b^3*d^2*e^2 + 13*(e*x + d)^(3/2)*A*b^4*d^2*e^2 + 27*sqrt(e*x + d)*B*a*b^3*d^3*e^2 - 11*sqrt(e*x + d)*A*b^4*d^3*e^2 + 38*(e*x + d)^(3/2)*B*a^2*b^2*d*e^3 - 26*(e*x + d)^(3/2)*A*a*b^3*d*e^3 - 57*sqrt(e*x + d)*B*a^2*b^2*d^2*e^3 + 33*sqrt(e*x + d)*A*a*b^3*d^2*e^3 - 17*(e*x + d)^(3/2)*B*a^3*b*e^4 + 13*(e*x + d)^(3/2)*A*a^2*b^2*e^4 + 49*sqrt(e*x + d)*B*a^3*b*d*e^4 - 33*sqrt(e*x + d)*A*a^2*b^2*d*e^4 - 15*sqrt(e*x + d)*B*a^4*e^5 + 11*sqrt(e*x + d)*A*a^3*b*e^5)/(((e*x + d)*b - b*d + a*e)^2*b^5) + 2/15*(3*(e*x + d)^(5/2)*B*b^12*e + 10*(e*x + d)^(3/2)*B*b^12*d*e + 45*sqrt(e*x + d)*B*b^12*d^2*e - 15*(e*x + d)^(3/2)*B*a*b^11*e^2 + 5*(e*x + d)^(3/2)*A*b^12*e^2 - 135*sqrt(e*x + d)*B*a*b^11*d*e^2 + 45*sqrt(e*x + d)*A*b^12*d*e^2 + 90*sqrt(e*x + d)*B*a^2*b^10*e^3 - 45*sqrt(e*x + d)*A*a*b^11*e^3)/b^15
```

Mupad [B] (verification not implemented)

Time = 1.05 (sec) , antiderivative size = 561, normalized size of antiderivative = 2.38

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^3} dx = \left(\frac{2Ae^2 - 2Bde}{3b^3} + \frac{2Be(3b^3d - 3ab^2e)}{3b^6} \right) (d + ex)^{3/2} + \left(\frac{\left(\frac{2Ae^2 - 2Bde}{b^3} + \frac{2Be(3b^3d - 3ab^2e)}{b^6} \right) (3b^3d - 3ab^2e) - \frac{6Be(ae - bd)^2}{b^5}}{b^3} \right) \sqrt{d + ex} - \frac{(d + ex)^{3/2} \left(-\frac{17Ba^3be^4}{4} + \frac{19Ba^2b^2de^3}{2} + \frac{13Aa^2b^2e^4}{4} - \frac{25Bab^3d^2e^2}{4} - \frac{13Aab^3de^3}{2} + Bb^4d^3e + \frac{13Ab^4d^2e^2}{4} \right) - 7e \operatorname{atan} \left(\frac{\sqrt{b}e(ae - bd)^{3/2} \sqrt{d + ex} (5Abe - 9Bae + 4Bbd)}{-9Ba^3e^4 + 22Ba^2bde^3 + 5Aa^2be^4 - 17Bab^2d^2e^2 - 10Aab^2de^3 + 4Bb^3d^3e + 5Ab^3d^2e^2} \right) (ae - bd)^{3/2} (5Abe - 9Bae + 4Bbd)}{4b^{11/2}}$$

input

```
int(((A + B*x)*(d + e*x)^(7/2))/(a + b*x)^3,x)
```

output

```
((2*A*e^2 - 2*B*d*e)/(3*b^3) + (2*B*e*(3*b^3*d - 3*a*b^2*e))/(3*b^6))*(d +
e*x)^(3/2) + (((2*A*e^2 - 2*B*d*e)/b^3 + (2*B*e*(3*b^3*d - 3*a*b^2*e))/b
^6)*(3*b^3*d - 3*a*b^2*e))/b^3 - (6*B*e*(a*e - b*d)^2)/b^5*(d + e*x)^(1/2
) - ((d + e*x)^(3/2)*(B*b^4*d^3*e - (17*B*a^3*b*e^4)/4 + (13*A*a^2*b^2*e^4
)/4 + (13*A*b^4*d^2*e^2)/4 - (25*B*a*b^3*d^2*e^2)/4 + (19*B*a^2*b^2*d*e^3)
/2 - (13*A*a*b^3*d*e^3)/2) - (d + e*x)^(1/2)*((15*B*a^4*e^5)/4 - (11*A*a^3
*b*e^5)/4 + B*b^4*d^4*e + (11*A*b^4*d^3*e^2)/4 - (33*A*a*b^3*d^2*e^3)/4 +
(33*A*a^2*b^2*d*e^4)/4 - (27*B*a*b^3*d^3*e^2)/4 + (57*B*a^2*b^2*d^2*e^3)/4
- (49*B*a^3*b*d*e^4)/4))/(b^7*(d + e*x)^2 - (2*b^7*d - 2*a*b^6*e)*(d + e*
x) + b^7*d^2 + a^2*b^5*e^2 - 2*a*b^6*d*e) + (2*B*e*(d + e*x)^(5/2))/(5*b^3
) + (7*e*atan((b^(1/2)*e*(a*e - b*d)^(3/2)*(d + e*x)^(1/2)*(5*A*b*e - 9*B*
a*e + 4*B*b*d)))/(5*A*a^2*b*e^4 - 9*B*a^3*e^4 + 4*B*b^3*d^3*e + 5*A*b^3*d^2
*e^2 - 17*B*a*b^2*d^2*e^2 - 10*A*a*b^2*d*e^3 + 22*B*a^2*b*d*e^3))*(a*e - b
*d)^(3/2)*(5*A*b*e - 9*B*a*e + 4*B*b*d))/(4*b^(11/2))
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 449, normalized size of antiderivative = 1.90

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^3} dx = \frac{-105\sqrt{b}\sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b}\sqrt{ae-bd}}\right) a^3 e^3 + 210\sqrt{b}\sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b}\sqrt{ae-bd}}\right)}{(a + bx)^3}$$

input

```
int((B*x+A)*(e*x+d)^(7/2)/(b*x+a)^3,x)
```

output

```
( - 105*sqrt(b)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e -
b*d)))*a**3*e**3 + 210*sqrt(b)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sq
rt(b)*sqrt(a*e - b*d)))*a**2*b*d*e**2 - 105*sqrt(b)*sqrt(a*e - b*d)*atan((
sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))*a**2*b*e**3*x - 105*sqrt(b)*sq
rt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))*a*b**2*d**
2*e + 210*sqrt(b)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e
- b*d)))*a*b**2*d*e**2*x - 105*sqrt(b)*sqrt(a*e - b*d)*atan((sqrt(d + e*x
)*b)/(sqrt(b)*sqrt(a*e - b*d)))*b**3*d**2*e*x + 105*sqrt(d + e*x)*a**3*b*e
**3 - 245*sqrt(d + e*x)*a**2*b**2*d*e**2 + 70*sqrt(d + e*x)*a**2*b**2*e**3
*x + 161*sqrt(d + e*x)*a*b**3*d**2*e - 168*sqrt(d + e*x)*a*b**3*d*e**2*x -
14*sqrt(d + e*x)*a*b**3*e**3*x**2 - 15*sqrt(d + e*x)*b**4*d**3 + 116*sqrt
(d + e*x)*b**4*d**2*e*x + 32*sqrt(d + e*x)*b**4*d*e**2*x**2 + 6*sqrt(d + e
*x)*b**4*e**3*x**3)/(15*b**5*(a + b*x))
```

3.165 $\int \frac{(A+Bx)(d+ex)^{5/2}}{(a+bx)^3} dx$

Optimal result	1511
Mathematica [A] (verified)	1512
Rubi [A] (verified)	1512
Maple [A] (verified)	1515
Fricas [A] (verification not implemented)	1516
Sympy [F(-1)]	1517
Maxima [F(-2)]	1517
Giac [B] (verification not implemented)	1518
Mupad [B] (verification not implemented)	1519
Reduce [B] (verification not implemented)	1519

Optimal result

Integrand size = 22, antiderivative size = 196

$$\int \frac{(A+Bx)(d+ex)^{5/2}}{(a+bx)^3} dx = \frac{e(8bBd+5Abe-13aBe)\sqrt{d+ex}}{2b^4} - \frac{(bd-ae)(4bBd+5Abe-9aBe)\sqrt{d+ex}}{4b^4(a+bx)} + \frac{2Be(d+ex)^{3/2}}{3b^3} - \frac{(Ab-aB)(d+ex)^{5/2}}{2b^2(a+bx)^2} - \frac{5e\sqrt{bd-ae}(4bBd+3Abe-7aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{4b^{9/2}}$$

output `1/2*e*(5*A*b*e-13*B*a*e+8*B*b*d)*(e*x+d)^(1/2)/b^4-1/4*(-a*e+b*d)*(5*A*b*e-9*B*a*e+4*B*b*d)*(e*x+d)^(1/2)/b^4/(b*x+a)+2/3*B*e*(e*x+d)^(3/2)/b^3-1/2*(A*b-B*a)*(e*x+d)^(5/2)/b^2/(b*x+a)^2-5/4*e*(-a*e+b*d)^(1/2)*(3*A*b*e-7*B*a*e+4*B*b*d)*arctanh(b^(1/2)*(e*x+d)^(1/2)/(-a*e+b*d)^(1/2))/b^(9/2)`

Mathematica [A] (verified)

Time = 0.62 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.09

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^3} dx =$$

$$\frac{\sqrt{d + ex}(3Ab(-15a^2e^2 + 5abe(d - 5ex) + b^2(2d^2 + 9dex - 8e^2x^2)) + B(105a^3e^2 + 5a^2be(-19d + 35ex) - 4b^3x(-3d^2 + 14d*ex + 2e^2x^2) + a*b^2*(6d^2 - 163d*ex + 56e^2x^2)))}{12b^4(a + bx)^2} - \frac{5e\sqrt{-bd + ae}(4bBd + 3Abe - 7aBe) \arctan\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right)}{4b^{9/2}}$$

input

```
Integrate[((A + B*x)*(d + e*x)^(5/2))/(a + b*x)^3,x]
```

output

```
-1/12*(Sqrt[d + e*x]*(3*A*b*(-15*a^2*e^2 + 5*a*b*e*(d - 5*e*x) + b^2*(2*d^2 + 9*d*e*x - 8*e^2*x^2)) + B*(105*a^3*e^2 + 5*a^2*b*e*(-19*d + 35*e*x) - 4*b^3*x*(-3*d^2 + 14*d*e*x + 2*e^2*x^2) + a*b^2*(6*d^2 - 163*d*e*x + 56*e^2*x^2)))/(b^4*(a + b*x)^2) - (5*e*Sqrt[-(b*d) + a*e]*(4*b*B*d + 3*A*b*e - 7*a*B*e)*ArcTan[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[-(b*d) + a*e]])/(4*b^(9/2))
```

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {87, 51, 60, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^3} dx$$

$$\downarrow 87$$

$$\frac{(-7aBe + 3Abe + 4bBd) \int \frac{(d+ex)^{5/2}}{(a+bx)^2} dx}{4b(bd - ae)} - \frac{(d + ex)^{7/2}(Ab - aB)}{2b(a + bx)^2(bd - ae)}$$

$$\downarrow 51$$

$$\begin{aligned}
 & \frac{(-7aBe + 3Abe + 4bBd) \left(\frac{5e \int \frac{(d+ex)^{3/2}}{a+bx} dx}{2b} - \frac{(d+ex)^{5/2}}{b(a+bx)} \right)}{4b(bd - ae)} - \frac{(d+ex)^{7/2}(Ab - aB)}{2b(a+bx)^2(bd - ae)} \\
 & \quad \downarrow 60 \\
 & \frac{(-7aBe + 3Abe + 4bBd) \left(\frac{5e \left(\frac{(bd-ae) \int \frac{\sqrt{d+ex}}{a+bx} dx}{b} + \frac{2(d+ex)^{3/2}}{3b} \right)}{2b} - \frac{(d+ex)^{5/2}}{b(a+bx)} \right)}{4b(bd - ae)} - \\
 & \quad \frac{(d+ex)^{7/2}(Ab - aB)}{2b(a+bx)^2(bd - ae)} \\
 & \quad \downarrow 60 \\
 & \frac{(-7aBe + 3Abe + 4bBd) \left(\frac{5e \left(\frac{(bd-ae) \left(\frac{1}{(a+bx)\sqrt{d+ex}} + \frac{2\sqrt{d+ex}}{b} \right)}{b} + \frac{2(d+ex)^{3/2}}{3b} \right)}{2b} - \frac{(d+ex)^{5/2}}{b(a+bx)} \right)}{4b(bd - ae)} - \\
 & \quad \frac{(d+ex)^{7/2}(Ab - aB)}{2b(a+bx)^2(bd - ae)} \\
 & \quad \downarrow 73 \\
 & \frac{(-7aBe + 3Abe + 4bBd) \left(\frac{5e \left(\frac{(bd-ae) \left(\frac{2(bd-ae) \int \frac{1}{a + \frac{b(d+ex)}{e} - \frac{bd}{e}} d\sqrt{d+ex}}{b} + \frac{2\sqrt{d+ex}}{b} \right)}{b} + \frac{2(d+ex)^{3/2}}{3b} \right)}{2b} - \frac{(d+ex)^{5/2}}{b(a+bx)} \right)}{4b(bd - ae)} - \\
 & \quad \frac{(d+ex)^{7/2}(Ab - aB)}{2b(a+bx)^2(bd - ae)} \\
 & \quad \downarrow 221
 \end{aligned}$$

$$\begin{aligned}
 & \left(\frac{(-7aBe + 3Abe + 4bBd) \left(\frac{5e \left(\frac{(bd-ae) \left(\frac{2\sqrt{d+ex}}{b} - \frac{2\sqrt{bd-ae} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{3/2}} \right)}{b} + \frac{2(d+ex)^{3/2}}{3b} \right)}{2b} - \frac{(d+ex)^{5/2}}{b(a+bx)} \right)}{4b(bd-ae)} \right. \\
 & \left. \frac{(d+ex)^{7/2}(Ab-aB)}{2b(a+bx)^2(bd-ae)} \right)
 \end{aligned}$$

input `Int[((A + B*x)*(d + e*x)^(5/2))/(a + b*x)^3,x]`

output `-1/2*((A*b - a*B)*(d + e*x)^(7/2))/(b*(b*d - a*e)*(a + b*x)^2) + ((4*b*B*d + 3*A*b*e - 7*a*B*e)*(-(d + e*x)^(5/2)/(b*(a + b*x))) + (5*e*((2*(d + e*x)^(3/2))/(3*b) + ((b*d - a*e)*((2*sqrt[d + e*x])/b - (2*sqrt[b*d - a*e]*ArcTanh[(sqrt[b]*sqrt[d + e*x])/sqrt[b*d - a*e]])/b^(3/2))/b))/(2*b)))/(4*b*(b*d - a*e))`

Defintions of rubi rules used

rule 51 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))] Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(- (b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Maple [A] (verified)

Time = 0.59 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.03

method	result
risch	$\frac{2e(ebBx+3Abe-9Bae+7Bbd)\sqrt{ex+d}}{3b^4} - \frac{(2ae-2db)e \left(\frac{(-\frac{9}{8}Ab^2e+\frac{13}{8}Babe-\frac{1}{2}b^2Bd)(ex+d)^{\frac{3}{2}} + (-\frac{7}{8}Aabe^2+\frac{7}{8}Ab^2de+\frac{1}{8}A^2e^2)}{(ex+d)b+ae-db} \right)}{(ex+d)b+ae-db}$
pseudoelliptic	$-\frac{15 \left((Ae+\frac{4Bd}{3})b-\frac{7Bae}{3} \right) (bx+a)^2 (ae-db)e \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{4} + \frac{15\sqrt{ex+d}\sqrt{(ae-db)b} \left(\frac{8(\frac{Bx}{3}+A)x^2e^2}{15} - \frac{3(-\frac{56Bx}{27}+A)}{5} \right)}{\sqrt{(ae-db)b}b^4}$
derivativedivides	$2e \left(\frac{\frac{bB(ex+d)^{\frac{3}{2}}}{3} + Abe\sqrt{ex+d} - 3Bae\sqrt{ex+d} + 2Bbd\sqrt{ex+d}}{b^4} - \frac{(-\frac{9}{8}b^2e^2aA + \frac{9}{8}Ab^3de + \frac{13}{8}Ba^2be^2 - \frac{17}{8}Bab^2de + \frac{1}{2}b^3Ba^2e^2)}{b^4} \right)$
default	$2e \left(\frac{\frac{bB(ex+d)^{\frac{3}{2}}}{3} + Abe\sqrt{ex+d} - 3Bae\sqrt{ex+d} + 2Bbd\sqrt{ex+d}}{b^4} - \frac{(-\frac{9}{8}b^2e^2aA + \frac{9}{8}Ab^3de + \frac{13}{8}Ba^2be^2 - \frac{17}{8}Bab^2de + \frac{1}{2}b^3Ba^2e^2)}{b^4} \right)$

input `int((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^3,x,method=_RETURNVERBOSE)`

output `2/3*e*(B*b*e*x+3*A*b*e-9*B*a*e+7*B*b*d)*(e*x+d)^(1/2)/b^4-1/b^4*(2*a*e-2*b*d)*e*(((-9/8*A*b^2*e+13/8*B*a*b*e-1/2*b^2*B*d)*(e*x+d)^(3/2)+(-7/8*A*a*b*e^2+7/8*A*b^2*d*e+11/8*B*a^2*e^2-15/8*B*a*b*d*e+1/2*b^2*B*d^2)*(e*x+d)^(1/2)))/((e*x+d)*b+a*e-d*b)^2+5/8*(3*A*b*e-7*B*a*e+4*B*b*d)/((a*e-b*d)*b)^(1/2)*arctan(b*(e*x+d)^(1/2)/((a*e-b*d)*b)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 680, normalized size of antiderivative = 3.47

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^3} dx = \frac{15(4Ba^2bde - (7Ba^3 - 3Aa^2b)e^2 + (4Bb^3de - (7Bab^2 - 3Ab^3)e^2)x^2 + 2(4Bab^2de - (7Ba^2b - 3Aa^2b)e^2)x + (4Bb^3de - (7Bab^2 - 3Ab^3)e^2))}{(a + bx)^3} + \frac{15(4Ba^2bde - (7Ba^3 - 3Aa^2b)e^2 + (4Bb^3de - (7Bab^2 - 3Ab^3)e^2)x^2 + 2(4Bab^2de - (7Ba^2b - 3Aa^2b)e^2)x + (4Bb^3de - (7Bab^2 - 3Ab^3)e^2))}{(a + bx)^3} + \frac{15(4Ba^2bde - (7Ba^3 - 3Aa^2b)e^2 + (4Bb^3de - (7Bab^2 - 3Ab^3)e^2))}{(a + bx)^3}$$

input `integrate((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^3,x, algorithm="fricas")`

output

```
[1/24*(15*(4*B*a^2*b*d*e - (7*B*a^3 - 3*A*a^2*b)*e^2 + (4*B*b^3*d*e - (7*B*a*b^2 - 3*A*b^3)*e^2)*x^2 + 2*(4*B*a*b^2*d*e - (7*B*a^2*b - 3*A*a*b^2)*e^2)*x)*sqrt((b*d - a*e)/b)*log((b*e*x + 2*b*d - a*e - 2*sqrt(e*x + d)*b*sqrt((b*d - a*e)/b))/(b*x + a)) + 2*(8*B*b^3*e^2*x^3 - 6*(B*a*b^2 + A*b^3)*d^2 + 5*(19*B*a^2*b - 3*A*a*b^2)*d*e - 15*(7*B*a^3 - 3*A*a^2*b)*e^2 + 8*(7*B*b^3*d*e - (7*B*a*b^2 - 3*A*b^3)*e^2)*x^2 - (12*B*b^3*d^2 - (163*B*a*b^2 - 27*A*b^3)*d*e + 25*(7*B*a^2*b - 3*A*a*b^2)*e^2)*x)*sqrt(e*x + d))/(b^6*x^2 + 2*a*b^5*x + a^2*b^4), -1/12*(15*(4*B*a^2*b*d*e - (7*B*a^3 - 3*A*a^2*b)*e^2 + (4*B*b^3*d*e - (7*B*a*b^2 - 3*A*b^3)*e^2)*x^2 + 2*(4*B*a*b^2*d*e - (7*B*a^2*b - 3*A*a*b^2)*e^2)*x)*sqrt(-(b*d - a*e)/b)*arctan(-sqrt(e*x + d)*b*sqrt(-(b*d - a*e)/b)/(b*d - a*e)) - (8*B*b^3*e^2*x^3 - 6*(B*a*b^2 + A*b^3)*d^2 + 5*(19*B*a^2*b - 3*A*a*b^2)*d*e - 15*(7*B*a^3 - 3*A*a^2*b)*e^2 + 8*(7*B*b^3*d*e - (7*B*a*b^2 - 3*A*b^3)*e^2)*x^2 - (12*B*b^3*d^2 - (163*B*a*b^2 - 27*A*b^3)*d*e + 25*(7*B*a^2*b - 3*A*a*b^2)*e^2)*x)*sqrt(e*x + d))/(b^6*x^2 + 2*a*b^5*x + a^2*b^4)]
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^3} dx = \text{Timed out}$$

input

```
integrate((B*x+A)*(e*x+d)**(5/2)/(b*x+a)**3,x)
```

output

Timed out

Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^3} dx = \text{Exception raised: ValueError}$$

input

```
integrate((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^3,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for m
ore detail
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 390 vs. $2(168) = 336$.

Time = 0.13 (sec) , antiderivative size = 390, normalized size of antiderivative = 1.99

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^3} dx = \frac{5(4Bb^2d^2e - 11Babde^2 + 3Ab^2de^2 + 7Ba^2e^3 - 3Aabe^3) \arctan\left(\frac{\sqrt{ex+db}}{\sqrt{-b^2d+abe}}\right) + 4(ex+d)^{\frac{3}{2}}Bb^3d^2e - 4\sqrt{ex+d}Bb^3d^3e - 17(ex+d)^{\frac{3}{2}}Bab^2de^2 + 9(ex+d)^{\frac{3}{2}}Ab^3de^2 + 19\sqrt{ex+d}Bab^2de^2 + 2\left((ex+d)^{\frac{3}{2}}Bb^6e + 6\sqrt{ex+d}Bb^6de - 9\sqrt{ex+d}Bab^5e^2 + 3\sqrt{ex+d}Ab^6e^2\right)}{4\sqrt{-b^2d+abe} + 3b^9}$$

input

```
integrate((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^3,x, algorithm="giac")
```

output

```
5/4*(4*B*b^2*d^2*e - 11*B*a*b*d*e^2 + 3*A*b^2*d*e^2 + 7*B*a^2*e^3 - 3*A*a*
b*e^3)*arctan(sqrt(e*x + d)*b/sqrt(-b^2*d + a*b*e))/(sqrt(-b^2*d + a*b*e)*
b^4) - 1/4*(4*(e*x + d)^(3/2)*B*b^3*d^2*e - 4*sqrt(e*x + d)*B*b^3*d^3*e -
17*(e*x + d)^(3/2)*B*a*b^2*d*e^2 + 9*(e*x + d)^(3/2)*A*b^3*d*e^2 + 19*sqrt
(e*x + d)*B*a*b^2*d^2*e^2 - 7*sqrt(e*x + d)*A*b^3*d^2*e^2 + 13*(e*x + d)^(
3/2)*B*a^2*b*e^3 - 9*(e*x + d)^(3/2)*A*a*b^2*e^3 - 26*sqrt(e*x + d)*B*a^2*
b*d*e^3 + 14*sqrt(e*x + d)*A*a*b^2*d*e^3 + 11*sqrt(e*x + d)*B*a^3*e^4 - 7*
sqrt(e*x + d)*A*a^2*b*e^4)/(((e*x + d)*b - b*d + a*e)^2*b^4) + 2/3*((e*x +
d)^(3/2)*B*b^6*e + 6*sqrt(e*x + d)*B*b^6*d*e - 9*sqrt(e*x + d)*B*a*b^5*e^
2 + 3*sqrt(e*x + d)*A*b^6*e^2)/b^9
```

Mupad [B] (verification not implemented)

Time = 1.09 (sec) , antiderivative size = 325, normalized size of antiderivative = 1.66

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^3} dx = \left(\frac{2Ae^2 - 2Bde}{b^3} + \frac{2Be(3b^3d - 3ab^2e)}{b^6} \right) \sqrt{d + ex} - \frac{(d + ex)^{3/2} \left(\frac{13Ba^2be^3}{4} - \frac{17Bab^2de^2}{4} - \frac{9Aab^2e^3}{4} + Bb^3d^2e + \frac{9Ab^3de^2}{4} \right) - \sqrt{d + ex} \left(-\frac{11Ba^3e^4}{4} + \frac{13Ba^2bd}{2} \right)}{b^6(d + ex)^2 - (2b^6d - 2ab^5e)(d + ex) + b^6d^2 + a^2b^5} + \frac{2Be(d + ex)^{3/2}}{3b^3} + \frac{e \operatorname{atan}\left(\frac{\sqrt{b}\sqrt{d+ex}1i}{\sqrt{bd-ae}}\right) \sqrt{bd-ae}(3Abe - 7Bae + 4Bbd) 5i}{4b^{9/2}}$$

input `int(((A + B*x)*(d + e*x)^(5/2))/(a + b*x)^3,x)`output
$$\left(\frac{(2Ae^2 - 2Bde)}{b^3} + \frac{(2Bee(3b^3d - 3ab^2e))}{b^6} \right) (d + ex)^{1/2} - \frac{(d + ex)^{3/2} \left(\frac{13Ba^2be^3}{4} - \frac{9Aa^2be^3}{4} + \frac{9Aab^3de^2}{4} + Bb^3d^2e - \frac{17Bab^2de^2}{4} \right) - (d + ex)^{1/2} \left(\frac{7Aa^2be^4}{4} - \frac{11Ba^3e^4}{4} + Bb^3d^3e + \frac{7Aab^3d^2e^2}{4} - \frac{19Baa^2b^2d^2e^2}{4} - \frac{7Aa^2b^2de^3}{2} + \frac{13Ba^2b^2de^3}{2} \right)}{b^6(d + ex)^2 - (2b^6d - 2ab^5e)(d + ex) + b^6d^2 + a^2b^4e^2 - 2a^2b^5de} + \frac{2Bee(d + ex)^{3/2}}{(3b^3)} + \frac{e \operatorname{atan}\left(\frac{b^{1/2}(d + ex)^{1/2}}{b^2d - ae}\right) (b^2d - ae)^{1/2} (3Abe - 7Bae + 4Bbd) 5i}{(4b^{9/2})}$$
Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.39

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^3} dx = \frac{15\sqrt{b}\sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b}\sqrt{ae-bd}}\right) a^2e^2 - 15\sqrt{b}\sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b}\sqrt{ae-bd}}\right) abd}{(a + bx)^3}$$

input `int((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^3,x)`

output

```
(15*sqrt(b)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d
)))**2**e**2 - 15*sqrt(b)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)
*sqrt(a*e - b*d)))*a*b*d*e + 15*sqrt(b)*sqrt(a*e - b*d)*atan((sqrt(d + e*x
)*b)/(sqrt(b)*sqrt(a*e - b*d)))*a*b*e**2*x - 15*sqrt(b)*sqrt(a*e - b*d)*at
an((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))*b**2*d*e*x - 15*sqrt(d + e
*x)*a**2*b*e**2 + 20*sqrt(d + e*x)*a*b**2*d*e - 10*sqrt(d + e*x)*a*b**2*e*
*2*x - 3*sqrt(d + e*x)*b**3*d**2 + 14*sqrt(d + e*x)*b**3*d*e*x + 2*sqrt(d
+ e*x)*b**3*e**2*x**2)/(3*b**4*(a + b*x))
```

3.166 $\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^3} dx$

Optimal result	1521
Mathematica [A] (verified)	1521
Rubi [A] (verified)	1522
Maple [A] (verified)	1524
Fricas [B] (verification not implemented)	1525
Sympy [F(-1)]	1526
Maxima [F(-2)]	1526
Giac [A] (verification not implemented)	1527
Mupad [B] (verification not implemented)	1527
Reduce [B] (verification not implemented)	1528

Optimal result

Integrand size = 22, antiderivative size = 152

$$\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^3} dx = \frac{2Be\sqrt{d+ex}}{b^3} - \frac{(4bBd+3Abe-7aBe)\sqrt{d+ex}}{4b^3(a+bx)} - \frac{(Ab-aB)(d+ex)^{3/2}}{2b^2(a+bx)^2} - \frac{3e(4bBd+Abe-5aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{4b^{7/2}\sqrt{bd-ae}}$$

output

```
2*B*e*(e*x+d)^(1/2)/b^3-1/4*(3*A*b*e-7*B*a*e+4*B*b*d)*(e*x+d)^(1/2)/b^3/(b*x+a)-1/2*(A*b-B*a)*(e*x+d)^(3/2)/b^2/(b*x+a)^2-3/4*e*(A*b*e-5*B*a*e+4*B*b*d)*arctanh(b^(1/2)*(e*x+d)^(1/2)/(-a*e+b*d)^(1/2))/b^(7/2)/(-a*e+b*d)^(1/2)
```

Mathematica [A] (verified)

Time = 0.55 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.91

$$\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^3} dx = \frac{\sqrt{d+ex}(Ab(2bd+3ae+5bex)+B(-15a^2e+ab(2d-25ex)+4b^2x(d-2ex)))}{4b^3(a+bx)^2} + \frac{3e(4bBd+Abe-5aBe)\operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right)}{4b^{7/2}\sqrt{-bd+ae}}$$

input `Integrate[((A + B*x)*(d + e*x)^(3/2))/(a + b*x)^3,x]`

output `-1/4*(Sqrt[d + e*x]*(A*b*(2*b*d + 3*a*e + 5*b*e*x) + B*(-15*a^2*e + a*b*(2*d - 25*e*x) + 4*b^2*x*(d - 2*e*x)))/(b^3*(a + b*x)^2) + (3*e*(4*b*B*d + A*b*e - 5*a*B*e)*ArcTan[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[-(b*d) + a*e]])/(4*b^(7/2)*Sqrt[-(b*d) + a*e])`

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.09, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.227$, Rules used = {87, 51, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^3} dx \\
 & \quad \downarrow 87 \\
 & \frac{(-5aBe + Abe + 4bBd) \int \frac{(d+ex)^{3/2}}{(a+bx)^2} dx}{4b(bd - ae)} - \frac{(d + ex)^{5/2}(Ab - aB)}{2b(a + bx)^2(bd - ae)} \\
 & \quad \downarrow 51 \\
 & \frac{(-5aBe + Abe + 4bBd) \left(\frac{3e \int \frac{\sqrt{d+ex}}{a+bx} dx}{2b} - \frac{(d+ex)^{3/2}}{b(a+bx)} \right)}{4b(bd - ae)} - \frac{(d + ex)^{5/2}(Ab - aB)}{2b(a + bx)^2(bd - ae)} \\
 & \quad \downarrow 60 \\
 & \frac{(-5aBe + Abe + 4bBd) \left(\frac{3e \left(\frac{(bd-ae) \int \frac{1}{(a+bx)\sqrt{d+ex}} dx}{b} + \frac{2\sqrt{d+ex}}{b} \right)}{2b} - \frac{(d+ex)^{3/2}}{b(a+bx)} \right)}{4b(bd - ae)} - \frac{(d+ex)^{5/2}(Ab - aB)}{2b(a + bx)^2(bd - ae)} \\
 & \quad \downarrow 73
 \end{aligned}$$

$$\begin{aligned}
 & \frac{(-5aBe + Abe + 4bBd) \left(\frac{3e \left(\frac{2(bd-ae) \int \frac{1}{a + \frac{b(d+ex) - bd}{e}} d\sqrt{d+ex} + \frac{2\sqrt{d+ex}}{b}} \right)}{2b} - \frac{(d+ex)^{3/2}}{b(a+bx)} \right)}{4b(bd - ae)} \\
 & \frac{(d + ex)^{5/2}(Ab - aB)}{2b(a + bx)^2(bd - ae)} \\
 & \quad \downarrow \text{221} \\
 & \frac{(-5aBe + Abe + 4bBd) \left(\frac{3e \left(\frac{2\sqrt{d+ex}}{b} - \frac{2\sqrt{bd-ae} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{3/2}} \right)}{2b} - \frac{(d+ex)^{3/2}}{b(a+bx)} \right)}{4b(bd - ae)} \\
 & \frac{(d + ex)^{5/2}(Ab - aB)}{2b(a + bx)^2(bd - ae)}
 \end{aligned}$$

input `Int[((A + B*x)*(d + e*x)^(3/2))/(a + b*x)^3,x]`

output `-1/2*((A*b - a*B)*(d + e*x)^(5/2))/(b*(b*d - a*e)*(a + b*x)^2) + ((4*b*B*d + A*b*e - 5*a*B*e)*(-(d + e*x)^(3/2)/(b*(a + b*x))) + (3*e*((2*sqrt[d + e*x])/b - (2*sqrt[b*d - a*e]*ArcTanh[(sqrt[b]*sqrt[d + e*x])/sqrt[b*d - a*e]])/b^(3/2)))/(2*b))/(4*b*(b*d - a*e))`

Defintions of rubi rules used

rule 51 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))] Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]`

- rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.94

method	result
pseudoelliptic	$\frac{3 \left(-(bx+a)^2 ((Ae+4Bd)b-5Bae)e \arctan \left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}} \right) + \sqrt{ex+d} \left(\frac{(5(-\frac{8Bx}{5}+A)xe+2d(2Bx+A))b^2}{3} + a \left(\left(-\frac{25Bx}{3} + \dots \right) \right) \right)}{4\sqrt{(ae-db)b}b^3(bx+a)^2}$
derivativedivides	$2e \left(\frac{B\sqrt{ex+d}}{b^3} + \frac{(-\frac{5}{8}Ab^2e + \frac{9}{8}Babe - \frac{1}{2}b^2Bd)(ex+d)^{\frac{3}{2}} + (-\frac{3}{8}Aabe^2 + \frac{3}{8}Ab^2de + \frac{7}{8}Ba^2e^2 - \frac{11}{8}Babde + \frac{1}{2}b^2Bd^2)\sqrt{ex+d}}{((ex+d)b+ae-db)^2} + \dots \right)$
default	$2e \left(\frac{B\sqrt{ex+d}}{b^3} + \frac{(-\frac{5}{8}Ab^2e + \frac{9}{8}Babe - \frac{1}{2}b^2Bd)(ex+d)^{\frac{3}{2}} + (-\frac{3}{8}Aabe^2 + \frac{3}{8}Ab^2de + \frac{7}{8}Ba^2e^2 - \frac{11}{8}Babde + \frac{1}{2}b^2Bd^2)\sqrt{ex+d}}{((ex+d)b+ae-db)^2} + \dots \right)$
risch	$\frac{2Be\sqrt{ex+d}}{b^3} + \frac{2e \left(\frac{(-\frac{5}{8}Ab^2e + \frac{9}{8}Babe - \frac{1}{2}b^2Bd)(ex+d)^{\frac{3}{2}} + (-\frac{3}{8}Aabe^2 + \frac{3}{8}Ab^2de + \frac{7}{8}Ba^2e^2 - \frac{11}{8}Babde + \frac{1}{2}b^2Bd^2)\sqrt{ex+d}}{((ex+d)b+ae-db)^2} + \dots \right)}{b^3}$

```
input int((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^3,x,method=_RETURNVERBOSE)
```

```
output -3/4*(-(b*x+a)^2*((A*e+4*B*d)*b-5*B*a*e)*e*arctan(b*(e*x+d)^(1/2)/((a*e-b*d)*b)^(1/2))+
(e*x+d)^(1/2)*(1/3*(5*(-8/5*B*x+A)*x*e+2*d*(2*B*x+A))*b^2+a*((-25/3*B*x+A)*e+2/3*B*d)*b-5*B*a^2*e)*((a*e-b*d)*b)^(1/2)/((a*e-b*d)*b)^(1/2)/b^3/(b*x+a)^2
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 345 vs. 2(130) = 260.

Time = 0.13 (sec) , antiderivative size = 703, normalized size of antiderivative = 4.62

$$\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^3} dx = \left[\frac{3(4Ba^2bde - (5Ba^3 - Aa^2b)e^2 + (4Bb^3de - (5Bab^2 - Ab^3)e^2)x^2 + 2(4 \dots)}{\dots} \right]$$

```
input integrate((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^3,x, algorithm="fricas")
```

output

```
[1/8*(3*(4*B*a^2*b*d*e - (5*B*a^3 - A*a^2*b)*e^2 + (4*B*b^3*d*e - (5*B*a*b^2 - A*b^3)*e^2)*x^2 + 2*(4*B*a*b^2*d*e - (5*B*a^2*b - A*a*b^2)*e^2)*x)*sqrt(b^2*d - a*b*e)*log((b*e*x + 2*b*d - a*e - 2*sqrt(b^2*d - a*b*e)*sqrt(e*x + d))/(b*x + a)) - 2*(2*(B*a*b^3 + A*b^4)*d^2 - (17*B*a^2*b^2 - A*a*b^3)*d*e + 3*(5*B*a^3*b - A*a^2*b^2)*e^2 - 8*(B*b^4*d*e - B*a*b^3*e^2)*x^2 + (4*B*b^4*d^2 - (29*B*a*b^3 - 5*A*b^4)*d*e + 5*(5*B*a^2*b^2 - A*a*b^3)*e^2)*x)*sqrt(e*x + d)/(a^2*b^5*d - a^3*b^4*e + (b^7*d - a*b^6*e)*x^2 + 2*(a*b^6*d - a^2*b^5*e)*x), 1/4*(3*(4*B*a^2*b*d*e - (5*B*a^3 - A*a^2*b)*e^2 + (4*B*b^3*d*e - (5*B*a*b^2 - A*b^3)*e^2)*x^2 + 2*(4*B*a*b^2*d*e - (5*B*a^2*b - A*a*b^2)*e^2)*x)*sqrt(-b^2*d + a*b*e)*arctan(sqrt(-b^2*d + a*b*e)*sqrt(e*x + d)/(b*e*x + b*d)) - (2*(B*a*b^3 + A*b^4)*d^2 - (17*B*a^2*b^2 - A*a*b^3)*d*e + 3*(5*B*a^3*b - A*a^2*b^2)*e^2 - 8*(B*b^4*d*e - B*a*b^3*e^2)*x^2 + (4*B*b^4*d^2 - (29*B*a*b^3 - 5*A*b^4)*d*e + 5*(5*B*a^2*b^2 - A*a*b^3)*e^2)*x)*sqrt(e*x + d)/(a^2*b^5*d - a^3*b^4*e + (b^7*d - a*b^6*e)*x^2 + 2*(a*b^6*d - a^2*b^5*e)*x)]
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^3} dx = \text{Timed out}$$

input

```
integrate((B*x+A)*(e*x+d)**(3/2)/(b*x+a)**3,x)
```

output

Timed out

Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^3} dx = \text{Exception raised: ValueError}$$

input

```
integrate((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^3,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for m
ore detail
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.49

$$\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^3} dx = \frac{2\sqrt{ex+d}Be}{b^3} + \frac{3(4Bbde - 5Bae^2 + Abe^2) \arctan\left(\frac{\sqrt{ex+db}}{\sqrt{-b^2d+abe}}\right)}{4\sqrt{-b^2d+abe}b^3} - \frac{4(ex+d)^{\frac{3}{2}}Bb^2de - 4\sqrt{ex+d}Bb^2d^2e - 9(ex+d)^{\frac{3}{2}}Babe^2 + 5(ex+d)^{\frac{3}{2}}Ab^2e^2 + 11\sqrt{ex+d}Babde^2 - 3}{4((ex+d)b - bd + ae)^2b^3}$$

input

```
integrate((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^3,x, algorithm="giac")
```

output

```
2*sqrt(e*x + d)*B*e/b^3 + 3/4*(4*B*b*d*e - 5*B*a*e^2 + A*b*e^2)*arctan(sqrt
t(e*x + d)*b/sqrt(-b^2*d + a*b*e))/(sqrt(-b^2*d + a*b*e)*b^3) - 1/4*(4*(e*
x + d)^(3/2)*B*b^2*d*e - 4*sqrt(e*x + d)*B*b^2*d^2*e - 9*(e*x + d)^(3/2)*B
*a*b*e^2 + 5*(e*x + d)^(3/2)*A*b^2*e^2 + 11*sqrt(e*x + d)*B*a*b*d*e^2 - 3*
sqrt(e*x + d)*A*b^2*d*e^2 - 7*sqrt(e*x + d)*B*a^2*e^3 + 3*sqrt(e*x + d)*A*
a*b*e^3)/(((e*x + d)*b - b*d + a*e)^2*b^3)
```

Mupad [B] (verification not implemented)

Time = 1.12 (sec) , antiderivative size = 256, normalized size of antiderivative = 1.68

$$\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^3} dx = \frac{\sqrt{d+ex} \left(\frac{7Ba^2e^3}{4} - \frac{11Babde^2}{4} - \frac{3Aabe^3}{4} + Bb^2d^2e + \frac{3Ab^2de^2}{4} \right) - (d+ex)^3}{b^5(d+ex)^2 - (2b^5d - 2ab^4e)(d+ex) + b^5d^2 + a^2b^3} + \frac{2Be\sqrt{d+ex}}{b^3} + \frac{3e \operatorname{atan}\left(\frac{\sqrt{b}\sqrt{d+ex}(Abe-5Bae+4Bbd)}{\sqrt{ae-bd}(Abe^2-5Bae^2+4Bbde)}\right) (Abe-5Bae+4Bbd)}{4b^{7/2}\sqrt{ae-bd}}$$

input `int(((A + B*x)*(d + e*x)^(3/2))/(a + b*x)^3,x)`

output `((d + e*x)^(1/2)*((7*B*a^2*e^3)/4 - (3*A*a*b*e^3)/4 + (3*A*b^2*d*e^2)/4 + B*b^2*d^2*e - (11*B*a*b*d*e^2)/4) - (d + e*x)^(3/2)*((5*A*b^2*e^2)/4 - (9*B*a*b*e^2)/4 + B*b^2*d*e))/(b^5*(d + e*x)^2 - (2*b^5*d - 2*a*b^4*e)*(d + e*x) + b^5*d^2 + a^2*b^3*e^2 - 2*a*b^4*d*e) + (2*B*e*(d + e*x)^(1/2))/b^3 + (3*e*atan((b^(1/2)*e*(d + e*x)^(1/2)*(A*b*e - 5*B*a*e + 4*B*b*d))/((a*e - b*d)^(1/2)*(A*b*e^2 - 5*B*a*e^2 + 4*B*b*d*e)))*(A*b*e - 5*B*a*e + 4*B*b*d))/(4*b^(7/2)*(a*e - b*d)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.84

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^3} dx = \frac{-3\sqrt{b}\sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b}\sqrt{ae-bd}}\right) ae - 3\sqrt{b}\sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b}\sqrt{ae-bd}}\right) bex +}{b^3 (bx + a)}$$

input `int((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^3,x)`

output `(- 3*sqrt(b)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))*a*e - 3*sqrt(b)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))*b*e*x + 3*sqrt(d + e*x)*a*b*e - sqrt(d + e*x)*b**2*d + 2*sqrt(d + e*x)*b**2*e*x)/(b**3*(a + b*x))`

3.167 $\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^3} dx$

Optimal result	1529
Mathematica [A] (verified)	1529
Rubi [A] (verified)	1530
Maple [A] (verified)	1532
Fricas [B] (verification not implemented)	1532
Sympy [F(-1)]	1533
Maxima [F(-2)]	1533
Giac [A] (verification not implemented)	1534
Mupad [B] (verification not implemented)	1534
Reduce [B] (verification not implemented)	1535

Optimal result

Integrand size = 22, antiderivative size = 146

$$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^3} dx = -\frac{(Ab-aB)\sqrt{d+ex}}{2b^2(a+bx)^2} - \frac{(4bBd+Abe-5aBe)\sqrt{d+ex}}{4b^2(bd-ae)(a+bx)} - \frac{e(4bBd-Abe-3aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{4b^{5/2}(bd-ae)^{3/2}}$$

output

```
-1/2*(A*b-B*a)*(e*x+d)^(1/2)/b^2/(b*x+a)^2-1/4*(A*b*e-5*B*a*e+4*B*b*d)*(e*x+d)^(1/2)/b^2/(-a*e+b*d)/(b*x+a)-1/4*e*(-A*b*e-3*B*a*e+4*B*b*d)*arctanh(b^(1/2)*(e*x+d)^(1/2)/(-a*e+b*d)^(1/2))/b^(5/2)/(-a*e+b*d)^(3/2)
```

Mathematica [A] (verified)

Time = 0.61 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.98

$$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^3} dx = \frac{\sqrt{b}\sqrt{d+ex}(Ab(2bd-ae+bx)+B(-3a^2e+4b^2dx+ab(2d-5ex)))}{(-bd+ae)(a+bx)^2} + \frac{e(-4bBd+Abe+3aBe)\operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right)}{(-bd+ae)^{3/2}}}{4b^{5/2}}$$

input `Integrate[((A + B*x)*Sqrt[d + e*x])/(a + b*x)^3,x]`

output `((Sqrt[b]*Sqrt[d + e*x]*(A*b*(2*b*d - a*e + b*e*x) + B*(-3*a^2*e + 4*b^2*d*x + a*b*(2*d - 5*e*x)))/((-b*d) + a*e)*(a + b*x)^2 + (e*(-4*b*B*d + A*b*e + 3*a*B*e)*ArcTan[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[-(b*d) + a*e]])/(-(b*d) + a*e)^(3/2))/(4*b^(5/2))`

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.99, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {87, 51, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^3} dx \\
 & \quad \downarrow 87 \\
 & \frac{(-3aBe - Abe + 4bBd) \int \frac{\sqrt{d+ex}}{(a+bx)^2} dx}{4b(bd - ae)} - \frac{(d + ex)^{3/2}(Ab - aB)}{2b(a + bx)^2(bd - ae)} \\
 & \quad \downarrow 51 \\
 & \frac{(-3aBe - Abe + 4bBd) \left(\frac{e \int \frac{1}{(a+bx)\sqrt{d+ex}} dx}{2b} - \frac{\sqrt{d+ex}}{b(a+bx)} \right)}{4b(bd - ae)} - \frac{(d + ex)^{3/2}(Ab - aB)}{2b(a + bx)^2(bd - ae)} \\
 & \quad \downarrow 73 \\
 & \frac{(-3aBe - Abe + 4bBd) \left(\frac{\int \frac{1}{a + \frac{b(d+ex)}{e} - \frac{bd}{e}} d\sqrt{d+ex}}{b} - \frac{\sqrt{d+ex}}{b(a+bx)} \right)}{4b(bd - ae)} - \frac{(d + ex)^{3/2}(Ab - aB)}{2b(a + bx)^2(bd - ae)} \\
 & \quad \downarrow 221 \\
 & \frac{(-3aBe - Abe + 4bBd) \left(-\frac{e \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{b^{3/2}\sqrt{bd-ae}} - \frac{\sqrt{d+ex}}{b(a+bx)} \right)}{4b(bd - ae)} - \frac{(d + ex)^{3/2}(Ab - aB)}{2b(a + bx)^2(bd - ae)}
 \end{aligned}$$

input `Int[((A + B*x)*Sqrt[d + e*x])/(a + b*x)^3,x]`

output `-1/2*((A*b - a*B)*(d + e*x)^(3/2))/(b*(b*d - a*e)*(a + b*x)^2) + ((4*b*B*d - A*b*e - 3*a*B*e)*(-(Sqrt[d + e*x]/(b*(a + b*x))) - (e*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[b*d - a*e]])/(b^(3/2)*Sqrt[b*d - a*e]))/(4*b*(b*d - a*e))`

Defintions of rubi rules used

rule 51 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))] Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.01

method	result
pseudoelliptic	$-\frac{(bx+a)^2 e((Ae-4Bd)b+3Bae) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right) + \sqrt{ex+d} \sqrt{(ae-db)b} ((-Aex-2d(2Bx+A))b^2+a((5Bx+A)))}{4\sqrt{(ae-db)b} b^2 (ae-db)(bx+a)^2}$
derivativedivides	$2e \left(\frac{\frac{(Abe-5Bae+4Bbd)(ex+d)^{\frac{3}{2}}}{8(ae-db)b} - \frac{(Abe+3Bae-4Bbd)\sqrt{ex+d}}{8b^2}}{(ex+d)b+ae-db)^2} + \frac{(Abe+3Bae-4Bbd) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{8(ae-db)b^2 \sqrt{(ae-db)b}} \right)$
default	$2e \left(\frac{\frac{(Abe-5Bae+4Bbd)(ex+d)^{\frac{3}{2}}}{8(ae-db)b} - \frac{(Abe+3Bae-4Bbd)\sqrt{ex+d}}{8b^2}}{(ex+d)b+ae-db)^2} + \frac{(Abe+3Bae-4Bbd) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{8(ae-db)b^2 \sqrt{(ae-db)b}} \right)$

input `int((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^3,x,method=_RETURNVERBOSE)`

output `-1/4*(-(b*x+a)^2*e*((A*e-4*B*d)*b+3*B*a*e)*arctan(b*(e*x+d)^(1/2)/((a*e-b*d)*b)^(1/2))+e*(x+d)^(1/2)*((a*e-b*d)*b)^(1/2)*((-A*e*x-2*d*(2*B*x+A))*b^2+a*((5*B*x+A)*e-2*B*d)*b+3*B*a^2*e)/((a*e-b*d)*b)^(1/2)/b^2/(a*e-b*d)/(b*x+a)^2`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 354 vs. 2(126) = 252.

Time = 0.10 (sec) , antiderivative size = 721, normalized size of antiderivative = 4.94

$$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^3} dx = \left[\frac{(4Ba^2bde - (3Ba^3 + Aa^2b)e^2 + (4Bb^3de - (3Bab^2 + Ab^3)e^2)x^2 + 2(4Bab^2de - (3Ba^2b + Aab^2)e^2))}{8(a^2b^5)} \right]$$

input `integrate((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^3,x,algorithm="fricas")`

output

```
[1/8*((4*B*a^2*b*d*e - (3*B*a^3 + A*a^2*b)*e^2 + (4*B*b^3*d*e - (3*B*a*b^2 + A*b^3)*e^2)*x^2 + 2*(4*B*a*b^2*d*e - (3*B*a^2*b + A*a*b^2)*e^2)*x)*sqrt(b^2*d - a*b*e)*log((b*e*x + 2*b*d - a*e - 2*sqrt(b^2*d - a*b*e)*sqrt(e*x + d))/(b*x + a)) - 2*(2*(B*a*b^3 + A*b^4)*d^2 - (5*B*a^2*b^2 + 3*A*a*b^3)*d*e + (3*B*a^3*b + A*a^2*b^2)*e^2 + (4*B*b^4*d^2 - (9*B*a*b^3 - A*b^4)*d*e + (5*B*a^2*b^2 - A*a*b^3)*e^2)*x)*sqrt(e*x + d))/(a^2*b^5*d^2 - 2*a^3*b^4*d*e + a^4*b^3*e^2 + (b^7*d^2 - 2*a*b^6*d*e + a^2*b^5*e^2)*x^2 + 2*(a*b^6*d^2 - 2*a^2*b^5*d*e + a^3*b^4*e^2)*x), 1/4*((4*B*a^2*b*d*e - (3*B*a^3 + A*a^2*b)*e^2 + (4*B*b^3*d*e - (3*B*a*b^2 + A*b^3)*e^2)*x^2 + 2*(4*B*a*b^2*d*e - (3*B*a^2*b + A*a*b^2)*e^2)*x)*sqrt(-b^2*d + a*b*e)*arctan(sqrt(-b^2*d + a*b*e)*sqrt(e*x + d)/(b*e*x + b*d)) - (2*(B*a*b^3 + A*b^4)*d^2 - (5*B*a^2*b^2 + 3*A*a*b^3)*d*e + (3*B*a^3*b + A*a^2*b^2)*e^2 + (4*B*b^4*d^2 - (9*B*a*b^3 - A*b^4)*d*e + (5*B*a^2*b^2 - A*a*b^3)*e^2)*x)*sqrt(e*x + d))/(a^2*b^5*d^2 - 2*a^3*b^4*d*e + a^4*b^3*e^2 + (b^7*d^2 - 2*a*b^6*d*e + a^2*b^5*e^2)*x^2 + 2*(a*b^6*d^2 - 2*a^2*b^5*d*e + a^3*b^4*e^2)*x)]
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^3} dx = \text{Timed out}$$

input

```
integrate((B*x+A)*(e*x+d)**(1/2)/(b*x+a)**3,x)
```

output

Timed out

Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^3} dx = \text{Exception raised: ValueError}$$

input

```
integrate((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^3,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for m
ore detail
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.61

$$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^3} dx = \frac{(4Bbde - 3Bae^2 - Abe^2) \arctan\left(\frac{\sqrt{ex+db}}{\sqrt{-b^2d+abe}}\right)}{4(b^3d - ab^2e)\sqrt{-b^2d+abe}} - \frac{4(ex+d)^{\frac{3}{2}}Bb^2de - 4\sqrt{ex+d}Bb^2d^2e - 5(ex+d)^{\frac{3}{2}}Babe^2 + (ex+d)^{\frac{3}{2}}Ab^2e^2 + 7\sqrt{ex+d}Babde^2 + \sqrt{ex+d}A^2b^2e^2}{4(b^3d - ab^2e)((ex+d)b - bd + ae)^2}$$

input

```
integrate((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^3,x, algorithm="giac")
```

output

```
1/4*(4*B*b*d*e - 3*B*a*e^2 - A*b*e^2)*arctan(sqrt(e*x + d)*b/sqrt(-b^2*d +
a*b*e))/((b^3*d - a*b^2*e)*sqrt(-b^2*d + a*b*e)) - 1/4*(4*(e*x + d)^(3/2)
*B*b^2*d*e - 4*sqrt(e*x + d)*B*b^2*d^2*e - 5*(e*x + d)^(3/2)*B*a*b*e^2 + (
e*x + d)^(3/2)*A*b^2*e^2 + 7*sqrt(e*x + d)*B*a*b*d*e^2 + sqrt(e*x + d)*A*b
^2*d*e^2 - 3*sqrt(e*x + d)*B*a^2*e^3 - sqrt(e*x + d)*A*a*b*e^3)/((b^3*d -
a*b^2*e)*((e*x + d)*b - b*d + a*e)^2)
```

Mupad [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.52

$$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^3} dx = \frac{e \operatorname{atan}\left(\frac{\sqrt{be}\sqrt{d+ex}(Abe+3Bae-4Bbd)}{\sqrt{ae-bd}(Abe^2+3Bae^2-4Bbde)}\right) (Abe+3Bae-4Bbd)}{4b^{5/2}(ae-bd)^{3/2}} - \frac{\frac{\sqrt{d+ex}(Abe^2+3Bae^2-4Bbde)}{4b^2} - \frac{(d+ex)^{3/2}(Abe^2-5Bae^2+4Bbde)}{4b(ae-bd)}}{b^2(d+ex)^2 - (2b^2d - 2abe)(d+ex) + a^2e^2 + b^2d^2 - 2abde}$$

input `int(((A + B*x)*(d + e*x)^(1/2))/(a + b*x)^3,x)`

output
$$\frac{(e \operatorname{atan}\left(\frac{b^{1/2} e (d + e x)^{1/2} (A b e + 3 B a e - 4 B b d)}{(a e - b d)^{1/2} (A b e^2 + 3 B a e^2 - 4 B b d e)}\right) + (A b e + 3 B a e - 4 B b d) / (4 b^{5/2} (a e - b d)^{3/2}) - ((d + e x)^{1/2} (A b e^2 + 3 B a e^2 - 4 B b d e)) / (4 b^2) - ((d + e x)^{3/2} (A b e^2 - 5 B a e^2 + 4 B b d e)) / (4 b (a e - b d))}{(b^2 (d + e x)^2 - (2 b^2 d - 2 a b e) (d + e x) + a^2 e^2 + b^2 d^2 - 2 a b d e)}$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.88

$$\int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^3} dx$$

$$= \frac{\sqrt{b} \sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b} \sqrt{ae-bd}}\right) ae + \sqrt{b} \sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b} \sqrt{ae-bd}}\right) bex - \sqrt{ex + d} abe + \sqrt{ex + d} b^2 d}{b^2 (abex - b^2 dx + a^2 e - abd)}$$

input `int((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^3,x)`

output
$$\frac{(\operatorname{sqrt}(b) \operatorname{sqrt}(a e - b d) \operatorname{atan}\left(\frac{\operatorname{sqrt}(d + e x) b}{\operatorname{sqrt}(b) \operatorname{sqrt}(a e - b d)}\right)) a e + \operatorname{sqrt}(b) \operatorname{sqrt}(a e - b d) \operatorname{atan}\left(\frac{\operatorname{sqrt}(d + e x) b}{\operatorname{sqrt}(b) \operatorname{sqrt}(a e - b d)}\right) b e x - \operatorname{sqrt}(d + e x) a b e + \operatorname{sqrt}(d + e x) b^2 d}{b^2 (a^2 e - a b d + a b e x - b^2 d x)}$$

3.168 $\int \frac{A+Bx}{(a+bx)^3\sqrt{d+ex}} dx$

Optimal result	1536
Mathematica [A] (verified)	1536
Rubi [A] (verified)	1537
Maple [A] (verified)	1539
Fricas [B] (verification not implemented)	1539
Sympy [F(-1)]	1540
Maxima [F(-2)]	1540
Giac [A] (verification not implemented)	1541
Mupad [B] (verification not implemented)	1541
Reduce [B] (verification not implemented)	1542

Optimal result

Integrand size = 22, antiderivative size = 157

$$\int \frac{A+Bx}{(a+bx)^3\sqrt{d+ex}} dx = -\frac{(Ab-aB)\sqrt{d+ex}}{2b(bd-ae)(a+bx)^2} - \frac{(4bBd-3Abe-aBe)\sqrt{d+ex}}{4b(bd-ae)^2(a+bx)} + \frac{e(4bBd-3Abe-aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{4b^{3/2}(bd-ae)^{5/2}}$$

output

```
-1/2*(A*b-B*a)*(e*x+d)^(1/2)/b/(-a*e+b*d)/(b*x+a)^2-1/4*(-3*A*b*e-B*a*e+4*B*b*d)*(e*x+d)^(1/2)/b/(-a*e+b*d)^2/(b*x+a)+1/4*e*(-3*A*b*e-B*a*e+4*B*b*d)*arctanh(b^(1/2)*(e*x+d)^(1/2)/(-a*e+b*d)^(1/2))/b^(3/2)/(-a*e+b*d)^(5/2)
```

Mathematica [A] (verified)

Time = 0.63 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.92

$$\int \frac{A+Bx}{(a+bx)^3\sqrt{d+ex}} dx = \frac{\sqrt{b}\sqrt{d+ex}(Ab(-2bd+5ae+3bex)-B(a^2e+4b^2dx+ab(2d-ex)))}{(bd-ae)^2(a+bx)^2} + \frac{e(-4bBd+3Abe+aBe)\arctan\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right)}{(-bd+ae)^{5/2}}$$

$4b^{3/2}$

input `Integrate[(A + B*x)/((a + b*x)^3*Sqrt[d + e*x]),x]`

output `((Sqrt[b]*Sqrt[d + e*x]*(A*b*(-2*b*d + 5*a*e + 3*b*e*x) - B*(a^2*e + 4*b^2*d*x + a*b*(2*d - e*x)))/((b*d - a*e)^2*(a + b*x)^2) + (e*(-4*b*B*d + 3*A*b*e + a*B*e)*ArcTan[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[-(b*d) + a*e]]/(-(b*d) + a*e)^(5/2))/(4*b^(3/2))`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.96, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {87, 52, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + Bx}{(a + bx)^3 \sqrt{d + ex}} dx \\
 & \quad \downarrow 87 \\
 & \frac{(-aBe - 3Abe + 4bBd) \int \frac{1}{(a+bx)^2 \sqrt{d+ex}} dx}{4b(bd - ae)} - \frac{\sqrt{d + ex}(Ab - aB)}{2b(a + bx)^2(bd - ae)} \\
 & \quad \downarrow 52 \\
 & \frac{(-aBe - 3Abe + 4bBd) \left(-\frac{e \int \frac{1}{(a+bx)\sqrt{d+ex}} dx}{2(bd-ae)} - \frac{\sqrt{d+ex}}{(a+bx)(bd-ae)} \right)}{4b(bd - ae)} - \frac{\sqrt{d + ex}(Ab - aB)}{2b(a + bx)^2(bd - ae)} \\
 & \quad \downarrow 73 \\
 & \frac{(-aBe - 3Abe + 4bBd) \left(-\frac{\int \frac{1}{a + \frac{b(d+ex) - bd}{e}} d\sqrt{d+ex}}{bd-ae} - \frac{\sqrt{d+ex}}{(a+bx)(bd-ae)} \right)}{4b(bd - ae)} - \frac{\sqrt{d + ex}(Ab - aB)}{2b(a + bx)^2(bd - ae)} \\
 & \quad \downarrow 221 \\
 & \frac{(-aBe - 3Abe + 4bBd) \left(\frac{e \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{\sqrt{b}(bd-ae)^{3/2}} - \frac{\sqrt{d+ex}}{(a+bx)(bd-ae)} \right)}{4b(bd - ae)} - \frac{\sqrt{d + ex}(Ab - aB)}{2b(a + bx)^2(bd - ae)}
 \end{aligned}$$

input `Int[(A + B*x)/((a + b*x)^3*Sqrt[d + e*x]),x]`

output `-1/2*((A*b - a*B)*Sqrt[d + e*x])/(b*(b*d - a*e)*(a + b*x)^2) + ((4*b*B*d - 3*A*b*e - a*B*e)*(-(Sqrt[d + e*x]/((b*d - a*e)*(a + b*x))) + (e*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[b*d - a*e]])/(Sqrt[b]*(b*d - a*e)^(3/2))))/(4*b*(b*d - a*e))`

Defintions of rubi rules used

rule 52 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.94

method	result
pseudoelliptic	$\frac{3(bx+a)^2 \left((Ae - \frac{4Bd}{3})b + \frac{Bae}{3} \right) e \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right) + 5\sqrt{ex+d} \sqrt{(ae-db)b} \left(\frac{(3Aex-2d(2Bx+A))b^2}{5} + a \left(\left(\frac{Bx}{5} + A\right)e - \frac{2Bd}{5} \right) b - \dots \right)}{(ae-db)^2 \sqrt{(ae-db)b} b(bx+a)^2}$
derivativedivides	$2e \left(\frac{\left(\frac{(3Abe+Bae-4Bbd)(ex+d)^{\frac{3}{2}}}{8a^2e^2-16abde+8b^2d^2} + \frac{(5Abe-Bae-4Bbd)\sqrt{ex+d}}{8(ae-db)b} \right)}{((ex+d)b+ae-db)^2} + \frac{(3Abe+Bae-4Bbd) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{8(a^2e^2-2abde+b^2d^2)b\sqrt{(ae-db)b}} \right)$
default	$2e \left(\frac{\left(\frac{(3Abe+Bae-4Bbd)(ex+d)^{\frac{3}{2}}}{8a^2e^2-16abde+8b^2d^2} + \frac{(5Abe-Bae-4Bbd)\sqrt{ex+d}}{8(ae-db)b} \right)}{((ex+d)b+ae-db)^2} + \frac{(3Abe+Bae-4Bbd) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{8(a^2e^2-2abde+b^2d^2)b\sqrt{(ae-db)b}} \right)$

input `int((B*x+A)/(b*x+a)^3/(e*x+d)^(1/2),x,method=_RETURNVERBOSE)`

output
$$\frac{5}{4} \frac{(a^2e-b^2d)^2 (3/5(bx+a)^2 ((Ae-4/3Bd)b + 1/3Bae) e \arctan(b\sqrt{ex+d}/\sqrt{(ae-db)b}) + (e^2x+d)\sqrt{(ae-db)b} ((1/5Bx+A)e - 2/5Bd)b - 1/5Bae^2)}{(a^2e-b^2d)^2 (b^2x+ab+ae)^2}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 397 vs. 2(137) = 274.

Time = 0.15 (sec) , antiderivative size = 808, normalized size of antiderivative = 5.15

$$\int \frac{A + Bx}{(a + bx)^3 \sqrt{d + ex}} dx = \text{Too large to display}$$

input `integrate((B*x+A)/(b*x+a)^3/(e*x+d)^(1/2),x, algorithm="fricas")`

output

```
[ -1/8*((4*B*a^2*b*d*e - (B*a^3 + 3*A*a^2*b)*e^2 + (4*B*b^3*d*e - (B*a*b^2 + 3*A*b^3)*e^2)*x^2 + 2*(4*B*a*b^2*d*e - (B*a^2*b + 3*A*a*b^2)*e^2)*x)*sqrt(b^2*d - a*b*e)*log((b*e*x + 2*b*d - a*e - 2*sqrt(b^2*d - a*b*e)*sqrt(e*x + d))/(b*x + a)) + 2*(2*(B*a*b^3 + A*b^4)*d^2 - (B*a^2*b^2 + 7*A*a*b^3)*d*e - (B*a^3*b - 5*A*a^2*b^2)*e^2 + (4*B*b^4*d^2 - (5*B*a*b^3 + 3*A*b^4)*d*e + (B*a^2*b^2 + 3*A*a*b^3)*e^2)*x)*sqrt(e*x + d))/(a^2*b^5*d^3 - 3*a^3*b^4*d^2*e + 3*a^4*b^3*d*e^2 - a^5*b^2*e^3 + (b^7*d^3 - 3*a*b^6*d^2*e + 3*a^2*b^5*d*e^2 - a^3*b^4*e^3)*x^2 + 2*(a*b^6*d^3 - 3*a^2*b^5*d^2*e + 3*a^3*b^4*d*e^2 - a^4*b^3*e^3)*x), -1/4*((4*B*a^2*b*d*e - (B*a^3 + 3*A*a^2*b)*e^2 + (4*B*b^3*d*e - (B*a*b^2 + 3*A*b^3)*e^2)*x^2 + 2*(4*B*a*b^2*d*e - (B*a^2*b + 3*A*a*b^2)*e^2)*x)*sqrt(-b^2*d + a*b*e)*arctan(sqrt(-b^2*d + a*b*e)*sqrt(e*x + d)/(b*e*x + b*d)) + (2*(B*a*b^3 + A*b^4)*d^2 - (B*a^2*b^2 + 7*A*a*b^3)*d*e - (B*a^3*b - 5*A*a^2*b^2)*e^2 + (4*B*b^4*d^2 - (5*B*a*b^3 + 3*A*b^4)*d*e + (B*a^2*b^2 + 3*A*a*b^3)*e^2)*x)*sqrt(e*x + d))/(a^2*b^5*d^3 - 3*a^3*b^4*d^2*e + 3*a^4*b^3*d*e^2 - a^5*b^2*e^3 + (b^7*d^3 - 3*a*b^6*d^2*e + 3*a^2*b^5*d*e^2 - a^3*b^4*e^3)*x^2 + 2*(a*b^6*d^3 - 3*a^2*b^5*d^2*e + 3*a^3*b^4*d*e^2 - a^4*b^3*e^3)*x]
```

Sympy [F(-1)]

Timed out.

$$\int \frac{A + Bx}{(a + bx)^3 \sqrt{d + ex}} dx = \text{Timed out}$$

input

```
integrate((B*x+A)/(b*x+a)**3/(e*x+d)**(1/2),x)
```

output

Timed out

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{(a + bx)^3 \sqrt{d + ex}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((B*x+A)/(b*x+a)^3/(e*x+d)^(1/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for m
ore detail
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 258, normalized size of antiderivative = 1.64

$$\int \frac{A + Bx}{(a + bx)^3 \sqrt{d + ex}} dx = -\frac{(4Bbde - Bae^2 - 3Abe^2) \arctan\left(\frac{\sqrt{ex+db}}{\sqrt{-b^2d+abe}}\right)}{4(b^3d^2 - 2ab^2de + a^2be^2)\sqrt{-b^2d+abe}} - \frac{4(ex+d)^{\frac{3}{2}}Bb^2de - 4\sqrt{ex+d}Bb^2d^2e - (ex+d)^{\frac{3}{2}}Babe^2 - 3(ex+d)^{\frac{3}{2}}Ab^2e^2 + 3\sqrt{ex+d}Babde^2 + 5}{4(b^3d^2 - 2ab^2de + a^2be^2)((ex+d)b - bd + ae)^2}$$

input

```
integrate((B*x+A)/(b*x+a)^3/(e*x+d)^(1/2),x, algorithm="giac")
```

output

```
-1/4*(4*B*b*d*e - B*a*e^2 - 3*A*b*e^2)*arctan(sqrt(e*x + d)*b/sqrt(-b^2*d
+ a*b*e))/((b^3*d^2 - 2*a*b^2*d*e + a^2*b*e^2)*sqrt(-b^2*d + a*b*e)) - 1/4
*(4*(e*x + d)^(3/2)*B*b^2*d*e - 4*sqrt(e*x + d)*B*b^2*d^2*e - (e*x + d)^(3
/2)*B*a*b*e^2 - 3*(e*x + d)^(3/2)*A*b^2*e^2 + 3*sqrt(e*x + d)*B*a*b*d*e^2
+ 5*sqrt(e*x + d)*A*b^2*d*e^2 + sqrt(e*x + d)*B*a^2*e^3 - 5*sqrt(e*x + d)*
A*a*b*e^3)/((b^3*d^2 - 2*a*b^2*d*e + a^2*b*e^2)*((e*x + d)*b - b*d + a*e)^
2)
```

Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.45

$$\int \frac{A + Bx}{(a + bx)^3 \sqrt{d + ex}} dx = \frac{(d+ex)^{3/2} (3Abe^2 + Bae^2 - 4Bbde)}{4(ae-bd)^2} - \frac{\sqrt{d+ex} (Bae^2 - 5Abe^2 + 4Bbde)}{4b(ae-bd)} + \frac{e \operatorname{atan}\left(\frac{\sqrt{be}\sqrt{d+ex}(3Abe+Bae-4Bbd)}{\sqrt{ae-bd}(3Abe^2+Bae^2-4Bbde)}\right) (3Abe + Bae - 4Bbd)}{4b^{3/2}(ae-bd)^{5/2}}$$

input `int((A + B*x)/((a + b*x)^3*(d + e*x)^(1/2)),x)`

output `((((d + e*x)^(3/2)*(3*A*b*e^2 + B*a*e^2 - 4*B*b*d*e))/(4*(a*e - b*d)^2) - ((d + e*x)^(1/2)*(B*a*e^2 - 5*A*b*e^2 + 4*B*b*d*e))/(4*b*(a*e - b*d)))/(b^2*(d + e*x)^2 - (2*b^2*d - 2*a*b*e)*(d + e*x) + a^2*e^2 + b^2*d^2 - 2*a*b*d*e) + (e*atan((b^(1/2)*e*(d + e*x)^(1/2)*(3*A*b*e + B*a*e - 4*B*b*d))/(a*e - b*d)^(1/2)*(3*A*b*e^2 + B*a*e^2 - 4*B*b*d*e)))*(3*A*b*e + B*a*e - 4*B*b*d))/(4*b^(3/2)*(a*e - b*d)^(5/2))`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.99

$$\int \frac{A + Bx}{(a + bx)^3 \sqrt{d + ex}} dx$$

$$= \frac{\sqrt{b} \sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b} \sqrt{ae-bd}}\right) ae + \sqrt{b} \sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b} \sqrt{ae-bd}}\right) be x + \sqrt{ex + d} abe - \sqrt{ex + d} b^2 d}{b(a^2 b e^2 x - 2a b^2 d e x + b^3 d^2 x + a^3 e^2 - 2a^2 b d e + a b^2 d^2)}$$

input `int((B*x+A)/(b*x+a)^3/(e*x+d)^(1/2),x)`

output `(sqrt(b)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))*a*e + sqrt(b)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))*b*e*x + sqrt(d + e*x)*a*b*e - sqrt(d + e*x)*b**2*d)/(b*(a**3*e**2 - 2*a**2*b*d*e + a**2*b*e**2*x + a*b**2*d**2 - 2*a*b**2*d*e*x + b**3*d**2*x))`

3.169 $\int \frac{A+Bx}{(a+bx)^3(d+ex)^{3/2}} dx$

Optimal result	1543
Mathematica [A] (verified)	1543
Rubi [A] (verified)	1544
Maple [A] (verified)	1546
Fricas [B] (verification not implemented)	1547
Sympy [F(-1)]	1548
Maxima [F(-2)]	1549
Giac [B] (verification not implemented)	1549
Mupad [B] (verification not implemented)	1550
Reduce [B] (verification not implemented)	1550

Optimal result

Integrand size = 22, antiderivative size = 180

$$\int \frac{A+Bx}{(a+bx)^3(d+ex)^{3/2}} dx = -\frac{2e(Bd-Ae)}{(bd-ae)^3\sqrt{d+ex}} - \frac{(Ab-aB)\sqrt{d+ex}}{2(bd-ae)^2(a+bx)^2}$$

$$- \frac{(4bBd-7Abe+3aBe)\sqrt{d+ex}}{4(bd-ae)^3(a+bx)} + \frac{3e(4bBd-5Abe+aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{4\sqrt{b}(bd-ae)^{7/2}}$$

output

```

-2*e*(-A*e+B*d)/(-a*e+b*d)^3/(e*x+d)^(1/2)-1/2*(A*b-B*a)*(e*x+d)^(1/2)/(-a
*e+b*d)^2/(b*x+a)^2-1/4*(-7*A*b*e+3*B*a*e+4*B*b*d)*(e*x+d)^(1/2)/(-a*e+b*d
)^3/(b*x+a)+3/4*e*(-5*A*b*e+B*a*e+4*B*b*d)*arctanh(b^(1/2)*(e*x+d)^(1/2)/(
-a*e+b*d)^(1/2))/b^(1/2)/(-a*e+b*d)^(7/2)
    
```

Mathematica [A] (verified)

Time = 0.96 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.08

$$\int \frac{A+Bx}{(a+bx)^3(d+ex)^{3/2}} dx = \frac{1}{4} \left(\frac{-B(4b^2dx(d+3ex) + a^2e(13d+5ex) + ab(2d^2+21dex+3e^2x^2)) + A((bd-ae)^3(a+bx)^2\sqrt{d+ex})}{(bd-ae)^3(a+bx)^2\sqrt{d+ex}} \right.$$

$$\left. + \frac{3e(4bBd-5Abe+aBe)\arctan\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right)}{\sqrt{b}(-bd+ae)^{7/2}} \right)$$

input `Integrate[(A + B*x)/((a + b*x)^3*(d + e*x)^(3/2)),x]`

output `((-(B*(4*b^2*d*x*(d + 3*e*x) + a^2*e*(13*d + 5*e*x) + a*b*(2*d^2 + 21*d*e*x + 3*e^2*x^2))) + A*(8*a^2*e^2 + a*b*e*(9*d + 25*e*x) + b^2*(-2*d^2 + 5*d*e*x + 15*e^2*x^2)))/((b*d - a*e)^3*(a + b*x)^2*sqrt[d + e*x]) + (3*e*(4*b*B*d - 5*A*b*e + a*B*e)*ArcTan[(sqrt[b]*sqrt[d + e*x])/sqrt[-(b*d) + a*e]])/(sqrt[b]*(-(b*d) + a*e)^(7/2))/4`

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.04, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.227$, Rules used = {87, 52, 61, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + Bx}{(a + bx)^3(d + ex)^{3/2}} dx \\
 & \quad \downarrow 87 \\
 & \frac{(aBe - 5Abe + 4bBd) \int \frac{1}{(a+bx)^2(d+ex)^{3/2}} dx}{4b(bd - ae)} - \frac{Ab - aB}{2b(a + bx)^2\sqrt{d + ex}(bd - ae)} \\
 & \quad \downarrow 52 \\
 & \frac{(aBe - 5Abe + 4bBd) \left(-\frac{3e \int \frac{1}{(a+bx)(d+ex)^{3/2}} dx}{2(bd-ae)} - \frac{1}{(a+bx)\sqrt{d+ex}(bd-ae)} \right)}{4b(bd - ae)} - \frac{Ab - aB}{2b(a + bx)^2\sqrt{d + ex}(bd - ae)} \\
 & \quad \downarrow 61 \\
 & \frac{(aBe - 5Abe + 4bBd) \left(-\frac{3e \left(\frac{b \int \frac{1}{(a+bx)\sqrt{d+ex}} dx}{bd-ae} + \frac{2}{\sqrt{d+ex}(bd-ae)} \right)}{2(bd-ae)} - \frac{1}{(a+bx)\sqrt{d+ex}(bd-ae)} \right)}{4b(bd - ae)} - \frac{Ab - aB}{2b(a + bx)^2\sqrt{d + ex}(bd - ae)}
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow 73 \\
 (aBe - 5Abe + 4bBd) \left(-\frac{3e \left(\frac{2b \int \frac{1}{a + \frac{b(d+ex) - bd}{e(bd-ae)}} d\sqrt{d+ex} + \frac{2}{\sqrt{d+ex}(bd-ae)}} \right)}{2(bd-ae)} - \frac{1}{(a+bx)\sqrt{d+ex}(bd-ae)} \right) \\
 \hline
 \frac{4b(bd - ae)}{Ab - aB} \\
 \frac{2b(a + bx)^2 \sqrt{d + ex}(bd - ae)}{2b(a + bx)^2 \sqrt{d + ex}(bd - ae)} \\
 \downarrow 221 \\
 (aBe - 5Abe + 4bBd) \left(-\frac{3e \left(\frac{2}{\sqrt{d+ex}(bd-ae)} - \frac{2\sqrt{b}\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{(bd-ae)^{3/2}} \right)}{2(bd-ae)} - \frac{1}{(a+bx)\sqrt{d+ex}(bd-ae)} \right) \\
 \hline
 \frac{4b(bd - ae)}{Ab - aB} \\
 \frac{2b(a + bx)^2 \sqrt{d + ex}(bd - ae)}{2b(a + bx)^2 \sqrt{d + ex}(bd - ae)}
 \end{array}$$

input `Int[(A + B*x)/((a + b*x)^3*(d + e*x)^(3/2)),x]`

output `-1/2*(A*b - a*B)/(b*(b*d - a*e)*(a + b*x)^2*Sqrt[d + e*x]) + ((4*b*B*d - 5*A*b*e + a*B*e)*(-1/((b*d - a*e)*(a + b*x)*Sqrt[d + e*x])) - (3*e*(2/((b*d - a*e)*Sqrt[d + e*x]) - (2*Sqrt[b]*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[b*d - a*e]])/(b*d - a*e)^(3/2)))/(2*(b*d - a*e)))/(4*b*(b*d - a*e))`

Defintions of rubi rules used

rule 52 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 61 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.08

method	result
derivativedivides	$2e \left(-\frac{\left(\frac{7}{8}Ab^2e - \frac{3}{8}Babe - \frac{1}{2}b^2Bd\right)(ex+d)^{\frac{3}{2}} + \left(\frac{9}{8}Ab^2e^2 - \frac{9}{8}Ab^2de - \frac{5}{8}Ba^2e^2 + \frac{1}{8}Babde + \frac{1}{2}b^2Bd^2\right)\sqrt{ex+d}}{\left((ex+d)b+ae-db\right)^2} + \frac{3(5Abe - Bae - 4Bbd)}{8\sqrt{ae-db}} \right) \frac{1}{(ae-db)^3}$
default	$2e \left(-\frac{\left(\frac{7}{8}Ab^2e - \frac{3}{8}Babe - \frac{1}{2}b^2Bd\right)(ex+d)^{\frac{3}{2}} + \left(\frac{9}{8}Ab^2e^2 - \frac{9}{8}Ab^2de - \frac{5}{8}Ba^2e^2 + \frac{1}{8}Babde + \frac{1}{2}b^2Bd^2\right)\sqrt{ex+d}}{\left((ex+d)b+ae-db\right)^2} + \frac{3(5Abe - Bae - 4Bbd)}{8\sqrt{ae-db}} \right) \frac{1}{(ae-db)^3}$
pseudoelliptic	$2 \left(\frac{15(bx+a)^2 e^{\sqrt{ex+d}} \left((Ae - \frac{4Bd}{5})b - \frac{Bae}{5} \right) \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{8} + \sqrt{(ae-db)b} \left(\left(\frac{15Ae^2x^2}{8} + \frac{5xd(-\frac{12Bx}{5} + A)e}{8} - \frac{d^2(2Bd - Ae)}{4} \right) \sqrt{ex+d} \sqrt{(ae-db)b} (bx+a)^2 \right) \right) \frac{1}{(ae-db)^3}$

```
input int((B*x+A)/(b*x+a)^3/(e*x+d)^(3/2), x, method=_RETURNVERBOSE)
```

```
output 2*e*(-1/(a*e-b*d)^3*(((7/8*A*b^2*e-3/8*B*a*b*e-1/2*b^2*B*d)*(e*x+d)^(3/2)+(9/8*A*a*b*e^2-9/8*A*b^2*d*e-5/8*B*a^2*e^2+1/8*B*a*b*d*e+1/2*b^2*B*d^2)*(e*x+d)^(1/2))/((e*x+d)*b+a*e-d*b)^2+3/8*(5*A*b*e-B*a*e-4*B*b*d)/((a*e-b*d)*b)^(1/2)*arctan(b*(e*x+d)^(1/2)/((a*e-b*d)*b)^(1/2))-1/(a*e-b*d)^3*(A*e-B*d)/(e*x+d)^(1/2))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 698 vs. 2(158) = 316.
 Time = 0.17 (sec) , antiderivative size = 1410, normalized size of antiderivative = 7.83

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^{3/2}} dx = \text{Too large to display}$$

```
input integrate((B*x+A)/(b*x+a)^3/(e*x+d)^(3/2), x, algorithm="fricas")
```


output

```
[1/8*(3*(4*B*a^2*b*d^2*e + (B*a^3 - 5*A*a^2*b)*d*e^2 + (4*B*b^3*d^2*e + (B
*a*b^2 - 5*A*b^3)*e^3)*x^3 + (4*B*b^3*d^2*e + (9*B*a*b^2 - 5*A*b^3)*d*e^2
+ 2*(B*a^2*b - 5*A*a*b^2)*e^3)*x^2 + (8*B*a*b^2*d^2*e + 2*(3*B*a^2*b - 5*A
*a*b^2)*d*e^2 + (B*a^3 - 5*A*a^2*b)*e^3)*x)*sqrt(b^2*d - a*b*e)*log((b*e*x
+ 2*b*d - a*e + 2*sqrt(b^2*d - a*b*e)*sqrt(e*x + d))/(b*x + a)) - 2*(8*A*
a^3*b*e^3 + 2*(B*a*b^3 + A*b^4)*d^3 + 11*(B*a^2*b^2 - A*a*b^3)*d^2*e - (13
*B*a^3*b - A*a^2*b^2)*d*e^2 + 3*(4*B*b^4*d^2*e - (3*B*a*b^3 + 5*A*b^4)*d*e
^2 - (B*a^2*b^2 - 5*A*a*b^3)*e^3)*x^2 + (4*B*b^4*d^3 + (17*B*a*b^3 - 5*A*b
^4)*d^2*e - 4*(4*B*a^2*b^2 + 5*A*a*b^3)*d*e^2 - 5*(B*a^3*b - 5*A*a^2*b^2)*
e^3)*x)*sqrt(e*x + d)/(a^2*b^5*d^5 - 4*a^3*b^4*d^4*e + 6*a^4*b^3*d^3*e^2
- 4*a^5*b^2*d^2*e^3 + a^6*b*d*e^4 + (b^7*d^4*e - 4*a*b^6*d^3*e^2 + 6*a^2*b
^5*d^2*e^3 - 4*a^3*b^4*d*e^4 + a^4*b^3*e^5)*x^3 + (b^7*d^5 - 2*a*b^6*d^4*e
- 2*a^2*b^5*d^3*e^2 + 8*a^3*b^4*d^2*e^3 - 7*a^4*b^3*d*e^4 + 2*a^5*b^2*e^5
)*x^2 + (2*a*b^6*d^5 - 7*a^2*b^5*d^4*e + 8*a^3*b^4*d^3*e^2 - 2*a^4*b^3*d^2
*e^3 - 2*a^5*b^2*d*e^4 + a^6*b*e^5)*x), -1/4*(3*(4*B*a^2*b*d^2*e + (B*a^3
- 5*A*a^2*b)*d*e^2 + (4*B*b^3*d^2*e + (B*a*b^2 - 5*A*b^3)*e^3)*x^3 + (4*B*
b^3*d^2*e + (9*B*a*b^2 - 5*A*b^3)*d*e^2 + 2*(B*a^2*b - 5*A*a*b^2)*e^3)*x^2
+ (8*B*a*b^2*d^2*e + 2*(3*B*a^2*b - 5*A*a*b^2)*d*e^2 + (B*a^3 - 5*A*a^2*b
)*e^3)*x)*sqrt(-b^2*d + a*b*e)*arctan(sqrt(-b^2*d + a*b*e)*sqrt(e*x + d)/(
b*e*x + b*d)) + (8*A*a^3*b*e^3 + 2*(B*a*b^3 + A*b^4)*d^3 + 11*(B*a^2*b^...
```

Sympy [F(-1)]

Timed out.

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^{3/2}} dx = \text{Timed out}$$

input

```
integrate((B*x+A)/(b*x+a)**3/(e*x+d)**(3/2), x)
```

output

Timed out

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^{3/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^3/(e*x+d)^(3/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 340 vs. 2(158) = 316.

Time = 0.13 (sec) , antiderivative size = 340, normalized size of antiderivative = 1.89

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^{3/2}} dx = -\frac{3(4Bbde + Bae^2 - 5Abe^2) \arctan\left(\frac{\sqrt{ex+db}}{\sqrt{-b^2d+abe}}\right) + 2(Bde - Ae^2)}{(b^3d^3 - 3ab^2d^2e + 3a^2bde^2 - a^3e^3)\sqrt{ex+d}} - \frac{4(ex+d)^{\frac{3}{2}}Bb^2de - 4\sqrt{ex+d}Bb^2d^2e + 3(ex+d)^{\frac{3}{2}}Babe^2 - 7(ex+d)^{\frac{3}{2}}Ab^2e^2 - \sqrt{ex+d}Babde^2 + 9\sqrt{ex+d}Ae^2}{4(b^3d^3 - 3ab^2d^2e + 3a^2bde^2 - a^3e^3)((ex+d)b - bd + a^2e)}$$

input `integrate((B*x+A)/(b*x+a)^3/(e*x+d)^(3/2),x, algorithm="giac")`

output
$$-3/4*(4*B*b*d*e + B*a*e^2 - 5*A*b*e^2)*\arctan(\sqrt{e*x + d}*b/\sqrt{-b^2*d + a*b*e})/((b^3*d^3 - 3*a*b^2*d^2*e + 3*a^2*b*d*e^2 - a^3*e^3)*\sqrt{-b^2*d + a*b*e}) - 2*(B*d*e - A*e^2)/((b^3*d^3 - 3*a*b^2*d^2*e + 3*a^2*b*d*e^2 - a^3*e^3)*\sqrt{e*x + d}) - 1/4*(4*(e*x + d)^(3/2)*B*b^2*d*e - 4*\sqrt{e*x + d}*B*b^2*d^2*e + 3*(e*x + d)^(3/2)*B*a*b*e^2 - 7*(e*x + d)^(3/2)*A*b^2*e^2 - \sqrt{e*x + d}*B*a*b*d*e^2 + 9*\sqrt{e*x + d}*A*b^2*d*e^2 + 5*\sqrt{e*x + d}*B*a^2*e^3 - 9*\sqrt{e*x + d}*A*a*b*e^3)/((b^3*d^3 - 3*a*b^2*d^2*e + 3*a^2*b*d*e^2 - a^3*e^3)*((e*x + d)*b - b*d + a^2*e))$$

Mupad [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 296, normalized size of antiderivative = 1.64

$$\int \frac{A + Bx}{(a + bx)^3 (d + ex)^{3/2}} dx = \frac{\frac{5(d+ex)(Bae^2 - 5Abe^2 + 4Bbde)}{4(ae-bd)^2} - \frac{2(Ae^2 - Bde)}{ae-bd} + \frac{3b(d+ex)^2(Bae^2 - 5Abe^2 + 4Bbde)}{4(ae-bd)^3}}{b^2(d+ex)^{5/2} - (2b^2d - 2abe)(d+ex)^{3/2} + \sqrt{d+ex}(a^2e^2 - 2abde + b^2d^2)} + \frac{3e \operatorname{atan}\left(\frac{3\sqrt{b}\sqrt{d+ex}(Bae - 5Abe + 4Bbd)(a^3e^3 - 3a^2bde^2 + 3ab^2d^2e - b^3d^3)}{(ae-bd)^{7/2}(3Bae^2 - 15Abe^2 + 12Bbde)}\right)}{4\sqrt{b}(ae-bd)^{7/2}} (Bae - 5Abe + 4Bbd)$$

input `int((A + B*x)/((a + b*x)^3*(d + e*x)^(3/2)),x)`output `((5*(d + e*x)*(B*a*e^2 - 5*A*b*e^2 + 4*B*b*d*e))/(4*(a*e - b*d)^2) - (2*(A*e^2 - B*d*e))/(a*e - b*d) + (3*b*(d + e*x)^2*(B*a*e^2 - 5*A*b*e^2 + 4*B*b*d*e))/(4*(a*e - b*d)^3))/(b^2*(d + e*x)^(5/2) - (2*b^2*d - 2*a*b*e)*(d + e*x)^(3/2) + (d + e*x)^(1/2)*(a^2*e^2 + b^2*d^2 - 2*a*b*d*e)) + (3*e*atan((3*b^(1/2)*e*(d + e*x)^(1/2)*(B*a*e - 5*A*b*e + 4*B*b*d)*(a^3*e^3 - b^3*d^3 + 3*a*b^2*d^2*e - 3*a^2*b*d*e^2)))/((a*e - b*d)^(7/2)*(3*B*a*e^2 - 15*A*b*e^2 + 12*B*b*d*e)))*(B*a*e - 5*A*b*e + 4*B*b*d))/(4*b^(1/2)*(a*e - b*d)^(7/2))`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.22

$$\int \frac{A + Bx}{(a + bx)^3 (d + ex)^{3/2}} dx = \frac{-3\sqrt{b}\sqrt{ex+d}\sqrt{ae-bd}\operatorname{atan}\left(\frac{\sqrt{ex+d}b}{\sqrt{b}\sqrt{ae-bd}}\right)ae - 3\sqrt{b}\sqrt{ex+d}\sqrt{ae-bd}\operatorname{atan}\left(\frac{\sqrt{ex+d}b}{\sqrt{b}\sqrt{ae-bd}}\right)}{\sqrt{ex+d}(a^3be^3x - 3a^2b^2de^2x + 3ab^3d^2ex - b^4d^3x + a^4d^3)}$$

input `int((B*x+A)/(b*x+a)^3/(e*x+d)^(3/2),x)`output `(- 3*sqrt(b)*sqrt(d + e*x)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))*a*e - 3*sqrt(b)*sqrt(d + e*x)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))*b*e*x - 2*a**2*e**2 + a*b*d*e - 3*a*b*e**2*x + b**2*d**2 + 3*b**2*d*e*x)/(sqrt(d + e*x)*(a**4*e**3 - 3*a**3*b*d*e**2 + a**3*b*e**3*x + 3*a**2*b**2*d**2*e - 3*a**2*b**2*d*e**2*x - a*b**3*d**3 + 3*a*b**3*d**2*e*x - b**4*d**3*x))`

3.170 $\int \frac{A+Bx}{(a+bx)^3(d+ex)^{5/2}} dx$

Optimal result	1551
Mathematica [A] (verified)	1552
Rubi [A] (verified)	1552
Maple [A] (verified)	1555
Fricas [B] (verification not implemented)	1556
Sympy [F(-1)]	1557
Maxima [F(-2)]	1557
Giac [B] (verification not implemented)	1557
Mupad [B] (verification not implemented)	1558
Reduce [B] (verification not implemented)	1559

Optimal result

Integrand size = 22, antiderivative size = 222

$$\int \frac{A+Bx}{(a+bx)^3(d+ex)^{5/2}} dx = -\frac{2e(Bd-Ae)}{3(bd-ae)^3(d+ex)^{3/2}} - \frac{2e(2bBd-3Abe+aBe)}{(bd-ae)^4\sqrt{d+ex}}$$

$$- \frac{b(Ab-aB)\sqrt{d+ex}}{2(bd-ae)^3(a+bx)^2} - \frac{b(4bBd-11Abe+7aBe)\sqrt{d+ex}}{4(bd-ae)^4(a+bx)}$$

$$+ \frac{5\sqrt{be}(4bBd-7Abe+3aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{4(bd-ae)^{9/2}}$$

output

```
-2/3*e*(-A*e+B*d)/(-a*e+b*d)^3/(e*x+d)^(3/2)-2*e*(-3*A*b*e+B*a*e+2*B*b*d)/
(-a*e+b*d)^4/(e*x+d)^(1/2)-1/2*b*(A*b-B*a)*(e*x+d)^(1/2)/(-a*e+b*d)^3/(b*x
+a)^2-1/4*b*(-11*A*b*e+7*B*a*e+4*B*b*d)*(e*x+d)^(1/2)/(-a*e+b*d)^4/(b*x+a)
+5/4*b^(1/2)*e*(-7*A*b*e+3*B*a*e+4*B*b*d)*arctanh(b^(1/2)*(e*x+d)^(1/2)/(-
a*e+b*d)^(1/2))/(-a*e+b*d)^(9/2)
```

Mathematica [A] (verified)

Time = 0.99 (sec) , antiderivative size = 291, normalized size of antiderivative = 1.31

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^{5/2}} dx = \frac{-B(8a^3e^2(2d + 3ex) + 4b^3dx(3d^2 + 20dex + 15e^2x^2) + a^2be(83d^2 + 134dex) + 5\sqrt{be}(4bBd - 7Abe + 3aBe) \arctan\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right)}{4(-bd + ae)^{9/2}}$$

input

```
Integrate[(A + B*x)/((a + b*x)^3*(d + e*x)^(5/2)),x]
```

output

```
(-(B*(8*a^3*e^2*(2*d + 3*e*x) + 4*b^3*d*x*(3*d^2 + 20*d*e*x + 15*e^2*x^2) + a^2*b*e*(83*d^2 + 134*d*e*x + 75*e^2*x^2) + a*b^2*(6*d^3 + 145*d^2*e*x + 160*d*e^2*x^2 + 45*e^3*x^3))) + A*(-8*a^3*e^3 + 8*a^2*b*e^2*(10*d + 7*e*x) + a*b^2*e*(39*d^2 + 238*d*e*x + 175*e^2*x^2) + b^3*(-6*d^3 + 21*d^2*e*x + 140*d*e^2*x^2 + 105*e^3*x^3)))/(12*(b*d - a*e)^4*(a + b*x)^2*(d + e*x)^(3/2)) - (5*sqrt[b]*e*(4*b*B*d - 7*A*b*e + 3*a*B*e)*ArcTan[(sqrt[b]*sqrt[d + e*x])/sqrt[-(b*d) + a*e]])/(4*(-(b*d) + a*e)^(9/2))
```

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.01, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {87, 52, 61, 61, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^{5/2}} dx$$

$$\downarrow 87$$

$$\frac{(3aBe - 7Abe + 4bBd) \int \frac{1}{(a+bx)^2(d+ex)^{5/2}} dx}{4b(bd - ae)} - \frac{Ab - aB}{2b(a + bx)^2(d + ex)^{3/2}(bd - ae)}$$

$$\downarrow 52$$

$$\begin{aligned}
 & \frac{(3aBe - 7Abe + 4bBd) \left(-\frac{5e \int \frac{1}{(a+bx)(d+ex)^{5/2}} dx}{2(bd-ae)} - \frac{1}{(a+bx)(d+ex)^{3/2}(bd-ae)} \right)}{\frac{4b(bd-ae)}{Ab-aB}} \\
 & \frac{4b(bd-ae)}{2b(a+bx)^2(d+ex)^{3/2}(bd-ae)} \\
 & \quad \downarrow \text{61} \\
 & \frac{(3aBe - 7Abe + 4bBd) \left(-\frac{5e \left(\frac{b \int \frac{1}{(a+bx)(d+ex)^{3/2}} dx}{bd-ae} + \frac{2}{3(d+ex)^{3/2}(bd-ae)} \right)}{2(bd-ae)} - \frac{1}{(a+bx)(d+ex)^{3/2}(bd-ae)} \right)}{\frac{4b(bd-ae)}{Ab-aB}} \\
 & \frac{4b(bd-ae)}{2b(a+bx)^2(d+ex)^{3/2}(bd-ae)} \\
 & \quad \downarrow \text{61} \\
 & \frac{(3aBe - 7Abe + 4bBd) \left(-\frac{5e \left(\frac{b \left(\frac{b \int \frac{1}{(a+bx)\sqrt{d+ex}} dx}{bd-ae} + \frac{2}{\sqrt{d+ex}(bd-ae)} \right)}{bd-ae} + \frac{2}{3(d+ex)^{3/2}(bd-ae)} \right)}{2(bd-ae)} - \frac{1}{(a+bx)(d+ex)^{3/2}(bd-ae)} \right)}{\frac{4b(bd-ae)}{Ab-aB}} \\
 & \frac{4b(bd-ae)}{2b(a+bx)^2(d+ex)^{3/2}(bd-ae)} \\
 & \quad \downarrow \text{73} \\
 & \frac{(3aBe - 7Abe + 4bBd) \left(-\frac{5e \left(\frac{b \left(\frac{2b \int \frac{1}{a+b(d+ex)} - \frac{bd}{e(bd-ae)} d\sqrt{d+ex}}{bd-ae} + \frac{2}{\sqrt{d+ex}(bd-ae)} \right)}{bd-ae} + \frac{2}{3(d+ex)^{3/2}(bd-ae)} \right)}{2(bd-ae)} - \frac{1}{(a+bx)(d+ex)^{3/2}(bd-ae)} \right)}{\frac{4b(bd-ae)}{Ab-aB}} \\
 & \frac{4b(bd-ae)}{2b(a+bx)^2(d+ex)^{3/2}(bd-ae)}
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow 221 \\
 (3aBe - 7Abe + 4bBd) \left(\frac{5e \left(\frac{b \left(\frac{2}{\sqrt{d+ex}(bd-ae)} - \frac{2\sqrt{b}\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{(bd-ae)^{3/2}} \right)}{bd-ae} + \frac{2}{3(d+ex)^{3/2}(bd-ae)} \right)}{2(bd-ae)} - \frac{1}{(a+bx)(d+ex)^{3/2}(bd-ae)} \right) \\
 \hline
 \frac{4b(bd-ae)}{Ab - aB} \\
 \frac{2b(a+bx)^2(d+ex)^{3/2}(bd-ae)}{2b(a+bx)^2(d+ex)^{3/2}(bd-ae)}
 \end{array}$$

input `Int[(A + B*x)/((a + b*x)^3*(d + e*x)^(5/2)), x]`

output `-1/2*(A*b - a*B)/(b*(b*d - a*e)*(a + b*x)^2*(d + e*x)^(3/2)) + ((4*b*B*d - 7*A*b*e + 3*a*B*e)*(-1/((b*d - a*e)*(a + b*x)*(d + e*x)^(3/2))) - (5*e*(2/(3*(b*d - a*e)*(d + e*x)^(3/2)) + (b*(2/((b*d - a*e)*Sqrt[d + e*x]) - (2*Sqrt[b]*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[b*d - a*e]]))/(b*d - a*e)^(3/2)))/(b*d - a*e)))/(2*(b*d - a*e)))/(4*b*(b*d - a*e))`

Defintions of rubi rules used

rule 52 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 61 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 $\text{Int}[(a_.) + (b_.)(x_)^{(m_)}((c_.) + (d_.)(x_)^{(n_)}, x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \text{ Subst}[\text{Int}[x^{(p*(m+1)-1)}(c - a*(d/b) + d*(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 87 $\text{Int}[(a_.) + (b_.)(x_)*((c_.) + (d_.)(x_)^{(n_.)}((e_.) + (f_.)(x_)^{(p_.)}, x_] \rightarrow \text{Simp}[(-b*e - a*f)*(c + d*x)^{(n+1)}*(e + f*x)^{(p+1)}/(f*(p+1)*(c*f - d*e)), x] - \text{Simp}[(a*d*f*(n+p+2) - b*(d*e*(n+1) + c*f*(p+1))]/(f*(p+1)*(c*f - d*e)) \text{ Int}[(c + d*x)^n*(e + f*x)^{(p+1)}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n\}, x] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ (!\text{LtQ}[n, -1] \ || \ \text{IntegerQ}[p] \ || \ !(\text{IntegerQ}[n] \ || \ !(\text{EqQ}[e, 0] \ || \ !(\text{EqQ}[c, 0] \ || \ \text{LtQ}[p, n])))$

rule 221 $\text{Int}[(a_) + (b_.)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$

Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.03

method	result
derivativedivides	$2e \left(\frac{b \left(\frac{(\frac{11}{8}Ab^2e - \frac{7}{8}Babe - \frac{1}{2}b^2Bd)(ex+d)^{\frac{3}{2}} + (\frac{13}{8}Aab e^2 - \frac{13}{8}Ab^2de - \frac{9}{8}Ba^2e^2 + \frac{5}{8}Babde + \frac{1}{2}b^2Ba^2)\sqrt{ex+d}}{((ex+d)b+ae-db)^2} + \frac{5(7Abe-3Bae-3Bbd)}{2((ex+d)b+ae-db)} \right)}{(ae-db)^4} \right)$
default	$2e \left(\frac{b \left(\frac{(\frac{11}{8}Ab^2e - \frac{7}{8}Babe - \frac{1}{2}b^2Bd)(ex+d)^{\frac{3}{2}} + (\frac{13}{8}Aab e^2 - \frac{13}{8}Ab^2de - \frac{9}{8}Ba^2e^2 + \frac{5}{8}Babde + \frac{1}{2}b^2Ba^2)\sqrt{ex+d}}{((ex+d)b+ae-db)^2} + \frac{5(7Abe-3Bae-3Bbd)}{2((ex+d)b+ae-db)} \right)}{(ae-db)^4} \right)$
pseudoelliptic	$2 \left(-\frac{105((Ae - \frac{4Bd}{7})b - \frac{3Bae}{7})(ex+d)^{\frac{3}{2}}b(bx+a)^2 e \arctan\left(\frac{b\sqrt{ex+d}}{\sqrt{(ae-db)b}}\right)}{8} + \sqrt{(ae-db)b} \left(-\frac{105Ae^3x^3}{8} - \frac{35(-\frac{3Bx}{7}+A)x^2}{2} \right) \right)$

input $\text{int}((B*x+A)/(b*x+a)^3/(e*x+d)^{(5/2)}, x, \text{method}=_RETURNVERBOSE)$

output

```
2*e*(1/(a*e-b*d)^4*b*((11/8*A*b^2*e-7/8*B*a*b*e-1/2*b^2*B*d)*(e*x+d)^(3/2)
)+(13/8*A*a*b*e^2-13/8*A*b^2*d*e-9/8*B*a^2*e^2+5/8*B*a*b*d*e+1/2*b^2*B*d^2
)*(e*x+d)^(1/2))/((e*x+d)*b+a*e-d*b)^2+5/8*(7*A*b*e-3*B*a*e-4*B*b*d)/((a*e
-b*d)*b)^(1/2)*arctan(b*(e*x+d)^(1/2)/((a*e-b*d)*b)^(1/2))-1/3/(a*e-b*d)^
3*(A*e-B*d)/(e*x+d)^(3/2)-1/(a*e-b*d)^4*(-3*A*b*e+B*a*e+2*B*b*d)/(e*x+d)^(
1/2))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 863 vs. $2(196) = 392$.

Time = 0.19 (sec) , antiderivative size = 1756, normalized size of antiderivative = 7.91

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^{5/2}} dx = \text{Too large to display}$$

input

```
integrate((B*x+A)/(b*x+a)^3/(e*x+d)^(5/2),x, algorithm="fricas")
```

output

```
[-1/24*(15*(4*B*a^2*b*d^3*e + (3*B*a^3 - 7*A*a^2*b)*d^2*e^2 + (4*B*b^3*d*e
^3 + (3*B*a*b^2 - 7*A*b^3)*e^4)*x^4 + 2*(4*B*b^3*d^2*e^2 + 7*(B*a*b^2 - A
b^3)*d*e^3 + (3*B*a^2*b - 7*A*a*b^2)*e^4)*x^3 + (4*B*b^3*d^3*e + (19*B*a*b
^2 - 7*A*b^3)*d^2*e^2 + 4*(4*B*a^2*b - 7*A*a*b^2)*d*e^3 + (3*B*a^3 - 7*A*a
^2*b)*e^4)*x^2 + 2*(4*B*a*b^2*d^3*e + 7*(B*a^2*b - A*a*b^2)*d^2*e^2 + (3*B
*a^3 - 7*A*a^2*b)*d*e^3)*x)*sqrt(b/(b*d - a*e))*log((b*e*x + 2*b*d - a*e -
2*(b*d - a*e)*sqrt(e*x + d)*sqrt(b/(b*d - a*e)))/(b*x + a)) + 2*(8*A*a^3*
e^3 + 6*(B*a*b^2 + A*b^3)*d^3 + (83*B*a^2*b - 39*A*a*b^2)*d^2*e + 16*(B*a^
3 - 5*A*a^2*b)*d*e^2 + 15*(4*B*b^3*d*e^2 + (3*B*a*b^2 - 7*A*b^3)*e^3)*x^3
+ 5*(16*B*b^3*d^2*e + 4*(8*B*a*b^2 - 7*A*b^3)*d*e^2 + 5*(3*B*a^2*b - 7*A*a
*b^2)*e^3)*x^2 + (12*B*b^3*d^3 + (145*B*a*b^2 - 21*A*b^3)*d^2*e + 2*(67*B*
a^2*b - 119*A*a*b^2)*d*e^2 + 8*(3*B*a^3 - 7*A*a^2*b)*e^3)*x)*sqrt(e*x + d)
)/(a^2*b^4*d^6 - 4*a^3*b^3*d^5*e + 6*a^4*b^2*d^4*e^2 - 4*a^5*b*d^3*e^3 + a
^6*d^2*e^4 + (b^6*d^4*e^2 - 4*a*b^5*d^3*e^3 + 6*a^2*b^4*d^2*e^4 - 4*a^3*b^
3*d*e^5 + a^4*b^2*e^6)*x^4 + 2*(b^6*d^5*e - 3*a*b^5*d^4*e^2 + 2*a^2*b^4*d^
3*e^3 + 2*a^3*b^3*d^2*e^4 - 3*a^4*b^2*d*e^5 + a^5*b*e^6)*x^3 + (b^6*d^6 -
9*a^2*b^4*d^4*e^2 + 16*a^3*b^3*d^3*e^3 - 9*a^4*b^2*d^2*e^4 + a^6*e^6)*x^2
+ 2*(a*b^5*d^6 - 3*a^2*b^4*d^5*e + 2*a^3*b^3*d^4*e^2 + 2*a^4*b^2*d^3*e^3 -
3*a^5*b*d^2*e^4 + a^6*d*e^5)*x), -1/12*(15*(4*B*a^2*b*d^3*e + (3*B*a^3 -
7*A*a^2*b)*d^2*e^2 + (4*B*b^3*d*e^3 + (3*B*a*b^2 - 7*A*b^3)*e^4)*x^4 + ...
```

Sympy [F(-1)]

Timed out.

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^{5/2}} dx = \text{Timed out}$$

input `integrate((B*x+A)/(b*x+a)**3/(e*x+d)**(5/2),x)`

output `Timed out`

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^3/(e*x+d)^(5/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail`

Giac [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 446 vs. $2(196) = 392$.

Time = 0.13 (sec) , antiderivative size = 446, normalized size of antiderivative = 2.01

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^{5/2}} dx =$$

$$\frac{5(4Bb^2de + 3Babe^2 - 7Ab^2e^2) \arctan\left(\frac{\sqrt{ex+db}}{\sqrt{-b^2d+abe}}\right)}{4(b^4d^4 - 4ab^3d^3e + 6a^2b^2d^2e^2 - 4a^3bde^3 + a^4e^4)\sqrt{-b^2d+abe}}$$

$$\frac{2(6(ex+d)Bbde + Bbd^2e + 3(ex+d)Bae^2 - 9(ex+d)Abe^2 - Bade^2 - Abde^2 + Aae^3)}{3(b^4d^4 - 4ab^3d^3e + 6a^2b^2d^2e^2 - 4a^3bde^3 + a^4e^4)(ex+d)^{3/2}}$$

$$\frac{4(ex+d)^{3/2}Bb^3de - 4\sqrt{ex+d}Bb^3d^2e + 7(ex+d)^{3/2}Bab^2e^2 - 11(ex+d)^{3/2}Ab^3e^2 - 5\sqrt{ex+d}Bab^2de^2 + 4(b^4d^4 - 4ab^3d^3e + 6a^2b^2d^2e^2 - 4a^3bde^3 + a^4e^4)((ex+d)^{3/2} + \sqrt{ex+d})}{4(b^4d^4 - 4ab^3d^3e + 6a^2b^2d^2e^2 - 4a^3bde^3 + a^4e^4)((ex+d)^{3/2} + \sqrt{ex+d})}$$

input

```
integrate((B*x+A)/(b*x+a)^3/(e*x+d)^(5/2),x, algorithm="giac")
```

output

```
-5/4*(4*B*b^2*d*e + 3*B*a*b*e^2 - 7*A*b^2*e^2)*arctan(sqrt(e*x + d)*b/sqrt(-b^2*d + a*b*e))/((b^4*d^4 - 4*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2 - 4*a^3*b*d*e^3 + a^4*e^4)*sqrt(-b^2*d + a*b*e)) - 2/3*(6*(e*x + d)*B*b*d^2*e + B*b*d^2*e + 3*(e*x + d)*B*a*e^2 - 9*(e*x + d)*A*b*e^2 - B*a*d*e^2 - A*b*d*e^2 + A*a*e^3)/((b^4*d^4 - 4*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2 - 4*a^3*b*d*e^3 + a^4*e^4)*(e*x + d)^(3/2)) - 1/4*(4*(e*x + d)^(3/2)*B*b^3*d*e - 4*sqrt(e*x + d)*B*b^3*d^2*e + 7*(e*x + d)^(3/2)*B*a*b^2*e^2 - 11*(e*x + d)^(3/2)*A*b^3*e^2 - 5*sqrt(e*x + d)*B*a*b^2*d*e^2 + 13*sqrt(e*x + d)*A*b^3*d*e^2 + 9*sqrt(e*x + d)*B*a^2*b*e^3 - 13*sqrt(e*x + d)*A*a*b^2*e^3)/((b^4*d^4 - 4*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2 - 4*a^3*b*d*e^3 + a^4*e^4)*(e*x + d)*b - b*d + a*e)^2)
```

Mupad [B] (verification not implemented)

Time = 1.43 (sec) , antiderivative size = 361, normalized size of antiderivative = 1.63

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^{5/2}} dx =$$

$$\frac{\frac{2(Ae^2 - Bde)}{3(ae - bd)} + \frac{2(d+ex)(3Bae^2 - 7Abe^2 + 4Bbde)}{3(ae - bd)^2} + \frac{25(d+ex)^2(-7Ab^2e^2 + 4Bdb^2e + 3Babe^2)}{12(ae - bd)^3} + \frac{5b^2(d+ex)^3(3Bae^2 - 7Abe^2 + 4Bbde)}{4(ae - bd)^4}}{b^2(d+ex)^{7/2} - (2b^2d - 2abe)(d+ex)^{5/2} + (d+ex)^{3/2}(a^2e^2 - 2abde + b^2d^2)}$$

$$\frac{5\sqrt{b}e \operatorname{atan}\left(\frac{\sqrt{b}e\sqrt{d+ex}(3Bae - 7Abe + 4Bbd)(a^4e^4 - 4a^3bde^3 + 6a^2b^2d^2e^2 - 4ab^3d^3e + b^4d^4)}{(ae - bd)^{9/2}(3Bae^2 - 7Abe^2 + 4Bbde)}\right)}{4(ae - bd)^{9/2}}(3Bae - 7Abe + 4Bbd)}{4(ae - bd)^{9/2}}$$

input `int((A + B*x)/((a + b*x)^3*(d + e*x)^(5/2)),x)`

output
$$- \frac{((2*(A*e^2 - B*d*e))/(3*(a*e - b*d)) + (2*(d + e*x)*(3*B*a*e^2 - 7*A*b*e^2 + 4*B*b*d*e))/(3*(a*e - b*d)^2) + (25*(d + e*x)^2*(3*B*a*b*e^2 - 7*A*b^2*e^2 + 4*B*b^2*d*e))/(12*(a*e - b*d)^3) + (5*b^2*(d + e*x)^3*(3*B*a*e^2 - 7*A*b*e^2 + 4*B*b*d*e))/(4*(a*e - b*d)^4)}{(b^2*(d + e*x)^{7/2} - (2*b^2*d - 2*a*b*e)*(d + e*x)^{5/2} + (d + e*x)^{3/2}*(a^2*e^2 + b^2*d^2 - 2*a*b*d*e)) - (5*b^{1/2}*e*atan((b^{1/2}*e*(d + e*x)^{1/2}*(3*B*a*e - 7*A*b*e + 4*B*b*d)*(a^4*e^4 + b^4*d^4 + 6*a^2*b^2*d^2*e^2 - 4*a*b^3*d^3*e - 4*a^3*b*d*e^3)))/((a*e - b*d)^{9/2}*(3*B*a*e^2 - 7*A*b*e^2 + 4*B*b*d*e)))*(3*B*a*e - 7*A*b*e + 4*B*b*d)/(4*(a*e - b*d)^{9/2})}$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 493, normalized size of antiderivative = 2.22

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^{5/2}} dx = \frac{15\sqrt{b}\sqrt{ex + d}\sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b}\sqrt{ae-bd}}\right) abde + 15\sqrt{b}\sqrt{ex + d}\sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b}\sqrt{ae-bd}}\right) abde + 15\sqrt{b}\sqrt{ex + d}\sqrt{ae - bd} \operatorname{atan}\left(\frac{\sqrt{ex+db}}{\sqrt{b}\sqrt{ae-bd}}\right) abde}{3\sqrt{ex + d} (a^4 b e^5 x^2 - 4 a^3 b^2 e^4 x + 3 a^2 b^3 d e^3 - 2 a b^4 d^2 e^2 + b^5 d^3)}$$

input `int((B*x+A)/(b*x+a)^3/(e*x+d)^(5/2),x)`

output
$$\frac{(15*\sqrt{b}*\sqrt{d + e*x}*\sqrt{a*e - b*d}*atan((\sqrt{d + e*x}*b)/(\sqrt{b}*\sqrt{a*e - b*d}))*a*b*d*e + 15*\sqrt{b}*\sqrt{d + e*x}*\sqrt{a*e - b*d}*atan((\sqrt{d + e*x}*b)/(\sqrt{b}*\sqrt{a*e - b*d}))*a*b*e**2*x + 15*\sqrt{b}*\sqrt{d + e*x}*\sqrt{a*e - b*d}*atan((\sqrt{d + e*x}*b)/(\sqrt{b}*\sqrt{a*e - b*d}))*b**2*d*e*x + 15*\sqrt{b}*\sqrt{d + e*x}*\sqrt{a*e - b*d}*atan((\sqrt{d + e*x}*b)/(\sqrt{b}*\sqrt{a*e - b*d}))*b**2*e**2*x**2 - 2*a**3*e**3 + 16*a**2*b*d*e**2 + 10*a**2*b*e**3*x - 11*a*b**2*d**2*e + 10*a*b**2*d*e**2*x + 15*a*b**2*e**3*x**2 - 3*b**3*d**3 - 20*b**3*d**2*e*x - 15*b**3*d*e**2*x**2)/(3*\sqrt{d + e*x}*(a**5*d*e**4 + a**5*e**5*x - 4*a**4*b*d**2*e**3 - 3*a**4*b*d*e**4*x + a**4*b*e**5*x**2 + 6*a**3*b**2*d**3*e**2 + 2*a**3*b**2*d**2*e**3*x - 4*a**3*b**2*d*e**4*x**2 - 4*a**2*b**3*d**4*e + 2*a**2*b**3*d**3*e**2*x + 6*a**2*b**3*d**2*e**3*x**2 + a*b**4*d**5 - 3*a*b**4*d**4*e*x - 4*a*b**4*d**3*e**2*x**2 + b**5*d**5*x + b**5*d**4*e*x**2))}$$

3.171 $\int \frac{A+Bx}{(a+bx)^3(d+ex)^{7/2}} dx$

Optimal result	1560
Mathematica [A] (verified)	1561
Rubi [A] (verified)	1561
Maple [A] (verified)	1565
Fricas [B] (verification not implemented)	1566
Sympy [F(-1)]	1567
Maxima [F(-2)]	1568
Giac [B] (verification not implemented)	1568
Mupad [B] (verification not implemented)	1569
Reduce [B] (verification not implemented)	1570

Optimal result

Integrand size = 22, antiderivative size = 265

$$\int \frac{A+Bx}{(a+bx)^3(d+ex)^{7/2}} dx = -\frac{2e(Bd-Ae)}{5(bd-ae)^3(d+ex)^{5/2}} - \frac{2e(2bBd-3Abe+aBe)}{3(bd-ae)^4(d+ex)^{3/2}} - \frac{6be(bBd-2Abe+aBe)}{(bd-ae)^5\sqrt{d+ex}} - \frac{b^2(Ab-aB)\sqrt{d+ex}}{2(bd-ae)^4(a+bx)^2} - \frac{b^2(4bBd-15Abe+11aBe)\sqrt{d+ex}}{4(bd-ae)^5(a+bx)} + \frac{7b^{3/2}e(4bBd-9Abe+5aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{4(bd-ae)^{11/2}}$$

output

```
-2/5*e*(-A*e+B*d)/(-a*e+b*d)^3/(e*x+d)^(5/2)-2/3*e*(-3*A*b*e+B*a*e+2*B*b*d)/(-a*e+b*d)^4/(e*x+d)^(3/2)-6*b*e*(-2*A*b*e+B*a*e+B*b*d)/(-a*e+b*d)^5/(e*x+d)^(1/2)-1/2*b^2*(A*b-B*a)*(e*x+d)^(1/2)/(-a*e+b*d)^4/(b*x+a)^2-1/4*b^2*(-15*A*b*e+11*B*a*e+4*B*b*d)*(e*x+d)^(1/2)/(-a*e+b*d)^5/(b*x+a)+7/4*b^(3/2)*e*(-9*A*b*e+5*B*a*e+4*B*b*d)*arctanh(b^(1/2)*(e*x+d)^(1/2)/(-a*e+b*d)^(1/2))/(-a*e+b*d)^(11/2)
```

Mathematica [A] (verified)

Time = 1.04 (sec) , antiderivative size = 408, normalized size of antiderivative = 1.54

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^{7/2}} dx = \frac{3A(8a^4e^4 - 8a^3be^3(7d + 3ex) + 24a^2b^2e^2(12d^2 + 17dex + 7e^2x^2) + ab^3e(85d^3 + 831d^2e*x + 1239d*e^2*x^2 + 525e^3*x^3) + b^4*(-10*d^4 + 45*d^3*e*x + 483*d^2*e^2*x^2 + 735*d*e^3*x^3 + 315*e^4*x^4)) + B*(8*a^4*e^3*(2*d + 5*e*x) - 8*a^3*b*e^2*(34*d^2 + 81*d*e*x + 35*e^2*x^2) - 4*b^4*d*x*(15*d^3 + 161*d^2*e*x + 245*d*e^2*x^2 + 105*e^3*x^3) - a^2*b^2*e*(659*d^3 + 1929*d^2*e*x + 2289*d*e^2*x^2 + 875*e^3*x^3) - a*b^3*(30*d^4 + 1183*d^3*e*x + 2457*d^2*e^2*x^2 + 1925*d*e^3*x^3 + 525*e^4*x^4))}{60*(b*d - a*e)^5*(a + b*x)^2*(d + e*x)^{5/2}} + \frac{7b^{3/2}e(4bBd - 9Abe + 5aBe) \arctan\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{-bd+ae}}\right)}{4(-bd + ae)^{11/2}}$$

input

```
Integrate[(A + B*x)/((a + b*x)^3*(d + e*x)^(7/2)),x]
```

output

```
(3*A*(8*a^4*e^4 - 8*a^3*b*e^3*(7*d + 3*e*x) + 24*a^2*b^2*e^2*(12*d^2 + 17*d*e*x + 7*e^2*x^2) + a*b^3*e*(85*d^3 + 831*d^2*e*x + 1239*d*e^2*x^2 + 525*e^3*x^3) + b^4*(-10*d^4 + 45*d^3*e*x + 483*d^2*e^2*x^2 + 735*d*e^3*x^3 + 315*e^4*x^4)) + B*(8*a^4*e^3*(2*d + 5*e*x) - 8*a^3*b*e^2*(34*d^2 + 81*d*e*x + 35*e^2*x^2) - 4*b^4*d*x*(15*d^3 + 161*d^2*e*x + 245*d*e^2*x^2 + 105*e^3*x^3) - a^2*b^2*e*(659*d^3 + 1929*d^2*e*x + 2289*d*e^2*x^2 + 875*e^3*x^3) - a*b^3*(30*d^4 + 1183*d^3*e*x + 2457*d^2*e^2*x^2 + 1925*d*e^3*x^3 + 525*e^4*x^4)))/(60*(b*d - a*e)^5*(a + b*x)^2*(d + e*x)^(5/2)) + (7*b^(3/2)*e*(4*b*B*d - 9*A*b*e + 5*a*B*e)*ArcTan[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[-(b*d) + a*e]])/(4*(-(b*d) + a*e)^(11/2))
```

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 260, normalized size of antiderivative = 0.98, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.318$, Rules used = {87, 52, 61, 61, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^{7/2}} dx$$

↓ 87

$$\frac{(5aBe - 9Abe + 4bBd) \int \frac{1}{(a+bx)^2(d+ex)^{7/2}} dx}{4b(bd - ae)} - \frac{Ab - aB}{2b(a + bx)^2(d + ex)^{5/2}(bd - ae)}$$

$$\begin{array}{c}
 \downarrow 52 \\
 \frac{(5aBe - 9Abe + 4bBd) \left(-\frac{7e \int \frac{1}{(a+bx)(d+ex)^{7/2}} dx}{2(bd-ae)} - \frac{1}{(a+bx)(d+ex)^{5/2}(bd-ae)} \right)}{\frac{4b(bd-ae)}{Ab-aB} \frac{1}{2b(a+bx)^2(d+ex)^{5/2}(bd-ae)}} \\
 \downarrow 61 \\
 \frac{(5aBe - 9Abe + 4bBd) \left(-\frac{7e \left(\frac{b \int \frac{1}{(a+bx)(d+ex)^{5/2}} dx}{bd-ae} + \frac{2}{5(d+ex)^{5/2}(bd-ae)} \right)}{2(bd-ae)} - \frac{1}{(a+bx)(d+ex)^{5/2}(bd-ae)} \right)}{\frac{4b(bd-ae)}{Ab-aB} \frac{1}{2b(a+bx)^2(d+ex)^{5/2}(bd-ae)}} \\
 \downarrow 61 \\
 \frac{(5aBe - 9Abe + 4bBd) \left(-\frac{7e \left(b \left(\frac{\int \frac{1}{(a+bx)(d+ex)^{3/2}} dx}{bd-ae} + \frac{2}{3(d+ex)^{3/2}(bd-ae)} \right) + \frac{2}{5(d+ex)^{5/2}(bd-ae)} \right)}{2(bd-ae)} - \frac{1}{(a+bx)(d+ex)^{5/2}(bd-ae)} \right)}{\frac{4b(bd-ae)}{Ab-aB} \frac{1}{2b(a+bx)^2(d+ex)^{5/2}(bd-ae)}} \\
 \downarrow 61
 \end{array}$$

$$(5aBe - 9Abe + 4bBd) \left[\frac{7e \left(\frac{b \int \frac{1}{(a+bx)\sqrt{d+ex}} dx}{bd-ae} + \frac{2}{\sqrt{d+ex}(bd-ae)} \right)}{bd-ae} + \frac{2}{3(d+ex)^{3/2}(bd-ae)} \right] - \frac{2}{5(d+ex)^{5/2}(bd-ae)} - \frac{1}{(a+bx)(d+ex)}$$

$$\frac{4b(bd - ae)}{2b(a + bx)^2(d + ex)^{5/2}(bd - ae)} \frac{Ab - aB}{2b(a + bx)^2(d + ex)^{5/2}(bd - ae)}$$

↓ 73

$$(5aBe - 9Abe + 4bBd) \left[\frac{7e \left(\frac{b \int \frac{1}{a + \frac{b(d+ex)}{e} - \frac{bd}{e}} d\sqrt{d+ex}}{bd-ae} + \frac{2}{\sqrt{d+ex}(bd-ae)} \right)}{bd-ae} + \frac{2}{3(d+ex)^{3/2}(bd-ae)} \right] - \frac{2}{5(d+ex)^{5/2}(bd-ae)} - \frac{1}{(a+bx)(d+ex)}$$

$$\frac{4b(bd - ae)}{2b(a + bx)^2(d + ex)^{5/2}(bd - ae)} \frac{Ab - aB}{2b(a + bx)^2(d + ex)^{5/2}(bd - ae)}$$

↓ 221

$$\frac{(5aBe - 9Abe + 4bBd) \left(\frac{7e \left(\frac{b \left(\frac{2}{\sqrt{d+ex}(bd-ae)} - \frac{2\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{bd-ae}}\right)}{(bd-ae)^{3/2}} \right)}{bd-ae} + \frac{2}{3(d+ex)^{3/2}(bd-ae)} \right)}{bd-ae} + \frac{2}{5(d+ex)^{5/2}(bd-ae)} \right)}{2(bd-ae)} - \frac{Ab - aB}{2b(a+bx)^2(d+ex)^{5/2}(bd-ae)} \right)}{4b(bd-ae)}$$

```
input Int[(A + B*x)/((a + b*x)^3*(d + e*x)^(7/2)), x]
```

```
output -1/2*(A*b - a*B)/(b*(b*d - a*e)*(a + b*x)^2*(d + e*x)^(5/2)) + ((4*b*B*d - 9*A*b*e + 5*a*B*e)*(-1/((b*d - a*e)*(a + b*x)*(d + e*x)^(5/2))) - (7*e*(2/(5*(b*d - a*e)*(d + e*x)^(5/2)) + (b*(2/(3*(b*d - a*e)*(d + e*x)^(3/2)) + (b*(2/((b*d - a*e)*Sqrt[d + e*x]) - (2*Sqrt[b]*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/Sqrt[b*d - a*e]])/(b*d - a*e)^(3/2)))/(b*d - a*e)))/(2*(b*d - a*e))))/(4*b*(b*d - a*e))
```

Defintions of rubi rules used

```
rule 52 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]
```

rule 61 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [A] (verified)

Time = 0.58 (sec) , antiderivative size = 268, normalized size of antiderivative = 1.01

method	result
derivativedivides	$2e \left(\frac{b^2 \left(\frac{15}{8} A b^2 e - \frac{11}{8} B a b e - \frac{1}{2} b^2 B d \right) (e x + d)^{\frac{3}{2}} + \left(\frac{17}{8} A a b e^2 - \frac{17}{8} A b^2 d e - \frac{13}{8} B a^2 e^2 + \frac{9}{8} B a b d e + \frac{1}{2} b^2 B d^2 \right) \sqrt{e x + d}}{\left((e x + d) b + a e - d b \right)^2} + \frac{7(9 A b e - \dots)}{(a e - d b)^5} \right)$
default	$2e \left(\frac{b^2 \left(\frac{15}{8} A b^2 e - \frac{11}{8} B a b e - \frac{1}{2} b^2 B d \right) (e x + d)^{\frac{3}{2}} + \left(\frac{17}{8} A a b e^2 - \frac{17}{8} A b^2 d e - \frac{13}{8} B a^2 e^2 + \frac{9}{8} B a b d e + \frac{1}{2} b^2 B d^2 \right) \sqrt{e x + d}}{\left((e x + d) b + a e - d b \right)^2} + \frac{7(9 A b e - \dots)}{(a e - d b)^5} \right)$
pseudoelliptic	$2 \left(\frac{315(e x + d)^{\frac{5}{2}} b^2 (b x + a)^2 \left(A e - \frac{4 B d}{9} \right) b - \frac{5 B a e}{9}}{8} e \arctan \left(\frac{b \sqrt{e x + d}}{\sqrt{(a e - d b) b}} \right) + \sqrt{(a e - d b) b} \left(\left(\frac{315 A e^4 x^4}{8} + \frac{735 \left(-\frac{4 B x}{21} + A \right) x^3 d}{8} \right) \right) \right)$

```
input int((B*x+A)/(b*x+a)^3/(e*x+d)^(7/2),x,method=_RETURNVERBOSE)
```

```
output 2*e*(-1/(a*e-b*d))^5*b^2*(((15/8*A*b^2*e-11/8*B*a*b*e-1/2*b^2*B*d)*(e*x+d)^(3/2)+(17/8*A*a*b*e^2-17/8*A*b^2*d*e-13/8*B*a^2*e^2+9/8*B*a*b*d*e+1/2*b^2*B*d^2)*(e*x+d)^(1/2))/((e*x+d)*b+a*e-d*b)^2+7/8*(9*A*b*e-5*B*a*e-4*B*b*d)/((a*e-b*d)*b)^(1/2)*arctan(b*(e*x+d)^(1/2)/((a*e-b*d)*b)^(1/2))-1/5*(A*e-B*d)/(a*e-b*d)^3/(e*x+d)^(5/2)-1/3*(-3*A*b*e+B*a*e+2*B*b*d)/(a*e-b*d)^4/(e*x+d)^(3/2)-3*b*(2*A*b*e-B*a*e-B*b*d)/(a*e-b*d)^5/(e*x+d)^(1/2))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1314 vs. 2(235) = 470.

Time = 0.43 (sec) , antiderivative size = 2657, normalized size of antiderivative = 10.03

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^{7/2}} dx = \text{Too large to display}$$

```
input integrate((B*x+A)/(b*x+a)^3/(e*x+d)^(7/2),x, algorithm="fricas")
```

output

```
[1/120*(105*(4*B*a^2*b^2*d^4*e + (5*B*a^3*b - 9*A*a^2*b^2)*d^3*e^2 + (4*B*
b^4*d*e^4 + (5*B*a*b^3 - 9*A*b^4)*e^5)*x^5 + (12*B*b^4*d^2*e^3 + (23*B*a*b
^3 - 27*A*b^4)*d*e^4 + 2*(5*B*a^2*b^2 - 9*A*a*b^3)*e^5)*x^4 + (12*B*b^4*d^
3*e^2 + 3*(13*B*a*b^3 - 9*A*b^4)*d^2*e^3 + 2*(17*B*a^2*b^2 - 27*A*a*b^3)*d
*e^4 + (5*B*a^3*b - 9*A*a^2*b^2)*e^5)*x^3 + (4*B*b^4*d^4*e + (29*B*a*b^3 -
9*A*b^4)*d^3*e^2 + 6*(7*B*a^2*b^2 - 9*A*a*b^3)*d^2*e^3 + 3*(5*B*a^3*b - 9
*A*a^2*b^2)*d*e^4)*x^2 + (8*B*a*b^3*d^4*e + 2*(11*B*a^2*b^2 - 9*A*a*b^3)*d
^3*e^2 + 3*(5*B*a^3*b - 9*A*a^2*b^2)*d^2*e^3)*x)*sqrt(b/(b*d - a*e))*log((
b*e*x + 2*b*d - a*e + 2*(b*d - a*e)*sqrt(e*x + d)*sqrt(b/(b*d - a*e)))/(b*
x + a)) + 2*(24*A*a^4*e^4 - 30*(B*a*b^3 + A*b^4)*d^4 - (659*B*a^2*b^2 - 25
5*A*a*b^3)*d^3*e - 16*(17*B*a^3*b - 54*A*a^2*b^2)*d^2*e^2 + 8*(2*B*a^4 - 2
1*A*a^3*b)*d*e^3 - 105*(4*B*b^4*d*e^3 + (5*B*a*b^3 - 9*A*b^4)*e^4)*x^4 - 3
5*(28*B*b^4*d^2*e^2 + (55*B*a*b^3 - 63*A*b^4)*d*e^3 + 5*(5*B*a^2*b^2 - 9*A
*a*b^3)*e^4)*x^3 - 7*(92*B*b^4*d^3*e + 9*(39*B*a*b^3 - 23*A*b^4)*d^2*e^2 +
3*(109*B*a^2*b^2 - 177*A*a*b^3)*d*e^3 + 8*(5*B*a^3*b - 9*A*a^2*b^2)*e^4)*
x^2 - (60*B*b^4*d^4 + (1183*B*a*b^3 - 135*A*b^4)*d^3*e + 3*(643*B*a^2*b^2
- 831*A*a*b^3)*d^2*e^2 + 72*(9*B*a^3*b - 17*A*a^2*b^2)*d*e^3 - 8*(5*B*a^4
- 9*A*a^3*b)*e^4)*x)*sqrt(e*x + d))/(a^2*b^5*d^8 - 5*a^3*b^4*d^7*e + 10*a^
4*b^3*d^6*e^2 - 10*a^5*b^2*d^5*e^3 + 5*a^6*b*d^4*e^4 - a^7*d^3*e^5 + (b^7*
d^5*e^3 - 5*a*b^6*d^4*e^4 + 10*a^2*b^5*d^3*e^5 - 10*a^3*b^4*d^2*e^6 + 5...
```

Sympy [F(-1)]

Timed out.

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^{7/2}} dx = \text{Timed out}$$

input

```
integrate((B*x+A)/(b*x+a)**3/(e*x+d)**(7/2),x)
```

output

Timed out

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^{7/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^3/(e*x+d)^(7/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 609 vs. 2(235) = 470.

Time = 0.14 (sec) , antiderivative size = 609, normalized size of antiderivative = 2.30

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^{7/2}} dx =$$

$$\frac{7(4Bb^3de + 5Bab^2e^2 - 9Ab^3e^2) \arctan\left(\frac{\sqrt{ex+db}}{\sqrt{-b^2d+abe}}\right)}{4(b^5d^5 - 5ab^4d^4e + 10a^2b^3d^3e^2 - 10a^3b^2d^2e^3 + 5a^4bde^4 - a^5e^5)\sqrt{-b^2d+abe}}$$

$$\frac{4(ex+d)^{\frac{3}{2}}Bb^4de - 4\sqrt{ex+d}Bb^4d^2e + 11(ex+d)^{\frac{3}{2}}Bab^3e^2 - 15(ex+d)^{\frac{3}{2}}Ab^4e^2 - 9\sqrt{ex+d}Bab^3de^2}{4(b^5d^5 - 5ab^4d^4e + 10a^2b^3d^3e^2 - 10a^3b^2d^2e^3 + 5a^4bde^4 - a^5e^5)}$$

$$\frac{2(45(ex+d)^2Bb^2de + 10(ex+d)Bb^2d^2e + 3Bb^2d^3e + 45(ex+d)^2Babe^2 - 90(ex+d)^2Ab^2e^2 - 5(ex+d)Bab^2e^2}{15(b^5d^5 - 5ab^4d^4e + 10a^2b^3d^3e^2 - 10a^3b^2d^2e^3 + 5a^4bde^4 - a^5e^5)}$$

input `integrate((B*x+A)/(b*x+a)^3/(e*x+d)^(7/2),x, algorithm="giac")`

output

```
-7/4*(4*B*b^3*d*e + 5*B*a*b^2*e^2 - 9*A*b^3*e^2)*arctan(sqrt(e*x + d)*b/sqrt(-b^2*d + a*b*e))/((b^5*d^5 - 5*a*b^4*d^4*e + 10*a^2*b^3*d^3*e^2 - 10*a^3*b^2*d^2*e^3 + 5*a^4*b*d*e^4 - a^5*e^5)*sqrt(-b^2*d + a*b*e)) - 1/4*(4*(e*x + d)^(3/2)*B*b^4*d*e - 4*sqrt(e*x + d)*B*b^4*d^2*e + 11*(e*x + d)^(3/2)*B*a*b^3*e^2 - 15*(e*x + d)^(3/2)*A*b^4*e^2 - 9*sqrt(e*x + d)*B*a*b^3*d*e^2 + 17*sqrt(e*x + d)*A*b^4*d*e^2 + 13*sqrt(e*x + d)*B*a^2*b^2*e^3 - 17*sqrt(e*x + d)*A*a*b^3*e^3)/((b^5*d^5 - 5*a*b^4*d^4*e + 10*a^2*b^3*d^3*e^2 - 10*a^3*b^2*d^2*e^3 + 5*a^4*b*d*e^4 - a^5*e^5)*((e*x + d)*b - b*d + a*e)^2) - 2/15*(45*(e*x + d)^2*B*b^2*d*e + 10*(e*x + d)*B*b^2*d^2*e + 3*B*b^2*d^3*e + 45*(e*x + d)^2*B*a*b*e^2 - 90*(e*x + d)^2*A*b^2*e^2 - 5*(e*x + d)*B*a*b*d*e^2 - 15*(e*x + d)*A*b^2*d*e^2 - 6*B*a*b*d^2*e^2 - 3*A*b^2*d^2*e^2 - 5*(e*x + d)*B*a^2*e^3 + 15*(e*x + d)*A*a*b*e^3 + 3*B*a^2*d*e^3 + 6*A*a*b*d*e^3 - 3*A*a^2*e^4)/((b^5*d^5 - 5*a*b^4*d^4*e + 10*a^2*b^3*d^3*e^2 - 10*a^3*b^2*d^2*e^3 + 5*a^4*b*d*e^4 - a^5*e^5)*(e*x + d)^(5/2))
```

Mupad [B] (verification not implemented)

Time = 1.32 (sec) , antiderivative size = 418, normalized size of antiderivative = 1.58

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^{7/2}} dx = \frac{35(d+ex)^3(-9Ab^3e^2+4Bdb^3e+5Bab^2e^2)}{12(ae-bd)^4} - \frac{2(Ae^2-Bde)}{5(ae-bd)} - \frac{2(d+ex)(5Bae^2-9Abe^2+4Bbd)}{15(ae-bd)^2} + \frac{b^2(d+ex)^{9/2} - (2b^2d - 2abe)(d+ex)^{7/2}}{4(ae-bd)^{11/2}} + \frac{7b^{3/2}e \operatorname{atan}\left(\frac{\sqrt{be}\sqrt{d+ex}(5Bae-9Abe+4Bbd)(a^5e^5-5a^4bde^4+10a^3b^2d^2e^3-10a^2b^3d^3e^2+5ab^4d^4e-b^5d^5)}{(ae-bd)^{11/2}(5Bae^2-9Abe^2+4Bbde)}\right)}{4(ae-bd)^{11/2}}(5Bae-9Abe+4Bbd)$$

input

```
int((A + B*x)/((a + b*x)^3*(d + e*x)^(7/2)),x)
```

output

```
((35*(d + e*x)^3*(4*B*b^3*d*e - 9*A*b^3*e^2 + 5*B*a*b^2*e^2))/(12*(a*e - b*d)^4) - (2*(A*e^2 - B*d*e))/(5*(a*e - b*d)) - (2*(d + e*x)*(5*B*a*e^2 - 9*A*b*e^2 + 4*B*b*d*e))/(15*(a*e - b*d)^2) + (7*b^3*(d + e*x)^4*(5*B*a*e^2 - 9*A*b*e^2 + 4*B*b*d*e))/(4*(a*e - b*d)^5) + (14*b*(d + e*x)^2*(5*B*a*e^2 - 9*A*b*e^2 + 4*B*b*d*e))/(15*(a*e - b*d)^3))/(b^2*(d + e*x)^(9/2) - (2*b^2*d - 2*a*b*e)*(d + e*x)^(7/2) + (d + e*x)^(5/2)*(a^2*e^2 + b^2*d^2 - 2*a*b*d*e)) + (7*b^(3/2)*e*atan((b^(1/2)*e*(d + e*x)^(1/2)*(5*B*a*e - 9*A*b*e + 4*B*b*d)*(a^5*e^5 - b^5*d^5 - 10*a^2*b^3*d^3*e^2 + 10*a^3*b^2*d^2*e^3 + 5*a*b^4*d^4*e - 5*a^4*b*d*e^4))/((a*e - b*d)^(11/2)*(5*B*a*e^2 - 9*A*b*e^2 + 4*B*b*d*e)))*(5*B*a*e - 9*A*b*e + 4*B*b*d))/(4*(a*e - b*d)^(11/2))
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 804, normalized size of antiderivative = 3.03

$$\int \frac{A + Bx}{(a + bx)^3(d + ex)^{7/2}} dx = \frac{-105\sqrt{b}\sqrt{ex+d}\sqrt{ae-bd}\operatorname{atan}\left(\frac{\sqrt{ex+d}b}{\sqrt{b}\sqrt{ae-bd}}\right)ab^2d^2e - 210\sqrt{b}\sqrt{ex+d}\sqrt{ae-bd}}{(a+bx)^3(d+ex)^{7/2}}$$

input `int((B*x+A)/(b*x+a)^3/(e*x+d)^(7/2),x)`

output

```
( - 105*sqrt(b)*sqrt(d + e*x)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))*a*b**2*d**2*e - 210*sqrt(b)*sqrt(d + e*x)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))*a*b**2*d*e**2*x - 105*sqrt(b)*sqrt(d + e*x)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))*a*b**2*e**3*x**2 - 105*sqrt(b)*sqrt(d + e*x)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))*b**3*d**2*e*x - 210*sqrt(b)*sqrt(d + e*x)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))*b**3*d*e**2*x**2 - 105*sqrt(b)*sqrt(d + e*x)*sqrt(a*e - b*d)*atan((sqrt(d + e*x)*b)/(sqrt(b)*sqrt(a*e - b*d)))*b**3*e**3*x**3 - 6*a**4*e**4 + 38*a**3*b*d*e**3 + 14*a**3*b*e**4*x - 148*a**2*b**2*d**2*e**2 - 182*a**2*b**2*d*e**3*x - 70*a**2*b**2*e**4*x**2 + 101*a*b**3*d**3*e + 7*a*b**3*d**2*e**2*x - 175*a*b**3*d*e**3*x**2 - 105*a*b**3*e**4*x**3 + 15*b**4*d**4 + 161*b**4*d**3*e*x + 245*b**4*d**2*e**2*x**2 + 105*b**4*d*e**3*x**3)/(15*sqrt(d + e*x)*(a**6*d**2*e**5 + 2*a**6*d*e**6*x + a**6*e**7*x**2 - 5*a**5*b*d**3*e**4 - 9*a**5*b*d**2*e**5*x - 3*a**5*b*d*e**6*x**2 + a**5*b*e**7*x**3 + 10*a**4*b**2*d**4*e**3 + 15*a**4*b**2*d**3*e**4*x - 5*a**4*b**2*d*e**6*x**3 - 10*a**3*b**3*d**5*e**2 - 10*a**3*b**3*d**4*e**3*x + 10*a**3*b**3*d**3*e**4*x**2 + 10*a**3*b**3*d**2*e**5*x**3 + 5*a**2*b**4*d**6*e - 15*a**2*b**4*d**4*e**3*x**2 - 10*a**2*b**4*d**3*e**4*x**3 - a*b**5*d**7 + 3*a*b**5*d**6*e*x + 9*a*b**5*d**5*e**2*x**2 + 5*a*b**5*d**4*e**3*x**3 - ...)
```

3.172 $\int \sqrt{a+bx}(A+Bx)(d+ex)^{5/2} dx$

Optimal result	1571
Mathematica [A] (verified)	1572
Rubi [A] (verified)	1572
Maple [B] (verified)	1576
Fricas [A] (verification not implemented)	1577
Sympy [F]	1578
Maxima [F(-2)]	1579
Giac [B] (verification not implemented)	1579
Mupad [F(-1)]	1580
Reduce [B] (verification not implemented)	1581

Optimal result

Integrand size = 24, antiderivative size = 304

$$\int \sqrt{a+bx}(A+Bx)(d+ex)^{5/2} dx =$$

$$\begin{aligned} & - \frac{(bd-ae)^3(3bBd-10Abe+7aBe)\sqrt{a+bx}\sqrt{d+ex}}{128b^4e^2} \\ & - \frac{(bd-ae)^2(3bBd-10Abe+7aBe)(a+bx)^{3/2}\sqrt{d+ex}}{64b^4e} \\ & - \frac{(bd-ae)(3bBd-10Abe+7aBe)(a+bx)^{3/2}(d+ex)^{3/2}}{48b^3e} \\ & - \frac{(3bBd-10Abe+7aBe)(a+bx)^{3/2}(d+ex)^{5/2}}{40b^2e} + \frac{B(a+bx)^{3/2}(d+ex)^{7/2}}{5be} \\ & + \frac{(bd-ae)^4(3bBd-10Abe+7aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{128b^{9/2}e^{5/2}} \end{aligned}$$

output

```
-1/128*(-a*e+b*d)^3*(-10*A*b*e+7*B*a*e+3*B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(1/2)
)/b^4/e^2-1/64*(-a*e+b*d)^2*(-10*A*b*e+7*B*a*e+3*B*b*d)*(b*x+a)^(3/2)*(e*x
+d)^(1/2)/b^4/e-1/48*(-a*e+b*d)*(-10*A*b*e+7*B*a*e+3*B*b*d)*(b*x+a)^(3/2)*
(e*x+d)^(3/2)/b^3/e-1/40*(-10*A*b*e+7*B*a*e+3*B*b*d)*(b*x+a)^(3/2)*(e*x+d)
^(5/2)/b^2/e+1/5*B*(b*x+a)^(3/2)*(e*x+d)^(7/2)/b/e+1/128*(-a*e+b*d)^4*(-10
*A*b*e+7*B*a*e+3*B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)^(1/2
))/b^(9/2)/e^(5/2)
```


Mathematica [A] (verified)

Time = 0.71 (sec) , antiderivative size = 322, normalized size of antiderivative = 1.06

$$\int \sqrt{a+bx}(A+Bx)(d+ex)^{5/2} dx = \frac{\sqrt{a+bx}\sqrt{d+ex}(-105a^4Be^4 + 10a^3be^3(34Bd + 15Ae + 7Bex) - 2a^2b^2e^2(25Ae(11d + 2ex) + (bd - ae)^4(3bBd - 10Abe + 7aBe))\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{128b^{9/2}e^{5/2}}$$

input `Integrate[Sqrt[a + b*x]*(A + B*x)*(d + e*x)^(5/2), x]`

output `(Sqrt[a + b*x]*Sqrt[d + e*x]*(-105*a^4*B*e^4 + 10*a^3*b*e^3*(34*B*d + 15*A*e + 7*B*e*x) - 2*a^2*b^2*e^2*(25*A*e*(11*d + 2*e*x) + B*(173*d^2 + 111*d*e*x + 28*e^2*x^2)) + 2*a*b^3*e*(5*A*e*(73*d^2 + 36*d*e*x + 8*e^2*x^2) + B*(30*d^3 + 109*d^2*e*x + 88*d*e^2*x^2 + 24*e^3*x^3)) + b^4*(10*A*e*(15*d^3 + 118*d^2*e*x + 136*d*e^2*x^2 + 48*e^3*x^3) + B*(-45*d^4 + 30*d^3*e*x + 74*d^2*e^2*x^2 + 1008*d*e^3*x^3 + 384*e^4*x^4)))/(1920*b^4*e^2) + ((b*d - a*e)^4*(3*b*B*d - 10*A*b*e + 7*a*B*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(128*b^(9/2)*e^(5/2))`

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 252, normalized size of antiderivative = 0.83, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.292$, Rules used = {90, 60, 60, 60, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{a+bx}(A+Bx)(d+ex)^{5/2} dx$$

↓ 90

$$\frac{B(a+bx)^{3/2}(d+ex)^{7/2}}{5be} - \frac{(7aBe - 10Abe + 3bBd) \int \sqrt{a+bx}(d+ex)^{5/2} dx}{10be}$$

$$\begin{array}{c}
 \downarrow 60 \\
 \frac{B(a+bx)^{3/2}(d+ex)^{7/2}}{5be} - \\
 (7aBe - 10Abe + 3bBd) \left(\frac{5(bd-ae) \int \sqrt{a+bx}(d+ex)^{3/2} dx}{8b} + \frac{(a+bx)^{3/2}(d+ex)^{5/2}}{4b} \right) \\
 \hline
 10be \\
 \downarrow 60 \\
 \frac{B(a+bx)^{3/2}(d+ex)^{7/2}}{5be} - \\
 (7aBe - 10Abe + 3bBd) \left(\frac{5(bd-ae) \left(\frac{(bd-ae) \int \sqrt{a+bx} \sqrt{d+ex} dx}{2b} + \frac{(a+bx)^{3/2}(d+ex)^{3/2}}{3b} \right)}{8b} + \frac{(a+bx)^{3/2}(d+ex)^{5/2}}{4b} \right) \\
 \hline
 10be \\
 \downarrow 60 \\
 \frac{B(a+bx)^{3/2}(d+ex)^{7/2}}{5be} - \\
 (7aBe - 10Abe + 3bBd) \left(\frac{5(bd-ae) \left(\frac{(bd-ae) \left(\frac{\int \sqrt{a+bx} dx}{4b} + \frac{(a+bx)^{3/2} \sqrt{d+ex}}{2b} \right)}{2b} + \frac{(a+bx)^{3/2}(d+ex)^{3/2}}{3b} \right)}{8b} + \frac{(a+bx)^{3/2}(d+ex)^{5/2}}{4b} \right) \\
 \hline
 10be \\
 \downarrow 60 \\
 \frac{B(a+bx)^{3/2}(d+ex)^{7/2}}{5be} - \\
 (7aBe - 10Abe + 3bBd) \left(\frac{5(bd-ae) \left(\frac{(bd-ae) \left(\frac{\sqrt{a+bx} \sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{\sqrt{a+bx} \sqrt{d+ex}} dx}{2e} \right)}{4b} + \frac{(a+bx)^{3/2} \sqrt{d+ex}}{2b} \right)}{2b} + \frac{(a+bx)^{3/2}(d+ex)^{5/2}}{3b} \right) \\
 \hline
 10be
 \end{array}$$

$$\begin{array}{c}
 \downarrow 66 \\
 \frac{B(a+bx)^{3/2}(d+ex)^{7/2}}{5be} - \\
 \left(\frac{(bd-ae) \left(\frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}} \right)}{(bd-ae) \frac{\sqrt{a+bx}\sqrt{d+ex}}{e}} + \frac{(a+bx)^{3/2}\sqrt{d+ex}}{2b} \right)}{5(bd-ae) \frac{2b}{2b}} + \frac{(a+bx)^{3/2}}{3b} \\
 \frac{(7aBe - 10Abe + 3bBd)}{8b} \\
 \hline
 10be \\
 \downarrow 221
 \end{array}$$

$$\frac{B(a+bx)^{3/2}(d+ex)^{7/2}}{5be} - \frac{(7aBe - 10Abe + 3bBd)}{10be} \left(\frac{(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{\sqrt{be}^{3/2}} \right) + \frac{(a+bx)^{3/2}\sqrt{d+ex}}{2b}}{2b} \right)$$

```
input Int[Sqrt[a + b*x]*(A + B*x)*(d + e*x)^(5/2), x]
```

```
output (B*(a + b*x)^(3/2)*(d + e*x)^(7/2))/(5*b*e) - ((3*b*B*d - 10*A*b*e + 7*a*B
*e)*(((a + b*x)^(3/2)*(d + e*x)^(5/2))/(4*b) + (5*(b*d - a*e)*(((a + b*x)^(
3/2)*(d + e*x)^(3/2))/(3*b) + ((b*d - a*e)*(((a + b*x)^(3/2)*Sqrt[d + e*x
])/ (2*b) + ((b*d - a*e)*((Sqrt[a + b*x]*Sqrt[d + e*x])/e - ((b*d - a*e)*Ar
cTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(Sqrt[b]*e^(3/2)))
)/(4*b)))/(2*b)))/(8*b)))/(10*b*e)
```

Defintions of rubi rules used

```
rule 60 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(
b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!Integer
Q[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinear
Q[a, b, c, d, m, n, x]
```

rule 66 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[
2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; Fre
eQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 90 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p
), x] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))),
x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p
+ 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n,
p}, x] && NeQ[n + p + 2, 0]`

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1371 vs. $2(260) = 520$.

Time = 0.27 (sec) , antiderivative size = 1372, normalized size of antiderivative = 4.51

method	result	size
default	Expression too large to display	1372

input `int((b*x+a)^(1/2)*(B*x+A)*(e*x+d)^(5/2),x,method=_RETURNVERBOSE)`

output

```

-1/3840*(b*x+a)^(1/2)*(e*x+d)^(1/2)*(-768*B*b^4*e^4*x^4*((e*x+d)*(b*x+a))^(
(1/2)*(b*e)^(1/2)-960*A*b^4*e^4*x^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-96
*B*a*b^3*e^4*x^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-2016*B*b^4*d*e^3*x^3*
((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-160*A*a*b^3*e^4*x^2*((e*x+d)*(b*x+a))^(
(1/2)*(b*e)^(1/2)-1460*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b^3*d^2*e^2
-680*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^3*b*d*e^3+692*B*((e*x+d)*(b*x
+a))^(1/2)*(b*e)^(1/2)*a^2*b^2*d^2*e^2-120*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)
^(1/2)*a*b^3*d^3*e+210*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^4*e^4+90*B*
((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^4*d^4-352*B*a*b^3*d*e^3*x^2*((e*x+d)
*(b*x+a))^(1/2)*(b*e)^(1/2)-720*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b^
3*d*e^3*x+444*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^2*b^2*d*e^3*x-436*B*
((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b^3*d^2*e^2*x+150*A*ln(1/2*(2*b*e*x+
2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^4*b*e^5+150*
A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/
2))*b^5*d^4*e+900*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+
a*e+d*b)/(b*e)^(1/2))*a^2*b^3*d^2*e^3-600*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*
x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^4*d^3*e^2+375*B*ln(1/2*(
2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^4*b*
d*e^4-450*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)
/(b*e)^(1/2))*a^3*b^2*d^2*e^3+150*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))...

```

Fricas [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 1044, normalized size of antiderivative = 3.43

$$\int \sqrt{a+bx}(A+Bx)(d+ex)^{5/2} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^(1/2)*(B*x+A)*(e*x+d)^(5/2),x, algorithm="fricas")
```

output

```

[-1/7680*(15*(3*B*b^5*d^5 - 5*(B*a*b^4 + 2*A*b^5)*d^4*e - 10*(B*a^2*b^3 -
4*A*a*b^4)*d^3*e^2 + 30*(B*a^3*b^2 - 2*A*a^2*b^3)*d^2*e^3 - 5*(5*B*a^4*b -
8*A*a^3*b^2)*d*e^4 + (7*B*a^5 - 10*A*a^4*b)*e^5)*sqrt(b*e)*log(8*b^2*e^2*x^
x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 - 4*(2*b*e*x + b*d + a*e)*sqrt(b*e)*sq
rt(b*x + a)*sqrt(e*x + d) + 8*(b^2*d*e + a*b*e^2)*x) - 4*(384*B*b^5*e^5*x^
4 - 45*B*b^5*d^4*e + 30*(2*B*a*b^4 + 5*A*b^5)*d^3*e^2 - 2*(173*B*a^2*b^3 -
365*A*a*b^4)*d^2*e^3 + 10*(34*B*a^3*b^2 - 55*A*a^2*b^3)*d*e^4 - 15*(7*B*a^
4*b - 10*A*a^3*b^2)*e^5 + 48*(21*B*b^5*d*e^4 + (B*a*b^4 + 10*A*b^5)*e^5)*
x^3 + 8*(93*B*b^5*d^2*e^3 + 2*(11*B*a*b^4 + 85*A*b^5)*d*e^4 - (7*B*a^2*b^3
- 10*A*a*b^4)*e^5)*x^2 + 2*(15*B*b^5*d^3*e^2 + (109*B*a*b^4 + 590*A*b^5)*
d^2*e^3 - 3*(37*B*a^2*b^3 - 60*A*a*b^4)*d*e^4 + 5*(7*B*a^3*b^2 - 10*A*a^2*
b^3)*e^5)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b^5*e^3), -1/3840*(15*(3*B*b^5*
d^5 - 5*(B*a*b^4 + 2*A*b^5)*d^4*e - 10*(B*a^2*b^3 - 4*A*a*b^4)*d^3*e^2 + 3
0*(B*a^3*b^2 - 2*A*a^2*b^3)*d^2*e^3 - 5*(5*B*a^4*b - 8*A*a^3*b^2)*d*e^4 +
(7*B*a^5 - 10*A*a^4*b)*e^5)*sqrt(-b*e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sq
rt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*e^2*x^2 + a*b*d*e + (b^2*d*e + a
*b*e^2)*x)) - 2*(384*B*b^5*e^5*x^4 - 45*B*b^5*d^4*e + 30*(2*B*a*b^4 + 5*A*
b^5)*d^3*e^2 - 2*(173*B*a^2*b^3 - 365*A*a*b^4)*d^2*e^3 + 10*(34*B*a^3*b^2
- 55*A*a^2*b^3)*d*e^4 - 15*(7*B*a^4*b - 10*A*a^3*b^2)*e^5 + 48*(21*B*b^5*d
*e^4 + (B*a*b^4 + 10*A*b^5)*e^5)*x^3 + 8*(93*B*b^5*d^2*e^3 + 2*(11*B*a...

```

Sympy [F]

$$\int \sqrt{a + bx}(A + Bx)(d + ex)^{5/2} dx = \int (A + Bx) \sqrt{a + bx}(d + ex)^{5/2} dx$$

input

```
integrate((b*x+a)**(1/2)*(B*x+A)*(e*x+d)**(5/2),x)
```

output

```
Integral((A + B*x)*sqrt(a + b*x)*(d + e*x)**(5/2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \sqrt{a+bx}(A+Bx)(d+ex)^{5/2} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(1/2)*(B*x+A)*(e*x+d)^(5/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2501 vs. 2(260) = 520.

Time = 0.47 (sec) , antiderivative size = 2501, normalized size of antiderivative = 8.23

$$\int \sqrt{a+bx}(A+Bx)(d+ex)^{5/2} dx = \text{Too large to display}$$

input `integrate((b*x+a)^(1/2)*(B*x+A)*(e*x+d)^(5/2),x, algorithm="giac")`

output

```
-1/1920*(1920*((b^2*d - a*b*e)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e))/sqrt(b*e) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))*sqrt(b*x + a)*A*a*d^2*abs(b)/b^2 - 20*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*(b*x + a)*(4*(b*x + a)*(6*(b*x + a)/b^3 + (b^12*d*e^5 - 25*a*b^11*e^6)/(b^14*e^6)) - (5*b^13*d^2*e^4 + 14*a*b^12*d*e^5 - 163*a^2*b^11*e^6)/(b^14*e^6)) + 3*(5*b^14*d^3*e^3 + 9*a*b^13*d^2*e^4 + 15*a^2*b^12*d*e^5 - 93*a^3*b^11*e^6)/(b^14*e^6))*sqrt(b*x + a) + 3*(5*b^4*d^4 + 4*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2 + 20*a^3*b*d*e^3 - 35*a^4*e^4)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*b^2*e^3))*B*d*e*abs(b)/b - 10*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*(b*x + a)*(4*(b*x + a)*(6*(b*x + a)/b^3 + (b^12*d*e^5 - 25*a*b^11*e^6)/(b^14*e^6)) - (5*b^13*d^2*e^4 + 14*a*b^12*d*e^5 - 163*a^2*b^11*e^6)/(b^14*e^6)) + 3*(5*b^14*d^3*e^3 + 9*a*b^13*d^2*e^4 + 15*a^2*b^12*d*e^5 - 93*a^3*b^11*e^6)/(b^14*e^6))*sqrt(b*x + a) + 3*(5*b^4*d^4 + 4*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2 + 20*a^3*b*d*e^3 - 35*a^4*e^4)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*b^2*e^3))*B*a*e^2*abs(b)/b^2 - 10*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*(b*x + a)*(4*(b*x + a)*(6*(b*x + a)/b^3 + (b^12*d*e^5 - 25*a*b^11*e^6)/(b^14*e^6)) - (5*b^13*d^2*e^4 + 14*a*b^12*d*e^5 - 163*a^2*b^11*e^6)/(b^14*e^6)) + 3*(5*b^14*d^3*e^3 + 9*a*b^13*d^2*e^4 + 15*a^2*b^12*d*e^5 - 93*a^3*b^11*e^6)/(b^14*e^6))*sqrt(b*x + a) + 3*(...
```

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a+bx}(A+Bx)(d+ex)^{5/2} dx = \int (A+Bx) \sqrt{a+bx} (d+ex)^{5/2} dx$$

input

```
int((A + B*x)*(a + b*x)^(1/2)*(d + e*x)^(5/2), x)
```

output

```
int((A + B*x)*(a + b*x)^(1/2)*(d + e*x)^(5/2), x)
```

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 651, normalized size of antiderivative = 2.14

$$\int \sqrt{a+bx}(A+Bx)(d+ex)^{5/2} dx = \frac{15\sqrt{ex+d}\sqrt{bx+a}a^4be^5 - 70\sqrt{ex+d}\sqrt{bx+a}a^3b^2de^4 - 10\sqrt{ex+d}\sqrt{bx+a}a^3b^2e^5x + 128\sqrt{ex+d}\sqrt{bx+a}a^3b^2de^4x - 10\sqrt{ex+d}\sqrt{bx+a}a^3b^2e^5x^2 + 46\sqrt{ex+d}\sqrt{bx+a}a^3b^2de^4x^2 + 8\sqrt{ex+d}\sqrt{bx+a}a^3b^2e^5x^3 + 70\sqrt{ex+d}\sqrt{bx+a}a^3b^2de^4x^3 + 466\sqrt{ex+d}\sqrt{bx+a}a^3b^2e^5x^4 + 512\sqrt{ex+d}\sqrt{bx+a}a^3b^2de^4x^4 + 176\sqrt{ex+d}\sqrt{bx+a}a^3b^2e^5x^5 - 15\sqrt{ex+d}\sqrt{bx+a}a^3b^2de^4e + 10\sqrt{ex+d}\sqrt{bx+a}a^3b^2e^5de^3x^2 + 248\sqrt{ex+d}\sqrt{bx+a}a^3b^2de^4x^3 + 128\sqrt{ex+d}\sqrt{bx+a}a^3b^2e^5x^4 - 15\sqrt{ex+d}\sqrt{bx+a}a^3b^2de^4e^5 + 75\sqrt{ex+d}\sqrt{bx+a}a^3b^2de^4e^5x^2 - 150\sqrt{ex+d}\sqrt{bx+a}a^3b^2de^4e^5x^3 + 150\sqrt{ex+d}\sqrt{bx+a}a^3b^2de^4e^5x^4 - 75\sqrt{ex+d}\sqrt{bx+a}a^3b^2de^4e^5x^5}{(640b^4e^3)}$$

input

```
int((b*x+a)^(1/2)*(B*x+A)*(e*x+d)^(5/2),x)
```

output

```
(15*sqrt(d + e*x)*sqrt(a + b*x)*a**4*b*e**5 - 70*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b**2*d*e**4 - 10*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b**2*e**5*x + 128*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**3*d**2*e**3 + 46*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**3*d*e**4*x + 8*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**3*e**5*x**2 + 70*sqrt(d + e*x)*sqrt(a + b*x)*a*b**4*d**3*e**2 + 466*sqrt(d + e*x)*sqrt(a + b*x)*a*b**4*d**2*e**3*x + 512*sqrt(d + e*x)*sqrt(a + b*x)*a*b**4*d*e**4*x**2 + 176*sqrt(d + e*x)*sqrt(a + b*x)*a*b**4*e**5*x**3 - 15*sqrt(d + e*x)*sqrt(a + b*x)*b**5*d**4*e + 10*sqrt(d + e*x)*sqrt(a + b*x)*b**5*d**3*e**2*x + 248*sqrt(d + e*x)*sqrt(a + b*x)*b**5*d**2*e**3*x**2 + 336*sqrt(d + e*x)*sqrt(a + b*x)*b**5*d*e**4*x**3 + 128*sqrt(d + e*x)*sqrt(a + b*x)*b**5*e**5*x**4 - 15*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**5*e**5 + 75*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**4*b*d*e**4 - 150*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**3*b**2*d**2*e**3 + 150*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**2*b**3*d**3*e**2 - 75*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*b**4*d**4*e + 15*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**5*d**5)/(640*b**4*e**3)
```

3.173 $\int \sqrt{a + bx}(A + Bx)(d + ex)^{3/2} dx$

Optimal result	1582
Mathematica [A] (verified)	1583
Rubi [A] (verified)	1583
Maple [B] (verified)	1586
Fricas [A] (verification not implemented)	1587
Sympy [F]	1588
Maxima [F(-2)]	1588
Giac [B] (verification not implemented)	1589
Mupad [F(-1)]	1590
Reduce [B] (verification not implemented)	1590

Optimal result

Integrand size = 24, antiderivative size = 250

$$\int \sqrt{a + bx}(A + Bx)(d + ex)^{3/2} dx =$$

$$-\frac{(bd - ae)^2(3bBd - 8Abe + 5aBe)\sqrt{a + bx}\sqrt{d + ex}}{64b^3e^2}$$

$$-\frac{(bd - ae)(3bBd - 8Abe + 5aBe)(a + bx)^{3/2}\sqrt{d + ex}}{32b^3e}$$

$$-\frac{(3bBd - 8Abe + 5aBe)(a + bx)^{3/2}(d + ex)^{3/2}}{24b^2e} + \frac{B(a + bx)^{3/2}(d + ex)^{5/2}}{4be}$$

$$+ \frac{(bd - ae)^3(3bBd - 8Abe + 5aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{64b^{7/2}e^{5/2}}$$

output

```
-1/64*(-a*e+b*d)^2*(-8*A*b*e+5*B*a*e+3*B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(1/2)/
b^3/e^2-1/32*(-a*e+b*d)*(-8*A*b*e+5*B*a*e+3*B*b*d)*(b*x+a)^(3/2)*(e*x+d)^(
1/2)/b^3/e-1/24*(-8*A*b*e+5*B*a*e+3*B*b*d)*(b*x+a)^(3/2)*(e*x+d)^(3/2)/b^2
/e+1/4*B*(b*x+a)^(3/2)*(e*x+d)^(5/2)/b/e+1/64*(-a*e+b*d)^3*(-8*A*b*e+5*B*a
*e+3*B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)^(1/2))/b^(7/2)/e
^(5/2)
```

Mathematica [A] (verified)

Time = 0.65 (sec) , antiderivative size = 231, normalized size of antiderivative = 0.92

$$\int \sqrt{a+bx}(A+Bx)(d+ex)^{3/2} dx = \frac{\sqrt{a+bx}\sqrt{d+ex}(15a^3Be^3 - a^2be^2(31Bd + 24Ae + 10Bex) + ab^2e(16Ae(4d + ex) + B(9d^2 + 20de + 8e^2x^2)) + b^3(8Ae(3d^2 + 14de + 8e^2x^2) + B(-9d^3 + 6d^2ex + 72de^2x^2 + 48e^3x^3)))}{192b^3e^2} + \frac{(bd - ae)^3(3bBd - 8Abe + 5aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{64b^{7/2}e^{5/2}}$$

input `Integrate[Sqrt[a + b*x]*(A + B*x)*(d + e*x)^(3/2), x]`

output `(Sqrt[a + b*x]*Sqrt[d + e*x]*(15*a^3*B*e^3 - a^2*b*e^2*(31*B*d + 24*A*e + 10*B*e*x) + a*b^2*e*(16*A*e*(4*d + e*x) + B*(9*d^2 + 20*d*e*x + 8*e^2*x^2)) + b^3*(8*A*e*(3*d^2 + 14*d*e*x + 8*e^2*x^2) + B*(-9*d^3 + 6*d^2*e*x + 72*d*e^2*x^2 + 48*e^3*x^3)))/(192*b^3*e^2) + ((b*d - a*e)^3*(3*b*B*d - 8*A*b*e + 5*a*B*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(64*b^(7/2)*e^(5/2))`

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 211, normalized size of antiderivative = 0.84, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {90, 60, 60, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{a+bx}(A+Bx)(d+ex)^{3/2} dx$$

$$\downarrow 90$$

$$\frac{B(a+bx)^{3/2}(d+ex)^{5/2}}{4be} - \frac{(5aBe - 8Abe + 3bBd) \int \sqrt{a+bx}(d+ex)^{3/2} dx}{8be}$$

$$\downarrow 60$$

$$\begin{aligned}
 & \frac{B(a+bx)^{3/2}(d+ex)^{5/2}}{(5aBe - 8Abe + 3bBd) \left(\frac{4be}{2b} \frac{(bd-ae) \int \sqrt{a+bx} \sqrt{d+ex} dx}{\sqrt{d+ex}} + \frac{(a+bx)^{3/2}(d+ex)^{3/2}}{3b} \right)}{8be} \\
 & \quad \downarrow 60 \\
 & \frac{B(a+bx)^{3/2}(d+ex)^{5/2}}{(5aBe - 8Abe + 3bBd) \left(\frac{4be}{2b} \frac{(bd-ae) \left(\frac{(bd-ae) \int \frac{\sqrt{a+bx} dx}{\sqrt{d+ex}} + \frac{(a+bx)^{3/2} \sqrt{d+ex}}{2b} \right)}{\sqrt{d+ex}} + \frac{(a+bx)^{3/2}(d+ex)^{3/2}}{3b} \right)}{8be} \\
 & \quad \downarrow 60 \\
 & \frac{B(a+bx)^{3/2}(d+ex)^{5/2}}{(5aBe - 8Abe + 3bBd) \left(\frac{4be}{2b} \frac{(bd-ae) \left(\frac{\sqrt{a+bx} \sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{\sqrt{a+bx} \sqrt{d+ex}} dx}{2e} \right)}{\sqrt{d+ex}} + \frac{(a+bx)^{3/2} \sqrt{d+ex}}{2b} \right)}{8be} + \frac{(a+bx)^{3/2}(d+ex)^{3/2}}{3b} \\
 & \quad \downarrow 66 \\
 & \frac{B(a+bx)^{3/2}(d+ex)^{5/2}}{(5aBe - 8Abe + 3bBd) \left(\frac{4be}{2b} \frac{(bd-ae) \left(\frac{\sqrt{a+bx} \sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{e} \right)}{\sqrt{d+ex}} + \frac{(a+bx)^{3/2} \sqrt{d+ex}}{2b} \right)}{8be} + \frac{(a+bx)^{3/2}(d+ex)^{3/2}}{3b} \\
 & \quad \downarrow 221
 \end{aligned}$$

$$\frac{B(a+bx)^{3/2}(d+ex)^{5/2}}{4be} - \frac{(5aBe - 8Abe + 3bBd) \left(\frac{(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{\sqrt{be}^{3/2}} \right)}{4b} + \frac{(a+bx)^{3/2}\sqrt{d+ex}}{2b} \right)}{2b} + \frac{(a+bx)^{3/2}(d+ex)^{5/2}}{3b}}{8be}$$

input `Int[Sqrt[a + b*x]*(A + B*x)*(d + e*x)^(3/2), x]`

output `(B*(a + b*x)^(3/2)*(d + e*x)^(5/2))/(4*b*e) - ((3*b*B*d - 8*A*b*e + 5*a*B*e)*(((a + b*x)^(3/2)*(d + e*x)^(3/2))/(3*b) + ((b*d - a*e)*((a + b*x)^(3/2)*Sqrt[d + e*x])/(2*b) + ((b*d - a*e)*((Sqrt[a + b*x]*Sqrt[d + e*x])/e - ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(Sqrt[b]*e^(3/2)))))/(4*b))/(2*b))/(8*b*e)`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 967 vs. $2(212) = 424$.

Time = 0.26 (sec) , antiderivative size = 968, normalized size of antiderivative = 3.87

method	result
default	$\frac{\sqrt{bx+a}\sqrt{ex+d}\left(96Bb^3e^3x^3\sqrt{(ex+d)(bx+a)}\sqrt{be}+128Ab^3e^3x^2\sqrt{(ex+d)(bx+a)}\sqrt{be}+144Bb^3de^2x^2\sqrt{(ex+d)(bx+a)}\sqrt{be}-15B\right)}{\dots}$

input `int((b*x+a)^(1/2)*(B*x+A)*(e*x+d)^(3/2),x,method=_RETURNVERBOSE)`

output

```

1/384*(b*x+a)^(1/2)*(e*x+d)^(1/2)*(96*B*b^3*e^3*x^3*((e*x+d)*(b*x+a))^(1/2)
)*(b*e)^(1/2)+128*A*b^3*e^3*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+144*B*
b^3*d*e^2*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-15*B*ln(1/2*(2*b*e*x+2*(
(e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^4*e^4+9*B*ln(1/
2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^4
*d^4+40*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b^2*d*e^2*x-72*A*ln(1/2*(2
*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*b^2
*d*e^3+72*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)
/(b*e)^(1/2))*a*b^3*d^2*e^2+32*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b^2
*e^3*x+224*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^3*d*e^2*x-20*B*((e*x+d)
*(b*x+a))^(1/2)*(b*e)^(1/2)*a^2*b*e^3*x+12*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)
^(1/2)*b^3*d^2*e*x+30*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^3*e^3-18*B*((
e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^3*d^3+24*A*ln(1/2*(2*b*e*x+2*((e*x+d)
*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^3*b*e^4-24*A*ln(1/2*(2
*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^4*d^3
*e+128*A*a*b^2*d*e^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+16*B*a*b^2*e^3*x^
2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-12*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x
+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^3*d^3*e-48*A*((e*x+d)*(b
x+a))^(1/2)*(b*e)^(1/2)*a^2*b*e^3+48*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)
*b^3*d^2*e+36*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a...

```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 766, normalized size of antiderivative = 3.06

$$\int \sqrt{a+bx}(A+Bx)(d+ex)^{3/2} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^(1/2)*(B*x+A)*(e*x+d)^(3/2),x, algorithm="fricas")
```


output

```
[1/768*(3*(3*B*b^4*d^4 - 4*(B*a*b^3 + 2*A*b^4)*d^3*e - 6*(B*a^2*b^2 - 4*A*
a*b^3)*d^2*e^2 + 12*(B*a^3*b - 2*A*a^2*b^2)*d*e^3 - (5*B*a^4 - 8*A*a^3*b)*
e^4)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*b*
e*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x + a)*sqrt(e*x + d) + 8*(b^2*d*e + a*b*
e^2)*x) + 4*(48*B*b^4*e^4*x^3 - 9*B*b^4*d^3*e + 3*(3*B*a*b^3 + 8*A*b^4)*d^
2*e^2 - (31*B*a^2*b^2 - 64*A*a*b^3)*d*e^3 + 3*(5*B*a^3*b - 8*A*a^2*b^2)*e^
4 + 8*(9*B*b^4*d*e^3 + (B*a*b^3 + 8*A*b^4)*e^4)*x^2 + 2*(3*B*b^4*d^2*e^2 +
2*(5*B*a*b^3 + 28*A*b^4)*d*e^3 - (5*B*a^2*b^2 - 8*A*a*b^3)*e^4)*x)*sqrt(b
*x + a)*sqrt(e*x + d))/(b^4*e^3), -1/384*(3*(3*B*b^4*d^4 - 4*(B*a*b^3 + 2*
A*b^4)*d^3*e - 6*(B*a^2*b^2 - 4*A*a*b^3)*d^2*e^2 + 12*(B*a^3*b - 2*A*a^2*b
^2)*d*e^3 - (5*B*a^4 - 8*A*a^3*b)*e^4)*sqrt(-b*e)*arctan(1/2*(2*b*e*x + b*
d + a*e)*sqrt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*e^2*x^2 + a*b*d*e + (
b^2*d*e + a*b*e^2)*x)) - 2*(48*B*b^4*e^4*x^3 - 9*B*b^4*d^3*e + 3*(3*B*a*b^
3 + 8*A*b^4)*d^2*e^2 - (31*B*a^2*b^2 - 64*A*a*b^3)*d*e^3 + 3*(5*B*a^3*b -
8*A*a^2*b^2)*e^4 + 8*(9*B*b^4*d*e^3 + (B*a*b^3 + 8*A*b^4)*e^4)*x^2 + 2*(3*
B*b^4*d^2*e^2 + 2*(5*B*a*b^3 + 28*A*b^4)*d*e^3 - (5*B*a^2*b^2 - 8*A*a*b^3)
*e^4)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b^4*e^3)]
```

Sympy [F]

$$\int \sqrt{a+bx}(A+Bx)(d+ex)^{3/2} dx = \int (A+Bx) \sqrt{a+bx}(d+ex)^{\frac{3}{2}} dx$$

input

```
integrate((b*x+a)**(1/2)*(B*x+A)*(e*x+d)**(3/2),x)
```

output

```
Integral((A + B*x)*sqrt(a + b*x)*(d + e*x)**(3/2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \sqrt{a+bx}(A+Bx)(d+ex)^{3/2} dx = \text{Exception raised: ValueError}$$

input

```
integrate((b*x+a)^(1/2)*(B*x+A)*(e*x+d)^(3/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1366 vs. $2(212) = 424$.

Time = 0.32 (sec) , antiderivative size = 1366, normalized size of antiderivative = 5.46

$$\int \sqrt{a+bx}(A+Bx)(d+ex)^{3/2} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^(1/2)*(B*x+A)*(e*x+d)^(3/2),x, algorithm="giac")
```

output

```
-1/192*(192*((b^2*d - a*b*e)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d
+ (b*x + a)*b*e - a*b*e)))/sqrt(b*e) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e
)*sqrt(b*x + a))*A*a*d*abs(b)/b^2 - (sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(
2*(b*x + a)*(4*(b*x + a)*(6*(b*x + a)/b^3 + (b^12*d*e^5 - 25*a*b^11*e^6)/(
b^14*e^6)) - (5*b^13*d^2*e^4 + 14*a*b^12*d*e^5 - 163*a^2*b^11*e^6)/(b^14*e
^6)) + 3*(5*b^14*d^3*e^3 + 9*a*b^13*d^2*e^4 + 15*a^2*b^12*d*e^5 - 93*a^3*b
^11*e^6)/(b^14*e^6))*sqrt(b*x + a) + 3*(5*b^4*d^4 + 4*a*b^3*d^3*e + 6*a^2*
b^2*d^2*e^2 + 20*a^3*b*d*e^3 - 35*a^4*e^4)*log(abs(-sqrt(b*e)*sqrt(b*x + a
) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*b^2*e^3))*B*e*abs(b)/
b - 48*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*b*x + 2*a + (b*d*e - 5*a*e^
2)/e^2)*sqrt(b*x + a) + (b^3*d^2 + 2*a*b^2*d*e - 3*a^2*b*e^2)*log(abs(-sqr
t(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*e)
)*B*a*d*abs(b)/b^3 - 48*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*b*x + 2*a
+ (b*d*e - 5*a*e^2)/e^2)*sqrt(b*x + a) + (b^3*d^2 + 2*a*b^2*d*e - 3*a^2*b*
e^2)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e
)))/(sqrt(b*e)*e))*A*d*abs(b)/b^2 - 48*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e
)*(2*b*x + 2*a + (b*d*e - 5*a*e^2)/e^2)*sqrt(b*x + a) + (b^3*d^2 + 2*a*b^2
*d*e - 3*a^2*b*e^2)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x +
a)*b*e - a*b*e)))/(sqrt(b*e)*e))*A*a*e*abs(b)/b^3 - 8*(sqrt(b^2*d + (b*x
+ a)*b*e - a*b*e)*(2*(4*b*x + 4*a + (b*d*e^3 - 13*a*e^4)/e^4)*(b*x + a)...
```

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a+bx}(A+Bx)(d+ex)^{3/2} dx = \int (A+Bx) \sqrt{a+bx} (d+ex)^{3/2} dx$$

input `int((A + B*x)*(a + b*x)^(1/2)*(d + e*x)^(3/2), x)`

output `int((A + B*x)*(a + b*x)^(1/2)*(d + e*x)^(3/2), x)`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 471, normalized size of antiderivative = 1.88

$$\int \sqrt{a+bx}(A+Bx)(d+ex)^{3/2} dx = \frac{-3\sqrt{ex+d}\sqrt{bx+a}a^3be^4 + 11\sqrt{ex+d}\sqrt{bx+a}a^2b^2de^3 + 2\sqrt{ex+d}\sqrt{bx+a}a^2b^2e^4x + 11\sqrt{ex+d}\sqrt{bx+a}a^3be^4 + 11\sqrt{ex+d}\sqrt{bx+a}a^2b^2de^3 + 2\sqrt{ex+d}\sqrt{bx+a}a^2b^2e^4x + 11\sqrt{ex+d}\sqrt{bx+a}a^3be^4}{(d+ex)^{3/2}}$$

input `int((b*x+a)^(1/2)*(B*x+A)*(e*x+d)^(3/2), x)`

output `(- 3*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b*e**4 + 11*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**2*d*e**3 + 2*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**2*e**4*x + 11*sqrt(d + e*x)*sqrt(a + b*x)*a*b**3*d**2*e**2 + 44*sqrt(d + e*x)*sqrt(a + b*x)*a*b**3*d*e**3*x + 24*sqrt(d + e*x)*sqrt(a + b*x)*a*b**3*e**4*x**2 - 3*sqrt(d + e*x)*sqrt(a + b*x)*b**4*d**3*e + 2*sqrt(d + e*x)*sqrt(a + b*x)*b**4*d**2*e**2*x + 24*sqrt(d + e*x)*sqrt(a + b*x)*b**4*d*e**3*x**2 + 16*sqrt(d + e*x)*sqrt(a + b*x)*b**4*e**4*x**3 + 3*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**4*e**4 - 12*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**3*b*d*e**3 + 18*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**2*b**2*d**2*e**2 - 12*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*b**3*d**3*e + 3*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**4*d**4)/(64*b**3*e**3)`

3.174 $\int \sqrt{a + bx}(A + Bx)\sqrt{d + ex} dx$

Optimal result	1591
Mathematica [A] (verified)	1592
Rubi [A] (verified)	1592
Maple [B] (verified)	1594
Fricas [A] (verification not implemented)	1595
Sympy [F]	1596
Maxima [F(-2)]	1596
Giac [B] (verification not implemented)	1597
Mupad [B] (verification not implemented)	1597
Reduce [B] (verification not implemented)	1598

Optimal result

Integrand size = 24, antiderivative size = 190

$$\int \sqrt{a + bx}(A + Bx)\sqrt{d + ex} dx = -\frac{(bd - ae)(bBd - 2Abe + aBe)\sqrt{a + bx}\sqrt{d + ex}}{8b^2e^2} - \frac{(bBd - 2Abe + aBe)(a + bx)^{3/2}\sqrt{d + ex}}{4b^2e} + \frac{B(a + bx)^{3/2}(d + ex)^{3/2}}{3be} + \frac{(bd - ae)^2(bBd - 2Abe + aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a + bx}}{\sqrt{b}\sqrt{d + ex}}\right)}{8b^{5/2}e^{5/2}}$$

output

```
-1/8*(-a*e+b*d)*(-2*A*b*e+B*a*e+B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(1/2)/b^2/e^2
-1/4*(-2*A*b*e+B*a*e+B*b*d)*(b*x+a)^(3/2)*(e*x+d)^(1/2)/b^2/e+1/3*B*(b*x+a)^(3/2)*(e*x+d)^(3/2)/b/e+1/8*(-a*e+b*d)^2*(-2*A*b*e+B*a*e+B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)^(1/2))/b^(5/2)/e^(5/2)
```

Mathematica [A] (verified)

Time = 0.39 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.84

$$\int \sqrt{a+bx}(A+Bx)\sqrt{d+ex} dx$$

$$= \frac{\sqrt{a+bx}\sqrt{d+ex}(-3a^2Be^2 + 2abe(3Ae + B(d+ex)) + b^2(6Ae(d+2ex) + B(-3d^2 + 2dex + 8e^2x^2)))}{24b^2e^2}$$

$$+ \frac{(bd-ae)^2(bBd - 2Abe + aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{e}\sqrt{a+bx}}\right)}{8b^{5/2}e^{5/2}}$$

input `Integrate[Sqrt[a + b*x]*(A + B*x)*Sqrt[d + e*x], x]`

output `(Sqrt[a + b*x]*Sqrt[d + e*x]*(-3*a^2*B*e^2 + 2*a*b*e*(3*A*e + B*(d + e*x)) + b^2*(6*A*e*(d + 2*e*x) + B*(-3*d^2 + 2*d*e*x + 8*e^2*x^2)))/(24*b^2*e^2) + ((b*d - a*e)^2*(b*B*d - 2*A*b*e + a*B*e)*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/(Sqrt[e]*Sqrt[a + b*x])])/(8*b^(5/2)*e^(5/2))`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 170, normalized size of antiderivative = 0.89, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$, Rules used = {90, 60, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{a+bx}(A+Bx)\sqrt{d+ex} dx$$

$$\downarrow 90$$

$$\frac{(2Abe - B(ae + bd)) \int \sqrt{a+bx}\sqrt{d+ex} dx}{2be} + \frac{B(a+bx)^{3/2}(d+ex)^{3/2}}{3be}$$

$$\downarrow 60$$

$$\frac{(2Abe - B(ae + bd)) \left(\frac{(bd-ae) \int \frac{\sqrt{a+bx}}{\sqrt{d+ex}} dx}{4b} + \frac{(a+bx)^{3/2}\sqrt{d+ex}}{2b} \right)}{2be} + \frac{B(a+bx)^{3/2}(d+ex)^{3/2}}{3be}$$

$$\begin{aligned}
 & \downarrow 60 \\
 & (2Abe - B(ae + bd)) \left(\frac{(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{2e} \right)}{4b} + \frac{(a+bx)^{3/2}\sqrt{d+ex}}{2b} \right) \\
 & \hline
 & \frac{B(a+bx)^{3/2}(d+ex)^{3/2}}{3be} \\
 & \downarrow 66 \\
 & (2Abe - B(ae + bd)) \left(\frac{(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{e} \right)}{4b} + \frac{(a+bx)^{3/2}\sqrt{d+ex}}{2b} \right) \\
 & \hline
 & \frac{B(a+bx)^{3/2}(d+ex)^{3/2}}{3be} \\
 & \downarrow 221 \\
 & (2Abe - B(ae + bd)) \left(\frac{(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \operatorname{arctanh} \left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}} \right)}{\sqrt{be}^{3/2}} \right)}{4b} + \frac{(a+bx)^{3/2}\sqrt{d+ex}}{2b} \right) \\
 & \hline
 & \frac{B(a+bx)^{3/2}(d+ex)^{3/2}}{3be}
 \end{aligned}$$

input `Int[Sqrt[a + b*x]*(A + B*x)*Sqrt[d + e*x],x]`

output `(B*(a + b*x)^(3/2)*(d + e*x)^(3/2))/(3*b*e) + ((2*A*b*e - B*(b*d + a*e))*((a + b*x)^(3/2)*Sqrt[d + e*x])/(2*b) + ((b*d - a*e)*((Sqrt[a + b*x]*Sqrt[d + e*x])/e - ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(Sqrt[b]*e^(3/2))))/(4*b))/(2*b*e)`

Definitions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 635 vs. $2(158) = 316$.

Time = 0.26 (sec) , antiderivative size = 636, normalized size of antiderivative = 3.35

method	result
default	$-\frac{\sqrt{bx+a}\sqrt{ex+d}\left(-16Bb^2e^2x^2\sqrt{(ex+d)(bx+a)}\sqrt{be}+6A\ln\left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}}\right)\right)a^2be^3-12A\ln\left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}}\right)}{2}$

input `int((b*x+a)^(1/2)*(B*x+A)*(e*x+d)^(1/2),x,method=_RETURNVERBOSE)`

output

```

-1/48*(b*x+a)^(1/2)*(e*x+d)^(1/2)*(-16*B*b^2*e^2*x^2*((e*x+d)*(b*x+a))^(1/2)
*(b*e)^(1/2)+6*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a
*e+d*b)/(b*e)^(1/2))*a^2*b*e^3-12*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)
*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^2*d*e^2+6*A*ln(1/2*(2*b*e*x+2*((e
*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^3*d^2*e-24*A*((e
*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^2*e^2*x-3*B*ln(1/2*(2*b*e*x+2*((e*x+d)*
(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^3*e^3+3*B*ln(1/2*(2*b*
e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*b*d*e^2
+3*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(
1/2))*a*b^2*d^2*e-3*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)
+a*e+d*b)/(b*e)^(1/2))*b^3*d^3-4*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a
*b*e^2*x-4*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^2*d*e*x-12*A*((e*x+d)*
(b*x+a))^(1/2)*(b*e)^(1/2)*a*b*e^2-12*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)
*b^2*d*e+6*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^2*e^2-4*B*((e*x+d)*(b*x
+a))^(1/2)*(b*e)^(1/2)*a*b*d*e+6*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^2
*d^2)/((e*x+d)*(b*x+a))^(1/2)/b^2/e^2/(b*e)^(1/2)

```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 528, normalized size of antiderivative = 2.78

$$\int \sqrt{a+bx}(A+Bx)\sqrt{d+ex} dx$$

$$= \left[\frac{3(Bb^3d^3 - (Bab^2 + 2Ab^3)d^2e - (Ba^2b - 4Aab^2)de^2 + (Ba^3 - 2Aa^2b)e^3)\sqrt{be} \log\left(8b^2e^2x^2 + b^2d^2 + \dots\right)}{3(Bb^3d^3 - (Bab^2 + 2Ab^3)d^2e - (Ba^2b - 4Aab^2)de^2 + (Ba^3 - 2Aa^2b)e^3)\sqrt{-be} \arctan\left(\frac{(2bex+bd+ae)\sqrt{be}}{2(b^2e^2x^2+abd)}\right)} \right]$$

input

```
integrate((b*x+a)^(1/2)*(B*x+A)*(e*x+d)^(1/2),x, algorithm="fricas")
```


output

```
[-1/96*(3*(B*b^3*d^3 - (B*a*b^2 + 2*A*b^3)*d^2*e - (B*a^2*b - 4*A*a*b^2)*d
*e^2 + (B*a^3 - 2*A*a^2*b)*e^3)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*
a*b*d*e + a^2*e^2 - 4*(2*b*e*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x + a)*sqrt(e
*x + d) + 8*(b^2*d*e + a*b*e^2)*x) - 4*(8*B*b^3*e^3*x^2 - 3*B*b^3*d^2*e +
2*(B*a*b^2 + 3*A*b^3)*d*e^2 - 3*(B*a^2*b - 2*A*a*b^2)*e^3 + 2*(B*b^3*d*e^2
+ (B*a*b^2 + 6*A*b^3)*e^3)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b^3*e^3), -1/
48*(3*(B*b^3*d^3 - (B*a*b^2 + 2*A*b^3)*d^2*e - (B*a^2*b - 4*A*a*b^2)*d*e^2
+ (B*a^3 - 2*A*a^2*b)*e^3)*sqrt(-b*e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sq
rt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*e^2*x^2 + a*b*d*e + (b^2*d*e + a
*b*e^2)*x)) - 2*(8*B*b^3*e^3*x^2 - 3*B*b^3*d^2*e + 2*(B*a*b^2 + 3*A*b^3)*d
*e^2 - 3*(B*a^2*b - 2*A*a*b^2)*e^3 + 2*(B*b^3*d*e^2 + (B*a*b^2 + 6*A*b^3)*
e^3)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b^3*e^3)]
```

Sympy [F]

$$\int \sqrt{a+bx}(A+Bx)\sqrt{d+ex} dx = \int (A+Bx)\sqrt{a+bx}\sqrt{d+ex} dx$$

input

```
integrate((b*x+a)**(1/2)*(B*x+A)*(e*x+d)**(1/2),x)
```

output

```
Integral((A + B*x)*sqrt(a + b*x)*sqrt(d + e*x), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \sqrt{a+bx}(A+Bx)\sqrt{d+ex} dx = \text{Exception raised: ValueError}$$

input

```
integrate((b*x+a)^(1/2)*(B*x+A)*(e*x+d)^(1/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 557 vs. $2(158) = 316$.

Time = 0.20 (sec) , antiderivative size = 557, normalized size of antiderivative = 2.93

$$\int \sqrt{a+bx}(A+Bx)\sqrt{d+ex} dx = \frac{24 \left(\frac{(b^2d-abe) \log\left(\frac{-\sqrt{be}\sqrt{bx+a} + \sqrt{b^2d+(bx+a)be-abe}}{\sqrt{be}}\right) - \sqrt{b^2d+(bx+a)be-abe}\sqrt{bx+a}}{b^2} \right) Aa|b| - 6 \left(\sqrt{b^2d+(bx+a)be-abe} (2bx+2a + \frac{bde}{\sqrt{b^2d+(bx+a)be-abe}}) \right)}{b^2}$$

input `integrate((b*x+a)^(1/2)*(B*x+A)*(e*x+d)^(1/2),x, algorithm="giac")`

output `-1/24*(24*((b^2*d - a*b*e)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e))/sqrt(b*e) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*sqrt(b*x + a))*A*a*abs(b)/b^2 - 6*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*b*x + 2*a + (b*d*e - 5*a*e^2)/e^2)*sqrt(b*x + a) + (b^3*d^2 + 2*a*b^2*d*e - 3*a^2*b*e^2)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*e))*B*a*abs(b)/b^3 - 6*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*b*x + 2*a + (b*d*e - 5*a*e^2)/e^2)*sqrt(b*x + a) + (b^3*d^2 + 2*a*b^2*d*e - 3*a^2*b*e^2)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*e))*A*abs(b)/b^2 - (sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*(4*b*x + 4*a + (b*d*e^3 - 13*a*e^4)/e^4)*(b*x + a) - 3*(b^2*d^2*e^2 + 2*a*b*d*e^3 - 11*a^2*e^4)/e^4)*sqrt(b*x + a) - 3*(b^4*d^3 + a*b^3*d^2*e + 3*a^2*b^2*d*e^2 - 5*a^3*b*e^3)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*e^2))*B*abs(b)/b^3)/b`

Mupad [B] (verification not implemented)

Time = 44.92 (sec) , antiderivative size = 1207, normalized size of antiderivative = 6.35

$$\int \sqrt{a+bx}(A+Bx)\sqrt{d+ex} dx = \text{Too large to display}$$

input `int((A + B*x)*(a + b*x)^(1/2)*(d + e*x)^(1/2),x)`

output

```

A*(x/2 + (a*e + b*d)/(4*b*e))*(a + b*x)^(1/2)*(d + e*x)^(1/2) - (((a + b*x)^(1/2) - a^(1/2))*((B*b^6*d^3)/4 + (B*a^3*b^3*e^3)/4 - (B*a^2*b^4*d*e^2)/4 - (B*a*b^5*d^2*e)/4))/(e^8*((d + e*x)^(1/2) - d^(1/2))) - (((a + b*x)^(1/2) - a^(1/2))^3*((17*B*b^5*d^3)/12 + (17*B*a^3*b^2*e^3)/12 + (101*B*a^2*b^3*d*e^2)/4 + (101*B*a*b^4*d^2*e)/4))/(e^7*((d + e*x)^(1/2) - d^(1/2))^3) - (((a + b*x)^(1/2) - a^(1/2))^7*((19*B*a^3*e^3)/2 + (19*B*b^3*d^3)/2 + (269*B*a*b^2*d^2*e)/2 + (269*B*a^2*b*d*e^2)/2))/(e^5*((d + e*x)^(1/2) - d^(1/2))^7) - (((a + b*x)^(1/2) - a^(1/2))^5*((19*B*b^4*d^3)/2 + (19*B*a^3*b*e^3)/2 + (269*B*a^2*b^2*d*e^2)/2 + (269*B*a*b^3*d^2*e)/2))/(e^6*((d + e*x)^(1/2) - d^(1/2))^5) + (((a + b*x)^(1/2) - a^(1/2))^11*((B*a^3*e^3)/4 + (B*b^3*d^3)/4 - (B*a*b^2*d^2*e)/4 - (B*a^2*b*d*e^2)/4))/(b^2*e^3*((d + e*x)^(1/2) - d^(1/2))^11) - (((a + b*x)^(1/2) - a^(1/2))^9*((17*B*a^3*e^3)/12 + (17*B*b^3*d^3)/12 + (101*B*a*b^2*d^2*e)/4 + (101*B*a^2*b*d*e^2)/4))/(b*e^4*((d + e*x)^(1/2) - d^(1/2))^9) + (a^(1/2)*d^(1/2))*((a + b*x)^(1/2) - a^(1/2))^4*(32*B*b^4*d^2 + 32*B*a^2*b^2*e^2 + 96*B*a*b^3*d*e))/(e^6*((d + e*x)^(1/2) - d^(1/2))^4) + (8*B*a^(3/2)*d^(3/2))*((a + b*x)^(1/2) - a^(1/2))^10/(e^2*((d + e*x)^(1/2) - d^(1/2))^10) + (a^(1/2)*d^(1/2))*((a + b*x)^(1/2) - a^(1/2))^6*(64*B*b^3*d^2 + 64*B*a^2*b*e^2 + (656*B*a*b^2*d*e)/3))/(e^5*((d + e*x)^(1/2) - d^(1/2))^6) + (a^(1/2)*d^(1/2))*((a + b*x)^(1/2) - a^(1/2))^8*(32*B*a^2*e^2 + 32*B*b^2*d^2 + 96*B*a*b*d*e))/(e^4*((d + e*x)^(1/2) - d^(1/2))^8)

```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 320, normalized size of antiderivative = 1.68

$$\int \sqrt{a+bx}(A+Bx)\sqrt{d+ex} dx$$

$$= \frac{3\sqrt{ex+d}\sqrt{bx+a}a^2be^3 + 8\sqrt{ex+d}\sqrt{bx+a}ab^2de^2 + 14\sqrt{ex+d}\sqrt{bx+a}ab^2e^3x - 3\sqrt{ex+d}\sqrt{bx+a}a^2be^3 + 8\sqrt{ex+d}\sqrt{bx+a}ab^2de^2 + 14\sqrt{ex+d}\sqrt{bx+a}ab^2e^3x - 3\sqrt{ex+d}\sqrt{bx+a}a^2be^3}{e^4}$$

input

```
int((b*x+a)^(1/2)*(B*x+A)*(e*x+d)^(1/2),x)
```

output

```
(3*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b*e**3 + 8*sqrt(d + e*x)*sqrt(a + b*x)
*a*b**2*d*e**2 + 14*sqrt(d + e*x)*sqrt(a + b*x)*a*b**2*e**3*x - 3*sqrt(d +
e*x)*sqrt(a + b*x)*b**3*d**2*e + 2*sqrt(d + e*x)*sqrt(a + b*x)*b**3*d*e**
2*x + 8*sqrt(d + e*x)*sqrt(a + b*x)*b**3*e**3*x**2 - 3*sqrt(e)*sqrt(b)*log
((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**3*e**
3 + 9*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/
sqrt(a*e - b*d))*a**2*b*d*e**2 - 9*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b
*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*b**2*d**2*e + 3*sqrt(e)*sq
rt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))
*b**3*d**3)/(24*b**2*e**3)
```

3.175 $\int \frac{\sqrt{a+bx}(A+Bx)}{\sqrt{d+ex}} dx$

Optimal result	1600
Mathematica [A] (verified)	1600
Rubi [A] (verified)	1601
Maple [B] (verified)	1603
Fricas [A] (verification not implemented)	1603
Sympy [F]	1604
Maxima [F(-2)]	1604
Giac [A] (verification not implemented)	1605
Mupad [B] (verification not implemented)	1605
Reduce [B] (verification not implemented)	1606

Optimal result

Integrand size = 24, antiderivative size = 140

$$\int \frac{\sqrt{a+bx}(A+Bx)}{\sqrt{d+ex}} dx = -\frac{(3bBd - 4Abe + aBe)\sqrt{a+bx}\sqrt{d+ex}}{4be^2} + \frac{B(a+bx)^{3/2}\sqrt{d+ex}}{2be} + \frac{(bd - ae)(3bBd - 4Abe + aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{4b^{3/2}e^{5/2}}$$

```
output -1/4*(-4*A*b*e+B*a*e+3*B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(1/2)/b/e^2+1/2*B*(b*x+a)^(3/2)*(e*x+d)^(1/2)/b/e+1/4*(-a*e+b*d)*(-4*A*b*e+B*a*e+3*B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)^(1/2))/b^(3/2)/e^(5/2)
```

Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.84

$$\int \frac{\sqrt{a+bx}(A+Bx)}{\sqrt{d+ex}} dx = \frac{b\sqrt{a+bx}\sqrt{d+ex}(aBe + b(-3Bd + 4Ae + 2Bex)) + \sqrt{\frac{b}{e}}(bd - ae)(-3bBd + 4Abe - aBe) \log\left(\sqrt{a+bx}\sqrt{d+ex}\right)}{4b^2e^2}$$

input `Integrate[(Sqrt[a + b*x]*(A + B*x))/Sqrt[d + e*x],x]`

output `(b*Sqrt[a + b*x]*Sqrt[d + e*x]*(a*B*e + b*(-3*B*d + 4*A*e + 2*B*e*x)) + Sqrt[b/e]*(b*d - a*e)*(-3*b*B*d + 4*A*b*e - a*B*e)*Log[Sqrt[a + b*x] - Sqrt[b/e]*Sqrt[d + e*x]])/(4*b^2*e^2)`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.91, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {90, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{a+bx}(A+Bx)}{\sqrt{d+ex}} dx \\
 & \quad \downarrow \text{90} \\
 & \frac{B(a+bx)^{3/2}\sqrt{d+ex}}{2be} - \frac{(aBe - 4Abe + 3bBd) \int \frac{\sqrt{a+bx}}{\sqrt{d+ex}} dx}{4be} \\
 & \quad \downarrow \text{60} \\
 & \frac{B(a+bx)^{3/2}\sqrt{d+ex}}{2be} - \frac{(aBe - 4Abe + 3bBd) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{2e} \right)}{4be} \\
 & \quad \downarrow \text{66} \\
 & \frac{B(a+bx)^{3/2}\sqrt{d+ex}}{2be} - \frac{(aBe - 4Abe + 3bBd) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{e} \right)}{4be} \\
 & \quad \downarrow \text{221} \\
 & \frac{B(a+bx)^{3/2}\sqrt{d+ex}}{2be} - \frac{(aBe - 4Abe + 3bBd) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{\sqrt{be}^{3/2}} \right)}{4be}
 \end{aligned}$$

input `Int[(Sqrt[a + b*x]*(A + B*x))/Sqrt[d + e*x],x]`

output `(B*(a + b*x)^(3/2)*Sqrt[d + e*x])/(2*b*e) - ((3*b*B*d - 4*A*b*e + a*B*e)*
(Sqrt[a + b*x]*Sqrt[d + e*x])/e - ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b
*x])/(Sqrt[b]*Sqrt[d + e*x])])/(Sqrt[b]*e^(3/2)))/(4*b*e)`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(
b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!Integer
Q[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinear
Q[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[
2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; Fre
eQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
.), x] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))),
x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p
+ 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n,
p}, x] && NeQ[n + p + 2, 0]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 375 vs. 2(114) = 228.

Time = 0.26 (sec) , antiderivative size = 376, normalized size of antiderivative = 2.69

method	result
default	$\frac{\sqrt{bx+a}\sqrt{ex+d}\left(4A\ln\left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}}\right)abe^2-4A\ln\left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}}\right)b^2de+4B\sqrt{(ex+d)(bx+a)}\right)}{\dots}$

input

```
int((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
1/8*(b*x+a)^(1/2)*(e*x+d)^(1/2)*(4*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b*e^2-4*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^2*d*e+4*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b*e*x-B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*e^2-2*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b*d*e+3*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^2*d^2+8*A*(b*e)^(1/2)*((e*x+d)*(b*x+a))^(1/2)*b*e+2*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*e-6*B*(b*e)^(1/2)*((e*x+d)*(b*x+a))^(1/2)*b*d)/((e*x+d)*(b*x+a))^(1/2)/e^2/b/(b*e)^(1/2)
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 364, normalized size of antiderivative = 2.60

$$\int \frac{\sqrt{a+bx}(A+Bx)}{\sqrt{d+ex}} dx = \frac{(3Bb^2d^2 - 2(Bab + 2Ab^2)de - (Ba^2 - 4Aab)e^2)\sqrt{be} \log\left(8b^2e^2x^2 + b^2d^2 + 6abde + a^2e^2 + 4(2bex + \dots)\right)}{8b^2e^3} - \frac{(3Bb^2d^2 - 2(Bab + 2Ab^2)de - (Ba^2 - 4Aab)e^2)\sqrt{-be} \arctan\left(\frac{(2bex+bd+ae)\sqrt{-be}\sqrt{bx+a}\sqrt{ex+d}}{2(b^2e^2x^2+abde+(b^2de+abe^2)x)}\right) - 2(2E \dots)}{8b^2e^3}$$

input

```
integrate((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(1/2),x, algorithm="fricas")
```


output

```
[1/16*((3*B*b^2*d^2 - 2*(B*a*b + 2*A*b^2)*d*e - (B*a^2 - 4*A*a*b)*e^2)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*b*e*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x + a)*sqrt(e*x + d) + 8*(b^2*d*e + a*b*e^2)*x) + 4*(2*B*b^2*e^2*x - 3*B*b^2*d*e + (B*a*b + 4*A*b^2)*e^2)*sqrt(b*x + a)*sqrt(e*x + d))/(b^2*e^3), -1/8*((3*B*b^2*d^2 - 2*(B*a*b + 2*A*b^2)*d*e - (B*a^2 - 4*A*a*b)*e^2)*sqrt(-b*e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*e^2*x^2 + a*b*d*e + (b^2*d*e + a*b*e^2)*x)) - 2*(2*B*b^2*e^2*x - 3*B*b^2*d*e + (B*a*b + 4*A*b^2)*e^2)*sqrt(b*x + a)*sqrt(e*x + d))/(b^2*e^3)]
```

Sympy [F]

$$\int \frac{\sqrt{a+bx}(A+Bx)}{\sqrt{d+ex}} dx = \int \frac{(A+Bx)\sqrt{a+bx}}{\sqrt{d+ex}} dx$$

input

```
integrate((b*x+a)**(1/2)*(B*x+A)/(e*x+d)**(1/2),x)
```

output

```
Integral((A + B*x)*sqrt(a + b*x)/sqrt(d + e*x), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a+bx}(A+Bx)}{\sqrt{d+ex}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(1/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e
```

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.29

$$\int \frac{\sqrt{a+bx}(A+Bx)}{\sqrt{d+ex}} dx$$

$$= \frac{\left(\sqrt{b^2d+(bx+a)be} - abe\sqrt{bx+a}\left(\frac{2(bx+a)B}{b^2e} - \frac{3Bb^3de+Bab^2e^2-4Ab^3e^2}{b^4e^3}\right) - \frac{(3Bb^2d^2-2Babde-4Ab^2de-Ba^2e^2+4Aa^2e)}{b^4e^3}\right)}{4|b|}$$

input `integrate((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(1/2),x, algorithm="giac")`

output `1/4*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*sqrt(b*x + a)*(2*(b*x + a)*B/(b^2*e) - (3*B*b^3*d*e + B*a*b^2*e^2 - 4*A*b^3*e^2)/(b^4*e^3)) - (3*B*b^2*d^2 - 2*B*a*b*d*e - 4*A*b^2*d*e - B*a^2*e^2 + 4*A*a*b*e^2)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*b*e^2))*b/abs(b)`

Mupad [B] (verification not implemented)

Time = 14.50 (sec) , antiderivative size = 872, normalized size of antiderivative = 6.23

$$\int \frac{\sqrt{a+bx}(A+Bx)}{\sqrt{d+ex}} dx = \text{Too large to display}$$

input `int(((A + B*x)*(a + b*x)^(1/2))/(d + e*x)^(1/2),x)`

output

```

((((a + b*x)^(1/2) - a^(1/2))*((B*a^2*b^2*e^2)/2 - (3*B*b^4*d^2)/2 + B*a*b
^3*d*e))/(e^6*((d + e*x)^(1/2) - d^(1/2))) + (((a + b*x)^(1/2) - a^(1/2))^
3*((11*B*b^3*d^2)/2 + (7*B*a^2*b*e^2)/2 + 23*B*a*b^2*d*e))/(e^5*((d + e*x)
^(1/2) - d^(1/2))^3) + (((a + b*x)^(1/2) - a^(1/2))^5*((7*B*a^2*e^2)/2 + (
11*B*b^2*d^2)/2 + 23*B*a*b*d*e))/(e^4*((d + e*x)^(1/2) - d^(1/2))^5) + (((
a + b*x)^(1/2) - a^(1/2))^7*((B*a^2*e^2)/2 - (3*B*b^2*d^2)/2 + B*a*b*d*e))
/(b*e^3*((d + e*x)^(1/2) - d^(1/2))^7) - (a^(1/2)*d^(1/2)*((a + b*x)^(1/2)
- a^(1/2))^4*(32*B*b^2*d + 16*B*a*b*e))/(e^4*((d + e*x)^(1/2) - d^(1/2))^
4) - (8*B*a^(3/2)*d^(1/2)*((a + b*x)^(1/2) - a^(1/2))^6)/(e^2*((d + e*x)^(
1/2) - d^(1/2))^6) - (8*B*a^(3/2)*b^2*d^(1/2)*((a + b*x)^(1/2) - a^(1/2))^
2)/(e^4*((d + e*x)^(1/2) - d^(1/2))^2)/(((a + b*x)^(1/2) - a^(1/2))^8/((d
+ e*x)^(1/2) - d^(1/2))^8 + b^4/e^4 - (4*b^3*((a + b*x)^(1/2) - a^(1/2))^
2)/(e^3*((d + e*x)^(1/2) - d^(1/2))^2) + (6*b^2*((a + b*x)^(1/2) - a^(1/2)
)^4)/(e^2*((d + e*x)^(1/2) - d^(1/2))^4) - (4*b*((a + b*x)^(1/2) - a^(1/2)
)^6)/(e*((d + e*x)^(1/2) - d^(1/2))^6)) + (((a + b*x)^(1/2) - a^(1/2))*(2
*A*b^2*d + 2*A*a*b*e))/(e^3*((d + e*x)^(1/2) - d^(1/2))) + ((2*A*a*e + 2*A
*b*d)*((a + b*x)^(1/2) - a^(1/2))^3)/(e^2*((d + e*x)^(1/2) - d^(1/2))^3) -
(8*A*a^(1/2)*b*d^(1/2)*((a + b*x)^(1/2) - a^(1/2))^2)/(e^2*((d + e*x)^(1/
2) - d^(1/2))^2)/(((a + b*x)^(1/2) - a^(1/2))^4/((d + e*x)^(1/2) - d^(1/2
))^4 + b^2/e^2 - (2*b*((a + b*x)^(1/2) - a^(1/2))^2)/(e*((d + e*x)^(1/2...

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.41

$$\int \frac{\sqrt{a+bx}(A+Bx)}{\sqrt{d+ex}} dx$$

$$= \frac{5\sqrt{ex+d}\sqrt{bx+a}abe^2 - 3\sqrt{ex+d}\sqrt{bx+a}b^2de + 2\sqrt{ex+d}\sqrt{bx+a}b^2e^2x + 3\sqrt{e}\sqrt{b}\log\left(\frac{\sqrt{e}\sqrt{bx+a} + \sqrt{ae-b}}{\sqrt{ae-b}}\right)}{4be^3}$$

input

```
int((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(1/2),x)
```

output

```
(5*sqrt(d + e*x)*sqrt(a + b*x)*a*b*e**2 - 3*sqrt(d + e*x)*sqrt(a + b*x)*b*
*2*d*e + 2*sqrt(d + e*x)*sqrt(a + b*x)*b**2*e**2*x + 3*sqrt(e)*sqrt(b)*log
((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**2*e**
2 - 6*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/
sqrt(a*e - b*d))*a*b*d*e + 3*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) +
sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**2*d**2)/(4*b*e**3)
```

3.176 $\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{3/2}} dx$

Optimal result	1608
Mathematica [A] (verified)	1608
Rubi [A] (verified)	1609
Maple [B] (verified)	1611
Fricas [A] (verification not implemented)	1611
Sympy [F]	1612
Maxima [F(-2)]	1612
Giac [A] (verification not implemented)	1613
Mupad [F(-1)]	1613
Reduce [B] (verification not implemented)	1614

Optimal result

Integrand size = 24, antiderivative size = 113

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{3/2}} dx = \frac{2(Bd - Ae)\sqrt{a+bx}}{e^2\sqrt{d+ex}} + \frac{B\sqrt{a+bx}\sqrt{d+ex}}{e^2} - \frac{(3bBd - 2Abe - aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{\sqrt{b}e^{5/2}}$$

output

```
2*(-A*e+B*d)*(b*x+a)^(1/2)/e^2/(e*x+d)^(1/2)+B*(b*x+a)^(1/2)*(e*x+d)^(1/2)
/e^2-(-2*A*b*e-B*a*e+3*B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)
)^(1/2))/b^(1/2)/e^(5/2)
```

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.81

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{3/2}} dx = \frac{\sqrt{a+bx}(3Bd - 2Ae + Bex)}{e^2\sqrt{d+ex}} + \frac{(-3bBd + 2Abe + aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{\sqrt{b}e^{5/2}}$$

input `Integrate[(Sqrt[a + b*x]*(A + B*x))/(d + e*x)^(3/2),x]`

output `(Sqrt[a + b*x]*(3*B*d - 2*A*e + B*e*x))/(e^2*Sqrt[d + e*x]) + ((-3*b*B*d + 2*A*b*e + a*B*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(Sqrt[b]*e^(5/2))`

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.28, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {87, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{3/2}} dx \\
 & \quad \downarrow 87 \\
 & \frac{(-aBe - 2Abe + 3bBd) \int \frac{\sqrt{a+bx}}{\sqrt{d+ex}} dx}{e(bd - ae)} - \frac{2(a+bx)^{3/2}(Bd - Ae)}{e\sqrt{d+ex}(bd - ae)} \\
 & \quad \downarrow 60 \\
 & \frac{(-aBe - 2Abe + 3bBd) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{2e} \right)}{e(bd - ae)} - \frac{2(a+bx)^{3/2}(Bd - Ae)}{e\sqrt{d+ex}(bd - ae)} \\
 & \quad \downarrow 66 \\
 & \frac{(-aBe - 2Abe + 3bBd) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{e} \right)}{e(bd - ae)} - \frac{2(a+bx)^{3/2}(Bd - Ae)}{e\sqrt{d+ex}(bd - ae)} \\
 & \quad \downarrow 221 \\
 & \frac{(-aBe - 2Abe + 3bBd) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{\sqrt{be}^{3/2}} \right)}{e(bd - ae)} - \frac{2(a+bx)^{3/2}(Bd - Ae)}{e\sqrt{d+ex}(bd - ae)}
 \end{aligned}$$

input `Int[(Sqrt[a + b*x]*(A + B*x))/(d + e*x)^(3/2),x]`

output `(-2*(B*d - A*e)*(a + b*x)^(3/2))/(e*(b*d - a*e)*Sqrt[d + e*x]) + ((3*b*B*d - 2*A*b*e - a*B*e)*((Sqrt[a + b*x]*Sqrt[d + e*x])/e - ((b*d - a*e)*ArcTan[h[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])]/(Sqrt[b]*e^(3/2))))/(e*(b*d - a*e))`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || (EqQ[e, 0] || (EqQ[c, 0] || LtQ[p, n]))))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 385 vs. $2(93) = 186$.

Time = 0.26 (sec) , antiderivative size = 386, normalized size of antiderivative = 3.42

method	result
default	$\frac{\sqrt{bx+a} \left(2A \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) b e^2 x + B \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) a e^2 x - 3B \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) \right)}{\dots}$

input `int((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(3/2),x,method=_RETURNVERBOSE)`

output

```
1/2*(b*x+a)^(1/2)*(2*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b*e^2*x+B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*e^2*x-3*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b*d*e*x+2*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b*d*e+B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*d*e-3*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b*d^2+2*B*e*x*(b*e)^(1/2)*((e*x+d)*(b*x+a))^(1/2)-4*A*e*(b*e)^(1/2)*((e*x+d)*(b*x+a))^(1/2)+6*B*d*(b*e)^(1/2)*((e*x+d)*(b*x+a))^(1/2))/(b*e)^(1/2)/((e*x+d)*(b*x+a))^(1/2)/e^2/(e*x+d)^(1/2)
```

Fricas [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 358, normalized size of antiderivative = 3.17

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{3/2}} dx = \left[-\frac{(3Bbd^2 - (Ba + 2Ab)de + (3Bbde - (Ba + 2Ab)e^2)x)\sqrt{be} \log(8b^2e^2x^2 + \dots)}{\dots} \right]$$

input `integrate((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(3/2),x, algorithm="fricas")`

output

```
[-1/4*((3*B*b*d^2 - (B*a + 2*A*b)*d*e + (3*B*b*d*e - (B*a + 2*A*b)*e^2)*x)
*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*b*e*x
+ b*d + a*e)*sqrt(b*e)*sqrt(b*x + a)*sqrt(e*x + d) + 8*(b^2*d*e + a*b*e^2)
*x) - 4*(B*b*e^2*x + 3*B*b*d*e - 2*A*b*e^2)*sqrt(b*x + a)*sqrt(e*x + d))/(
b*e^4*x + b*d*e^3), 1/2*((3*B*b*d^2 - (B*a + 2*A*b)*d*e + (3*B*b*d*e - (B*
a + 2*A*b)*e^2)*x)*sqrt(-b*e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(-b*e)*
sqrt(b*x + a)*sqrt(e*x + d)/(b^2*e^2*x^2 + a*b*d*e + (b^2*d*e + a*b*e^2)*x
)) + 2*(B*b*e^2*x + 3*B*b*d*e - 2*A*b*e^2)*sqrt(b*x + a)*sqrt(e*x + d)/(b
*e^4*x + b*d*e^3)]
```

Sympy [F]

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{3/2}} dx = \int \frac{(A+Bx)\sqrt{a+bx}}{(d+ex)^{\frac{3}{2}}} dx$$

input

```
integrate((b*x+a)**(1/2)*(B*x+A)/(e*x+d)**(3/2),x)
```

output

```
Integral((A + B*x)*sqrt(a + b*x)/(d + e*x)**(3/2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{3/2}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(3/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f
or more de
```

Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.39

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{3/2}} dx = \frac{\sqrt{bx+a} \left(\frac{(bx+a)B|b|}{be} + \frac{3Bb^2de|b|-Babe^2|b|-2Ab^2e^2|b|}{b^2e^3} \right)}{\sqrt{b^2d+(bx+a)be-abe}} + \frac{(3Bbd|b|-Bae|b|-2Abe|b|) \log \left(\left| -\sqrt{be}\sqrt{bx+a} + \sqrt{b^2d+(bx+a)be-abe} \right| \right)}{\sqrt{bebe^2}}$$

input `integrate((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(3/2),x, algorithm="giac")`

output `sqrt(b*x + a)*((b*x + a)*B*abs(b)/(b*e) + (3*B*b^2*d*e*abs(b) - B*a*b*e^2*abs(b) - 2*A*b^2*e^2*abs(b))/(b^2*e^3))/sqrt(b^2*d + (b*x + a)*b*e - a*b*e) + (3*B*b*d*abs(b) - B*a*e*abs(b) - 2*A*b*e*abs(b))*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*b*e^2)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{3/2}} dx = \int \frac{(A+Bx)\sqrt{a+bx}}{(d+ex)^{3/2}} dx$$

input `int(((A + B*x)*(a + b*x)^(1/2))/(d + e*x)^(3/2), x)`

output `int(((A + B*x)*(a + b*x)^(1/2))/(d + e*x)^(3/2), x)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 275, normalized size of antiderivative = 2.43

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{3/2}} dx = \frac{-8\sqrt{ex+d}\sqrt{bx+a}ae^2 + 12\sqrt{ex+d}\sqrt{bx+a}bde + 4\sqrt{ex+d}\sqrt{bx+a}be^2x}{(d+ex)^{3/2}}$$

input `int((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(3/2),x)`

output

```
( - 8*sqrt(d + e*x)*sqrt(a + b*x)*a*e**2 + 12*sqrt(d + e*x)*sqrt(a + b*x)*
b*d*e + 4*sqrt(d + e*x)*sqrt(a + b*x)*b*e**2*x + 12*sqrt(e)*sqrt(b)*log((s
qrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*d*e + 12*
sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a
*e - b*d))*a*e**2*x - 12*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt
(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b*d**2 - 12*sqrt(e)*sqrt(b)*log((sqrt(
e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b*d*e*x - 9*sqr
t(e)*sqrt(b)*a*d*e - 9*sqrt(e)*sqrt(b)*a*e**2*x + 9*sqrt(e)*sqrt(b)*b*d**2
+ 9*sqrt(e)*sqrt(b)*b*d*e*x)/(4*e**3*(d + e*x))
```

3.177 $\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{5/2}} dx$

Optimal result	1615
Mathematica [A] (verified)	1615
Rubi [A] (verified)	1616
Maple [B] (verified)	1618
Fricas [B] (verification not implemented)	1618
Sympy [F]	1619
Maxima [F(-2)]	1620
Giac [B] (verification not implemented)	1620
Mupad [F(-1)]	1621
Reduce [B] (verification not implemented)	1621

Optimal result

Integrand size = 24, antiderivative size = 111

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{5/2}} dx = -\frac{2(Bd - Ae)(a+bx)^{3/2}}{3e(bd - ae)(d+ex)^{3/2}} - \frac{2B\sqrt{a+bx}}{e^2\sqrt{d+ex}} + \frac{2\sqrt{b}\text{Barctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{e^{5/2}}$$

output

```
-2/3*(-A*e+B*d)*(b*x+a)^(3/2)/e/(-a*e+b*d)/(e*x+d)^(3/2)-2*B*(b*x+a)^(1/2)
/e^2/(e*x+d)^(1/2)+2*b^(1/2)*B*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+
d)^(1/2))/e^(5/2)
```

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.05

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{5/2}} dx = -\frac{2\sqrt{a+bx}(Abe^2x - bBd(3d + 4ex) + ae(2Bd + Ae + 3Bex))}{3e^2(-bd + ae)(d+ex)^{3/2}} + \frac{2\sqrt{b}\text{Barctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{e^{5/2}}$$

input `Integrate[(Sqrt[a + b*x]*(A + B*x))/(d + e*x)^(5/2),x]`

output `(-2*Sqrt[a + b*x]*(A*b*e^2*x - b*B*d*(3*d + 4*e*x) + a*e*(2*B*d + A*e + 3*B*e*x))/(3*e^2*(-(b*d) + a*e)*(d + e*x)^(3/2)) + (2*Sqrt[b]*B*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/e^(5/2)`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.04, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {87, 57, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{5/2}} dx \\
 & \quad \downarrow 87 \\
 & \frac{B \int \frac{\sqrt{a+bx}}{(d+ex)^{3/2}} dx}{e} - \frac{2(a+bx)^{3/2}(Bd-Ae)}{3e(d+ex)^{3/2}(bd-ae)} \\
 & \quad \downarrow 57 \\
 & \frac{B \left(\frac{b \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{e} - \frac{2\sqrt{a+bx}}{e\sqrt{d+ex}} \right)}{e} - \frac{2(a+bx)^{3/2}(Bd-Ae)}{3e(d+ex)^{3/2}(bd-ae)} \\
 & \quad \downarrow 66 \\
 & \frac{B \left(\frac{2b \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{e} - \frac{2\sqrt{a+bx}}{e\sqrt{d+ex}} \right)}{e} - \frac{2(a+bx)^{3/2}(Bd-Ae)}{3e(d+ex)^{3/2}(bd-ae)} \\
 & \quad \downarrow 221 \\
 & \frac{B \left(\frac{2\sqrt{b} \operatorname{arctanh} \left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}} \right)}{e^{3/2}} - \frac{2\sqrt{a+bx}}{e\sqrt{d+ex}} \right)}{e} - \frac{2(a+bx)^{3/2}(Bd-Ae)}{3e(d+ex)^{3/2}(bd-ae)}
 \end{aligned}$$

input `Int[(Sqrt[a + b*x]*(A + B*x))/(d + e*x)^(5/2),x]`

output `(-2*(B*d - A*e)*(a + b*x)^(3/2))/(3*e*(b*d - a*e)*(d + e*x)^(3/2)) + (B*((-2*Sqrt[a + b*x])/(e*Sqrt[d + e*x]) + (2*Sqrt[b]*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])]))/e^(3/2))/e`

Defintions of rubi rules used

rule 57 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))]
Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 502 vs. $2(89) = 178$.

Time = 0.27 (sec) , antiderivative size = 503, normalized size of antiderivative = 4.53

method	result
default	$-\left(-3B \ln\left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}}\right)abe^3x^2+3B \ln\left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}}\right)b^2de^2x^2-6B \ln\left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}}\right)\right)$

input

```
int((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(5/2),x,method=_RETURNVERBOSE)
```

output

```
-1/3*(-3*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/
(b*e)^(1/2))*a*b*e^3*x^2+3*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*
e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^2*d*e^2*x^2-6*B*ln(1/2*(2*b*e*x+2*((e*x+d
)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b*d*e^2*x+6*B*ln(1/2*
(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^2*d
^2*e*x+2*A*b*e^2*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-3*B*ln(1/2*(2*b*e*x
+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b*d^2*e+3*B
*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2
))*b^2*d^3+6*B*a*e^2*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-8*B*b*d*e*x*((e
*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+2*A*a*e^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(
1/2)+4*B*a*d*e*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-6*B*b*d^2*((e*x+d)*(b*x
+a))^(1/2)*(b*e)^(1/2))*(b*x+a)^(1/2)/(b*e)^(1/2)/(a*e-b*d)/((e*x+d)*(b*x+
a))^(1/2)/e^2/(e*x+d)^(3/2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 245 vs. $2(89) = 178$.

Time = 0.67 (sec) , antiderivative size = 515, normalized size of antiderivative = 4.64

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{5/2}} dx = \frac{3(Bbd^3 - Bad^2e + (Bbde^2 - Bae^3)x^2 + 2(Bbd^2e - Bade^2)x)\sqrt{\frac{b}{e}} \log\left(8b^2e^2\right) + 3(Bbd^3 - Bad^2e + (Bbde^2 - Bae^3)x^2 + 2(Bbd^2e - Bade^2)x)\sqrt{-\frac{b}{e}} \arctan\left(\frac{(2bex+bd+ae)\sqrt{bx+a}\sqrt{ex+d}\sqrt{-\frac{b}{e}}}{2(b^2ex^2+abd+(b^2d+abe)x)}\right)}{3(bd^3e^2 - ad^2e^3 + (bde^4 - ae^5)x^2 + 2(bd^2e^3 - ade^4)x)}$$

input `integrate((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(5/2),x, algorithm="fricas")`

output `[1/6*(3*(B*b*d^3 - B*a*d^2*e + (B*b*d*e^2 - B*a*e^3)*x^2 + 2*(B*b*d^2*e - B*a*d*e^2)*x)*sqrt(b/e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*b*e^2*x + b*d*e + a*e^2)*sqrt(b*x + a)*sqrt(e*x + d)*sqrt(b/e) + 8*(b^2*d*e + a*b*e^2)*x) - 4*(3*B*b*d^2 - 2*B*a*d*e - A*a*e^2 + (4*B*b*d*e - (3*B*a + A*b)*e^2)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b*d^3*e^2 - a*d^2*e^3 + (b*d*e^4 - a*e^5)*x^2 + 2*(b*d^2*e^3 - a*d*e^4)*x), -1/3*(3*(B*b*d^3 - B*a*d^2*e + (B*b*d*e^2 - B*a*e^3)*x^2 + 2*(B*b*d^2*e - B*a*d*e^2)*x)*sqrt(-b/e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(b*x + a)*sqrt(e*x + d)*sqrt(-b/e)/(b^2*e*x^2 + a*b*d + (b^2*d + a*b*e)*x)) + 2*(3*B*b*d^2 - 2*B*a*d*e - A*a*e^2 + (4*B*b*d*e - (3*B*a + A*b)*e^2)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b*d^3*e^2 - a*d^2*e^3 + (b*d*e^4 - a*e^5)*x^2 + 2*(b*d^2*e^3 - a*d*e^4)*x]`

Sympy [F]

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{5/2}} dx = \int \frac{(A+Bx)\sqrt{a+bx}}{(d+ex)^{5/2}} dx$$

input `integrate((b*x+a)**(1/2)*(B*x+A)/(e*x+d)**(5/2),x)`

output `Integral((A + B*x)*sqrt(a + b*x)/(d + e*x)**(5/2), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(5/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f or more de

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 205 vs. 2(89) = 178.

Time = 0.18 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.85

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{5/2}} dx = \frac{2B|b| \log \left(\left| -\sqrt{be}\sqrt{bx+a} + \sqrt{b^2d+(bx+a)be-abe} \right| \right)}{\sqrt{bee^2}} \frac{2\sqrt{bx+a} \left(\frac{(4Bb^4de^2|b|-3Bab^3e^3|b|-Ab^4e^3|b|)(bx+a)}{b^3de^3-ab^2e^4} + \frac{3(Bb^5d^2e|b|-2Bab^4de^2|b|+Ba^2b^3e^3|b|)}{b^3de^3-ab^2e^4} \right)}{3(b^2d+(bx+a)be-abe)^{\frac{3}{2}}}$$

input `integrate((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(5/2),x, algorithm="giac")`

output `-2*B*abs(b)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*e^2) - 2/3*sqrt(b*x + a)*((4*B*b^4*d*e^2*abs(b) - 3*B*a*b^3*e^3*abs(b) - A*b^4*e^3*abs(b))*(b*x + a)/(b^3*d*e^3 - a*b^2*e^4) + 3*(B*b^5*d^2*e*abs(b) - 2*B*a*b^4*d*e^2*abs(b) + B*a^2*b^3*e^3*abs(b))/(b^3*d*e^3 - a*b^2*e^4))/(b^2*d + (b*x + a)*b*e - a*b*e)^(3/2)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{5/2}} dx = \int \frac{(A+Bx)\sqrt{a+bx}}{(d+ex)^{5/2}} dx$$

input `int(((A + B*x)*(a + b*x)^(1/2))/(d + e*x)^(5/2), x)`

output `int(((A + B*x)*(a + b*x)^(1/2))/(d + e*x)^(5/2), x)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.86

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{5/2}} dx = \frac{-\frac{2\sqrt{ex+d}\sqrt{bx+ae^2}}{3} - 2\sqrt{ex+d}\sqrt{bx+a}bde - \frac{8\sqrt{ex+d}\sqrt{bx+ae^2}x}{3} + 2\sqrt{e}\sqrt{b}\log\left(\frac{\sqrt{e}\sqrt{a+bx} + \sqrt{b}\sqrt{d+ex}}{\sqrt{ae-bd}}\right)}{(d+ex)^{5/2}}$$

input `int((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(5/2), x)`

output `(2*(-sqrt(d + e*x)*sqrt(a + b*x)*a*e**2 - 3*sqrt(d + e*x)*sqrt(a + b*x)*b*d*e - 4*sqrt(d + e*x)*sqrt(a + b*x)*b*e**2*x + 3*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b*d**2 + 6*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b*d*e*x + 3*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b*e**2*x**2))/(3*e**3*(d**2 + 2*d*e*x + e**2*x**2))`

3.178 $\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{7/2}} dx$

Optimal result	1622
Mathematica [A] (verified)	1622
Rubi [A] (verified)	1623
Maple [A] (verified)	1624
Fricas [B] (verification not implemented)	1624
Sympy [F]	1625
Maxima [F(-2)]	1625
Giac [B] (verification not implemented)	1626
Mupad [B] (verification not implemented)	1626
Reduce [B] (verification not implemented)	1627

Optimal result

Integrand size = 24, antiderivative size = 95

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{7/2}} dx = -\frac{2(Bd - Ae)(a+bx)^{3/2}}{5e(bd - ae)(d+ex)^{5/2}} + \frac{2(3bBd + 2Abe - 5aBe)(a+bx)^{3/2}}{15e(bd - ae)^2(d+ex)^{3/2}}$$

output `-2/5*(-A*e+B*d)*(b*x+a)^(3/2)/e/(-a*e+b*d)/(e*x+d)^(5/2)+2/15*(2*A*b*e-5*B*a*e+3*B*b*d)*(b*x+a)^(3/2)/e/(-a*e+b*d)^2/(e*x+d)^(3/2)`

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.69

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{7/2}} dx = \frac{2(a+bx)^{3/2}(B(-2ad + 3bdx - 5aex) + A(5bd - 3ae + 2bex))}{15(bd - ae)^2(d+ex)^{5/2}}$$

input `Integrate[(Sqrt[a + b*x]*(A + B*x))/(d + e*x)^(7/2), x]`

output `(2*(a + b*x)^(3/2)*(B*(-2*a*d + 3*b*d*x - 5*a*e*x) + A*(5*b*d - 3*a*e + 2*b*e*x)))/(15*(b*d - a*e)^2*(d + e*x)^(5/2))`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {87, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{7/2}} dx$$

↓ 87

$$\frac{(-5aBe + 2Abe + 3bBd) \int \frac{\sqrt{a+bx}}{(d+ex)^{5/2}} dx}{5e(bd - ae)} - \frac{2(a+bx)^{3/2}(Bd - Ae)}{5e(d+ex)^{5/2}(bd - ae)}$$

↓ 48

$$\frac{2(a+bx)^{3/2}(-5aBe + 2Abe + 3bBd)}{15e(d+ex)^{3/2}(bd - ae)^2} - \frac{2(a+bx)^{3/2}(Bd - Ae)}{5e(d+ex)^{5/2}(bd - ae)}$$

input `Int[(Sqrt[a + b*x]*(A + B*x))/(d + e*x)^(7/2),x]`

output `(-2*(B*d - A*e)*(a + b*x)^(3/2))/(5*e*(b*d - a*e)*(d + e*x)^(5/2)) + (2*(3*b*B*d + 2*A*b*e - 5*a*B*e)*(a + b*x)^(3/2))/(15*e*(b*d - a*e)^2*(d + e*x)^(3/2))`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 87

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))
```

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.64

method	result	size
default	$-\frac{2(bx+a)^{\frac{3}{2}}(-2Abe x+5Baex-3Bbdx+3Aae-5Abd+2Bad)}{15(ex+d)^{\frac{5}{2}}(ae-db)^2}$	61
gospers	$-\frac{2(bx+a)^{\frac{3}{2}}(-2Abe x+5Baex-3Bbdx+3Aae-5Abd+2Bad)}{15(ex+d)^{\frac{5}{2}}(a^2e^2-2abde+b^2d^2)}$	74
orering	$-\frac{2(bx+a)^{\frac{3}{2}}(-2Abe x+5Baex-3Bbdx+3Aae-5Abd+2Bad)}{15(ex+d)^{\frac{5}{2}}(a^2e^2-2abde+b^2d^2)}$	74

input

```
int((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(7/2),x,method=_RETURNVERBOSE)
```

output

```
-2/15*(b*x+a)^(3/2)/(e*x+d)^(5/2)*(-2*A*b*e*x+5*B*a*e*x-3*B*b*d*x+3*A*a*e-5*A*b*d+2*B*a*d)/(a*e-b*d)^2
```

Fricas [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 221 vs. $2(83) = 166$.

Time = 1.88 (sec) , antiderivative size = 221, normalized size of antiderivative = 2.33

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{7/2}} dx =$$

$$\frac{2(3Aa^2e - (3Bb^2d - (5Bab - 2Ab^2)e)x^2 + (2Ba^2 - 5Aab)d - ((Bab + 5Ab^2)d - (5Ba^2 + Aab)e)}{15(b^2d^5 - 2abd^4e + a^2d^3e^2 + (b^2d^2e^3 - 2abde^4 + a^2e^5)x^3 + 3(b^2d^3e^2 - 2abd^2e^3 + a^2de^4)x^2 + 3(b^2d^4e -$$

input

```
integrate((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(7/2),x, algorithm="fricas")
```

output

```
-2/15*(3*A*a^2*e - (3*B*b^2*d - (5*B*a*b - 2*A*b^2)*e)*x^2 + (2*B*a^2 - 5*
A*a*b)*d - ((B*a*b + 5*A*b^2)*d - (5*B*a^2 + A*a*b)*e)*x)*sqrt(b*x + a)*sq
rt(e*x + d)/(b^2*d^5 - 2*a*b*d^4*e + a^2*d^3*e^2 + (b^2*d^2*e^3 - 2*a*b*d*
e^4 + a^2*e^5)*x^3 + 3*(b^2*d^3*e^2 - 2*a*b*d^2*e^3 + a^2*d*e^4)*x^2 + 3*(
b^2*d^4*e - 2*a*b*d^3*e^2 + a^2*d^2*e^3)*x)
```

Sympy [F]

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{7/2}} dx = \int \frac{(A+Bx)\sqrt{a+bx}}{(d+ex)^{7/2}} dx$$

input

```
integrate((b*x+a)**(1/2)*(B*x+A)/(e*x+d)**(7/2),x)
```

output

```
Integral((A + B*x)*sqrt(a + b*x)/(d + e*x)**(7/2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{7/2}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(7/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f
or more de
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 191 vs. $2(83) = 166$.

Time = 0.22 (sec) , antiderivative size = 191, normalized size of antiderivative = 2.01

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{7/2}} dx = \frac{2(bx+a)^{\frac{3}{2}} \left(\frac{(3Bb^6de^2|b|-5Bab^5e^3|b|+2Ab^6e^3|b|)(bx+a)}{b^4d^2e^2-2ab^3de^3+a^2b^2e^4} - \frac{5(Bab^6de^2|b|-Ab^7de^2|b|-Ba^2b^5e^3|b|+A^2b^6e^3|b|)}{b^4d^2e^2-2ab^3de^3+a^2b^2e^4} \right)}{15(b^2d+(bx+a)be-abe)^{\frac{5}{2}}}$$

input `integrate((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(7/2),x, algorithm="giac")`

output
$$\frac{2/15*(b*x + a)^{(3/2)*((3*B*b^6*d*e^2*abs(b) - 5*B*a*b^5*e^3*abs(b) + 2*A*b^6*e^3*abs(b))*(b*x + a)/(b^4*d^2*e^2 - 2*a*b^3*d*e^3 + a^2*b^2*e^4) - 5*(B*a*b^6*d*e^2*abs(b) - A*b^7*d*e^2*abs(b) - B*a^2*b^5*e^3*abs(b) + A*a*b^6*e^3*abs(b))/(b^4*d^2*e^2 - 2*a*b^3*d*e^3 + a^2*b^2*e^4))/(b^2*d + (b*x + a)*b*e - a*b*e)^{(5/2)}$$

Mupad [B] (verification not implemented)

Time = 1.36 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.88

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{7/2}} dx = \frac{\sqrt{d+ex} \left(\frac{x^2\sqrt{a+bx}(4Ab^2e+6Bb^2d-10Babe)}{15e^3(ae-bd)^2} - \frac{\sqrt{a+bx}(6Aa^2e+4Ba^2d-10Aabd)}{15e^3(ae-bd)^2} \right)}{x^3 + \frac{d^3}{e^3} + \frac{3dx^2}{e} + \frac{3d^2x}{e^2}} + x\sqrt{\frac{a+bx}{d+ex}}$$

input `int(((A + B*x)*(a + b*x)^(1/2))/(d + e*x)^(7/2),x)`

output
$$\frac{((d + e*x)^{(1/2))*((x^2*(a + b*x)^{(1/2))*(4*A*b^2*e + 6*B*b^2*d - 10*B*a*b*e))/(15*e^3*(a*e - b*d)^2) - ((a + b*x)^{(1/2))*(6*A*a^2*e + 4*B*a^2*d - 10*A*a*b*d))/(15*e^3*(a*e - b*d)^2) + (x*(a + b*x)^{(1/2))*(10*A*b^2*d - 10*B*a^2*e - 2*A*a*b*e + 2*B*a*b*d))/(15*e^3*(a*e - b*d)^2)}}{(x^3 + d^3/e^3 + (3*d*x^2)/e + (3*d^2*x)/e^2)}$$

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 199, normalized size of antiderivative = 2.09

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{7/2}} dx = \frac{-\frac{2\sqrt{ex+d}\sqrt{bx+a}a^2e^3}{5} - \frac{4\sqrt{ex+d}\sqrt{bx+a}abe^3x}{5} - \frac{2\sqrt{ex+d}\sqrt{bx+a}b^2e^3x^2}{5} - \frac{2\sqrt{e}\sqrt{b}b^2d^3}{5} - \frac{6\sqrt{e}\sqrt{b}bd^2x}{5}}{e^3(ae^4x^3 - bde^3x^3 + 3ade^3x^2 - 3bd^2e^2x^2 + 3ad^2e^2x - 3bd^2)}$$

input `int((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(7/2),x)`

output

```
(2*( - sqrt(d + e*x)*sqrt(a + b*x)*a**2*e**3 - 2*sqrt(d + e*x)*sqrt(a + b*x)*a*b*e**3*x - sqrt(d + e*x)*sqrt(a + b*x)*b**2*e**3*x**2 - sqrt(e)*sqrt(b)*b**2*d**3 - 3*sqrt(e)*sqrt(b)*b**2*d**2*e*x - 3*sqrt(e)*sqrt(b)*b**2*d**e**2*x**2 - sqrt(e)*sqrt(b)*b**2*e**3*x**3))/(5*e**3*(a*d**3*e + 3*a*d**2*e**2*x + 3*a*d*e**3*x**2 + a*e**4*x**3 - b*d**4 - 3*b*d**3*e*x - 3*b*d**2*e**2*x**2 - b*d*e**3*x**3))
```


3.179 $\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{9/2}} dx$

Optimal result	1628
Mathematica [A] (verified)	1628
Rubi [A] (verified)	1629
Maple [A] (verified)	1630
Fricas [B] (verification not implemented)	1631
Sympy [F]	1631
Maxima [F(-2)]	1632
Giac [B] (verification not implemented)	1632
Mupad [B] (verification not implemented)	1633
Reduce [B] (verification not implemented)	1633

Optimal result

Integrand size = 24, antiderivative size = 147

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{9/2}} dx = -\frac{2(Bd - Ae)(a+bx)^{3/2}}{7e(bd - ae)(d+ex)^{7/2}} + \frac{2(3bBd + 4Abe - 7aBe)(a+bx)^{3/2}}{35e(bd - ae)^2(d+ex)^{5/2}} + \frac{4b(3bBd + 4Abe - 7aBe)(a+bx)^{3/2}}{105e(bd - ae)^3(d+ex)^{3/2}}$$

output `-2/7*(-A*e+B*d)*(b*x+a)^(3/2)/e/(-a*e+b*d)/(e*x+d)^(7/2)+2/35*(4*A*b*e-7*B*a*e+3*B*b*d)*(b*x+a)^(3/2)/e/(-a*e+b*d)^2/(e*x+d)^(5/2)+4/105*b*(4*A*b*e-7*B*a*e+3*B*b*d)*(b*x+a)^(3/2)/e/(-a*e+b*d)^3/(e*x+d)^(3/2)`

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.91

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{9/2}} dx = \frac{2(a+bx)^{3/2} \left(35Ab^2 - 35abB - \frac{15Bde(a+bx)^2}{(d+ex)^2} + \frac{15Ae^2(a+bx)^2}{(d+ex)^2} + \frac{21bBd(a+bx)}{d+ex} - \frac{42Abd}{d} \right)}{105(bd - ae)^3(d+ex)^{3/2}}$$

input `Integrate[(Sqrt[a + b*x]*(A + B*x))/(d + e*x)^(9/2),x]`

output

$$(2*(a + b*x)^{(3/2)}*(35*A*b^2 - 35*a*b*B - (15*B*d*e*(a + b*x)^2)/(d + e*x)^2 + (15*A*e^2*(a + b*x)^2)/(d + e*x)^2 + (21*b*B*d*(a + b*x))/(d + e*x) - (42*A*b*e*(a + b*x))/(d + e*x) + (21*a*B*e*(a + b*x))/(d + e*x))/(105*(b*d - a*e)^3*(d + e*x)^{(3/2)})$$

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.97, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {87, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{9/2}} dx$$

↓ 87

$$\frac{(-7aBe + 4Abe + 3bBd) \int \frac{\sqrt{a+bx}}{(d+ex)^{7/2}} dx}{7e(bd - ae)} - \frac{2(a+bx)^{3/2}(Bd - Ae)}{7e(d+ex)^{7/2}(bd - ae)}$$

↓ 55

$$\frac{(-7aBe + 4Abe + 3bBd) \left(\frac{2b \int \frac{\sqrt{a+bx}}{(d+ex)^{5/2}} dx}{5(bd - ae)} + \frac{2(a+bx)^{3/2}}{5(d+ex)^{5/2}(bd - ae)} \right)}{7e(bd - ae)} - \frac{2(a+bx)^{3/2}(Bd - Ae)}{7e(d+ex)^{7/2}(bd - ae)}$$

↓ 48

$$\left(\frac{4b(a+bx)^{3/2}}{15(d+ex)^{3/2}(bd - ae)^2} + \frac{2(a+bx)^{3/2}}{5(d+ex)^{5/2}(bd - ae)} \right) \frac{(-7aBe + 4Abe + 3bBd)}{7e(bd - ae)} - \frac{2(a+bx)^{3/2}(Bd - Ae)}{7e(d+ex)^{7/2}(bd - ae)}$$

input

```
Int[(Sqrt[a + b*x]*(A + B*x))/(d + e*x)^(9/2), x]
```

output

$$(-2*(B*d - A*e)*(a + b*x)^{(3/2)})/(7*e*(b*d - a*e)*(d + e*x)^{(7/2)}) + ((3*b*B*d + 4*A*b*e - 7*a*B*e)*((2*(a + b*x)^{(3/2)})/(5*(b*d - a*e)*(d + e*x)^{(5/2)})) + (4*b*(a + b*x)^{(3/2)})/(15*(b*d - a*e)^2*(d + e*x)^{(3/2)}))/ (7*e*(b*d - a*e))$$

Defintions of rubi rules used

```
rule 48 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 55 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
lerQ[n, 1])
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && ( !LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.01

method	result
default	$-\frac{2(bx+a)^{\frac{3}{2}}(8Ab^2e^2x^2-14Babe^2x^2+6Bb^2dex^2-12Aabe^2x+28Ab^2dex+21Ba^2e^2x-58Babdex+21b^2Bd^2x+15a^2Ae^2-42Aab)}{105(ex+d)^{\frac{7}{2}}(ae-db)^3}$
gospers	$-\frac{2(bx+a)^{\frac{3}{2}}(8Ab^2e^2x^2-14Babe^2x^2+6Bb^2dex^2-12Aabe^2x+28Ab^2dex+21Ba^2e^2x-58Babdex+21b^2Bd^2x+15a^2Ae^2-42Aab)}{105(ex+d)^{\frac{7}{2}}(a^3e^3-3a^2be^2d+3ab^2d^2e-b^3d^3)}$
orering	$-\frac{2(bx+a)^{\frac{3}{2}}(8Ab^2e^2x^2-14Babe^2x^2+6Bb^2dex^2-12Aabe^2x+28Ab^2dex+21Ba^2e^2x-58Babdex+21b^2Bd^2x+15a^2Ae^2-42Aab)}{105(ex+d)^{\frac{7}{2}}(a^3e^3-3a^2be^2d+3ab^2d^2e-b^3d^3)}$

```
input int((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(9/2), x, method=_RETURNVERBOSE)
```


Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{9/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(9/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f or more de`

Giac [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 379 vs. $2(129) = 258$.

Time = 0.26 (sec) , antiderivative size = 379, normalized size of antiderivative = 2.58

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{9/2}} dx = \frac{2 \left((bx+a) \left(\frac{2(3Bb^8de^4|b|-7Bab^7e^5|b|+4Ab^8e^5|b|)(bx+a)}{b^5d^3e^3-3ab^4d^2e^4+3a^2b^3de^5-a^3b^2e^6} + \frac{7(3Bb^9d^2e^3|b|-10Bab^8de^4|b|+4Ab^9d^2e^3|b|-10Bab^8de^4|b|+4Ab^9d^2e^3|b|)}{b^5d^3e^3-3ab^4d^2e^4+3a^2b^3de^5-a^3b^2e^6} \right) \right)}{(d+ex)^{9/2}}$$

10

input `integrate((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(9/2),x, algorithm="giac")`

output `2/105*((b*x + a)*(2*(3*B*b^8*d*e^4*abs(b) - 7*B*a*b^7*e^5*abs(b) + 4*A*b^8*e^5*abs(b))*(b*x + a)/(b^5*d^3*e^3 - 3*a*b^4*d^2*e^4 + 3*a^2*b^3*d*e^5 - a^3*b^2*e^6) + 7*(3*B*b^9*d^2*e^3*abs(b) - 10*B*a*b^8*d*e^4*abs(b) + 4*A*b^9*d*e^4*abs(b) + 7*B*a^2*b^7*e^5*abs(b) - 4*A*a*b^8*e^5*abs(b))/(b^5*d^3*e^3 - 3*a*b^4*d^2*e^4 + 3*a^2*b^3*d*e^5 - a^3*b^2*e^6)) - 35*(B*a*b^9*d^2*e^3*abs(b) - A*b^10*d^2*e^3*abs(b) - 2*B*a^2*b^8*d*e^4*abs(b) + 2*A*a*b^9*d*e^4*abs(b) + B*a^3*b^7*e^5*abs(b) - A*a^2*b^8*e^5*abs(b))/(b^5*d^3*e^3 - 3*a*b^4*d^2*e^4 + 3*a^2*b^3*d*e^5 - a^3*b^2*e^6))*(b*x + a)^(3/2)/(b^2*d + (b*x + a)*b*e - a*b*e)^(7/2)`

Mupad [B] (verification not implemented)

Time = 1.52 (sec) , antiderivative size = 295, normalized size of antiderivative = 2.01

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{9/2}} dx = \frac{\sqrt{d+ex} \left(\frac{\sqrt{a+bx}(12Ba^3de+30Aa^3e^2-28Ba^2bd^2-84Aa^2bde+70Aab^2d^2)}{105e^4(ae-bd)^3} + \frac{x\sqrt{a+bx}(42Ba^3e^2-104Ba^2bde+6Aa^2be^2+105e^4(ae-bd))}{105e^4(ae-bd)^3} \right)}{x^4 + \frac{d^4}{e^4} + \frac{4dx^3}{e} + \dots}$$

input `int(((A + B*x)*(a + b*x)^(1/2))/(d + e*x)^(9/2),x)`output `-((d + e*x)^(1/2)*((a + b*x)^(1/2)*(30*A*a^3*e^2 + 12*B*a^3*d*e + 70*A*a*b^2*d^2 - 28*B*a^2*b*d^2 - 84*A*a^2*b*d*e))/(105*e^4*(a*e - b*d)^3) + (x*(a + b*x)^(1/2)*(70*A*b^3*d^2 + 42*B*a^3*e^2 + 6*A*a^2*b*e^2 + 14*B*a*b^2*d^2 - 28*A*a*b^2*d*e - 104*B*a^2*b*d*e))/(105*e^4*(a*e - b*d)^3) + (4*b^2*x^3*(a + b*x)^(1/2)*(4*A*b*e - 7*B*a*e + 3*B*b*d))/(105*e^3*(a*e - b*d)^3) - (2*b*x^2*(a*e - 7*b*d)*(a + b*x)^(1/2)*(4*A*b*e - 7*B*a*e + 3*B*b*d))/(105*e^4*(a*e - b*d)^3)))/(x^4 + d^4/e^4 + (4*d*x^3)/e + (4*d^3*x)/e^3 + (6*d^2*x^2)/e^2)`**Reduce [B] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 414, normalized size of antiderivative = 2.82

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{9/2}} dx = \frac{-\frac{2\sqrt{ex+d}\sqrt{bx+a}a^3e^4}{7} + \frac{2\sqrt{ex+d}\sqrt{bx+a}a^2bde^3}{5} - \frac{16\sqrt{ex+d}\sqrt{bx+a}a^2be^4x}{35} + \frac{4\sqrt{ex+d}\sqrt{bx+a}ab^2}{5}}{e^3(a^2e^6x^4 - 2abde^5x^4 + b^2d^2e^4x^4 + 4a^2de^5x^3 - 8abd^2)}$$

input `int((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(9/2),x)`

output

```
(2*( - 5*sqrt(d + e*x)*sqrt(a + b*x)*a**3*e**4 + 7*sqrt(d + e*x)*sqrt(a +
b*x)*a**2*b*d*e**3 - 8*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b*e**4*x + 14*sqrt
(d + e*x)*sqrt(a + b*x)*a*b**2*d*e**3*x - sqrt(d + e*x)*sqrt(a + b*x)*a*b*
*2*e**4*x**2 + 7*sqrt(d + e*x)*sqrt(a + b*x)*b**3*d*e**3*x**2 + 2*sqrt(d +
e*x)*sqrt(a + b*x)*b**3*e**4*x**3 - 2*sqrt(e)*sqrt(b)*b**3*d**4 - 8*sqrt(
e)*sqrt(b)*b**3*d**3*e*x - 12*sqrt(e)*sqrt(b)*b**3*d**2*e**2*x**2 - 8*sqrt
(e)*sqrt(b)*b**3*d*e**3*x**3 - 2*sqrt(e)*sqrt(b)*b**3*e**4*x**4))/(35*e**3
*(a**2*d**4*e**2 + 4*a**2*d**3*e**3*x + 6*a**2*d**2*e**4*x**2 + 4*a**2*d*e
**5*x**3 + a**2*e**6*x**4 - 2*a*b*d**5*e - 8*a*b*d**4*e**2*x - 12*a*b*d**3
*e**3*x**2 - 8*a*b*d**2*e**4*x**3 - 2*a*b*d*e**5*x**4 + b**2*d**6 + 4*b**2
*d**5*e*x + 6*b**2*d**4*e**2*x**2 + 4*b**2*d**3*e**3*x**3 + b**2*d**2*e**4
*x**4))
```

3.180 $\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{11/2}} dx$

Optimal result	1635
Mathematica [A] (verified)	1636
Rubi [A] (verified)	1636
Maple [A] (verified)	1638
Fricas [B] (verification not implemented)	1639
Sympy [F]	1639
Maxima [F(-2)]	1640
Giac [B] (verification not implemented)	1640
Mupad [B] (verification not implemented)	1641
Reduce [B] (verification not implemented)	1642

Optimal result

Integrand size = 24, antiderivative size = 198

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{11/2}} dx = -\frac{2(Bd - Ae)(a+bx)^{3/2}}{9e(bd - ae)(d+ex)^{9/2}} + \frac{2(bBd + 2Abe - 3aBe)(a+bx)^{3/2}}{21e(bd - ae)^2(d+ex)^{7/2}} + \frac{8b(bBd + 2Abe - 3aBe)(a+bx)^{3/2}}{105e(bd - ae)^3(d+ex)^{5/2}} + \frac{16b^2(bBd + 2Abe - 3aBe)(a+bx)^{3/2}}{315e(bd - ae)^4(d+ex)^{3/2}}$$

output

```
-2/9*(-A*e+B*d)*(b*x+a)^(3/2)/e/(-a*e+b*d)/(e*x+d)^(9/2)+2/21*(2*A*b*e-3*B*a*e+B*b*d)*(b*x+a)^(3/2)/e/(-a*e+b*d)^2/(e*x+d)^(7/2)+8/105*b*(2*A*b*e-3*B*a*e+B*b*d)*(b*x+a)^(3/2)/e/(-a*e+b*d)^3/(e*x+d)^(5/2)+16/315*b^2*(2*A*b*e-3*B*a*e+B*b*d)*(b*x+a)^(3/2)/e/(-a*e+b*d)^4/(e*x+d)^(3/2)
```


Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.01

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{11/2}} dx = \frac{2(a+bx)^{3/2}(35Bde^2(a+bx)^3 - 35Ae^3(a+bx)^3 - 90bBde(a+bx)^2(d+ex) + 135A^2e^2(a+bx)^2(d+ex) + 135A^2b^2e^2(a+bx)^2(d+ex) - 45a^2B^2e^2(a+bx)^2(d+ex) + 63b^2B^2d^2(a+bx)(d+ex)^2 - 189A^2b^2e^2(a+bx)(d+ex)^2 + 126a^2b^2B^2e^2(a+bx)(d+ex)^2 + 105A^2b^3(d+ex)^3 - 105a^2b^2B^2(d+ex)^3)}{(315(bd-ae)^4(d+ex)^{9/2})}$$

input

```
Integrate[(Sqrt[a + b*x]*(A + B*x))/(d + e*x)^(11/2),x]
```

output

```
(2*(a + b*x)^(3/2)*(35*B*d*e^2*(a + b*x)^3 - 35*A*e^3*(a + b*x)^3 - 90*b*B*d*e*(a + b*x)^2*(d + e*x) + 135*A*b*e^2*(a + b*x)^2*(d + e*x) - 45*a*B*e^2*(a + b*x)^2*(d + e*x) + 63*b^2*B*d*(a + b*x)*(d + e*x)^2 - 189*A*b^2*e*(a + b*x)*(d + e*x)^2 + 126*a*b*B*e*(a + b*x)*(d + e*x)^2 + 105*A*b^3*(d + e*x)^3 - 105*a*b^2*B*(d + e*x)^3)/(315*(b*d - a*e)^4*(d + e*x)^(9/2))
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 190, normalized size of antiderivative = 0.96, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {87, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{11/2}} dx \\ & \quad \downarrow 87 \\ & \frac{(-3aBe + 2Abe + bBd) \int \frac{\sqrt{a+bx}}{(d+ex)^{9/2}} dx}{3e(bd-ae)} - \frac{2(a+bx)^{3/2}(Bd-Ae)}{9e(d+ex)^{9/2}(bd-ae)} \\ & \quad \downarrow 55 \\ & \frac{(-3aBe + 2Abe + bBd) \left(\frac{4b \int \frac{\sqrt{a+bx}}{(d+ex)^{7/2}} dx}{7(bd-ae)} + \frac{2(a+bx)^{3/2}}{7(d+ex)^{7/2}(bd-ae)} \right)}{3e(bd-ae)} - \frac{2(a+bx)^{3/2}(Bd-Ae)}{9e(d+ex)^{9/2}(bd-ae)} \\ & \quad \downarrow 55 \end{aligned}$$

$$\begin{aligned}
 & \frac{(-3aBe + 2Abe + bBd) \left(\frac{4b \left(\frac{2b \int \frac{\sqrt{a+bx}}{(d+ex)^{5/2}} dx + \frac{2(a+bx)^{3/2}}{5(d+ex)^{5/2}(bd-ae)} \right)}{7(bd-ae)} + \frac{2(a+bx)^{3/2}}{7(d+ex)^{7/2}(bd-ae)} \right)}{3e(bd-ae)} \\
 & \frac{2(a+bx)^{3/2}(Bd-Ae)}{9e(d+ex)^{9/2}(bd-ae)} \\
 & \quad \downarrow 48 \\
 & \frac{\left(\frac{2(a+bx)^{3/2}}{7(d+ex)^{7/2}(bd-ae)} + \frac{4b \left(\frac{4b(a+bx)^{3/2}}{15(d+ex)^{3/2}(bd-ae)^2} + \frac{2(a+bx)^{3/2}}{5(d+ex)^{5/2}(bd-ae)} \right)}{7(bd-ae)} \right) (-3aBe + 2Abe + bBd)}{3e(bd-ae)} \\
 & \frac{2(a+bx)^{3/2}(Bd-Ae)}{9e(d+ex)^{9/2}(bd-ae)}
 \end{aligned}$$

input `Int[(Sqrt[a + b*x]*(A + B*x))/(d + e*x)^(11/2),x]`

output `(-2*(B*d - A*e)*(a + b*x)^(3/2))/(9*e*(b*d - a*e)*(d + e*x)^(9/2)) + ((b*B*d + 2*A*b*e - 3*a*B*e)*((2*(a + b*x)^(3/2))/(7*(b*d - a*e)*(d + e*x)^(7/2))) + (4*b*((2*(a + b*x)^(3/2))/(5*(b*d - a*e)*(d + e*x)^(5/2)) + (4*b*(a + b*x)^(3/2))/(15*(b*d - a*e)^2*(d + e*x)^(3/2))))/(7*(b*d - a*e)))/(3*e*(b*d - a*e))`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 55 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(Simplify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.42

method	result
default	$-\frac{2(bx+a)^{\frac{3}{2}}(-16Ab^3e^3x^3+24Bab^2e^3x^3-8Bb^3de^2x^3+24Aab^2e^3x^2-72Ab^3de^2x^2-36Ba^2be^3x^2+120Bab^2de^2x^2-36Bb^3d^2e^2x^2-36Bb^3d^2e^2x^2-36Bb^3d^2e^2x^2)}{2(bx+a)^{\frac{3}{2}}(-16Ab^3e^3x^3+24Bab^2e^3x^3-8Bb^3de^2x^3+24Aab^2e^3x^2-72Ab^3de^2x^2-36Ba^2be^3x^2+120Bab^2de^2x^2-36Bb^3d^2e^2x^2-36Bb^3d^2e^2x^2-36Bb^3d^2e^2x^2)}$
gosper	$-\frac{2(bx+a)^{\frac{3}{2}}(-16Ab^3e^3x^3+24Bab^2e^3x^3-8Bb^3de^2x^3+24Aab^2e^3x^2-72Ab^3de^2x^2-36Ba^2be^3x^2+120Bab^2de^2x^2-36Bb^3d^2e^2x^2-36Bb^3d^2e^2x^2-36Bb^3d^2e^2x^2)}{2(bx+a)^{\frac{3}{2}}(-16Ab^3e^3x^3+24Bab^2e^3x^3-8Bb^3de^2x^3+24Aab^2e^3x^2-72Ab^3de^2x^2-36Ba^2be^3x^2+120Bab^2de^2x^2-36Bb^3d^2e^2x^2-36Bb^3d^2e^2x^2-36Bb^3d^2e^2x^2)}$
orering	$-\frac{2(bx+a)^{\frac{3}{2}}(-16Ab^3e^3x^3+24Bab^2e^3x^3-8Bb^3de^2x^3+24Aab^2e^3x^2-72Ab^3de^2x^2-36Ba^2be^3x^2+120Bab^2de^2x^2-36Bb^3d^2e^2x^2-36Bb^3d^2e^2x^2-36Bb^3d^2e^2x^2)}{2(bx+a)^{\frac{3}{2}}(-16Ab^3e^3x^3+24Bab^2e^3x^3-8Bb^3de^2x^3+24Aab^2e^3x^2-72Ab^3de^2x^2-36Ba^2be^3x^2+120Bab^2de^2x^2-36Bb^3d^2e^2x^2-36Bb^3d^2e^2x^2-36Bb^3d^2e^2x^2)}$

input `int((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(11/2), x, method=_RETURNVERBOSE)`

output
$$-\frac{2}{315}(bx+a)^{\frac{3}{2}}(e*x+d)^{\frac{9}{2}}(-16A*b^3*e^3*x^3+24*B*a*b^2*e^3*x^3-8*B*b^3*d*e^2*x^3+24*A*a*b^2*e^3*x^2-72*A*b^3*d*e^2*x^2-36*B*a^2*b*e^3*x^2+120*B*a*b^2*d*e^2*x^2-36*B*b^3*d^2*e*x^2-30*A*a^2*b*e^3*x+108*A*a*b^2*d*e^2*x-126*A*b^3*d^2*e*x+45*B*a^3*e^3*x-177*B*a^2*b*d*e^2*x+243*B*a*b^2*d^2*e*x-63*B*b^3*d^3*x+35*A*a^3*e^3-135*A*a^2*b*d*e^2+189*A*a*b^2*d^2*e-105*A*b^3*d^3+10*B*a^3*d*e^2-36*B*a^2*b*d^2*e+42*B*a*b^2*d^3)/(a*e-b*d)^4$$

output `Integral((A + B*x)*sqrt(a + b*x)/(d + e*x)**(11/2), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a + bx}(A + Bx)}{(d + ex)^{11/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(11/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f or more de`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 620 vs. $2(174) = 348$.

Time = 0.36 (sec) , antiderivative size = 620, normalized size of antiderivative = 3.13

$$\int \frac{\sqrt{a + bx}(A + Bx)}{(d + ex)^{11/2}} dx = \frac{2 \left(\left(4(bx + a) \left(\frac{2(Bb^{10}de^6|b| - 3Bab^9e^7|b| + 2Ab^{10}e^7|b|)(bx+a)}{b^6d^4e^4 - 4ab^5d^3e^5 + 6a^2b^4d^2e^6 - 4a^3b^3de^7 + a^4b^2e^8} + \frac{9(Bb^{11}d^2e^5|b| - 4Bab^{10}de^6|b|)}{b^6d^4e^4 - 4ab^5d^3e^5} \right) \right)}{b^6d^4e^4 - 4ab^5d^3e^5 + 6a^2b^4d^2e^6 - 4a^3b^3de^7 + a^4b^2e^8}$$

input `integrate((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(11/2),x, algorithm="giac")`

Reduce [B] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 702, normalized size of antiderivative = 3.55

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{11/2}} dx = \frac{-\frac{2\sqrt{ex+d}\sqrt{bx+a}a^4e^5}{9} + \frac{4\sqrt{ex+d}\sqrt{bx+a}a^3bde^4}{7} - \frac{20\sqrt{ex+d}\sqrt{bx+a}a^3be^5x}{63} - \frac{2\sqrt{ex+d}\sqrt{bx+a}a^2b^2e^5}{5}}{e^3(a^3e^8x^5 - 3a^2bde^7x^5 + 3ab^2d^2e^6x^5 - b^3d^3e^5x^5 + 5a^3de^4x^5)}$$

input `int((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(11/2),x)`

output

```
(2*(-35*sqrt(d+e*x)*sqrt(a+b*x)*a**4*e**5 + 90*sqrt(d+e*x)*sqrt(a+b*x)*a**3*b*d*e**4 - 50*sqrt(d+e*x)*sqrt(a+b*x)*a**3*b*e**5*x - 63*sqrt(d+e*x)*sqrt(a+b*x)*a**2*b**2*d**2*e**3 + 144*sqrt(d+e*x)*sqrt(a+b*x)*a**2*b**2*d*e**4*x - 3*sqrt(d+e*x)*sqrt(a+b*x)*a**2*b**2*e**5*x**2 - 126*sqrt(d+e*x)*sqrt(a+b*x)*a*b**3*d**2*e**3*x + 18*sqrt(d+e*x)*sqrt(a+b*x)*a*b**3*d*e**4*x**2 + 4*sqrt(d+e*x)*sqrt(a+b*x)*a*b**3*e**5*x**3 - 63*sqrt(d+e*x)*sqrt(a+b*x)*b**4*d**2*e**3*x**2 - 36*sqrt(d+e*x)*sqrt(a+b*x)*b**4*d*e**4*x**3 - 8*sqrt(d+e*x)*sqrt(a+b*x)*b**4*e**5*x**4 + 8*sqrt(e)*sqrt(b)*b**4*d**5 + 40*sqrt(e)*sqrt(b)*b**4*d**4*e*x + 80*sqrt(e)*sqrt(b)*b**4*d**3*e**2*x**2 + 80*sqrt(e)*sqrt(b)*b**4*d**2*e**3*x**3 + 40*sqrt(e)*sqrt(b)*b**4*d*e**4*x**4 + 8*sqrt(e)*sqrt(b)*b**4*e**5*x**5)/(315*e**3*(a**3*d**5*e**3 + 5*a**3*d**4*e**4*x + 10*a**3*d**3*e**5*x**2 + 10*a**3*d**2*e**6*x**3 + 5*a**3*d*e**7*x**4 + a**3*e**8*x**5 - 3*a**2*b*d**6*e**2 - 15*a**2*b*d**5*e**3*x - 30*a**2*b*d**4*e**4*x**2 - 30*a**2*b*d**3*e**5*x**3 - 15*a**2*b*d**2*e**6*x**4 - 3*a**2*b*d*e**7*x**5 + 3*a*b**2*d**7*e + 15*a*b**2*d**6*e**2*x + 30*a*b**2*d**5*e**3*x**2 + 30*a*b**2*d**4*e**4*x**3 + 15*a*b**2*d**3*e**5*x**4 + 3*a*b**2*d**2*e**6*x**5 - b**3*d**8 - 5*b**3*d**7*e*x - 10*b**3*d**6*e**2*x**2 - 10*b**3*d**5*e**3*x**3 - 5*b**3*d**4*e**4*x**4 - b**3*d**3*e**5*x**5))
```

3.181 $\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{13/2}} dx$

Optimal result	1643
Mathematica [A] (verified)	1644
Rubi [A] (verified)	1644
Maple [A] (verified)	1647
Fricas [B] (verification not implemented)	1647
Sympy [F]	1648
Maxima [F(-2)]	1649
Giac [B] (verification not implemented)	1649
Mupad [B] (verification not implemented)	1650
Reduce [B] (verification not implemented)	1651

Optimal result

Integrand size = 24, antiderivative size = 255

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{13/2}} dx = -\frac{2(Bd - Ae)(a+bx)^{3/2}}{11e(bd - ae)(d+ex)^{11/2}} + \frac{2(3bBd + 8Abe - 11aBe)(a+bx)^{3/2}}{99e(bd - ae)^2(d+ex)^{9/2}} + \frac{4b(3bBd + 8Abe - 11aBe)(a+bx)^{3/2}}{231e(bd - ae)^3(d+ex)^{7/2}} + \frac{16b^2(3bBd + 8Abe - 11aBe)(a+bx)^{3/2}}{1155e(bd - ae)^4(d+ex)^{5/2}} + \frac{32b^3(3bBd + 8Abe - 11aBe)(a+bx)^{3/2}}{3465e(bd - ae)^5(d+ex)^{3/2}}$$

output

```
-2/11*(-A*e+B*d)*(b*x+a)^(3/2)/e/(-a*e+b*d)/(e*x+d)^(11/2)+2/99*(8*A*b*e-1
1*B*a*e+3*B*b*d)*(b*x+a)^(3/2)/e/(-a*e+b*d)^2/(e*x+d)^(9/2)+4/231*b*(8*A*b
*e-11*B*a*e+3*B*b*d)*(b*x+a)^(3/2)/e/(-a*e+b*d)^3/(e*x+d)^(7/2)+16/1155*b^
2*(8*A*b*e-11*B*a*e+3*B*b*d)*(b*x+a)^(3/2)/e/(-a*e+b*d)^4/(e*x+d)^(5/2)+32
/3465*b^3*(8*A*b*e-11*B*a*e+3*B*b*d)*(b*x+a)^(3/2)/e/(-a*e+b*d)^5/(e*x+d)^(
3/2)
```


Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.06

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{13/2}} dx = \frac{2(a+bx)^{3/2}(-315Bde^3(a+bx)^4 + 315Ae^4(a+bx)^4 + 1155bBde^2(a+bx)^3(d+ex) - 1540A^2b^2e^3(a+bx)^3(d+ex) + 385a^2B^2e^3(a+bx)^3(d+ex) - 1485b^2B^2de^2(a+bx)^2(d+ex)^2 + 2970A^2b^2e^2(a+bx)^2(d+ex)^2 - 1485a^2b^2B^2e^2(a+bx)^2(d+ex)^2 + 693b^3B^2d(a+bx)(d+ex)^3 - 2772A^2b^3e(a+bx)(d+ex)^3 + 2079a^2b^2B^2e(a+bx)(d+ex)^3 + 1155A^2b^4(d+ex)^4 - 1155a^2b^3B^2(d+ex)^4)/(3465*(b*d - a*e)^5*(d+ex)^{(11/2)})$$

input

```
Integrate[(Sqrt[a + b*x]*(A + B*x))/(d + e*x)^(13/2),x]
```

output

```
(2*(a + b*x)^(3/2)*(-315*B*d*e^3*(a + b*x)^4 + 315*A*e^4*(a + b*x)^4 + 1155*b*B*d*e^2*(a + b*x)^3*(d + e*x) - 1540*A^2*b^2*e^3*(a + b*x)^3*(d + e*x) + 385*a^2*B^2*e^3*(a + b*x)^3*(d + e*x) - 1485*b^2*B^2*d*e^2*(a + b*x)^2*(d + e*x)^2 + 2970*A^2*b^2*e^2*(a + b*x)^2*(d + e*x)^2 - 1485*a^2*b^2*B^2*e^2*(a + b*x)^2*(d + e*x)^2 + 693*b^3*B^2*d*(a + b*x)*(d + e*x)^3 - 2772*A^2*b^3*e*(a + b*x)*(d + e*x)^3 + 2079*a^2*b^2*B^2*e*(a + b*x)*(d + e*x)^3 + 1155*A^2*b^4*(d + e*x)^4 - 1155*a^2*b^3*B^2*(d + e*x)^4)/(3465*(b*d - a*e)^5*(d + e*x)^(11/2))
```

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 239, normalized size of antiderivative = 0.94, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$, Rules used = {87, 55, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{13/2}} dx$$

$$\downarrow 87$$

$$\frac{(-11aBe + 8Abe + 3bBd) \int \frac{\sqrt{a+bx}}{(d+ex)^{11/2}} dx}{11e(bd - ae)} - \frac{2(a+bx)^{3/2}(Bd - Ae)}{11e(d+ex)^{11/2}(bd - ae)}$$

$$\downarrow 55$$

$$\frac{(-11aBe + 8Abe + 3bBd) \left(\frac{2b \int \frac{\sqrt{a+bx}}{(d+ex)^{9/2}} dx}{3(bd - ae)} + \frac{2(a+bx)^{3/2}}{9(d+ex)^{9/2}(bd - ae)} \right)}{11e(bd - ae)} - \frac{2(a+bx)^{3/2}(Bd - Ae)}{11e(d+ex)^{11/2}(bd - ae)}$$

$$\begin{aligned}
 & \downarrow 55 \\
 & (-11aBe + 8Abe + 3bBd) \left(\frac{2b \left(\frac{4b \int \frac{\sqrt{a+bx}}{(d+ex)^{7/2}} dx}{7(bd-ae)} + \frac{2(a+bx)^{3/2}}{7(d+ex)^{7/2}(bd-ae)} \right)}{3(bd-ae)} + \frac{2(a+bx)^{3/2}}{9(d+ex)^{9/2}(bd-ae)} \right)
 \end{aligned}$$

$$\frac{11e(bd - ae)}{11e(d + ex)^{11/2}(bd - ae)}$$

↓ 55

$$\begin{aligned}
 & (-11aBe + 8Abe + 3bBd) \left(\frac{2b \left(\frac{4b \left(\frac{2b \int \frac{\sqrt{a+bx}}{(d+ex)^{5/2}} dx}{5(bd-ae)} + \frac{2(a+bx)^{3/2}}{5(d+ex)^{5/2}(bd-ae)} \right)}{7(bd-ae)} + \frac{2(a+bx)^{3/2}}{7(d+ex)^{7/2}(bd-ae)} \right)}{3(bd-ae)} + \frac{2(a+bx)^{3/2}}{9(d+ex)^{9/2}(bd-ae)} \right)
 \end{aligned}$$

$$\frac{11e(bd - ae)}{11e(d + ex)^{11/2}(bd - ae)}$$

↓ 48

$$\left(\frac{2(a+bx)^{3/2}}{9(d+ex)^{9/2}(bd-ae)} + \frac{2b \left(\frac{2(a+bx)^{3/2}}{7(d+ex)^{7/2}(bd-ae)} + \frac{4b \left(\frac{4b(a+bx)^{3/2}}{15(d+ex)^{3/2}(bd-ae)^2} + \frac{2(a+bx)^{3/2}}{5(d+ex)^{5/2}(bd-ae)} \right)}{7(bd-ae)} \right)}{3(bd-ae)} \right) (-11aBe + 8Abe + 3bBd)$$

$$\frac{11e(bd - ae)}{11e(d + ex)^{11/2}(bd - ae)}$$

input

Int[(Sqrt[a + b*x]*(A + B*x))/(d + e*x)^(13/2),x]

output

$$\begin{aligned} & (-2*(B*d - A*e)*(a + b*x)^{(3/2)})/(11*e*(b*d - a*e)*(d + e*x)^{(11/2)}) + ((3 \\ & *b*B*d + 8*A*b*e - 11*a*B*e)*((2*(a + b*x)^{(3/2)})/(9*(b*d - a*e)*(d + e*x) \\ & ^{(9/2)}) + (2*b*((2*(a + b*x)^{(3/2)})/(7*(b*d - a*e)*(d + e*x)^{(7/2)}) + (4*b \\ & *((2*(a + b*x)^{(3/2)})/(5*(b*d - a*e)*(d + e*x)^{(5/2)}) + (4*b*(a + b*x)^{(3/ \\ & 2))/(15*(b*d - a*e)^2*(d + e*x)^{(3/2)})))/(7*(b*d - a*e)))/(3*(b*d - a*e)) \\ &))/(11*e*(b*d - a*e)) \end{aligned}$$

Defintions of rubi rules used

rule 48

$$\text{Int}[(a_.) + (b_.)*(x_)^{(m_.)}*((c_.) + (d_.)*(x_)^{(n_.)}), x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^{(n + 1)})/((b*c - a*d)*(m + 1))], x] \text{ ; FreeQ}[\{a, b, c, d, m, n\}, x] \ \&\& \ \text{EqQ}[m + n + 2, 0] \ \&\& \ \text{NeQ}[m, -1]$$

rule 55

$$\begin{aligned} & \text{Int}[(a_.) + (b_.)*(x_)^{(m_.)}*((c_.) + (d_.)*(x_)^{(n_.)}), x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^{(n + 1)})/((b*c - a*d)*(m + 1))], x] \\ & - \text{Simp}[d*(\text{Simplify}[m + n + 2]/((b*c - a*d)*(m + 1))) \ \text{Int}[(a + b*x)^{\text{Simplify}[m + 1]}*(c + d*x)^n, x], x] \text{ ; FreeQ}[\{a, b, c, d, m, n\}, x] \ \&\& \ \text{ILtQ}[\text{Simplify}[m + n + 2], 0] \ \&\& \ \text{NeQ}[m, -1] \ \&\& \ \text{!(LtQ}[m, -1] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ (\text{EqQ}[a, 0] \ || \ (\text{NeQ}[c, 0] \ \&\& \ \text{LtQ}[m - n, 0] \ \&\& \ \text{IntegerQ}[n]))) \ \&\& \ (\text{SumSimplerQ}[m, 1] \ || \ \text{!SumSimplerQ}[n, 1]) \end{aligned}$$

rule 87

$$\begin{aligned} & \text{Int}[(a_.) + (b_.)*(x_)^{(m_.)}*((c_.) + (d_.)*(x_)^{(n_.)}*((e_.) + (f_.)*(x_)^{(p_.)}), x_Symbol] \rightarrow \text{Simp}[(-b*e - a*f)*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)})/(f*(p + 1)*(c*f - d*e))], x] \\ & - \text{Simp}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) \ \text{Int}[(c + d*x)^n*(e + f*x)^{(p + 1)}, x], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, n\}, x] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ (\text{!LtQ}[n, -1] \ || \ \text{IntegerQ}[p] \ || \ \text{!(IntegerQ}[n] \ || \ \text{!(EqQ}[e, 0] \ || \ \text{!(EqQ}[c, 0] \ || \ \text{LtQ}[p, n])))) \end{aligned}$$

Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 449, normalized size of antiderivative = 1.76

method	result
default	$-\frac{2(bx+a)^{\frac{3}{2}}(128Ab^4e^4x^4-176Ba b^3e^4x^4+48Bb^4de^3x^4-192Aab^3e^4x^3+704Ab^4de^3x^3+264Ba^2b^2e^4x^3-1040Bab^3de^3x^3+264A^2b^2e^4x^2-1056Aab^3de^3x^2+1584Aab^4d^2e^2x^2-330Bba^3be^4x^2+1542Bba^2b^2de^3x^2-2574Bba^3bd^2e^2x^2+594Bb^4d^3e^2x^2-280Aa^3be^4x+1320Aa^2b^2de^3x-2376Aa^3bd^2e^2x+1848Ab^4d^3e^2x+385Bba^4e^4x-1920Bba^3bd^2e^3x+3762Bba^2b^2d^2e^2x-3432Bba^3bd^3e^2x+693Bb^4d^4x+315Aa^4e^4-1540Aa^3bd^2e^3+2970Aa^2b^2d^2e^2-2772Aa^3bd^3e+1155Ab^4d^4+70Bba^4de^3-330Bba^3bd^2e^2+594Bba^2b^2d^3e-462Bba^3bd^4)/(a^2e-b^2d)^5$
gosper	$-\frac{2(bx+a)^{\frac{3}{2}}(128Ab^4e^4x^4-176Ba b^3e^4x^4+48Bb^4de^3x^4-192Aab^3e^4x^3+704Ab^4de^3x^3+264Ba^2b^2e^4x^3-1040Bab^3de^3x^3+264A^2b^2e^4x^2-1056Aab^3de^3x^2+1584Aab^4d^2e^2x^2-330Bba^3be^4x^2+1542Bba^2b^2de^3x^2-2574Bba^3bd^2e^2x^2+594Bb^4d^3e^2x^2-280Aa^3be^4x+1320Aa^2b^2de^3x-2376Aa^3bd^2e^2x+1848Ab^4d^3e^2x+385Bba^4e^4x-1920Bba^3bd^2e^3x+3762Bba^2b^2d^2e^2x-3432Bba^3bd^3e^2x+693Bb^4d^4x+315Aa^4e^4-1540Aa^3bd^2e^3+2970Aa^2b^2d^2e^2-2772Aa^3bd^3e+1155Ab^4d^4+70Bba^4de^3-330Bba^3bd^2e^2+594Bba^2b^2d^3e-462Bba^3bd^4)/(a^2e-b^2d)^5$
orering	$-\frac{2(bx+a)^{\frac{3}{2}}(128Ab^4e^4x^4-176Ba b^3e^4x^4+48Bb^4de^3x^4-192Aab^3e^4x^3+704Ab^4de^3x^3+264Ba^2b^2e^4x^3-1040Bab^3de^3x^3+264A^2b^2e^4x^2-1056Aab^3de^3x^2+1584Aab^4d^2e^2x^2-330Bba^3be^4x^2+1542Bba^2b^2de^3x^2-2574Bba^3bd^2e^2x^2+594Bb^4d^3e^2x^2-280Aa^3be^4x+1320Aa^2b^2de^3x-2376Aa^3bd^2e^2x+1848Ab^4d^3e^2x+385Bba^4e^4x-1920Bba^3bd^2e^3x+3762Bba^2b^2d^2e^2x-3432Bba^3bd^3e^2x+693Bb^4d^4x+315Aa^4e^4-1540Aa^3bd^2e^3+2970Aa^2b^2d^2e^2-2772Aa^3bd^3e+1155Ab^4d^4+70Bba^4de^3-330Bba^3bd^2e^2+594Bba^2b^2d^3e-462Bba^3bd^4)/(a^2e-b^2d)^5$

input `int((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(13/2),x,method=_RETURNVERBOSE)`

output $-\frac{2}{3465}(b*x+a)^{\frac{3}{2}}/(e*x+d)^{\frac{11}{2}}*(128*A*b^4*e^4*x^4-176*B*a*b^3*e^4*x^4+48*B*b^4*d*e^3*x^4-192*A*a*b^3*e^4*x^3+704*A*b^4*d*e^3*x^3+264*B*a^2*b^2*e^4*x^3-1040*B*a*b^3*d*e^3*x^3+264*B*b^4*d^2*e^2*x^3+240*A*a^2*b^2*e^4*x^2-1056*A*a*b^3*d*e^3*x^2+1584*A*b^4*d^2*e^2*x^2-330*B*a^3*b*e^4*x^2+1542*B*a^2*b^2*d*e^3*x^2-2574*B*a*b^3*d^2*e^2*x^2+594*B*b^4*d^3*e^2*x^2-280*A*a^3*b*e^4*x+1320*A*a^2*b^2*d*e^3*x-2376*A*a*b^3*d^2*e^2*x+1848*A*b^4*d^3*e^2*x+385*B*a^4*e^4*x-1920*B*a^3*b*d^2*e^3*x+3762*B*a^2*b^2*d^2*e^2*x-3432*B*a*b^3*d^3*e^2*x+693*B*b^4*d^4*x+315*A*a^4*e^4-1540*A*a^3*b*d^2*e^3+2970*A*a^2*b^2*d^2*e^2-2772*A*a*b^3*d^3*e+1155*A*b^4*d^4+70*B*a^4*d*e^3-330*B*a^3*b*d^2*e^2+594*B*a^2*b^2*d^3*e-462*B*a*b^3*d^4)/(a^2e-b^2d)^5$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1045 vs. 2(225) = 450.

Time = 64.58 (sec) , antiderivative size = 1045, normalized size of antiderivative = 4.10

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{13/2}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(13/2),x, algorithm="fricas")`

output

```

2/3465*(315*A*a^5*e^4 + 16*(3*B*b^5*d*e^3 - (11*B*a*b^4 - 8*A*b^5)*e^4)*x^
5 - 231*(2*B*a^2*b^3 - 5*A*a*b^4)*d^4 + 198*(3*B*a^3*b^2 - 14*A*a^2*b^3)*d
^3*e - 330*(B*a^4*b - 9*A*a^3*b^2)*d^2*e^2 + 70*(B*a^5 - 22*A*a^4*b)*d*e^3
+ 8*(33*B*b^5*d^2*e^2 - 4*(31*B*a*b^4 - 22*A*b^5)*d*e^3 + (11*B*a^2*b^3 -
8*A*a*b^4)*e^4)*x^4 + 2*(297*B*b^5*d^3*e - 33*(35*B*a*b^4 - 24*A*b^5)*d^2
*e^2 + (251*B*a^2*b^3 - 176*A*a*b^4)*d*e^3 - 3*(11*B*a^3*b^2 - 8*A*a^2*b^3
)*e^4)*x^3 + (693*B*b^5*d^4 - 66*(43*B*a*b^4 - 28*A*b^5)*d^3*e + 396*(3*B*
a^2*b^3 - 2*A*a*b^4)*d^2*e^2 - 6*(63*B*a^3*b^2 - 44*A*a^2*b^3)*d*e^3 + 5*(
11*B*a^4*b - 8*A*a^3*b^2)*e^4)*x^2 + (231*(B*a*b^4 + 5*A*b^5)*d^4 - 66*(43
*B*a^2*b^3 + 14*A*a*b^4)*d^3*e + 66*(52*B*a^3*b^2 + 9*A*a^2*b^3)*d^2*e^2 -
10*(185*B*a^4*b + 22*A*a^3*b^2)*d*e^3 + 35*(11*B*a^5 + A*a^4*b)*e^4)*x)*s
qrt(b*x + a)*sqrt(e*x + d)/(b^5*d^11 - 5*a*b^4*d^10*e + 10*a^2*b^3*d^9*e^2
- 10*a^3*b^2*d^8*e^3 + 5*a^4*b*d^7*e^4 - a^5*d^6*e^5 + (b^5*d^5*e^6 - 5*a
*b^4*d^4*e^7 + 10*a^2*b^3*d^3*e^8 - 10*a^3*b^2*d^2*e^9 + 5*a^4*b*d*e^10 -
a^5*e^11)*x^6 + 6*(b^5*d^6*e^5 - 5*a*b^4*d^5*e^6 + 10*a^2*b^3*d^4*e^7 - 10
*a^3*b^2*d^3*e^8 + 5*a^4*b*d^2*e^9 - a^5*d*e^10)*x^5 + 15*(b^5*d^7*e^4 - 5
*a*b^4*d^6*e^5 + 10*a^2*b^3*d^5*e^6 - 10*a^3*b^2*d^4*e^7 + 5*a^4*b*d^3*e^8
- a^5*d^2*e^9)*x^4 + 20*(b^5*d^8*e^3 - 5*a*b^4*d^7*e^4 + 10*a^2*b^3*d^6*e
^5 - 10*a^3*b^2*d^5*e^6 + 5*a^4*b*d^4*e^7 - a^5*d^3*e^8)*x^3 + 15*(b^5*d^9
*e^2 - 5*a*b^4*d^8*e^3 + 10*a^2*b^3*d^7*e^4 - 10*a^3*b^2*d^6*e^5 + 5*a^...

```

Sympy [F]

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{13/2}} dx = \int \frac{(A+Bx)\sqrt{a+bx}}{(d+ex)^{\frac{13}{2}}} dx$$

input

```
integrate((b*x+a)**(1/2)*(B*x+A)/(e*x+d)**(13/2),x)
```

output

```
Integral((A + B*x)*sqrt(a + b*x)/(d + e*x)**(13/2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{13/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(13/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f or more de`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 937 vs. 2(225) = 450.

Time = 0.41 (sec) , antiderivative size = 937, normalized size of antiderivative = 3.67

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{13/2}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(13/2),x, algorithm="giac")`

output

```

-((d + e*x)^(1/2)*(((a + b*x)^(1/2)*(630*A*a^5*e^4 + 2310*A*a*b^4*d^4 + 14
0*B*a^5*d*e^3 - 924*B*a^2*b^3*d^4 - 5544*A*a^2*b^3*d^3*e + 1188*B*a^3*b^2*
d^3*e - 660*B*a^4*b*d^2*e^2 + 5940*A*a^3*b^2*d^2*e^2 - 3080*A*a^4*b*d*e^3)
)/(3465*e^6*(a*e - b*d)^5) + (x*(a + b*x)^(1/2)*(2310*A*b^5*d^4 + 770*B*a^
5*e^4 + 70*A*a^4*b*e^4 + 462*B*a*b^4*d^4 - 440*A*a^3*b^2*d*e^3 - 5676*B*a^
2*b^3*d^3*e + 1188*A*a^2*b^3*d^2*e^2 + 6864*B*a^3*b^2*d^2*e^2 - 1848*A*a*b
^4*d^3*e - 3700*B*a^4*b*d*e^3))/(3465*e^6*(a*e - b*d)^5) + (32*b^4*x^5*(a
+ b*x)^(1/2)*(8*A*b*e - 11*B*a*e + 3*B*b*d))/(3465*e^3*(a*e - b*d)^5) - (1
6*b^3*x^4*(a*e - 11*b*d)*(a + b*x)^(1/2)*(8*A*b*e - 11*B*a*e + 3*B*b*d))/(
3465*e^4*(a*e - b*d)^5) + (4*b^2*x^3*(a + b*x)^(1/2)*(3*a^2*e^2 + 99*b^2*d
^2 - 22*a*b*d*e)*(8*A*b*e - 11*B*a*e + 3*B*b*d))/(3465*e^5*(a*e - b*d)^5)
- (2*b*x^2*(a + b*x)^(1/2)*(8*A*b*e - 11*B*a*e + 3*B*b*d)*(5*a^3*e^3 - 231
*b^3*d^3 + 99*a*b^2*d^2*e - 33*a^2*b*d*e^2))/(3465*e^6*(a*e - b*d)^5))/(x
^6 + d^6/e^6 + (6*d*x^5)/e + (6*d^5*x)/e^5 + (15*d^2*x^4)/e^2 + (20*d^3*x^
3)/e^3 + (15*d^4*x^2)/e^4)

```

Reduce [B] (verification not implemented)

Time = 0.53 (sec) , antiderivative size = 1049, normalized size of antiderivative = 4.11

$$\int \frac{\sqrt{a+bx}(A+Bx)}{(d+ex)^{13/2}} dx = \text{Too large to display}$$

input

```
int((b*x+a)^(1/2)*(B*x+A)/(e*x+d)^(13/2),x)
```


output

```
(2*( - 105*sqrt(d + e*x)*sqrt(a + b*x)*a**5*e**6 + 385*sqrt(d + e*x)*sqrt(
a + b*x)*a**4*b*d*e**5 - 140*sqrt(d + e*x)*sqrt(a + b*x)*a**4*b*e**6*x - 4
95*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b**2*d**2*e**4 + 550*sqrt(d + e*x)*sqr
t(a + b*x)*a**3*b**2*d*e**5*x - 5*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b**2*e*
*6*x**2 + 231*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**3*d**3*e**3 - 792*sqrt(d
+ e*x)*sqrt(a + b*x)*a**2*b**3*d**2*e**4*x + 33*sqrt(d + e*x)*sqrt(a + b*
x)*a**2*b**3*d*e**5*x**2 + 6*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**3*e**6*x*
*3 + 462*sqrt(d + e*x)*sqrt(a + b*x)*a*b**4*d**3*e**3*x - 99*sqrt(d + e*x)
*sqrt(a + b*x)*a*b**4*d**2*e**4*x**2 - 44*sqrt(d + e*x)*sqrt(a + b*x)*a*b*
*4*d*e**5*x**3 - 8*sqrt(d + e*x)*sqrt(a + b*x)*a*b**4*e**6*x**4 + 231*sqrt
(d + e*x)*sqrt(a + b*x)*b**5*d**3*e**3*x**2 + 198*sqrt(d + e*x)*sqrt(a + b
*x)*b**5*d**2*e**4*x**3 + 88*sqrt(d + e*x)*sqrt(a + b*x)*b**5*d*e**5*x**4
+ 16*sqrt(d + e*x)*sqrt(a + b*x)*b**5*e**6*x**5 - 16*sqrt(e)*sqrt(b)*b**5*
d**6 - 96*sqrt(e)*sqrt(b)*b**5*d**5*e*x - 240*sqrt(e)*sqrt(b)*b**5*d**4*e*
*2*x**2 - 320*sqrt(e)*sqrt(b)*b**5*d**3*e**3*x**3 - 240*sqrt(e)*sqrt(b)*b*
*5*d**2*e**4*x**4 - 96*sqrt(e)*sqrt(b)*b**5*d*e**5*x**5 - 16*sqrt(e)*sqrt(
b)*b**5*e**6*x**6))/(1155*e**3*(a**4*d**6*e**4 + 6*a**4*d**5*e**5*x + 15*a
**4*d**4*e**6*x**2 + 20*a**4*d**3*e**7*x**3 + 15*a**4*d**2*e**8*x**4 + 6*a
**4*d*e**9*x**5 + a**4*e**10*x**6 - 4*a**3*b*d**7*e**3 - 24*a**3*b*d**6*e*
*4*x - 60*a**3*b*d**5*e**5*x**2 - 80*a**3*b*d**4*e**6*x**3 - 60*a**3*b*...
```

3.182 $\int (a + bx)^{3/2}(A + Bx)(d + ex)^{5/2} dx$

Optimal result	1653
Mathematica [A] (verified)	1654
Rubi [A] (verified)	1655
Maple [B] (verified)	1661
Fricas [B] (verification not implemented)	1662
Sympy [F]	1663
Maxima [F(-2)]	1663
Giac [B] (verification not implemented)	1663
Mupad [F(-1)]	1664
Reduce [B] (verification not implemented)	1665

Optimal result

Integrand size = 24, antiderivative size = 358

$$\begin{aligned}
 & \int (a + bx)^{3/2}(A + Bx)(d + ex)^{5/2} dx = \frac{(bd - ae)^4(5bBd - 12Abe + 7aBe)\sqrt{a + bx}\sqrt{d + ex}}{512b^4e^3} \\
 & - \frac{(bd - ae)^3(5bBd - 12Abe + 7aBe)(a + bx)^{3/2}\sqrt{d + ex}}{768b^4e^2} \\
 & - \frac{(bd - ae)^2(5bBd - 12Abe + 7aBe)(a + bx)^{5/2}\sqrt{d + ex}}{192b^4e} \\
 & - \frac{(bd - ae)(5bBd - 12Abe + 7aBe)(a + bx)^{5/2}(d + ex)^{3/2}}{96b^3e} \\
 & - \frac{(5bBd - 12Abe + 7aBe)(a + bx)^{5/2}(d + ex)^{5/2}}{60b^2e} + \frac{B(a + bx)^{5/2}(d + ex)^{7/2}}{6be} \\
 & - \frac{(bd - ae)^5(5bBd - 12Abe + 7aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{512b^{9/2}e^{7/2}}
 \end{aligned}$$

output

$$\begin{aligned} & 1/512*(-a*e+b*d)^4*(-12*A*b*e+7*B*a*e+5*B*b*d)*(b*x+a)^{(1/2)}*(e*x+d)^{(1/2)} \\ & /b^4/e^3-1/768*(-a*e+b*d)^3*(-12*A*b*e+7*B*a*e+5*B*b*d)*(b*x+a)^{(3/2)}*(e*x \\ & +d)^{(1/2)}/b^4/e^2-1/192*(-a*e+b*d)^2*(-12*A*b*e+7*B*a*e+5*B*b*d)*(b*x+a)^{(5/2)} \\ & *(e*x+d)^{(1/2)}/b^4/e-1/96*(-a*e+b*d)*(-12*A*b*e+7*B*a*e+5*B*b*d)*(b*x+a)^{(5/2)} \\ & *(e*x+d)^{(3/2)}/b^3/e-1/60*(-12*A*b*e+7*B*a*e+5*B*b*d)*(b*x+a)^{(5/2)} \\ & *(e*x+d)^{(5/2)}/b^2/e+1/6*B*(b*x+a)^{(5/2)}*(e*x+d)^{(7/2)}/b/e-1/512*(-a*e+b*d)^5 \\ & *(-12*A*b*e+7*B*a*e+5*B*b*d)*\operatorname{arctanh}(e^{(1/2)}*(b*x+a)^{(1/2)}/(e*x+d)^{(1/2)})/b^{(9/2)}/e^{(7/2)} \end{aligned}$$

Mathematica [A] (verified)

Time = 1.01 (sec) , antiderivative size = 432, normalized size of antiderivative = 1.21

$$\begin{aligned} & \int (a + bx)^{3/2} (A + Bx) (d + ex)^{5/2} dx = \frac{\sqrt{a + bx} \sqrt{d + ex} (-105a^5 B e^5 + 5a^4 b e^4 (83Bd + 36Ae + 14Bex) - 2a^3 b^2 e^3 (60Ae(7d + ex) + \\ & (bd - ae)^5 (-5bBd + 12Abe - 7aBe) \operatorname{arctanh}\left(\frac{\sqrt{b} \sqrt{d+ex}}{\sqrt{e} \sqrt{a+bx}}\right))}{512b^{9/2} e^{7/2}} \end{aligned}$$

input

Integrate[(a + b*x)^(3/2)*(A + B*x)*(d + e*x)^(5/2),x]

output

$$\begin{aligned} & (\operatorname{Sqrt}[a + b*x]*\operatorname{Sqrt}[d + e*x]*(-105*a^5*B*e^5 + 5*a^4*b*e^4*(83*B*d + 36*A* \\ & e + 14*B*e*x) - 2*a^3*b^2*e^3*(60*A*e*(7*d + e*x) + B*(273*d^2 + 136*d*e*x \\ & + 28*e^2*x^2)) + 6*a^2*b^3*e^2*(4*A*e*(64*d^2 + 23*d*e*x + 4*e^2*x^2) + B \\ & *(25*d^3 + 58*d^2*e*x + 36*d*e^2*x^2 + 8*e^3*x^3)) + a*b^4*e*(24*A*e*(35*d^3 \\ & + 233*d^2*e*x + 256*d*e^2*x^2 + 88*e^3*x^3) + B*(-245*d^4 + 160*d^3*e*x \\ & + 3384*d^2*e^2*x^2 + 4448*d*e^3*x^3 + 1664*e^4*x^4)) + b^5*(12*A*e*(-15*d^4 \\ & + 10*d^3*e*x + 248*d^2*e^2*x^2 + 336*d*e^3*x^3 + 128*e^4*x^4) + 5*B*(15 \\ & *d^5 - 10*d^4*e*x + 8*d^3*e^2*x^2 + 432*d^2*e^3*x^3 + 640*d*e^4*x^4 + 256* \\ & e^5*x^5)))/(7680*b^4*e^3) + ((b*d - a*e)^5*(-5*b*B*d + 12*A*b*e - 7*a*B*e) \\ &)*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*\operatorname{Sqrt}[d + e*x])/(\operatorname{Sqrt}[e]*\operatorname{Sqrt}[a + b*x])]/(512*b^{(9/2)}*e^{(7/2)}) \end{aligned}$$

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 293, normalized size of antiderivative = 0.82, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {90, 60, 60, 60, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + bx)^{3/2}(A + Bx)(d + ex)^{5/2} dx \\
 & \quad \downarrow \text{90} \\
 & \frac{B(a + bx)^{5/2}(d + ex)^{7/2}}{6be} - \frac{(7aBe - 12Abe + 5bBd) \int (a + bx)^{3/2}(d + ex)^{5/2} dx}{12be} \\
 & \quad \downarrow \text{60} \\
 & \frac{B(a + bx)^{5/2}(d + ex)^{7/2}}{6be} - \\
 & \frac{(7aBe - 12Abe + 5bBd) \left(\frac{(bd-ae) \int (a+bx)^{3/2}(d+ex)^{3/2} dx}{2b} + \frac{(a+bx)^{5/2}(d+ex)^{5/2}}{5b} \right)}{12be} \\
 & \quad \downarrow \text{60} \\
 & \frac{B(a + bx)^{5/2}(d + ex)^{7/2}}{6be} - \\
 & \frac{(7aBe - 12Abe + 5bBd) \left(\frac{(bd-ae) \left(\frac{3(bd-ae) \int (a+bx)^{3/2} \sqrt{d+ex} dx}{8b} + \frac{(a+bx)^{5/2}(d+ex)^{3/2}}{4b} \right)}{2b} + \frac{(a+bx)^{5/2}(d+ex)^{5/2}}{5b} \right)}{12be} \\
 & \quad \downarrow \text{60} \\
 & \frac{B(a + bx)^{5/2}(d + ex)^{7/2}}{6be} - \\
 & \frac{(7aBe - 12Abe + 5bBd) \left(\frac{(bd-ae) \left(\frac{3(bd-ae) \left(\frac{(bd-ae) \int (a+bx)^{3/2} dx}{6b \sqrt{d+ex}} + \frac{(a+bx)^{5/2} \sqrt{d+ex}}{3b} \right)}{8b} + \frac{(a+bx)^{5/2}(d+ex)^{3/2}}{4b} \right)}{2b} + \frac{(a+bx)^{5/2}(d+ex)^{5/2}}{5b} \right)}{12be} \\
 & \quad \downarrow \text{60} \\
 & \frac{B(a + bx)^{5/2}(d + ex)^{7/2}}{6be} - \\
 & \frac{(7aBe - 12Abe + 5bBd) \left(\frac{(bd-ae) \left(\frac{3(bd-ae) \left(\frac{(bd-ae) \int (a+bx)^{3/2} dx}{6b \sqrt{d+ex}} + \frac{(a+bx)^{5/2} \sqrt{d+ex}}{3b} \right)}{8b} + \frac{(a+bx)^{5/2}(d+ex)^{3/2}}{4b} \right)}{2b} + \frac{(a+bx)^{5/2}(d+ex)^{5/2}}{5b} \right)}{12be}
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow 60 \\
 \frac{B(a+bx)^{5/2}(d+ex)^{7/2}}{6be} - \\
 \left((bd-ae) \left(\frac{3(bd-ae) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \int \frac{\sqrt{a+bx}}{\sqrt{d+ex}} dx}{4e} \right) + \frac{(a+bx)^{5/2}\sqrt{d+ex}}{3b}}{6b} \right) + \frac{(a+bx)^{5/2}(d+ex)}{4b} \right) \\
 (7aBe - 12Abe + 5bBd) \frac{\hspace{15em}}{2b} \\
 \hline
 \hspace{15em} 12be \\
 \downarrow 60
 \end{array}$$

$$\frac{B(a+bx)^{5/2}(d+ex)^{7/2}}{6be} - \left(\frac{(bd-ae)}{3(bd-ae)} \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae)}{6b} \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{4e} \right) \right) + (a+bx) \right) \frac{(bd-ae)}{8b} - \frac{(7aBe - 12Abe + 5bBd)}{2b}$$

12be

↓ 66

$$\begin{aligned}
 & \frac{B(a+bx)^{5/2}(d+ex)^{7/2}}{6be} - \\
 & \left(\begin{aligned}
 & \left(\begin{aligned}
 & \left(\begin{aligned}
 & \frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae)}{4e} \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{b-\frac{e(a+bx)}{d+ex}} d\sqrt{a+bx}}{\sqrt{d+ex}} \right) \right) \\
 & \frac{3(bd-ae)}{6b} \\
 & \frac{(bd-ae)}{8b} \\
 & \frac{(7aBe - 12Abe + 5bBd)}{2b}
 \end{aligned} \right) \\
 & \frac{12be}{12be}
 \end{aligned} \right)
 \end{aligned}
 \end{aligned}
 \end{aligned}$$

$$\begin{array}{l}
 \downarrow 221 \\
 \frac{B(a+bx)^{5/2}(d+ex)^{7/2}}{6be} - \\
 \left(\begin{array}{l}
 \frac{(bd-ae)}{3(bd-ae)} \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae)}{4e} \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{\sqrt{be}^{3/2}} \right) \right) \\
 \frac{(bd-ae)}{8b} \\
 \frac{(7aBe - 12Abe + 5bBd)}{2b}
 \end{array} \right) \\
 \hline
 12be
 \end{array}$$

input `Int[(a + b*x)^(3/2)*(A + B*x)*(d + e*x)^(5/2),x]`

output

$$\frac{(B(a + bx)^{5/2}(d + ex)^{7/2})/(6b^2e) - ((5b^2d - 12Abe + 7A^2e) * ((a + bx)^{5/2}(d + ex)^{5/2})/(5b) + ((bd - ae) * ((a + bx)^{5/2}(d + ex)^{3/2})/(4b) + (3(bd - ae) * ((a + bx)^{5/2} \sqrt{d + ex})) / (3b) + ((bd - ae) * ((a + bx)^{3/2} \sqrt{d + ex})) / (2e) - (3(bd - ae) * (\sqrt{a + bx} \sqrt{d + ex})) / e - ((bd - ae) * \operatorname{ArcTanh}[\sqrt{e} \sqrt{a + bx}] / (\sqrt{b} \sqrt{d + ex})) / (\sqrt{b} e^{3/2})) / (4e)) / (6b)) / (8b)) / (2b)) / (12b^2e)}$$

Definitions of rubi rules used

rule 60

$$\operatorname{Int}[(a + bx)^m (c + dx)^n, x] \rightarrow \operatorname{Simp}[(a + bx)^{m+1} (c + dx)^n / (b(m+n+1)), x] + \operatorname{Simp}[n(b^2c - a^2d) / (b(m+n+1)) \operatorname{Int}[(a + bx)^m (c + dx)^{n-1}, x], x] /;$$

FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m+n+1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m-n, 0]))) && !ILtQ[m+n+2, 0] && IntLinearQ[a, b, c, d, m, n, x]

rule 66

$$\operatorname{Int}[1/(\sqrt{a + bx} \sqrt{c + dx}), x] \rightarrow \operatorname{Simp}[2 \operatorname{Subst}[\operatorname{Int}[1/(b - dx^2), x], x, \sqrt{a + bx} / \sqrt{c + dx}], x] /;$$

FreeQ[{a, b, c, d}, x] && !GtQ[c - a(d/b), 0]

rule 90

$$\operatorname{Int}[(a + bx)^m (c + dx)^n (e + fx)^p, x] \rightarrow \operatorname{Simp}[b(c + dx)^{n+1} (e + fx)^{p+1} / (d^2 f (n + p + 2)), x] + \operatorname{Simp}[(a d^2 f (n + p + 2) - b(d^2 e (n + 1) + c f (p + 1))] / (d^2 f (n + p + 2)) \operatorname{Int}[(c + dx)^n (e + fx)^p, x], x] /;$$

FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]

rule 221

$$\operatorname{Int}[(a + bx)^{-2}, x] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2]/a) \operatorname{ArcTanh}[x / \operatorname{Rt}[-a/b, 2]], x] /;$$

FreeQ[{a, b}, x] && NegQ[a/b]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1847 vs. $2(308) = 616$.

Time = 0.27 (sec) , antiderivative size = 1848, normalized size of antiderivative = 5.16

method	result	size
default	Expression too large to display	1848

input `int((b*x+a)^(3/2)*(B*x+A)*(e*x+d)^(5/2),x,method=_RETURNVERBOSE)`

output

```
-1/15360*(b*x+a)^(1/2)*(e*x+d)^(1/2)*(240*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^3*b^2*e^5*x-240*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^5*d^3*e^2*x-140*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^4*b*e^5*x+100*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^5*d^4*e*x-8896*B*a*b^4*d*e^4*x^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-12288*A*a*b^4*d*e^4*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-432*B*a^2*b^3*d*e^4*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-6768*B*a*b^4*d^2*e^3*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-1104*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^2*b^3*d*e^4*x-11184*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b^4*d^2*e^3*x+544*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^3*b^2*d*e^4*x-696*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^2*b^3*d^2*e^3*x-320*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b^4*d^3*e^2*x+900*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^5*d^4*e^2+450*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^5*b*d*e^5-2560*B*b^5*e^5*x^5*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-3072*A*b^5*e^5*x^4*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-4320*B*b^5*d^2*e^3*x^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+180*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^5*b*e^6-180*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^6*d^5*e+210*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^5*e^5-150*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^5*d^5+300*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 685 vs. $2(308) = 616$.

Time = 0.16 (sec) , antiderivative size = 1384, normalized size of antiderivative = 3.87

$$\int (a + bx)^{3/2}(A + Bx)(d + ex)^{5/2} dx = \text{Too large to display}$$

input `integrate((b*x+a)^(3/2)*(B*x+A)*(e*x+d)^(5/2),x, algorithm="fricas")`

output

```
[1/30720*(15*(5*B*b^6*d^6 - 6*(3*B*a*b^5 + 2*A*b^6)*d^5*e + 15*(B*a^2*b^4
+ 4*A*a*b^5)*d^4*e^2 + 20*(B*a^3*b^3 - 6*A*a^2*b^4)*d^3*e^3 - 15*(3*B*a^4*
b^2 - 8*A*a^3*b^3)*d^2*e^4 + 30*(B*a^5*b - 2*A*a^4*b^2)*d*e^5 - (7*B*a^6 -
12*A*a^5*b)*e^6)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*
e^2 - 4*(2*b*e*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x + a)*sqrt(e*x + d) + 8*(b
^2*d*e + a*b*e^2)*x) + 4*(1280*B*b^6*e^6*x^5 + 75*B*b^6*d^5*e - 5*(49*B*a*
b^5 + 36*A*b^6)*d^4*e^2 + 30*(5*B*a^2*b^4 + 28*A*a*b^5)*d^3*e^3 - 6*(91*B*
a^3*b^3 - 256*A*a^2*b^4)*d^2*e^4 + 5*(83*B*a^4*b^2 - 168*A*a^3*b^3)*d*e^5
- 15*(7*B*a^5*b - 12*A*a^4*b^2)*e^6 + 128*(25*B*b^6*d*e^5 + (13*B*a*b^5 +
12*A*b^6)*e^6)*x^4 + 16*(135*B*b^6*d^2*e^4 + 2*(139*B*a*b^5 + 126*A*b^6)*d
*e^5 + 3*(B*a^2*b^4 + 44*A*a*b^5)*e^6)*x^3 + 8*(5*B*b^6*d^3*e^3 + 3*(141*B
*a*b^5 + 124*A*b^6)*d^2*e^4 + 3*(9*B*a^2*b^4 + 256*A*a*b^5)*d*e^5 - (7*B*a
^3*b^3 - 12*A*a^2*b^4)*e^6)*x^2 - 2*(25*B*b^6*d^4*e^2 - 20*(4*B*a*b^5 + 3*
A*b^6)*d^3*e^3 - 6*(29*B*a^2*b^4 + 466*A*a*b^5)*d^2*e^4 + 4*(34*B*a^3*b^3
- 69*A*a^2*b^4)*d*e^5 - 5*(7*B*a^4*b^2 - 12*A*a^3*b^3)*e^6)*x)*sqrt(b*x +
a)*sqrt(e*x + d)/(b^5*e^4), 1/15360*(15*(5*B*b^6*d^6 - 6*(3*B*a*b^5 + 2*A*
b^6)*d^5*e + 15*(B*a^2*b^4 + 4*A*a*b^5)*d^4*e^2 + 20*(B*a^3*b^3 - 6*A*a^2
*b^4)*d^3*e^3 - 15*(3*B*a^4*b^2 - 8*A*a^3*b^3)*d^2*e^4 + 30*(B*a^5*b - 2*A
*a^4*b^2)*d*e^5 - (7*B*a^6 - 12*A*a^5*b)*e^6)*sqrt(-b*e)*arctan(1/2*(2*b*e
*x + b*d + a*e)*sqrt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*e^2*x^2 + a...
```

Sympy [F]

$$\int (a + bx)^{3/2} (A + Bx) (d + ex)^{5/2} dx = \int (A + Bx) (a + bx)^{3/2} (d + ex)^{5/2} dx$$

input `integrate((b*x+a)**(3/2)*(B*x+A)*(e*x+d)**(5/2),x)`

output `Integral((A + B*x)*(a + b*x)**(3/2)*(d + e*x)**(5/2), x)`

Maxima [F(-2)]

Exception generated.

$$\int (a + bx)^{3/2} (A + Bx) (d + ex)^{5/2} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(3/2)*(B*x+A)*(e*x+d)^(5/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4461 vs. 2(308) = 616.

Time = 0.68 (sec) , antiderivative size = 4461, normalized size of antiderivative = 12.46

$$\int (a + bx)^{3/2} (A + Bx) (d + ex)^{5/2} dx = \text{Too large to display}$$

input `integrate((b*x+a)^(3/2)*(B*x+A)*(e*x+d)^(5/2),x, algorithm="giac")`

output

```

1/7680*(40*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*(b*x + a)*(4*(b*x + a)*
(6*(b*x + a)/b^3 + (b^12*d*e^5 - 25*a*b^11*e^6)/(b^14*e^6)) - (5*b^13*d^2*
e^4 + 14*a*b^12*d*e^5 - 163*a^2*b^11*e^6)/(b^14*e^6)) + 3*(5*b^14*d^3*e^3
+ 9*a*b^13*d^2*e^4 + 15*a^2*b^12*d*e^5 - 93*a^3*b^11*e^6)/(b^14*e^6))*sqrt
(b*x + a) + 3*(5*b^4*d^4 + 4*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2 + 20*a^3*b*d*
e^3 - 35*a^4*e^4)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a
)*b*e - a*b*e)))/(sqrt(b*e)*b^2*e^3))*B*d^2*abs(b) - 7680*((b^2*d - a*b*e)
*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/
sqrt(b*e) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*sqrt(b*x + a))*A*a^2*d^2*a
bs(b)/b^2 + 80*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*(b*x + a)*(4*(b*x +
a)*(6*(b*x + a)/b^3 + (b^12*d*e^5 - 25*a*b^11*e^6)/(b^14*e^6)) - (5*b^13*
d^2*e^4 + 14*a*b^12*d*e^5 - 163*a^2*b^11*e^6)/(b^14*e^6)) + 3*(5*b^14*d^3*
e^3 + 9*a*b^13*d^2*e^4 + 15*a^2*b^12*d*e^5 - 93*a^3*b^11*e^6)/(b^14*e^6))*
sqrt(b*x + a) + 3*(5*b^4*d^4 + 4*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2 + 20*a^3*
b*d*e^3 - 35*a^4*e^4)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x
+ a)*b*e - a*b*e)))/(sqrt(b*e)*b^2*e^3))*A*d*e*abs(b) + 160*(sqrt(b^2*d +
(b*x + a)*b*e - a*b*e)*(2*(b*x + a)*(4*(b*x + a)*(6*(b*x + a)/b^3 + (b^12
*d*e^5 - 25*a*b^11*e^6)/(b^14*e^6)) - (5*b^13*d^2*e^4 + 14*a*b^12*d*e^5 -
163*a^2*b^11*e^6)/(b^14*e^6)) + 3*(5*b^14*d^3*e^3 + 9*a*b^13*d^2*e^4 + 15*
a^2*b^12*d*e^5 - 93*a^3*b^11*e^6)/(b^14*e^6))*sqrt(b*x + a) + 3*(5*b^4*...

```

Mupad [F(-1)]

Timed out.

$$\int (a + bx)^{3/2} (A + Bx) (d + ex)^{5/2} dx = \int (A + Bx) (a + bx)^{3/2} (d + ex)^{5/2} dx$$

input

```
int((A + B*x)*(a + b*x)^(3/2)*(d + e*x)^(5/2),x)
```

output

```
int((A + B*x)*(a + b*x)^(3/2)*(d + e*x)^(5/2), x)
```

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 860, normalized size of antiderivative = 2.40

$$\int (a + bx)^{3/2}(A + Bx)(d + ex)^{5/2} dx = \text{Too large to display}$$

input `int((b*x+a)^(3/2)*(B*x+A)*(e*x+d)^(5/2),x)`

output

```
(15*sqrt(d + e*x)*sqrt(a + b*x)*a**5*b*e**6 - 85*sqrt(d + e*x)*sqrt(a + b*x)*a**4*b**2*d*e**5 - 10*sqrt(d + e*x)*sqrt(a + b*x)*a**4*b**2*e**6*x + 198*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b**3*d**2*e**4 + 56*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b**3*d*e**5*x + 8*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b**3*e**6*x**2 + 198*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**4*d**3*e**3 + 1188*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**4*d**2*e**4*x + 1272*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**4*d*e**5*x**2 + 432*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**4*e**6*x**3 - 85*sqrt(d + e*x)*sqrt(a + b*x)*a*b**5*d**4*e**2 + 56*sqrt(d + e*x)*sqrt(a + b*x)*a*b**5*d**3*e**3*x + 1272*sqrt(d + e*x)*sqrt(a + b*x)*a*b**5*d**2*e**4*x**2 + 1696*sqrt(d + e*x)*sqrt(a + b*x)*a*b**5*d*e**5*x**3 + 640*sqrt(d + e*x)*sqrt(a + b*x)*a*b**5*e**6*x**4 + 15*sqrt(d + e*x)*sqrt(a + b*x)*b**6*d**5*e - 10*sqrt(d + e*x)*sqrt(a + b*x)*b**6*d**4*e**2*x + 8*sqrt(d + e*x)*sqrt(a + b*x)*b**6*d**3*e**3*x**2 + 432*sqrt(d + e*x)*sqrt(a + b*x)*b**6*d**2*e**4*x**3 + 640*sqrt(d + e*x)*sqrt(a + b*x)*b**6*d*e**5*x**4 + 256*sqrt(d + e*x)*sqrt(a + b*x)*b**6*e**6*x**5 - 15*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**6*e**6 + 90*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**5*b*d*e**5 - 225*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**4*b**2*d**2*e**4 + 300*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))...
```

3.183 $\int (a + bx)^{3/2}(A + Bx)(d + ex)^{3/2} dx$

Optimal result	1666
Mathematica [A] (verified)	1667
Rubi [A] (verified)	1667
Maple [B] (verified)	1671
Fricas [B] (verification not implemented)	1672
Sympy [F]	1673
Maxima [F(-2)]	1674
Giac [B] (verification not implemented)	1674
Mupad [F(-1)]	1675
Reduce [B] (verification not implemented)	1676

Optimal result

Integrand size = 24, antiderivative size = 294

$$\int (a + bx)^{3/2}(A + Bx)(d + ex)^{3/2} dx = \frac{3(bd - ae)^3(bBd - 2Abe + aBe)\sqrt{a + bx}\sqrt{d + ex}}{128b^3e^3} - \frac{(bd - ae)^2(bBd - 2Abe + aBe)(a + bx)^{3/2}\sqrt{d + ex}}{64b^3e^2} - \frac{(bd - ae)(bBd - 2Abe + aBe)(a + bx)^{5/2}\sqrt{d + ex}}{16b^3e} - \frac{(bBd - 2Abe + aBe)(a + bx)^{5/2}(d + ex)^{3/2}}{8b^2e} + \frac{B(a + bx)^{5/2}(d + ex)^{5/2}}{5be} - \frac{3(bd - ae)^4(bBd - 2Abe + aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{128b^{7/2}e^{7/2}}$$

output

```
3/128*(-a*e+b*d)^3*(-2*A*b*e+B*a*e+B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(1/2)/b^3/e^3-1/64*(-a*e+b*d)^2*(-2*A*b*e+B*a*e+B*b*d)*(b*x+a)^(3/2)*(e*x+d)^(1/2)/b^3/e^2-1/16*(-a*e+b*d)*(-2*A*b*e+B*a*e+B*b*d)*(b*x+a)^(5/2)*(e*x+d)^(1/2)/b^3/e-1/8*(-2*A*b*e+B*a*e+B*b*d)*(b*x+a)^(5/2)*(e*x+d)^(3/2)/b^2/e+1/5*B*(b*x+a)^(5/2)*(e*x+d)^(5/2)/b/e-3/128*(-a*e+b*d)^4*(-2*A*b*e+B*a*e+B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)^(1/2))/b^(7/2)/e^(7/2)
```

Mathematica [A] (verified)

Time = 0.72 (sec) , antiderivative size = 319, normalized size of antiderivative = 1.09

$$\int (a + bx)^{3/2} (A + Bx) (d + ex)^{3/2} dx = \frac{\sqrt{a + bx} \sqrt{d + ex} (15a^4 B e^4 - 10a^3 b e^3 (4Bd + 3Ae + Bex) + 2a^2 b^2 e^2 (5Ae(11d + 2ex) + B(9d + ex))) + 3(bd - ae)^4 (bBd - 2Abe + aBe) \operatorname{arctanh}\left(\frac{\sqrt{b} \sqrt{d + ex}}{\sqrt{e} \sqrt{a + bx}}\right)}{128b^{7/2} e^{7/2}}$$

input `Integrate[(a + b*x)^(3/2)*(A + B*x)*(d + e*x)^(3/2),x]`

output `(Sqrt[a + b*x]*Sqrt[d + e*x]*(15*a^4*B*e^4 - 10*a^3*b*e^3*(4*B*d + 3*A*e + B*e*x) + 2*a^2*b^2*e^2*(5*A*e*(11*d + 2*e*x) + B*(9*d^2 + 13*d*e*x + 4*e^2*x^2)) + 2*a*b^3*e*(5*A*e*(11*d^2 + 44*d*e*x + 24*e^2*x^2) + B*(-20*d^3 + 13*d^2*e*x + 136*d*e^2*x^2 + 88*e^3*x^3)) + b^4*(10*A*e*(-3*d^3 + 2*d^2*e*x + 24*d*e^2*x^2 + 16*e^3*x^3) + B*(15*d^4 - 10*d^3*e*x + 8*d^2*e^2*x^2 + 176*d*e^3*x^3 + 128*e^4*x^4)))/(640*b^3*e^3) - (3*(b*d - a*e)^4*(b*B*d - 2*A*b*e + a*B*e)*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/(Sqrt[e]*Sqrt[a + b*x])])/(128*b^(7/2)*e^(7/2))`

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 252, normalized size of antiderivative = 0.86, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.292$, Rules used = {90, 60, 60, 60, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^{3/2} (A + Bx) (d + ex)^{3/2} dx$$

↓ 90

$$\frac{(2Abe - B(ae + bd)) \int (a + bx)^{3/2} (d + ex)^{3/2} dx}{2be} + \frac{B(a + bx)^{5/2} (d + ex)^{5/2}}{5be}$$

$$\begin{aligned}
 & \downarrow 60 \\
 & \frac{(2Abe - B(ae + bd)) \left(\frac{3(bd-ae) \int (a+bx)^{3/2} \sqrt{d+ex} dx}{8b} + \frac{(a+bx)^{5/2} (d+ex)^{3/2}}{4b} \right)}{B(a+bx)^{5/2} (d+ex)^{5/2}} + \\
 & \qquad \qquad \qquad \frac{2be}{5be} \\
 & \downarrow 60 \\
 & \frac{(2Abe - B(ae + bd)) \left(\frac{3(bd-ae) \left(\frac{(bd-ae) \int \frac{(a+bx)^{3/2}}{\sqrt{d+ex}} dx}{6b} + \frac{(a+bx)^{5/2} \sqrt{d+ex}}{3b} \right)}{8b} + \frac{(a+bx)^{5/2} (d+ex)^{3/2}}{4b} \right)}{B(a+bx)^{5/2} (d+ex)^{5/2}} + \\
 & \qquad \qquad \qquad \frac{2be}{5be} \\
 & \downarrow 60 \\
 & \frac{(2Abe - B(ae + bd)) \left(\frac{3(bd-ae) \left(\frac{(bd-ae) \left(\frac{(a+bx)^{3/2} \sqrt{d+ex}}{2e} - \frac{3(bd-ae) \int \frac{\sqrt{a+bx}}{\sqrt{d+ex}} dx}{4e} \right)}{6b} + \frac{(a+bx)^{5/2} \sqrt{d+ex}}{3b} \right)}{8b} + \frac{(a+bx)^{5/2} (d+ex)^{3/2}}{4b} \right)}{B(a+bx)^{5/2} (d+ex)^{5/2}} + \\
 & \qquad \qquad \qquad \frac{2be}{5be} \\
 & \downarrow 60
 \end{aligned}$$

$$(2Abe - B(ae + bd)) \left(\frac{3(bd-ae) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{2e} \right)}{4e} \right)}{6b} + \frac{(a+bx)^{5/2}\sqrt{d+ex}}{3b} \right)}{8b}$$

$$\frac{B(a+bx)^{5/2}(d+ex)^{5/2}}{5be} \quad 2be$$

↓ 66

$$(2Abe - B(ae + bd)) \left(\frac{3(bd-ae) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d\sqrt{a+bx}}{e} \right)}{4e} \right)}{6b} + \frac{(a+bx)^{5/2}\sqrt{d+ex}}{3b} \right)}{8b}$$

$$\frac{B(a+bx)^{5/2}(d+ex)^{5/2}}{5be} \quad 2be$$

221

$$\begin{aligned}
 & \left(\frac{(bd-ae) \left(\frac{(a+bx)^{3/2} \sqrt{d+ex}}{2e} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx} \sqrt{d+ex}}{e} - \frac{(bd-ae) \operatorname{arctanh} \left(\frac{\sqrt{e} \sqrt{a+bx}}{\sqrt{b} \sqrt{d+ex}} \right)}{\sqrt{b} e^{3/2}} \right)}{4e} \right)}{6b} + \frac{(a+bx)^{5/2} \sqrt{d+ex}}{3b} \right) \\
 & \frac{(2Abe - B(ae + bd))}{8b} \\
 & \frac{B(a+bx)^{5/2} (d+ex)^{5/2}}{5be}
 \end{aligned}$$

```
input Int[(a + b*x)^(3/2)*(A + B*x)*(d + e*x)^(3/2),x]
```

```
output (B*(a + b*x)^(5/2)*(d + e*x)^(5/2))/(5*b*e) + ((2*A*b*e - B*(b*d + a*e))*
((a + b*x)^(5/2)*(d + e*x)^(3/2))/(4*b) + (3*(b*d - a*e)*((a + b*x)^(5/2)
*Sqrt[d + e*x])/(3*b) + ((b*d - a*e)*((a + b*x)^(3/2)*Sqrt[d + e*x])/(2*
e) - (3*(b*d - a*e)*((Sqrt[a + b*x]*Sqrt[d + e*x])/e - ((b*d - a*e)*ArcTanh
[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(Sqrt[b]*e^(3/2))))/(4*
e)))/(6*b))/(8*b))/(2*b*e)
```

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1371 vs. $2(250) = 500$.

Time = 0.27 (sec) , antiderivative size = 1372, normalized size of antiderivative = 4.67

method	result	size
default	Expression too large to display	1372

input `int((b*x+a)^(3/2)*(B*x+A)*(e*x+d)^(3/2),x,method=_RETURNVERBOSE)`

output

```

1/1280*(b*x+a)^(1/2)*(e*x+d)^(1/2)*(256*B*b^4*e^4*x^4*((e*x+d)*(b*x+a))^(1
/2)*(b*e)^(1/2)+320*A*b^4*e^4*x^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+352*
B*a*b^3*e^4*x^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+352*B*b^4*d*e^3*x^3*((
e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+480*A*a*b^3*e^4*x^2*((e*x+d)*(b*x+a))^(1
/2)*(b*e)^(1/2)+220*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b^3*d^2*e^2-80
*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^3*b*d*e^3+36*B*((e*x+d)*(b*x+a))^(
1/2)*(b*e)^(1/2)*a^2*b^2*d^2*e^2-80*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)
*a*b^3*d^3*e+30*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^4*e^4+30*B*((e*x+d
)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^4*d^4+544*B*a*b^3*d*e^3*x^2*((e*x+d)*(b*x+a
))^(1/2)*(b*e)^(1/2)+880*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b^3*d*e^3
*x+52*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^2*b^2*d*e^3*x+52*B*((e*x+d)*
(b*x+a))^(1/2)*(b*e)^(1/2)*a*b^3*d^2*e^2*x+30*A*ln(1/2*(2*b*e*x+2*((e*x+d)
*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^4*b*e^5+30*A*ln(1/2*(2
*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^5*d^4
*e+180*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b
*e)^(1/2))*a^2*b^3*d^2*e^3-120*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)
*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^4*d^3*e^2+45*B*ln(1/2*(2*b*e*x+2*((
e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^4*b*d*e^4-30*B*ln
(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))
*a^3*b^2*d^2*e^3-30*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 511 vs. $2(250) = 500$.

Time = 0.16 (sec) , antiderivative size = 1036, normalized size of antiderivative = 3.52

$$\int (a + bx)^{3/2} (A + Bx)(d + ex)^{3/2} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^(3/2)*(B*x+A)*(e*x+d)^(3/2),x, algorithm="fricas")
```

output

```

[-1/2560*(15*(B*b^5*d^5 - (3*B*a*b^4 + 2*A*b^5)*d^4*e + 2*(B*a^2*b^3 + 4*A
*a*b^4)*d^3*e^2 + 2*(B*a^3*b^2 - 6*A*a^2*b^3)*d^2*e^3 - (3*B*a^4*b - 8*A*a
^3*b^2)*d*e^4 + (B*a^5 - 2*A*a^4*b)*e^5)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2
*d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*b*e*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x +
a)*sqrt(e*x + d) + 8*(b^2*d*e + a*b*e^2)*x) - 4*(128*B*b^5*e^5*x^4 + 15*B*
b^5*d^4*e - 10*(4*B*a*b^4 + 3*A*b^5)*d^3*e^2 + 2*(9*B*a^2*b^3 + 55*A*a*b^4
)*d^2*e^3 - 10*(4*B*a^3*b^2 - 11*A*a^2*b^3)*d*e^4 + 15*(B*a^4*b - 2*A*a^3*
b^2)*e^5 + 16*(11*B*b^5*d*e^4 + (11*B*a*b^4 + 10*A*b^5)*e^5)*x^3 + 8*(B*b^
5*d^2*e^3 + 2*(17*B*a*b^4 + 15*A*b^5)*d*e^4 + (B*a^2*b^3 + 30*A*a*b^4)*e^5
)*x^2 - 2*(5*B*b^5*d^3*e^2 - (13*B*a*b^4 + 10*A*b^5)*d^2*e^3 - (13*B*a^2*b
^3 + 220*A*a*b^4)*d*e^4 + 5*(B*a^3*b^2 - 2*A*a^2*b^3)*e^5)*x)*sqrt(b*x + a
)*sqrt(e*x + d))/(b^4*e^4), 1/1280*(15*(B*b^5*d^5 - (3*B*a*b^4 + 2*A*b^5)*
d^4*e + 2*(B*a^2*b^3 + 4*A*a*b^4)*d^3*e^2 + 2*(B*a^3*b^2 - 6*A*a^2*b^3)*d^
2*e^3 - (3*B*a^4*b - 8*A*a^3*b^2)*d*e^4 + (B*a^5 - 2*A*a^4*b)*e^5)*sqrt(-b
*e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d
))/(b^2*e^2*x^2 + a*b*d*e + (b^2*d*e + a*b*e^2)*x)) + 2*(128*B*b^5*e^5*x^4
+ 15*B*b^5*d^4*e - 10*(4*B*a*b^4 + 3*A*b^5)*d^3*e^2 + 2*(9*B*a^2*b^3 + 55*
A*a*b^4)*d^2*e^3 - 10*(4*B*a^3*b^2 - 11*A*a^2*b^3)*d*e^4 + 15*(B*a^4*b - 2
*A*a^3*b^2)*e^5 + 16*(11*B*b^5*d*e^4 + (11*B*a*b^4 + 10*A*b^5)*e^5)*x^3 +
8*(B*b^5*d^2*e^3 + 2*(17*B*a*b^4 + 15*A*b^5)*d*e^4 + (B*a^2*b^3 + 30*A*...

```

Sympy [F]

$$\int (a + bx)^{3/2} (A + Bx) (d + ex)^{3/2} dx = \int (A + Bx) (a + bx)^{\frac{3}{2}} (d + ex)^{\frac{3}{2}} dx$$

input

```
integrate((b*x+a)**(3/2)*(B*x+A)*(e*x+d)**(3/2),x)
```

output

```
Integral((A + B*x)*(a + b*x)**(3/2)*(d + e*x)**(3/2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int (a + bx)^{3/2}(A + Bx)(d + ex)^{3/2} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(3/2)*(B*x+A)*(e*x+d)^(3/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2484 vs. 2(250) = 500.

Time = 0.45 (sec) , antiderivative size = 2484, normalized size of antiderivative = 8.45

$$\int (a + bx)^{3/2}(A + Bx)(d + ex)^{3/2} dx = \text{Too large to display}$$

input `integrate((b*x+a)^(3/2)*(B*x+A)*(e*x+d)^(3/2),x, algorithm="giac")`

output

```

1/1920*(10*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*(b*x + a)*(4*(b*x + a)*
(6*(b*x + a)/b^3 + (b^12*d*e^5 - 25*a*b^11*e^6)/(b^14*e^6)) - (5*b^13*d^2*
e^4 + 14*a*b^12*d*e^5 - 163*a^2*b^11*e^6)/(b^14*e^6)) + 3*(5*b^14*d^3*e^3
+ 9*a*b^13*d^2*e^4 + 15*a^2*b^12*d*e^5 - 93*a^3*b^11*e^6)/(b^14*e^6))*sqrt
(b*x + a) + 3*(5*b^4*d^4 + 4*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2 + 20*a^3*b*d*
e^3 - 35*a^4*e^4)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a
)*b*e - a*b*e)))/(sqrt(b*e)*b^2*e^3))*B*d*abs(b) - 1920*((b^2*d - a*b*e)*l
og(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/sq
rt(b*e) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*sqrt(b*x + a))*A*a^2*d*abs(b
)/b^2 + 10*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*(b*x + a)*(4*(b*x + a)*
(6*(b*x + a)/b^3 + (b^12*d*e^5 - 25*a*b^11*e^6)/(b^14*e^6)) - (5*b^13*d^2*
e^4 + 14*a*b^12*d*e^5 - 163*a^2*b^11*e^6)/(b^14*e^6)) + 3*(5*b^14*d^3*e^3
+ 9*a*b^13*d^2*e^4 + 15*a^2*b^12*d*e^5 - 93*a^3*b^11*e^6)/(b^14*e^6))*sqrt
(b*x + a) + 3*(5*b^4*d^4 + 4*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2 + 20*a^3*b*d*
e^3 - 35*a^4*e^4)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a
)*b*e - a*b*e)))/(sqrt(b*e)*b^2*e^3))*A*e*abs(b) + 20*(sqrt(b^2*d + (b*x +
a)*b*e - a*b*e)*(2*(b*x + a)*(4*(b*x + a)*(6*(b*x + a)/b^3 + (b^12*d*e^5
- 25*a*b^11*e^6)/(b^14*e^6)) - (5*b^13*d^2*e^4 + 14*a*b^12*d*e^5 - 163*a^2
*b^11*e^6)/(b^14*e^6)) + 3*(5*b^14*d^3*e^3 + 9*a*b^13*d^2*e^4 + 15*a^2*b^1
2*d*e^5 - 93*a^3*b^11*e^6)/(b^14*e^6))*sqrt(b*x + a) + 3*(5*b^4*d^4 + 4...

```

Mupad [F(-1)]

Timed out.

$$\int (a + bx)^{3/2} (A + Bx) (d + ex)^{3/2} dx = \int (A + Bx) (a + bx)^{3/2} (d + ex)^{3/2} dx$$

input

```
int((A + B*x)*(a + b*x)^(3/2)*(d + e*x)^(3/2),x)
```

output

```
int((A + B*x)*(a + b*x)^(3/2)*(d + e*x)^(3/2), x)
```


Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 651, normalized size of antiderivative = 2.21

$$\int (a + bx)^{3/2} (A + Bx) (d + ex)^{3/2} dx = \frac{-15\sqrt{ex + d}\sqrt{bx + a}a^4b e^5 + 70\sqrt{ex + d}\sqrt{bx + a}a^3b^2d e^4 + 10\sqrt{ex + d}\sqrt{bx + a}a^3b^2e^5x + \dots}{\dots}$$

input

```
int((b*x+a)^(3/2)*(B*x+A)*(e*x+d)^(3/2),x)
```

output

```
( - 15*sqrt(d + e*x)*sqrt(a + b*x)*a**4*b*e**5 + 70*sqrt(d + e*x)*sqrt(a +
b*x)*a**3*b**2*d*e**4 + 10*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b**2*e**5*x +
128*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**3*d**2*e**3 + 466*sqrt(d + e*x)*s
qrt(a + b*x)*a**2*b**3*d*e**4*x + 248*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**
3*e**5*x**2 - 70*sqrt(d + e*x)*sqrt(a + b*x)*a*b**4*d**3*e**2 + 46*sqrt(d
+ e*x)*sqrt(a + b*x)*a*b**4*d**2*e**3*x + 512*sqrt(d + e*x)*sqrt(a + b*x)*
a*b**4*d*e**4*x**2 + 336*sqrt(d + e*x)*sqrt(a + b*x)*a*b**4*e**5*x**3 + 15
*sqrt(d + e*x)*sqrt(a + b*x)*b**5*d**4*e - 10*sqrt(d + e*x)*sqrt(a + b*x)*
b**5*d**3*e**2*x + 8*sqrt(d + e*x)*sqrt(a + b*x)*b**5*d**2*e**3*x**2 + 176
*sqrt(d + e*x)*sqrt(a + b*x)*b**5*d*e**4*x**3 + 128*sqrt(d + e*x)*sqrt(a +
b*x)*b**5*e**5*x**4 + 15*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqr
t(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**5*e**5 - 75*sqrt(e)*sqrt(b)*log((s
qrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**4*b*d*e*
**4 + 150*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x
))/sqrt(a*e - b*d))*a**3*b**2*d**2*e**3 - 150*sqrt(e)*sqrt(b)*log((sqrt(e)
*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**2*b**3*d**3*e*
**2 + 75*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x
))/sqrt(a*e - b*d))*a*b**4*d**4*e - 15*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a
+ b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**5*d**5)/(640*b**3*e**4
)
```

3.184 $\int (a + bx)^{3/2} (A + Bx) \sqrt{d + ex} dx$

Optimal result	1677
Mathematica [A] (verified)	1678
Rubi [A] (verified)	1678
Maple [B] (verified)	1681
Fricas [A] (verification not implemented)	1682
Sympy [F]	1683
Maxima [F(-2)]	1683
Giac [B] (verification not implemented)	1684
Mupad [F(-1)]	1685
Reduce [B] (verification not implemented)	1685

Optimal result

Integrand size = 24, antiderivative size = 250

$$\int (a + bx)^{3/2} (A + Bx) \sqrt{d + ex} dx = \frac{(bd - ae)^2 (5bBd - 8Abe + 3aBe) \sqrt{a + bx} \sqrt{d + ex}}{64b^2 e^3} - \frac{(bd - ae) (5bBd - 8Abe + 3aBe) (a + bx)^{3/2} \sqrt{d + ex}}{96b^2 e^2} - \frac{(5bBd - 8Abe + 3aBe) (a + bx)^{5/2} \sqrt{d + ex}}{24b^2 e} + \frac{B(a + bx)^{5/2} (d + ex)^{3/2}}{4be} - \frac{(bd - ae)^3 (5bBd - 8Abe + 3aBe) \operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{64b^{5/2} e^{7/2}}$$

output

```
1/64*(-a*e+b*d)^2*(-8*A*b*e+3*B*a*e+5*B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(1/2)/b
^2/e^3-1/96*(-a*e+b*d)*(-8*A*b*e+3*B*a*e+5*B*b*d)*(b*x+a)^(3/2)*(e*x+d)^(1
/2)/b^2/e^2-1/24*(-8*A*b*e+3*B*a*e+5*B*b*d)*(b*x+a)^(5/2)*(e*x+d)^(1/2)/b^
2/e+1/4*B*(b*x+a)^(5/2)*(e*x+d)^(3/2)/b/e-1/64*(-a*e+b*d)^3*(-8*A*b*e+3*B*
a*e+5*B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)^(1/2))/b^(5/2)/
e^(7/2)
```

Mathematica [A] (verified)

Time = 0.59 (sec) , antiderivative size = 232, normalized size of antiderivative = 0.93

$$\int (a + bx)^{3/2} (A + Bx) \sqrt{d + ex} dx = \frac{\sqrt{a + bx} \sqrt{d + ex} (-9a^3 B e^3 + 3a^2 b e^2 (3Bd + 8Ae + 2Bex) + ab^2 e (16Ae(4d + 7ex) + B(-31d^2 + 20d e x + 72e^2 x^2)) + b^3 (8Ae(-3d^2 + 2d e x + 8e^2 x^2) + B(15d^3 - 10d^2 e x + 8d e^2 x^2 + 48e^3 x^3)))}{64b^{5/2} e^{7/2}} + \frac{(bd - ae)^3 (-5bBd + 8Abe - 3aBe) \operatorname{arctanh}\left(\frac{\sqrt{b} \sqrt{d+ex}}{\sqrt{e} \sqrt{a+bx}}\right)}{64b^{5/2} e^{7/2}}$$

input `Integrate[(a + b*x)^(3/2)*(A + B*x)*Sqrt[d + e*x], x]`

output `(Sqrt[a + b*x]*Sqrt[d + e*x]*(-9*a^3*B*e^3 + 3*a^2*b*e^2*(3*B*d + 8*A*e + 2*B*e*x) + a*b^2*e*(16*A*e*(4*d + 7*e*x) + B*(-31*d^2 + 20*d*e*x + 72*e^2*x^2)) + b^3*(8*A*e*(-3*d^2 + 2*d*e*x + 8*e^2*x^2) + B*(15*d^3 - 10*d^2*e*x + 8*d*e^2*x^2 + 48*e^3*x^3)))/(192*b^2*e^3) + ((b*d - a*e)^3*(-5*b*B*d + 8*A*b*e - 3*a*B*e)*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/(Sqrt[e]*Sqrt[a + b*x])])/(64*b^(5/2)*e^(7/2))`

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 211, normalized size of antiderivative = 0.84, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {90, 60, 60, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^{3/2} (A + Bx) \sqrt{d + ex} dx$$

$$\downarrow 90$$

$$\frac{B(a + bx)^{5/2} (d + ex)^{3/2}}{4be} - \frac{(3aBe - 8Abe + 5bBd) \int (a + bx)^{3/2} \sqrt{d + ex} dx}{8be}$$

$$\downarrow 60$$

$$\frac{B(a+bx)^{5/2}(d+ex)^{3/2}}{4be} - \frac{(3aBe - 8Abe + 5bBd) \left(\frac{(bd-ae) \int \frac{(a+bx)^{3/2}}{\sqrt{d+ex}} dx}{6b} + \frac{(a+bx)^{5/2}\sqrt{d+ex}}{3b} \right)}{8be}$$

↓ 60

$$\frac{B(a+bx)^{5/2}(d+ex)^{3/2}}{4be} - \frac{(3aBe - 8Abe + 5bBd) \left(\frac{(bd-ae) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \int \frac{\sqrt{a+bx}}{\sqrt{d+ex}} dx}{4e} \right)}{6b} + \frac{(a+bx)^{5/2}\sqrt{d+ex}}{3b} \right)}{8be}$$

8be

↓ 60

$$\frac{B(a+bx)^{5/2}(d+ex)^{3/2}}{4be} - \frac{(3aBe - 8Abe + 5bBd) \left(\frac{(bd-ae) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{2e} \right)}{4e} \right)}{6b} + \frac{(a+bx)^{5/2}\sqrt{d+ex}}{3b} \right)}{8be}$$

8be

↓ 66

$$\frac{B(a+bx)^{5/2}(d+ex)^{3/2}}{4be} - \frac{(3aBe - 8Abe + 5bBd) \left(\frac{(bd-ae) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}} \right)}{4e} \right)}{6b} + \frac{(a+bx)^{5/2}\sqrt{d+ex}}{3b} \right)}{8be}$$

8be

↓ 221

$$\frac{B(a+bx)^{5/2}(d+ex)^{3/2}}{4be} - \frac{(3aBe - 8Abe + 5bBd) \left(\frac{(bd-ae) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{\sqrt{be}^{3/2}} \right)}{4e} \right)}{6b} \right)}{8be} + \frac{(a+bx)^{5/2}\sqrt{d+ex}}{3b}$$

input `Int[(a + b*x)^(3/2)*(A + B*x)*Sqrt[d + e*x], x]`

output `(B*(a + b*x)^(5/2)*(d + e*x)^(3/2))/(4*b*e) - ((5*b*B*d - 8*A*b*e + 3*a*B*e)*(((a + b*x)^(5/2)*Sqrt[d + e*x])/(3*b) + ((b*d - a*e)*(((a + b*x)^(3/2)*Sqrt[d + e*x])/(2*e) - (3*(b*d - a*e)*((Sqrt[a + b*x]*Sqrt[d + e*x])/e - ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(Sqrt[b]*e^(3/2)))))/(4*e)))/(6*b)))/(8*b*e)`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 967 vs. $2(212) = 424$.

Time = 0.26 (sec) , antiderivative size = 968, normalized size of antiderivative = 3.87

method	result
default	$-\frac{\sqrt{bx+a}\sqrt{ex+d}\left(-96Bb^3e^3x^3\sqrt{(ex+d)(bx+a)}\sqrt{be}-128Ab^3e^3x^2\sqrt{(ex+d)(bx+a)}\sqrt{be}-16Bb^3de^2x^2\sqrt{(ex+d)(bx+a)}\sqrt{be}-9\right)}{\dots}$

input `int((b*x+a)^(3/2)*(B*x+A)*(e*x+d)^(1/2),x,method=_RETURNVERBOSE)`

output

```

-1/384*(b*x+a)^(1/2)*(e*x+d)^(1/2)*(-96*B*b^3*e^3*x^3*((e*x+d)*(b*x+a))^(1
/2)*(b*e)^(1/2)-128*A*b^3*e^3*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-16*B
*b^3*d*e^2*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-9*B*ln(1/2*(2*b*e*x+2*(
(e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^4*e^4+15*B*ln(1
/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^
4*d^4-40*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b^2*d*e^2*x-72*A*ln(1/2*(
2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*b^
2*d*e^3+72*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b
)/(b*e)^(1/2))*a*b^3*d^2*e^2-224*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b
^2*e^3*x-32*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^3*d*e^2*x-12*B*((e*x+d
)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^2*b*e^3*x+20*B*((e*x+d)*(b*x+a))^(1/2)*(b*e
)^(1/2)*b^3*d^2*e*x+18*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^3*e^3-30*B*
((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^3*d^3+24*A*ln(1/2*(2*b*e*x+2*((e*x+d
)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^3*b*e^4-24*A*ln(1/2*(
2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^4*d^
3*e-128*A*a*b^2*d*e^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-144*B*a*b^2*e^3*
x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-36*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b
*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^3*d^3*e-48*A*((e*x+d)*(
b*x+a))^(1/2)*(b*e)^(1/2)*a^2*b*e^3+48*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/
2)*b^3*d^2*e+12*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)...

```

Fricas [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 766, normalized size of antiderivative = 3.06

$$\int (a + bx)^{3/2} (A + Bx) \sqrt{d + ex} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^(3/2)*(B*x+A)*(e*x+d)^(1/2),x, algorithm="fricas")
```

output

```
[1/768*(3*(5*B*b^4*d^4 - 4*(3*B*a*b^3 + 2*A*b^4)*d^3*e + 6*(B*a^2*b^2 + 4*
A*a*b^3)*d^2*e^2 + 4*(B*a^3*b - 6*A*a^2*b^2)*d*e^3 - (3*B*a^4 - 8*A*a^3*b)
*e^4)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 - 4*(2*b
*e*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x + a)*sqrt(e*x + d) + 8*(b^2*d*e + a*b
*e^2)*x) + 4*(48*B*b^4*e^4*x^3 + 15*B*b^4*d^3*e - (31*B*a*b^3 + 24*A*b^4)*
d^2*e^2 + (9*B*a^2*b^2 + 64*A*a*b^3)*d*e^3 - 3*(3*B*a^3*b - 8*A*a^2*b^2)*e
^4 + 8*(B*b^4*d*e^3 + (9*B*a*b^3 + 8*A*b^4)*e^4)*x^2 - 2*(5*B*b^4*d^2*e^2
- 2*(5*B*a*b^3 + 4*A*b^4)*d*e^3 - (3*B*a^2*b^2 + 56*A*a*b^3)*e^4)*x)*sqrt(
b*x + a)*sqrt(e*x + d))/(b^3*e^4), 1/384*(3*(5*B*b^4*d^4 - 4*(3*B*a*b^3 +
2*A*b^4)*d^3*e + 6*(B*a^2*b^2 + 4*A*a*b^3)*d^2*e^2 + 4*(B*a^3*b - 6*A*a^2*
b^2)*d*e^3 - (3*B*a^4 - 8*A*a^3*b)*e^4)*sqrt(-b*e)*arctan(1/2*(2*b*e*x + b
*d + a*e)*sqrt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*e^2*x^2 + a*b*d*e +
(b^2*d*e + a*b*e^2)*x)) + 2*(48*B*b^4*e^4*x^3 + 15*B*b^4*d^3*e - (31*B*a*b
^3 + 24*A*b^4)*d^2*e^2 + (9*B*a^2*b^2 + 64*A*a*b^3)*d*e^3 - 3*(3*B*a^3*b -
8*A*a^2*b^2)*e^4 + 8*(B*b^4*d*e^3 + (9*B*a*b^3 + 8*A*b^4)*e^4)*x^2 - 2*(5
*B*b^4*d^2*e^2 - 2*(5*B*a*b^3 + 4*A*b^4)*d*e^3 - (3*B*a^2*b^2 + 56*A*a*b^3
)*e^4)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b^3*e^4)]
```

Sympy [F]

$$\int (a + bx)^{3/2} (A + Bx) \sqrt{d + ex} dx = \int (A + Bx) (a + bx)^{3/2} \sqrt{d + ex} dx$$

input

```
integrate((b*x+a)**(3/2)*(B*x+A)*(e*x+d)**(1/2),x)
```

output

```
Integral((A + B*x)*(a + b*x)**(3/2)*sqrt(d + e*x), x)
```

Maxima [F(-2)]

Exception generated.

$$\int (a + bx)^{3/2} (A + Bx) \sqrt{d + ex} dx = \text{Exception raised: ValueError}$$

input

```
integrate((b*x+a)**(3/2)*(B*x+A)*(e*x+d)**(1/2),x, algorithm="maxima")
```


output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1036 vs. $2(212) = 424$.

Time = 0.28 (sec) , antiderivative size = 1036, normalized size of antiderivative = 4.14

$$\int (a + bx)^{3/2}(A + Bx)\sqrt{d + ex} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^(3/2)*(B*x+A)*(e*x+d)^(1/2),x, algorithm="giac")
```

output

```
1/192*((sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*(b*x + a)*(4*(b*x + a)*(6*(
b*x + a)/b^3 + (b^12*d*e^5 - 25*a*b^11*e^6)/(b^14*e^6)) - (5*b^13*d^2*e^4
+ 14*a*b^12*d*e^5 - 163*a^2*b^11*e^6)/(b^14*e^6)) + 3*(5*b^14*d^3*e^3 + 9*
a*b^13*d^2*e^4 + 15*a^2*b^12*d*e^5 - 93*a^3*b^11*e^6)/(b^14*e^6))*sqrt(b*x
+ a) + 3*(5*b^4*d^4 + 4*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2 + 20*a^3*b*d*e^3
- 35*a^4*e^4)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*
e - a*b*e)))/(sqrt(b*e)*b^2*e^3))*B*abs(b) - 192*((b^2*d - a*b*e)*log(abs(
-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/sqrt(b*e)
- sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*sqrt(b*x + a))*A*a^2*abs(b)/b^2 + 4
8*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*b*x + 2*a + (b*d*e - 5*a*e^2)/e^
2)*sqrt(b*x + a) + (b^3*d^2 + 2*a*b^2*d*e - 3*a^2*b*e^2)*log(abs(-sqrt(b*e)
)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*e))*B*a
^2*abs(b)/b^3 + 96*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*b*x + 2*a + (b*
d*e - 5*a*e^2)/e^2)*sqrt(b*x + a) + (b^3*d^2 + 2*a*b^2*d*e - 3*a^2*b*e^2)*
log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(
sqrt(b*e)*e))*A*a*abs(b)/b^2 + 16*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*
(4*b*x + 4*a + (b*d*e^3 - 13*a*e^4)/e^4)*(b*x + a) - 3*(b^2*d^2*e^2 + 2*a*
b*d*e^3 - 11*a^2*e^4)/e^4)*sqrt(b*x + a) - 3*(b^4*d^3 + a*b^3*d^2*e + 3*a^
2*b^2*d*e^2 - 5*a^3*b*e^3)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d +
(b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*e^2))*B*a*abs(b)/b^3 + 8*(sqrt(b^2...
```

Mupad [F(-1)]

Timed out.

$$\int (a + bx)^{3/2} (A + Bx) \sqrt{d + ex} dx = \int (A + Bx) (a + bx)^{3/2} \sqrt{d + ex} dx$$

input `int((A + B*x)*(a + b*x)^(3/2)*(d + e*x)^(1/2), x)`

output `int((A + B*x)*(a + b*x)^(3/2)*(d + e*x)^(1/2), x)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 471, normalized size of antiderivative = 1.88

$$\int (a + bx)^{3/2} (A + Bx) \sqrt{d + ex} dx = \frac{15\sqrt{ex + d} \sqrt{bx + a} a^3 b e^4 + 73\sqrt{ex + d} \sqrt{bx + a} a^2 b^2 d e^3 + 118\sqrt{ex + d} \sqrt{bx + a} a^2 b^2 e^2 + 55\sqrt{ex + d} \sqrt{bx + a} a b^3 d e^2 + 36\sqrt{ex + d} \sqrt{bx + a} a b^3 d e^2 + 15\sqrt{ex + d} \sqrt{bx + a} b^4 d^3 e - 10\sqrt{ex + d} \sqrt{bx + a} b^4 d^3 e^2 + 8\sqrt{ex + d} \sqrt{bx + a} b^4 d^3 e^3 + 48\sqrt{ex + d} \sqrt{bx + a} b^4 d^3 e^4 - 15\sqrt{e} \sqrt{b} \log((\sqrt{e} \sqrt{a + bx} + \sqrt{b} \sqrt{d + ex}) / \sqrt{a e - b d}) a^4 e^4 + 60\sqrt{e} \sqrt{b} \log((\sqrt{e} \sqrt{a + bx} + \sqrt{b} \sqrt{d + ex}) / \sqrt{a e - b d}) a^3 b d e^3 - 90\sqrt{e} \sqrt{b} \log((\sqrt{e} \sqrt{a + bx} + \sqrt{b} \sqrt{d + ex}) / \sqrt{a e - b d}) a^2 b^2 d^2 e^2 + 60\sqrt{e} \sqrt{b} \log((\sqrt{e} \sqrt{a + bx} + \sqrt{b} \sqrt{d + ex}) / \sqrt{a e - b d}) a b^3 d^3 e - 15\sqrt{e} \sqrt{b} \log((\sqrt{e} \sqrt{a + bx} + \sqrt{b} \sqrt{d + ex}) / \sqrt{a e - b d}) b^4 d^4 / (192 b^2 e^4)$$

input `int((b*x+a)^(3/2)*(B*x+A)*(e*x+d)^(1/2), x)`

output `(15*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b*e**4 + 73*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**2*d*e**3 + 118*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**2*e**4*x - 55*sqrt(d + e*x)*sqrt(a + b*x)*a*b**3*d**2*e**2 + 36*sqrt(d + e*x)*sqrt(a + b*x)*a*b**3*d*e**3*x + 136*sqrt(d + e*x)*sqrt(a + b*x)*a*b**3*e**4*x**2 + 15*sqrt(d + e*x)*sqrt(a + b*x)*b**4*d**3*e - 10*sqrt(d + e*x)*sqrt(a + b*x)*b**4*d**2*e**2*x + 8*sqrt(d + e*x)*sqrt(a + b*x)*b**4*d*e**3*x**2 + 48*sqrt(d + e*x)*sqrt(a + b*x)*b**4*e**4*x**3 - 15*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**4*e**4 + 60*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**3*b*d*e**3 - 90*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**2*b**2*d**2*e**2 + 60*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*b**3*d**3*e - 15*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**4*d**4/(192*b**2*e**4)`

3.185 $\int \frac{(a+bx)^{3/2}(A+Bx)}{\sqrt{d+ex}} dx$

Optimal result	1686
Mathematica [A] (verified)	1687
Rubi [A] (verified)	1687
Maple [B] (verified)	1689
Fricas [A] (verification not implemented)	1690
Sympy [F]	1691
Maxima [F(-2)]	1691
Giac [A] (verification not implemented)	1692
Mupad [F(-1)]	1692
Reduce [B] (verification not implemented)	1693

Optimal result

Integrand size = 24, antiderivative size = 193

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{\sqrt{d+ex}} dx = \frac{(bd-ae)(5bBd-6Abe+aBe)\sqrt{a+bx}\sqrt{d+ex}}{8be^3} - \frac{(5bBd-6Abe+aBe)(a+bx)^{3/2}\sqrt{d+ex}}{12be^2} + \frac{B(a+bx)^{5/2}\sqrt{d+ex}}{3be} - \frac{(bd-ae)^2(5bBd-6Abe+aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{8b^{3/2}e^{7/2}}$$

output

```
1/8*(-a*e+b*d)*(-6*A*b*e+B*a*e+5*B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(1/2)/b/e^3-
1/12*(-6*A*b*e+B*a*e+5*B*b*d)*(b*x+a)^(3/2)*(e*x+d)^(1/2)/b/e^2+1/3*B*(b*x
+a)^(5/2)*(e*x+d)^(1/2)/b/e-1/8*(-a*e+b*d)^2*(-6*A*b*e+B*a*e+5*B*b*d)*arct
anh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)^(1/2))/b^(3/2)/e^(7/2)
```


$$\frac{(aBe - 6Abe + 5bBd) \left(\frac{B(a+bx)^{5/2}\sqrt{d+ex}}{3be} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{2e} \right)}{4e} \right)}{6be}$$

↓ 66

$$\frac{(aBe - 6Abe + 5bBd) \left(\frac{B(a+bx)^{5/2}\sqrt{d+ex}}{3be} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{e} \right)}{4e} \right)}{6be}$$

↓ 221

$$\frac{(aBe - 6Abe + 5bBd) \left(\frac{B(a+bx)^{5/2}\sqrt{d+ex}}{3be} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \operatorname{arctanh} \left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}} \right)}{\sqrt{be}^{3/2}} \right)}{4e} \right)}{6be}$$

input `Int[((a + b*x)^(3/2)*(A + B*x))/Sqrt[d + e*x],x]`

output `(B*(a + b*x)^(5/2)*Sqrt[d + e*x])/(3*b*e) - ((5*b*B*d - 6*A*b*e + a*B*e)*((a + b*x)^(3/2)*Sqrt[d + e*x])/(2*e) - (3*(b*d - a*e)*((Sqrt[a + b*x]*Sqrt[d + e*x])/e - ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(Sqrt[b]*e^(3/2))))/(4*e))/(6*b*e)`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 635 vs. $2(161) = 322$.

Time = 0.27 (sec) , antiderivative size = 636, normalized size of antiderivative = 3.30

method	result
default	$\frac{\sqrt{bx+a}\sqrt{ex+d}\left(16Bb^2e^2x^2\sqrt{(ex+d)(bx+a)}\sqrt{be}+18A\ln\left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}}\right)a^2be^3-36A\ln\left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}}\right)\right)}{\dots}$

input `int((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(1/2),x,method=_RETURNVERBOSE)`

output

```

1/48*(b*x+a)^(1/2)*(e*x+d)^(1/2)*(16*B*b^2*e^2*x^2*((e*x+d)*(b*x+a))^(1/2)
*(b*e)^(1/2)+18*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*
e+d*b)/(b*e)^(1/2))*a^2*b*e^3-36*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)
*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^2*d*e^2+18*A*ln(1/2*(2*b*e*x+2*((
e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^3*d^2*e+24*A*((e
*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^2*e^2*x-3*B*ln(1/2*(2*b*e*x+2*((e*x+d)*
(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^3*e^3-9*B*ln(1/2*(2*b*e
*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*b*d*e^2
+27*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)
^(1/2))*a*b^2*d^2*e-15*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(
1/2)+a*e+d*b)/(b*e)^(1/2))*b^3*d^3+28*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)
)*a*b*e^2*x-20*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^2*d*e*x+60*A*((e*x+
d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b*e^2-36*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(
1/2)*b^2*d*e+6*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^2*e^2-44*B*((e*x+d)
*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b*d*e+30*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/
2)*b^2*d^2)/b/e^3/((e*x+d)*(b*x+a))^(1/2)/(b*e)^(1/2)

```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 540, normalized size of antiderivative = 2.80

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{\sqrt{d+ex}} dx = \left[-\frac{3(5Bb^3d^3 - 3(3Bab^2 + 2Ab^3)d^2e + 3(Ba^2b + 4Aab^2)de^2 + (Ba^3 - 6Aa^2b))}{\dots} \right]$$

input

```
integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(1/2),x, algorithm="fricas")
```

output

```
[-1/96*(3*(5*B*b^3*d^3 - 3*(3*B*a*b^2 + 2*A*b^3)*d^2*e + 3*(B*a^2*b + 4*A*
a*b^2)*d*e^2 + (B*a^3 - 6*A*a^2*b)*e^3)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*
d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*b*e*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x + a
)*sqrt(e*x + d) + 8*(b^2*d*e + a*b*e^2)*x) - 4*(8*B*b^3*e^3*x^2 + 15*B*b^3
*d^2*e - 2*(11*B*a*b^2 + 9*A*b^3)*d*e^2 + 3*(B*a^2*b + 10*A*a*b^2)*e^3 - 2
*(5*B*b^3*d*e^2 - (7*B*a*b^2 + 6*A*b^3)*e^3)*x)*sqrt(b*x + a)*sqrt(e*x + d
))/b^2*e^4, 1/48*(3*(5*B*b^3*d^3 - 3*(3*B*a*b^2 + 2*A*b^3)*d^2*e + 3*(B*
a^2*b + 4*A*a*b^2)*d*e^2 + (B*a^3 - 6*A*a^2*b)*e^3)*sqrt(-b*e)*arctan(1/2*
(2*b*e*x + b*d + a*e)*sqrt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*e^2*x^2
+ a*b*d*e + (b^2*d*e + a*b*e^2)*x)) + 2*(8*B*b^3*e^3*x^2 + 15*B*b^3*d^2*e
- 2*(11*B*a*b^2 + 9*A*b^3)*d*e^2 + 3*(B*a^2*b + 10*A*a*b^2)*e^3 - 2*(5*B*b
^3*d*e^2 - (7*B*a*b^2 + 6*A*b^3)*e^3)*x)*sqrt(b*x + a)*sqrt(e*x + d))/b^2
*e^4]
```

Sympy [F]

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{\sqrt{d + ex}} dx = \int \frac{(A + Bx)(a + bx)^{\frac{3}{2}}}{\sqrt{d + ex}} dx$$

input

```
integrate((b*x+a)**(3/2)*(B*x+A)/(e*x+d)**(1/2),x)
```

output

```
Integral((A + B*x)*(a + b*x)**(3/2)/sqrt(d + e*x), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{\sqrt{d + ex}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(1/2),x, algorithm="maxima")
```


output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for m
ore detail
```

Giac [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.47

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{\sqrt{d+ex}} dx = \frac{\left(\sqrt{b^2d+(bx+a)be-abe\sqrt{bx+a}}\left(2(bx+a)\left(\frac{4(bx+a)B}{b^2e} - \frac{5Bb^3de^3+Bab^2e^4-6Aa^2b^3e^4}{b^4e^5}\right)\right)\right)}{\dots}$$

input

```
integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(1/2),x, algorithm="giac")
```

output

```
1/24*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*sqrt(b*x + a)*(2*(b*x + a)*(4*(b
*x + a)*B/(b^2*e) - (5*B*b^3*d*e^3 + B*a*b^2*e^4 - 6*A*b^3*e^4)/(b^4*e^5))
+ 3*(5*B*b^4*d^2*e^2 - 4*B*a*b^3*d*e^3 - 6*A*b^4*d*e^3 - B*a^2*b^2*e^4 +
6*A*a*b^3*e^4)/(b^4*e^5)) + 3*(5*B*b^3*d^3 - 9*B*a*b^2*d^2*e - 6*A*b^3*d^2
*e + 3*B*a^2*b*d*e^2 + 12*A*a*b^2*d*e^2 + B*a^3*e^3 - 6*A*a^2*b*e^3)*log(a
bs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(
b*e)*b*e^3))*b/abs(b)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{\sqrt{d+ex}} dx = \int \frac{(A+Bx)(a+bx)^{3/2}}{\sqrt{d+ex}} dx$$

input

```
int(((A + B*x)*(a + b*x)^(3/2))/(d + e*x)^(1/2), x)
```

output

```
int(((A + B*x)*(a + b*x)^(3/2))/(d + e*x)^(1/2), x)
```

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 320, normalized size of antiderivative = 1.66

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{\sqrt{d+ex}} dx = \frac{33\sqrt{ex+d}\sqrt{bx+a}a^2be^3 - 40\sqrt{ex+d}\sqrt{bx+a}ab^2de^2 + 26\sqrt{ex+d}\sqrt{bx+a}a^2be^2 - 40\sqrt{ex+d}\sqrt{bx+a}ab^2de + 26\sqrt{ex+d}\sqrt{bx+a}a^2be - 40\sqrt{ex+d}\sqrt{bx+a}ab^2d + 26\sqrt{ex+d}\sqrt{bx+a}a^2b}{24b^3e^4}$$

input `int((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(1/2),x)`

output

```
(33*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b*e**3 - 40*sqrt(d + e*x)*sqrt(a + b*x)*a*b**2*d*e**2 + 26*sqrt(d + e*x)*sqrt(a + b*x)*a*b**2*e**3*x + 15*sqrt(d + e*x)*sqrt(a + b*x)*b**3*d**2*e - 10*sqrt(d + e*x)*sqrt(a + b*x)*b**3*d*e**2*x + 8*sqrt(d + e*x)*sqrt(a + b*x)*b**3*e**3*x**2 + 15*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**3*e**3 - 45*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**2*b*d*e**2 + 45*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*b**2*d**2*e - 15*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**3*d**3)/(24*b*e**4)
```

$$3.186 \quad \int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{3/2}} dx$$

Optimal result	1694
Mathematica [A] (verified)	1695
Rubi [A] (verified)	1695
Maple [B] (verified)	1697
Fricas [B] (verification not implemented)	1698
Sympy [F]	1699
Maxima [F(-2)]	1699
Giac [A] (verification not implemented)	1700
Mupad [F(-1)]	1700
Reduce [B] (verification not implemented)	1701

Optimal result

Integrand size = 24, antiderivative size = 167

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{3/2}} dx = \frac{2(Bd - Ae)(a+bx)^{3/2}}{e^2\sqrt{d+ex}} - \frac{3(5bBd - 4Abe - aBe)\sqrt{a+bx}\sqrt{d+ex}}{4e^3} + \frac{B(a+bx)^{3/2}\sqrt{d+ex}}{2e^2} + \frac{3(bd - ae)(5bBd - 4Abe - aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{4\sqrt{b}e^{7/2}}$$

output

```
2*(-A*e+B*d)*(b*x+a)^(3/2)/e^2/(e*x+d)^(1/2)-3/4*(-4*A*b*e-B*a*e+5*B*b*d)*
(b*x+a)^(1/2)*(e*x+d)^(1/2)/e^3+1/2*B*(b*x+a)^(3/2)*(e*x+d)^(1/2)/e^2+3/4*
(-a*e+b*d)*(-4*A*b*e-B*a*e+5*B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/
(e*x+d)^(1/2))/b^(1/2)/e^(7/2)
```

Mathematica [A] (verified)

Time = 1.14 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.99

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{3/2}} dx = \frac{\sqrt{e}\sqrt{a+bx}(4Abe(3d+ex)+ae(13Bd-8Ae+5Bex)+bB(-15d^2-5dex+2e^2x^2))}{\sqrt{d+ex}} - \frac{6(bd-ae)(5bBd-4Ab)}{4e^{7/2}}$$

input

```
Integrate[((a + b*x)^(3/2)*(A + B*x))/(d + e*x)^(3/2),x]
```

output

```
((Sqrt[e]*Sqrt[a + b*x]*(4*A*b*e*(3*d + e*x) + a*e*(13*B*d - 8*A*e + 5*B*e*x) + b*B*(-15*d^2 - 5*d*e*x + 2*e^2*x^2)))/Sqrt[d + e*x] - (6*(b*d - a*e)*(5*b*B*d - 4*A*b*e - a*B*e)*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/(Sqrt[e]*(Sqrt[a - (b*d)/e] - Sqrt[a + b*x]))])/Sqrt[b])/(4*e^(7/2))
```

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.11, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$, Rules used = {87, 60, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{3/2}} dx$$

↓ 87

$$\frac{(-aBe - 4Abe + 5bBd) \int \frac{(a+bx)^{3/2}}{\sqrt{d+ex}} dx}{e(bd - ae)} - \frac{2(a + bx)^{5/2}(Bd - Ae)}{e\sqrt{d + ex}(bd - ae)}$$

↓ 60

$$\frac{(-aBe - 4Abe + 5bBd) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \int \frac{\sqrt{a+bx}}{\sqrt{d+ex}} dx}{4e} \right)}{e(bd - ae)} - \frac{2(a + bx)^{5/2}(Bd - Ae)}{e\sqrt{d + ex}(bd - ae)}$$

↓ 60

$$\begin{aligned}
 & \frac{(-aBe - 4Abe + 5bBd) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{2e} \right)}{4e} \right)}{e(bd-ae)} \\
 & \frac{2(a+bx)^{5/2}(Bd-Ae)}{e\sqrt{d+ex}(bd-ae)} \\
 & \quad \downarrow \text{66} \\
 & \frac{(-aBe - 4Abe + 5bBd) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d\sqrt{d+ex}}{e} \right)}{4e} \right)}{e(bd-ae)} \\
 & \frac{2(a+bx)^{5/2}(Bd-Ae)}{e\sqrt{d+ex}(bd-ae)} \\
 & \quad \downarrow \text{221} \\
 & \frac{(-aBe - 4Abe + 5bBd) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{\sqrt{be}^{3/2}} \right)}{4e} \right)}{e(bd-ae)} \\
 & \frac{2(a+bx)^{5/2}(Bd-Ae)}{e\sqrt{d+ex}(bd-ae)}
 \end{aligned}$$

input `Int[((a + b*x)^(3/2)*(A + B*x))/(d + e*x)^(3/2),x]`

output `(-2*(B*d - A*e)*(a + b*x)^(5/2))/(e*(b*d - a*e)*Sqrt[d + e*x]) + ((5*b*B*d - 4*A*b*e - a*B*e)*((a + b*x)^(3/2)*Sqrt[d + e*x])/(2*e) - (3*(b*d - a*e))*((Sqrt[a + b*x]*Sqrt[d + e*x])/e - ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(Sqrt[b]*e^(3/2))))/(4*e))/(e*(b*d - a*e))`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 739 vs. $2(137) = 274$.

Time = 0.26 (sec) , antiderivative size = 740, normalized size of antiderivative = 4.43

method	result
default	$\frac{\sqrt{bx+a} \left(12A \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) abe^3x - 12A \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) b^2de^2x + 3B \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) \right)}{\dots}$

input `int((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(3/2),x,method=_RETURNVERBOSE)`

output

```
[1/16*(3*(5*B*b^2*d^3 - 2*(3*B*a*b + 2*A*b^2)*d^2*e + (B*a^2 + 4*A*a*b)*d*
e^2 + (5*B*b^2*d^2*e - 2*(3*B*a*b + 2*A*b^2)*d*e^2 + (B*a^2 + 4*A*a*b)*e^3
)*x)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*b*
e*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x + a)*sqrt(e*x + d) + 8*(b^2*d*e + a*b*
e^2)*x) + 4*(2*B*b^2*e^3*x^2 - 15*B*b^2*d^2*e - 8*A*a*b*e^3 + (13*B*a*b +
12*A*b^2)*d*e^2 - (5*B*b^2*d*e^2 - (5*B*a*b + 4*A*b^2)*e^3)*x)*sqrt(b*x +
a)*sqrt(e*x + d))/(b*e^5*x + b*d*e^4), -1/8*(3*(5*B*b^2*d^3 - 2*(3*B*a*b +
2*A*b^2)*d^2*e + (B*a^2 + 4*A*a*b)*d*e^2 + (5*B*b^2*d^2*e - 2*(3*B*a*b +
2*A*b^2)*d*e^2 + (B*a^2 + 4*A*a*b)*e^3)*x)*sqrt(-b*e)*arctan(1/2*(2*b*e*x
+ b*d + a*e)*sqrt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*e^2*x^2 + a*b*d*e
+ (b^2*d*e + a*b*e^2)*x)) - 2*(2*B*b^2*e^3*x^2 - 15*B*b^2*d^2*e - 8*A*a*b
*e^3 + (13*B*a*b + 12*A*b^2)*d*e^2 - (5*B*b^2*d*e^2 - (5*B*a*b + 4*A*b^2)*
e^3)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b*e^5*x + b*d*e^4)]
```

Sympy [F]

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{3/2}} dx = \int \frac{(A + Bx)(a + bx)^{\frac{3}{2}}}{(d + ex)^{\frac{3}{2}}} dx$$

input

```
integrate((b*x+a)**(3/2)*(B*x+A)/(e*x+d)**(3/2), x)
```

output

```
Integral((A + B*x)*(a + b*x)**(3/2)/(d + e*x)**(3/2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{3/2}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(3/2), x, algorithm="maxima")
```


output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f
or more de
```

Giac [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.62

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{3/2}} dx = \frac{\sqrt{bx+a} \left((bx+a) \left(\frac{2(bx+a)B|b|}{be} - \frac{5Bb^2de^3|b|-Babe^4|b|-4Ab^2e^4|b|}{b^2e^5} \right) - \frac{3(5Bb^3d^2e^2|b|-6Babde|b|-4Ab^2de|b|+Ba^2e^2|b|+4Aabe^2|b|)}{4\sqrt{b^2d+(bx+a)be-abe}} \right) - \frac{3(5Bb^2d^2|b|-6Babde|b|-4Ab^2de|b|+Ba^2e^2|b|+4Aabe^2|b|) \log \left(\left| -\sqrt{be}\sqrt{bx+a} + \sqrt{b^2d+(bx+a)be-abe} \right| \right)}{4\sqrt{be}be^3}$$

input

```
integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(3/2),x, algorithm="giac")
```

output

```
1/4*sqrt(b*x + a)*((b*x + a)*(2*(b*x + a)*B*abs(b)/(b*e) - (5*B*b^2*d*e^3*
abs(b) - B*a*b*e^4*abs(b) - 4*A*b^2*e^4*abs(b))/(b^2*e^5)) - 3*(5*B*b^3*d^
2*e^2*abs(b) - 6*B*a*b^2*d*e^3*abs(b) - 4*A*b^3*d*e^3*abs(b) + B*a^2*b*e^4
*abs(b) + 4*A*a*b^2*e^4*abs(b))/(b^2*e^5))/sqrt(b^2*d + (b*x + a)*b*e - a*
b*e) - 3/4*(5*B*b^2*d^2*abs(b) - 6*B*a*b*d*e*abs(b) - 4*A*b^2*d*e*abs(b) +
B*a^2*e^2*abs(b) + 4*A*a*b*e^2*abs(b))*log(abs(-sqrt(b*e)*sqrt(b*x + a) +
sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*b*e^3)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{3/2}} dx = \int \frac{(A+Bx)(a+bx)^{3/2}}{(d+ex)^{3/2}} dx$$

input

```
int(((A + B*x)*(a + b*x)^(3/2))/(d + e*x)^(3/2),x)
```

output

```
int(((A + B*x)*(a + b*x)^(3/2))/(d + e*x)^(3/2), x)
```

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 485, normalized size of antiderivative = 2.90

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{3/2}} dx = \frac{-8\sqrt{ex+d}\sqrt{bx+a}a^2e^3 + 25\sqrt{ex+d}\sqrt{bx+a}abde^2 + 9\sqrt{ex+d}\sqrt{bx+a}}$$

input `int((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(3/2),x)`

output

```
( - 8*sqrt(d + e*x)*sqrt(a + b*x)*a**2*e**3 + 25*sqrt(d + e*x)*sqrt(a + b*x)*a*b*d*e**2 + 9*sqrt(d + e*x)*sqrt(a + b*x)*a*b*e**3*x - 15*sqrt(d + e*x)*sqrt(a + b*x)*b**2*d**2*e - 5*sqrt(d + e*x)*sqrt(a + b*x)*b**2*d*e**2*x + 2*sqrt(d + e*x)*sqrt(a + b*x)*b**2*e**3*x**2 + 15*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**2*d*e**2 + 15*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**2*e**3*x - 30*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*b*d**2*e - 30*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*b*d*e**2*x + 15*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**2*d**3 + 15*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**2*d**2*e*x - 10*sqrt(e)*sqrt(b)*a**2*d*e**2 - 10*sqrt(e)*sqrt(b)*a**2*e**3*x + 20*sqrt(e)*sqrt(b)*a*b*d**2*e + 20*sqrt(e)*sqrt(b)*a*b*d*e**2*x - 10*sqrt(e)*sqrt(b)*b**2*d**3 - 10*sqrt(e)*sqrt(b)*b**2*d**2*e*x)/(4*e**4*(d + e*x))
```

3.187 $\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{5/2}} dx$

Optimal result	1702
Mathematica [A] (verified)	1703
Rubi [A] (verified)	1703
Maple [B] (verified)	1706
Fricas [A] (verification not implemented)	1707
Sympy [F]	1707
Maxima [F(-2)]	1708
Giac [B] (verification not implemented)	1708
Mupad [F(-1)]	1709
Reduce [B] (verification not implemented)	1709

Optimal result

Integrand size = 24, antiderivative size = 155

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{5/2}} dx = \frac{2(Bd - Ae)(a+bx)^{3/2}}{3e^2(d+ex)^{3/2}} + \frac{2(2bBd - Abe - aBe)\sqrt{a+bx}}{e^3\sqrt{d+ex}} + \frac{bB\sqrt{a+bx}\sqrt{d+ex}}{e^3} - \frac{\sqrt{b}(5bBd - 2Abe - 3aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{e^{7/2}}$$

output

```
2/3*(-A*e+B*d)*(b*x+a)^(3/2)/e^2/(e*x+d)^(3/2)+2*(-A*b*e-B*a*e+2*B*b*d)*(b
*x+a)^(1/2)/e^3/(e*x+d)^(1/2)+b*B*(b*x+a)^(1/2)*(e*x+d)^(1/2)/e^3-b^(1/2)*
(-2*A*b*e-3*B*a*e+5*B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)^(
1/2))/e^(7/2)
```

Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.88

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{5/2}} dx = \frac{\sqrt{a+bx}(-2Abe(3d+4ex) - 2ae(2Bd+ Ae+ 3Bex) + bB(15d^2 + 20dex + 3e^2x^2))}{3e^3(d+ex)^{3/2}} + \frac{\sqrt{b}(-5bBd + 2Abe + 3aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{e}\sqrt{a+bx}}\right)}{e^{7/2}}$$

input

```
Integrate[((a + b*x)^(3/2)*(A + B*x))/(d + e*x)^(5/2),x]
```

output

```
(Sqrt[a + b*x]*(-2*A*b*e*(3*d + 4*e*x) - 2*a*e*(2*B*d + A*e + 3*B*e*x) + b*B*(15*d^2 + 20*d*e*x + 3*e^2*x^2)))/(3*e^3*(d + e*x)^(3/2)) + (Sqrt[b]*(-5*b*B*d + 2*A*b*e + 3*a*B*e)*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/(Sqrt[e]*Sqrt[a + b*x])])/e^(7/2)
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.16, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$, Rules used = {87, 57, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{5/2}} dx$$

$$\downarrow 87$$

$$\frac{(-3aBe - 2Abe + 5bBd) \int \frac{(a+bx)^{3/2}}{(d+ex)^{3/2}} dx}{3e(bd - ae)} - \frac{2(a+bx)^{5/2}(Bd - Ae)}{3e(d+ex)^{3/2}(bd - ae)}$$

$$\downarrow 57$$

$$\frac{(-3aBe - 2Abe + 5bBd) \left(\frac{3b \int \frac{\sqrt{a+bx}}{\sqrt{d+ex}} dx}{e} - \frac{2(a+bx)^{3/2}}{e\sqrt{d+ex}} \right)}{3e(bd - ae)} - \frac{2(a+bx)^{5/2}(Bd - Ae)}{3e(d+ex)^{3/2}(bd - ae)}$$

$$\begin{aligned}
 & \downarrow 60 \\
 & \frac{(-3aBe - 2Abe + 5bBd) \left(\frac{3b \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{2e} \right)}{e} - \frac{2(a+bx)^{3/2}}{e\sqrt{d+ex}} \right)}{3e(bd - ae)} \\
 & \frac{2(a + bx)^{5/2}(Bd - Ae)}{3e(d + ex)^{3/2}(bd - ae)} \\
 & \downarrow 66 \\
 & \frac{(-3aBe - 2Abe + 5bBd) \left(\frac{3b \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{e} \right)}{e} - \frac{2(a+bx)^{3/2}}{e\sqrt{d+ex}} \right)}{3e(bd - ae)} \\
 & \frac{2(a + bx)^{5/2}(Bd - Ae)}{3e(d + ex)^{3/2}(bd - ae)} \\
 & \downarrow 221 \\
 & \frac{(-3aBe - 2Abe + 5bBd) \left(\frac{3b \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \operatorname{arctanh} \left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}} \right)}{\sqrt{be}^{3/2}} \right)}{e} - \frac{2(a+bx)^{3/2}}{e\sqrt{d+ex}} \right)}{3e(bd - ae)} \\
 & \frac{2(a + bx)^{5/2}(Bd - Ae)}{3e(d + ex)^{3/2}(bd - ae)}
 \end{aligned}$$

input `Int[((a + b*x)^(3/2)*(A + B*x))/(d + e*x)^(5/2),x]`

output `(-2*(B*d - A*e)*(a + b*x)^(5/2))/(3*e*(b*d - a*e)*(d + e*x)^(3/2)) + ((5*b*B*d - 2*A*b*e - 3*a*B*e)*((-2*(a + b*x)^(3/2))/(e*Sqrt[d + e*x]) + (3*b*(Sqrt[a + b*x]*Sqrt[d + e*x])/e - ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(Sqrt[b]*e^(3/2))))/e)/(3*e*(b*d - a*e))`

Definitions of rubi rules used

rule 57 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))]
 Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] &&
 GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] &&
 (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1))]
 Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] &&
 NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) &&
 !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)]
 Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || (EqQ[e, 0] || (EqQ[c, 0] || LtQ[p, n]))))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 697 vs. $2(129) = 258$.

Time = 0.27 (sec) , antiderivative size = 698, normalized size of antiderivative = 4.50

method	result
default	$\sqrt{bx+a} \left(6A \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) b^2 e^3 x^2 + 9B \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) ab e^3 x^2 - 15B \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) \right)$

input

```
int((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(5/2),x,method=_RETURNVERBOSE)
```

output

```
1/6*(b*x+a)^(1/2)*(6*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^2*e^3*x^2+9*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b*e^3*x^2-15*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^2*d*e^2*x^2+12*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^2*d*e^2*x+18*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b*d*e^2*x-30*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^2*d^2*e*x+6*B*b*e^2*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+6*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^2*d^2*e-16*A*b*e^2*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+9*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b*d^2*e-15*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^2*d^3-12*B*a*e^2*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+40*B*b*d*e*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-4*A*a*e^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-12*A*b*d*e*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-8*B*a*d*e*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+30*B*b*d^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2))/(b*e)^(1/2)/((e*x+d)*(b*x+a))^(1/2)/e^3/(e*x+d)^(3/2)
```

Fricas [A] (verification not implemented)

Time = 0.92 (sec) , antiderivative size = 537, normalized size of antiderivative = 3.46

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{5/2}} dx = \left[-\frac{3(5Bbd^3 - (3Ba + 2Ab)d^2e + (5Bbde^2 - (3Ba + 2Ab)e^3)x^2 + 2(5Bbde^2 - (3Ba + 2Ab)e^3)x + 2(5Bbde^2 - (3Ba + 2Ab)e^3))}{(d+ex)^{5/2}} \right]$$

input `integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(5/2),x, algorithm="fricas")`

output `[-1/12*(3*(5*B*b*d^3 - (3*B*a + 2*A*b)*d^2*e + (5*B*b*d*e^2 - (3*B*a + 2*A*b)*b)*e^3)*x^2 + 2*(5*B*b*d^2*e - (3*B*a + 2*A*b)*d*e^2)*x)*sqrt(b/e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*b*e^2*x + b*d*e + a*e^2))*sqrt(b*x + a)*sqrt(e*x + d)*sqrt(b/e) + 8*(b^2*d*e + a*b*e^2)*x - 4*(3*B*b*e^2*x^2 + 15*B*b*d^2 - 2*A*a*e^2 - 2*(2*B*a + 3*A*b)*d*e + 2*(10*B*b*d*e - (3*B*a + 4*A*b)*e^2)*x)*sqrt(b*x + a)*sqrt(e*x + d)/(e^5*x^2 + 2*d*e^4*x + d^2*e^3), 1/6*(3*(5*B*b*d^3 - (3*B*a + 2*A*b)*d^2*e + (5*B*b*d*e^2 - (3*B*a + 2*A*b)*e^3)*x^2 + 2*(5*B*b*d^2*e - (3*B*a + 2*A*b)*d*e^2)*x)*sqrt(-b/e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(b*x + a)*sqrt(e*x + d)*sqrt(-b/e)/(b^2*e*x^2 + a*b*d + (b^2*d + a*b*e)*x)) + 2*(3*B*b*e^2*x^2 + 15*B*b*d^2 - 2*A*a*e^2 - 2*(2*B*a + 3*A*b)*d*e + 2*(10*B*b*d*e - (3*B*a + 4*A*b)*e^2)*x)*sqrt(b*x + a)*sqrt(e*x + d)/(e^5*x^2 + 2*d*e^4*x + d^2*e^3)]`

Sympy [F]

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{5/2}} dx = \int \frac{(A+Bx)(a+bx)^{3/2}}{(d+ex)^{5/2}} dx$$

input `integrate((b*x+a)**(3/2)*(B*x+A)/(e*x+d)**(5/2),x)`

output `Integral((A + B*x)*(a + b*x)**(3/2)/(d + e*x)**(5/2), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(5/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f or more de

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 369 vs. 2(129) = 258.

Time = 0.21 (sec) , antiderivative size = 369, normalized size of antiderivative = 2.38

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{5/2}} dx = \frac{\left((bx + a) \left(\frac{3(Bb^5de^4|b| - Bab^4e^5|b|)(bx+a)}{b^4de^5 - ab^3e^6} + \frac{4(5Bb^6d^2e^3|b| - 8Bab^5de^4|b| - 2Ab^6de^4|b| + 3Ba^2b^6)}{b^4de^5 - ab^3e^6} \right) \right.}{(5Bbd|b| - 3Bae|b| - 2Abe|b|) \log \left(\left| -\sqrt{be}\sqrt{bx+a} + \sqrt{b^2d + (bx+a)be - abe} \right| \right)} + \frac{\left. \right)}{\sqrt{bee^3}}$$

input `integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(5/2),x, algorithm="giac")`

output $\frac{1}{3} \left((bx + a) \left(3(Bb^5de^4|b| - Bab^4e^5|b|)(bx+a) / (b^4de^5 - ab^3e^6) + 4(5Bb^6d^2e^3|b| - 8Bab^5de^4|b| - 2Ab^6de^4|b| + 3Ba^2b^6) / (b^4de^5 - ab^3e^6) \right) \right. + \frac{(5Bbd|b| - 3Bae|b| - 2Abe|b|) \log \left(\left| -\sqrt{be}\sqrt{bx+a} + \sqrt{b^2d + (bx+a)be - abe} \right| \right)}{\sqrt{bee^3}} \left. \right)$

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{5/2}} dx = \int \frac{(A + Bx)(a + bx)^{3/2}}{(d + ex)^{5/2}} dx$$

input `int(((A + B*x)*(a + b*x)^(3/2))/(d + e*x)^(5/2),x)`

output `int(((A + B*x)*(a + b*x)^(3/2))/(d + e*x)^(5/2), x)`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 504, normalized size of antiderivative = 3.25

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{5/2}} dx = \frac{-4\sqrt{ex + d}\sqrt{bx + a}a^2e^3 - 20\sqrt{ex + d}\sqrt{bx + a}abd e^2 - 28\sqrt{ex + d}\sqrt{bx + a}}{(d + ex)^{5/2}}$$

input `int((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(5/2),x)`

output `(- 4*sqrt(d + e*x)*sqrt(a + b*x)*a**2*e**3 - 20*sqrt(d + e*x)*sqrt(a + b*x)*a*b*d*e**2 - 28*sqrt(d + e*x)*sqrt(a + b*x)*a*b*e**3*x + 30*sqrt(d + e*x)*sqrt(a + b*x)*b**2*d**2*e + 40*sqrt(d + e*x)*sqrt(a + b*x)*b**2*d*e**2*x + 6*sqrt(d + e*x)*sqrt(a + b*x)*b**2*e**3*x**2 + 30*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*b*d**2*e + 60*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*b*d*e**2*x + 30*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*b*e**3*x**2 - 30*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**2*d**3 - 60*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**2*d**2*e*x - 30*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**2*d*e**2*x**2 + 5*sqrt(e)*sqrt(b)*a*b*d**2*e + 10*sqrt(e)*sqrt(b)*a*b*d*e**2*x + 5*sqrt(e)*sqrt(b)*a*b*e**3*x**2 - 5*sqrt(e)*sqrt(b)*b**2*d**3 - 10*sqrt(e)*sqrt(b)*b**2*d**2*e*x - 5*sqrt(e)*sqrt(b)*b**2*d*e**2*x**2)/(6*e**4*(d**2 + 2*d*e*x + e**2*x**2))`

3.188 $\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{7/2}} dx$

Optimal result	1710
Mathematica [A] (verified)	1710
Rubi [A] (verified)	1711
Maple [B] (verified)	1713
Fricas [B] (verification not implemented)	1714
Sympy [F]	1715
Maxima [F(-2)]	1715
Giac [B] (verification not implemented)	1716
Mupad [F(-1)]	1717
Reduce [B] (verification not implemented)	1717

Optimal result

Integrand size = 24, antiderivative size = 138

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{7/2}} dx = -\frac{2(Bd - Ae)(a+bx)^{5/2}}{5e(bd - ae)(d+ex)^{5/2}} - \frac{2B(a+bx)^{3/2}}{3e^2(d+ex)^{3/2}} - \frac{2bB\sqrt{a+bx}}{e^3\sqrt{d+ex}} + \frac{2b^{3/2} \text{Barctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{e^{7/2}}$$

output
$$-2/5*(-A*e+B*d)*(b*x+a)^{(5/2)}/e/(-a*e+b*d)/(e*x+d)^{(5/2)}-2/3*B*(b*x+a)^{(3/2)}/e^2/(e*x+d)^{(3/2)}-2*b*B*(b*x+a)^{(1/2)}/e^3/(e*x+d)^{(1/2)}+2*b^{(3/2)}*B*\text{arc tanh}(e^{(1/2)}*(b*x+a)^{(1/2)}/b^{(1/2)}/(e*x+d)^{(1/2)})/e^{(7/2)}$$

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.24

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{7/2}} dx = \frac{2(a+bx)^{5/2} \left(-3Bde^2 + 3Ae^3 - \frac{5bBde(d+ex)}{a+bx} + \frac{5aBe^2(d+ex)}{a+bx} - \frac{15b^2Bd(d+ex)^2}{(a+bx)^2} + \frac{15abBe(d+ex)^2}{(a+bx)^2} \right)}{15e^3(-bd + ae)(d+ex)^{5/2}} + \frac{2b^{3/2} \text{Barctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{e}\sqrt{a+bx}}\right)}{e^{7/2}}$$

input `Integrate[((a + b*x)^(3/2)*(A + B*x))/(d + e*x)^(7/2),x]`

output $(-2*(a + b*x)^(5/2)*(-3*B*d*e^2 + 3*A*e^3 - (5*b*B*d*e*(d + e*x))/(a + b*x) + (5*a*B*e^2*(d + e*x))/(a + b*x) - (15*b^2*B*d*(d + e*x)^2)/(a + b*x)^2 + (15*a*b*B*e*(d + e*x)^2)/(a + b*x)^2)/(15*e^3*(-(b*d) + a*e)*(d + e*x)^(5/2)) + (2*b^(3/2)*B*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/(Sqrt[e]*Sqrt[a + b*x])])/e^(7/2)$

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.06, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$, Rules used = {87, 57, 57, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{7/2}} dx$$

$$\downarrow 87$$

$$\frac{B \int \frac{(a+bx)^{3/2}}{(d+ex)^{5/2}} dx}{e} - \frac{2(a+bx)^{5/2}(Bd - Ae)}{5e(d+ex)^{5/2}(bd - ae)}$$

$$\downarrow 57$$

$$\frac{B \left(\frac{b \int \frac{\sqrt{a+bx}}{(d+ex)^{3/2}} dx}{e} - \frac{2(a+bx)^{3/2}}{3e(d+ex)^{3/2}} \right)}{e} - \frac{2(a+bx)^{5/2}(Bd - Ae)}{5e(d+ex)^{5/2}(bd - ae)}$$

$$\downarrow 57$$

$$\frac{B \left(\frac{b \left(\frac{b \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{e} - \frac{2\sqrt{a+bx}}{e\sqrt{d+ex}} \right)}{e} - \frac{2(a+bx)^{3/2}}{3e(d+ex)^{3/2}} \right)}{e} - \frac{2(a+bx)^{5/2}(Bd - Ae)}{5e(d+ex)^{5/2}(bd - ae)}$$

$$\downarrow 66$$

$$B \left(\frac{b \left(\frac{2b \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}} - \frac{2\sqrt{a+bx}}{e\sqrt{d+ex}} \right)}{e} - \frac{2(a+bx)^{3/2}}{3e(d+ex)^{3/2}} \right)}{e} - \frac{2(a+bx)^{5/2}(Bd - Ae)}{5e(d+ex)^{5/2}(bd - ae)} \right)$$

↓ 221

$$B \left(\frac{b \left(\frac{2\sqrt{b} \operatorname{arctanh} \left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}} \right) - \frac{2\sqrt{a+bx}}{e\sqrt{d+ex}} \right)}{e^{3/2}} - \frac{2(a+bx)^{3/2}}{3e(d+ex)^{3/2}} \right)}{e} - \frac{2(a+bx)^{5/2}(Bd - Ae)}{5e(d+ex)^{5/2}(bd - ae)} \right)$$

input `Int[((a + b*x)^(3/2)*(A + B*x))/(d + e*x)^(7/2),x]`

output `(-2*(B*d - A*e)*(a + b*x)^(5/2))/(5*e*(b*d - a*e)*(d + e*x)^(5/2)) + (B*((-2*(a + b*x)^(3/2))/(3*e*(d + e*x)^(3/2)) + (b*((-2*sqrt[a + b*x])/(e*sqrt[d + e*x]) + (2*sqrt[b]*ArcTanh[(sqrt[e]*sqrt[a + b*x])/(sqrt[b]*sqrt[d + e*x])])))/e^(3/2)))/e`

Defintions of rubi rules used

rule 57 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))]`
`Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(IntegerQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(sqrt[(a_) + (b_.)*(x_)]*sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, sqrt[a + b*x]/sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

output

```

-1/15*(b*x+a)^(1/2)*(-15*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)
^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^2*e^4*x^3+15*B*ln(1/2*(2*b*e*x+2*((e*x+d)
*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^3*d*e^3*x^3-45*B*ln(1/
2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b
^2*d*e^3*x^2+45*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*
e+d*b)/(b*e)^(1/2))*b^3*d^2*e^2*x^2+6*A*b^2*e^3*x^2*((e*x+d)*(b*x+a))^(1/2)
)*(b*e)^(1/2)-45*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a
*e+d*b)/(b*e)^(1/2))*a*b^2*d^2*e^2*x+45*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+
a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^3*d^3*e*x+40*B*a*b*e^3*x^2*(
(e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-46*B*b^2*d*e^2*x^2*((e*x+d)*(b*x+a))^(1
/2)*(b*e)^(1/2)+12*A*a*b*e^3*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-15*B*ln
(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*
a*b^2*d^3*e+15*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e
+d*b)/(b*e)^(1/2))*b^3*d^4+10*B*a^2*e^3*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1
/2)+48*B*a*b*d*e^2*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-70*B*b^2*d^2*e*x*
((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+6*A*a^2*e^3*((e*x+d)*(b*x+a))^(1/2)*(b
*e)^(1/2)+4*B*a^2*d*e^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+20*B*a*b*d^2*e
*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-30*B*b^2*d^3*((e*x+d)*(b*x+a))^(1/2)*
(b*e)^(1/2))/((e*x+d)*(b*x+a))^(1/2)/(a*e-b*d)/(b*e)^(1/2)/(e*x+d)^(5/2)/e
^3

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 371 vs. $2(110) = 220$.

Time = 2.54 (sec) , antiderivative size = 767, normalized size of antiderivative = 5.56

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{7/2}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(7/2),x, algorithm="fricas")
```

output

```
[1/30*(15*(B*b^2*d^4 - B*a*b*d^3*e + (B*b^2*d*e^3 - B*a*b*e^4)*x^3 + 3*(B*b^2*d^2*e^2 - B*a*b*d*e^3)*x^2 + 3*(B*b^2*d^3*e - B*a*b*d^2*e^2)*x)*sqrt(b/e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*b*e^2*x + b*d*e + a*e^2)*sqrt(b*x + a)*sqrt(e*x + d)*sqrt(b/e) + 8*(b^2*d*e + a*b*e^2)*x) - 4*(15*B*b^2*d^3 - 10*B*a*b*d^2*e - 2*B*a^2*d*e^2 - 3*A*a^2*e^3 + (23*B*b^2*d*e^2 - (20*B*a*b + 3*A*b^2)*e^3)*x^2 + (35*B*b^2*d^2*e - 24*B*a*b*d*e^2 - (5*B*a^2 + 6*A*a*b)*e^3)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b*d^4*e^3 - a*d^3*e^4 + (b*d*e^6 - a*e^7)*x^3 + 3*(b*d^2*e^5 - a*d*e^6)*x^2 + 3*(b*d^3*e^4 - a*d^2*e^5)*x), -1/15*(15*(B*b^2*d^4 - B*a*b*d^3*e + (B*b^2*d*e^3 - B*a*b*e^4)*x^3 + 3*(B*b^2*d^2*e^2 - B*a*b*d*e^3)*x^2 + 3*(B*b^2*d^3*e - B*a*b*d^2*e^2)*x)*sqrt(-b/e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(b*x + a)*sqrt(e*x + d)*sqrt(-b/e)/(b^2*e*x^2 + a*b*d + (b^2*d + a*b*e)*x)) + 2*(15*B*b^2*d^3 - 10*B*a*b*d^2*e - 2*B*a^2*d*e^2 - 3*A*a^2*e^3 + (23*B*b^2*d*e^2 - (20*B*a*b + 3*A*b^2)*e^3)*x^2 + (35*B*b^2*d^2*e - 24*B*a*b*d*e^2 - (5*B*a^2 + 6*A*a*b)*e^3)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b*d^4*e^3 - a*d^3*e^4 + (b*d*e^6 - a*e^7)*x^3 + 3*(b*d^2*e^5 - a*d*e^6)*x^2 + 3*(b*d^3*e^4 - a*d^2*e^5)*x)]
```

Sympy [F]

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{7/2}} dx = \int \frac{(A + Bx)(a + bx)^{3/2}}{(d + ex)^{7/2}} dx$$

input

```
integrate((b*x+a)**(3/2)*(B*x+A)/(e*x+d)**(7/2), x)
```

output

```
Integral((A + B*x)*(a + b*x)**(3/2)/(d + e*x)**(7/2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{7/2}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(7/2), x, algorithm="maxima")
```


output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f
or more de
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 398 vs. $2(110) = 220$.

Time = 0.26 (sec) , antiderivative size = 398, normalized size of antiderivative = 2.88

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{7/2}} dx =$$

$$\frac{2Bb|b| \log\left(\left|-\sqrt{be}\sqrt{bx+a} + \sqrt{b^2d+(bx+a)be-abe}\right|\right)}{\sqrt{bee^3}}$$

$$- \frac{2\left((bx+a)\left(\frac{(23Bb^7d^2e^4|b|-43Bab^6de^5|b|-3Ab^7de^5|b|+20Ba^2b^5e^6|b|+3Aab^6e^6|b|)(bx+a)}{b^4d^2e^5-2ab^3de^6+a^2b^2e^7} + \frac{35(Bb^8d^3e^3|b|-3Bab^7d^2e^4|b|+3Ba^2b^6e^5|b|)(bx+a)}{b^4d^2e^5-2ab^3de^6+a^2b^2e^7}\right)}{15(b^2d+(bx+a)be-ae^2)}$$

input

```
integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(7/2),x, algorithm="giac")
```

output

```
-2*B*b*abs(b)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*
e - a*b*e)))/(sqrt(b*e)*e^3) - 2/15*((b*x + a)*((23*B*b^7*d^2*e^4*abs(b) -
43*B*a*b^6*d*e^5*abs(b) - 3*A*b^7*d*e^5*abs(b) + 20*B*a^2*b^5*e^6*abs(b)
+ 3*A*a*b^6*e^6*abs(b))*(b*x + a)/(b^4*d^2*e^5 - 2*a*b^3*d*e^6 + a^2*b^2*e
^7) + 35*(B*b^8*d^3*e^3*abs(b) - 3*B*a*b^7*d^2*e^4*abs(b) + 3*B*a^2*b^6*d*
e^5*abs(b) - B*a^3*b^5*e^6*abs(b))/(b^4*d^2*e^5 - 2*a*b^3*d*e^6 + a^2*b^2*
e^7)) + 15*(B*b^9*d^4*e^2*abs(b) - 4*B*a*b^8*d^3*e^3*abs(b) + 6*B*a^2*b^7*
d^2*e^4*abs(b) - 4*B*a^3*b^6*d*e^5*abs(b) + B*a^4*b^5*e^6*abs(b))/(b^4*d^2
*e^5 - 2*a*b^3*d*e^6 + a^2*b^2*e^7))*sqrt(b*x + a)/(b^2*d + (b*x + a)*b*e
- a*b*e)^(5/2)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{7/2}} dx = \int \frac{(A + Bx)(a + bx)^{3/2}}{(d + ex)^{7/2}} dx$$

input `int(((A + B*x)*(a + b*x)^(3/2))/(d + e*x)^(7/2), x)`

output `int(((A + B*x)*(a + b*x)^(3/2))/(d + e*x)^(7/2), x)`

Reduce [B] (verification not implemented)

Time = 0.71 (sec) , antiderivative size = 403, normalized size of antiderivative = 2.92

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{7/2}} dx = \frac{-\frac{2\sqrt{ex+d}\sqrt{bx+a}a^2e^3}{5} - \frac{2\sqrt{ex+d}\sqrt{bx+a}abde^2}{3} - \frac{22\sqrt{ex+d}\sqrt{bx+a}abe^3x}{15} - 2\sqrt{ex+d}\sqrt{bx+a}}{(d + ex)^{7/2}}$$

input `int((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(7/2), x)`

output `(2*(- 3*sqrt(d + e*x)*sqrt(a + b*x)*a**2*e**3 - 5*sqrt(d + e*x)*sqrt(a + b*x)*a*b*d*e**2 - 11*sqrt(d + e*x)*sqrt(a + b*x)*a*b*e**3*x - 15*sqrt(d + e*x)*sqrt(a + b*x)*b**2*d**2*e - 35*sqrt(d + e*x)*sqrt(a + b*x)*b**2*d*e**2*x - 23*sqrt(d + e*x)*sqrt(a + b*x)*b**2*e**3*x**2 + 15*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**2*d**3 + 45*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**2*d**2*e*x + 45*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**2*d*e**2*x**2 + 15*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**2*e**3*x**3 + 5*sqrt(e)*sqrt(b)*b**2*d**3 + 15*sqrt(e)*sqrt(b)*b**2*d**2*e*x + 15*sqrt(e)*sqrt(b)*b**2*d*e**2*x**2 + 5*sqrt(e)*sqrt(b)*b**2*e**3*x**3)/(15*e**4*(d**3 + 3*d**2*e*x + 3*d*e**2*x**2 + e**3*x**3))`

3.189 $\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{9/2}} dx$

Optimal result	1718
Mathematica [A] (verified)	1718
Rubi [A] (verified)	1719
Maple [A] (verified)	1720
Fricas [B] (verification not implemented)	1721
Sympy [F]	1721
Maxima [F(-2)]	1722
Giac [B] (verification not implemented)	1722
Mupad [B] (verification not implemented)	1723
Reduce [B] (verification not implemented)	1723

Optimal result

Integrand size = 24, antiderivative size = 95

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{9/2}} dx = -\frac{2(Bd - Ae)(a+bx)^{5/2}}{7e(bd - ae)(d+ex)^{7/2}} + \frac{2(5bBd + 2Abe - 7aBe)(a+bx)^{5/2}}{35e(bd - ae)^2(d+ex)^{5/2}}$$

output `-2/7*(-A*e+B*d)*(b*x+a)^(5/2)/e/(-a*e+b*d)/(e*x+d)^(7/2)+2/35*(2*A*b*e-7*B*a*e+5*B*b*d)*(b*x+a)^(5/2)/e/(-a*e+b*d)^2/(e*x+d)^(5/2)`

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.69

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{9/2}} dx = \frac{2(a+bx)^{5/2}(B(-2ad + 5bdx - 7aex) + A(7bd - 5ae + 2bex))}{35(bd - ae)^2(d+ex)^{7/2}}$$

input `Integrate[((a + b*x)^(3/2)*(A + B*x))/(d + e*x)^(9/2),x]`

output

$$(2*(a + b*x)^(5/2)*(B*(-2*a*d + 5*b*d*x - 7*a*e*x) + A*(7*b*d - 5*a*e + 2*b*e*x)))/(35*(b*d - a*e)^2*(d + e*x)^(7/2))$$

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {87, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{9/2}} dx$$

$$\downarrow 87$$

$$\frac{(-7aBe + 2Abe + 5bBd) \int \frac{(a+bx)^{3/2}}{(d+ex)^{7/2}} dx}{7e(bd - ae)} - \frac{2(a + bx)^{5/2}(Bd - Ae)}{7e(d + ex)^{7/2}(bd - ae)}$$

$$\downarrow 48$$

$$\frac{2(a + bx)^{5/2}(-7aBe + 2Abe + 5bBd)}{35e(d + ex)^{5/2}(bd - ae)^2} - \frac{2(a + bx)^{5/2}(Bd - Ae)}{7e(d + ex)^{7/2}(bd - ae)}$$

input

$$\text{Int}[(a + b*x)^(3/2)*(A + B*x)/(d + e*x)^(9/2), x]$$

output

$$(-2*(B*d - A*e)*(a + b*x)^(5/2))/(7*e*(b*d - a*e)*(d + e*x)^(7/2)) + (2*(5*b*B*d + 2*A*b*e - 7*a*B*e)*(a + b*x)^(5/2))/(35*e*(b*d - a*e)^2*(d + e*x)^(5/2))$$

Definitions of rubi rules used

rule 48

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[[a + b*x]^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

rule 87

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.78

method	result	size
gospers	$-\frac{2(bx+a)^{\frac{5}{2}}(-2Abex+7Bae x-5Bbdx+5Aae-7Abd+2Bad)}{35(ex+d)^{\frac{7}{2}}(a^2e^2-2abde+b^2d^2)}$	74
orering	$-\frac{2(bx+a)^{\frac{5}{2}}(-2Abex+7Bae x-5Bbdx+5Aae-7Abd+2Bad)}{35(ex+d)^{\frac{7}{2}}(a^2e^2-2abde+b^2d^2)}$	74
default	$-\frac{2(-2A b^2 e x^2+7B a b e x^2-5B b^2 d x^2+3A a b e x-7A b^2 d x+7B a^2 e x-3B a b d x+5a^2 A e-7A a b d+2B a^2 d)(b x+a)^{\frac{3}{2}}}{35(ex+d)^{\frac{7}{2}}(ae-db)^2}$	107

input

```
int((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(9/2),x,method=_RETURNVERBOSE)
```

output

```
-2/35*(b*x+a)^(5/2)*(-2*A*b*e*x+7*B*a*e*x-5*B*b*d*x+5*A*a*e-7*A*b*d+2*B*a*
d)/(e*x+d)^(7/2)/(a^2*e^2-2*a*b*d*e+b^2*d^2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 306 vs. $2(83) = 166$.

Time = 5.94 (sec) , antiderivative size = 306, normalized size of antiderivative = 3.22

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{9/2}} dx =$$

$$-\frac{2(5Aa^3e - (5Bb^3d - (7Bab^2 - 2Ab^3)e)x^3 - ((8Bab^2 + 7Ab^3)d - (14Ba^2b + Aab^2)e)x^2 + (2Ba^3 - 7Aa^2b)d - ((B^2a^2b + 14A^2a^2b^2)d - (7B^2a^3 + 8A^2a^2b^2)e)x) \sqrt{bx + a} \sqrt{ex + d}}{35(b^2d^6 - 2abd^5e + a^2d^4e^2 + (b^2d^2e^4 - 2abde^5 + a^2e^6)x^4 + 4(b^2d^3e^3 - 2abd^2e^4 + a^2de^5)x^3 + 6(b^2d^4e^2 - 2a^2bd^3e^3 + a^2d^2e^4)x^2 + 4(b^2d^5e - 2a^2bd^4e^2 + a^2d^3e^3)x}$$

input `integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(9/2),x, algorithm="fricas")`

output `-2/35*(5*A*a^3*e - (5*B*b^3*d - (7*B*a*b^2 - 2*A*b^3)*e)*x^3 - ((8*B*a*b^2 + 7*A*b^3)*d - (14*B*a^2*b + A*a*b^2)*e)*x^2 + (2*B*a^3 - 7*A*a^2*b)*d - ((B*a^2*b + 14*A*a*b^2)*d - (7*B*a^3 + 8*A*a^2*b)*e)*x)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*d^6 - 2*a*b*d^5*e + a^2*d^4*e^2 + (b^2*d^2*e^4 - 2*a*b*d*e^5 + a^2*e^6)*x^4 + 4*(b^2*d^3*e^3 - 2*a*b*d^2*e^4 + a^2*d*e^5)*x^3 + 6*(b^2*d^4*e^2 - 2*a*b*d^3*e^3 + a^2*d^2*e^4)*x^2 + 4*(b^2*d^5*e - 2*a*b*d^4*e^2 + a^2*d^3*e^3)*x)`

Sympy [F]

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{9/2}} dx = \int \frac{(A + Bx)(a + bx)^{\frac{3}{2}}}{(d + ex)^{\frac{9}{2}}} dx$$

input `integrate((b*x+a)**(3/2)*(B*x+A)/(e*x+d)**(9/2),x)`

output `Integral((A + B*x)*(a + b*x)**(3/2)/(d + e*x)**(9/2), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{9/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(9/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f or more de

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 285 vs. 2(83) = 166.

Time = 0.31 (sec) , antiderivative size = 285, normalized size of antiderivative = 3.00

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{9/2}} dx = \frac{2(bx+a)^{5/2} \left(\frac{(5Bb^9d^2e^3|b|-12Bab^8de^4|b|+2Ab^9de^4|b|+7Ba^2b^7e^5|b|-2Aab^8e^5|b|)(bx+a)}{b^5d^3e^3-3ab^4d^2e^4+3a^2b^3de^5-a^3b^2e^6} - \frac{7(B}{35(b^2d+(bx+a)be -$$

input `integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(9/2),x, algorithm="giac")`

output
$$\frac{2/35*(b*x + a)^{(5/2)*((5*B*b^9*d^2*e^3*abs(b) - 12*B*a*b^8*d*e^4*abs(b) + 2*A*b^9*d*e^4*abs(b) + 7*B*a^2*b^7*e^5*abs(b) - 2*A*a*b^8*e^5*abs(b))*(b*x + a)/(b^5*d^3*e^3 - 3*a*b^4*d^2*e^4 + 3*a^2*b^3*d*e^5 - a^3*b^2*e^6) - 7*(B*a*b^9*d^2*e^3*abs(b) - A*b^10*d^2*e^3*abs(b) - 2*B*a^2*b^8*d*e^4*abs(b) + 2*A*a*b^9*d*e^4*abs(b) + B*a^3*b^7*e^5*abs(b) - A*a^2*b^8*e^5*abs(b))/(b^5*d^3*e^3 - 3*a*b^4*d^2*e^4 + 3*a^2*b^3*d*e^5 - a^3*b^2*e^6))/(b^2*d + (b*x + a)*b*e - a*b*e)^{(7/2)}$$

Mupad [B] (verification not implemented)

Time = 1.46 (sec) , antiderivative size = 257, normalized size of antiderivative = 2.71

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{9/2}} dx = \frac{\sqrt{d+ex} \left(\frac{\sqrt{a+bx}(10Aa^3e+4Ba^3d-14Aa^2bd)}{35e^4(ae-bd)^2} - \frac{x^3\sqrt{a+bx}(4Ab^3e+10Bb^3d-14Bab^2e)}{35e^4(ae-bd)^2} + \frac{x\sqrt{a+bx}(14Ba^3e-28Aab^2d+14Bb^3d-14Aab^2e)}{35e^4(ae-bd)^2} \right)}{x^4 + \frac{d^4}{e^4} + \frac{4dx^3}{e} + \frac{4d^3x}{e^3} + \frac{6d^2x^2}{e^2}}$$

input

```
int(((A + B*x)*(a + b*x)^(3/2))/(d + e*x)^(9/2), x)
```

output

```

-((d + e*x)^(1/2)*(((a + b*x)^(1/2)*(10*A*a^3*e + 4*B*a^3*d - 14*A*a^2*b*d
))/ (35*e^4*(a*e - b*d)^2) - (x^3*(a + b*x)^(1/2)*(4*A*b^3*e + 10*B*b^3*d -
14*B*a*b^2*e))/ (35*e^4*(a*e - b*d)^2) + (x*(a + b*x)^(1/2)*(14*B*a^3*e -
28*A*a*b^2*d + 16*A*a^2*b*e - 2*B*a^2*b*d))/ (35*e^4*(a*e - b*d)^2) - (x^2*
(a + b*x)^(1/2)*(14*A*b^3*d - 2*A*a*b^2*e + 16*B*a*b^2*d - 28*B*a^2*b*e))/
(35*e^4*(a*e - b*d)^2)))/(x^4 + d^4/e^4 + (4*d*x^3)/e + (4*d^3*x)/e^3 + (6
*d^2*x^2)/e^2)

```

Reduce [B] (verification not implemented)

Time = 0.44 (sec) , antiderivative size = 267, normalized size of antiderivative = 2.81

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{9/2}} dx = \frac{-\frac{2\sqrt{ex+d}\sqrt{bx+a}a^3e^4}{7} - \frac{6\sqrt{ex+d}\sqrt{bx+a}a^2be^4x}{7} - \frac{6\sqrt{ex+d}\sqrt{bx+a}ab^2e^4x^2}{7} - \frac{2\sqrt{ex+d}\sqrt{bx+a}b^3e^4x^3}{7}}{e^4(ae^5x^4 - bde^4x^4 + 4ade^4x^3 - 4bd^2e^3x^3 + 6ad^2e^2x^2)}$$

input

```
int((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(9/2), x)
```

output

```

(2*( - sqrt(d + e*x)*sqrt(a + b*x)*a**3*e**4 - 3*sqrt(d + e*x)*sqrt(a + b*
x)*a**2*b*e**4*x - 3*sqrt(d + e*x)*sqrt(a + b*x)*a*b**2*e**4*x**2 - sqrt(d
+ e*x)*sqrt(a + b*x)*b**3*e**4*x**3 - sqrt(e)*sqrt(b)*b**3*d**4 - 4*sqrt(
e)*sqrt(b)*b**3*d**3*e*x - 6*sqrt(e)*sqrt(b)*b**3*d**2*e**2*x**2 - 4*sqrt(
e)*sqrt(b)*b**3*d*e**3*x**3 - sqrt(e)*sqrt(b)*b**3*e**4*x**4))/(7*e**4*(a
*d**4*e + 4*a*d**3*e**2*x + 6*a*d**2*e**3*x**2 + 4*a*d*e**4*x**3 + a*e**5*x
**4 - b*d**5 - 4*b*d**4*e*x - 6*b*d**3*e**2*x**2 - 4*b*d**2*e**3*x**3 - b*
d*e**4*x**4))

```


3.190 $\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{11/2}} dx$

Optimal result	1724
Mathematica [A] (verified)	1724
Rubi [A] (verified)	1725
Maple [A] (verified)	1727
Fricas [B] (verification not implemented)	1727
Sympy [F]	1728
Maxima [F(-2)]	1728
Giac [B] (verification not implemented)	1729
Mupad [B] (verification not implemented)	1730
Reduce [B] (verification not implemented)	1730

Optimal result

Integrand size = 24, antiderivative size = 147

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{11/2}} dx = -\frac{2(Bd - Ae)(a+bx)^{5/2}}{9e(bd - ae)(d+ex)^{9/2}} + \frac{2(5bBd + 4Abe - 9aBe)(a+bx)^{5/2}}{63e(bd - ae)^2(d+ex)^{7/2}} + \frac{4b(5bBd + 4Abe - 9aBe)(a+bx)^{5/2}}{315e(bd - ae)^3(d+ex)^{5/2}}$$

output `-2/9*(-A*e+B*d)*(b*x+a)^(5/2)/e/(-a*e+b*d)/(e*x+d)^(9/2)+2/63*(4*A*b*e-9*B*a*e+5*B*b*d)*(b*x+a)^(5/2)/e/(-a*e+b*d)^2/(e*x+d)^(7/2)+4/315*b*(4*A*b*e-9*B*a*e+5*B*b*d)*(b*x+a)^(5/2)/e/(-a*e+b*d)^3/(e*x+d)^(5/2)`

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.91

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{11/2}} dx = \frac{2(a+bx)^{9/2} \left(-35Bde + 35Ae^2 + \frac{45bBd(d+ex)}{a+bx} - \frac{90Abe(d+ex)}{a+bx} + \frac{45aBe(d+ex)}{a+bx} + 63e^2 \right)}{315(bd - ae)^3(d+ex)^{9/2}}$$

input `Integrate[((a + b*x)^(3/2)*(A + B*x))/(d + e*x)^(11/2),x]`

output

$$(2*(a + b*x)^(9/2)*(-35*B*d*e + 35*A*e^2 + (45*b*B*d*(d + e*x))/(a + b*x) - (90*A*b*e*(d + e*x))/(a + b*x) + (45*a*B*e*(d + e*x))/(a + b*x) + (63*A*b^2*(d + e*x)^2)/(a + b*x)^2 - (63*a*b*B*(d + e*x)^2)/(a + b*x)^2))/(315*(b*d - a*e)^3*(d + e*x)^(9/2))$$

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.97, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {87, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{11/2}} dx$$

$$\downarrow 87$$

$$\frac{(-9aBe + 4Abe + 5bBd) \int \frac{(a+bx)^{3/2}}{(d+ex)^{9/2}} dx}{9e(bd - ae)} - \frac{2(a + bx)^{5/2}(Bd - Ae)}{9e(d + ex)^{9/2}(bd - ae)}$$

$$\downarrow 55$$

$$\frac{(-9aBe + 4Abe + 5bBd) \left(\frac{2b \int \frac{(a+bx)^{3/2}}{(d+ex)^{7/2}} dx}{7(bd-ae)} + \frac{2(a+bx)^{5/2}}{7(d+ex)^{7/2}(bd-ae)} \right)}{9e(bd - ae)} - \frac{2(a + bx)^{5/2}(Bd - Ae)}{9e(d + ex)^{9/2}(bd - ae)}$$

$$\downarrow 48$$

$$\frac{\left(\frac{4b(a+bx)^{5/2}}{35(d+ex)^{5/2}(bd-ae)^2} + \frac{2(a+bx)^{5/2}}{7(d+ex)^{7/2}(bd-ae)} \right) (-9aBe + 4Abe + 5bBd)}{9e(bd - ae)} - \frac{2(a + bx)^{5/2}(Bd - Ae)}{9e(d + ex)^{9/2}(bd - ae)}$$

input

$$\text{Int}[(a + b*x)^(3/2)*(A + B*x)/(d + e*x)^(11/2), x]$$

output

$$\begin{aligned} & (-2*(B*d - A*e)*(a + b*x)^{(5/2)})/(9*e*(b*d - a*e)*(d + e*x)^{(9/2)}) + ((5*b \\ & *B*d + 4*A*b*e - 9*a*B*e)*((2*(a + b*x)^{(5/2)})/(7*(b*d - a*e)*(d + e*x)^{(7 \\ & /2)}) + (4*b*(a + b*x)^{(5/2)})/(35*(b*d - a*e)^2*(d + e*x)^{(5/2)}))/9*e*(b* \\ & d - a*e) \end{aligned}$$

Defintions of rubi rules used

rule 48

$$\text{Int}[(a_.) + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp} \\ [(a + b*x)^{(m + 1)*((c + d*x)^{(n + 1)})/(b*c - a*d)*(m + 1))], x] /; \text{FreeQ}\{ \\ a, b, c, d, m, n\}, x\} \ \&\& \ \text{EqQ}[m + n + 2, 0] \ \&\& \ \text{NeQ}[m, -1]$$

rule 55

$$\text{Int}[(a_.) + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[\\ (a + b*x)^{(m + 1)*((c + d*x)^{(n + 1)})/(b*c - a*d)*(m + 1))], x] - \text{Simp}[d*(S \\ \text{implify}[m + n + 2]/((b*c - a*d)*(m + 1))) \ \text{Int}[(a + b*x)^{\text{Simplify}[m + 1]}* \\ c + d*x)^n, x], x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x\} \ \&\& \ \text{ILtQ}[\text{Simplify}[m + n + \\ 2], 0] \ \&\& \ \text{NeQ}[m, -1] \ \&\& \ !(\text{LtQ}[m, -1] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ (\text{EqQ}[a, 0] \ || \ (\text{NeQ}[\\ c, 0] \ \&\& \ \text{LtQ}[m - n, 0] \ \&\& \ \text{IntegerQ}[n]))) \ \&\& \ (\text{SumSimplerQ}[m, 1] \ || \ !\text{SumSimp} \\ \text{lerQ}[n, 1])$$

rule 87

$$\text{Int}[(a_.) + (b_.)*(x_.)*((c_.) + (d_.)*(x_.))^{(n_.)*((e_.) + (f_.)*(x_.))^{(p \\ _.)}, x_] \rightarrow \text{Simp}[(-b*e - a*f)*(c + d*x)^{(n + 1)*((e + f*x)^{(p + 1)})/(f*(p \\ + 1)*(c*f - d*e))], x] - \text{Simp}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p \\ + 1)))/(f*(p + 1)*(c*f - d*e)) \ \text{Int}[(c + d*x)^n*(e + f*x)^{(p + 1)}, x], x] \\ /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x\} \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ (!\text{LtQ}[n, -1] \ || \ \text{Intege} \\ \text{rQ}[p] \ || \ !(\text{IntegerQ}[n] \ || \ !(\text{EqQ}[e, 0] \ || \ !(\text{EqQ}[c, 0] \ || \ \text{LtQ}[p, n])))$$

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.20

method	result
gospers	$-\frac{2(bx+a)^{\frac{5}{2}}(8Ab^2e^2x^2-18Bab^2e^2x^2+10Bb^2dex^2-20Aab^2e^2x+36Ab^2dex+45Ba^2e^2x-106Babdex+45b^2Bd^2x+35a^2Ae^2-90A^2a^2b^2e^2)}{315(ex+d)^{\frac{9}{2}}(a^3e^3-3a^2be^2d+3ab^2d^2e-b^3d^3)}$
orering	$-\frac{2(bx+a)^{\frac{5}{2}}(8Ab^2e^2x^2-18Bab^2e^2x^2+10Bb^2dex^2-20Aab^2e^2x+36Ab^2dex+45Ba^2e^2x-106Babdex+45b^2Bd^2x+35a^2Ae^2-90A^2a^2b^2e^2)}{315(ex+d)^{\frac{9}{2}}(a^3e^3-3a^2be^2d+3ab^2d^2e-b^3d^3)}$
default	$-\frac{2(8Ab^3e^2x^3-18Bab^2e^2x^3+10Bb^3dex^3-12Aab^2e^2x^2+36Ab^3dex^2+27Ba^2be^2x^2-96Bab^2dex^2+45Bb^3d^2x^2+15Aa^2be^2x-90A^2a^2b^2e^2)}{315(ex+d)^{\frac{9}{2}}(a^3e^3-3a^2be^2d+3ab^2d^2e-b^3d^3)}$

input

```
int((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(11/2),x,method=_RETURNVERBOSE)
```

output

```
-2/315*(b*x+a)^(5/2)*(8*A*b^2*e^2*x^2-18*B*a*b*e^2*x^2+10*B*b^2*d*e*x^2-20*A*a*b*e^2*x+36*A*b^2*d*e*x+45*B*a^2*e^2*x-106*B*a*b*d*e*x+45*B*b^2*d^2*x+35*A*a^2*e^2-90*A*a*b*d*e+63*A*b^2*d^2+10*B*a^2*d*e-18*B*a*b*d^2)/(e*x+d)^(9/2)/(a^3*e^3-3*a^2*b*d*e^2+3*a*b^2*d^2*e-b^3*d^3)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 567 vs. 2(129) = 258.

Time = 28.00 (sec) , antiderivative size = 567, normalized size of antiderivative = 3.86

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{11/2}} dx = \frac{2(35Aa^4e^2 + 2(5Bb^4de - (9Bab^3 - 4Ab^4)e^2)x^4 + (45Bb^4d^2 - 2(43Bab^3 - 35Aa^2e^2 - 90Aa^2b^2e^2 - 63Aa^2b^2d^2 + 10Bb^3d^2 + 10Bb^3d^2e - 18Bb^3d^2e^2 - 96Bb^3d^2e^2 + 45Bb^3d^2e^2 - a^3d^5e^3 + \dots))x^4 + \dots}{315(b^3d^8 - 3ab^2d^7e + 3a^2bd^6e^2 - a^3d^5e^3 + \dots)}$$

input

```
integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(11/2),x, algorithm="fricas")
```

output

```
2/315*(35*A*a^4*e^2 + 2*(5*B*b^4*d*e - (9*B*a*b^3 - 4*A*b^4)*e^2)*x^4 + (4
5*B*b^4*d^2 - 2*(43*B*a*b^3 - 18*A*b^4)*d*e + (9*B*a^2*b^2 - 4*A*a*b^3)*e^
2)*x^3 - 9*(2*B*a^3*b - 7*A*a^2*b^2)*d^2 + 10*(B*a^4 - 9*A*a^3*b)*d*e + 3*
(3*(8*B*a*b^3 + 7*A*b^4)*d^2 - 2*(32*B*a^2*b^2 + 3*A*a*b^3)*d*e + (24*B*a^
3*b + A*a^2*b^2)*e^2)*x^2 + (9*(B*a^2*b^2 + 14*A*a*b^3)*d^2 - 2*(43*B*a^3*
b + 72*A*a^2*b^2)*d*e + 5*(9*B*a^4 + 10*A*a^3*b)*e^2)*x)*sqrt(b*x + a)*sqr
t(e*x + d)/(b^3*d^8 - 3*a*b^2*d^7*e + 3*a^2*b*d^6*e^2 - a^3*d^5*e^3 + (b^3
*d^3*e^5 - 3*a*b^2*d^2*e^6 + 3*a^2*b*d*e^7 - a^3*e^8)*x^5 + 5*(b^3*d^4*e^4
- 3*a*b^2*d^3*e^5 + 3*a^2*b*d^2*e^6 - a^3*d*e^7)*x^4 + 10*(b^3*d^5*e^3 -
3*a*b^2*d^4*e^4 + 3*a^2*b*d^3*e^5 - a^3*d^2*e^6)*x^3 + 10*(b^3*d^6*e^2 - 3
*a*b^2*d^5*e^3 + 3*a^2*b*d^4*e^4 - a^3*d^3*e^5)*x^2 + 5*(b^3*d^7*e - 3*a*b
^2*d^6*e^2 + 3*a^2*b*d^5*e^3 - a^3*d^4*e^4)*x)
```

Sympy [F]

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{11/2}} dx = \int \frac{(A + Bx)(a + bx)^{\frac{3}{2}}}{(d + ex)^{\frac{11}{2}}} dx$$

input

```
integrate((b*x+a)**(3/2)*(B*x+A)/(e*x+d)**(11/2),x)
```

output

```
Integral((A + B*x)*(a + b*x)**(3/2)/(d + e*x)**(11/2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{11/2}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(11/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f
or more de
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 516 vs. $2(129) = 258$.

Time = 0.45 (sec) , antiderivative size = 516, normalized size of antiderivative = 3.51

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{11/2}} dx = \frac{2 \left((bx+a) \left(\frac{2(5Bb^{11}d^2e^5|b|-14Bab^{10}de^6|b|+4Ab^{11}de^6|b|+9Ba^2b^9e^7|b|-4Aab^{10}e^7|b|)(bx+a)}{b^6d^4e^4-4ab^5d^3e^5+6a^2b^4d^2e^6-4a^3b^3de^7+a^4b^2e^8} \right) \right)}{b^6d^4e^4-4ab^5d^3e^5+6a^2b^4d^2e^6-4a^3b^3de^7+a^4b^2e^8} +$$

input

```
integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(11/2),x, algorithm="giac")
```

output

```
2/315*((b*x + a)*(2*(5*B*b^11*d^2*e^5*abs(b) - 14*B*a*b^10*d*e^6*abs(b) +
4*A*b^11*d*e^6*abs(b) + 9*B*a^2*b^9*e^7*abs(b) - 4*A*a*b^10*e^7*abs(b))*(b
*x + a)/(b^6*d^4*e^4 - 4*a*b^5*d^3*e^5 + 6*a^2*b^4*d^2*e^6 - 4*a^3*b^3*d*e
^7 + a^4*b^2*e^8) + 9*(5*B*b^12*d^3*e^4*abs(b) - 19*B*a*b^11*d^2*e^5*abs(b
) + 4*A*b^12*d^2*e^5*abs(b) + 23*B*a^2*b^10*d*e^6*abs(b) - 8*A*a*b^11*d*e^
6*abs(b) - 9*B*a^3*b^9*e^7*abs(b) + 4*A*a^2*b^10*e^7*abs(b))/(b^6*d^4*e^4
- 4*a*b^5*d^3*e^5 + 6*a^2*b^4*d^2*e^6 - 4*a^3*b^3*d*e^7 + a^4*b^2*e^8)) -
63*(B*a*b^12*d^3*e^4*abs(b) - A*b^13*d^3*e^4*abs(b) - 3*B*a^2*b^11*d^2*e^5
*abs(b) + 3*A*a*b^12*d^2*e^5*abs(b) + 3*B*a^3*b^10*d*e^6*abs(b) - 3*A*a^2*
b^11*d*e^6*abs(b) - B*a^4*b^9*e^7*abs(b) + A*a^3*b^10*e^7*abs(b))/(b^6*d^4
*e^4 - 4*a*b^5*d^3*e^5 + 6*a^2*b^4*d^2*e^6 - 4*a^3*b^3*d*e^7 + a^4*b^2*e^8
))* (b*x + a)^(5/2)/(b^2*d + (b*x + a)*b*e - a*b*e)^(9/2)
```

Mupad [B] (verification not implemented)

Time = 1.65 (sec) , antiderivative size = 402, normalized size of antiderivative = 2.73

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{11/2}} dx = \frac{\sqrt{d+ex} \left(\frac{\sqrt{a+bx} (20Ba^4de+70Aa^4e^2-36Ba^3bd^2-180Aa^3bde+126Aa^2b^2d^2)}{315e^5(ae-bd)^3} + \frac{x^2\sqrt{a+bx} (144Ba^3be^2-384Ba^2b^2de+6Aa^2b^2d^2)}{315e^5} \right)}{1}$$

input `int(((A + B*x)*(a + b*x)^(3/2))/(d + e*x)^(11/2),x)`

output

$$\begin{aligned} & -((d + e*x)^{(1/2)} * (((a + b*x)^{(1/2)} * (70*A*a^4*e^2 + 20*B*a^4*d*e - 36*B*a^3*b*d^2 + 126*A*a^2*b^2*d^2 - 180*A*a^3*b*d*e)) / (315*e^5*(a*e - b*d)^3) + \\ & (x^2*(a + b*x)^{(1/2)} * (126*A*b^4*d^2 + 144*B*a*b^3*d^2 + 144*B*a^3*b*e^2 + 6*A*a^2*b^2*e^2 - 36*A*a*b^3*d*e - 384*B*a^2*b^2*d*e)) / (315*e^5*(a*e - b*d)^3) + \\ & (x*(a + b*x)^{(1/2)} * (90*B*a^4*e^2 + 252*A*a*b^3*d^2 + 100*A*a^3*b*e^2 + 18*B*a^2*b^2*d^2 - 172*B*a^3*b*d*e - 288*A*a^2*b^2*d*e)) / (315*e^5*(a*e - b*d)^3) + \\ & (4*b^3*x^4*(a + b*x)^{(1/2)} * (4*A*b*e - 9*B*a*e + 5*B*b*d)) / (315*e^4*(a*e - b*d)^3) - \\ & (2*b^2*x^3*(a*e - 9*b*d) * (a + b*x)^{(1/2)} * (4*A*b*e - 9*B*a*e + 5*B*b*d)) / (315*e^5*(a*e - b*d)^3)) / (x^5 + d^5/e^5 + (5*d*x^4)/e + (5*d^4*x)/e^4 + (10*d^2*x^3)/e^2 + (10*d^3*x^2)/e^3) \end{aligned}$$
Reduce [B] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 526, normalized size of antiderivative = 3.58

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{11/2}} dx = \frac{-\frac{2\sqrt{ex+d}\sqrt{bx+a}a^4e^5}{9} + \frac{2\sqrt{ex+d}\sqrt{bx+a}a^3bde^4}{7} - \frac{38\sqrt{ex+d}\sqrt{bx+a}a^3be^5x}{63} + \frac{6\sqrt{ex+d}\sqrt{bx+a}}{7}}{e^4(a^2e^7x^5 - 2abd e^6x^5 + b^2d^2e^5x^5 + 5a^2d e^6x^4 - 5abd e^5x^4 + b^2d^2e^4x^4 + 5a^2d e^5x^3 - 5abd e^4x^3 + b^2d^2e^3x^3 + 5a^2d e^4x^2 - 5abd e^3x^2 + b^2d^2e^2x^2 + 5a^2d e^3x - 5abd e^2x + b^2d^2e x + 5a^2d e^2 - 5abd e + b^2d^2)}$$

input `int((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(11/2),x)`

output

```
(2*( - 7*sqrt(d + e*x)*sqrt(a + b*x)*a**4*e**5 + 9*sqrt(d + e*x)*sqrt(a +
b*x)*a**3*b*d*e**4 - 19*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b*e**5*x + 27*sq
r
t(d + e*x)*sqrt(a + b*x)*a**2*b**2*d*e**4*x - 15*sqrt(d + e*x)*sqrt(a + b*
x)*a**2*b**2*e**5*x**2 + 27*sqrt(d + e*x)*sqrt(a + b*x)*a*b**3*d*e**4*x**2
- sqrt(d + e*x)*sqrt(a + b*x)*a*b**3*e**5*x**3 + 9*sqrt(d + e*x)*sqrt(a +
b*x)*b**4*d*e**4*x**3 + 2*sqrt(d + e*x)*sqrt(a + b*x)*b**4*e**5*x**4 - 2*
sqrt(e)*sqrt(b)*b**4*d**5 - 10*sqrt(e)*sqrt(b)*b**4*d**4*e*x - 20*sqrt(e)*
sqrt(b)*b**4*d**3*e**2*x**2 - 20*sqrt(e)*sqrt(b)*b**4*d**2*e**3*x**3 - 10*
sqrt(e)*sqrt(b)*b**4*d*e**4*x**4 - 2*sqrt(e)*sqrt(b)*b**4*e**5*x**5))/(63*
e**4*(a**2*d**5*e**2 + 5*a**2*d**4*e**3*x + 10*a**2*d**3*e**4*x**2 + 10*a*
*2*d**2*e**5*x**3 + 5*a**2*d*e**6*x**4 + a**2*e**7*x**5 - 2*a*b*d**6*e - 1
0*a*b*d**5*e**2*x - 20*a*b*d**4*e**3*x**2 - 20*a*b*d**3*e**4*x**3 - 10*a*b
*d**2*e**5*x**4 - 2*a*b*d*e**6*x**5 + b**2*d**7 + 5*b**2*d**6*e*x + 10*b**
2*d**5*e**2*x**2 + 10*b**2*d**4*e**3*x**3 + 5*b**2*d**3*e**4*x**4 + b**2*d
**2*e**5*x**5))
```


3.191 $\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{13/2}} dx$

Optimal result	1732
Mathematica [A] (verified)	1733
Rubi [A] (verified)	1733
Maple [A] (verified)	1735
Fricas [B] (verification not implemented)	1736
Sympy [F]	1736
Maxima [F(-2)]	1737
Giac [B] (verification not implemented)	1737
Mupad [B] (verification not implemented)	1738
Reduce [B] (verification not implemented)	1739

Optimal result

Integrand size = 24, antiderivative size = 201

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{13/2}} dx = -\frac{2(Bd - Ae)(a+bx)^{5/2}}{11e(bd - ae)(d+ex)^{11/2}} + \frac{2(5bBd + 6Abe - 11aBe)(a+bx)^{5/2}}{99e(bd - ae)^2(d+ex)^{9/2}} + \frac{8b(5bBd + 6Abe - 11aBe)(a+bx)^{5/2}}{693e(bd - ae)^3(d+ex)^{7/2}} + \frac{16b^2(5bBd + 6Abe - 11aBe)(a+bx)^{5/2}}{3465e(bd - ae)^4(d+ex)^{5/2}}$$

output

```
-2/11*(-A*e+B*d)*(b*x+a)^(5/2)/e/(-a*e+b*d)/(e*x+d)^(11/2)+2/99*(6*A*b*e-1
1*B*a*e+5*B*b*d)*(b*x+a)^(5/2)/e/(-a*e+b*d)^2/(e*x+d)^(9/2)+8/693*b*(6*A*b
*e-11*B*a*e+5*B*b*d)*(b*x+a)^(5/2)/e/(-a*e+b*d)^3/(e*x+d)^(7/2)+16/3465*b^
2*(6*A*b*e-11*B*a*e+5*B*b*d)*(b*x+a)^(5/2)/e/(-a*e+b*d)^4/(e*x+d)^(5/2)
```

Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 199, normalized size of antiderivative = 0.99

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{13/2}} dx = \frac{2(a+bx)^{5/2} (315Bde^2(a+bx)^3 - 315Ae^3(a+bx)^3 - 770bBde(a+bx)^2(d+ex) + 1155A^2b^2e^2(a+bx)^2(d+ex) - 385A^2B^2e^2(a+bx)^2(d+ex) + 495b^2B^2d^2(a+bx)(d+ex)^2 - 1485A^2b^2e^2(a+bx)(d+ex)^2 + 990a^2b^2B^2e^2(a+bx)(d+ex)^2 + 693A^2b^3(d+ex)^3 - 693a^2b^2B^2(d+ex)^3)}{(3465(bd-ae)^4(d+ex)^{11/2}}$$

input

```
Integrate[((a + b*x)^(3/2)*(A + B*x))/(d + e*x)^(13/2),x]
```

output

```
(2*(a + b*x)^(5/2)*(315*B*d*e^2*(a + b*x)^3 - 315*A*e^3*(a + b*x)^3 - 770*
b*B*d*e*(a + b*x)^2*(d + e*x) + 1155*A*b*e^2*(a + b*x)^2*(d + e*x) - 385*A
*B*e^2*(a + b*x)^2*(d + e*x) + 495*b^2*B*d*(a + b*x)*(d + e*x)^2 - 1485*A*
b^2*e*(a + b*x)*(d + e*x)^2 + 990*a*b*B*e*(a + b*x)*(d + e*x)^2 + 693*A*b^
3*(d + e*x)^3 - 693*a*b^2*B*(d + e*x)^3))/(3465*(b*d - a*e)^4*(d + e*x)^(1
1/2))
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.95, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {87, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{13/2}} dx$$

$$\downarrow 87$$

$$\frac{(-11aBe + 6Abe + 5bBd) \int \frac{(a+bx)^{3/2}}{(d+ex)^{11/2}} dx}{11e(bd-ae)} - \frac{2(a+bx)^{5/2}(Bd-Ae)}{11e(d+ex)^{11/2}(bd-ae)}$$

$$\downarrow 55$$

$$\frac{(-11aBe + 6Abe + 5bBd) \left(\frac{4b \int \frac{(a+bx)^{3/2}}{(d+ex)^{9/2}} dx}{9(bd-ae)} + \frac{2(a+bx)^{5/2}}{9(d+ex)^{9/2}(bd-ae)} \right)}{11e(bd-ae)} - \frac{2(a+bx)^{5/2}(Bd-Ae)}{11e(d+ex)^{11/2}(bd-ae)}$$

$$\begin{array}{c}
 \downarrow 55 \\
 (-11aBe + 6Abe + 5bBd) \left(\frac{4b \left(\frac{2b \int \frac{(a+bx)^{3/2} dx}{(d+ex)^{7/2}} + \frac{2(a+bx)^{5/2}}{7(d+ex)^{7/2}(bd-ae)} \right)}{9(bd-ae)} + \frac{2(a+bx)^{5/2}}{9(d+ex)^{9/2}(bd-ae)} \right) \\
 \hline
 \frac{11e(bd-ae)}{2(a+bx)^{5/2}(Bd-Ae)} \\
 \frac{2(a+bx)^{5/2}(Bd-Ae)}{11e(d+ex)^{11/2}(bd-ae)} \\
 \downarrow 48 \\
 \left(\frac{2(a+bx)^{5/2}}{9(d+ex)^{9/2}(bd-ae)} + \frac{4b \left(\frac{4b(a+bx)^{5/2}}{35(d+ex)^{5/2}(bd-ae)^2} + \frac{2(a+bx)^{5/2}}{7(d+ex)^{7/2}(bd-ae)} \right)}{9(bd-ae)} \right) (-11aBe + 6Abe + 5bBd) \\
 \hline
 \frac{11e(bd-ae)}{2(a+bx)^{5/2}(Bd-Ae)} \\
 \frac{2(a+bx)^{5/2}(Bd-Ae)}{11e(d+ex)^{11/2}(bd-ae)}
 \end{array}$$

input `Int[((a + b*x)^(3/2)*(A + B*x))/(d + e*x)^(13/2),x]`

output `(-2*(B*d - A*e)*(a + b*x)^(5/2))/(11*e*(b*d - a*e)*(d + e*x)^(11/2)) + ((5*b*B*d + 6*A*b*e - 11*a*B*e)*((2*(a + b*x)^(5/2))/(9*(b*d - a*e)*(d + e*x)^(9/2)) + (4*b*((2*(a + b*x)^(5/2))/(7*(b*d - a*e)*(d + e*x)^(7/2)) + (4*b*(a + b*x)^(5/2))/(35*(b*d - a*e)^2*(d + e*x)^(5/2))))/(9*(b*d - a*e)))/(11*e*(b*d - a*e))`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 55 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(Simplify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 322, normalized size of antiderivative = 1.60

method	result
gospers	$-\frac{2(bx+a)^{\frac{5}{2}}(-48Ab^3e^3x^3+88Bab^2e^3x^3-40Bb^3de^2x^3+120Aab^2e^3x^2-264Ab^3de^2x^2-220Ba^2be^3x^2+584Bab^2de^2x^2-220Bb^3de^2x^2)}{...}$
orering	$-\frac{2(bx+a)^{\frac{5}{2}}(-48Ab^3e^3x^3+88Bab^2e^3x^3-40Bb^3de^2x^3+120Aab^2e^3x^2-264Ab^3de^2x^2-220Ba^2be^3x^2+584Bab^2de^2x^2-220Bb^3de^2x^2)}{...}$
default	$-\frac{2(-48Ab^4e^3x^4+88Bab^3e^3x^4-40Bb^4de^2x^4+72Aab^3e^3x^3-264Ab^4de^2x^3-132Ba^2b^2e^3x^3+544Bab^3de^2x^3-220Bb^4de^2x^3-220Bb^4de^2x^3)}{...}$

input `int((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(13/2), x, method=_RETURNVERBOSE)`

output
$$-\frac{2}{3465}(b*x+a)^{\frac{5}{2}}*(-48*A*b^3*e^3*x^3+88*B*a*b^2*e^3*x^3-40*B*b^3*d*e^2*x^3+120*A*a*b^2*e^3*x^2-264*A*b^3*d*e^2*x^2-220*B*a^2*b*e^3*x^2+584*B*a*b^2*d*e^2*x^2-220*B*b^3*d^2*e*x^2-210*A*a^2*b*e^3*x+660*A*a*b^2*d*e^2*x-594*A*b^3*d^2*e*x+385*B*a^3*e^3*x-1385*B*a^2*b*d*e^2*x+1639*B*a*b^2*d^2*e*x-495*B*b^3*d^3*x+315*A*a^3*e^3-1155*A*a^2*b*d*e^2+1485*A*a*b^2*d^2*e-693*A*b^3*d^3+70*B*a^3*d*e^2-220*B*a^2*b*d^2*e+198*B*a*b^2*d^3)/(e*x+d)^(11/2)/(a^4*e^4-4*a^3*b*d*e^3+6*a^2*b^2*d^2*e^2-4*a*b^3*d^3*e+b^4*d^4)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 880 vs. $2(177) = 354$.

Time = 64.04 (sec) , antiderivative size = 880, normalized size of antiderivative = 4.38

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{13/2}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(13/2),x, algorithm="fricas")`

output

```
-2/3465*(315*A*a^5*e^3 - 8*(5*B*b^5*d*e^2 - (11*B*a*b^4 - 6*A*b^5)*e^3)*x^
5 - 4*(55*B*b^5*d^2*e - 6*(21*B*a*b^4 - 11*A*b^5)*d*e^2 + (11*B*a^2*b^3 -
6*A*a*b^4)*e^3)*x^4 + 99*(2*B*a^3*b^2 - 7*A*a^2*b^3)*d^3 - 55*(4*B*a^4*b -
27*A*a^3*b^2)*d^2*e + 35*(2*B*a^5 - 33*A*a^4*b)*d*e^2 - (495*B*b^5*d^3 -
11*(109*B*a*b^4 - 54*A*b^5)*d^2*e + (257*B*a^2*b^3 - 132*A*a*b^4)*d*e^2 -
3*(11*B*a^3*b^2 - 6*A*a^2*b^3)*e^3)*x^3 - (99*(8*B*a*b^4 + 7*A*b^5)*d^3 -
33*(86*B*a^2*b^3 + 9*A*a*b^4)*d^2*e + (2116*B*a^3*b^2 + 99*A*a^2*b^3)*d*e^
2 - 5*(110*B*a^4*b + 3*A*a^3*b^2)*e^3)*x^2 - (99*(B*a^2*b^3 + 14*A*a*b^4)*
d^3 - 11*(109*B*a^3*b^2 + 216*A*a^2*b^3)*d^2*e + 15*(83*B*a^4*b + 110*A*a^
3*b^2)*d*e^2 - 35*(11*B*a^5 + 12*A*a^4*b)*e^3)*x)*sqrt(b*x + a)*sqrt(e*x +
d)/(b^4*d^10 - 4*a*b^3*d^9*e + 6*a^2*b^2*d^8*e^2 - 4*a^3*b*d^7*e^3 + a^4*
d^6*e^4 + (b^4*d^4*e^6 - 4*a*b^3*d^3*e^7 + 6*a^2*b^2*d^2*e^8 - 4*a^3*b*d*e^
9 + a^4*e^10)*x^6 + 6*(b^4*d^5*e^5 - 4*a*b^3*d^4*e^6 + 6*a^2*b^2*d^3*e^7
- 4*a^3*b*d^2*e^8 + a^4*d*e^9)*x^5 + 15*(b^4*d^6*e^4 - 4*a*b^3*d^5*e^5 + 6
*a^2*b^2*d^4*e^6 - 4*a^3*b*d^3*e^7 + a^4*d^2*e^8)*x^4 + 20*(b^4*d^7*e^3 -
4*a*b^3*d^6*e^4 + 6*a^2*b^2*d^5*e^5 - 4*a^3*b*d^4*e^6 + a^4*d^3*e^7)*x^3 +
15*(b^4*d^8*e^2 - 4*a*b^3*d^7*e^3 + 6*a^2*b^2*d^6*e^4 - 4*a^3*b*d^5*e^5 +
a^4*d^4*e^6)*x^2 + 6*(b^4*d^9*e - 4*a*b^3*d^8*e^2 + 6*a^2*b^2*d^7*e^3 - 4
*a^3*b*d^6*e^4 + a^4*d^5*e^5)*x)
```

Sympy [F]

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{13/2}} dx = \int \frac{(A + Bx)(a + bx)^{\frac{3}{2}}}{(d + ex)^{\frac{13}{2}}} dx$$

input `integrate((b*x+a)**(3/2)*(B*x+A)/(e*x+d)**(13/2),x)`

output `Integral((A + B*x)*(a + b*x)**(3/2)/(d + e*x)**(13/2), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{13/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(13/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f or more de`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 815 vs. $2(177) = 354$.

Time = 0.50 (sec) , antiderivative size = 815, normalized size of antiderivative = 4.05

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{13/2}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(13/2),x, algorithm="giac")`

output

```

2/3465*((4*(b*x + a)*(2*(5*B*b^13*d^2*e^7*abs(b) - 16*B*a*b^12*d*e^8*abs(b)
) + 6*A*b^13*d*e^8*abs(b) + 11*B*a^2*b^11*e^9*abs(b) - 6*A*a*b^12*e^9*abs(
b))*(b*x + a)/(b^7*d^5*e^5 - 5*a*b^6*d^4*e^6 + 10*a^2*b^5*d^3*e^7 - 10*a^3
*b^4*d^2*e^8 + 5*a^4*b^3*d*e^9 - a^5*b^2*e^10) + 11*(5*B*b^14*d^3*e^6*abs(
b) - 21*B*a*b^13*d^2*e^7*abs(b) + 6*A*b^14*d^2*e^7*abs(b) + 27*B*a^2*b^12*
d*e^8*abs(b) - 12*A*a*b^13*d*e^8*abs(b) - 11*B*a^3*b^11*e^9*abs(b) + 6*A*a
^2*b^12*e^9*abs(b)))/(b^7*d^5*e^5 - 5*a*b^6*d^4*e^6 + 10*a^2*b^5*d^3*e^7 -
10*a^3*b^4*d^2*e^8 + 5*a^4*b^3*d*e^9 - a^5*b^2*e^10)) + 99*(5*B*b^15*d^4*e
^5*abs(b) - 26*B*a*b^14*d^3*e^6*abs(b) + 6*A*b^15*d^3*e^6*abs(b) + 48*B*a^
2*b^13*d^2*e^7*abs(b) - 18*A*a*b^14*d^2*e^7*abs(b) - 38*B*a^3*b^12*d*e^8*a
bs(b) + 18*A*a^2*b^13*d*e^8*abs(b) + 11*B*a^4*b^11*e^9*abs(b) - 6*A*a^3*b^
12*e^9*abs(b)))/(b^7*d^5*e^5 - 5*a*b^6*d^4*e^6 + 10*a^2*b^5*d^3*e^7 - 10*a^
3*b^4*d^2*e^8 + 5*a^4*b^3*d*e^9 - a^5*b^2*e^10))*(b*x + a) - 693*(B*a*b^15
*d^4*e^5*abs(b) - A*b^16*d^4*e^5*abs(b) - 4*B*a^2*b^14*d^3*e^6*abs(b) + 4*
A*a*b^15*d^3*e^6*abs(b) + 6*B*a^3*b^13*d^2*e^7*abs(b) - 6*A*a^2*b^14*d^2*e
^7*abs(b) - 4*B*a^4*b^12*d*e^8*abs(b) + 4*A*a^3*b^13*d*e^8*abs(b) + B*a^5*
b^11*e^9*abs(b) - A*a^4*b^12*e^9*abs(b))/(b^7*d^5*e^5 - 5*a*b^6*d^4*e^6 +
10*a^2*b^5*d^3*e^7 - 10*a^3*b^4*d^2*e^8 + 5*a^4*b^3*d*e^9 - a^5*b^2*e^10))
*(b*x + a)^(5/2)/(b^2*d + (b*x + a)*b*e - a*b*e)^(11/2)

```

Mupad [B] (verification not implemented)

Time = 1.81 (sec) , antiderivative size = 570, normalized size of antiderivative = 2.84

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{13/2}} dx =$$

$$\frac{\sqrt{d + ex} \left(\frac{\sqrt{a+bx} (140 B a^5 d e^2 + 630 A a^5 e^3 - 440 B a^4 b d^2 e - 2310 A a^4 b d e^2 + 396 B a^3 b^2 d^3 + 2970 A a^3 b^2 d^2 e - 1386 A a^2 b^3 d^3)}{3465 e^6 (a - b d)^4} + \frac{x \sqrt{d + ex}}{3465 e^6 (a - b d)^4} \right)}{(d + ex)^{13/2}}$$

input

```
int(((A + B*x)*(a + b*x)^(3/2))/(d + e*x)^(13/2),x)
```

output

```

-((d + e*x)^(1/2)*(((a + b*x)^(1/2)*(630*A*a^5*e^3 + 140*B*a^5*d*e^2 - 138
6*A*a^2*b^3*d^3 + 396*B*a^3*b^2*d^3 + 2970*A*a^3*b^2*d^2*e - 2310*A*a^4*b*
d*e^2 - 440*B*a^4*b*d^2*e)))/(3465*e^6*(a*e - b*d)^4) + (x*(a + b*x)^(1/2)*
(770*B*a^5*e^3 - 2772*A*a*b^4*d^3 + 840*A*a^4*b*e^3 - 198*B*a^2*b^3*d^3 +
4752*A*a^2*b^3*d^2*e - 3300*A*a^3*b^2*d*e^2 + 2398*B*a^3*b^2*d^2*e - 2490*
B*a^4*b*d*e^2))/(3465*e^6*(a*e - b*d)^4) - (x^2*(a + b*x)^(1/2)*(1386*A*b^
5*d^3 + 1584*B*a*b^4*d^3 - 1100*B*a^4*b*e^3 - 30*A*a^3*b^2*e^3 + 198*A*a^2
*b^3*d*e^2 - 5676*B*a^2*b^3*d^2*e + 4232*B*a^3*b^2*d*e^2 - 594*A*a*b^4*d^2
*e))/(3465*e^6*(a*e - b*d)^4) - (16*b^4*x^5*(a + b*x)^(1/2)*(6*A*b*e - 11*
B*a*e + 5*B*b*d))/(3465*e^4*(a*e - b*d)^4) + (8*b^3*x^4*(a*e - 11*b*d)*(a
+ b*x)^(1/2)*(6*A*b*e - 11*B*a*e + 5*B*b*d))/(3465*e^5*(a*e - b*d)^4) - (2
*b^2*x^3*(a + b*x)^(1/2)*(3*a^2*e^2 + 99*b^2*d^2 - 22*a*b*d*e)*(6*A*b*e -
11*B*a*e + 5*B*b*d))/(3465*e^6*(a*e - b*d)^4)))/(x^6 + d^6/e^6 + (6*d*x^5)
/e + (6*d^5*x)/e^5 + (15*d^2*x^4)/e^2 + (20*d^3*x^3)/e^3 + (15*d^4*x^2)/e^
4)

```

Reduce [B] (verification not implemented)

Time = 0.61 (sec) , antiderivative size = 860, normalized size of antiderivative = 4.28

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{13/2}} dx = \text{Too large to display}$$

input

```
int((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(13/2),x)
```


output

```
(2*( - 63*sqrt(d + e*x)*sqrt(a + b*x)*a**5*e**6 + 154*sqrt(d + e*x)*sqrt(a
+ b*x)*a**4*b*d*e**5 - 161*sqrt(d + e*x)*sqrt(a + b*x)*a**4*b*e**6*x - 99
*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b**2*d**2*e**4 + 418*sqrt(d + e*x)*sqrt(
a + b*x)*a**3*b**2*d*e**5*x - 113*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b**2*e*
**6*x**2 - 297*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**3*d**2*e**4*x + 330*sqrt
(d + e*x)*sqrt(a + b*x)*a**2*b**3*d*e**5*x**2 - 3*sqrt(d + e*x)*sqrt(a + b
*x)*a**2*b**3*e**6*x**3 - 297*sqrt(d + e*x)*sqrt(a + b*x)*a*b**4*d**2*e**4
*x**2 + 22*sqrt(d + e*x)*sqrt(a + b*x)*a*b**4*d*e**5*x**3 + 4*sqrt(d + e*x
)*sqrt(a + b*x)*a*b**4*e**6*x**4 - 99*sqrt(d + e*x)*sqrt(a + b*x)*b**5*d**
2*e**4*x**3 - 44*sqrt(d + e*x)*sqrt(a + b*x)*b**5*d*e**5*x**4 - 8*sqrt(d +
e*x)*sqrt(a + b*x)*b**5*e**6*x**5 + 8*sqrt(e)*sqrt(b)*b**5*d**6 + 48*sqrt
(e)*sqrt(b)*b**5*d**5*e*x + 120*sqrt(e)*sqrt(b)*b**5*d**4*e**2*x**2 + 160*
sqrt(e)*sqrt(b)*b**5*d**3*e**3*x**3 + 120*sqrt(e)*sqrt(b)*b**5*d**2*e**4*x
**4 + 48*sqrt(e)*sqrt(b)*b**5*d*e**5*x**5 + 8*sqrt(e)*sqrt(b)*b**5*e**6*x*
*6))/(693*e**4*(a**3*d**6*e**3 + 6*a**3*d**5*e**4*x + 15*a**3*d**4*e**5*x*
*2 + 20*a**3*d**3*e**6*x**3 + 15*a**3*d**2*e**7*x**4 + 6*a**3*d*e**8*x**5
+ a**3*e**9*x**6 - 3*a**2*b*d**7*e**2 - 18*a**2*b*d**6*e**3*x - 45*a**2*b*
d**5*e**4*x**2 - 60*a**2*b*d**4*e**5*x**3 - 45*a**2*b*d**3*e**6*x**4 - 18*
a**2*b*d**2*e**7*x**5 - 3*a**2*b*d*e**8*x**6 + 3*a*b**2*d**8*e + 18*a*b**2
*d**7*e**2*x + 45*a*b**2*d**6*e**3*x**2 + 60*a*b**2*d**5*e**4*x**3 + 45...
```

3.192 $\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{15/2}} dx$

Optimal result	1741
Mathematica [A] (verified)	1742
Rubi [A] (verified)	1742
Maple [B] (verified)	1744
Fricas [B] (verification not implemented)	1745
Sympy [F(-1)]	1746
Maxima [F(-2)]	1747
Giac [B] (verification not implemented)	1747
Mupad [B] (verification not implemented)	1748
Reduce [B] (verification not implemented)	1749

Optimal result

Integrand size = 24, antiderivative size = 255

$$\int \frac{(a+bx)^{3/2}(A+Bx)}{(d+ex)^{15/2}} dx = -\frac{2(Bd - Ae)(a+bx)^{5/2}}{13e(bd - ae)(d+ex)^{13/2}} + \frac{2(5bBd + 8Abe - 13aBe)(a+bx)^{5/2}}{143e(bd - ae)^2(d+ex)^{11/2}} + \frac{4b(5bBd + 8Abe - 13aBe)(a+bx)^{5/2}}{429e(bd - ae)^3(d+ex)^{9/2}} + \frac{16b^2(5bBd + 8Abe - 13aBe)(a+bx)^{5/2}}{3003e(bd - ae)^4(d+ex)^{7/2}} + \frac{32b^3(5bBd + 8Abe - 13aBe)(a+bx)^{5/2}}{15015e(bd - ae)^5(d+ex)^{5/2}}$$

output

```
-2/13*(-A*e+B*d)*(b*x+a)^(5/2)/e/(-a*e+b*d)/(e*x+d)^(13/2)+2/143*(8*A*b*e-13*B*a*e+5*B*b*d)*(b*x+a)^(5/2)/e/(-a*e+b*d)^2/(e*x+d)^(11/2)+4/429*b*(8*A*b*e-13*B*a*e+5*B*b*d)*(b*x+a)^(5/2)/e/(-a*e+b*d)^3/(e*x+d)^(9/2)+16/3003*b^2*(8*A*b*e-13*B*a*e+5*B*b*d)*(b*x+a)^(5/2)/e/(-a*e+b*d)^4/(e*x+d)^(7/2)+32/15015*b^3*(8*A*b*e-13*B*a*e+5*B*b*d)*(b*x+a)^(5/2)/e/(-a*e+b*d)^5/(e*x+d)^(5/2)
```


$$\begin{aligned} & \downarrow 55 \\ & (-13aBe + 8Abe + 5bBd) \left(\frac{6b \left(\frac{4b \int \frac{(a+bx)^{3/2}}{(d+ex)^{9/2}} dx}{9(bd-ae)} + \frac{2(a+bx)^{5/2}}{9(d+ex)^{9/2}(bd-ae)} \right)}{11(bd-ae)} + \frac{2(a+bx)^{5/2}}{11(d+ex)^{11/2}(bd-ae)} \right) \end{aligned}$$

$$\frac{13e(bd-ae)}{2(a+bx)^{5/2}(Bd-Ae)} \frac{2(a+bx)^{5/2}(Bd-Ae)}{13e(d+ex)^{13/2}(bd-ae)}$$

↓ 55

$$\begin{aligned} & (-13aBe + 8Abe + 5bBd) \left(\frac{6b \left(\frac{4b \left(\frac{2b \int \frac{(a+bx)^{3/2}}{(d+ex)^{7/2}} dx}{7(bd-ae)} + \frac{2(a+bx)^{5/2}}{7(d+ex)^{7/2}(bd-ae)} \right)}{9(bd-ae)} + \frac{2(a+bx)^{5/2}}{9(d+ex)^{9/2}(bd-ae)} \right)}{11(bd-ae)} + \frac{2(a+bx)^{5/2}}{11(d+ex)^{11/2}(bd-ae)} \right) \end{aligned}$$

$$\frac{13e(bd-ae)}{2(a+bx)^{5/2}(Bd-Ae)} \frac{2(a+bx)^{5/2}(Bd-Ae)}{13e(d+ex)^{13/2}(bd-ae)}$$

↓ 48

$$\begin{aligned} & \left(\frac{2(a+bx)^{5/2}}{11(d+ex)^{11/2}(bd-ae)} + \frac{6b \left(\frac{2(a+bx)^{5/2}}{9(d+ex)^{9/2}(bd-ae)} + \frac{4b \left(\frac{4b(a+bx)^{5/2}}{35(d+ex)^{5/2}(bd-ae)^2} + \frac{2(a+bx)^{5/2}}{7(d+ex)^{7/2}(bd-ae)} \right)}{9(bd-ae)} \right)}{11(bd-ae)} \right) (-13aBe + 8Abe + 5bBd) \end{aligned}$$

$$\frac{13e(bd-ae)}{2(a+bx)^{5/2}(Bd-Ae)} \frac{2(a+bx)^{5/2}(Bd-Ae)}{13e(d+ex)^{13/2}(bd-ae)}$$

input `Int[((a + b*x)^(3/2)*(A + B*x))/(d + e*x)^(15/2),x]`

output

$$\begin{aligned} & (-2*(B*d - A*e)*(a + b*x)^{(5/2)})/(13*e*(b*d - a*e)*(d + e*x)^{(13/2)}) + ((5 \\ & *b*B*d + 8*A*b*e - 13*a*B*e)*((2*(a + b*x)^{(5/2)})/(11*(b*d - a*e)*(d + e*x) \\ &)^{(11/2)}) + (6*b*((2*(a + b*x)^{(5/2)})/(9*(b*d - a*e)*(d + e*x)^{(9/2)}) + (4 \\ & *b*((2*(a + b*x)^{(5/2)})/(7*(b*d - a*e)*(d + e*x)^{(7/2)}) + (4*b*(a + b*x)^{(5/2)}) \\ &)/(35*(b*d - a*e)^2*(d + e*x)^{(5/2)})))/(9*(b*d - a*e)))/(11*(b*d - a* \\ & e)))/(13*e*(b*d - a*e)) \end{aligned}$$

Defintions of rubi rules used

rule 48

$$\text{Int}[(a_.) + (b_.)*(x_)^{(m_.)}*((c_.) + (d_.)*(x_)^{(n_.)}), x_Symbol] \rightarrow \text{Simp} \\ [(a + b*x)^{(m + 1)}*((c + d*x)^{(n + 1)})/((b*c - a*d)*(m + 1))], x] \text{ ; FreeQ}[\{ \\ a, b, c, d, m, n\}, x] \ \&\& \text{EqQ}[m + n + 2, 0] \ \&\& \text{NeQ}[m, -1]$$

rule 55

$$\text{Int}[(a_.) + (b_.)*(x_)^{(m_.)}*((c_.) + (d_.)*(x_)^{(n_.)}), x_Symbol] \rightarrow \text{Simp} [\\ (a + b*x)^{(m + 1)}*((c + d*x)^{(n + 1)})/((b*c - a*d)*(m + 1))], x] - \text{Simp}[d*(S \\ \text{simplify}[m + n + 2]/((b*c - a*d)*(m + 1))) \ \text{Int}[(a + b*x)^{\text{Simplify}[m + 1]}*(\\ c + d*x)^n, x], x] \text{ ; FreeQ}[\{a, b, c, d, m, n\}, x] \ \&\& \text{ILtQ}[\text{Simplify}[m + n + \\ 2], 0] \ \&\& \text{NeQ}[m, -1] \ \&\& \text{!(LtQ}[m, -1] \ \&\& \text{LtQ}[n, -1] \ \&\& (\text{EqQ}[a, 0] \ || (\text{NeQ}[\\ c, 0] \ \&\& \text{LtQ}[m - n, 0] \ \&\& \text{IntegerQ}[n]))) \ \&\& (\text{SumSimplerQ}[m, 1] \ || \ \text{!SumSimp} \\ \text{lerQ}[n, 1])$$

rule 87

$$\text{Int}[(a_.) + (b_.)*(x_)^{(m_.)}*((c_.) + (d_.)*(x_)^{(n_.)}*((e_.) + (f_.)*(x_)^{(p} \\ _.), x_] \rightarrow \text{Simp}[(-b*e - a*f)*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)})/(f*(p \\ + 1)*(c*f - d*e))], x] - \text{Simp}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p \\ + 1)))/(f*(p + 1)*(c*f - d*e)) \ \text{Int}[(c + d*x)^n*(e + f*x)^{(p + 1)}, x], x] \\ \text{ ; FreeQ}[\{a, b, c, d, e, f, n\}, x] \ \&\& \text{LtQ}[p, -1] \ \&\& (\text{!LtQ}[n, -1] \ || \ \text{Intege} \\ \text{rQ}[p] \ || \ \text{!(IntegerQ}[n] \ || \ \text{!(EqQ}[e, 0] \ || \ \text{!(EqQ}[c, 0] \ || \ \text{LtQ}[p, n]))))$$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 504 vs. $2(225) = 450$.

Time = 0.28 (sec) , antiderivative size = 505, normalized size of antiderivative = 1.98

output

```

2/15015*(1155*A*a^6*e^4 + 16*(5*B*b^6*d*e^3 - (13*B*a*b^5 - 8*A*b^6)*e^4)*
x^6 + 8*(65*B*b^6*d^2*e^2 - 2*(87*B*a*b^5 - 52*A*b^6)*d*e^3 + (13*B*a^2*b^
4 - 8*A*a*b^5)*e^4)*x^5 - 429*(2*B*a^3*b^3 - 7*A*a^2*b^4)*d^4 + 1430*(B*a^
4*b^2 - 6*A*a^3*b^3)*d^3*e - 910*(B*a^5*b - 11*A*a^4*b^2)*d^2*e^2 + 210*(B
*a^6 - 26*A*a^5*b)*d*e^3 + 2*(715*B*b^6*d^3*e - 13*(153*B*a*b^5 - 88*A*b^6
)*d^2*e^2 + (353*B*a^2*b^4 - 208*A*a*b^5)*d*e^3 - 3*(13*B*a^3*b^3 - 8*A*a^
2*b^4)*e^4)*x^4 + (2145*B*b^6*d^4 - 572*(11*B*a*b^5 - 6*A*b^6)*d^3*e + 26*
(79*B*a^2*b^4 - 44*A*a*b^5)*d^2*e^2 - 4*(133*B*a^3*b^3 - 78*A*a^2*b^4)*d*e
^3 + 5*(13*B*a^4*b^2 - 8*A*a^3*b^3)*e^4)*x^3 + (429*(8*B*a*b^5 + 7*A*b^6)*
d^4 - 1716*(9*B*a^2*b^4 + A*a*b^5)*d^3*e + 26*(662*B*a^3*b^3 + 33*A*a^2*b^
4)*d^2*e^2 - 20*(447*B*a^4*b^2 + 13*A*a^3*b^3)*d*e^3 + 35*(52*B*a^5*b + A
*a^4*b^2)*e^4)*x^2 + (429*(B*a^2*b^4 + 14*A*a*b^5)*d^4 - 572*(11*B*a^3*b^3
+ 24*A*a^2*b^4)*d^3*e + 650*(15*B*a^4*b^2 + 22*A*a^3*b^3)*d^2*e^2 - 140*(4
3*B*a^5*b + 52*A*a^4*b^2)*d*e^3 + 105*(13*B*a^6 + 14*A*a^5*b)*e^4)*x)*sqrt
(b*x + a)*sqrt(e*x + d)/(b^5*d^12 - 5*a*b^4*d^11*e + 10*a^2*b^3*d^10*e^2 -
10*a^3*b^2*d^9*e^3 + 5*a^4*b*d^8*e^4 - a^5*d^7*e^5 + (b^5*d^5*e^7 - 5*a*b
^4*d^4*e^8 + 10*a^2*b^3*d^3*e^9 - 10*a^3*b^2*d^2*e^10 + 5*a^4*b*d*e^11 - a
^5*e^12)*x^7 + 7*(b^5*d^6*e^6 - 5*a*b^4*d^5*e^7 + 10*a^2*b^3*d^4*e^8 - 10
a^3*b^2*d^3*e^9 + 5*a^4*b*d^2*e^10 - a^5*d*e^11)*x^6 + 21*(b^5*d^7*e^5 - 5
*a*b^4*d^6*e^6 + 10*a^2*b^3*d^5*e^7 - 10*a^3*b^2*d^4*e^8 + 5*a^4*b*d^3*...

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{15/2}} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**(3/2)*(B*x+A)/(e*x+d)**(15/2),x)
```

output

Timed out

Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{15/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(15/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f or more de`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1168 vs. 2(225) = 450.

Time = 0.67 (sec) , antiderivative size = 1168, normalized size of antiderivative = 4.58

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{15/2}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(15/2),x, algorithm="giac")`

output

```

2/15015*((2*(4*(b*x + a))*(2*(5*B*b^15*d^2*e^9*abs(b) - 18*B*a*b^14*d*e^10*
abs(b) + 8*A*b^15*d*e^10*abs(b) + 13*B*a^2*b^13*e^11*abs(b) - 8*A*a*b^14*e
^11*abs(b)))*(b*x + a)/(b^8*d^6*e^6 - 6*a*b^7*d^5*e^7 + 15*a^2*b^6*d^4*e^8
- 20*a^3*b^5*d^3*e^9 + 15*a^4*b^4*d^2*e^10 - 6*a^5*b^3*d*e^11 + a^6*b^2*e^
12) + 13*(5*B*b^16*d^3*e^8*abs(b) - 23*B*a*b^15*d^2*e^9*abs(b) + 8*A*b^16*
d^2*e^9*abs(b) + 31*B*a^2*b^14*d*e^10*abs(b) - 16*A*a*b^15*d*e^10*abs(b) -
13*B*a^3*b^13*e^11*abs(b) + 8*A*a^2*b^14*e^11*abs(b)))/(b^8*d^6*e^6 - 6*a*
b^7*d^5*e^7 + 15*a^2*b^6*d^4*e^8 - 20*a^3*b^5*d^3*e^9 + 15*a^4*b^4*d^2*e^1
0 - 6*a^5*b^3*d*e^11 + a^6*b^2*e^12)) + 143*(5*B*b^17*d^4*e^7*abs(b) - 28*
B*a*b^16*d^3*e^8*abs(b) + 8*A*b^17*d^3*e^8*abs(b) + 54*B*a^2*b^15*d^2*e^9*
abs(b) - 24*A*a*b^16*d^2*e^9*abs(b) - 44*B*a^3*b^14*d*e^10*abs(b) + 24*A*a
^2*b^15*d*e^10*abs(b) + 13*B*a^4*b^13*e^11*abs(b) - 8*A*a^3*b^14*e^11*abs(
b))/(b^8*d^6*e^6 - 6*a*b^7*d^5*e^7 + 15*a^2*b^6*d^4*e^8 - 20*a^3*b^5*d^3*e
^9 + 15*a^4*b^4*d^2*e^10 - 6*a^5*b^3*d*e^11 + a^6*b^2*e^12))*(b*x + a) + 4
29*(5*B*b^18*d^5*e^6*abs(b) - 33*B*a*b^17*d^4*e^7*abs(b) + 8*A*b^18*d^4*e^
7*abs(b) + 82*B*a^2*b^16*d^3*e^8*abs(b) - 32*A*a*b^17*d^3*e^8*abs(b) - 98*
B*a^3*b^15*d^2*e^9*abs(b) + 48*A*a^2*b^16*d^2*e^9*abs(b) + 57*B*a^4*b^14*d
*e^10*abs(b) - 32*A*a^3*b^15*d*e^10*abs(b) - 13*B*a^5*b^13*e^11*abs(b) + 8
*A*a^4*b^14*e^11*abs(b))/(b^8*d^6*e^6 - 6*a*b^7*d^5*e^7 + 15*a^2*b^6*d^4*e
^8 - 20*a^3*b^5*d^3*e^9 + 15*a^4*b^4*d^2*e^10 - 6*a^5*b^3*d*e^11 + a^6*...

```

Mupad [B] (verification not implemented)

Time = 2.07 (sec) , antiderivative size = 752, normalized size of antiderivative = 2.95

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{15/2}} dx =$$

$$\frac{\sqrt{d + ex} \left(\frac{\sqrt{a+bx} (420 B a^6 d e^3 + 2310 A a^6 e^4 - 1820 B a^5 b d^2 e^2 - 10920 A a^5 b d e^3 + 2860 B a^4 b^2 d^3 e + 20020 A a^4 b^2 d^2 e^2 - 1716 B a^3 b^3 d^2 e - 15015 e^7 (a e - b d)^5)}{15015 e^7 (a e - b d)^5} \right)}{15015 e^7 (a e - b d)^5}$$

input

```
int(((A + B*x)*(a + b*x)^(3/2))/(d + e*x)^(15/2),x)
```

output

```

-((d + e*x)^(1/2)*(((a + b*x)^(1/2)*(2310*A*a^6*e^4 + 420*B*a^6*d*e^3 + 60
06*A*a^2*b^4*d^4 - 1716*B*a^3*b^3*d^4 - 17160*A*a^3*b^3*d^3*e + 2860*B*a^4
*b^2*d^3*e - 1820*B*a^5*b*d^2*e^2 + 20020*A*a^4*b^2*d^2*e^2 - 10920*A*a^5*
b*d*e^3)))/(15015*e^7*(a*e - b*d)^5) + (x*(a + b*x)^(1/2)*(2730*B*a^6*e^4 +
12012*A*a*b^5*d^4 + 2940*A*a^5*b*e^4 + 858*B*a^2*b^4*d^4 - 27456*A*a^2*b^
4*d^3*e - 14560*A*a^4*b^2*d*e^3 - 12584*B*a^3*b^3*d^3*e + 28600*A*a^3*b^3*
d^2*e^2 + 19500*B*a^4*b^2*d^2*e^2 - 12040*B*a^5*b*d*e^3))/(15015*e^7*(a*e
- b*d)^5) + (x^2*(a + b*x)^(1/2)*(6006*A*b^6*d^4 + 6864*B*a*b^5*d^4 + 3640
*B*a^5*b*e^4 + 70*A*a^4*b^2*e^4 - 520*A*a^3*b^3*d*e^3 - 30888*B*a^2*b^4*d^
3*e - 17880*B*a^4*b^2*d*e^3 + 1716*A*a^2*b^4*d^2*e^2 + 34424*B*a^3*b^3*d^2
*e^2 - 3432*A*a*b^5*d^3*e))/(15015*e^7*(a*e - b*d)^5) + (32*b^5*x^6*(a + b
*x)^(1/2)*(8*A*b*e - 13*B*a*e + 5*B*b*d))/(15015*e^4*(a*e - b*d)^5) - (2*b
^2*x^3*(a + b*x)^(1/2)*(8*A*b*e - 13*B*a*e + 5*B*b*d)*(5*a^3*e^3 - 429*b^3
*d^3 + 143*a*b^2*d^2*e - 39*a^2*b*d*e^2))/(15015*e^7*(a*e - b*d)^5) - (16*
b^4*x^5*(a*e - 13*b*d)*(a + b*x)^(1/2)*(8*A*b*e - 13*B*a*e + 5*B*b*d))/(15
015*e^5*(a*e - b*d)^5) + (4*b^3*x^4*(a + b*x)^(1/2)*(3*a^2*e^2 + 143*b^2*d
^2 - 26*a*b*d*e)*(8*A*b*e - 13*B*a*e + 5*B*b*d))/(15015*e^6*(a*e - b*d)^5)
))/(x^7 + d^7/e^7 + (7*d*x^6)/e + (7*d^6*x)/e^6 + (21*d^2*x^5)/e^2 + (35*d
^3*x^4)/e^3 + (35*d^4*x^3)/e^4 + (21*d^5*x^2)/e^5)

```

Reduce [B] (verification not implemented)

Time = 1.18 (sec) , antiderivative size = 1253, normalized size of antiderivative = 4.91

$$\int \frac{(a + bx)^{3/2}(A + Bx)}{(d + ex)^{15/2}} dx = \text{Too large to display}$$

input

```
int((b*x+a)^(3/2)*(B*x+A)/(e*x+d)^(15/2),x)
```

output

```
(2*( - 231*sqrt(d + e*x)*sqrt(a + b*x)*a**6*e**7 + 819*sqrt(d + e*x)*sqrt(
a + b*x)*a**5*b*d*e**6 - 567*sqrt(d + e*x)*sqrt(a + b*x)*a**5*b*e**7*x - 1
001*sqrt(d + e*x)*sqrt(a + b*x)*a**4*b**2*d**2*e**5 + 2093*sqrt(d + e*x)*s
qrt(a + b*x)*a**4*b**2*d*e**6*x - 371*sqrt(d + e*x)*sqrt(a + b*x)*a**4*b**
2*e**7*x**2 + 429*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b**3*d**3*e**4 - 2717*s
qrt(d + e*x)*sqrt(a + b*x)*a**3*b**3*d**2*e**5*x + 1469*sqrt(d + e*x)*sqrt
(a + b*x)*a**3*b**3*d*e**6*x**2 - 5*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b**3*
e**7*x**3 + 1287*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**4*d**3*e**4*x - 2145*
sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**4*d**2*e**5*x**2 + 39*sqrt(d + e*x)*sq
rt(a + b*x)*a**2*b**4*d*e**6*x**3 + 6*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**
4*e**7*x**4 + 1287*sqrt(d + e*x)*sqrt(a + b*x)*a*b**5*d**3*e**4*x**2 - 143
*sqrt(d + e*x)*sqrt(a + b*x)*a*b**5*d**2*e**5*x**3 - 52*sqrt(d + e*x)*sqrt
(a + b*x)*a*b**5*d*e**6*x**4 - 8*sqrt(d + e*x)*sqrt(a + b*x)*a*b**5*e**7*x
**5 + 429*sqrt(d + e*x)*sqrt(a + b*x)*b**6*d**3*e**4*x**3 + 286*sqrt(d + e
*x)*sqrt(a + b*x)*b**6*d**2*e**5*x**4 + 104*sqrt(d + e*x)*sqrt(a + b*x)*b*
**6*d*e**6*x**5 + 16*sqrt(d + e*x)*sqrt(a + b*x)*b**6*e**7*x**6 - 16*sqrt(e
)*sqrt(b)*b**6*d**7 - 112*sqrt(e)*sqrt(b)*b**6*d**6*e*x - 336*sqrt(e)*sqrt
(b)*b**6*d**5*e**2*x**2 - 560*sqrt(e)*sqrt(b)*b**6*d**4*e**3*x**3 - 560*sq
rt(e)*sqrt(b)*b**6*d**3*e**4*x**4 - 336*sqrt(e)*sqrt(b)*b**6*d**2*e**5*x**
5 - 112*sqrt(e)*sqrt(b)*b**6*d*e**6*x**6 - 16*sqrt(e)*sqrt(b)*b**6*e**7...
```

3.193 $\int (a + bx)^{5/2}(A + Bx)(d + ex)^{5/2} dx$

Optimal result	1751
Mathematica [A] (verified)	1752
Rubi [A] (verified)	1753
Maple [B] (verified)	1762
Fricas [B] (verification not implemented)	1763
Sympy [F]	1764
Maxima [F(-2)]	1764
Giac [B] (verification not implemented)	1764
Mupad [F(-1)]	1765
Reduce [B] (verification not implemented)	1766

Optimal result

Integrand size = 24, antiderivative size = 398

$$\begin{aligned}
 & \int (a + bx)^{5/2}(A + Bx)(d + ex)^{5/2} dx = \\
 & - \frac{5(bd - ae)^5(bBd - 2Abe + aBe)\sqrt{a + bx}\sqrt{d + ex}}{1024b^4e^4} \\
 & + \frac{5(bd - ae)^4(bBd - 2Abe + aBe)(a + bx)^{3/2}\sqrt{d + ex}}{1536b^4e^3} \\
 & - \frac{(bd - ae)^3(bBd - 2Abe + aBe)(a + bx)^{5/2}\sqrt{d + ex}}{384b^4e^2} \\
 & - \frac{(bd - ae)^2(bBd - 2Abe + aBe)(a + bx)^{7/2}\sqrt{d + ex}}{64b^4e} \\
 & - \frac{(bd - ae)(bBd - 2Abe + aBe)(a + bx)^{7/2}(d + ex)^{3/2}}{24b^3e} \\
 & - \frac{(bBd - 2Abe + aBe)(a + bx)^{7/2}(d + ex)^{5/2}}{12b^2e} + \frac{B(a + bx)^{7/2}(d + ex)^{7/2}}{7be} \\
 & + \frac{5(bd - ae)^6(bBd - 2Abe + aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{1024b^{9/2}e^{9/2}}
 \end{aligned}$$

output

```
-5/1024*(-a*e+b*d)^5*(-2*A*b*e+B*a*e+B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(1/2)/b^4/e^4+5/1536*(-a*e+b*d)^4*(-2*A*b*e+B*a*e+B*b*d)*(b*x+a)^(3/2)*(e*x+d)^(1/2)/b^4/e^3-1/384*(-a*e+b*d)^3*(-2*A*b*e+B*a*e+B*b*d)*(b*x+a)^(5/2)*(e*x+d)^(1/2)/b^4/e^2-1/64*(-a*e+b*d)^2*(-2*A*b*e+B*a*e+B*b*d)*(b*x+a)^(7/2)*(e*x+d)^(1/2)/b^4/e-1/24*(-a*e+b*d)*(-2*A*b*e+B*a*e+B*b*d)*(b*x+a)^(7/2)*(e*x+d)^(3/2)/b^3/e-1/12*(-2*A*b*e+B*a*e+B*b*d)*(b*x+a)^(7/2)*(e*x+d)^(5/2)/b^2/e+1/7*B*(b*x+a)^(7/2)*(e*x+d)^(7/2)/b/e+5/1024*(-a*e+b*d)^6*(-2*A*b*e+B*a*e+B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)^(1/2))/b^(9/2)/e^(9/2)
```

Mathematica [A] (verified)

Time = 1.20 (sec) , antiderivative size = 514, normalized size of antiderivative = 1.29

$$\int (a + bx)^{5/2} (A + Bx) (d + ex)^{5/2} dx = \frac{(bd - ae)^6 \left(-\sqrt{b}\sqrt{e}\sqrt{a+bx}\sqrt{d+ex} (105bBde^6(a+bx)^6 - 210Abe^7(a+bx)^6 + 105aBe^7(a+bx)^6 - 700b^2Bde^5(a+bx)^5(d+ex) + ex)^{5/2} \right)}{21504b^{9/2}e^{9/2}}$$

input

```
Integrate[(a + b*x)^(5/2)*(A + B*x)*(d + e*x)^(5/2),x]
```

output

```
((b*d - a*e)^6*(-((Sqrt[b]*Sqrt[e]*Sqrt[a + b*x]*Sqrt[d + e*x]*(105*b*B*d*e^6*(a + b*x)^6 - 210*A*b*e^7*(a + b*x)^6 + 105*a*B*e^7*(a + b*x)^6 - 700*b^2*B*d*e^5*(a + b*x)^5*(d + e*x) + 1400*A*b^2*e^6*(a + b*x)^5*(d + e*x) - 700*a*b*B*e^6*(a + b*x)^5*(d + e*x) + 1981*b^3*B*d*e^4*(a + b*x)^4*(d + e*x)^2 - 3962*A*b^3*e^5*(a + b*x)^4*(d + e*x)^2 + 1981*a*b^2*B*e^5*(a + b*x)^4*(d + e*x)^2 + 3072*b^4*B*d*e^3*(a + b*x)^3*(d + e*x)^3 - 3072*a*b^3*B*e^4*(a + b*x)^3*(d + e*x)^3 - 1981*b^5*B*d*e^2*(a + b*x)^2*(d + e*x)^4 + 3962*A*b^5*e^3*(a + b*x)^2*(d + e*x)^4 - 1981*a*b^4*B*e^3*(a + b*x)^2*(d + e*x)^4 + 700*b^6*B*d*e*(a + b*x)*(d + e*x)^5 - 1400*A*b^6*e^2*(a + b*x)*(d + e*x)^5 + 700*a*b^5*B*e^2*(a + b*x)*(d + e*x)^5 - 105*b^7*B*d*(d + e*x)^6 + 210*A*b^7*e*(d + e*x)^6 - 105*a*b^6*B*e*(d + e*x)^6)))/(- (b*d) + a*e)^7) + 105*(b*B*d - 2*A*b*e + a*B*e)*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/(Sqrt[e]*Sqrt[a + b*x])])/(21504*b^(9/2)*e^(9/2))
```

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 334, normalized size of antiderivative = 0.84, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {90, 60, 60, 60, 60, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + bx)^{5/2} (A + Bx) (d + ex)^{5/2} dx \\
 & \quad \downarrow 90 \\
 & \frac{(2Abe - B(ae + bd)) \int (a + bx)^{5/2} (d + ex)^{5/2} dx}{2be} + \frac{B(a + bx)^{7/2} (d + ex)^{7/2}}{7be} \\
 & \quad \downarrow 60 \\
 & \frac{(2Abe - B(ae + bd)) \left(\frac{5(bd - ae) \int (a + bx)^{5/2} (d + ex)^{3/2} dx}{12b} + \frac{(a + bx)^{7/2} (d + ex)^{5/2}}{6b} \right)}{2be} + \\
 & \quad \frac{B(a + bx)^{7/2} (d + ex)^{7/2}}{7be} \\
 & \quad \downarrow 60 \\
 & \frac{(2Abe - B(ae + bd)) \left(\frac{5(bd - ae) \left(\frac{3(bd - ae) \int (a + bx)^{5/2} \sqrt{d + ex} dx}{10b} + \frac{(a + bx)^{7/2} (d + ex)^{3/2}}{5b} \right)}{12b} + \frac{(a + bx)^{7/2} (d + ex)^{5/2}}{6b} \right)}{2be} + \\
 & \quad \frac{B(a + bx)^{7/2} (d + ex)^{7/2}}{7be} \\
 & \quad \downarrow 60
 \end{aligned}$$

$$(2Abe - B(ae + bd)) \left(\frac{5(bd-ae) \left(\frac{3(bd-ae) \int \frac{(a+bx)^{5/2}}{\sqrt{d+ex}} dx + \frac{(a+bx)^{7/2} \sqrt{d+ex}}{4b}}{10b} \right) + \frac{(a+bx)^{7/2} (d+ex)^{3/2}}{5b}}{12b} \right) + \frac{(a+bx)^{7/2} (d+ex)^{5/2}}{6b}$$

$$\frac{B(a+bx)^{7/2} (d+ex)^{7/2}}{7be}$$

↓ 60

$$(2Abe - B(ae + bd)) \left(\frac{5(bd-ae) \left(\frac{3(bd-ae) \left(\frac{(a+bx)^{5/2} \sqrt{d+ex}}{3e} - \frac{5(bd-ae) \int \frac{(a+bx)^{3/2}}{\sqrt{d+ex}} dx}{6e} \right) + \frac{(a+bx)^{7/2} \sqrt{d+ex}}{4b}}{10b} \right) + \frac{(a+bx)^{7/2} (d+ex)^{5/2}}{5b}}{12b} \right)$$

$$\frac{B(a+bx)^{7/2} (d+ex)^{7/2}}{7be} \quad 2be$$

↓ 60

$$\begin{aligned}
 & \left(\frac{(bd-ae) \left(\frac{(a+bx)^{5/2} \sqrt{d+ex}}{3e} - \frac{5(bd-ae)}{6e} \left(\frac{(a+bx)^{3/2} \sqrt{d+ex}}{2e} - \frac{3(bd-ae) \int \frac{\sqrt{a+bx}}{\sqrt{d+ex}} dx}{4e} \right) \right)}{3(bd-ae) \cdot 8b} + \frac{(a+bx)^{7/2}}{4b} \right) \\
 & \frac{5(bd-ae)}{10b} \\
 & \frac{(2Abe - B(ae + bd))}{12b} \\
 & \frac{B(a+bx)^{7/2}(d+ex)^{7/2}}{7be} \\
 & \frac{2be}{7be} \\
 & \downarrow 60
 \end{aligned}$$

$$\begin{aligned}
 & \left(\frac{(a+bx)^{5/2} \sqrt{d+ex}}{3e} - \frac{(bd-ae)}{6e} \left(\frac{(a+bx)^{3/2} \sqrt{d+ex}}{2e} - \frac{3(bd-ae)}{4e} \left(\frac{\sqrt{a+bx} \sqrt{d+ex}}{e} - \frac{(bd-ae)}{4e} \right) \right) \right) \\
 & \quad \frac{3(bd-ae)}{8b} \\
 & \quad \frac{5(bd-ae)}{10b} \\
 & \quad (2Abe - B(ae + bd)) \frac{12b}{12b}
 \end{aligned}$$

↓ 66

↓ 221

$$\begin{aligned}
 & \left(\frac{(bd-ae)}{3e} \left(\frac{(a+bx)^{5/2} \sqrt{d+ex}}{3e} - \frac{5(bd-ae)}{2e} \left(\frac{(a+bx)^{3/2} \sqrt{d+ex}}{2e} - \frac{3(bd-ae)}{e} \left(\frac{\sqrt{a+bx} \sqrt{d+ex}}{e} - \frac{(bd-ae) \arctan\left(\frac{\sqrt{a+bx} \sqrt{d+ex}}{e}\right)}{4e} \right) \right) \right) \right) \\
 & \frac{3(bd-ae)}{8b} \\
 & \frac{5(bd-ae)}{10b} \\
 & (2Abe - B(ae + bd)) \frac{12b}{}
 \end{aligned}$$

input `Int[(a + b*x)^(5/2)*(A + B*x)*(d + e*x)^(5/2),x]`

output `(B*(a + b*x)^(7/2)*(d + e*x)^(7/2))/(7*b*e) + ((2*A*b*e - B*(b*d + a*e))*((a + b*x)^(7/2)*(d + e*x)^(5/2))/(6*b) + (5*(b*d - a*e)*((a + b*x)^(7/2)*(d + e*x)^(3/2))/(5*b) + (3*(b*d - a*e)*((a + b*x)^(7/2)*Sqrt[d + e*x])/(4*b) + ((b*d - a*e)*((a + b*x)^(5/2)*Sqrt[d + e*x])/(3*e) - (5*(b*d - a*e)*((a + b*x)^(3/2)*Sqrt[d + e*x])/(2*e) - (3*(b*d - a*e)*((Sqrt[a + b*x]*Sqrt[d + e*x])/e - ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(Sqrt[b]*e^(3/2))))/(4*e)))/(6*e)))/(8*b)))/(10*b)))/(12*b)))/(2*b*e)`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2395 vs. $2(342) = 684$.

Time = 0.29 (sec) , antiderivative size = 2396, normalized size of antiderivative = 6.02

method	result	size
default	Expression too large to display	2396

input `int((b*x+a)^(5/2)*(B*x+A)*(e*x+d)^(5/2),x,method=_RETURNVERBOSE)`

output

```
-1/43008*(b*x+a)^(1/2)*(e*x+d)^(1/2)*(210*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^6*b*e^7+210*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^7*d^6*e-5544*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^2*b^4*d^3*e^3-19680*B*a^2*b^4*d^2*e^4*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-512*B*a*b^5*d^3*e^3*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-37376*B*a*b^5*d*e^5*x^4*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-47488*A*a*b^5*d*e^5*x^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-25504*B*a^2*b^4*d*e^5*x^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-25504*B*a*b^5*d^2*e^4*x^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-35616*A*a^2*b^4*d*e^5*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-35616*A*a*b^5*d^2*e^4*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-512*B*a^3*b^3*d*e^5*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-1016*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^2*b^4*d^3*e^3*x+644*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b^5*d^4*e^2*x-1568*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^3*b^3*d*e^5*x-33264*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^2*b^4*d^2*e^4*x-1568*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b^5*d^3*e^3*x+644*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^4*b^2*d*e^5*x-1016*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^3*b^3*d^2*e^4*x-1260*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^5*b^2*d*e^6+3150*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^4*b^3*d^2*e^5-4200*A*ln(1/2*(2*b*e*x+2*((e*x...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 872 vs. $2(342) = 684$.

Time = 0.18 (sec) , antiderivative size = 1758, normalized size of antiderivative = 4.42

$$\int (a + bx)^{5/2} (A + Bx)(d + ex)^{5/2} dx = \text{Too large to display}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)*(e*x+d)^(5/2),x, algorithm="fricas")`

output

```
[-1/86016*(105*(B*b^7*d^7 - (5*B*a*b^6 + 2*A*b^7)*d^6*e + 3*(3*B*a^2*b^5 +
4*A*a*b^6)*d^5*e^2 - 5*(B*a^3*b^4 + 6*A*a^2*b^5)*d^4*e^3 - 5*(B*a^4*b^3 -
8*A*a^3*b^4)*d^3*e^4 + 3*(3*B*a^5*b^2 - 10*A*a^4*b^3)*d^2*e^5 - (5*B*a^6*b
b - 12*A*a^5*b^2)*d*e^6 + (B*a^7 - 2*A*a^6*b)*e^7)*sqrt(b*e)*log(8*b^2*e^2
*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 - 4*(2*b*e*x + b*d + a*e)*sqrt(b*e)*s
qrt(b*x + a)*sqrt(e*x + d) + 8*(b^2*d*e + a*b*e^2)*x) - 4*(3072*B*b^7*e^7*x
x^6 - 105*B*b^7*d^6*e + 70*(7*B*a*b^6 + 3*A*b^7)*d^5*e^2 - 7*(113*B*a^2*b^
5 + 170*A*a*b^6)*d^4*e^3 + 12*(25*B*a^3*b^4 + 231*A*a^2*b^5)*d^3*e^4 - 7*(
113*B*a^4*b^3 - 396*A*a^3*b^4)*d^2*e^5 + 70*(7*B*a^5*b^2 - 17*A*a^4*b^3)*d
*e^6 - 105*(B*a^6*b - 2*A*a^5*b^2)*e^7 + 256*(29*B*b^7*d*e^6 + (29*B*a*b^6
+ 14*A*b^7)*e^7)*x^5 + 128*(37*B*b^7*d^2*e^5 + 2*(73*B*a*b^6 + 35*A*b^7)*
d*e^6 + (37*B*a^2*b^5 + 70*A*a*b^6)*e^7)*x^4 + 16*(3*B*b^7*d^3*e^4 + (797*
B*a*b^6 + 378*A*b^7)*d^2*e^5 + (797*B*a^2*b^5 + 1484*A*a*b^6)*d*e^6 + 3*(B
*a^3*b^4 + 126*A*a^2*b^5)*e^7)*x^3 - 8*(7*B*b^7*d^4*e^3 - 2*(16*B*a*b^6 +
7*A*b^7)*d^3*e^4 - 6*(205*B*a^2*b^5 + 371*A*a*b^6)*d^2*e^5 - 2*(16*B*a^3*b
^4 + 1113*A*a^2*b^5)*d*e^6 + 7*(B*a^4*b^3 - 2*A*a^3*b^4)*e^7)*x^2 + 2*(35*
B*b^7*d^5*e^2 - 7*(23*B*a*b^6 + 10*A*b^7)*d^4*e^3 + 2*(127*B*a^2*b^5 + 196
*A*a*b^6)*d^3*e^4 + 2*(127*B*a^3*b^4 + 4158*A*a^2*b^5)*d^2*e^5 - 7*(23*B*a
^4*b^3 - 56*A*a^3*b^4)*d*e^6 + 35*(B*a^5*b^2 - 2*A*a^4*b^3)*e^7)*x)*sqrt(b
*x + a)*sqrt(e*x + d))/(b^5*e^5), -1/43008*(105*(B*b^7*d^7 - (5*B*a*b^6...
```


Sympy [F]

$$\int (a + bx)^{5/2} (A + Bx) (d + ex)^{5/2} dx = \int (A + Bx) (a + bx)^{5/2} (d + ex)^{5/2} dx$$

input `integrate((b*x+a)**(5/2)*(B*x+A)*(e*x+d)**(5/2),x)`

output `Integral((A + B*x)*(a + b*x)**(5/2)*(d + e*x)**(5/2), x)`

Maxima [F(-2)]

Exception generated.

$$\int (a + bx)^{5/2} (A + Bx) (d + ex)^{5/2} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)*(e*x+d)^(5/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6983 vs. $2(342) = 684$.

Time = 0.95 (sec) , antiderivative size = 6983, normalized size of antiderivative = 17.55

$$\int (a + bx)^{5/2} (A + Bx) (d + ex)^{5/2} dx = \text{Too large to display}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)*(e*x+d)^(5/2),x, algorithm="giac")`

output

```

1/107520*(1680*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e))*(2*(b*x + a)*(4*(b*x +
a)*(6*(b*x + a)/b^3 + (b^12*d*e^5 - 25*a*b^11*e^6)/(b^14*e^6)) - (5*b^13*
d^2*e^4 + 14*a*b^12*d*e^5 - 163*a^2*b^11*e^6)/(b^14*e^6)) + 3*(5*b^14*d^3*
e^3 + 9*a*b^13*d^2*e^4 + 15*a^2*b^12*d*e^5 - 93*a^3*b^11*e^6)/(b^14*e^6))*
sqrt(b*x + a) + 3*(5*b^4*d^4 + 4*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2 + 20*a^3*
b*d*e^3 - 35*a^4*e^4)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x
+ a)*b*e - a*b*e)))/(sqrt(b*e)*b^2*e^3))*B*a*d^2*abs(b) - 107520*((b^2*d
- a*b*e)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a
*b*e)))/sqrt(b*e) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*sqrt(b*x + a))*A*a
^3*d^2*abs(b)/b^2 + 560*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*(b*x + a)*
(4*(b*x + a)*(6*(b*x + a)/b^3 + (b^12*d*e^5 - 25*a*b^11*e^6)/(b^14*e^6)) -
(5*b^13*d^2*e^4 + 14*a*b^12*d*e^5 - 163*a^2*b^11*e^6)/(b^14*e^6)) + 3*(5*
b^14*d^3*e^3 + 9*a*b^13*d^2*e^4 + 15*a^2*b^12*d*e^5 - 93*a^3*b^11*e^6)/(b^
14*e^6))*sqrt(b*x + a) + 3*(5*b^4*d^4 + 4*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2
+ 20*a^3*b*d*e^3 - 35*a^4*e^4)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2
*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*b^2*e^3))*A*b*d^2*abs(b) + 3360*(
sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*(b*x + a)*(4*(b*x + a)*(6*(b*x + a)
/b^3 + (b^12*d*e^5 - 25*a*b^11*e^6)/(b^14*e^6)) - (5*b^13*d^2*e^4 + 14*a*b
^12*d*e^5 - 163*a^2*b^11*e^6)/(b^14*e^6)) + 3*(5*b^14*d^3*e^3 + 9*a*b^13*d
^2*e^4 + 15*a^2*b^12*d*e^5 - 93*a^3*b^11*e^6)/(b^14*e^6))*sqrt(b*x + a)...

```

Mupad [F(-1)]

Timed out.

$$\int (a + bx)^{5/2} (A + Bx) (d + ex)^{5/2} dx = \int (A + Bx) (a + bx)^{5/2} (d + ex)^{5/2} dx$$

input

```
int((A + B*x)*(a + b*x)^(5/2)*(d + e*x)^(5/2),x)
```

output

```
int((A + B*x)*(a + b*x)^(5/2)*(d + e*x)^(5/2), x)
```

Reduce [B] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 1098, normalized size of antiderivative = 2.76

$$\int (a + bx)^{5/2} (A + Bx)(d + ex)^{5/2} dx = \text{Too large to display}$$

input `int((b*x+a)^(5/2)*(B*x+A)*(e*x+d)^(5/2),x)`

output

```
(105*sqrt(d + e*x)*sqrt(a + b*x)*a**6*b*e**7 - 700*sqrt(d + e*x)*sqrt(a +
b*x)*a**5*b**2*d*e**6 - 70*sqrt(d + e*x)*sqrt(a + b*x)*a**5*b**2*e**7*x +
1981*sqrt(d + e*x)*sqrt(a + b*x)*a**4*b**3*d**2*e**5 + 462*sqrt(d + e*x)*s
qrt(a + b*x)*a**4*b**3*d*e**6*x + 56*sqrt(d + e*x)*sqrt(a + b*x)*a**4*b**3
*e**7*x**2 + 3072*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b**4*d**3*e**4 + 17140*
sqrt(d + e*x)*sqrt(a + b*x)*a**3*b**4*d**2*e**5*x + 18064*sqrt(d + e*x)*sq
rt(a + b*x)*a**3*b**4*d*e**6*x**2 + 6096*sqrt(d + e*x)*sqrt(a + b*x)*a**3*
b**4*e**7*x**3 - 1981*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**5*d**4*e**3 + 12
92*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**5*d**3*e**4*x + 27648*sqrt(d + e*x)
*sqrt(a + b*x)*a**2*b**5*d**2*e**5*x**2 + 36496*sqrt(d + e*x)*sqrt(a + b*x
)*a**2*b**5*d*e**6*x**3 + 13696*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**5*e**7
*x**4 + 700*sqrt(d + e*x)*sqrt(a + b*x)*a*b**6*d**5*e**2 - 462*sqrt(d + e
*x)*sqrt(a + b*x)*a*b**6*d**4*e**3*x + 368*sqrt(d + e*x)*sqrt(a + b*x)*a*b
**6*d**3*e**4*x**2 + 18800*sqrt(d + e*x)*sqrt(a + b*x)*a*b**6*d**2*e**5*x**
3 + 27648*sqrt(d + e*x)*sqrt(a + b*x)*a*b**6*d*e**6*x**4 + 11008*sqrt(d +
e*x)*sqrt(a + b*x)*a*b**6*e**7*x**5 - 105*sqrt(d + e*x)*sqrt(a + b*x)*b**7
*d**6*e + 70*sqrt(d + e*x)*sqrt(a + b*x)*b**7*d**5*e**2*x - 56*sqrt(d + e
*x)*sqrt(a + b*x)*b**7*d**4*e**3*x**2 + 48*sqrt(d + e*x)*sqrt(a + b*x)*b**7
*d**3*e**4*x**3 + 4736*sqrt(d + e*x)*sqrt(a + b*x)*b**7*d**2*e**5*x**4 + 7
424*sqrt(d + e*x)*sqrt(a + b*x)*b**7*d*e**6*x**5 + 3072*sqrt(d + e*x)*s...
```

3.194 $\int (a + bx)^{5/2}(A + Bx)(d + ex)^{3/2} dx$

Optimal result	1767
Mathematica [A] (verified)	1768
Rubi [A] (verified)	1769
Maple [B] (verified)	1775
Fricas [B] (verification not implemented)	1776
Sympy [F]	1777
Maxima [F(-2)]	1777
Giac [B] (verification not implemented)	1777
Mupad [F(-1)]	1778
Reduce [B] (verification not implemented)	1779

Optimal result

Integrand size = 24, antiderivative size = 358

$$\begin{aligned}
 & \int (a + bx)^{5/2}(A + Bx)(d + ex)^{3/2} dx = \\
 & - \frac{(bd - ae)^4(7bBd - 12Abe + 5aBe)\sqrt{a + bx}\sqrt{d + ex}}{512b^3e^4} \\
 & + \frac{(bd - ae)^3(7bBd - 12Abe + 5aBe)(a + bx)^{3/2}\sqrt{d + ex}}{768b^3e^3} \\
 & - \frac{(bd - ae)^2(7bBd - 12Abe + 5aBe)(a + bx)^{5/2}\sqrt{d + ex}}{960b^3e^2} \\
 & - \frac{(bd - ae)(7bBd - 12Abe + 5aBe)(a + bx)^{7/2}\sqrt{d + ex}}{160b^3e} \\
 & - \frac{(7bBd - 12Abe + 5aBe)(a + bx)^{7/2}(d + ex)^{3/2}}{60b^2e} + \frac{B(a + bx)^{7/2}(d + ex)^{5/2}}{6be} \\
 & + \frac{(bd - ae)^5(7bBd - 12Abe + 5aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{512b^{7/2}e^{9/2}}
 \end{aligned}$$

output

```
-1/512*(-a*e+b*d)^4*(-12*A*b*e+5*B*a*e+7*B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(1/2)
)/b^3/e^4+1/768*(-a*e+b*d)^3*(-12*A*b*e+5*B*a*e+7*B*b*d)*(b*x+a)^(3/2)*(e*
x+d)^(1/2)/b^3/e^3-1/960*(-a*e+b*d)^2*(-12*A*b*e+5*B*a*e+7*B*b*d)*(b*x+a)^(
5/2)*(e*x+d)^(1/2)/b^3/e^2-1/160*(-a*e+b*d)*(-12*A*b*e+5*B*a*e+7*B*b*d)*
(b*x+a)^(7/2)*(e*x+d)^(1/2)/b^3/e-1/60*(-12*A*b*e+5*B*a*e+7*B*b*d)*(b*x+a)^(
7/2)*(e*x+d)^(3/2)/b^2/e+1/6*B*(b*x+a)^(7/2)*(e*x+d)^(5/2)/b/e+1/512*(-a*
e+b*d)^5*(-12*A*b*e+5*B*a*e+7*B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)
/(e*x+d)^(1/2))/b^(7/2)/e^(9/2)
```

Mathematica [A] (verified)

Time = 0.95 (sec) , antiderivative size = 431, normalized size of antiderivative = 1.20

$$\int (a + bx)^{5/2} (A + Bx) (d + ex)^{3/2} dx = \frac{\sqrt{a + bx} \sqrt{d + ex} (75a^5 B e^5 - 5a^4 b e^4 (49Bd + 36Ae + 10Bex) + 10a^3 b^2 e^3 (12Ae(7d + ex) + B(bd - ae)^5 (7bBd - 12Abe + 5aBe) \operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right))}{512b^{7/2}e^{9/2}}$$

input

```
Integrate[(a + b*x)^(5/2)*(A + B*x)*(d + e*x)^(3/2),x]
```

output

```
(Sqrt[a + b*x]*Sqrt[d + e*x]*(75*a^5*B*e^5 - 5*a^4*b*e^4*(49*B*d + 36*A*e
+ 10*B*e*x) + 10*a^3*b^2*e^3*(12*A*e*(7*d + e*x) + B*(15*d^2 + 16*d*e*x +
4*e^2*x^2)) + 6*a^2*b^3*e^2*(4*A*e*(64*d^2 + 233*d*e*x + 124*e^2*x^2) + B*
(-91*d^3 + 58*d^2*e*x + 564*d*e^2*x^2 + 360*e^3*x^3)) + a*b^4*e*(24*A*e*(-
35*d^3 + 23*d^2*e*x + 256*d*e^2*x^2 + 168*e^3*x^3) + B*(415*d^4 - 272*d^3*
e*x + 216*d^2*e^2*x^2 + 4448*d*e^3*x^3 + 3200*e^4*x^4)) + b^5*(12*A*e*(15*
d^4 - 10*d^3*e*x + 8*d^2*e^2*x^2 + 176*d*e^3*x^3 + 128*e^4*x^4) + B*(-105*
d^5 + 70*d^4*e*x - 56*d^3*e^2*x^2 + 48*d^2*e^3*x^3 + 1664*d*e^4*x^4 + 1280
*e^5*x^5))))/(7680*b^3*e^4) + ((b*d - a*e)^5*(7*b*B*d - 12*A*b*e + 5*a*B*e)
)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])]/(512*b^(7/2)*e
^(9/2))
```

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 293, normalized size of antiderivative = 0.82, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {90, 60, 60, 60, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + bx)^{5/2} (A + Bx) (d + ex)^{3/2} dx \\
 & \quad \downarrow 90 \\
 & \frac{B(a + bx)^{7/2} (d + ex)^{5/2}}{6be} - \frac{(5aBe - 12Abe + 7bBd) \int (a + bx)^{5/2} (d + ex)^{3/2} dx}{12be} \\
 & \quad \downarrow 60 \\
 & \frac{B(a + bx)^{7/2} (d + ex)^{5/2}}{6be} - \\
 & \frac{(5aBe - 12Abe + 7bBd) \left(\frac{3(bd - ae) \int (a + bx)^{5/2} \sqrt{d + ex} dx}{10b} + \frac{(a + bx)^{7/2} (d + ex)^{3/2}}{5b} \right)}{12be} \\
 & \quad \downarrow 60 \\
 & \frac{B(a + bx)^{7/2} (d + ex)^{5/2}}{6be} - \\
 & \frac{(5aBe - 12Abe + 7bBd) \left(\frac{3(bd - ae) \left(\frac{(bd - ae) \int \frac{(a + bx)^{5/2}}{\sqrt{d + ex}} dx}{8b} + \frac{(a + bx)^{7/2} \sqrt{d + ex}}{4b} \right)}{10b} + \frac{(a + bx)^{7/2} (d + ex)^{3/2}}{5b} \right)}{12be} \\
 & \quad \downarrow 60
 \end{aligned}$$

$$\begin{array}{l}
 \frac{B(a+bx)^{7/2}(d+ex)^{5/2}}{6be} - \\
 (5aBe - 12Abe + 7bBd) \left(\frac{3(bd-ae) \left(\frac{(bd-ae) \left(\frac{(a+bx)^{5/2}\sqrt{d+ex}}{3e} - \frac{5(bd-ae) \int \frac{(a+bx)^{3/2}}{\sqrt{d+ex}} dx}{6e} \right)}{8b} + \frac{(a+bx)^{7/2}\sqrt{d+ex}}{4b} \right)}{10b} \right) + \frac{(a+bx)^{7/2}(d+ex)^{3/2}}{5b} \\
 \hline
 12be
 \end{array}$$

↓ 60

$$\begin{array}{l}
 \frac{B(a+bx)^{7/2}(d+ex)^{5/2}}{6be} - \\
 (5aBe - 12Abe + 7bBd) \left(\frac{3(bd-ae) \left(\frac{(bd-ae) \left(\frac{(a+bx)^{5/2}\sqrt{d+ex}}{3e} - \frac{5(bd-ae) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \int \frac{\sqrt{a+bx}}{\sqrt{d+ex}} dx}{4e} \right)}{6e} \right)}{8b} + \frac{(a+bx)^{7/2}\sqrt{d+ex}}{4b} \right)}{10b} \right) \\
 \hline
 12be
 \end{array}$$

↓ 60

$$\frac{B(a+bx)^{7/2}(d+ex)^{5/2}}{6be} - \left(\frac{(bd-ae)}{3(bd-ae)} \left(\frac{(a+bx)^{5/2}\sqrt{d+ex}}{3e} - \frac{5(bd-ae)}{2e} \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae)}{4e} \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae)\int\frac{\sqrt{a+bx}}{2e}}{4e} \right) \right) \right) \right) \frac{3(bd-ae)}{8b} + \frac{(5aBe - 12Abe + 7bBd)}{10b}$$

12be

$$\begin{aligned}
 & \frac{B(a+bx)^{7/2}(d+ex)^{5/2}}{6be} - \\
 & \left(\frac{(bd-ae)}{3e} \left(\frac{(a+bx)^{5/2}\sqrt{d+ex}}{3e} - \frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} \right) - \frac{(bd-ae) \int \frac{1}{b-\frac{d+ex}{e}}}{4e} \right) - \\
 & \frac{3(bd-ae)}{8b} - \\
 & \frac{(5aBe - 12Abe + 7bBd)}{10b} - \\
 & \frac{12be}{12be}
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow 221 \\
 \frac{B(a+bx)^{7/2}(d+ex)^{5/2}}{6be} - \\
 \left(\frac{(bd-ae) \left(\frac{(a+bx)^{5/2}\sqrt{d+ex}}{3e} - \frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} \right)}{3(bd-ae)} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae)\arctan\left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{\sqrt{be}}\right)}{4e} \right)}{8b} \right) \\
 \frac{(5aBe - 12Abe + 7bBd)}{10b} \\
 \hline
 12be
 \end{array}$$

input `Int[(a + b*x)^(5/2)*(A + B*x)*(d + e*x)^(3/2),x]`

output

$$\begin{aligned} & (B*(a + b*x)^{(7/2)}*(d + e*x)^{(5/2)})/(6*b*e) - ((7*b*B*d - 12*A*b*e + 5*a*B \\ & *e)*(((a + b*x)^{(7/2)}*(d + e*x)^{(3/2)})/(5*b) + (3*(b*d - a*e)*((a + b*x)^{ \\ & (7/2)*Sqrt[d + e*x])/(4*b) + ((b*d - a*e)*((a + b*x)^{(5/2)*Sqrt[d + e*x]) \\ & / (3*e) - (5*(b*d - a*e)*((a + b*x)^{(3/2)*Sqrt[d + e*x])/(2*e) - (3*(b*d - \\ & a*e)*((Sqrt[a + b*x]*Sqrt[d + e*x])/e - ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqr \\ & t[a + b*x])/(Sqrt[b]*Sqrt[d + e*x]))/(Sqrt[b]*e^{(3/2)})))/(4*e)))/(6*e)))/(\\ & (8*b)))/(10*b)))/(12*b*e) \end{aligned}$$
Defintions of rubi rules used

rule 60

$$\begin{aligned} & \text{Int}[(a_.) + (b_.)*(x_.)]^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \text{ :> } \text{Simp}[\\ & (a + b*x)^{(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + \text{Simp}[n*((b*c - a*d)/(\\ & b*(m + n + 1)) \quad \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] \text{ /; } \text{FreeQ}\{a, b, \\ & c, d\}, x] \&\& \text{GtQ}[n, 0] \&\& \text{NeQ}[m + n + 1, 0] \&\& !(\text{IGtQ}[m, 0] \&\& (!\text{Integer} \\ & \text{Q}[n] \text{ || } (\text{GtQ}[m, 0] \&\& \text{LtQ}[m - n, 0])) \&\& !\text{ILtQ}[m + n + 2, 0] \&\& \text{IntLinear} \\ & \text{Q}[a, b, c, d, m, n, x] \end{aligned}$$

rule 66

$$\begin{aligned} & \text{Int}[1/(Sqrt[(a_) + (b_.)*(x_.)]*Sqrt[(c_) + (d_.)*(x_.)]), x_Symbol] \text{ :> } \text{Simp}[\\ & 2 \quad \text{Subst}[\text{Int}[1/(b - d*x^2), x], x, \text{Sqrt}[a + b*x]/\text{Sqrt}[c + d*x]], x] \text{ /; } \text{Fre} \\ & \text{eQ}\{a, b, c, d\}, x] \&\& !\text{GtQ}[c - a*(d/b), 0] \end{aligned}$$

rule 90

$$\begin{aligned} & \text{Int}[(a_.) + (b_.)*(x_.)*((c_.) + (d_.)*(x_.))^{(n_.)*((e_.) + (f_.)*(x_.))^{(p} \\ & _.), x_] \text{ :> } \text{Simp}[b*(c + d*x)^{(n + 1)*((e + f*x)^{(p + 1)/(d*f*(n + p + 2))}, \\ & x] + \text{Simp}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p \\ & + 2)) \quad \text{Int}[(c + d*x)^n*(e + f*x)^p, x], x] \text{ /; } \text{FreeQ}\{a, b, c, d, e, f, n, \\ & p\}, x] \&\& \text{NeQ}[n + p + 2, 0] \end{aligned}$$

rule 221

$$\begin{aligned} & \text{Int}[(a_) + (b_.)*(x_)^2]^{(-1)}, x_Symbol] \text{ :> } \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x \\ & / \text{Rt}[-a/b, 2]], x] \text{ /; } \text{FreeQ}\{a, b\}, x] \&\& \text{NegQ}[a/b] \end{aligned}$$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1847 vs. $2(308) = 616$.

Time = 0.26 (sec) , antiderivative size = 1848, normalized size of antiderivative = 5.16

method	result	size
default	Expression too large to display	1848

input `int((b*x+a)^(5/2)*(B*x+A)*(e*x+d)^(3/2),x,method=_RETURNVERBOSE)`

output

```
1/15360*(b*x+a)^(1/2)*(e*x+d)^(1/2)*(240*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(
1/2)*a^3*b^2*e^5*x-240*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^5*d^3*e^2*x
-100*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^4*b*e^5*x+140*B*((e*x+d)*(b*x
+a))^(1/2)*(b*e)^(1/2)*b^5*d^4*e*x+8896*B*a*b^4*d*e^4*x^3*((e*x+d)*(b*x+a)
)^(1/2)*(b*e)^(1/2)+12288*A*a*b^4*d*e^4*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(
1/2)+6768*B*a^2*b^3*d*e^4*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+432*B*a
*b^4*d^2*e^3*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+11184*A*((e*x+d)*(b*x
+a))^(1/2)*(b*e)^(1/2)*a^2*b^3*d*e^4*x+1104*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)
^(1/2)*a*b^4*d^2*e^3*x+320*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^3*b^2*
d*e^4*x+696*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^2*b^3*d^2*e^3*x-544*B*
((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b^4*d^3*e^2*x+900*A*ln(1/2*(2*b*e*x+
2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^5*d^4*e^2+
270*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)
^(1/2))*a^5*b*d*e^5+2560*B*b^5*e^5*x^5*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)
+3072*A*b^5*e^5*x^4*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+96*B*b^5*d^2*e^3*x
^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+180*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b
*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^5*b*e^6-180*A*ln(1/2*(2*b
*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^6*d^5*e
+150*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^5*e^5-210*B*((e*x+d)*(b*x+a))
^(1/2)*(b*e)^(1/2)*b^5*d^5-300*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 687 vs. $2(308) = 616$.

Time = 0.17 (sec) , antiderivative size = 1388, normalized size of antiderivative = 3.88

$$\int (a + bx)^{5/2}(A + Bx)(d + ex)^{3/2} dx = \text{Too large to display}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)*(e*x+d)^(3/2),x, algorithm="fricas")`

output

```
[1/30720*(15*(7*B*b^6*d^6 - 6*(5*B*a*b^5 + 2*A*b^6)*d^5*e + 15*(3*B*a^2*b^4 + 4*A*a*b^5)*d^4*e^2 - 20*(B*a^3*b^3 + 6*A*a^2*b^4)*d^3*e^3 - 15*(B*a^4*b^2 - 8*A*a^3*b^3)*d^2*e^4 + 6*(3*B*a^5*b - 10*A*a^4*b^2)*d*e^5 - (5*B*a^6 - 12*A*a^5*b)*e^6)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*b*e*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x + a)*sqrt(e*x + d) + 8*(b^2*d*e + a*b*e^2)*x) + 4*(1280*B*b^6*e^6*x^5 - 105*B*b^6*d^5*e + 5*(83*B*a*b^5 + 36*A*b^6)*d^4*e^2 - 42*(13*B*a^2*b^4 + 20*A*a*b^5)*d^3*e^3 + 6*(2*5*B*a^3*b^3 + 256*A*a^2*b^4)*d^2*e^4 - 35*(7*B*a^4*b^2 - 24*A*a^3*b^3)*d*e^5 + 15*(5*B*a^5*b - 12*A*a^4*b^2)*e^6 + 128*(13*B*b^6*d*e^5 + (25*B*a*b^5 + 12*A*b^6)*e^6)*x^4 + 16*(3*B*b^6*d^2*e^4 + 2*(139*B*a*b^5 + 66*A*b^6)*d*e^5 + 9*(15*B*a^2*b^4 + 28*A*a*b^5)*e^6)*x^3 - 8*(7*B*b^6*d^3*e^3 - 3*(9*B*a*b^5 + 4*A*b^6)*d^2*e^4 - 3*(141*B*a^2*b^4 + 256*A*a*b^5)*d*e^5 - (5*B*a^3*b^3 + 372*A*a^2*b^4)*e^6)*x^2 + 2*(35*B*b^6*d^4*e^2 - 4*(34*B*a*b^5 + 15*A*b^6)*d^3*e^3 + 6*(29*B*a^2*b^4 + 46*A*a*b^5)*d^2*e^4 + 4*(20*B*a^3*b^3 + 699*A*a^2*b^4)*d*e^5 - 5*(5*B*a^4*b^2 - 12*A*a^3*b^3)*e^6)*x)*sqrt(b*x + a)*sqrt(e*x + d)/(b^4*e^5), -1/15360*(15*(7*B*b^6*d^6 - 6*(5*B*a*b^5 + 2*A*b^6)*d^5*e + 15*(3*B*a^2*b^4 + 4*A*a*b^5)*d^4*e^2 - 20*(B*a^3*b^3 + 6*A*a^2*b^4)*d^3*e^3 - 15*(B*a^4*b^2 - 8*A*a^3*b^3)*d^2*e^4 + 6*(3*B*a^5*b - 10*A*a^4*b^2)*d*e^5 - (5*B*a^6 - 12*A*a^5*b)*e^6)*sqrt(-b*e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*e^2*x...
```

Sympy [F]

$$\int (a + bx)^{5/2} (A + Bx) (d + ex)^{3/2} dx = \int (A + Bx) (a + bx)^{5/2} (d + ex)^{3/2} dx$$

input `integrate((b*x+a)**(5/2)*(B*x+A)*(e*x+d)**(3/2),x)`

output `Integral((A + B*x)*(a + b*x)**(5/2)*(d + e*x)**(3/2), x)`

Maxima [F(-2)]

Exception generated.

$$\int (a + bx)^{5/2} (A + Bx) (d + ex)^{3/2} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)*(e*x+d)^(3/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3955 vs. 2(308) = 616.

Time = 0.61 (sec) , antiderivative size = 3955, normalized size of antiderivative = 11.05

$$\int (a + bx)^{5/2} (A + Bx) (d + ex)^{3/2} dx = \text{Too large to display}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)*(e*x+d)^(3/2),x, algorithm="giac")`

output

```

1/7680*(120*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e))*(2*(b*x + a)*(4*(b*x + a)
*(6*(b*x + a)/b^3 + (b^12*d*e^5 - 25*a*b^11*e^6)/(b^14*e^6)) - (5*b^13*d^2
*e^4 + 14*a*b^12*d*e^5 - 163*a^2*b^11*e^6)/(b^14*e^6)) + 3*(5*b^14*d^3*e^3
+ 9*a*b^13*d^2*e^4 + 15*a^2*b^12*d*e^5 - 93*a^3*b^11*e^6)/(b^14*e^6))*sqr
t(b*x + a) + 3*(5*b^4*d^4 + 4*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2 + 20*a^3*b*d
*e^3 - 35*a^4*e^4)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x +
a)*b*e - a*b*e)))/(sqrt(b*e)*b^2*e^3))*B*a*d*abs(b) - 7680*((b^2*d - a*b*e
)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))
/sqrt(b*e) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*sqrt(b*x + a))*A*a^3*d*ab
s(b)/b^2 + 40*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e))*(2*(b*x + a)*(4*(b*x +
a)*(6*(b*x + a)/b^3 + (b^12*d*e^5 - 25*a*b^11*e^6)/(b^14*e^6)) - (5*b^13*d^2
*e^4 + 14*a*b^12*d*e^5 - 163*a^2*b^11*e^6)/(b^14*e^6)) + 3*(5*b^14*d^3*e^3
+ 9*a*b^13*d^2*e^4 + 15*a^2*b^12*d*e^5 - 93*a^3*b^11*e^6)/(b^14*e^6))*s
qrt(b*x + a) + 3*(5*b^4*d^4 + 4*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2 + 20*a^3*b
*d*e^3 - 35*a^4*e^4)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x
+ a)*b*e - a*b*e)))/(sqrt(b*e)*b^2*e^3))*A*b*d*abs(b) + 120*(sqrt(b^2*d +
(b*x + a)*b*e - a*b*e))*(2*(b*x + a)*(4*(b*x + a)*(6*(b*x + a)/b^3 + (b^12*
d*e^5 - 25*a*b^11*e^6)/(b^14*e^6)) - (5*b^13*d^2*e^4 + 14*a*b^12*d*e^5 - 1
63*a^2*b^11*e^6)/(b^14*e^6)) + 3*(5*b^14*d^3*e^3 + 9*a*b^13*d^2*e^4 + 15*a
^2*b^12*d*e^5 - 93*a^3*b^11*e^6)/(b^14*e^6))*sqrt(b*x + a) + 3*(5*b^4*d...

```

Mupad [F(-1)]

Timed out.

$$\int (a + bx)^{5/2} (A + Bx) (d + ex)^{3/2} dx = \int (A + Bx) (a + bx)^{5/2} (d + ex)^{3/2} dx$$

input

```
int((A + B*x)*(a + b*x)^(5/2)*(d + e*x)^(3/2), x)
```

output

```
int((A + B*x)*(a + b*x)^(5/2)*(d + e*x)^(3/2), x)
```

Reduce [B] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 860, normalized size of antiderivative = 2.40

$$\int (a + bx)^{5/2}(A + Bx)(d + ex)^{3/2} dx = \text{Too large to display}$$

input `int((b*x+a)^(5/2)*(B*x+A)*(e*x+d)^(3/2),x)`

output

```
( - 105*sqrt(d + e*x)*sqrt(a + b*x)*a**5*b*e**6 + 595*sqrt(d + e*x)*sqrt(a
+ b*x)*a**4*b**2*d*e**5 + 70*sqrt(d + e*x)*sqrt(a + b*x)*a**4*b**2*e**6*x
+ 1686*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b**3*d**2*e**4 + 5752*sqrt(d + e
*x)*sqrt(a + b*x)*a**3*b**3*d*e**5*x + 3016*sqrt(d + e*x)*sqrt(a + b*x)*a**
3*b**3*e**6*x**2 - 1386*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**4*d**3*e**3 +
900*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**4*d**2*e**4*x + 9528*sqrt(d + e*x)
*sqrt(a + b*x)*a**2*b**4*d*e**5*x**2 + 6192*sqrt(d + e*x)*sqrt(a + b*x)*a
**2*b**4*e**6*x**3 + 595*sqrt(d + e*x)*sqrt(a + b*x)*a*b**5*d**4*e**2 - 392
*sqrt(d + e*x)*sqrt(a + b*x)*a*b**5*d**3*e**3*x + 312*sqrt(d + e*x)*sqrt(a
+ b*x)*a*b**5*d**2*e**4*x**2 + 6560*sqrt(d + e*x)*sqrt(a + b*x)*a*b**5*d*
e**5*x**3 + 4736*sqrt(d + e*x)*sqrt(a + b*x)*a*b**5*e**6*x**4 - 105*sqrt(d
+ e*x)*sqrt(a + b*x)*b**6*d**5*e + 70*sqrt(d + e*x)*sqrt(a + b*x)*b**6*d*
**4*e**2*x - 56*sqrt(d + e*x)*sqrt(a + b*x)*b**6*d**3*e**3*x**2 + 48*sqrt(d
+ e*x)*sqrt(a + b*x)*b**6*d**2*e**4*x**3 + 1664*sqrt(d + e*x)*sqrt(a + b*
x)*b**6*d*e**5*x**4 + 1280*sqrt(d + e*x)*sqrt(a + b*x)*b**6*e**6*x**5 + 10
5*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt
(a*e - b*d))*a**6*e**6 - 630*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) +
sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**5*b*d*e**5 + 1575*sqrt(e)*sqrt(
b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a
**4*b**2*d**2*e**4 - 2100*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + s...
```


3.195 $\int (a + bx)^{5/2} (A + Bx) \sqrt{d + ex} dx$

Optimal result	1780
Mathematica [A] (verified)	1781
Rubi [A] (verified)	1781
Maple [B] (verified)	1785
Fricas [A] (verification not implemented)	1786
Sympy [F]	1787
Maxima [F(-2)]	1788
Giac [B] (verification not implemented)	1788
Mupad [F(-1)]	1789
Reduce [B] (verification not implemented)	1790

Optimal result

Integrand size = 24, antiderivative size = 304

$$\int (a + bx)^{5/2} (A + Bx) \sqrt{d + ex} dx =$$

$$-\frac{(bd - ae)^3 (7bBd - 10Abe + 3aBe) \sqrt{a + bx} \sqrt{d + ex}}{128b^2e^4}$$

$$+ \frac{(bd - ae)^2 (7bBd - 10Abe + 3aBe) (a + bx)^{3/2} \sqrt{d + ex}}{192b^2e^3}$$

$$- \frac{(bd - ae) (7bBd - 10Abe + 3aBe) (a + bx)^{5/2} \sqrt{d + ex}}{240b^2e^2}$$

$$- \frac{(7bBd - 10Abe + 3aBe) (a + bx)^{7/2} \sqrt{d + ex}}{40b^2e} + \frac{B(a + bx)^{7/2} (d + ex)^{3/2}}{5be}$$

$$+ \frac{(bd - ae)^4 (7bBd - 10Abe + 3aBe) \operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{128b^{5/2}e^{9/2}}$$

output

```
-1/128*(-a*e+b*d)^3*(-10*A*b*e+3*B*a*e+7*B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(1/2)
)/b^2/e^4+1/192*(-a*e+b*d)^2*(-10*A*b*e+3*B*a*e+7*B*b*d)*(b*x+a)^(3/2)*(e*
x+d)^(1/2)/b^2/e^3-1/240*(-a*e+b*d)*(-10*A*b*e+3*B*a*e+7*B*b*d)*(b*x+a)^(5
/2)*(e*x+d)^(1/2)/b^2/e^2-1/40*(-10*A*b*e+3*B*a*e+7*B*b*d)*(b*x+a)^(7/2)*(
e*x+d)^(1/2)/b^2/e+1/5*B*(b*x+a)^(7/2)*(e*x+d)^(3/2)/b/e+1/128*(-a*e+b*d)^
4*(-10*A*b*e+3*B*a*e+7*B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d
)^(1/2))/b^(5/2)/e^(9/2)
```

Mathematica [A] (verified)

Time = 0.74 (sec) , antiderivative size = 321, normalized size of antiderivative = 1.06

$$\int (a + bx)^{5/2} (A + Bx) \sqrt{d + ex} dx = \frac{\sqrt{a + bx} \sqrt{d + ex} (-45a^4 B e^4 + 30a^3 b e^3 (2Bd + 5Ae + Bex) + 2a^2 b^2 e^2 (5Ae(73d + 118e) + B(-173d^2 + 109d e + 372e^2 x^2)) + 2a b^3 e (5Ae(-55d^2 + 36d e + 136e^2 x^2) + B(170d^3 - 111d^2 e + 88d e^2 x^2 + 504e^3 x^3)) + b^4 (10Ae(15d^3 - 10d^2 e + 8d e^2 x^2 + 48e^3 x^3) + B(-105d^4 + 70d^3 e + 56d^2 e^2 x^2 + 48d e^3 x^3 + 384e^4 x^4)))}{128b^5/2 e^{9/2}} + \frac{(bd - ae)^4 (7bBd - 10Abe + 3aBe) \operatorname{arctanh}\left(\frac{\sqrt{b} \sqrt{d+ex}}{\sqrt{e} \sqrt{a+bx}}\right)}{128b^5/2 e^{9/2}}$$

input `Integrate[(a + b*x)^(5/2)*(A + B*x)*Sqrt[d + e*x], x]`

output
$$\frac{(\operatorname{Sqrt}[a + b*x] \operatorname{Sqrt}[d + e*x] * (-45*a^4*B*e^4 + 30*a^3*b*e^3*(2*B*d + 5*A*e + B*e*x) + 2*a^2*b^2*e^2*(5*A*e*(73*d + 118*e*x) + B*(-173*d^2 + 109*d*e*x + 372*e^2*x^2)) + 2*a*b^3*e*(5*A*e*(-55*d^2 + 36*d*e*x + 136*e^2*x^2) + B*(170*d^3 - 111*d^2*e*x + 88*d*e^2*x^2 + 504*e^3*x^3)) + b^4*(10*A*e*(15*d^3 - 10*d^2*e*x + 8*d*e^2*x^2 + 48*e^3*x^3) + B*(-105*d^4 + 70*d^3*e*x - 56*d^2*e^2*x^2 + 48*d*e^3*x^3 + 384*e^4*x^4))))}{(1920*b^2*e^4) + ((b*d - a*e)^4*(7*b*B*d - 10*A*b*e + 3*a*B*e)*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*\operatorname{Sqrt}[d + e*x])]/(\operatorname{Sqrt}[e]*\operatorname{Sqrt}[a + b*x]))}{(128*b^(5/2)*e^(9/2))}$$

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 252, normalized size of antiderivative = 0.83, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.292$, Rules used = {90, 60, 60, 60, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^{5/2} (A + Bx) \sqrt{d + ex} dx$$

↓ 90

$$\frac{B(a + bx)^{7/2} (d + ex)^{3/2}}{5be} - \frac{(3aBe - 10Abe + 7bBd) \int (a + bx)^{5/2} \sqrt{d + ex} dx}{10be}$$

$$\begin{aligned}
 & \downarrow 60 \\
 & \frac{B(a+bx)^{7/2}(d+ex)^{3/2}}{5be} - \frac{(3aBe - 10Abe + 7bBd) \left(\frac{(bd-ae) \int \frac{(a+bx)^{5/2}}{\sqrt{d+ex}} dx}{8b} + \frac{(a+bx)^{7/2} \sqrt{d+ex}}{4b} \right)}{10be} \\
 & \downarrow 60 \\
 & \frac{B(a+bx)^{7/2}(d+ex)^{3/2}}{5be} - \frac{(3aBe - 10Abe + 7bBd) \left(\frac{(bd-ae) \left(\frac{(a+bx)^{5/2} \sqrt{d+ex}}{3e} - \frac{5(bd-ae) \int \frac{(a+bx)^{3/2}}{\sqrt{d+ex}} dx}{6e} \right)}{8b} + \frac{(a+bx)^{7/2} \sqrt{d+ex}}{4b} \right)}{10be} \\
 & \downarrow 60 \\
 & \frac{B(a+bx)^{7/2}(d+ex)^{3/2}}{5be} - \frac{(3aBe - 10Abe + 7bBd) \left(\frac{(bd-ae) \left(\frac{(a+bx)^{5/2} \sqrt{d+ex}}{3e} - \frac{5(bd-ae) \left(\frac{(a+bx)^{3/2} \sqrt{d+ex}}{2e} - \frac{3(bd-ae) \int \frac{\sqrt{a+bx}}{\sqrt{d+ex}} dx}{4e} \right)}{6e} \right)}{8b} + \frac{(a+bx)^{7/2} \sqrt{d+ex}}{4b} \right)}{10be} \\
 & \downarrow 60
 \end{aligned}$$

$$\begin{array}{l}
 \frac{B(a+bx)^{7/2}(d+ex)^{3/2}}{5be} - \\
 (bd-ae) \left(\frac{(a+bx)^{5/2}\sqrt{d+ex}}{3e} - \frac{5(bd-ae) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae)}{4e} \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{2e} \right)}{6e} \right)}{8b} \right) \\
 (3aBe - 10Abe + 7bBd)
 \end{array}$$

10be

↓ 66

$$\begin{array}{l}
 \frac{B(a+bx)^{7/2}(d+ex)^{3/2}}{5be} - \\
 (bd-ae) \left(\frac{(a+bx)^{5/2}\sqrt{d+ex}}{3e} - \frac{5(bd-ae) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}} dx}{e} \right)}{4e} \right)}{6e} \right)}{8b} \\
 (3aBe - 10Abe + 7bBd)
 \end{array}$$

10be

↓ 221

$$\frac{(3aBe - 10Abe + 7bBd) \left(\frac{B(a+bx)^{7/2}(d+ex)^{3/2}}{5be} - \frac{(bd-ae) \left(\frac{(a+bx)^{5/2}\sqrt{d+ex}}{3e} - \frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{\sqrt{be}^{3/2}} \right)}{4e} \right)}{6e} \right)}{8b} \right)}{10be}$$

```
input Int[(a + b*x)^(5/2)*(A + B*x)*Sqrt[d + e*x], x]
```

```
output (B*(a + b*x)^(7/2)*(d + e*x)^(3/2))/(5*b*e) - ((7*b*B*d - 10*A*b*e + 3*a*B*e)*(((a + b*x)^(7/2)*Sqrt[d + e*x])/(4*b) + ((b*d - a*e)*(((a + b*x)^(5/2)*Sqrt[d + e*x])/(3*e) - (5*(b*d - a*e)*(((a + b*x)^(3/2)*Sqrt[d + e*x])/(2*e) - (3*(b*d - a*e)*((Sqrt[a + b*x]*Sqrt[d + e*x])/e - ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(Sqrt[b]*e^(3/2)))))/(4*e)))/(6*e)))/(8*b)))/(10*b*e)
```

Defintions of rubi rules used

```
rule 60 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

rule 66 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[
2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; Fre
eQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 90 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p
), x] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))),
x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p
+ 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n,
p}, x] && NeQ[n + p + 2, 0]`

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1371 vs. $2(260) = 520$.

Time = 0.26 (sec) , antiderivative size = 1372, normalized size of antiderivative = 4.51

method	result	size
default	Expression too large to display	1372

input `int((b*x+a)^(5/2)*(B*x+A)*(e*x+d)^(1/2),x,method=_RETURNVERBOSE)`

output

```

-1/3840*(b*x+a)^(1/2)*(e*x+d)^(1/2)*(-768*B*b^4*e^4*x^4*((e*x+d)*(b*x+a))^(
(1/2)*(b*e)^(1/2)-960*A*b^4*e^4*x^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-20
16*B*a*b^3*e^4*x^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-96*B*b^4*d*e^3*x^3*
((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-2720*A*a*b^3*e^4*x^2*((e*x+d)*(b*x+a))
^(1/2)*(b*e)^(1/2)+1100*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b^3*d^2*e^
2-120*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^3*b*d*e^3+692*B*((e*x+d)*(b*
x+a))^(1/2)*(b*e)^(1/2)*a^2*b^2*d^2*e^2-680*B*((e*x+d)*(b*x+a))^(1/2)*(b*e
)^(1/2)*a*b^3*d^3*e+90*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^4*e^4+210*B
*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^4*d^4-352*B*a*b^3*d*e^3*x^2*((e*x+d
)*(b*x+a))^(1/2)*(b*e)^(1/2)-720*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b
^3*d*e^3*x-436*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^2*b^2*d*e^3*x+444*B
*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b^3*d^2*e^2*x+150*A*ln(1/2*(2*b*e*x
+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^4*b*e^5+150
*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1
/2))*b^5*d^4*e+900*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)
+a*e+d*b)/(b*e)^(1/2))*a^2*b^3*d^2*e^3-600*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b
*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^4*d^3*e^2+75*B*ln(1/2*(
2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^4*b*
d*e^4+150*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)
/(b*e)^(1/2))*a^3*b^2*d^2*e^3-450*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))...

```

Fricas [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 1046, normalized size of antiderivative = 3.44

$$\int (a + bx)^{5/2} (A + Bx) \sqrt{d + ex} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^(5/2)*(B*x+A)*(e*x+d)^(1/2),x, algorithm="fricas")
```

output

```

[-1/7680*(15*(7*B*b^5*d^5 - 5*(5*B*a*b^4 + 2*A*b^5)*d^4*e + 10*(3*B*a^2*b^3 + 4*A*a*b^4)*d^3*e^2 - 10*(B*a^3*b^2 + 6*A*a^2*b^3)*d^2*e^3 - 5*(B*a^4*b - 8*A*a^3*b^2)*d*e^4 + (3*B*a^5 - 10*A*a^4*b)*e^5)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 - 4*(2*b*e*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x + a)*sqrt(e*x + d) + 8*(b^2*d*e + a*b*e^2)*x) - 4*(384*B*b^5*e^5*x^4 - 105*B*b^5*d^4*e + 10*(34*B*a*b^4 + 15*A*b^5)*d^3*e^2 - 2*(173*B*a^2*b^3 + 275*A*a*b^4)*d^2*e^3 + 10*(6*B*a^3*b^2 + 73*A*a^2*b^3)*d*e^4 - 15*(3*B*a^4*b - 10*A*a^3*b^2)*e^5 + 48*(B*b^5*d*e^4 + (21*B*a*b^4 + 10*A*b^5)*e^5)*x^3 - 8*(7*B*b^5*d^2*e^3 - 2*(11*B*a*b^4 + 5*A*b^5)*d*e^4 - (93*B*a^2*b^3 + 170*A*a*b^4)*e^5)*x^2 + 2*(35*B*b^5*d^3*e^2 - (111*B*a*b^4 + 50*A*b^5)*d^2*e^3 + (109*B*a^2*b^3 + 180*A*a*b^4)*d*e^4 + 5*(3*B*a^3*b^2 + 118*A*a^2*b^3)*e^5)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b^3*e^5), -1/3840*(15*(7*B*b^5*d^5 - 5*(5*B*a*b^4 + 2*A*b^5)*d^4*e + 10*(3*B*a^2*b^3 + 4*A*a*b^4)*d^3*e^2 - 10*(B*a^3*b^2 + 6*A*a^2*b^3)*d^2*e^3 - 5*(B*a^4*b - 8*A*a^3*b^2)*d*e^4 + (3*B*a^5 - 10*A*a^4*b)*e^5)*sqrt(-b*e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*e^2*x^2 + a*b*d*e + (b^2*d*e + a*b*e^2)*x)) - 2*(384*B*b^5*e^5*x^4 - 105*B*b^5*d^4*e + 10*(34*B*a*b^4 + 15*A*b^5)*d^3*e^2 - 2*(173*B*a^2*b^3 + 275*A*a*b^4)*d^2*e^3 + 10*(6*B*a^3*b^2 + 73*A*a^2*b^3)*d*e^4 - 15*(3*B*a^4*b - 10*A*a^3*b^2)*e^5 + 48*(B*b^5*d*e^4 + (21*B*a*b^4 + 10*A*b^5)*e^5)*x^3 - 8*(7*B*b^5*d^2*e^3 - 2*(...

```

Sympy [F]

$$\int (a + bx)^{5/2} (A + Bx) \sqrt{d + ex} dx = \int (A + Bx) (a + bx)^{5/2} \sqrt{d + ex} dx$$

input

```
integrate((b*x+a)**(5/2)*(B*x+A)*(e*x+d)**(1/2),x)
```

output

```
Integral((A + B*x)*(a + b*x)**(5/2)*sqrt(d + e*x), x)
```


Maxima [F(-2)]

Exception generated.

$$\int (a + bx)^{5/2} (A + Bx) \sqrt{d + ex} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)*(e*x+d)^(1/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1672 vs. 2(260) = 520.

Time = 0.34 (sec) , antiderivative size = 1672, normalized size of antiderivative = 5.50

$$\int (a + bx)^{5/2} (A + Bx) \sqrt{d + ex} dx = \text{Too large to display}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)*(e*x+d)^(1/2),x, algorithm="giac")`

output

```

1/1920*(30*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*(b*x + a)*(4*(b*x + a)*
(6*(b*x + a)/b^3 + (b^12*d*e^5 - 25*a*b^11*e^6)/(b^14*e^6)) - (5*b^13*d^2*
e^4 + 14*a*b^12*d*e^5 - 163*a^2*b^11*e^6)/(b^14*e^6)) + 3*(5*b^14*d^3*e^3
+ 9*a*b^13*d^2*e^4 + 15*a^2*b^12*d*e^5 - 93*a^3*b^11*e^6)/(b^14*e^6))*sqrt
(b*x + a) + 3*(5*b^4*d^4 + 4*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2 + 20*a^3*b*d*
e^3 - 35*a^4*e^4)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)
)*b*e - a*b*e)))/(sqrt(b*e)*b^2*e^3))*B*a*abs(b) - 1920*((b^2*d - a*b*e)*l
og(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/sq
rt(b*e) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*sqrt(b*x + a))*A*a^3*abs(b)/
b^2 + 10*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*(b*x + a)*(4*(b*x + a)*(6
*(b*x + a)/b^3 + (b^12*d*e^5 - 25*a*b^11*e^6)/(b^14*e^6)) - (5*b^13*d^2*e^
4 + 14*a*b^12*d*e^5 - 163*a^2*b^11*e^6)/(b^14*e^6)) + 3*(5*b^14*d^3*e^3 +
9*a*b^13*d^2*e^4 + 15*a^2*b^12*d*e^5 - 93*a^3*b^11*e^6)/(b^14*e^6))*sqrt(b
*x + a) + 3*(5*b^4*d^4 + 4*a*b^3*d^3*e + 6*a^2*b^2*d^2*e^2 + 20*a^3*b*d*e^
3 - 35*a^4*e^4)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*
b*e - a*b*e)))/(sqrt(b*e)*b^2*e^3))*A*b*abs(b) + 480*(sqrt(b^2*d + (b*x +
a)*b*e - a*b*e)*(2*b*x + 2*a + (b*d*e - 5*a*e^2)/e^2)*sqrt(b*x + a) + (b^3
*d^2 + 2*a*b^2*d*e - 3*a^2*b*e^2)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(
b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*e))*B*a^3*abs(b)/b^3 + 1440*(s
qrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*b*x + 2*a + (b*d*e - 5*a*e^2)/e^2...

```

Mupad [F(-1)]

Timed out.

$$\int (a + bx)^{5/2} (A + Bx) \sqrt{d + ex} dx = \int (A + Bx) (a + bx)^{5/2} \sqrt{d + ex} dx$$

input

```
int((A + B*x)*(a + b*x)^(5/2)*(d + e*x)^(1/2),x)
```

output

```
int((A + B*x)*(a + b*x)^(5/2)*(d + e*x)^(1/2), x)
```


3.196 $\int \frac{(a+bx)^{5/2}(A+Bx)}{\sqrt{d+ex}} dx$

Optimal result	1791
Mathematica [A] (verified)	1792
Rubi [A] (verified)	1792
Maple [B] (verified)	1795
Fricas [A] (verification not implemented)	1796
Sympy [F]	1797
Maxima [F(-2)]	1798
Giac [A] (verification not implemented)	1798
Mupad [F(-1)]	1799
Reduce [B] (verification not implemented)	1799

Optimal result

Integrand size = 24, antiderivative size = 246

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{\sqrt{d+ex}} dx =$$

$$\frac{5(bd-ae)^2(7bBd-8Abe+aBe)\sqrt{a+bx}\sqrt{d+ex}}{64be^4}$$

$$+ \frac{5(bd-ae)(7bBd-8Abe+aBe)(a+bx)^{3/2}\sqrt{d+ex}}{96be^3}$$

$$- \frac{(7bBd-8Abe+aBe)(a+bx)^{5/2}\sqrt{d+ex}}{24be^2} + \frac{B(a+bx)^{7/2}\sqrt{d+ex}}{4be}$$

$$+ \frac{5(bd-ae)^3(7bBd-8Abe+aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{64b^{3/2}e^{9/2}}$$

output

```
-5/64*(-a*e+b*d)^2*(-8*A*b*e+B*a*e+7*B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(1/2)/b/
e^4+5/96*(-a*e+b*d)*(-8*A*b*e+B*a*e+7*B*b*d)*(b*x+a)^(3/2)*(e*x+d)^(1/2)/
/e^3-1/24*(-8*A*b*e+B*a*e+7*B*b*d)*(b*x+a)^(5/2)*(e*x+d)^(1/2)/b/e^2+1/4*B
*(b*x+a)^(7/2)*(e*x+d)^(1/2)/b/e+5/64*(-a*e+b*d)^3*(-8*A*b*e+B*a*e+7*B*b*d
)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)^(1/2))/b^(3/2)/e^(9/2)
```

Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 230, normalized size of antiderivative = 0.93

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{\sqrt{d + ex}} dx = \frac{\sqrt{a + bx}\sqrt{d + ex}(15a^3Be^3 + a^2be^2(-191Bd + 264Ae + 118Bex) + ab^2e(16.5(bd - ae)^3(7bBd - 8Abe + aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{e}\sqrt{a+bx}}\right) + \frac{5(bd - ae)^3(7bBd - 8Abe + aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{e}\sqrt{a+bx}}\right)}{64b^{3/2}e^{9/2}}$$

input

```
Integrate[((a + b*x)^(5/2)*(A + B*x))/Sqrt[d + e*x],x]
```

output

```
(Sqrt[a + b*x]*Sqrt[d + e*x]*(15*a^3*B*e^3 + a^2*b*e^2*(-191*B*d + 264*A*e + 118*B*e*x) + a*b^2*e*(16*A*e*(-20*d + 13*e*x) + B*(265*d^2 - 172*d*e*x + 136*e^2*x^2)) + b^3*(8*A*e*(15*d^2 - 10*d*e*x + 8*e^2*x^2) + B*(-105*d^3 + 70*d^2*e*x - 56*d*e^2*x^2 + 48*e^3*x^3)))/(192*b*e^4) + (5*(b*d - a*e)^3*(7*b*B*d - 8*A*b*e + a*B*e)*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/(Sqrt[e]*Sqrt[a + b*x])])/(64*b^(3/2)*e^(9/2))
```

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 210, normalized size of antiderivative = 0.85, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {90, 60, 60, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{\sqrt{d + ex}} dx$$

↓ 90

$$\frac{B(a + bx)^{7/2}\sqrt{d + ex}}{4be} - \frac{(aBe - 8Abe + 7bBd) \int \frac{(a+bx)^{5/2}}{\sqrt{d+ex}} dx}{8be}$$

↓ 60

$$\begin{aligned}
 & \frac{B(a+bx)^{7/2}\sqrt{d+ex}}{4be} - \frac{(aBe - 8Abe + 7bBd) \left(\frac{(a+bx)^{5/2}\sqrt{d+ex}}{3e} - \frac{5(bd-ae) \int \frac{(a+bx)^{3/2}}{\sqrt{d+ex}} dx}{6e} \right)}{8be} \\
 & \quad \downarrow 60 \\
 & \frac{B(a+bx)^{7/2}\sqrt{d+ex}}{4be} - \frac{(aBe - 8Abe + 7bBd) \left(\frac{(a+bx)^{5/2}\sqrt{d+ex}}{3e} - \frac{5(bd-ae) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \int \frac{\sqrt{a+bx}}{\sqrt{d+ex}} dx}{4e} \right)}{6e} \right)}{8be} \\
 & \quad \downarrow 60 \\
 & \frac{B(a+bx)^{7/2}\sqrt{d+ex}}{4be} - \frac{(aBe - 8Abe + 7bBd) \left(\frac{(a+bx)^{5/2}\sqrt{d+ex}}{3e} - \frac{5(bd-ae) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{2e} \right)}{4e} \right)}{6e} \right)}{8be} \\
 & \quad \downarrow 66 \\
 & \frac{B(a+bx)^{7/2}\sqrt{d+ex}}{4be} - \frac{(aBe - 8Abe + 7bBd) \left(\frac{(a+bx)^{5/2}\sqrt{d+ex}}{3e} - \frac{5(bd-ae) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}} dx}{e} \right)}{4e} \right)}{6e} \right)}{8be} \\
 & \quad \downarrow 221
 \end{aligned}$$

$$\frac{B(a+bx)^{7/2}\sqrt{d+ex}}{4be} - \frac{(aBe - 8Abe + 7bBd) \left(\frac{(a+bx)^{5/2}\sqrt{d+ex}}{3e} - \frac{5(bd-ae) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{\sqrt{be}^{3/2}} \right)}{4e} \right)}{6e} \right)}{8be}$$

```
input Int[((a + b*x)^(5/2)*(A + B*x))/Sqrt[d + e*x], x]
```

```
output (B*(a + b*x)^(7/2)*Sqrt[d + e*x])/(4*b*e) - ((7*b*B*d - 8*A*b*e + a*B*e)*((a + b*x)^(5/2)*Sqrt[d + e*x])/(3*e) - (5*(b*d - a*e)*((a + b*x)^(3/2)*Sqrt[d + e*x])/(2*e) - (3*(b*d - a*e)*((Sqrt[a + b*x]*Sqrt[d + e*x])/e - ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(Sqrt[b]*e^(3/2))))/(4*e))/(6*e))/(8*b*e)
```

Defintions of rubi rules used

```
rule 60 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

```
rule 66 Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]
```

rule 90

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p
_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))),
x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p
+ 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n,
p}, x] && NeQ[n + p + 2, 0]
```

rule 221

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 967 vs. $2(208) = 416$.

Time = 0.26 (sec) , antiderivative size = 968, normalized size of antiderivative = 3.93

method	result
default	$\frac{\sqrt{bx+a}\sqrt{ex+d}\left(96Bb^3e^3x^3\sqrt{(ex+d)(bx+a)}\sqrt{be}+128Ab^3e^3x^2\sqrt{(ex+d)(bx+a)}\sqrt{be}-112Bb^3de^2x^2\sqrt{(ex+d)(bx+a)}\sqrt{be}-15B\right)}{\dots}$

input

```
int((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(1/2),x,method=_RETURNVERBOSE)
```


output

```

1/384*(b*x+a)^(1/2)*(e*x+d)^(1/2)*(96*B*b^3*e^3*x^3*((e*x+d)*(b*x+a))^(1/2)
)*(b*e)^(1/2)+128*A*b^3*e^3*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-112*B*
b^3*d*e^2*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-15*B*ln(1/2*(2*b*e*x+2*(
(e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^4*e^4+105*B*ln(
1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b
^4*d^4-344*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b^2*d*e^2*x-360*A*ln(1/
2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2
*b^2*d*e^3+360*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e
+d*b)/(b*e)^(1/2))*a*b^3*d^2*e^2+416*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)
*a*b^2*e^3*x-160*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^3*d*e^2*x+236*B*(
(e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^2*b*e^3*x+140*B*((e*x+d)*(b*x+a))^(1/
2)*(b*e)^(1/2)*b^3*d^2*e*x+30*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^3*e^
3-210*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^3*d^3+120*A*ln(1/2*(2*b*e*x+
2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^3*b*e^4-120*
A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/
2))*b^4*d^3*e-640*A*a*b^2*d*e^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+272*B*
a*b^2*e^3*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-300*B*ln(1/2*(2*b*e*x+2*
((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^3*d^3*e+528*
A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^2*b*e^3+240*A*((e*x+d)*(b*x+a))^(1
/2)*(b*e)^(1/2)*b^3*d^2*e-60*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)...

```

Fricas [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 770, normalized size of antiderivative = 3.13

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{\sqrt{d + ex}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(1/2),x, algorithm="fricas")
```

output

```
[1/768*(15*(7*B*b^4*d^4 - 4*(5*B*a*b^3 + 2*A*b^4)*d^3*e + 6*(3*B*a^2*b^2 +
4*A*a*b^3)*d^2*e^2 - 4*(B*a^3*b + 6*A*a^2*b^2)*d*e^3 - (B*a^4 - 8*A*a^3*b
)*e^4)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*
b*e*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x + a)*sqrt(e*x + d) + 8*(b^2*d*e + a*
b*e^2)*x) + 4*(48*B*b^4*e^4*x^3 - 105*B*b^4*d^3*e + 5*(53*B*a*b^3 + 24*A*b
^4)*d^2*e^2 - (191*B*a^2*b^2 + 320*A*a*b^3)*d*e^3 + 3*(5*B*a^3*b + 88*A*a^
2*b^2)*e^4 - 8*(7*B*b^4*d*e^3 - (17*B*a*b^3 + 8*A*b^4)*e^4)*x^2 + 2*(35*B*
b^4*d^2*e^2 - 2*(43*B*a*b^3 + 20*A*b^4)*d*e^3 + (59*B*a^2*b^2 + 104*A*a*b^
3)*e^4)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b^2*e^5), -1/384*(15*(7*B*b^4*d^4
- 4*(5*B*a*b^3 + 2*A*b^4)*d^3*e + 6*(3*B*a^2*b^2 + 4*A*a*b^3)*d^2*e^2 - 4
*(B*a^3*b + 6*A*a^2*b^2)*d*e^3 - (B*a^4 - 8*A*a^3*b)*e^4)*sqrt(-b*e)*arcta
n(1/2*(2*b*e*x + b*d + a*e)*sqrt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d))/(b^2*e^
2*x^2 + a*b*d*e + (b^2*d*e + a*b*e^2)*x) - 2*(48*B*b^4*e^4*x^3 - 105*B*b^
4*d^3*e + 5*(53*B*a*b^3 + 24*A*b^4)*d^2*e^2 - (191*B*a^2*b^2 + 320*A*a*b^3
)*d*e^3 + 3*(5*B*a^3*b + 88*A*a^2*b^2)*e^4 - 8*(7*B*b^4*d*e^3 - (17*B*a*b^
3 + 8*A*b^4)*e^4)*x^2 + 2*(35*B*b^4*d^2*e^2 - 2*(43*B*a*b^3 + 20*A*b^4)*d*
e^3 + (59*B*a^2*b^2 + 104*A*a*b^3)*e^4)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b
^2*e^5)]
```

SymPy [F]

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{\sqrt{d + ex}} dx = \int \frac{(A + Bx)(a + bx)^{5/2}}{\sqrt{d + ex}} dx$$

input

```
integrate((b*x+a)**(5/2)*(B*x+A)/(e*x+d)**(1/2),x)
```

output

```
Integral((A + B*x)*(a + b*x)**(5/2)/sqrt(d + e*x), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{\sqrt{d + ex}} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(1/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail

Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 415, normalized size of antiderivative = 1.69

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{\sqrt{d + ex}} dx = \frac{\left(\sqrt{b^2d + (bx + a)be - a*be} \left(2(bx + a) \left(4(bx + a) \left(\frac{6(bx+a)B}{b^2e} - \frac{7Bb^3de^5 + Bab^2e^6}{b^4e^7} \right) \right) \right) \right)}{\dots}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(1/2),x, algorithm="giac")`

output `1/192*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*(b*x + a)*(4*(b*x + a)*(6*(b*x + a)*B/(b^2*e) - (7*B*b^3*d*e^5 + B*a*b^2*e^6 - 8*A*b^3*e^6)/(b^4*e^7)) + 5*(7*B*b^4*d^2*e^4 - 6*B*a*b^3*d*e^5 - 8*A*b^4*d*e^5 - B*a^2*b^2*e^6 + 8*A*a*b^3*e^6)/(b^4*e^7)) - 15*(7*B*b^5*d^3*e^3 - 13*B*a*b^4*d^2*e^4 - 8*A*b^5*d^2*e^4 + 5*B*a^2*b^3*d*e^5 + 16*A*a*b^4*d*e^5 + B*a^3*b^2*e^6 - 8*A*a^2*b^3*e^6)/(b^4*e^7))*sqrt(b*x + a) - 15*(7*B*b^4*d^4 - 20*B*a*b^3*d^3*e - 8*A*b^4*d^3*e + 18*B*a^2*b^2*d^2*e^2 + 24*A*a*b^3*d^2*e^2 - 4*B*a^3*b*d*e^3 - 24*A*a^2*b^2*d*e^3 - B*a^4*e^4 + 8*A*a^3*b*e^4)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*b*e^4)*b/abs(b)`

3.197 $\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{3/2}} dx$

Optimal result	1800
Mathematica [A] (verified)	1801
Rubi [A] (verified)	1801
Maple [B] (verified)	1804
Fricas [B] (verification not implemented)	1805
Sympy [F]	1806
Maxima [F(-2)]	1807
Giac [B] (verification not implemented)	1807
Mupad [F(-1)]	1808
Reduce [B] (verification not implemented)	1808

Optimal result

Integrand size = 24, antiderivative size = 218

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{3/2}} dx = \frac{2(Bd - Ae)(a+bx)^{5/2}}{e^2\sqrt{d+ex}} + \frac{5(bd - ae)(7bBd - 6Abe - aBe)\sqrt{a+bx}\sqrt{d+ex}}{8e^4} - \frac{5(7bBd - 6Abe - aBe)(a+bx)^{3/2}\sqrt{d+ex}}{12e^3} + \frac{B(a+bx)^{5/2}\sqrt{d+ex}}{3e^2} - \frac{5(bd - ae)^2(7bBd - 6Abe - aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{8\sqrt{b}e^{9/2}}$$

output

```
2*(-A*e+B*d)*(b*x+a)^(5/2)/e^2/(e*x+d)^(1/2)+5/8*(-a*e+b*d)*(-6*A*b*e-B*a*
e+7*B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(1/2)/e^4-5/12*(-6*A*b*e-B*a*e+7*B*b*d)*
(b*x+a)^(3/2)*(e*x+d)^(1/2)/e^3+1/3*B*(b*x+a)^(5/2)*(e*x+d)^(1/2)/e^2-5/8*
(-a*e+b*d)^2*(-6*A*b*e-B*a*e+7*B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)
/(e*x+d)^(1/2))/b^(1/2)/e^(9/2)
```

Mathematica [A] (verified)

Time = 0.56 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.00

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{3/2}} dx = \frac{\sqrt{a+bx}(3a^2e^2(27Bd-16Ae+11Bex) + 2abe(3Ae(25d+9ex) + B(-95d^2 - 34d*ex + 13e^2*x^2))) + b^2(6Ae*(-15d^2 - 5d*ex + 2e^2*x^2) + B(105d^3 + 35d^2*ex - 14d*e^2*x^2 + 8e^3*x^3))}{24e^4\sqrt{d+ex}} + \frac{5(bd-ae)^2(-7bBd+6Abe+aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{8\sqrt{b}e^{9/2}}$$

input `Integrate[((a + b*x)^(5/2)*(A + B*x))/(d + e*x)^(3/2),x]`

output `(Sqrt[a + b*x]*(3*a^2*e^2*(27*B*d - 16*A*e + 11*B*e*x) + 2*a*b*e*(3*A*e*(25*d + 9*e*x) + B*(-95*d^2 - 34*d*e*x + 13*e^2*x^2))) + b^2*(6*A*e*(-15*d^2 - 5*d*e*x + 2*e^2*x^2) + B*(105*d^3 + 35*d^2*e*x - 14*d*e^2*x^2 + 8*e^3*x^3)))/(24*e^4*Sqrt[d + e*x]) + (5*(b*d - a*e)^2*(-7*b*B*d + 6*A*b*e + a*B*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(8*Sqrt[b]*e^(9/2))`

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.04, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {87, 60, 60, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{3/2}} dx$$

$$\downarrow 87$$

$$\frac{(-aBe - 6Abe + 7bBd) \int \frac{(a+bx)^{5/2}}{\sqrt{d+ex}} dx}{e(bd-ae)} - \frac{2(a+bx)^{7/2}(Bd-Ae)}{e\sqrt{d+ex}(bd-ae)}$$

$$\downarrow 60$$

$$\frac{(-aBe - 6Abe + 7bBd) \left(\frac{(a+bx)^{5/2}\sqrt{d+ex}}{3e} - \frac{5(bd-ae) \int \frac{(a+bx)^{3/2}}{\sqrt{d+ex}} dx}{6e} \right)}{e(bd - ae)} - \frac{2(a + bx)^{7/2}(Bd - Ae)}{e\sqrt{d + ex}(bd - ae)}$$

↓ 60

$$\frac{(-aBe - 6Abe + 7bBd) \left(\frac{(a+bx)^{5/2}\sqrt{d+ex}}{3e} - \frac{5(bd-ae) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \int \frac{\sqrt{a+bx}}{\sqrt{d+ex}} dx}{4e} \right)}{6e} \right)}{e(bd - ae)}$$

$$\frac{2(a + bx)^{7/2}(Bd - Ae)}{e\sqrt{d + ex}(bd - ae)}$$

↓ 60

$$\frac{(-aBe - 6Abe + 7bBd) \left(\frac{(a+bx)^{5/2}\sqrt{d+ex}}{3e} - \frac{5(bd-ae) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{2e} \right)}{4e} \right)}{6e} \right)}{e(bd - ae)}$$

$$\frac{2(a + bx)^{7/2}(Bd - Ae)}{e\sqrt{d + ex}(bd - ae)}$$

↓ 66

$$\frac{(-aBe - 6Abe + 7bBd) \left(\frac{(a+bx)^{5/2}\sqrt{d+ex}}{3e} - \frac{5(bd-ae) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}} dx}{e} \right)}{4e} \right)}{6e} \right)}{e(bd - ae)}$$

$$\frac{2(a + bx)^{7/2}(Bd - Ae)}{e\sqrt{d + ex}(bd - ae)}$$

$$\begin{aligned}
 & \downarrow 221 \\
 & (-aBe - 6Abe + 7bBd) \left(\frac{(a+bx)^{5/2}\sqrt{d+ex}}{3e} - \frac{5(bd-ae) \left(\frac{(a+bx)^{3/2}\sqrt{d+ex}}{2e} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{\sqrt{be}^{3/2}} \right)}{4e} \right)}{6e} \right) \\
 & \frac{e(bd-ae)}{2(a+bx)^{7/2}(Bd-Ae)} \\
 & \frac{e\sqrt{d+ex}(bd-ae)}{e\sqrt{d+ex}(bd-ae)}
 \end{aligned}$$

input

```
Int[((a + b*x)^(5/2)*(A + B*x))/(d + e*x)^(3/2), x]
```

output

```
(-2*(B*d - A*e)*(a + b*x)^(7/2))/(e*(b*d - a*e)*Sqrt[d + e*x]) + ((7*b*B*d - 6*A*b*e - a*B*e)*(((a + b*x)^(5/2)*Sqrt[d + e*x])/(3*e) - (5*(b*d - a*e))*(((a + b*x)^(3/2)*Sqrt[d + e*x])/(2*e) - (3*(b*d - a*e))*((Sqrt[a + b*x]*Sqrt[d + e*x])/e - ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(Sqrt[b]*e^(3/2))))/(4*e)))/(6*e))/(e*(b*d - a*e))
```

Defintions of rubi rules used

rule 60

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

rule 66

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]
```


rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1183 vs. $2(182) = 364$.

Time = 0.27 (sec) , antiderivative size = 1184, normalized size of antiderivative = 5.43

method	result	size
default	Expression too large to display	1184

input `int((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(3/2),x,method=_RETURNVERBOSE)`

output

```

1/48*(b*x+a)^(1/2)*(-96*A*a^2*e^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+210*
B*b^2*d^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-136*B*a*b*d*e^2*x*((e*x+d)*
(b*x+a))^(1/2)*(b*e)^(1/2)+15*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*
(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^3*e^4*x+24*A*b^2*e^3*x^2*((e*x+d)*(b*x+
a))^(1/2)*(b*e)^(1/2)-105*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e
)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^3*d^3*e*x+225*B*ln(1/2*(2*b*e*x+2*((e*x+d)
*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^2*d^3*e+66*B*a^2*e^3
*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+162*B*a^2*d*e^2*((e*x+d)*(b*x+a))^(
1/2)*(b*e)^(1/2)+225*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/
2)+a*e+d*b)/(b*e)^(1/2))*a*b^2*d^2*e^2*x+52*B*a*b*e^3*x^2*((e*x+d)*(b*x+a)
)^(1/2)*(b*e)^(1/2)-28*B*b^2*d*e^2*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)
+108*A*a*b*e^3*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+70*B*b^2*d^2*e*x*((e
x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-60*A*b^2*d*e^2*x*((e*x+d)*(b*x+a))^(1/2)*
(b*e)^(1/2)+300*A*a*b*d*e^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-180*A*ln(1/
2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b
^2*d*e^3*x-135*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e
+d*b)/(b*e)^(1/2))*a^2*b*d*e^3*x-105*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))
^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^3*d^4-380*B*a*b*d^2*e*((e*x+d)*
(b*x+a))^(1/2)*(b*e)^(1/2)+90*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*
(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^3*d^3*e+15*B*ln(1/2*(2*b*e*x+2*((e...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 422 vs. $2(182) = 364$.

Time = 0.75 (sec) , antiderivative size = 858, normalized size of antiderivative = 3.94

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{3/2}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(3/2),x, algorithm="fricas")
```

output

```

[-1/96*(15*(7*B*b^3*d^4 - 3*(5*B*a*b^2 + 2*A*b^3)*d^3*e + 3*(3*B*a^2*b + 4
*A*a*b^2)*d^2*e^2 - (B*a^3 + 6*A*a^2*b)*d*e^3 + (7*B*b^3*d^3*e - 3*(5*B*a*
b^2 + 2*A*b^3)*d^2*e^2 + 3*(3*B*a^2*b + 4*A*a*b^2)*d*e^3 - (B*a^3 + 6*A*a^
2*b)*e^4)*x)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 +
4*(2*b*e*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x + a)*sqrt(e*x + d) + 8*(b^2*d*
e + a*b*e^2)*x) - 4*(8*B*b^3*e^4*x^3 + 105*B*b^3*d^3*e - 48*A*a^2*b*e^4 -
10*(19*B*a*b^2 + 9*A*b^3)*d^2*e^2 + 3*(27*B*a^2*b + 50*A*a*b^2)*d*e^3 - 2*
(7*B*b^3*d*e^3 - (13*B*a*b^2 + 6*A*b^3)*e^4)*x^2 + (35*B*b^3*d^2*e^2 - 2*(
34*B*a*b^2 + 15*A*b^3)*d*e^3 + 3*(11*B*a^2*b + 18*A*a*b^2)*e^4)*x)*sqrt(b*
x + a)*sqrt(e*x + d))/(b*e^6*x + b*d*e^5), 1/48*(15*(7*B*b^3*d^4 - 3*(5*B*
a*b^2 + 2*A*b^3)*d^3*e + 3*(3*B*a^2*b + 4*A*a*b^2)*d^2*e^2 - (B*a^3 + 6*A*
a^2*b)*d*e^3 + (7*B*b^3*d^3*e - 3*(5*B*a*b^2 + 2*A*b^3)*d^2*e^2 + 3*(3*B*a
^2*b + 4*A*a*b^2)*d*e^3 - (B*a^3 + 6*A*a^2*b)*e^4)*x)*sqrt(-b*e)*arctan(1/
2*(2*b*e*x + b*d + a*e)*sqrt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*e^2*x^
2 + a*b*d*e + (b^2*d*e + a*b*e^2)*x)) + 2*(8*B*b^3*e^4*x^3 + 105*B*b^3*d^3
*e - 48*A*a^2*b*e^4 - 10*(19*B*a*b^2 + 9*A*b^3)*d^2*e^2 + 3*(27*B*a^2*b +
50*A*a*b^2)*d*e^3 - 2*(7*B*b^3*d*e^3 - (13*B*a*b^2 + 6*A*b^3)*e^4)*x^2 + (
35*B*b^3*d^2*e^2 - 2*(34*B*a*b^2 + 15*A*b^3)*d*e^3 + 3*(11*B*a^2*b + 18*A*
a*b^2)*e^4)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b*e^6*x + b*d*e^5)]

```

Sympy [F]

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{3/2}} dx = \int \frac{(A + Bx)(a + bx)^{5/2}}{(d + ex)^{3/2}} dx$$

input

```
integrate((b*x+a)**(5/2)*(B*x+A)/(e*x+d)**(3/2),x)
```

output

```
Integral((A + B*x)*(a + b*x)**(5/2)/(d + e*x)**(3/2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{3/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(3/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 419 vs. 2(182) = 364.

Time = 0.20 (sec) , antiderivative size = 419, normalized size of antiderivative = 1.92

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{3/2}} dx = \frac{\left((bx + a) \left(2 (bx + a) \left(\frac{4(bx+a)B|b|}{be} - \frac{7Bb^2de^5|b| - Babe^6|b| - 6Ab^2e^6|b|}{b^2e^7} \right) + \frac{5(7Bb^3d^2e^4|b|}{5(7Bb^3d^3|b| - 15Bab^2d^2e|b| - 6Ab^3d^2e|b| + 9Ba^2bde^2|b| + 12Aab^2de^2|b| - Ba^3e^3|b| - 6Aa^2be^3|b|) \log + \frac{5(7Bb^3d^2e^4|b|}{8\sqrt{be}e^4} \right)}{8\sqrt{be}e^4}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(3/2),x, algorithm="giac")`

output

```
1/24*((b*x + a)*(2*(b*x + a)*(4*(b*x + a)*B*abs(b)/(b*e) - (7*B*b^2*d*e^5*
abs(b) - B*a*b*e^6*abs(b) - 6*A*b^2*e^6*abs(b)))/(b^2*e^7)) + 5*(7*B*b^3*d^
2*e^4*abs(b) - 8*B*a*b^2*d*e^5*abs(b) - 6*A*b^3*d*e^5*abs(b) + B*a^2*b*e^6
*abs(b) + 6*A*a*b^2*e^6*abs(b)))/(b^2*e^7) + 15*(7*B*b^4*d^3*e^3*abs(b) -
15*B*a*b^3*d^2*e^4*abs(b) - 6*A*b^4*d^2*e^4*abs(b) + 9*B*a^2*b^2*d*e^5*abs
(b) + 12*A*a*b^3*d*e^5*abs(b) - B*a^3*b*e^6*abs(b) - 6*A*a^2*b^2*e^6*abs(b)
))/ (b^2*e^7)*sqrt(b*x + a)/sqrt(b^2*d + (b*x + a)*b*e - a*b*e) + 5/8*(7*B
*b^3*d^3*abs(b) - 15*B*a*b^2*d^2*e*abs(b) - 6*A*b^3*d^2*e*abs(b) + 9*B*a^2
*b*d*e^2*abs(b) + 12*A*a*b^2*d*e^2*abs(b) - B*a^3*e^3*abs(b) - 6*A*a^2*b*e
^3*abs(b))*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e -
a*b*e)))/(sqrt(b*e)*b*e^4)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{3/2}} dx = \int \frac{(A + Bx)(a + bx)^{5/2}}{(d + ex)^{3/2}} dx$$

input

```
int(((A + B*x)*(a + b*x)^(5/2))/(d + e*x)^(3/2), x)
```

output

```
int(((A + B*x)*(a + b*x)^(5/2))/(d + e*x)^(3/2), x)
```

Reduce [B] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 724, normalized size of antiderivative = 3.32

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{3/2}} dx = \text{Too large to display}$$

input

```
int((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(3/2), x)
```

output

```
( - 384*sqrt(d + e*x)*sqrt(a + b*x)*a**3*e**4 + 1848*sqrt(d + e*x)*sqrt(a
+ b*x)*a**2*b*d*e**3 + 696*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b*e**4*x - 224
0*sqrt(d + e*x)*sqrt(a + b*x)*a*b**2*d**2*e**2 - 784*sqrt(d + e*x)*sqrt(a
+ b*x)*a*b**2*d*e**3*x + 304*sqrt(d + e*x)*sqrt(a + b*x)*a*b**2*e**4*x**2
+ 840*sqrt(d + e*x)*sqrt(a + b*x)*b**3*d**3*e + 280*sqrt(d + e*x)*sqrt(a +
b*x)*b**3*d**2*e**2*x - 112*sqrt(d + e*x)*sqrt(a + b*x)*b**3*d*e**3*x**2
+ 64*sqrt(d + e*x)*sqrt(a + b*x)*b**3*e**4*x**3 + 840*sqrt(e)*sqrt(b)*log(
(sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**3*d*e*
*3 + 840*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x
))/sqrt(a*e - b*d))*a**3*e**4*x - 2520*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a
+ b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**2*b*d**2*e**2 - 2520*
sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a
*e - b*d))*a**2*b*d*e**3*x + 2520*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*
x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*b**2*d**3*e + 2520*sqrt(e)*
sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d
))*a*b**2*d**2*e**2*x - 840*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + s
qrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**3*d**4 - 840*sqrt(e)*sqrt(b)*log
((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**3*d**
3*e*x - 525*sqrt(e)*sqrt(b)*a**3*d*e**3 - 525*sqrt(e)*sqrt(b)*a**3*e**4*x
+ 1575*sqrt(e)*sqrt(b)*a**2*b*d**2*e**2 + 1575*sqrt(e)*sqrt(b)*a**2*b*d...
```

3.198 $\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{5/2}} dx$

Optimal result	1810
Mathematica [C] (verified)	1811
Rubi [A] (verified)	1811
Maple [B] (verified)	1814
Fricas [B] (verification not implemented)	1815
Sympy [F]	1816
Maxima [F(-2)]	1817
Giac [B] (verification not implemented)	1817
Mupad [F(-1)]	1818
Reduce [B] (verification not implemented)	1818

Optimal result

Integrand size = 24, antiderivative size = 212

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{5/2}} dx = \frac{2(Bd - Ae)(a+bx)^{5/2}}{3e^2(d+ex)^{3/2}} + \frac{2(8bBd - 5Abe - 3aBe)(a+bx)^{3/2}}{3e^3\sqrt{d+ex}} - \frac{5b(7bBd - 4Abe - 3aBe)\sqrt{a+bx}\sqrt{d+ex}}{4e^4} + \frac{bB(a+bx)^{3/2}\sqrt{d+ex}}{2e^3} + \frac{5\sqrt{b}(bd - ae)(7bBd - 4Abe - 3aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{4e^{9/2}}$$

output

```
2/3*(-A*e+B*d)*(b*x+a)^(5/2)/e^2/(e*x+d)^(3/2)+2/3*(-5*A*b*e-3*B*a*e+8*B*b
*d)*(b*x+a)^(3/2)/e^3/(e*x+d)^(1/2)-5/4*b*(-4*A*b*e-3*B*a*e+7*B*b*d)*(b*x+
a)^(1/2)*(e*x+d)^(1/2)/e^4+1/2*b*B*(b*x+a)^(3/2)*(e*x+d)^(1/2)/e^3+5/4*b^(
1/2)*(-a*e+b*d)*(-4*A*b*e-3*B*a*e+7*B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b
^(1/2)/(e*x+d)^(1/2))/e^(9/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 10.10 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.53

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{5/2}} dx = \frac{2(a+bx)^{7/2} \left(7Bd - 7Ae - \frac{(7bBd - 4Abe - 3aBe) \left(\frac{b(d+ex)}{bd-ae} \right)^{3/2} \text{Hypergeometric2F1} \left(\frac{3}{2}, \frac{7}{2}, \frac{9}{2}, \frac{b(d+ex)}{bd-ae} \right)}{b} \right)}{21e(-bd+ae)(d+ex)^{3/2}}$$

input

```
Integrate[((a + b*x)^(5/2)*(A + B*x))/(d + e*x)^(5/2), x]
```

output

```
(2*(a + b*x)^(7/2)*(7*B*d - 7*A*e - ((7*b*B*d - 4*A*b*e - 3*a*B*e)*((b*(d + e*x))/(b*d - a*e))^(3/2)*Hypergeometric2F1[3/2, 7/2, 9/2, (e*(a + b*x))/(-b*d) + a*e])/b)/(21*e*(-b*d) + a*e)*(d + e*x)^(3/2))
```

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.04, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {87, 57, 60, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{5/2}} dx \\ & \quad \downarrow 87 \\ & \frac{(-3aBe - 4Abe + 7bBd) \int \frac{(a+bx)^{5/2}}{(d+ex)^{3/2}} dx}{3e(bd-ae)} - \frac{2(a+bx)^{7/2}(Bd-Ae)}{3e(d+ex)^{3/2}(bd-ae)} \\ & \quad \downarrow 57 \\ & \frac{(-3aBe - 4Abe + 7bBd) \left(\frac{5b \int \frac{(a+bx)^{3/2}}{\sqrt{d+ex}} dx}{e} - \frac{2(a+bx)^{5/2}}{e\sqrt{d+ex}} \right)}{3e(bd-ae)} - \frac{2(a+bx)^{7/2}(Bd-Ae)}{3e(d+ex)^{3/2}(bd-ae)} \end{aligned}$$

$$\begin{array}{c}
 \downarrow 60 \\
 (-3aBe - 4Abe + 7bBd) \left(\frac{5b \left(\frac{(a+bx)^{3/2} \sqrt{d+ex}}{2e} - \frac{3(bd-ae) \int \frac{\sqrt{a+bx}}{\sqrt{d+ex}} dx}{4e} \right)}{e} - \frac{2(a+bx)^{5/2}}{e\sqrt{d+ex}} \right) \\
 \hline
 \frac{3e(bd - ae)}{2(a + bx)^{7/2}(Bd - Ae)} \\
 \frac{3e(d + ex)^{3/2}(bd - ae)}{2(a + bx)^{7/2}(Bd - Ae)} \\
 \downarrow 60 \\
 (-3aBe - 4Abe + 7bBd) \left(\frac{5b \left(\frac{(a+bx)^{3/2} \sqrt{d+ex}}{2e} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx} \sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{\sqrt{a+bx} \sqrt{d+ex}} dx}{2e} \right)}{4e} \right)}{e} - \frac{2(a+bx)^{5/2}}{e\sqrt{d+ex}} \right) \\
 \hline
 \frac{3e(bd - ae)}{2(a + bx)^{7/2}(Bd - Ae)} \\
 \frac{3e(d + ex)^{3/2}(bd - ae)}{2(a + bx)^{7/2}(Bd - Ae)} \\
 \downarrow 66 \\
 (-3aBe - 4Abe + 7bBd) \left(\frac{5b \left(\frac{(a+bx)^{3/2} \sqrt{d+ex}}{2e} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx} \sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{e} \right)}{4e} \right)}{e} - \frac{2(a+bx)^{5/2}}{e\sqrt{d+ex}} \right) \\
 \hline
 \frac{3e(bd - ae)}{2(a + bx)^{7/2}(Bd - Ae)} \\
 \frac{3e(d + ex)^{3/2}(bd - ae)}{2(a + bx)^{7/2}(Bd - Ae)} \\
 \downarrow 221
 \end{array}$$

$$\begin{aligned}
 & \left((-3aBe - 4Abe + 7bBd) \frac{5b \left(\frac{(a+bx)^{3/2} \sqrt{d+ex}}{2e} - \frac{3(bd-ae) \left(\frac{\sqrt{a+bx} \sqrt{d+ex}}{e} - \frac{(bd-ae) \operatorname{arctanh} \left(\frac{\sqrt{e} \sqrt{a+bx}}{\sqrt{b} \sqrt{d+ex}} \right)}{\sqrt{be}^{3/2}} \right)}{4e} \right)}{e} - \frac{2(a+bx)^{5/2}}{e \sqrt{d+ex}} \right) \\
 & \frac{3e(bd-ae)}{2(a+bx)^{7/2}(Bd-Ae)} \\
 & \frac{3e(d+ex)^{3/2}(bd-ae)}{3e(d+ex)^{3/2}(bd-ae)}
 \end{aligned}$$

input `Int[((a + b*x)^(5/2)*(A + B*x))/(d + e*x)^(5/2),x]`

output `(-2*(B*d - A*e)*(a + b*x)^(7/2))/(3*e*(b*d - a*e)*(d + e*x)^(3/2)) + ((7*b*B*d - 4*A*b*e - 3*a*B*e)*((-2*(a + b*x)^(5/2))/(e*Sqrt[d + e*x]) + (5*b*((a + b*x)^(3/2)*Sqrt[d + e*x])/(2*e) - (3*(b*d - a*e)*((Sqrt[a + b*x]*Sqrt[d + e*x])/e - ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(Sqrt[b]*e^(3/2))))/(4*e)))/e)/(3*e*(b*d - a*e))`

Defintions of rubi rules used

rule 57 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))] Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)))] Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[
2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; Fre
eQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p
), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))`

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1249 vs. $2(174) = 348$.

Time = 0.27 (sec) , antiderivative size = 1250, normalized size of antiderivative = 5.90

method	result	size
default	Expression too large to display	1250

input `int((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(5/2), x, method=_RETURNVERBOSE)`

output

```

1/24*(b*x+a)^(1/2)*(-16*A*a^2*e^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-210*
B*b^2*d^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+316*B*a*b*d*e^2*x*((e*x+d)*
(b*x+a))^(1/2)*(b*e)^(1/2)+105*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*
(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^3*d^2*e^2*x^2+24*A*b^2*e^3*x^2*((e*x+d)
*(b*x+a))^(1/2)*(b*e)^(1/2)+210*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)
*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^3*d^3*e*x-150*B*ln(1/2*(2*b*e*x+2*(
(e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^2*d^3*e-48*B*
a^2*e^3*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-32*B*a^2*d*e^2*((e*x+d)*(b*x
+a))^(1/2)*(b*e)^(1/2)-150*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*
e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^2*d*e^3*x^2-300*B*ln(1/2*(2*b*e*x+2*(e
*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^2*d^2*e^2*x+54*
B*a*b*e^3*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-42*B*b^2*d*e^2*x^2*((e*x
+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-112*A*a*b*e^3*x*((e*x+d)*(b*x+a))^(1/2)*(b*
e)^(1/2)-280*B*b^2*d^2*e*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+160*A*b^2*d
*e^2*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-80*A*a*b*d*e^2*((e*x+d)*(b*x+a)
)^(1/2)*(b*e)^(1/2)+120*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(
1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^2*d*e^3*x+90*B*ln(1/2*(2*b*e*x+2*((e*x+d)*
(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*b*d*e^3*x+105*B*ln(1/
2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^3
*d^4+230*B*a*b*d^2*e*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+60*A*ln(1/2*(2...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 415 vs. $2(174) = 348$.

Time = 1.31 (sec) , antiderivative size = 855, normalized size of antiderivative = 4.03

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{5/2}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(5/2),x, algorithm="fricas")
```

output

```
[1/48*(15*(7*B*b^2*d^4 - 2*(5*B*a*b + 2*A*b^2)*d^3*e + (3*B*a^2 + 4*A*a*b)
*d^2*e^2 + (7*B*b^2*d^2*e^2 - 2*(5*B*a*b + 2*A*b^2)*d*e^3 + (3*B*a^2 + 4*A
*a*b)*e^4)*x^2 + 2*(7*B*b^2*d^3*e - 2*(5*B*a*b + 2*A*b^2)*d^2*e^2 + (3*B*a
^2 + 4*A*a*b)*d*e^3)*x)*sqrt(b/e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e
+ a^2*e^2 + 4*(2*b*e^2*x + b*d*e + a*e^2)*sqrt(b*x + a)*sqrt(e*x + d)*sqrt
(b/e) + 8*(b^2*d*e + a*b*e^2)*x) + 4*(6*B*b^2*e^3*x^3 - 105*B*b^2*d^3 - 8*
A*a^2*e^3 + 5*(23*B*a*b + 12*A*b^2)*d^2*e - 8*(2*B*a^2 + 5*A*a*b)*d*e^2 -
3*(7*B*b^2*d*e^2 - (9*B*a*b + 4*A*b^2)*e^3)*x^2 - 2*(70*B*b^2*d^2*e - (79*
B*a*b + 40*A*b^2)*d*e^2 + 4*(3*B*a^2 + 7*A*a*b)*e^3)*x)*sqrt(b*x + a)*sqrt
(e*x + d))/(e^6*x^2 + 2*d*e^5*x + d^2*e^4), -1/24*(15*(7*B*b^2*d^4 - 2*(5*
B*a*b + 2*A*b^2)*d^3*e + (3*B*a^2 + 4*A*a*b)*d^2*e^2 + (7*B*b^2*d^2*e^2 -
2*(5*B*a*b + 2*A*b^2)*d*e^3 + (3*B*a^2 + 4*A*a*b)*e^4)*x^2 + 2*(7*B*b^2*d^
3*e - 2*(5*B*a*b + 2*A*b^2)*d^2*e^2 + (3*B*a^2 + 4*A*a*b)*d*e^3)*x)*sqrt(-
b/e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(b*x + a)*sqrt(e*x + d)*sqrt(-b/
e)/(b^2*e*x^2 + a*b*d + (b^2*d + a*b*e)*x)) - 2*(6*B*b^2*e^3*x^3 - 105*B*b
^2*d^3 - 8*A*a^2*e^3 + 5*(23*B*a*b + 12*A*b^2)*d^2*e - 8*(2*B*a^2 + 5*A*a*
b)*d*e^2 - 3*(7*B*b^2*d*e^2 - (9*B*a*b + 4*A*b^2)*e^3)*x^2 - 2*(70*B*b^2*d
^2*e - (79*B*a*b + 40*A*b^2)*d*e^2 + 4*(3*B*a^2 + 7*A*a*b)*e^3)*x)*sqrt(b*
x + a)*sqrt(e*x + d))/(e^6*x^2 + 2*d*e^5*x + d^2*e^4)]
```

Sympy [F]

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{5/2}} dx = \int \frac{(A + Bx)(a + bx)^{5/2}}{(d + ex)^{5/2}} dx$$

input

```
integrate((b*x+a)**(5/2)*(B*x+A)/(e*x+d)**(5/2),x)
```

output

```
Integral((A + B*x)*(a + b*x)**(5/2)/(d + e*x)**(5/2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(5/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f or more de`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 564 vs. 2(174) = 348.

Time = 0.24 (sec) , antiderivative size = 564, normalized size of antiderivative = 2.66

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{5/2}} dx = \frac{\left(\left(3(bx+a) \left(\frac{2(Bb^5de^6|b|-Bab^4e^7|b|)(bx+a)}{b^4de^7-ab^3e^8} - \frac{7Bb^6d^2e^5|b|-10Bab^5de^6|b|-4Ab^6de^6|b|+3Ba^6de^6|b|}{b^4de^7-ab^3e^8} \right) \right) \log \left(\left| -\sqrt{be}\sqrt{bx+a} + \sqrt{b^2d+(bx+a)} \right| \right)}{4\sqrt{bee^4}}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(5/2),x, algorithm="giac")`

output

```
1/12*((3*(b*x + a)*(2*(B*b^5*d*e^6*abs(b) - B*a*b^4*e^7*abs(b))*(b*x + a)/
(b^4*d*e^7 - a*b^3*e^8) - (7*B*b^6*d^2*e^5*abs(b) - 10*B*a*b^5*d*e^6*abs(b)
) - 4*A*b^6*d*e^6*abs(b) + 3*B*a^2*b^4*e^7*abs(b) + 4*A*a*b^5*e^7*abs(b))/
(b^4*d*e^7 - a*b^3*e^8)) - 20*(7*B*b^7*d^3*e^4*abs(b) - 17*B*a*b^6*d^2*e^5
*abs(b) - 4*A*b^7*d^2*e^5*abs(b) + 13*B*a^2*b^5*d*e^6*abs(b) + 8*A*a*b^6*d
*e^6*abs(b) - 3*B*a^3*b^4*e^7*abs(b) - 4*A*a^2*b^5*e^7*abs(b))/(b^4*d*e^7
- a*b^3*e^8)*(b*x + a) - 15*(7*B*b^8*d^4*e^3*abs(b) - 24*B*a*b^7*d^3*e^4*
abs(b) - 4*A*b^8*d^3*e^4*abs(b) + 30*B*a^2*b^6*d^2*e^5*abs(b) + 12*A*a*b^7
*d^2*e^5*abs(b) - 16*B*a^3*b^5*d*e^6*abs(b) - 12*A*a^2*b^6*d*e^6*abs(b) +
3*B*a^4*b^4*e^7*abs(b) + 4*A*a^3*b^5*e^7*abs(b))/(b^4*d*e^7 - a*b^3*e^8))*
sqrt(b*x + a)/(b^2*d + (b*x + a)*b*e - a*b*e)^(3/2) - 5/4*(7*B*b^2*d^2*abs
(b) - 10*B*a*b*d*e*abs(b) - 4*A*b^2*d*e*abs(b) + 3*B*a^2*e^2*abs(b) + 4*A*
a*b*e^2*abs(b))*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*
b*e - a*b*e)))/(sqrt(b*e)*e^4)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{5/2}} dx = \int \frac{(A + Bx)(a + bx)^{5/2}}{(d + ex)^{5/2}} dx$$

input

```
int(((A + B*x)*(a + b*x)^(5/2))/(d + e*x)^(5/2),x)
```

output

```
int(((A + B*x)*(a + b*x)^(5/2))/(d + e*x)^(5/2), x)
```

Reduce [B] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 817, normalized size of antiderivative = 3.85

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{5/2}} dx = \text{Too large to display}$$

input

```
int((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(5/2),x)
```

output

```
( - 64*sqrt(d + e*x)*sqrt(a + b*x)*a**3*e**4 - 448*sqrt(d + e*x)*sqrt(a +
b*x)*a**2*b*d*e**3 - 640*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b*e**4*x + 1400*
sqrt(d + e*x)*sqrt(a + b*x)*a*b**2*d**2*e**2 + 1904*sqrt(d + e*x)*sqrt(a +
b*x)*a*b**2*d*e**3*x + 312*sqrt(d + e*x)*sqrt(a + b*x)*a*b**2*e**4*x**2 -
840*sqrt(d + e*x)*sqrt(a + b*x)*b**3*d**3*e - 1120*sqrt(d + e*x)*sqrt(a +
b*x)*b**3*d**2*e**2*x - 168*sqrt(d + e*x)*sqrt(a + b*x)*b**3*d*e**3*x**2
+ 48*sqrt(d + e*x)*sqrt(a + b*x)*b**3*e**4*x**3 + 840*sqrt(e)*sqrt(b)*log(
(sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**2*b*d*
**2*e**2 + 1680*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d
+ e*x))/sqrt(a*e - b*d))*a**2*b*d*e**3*x + 840*sqrt(e)*sqrt(b)*log((sqrt(
e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**2*b*e**4*x**
2 - 1680*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x
))/sqrt(a*e - b*d))*a*b**2*d**3*e - 3360*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt
(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*b**2*d**2*e**2*x - 1
680*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sq
rt(a*e - b*d))*a*b**2*d*e**3*x**2 + 840*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(
a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**3*d**4 + 1680*sqrt(e
)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b
*d))*b**3*d**3*e*x + 840*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt
(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**3*d**2*e**2*x**2 + 175*sqrt(e)*s...
```


3.199
$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{7/2}} dx$$

Optimal result	1820
Mathematica [A] (verified)	1821
Rubi [A] (verified)	1821
Maple [B] (verified)	1824
Fricas [B] (verification not implemented)	1825
Sympy [F]	1826
Maxima [F(-2)]	1827
Giac [B] (verification not implemented)	1827
Mupad [F(-1)]	1828
Reduce [B] (verification not implemented)	1829

Optimal result

Integrand size = 24, antiderivative size = 199

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{7/2}} dx = \frac{2(Bd - Ae)(a+bx)^{5/2}}{5e^2(d+ex)^{5/2}} + \frac{2(2bBd - Abe - aBe)(a+bx)^{3/2}}{3e^3(d+ex)^{3/2}} + \frac{2b(3bBd - Abe - 2aBe)\sqrt{a+bx}}{e^4\sqrt{d+ex}} + \frac{b^2B\sqrt{a+bx}\sqrt{d+ex}}{e^4} - \frac{b^{3/2}(7bBd - 2Abe - 5aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{e^{9/2}}$$

output
$$\begin{aligned} & 2/5*(-A*e+B*d)*(b*x+a)^{(5/2)}/e^2/(e*x+d)^{(5/2)}+2/3*(-A*b*e-B*a*e+2*B*b*d)* \\ & (b*x+a)^{(3/2)}/e^3/(e*x+d)^{(3/2)}+2*b*(-A*b*e-2*B*a*e+3*B*b*d)*(b*x+a)^{(1/2)} \\ & /e^4/(e*x+d)^{(1/2)}+b^2*B*(b*x+a)^{(1/2)*(e*x+d)^{(1/2)}/e^4-b^{(3/2)*(-2*A*b*e} \\ & -5*B*a*e+7*B*b*d)*\operatorname{arctanh}(e^{(1/2)*(b*x+a)^{(1/2)}/b^{(1/2)/(e*x+d)^{(1/2)}}/e^{(} \\ & 9/2) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.03

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{7/2}} dx = \frac{\sqrt{a+bx}(2a^2e^2(2Bd+3Ae+5Bex) + 2abe(Ae(5d+11ex) + B(20d^2+49dex+35e^2x^2)) + b^2(2Ae(15d^2+11dex+5e^2x^2) + B(20d^2+49dex+35e^2x^2)))}{15e^4(d+ex)^{5/2}} + \frac{b^{3/2}(-7bBd+2Abe+5aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{e}\sqrt{a+bx}}\right)}{e^{9/2}}$$

input `Integrate[((a + b*x)^(5/2)*(A + B*x))/(d + e*x)^(7/2),x]`

output `-1/15*(Sqrt[a + b*x]*(2*a^2*e^2*(2*B*d + 3*A*e + 5*B*e*x) + 2*a*b*e*(A*e*(5*d + 11*e*x) + B*(20*d^2 + 49*d*e*x + 35*e^2*x^2)) + b^2*(2*A*e*(15*d^2 + 35*d*e*x + 23*e^2*x^2) - B*(105*d^3 + 245*d^2*e*x + 161*d*e^2*x^2 + 15*e^3*x^3)))/(e^4*(d + e*x)^(5/2)) + (b^(3/2)*(-7*b*B*d + 2*A*b*e + 5*a*B*e)*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/(Sqrt[e]*Sqrt[a + b*x])])/e^(9/2)`

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.08, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {87, 57, 57, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{7/2}} dx$$

$$\downarrow 87$$

$$\frac{(-5aBe - 2Abe + 7bBd) \int \frac{(a+bx)^{5/2}}{(d+ex)^{5/2}} dx}{5e(bd - ae)} - \frac{2(a+bx)^{7/2}(Bd - Ae)}{5e(d+ex)^{5/2}(bd - ae)}$$

$$\downarrow 57$$

$$\begin{aligned}
 & \frac{(-5aBe - 2Abe + 7bBd) \left(\frac{5b \int \frac{(a+bx)^{3/2} dx}{(d+ex)^{3/2}} - \frac{2(a+bx)^{5/2}}{3e(d+ex)^{3/2}}}{3e} \right)}{5e(bd - ae)} - \frac{2(a+bx)^{7/2}(Bd - Ae)}{5e(d+ex)^{5/2}(bd - ae)} \\
 & \quad \downarrow 57 \\
 & \frac{(-5aBe - 2Abe + 7bBd) \left(\frac{5b \left(\frac{3b \int \frac{\sqrt{a+bx}}{\sqrt{d+ex}} dx}{e} - \frac{2(a+bx)^{3/2}}{e\sqrt{d+ex}} \right)}{3e} - \frac{2(a+bx)^{5/2}}{3e(d+ex)^{3/2}} \right)}{5e(bd - ae)} - \frac{2(a+bx)^{7/2}(Bd - Ae)}{5e(d+ex)^{5/2}(bd - ae)} \\
 & \quad \downarrow 60 \\
 & \frac{(-5aBe - 2Abe + 7bBd) \left(\frac{5b \left(\frac{3b \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{2e} \right)}{e} - \frac{2(a+bx)^{3/2}}{e\sqrt{d+ex}} \right)}{3e} - \frac{2(a+bx)^{5/2}}{3e(d+ex)^{3/2}} \right)}{5e(bd - ae)} - \frac{2(a+bx)^{7/2}(Bd - Ae)}{5e(d+ex)^{5/2}(bd - ae)} \\
 & \quad \downarrow 66 \\
 & \frac{(-5aBe - 2Abe + 7bBd) \left(\frac{5b \left(\frac{3b \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{e} \right)}{e} - \frac{2(a+bx)^{3/2}}{e\sqrt{d+ex}} \right)}{3e} - \frac{2(a+bx)^{5/2}}{3e(d+ex)^{3/2}} \right)}{5e(bd - ae)} - \frac{2(a+bx)^{7/2}(Bd - Ae)}{5e(d+ex)^{5/2}(bd - ae)}
 \end{aligned}$$

↓ 221

$$(-5aBe - 2Abe + 7bBd) \left(\frac{5b \left(\frac{3b \left(\frac{\sqrt{a+bx}\sqrt{d+ex}}{e} - \frac{(bd-ae)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{\sqrt{be}^{3/2}} \right)}{e} - \frac{2(a+bx)^{3/2}}{e\sqrt{d+ex}} \right)}{3e} - \frac{2(a+bx)^{5/2}}{3e(d+ex)^{3/2}} \right)}{5e(bd-ae) + \frac{2(a+bx)^{7/2}(Bd-Ae)}{5e(d+ex)^{5/2}(bd-ae)}}$$

input `Int[((a + b*x)^(5/2)*(A + B*x))/(d + e*x)^(7/2), x]`

output `(-2*(B*d - A*e)*(a + b*x)^(7/2))/(5*e*(b*d - a*e)*(d + e*x)^(5/2)) + ((7*b*B*d - 2*A*b*e - 5*a*B*e)*((-2*(a + b*x)^(5/2))/(3*e*(d + e*x)^(3/2)) + (5*b*((-2*(a + b*x)^(3/2))/(e*Sqrt[d + e*x]) + (3*b*((Sqrt[a + b*x]*Sqrt[d + e*x])/e - ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])]))/(Sqrt[b]*e^(3/2))))/e)/(3*e))/(5*e*(b*d - a*e))`

Defintions of rubi rules used

rule 57 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))] Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1091 vs. $2(167) = 334$.

Time = 0.27 (sec) , antiderivative size = 1092, normalized size of antiderivative = 5.49

method	result	size
default	Expression too large to display	1092

input `int((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(7/2),x,method=_RETURNVERBOSE)`

output

```

1/30*(30*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/
(b*e)^(1/2))*b^3*e^4*x^3-12*A*a^2*e^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+
210*B*b^2*d^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-196*B*a*b*d*e^2*x*((e*x+
d)*(b*x+a))^(1/2)*(b*e)^(1/2)+75*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/
2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^2*e^4*x^3-105*B*ln(1/2*(2*b*e*x+2
*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^3*d*e^3*x^3-3
15*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(
1/2))*b^3*d^2*e^2*x^2-92*A*b^2*e^3*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)
)-315*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*
e)^(1/2))*b^3*d^3*e*x+75*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)
^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^2*d^3*e-20*B*a^2*e^3*x*((e*x+d)*(b*x+a))^(
1/2)*(b*e)^(1/2)-8*B*a^2*d*e^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+225*B*
ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2)
)*a*b^2*d*e^3*x^2+225*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1
/2)+a*e+d*b)/(b*e)^(1/2))*a*b^2*d^2*e^2*x-140*B*a*b*e^3*x^2*((e*x+d)*(b*x+
a))^(1/2)*(b*e)^(1/2)+322*B*b^2*d*e^2*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1
/2)-44*A*a*b*e^3*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+490*B*b^2*d^2*e*x*((
e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-140*A*b^2*d*e^2*x*((e*x+d)*(b*x+a))^(1/
2)*(b*e)^(1/2)-20*A*a*b*d*e^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-105*B*ln
(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2)...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 405 vs. $2(167) = 334$.

Time = 3.22 (sec) , antiderivative size = 835, normalized size of antiderivative = 4.20

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{7/2}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(7/2),x, algorithm="fricas")
```

output

```

[-1/60*(15*(7*B*b^2*d^4 - (5*B*a*b + 2*A*b^2)*d^3*e + (7*B*b^2*d*e^3 - (5*
B*a*b + 2*A*b^2)*e^4)*x^3 + 3*(7*B*b^2*d^2*e^2 - (5*B*a*b + 2*A*b^2)*d*e^3
)*x^2 + 3*(7*B*b^2*d^3*e - (5*B*a*b + 2*A*b^2)*d^2*e^2)*x)*sqrt(b/e)*log(8
*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*b*e^2*x + b*d*e + a*e^
2)*sqrt(b*x + a)*sqrt(e*x + d)*sqrt(b/e) + 8*(b^2*d*e + a*b*e^2)*x) - 4*(1
5*B*b^2*e^3*x^3 + 105*B*b^2*d^3 - 6*A*a^2*e^3 - 10*(4*B*a*b + 3*A*b^2)*d^2
*e - 2*(2*B*a^2 + 5*A*a*b)*d*e^2 + (161*B*b^2*d*e^2 - 2*(35*B*a*b + 23*A*b
^2)*e^3)*x^2 + (245*B*b^2*d^2*e - 14*(7*B*a*b + 5*A*b^2)*d*e^2 - 2*(5*B*a^
2 + 11*A*a*b)*e^3)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(e^7*x^3 + 3*d*e^6*x^2
+ 3*d^2*e^5*x + d^3*e^4), 1/30*(15*(7*B*b^2*d^4 - (5*B*a*b + 2*A*b^2)*d^3*
e + (7*B*b^2*d*e^3 - (5*B*a*b + 2*A*b^2)*e^4)*x^3 + 3*(7*B*b^2*d^2*e^2 - (
5*B*a*b + 2*A*b^2)*d*e^3)*x^2 + 3*(7*B*b^2*d^3*e - (5*B*a*b + 2*A*b^2)*d^2
*e^2)*x)*sqrt(-b/e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(b*x + a)*sqrt(e*
x + d)*sqrt(-b/e)/(b^2*e*x^2 + a*b*d + (b^2*d + a*b*e)*x)) + 2*(15*B*b^2*e
^3*x^3 + 105*B*b^2*d^3 - 6*A*a^2*e^3 - 10*(4*B*a*b + 3*A*b^2)*d^2*e - 2*(2
*B*a^2 + 5*A*a*b)*d*e^2 + (161*B*b^2*d*e^2 - 2*(35*B*a*b + 23*A*b^2)*e^3)*
x^2 + (245*B*b^2*d^2*e - 14*(7*B*a*b + 5*A*b^2)*d*e^2 - 2*(5*B*a^2 + 11*A*
a*b)*e^3)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(e^7*x^3 + 3*d*e^6*x^2 + 3*d^2*
e^5*x + d^3*e^4)]

```

Sympy [F]

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{7/2}} dx = \int \frac{(A + Bx)(a + bx)^{5/2}}{(d + ex)^{7/2}} dx$$

input

```
integrate((b*x+a)**(5/2)*(B*x+A)/(e*x+d)**(7/2),x)
```

output

```
Integral((A + B*x)*(a + b*x)**(5/2)/(d + e*x)**(7/2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{7/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(7/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f or more de`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 707 vs. 2(167) = 334.

Time = 0.30 (sec) , antiderivative size = 707, normalized size of antiderivative = 3.55

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{7/2}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(7/2),x, algorithm="giac")`

output

```

1/15*(((b*x + a)*(15*(B*b^9*d^2*e^6*abs(b) - 2*B*a*b^8*d*e^7*abs(b) + B*a^
2*b^7*e^8*abs(b))*(b*x + a)/(b^6*d^2*e^7 - 2*a*b^5*d*e^8 + a^2*b^4*e^9) +
23*(7*B*b^10*d^3*e^5*abs(b) - 19*B*a*b^9*d^2*e^6*abs(b) - 2*A*b^10*d^2*e^6
*abs(b) + 17*B*a^2*b^8*d*e^7*abs(b) + 4*A*a*b^9*d*e^7*abs(b) - 5*B*a^3*b^7
*e^8*abs(b) - 2*A*a^2*b^8*e^8*abs(b))/(b^6*d^2*e^7 - 2*a*b^5*d*e^8 + a^2*b
^4*e^9)) + 35*(7*B*b^11*d^4*e^4*abs(b) - 26*B*a*b^10*d^3*e^5*abs(b) - 2*A*
b^11*d^3*e^5*abs(b) + 36*B*a^2*b^9*d^2*e^6*abs(b) + 6*A*a*b^10*d^2*e^6*abs
(b) - 22*B*a^3*b^8*d*e^7*abs(b) - 6*A*a^2*b^9*d*e^7*abs(b) + 5*B*a^4*b^7*e
^8*abs(b) + 2*A*a^3*b^8*e^8*abs(b))/(b^6*d^2*e^7 - 2*a*b^5*d*e^8 + a^2*b^4
*e^9))*(b*x + a) + 15*(7*B*b^12*d^5*e^3*abs(b) - 33*B*a*b^11*d^4*e^4*abs(b)
) - 2*A*b^12*d^4*e^4*abs(b) + 62*B*a^2*b^10*d^3*e^5*abs(b) + 8*A*a*b^11*d^
3*e^5*abs(b) - 58*B*a^3*b^9*d^2*e^6*abs(b) - 12*A*a^2*b^10*d^2*e^6*abs(b)
+ 27*B*a^4*b^8*d*e^7*abs(b) + 8*A*a^3*b^9*d*e^7*abs(b) - 5*B*a^5*b^7*e^8*a
bs(b) - 2*A*a^4*b^8*e^8*abs(b))/(b^6*d^2*e^7 - 2*a*b^5*d*e^8 + a^2*b^4*e^9
))*sqrt(b*x + a)/(b^2*d + (b*x + a)*b*e - a*b*e)^(5/2) + (7*B*b^2*d*abs(b)
- 5*B*a*b*e*abs(b) - 2*A*b^2*e*abs(b))*log(abs(-sqrt(b*e)*sqrt(b*x + a) +
sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*e^4)

```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{7/2}} dx = \int \frac{(A + Bx)(a + bx)^{5/2}}{(d + ex)^{7/2}} dx$$

input

```
int(((A + B*x)*(a + b*x)^(5/2))/(d + e*x)^(7/2), x)
```

output

```
int(((A + B*x)*(a + b*x)^(5/2))/(d + e*x)^(7/2), x)
```

Reduce [B] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 766, normalized size of antiderivative = 3.85

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{7/2}} dx = \text{Too large to display}$$

input `int((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(7/2),x)`

output

```
( - 24*sqrt(d + e*x)*sqrt(a + b*x)*a**3*e**4 - 56*sqrt(d + e*x)*sqrt(a + b
*x)*a**2*b*d*e**3 - 128*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b*e**4*x - 280*sq
rt(d + e*x)*sqrt(a + b*x)*a*b**2*d**2*e**2 - 672*sqrt(d + e*x)*sqrt(a + b*
x)*a*b**2*d*e**3*x - 464*sqrt(d + e*x)*sqrt(a + b*x)*a*b**2*e**4*x**2 + 42
0*sqrt(d + e*x)*sqrt(a + b*x)*b**3*d**3*e + 980*sqrt(d + e*x)*sqrt(a + b*x
)*b**3*d**2*e**2*x + 644*sqrt(d + e*x)*sqrt(a + b*x)*b**3*d*e**3*x**2 + 60
*sqrt(d + e*x)*sqrt(a + b*x)*b**3*e**4*x**3 + 420*sqrt(e)*sqrt(b)*log((sqr
t(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*b**2*d**3*e
+ 1260*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x)
)/sqrt(a*e - b*d))*a*b**2*d**2*e**2*x + 1260*sqrt(e)*sqrt(b)*log((sqrt(e)*
sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*b**2*d*e**3*x**2
+ 420*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))
/sqrt(a*e - b*d))*a*b**2*e**4*x**3 - 420*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt
(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**3*d**4 - 1260*sqrt(
e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e -
b*d))*b**3*d**3*e*x - 1260*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sq
rt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**3*d**2*e**2*x**2 - 420*sqrt(e)*sq
rt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))
*b**3*d*e**3*x**3 + 203*sqrt(e)*sqrt(b)*a*b**2*d**3*e + 609*sqrt(e)*sqrt(b
)*a*b**2*d**2*e**2*x + 609*sqrt(e)*sqrt(b)*a*b**2*d*e**3*x**2 + 203*sqr...
```

3.200 $\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{9/2}} dx$

Optimal result	1830
Mathematica [A] (verified)	1830
Rubi [A] (verified)	1831
Maple [B] (verified)	1834
Fricas [B] (verification not implemented)	1835
Sympy [F]	1836
Maxima [F(-2)]	1836
Giac [B] (verification not implemented)	1836
Mupad [F(-1)]	1837
Reduce [B] (verification not implemented)	1837

Optimal result

Integrand size = 24, antiderivative size = 167

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{9/2}} dx = -\frac{2(Bd - Ae)(a+bx)^{7/2}}{7e(bd - ae)(d+ex)^{7/2}} - \frac{2B(a+bx)^{5/2}}{5e^2(d+ex)^{5/2}} - \frac{2bB(a+bx)^{3/2}}{3e^3(d+ex)^{3/2}} - \frac{2b^2B\sqrt{a+bx}}{e^4\sqrt{d+ex}} + \frac{2b^{5/2}B\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{e^{9/2}}$$

output

```
-2/7*(-A*e+B*d)*(b*x+a)^(7/2)/e/(-a*e+b*d)/(e*x+d)^(7/2)-2/5*B*(b*x+a)^(5/2)/e^2/(e*x+d)^(5/2)-2/3*b*B*(b*x+a)^(3/2)/e^3/(e*x+d)^(3/2)-2*b^2*B*(b*x+a)^(1/2)/e^4/(e*x+d)^(1/2)+2*b^(5/2)*B*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)^(1/2))/e^(9/2)
```

Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.29

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{9/2}} dx = \frac{2\sqrt{a+bx}(-15Bde^3(a+bx)^3 + 15Ae^4(a+bx)^3 - 21bBde^2(a+bx)^2(d+ex) + 21aBe^3(a+bx)^2(d+ex) + 105e^4(-bd+ax))}{105e^4(-bd+ax)^{5/2}} + \frac{2b^{5/2}B\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{e}\sqrt{a+bx}}\right)}{e^{9/2}}$$

input `Integrate[((a + b*x)^(5/2)*(A + B*x))/(d + e*x)^(9/2),x]`

output `(-2*sqrt[a + b*x]*(-15*B*d*e^3*(a + b*x)^3 + 15*A*e^4*(a + b*x)^3 - 21*b*B*d*e^2*(a + b*x)^2*(d + e*x) + 21*a*B*e^3*(a + b*x)^2*(d + e*x) - 35*b^2*B*d*e*(a + b*x)*(d + e*x)^2 + 35*a*b*B*e^2*(a + b*x)*(d + e*x)^2 - 105*b^3*B*d*(d + e*x)^3 + 105*a*b^2*B*e*(d + e*x)^3)/(105*e^4*(-(b*d) + a*e)*(d + e*x)^(7/2)) + (2*b^(5/2)*B*ArcTanh[(sqrt[b]*sqrt[d + e*x])/(sqrt[e]*sqrt[a + b*x])])/e^(9/2)`

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.06, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {87, 57, 57, 57, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{9/2}} dx \\
 & \quad \downarrow 87 \\
 & \frac{B \int \frac{(a+bx)^{5/2} dx}{(d+ex)^{7/2}}}{e} - \frac{2(a+bx)^{7/2}(Bd - Ae)}{7e(d+ex)^{7/2}(bd - ae)} \\
 & \quad \downarrow 57 \\
 & \frac{B \left(\frac{b \int \frac{(a+bx)^{3/2} dx}{(d+ex)^{5/2}}}{e} - \frac{2(a+bx)^{5/2}}{5e(d+ex)^{5/2}} \right)}{e} - \frac{2(a+bx)^{7/2}(Bd - Ae)}{7e(d+ex)^{7/2}(bd - ae)} \\
 & \quad \downarrow 57 \\
 & \frac{B \left(\frac{b \left(\frac{b \int \frac{\sqrt{a+bx}}{(d+ex)^{3/2}} dx}{e} - \frac{2(a+bx)^{3/2}}{3e(d+ex)^{3/2}} \right)}{e} - \frac{2(a+bx)^{5/2}}{5e(d+ex)^{5/2}} \right)}{e} - \frac{2(a+bx)^{7/2}(Bd - Ae)}{7e(d+ex)^{7/2}(bd - ae)}
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow 57 \\
 B \left(\frac{b \left(\frac{b \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{e} - \frac{2\sqrt{a+bx}}{e\sqrt{d+ex}} \right) - \frac{2(a+bx)^{3/2}}{3e(d+ex)^{3/2}}}{e} - \frac{2(a+bx)^{5/2}}{5e(d+ex)^{5/2}} \right) \\
 \hline
 e - \frac{2(a+bx)^{7/2}(Bd - Ae)}{7e(d+ex)^{7/2}(bd - ae)}
 \end{array}$$

$$\begin{array}{c}
 \downarrow 66 \\
 B \left(\frac{b \left(\frac{2b \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{e} - \frac{2\sqrt{a+bx}}{e\sqrt{d+ex}} \right) - \frac{2(a+bx)^{3/2}}{3e(d+ex)^{3/2}}}{e} - \frac{2(a+bx)^{5/2}}{5e(d+ex)^{5/2}} \right) \\
 \hline
 e - \frac{2(a+bx)^{7/2}(Bd - Ae)}{7e(d+ex)^{7/2}(bd - ae)}
 \end{array}$$

$$\begin{array}{c}
 \downarrow 221 \\
 B \left(\frac{b \left(\frac{2\sqrt{b} \operatorname{arctanh} \left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}} \right)}{e^{3/2}} - \frac{2\sqrt{a+bx}}{e\sqrt{d+ex}} \right) - \frac{2(a+bx)^{3/2}}{3e(d+ex)^{3/2}}}{e} - \frac{2(a+bx)^{5/2}}{5e(d+ex)^{5/2}} \right) \\
 \hline
 e - \frac{2(a+bx)^{7/2}(Bd - Ae)}{7e(d+ex)^{7/2}(bd - ae)}
 \end{array}$$

input `Int[((a + b*x)^(5/2)*(A + B*x))/(d + e*x)^(9/2),x]`

output

```
(-2*(B*d - A*e)*(a + b*x)^(7/2))/(7*e*(b*d - a*e)*(d + e*x)^(7/2)) + (B*((-2*(a + b*x)^(5/2))/(5*e*(d + e*x)^(5/2)) + (b*((-2*(a + b*x)^(3/2))/(3*e*(d + e*x)^(3/2)) + (b*((-2*Sqrt[a + b*x])/(e*Sqrt[d + e*x]) + (2*Sqrt[b]*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/e^(3/2)))/e))/e
```

Defintions of rubi rules used

rule 57

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]
```

rule 66

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]
```

rule 87

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))
```

rule 221

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1088 vs. $2(133) = 266$.

Time = 0.27 (sec) , antiderivative size = 1089, normalized size of antiderivative = 6.52

method	result	size
default	Expression too large to display	1089

input `int((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(9/2),x,method=_RETURNVERBOSE)`

output

```
-1/105*(b*x+a)^(1/2)*(12*B*a^3*d*e^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-1
05*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(
1/2))*a*b^3*e^5*x^4+105*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)
^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^4*d*e^4*x^4+420*B*ln(1/2*(2*b*e*x+2*((e*x+d)
)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^4*d^2*e^3*x^3+630*B*ln
(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))
*b^4*d^3*e^2*x^2+420*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/
2)+a*e+d*b)/(b*e)^(1/2))*b^4*d^4*e*x-105*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x
+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^3*d^4*e+568*B*a*b^2*d*e^3
*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+92*B*a^2*b*d*e^3*x*((e*x+d)*(b*x+
a))^(1/2)*(b*e)^(1/2)+476*B*a*b^2*d^2*e^2*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(
1/2)+30*A*a^3*e^4*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-210*B*b^3*d^4*((e*x
+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+30*A*b^3*e^4*x^3*((e*x+d)*(b*x+a))^(1/2)*(b
*e)^(1/2)+42*B*a^3*e^4*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+322*B*a*b^2*e
^4*x^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-352*B*b^3*d*e^3*x^3*((e*x+d)*(b
*x+a))^(1/2)*(b*e)^(1/2)+90*A*a*b^2*e^4*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(
1/2)+154*B*a^2*b*e^4*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-812*B*b^3*d^
2*e^2*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+90*A*a^2*b*e^4*x*((e*x+d)*(b
*x+a))^(1/2)*(b*e)^(1/2)-700*B*b^3*d^3*e*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(
1/2)+28*B*a^2*b*d^2*e^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+140*B*a*b^2...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 514 vs. $2(133) = 266$.

Time = 8.13 (sec) , antiderivative size = 1053, normalized size of antiderivative = 6.31

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{9/2}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(9/2),x, algorithm="fricas")`

output

```
[1/210*(105*(B*b^3*d^5 - B*a*b^2*d^4*e + (B*b^3*d*e^4 - B*a*b^2*e^5)*x^4 +
4*(B*b^3*d^2*e^3 - B*a*b^2*d*e^4)*x^3 + 6*(B*b^3*d^3*e^2 - B*a*b^2*d^2*e^
3)*x^2 + 4*(B*b^3*d^4*e - B*a*b^2*d^3*e^2)*x)*sqrt(b/e)*log(8*b^2*e^2*x^2
+ b^2*d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*b*e^2*x + b*d*e + a*e^2)*sqrt(b*x +
a)*sqrt(e*x + d)*sqrt(b/e) + 8*(b^2*d*e + a*b*e^2)*x) - 4*(105*B*b^3*d^4
- 70*B*a*b^2*d^3*e - 14*B*a^2*b*d^2*e^2 - 6*B*a^3*d*e^3 - 15*A*a^3*e^4 + (
176*B*b^3*d*e^3 - (161*B*a*b^2 + 15*A*b^3)*e^4)*x^3 + (406*B*b^3*d^2*e^2 -
284*B*a*b^2*d*e^3 - (77*B*a^2*b + 45*A*a*b^2)*e^4)*x^2 + (350*B*b^3*d^3*e
- 238*B*a*b^2*d^2*e^2 - 46*B*a^2*b*d*e^3 - 3*(7*B*a^3 + 15*A*a^2*b)*e^4)*
x)*sqrt(b*x + a)*sqrt(e*x + d))/(b*d^5*e^4 - a*d^4*e^5 + (b*d*e^8 - a*e^9)
*x^4 + 4*(b*d^2*e^7 - a*d*e^8)*x^3 + 6*(b*d^3*e^6 - a*d^2*e^7)*x^2 + 4*(b*
d^4*e^5 - a*d^3*e^6)*x), -1/105*(105*(B*b^3*d^5 - B*a*b^2*d^4*e + (B*b^3*d
*e^4 - B*a*b^2*e^5)*x^4 + 4*(B*b^3*d^2*e^3 - B*a*b^2*d*e^4)*x^3 + 6*(B*b^3
*d^3*e^2 - B*a*b^2*d^2*e^3)*x^2 + 4*(B*b^3*d^4*e - B*a*b^2*d^3*e^2)*x)*sqr
t(-b/e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(b*x + a)*sqrt(e*x + d)*sqrt(
-b/e)/(b^2*e*x^2 + a*b*d + (b^2*d + a*b*e)*x)) + 2*(105*B*b^3*d^4 - 70*B*a
*b^2*d^3*e - 14*B*a^2*b*d^2*e^2 - 6*B*a^3*d*e^3 - 15*A*a^3*e^4 + (176*B*b^
3*d*e^3 - (161*B*a*b^2 + 15*A*b^3)*e^4)*x^3 + (406*B*b^3*d^2*e^2 - 284*B*a
*b^2*d*e^3 - (77*B*a^2*b + 45*A*a*b^2)*e^4)*x^2 + (350*B*b^3*d^3*e - 238*B
*a*b^2*d^2*e^2 - 46*B*a^2*b*d*e^3 - 3*(7*B*a^3 + 15*A*a^2*b)*e^4)*x)*sq...
```


Sympy [F]

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{9/2}} dx = \int \frac{(A + Bx)(a + bx)^{5/2}}{(d + ex)^{9/2}} dx$$

input `integrate((b*x+a)**(5/2)*(B*x+A)/(e*x+d)**(9/2),x)`

output `Integral((A + B*x)*(a + b*x)**(5/2)/(d + e*x)**(9/2), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{9/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(9/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f or more de`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 678 vs. 2(133) = 266.

Time = 0.40 (sec) , antiderivative size = 678, normalized size of antiderivative = 4.06

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{9/2}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(9/2),x, algorithm="giac")`

output

```

-2*B*b^2*abs(b)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*
b*e - a*b*e)))/(sqrt(b*e)*e^4) - 2/105*(((b*x + a)*((176*B*b^10*d^3*e^6*ab
s(b) - 513*B*a*b^9*d^2*e^7*abs(b) - 15*A*b^10*d^2*e^7*abs(b) + 498*B*a^2*b
^8*d*e^8*abs(b) + 30*A*a*b^9*d*e^8*abs(b) - 161*B*a^3*b^7*e^9*abs(b) - 15*
A*a^2*b^8*e^9*abs(b))*(b*x + a)/(b^5*d^3*e^7 - 3*a*b^4*d^2*e^8 + 3*a^2*b^3
*d*e^9 - a^3*b^2*e^10) + 406*(B*b^11*d^4*e^5*abs(b) - 4*B*a*b^10*d^3*e^6*a
bs(b) + 6*B*a^2*b^9*d^2*e^7*abs(b) - 4*B*a^3*b^8*d*e^8*abs(b) + B*a^4*b^7*
e^9*abs(b))/(b^5*d^3*e^7 - 3*a*b^4*d^2*e^8 + 3*a^2*b^3*d*e^9 - a^3*b^2*e^1
0)) + 350*(B*b^12*d^5*e^4*abs(b) - 5*B*a*b^11*d^4*e^5*abs(b) + 10*B*a^2*b^
10*d^3*e^6*abs(b) - 10*B*a^3*b^9*d^2*e^7*abs(b) + 5*B*a^4*b^8*d*e^8*abs(b)
- B*a^5*b^7*e^9*abs(b))/(b^5*d^3*e^7 - 3*a*b^4*d^2*e^8 + 3*a^2*b^3*d*e^9
- a^3*b^2*e^10))*(b*x + a) + 105*(B*b^13*d^6*e^3*abs(b) - 6*B*a*b^12*d^5*e
^4*abs(b) + 15*B*a^2*b^11*d^4*e^5*abs(b) - 20*B*a^3*b^10*d^3*e^6*abs(b) +
15*B*a^4*b^9*d^2*e^7*abs(b) - 6*B*a^5*b^8*d*e^8*abs(b) + B*a^6*b^7*e^9*abs
(b))/(b^5*d^3*e^7 - 3*a*b^4*d^2*e^8 + 3*a^2*b^3*d*e^9 - a^3*b^2*e^10))*sqr
t(b*x + a)/(b^2*d + (b*x + a)*b*e - a*b*e)^(7/2)

```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{9/2}} dx = \int \frac{(A+Bx)(a+bx)^{5/2}}{(d+ex)^{9/2}} dx$$

input

```
int(((A + B*x)*(a + b*x)^(5/2))/(d + e*x)^(9/2), x)
```

output

```
int(((A + B*x)*(a + b*x)^(5/2))/(d + e*x)^(9/2), x)
```

Reduce [B] (verification not implemented)

Time = 94.66 (sec) , antiderivative size = 583, normalized size of antiderivative = 3.49

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{9/2}} dx = \frac{-\frac{2\sqrt{ex+d}\sqrt{bx+a}a^3e^4}{7} - \frac{2\sqrt{ex+d}\sqrt{bx+a}a^2bde^3}{5} - \frac{44\sqrt{ex+d}\sqrt{bx+a}a^2be^4x}{35} - \frac{2\sqrt{ex+d}\sqrt{bx+a}}{3}}$$

input

```
int((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(9/2), x)
```

output

```
(2*( - 15*sqrt(d + e*x)*sqrt(a + b*x)*a**3*e**4 - 21*sqrt(d + e*x)*sqrt(a
+ b*x)*a**2*b*d*e**3 - 66*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b*e**4*x - 35*s
qrt(d + e*x)*sqrt(a + b*x)*a*b**2*d**2*e**2 - 112*sqrt(d + e*x)*sqrt(a + b
*x)*a*b**2*d*e**3*x - 122*sqrt(d + e*x)*sqrt(a + b*x)*a*b**2*e**4*x**2 - 1
05*sqrt(d + e*x)*sqrt(a + b*x)*b**3*d**3*e - 350*sqrt(d + e*x)*sqrt(a + b*
x)*b**3*d**2*e**2*x - 406*sqrt(d + e*x)*sqrt(a + b*x)*b**3*d*e**3*x**2 - 1
76*sqrt(d + e*x)*sqrt(a + b*x)*b**3*e**4*x**3 + 105*sqrt(e)*sqrt(b)*log((s
qrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**3*d**4 +
420*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/s
qrt(a*e - b*d))*b**3*d**3*e*x + 630*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a +
b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**3*d**2*e**2*x**2 + 420*s
qrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*
e - b*d))*b**3*d*e**3*x**3 + 105*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x
) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**3*e**4*x**4 + 56*sqrt(e)*sq
rt(b)*b**3*d**4 + 224*sqrt(e)*sqrt(b)*b**3*d**3*e*x + 336*sqrt(e)*sqrt(b)*
b**3*d**2*e**2*x**2 + 224*sqrt(e)*sqrt(b)*b**3*d*e**3*x**3 + 56*sqrt(e)*sq
rt(b)*b**3*e**4*x**4)/(105*e**5*(d**4 + 4*d**3*e*x + 6*d**2*e**2*x**2 + 4
*d*e**3*x**3 + e**4*x**4))
```

3.201 $\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{11/2}} dx$

Optimal result	1839
Mathematica [A] (verified)	1839
Rubi [A] (verified)	1840
Maple [A] (verified)	1841
Fricas [B] (verification not implemented)	1842
Sympy [F]	1842
Maxima [F(-2)]	1843
Giac [B] (verification not implemented)	1843
Mupad [B] (verification not implemented)	1844
Reduce [B] (verification not implemented)	1844

Optimal result

Integrand size = 24, antiderivative size = 95

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{11/2}} dx = -\frac{2(Bd - Ae)(a+bx)^{7/2}}{9e(bd - ae)(d+ex)^{9/2}} + \frac{2(7bBd + 2Abe - 9aBe)(a+bx)^{7/2}}{63e(bd - ae)^2(d+ex)^{7/2}}$$

output `-2/9*(-A*e+B*d)*(b*x+a)^(7/2)/e/(-a*e+b*d)/(e*x+d)^(9/2)+2/63*(2*A*b*e-9*B*a*e+7*B*b*d)*(b*x+a)^(7/2)/e/(-a*e+b*d)^2/(e*x+d)^(7/2)`

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.69

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{11/2}} dx = \frac{2(a+bx)^{7/2}(B(-2ad + 7bdx - 9aex) + A(9bd - 7ae + 2bex))}{63(bd - ae)^2(d+ex)^{9/2}}$$

input `Integrate[((a + b*x)^(5/2)*(A + B*x))/(d + e*x)^(11/2),x]`

output

$$(2*(a + b*x)^{(7/2)}*(B*(-2*a*d + 7*b*d*x - 9*a*e*x) + A*(9*b*d - 7*a*e + 2*b*e*x)))/(63*(b*d - a*e)^2*(d + e*x)^{(9/2)})$$

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {87, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{11/2}} dx$$

$$\downarrow 87$$

$$\frac{(-9aBe + 2Abe + 7bBd) \int \frac{(a+bx)^{5/2}}{(d+ex)^{9/2}} dx}{9e(bd - ae)} - \frac{2(a + bx)^{7/2}(Bd - Ae)}{9e(d + ex)^{9/2}(bd - ae)}$$

$$\downarrow 48$$

$$\frac{2(a + bx)^{7/2}(-9aBe + 2Abe + 7bBd)}{63e(d + ex)^{7/2}(bd - ae)^2} - \frac{2(a + bx)^{7/2}(Bd - Ae)}{9e(d + ex)^{9/2}(bd - ae)}$$

input

$$\text{Int}[(a + b*x)^{(5/2)}*(A + B*x)/(d + e*x)^{(11/2)}, x]$$

output

$$(-2*(B*d - A*e)*(a + b*x)^{(7/2)})/(9*e*(b*d - a*e)*(d + e*x)^{(9/2)}) + (2*(7*b*B*d + 2*A*b*e - 9*a*B*e)*(a + b*x)^{(7/2)})/(63*e*(b*d - a*e)^2*(d + e*x)^{(7/2)})$$

Definitions of rubi rules used

rule 48

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[[a + b*x]^(m + 1)*[(c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))], x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

rule 87

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))
```

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.78

method	result
gospers	$-\frac{2(bx+a)^{\frac{7}{2}}(-2Abex+9Baeax-7Bbdx+7Aae-9Abd+2Bad)}{63(ex+d)^{\frac{9}{2}}(a^2e^2-2abde+b^2d^2)}$
orering	$-\frac{2(bx+a)^{\frac{7}{2}}(-2Abex+9Baeax-7Bbdx+7Aae-9Abd+2Bad)}{63(ex+d)^{\frac{9}{2}}(a^2e^2-2abde+b^2d^2)}$
default	$-\frac{2(-2A^3b^3ex^3+9Ba^2b^2ex^3-7Bb^3dx^3+3Aa^2b^2ex^2-9Ab^3dx^2+18Ba^2bex^2-12Ba^2bdx^2+12Aa^2bex-18Aa^2bdx+9Ba^3ex-3A^3b^3d)}{63(ex+d)^{\frac{9}{2}}(ae-db)^2}$

input

```
int((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(11/2),x,method=_RETURNVERBOSE)
```

output

```
-2/63*(b*x+a)^(7/2)*(-2*A*b*e*x+9*B*a*e*x-7*B*b*d*x+7*A*a*e-9*A*b*d+2*B*a*
d)/(e*x+d)^(9/2)/(a^2*e^2-2*a*b*d*e+b^2*d^2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 391 vs. $2(83) = 166$.

Time = 28.89 (sec) , antiderivative size = 391, normalized size of antiderivative = 4.12

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{11/2}} dx = \frac{2(7Aa^4e - (7Bb^4d - (9Bab^3 - 2Ab^4)e)x^4 - ((19Bab^3 + 9Ab^4)d - (27Ba^2b^2 + Aab^3)e)x^3 - 3((5Ba^3b^3)d - (9Bba^3b + 5Aa^2b^2)e)x^2 + (2Ba^4 - 9Aa^3b)d - ((Bba^3b + 27Aa^2b^2)d - (9Ba^4 + 19Aa^3b)e)x)\sqrt{bx+a}\sqrt{ex+d}}{63(b^2d^7 - 2abd^6e + a^2d^5e^2 + (b^2d^2e^5 - 2abde^6 + a^2e^7)x^5 + 5(b^2d^3e^4 - 2abd^2e^5 + a^2de^6)x^4 + 10(b^2d^4e^3 - 2a^2bd^3e^4 + a^2d^2e^5)x^3 + 10(b^2d^5e^2 - 2a^2bd^4e^3 + a^2d^3e^4)x^2 + 5(b^2d^6e - 2a^2bd^5e^2 + a^2d^4e^3)x)}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(11/2),x, algorithm="fricas")`

output `-2/63*(7*A*a^4*e - (7*B*b^4*d - (9*B*a*b^3 - 2*A*b^4)*e)*x^4 - ((19*B*a*b^3 + 9*A*b^4)*d - (27*B*a^2*b^2 + A*a*b^3)*e)*x^3 - 3*((5*B*a^2*b^2 + 9*A*a*b^3)*d - (9*B*a^3*b + 5*A*a^2*b^2)*e)*x^2 + (2*B*a^4 - 9*A*a^3*b)*d - ((B*a^3*b + 27*A*a^2*b^2)*d - (9*B*a^4 + 19*A*a^3*b)*e)*x)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*d^7 - 2*a*b*d^6*e + a^2*d^5*e^2 + (b^2*d^2*e^5 - 2*a*b*d*e^6 + a^2*e^7)*x^5 + 5*(b^2*d^3*e^4 - 2*a*b*d^2*e^5 + a^2*d*e^6)*x^4 + 10*(b^2*d^4*e^3 - 2*a*b*d^3*e^4 + a^2*d^2*e^5)*x^3 + 10*(b^2*d^5*e^2 - 2*a*b*d^4*e^3 + a^2*d^3*e^4)*x^2 + 5*(b^2*d^6*e - 2*a*b*d^5*e^2 + a^2*d^4*e^3)*x)`

Sympy [F]

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{11/2}} dx = \int \frac{(A+Bx)(a+bx)^{\frac{5}{2}}}{(d+ex)^{\frac{11}{2}}} dx$$

input `integrate((b*x+a)**(5/2)*(B*x+A)/(e*x+d)**(11/2),x)`

output `Integral((A + B*x)*(a + b*x)**(5/2)/(d + e*x)**(11/2), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{11/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(11/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f or more de

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 379 vs. 2(83) = 166.

Time = 0.56 (sec) , antiderivative size = 379, normalized size of antiderivative = 3.99

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{11/2}} dx = \frac{2(bx+a)^{7/2} \left(\frac{(7Bb^{12}d^3e^4|b| - 23Bab^{11}d^2e^5|b| + 2Ab^{12}d^2e^5|b| + 25Ba^2b^{10}de^6|b| - 4Aab^{11}de^6|b| - 9Ba^2b^9e^7|b| + 2A^2a^2b^{10}e^7|b|)}{b^6d^4e^4 - 4ab^5d^3e^5 + 6a^2b^4d^2e^6 - 4a^3b^3de^7 + a^4b^2e^8} \right)}{(d+ex)^{11/2}}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(11/2),x, algorithm="giac")`

output
$$\frac{2/63*(b*x + a)^{(7/2)*((7*B*b^{12}*d^3*e^4*abs(b) - 23*B*a*b^{11}*d^2*e^5*abs(b) + 2*A*b^{12}*d^2*e^5*abs(b) + 25*B*a^2*b^{10}*d*e^6*abs(b) - 4*A*a*b^{11}*d*e^6*abs(b) - 9*B*a^3*b^9*e^7*abs(b) + 2*A*a^2*b^{10}*e^7*abs(b)))*(b*x + a)/(b^6*d^4*e^4 - 4*a*b^5*d^3*e^5 + 6*a^2*b^4*d^2*e^6 - 4*a^3*b^3*d*e^7 + a^4*b^2*e^8) - 9*(B*a*b^{12}*d^3*e^4*abs(b) - A*b^{13}*d^3*e^4*abs(b) - 3*B*a^2*b^{11}*d^2*e^5*abs(b) + 3*A*a*b^{12}*d^2*e^5*abs(b) + 3*B*a^3*b^{10}*d*e^6*abs(b) - 3*A*a^2*b^{11}*d*e^6*abs(b) - B*a^4*b^9*e^7*abs(b) + A*a^3*b^{10}*e^7*abs(b))}{(b^6*d^4*e^4 - 4*a*b^5*d^3*e^5 + 6*a^2*b^4*d^2*e^6 - 4*a^3*b^3*d*e^7 + a^4*b^2*e^8)}/(b^2*d + (b*x + a)*b*e - a*b*e)^{(9/2)}$$

Mupad [B] (verification not implemented)

Time = 1.70 (sec) , antiderivative size = 325, normalized size of antiderivative = 3.42

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{11/2}} dx = \frac{\sqrt{d+ex} \left(\frac{x^3 \sqrt{a+bx} (18Ab^4d - 2Aab^3e + 38Bab^3d - 54Ba^2b^2e)}{63e^5(ae-bd)^2} - \frac{\sqrt{a+bx} (14Aa^4e + 4Ba^4)}{63e^5(ae-bd)} \right)}{(d+ex)^{11/2}}$$

input `int(((A + B*x)*(a + b*x)^(5/2))/(d + e*x)^(11/2),x)`output
$$\begin{aligned} & ((d+e*x)^{(1/2)}*((x^3*(a+b*x)^{(1/2)}*(18*A*b^4*d - 2*A*a*b^3*e + 38*B*a*b^3*d - 54*B*a^2*b^2*e))/(63*e^5*(a*e - b*d)^2) - ((a+b*x)^{(1/2)}*(14*A*a^4*e + 4*B*a^4*d - 18*A*a^3*b*d))/(63*e^5*(a*e - b*d)^2) + (x^4*(a+b*x)^{(1/2)}*(4*A*b^4*e + 14*B*b^4*d - 18*B*a*b^3*e))/(63*e^5*(a*e - b*d)^2) - (x*(a+b*x)^{(1/2)}*(18*B*a^4*e + 38*A*a^3*b*e - 2*B*a^3*b*d - 54*A*a^2*b^2*d))/(63*e^5*(a*e - b*d)^2) + (2*a*b*x^2*(a+b*x)^{(1/2)}*(9*A*b^2*d - 9*B*a^2*e - 5*A*a*b*e + 5*B*a*b*d))/(21*e^5*(a*e - b*d)^2))/((x^5 + d^5/e^5 + (5*d*x^4)/e + (5*d^4*x)/e^4 + (10*d^2*x^3)/e^2 + (10*d^3*x^2)/e^3) \end{aligned}$$
Reduce [B] (verification not implemented)

Time = 2.29 (sec) , antiderivative size = 335, normalized size of antiderivative = 3.53

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{11/2}} dx = \frac{-\frac{2\sqrt{ex+d}\sqrt{bx+a}a^4e^5}{9} - \frac{8\sqrt{ex+d}\sqrt{bx+a}a^3be^5x}{9} - \frac{4\sqrt{ex+d}\sqrt{bx+a}a^2b^2e^5x^2}{3} - \frac{8\sqrt{ex+d}\sqrt{bx+a}}{9}}{e^5(ae^6x^5 - bde^5x^5 + 5ade^5x^4 - 5bd^2e^4x^4 + \dots)}$$

input `int((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(11/2),x)`

output

```
(2*( - sqrt(d + e*x)*sqrt(a + b*x)*a**4*e**5 - 4*sqrt(d + e*x)*sqrt(a + b*
x)*a**3*b*e**5*x - 6*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**2*e**5*x**2 - 4*s
qrt(d + e*x)*sqrt(a + b*x)*a*b**3*e**5*x**3 - sqrt(d + e*x)*sqrt(a + b*x)*
b**4*e**5*x**4 - sqrt(e)*sqrt(b)*b**4*d**5 - 5*sqrt(e)*sqrt(b)*b**4*d**4*e
*x - 10*sqrt(e)*sqrt(b)*b**4*d**3*e**2*x**2 - 10*sqrt(e)*sqrt(b)*b**4*d**2
*e**3*x**3 - 5*sqrt(e)*sqrt(b)*b**4*d*e**4*x**4 - sqrt(e)*sqrt(b)*b**4*e**
5*x**5))/(9*e**5*(a*d**5*e + 5*a*d**4*e**2*x + 10*a*d**3*e**3*x**2 + 10*a*
d**2*e**4*x**3 + 5*a*d*e**5*x**4 + a*e**6*x**5 - b*d**6 - 5*b*d**5*e*x - 1
0*b*d**4*e**2*x**2 - 10*b*d**3*e**3*x**3 - 5*b*d**2*e**4*x**4 - b*d*e**5*x
**5))
```

3.202 $\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{13/2}} dx$

Optimal result	1846
Mathematica [A] (verified)	1846
Rubi [A] (verified)	1847
Maple [A] (verified)	1849
Fricas [B] (verification not implemented)	1849
Sympy [F(-1)]	1850
Maxima [F(-2)]	1850
Giac [B] (verification not implemented)	1851
Mupad [B] (verification not implemented)	1852
Reduce [B] (verification not implemented)	1852

Optimal result

Integrand size = 24, antiderivative size = 147

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{13/2}} dx = -\frac{2(Bd - Ae)(a+bx)^{7/2}}{11e(bd - ae)(d+ex)^{11/2}} + \frac{2(7bBd + 4Abe - 11aBe)(a+bx)^{7/2}}{99e(bd - ae)^2(d+ex)^{9/2}} + \frac{4b(7bBd + 4Abe - 11aBe)(a+bx)^{7/2}}{693e(bd - ae)^3(d+ex)^{7/2}}$$

output `-2/11*(-A*e+B*d)*(b*x+a)^(7/2)/e/(-a*e+b*d)/(e*x+d)^(11/2)+2/99*(4*A*b*e-11*B*a*e+7*B*b*d)*(b*x+a)^(7/2)/e/(-a*e+b*d)^2/(e*x+d)^(9/2)+4/693*b*(4*A*b*e-11*B*a*e+7*B*b*d)*(b*x+a)^(7/2)/e/(-a*e+b*d)^3/(e*x+d)^(7/2)`

Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.91

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{13/2}} dx = \frac{2(a+bx)^{11/2} \left(-63Bde + 63Ae^2 + \frac{77bBd(d+ex)}{a+bx} - \frac{154Abe(d+ex)}{a+bx} + \frac{77aBe(d+ex)}{a+bx} \right)}{693(bd - ae)^3(d+ex)^{11/2}}$$

input `Integrate[((a + b*x)^(5/2)*(A + B*x))/(d + e*x)^(13/2),x]`

output

```
(2*(a + b*x)^(11/2)*(-63*B*d*e + 63*A*e^2 + (77*b*B*d*(d + e*x))/(a + b*x)
- (154*A*b*e*(d + e*x))/(a + b*x) + (77*a*B*e*(d + e*x))/(a + b*x) + (99*
A*b^2*(d + e*x)^2)/(a + b*x)^2 - (99*a*b*B*(d + e*x)^2)/(a + b*x)^2))/(693
*(b*d - a*e)^3*(d + e*x)^(11/2))
```

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.97, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {87, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{13/2}} dx$$

$$\downarrow 87$$

$$\frac{(-11aBe + 4Abe + 7bBd) \int \frac{(a+bx)^{5/2}}{(d+ex)^{11/2}} dx}{11e(bd - ae)} - \frac{2(a + bx)^{7/2}(Bd - Ae)}{11e(d + ex)^{11/2}(bd - ae)}$$

$$\downarrow 55$$

$$\frac{(-11aBe + 4Abe + 7bBd) \left(\frac{2b \int \frac{(a+bx)^{5/2}}{(d+ex)^{9/2}} dx}{9(bd - ae)} + \frac{2(a+bx)^{7/2}}{9(d+ex)^{9/2}(bd - ae)} \right)}{11e(bd - ae)} - \frac{2(a + bx)^{7/2}(Bd - Ae)}{11e(d + ex)^{11/2}(bd - ae)}$$

$$\downarrow 48$$

$$\frac{\left(\frac{4b(a+bx)^{7/2}}{63(d+ex)^{7/2}(bd - ae)^2} + \frac{2(a+bx)^{7/2}}{9(d+ex)^{9/2}(bd - ae)} \right) (-11aBe + 4Abe + 7bBd)}{11e(bd - ae)} - \frac{2(a + bx)^{7/2}(Bd - Ae)}{11e(d + ex)^{11/2}(bd - ae)}$$

input

```
Int[((a + b*x)^(5/2)*(A + B*x))/(d + e*x)^(13/2), x]
```

output

```
(-2*(B*d - A*e)*(a + b*x)^(7/2))/(11*e*(b*d - a*e)*(d + e*x)^(11/2)) + ((7
*b*B*d + 4*A*b*e - 11*a*B*e)*((2*(a + b*x)^(7/2))/(9*(b*d - a*e)*(d + e*x)
^(9/2)) + (4*b*(a + b*x)^(7/2))/(63*(b*d - a*e)^2*(d + e*x)^(7/2)))/(11*e
*(b*d - a*e))
```

Defintions of rubi rules used

rule 48

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

rule 55

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
lerQ[n, 1])
```

rule 87

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && ( !LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.20

method	result
gospers	$-\frac{2(bx+a)^{\frac{7}{2}}(8A^2b^2e^2x^2-22Bab^2e^2x^2+14B^2b^2de^2x^2-28Aab^2e^2x+44A^2b^2dex+77B^2a^2e^2x-170Babdex+77b^2B^2d^2x+63a^2Ae^2-154A^2a^2b^2e^2+99A^2b^2d^2+14B^2a^2d^2e-22B^2a^2b^2d^2)}{693(ex+d)^{\frac{11}{2}}(a^3e^3-3a^2be^2d+3ab^2d^2e-b^3d^3)}$
orering	$-\frac{2(bx+a)^{\frac{7}{2}}(8A^2b^2e^2x^2-22Bab^2e^2x^2+14B^2b^2de^2x^2-28Aab^2e^2x+44A^2b^2dex+77B^2a^2e^2x-170Babdex+77b^2B^2d^2x+63a^2Ae^2-154A^2a^2b^2e^2+99A^2b^2d^2+14B^2a^2d^2e-22B^2a^2b^2d^2)}{693(ex+d)^{\frac{11}{2}}(a^3e^3-3a^2be^2d+3ab^2d^2e-b^3d^3)}$
default	$-\frac{2(8A^4b^4e^2x^4-22B^2a^2b^3e^2x^4+14B^2b^4de^2x^4-12A^2a^2b^3e^2x^3+44A^2b^4de^2x^3+33B^2a^2b^2e^2x^3-142B^2a^2b^3de^2x^3+77B^2b^4d^2x^3+15A^2a^2b^2e^2x^2-22B^2a^2b^3de^2x^2+14B^2b^4d^2e^2x^2-28A^2a^2b^3e^2x+44A^2b^4dex+77B^2a^2e^2x-170B^2abdex+77b^2B^2d^2x+63A^2a^2e^2-154A^2a^2b^2e^2+99A^2b^2d^2+14B^2a^2d^2e-22B^2a^2b^2d^2)}{693(ex+d)^{\frac{11}{2}}(a^3e^3-3a^2be^2d+3ab^2d^2e-b^3d^3)}$

input `int((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(13/2),x,method=_RETURNVERBOSE)`

output
$$-\frac{2}{693}(b*x+a)^{\frac{7}{2}}*(8*A*b^2*e^2*x^2-22*B*a*b*e^2*x^2+14*B*b^2*d*e*x^2-28*A*a*b*e^2*x+44*A*b^2*d*e*x+77*B*a^2*e^2*x-170*B*a*b*d*e*x+77*B*b^2*d^2*x+63*A*a^2*e^2-154*A*a*b*d*e+99*A*b^2*d^2+14*B*a^2*d^2e-22*B*a*b*d^2)/(e*x+d)^{\frac{11}{2}}/(a^3*e^3-3*a^2*b*d*e^2+3*a*b^2*d^2e-b^3*d^3)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 693 vs. 2(129) = 258.

Time = 66.66 (sec) , antiderivative size = 693, normalized size of antiderivative = 4.71

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{13/2}} dx = \frac{2(63Aa^5e^2+2(7Bb^5de-(11Bab^4-4Ab^5)e^2)x^5+(77Bb^5d^2-4(32Bab^4d-3Aa^2b^2d^2e-b^3d^3)))}{693(b^3d^9-3a^2b^2d^8+3abd^7-3a^2b^2d^6+3a^2bd^5-3a^3d^4)}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(13/2),x, algorithm="fricas")`

output

```
2/693*(63*A*a^5*e^2 + 2*(7*B*b^5*d*e - (11*B*a*b^4 - 4*A*b^5)*e^2)*x^5 + (
77*B*b^5*d^2 - 4*(32*B*a*b^4 - 11*A*b^5)*d*e + (11*B*a^2*b^3 - 4*A*a*b^4)*
e^2)*x^4 + (11*(19*B*a*b^4 + 9*A*b^5)*d^2 - 2*(227*B*a^2*b^3 + 11*A*a*b^4)
*d*e + 3*(55*B*a^3*b^2 + A*a^2*b^3)*e^2)*x^3 - 11*(2*B*a^4*b - 9*A*a^3*b^2
)*d^2 + 14*(B*a^5 - 11*A*a^4*b)*d*e + (33*(5*B*a^2*b^3 + 9*A*a*b^4)*d^2 -
2*(227*B*a^3*b^2 + 165*A*a^2*b^3)*d*e + (209*B*a^4*b + 113*A*a^3*b^2)*e^2)
*x^2 + (11*(B*a^3*b^2 + 27*A*a^2*b^3)*d^2 - 2*(64*B*a^4*b + 209*A*a^3*b^2)
*d*e + 7*(11*B*a^5 + 23*A*a^4*b)*e^2)*x)*sqrt(b*x + a)*sqrt(e*x + d)/(b^3*
d^9 - 3*a*b^2*d^8*e + 3*a^2*b*d^7*e^2 - a^3*d^6*e^3 + (b^3*d^3*e^6 - 3*a*b
^2*d^2*e^7 + 3*a^2*b*d*e^8 - a^3*e^9)*x^6 + 6*(b^3*d^4*e^5 - 3*a*b^2*d^3*e
^6 + 3*a^2*b*d^2*e^7 - a^3*d*e^8)*x^5 + 15*(b^3*d^5*e^4 - 3*a*b^2*d^4*e^5
+ 3*a^2*b*d^3*e^6 - a^3*d^2*e^7)*x^4 + 20*(b^3*d^6*e^3 - 3*a*b^2*d^5*e^4 +
3*a^2*b*d^4*e^5 - a^3*d^3*e^6)*x^3 + 15*(b^3*d^7*e^2 - 3*a*b^2*d^6*e^3 +
3*a^2*b*d^5*e^4 - a^3*d^4*e^5)*x^2 + 6*(b^3*d^8*e - 3*a*b^2*d^7*e^2 + 3*a
^2*b*d^6*e^3 - a^3*d^5*e^4)*x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{13/2}} dx = \text{Timed out}$$

input

```
integrate((b*x+a)**(5/2)*(B*x+A)/(e*x+d)**(13/2),x)
```

output

Timed out

Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{13/2}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(13/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f
or more de
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 663 vs. $2(129) = 258$.

Time = 0.57 (sec) , antiderivative size = 663, normalized size of antiderivative = 4.51

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{13/2}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(13/2),x, algorithm="giac")
```

output

```
2/693*((b*x + a)*(2*(7*B*b^14*d^3*e^6*abs(b) - 25*B*a*b^13*d^2*e^7*abs(b)
+ 4*A*b^14*d^2*e^7*abs(b) + 29*B*a^2*b^12*d*e^8*abs(b) - 8*A*a*b^13*d*e^8*
abs(b) - 11*B*a^3*b^11*e^9*abs(b) + 4*A*a^2*b^12*e^9*abs(b))*(b*x + a)/(b^
7*d^5*e^5 - 5*a*b^6*d^4*e^6 + 10*a^2*b^5*d^3*e^7 - 10*a^3*b^4*d^2*e^8 + 5*
a^4*b^3*d*e^9 - a^5*b^2*e^10) + 11*(7*B*b^15*d^4*e^5*abs(b) - 32*B*a*b^14*
d^3*e^6*abs(b) + 4*A*b^15*d^3*e^6*abs(b) + 54*B*a^2*b^13*d^2*e^7*abs(b) -
12*A*a*b^14*d^2*e^7*abs(b) - 40*B*a^3*b^12*d*e^8*abs(b) + 12*A*a^2*b^13*d*
e^8*abs(b) + 11*B*a^4*b^11*e^9*abs(b) - 4*A*a^3*b^12*e^9*abs(b))/(b^7*d^5*
e^5 - 5*a*b^6*d^4*e^6 + 10*a^2*b^5*d^3*e^7 - 10*a^3*b^4*d^2*e^8 + 5*a^4*b^
3*d*e^9 - a^5*b^2*e^10)) - 99*(B*a*b^15*d^4*e^5*abs(b) - A*b^16*d^4*e^5*ab
s(b) - 4*B*a^2*b^14*d^3*e^6*abs(b) + 4*A*a*b^15*d^3*e^6*abs(b) + 6*B*a^3*b
^13*d^2*e^7*abs(b) - 6*A*a^2*b^14*d^2*e^7*abs(b) - 4*B*a^4*b^12*d*e^8*abs(
b) + 4*A*a^3*b^13*d*e^8*abs(b) + B*a^5*b^11*e^9*abs(b) - A*a^4*b^12*e^9*ab
s(b))/(b^7*d^5*e^5 - 5*a*b^6*d^4*e^6 + 10*a^2*b^5*d^3*e^7 - 10*a^3*b^4*d^2
*e^8 + 5*a^4*b^3*d*e^9 - a^5*b^2*e^10))*(b*x + a)^(7/2)/(b^2*d + (b*x + a)
*b*e - a*b*e)^(11/2)
```


Mupad [B] (verification not implemented)

Time = 1.91 (sec) , antiderivative size = 509, normalized size of antiderivative = 3.46

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{13/2}} dx = \frac{\sqrt{d+ex} \left(\frac{\sqrt{a+bx}(28Ba^5de+126Aa^5e^2-44Ba^4bd^2-308Aa^4bde+198Aa^3b^2d^2)}{693e^6(ae-bd)^3} + \frac{x\sqrt{a+bx}(154Ba^5e^2-256Ba^4bde+322Aa^4b^2d^2)}{693e^6(ae-bd)^3} \right)}{1}$$

input `int(((A + B*x)*(a + b*x)^(5/2))/(d + e*x)^(13/2),x)`output
$$\begin{aligned} & -((d + e*x)^{(1/2)} * (((a + b*x)^{(1/2)} * (126*A*a^5*e^2 + 28*B*a^5*d*e - 44*B*a^4*b*d^2 + 198*A*a^3*b^2*d^2 - 308*A*a^4*b*d*e)) / (693*e^6*(a*e - b*d)^3) + \\ & (x*(a + b*x)^{(1/2)} * (154*B*a^5*e^2 + 322*A*a^4*b*e^2 + 594*A*a^2*b^3*d^2 + 22*B*a^3*b^2*d^2 - 256*B*a^4*b*d*e - 836*A*a^3*b^2*d*e)) / (693*e^6*(a*e - b*d)^3) + \\ & (x^2*(a + b*x)^{(1/2)} * (594*A*a*b^4*d^2 + 418*B*a^4*b*e^2 + 226*A*a^3*b^2*e^2 + 330*B*a^2*b^3*d^2 - 660*A*a^2*b^3*d*e - 908*B*a^3*b^2*d*e)) / \\ & (693*e^6*(a*e - b*d)^3) + (x^3*(a + b*x)^{(1/2)} * (198*A*b^5*d^2 + 418*B*a*b^4*d^2 + 6*A*a^2*b^3*e^2 + 330*B*a^3*b^2*e^2 - 44*A*a*b^4*d*e - 908*B*a^2*b^3*d*e)) / \\ & (693*e^6*(a*e - b*d)^3) + (4*b^4*x^5*(a + b*x)^{(1/2)} * (4*A*b*e - 11*B*a*e + 7*B*b*d)) / (693*e^5*(a*e - b*d)^3) - (2*b^3*x^4*(a*e - 11*b*d)*(a + b*x)^{(1/2)} * (4*A*b*e - 11*B*a*e + 7*B*b*d)) / (693*e^6*(a*e - b*d)^3)) / (x^6 + d^6/e^6 + (6*d*x^5)/e + (6*d^5*x)/e^5 + (15*d^2*x^4)/e^2 + (20*d^3*x^3)/e^3 + (15*d^4*x^2)/e^4) \end{aligned}$$
Reduce [B] (verification not implemented)

Time = 0.57 (sec) , antiderivative size = 638, normalized size of antiderivative = 4.34

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{13/2}} dx = \frac{-\frac{2\sqrt{ex+d}\sqrt{bx+a}a^5e^6}{11} + \frac{2\sqrt{ex+d}\sqrt{bx+a}a^4bde^5}{9} - \frac{68\sqrt{ex+d}\sqrt{bx+a}a^4be^6x}{99} + \frac{8\sqrt{ex+d}\sqrt{bx+a}}{9}}{e^5(a^2e^8x^6 - 2abde^7x^6 + b^2d^2e^6x^6)}$$

input `int((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(13/2),x)`

output

```
(2*( - 9*sqrt(d + e*x)*sqrt(a + b*x)*a**5*e**6 + 11*sqrt(d + e*x)*sqrt(a +
b*x)*a**4*b*d*e**5 - 34*sqrt(d + e*x)*sqrt(a + b*x)*a**4*b*e**6*x + 44*sq
rt(d + e*x)*sqrt(a + b*x)*a**3*b**2*d*e**5*x - 46*sqrt(d + e*x)*sqrt(a + b
*x)*a**3*b**2*e**6*x**2 + 66*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**3*d*e**5*
x**2 - 24*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**3*e**6*x**3 + 44*sqrt(d + e*
x)*sqrt(a + b*x)*a*b**4*d*e**5*x**3 - sqrt(d + e*x)*sqrt(a + b*x)*a*b**4*e
**6*x**4 + 11*sqrt(d + e*x)*sqrt(a + b*x)*b**5*d*e**5*x**4 + 2*sqrt(d + e*
x)*sqrt(a + b*x)*b**5*e**6*x**5 - 2*sqrt(e)*sqrt(b)*b**5*d**6 - 12*sqrt(e)
*sqrt(b)*b**5*d**5*e*x - 30*sqrt(e)*sqrt(b)*b**5*d**4*e**2*x**2 - 40*sqrt(
e)*sqrt(b)*b**5*d**3*e**3*x**3 - 30*sqrt(e)*sqrt(b)*b**5*d**2*e**4*x**4 -
12*sqrt(e)*sqrt(b)*b**5*d*e**5*x**5 - 2*sqrt(e)*sqrt(b)*b**5*e**6*x**6))/(
99*e**5*(a**2*d**6*e**2 + 6*a**2*d**5*e**3*x + 15*a**2*d**4*e**4*x**2 + 20
*a**2*d**3*e**5*x**3 + 15*a**2*d**2*e**6*x**4 + 6*a**2*d*e**7*x**5 + a**2*
e**8*x**6 - 2*a*b*d**7*e - 12*a*b*d**6*e**2*x - 30*a*b*d**5*e**3*x**2 - 40
*a*b*d**4*e**4*x**3 - 30*a*b*d**3*e**5*x**4 - 12*a*b*d**2*e**6*x**5 - 2*a*
b*d*e**7*x**6 + b**2*d**8 + 6*b**2*d**7*e*x + 15*b**2*d**6*e**2*x**2 + 20*
b**2*d**5*e**3*x**3 + 15*b**2*d**4*e**4*x**4 + 6*b**2*d**3*e**5*x**5 + b**
2*d**2*e**6*x**6))
```

3.203 $\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{15/2}} dx$

Optimal result	1854
Mathematica [A] (verified)	1855
Rubi [A] (verified)	1855
Maple [A] (verified)	1857
Fricas [B] (verification not implemented)	1858
Sympy [F(-1)]	1859
Maxima [F(-2)]	1859
Giac [B] (verification not implemented)	1859
Mupad [B] (verification not implemented)	1860
Reduce [B] (verification not implemented)	1861

Optimal result

Integrand size = 24, antiderivative size = 201

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{15/2}} dx = -\frac{2(Bd - Ae)(a+bx)^{7/2}}{13e(bd - ae)(d+ex)^{13/2}} + \frac{2(7bBd + 6Abe - 13aBe)(a+bx)^{7/2}}{143e(bd - ae)^2(d+ex)^{11/2}} + \frac{8b(7bBd + 6Abe - 13aBe)(a+bx)^{7/2}}{1287e(bd - ae)^3(d+ex)^{9/2}} + \frac{16b^2(7bBd + 6Abe - 13aBe)(a+bx)^{7/2}}{9009e(bd - ae)^4(d+ex)^{7/2}}$$

output

```
-2/13*(-A*e+B*d)*(b*x+a)^(7/2)/e/(-a*e+b*d)/(e*x+d)^(13/2)+2/143*(6*A*b*e-13*B*a*e+7*B*b*d)*(b*x+a)^(7/2)/e/(-a*e+b*d)^2/(e*x+d)^(11/2)+8/1287*b*(6*A*b*e-13*B*a*e+7*B*b*d)*(b*x+a)^(7/2)/e/(-a*e+b*d)^3/(e*x+d)^(9/2)+16/9009*b^2*(6*A*b*e-13*B*a*e+7*B*b*d)*(b*x+a)^(7/2)/e/(-a*e+b*d)^4/(e*x+d)^(7/2)
```

Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 199, normalized size of antiderivative = 0.99

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{15/2}} dx = \frac{2(a+bx)^{7/2} (693Bde^2(a+bx)^3 - 693Ae^3(a+bx)^3 - 1638bBde(a+bx)^2(d+ex) + 2457A*b*e^2(a+bx)^2(d+ex) - 819*a*B*e^2(a+bx)^2(d+ex) + 1001*b^2*B*d*(a+bx)*(d+ex)^2 - 3003*A*b^2*e*(a+bx)*(d+ex)^2 + 2002*a*b*B*e*(a+bx)*(d+ex)^2 + 1287*A*b^3*(d+ex)^3 - 1287*a*b^2*B*(d+ex)^3)}{(9009*(b*d - a*e)^4*(d+ex)^{13/2}}$$

input

```
Integrate[((a + b*x)^(5/2)*(A + B*x))/(d + e*x)^(15/2),x]
```

output

```
(2*(a + b*x)^(7/2)*(693*B*d*e^2*(a + b*x)^3 - 693*A*e^3*(a + b*x)^3 - 1638
*b*B*d*e*(a + b*x)^2*(d + e*x) + 2457*A*b*e^2*(a + b*x)^2*(d + e*x) - 819*
a*B*e^2*(a + b*x)^2*(d + e*x) + 1001*b^2*B*d*(a + b*x)*(d + e*x)^2 - 3003*
A*b^2*e*(a + b*x)*(d + e*x)^2 + 2002*a*b*B*e*(a + b*x)*(d + e*x)^2 + 1287*
A*b^3*(d + e*x)^3 - 1287*a*b^2*B*(d + e*x)^3))/(9009*(b*d - a*e)^4*(d + e*
x)^(13/2))
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.95, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {87, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{15/2}} dx$$

$$\downarrow 87$$

$$\frac{(-13aBe + 6Abe + 7bBd) \int \frac{(a+bx)^{5/2}}{(d+ex)^{13/2}} dx}{13e(bd - ae)} - \frac{2(a+bx)^{7/2}(Bd - Ae)}{13e(d+ex)^{13/2}(bd - ae)}$$

$$\downarrow 55$$

$$\frac{(-13aBe + 6Abe + 7bBd) \left(\frac{4b \int \frac{(a+bx)^{5/2}}{(d+ex)^{11/2}} dx}{11(bd - ae)} + \frac{2(a+bx)^{7/2}}{11(d+ex)^{11/2}(bd - ae)} \right)}{13e(bd - ae)} - \frac{2(a+bx)^{7/2}(Bd - Ae)}{13e(d+ex)^{13/2}(bd - ae)}$$

$$\begin{array}{c}
 \downarrow 55 \\
 (-13aBe + 6Abe + 7bBd) \left(\frac{4b \left(\frac{2b \int \frac{(a+bx)^{5/2}}{(d+ex)^{9/2}} dx}{9(bd-ae)} + \frac{2(a+bx)^{7/2}}{9(d+ex)^{9/2}(bd-ae)} \right)}{11(bd-ae)} + \frac{2(a+bx)^{7/2}}{11(d+ex)^{11/2}(bd-ae)} \right) \\
 \hline
 \frac{13e(bd-ae)}{2(a+bx)^{7/2}(Bd-Ae)} \\
 \frac{2(a+bx)^{7/2}(Bd-Ae)}{13e(d+ex)^{13/2}(bd-ae)} \\
 \downarrow 48 \\
 \left(\frac{2(a+bx)^{7/2}}{11(d+ex)^{11/2}(bd-ae)} + \frac{4b \left(\frac{4b(a+bx)^{7/2}}{63(d+ex)^{7/2}(bd-ae)^2} + \frac{2(a+bx)^{7/2}}{9(d+ex)^{9/2}(bd-ae)} \right)}{11(bd-ae)} \right) (-13aBe + 6Abe + 7bBd) \\
 \hline
 \frac{13e(bd-ae)}{2(a+bx)^{7/2}(Bd-Ae)} \\
 \frac{2(a+bx)^{7/2}(Bd-Ae)}{13e(d+ex)^{13/2}(bd-ae)}
 \end{array}$$

input `Int[((a + b*x)^(5/2)*(A + B*x))/(d + e*x)^(15/2),x]`

output `(-2*(B*d - A*e)*(a + b*x)^(7/2))/(13*e*(b*d - a*e)*(d + e*x)^(13/2)) + ((7*b*B*d + 6*A*b*e - 13*a*B*e)*((2*(a + b*x)^(7/2))/(11*(b*d - a*e)*(d + e*x)^(11/2)) + (4*b*((2*(a + b*x)^(7/2))/(9*(b*d - a*e)*(d + e*x)^(9/2)) + (4*b*(a + b*x)^(7/2))/(63*(b*d - a*e)^2*(d + e*x)^(7/2)))/(11*(b*d - a*e)))/(13*e*(b*d - a*e))`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1047 vs. $2(177) = 354$.

Time = 138.52 (sec) , antiderivative size = 1047, normalized size of antiderivative = 5.21

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{15/2}} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(15/2),x, algorithm="fricas")
```

output

```
-2/9009*(693*A*a^6*e^3 - 8*(7*B*b^6*d*e^2 - (13*B*a*b^5 - 6*A*b^6)*e^3)*x^
6 - 4*(91*B*b^6*d^2*e - 2*(88*B*a*b^5 - 39*A*b^6)*d*e^2 + (13*B*a^2*b^4 -
6*A*a*b^5)*e^3)*x^5 - (1001*B*b^6*d^3 - 13*(157*B*a*b^5 - 66*A*b^6)*d^2*e
+ (359*B*a^2*b^4 - 156*A*a*b^5)*d*e^2 - 3*(13*B*a^3*b^3 - 6*A*a^2*b^4)*e^3
)*x^4 + 143*(2*B*a^4*b^2 - 9*A*a^3*b^3)*d^3 - 91*(4*B*a^5*b - 33*A*a^4*b^2
)*d^2*e + 63*(2*B*a^6 - 39*A*a^5*b)*d*e^2 - (143*(19*B*a*b^5 + 9*A*b^6)*d^
3 - 13*(611*B*a^2*b^4 + 33*A*a*b^5)*d^2*e + (5735*B*a^3*b^3 + 117*A*a^2*b^
4)*d*e^2 - (1469*B*a^4*b^2 + 15*A*a^3*b^3)*e^3)*x^3 - (429*(5*B*a^2*b^4 +
9*A*a*b^5)*d^3 - 13*(611*B*a^3*b^3 + 495*A*a^2*b^4)*d^2*e + (7171*B*a^4*b^
2 + 4407*A*a^3*b^3)*d*e^2 - 7*(299*B*a^5*b + 159*A*a^4*b^2)*e^3)*x^2 - (14
3*(B*a^3*b^3 + 27*A*a^2*b^4)*d^3 - 13*(157*B*a^4*b^2 + 627*A*a^3*b^3)*d^2*
e + 7*(347*B*a^5*b + 897*A*a^4*b^2)*d*e^2 - 63*(13*B*a^6 + 27*A*a^5*b)*e^3
)*x)*sqrt(b*x + a)*sqrt(e*x + d)/(b^4*d^11 - 4*a*b^3*d^10*e + 6*a^2*b^2*d^
9*e^2 - 4*a^3*b*d^8*e^3 + a^4*d^7*e^4 + (b^4*d^4*e^7 - 4*a*b^3*d^3*e^8 + 6
*a^2*b^2*d^2*e^9 - 4*a^3*b*d*e^10 + a^4*e^11)*x^7 + 7*(b^4*d^5*e^6 - 4*a*b
^3*d^4*e^7 + 6*a^2*b^2*d^3*e^8 - 4*a^3*b*d^2*e^9 + a^4*d*e^10)*x^6 + 21*(b
^4*d^6*e^5 - 4*a*b^3*d^5*e^6 + 6*a^2*b^2*d^4*e^7 - 4*a^3*b*d^3*e^8 + a^4*d
^2*e^9)*x^5 + 35*(b^4*d^7*e^4 - 4*a*b^3*d^6*e^5 + 6*a^2*b^2*d^5*e^6 - 4*a^
3*b*d^4*e^7 + a^4*d^3*e^8)*x^4 + 35*(b^4*d^8*e^3 - 4*a*b^3*d^7*e^4 + 6*a^2
*b^2*d^6*e^5 - 4*a^3*b*d^5*e^6 + a^4*d^4*e^7)*x^3 + 21*(b^4*d^9*e^2 - 4...
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{15/2}} dx = \text{Timed out}$$

input `integrate((b*x+a)**(5/2)*(B*x+A)/(e*x+d)**(15/2),x)`

output Timed out

Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{15/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(15/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f or more de

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1003 vs. 2(177) = 354.

Time = 0.80 (sec) , antiderivative size = 1003, normalized size of antiderivative = 4.99

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{15/2}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(15/2),x, algorithm="giac")`

output

```
2/9009*((4*(b*x + a)*(2*(7*B*b^16*d^3*e^8*abs(b) - 27*B*a*b^15*d^2*e^9*abs(b) + 6*A*b^16*d^2*e^9*abs(b) + 33*B*a^2*b^14*d*e^10*abs(b) - 12*A*a*b^15*d*e^10*abs(b) - 13*B*a^3*b^13*e^11*abs(b) + 6*A*a^2*b^14*e^11*abs(b))*(b*x + a)/(b^8*d^6*e^6 - 6*a*b^7*d^5*e^7 + 15*a^2*b^6*d^4*e^8 - 20*a^3*b^5*d^3*e^9 + 15*a^4*b^4*d^2*e^10 - 6*a^5*b^3*d*e^11 + a^6*b^2*e^12) + 13*(7*B*b^17*d^4*e^7*abs(b) - 34*B*a*b^16*d^3*e^8*abs(b) + 6*A*b^17*d^3*e^8*abs(b) + 60*B*a^2*b^15*d^2*e^9*abs(b) - 18*A*a*b^16*d^2*e^9*abs(b) - 46*B*a^3*b^14*d*e^10*abs(b) + 18*A*a^2*b^15*d*e^10*abs(b) + 13*B*a^4*b^13*e^11*abs(b) - 6*A*a^3*b^14*e^11*abs(b))/(b^8*d^6*e^6 - 6*a*b^7*d^5*e^7 + 15*a^2*b^6*d^4*e^8 - 20*a^3*b^5*d^3*e^9 + 15*a^4*b^4*d^2*e^10 - 6*a^5*b^3*d*e^11 + a^6*b^2*e^12)) + 143*(7*B*b^18*d^5*e^6*abs(b) - 41*B*a*b^17*d^4*e^7*abs(b) + 6*A*b^18*d^4*e^7*abs(b) + 94*B*a^2*b^16*d^3*e^8*abs(b) - 24*A*a*b^17*d^3*e^8*abs(b) - 106*B*a^3*b^15*d^2*e^9*abs(b) + 36*A*a^2*b^16*d^2*e^9*abs(b) + 59*B*a^4*b^14*d*e^10*abs(b) - 24*A*a^3*b^15*d*e^10*abs(b) - 13*B*a^5*b^13*e^11*abs(b) + 6*A*a^4*b^14*e^11*abs(b))/(b^8*d^6*e^6 - 6*a*b^7*d^5*e^7 + 15*a^2*b^6*d^4*e^8 - 20*a^3*b^5*d^3*e^9 + 15*a^4*b^4*d^2*e^10 - 6*a^5*b^3*d*e^11 + a^6*b^2*e^12))*(b*x + a) - 1287*(B*a*b^18*d^5*e^6*abs(b) - A*b^19*d^5*e^6*abs(b) - 5*B*a^2*b^17*d^4*e^7*abs(b) + 5*A*a*b^18*d^4*e^7*abs(b) + 10*B*a^3*b^16*d^3*e^8*abs(b) - 10*A*a^2*b^17*d^3*e^8*abs(b) - 10*B*a^4*b^15*d^2*e^9*abs(b) + 10*A*a^3*b^16*d^2*e^9*abs(b) + 5*B*a^5*b^14*d*e^10*a...
```

Mupad [B] (verification not implemented)

Time = 2.13 (sec) , antiderivative size = 706, normalized size of antiderivative = 3.51

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{15/2}} dx = \frac{\sqrt{d + ex} \left(\frac{x^3 \sqrt{a+bx} (-2938 B a^4 b^2 e^3 + 11470 B a^3 b^3 d e^2 - 30 A a^3 b^3 e^3 - 15886 B a^2 b^4 d^2 e + 234 A a^2 b^4 d^2 e^2 - 15886 B a^2 b^4 d^2 e + 234 A a^2 b^4 d^2 e^2)}{9009 e^7 (a e - b d)^4} \right)}{(d + ex)^{15/2}}$$

input

```
int(((A + B*x)*(a + b*x)^(5/2))/(d + e*x)^(15/2),x)
```

output

```

((d + e*x)^(1/2)*((x^3*(a + b*x)^(1/2)*(2574*A*b^6*d^3 + 5434*B*a*b^5*d^3
- 30*A*a^3*b^3*e^3 - 2938*B*a^4*b^2*e^3 + 234*A*a^2*b^4*d*e^2 - 15886*B*a^
2*b^4*d^2*e + 11470*B*a^3*b^3*d*e^2 - 858*A*a*b^5*d^2*e))/(9009*e^7*(a*e -
b*d)^4) - ((a + b*x)^(1/2)*(1386*A*a^6*e^3 + 252*B*a^6*d*e^2 - 2574*A*a^3
*b^3*d^3 + 572*B*a^4*b^2*d^3 + 6006*A*a^4*b^2*d^2*e - 4914*A*a^5*b*d*e^2 -
728*B*a^5*b*d^2*e))/(9009*e^7*(a*e - b*d)^4) - (x*(a + b*x)^(1/2)*(1638*B
*a^6*e^3 + 3402*A*a^5*b*e^3 - 7722*A*a^2*b^4*d^3 - 286*B*a^3*b^3*d^3 + 163
02*A*a^3*b^3*d^2*e - 12558*A*a^4*b^2*d*e^2 + 4082*B*a^4*b^2*d^2*e - 4858*B
*a^5*b*d*e^2))/(9009*e^7*(a*e - b*d)^4) + (x^2*(a + b*x)^(1/2)*(7722*A*a*b
^5*d^3 - 4186*B*a^5*b*e^3 - 2226*A*a^4*b^2*e^3 + 4290*B*a^2*b^4*d^3 - 1287
0*A*a^2*b^4*d^2*e + 8814*A*a^3*b^3*d*e^2 - 15886*B*a^3*b^3*d^2*e + 14342*B
*a^4*b^2*d*e^2))/(9009*e^7*(a*e - b*d)^4) + (16*b^5*x^6*(a + b*x)^(1/2)*(6
*A*b*e - 13*B*a*e + 7*B*b*d))/(9009*e^5*(a*e - b*d)^4) - (8*b^4*x^5*(a*e -
13*b*d)*(a + b*x)^(1/2)*(6*A*b*e - 13*B*a*e + 7*B*b*d))/(9009*e^6*(a*e -
b*d)^4) + (2*b^3*x^4*(a + b*x)^(1/2)*(3*a^2*e^2 + 143*b^2*d^2 - 26*a*b*d*e
)*(6*A*b*e - 13*B*a*e + 7*B*b*d))/(9009*e^7*(a*e - b*d)^4)))/(x^7 + d^7/e^
7 + (7*d*x^6)/e + (7*d^6*x)/e^6 + (21*d^2*x^5)/e^2 + (35*d^3*x^4)/e^3 + (3
5*d^4*x^3)/e^4 + (21*d^5*x^2)/e^5)

```

Reduce [B] (verification not implemented)

Time = 1.06 (sec) , antiderivative size = 1018, normalized size of antiderivative = 5.06

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{15/2}} dx = \text{Too large to display}$$

input

```
int((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(15/2),x)
```

output

```
(2*( - 99*sqrt(d + e*x)*sqrt(a + b*x)*a**6*e**7 + 234*sqrt(d + e*x)*sqrt(a
+ b*x)*a**5*b*d*e**6 - 360*sqrt(d + e*x)*sqrt(a + b*x)*a**5*b*e**7*x - 14
3*sqrt(d + e*x)*sqrt(a + b*x)*a**4*b**2*d**2*e**5 + 884*sqrt(d + e*x)*sqrt
(a + b*x)*a**4*b**2*d*e**6*x - 458*sqrt(d + e*x)*sqrt(a + b*x)*a**4*b**2*e
**7*x**2 - 572*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b**3*d**2*e**5*x + 1196*sq
rt(d + e*x)*sqrt(a + b*x)*a**3*b**3*d*e**6*x**2 - 212*sqrt(d + e*x)*sqrt(a
+ b*x)*a**3*b**3*e**7*x**3 - 858*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**4*d*
**2*e**5*x**2 + 624*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**4*d*e**6*x**3 - 3*s
qrt(d + e*x)*sqrt(a + b*x)*a**2*b**4*e**7*x**4 - 572*sqrt(d + e*x)*sqrt(a
+ b*x)*a*b**5*d**2*e**5*x**3 + 26*sqrt(d + e*x)*sqrt(a + b*x)*a*b**5*d*e**
6*x**4 + 4*sqrt(d + e*x)*sqrt(a + b*x)*a*b**5*e**7*x**5 - 143*sqrt(d + e*x
)*sqrt(a + b*x)*b**6*d**2*e**5*x**4 - 52*sqrt(d + e*x)*sqrt(a + b*x)*b**6*
d*e**6*x**5 - 8*sqrt(d + e*x)*sqrt(a + b*x)*b**6*e**7*x**6 + 8*sqrt(e)*sq
rt(b)*b**6*d**7 + 56*sqrt(e)*sqrt(b)*b**6*d**6*e*x + 168*sqrt(e)*sqrt(b)*b
**6*d**5*e**2*x**2 + 280*sqrt(e)*sqrt(b)*b**6*d**4*e**3*x**3 + 280*sqrt(e)*
sqrt(b)*b**6*d**3*e**4*x**4 + 168*sqrt(e)*sqrt(b)*b**6*d**2*e**5*x**5 + 56
*sqrt(e)*sqrt(b)*b**6*d*e**6*x**6 + 8*sqrt(e)*sqrt(b)*b**6*e**7*x**7))/(12
87*e**5*(a**3*d**7*e**3 + 7*a**3*d**6*e**4*x + 21*a**3*d**5*e**5*x**2 + 35
*a**3*d**4*e**6*x**3 + 35*a**3*d**3*e**7*x**4 + 21*a**3*d**2*e**8*x**5 + 7
*a**3*d*e**9*x**6 + a**3*e**10*x**7 - 3*a**2*b*d**8*e**2 - 21*a**2*b*d*...
```

3.204 $\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{17/2}} dx$

Optimal result	1863
Mathematica [A] (verified)	1864
Rubi [A] (verified)	1864
Maple [B] (verified)	1867
Fricas [F(-1)]	1867
Sympy [F(-1)]	1868
Maxima [F(-2)]	1868
Giac [B] (verification not implemented)	1868
Mupad [B] (verification not implemented)	1869
Reduce [B] (verification not implemented)	1870

Optimal result

Integrand size = 24, antiderivative size = 255

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{17/2}} dx = -\frac{2(Bd - Ae)(a+bx)^{7/2}}{15e(bd - ae)(d+ex)^{15/2}} + \frac{2(7bBd + 8Abe - 15aBe)(a+bx)^{7/2}}{195e(bd - ae)^2(d+ex)^{13/2}} + \frac{4b(7bBd + 8Abe - 15aBe)(a+bx)^{7/2}}{715e(bd - ae)^3(d+ex)^{11/2}} + \frac{16b^2(7bBd + 8Abe - 15aBe)(a+bx)^{7/2}}{6435e(bd - ae)^4(d+ex)^{9/2}} + \frac{32b^3(7bBd + 8Abe - 15aBe)(a+bx)^{7/2}}{45045e(bd - ae)^5(d+ex)^{7/2}}$$

output

```
-2/15*(-A*e+B*d)*(b*x+a)^(7/2)/e/(-a*e+b*d)/(e*x+d)^(15/2)+2/195*(8*A*b*e-15*B*a*e+7*B*b*d)*(b*x+a)^(7/2)/e/(-a*e+b*d)^2/(e*x+d)^(13/2)+4/715*b*(8*A*b*e-15*B*a*e+7*B*b*d)*(b*x+a)^(7/2)/e/(-a*e+b*d)^3/(e*x+d)^(11/2)+16/6435*b^2*(8*A*b*e-15*B*a*e+7*B*b*d)*(b*x+a)^(7/2)/e/(-a*e+b*d)^4/(e*x+d)^(9/2)+32/45045*b^3*(8*A*b*e-15*B*a*e+7*B*b*d)*(b*x+a)^(7/2)/e/(-a*e+b*d)^5/(e*x+d)^(7/2)
```

Mathematica [A] (verified)

Time = 0.40 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.06

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{17/2}} dx = \frac{2(a+bx)^{7/2}(-3003Bde^3(a+bx)^4 + 3003Ae^4(a+bx)^4 + 10395bBde^2(a+bx)^3 + 13860A^2bde^3(a+bx)^3 + 3465a^2Bde^3(a+bx)^3 + 12285a^2B^2de^2(a+bx)^2 + 24570A^2b^2e^2(a+bx)^2 + 12285a^2b^2Bde^2(a+bx)^2 + 5005b^3B^2d(a+bx)(d+ex)^3 - 20020A^2b^3e(a+bx)(d+ex)^3 + 15015a^2b^2B^2e(a+bx)(d+ex)^3 + 6435A^2b^4(d+ex)^4 - 6435a^2b^3B^2(d+ex)^4)}{(45045(bd-ae)^5(d+ex)^{15/2}}$$

input

```
Integrate[((a + b*x)^(5/2)*(A + B*x))/(d + e*x)^(17/2),x]
```

output

```
(2*(a + b*x)^(7/2)*(-3003*B*d*e^3*(a + b*x)^4 + 3003*A*e^4*(a + b*x)^4 + 10395*b*B*d*e^2*(a + b*x)^3*(d + e*x) - 13860*A*b*e^3*(a + b*x)^3*(d + e*x) + 3465*a*B*e^3*(a + b*x)^3*(d + e*x) - 12285*b^2*B*d*e*(a + b*x)^2*(d + e*x)^2 + 24570*A*b^2*e^2*(a + b*x)^2*(d + e*x)^2 - 12285*a*b*B*e^2*(a + b*x)^2*(d + e*x)^2 + 5005*b^3*B*d*(a + b*x)*(d + e*x)^3 - 20020*A*b^3*e*(a + b*x)*(d + e*x)^3 + 15015*a*b^2*B*e*(a + b*x)*(d + e*x)^3 + 6435*A*b^4*(d + e*x)^4 - 6435*a*b^3*B*(d + e*x)^4)/(45045*(b*d - a*e)^5*(d + e*x)^(15/2))
```

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 239, normalized size of antiderivative = 0.94, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$, Rules used = {87, 55, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a+bx)^{5/2}(A+Bx)}{(d+ex)^{17/2}} dx$$

$$\downarrow 87$$

$$\frac{(-15aBe + 8Abe + 7bBd) \int \frac{(a+bx)^{5/2}}{(d+ex)^{15/2}} dx}{15e(bd-ae)} - \frac{2(a+bx)^{7/2}(Bd-Ae)}{15e(d+ex)^{15/2}(bd-ae)}$$

$$\downarrow 55$$

$$\frac{(-15aBe + 8Abe + 7bBd) \left(\frac{6b \int \frac{(a+bx)^{5/2}}{(d+ex)^{13/2}} dx}{13(bd-ae)} + \frac{2(a+bx)^{7/2}}{13(d+ex)^{13/2}(bd-ae)} \right)}{15e(bd-ae)} - \frac{2(a+bx)^{7/2}(Bd-Ae)}{15e(d+ex)^{15/2}(bd-ae)}$$

↓ 55

$$\frac{(-15aBe + 8Abe + 7bBd) \left(\frac{6b \left(\frac{4b \int \frac{(a+bx)^{5/2}}{(d+ex)^{11/2}} dx}{11(bd-ae)} + \frac{2(a+bx)^{7/2}}{11(d+ex)^{11/2}(bd-ae)} \right)}{13(bd-ae)} + \frac{2(a+bx)^{7/2}}{13(d+ex)^{13/2}(bd-ae)} \right)}{15e(bd-ae)}$$

$$\frac{2(a+bx)^{7/2}(Bd-Ae)}{15e(d+ex)^{15/2}(bd-ae)}$$

↓ 55

$$\frac{(-15aBe + 8Abe + 7bBd) \left(\frac{6b \left(\frac{4b \left(\frac{2b \int \frac{(a+bx)^{5/2}}{(d+ex)^{9/2}} dx}{9(bd-ae)} + \frac{2(a+bx)^{7/2}}{9(d+ex)^{9/2}(bd-ae)} \right)}{11(bd-ae)} + \frac{2(a+bx)^{7/2}}{11(d+ex)^{11/2}(bd-ae)} \right)}{13(bd-ae)} + \frac{2(a+bx)^{7/2}}{13(d+ex)^{13/2}(bd-ae)} \right)}{15e(bd-ae)}$$

$$\frac{2(a+bx)^{7/2}(Bd-Ae)}{15e(d+ex)^{15/2}(bd-ae)}$$

↓ 48

$$\left(\frac{2(a+bx)^{7/2}}{13(d+ex)^{13/2}(bd-ae)} + \frac{6b \left(\frac{2(a+bx)^{7/2}}{11(d+ex)^{11/2}(bd-ae)} + \frac{4b \left(\frac{4b(a+bx)^{7/2}}{63(d+ex)^{7/2}(bd-ae)^2} + \frac{2(a+bx)^{7/2}}{9(d+ex)^{9/2}(bd-ae)} \right)}{11(bd-ae)} \right)}{13(bd-ae)} \right) (-15aBe + 8Abe + 7bBd)$$

$$\frac{2(a+bx)^{7/2}(Bd-Ae)}{15e(d+ex)^{15/2}(bd-ae)}$$

input `Int[((a + b*x)^(5/2)*(A + B*x))/(d + e*x)^(17/2),x]`

output `(-2*(B*d - A*e)*(a + b*x)^(7/2))/(15*e*(b*d - a*e)*(d + e*x)^(15/2)) + ((7*b*B*d + 8*A*b*e - 15*a*B*e)*((2*(a + b*x)^(7/2))/(13*(b*d - a*e)*(d + e*x)^(13/2)) + (6*b*((2*(a + b*x)^(7/2))/(11*(b*d - a*e)*(d + e*x)^(11/2)) + (4*b*((2*(a + b*x)^(7/2))/(9*(b*d - a*e)*(d + e*x)^(9/2)) + (4*b*(a + b*x)^(7/2))/(63*(b*d - a*e)^2*(d + e*x)^(7/2))))/(11*(b*d - a*e)))/(13*(b*d - a*e)))/(15*e*(b*d - a*e))`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 55 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(Simplify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{17/2}} dx = \text{Timed out}$$

input `integrate((b*x+a)**(5/2)*(B*x+A)/(e*x+d)**(17/2),x)`

output `Timed out`

Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{17/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(17/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f or more de`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1413 vs. 2(225) = 450.

Time = 1.06 (sec) , antiderivative size = 1413, normalized size of antiderivative = 5.54

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{17/2}} dx = \text{Too large to display}$$

input `integrate((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(17/2),x, algorithm="giac")`

output

```

2/45045*((2*(4*(b*x + a))*(2*(7*B*b^18*d^3*e^10*abs(b) - 29*B*a*b^17*d^2*e^
11*abs(b) + 8*A*b^18*d^2*e^11*abs(b) + 37*B*a^2*b^16*d*e^12*abs(b) - 16*A*
a*b^17*d*e^12*abs(b) - 15*B*a^3*b^15*e^13*abs(b) + 8*A*a^2*b^16*e^13*abs(b)
))*(b*x + a)/(b^9*d^7*e^7 - 7*a*b^8*d^6*e^8 + 21*a^2*b^7*d^5*e^9 - 35*a^3*
b^6*d^4*e^10 + 35*a^4*b^5*d^3*e^11 - 21*a^5*b^4*d^2*e^12 + 7*a^6*b^3*d*e^1
3 - a^7*b^2*e^14) + 15*(7*B*b^19*d^4*e^9*abs(b) - 36*B*a*b^18*d^3*e^10*abs
(b) + 8*A*b^19*d^3*e^10*abs(b) + 66*B*a^2*b^17*d^2*e^11*abs(b) - 24*A*a*b^
18*d^2*e^11*abs(b) - 52*B*a^3*b^16*d*e^12*abs(b) + 24*A*a^2*b^17*d*e^12*ab
s(b) + 15*B*a^4*b^15*e^13*abs(b) - 8*A*a^3*b^16*e^13*abs(b))/(b^9*d^7*e^7
- 7*a*b^8*d^6*e^8 + 21*a^2*b^7*d^5*e^9 - 35*a^3*b^6*d^4*e^10 + 35*a^4*b^5*
d^3*e^11 - 21*a^5*b^4*d^2*e^12 + 7*a^6*b^3*d*e^13 - a^7*b^2*e^14)) + 195*(
7*B*b^20*d^5*e^8*abs(b) - 43*B*a*b^19*d^4*e^9*abs(b) + 8*A*b^20*d^4*e^9*ab
s(b) + 102*B*a^2*b^18*d^3*e^10*abs(b) - 32*A*a*b^19*d^3*e^10*abs(b) - 118*
B*a^3*b^17*d^2*e^11*abs(b) + 48*A*a^2*b^18*d^2*e^11*abs(b) + 67*B*a^4*b^16
*d*e^12*abs(b) - 32*A*a^3*b^17*d*e^12*abs(b) - 15*B*a^5*b^15*e^13*abs(b) +
8*A*a^4*b^16*e^13*abs(b))/(b^9*d^7*e^7 - 7*a*b^8*d^6*e^8 + 21*a^2*b^7*d^5
*e^9 - 35*a^3*b^6*d^4*e^10 + 35*a^4*b^5*d^3*e^11 - 21*a^5*b^4*d^2*e^12 + 7
*a^6*b^3*d*e^13 - a^7*b^2*e^14))*(b*x + a) + 715*(7*B*b^21*d^6*e^7*abs(b)
- 50*B*a*b^20*d^5*e^8*abs(b) + 8*A*b^21*d^5*e^8*abs(b) + 145*B*a^2*b^19*d^
4*e^9*abs(b) - 40*A*a*b^20*d^4*e^9*abs(b) - 220*B*a^3*b^18*d^3*e^10*abs...

```

Mupad [B] (verification not implemented)

Time = 2.55 (sec) , antiderivative size = 917, normalized size of antiderivative = 3.60

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{17/2}} dx = \text{Too large to display}$$

input

```
int(((A + B*x)*(a + b*x)^(5/2))/(d + e*x)^(17/2),x)
```

output

```

-((d + e*x)^(1/2)*(((a + b*x)^(1/2)*(6006*A*a^7*e^4 + 924*B*a^7*d*e^3 + 12
870*A*a^3*b^4*d^4 - 2860*B*a^4*b^3*d^4 - 40040*A*a^4*b^3*d^3*e + 5460*B*a^
5*b^2*d^3*e - 3780*B*a^6*b*d^2*e^2 + 49140*A*a^5*b^2*d^2*e^2 - 27720*A*a^6
*b*d*e^3)))/(45045*e^8*(a*e - b*d)^5) + (x^2*(a + b*x)^(1/2)*(38610*A*a*b^6
*d^4 + 17010*B*a^6*b*e^4 + 8946*A*a^5*b^2*e^4 + 21450*B*a^2*b^5*d^4 - 8580
0*A*a^2*b^5*d^3*e - 44520*A*a^4*b^3*d*e^3 - 99840*B*a^3*b^4*d^3*e - 77616*
B*a^5*b^2*d*e^3 + 88140*A*a^3*b^4*d^2*e^2 + 133620*B*a^4*b^3*d^2*e^2))/(45
045*e^8*(a*e - b*d)^5) + (x^3*(a + b*x)^(1/2)*(12870*A*b^7*d^4 + 27170*B*a
*b^6*d^4 + 70*A*a^4*b^3*e^4 + 11130*B*a^5*b^2*e^4 - 600*A*a^3*b^4*d*e^3 -
99840*B*a^2*b^5*d^3*e - 55120*B*a^4*b^3*d*e^3 + 2340*A*a^2*b^5*d^2*e^2 + 1
07700*B*a^3*b^4*d^2*e^2 - 5720*A*a*b^6*d^3*e))/(45045*e^8*(a*e - b*d)^5) +
(x*(a + b*x)^(1/2)*(6930*B*a^7*e^4 + 14322*A*a^6*b*e^4 + 38610*A*a^2*b^5*
d^4 + 1430*B*a^3*b^4*d^4 - 108680*A*a^3*b^4*d^3*e - 68040*A*a^5*b^2*d*e^3
- 24180*B*a^4*b^3*d^3*e + 125580*A*a^4*b^3*d^2*e^2 + 42840*B*a^5*b^2*d^2*
e^2 - 28812*B*a^6*b*d*e^3))/(45045*e^8*(a*e - b*d)^5) + (32*b^6*x^7*(a + b*
x)^(1/2)*(8*A*b*e - 15*B*a*e + 7*B*b*d))/(45045*e^5*(a*e - b*d)^5) - (2*b^
3*x^4*(a + b*x)^(1/2)*(8*A*b*e - 15*B*a*e + 7*B*b*d)*(a^3*e^3 - 143*b^3*d^
3 + 39*a*b^2*d^2*e - 9*a^2*b*d*e^2))/(9009*e^8*(a*e - b*d)^5) - (16*b^5*x^
6*(a*e - 15*b*d)*(a + b*x)^(1/2)*(8*A*b*e - 15*B*a*e + 7*B*b*d))/(45045*e^
6*(a*e - b*d)^5) + (4*b^4*x^5*(a + b*x)^(1/2)*(a^2*e^2 + 65*b^2*d^2 - 1...

```

Reduce [B] (verification not implemented)

Time = 2.06 (sec) , antiderivative size = 1457, normalized size of antiderivative = 5.71

$$\int \frac{(a + bx)^{5/2}(A + Bx)}{(d + ex)^{17/2}} dx = \text{Too large to display}$$

input

```
int((b*x+a)^(5/2)*(B*x+A)/(e*x+d)^(17/2),x)
```

output

```
(2*( - 429*sqrt(d + e*x)*sqrt(a + b*x)*a**7*e**8 + 1485*sqrt(d + e*x)*sqrt
(a + b*x)*a**6*b*d*e**7 - 1518*sqrt(d + e*x)*sqrt(a + b*x)*a**6*b*e**8*x -
1755*sqrt(d + e*x)*sqrt(a + b*x)*a**5*b**2*d**2*e**6 + 5400*sqrt(d + e*x)
*sqrt(a + b*x)*a**5*b**2*d*e**7*x - 1854*sqrt(d + e*x)*sqrt(a + b*x)*a**5*
b**2*e**8*x**2 + 715*sqrt(d + e*x)*sqrt(a + b*x)*a**4*b**3*d**3*e**5 - 663
0*sqrt(d + e*x)*sqrt(a + b*x)*a**4*b**3*d**2*e**6*x + 6870*sqrt(d + e*x)*s
qrt(a + b*x)*a**4*b**3*d*e**7*x**2 - 800*sqrt(d + e*x)*sqrt(a + b*x)*a**4*
b**3*e**8*x**3 + 2860*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b**4*d**3*e**5*x -
8970*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b**4*d**2*e**6*x**2 + 3180*sqrt(d +
e*x)*sqrt(a + b*x)*a**3*b**4*d*e**7*x**3 - 5*sqrt(d + e*x)*sqrt(a + b*x)*a
**3*b**4*e**8*x**4 + 4290*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**5*d**3*e**5*
x**2 - 4680*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**5*d**2*e**6*x**3 + 45*sqrt
(d + e*x)*sqrt(a + b*x)*a**2*b**5*d*e**7*x**4 + 6*sqrt(d + e*x)*sqrt(a + b
*x)*a**2*b**5*e**8*x**5 + 2860*sqrt(d + e*x)*sqrt(a + b*x)*a*b**6*d**3*e**
5*x**3 - 195*sqrt(d + e*x)*sqrt(a + b*x)*a*b**6*d**2*e**6*x**4 - 60*sqrt(d
+ e*x)*sqrt(a + b*x)*a*b**6*d*e**7*x**5 - 8*sqrt(d + e*x)*sqrt(a + b*x)*a
*b**6*e**8*x**6 + 715*sqrt(d + e*x)*sqrt(a + b*x)*b**7*d**3*e**5*x**4 + 39
0*sqrt(d + e*x)*sqrt(a + b*x)*b**7*d**2*e**6*x**5 + 120*sqrt(d + e*x)*sqrt
(a + b*x)*b**7*d*e**7*x**6 + 16*sqrt(d + e*x)*sqrt(a + b*x)*b**7*e**8*x**7
- 16*sqrt(e)*sqrt(b)*b**7*d**8 - 128*sqrt(e)*sqrt(b)*b**7*d**7*e*x - 4...
```

3.205 $\int \frac{(A+Bx)(d+ex)^{5/2}}{\sqrt{a+bx}} dx$

Optimal result	1872
Mathematica [A] (verified)	1873
Rubi [A] (verified)	1873
Maple [B] (verified)	1876
Fricas [A] (verification not implemented)	1877
Sympy [F]	1878
Maxima [F(-2)]	1879
Giac [B] (verification not implemented)	1879
Mupad [F(-1)]	1880
Reduce [B] (verification not implemented)	1881

Optimal result

Integrand size = 24, antiderivative size = 246

$$\int \frac{(A+Bx)(d+ex)^{5/2}}{\sqrt{a+bx}} dx =$$

$$\frac{5(bd-ae)^2(bBd-8Abe+7aBe)\sqrt{a+bx}\sqrt{d+ex}}{64b^4e}$$

$$- \frac{5(bd-ae)(bBd-8Abe+7aBe)\sqrt{a+bx}(d+ex)^{3/2}}{96b^3e}$$

$$- \frac{(bBd-8Abe+7aBe)\sqrt{a+bx}(d+ex)^{5/2}}{24b^2e} + \frac{B\sqrt{a+bx}(d+ex)^{7/2}}{4be}$$

$$- \frac{5(bd-ae)^3(bBd-8Abe+7aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{64b^{9/2}e^{3/2}}$$

output

```
-5/64*(-a*e+b*d)^2*(-8*A*b*e+7*B*a*e+B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(1/2)/b^
4/e-5/96*(-a*e+b*d)*(-8*A*b*e+7*B*a*e+B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(3/2)/b
^3/e-1/24*(-8*A*b*e+7*B*a*e+B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(5/2)/b^2/e+1/4*B
*(b*x+a)^(1/2)*(e*x+d)^(7/2)/b/e-5/64*(-a*e+b*d)^3*(-8*A*b*e+7*B*a*e+B*b*d
)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)^(1/2))/b^(9/2)/e^(3/2)
```


$$\begin{array}{c}
 \frac{B\sqrt{a+bx}(d+ex)^{7/2}}{4be} - \frac{(7aBe - 8Abe + bBd) \left(\frac{5(bd-ae) \int \frac{(d+ex)^{3/2}}{\sqrt{a+bx}} dx}{6b} + \frac{\sqrt{a+bx}(d+ex)^{5/2}}{3b} \right)}{8be} \\
 \downarrow 60 \\
 \frac{B\sqrt{a+bx}(d+ex)^{7/2}}{4be} - \frac{(7aBe - 8Abe + bBd) \left(\frac{5(bd-ae) \left(\frac{3(bd-ae) \int \frac{\sqrt{d+ex}}{\sqrt{a+bx}} dx}{4b} + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{6b} + \frac{\sqrt{a+bx}(d+ex)^{5/2}}{3b} \right)}{8be} \\
 \downarrow 60 \\
 \frac{B\sqrt{a+bx}(d+ex)^{7/2}}{4be} - \frac{(7aBe - 8Abe + bBd) \left(\frac{5(bd-ae) \left(\frac{3(bd-ae) \left(\frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{4b} + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{4b} + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{6b} + \frac{\sqrt{a+bx}(d+ex)^{5/2}}{3b} \right)}{8be} \\
 \downarrow 66 \\
 \frac{B\sqrt{a+bx}(d+ex)^{7/2}}{4be} - \frac{(7aBe - 8Abe + bBd) \left(\frac{5(bd-ae) \left(\frac{3(bd-ae) \left(\frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{4b} + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{4b} + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{6b} + \frac{\sqrt{a+bx}(d+ex)^{5/2}}{3b} \right)}{8be} \\
 \downarrow 221
 \end{array}$$

$$\frac{B\sqrt{a+bx}(d+ex)^{7/2}}{4be} - \frac{(7aBe - 8Abe + bBd) \left(\frac{5(bd-ae) \left(\frac{3(bd-ae) \operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right) + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b}\right)}{4b} + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{6b} + \frac{\sqrt{a+bx}(d+ex)^{5/2}}{3b} \right)}{8be}$$

input `Int[((A + B*x)*(d + e*x)^(5/2))/Sqrt[a + b*x],x]`

output `(B*Sqrt[a + b*x]*(d + e*x)^(7/2))/(4*b*e) - ((b*B*d - 8*A*b*e + 7*a*B*e)*(Sqrt[a + b*x]*(d + e*x)^(5/2))/(3*b) + (5*(b*d - a*e)*((Sqrt[a + b*x]*(d + e*x)^(3/2))/(2*b) + (3*(b*d - a*e)*((Sqrt[a + b*x]*Sqrt[d + e*x])/b + ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])]))/(b^(3/2)*Sqrt[e])))/(4*b)))/(6*b)))/(8*b*e)`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_.), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_.)]*Sqrt[(c_) + (d_.)*(x_.)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 967 vs. $2(208) = 416$.

Time = 0.27 (sec) , antiderivative size = 968, normalized size of antiderivative = 3.93

method	result
default	$-\frac{\sqrt{ex+d}\sqrt{bx+a}\left(-96Bb^3e^3x^3\sqrt{(ex+d)(bx+a)}\sqrt{be}-128Ab^3e^3x^2\sqrt{(ex+d)(bx+a)}\sqrt{be}-272Bb^3de^2x^2\sqrt{(ex+d)(bx+a)}\sqrt{be}-\dots\right)}{\dots}$

input `int((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^(1/2),x,method=_RETURNVERBOSE)`

output

```

-1/384*(e*x+d)^(1/2)*(b*x+a)^(1/2)*(-96*B*b^3*e^3*x^3*((e*x+d)*(b*x+a))^(1
/2)*(b*e)^(1/2)-128*A*b^3*e^3*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-272*
B*b^3*d*e^2*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-105*B*ln(1/2*(2*b*e*x+
2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^4*e^4+15*B*ln
(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))
*b^4*d^4+344*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b^2*d*e^2*x-360*A*ln(
1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a
^2*b^2*d*e^3+360*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a
*e+d*b)/(b*e)^(1/2))*a*b^3*d^2*e^2+160*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/
2)*a*b^2*e^3*x-416*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^3*d*e^2*x-140*B
*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^2*b*e^3*x-236*B*((e*x+d)*(b*x+a))^(
1/2)*(b*e)^(1/2)*b^3*d^2*e*x+210*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^3
*e^3-30*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^3*d^3+120*A*ln(1/2*(2*b*e*
x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^3*b*e^4-12
0*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(
1/2))*b^4*d^3*e+640*A*a*b^2*d*e^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+112*
B*a*b^2*e^3*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+60*B*ln(1/2*(2*b*e*x+2
*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^3*d^3*e-240
*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^2*b*e^3-528*A*((e*x+d)*(b*x+a))^(
1/2)*(b*e)^(1/2)*b^3*d^2*e+300*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1...

```

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 772, normalized size of antiderivative = 3.14

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{\sqrt{a + bx}} dx = \text{Too large to display}$$

input

```
integrate((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^(1/2),x, algorithm="fricas")
```

output

```
[1/768*(15*(B*b^4*d^4 + 4*(B*a*b^3 - 2*A*b^4)*d^3*e - 6*(3*B*a^2*b^2 - 4*A*a*b^3)*d^2*e^2 + 4*(5*B*a^3*b - 6*A*a^2*b^2)*d*e^3 - (7*B*a^4 - 8*A*a^3*b)*e^4)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 - 4*(2*b*e*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x + a)*sqrt(e*x + d) + 8*(b^2*d*e + a*b*e^2)*x) + 4*(48*B*b^4*e^4*x^3 + 15*B*b^4*d^3*e - (191*B*a*b^3 - 264*A*b^4)*d^2*e^2 + 5*(53*B*a^2*b^2 - 64*A*a*b^3)*d*e^3 - 15*(7*B*a^3*b - 8*A*a^2*b^2)*e^4 + 8*(17*B*b^4*d*e^3 - (7*B*a*b^3 - 8*A*b^4)*e^4)*x^2 + 2*(59*B*b^4*d^2*e^2 - 2*(43*B*a*b^3 - 52*A*b^4)*d*e^3 + 5*(7*B*a^2*b^2 - 8*A*a*b^3)*e^4)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b^5*e^2), 1/384*(15*(B*b^4*d^4 + 4*(B*a*b^3 - 2*A*b^4)*d^3*e - 6*(3*B*a^2*b^2 - 4*A*a*b^3)*d^2*e^2 + 4*(5*B*a^3*b - 6*A*a^2*b^2)*d*e^3 - (7*B*a^4 - 8*A*a^3*b)*e^4)*sqrt(-b*e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*e^2*x^2 + a*b*d*e + (b^2*d*e + a*b*e^2)*x)) + 2*(48*B*b^4*e^4*x^3 + 15*B*b^4*d^3*e - (191*B*a*b^3 - 264*A*b^4)*d^2*e^2 + 5*(53*B*a^2*b^2 - 64*A*a*b^3)*d*e^3 - 15*(7*B*a^3*b - 8*A*a^2*b^2)*e^4 + 8*(17*B*b^4*d*e^3 - (7*B*a*b^3 - 8*A*b^4)*e^4)*x^2 + 2*(59*B*b^4*d^2*e^2 - 2*(43*B*a*b^3 - 52*A*b^4)*d*e^3 + 5*(7*B*a^2*b^2 - 8*A*a*b^3)*e^4)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b^5*e^2)]
```

Sympy [F]

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{\sqrt{a + bx}} dx = \int \frac{(A + Bx)(d + ex)^{5/2}}{\sqrt{a + bx}} dx$$

input

```
integrate((B*x+A)*(e*x+d)**(5/2)/(b*x+a)**(1/2),x)
```

output

```
Integral((A + B*x)*(d + e*x)**(5/2)/sqrt(a + b*x), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{\sqrt{a + bx}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^(1/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1048 vs. 2(208) = 416.

Time = 0.28 (sec) , antiderivative size = 1048, normalized size of antiderivative = 4.26

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{\sqrt{a + bx}} dx = \text{Too large to display}$$

input `integrate((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^(1/2),x, algorithm="giac")`

output

```

-1/192*(192*((b^2*d - a*b*e)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d
+ (b*x + a)*b*e - a*b*e)))/sqrt(b*e) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e
)*sqrt(b*x + a))*A*d^2*abs(b)/b^2 - (sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(
2*(b*x + a)*(4*(b*x + a)*(6*(b*x + a)/b^3 + (b^12*d*e^5 - 25*a*b^11*e^6)/(
b^14*e^6)) - (5*b^13*d^2*e^4 + 14*a*b^12*d*e^5 - 163*a^2*b^11*e^6)/(b^14*e
^6)) + 3*(5*b^14*d^3*e^3 + 9*a*b^13*d^2*e^4 + 15*a^2*b^12*d*e^5 - 93*a^3*b
^11*e^6)/(b^14*e^6))*sqrt(b*x + a) + 3*(5*b^4*d^4 + 4*a*b^3*d^3*e + 6*a^2*
b^2*d^2*e^2 + 20*a^3*b*d*e^3 - 35*a^4*e^4)*log(abs(-sqrt(b*e)*sqrt(b*x + a
) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*b^2*e^3))*B*e^2*abs(b
)/b^2 - 48*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*b*x + 2*a + (b*d*e - 5*
a*e^2)/e^2)*sqrt(b*x + a) + (b^3*d^2 + 2*a*b^2*d*e - 3*a^2*b*e^2)*log(abs(
-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e
)*e))*B*d^2*abs(b)/b^3 - 96*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*b*x +
2*a + (b*d*e - 5*a*e^2)/e^2)*sqrt(b*x + a) + (b^3*d^2 + 2*a*b^2*d*e - 3*a^
2*b*e^2)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a
*b*e)))/(sqrt(b*e)*e))*A*d*e*abs(b)/b^3 - 16*(sqrt(b^2*d + (b*x + a)*b*e -
a*b*e)*(2*(4*b*x + 4*a + (b*d*e^3 - 13*a*e^4)/e^4)*(b*x + a) - 3*(b^2*d^2
*e^2 + 2*a*b*d*e^3 - 11*a^2*e^4)/e^4)*sqrt(b*x + a) - 3*(b^4*d^3 + a*b^3*d
^2*e + 3*a^2*b^2*d*e^2 - 5*a^3*b*e^3)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + s
qrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*e^2))*B*d*e*abs(b)/b^4 ...

```

Mupad [F(-1)]

Timed out.

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{\sqrt{a + bx}} dx = \int \frac{(A + Bx)(d + ex)^{5/2}}{\sqrt{a + bx}} dx$$

input

```
int(((A + B*x)*(d + e*x)^(5/2))/(a + b*x)^(1/2),x)
```

output

```
int(((A + B*x)*(d + e*x)^(5/2))/(a + b*x)^(1/2), x)
```


3.206 $\int \frac{(A+Bx)(d+ex)^{3/2}}{\sqrt{a+bx}} dx$

Optimal result	1882
Mathematica [A] (verified)	1883
Rubi [A] (verified)	1883
Maple [B] (verified)	1885
Fricas [A] (verification not implemented)	1886
Sympy [F]	1887
Maxima [F(-2)]	1887
Giac [B] (verification not implemented)	1888
Mupad [F(-1)]	1889
Reduce [B] (verification not implemented)	1889

Optimal result

Integrand size = 24, antiderivative size = 193

$$\int \frac{(A+Bx)(d+ex)^{3/2}}{\sqrt{a+bx}} dx = -\frac{(bd-ae)(bBd-6Abe+5aBe)\sqrt{a+bx}\sqrt{d+ex}}{8b^3e} - \frac{(bBd-6Abe+5aBe)\sqrt{a+bx}(d+ex)^{3/2}}{12b^2e} + \frac{B\sqrt{a+bx}(d+ex)^{5/2}}{3be} - \frac{(bd-ae)^2(bBd-6Abe+5aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{8b^{7/2}e^{3/2}}$$

output

```
-1/8*(-a*e+b*d)*(-6*A*b*e+5*B*a*e+B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(1/2)/b^3/e
-1/12*(-6*A*b*e+5*B*a*e+B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(3/2)/b^2/e+1/3*B*(b*
x+a)^(1/2)*(e*x+d)^(5/2)/b/e-1/8*(-a*e+b*d)^2*(-6*A*b*e+5*B*a*e+B*b*d)*arc
tanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)^(1/2))/b^(7/2)/e^(3/2)
```

Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.85

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{\sqrt{a + bx}} dx = \frac{\sqrt{a + bx}\sqrt{d + ex}(15a^2Be^2 - 2abe(11Bd + 9Ae + 5Bex) + b^2(6Ae(5d + 2ex) + (bd - ae)^2(bBd - 6Abe + 5aBe))\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{24b^3e} - \frac{8b^{7/2}e^{3/2}}{8b^{7/2}e^{3/2}}$$

input

```
Integrate[((A + B*x)*(d + e*x)^(3/2))/Sqrt[a + b*x],x]
```

output

```
(Sqrt[a + b*x]*Sqrt[d + e*x]*(15*a^2*B*e^2 - 2*a*b*e*(11*B*d + 9*A*e + 5*B*e*x) + b^2*(6*A*e*(5*d + 2*e*x) + B*(3*d^2 + 14*d*e*x + 8*e^2*x^2)))/(24*b^3*e) - ((b*d - a*e)^2*(b*B*d - 6*A*b*e + 5*a*B*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(8*b^(7/2)*e^(3/2))
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.87, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$, Rules used = {90, 60, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{\sqrt{a + bx}} dx$$

↓ 90

$$\frac{B\sqrt{a + bx}(d + ex)^{5/2}}{3be} - \frac{(5aBe - 6Abe + bBd) \int \frac{(d+ex)^{3/2}}{\sqrt{a+bx}} dx}{6be}$$

↓ 60

$$\frac{B\sqrt{a + bx}(d + ex)^{5/2}}{3be} - \frac{(5aBe - 6Abe + bBd) \left(\frac{3(bd-ae) \int \frac{\sqrt{d+ex}}{\sqrt{a+bx}} dx}{4b} + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{6be}$$

↓ 60

$$\frac{(5aBe - 6Abe + bBd) \left(\frac{B\sqrt{a+bx}(d+ex)^{5/2}}{3be} - \frac{3(bd-ae) \left(\frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{2b} + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{4b} + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{6be}$$

↓ 66

$$\frac{(5aBe - 6Abe + bBd) \left(\frac{B\sqrt{a+bx}(d+ex)^{5/2}}{3be} - \frac{3(bd-ae) \left(\frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{4b} + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{4b} + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{6be}$$

↓ 221

$$\frac{(5aBe - 6Abe + bBd) \left(\frac{B\sqrt{a+bx}(d+ex)^{5/2}}{3be} - \frac{3(bd-ae) \left(\frac{(bd-ae) \operatorname{arctanh} \left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}} \right) + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{4b} + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{6be}$$

input `Int[((A + B*x)*(d + e*x)^(3/2))/Sqrt[a + b*x], x]`

output `(B*Sqrt[a + b*x]*(d + e*x)^(5/2))/(3*b*e) - ((b*B*d - 6*A*b*e + 5*a*B*e)*((Sqrt[a + b*x]*(d + e*x)^(3/2))/(2*b) + (3*(b*d - a*e)*((Sqrt[a + b*x]*Sqrt[d + e*x])/b + ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])]))/(b^(3/2)*Sqrt[e])))/(4*b))/(6*b*e)`

Definitions of rubi rules used

rule 60

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(
b*(m + n + 1))) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!Integer
Q[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinear
Q[a, b, c, d, m, n, x]
```

rule 66

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[
2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; Fre
eQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]
```

rule 90

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))),
x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p
+ 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n,
p}, x] && NeQ[n + p + 2, 0]
```

rule 221

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 635 vs. $2(161) = 322$.

Time = 0.26 (sec) , antiderivative size = 636, normalized size of antiderivative = 3.30

method	result
default	$\frac{\sqrt{ex+d}\sqrt{bx+a}\left(16Bb^2e^2x^2\sqrt{(ex+d)(bx+a)}\sqrt{be}+18A\ln\left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}}\right)a^2be^3-36A\ln\left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}}\right)\right)}{\dots}$

input

```
int((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^(1/2),x,method=_RETURNVERBOSE)
```

output

```

1/48*(e*x+d)^(1/2)*(b*x+a)^(1/2)*(16*B*b^2*e^2*x^2*((e*x+d)*(b*x+a))^(1/2)
*(b*e)^(1/2)+18*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*
e+d*b)/(b*e)^(1/2))*a^2*b*e^3-36*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)
*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^2*d*e^2+18*A*ln(1/2*(2*b*e*x+2*((
e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^3*d^2*e+24*A*((e
*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^2*e^2*x-15*B*ln(1/2*(2*b*e*x+2*((e*x+d)
*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^3*e^3+27*B*ln(1/2*(2*b
*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*b*d*e
^2-9*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e
)^(1/2))*a*b^2*d^2*e-3*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(
1/2)+a*e+d*b)/(b*e)^(1/2))*b^3*d^3-20*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)
)*a*b*e^2*x+28*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*b^2*d*e*x-36*A*((e*x+
d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b*e^2+60*A*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(
1/2)*b^2*d*e+30*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)*a^2*e^2-44*B*((e*x+d)
)*(b*x+a))^(1/2)*(b*e)^(1/2)*a*b*d*e+6*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/
2)*b^2*d^2)/b^3/e/((e*x+d)*(b*x+a))^(1/2)/(b*e)^(1/2)

```

Fricas [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 542, normalized size of antiderivative = 2.81

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{\sqrt{a + bx}} dx = \left[-\frac{3(Bb^3d^3 + 3(Bab^2 - 2Ab^3)d^2e - 3(3Ba^2b - 4Aab^2)de^2 + (5Ba^3 - 6Aa^2b)de^3 + (3Aa^2d - 3Aab^2)e^3 + (3Aa^2d - 3Aab^2)e^3 + (3Aa^2d - 3Aab^2)e^3 + (3Aa^2d - 3Aab^2)e^3)}{b^3} \right]$$

input

```
integrate((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^(1/2),x, algorithm="fricas")
```

output

```
[-1/96*(3*(B*b^3*d^3 + 3*(B*a*b^2 - 2*A*b^3)*d^2*e - 3*(3*B*a^2*b - 4*A*a*b^2)*d*e^2 + (5*B*a^3 - 6*A*a^2*b)*e^3)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*b*e*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x + a)*sqrt(e*x + d) + 8*(b^2*d*e + a*b*e^2)*x) - 4*(8*B*b^3*e^3*x^2 + 3*B*b^3*d^2*e - 2*(11*B*a*b^2 - 15*A*b^3)*d*e^2 + 3*(5*B*a^2*b - 6*A*a*b^2)*e^3 + 2*(7*B*b^3*d*e^2 - (5*B*a*b^2 - 6*A*b^3)*e^3)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b^4*e^2), 1/48*(3*(B*b^3*d^3 + 3*(B*a*b^2 - 2*A*b^3)*d^2*e - 3*(3*B*a^2*b - 4*A*a*b^2)*d*e^2 + (5*B*a^3 - 6*A*a^2*b)*e^3)*sqrt(-b*e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*e^2*x^2 + a*b*d*e + (b^2*d*e + a*b*e^2)*x)) + 2*(8*B*b^3*e^3*x^2 + 3*B*b^3*d^2*e - 2*(11*B*a*b^2 - 15*A*b^3)*d*e^2 + 3*(5*B*a^2*b - 6*A*a*b^2)*e^3 + 2*(7*B*b^3*d*e^2 - (5*B*a*b^2 - 6*A*b^3)*e^3)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b^4*e^2)]
```

Sympy [F]

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{\sqrt{a + bx}} dx = \int \frac{(A + Bx)(d + ex)^{\frac{3}{2}}}{\sqrt{a + bx}} dx$$

input

```
integrate((B*x+A)*(e*x+d)**(3/2)/(b*x+a)**(1/2),x)
```

output

```
Integral((A + B*x)*(d + e*x)**(3/2)/sqrt(a + b*x), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{\sqrt{a + bx}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^(1/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for m
ore detail
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 559 vs. $2(161) = 322$.

Time = 0.20 (sec) , antiderivative size = 559, normalized size of antiderivative = 2.90

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{\sqrt{a + bx}} dx = \frac{24 \left(\frac{(b^2 d - a b e) \log \left(\left| \frac{-\sqrt{b e} \sqrt{b x + a} + \sqrt{b^2 d + (b x + a) b e - a b e}}{\sqrt{b e}} \right| - \sqrt{b^2 d + (b x + a) b e - a b e} \sqrt{b x + a} \right)}{\sqrt{b e}} - \sqrt{b^2 d + (b x + a) b e - a b e} \sqrt{b x + a} \right) A d | b|}{b^2} - \frac{6 \left(\sqrt{b^2 d + (b x + a) b e - a b e} \left(2 b x + 2 a + \frac{b d e - 5 a^2}{e^2} \right) \right)}{b^2}$$

input

```
integrate((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^(1/2),x, algorithm="giac")
```

output

```
-1/24*(24*((b^2*d - a*b*e)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d +
(b*x + a)*b*e - a*b*e)))/sqrt(b*e) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*
sqrt(b*x + a))*A*d*abs(b)/b^2 - 6*(sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*
b*x + 2*a + (b*d*e - 5*a*e^2)/e^2)*sqrt(b*x + a) + (b^3*d^2 + 2*a*b^2*d*e
- 3*a^2*b*e^2)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b
*e - a*b*e)))/(sqrt(b*e)*e))*B*d*abs(b)/b^3 - 6*(sqrt(b^2*d + (b*x + a)*b*
e - a*b*e)*(2*b*x + 2*a + (b*d*e - 5*a*e^2)/e^2)*sqrt(b*x + a) + (b^3*d^2
+ 2*a*b^2*d*e - 3*a^2*b*e^2)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d
+ (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*e))*A*e*abs(b)/b^3 - (sqrt(b^2*d +
(b*x + a)*b*e - a*b*e)*(2*(4*b*x + 4*a + (b*d*e^3 - 13*a*e^4)/e^4)*(b*x +
a) - 3*(b^2*d^2*e^2 + 2*a*b*d*e^3 - 11*a^2*e^4)/e^4)*sqrt(b*x + a) - 3*(b^
4*d^3 + a*b^3*d^2*e + 3*a^2*b^2*d*e^2 - 5*a^3*b*e^3)*log(abs(-sqrt(b*e)*sq
rt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*e^2))*B*e*a
bs(b)/b^4)/b
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{\sqrt{a + bx}} dx = \int \frac{(A + Bx)(d + ex)^{3/2}}{\sqrt{a + bx}} dx$$

input `int(((A + B*x)*(d + e*x)^(3/2))/(a + b*x)^(1/2),x)`

output `int(((A + B*x)*(d + e*x)^(3/2))/(a + b*x)^(1/2), x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 320, normalized size of antiderivative = 1.66

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{\sqrt{a + bx}} dx = \frac{-3\sqrt{ex + d}\sqrt{bx + a}a^2be^3 + 8\sqrt{ex + d}\sqrt{bx + a}ab^2de^2 + 2\sqrt{ex + d}\sqrt{bx + a}}{\sqrt{a + bx}}$$

input `int((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^(1/2),x)`

output `(- 3*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b*e**3 + 8*sqrt(d + e*x)*sqrt(a + b*x)*a*b**2*d*e**2 + 2*sqrt(d + e*x)*sqrt(a + b*x)*a*b**2*e**3*x + 3*sqrt(d + e*x)*sqrt(a + b*x)*b**3*d**2*e + 14*sqrt(d + e*x)*sqrt(a + b*x)*b**3*d*e**2*x + 8*sqrt(d + e*x)*sqrt(a + b*x)*b**3*e**3*x**2 + 3*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**3*e**3 - 9*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**2*b*d*e**2 + 9*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*b**2*d**2*e - 3*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**3*d**3)/(24*b**3*e**2)`

3.207 $\int \frac{(A+Bx)\sqrt{d+ex}}{\sqrt{a+bx}} dx$

Optimal result	1890
Mathematica [A] (verified)	1891
Rubi [A] (verified)	1891
Maple [B] (verified)	1893
Fricas [A] (verification not implemented)	1894
Sympy [F]	1894
Maxima [F(-2)]	1895
Giac [B] (verification not implemented)	1895
Mupad [B] (verification not implemented)	1896
Reduce [B] (verification not implemented)	1896

Optimal result

Integrand size = 24, antiderivative size = 140

$$\int \frac{(A+Bx)\sqrt{d+ex}}{\sqrt{a+bx}} dx = -\frac{(bBd - 4Abe + 3aBe)\sqrt{a+bx}\sqrt{d+ex}}{4b^2e} + \frac{B\sqrt{a+bx}(d+ex)^{3/2}}{2be} - \frac{(bd - ae)(bBd - 4Abe + 3aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{4b^{5/2}e^{3/2}}$$

output

```
-1/4*(-4*A*b*e+3*B*a*e+B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(1/2)/b^2/e+1/2*B*(b*x+a)^(1/2)*(e*x+d)^(3/2)/b/e-1/4*(-a*e+b*d)*(-4*A*b*e+3*B*a*e+B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)^(1/2))/b^(5/2)/e^(3/2)
```

Mathematica [A] (verified)

Time = 0.65 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.96

$$\int \frac{(A + Bx)\sqrt{d + ex}}{\sqrt{a + bx}} dx$$

$$= \frac{\sqrt{a + bx}\sqrt{d + ex}(4Abe - 3aBe + bB(d + 2ex))}{4b^2e}$$

$$+ \frac{(bd - ae)(bBd - 4Abe + 3aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{e}\left(\sqrt{a-\frac{bd}{e}}-\sqrt{a+bx}\right)}\right)}{2b^{5/2}e^{3/2}}$$

input

```
Integrate[((A + B*x)*Sqrt[d + e*x])/Sqrt[a + b*x],x]
```

output

```
(Sqrt[a + b*x]*Sqrt[d + e*x]*(4*A*b*e - 3*a*B*e + b*B*(d + 2*e*x)))/(4*b^2*e) + ((b*d - a*e)*(b*B*d - 4*A*b*e + 3*a*B*e)*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/(Sqrt[e]*(Sqrt[a - (b*d)/e] - Sqrt[a + b*x]))])/(2*b^(5/2)*e^(3/2))
```

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.91, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {90, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)\sqrt{d + ex}}{\sqrt{a + bx}} dx$$

$$\downarrow 90$$

$$\frac{B\sqrt{a + bx}(d + ex)^{3/2}}{2be} - \frac{(3aBe - 4Abe + bBd) \int \frac{\sqrt{d+ex}}{\sqrt{a+bx}} dx}{4be}$$

$$\downarrow 60$$

$$\frac{B\sqrt{a+bx}(d+ex)^{3/2}}{2be} - \frac{(3aBe - 4Abe + bBd) \left(\frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{2b} + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{4be}$$

↓ 66

$$\frac{B\sqrt{a+bx}(d+ex)^{3/2}}{2be} - \frac{(3aBe - 4Abe + bBd) \left(\frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d\frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{b} + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{4be}$$

↓ 221

$$\frac{B\sqrt{a+bx}(d+ex)^{3/2}}{2be} - \frac{(3aBe - 4Abe + bBd) \left(\frac{(bd-ae) \operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{b^{3/2}\sqrt{e}} + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{4be}$$

input `Int[((A + B*x)*Sqrt[d + e*x])/Sqrt[a + b*x], x]`

output `(B*Sqrt[a + b*x]*(d + e*x)^(3/2))/(2*b*e) - ((b*B*d - 4*A*b*e + 3*a*B*e)*(Sqrt[a + b*x]*Sqrt[d + e*x])/b + ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(b^(3/2)*Sqrt[e]))/(4*b*e)`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 374 vs. $2(114) = 228$.

Time = 0.27 (sec) , antiderivative size = 375, normalized size of antiderivative = 2.68

method	result
default	$-\frac{\sqrt{ex+d}\sqrt{bx+a}\left(4A\ln\left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}}\right)abe^2-4A\ln\left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}}\right)b^2de-3B\ln\left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}}\right)\right)}{\dots}$

input `int((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^(1/2),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -1/8*(e*x+d)^{(1/2)}*(b*x+a)^{(1/2)}*(4*A*\ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^{(1/2)}*(b*e)^{(1/2)+a*e+d*b})/(b*e)^{(1/2)})*a*b*e^{-2-4*A*\ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^{(1/2)}*(b*e)^{(1/2)+a*e+d*b})/(b*e)^{(1/2)})} \\ & *b^2*d*e^{-3*B*\ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^{(1/2)}*(b*e)^{(1/2)+a*e+d*b})/(b*e)^{(1/2)})} \\ & *a^2*e^{-2+2*B*\ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^{(1/2)}*(b*e)^{(1/2)+a*e+d*b})/(b*e)^{(1/2)})} \\ & *a*b*d*e+B*\ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^{(1/2)}*(b*e)^{(1/2)+a*e+d*b})/(b*e)^{(1/2)}) \\ & *b^2*d^{-2-4*B*((e*x+d)*(b*x+a))^{(1/2)}*(b*e)^{(1/2)}*b*e*x-8*A*(b*e)^{(1/2)}*((e*x+d)*(b*x+a))^{(1/2)}*b*e+6*B*((e*x+d)*(b*x+a))^{(1/2)}*(b*e)^{(1/2)} \\ & *a*e^{-2*B*(b*e)^{(1/2)}*((e*x+d)*(b*x+a))^{(1/2)}*b*d}/((e*x+d)*(b*x+a))^{(1/2)}/b^2/e/(b*e)^{(1/2)} \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 366, normalized size of antiderivative = 2.61

$$\int \frac{(A + Bx)\sqrt{d + ex}}{\sqrt{a + bx}} dx$$

$$= \left[\frac{(Bb^2d^2 + 2(Bab - 2Ab^2)de - (3Ba^2 - 4Aab)e^2)\sqrt{be} \log\left(8b^2e^2x^2 + b^2d^2 + 6abde + a^2e^2 - 4(2bex + \dots)\right)}{\dots} \right]$$

input `integrate((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^(1/2),x, algorithm="fricas")`

output `[1/16*((B*b^2*d^2 + 2*(B*a*b - 2*A*b^2)*d*e - (3*B*a^2 - 4*A*a*b)*e^2)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 - 4*(2*b*e*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x + a)*sqrt(e*x + d) + 8*(b^2*d*e + a*b*e^2)*x + 4*(2*B*b^2*e^2*x + B*b^2*d*e - (3*B*a*b - 4*A*b^2)*e^2)*sqrt(b*x + a)*sqrt(e*x + d))/(b^3*e^2), 1/8*((B*b^2*d^2 + 2*(B*a*b - 2*A*b^2)*d*e - (3*B*a^2 - 4*A*a*b)*e^2)*sqrt(-b*e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*e^2*x^2 + a*b*d*e + (b^2*d*e + a*b*e^2)*x)) + 2*(2*B*b^2*e^2*x + B*b^2*d*e - (3*B*a*b - 4*A*b^2)*e^2)*sqrt(b*x + a)*sqrt(e*x + d))/(b^3*e^2)]`

Sympy [F]

$$\int \frac{(A + Bx)\sqrt{d + ex}}{\sqrt{a + bx}} dx = \int \frac{(A + Bx)\sqrt{d + ex}}{\sqrt{a + bx}} dx$$

input `integrate((B*x+A)*(e*x+d)**(1/2)/(b*x+a)**(1/2),x)`

output `Integral((A + B*x)*sqrt(d + e*x)/sqrt(a + b*x), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + Bx)\sqrt{d + ex}}{\sqrt{a + bx}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^(1/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 233 vs. 2(114) = 228.

Time = 0.16 (sec) , antiderivative size = 233, normalized size of antiderivative = 1.66

$$\int \frac{(A + Bx)\sqrt{d + ex}}{\sqrt{a + bx}} dx = \frac{4 \left(\frac{(b^2 d - a b e) \log\left(\left| \frac{-\sqrt{b e} \sqrt{b x + a} + \sqrt{b^2 d + (b x + a) b e - a b e}}{\sqrt{b e}} \right| \right) - \sqrt{b^2 d + (b x + a) b e - a b e} \sqrt{b x + a}}{b^2} \right) A |b| - \left(\sqrt{b^2 d + (b x + a) b e - a b e} \left(2 b x + 2 a + \frac{b d e - 5 a e^2}{e^2} \right) \right)}{4 b}$$

input `integrate((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^(1/2),x, algorithm="giac")`

output `-1/4*(4*((b^2*d - a*b*e)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/sqrt(b*e) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*sqrt(b*x + a))*A*abs(b)/b^2 - (sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*(2*b*x + 2*a + (b*d*e - 5*a*e^2)/e^2)*sqrt(b*x + a) + (b^3*d^2 + 2*a*b^2*d*e - 3*a^2*b*e^2)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*e))*B*abs(b)/b^3)/b`

Mupad [B] (verification not implemented)

Time = 15.87 (sec) , antiderivative size = 866, normalized size of antiderivative = 6.19

$$\int \frac{(A + Bx)\sqrt{d + ex}}{\sqrt{a + bx}} dx = \text{Too large to display}$$

input `int(((A + B*x)*(d + e*x)^(1/2))/(a + b*x)^(1/2),x)`

output

```

((((a + b*x)^(1/2) - a^(1/2))^3*((11*B*a^2*e^2)/2 + (7*B*b^2*d^2)/2 + 23*B
*a*b*d*e))/(e^4*((d + e*x)^(1/2) - d^(1/2))^3) + (((a + b*x)^(1/2) - a^(1/
2))*((B*b^3*d^2)/2 - (3*B*a^2*b*e^2)/2 + B*a*b^2*d*e))/(e^5*((d + e*x)^(1/
2) - d^(1/2))) + (((a + b*x)^(1/2) - a^(1/2))^7*((B*b^2*d^2)/2 - (3*B*a^2*
e^2)/2 + B*a*b*d*e))/(b^2*e^2*((d + e*x)^(1/2) - d^(1/2))^7) + (((a + b*x)
^(1/2) - a^(1/2))^5*((11*B*a^2*e^2)/2 + (7*B*b^2*d^2)/2 + 23*B*a*b*d*e))/(
b*e^3*((d + e*x)^(1/2) - d^(1/2))^5) - (a^(1/2)*d^(1/2)*(32*B*a*e + 16*B*b
*d)*((a + b*x)^(1/2) - a^(1/2))^4)/(e^3*((d + e*x)^(1/2) - d^(1/2))^4) - (
8*B*a^(1/2)*d^(3/2)*((a + b*x)^(1/2) - a^(1/2))^6)/(e^2*((d + e*x)^(1/2) -
d^(1/2))^6) - (8*B*a^(1/2)*b^2*d^(3/2)*((a + b*x)^(1/2) - a^(1/2))^2)/(e^
4*((d + e*x)^(1/2) - d^(1/2))^2))/(((a + b*x)^(1/2) - a^(1/2))^8/((d + e*x)
^(1/2) - d^(1/2))^8 + b^4/e^4 - (4*b^3*((a + b*x)^(1/2) - a^(1/2))^2)/(e^
3*((d + e*x)^(1/2) - d^(1/2))^2) + (6*b^2*((a + b*x)^(1/2) - a^(1/2))^4)/(
e^2*((d + e*x)^(1/2) - d^(1/2))^4) - (4*b*((a + b*x)^(1/2) - a^(1/2))^6)/(
e*((d + e*x)^(1/2) - d^(1/2))^6)) + (((2*A*a*e + 2*A*b*d)*((a + b*x)^(1/2)
- a^(1/2)))/(e^2*((d + e*x)^(1/2) - d^(1/2))) + ((2*A*a*e + 2*A*b*d)*((a
+ b*x)^(1/2) - a^(1/2))^3)/(b*e*((d + e*x)^(1/2) - d^(1/2))^3) - (8*A*a^(1
/2)*d^(1/2)*((a + b*x)^(1/2) - a^(1/2))^2)/(e*((d + e*x)^(1/2) - d^(1/2))^
2))/(((a + b*x)^(1/2) - a^(1/2))^4/((d + e*x)^(1/2) - d^(1/2))^4 + b^2/e^2
- (2*b*((a + b*x)^(1/2) - a^(1/2))^2)/(e*((d + e*x)^(1/2) - d^(1/2))^2...

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.40

$$\int \frac{(A + Bx)\sqrt{d + ex}}{\sqrt{a + bx}} dx$$

$$= \frac{\sqrt{ex + d}\sqrt{bx + a}abe^2 + \sqrt{ex + d}\sqrt{bx + a}b^2de + 2\sqrt{ex + d}\sqrt{bx + a}b^2e^2x - \sqrt{e}\sqrt{b}\log\left(\frac{\sqrt{e}\sqrt{bx+a} + \sqrt{b}\sqrt{e}}{\sqrt{ae-bd}}\right)}{4b^2e^2}$$

input `int((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^(1/2),x)`

output `(sqrt(d + e*x)*sqrt(a + b*x)*a*b*e**2 + sqrt(d + e*x)*sqrt(a + b*x)*b**2*d
*e + 2*sqrt(d + e*x)*sqrt(a + b*x)*b**2*e**2*x - sqrt(e)*sqrt(b)*log((sqrt
(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**2*e**2 + 2*
sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a
*e - b*d))*a*b*d*e - sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*
sqrt(d + e*x))/sqrt(a*e - b*d))*b**2*d**2)/(4*b**2*e**2)`

3.208 $\int \frac{A+Bx}{\sqrt{a+bx}\sqrt{d+ex}} dx$

Optimal result	1898
Mathematica [A] (verified)	1898
Rubi [A] (verified)	1899
Maple [B] (verified)	1900
Fricas [A] (verification not implemented)	1900
Sympy [F]	1901
Maxima [F(-2)]	1901
Giac [A] (verification not implemented)	1902
Mupad [B] (verification not implemented)	1902
Reduce [B] (verification not implemented)	1903

Optimal result

Integrand size = 24, antiderivative size = 83

$$\int \frac{A + Bx}{\sqrt{a + bx}\sqrt{d + ex}} dx = \frac{B\sqrt{a + bx}\sqrt{d + ex}}{be} - \frac{(bBd - 2Abe + aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{b^{3/2}e^{3/2}}$$

output

$$B*(b*x+a)^{(1/2)}*(e*x+d)^{(1/2)}/b/e - (-2*A*b*e+B*a*e+B*b*d)*\operatorname{arctanh}(e^{(1/2)}*(b*x+a)^{(1/2)}/b^{(1/2)}/(e*x+d)^{(1/2)})/b^{(3/2)}/e^{(3/2)}$$

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.00

$$\int \frac{A + Bx}{\sqrt{a + bx}\sqrt{d + ex}} dx = \frac{B\sqrt{a + bx}\sqrt{d + ex}}{be} - \frac{(bBd - 2Abe + aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{e}\sqrt{a+bx}}\right)}{b^{3/2}e^{3/2}}$$

input

`Integrate[(A + B*x)/(Sqrt[a + b*x]*Sqrt[d + e*x]),x]`

output

$$(B*\operatorname{Sqrt}[a + b*x]*\operatorname{Sqrt}[d + e*x])/(b*e) - ((b*B*d - 2*A*b*e + a*B*e)*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*\operatorname{Sqrt}[d + e*x])/(\operatorname{Sqrt}[e]*\operatorname{Sqrt}[a + b*x])])/(b^{(3/2)}*e^{(3/2)})$$

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.01, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {90, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{\sqrt{a + bx}\sqrt{d + ex}} dx$$

$$\downarrow 90$$

$$\frac{(2Abe - B(ae + bd)) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{2be} + \frac{B\sqrt{a + bx}\sqrt{d + ex}}{be}$$

$$\downarrow 66$$

$$\frac{(2Abe - B(ae + bd)) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{be} + \frac{B\sqrt{a + bx}\sqrt{d + ex}}{be}$$

$$\downarrow 221$$

$$\frac{(2Abe - B(ae + bd)) \operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{b^{3/2}e^{3/2}} + \frac{B\sqrt{a + bx}\sqrt{d + ex}}{be}$$

input `Int[(A + B*x)/(Sqrt[a + b*x]*Sqrt[d + e*x]), x]`

output `(B*Sqrt[a + b*x]*Sqrt[d + e*x])/(b*e) + ((2*A*b*e - B*(b*d + a*e))*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(b^(3/2)*e^(3/2))`

Defintions of rubi rules used

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`


```
rule 90 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 197 vs. 2(67) = 134.

Time = 0.29 (sec) , antiderivative size = 198, normalized size of antiderivative = 2.39

method	result
default	$\frac{\left(2A \ln\left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}}\right)be - B \ln\left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}}\right)ae - B \ln\left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}}\right)}{2e\sqrt{be}b\sqrt{(ex+d)(bx+a)}}$

```
input int((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/2*(2*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b*e-B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*e-B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b*d+2*B*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2))*(b*x+a)^(1/2)*(e*x+d)^(1/2)/e/(b*e)^(1/2)/b/((e*x+d)*(b*x+a))^(1/2)
```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 251, normalized size of antiderivative = 3.02

$$\int \frac{A + Bx}{\sqrt{a + bx}\sqrt{d + ex}} dx = \frac{4\sqrt{bx+a}\sqrt{ex+d}Bbe - (Bbd + (Ba - 2Ab)e)\sqrt{be} \log\left(8b^2e^2x^2 + b^2d^2 + 6abde + a^2e^2 + 4(2bex + \dots)\right)}{4b^2e^2}$$

input `integrate((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(1/2),x, algorithm="fricas")`

output `[1/4*(4*sqrt(b*x + a)*sqrt(e*x + d)*B*b*e - (B*b*d + (B*a - 2*A*b)*e)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*b*e*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x + a)*sqrt(e*x + d) + 8*(b^2*d*e + a*b*e^2)*x))/ (b^2*e^2), 1/2*(2*sqrt(b*x + a)*sqrt(e*x + d)*B*b*e + (B*b*d + (B*a - 2*A*b)*e)*sqrt(-b*e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*e^2*x^2 + a*b*d*e + (b^2*d*e + a*b*e^2)*x)))/(b^2*e^2)
]`

Sympy [F]

$$\int \frac{A + Bx}{\sqrt{a + bx}\sqrt{d + ex}} dx = \int \frac{A + Bx}{\sqrt{a + bx}\sqrt{d + ex}} dx$$

input `integrate((B*x+A)/(b*x+a)**(1/2)/(e*x+d)**(1/2),x)`

output `Integral((A + B*x)/(sqrt(a + b*x)*sqrt(d + e*x)), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{\sqrt{a + bx}\sqrt{d + ex}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(1/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.29

$$\int \frac{A + Bx}{\sqrt{a + bx}\sqrt{d + ex}} dx$$

$$= \frac{b \left(\frac{(Bbd + Bae - 2Abe) \log\left(\left| \frac{-\sqrt{be}\sqrt{bx+a} + \sqrt{b^2d + (bx+a)be - abe}}{\sqrt{be}} \right|\right) + \frac{\sqrt{b^2d + (bx+a)be - abe}\sqrt{bx+a}B}{b^2e}}{\sqrt{be}} \right)}{|b|}$$

input `integrate((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(1/2),x, algorithm="giac")`output `b*((B*b*d + B*a*e - 2*A*b*e)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*b*e) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*sqrt(b*x + a)*B/(b^2*e))/abs(b)`**Mupad [B] (verification not implemented)**

Time = 3.85 (sec) , antiderivative size = 311, normalized size of antiderivative = 3.75

$$\int \frac{A + Bx}{\sqrt{a + bx}\sqrt{d + ex}} dx$$

$$= \frac{\frac{(2Bae + 2Bbd)(\sqrt{a+bx}-\sqrt{a})}{e^3(\sqrt{d+ex}-\sqrt{d})} + \frac{(2Bae + 2Bbd)(\sqrt{a+bx}-\sqrt{a})^3}{be^2(\sqrt{d+ex}-\sqrt{d})^3} - \frac{8B\sqrt{a}\sqrt{d}(\sqrt{a+bx}-\sqrt{a})^2}{e^2(\sqrt{d+ex}-\sqrt{d})^2}}{\frac{(\sqrt{a+bx}-\sqrt{a})^4}{(\sqrt{d+ex}-\sqrt{d})^4} + \frac{b^2}{e^2} - \frac{2b(\sqrt{a+bx}-\sqrt{a})^2}{e(\sqrt{d+ex}-\sqrt{d})^2}}$$

$$- \frac{4A \operatorname{atan}\left(\frac{b(\sqrt{d+ex}-\sqrt{d})}{\sqrt{-be}(\sqrt{a+bx}-\sqrt{a})}\right)}{\sqrt{-be}} - \frac{2B \operatorname{atanh}\left(\frac{\sqrt{e}(\sqrt{a+bx}-\sqrt{a})}{\sqrt{b}(\sqrt{d+ex}-\sqrt{d})}\right)(ae + bd)}{b^{3/2}e^{3/2}}$$

input `int((A + B*x)/((a + b*x)^(1/2)*(d + e*x)^(1/2)),x)`

output

```
((2*B*a*e + 2*B*b*d)*((a + b*x)^(1/2) - a^(1/2)))/(e^3*((d + e*x)^(1/2) - d^(1/2))) + ((2*B*a*e + 2*B*b*d)*((a + b*x)^(1/2) - a^(1/2))^3)/(b*e^2*((d + e*x)^(1/2) - d^(1/2))^3) - (8*B*a^(1/2)*d^(1/2)*((a + b*x)^(1/2) - a^(1/2))^2)/(e^2*((d + e*x)^(1/2) - d^(1/2))^2)/(((a + b*x)^(1/2) - a^(1/2))^4/((d + e*x)^(1/2) - d^(1/2))^4 + b^2/e^2 - (2*b*((a + b*x)^(1/2) - a^(1/2))^2)/(e*((d + e*x)^(1/2) - d^(1/2))^2)) - (4*A*atan((b*((d + e*x)^(1/2) - d^(1/2)))/((-b*e)^(1/2)*((a + b*x)^(1/2) - a^(1/2)))))/((-b*e)^(1/2) - (2*B*atanh((e^(1/2)*((a + b*x)^(1/2) - a^(1/2)))/(b^(1/2)*((d + e*x)^(1/2) - d^(1/2))))*(a*e + b*d))/(b^(3/2)*e^(3/2))
```

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.23

$$\int \frac{A + Bx}{\sqrt{a + bx}\sqrt{d + ex}} dx$$

$$= \frac{\sqrt{ex + d}\sqrt{bx + a}be + \sqrt{e}\sqrt{b}\log\left(\frac{\sqrt{e}\sqrt{bx+a} + \sqrt{b}\sqrt{ex+d}}{\sqrt{ae-bd}}\right)ae - \sqrt{e}\sqrt{b}\log\left(\frac{\sqrt{e}\sqrt{bx+a} + \sqrt{b}\sqrt{ex+d}}{\sqrt{ae-bd}}\right)bd}{be^2}$$

input

```
int((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(1/2),x)
```

output

```
(sqrt(d + e*x)*sqrt(a + b*x)*b*e + sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*e - sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b*d)/(b*e**2)
```

3.209 $\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{3/2}} dx$

Optimal result	1904
Mathematica [A] (verified)	1904
Rubi [A] (verified)	1905
Maple [B] (verified)	1906
Fricas [B] (verification not implemented)	1907
Sympy [F]	1907
Maxima [F(-2)]	1908
Giac [A] (verification not implemented)	1908
Mupad [F(-1)]	1909
Reduce [B] (verification not implemented)	1909

Optimal result

Integrand size = 24, antiderivative size = 85

$$\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{3/2}} dx = -\frac{2(Bd-Ae)\sqrt{a+bx}}{e(bd-ae)\sqrt{d+ex}} + \frac{2B\text{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{\sqrt{be}^{3/2}}$$

output

```
-2*(-A*e+B*d)*(b*x+a)^(1/2)/e/(-a*e+b*d)/(e*x+d)^(1/2)+2*B*arctanh(e^(1/2)
*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)^(1/2))/b^(1/2)/e^(3/2)
```

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00

$$\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{3/2}} dx = -\frac{2(-Bd+ Ae)\sqrt{a+bx}}{e(-bd+ ae)\sqrt{d+ex}} + \frac{2B\text{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{\sqrt{be}^{3/2}}$$

input

```
Integrate[(A + B*x)/(Sqrt[a + b*x]*(d + e*x)^(3/2)),x]
```

output

```
(-2*(-(B*d) + A*e)*Sqrt[a + b*x])/(e*(-(b*d) + a*e)*Sqrt[d + e*x]) + (2*B*
ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(Sqrt[b]*e^(3/2)
)
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {87, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{3/2}} dx$$

$$\downarrow 87$$

$$\frac{B \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{e} - \frac{2\sqrt{a + bx}(Bd - Ae)}{e\sqrt{d + ex}(bd - ae)}$$

$$\downarrow 66$$

$$\frac{2B \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d\frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{e} - \frac{2\sqrt{a + bx}(Bd - Ae)}{e\sqrt{d + ex}(bd - ae)}$$

$$\downarrow 221$$

$$\frac{2B \operatorname{Arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{\sqrt{b}e^{3/2}} - \frac{2\sqrt{a + bx}(Bd - Ae)}{e\sqrt{d + ex}(bd - ae)}$$

input `Int[(A + B*x)/(Sqrt[a + b*x]*(d + e*x)^(3/2)),x]`

output `(-2*(B*d - A*e)*Sqrt[a + b*x])/(e*(b*d - a*e)*Sqrt[d + e*x]) + (2*B*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(Sqrt[b]*e^(3/2))`

Defintions of rubi rules used

```
rule 66 Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[
2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; Fre
eQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 277 vs. 2(69) = 138.

Time = 0.30 (sec) , antiderivative size = 278, normalized size of antiderivative = 3.27

method	result
default	$\frac{\left(B \ln \left(\frac{2be x + 2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) a e^2 x - B \ln \left(\frac{2be x + 2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) b d e x + B \ln \left(\frac{2be x + 2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) \sqrt{be} (ae - db) \sqrt{(ex+d)(bx+a)}}{\sqrt{be} (ae - db) \sqrt{(ex+d)(bx+a)}} \right)}{\sqrt{be} (ae - db) \sqrt{(ex+d)(bx+a)}}$

```
input int((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(3/2),x,method=_RETURNVERBOSE)
```

```
output (B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1
/2))*a*e^2*x-B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d
*b)/(b*e)^(1/2))*b*d*e*x+B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)
^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*d*e-B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(
1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b*d^2-2*A*e*(b*e)^(1/2)*((e*x+d)*(b
*x+a))^(1/2)+2*B*d*(b*e)^(1/2)*((e*x+d)*(b*x+a))^(1/2)*(b*x+a)^(1/2)/(b*e
)^(1/2)/(a*e-b*d)/((e*x+d)*(b*x+a))^(1/2)/e/(e*x+d)^(1/2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 174 vs. $2(69) = 138$.

Time = 0.31 (sec) , antiderivative size = 362, normalized size of antiderivative = 4.26

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{3/2}} dx = \frac{\left[(Bbd^2 - Bade + (Bbde - Bae^2)x)\sqrt{be} \log \left(8b^2e^2x^2 + b^2d^2 + 6abde + a^2e^2 \right) \right.}{2(b^2d^2e^2 - abde^3 + (b^2de^3 - abe^4)x)} + \frac{(Bbd^2 - Bade + (Bbde - Bae^2)x)\sqrt{-be} \arctan \left(\frac{(2bex + bd + ae)\sqrt{-be}\sqrt{bx+a}\sqrt{ex+d}}{2(b^2e^2x^2 + abde + (b^2de + abe^2)x)} \right) + 2(Bbde - Abe^2)\sqrt{bx+a}}{b^2d^2e^2 - abde^3 + (b^2de^3 - abe^4)x}$$

input `integrate((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(3/2),x, algorithm="fricas")`

output `[1/2*((B*b*d^2 - B*a*d*e + (B*b*d*e - B*a*e^2)*x)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*b*e*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x + a)*sqrt(e*x + d) + 8*(b^2*d*e + a*b*e^2)*x) - 4*(B*b*d*e - A*b*e^2)*sqrt(b*x + a)*sqrt(e*x + d))/(b^2*d^2*e^2 - a*b*d*e^3 + (b^2*d*e^3 - a*b*e^4)*x), -((B*b*d^2 - B*a*d*e + (B*b*d*e - B*a*e^2)*x)*sqrt(-b*e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*e^2*x^2 + a*b*d*e + (b^2*d*e + a*b*e^2)*x)) + 2*(B*b*d*e - A*b*e^2)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*d^2*e^2 - a*b*d*e^3 + (b^2*d*e^3 - a*b*e^4)*x)]`

Sympy [F]

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{3/2}} dx = \int \frac{A + Bx}{\sqrt{a + bx} (d + ex)^{\frac{3}{2}}} dx$$

input `integrate((B*x+A)/(b*x+a)**(1/2)/(e*x+d)**(3/2),x)`

output `Integral((A + B*x)/(sqrt(a + b*x)*(d + e*x)**(3/2)), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{3/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(3/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

Giac [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.42

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{3/2}} dx =$$

$$\frac{2 B|b| \log \left(\left| -\sqrt{be}\sqrt{bx + a} + \sqrt{b^2d + (bx + a)be - abe} \right| \right)}{\sqrt{bebe}}$$

$$- \frac{2 (Bb^2d|b| - Ab^2e|b|)\sqrt{bx + a}}{(b^3de - ab^2e^2)\sqrt{b^2d + (bx + a)be - abe}}$$

input `integrate((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(3/2),x, algorithm="giac")`

output `-2*B*abs(b)*log(abs(-sqrt(b*e)*sqrt(b*x + a) + sqrt(b^2*d + (b*x + a)*b*e - a*b*e)))/(sqrt(b*e)*b*e) - 2*(B*b^2*d*abs(b) - A*b^2*e*abs(b))*sqrt(b*x + a)/((b^3*d*e - a*b^2*e^2)*sqrt(b^2*d + (b*x + a)*b*e - a*b*e))`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{3/2}} dx = \int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{3/2}} dx$$

input `int((A + B*x)/((a + b*x)^(1/2)*(d + e*x)^(3/2)),x)`

output `int((A + B*x)/((a + b*x)^(1/2)*(d + e*x)^(3/2)), x)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.41

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{3/2}} dx = \frac{-2\sqrt{ex + d}\sqrt{bx + a}e + 2\sqrt{e}\sqrt{b}\log\left(\frac{\sqrt{e}\sqrt{bx+a} + \sqrt{b}\sqrt{ex+d}}{\sqrt{ae-bd}}\right)d + 2\sqrt{e}\sqrt{b}\log\left(\frac{\sqrt{e}\sqrt{bx+a} + \sqrt{b}\sqrt{ex+d}}{\sqrt{ae-bd}}\right)e*x - \sqrt{e}\sqrt{b}d - \sqrt{e}\sqrt{b}e*x}{e^2(ex + d)}$$

input `int((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(3/2),x)`

output `(2*(- sqrt(d + e*x)*sqrt(a + b*x)*e + sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*d + sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*e*x - sqrt(e)*sqrt(b)*d - sqrt(e)*sqrt(b)*e*x)/(e**2*(d + e*x))`

3.210 $\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{5/2}} dx$

Optimal result	1910
Mathematica [A] (verified)	1910
Rubi [A] (verified)	1911
Maple [A] (verified)	1912
Fricas [A] (verification not implemented)	1913
Sympy [F]	1913
Maxima [F(-2)]	1913
Giac [B] (verification not implemented)	1914
Mupad [B] (verification not implemented)	1914
Reduce [B] (verification not implemented)	1915

Optimal result

Integrand size = 24, antiderivative size = 94

$$\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{5/2}} dx = -\frac{2(Bd-Ae)\sqrt{a+bx}}{3e(bd-ae)(d+ex)^{3/2}} + \frac{2(bBd+2Abe-3aBe)\sqrt{a+bx}}{3e(bd-ae)^2\sqrt{d+ex}}$$

```
output -2/3*(-A*e+B*d)*(b*x+a)^(1/2)/e/(-a*e+b*d)/(e*x+d)^(3/2)+2/3*(2*A*b*e-3*B*
a*e+B*b*d)*(b*x+a)^(1/2)/e/(-a*e+b*d)^2/(e*x+d)^(1/2)
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.69

$$\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{5/2}} dx = \frac{2\sqrt{a+bx}(B(-2ad+bdx-3aex)+A(3bd-ae+2bex))}{3(bd-ae)^2(d+ex)^{3/2}}$$

```
input Integrate[(A + B*x)/(Sqrt[a + b*x]*(d + e*x)^(5/2)),x]
```

```
output (2*Sqrt[a + b*x]*(B*(-2*a*d + b*d*x - 3*a*e*x) + A*(3*b*d - a*e + 2*b*e*x))
)/ (3*(b*d - a*e)^2*(d + e*x)^(3/2))
```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {87, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{5/2}} dx$$

$$\downarrow 87$$

$$\frac{(-3aBe + 2Abe + bBd) \int \frac{1}{\sqrt{a+bx}(d+ex)^{3/2}} dx}{3e(bd - ae)} - \frac{2\sqrt{a + bx}(Bd - Ae)}{3e(d + ex)^{3/2}(bd - ae)}$$

$$\downarrow 48$$

$$\frac{2\sqrt{a + bx}(-3aBe + 2Abe + bBd)}{3e\sqrt{d + ex}(bd - ae)^2} - \frac{2\sqrt{a + bx}(Bd - Ae)}{3e(d + ex)^{3/2}(bd - ae)}$$

input `Int[(A + B*x)/(Sqrt[a + b*x]*(d + e*x)^(5/2)),x]`

output `(-2*(B*d - A*e)*Sqrt[a + b*x])/(3*e*(b*d - a*e)*(d + e*x)^(3/2)) + (2*(b*B*d + 2*A*b*e - 3*a*B*e)*Sqrt[a + b*x])/(3*e*(b*d - a*e)^2*Sqrt[d + e*x])`

Definitions of rubi rules used

rule 48

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

rule 87

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e)), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.64

method	result	size
default	$-\frac{2\sqrt{bx+a}(-2Abex+3Baex-Bbdx+Aae-3Abd+2Bad)}{3(ex+d)^{\frac{3}{2}}(ae-db)^2}$	60
gosper	$-\frac{2\sqrt{bx+a}(-2Abex+3Baex-Bbdx+Aae-3Abd+2Bad)}{3(ex+d)^{\frac{3}{2}}(a^2e^2-2abde+b^2d^2)}$	73
orering	$-\frac{2\sqrt{bx+a}(-2Abex+3Baex-Bbdx+Aae-3Abd+2Bad)}{3(ex+d)^{\frac{3}{2}}(a^2e^2-2abde+b^2d^2)}$	73

input

```
int((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(5/2),x,method=_RETURNVERBOSE)
```

output

```
-2/3*(b*x+a)^(1/2)*(-2*A*b*e*x+3*B*a*e*x-B*b*d*x+A*a*e-3*A*b*d+2*B*a*d)/(e
*x+d)^(3/2)/(a*e-b*d)^2
```

Fricas [A] (verification not implemented)

Time = 0.59 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.49

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{5/2}} dx = \frac{2(Aae + (2Ba - 3Ab)d - (Bbd - (3Ba - 2Ab)e)x)\sqrt{bx + a}\sqrt{ex + d}}{3(b^2d^4 - 2abd^3e + a^2d^2e^2 + (b^2d^2e^2 - 2abde^3 + a^2e^4)x^2 + 2(b^2d^3e - 2abd^2e^2 + a^2de^3)x)}$$

input `integrate((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(5/2),x, algorithm="fricas")`

output `-2/3*(A*a*e + (2*B*a - 3*A*b)*d - (B*b*d - (3*B*a - 2*A*b)*e)*x)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*d^4 - 2*a*b*d^3*e + a^2*d^2*e^2 + (b^2*d^2*e^2 - 2*a*b*d*e^3 + a^2*e^4)*x^2 + 2*(b^2*d^3*e - 2*a*b*d^2*e^2 + a^2*d*e^3)*x)`

Sympy [F]

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{5/2}} dx = \int \frac{A + Bx}{\sqrt{a + bx} (d + ex)^{\frac{5}{2}}} dx$$

input `integrate((B*x+A)/(b*x+a)**(1/2)/(e*x+d)**(5/2),x)`

output `Integral((A + B*x)/(sqrt(a + b*x)*(d + e*x)**(5/2)), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(5/2),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f
or more de
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 180 vs. $2(82) = 164$.

Time = 0.17 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.91

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{5/2}} dx = \frac{2\sqrt{bx + a} \left(\frac{(Bb^4de|b| - 3Bab^3e^2|b| + 2Ab^4e^2|b|)(bx + a)}{b^4d^2e - 2ab^3de^2 + a^2b^2e^3} - \frac{3(Bab^4de|b| - Ab^5de|b| - Ba^2b^3e^2|b| + Aab^4e^2|b|)}{b^4d^2e - 2ab^3de^2 + a^2b^2e^3} \right)}{3(b^2d + (bx + a)be - abe)^{3/2}}$$

input

```
integrate((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(5/2),x, algorithm="giac")
```

output

```
2/3*sqrt(b*x + a)*((B*b^4*d*e*abs(b) - 3*B*a*b^3*e^2*abs(b) + 2*A*b^4*e^2*
abs(b))*(b*x + a)/(b^4*d^2*e - 2*a*b^3*d*e^2 + a^2*b^2*e^3) - 3*(B*a*b^4*d
*e*abs(b) - A*b^5*d*e*abs(b) - B*a^2*b^3*e^2*abs(b) + A*a*b^4*e^2*abs(b))/
(b^4*d^2*e - 2*a*b^3*d*e^2 + a^2*b^2*e^3))/(b^2*d + (b*x + a)*b*e - a*b*e)
^(3/2)
```

Mupad [B] (verification not implemented)

Time = 1.50 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.80

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{5/2}} dx = \frac{\sqrt{d + ex} \left(\frac{x(6Ab^2d - 6Ba^2e + 2Aabe - 2Babd)}{3e^2(ae - bd)^2} - \frac{2Aa^2e + 4Ba^2d - 6Aabd}{3e^2(ae - bd)^2} + \frac{x^2(4Ab^2e + 2Aab^2)}{3e^2(ae - bd)^2} \right)}{x^2\sqrt{a + bx} + \frac{d^2\sqrt{a + bx}}{e^2} + \frac{2dx\sqrt{a + bx}}{e}}$$

input

```
int((A + B*x)/((a + b*x)^(1/2)*(d + e*x)^(5/2)),x)
```

output

```
((d + e*x)^(1/2)*((x*(6*A*b^2*d - 6*B*a^2*e + 2*A*a*b*e - 2*B*a*b*d))/(3*e^2*(a*e - b*d)^2) - (2*A*a^2*e + 4*B*a^2*d - 6*A*a*b*d)/(3*e^2*(a*e - b*d)^2) + (x^2*(4*A*b^2*e + 2*B*b^2*d - 6*B*a*b*e))/(3*e^2*(a*e - b*d)^2)))/(x^2*(a + b*x)^(1/2) + (d^2*(a + b*x)^(1/2))/e^2 + (2*d*x*(a + b*x)^(1/2))/e)
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.33

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{5/2}} dx = \frac{-\frac{2\sqrt{ex+d}\sqrt{bx+ae^2}}{3} - \frac{2\sqrt{ex+d}\sqrt{bx+abe^2x}}{3} - \frac{2\sqrt{e}\sqrt{b}bd^2}{3} - \frac{4\sqrt{e}\sqrt{b}bde^2x}{3} - \frac{2\sqrt{e}\sqrt{b}be^2x^2}{3}}{e^2(ae^3x^2 - bde^2x^2 + 2ade^2x - 2bd^2ex + ad^2e - bd^3)}$$

input

```
int((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(5/2),x)
```

output

```
(2*( - sqrt(d + e*x)*sqrt(a + b*x)*a*e**2 - sqrt(d + e*x)*sqrt(a + b*x)*b*e**2*x - sqrt(e)*sqrt(b)*b*d**2 - 2*sqrt(e)*sqrt(b)*b*d*e*x - sqrt(e)*sqrt(b)*b*e**2*x**2))/(3*e**2*(a*d**2*e + 2*a*d*e**2*x + a*e**3*x**2 - b*d**3 - 2*b*d**2*e*x - b*d*e**2*x**2))
```


3.211 $\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{7/2}} dx$

Optimal result	1916
Mathematica [A] (verified)	1916
Rubi [A] (verified)	1917
Maple [A] (verified)	1918
Fricas [B] (verification not implemented)	1919
Sympy [F]	1919
Maxima [F(-2)]	1920
Giac [B] (verification not implemented)	1920
Mupad [B] (verification not implemented)	1921
Reduce [B] (verification not implemented)	1921

Optimal result

Integrand size = 24, antiderivative size = 145

$$\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{7/2}} dx = -\frac{2(Bd-Ae)\sqrt{a+bx}}{5e(bd-ae)(d+ex)^{5/2}} + \frac{2(bBd+4Abe-5aBe)\sqrt{a+bx}}{15e(bd-ae)^2(d+ex)^{3/2}} + \frac{4b(bBd+4Abe-5aBe)\sqrt{a+bx}}{15e(bd-ae)^3\sqrt{d+ex}}$$

output

```
-2/5*(-A*e+B*d)*(b*x+a)^(1/2)/e/(-a*e+b*d)/(e*x+d)^(5/2)+2/15*(4*A*b*e-5*B
*a*e+B*b*d)*(b*x+a)^(1/2)/e/(-a*e+b*d)^2/(e*x+d)^(3/2)+4/15*b*(4*A*b*e-5*B
*a*e+B*b*d)*(b*x+a)^(1/2)/e/(-a*e+b*d)^3/(e*x+d)^(1/2)
```

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.92

$$\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{7/2}} dx = \frac{2\sqrt{a+bx}\left(15Ab^2 - 15abB - \frac{3Bde(a+bx)^2}{(d+ex)^2} + \frac{3Ae^2(a+bx)^2}{(d+ex)^2} + \frac{5bBd(a+bx)}{d+ex} - \frac{10Abe(a+bx)}{d+ex}\right)}{15(bd-ae)^3\sqrt{d+ex}}$$

input

```
Integrate[(A + B*x)/(Sqrt[a + b*x]*(d + e*x)^(7/2)),x]
```

output

$$(2*\text{Sqrt}[a + b*x]*(15*A*b^2 - 15*a*b*B - (3*B*d*e*(a + b*x)^2)/(d + e*x)^2 + (3*A*e^2*(a + b*x)^2)/(d + e*x)^2 + (5*b*B*d*(a + b*x))/(d + e*x) - (10*A*b*e*(a + b*x))/(d + e*x) + (5*a*B*e*(a + b*x))/(d + e*x))/(15*(b*d - a*e)^3*\text{Sqrt}[d + e*x])$$
Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.98, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {87, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{7/2}} dx$$

$$\downarrow 87$$

$$\frac{(-5aBe + 4Abe + bBd) \int \frac{1}{\sqrt{a+bx}(d+ex)^{5/2}} dx}{5e(bd - ae)} - \frac{2\sqrt{a + bx}(Bd - Ae)}{5e(d + ex)^{5/2}(bd - ae)}$$

$$\downarrow 55$$

$$\frac{(-5aBe + 4Abe + bBd) \left(\frac{2b \int \frac{1}{\sqrt{a+bx}(d+ex)^{3/2}} dx}{3(bd - ae)} + \frac{2\sqrt{a+bx}}{3(d+ex)^{3/2}(bd - ae)} \right)}{5e(bd - ae)} - \frac{2\sqrt{a + bx}(Bd - Ae)}{5e(d + ex)^{5/2}(bd - ae)}$$

$$\downarrow 48$$

$$\frac{\left(\frac{4b\sqrt{a+bx}}{3\sqrt{d+ex}(bd - ae)^2} + \frac{2\sqrt{a+bx}}{3(d+ex)^{3/2}(bd - ae)} \right) (-5aBe + 4Abe + bBd)}{5e(bd - ae)} - \frac{2\sqrt{a + bx}(Bd - Ae)}{5e(d + ex)^{5/2}(bd - ae)}$$

input

$$\text{Int}[(A + B*x)/(\text{Sqrt}[a + b*x]*(d + e*x)^(7/2)), x]$$

output

$$(-2*(B*d - A*e)*\text{Sqrt}[a + b*x])/(5*e*(b*d - a*e)*(d + e*x)^(5/2)) + ((b*B*d + 4*A*b*e - 5*a*B*e)*((2*\text{Sqrt}[a + b*x])/(3*(b*d - a*e)*(d + e*x)^(3/2)) + (4*b*\text{Sqrt}[a + b*x])/(3*(b*d - a*e)^2*\text{Sqrt}[d + e*x]))/(5*e*(b*d - a*e))$$

Defintions of rubi rules used

rule 48 $\text{Int}[(a_.) + (b_.)(x_)^{(m_.)}((c_.) + (d_.)(x_)^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}((c + d*x)^{(n + 1)}((b*c - a*d)*(m + 1))), x] \text{ /; FreeQ}\{a, b, c, d, m, n\}, x] \ \&\& \ \text{EqQ}[m + n + 2, 0] \ \&\& \ \text{NeQ}[m, -1]$

rule 55 $\text{Int}[(a_.) + (b_.)(x_)^{(m_.)}((c_.) + (d_.)(x_)^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}((c + d*x)^{(n + 1)}((b*c - a*d)*(m + 1))), x] - \text{Simp}[d*(\text{Simplify}[m + n + 2]/((b*c - a*d)*(m + 1))) \ \text{Int}[(a + b*x)^{\text{Simplify}[m + 1]}(c + d*x)^n, x], x] \text{ /; FreeQ}\{a, b, c, d, m, n\}, x] \ \&\& \ \text{ILtQ}[\text{Simplify}[m + n + 2], 0] \ \&\& \ \text{NeQ}[m, -1] \ \&\& \ !(\text{LtQ}[m, -1] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ (\text{EqQ}[a, 0] \ || \ (\text{NeQ}[c, 0] \ \&\& \ \text{LtQ}[m - n, 0] \ \&\& \ \text{IntegerQ}[n]))) \ \&\& \ (\text{SumSimplerQ}[m, 1] \ || \ !\text{SumSimplerQ}[n, 1])$

rule 87 $\text{Int}[(a_.) + (b_.)(x_)*((c_.) + (d_.)(x_)^{(n_.)}((e_.) + (f_.)(x_)^{(p_.)}, x_] \rightarrow \text{Simp}[(-b*e - a*f)*(c + d*x)^{(n + 1)}((e + f*x)^{(p + 1)}(f*(p + 1)*(c*f - d*e))), x] - \text{Simp}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1))]/(f*(p + 1)*(c*f - d*e)) \ \text{Int}[(c + d*x)^n*(e + f*x)^{(p + 1)}, x], x] \text{ /; FreeQ}\{a, b, c, d, e, f, n\}, x] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ (!\text{LtQ}[n, -1] \ || \ \text{IntegerQ}[p] \ || \ !(\text{IntegerQ}[n] \ || \ !(\text{EqQ}[e, 0] \ || \ !(\text{EqQ}[c, 0] \ || \ \text{LtQ}[p, n])))$

Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.03

method	result
default	$-\frac{2\sqrt{bx+a}(8Ab^2e^2x^2-10Babe^2x^2+2Bb^2dex^2-4Aabe^2x+20Ab^2dex+5Ba^2e^2x-26Babdex+5b^2Bd^2x+3a^2Ae^2-10Aabde+15(ex+d)^{\frac{5}{2}}(ae-db)^3)}{15(ex+d)^{\frac{5}{2}}(a^3e^3-3a^2be^2d+3ab^2d^2e-b^3d^3)}$
gospers	$-\frac{2\sqrt{bx+a}(8Ab^2e^2x^2-10Babe^2x^2+2Bb^2dex^2-4Aabe^2x+20Ab^2dex+5Ba^2e^2x-26Babdex+5b^2Bd^2x+3a^2Ae^2-10Aabde+15(ex+d)^{\frac{5}{2}}(a^3e^3-3a^2be^2d+3ab^2d^2e-b^3d^3)}{15(ex+d)^{\frac{5}{2}}(a^3e^3-3a^2be^2d+3ab^2d^2e-b^3d^3)}$
orering	$-\frac{2\sqrt{bx+a}(8Ab^2e^2x^2-10Babe^2x^2+2Bb^2dex^2-4Aabe^2x+20Ab^2dex+5Ba^2e^2x-26Babdex+5b^2Bd^2x+3a^2Ae^2-10Aabde+15(ex+d)^{\frac{5}{2}}(a^3e^3-3a^2be^2d+3ab^2d^2e-b^3d^3)}{15(ex+d)^{\frac{5}{2}}(a^3e^3-3a^2be^2d+3ab^2d^2e-b^3d^3)}$

input

$$\text{int}((B*x+A)/(b*x+a)^{(1/2)}/(e*x+d)^{(7/2)}, x, \text{method}=_RETURNVERBOSE)$$

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{7/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(7/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f or more de

Giac [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 377 vs. $2(127) = 254$.

Time = 0.20 (sec) , antiderivative size = 377, normalized size of antiderivative = 2.60

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{7/2}} dx = \frac{2 \left((bx + a) \left(\frac{2(Bb^6de^3|b| - 5Bab^5e^4|b| + 4Ab^6e^4|b|)(bx+a)}{b^5d^3e^2 - 3ab^4d^2e^3 + 3a^2b^3de^4 - a^3b^2e^5} + \frac{5(Bb^7d^2e^2|b| - 6Bab^6de^3|b| + 4Ab^7de^3|b|)}{b^5d^3e^2 - 3ab^4d^2e^3 + 3a^2b^3de^4 - a^3b^2e^5} \right) \right)}{b^5d^3e^2 - 3ab^4d^2e^3 + 3a^2b^3de^4 - a^3b^2e^5}$$

15

input `integrate((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(7/2),x, algorithm="giac")`

output
$$\frac{2}{15} \cdot ((bx + a) \cdot (2 \cdot (B \cdot b^6 \cdot d \cdot e^3 \cdot \text{abs}(b) - 5 \cdot B \cdot a \cdot b^5 \cdot e^4 \cdot \text{abs}(b) + 4 \cdot A \cdot b^6 \cdot e^4 \cdot \text{abs}(b)) \cdot (bx + a) / (b^5 \cdot d^3 \cdot e^2 - 3 \cdot a \cdot b^4 \cdot d^2 \cdot e^3 + 3 \cdot a^2 \cdot b^3 \cdot d \cdot e^4 - a^3 \cdot b^2 \cdot e^5) + 5 \cdot (B \cdot b^7 \cdot d^2 \cdot e^2 \cdot \text{abs}(b) - 6 \cdot B \cdot a \cdot b^6 \cdot d \cdot e^3 \cdot \text{abs}(b) + 4 \cdot A \cdot b^7 \cdot d \cdot e^3 \cdot \text{abs}(b) + 5 \cdot B \cdot a^2 \cdot b^5 \cdot e^4 \cdot \text{abs}(b) - 4 \cdot A \cdot a \cdot b^6 \cdot e^4 \cdot \text{abs}(b)) / (b^5 \cdot d^3 \cdot e^2 - 3 \cdot a \cdot b^4 \cdot d^2 \cdot e^3 + 3 \cdot a^2 \cdot b^3 \cdot d \cdot e^4 - a^3 \cdot b^2 \cdot e^5)) - 15 \cdot (B \cdot a \cdot b^7 \cdot d^2 \cdot e^2 \cdot \text{abs}(b) - A \cdot b^8 \cdot d^2 \cdot e^2 \cdot \text{abs}(b) - 2 \cdot B \cdot a^2 \cdot b^6 \cdot d \cdot e^3 \cdot \text{abs}(b) + 2 \cdot A \cdot a \cdot b^7 \cdot d \cdot e^3 \cdot \text{abs}(b) + B \cdot a^3 \cdot b^5 \cdot e^4 \cdot \text{abs}(b) - A \cdot a^2 \cdot b^6 \cdot e^4 \cdot \text{abs}(b)) / (b^5 \cdot d^3 \cdot e^2 - 3 \cdot a \cdot b^4 \cdot d^2 \cdot e^3 + 3 \cdot a^2 \cdot b^3 \cdot d \cdot e^4 - a^3 \cdot b^2 \cdot e^5)) \cdot \text{sqrt}(bx + a) / (b^2 \cdot d + (bx + a) \cdot b \cdot e - a \cdot b \cdot e)^{(5/2)}$$

Mupad [B] (verification not implemented)

Time = 1.61 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.95

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{7/2}} dx = \frac{\sqrt{d + ex} \left(\frac{4Ba^3de + 6Aa^3e^2 - 20Ba^2bd^2 - 20Aa^2bde + 30Aab^2d^2}{15e^3(ae - bd)^3} + \frac{x(10Ba^3e^2 - 48Ba^2bde - 2Aa^2be^2 - 10Bab^2d^2 + 20Aab^2d^2)}{15e^3(ae - bd)^3} \right)}{x^3 \sqrt{a + bx} + \frac{d^3 \sqrt{a + bx}}{e^3} + \frac{3dx^2 \sqrt{a + bx}}{e} + 3dx}$$

input `int((A + B*x)/((a + b*x)^(1/2)*(d + e*x)^(7/2)),x)`output `-((d + e*x)^(1/2)*((6*A*a^3*e^2 + 4*B*a^3*d*e + 30*A*a*b^2*d^2 - 20*B*a^2*b*d^2 - 20*A*a^2*b*d*e)/(15*e^3*(a*e - b*d)^3) + (x*(30*A*b^3*d^2 + 10*B*a^3*e^2 - 2*A*a^2*b*e^2 - 10*B*a*b^2*d^2 + 20*A*a*b^2*d*e - 48*B*a^2*b*d*e))/(15*e^3*(a*e - b*d)^3) + (4*b^2*x^3*(4*A*b*e - 5*B*a*e + B*b*d))/(15*e^2*(a*e - b*d)^3) + (2*b*x^2*(a*e + 5*b*d)*(4*A*b*e - 5*B*a*e + B*b*d))/(15*e^3*(a*e - b*d)^3)))/(x^3*(a + b*x)^(1/2) + (d^3*(a + b*x)^(1/2))/e^3 + (3*d*x^2*(a + b*x)^(1/2))/e + (3*d^2*x*(a + b*x)^(1/2))/e^2)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 302, normalized size of antiderivative = 2.08

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{7/2}} dx = \frac{-\frac{2\sqrt{ex+d}\sqrt{bx+a}a^2e^3}{5} + \frac{2\sqrt{ex+d}\sqrt{bx+a}abde^2}{3} - \frac{2\sqrt{ex+d}\sqrt{bx+a}abe^3x}{15} + \frac{2\sqrt{ex+d}\sqrt{bx+a}b^2de^2}{3}}{e^2(a^2e^5x^3 - 2abd^2e^4x^3 + b^2d^2e^3x^3 + 3a^2de^4x^2 - 6abd^2e^3x^2 + 3b^2d^2e^2x^2)}$$

input `int((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(7/2),x)`

output

```
(2*( - 3*sqrt(d + e*x)*sqrt(a + b*x)*a**2*e**3 + 5*sqrt(d + e*x)*sqrt(a +
b*x)*a*b*d*e**2 - sqrt(d + e*x)*sqrt(a + b*x)*a*b*e**3*x + 5*sqrt(d + e*x)
*sqrt(a + b*x)*b**2*d*e**2*x + 2*sqrt(d + e*x)*sqrt(a + b*x)*b**2*e**3*x**
2 - 2*sqrt(e)*sqrt(b)*b**2*d**3 - 6*sqrt(e)*sqrt(b)*b**2*d**2*e*x - 6*sqrt
(e)*sqrt(b)*b**2*d*e**2*x**2 - 2*sqrt(e)*sqrt(b)*b**2*e**3*x**3))/(15*e**2
*(a**2*d**3*e**2 + 3*a**2*d**2*e**3*x + 3*a**2*d*e**4*x**2 + a**2*e**5*x**
3 - 2*a*b*d**4*e - 6*a*b*d**3*e**2*x - 6*a*b*d**2*e**3*x**2 - 2*a*b*d*e**4
*x**3 + b**2*d**5 + 3*b**2*d**4*e*x + 3*b**2*d**3*e**2*x**2 + b**2*d**2*e
**3*x**3))
```

3.212 $\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{9/2}} dx$

Optimal result	1923
Mathematica [A] (verified)	1924
Rubi [A] (verified)	1924
Maple [A] (verified)	1926
Fricas [B] (verification not implemented)	1926
Sympy [F]	1927
Maxima [F(-2)]	1927
Giac [B] (verification not implemented)	1928
Mupad [B] (verification not implemented)	1929
Reduce [B] (verification not implemented)	1929

Optimal result

Integrand size = 24, antiderivative size = 198

$$\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{9/2}} dx = -\frac{2(Bd-Ae)\sqrt{a+bx}}{7e(bd-ae)(d+ex)^{7/2}} + \frac{2(bBd+6Abe-7aBe)\sqrt{a+bx}}{35e(bd-ae)^2(d+ex)^{5/2}} + \frac{8b(bBd+6Abe-7aBe)\sqrt{a+bx}}{105e(bd-ae)^3(d+ex)^{3/2}} + \frac{16b^2(bBd+6Abe-7aBe)\sqrt{a+bx}}{105e(bd-ae)^4\sqrt{d+ex}}$$

```
output -2/7*(-A*e+B*d)*(b*x+a)^(1/2)/e/(-a*e+b*d)/(e*x+d)^(7/2)+2/35*(6*A*b*e-7*B
*a*e+B*b*d)*(b*x+a)^(1/2)/e/(-a*e+b*d)^2/(e*x+d)^(5/2)+8/105*b*(6*A*b*e-7*
B*a*e+B*b*d)*(b*x+a)^(1/2)/e/(-a*e+b*d)^3/(e*x+d)^(3/2)+16/105*b^2*(6*A*b*
e-7*B*a*e+B*b*d)*(b*x+a)^(1/2)/e/(-a*e+b*d)^4/(e*x+d)^(1/2)
```


Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.01

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{9/2}} dx = \frac{2\sqrt{a + bx}(15Bde^2(a + bx)^3 - 15Ae^3(a + bx)^3 - 42bBde(a + bx)^2(d + ex) + 63A^2e^2(a + bx)^2(d + ex) - 21a^2Bde^2(a + bx)(d + ex) + 35b^2Bd(a + bx)(d + ex)^2 - 105A^2b^2e^2(a + bx)(d + ex)^2 + 70a^2bBde^2(a + bx)(d + ex)^2 + 105A^2b^3(d + ex)^3 - 105a^2b^2B(d + ex)^3)}{(105(bd - ae)^4(d + ex)^{7/2})}$$

input

```
Integrate[(A + B*x)/(Sqrt[a + b*x]*(d + e*x)^(9/2)),x]
```

output

```
(2*Sqrt[a + b*x]*(15*B*d*e^2*(a + b*x)^3 - 15*A*e^3*(a + b*x)^3 - 42*b*B*d
*e*(a + b*x)^2*(d + e*x) + 63*A*b*e^2*(a + b*x)^2*(d + e*x) - 21*a*B*e^2*(
a + b*x)^2*(d + e*x) + 35*b^2*B*d*(a + b*x)*(d + e*x)^2 - 105*A*b^2*e*(a +
b*x)*(d + e*x)^2 + 70*a*b*B*e*(a + b*x)*(d + e*x)^2 + 105*A*b^3*(d + e*x)
^3 - 105*a*b^2*B*(d + e*x)^3))/(105*(b*d - a*e)^4*(d + e*x)^(7/2))
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 190, normalized size of antiderivative = 0.96,
 number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules
 used = {87, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{9/2}} dx$$

$$\downarrow 87$$

$$\frac{(-7aBe + 6Abe + bBd) \int \frac{1}{\sqrt{a + bx}(d + ex)^{7/2}} dx}{7e(bd - ae)} - \frac{2\sqrt{a + bx}(Bd - Ae)}{7e(d + ex)^{7/2}(bd - ae)}$$

$$\downarrow 55$$

$$\frac{(-7aBe + 6Abe + bBd) \left(\frac{4b \int \frac{1}{\sqrt{a + bx}(d + ex)^{5/2}} dx}{5(bd - ae)} + \frac{2\sqrt{a + bx}}{5(d + ex)^{5/2}(bd - ae)} \right)}{7e(bd - ae)} - \frac{2\sqrt{a + bx}(Bd - Ae)}{7e(d + ex)^{7/2}(bd - ae)}$$

$$\downarrow 55$$

$$\begin{aligned}
 & \frac{(-7aBe + 6Abe + bBd) \left(\frac{4b \left(\frac{2b \int \frac{1}{\sqrt{a+bx}(d+ex)^{3/2}} dx}{3(bd-ae)} + \frac{2\sqrt{a+bx}}{3(d+ex)^{3/2}(bd-ae)} \right)}{5(bd-ae)} + \frac{2\sqrt{a+bx}}{5(d+ex)^{5/2}(bd-ae)} \right)}{\frac{7e(bd-ae)}{2\sqrt{a+bx}(Bd-Ae)} \frac{1}{7e(d+ex)^{7/2}(bd-ae)}}} \\
 & \quad \downarrow 48 \\
 & \frac{\left(\frac{4b \left(\frac{4b\sqrt{a+bx}}{3\sqrt{d+ex}(bd-ae)^2} + \frac{2\sqrt{a+bx}}{3(d+ex)^{3/2}(bd-ae)} \right)}{5(bd-ae)} + \frac{2\sqrt{a+bx}}{5(d+ex)^{5/2}(bd-ae)} \right) (-7aBe + 6Abe + bBd)}{\frac{7e(bd-ae)}{2\sqrt{a+bx}(Bd-Ae)} \frac{1}{7e(d+ex)^{7/2}(bd-ae)}}}
 \end{aligned}$$

input `Int[(A + B*x)/(Sqrt[a + b*x]*(d + e*x)^(9/2)),x]`

output `(-2*(B*d - A*e)*Sqrt[a + b*x])/(7*e*(b*d - a*e)*(d + e*x)^(7/2)) + ((b*B*d + 6*A*b*e - 7*a*B*e)*((2*Sqrt[a + b*x])/(5*(b*d - a*e)*(d + e*x)^(5/2)) + (4*b*((2*Sqrt[a + b*x])/(3*(b*d - a*e)*(d + e*x)^(3/2)) + (4*b*Sqrt[a + b*x])/(3*(b*d - a*e)^2*Sqrt[d + e*x])))/(5*(b*d - a*e)))/(7*e*(b*d - a*e))`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 55 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(Simplify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])`

rule 87

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] :> Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))
```

Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.42

method	result
default	$-\frac{2\sqrt{bx+a}(-48Ab^3e^3x^3+56Bab^2e^3x^3-8Bb^3de^2x^3+24Aab^2e^3x^2-168Ab^3de^2x^2-28Ba^2be^3x^2+200Bab^2de^2x^2-28Bb^3d^2e^2x^2)}{(e*x+d)^{9/2}}$
gospers	$-\frac{2\sqrt{bx+a}(-48Ab^3e^3x^3+56Bab^2e^3x^3-8Bb^3de^2x^3+24Aab^2e^3x^2-168Ab^3de^2x^2-28Ba^2be^3x^2+200Bab^2de^2x^2-28Bb^3d^2e^2x^2)}{(e*x+d)^{9/2}}$
orering	$-\frac{2\sqrt{bx+a}(-48Ab^3e^3x^3+56Bab^2e^3x^3-8Bb^3de^2x^3+24Aab^2e^3x^2-168Ab^3de^2x^2-28Ba^2be^3x^2+200Bab^2de^2x^2-28Bb^3d^2e^2x^2)}{(e*x+d)^{9/2}}$

input

```
int((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(9/2),x,method=_RETURNVERBOSE)
```

output

```
-2/105*(b*x+a)^(1/2)*(-48*A*b^3*e^3*x^3+56*B*a*b^2*e^3*x^3-8*B*b^3*d*e^2*x^3+24*A*a*b^2*e^3*x^2-168*A*b^3*d*e^2*x^2-28*B*a^2*b*e^3*x^2+200*B*a*b^2*d*e^2*x^2-28*B*b^3*d^2*e*x^2-18*A*a^2*b*e^3*x+84*A*a*b^2*d*e^2*x-210*A*b^3*d^2*e*x+21*B*a^3*e^3*x-101*B*a^2*b*d*e^2*x+259*B*a*b^2*d^2*e*x-35*B*b^3*d^3*x+15*A*a^3*e^3-63*A*a^2*b*d*e^2+105*A*a*b^2*d^2*e-105*A*b^3*d^3+6*B*a^3*d*e^2-28*B*a^2*b*d^2*e+70*B*a*b^2*d^3)/(e*x+d)^(7/2)/(a*e-b*d)^4
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 547 vs. 2(174) = 348.

Time = 6.23 (sec) , antiderivative size = 547, normalized size of antiderivative = 2.76

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{9/2}} dx = \frac{2(15Aa^3e^3 + 35(2Bab^2 - 3Ab^3)d^3 - 7(4Ba^2b - 15Aab^2)d^2e + 3(2Ba^3 - 21Aa^2b)de^2 - 8(Bb^3de^2 - 4a^3b^3d^2e^2))}{105(b^4d^8 - 4ab^3d^7e + 6a^2b^2d^6e^2 - 4a^3bd^5e^3 + a^4d^4e^4 + (b^4d^4e^4 - 4ab^3d^3e^5 + 6a^2b^2d^2e^6 - 4a^3bd^2e^7))^{1/2}}$$

input `integrate((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(9/2),x, algorithm="fricas")`

output
$$\begin{aligned} & -2/105*(15*A*a^3*e^3 + 35*(2*B*a*b^2 - 3*A*b^3)*d^3 - 7*(4*B*a^2*b - 15*A* \\ & a*b^2)*d^2*e + 3*(2*B*a^3 - 21*A*a^2*b)*d*e^2 - 8*(B*b^3*d*e^2 - (7*B*a*b^2 \\ & - 6*A*b^3)*e^3)*x^3 - 4*(7*B*b^3*d^2*e - 2*(25*B*a*b^2 - 21*A*b^3)*d*e^2 \\ & + (7*B*a^2*b - 6*A*a*b^2)*e^3)*x^2 - (35*B*b^3*d^3 - 7*(37*B*a*b^2 - 30*A \\ & *b^3)*d^2*e + (101*B*a^2*b - 84*A*a*b^2)*d*e^2 - 3*(7*B*a^3 - 6*A*a^2*b)*e \\ & ^3)*x)*\text{sqrt}(b*x + a)*\text{sqrt}(e*x + d)/(b^4*d^8 - 4*a*b^3*d^7*e + 6*a^2*b^2*d^ \\ & 6*e^2 - 4*a^3*b*d^5*e^3 + a^4*d^4*e^4 + (b^4*d^4*e^4 - 4*a*b^3*d^3*e^5 + 6 \\ & *a^2*b^2*d^2*e^6 - 4*a^3*b*d*e^7 + a^4*e^8)*x^4 + 4*(b^4*d^5*e^3 - 4*a*b^3 \\ & *d^4*e^4 + 6*a^2*b^2*d^3*e^5 - 4*a^3*b*d^2*e^6 + a^4*d*e^7)*x^3 + 6*(b^4*d \\ & ^6*e^2 - 4*a*b^3*d^5*e^3 + 6*a^2*b^2*d^4*e^4 - 4*a^3*b*d^3*e^5 + a^4*d^2*e \\ & ^6)*x^2 + 4*(b^4*d^7*e - 4*a*b^3*d^6*e^2 + 6*a^2*b^2*d^5*e^3 - 4*a^3*b*d^4 \\ & *e^4 + a^4*d^3*e^5)*x \end{aligned}$$

Sympy [F]

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{9/2}} dx = \int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{\frac{9}{2}}} dx$$

input `integrate((B*x+A)/(b*x+a)**(1/2)/(e*x+d)**(9/2),x)`

output `Integral((A + B*x)/(sqrt(a + b*x)*(d + e*x)**(9/2)), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{9/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(9/2),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f
or more de
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 620 vs. $2(174) = 348$.

Time = 0.23 (sec) , antiderivative size = 620, normalized size of antiderivative = 3.13

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{9/2}} dx = \frac{2 \left(\left(4(bx + a) \left(\frac{2(Bb^8de^5|b| - 7Bab^7e^6|b| + 6Ab^8e^6|b|)(bx+a)}{b^6d^4e^3 - 4ab^5d^3e^4 + 6a^2b^4d^2e^5 - 4a^3b^3de^6 + a^4b^2e^7} + \frac{7(Bb^9d^2e^4|b| - 8Bab^8de^5|b| - 7B^2b^8d^2e^4|b| + 6B^2ab^7de^5|b| - 5B^2a^2b^6d^2e^6|b| + 4B^2a^3b^5d^2e^7|b| - 3B^2a^4b^4d^2e^8|b| + 2B^2a^5b^3d^2e^9|b| - B^2a^6b^2d^2e^{10}|b| + B^2a^7b^2d^2e^{11}|b|) \right) \right)}{b^6d^4e^3 - 4ab^5d^3e^4 + 6a^2b^4d^2e^5 - 4a^3b^3de^6 + a^4b^2e^7} + \frac{7(Bb^9d^2e^4|b| - 8Bab^8de^5|b| - 7B^2b^8d^2e^4|b| + 6B^2ab^7de^5|b| - 5B^2a^2b^6d^2e^6|b| + 4B^2a^3b^5d^2e^7|b| - 3B^2a^4b^4d^2e^8|b| + 2B^2a^5b^3d^2e^9|b| - B^2a^6b^2d^2e^{10}|b| + B^2a^7b^2d^2e^{11}|b|)}{b^6d^4e^3 - 4ab^5d^3e^4 + 6a^2b^4d^2e^5 - 4a^3b^3de^6 + a^4b^2e^7}$$

input

```
integrate((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(9/2),x, algorithm="giac")
```

output

```
2/105*((4*(b*x + a)*(2*(B*b^8*d*e^5*abs(b) - 7*B*a*b^7*e^6*abs(b) + 6*A*b^
8*e^6*abs(b))*(b*x + a)/(b^6*d^4*e^3 - 4*a*b^5*d^3*e^4 + 6*a^2*b^4*d^2*e^5
- 4*a^3*b^3*d*e^6 + a^4*b^2*e^7) + 7*(B*b^9*d^2*e^4*abs(b) - 8*B*a*b^8*d*
e^5*abs(b) + 6*A*b^9*d*e^5*abs(b) + 7*B*a^2*b^7*e^6*abs(b) - 6*A*a*b^8*e^6
*abs(b)))/(b^6*d^4*e^3 - 4*a*b^5*d^3*e^4 + 6*a^2*b^4*d^2*e^5 - 4*a^3*b^3*d*
e^6 + a^4*b^2*e^7)) + 35*(B*b^10*d^3*e^3*abs(b) - 9*B*a*b^9*d^2*e^4*abs(b)
+ 6*A*b^10*d^2*e^4*abs(b) + 15*B*a^2*b^8*d*e^5*abs(b) - 12*A*a*b^9*d*e^5*
abs(b) - 7*B*a^3*b^7*e^6*abs(b) + 6*A*a^2*b^8*e^6*abs(b))/(b^6*d^4*e^3 - 4
*a*b^5*d^3*e^4 + 6*a^2*b^4*d^2*e^5 - 4*a^3*b^3*d*e^6 + a^4*b^2*e^7))*(b*x
+ a) - 105*(B*a*b^10*d^3*e^3*abs(b) - A*b^11*d^3*e^3*abs(b) - 3*B*a^2*b^9*
d^2*e^4*abs(b) + 3*A*a*b^10*d^2*e^4*abs(b) + 3*B*a^3*b^8*d*e^5*abs(b) - 3*
A*a^2*b^9*d*e^5*abs(b) - B*a^4*b^7*e^6*abs(b) + A*a^3*b^8*e^6*abs(b))/(b^6
*d^4*e^3 - 4*a*b^5*d^3*e^4 + 6*a^2*b^4*d^2*e^5 - 4*a^3*b^3*d*e^6 + a^4*b^2
*e^7))*sqrt(b*x + a)/(b^2*d + (b*x + a)*b*e - a*b*e)^(7/2)
```

Mupad [B] (verification not implemented)

Time = 1.83 (sec) , antiderivative size = 419, normalized size of antiderivative = 2.12

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{9/2}} dx = \frac{\sqrt{d + ex} \left(\frac{x(-42Ba^4e^3 + 190Ba^3bde^2 + 6Aa^3be^3 - 462Ba^2b^2d^2e - 42Aa^2b^2de^2 - 70Bab^3d^3 + 105e^4(ae - bd)^4}{105e^4(ae - bd)^4} \right)}{105e^4(ae - bd)^4}$$

input

```
int((A + B*x)/((a + b*x)^(1/2)*(d + e*x)^(9/2)),x)
```

output

```
((d + e*x)^(1/2)*((x*(210*A*b^4*d^3 - 42*B*a^4*e^3 + 6*A*a^3*b*e^3 - 70*B*
a*b^3*d^3 - 42*A*a^2*b^2*d*e^2 - 462*B*a^2*b^2*d^2*e + 210*A*a*b^3*d^2*e +
190*B*a^3*b*d*e^2))/(105*e^4*(a*e - b*d)^4) - (30*A*a^4*e^3 - 210*A*a*b^3
*d^3 + 12*B*a^4*d*e^2 + 140*B*a^2*b^2*d^3 + 210*A*a^2*b^2*d^2*e - 126*A*a^
3*b*d*e^2 - 56*B*a^3*b*d^2*e)/(105*e^4*(a*e - b*d)^4) + (16*b^3*x^4*(6*A*b
*e - 7*B*a*e + B*b*d))/(105*e^2*(a*e - b*d)^4) + (8*b^2*x^3*(a*e + 7*b*d)*
(6*A*b*e - 7*B*a*e + B*b*d))/(105*e^3*(a*e - b*d)^4) + (2*b*x^2*(35*b^2*d^
2 - a^2*e^2 + 14*a*b*d*e)*(6*A*b*e - 7*B*a*e + B*b*d))/(105*e^4*(a*e - b*d
)^4)))/(x^4*(a + b*x)^(1/2) + (d^4*(a + b*x)^(1/2))/e^4 + (6*d^2*x^2*(a +
b*x)^(1/2))/e^2 + (4*d*x^3*(a + b*x)^(1/2))/e + (4*d^3*x*(a + b*x)^(1/2))/
e^3)
```

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 544, normalized size of antiderivative = 2.75

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{9/2}} dx = \frac{-\frac{2\sqrt{ex+d}\sqrt{bx+a}a^3e^4}{7} + \frac{4\sqrt{ex+d}\sqrt{bx+a}a^2bde^3}{5} - \frac{2\sqrt{ex+d}\sqrt{bx+a}a^2be^4x}{35} - \frac{2\sqrt{ex+d}\sqrt{bx+a}ab^2e^4}{3}}{e^2(a^3e^7x^4 - 3a^2bde^6x^4 + 3ab^2d^2e^5x^4 - b^3d^3e^4x^4 + 4a^3de^6x^3 - 12a^2bd^2e^5x^3 + 12a^2bd^2e^5x^3 - 12a^2bd^2e^5x^3)}$$

input

```
int((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(9/2),x)
```

output

```
(2*( - 15*sqrt(d + e*x)*sqrt(a + b*x)*a**3*e**4 + 42*sqrt(d + e*x)*sqrt(a
+ b*x)*a**2*b*d*e**3 - 3*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b*e**4*x - 35*sq
rt(d + e*x)*sqrt(a + b*x)*a*b**2*d**2*e**2 + 14*sqrt(d + e*x)*sqrt(a + b*x
)*a*b**2*d*e**3*x + 4*sqrt(d + e*x)*sqrt(a + b*x)*a*b**2*e**4*x**2 - 35*sq
rt(d + e*x)*sqrt(a + b*x)*b**3*d**2*e**2*x - 28*sqrt(d + e*x)*sqrt(a + b*x
)*b**3*d*e**3*x**2 - 8*sqrt(d + e*x)*sqrt(a + b*x)*b**3*e**4*x**3 + 8*sqrt
(e)*sqrt(b)*b**3*d**4 + 32*sqrt(e)*sqrt(b)*b**3*d**3*e*x + 48*sqrt(e)*sqrt
(b)*b**3*d**2*e**2*x**2 + 32*sqrt(e)*sqrt(b)*b**3*d*e**3*x**3 + 8*sqrt(e)*
sqrt(b)*b**3*e**4*x**4))/(105*e**2*(a**3*d**4*e**3 + 4*a**3*d**3*e**4*x +
6*a**3*d**2*e**5*x**2 + 4*a**3*d*e**6*x**3 + a**3*e**7*x**4 - 3*a**2*b*d**
5*e**2 - 12*a**2*b*d**4*e**3*x - 18*a**2*b*d**3*e**4*x**2 - 12*a**2*b*d**2
*e**5*x**3 - 3*a**2*b*d*e**6*x**4 + 3*a*b**2*d**6*e + 12*a*b**2*d**5*e**2*
x + 18*a*b**2*d**4*e**3*x**2 + 12*a*b**2*d**3*e**4*x**3 + 3*a*b**2*d**2*e
**5*x**4 - b**3*d**7 - 4*b**3*d**6*e*x - 6*b**3*d**5*e**2*x**2 - 4*b**3*d**
4*e**3*x**3 - b**3*d**3*e**4*x**4))
```

3.213 $\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{11/2}} dx$

Optimal result	1931
Mathematica [A] (verified)	1932
Rubi [A] (verified)	1932
Maple [B] (verified)	1934
Fricas [B] (verification not implemented)	1935
Sympy [F]	1936
Maxima [F(-2)]	1937
Giac [B] (verification not implemented)	1937
Mupad [B] (verification not implemented)	1938
Reduce [B] (verification not implemented)	1939

Optimal result

Integrand size = 24, antiderivative size = 251

$$\int \frac{A+Bx}{\sqrt{a+bx}(d+ex)^{11/2}} dx = -\frac{2(Bd-Ae)\sqrt{a+bx}}{9e(bd-ae)(d+ex)^{9/2}} + \frac{2(bBd+8Abe-9aBe)\sqrt{a+bx}}{63e(bd-ae)^2(d+ex)^{7/2}} + \frac{4b(bBd+8Abe-9aBe)\sqrt{a+bx}}{105e(bd-ae)^3(d+ex)^{5/2}} + \frac{16b^2(bBd+8Abe-9aBe)\sqrt{a+bx}}{315e(bd-ae)^4(d+ex)^{3/2}} + \frac{32b^3(bBd+8Abe-9aBe)\sqrt{a+bx}}{315e(bd-ae)^5\sqrt{d+ex}}$$

output

```
-2/9*(-A*e+B*d)*(b*x+a)^(1/2)/e/(-a*e+b*d)/(e*x+d)^(9/2)+2/63*(8*A*b*e-9*B
*a*e+B*b*d)*(b*x+a)^(1/2)/e/(-a*e+b*d)^2/(e*x+d)^(7/2)+4/105*b*(8*A*b*e-9*
B*a*e+B*b*d)*(b*x+a)^(1/2)/e/(-a*e+b*d)^3/(e*x+d)^(5/2)+16/315*b^2*(8*A*b*
e-9*B*a*e+B*b*d)*(b*x+a)^(1/2)/e/(-a*e+b*d)^4/(e*x+d)^(3/2)+32/315*b^3*(8*
A*b*e-9*B*a*e+B*b*d)*(b*x+a)^(1/2)/e/(-a*e+b*d)^5/(e*x+d)^(1/2)
```


Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.08

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{11/2}} dx = \frac{2\sqrt{a + bx}(-35Bde^3(a + bx)^4 + 35Ae^4(a + bx)^4 + 135bBde^2(a + bx)^3(d + ex) + 189A^2b^2e^3(a + bx)^3(d + ex) - 189A^2b^2e^3(a + bx)^3(d + ex) + 45a^2B^2e^3(a + bx)^3(d + ex) - 189b^2B^2d^2e^3(a + bx)^2(d + ex)^2 + 378A^2b^2e^2(a + bx)^2(d + ex)^2 - 189a^2B^2e^2(a + bx)^2(d + ex)^2 + 105b^3B^2d^2(a + bx)(d + ex)^3 - 420A^2b^3e^3(a + bx)(d + ex)^3 + 315a^2b^2B^2e^3(a + bx)(d + ex)^3 + 315A^2b^4(d + ex)^4 - 315a^2b^3B^2(d + ex)^4)}{(315*(b*d - a*e)^5*(d + e*x)^(9/2))}$$

input `Integrate[(A + B*x)/(Sqrt[a + b*x]*(d + e*x)^(11/2)),x]`

output $(2*\text{Sqrt}[a + b*x]*(-35*B*d*e^3*(a + b*x)^4 + 35*A*e^4*(a + b*x)^4 + 135*b*B*d*e^2*(a + b*x)^3*(d + e*x) - 180*A*b*e^3*(a + b*x)^3*(d + e*x) + 45*a*B*e^3*(a + b*x)^3*(d + e*x) - 189*b^2*B*d*e*(a + b*x)^2*(d + e*x)^2 + 378*A*b^2*e^2*(a + b*x)^2*(d + e*x)^2 - 189*a*b*B*e^2*(a + b*x)^2*(d + e*x)^2 + 105*b^3*B*d*(a + b*x)*(d + e*x)^3 - 420*A*b^3*e*(a + b*x)*(d + e*x)^3 + 315*a*b^2*B*e*(a + b*x)*(d + e*x)^3 + 315*A*b^4*(d + e*x)^4 - 315*a*b^3*B*(d + e*x)^4)/(315*(b*d - a*e)^5*(d + e*x)^(9/2))$

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 238, normalized size of antiderivative = 0.95, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$, Rules used = {87, 55, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{11/2}} dx$$

$$\downarrow 87$$

$$\frac{(-9aBe + 8Abe + bBd) \int \frac{1}{\sqrt{a+bx}(d+ex)^{9/2}} dx}{9e(bd - ae)} - \frac{2\sqrt{a + bx}(Bd - Ae)}{9e(d + ex)^{9/2}(bd - ae)}$$

$$\downarrow 55$$

$$\frac{(-9aBe + 8Abe + bBd) \left(\frac{6b \int \frac{1}{\sqrt{a+bx}(d+ex)^{7/2}} dx}{7(bd - ae)} + \frac{2\sqrt{a+bx}}{7(d+ex)^{7/2}(bd - ae)} \right)}{9e(bd - ae)} - \frac{2\sqrt{a + bx}(Bd - Ae)}{9e(d + ex)^{9/2}(bd - ae)}$$

$$\begin{aligned}
 & \downarrow 55 \\
 & (-9aBe + 8Abe + bBd) \left(\frac{6b \left(\frac{4b \int \frac{1}{\sqrt{a+bx}(d+ex)^{5/2}} dx + \frac{2\sqrt{a+bx}}{5(d+ex)^{5/2}(bd-ae)}}{5(bd-ae)} + \frac{2\sqrt{a+bx}}{7(d+ex)^{7/2}(bd-ae)} \right)}{7(bd-ae)} \right) \\
 & \hline
 & \frac{9e(bd-ae)}{2\sqrt{a+bx}(Bd-Ae)} \\
 & \frac{9e(d+ex)^{9/2}(bd-ae)}{9e(d+ex)^{9/2}(bd-ae)}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 55 \\
 & (-9aBe + 8Abe + bBd) \left(\frac{6b \left(\frac{4b \left(\frac{2b \int \frac{1}{\sqrt{a+bx}(d+ex)^{3/2}} dx + \frac{2\sqrt{a+bx}}{3(d+ex)^{3/2}(bd-ae)} \right)}{5(bd-ae)} + \frac{2\sqrt{a+bx}}{5(d+ex)^{5/2}(bd-ae)} \right)}{7(bd-ae)} + \frac{2\sqrt{a+bx}}{7(d+ex)^{7/2}(bd-ae)} \right) \\
 & \hline
 & \frac{9e(bd-ae)}{2\sqrt{a+bx}(Bd-Ae)} \\
 & \frac{9e(d+ex)^{9/2}(bd-ae)}{9e(d+ex)^{9/2}(bd-ae)}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 48 \\
 & \left(\frac{6b \left(\frac{4b \left(\frac{4b\sqrt{a+bx}}{3\sqrt{d+ex}(bd-ae)^2} + \frac{2\sqrt{a+bx}}{3(d+ex)^{3/2}(bd-ae)} \right)}{5(bd-ae)} + \frac{2\sqrt{a+bx}}{5(d+ex)^{5/2}(bd-ae)} \right)}{7(bd-ae)} + \frac{2\sqrt{a+bx}}{7(d+ex)^{7/2}(bd-ae)} \right) (-9aBe + 8Abe + bBd) \\
 & \hline
 & \frac{9e(bd-ae)}{2\sqrt{a+bx}(Bd-Ae)} \\
 & \frac{9e(d+ex)^{9/2}(bd-ae)}{9e(d+ex)^{9/2}(bd-ae)}
 \end{aligned}$$

input

Int[(A + B*x)/(Sqrt[a + b*x]*(d + e*x)^(11/2)),x]

output

```
(-2*(B*d - A*e)*Sqrt[a + b*x])/(9*e*(b*d - a*e)*(d + e*x)^(9/2)) + ((b*B*d
+ 8*A*b*e - 9*a*B*e)*((2*Sqrt[a + b*x])/(7*(b*d - a*e)*(d + e*x)^(7/2)) +
(6*b*((2*Sqrt[a + b*x])/(5*(b*d - a*e)*(d + e*x)^(5/2)) + (4*b*((2*Sqrt[a
+ b*x])/(3*(b*d - a*e)*(d + e*x)^(3/2)) + (4*b*Sqrt[a + b*x])/(3*(b*d - a
*e)^2*Sqrt[d + e*x])))/(5*(b*d - a*e)))/(7*(b*d - a*e)))/(9*e*(b*d - a*e
))
```

Defintions of rubi rules used

rule 48

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

rule 55

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
lerQ[n, 1])
```

rule 87

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && ( !LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 448 vs. $2(221) = 442$.

Time = 0.34 (sec) , antiderivative size = 449, normalized size of antiderivative = 1.79

output

```

2/315*(35*A*a^4*e^4 - 105*(2*B*a*b^3 - 3*A*b^4)*d^4 + 42*(3*B*a^2*b^2 - 10
*A*a*b^3)*d^3*e - 54*(B*a^3*b - 7*A*a^2*b^2)*d^2*e^2 + 10*(B*a^4 - 18*A*a^
3*b)*d*e^3 + 16*(B*b^4*d*e^3 - (9*B*a*b^3 - 8*A*b^4)*e^4)*x^4 + 8*(9*B*b^4
*d^2*e^2 - 2*(41*B*a*b^3 - 36*A*b^4)*d*e^3 + (9*B*a^2*b^2 - 8*A*a*b^3)*e^4
)*x^3 + 6*(21*B*b^4*d^3*e - 3*(65*B*a*b^3 - 56*A*b^4)*d^2*e^2 + (55*B*a^2*
b^2 - 48*A*a*b^3)*d*e^3 - (9*B*a^3*b - 8*A*a^2*b^2)*e^4)*x^2 + (105*B*b^4*
d^4 - 168*(6*B*a*b^3 - 5*A*b^4)*d^3*e + 18*(33*B*a^2*b^2 - 28*A*a*b^3)*d^2
*e^2 - 8*(31*B*a^3*b - 27*A*a^2*b^2)*d*e^3 + 5*(9*B*a^4 - 8*A*a^3*b)*e^4)*
x)*sqrt(b*x + a)*sqrt(e*x + d)/(b^5*d^10 - 5*a*b^4*d^9*e + 10*a^2*b^3*d^8*
e^2 - 10*a^3*b^2*d^7*e^3 + 5*a^4*b*d^6*e^4 - a^5*d^5*e^5 + (b^5*d^5*e^5 -
5*a*b^4*d^4*e^6 + 10*a^2*b^3*d^3*e^7 - 10*a^3*b^2*d^2*e^8 + 5*a^4*b*d*e^9
- a^5*e^10)*x^5 + 5*(b^5*d^6*e^4 - 5*a*b^4*d^5*e^5 + 10*a^2*b^3*d^4*e^6 -
10*a^3*b^2*d^3*e^7 + 5*a^4*b*d^2*e^8 - a^5*d*e^9)*x^4 + 10*(b^5*d^7*e^3 -
5*a*b^4*d^6*e^4 + 10*a^2*b^3*d^5*e^5 - 10*a^3*b^2*d^4*e^6 + 5*a^4*b*d^3*e^
7 - a^5*d^2*e^8)*x^3 + 10*(b^5*d^8*e^2 - 5*a*b^4*d^7*e^3 + 10*a^2*b^3*d^6*
e^4 - 10*a^3*b^2*d^5*e^5 + 5*a^4*b*d^4*e^6 - a^5*d^3*e^7)*x^2 + 5*(b^5*d^9
*e - 5*a*b^4*d^8*e^2 + 10*a^2*b^3*d^7*e^3 - 10*a^3*b^2*d^6*e^4 + 5*a^4*b*d
^5*e^5 - a^5*d^4*e^6)*x)

```

SymPy [F]

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{11/2}} dx = \int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{\frac{11}{2}}} dx$$

input

```
integrate((B*x+A)/(b*x+a)**(1/2)/(e*x+d)**(11/2),x)
```

output

```
Integral((A + B*x)/(sqrt(a + b*x)*(d + e*x)**(11/2)), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{11/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(11/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f or more de

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 933 vs. 2(221) = 442.

Time = 0.31 (sec) , antiderivative size = 933, normalized size of antiderivative = 3.72

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{11/2}} dx = \text{Too large to display}$$

input `integrate((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(11/2),x, algorithm="giac")`

output

```

2/315*((2*(4*(b*x + a)*(2*(B*b^10*d*e^7*abs(b) - 9*B*a*b^9*e^8*abs(b) + 8*
A*b^10*e^8*abs(b))*(b*x + a)/(b^7*d^5*e^4 - 5*a*b^6*d^4*e^5 + 10*a^2*b^5*d
^3*e^6 - 10*a^3*b^4*d^2*e^7 + 5*a^4*b^3*d*e^8 - a^5*b^2*e^9) + 9*(B*b^11*d
^2*e^6*abs(b) - 10*B*a*b^10*d*e^7*abs(b) + 8*A*b^11*d*e^7*abs(b) + 9*B*a^2
*b^9*e^8*abs(b) - 8*A*a*b^10*e^8*abs(b))/(b^7*d^5*e^4 - 5*a*b^6*d^4*e^5 +
10*a^2*b^5*d^3*e^6 - 10*a^3*b^4*d^2*e^7 + 5*a^4*b^3*d*e^8 - a^5*b^2*e^9))
+ 63*(B*b^12*d^3*e^5*abs(b) - 11*B*a*b^11*d^2*e^6*abs(b) + 8*A*b^12*d^2*e^
6*abs(b) + 19*B*a^2*b^10*d*e^7*abs(b) - 16*A*a*b^11*d*e^7*abs(b) - 9*B*a^3
*b^9*e^8*abs(b) + 8*A*a^2*b^10*e^8*abs(b))/(b^7*d^5*e^4 - 5*a*b^6*d^4*e^5
+ 10*a^2*b^5*d^3*e^6 - 10*a^3*b^4*d^2*e^7 + 5*a^4*b^3*d*e^8 - a^5*b^2*e^9)
)* (b*x + a) + 105*(B*b^13*d^4*e^4*abs(b) - 12*B*a*b^12*d^3*e^5*abs(b) + 8*
A*b^13*d^3*e^5*abs(b) + 30*B*a^2*b^11*d^2*e^6*abs(b) - 24*A*a*b^12*d^2*e^6
*abs(b) - 28*B*a^3*b^10*d*e^7*abs(b) + 24*A*a^2*b^11*d*e^7*abs(b) + 9*B*a^
4*b^9*e^8*abs(b) - 8*A*a^3*b^10*e^8*abs(b))/(b^7*d^5*e^4 - 5*a*b^6*d^4*e^5
+ 10*a^2*b^5*d^3*e^6 - 10*a^3*b^4*d^2*e^7 + 5*a^4*b^3*d*e^8 - a^5*b^2*e^9
))* (b*x + a) - 315*(B*a*b^13*d^4*e^4*abs(b) - A*b^14*d^4*e^4*abs(b) - 4*B*
a^2*b^12*d^3*e^5*abs(b) + 4*A*a*b^13*d^3*e^5*abs(b) + 6*B*a^3*b^11*d^2*e^6
*abs(b) - 6*A*a^2*b^12*d^2*e^6*abs(b) - 4*B*a^4*b^10*d*e^7*abs(b) + 4*A*a^
3*b^11*d*e^7*abs(b) + B*a^5*b^9*e^8*abs(b) - A*a^4*b^10*e^8*abs(b))/(b^7*d
^5*e^4 - 5*a*b^6*d^4*e^5 + 10*a^2*b^5*d^3*e^6 - 10*a^3*b^4*d^2*e^7 + 5*...

```

Mupad [B] (verification not implemented)

Time = 1.97 (sec) , antiderivative size = 570, normalized size of antiderivative = 2.27

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{11/2}} dx = \frac{\sqrt{d + ex} \left(\frac{20 B a^5 d e^3 + 70 A a^5 e^4 - 108 B a^4 b d^2 e^2 - 360 A a^4 b d e^3 + 252 B a^3 b^2 d^3 e + 756 A a^3 b^2 d^2 e^2 - 420 B a^2 b^3 d^4 - 840 A a^2 b^3 d^3 e + 630 A a b^4 d^5}{315 e^5 (a e - b d)^5} \right)}{\sqrt{a + bx}(d + ex)^{11/2}}$$

input

```
int((A + B*x)/((a + b*x)^(1/2)*(d + e*x)^(11/2)),x)
```

output

```

-((d + e*x)^(1/2)*((70*A*a^5*e^4 + 630*A*a*b^4*d^4 + 20*B*a^5*d*e^3 - 420*
B*a^2*b^3*d^4 - 840*A*a^2*b^3*d^3*e + 252*B*a^3*b^2*d^3*e - 108*B*a^4*b*d^
2*e^2 + 756*A*a^3*b^2*d^2*e^2 - 360*A*a^4*b*d*e^3)/(315*e^5*(a*e - b*d)^5)
+ (x*(630*A*b^5*d^4 + 90*B*a^5*e^4 - 10*A*a^4*b*e^4 - 210*B*a*b^4*d^4 + 7
2*A*a^3*b^2*d*e^3 - 1764*B*a^2*b^3*d^3*e - 252*A*a^2*b^3*d^2*e^2 + 1080*B*
a^3*b^2*d^2*e^2 + 840*A*a*b^4*d^3*e - 476*B*a^4*b*d*e^3))/(315*e^5*(a*e -
b*d)^5) + (32*b^4*x^5*(8*A*b*e - 9*B*a*e + B*b*d))/(315*e^2*(a*e - b*d)^5)
+ (16*b^3*x^4*(a*e + 9*b*d)*(8*A*b*e - 9*B*a*e + B*b*d))/(315*e^3*(a*e -
b*d)^5) + (4*b^2*x^3*(63*b^2*d^2 - a^2*e^2 + 18*a*b*d*e)*(8*A*b*e - 9*B*a*
e + B*b*d))/(315*e^4*(a*e - b*d)^5) + (2*b*x^2*(8*A*b*e - 9*B*a*e + B*b*d)
*(a^3*e^3 + 105*b^3*d^3 + 63*a*b^2*d^2*e - 9*a^2*b*d*e^2))/(315*e^5*(a*e -
b*d)^5)))/(x^5*(a + b*x)^(1/2) + (d^5*(a + b*x)^(1/2))/e^5 + (10*d^2*x^3*
(a + b*x)^(1/2))/e^2 + (10*d^3*x^2*(a + b*x)^(1/2))/e^3 + (5*d*x^4*(a + b*
x)^(1/2))/e + (5*d^4*x*(a + b*x)^(1/2))/e^4)

```

Reduce [B] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 845, normalized size of antiderivative = 3.37

$$\int \frac{A + Bx}{\sqrt{a + bx}(d + ex)^{11/2}} dx = \text{Too large to display}$$

input

```
int((B*x+A)/(b*x+a)^(1/2)/(e*x+d)^(11/2),x)
```


output

```
(2*( - 35*sqrt(d + e*x)*sqrt(a + b*x)*a**4*e**5 + 135*sqrt(d + e*x)*sqrt(a
+ b*x)*a**3*b*d*e**4 - 5*sqrt(d + e*x)*sqrt(a + b*x)*a**3*b*e**5*x - 189*
sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**2*d**2*e**3 + 27*sqrt(d + e*x)*sqrt(a
+ b*x)*a**2*b**2*d*e**4*x + 6*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b**2*e**5*x
**2 + 105*sqrt(d + e*x)*sqrt(a + b*x)*a*b**3*d**3*e**2 - 63*sqrt(d + e*x)*
sqrt(a + b*x)*a*b**3*d**2*e**3*x - 36*sqrt(d + e*x)*sqrt(a + b*x)*a*b**3*d
*e**4*x**2 - 8*sqrt(d + e*x)*sqrt(a + b*x)*a*b**3*e**5*x**3 + 105*sqrt(d +
e*x)*sqrt(a + b*x)*b**4*d**3*e**2*x + 126*sqrt(d + e*x)*sqrt(a + b*x)*b**
4*d**2*e**3*x**2 + 72*sqrt(d + e*x)*sqrt(a + b*x)*b**4*d*e**4*x**3 + 16*sq
rt(d + e*x)*sqrt(a + b*x)*b**4*e**5*x**4 - 16*sqrt(e)*sqrt(b)*b**4*d**5 -
80*sqrt(e)*sqrt(b)*b**4*d**4*e*x - 160*sqrt(e)*sqrt(b)*b**4*d**3*e**2*x**2
- 160*sqrt(e)*sqrt(b)*b**4*d**2*e**3*x**3 - 80*sqrt(e)*sqrt(b)*b**4*d*e**
4*x**4 - 16*sqrt(e)*sqrt(b)*b**4*e**5*x**5))/(315*e**2*(a**4*d**5*e**4 + 5
*a**4*d**4*e**5*x + 10*a**4*d**3*e**6*x**2 + 10*a**4*d**2*e**7*x**3 + 5*a
**4*d*e**8*x**4 + a**4*e**9*x**5 - 4*a**3*b*d**6*e**3 - 20*a**3*b*d**5*e**4
*x - 40*a**3*b*d**4*e**5*x**2 - 40*a**3*b*d**3*e**6*x**3 - 20*a**3*b*d**2*
e**7*x**4 - 4*a**3*b*d*e**8*x**5 + 6*a**2*b**2*d**7*e**2 + 30*a**2*b**2*d
**6*e**3*x + 60*a**2*b**2*d**5*e**4*x**2 + 60*a**2*b**2*d**4*e**5*x**3 + 30
*a**2*b**2*d**3*e**6*x**4 + 6*a**2*b**2*d**2*e**7*x**5 - 4*a*b**3*d**8*e -
20*a*b**3*d**7*e**2*x - 40*a*b**3*d**6*e**3*x**2 - 40*a*b**3*d**5*e**4...
```

3.214 $\int \frac{(A+Bx)(d+ex)^{5/2}}{(a+bx)^{3/2}} dx$

Optimal result	1941
Mathematica [A] (verified)	1942
Rubi [A] (verified)	1942
Maple [B] (verified)	1945
Fricas [B] (verification not implemented)	1946
Sympy [F]	1947
Maxima [F(-2)]	1948
Giac [B] (verification not implemented)	1948
Mupad [F(-1)]	1949
Reduce [B] (verification not implemented)	1949

Optimal result

Integrand size = 24, antiderivative size = 215

$$\int \frac{(A+Bx)(d+ex)^{5/2}}{(a+bx)^{3/2}} dx = \frac{5(bd-ae)(bBd+6Abe-7aBe)\sqrt{a+bx}\sqrt{d+ex}}{8b^4} + \frac{5(bBd+6Abe-7aBe)\sqrt{a+bx}(d+ex)^{3/2}}{12b^3} - \frac{2(Ab-aB)(d+ex)^{5/2}}{b^2\sqrt{a+bx}} + \frac{B\sqrt{a+bx}(d+ex)^{5/2}}{3b^2} + \frac{5(bd-ae)^2(bBd+6Abe-7aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{8b^{9/2}\sqrt{e}}$$

output

```
5/8*(-a*e+b*d)*(6*A*b*e-7*B*a*e+B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(1/2)/b^4+5/12*(6*A*b*e-7*B*a*e+B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(3/2)/b^3-2*(A*b-B*a)*(e*x+d)^(5/2)/b^2/(b*x+a)^(1/2)+1/3*B*(b*x+a)^(1/2)*(e*x+d)^(5/2)/b^2+5/8*(-a*e+b*d)^2*(6*A*b*e-7*B*a*e+B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)^(1/2))/b^(9/2)/e^(1/2)
```

Mathematica [A] (verified)

Time = 0.60 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.01

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^{3/2}} dx = \frac{\sqrt{d + ex}(-6Ab(15a^2e^2 + 5abe(-5d + ex) + b^2(8d^2 - 9dex - 2e^2x^2)) + B(15bd - ae)^2(bBd + 6Abe - 7aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{e}\sqrt{a+bx}}\right)}{8b^{9/2}\sqrt{e}}$$

input `Integrate[((A + B*x)*(d + e*x)^(5/2))/(a + b*x)^(3/2), x]`

output `(Sqrt[d + e*x]*(-6*A*b*(15*a^2*e^2 + 5*a*b*e*(-5*d + e*x) + b^2*(8*d^2 - 9*d*e*x - 2*e^2*x^2)) + B*(105*a^3*e^2 + 5*a^2*b*e*(-38*d + 7*e*x) + a*b^2*(81*d^2 - 68*d*e*x - 14*e^2*x^2) + b^3*x*(33*d^2 + 26*d*e*x + 8*e^2*x^2)))/(24*b^4*Sqrt[a + b*x]) + (5*(b*d - a*e)^2*(b*B*d + 6*A*b*e - 7*a*B*e)*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/(Sqrt[e]*Sqrt[a + b*x])])/(8*b^(9/2)*Sqrt[e])`

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.05, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {87, 60, 60, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^{3/2}} dx$$

$$\downarrow 87$$

$$\frac{(-7aBe + 6Abe + bBd) \int \frac{(d+ex)^{5/2}}{\sqrt{a+bx}} dx}{b(bd - ae)} - \frac{2(d + ex)^{7/2}(Ab - aB)}{b\sqrt{a + bx}(bd - ae)}$$

$$\downarrow 60$$

$$\frac{(-7aBe + 6Abe + bBd) \left(\frac{5(bd-ae) \int \frac{(d+ex)^{3/2}}{\sqrt{a+bx}} dx}{6b} + \frac{\sqrt{a+bx}(d+ex)^{5/2}}{3b} \right)}{b(bd-ae)} - \frac{2(d+ex)^{7/2}(Ab-aB)}{b\sqrt{a+bx}(bd-ae)}$$

↓ 60

$$\frac{(-7aBe + 6Abe + bBd) \left(\frac{5(bd-ae) \left(\frac{3(bd-ae) \int \frac{\sqrt{d+ex}}{\sqrt{a+bx}} dx}{4b} + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{6b} + \frac{\sqrt{a+bx}(d+ex)^{5/2}}{3b} \right)}{b(bd-ae)}$$

$$\frac{2(d+ex)^{7/2}(Ab-aB)}{b\sqrt{a+bx}(bd-ae)}$$

↓ 60

$$\frac{(-7aBe + 6Abe + bBd) \left(\frac{5(bd-ae) \left(\frac{3(bd-ae) \left(\frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{4b} + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{6b} + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{6b} + \frac{\sqrt{a+bx}(d+ex)^{5/2}}{3b} \right)}{b(bd-ae)}$$

$$\frac{2(d+ex)^{7/2}(Ab-aB)}{b\sqrt{a+bx}(bd-ae)}$$

↓ 66

$$\frac{(-7aBe + 6Abe + bBd) \left(\frac{5(bd-ae) \left(\frac{3(bd-ae) \left(\frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{4b} + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{6b} + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{6b} + \frac{\sqrt{a+bx}(d+ex)^{5/2}}{3b} \right)}{b(bd-ae)}$$

$$\frac{2(d+ex)^{7/2}(Ab-aB)}{b\sqrt{a+bx}(bd-ae)}$$

↓ 221

$$(-7aBe + 6Abe + bBd) \left(\frac{5(bd-ae) \left(\frac{3(bd-ae) \left(\frac{(bd-ae) \operatorname{arctanh} \left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}} \right) + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{b^{3/2}\sqrt{e}} + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{4b} \right)}{6b} \right) + \frac{\sqrt{a+bx}(d+ex)^5}{3b}$$

$$\frac{b(bd-ae) 2(d+ex)^{7/2}(Ab-aB)}{b\sqrt{a+bx}(bd-ae)}$$

input

```
Int[((A + B*x)*(d + e*x)^(5/2))/(a + b*x)^(3/2), x]
```

output

```
(-2*(A*b - a*B)*(d + e*x)^(7/2))/(b*(b*d - a*e)*Sqrt[a + b*x]) + ((b*B*d + 6*A*b*e - 7*a*B*e)*((Sqrt[a + b*x]*(d + e*x)^(5/2))/(3*b) + (5*(b*d - a*e)*((Sqrt[a + b*x]*(d + e*x)^(3/2))/(2*b) + (3*(b*d - a*e)*((Sqrt[a + b*x]*Sqrt[d + e*x])/b + ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(b^(3/2)*Sqrt[e])))/(4*b)))/(6*b)))/(b*(b*d - a*e))
```

Defintions of rubi rules used

rule 60

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

rule 66

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]
```

rule 87

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p
_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && ( !LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

rule 221

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1183 vs. $2(179) = 358$.

Time = 0.28 (sec) , antiderivative size = 1184, normalized size of antiderivative = 5.51

method	result	size
default	Expression too large to display	1184

input

```
int((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^(3/2),x,method=_RETURNVERBOSE)
```

output

```

1/48*(e*x+d)^(1/2)*(-28*B*a*b^2*e^2*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)
)+52*B*b^3*d*e*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-60*A*a*b^2*e^2*x*((
e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+108*A*b^3*d*e*x*((e*x+d)*(b*x+a))^(1/2)*
(b*e)^(1/2)+70*B*a^2*b*e^2*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+90*A*ln(1
/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^
3*b*e^3+15*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b
)/(b*e)^(1/2))*a*b^3*d^3-136*B*a*b^2*d*e*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(
1/2)+300*A*a*b^2*d*e*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-380*B*a^2*b*d*e*((
e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-180*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a
))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^3*d*e^2*x+225*B*ln(1/2*(2*b
*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*b^2*d
*e^2*x-135*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b
)/(b*e)^(1/2))*a*b^3*d^2*e*x-180*A*a^2*b*e^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)
^(1/2)+162*B*a*b^2*d^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+90*A*ln(1/2*(2*
b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*b^2*
e^3*x+90*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/
(b*e)^(1/2))*b^4*d^2*e*x-105*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*
(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^3*b*e^3*x-180*A*ln(1/2*(2*b*e*x+2*((e*x
+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*b^2*d*e^2+90*A*ln
(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2)...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 429 vs. $2(179) = 358$.

Time = 0.53 (sec) , antiderivative size = 872, normalized size of antiderivative = 4.06

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^{3/2}} dx = \text{Too large to display}$$

input

```
integrate((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^(3/2),x, algorithm="fricas")
```

output

```
[1/96*(15*(B*a*b^3*d^3 - 3*(3*B*a^2*b^2 - 2*A*a*b^3)*d^2*e + 3*(5*B*a^3*b
- 4*A*a^2*b^2)*d*e^2 - (7*B*a^4 - 6*A*a^3*b)*e^3 + (B*b^4*d^3 - 3*(3*B*a*b
^3 - 2*A*b^4)*d^2*e + 3*(5*B*a^2*b^2 - 4*A*a*b^3)*d*e^2 - (7*B*a^3*b - 6*A
*a^2*b^2)*e^3)*x)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*
e^2 + 4*(2*b*e*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x + a)*sqrt(e*x + d) + 8*(b
^2*d*e + a*b*e^2)*x) + 4*(8*B*b^4*e^3*x^3 + 3*(27*B*a*b^3 - 16*A*b^4)*d^2*
e - 10*(19*B*a^2*b^2 - 15*A*a*b^3)*d*e^2 + 15*(7*B*a^3*b - 6*A*a^2*b^2)*e^
3 + 2*(13*B*b^4*d*e^2 - (7*B*a*b^3 - 6*A*b^4)*e^3)*x^2 + (33*B*b^4*d^2*e -
2*(34*B*a*b^3 - 27*A*b^4)*d*e^2 + 5*(7*B*a^2*b^2 - 6*A*a*b^3)*e^3)*x)*sqr
t(b*x + a)*sqrt(e*x + d))/(b^6*e*x + a*b^5*e), -1/48*(15*(B*a*b^3*d^3 - 3*
(3*B*a^2*b^2 - 2*A*a*b^3)*d^2*e + 3*(5*B*a^3*b - 4*A*a^2*b^2)*d*e^2 - (7*B
*a^4 - 6*A*a^3*b)*e^3 + (B*b^4*d^3 - 3*(3*B*a*b^3 - 2*A*b^4)*d^2*e + 3*(5*
B*a^2*b^2 - 4*A*a*b^3)*d*e^2 - (7*B*a^3*b - 6*A*a^2*b^2)*e^3)*x)*sqrt(-b*e
)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d)/
(b^2*e^2*x^2 + a*b*d*e + (b^2*d*e + a*b*e^2)*x)) - 2*(8*B*b^4*e^3*x^3 + 3*
(27*B*a*b^3 - 16*A*b^4)*d^2*e - 10*(19*B*a^2*b^2 - 15*A*a*b^3)*d*e^2 + 15*
(7*B*a^3*b - 6*A*a^2*b^2)*e^3 + 2*(13*B*b^4*d*e^2 - (7*B*a*b^3 - 6*A*b^4)*
e^3)*x^2 + (33*B*b^4*d^2*e - 2*(34*B*a*b^3 - 27*A*b^4)*d*e^2 + 5*(7*B*a^2*
b^2 - 6*A*a*b^3)*e^3)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b^6*e*x + a*b^5*e)]
```

Sympy [F]

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^{3/2}} dx = \int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^{3/2}} dx$$

input

```
integrate((B*x+A)*(e*x+d)**(5/2)/(b*x+a)**(3/2),x)
```

output

```
Integral((A + B*x)*(d + e*x)**(5/2)/(a + b*x)**(3/2), x)
```


Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^{3/2}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^(3/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for m
ore detail
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 479 vs. 2(179) = 358.

Time = 0.39 (sec) , antiderivative size = 479, normalized size of antiderivative = 2.23

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^{3/2}} dx = \frac{1}{24} \sqrt{b^2d + (bx + a)be - abe} \sqrt{bx + a} \left(2(bx + a) \left(\frac{4(bx + a)Be^2|b|}{b^6} + \frac{13Bb}{b^6} \right. \right. \\ \left. \left. 5(Bb^3d^3|b| - 9Bab^2d^2e|b| + 6Ab^3d^2e|b| + 15Ba^2bde^2|b| - 12Aab^2de^2|b| - 7Ba^3e^3|b| + 6Aa^2be^3|b|) \log \right. \right. \\ \left. \left. + \frac{16\sqrt{beb^5}}{\left(b^2d - abe - \left(\sqrt{be}\sqrt{bx + a} - \sqrt{b^2d + (bx + a)be - abe} \right)^2 \right) \sqrt{beb^4}} \right. \right. \\ \left. \left. 4(Bab^3d^3e|b| - Ab^4d^3e|b| - 3Ba^2b^2d^2e^2|b| + 3Aab^3d^2e^2|b| + 3Ba^3bde^3|b| - 3Aa^2b^2de^3|b| - Ba^4e^4|b| + \right. \right. \\ \left. \left. \right. \right)$$

input

```
integrate((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^(3/2),x, algorithm="giac")
```

output

```
1/24*sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*sqrt(b*x + a)*(2*(b*x + a)*(4*(b*x + a)*B*e^2*abs(b)/b^6 + (13*B*b^18*d*e^5*abs(b) - 19*B*a*b^17*e^6*abs(b) + 6*A*b^18*e^6*abs(b))/(b^23*e^4)) + 3*(11*B*b^19*d^2*e^4*abs(b) - 40*B*a*b^18*d*e^5*abs(b) + 18*A*b^19*d*e^5*abs(b) + 29*B*a^2*b^17*e^6*abs(b) - 18*A*a*b^18*e^6*abs(b))/(b^23*e^4) - 5/16*(B*b^3*d^3*abs(b) - 9*B*a*b^2*d^2*e*abs(b) + 6*A*b^3*d^2*e*abs(b) + 15*B*a^2*b*d*e^2*abs(b) - 12*A*a*b^2*d*e^2*abs(b) - 7*B*a^3*e^3*abs(b) + 6*A*a^2*b*e^3*abs(b))*log((sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2)/(sqrt(b*e)*b^5) + 4*(B*a*b^3*d^3*e*abs(b) - A*b^4*d^3*e*abs(b) - 3*B*a^2*b^2*d^2*e^2*abs(b) + 3*A*a*b^3*d^2*e^2*abs(b) + 3*B*a^3*b*d*e^3*abs(b) - 3*A*a^2*b^2*d*e^3*abs(b) - B*a^4*e^4*abs(b) + A*a^3*b*e^4*abs(b))/((b^2*d - a*b*e - (sqrt(b*e))*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2)*sqrt(b*e)*b^4
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^{3/2}} dx = \int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^{3/2}} dx$$

input

```
int(((A + B*x)*(d + e*x)^(5/2))/(a + b*x)^(3/2), x)
```

output

```
int(((A + B*x)*(d + e*x)^(5/2))/(a + b*x)^(3/2), x)
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 320, normalized size of antiderivative = 1.49

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^{3/2}} dx = \frac{15\sqrt{ex + d}\sqrt{bx + a}a^2be^3 - 40\sqrt{ex + d}\sqrt{bx + a}ab^2de^2 - 10\sqrt{ex + d}\sqrt{bx + a}a^2be^2 + 10\sqrt{ex + d}\sqrt{bx + a}ab^2de - 10\sqrt{ex + d}\sqrt{bx + a}a^2be - 10\sqrt{ex + d}\sqrt{bx + a}ab^2d - 10\sqrt{ex + d}\sqrt{bx + a}a^2b - 10\sqrt{ex + d}\sqrt{bx + a}ab^2 - 10\sqrt{ex + d}\sqrt{bx + a}a^2 - 10\sqrt{ex + d}\sqrt{bx + a}b^2 - 10\sqrt{ex + d}\sqrt{bx + a}a - 10\sqrt{ex + d}\sqrt{bx + a}b - 10\sqrt{ex + d}\sqrt{bx + a}}{(a + bx)^{3/2}}$$

input

```
int((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^(3/2), x)
```

output

```
(15*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b*e**3 - 40*sqrt(d + e*x)*sqrt(a + b*x)*a*b**2*d*e**2 - 10*sqrt(d + e*x)*sqrt(a + b*x)*a*b**2*e**3*x + 33*sqrt(d + e*x)*sqrt(a + b*x)*b**3*d**2*e + 26*sqrt(d + e*x)*sqrt(a + b*x)*b**3*d*e**2*x + 8*sqrt(d + e*x)*sqrt(a + b*x)*b**3*e**3*x**2 - 15*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**3*e**3 + 45*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**2*b*d*e**2 - 45*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*b**2*d**2*e + 15*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**3*d**3)/(24*b**4*e)
```

3.215 $\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^{3/2}} dx$

Optimal result	1951
Mathematica [A] (verified)	1952
Rubi [A] (verified)	1952
Maple [B] (verified)	1954
Fricas [B] (verification not implemented)	1955
Sympy [F]	1956
Maxima [F(-2)]	1956
Giac [B] (verification not implemented)	1957
Mupad [F(-1)]	1958
Reduce [B] (verification not implemented)	1958

Optimal result

Integrand size = 24, antiderivative size = 165

$$\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^{3/2}} dx = \frac{3(bBd+4Abe-5aBe)\sqrt{a+bx}\sqrt{d+ex}}{4b^3} - \frac{2(Ab-aB)(d+ex)^{3/2}}{b^2\sqrt{a+bx}} + \frac{B\sqrt{a+bx}(d+ex)^{3/2}}{2b^2} + \frac{3(bd-ae)(bBd+4Abe-5aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{4b^{7/2}\sqrt{e}}$$

output

```
3/4*(4*A*b*e-5*B*a*e+B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(1/2)/b^3-2*(A*b-B*a)*(e*x+d)^(3/2)/b^2/(b*x+a)^(1/2)+1/2*B*(b*x+a)^(1/2)*(e*x+d)^(3/2)/b^2+3/4*(-a*e+b*d)*(4*A*b*e-5*B*a*e+B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)^(1/2))/b^(7/2)/e^(1/2)
```

Mathematica [A] (verified)

Time = 0.98 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.99

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^{3/2}} dx = \frac{\sqrt{b}\sqrt{d+ex}(4Ab(-2bd+3ae+box)+B(-15a^2e+ab(13d-5ex)+b^2x(5d+2ex)))}{\sqrt{a+bx}} + \frac{6(-bd+ae)(bBd+4Ab)}{4b^{7/2}}$$

input `Integrate[((A + B*x)*(d + e*x)^(3/2))/(a + b*x)^(3/2),x]`

output `((Sqrt[b]*Sqrt[d + e*x]*(4*A*b*(-2*b*d + 3*a*e + b*e*x) + B*(-15*a^2*e + a*b*(13*d - 5*e*x) + b^2*x*(5*d + 2*e*x)))/Sqrt[a + b*x] + (6*(-(b*d) + a*e)*(b*B*d + 4*A*b*e - 5*a*B*e)*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/(Sqrt[e]*(Sqrt[a - (b*d)/e] - Sqrt[a + b*x])]))/Sqrt[e])/(4*b^(7/2))`

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.12, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$, Rules used = {87, 60, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^{3/2}} dx \\ & \quad \downarrow 87 \\ & \frac{(-5aBe + 4Abe + bBd) \int \frac{(d+ex)^{3/2}}{\sqrt{a+bx}} dx}{b(bd - ae)} - \frac{2(d + ex)^{5/2}(Ab - aB)}{b\sqrt{a + bx}(bd - ae)} \\ & \quad \downarrow 60 \\ & \frac{(-5aBe + 4Abe + bBd) \left(\frac{3(bd - ae) \int \frac{\sqrt{d+ex}}{\sqrt{a+bx}} dx}{4b} + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{b(bd - ae)} - \frac{2(d + ex)^{5/2}(Ab - aB)}{b\sqrt{a + bx}(bd - ae)} \\ & \quad \downarrow 60 \end{aligned}$$

$$\begin{aligned}
 & \frac{(-5aBe + 4Abe + bBd) \left(\frac{3(bd-ae) \left(\frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{2b} + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{4b} + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{b(bd-ae)} \\
 & \frac{2(d+ex)^{5/2}(Ab-aB)}{b\sqrt{a+bx}(bd-ae)} \\
 & \quad \downarrow 66 \\
 & \frac{(-5aBe + 4Abe + bBd) \left(\frac{3(bd-ae) \left(\frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{b} + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{4b} + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{b(bd-ae)} \\
 & \frac{2(d+ex)^{5/2}(Ab-aB)}{b\sqrt{a+bx}(bd-ae)} \\
 & \quad \downarrow 221 \\
 & \frac{(-5aBe + 4Abe + bBd) \left(\frac{3(bd-ae) \left(\frac{(bd-ae) \operatorname{arctanh} \left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}} \right) + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{b^{3/2}\sqrt{e}} + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{b(bd-ae)} \\
 & \frac{2(d+ex)^{5/2}(Ab-aB)}{b\sqrt{a+bx}(bd-ae)}
 \end{aligned}$$

input `Int[((A + B*x)*(d + e*x)^(3/2))/(a + b*x)^(3/2),x]`

output `(-2*(A*b - a*B)*(d + e*x)^(5/2))/(b*(b*d - a*e)*Sqrt[a + b*x]) + ((b*B*d + 4*A*b*e - 5*a*B*e)*((Sqrt[a + b*x]*(d + e*x)^(3/2))/(2*b) + (3*(b*d - a*e))*((Sqrt[a + b*x]*Sqrt[d + e*x])/b + ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(b^(3/2)*Sqrt[e])))/(4*b))/(b*(b*d - a*e))`

Definitions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 739 vs. $2(135) = 270$.

Time = 0.27 (sec) , antiderivative size = 740, normalized size of antiderivative = 4.48

method	result
default	$-\frac{\sqrt{ex+d} \left(12A \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) a b^2 e^2 x - 12A \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) b^3 dex - 15B \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) \right)}{b^3 dex - 15B \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right)}$

input `int((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^(3/2),x,method=_RETURNVERBOSE)`

output

```

-1/8*(e*x+d)^(1/2)*(12*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(
1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^2*e^2*x-12*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*
x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^3*d*e*x-15*B*ln(1/2*(2*b*e
*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*b*e^2*x
+18*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)
^(1/2))*a*b^2*d*e*x-3*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1
/2)+a*e+d*b)/(b*e)^(1/2))*b^3*d^2*x-4*B*b^2*e*x^2*(b*e)^(1/2)*((e*x+d)*(b*
x+a))^(1/2)+12*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e
+d*b)/(b*e)^(1/2))*a^2*b*e^2-12*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2
)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^2*d*e-8*A*b^2*e*x*(b*e)^(1/2)*((e*
x+d)*(b*x+a))^(1/2)-15*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(
1/2)+a*e+d*b)/(b*e)^(1/2))*a^3*e^2+18*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a)
)^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*b*d*e-3*B*ln(1/2*(2*b*e*x+2*
((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^2*d^2+10*B*a
*b*e*x*(b*e)^(1/2)*((e*x+d)*(b*x+a))^(1/2)-10*B*b^2*d*x*(b*e)^(1/2)*((e*x+
d)*(b*x+a))^(1/2)-24*A*a*b*e*(b*e)^(1/2)*((e*x+d)*(b*x+a))^(1/2)+16*A*b^2*
d*(b*e)^(1/2)*((e*x+d)*(b*x+a))^(1/2)+30*B*a^2*e*(b*e)^(1/2)*((e*x+d)*(b*x
+a))^(1/2)-26*B*a*b*d*(b*e)^(1/2)*((e*x+d)*(b*x+a))^(1/2))/((e*x+d)*(b*x+a
))^(1/2)/(b*e)^(1/2)/(b*x+a)^(1/2)/b^3

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 285 vs. $2(135) = 270$.

Time = 0.53 (sec) , antiderivative size = 584, normalized size of antiderivative = 3.54

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^{3/2}} dx = \left[-\frac{3(Bab^2d^2 - 2(3Ba^2b - 2Aab^2)de + (5Ba^3 - 4Aa^2b)e^2 + (Bb^3d^2 - 2(3$$

input

```
integrate((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^(3/2),x, algorithm="fricas")
```


output

```
[-1/16*(3*(B*a*b^2*d^2 - 2*(3*B*a^2*b - 2*A*a*b^2)*d*e + (5*B*a^3 - 4*A*a^2*b)*e^2 + (B*b^3*d^2 - 2*(3*B*a*b^2 - 2*A*b^3)*d*e + (5*B*a^2*b - 4*A*a*b^2)*e^2)*x)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 - 4*(2*b*e*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x + a)*sqrt(e*x + d) + 8*(b^2*d*e + a*b*e^2)*x) - 4*(2*B*b^3*e^2*x^2 + (13*B*a*b^2 - 8*A*b^3)*d*e - 3*(5*B*a^2*b - 4*A*a*b^2)*e^2 + (5*B*b^3*d*e - (5*B*a*b^2 - 4*A*b^3)*e^2)*x)*sqrt(b*x + a)*sqrt(e*x + d)/(b^5*e*x + a*b^4*e), -1/8*(3*(B*a*b^2*d^2 - 2*(3*B*a^2*b - 2*A*a*b^2)*d*e + (5*B*a^3 - 4*A*a^2*b)*e^2 + (B*b^3*d^2 - 2*(3*B*a*b^2 - 2*A*b^3)*d*e + (5*B*a^2*b - 4*A*a*b^2)*e^2)*x)*sqrt(-b*e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*e^2*x^2 + a*b*d*e + (b^2*d*e + a*b*e^2)*x)) - 2*(2*B*b^3*e^2*x^2 + (13*B*a*b^2 - 8*A*b^3)*d*e - 3*(5*B*a^2*b - 4*A*a*b^2)*e^2 + (5*B*b^3*d*e - (5*B*a*b^2 - 4*A*b^3)*e^2)*x)*sqrt(b*x + a)*sqrt(e*x + d)/(b^5*e*x + a*b^4*e)]
```

Sympy [F]

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^{3/2}} dx = \int \frac{(A + Bx)(d + ex)^{\frac{3}{2}}}{(a + bx)^{\frac{3}{2}}} dx$$

input

```
integrate((B*x+A)*(e*x+d)**(3/2)/(b*x+a)**(3/2),x)
```

output

```
Integral((A + B*x)*(d + e*x)**(3/2)/(a + b*x)**(3/2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^{3/2}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^(3/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for m
ore detail
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 327 vs. $2(135) = 270$.

Time = 0.27 (sec) , antiderivative size = 327, normalized size of antiderivative = 1.98

$$\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^{3/2}} dx = \frac{1}{4} \sqrt{b^2d+(bx+a)be-abe} \sqrt{bx+a} \left(\frac{2(bx+a)Be|b|}{b^5} + \frac{5Bb^{10}de^2|b|-9Ba}{b^{14}} \right. \\ \left. + \frac{3(Bb^2d^2|b|-6Babde|b|+4Ab^2de|b|+5Ba^2e^2|b|-4Aabe^2|b|) \log\left(\left(\sqrt{be}\sqrt{bx+a}-\sqrt{b^2d+(bx+a)be-abe}\right)\right)}{8\sqrt{beb^4}} \right) \\ + \frac{4(Bab^2d^2e|b|-Ab^3d^2e|b|-2Ba^2bde^2|b|+2Aab^2de^2|b|+Ba^3e^3|b|-Aa^2be^3|b|)}{\left(b^2d-abe-\left(\sqrt{be}\sqrt{bx+a}-\sqrt{b^2d+(bx+a)be-abe}\right)^2\right)\sqrt{beb^3}}$$

input

```
integrate((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^(3/2),x, algorithm="giac")
```

output

```
1/4*sqrt(b^2*d+(b*x+a)*b*e-a*b*e)*sqrt(b*x+a)*(2*(b*x+a)*B*e*abs
(b)/b^5+(5*B*b^10*d*e^2*abs(b)-9*B*a*b^9*e^3*abs(b)+4*A*b^10*e^3*abs
(b))/(b^14*e^2))-3/8*(B*b^2*d^2*abs(b)-6*B*a*b*d*e*abs(b)+4*A*b^2*d*
e*abs(b)+5*B*a^2*e^2*abs(b)-4*A*a*b*e^2*abs(b))*log((sqrt(b*e)*sqrt(b*
x+a)-sqrt(b^2*d+(b*x+a)*b*e-a*b*e))^2/(sqrt(b*e)*b^4)+4*(B*a*
b^2*d^2*e*abs(b)-A*b^3*d^2*e*abs(b)-2*B*a^2*b*d*e^2*abs(b)+2*A*a*b^2
*d*e^2*abs(b)+B*a^3*e^3*abs(b)-A*a^2*b*e^3*abs(b)))/((b^2*d-a*b*e-(
sqrt(b*e)*sqrt(b*x+a)-sqrt(b^2*d+(b*x+a)*b*e-a*b*e))^2)*sqrt(b*e
)*b^3)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^{3/2}} dx = \int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^{3/2}} dx$$

input `int(((A + B*x)*(d + e*x)^(3/2))/(a + b*x)^(3/2), x)`

output `int(((A + B*x)*(d + e*x)^(3/2))/(a + b*x)^(3/2), x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.20

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^{3/2}} dx = \frac{-3\sqrt{ex + d}\sqrt{bx + a}abe^2 + 5\sqrt{ex + d}\sqrt{bx + a}b^2de + 2\sqrt{ex + d}\sqrt{bx + a}b^2}{(a + bx)^{3/2}}$$

input `int((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^(3/2), x)`

output `(- 3*sqrt(d + e*x)*sqrt(a + b*x)*a*b*e**2 + 5*sqrt(d + e*x)*sqrt(a + b*x)*b**2*d*e + 2*sqrt(d + e*x)*sqrt(a + b*x)*b**2*e**2*x + 3*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**2*e**2 - 6*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*b*d*e + 3*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**2*d**2)/(4*b**3*e)`

3.216 $\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^{3/2}} dx$

Optimal result	1959
Mathematica [A] (verified)	1959
Rubi [A] (verified)	1960
Maple [B] (verified)	1962
Fricas [A] (verification not implemented)	1962
Sympy [F]	1963
Maxima [F(-2)]	1963
Giac [B] (verification not implemented)	1964
Mupad [F(-1)]	1964
Reduce [B] (verification not implemented)	1965

Optimal result

Integrand size = 24, antiderivative size = 111

$$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^{3/2}} dx = -\frac{2(Ab-aB)\sqrt{d+ex}}{b^2\sqrt{a+bx}} + \frac{B\sqrt{a+bx}\sqrt{d+ex}}{b^2} + \frac{(bBd+2Abe-3aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{b^{5/2}\sqrt{e}}$$

output

```
-2*(A*b-B*a)*(e*x+d)^(1/2)/b^2/(b*x+a)^(1/2)+B*(b*x+a)^(1/2)*(e*x+d)^(1/2)
/b^2+(2*A*b*e-3*B*a*e+B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)
^(1/2))/b^(5/2)/e^(1/2)
```

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.83

$$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^{3/2}} dx = \frac{(-2Ab+3aB+bBx)\sqrt{d+ex}}{b^2\sqrt{a+bx}} + \frac{(bBd+2Abe-3aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{e}\sqrt{a+bx}}\right)}{b^{5/2}\sqrt{e}}$$

input `Integrate[((A + B*x)*Sqrt[d + e*x])/(a + b*x)^(3/2),x]`

output `((-2*A*b + 3*a*B + b*B*x)*Sqrt[d + e*x])/(b^2*Sqrt[a + b*x]) + ((b*B*d + 2*A*b*e - 3*a*B*e)*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/(Sqrt[e]*Sqrt[a + b*x])])/(b^(5/2)*Sqrt[e])`

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.29, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {87, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^{3/2}} dx \\
 & \quad \downarrow 87 \\
 & \frac{(-3aBe + 2Abe + bBd) \int \frac{\sqrt{d+ex}}{\sqrt{a+bx}} dx}{b(bd - ae)} - \frac{2(d + ex)^{3/2}(Ab - aB)}{b\sqrt{a + bx}(bd - ae)} \\
 & \quad \downarrow 60 \\
 & \frac{(-3aBe + 2Abe + bBd) \left(\frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{2b} + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{b(bd - ae)} - \frac{2(d + ex)^{3/2}(Ab - aB)}{b\sqrt{a + bx}(bd - ae)} \\
 & \quad \downarrow 66 \\
 & \frac{(-3aBe + 2Abe + bBd) \left(\frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{b} + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{b(bd - ae)} - \frac{2(d + ex)^{3/2}(Ab - aB)}{b\sqrt{a + bx}(bd - ae)} \\
 & \quad \downarrow 221 \\
 & \frac{(-3aBe + 2Abe + bBd) \left(\frac{(bd-ae) \operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{b^{3/2}\sqrt{e}} + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{b(bd - ae)} - \frac{2(d + ex)^{3/2}(Ab - aB)}{b\sqrt{a + bx}(bd - ae)}
 \end{aligned}$$

input `Int[((A + B*x)*Sqrt[d + e*x])/(a + b*x)^(3/2),x]`

output `(-2*(A*b - a*B)*(d + e*x)^(3/2))/(b*(b*d - a*e)*Sqrt[a + b*x]) + ((b*B*d + 2*A*b*e - 3*a*B*e)*((Sqrt[a + b*x]*Sqrt[d + e*x])/b + ((b*d - a*e)*ArcTan h[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(b^(3/2)*Sqrt[e])))/(b*(b*d - a*e))`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || (EqQ[e, 0] || (EqQ[c, 0] || LtQ[p, n]))))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 385 vs. $2(91) = 182$.

Time = 0.27 (sec) , antiderivative size = 386, normalized size of antiderivative = 3.48

method	result
default	$\frac{\sqrt{ex+d} \left(2A \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) b^2 ex - 3B \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) abex + B \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) \right)}{\dots}$

input `int((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^(3/2),x,method=_RETURNVERBOSE)`

output

```
1/2*(e*x+d)^(1/2)*(2*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^2*e*x-3*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b*e*x+B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^2*d*x+2*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b*e-3*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*e+B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b*d+2*B*b*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-4*A*b*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+6*B*a*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2))/(b*e)^(1/2)/((e*x+d)*(b*x+a))^(1/2)/b^2/(b*x+a)^(1/2)
```

Fricas [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 366, normalized size of antiderivative = 3.30

$$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^{3/2}} dx = \frac{\left((Babd - (3Ba^2 - 2Aab)e + (Bb^2d - (3Bab - 2Ab^2)e)x)\sqrt{be} \log \left(8b^2e^2x^2 - \dots \right) \right)}{2(b^4ex + ab^3e)} - \frac{\left((Babd - (3Ba^2 - 2Aab)e + (Bb^2d - (3Bab - 2Ab^2)e)x)\sqrt{-be} \arctan \left(\frac{(2bex+bd+ae)\sqrt{-be}\sqrt{bx+a}\sqrt{ex+d}}{2(b^2e^2x^2+abde+(b^2de+abe^2)x)} \right) \right)}{2(b^4ex + ab^3e)}$$

input `integrate((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^(3/2),x, algorithm="fricas")`

output

```
[1/4*((B*a*b*d - (3*B*a^2 - 2*A*a*b)*e + (B*b^2*d - (3*B*a*b - 2*A*b^2)*e)
*x)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*b*e
*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x + a)*sqrt(e*x + d) + 8*(b^2*d*e + a*b*e
^2)*x) + 4*(B*b^2*e*x + (3*B*a*b - 2*A*b^2)*e)*sqrt(b*x + a)*sqrt(e*x + d)
)/(b^4*e*x + a*b^3*e), -1/2*((B*a*b*d - (3*B*a^2 - 2*A*a*b)*e + (B*b^2*d -
(3*B*a*b - 2*A*b^2)*e)*x)*sqrt(-b*e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqr
t(-b*e)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*e^2*x^2 + a*b*d*e + (b^2*d*e + a*
b*e^2)*x)) - 2*(B*b^2*e*x + (3*B*a*b - 2*A*b^2)*e)*sqrt(b*x + a)*sqrt(e*x
+ d))/(b^4*e*x + a*b^3*e)]
```

Sympy [F]

$$\int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^{3/2}} dx = \int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^{\frac{3}{2}}} dx$$

input

```
integrate((B*x+A)*(e*x+d)**(1/2)/(b*x+a)**(3/2),x)
```

output

```
Integral((A + B*x)*sqrt(d + e*x)/(a + b*x)**(3/2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^{3/2}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^(3/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for m
ore detail
```


Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 210 vs. 2(91) = 182.

Time = 0.20 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.89

$$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^{3/2}} dx = \frac{\sqrt{b^2d+(bx+a)be-abe}\sqrt{bx+a}B|b|}{b^4} - \frac{(Bbd|b|-3Bae|b|+2Abe|b|)\log\left(\left(\sqrt{be}\sqrt{bx+a}-\sqrt{b^2d+(bx+a)be-abe}\right)^2\right)}{2\sqrt{beb^3}} + \frac{4(Babde|b|-Ab^2de|b|-Ba^2e^2|b|+Aabe^2|b|)}{\left(b^2d-abe-\left(\sqrt{be}\sqrt{bx+a}-\sqrt{b^2d+(bx+a)be-abe}\right)^2\right)\sqrt{beb^2}}$$

input `integrate((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^(3/2),x, algorithm="giac")`

output `sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*sqrt(b*x + a)*B*abs(b)/b^4 - 1/2*(B*b*d*abs(b) - 3*B*a*e*abs(b) + 2*A*b*e*abs(b))*log((sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2)/(sqrt(b*e)*b^3) + 4*(B*a*b*d*e*abs(b) - A*b^2*d*e*abs(b) - B*a^2*e^2*abs(b) + A*a*b*e^2*abs(b))/((b^2*d - a*b*e - (sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2)*sqrt(b*e)*b^2)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^{3/2}} dx = \int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^{3/2}} dx$$

input `int(((A + B*x)*(d + e*x)^(1/2))/(a + b*x)^(3/2),x)`

output `int(((A + B*x)*(d + e*x)^(1/2))/(a + b*x)^(3/2), x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.92

$$\int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^{3/2}} dx = \frac{\sqrt{ex + d} \sqrt{bx + a} be - \sqrt{e} \sqrt{b} \log\left(\frac{\sqrt{e} \sqrt{bx+a} + \sqrt{b} \sqrt{ex+d}}{\sqrt{ae-bd}}\right) ae + \sqrt{e} \sqrt{b} \log\left(\frac{\sqrt{e} \sqrt{bx+a}}{\sqrt{a}}\right)}{b^2 e}$$

input `int((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^(3/2),x)`output `(sqrt(d + e*x)*sqrt(a + b*x)*b*e - sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*e + sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b*d)/(b**2*e)`

3.217 $\int \frac{A+Bx}{(a+bx)^{3/2}\sqrt{d+ex}} dx$

Optimal result	1966
Mathematica [A] (verified)	1966
Rubi [A] (verified)	1967
Maple [B] (verified)	1968
Fricas [B] (verification not implemented)	1969
Sympy [F]	1969
Maxima [F(-2)]	1970
Giac [A] (verification not implemented)	1970
Mupad [F(-1)]	1971
Reduce [B] (verification not implemented)	1971

Optimal result

Integrand size = 24, antiderivative size = 85

$$\int \frac{A + Bx}{(a + bx)^{3/2}\sqrt{d + ex}} dx = -\frac{2(Ab - aB)\sqrt{d + ex}}{b(bd - ae)\sqrt{a + bx}} + \frac{2B\text{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{b^{3/2}\sqrt{e}}$$

output `-2*(A*b-B*a)*(e*x+d)^(1/2)/b/(-a*e+b*d)/(b*x+a)^(1/2)+2*B*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)^(1/2))/b^(3/2)/e^(1/2)`

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00

$$\int \frac{A + Bx}{(a + bx)^{3/2}\sqrt{d + ex}} dx = -\frac{2(Ab - aB)\sqrt{d + ex}}{b(bd - ae)\sqrt{a + bx}} + \frac{2B\text{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{e}\sqrt{a+bx}}\right)}{b^{3/2}\sqrt{e}}$$

input `Integrate[(A + B*x)/((a + b*x)^(3/2)*Sqrt[d + e*x]),x]`

output `(-2*(A*b - a*B)*Sqrt[d + e*x])/(b*(b*d - a*e)*Sqrt[a + b*x]) + (2*B*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/(Sqrt[e]*Sqrt[a + b*x])])/(b^(3/2)*Sqrt[e])`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {87, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^{3/2} \sqrt{d + ex}} dx$$

$$\downarrow 87$$

$$\frac{B \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{b} - \frac{2\sqrt{d+ex}(Ab - aB)}{b\sqrt{a+bx}(bd - ae)}$$

$$\downarrow 66$$

$$\frac{2B \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d\frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{b} - \frac{2\sqrt{d+ex}(Ab - aB)}{b\sqrt{a+bx}(bd - ae)}$$

$$\downarrow 221$$

$$\frac{2B \operatorname{Arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{b^{3/2}\sqrt{e}} - \frac{2\sqrt{d+ex}(Ab - aB)}{b\sqrt{a+bx}(bd - ae)}$$

input

```
Int[(A + B*x)/((a + b*x)^(3/2)*Sqrt[d + e*x]),x]
```

output

```
(-2*(A*b - a*B)*Sqrt[d + e*x])/(b*(b*d - a*e)*Sqrt[a + b*x]) + (2*B*ArcTan  
h[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(b^(3/2)*Sqrt[e])
```

Definitions of rubi rules used

rule 66

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[
  2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; Free
  eQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]
```

rule 87

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Integer
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

rule 221

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 277 vs. $2(69) = 138$.

Time = 0.30 (sec) , antiderivative size = 278, normalized size of antiderivative = 3.27

method	result
default	$\frac{\sqrt{ex+d} \left(B \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) abex - B \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) b^2 dx + B \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) \sqrt{be} (ae-db) \sqrt{(ex+d)(bx+a)}}{\sqrt{be} (ae-db) \sqrt{(ex+d)(bx+a)}} \right)$

input

```
int((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
(e*x+d)^(1/2)*(B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e
+d*b)/(b*e)^(1/2))*a*b*e*x-B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*
e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^2*d*x+B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a
))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*e-B*ln(1/2*(2*b*e*x+2*((e*x
+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b*d+2*A*b*((e*x+d)*
(b*x+a))^(1/2)*(b*e)^(1/2)-2*B*a*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2))/(b*e
)^(1/2)/(a*e-b*d)/((e*x+d)*(b*x+a))^(1/2)/b/(b*x+a)^(1/2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 173 vs. $2(69) = 138$.

Time = 0.25 (sec) , antiderivative size = 360, normalized size of antiderivative = 4.24

$$\int \frac{A + Bx}{(a + bx)^{3/2} \sqrt{d + ex}} dx = \frac{4(Bab - Ab^2)\sqrt{bx + a}\sqrt{ex + d}e + (Babd - Ba^2e + (Bb^2d - Babe)x)\sqrt{be} \log\left(\frac{8b^2e^2x^2 + b^2d^2 + 6a*b*d*e + a^2e^2 + 4(2b*ex + b*d + a*e)\sqrt{b*e}\sqrt{b*x + a}\sqrt{e*x + d} + 8(b^2*d*e + a*b*e^2)*x}{(a*b^3*d*e - a^2*b^2*e^2 + (b^4*d*e - a*b^3*e^2)*x)}\right) + (2*(B*a*b - A*b^2)*\sqrt{b*x + a}\sqrt{e*x + d}*e - (B*a*b*d - B*a^2*e + (B*b^2*d - B*a*b*e)*x)*\sqrt{-b*e}*\arctan(1/2*(2*b*e*x + b*d + a*e)*\sqrt{-b*e}\sqrt{b*x + a}\sqrt{e*x + d}/(b^2*e^2*x^2 + a*b*d*e + (b^2*d*e + a*b*e^2)*x))}{(a*b^3*d*e - a^2*b^2*e^2 + (b^4*d*e - a*b^3*e^2)*x)}$$

input `integrate((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(1/2),x, algorithm="fricas")`

output `[1/2*(4*(B*a*b - A*b^2)*sqrt(b*x + a)*sqrt(e*x + d)*e + (B*a*b*d - B*a^2*e + (B*b^2*d - B*a*b*e)*x)*sqrt(b*e)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*b*e*x + b*d + a*e)*sqrt(b*e)*sqrt(b*x + a)*sqrt(e*x + d) + 8*(b^2*d*e + a*b*e^2)*x))/(a*b^3*d*e - a^2*b^2*e^2 + (b^4*d*e - a*b^3*e^2)*x), (2*(B*a*b - A*b^2)*sqrt(b*x + a)*sqrt(e*x + d)*e - (B*a*b*d - B*a^2*e + (B*b^2*d - B*a*b*e)*x)*sqrt(-b*e)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(-b*e)*sqrt(b*x + a)*sqrt(e*x + d)/(b^2*e^2*x^2 + a*b*d*e + (b^2*d*e + a*b*e^2)*x)))/(a*b^3*d*e - a^2*b^2*e^2 + (b^4*d*e - a*b^3*e^2)*x)]`

Sympy [F]

$$\int \frac{A + Bx}{(a + bx)^{3/2} \sqrt{d + ex}} dx = \int \frac{A + Bx}{(a + bx)^{\frac{3}{2}} \sqrt{d + ex}} dx$$

input `integrate((B*x+A)/(b*x+a)**(3/2)/(e*x+d)**(1/2),x)`

output `Integral((A + B*x)/((a + b*x)**(3/2)*sqrt(d + e*x)), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{(a + bx)^{3/2} \sqrt{d + ex}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(1/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.54

$$\int \frac{A + Bx}{(a + bx)^{3/2} \sqrt{d + ex}} dx =$$

$$\frac{B \log \left(\left(\sqrt{be} \sqrt{bx + a} - \sqrt{b^2 d + (bx + a)be - abe} \right)^2 \right)}{\sqrt{be} |b|}$$

$$+ \frac{4 \left(\sqrt{be} Ba - \sqrt{be} Ab \right)}{\left(b^2 d - abe - \left(\sqrt{be} \sqrt{bx + a} - \sqrt{b^2 d + (bx + a)be - abe} \right)^2 \right) |b|}$$

input `integrate((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(1/2),x, algorithm="giac")`

output `-B*log((sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2)/(sqrt(b*e)*abs(b)) + 4*(sqrt(b*e)*B*a - sqrt(b*e)*A*b)/((b^2*d - a*b*e - (sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2)*abs(b))`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx}{(a + bx)^{3/2} \sqrt{d + ex}} dx = \int \frac{A + Bx}{(a + bx)^{3/2} \sqrt{d + ex}} dx$$

input `int((A + B*x)/((a + b*x)^(3/2)*(d + e*x)^(1/2)),x)`

output `int((A + B*x)/((a + b*x)^(3/2)*(d + e*x)^(1/2)), x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.52

$$\int \frac{A + Bx}{(a + bx)^{3/2} \sqrt{d + ex}} dx = \frac{2\sqrt{e} \sqrt{b} \log\left(\frac{\sqrt{e} \sqrt{bx+a} + \sqrt{b} \sqrt{ex+d}}{\sqrt{ae-bd}}\right)}{be}$$

input `int((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(1/2),x)`

output `(2*sqrt(e)*sqrt(b)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d)))/(b*e)`

3.218 $\int \frac{A+Bx}{(a+bx)^{3/2}(d+ex)^{3/2}} dx$

Optimal result	1972
Mathematica [A] (verified)	1972
Rubi [A] (verified)	1973
Maple [A] (verified)	1974
Fricas [A] (verification not implemented)	1975
Sympy [F]	1975
Maxima [F(-2)]	1975
Giac [B] (verification not implemented)	1976
Mupad [B] (verification not implemented)	1976
Reduce [B] (verification not implemented)	1977

Optimal result

Integrand size = 24, antiderivative size = 89

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{3/2}} dx = -\frac{2(Ab - aB)}{b(bd - ae)\sqrt{a + bx}\sqrt{d + ex}} + \frac{2(bBd - 2Abe + aBe)\sqrt{a + bx}}{b(bd - ae)^2\sqrt{d + ex}}$$

output

```
(-2*A*b+2*B*a)/b/(-a*e+b*d)/(b*x+a)^(1/2)/(e*x+d)^(1/2)+2*(-2*A*b*e+B*a*e+B*b*d)*(b*x+a)^(1/2)/b/(-a*e+b*d)^2/(e*x+d)^(1/2)
```

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.69

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{3/2}} dx = \frac{2B(2ad + bdx + aex) - 2A(ae + b(d + 2ex))}{(bd - ae)^2\sqrt{a + bx}\sqrt{d + ex}}$$

input

```
Integrate[(A + B*x)/((a + b*x)^(3/2)*(d + e*x)^(3/2)),x]
```

output

```
(2*B*(2*a*d + b*d*x + a*e*x) - 2*A*(a*e + b*(d + 2*e*x)))/((b*d - a*e)^2*Sqrt[a + b*x]*Sqrt[d + e*x])
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {87, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{3/2}} dx$$

$$\downarrow 87$$

$$-\frac{(aBe - 2Abe + bBd) \int \frac{1}{(a+bx)^{3/2}\sqrt{d+ex}} dx}{e(bd - ae)} - \frac{2(Bd - Ae)}{e\sqrt{a + bx}\sqrt{d + ex}(bd - ae)}$$

$$\downarrow 48$$

$$\frac{2\sqrt{d + ex}(aBe - 2Abe + bBd)}{e\sqrt{a + bx}(bd - ae)^2} - \frac{2(Bd - Ae)}{e\sqrt{a + bx}\sqrt{d + ex}(bd - ae)}$$

input

```
Int[(A + B*x)/((a + b*x)^(3/2)*(d + e*x)^(3/2)),x]
```

output

```
(-2*(B*d - A*e))/(e*(b*d - a*e)*Sqrt[a + b*x]*Sqrt[d + e*x]) + (2*(b*B*d - 2*A*b*e + a*B*e)*Sqrt[d + e*x])/(e*(b*d - a*e)^2*Sqrt[a + b*x])
```

Definitions of rubi rules used

rule 48

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

rule 87

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.66

method	result	size
default	$-\frac{2(2Abe x - Baex - Bbdx + Aae + Abd - 2Bad)}{(ae - db)^2 \sqrt{ex + d} \sqrt{bx + a}}$	59
gosper	$-\frac{2(2Abe x - Baex - Bbdx + Aae + Abd - 2Bad)}{\sqrt{ex + d} \sqrt{bx + a} (a^2 e^2 - 2abde + b^2 d^2)}$	72
orering	$-\frac{2(2Abe x - Baex - Bbdx + Aae + Abd - 2Bad)}{\sqrt{ex + d} \sqrt{bx + a} (a^2 e^2 - 2abde + b^2 d^2)}$	72

input

```
int((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(3/2),x,method=_RETURNVERBOSE)
```

output

```
-2*(2*A*b*e*x-B*a*e*x-B*b*d*x+A*a*e+A*b*d-2*B*a*d)/(a*e-b*d)^2/(e*x+d)^(1/
2)/(b*x+a)^(1/2)
```

Fricas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.66

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{3/2}} dx = \frac{2(Aae - (2Ba - Ab)d - (Bbd + (Ba - 2Ab)e)x)\sqrt{bx + a}\sqrt{ex + d}}{ab^2d^3 - 2a^2bd^2e + a^3de^2 + (b^3d^2e - 2ab^2de^2 + a^2be^3)x^2 + (b^3d^3 - ab^2d^2e - a^2bde^2 + a^3e^3)x}$$

input `integrate((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(3/2),x, algorithm="fricas")`

output `-2*(A*a*e - (2*B*a - A*b)*d - (B*b*d + (B*a - 2*A*b)*e)*x)*sqrt(b*x + a)*s
qrt(e*x + d)/(a*b^2*d^3 - 2*a^2*b*d^2*e + a^3*d*e^2 + (b^3*d^2*e - 2*a*b^2
*d*e^2 + a^2*b*e^3)*x^2 + (b^3*d^3 - a*b^2*d^2*e - a^2*b*d*e^2 + a^3*e^3)*
x)`

Sympy [F]

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{3/2}} dx = \int \frac{A + Bx}{(a + bx)^{\frac{3}{2}}(d + ex)^{\frac{3}{2}}} dx$$

input `integrate((B*x+A)/(b*x+a)**(3/2)/(e*x+d)**(3/2),x)`

output `Integral((A + B*x)/((a + b*x)**(3/2)*(d + e*x)**(3/2)), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{3/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(3/2),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for m
ore detail
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 271 vs. $2(81) = 162$.

Time = 0.19 (sec) , antiderivative size = 271, normalized size of antiderivative = 3.04

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{3/2}} dx = \frac{2(Bb^2d - Ab^2e)\sqrt{bx + a}}{(b^2d^2|b| - 2abde|b| + a^2e^2|b|)\sqrt{b^2d + (bx + a)be - abe}} + \frac{4(B^2a^2b^3e - 2ABab^4e + A^2b^5e)}{\left(\sqrt{be}Bab^3d - \sqrt{be}Ab^4d - \sqrt{be}Ba^2b^2e + \sqrt{be}Aab^3e - \sqrt{be}\left(\sqrt{be}\sqrt{bx + a} - \sqrt{b^2d + (bx + a)be - abe}\right)^2\right)^2}$$

input

```
integrate((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(3/2),x, algorithm="giac")
```

output

```
2*(B*b^2*d - A*b^2*e)*sqrt(b*x + a)/((b^2*d^2*abs(b) - 2*a*b*d*e*abs(b) +
a^2*e^2*abs(b))*sqrt(b^2*d + (b*x + a)*b*e - a*b*e)) + 4*(B^2*a^2*b^3*e -
2*A*B*a*b^4*e + A^2*b^5*e)/((sqrt(b*e)*B*a*b^3*d - sqrt(b*e)*A*b^4*d - sq
rt(b*e)*B*a^2*b^2*e + sqrt(b*e)*A*a*b^3*e - sqrt(b*e)*(sqrt(b*e)*sqrt(b*x +
a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*B*a*b + sqrt(b*e)*(sqrt(b*e)*
sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*A*b^2)*(b*d*abs(b)
- a*e*abs(b)))
```

Mupad [B] (verification not implemented)

Time = 1.47 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.08

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{3/2}} dx = -\frac{\left(\frac{2Aae+2Abd-4Bad}{e(ae-bd)^2} - \frac{x(2Bae-4Abe+2Bbd)}{e(ae-bd)^2}\right)\sqrt{d+ex}}{x\sqrt{a+bx} + \frac{d\sqrt{a+bx}}{e}}$$

input

```
int((A + B*x)/((a + b*x)^(3/2)*(d + e*x)^(3/2)),x)
```

output

```
-(((2*A*a*e + 2*A*b*d - 4*B*a*d)/(e*(a*e - b*d)^2) - (x*(2*B*a*e - 4*A*b*e
+ 2*B*b*d))/(e*(a*e - b*d)^2))*(d + e*x)^(1/2))/(x*(a + b*x)^(1/2) + (d*(
a + b*x)^(1/2))/e)
```

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.67

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{3/2}} dx = \frac{-2\sqrt{ex + d}\sqrt{bx + a}e - 2\sqrt{e}\sqrt{b}d - 2\sqrt{e}\sqrt{b}ex}{e(ae^2x - bdex + ade - bd^2)}$$

input

```
int((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(3/2),x)
```

output

```
( - 2*(sqrt(d + e*x)*sqrt(a + b*x)*e + sqrt(e)*sqrt(b)*d + sqrt(e)*sqrt(b)
*e*x))/(e*(a*d*e + a*e**2*x - b*d**2 - b*d*e*x))
```

3.219 $\int \frac{A+Bx}{(a+bx)^{3/2}(d+ex)^{5/2}} dx$

Optimal result	1978
Mathematica [A] (verified)	1978
Rubi [A] (verified)	1979
Maple [A] (verified)	1980
Fricas [B] (verification not implemented)	1981
Sympy [F]	1981
Maxima [F(-2)]	1982
Giac [B] (verification not implemented)	1982
Mupad [B] (verification not implemented)	1983
Reduce [B] (verification not implemented)	1984

Optimal result

Integrand size = 24, antiderivative size = 139

$$\int \frac{A+Bx}{(a+bx)^{3/2}(d+ex)^{5/2}} dx = -\frac{2(Ab-aB)}{b(bd-ae)\sqrt{a+bx}(d+ex)^{3/2}} + \frac{2(bBd-4Abe+3aBe)\sqrt{a+bx}}{3b(bd-ae)^2(d+ex)^{3/2}} + \frac{4(bBd-4Abe+3aBe)\sqrt{a+bx}}{3(bd-ae)^3\sqrt{d+ex}}$$

output

```
(-2*A*b+2*B*a)/b/(-a*e+b*d)/(b*x+a)^(1/2)/(e*x+d)^(3/2)+2/3*(-4*A*b*e+3*B*a*e+B*b*d)*(b*x+a)^(1/2)/b/(-a*e+b*d)^2/(e*x+d)^(3/2)+4/3*(-4*A*b*e+3*B*a*e+B*b*d)*(b*x+a)^(1/2)/(-a*e+b*d)^3/(e*x+d)^(1/2)
```

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.96

$$\int \frac{A+Bx}{(a+bx)^{3/2}(d+ex)^{5/2}} dx = \frac{2(a+bx)^{3/2} \left(Bde - Ae^2 - \frac{3bBd(d+ex)}{a+bx} + \frac{6Abe(d+ex)}{a+bx} - \frac{3aBe(d+ex)}{a+bx} + \frac{3Ab^2(d+ex)^2}{(a+bx)^2} - \frac{3abB(d+ex)^2}{(a+bx)^2} \right)}{3(bd-ae)^3(d+ex)^{3/2}}$$

input

```
Integrate[(A + B*x)/((a + b*x)^(3/2)*(d + e*x)^(5/2)),x]
```

output

$$(-2*(a + b*x)^{(3/2)}*(B*d*e - A*e^2 - (3*b*B*d*(d + e*x))/(a + b*x) + (6*A*b*e*(d + e*x))/(a + b*x) - (3*a*B*e*(d + e*x))/(a + b*x) + (3*A*b^2*(d + e*x)^2)/(a + b*x)^2 - (3*a*b*B*(d + e*x)^2)/(a + b*x)^2))/(3*(b*d - a*e)^3*(d + e*x)^{(3/2)})$$
Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.99, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {87, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{5/2}} dx$$

$$\downarrow 87$$

$$\frac{(3aBe - 4Abe + bBd) \int \frac{1}{\sqrt{a+bx}(d+ex)^{5/2}} dx}{b(bd - ae)} - \frac{2(Ab - aB)}{b\sqrt{a + bx}(d + ex)^{3/2}(bd - ae)}$$

$$\downarrow 55$$

$$\frac{(3aBe - 4Abe + bBd) \left(\frac{2b \int \frac{1}{\sqrt{a+bx}(d+ex)^{3/2}} dx}{3(bd - ae)} + \frac{2\sqrt{a+bx}}{3(d+ex)^{3/2}(bd - ae)} \right)}{b(bd - ae)} - \frac{2(Ab - aB)}{b\sqrt{a + bx}(d + ex)^{3/2}(bd - ae)}$$

$$\downarrow 48$$

$$\frac{\left(\frac{4b\sqrt{a+bx}}{3\sqrt{d+ex}(bd - ae)^2} + \frac{2\sqrt{a+bx}}{3(d+ex)^{3/2}(bd - ae)} \right) (3aBe - 4Abe + bBd)}{b(bd - ae)} - \frac{2(Ab - aB)}{b\sqrt{a + bx}(d + ex)^{3/2}(bd - ae)}$$

input

$$\text{Int}[(A + B*x)/((a + b*x)^{(3/2)}*(d + e*x)^{(5/2))}, x]$$

output
$$\frac{(-2*(A*b - a*B))/(b*(b*d - a*e)*\text{Sqrt}[a + b*x]*(d + e*x)^{(3/2)}) + ((b*B*d - 4*A*b*e + 3*a*B*e)*((2*\text{Sqrt}[a + b*x]))/(3*(b*d - a*e)*(d + e*x)^{(3/2)}) + (4*b*\text{Sqrt}[a + b*x]))/(3*(b*d - a*e)^2*\text{Sqrt}[d + e*x]))}{(b*(b*d - a*e))}$$

Defintions of rubi rules used

rule 48
$$\text{Int}[(a + b*x)^m * (c + d*x)^n, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{m+1} * (c + d*x)^{n+1} / ((b*c - a*d) * (m+1)), x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x \ \&\& \ \text{EqQ}[m + n + 2, 0] \ \&\& \ \text{NeQ}[m, -1]$$

rule 55
$$\text{Int}[(a + b*x)^m * (c + d*x)^n, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{m+1} * (c + d*x)^{n+1} / ((b*c - a*d) * (m+1)), x] - \text{Simp}[d * (\text{Simplify}[m + n + 2] / ((b*c - a*d) * (m+1))) \ \text{Int}[(a + b*x)^{\text{Simplify}[m+1]} * (c + d*x)^n, x], x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x \ \&\& \ \text{ILtQ}[\text{Simplify}[m + n + 2], 0] \ \&\& \ \text{NeQ}[m, -1] \ \&\& \ !(\text{LtQ}[m, -1] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ (\text{EqQ}[a, 0] \ || \ (\text{NeQ}[c, 0] \ \&\& \ \text{LtQ}[m - n, 0] \ \&\& \ \text{IntegerQ}[n]))) \ \&\& \ (\text{SumSimplerQ}[m, 1] \ || \ !\text{SumSimplerQ}[n, 1])$$

rule 87
$$\text{Int}[(a + b*x)^m * (c + d*x)^n * (e + f*x)^p, x] \rightarrow \text{Simp}[(-b*e - a*f) * (c + d*x)^{n+1} * (e + f*x)^{p+1} / (f * (p+1) * (c*f - d*e)), x] - \text{Simp}[(a*d*f * (n + p + 2) - b * (d*e * (n + 1) + c*f * (p + 1))) / (f * (p + 1) * (c*f - d*e)) \ \text{Int}[(c + d*x)^n * (e + f*x)^{p+1}, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ (!\text{LtQ}[n, -1] \ || \ \text{IntegerQ}[p] \ || \ !(\text{IntegerQ}[n] \ || \ !(\text{EqQ}[e, 0] \ || \ !(\text{EqQ}[c, 0] \ || \ \text{LtQ}[p, n])))$$

Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.06

method	result
default	$-\frac{2(-8A^2b^2e^2x^2+6Bab^2e^2x^2+2B^2b^2dex^2-4Aab^2e^2x-12A^2b^2dex+3Ba^2e^2x+10Babdex+3b^2Bd^2x+a^2Ae^2-6Aabde-3Ab^2d^2+2A^2e^2)}{3(ex+d)^{\frac{3}{2}}\sqrt{bx+a}(ae-db)^3}$
gospers	$-\frac{2(-8A^2b^2e^2x^2+6Bab^2e^2x^2+2B^2b^2dex^2-4Aab^2e^2x-12A^2b^2dex+3Ba^2e^2x+10Babdex+3b^2Bd^2x+a^2Ae^2-6Aabde-3Ab^2d^2+2A^2e^2)}{3(ex+d)^{\frac{3}{2}}\sqrt{bx+a}(a^3e^3-3a^2be^2d+3ab^2d^2e-b^3d^3)}$
orering	$-\frac{2(-8A^2b^2e^2x^2+6Bab^2e^2x^2+2B^2b^2dex^2-4Aab^2e^2x-12A^2b^2dex+3Ba^2e^2x+10Babdex+3b^2Bd^2x+a^2Ae^2-6Aabde-3Ab^2d^2+2A^2e^2)}{3(ex+d)^{\frac{3}{2}}\sqrt{bx+a}(a^3e^3-3a^2be^2d+3ab^2d^2e-b^3d^3)}$

input `int((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(5/2),x,method=_RETURNVERBOSE)`

output
$$-\frac{2}{3} \frac{(-8A^2b^2e^2x^2 + 6B^2a^2be^2x^2 + 2B^2b^2d^2e^2x - 4A^2a^2be^2x - 12A^2b^2d^2e^2x + 3B^2a^2e^2x + 10B^2a^2bd^2e^2x + 3B^2b^2d^2x + A^2a^2e^2 - 6A^2a^2bd^2e - 3A^2b^2d^2 + 2B^2a^2d^2e + 6B^2a^2bd^2)}{(e*x+d)^{(3/2)}(b*x+a)^{(1/2)}(a*e-b*d)^3}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 336 vs. $2(123) = 246$.

Time = 1.43 (sec) , antiderivative size = 336, normalized size of antiderivative = 2.42

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{5/2}} dx = \frac{2(Aa^2e^2 + 3(2Bab - Ab^2)d^2 + 2(Ba^2 - 3Aab)de + 2(Bb^2d^2e + (3B^2a^2b - 4A^2b^2)e^2)x^2 + (3B^2b^2d^2 + 2(5B^2a^2b - 6A^2b^2)d^2e + (3B^2a^2 - 4A^2a^2b)e^2)x) \sqrt{bx + a} \sqrt{ex + d}}{3(ab^3d^5 - 3a^2b^2d^4e + 3a^3bd^3e^2 - a^4d^2e^3 + (b^4d^3e^2 - 3ab^3d^2e^3 + 3a^2b^2de^4)x^5 - 3a^2b^2d^4e + 3a^3b^2d^3e^2 - a^4d^2e^3 + (b^4d^3e^2 - 3a^2b^3d^2e^3 + 3a^2b^2d^2e^4 - a^3b^2e^5)x^3 + (2b^4d^4e - 5a^2b^3d^3e^2 + 3a^2b^2d^2e^3 + a^3b^2d^2e^4 - a^4e^5)x^2 + (b^4d^5 - a^2b^3d^4e - 3a^2b^2d^3e^2 + 5a^3b^2d^2e^3 - 2a^4d^2e^4)x}$$

input `integrate((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(5/2),x, algorithm="fricas")`

output
$$\frac{2/3(A^2a^2e^2 + 3(2B^2a^2b - A^2b^2)d^2 + 2(B^2a^2 - 3A^2a^2b)d^2e + 2(B^2b^2d^2e + (3B^2a^2b - 4A^2b^2)e^2)x^2 + (3B^2b^2d^2 + 2(5B^2a^2b - 6A^2b^2)d^2e + (3B^2a^2 - 4A^2a^2b)e^2)x) \sqrt{bx + a} \sqrt{ex + d}}{(a^2b^3d^5 - 3a^2b^2d^4e + 3a^3b^2d^3e^2 - a^4d^2e^3 + (b^4d^3e^2 - 3a^2b^3d^2e^3 + 3a^2b^2d^2e^4 - a^3b^2e^5)x^3 + (2b^4d^4e - 5a^2b^3d^3e^2 + 3a^2b^2d^2e^3 + a^3b^2d^2e^4 - a^4e^5)x^2 + (b^4d^5 - a^2b^3d^4e - 3a^2b^2d^3e^2 + 5a^3b^2d^2e^3 - 2a^4d^2e^4)x}$$

Sympy [F]

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{5/2}} dx = \int \frac{A + Bx}{(a + bx)^{\frac{3}{2}}(d + ex)^{\frac{5}{2}}} dx$$

input `integrate((B*x+A)/(b*x+a)**(3/2)/(e*x+d)**(5/2),x)`

output `Integral((A + B*x)/((a + b*x)**(3/2)*(d + e*x)**(5/2)), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(5/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f or more de

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 616 vs. 2(123) = 246.

Time = 0.24 (sec) , antiderivative size = 616, normalized size of antiderivative = 4.43

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{5/2}} dx = \frac{2\sqrt{bx+a} \left(\frac{(2Bb^6d^3e^2|b| - Bab^5d^2e^3|b| - 5Ab^6d^2e^3|b| - 4Ba^2b^4de^4|b| + 10Aab^5de^4|b| + 3Ba^3b^3e^5|b|}{b^7d^5e - 5ab^6d^4e^2 + 10a^2b^5d^3e^3 - 10a^3b^4d^2e^4 + 5a^4b^3de^5 - a^5b^2e^6} \right)}{4(B^2a^2b^5e - 2ABab^6e} + \frac{\left(\sqrt{be}Bab^4d - \sqrt{be}Ab^5d - \sqrt{be}Ba^2b^3e + \sqrt{be}Aab^4e - \sqrt{be} \left(\sqrt{be}\sqrt{bx+a} - \sqrt{b^2d + (bx+a)be - abe} \right)^2 \right)}{E}$$

input `integrate((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(5/2),x, algorithm="giac")`

output

```

2/3*sqrt(b*x + a)*((2*B*b^6*d^3*e^2*abs(b) - B*a*b^5*d^2*e^3*abs(b) - 5*A*
b^6*d^2*e^3*abs(b) - 4*B*a^2*b^4*d*e^4*abs(b) + 10*A*a*b^5*d*e^4*abs(b) +
3*B*a^3*b^3*e^5*abs(b) - 5*A*a^2*b^4*e^5*abs(b))*(b*x + a)/(b^7*d^5*e - 5*
a*b^6*d^4*e^2 + 10*a^2*b^5*d^3*e^3 - 10*a^3*b^4*d^2*e^4 + 5*a^4*b^3*d*e^5
- a^5*b^2*e^6) + 3*(B*b^7*d^4*e*abs(b) - 2*B*a*b^6*d^3*e^2*abs(b) - 2*A*b^
7*d^3*e^2*abs(b) + 6*A*a*b^6*d^2*e^3*abs(b) + 2*B*a^3*b^4*d*e^4*abs(b) - 6
*A*a^2*b^5*d*e^4*abs(b) - B*a^4*b^3*e^5*abs(b) + 2*A*a^3*b^4*e^5*abs(b))/(
b^7*d^5*e - 5*a*b^6*d^4*e^2 + 10*a^2*b^5*d^3*e^3 - 10*a^3*b^4*d^2*e^4 + 5*
a^4*b^3*d*e^5 - a^5*b^2*e^6))/(b^2*d + (b*x + a)*b*e - a*b*e)^(3/2) + 4*(B
^2*a^2*b^5*e - 2*A*B*a*b^6*e + A^2*b^7*e)/((sqrt(b*e)*B*a*b^4*d - sqrt(b*e
)*A*b^5*d - sqrt(b*e)*B*a^2*b^3*e + sqrt(b*e)*A*a*b^4*e - sqrt(b*e)*(sqrt(
b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*B*a*b^2 + sqrt
(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*A*
b^3)*(b^2*d^2*abs(b) - 2*a*b*d*e*abs(b) + a^2*e^2*abs(b)))

```

Mupad [B] (verification not implemented)

Time = 1.61 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.31

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{5/2}} dx = \frac{\sqrt{d + ex} \left(\frac{4Ba^2de + 2Aa^2e^2 + 12Babd^2 - 12Aabde - 6Ab^2d^2}{3e^2(ae - bd)^3} + \frac{2x(ae + 3bd)(3Bae - 4Abe + Bbd)}{3e^2(ae - bd)^3} + \frac{4bx^2(3Bae - 4Abe + Bbd)}{3e(ae - bd)^3} \right)}{x^2 \sqrt{a + bx} + \frac{d^2 \sqrt{a + bx}}{e^2} + \frac{2dx \sqrt{a + bx}}{e}}$$

input

```
int((A + B*x)/((a + b*x)^(3/2)*(d + e*x)^(5/2)),x)
```

output

```

-((d + e*x)^(1/2))*((2*A*a^2*e^2 - 6*A*b^2*d^2 + 12*B*a*b*d^2 + 4*B*a^2*d*e
- 12*A*a*b*d*e)/(3*e^2*(a*e - b*d)^3) + (2*x*(a*e + 3*b*d)*(3*B*a*e - 4*A
*b*e + B*b*d))/(3*e^2*(a*e - b*d)^3) + (4*b*x^2*(3*B*a*e - 4*A*b*e + B*b*d
))/(3*e*(a*e - b*d)^3))/(x^2*(a + b*x)^(1/2) + (d^2*(a + b*x)^(1/2))/e^2
+ (2*d*x*(a + b*x)^(1/2))/e)

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.34

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{5/2}} dx = \frac{-\frac{2\sqrt{ex+d}\sqrt{bx+a}ae^2}{3} + 2\sqrt{ex+d}\sqrt{bx+a}bde + \frac{4\sqrt{ex+d}\sqrt{bx+a}be^2x}{3} - \frac{4\sqrt{e}\sqrt{b}bd^2}{3}}{e(a^2e^4x^2 - 2abd e^3x^2 + b^2d^2e^2x^2 + 2a^2d e^3x - 4abd^2e^2x + 2b^2d^3ex + a^2)}$$

input `int((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(5/2),x)`output `(2*(- sqrt(d + e*x)*sqrt(a + b*x)*a**2 + 3*sqrt(d + e*x)*sqrt(a + b*x)*b*d*e + 2*sqrt(d + e*x)*sqrt(a + b*x)*b*e**2*x - 2*sqrt(e)*sqrt(b)*b*d**2 - 4*sqrt(e)*sqrt(b)*b*d*e*x - 2*sqrt(e)*sqrt(b)*b*e**2*x**2))/(3*e*(a**2*d**2*e**2 + 2*a**2*d*e**3*x + a**2*e**4*x**2 - 2*a*b*d**3*e - 4*a*b*d**2*e**2*x - 2*a*b*d*e**3*x**2 + b**2*d**4 + 2*b**2*d**3*e*x + b**2*d**2*e**2*x**2))`

3.220 $\int \frac{A+Bx}{(a+bx)^{3/2}(d+ex)^{7/2}} dx$

Optimal result	1985
Mathematica [A] (verified)	1986
Rubi [A] (verified)	1986
Maple [A] (verified)	1988
Fricas [B] (verification not implemented)	1989
Sympy [F]	1989
Maxima [F(-2)]	1990
Giac [B] (verification not implemented)	1990
Mupad [B] (verification not implemented)	1991
Reduce [B] (verification not implemented)	1992

Optimal result

Integrand size = 24, antiderivative size = 187

$$\int \frac{A+Bx}{(a+bx)^{3/2}(d+ex)^{7/2}} dx = -\frac{2(Ab-aB)}{b(bd-ae)\sqrt{a+bx}(d+ex)^{5/2}} + \frac{2(bBd-6Abe+5aBe)\sqrt{a+bx}}{5b(bd-ae)^2(d+ex)^{5/2}} + \frac{8(bBd-6Abe+5aBe)\sqrt{a+bx}}{15(bd-ae)^3(d+ex)^{3/2}} + \frac{16b(bBd-6Abe+5aBe)\sqrt{a+bx}}{15(bd-ae)^4\sqrt{d+ex}}$$

output

```
(-2*A*b+2*B*a)/b/(-a*e+b*d)/(b*x+a)^(1/2)/(e*x+d)^(5/2)+2/5*(-6*A*b*e+5*B*a*e+B*b*d)*(b*x+a)^(1/2)/b/(-a*e+b*d)^2/(e*x+d)^(5/2)+8/15*(-6*A*b*e+5*B*a*e+B*b*d)*(b*x+a)^(1/2)/(-a*e+b*d)^3/(e*x+d)^(3/2)+16/15*b*(-6*A*b*e+5*B*a*e+B*b*d)*(b*x+a)^(1/2)/(-a*e+b*d)^4/(e*x+d)^(1/2)
```

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.06

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{7/2}} dx = \frac{2(-3Bde^2(a + bx)^3 + 3Ae^3(a + bx)^3 + 10bBde(a + bx)^2(d + ex) - 15Abe^2(a + bx)^2(d + ex) + 5aBe^2(a + bx)^2(d + ex))}{15(bd - ae)^4 \sqrt{a + bx} (d + ex)^{5/2}}$$

input

```
Integrate[(A + B*x)/((a + b*x)^(3/2)*(d + e*x)^(7/2)),x]
```

output

```
(-2*(-3*B*d*e^2*(a + b*x)^3 + 3*A*e^3*(a + b*x)^3 + 10*b*B*d*e*(a + b*x)^2*(d + e*x) - 15*A*b*e^2*(a + b*x)^2*(d + e*x) + 5*a*B*e^2*(a + b*x)^2*(d + e*x) - 15*b^2*B*d*(a + b*x)*(d + e*x)^2 + 45*A*b^2*e*(a + b*x)*(d + e*x)^2 - 30*a*b*B*e*(a + b*x)*(d + e*x)^2 + 15*A*b^3*(d + e*x)^3 - 15*a*b^2*B*(d + e*x)^3))/(15*(b*d - a*e)^4*sqrt[a + b*x]*(d + e*x)^(5/2))
```

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 185, normalized size of antiderivative = 0.99, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {87, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{7/2}} dx$$

$$\downarrow 87$$

$$\frac{(5aBe - 6Abe + bBd) \int \frac{1}{\sqrt{a+bx}(d+ex)^{7/2}} dx}{b(bd - ae)} - \frac{2(Ab - aB)}{b\sqrt{a + bx}(d + ex)^{5/2}(bd - ae)}$$

$$\downarrow 55$$

$$\begin{aligned}
 & \frac{(5aBe - 6Abe + bBd) \left(\frac{4b \int \frac{1}{\sqrt{a+bx}(d+ex)^{5/2}} dx}{5(bd-ae)} + \frac{2\sqrt{a+bx}}{5(d+ex)^{5/2}(bd-ae)} \right)}{\frac{b(bd-ae)}{2(Ab-aB)} \frac{1}{b\sqrt{a+bx}(d+ex)^{5/2}(bd-ae)}} \\
 & \quad \downarrow 55 \\
 & \frac{(5aBe - 6Abe + bBd) \left(\frac{4b \left(\frac{2b \int \frac{1}{\sqrt{a+bx}(d+ex)^{3/2}} dx}{3(bd-ae)} + \frac{2\sqrt{a+bx}}{3(d+ex)^{3/2}(bd-ae)} \right)}{5(bd-ae)} + \frac{2\sqrt{a+bx}}{5(d+ex)^{5/2}(bd-ae)} \right)}{\frac{b(bd-ae)}{2(Ab-aB)} \frac{1}{b\sqrt{a+bx}(d+ex)^{5/2}(bd-ae)}} \\
 & \quad \downarrow 48 \\
 & \frac{\left(\frac{4b \left(\frac{4b\sqrt{a+bx}}{3\sqrt{d+ex}(bd-ae)^2} + \frac{2\sqrt{a+bx}}{3(d+ex)^{3/2}(bd-ae)} \right)}{5(bd-ae)} + \frac{2\sqrt{a+bx}}{5(d+ex)^{5/2}(bd-ae)} \right) (5aBe - 6Abe + bBd)}{\frac{b(bd-ae)}{2(Ab-aB)} \frac{1}{b\sqrt{a+bx}(d+ex)^{5/2}(bd-ae)}}
 \end{aligned}$$

```
input Int[(A + B*x)/((a + b*x)^(3/2)*(d + e*x)^(7/2)),x]
```

```
output (-2*(A*b - a*B))/(b*(b*d - a*e)*Sqrt[a + b*x]*(d + e*x)^(5/2)) + ((b*B*d - 6*A*b*e + 5*a*B*e)*((2*Sqrt[a + b*x])/(5*(b*d - a*e)*(d + e*x)^(5/2)) + (4*b*((2*Sqrt[a + b*x])/(3*(b*d - a*e)*(d + e*x)^(3/2)) + (4*b*Sqrt[a + b*x])/((3*(b*d - a*e)^2*Sqrt[d + e*x])))/(5*(b*d - a*e))))/(b*(b*d - a*e))
```

Defintions of rubi rules used

```
rule 48 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```


Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{7/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(7/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f or more de

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1353 vs. 2(165) = 330.

Time = 0.37 (sec) , antiderivative size = 1353, normalized size of antiderivative = 7.24

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{7/2}} dx = \text{Too large to display}$$

input `integrate((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(7/2),x, algorithm="giac")`

output

```
4*(B^2*a^2*b^7*e - 2*A*B*a*b^8*e + A^2*b^9*e)/((sqrt(b*e)*B*a*b^5*d - sqrt
(b*e)*A*b^6*d - sqrt(b*e)*B*a^2*b^4*e + sqrt(b*e)*A*a*b^5*e - sqrt(b*e)*(s
qrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*B*a*b^3 +
sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^
2*A*b^4)*(b^3*d^3*abs(b) - 3*a*b^2*d^2*e*abs(b) + 3*a^2*b*d*e^2*abs(b) - a
^3*e^3*abs(b))) + 2/15*((b*x + a)*((8*B*b^13*d^6*e^4 - 15*B*a*b^12*d^5*e^5
- 33*A*b^13*d^5*e^5 - 45*B*a^2*b^11*d^4*e^6 + 165*A*a*b^12*d^4*e^6 + 170*
B*a^3*b^10*d^3*e^7 - 330*A*a^2*b^11*d^3*e^7 - 210*B*a^4*b^9*d^2*e^8 + 330*
A*a^3*b^10*d^2*e^8 + 117*B*a^5*b^8*d*e^9 - 165*A*a^4*b^9*d*e^9 - 25*B*a^6*
b^7*e^10 + 33*A*a^5*b^8*e^10)*(b*x + a)/(b^11*d^9*e^2*abs(b) - 9*a*b^10*d^
8*e^3*abs(b) + 36*a^2*b^9*d^7*e^4*abs(b) - 84*a^3*b^8*d^6*e^5*abs(b) + 126
*a^4*b^7*d^5*e^6*abs(b) - 126*a^5*b^6*d^4*e^7*abs(b) + 84*a^6*b^5*d^3*e^8*
abs(b) - 36*a^7*b^4*d^2*e^9*abs(b) + 9*a^8*b^3*d*e^10*abs(b) - a^9*b^2*e^1
1*abs(b)) + 5*(4*B*b^14*d^7*e^3 - 13*B*a*b^13*d^6*e^4 - 15*A*b^14*d^6*e^4
- 6*B*a^2*b^12*d^5*e^5 + 90*A*a*b^13*d^5*e^5 + 85*B*a^3*b^11*d^4*e^6 - 225
*A*a^2*b^12*d^4*e^6 - 160*B*a^4*b^10*d^3*e^7 + 300*A*a^3*b^11*d^3*e^7 + 14
1*B*a^5*b^9*d^2*e^8 - 225*A*a^4*b^10*d^2*e^8 - 62*B*a^6*b^8*d*e^9 + 90*A*a
^5*b^9*d*e^9 + 11*B*a^7*b^7*e^10 - 15*A*a^6*b^8*e^10)/(b^11*d^9*e^2*abs(b)
- 9*a*b^10*d^8*e^3*abs(b) + 36*a^2*b^9*d^7*e^4*abs(b) - 84*a^3*b^8*d^6*e^
5*abs(b) + 126*a^4*b^7*d^5*e^6*abs(b) - 126*a^5*b^6*d^4*e^7*abs(b) + 84...
```

Mupad [B] (verification not implemented)

Time = 1.79 (sec) , antiderivative size = 288, normalized size of antiderivative = 1.54

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{7/2}} dx = \frac{\sqrt{d + ex} \left(\frac{2x(-a^2 e^2 + 10 a b d e + 15 b^2 d^2)(5 B a e - 6 A b e + B b d)}{15 e^3 (a e - b d)^4} - \frac{4 B a^3 d e^2 + 6 A a^3 e^3 - 40 B}{x^3 \sqrt{a + b x}} \right)}{x^3 \sqrt{a + b x}}$$

input

```
int((A + B*x)/((a + b*x)^(3/2)*(d + e*x)^(7/2)),x)
```

output

```
((d + e*x)^(1/2)*((2*x*(15*b^2*d^2 - a^2*e^2 + 10*a*b*d*e)*(5*B*a*e - 6*A*
b*e + B*b*d))/(15*e^3*(a*e - b*d)^4) - (6*A*a^3*e^3 + 30*A*b^3*d^3 - 60*B*
a*b^2*d^3 + 4*B*a^3*d*e^2 + 90*A*a*b^2*d^2*e - 30*A*a^2*b*d*e^2 - 40*B*a^2
*b*d^2*e))/(15*e^3*(a*e - b*d)^4) + (16*b^2*x^3*(5*B*a*e - 6*A*b*e + B*b*d)
)/(15*e*(a*e - b*d)^4) + (8*b*x^2*(a*e + 5*b*d)*(5*B*a*e - 6*A*b*e + B*b*d)
))/(15*e^2*(a*e - b*d)^4))/(x^3*(a + b*x)^(1/2) + (d^3*(a + b*x)^(1/2))/e
^3 + (3*d*x^2*(a + b*x)^(1/2))/e + (3*d^2*x*(a + b*x)^(1/2))/e^2)
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 388, normalized size of antiderivative = 2.07

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{7/2}} dx = \frac{-\frac{2\sqrt{ex+d}\sqrt{bx+a}a^2e^3}{5} + \frac{4\sqrt{ex+d}\sqrt{bx+a}abde^2}{3} + \frac{8\sqrt{ex+d}\sqrt{bx+a}abe^3x}{15} - 2\sqrt{ex+d}\sqrt{bx+a}}{e(a^3e^6x^3 - 3a^2bde^5x^3 + 3ab^2d^2e^4x^3 - b^3d^3e^3x^3 + 3a^3de^5x^2 - 9a^2bd^2e^4x^2)}$$

input `int((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(7/2),x)`

output

```
(2*( - 3*sqrt(d + e*x)*sqrt(a + b*x)*a**2*e**3 + 10*sqrt(d + e*x)*sqrt(a +
b*x)*a*b*d*e**2 + 4*sqrt(d + e*x)*sqrt(a + b*x)*a*b*e**3*x - 15*sqrt(d +
e*x)*sqrt(a + b*x)*b**2*d**2*e - 20*sqrt(d + e*x)*sqrt(a + b*x)*b**2*d*e**
2*x - 8*sqrt(d + e*x)*sqrt(a + b*x)*b**2*e**3*x**2 + 8*sqrt(e)*sqrt(b)*b**
2*d**3 + 24*sqrt(e)*sqrt(b)*b**2*d**2*e*x + 24*sqrt(e)*sqrt(b)*b**2*d*e**2
*x**2 + 8*sqrt(e)*sqrt(b)*b**2*e**3*x**3))/(15*e*(a**3*d**3*e**3 + 3*a**3*
d**2*e**4*x + 3*a**3*d*e**5*x**2 + a**3*e**6*x**3 - 3*a**2*b*d**4*e**2 - 9
*a**2*b*d**3*e**3*x - 9*a**2*b*d**2*e**4*x**2 - 3*a**2*b*d*e**5*x**3 + 3*a
*b**2*d**5*e + 9*a*b**2*d**4*e**2*x + 9*a*b**2*d**3*e**3*x**2 + 3*a*b**2*d
**2*e**4*x**3 - b**3*d**6 - 3*b**3*d**5*e*x - 3*b**3*d**4*e**2*x**2 - b**3
*d**3*e**3*x**3))
```

3.221 $\int \frac{A+Bx}{(a+bx)^{3/2}(d+ex)^{9/2}} dx$

Optimal result	1993
Mathematica [A] (verified)	1994
Rubi [A] (verified)	1994
Maple [B] (verified)	1996
Fricas [B] (verification not implemented)	1997
Sympy [F]	1998
Maxima [F(-2)]	1999
Giac [B] (verification not implemented)	1999
Mupad [B] (verification not implemented)	2000
Reduce [B] (verification not implemented)	2001

Optimal result

Integrand size = 24, antiderivative size = 237

$$\int \frac{A+Bx}{(a+bx)^{3/2}(d+ex)^{9/2}} dx = -\frac{2(Ab-aB)}{b(bd-ae)\sqrt{a+bx}(d+ex)^{7/2}} + \frac{2(bBd-8Abe+7aBe)\sqrt{a+bx}}{7b(bd-ae)^2(d+ex)^{7/2}} + \frac{12(bBd-8Abe+7aBe)\sqrt{a+bx}}{35(bd-ae)^3(d+ex)^{5/2}} + \frac{16b(bBd-8Abe+7aBe)\sqrt{a+bx}}{35(bd-ae)^4(d+ex)^{3/2}} + \frac{32b^2(bBd-8Abe+7aBe)\sqrt{a+bx}}{35(bd-ae)^5\sqrt{d+ex}}$$

output

```
(-2*A*b+2*B*a)/b/(-a*e+b*d)/(b*x+a)^(1/2)/(e*x+d)^(7/2)+2/7*(-8*A*b*e+7*B*a*e+B*b*d)*(b*x+a)^(1/2)/b/(-a*e+b*d)^2/(e*x+d)^(7/2)+12/35*(-8*A*b*e+7*B*a*e+B*b*d)*(b*x+a)^(1/2)/(-a*e+b*d)^3/(e*x+d)^(5/2)+16/35*b*(-8*A*b*e+7*B*a*e+B*b*d)*(b*x+a)^(1/2)/(-a*e+b*d)^4/(e*x+d)^(3/2)+32/35*b^2*(-8*A*b*e+7*B*a*e+B*b*d)*(b*x+a)^(1/2)/(-a*e+b*d)^5/(e*x+d)^(1/2)
```

Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.14

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{9/2}} dx = \frac{2(5Bde^3(a + bx)^4 - 5Ae^4(a + bx)^4 - 21bBde^2(a + bx)^3(d + ex) + 28Abe^3(a + bx)^3(d + ex) - 7aBe^3(a$$

input `Integrate[(A + B*x)/((a + b*x)^(3/2)*(d + e*x)^(9/2)),x]`

output `(-2*(5*B*d*e^3*(a + b*x)^4 - 5*A*e^4*(a + b*x)^4 - 21*b*B*d*e^2*(a + b*x)^3*(d + e*x) + 28*A*b*e^3*(a + b*x)^3*(d + e*x) - 7*a*B*e^3*(a + b*x)^3*(d + e*x) + 35*b^2*B*d*e*(a + b*x)^2*(d + e*x)^2 - 70*A*b^2*e^2*(a + b*x)^2*(d + e*x)^2 + 35*a*b*B*e^2*(a + b*x)^2*(d + e*x)^2 - 35*b^3*B*d*(a + b*x)*(d + e*x)^3 + 140*A*b^3*e*(a + b*x)*(d + e*x)^3 - 105*a*b^2*B*e*(a + b*x)*(d + e*x)^3 + 35*A*b^4*(d + e*x)^4 - 35*a*b^3*B*(d + e*x)^4))/(35*(b*d - a*e)^5*sqrt[a + b*x]*(d + e*x)^(7/2))`

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 233, normalized size of antiderivative = 0.98, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$, Rules used = {87, 55, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{9/2}} dx$$

↓ 87

$$\frac{(7aBe - 8Abe + bBd) \int \frac{1}{\sqrt{a+bx}(d+ex)^{9/2}} dx}{b(bd - ae)} - \frac{2(Ab - aB)}{b\sqrt{a + bx}(d + ex)^{7/2}(bd - ae)}$$

↓ 55

$$\begin{aligned}
 & \frac{(7aBe - 8Abe + bBd) \left(\frac{6b \int \frac{1}{\sqrt{a+bx}(d+ex)^{7/2}} dx}{7(bd-ae)} + \frac{2\sqrt{a+bx}}{7(d+ex)^{7/2}(bd-ae)} \right)}{\frac{b(bd-ae)}{2(Ab-aB)} \frac{1}{b\sqrt{a+bx}(d+ex)^{7/2}(bd-ae)}} \\
 & \quad \downarrow 55 \\
 & \frac{(7aBe - 8Abe + bBd) \left(\frac{6b \left(\frac{4b \int \frac{1}{\sqrt{a+bx}(d+ex)^{5/2}} dx}{5(bd-ae)} + \frac{2\sqrt{a+bx}}{5(d+ex)^{5/2}(bd-ae)} \right)}{7(bd-ae)} + \frac{2\sqrt{a+bx}}{7(d+ex)^{7/2}(bd-ae)} \right)}{\frac{b(bd-ae)}{2(Ab-aB)} \frac{1}{b\sqrt{a+bx}(d+ex)^{7/2}(bd-ae)}} \\
 & \quad \downarrow 55 \\
 & \frac{(7aBe - 8Abe + bBd) \left(\frac{6b \left(\frac{4b \left(\frac{2b \int \frac{1}{\sqrt{a+bx}(d+ex)^{3/2}} dx}{3(bd-ae)} + \frac{2\sqrt{a+bx}}{3(d+ex)^{3/2}(bd-ae)} \right)}{5(bd-ae)} + \frac{2\sqrt{a+bx}}{5(d+ex)^{5/2}(bd-ae)} \right)}{7(bd-ae)} + \frac{2\sqrt{a+bx}}{7(d+ex)^{7/2}(bd-ae)} \right)}{\frac{b(bd-ae)}{2(Ab-aB)} \frac{1}{b\sqrt{a+bx}(d+ex)^{7/2}(bd-ae)}} \\
 & \quad \downarrow 48 \\
 & \left(\frac{6b \left(\frac{4b \left(\frac{4b\sqrt{a+bx}}{3\sqrt{d+ex}(bd-ae)^2} + \frac{2\sqrt{a+bx}}{3(d+ex)^{3/2}(bd-ae)} \right)}{5(bd-ae)} + \frac{2\sqrt{a+bx}}{5(d+ex)^{5/2}(bd-ae)} \right)}{7(bd-ae)} + \frac{2\sqrt{a+bx}}{7(d+ex)^{7/2}(bd-ae)} \right) (7aBe - 8Abe + bBd) \\
 & \quad \frac{b(bd-ae)}{2(Ab-aB)} \frac{1}{b\sqrt{a+bx}(d+ex)^{7/2}(bd-ae)}
 \end{aligned}$$

input

```
Int[(A + B*x)/((a + b*x)^(3/2)*(d + e*x)^(9/2)),x]
```


output

$$\begin{aligned} & (-2*(A*b - a*B))/(b*(b*d - a*e)*\text{Sqrt}[a + b*x]*(d + e*x)^{(7/2)}) + ((b*B*d - \\ & 8*A*b*e + 7*a*B*e)*((2*\text{Sqrt}[a + b*x])/(7*(b*d - a*e)*(d + e*x)^{(7/2)})) + (\\ & 6*b*((2*\text{Sqrt}[a + b*x])/(5*(b*d - a*e)*(d + e*x)^{(5/2)})) + (4*b*((2*\text{Sqrt}[a + \\ & b*x])/(3*(b*d - a*e)*(d + e*x)^{(3/2)})) + (4*b*\text{Sqrt}[a + b*x])/(3*(b*d - a*e) \\ &)^2*\text{Sqrt}[d + e*x]))/(5*(b*d - a*e)))/(7*(b*d - a*e)))/(b*(b*d - a*e)) \end{aligned}$$

Defintions of rubi rules used

rule 48

$$\text{Int}[(a + b*x)^m * (c + d*x)^n, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{m+1} * (c + d*x)^{n+1} / ((b*c - a*d)*(m+1)), x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x\} \ \&\& \ \text{EqQ}[m + n + 2, 0] \ \&\& \ \text{NeQ}[m, -1]$$

rule 55

$$\begin{aligned} & \text{Int}[(a + b*x)^m * (c + d*x)^n, x_Symbol] \rightarrow \text{Simp}[\\ & (a + b*x)^{m+1} * (c + d*x)^{n+1} / ((b*c - a*d)*(m+1)), x] - \text{Simp}[d * (S \\ & \text{implify}[m + n + 2] / ((b*c - a*d)*(m+1))) \ \text{Int}[(a + b*x)^{\text{Simplify}[m + 1]} * \\ & (c + d*x)^n, x], x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x\} \ \&\& \ \text{ILtQ}[\text{Simplify}[m + n + \\ & 2], 0] \ \&\& \ \text{NeQ}[m, -1] \ \&\& \ !(\text{LtQ}[m, -1] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ (\text{EqQ}[a, 0] \ || \ (\text{NeQ}[\\ & c, 0] \ \&\& \ \text{LtQ}[m - n, 0] \ \&\& \ \text{IntegerQ}[n]))) \ \&\& \ (\text{SumSimplerQ}[m, 1] \ || \ !\text{SumSimp \\ & lerQ}[n, 1]) \end{aligned}$$

rule 87

$$\begin{aligned} & \text{Int}[(a + b*x)^m * (c + d*x)^n * (e + f*x)^p, x_Symbol] \rightarrow \text{Simp}[\\ & (-b*e - a*f) * (c + d*x)^{n+1} * (e + f*x)^{p+1} / (f * (p \\ & + 1) * (c*f - d*e)), x] - \text{Simp}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p \\ & + 1)) / (f * (p + 1) * (c*f - d*e)) \ \text{Int}[(c + d*x)^n * (e + f*x)^{p+1}, x], x] \\ & /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x\} \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ (!\text{LtQ}[n, -1] \ || \ \text{Intege \\ & rQ}[p] \ || \ !(\text{IntegerQ}[n] \ || \ !(\text{EqQ}[e, 0] \ || \ !(\text{EqQ}[c, 0] \ || \ \text{LtQ}[p, n]))) \end{aligned}$$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 448 vs. $2(209) = 418$.

Time = 0.34 (sec) , antiderivative size = 449, normalized size of antiderivative = 1.89

method	result
default	$-\frac{2(-128Ab^4e^4x^4+112Bab^3e^4x^4+16Bb^4de^3x^4-64Aab^3e^4x^3-448Ab^4de^3x^3+56Ba^2b^2e^4x^3+400Bab^3de^3x^3+56Bb^4d^2e^2x^3}{(b^2x+a)^{5/2}(ex+d)^{9/2}}$
gospers	$-\frac{2(-128Ab^4e^4x^4+112Bab^3e^4x^4+16Bb^4de^3x^4-64Aab^3e^4x^3-448Ab^4de^3x^3+56Ba^2b^2e^4x^3+400Bab^3de^3x^3+56Bb^4d^2e^2x^3}{(b^2x+a)^{5/2}(ex+d)^{9/2}}$
orering	$-\frac{2(-128Ab^4e^4x^4+112Bab^3e^4x^4+16Bb^4de^3x^4-64Aab^3e^4x^3-448Ab^4de^3x^3+56Ba^2b^2e^4x^3+400Bab^3de^3x^3+56Bb^4d^2e^2x^3}{(b^2x+a)^{5/2}(ex+d)^{9/2}}$

input `int((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(9/2),x,method=_RETURNVERBOSE)`

output
$$-\frac{2}{35}(-128A^2b^4e^4x^4+112B^2a^2b^3e^4x^4+16B^2b^4de^3x^4-64A^2a^2b^3e^4x^3-448A^2Ab^4de^3x^3+56B^2Ba^2b^2e^4x^3+400B^2Bab^3de^3x^3+56B^2b^4d^2e^2x^3+16A^2a^2b^2e^4x^2-224A^2a^2b^3de^3x^2-560A^2Ab^4d^2e^2x^2-14B^2a^2b^3e^4x^2+194B^2a^2b^2de^3x^2+518B^2Ba^2b^3d^2e^2x^2+70B^2b^4d^3e^3x^2-8A^2a^3b^3e^4x+56A^2a^2b^2de^3x-280A^2a^2b^3d^2e^2x-280A^2Ab^4d^3e^3x+7B^2a^4e^4x-48B^2a^3b^3de^3x+238B^2a^2b^2d^2e^2x+280B^2a^2b^3d^3e^3x+35B^2b^4d^4x+5A^2a^4e^4-28A^2a^3b^3de^3+70A^2a^2b^2d^2e^2-140A^2a^2b^3d^3e^3-35A^2Ab^4d^4+2B^2a^4de^3-14B^2a^3b^3d^2e^2+70B^2a^2b^2d^3e^3+70B^2a^2b^3d^4)/(e*x+d)^(7/2)/(b*x+a)^(1/2)/(a*e-b*d)^5$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 887 vs. $2(209) = 418$.

Time = 17.48 (sec) , antiderivative size = 887, normalized size of antiderivative = 3.74

$$\int \frac{A+Bx}{(a+bx)^{3/2}(d+ex)^{9/2}} dx = \text{Too large to display}$$

input `integrate((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(9/2),x, algorithm="fricas")`

output

```

2/35*(5*A*a^4*e^4 + 35*(2*B*a*b^3 - A*b^4)*d^4 + 70*(B*a^2*b^2 - 2*A*a*b^3
)*d^3*e - 14*(B*a^3*b - 5*A*a^2*b^2)*d^2*e^2 + 2*(B*a^4 - 14*A*a^3*b)*d*e^
3 + 16*(B*b^4*d*e^3 + (7*B*a*b^3 - 8*A*b^4)*e^4)*x^4 + 8*(7*B*b^4*d^2*e^2
+ 2*(25*B*a*b^3 - 28*A*b^4)*d*e^3 + (7*B*a^2*b^2 - 8*A*a*b^3)*e^4)*x^3 + 2
*(35*B*b^4*d^3*e + 7*(37*B*a*b^3 - 40*A*b^4)*d^2*e^2 + (97*B*a^2*b^2 - 112
*A*a*b^3)*d*e^3 - (7*B*a^3*b - 8*A*a^2*b^2)*e^4)*x^2 + (35*B*b^4*d^4 + 280
*(B*a*b^3 - A*b^4)*d^3*e + 14*(17*B*a^2*b^2 - 20*A*a*b^3)*d^2*e^2 - 8*(6*B
*a^3*b - 7*A*a^2*b^2)*d*e^3 + (7*B*a^4 - 8*A*a^3*b)*e^4)*x)*sqrt(b*x + a)*
sqrt(e*x + d)/(a*b^5*d^9 - 5*a^2*b^4*d^8*e + 10*a^3*b^3*d^7*e^2 - 10*a^4*b
^2*d^6*e^3 + 5*a^5*b*d^5*e^4 - a^6*d^4*e^5 + (b^6*d^5*e^4 - 5*a*b^5*d^4*e^
5 + 10*a^2*b^4*d^3*e^6 - 10*a^3*b^3*d^2*e^7 + 5*a^4*b^2*d*e^8 - a^5*b*e^9)
*x^5 + (4*b^6*d^6*e^3 - 19*a*b^5*d^5*e^4 + 35*a^2*b^4*d^4*e^5 - 30*a^3*b^3
*d^3*e^6 + 10*a^4*b^2*d^2*e^7 + a^5*b*d*e^8 - a^6*e^9)*x^4 + 2*(3*b^6*d^7*
e^2 - 13*a*b^5*d^6*e^3 + 20*a^2*b^4*d^5*e^4 - 10*a^3*b^3*d^4*e^5 - 5*a^4*b
^2*d^3*e^6 + 7*a^5*b*d^2*e^7 - 2*a^6*d*e^8)*x^3 + 2*(2*b^6*d^8*e - 7*a*b^5
*d^7*e^2 + 5*a^2*b^4*d^6*e^3 + 10*a^3*b^3*d^5*e^4 - 20*a^4*b^2*d^4*e^5 + 1
3*a^5*b*d^3*e^6 - 3*a^6*d^2*e^7)*x^2 + (b^6*d^9 - a*b^5*d^8*e - 10*a^2*b^4
*d^7*e^2 + 30*a^3*b^3*d^6*e^3 - 35*a^4*b^2*d^5*e^4 + 19*a^5*b*d^4*e^5 - 4*
a^6*d^3*e^6)*x)

```

Sympy [F]

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{9/2}} dx = \int \frac{A + Bx}{(a + bx)^{\frac{3}{2}}(d + ex)^{\frac{9}{2}}} dx$$

input

```
integrate((B*x+A)/(b*x+a)**(3/2)/(e*x+d)**(9/2),x)
```

output

```
Integral((A + B*x)/((a + b*x)**(3/2)*(d + e*x)**(9/2)), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{9/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(9/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f or more de

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2645 vs. 2(209) = 418.

Time = 1.00 (sec) , antiderivative size = 2645, normalized size of antiderivative = 11.16

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{9/2}} dx = \text{Too large to display}$$

input `integrate((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(9/2),x, algorithm="giac")`

output

```

4*(B^2*a^2*b^9*e - 2*A*B*a*b^10*e + A^2*b^11*e)/((sqrt(b*e)*B*a*b^6*d - sq
rt(b*e)*A*b^7*d - sqrt(b*e)*B*a^2*b^5*e + sqrt(b*e)*A*a*b^6*e - sqrt(b*e)*
(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*B*a*b^4
+ sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e)
)^2*A*b^5)*(b^4*d^4*abs(b) - 4*a*b^3*d^3*e*abs(b) + 6*a^2*b^2*d^2*e^2*abs(
b) - 4*a^3*b*d*e^3*abs(b) + a^4*e^4*abs(b))) + 2/35*(((b*x + a)*((16*B*b^1
9*d^10*e^6*abs(b) - 67*B*a*b^18*d^9*e^7*abs(b) - 93*A*b^19*d^9*e^7*abs(b)
- 117*B*a^2*b^17*d^8*e^8*abs(b) + 837*A*a*b^18*d^8*e^8*abs(b) + 1428*B*a^3
*b^16*d^7*e^9*abs(b) - 3348*A*a^2*b^17*d^7*e^9*abs(b) - 4452*B*a^4*b^15*d^
6*e^10*abs(b) + 7812*A*a^3*b^16*d^6*e^10*abs(b) + 7686*B*a^5*b^14*d^5*e^11
*abs(b) - 11718*A*a^4*b^15*d^5*e^11*abs(b) - 8358*B*a^6*b^13*d^4*e^12*abs(
b) + 11718*A*a^5*b^14*d^4*e^12*abs(b) + 5892*B*a^7*b^12*d^3*e^13*abs(b) -
7812*A*a^6*b^13*d^3*e^13*abs(b) - 2628*B*a^8*b^11*d^2*e^14*abs(b) + 3348*A
*a^7*b^12*d^2*e^14*abs(b) + 677*B*a^9*b^10*d*e^15*abs(b) - 837*A*a^8*b^11*
d*e^15*abs(b) - 77*B*a^10*b^9*e^16*abs(b) + 93*A*a^9*b^10*e^16*abs(b))*(b*
x + a)/(b^18*d^14*e^3 - 14*a*b^17*d^13*e^4 + 91*a^2*b^16*d^12*e^5 - 364*a^
3*b^15*d^11*e^6 + 1001*a^4*b^14*d^10*e^7 - 2002*a^5*b^13*d^9*e^8 + 3003*a^
6*b^12*d^8*e^9 - 3432*a^7*b^11*d^7*e^10 + 3003*a^8*b^10*d^6*e^11 - 2002*a^
9*b^9*d^5*e^12 + 1001*a^10*b^8*d^4*e^13 - 364*a^11*b^7*d^3*e^14 + 91*a^12*
b^6*d^2*e^15 - 14*a^13*b^5*d*e^16 + a^14*b^4*e^17) + 28*(2*B*b^20*d^11*...

```

Mupad [B] (verification not implemented)

Time = 1.98 (sec) , antiderivative size = 409, normalized size of antiderivative = 1.73

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{9/2}} dx = \frac{\sqrt{d + ex} \left(\frac{4Ba^4de^3 + 10Aa^4e^4 - 28Ba^3bd^2e^2 - 56Aa^3bde^3 + 140Ba^2b^2d^3e + 140Aa^2b^2d^2e^2 + 140Ba^2b^3d^4 - 280Aab^3d^3e - 70Ab^4}{35e^4(ae - bd)^5} \right)}{x^4}$$

input

```
int((A + B*x)/((a + b*x)^(3/2)*(d + e*x)^(9/2)),x)
```

output

```

-((d + e*x)^(1/2)*((10*A*a^4*e^4 - 70*A*b^4*d^4 + 140*B*a*b^3*d^4 + 4*B*a^
4*d*e^3 + 140*B*a^2*b^2*d^3*e - 28*B*a^3*b*d^2*e^2 + 140*A*a^2*b^2*d^2*e^2
- 280*A*a*b^3*d^3*e - 56*A*a^3*b*d*e^3)/(35*e^4*(a*e - b*d)^5) + (32*b^3*
x^4*(7*B*a*e - 8*A*b*e + B*b*d))/(35*e*(a*e - b*d)^5) + (2*x*(7*B*a*e - 8*
A*b*e + B*b*d)*(a^3*e^3 + 35*b^3*d^3 + 35*a*b^2*d^2*e - 7*a^2*b*d*e^2))/(3
5*e^4*(a*e - b*d)^5) + (16*b^2*x^3*(a*e + 7*b*d)*(7*B*a*e - 8*A*b*e + B*b*
d))/(35*e^2*(a*e - b*d)^5) + (4*b*x^2*(35*b^2*d^2 - a^2*e^2 + 14*a*b*d*e)*
(7*B*a*e - 8*A*b*e + B*b*d))/(35*e^3*(a*e - b*d)^5)))/(x^4*(a + b*x)^(1/2)
+ (d^4*(a + b*x)^(1/2))/e^4 + (6*d^2*x^2*(a + b*x)^(1/2))/e^2 + (4*d*x^3*
(a + b*x)^(1/2))/e + (4*d^3*x*(a + b*x)^(1/2))/e^3

```

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 643, normalized size of antiderivative = 2.71

$$\int \frac{A + Bx}{(a + bx)^{3/2}(d + ex)^{9/2}} dx = \frac{-\frac{2\sqrt{ex+d}\sqrt{bx+a}a^3e^4}{7} + \frac{6\sqrt{ex+d}\sqrt{bx+a}a^2bde^3}{5} + \frac{12\sqrt{ex+d}\sqrt{bx+a}a^2be^4x}{35} - 2\sqrt{ex}}{e(a^4e^8x^4 - 4a^3bde^7x^4 + 6a^2b^2d^2e^6x^4 - 4ab^3d^3e^5x^4 + b^4d^4e^4x^4 + 4a^4de^7x^3)}$$

input

```
int((B*x+A)/(b*x+a)^(3/2)/(e*x+d)^(9/2),x)
```

output

```

(2*(- 5*sqrt(d + e*x)*sqrt(a + b*x)*a**3*e**4 + 21*sqrt(d + e*x)*sqrt(a +
b*x)*a**2*b*d*e**3 + 6*sqrt(d + e*x)*sqrt(a + b*x)*a**2*b*e**4*x - 35*sqr
t(d + e*x)*sqrt(a + b*x)*a*b**2*d**2*e**2 - 28*sqrt(d + e*x)*sqrt(a + b*x)
*a*b**2*d*e**3*x - 8*sqrt(d + e*x)*sqrt(a + b*x)*a*b**2*e**4*x**2 + 35*sqr
t(d + e*x)*sqrt(a + b*x)*b**3*d**3*e + 70*sqrt(d + e*x)*sqrt(a + b*x)*b**3
*d**2*e**2*x + 56*sqrt(d + e*x)*sqrt(a + b*x)*b**3*d*e**3*x**2 + 16*sqrt(d
+ e*x)*sqrt(a + b*x)*b**3*e**4*x**3 - 16*sqrt(e)*sqrt(b)*b**3*d**4 - 64*sq
rt(e)*sqrt(b)*b**3*d**3*e*x - 96*sqrt(e)*sqrt(b)*b**3*d**2*e**2*x**2 - 64
*sqrt(e)*sqrt(b)*b**3*d*e**3*x**3 - 16*sqrt(e)*sqrt(b)*b**3*e**4*x**4))/(3
5*e*(a**4*d**4*e**4 + 4*a**4*d**3*e**5*x + 6*a**4*d**2*e**6*x**2 + 4*a**4*
d*e**7*x**3 + a**4*e**8*x**4 - 4*a**3*b*d**5*e**3 - 16*a**3*b*d**4*e**4*x
- 24*a**3*b*d**3*e**5*x**2 - 16*a**3*b*d**2*e**6*x**3 - 4*a**3*b*d*e**7*x*
**4 + 6*a**2*b**2*d**6*e**2 + 24*a**2*b**2*d**5*e**3*x + 36*a**2*b**2*d**4*
e**4*x**2 + 24*a**2*b**2*d**3*e**5*x**3 + 6*a**2*b**2*d**2*e**6*x**4 - 4*a
*b**3*d**7*e - 16*a*b**3*d**6*e**2*x - 24*a*b**3*d**5*e**3*x**2 - 16*a*b**
3*d**4*e**4*x**3 - 4*a*b**3*d**3*e**5*x**4 + b**4*d**8 + 4*b**4*d**7*e*x +
6*b**4*d**6*e**2*x**2 + 4*b**4*d**5*e**3*x**3 + b**4*d**4*e**4*x**4))

```

3.222 $\int \frac{(A+Bx)(d+ex)^{7/2}}{(a+bx)^{5/2}} dx$

Optimal result	2002
Mathematica [A] (verified)	2003
Rubi [A] (verified)	2003
Maple [B] (verified)	2008
Fricas [B] (verification not implemented)	2009
Sympy [F]	2010
Maxima [F(-2)]	2010
Giac [B] (verification not implemented)	2010
Mupad [F(-1)]	2011
Reduce [B] (verification not implemented)	2012

Optimal result

Integrand size = 24, antiderivative size = 261

$$\int \frac{(A+Bx)(d+ex)^{7/2}}{(a+bx)^{5/2}} dx = \frac{35e(bd-ae)(bBd+2Abe-3aBe)\sqrt{a+bx}\sqrt{d+ex}}{8b^5} + \frac{35e(bBd+2Abe-3aBe)\sqrt{a+bx}(d+ex)^{3/2}}{12b^4} - \frac{2(3bBd+7Abe-10aBe)(d+ex)^{5/2}}{3b^3\sqrt{a+bx}} + \frac{Be\sqrt{a+bx}(d+ex)^{5/2}}{3b^3} - \frac{2(Ab-aB)(d+ex)^{7/2}}{3b^2(a+bx)^{3/2}} + \frac{35\sqrt{e}(bd-ae)^2(bBd+2Abe-3aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{8b^{11/2}}$$

output

```
35/8*e*(-a*e+b*d)*(2*A*b*e-3*B*a*e+B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(1/2)/b^5+
35/12*e*(2*A*b*e-3*B*a*e+B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(3/2)/b^4-2/3*(7*A*b
*e-10*B*a*e+3*B*b*d)*(e*x+d)^(5/2)/b^3/(b*x+a)^(1/2)+1/3*B*e*(b*x+a)^(1/2)
*(e*x+d)^(5/2)/b^3-2/3*(A*b-B*a)*(e*x+d)^(7/2)/b^2/(b*x+a)^(3/2)+35/8*e^(1
/2)*(-a*e+b*d)^2*(2*A*b*e-3*B*a*e+B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(
1/2)/(e*x+d)^(1/2))/b^(11/2)
```

Mathematica [A] (verified)

Time = 0.73 (sec) , antiderivative size = 313, normalized size of antiderivative = 1.20

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^{5/2}} dx = \frac{\sqrt{d + ex}(-2Ab(105a^3e^3 + 35a^2be^2(-5d + 4ex) + 7ab^2e(8d^2 - 34dex + 3e^2x^2) + 35\sqrt{e}(bd - ae)^2(bBd + 2Abe - 3aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{e}\sqrt{a+bx}}\right))}{8b^{11/2}}$$

input

```
Integrate[((A + B*x)*(d + e*x)^(7/2))/(a + b*x)^(5/2), x]
```

output

```
(Sqrt[d + e*x]*(-2*A*b*(105*a^3*e^3 + 35*a^2*b*e^2*(-5*d + 4*e*x) + 7*a*b^2*e*(8*d^2 - 34*d*e*x + 3*e^2*x^2) + b^3*(8*d^3 + 80*d^2*e*x - 39*d*e^2*x^2 - 6*e^3*x^3)) + B*(315*a^4*e^3 + 210*a^3*b*e^2*(-3*d + 2*e*x) + 7*a^2*b^2*e*(49*d^2 - 122*d*e*x + 9*e^2*x^2) + b^4*x*(-48*d^3 + 87*d^2*e*x + 38*d*e^2*x^2 + 8*e^3*x^3) - 2*a*b^3*(16*d^3 - 239*d^2*e*x + 69*d*e^2*x^2 + 9*e^3*x^3)))/(24*b^5*(a + b*x)^(3/2)) + (35*Sqrt[e]*(b*d - a*e)^2*(b*B*d + 2*A*b*e - 3*a*B*e)*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/(Sqrt[e]*Sqrt[a + b*x])])/(8*b^(11/2))
```

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 257, normalized size of antiderivative = 0.98, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.292$, Rules used = {87, 57, 60, 60, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^{5/2}} dx$$

$$\downarrow 87$$

$$\frac{(-3aBe + 2Abe + bBd) \int \frac{(d+ex)^{7/2}}{(a+bx)^{3/2}} dx}{b(bd - ae)} - \frac{2(d + ex)^{9/2}(Ab - aB)}{3b(a + bx)^{3/2}(bd - ae)}$$

$$\downarrow 57$$

$$\frac{(-3aBe + 2Abe + bBd) \left(\frac{7e \int \frac{(d+ex)^{5/2}}{\sqrt{a+bx}} dx}{b} - \frac{2(d+ex)^{7/2}}{b\sqrt{a+bx}} \right)}{b(bd - ae)} - \frac{2(d+ex)^{9/2}(Ab - aB)}{3b(a+bx)^{3/2}(bd - ae)}$$

↓ 60

$$\frac{(-3aBe + 2Abe + bBd) \left(\frac{7e \left(\frac{5(bd-ae) \int \frac{(d+ex)^{3/2}}{\sqrt{a+bx}} dx}{6b} + \frac{\sqrt{a+bx}(d+ex)^{5/2}}{3b} \right)}{b} - \frac{2(d+ex)^{7/2}}{b\sqrt{a+bx}} \right)}{b(bd - ae)}$$

$$\frac{2(d+ex)^{9/2}(Ab - aB)}{3b(a+bx)^{3/2}(bd - ae)}$$

↓ 60

$$\frac{(-3aBe + 2Abe + bBd) \left(\frac{7e \left(\frac{5(bd-ae) \left(\frac{3(bd-ae) \int \frac{\sqrt{d+ex}}{\sqrt{a+bx}} dx}{4b} + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{6b} + \frac{\sqrt{a+bx}(d+ex)^{5/2}}{3b} \right)}{b} - \frac{2(d+ex)^{7/2}}{b\sqrt{a+bx}} \right)}{b(bd - ae)}$$

$$\frac{2(d+ex)^{9/2}(Ab - aB)}{3b(a+bx)^{3/2}(bd - ae)}$$

↓ 60

$$\left(\frac{(-3aBe + 2Abe + bBd) \left(\frac{5(bd-ae) \left(\frac{3(bd-ae) \left(\frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{4b} + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right) + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{6b} \right) + \frac{\sqrt{a+bx}(d+ex)^{5/2}}{3b}}{b} \right)$$

$$\frac{b(bd - ae)}{3b(a + bx)^{3/2}(bd - ae)} \cdot \frac{2(d + ex)^{9/2}(Ab - aB)}{b(bd - ae)}$$

66

$$\left(\frac{(-3aBe + 2Abe + bBd) \left(\frac{5(bd-ae) \left(\frac{3(bd-ae) \left(\frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{4b} + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right) + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{6b} \right) + \frac{\sqrt{a+bx}(d+ex)^{5/2}}{3b}}{b} \right)$$

$$\frac{b(bd - ae)}{3b(a + bx)^{3/2}(bd - ae)} \cdot \frac{2(d + ex)^{9/2}(Ab - aB)}{b(bd - ae)}$$

221

$$\begin{aligned}
 & \left(\frac{5(bd-ae)}{7e} \left(\frac{3(bd-ae) \left(\frac{(bd-ae) \operatorname{arctanh} \left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}} \right) + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{4b} + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right) + \frac{\sqrt{a+bx}(d+ex)^{5/2}}{3b} \right) \right. \\
 & \left. \frac{(-3aBe + 2Abe + bBd)}{b} \right) \\
 & \frac{2(d+ex)^{9/2}(Ab - aB)}{3b(a+bx)^{3/2}(bd-ae)}
 \end{aligned}$$

input

```
Int[((A + B*x)*(d + e*x)^(7/2))/(a + b*x)^(5/2),x]
```

output

```
(-2*(A*b - a*B)*(d + e*x)^(9/2))/(3*b*(b*d - a*e)*(a + b*x)^(3/2)) + ((b*B*d + 2*A*b*e - 3*a*B*e)*((-2*(d + e*x)^(7/2))/(b*Sqrt[a + b*x]) + (7*e*(Sqrt[a + b*x]*(d + e*x)^(5/2))/(3*b) + (5*(b*d - a*e)*((Sqrt[a + b*x]*(d + e*x)^(3/2))/(2*b) + (3*(b*d - a*e)*((Sqrt[a + b*x]*Sqrt[d + e*x])/b + ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/(b^(3/2)*Sqrt[e])))/(4*b)))/(6*b))/b)/(b*(b*d - a*e))
```

Definitions of rubi rules used

rule 57 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))]
 Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] &&
 GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] &&
 (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1))]
 Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] ||
 (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)]
 Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || (EqQ[e, 0] || (EqQ[c, 0] || LtQ[p, n]))))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1881 vs. $2(217) = 434$.

Time = 0.29 (sec) , antiderivative size = 1882, normalized size of antiderivative = 7.21

method	result	size
default	Expression too large to display	1882

input `int((B*x+A)*(e*x+d)^(7/2)/(b*x+a)^(5/2),x,method=_RETURNVERBOSE)`

output

```
1/48*(e*x+d)^(1/2)*(-560*A*a^2*b^2*e^3*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-320*A*b^4*d^2*e*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+840*B*a^3*b*e^3*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-315*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^5*e^4-276*B*a*b^3*d*e^2*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+952*A*a*b^3*d*e^2*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-1708*B*a^2*b^2*d*e^2*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+956*B*a*b^3*d^2*e*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+420*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^3*b^2*e^4*x-630*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^4*b*e^4*x-420*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^3*b^2*d*e^3+210*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*b^3*d^2*e^2+735*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^4*b*d*e^3-525*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^3*b^2*d^2*e^2+105*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*b^3*d^3*e-32*A*b^4*d^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+630*B*a^4*e^3*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+210*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^4*b*e^4-420*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^4*d*e^3*x^2+735*B*ln(1/2...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 623 vs. $2(217) = 434$.

Time = 1.46 (sec) , antiderivative size = 1271, normalized size of antiderivative = 4.87

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^{5/2}} dx = \text{Too large to display}$$

input `integrate((B*x+A)*(e*x+d)^(7/2)/(b*x+a)^(5/2),x, algorithm="fricas")`

output

```
[1/96*(105*(B*a^2*b^3*d^3 - (5*B*a^3*b^2 - 2*A*a^2*b^3)*d^2*e + (7*B*a^4*b
- 4*A*a^3*b^2)*d*e^2 - (3*B*a^5 - 2*A*a^4*b)*e^3 + (B*b^5*d^3 - (5*B*a*b^
4 - 2*A*b^5)*d^2*e + (7*B*a^2*b^3 - 4*A*a*b^4)*d*e^2 - (3*B*a^3*b^2 - 2*A*
a^2*b^3)*e^3)*x^2 + 2*(B*a*b^4*d^3 - (5*B*a^2*b^3 - 2*A*a*b^4)*d^2*e + (7*
B*a^3*b^2 - 4*A*a^2*b^3)*d*e^2 - (3*B*a^4*b - 2*A*a^3*b^2)*e^3)*x)*sqrt(e/
b)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*b^2*e*x + b^2*
d + a*b*e)*sqrt(b*x + a)*sqrt(e*x + d)*sqrt(e/b) + 8*(b^2*d*e + a*b*e^2)*x
) + 4*(8*B*b^4*e^3*x^4 - 16*(2*B*a*b^3 + A*b^4)*d^3 + 7*(49*B*a^2*b^2 - 16
*A*a*b^3)*d^2*e - 70*(9*B*a^3*b - 5*A*a^2*b^2)*d*e^2 + 105*(3*B*a^4 - 2*A*
a^3*b)*e^3 + 2*(19*B*b^4*d*e^2 - 3*(3*B*a*b^3 - 2*A*b^4)*e^3)*x^3 + 3*(29*
B*b^4*d^2*e - 2*(23*B*a*b^3 - 13*A*b^4)*d*e^2 + 7*(3*B*a^2*b^2 - 2*A*a*b^3
)*e^3)*x^2 - 2*(24*B*b^4*d^3 - (239*B*a*b^3 - 80*A*b^4)*d^2*e + 7*(61*B*a^
2*b^2 - 34*A*a*b^3)*d*e^2 - 70*(3*B*a^3*b - 2*A*a^2*b^2)*e^3)*x)*sqrt(b*x
+ a)*sqrt(e*x + d))/(b^7*x^2 + 2*a*b^6*x + a^2*b^5), -1/48*(105*(B*a^2*b^3
*d^3 - (5*B*a^3*b^2 - 2*A*a^2*b^3)*d^2*e + (7*B*a^4*b - 4*A*a^3*b^2)*d*e^2
- (3*B*a^5 - 2*A*a^4*b)*e^3 + (B*b^5*d^3 - (5*B*a*b^4 - 2*A*b^5)*d^2*e +
(7*B*a^2*b^3 - 4*A*a*b^4)*d*e^2 - (3*B*a^3*b^2 - 2*A*a^2*b^3)*e^3)*x^2 + 2
*(B*a*b^4*d^3 - (5*B*a^2*b^3 - 2*A*a*b^4)*d^2*e + (7*B*a^3*b^2 - 4*A*a^2*b
^3)*d*e^2 - (3*B*a^4*b - 2*A*a^3*b^2)*e^3)*x)*sqrt(-e/b)*arctan(1/2*(2*b*e
*x + b*d + a*e)*sqrt(b*x + a)*sqrt(e*x + d)*sqrt(-e/b)/(b*e^2*x^2 + a*d...
```

Sympy [F]

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^{5/2}} dx = \int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^{5/2}} dx$$

input `integrate((B*x+A)*(e*x+d)**(7/2)/(b*x+a)**(5/2),x)`

output `Integral((A + B*x)*(d + e*x)**(7/2)/(a + b*x)**(5/2), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)*(e*x+d)^(7/2)/(b*x+a)^(5/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1814 vs. 2(217) = 434.

Time = 0.65 (sec) , antiderivative size = 1814, normalized size of antiderivative = 6.95

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^{5/2}} dx = \text{Too large to display}$$

input `integrate((B*x+A)*(e*x+d)^(7/2)/(b*x+a)^(5/2),x, algorithm="giac")`

output

```

1/24*sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*sqrt(b*x + a)*(2*(b*x + a)*(4*(b*
x + a)*B*e^3*abs(b)/b^7 + (19*B*b^21*d*e^6*abs(b) - 25*B*a*b^20*e^7*abs(b)
+ 6*A*b^21*e^7*abs(b))/(b^27*e^4)) + 3*(29*B*b^22*d^2*e^5*abs(b) - 84*B*a
*b^21*d*e^6*abs(b) + 26*A*b^22*d*e^6*abs(b) + 55*B*a^2*b^20*e^7*abs(b) - 2
6*A*a*b^21*e^7*abs(b))/(b^27*e^4) - 35/16*(sqrt(b*e)*B*b^3*d^3*abs(b) - 5
*sqrt(b*e)*B*a*b^2*d^2*e*abs(b) + 2*sqrt(b*e)*A*b^3*d^2*e*abs(b) + 7*sqrt(
b*e)*B*a^2*b*d*e^2*abs(b) - 4*sqrt(b*e)*A*a*b^2*d*e^2*abs(b) - 3*sqrt(b*e)
*B*a^3*e^3*abs(b) + 2*sqrt(b*e)*A*a^2*b*e^3*abs(b))*log((sqrt(b*e)*sqrt(b*
x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2)/b^7 - 4/3*(3*sqrt(b*e)*B*
b^8*d^6*abs(b) - 28*sqrt(b*e)*B*a*b^7*d^5*e*abs(b) + 10*sqrt(b*e)*A*b^8*d^
5*e*abs(b) + 95*sqrt(b*e)*B*a^2*b^6*d^4*e^2*abs(b) - 50*sqrt(b*e)*A*a*b^7*
d^4*e^2*abs(b) - 160*sqrt(b*e)*B*a^3*b^5*d^3*e^3*abs(b) + 100*sqrt(b*e)*A*
a^2*b^6*d^3*e^3*abs(b) + 145*sqrt(b*e)*B*a^4*b^4*d^2*e^4*abs(b) - 100*sqrt
(b*e)*A*a^3*b^5*d^2*e^4*abs(b) - 68*sqrt(b*e)*B*a^5*b^3*d*e^5*abs(b) + 50*
sqrt(b*e)*A*a^4*b^4*d*e^5*abs(b) + 13*sqrt(b*e)*B*a^6*b^2*e^6*abs(b) - 10*
sqrt(b*e)*A*a^5*b^3*e^6*abs(b) - 6*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sq
rt(b^2*d + (b*x + a)*b*e - a*b*e))^2*B*b^6*d^5*abs(b) + 48*sqrt(b*e)*(sqrt
(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*B*a*b^5*d^4*e
*abs(b) - 18*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b
*e - a*b*e))^2*A*b^6*d^4*e*abs(b) - 132*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x +...

```

Mupad [F(-1)]

Timed out.

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^{5/2}} dx = \int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^{5/2}} dx$$

input

```
int(((A + B*x)*(d + e*x)^(7/2))/(a + b*x)^(5/2), x)
```

output

```
int(((A + B*x)*(d + e*x)^(7/2))/(a + b*x)^(5/2), x)
```


Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 461, normalized size of antiderivative = 1.77

$$\int \frac{(A + Bx)(d + ex)^{7/2}}{(a + bx)^{5/2}} dx = \frac{-840\sqrt{e}\sqrt{b}\sqrt{bx+a}\log\left(\frac{\sqrt{e}\sqrt{bx+a}+\sqrt{b}\sqrt{ex+d}}{\sqrt{ae-bd}}\right)a^3e^3 + 2520\sqrt{e}\sqrt{b}\sqrt{bx+a}\log$$

input `int((B*x+A)*(e*x+d)^(7/2)/(b*x+a)^(5/2),x)`

output

```
( - 840*sqrt(e)*sqrt(b)*sqrt(a + b*x)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)
*sqrt(d + e*x))/sqrt(a*e - b*d))*a**3*e**3 + 2520*sqrt(e)*sqrt(b)*sqrt(a +
b*x)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))
*a**2*b*d*e**2 - 2520*sqrt(e)*sqrt(b)*sqrt(a + b*x)*log((sqrt(e)*sqrt(a +
b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*b**2*d**2*e + 840*sqrt(e)
*sqrt(b)*sqrt(a + b*x)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))
/sqrt(a*e - b*d))*b**3*d**3 + 525*sqrt(e)*sqrt(b)*sqrt(a + b*x)*a**3*e**3
- 1575*sqrt(e)*sqrt(b)*sqrt(a + b*x)*a**2*b*d*e**2 + 1575*sqrt(e)*sqrt(b)*
sqrt(a + b*x)*a*b**2*d**2*e - 525*sqrt(e)*sqrt(b)*sqrt(a + b*x)*b**3*d**3
+ 840*sqrt(d + e*x)*a**3*b*e**3 - 2240*sqrt(d + e*x)*a**2*b**2*d*e**2 + 28
0*sqrt(d + e*x)*a**2*b**2*e**3*x + 1848*sqrt(d + e*x)*a*b**3*d**2*e - 784*
sqrt(d + e*x)*a*b**3*d*e**2*x - 112*sqrt(d + e*x)*a*b**3*e**3*x**2 - 384*s
qrt(d + e*x)*b**4*d**3 + 696*sqrt(d + e*x)*b**4*d**2*e*x + 304*sqrt(d + e*
x)*b**4*d*e**2*x**2 + 64*sqrt(d + e*x)*b**4*e**3*x**3)/(192*sqrt(a + b*x)*
b**5)
```

3.223
$$\int \frac{(A+Bx)(d+ex)^{5/2}}{(a+bx)^{5/2}} dx$$

Optimal result	2013
Mathematica [A] (verified)	2014
Rubi [A] (verified)	2014
Maple [B] (verified)	2017
Fricas [B] (verification not implemented)	2018
Sympy [F]	2019
Maxima [F(-2)]	2020
Giac [B] (verification not implemented)	2020
Mupad [F(-1)]	2021
Reduce [B] (verification not implemented)	2022

Optimal result

Integrand size = 24, antiderivative size = 212

$$\int \frac{(A+Bx)(d+ex)^{5/2}}{(a+bx)^{5/2}} dx = \frac{5e(3bBd+4Abe-7aBe)\sqrt{a+bx}\sqrt{d+ex}}{4b^4} - \frac{2(3bBd+5Abe-8aBe)(d+ex)^{3/2}}{3b^3\sqrt{a+bx}} + \frac{Be\sqrt{a+bx}(d+ex)^{3/2}}{2b^3} - \frac{2(Ab-aB)(d+ex)^{5/2}}{3b^2(a+bx)^{3/2}} + \frac{5\sqrt{e}(bd-ae)(3bBd+4Abe-7aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{4b^{9/2}}$$

output

```
5/4*e*(4*A*b*e-7*B*a*e+3*B*b*d)*(b*x+a)^(1/2)*(e*x+d)^(1/2)/b^4-2/3*(5*A*b
*e-8*B*a*e+3*B*b*d)*(e*x+d)^(3/2)/b^3/(b*x+a)^(1/2)+1/2*B*e*(b*x+a)^(1/2)*
(e*x+d)^(3/2)/b^3-2/3*(A*b-B*a)*(e*x+d)^(5/2)/b^2/(b*x+a)^(3/2)+5/4*e^(1/2
)*(-a*e+b*d)*(4*A*b*e-7*B*a*e+3*B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/
2)/(e*x+d)^(1/2))/b^(9/2)
```

Mathematica [A] (verified)

Time = 1.33 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.11

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^{5/2}} dx = \frac{\sqrt{d + ex}(B(-105a^3e^2 + 5a^2be(23d - 28ex) + ab^2(-16d^2 + 158dex - 21e^2x^2)) + 5\sqrt{e}(-bd + ae)(3bBd + 4Abe - 7aBe)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{e}\left(\sqrt{a-\frac{bd}{e}}-\sqrt{a+bx}\right)}\right)}{2b^{9/2}}$$

input

```
Integrate[((A + B*x)*(d + e*x)^(5/2))/(a + b*x)^(5/2), x]
```

output

```
(Sqrt[d + e*x]*(B*(-105*a^3*e^2 + 5*a^2*b*e*(23*d - 28*e*x) + a*b^2*(-16*d^2 + 158*d*e*x - 21*e^2*x^2)) + 3*b^3*x*(-8*d^2 + 9*d*e*x + 2*e^2*x^2)) + 4*A*b*(15*a^2*e^2 - 10*a*b*e*(d - 2*e*x) + b^2*(-2*d^2 - 14*d*e*x + 3*e^2*x^2)))/(12*b^4*(a + b*x)^(3/2)) + (5*Sqrt[e]*(-(b*d) + a*e)*(3*b*B*d + 4*A*b*e - 7*a*B*e)*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/(Sqrt[e]*(Sqrt[a - (b*d)/e] - Sqrt[a + b*x]))])/(2*b^(9/2))
```

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.04, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {87, 57, 60, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^{5/2}} dx$$

$$\downarrow 87$$

$$\frac{(-7aBe + 4Abe + 3bBd) \int \frac{(d+ex)^{5/2}}{(a+bx)^{3/2}} dx}{3b(bd - ae)} - \frac{2(d + ex)^{7/2}(Ab - aB)}{3b(a + bx)^{3/2}(bd - ae)}$$

$$\downarrow 57$$

$$\begin{aligned}
 & \frac{(-7aBe + 4Abe + 3bBd) \left(\frac{5e \int \frac{(d+ex)^{3/2}}{\sqrt{a+bx}} dx}{b} - \frac{2(d+ex)^{5/2}}{b\sqrt{a+bx}} \right)}{3b(bd - ae)} - \frac{2(d+ex)^{7/2}(Ab - aB)}{3b(a+bx)^{3/2}(bd - ae)} \\
 & \quad \downarrow 60 \\
 & \frac{(-7aBe + 4Abe + 3bBd) \left(\frac{5e \left(\frac{3(bd-ae) \int \frac{\sqrt{d+ex}}{\sqrt{a+bx}} dx}{4b} + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{b} - \frac{2(d+ex)^{5/2}}{b\sqrt{a+bx}} \right)}{3b(bd - ae)} - \frac{2(d+ex)^{7/2}(Ab - aB)}{3b(a+bx)^{3/2}(bd - ae)} \\
 & \quad \downarrow 60 \\
 & \frac{(-7aBe + 4Abe + 3bBd) \left(\frac{5e \left(\frac{3(bd-ae) \left(\frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{4b} + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right) + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{b} - \frac{2(d+ex)^{5/2}}{b\sqrt{a+bx}} \right)}{3b(bd - ae)} - \frac{2(d+ex)^{7/2}(Ab - aB)}{3b(a+bx)^{3/2}(bd - ae)} \\
 & \quad \downarrow 66 \\
 & \frac{(-7aBe + 4Abe + 3bBd) \left(\frac{5e \left(\frac{3(bd-ae) \left(\frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}} + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right) + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{b} - \frac{2(d+ex)^{5/2}}{b\sqrt{a+bx}} \right)}{3b(bd - ae)} - \frac{2(d+ex)^{7/2}(Ab - aB)}{3b(a+bx)^{3/2}(bd - ae)} \\
 & \quad \downarrow 221
 \end{aligned}$$

$$\begin{aligned}
 & \left(\frac{(-7aBe + 4Abe + 3bBd)}{b} \left(\frac{5e \left(\frac{3(bd-ae) \left(\frac{(bd-ae) \operatorname{arctanh} \left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}} \right) + \sqrt{a+bx}\sqrt{d+ex}}{b^{3/2}\sqrt{e}} \right) + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{4b} + \frac{\sqrt{a+bx}(d+ex)^{3/2}}{2b} \right)}{b} \right) - \frac{2(d+ex)^{5/2}}{b\sqrt{a+bx}} \right) \\
 & \frac{3b(bd-ae)}{2(d+ex)^{7/2}(Ab-aB)} \\
 & \frac{3b(a+bx)^{3/2}(bd-ae)}{3b(a+bx)^{3/2}(bd-ae)}
 \end{aligned}$$

input `Int[((A + B*x)*(d + e*x)^(5/2))/(a + b*x)^(5/2), x]`

output `(-2*(A*b - a*B)*(d + e*x)^(7/2))/(3*b*(b*d - a*e)*(a + b*x)^(3/2)) + ((3*b*B*d + 4*A*b*e - 7*a*B*e)*((-2*(d + e*x)^(5/2))/(b*Sqrt[a + b*x]) + (5*e*(Sqrt[a + b*x]*(d + e*x)^(3/2))/(2*b) + (3*(b*d - a*e)*((Sqrt[a + b*x]*Sqrt[d + e*x])/b + ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])]))/(b^(3/2)*Sqrt[e]))/(4*b))/b)/(3*b*(b*d - a*e))`

Defintions of rubi rules used

rule 57 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))] Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[
2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; Fre
eQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p
), x] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))`

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1249 vs. $2(174) = 348$.

Time = 0.27 (sec) , antiderivative size = 1250, normalized size of antiderivative = 5.90

method	result	size
default	Expression too large to display	1250

input `int((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^(5/2), x, method=_RETURNVERBOSE)`

output

```

-1/24*(e*x+d)^(1/2)*(60*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^3*e^3*x^2-60*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^4*d*e^2*x^2-105*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*b^2*e^3*x^2-45*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^4*d^2*e*x^2+42*B*a*b^2*e^2*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-54*B*b^3*d*e*x^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-160*A*a*b^2*e^2*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+112*A*b^3*d*e*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+280*B*a^2*b*e^2*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+60*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^3*b*e^3-316*B*a*b^2*d*e*x*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+80*A*a*b^2*d*e*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-230*B*a^2*b*d*e*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)-120*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^3*d*e^2*x+300*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*b^2*d*e^2*x-90*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^3*d^2*e*x+150*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^3*d*e^2*x^2-120*A*a^2*b*e^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+32*B*a*b^2*d^2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+120*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 431 vs. $2(174) = 348$.

Time = 1.26 (sec) , antiderivative size = 887, normalized size of antiderivative = 4.18

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^{5/2}} dx = \text{Too large to display}$$

input

```
integrate((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^(5/2),x, algorithm="fricas")
```

output

```
[-1/48*(15*(3*B*a^2*b^2*d^2 - 2*(5*B*a^3*b - 2*A*a^2*b^2)*d*e + (7*B*a^4 -
4*A*a^3*b)*e^2 + (3*B*b^4*d^2 - 2*(5*B*a*b^3 - 2*A*b^4)*d*e + (7*B*a^2*b^
2 - 4*A*a*b^3)*e^2)*x^2 + 2*(3*B*a*b^3*d^2 - 2*(5*B*a^2*b^2 - 2*A*a*b^3)*d
*e + (7*B*a^3*b - 4*A*a^2*b^2)*e^2)*x)*sqrt(e/b)*log(8*b^2*e^2*x^2 + b^2*d
^2 + 6*a*b*d*e + a^2*e^2 - 4*(2*b^2*e*x + b^2*d + a*b*e)*sqrt(b*x + a)*sqr
t(e*x + d)*sqrt(e/b) + 8*(b^2*d*e + a*b*e^2)*x) - 4*(6*B*b^3*e^2*x^3 - 8*(
2*B*a*b^2 + A*b^3)*d^2 + 5*(23*B*a^2*b - 8*A*a*b^2)*d*e - 15*(7*B*a^3 - 4*
A*a^2*b)*e^2 + 3*(9*B*b^3*d*e - (7*B*a*b^2 - 4*A*b^3)*e^2)*x^2 - 2*(12*B*b
^3*d^2 - (79*B*a*b^2 - 28*A*b^3)*d*e + 10*(7*B*a^2*b - 4*A*a*b^2)*e^2)*x)*
sqrt(b*x + a)*sqrt(e*x + d))/(b^6*x^2 + 2*a*b^5*x + a^2*b^4), -1/24*(15*(3
*B*a^2*b^2*d^2 - 2*(5*B*a^3*b - 2*A*a^2*b^2)*d*e + (7*B*a^4 - 4*A*a^3*b)*e
^2 + (3*B*b^4*d^2 - 2*(5*B*a*b^3 - 2*A*b^4)*d*e + (7*B*a^2*b^2 - 4*A*a*b^3
)*e^2)*x^2 + 2*(3*B*a*b^3*d^2 - 2*(5*B*a^2*b^2 - 2*A*a*b^3)*d*e + (7*B*a^3
*b - 4*A*a^2*b^2)*e^2)*x)*sqrt(-e/b)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt
(b*x + a)*sqrt(e*x + d)*sqrt(-e/b)/(b*e^2*x^2 + a*d*e + (b*d*e + a*e^2)*x)
) - 2*(6*B*b^3*e^2*x^3 - 8*(2*B*a*b^2 + A*b^3)*d^2 + 5*(23*B*a^2*b - 8*A*a
*b^2)*d*e - 15*(7*B*a^3 - 4*A*a^2*b)*e^2 + 3*(9*B*b^3*d*e - (7*B*a*b^2 - 4
*A*b^3)*e^2)*x^2 - 2*(12*B*b^3*d^2 - (79*B*a*b^2 - 28*A*b^3)*d*e + 10*(7*B
*a^2*b - 4*A*a*b^2)*e^2)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b^6*x^2 + 2*a*b^
5*x + a^2*b^4)]
```

SymPy [F]

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^{5/2}} dx = \int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^{5/2}} dx$$

input

```
integrate((B*x+A)*(e*x+d)**(5/2)/(b*x+a)**(5/2),x)
```

output

```
Integral((A + B*x)*(d + e*x)**(5/2)/(a + b*x)**(5/2), x)
```


Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^(5/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1401 vs. 2(174) = 348.

Time = 0.49 (sec) , antiderivative size = 1401, normalized size of antiderivative = 6.61

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^{5/2}} dx = \text{Too large to display}$$

input `integrate((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^(5/2),x, algorithm="giac")`

output

```

1/4*sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*sqrt(b*x + a)*(2*(b*x + a)*B*e^2*a
bs(b)/b^6 + (9*B*b^12*d*e^3*abs(b) - 13*B*a*b^11*e^4*abs(b) + 4*A*b^12*e^4
*abs(b))/(b^17*e^2)) - 5/8*(3*sqrt(b*e)*B*b^2*d^2*abs(b) - 10*sqrt(b*e)*B*
a*b*d*e*abs(b) + 4*sqrt(b*e)*A*b^2*d*e*abs(b) + 7*sqrt(b*e)*B*a^2*e^2*abs(
b) - 4*sqrt(b*e)*A*a*b*e^2*abs(b))*log((sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2
*d + (b*x + a)*b*e - a*b*e))^2)/b^6 - 4/3*(3*sqrt(b*e)*B*b^7*d^5*abs(b) -
22*sqrt(b*e)*B*a*b^6*d^4*e*abs(b) + 7*sqrt(b*e)*A*b^7*d^4*e*abs(b) + 58*sq
rt(b*e)*B*a^2*b^5*d^3*e^2*abs(b) - 28*sqrt(b*e)*A*a*b^6*d^3*e^2*abs(b) - 7
2*sqrt(b*e)*B*a^3*b^4*d^2*e^3*abs(b) + 42*sqrt(b*e)*A*a^2*b^5*d^2*e^3*abs(
b) + 43*sqrt(b*e)*B*a^4*b^3*d*e^4*abs(b) - 28*sqrt(b*e)*A*a^3*b^4*d*e^4*ab
s(b) - 10*sqrt(b*e)*B*a^5*b^2*e^5*abs(b) + 7*sqrt(b*e)*A*a^4*b^3*e^5*abs(b
) - 6*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*
b*e))^2*B*b^5*d^4*abs(b) + 36*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^
2*d + (b*x + a)*b*e - a*b*e))^2*B*a*b^4*d^3*e*abs(b) - 12*sqrt(b*e)*(sqrt(
b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*A*b^5*d^3*e*ab
s(b) - 72*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e
- a*b*e))^2*B*a^2*b^3*d^2*e^2*abs(b) + 36*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x +
a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*A*a*b^4*d^2*e^2*abs(b) + 60*sq
rt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*
B*a^3*b^2*d*e^3*abs(b) - 36*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b...

```

Mupad [F(-1)]

Timed out.

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^{5/2}} dx = \int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^{5/2}} dx$$

input

```
int(((A + B*x)*(d + e*x)^(5/2))/(a + b*x)^(5/2), x)
```

output

```
int(((A + B*x)*(d + e*x)^(5/2))/(a + b*x)^(5/2), x)
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 304, normalized size of antiderivative = 1.43

$$\int \frac{(A + Bx)(d + ex)^{5/2}}{(a + bx)^{5/2}} dx = \frac{15\sqrt{e}\sqrt{b}\sqrt{bx+a}\log\left(\frac{\sqrt{e}\sqrt{bx+a}+\sqrt{b}\sqrt{ex+d}}{\sqrt{ae-bd}}\right) a^2 e^2 - 30\sqrt{e}\sqrt{b}\sqrt{bx+a}\log\left(\frac{\sqrt{e}\sqrt{bx+a}+\sqrt{b}\sqrt{ex+d}}{\sqrt{ae-bd}}\right)}{(a + bx)^{5/2}}$$

input `int((B*x+A)*(e*x+d)^(5/2)/(b*x+a)^(5/2),x)`

output

```
(15*sqrt(e)*sqrt(b)*sqrt(a + b*x)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a**2*e**2 - 30*sqrt(e)*sqrt(b)*sqrt(a + b*x)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*b*d*e + 15*sqrt(e)*sqrt(b)*sqrt(a + b*x)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b**2*d**2 - 10*sqrt(e)*sqrt(b)*sqrt(a + b*x)*a**2*e**2 + 20*sqrt(e)*sqrt(b)*sqrt(a + b*x)*a*b*d*e - 10*sqrt(e)*sqrt(b)*sqrt(a + b*x)*b**2*d**2 - 15*sqrt(d + e*x)*a**2*b*e**2 + 25*sqrt(d + e*x)*a*b**2*d*e - 5*sqrt(d + e*x)*a*b**2*e**2*x - 8*sqrt(d + e*x)*b**3*d**2 + 9*sqrt(d + e*x)*b**3*d*e*x + 2*sqrt(d + e*x)*b**3*e**2*x**2)/(4*sqrt(a + b*x)*b**4)
```

3.224
$$\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^{5/2}} dx$$

Optimal result	2023
Mathematica [A] (verified)	2024
Rubi [A] (verified)	2024
Maple [B] (verified)	2027
Fricas [B] (verification not implemented)	2028
Sympy [F]	2028
Maxima [F(-2)]	2029
Giac [B] (verification not implemented)	2029
Mupad [F(-1)]	2030
Reduce [B] (verification not implemented)	2031

Optimal result

Integrand size = 24, antiderivative size = 152

$$\int \frac{(A+Bx)(d+ex)^{3/2}}{(a+bx)^{5/2}} dx = -\frac{2(bBd + Abe - 2aBe)\sqrt{d+ex}}{b^3\sqrt{a+bx}} + \frac{Be\sqrt{a+bx}\sqrt{d+ex}}{b^3} - \frac{2(Ab - aB)(d+ex)^{3/2}}{3b^2(a+bx)^{3/2}} + \frac{\sqrt{e}(3bBd + 2Abe - 5aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{b^{7/2}}$$

output

```
-2*(A*b*e-2*B*a*e+B*b*d)*(e*x+d)^(1/2)/b^3/(b*x+a)^(1/2)+B*e*(b*x+a)^(1/2)
*(e*x+d)^(1/2)/b^3-2/3*(A*b-B*a)*(e*x+d)^(3/2)/b^2/(b*x+a)^(3/2)+e^(1/2)*
(2*A*b*e-5*B*a*e+3*B*b*d)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)^(1/
2))/b^(7/2)
```

Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.88

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^{5/2}} dx = \frac{\sqrt{d + ex}(B(-15a^2e + 4ab(d - 5ex) - 3b^2x(-2d + ex)) + 2Ab(3ae + b(d + 4ex)))}{3b^3(a + bx)^{3/2}} + \frac{\sqrt{e}(3bBd + 2Abe - 5aBe)\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{b^{7/2}}$$

input

```
Integrate[((A + B*x)*(d + e*x)^(3/2))/(a + b*x)^(5/2), x]
```

output

```
-1/3*(Sqrt[d + e*x]*(B*(-15*a^2*e + 4*a*b*(d - 5*e*x) - 3*b^2*x*(-2*d + e*x)) + 2*A*b*(3*a*e + b*(d + 4*e*x)))/(b^3*(a + b*x)^(3/2)) + (Sqrt[e]*(3*b*B*d + 2*A*b*e - 5*a*B*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/b^(7/2)
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.18, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$, Rules used = {87, 57, 60, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^{5/2}} dx$$

$$\downarrow 87$$

$$\frac{(-5aBe + 2Abe + 3bBd) \int \frac{(d+ex)^{3/2}}{(a+bx)^{3/2}} dx}{3b(bd - ae)} - \frac{2(d + ex)^{5/2}(Ab - aB)}{3b(a + bx)^{3/2}(bd - ae)}$$

$$\downarrow 57$$

$$\frac{(-5aBe + 2Abe + 3bBd) \left(\frac{3e \int \frac{\sqrt{d+ex}}{\sqrt{a+bx}} dx}{b} - \frac{2(d+ex)^{3/2}}{b\sqrt{a+bx}} \right)}{3b(bd - ae)} - \frac{2(d+ex)^{5/2}(Ab - aB)}{3b(a+bx)^{3/2}(bd - ae)}$$

↓ 60

$$\frac{(-5aBe + 2Abe + 3bBd) \left(\frac{3e \left(\frac{(bd-ae) \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{2b} + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{b} - \frac{2(d+ex)^{3/2}}{b\sqrt{a+bx}} \right)}{3b(bd - ae)} - \frac{2(d+ex)^{5/2}(Ab - aB)}{3b(a+bx)^{3/2}(bd - ae)}$$

↓ 66

$$\frac{(-5aBe + 2Abe + 3bBd) \left(\frac{3e \left(\frac{(bd-ae) \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}} + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{b} - \frac{2(d+ex)^{3/2}}{b\sqrt{a+bx}} \right)}{3b(bd - ae)} - \frac{2(d+ex)^{5/2}(Ab - aB)}{3b(a+bx)^{3/2}(bd - ae)}$$

↓ 221

$$\frac{(-5aBe + 2Abe + 3bBd) \left(\frac{3e \left(\frac{(bd-ae) \operatorname{arctanh} \left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}} \right) + \frac{\sqrt{a+bx}\sqrt{d+ex}}{b} \right)}{b^{3/2}\sqrt{e}} - \frac{2(d+ex)^{3/2}}{b\sqrt{a+bx}} \right)}{3b(bd - ae)} - \frac{2(d+ex)^{5/2}(Ab - aB)}{3b(a+bx)^{3/2}(bd - ae)}$$

input

```
Int[((A + B*x)*(d + e*x)^(3/2))/(a + b*x)^(5/2), x]
```

output

```
(-2*(A*b - a*B)*(d + e*x)^(5/2))/(3*b*(b*d - a*e)*(a + b*x)^(3/2)) + ((3*b*B*d + 2*A*b*e - 5*a*B*e)*((-2*(d + e*x)^(3/2))/(b*Sqrt[a + b*x]) + (3*e*(Sqrt[a + b*x]*Sqrt[d + e*x])/b + ((b*d - a*e)*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])]))/(b^3/2*Sqrt[e]))/(3*b*(b*d - a*e))
```

Definitions of rubi rules used

rule 57 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))]
 Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] &&
 GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] &&
 (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1))]
 Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)]
 Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || (EqQ[e, 0] || (EqQ[c, 0] || LtQ[p, n]))))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 697 vs. $2(126) = 252$.

Time = 0.27 (sec) , antiderivative size = 698, normalized size of antiderivative = 4.59

method	result
default	$\frac{\sqrt{ex+d} \left(6A \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) b^3 e^2 x^2 - 15B \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) a b^2 e^2 x^2 + 9B \ln \left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}} \right) \right)}{\dots}$

input

```
int((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^(5/2),x,method=_RETURNVERBOSE)
```

output

```
1/6*(e*x+d)^(1/2)*(6*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^3*e^2*x^2-15*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^2*e^2*x^2+9*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*b^3*d*e*x^2+12*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^2*e^2*x-30*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*b*e^2*x+18*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a*b^2*d*e*x+6*B*b^2*e*x^2*(b*e)^(1/2)*((e*x+d)*(b*x+a))^(1/2)+6*A*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*b*e^2-16*A*b^2*e*x*(b*e)^(1/2)*((e*x+d)*(b*x+a))^(1/2)-15*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^3*e^2+9*B*ln(1/2*(2*b*e*x+2*((e*x+d)*(b*x+a))^(1/2)*(b*e)^(1/2)+a*e+d*b)/(b*e)^(1/2))*a^2*b*d*e+40*B*a*b*e*x*(b*e)^(1/2)*((e*x+d)*(b*x+a))^(1/2)-12*B*b^2*d*x*(b*e)^(1/2)*((e*x+d)*(b*x+a))^(1/2)-12*A*a*b*e*(b*e)^(1/2)*((e*x+d)*(b*x+a))^(1/2)-4*A*b^2*d*(b*e)^(1/2)*((e*x+d)*(b*x+a))^(1/2)+30*B*a^2*e*(b*e)^(1/2)*((e*x+d)*(b*x+a))^(1/2)-8*B*a*b*d*(b*e)^(1/2)*((e*x+d)*(b*x+a))^(1/2)/(b*e)^(1/2)/((e*x+d)*(b*x+a))^(1/2)/b^3/(b*x+a)^(3/2)
```


Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 268 vs. $2(126) = 252$.

Time = 0.98 (sec) , antiderivative size = 561, normalized size of antiderivative = 3.69

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^{5/2}} dx = \left[\frac{3(3Ba^2bd + (3Bb^3d - (5Bab^2 - 2Ab^3)e)x^2 - (5Ba^3 - 2Aa^2b)e + 2(3Bb^2d - 2Aab^2)e)x\sqrt{e/b} \log(8b^2e^2x^2 + b^2d^2 + 6abde + a^2e^2 + 4(2b^2ex + b^2d + abe)\sqrt{bx+a})\sqrt{ex+d}\sqrt{e/b} + 8(b^2de + abe^2)x) + 4(3Bb^2ex^2 - 2(2Bab + Ab^2)d + 3(5Ba^2 - 2Aab)e - 2(3Bb^2d - 2(5Bab - 2Ab^2)e)x)\sqrt{bx+a}\sqrt{ex+d}}{(b^5x^2 + 2ab^4x + a^2b^3)}, -\frac{1}{6}(3(3Ba^2bd + (3Bb^3d - (5Bab^2 - 2Ab^3)e)x^2 - (5Ba^3 - 2Aa^2b)e + 2(3Bb^2d - 2(5Bab - 2Ab^2)e)x)\sqrt{-e/b})\arctan(1/2(2bex + bd + ae)\sqrt{bx+a})\sqrt{ex+d}\sqrt{-e/b}}{(b^5x^2 + 2ab^4x + a^2b^3)} \right]$$

input `integrate((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^(5/2),x, algorithm="fricas")`

output `[1/12*(3*(3*B*a^2*b*d + (3*B*b^3*d - (5*B*a*b^2 - 2*A*b^3)*e)*x^2 - (5*B*a^3 - 2*A*a^2*b)*e + 2*(3*B*a*b^2*d - (5*B*a^2*b - 2*A*a*b^2)*e)*x)*sqrt(e/b)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*b^2*e*x + b^2*d + a*b*e)*sqrt(b*x + a)*sqrt(e*x + d)*sqrt(e/b) + 8*(b^2*d*e + a*b*e^2)*x) + 4*(3*B*b^2*e*x^2 - 2*(2*B*a*b + A*b^2)*d + 3*(5*B*a^2 - 2*A*a*b)*e - 2*(3*B*b^2*d - 2*(5*B*a*b - 2*A*b^2)*e)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b^5*x^2 + 2*a*b^4*x + a^2*b^3), -1/6*(3*(3*B*a^2*b*d + (3*B*b^3*d - (5*B*a*b^2 - 2*A*b^3)*e)*x^2 - (5*B*a^3 - 2*A*a^2*b)*e + 2*(3*B*a*b^2*d - (5*B*a^2*b - 2*A*a*b^2)*e)*x)*sqrt(-e/b)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(b*x + a)*sqrt(e*x + d)*sqrt(-e/b)/(b*e^2*x^2 + a*d*e + (b*d*e + a*e^2)*x)) - 2*(3*B*b^2*e*x^2 - 2*(2*B*a*b + A*b^2)*d + 3*(5*B*a^2 - 2*A*a*b)*e - 2*(3*B*b^2*d - 2*(5*B*a*b - 2*A*b^2)*e)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(b^5*x^2 + 2*a*b^4*x + a^2*b^3)]`

Sympy [F]

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^{5/2}} dx = \int \frac{(A + Bx)(d + ex)^{\frac{3}{2}}}{(a + bx)^{\frac{5}{2}}} dx$$

input `integrate((B*x+A)*(e*x+d)**(3/2)/(b*x+a)**(5/2),x)`

output `Integral((A + B*x)*(d + e*x)**(3/2)/(a + b*x)**(5/2), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^{5/2}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^(5/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for m
ore detail
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1019 vs. 2(126) = 252.

Time = 0.34 (sec) , antiderivative size = 1019, normalized size of antiderivative = 6.70

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^{5/2}} dx = \text{Too large to display}$$

input

```
integrate((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^(5/2),x, algorithm="giac")
```

output

```

sqrt(b^2*d + (b*x + a)*b*e - a*b*e)*sqrt(b*x + a)*B*e*abs(b)/b^5 - 1/2*(3*
sqrt(b*e)*B*b*d*abs(b) - 5*sqrt(b*e)*B*a*e*abs(b) + 2*sqrt(b*e)*A*b*e*abs(
b))*log((sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2)
/b^5 - 4/3*(3*sqrt(b*e)*B*b^6*d^4*abs(b) - 16*sqrt(b*e)*B*a*b^5*d^3*e*abs(
b) + 4*sqrt(b*e)*A*b^6*d^3*e*abs(b) + 30*sqrt(b*e)*B*a^2*b^4*d^2*e^2*abs(b
) - 12*sqrt(b*e)*A*a*b^5*d^2*e^2*abs(b) - 24*sqrt(b*e)*B*a^3*b^3*d*e^3*abs
(b) + 12*sqrt(b*e)*A*a^2*b^4*d*e^3*abs(b) + 7*sqrt(b*e)*B*a^4*b^2*e^4*abs(
b) - 4*sqrt(b*e)*A*a^3*b^3*e^4*abs(b) - 6*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x +
a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*B*b^4*d^3*abs(b) + 24*sqrt(b*e
)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*B*a*b^
3*d^2*e*abs(b) - 6*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x
+ a)*b*e - a*b*e))^2*A*b^4*d^2*e*abs(b) - 30*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x
+ a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*B*a^2*b^2*d*e^2*abs(b) + 12
*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))
^2*A*a*b^3*d*e^2*abs(b) + 12*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2
*d + (b*x + a)*b*e - a*b*e))^2*B*a^3*b*e^3*abs(b) - 6*sqrt(b*e)*(sqrt(b*e)
*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*A*a^2*b^2*e^3*abs(
b) + 3*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e -
a*b*e))^4*B*b^2*d^2*abs(b) - 12*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(
b^2*d + (b*x + a)*b*e - a*b*e))^4*B*a*b*d*e*abs(b) + 6*sqrt(b*e)*(sqrt(b...

```

Mupad [F(-1)]

Timed out.

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^{5/2}} dx = \int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^{5/2}} dx$$

input

```
int(((A + B*x)*(d + e*x)^(3/2))/(a + b*x)^(5/2), x)
```

output

```
int(((A + B*x)*(d + e*x)^(3/2))/(a + b*x)^(5/2), x)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.12

$$\int \frac{(A + Bx)(d + ex)^{3/2}}{(a + bx)^{5/2}} dx = \frac{-12\sqrt{e}\sqrt{b}\sqrt{bx+a}\log\left(\frac{\sqrt{e}\sqrt{bx+a}+\sqrt{b}\sqrt{ex+d}}{\sqrt{ae-bd}}\right)ae + 12\sqrt{e}\sqrt{b}\sqrt{bx+a}\log\left(\frac{\sqrt{e}\sqrt{bx+a}+\sqrt{b}\sqrt{ex+d}}{\sqrt{ae-bd}}\right)}{(a + bx)^{5/2}}$$

input `int((B*x+A)*(e*x+d)^(3/2)/(b*x+a)^(5/2),x)`output `(- 12*sqrt(e)*sqrt(b)*sqrt(a + b*x)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*a*e + 12*sqrt(e)*sqrt(b)*sqrt(a + b*x)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d))*b*d + 9*sqrt(e)*sqrt(b)*sqrt(a + b*x)*a*e - 9*sqrt(e)*sqrt(b)*sqrt(a + b*x)*b*d + 12*sqrt(d + e*x)*a*b*e - 8*sqrt(d + e*x)*b**2*d + 4*sqrt(d + e*x)*b**2*e*x)/(4*sqrt(a + b*x)*b**3)`

3.225 $\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^{5/2}} dx$

Optimal result	2032
Mathematica [A] (verified)	2032
Rubi [A] (verified)	2033
Maple [B] (verified)	2035
Fricas [B] (verification not implemented)	2035
Sympy [F]	2036
Maxima [F(-2)]	2037
Giac [B] (verification not implemented)	2037
Mupad [F(-1)]	2038
Reduce [B] (verification not implemented)	2038

Optimal result

Integrand size = 24, antiderivative size = 111

$$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^{5/2}} dx = -\frac{2B\sqrt{d+ex}}{b^2\sqrt{a+bx}} - \frac{2(Ab-aB)(d+ex)^{3/2}}{3b(bd-ae)(a+bx)^{3/2}} + \frac{2B\sqrt{e}\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}}\right)}{b^{5/2}}$$

output

```
-2*B*(e*x+d)^(1/2)/b^2/(b*x+a)^(1/2)-2/3*(A*b-B*a)*(e*x+d)^(3/2)/b/(-a*e+b*d)/(b*x+a)^(3/2)+2*B*e^(1/2)*arctanh(e^(1/2)*(b*x+a)^(1/2)/b^(1/2)/(e*x+d)^(1/2))/b^(5/2)
```

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.05

$$\int \frac{(A+Bx)\sqrt{d+ex}}{(a+bx)^{5/2}} dx = -\frac{2\sqrt{d+ex}(Ab^2(d+ex)+B(-3a^2e+3b^2dx+2ab(d-2ex)))}{3b^2(bd-ae)(a+bx)^{3/2}} + \frac{2B\sqrt{e}\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{d+ex}}{\sqrt{e}\sqrt{a+bx}}\right)}{b^{5/2}}$$

input `Integrate[((A + B*x)*Sqrt[d + e*x])/(a + b*x)^(5/2),x]`

output `(-2*Sqrt[d + e*x]*(A*b^2*(d + e*x) + B*(-3*a^2*e + 3*b^2*d*x + 2*a*b*(d - 2*e*x)))/(3*b^2*(b*d - a*e)*(a + b*x)^(3/2)) + (2*B*Sqrt[e]*ArcTanh[(Sqrt[b]*Sqrt[d + e*x])/(Sqrt[e]*Sqrt[a + b*x])])/b^(5/2)`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.04, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {87, 57, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^{5/2}} dx \\
 & \quad \downarrow 87 \\
 & \frac{B \int \frac{\sqrt{d+ex}}{(a+bx)^{3/2}} dx}{b} - \frac{2(d+ex)^{3/2}(Ab - aB)}{3b(a+bx)^{3/2}(bd - ae)} \\
 & \quad \downarrow 57 \\
 & \frac{B \left(\frac{e \int \frac{1}{\sqrt{a+bx}\sqrt{d+ex}} dx}{b} - \frac{2\sqrt{d+ex}}{b\sqrt{a+bx}} \right)}{b} - \frac{2(d+ex)^{3/2}(Ab - aB)}{3b(a+bx)^{3/2}(bd - ae)} \\
 & \quad \downarrow 66 \\
 & \frac{B \left(\frac{2e \int \frac{1}{b - \frac{e(a+bx)}{d+ex}} d \frac{\sqrt{a+bx}}{\sqrt{d+ex}}}{b} - \frac{2\sqrt{d+ex}}{b\sqrt{a+bx}} \right)}{b} - \frac{2(d+ex)^{3/2}(Ab - aB)}{3b(a+bx)^{3/2}(bd - ae)} \\
 & \quad \downarrow 221 \\
 & \frac{B \left(\frac{2\sqrt{e} \operatorname{arctanh} \left(\frac{\sqrt{e}\sqrt{a+bx}}{\sqrt{b}\sqrt{d+ex}} \right)}{b^{3/2}} - \frac{2\sqrt{d+ex}}{b\sqrt{a+bx}} \right)}{b} - \frac{2(d+ex)^{3/2}(Ab - aB)}{3b(a+bx)^{3/2}(bd - ae)}
 \end{aligned}$$

input `Int[((A + B*x)*Sqrt[d + e*x])/(a + b*x)^(5/2),x]`

output `(-2*(A*b - a*B)*(d + e*x)^(3/2))/(3*b*(b*d - a*e)*(a + b*x)^(3/2)) + (B*((-2*Sqrt[d + e*x])/(b*Sqrt[a + b*x]) + (2*Sqrt[e]*ArcTanh[(Sqrt[e]*Sqrt[a + b*x])/(Sqrt[b]*Sqrt[d + e*x])])/b^(3/2)))/b`

Defintions of rubi rules used

rule 57 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))]
Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 502 vs. $2(89) = 178$.

Time = 0.27 (sec) , antiderivative size = 503, normalized size of antiderivative = 4.53

method	result
default	$\left(3B \ln\left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}}\right) a b^2 e^2 x^2 - 3B \ln\left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}}\right) b^3 d e x^2 + 6B \ln\left(\frac{2bex+2\sqrt{(ex+d)(bx+a)}\sqrt{be+ae+db}}{2\sqrt{be}}\right) \right)$

input `int((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^(5/2),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & \frac{1}{3} * (3 * B * \ln(1/2 * (2 * b * e * x + 2 * ((e * x + d) * (b * x + a))^{1/2} * (b * e)^{1/2} + a * e + d * b) / (b * e)^{1/2}) * a * b^2 * e^{-2} * x^2 - 3 * B * \ln(1/2 * (2 * b * e * x + 2 * ((e * x + d) * (b * x + a))^{1/2} * (b * e)^{1/2} + a * e + d * b) / (b * e)^{1/2}) * b^3 * d * e * x^2 + 6 * B * \ln(1/2 * (2 * b * e * x + 2 * ((e * x + d) * (b * x + a))^{1/2} * (b * e)^{1/2} + a * e + d * b) / (b * e)^{1/2}) * a^2 * b * e^{-2} * x - 6 * B * \ln(1/2 * (2 * b * e * x + 2 * ((e * x + d) * (b * x + a))^{1/2} * (b * e)^{1/2} + a * e + d * b) / (b * e)^{1/2}) * a * b^2 * d * e * x + 2 * A * b^2 * e * x * (b * e)^{1/2} * ((e * x + d) * (b * x + a))^{1/2} + 3 * B * \ln(1/2 * (2 * b * e * x + 2 * ((e * x + d) * (b * x + a))^{1/2} * (b * e)^{1/2} + a * e + d * b) / (b * e)^{1/2}) * a^3 * e^{-2} - 3 * B * \ln(1/2 * (2 * b * e * x + 2 * ((e * x + d) * (b * x + a))^{1/2} * (b * e)^{1/2} + a * e + d * b) / (b * e)^{1/2}) * a^2 * b * d * e - 8 * B * a * b * e * x * (b * e)^{1/2} * ((e * x + d) * (b * x + a))^{1/2} + 6 * B * b^2 * d * x * (b * e)^{1/2} * ((e * x + d) * (b * x + a))^{1/2} + 2 * A * b^2 * d * (b * e)^{1/2} * ((e * x + d) * (b * x + a))^{1/2} - 6 * B * a^2 * e * (b * e)^{1/2} * ((e * x + d) * (b * x + a))^{1/2} + 4 * B * a * b * d * (b * e)^{1/2} * ((e * x + d) * (b * x + a))^{1/2} * (e * x + d)^{1/2} / (b * e)^{1/2} / (a * e - b * d) / ((e * x + d) * (b * x + a))^{1/2} / b^2 / (b * x + a)^{3/2} \end{aligned}$$
Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 250 vs. $2(89) = 178$.

Time = 0.67 (sec) , antiderivative size = 525, normalized size of antiderivative = 4.73

$$\int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^{5/2}} dx = \frac{3(Ba^2bd - Ba^3e + (Bb^3d - Bab^2e)x^2 + 2(Bab^2d - Ba^2be)x)\sqrt{\frac{e}{b}} \log(8b^2e^2) + 3(Ba^2bd - Ba^3e + (Bb^3d - Bab^2e)x^2 + 2(Bab^2d - Ba^2be)x)\sqrt{-\frac{e}{b}} \arctan\left(\frac{(2beax + bd + ae)\sqrt{bx+a}\sqrt{ex+d}\sqrt{-\frac{e}{b}}}{2(be^2x^2 + ade + (bde + ae^2)x)}\right)}{3(a^2b^3d - a^3b^2e + (b^5d - ab^4e)x^2 + 2(b^5d - ab^4e)x)}$$

input `integrate((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^(5/2),x, algorithm="fricas")`

output `[1/6*(3*(B*a^2*b*d - B*a^3*e + (B*b^3*d - B*a*b^2*e)*x^2 + 2*(B*a*b^2*d - B*a^2*b*e)*x)*sqrt(e/b)*log(8*b^2*e^2*x^2 + b^2*d^2 + 6*a*b*d*e + a^2*e^2 + 4*(2*b^2*e*x + b^2*d + a*b*e)*sqrt(b*x + a)*sqrt(e*x + d)*sqrt(e/b) + 8*(b^2*d*e + a*b*e^2)*x) + 4*(3*B*a^2*e - (2*B*a*b + A*b^2)*d - (3*B*b^2*d - (4*B*a*b - A*b^2)*e)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(a^2*b^3*d - a^3*b^2*e + (b^5*d - a*b^4*e)*x^2 + 2*(a*b^4*d - a^2*b^3*e)*x), -1/3*(3*(B*a^2*b*d - B*a^3*e + (B*b^3*d - B*a*b^2*e)*x^2 + 2*(B*a*b^2*d - B*a^2*b*e)*x)*sqrt(-e/b)*arctan(1/2*(2*b*e*x + b*d + a*e)*sqrt(b*x + a)*sqrt(e*x + d)*sqrt(-e/b)/(b*e^2*x^2 + a*d*e + (b*d*e + a*e^2)*x)) - 2*(3*B*a^2*e - (2*B*a*b + A*b^2)*d - (3*B*b^2*d - (4*B*a*b - A*b^2)*e)*x)*sqrt(b*x + a)*sqrt(e*x + d))/(a^2*b^3*d - a^3*b^2*e + (b^5*d - a*b^4*e)*x^2 + 2*(a*b^4*d - a^2*b^3*e)*x)]`

Sympy [F]

$$\int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^{5/2}} dx = \int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^{5/2}} dx$$

input `integrate((B*x+A)*(e*x+d)**(1/2)/(b*x+a)**(5/2),x)`

output `Integral((A + B*x)*sqrt(d + e*x)/(a + b*x)**(5/2), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^(5/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 558 vs. 2(89) = 178.

Time = 0.23 (sec) , antiderivative size = 558, normalized size of antiderivative = 5.03

$$\int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^{5/2}} dx =$$

$$\frac{\sqrt{be}B|b| \log \left(\left(\sqrt{be}\sqrt{bx + a} - \sqrt{b^2d + (bx + a)be - abe} \right)^2 \right)}{b^4}$$

$$- \frac{4 \left(3\sqrt{be}Bb^5d^3|b| - 10\sqrt{be}Bab^4d^2e|b| + \sqrt{be}Ab^5d^2e|b| + 11\sqrt{be}Ba^2b^3de^2|b| - 2\sqrt{be}Aab^4de^2|b| - 4\sqrt{be} \right)}{b^4}$$

input `integrate((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^(5/2),x, algorithm="giac")`

output

```
-sqrt(b*e)*B*abs(b)*log((sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*
b*e - a*b*e))^2)/b^4 - 4/3*(3*sqrt(b*e)*B*b^5*d^3*abs(b) - 10*sqrt(b*e)*B*
a*b^4*d^2*e*abs(b) + sqrt(b*e)*A*b^5*d^2*e*abs(b) + 11*sqrt(b*e)*B*a^2*b^3
*d*e^2*abs(b) - 2*sqrt(b*e)*A*a*b^4*d*e^2*abs(b) - 4*sqrt(b*e)*B*a^3*b^2*e
^3*abs(b) + sqrt(b*e)*A*a^2*b^3*e^3*abs(b) - 6*sqrt(b*e)*(sqrt(b*e)*sqrt(b
*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*B*b^3*d^2*abs(b) + 12*sq
rt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*B
*a*b^2*d*e*abs(b) - 6*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b
*x + a)*b*e - a*b*e))^2*B*a^2*b*e^2*abs(b) + 3*sqrt(b*e)*(sqrt(b*e)*sqrt(b
*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^4*B*b*d*abs(b) - 6*sqrt(b*e
)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^4*B*a*e*
abs(b) + 3*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e
- a*b*e))^4*A*b*e*abs(b))/((b^2*d - a*b*e - (sqrt(b*e)*sqrt(b*x + a) - sq
rt(b^2*d + (b*x + a)*b*e - a*b*e))^2)^3*b^3)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^{5/2}} dx = \int \frac{(A + Bx) \sqrt{d + ex}}{(a + bx)^{5/2}} dx$$

input

```
int(((A + B*x)*(d + e*x)^(1/2))/(a + b*x)^(5/2), x)
```

output

```
int(((A + B*x)*(d + e*x)^(1/2))/(a + b*x)^(5/2), x)
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.70

$$\int \frac{(A + Bx)\sqrt{d + ex}}{(a + bx)^{5/2}} dx = \frac{2\sqrt{e} \sqrt{b} \sqrt{bx + a} \log\left(\frac{\sqrt{e} \sqrt{bx+a} + \sqrt{b} \sqrt{ex+d}}{\sqrt{ae-bd}}\right) - 2\sqrt{e} \sqrt{b} \sqrt{bx + a} - 2\sqrt{ex + d} b}{\sqrt{bx + a} b^2}$$

input

```
int((B*x+A)*(e*x+d)^(1/2)/(b*x+a)^(5/2), x)
```

output

```
(2*(sqrt(e)*sqrt(b)*sqrt(a + b*x)*log((sqrt(e)*sqrt(a + b*x) + sqrt(b)*sqrt(d + e*x))/sqrt(a*e - b*d)) - sqrt(e)*sqrt(b)*sqrt(a + b*x) - sqrt(d + e*x)*b))/(sqrt(a + b*x)*b**2)
```

3.226 $\int \frac{A+Bx}{(a+bx)^{5/2}\sqrt{d+ex}} dx$

Optimal result	2040
Mathematica [A] (verified)	2040
Rubi [A] (verified)	2041
Maple [A] (verified)	2042
Fricas [A] (verification not implemented)	2043
Sympy [F]	2043
Maxima [F(-2)]	2043
Giac [B] (verification not implemented)	2044
Mupad [B] (verification not implemented)	2044
Reduce [B] (verification not implemented)	2045

Optimal result

Integrand size = 24, antiderivative size = 95

$$\int \frac{A+Bx}{(a+bx)^{5/2}\sqrt{d+ex}} dx = -\frac{2(Ab-aB)\sqrt{d+ex}}{3b(bd-ae)(a+bx)^{3/2}} - \frac{2(3bBd-2Abe-aBe)\sqrt{d+ex}}{3b(bd-ae)^2\sqrt{a+bx}}$$

output

```
-2/3*(A*b-B*a)*(e*x+d)^(1/2)/b/(-a*e+b*d)/(b*x+a)^(3/2)-2/3*(-2*A*b*e-B*a*
e+3*B*b*d)*(e*x+d)^(1/2)/b/(-a*e+b*d)^2/(b*x+a)^(1/2)
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.67

$$\int \frac{A+Bx}{(a+bx)^{5/2}\sqrt{d+ex}} dx = -\frac{2\sqrt{d+ex}(-3aAe+Ab(d-2ex)+B(2ad+3bdx-ae))}{3(bd-ae)^2(a+bx)^{3/2}}$$

input

```
Integrate[(A + B*x)/((a + b*x)^(5/2)*Sqrt[d + e*x]),x]
```

output

```
(-2*Sqrt[d + e*x]*(-3*a*A*e + A*b*(d - 2*e*x) + B*(2*a*d + 3*b*d*x - a*e*x
)))/(3*(b*d - a*e)^2*(a + b*x)^(3/2))
```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {87, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^{5/2} \sqrt{d + ex}} dx$$

$$\downarrow 87$$

$$\frac{(-aBe - 2Abe + 3bBd) \int \frac{1}{(a+bx)^{3/2} \sqrt{d+ex}} dx}{3b(bd - ae)} - \frac{2\sqrt{d + ex}(Ab - aB)}{3b(a + bx)^{3/2}(bd - ae)}$$

$$\downarrow 48$$

$$-\frac{2\sqrt{d + ex}(Ab - aB)}{3b(a + bx)^{3/2}(bd - ae)} - \frac{2\sqrt{d + ex}(-aBe - 2Abe + 3bBd)}{3b\sqrt{a + bx}(bd - ae)^2}$$

input

```
Int[(A + B*x)/((a + b*x)^(5/2)*Sqrt[d + e*x]),x]
```

output

```
(-2*(A*b - a*B)*Sqrt[d + e*x])/(3*b*(b*d - a*e)*(a + b*x)^(3/2)) - (2*(3*b*B*d - 2*A*b*e - a*B*e)*Sqrt[d + e*x])/(3*b*(b*d - a*e)^2*Sqrt[a + b*x])
```

Definitions of rubi rules used

rule 48

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

rule 87

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(- (b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.63

method	result	size
default	$\frac{2\sqrt{ex+d}(2Abe x+Baex-3Bbdx+3Aae-Abd-2Bad)}{3(bx+a)^{\frac{3}{2}}(ae-db)^2}$	60
gospers	$\frac{2\sqrt{ex+d}(2Abe x+Baex-3Bbdx+3Aae-Abd-2Bad)}{3(bx+a)^{\frac{3}{2}}(a^2e^2-2abde+b^2d^2)}$	73
orering	$\frac{2\sqrt{ex+d}(2Abe x+Baex-3Bbdx+3Aae-Abd-2Bad)}{3(bx+a)^{\frac{3}{2}}(a^2e^2-2abde+b^2d^2)}$	73

input

```
int((B*x+A)/(b*x+a)^(5/2)/(e*x+d)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
2/3*(e*x+d)^(1/2)*(2*A*b*e*x+B*a*e*x-3*B*b*d*x+3*A*a*e-A*b*d-2*B*a*d)/(b*x
+a)^(3/2)/(a*e-b*d)^2
```

Fricas [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.48

$$\int \frac{A + Bx}{(a + bx)^{5/2} \sqrt{d + ex}} dx = \frac{2(3Aae - (2Ba + Ab)d - (3Bbd - (Ba + 2Ab)e)x) \sqrt{bx + a} \sqrt{ex} - 3(a^2b^2d^2 - 2a^3bde + a^4e^2 + (b^4d^2 - 2ab^3de + a^2b^2e^2)x^2 + 2(ab^3d^2 - 2a^2b^2de + a^3b^2e^2)x)}{3(a^2b^2d^2 - 2a^3bde + a^4e^2 + (b^4d^2 - 2ab^3de + a^2b^2e^2)x^2 + 2(ab^3d^2 - 2a^2b^2de + a^3b^2e^2)x)}$$

input `integrate((B*x+A)/(b*x+a)^(5/2)/(e*x+d)^(1/2),x, algorithm="fricas")`

output `2/3*(3*A*a*e - (2*B*a + A*b)*d - (3*B*b*d - (B*a + 2*A*b)*e)*x)*sqrt(b*x + a)*sqrt(e*x + d)/(a^2*b^2*d^2 - 2*a^3*b*d*e + a^4*e^2 + (b^4*d^2 - 2*a*b^3*d*e + a^2*b^2*e^2)*x^2 + 2*(a*b^3*d^2 - 2*a^2*b^2*d*e + a^3*b^2*e^2)*x)`

Sympy [F]

$$\int \frac{A + Bx}{(a + bx)^{5/2} \sqrt{d + ex}} dx = \int \frac{A + Bx}{(a + bx)^{\frac{5}{2}} \sqrt{d + ex}} dx$$

input `integrate((B*x+A)/(b*x+a)**(5/2)/(e*x+d)**(1/2),x)`

output `Integral((A + B*x)/((a + b*x)**(5/2)*sqrt(d + e*x)), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{(a + bx)^{5/2} \sqrt{d + ex}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^(5/2)/(e*x+d)^(1/2),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for m
ore detail
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 282 vs. $2(83) = 166$.

Time = 0.17 (sec) , antiderivative size = 282, normalized size of antiderivative = 2.97

$$\int \frac{A + Bx}{(a + bx)^{5/2} \sqrt{d + ex}} dx =$$

$$\frac{4 \left(3 \sqrt{be} B b^4 d^2 - 4 \sqrt{be} B a b^3 d e - 2 \sqrt{be} A b^4 d e + \sqrt{be} B a^2 b^2 e^2 + 2 \sqrt{be} A a b^3 e^2 - 6 \sqrt{be} \left(\sqrt{be} \sqrt{bx + a} - \sqrt{be} \sqrt{bx + a} \right) \right)}{3 \left(b^2 d - a b \right)}$$

input

```
integrate((B*x+A)/(b*x+a)^(5/2)/(e*x+d)^(1/2),x, algorithm="giac")
```

output

```
-4/3*(3*sqrt(b*e)*B*b^4*d^2 - 4*sqrt(b*e)*B*a*b^3*d*e - 2*sqrt(b*e)*A*b^4*
d*e + sqrt(b*e)*B*a^2*b^2*e^2 + 2*sqrt(b*e)*A*a*b^3*e^2 - 6*sqrt(b*e)*(sqr
t(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*B*b^2*d + 6*
sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^
2*A*b^2*e + 3*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*
b*e - a*b*e))^4*B)/((b^2*d - a*b*e - (sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d
+ (b*x + a)*b*e - a*b*e))^2)^3*abs(b))
```

Mupad [B] (verification not implemented)

Time = 1.56 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.02

$$\int \frac{A + Bx}{(a + bx)^{5/2} \sqrt{d + ex}} dx = - \frac{\left(\frac{2Abd - 6Aae + 4Bad}{3b(ae - bd)^2} - \frac{x(4Abe + 2Bae - 6Bbd)}{3b(ae - bd)^2} \right) \sqrt{d + ex}}{x \sqrt{a + bx} + \frac{a \sqrt{a + bx}}{b}}$$

input

```
int((A + B*x)/((a + b*x)^(5/2)*(d + e*x)^(1/2)),x)
```

output

```
-(((2*A*b*d - 6*A*a*e + 4*B*a*d)/(3*b*(a*e - b*d)^2) - (x*(4*A*b*e + 2*B*a
*e - 6*B*b*d))/(3*b*(a*e - b*d)^2))*(d + e*x)^(1/2))/(x*(a + b*x)^(1/2) +
(a*(a + b*x)^(1/2))/b)
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.45

$$\int \frac{A + Bx}{(a + bx)^{5/2} \sqrt{d + ex}} dx = \frac{2\sqrt{e} \sqrt{b} \sqrt{bx + a} + 2\sqrt{ex + d} b}{\sqrt{bx + a} b (ae - bd)}$$

input

```
int((B*x+A)/(b*x+a)^(5/2)/(e*x+d)^(1/2),x)
```

output

```
(2*(sqrt(e)*sqrt(b)*sqrt(a + b*x) + sqrt(d + e*x)*b))/(sqrt(a + b*x)*b*(a*
e - b*d))
```

3.227 $\int \frac{A+Bx}{(a+bx)^{5/2}(d+ex)^{3/2}} dx$

Optimal result	2046
Mathematica [A] (verified)	2046
Rubi [A] (verified)	2047
Maple [A] (verified)	2048
Fricas [B] (verification not implemented)	2049
Sympy [F]	2049
Maxima [F(-2)]	2050
Giac [B] (verification not implemented)	2050
Mupad [B] (verification not implemented)	2051
Reduce [B] (verification not implemented)	2052

Optimal result

Integrand size = 24, antiderivative size = 145

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{3/2}} dx = -\frac{2(Ab - aB)}{3b(bd - ae)(a + bx)^{3/2}\sqrt{d + ex}} - \frac{2(3bBd - 4Abe + aBe)}{3b(bd - ae)^2\sqrt{a + bx}\sqrt{d + ex}} - \frac{4e(3bBd - 4Abe + aBe)\sqrt{a + bx}}{3b(bd - ae)^3\sqrt{d + ex}}$$

output

```
1/3*(-2*A*b+2*B*a)/b/(-a*e+b*d)/(b*x+a)^(3/2)/(e*x+d)^(1/2)-2/3*(-4*A*b*e+
B*a*e+3*B*b*d)/b/(-a*e+b*d)^2/(b*x+a)^(1/2)/(e*x+d)^(1/2)-4/3*e*(-4*A*b*e+
B*a*e+3*B*b*d)*(b*x+a)^(1/2)/b/(-a*e+b*d)^3/(e*x+d)^(1/2)
```

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.86

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{3/2}} dx = \frac{2(A(-3a^2e^2 - 6abe(d + 2ex) + b^2(d^2 - 4dex - 8e^2x^2)) + B(3a^2e(2d + ex) + 3b^2dx(d + 2ex) + 2ab(d^2 - 3(bd - ae)^3(a + bx)^{3/2}\sqrt{d + ex}))}{3(bd - ae)^3(a + bx)^{3/2}\sqrt{d + ex}}$$

input

```
Integrate[(A + B*x)/((a + b*x)^(5/2)*(d + e*x)^(3/2)),x]
```

output

$$\frac{(-2*(A*(-3*a^2*e^2 - 6*a*b*e*(d + 2*e*x) + b^2*(d^2 - 4*d*e*x - 8*e^2*x^2)) + B*(3*a^2*e*(2*d + e*x) + 3*b^2*d*x*(d + 2*e*x) + 2*a*b*(d^2 + 5*d*e*x + e^2*x^2)))/(3*(b*d - a*e)^3*(a + b*x)^(3/2)*\text{Sqrt}[d + e*x])$$

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.95, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {87, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{3/2}} dx$$

↓ 87

$$\frac{(aBe - 4Abe + 3bBd) \int \frac{1}{(a+bx)^{5/2}\sqrt{d+ex}} dx}{e(bd - ae)} - \frac{2(Bd - Ae)}{e(a + bx)^{3/2}\sqrt{d + ex}(bd - ae)}$$

↓ 55

$$\frac{(aBe - 4Abe + 3bBd) \left(-\frac{2e \int \frac{1}{(a+bx)^{3/2}\sqrt{d+ex}} dx}{3(bd-ae)} - \frac{2\sqrt{d+ex}}{3(a+bx)^{3/2}(bd-ae)} \right)}{e(bd - ae) \frac{2(Bd - Ae)}{e(a + bx)^{3/2}\sqrt{d + ex}(bd - ae)}}$$

↓ 48

$$\frac{2(Bd - Ae)}{e(a + bx)^{3/2}\sqrt{d + ex}(bd - ae)} - \frac{\left(\frac{4e\sqrt{d+ex}}{3\sqrt{a+bx}(bd-ae)^2} - \frac{2\sqrt{d+ex}}{3(a+bx)^{3/2}(bd-ae)} \right) (aBe - 4Abe + 3bBd)}{e(bd - ae)}$$

input

$$\text{Int}[(A + B*x)/((a + b*x)^(5/2)*(d + e*x)^(3/2)), x]$$

output

$$\frac{(-2*(B*d - A*e))/(e*(b*d - a*e)*(a + b*x)^(3/2)*\text{Sqrt}[d + e*x]) - ((3*b*B*d - 4*A*b*e + a*B*e)*((-2*\text{Sqrt}[d + e*x])/(3*(b*d - a*e)*(a + b*x)^(3/2)) + (4*e*\text{Sqrt}[d + e*x])/(3*(b*d - a*e)^2*\text{Sqrt}[a + b*x]))/(e*(b*d - a*e))$$

Defintions of rubi rules used

```
rule 48 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 55 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
lerQ[n, 1])
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && ( !LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.03

method	result
default	$-\frac{2(8Ab^2e^2x^2-2Babe^2x^2-6Bb^2dex^2+12Aabe^2x+4Ab^2dex-3Ba^2e^2x-10Babdex-3b^2Bd^2x+3a^2Ae^2+6Aabde-Ab^2d^2-6B}{3\sqrt{ex+d}(bx+a)^{\frac{3}{2}}(ae-db)^3}$
gosper	$-\frac{2(8Ab^2e^2x^2-2Babe^2x^2-6Bb^2dex^2+12Aabe^2x+4Ab^2dex-3Ba^2e^2x-10Babdex-3b^2Bd^2x+3a^2Ae^2+6Aabde-Ab^2d^2-6B}{3\sqrt{ex+d}(bx+a)^{\frac{3}{2}}(a^3e^3-3a^2be^2d+3ab^2d^2e-b^3d^3)}$
orering	$-\frac{2(8Ab^2e^2x^2-2Babe^2x^2-6Bb^2dex^2+12Aabe^2x+4Ab^2dex-3Ba^2e^2x-10Babdex-3b^2Bd^2x+3a^2Ae^2+6Aabde-Ab^2d^2-6B}{3\sqrt{ex+d}(bx+a)^{\frac{3}{2}}(a^3e^3-3a^2be^2d+3ab^2d^2e-b^3d^3)}$

```
input int((B*x+A)/(b*x+a)^(5/2)/(e*x+d)^(3/2),x,method=_RETURNVERBOSE)
```

output

```
-2/3*(8*A*b^2*e^2*x^2-2*B*a*b*e^2*x^2-6*B*b^2*d*e*x^2+12*A*a*b*e^2*x+4*A*b^2*d*e*x-3*B*a^2*e^2*x-10*B*a*b*d*e*x-3*B*b^2*d^2*x+3*A*a^2*e^2+6*A*a*b*d*e-A*b^2*d^2-6*B*a^2*d*e-2*B*a*b*d^2)/(e*x+d)^(1/2)/(b*x+a)^(3/2)/(a*e-b*d)^3
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 337 vs. $2(127) = 254$.

Time = 0.95 (sec) , antiderivative size = 337, normalized size of antiderivative = 2.32

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{3/2}} dx = \frac{2(3Aa^2e^2 - (2Bab + Ab^2)d^2 - 6(Ba^2 - Aab)de - 2(3a^2b^3d^4 - 3a^3b^2d^3e + 3a^4bd^2e^2 - a^5de^3 + (b^5d^3e - 3ab^4d^2e^2 + 3a^2b^3de^3))}{3(a^2b^3d^4 - 3a^3b^2d^3e + 3a^4bd^2e^2 - a^5de^3 + (b^5d^3e - 3ab^4d^2e^2 + 3a^2b^3de^3)}$$

input

```
integrate((B*x+A)/(b*x+a)^(5/2)/(e*x+d)^(3/2),x, algorithm="fricas")
```

output

```
2/3*(3*A*a^2*e^2 - (2*B*a*b + A*b^2)*d^2 - 6*(B*a^2 - A*a*b)*d*e - 2*(3*B*b^2*d*e + (B*a*b - 4*A*b^2)*e^2)*x^2 - (3*B*b^2*d^2 + 2*(5*B*a*b - 2*A*b^2)*d*e + 3*(B*a^2 - 4*A*a*b)*e^2)*x)*sqrt(b*x + a)*sqrt(e*x + d)/(a^2*b^3*d^4 - 3*a^3*b^2*d^3*e + 3*a^4*b*d^2*e^2 - a^5*d*e^3 + (b^5*d^3*e - 3*a*b^4*d^2*e^2 + 3*a^2*b^3*d*e^3 - a^3*b^2*e^4)*x^3 + (b^5*d^4 - a*b^4*d^3*e - 3*a^2*b^3*d^2*e^2 + 5*a^3*b^2*d*e^3 - 2*a^4*b*e^4)*x^2 + (2*a*b^4*d^4 - 5*a^2*b^3*d^3*e + 3*a^3*b^2*d^2*e^2 + a^4*b*d*e^3 - a^5*e^4)*x)
```

Sympy [F]

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{3/2}} dx = \int \frac{A + Bx}{(a + bx)^{\frac{5}{2}}(d + ex)^{\frac{3}{2}}} dx$$

input

```
integrate((B*x+A)/(b*x+a)**(5/2)/(e*x+d)**(3/2),x)
```

output

```
Integral((A + B*x)/((a + b*x)**(5/2)*(d + e*x)**(3/2)), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{3/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^(5/2)/(e*x+d)^(3/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 610 vs. 2(127) = 254.

Time = 0.31 (sec) , antiderivative size = 610, normalized size of antiderivative = 4.21

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{3/2}} dx =$$

$$\frac{2(Bb^2de - Ab^2e^2)\sqrt{bx + a}}{(b^3d^3|b| - 3ab^2d^2e|b| + 3a^2bde^2|b| - a^3e^3|b|)\sqrt{b^2d + (bx + a)be - abe}}$$

$$4 \left(3\sqrt{be}Bb^6d^3 - 4\sqrt{be}Bab^5d^2e - 5\sqrt{be}Ab^6d^2e - \sqrt{be}Ba^2b^4de^2 + 10\sqrt{be}Aab^5de^2 + 2\sqrt{be}Ba^3b^3e^3 - 5 \right)$$

input `integrate((B*x+A)/(b*x+a)^(5/2)/(e*x+d)^(3/2),x, algorithm="giac")`

output

```

-2*(B*b^2*d*e - A*b^2*e^2)*sqrt(b*x + a)/((b^3*d^3*abs(b) - 3*a*b^2*d^2*e*
abs(b) + 3*a^2*b*d*e^2*abs(b) - a^3*e^3*abs(b))*sqrt(b^2*d + (b*x + a)*b*e
- a*b*e)) - 4/3*(3*sqrt(b*e)*B*b^6*d^3 - 4*sqrt(b*e)*B*a*b^5*d^2*e - 5*sq
rt(b*e)*A*b^6*d^2*e - sqrt(b*e)*B*a^2*b^4*d*e^2 + 10*sqrt(b*e)*A*a*b^5*d*e
^2 + 2*sqrt(b*e)*B*a^3*b^3*e^3 - 5*sqrt(b*e)*A*a^2*b^4*e^3 - 6*sqrt(b*e)*(
sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*B*b^4*d^2
+ 12*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*
b*e))^2*A*b^4*d*e + 6*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b
*x + a)*b*e - a*b*e))^2*B*a^2*b^2*e^2 - 12*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x +
a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*A*a*b^3*e^2 + 3*sqrt(b*e)*(sq
rt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^4*B*b^2*d - 3
*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))
^4*A*b^2*e)/((b^2*d^2*abs(b) - 2*a*b*d*e*abs(b) + a^2*e^2*abs(b))*(b^2*d -
a*b*e - (sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2
)^3)

```

Mupad [B] (verification not implemented)

Time = 1.70 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.33

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{3/2}} dx = \frac{\sqrt{d + ex} \left(\frac{4x^2(Bae - 4Abe + 3Bbd)}{3(ae - bd)^3} + \frac{12Ba^2de - 6Aa^2e^2 + 4Babd^2 - 12Aabde + 2Ab^2d^2}{3be(ae - bd)^3} \right)}{x^2 \sqrt{a + bx} + \frac{ad\sqrt{a+bx}}{be} + \frac{x(ae+bd)\sqrt{a+bx}}{be}}$$

input

```
int((A + B*x)/((a + b*x)^(5/2)*(d + e*x)^(3/2)),x)
```

output

```

((d + e*x)^(1/2)*((4*x^2*(B*a*e - 4*A*b*e + 3*B*b*d))/(3*(a*e - b*d)^3) +
(2*A*b^2*d^2 - 6*A*a^2*e^2 + 4*B*a*b*d^2 + 12*B*a^2*d*e - 12*A*a*b*d*e)/(3
*b*e*(a*e - b*d)^3) + (2*x*(3*a*e + b*d)*(B*a*e - 4*A*b*e + 3*B*b*d))/(3*b
*e*(a*e - b*d)^3))/((x^2*(a + b*x)^(1/2) + (a*d*(a + b*x)^(1/2))/(b*e) + (
x*(a*e + b*d)*(a + b*x)^(1/2))/(b*e))

```


Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.83

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{3/2}} dx = \frac{-4\sqrt{e}\sqrt{b}\sqrt{bx+a}d - 4\sqrt{e}\sqrt{b}\sqrt{bx+a}ex - 2\sqrt{ex+d}ae - 2\sqrt{ex+d}bd - 2\sqrt{ex+d}d^2e}{\sqrt{bx+a}(a^2e^3x - 2abd e^2x + b^2d^2ex + a^2de^2 - 2abd^2e + b^2d^2)}$$

input `int((B*x+A)/(b*x+a)^(5/2)/(e*x+d)^(3/2),x)`output `(2*(- 2*sqrt(e)*sqrt(b)*sqrt(a + b*x)*d - 2*sqrt(e)*sqrt(b)*sqrt(a + b*x)*e*x - sqrt(d + e*x)*a*e - sqrt(d + e*x)*b*d - 2*sqrt(d + e*x)*b*e*x))/(sqrt(a + b*x)*(a**2*d*e**2 + a**2*e**3*x - 2*a*b*d**2*e - 2*a*b*d*e**2*x + b**2*d**3 + b**2*d**2*e*x))`

3.228 $\int \frac{A+Bx}{(a+bx)^{5/2}(d+ex)^{5/2}} dx$

Optimal result 2053
 Mathematica [A] (verified) 2054
 Rubi [A] (verified) 2054
 Maple [A] (verified) 2056
 Fracas [B] (verification not implemented) 2057
 Sympy [F] 2057
 Maxima [F(-2)] 2058
 Giac [B] (verification not implemented) 2058
 Mupad [B] (verification not implemented) 2059
 Reduce [B] (verification not implemented) 2060

Optimal result

Integrand size = 24, antiderivative size = 192

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{5/2}} dx = -\frac{2(Ab - aB)}{3b(bd - ae)(a + bx)^{3/2}(d + ex)^{3/2}} - \frac{2(bBd - 2Abe + aBe)}{b(bd - ae)^2\sqrt{a + bx}(d + ex)^{3/2}} + \frac{8e(2Abe - B(bd + ae))\sqrt{a + bx}}{3b(bd - ae)^3(d + ex)^{3/2}} + \frac{16e(2Abe - B(bd + ae))\sqrt{a + bx}}{3(bd - ae)^4\sqrt{d + ex}}$$

output

```
1/3*(-2*A*b+2*B*a)/b/(-a*e+b*d)/(b*x+a)^(3/2)/(e*x+d)^(3/2)-2*(-2*A*b*e+B*
a*e+B*b*d)/b/(-a*e+b*d)^2/(b*x+a)^(1/2)/(e*x+d)^(3/2)+8/3*e*(2*A*b*e-B*(a*
e+b*d))*(b*x+a)^(1/2)/b/(-a*e+b*d)^3/(e*x+d)^(3/2)+16/3*e*(2*A*b*e-B*(a*e+
b*d))*(b*x+a)^(1/2)/(-a*e+b*d)^4/(e*x+d)^(1/2)
```

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.03

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{5/2}} dx = \frac{2(-Bde^2(a + bx)^3 + Ae^3(a + bx)^3 + 6bBde(a + bx)^2(d + ex) - 9Abe^2(a + bx)^2(d + ex) + 3aBe^2(a + bx)(d + ex) - 3abd^2)}{3(bd - ae)^2(a + bx)^{3/2}(d + ex)^{3/2}}$$

input

```
Integrate[(A + B*x)/((a + b*x)^(5/2)*(d + e*x)^(5/2)),x]
```

output

```
(-2*(-(B*d*e^2*(a + b*x)^3) + A*e^3*(a + b*x)^3 + 6*b*B*d*e*(a + b*x)^2*(d + e*x) - 9*A*b*e^2*(a + b*x)^2*(d + e*x) + 3*a*B*e^2*(a + b*x)^2*(d + e*x) + 3*b^2*B*d*(a + b*x)*(d + e*x)^2 - 9*A*b^2*e*(a + b*x)*(d + e*x)^2 + 6*a*b*B*e*(a + b*x)*(d + e*x)^2 + A*b^3*(d + e*x)^3 - a*b^2*B*(d + e*x)^3))/(3*(b*d - a*e)^4*(a + b*x)^(3/2)*(d + e*x)^(3/2))
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 183, normalized size of antiderivative = 0.95, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {87, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{5/2}} dx$$

↓ 87

$$\frac{(aBe - 2Abe + bBd) \int \frac{1}{(a+bx)^{5/2}(d+ex)^{3/2}} dx - \frac{2(Bd - Ae)}{3e(a + bx)^{3/2}(d + ex)^{3/2}(bd - ae)}}{e(bd - ae)}$$

↓ 55

$$\begin{aligned}
 & \frac{(aBe - 2Abe + bBd) \left(-\frac{4e \int \frac{1}{(a+bx)^{3/2}(d+ex)^{3/2}} dx}{3(bd-ae)} - \frac{2}{3(a+bx)^{3/2}\sqrt{d+ex}(bd-ae)} \right)}{\frac{e(bd-ae)}{2(Bd-Ae)} \frac{1}{3e(a+bx)^{3/2}(d+ex)^{3/2}(bd-ae)}} \\
 & \quad \downarrow 55 \\
 & \frac{(aBe - 2Abe + bBd) \left(-\frac{4e \left(-\frac{2e \int \frac{1}{\sqrt{a+bx}(d+ex)^{3/2}} dx}{bd-ae} - \frac{2}{\sqrt{a+bx}\sqrt{d+ex}(bd-ae)} \right)}{3(bd-ae)} - \frac{2}{3(a+bx)^{3/2}\sqrt{d+ex}(bd-ae)} \right)}{\frac{e(bd-ae)}{2(Bd-Ae)} \frac{1}{3e(a+bx)^{3/2}(d+ex)^{3/2}(bd-ae)}} \\
 & \quad \downarrow 48 \\
 & \frac{\left(-\frac{4e \left(-\frac{4e\sqrt{a+bx}}{\sqrt{d+ex}(bd-ae)^2} - \frac{2}{\sqrt{a+bx}\sqrt{d+ex}(bd-ae)} \right)}{3(bd-ae)} - \frac{2}{3(a+bx)^{3/2}\sqrt{d+ex}(bd-ae)} \right) (aBe - 2Abe + bBd)}{e(bd-ae)}
 \end{aligned}$$

input `Int[(A + B*x)/((a + b*x)^(5/2)*(d + e*x)^(5/2)),x]`

output `(-2*(B*d - A*e))/(3*e*(b*d - a*e)*(a + b*x)^(3/2)*(d + e*x)^(3/2)) - ((b*B*d - 2*A*b*e + a*B*e)*(-2/(3*(b*d - a*e)*(a + b*x)^(3/2)*Sqrt[d + e*x]) - (4*e*(-2/((b*d - a*e)*Sqrt[a + b*x]*Sqrt[d + e*x]) - (4*e*Sqrt[a + b*x])/(b*d - a*e)^2*Sqrt[d + e*x])))/(3*(b*d - a*e)))/(e*(b*d - a*e))`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 565 vs. $2(170) = 340$.

Time = 3.15 (sec) , antiderivative size = 565, normalized size of antiderivative = 2.94

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{5/2}} dx = \frac{2(Aa^3e^3 + (2Bab^2 + Ab^3)d^3 + 3(4Ba^2b - 3Aab^2)d^2e + (2Ba^3 - 9Aa^2b)de^2 + 8A^2b^2e^3 + 3A^2b^2d^2e + 3A^2b^2d^2e^2 + 3A^2b^2d^2e^3 + 3A^2b^2d^2e^4 + 3A^2b^2d^2e^5 + 3A^2b^2d^2e^6 + 3A^2b^2d^2e^7 + 3A^2b^2d^2e^8 + 3A^2b^2d^2e^9 + 3A^2b^2d^2e^{10})}{3(a^2b^4d^6 - 4a^3b^3d^5e + 6a^4b^2d^4e^2 - 4a^5bd^3e^3 + a^6d^2e^4 + (b^6d^4e^2 - 4ab^5d^3e^3 + 6a^2b^4d^2e^4 - 4a^3b^3de^5 + 3a^4b^2d^2e^6 - 4a^5bd^3e^7 + a^6d^2e^8 + b^6d^4e^{10} - 4ab^5d^3e^{11} + 6a^2b^4d^2e^{12} - 4a^3b^3de^{13} + 3a^4b^2d^2e^{14} - 4a^5bd^3e^{15} + a^6d^2e^{16})}$$

input `integrate((B*x+A)/(b*x+a)^(5/2)/(e*x+d)^(5/2),x, algorithm="fricas")`

output

```
-2/3*(A*a^3*e^3 + (2*B*a*b^2 + A*b^3)*d^3 + 3*(4*B*a^2*b - 3*A*a*b^2)*d^2*
e + (2*B*a^3 - 9*A*a^2*b)*d*e^2 + 8*(B*b^3*d*e^2 + (B*a*b^2 - 2*A*b^3)*e^3
)*x^3 + 12*(B*b^3*d^2*e + 2*(B*a*b^2 - A*b^3)*d*e^2 + (B*a^2*b - 2*A*a*b^2
)*e^3)*x^2 + 3*(B*b^3*d^3 + (7*B*a*b^2 - 2*A*b^3)*d^2*e + (7*B*a^2*b - 12*
A*a*b^2)*d*e^2 + (B*a^3 - 2*A*a^2*b)*e^3)*x)*sqrt(b*x + a)*sqrt(e*x + d)/(
a^2*b^4*d^6 - 4*a^3*b^3*d^5*e + 6*a^4*b^2*d^4*e^2 - 4*a^5*b*d^3*e^3 + a^6*
d^2*e^4 + (b^6*d^4*e^2 - 4*a*b^5*d^3*e^3 + 6*a^2*b^4*d^2*e^4 - 4*a^3*b^3*d
*e^5 + a^4*b^2*e^6)*x^4 + 2*(b^6*d^5*e - 3*a*b^5*d^4*e^2 + 2*a^2*b^4*d^3*e
^3 + 2*a^3*b^3*d^2*e^4 - 3*a^4*b^2*d*e^5 + a^5*b*e^6)*x^3 + (b^6*d^6 - 9*a
^2*b^4*d^4*e^2 + 16*a^3*b^3*d^3*e^3 - 9*a^4*b^2*d^2*e^4 + a^6*e^6)*x^2 + 2
*(a*b^5*d^6 - 3*a^2*b^4*d^5*e + 2*a^3*b^3*d^4*e^2 + 2*a^4*b^2*d^3*e^3 - 3*
a^5*b*d^2*e^4 + a^6*d*e^5)*x)
```

Sympy [F]

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{5/2}} dx = \int \frac{A + Bx}{(a + bx)^{\frac{5}{2}}(d + ex)^{\frac{5}{2}}} dx$$

input `integrate((B*x+A)/(b*x+a)**(5/2)/(e*x+d)**(5/2),x)`

output

```
Integral((A + B*x)/((a + b*x)**(5/2)*(d + e*x)**(5/2)), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^(5/2)/(e*x+d)^(5/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*e-b*d>0)', see `assume?` for more detail)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1183 vs. 2(170) = 340.

Time = 0.46 (sec) , antiderivative size = 1183, normalized size of antiderivative = 6.16

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{5/2}} dx = \text{Too large to display}$$

input `integrate((B*x+A)/(b*x+a)^(5/2)/(e*x+d)^(5/2),x, algorithm="giac")`

output

```
-2/3*sqrt(b*x + a)*((5*B*b^7*d^4*e^3*abs(b) - 12*B*a*b^6*d^3*e^4*abs(b) -
8*A*b^7*d^3*e^4*abs(b) + 6*B*a^2*b^5*d^2*e^5*abs(b) + 24*A*a*b^6*d^2*e^5*a
bs(b) + 4*B*a^3*b^4*d*e^6*abs(b) - 24*A*a^2*b^5*d*e^6*abs(b) - 3*B*a^4*b^3
*e^7*abs(b) + 8*A*a^3*b^4*e^7*abs(b))*(b*x + a)/(b^9*d^7*e - 7*a*b^8*d^6*e
^2 + 21*a^2*b^7*d^5*e^3 - 35*a^3*b^6*d^4*e^4 + 35*a^4*b^5*d^3*e^5 - 21*a^5
*b^4*d^2*e^6 + 7*a^6*b^3*d*e^7 - a^7*b^2*e^8) + 3*(2*B*b^8*d^5*e^2*abs(b)
- 7*B*a*b^7*d^4*e^3*abs(b) - 3*A*b^8*d^4*e^3*abs(b) + 8*B*a^2*b^6*d^3*e^4*
abs(b) + 12*A*a*b^7*d^3*e^4*abs(b) - 2*B*a^3*b^5*d^2*e^5*abs(b) - 18*A*a^2
*b^6*d^2*e^5*abs(b) - 2*B*a^4*b^4*d*e^6*abs(b) + 12*A*a^3*b^5*d*e^6*abs(b)
+ B*a^5*b^3*e^7*abs(b) - 3*A*a^4*b^4*e^7*abs(b))/(b^9*d^7*e - 7*a*b^8*d^6
*e^2 + 21*a^2*b^7*d^5*e^3 - 35*a^3*b^6*d^4*e^4 + 35*a^4*b^5*d^3*e^5 - 21*a
^5*b^4*d^2*e^6 + 7*a^6*b^3*d*e^7 - a^7*b^2*e^8))/(b^2*d + (b*x + a)*b*e -
a*b*e)^(3/2) - 4/3*(3*sqrt(b*e)*B*b^7*d^3 - sqrt(b*e)*B*a*b^6*d^2*e - 8*sq
rt(b*e)*A*b^7*d^2*e - 7*sqrt(b*e)*B*a^2*b^5*d*e^2 + 16*sqrt(b*e)*A*a*b^6*d
*e^2 + 5*sqrt(b*e)*B*a^3*b^4*e^3 - 8*sqrt(b*e)*A*a^2*b^5*e^3 - 6*sqrt(b*e)
*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*B*b^5*d
^2 - 6*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d + (b*x + a)*b*e - a
*b*e))^2*B*a*b^4*d*e + 18*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x + a) - sqrt(b^2*d
+ (b*x + a)*b*e - a*b*e))^2*A*b^5*d*e + 12*sqrt(b*e)*(sqrt(b*e)*sqrt(b*x +
a) - sqrt(b^2*d + (b*x + a)*b*e - a*b*e))^2*B*a^2*b^3*e^2 - 18*sqrt(b*...
```

Mupad [B] (verification not implemented)

Time = 1.87 (sec) , antiderivative size = 304, normalized size of antiderivative = 1.58

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{5/2}} dx = \frac{\sqrt{d + ex} \left(\frac{16bx^3(Bae - 2Abe + Bbd)}{3(ae - bd)^4} + \frac{4Ba^3de^2 + 2Aa^3e^3 + 24Ba^2bd^2e - 18Aa^2bde^2 + 4Bab^2d^3 - 18Aab^2d^2e + 2Ab^3d^3}{3be^2(ae - bd)^4} + 8 \right)}{x^3 \sqrt{a + bx} + \frac{ad^2 \sqrt{a + bx}}{be^2} + \frac{x^2(ae + 2bd) \sqrt{a + bx}}{be} + \frac{dx(2a + b)}{e}}$$

input

```
int((A + B*x)/((a + b*x)^(5/2)*(d + e*x)^(5/2)),x)
```


output

```

-((d + e*x)^(1/2)*((16*b*x^3*(B*a*e - 2*A*b*e + B*b*d))/(3*(a*e - b*d)^4)
+ (2*A*a^3*e^3 + 2*A*b^3*d^3 + 4*B*a*b^2*d^3 + 4*B*a^3*d*e^2 - 18*A*a*b^2*
d^2*e - 18*A*a^2*b*d*e^2 + 24*B*a^2*b*d^2*e)/(3*b*e^2*(a*e - b*d)^4) + (8*
x^2*(a*e + b*d)*(B*a*e - 2*A*b*e + B*b*d))/(e*(a*e - b*d)^4) + (2*x*(a^2*e
^2 + b^2*d^2 + 6*a*b*d*e)*(B*a*e - 2*A*b*e + B*b*d))/(b*e^2*(a*e - b*d)^4)
))/((x^3*(a + b*x)^(1/2) + (a*d^2*(a + b*x)^(1/2))/(b*e^2) + (x^2*(a*e + 2*
b*d)*(a + b*x)^(1/2))/(b*e) + (d*x*(2*a*e + b*d)*(a + b*x)^(1/2))/(b*e^2))

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 288, normalized size of antiderivative = 1.50

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{5/2}} dx = \frac{-\frac{16\sqrt{e}\sqrt{b}\sqrt{bx+abd^2}}{3} - \frac{32\sqrt{e}\sqrt{b}\sqrt{bx+abdex}}{3} - \frac{16\sqrt{e}\sqrt{b}\sqrt{bx+abe^2x^2}}{3} - \frac{2\sqrt{ex+da^2e^2}}{3} + 4\sqrt{ex+da^2e^2}}{\sqrt{bx+a}(a^3e^5x^2 - 3a^2bde^4x^2 + 3ab^2d^2e^3x^2 - b^3d^3e^2x^2 + 2a^3de^4x - b^3d^3e^2x^2)}$$

input

```
int((B*x+A)/(b*x+a)^(5/2)/(e*x+d)^(5/2),x)
```

output

```

(2*( - 8*sqrt(e)*sqrt(b)*sqrt(a + b*x)*b*d**2 - 16*sqrt(e)*sqrt(b)*sqrt(a
+ b*x)*b*d*e*x - 8*sqrt(e)*sqrt(b)*sqrt(a + b*x)*b*e**2*x**2 - sqrt(d + e*
x)*a**2*e**2 + 6*sqrt(d + e*x)*a*b*d*e + 4*sqrt(d + e*x)*a*b*e**2*x + 3*sq
rt(d + e*x)*b**2*d**2 + 12*sqrt(d + e*x)*b**2*d*e*x + 8*sqrt(d + e*x)*b**2
*e**2*x**2))/(3*sqrt(a + b*x)*(a**3*d**2*e**3 + 2*a**3*d*e**4*x + a**3*e**
5*x**2 - 3*a**2*b*d**3*e**2 - 6*a**2*b*d**2*e**3*x - 3*a**2*b*d*e**4*x**2
+ 3*a*b**2*d**4*e + 6*a*b**2*d**3*e**2*x + 3*a*b**2*d**2*e**3*x**2 - b**3*
d**5 - 2*b**3*d**4*e*x - b**3*d**3*e**2*x**2))

```

3.229 $\int \frac{A+Bx}{(a+bx)^{5/2}(d+ex)^{7/2}} dx$

Optimal result	2061
Mathematica [A] (verified)	2062
Rubi [A] (verified)	2062
Maple [B] (verified)	2064
Fricas [B] (verification not implemented)	2065
Sympy [F]	2066
Maxima [F(-2)]	2067
Giac [B] (verification not implemented)	2067
Mupad [B] (verification not implemented)	2068
Reduce [B] (verification not implemented)	2069

Optimal result

Integrand size = 24, antiderivative size = 246

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{7/2}} dx = -\frac{2(Ab - aB)}{3b(bd - ae)(a + bx)^{3/2}(d + ex)^{5/2}} - \frac{2(3bBd - 8Abe + 5aBe)}{3b(bd - ae)^2\sqrt{a + bx}(d + ex)^{5/2}} - \frac{4e(3bBd - 8Abe + 5aBe)\sqrt{a + bx}}{5b(bd - ae)^3(d + ex)^{5/2}} - \frac{16e(3bBd - 8Abe + 5aBe)\sqrt{a + bx}}{15(bd - ae)^4(d + ex)^{3/2}} - \frac{32be(3bBd - 8Abe + 5aBe)\sqrt{a + bx}}{15(bd - ae)^5\sqrt{d + ex}}$$

output

```
1/3*(-2*A*b+2*B*a)/b/(-a*e+b*d)/(b*x+a)^(3/2)/(e*x+d)^(5/2)-2/3*(-8*A*b*e+
5*B*a*e+3*B*b*d)/b/(-a*e+b*d)^2/(b*x+a)^(1/2)/(e*x+d)^(5/2)-4/5*e*(-8*A*b*
e+5*B*a*e+3*B*b*d)*(b*x+a)^(1/2)/b/(-a*e+b*d)^3/(e*x+d)^(5/2)-16/15*e*(-8*
A*b*e+5*B*a*e+3*B*b*d)*(b*x+a)^(1/2)/(-a*e+b*d)^4/(e*x+d)^(3/2)-32/15*b*e*
(-8*A*b*e+5*B*a*e+3*B*b*d)*(b*x+a)^(1/2)/(-a*e+b*d)^5/(e*x+d)^(1/2)
```

Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.10

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{7/2}} dx = \frac{2(3Bde^3(a + bx)^4 - 3Ae^4(a + bx)^4 - 15bBde^2(a + bx)^3(d + ex) + 20Abe^3(a + bx)^3(d + ex) - 5aBe^3(a$$

input

```
Integrate[(A + B*x)/((a + b*x)^(5/2)*(d + e*x)^(7/2)),x]
```

output

```
(-2*(3*B*d*e^3*(a + b*x)^4 - 3*A*e^4*(a + b*x)^4 - 15*b*B*d*e^2*(a + b*x)^3*(d + e*x) + 20*A*b*e^3*(a + b*x)^3*(d + e*x) - 5*a*B*e^3*(a + b*x)^3*(d + e*x) + 45*b^2*B*d*e*(a + b*x)^2*(d + e*x)^2 - 90*A*b^2*e^2*(a + b*x)^2*(d + e*x)^2 + 45*a*b*B*e^2*(a + b*x)^2*(d + e*x)^2 + 15*b^3*B*d*(a + b*x)*(d + e*x)^3 - 60*A*b^3*e*(a + b*x)*(d + e*x)^3 + 45*a*b^2*B*e*(a + b*x)*(d + e*x)^3 + 5*A*b^4*(d + e*x)^4 - 5*a*b^3*B*(d + e*x)^4))/(15*(b*d - a*e)^5*(a + b*x)^(3/2)*(d + e*x)^(5/2))
```

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 235, normalized size of antiderivative = 0.96, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$, Rules used = {87, 55, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{7/2}} dx$$

$$\downarrow 87$$

$$\frac{(5aBe - 8Abe + 3bBd) \int \frac{1}{(a+bx)^{3/2}(d+ex)^{7/2}} dx}{3b(bd - ae)} - \frac{2(Ab - aB)}{3b(a + bx)^{3/2}(d + ex)^{5/2}(bd - ae)}$$

$$\downarrow 55$$

$$\begin{aligned}
 & \frac{(5aBe - 8Abe + 3bBd) \left(-\frac{6e \int \frac{1}{\sqrt{a+bx}(d+ex)^{7/2}} dx}{bd-ae} - \frac{2}{\sqrt{a+bx}(d+ex)^{5/2}(bd-ae)} \right)}{\frac{3b(bd - ae)}{2(Ab - aB)}} \\
 & \qquad \qquad \qquad \frac{3b(a + bx)^{3/2}(d + ex)^{5/2}(bd - ae)}{\downarrow 55} \\
 & \frac{(5aBe - 8Abe + 3bBd) \left(-\frac{6e \left(\frac{4b \int \frac{1}{\sqrt{a+bx}(d+ex)^{5/2}} dx}{5(bd-ae)} + \frac{2\sqrt{a+bx}}{5(d+ex)^{5/2}(bd-ae)} \right)}{bd-ae} - \frac{2}{\sqrt{a+bx}(d+ex)^{5/2}(bd-ae)} \right)}{\frac{3b(bd - ae)}{2(Ab - aB)}} \\
 & \qquad \qquad \qquad \frac{3b(a + bx)^{3/2}(d + ex)^{5/2}(bd - ae)}{\downarrow 55} \\
 & \frac{(5aBe - 8Abe + 3bBd) \left(-\frac{6e \left(\frac{4b \left(\frac{2b \int \frac{1}{\sqrt{a+bx}(d+ex)^{3/2}} dx}{3(bd-ae)} + \frac{2\sqrt{a+bx}}{3(d+ex)^{3/2}(bd-ae)} \right)}{5(bd-ae)} + \frac{2\sqrt{a+bx}}{5(d+ex)^{5/2}(bd-ae)} \right)}{bd-ae} - \frac{2}{\sqrt{a+bx}(d+ex)^{5/2}(bd-ae)} \right)}{\frac{3b(bd - ae)}{2(Ab - aB)}} \\
 & \qquad \qquad \qquad \frac{3b(a + bx)^{3/2}(d + ex)^{5/2}(bd - ae)}{\downarrow 48} \\
 & \left(-\frac{6e \left(\frac{4b \left(\frac{4b\sqrt{a+bx}}{3\sqrt{d+ex}(bd-ae)^2} + \frac{2\sqrt{a+bx}}{3(d+ex)^{3/2}(bd-ae)} \right)}{5(bd-ae)} + \frac{2\sqrt{a+bx}}{5(d+ex)^{5/2}(bd-ae)} \right)}{bd-ae} - \frac{2}{\sqrt{a+bx}(d+ex)^{5/2}(bd-ae)} \right) (5aBe - 8Abe + 3bBd) \\
 & \qquad \qquad \qquad \frac{3b(bd - ae)}{2(Ab - aB)} \\
 & \qquad \qquad \qquad \frac{3b(a + bx)^{3/2}(d + ex)^{5/2}(bd - ae)}{\downarrow 48}
 \end{aligned}$$

input

```
Int[(A + B*x)/((a + b*x)^(5/2)*(d + e*x)^(7/2)),x]
```

output

$$\begin{aligned} & \frac{(-2(A*b - a*B))/(3*b*(b*d - a*e)*(a + b*x)^{(3/2)}*(d + e*x)^{(5/2)}) + ((3*b*B*d - 8*A*b*e + 5*a*B*e)*(-2/((b*d - a*e)*\text{Sqrt}[a + b*x]*(d + e*x)^{(5/2)})) - (6*e*((2*\text{Sqrt}[a + b*x])/((5*(b*d - a*e)*(d + e*x)^{(5/2)})) + (4*b*((2*\text{Sqrt}[a + b*x])/((3*(b*d - a*e)*(d + e*x)^{(3/2)})) + (4*b*\text{Sqrt}[a + b*x])/((3*(b*d - a*e)^2*\text{Sqrt}[d + e*x])))/((5*(b*d - a*e))))/(b*d - a*e)))/(3*b*(b*d - a*e)) \end{aligned}$$

Defintions of rubi rules used

rule 48

$$\text{Int}[(a + b*x)^m * (c + d*x)^n, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{m+1} * (c + d*x)^{n+1} / ((b*c - a*d) * (m+1)), x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x \ \&\& \ \text{EqQ}[m + n + 2, 0] \ \&\& \ \text{NeQ}[m, -1]$$

rule 55

$$\begin{aligned} & \text{Int}[(a + b*x)^m * (c + d*x)^n, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{m+1} * (c + d*x)^{n+1} / ((b*c - a*d) * (m+1)), x] - \text{Simp}[d * (\text{Simplify}[m + n + 2] / ((b*c - a*d) * (m+1))) \ \text{Int}[(a + b*x)^{\text{Simplify}[m+1]} * (c + d*x)^n, x], x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x \ \&\& \ \text{ILtQ}[\text{Simplify}[m + n + 2], 0] \ \&\& \ \text{NeQ}[m, -1] \ \&\& \ !(\text{LtQ}[m, -1] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ (\text{EqQ}[a, 0] \ || \ (\text{NeQ}[c, 0] \ \&\& \ \text{LtQ}[m - n, 0] \ \&\& \ \text{IntegerQ}[n]))) \ \&\& \ (\text{SumSimplerQ}[m, 1] \ || \ !\text{SumSimplerQ}[n, 1]) \end{aligned}$$

rule 87

$$\begin{aligned} & \text{Int}[(a + b*x)^m * (c + d*x)^n * (e + f*x)^p, x_Symbol] \rightarrow \text{Simp}[(-b*e - a*f) * (c + d*x)^{n+1} * (e + f*x)^{p+1} / (f * (p+1) * (c*f - d*e)), x] - \text{Simp}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1))) / (f * (p + 1) * (c*f - d*e)) \ \text{Int}[(c + d*x)^n * (e + f*x)^{p+1}, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ (!\text{LtQ}[n, -1] \ || \ \text{IntegerQ}[p] \ || \ !(\text{IntegerQ}[n] \ || \ !(\text{EqQ}[e, 0] \ || \ (\text{EqQ}[c, 0] \ || \ \text{LtQ}[p, n]))) \end{aligned}$$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 448 vs. $2(217) = 434$.

Time = 0.35 (sec) , antiderivative size = 449, normalized size of antiderivative = 1.83

method	result
default	$-\frac{2(128A b^4 e^4 x^4 - 80B a b^3 e^4 x^4 - 48B b^4 d e^3 x^4 + 192A a b^3 e^4 x^3 + 320A b^4 d e^3 x^3 - 120B a^2 b^2 e^4 x^3 - 272B a b^3 d e^3 x^3 - 120B b^4 d^2 e^2 x^3}{(b^2 x^2 + 2b d x + d^2)^{5/2} (e x + d)^{7/2}}$
gospers	$-\frac{2(128A b^4 e^4 x^4 - 80B a b^3 e^4 x^4 - 48B b^4 d e^3 x^4 + 192A a b^3 e^4 x^3 + 320A b^4 d e^3 x^3 - 120B a^2 b^2 e^4 x^3 - 272B a b^3 d e^3 x^3 - 120B b^4 d^2 e^2 x^3}{(b^2 x^2 + 2b d x + d^2)^{5/2} (e x + d)^{7/2}}$
orering	$-\frac{2(128A b^4 e^4 x^4 - 80B a b^3 e^4 x^4 - 48B b^4 d e^3 x^4 + 192A a b^3 e^4 x^3 + 320A b^4 d e^3 x^3 - 120B a^2 b^2 e^4 x^3 - 272B a b^3 d e^3 x^3 - 120B b^4 d^2 e^2 x^3}{(b^2 x^2 + 2b d x + d^2)^{5/2} (e x + d)^{7/2}}$

```
input int((B*x+A)/(b*x+a)^(5/2)/(e*x+d)^(7/2),x,method=_RETURNVERBOSE)
```

```
output -2/15*(128*A*b^4*e^4*x^4-80*B*a*b^3*e^4*x^4-48*B*b^4*d*e^3*x^4+192*A*a*b^3
*e^4*x^3+320*A*b^4*d*e^3*x^3-120*B*a^2*b^2*e^4*x^3-272*B*a*b^3*d*e^3*x^3-1
20*B*b^4*d^2*e^2*x^3+48*A*a^2*b^2*e^4*x^2+480*A*a*b^3*d*e^3*x^2+240*A*b^4*
d^2*e^2*x^2-30*B*a^3*b*e^4*x^2-318*B*a^2*b^2*d*e^3*x^2-330*B*a*b^3*d^2*e^2
*x^2-90*B*b^4*d^3*e*x^2-8*A*a^3*b*e^4*x+120*A*a^2*b^2*d*e^3*x+360*A*a*b^3*
d^2*e^2*x+40*A*b^4*d^3*e*x+5*B*a^4*e^4*x-72*B*a^3*b*d*e^3*x-270*B*a^2*b^2*
d^2*e^2*x-160*B*a*b^3*d^3*e*x-15*B*b^4*d^4*x+3*A*a^4*e^4-20*A*a^3*b*d*e^3+
90*A*a^2*b^2*d^2*e^2+60*A*a*b^3*d^3*e-5*A*b^4*d^4+2*B*a^4*d*e^3-30*B*a^3*b
*d^2*e^2-90*B*a^2*b^2*d^3*e-10*B*a*b^3*d^4)/(e*x+d)^(5/2)/(b*x+a)^(3/2)/(a
*e-b*d)^5
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 916 vs. 2(216) = 432.

Time = 24.16 (sec) , antiderivative size = 916, normalized size of antiderivative = 3.72

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{7/2}} dx = \text{Too large to display}$$

```
input integrate((B*x+A)/(b*x+a)^(5/2)/(e*x+d)^(7/2),x, algorithm="fricas")
```

output

```

2/15*(3*A*a^4*e^4 - 5*(2*B*a*b^3 + A*b^4)*d^4 - 30*(3*B*a^2*b^2 - 2*A*a*b^
3)*d^3*e - 30*(B*a^3*b - 3*A*a^2*b^2)*d^2*e^2 + 2*(B*a^4 - 10*A*a^3*b)*d*e
^3 - 16*(3*B*b^4*d*e^3 + (5*B*a*b^3 - 8*A*b^4)*e^4)*x^4 - 8*(15*B*b^4*d^2*
e^2 + 2*(17*B*a*b^3 - 20*A*b^4)*d*e^3 + 3*(5*B*a^2*b^2 - 8*A*a*b^3)*e^4)*x
^3 - 6*(15*B*b^4*d^3*e + 5*(11*B*a*b^3 - 8*A*b^4)*d^2*e^2 + (53*B*a^2*b^2
- 80*A*a*b^3)*d*e^3 + (5*B*a^3*b - 8*A*a^2*b^2)*e^4)*x^2 - (15*B*b^4*d^4 +
40*(4*B*a*b^3 - A*b^4)*d^3*e + 90*(3*B*a^2*b^2 - 4*A*a*b^3)*d^2*e^2 + 24*
(3*B*a^3*b - 5*A*a^2*b^2)*d*e^3 - (5*B*a^4 - 8*A*a^3*b)*e^4)*x)*sqrt(b*x +
a)*sqrt(e*x + d)/(a^2*b^5*d^8 - 5*a^3*b^4*d^7*e + 10*a^4*b^3*d^6*e^2 - 10
*a^5*b^2*d^5*e^3 + 5*a^6*b*d^4*e^4 - a^7*d^3*e^5 + (b^7*d^5*e^3 - 5*a*b^6*
d^4*e^4 + 10*a^2*b^5*d^3*e^5 - 10*a^3*b^4*d^2*e^6 + 5*a^4*b^3*d*e^7 - a^5*
b^2*e^8)*x^5 + (3*b^7*d^6*e^2 - 13*a*b^6*d^5*e^3 + 20*a^2*b^5*d^4*e^4 - 10
*a^3*b^4*d^3*e^5 - 5*a^4*b^3*d^2*e^6 + 7*a^5*b^2*d*e^7 - 2*a^6*b*e^8)*x^4
+ (3*b^7*d^7*e - 9*a*b^6*d^6*e^2 + a^2*b^5*d^5*e^3 + 25*a^3*b^4*d^4*e^4 -
35*a^4*b^3*d^3*e^5 + 17*a^5*b^2*d^2*e^6 - a^6*b*d*e^7 - a^7*e^8)*x^3 + (b^
7*d^8 + a*b^6*d^7*e - 17*a^2*b^5*d^6*e^2 + 35*a^3*b^4*d^5*e^3 - 25*a^4*b^3
*d^4*e^4 - a^5*b^2*d^3*e^5 + 9*a^6*b*d^2*e^6 - 3*a^7*d*e^7)*x^2 + (2*a*b^6
*d^8 - 7*a^2*b^5*d^7*e + 5*a^3*b^4*d^6*e^2 + 10*a^4*b^3*d^5*e^3 - 20*a^5*b
^2*d^4*e^4 + 13*a^6*b*d^3*e^5 - 3*a^7*d^2*e^6)*x)

```

SymPy [F]

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{7/2}} dx = \int \frac{A + Bx}{(a + bx)^{\frac{5}{2}}(d + ex)^{\frac{7}{2}}} dx$$

input

```
integrate((B*x+A)/(b*x+a)**(5/2)/(e*x+d)**(7/2),x)
```

output

```
Integral((A + B*x)/((a + b*x)**(5/2)*(d + e*x)**(7/2)), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{7/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*x+A)/(b*x+a)^(5/2)/(e*x+d)^(7/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e*(a*e-b*d)>0)', see `assume?` f or more de

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2081 vs. 2(216) = 432.

Time = 1.00 (sec) , antiderivative size = 2081, normalized size of antiderivative = 8.46

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{7/2}} dx = \text{Too large to display}$$

input `integrate((B*x+A)/(b*x+a)^(5/2)/(e*x+d)^(7/2),x, algorithm="giac")`

output

```
((d + e*x)^(1/2)*((32*b^2*x^4*(5*B*a*e - 8*A*b*e + 3*B*b*d))/(15*(a*e - b*d)^5) + (10*A*b^4*d^4 - 6*A*a^4*e^4 + 20*B*a*b^3*d^4 - 4*B*a^4*d*e^3 + 180*B*a^2*b^2*d^3*e + 60*B*a^3*b*d^2*e^2 - 180*A*a^2*b^2*d^2*e^2 - 120*A*a*b^3*d^3*e + 40*A*a^3*b*d*e^3)/(15*b*e^3*(a*e - b*d)^5) + (4*x^2*(a^2*e^2 + 5*b^2*d^2 + 10*a*b*d*e)*(5*B*a*e - 8*A*b*e + 3*B*b*d))/(5*e^2*(a*e - b*d)^5) + (2*x*(5*B*a*e - 8*A*b*e + 3*B*b*d)*(5*b^3*d^3 - a^3*e^3 + 45*a*b^2*d^2*e + 15*a^2*b*d*e^2))/(15*b*e^3*(a*e - b*d)^5) + (16*b*x^3*(3*a*e + 5*b*d)*(5*B*a*e - 8*A*b*e + 3*B*b*d))/(15*e*(a*e - b*d)^5))/(x^4*(a + b*x)^(1/2) + (a*d^3*(a + b*x)^(1/2))/(b*e^3) + (x^3*(a*e + 3*b*d)*(a + b*x)^(1/2))/(b*e) + (3*d*x^2*(a*e + b*d)*(a + b*x)^(1/2))/(b*e^2) + (d^2*x*(3*a*e + b*d)*(a + b*x)^(1/2))/(b*e^3))
```

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 514, normalized size of antiderivative = 2.09

$$\int \frac{A + Bx}{(a + bx)^{5/2}(d + ex)^{7/2}} dx = \frac{32\sqrt{e}\sqrt{b}\sqrt{bx+a}b^2d^3}{5} + \frac{96\sqrt{e}\sqrt{b}\sqrt{bx+a}b^2d^2ex}{5} + \frac{96\sqrt{e}\sqrt{b}\sqrt{bx+a}b^2de^2x^2}{5} + \frac{32\sqrt{e}\sqrt{b}\sqrt{bx+a}b^2d^3ex^3}{5} + \dots$$

input

```
int((B*x+A)/(b*x+a)^(5/2)/(e*x+d)^(7/2),x)
```

output

```
(2*(16*sqrt(e)*sqrt(b)*sqrt(a + b*x)*b**2*d**3 + 48*sqrt(e)*sqrt(b)*sqrt(a + b*x)*b**2*d**2*e*x + 48*sqrt(e)*sqrt(b)*sqrt(a + b*x)*b**2*d*e**2*x**2 + 16*sqrt(e)*sqrt(b)*sqrt(a + b*x)*b**2*e**3*x**3 - sqrt(d + e*x)*a**3*e**3 + 5*sqrt(d + e*x)*a**2*b*d*e**2 + 2*sqrt(d + e*x)*a**2*b*e**3*x - 15*sqrt(d + e*x)*a*b**2*d**2*e - 20*sqrt(d + e*x)*a*b**2*d*e**2*x - 8*sqrt(d + e*x)*a*b**2*e**3*x**2 - 5*sqrt(d + e*x)*b**3*d**3 - 30*sqrt(d + e*x)*b**3*d**2*e*x - 40*sqrt(d + e*x)*b**3*d*e**2*x**2 - 16*sqrt(d + e*x)*b**3*e**3*x**3)/(5*sqrt(a + b*x)*(a**4*d**3*e**4 + 3*a**4*d**2*e**5*x + 3*a**4*d*e**6*x**2 + a**4*e**7*x**3 - 4*a**3*b*d**4*e**3 - 12*a**3*b*d**3*e**4*x - 12*a**3*b*d**2*e**5*x**2 - 4*a**3*b*d*e**6*x**3 + 6*a**2*b**2*d**5*e**2 + 18*a**2*b**2*d**4*e**3*x + 18*a**2*b**2*d**3*e**4*x**2 + 6*a**2*b**2*d**2*e**5*x**3 - 4*a*b**3*d**6*e - 12*a*b**3*d**5*e**2*x - 12*a*b**3*d**4*e**3*x**2 - 4*a*b**3*d**3*e**4*x**3 + b**4*d**7 + 3*b**4*d**6*e*x + 3*b**4*d**5*e**2*x**2 + b**4*d**4*e**3*x**3))
```

3.230 $\int (a + bx)^3 (A + Bx)(d + ex)^m dx$

Optimal result	2070
Mathematica [A] (verified)	2071
Rubi [A] (verified)	2071
Maple [B] (verified)	2072
Fricas [B] (verification not implemented)	2074
Sympy [B] (verification not implemented)	2075
Maxima [B] (verification not implemented)	2076
Giac [B] (verification not implemented)	2077
Mupad [B] (verification not implemented)	2078
Reduce [B] (verification not implemented)	2078

Optimal result

Integrand size = 20, antiderivative size = 186

$$\int (a + bx)^3 (A + Bx)(d + ex)^m dx = \frac{(bd - ae)^3 (Bd - Ae)(d + ex)^{1+m}}{e^5(1 + m)} - \frac{(bd - ae)^2 (4bBd - 3Abe - aBe)(d + ex)^{2+m}}{e^5(2 + m)} + \frac{3b(bd - ae)(2bBd - Abe - aBe)(d + ex)^{3+m}}{e^5(3 + m)} - \frac{b^2(4bBd - Abe - 3aBe)(d + ex)^{4+m}}{e^5(4 + m)} + \frac{b^3 B(d + ex)^{5+m}}{e^5(5 + m)}$$

output

```
(-a*e+b*d)^3*(-A*e+B*d)*(e*x+d)^(1+m)/e^5/(1+m)-(-a*e+b*d)^2*(-3*A*b*e-B*a
*e+4*B*b*d)*(e*x+d)^(2+m)/e^5/(2+m)+3*b*(-a*e+b*d)*(-A*b*e-B*a*e+2*B*b*d)*
(e*x+d)^(3+m)/e^5/(3+m)-b^2*(-A*b*e-3*B*a*e+4*B*b*d)*(e*x+d)^(4+m)/e^5/(4+
m)+b^3*B*(e*x+d)^(5+m)/e^5/(5+m)
```

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.89

$$\int (a + bx)^3(A + Bx)(d + ex)^m dx$$

$$= \frac{(d + ex)^{1+m} \left(\frac{(bd-ae)^3(Bd-Ae)}{1+m} - \frac{(bd-ae)^2(4bBd-3Abe-aBe)(d+ex)}{2+m} + \frac{3b(bd-ae)(2bBd-Abe-aBe)(d+ex)^2}{3+m} - \frac{b^2(4bBd-Abe-aBe)(d+ex)^3}{4+m} + \frac{b^3B(d+ex)^4}{5+m} \right)}{e^5}$$

input

```
Integrate[(a + b*x)^3*(A + B*x)*(d + e*x)^m,x]
```

output

```
((d + e*x)^(1 + m)*(((b*d - a*e)^3*(B*d - A*e))/(1 + m) - ((b*d - a*e)^2*(4*b*B*d - 3*A*b*e - a*B*e)*(d + e*x))/(2 + m) + (3*b*(b*d - a*e)*(2*b*B*d - A*b*e - a*B*e)*(d + e*x)^2)/(3 + m) - (b^2*(4*b*B*d - A*b*e - 3*a*B*e)*(d + e*x)^3)/(4 + m) + (b^3*B*(d + e*x)^4)/(5 + m)))/e^5
```

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^3(A + Bx)(d + ex)^m dx$$

$$\downarrow 86$$

$$\int \left(\frac{b^2(d + ex)^{m+3}(3aBe + Abe - 4bBd)}{e^4} + \frac{(ae - bd)^3(Ae - Bd)(d + ex)^m}{e^4} + \frac{(ae - bd)^2(d + ex)^{m+1}(aBe + 3aBe - 4bBd)}{e^4} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
& - \frac{b^2(d+ex)^{m+4}(-3aBe - Abe + 4bBd)}{e^5(m+4)} + \frac{(bd-ae)^3(Bd-Ae)(d+ex)^{m+1}}{e^5(m+1)} - \\
& \frac{(bd-ae)^2(d+ex)^{m+2}(-aBe - 3Abe + 4bBd)}{e^5(m+2)} + \\
& \frac{3b(bd-ae)(d+ex)^{m+3}(-aBe - Abe + 2bBd)}{e^5(m+3)} + \frac{b^3B(d+ex)^{m+5}}{e^5(m+5)}
\end{aligned}$$

input `Int[(a + b*x)^3*(A + B*x)*(d + e*x)^m,x]`

output `((b*d - a*e)^3*(B*d - A*e)*(d + e*x)^(1 + m))/(e^5*(1 + m)) - ((b*d - a*e)^2*(4*b*B*d - 3*A*b*e - a*B*e)*(d + e*x)^(2 + m))/(e^5*(2 + m)) + (3*b*(b*d - a*e)*(2*b*B*d - A*b*e - a*B*e)*(d + e*x)^(3 + m))/(e^5*(3 + m)) - (b^2*(4*b*B*d - A*b*e - 3*a*B*e)*(d + e*x)^(4 + m))/(e^5*(4 + m)) + (b^3*B*(d + e*x)^(5 + m))/(e^5*(5 + m))`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1164 vs. 2(186) = 372.

Time = 0.31 (sec) , antiderivative size = 1165, normalized size of antiderivative = 6.26

method	result	size
norman	Expression too large to display	1165
gosper	Expression too large to display	1270
oring	Expression too large to display	1273
risch	Expression too large to display	1712
parallelrisc	Expression too large to display	2612

```
input int((b*x+a)^3*(B*x+A)*(e*x+d)^m,x,method=_RETURNVERBOSE)
```

```
output b^3*B/(5+m)*x^5*exp(m*ln(e*x+d))+d*(A*a^3*e^4*m^4+14*A*a^3*e^4*m^3-3*A*a^2
*b*d*e^3*m^3-B*a^3*d*e^3*m^3+71*A*a^3*e^4*m^2-36*A*a^2*b*d*e^3*m^2+6*A*a*b
^2*d^2*e^2*m^2-12*B*a^3*d*e^3*m^2+6*B*a^2*b*d^2*e^2*m^2+154*A*a^3*e^4*m-14
1*A*a^2*b*d*e^3*m+54*A*a*b^2*d^2*e^2*m-6*A*b^3*d^3*e*m-47*B*a^3*d*e^3*m+54
*B*a^2*b*d^2*e^2*m-18*B*a*b^2*d^3*e*m+120*A*a^3*e^4-180*A*a^2*b*d*e^3+120*
A*a*b^2*d^2*e^2-30*A*b^3*d^3*e-60*B*a^3*d*e^3+120*B*a^2*b*d^2*e^2-90*B*a*b
^2*d^3*e+24*B*b^3*d^4)/e^5/(m^5+15*m^4+85*m^3+225*m^2+274*m+120)*exp(m*ln(
e*x+d))+(3*A*a^2*b*e^3*m^3+3*A*a*b^2*d*e^2*m^3+B*a^3*e^3*m^3+3*B*a^2*b*d*e
^2*m^3+36*A*a^2*b*e^3*m^2+27*A*a*b^2*d*e^2*m^2-3*A*b^3*d^2*e*m^2+12*B*a^3*
e^3*m^2+27*B*a^2*b*d*e^2*m^2-9*B*a*b^2*d^2*e*m^2+141*A*a^2*b*e^3*m+60*A*a*
b^2*d*e^2*m-15*A*b^3*d^2*e*m+47*B*a^3*e^3*m+60*B*a^2*b*d*e^2*m-45*B*a*b^2*
d^2*e*m+12*B*b^3*d^3*m+180*A*a^2*b*e^3+60*B*a^3*e^3)/e^3/(m^4+14*m^3+71*m^
2+154*m+120)*x^2*exp(m*ln(e*x+d))+(A*a^3*e^4*m^4+3*A*a^2*b*d*e^3*m^4+B*a^3
*d*e^3*m^4+14*A*a^3*e^4*m^3+36*A*a^2*b*d*e^3*m^3-6*A*a*b^2*d^2*e^2*m^3+12*
B*a^3*d*e^3*m^3-6*B*a^2*b*d^2*e^2*m^3+71*A*a^3*e^4*m^2+141*A*a^2*b*d*e^3*m
^2-54*A*a*b^2*d^2*e^2*m^2+6*A*b^3*d^3*e*m^2+47*B*a^3*d*e^3*m^2-54*B*a^2*b*
d^2*e^2*m^2+18*B*a*b^2*d^3*e*m^2+154*A*a^3*e^4*m+180*A*a^2*b*d*e^3*m-120*A
*a*b^2*d^2*e^2*m+30*A*b^3*d^3*e*m+60*B*a^3*d*e^3*m-120*B*a^2*b*d^2*e^2*m+9
0*B*a*b^2*d^3*e*m-24*B*b^3*d^4*m+120*A*a^3*e^4)/e^4/(m^5+15*m^4+85*m^3+225
*m^2+274*m+120)*x*exp(m*ln(e*x+d))+(A*b*e*m+3*B*a*e*m+B*b*d*m+5*A*b*e+1...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1282 vs. $2(186) = 372$.

Time = 0.13 (sec) , antiderivative size = 1282, normalized size of antiderivative = 6.89

$$\int (a + bx)^3 (A + Bx)(d + ex)^m dx = \text{Too large to display}$$

input `integrate((b*x+a)^3*(B*x+A)*(e*x+d)^m,x, algorithm="fricas")`

output

```
(A*a^3*d*e^4*m^4 + 24*B*b^3*d^5 + 120*A*a^3*d*e^4 - 30*(3*B*a*b^2 + A*b^3)
*d^4*e + 120*(B*a^2*b + A*a*b^2)*d^3*e^2 - 60*(B*a^3 + 3*A*a^2*b)*d^2*e^3
+ (B*b^3*e^5*m^4 + 10*B*b^3*e^5*m^3 + 35*B*b^3*e^5*m^2 + 50*B*b^3*e^5*m
+ 24*B*b^3*e^5)*x^5 + (30*(3*B*a*b^2 + A*b^3)*e^5 + (B*b^3*d*e^4 + (3*B*a*b^
2 + A*b^3)*e^5)*m^4 + (6*B*b^3*d*e^4 + 11*(3*B*a*b^2 + A*b^3)*e^5)*m^3 + (
11*B*b^3*d*e^4 + 41*(3*B*a*b^2 + A*b^3)*e^5)*m^2 + (6*B*b^3*d*e^4 + 61*(3*
B*a*b^2 + A*b^3)*e^5)*m)*x^4 + (14*A*a^3*d*e^4 - (B*a^3 + 3*A*a^2*b)*d^2*e
^3)*m^3 + (120*(B*a^2*b + A*a*b^2)*e^5 + ((3*B*a*b^2 + A*b^3)*d*e^4 + 3*(B
*a^2*b + A*a*b^2)*e^5)*m^4 - 4*(B*b^3*d^2*e^3 - 2*(3*B*a*b^2 + A*b^3)*d*e^
4 - 9*(B*a^2*b + A*a*b^2)*e^5)*m^3 - (12*B*b^3*d^2*e^3 - 17*(3*B*a*b^2 + A
*b^3)*d*e^4 - 147*(B*a^2*b + A*a*b^2)*e^5)*m^2 - 2*(4*B*b^3*d^2*e^3 - 5*(3
*B*a*b^2 + A*b^3)*d*e^4 - 117*(B*a^2*b + A*a*b^2)*e^5)*m)*x^3 + (71*A*a^3*
d*e^4 + 6*(B*a^2*b + A*a*b^2)*d^3*e^2 - 12*(B*a^3 + 3*A*a^2*b)*d^2*e^3)*m^
2 + (60*(B*a^3 + 3*A*a^2*b)*e^5 + (3*(B*a^2*b + A*a*b^2)*d*e^4 + (B*a^3 +
3*A*a^2*b)*e^5)*m^4 - (3*(3*B*a*b^2 + A*b^3)*d^2*e^3 - 30*(B*a^2*b + A*a*b
^2)*d*e^4 - 13*(B*a^3 + 3*A*a^2*b)*e^5)*m^3 + (12*B*b^3*d^3*e^2 - 18*(3*B*
a*b^2 + A*b^3)*d^2*e^3 + 87*(B*a^2*b + A*a*b^2)*d*e^4 + 59*(B*a^3 + 3*A*a^
2*b)*e^5)*m^2 + (12*B*b^3*d^3*e^2 - 15*(3*B*a*b^2 + A*b^3)*d^2*e^3 + 60*(B
*a^2*b + A*a*b^2)*d*e^4 + 107*(B*a^3 + 3*A*a^2*b)*e^5)*m)*x^2 + (154*A*a^3
*d*e^4 - 6*(3*B*a*b^2 + A*b^3)*d^4*e + 54*(B*a^2*b + A*a*b^2)*d^3*e^2 - ...
```


Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 620 vs. $2(186) = 372$.

Time = 0.08 (sec) , antiderivative size = 620, normalized size of antiderivative = 3.33

$$\int (a + bx)^3 (A + Bx)(d + ex)^m dx = \frac{(e^2(m+1)x^2 + demx - d^2)(ex + d)^m Ba^3}{(m^2 + 3m + 2)e^2} + \frac{3(e^2(m+1)x^2 + demx - d^2)(ex + d)^m Aa^2b}{(m^2 + 3m + 2)e^2} + \frac{(ex + d)^{m+1} Aa^3}{e(m+1)} + \frac{3((m^2 + 3m + 2)e^3x^3 + (m^2 + m)de^2x^2 - 2d^2emx + 2d^3)(ex + d)^m Ba^2b}{(m^3 + 6m^2 + 11m + 6)e^3} + \frac{3((m^2 + 3m + 2)e^3x^3 + (m^2 + m)de^2x^2 - 2d^2emx + 2d^3)(ex + d)^m Aab^2}{(m^3 + 6m^2 + 11m + 6)e^3} + \frac{3((m^3 + 6m^2 + 11m + 6)e^4x^4 + (m^3 + 3m^2 + 2m)de^3x^3 - 3(m^2 + m)d^2e^2x^2 + 6d^3emx - 6d^4)(ex + d)^m B^2a^2b^2}{(m^4 + 10m^3 + 35m^2 + 50m + 24)e^4} + \frac{((m^3 + 6m^2 + 11m + 6)e^4x^4 + (m^3 + 3m^2 + 2m)de^3x^3 - 3(m^2 + m)d^2e^2x^2 + 6d^3emx - 6d^4)(ex + d)^m A^2b^2}{(m^4 + 10m^3 + 35m^2 + 50m + 24)e^4} + \frac{((m^4 + 10m^3 + 35m^2 + 50m + 24)e^5x^5 + (m^4 + 6m^3 + 11m^2 + 6m)de^4x^4 - 4(m^3 + 3m^2 + 2m)d^3e^3x^3 + 12(m^2 + m)d^4e^2x^2 - 24d^4emx + 24d^5)(ex + d)^m B^2b^3}{(m^5 + 15m^4 + 85m^3 + 225m^2 + 274m + 120)e^5}$$

input `integrate((b*x+a)^3*(B*x+A)*(e*x+d)^m,x, algorithm="maxima")`

output `(e^2*(m + 1)*x^2 + d*e*m*x - d^2)*(e*x + d)^m*B*a^3/((m^2 + 3*m + 2)*e^2) + 3*(e^2*(m + 1)*x^2 + d*e*m*x - d^2)*(e*x + d)^m*A*a^2*b/((m^2 + 3*m + 2)*e^2) + (e*x + d)^(m + 1)*A*a^3/(e*(m + 1)) + 3*((m^2 + 3*m + 2)*e^3*x^3 + (m^2 + m)*d*e^2*x^2 - 2*d^2*e*m*x + 2*d^3)*(e*x + d)^m*B*a^2*b/((m^3 + 6*m^2 + 11*m + 6)*e^3) + 3*((m^2 + 3*m + 2)*e^3*x^3 + (m^2 + m)*d*e^2*x^2 - 2*d^2*e*m*x + 2*d^3)*(e*x + d)^m*A*a*b^2/((m^3 + 6*m^2 + 11*m + 6)*e^3) + 3*((m^3 + 6*m^2 + 11*m + 6)*e^4*x^4 + (m^3 + 3*m^2 + 2*m)*d*e^3*x^3 - 3*(m^2 + m)*d^2*e^2*x^2 + 6*d^3*e*m*x - 6*d^4)*(e*x + d)^m*B*a*b^2/((m^4 + 10*m^3 + 35*m^2 + 50*m + 24)*e^4) + ((m^3 + 6*m^2 + 11*m + 6)*e^4*x^4 + (m^3 + 3*m^2 + 2*m)*d*e^3*x^3 - 3*(m^2 + m)*d^2*e^2*x^2 + 6*d^3*e*m*x - 6*d^4)*(e*x + d)^m*A*b^3/((m^4 + 10*m^3 + 35*m^2 + 50*m + 24)*e^4) + ((m^4 + 10*m^3 + 35*m^2 + 50*m + 24)*e^5*x^5 + (m^4 + 6*m^3 + 11*m^2 + 6*m)*d*e^4*x^4 - 4*(m^3 + 3*m^2 + 2*m)*d^2*e^3*x^3 + 12*(m^2 + m)*d^3*e^2*x^2 - 24*d^4*e*m*x + 24*d^5)*(e*x + d)^m*B*b^3/((m^5 + 15*m^4 + 85*m^3 + 225*m^2 + 274*m + 120)*e^5)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2519 vs. $2(186) = 372$.

Time = 0.14 (sec) , antiderivative size = 2519, normalized size of antiderivative = 13.54

$$\int (a + bx)^3 (A + Bx)(d + ex)^m dx = \text{Too large to display}$$

input `integrate((b*x+a)^3*(B*x+A)*(e*x+d)^m,x, algorithm="giac")`

output

```
((e*x + d)^m*B*b^3*e^5*m^4*x^5 + (e*x + d)^m*B*b^3*d*e^4*m^4*x^4 + 3*(e*x + d)^m*B*a*b^2*e^5*m^4*x^4 + (e*x + d)^m*A*b^3*e^5*m^4*x^4 + 10*(e*x + d)^m*B*b^3*e^5*m^3*x^5 + 3*(e*x + d)^m*B*a*b^2*d*e^4*m^4*x^3 + (e*x + d)^m*A*b^3*d*e^4*m^4*x^3 + 3*(e*x + d)^m*B*a^2*b*e^5*m^4*x^3 + 3*(e*x + d)^m*A*a*b^2*e^5*m^4*x^3 + 6*(e*x + d)^m*B*b^3*d*e^4*m^3*x^4 + 33*(e*x + d)^m*B*a*b^2*e^5*m^3*x^4 + 11*(e*x + d)^m*A*b^3*e^5*m^3*x^4 + 35*(e*x + d)^m*B*b^3*e^5*m^2*x^5 + 3*(e*x + d)^m*B*a^2*b*d*e^4*m^4*x^2 + 3*(e*x + d)^m*A*a*b^2*d*e^4*m^4*x^2 + (e*x + d)^m*B*a^3*e^5*m^4*x^2 + 3*(e*x + d)^m*A*a^2*b*e^5*m^4*x^2 - 4*(e*x + d)^m*B*b^3*d^2*e^3*m^3*x^3 + 24*(e*x + d)^m*B*a*b^2*d*e^4*m^3*x^3 + 8*(e*x + d)^m*A*b^3*d*e^4*m^3*x^3 + 36*(e*x + d)^m*B*a^2*b*e^5*m^3*x^3 + 36*(e*x + d)^m*A*a*b^2*e^5*m^3*x^3 + 11*(e*x + d)^m*B*b^3*d*e^4*m^2*x^4 + 123*(e*x + d)^m*B*a*b^2*e^5*m^2*x^4 + 41*(e*x + d)^m*A*b^3*e^5*m^2*x^4 + 50*(e*x + d)^m*B*b^3*e^5*m*x^5 + (e*x + d)^m*B*a^3*d*e^4*m^4*x + 3*(e*x + d)^m*A*a^2*b*d*e^4*m^4*x + (e*x + d)^m*A*a^3*e^5*m^4*x - 9*(e*x + d)^m*B*a*b^2*d^2*e^3*m^3*x^2 - 3*(e*x + d)^m*A*b^3*d^2*e^3*m^3*x^2 + 30*(e*x + d)^m*B*a^2*b*d*e^4*m^3*x^2 + 30*(e*x + d)^m*A*a*b^2*d*e^4*m^3*x^2 + 13*(e*x + d)^m*B*a^3*e^5*m^3*x^2 + 39*(e*x + d)^m*A*a^2*b*e^5*m^3*x^2 - 12*(e*x + d)^m*B*b^3*d^2*e^3*m^2*x^3 + 51*(e*x + d)^m*B*a*b^2*d*e^4*m^2*x^3 + 17*(e*x + d)^m*A*b^3*d*e^4*m^2*x^3 + 147*(e*x + d)^m*B*a^2*b*e^5*m^2*x^3 + 147*(e*x + d)^m*A*a*b^2*e^5*m^2*x^3 + 6*(e*x + d)^m*B*b^3*d*e^4*m*x...
```

Mupad [B] (verification not implemented)

Time = 1.50 (sec) , antiderivative size = 1273, normalized size of antiderivative = 6.84

$$\int (a + bx)^3(A + Bx)(d + ex)^m dx = \text{Too large to display}$$

input `int((A + B*x)*(a + b*x)^3*(d + e*x)^m,x)`

output

```
((d + e*x)^m*(24*B*b^3*d^5 + 120*A*a^3*d*e^4 - 30*A*b^3*d^4*e - 60*B*a^3*d^2*e^3 + 120*A*a*b^2*d^3*e^2 - 180*A*a^2*b*d^2*e^3 + 120*B*a^2*b*d^3*e^2 + 71*A*a^3*d*e^4*m^2 + 14*A*a^3*d*e^4*m^3 + A*a^3*d*e^4*m^4 - 47*B*a^3*d^2*e^3*m - 12*B*a^3*d^2*e^3*m^2 - B*a^3*d^2*e^3*m^3 - 90*B*a*b^2*d^4*e + 154*A*a^3*d*e^4*m - 6*A*b^3*d^4*e*m + 6*A*a*b^2*d^3*e^2*m^2 - 36*A*a^2*b*d^2*e^3*m^2 - 3*A*a^2*b*d^2*e^3*m^3 + 6*B*a^2*b*d^3*e^2*m^2 - 18*B*a*b^2*d^4*e*m + 54*A*a*b^2*d^3*e^2*m - 141*A*a^2*b*d^2*e^3*m + 54*B*a^2*b*d^3*e^2*m))/
(e^5*(274*m + 225*m^2 + 85*m^3 + 15*m^4 + m^5 + 120)) + (x*(d + e*x)^m*(120*A*a^3*e^5 + 154*A*a^3*e^5*m + 71*A*a^3*e^5*m^2 + 14*A*a^3*e^5*m^3 + A*a^3*e^5*m^4 + 30*A*b^3*d^3*e^2*m + 47*B*a^3*d*e^4*m^2 + 12*B*a^3*d*e^4*m^3 + B*a^3*d*e^4*m^4 + 6*A*b^3*d^3*e^2*m^2 + 60*B*a^3*d*e^4*m - 24*B*b^3*d^4*e*m - 54*A*a*b^2*d^2*e^3*m^2 - 6*A*a*b^2*d^2*e^3*m^3 + 18*B*a*b^2*d^3*e^2*m^2 - 54*B*a^2*b*d^2*e^3*m^2 - 6*B*a^2*b*d^2*e^3*m^3 + 180*A*a^2*b*d*e^4*m - 120*A*a*b^2*d^2*e^3*m + 141*A*a^2*b*d*e^4*m^2 + 36*A*a^2*b*d*e^4*m^3 + 3*A*a^2*b*d*e^4*m^4 + 90*B*a*b^2*d^3*e^2*m - 120*B*a^2*b*d^2*e^3*m))/
(e^5*(274*m + 225*m^2 + 85*m^3 + 15*m^4 + m^5 + 120)) + (x^2*(m + 1)*(d + e*x)^m*(60*B*a^3*e^3 + 180*A*a^2*b*e^3 + 47*B*a^3*e^3*m + 12*B*b^3*d^3*m + 12*B*a^3*e^3*m^2 + B*a^3*e^3*m^3 + 36*A*a^2*b*e^3*m^2 + 3*A*a^2*b*e^3*m^3 - 3*A*b^3*d^2*e*m^2 + 141*A*a^2*b*e^3*m - 15*A*b^3*d^2*e*m + 60*A*a*b^2*d*e^2*m - 45*B*a*b^2*d^2*e*m + 60*B*a^2*b*d*e^2*m + 27*A*a*b^2*d*e^2*m^2 + 3*A...
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 1028, normalized size of antiderivative = 5.53

$$\int (a + bx)^3(A + Bx)(d + ex)^m dx = \text{Too large to display}$$

input `int((b*x+a)^3*(B*x+A)*(e*x+d)^m,x)`

output

```

((d + e*x)**m*(a**4*d***4*m**4 + 14*a**4*d***4*m**3 + 71*a**4*d***4*m**
2 + 154*a**4*d***4*m + 120*a**4*d***4 + a**4*e**5*m**4*x + 14*a**4*e**5*
m**3*x + 71*a**4*e**5*m**2*x + 154*a**4*e**5*m*x + 120*a**4*e**5*x - 4*a**
3*b*d**2*e**3*m**3 - 48*a**3*b*d**2*e**3*m**2 - 188*a**3*b*d**2*e**3*m - 2
40*a**3*b*d**2*e**3 + 4*a**3*b*d***4*m**4*x + 48*a**3*b*d***4*m**3*x + 1
88*a**3*b*d***4*m**2*x + 240*a**3*b*d***4*m*x + 4*a**3*b*e**5*m**4*x**2
+ 52*a**3*b*e**5*m**3*x**2 + 236*a**3*b*e**5*m**2*x**2 + 428*a**3*b*e**5*m
*x**2 + 240*a**3*b*e**5*x**2 + 12*a**2*b**2*d**3*e**2*m**2 + 108*a**2*b**2
*d**3*e**2*m + 240*a**2*b**2*d**3*e**2 - 12*a**2*b**2*d**2*e**3*m**3*x - 1
08*a**2*b**2*d**2*e**3*m**2*x - 240*a**2*b**2*d**2*e**3*m*x + 6*a**2*b**2*
d***4*m**4*x**2 + 60*a**2*b**2*d***4*m**3*x**2 + 174*a**2*b**2*d***4*m**
*2*x**2 + 120*a**2*b**2*d***4*m*x**2 + 6*a**2*b**2*e**5*m**4*x**3 + 72*a*
*2*b**2*e**5*m**3*x**3 + 294*a**2*b**2*e**5*m**2*x**3 + 468*a**2*b**2*e**5
*m*x**3 + 240*a**2*b**2*e**5*x**3 - 24*a*b**3*d**4*e*m - 120*a*b**3*d**4*e
+ 24*a*b**3*d**3*e**2*m**2*x + 120*a*b**3*d**3*e**2*m*x - 12*a*b**3*d**2*
e**3*m**3*x**2 - 72*a*b**3*d**2*e**3*m**2*x**2 - 60*a*b**3*d**2*e**3*m*x**
2 + 4*a*b**3*d***4*m**4*x**3 + 32*a*b**3*d***4*m**3*x**3 + 68*a*b**3*d*
**4*m**2*x**3 + 40*a*b**3*d***4*m*x**3 + 4*a*b**3*e**5*m**4*x**4 + 44*a*b
**3*e**5*m**3*x**4 + 164*a*b**3*e**5*m**2*x**4 + 244*a*b**3*e**5*m*x**4 +
120*a*b**3*e**5*x**4 + 24*b**4*d**5 - 24*b**4*d**4*e*m*x + 12*b**4*d**3...

```

3.231 $\int (a + bx)^2 (A + Bx)(d + ex)^m dx$

Optimal result	2080
Mathematica [A] (verified)	2081
Rubi [A] (verified)	2081
Maple [B] (verified)	2082
Fricas [B] (verification not implemented)	2083
Sympy [B] (verification not implemented)	2084
Maxima [B] (verification not implemented)	2085
Giac [B] (verification not implemented)	2086
Mupad [B] (verification not implemented)	2087
Reduce [B] (verification not implemented)	2088

Optimal result

Integrand size = 20, antiderivative size = 138

$$\int (a + bx)^2 (A + Bx)(d + ex)^m dx = -\frac{(bd - ae)^2 (Bd - Ae)(d + ex)^{1+m}}{e^4(1 + m)} + \frac{(bd - ae)(3bBd - 2Abe - aBe)(d + ex)^{2+m}}{e^4(2 + m)} - \frac{b(3bBd - Abe - 2aBe)(d + ex)^{3+m}}{e^4(3 + m)} + \frac{b^2 B(d + ex)^{4+m}}{e^4(4 + m)}$$

output

```
-(-a*e+b*d)^2*(-A*e+B*d)*(e*x+d)^(1+m)/e^4/(1+m)+(-a*e+b*d)*(-2*A*b*e-B*a*
e+3*B*b*d)*(e*x+d)^(2+m)/e^4/(2+m)-b*(-A*b*e-2*B*a*e+3*B*b*d)*(e*x+d)^(3+m
)/e^4/(3+m)+b^2*B*(e*x+d)^(4+m)/e^4/(4+m)
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.88

$$\int (a + bx)^2 (A + Bx)(d + ex)^m dx$$

$$= \frac{(d + ex)^{1+m} \left(-\frac{(bd-ae)^2(Bd-Ae)}{1+m} + \frac{(bd-ae)(3bBd-2Abe-aBe)(d+ex)}{2+m} - \frac{b(3bBd-Abe-2aBe)(d+ex)^2}{3+m} + \frac{b^2B(d+ex)^3}{4+m} \right)}{e^4}$$

input

```
Integrate[(a + b*x)^2*(A + B*x)*(d + e*x)^m,x]
```

output

```
((d + e*x)^(1 + m)*(-(((b*d - a*e)^2*(B*d - A*e))/(1 + m)) + ((b*d - a*e)*(3*b*B*d - 2*A*b*e - a*B*e)*(d + e*x))/(2 + m) - (b*(3*b*B*d - A*b*e - 2*a*B*e)*(d + e*x)^2)/(3 + m) + (b^2*B*(d + e*x)^3)/(4 + m)))/e^4
```

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^2 (A + Bx)(d + ex)^m dx$$

$$\downarrow 86$$

$$\int \left(\frac{(ae - bd)^2 (Ae - Bd)(d + ex)^m}{e^3} + \frac{(ae - bd)(d + ex)^{m+1} (aBe + 2Abe - 3bBd)}{e^3} + \frac{b(d + ex)^{m+2} (2aBe + Ab)}{e^3} \right)$$

$$\downarrow 2009$$

$$-\frac{(bd - ae)^2 (Bd - Ae)(d + ex)^{m+1}}{e^4(m + 1)} + \frac{(bd - ae)(d + ex)^{m+2} (-aBe - 2Abe + 3bBd)}{e^4(m + 2)} - \frac{b(d + ex)^{m+3} (-2aBe - Abe + 3bBd)}{e^4(m + 3)} + \frac{b^2 B(d + ex)^{m+4}}{e^4(m + 4)}$$

output

```
(A*a^2*d*e^3*m^3 - 6*B*b^2*d^4 + 24*A*a^2*d*e^3 + 8*(2*B*a*b + A*b^2)*d^3*
e - 12*(B*a^2 + 2*A*a*b)*d^2*e^2 + (B*b^2*e^4*m^3 + 6*B*b^2*e^4*m^2 + 11*B
*b^2*e^4*m + 6*B*b^2*e^4)*x^4 + (8*(2*B*a*b + A*b^2)*e^4 + (B*b^2*d*e^3 +
(2*B*a*b + A*b^2)*e^4)*m^3 + (3*B*b^2*d*e^3 + 7*(2*B*a*b + A*b^2)*e^4)*m^2
+ 2*(B*b^2*d*e^3 + 7*(2*B*a*b + A*b^2)*e^4)*m)*x^3 + (9*A*a^2*d*e^3 - (B*
a^2 + 2*A*a*b)*d^2*e^2)*m^2 + (12*(B*a^2 + 2*A*a*b)*e^4 + ((2*B*a*b + A*b^
2)*d*e^3 + (B*a^2 + 2*A*a*b)*e^4)*m^3 - (3*B*b^2*d^2*e^2 - 5*(2*B*a*b + A*
b^2)*d*e^3 - 8*(B*a^2 + 2*A*a*b)*e^4)*m^2 - (3*B*b^2*d^2*e^2 - 4*(2*B*a*b
+ A*b^2)*d*e^3 - 19*(B*a^2 + 2*A*a*b)*e^4)*m)*x^2 + (26*A*a^2*d*e^3 + 2*(2
*B*a*b + A*b^2)*d^3*e - 7*(B*a^2 + 2*A*a*b)*d^2*e^2)*m + (24*A*a^2*e^4 + (
A*a^2*e^4 + (B*a^2 + 2*A*a*b)*d*e^3)*m^3 + (9*A*a^2*e^4 - 2*(2*B*a*b + A*b
^2)*d^2*e^2 + 7*(B*a^2 + 2*A*a*b)*d*e^3)*m^2 + 2*(3*B*b^2*d^3*e + 13*A*a^2
*e^4 - 4*(2*B*a*b + A*b^2)*d^2*e^2 + 6*(B*a^2 + 2*A*a*b)*d*e^3)*m)*x)*(e*x
+ d)^m/(e^4*m^4 + 10*e^4*m^3 + 35*e^4*m^2 + 50*e^4*m + 24*e^4)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6186 vs. $2(126) = 252$.

Time = 1.37 (sec) , antiderivative size = 6186, normalized size of antiderivative = 44.83

$$\int (a + bx)^2 (A + Bx)(d + ex)^m dx = \text{Too large to display}$$

input

```
integrate((b*x+a)**2*(B*x+A)*(e*x+d)**m,x)
```

output

```
Piecewise((d**m*(A*a**2*x + A*a*b*x**2 + A*b**2*x**3/3 + B*a**2*x**2/2 + 2
*B*a*b*x**3/3 + B*b**2*x**4/4), Eq(e, 0)), (-2*A*a**2*e**3/(6*d**3*e**4 +
18*d**2*e**5*x + 18*d*e**6*x**2 + 6*e**7*x**3) - 2*A*a*b*d*e**2/(6*d**3*e
**4 + 18*d**2*e**5*x + 18*d*e**6*x**2 + 6*e**7*x**3) - 6*A*a*b*e**3*x/(6*d
**3*e**4 + 18*d**2*e**5*x + 18*d*e**6*x**2 + 6*e**7*x**3) - 2*A*b**2*d**2*e
/(6*d**3*e**4 + 18*d**2*e**5*x + 18*d*e**6*x**2 + 6*e**7*x**3) - 6*A*b**2*
d*e**2*x/(6*d**3*e**4 + 18*d**2*e**5*x + 18*d*e**6*x**2 + 6*e**7*x**3) - 6
*A*b**2*e**3*x**2/(6*d**3*e**4 + 18*d**2*e**5*x + 18*d*e**6*x**2 + 6*e**7*
x**3) - B*a**2*d*e**2/(6*d**3*e**4 + 18*d**2*e**5*x + 18*d*e**6*x**2 + 6*e
**7*x**3) - 3*B*a**2*e**3*x/(6*d**3*e**4 + 18*d**2*e**5*x + 18*d*e**6*x**2
+ 6*e**7*x**3) - 4*B*a*b*d**2*e/(6*d**3*e**4 + 18*d**2*e**5*x + 18*d*e**6
*x**2 + 6*e**7*x**3) - 12*B*a*b*d*e**2*x/(6*d**3*e**4 + 18*d**2*e**5*x + 1
8*d*e**6*x**2 + 6*e**7*x**3) - 12*B*a*b*e**3*x**2/(6*d**3*e**4 + 18*d**2*e
**5*x + 18*d*e**6*x**2 + 6*e**7*x**3) + 6*B*b**2*d**3*log(d/e + x)/(6*d**3
*e**4 + 18*d**2*e**5*x + 18*d*e**6*x**2 + 6*e**7*x**3) + 11*B*b**2*d**3/(6
*d**3*e**4 + 18*d**2*e**5*x + 18*d*e**6*x**2 + 6*e**7*x**3) + 18*B*b**2*d*
**2*e*x*log(d/e + x)/(6*d**3*e**4 + 18*d**2*e**5*x + 18*d*e**6*x**2 + 6*e
**7*x**3) + 27*B*b**2*d**2*e*x/(6*d**3*e**4 + 18*d**2*e**5*x + 18*d*e**6*x
**2 + 6*e**7*x**3) + 18*B*b**2*d*e**2*x**2*log(d/e + x)/(6*d**3*e**4 + 18*d
**2*e**5*x + 18*d*e**6*x**2 + 6*e**7*x**3) + 18*B*b**2*d*e**2*x**2/(6*d...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 364 vs. $2(138) = 276$.

Time = 0.05 (sec) , antiderivative size = 364, normalized size of antiderivative = 2.64

$$\int (a + bx)^2(A + Bx)(d + ex)^m dx = \frac{(e^2(m+1)x^2 + demx - d^2)(ex + d)^m Ba^2}{(m^2 + 3m + 2)e^2} + \frac{2(e^2(m+1)x^2 + demx - d^2)(ex + d)^m Aab}{(m^2 + 3m + 2)e^2} + \frac{(ex + d)^{m+1} Aa^2}{e(m+1)} + \frac{2((m^2 + 3m + 2)e^3x^3 + (m^2 + m)de^2x^2 - 2d^2emx + 2d^3)(ex + d)^m Bab}{(m^3 + 6m^2 + 11m + 6)e^3} + \frac{((m^2 + 3m + 2)e^3x^3 + (m^2 + m)de^2x^2 - 2d^2emx + 2d^3)(ex + d)^m Ab^2}{(m^3 + 6m^2 + 11m + 6)e^3} + \frac{((m^3 + 6m^2 + 11m + 6)e^4x^4 + (m^3 + 3m^2 + 2m)de^3x^3 - 3(m^2 + m)d^2e^2x^2 + 6d^3emx - 6d^4)(ex + d)^m}{(m^4 + 10m^3 + 35m^2 + 50m + 24)e^4}$$

input

```
integrate((b*x+a)^2*(B*x+A)*(e*x+d)^m,x, algorithm="maxima")
```

output

```
(e^2*(m + 1)*x^2 + d*e*m*x - d^2)*(e*x + d)^m*B*a^2/((m^2 + 3*m + 2)*e^2)
+ 2*(e^2*(m + 1)*x^2 + d*e*m*x - d^2)*(e*x + d)^m*A*a*b/((m^2 + 3*m + 2)*e
^2) + (e*x + d)^(m + 1)*A*a^2/(e*(m + 1)) + 2*((m^2 + 3*m + 2)*e^3*x^3 + (
m^2 + m)*d*e^2*x^2 - 2*d^2*e*m*x + 2*d^3)*(e*x + d)^m*B*a*b/((m^3 + 6*m^2
+ 11*m + 6)*e^3) + ((m^2 + 3*m + 2)*e^3*x^3 + (m^2 + m)*d*e^2*x^2 - 2*d^2*
e*m*x + 2*d^3)*(e*x + d)^m*A*b^2/((m^3 + 6*m^2 + 11*m + 6)*e^3) + ((m^3 +
6*m^2 + 11*m + 6)*e^4*x^4 + (m^3 + 3*m^2 + 2*m)*d*e^3*x^3 - 3*(m^2 + m)*d^
2*e^2*x^2 + 6*d^3*e*m*x - 6*d^4)*(e*x + d)^m*B*b^2/((m^4 + 10*m^3 + 35*m^2
+ 50*m + 24)*e^4)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1261 vs. $2(138) = 276$.

Time = 0.13 (sec) , antiderivative size = 1261, normalized size of antiderivative = 9.14

$$\int (a + bx)^2 (A + Bx)(d + ex)^m dx = \text{Too large to display}$$

input

```
integrate((b*x+a)^2*(B*x+A)*(e*x+d)^m,x, algorithm="giac")
```


output

```

((d + e*x)^m*(24*A*a^2*d*e^3 - 6*B*b^2*d^4 + 8*A*b^2*d^3*e - 12*B*a^2*d^2*
e^2 + 9*A*a^2*d*e^3*m^2 + A*a^2*d*e^3*m^3 - 7*B*a^2*d^2*e^2*m + 16*B*a*b*d
^3*e - B*a^2*d^2*e^2*m^2 - 24*A*a*b*d^2*e^2 + 26*A*a^2*d*e^3*m + 2*A*b^2*d
^3*e*m - 14*A*a*b*d^2*e^2*m - 2*A*a*b*d^2*e^2*m^2 + 4*B*a*b*d^3*e*m))/(e^4
*(50*m + 35*m^2 + 10*m^3 + m^4 + 24)) + (x*(d + e*x)^m*(24*A*a^2*e^4 + 26*
A*a^2*e^4*m + 9*A*a^2*e^4*m^2 + A*a^2*e^4*m^3 - 8*A*b^2*d^2*e^2*m + 7*B*a^
2*d*e^3*m^2 + B*a^2*d*e^3*m^3 - 2*A*b^2*d^2*e^2*m^2 + 12*B*a^2*d*e^3*m + 6
*B*b^2*d^3*e*m + 14*A*a*b*d*e^3*m^2 + 2*A*a*b*d*e^3*m^3 - 16*B*a*b*d^2*e^2
*m - 4*B*a*b*d^2*e^2*m^2 + 24*A*a*b*d*e^3*m))/(e^4*(50*m + 35*m^2 + 10*m^3
+ m^4 + 24)) + (x^2*(m + 1)*(d + e*x)^m*(12*B*a^2*e^2 + 24*A*a*b*e^2 + 7*
B*a^2*e^2*m - 3*B*b^2*d^2*m + B*a^2*e^2*m^2 + 14*A*a*b*e^2*m + 4*A*b^2*d*e
*m + 2*A*a*b*e^2*m^2 + A*b^2*d*e*m^2 + 8*B*a*b*d*e*m + 2*B*a*b*d*e*m^2))/(
e^2*(50*m + 35*m^2 + 10*m^3 + m^4 + 24)) + (B*b^2*x^4*(d + e*x)^m*(11*m +
6*m^2 + m^3 + 6))/(50*m + 35*m^2 + 10*m^3 + m^4 + 24) + (b*x^3*(d + e*x)^m
*(3*m + m^2 + 2)*(4*A*b*e + 8*B*a*e + A*b*e*m + 2*B*a*e*m + B*b*d*m))/(e*(
50*m + 35*m^2 + 10*m^3 + m^4 + 24))

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 546, normalized size of antiderivative = 3.96

$$\int (a + bx)^2 (A + Bx)(d + ex)^m dx$$

$$= \frac{(ex + d)^m (b^3 e^4 m^3 x^4 + 3a b^2 e^4 m^3 x^3 + b^3 d e^3 m^3 x^3 + 6b^3 e^4 m^2 x^4 + 3a^2 b e^4 m^3 x^2 + 3a b^2 d e^3 m^3 x^2 + 21a b^2 d e^3 m^3 x^2 + 21a b^2 d e^3 m^3 x^2 + 21a b^2 d e^3 m^3 x^2)}{(50m + 35m^2 + 10m^3 + m^4 + 24)}$$

input

```
int((b*x+a)^2*(B*x+A)*(e*x+d)^m,x)
```

output

```

((d + e*x)**m*(a**3*d*e**3*m**3 + 9*a**3*d*e**3*m**2 + 26*a**3*d*e**3*m +
24*a**3*d*e**3 + a**3*e**4*m**3*x + 9*a**3*e**4*m**2*x + 26*a**3*e**4*m*x
+ 24*a**3*e**4*x - 3*a**2*b*d**2*e**2*m**2 - 21*a**2*b*d**2*e**2*m - 36*a*
**2*b*d**2*e**2 + 3*a**2*b*d*e**3*m**3*x + 21*a**2*b*d*e**3*m**2*x + 36*a**
2*b*d*e**3*m*x + 3*a**2*b*e**4*m**3*x**2 + 24*a**2*b*e**4*m**2*x**2 + 57*a
**2*b*e**4*m*x**2 + 36*a**2*b*e**4*x**2 + 6*a*b**2*d**3*e*m + 24*a*b**2*d*
*3*e - 6*a*b**2*d**2*e**2*m**2*x - 24*a*b**2*d**2*e**2*m*x + 3*a*b**2*d*e*
*3*m**3*x**2 + 15*a*b**2*d*e**3*m**2*x**2 + 12*a*b**2*d*e**3*m*x**2 + 3*a*
b**2*e**4*m**3*x**3 + 21*a*b**2*e**4*m**2*x**3 + 42*a*b**2*e**4*m*x**3 + 2
4*a*b**2*e**4*x**3 - 6*b**3*d**4 + 6*b**3*d**3*e*m*x - 3*b**3*d**2*e**2*m*
*2*x**2 - 3*b**3*d**2*e**2*m*x**2 + b**3*d*e**3*m**3*x**3 + 3*b**3*d*e**3*
m**2*x**3 + 2*b**3*d*e**3*m*x**3 + b**3*e**4*m**3*x**4 + 6*b**3*e**4*m**2*
x**4 + 11*b**3*e**4*m*x**4 + 6*b**3*e**4*x**4))/(e**4*(m**4 + 10*m**3 + 35
*m**2 + 50*m + 24))

```

3.232 $\int (a + bx)(A + Bx)(d + ex)^m dx$

Optimal result	2090
Mathematica [A] (verified)	2090
Rubi [A] (verified)	2091
Maple [B] (verified)	2092
Fricas [B] (verification not implemented)	2093
Sympy [B] (verification not implemented)	2093
Maxima [A] (verification not implemented)	2094
Giac [B] (verification not implemented)	2095
Mupad [B] (verification not implemented)	2096
Reduce [B] (verification not implemented)	2096

Optimal result

Integrand size = 18, antiderivative size = 90

$$\int (a + bx)(A + Bx)(d + ex)^m dx = \frac{(bd - ae)(Bd - Ae)(d + ex)^{1+m}}{e^3(1 + m)} - \frac{(2bBd - Abe - aBe)(d + ex)^{2+m}}{e^3(2 + m)} + \frac{bB(d + ex)^{3+m}}{e^3(3 + m)}$$

output

$(-a*e+b*d)*(-A*e+B*d)*(e*x+d)^(1+m)/e^3/(1+m)-(-A*b*e-B*a*e+2*B*b*d)*(e*x+d)^(2+m)/e^3/(2+m)+b*B*(e*x+d)^(3+m)/e^3/(3+m)$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.88

$$\int (a + bx)(A + Bx)(d + ex)^m dx = \frac{(d + ex)^{1+m} \left(\frac{(bd - ae)(Bd - Ae)}{1 + m} - \frac{(2bBd - Abe - aBe)(d + ex)}{2 + m} + \frac{bB(d + ex)^2}{3 + m} \right)}{e^3}$$

input

`Integrate[(a + b*x)*(A + B*x)*(d + e*x)^m,x]`

output

$$\frac{((d + ex)^{(1 + m)} * ((b*d - a*e)*(B*d - A*e))/(1 + m) - ((2*b*B*d - A*b*e - a*B*e)*(d + ex))/(2 + m) + (b*B*(d + ex)^2)/(3 + m))}{e^3}$$

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)(A + Bx)(d + ex)^m dx$$

↓ 86

$$\int \left(\frac{(ae - bd)(Ae - Bd)(d + ex)^m}{e^2} + \frac{(d + ex)^{m+1}(aBe + Abe - 2bBd)}{e^2} + \frac{bB(d + ex)^{m+2}}{e^2} \right) dx$$

↓ 2009

$$\frac{(bd - ae)(Bd - Ae)(d + ex)^{m+1}}{e^3(m + 1)} - \frac{(d + ex)^{m+2}(-aBe - Abe + 2bBd)}{e^3(m + 2)} + \frac{bB(d + ex)^{m+3}}{e^3(m + 3)}$$

input

$$\text{Int}[(a + b*x)*(A + B*x)*(d + e*x)^m, x]$$

output

$$\frac{((b*d - a*e)*(B*d - A*e)*(d + e*x)^{(1 + m)})}{(e^3*(1 + m))} - \frac{((2*b*B*d - A*b*e - a*B*e)*(d + e*x)^{(2 + m)})}{(e^3*(2 + m))} + \frac{(b*B*(d + e*x)^{(3 + m)})}{(e^3*(3 + m))}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 256 vs. $2(90) = 180$.

Time = 0.09 (sec) , antiderivative size = 256, normalized size of antiderivative = 2.84

$$\int (a + bx)(A + Bx)(d + ex)^m dx$$

$$= \frac{(Aade^2m^2 + 2Bbd^3 + 6Aade^2 - 3(Ba + Ab)d^2e + (Bbe^3m^2 + 3Bbe^3m + 2Bbe^3)x^3 + (3(Ba + Ab)e^3$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)^m,x, algorithm="fricas")`

output $(A*a*d*e^2*m^2 + 2*B*b*d^3 + 6*A*a*d*e^2 - 3*(B*a + A*b)*d^2*e + (B*b*e^3*m^2 + 3*B*b*e^3*m + 2*B*b*e^3)*x^3 + (3*(B*a + A*b)*e^3 + (B*b*d*e^2 + (B*a + A*b)*e^3)*m^2 + (B*b*d*e^2 + 4*(B*a + A*b)*e^3)*m)*x^2 + (5*A*a*d*e^2 - (B*a + A*b)*d^2*e)*m + (6*A*a*e^3 + (A*a*e^3 + (B*a + A*b)*d*e^2)*m^2 - (2*B*b*d^2*e - 5*A*a*e^3 - 3*(B*a + A*b)*d*e^2)*m)*x)*(e*x + d)^m/(e^3*m^3 + 6*e^3*m^2 + 11*e^3*m + 6*e^3)$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1982 vs. $2(78) = 156$.

Time = 0.65 (sec) , antiderivative size = 1982, normalized size of antiderivative = 22.02

$$\int (a + bx)(A + Bx)(d + ex)^m dx = \text{Too large to display}$$

input `integrate((b*x+a)*(B*x+A)*(e*x+d)**m,x)`

output

```
Piecewise((d**m*(A*a*x + A*b*x**2/2 + B*a*x**2/2 + B*b*x**3/3), Eq(e, 0)),
(-A*a*e**2/(2*d**2*e**3 + 4*d*e**4*x + 2*e**5*x**2) - A*b*d*e/(2*d**2*e**
3 + 4*d*e**4*x + 2*e**5*x**2) - 2*A*b*e**2*x/(2*d**2*e**3 + 4*d*e**4*x + 2
*e**5*x**2) - B*a*d*e/(2*d**2*e**3 + 4*d*e**4*x + 2*e**5*x**2) - 2*B*a*e**
2*x/(2*d**2*e**3 + 4*d*e**4*x + 2*e**5*x**2) + 2*B*b*d**2*log(d/e + x)/(2*
d**2*e**3 + 4*d*e**4*x + 2*e**5*x**2) + 3*B*b*d**2/(2*d**2*e**3 + 4*d*e**4
*x + 2*e**5*x**2) + 4*B*b*d*e*x*log(d/e + x)/(2*d**2*e**3 + 4*d*e**4*x + 2
*e**5*x**2) + 4*B*b*d*e*x/(2*d**2*e**3 + 4*d*e**4*x + 2*e**5*x**2) + 2*B*b
*e**2*x**2*log(d/e + x)/(2*d**2*e**3 + 4*d*e**4*x + 2*e**5*x**2), Eq(m, -3
)), (-A*a*e**2/(d*e**3 + e**4*x) + A*b*d*e*log(d/e + x)/(d*e**3 + e**4*x)
+ A*b*d*e/(d*e**3 + e**4*x) + A*b*e**2*x*log(d/e + x)/(d*e**3 + e**4*x) +
B*a*d*e*log(d/e + x)/(d*e**3 + e**4*x) + B*a*d*e/(d*e**3 + e**4*x) + B*a*e
**2*x*log(d/e + x)/(d*e**3 + e**4*x) - 2*B*b*d**2*log(d/e + x)/(d*e**3 + e
**4*x) - 2*B*b*d**2/(d*e**3 + e**4*x) - 2*B*b*d*e*x*log(d/e + x)/(d*e**3 +
e**4*x) + B*b*e**2*x**2/(d*e**3 + e**4*x), Eq(m, -2)), (A*a*log(d/e + x)/
e - A*b*d*log(d/e + x)/e**2 + A*b*x/e - B*a*d*log(d/e + x)/e**2 + B*a*x/e
+ B*b*d**2*log(d/e + x)/e**3 - B*b*d*x/e**2 + B*b*x**2/(2*e), Eq(m, -1)),
(A*a*d*e**2*m**2*(d + e*x)**m/(e**3*m**3 + 6*e**3*m**2 + 11*e**3*m + 6*e**
3) + 5*A*a*d*e**2*m*(d + e*x)**m/(e**3*m**3 + 6*e**3*m**2 + 11*e**3*m + 6*
e**3) + 6*A*a*d*e**2*(d + e*x)**m/(e**3*m**3 + 6*e**3*m**2 + 11*e**3*m ...
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.99

$$\int (a + bx)(A + Bx)(d + ex)^m dx$$

$$= \frac{(e^2(m+1)x^2 + demx - d^2)(ex + d)^m Ba}{(m^2 + 3m + 2)e^2}$$

$$+ \frac{(e^2(m+1)x^2 + demx - d^2)(ex + d)^m Ab}{(m^2 + 3m + 2)e^2} + \frac{(ex + d)^{m+1} Aa}{e(m+1)}$$

$$+ \frac{((m^2 + 3m + 2)e^3x^3 + (m^2 + m)de^2x^2 - 2d^2emx + 2d^3)(ex + d)^m Bb}{(m^3 + 6m^2 + 11m + 6)e^3}$$

input

```
integrate((b*x+a)*(B*x+A)*(e*x+d)^m,x, algorithm="maxima")
```


Mupad [B] (verification not implemented)

Time = 1.21 (sec) , antiderivative size = 259, normalized size of antiderivative = 2.88

$$\int (a + bx)(A + Bx)(d + ex)^m dx$$

$$= (d + ex)^m \left(\frac{x(6Aae^3 + 5Aae^3m + Aae^3m^2 + 3Abde^2m + 3Bade^2m - 2Bbd^2em + Abde^2m)}{e^3(m^3 + 6m^2 + 11m + 6)} \right.$$

$$+ \frac{d(6Aae^2 + 2Bbd^2 + 5Aae^2m + Aae^2m^2 - 3Abde - 3Bade - Abdem - Badem)}{e^3(m^3 + 6m^2 + 11m + 6)}$$

$$+ \frac{Bbx^3(m^2 + 3m + 2)}{m^3 + 6m^2 + 11m + 6}$$

$$\left. + \frac{x^2(m + 1)(3Abe + 3Bae + Abem + Baem + Bbdm)}{e(m^3 + 6m^2 + 11m + 6)} \right)$$

input `int((A + B*x)*(a + b*x)*(d + e*x)^m,x)`output `(d + e*x)^m*((x*(6*A*a*e^3 + 5*A*a*e^3*m + A*a*e^3*m^2 + 3*A*b*d*e^2*m + 3*B*a*d*e^2*m - 2*B*b*d^2*e*m + A*b*d*e^2*m^2 + B*a*d*e^2*m^2))/(e^3*(11*m + 6*m^2 + m^3 + 6)) + (d*(6*A*a*e^2 + 2*B*b*d^2 + 5*A*a*e^2*m + A*a*e^2*m^2 - 3*A*b*d*e - 3*B*a*d*e - A*b*d*e*m - B*a*d*e*m))/(e^3*(11*m + 6*m^2 + m^3 + 6)) + (B*b*x^3*(3*m + m^2 + 2))/(11*m + 6*m^2 + m^3 + 6) + (x^2*(m + 1)*(3*A*b*e + 3*B*a*e + A*b*e*m + B*a*e*m + B*b*d*m))/(e*(11*m + 6*m^2 + m^3 + 6)))`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 241, normalized size of antiderivative = 2.68

$$\int (a + bx)(A + Bx)(d + ex)^m dx$$

$$= \frac{(ex + d)^m (b^2e^3m^2x^3 + 2abe^3m^2x^2 + b^2de^2m^2x^2 + 3b^2e^3mx^3 + a^2e^3m^2x + 2abde^2m^2x + 8abe^3mx^2 +$$

input `int((b*x+a)*(B*x+A)*(e*x+d)^m,x)`

output

```
((d + e*x)**m*(a**2*d*e**2*m**2 + 5*a**2*d*e**2*m + 6*a**2*d*e**2 + a**2*e
**3*m**2*x + 5*a**2*e**3*m*x + 6*a**2*e**3*x - 2*a*b*d**2*e*m - 6*a*b*d**2
*e + 2*a*b*d*e**2*m**2*x + 6*a*b*d*e**2*m*x + 2*a*b*e**3*m**2*x**2 + 8*a*b
*e**3*m*x**2 + 6*a*b*e**3*x**2 + 2*b**2*d**3 - 2*b**2*d**2*e*m*x + b**2*d*
e**2*m**2*x**2 + b**2*d*e**2*m*x**2 + b**2*e**3*m**2*x**3 + 3*b**2*e**3*m*
x**3 + 2*b**2*e**3*x**3))/(e**3*(m**3 + 6*m**2 + 11*m + 6))
```

3.233 $\int (A + Bx)(d + ex)^m dx$

Optimal result	2098
Mathematica [A] (verified)	2098
Rubi [A] (verified)	2099
Maple [A] (verified)	2100
Fricas [A] (verification not implemented)	2100
Sympy [B] (verification not implemented)	2101
Maxima [A] (verification not implemented)	2101
Giac [B] (verification not implemented)	2102
Mupad [B] (verification not implemented)	2102
Reduce [B] (verification not implemented)	2103

Optimal result

Integrand size = 13, antiderivative size = 47

$$\int (A + Bx)(d + ex)^m dx = -\frac{(Bd - Ae)(d + ex)^{1+m}}{e^2(1+m)} + \frac{B(d + ex)^{2+m}}{e^2(2+m)}$$

output

```
-(-A*e+B*d)*(e*x+d)^(1+m)/e^2/(1+m)+B*(e*x+d)^(2+m)/e^2/(2+m)
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.87

$$\int (A + Bx)(d + ex)^m dx = \frac{(d + ex)^{1+m}(-Bd + Ae(2 + m) + Be(1 + m)x)}{e^2(1 + m)(2 + m)}$$

input

```
Integrate[(A + B*x)*(d + e*x)^m,x]
```

output

```
((d + e*x)^(1 + m)*(-(B*d) + A*e*(2 + m) + B*e*(1 + m)*x))/(e^2*(1 + m)*(2 + m))
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (A + Bx)(d + ex)^m dx$$

$$\downarrow 53$$

$$\int \left(\frac{(Ae - Bd)(d + ex)^m}{e} + \frac{B(d + ex)^{m+1}}{e} \right) dx$$

$$\downarrow 2009$$

$$\frac{B(d + ex)^{m+2}}{e^2(m + 2)} - \frac{(Bd - Ae)(d + ex)^{m+1}}{e^2(m + 1)}$$

input `Int[(A + B*x)*(d + e*x)^m,x]`

output `-(((B*d - A*e)*(d + e*x)^(1 + m))/(e^2*(1 + m))) + (B*(d + e*x)^(2 + m))/(e^2*(2 + m))`

Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.98

method	result
gospers	$\frac{(ex+d)^{1+m}(Bemx+Aem+Bex+2Ae-Bd)}{e^2(m^2+3m+2)}$
orering	$\frac{(ex+d)^m(Bemx+Aem+Bex+2Ae-Bd)(ex+d)}{e^2(m^2+3m+2)}$
risch	$\frac{(x^2 B e^2 m + A e^2 m x + B d e m x + x^2 B e^2 + A d e m + 2 A e^2 x + 2 A d e - B d^2)(ex+d)^m}{e^2(2+m)(1+m)}$
norman	$\frac{B x^2 e^{m \ln(ex+d)}}{2+m} + \frac{d(Aem+2Ae-Bd)e^{m \ln(ex+d)}}{e^2(m^2+3m+2)} + \frac{(Aem+Bdm+2Ae)x e^{m \ln(ex+d)}}{e(m^2+3m+2)}$
parallelrisch	$\frac{B x^2 (ex+d)^m d e^2 m + A x (ex+d)^m d e^2 m + B x^2 (ex+d)^m d e^2 + B x (ex+d)^m d^2 e m + 2 A x (ex+d)^m d e^2 + A (ex+d)^m d^2 e m + 2 A (ex+d)^m d e^2}{(2+m)(1+m)e^2 d}$

input `int((B*x+A)*(e*x+d)^m,x,method=_RETURNVERBOSE)`output `1/e^2*(e*x+d)^(1+m)/(m^2+3*m+2)*(B*e*m*x+A*e*m+B*e*x+2*A*e-B*d)`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.77

$$\int (A + Bx)(d + ex)^m dx$$

$$= \frac{(Adem - Bd^2 + 2Ade + (Be^2m + Be^2)x^2 + (2Ae^2 + (Bde + Ae^2)m)x)(ex + d)^m}{e^2m^2 + 3e^2m + 2e^2}$$

input `integrate((B*x+A)*(e*x+d)^m,x, algorithm="fricas")`output `(A*d*e*m - B*d^2 + 2*A*d*e + (B*e^2*m + B*e^2)*x^2 + (2*A*e^2 + (B*d*e + A*e^2)*m)*x)*(e*x + d)^m/(e^2*m^2 + 3*e^2*m + 2*e^2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 377 vs. $2(37) = 74$.

Time = 0.34 (sec) , antiderivative size = 377, normalized size of antiderivative = 8.02

$$\int (A + Bx)(d + ex)^m dx$$

$$= \begin{cases} d^m \left(Ax + \frac{Bx^2}{2} \right) \\ -\frac{Ae}{de^2 + e^3x} + \frac{Bd \log\left(\frac{d}{e} + x\right)}{de^2 + e^3x} + \frac{Bd}{de^2 + e^3x} + \frac{Bex \log\left(\frac{d}{e} + x\right)}{de^2 + e^3x} \\ \frac{A \log\left(\frac{d}{e} + x\right)}{e} - \frac{Bd \log\left(\frac{d}{e} + x\right)}{e^2} + \frac{Bx}{e} \\ \frac{Adem(d+ex)^m}{e^2m^2+3e^2m+2e^2} + \frac{2Ade(d+ex)^m}{e^2m^2+3e^2m+2e^2} + \frac{Ae^2mx(d+ex)^m}{e^2m^2+3e^2m+2e^2} + \frac{2Ae^2x(d+ex)^m}{e^2m^2+3e^2m+2e^2} - \frac{Bd^2(d+ex)^m}{e^2m^2+3e^2m+2e^2} + \frac{Bdemx(d+ex)^m}{e^2m^2+3e^2m+2e^2} + \frac{Bex^2(d+ex)^m}{e^2m^2+3e^2m+2e^2} \end{cases}$$

input `integrate((B*x+A)*(e*x+d)**m,x)`

output

```
Piecewise((d**m*(A*x + B*x**2/2), Eq(e, 0)), (-A*e/(d*e**2 + e**3*x) + B*d
*log(d/e + x)/(d*e**2 + e**3*x) + B*d/(d*e**2 + e**3*x) + B*e*x*log(d/e +
x)/(d*e**2 + e**3*x), Eq(m, -2)), (A*log(d/e + x)/e - B*d*log(d/e + x)/e**
2 + B*x/e, Eq(m, -1)), (A*d*e*m*(d + e*x)**m/(e**2*m**2 + 3*e**2*m + 2*e**
2) + 2*A*d*e*(d + e*x)**m/(e**2*m**2 + 3*e**2*m + 2*e**2) + A*e**2*m*x*(d
+ e*x)**m/(e**2*m**2 + 3*e**2*m + 2*e**2) + 2*A*e**2*x*(d + e*x)**m/(e**2*
m**2 + 3*e**2*m + 2*e**2) - B*d**2*(d + e*x)**m/(e**2*m**2 + 3*e**2*m + 2*
e**2) + B*d*e*m*x*(d + e*x)**m/(e**2*m**2 + 3*e**2*m + 2*e**2) + B*e**2*m*
x**2*(d + e*x)**m/(e**2*m**2 + 3*e**2*m + 2*e**2) + B*e**2*x**2*(d + e*x)*
*m/(e**2*m**2 + 3*e**2*m + 2*e**2), True))
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.34

$$\int (A + Bx)(d + ex)^m dx = \frac{(e^2(m+1)x^2 + demx - d^2)(ex + d)^m B}{(m^2 + 3m + 2)e^2} + \frac{(ex + d)^{m+1} A}{e(m+1)}$$

input `integrate((B*x+A)*(e*x+d)^m,x, algorithm="maxima")`

output

```
(e^2*(m + 1)*x^2 + d*e*m*x - d^2)*(e*x + d)^m*B/((m^2 + 3*m + 2)*e^2) + (e*x + d)^(m + 1)*A/(e*(m + 1))
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 132 vs. $2(47) = 94$.

Time = 0.12 (sec) , antiderivative size = 132, normalized size of antiderivative = 2.81

$$\int (A + Bx)(d + ex)^m dx = \frac{(ex + d)^m B e^2 m x^2 + (ex + d)^m B d e m x + (ex + d)^m A e^2 m x + (ex + d)^m B e^2 x^2 + (ex + d)^m A d e m + 2(e^2 m^2 + 3 e^2 m + 2 e^2)}{e^2 m^2 + 3 e^2 m + 2 e^2}$$

input

```
integrate((B*x+A)*(e*x+d)^m,x, algorithm="giac")
```

output

```
((e*x + d)^m*B*e^2*m*x^2 + (e*x + d)^m*B*d*e*m*x + (e*x + d)^m*A*e^2*m*x + (e*x + d)^m*B*e^2*x^2 + (e*x + d)^m*A*d*e*m + 2*(e*x + d)^m*A*e^2*x - (e*x + d)^m*B*d^2 + 2*(e*x + d)^m*A*d*e)/(e^2*m^2 + 3*e^2*m + 2*e^2)
```

Mupad [B] (verification not implemented)

Time = 1.19 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.87

$$\int (A + Bx)(d + ex)^m dx = (d + ex)^m \left(\frac{x(2Ae^2 + Ae^2m + Bdem)}{e^2(m^2 + 3m + 2)} + \frac{Bx^2(m + 1)}{m^2 + 3m + 2} + \frac{d(2Ae - Bd + Aem)}{e^2(m^2 + 3m + 2)} \right)$$

input

```
int((A + B*x)*(d + e*x)^m,x)
```

output

```
(d + e*x)^m*((x*(2*A*e^2 + A*e^2*m + B*d*e*m))/(e^2*(3*m + m^2 + 2)) + (B*x^2*(m + 1))/(3*m + m^2 + 2) + (d*(2*A*e - B*d + A*e*m))/(e^2*(3*m + m^2 + 2)))
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.60

$$\int (A + Bx)(d + ex)^m dx$$

$$= \frac{(ex + d)^m (be^2mx^2 + ae^2mx + bdemx + be^2x^2 + adem + 2ae^2x + 2ade - bd^2)}{e^2(m^2 + 3m + 2)}$$

input `int((B*x+A)*(e*x+d)^m,x)`output `((d + e*x)**m*(a*d*e*m + 2*a*d*e + a*e**2*m*x + 2*a*e**2*x - b*d**2 + b*d*e*m*x + b*e**2*m*x**2 + b*e**2*x**2))/(e**2*(m**2 + 3*m + 2))`

3.234 $\int \frac{(A+Bx)(d+ex)^m}{a+bx} dx$

Optimal result	2104
Mathematica [A] (verified)	2104
Rubi [A] (verified)	2105
Maple [F]	2106
Fricas [F]	2106
Sympy [F]	2107
Maxima [F]	2107
Giac [F]	2107
Mupad [F(-1)]	2108
Reduce [B] (verification not implemented)	2108

Optimal result

Integrand size = 20, antiderivative size = 85

$$\int \frac{(A+Bx)(d+ex)^m}{a+bx} dx = \frac{B(d+ex)^{1+m}}{be(1+m)} - \frac{(Ab-aB)(d+ex)^{1+m} \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, \frac{b(d+ex)}{bd-ae}\right)}{b(bd-ae)(1+m)}$$

output

```
B*(e*x+d)^(1+m)/b/e/(1+m)-(A*b-B*a)*(e*x+d)^(1+m)*hypergeom([1, 1+m], [2+m], b*(e*x+d)/(-a*e+b*d))/b/(-a*e+b*d)/(1+m)
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.92

$$\int \frac{(A+Bx)(d+ex)^m}{a+bx} dx = \frac{(d+ex)^{1+m} \left(B(bd-ae) + (-Abe+aBe) \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, \frac{b(d+ex)}{bd-ae}\right) \right)}{be(bd-ae)(1+m)}$$

input `Integrate[((A + B*x)*(d + e*x)^m)/(a + b*x), x]`

output `((d + e*x)^(1 + m)*(B*(b*d - a*e) + (-A*b*e) + a*B*e)*Hypergeometric2F1[1, 1 + m, 2 + m, (b*(d + e*x))/(b*d - a*e)])/(b*e*(b*d - a*e)*(1 + m))`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {90, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)^m}{a + bx} dx$$

$$\downarrow 90$$

$$\frac{(Ab - aB) \int \frac{(d+ex)^m}{a+bx} dx}{b} + \frac{B(d + ex)^{m+1}}{be(m + 1)}$$

$$\downarrow 78$$

$$\frac{B(d + ex)^{m+1}}{be(m + 1)} - \frac{(Ab - aB)(d + ex)^{m+1} \text{Hypergeometric2F1}\left(1, m + 1, m + 2, \frac{b(d+ex)}{bd-ae}\right)}{b(m + 1)(bd - ae)}$$

input `Int[((A + B*x)*(d + e*x)^m)/(a + b*x), x]`

output `(B*(d + e*x)^(1 + m))/(b*e*(1 + m)) - ((A*b - a*B)*(d + e*x)^(1 + m)*Hypergeometric2F1[1, 1 + m, 2 + m, (b*(d + e*x))/(b*d - a*e)]/(b*(b*d - a*e)*(1 + m))`

Definitions of rubi rules used

rule 78 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 90 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`

Maple [F]

$$\int \frac{(Bx + A)(ex + d)^m}{bx + a} dx$$

input `int((B*x+A)*(e*x+d)^m/(b*x+a), x)`

output `int((B*x+A)*(e*x+d)^m/(b*x+a), x)`

Fricas [F]

$$\int \frac{(A + Bx)(d + ex)^m}{a + bx} dx = \int \frac{(Bx + A)(ex + d)^m}{bx + a} dx$$

input `integrate((B*x+A)*(e*x+d)^m/(b*x+a), x, algorithm="fricas")`

output `integral((B*x + A)*(e*x + d)^m/(b*x + a), x)`

Sympy [F]

$$\int \frac{(A + Bx)(d + ex)^m}{a + bx} dx = \int \frac{(A + Bx)(d + ex)^m}{a + bx} dx$$

input `integrate((B*x+A)*(e*x+d)**m/(b*x+a), x)`

output `Integral((A + B*x)*(d + e*x)**m/(a + b*x), x)`

Maxima [F]

$$\int \frac{(A + Bx)(d + ex)^m}{a + bx} dx = \int \frac{(Bx + A)(ex + d)^m}{bx + a} dx$$

input `integrate((B*x+A)*(e*x+d)^m/(b*x+a), x, algorithm="maxima")`

output `integrate((B*x + A)*(e*x + d)^m/(b*x + a), x)`

Giac [F]

$$\int \frac{(A + Bx)(d + ex)^m}{a + bx} dx = \int \frac{(Bx + A)(ex + d)^m}{bx + a} dx$$

input `integrate((B*x+A)*(e*x+d)^m/(b*x+a), x, algorithm="giac")`

output `integrate((B*x + A)*(e*x + d)^m/(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(A + Bx)(d + ex)^m}{a + bx} dx = \int \frac{(A + Bx) (d + ex)^m}{a + bx} dx$$

input `int(((A + B*x)*(d + e*x)^m)/(a + b*x), x)`output `int(((A + B*x)*(d + e*x)^m)/(a + b*x), x)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.25

$$\int \frac{(A + Bx)(d + ex)^m}{a + bx} dx = \frac{(ex + d)^m (ex + d)}{e(m + 1)}$$

input `int((B*x+A)*(e*x+d)^m/(b*x+a), x)`output `((d + e*x)**m*(d + e*x))/(e*(m + 1))`

3.235 $\int \frac{(A+Bx)(d+ex)^m}{(a+bx)^2} dx$

Optimal result	2109
Mathematica [A] (verified)	2109
Rubi [A] (verified)	2110
Maple [F]	2111
Fricas [F]	2111
Sympy [F]	2112
Maxima [F]	2112
Giac [F]	2112
Mupad [F(-1)]	2113
Reduce [F]	2113

Optimal result

Integrand size = 20, antiderivative size = 104

$$\int \frac{(A+Bx)(d+ex)^m}{(a+bx)^2} dx = \frac{B(d+ex)^{1+m}}{bem(a+bx)} - \frac{(aBe(1+m) - b(Bd+Aem))(d+ex)^{1+m} \text{Hypergeometric2F1}\left(2, 1+m, 2+m, \frac{b(d+ex)}{bd-ae}\right)}{b(bd-ae)^2 m(1+m)}$$

output

```
B*(e*x+d)^(1+m)/b/e/m/(b*x+a)-(a*B*e*(1+m)-b*(A*e*m+B*d))*(e*x+d)^(1+m)*hy
geom([2, 1+m], [2+m], b*(e*x+d)/(-a*e+b*d))/b/(-a*e+b*d)^2/m/(1+m)
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.95

$$\int \frac{(A+Bx)(d+ex)^m}{(a+bx)^2} dx = \frac{(d+ex)^{1+m} \left(-\frac{(Ab-aB)(bd-ae)}{a+bx} + \frac{(aBe(1+m)-b(Bd+Aem)) \text{Hypergeometric2F1}\left(1, 1+m, 2+m, \frac{b(d+ex)}{bd-ae}\right)}{1+m} \right)}{b(bd-ae)^2}$$

input

```
Integrate[((A + B*x)*(d + e*x)^m)/(a + b*x)^2,x]
```

output

$$\frac{((d + ex)^{(1 + m)} * (-(((A*b - a*B)*(b*d - a*e))/(a + b*x)) + ((a*B*e*(1 + m) - b*(B*d + A*e*m))*Hypergeometric2F1[1, 1 + m, 2 + m, (b*(d + e*x))/(b*d - a*e)])/(1 + m)))/(b*(b*d - a*e)^2}$$

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.08, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {87, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)^m}{(a + bx)^2} dx$$

$$\downarrow 87$$

$$\frac{(aBe(m + 1) - b(Aem + Bd)) \int \frac{(d+ex)^m}{a+bx} dx}{b(bd - ae)} - \frac{(Ab - aB)(d + ex)^{m+1}}{b(a + bx)(bd - ae)}$$

$$\downarrow 78$$

$$\frac{(d + ex)^{m+1}(aBe(m + 1) - b(Aem + Bd)) \text{Hypergeometric2F1}\left(1, m + 1, m + 2, \frac{b(d+ex)}{bd-ae}\right)}{\frac{b(m + 1)(bd - ae)^2}{(Ab - aB)(d + ex)^{m+1}} b(a + bx)(bd - ae)}$$

input

$$\text{Int}[(A + B*x)*(d + e*x)^m/(a + b*x)^2, x]$$

output

$$-(((A*b - a*B)*(d + e*x)^{(1 + m)})/(b*(b*d - a*e)*(a + b*x))) + ((a*B*e*(1 + m) - b*(B*d + A*e*m))*(d + e*x)^{(1 + m)}*Hypergeometric2F1[1, 1 + m, 2 + m, (b*(d + e*x))/(b*d - a*e)]/(b*(b*d - a*e)^2*(1 + m))$$

Definitions of rubi rules used

rule 78 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

Maple [F]

$$\int \frac{(Bx + A)(ex + d)^m}{(bx + a)^2} dx$$

input `int((B*x+A)*(e*x+d)^m/(b*x+a)^2,x)`

output `int((B*x+A)*(e*x+d)^m/(b*x+a)^2,x)`

Fricas [F]

$$\int \frac{(A + Bx)(d + ex)^m}{(a + bx)^2} dx = \int \frac{(Bx + A)(ex + d)^m}{(bx + a)^2} dx$$

input `integrate((B*x+A)*(e*x+d)^m/(b*x+a)^2,x, algorithm="fricas")`

output `integral((B*x + A)*(e*x + d)^m/(b^2*x^2 + 2*a*b*x + a^2), x)`

Sympy [F]

$$\int \frac{(A + Bx)(d + ex)^m}{(a + bx)^2} dx = \int \frac{(A + Bx)(d + ex)^m}{(a + bx)^2} dx$$

input `integrate((B*x+A)*(e*x+d)**m/(b*x+a)**2,x)`

output `Integral((A + B*x)*(d + e*x)**m/(a + b*x)**2, x)`

Maxima [F]

$$\int \frac{(A + Bx)(d + ex)^m}{(a + bx)^2} dx = \int \frac{(Bx + A)(ex + d)^m}{(bx + a)^2} dx$$

input `integrate((B*x+A)*(e*x+d)^m/(b*x+a)^2,x, algorithm="maxima")`

output `integrate((B*x + A)*(e*x + d)^m/(b*x + a)^2, x)`

Giac [F]

$$\int \frac{(A + Bx)(d + ex)^m}{(a + bx)^2} dx = \int \frac{(Bx + A)(ex + d)^m}{(bx + a)^2} dx$$

input `integrate((B*x+A)*(e*x+d)^m/(b*x+a)^2,x, algorithm="giac")`

output `integrate((B*x + A)*(e*x + d)^m/(b*x + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(A + Bx)(d + ex)^m}{(a + bx)^2} dx = \int \frac{(A + Bx)(d + ex)^m}{(a + bx)^2} dx$$

input `int(((A + B*x)*(d + e*x)^m)/(a + b*x)^2,x)`output `int(((A + B*x)*(d + e*x)^m)/(a + b*x)^2, x)`**Reduce [F]**

$$\int \frac{(A + Bx)(d + ex)^m}{(a + bx)^2} dx$$

$$= \frac{(ex + d)^m d + \left(\int \frac{(ex+d)^m x}{be x^2 + aex + bdx + ad} dx \right) a e^2 m - \left(\int \frac{(ex+d)^m x}{be x^2 + aex + bdx + ad} dx \right) b d e m}{a e m}$$

input `int((B*x+A)*(e*x+d)^m/(b*x+a)^2,x)`output `((d + e*x)**m*d + int(((d + e*x)**m*x)/(a*d + a*e*x + b*d*x + b*e*x**2),x) *a*e**2*m - int(((d + e*x)**m*x)/(a*d + a*e*x + b*d*x + b*e*x**2),x)*b*d*e *m)/(a*e*m)`

3.236 $\int \frac{(A+Bx)(d+ex)^m}{(a+bx)^3} dx$

Optimal result	2114
Mathematica [A] (verified)	2114
Rubi [A] (verified)	2115
Maple [F]	2116
Fricas [F]	2116
Sympy [F]	2117
Maxima [F]	2117
Giac [F]	2117
Mupad [F(-1)]	2118
Reduce [F]	2118

Optimal result

Integrand size = 20, antiderivative size = 118

$$\int \frac{(A+Bx)(d+ex)^m}{(a+bx)^3} dx = -\frac{B(d+ex)^{1+m}}{be(1-m)(a+bx)^2} + \frac{e(2bBd - Abe(1-m) - aBe(1+m))(d+ex)^{1+m} \text{Hypergeometric2F1}\left(3, 1+m, 2+m, \frac{b(d+ex)}{bd-ae}\right)}{b(bd-ae)^3(1-m)(1+m)}$$

output

```
-B*(e*x+d)^(1+m)/b/e/(1-m)/(b*x+a)^2+e*(2*B*b*d-A*b*e*(1-m)-a*B*e*(1+m))*(e*x+d)^(1+m)*hypergeom([3, 1+m],[2+m],b*(e*x+d)/(-a*e+b*d))/b/(-a*e+b*d)^3/(1-m)/(1+m)
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.90

$$\int \frac{(A+Bx)(d+ex)^m}{(a+bx)^3} dx = \frac{(d+ex)^{1+m} \left(\frac{-Ab+aB}{(a+bx)^2} + \frac{e(2bBd+Abe(-1+m)-aBe(1+m)) \text{Hypergeometric2F1}\left(2, 1+m, 2+m, \frac{b(d+ex)}{bd-ae}\right)}{(bd-ae)^2(1+m)} \right)}{2b(bd-ae)}$$

input `Integrate[((A + B*x)*(d + e*x)^m)/(a + b*x)^3,x]`

output `((d + e*x)^(1 + m)*((-A*b) + a*B)/(a + b*x)^2 + (e*(2*b*B*d + A*b*e*(-1 + m) - a*B*e*(1 + m))*Hypergeometric2F1[2, 1 + m, 2 + m, (b*(d + e*x))/(b*d - a*e)])/((b*d - a*e)^2*(1 + m)))/(2*b*(b*d - a*e))`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.04, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {87, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx)(d + ex)^m}{(a + bx)^3} dx$$

$$\downarrow 87$$

$$\frac{(-aBe(m + 1) - Abe(1 - m) + 2bBd) \int \frac{(d+ex)^m}{(a+bx)^2} dx}{2b(bd - ae)} - \frac{(Ab - aB)(d + ex)^{m+1}}{2b(a + bx)^2(bd - ae)}$$

$$\downarrow 78$$

$$\frac{e(d + ex)^{m+1}(-aBe(m + 1) - Abe(1 - m) + 2bBd) \text{Hypergeometric2F1}\left(2, m + 1, m + 2, \frac{b(d+ex)}{bd-ae}\right)}{2b(m + 1)(bd - ae)^3} - \frac{(Ab - aB)(d + ex)^{m+1}}{2b(a + bx)^2(bd - ae)}$$

input `Int[((A + B*x)*(d + e*x)^m)/(a + b*x)^3,x]`

output `-1/2*((A*b - a*B)*(d + e*x)^(1 + m))/(b*(b*d - a*e)*(a + b*x)^2) + (e*(2*b*B*d - A*b*e*(1 - m) - a*B*e*(1 + m))*(d + e*x)^(1 + m)*Hypergeometric2F1[2, 1 + m, 2 + m, (b*(d + e*x))/(b*d - a*e)])/(2*b*(b*d - a*e)^3*(1 + m))`

Definitions of rubi rules used

rule 78

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b
*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m
+ 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x]
&& !IntegerQ[m] && IntegerQ[n]
```

rule 87

```
Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p
_), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

Maple [F]

$$\int \frac{(Bx + A)(ex + d)^m}{(bx + a)^3} dx$$

input

```
int((B*x+A)*(e*x+d)^m/(b*x+a)^3,x)
```

output

```
int((B*x+A)*(e*x+d)^m/(b*x+a)^3,x)
```

Fricas [F]

$$\int \frac{(A + Bx)(d + ex)^m}{(a + bx)^3} dx = \int \frac{(Bx + A)(ex + d)^m}{(bx + a)^3} dx$$

input

```
integrate((B*x+A)*(e*x+d)^m/(b*x+a)^3,x, algorithm="fricas")
```

output

```
integral((B*x + A)*(e*x + d)^m/(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3),
x)
```

Sympy [F]

$$\int \frac{(A + Bx)(d + ex)^m}{(a + bx)^3} dx = \int \frac{(A + Bx)(d + ex)^m}{(a + bx)^3} dx$$

input `integrate((B*x+A)*(e*x+d)**m/(b*x+a)**3,x)`

output `Integral((A + B*x)*(d + e*x)**m/(a + b*x)**3, x)`

Maxima [F]

$$\int \frac{(A + Bx)(d + ex)^m}{(a + bx)^3} dx = \int \frac{(Bx + A)(ex + d)^m}{(bx + a)^3} dx$$

input `integrate((B*x+A)*(e*x+d)^m/(b*x+a)^3,x, algorithm="maxima")`

output `integrate((B*x + A)*(e*x + d)^m/(b*x + a)^3, x)`

Giac [F]

$$\int \frac{(A + Bx)(d + ex)^m}{(a + bx)^3} dx = \int \frac{(Bx + A)(ex + d)^m}{(bx + a)^3} dx$$

input `integrate((B*x+A)*(e*x+d)^m/(b*x+a)^3,x, algorithm="giac")`

output `integrate((B*x + A)*(e*x + d)^m/(b*x + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(A + Bx)(d + ex)^m}{(a + bx)^3} dx = \int \frac{(A + Bx)(d + ex)^m}{(a + bx)^3} dx$$

input `int(((A + B*x)*(d + e*x)^m)/(a + b*x)^3,x)`output `int(((A + B*x)*(d + e*x)^m)/(a + b*x)^3, x)`**Reduce [F]**

$$\int \frac{(A + Bx)(d + ex)^m}{(a + bx)^3} dx = \text{Too large to display}$$

input `int((B*x+A)*(e*x+d)^m/(b*x+a)^3,x)`

output

```

((d + e*x)**m*d + int(((d + e*x)**m*x)/(a**3*d*e*m + a**3*e**2*m*x - a**2*
b*d**2 + 2*a**2*b*d*e*m*x - a**2*b*d*e*x + 2*a**2*b*e**2*m*x**2 - 2*a*b**2
*d**2*x + a*b**2*d*e*m*x**2 - 2*a*b**2*d*e*x**2 + a*b**2*e**2*m*x**3 - b**
3*d**2*x**2 - b**3*d*e*x**3),x)*a**3*e**3*m**2 - int(((d + e*x)**m*x)/(a**
3*d*e*m + a**3*e**2*m*x - a**2*b*d**2 + 2*a**2*b*d*e*m*x - a**2*b*d*e*x +
2*a**2*b*e**2*m*x**2 - 2*a*b**2*d**2*x + a*b**2*d*e*m*x**2 - 2*a*b**2*d*e*
x**2 + a*b**2*e**2*m*x**3 - b**3*d**2*x**2 - b**3*d*e*x**3),x)*a**2*b*d*e*
**2*m**2 - int(((d + e*x)**m*x)/(a**3*d*e*m + a**3*e**2*m*x - a**2*b*d**2 +
2*a**2*b*d*e*m*x - a**2*b*d*e*x + 2*a**2*b*e**2*m*x**2 - 2*a*b**2*d**2*x
+ a*b**2*d*e*m*x**2 - 2*a*b**2*d*e*x**2 + a*b**2*e**2*m*x**3 - b**3*d**2*x
**2 - b**3*d*e*x**3),x)*a**2*b*d*e**2*m + int(((d + e*x)**m*x)/(a**3*d*e*m
+ a**3*e**2*m*x - a**2*b*d**2 + 2*a**2*b*d*e*m*x - a**2*b*d*e*x + 2*a**2*
b*e**2*m*x**2 - 2*a*b**2*d**2*x + a*b**2*d*e*m*x**2 - 2*a*b**2*d*e*x**2 +
a*b**2*e**2*m*x**3 - b**3*d**2*x**2 - b**3*d*e*x**3),x)*a**2*b*e**3*m**2*x
+ int(((d + e*x)**m*x)/(a**3*d*e*m + a**3*e**2*m*x - a**2*b*d**2 + 2*a**2
*b*d*e*m*x - a**2*b*d*e*x + 2*a**2*b*e**2*m*x**2 - 2*a*b**2*d**2*x + a*b**
2*d*e*m*x**2 - 2*a*b**2*d*e*x**2 + a*b**2*e**2*m*x**3 - b**3*d**2*x**2 - b
**3*d*e*x**3),x)*a*b**2*d**2*e*m - int(((d + e*x)**m*x)/(a**3*d*e*m + a**3
*e**2*m*x - a**2*b*d**2 + 2*a**2*b*d*e*m*x - a**2*b*d*e*x + 2*a**2*b*e**2*
m*x**2 - 2*a*b**2*d**2*x + a*b**2*d*e*m*x**2 - 2*a*b**2*d*e*x**2 + a*b*...

```

3.237 $\int (1 - 2x)(2 + 3x)^m(3 + 5x)^3 dx$

Optimal result	2120
Mathematica [A] (verified)	2120
Rubi [A] (verified)	2121
Maple [C] (verified)	2122
Fricas [B] (verification not implemented)	2123
Sympy [B] (verification not implemented)	2123
Maxima [B] (verification not implemented)	2124
Giac [B] (verification not implemented)	2125
Mupad [B] (verification not implemented)	2126
Reduce [B] (verification not implemented)	2127

Optimal result

Integrand size = 20, antiderivative size = 91

$$\int (1 - 2x)(2 + 3x)^m(3 + 5x)^3 dx = -\frac{7(2 + 3x)^{1+m}}{243(1 + m)} + \frac{107(2 + 3x)^{2+m}}{243(2 + m)} - \frac{185(2 + 3x)^{3+m}}{81(3 + m)} + \frac{1025(2 + 3x)^{4+m}}{243(4 + m)} - \frac{250(2 + 3x)^{5+m}}{243(5 + m)}$$

output

```
-7*(2+3*x)^(1+m)/(243+243*m)+107*(2+3*x)^(2+m)/(486+243*m)-185*(2+3*x)^(3+m)/(243+81*m)+1025*(2+3*x)^(4+m)/(972+243*m)-250*(2+3*x)^(5+m)/(1215+243*m)
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.82

$$\int (1 - 2x)(2 + 3x)^m(3 + 5x)^3 dx = \frac{1}{243}(2 + 3x)^{1+m} \left(-\frac{7}{1 + m} + \frac{107(2 + 3x)}{2 + m} - \frac{555(2 + 3x)^2}{3 + m} + \frac{1025(2 + 3x)^3}{4 + m} - \frac{250(2 + 3x)^4}{5 + m} \right)$$

input `Integrate[(1 - 2*x)*(2 + 3*x)^m*(3 + 5*x)^3,x]`

output `((2 + 3*x)^(1 + m)*(-7/(1 + m) + (107*(2 + 3*x))/(2 + m) - (555*(2 + 3*x)^2)/(3 + m) + (1025*(2 + 3*x)^3)/(4 + m) - (250*(2 + 3*x)^4)/(5 + m))/243`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (1 - 2x)(5x + 3)^3(3x + 2)^m dx$$

$$\downarrow 86$$

$$\int \left(-\frac{7}{81}(3x + 2)^m + \frac{107}{81}(3x + 2)^{m+1} - \frac{185}{27}(3x + 2)^{m+2} + \frac{1025}{81}(3x + 2)^{m+3} - \frac{250}{81}(3x + 2)^{m+4} \right) dx$$

$$\downarrow 2009$$

$$-\frac{7(3x + 2)^{m+1}}{243(m + 1)} + \frac{107(3x + 2)^{m+2}}{243(m + 2)} - \frac{185(3x + 2)^{m+3}}{81(m + 3)} + \frac{1025(3x + 2)^{m+4}}{243(m + 4)} - \frac{250(3x + 2)^{m+5}}{243(m + 5)}$$

input `Int[(1 - 2*x)*(2 + 3*x)^m*(3 + 5*x)^3,x]`

output `(-7*(2 + 3*x)^(1 + m))/(243*(1 + m)) + (107*(2 + 3*x)^(2 + m))/(243*(2 + m)) - (185*(2 + 3*x)^(3 + m))/(81*(3 + m)) + (1025*(2 + 3*x)^(4 + m))/(243*(4 + m)) - (250*(2 + 3*x)^(5 + m))/(243*(5 + m))`

Defintions of rubi rules used

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3.

Time = 0.28 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.11

method	result
meijerg	$27 \cdot 2^m x \operatorname{hypergeom}\left(\left[1, -m\right], \left[2\right], -\frac{3x}{2}\right) + 81 \cdot 2^{m-1} x^2 \operatorname{hypergeom}\left(\left[2, -m\right], \left[3\right], -\frac{3x}{2}\right) - 15 \cdot 2^m$
gosper	$-\frac{(2+3x)^{1+m} (6750m^4x^4 + 8775m^4x^3 + 67500m^3x^4 + 1215m^4x^2 + 78525m^3x^3 + 236250m^2x^4 - 2187m^4x - 2970m^3x^2 + 251775m^2x^3 + 337500m^3x^4)}{(6750m^4x^4 + 8775m^4x^3 + 67500m^3x^4 + 1215m^4x^2 + 78525m^3x^3 + 236250m^2x^4 - 2187m^4x - 2970m^3x^2 + 251775m^2x^3 + 337500m^3x^4)}$
orering	$-\frac{(20250m^4x^5 + 39825m^4x^4 + 202500m^3x^5 + 21195m^4x^3 + 370575m^3x^4 + 708750m^2x^5 - 4131m^4x^2 + 148140m^3x^3 + 1227825m^2x^4 - 1052076m^3x^5 - 97600(2+3x)^m m - 79650x^4(2+3x)^m m^4 + 1417500x^5(2+3x)^m m^5)}{(20250m^4x^5 + 39825m^4x^4 + 202500m^3x^5 + 21195m^4x^3 + 370575m^3x^4 + 708750m^2x^5 - 4131m^4x^2 + 148140m^3x^3 + 1227825m^2x^4 - 1052076m^3x^5 - 97600(2+3x)^m m - 79650x^4(2+3x)^m m^4 + 1417500x^5(2+3x)^m m^5)}$
risch	$-\frac{250x^5 e^{m \ln(2+3x)}}{5+m} + \frac{2(729m^4 + 8748m^3 + 33183m^2 + 49620m + 24400)e^{m \ln(2+3x)}}{81(m^5 + 15m^4 + 85m^3 + 225m^2 + 274m + 120)} - \frac{25(59m + 195)x^4 e^{m \ln(2+3x)}}{3(m^2 + 9m + 20)} - \frac{5000x^5(2+3x)^m m^3 + 40500x^5(2+3x)^m m^4 - 1052076x(2+3x)^m m - 97600(2+3x)^m m + 79650x^4(2+3x)^m m^4 + 1417500x^5(2+3x)^m m^5}{(20250m^4x^5 + 39825m^4x^4 + 202500m^3x^5 + 21195m^4x^3 + 370575m^3x^4 + 708750m^2x^5 - 4131m^4x^2 + 148140m^3x^3 + 1227825m^2x^4 - 1052076m^3x^5 - 97600(2+3x)^m m - 79650x^4(2+3x)^m m^4 + 1417500x^5(2+3x)^m m^5)}$
norman	$-\frac{250x^5 e^{m \ln(2+3x)}}{5+m} + \frac{2(729m^4 + 8748m^3 + 33183m^2 + 49620m + 24400)e^{m \ln(2+3x)}}{81(m^5 + 15m^4 + 85m^3 + 225m^2 + 274m + 120)} - \frac{25(59m + 195)x^4 e^{m \ln(2+3x)}}{3(m^2 + 9m + 20)} - \frac{5000x^5(2+3x)^m m^3 + 40500x^5(2+3x)^m m^4 - 1052076x(2+3x)^m m - 97600(2+3x)^m m + 79650x^4(2+3x)^m m^4 + 1417500x^5(2+3x)^m m^5}{(20250m^4x^5 + 39825m^4x^4 + 202500m^3x^5 + 21195m^4x^3 + 370575m^3x^4 + 708750m^2x^5 - 4131m^4x^2 + 148140m^3x^3 + 1227825m^2x^4 - 1052076m^3x^5 - 97600(2+3x)^m m - 79650x^4(2+3x)^m m^4 + 1417500x^5(2+3x)^m m^5)}$
parallelrisch	$-\frac{405000x^5(2+3x)^m m^3 + 40500x^5(2+3x)^m m^4 - 1052076x(2+3x)^m m - 97600(2+3x)^m m + 79650x^4(2+3x)^m m^4 + 1417500x^5(2+3x)^m m^5}{(20250m^4x^5 + 39825m^4x^4 + 202500m^3x^5 + 21195m^4x^3 + 370575m^3x^4 + 708750m^2x^5 - 4131m^4x^2 + 148140m^3x^3 + 1227825m^2x^4 - 1052076m^3x^5 - 97600(2+3x)^m m - 79650x^4(2+3x)^m m^4 + 1417500x^5(2+3x)^m m^5)}$

```
input int((1-2*x)*(2+3*x)^m*(3+5*x)^3,x,method=_RETURNVERBOSE)
```

```
output 27*2^m*x*hypergeom([1, -m], [2], -3/2*x)+81*2^(m-1)*x^2*hypergeom([2, -m], [3], -3/2*x)-15*2^m*x^3*hypergeom([3, -m], [4], -3/2*x)-325*2^(-2+m)*x^4*hypergeom([4, -m], [5], -3/2*x)-25*2^(1+m)*x^5*hypergeom([5, -m], [6], -3/2*x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 175 vs. $2(81) = 162$.

Time = 0.10 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.92

$$\int (1 - 2x)(2 + 3x)^m(3 + 5x)^3 dx = \frac{(20250(m^4 + 10m^3 + 35m^2 + 50m + 24)x^5 + 675(59m^4 + 549m^3 + 1819m^2 + 2499m + 1170)x^4 - 1458m^4 + 45(471m^4 + 3292m^3 + 199m^2 + 8618m + 3240)x^3 - 17496m^3 - 9(459m^4 + 10677m^3 + 50581m^2 + 84103m + 43740)x^2 - 66366m^2 - 3(2187m^4 + 28782m^3 + 114405m^2 + 175346m + 87480)x - 99240m - 48800)(3x + 2)^m}{(m^5 + 15m^4 + 85m^3 + 225m^2 + 274m + 120)}$$

input `integrate((1-2*x)*(2+3*x)^m*(3+5*x)^3,x, algorithm="fricas")`

output `-1/81*(20250*(m^4 + 10*m^3 + 35*m^2 + 50*m + 24)*x^5 + 675*(59*m^4 + 549*m^3 + 1819*m^2 + 2499*m + 1170)*x^4 - 1458*m^4 + 45*(471*m^4 + 3292*m^3 + 199*m^2 + 8618*m + 3240)*x^3 - 17496*m^3 - 9*(459*m^4 + 10677*m^3 + 50581*m^2 + 84103*m + 43740)*x^2 - 66366*m^2 - 3*(2187*m^4 + 28782*m^3 + 114405*m^2 + 175346*m + 87480)*x - 99240*m - 48800)*(3*x + 2)^m/(m^5 + 15*m^4 + 85*m^3 + 225*m^2 + 274*m + 120)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1822 vs. $2(75) = 150$.

Time = 0.59 (sec) , antiderivative size = 1822, normalized size of antiderivative = 20.02

$$\int (1 - 2x)(2 + 3x)^m(3 + 5x)^3 dx = \text{Too large to display}$$

input `integrate((1-2*x)*(2+3*x)**m*(3+5*x)**3,x)`

output

```
Piecewise((-243000*x**4*log(3*x + 2)/(236196*x**4 + 629856*x**3 + 629856*x**2 + 279936*x + 46656) - 648000*x**3*log(3*x + 2)/(236196*x**4 + 629856*x**3 + 629856*x**2 + 279936*x + 46656) - 332100*x**3/(236196*x**4 + 629856*x**3 + 629856*x**2 + 279936*x + 46656) - 648000*x**2*log(3*x + 2)/(236196*x**4 + 629856*x**3 + 629856*x**2 + 279936*x + 46656) - 634230*x**2/(236196*x**4 + 629856*x**3 + 629856*x**2 + 279936*x + 46656) - 288000*x*log(3*x + 2)/(236196*x**4 + 629856*x**3 + 629856*x**2 + 279936*x + 46656) - 404124*x/(236196*x**4 + 629856*x**3 + 629856*x**2 + 279936*x + 46656) - 48000*log(3*x + 2)/(236196*x**4 + 629856*x**3 + 629856*x**2 + 279936*x + 46656) - 85915/(236196*x**4 + 629856*x**3 + 629856*x**2 + 279936*x + 46656), Eq(m, -5)), (-121500*x**4/(39366*x**3 + 78732*x**2 + 52488*x + 11664) + 166050*x**3*log(3*x + 2)/(39366*x**3 + 78732*x**2 + 52488*x + 11664) + 332100*x**2*log(3*x + 2)/(39366*x**3 + 78732*x**2 + 52488*x + 11664) + 353970*x**2/(39366*x**3 + 78732*x**2 + 52488*x + 11664) + 221400*x*log(3*x + 2)/(39366*x**3 + 78732*x**2 + 52488*x + 11664) + 326997*x/(39366*x**3 + 78732*x**2 + 52488*x + 11664) + 49200*log(3*x + 2)/(39366*x**3 + 78732*x**2 + 52488*x + 11664) + 84692/(39366*x**3 + 78732*x**2 + 52488*x + 11664), Eq(m, -4)), (-6750*x**4/(1458*x**2 + 1944*x + 648) + 450*x**3/(1458*x**2 + 1944*x + 648) - 3330*x**2*log(3*x + 2)/(1458*x**2 + 1944*x + 648) - 4440*x*log(3*x + 2)/(1458*x**2 + 1944*x + 648) - 8814*x/(1458*x**2 + 1944*x + 648) - 1480*...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 295 vs. $2(81) = 162$.

Time = 0.04 (sec) , antiderivative size = 295, normalized size of antiderivative = 3.24

$$\int (1 - 2x)(2 + 3x)^m(3 + 5x)^3 dx =$$

$$\frac{250(81(m^4 + 10m^3 + 35m^2 + 50m + 24)x^5 + 54(m^4 + 6m^3 + 11m^2 + 6m)x^4 - 144(m^3 + 3m^2 + 12m + 12)x^3 + 325(27(m^3 + 6m^2 + 11m + 6)x^4 + 18(m^3 + 3m^2 + 2m)x^3 - 36(m^2 + m)x^2 + 48mx - 32)(3x + 2))}{81(m^5 + 15m^4 + 85m^3 + 225m^2 + 274m + 12)}$$

$$\frac{5(27(m^2 + 3m + 2)x^3 + 18(m^2 + m)x^2 - 24mx + 16)(3x + 2)^m}{27(m^4 + 10m^3 + 35m^2 + 50m + 24)}$$

$$\frac{9(9(m + 1)x^2 + 6mx - 4)(3x + 2)^m}{3(m^3 + 6m^2 + 11m + 6)} + \frac{9(3x + 2)^{m+1}}{m + 1}$$

input

```
integrate((1-2*x)*(2+3*x)^m*(3+5*x)^3,x, algorithm="maxima")
```

output

```
-250/81*(81*(m^4 + 10*m^3 + 35*m^2 + 50*m + 24)*x^5 + 54*(m^4 + 6*m^3 + 11
*m^2 + 6*m)*x^4 - 144*(m^3 + 3*m^2 + 2*m)*x^3 + 288*(m^2 + m)*x^2 - 384*m*
x + 256)*(3*x + 2)^m/(m^5 + 15*m^4 + 85*m^3 + 225*m^2 + 274*m + 120) - 325
/27*(27*(m^3 + 6*m^2 + 11*m + 6)*x^4 + 18*(m^3 + 3*m^2 + 2*m)*x^3 - 36*(m^
2 + m)*x^2 + 48*m*x - 32)*(3*x + 2)^m/(m^4 + 10*m^3 + 35*m^2 + 50*m + 24)
- 5/3*(27*(m^2 + 3*m + 2)*x^3 + 18*(m^2 + m)*x^2 - 24*m*x + 16)*(3*x + 2)^
m/(m^3 + 6*m^2 + 11*m + 6) + 9*(9*(m + 1)*x^2 + 6*m*x - 4)*(3*x + 2)^m/(m^
2 + 3*m + 2) + 9*(3*x + 2)^(m + 1)/(m + 1)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 423 vs. $2(81) = 162$.

Time = 0.12 (sec) , antiderivative size = 423, normalized size of antiderivative = 4.65

$$\int (1 - 2x)(2 + 3x)^m(3 + 5x)^3 dx =$$

$$\frac{20250 m^4 (3x + 2)^m x^5 + 39825 m^4 (3x + 2)^m x^4 + 202500 m^3 (3x + 2)^m x^5 + 21195 m^4 (3x + 2)^m x^3 +$$

input

```
integrate((1-2*x)*(2+3*x)^m*(3+5*x)^3,x, algorithm="giac")
```

output

```
-1/81*(20250*m^4*(3*x + 2)^m*x^5 + 39825*m^4*(3*x + 2)^m*x^4 + 202500*m^3*
(3*x + 2)^m*x^5 + 21195*m^4*(3*x + 2)^m*x^3 + 370575*m^3*(3*x + 2)^m*x^4 +
708750*m^2*(3*x + 2)^m*x^5 - 4131*m^4*(3*x + 2)^m*x^2 + 148140*m^3*(3*x +
2)^m*x^3 + 1227825*m^2*(3*x + 2)^m*x^4 + 1012500*m*(3*x + 2)^m*x^5 - 6561
*m^4*(3*x + 2)^m*x - 96093*m^3*(3*x + 2)^m*x^2 + 368955*m^2*(3*x + 2)^m*x^
3 + 1686825*m*(3*x + 2)^m*x^4 + 486000*(3*x + 2)^m*x^5 - 1458*m^4*(3*x + 2
)^m - 86346*m^3*(3*x + 2)^m*x - 455229*m^2*(3*x + 2)^m*x^2 + 387810*m*(3*x
+ 2)^m*x^3 + 789750*(3*x + 2)^m*x^4 - 17496*m^3*(3*x + 2)^m - 343215*m^2*
(3*x + 2)^m*x - 756927*m*(3*x + 2)^m*x^2 + 145800*(3*x + 2)^m*x^3 - 66366*
m^2*(3*x + 2)^m - 526038*m*(3*x + 2)^m*x - 393660*(3*x + 2)^m*x^2 - 99240*
m*(3*x + 2)^m - 262440*(3*x + 2)^m*x - 48800*(3*x + 2)^m)/(m^5 + 15*m^4 +
85*m^3 + 225*m^2 + 274*m + 120)
```

Mupad [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 313, normalized size of antiderivative = 3.44

$$\int (1-2x)(2+3x)^m(3+5x)^3 dx$$

$$= (3x+2)^m \left(\frac{1458m^4 + 17496m^3 + 66366m^2 + 99240m + 48800}{81m^5 + 1215m^4 + 6885m^3 + 18225m^2 + 22194m + 9720} \right. \\ - \frac{x^3(21195m^4 + 148140m^3 + 368955m^2 + 387810m + 145800)}{81m^5 + 1215m^4 + 6885m^3 + 18225m^2 + 22194m + 9720} \\ + \frac{x^2(4131m^4 + 96093m^3 + 455229m^2 + 756927m + 393660)}{81m^5 + 1215m^4 + 6885m^3 + 18225m^2 + 22194m + 9720} \\ - \frac{x^5(20250m^4 + 202500m^3 + 708750m^2 + 1012500m + 486000)}{81m^5 + 1215m^4 + 6885m^3 + 18225m^2 + 22194m + 9720} \\ - \frac{x^4(39825m^4 + 370575m^3 + 1227825m^2 + 1686825m + 789750)}{81m^5 + 1215m^4 + 6885m^3 + 18225m^2 + 22194m + 9720} \\ \left. + \frac{x(6561m^4 + 86346m^3 + 343215m^2 + 526038m + 262440)}{81m^5 + 1215m^4 + 6885m^3 + 18225m^2 + 22194m + 9720} \right)$$

input `int(-(2*x - 1)*(3*x + 2)^m*(5*x + 3)^3,x)`output `(3*x + 2)^m*((99240*m + 66366*m^2 + 17496*m^3 + 1458*m^4 + 48800)/(22194*m + 18225*m^2 + 6885*m^3 + 1215*m^4 + 81*m^5 + 9720) - (x^3*(387810*m + 368955*m^2 + 148140*m^3 + 21195*m^4 + 145800))/(22194*m + 18225*m^2 + 6885*m^3 + 1215*m^4 + 81*m^5 + 9720) + (x^2*(756927*m + 455229*m^2 + 96093*m^3 + 4131*m^4 + 393660))/(22194*m + 18225*m^2 + 6885*m^3 + 1215*m^4 + 81*m^5 + 9720) - (x^5*(1012500*m + 708750*m^2 + 202500*m^3 + 20250*m^4 + 486000))/(22194*m + 18225*m^2 + 6885*m^3 + 1215*m^4 + 81*m^5 + 9720) - (x^4*(1686825*m + 1227825*m^2 + 370575*m^3 + 39825*m^4 + 789750))/(22194*m + 18225*m^2 + 6885*m^3 + 1215*m^4 + 81*m^5 + 9720) + (x*(526038*m + 343215*m^2 + 86346*m^3 + 6561*m^4 + 262440))/(22194*m + 18225*m^2 + 6885*m^3 + 1215*m^4 + 81*m^5 + 9720))`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 220, normalized size of antiderivative = 2.42

$$\int (1 - 2x)(2 + 3x)^m(3 + 5x)^3 dx$$

$$= \frac{(3x + 2)^m (-20250m^4x^5 - 39825m^4x^4 - 202500m^3x^5 - 21195m^4x^3 - 370575m^3x^4 - 708750m^2x^5 + 4131m^4x^2 + 6561m^4x + 1458m^4 - 202500m^3x^5 - 370575m^3x^4 - 148140m^3x^3 + 96093m^3x^2 + 86346m^3x + 17496m^3 - 708750m^2x^5 - 1227825m^2x^4 - 368955m^2x^3 + 455229m^2x^2 + 343215m^2x + 66366m^2 - 1012500mx^5 - 1686825mx^4 - 387810mx^3 + 756927mx^2 + 526038mx + 99240m - 486000x^5 - 789750x^4 - 145800x^3 + 393660x^2 + 262440x + 48800)}{(81(m^5 + 15m^4 + 85m^3 + 225m^2 + 274m + 120))}$$

input `int((1-2*x)*(2+3*x)^m*(3+5*x)^3,x)`output `((3*x + 2)**m*(- 20250*m**4*x**5 - 39825*m**4*x**4 - 21195*m**4*x**3 + 4131*m**4*x**2 + 6561*m**4*x + 1458*m**4 - 202500*m**3*x**5 - 370575*m**3*x**4 - 148140*m**3*x**3 + 96093*m**3*x**2 + 86346*m**3*x + 17496*m**3 - 708750*m**2*x**5 - 1227825*m**2*x**4 - 368955*m**2*x**3 + 455229*m**2*x**2 + 343215*m**2*x + 66366*m**2 - 1012500*m*x**5 - 1686825*m*x**4 - 387810*m*x**3 + 756927*m*x**2 + 526038*m*x + 99240*m - 486000*x**5 - 789750*x**4 - 145800*x**3 + 393660*x**2 + 262440*x + 48800))/(81*(m**5 + 15*m**4 + 85*m**3 + 225*m**2 + 274*m + 120))`

3.238 $\int (1 - 2x)(2 + 3x)^m(3 + 5x)^2 dx$

Optimal result	2128
Mathematica [A] (verified)	2128
Rubi [A] (verified)	2129
Maple [C] (verified)	2130
Fricas [A] (verification not implemented)	2130
Sympy [B] (verification not implemented)	2131
Maxima [B] (verification not implemented)	2132
Giac [B] (verification not implemented)	2132
Mupad [B] (verification not implemented)	2133
Reduce [B] (verification not implemented)	2134

Optimal result

Integrand size = 20, antiderivative size = 73

$$\int (1 - 2x)(2 + 3x)^m(3 + 5x)^2 dx = \frac{7(2 + 3x)^{1+m}}{81(1 + m)} - \frac{8(2 + 3x)^{2+m}}{9(2 + m)} + \frac{65(2 + 3x)^{3+m}}{27(3 + m)} - \frac{50(2 + 3x)^{4+m}}{81(4 + m)}$$

output

```
7*(2+3*x)^(1+m)/(81+81*m)-8*(2+3*x)^(2+m)/(18+9*m)+65*(2+3*x)^(3+m)/(81+27*m)-50*(2+3*x)^(4+m)/(324+81*m)
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.84

$$\int (1 - 2x)(2 + 3x)^m(3 + 5x)^2 dx = \frac{1}{81}(2 + 3x)^{1+m} \left(\frac{7}{1 + m} - \frac{72(2 + 3x)}{2 + m} + \frac{195(2 + 3x)^2}{3 + m} - \frac{50(2 + 3x)^3}{4 + m} \right)$$

input

```
Integrate[(1 - 2*x)*(2 + 3*x)^m*(3 + 5*x)^2,x]
```

output
$$\frac{((2 + 3x)^{(1 + m)} \cdot (7/(1 + m) - (72 \cdot (2 + 3x))/(2 + m) + (195 \cdot (2 + 3x)^2)/(3 + m) - (50 \cdot (2 + 3x)^3)/(4 + m)))/81}$$

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (1 - 2x)(5x + 3)^2(3x + 2)^m dx$$

↓ 86

$$\int \left(\frac{7}{27}(3x + 2)^m - \frac{8}{3}(3x + 2)^{m+1} + \frac{65}{9}(3x + 2)^{m+2} - \frac{50}{27}(3x + 2)^{m+3} \right) dx$$

↓ 2009

$$\frac{7(3x + 2)^{m+1}}{81(m + 1)} - \frac{8(3x + 2)^{m+2}}{9(m + 2)} + \frac{65(3x + 2)^{m+3}}{27(m + 3)} - \frac{50(3x + 2)^{m+4}}{81(m + 4)}$$

input `Int[(1 - 2*x)*(2 + 3*x)^m*(3 + 5*x)^2,x]`

output
$$\frac{7 \cdot (2 + 3x)^{(1 + m)}}{81 \cdot (1 + m)} - \frac{8 \cdot (2 + 3x)^{(2 + m)}}{9 \cdot (2 + m)} + \frac{65 \cdot (2 + 3x)^{(3 + m)}}{27 \cdot (3 + m)} - \frac{50 \cdot (2 + 3x)^{(4 + m)}}{81 \cdot (4 + m)}$$

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;`
`FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

output

```
-1/27*(1350*(m^3 + 6*m^2 + 11*m + 6)*x^4 + 45*(41*m^3 + 207*m^2 + 334*m +
168)*x^3 - 162*m^3 + 18*(17*m^3 - 69*m^2 - 302*m - 216)*x^2 - 1314*m^2 - 3
*(153*m^3 + 1513*m^2 + 3290*m + 1944)*x - 2644*m - 1520)*(3*x + 2)^m/(m^4
+ 10*m^3 + 35*m^2 + 50*m + 24)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1017 vs. $2(60) = 120$.

Time = 0.43 (sec) , antiderivative size = 1017, normalized size of antiderivative = 13.93

$$\int (1 - 2x)(2 + 3x)^m(3 + 5x)^2 dx = \text{Too large to display}$$

input

```
integrate((1-2*x)*(2+3*x)**m*(3+5*x)**2,x)
```

output

```
Piecewise((-4050*x**3*log(3*x + 2)/(6561*x**3 + 13122*x**2 + 8748*x + 1944
) - 8100*x**2*log(3*x + 2)/(6561*x**3 + 13122*x**2 + 8748*x + 1944) - 5265
*x**2/(6561*x**3 + 13122*x**2 + 8748*x + 1944) - 5400*x*log(3*x + 2)/(6561
*x**3 + 13122*x**2 + 8748*x + 1944) - 6696*x/(6561*x**3 + 13122*x**2 + 874
8*x + 1944) - 1200*log(3*x + 2)/(6561*x**3 + 13122*x**2 + 8748*x + 1944) -
2131/(6561*x**3 + 13122*x**2 + 8748*x + 1944), Eq(m, -4)), (-900*x**3/(48
6*x**2 + 648*x + 216) + 1170*x**2*log(3*x + 2)/(486*x**2 + 648*x + 216) +
1560*x*log(3*x + 2)/(486*x**2 + 648*x + 216) + 1344*x/(486*x**2 + 648*x +
216) + 520*log(3*x + 2)/(486*x**2 + 648*x + 216) + 627/(486*x**2 + 648*x +
216), Eq(m, -3)), (-75*x**3/(27*x + 18) + 45*x**2/(27*x + 18) - 24*x*log(
3*x + 2)/(27*x + 18) - 16*log(3*x + 2)/(27*x + 18) - 43/(27*x + 18), Eq(m,
-2)), (-50*x**3/9 - 5*x**2/18 + 118*x/27 + 7*log(3*x + 2)/81, Eq(m, -1)),
(-1350*m**3*x**4*(3*x + 2)**m/(27*m**4 + 270*m**3 + 945*m**2 + 1350*m + 6
48) - 1845*m**3*x**3*(3*x + 2)**m/(27*m**4 + 270*m**3 + 945*m**2 + 1350*m
+ 648) - 306*m**3*x**2*(3*x + 2)**m/(27*m**4 + 270*m**3 + 945*m**2 + 1350*
m + 648) + 459*m**3*x*(3*x + 2)**m/(27*m**4 + 270*m**3 + 945*m**2 + 1350*
m + 648) + 162*m**3*(3*x + 2)**m/(27*m**4 + 270*m**3 + 945*m**2 + 1350*m +
648) - 8100*m**2*x**4*(3*x + 2)**m/(27*m**4 + 270*m**3 + 945*m**2 + 1350*
m + 648) - 9315*m**2*x**3*(3*x + 2)**m/(27*m**4 + 270*m**3 + 945*m**2 + 135
0*m + 648) + 1242*m**2*x**2*(3*x + 2)**m/(27*m**4 + 270*m**3 + 945*m**2...
```


Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 183 vs. $2(65) = 130$.

Time = 0.04 (sec) , antiderivative size = 183, normalized size of antiderivative = 2.51

$$\int (1 - 2x)(2 + 3x)^m(3 + 5x)^2 dx =$$

$$\frac{50(27(m^3 + 6m^2 + 11m + 6)x^4 + 18(m^3 + 3m^2 + 2m)x^3 - 36(m^2 + m)x^2 + 48mx - 32)(3x + 2)}{27(m^4 + 10m^3 + 35m^2 + 50m + 24)}$$

$$- \frac{35(27(m^2 + 3m + 2)x^3 + 18(m^2 + m)x^2 - 24mx + 16)(3x + 2)^m}{27(m^3 + 6m^2 + 11m + 6)}$$

$$+ \frac{4(9(m + 1)x^2 + 6mx - 4)(3x + 2)^m}{3(m^2 + 3m + 2)} + \frac{3(3x + 2)^{m+1}}{m + 1}$$

input `integrate((1-2*x)*(2+3*x)^m*(3+5*x)^2,x, algorithm="maxima")`

output `-50/27*(27*(m^3 + 6*m^2 + 11*m + 6)*x^4 + 18*(m^3 + 3*m^2 + 2*m)*x^3 - 36*(m^2 + m)*x^2 + 48*m*x - 32)*(3*x + 2)^m/(m^4 + 10*m^3 + 35*m^2 + 50*m + 24) - 35/27*(27*(m^2 + 3*m + 2)*x^3 + 18*(m^2 + m)*x^2 - 24*m*x + 16)*(3*x + 2)^m/(m^3 + 6*m^2 + 11*m + 6) + 4/3*(9*(m + 1)*x^2 + 6*m*x - 4)*(3*x + 2)^m/(m^2 + 3*m + 2) + 3*(3*x + 2)^(m + 1)/(m + 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 278 vs. $2(65) = 130$.

Time = 0.11 (sec) , antiderivative size = 278, normalized size of antiderivative = 3.81

$$\int (1 - 2x)(2 + 3x)^m(3 + 5x)^2 dx =$$

$$\frac{1350 m^3(3x + 2)^m x^4 + 1845 m^3(3x + 2)^m x^3 + 8100 m^2(3x + 2)^m x^4 + 306 m^3(3x + 2)^m x^2 + 9315 m^3(3x + 2)^m x + 1845 m^3(3x + 2)^m}{27(m^4 + 10m^3 + 35m^2 + 50m + 24)}$$

input `integrate((1-2*x)*(2+3*x)^m*(3+5*x)^2,x, algorithm="giac")`

output

```
-1/27*(1350*m^3*(3*x + 2)^m*x^4 + 1845*m^3*(3*x + 2)^m*x^3 + 8100*m^2*(3*x
+ 2)^m*x^4 + 306*m^3*(3*x + 2)^m*x^2 + 9315*m^2*(3*x + 2)^m*x^3 + 14850*m
*(3*x + 2)^m*x^4 - 459*m^3*(3*x + 2)^m*x - 1242*m^2*(3*x + 2)^m*x^2 + 1503
0*m*(3*x + 2)^m*x^3 + 8100*(3*x + 2)^m*x^4 - 162*m^3*(3*x + 2)^m - 4539*m^
2*(3*x + 2)^m*x - 5436*m*(3*x + 2)^m*x^2 + 7560*(3*x + 2)^m*x^3 - 1314*m^2
*(3*x + 2)^m - 9870*m*(3*x + 2)^m*x - 3888*(3*x + 2)^m*x^2 - 2644*m*(3*x +
2)^m - 5832*(3*x + 2)^m*x - 1520*(3*x + 2)^m)/(m^4 + 10*m^3 + 35*m^2 + 50
*m + 24)
```

Mupad [B] (verification not implemented)

Time = 1.18 (sec) , antiderivative size = 211, normalized size of antiderivative = 2.89

$$\int (1 - 2x)(2 + 3x)^m(3 + 5x)^2 dx$$

$$= (3x + 2)^m \left(\frac{162m^3 + 1314m^2 + 2644m + 1520}{27m^4 + 270m^3 + 945m^2 + 1350m + 648} \right. \\ \left. + \frac{x(459m^3 + 4539m^2 + 9870m + 5832)}{27m^4 + 270m^3 + 945m^2 + 1350m + 648} \right. \\ \left. + \frac{x^2(-306m^3 + 1242m^2 + 5436m + 3888)}{27m^4 + 270m^3 + 945m^2 + 1350m + 648} \right. \\ \left. - \frac{x^4(1350m^3 + 8100m^2 + 14850m + 8100)}{27m^4 + 270m^3 + 945m^2 + 1350m + 648} \right. \\ \left. - \frac{x^3(1845m^3 + 9315m^2 + 15030m + 7560)}{27m^4 + 270m^3 + 945m^2 + 1350m + 648} \right)$$

input

```
int(-(2*x - 1)*(3*x + 2)^m*(5*x + 3)^2,x)
```

output

```
(3*x + 2)^m*((2644*m + 1314*m^2 + 162*m^3 + 1520)/(1350*m + 945*m^2 + 270*
m^3 + 27*m^4 + 648) + (x*(9870*m + 4539*m^2 + 459*m^3 + 5832))/(1350*m + 9
45*m^2 + 270*m^3 + 27*m^4 + 648) + (x^2*(5436*m + 1242*m^2 - 306*m^3 + 388
8))/(1350*m + 945*m^2 + 270*m^3 + 27*m^4 + 648) - (x^4*(14850*m + 8100*m^2
+ 1350*m^3 + 8100))/(1350*m + 945*m^2 + 270*m^3 + 27*m^4 + 648) - (x^3*(1
5030*m + 9315*m^2 + 1845*m^3 + 7560))/(1350*m + 945*m^2 + 270*m^3 + 27*m^4
+ 648))
```

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.99

$$\int (1 - 2x)(2 + 3x)^m(3 + 5x)^2 dx$$

$$= \frac{(3x + 2)^m (-1350m^3x^4 - 1845m^3x^3 - 8100m^2x^4 - 306m^3x^2 - 9315m^2x^3 - 14850mx^4 + 459m^3x + 12)}{27m^4}$$

input `int((1-2*x)*(2+3*x)^m*(3+5*x)^2,x)`output `((3*x + 2)**m*(- 1350*m**3*x**4 - 1845*m**3*x**3 - 306*m**3*x**2 + 459*m**3*x + 162*m**3 - 8100*m**2*x**4 - 9315*m**2*x**3 + 1242*m**2*x**2 + 4539*m**2*x + 1314*m**2 - 14850*m*x**4 - 15030*m*x**3 + 5436*m*x**2 + 9870*m*x + 2644*m - 8100*x**4 - 7560*x**3 + 3888*x**2 + 5832*x + 1520))/(27*(m**4 + 10*m**3 + 35*m**2 + 50*m + 24))`

3.239 $\int (1 - 2x)(2 + 3x)^m(3 + 5x) dx$

Optimal result	2135
Mathematica [A] (verified)	2135
Rubi [A] (verified)	2136
Maple [C] (verified)	2137
Fricas [A] (verification not implemented)	2137
Sympy [B] (verification not implemented)	2138
Maxima [B] (verification not implemented)	2139
Giac [B] (verification not implemented)	2139
Mupad [B] (verification not implemented)	2140
Reduce [B] (verification not implemented)	2140

Optimal result

Integrand size = 18, antiderivative size = 55

$$\int (1 - 2x)(2 + 3x)^m(3 + 5x) dx = -\frac{7(2 + 3x)^{1+m}}{27(1 + m)} + \frac{37(2 + 3x)^{2+m}}{27(2 + m)} - \frac{10(2 + 3x)^{3+m}}{27(3 + m)}$$

output

```
-7*(2+3*x)^(1+m)/(27+27*m)+37*(2+3*x)^(2+m)/(54+27*m)-10*(2+3*x)^(3+m)/(81+27*m)
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.85

$$\int (1 - 2x)(2 + 3x)^m(3 + 5x) dx = \frac{1}{27}(2 + 3x)^{1+m} \left(-\frac{7}{1 + m} + \frac{37(2 + 3x)}{2 + m} - \frac{10(2 + 3x)^2}{3 + m} \right)$$

input

```
Integrate[(1 - 2*x)*(2 + 3*x)^m*(3 + 5*x), x]
```

output

```
((2 + 3*x)^(1 + m)*(-7/(1 + m) + (37*(2 + 3*x))/(2 + m) - (10*(2 + 3*x)^2)/(3 + m))/27
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (1 - 2x)(5x + 3)(3x + 2)^m dx$$

$$\downarrow 86$$

$$\int \left(-\frac{7}{9}(3x + 2)^m + \frac{37}{9}(3x + 2)^{m+1} - \frac{10}{9}(3x + 2)^{m+2} \right) dx$$

$$\downarrow 2009$$

$$-\frac{7(3x + 2)^{m+1}}{27(m + 1)} + \frac{37(3x + 2)^{m+2}}{27(m + 2)} - \frac{10(3x + 2)^{m+3}}{27(m + 3)}$$

input `Int[(1 - 2*x)*(2 + 3*x)^m*(3 + 5*x), x]`

output `(-7*(2 + 3*x)^(1 + m))/(27*(1 + m)) + (37*(2 + 3*x)^(2 + m))/(27*(2 + m)) - (10*(2 + 3*x)^(3 + m))/(27*(3 + m))`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3.

Time = 0.17 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.11

method	result
meijerg	$3 \cdot 2^m x \operatorname{hypergeom}\left(\left[1, -m\right], \left[2\right], -\frac{3x}{2}\right) - 2^{m-1} x^2 \operatorname{hypergeom}\left(\left[2, -m\right], \left[3\right], -\frac{3x}{2}\right) - \frac{5 \cdot 2^{1+m} x^3 \operatorname{hypergeom}\left(\left[3, -m\right], \left[4\right], -\frac{3x}{2}\right)}{27(m^3 + 6m^2 + 11m + 6)}$
gospers	$-\frac{(2+3x)^{1+m} (90m^2x^2+9m^2x+270m^2x^2-27m^2-84xm+180x^2-141m-93x-100)}{27(m^3+6m^2+11m+6)}$
orering	$\frac{(90m^2x^2+9m^2x+270m^2x^2-27m^2-84xm+180x^2-141m-93x-100)(2+3x)(1-2x)(2+3x)^m}{27(m^3+6m^2+11m+6)(-1+2x)}$
risch	$-\frac{(270m^2x^3+207m^2x^2+810m^2x^3-63m^2x+288m^2x^2+540x^3-54m^2-591xm+81x^2-282m-486x-200)(2+3x)^m}{27(2+m)(3+m)(1+m)}$
norman	$-\frac{10x^3 e^{m \ln(2+3x)}}{3+m} + \frac{2(27m^2+141m+100)e^{m \ln(2+3x)}}{27(m^3+6m^2+11m+6)} - \frac{(23m+9)x^2 e^{m \ln(2+3x)}}{3(m^2+5m+6)} + \frac{(21m^2+197m+162)x e^{m \ln(2+3x)}}{9m^3+54m^2+99m+54}$
parallelrisch	$-\frac{540x^3(2+3x)^m m^2+1620x^3(2+3x)^m m+414x^2(2+3x)^m m^2+1080(2+3x)^m x^3+576x^2(2+3x)^m m-126x(2+3x)^m m^2+162(2+3x)^m m^2}{54(m^3+6m^2+11m+6)}$

input `int((1-2*x)*(2+3*x)^m*(3+5*x),x,method=_RETURNVERBOSE)`

output `3*2^m*x*hypergeom([1, -m], [2], -3/2*x)-2^(m-1)*x^2*hypergeom([2, -m], [3], -3/2*x)-5/3*2^(1+m)*x^3*hypergeom([3, -m], [4], -3/2*x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.36

$$\int (1 - 2x)(2 + 3x)^m(3 + 5x) dx = -\frac{(270(m^2 + 3m + 2)x^3 + 9(23m^2 + 32m + 9)x^2 - 54m^2 - 3(21m^2 + 197m + 162)x - 282m - 200)(2+3x)^m}{27(m^3 + 6m^2 + 11m + 6)}$$

input `integrate((1-2*x)*(2+3*x)^m*(3+5*x),x, algorithm="fricas")`

output `-1/27*(270*(m^2 + 3*m + 2)*x^3 + 9*(23*m^2 + 32*m + 9)*x^2 - 54*m^2 - 3*(21*m^2 + 197*m + 162)*x - 282*m - 200)*(3*x + 2)^m/(m^3 + 6*m^2 + 11*m + 6)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 488 vs. $2(44) = 88$.

Time = 0.32 (sec) , antiderivative size = 488, normalized size of antiderivative = 8.87

$$\int (1 - 2x)(2 + 3x)^m(3 + 5x) dx$$

$$= \begin{cases} -\frac{180x^2 \log(3x+2)}{486x^2+648x+216} - \frac{240x \log(3x+2)}{486x^2+648x+216} - \frac{222x}{486x^2+648x+216} - \frac{80 \log(3x+2)}{486x^2+648x+216} - \frac{141}{486x^2+648x+216} \\ -\frac{90x^2}{81x+54} + \frac{111x \log(3x+2)}{81x+54} + \frac{74 \log(3x+2)}{81x+54} + \frac{47}{81x+54} \\ -\frac{5x^2}{3} + \frac{17x}{9} - \frac{7 \log(3x+2)}{27} \\ -\frac{270m^2x^3(3x+2)^m}{27m^3+162m^2+297m+162} - \frac{207m^2x^2(3x+2)^m}{27m^3+162m^2+297m+162} + \frac{63m^2x(3x+2)^m}{27m^3+162m^2+297m+162} + \frac{54m^2(3x+2)^m}{27m^3+162m^2+297m+162} - \frac{810m}{27m^3+162m^2+297m+162} \end{cases}$$

input `integrate((1-2*x)*(2+3*x)**m*(3+5*x), x)`

output `Piecewise((-180*x**2*log(3*x + 2)/(486*x**2 + 648*x + 216) - 240*x*log(3*x + 2)/(486*x**2 + 648*x + 216) - 222*x/(486*x**2 + 648*x + 216) - 80*log(3*x + 2)/(486*x**2 + 648*x + 216) - 141/(486*x**2 + 648*x + 216), Eq(m, -3)), (-90*x**2/(81*x + 54) + 111*x*log(3*x + 2)/(81*x + 54) + 74*log(3*x + 2)/(81*x + 54) + 47/(81*x + 54), Eq(m, -2)), (-5*x**2/3 + 17*x/9 - 7*log(3*x + 2)/27, Eq(m, -1)), (-270*m**2*x**3*(3*x + 2)**m/(27*m**3 + 162*m**2 + 297*m + 162) - 207*m**2*x**2*(3*x + 2)**m/(27*m**3 + 162*m**2 + 297*m + 162) + 63*m**2*x*(3*x + 2)**m/(27*m**3 + 162*m**2 + 297*m + 162) + 54*m**2*(3*x + 2)**m/(27*m**3 + 162*m**2 + 297*m + 162) - 810*m*x**3*(3*x + 2)**m/(27*m**3 + 162*m**2 + 297*m + 162) - 288*m*x**2*(3*x + 2)**m/(27*m**3 + 162*m**2 + 297*m + 162) + 591*m*x*(3*x + 2)**m/(27*m**3 + 162*m**2 + 297*m + 162) + 282*m*(3*x + 2)**m/(27*m**3 + 162*m**2 + 297*m + 162) - 540*x**3*(3*x + 2)**m/(27*m**3 + 162*m**2 + 297*m + 162) - 81*x**2*(3*x + 2)**m/(27*m**3 + 162*m**2 + 297*m + 162) + 486*x*(3*x + 2)**m/(27*m**3 + 162*m**2 + 297*m + 162) + 200*(3*x + 2)**m/(27*m**3 + 162*m**2 + 297*m + 162), True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 102 vs. $2(49) = 98$.

Time = 0.03 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.85

$$\int (1-2x)(2+3x)^m(3+5x) dx$$

$$= -\frac{10(27(m^2+3m+2)x^3+18(m^2+m)x^2-24mx+16)(3x+2)^m}{27(m^3+6m^2+11m+6)}$$

$$-\frac{(9(m+1)x^2+6mx-4)(3x+2)^m}{9(m^2+3m+2)} + \frac{(3x+2)^{m+1}}{m+1}$$

input `integrate((1-2*x)*(2+3*x)^m*(3+5*x),x, algorithm="maxima")`

output `-10/27*(27*(m^2 + 3*m + 2)*x^3 + 18*(m^2 + m)*x^2 - 24*m*x + 16)*(3*x + 2)^m/(m^3 + 6*m^2 + 11*m + 6) - 1/9*(9*(m + 1)*x^2 + 6*m*x - 4)*(3*x + 2)^m/(m^2 + 3*m + 2) + (3*x + 2)^(m + 1)/(m + 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 163 vs. $2(49) = 98$.

Time = 0.12 (sec) , antiderivative size = 163, normalized size of antiderivative = 2.96

$$\int (1-2x)(2+3x)^m(3+5x) dx =$$

$$-\frac{270m^2(3x+2)^m x^3 + 207m^2(3x+2)^m x^2 + 810m(3x+2)^m x^3 - 63m^2(3x+2)^m x + 288m(3x+2)^m}{(m^3 + 6m^2 + 11m + 6)}$$

input `integrate((1-2*x)*(2+3*x)^m*(3+5*x),x, algorithm="giac")`

output `-1/27*(270*m^2*(3*x + 2)^m*x^3 + 207*m^2*(3*x + 2)^m*x^2 + 810*m*(3*x + 2)^m*x^3 - 63*m^2*(3*x + 2)^m*x + 288*m*(3*x + 2)^m*x^2 + 540*(3*x + 2)^m*x^3 - 54*m^2*(3*x + 2)^m - 591*m*(3*x + 2)^m*x + 81*(3*x + 2)^m*x^2 - 282*m*(3*x + 2)^m - 486*(3*x + 2)^m*x - 200*(3*x + 2)^m)/(m^3 + 6*m^2 + 11*m + 6)`

Mupad [B] (verification not implemented)

Time = 1.17 (sec) , antiderivative size = 130, normalized size of antiderivative = 2.36

$$\int (1-2x)(2+3x)^m(3+5x) dx = (3x+2)^m \left(\frac{54m^2 + 282m + 200}{27m^3 + 162m^2 + 297m + 162} + \frac{x(63m^2 + 591m + 486)}{27m^3 + 162m^2 + 297m + 162} - \frac{x^2(207m^2 + 288m + 81)}{27m^3 + 162m^2 + 297m + 162} - \frac{x^3(270m^2 + 810m + 540)}{27m^3 + 162m^2 + 297m + 162} \right)$$

input `int(-(2*x - 1)*(3*x + 2)^m*(5*x + 3), x)`output `(3*x + 2)^m*((282*m + 54*m^2 + 200)/(297*m + 162*m^2 + 27*m^3 + 162) + (x*(591*m + 63*m^2 + 486))/(297*m + 162*m^2 + 27*m^3 + 162) - (x^2*(288*m + 207*m^2 + 81))/(297*m + 162*m^2 + 27*m^3 + 162) - (x^3*(810*m + 270*m^2 + 540))/(297*m + 162*m^2 + 27*m^3 + 162))`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.56

$$\int (1-2x)(2+3x)^m(3+5x) dx = \frac{(3x+2)^m(-270m^2x^3 - 207m^2x^2 - 810mx^3 + 63m^2x - 288mx^2 - 540x^3 + 54m^2 + 591mx - 81x^2 + 200)}{27m^3 + 162m^2 + 297m + 162}$$

input `int((1-2*x)*(2+3*x)^m*(3+5*x), x)`output `((3*x + 2)**m*(- 270*m**2*x**3 - 207*m**2*x**2 + 63*m**2*x + 54*m**2 - 810*m*x**3 - 288*m*x**2 + 591*m*x + 282*m - 540*x**3 - 81*x**2 + 486*x + 200))/(27*(m**3 + 6*m**2 + 11*m + 6))`

3.240 $\int \frac{(1-2x)(2+3x)^m}{3+5x} dx$

Optimal result	2141
Mathematica [A] (verified)	2141
Rubi [A] (verified)	2142
Maple [F]	2143
Fricas [F]	2143
Sympy [F]	2143
Maxima [F]	2144
Giac [F]	2144
Mupad [F(-1)]	2144
Reduce [F]	2145

Optimal result

Integrand size = 20, antiderivative size = 52

$$\int \frac{(1-2x)(2+3x)^m}{3+5x} dx = -\frac{2(2+3x)^{1+m}}{15(1+m)} - \frac{11(2+3x)^{1+m} \text{Hypergeometric2F1}(1, 1+m, 2+m, 5(2+3x))}{5(1+m)}$$

output

```
-2*(2+3*x)^(1+m)/(15+15*m)-11*(2+3*x)^(1+m)*hypergeom([1, 1+m], [2+m], 10+15*x)/(5+5*m)
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.71

$$\int \frac{(1-2x)(2+3x)^m}{3+5x} dx = -\frac{(2+3x)^{1+m}(2+33 \text{Hypergeometric2F1}(1, 1+m, 2+m, 5(2+3x)))}{15(1+m)}$$

input

```
Integrate[((1 - 2*x)*(2 + 3*x)^m)/(3 + 5*x), x]
```

output

$$-1/15*((2 + 3*x)^(1 + m)*(2 + 33*Hypergeometric2F1[1, 1 + m, 2 + m, 5*(2 + 3*x)])))/(1 + m)$$

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {90, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1 - 2x)(3x + 2)^m}{5x + 3} dx$$

↓ 90

$$\frac{11}{5} \int \frac{(3x + 2)^m}{5x + 3} dx - \frac{2(3x + 2)^{m+1}}{15(m + 1)}$$

↓ 78

$$-\frac{11(3x + 2)^{m+1} \text{Hypergeometric2F1}(1, m + 1, m + 2, 5(3x + 2))}{5(m + 1)} - \frac{2(3x + 2)^{m+1}}{15(m + 1)}$$

input

$$\text{Int}[(1 - 2*x)*(2 + 3*x)^m/(3 + 5*x), x]$$

output

$$(-2*(2 + 3*x)^(1 + m))/(15*(1 + m)) - (11*(2 + 3*x)^(1 + m)*Hypergeometric2F1[1, 1 + m, 2 + m, 5*(2 + 3*x)])/(5*(1 + m))$$

Defintions of rubi rules used

rule 78

$$\text{Int}[(a_ + (b_)*(x_))^(m_)*((c_ + (d_)*(x_))^(n_), x_Symbol] \rightarrow \text{Simp}[(b *c - a*d)^(n+1)*(a + b*x)^(m+1)/(b^(n+1)*(m+1))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(a + b*x)/(b*c - a*d)], x] /; \text{FreeQ}\{a, b, c, d, m\}, x] \&\& \text{IntegerQ}[m] \&\& \text{IntegerQ}[n]$$

rule 90

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p
_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))),
x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p
+ 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n,
p}, x] && NeQ[n + p + 2, 0]
```

Maple [F]

$$\int \frac{(1 - 2x)(2 + 3x)^m}{3 + 5x} dx$$

input

```
int((1-2*x)*(2+3*x)^m/(3+5*x),x)
```

output

```
int((1-2*x)*(2+3*x)^m/(3+5*x),x)
```

Fricas [F]

$$\int \frac{(1 - 2x)(2 + 3x)^m}{3 + 5x} dx = \int -\frac{(3x + 2)^m(2x - 1)}{5x + 3} dx$$

input

```
integrate((1-2*x)*(2+3*x)^m/(3+5*x),x, algorithm="fricas")
```

output

```
integral(-(3*x + 2)^m*(2*x - 1)/(5*x + 3), x)
```

Sympy [F]

$$\int \frac{(1 - 2x)(2 + 3x)^m}{3 + 5x} dx = - \int \left(-\frac{(3x + 2)^m}{5x + 3} \right) dx - \int \frac{2x(3x + 2)^m}{5x + 3} dx$$

input

```
integrate((1-2*x)*(2+3*x)**m/(3+5*x),x)
```

output `-Integral(-(3*x + 2)**m/(5*x + 3), x) - Integral(2*x*(3*x + 2)**m/(5*x + 3), x)`

Maxima [F]

$$\int \frac{(1 - 2x)(2 + 3x)^m}{3 + 5x} dx = \int -\frac{(3x + 2)^m(2x - 1)}{5x + 3} dx$$

input `integrate((1-2*x)*(2+3*x)^m/(3+5*x),x, algorithm="maxima")`

output `-integrate((3*x + 2)^m*(2*x - 1)/(5*x + 3), x)`

Giac [F]

$$\int \frac{(1 - 2x)(2 + 3x)^m}{3 + 5x} dx = \int -\frac{(3x + 2)^m(2x - 1)}{5x + 3} dx$$

input `integrate((1-2*x)*(2+3*x)^m/(3+5*x),x, algorithm="giac")`

output `integrate(-(3*x + 2)^m*(2*x - 1)/(5*x + 3), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(1 - 2x)(2 + 3x)^m}{3 + 5x} dx = -\int \frac{(2x - 1)(3x + 2)^m}{5x + 3} dx$$

input `int(-((2*x - 1)*(3*x + 2)^m)/(5*x + 3),x)`

output `-int(((2*x - 1)*(3*x + 2)^m)/(5*x + 3), x)`

Reduce [F]

$$\int \frac{(1-2x)(2+3x)^m}{3+5x} dx$$

$$= \frac{-18(3x+2)^m mx + 10(3x+2)^m m + 22(3x+2)^m - 33 \left(\int \frac{(3x+2)^m x}{15x^2+19x+6} dx \right) m^2 - 33 \left(\int \frac{(3x+2)^m x}{15x^2+19x+6} dx \right) m}{45m(m+1)}$$

input `int((1-2*x)*(2+3*x)^m/(3+5*x),x)`

output `(- 18*(3*x + 2)**m*m*x + 10*(3*x + 2)**m*m + 22*(3*x + 2)**m - 33*int(((3*x + 2)**m*x)/(15*x**2 + 19*x + 6),x)*m**2 - 33*int(((3*x + 2)**m*x)/(15*x**2 + 19*x + 6),x)*m)/(45*m*(m + 1))`

3.241 $\int \frac{(1-2x)(2+3x)^m}{(3+5x)^2} dx$

Optimal result	2146
Mathematica [A] (verified)	2146
Rubi [A] (verified)	2147
Maple [F]	2148
Fricas [F]	2148
Sympy [F]	2149
Maxima [F]	2149
Giac [F]	2149
Mupad [F(-1)]	2150
Reduce [F]	2150

Optimal result

Integrand size = 20, antiderivative size = 65

$$\int \frac{(1-2x)(2+3x)^m}{(3+5x)^2} dx$$

$$= -\frac{2(2+3x)^{1+m}}{15m(3+5x)} - \frac{(2-33m)(2+3x)^{1+m} \operatorname{Hypergeometric2F1}(2, 1+m, 2+m, 5(2+3x))}{5m(1+m)}$$

output

```
-2/15*(2+3*x)^(1+m)/m/(3+5*x)-1/5*(2-33*m)*(2+3*x)^(1+m)*hypergeom([2, 1+m], [2+m], 10+15*x)/m/(1+m)
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.88

$$\int \frac{(1-2x)(2+3x)^m}{(3+5x)^2} dx = -\frac{(2+3x)^{1+m}(11(1+m) + (-2+33m)(3+5x) \operatorname{Hypergeometric2F1}(1, 1+m, 2+m, 5(2+3x)))}{5(1+m)(3+5x)}$$

input

```
Integrate[((1 - 2*x)*(2 + 3*x)^m)/(3 + 5*x)^2,x]
```

output

```
-1/5*((2 + 3*x)^(1 + m)*(11*(1 + m) + (-2 + 33*m)*(3 + 5*x)*Hypergeometric
2F1[1, 1 + m, 2 + m, 5*(2 + 3*x)])))/((1 + m)*(3 + 5*x))
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.91, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {87, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1-2x)(3x+2)^m}{(5x+3)^2} dx$$

$$\downarrow 87$$

$$-\frac{1}{5}(2-33m) \int \frac{(3x+2)^m}{5x+3} dx - \frac{11(3x+2)^{m+1}}{5(5x+3)}$$

$$\downarrow 78$$

$$\frac{(2-33m)(3x+2)^{m+1} \text{Hypergeometric2F1}(1, m+1, m+2, 5(3x+2))}{5(m+1)} - \frac{11(3x+2)^{m+1}}{5(5x+3)}$$

input

```
Int[((1 - 2*x)*(2 + 3*x)^m)/(3 + 5*x)^2,x]
```

output

```
(-11*(2 + 3*x)^(1 + m))/(5*(3 + 5*x)) + ((2 - 33*m)*(2 + 3*x)^(1 + m)*Hypergeometric2F1[1, 1 + m, 2 + m, 5*(2 + 3*x)])/(5*(1 + m))
```


Definitions of rubi rules used

rule 78

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] :> Simp[(b
*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m
+ 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x]
&& !IntegerQ[m] && IntegerQ[n]
```

rule 87

```
Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p
_), x_] :> Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

Maple [F]

$$\int \frac{(1-2x)(2+3x)^m}{(3+5x)^2} dx$$

input

```
int((1-2*x)*(2+3*x)^m/(3+5*x)^2,x)
```

output

```
int((1-2*x)*(2+3*x)^m/(3+5*x)^2,x)
```

Fricas [F]

$$\int \frac{(1-2x)(2+3x)^m}{(3+5x)^2} dx = \int -\frac{(3x+2)^m(2x-1)}{(5x+3)^2} dx$$

input

```
integrate((1-2*x)*(2+3*x)^m/(3+5*x)^2,x, algorithm="fricas")
```

output

```
integral(-(3*x + 2)^m*(2*x - 1)/(25*x^2 + 30*x + 9), x)
```

Sympy [F]

$$\int \frac{(1-2x)(2+3x)^m}{(3+5x)^2} dx = -\int \left(-\frac{(3x+2)^m}{25x^2+30x+9} \right) dx - \int \frac{2x(3x+2)^m}{25x^2+30x+9} dx$$

input `integrate((1-2*x)*(2+3*x)**m/(3+5*x)**2,x)`

output `-Integral(-(3*x + 2)**m/(25*x**2 + 30*x + 9), x) - Integral(2*x*(3*x + 2)**m/(25*x**2 + 30*x + 9), x)`

Maxima [F]

$$\int \frac{(1-2x)(2+3x)^m}{(3+5x)^2} dx = \int -\frac{(3x+2)^m(2x-1)}{(5x+3)^2} dx$$

input `integrate((1-2*x)*(2+3*x)^m/(3+5*x)^2,x, algorithm="maxima")`

output `-integrate((3*x + 2)^m*(2*x - 1)/(5*x + 3)^2, x)`

Giac [F]

$$\int \frac{(1-2x)(2+3x)^m}{(3+5x)^2} dx = \int -\frac{(3x+2)^m(2x-1)}{(5x+3)^2} dx$$

input `integrate((1-2*x)*(2+3*x)^m/(3+5*x)^2,x, algorithm="giac")`

output `integrate(-(3*x + 2)^m*(2*x - 1)/(5*x + 3)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(1-2x)(2+3x)^m}{(3+5x)^2} dx = - \int \frac{(2x-1)(3x+2)^m}{(5x+3)^2} dx$$

input `int(-((2*x - 1)*(3*x + 2)^m)/(5*x + 3)^2,x)`output `-int(((2*x - 1)*(3*x + 2)^m)/(5*x + 3)^2, x)`**Reduce [F]**

$$\int \frac{(1-2x)(2+3x)^m}{(3+5x)^2} dx$$

$$= \frac{-18(3x+2)^m mx + 10(3x+2)^m m + 20(3x+2)^m x + 12(3x+2)^m - 1485 \left(\int \frac{(3x+2)^m}{675m x^3 + 1260m x^2 - 750x^3 + 783} \right)}{675m x^3 + 1260m x^2 - 750x^3 + 783}$$

input `int(((1-2*x)*(2+3*x)^m)/(3+5*x)^2,x)`output `(- 18*(3*x + 2)**m*x + 10*(3*x + 2)**m*m + 20*(3*x + 2)**m*x + 12*(3*x + 2)**m - 1485*int(((3*x + 2)**m*x)/(675*m*x**3 + 1260*m*x**2 + 783*m*x + 162*m - 750*x**3 - 1400*x**2 - 870*x - 180),x)*m**3*x - 891*int(((3*x + 2)**m*x)/(675*m*x**3 + 1260*m*x**2 + 783*m*x + 162*m - 750*x**3 - 1400*x**2 - 870*x - 180),x)*m**3 + 1740*int(((3*x + 2)**m*x)/(675*m*x**3 + 1260*m*x**2 + 783*m*x + 162*m - 750*x**3 - 1400*x**2 - 870*x - 180),x)*m**2*x + 1044*int(((3*x + 2)**m*x)/(675*m*x**3 + 1260*m*x**2 + 783*m*x + 162*m - 750*x**3 - 1400*x**2 - 870*x - 180),x)*m**2 - 100*int(((3*x + 2)**m*x)/(675*m*x**3 + 1260*m*x**2 + 783*m*x + 162*m - 750*x**3 - 1400*x**2 - 870*x - 180),x)*m*x - 60*int(((3*x + 2)**m*x)/(675*m*x**3 + 1260*m*x**2 + 783*m*x + 162*m - 750*x**3 - 1400*x**2 - 870*x - 180),x)*m)/(5*m*(45*m*x + 27*m - 50*x - 30))`

3.242 $\int \frac{(1-2x)(2+3x)^m}{(3+5x)^3} dx$

Optimal result	2151
Mathematica [A] (verified)	2151
Rubi [A] (verified)	2152
Maple [F]	2153
Fricas [F]	2153
Sympy [F]	2154
Maxima [F]	2154
Giac [F]	2154
Mupad [F(-1)]	2155
Reduce [F]	2155

Optimal result

Integrand size = 20, antiderivative size = 70

$$\int \frac{(1-2x)(2+3x)^m}{(3+5x)^3} dx$$

$$= \frac{2(2+3x)^{1+m}}{15(1-m)(3+5x)^2} - \frac{3(37-33m)(2+3x)^{1+m} \operatorname{Hypergeometric2F1}(3, 1+m, 2+m, 5(2+3x))}{5(1-m^2)}$$

output 2/15*(2+3*x)^(1+m)/(1-m)/(3+5*x)^2-3*(37-33*m)*(2+3*x)^(1+m)*hypergeom([3, 1+m],[2+m],10+15*x)/(-5*m^2+5)

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.69

$$\int \frac{(1-2x)(2+3x)^m}{(3+5x)^3} dx$$

$$= \frac{1}{10}(2+3x)^{1+m} \left(-\frac{11}{(3+5x)^2} + \frac{3(-37+33m) \operatorname{Hypergeometric2F1}(2, 1+m, 2+m, 10+15x)}{1+m} \right)$$

input `Integrate[((1 - 2*x)*(2 + 3*x)^m)/(3 + 5*x)^3,x]`

output `((2 + 3*x)^(1 + m)*(-11/(3 + 5*x)^2 + (3*(-37 + 33*m)*Hypergeometric2F1[2, 1 + m, 2 + m, 10 + 15*x])/(1 + m)))/10`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.84, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {87, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1 - 2x)(3x + 2)^m}{(5x + 3)^3} dx$$

$$\downarrow 87$$

$$-\frac{1}{10}(37 - 33m) \int \frac{(3x + 2)^m}{(5x + 3)^2} dx - \frac{11(3x + 2)^{m+1}}{10(5x + 3)^2}$$

$$\downarrow 78$$

$$-\frac{3(37 - 33m)(3x + 2)^{m+1} \text{Hypergeometric2F1}(2, m + 1, m + 2, 5(3x + 2))}{10(m + 1)} - \frac{11(3x + 2)^{m+1}}{10(5x + 3)^2}$$

input `Int[((1 - 2*x)*(2 + 3*x)^m)/(3 + 5*x)^3,x]`

output `(-11*(2 + 3*x)^(1 + m))/(10*(3 + 5*x)^2) - (3*(37 - 33*m)*(2 + 3*x)^(1 + m)*Hypergeometric2F1[2, 1 + m, 2 + m, 5*(2 + 3*x)])/(10*(1 + m))`

Definitions of rubi rules used

rule 78 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

Maple [F]

$$\int \frac{(1 - 2x)(2 + 3x)^m}{(3 + 5x)^3} dx$$

input `int((1-2*x)*(2+3*x)^m/(3+5*x)^3,x)`

output `int((1-2*x)*(2+3*x)^m/(3+5*x)^3,x)`

Fricas [F]

$$\int \frac{(1 - 2x)(2 + 3x)^m}{(3 + 5x)^3} dx = \int -\frac{(3x + 2)^m(2x - 1)}{(5x + 3)^3} dx$$

input `integrate((1-2*x)*(2+3*x)^m/(3+5*x)^3,x, algorithm="fricas")`

output `integral(-(3*x + 2)^m*(2*x - 1)/(125*x^3 + 225*x^2 + 135*x + 27), x)`

Sympy [F]

$$\int \frac{(1-2x)(2+3x)^m}{(3+5x)^3} dx = - \int \left(-\frac{(3x+2)^m}{125x^3 + 225x^2 + 135x + 27} \right) dx - \int \frac{2x(3x+2)^m}{125x^3 + 225x^2 + 135x + 27} dx$$

input `integrate((1-2*x)*(2+3*x)**m/(3+5*x)**3,x)`

output `-Integral(-(3*x + 2)**m/(125*x**3 + 225*x**2 + 135*x + 27), x) - Integral(2*x*(3*x + 2)**m/(125*x**3 + 225*x**2 + 135*x + 27), x)`

Maxima [F]

$$\int \frac{(1-2x)(2+3x)^m}{(3+5x)^3} dx = \int -\frac{(3x+2)^m(2x-1)}{(5x+3)^3} dx$$

input `integrate((1-2*x)*(2+3*x)^m/(3+5*x)^3,x, algorithm="maxima")`

output `-integrate((3*x + 2)^m*(2*x - 1)/(5*x + 3)^3, x)`

Giac [F]

$$\int \frac{(1-2x)(2+3x)^m}{(3+5x)^3} dx = \int -\frac{(3x+2)^m(2x-1)}{(5x+3)^3} dx$$

input `integrate((1-2*x)*(2+3*x)^m/(3+5*x)^3,x, algorithm="giac")`

output `integrate(-(3*x + 2)^m*(2*x - 1)/(5*x + 3)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(1-2x)(2+3x)^m}{(3+5x)^3} dx = - \int \frac{(2x-1)(3x+2)^m}{(5x+3)^3} dx$$

input `int(-((2*x - 1)*(3*x + 2)^m)/(5*x + 3)^3,x)`output `-int(((2*x - 1)*(3*x + 2)^m)/(5*x + 3)^3, x)`**Reduce [F]**

$$\int \frac{(1-2x)(2+3x)^m}{(3+5x)^3} dx = \text{Too large to display}$$

input `int((1-2*x)*(2+3*x)^m/(3+5*x)^3,x)`

output

```
( - 18*(3*x + 2)**m*m*x + 10*(3*x + 2)**m*m + 40*(3*x + 2)**m*x + 2*(3*x +
2)**m - 7425*int(((3*x + 2)**m*x)/(3375*m**2*x**4 + 8325*m**2*x**3 + 7695
*m**2*x**2 + 3159*m**2*x + 486*m**2 - 10875*m*x**4 - 26825*m*x**3 - 24795*
m*x**2 - 10179*m*x - 1566*m + 7500*x**4 + 18500*x**3 + 17100*x**2 + 7020*x
+ 1080),x)*m**4*x**2 - 8910*int(((3*x + 2)**m*x)/(3375*m**2*x**4 + 8325*m
**2*x**3 + 7695*m**2*x**2 + 3159*m**2*x + 486*m**2 - 10875*m*x**4 - 26825*
m*x**3 - 24795*m*x**2 - 10179*m*x - 1566*m + 7500*x**4 + 18500*x**3 + 1710
0*x**2 + 7020*x + 1080),x)*m**4*x - 2673*int(((3*x + 2)**m*x)/(3375*m**2*x
**4 + 8325*m**2*x**3 + 7695*m**2*x**2 + 3159*m**2*x + 486*m**2 - 10875*m*x
**4 - 26825*m*x**3 - 24795*m*x**2 - 10179*m*x - 1566*m + 7500*x**4 + 18500
*x**3 + 17100*x**2 + 7020*x + 1080),x)*m**4 + 32250*int(((3*x + 2)**m*x)/(
3375*m**2*x**4 + 8325*m**2*x**3 + 7695*m**2*x**2 + 3159*m**2*x + 486*m**2
- 10875*m*x**4 - 26825*m*x**3 - 24795*m*x**2 - 10179*m*x - 1566*m + 7500*x
**4 + 18500*x**3 + 17100*x**2 + 7020*x + 1080),x)*m**3*x**2 + 38700*int(((
3*x + 2)**m*x)/(3375*m**2*x**4 + 8325*m**2*x**3 + 7695*m**2*x**2 + 3159*m*
*2*x + 486*m**2 - 10875*m*x**4 - 26825*m*x**3 - 24795*m*x**2 - 10179*m*x -
1566*m + 7500*x**4 + 18500*x**3 + 17100*x**2 + 7020*x + 1080),x)*m**3*x +
11610*int(((3*x + 2)**m*x)/(3375*m**2*x**4 + 8325*m**2*x**3 + 7695*m**2*x
**2 + 3159*m**2*x + 486*m**2 - 10875*m*x**4 - 26825*m*x**3 - 24795*m*x**2
- 10179*m*x - 1566*m + 7500*x**4 + 18500*x**3 + 17100*x**2 + 7020*x + 1...
```

3.243 $\int \frac{(c+dx)^{1+m}(e+fx)}{(a+bx)^2} dx$

Optimal result	2157
Mathematica [A] (verified)	2157
Rubi [A] (verified)	2158
Maple [F]	2159
Fricas [F]	2159
Sympy [F]	2160
Maxima [F]	2160
Giac [F]	2160
Mupad [F(-1)]	2161
Reduce [F]	2161

Optimal result

Integrand size = 22, antiderivative size = 108

$$\int \frac{(c+dx)^{1+m}(e+fx)}{(a+bx)^2} dx = \frac{f(c+dx)^{2+m}}{bd(1+m)(a+bx)} + \frac{(bcf+bde(1+m)-adf(2+m))(c+dx)^{2+m} \operatorname{Hypergeometric2F1}\left(2, 2+m, 3+m, \frac{b(c+dx)}{bc-ad}\right)}{b(bc-ad)^2(1+m)(2+m)}$$

output

```
f*(d*x+c)^(2+m)/b/d/(1+m)/(b*x+a)+(b*c*f+b*d*e*(1+m)-a*d*f*(2+m))*(d*x+c)^(2+m)*hypergeom([2, 2+m], [3+m], b*(d*x+c)/(-a*d+b*c))/b/(-a*d+b*c)^2/(1+m)/(2+m)
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.94

$$\int \frac{(c+dx)^{1+m}(e+fx)}{(a+bx)^2} dx = \frac{(c+dx)^{2+m} \left(-\frac{(bc-ad)(be-af)}{a+bx} - \frac{(bcf+bde(1+m)-adf(2+m)) \operatorname{Hypergeometric2F1}\left(1, 2+m, 3+m, \frac{b(c+dx)}{bc-ad}\right)}{2+m} \right)}{b(bc-ad)^2}$$

input `Integrate[((c + d*x)^(1 + m)*(e + f*x))/(a + b*x)^2,x]`

output `((c + d*x)^(2 + m)*(-(((b*c - a*d)*(b*e - a*f))/(a + b*x)) - ((b*c*f + b*d*e*(1 + m) - a*d*f*(2 + m))*Hypergeometric2F1[1, 2 + m, 3 + m, (b*(c + d*x))/(b*c - a*d)])/(2 + m)))/(b*(b*c - a*d)^2)`

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.06, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {87, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(e + fx)(c + dx)^{m+1}}{(a + bx)^2} dx$$

$$\downarrow 87$$

$$\frac{(-adf(m + 2) + bcf + bde(m + 1)) \int \frac{(c+dx)^{m+1}}{a+bx} dx}{b(bc - ad)} - \frac{(be - af)(c + dx)^{m+2}}{b(a + bx)(bc - ad)}$$

$$\downarrow 78$$

$$\frac{(c + dx)^{m+2}(-adf(m + 2) + bcf + bde(m + 1)) \text{Hypergeometric2F1}\left(1, m + 2, m + 3, \frac{b(c+dx)}{bc-ad}\right)}{\frac{b(m + 2)(bc - ad)^2}{(be - af)(c + dx)^{m+2}} b(a + bx)(bc - ad)}$$

input `Int[((c + d*x)^(1 + m)*(e + f*x))/(a + b*x)^2,x]`

output `-(((b*e - a*f)*(c + d*x)^(2 + m))/(b*(b*c - a*d)*(a + b*x))) - ((b*c*f + b*d*e*(1 + m) - a*d*f*(2 + m))*(c + d*x)^(2 + m)*Hypergeometric2F1[1, 2 + m, 3 + m, (b*(c + d*x))/(b*c - a*d)])/(b*(b*c - a*d)^2*(2 + m))`

Definitions of rubi rules used

rule 78 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

Maple [F]

$$\int \frac{(xd + c)^{1+m} (fx + e)}{(bx + a)^2} dx$$

input `int((d*x+c)^(1+m)*(f*x+e)/(b*x+a)^2,x)`

output `int((d*x+c)^(1+m)*(f*x+e)/(b*x+a)^2,x)`

Fricas [F]

$$\int \frac{(c + dx)^{1+m} (e + fx)}{(a + bx)^2} dx = \int \frac{(fx + e)(dx + c)^{m+1}}{(bx + a)^2} dx$$

input `integrate((d*x+c)^(1+m)*(f*x+e)/(b*x+a)^2,x, algorithm="fricas")`

output `integral((f*x + e)*(d*x + c)^(m + 1)/(b^2*x^2 + 2*a*b*x + a^2), x)`

Sympy [F]

$$\int \frac{(c + dx)^{1+m}(e + fx)}{(a + bx)^2} dx = \int \frac{(c + dx)^{m+1}(e + fx)}{(a + bx)^2} dx$$

input `integrate((d*x+c)**(1+m)*(f*x+e)/(b*x+a)**2,x)`

output `Integral((c + d*x)**(m + 1)*(e + f*x)/(a + b*x)**2, x)`

Maxima [F]

$$\int \frac{(c + dx)^{1+m}(e + fx)}{(a + bx)^2} dx = \int \frac{(fx + e)(dx + c)^{m+1}}{(bx + a)^2} dx$$

input `integrate((d*x+c)^(1+m)*(f*x+e)/(b*x+a)^2,x, algorithm="maxima")`

output `integrate((f*x + e)*(d*x + c)^(m + 1)/(b*x + a)^2, x)`

Giac [F]

$$\int \frac{(c + dx)^{1+m}(e + fx)}{(a + bx)^2} dx = \int \frac{(fx + e)(dx + c)^{m+1}}{(bx + a)^2} dx$$

input `integrate((d*x+c)^(1+m)*(f*x+e)/(b*x+a)^2,x, algorithm="giac")`

output `integrate((f*x + e)*(d*x + c)^(m + 1)/(b*x + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^{1+m}(e + fx)}{(a + bx)^2} dx = \int \frac{(e + fx)(c + dx)^{m+1}}{(a + bx)^2} dx$$

input `int(((e + f*x)*(c + d*x)^(m + 1))/(a + b*x)^2,x)`output `int(((e + f*x)*(c + d*x)^(m + 1))/(a + b*x)^2, x)`**Reduce [F]**

$$\int \frac{(c + dx)^{1+m}(e + fx)}{(a + bx)^2} dx = \text{too large to display}$$

input `int((d*x+c)^(1+m)*(f*x+e)/(b*x+a)^2,x)`

output

```

((c + d*x)**m*a**2*c*d*f*m + 2*(c + d*x)**m*a**2*c*d*f - (c + d*x)**m*a**2
*d**2*f*m**2*x - 2*(c + d*x)**m*a**2*d**2*f*m*x - 2*(c + d*x)**m*a*b*c**2*
f*m - (c + d*x)**m*a*b*c**2*f - (c + d*x)**m*a*b*c*d*e*m - (c + d*x)**m*a*
b*c*d*e + 2*(c + d*x)**m*a*b*c*d*f*m**2*x + 2*(c + d*x)**m*a*b*c*d*f*m*x +
2*(c + d*x)**m*a*b*c*d*f*x + (c + d*x)**m*a*b*d**2*e*m**2*x + (c + d*x)**
m*a*b*d**2*e*m*x + (c + d*x)**m*a*b*d**2*f*m**2*x**2 + (c + d*x)**m*b**2*c
**2*e*m**2 + (c + d*x)**m*b**2*c**2*e*m - 2*(c + d*x)**m*b**2*c**2*f*m*x -
(c + d*x)**m*b**2*c**2*f*x - (c + d*x)**m*b**2*c*d*e*m*x - (c + d*x)**m*b
**2*c*d*e*x - (c + d*x)**m*b**2*c*d*f*m*x**2 + int(((c + d*x)**m*x)/(a**3*
c*d*m + a**3*d**2*m*x - a**2*b*c**2 + 2*a**2*b*c*d*m*x - a**2*b*c*d*x + 2*
a**2*b*d**2*m*x**2 - 2*a*b**2*c**2*x + a*b**2*c*d*m*x**2 - 2*a*b**2*c*d*x*
*2 + a*b**2*d**2*m*x**3 - b**3*c**2*x**2 - b**3*c*d*x**3),x)*a**5*d**4*f*m
**4 + 3*int(((c + d*x)**m*x)/(a**3*c*d*m + a**3*d**2*m*x - a**2*b*c**2 + 2
*a**2*b*c*d*m*x - a**2*b*c*d*x + 2*a**2*b*d**2*m*x**2 - 2*a*b**2*c**2*x +
a*b**2*c*d*m*x**2 - 2*a*b**2*c*d*x**2 + a*b**2*d**2*m*x**3 - b**3*c**2*x**
2 - b**3*c*d*x**3),x)*a**5*d**4*f*m**3 + 2*int(((c + d*x)**m*x)/(a**3*c*d*
m + a**3*d**2*m*x - a**2*b*c**2 + 2*a**2*b*c*d*m*x - a**2*b*c*d*x + 2*a**2
*b*d**2*m*x**2 - 2*a*b**2*c**2*x + a*b**2*c*d*m*x**2 - 2*a*b**2*c*d*x**2 +
a*b**2*d**2*m*x**3 - b**3*c**2*x**2 - b**3*c*d*x**3),x)*a**5*d**4*f*m**2
- 2*int(((c + d*x)**m*x)/(a**3*c*d*m + a**3*d**2*m*x - a**2*b*c**2 + 2*...

```

3.244 $\int \frac{(c+dx)^m(e+fx)}{(a+bx)^2} dx$

Optimal result	2163
Mathematica [A] (verified)	2163
Rubi [A] (verified)	2164
Maple [F]	2165
Fricas [F]	2165
Sympy [F]	2166
Maxima [F]	2166
Giac [F]	2166
Mupad [F(-1)]	2167
Reduce [F]	2167

Optimal result

Integrand size = 20, antiderivative size = 104

$$\int \frac{(c+dx)^m(e+fx)}{(a+bx)^2} dx = \frac{f(c+dx)^{1+m}}{b d m (a+bx)} - \frac{(a d f (1+m) - b (c f + d e m))(c+dx)^{1+m} \operatorname{Hypergeometric2F1}\left(2, 1+m, 2+m, \frac{b(c+dx)}{bc-ad}\right)}{b(bc-ad)^2 m (1+m)}$$

output `f*(d*x+c)^(1+m)/b/d/m/(b*x+a)-(a*d*f*(1+m)-b*(d*e*m+c*f))*(d*x+c)^(1+m)*hypergeom([2, 1+m], [2+m], b*(d*x+c)/(-a*d+b*c))/b/(-a*d+b*c)^2/m/(1+m)`

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.95

$$\int \frac{(c+dx)^m(e+fx)}{(a+bx)^2} dx = \frac{(c+dx)^{1+m} \left(-\frac{(bc-ad)(be-af)}{a+bx} + \frac{(adf(1+m)-b(cf+dem)) \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, \frac{b(c+dx)}{bc-ad}\right)}{1+m} \right)}{b(bc-ad)^2}$$

input `Integrate[((c + d*x)^m*(e + f*x))/(a + b*x)^2,x]`

output

$$\frac{((c + dx)^{(1 + m)} * (-((b*c - a*d)*(b*e - a*f))/(a + b*x)) + ((a*d*f*(1 + m) - b*(c*f + d*e*m)) * \text{Hypergeometric2F1}[1, 1 + m, 2 + m, (b*(c + d*x))/(b*c - a*d]]) / (1 + m))}{b*(b*c - a*d)^2}$$
Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.08, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {87, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(e + fx)(c + dx)^m}{(a + bx)^2} dx$$

$$\downarrow 87$$

$$-\frac{(adf(m + 1) - b(cf + dem)) \int \frac{(c+dx)^m}{a+bx} dx}{b(bc - ad)} - \frac{(be - af)(c + dx)^{m+1}}{b(a + bx)(bc - ad)}$$

$$\downarrow 78$$

$$\frac{(c + dx)^{m+1}(adf(m + 1) - b(cf + dem)) \text{Hypergeometric2F1}\left(1, m + 1, m + 2, \frac{b(c+dx)}{bc-ad}\right)}{b(m + 1)(bc - ad)^2} - \frac{(be - af)(c + dx)^{m+1}}{b(a + bx)(bc - ad)}$$

input

$$\text{Int}[(c + d*x)^m*(e + f*x)/(a + b*x)^2, x]$$

output

$$-(((b*e - a*f)*(c + d*x)^{(1 + m)})/(b*(b*c - a*d)*(a + b*x))) + ((a*d*f*(1 + m) - b*(c*f + d*e*m))*(c + d*x)^{(1 + m)}*\text{Hypergeometric2F1}[1, 1 + m, 2 + m, (b*(c + d*x))/(b*c - a*d)]/(b*(b*c - a*d)^2*(1 + m)))$$

Definitions of rubi rules used

rule 78 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

Maple [F]

$$\int \frac{(xd + c)^m (fx + e)}{(bx + a)^2} dx$$

input `int((d*x+c)^m*(f*x+e)/(b*x+a)^2,x)`

output `int((d*x+c)^m*(f*x+e)/(b*x+a)^2,x)`

Fricas [F]

$$\int \frac{(c + dx)^m (e + fx)}{(a + bx)^2} dx = \int \frac{(fx + e)(dx + c)^m}{(bx + a)^2} dx$$

input `integrate((d*x+c)^m*(f*x+e)/(b*x+a)^2,x, algorithm="fricas")`

output `integral((f*x + e)*(d*x + c)^m/(b^2*x^2 + 2*a*b*x + a^2), x)`

Sympy [F]

$$\int \frac{(c + dx)^m (e + fx)}{(a + bx)^2} dx = \int \frac{(c + dx)^m (e + fx)}{(a + bx)^2} dx$$

input `integrate((d*x+c)**m*(f*x+e)/(b*x+a)**2,x)`

output `Integral((c + d*x)**m*(e + f*x)/(a + b*x)**2, x)`

Maxima [F]

$$\int \frac{(c + dx)^m (e + fx)}{(a + bx)^2} dx = \int \frac{(fx + e)(dx + c)^m}{(bx + a)^2} dx$$

input `integrate((d*x+c)^m*(f*x+e)/(b*x+a)^2,x, algorithm="maxima")`

output `integrate((f*x + e)*(d*x + c)^m/(b*x + a)^2, x)`

Giac [F]

$$\int \frac{(c + dx)^m (e + fx)}{(a + bx)^2} dx = \int \frac{(fx + e)(dx + c)^m}{(bx + a)^2} dx$$

input `integrate((d*x+c)^m*(f*x+e)/(b*x+a)^2,x, algorithm="giac")`

output `integrate((f*x + e)*(d*x + c)^m/(b*x + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^m (e + fx)}{(a + bx)^2} dx = \int \frac{(e + fx) (c + dx)^m}{(a + bx)^2} dx$$

input `int(((e + f*x)*(c + d*x)^m)/(a + b*x)^2,x)`output `int(((e + f*x)*(c + d*x)^m)/(a + b*x)^2, x)`**Reduce [F]**

$$\int \frac{(c + dx)^m (e + fx)}{(a + bx)^2} dx = \text{too large to display}$$

input `int((d*x+c)^m*(f*x+e)/(b*x+a)^2,x)`

output

```
( - (c + d*x)**m*a*c*f + (c + d*x)**m*a*d*f*m*x + (c + d*x)**m*b*c*e*m - (
c + d*x)**m*b*c*f*x - int(((c + d*x)**m*x)/(a**3*c*d*m + a**3*d**2*m*x - a
**2*b*c**2 + 2*a**2*b*c*d*m*x - a**2*b*c*d*x + 2*a**2*b*d**2*m*x**2 - 2*a*
b**2*c**2*x + a*b**2*c*d*m*x**2 - 2*a*b**2*c*d*x**2 + a*b**2*d**2*m*x**3 -
b**3*c**2*x**2 - b**3*c*d*x**3),x)*a**4*d**3*f*m**3 - int(((c + d*x)**m*x
)/(a**3*c*d*m + a**3*d**2*m*x - a**2*b*c**2 + 2*a**2*b*c*d*m*x - a**2*b*c*
d*x + 2*a**2*b*d**2*m*x**2 - 2*a*b**2*c**2*x + a*b**2*c*d*m*x**2 - 2*a*b**
2*c*d*x**2 + a*b**2*d**2*m*x**3 - b**3*c**2*x**2 - b**3*c*d*x**3),x)*a**4*
d**3*f*m**2 + int(((c + d*x)**m*x)/(a**3*c*d*m + a**3*d**2*m*x - a**2*b*c*
**2 + 2*a**2*b*c*d*m*x - a**2*b*c*d*x + 2*a**2*b*d**2*m*x**2 - 2*a*b**2*c**
2*x + a*b**2*c*d*m*x**2 - 2*a*b**2*c*d*x**2 + a*b**2*d**2*m*x**3 - b**3*c*
**2*x**2 - b**3*c*d*x**3),x)*a**3*b*c*d**2*f*m**3 + 3*int(((c + d*x)**m*x)/
(a**3*c*d*m + a**3*d**2*m*x - a**2*b*c**2 + 2*a**2*b*c*d*m*x - a**2*b*c*d*
x + 2*a**2*b*d**2*m*x**2 - 2*a*b**2*c**2*x + a*b**2*c*d*m*x**2 - 2*a*b**2*
c*d*x**2 + a*b**2*d**2*m*x**3 - b**3*c**2*x**2 - b**3*c*d*x**3),x)*a**3*b*
c*d**2*f*m**2 + int(((c + d*x)**m*x)/(a**3*c*d*m + a**3*d**2*m*x - a**2*b*
c**2 + 2*a**2*b*c*d*m*x - a**2*b*c*d*x + 2*a**2*b*d**2*m*x**2 - 2*a*b**2*c
**2*x + a*b**2*c*d*m*x**2 - 2*a*b**2*c*d*x**2 + a*b**2*d**2*m*x**3 - b**3*
c**2*x**2 - b**3*c*d*x**3),x)*a**3*b*c*d**2*f*m + int(((c + d*x)**m*x)/(a
**3*c*d*m + a**3*d**2*m*x - a**2*b*c**2 + 2*a**2*b*c*d*m*x - a**2*b*c*d...
```

3.245 $\int \frac{(c+dx)^{-1+m}(e+fx)}{(a+bx)^2} dx$

Optimal result	2169
Mathematica [A] (verified)	2169
Rubi [A] (verified)	2170
Maple [F]	2171
Fricas [F]	2171
Sympy [F]	2172
Maxima [F]	2172
Giac [F]	2172
Mupad [F(-1)]	2173
Reduce [F]	2173

Optimal result

Integrand size = 22, antiderivative size = 107

$$\int \frac{(c+dx)^{-1+m}(e+fx)}{(a+bx)^2} dx = -\frac{f(c+dx)^m}{bd(1-m)(a+bx)} - \frac{(bcf - bde(1-m) - adfm)(c+dx)^m \operatorname{Hypergeometric2F1}\left(2, m, 1+m, \frac{b(c+dx)}{bc-ad}\right)}{b(bc-ad)^2(1-m)m}$$

output

```
-f*(d*x+c)^m/b/d/(1-m)/(b*x+a)-(b*c*f-b*d*e*(1-m)-a*d*f*m)*(d*x+c)^m*hypergeom([2, m],[1+m],b*(d*x+c)/(-a*d+b*c))/b/(-a*d+b*c)^2/(1-m)/m
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.87

$$\int \frac{(c+dx)^{-1+m}(e+fx)}{(a+bx)^2} dx = \frac{(c+dx)^m \left(-\frac{(bc-ad)(be-af)}{a+bx} - \frac{(bcf+bde(-1+m)-adfm) \operatorname{Hypergeometric2F1}\left(1, m, 1+m, \frac{b(c+dx)}{bc-ad}\right)}{m} \right)}{b(bc-ad)^2}$$

input `Integrate[((c + d*x)^(-1 + m)*(e + f*x))/(a + b*x)^2,x]`

output `((c + d*x)^m*(-(((b*c - a*d)*(b*e - a*f))/(a + b*x)) - ((b*c*f + b*d*e*(-1 + m) - a*d*f*m)*Hypergeometric2F1[1, m, 1 + m, (b*(c + d*x))/(b*c - a*d)]/m))/(b*(b*c - a*d)^2)`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {87, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(e + fx)(c + dx)^{m-1}}{(a + bx)^2} dx$$

$$\downarrow 87$$

$$\frac{(-adf m + bcf - bde(1 - m)) \int \frac{(c+dx)^{m-1}}{a+bx} dx}{b(bc - ad)} - \frac{(be - af)(c + dx)^m}{b(a + bx)(bc - ad)}$$

$$\downarrow 78$$

$$\frac{(c + dx)^m (-adf m + bcf - bde(1 - m)) \text{Hypergeometric2F1}\left(1, m, m + 1, \frac{b(c+dx)}{bc-ad}\right)}{bm(bc - ad)^2} - \frac{(be - af)(c + dx)^m}{b(a + bx)(bc - ad)}$$

input `Int[((c + d*x)^(-1 + m)*(e + f*x))/(a + b*x)^2,x]`

output `-(((b*e - a*f)*(c + d*x)^m)/(b*(b*c - a*d)*(a + b*x))) - ((b*c*f - b*d*e*(1 - m) - a*d*f*m)*(c + d*x)^m*Hypergeometric2F1[1, m, 1 + m, (b*(c + d*x))/(b*c - a*d)]/(b*(b*c - a*d)^2*m)`

Defintions of rubi rules used

rule 78 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

Maple [F]

$$\int \frac{(xd + c)^{m-1} (fx + e)}{(bx + a)^2} dx$$

input `int((d*x+c)^(m-1)*(f*x+e)/(b*x+a)^2,x)`

output `int((d*x+c)^(m-1)*(f*x+e)/(b*x+a)^2,x)`

Fricas [F]

$$\int \frac{(c + dx)^{-1+m}(e + fx)}{(a + bx)^2} dx = \int \frac{(fx + e)(dx + c)^{m-1}}{(bx + a)^2} dx$$

input `integrate((d*x+c)^(-1+m)*(f*x+e)/(b*x+a)^2,x, algorithm="fricas")`

output `integral((f*x + e)*(d*x + c)^(m - 1)/(b^2*x^2 + 2*a*b*x + a^2), x)`

Sympy [F]

$$\int \frac{(c + dx)^{-1+m}(e + fx)}{(a + bx)^2} dx = \int \frac{(c + dx)^{m-1}(e + fx)}{(a + bx)^2} dx$$

input `integrate((d*x+c)**(-1+m)*(f*x+e)/(b*x+a)**2,x)`

output `Integral((c + d*x)**(m - 1)*(e + f*x)/(a + b*x)**2, x)`

Maxima [F]

$$\int \frac{(c + dx)^{-1+m}(e + fx)}{(a + bx)^2} dx = \int \frac{(fx + e)(dx + c)^{m-1}}{(bx + a)^2} dx$$

input `integrate((d*x+c)^(-1+m)*(f*x+e)/(b*x+a)^2,x, algorithm="maxima")`

output `integrate((f*x + e)*(d*x + c)^(m - 1)/(b*x + a)^2, x)`

Giac [F]

$$\int \frac{(c + dx)^{-1+m}(e + fx)}{(a + bx)^2} dx = \int \frac{(fx + e)(dx + c)^{m-1}}{(bx + a)^2} dx$$

input `integrate((d*x+c)^(-1+m)*(f*x+e)/(b*x+a)^2,x, algorithm="giac")`

output `integrate((f*x + e)*(d*x + c)^(m - 1)/(b*x + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^{-1+m}(e + fx)}{(a + bx)^2} dx = \int \frac{(e + fx)(c + dx)^{m-1}}{(a + bx)^2} dx$$

input `int(((e + f*x)*(c + d*x)^(m - 1))/(a + b*x)^2,x)`

output `int(((e + f*x)*(c + d*x)^(m - 1))/(a + b*x)^2, x)`

Reduce [F]

$$\int \frac{(c + dx)^{-1+m}(e + fx)}{(a + bx)^2} dx = \text{too large to display}$$

input `int((d*x+c)^(-1+m)*(f*x+e)/(b*x+a)^2,x)`

output

```

((c + d*x)**m*e + int(((c + d*x)**m*x)/(a**3*c*d*m + a**3*d**2*m*x - a**2*
b*c**2 + 2*a**2*b*c*d*m*x - a**2*b*c*d*x + 2*a**2*b*d**2*m*x**2 - 2*a*b**2
*c**2*x + a*b**2*c*d*m*x**2 - 2*a*b**2*c*d*x**2 + a*b**2*d**2*m*x**3 - b**
3*c**2*x**2 - b**3*c*d*x**3),x)*a**3*d**2*f*m**2 - 2*int(((c + d*x)**m*x)/
(a**3*c*d*m + a**3*d**2*m*x - a**2*b*c**2 + 2*a**2*b*c*d*m*x - a**2*b*c*d*
x + 2*a**2*b*d**2*m*x**2 - 2*a*b**2*c**2*x + a*b**2*c*d*m*x**2 - 2*a*b**2*
c*d*x**2 + a*b**2*d**2*m*x**3 - b**3*c**2*x**2 - b**3*c*d*x**3),x)*a**2*b*
c*d*f*m - int(((c + d*x)**m*x)/(a**3*c*d*m + a**3*d**2*m*x - a**2*b*c**2 +
2*a**2*b*c*d*m*x - a**2*b*c*d*x + 2*a**2*b*d**2*m*x**2 - 2*a*b**2*c**2*x
+ a*b**2*c*d*m*x**2 - 2*a*b**2*c*d*x**2 + a*b**2*d**2*m*x**3 - b**3*c**2*x
**2 - b**3*c*d*x**3),x)*a**2*b*d**2*e*m**2 + int(((c + d*x)**m*x)/(a**3*c*
d*m + a**3*d**2*m*x - a**2*b*c**2 + 2*a**2*b*c*d*m*x - a**2*b*c*d*x + 2*a*
**2*b*d**2*m*x**2 - 2*a*b**2*c**2*x + a*b**2*c*d*m*x**2 - 2*a*b**2*c*d*x**2
+ a*b**2*d**2*m*x**3 - b**3*c**2*x**2 - b**3*c*d*x**3),x)*a**2*b*d**2*e*m
+ int(((c + d*x)**m*x)/(a**3*c*d*m + a**3*d**2*m*x - a**2*b*c**2 + 2*a**2
*b*c*d*m*x - a**2*b*c*d*x + 2*a**2*b*d**2*m*x**2 - 2*a*b**2*c**2*x + a*b**
2*c*d*m*x**2 - 2*a*b**2*c*d*x**2 + a*b**2*d**2*m*x**3 - b**3*c**2*x**2 - b
**3*c*d*x**3),x)*a**2*b*d**2*f*m**2*x + int(((c + d*x)**m*x)/(a**3*c*d*m +
a**3*d**2*m*x - a**2*b*c**2 + 2*a**2*b*c*d*m*x - a**2*b*c*d*x + 2*a**2*b*
d**2*m*x**2 - 2*a*b**2*c**2*x + a*b**2*c*d*m*x**2 - 2*a*b**2*c*d*x**2 + ...

```

3.246 $\int \frac{(c+dx)^{1+m}(e+fx)}{(a+bx)^3} dx$

Optimal result	2175
Mathematica [A] (verified)	2175
Rubi [A] (verified)	2176
Maple [F]	2177
Fricas [F]	2177
Sympy [F(-2)]	2178
Maxima [F]	2178
Giac [F]	2178
Mupad [F(-1)]	2179
Reduce [F]	2179

Optimal result

Integrand size = 22, antiderivative size = 105

$$\int \frac{(c+dx)^{1+m}(e+fx)}{(a+bx)^3} dx = \frac{f(c+dx)^{2+m}}{bdm(a+bx)^2} + \frac{d(adf(2+m) - b(2cf + dem))(c+dx)^{2+m} \operatorname{Hypergeometric2F1}\left(3, 2+m, 3+m, \frac{b(c+dx)}{bc-ad}\right)}{b(bc-ad)^3 m(2+m)}$$

```
output f*(d*x+c)^(2+m)/b/d/m/(b*x+a)^2+d*(a*d*f*(2+m)-b*(d*e*m+2*c*f))*(d*x+c)^(2+m)*hypergeom([3, 2+m],[3+m],b*(d*x+c)/(-a*d+b*c))/b/(-a*d+b*c)^3/m/(2+m)
```

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.00

$$\int \frac{(c+dx)^{1+m}(e+fx)}{(a+bx)^3} dx = \frac{(c+dx)^{2+m} \left(\frac{-be+af}{(a+bx)^2} + \frac{d(-adf(2+m)+b(2cf+dem)) \operatorname{Hypergeometric2F1}\left(2, 2+m, 3+m, \frac{b(c+dx)}{bc-ad}\right)}{(bc-ad)^2(2+m)} \right)}{2b(bc-ad)}$$

```
input Integrate[((c + d*x)^(1 + m)*(e + f*x))/(a + b*x)^3,x]
```

output

```
((c + d*x)^(2 + m)*((-b*e) + a*f)/(a + b*x)^2 + (d*(-a*d*f*(2 + m)) + b*(2*c*f + d*e*m))*Hypergeometric2F1[2, 2 + m, 3 + m, (b*(c + d*x))/(b*c - a*d)]/(b*c - a*d)^2*(2 + m)))/(2*b*(b*c - a*d))
```

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.13, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {87, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(e + fx)(c + dx)^{m+1}}{(a + bx)^3} dx$$

$$\downarrow 87$$

$$-\frac{(adf(m+2) - b(2cf + dem)) \int \frac{(c+dx)^{m+1}}{(a+bx)^2} dx}{2b(bc - ad)} - \frac{(be - af)(c + dx)^{m+2}}{2b(a + bx)^2(bc - ad)}$$

$$\downarrow 78$$

$$-\frac{d(c + dx)^{m+2}(adf(m+2) - b(2cf + dem)) \text{Hypergeometric2F1}\left(2, m+2, m+3, \frac{b(c+dx)}{bc-ad}\right)}{2b(m+2)(bc - ad)^3 \frac{(be - af)(c + dx)^{m+2}}{2b(a + bx)^2(bc - ad)}}$$

input

```
Int[((c + d*x)^(1 + m)*(e + f*x))/(a + b*x)^3,x]
```

output

```
-1/2*((b*e - a*f)*(c + d*x)^(2 + m))/(b*(b*c - a*d)*(a + b*x)^2) - (d*(a*d*f*(2 + m) - b*(2*c*f + d*e*m))*(c + d*x)^(2 + m)*Hypergeometric2F1[2, 2 + m, 3 + m, (b*(c + d*x))/(b*c - a*d)]/(2*b*(b*c - a*d)^3*(2 + m))
```

Definitions of rubi rules used

rule 78 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

Maple [F]

$$\int \frac{(xd + c)^{1+m} (fx + e)}{(bx + a)^3} dx$$

input `int((d*x+c)^(1+m)*(f*x+e)/(b*x+a)^3,x)`

output `int((d*x+c)^(1+m)*(f*x+e)/(b*x+a)^3,x)`

Fricas [F]

$$\int \frac{(c + dx)^{1+m} (e + fx)}{(a + bx)^3} dx = \int \frac{(fx + e)(dx + c)^{m+1}}{(bx + a)^3} dx$$

input `integrate((d*x+c)^(1+m)*(f*x+e)/(b*x+a)^3,x, algorithm="fricas")`

output `integral((f*x + e)*(d*x + c)^(m + 1)/(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3), x)`

Sympy [F(-2)]

Exception generated.

$$\int \frac{(c + dx)^{1+m}(e + fx)}{(a + bx)^3} dx = \text{Exception raised: HeuristicGCDFailed}$$

input `integrate((d*x+c)**(1+m)*(f*x+e)/(b*x+a)**3,x)`

output `Exception raised: HeuristicGCDFailed >> no luck`

Maxima [F]

$$\int \frac{(c + dx)^{1+m}(e + fx)}{(a + bx)^3} dx = \int \frac{(fx + e)(dx + c)^{m+1}}{(bx + a)^3} dx$$

input `integrate((d*x+c)^(1+m)*(f*x+e)/(b*x+a)^3,x, algorithm="maxima")`

output `integrate((f*x + e)*(d*x + c)^(m + 1)/(b*x + a)^3, x)`

Giac [F]

$$\int \frac{(c + dx)^{1+m}(e + fx)}{(a + bx)^3} dx = \int \frac{(fx + e)(dx + c)^{m+1}}{(bx + a)^3} dx$$

input `integrate((d*x+c)^(1+m)*(f*x+e)/(b*x+a)^3,x, algorithm="giac")`

output `integrate((f*x + e)*(d*x + c)^(m + 1)/(b*x + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^{1+m}(e + fx)}{(a + bx)^3} dx = \int \frac{(e + fx)(c + dx)^{m+1}}{(a + bx)^3} dx$$

input `int(((e + f*x)*(c + d*x)^(m + 1))/(a + b*x)^3,x)`output `int(((e + f*x)*(c + d*x)^(m + 1))/(a + b*x)^3, x)`**Reduce [F]**

$$\int \frac{(c + dx)^{1+m}(e + fx)}{(a + bx)^3} dx = \text{too large to display}$$

input `int((d*x+c)^(1+m)*(f*x+e)/(b*x+a)^3,x)`

output

```

((c + d*x)**m*a**2*c*d*f*m + 2*(c + d*x)**m*a**2*c*d*f - (c + d*x)**m*a**2
*d**2*f*m**2*x - 2*(c + d*x)**m*a**2*d**2*f*m*x - 2*(c + d*x)**m*a*b*c**2*
f*m - (c + d*x)**m*a*b*c*d*e*m + 2*(c + d*x)**m*a*b*c*d*f*m**2*x + 2*(c +
d*x)**m*a*b*c*d*f*m*x + 4*(c + d*x)**m*a*b*c*d*f*x + (c + d*x)**m*a*b*d**2
*e*m**2*x + (c + d*x)**m*a*b*d**2*f*m**2*x**2 - (c + d*x)**m*a*b*d**2*f*m*
x**2 + (c + d*x)**m*b**2*c**2*e*m**2 - (c + d*x)**m*b**2*c**2*e*m - 4*(c +
d*x)**m*b**2*c**2*f*m*x - 2*(c + d*x)**m*b**2*c*d*e*m*x - 2*(c + d*x)**m*
b**2*c*d*f*m*x**2 + 2*(c + d*x)**m*b**2*c*d*f*x**2 + int(((c + d*x)**m*x)/
(a**4*c*d*m**2 - a**4*c*d*m + a**4*d**2*m**2*x - a**4*d**2*m*x - 2*a**3*b*
c**2*m + 2*a**3*b*c**2 + 3*a**3*b*c*d*m**2*x - 5*a**3*b*c*d*m*x + 2*a**3*b
*c*d*x + 3*a**3*b*d**2*m**2*x**2 - 3*a**3*b*d**2*m*x**2 - 6*a**2*b**2*c**2
*m*x + 6*a**2*b**2*c**2*x + 3*a**2*b**2*c*d*m**2*x**2 - 9*a**2*b**2*c*d*m*
x**2 + 6*a**2*b**2*c*d*x**2 + 3*a**2*b**2*d**2*m**2*x**3 - 3*a**2*b**2*d**
2*m*x**3 - 6*a*b**3*c**2*m*x**2 + 6*a*b**3*c**2*x**2 + a*b**3*c*d*m**2*x**
3 - 7*a*b**3*c*d*m*x**3 + 6*a*b**3*c*d*x**3 + a*b**3*d**2*m**2*x**4 - a*b*
**3*d**2*m*x**4 - 2*b**4*c**2*m*x**3 + 2*b**4*c**2*x**3 - 2*b**4*c*d*m*x**4
+ 2*b**4*c*d*x**4),x)*a**6*d**4*f*m**5 + 2*int(((c + d*x)**m*x)/(a**4*c*d
*m**2 - a**4*c*d*m + a**4*d**2*m**2*x - a**4*d**2*m*x - 2*a**3*b*c**2*m +
2*a**3*b*c**2 + 3*a**3*b*c*d*m**2*x - 5*a**3*b*c*d*m*x + 2*a**3*b*c*d*x +
3*a**3*b*d**2*m**2*x**2 - 3*a**3*b*d**2*m*x**2 - 6*a**2*b**2*c**2*m*x + ...

```

3.247 $\int \frac{(c+dx)^m(e+fx)}{(a+bx)^3} dx$

Optimal result	2181
Mathematica [A] (verified)	2181
Rubi [A] (verified)	2182
Maple [F]	2183
Fricas [F]	2183
Sympy [F]	2184
Maxima [F]	2184
Giac [F]	2184
Mupad [F(-1)]	2185
Reduce [F]	2185

Optimal result

Integrand size = 20, antiderivative size = 118

$$\int \frac{(c+dx)^m(e+fx)}{(a+bx)^3} dx = -\frac{f(c+dx)^{1+m}}{bd(1-m)(a+bx)^2} + \frac{d(2bcf - bde(1-m) - adf(1+m))(c+dx)^{1+m} \operatorname{Hypergeometric2F1}\left(3, 1+m, 2+m, \frac{b(c+dx)}{bc-ad}\right)}{b(bc-ad)^3(1-m)(1+m)}$$

output

```
-f*(d*x+c)^(1+m)/b/d/(1-m)/(b*x+a)^2+d*(2*b*c*f-b*d*e*(1-m)-a*d*f*(1+m))*
(d*x+c)^(1+m)*hypergeom([3, 1+m],[2+m],b*(d*x+c)/(-a*d+b*c))/b/(-a*d+b*c)^3
/(1-m)/(1+m)
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.90

$$\int \frac{(c+dx)^m(e+fx)}{(a+bx)^3} dx = \frac{(c+dx)^{1+m} \left(\frac{-be+af}{(a+bx)^2} + \frac{d(2bcf+bde(-1+m)-adf(1+m)) \operatorname{Hypergeometric2F1}\left(2, 1+m, 2+m, \frac{b(c+dx)}{bc-ad}\right)}{(bc-ad)^2(1+m)} \right)}{2b(bc-ad)}$$

input `Integrate[((c + d*x)^m*(e + f*x))/(a + b*x)^3,x]`

output `((c + d*x)^(1 + m)*((-b*e) + a*f)/(a + b*x)^2 + (d*(2*b*c*f + b*d*e*(-1 + m) - a*d*f*(1 + m))*Hypergeometric2F1[2, 1 + m, 2 + m, (b*(c + d*x))/(b*c - a*d)])/(b*c - a*d)^2*(1 + m)))/(2*b*(b*c - a*d))`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.04, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {87, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(e + fx)(c + dx)^m}{(a + bx)^3} dx$$

$$\downarrow 87$$

$$\frac{(-adf(m + 1) + 2bcf - bde(1 - m)) \int \frac{(c+dx)^m}{(a+bx)^2} dx}{2b(bc - ad)} - \frac{(be - af)(c + dx)^{m+1}}{2b(a + bx)^2(bc - ad)}$$

$$\downarrow 78$$

$$\frac{d(c + dx)^{m+1}(-adf(m + 1) + 2bcf - bde(1 - m)) \text{Hypergeometric2F1}\left(2, m + 1, m + 2, \frac{b(c+dx)}{bc-ad}\right)}{2b(m + 1)(bc - ad)^3} - \frac{(be - af)(c + dx)^{m+1}}{2b(a + bx)^2(bc - ad)}$$

input `Int[((c + d*x)^m*(e + f*x))/(a + b*x)^3,x]`

output `-1/2*((b*e - a*f)*(c + d*x)^(1 + m))/(b*(b*c - a*d)*(a + b*x)^2) + (d*(2*b*c*f - b*d*e*(1 - m) - a*d*f*(1 + m))*(c + d*x)^(1 + m)*Hypergeometric2F1[2, 1 + m, 2 + m, (b*(c + d*x))/(b*c - a*d)])/(2*b*(b*c - a*d)^3*(1 + m))`

Definitions of rubi rules used

rule 78

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b
*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m
+ 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x]
&& !IntegerQ[m] && IntegerQ[n]
```

rule 87

```
Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p
_), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

Maple [F]

$$\int \frac{(xd + c)^m (fx + e)}{(bx + a)^3} dx$$

input

```
int((d*x+c)^m*(f*x+e)/(b*x+a)^3,x)
```

output

```
int((d*x+c)^m*(f*x+e)/(b*x+a)^3,x)
```

Fricas [F]

$$\int \frac{(c + dx)^m (e + fx)}{(a + bx)^3} dx = \int \frac{(fx + e)(dx + c)^m}{(bx + a)^3} dx$$

input

```
integrate((d*x+c)^m*(f*x+e)/(b*x+a)^3,x, algorithm="fricas")
```

output

```
integral((f*x + e)*(d*x + c)^m/(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3),
x)
```

Sympy [F]

$$\int \frac{(c + dx)^m (e + fx)}{(a + bx)^3} dx = \int \frac{(c + dx)^m (e + fx)}{(a + bx)^3} dx$$

input `integrate((d*x+c)**m*(f*x+e)/(b*x+a)**3,x)`

output `Integral((c + d*x)**m*(e + f*x)/(a + b*x)**3, x)`

Maxima [F]

$$\int \frac{(c + dx)^m (e + fx)}{(a + bx)^3} dx = \int \frac{(fx + e)(dx + c)^m}{(bx + a)^3} dx$$

input `integrate((d*x+c)^m*(f*x+e)/(b*x+a)^3,x, algorithm="maxima")`

output `integrate((f*x + e)*(d*x + c)^m/(b*x + a)^3, x)`

Giac [F]

$$\int \frac{(c + dx)^m (e + fx)}{(a + bx)^3} dx = \int \frac{(fx + e)(dx + c)^m}{(bx + a)^3} dx$$

input `integrate((d*x+c)^m*(f*x+e)/(b*x+a)^3,x, algorithm="giac")`

output `integrate((f*x + e)*(d*x + c)^m/(b*x + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^m (e + fx)}{(a + bx)^3} dx = \int \frac{(e + fx) (c + dx)^m}{(a + bx)^3} dx$$

input `int(((e + f*x)*(c + d*x)^m)/(a + b*x)^3,x)`output `int(((e + f*x)*(c + d*x)^m)/(a + b*x)^3, x)`**Reduce [F]**

$$\int \frac{(c + dx)^m (e + fx)}{(a + bx)^3} dx = \text{too large to display}$$

input `int((d*x+c)^m*(f*x+e)/(b*x+a)^3,x)`

output

```
( - (c + d*x)**m*a*c*f + (c + d*x)**m*a*d*f*m*x + (c + d*x)**m*b*c*e*m - (
c + d*x)**m*b*c*e - 2*(c + d*x)**m*b*c*f*x - int(((c + d*x)**m*x)/(a**4*c*
d*m**2 - a**4*c*d*m + a**4*d**2*m**2*x - a**4*d**2*m*x - 2*a**3*b*c**2*m +
2*a**3*b*c**2 + 3*a**3*b*c*d*m**2*x - 5*a**3*b*c*d*m*x + 2*a**3*b*c*d*x +
3*a**3*b*d**2*m**2*x**2 - 3*a**3*b*d**2*m*x**2 - 6*a**2*b**2*c**2*m*x + 6
*a**2*b**2*c**2*x + 3*a**2*b**2*c*d*m**2*x**2 - 9*a**2*b**2*c*d*m*x**2 + 6
*a**2*b**2*c*d*x**2 + 3*a**2*b**2*d**2*m**2*x**3 - 3*a**2*b**2*d**2*m*x**3
- 6*a*b**3*c**2*m*x**2 + 6*a*b**3*c**2*x**2 + a*b**3*c*d*m**2*x**3 - 7*a*
b**3*c*d*m*x**3 + 6*a*b**3*c*d*x**3 + a*b**3*d**2*m**2*x**4 - a*b**3*d**2*
m*x**4 - 2*b**4*c**2*m*x**3 + 2*b**4*c**2*x**3 - 2*b**4*c*d*m*x**4 + 2*b**
4*c*d*x**4),x)*a**5*d**3*f*m**4 + int(((c + d*x)**m*x)/(a**4*c*d*m**2 - a*
**4*c*d*m + a**4*d**2*m**2*x - a**4*d**2*m*x - 2*a**3*b*c**2*m + 2*a**3*b*c
**2 + 3*a**3*b*c*d*m**2*x - 5*a**3*b*c*d*m*x + 2*a**3*b*c*d*x + 3*a**3*b*d
**2*m**2*x**2 - 3*a**3*b*d**2*m*x**2 - 6*a**2*b**2*c**2*m*x + 6*a**2*b**2*
c**2*x + 3*a**2*b**2*c*d*m**2*x**2 - 9*a**2*b**2*c*d*m*x**2 + 6*a**2*b**2*
c*d*x**2 + 3*a**2*b**2*d**2*m**2*x**3 - 3*a**2*b**2*d**2*m*x**3 - 6*a*b**3
*c**2*m*x**2 + 6*a*b**3*c**2*x**2 + a*b**3*c*d*m**2*x**3 - 7*a*b**3*c*d*m*
x**3 + 6*a*b**3*c*d*x**3 + a*b**3*d**2*m**2*x**4 - a*b**3*d**2*m*x**4 - 2*
b**4*c**2*m*x**3 + 2*b**4*c**2*x**3 - 2*b**4*c*d*m*x**4 + 2*b**4*c*d*x**4)
,x)*a**5*d**3*f*m**2 + int(((c + d*x)**m*x)/(a**4*c*d*m**2 - a**4*c*d*m...
```

3.248 $\int \frac{(c+dx)^{-1+m}(e+fx)}{(a+bx)^3} dx$

Optimal result	2187
Mathematica [A] (verified)	2187
Rubi [A] (verified)	2188
Maple [F]	2189
Fricas [F]	2189
Sympy [F]	2190
Maxima [F]	2190
Giac [F]	2190
Mupad [F(-1)]	2191
Reduce [F]	2191

Optimal result

Integrand size = 22, antiderivative size = 109

$$\int \frac{(c+dx)^{-1+m}(e+fx)}{(a+bx)^3} dx = -\frac{f(c+dx)^m}{bd(2-m)(a+bx)^2} + \frac{d(b(2cf-de(2-m))-adf m)(c+dx)^m \operatorname{Hypergeometric2F1}\left(3, m, 1+m, \frac{b(c+dx)}{bc-ad}\right)}{b(bc-ad)^3(2-m)m}$$

output

```
-f*(d*x+c)^m/b/d/(2-m)/(b*x+a)^2+d*(b*(2*c*f-d*e*(2-m))-a*d*f*m)*(d*x+c)^m
*hypergeom([3, m], [1+m], b*(d*x+c)/(-a*d+b*c))/b/(-a*d+b*c)^3/(2-m)/m
```

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.90

$$\int \frac{(c+dx)^{-1+m}(e+fx)}{(a+bx)^3} dx = \frac{(c+dx)^m \left(\frac{-be+af}{(a+bx)^2} + \frac{d(2bcf+bde(-2+m)-adf m) \operatorname{Hypergeometric2F1}\left(2, m, 1+m, \frac{b(c+dx)}{bc-ad}\right)}{(bc-ad)^2 m} \right)}{2b(bc-ad)}$$

input

```
Integrate[((c + d*x)^(-1 + m)*(e + f*x))/(a + b*x)^3,x]
```


output

$$\frac{((c + dx)^m((-b*e) + a*f)/(a + b*x)^2 + (d*(2*b*c*f + b*d*e*(-2 + m) - a*d*f*m)*Hypergeometric2F1[2, m, 1 + m, (b*(c + d*x))/(b*c - a*d)])/((b*c - a*d)^{2*m}))}{2*b*(b*c - a*d)}$$
Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.04, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {87, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(e + fx)(c + dx)^{m-1}}{(a + bx)^3} dx$$

↓ 87

$$\frac{(-adf m + 2bcf - bde(2 - m)) \int \frac{(c+dx)^{m-1}}{(a+bx)^2} dx}{2b(bc - ad)} - \frac{(be - af)(c + dx)^m}{2b(a + bx)^2(bc - ad)}$$

↓ 78

$$\frac{d(c + dx)^m(-adf m + 2bcf - bde(2 - m)) \text{Hypergeometric2F1}\left(2, m, m + 1, \frac{b(c+dx)}{bc-ad}\right) - \frac{2bm(bc - ad)^3}{(be - af)(c + dx)^m}}{2b(a + bx)^2(bc - ad)}$$

input

$$\text{Int}[\frac{(c + d*x)^{-1 + m}*(e + f*x)}{(a + b*x)^3}, x]$$

output

$$-1/2*((b*e - a*f)*(c + d*x)^m)/(b*(b*c - a*d)*(a + b*x)^2) + (d*(2*b*c*f - b*d*e*(2 - m) - a*d*f*m)*(c + d*x)^m*Hypergeometric2F1[2, m, 1 + m, (b*(c + d*x))/(b*c - a*d)])/(2*b*(b*c - a*d)^3*m)$$

Definitions of rubi rules used

rule 78

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b
*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m
+ 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x]
&& !IntegerQ[m] && IntegerQ[n]
```

rule 87

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))
```

Maple [F]

$$\int \frac{(xd + c)^{m-1} (fx + e)}{(bx + a)^3} dx$$

input

```
int((d*x+c)^(m-1)*(f*x+e)/(b*x+a)^3,x)
```

output

```
int((d*x+c)^(m-1)*(f*x+e)/(b*x+a)^3,x)
```

Fricas [F]

$$\int \frac{(c + dx)^{-1+m} (e + fx)}{(a + bx)^3} dx = \int \frac{(fx + e)(dx + c)^{m-1}}{(bx + a)^3} dx$$

input

```
integrate((d*x+c)^(-1+m)*(f*x+e)/(b*x+a)^3,x, algorithm="fricas")
```

output

```
integral((f*x + e)*(d*x + c)^(m - 1)/(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x +
a^3), x)
```

Sympy [F]

$$\int \frac{(c + dx)^{-1+m}(e + fx)}{(a + bx)^3} dx = \int \frac{(c + dx)^{m-1}(e + fx)}{(a + bx)^3} dx$$

input `integrate((d*x+c)**(-1+m)*(f*x+e)/(b*x+a)**3,x)`

output `Integral((c + d*x)**(m - 1)*(e + f*x)/(a + b*x)**3, x)`

Maxima [F]

$$\int \frac{(c + dx)^{-1+m}(e + fx)}{(a + bx)^3} dx = \int \frac{(fx + e)(dx + c)^{m-1}}{(bx + a)^3} dx$$

input `integrate((d*x+c)^(-1+m)*(f*x+e)/(b*x+a)^3,x, algorithm="maxima")`

output `integrate((f*x + e)*(d*x + c)^(m - 1)/(b*x + a)^3, x)`

Giac [F]

$$\int \frac{(c + dx)^{-1+m}(e + fx)}{(a + bx)^3} dx = \int \frac{(fx + e)(dx + c)^{m-1}}{(bx + a)^3} dx$$

input `integrate((d*x+c)^(-1+m)*(f*x+e)/(b*x+a)^3,x, algorithm="giac")`

output `integrate((f*x + e)*(d*x + c)^(m - 1)/(b*x + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^{-1+m}(e + fx)}{(a + bx)^3} dx = \int \frac{(e + fx)(c + dx)^{m-1}}{(a + bx)^3} dx$$

input `int(((e + f*x)*(c + d*x)^(m - 1))/(a + b*x)^3,x)`output `int(((e + f*x)*(c + d*x)^(m - 1))/(a + b*x)^3, x)`**Reduce [F]**

$$\int \frac{(c + dx)^{-1+m}(e + fx)}{(a + bx)^3} dx = \text{too large to display}$$

input `int((d*x+c)^(-1+m)*(f*x+e)/(b*x+a)^3,x)`

output

```

((c + d*x)**m*e + int(((c + d*x)**m*x)/(a**4*c*d*m + a**4*d**2*m*x - 2*a**
3*b*c**2 + 3*a**3*b*c*d*m*x - 2*a**3*b*c*d*x + 3*a**3*b*d**2*m*x**2 - 6*a**
*2*b**2*c**2*x + 3*a**2*b**2*c*d*m*x**2 - 6*a**2*b**2*c*d*x**2 + 3*a**2*b*
*2*d**2*m*x**3 - 6*a*b**3*c**2*x**2 + a*b**3*c*d*m*x**3 - 6*a*b**3*c*d*x**
3 + a*b**3*d**2*m*x**4 - 2*b**4*c**2*x**3 - 2*b**4*c*d*x**4),x)*a**4*d**2*
f*m**2 - 4*int(((c + d*x)**m*x)/(a**4*c*d*m + a**4*d**2*m*x - 2*a**3*b*c**
2 + 3*a**3*b*c*d*m*x - 2*a**3*b*c*d*x + 3*a**3*b*d**2*m*x**2 - 6*a**2*b**2
*c**2*x + 3*a**2*b**2*c*d*m*x**2 - 6*a**2*b**2*c*d*x**2 + 3*a**2*b**2*d**2
*m*x**3 - 6*a*b**3*c**2*x**2 + a*b**3*c*d*m*x**3 - 6*a*b**3*c*d*x**3 + a*b
**3*d**2*m*x**4 - 2*b**4*c**2*x**3 - 2*b**4*c*d*x**4),x)*a**3*b*c*d*f*m -
int(((c + d*x)**m*x)/(a**4*c*d*m + a**4*d**2*m*x - 2*a**3*b*c**2 + 3*a**3*
b*c*d*m*x - 2*a**3*b*c*d*x + 3*a**3*b*d**2*m*x**2 - 6*a**2*b**2*c**2*x + 3
*a**2*b**2*c*d*m*x**2 - 6*a**2*b**2*c*d*x**2 + 3*a**2*b**2*d**2*m*x**3 - 6
*a*b**3*c**2*x**2 + a*b**3*c*d*m*x**3 - 6*a*b**3*c*d*x**3 + a*b**3*d**2*m*
x**4 - 2*b**4*c**2*x**3 - 2*b**4*c*d*x**4),x)*a**3*b*d**2*e*m**2 + 2*int((
(c + d*x)**m*x)/(a**4*c*d*m + a**4*d**2*m*x - 2*a**3*b*c**2 + 3*a**3*b*c*d
*m*x - 2*a**3*b*c*d*x + 3*a**3*b*d**2*m*x**2 - 6*a**2*b**2*c**2*x + 3*a**2
*b**2*c*d*m*x**2 - 6*a**2*b**2*c*d*x**2 + 3*a**2*b**2*d**2*m*x**3 - 6*a*b
**3*c**2*x**2 + a*b**3*c*d*m*x**3 - 6*a*b**3*c*d*x**3 + a*b**3*d**2*m*x**4
- 2*b**4*c**2*x**3 - 2*b**4*c*d*x**4),x)*a**3*b*d**2*e*m + 2*int(((c + ...

```

3.249 $\int (a + bx)^m (c + dx)^n (e + fx) dx$

Optimal result	2193
Mathematica [A] (verified)	2193
Rubi [A] (verified)	2194
Maple [F]	2195
Fricas [F]	2196
Sympy [F(-2)]	2196
Maxima [F]	2196
Giac [F]	2197
Mupad [F(-1)]	2197
Reduce [F]	2197

Optimal result

Integrand size = 20, antiderivative size = 131

$$\int (a + bx)^m (c + dx)^n (e + fx) dx = \frac{f(a + bx)^{1+m} (c + dx)^{1+n}}{bd(2 + m + n)} - \frac{(bcf(1 + m) + adf(1 + n) - bde(2 + m + n))(a + bx)^{1+m} (c + dx)^{1+n} \text{Hypergeometric2F1}\left(1, 2 + m + n, 2 + m + n, \frac{b(c + dx)}{bc - ad}\right)}{bd(bc - ad)(1 + m)(2 + m + n)}$$

output

```
f*(b*x+a)^(1+m)*(d*x+c)^(1+n)/b/d/(2+m+n)-(b*c*f*(1+m)+a*d*f*(1+n)-b*d*e*(2+m+n)*(b*x+a)^(1+m)*(d*x+c)^(1+n)*hypergeom([1, 2+m+n],[2+m+n],-d*(b*x+a)/(-a*d+b*c))/b/d/(-a*d+b*c)/(1+m)/(2+m+n)
```

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.89

$$\int (a + bx)^m (c + dx)^n (e + fx) dx = \frac{(a + bx)^{1+m} (c + dx)^n \left(bf(c + dx) + \frac{(-bcf(1+m) - adf(1+n) + bde(2+m+n)) \left(\frac{b(c+dx)}{bc-ad}\right)^{-n} \text{Hypergeometric2F1}\left(1+m, -n, 2+m+n, \frac{b(c+dx)}{bc-ad}\right)}{1+m} \right)}{b^2 d(2 + m + n)}$$

input `Integrate[(a + b*x)^m*(c + d*x)^n*(e + f*x),x]`

output `((a + b*x)^(1 + m)*(c + d*x)^n*(b*f*(c + d*x) + ((-(b*c*f*(1 + m)) - a*d*f*(1 + n) + b*d*e*(2 + m + n))*Hypergeometric2F1[1 + m, -n, 2 + m, (d*(a + b*x))/(-(b*c) + a*d)])/((1 + m)*((b*(c + d*x))/(b*c - a*d))^n))/(b^2*d*(2 + m + n))`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.05, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {90, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx)(a + bx)^m(c + dx)^n dx$$

$$\downarrow 90$$

$$\left(e - \frac{f(ad(n+1) + bc(m+1))}{bd(m+n+2)} \right) \int (a + bx)^m(c + dx)^n dx + \frac{f(a + bx)^{m+1}(c + dx)^{n+1}}{bd(m+n+2)}$$

$$\downarrow 80$$

$$(c + dx)^n \left(\frac{b(c + dx)}{bc - ad} \right)^{-n} \left(e - \frac{f(ad(n+1) + bc(m+1))}{bd(m+n+2)} \right) \int (a + bx)^m \left(\frac{bc}{bc - ad} + \frac{bdx}{bc - ad} \right)^n dx + \frac{f(a + bx)^{m+1}(c + dx)^{n+1}}{bd(m+n+2)}$$

$$\downarrow 79$$

$$\frac{(a + bx)^{m+1}(c + dx)^n \left(\frac{b(c+dx)}{bc-ad} \right)^{-n} \left(e - \frac{f(ad(n+1)+bc(m+1))}{bd(m+n+2)} \right) \text{Hypergeometric2F1} \left(m+1, -n, m+2, -\frac{d(a+bx)}{bc-ad} \right)}{b(m+1)} + \frac{f(a + bx)^{m+1}(c + dx)^{n+1}}{bd(m+n+2)}$$

input `Int[(a + b*x)^m*(c + d*x)^n*(e + f*x),x]`

output

```
(f*(a + b*x)^(1 + m)*(c + d*x)^(1 + n))/(b*d*(2 + m + n)) + ((e - (f*(b*c*(1 + m) + a*d*(1 + n)))/(b*d*(2 + m + n)))*(a + b*x)^(1 + m)*(c + d*x)^n*Hypergeometric2F1[1 + m, -n, 2 + m, -((d*(a + b*x))/(b*c - a*d))]/(b*(1 + m)*((b*(c + d*x))/(b*c - a*d))^n)
```

Defintions of rubi rules used

rule 79

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))
```

rule 80

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(c + d*x)^FracPart[n]/((b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])
```

rule 90

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]
```

Maple [F]

$$\int (bx + a)^m (xd + c)^n (fx + e) dx$$

input

```
int((b*x+a)^m*(d*x+c)^n*(f*x+e),x)
```

output

```
int((b*x+a)^m*(d*x+c)^n*(f*x+e),x)
```


Fricas [F]

$$\int (a + bx)^m (c + dx)^n (e + fx) dx = \int (fx + e)(bx + a)^m (dx + c)^n dx$$

input `integrate((b*x+a)^m*(d*x+c)^n*(f*x+e),x, algorithm="fricas")`

output `integral((f*x + e)*(b*x + a)^m*(d*x + c)^n, x)`

Sympy [F(-2)]

Exception generated.

$$\int (a + bx)^m (c + dx)^n (e + fx) dx = \text{Exception raised: HeuristicGCDFailed}$$

input `integrate((b*x+a)**m*(d*x+c)**n*(f*x+e),x)`

output `Exception raised: HeuristicGCDFailed >> no luck`

Maxima [F]

$$\int (a + bx)^m (c + dx)^n (e + fx) dx = \int (fx + e)(bx + a)^m (dx + c)^n dx$$

input `integrate((b*x+a)^m*(d*x+c)^n*(f*x+e),x, algorithm="maxima")`

output `integrate((f*x + e)*(b*x + a)^m*(d*x + c)^n, x)`

Giac [F]

$$\int (a + bx)^m (c + dx)^n (e + fx) dx = \int (fx + e)(bx + a)^m (dx + c)^n dx$$

input `integrate((b*x+a)^m*(d*x+c)^n*(f*x+e),x, algorithm="giac")`

output `integrate((f*x + e)*(b*x + a)^m*(d*x + c)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + bx)^m (c + dx)^n (e + fx) dx = \int (e + fx) (a + bx)^m (c + dx)^n dx$$

input `int((e + f*x)*(a + b*x)^m*(c + d*x)^n,x)`

output `int((e + f*x)*(a + b*x)^m*(c + d*x)^n, x)`

Reduce [F]

$$\int (a + bx)^m (c + dx)^n (e + fx) dx = \text{too large to display}$$

input `int((b*x+a)^m*(d*x+c)^n*(f*x+e),x)`

output

```
( - (c + d*x)**n*(a + b*x)**m*a**2*c*d*f*m + (c + d*x)**n*(a + b*x)**m*a**
2*d**2*f*m*n*x - (c + d*x)**n*(a + b*x)**m*a*b*c**2*f*n + (c + d*x)**n*(a
+ b*x)**m*a*b*c*d*e*m**2 + 2*(c + d*x)**n*(a + b*x)**m*a*b*c*d*e*m*n + 2*(
c + d*x)**n*(a + b*x)**m*a*b*c*d*e*m + (c + d*x)**n*(a + b*x)**m*a*b*c*d*e
n**2 + 2*(c + d*x)**n*(a + b*x)**m*a*b*c*d*e*n + (c + d*x)**n*(a + b*x)**
m*a*b*c*d*f*m**2*x + (c + d*x)**n*(a + b*x)**m*a*b*c*d*f*n**2*x + (c + d*x
)**n*(a + b*x)**m*a*b*d**2*e*m*n*x + (c + d*x)**n*(a + b*x)**m*a*b*d**2*e*
n**2*x + 2*(c + d*x)**n*(a + b*x)**m*a*b*d**2*e*n*x + (c + d*x)**n*(a + b*
x)**m*a*b*d**2*f*m*n*x**2 + (c + d*x)**n*(a + b*x)**m*a*b*d**2*f*n**2*x**2
+ (c + d*x)**n*(a + b*x)**m*a*b*d**2*f*n*x**2 + (c + d*x)**n*(a + b*x)**m
*b**2*c**2*f*m*n*x + (c + d*x)**n*(a + b*x)**m*b**2*c*d*e*m**2*x + (c + d*
x)**n*(a + b*x)**m*b**2*c*d*e*m*n*x + 2*(c + d*x)**n*(a + b*x)**m*b**2*c*d
*e*m*x + (c + d*x)**n*(a + b*x)**m*b**2*c*d*f*m**2*x**2 + (c + d*x)**n*(a
+ b*x)**m*b**2*c*d*f*m*n*x**2 + (c + d*x)**n*(a + b*x)**m*b**2*c*d*f*m*x**
2 - int(((c + d*x)**n*(a + b*x)**m*x)/(a**2*c*d*m**2*n + 2*a**2*c*d*m*n**2
+ 3*a**2*c*d*m*n + a**2*c*d*n**3 + 3*a**2*c*d*n**2 + 2*a**2*c*d*n + a**2*
d**2*m**2*n*x + 2*a**2*d**2*m*n**2*x + 3*a**2*d**2*m*n*x + a**2*d**2*n**3*
x + 3*a**2*d**2*n**2*x + 2*a**2*d**2*n*x + a*b*c**2*m**3 + 2*a*b*c**2*m**2
*n + 3*a*b*c**2*m**2 + a*b*c**2*m*n**2 + 3*a*b*c**2*m*n + 2*a*b*c**2*m + a
*b*c*d*m**3*x + 3*a*b*c*d*m**2*n*x + 3*a*b*c*d*m**2*x + 3*a*b*c*d*m*n**...
```

3.250 $\int (a + bx)^p (c + dx)^{-2-p} (e + fx) dx$

Optimal result	2199
Mathematica [A] (verified)	2199
Rubi [A] (verified)	2200
Maple [F]	2201
Fricas [F]	2202
Sympy [F]	2202
Maxima [F]	2202
Giac [F]	2203
Mupad [F(-1)]	2203
Reduce [F]	2203

Optimal result

Integrand size = 24, antiderivative size = 109

$$\int (a + bx)^p (c + dx)^{-2-p} (e + fx) dx$$

$$= \frac{(de - cf)(a + bx)^{1+p} (c + dx)^{-1-p}}{d(bc - ad)(1 + p)}$$

$$+ \frac{f(a + bx)^{1+p} (c + dx)^{-p} \operatorname{Hypergeometric2F1}\left(1, 1, 1 - p, \frac{b(c+dx)}{bc-ad}\right)}{d(bc - ad)p}$$

output

```
(-c*f+d*e)*(b*x+a)^(p+1)*(d*x+c)^(-1-p)/d/(-a*d+b*c)/(p+1)+f*(b*x+a)^(p+1)
*hypergeom([1, 1], [1-p], b*(d*x+c)/(-a*d+b*c))/d/(-a*d+b*c)/p/((d*x+c)^p)
```

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.05

$$\int (a + bx)^p (c + dx)^{-2-p} (e + fx) dx$$

$$= \frac{(a + bx)^p (c + dx)^{-p} \left(\frac{d(de - cf)(a + bx)}{(bc - ad)(1 + p)(c + dx)} - \frac{f \left(\frac{d(a + bx)}{-bc + ad} \right)^{-p} \operatorname{Hypergeometric2F1}\left(-p, -p, 1 - p, \frac{b(c + dx)}{bc - ad}\right)}{p} \right)}{d^2}$$

input `Integrate[(a + b*x)^p*(c + d*x)^(-2 - p)*(e + f*x),x]`

output
$$\frac{((a + b*x)^p*((d*(d*e - c*f)*(a + b*x))/((b*c - a*d)*(1 + p)*(c + d*x)) - (f*Hypergeometric2F1[-p, -p, 1 - p, (b*(c + d*x))/(b*c - a*d]])/p*((d*(a + b*x))/(-b*c + a*d))^p)}{(d^2*(c + d*x)^p)}$$

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.14, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {88, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (e + fx)(a + bx)^p(c + dx)^{-p-2} dx \\ & \quad \downarrow 88 \\ & \frac{f \int (a + bx)^p(c + dx)^{-p-1} dx}{d} + \frac{(a + bx)^{p+1}(de - cf)(c + dx)^{-p-1}}{d(p + 1)(bc - ad)} \\ & \quad \downarrow 80 \\ & \frac{f(a + bx)^p \left(-\frac{d(a+bx)}{bc-ad}\right)^{-p} \int (c + dx)^{-p-1} \left(-\frac{bxd}{bc-ad} - \frac{ad}{bc-ad}\right)^p dx}{\frac{d}{(a + bx)^{p+1}(de - cf)(c + dx)^{-p-1}}} + \\ & \quad \frac{d}{d(p + 1)(bc - ad)} \\ & \quad \downarrow 79 \\ & \frac{(a + bx)^{p+1}(de - cf)(c + dx)^{-p-1}}{d(p + 1)(bc - ad)} - \\ & \frac{f(a + bx)^p(c + dx)^{-p} \left(-\frac{d(a+bx)}{bc-ad}\right)^{-p} \text{Hypergeometric2F1}\left(-p, -p, 1 - p, \frac{b(c+dx)}{bc-ad}\right)}{d^2 p} \end{aligned}$$

input `Int[(a + b*x)^p*(c + d*x)^(-2 - p)*(e + f*x),x]`

output
$$\frac{((d*e - c*f)*(a + b*x)^{(1 + p)}*(c + d*x)^{(-1 - p)})/(d*(b*c - a*d)*(1 + p)) - (f*(a + b*x)^p*Hypergeometric2F1[-p, -p, 1 - p, (b*(c + d*x))/(b*c - a*d)])/(d^2*p*(-((d*(a + b*x))/(b*c - a*d)))^p*(c + d*x)^p}$$

Defintions of rubi rules used

rule 79
$$\text{Int}[(a + b*x)^m*(c + d*x)^n, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{m+1}/(b*(m+1)*(b*(c + d*x)/(b*c - a*d))^n)*Hypergeometric2F1[-n, m+1, m+2, (-d)*(a + b*x)/(b*c - a*d)], x] /;$$
 FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

rule 80
$$\text{Int}[(a + b*x)^m*(c + d*x)^n, x_Symbol] \rightarrow \text{Simp}[(c + d*x)^{\text{FracPart}[n]}/((b/(b*c - a*d))^{\text{IntPart}[n]}*(b*((c + d*x)/(b*c - a*d)))^{\text{FracPart}[n]}) \text{Int}[(a + b*x)^m*\text{Simp}[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n, x], x] /;$$
 FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])

rule 88
$$\text{Int}[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x] \rightarrow \text{Simp}[(-b*e - a*f)*(c + d*x)^{n+1}*(e + f*x)^{p+1}/(f*(p+1)*(c*f - d*e)), x] - \text{Simp}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) \text{Int}[(c + d*x)^n*(e + f*x)^{\text{Simplify}[p + 1]}, x], x] /;$$
 FreeQ[{a, b, c, d, e, f, n, p}, x] && !RationalQ[p] && SumSimplerQ[p, 1]

Maple [F]

$$\int (bx + a)^p (xd + c)^{-2-p} (fx + e) dx$$

input
$$\text{int}((b*x+a)^p*(d*x+c)^{-2-p}*(f*x+e), x)$$

output
$$\text{int}((b*x+a)^p*(d*x+c)^{-2-p}*(f*x+e), x)$$

Fricas [F]

$$\int (a + bx)^p (c + dx)^{-2-p} (e + fx) dx = \int (fx + e)(bx + a)^p (dx + c)^{-p-2} dx$$

input `integrate((b*x+a)^p*(d*x+c)^(-2-p)*(f*x+e),x, algorithm="fricas")`

output `integral((f*x + e)*(b*x + a)^p*(d*x + c)^(-p - 2), x)`

Sympy [F]

$$\int (a + bx)^p (c + dx)^{-2-p} (e + fx) dx = \int (a + bx)^p (c + dx)^{-p-2} (e + fx) dx$$

input `integrate((b*x+a)**p*(d*x+c)**(-2-p)*(f*x+e),x)`

output `Integral((a + b*x)**p*(c + d*x)**(-p - 2)*(e + f*x), x)`

Maxima [F]

$$\int (a + bx)^p (c + dx)^{-2-p} (e + fx) dx = \int (fx + e)(bx + a)^p (dx + c)^{-p-2} dx$$

input `integrate((b*x+a)^p*(d*x+c)^(-2-p)*(f*x+e),x, algorithm="maxima")`

output `integrate((f*x + e)*(b*x + a)^p*(d*x + c)^(-p - 2), x)`

Giac [F]

$$\int (a + bx)^p (c + dx)^{-2-p} (e + fx) dx = \int (fx + e)(bx + a)^p (dx + c)^{-p-2} dx$$

input `integrate((b*x+a)^p*(d*x+c)^(-2-p)*(f*x+e),x, algorithm="giac")`

output `integrate((f*x + e)*(b*x + a)^p*(d*x + c)^(-p - 2), x)`

Mupad [F(-1)]

Timed out.

$$\int (a + bx)^p (c + dx)^{-2-p} (e + fx) dx = \int \frac{(e + fx)(a + bx)^p}{(c + dx)^{p+2}} dx$$

input `int(((e + f*x)*(a + b*x)^p)/(c + d*x)^(p + 2),x)`

output `int(((e + f*x)*(a + b*x)^p)/(c + d*x)^(p + 2), x)`

Reduce [F]

$$\begin{aligned} & \int (a + bx)^p (c + dx)^{-2-p} (e + fx) dx \\ &= \left(\int \frac{(bx + a)^p}{(dx + c)^p c^2 + 2(dx + c)^p c dx + (dx + c)^p d^2 x^2} dx \right) e \\ & \quad + \left(\int \frac{(bx + a)^p x}{(dx + c)^p c^2 + 2(dx + c)^p c dx + (dx + c)^p d^2 x^2} dx \right) f \end{aligned}$$

input `int((b*x+a)^p*(d*x+c)^(-2-p)*(f*x+e),x)`

output

```
int((a + b*x)**p/((c + d*x)**p*c**2 + 2*(c + d*x)**p*c*d*x + (c + d*x)**p*
d**2*x**2),x)*e + int(((a + b*x)**p*x)/((c + d*x)**p*c**2 + 2*(c + d*x)**p
*c*d*x + (c + d*x)**p*d**2*x**2),x)*f
```

3.251 $\int (a+bx)^m (ac-bcx)^n (ad(m-n) - bd(2+m+n)x) dx$

Optimal result	2205
Mathematica [A] (verified)	2205
Rubi [A] (verified)	2206
Maple [A] (verified)	2206
Fricas [A] (verification not implemented)	2207
Sympy [B] (verification not implemented)	2208
Maxima [A] (verification not implemented)	2208
Giac [A] (verification not implemented)	2209
Mupad [B] (verification not implemented)	2209
Reduce [B] (verification not implemented)	2210

Optimal result

Integrand size = 37, antiderivative size = 30

$$\int (a+bx)^m (ac-bcx)^n (ad(m-n) - bd(2+m+n)x) dx = \frac{d(a+bx)^{1+m} (ac-bcx)^{1+n}}{bc}$$

output `d*(b*x+a)^(1+m)*(-b*c*x+a*c)^(1+n)/b/c`

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\begin{aligned} & \int (a+bx)^m (ac-bcx)^n (ad(m-n) - bd(2+m+n)x) dx \\ &= \frac{d(a-bx)(c(a-bx))^n (a+bx)^{1+m}}{b} \end{aligned}$$

input `Integrate[(a + b*x)^m*(a*c - b*c*x)^n*(a*d*(m - n) - b*d*(2 + m + n)*x),x]`

output `(d*(a - b*x)*(c*(a - b*x))^n*(a + b*x)^(1 + m))/b`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.027$, Rules used = {83}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^m (ac - bcx)^n (ad(m - n) - bdx(m + n + 2)) dx$$

$$\downarrow 83$$

$$\frac{d(a + bx)^{m+1} (ac - bcx)^{n+1}}{bc}$$

input `Int[(a + b*x)^m*(a*c - b*c*x)^n*(a*d*(m - n) - b*d*(2 + m + n)*x),x]`

output `(d*(a + b*x)^(1 + m)*(a*c - b*c*x)^(1 + n))/(b*c)`

Defintions of rubi rules used

rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`

Maple [A] (verified)

Time = 0.89 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.07

method	result
gospers	$\frac{d(-bx+a)(bx+a)^{1+m}(-bcx+ac)^n}{b}$
paralelrisch	$-\frac{x^2(bx+a)^m(c(-bx+a))^n b^3 d - (bx+a)^m(c(-bx+a))^n a^2 bd}{b^2}$
oring	$\frac{(-bx+a)(bx+a)(bx+a)^m(-bcx+ac)^n(ad(m-n)-bd(2+m+n)x)}{b(-bmx-bxn+am-an-2bx)}$
risch	$\frac{(bx+a)^m d(-b^2x^2+a^2)c^n(-bx+a)^n e^{-\frac{i\pi \operatorname{csgn}(ic(-bx+a))n(-\operatorname{csgn}(ic(-bx+a))+\operatorname{csgn}(i(-bx+a)))}{2}}(-\operatorname{csgn}(ic(-bx+a))+\operatorname{csgn}(ic))}{b}$

input `int((b*x+a)^m*(-b*c*x+a*c)^n*(a*d*(m-n)-b*d*(2+m+n)*x),x,method=_RETURNVERBOSE)`

output `d/b*(-b*x+a)*(b*x+a)^(1+m)*(-b*c*x+a*c)^n`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.27

$$\int (a+bx)^m (ac-bcx)^n (ad(m-n)-bd(2+m+n)x) dx$$

$$= -\frac{(b^2dx^2-a^2d)(-bcx+ac)^n(bx+a)^m}{b}$$

input `integrate((b*x+a)^m*(-b*c*x+a*c)^n*(a*d*(m-n)-b*d*(2+m+n)*x),x,algorithm="fricas")`

output `-(b^2*d*x^2 - a^2*d)*(-b*c*x + a*c)^n*(b*x + a)^m/b`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 63 vs. $2(24) = 48$.

Time = 1.28 (sec) , antiderivative size = 63, normalized size of antiderivative = 2.10

$$\int (a + bx)^m (ac - bcx)^n (ad(m - n) - bd(2 + m + n)x) dx$$

$$= \begin{cases} \frac{a^2 d (a + bx)^m (ac - bcx)^n}{b} - b d x^2 (a + bx)^m (ac - bcx)^n & \text{for } b \neq 0 \\ a a^m dx (ac)^n (m - n) & \text{otherwise} \end{cases}$$

input `integrate((b*x+a)**m*(-b*c*x+a*c)**n*(a*d*(m-n)-b*d*(2+m+n)*x),x)`

output `Piecewise((a**2*d*(a + b*x)**m*(a*c - b*c*x)**n/b - b*d*x**2*(a + b*x)**m*(a*c - b*c*x)**n, Ne(b, 0)), (a*a**m*d*x*(a*c)**n*(m - n), True))`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.50

$$\int (a + bx)^m (ac - bcx)^n (ad(m - n) - bd(2 + m + n)x) dx$$

$$= -\frac{(b^2 c^n dx^2 - a^2 c^n d) e^{(m \log(bx+a) + n \log(-bx+a))}}{b}$$

input `integrate((b*x+a)^m*(-b*c*x+a*c)^n*(a*d*(m-n)-b*d*(2+m+n)*x),x, algorithm="maxima")`

output `-(b^2*c^n*d*x^2 - a^2*c^n*d)*e^(m*log(b*x + a) + n*log(-b*x + a))/b`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.87

$$\int (a + bx)^m (ac - bcx)^n (ad(m - n) - bd(2 + m + n)x) dx$$

$$= -\frac{(-bcx + ac)^n (bx + a)^m b^2 dx^2 - (-bcx + ac)^n (bx + a)^m a^2 d}{b}$$

input `integrate((b*x+a)^m*(-b*c*x+a*c)^n*(a*d*(m-n)-b*d*(2+m+n)*x),x, algorithm="giac")`

output `-((-b*c*x + a*c)^n*(b*x + a)^m*b^2*d*x^2 - (-b*c*x + a*c)^n*(b*x + a)^m*a^2*d)/b`

Mupad [B] (verification not implemented)

Time = 1.18 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.17

$$\int (a + bx)^m (ac - bcx)^n (ad(m - n) - bd(2 + m + n)x) dx$$

$$= \frac{d(a^2 - b^2 x^2) (ac - bcx)^n (a + bx)^m}{b}$$

input `int((a*c - b*c*x)^n*(a*d*(m - n) - b*d*x*(m + n + 2))*(a + b*x)^m,x)`

output `(d*(a^2 - b^2*x^2)*(a*c - b*c*x)^n*(a + b*x)^m)/b`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.17

$$\int (a + bx)^m (ac - bcx)^n (ad(m - n) - bd(2 + m + n)x) dx$$

$$= \frac{(bx + a)^m (-bcx + ac)^n d(-b^2x^2 + a^2)}{b}$$

input `int((b*x+a)^m*(-b*c*x+a*c)^n*(a*d*(m-n)-b*d*(2+m+n)*x),x)`output `((a + b*x)**m*(a*c - b*c*x)**n*d*(a**2 - b**2*x**2))/b`

3.252 $\int (a + bx)(c + dx)^n(e + fx)^{-n} dx$

Optimal result	2211
Mathematica [A] (verified)	2211
Rubi [A] (verified)	2212
Maple [F]	2213
Fricas [F]	2214
Sympy [F(-2)]	2214
Maxima [F]	2214
Giac [F]	2215
Mupad [F(-1)]	2215
Reduce [F]	2215

Optimal result

Integrand size = 22, antiderivative size = 125

$$\int (a + bx)(c + dx)^n(e + fx)^{-n} dx = \frac{b(c + dx)^{1+n}(e + fx)^{1-n}}{2df} + \frac{(2adf - bcf(1 - n) - bde(1 + n))(c + dx)^{1+n}(e + fx)^{1-n} \text{Hypergeometric2F1}\left(1, 2, 2 + n, -\frac{f(c+dx)}{de-cf}\right)}{2df(de - cf)(1 + n)}$$

output

```
1/2*b*(d*x+c)^(1+n)*(f*x+e)^(1-n)/d/f+1/2*(2*a*d*f-b*c*f*(1-n)-b*d*e*(1+n)
)*(d*x+c)^(1+n)*(f*x+e)^(1-n)*hypergeom([1, 2], [2+n], -f*(d*x+c)/(-c*f+d*e)
)/d/f/(-c*f+d*e)/(1+n)
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.87

$$\int (a + bx)(c + dx)^n(e + fx)^{-n} dx = \frac{(c + dx)^{1+n}(e + fx)^{-n} \left(bd(e + fx) - \frac{(-2adf - bcf(-1+n) + bde(1+n)) \left(\frac{d(e+fx)}{de-cf}\right)^n \text{Hypergeometric2F1}\left(n, 1+n, 2+n, \frac{f(c+dx)}{-de+cf}\right)}{1+n} \right)}{2d^2f}$$

input

```
Integrate[((a + b*x)*(c + d*x)^n)/(e + f*x)^n,x]
```


output

$$\left((c + dx)^{(1+n)} (bd(e+fx) - (-2ad - bcf(-1+n) + bde(1+n)) \left(\frac{d(e+fx)}{de-cf} \right)^n \text{Hypergeometric2F1}[n, 1+n, 2+n, (f(c+dx)/(-d*e) + c*f)] / (1+n)) \right) / (2d^2 f (e+fx)^n)$$
Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.07, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.136$, Rules used = {90, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)(c + dx)^n (e + fx)^{-n} dx$$

$$\downarrow 90$$

$$\frac{(2adf - bcf(1 - n) - bde(n + 1)) \int (c + dx)^n (e + fx)^{-n} dx}{2df} + \frac{b(c + dx)^{n+1} (e + fx)^{1-n}}{2df}$$

$$\downarrow 80$$

$$\frac{(e + fx)^{-n} \left(\frac{d(e+fx)}{de-cf} \right)^n (2adf - bcf(1 - n) - bde(n + 1)) \int (c + dx)^n \left(\frac{de}{de-cf} + \frac{dfx}{de-cf} \right)^{-n} dx}{2df} + \frac{b(c + dx)^{n+1} (e + fx)^{1-n}}{2df}$$

$$\downarrow 79$$

$$\frac{(c + dx)^{n+1} (e + fx)^{-n} \left(\frac{d(e+fx)}{de-cf} \right)^n (2adf - bcf(1 - n) - bde(n + 1)) \text{Hypergeometric2F1} \left(n, n + 1, n + 2, -\frac{f(c+dx)}{de-cf} \right)}{2d^2 f (n + 1)} + \frac{b(c + dx)^{n+1} (e + fx)^{1-n}}{2df}$$

input

$$\text{Int}[(a + b*x)*(c + d*x)^n/(e + f*x)^n, x]$$

output

```
(b*(c + d*x)^(1 + n)*(e + f*x)^(1 - n))/(2*d*f) + ((2*a*d*f - b*c*f*(1 - n) - b*d*e*(1 + n))*(c + d*x)^(1 + n)*((d*(e + f*x))/(d*e - c*f))^n*Hypergeometric2F1[n, 1 + n, 2 + n, -(f*(c + d*x)/(d*e - c*f))]/(2*d^2*f*(1 + n))*(e + f*x)^n)
```

Defintions of rubi rules used

rule 79

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))
```

rule 80

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(c + d*x)^FracPart[n]/((b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])
```

rule 90

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]
```

Maple [F]

$$\int (bx + a)(xd + c)^n (fx + e)^{-n} dx$$

input

```
int((b*x+a)*(d*x+c)^n/((f*x+e)^n),x)
```

output

```
int((b*x+a)*(d*x+c)^n/((f*x+e)^n),x)
```

Fricas [F]

$$\int (a + bx)(c + dx)^n (e + fx)^{-n} dx = \int \frac{(bx + a)(dx + c)^n}{(fx + e)^n} dx$$

input `integrate((b*x+a)*(d*x+c)^n/((f*x+e)^n),x, algorithm="fricas")`

output `integral((b*x + a)*(d*x + c)^n/(f*x + e)^n, x)`

Sympy [F(-2)]

Exception generated.

$$\int (a + bx)(c + dx)^n (e + fx)^{-n} dx = \text{Exception raised: HeuristicGCDFailed}$$

input `integrate((b*x+a)*(d*x+c)**n/((f*x+e)**n),x)`

output `Exception raised: HeuristicGCDFailed >> no luck`

Maxima [F]

$$\int (a + bx)(c + dx)^n (e + fx)^{-n} dx = \int \frac{(bx + a)(dx + c)^n}{(fx + e)^n} dx$$

input `integrate((b*x+a)*(d*x+c)^n/((f*x+e)^n),x, algorithm="maxima")`

output `integrate((b*x + a)*(d*x + c)^n/(f*x + e)^n, x)`

Giac [F]

$$\int (a + bx)(c + dx)^n (e + fx)^{-n} dx = \int \frac{(bx + a)(dx + c)^n}{(fx + e)^n} dx$$

input `integrate((b*x+a)*(d*x+c)^n/((f*x+e)^n),x, algorithm="giac")`

output `integrate((b*x + a)*(d*x + c)^n/(f*x + e)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + bx)(c + dx)^n (e + fx)^{-n} dx = \int \frac{(a + bx)(c + dx)^n}{(e + fx)^n} dx$$

input `int(((a + b*x)*(c + d*x)^n)/(e + f*x)^n,x)`

output `int(((a + b*x)*(c + d*x)^n)/(e + f*x)^n, x)`

Reduce [F]

$$\int (a + bx)(c + dx)^n (e + fx)^{-n} dx = \left(\int \frac{(dx + c)^n}{(fx + e)^n} dx \right) a + \left(\int \frac{(dx + c)^n x}{(fx + e)^n} dx \right) b$$

input `int((b*x+a)*(d*x+c)^n/((f*x+e)^n),x)`

output `int((c + d*x)**n/(e + f*x)**n,x)*a + int(((c + d*x)**n*x)/(e + f*x)**n,x)*b`

3.253 $\int (a + bx)(c + dx)^{-1+n}(e + fx)^{-n} dx$

Optimal result	2216
Mathematica [A] (verified)	2216
Rubi [A] (verified)	2217
Maple [F]	2218
Fricas [F]	2219
Sympy [F(-2)]	2219
Maxima [F]	2219
Giac [F]	2220
Mupad [F(-1)]	2220
Reduce [F]	2220

Optimal result

Integrand size = 24, antiderivative size = 110

$$\int (a + bx)(c + dx)^{-1+n}(e + fx)^{-n} dx = \frac{b(c + dx)^n(e + fx)^{1-n}}{df} + \frac{(adf - bcf(1 - n) - bden)(c + dx)^n(e + fx)^{1-n} \operatorname{Hypergeometric2F1}\left(1, 1, 1 + n, -\frac{f(c+dx)}{de-cf}\right)}{df(de - cf)n}$$

output

```
b*(d*x+c)^n*(f*x+e)^(1-n)/d/f+(a*d*f-b*c*f*(1-n)-b*d*e*n)*(d*x+c)^n*(f*x+e)^(1-n)*hypergeom([1, 1],[1+n],-f*(d*x+c)/(-c*f+d*e))/d/f/(-c*f+d*e)/n
```

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.13

$$\int (a + bx)(c + dx)^{-1+n}(e + fx)^{-n} dx = \frac{(c + dx)^n(e + fx)^{-n} \left(d(-bc + ad)(e + fx) + \frac{(-adf + bden + bc(f - fn))(c + dx) \left(\frac{d(e + fx)}{de - cf} \right)^n \operatorname{Hypergeometric2F1}(n, 1 + n, 2 + n, -\frac{d(e + fx)}{de - cf})}{1 + n} \right)}{d^2(de - cf)n}$$

input

```
Integrate[((a + b*x)*(c + d*x)^(-1 + n))/(e + f*x)^n,x]
```

output

$$\frac{((c + dx)^n (d(-bc) + ad)(e + fx) + ((-adf) + bde^n + bc(f - fn)) * (c + dx) * ((d(e + fx))/(de - cf))^n \text{Hypergeometric2F1}[n, 1 + n, 2 + n, (f(c + dx))/(-d(e) + cf)] / (1 + n)) / (d^2 * (de - cf) * n * (e + fx)^n)}$$
Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.38, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {88, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)(c + dx)^{n-1}(e + fx)^{-n} dx$$

$$\downarrow 88$$

$$\frac{(adf - bcf(1 - n) - bden) \int (c + dx)^n (e + fx)^{-n} dx}{dn(de - cf)} - \frac{(bc - ad)(c + dx)^n (e + fx)^{1-n}}{dn(de - cf)}$$

$$\downarrow 80$$

$$\frac{(e + fx)^{-n} \left(\frac{d(e+fx)}{de-cf}\right)^n (adf - bcf(1 - n) - bden) \int (c + dx)^n \left(\frac{de}{de-cf} + \frac{dfx}{de-cf}\right)^{-n} dx}{dn(de - cf) \frac{(bc - ad)(c + dx)^n (e + fx)^{1-n}}{dn(de - cf)}}$$

$$\downarrow 79$$

$$\frac{(c + dx)^{n+1} (e + fx)^{-n} \left(\frac{d(e+fx)}{de-cf}\right)^n (adf - bcf(1 - n) - bden) \text{Hypergeometric2F1}\left(n, n + 1, n + 2, -\frac{f(c+dx)}{de-cf}\right)}{\frac{d^2 n(n + 1)(de - cf)}{(bc - ad)(c + dx)^n (e + fx)^{1-n}} dn(de - cf)}$$

input

$$\text{Int}[\frac{(a + b*x)*(c + d*x)^{-1 + n}}{(e + f*x)^n}, x]$$

output
$$-\left(\frac{(b*c - a*d)*(c + d*x)^n*(e + f*x)^{(1-n)}}{d*(d*e - c*f)*n}\right) - \left(\frac{a*d*f - b*c*f*(1-n) - b*d*e*n}{d*(d*e - c*f)}\right) * (c + d*x)^{(1+n)} * \frac{d*(e + f*x)}{d*e - c*f} * \text{Hypergeometric2F1}[n, 1+n, 2+n, -\frac{f*(c + d*x)}{d*e - c*f}] / (d^2*(d*e - c*f)*n*(1+n)*(e + f*x)^n)$$

Defintions of rubi rules used

rule 79
$$\text{Int}[(a + b*x)^m * (c + d*x)^n, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{m+1} / (b*(m+1)*(b*(c - a*d))^n) * \text{Hypergeometric2F1}[-n, m+1, m+2, (-d)*(a + b*x)/(b*c - a*d)], x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x \&\& \text{!IntegerQ}[m] \&\& \text{!IntegerQ}[n] \&\& \text{GtQ}[b/(b*c - a*d), 0] \&\& (\text{RationalQ}[m] \parallel \text{!RationalQ}[n] \&\& \text{GtQ}[-d/(b*c - a*d), 0])$$

rule 80
$$\text{Int}[(a + b*x)^m * (c + d*x)^n, x_Symbol] \rightarrow \text{Simp}[(c + d*x)^{\text{FracPart}[n]} / (b*(b*c - a*d))^{\text{IntPart}[n]} * (b*(c + d*x)/(b*c - a*d))^{\text{FracPart}[n]} \text{Int}[(a + b*x)^m * \text{Simp}[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n, x], x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x \&\& \text{!IntegerQ}[m] \&\& \text{!IntegerQ}[n] \&\& (\text{RationalQ}[m] \parallel \text{!SimplerQ}[n + 1, m + 1])$$

rule 88
$$\text{Int}[(a + b*x)^m * (c + d*x)^n * (e + f*x)^p, x] \rightarrow \text{Simp}[-(b*e - a*f)*(c + d*x)^{n+1} * (e + f*x)^{p+1} / (f*(p+1)*(c*f - d*e)), x] - \text{Simp}[(a*d*f*(n+p+2) - b*(d*e*(n+1) + c*f*(p+1))] / (f*(p+1)*(c*f - d*e)) \text{Int}[(c + d*x)^n * (e + f*x)^{\text{Simplify}[p+1]}, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n, p\}, x \&\& \text{!RationalQ}[p] \&\& \text{SumSimplerQ}[p, 1]$$

Maple [F]

$$\int (bx + a)(xd + c)^{-1+n}(fx + e)^{-n} dx$$

input
$$\text{int}((b*x+a)*(d*x+c)^{-1+n}/((f*x+e)^n), x)$$

output
$$\text{int}((b*x+a)*(d*x+c)^{-1+n}/((f*x+e)^n), x)$$

Fricas [F]

$$\int (a + bx)(c + dx)^{-1+n}(e + fx)^{-n} dx = \int \frac{(bx + a)(dx + c)^{n-1}}{(fx + e)^n} dx$$

input `integrate((b*x+a)*(d*x+c)^(-1+n)/((f*x+e)^n),x, algorithm="fricas")`

output `integral((b*x + a)*(d*x + c)^(n - 1)/(f*x + e)^n, x)`

Sympy [F(-2)]

Exception generated.

$$\int (a + bx)(c + dx)^{-1+n}(e + fx)^{-n} dx = \text{Exception raised: HeuristicGCDFailed}$$

input `integrate((b*x+a)*(d*x+c)**(-1+n)/((f*x+e)**n),x)`

output `Exception raised: HeuristicGCDFailed >> no luck`

Maxima [F]

$$\int (a + bx)(c + dx)^{-1+n}(e + fx)^{-n} dx = \int \frac{(bx + a)(dx + c)^{n-1}}{(fx + e)^n} dx$$

input `integrate((b*x+a)*(d*x+c)^(-1+n)/((f*x+e)^n),x, algorithm="maxima")`

output `integrate((b*x + a)*(d*x + c)^(n - 1)/(f*x + e)^n, x)`

Giac [F]

$$\int (a + bx)(c + dx)^{-1+n}(e + fx)^{-n} dx = \int \frac{(bx + a)(dx + c)^{n-1}}{(fx + e)^n} dx$$

input `integrate((b*x+a)*(d*x+c)^(-1+n)/((f*x+e)^n),x, algorithm="giac")`

output `integrate((b*x + a)*(d*x + c)^(n - 1)/(f*x + e)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + bx)(c + dx)^{-1+n}(e + fx)^{-n} dx = \int \frac{(a + bx)(c + dx)^{n-1}}{(e + fx)^n} dx$$

input `int(((a + b*x)*(c + d*x)^(n - 1))/(e + f*x)^n,x)`

output `int(((a + b*x)*(c + d*x)^(n - 1))/(e + f*x)^n, x)`

Reduce [F]

$$\int (a + bx)(c + dx)^{-1+n}(e + fx)^{-n} dx = \left(\int \frac{(dx + c)^n}{(fx + e)^n c + (fx + e)^n dx} dx \right) a + \left(\int \frac{(dx + c)^n x}{(fx + e)^n c + (fx + e)^n dx} dx \right) b$$

input `int((b*x+a)*(d*x+c)^(-1+n)/((f*x+e)^n),x)`

output `int((c + d*x)**n/((e + f*x)**n*c + (e + f*x)**n*d*x),x)*a + int(((c + d*x)**n*x)/((e + f*x)**n*c + (e + f*x)**n*d*x),x)*b`

3.254 $\int (a + bx)(c + dx)^{-2+n}(e + fx)^{-n} dx$

Optimal result	2221
Mathematica [A] (verified)	2221
Rubi [A] (verified)	2222
Maple [F]	2223
Fricas [F]	2224
Sympy [F(-2)]	2224
Maxima [F]	2224
Giac [F]	2225
Mupad [F(-1)]	2225
Reduce [F]	2225

Optimal result

Integrand size = 24, antiderivative size = 110

$$\int (a + bx)(c + dx)^{-2+n}(e + fx)^{-n} dx$$

$$= \frac{(bc - ad)(c + dx)^{-1+n}(e + fx)^{1-n}}{d(de - cf)(1 - n)}$$

$$+ \frac{b(c + dx)^n(e + fx)^{1-n} \text{Hypergeometric2F1}\left(1, 1, 1 + n, -\frac{f(c+dx)}{de-cf}\right)}{d(de - cf)n}$$

output

```
(-a*d+b*c)*(d*x+c)^(-1+n)*(f*x+e)^(1-n)/d/(-c*f+d*e)/(1-n)+b*(d*x+c)^n*(f*x+e)^(1-n)*hypergeom([1, 1], [1+n], -f*(d*x+c)/(-c*f+d*e))/d/(-c*f+d*e)/n
```

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.05

$$\int (a + bx)(c + dx)^{-2+n}(e + fx)^{-n} dx$$

$$= \frac{(c + dx)^{-1+n}(e + fx)^{-n} \left(d^2 (be - af)(e + fx) - b(de - cf)^2 \left(\frac{d(e+fx)}{de-cf} \right)^n \text{Hypergeometric2F1}\left(-1 + n, -\right)}{d^2 f(-de + cf)(-1 + n)}$$

input `Integrate[((a + b*x)*(c + d*x)^(-2 + n))/(e + f*x)^n,x]`

output `((c + d*x)^(-1 + n)*(d^2*(b*e - a*f)*(e + f*x) - b*(d*e - c*f)^2*((d*(e + f*x))/(d*e - c*f))^n*Hypergeometric2F1[-1 + n, -1 + n, n, (f*(c + d*x))/(-(d*e) + c*f)]))/(d^2*f*(-(d*e) + c*f)*(-1 + n)*(e + f*x)^n)`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.06, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {88, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + bx)(c + dx)^{n-2}(e + fx)^{-n} dx \\
 & \quad \downarrow 88 \\
 & \frac{b \int (c + dx)^{n-1}(e + fx)^{-n} dx}{d} + \frac{(bc - ad)(c + dx)^{n-1}(e + fx)^{1-n}}{d(1 - n)(de - cf)} \\
 & \quad \downarrow 80 \\
 & \frac{b(e + fx)^{-n} \left(\frac{d(e+fx)}{de-cf} \right)^n \int (c + dx)^{n-1} \left(\frac{de}{de-cf} + \frac{dfx}{de-cf} \right)^{-n} dx}{\frac{(bc - ad)(c + dx)^{n-1}(e + fx)^{1-n}}{d(1 - n)(de - cf)}} + \\
 & \quad \downarrow 79 \\
 & \frac{(bc - ad)(c + dx)^{n-1}(e + fx)^{1-n}}{d(1 - n)(de - cf)} + \\
 & \frac{b(c + dx)^n(e + fx)^{-n} \left(\frac{d(e+fx)}{de-cf} \right)^n \text{Hypergeometric2F1} \left(n, n, n + 1, -\frac{f(c+dx)}{de-cf} \right)}{d^2 n}
 \end{aligned}$$

input `Int[((a + b*x)*(c + d*x)^(-2 + n))/(e + f*x)^n,x]`

output $((b*c - a*d)*(c + d*x)^{-1 + n}*(e + f*x)^{(1 - n)})/(d*(d*e - c*f)*(1 - n)) + (b*(c + d*x)^n*((d*(e + f*x))/(d*e - c*f))^n*Hypergeometric2F1[n, n, 1 + n, -((f*(c + d*x))/(d*e - c*f))]/(d^{2*n}*(e + f*x)^n)$

Defintions of rubi rules used

rule 79 $\text{Int}[(a + b*x)^m*(c + d*x)^n, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{m+1}/(b*(m+1)*(b*(c - a*d))^n)*Hypergeometric2F1[-n, m+1, m+2, (-d)*(a + b*x)/(b*c - a*d)], x] /;$ $\text{FreeQ}\{a, b, c, d, m, n\}, x$ $\&\& \text{!IntegerQ}[m] \&\& \text{!IntegerQ}[n] \&\& \text{GtQ}[b/(b*c - a*d), 0] \&\& (\text{RationalQ}[m] \parallel \text{!RationalQ}[n] \&\& \text{GtQ}[-d/(b*c - a*d), 0])$

rule 80 $\text{Int}[(a + b*x)^m*(c + d*x)^n, x_Symbol] \rightarrow \text{Simp}[(c + d*x)^{\text{FracPart}[n]}/((b/(b*c - a*d))^{\text{IntPart}[n]}*(b*((c + d*x)/(b*c - a*d)))^{\text{FracPart}[n]}) \text{Int}[(a + b*x)^m*\text{Simp}[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n, x], x] /;$ $\text{FreeQ}\{a, b, c, d, m, n\}, x$ $\&\& \text{!IntegerQ}[m] \&\& \text{!IntegerQ}[n] \&\& (\text{RationalQ}[m] \parallel \text{!SimplerQ}[n + 1, m + 1])$

rule 88 $\text{Int}[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x] \rightarrow \text{Simp}[-(b*e - a*f)*(c + d*x)^{n+1}*(e + f*x)^{p+1}/(f*(p+1)*(c*f - d*e)), x] - \text{Simp}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1))]/(f*(p + 1)*(c*f - d*e)) \text{Int}[(c + d*x)^n*(e + f*x)^{\text{Simplify}[p + 1]}, x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f, n, p\}, x$ $\&\& \text{!RationalQ}[p] \&\& \text{SumSimplerQ}[p, 1]$

Maple [F]

$$\int (bx + a)(xd + c)^{n-2}(fx + e)^{-n} dx$$

input $\text{int}((b*x+a)*(d*x+c)^{(n-2)}((f*x+e)^n), x)$

output $\text{int}((b*x+a)*(d*x+c)^{(n-2)}((f*x+e)^n), x)$

Fricas [F]

$$\int (a + bx)(c + dx)^{-2+n}(e + fx)^{-n} dx = \int \frac{(bx + a)(dx + c)^{n-2}}{(fx + e)^n} dx$$

input `integrate((b*x+a)*(d*x+c)^(-2+n)/((f*x+e)^n),x, algorithm="fricas")`

output `integral((b*x + a)*(d*x + c)^(n - 2)/(f*x + e)^n, x)`

Sympy [F(-2)]

Exception generated.

$$\int (a + bx)(c + dx)^{-2+n}(e + fx)^{-n} dx = \text{Exception raised: HeuristicGCDFailed}$$

input `integrate((b*x+a)*(d*x+c)**(-2+n)/((f*x+e)**n),x)`

output `Exception raised: HeuristicGCDFailed >> no luck`

Maxima [F]

$$\int (a + bx)(c + dx)^{-2+n}(e + fx)^{-n} dx = \int \frac{(bx + a)(dx + c)^{n-2}}{(fx + e)^n} dx$$

input `integrate((b*x+a)*(d*x+c)^(-2+n)/((f*x+e)^n),x, algorithm="maxima")`

output `integrate((b*x + a)*(d*x + c)^(n - 2)/(f*x + e)^n, x)`

Giac [F]

$$\int (a + bx)(c + dx)^{-2+n}(e + fx)^{-n} dx = \int \frac{(bx + a)(dx + c)^{n-2}}{(fx + e)^n} dx$$

input `integrate((b*x+a)*(d*x+c)^(-2+n)/((f*x+e)^n),x, algorithm="giac")`

output `integrate((b*x + a)*(d*x + c)^(n - 2)/(f*x + e)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + bx)(c + dx)^{-2+n}(e + fx)^{-n} dx = \int \frac{(a + bx)(c + dx)^{n-2}}{(e + fx)^n} dx$$

input `int(((a + b*x)*(c + d*x)^(n - 2))/(e + f*x)^n,x)`

output `int(((a + b*x)*(c + d*x)^(n - 2))/(e + f*x)^n, x)`

Reduce [F]

$$\begin{aligned} & \int (a + bx)(c + dx)^{-2+n}(e + fx)^{-n} dx \\ &= \left(\int \frac{(dx + c)^n}{(fx + e)^n c^2 + 2(fx + e)^n cdx + (fx + e)^n d^2x^2} dx \right) a \\ &+ \left(\int \frac{(dx + c)^n x}{(fx + e)^n c^2 + 2(fx + e)^n cdx + (fx + e)^n d^2x^2} dx \right) b \end{aligned}$$

input `int((b*x+a)*(d*x+c)^(-2+n)/((f*x+e)^n),x)`

output

```
int((c + d*x)**n/((e + f*x)**n*c**2 + 2*(e + f*x)**n*c*d*x + (e + f*x)**n*
d**2*x**2),x)*a + int(((c + d*x)**n*x)/((e + f*x)**n*c**2 + 2*(e + f*x)**n
*c*d*x + (e + f*x)**n*d**2*x**2),x)*b
```

3.255 $\int (a + bx)(c + dx)^{-3+n}(e + fx)^{-n} dx$

Optimal result	2227
Mathematica [A] (verified)	2227
Rubi [A] (verified)	2228
Maple [A] (verified)	2229
Fricas [B] (verification not implemented)	2230
Sympy [F(-2)]	2230
Maxima [F]	2231
Giac [B] (verification not implemented)	2231
Mupad [B] (verification not implemented)	2232
Reduce [F]	2233

Optimal result

Integrand size = 24, antiderivative size = 122

$$\int (a + bx)(c + dx)^{-3+n}(e + fx)^{-n} dx$$

$$= \frac{(bc - ad)(c + dx)^{-2+n}(e + fx)^{1-n}}{d(de - cf)(2 - n)} + \frac{(adf + bcf(1 - n) - bde(2 - n))(c + dx)^{-1+n}(e + fx)^{1-n}}{d(de - cf)^2(1 - n)(2 - n)}$$

output

$$\frac{(-a*d+b*c)*(d*x+c)^{-2+n}*(f*x+e)^{(1-n)}/d/(-c*f+d*e)/(2-n)+(a*d*f+b*c*f*(1-n)-b*d*e*(2-n))*(d*x+c)^{-1+n}*(f*x+e)^{(1-n)}/d/(-c*f+d*e)^2/(1-n)/(2-n)}{d(de - cf)^2(1 - n)(2 - n)}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.67

$$\int (a + bx)(c + dx)^{-3+n}(e + fx)^{-n} dx$$

$$= \frac{(c + dx)^{-2+n}(e + fx)^{1-n}(-acf(-2 + n) + ade(-1 + n) + adfx + bde(-2 + n)x - bc(e + f(-1 + n)x))}{(de - cf)^2(-2 + n)(-1 + n)}$$

input

$$\text{Integrate}[(a + b*x)*(c + d*x)^{-3 + n})/(e + f*x)^n, x]$$

output

```
((c + d*x)^(-2 + n)*(e + f*x)^(1 - n)*(-(a*c*f*(-2 + n)) + a*d*e*(-1 + n)
+ a*d*f*x + b*d*e*(-2 + n)*x - b*c*(e + f*(-1 + n)*x)))/((d*e - c*f)^2*(-2
+ n)*(-1 + n))
```

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {88, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)(c + dx)^{n-3}(e + fx)^{-n} dx$$

$$\downarrow 88$$

$$\frac{(bc - ad)(c + dx)^{n-2}(e + fx)^{1-n}}{d(2 - n)(de - cf)} - \frac{(adf + bcf(1 - n) - bde(2 - n)) \int (c + dx)^{n-2}(e + fx)^{-n} dx}{d(2 - n)(de - cf)}$$

$$\downarrow 48$$

$$\frac{(bc - ad)(c + dx)^{n-2}(e + fx)^{1-n}}{d(2 - n)(de - cf)} + \frac{(c + dx)^{n-1}(e + fx)^{1-n}(adf + bcf(1 - n) - bde(2 - n))}{d(1 - n)(2 - n)(de - cf)^2}$$

input

```
Int[((a + b*x)*(c + d*x)^(-3 + n))/(e + f*x)^n,x]
```

output

```
((b*c - a*d)*(c + d*x)^(-2 + n)*(e + f*x)^(1 - n))/(d*(d*e - c*f)*(2 - n))
+ ((a*d*f + b*c*f*(1 - n) - b*d*e*(2 - n))*(c + d*x)^(-1 + n)*(e + f*x)^(
1 - n))/(d*(d*e - c*f)^2*(1 - n)*(2 - n))
```

Defintions of rubi rules used

```
rule 48 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 88 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^Simplify[p + 1],
x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && !RationalQ[p] && SumSimpl
erQ[p, 1]
```

Maple [A] (verified)

Time = 0.58 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.32

method	result
gospers	$-\frac{(xd+c)^{n-2}(fx+e)(fx+e)^{-n}(bcfnx-bdenx+acfn-aden-adjx-bcfx+2bdex-2acf+ade+bce)}{c^2f^2n^2-2cdefn^2+d^2e^2n^2-3c^2f^2n+6cdefn-3d^2e^2n+2c^2f^2-4cdef+2d^2e^2}$
orering	$-\frac{(fx+e)(xd+c)(bcfnx-bdenx+acfn-aden-adjx-bcfx+2bdex-2acf+ade+bce)(xd+c)^{-3+n}(fx+e)^{-n}}{c^2f^2n^2-2cdefn^2+d^2e^2n^2-3c^2f^2n+6cdefn-3d^2e^2n+2c^2f^2-4cdef+2d^2e^2}$
parallelrisch	$(x^3(xd+c)^{-3+n}bd^3ef^2n-x^2(xd+c)^{-3+n}acd^2f^3n+x^2(xd+c)^{-3+n}ad^3ef^2n-x^2(xd+c)^{-3+n}bc^2df^3n+x^2(xd+c)^{-3+n}b$

```
input int((b*x+a)*(d*x+c)^(-3+n)/((f*x+e)^n),x,method=_RETURNVERBOSE)
```

```
output -(d*x+c)^(n-2)*(f*x+e)/((f*x+e)^n)/(c^2*f^2*n^2-2*c*d*e*f*n^2+d^2*e^2*n^2-
3*c^2*f^2*n+6*c*d*e*f*n-3*d^2*e^2*n+2*c^2*f^2-4*c*d*e*f+2*d^2*e^2)*(b*c*f*
n*x-b*d*e*n*x+a*c*f*n-a*d*e*n-a*d*f*x-b*c*f*x+2*b*d*e*x-2*a*c*f+a*d*e+b*c*
e)
```


Maxima [F]

$$\int (a + bx)(c + dx)^{-3+n}(e + fx)^{-n} dx = \int \frac{(bx + a)(dx + c)^{n-3}}{(fx + e)^n} dx$$

input `integrate((b*x+a)*(d*x+c)^(-3+n)/((f*x+e)^n),x, algorithm="maxima")`

output `integrate((b*x + a)*(d*x + c)^(n - 3)/(f*x + e)^n, x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1049 vs. $2(115) = 230$.

Time = 0.15 (sec) , antiderivative size = 1049, normalized size of antiderivative = 8.60

$$\int (a + bx)(c + dx)^{-3+n}(e + fx)^{-n} dx = \text{Too large to display}$$

input `integrate((b*x+a)*(d*x+c)^(-3+n)/((f*x+e)^n),x, algorithm="giac")`

output

```
(b*d^2*e*f*n*x^3*e^(n*log(d*x + c) - 3*log(d*x + c))/(f*x + e)^n - b*c*d*f
^2*n*x^3*e^(n*log(d*x + c) - 3*log(d*x + c))/(f*x + e)^n + b*d^2*e^2*n*x^2
*e^(n*log(d*x + c) - 3*log(d*x + c))/(f*x + e)^n + a*d^2*e*f*n*x^2*e^(n*lo
g(d*x + c) - 3*log(d*x + c))/(f*x + e)^n - b*c^2*f^2*n*x^2*e^(n*log(d*x +
c) - 3*log(d*x + c))/(f*x + e)^n - a*c*d*f^2*n*x^2*e^(n*log(d*x + c) - 3*l
og(d*x + c))/(f*x + e)^n - 2*b*d^2*e*f*x^3*e^(n*log(d*x + c) - 3*log(d*x +
c))/(f*x + e)^n + b*c*d*f^2*x^3*e^(n*log(d*x + c) - 3*log(d*x + c))/(f*x
+ e)^n + a*d^2*f^2*x^3*e^(n*log(d*x + c) - 3*log(d*x + c))/(f*x + e)^n + b
*c*d*e^2*n*x*e^(n*log(d*x + c) - 3*log(d*x + c))/(f*x + e)^n + a*d^2*e^2*n
*x*e^(n*log(d*x + c) - 3*log(d*x + c))/(f*x + e)^n - b*c^2*e*f*n*x*e^(n*lo
g(d*x + c) - 3*log(d*x + c))/(f*x + e)^n - a*c^2*f^2*n*x*e^(n*log(d*x + c)
- 3*log(d*x + c))/(f*x + e)^n - 2*b*d^2*e^2*x^2*e^(n*log(d*x + c) - 3*log
(d*x + c))/(f*x + e)^n - 2*b*c*d*e*f*x^2*e^(n*log(d*x + c) - 3*log(d*x + c
))/(f*x + e)^n + b*c^2*f^2*x^2*e^(n*log(d*x + c) - 3*log(d*x + c))/(f*x +
e)^n + 3*a*c*d*f^2*x^2*e^(n*log(d*x + c) - 3*log(d*x + c))/(f*x + e)^n + a
*c*d*e^2*n*e^(n*log(d*x + c) - 3*log(d*x + c))/(f*x + e)^n - a*c^2*e*f*n*e
^(n*log(d*x + c) - 3*log(d*x + c))/(f*x + e)^n - 3*b*c*d*e^2*x*e^(n*log(d*
x + c) - 3*log(d*x + c))/(f*x + e)^n - a*d^2*e^2*x*e^(n*log(d*x + c) - 3*l
og(d*x + c))/(f*x + e)^n + 2*a*c*d*e*f*x*e^(n*log(d*x + c) - 3*log(d*x + c
))/(f*x + e)^n + 2*a*c^2*f^2*x*e^(n*log(d*x + c) - 3*log(d*x + c))/(f*x...
```

Mupad [B] (verification not implemented)

Time = 1.29 (sec) , antiderivative size = 358, normalized size of antiderivative = 2.93

$$\int (a + bx)(c + dx)^{-3+n}(e + fx)^{-n} dx$$

$$= \frac{x(c + dx)^{n-3} (2ac^2f^2 - ad^2e^2 - 3bcde^2 - ac^2f^2n + ad^2e^2n + 2acdef + bcde^2n - bc^2efn)}{(e + fx)^n (cf - de)^2 (n^2 - 3n + 2)}$$

$$+ \frac{x^2(c + dx)^{n-3} (bc^2f^2 - 2bd^2e^2 + 3acdf^2 - bc^2f^2n + bd^2e^2n - 2bcdef - acdf^2n + ad^2efn)}{(e + fx)^n (cf - de)^2 (n^2 - 3n + 2)}$$

$$- \frac{ce(c + dx)^{n-3} (ade - 2acf + bce + acfn - aden)}{(e + fx)^n (cf - de)^2 (n^2 - 3n + 2)}$$

$$+ \frac{dfx^3(c + dx)^{n-3} (adf + bcf - 2bde - bcfn + bden)}{(e + fx)^n (cf - de)^2 (n^2 - 3n + 2)}$$

input

```
int(((a + b*x)*(c + d*x)^(n - 3))/(e + f*x)^n,x)
```

output

```
(x*(c + d*x)^(n - 3)*(2*a*c^2*f^2 - a*d^2*e^2 - 3*b*c*d*e^2 - a*c^2*f^2*n
+ a*d^2*e^2*n + 2*a*c*d*e*f + b*c*d*e^2*n - b*c^2*e*f*n))/((e + f*x)^n*(c*
f - d*e)^2*(n^2 - 3*n + 2)) + (x^2*(c + d*x)^(n - 3)*(b*c^2*f^2 - 2*b*d^2*
e^2 + 3*a*c*d*f^2 - b*c^2*f^2*n + b*d^2*e^2*n - 2*b*c*d*e*f - a*c*d*f^2*n
+ a*d^2*e*f*n))/((e + f*x)^n*(c*f - d*e)^2*(n^2 - 3*n + 2)) - (c*e*(c + d*
x)^(n - 3)*(a*d*e - 2*a*c*f + b*c*e + a*c*f*n - a*d*e*n))/((e + f*x)^n*(c*
f - d*e)^2*(n^2 - 3*n + 2)) + (d*f*x^3*(c + d*x)^(n - 3)*(a*d*f + b*c*f -
2*b*d*e - b*c*f*n + b*d*e*n))/((e + f*x)^n*(c*f - d*e)^2*(n^2 - 3*n + 2))
```

Reduce [F]

$$\int (a + bx)(c + dx)^{-3+n}(e + fx)^{-n} dx$$

$$= \left(\int \frac{(dx + c)^n}{(fx + e)^n c^3 + 3(fx + e)^n c^2 dx + 3(fx + e)^n c d^2 x^2 + (fx + e)^n d^3 x^3} dx \right) a$$

$$+ \left(\int \frac{(dx + c)^n x}{(fx + e)^n c^3 + 3(fx + e)^n c^2 dx + 3(fx + e)^n c d^2 x^2 + (fx + e)^n d^3 x^3} dx \right) b$$

input

```
int((b*x+a)*(d*x+c)^(-3+n)/((f*x+e)^n),x)
```

output

```
int((c + d*x)**n/((e + f*x)**n*c**3 + 3*(e + f*x)**n*c**2*d*x + 3*(e + f*x)
)**n*c*d**2*x**2 + (e + f*x)**n*d**3*x**3),x)*a + int(((c + d*x)**n*x)/((e
+ f*x)**n*c**3 + 3*(e + f*x)**n*c**2*d*x + 3*(e + f*x)**n*c*d**2*x**2 + (
e + f*x)**n*d**3*x**3),x)*b
```

3.256 $\int (a + bx)(c + dx)^{-4+n}(e + fx)^{-n} dx$

Optimal result	2234
Mathematica [A] (verified)	2235
Rubi [A] (verified)	2235
Maple [B] (verified)	2237
Fricas [B] (verification not implemented)	2238
Sympy [F(-2)]	2238
Maxima [F]	2239
Giac [F]	2239
Mupad [B] (verification not implemented)	2240
Reduce [F]	2241

Optimal result

Integrand size = 24, antiderivative size = 205

$$\begin{aligned} & \int (a + bx)(c + dx)^{-4+n}(e + fx)^{-n} dx \\ &= \frac{(bc - ad)(c + dx)^{-3+n}(e + fx)^{1-n}}{d(de - cf)(3 - n)} \\ &+ \frac{(2adf + bcf(1 - n) - bde(3 - n))(c + dx)^{-2+n}(e + fx)^{1-n}}{d(de - cf)^2(2 - n)(3 - n)} \\ &- \frac{f(2adf + bcf(1 - n) - bde(3 - n))(c + dx)^{-1+n}(e + fx)^{1-n}}{d(de - cf)^3(1 - n)(2 - n)(3 - n)} \end{aligned}$$

output

```
(-a*d+b*c)*(d*x+c)^(-3+n)*(f*x+e)^(1-n)/d/(-c*f+d*e)/(3-n)+(2*a*d*f+b*c*f*(1-n)-b*d*e*(3-n))*(d*x+c)^(-2+n)*(f*x+e)^(1-n)/d/(-c*f+d*e)^2/(2-n)/(3-n)-f*(2*a*d*f+b*c*f*(1-n)-b*d*e*(3-n))*(d*x+c)^(-1+n)*(f*x+e)^(1-n)/d/(-c*f+d*e)^3/(1-n)/(2-n)/(3-n)
```

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.55

$$\int (a + bx)(c + dx)^{-4+n}(e + fx)^{-n} dx$$

$$= \frac{(c + dx)^{-3+n}(e + fx)^{1-n} \left(-bc + ad + \frac{(2adf + bde(-3+n) - bcf(-1+n))(c+dx)(-cf(-2+n) + de(-1+n) + dfx)}{(de - cf)^2(-2+n)(-1+n)} \right)}{d(de - cf)(-3 + n)}$$

input

```
Integrate[((a + b*x)*(c + d*x)^(-4 + n))/(e + f*x)^n,x]
```

output

```
((c + d*x)^(-3 + n)*(e + f*x)^(1 - n)*(-(b*c) + a*d + ((2*a*d*f + b*d*e*(-3 + n) - b*c*f*(-1 + n))*(c + d*x)*(-(c*f*(-2 + n)) + d*e*(-1 + n) + d*f*x)))/((d*e - c*f)^2*(-2 + n)*(-1 + n)))/(d*(d*e - c*f)*(-3 + n))
```

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 183, normalized size of antiderivative = 0.89, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {88, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)(c + dx)^{n-4}(e + fx)^{-n} dx$$

$$\downarrow 88$$

$$\frac{(bc - ad)(c + dx)^{n-3}(e + fx)^{1-n}}{d(3 - n)(de - cf)} -$$

$$\frac{(2adf + bcf(1 - n) - bde(3 - n)) \int (c + dx)^{n-3}(e + fx)^{-n} dx}{d(3 - n)(de - cf)}$$

$$\downarrow 55$$

$$\frac{(bc - ad)(c + dx)^{n-3}(e + fx)^{1-n}}{d(3 - n)(de - cf)} - \frac{(2adf + bcf(1 - n) - bde(3 - n)) \left(-\frac{f \int (c+dx)^{n-2}(e+fx)^{-n} dx}{(2-n)(de-cf)} - \frac{(c+dx)^{n-2}(e+fx)^{1-n}}{(2-n)(de-cf)} \right)}{d(3 - n)(de - cf)}$$

↓ 48

$$\frac{(bc - ad)(c + dx)^{n-3}(e + fx)^{1-n}}{d(3 - n)(de - cf)} - \frac{\left(\frac{f(c+dx)^{n-1}(e+fx)^{1-n}}{(1-n)(2-n)(de-cf)^2} - \frac{(c+dx)^{n-2}(e+fx)^{1-n}}{(2-n)(de-cf)} \right) (2adf + bcf(1 - n) - bde(3 - n))}{d(3 - n)(de - cf)}$$

input `Int[((a + b*x)*(c + d*x)^(-4 + n))/(e + f*x)^n,x]`

output `((b*c - a*d)*(c + d*x)^(-3 + n)*(e + f*x)^(1 - n))/(d*(d*e - c*f)*(3 - n)) - ((2*a*d*f + b*c*f*(1 - n) - b*d*e*(3 - n))*(-((c + d*x)^(-2 + n)*(e + f*x)^(1 - n))/((d*e - c*f)*(2 - n))) + (f*(c + d*x)^(-1 + n)*(e + f*x)^(1 - n))/((d*e - c*f)^2*(1 - n)*(2 - n)))/(d*(d*e - c*f)*(3 - n))`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 55 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(Simplify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])`

rule 88

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p
_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^Simplify[p + 1],
x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && !RationalQ[p] && SumSimpl
erQ[p, 1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 505 vs. 2(205) = 410.

Time = 0.54 (sec) , antiderivative size = 506, normalized size of antiderivative = 2.47

method	result
gospers	$\frac{(xd+c)^{-3+n}(fx+e)(fx+e)^{-n}(bc^2f^2n^2x-2bcdefn^2x-bcdf^2nx^2+bd^2e^2n^2x+bd^2efnx^2+ac^2f^2n^2-2acdefn^2-2acd^2n^2)}{c^3f^3n^3-3c^2def^2n^3+3c^2d^2efn^3}$
orering	$\frac{(fx+e)(xd+c)(bc^2f^2n^2x-2bcdefn^2x-bcdf^2nx^2+bd^2e^2n^2x+bd^2efnx^2+ac^2f^2n^2-2acdefn^2-2acd^2nx+ad^2e^2n^2)}{c^3f^3n^3-3c^2def^2n^3+3c^2d^2efn^3}$
parallelrisch	Expression too large to display

input

```
int((b*x+a)*(d*x+c)^(-4+n)/((f*x+e)^n),x,method=_RETURNVERBOSE)
```

output

```
-(d*x+c)^(-3+n)*(f*x+e)/((f*x+e)^n)/(c^3*f^3*n^3-3*c^2*d*e*f^2*n^3+3*c*d^2
*e^2*f*n^3-d^3*e^3*n^3-6*c^3*f^3*n^2+18*c^2*d*e*f^2*n^2-18*c*d^2*e^2*f*n^2
+6*d^3*e^3*n^2+11*c^3*f^3*n-33*c^2*d*e*f^2*n+33*c*d^2*e^2*f*n-11*d^3*e^3*n
-6*c^3*f^3+18*c^2*d*e*f^2-18*c*d^2*e^2*f+6*d^3*e^3)*(b*c^2*f^2*n^2*x-2*b*c
*d*e*f*n^2*x-b*c*d*f^2*n*x^2+b*d^2*e^2*n^2*x+b*d^2*e*f*n*x^2+a*c^2*f^2*n^2
-2*a*c*d*e*f*n^2-2*a*c*d*f^2*n*x+a*d^2*e^2*n^2+2*a*d^2*e*f*n*x+2*a*d^2*f^2
*x^2-4*b*c^2*f^2*n*x+8*b*c*d*e*f*n*x+b*c*d*f^2*x^2-4*b*d^2*e^2*n*x-3*b*d^2
*e*f*x^2-5*a*c^2*f^2*n+8*a*c*d*e*f*n+6*a*c*d*f^2*x-3*a*d^2*e^2*n-2*a*d^2*e
*f*x+b*c^2*e*f*n+3*b*c^2*f^2*x-b*c*d*e^2*n-10*b*c*d*e*f*x+3*b*d^2*e^2*x+6*
a*c^2*f^2-6*a*c*d*e*f+2*a*d^2*e^2-3*b*c^2*e*f+b*c*d*e^2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 884 vs. $2(187) = 374$.

Time = 0.13 (sec) , antiderivative size = 884, normalized size of antiderivative = 4.31

$$\int (a + bx)(c + dx)^{-4+n}(e + fx)^{-n} dx = \text{Too large to display}$$

input `integrate((b*x+a)*(d*x+c)^(-4+n)/((f*x+e)^n),x, algorithm="fricas")`

output

```

-(6*a*c^3*e*f^2 - (3*b*d^3*e*f^2 - (b*c*d^2 + 2*a*d^3)*f^3 - (b*d^3*e*f^2
- b*c*d^2*f^3)*n)*x^4 + (b*c^2*d + 2*a*c*d^2)*e^3 - 3*(b*c^3 + 2*a*c^2*d)*
e^2*f - (12*b*c*d^2*e*f^2 - 4*(b*c^2*d + 2*a*c*d^2)*f^3 - (b*d^3*e^2*f - 2
*b*c*d^2*e*f^2 + b*c^2*d*f^3)*n^2 + (3*b*d^3*e^2*f - 2*(4*b*c*d^2 + a*d^3)
*e*f^2 + (5*b*c^2*d + 2*a*c*d^2)*f^3)*n)*x^3 + (a*c*d^2*e^3 - 2*a*c^2*d*e^
2*f + a*c^3*e*f^2)*n^2 + (3*b*d^3*e^3 - 9*b*c*d^2*e^2*f - 9*b*c^2*d*e*f^2
+ 3*(b*c^3 + 4*a*c^2*d)*f^3 + (b*d^3*e^3 - (b*c*d^2 - a*d^3)*e^2*f - (b*c^
2*d + 2*a*c*d^2)*e*f^2 + (b*c^3 + a*c^2*d)*f^3)*n^2 - (4*b*d^3*e^3 - (4*b*
c*d^2 - a*d^3)*e^2*f - 4*(b*c^2*d + 2*a*c*d^2)*e*f^2 + (4*b*c^3 + 7*a*c^2*
d)*f^3)*n)*x^2 - (5*a*c^3*e*f^2 + (b*c^2*d + 3*a*c*d^2)*e^3 - (b*c^3 + 8*a
*c^2*d)*e^2*f)*n + (6*a*c^2*d*e*f^2 + 6*a*c^3*f^3 + 2*(2*b*c*d^2 + a*d^3)*
e^3 - 6*(2*b*c^2*d + a*c*d^2)*e^2*f + (a*c^3*f^3 + (b*c*d^2 + a*d^3)*e^3 -
(2*b*c^2*d + a*c*d^2)*e^2*f + (b*c^3 - a*c^2*d)*e*f^2)*n^2 - (5*a*c^3*f^3
+ (5*b*c*d^2 + 3*a*d^3)*e^3 - (8*b*c^2*d + 7*a*c*d^2)*e^2*f + (3*b*c^3 -
a*c^2*d)*e*f^2)*n)*x*(d*x + c)^(n - 4)/((6*d^3*e^3 - 18*c*d^2*e^2*f + 18*
c^2*d*e*f^2 - 6*c^3*f^3 - (d^3*e^3 - 3*c*d^2*e^2*f + 3*c^2*d*e*f^2 - c^3*f^
3)*n^3 + 6*(d^3*e^3 - 3*c*d^2*e^2*f + 3*c^2*d*e*f^2 - c^3*f^3)*n^2 - 11*(
d^3*e^3 - 3*c*d^2*e^2*f + 3*c^2*d*e*f^2 - c^3*f^3)*n)*(f*x + e)^n

```

Sympy [F(-2)]

Exception generated.

$$\int (a + bx)(c + dx)^{-4+n}(e + fx)^{-n} dx = \text{Exception raised: HeuristicGCDFailed}$$

input `integrate((b*x+a)*(d*x+c)**(-4+n)/((f*x+e)**n),x)`

output Exception raised: HeuristicGCDFailed >> no luck

Maxima [F]

$$\int (a + bx)(c + dx)^{-4+n}(e + fx)^{-n} dx = \int \frac{(bx + a)(dx + c)^{n-4}}{(fx + e)^n} dx$$

input `integrate((b*x+a)*(d*x+c)^(-4+n)/((f*x+e)^n),x, algorithm="maxima")`

output `integrate((b*x + a)*(d*x + c)^(n - 4)/(f*x + e)^n, x)`

Giac [F]

$$\int (a + bx)(c + dx)^{-4+n}(e + fx)^{-n} dx = \int \frac{(bx + a)(dx + c)^{n-4}}{(fx + e)^n} dx$$

input `integrate((b*x+a)*(d*x+c)^(-4+n)/((f*x+e)^n),x, algorithm="giac")`

output `integrate((b*x + a)*(d*x + c)^(n - 4)/(f*x + e)^n, x)`

output

```

- (x*(c + d*x)^(n - 4)*(6*a*c^3*f^3 + 2*a*d^3*e^3 + a*c^3*f^3*n^2 + a*d^3*
e^3*n^2 + 4*b*c*d^2*e^3 - 5*a*c^3*f^3*n - 3*a*d^3*e^3*n - 6*a*c*d^2*e^2*f
+ 6*a*c^2*d*e*f^2 - 12*b*c^2*d*e^2*f - 5*b*c*d^2*e^3*n - 3*b*c^3*e*f^2*n +
b*c*d^2*e^3*n^2 + b*c^3*e*f^2*n^2 + 7*a*c*d^2*e^2*f*n + a*c^2*d*e*f^2*n +
8*b*c^2*d*e^2*f*n - a*c*d^2*e^2*f*n^2 - a*c^2*d*e*f^2*n^2 - 2*b*c^2*d*e^2
*f*n^2))/((e + f*x)^n*(c*f - d*e)^3*(11*n - 6*n^2 + n^3 - 6)) - (x^2*(c +
d*x)^(n - 4)*(3*b*c^3*f^3 + 3*b*d^3*e^3 + b*c^3*f^3*n^2 + b*d^3*e^3*n^2 +
12*a*c^2*d*f^3 - 4*b*c^3*f^3*n - 4*b*d^3*e^3*n - 9*b*c*d^2*e^2*f - 9*b*c^2
*d*e*f^2 - 7*a*c^2*d*f^3*n - a*d^3*e^2*f*n + a*c^2*d*f^3*n^2 + a*d^3*e^2*f
*n^2 + 8*a*c*d^2*e*f^2*n + 4*b*c*d^2*e^2*f*n + 4*b*c^2*d*e*f^2*n - 2*a*c*d
^2*e*f^2*n^2 - b*c*d^2*e^2*f*n^2 - b*c^2*d*e*f^2*n^2))/((e + f*x)^n*(c*f -
d*e)^3*(11*n - 6*n^2 + n^3 - 6)) - (c*e*(c + d*x)^(n - 4)*(6*a*c^2*f^2 +
2*a*d^2*e^2 + a*c^2*f^2*n^2 + a*d^2*e^2*n^2 + b*c*d*e^2 - 3*b*c^2*e*f - 5*
a*c^2*f^2*n - 3*a*d^2*e^2*n - 6*a*c*d*e*f - b*c*d*e^2*n + b*c^2*e*f*n - 2*
a*c*d*e*f*n^2 + 8*a*c*d*e*f*n))/((e + f*x)^n*(c*f - d*e)^3*(11*n - 6*n^2 +
n^3 - 6)) - (d^2*f^2*x^4*(c + d*x)^(n - 4)*(2*a*d*f + b*c*f - 3*b*d*e - b
*c*f*n + b*d*e*n))/((e + f*x)^n*(c*f - d*e)^3*(11*n - 6*n^2 + n^3 - 6)) -
(d*f*x^3*(c + d*x)^(n - 4)*(4*c*f - c*f*n + d*e*n)*(2*a*d*f + b*c*f - 3*b*
d*e - b*c*f*n + b*d*e*n))/((e + f*x)^n*(c*f - d*e)^3*(11*n - 6*n^2 + n^3 -
6))

```

Reduce [F]

$$\begin{aligned}
& \int (a + bx)(c + dx)^{-4+n}(e + fx)^{-n} dx \\
&= \left(\int \frac{(dx + c)^n}{(fx + e)^n c^4 + 4(fx + e)^n c^3 dx + 6(fx + e)^n c^2 d^2 x^2 + 4(fx + e)^n c d^3 x^3 + (fx + e)^n d^4 x^4} dx \right) a \\
&+ \left(\int \frac{(dx + c)^n x}{(fx + e)^n c^4 + 4(fx + e)^n c^3 dx + 6(fx + e)^n c^2 d^2 x^2 + 4(fx + e)^n c d^3 x^3 + (fx + e)^n d^4 x^4} dx \right) b
\end{aligned}$$

input

```
int((b*x+a)*(d*x+c)^(-4+n)/((f*x+e)^n),x)
```

output

```

int((c + d*x)**n/((e + f*x)**n*c**4 + 4*(e + f*x)**n*c**3*d*x + 6*(e + f*x
)**n*c**2*d**2*x**2 + 4*(e + f*x)**n*c*d**3*x**3 + (e + f*x)**n*d**4*x**4)
,x)*a + int(((c + d*x)**n*x)/((e + f*x)**n*c**4 + 4*(e + f*x)**n*c**3*d*x
+ 6*(e + f*x)**n*c**2*d**2*x**2 + 4*(e + f*x)**n*c*d**3*x**3 + (e + f*x)**
n*d**4*x**4),x)*b

```

3.257 $\int (a + bx)(c + dx)^{-5+n}(e + fx)^{-n} dx$

Optimal result	2242
Mathematica [A] (verified)	2243
Rubi [A] (verified)	2243
Maple [B] (verified)	2245
Fricas [B] (verification not implemented)	2246
Sympy [F(-2)]	2247
Maxima [F]	2248
Giac [F]	2248
Mupad [B] (verification not implemented)	2248
Reduce [F]	2249

Optimal result

Integrand size = 24, antiderivative size = 296

$$\begin{aligned} & \int (a + bx)(c + dx)^{-5+n}(e + fx)^{-n} dx \\ &= \frac{(bc - ad)(c + dx)^{-4+n}(e + fx)^{1-n}}{d(de - cf)(4 - n)} \\ &+ \frac{(3adf + bcf(1 - n) - bde(4 - n))(c + dx)^{-3+n}(e + fx)^{1-n}}{d(de - cf)^2(3 - n)(4 - n)} \\ &- \frac{2f(3adf + bcf(1 - n) - bde(4 - n))(c + dx)^{-2+n}(e + fx)^{1-n}}{d(de - cf)^3(2 - n)(3 - n)(4 - n)} \\ &+ \frac{2f^2(3adf + bcf(1 - n) - bde(4 - n))(c + dx)^{-1+n}(e + fx)^{1-n}}{d(de - cf)^4(1 - n)(2 - n)(3 - n)(4 - n)} \end{aligned}$$

output

```
(-a*d+b*c)*(d*x+c)^(-4+n)*(f*x+e)^(1-n)/d/(-c*f+d*e)/(4-n)+(3*a*d*f+b*c*f*(1-n)-b*d*e*(4-n))*(d*x+c)^(-3+n)*(f*x+e)^(1-n)/d/(-c*f+d*e)^2/(3-n)/(4-n)-2*f*(3*a*d*f+b*c*f*(1-n)-b*d*e*(4-n))*(d*x+c)^(-2+n)*(f*x+e)^(1-n)/d/(-c*f+d*e)^3/(2-n)/(3-n)/(4-n)+2*f^2*(3*a*d*f+b*c*f*(1-n)-b*d*e*(4-n))*(d*x+c)^(-1+n)*(f*x+e)^(1-n)/d/(-c*f+d*e)^4/(1-n)/(2-n)/(3-n)/(4-n)
```

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.56

$$\int (a + bx)(c + dx)^{-5+n}(e + fx)^{-n} dx$$

$$= \frac{(c + dx)^{-4+n}(e + fx)^{1-n} \left(-bc + ad + \frac{(3adf + bde(-4+n) - bcf(-1+n))(c + dx)(c^2 f^2 (6 - 5n + n^2) - 2cdf(-3+n)(e(-1+n) + fx))}{(de - cf)^3(-3+n)(-2+n)(-1+n)} \right)}{d(de - cf)(-4 + n)}$$

input

```
Integrate[((a + b*x)*(c + d*x)^(-5 + n))/(e + f*x)^n,x]
```

output

```
((c + d*x)^(-4 + n)*(e + f*x)^(1 - n)*(-(b*c) + a*d + ((3*a*d*f + b*d*e*(-4 + n) - b*c*f*(-1 + n))*(c + d*x)*(c^2*f^2*(6 - 5*n + n^2) - 2*c*d*f*(-3 + n)*(e*(-1 + n) + f*x) + d^2*(e^2*(2 - 3*n + n^2) + 2*e*f*(-1 + n)*x + 2*f^2*x^2)))/((d*e - c*f)^3*(-3 + n)*(-2 + n)*(-1 + n)))/(d*(d*e - c*f)*(-4 + n))
```

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 243, normalized size of antiderivative = 0.82, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {88, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)(c + dx)^{n-5}(e + fx)^{-n} dx$$

$$\downarrow 88$$

$$\frac{(bc - ad)(c + dx)^{n-4}(e + fx)^{1-n}}{d(4 - n)(de - cf)}$$

$$\frac{(3adf + bcf(1 - n) - bde(4 - n)) \int (c + dx)^{n-4}(e + fx)^{-n} dx}{d(4 - n)(de - cf)}$$

$$\downarrow 55$$

$$\begin{aligned}
 & \frac{(bc - ad)(c + dx)^{n-4}(e + fx)^{1-n}}{d(4 - n)(de - cf)} - \frac{(3adf + bcf(1 - n) - bde(4 - n)) \left(-\frac{2f \int (c+dx)^{n-3}(e+fx)^{-n} dx}{(3-n)(de-cf)} - \frac{(c+dx)^{n-3}(e+fx)^{1-n}}{(3-n)(de-cf)} \right)}{d(4 - n)(de - cf)} \\
 & \quad \downarrow 55 \\
 & \frac{(bc - ad)(c + dx)^{n-4}(e + fx)^{1-n}}{d(4 - n)(de - cf)} - \frac{(3adf + bcf(1 - n) - bde(4 - n)) \left(-\frac{2f \left(-\frac{f \int (c+dx)^{n-2}(e+fx)^{-n} dx}{(2-n)(de-cf)} - \frac{(c+dx)^{n-2}(e+fx)^{1-n}}{(2-n)(de-cf)} \right)}{(3-n)(de-cf)} - \frac{(c+dx)^{n-3}(e+fx)^{1-n}}{(3-n)(de-cf)} \right)}{d(4 - n)(de - cf)} \\
 & \quad \downarrow 48 \\
 & \frac{(bc - ad)(c + dx)^{n-4}(e + fx)^{1-n}}{d(4 - n)(de - cf)} - \frac{\left(-\frac{(c+dx)^{n-3}(e+fx)^{1-n}}{(3-n)(de-cf)} - \frac{2f \left(\frac{f(c+dx)^{n-1}(e+fx)^{1-n}}{(1-n)(2-n)(de-cf)^2} - \frac{(c+dx)^{n-2}(e+fx)^{1-n}}{(2-n)(de-cf)} \right)}{(3-n)(de-cf)} \right) (3adf + bcf(1 - n) - bde(4 - n))}{d(4 - n)(de - cf)}
 \end{aligned}$$

input `Int[((a + b*x)*(c + d*x)^(-5 + n))/(e + f*x)^n,x]`

output `((b*c - a*d)*(c + d*x)^(-4 + n)*(e + f*x)^(1 - n))/(d*(d*e - c*f)*(4 - n)) - (((3*a*d*f + b*c*f*(1 - n) - b*d*e*(4 - n))*(-(((c + d*x)^(-3 + n)*(e + f*x)^(1 - n))/(d*e - c*f)*(3 - n))) - (2*f*(-(((c + d*x)^(-2 + n)*(e + f*x)^(1 - n))/(d*e - c*f)*(2 - n))) + (f*(c + d*x)^(-1 + n)*(e + f*x)^(1 - n))/(d*e - c*f)^2*(1 - n)*(2 - n)))/(d*e - c*f)*(3 - n)))/(d*(d*e - c*f)*(4 - n))`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 55

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
lerQ[n, 1])
```

rule 88

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^Simplify[p + 1],
x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && !RationalQ[p] && SumSimpl
erQ[p, 1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1186 vs. $2(296) = 592$.

Time = 0.56 (sec) , antiderivative size = 1187, normalized size of antiderivative = 4.01

method	result	size
gospers	Expression too large to display	1187
orering	Expression too large to display	1192
parallelisch	Expression too large to display	3499

input

```
int((b*x+a)*(d*x+c)^(-5+n)/((f*x+e)^n),x,method=_RETURNVERBOSE)
```

output

```

-(d*x+c)^(-4+n)*(f*x+e)/((f*x+e)^n)/(c^4*f^4*n^4-4*c^3*d*e*f^3*n^4+6*c^2*d^2*e^2*f^2*n^4-4*c*d^3*e^3*f*n^4+d^4*e^4*n^4-10*c^4*f^4*n^3+40*c^3*d*e*f^3*n^3-60*c^2*d^2*e^2*f^2*n^3+40*c*d^3*e^3*f*n^3-10*d^4*e^4*n^3+35*c^4*f^4*n^2-140*c^3*d*e*f^3*n^2+210*c^2*d^2*e^2*f^2*n^2-140*c*d^3*e^3*f*n^2+35*d^4*e^4*n^2-50*c^4*f^4*n+200*c^3*d*e*f^3*n-300*c^2*d^2*e^2*f^2*n+200*c*d^3*e^3*f*n-50*d^4*e^4*n+24*c^4*f^4-96*c^3*d*e*f^3+144*c^2*d^2*e^2*f^2-96*c*d^3*e^3*f+24*d^4*e^4)*(b*c^3*f^3*n^3*x-3*b*c^2*d*e*f^2*n^3*x-2*b*c^2*d*f^3*n^2*x^2+3*b*c*d^2*e^2*f*n^3*x+4*b*c*d^2*e*f^2*n^2*x^2+2*b*c*d^2*f^3*n*x^3-b*d^3*e^3*n^3*x-2*b*d^3*e^2*f*n^2*x^2-2*b*d^3*e*f^2*n*x^3+a*c^3*f^3*n^3-3*a*c^2*d*e*f^2*n^3-3*a*c^2*d*f^3*n^2*x+3*a*c*d^2*e^2*f*n^3+6*a*c*d^2*e*f^2*n^2*x+6*a*c*d^2*f^3*n*x^2-a*d^3*e^3*n^3-3*a*d^3*e^2*f*n^2*x-6*a*d^3*e*f^2*n*x^2-6*a*d^3*f^3*x^3-8*b*c^3*f^3*n^2*x+23*b*c^2*d*e*f^2*n^2*x+10*b*c^2*d*f^3*n*x^2-22*b*c*d^2*e^2*f*n^2*x-20*b*c*d^2*e*f^2*n*x^2-2*b*c*d^2*f^3*x^3+7*b*d^3*e^3*n^2*x+10*b*d^3*e^2*f*n*x^2+8*b*d^3*e*f^2*x^3-9*a*c^3*f^3*n^2+24*a*c^2*d*e*f^2*n^2+21*a*c^2*d*f^3*n*x-21*a*c*d^2*e^2*f*n^2-30*a*c*d^2*e*f^2*n*x-24*a*c*d^2*f^3*x^2+6*a*d^3*e^3*n^2+9*a*d^3*e^2*f*n*x+6*a*d^3*e*f^2*x^2+b*c^3*e*f^2*n^2+19*b*c^3*f^3*n*x-2*b*c^2*d*e^2*f*n^2-58*b*c^2*d*e*f^2*n*x-8*b*c^2*d*f^3*x^2+b*c*d^2*e^3*n^2+53*b*c*d^2*e^2*f*n*x+34*b*c*d^2*e*f^2*x^2-14*b*d^3*e^3*n*x-8*b*d^3*e^2*f*x^2+26*a*c^3*f^3*n-57*a*c^2*d*e*f^2*n-36*a*c^2*d*f^3*x+42*a*c*d^2*e^2*f*n+24*a*c*d^2*e*f^2*x-11*a*d^3*e^3*n-6*a*...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1741 vs. $2(266) = 532$.

Time = 0.14 (sec) , antiderivative size = 1741, normalized size of antiderivative = 5.88

$$\int (a + bx)(c + dx)^{-5+n}(e + fx)^{-n} dx = \text{Too large to display}$$

input

```
integrate((b*x+a)*(d*x+c)^(-5+n)/((f*x+e)^n),x, algorithm="fricas")
```

output

```
(24*a*c^4*e*f^3 - 2*(4*b*d^4*e*f^3 - (b*c*d^3 + 3*a*d^4)*f^4 - (b*d^4*e*f^3 - b*c*d^3*f^4)*n)*x^5 - 2*(b*c^2*d^2 + 3*a*c*d^3)*e^4 + 8*(b*c^3*d + 3*a*c^2*d^2)*e^3*f - 12*(b*c^4 + 3*a*c^3*d)*e^2*f^2 - 2*(20*b*c*d^3*e*f^3 - 5*(b*c^2*d^2 + 3*a*c*d^3)*f^4 - (b*d^4*e^2*f^2 - 2*b*c*d^3*e*f^3 + b*c^2*d^2*f^4)*n^2 + (4*b*d^4*e^2*f^2 - (10*b*c*d^3 + 3*a*d^4)*e*f^3 + 3*(2*b*c^2*d^2 + a*c*d^3)*f^4)*n)*x^4 + (a*c*d^3*e^4 - 3*a*c^2*d^2*e^3*f + 3*a*c^3*d*e^2*f^2 - a*c^4*e*f^3)*n^3 - (80*b*c^2*d^2*e*f^3 - 20*(b*c^3*d + 3*a*c^2*d^2)*f^4 - (b*d^4*e^3*f - 3*b*c*d^3*e^2*f^2 + 3*b*c^2*d^2*e*f^3 - b*c^3*d*f^4)*n^3 + (5*b*d^4*e^3*f - (20*b*c*d^3 + 3*a*d^4)*e^2*f^2 + (25*b*c^2*d^2 + 6*a*c*d^3)*e*f^3 - (10*b*c^3*d + 3*a*c^2*d^2)*f^4)*n^2 - (4*b*d^4*e^3*f - (41*b*c*d^3 + 3*a*d^4)*e^2*f^2 + 6*(11*b*c^2*d^2 + 5*a*c*d^3)*e*f^3 - (29*b*c^3*d + 27*a*c^2*d^2)*f^4)*n)*x^3 + (9*a*c^4*e*f^3 - (b*c^2*d^2 + 6*a*c*d^3)*e^4 + (2*b*c^3*d + 21*a*c^2*d^2)*e^3*f - (b*c^4 + 24*a*c^3*d)*e^2*f^2)*n^2 - (8*b*d^4*e^4 - 32*b*c*d^3*e^3*f + 48*b*c^2*d^2*e^2*f^2 + 48*b*c^3*d*e*f^3 - 12*(b*c^4 + 5*a*c^3*d)*f^4 - (b*d^4*e^4 - 3*a*c*d^3*e^2*f^2 - (2*b*c*d^3 - a*d^4)*e^3*f + (2*b*c^3*d + 3*a*c^2*d^2)*e*f^3 - (b*c^4 + a*c^3*d)*f^4)*n^3 + (7*b*d^4*e^4 - (16*b*c*d^3 - 3*a*d^4)*e^3*f + 3*(b*c^2*d^2 - 6*a*c*d^3)*e^2*f^2 + (14*b*c^3*d + 27*a*c^2*d^2)*e*f^3 - 4*(2*b*c^4 + 3*a*c^3*d)*f^4)*n^2 - (14*b*d^4*e^4 - 2*(23*b*c*d^3 - a*d^4)*e^3*f + 15*(b*c^2*d^2 - a*c*d^3)*e^2*f^2 + 12*(3*b*c^3*d + 5*a*c^2*d^2)*e*f^3 - (19*...
```

Sympy [F(-2)]

Exception generated.

$$\int (a + bx)(c + dx)^{-5+n}(e + fx)^{-n} dx = \text{Exception raised: HeuristicGCDFailed}$$

input

```
integrate((b*x+a)*(d*x+c)**(-5+n)/((f*x+e)**n),x)
```

output

```
Exception raised: HeuristicGCDFailed >> no luck
```

Maxima [F]

$$\int (a + bx)(c + dx)^{-5+n}(e + fx)^{-n} dx = \int \frac{(bx + a)(dx + c)^{n-5}}{(fx + e)^n} dx$$

input `integrate((b*x+a)*(d*x+c)^(-5+n)/((f*x+e)^n),x, algorithm="maxima")`

output `integrate((b*x + a)*(d*x + c)^(n - 5)/(f*x + e)^n, x)`

Giac [F]

$$\int (a + bx)(c + dx)^{-5+n}(e + fx)^{-n} dx = \int \frac{(bx + a)(dx + c)^{n-5}}{(fx + e)^n} dx$$

input `integrate((b*x+a)*(d*x+c)^(-5+n)/((f*x+e)^n),x, algorithm="giac")`

output `integrate((b*x + a)*(d*x + c)^(n - 5)/(f*x + e)^n, x)`

Mupad [B] (verification not implemented)

Time = 2.37 (sec) , antiderivative size = 1657, normalized size of antiderivative = 5.60

$$\int (a + bx)(c + dx)^{-5+n}(e + fx)^{-n} dx = \text{Too large to display}$$

input `int(((a + b*x)*(c + d*x)^(n - 5))/(e + f*x)^n,x)`

output

```
(x*(c + d*x)^(n - 5)*(24*a*c^4*f^4 - 6*a*d^4*e^4 + 9*a*c^4*f^4*n^2 - 6*a*d^4*e^4*n^2 - a*c^4*f^4*n^3 + a*d^4*e^4*n^3 - 10*b*c*d^3*e^4 - 26*a*c^4*f^4*n + 11*a*d^4*e^4*n + 24*a*c*d^3*e^3*f + 24*a*c^3*d*e*f^3 + 17*b*c*d^3*e^4*n - 12*b*c^4*e*f^3*n + 40*b*c^2*d^2*e^3*f - 60*b*c^3*d*e^2*f^2 - 8*b*c*d^3*e^4*n^2 + b*c*d^3*e^4*n^3 + 7*b*c^4*e*f^3*n^2 - b*c^4*e*f^3*n^3 - 36*a*c^2*d^2*e^2*f^2 + 45*a*c^2*d^2*e^2*f^2*n + 23*b*c^2*d^2*e^3*f*n^2 - 22*b*c^3*d*e^2*f^2*n^2 - 3*b*c^2*d^2*e^3*f*n^3 + 3*b*c^3*d*e^2*f^2*n^3 - 40*a*c*d^3*e^3*f*n + 10*a*c^3*d*e*f^3*n - 9*a*c^2*d^2*e^2*f^2*n^2 + 18*a*c*d^3*e^3*f*n^2 - 12*a*c^3*d*e*f^3*n^2 - 2*a*c*d^3*e^3*f*n^3 + 2*a*c^3*d*e*f^3*n^3 - 60*b*c^2*d^2*e^3*f*n + 55*b*c^3*d*e^2*f^2*n))/((e + f*x)^n*(c*f - d*e)^4*(35*n^2 - 50*n - 10*n^3 + n^4 + 24)) - ((c + d*x)^(n - 5)*(2*b*c^2*d^2*e^4 + 12*b*c^4*e^2*f^2 + 6*a*c*d^3*e^4 - 24*a*c^4*e*f^3 - 8*b*c^3*d*e^3*f - 11*a*c*d^3*e^4*n + 26*a*c^4*e*f^3*n - 24*a*c^2*d^2*e^3*f + 36*a*c^3*d*e^2*f^2 + 6*a*c*d^3*e^4*n^2 - a*c*d^3*e^4*n^3 - 3*b*c^2*d^2*e^4*n - 9*a*c^4*e*f^3*n^2 + a*c^4*e*f^3*n^3 - 7*b*c^4*e^2*f^2*n + b*c^2*d^2*e^4*n^2 + b*c^4*e^2*f^2*n^2 - 21*a*c^2*d^2*e^3*f*n^2 + 24*a*c^3*d*e^2*f^2*n^2 + 3*a*c^2*d^2*e^3*f*n^3 - 3*a*c^3*d*e^2*f^2*n^3 + 10*b*c^3*d*e^3*f*n + 42*a*c^2*d^2*e^3*f*n - 57*a*c^3*d*e^2*f^2*n - 2*b*c^3*d*e^3*f*n^2))/((e + f*x)^n*(c*f - d*e)^4*(35*n^2 - 50*n - 10*n^3 + n^4 + 24)) + (x^2*(c + d*x)^(n - 5)*(12*b*c^4*f^4 - 8*b*d^4*e^4 + 8*b*c^4*f^4*n^2 - 7*b*d^4*e^4*n^2 - b*c^4*f^4*n...
```

Reduce [F]

$$\int (a + bx)(c + dx)^{-5+n}(e + fx)^{-n} dx$$

$$= \left(\int \frac{(dx + c)^n}{(fx + e)^n c^5 + 5(fx + e)^n c^4 dx + 10(fx + e)^n c^3 d^2 x^2 + 10(fx + e)^n c^2 d^3 x^3 + 5(fx + e)^n c d^4 x^4 + (dx + c)^n x} \right) + \left(\int \frac{(dx + c)^n}{(fx + e)^n c^5 + 5(fx + e)^n c^4 dx + 10(fx + e)^n c^3 d^2 x^2 + 10(fx + e)^n c^2 d^3 x^3 + 5(fx + e)^n c d^4 x^4} \right)$$

input

```
int((b*x+a)*(d*x+c)^(-5+n)/((f*x+e)^n),x)
```

output

```
int((c + d*x)**n/((e + f*x)**n*c**5 + 5*(e + f*x)**n*c**4*d*x + 10*(e + f*
x)**n*c**3*d**2*x**2 + 10*(e + f*x)**n*c**2*d**3*x**3 + 5*(e + f*x)**n*c*d
**4*x**4 + (e + f*x)**n*d**5*x**5),x)*a + int(((c + d*x)**n*x)/((e + f*x)*
**n*c**5 + 5*(e + f*x)**n*c**4*d*x + 10*(e + f*x)**n*c**3*d**2*x**2 + 10*(e
+ f*x)**n*c**2*d**3*x**3 + 5*(e + f*x)**n*c*d**4*x**4 + (e + f*x)**n*d**5
*x**5),x)*b
```

3.258 $\int (a + bx)^{-n}(c + dx)(e + fx)^n dx$

Optimal result	2251
Mathematica [A] (verified)	2251
Rubi [A] (verified)	2252
Maple [F]	2253
Fricas [F]	2254
Sympy [F(-2)]	2254
Maxima [F]	2254
Giac [F]	2255
Mupad [F(-1)]	2255
Reduce [F]	2255

Optimal result

Integrand size = 22, antiderivative size = 129

$$\int (a + bx)^{-n}(c + dx)(e + fx)^n dx = \frac{d(a + bx)^{1-n}(e + fx)^{1+n}}{2bf} + \frac{(2bcf - bde(1 - n) - adf(1 + n))(a + bx)^{1-n}(e + fx)^{1+n} \text{Hypergeometric2F1}\left(1, 2, 2 - n, -\frac{f(a+bx)}{be-af}\right)}{2bf(be - af)(1 - n)}$$

output

```
1/2*d*(b*x+a)^(1-n)*(f*x+e)^(1+n)/b/f+1/2*(2*b*c*f-b*d*e*(1-n)-a*d*f*(1+n)
)*(b*x+a)^(1-n)*(f*x+e)^(1+n)*hypergeom([1, 2], [2-n], -f*(b*x+a)/(-a*f+b*e)
)/b/f/(-a*f+b*e)/(1-n)
```

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.84

$$\int (a + bx)^{-n}(c + dx)(e + fx)^n dx = \frac{(a + bx)^{-n}(e + fx)^{1+n} \left(df(a + bx) + \frac{(2bcf + bde(-1+n) - adf(1+n)) \left(\frac{f(a+bx)}{-be+af} \right)^n \text{Hypergeometric2F1}\left(n, 1+n, 2+n, \frac{b(e+fx)}{be-af}\right)}{1+n} \right)}{2bf^2}$$

input

```
Integrate[((c + d*x)*(e + f*x)^n)/(a + b*x)^n,x]
```


output

```
((e + f*x)^(1 + n)*(d*f*(a + b*x) + ((2*b*c*f + b*d*e*(-1 + n) - a*d*f*(1 + n))*(f*(a + b*x))/(-(b*e) + a*f))^n*Hypergeometric2F1[n, 1 + n, 2 + n, (b*(e + f*x))/(b*e - a*f)]/(1 + n))/(2*b*f^2*(a + b*x)^n)
```

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.04, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.136$, Rules used = {90, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + dx)(a + bx)^{-n}(e + fx)^n dx$$

$$\downarrow 90$$

$$\frac{(-adf(n + 1) + 2bcf - bde(1 - n)) \int (a + bx)^{-n}(e + fx)^n dx}{2bf} + \frac{d(a + bx)^{1-n}(e + fx)^{n+1}}{2bf}$$

$$\downarrow 80$$

$$\frac{(a + bx)^{-n} \left(-\frac{f(a+bx)}{be-af}\right)^n (-adf(n + 1) + 2bcf - bde(1 - n)) \int (e + fx)^n \left(-\frac{bxf}{be-af} - \frac{af}{be-af}\right)^{-n} dx}{\frac{2bf}{d(a + bx)^{1-n}(e + fx)^{n+1}}} +$$

$$\downarrow 79$$

$$\frac{(a + bx)^{-n}(e + fx)^{n+1} \left(-\frac{f(a+bx)}{be-af}\right)^n (-adf(n + 1) + 2bcf - bde(1 - n)) \text{Hypergeometric2F1}\left(n, n + 1, n + 2, \frac{b}{b}\right)}{\frac{2bf^2(n + 1)}{d(a + bx)^{1-n}(e + fx)^{n+1}}}$$

input

```
Int[((c + d*x)*(e + f*x)^n)/(a + b*x)^n,x]
```

output

```
(d*(a + b*x)^(1 - n)*(e + f*x)^(1 + n))/(2*b*f) + ((2*b*c*f - b*d*e*(1 - n) - a*d*f*(1 + n))*(-(f*(a + b*x))/(b*e - a*f)))^n*(e + f*x)^(1 + n)*Hypergeometric2F1[n, 1 + n, 2 + n, (b*(e + f*x))/(b*e - a*f)]/(2*b*f^2*(1 + n)*(a + b*x)^n)
```

Defintions of rubi rules used

rule 79

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))
```

rule 80

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(c + d*x)^FracPart[n]/(b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])
```

rule 90

```
Int[((a_.) + (b_)*(x_))*((c_.) + (d_)*(x_))^(n_.)*((e_.) + (f_)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]
```

Maple [F]

$$\int (xd + c)(fx + e)^n (bx + a)^{-n} dx$$

input

```
int((d*x+c)*(f*x+e)^n/((b*x+a)^n),x)
```

output

```
int((d*x+c)*(f*x+e)^n/((b*x+a)^n),x)
```

Fricas [F]

$$\int (a + bx)^{-n}(c + dx)(e + fx)^n dx = \int \frac{(dx + c)(fx + e)^n}{(bx + a)^n} dx$$

input `integrate((d*x+c)*(f*x+e)^n/((b*x+a)^n),x, algorithm="fricas")`

output `integral((d*x + c)*(f*x + e)^n/(b*x + a)^n, x)`

Sympy [F(-2)]

Exception generated.

$$\int (a + bx)^{-n}(c + dx)(e + fx)^n dx = \text{Exception raised: HeuristicGCDFailed}$$

input `integrate((d*x+c)*(f*x+e)**n/((b*x+a)**n),x)`

output `Exception raised: HeuristicGCDFailed >> no luck`

Maxima [F]

$$\int (a + bx)^{-n}(c + dx)(e + fx)^n dx = \int \frac{(dx + c)(fx + e)^n}{(bx + a)^n} dx$$

input `integrate((d*x+c)*(f*x+e)^n/((b*x+a)^n),x, algorithm="maxima")`

output `integrate((d*x + c)*(f*x + e)^n/(b*x + a)^n, x)`

Giac [F]

$$\int (a + bx)^{-n} (c + dx) (e + fx)^n dx = \int \frac{(dx + c)(fx + e)^n}{(bx + a)^n} dx$$

input `integrate((d*x+c)*(f*x+e)^n/((b*x+a)^n),x, algorithm="giac")`

output `integrate((d*x + c)*(f*x + e)^n/(b*x + a)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + bx)^{-n} (c + dx) (e + fx)^n dx = \int \frac{(e + fx)^n (c + dx)}{(a + bx)^n} dx$$

input `int(((e + f*x)^n*(c + d*x))/(a + b*x)^n,x)`

output `int(((e + f*x)^n*(c + d*x))/(a + b*x)^n, x)`

Reduce [F]

$$\int (a + bx)^{-n} (c + dx) (e + fx)^n dx = \left(\int \frac{(fx + e)^n}{(bx + a)^n} dx \right) c + \left(\int \frac{(fx + e)^n x}{(bx + a)^n} dx \right) d$$

input `int((d*x+c)*(f*x+e)^n/((b*x+a)^n),x)`

output `int((e + f*x)**n/(a + b*x)**n,x)*c + int(((e + f*x)**n*x)/(a + b*x)**n,x)*d`

3.259 $\int (a + bx)^{-n}(c + dx)(e + fx)^{-1+n} dx$

Optimal result	2256
Mathematica [A] (verified)	2256
Rubi [A] (verified)	2257
Maple [F]	2258
Fricas [F]	2259
Sympy [F(-2)]	2259
Maxima [F]	2259
Giac [F]	2260
Mupad [F(-1)]	2260
Reduce [F]	2260

Optimal result

Integrand size = 24, antiderivative size = 110

$$\int (a + bx)^{-n}(c + dx)(e + fx)^{-1+n} dx = \frac{d(a + bx)^{1-n}(e + fx)^n}{bf} - \frac{(bcf - bde(1 - n) - adfn)(a + bx)^{1-n}(e + fx)^n \operatorname{Hypergeometric2F1}\left(1, 1, 1 + n, \frac{b(e+fx)}{be-af}\right)}{bf(be - af)n}$$

output

```
d*(b*x+a)^(1-n)*(f*x+e)^n/b/f-(b*c*f-b*d*e*(1-n)-a*d*f*n)*(b*x+a)^(1-n)*(f*x+e)^n*hypergeom([1, 1],[1+n],b*(f*x+e)/(-a*f+b*e))/b/f/(-a*f+b*e)/n
```

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.12

$$\int (a + bx)^{-n}(c + dx)(e + fx)^{-1+n} dx = \frac{(a + bx)^{-n}(e + fx)^n \left(f(-de + cf)(a + bx) + \frac{(-b(cf + de(-1 + n)) + adfn) \left(\frac{f(a + bx)}{-be + af} \right)^n (e + fx) \operatorname{Hypergeometric2F1}(n, 1 + n, 2, \frac{f(a + bx)}{-be + af})}{1 + n} \right)}{f^2(-be + af)n}$$

input

```
Integrate[((c + d*x)*(e + f*x)^(-1 + n))/(a + b*x)^n,x]
```

output

```
((e + f*x)^n*(f*(-(d*e) + c*f)*(a + b*x) + ((-(b*(c*f + d*e*(-1 + n))) + a
*d*f*n)*((f*(a + b*x))/(-(b*e) + a*f))^n*(e + f*x)*Hypergeometric2F1[n, 1
+ n, 2 + n, (b*(e + f*x))/(b*e - a*f)]/(1 + n)))/(f^2*(-(b*e) + a*f)*n*(a
+ b*x)^n)
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.36, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {88, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + dx)(a + bx)^{-n}(e + fx)^{n-1} dx$$

$$\downarrow 88$$

$$\frac{(-adf n + bcf - bde(1 - n)) \int (a + bx)^{-n}(e + fx)^n dx}{fn(be - af)} + \frac{(a + bx)^{1-n}(de - cf)(e + fx)^n}{fn(be - af)}$$

$$\downarrow 80$$

$$\frac{(a + bx)^{-n} \left(-\frac{f(a+bx)}{be-af}\right)^n (-adf n + bcf - bde(1 - n)) \int (e + fx)^n \left(-\frac{bxf}{be-af} - \frac{af}{be-af}\right)^{-n} dx}{fn(be - af)} + \frac{(a + bx)^{1-n}(de - cf)(e + fx)^n}{fn(be - af)}$$

$$\downarrow 79$$

$$\frac{(a + bx)^{-n}(e + fx)^{n+1} \left(-\frac{f(a+bx)}{be-af}\right)^n (-adf n + bcf - bde(1 - n)) \text{Hypergeometric2F1}\left(n, n + 1, n + 2, \frac{b(e+fx)}{be-af}\right)}{f^2 n(n + 1)(be - af)} + \frac{(a + bx)^{1-n}(de - cf)(e + fx)^n}{fn(be - af)}$$

input

```
Int[((c + d*x)*(e + f*x)^(-1 + n))/(a + b*x)^n, x]
```

output

```
((d*e - c*f)*(a + b*x)^(1 - n)*(e + f*x)^n)/(f*(b*e - a*f)*n) + ((b*c*f -
b*d*e*(1 - n) - a*d*f*n)*(-(f*(a + b*x))/(b*e - a*f)))^n*(e + f*x)^(1 + n)
)*Hypergeometric2F1[n, 1 + n, 2 + n, (b*(e + f*x))/(b*e - a*f)]/(f^2*(b*e
- a*f)*n*(1 + n)*(a + b*x)^n)
```

Defintions of rubi rules used

rule 79

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[((
a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n)*Hypergeometric2F1[-n, m + 1
, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x]
&& !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m]
|| !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))
```

rule 80

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(c
+ d*x)^FracPart[n]/((b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d))
^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)
), x]^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !Integ
erQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])
```

rule 88

```
Int[((a_.) + (b_)*(x_))*((c_.) + (d_)*(x_))^(n_.)*((e_.) + (f_)*(x_))^(p
_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^Simplify[p + 1],
x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && !RationalQ[p] && SumSimpl
erQ[p, 1]
```

Maple [F]

$$\int (xd + c)(fx + e)^{-1+n} (bx + a)^{-n} dx$$

input

```
int((d*x+c)*(f*x+e)^(-1+n)/((b*x+a)^n),x)
```

output

```
int((d*x+c)*(f*x+e)^(-1+n)/((b*x+a)^n),x)
```

Fricas [F]

$$\int (a + bx)^{-n} (c + dx) (e + fx)^{-1+n} dx = \int \frac{(dx + c)(fx + e)^{n-1}}{(bx + a)^n} dx$$

input `integrate((d*x+c)*(f*x+e)^(-1+n)/((b*x+a)^n),x, algorithm="fricas")`

output `integral((d*x + c)*(f*x + e)^(n - 1)/(b*x + a)^n, x)`

Sympy [F(-2)]

Exception generated.

$$\int (a + bx)^{-n} (c + dx) (e + fx)^{-1+n} dx = \text{Exception raised: HeuristicGCDFailed}$$

input `integrate((d*x+c)*(f*x+e)**(-1+n)/((b*x+a)**n),x)`

output `Exception raised: HeuristicGCDFailed >> no luck`

Maxima [F]

$$\int (a + bx)^{-n} (c + dx) (e + fx)^{-1+n} dx = \int \frac{(dx + c)(fx + e)^{n-1}}{(bx + a)^n} dx$$

input `integrate((d*x+c)*(f*x+e)^(-1+n)/((b*x+a)^n),x, algorithm="maxima")`

output `integrate((d*x + c)*(f*x + e)^(n - 1)/(b*x + a)^n, x)`

Giac [F]

$$\int (a + bx)^{-n} (c + dx) (e + fx)^{-1+n} dx = \int \frac{(dx + c)(fx + e)^{n-1}}{(bx + a)^n} dx$$

input `integrate((d*x+c)*(f*x+e)^(-1+n)/((b*x+a)^n),x, algorithm="giac")`

output `integrate((d*x + c)*(f*x + e)^(n - 1)/(b*x + a)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + bx)^{-n} (c + dx) (e + fx)^{-1+n} dx = \int \frac{(e + fx)^{n-1} (c + dx)}{(a + bx)^n} dx$$

input `int(((e + f*x)^(n - 1)*(c + d*x))/(a + b*x)^n,x)`

output `int(((e + f*x)^(n - 1)*(c + d*x))/(a + b*x)^n, x)`

Reduce [F]

$$\int (a + bx)^{-n} (c + dx) (e + fx)^{-1+n} dx = \left(\int \frac{(fx + e)^n}{(bx + a)^n e + (bx + a)^n fx} dx \right) c + \left(\int \frac{(fx + e)^n x}{(bx + a)^n e + (bx + a)^n fx} dx \right) d$$

input `int((d*x+c)*(f*x+e)^(-1+n)/((b*x+a)^n),x)`

output `int((e + f*x)**n/((a + b*x)**n*e + (a + b*x)**n*f*x),x)*c + int(((e + f*x)**n*x)/((a + b*x)**n*e + (a + b*x)**n*f*x),x)*d`

3.260 $\int (a + bx)^{-n}(c + dx)(e + fx)^{-2+n} dx$

Optimal result	2261
Mathematica [A] (verified)	2261
Rubi [A] (verified)	2262
Maple [F]	2263
Fricas [F]	2264
Sympy [F(-2)]	2264
Maxima [F]	2264
Giac [F]	2265
Mupad [F(-1)]	2265
Reduce [F]	2265

Optimal result

Integrand size = 24, antiderivative size = 111

$$\int (a + bx)^{-n}(c + dx)(e + fx)^{-2+n} dx$$

$$= -\frac{(de - cf)(a + bx)^{1-n}(e + fx)^{-1+n}}{f(be - af)(1 - n)}$$

$$- \frac{d(a + bx)^{1-n}(e + fx)^n \operatorname{Hypergeometric2F1}\left(1, 1, 1 + n, \frac{b(e+fx)}{be-af}\right)}{f(be - af)n}$$

output

```

-(-c*f+d*e)*(b*x+a)^(1-n)*(f*x+e)^(-1+n)/f/(-a*f+b*e)/(1-n)-d*(b*x+a)^(1-n)
)*(f*x+e)^n*hypergeom([1, 1],[1+n],b*(f*x+e)/(-a*f+b*e))/f/(-a*f+b*e)/n
    
```

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.95

$$\int (a + bx)^{-n}(c + dx)(e + fx)^{-2+n} dx$$

$$= \frac{(a + bx)^{-n}(e + fx)^n \left(\frac{f(de - cf)(a + bx)}{(be - af)(-1 + n)(e + fx)} + \frac{d \left(\frac{f(a + bx)}{-be + af} \right)^n \operatorname{Hypergeometric2F1}\left(n, n, 1 + n, \frac{b(e + fx)}{be - af}\right)}{n} \right)}{f^2}$$

input `Integrate[((c + d*x)*(e + f*x)^(-2 + n))/(a + b*x)^n,x]`

output `((e + f*x)^n*((f*(d*e - c*f)*(a + b*x))/((b*e - a*f)*(-1 + n)*(e + f*x)) + (d*((f*(a + b*x))/(-b*e) + a*f))^n*Hypergeometric2F1[n, n, 1 + n, (b*(e + f*x))/(b*e - a*f)]/n))/(f^2*(a + b*x)^n)`

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.06, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {88, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c + dx)(a + bx)^{-n}(e + fx)^{n-2} dx \\
 & \quad \downarrow 88 \\
 & \frac{d \int (a + bx)^{-n}(e + fx)^{n-1} dx}{f} - \frac{(a + bx)^{1-n}(de - cf)(e + fx)^{n-1}}{f(1 - n)(be - af)} \\
 & \quad \downarrow 80 \\
 & \frac{d(a + bx)^{-n} \left(-\frac{f(a+bx)}{be-af}\right)^n \int (e + fx)^{n-1} \left(-\frac{bxf}{be-af} - \frac{af}{be-af}\right)^{-n} dx}{\frac{f}{(a + bx)^{1-n}(de - cf)(e + fx)^{n-1}}} - \frac{f}{f(1 - n)(be - af)} \\
 & \quad \downarrow 79 \\
 & \frac{d(a + bx)^{-n}(e + fx)^n \left(-\frac{f(a+bx)}{be-af}\right)^n \text{Hypergeometric2F1}\left(n, n, n + 1, \frac{b(e+fx)}{be-af}\right)}{\frac{f^2 n}{(a + bx)^{1-n}(de - cf)(e + fx)^{n-1}}} - \frac{f}{f(1 - n)(be - af)}
 \end{aligned}$$

input `Int[((c + d*x)*(e + f*x)^(-2 + n))/(a + b*x)^n,x]`

output

```

-(((d*e - c*f)*(a + b*x)^(1 - n)*(e + f*x)^(-1 + n))/(f*(b*e - a*f)*(1 - n
))) + (d*(-((f*(a + b*x))/(b*e - a*f)))^n*(e + f*x)^n*Hypergeometric2F1[n,
n, 1 + n, (b*(e + f*x))/(b*e - a*f)]/(f^2*n*(a + b*x)^n)

```

Defintions of rubi rules used

rule 79

```

Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(((
a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n)*Hypergeometric2F1[-n, m + 1
, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x]
&& !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m]
|| !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

```

rule 80

```

Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(c
+ d*x)^FracPart[n]/((b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))
^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)
), x]^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !Integ
erQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])

```

rule 88

```

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e)), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^Simplify[p + 1],
x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && !RationalQ[p] && SumSimpl
erQ[p, 1]

```

Maple [F]

$$\int (xd + c)(fx + e)^{n-2}(bx + a)^{-n} dx$$

input

```
int((d*x+c)*(f*x+e)^(n-2)/((b*x+a)^n),x)
```

output

```
int((d*x+c)*(f*x+e)^(n-2)/((b*x+a)^n),x)
```

Fricas [F]

$$\int (a + bx)^{-n} (c + dx) (e + fx)^{-2+n} dx = \int \frac{(dx + c)(fx + e)^{n-2}}{(bx + a)^n} dx$$

input `integrate((d*x+c)*(f*x+e)^(-2+n)/((b*x+a)^n),x, algorithm="fricas")`

output `integral((d*x + c)*(f*x + e)^(n - 2)/(b*x + a)^n, x)`

Sympy [F(-2)]

Exception generated.

$$\int (a + bx)^{-n} (c + dx) (e + fx)^{-2+n} dx = \text{Exception raised: HeuristicGCDFailed}$$

input `integrate((d*x+c)*(f*x+e)**(-2+n)/((b*x+a)**n),x)`

output `Exception raised: HeuristicGCDFailed >> no luck`

Maxima [F]

$$\int (a + bx)^{-n} (c + dx) (e + fx)^{-2+n} dx = \int \frac{(dx + c)(fx + e)^{n-2}}{(bx + a)^n} dx$$

input `integrate((d*x+c)*(f*x+e)^(-2+n)/((b*x+a)^n),x, algorithm="maxima")`

output `integrate((d*x + c)*(f*x + e)^(n - 2)/(b*x + a)^n, x)`

Giac [F]

$$\int (a + bx)^{-n} (c + dx) (e + fx)^{-2+n} dx = \int \frac{(dx + c)(fx + e)^{n-2}}{(bx + a)^n} dx$$

input `integrate((d*x+c)*(f*x+e)^(-2+n)/((b*x+a)^n),x, algorithm="giac")`

output `integrate((d*x + c)*(f*x + e)^(n - 2)/(b*x + a)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + bx)^{-n} (c + dx) (e + fx)^{-2+n} dx = \int \frac{(e + fx)^{n-2} (c + dx)}{(a + bx)^n} dx$$

input `int(((e + f*x)^(n - 2)*(c + d*x))/(a + b*x)^n,x)`

output `int(((e + f*x)^(n - 2)*(c + d*x))/(a + b*x)^n, x)`

Reduce [F]

$$\begin{aligned} & \int (a + bx)^{-n} (c + dx) (e + fx)^{-2+n} dx \\ &= \left(\int \frac{(fx + e)^n}{(bx + a)^n e^2 + 2(bx + a)^n efx + (bx + a)^n f^2 x^2} dx \right) c \\ & \quad + \left(\int \frac{(fx + e)^n x}{(bx + a)^n e^2 + 2(bx + a)^n efx + (bx + a)^n f^2 x^2} dx \right) d \end{aligned}$$

input `int((d*x+c)*(f*x+e)^(-2+n)/((b*x+a)^n),x)`

output

```
int((e + f*x)**n/((a + b*x)**n*e**2 + 2*(a + b*x)**n*e*f*x + (a + b*x)**n*
f**2*x**2),x)*c + int(((e + f*x)**n*x)/((a + b*x)**n*e**2 + 2*(a + b*x)**n
*e*f*x + (a + b*x)**n*f**2*x**2),x)*d
```

3.261 $\int (a + bx)^{-n}(c + dx)(e + fx)^{-3+n} dx$

Optimal result	2267
Mathematica [A] (verified)	2267
Rubi [A] (verified)	2268
Maple [A] (verified)	2269
Fricas [B] (verification not implemented)	2270
Sympy [F(-2)]	2270
Maxima [F]	2271
Giac [B] (verification not implemented)	2271
Mupad [B] (verification not implemented)	2272
Reduce [F]	2273

Optimal result

Integrand size = 24, antiderivative size = 123

$$\int (a + bx)^{-n}(c + dx)(e + fx)^{-3+n} dx$$

$$= -\frac{(de - cf)(a + bx)^{1-n}(e + fx)^{-2+n}}{f(be - af)(2 - n)}$$

$$+ \frac{(bcf - adf(2 - n) + bd(e - en))(a + bx)^{1-n}(e + fx)^{-1+n}}{f(be - af)^2(1 - n)(2 - n)}$$

output

$$-(-c*f+d*e)*(b*x+a)^{(1-n)}*(f*x+e)^{(-2+n)}/f/(-a*f+b*e)/(2-n)+(b*c*f-a*d*f*(2-n)+b*d*(-e*n+e))*(b*x+a)^{(1-n)}*(f*x+e)^{(-1+n)}/f/(-a*f+b*e)^2/(1-n)/(2-n)$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.68

$$\int (a + bx)^{-n}(c + dx)(e + fx)^{-3+n} dx$$

$$= \frac{(a + bx)^{1-n}(e + fx)^{-2+n}(-ade + acf(-1 + n) + adf(-2 + n)x - bde(-1 + n)x + bc(-e(-2 + n) + f))}{(be - af)^2(-2 + n)(-1 + n)}$$

input

`Integrate[((c + d*x)*(e + f*x)^(-3 + n))/(a + b*x)^n,x]`

output

```
((a + b*x)^(1 - n)*(e + f*x)^(-2 + n)*(-(a*d*e) + a*c*f*(-1 + n) + a*d*f*(-2 + n)*x - b*d*e*(-1 + n)*x + b*c*(-(e*(-2 + n)) + f*x)))/((b*e - a*f)^2*(-2 + n)*(-1 + n))
```

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {88, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + dx)(a + bx)^{-n}(e + fx)^{n-3} dx$$

$$\downarrow 88$$

$$\frac{(-adf(2 - n) + bcf + bd(e - en)) \int (a + bx)^{-n}(e + fx)^{n-2} dx}{\frac{f(2 - n)(be - af)}{(a + bx)^{1-n}(de - cf)(e + fx)^{n-2}}}$$

$$\downarrow 48$$

$$\frac{(a + bx)^{1-n}(e + fx)^{n-1}(-adf(2 - n) + bcf + bd(e - en))}{f(1 - n)(2 - n)(be - af)^2} - \frac{(a + bx)^{1-n}(de - cf)(e + fx)^{n-2}}{f(2 - n)(be - af)}$$

input

```
Int[((c + d*x)*(e + f*x)^(-3 + n))/(a + b*x)^n,x]
```

output

```
-(((d*e - c*f)*(a + b*x)^(1 - n)*(e + f*x)^(-2 + n))/(f*(b*e - a*f)*(2 - n))) + ((b*c*f - a*d*f*(2 - n) + b*d*(e - e*n))*(a + b*x)^(1 - n)*(e + f*x)^(-1 + n))/(f*(b*e - a*f)^2*(1 - n)*(2 - n))
```


Maxima [F]

$$\int (a + bx)^{-n} (c + dx) (e + fx)^{-3+n} dx = \int \frac{(dx + c)(fx + e)^{n-3}}{(bx + a)^n} dx$$

input `integrate((d*x+c)*(f*x+e)^(-3+n)/(b*x+a)^n),x, algorithm="maxima")`

output `integrate((d*x + c)*(f*x + e)^(n - 3)/(b*x + a)^n, x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1050 vs. $2(115) = 230$.

Time = 0.15 (sec) , antiderivative size = 1050, normalized size of antiderivative = 8.54

$$\int (a + bx)^{-n} (c + dx) (e + fx)^{-3+n} dx = \text{Too large to display}$$

input `integrate((d*x+c)*(f*x+e)^(-3+n)/(b*x+a)^n),x, algorithm="giac")`

output

```

-(b^2*d*e*f*n*x^3*e^(n*log(f*x + e) - 3*log(f*x + e))/(b*x + a)^n - a*b*d*
f^2*n*x^3*e^(n*log(f*x + e) - 3*log(f*x + e))/(b*x + a)^n + b^2*d*e^2*n*x^
2*e^(n*log(f*x + e) - 3*log(f*x + e))/(b*x + a)^n + b^2*c*e*f*n*x^2*e^(n*log
og(f*x + e) - 3*log(f*x + e))/(b*x + a)^n - a*b*c*f^2*n*x^2*e^(n*log(f*x +
e) - 3*log(f*x + e))/(b*x + a)^n - a^2*d*f^2*n*x^2*e^(n*log(f*x + e) - 3*
log(f*x + e))/(b*x + a)^n - b^2*d*e*f*x^3*e^(n*log(f*x + e) - 3*log(f*x +
e))/(b*x + a)^n - b^2*c*f^2*x^3*e^(n*log(f*x + e) - 3*log(f*x + e))/(b*x +
a)^n + 2*a*b*d*f^2*x^3*e^(n*log(f*x + e) - 3*log(f*x + e))/(b*x + a)^n +
b^2*c*e^2*n*x*e^(n*log(f*x + e) - 3*log(f*x + e))/(b*x + a)^n + a*b*d*e^2*
n*x*e^(n*log(f*x + e) - 3*log(f*x + e))/(b*x + a)^n - a^2*d*e*f*n*x*e^(n*log
og(f*x + e) - 3*log(f*x + e))/(b*x + a)^n - a^2*c*f^2*n*x*e^(n*log(f*x + e
) - 3*log(f*x + e))/(b*x + a)^n - b^2*d*e^2*x^2*e^(n*log(f*x + e) - 3*log(
f*x + e))/(b*x + a)^n - 3*b^2*c*e*f*x^2*e^(n*log(f*x + e) - 3*log(f*x + e)
)/(b*x + a)^n + 2*a*b*d*e*f*x^2*e^(n*log(f*x + e) - 3*log(f*x + e))/(b*x +
a)^n + 2*a^2*d*f^2*x^2*e^(n*log(f*x + e) - 3*log(f*x + e))/(b*x + a)^n +
a*b*c*e^2*n*e^(n*log(f*x + e) - 3*log(f*x + e))/(b*x + a)^n - a^2*c*e*f*n*
e^(n*log(f*x + e) - 3*log(f*x + e))/(b*x + a)^n - 2*b^2*c*e^2*x*e^(n*log(f
*x + e) - 3*log(f*x + e))/(b*x + a)^n - 2*a*b*c*e*f*x*e^(n*log(f*x + e) -
3*log(f*x + e))/(b*x + a)^n + 3*a^2*d*e*f*x*e^(n*log(f*x + e) - 3*log(f*x
+ e))/(b*x + a)^n + a^2*c*f^2*x*e^(n*log(f*x + e) - 3*log(f*x + e))/(b*x...

```

Mupad [B] (verification not implemented)

Time = 1.26 (sec) , antiderivative size = 360, normalized size of antiderivative = 2.93

$$\begin{aligned}
 & \int (a + bx)^{-n} (c + dx) (e + fx)^{-3+n} dx \\
 &= \frac{bf^3x^3(e + fx)^{n-3}(bcf - 2adf + bde + adfn - bden)}{(af - be)^2(a + bx)^n(n^2 - 3n + 2)} \\
 & \quad - \frac{x^2(e + fx)^{n-3}(2a^2df^2 - b^2de^2 - 3b^2cef - a^2df^2n + b^2de^2n + 2abdef - abc f^2n + b^2cef n)}{(af - be)^2(a + bx)^n(n^2 - 3n + 2)} \\
 & \quad - \frac{ae(e + fx)^{n-3}(acf + ade - 2bce - acfn + bcn)}{(af - be)^2(a + bx)^n(n^2 - 3n + 2)} \\
 & \quad - \frac{x(e + fx)^{n-3}(a^2cf^2 - 2b^2ce^2 + 3a^2def - a^2cf^2n + b^2ce^2n - 2abcef + abde^2n - a^2defn)}{(af - be)^2(a + bx)^n(n^2 - 3n + 2)}
 \end{aligned}$$

input

```
int(((e + f*x)^(n - 3)*(c + d*x))/(a + b*x)^n,x)
```

output

```
(b*f*x^3*(e + f*x)^(n - 3)*(b*c*f - 2*a*d*f + b*d*e + a*d*f*n - b*d*e*n))/
((a*f - b*e)^2*(a + b*x)^n*(n^2 - 3*n + 2)) - (x^2*(e + f*x)^(n - 3)*(2*a^
2*d*f^2 - b^2*d*e^2 - 3*b^2*c*e*f - a^2*d*f^2*n + b^2*d*e^2*n + 2*a*b*d*e*
f - a*b*c*f^2*n + b^2*c*e*f*n))/((a*f - b*e)^2*(a + b*x)^n*(n^2 - 3*n + 2)
) - (a*e*(e + f*x)^(n - 3)*(a*c*f + a*d*e - 2*b*c*e - a*c*f*n + b*c*e*n))/
((a*f - b*e)^2*(a + b*x)^n*(n^2 - 3*n + 2)) - (x*(e + f*x)^(n - 3)*(a^2*c*
f^2 - 2*b^2*c*e^2 + 3*a^2*d*e*f - a^2*c*f^2*n + b^2*c*e^2*n - 2*a*b*c*e*f
+ a*b*d*e^2*n - a^2*d*e*f*n))/((a*f - b*e)^2*(a + b*x)^n*(n^2 - 3*n + 2))
```

Reduce [F]

$$\int (a + bx)^{-n} (c + dx) (e + fx)^{-3+n} dx$$

$$= \left(\int \frac{(fx + e)^n}{(bx + a)^n e^3 + 3(bx + a)^n e^2 fx + 3(bx + a)^n e f^2 x^2 + (bx + a)^n f^3 x^3} dx \right) c$$

$$+ \left(\int \frac{(fx + e)^n x}{(bx + a)^n e^3 + 3(bx + a)^n e^2 fx + 3(bx + a)^n e f^2 x^2 + (bx + a)^n f^3 x^3} dx \right) d$$

input

```
int((d*x+c)*(f*x+e)^(-3+n)/((b*x+a)^n),x)
```

output

```
int((e + f*x)**n/((a + b*x)**n*e**3 + 3*(a + b*x)**n*e**2*f*x + 3*(a + b*x)
)**n*e*f**2*x**2 + (a + b*x)**n*f**3*x**3),x)*c + int(((e + f*x)**n*x)/((a
+ b*x)**n*e**3 + 3*(a + b*x)**n*e**2*f*x + 3*(a + b*x)**n*e*f**2*x**2 + (
a + b*x)**n*f**3*x**3),x)*d
```

3.262 $\int (a + bx)^{-n}(c + dx)(e + fx)^{-4+n} dx$

Optimal result	2274
Mathematica [A] (verified)	2275
Rubi [A] (verified)	2275
Maple [B] (verified)	2277
Fricas [B] (verification not implemented)	2278
Sympy [F(-2)]	2278
Maxima [F]	2279
Giac [F]	2279
Mupad [B] (verification not implemented)	2280
Reduce [F]	2281

Optimal result

Integrand size = 24, antiderivative size = 205

$$\begin{aligned} & \int (a + bx)^{-n}(c + dx)(e + fx)^{-4+n} dx \\ &= -\frac{(de - cf)(a + bx)^{1-n}(e + fx)^{-3+n}}{f(be - af)(3 - n)} \\ & \quad + \frac{(2bcf - adf(3 - n) + bd(e - en))(a + bx)^{1-n}(e + fx)^{-2+n}}{f(be - af)^2(2 - n)(3 - n)} \\ & \quad + \frac{b(2bcf - adf(3 - n) + bd(e - en))(a + bx)^{1-n}(e + fx)^{-1+n}}{f(be - af)^3(1 - n)(2 - n)(3 - n)} \end{aligned}$$

output

```

-(-c*f+d*e)*(b*x+a)^(1-n)*(f*x+e)^(-3+n)/f/(-a*f+b*e)/(3-n)+(2*b*c*f-a*d*f
*(3-n)+b*d*(-e*n+e))*(b*x+a)^(1-n)*(f*x+e)^(-2+n)/f/(-a*f+b*e)^2/(2-n)/(3-
n)+b*(2*b*c*f-a*d*f*(3-n)+b*d*(-e*n+e))*(b*x+a)^(1-n)*(f*x+e)^(-1+n)/f/(-a
*f+b*e)^3/(1-n)/(2-n)/(3-n)

```


$$\frac{(-adf(3-n) + 2bcf + bde(1-n)) \left(\frac{b \int (a+bx)^{-n} (e+fx)^{n-2} dx}{(2-n)(be-af)} + \frac{(a+bx)^{1-n} (e+fx)^{n-2}}{(2-n)(be-af)} \right)}{\frac{f(3-n)(be-af)}{(a+bx)^{1-n}(de-cf)(e+fx)^{n-3}} \cdot \frac{f(3-n)(be-af)}{f(3-n)(be-af)}}}$$

↓ 48

$$\frac{\left(\frac{(a+bx)^{1-n} (e+fx)^{n-2}}{(2-n)(be-af)} + \frac{b(a+bx)^{1-n} (e+fx)^{n-1}}{(1-n)(2-n)(be-af)^2} \right) (-adf(3-n) + 2bcf + bde(1-n))}{\frac{f(3-n)(be-af)}{(a+bx)^{1-n}(de-cf)(e+fx)^{n-3}} \cdot \frac{f(3-n)(be-af)}{f(3-n)(be-af)}}$$

input `Int[((c + d*x)*(e + f*x)^(-4 + n))/(a + b*x)^n,x]`

output `-(((d*e - c*f)*(a + b*x)^(1 - n)*(e + f*x)^(-3 + n))/(f*(b*e - a*f)*(3 - n))) + ((2*b*c*f + b*d*e*(1 - n) - a*d*f*(3 - n))*(((a + b*x)^(1 - n)*(e + f*x)^(-2 + n))/((b*e - a*f)*(2 - n)) + (b*(a + b*x)^(1 - n)*(e + f*x)^(-1 + n))/((b*e - a*f)^2*(1 - n)*(2 - n))))/(f*(b*e - a*f)*(3 - n))`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 55 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(Simplify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])`

rule 88

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p
_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^Simplify[p + 1],
x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && !RationalQ[p] && SumSimpl
erQ[p, 1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 504 vs. 2(205) = 410.

Time = 0.54 (sec) , antiderivative size = 505, normalized size of antiderivative = 2.46

method	result
gospers	$\frac{(bx+a)(fx+e)^{-3+n}(bx+a)^{-n}(a^2d f^2n^2x-2abdef n^2x+abdf^2n x^2+b^2de^2n^2x-b^2defn x^2+a^2c f^2n^2-4a^2d f^2nx-2abcef n^2+2abc f^2nx-a^3 f^3n^3-3a^2be f^2n^3+3ab^2e^2n^3)}{a^3 f^3n^3-3a^2be f^2n^3+3ab^2e^2n^3}$
orering	$\frac{(bx+a)(fx+e)(a^2d f^2n^2x-2abdef n^2x+abdf^2n x^2+b^2de^2n^2x-b^2defn x^2+a^2c f^2n^2-4a^2d f^2nx-2abcef n^2+2abc f^2nx-a^3 f^3n^3-3a^2be f^2n^3+3ab^2e^2n^3)}{a^3 f^3n^3-3a^2be f^2n^3+3ab^2e^2n^3}$
parallelrisc	Expression too large to display

input

```
int((d*x+c)*(f*x+e)^(-4+n)/((b*x+a)^n),x,method=_RETURNVERBOSE)
```

output

```
(b*x+a)*(f*x+e)^(-3+n)/((b*x+a)^n)/(a^3*f^3*n^3-3*a^2*b*e*f^2*n^3+3*a*b^2*
e^2*f*n^3-b^3*e^3*n^3-6*a^3*f^3*n^2+18*a^2*b*e*f^2*n^2-18*a*b^2*e^2*f*n^2+
6*b^3*e^3*n^2+11*a^3*f^3*n-33*a^2*b*e*f^2*n+33*a*b^2*e^2*f*n-11*b^3*e^3*n-
6*a^3*f^3+18*a^2*b*e*f^2-18*a*b^2*e^2*f+6*b^3*e^3)*(a^2*d*f^2*n^2*x-2*a*b*
d*e*f*n^2*x+a*b*d*f^2*n*x^2+b^2*d*e^2*n^2*x-b^2*d*e*f*n*x^2+a^2*c*f^2*n^2-
4*a^2*d*f^2*n*x-2*a*b*c*e*f*n^2+2*a*b*c*f^2*n*x+8*a*b*d*e*f*n*x-3*a*b*d*f^
2*x^2+b^2*c*e^2*n^2-2*b^2*c*e*f*n*x+2*b^2*c*f^2*x^2-4*b^2*d*e^2*n*x+b^2*d*
e*f*x^2-3*a^2*c*f^2*n-a^2*d*e*f*n+3*a^2*d*f^2*x+8*a*b*c*e*f*n-2*a*b*c*f^2*
x+a*b*d*e^2*n-10*a*b*d*e*f*x-5*b^2*c*e^2*n+6*b^2*c*e*f*x+3*b^2*d*e^2*x+2*a
^2*c*f^2+a^2*d*e*f-6*a*b*c*e*f-3*a*b*d*e^2+6*b^2*c*e^2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 884 vs. $2(191) = 382$.

Time = 0.13 (sec) , antiderivative size = 884, normalized size of antiderivative = 4.31

$$\int (a + bx)^{-n}(c + dx)(e + fx)^{-4+n} dx = \text{Too large to display}$$

input `integrate((d*x+c)*(f*x+e)^(-4+n)/((b*x+a)^n),x, algorithm="fricas")`

output

```
(2*a^3*c*e*f^2 + (b^3*d*e*f^2 + (2*b^3*c - 3*a*b^2*d)*f^3 - (b^3*d*e*f^2 -
a*b^2*d*f^3)*n)*x^4 + 3*(2*a*b^2*c - a^2*b*d)*e^3 - (6*a^2*b*c - a^3*d)*e
^2*f + (4*b^3*d*e^2*f + 4*(2*b^3*c - 3*a*b^2*d)*e*f^2 + (b^3*d*e^2*f - 2*a
*b^2*d*e*f^2 + a^2*b*d*f^3)*n^2 - (5*b^3*d*e^2*f + 2*(b^3*c - 4*a*b^2*d)*e
*f^2 - (2*a*b^2*c - 3*a^2*b*d)*f^3)*n)*x^3 + (a*b^2*c*e^3 - 2*a^2*b*c*e^2*
f + a^3*c*e*f^2)*n^2 + (3*b^3*d*e^3 - 9*a^2*b*d*e*f^2 + 3*a^3*d*f^3 + 3*(4
*b^3*c - 3*a*b^2*d)*e^2*f + (b^3*d*e^3 + (b^3*c - a*b^2*d)*e^2*f - (2*a*b^
2*c + a^2*b*d)*e*f^2 + (a^2*b*c + a^3*d)*f^3)*n^2 - (4*b^3*d*e^3 + (7*b^3*
c - 4*a*b^2*d)*e^2*f - 4*(2*a*b^2*c + a^2*b*d)*e*f^2 + (a^2*b*c + 4*a^3*d)
*f^3)*n)*x^2 - (3*a^3*c*e*f^2 + (5*a*b^2*c - a^2*b*d)*e^3 - (8*a^2*b*c - a
^3*d)*e^2*f)*n + (6*b^3*c*e^3 + 2*a^3*c*f^3 + 6*(a*b^2*c - 2*a^2*b*d)*e^2*
f - 2*(3*a^2*b*c - 2*a^3*d)*e*f^2 + (a^3*c*f^3 + (b^3*c + a*b^2*d)*e^3 - (
a*b^2*c + 2*a^2*b*d)*e^2*f - (a^2*b*c - a^3*d)*e*f^2)*n^2 - (3*a^3*c*f^3 +
(5*b^3*c + 3*a*b^2*d)*e^3 - (a*b^2*c + 8*a^2*b*d)*e^2*f - (7*a^2*b*c - 5*
a^3*d)*e*f^2)*n)*x*(f*x + e)^(n - 4)/((6*b^3*e^3 - 18*a*b^2*e^2*f + 18*a^
2*b*e*f^2 - 6*a^3*f^3 - (b^3*e^3 - 3*a*b^2*e^2*f + 3*a^2*b*e*f^2 - a^3*f^3)
)*n^3 + 6*(b^3*e^3 - 3*a*b^2*e^2*f + 3*a^2*b*e*f^2 - a^3*f^3)*n^2 - 11*(b^
3*e^3 - 3*a*b^2*e^2*f + 3*a^2*b*e*f^2 - a^3*f^3)*n)*(b*x + a)^n)
```

Sympy [F(-2)]

Exception generated.

$$\int (a + bx)^{-n}(c + dx)(e + fx)^{-4+n} dx = \text{Exception raised: HeuristicGCDFailed}$$

input `integrate((d*x+c)*(f*x+e)**(-4+n)/((b*x+a)**n),x)`

output Exception raised: HeuristicGCDFailed >> no luck

Maxima [F]

$$\int (a + bx)^{-n}(c + dx)(e + fx)^{-4+n} dx = \int \frac{(dx + c)(fx + e)^{n-4}}{(bx + a)^n} dx$$

input `integrate((d*x+c)*(f*x+e)^(-4+n)/((b*x+a)^n),x, algorithm="maxima")`

output `integrate((d*x + c)*(f*x + e)^(n - 4)/(b*x + a)^n, x)`

Giac [F]

$$\int (a + bx)^{-n}(c + dx)(e + fx)^{-4+n} dx = \int \frac{(dx + c)(fx + e)^{n-4}}{(bx + a)^n} dx$$

input `integrate((d*x+c)*(f*x+e)^(-4+n)/((b*x+a)^n),x, algorithm="giac")`

output `integrate((d*x + c)*(f*x + e)^(n - 4)/(b*x + a)^n, x)`

Mupad [B] (verification not implemented)

Time = 1.66 (sec) , antiderivative size = 869, normalized size of antiderivative = 4.24

$$\begin{aligned}
& \int (a + bx)^{-n} (c + dx) (e + fx)^{-4+n} dx \\
& = \frac{x (e + fx)^{n-4} (da^3 e f^2 n^2 - 5da^3 e f^2 n + 4da^3 e f^2 + ca^3 f^3 n^2 - 3ca^3 f^3 n + 2ca^3 f^3 - 2da^2 b e^2 f}{} \\
& + \frac{x^2 (e + fx)^{n-4} (da^3 f^3 n^2 - 4da^3 f^3 n + 3da^3 f^3 - da^2 b e f^2 n^2 + 4da^2 b e f^2 n - 9da^2 b e f^2 + ca^2}{} \\
& + \frac{a e (e + fx)^{n-4} (-da^2 e f n + da^2 e f + ca^2 f^2 n^2 - 3ca^2 f^2 n + 2ca^2 f^2 + d a b e^2 n - 3 d a b e^2 - 2 c}{(a f - b e)^3 (a + b x)^n (n^3 - 6n^2 + 11n - 6)} \\
& + \frac{b^2 f^2 x^4 (e + fx)^{n-4} (2bcf - 3adf + bde + adfn - bden)}{(a f - b e)^3 (a + b x)^n (n^3 - 6n^2 + 11n - 6)} \\
& + \frac{b f x^3 (e + fx)^{n-4} (4be + a f n - ben) (2bcf - 3adf + bde + adfn - bden)}{(a f - b e)^3 (a + b x)^n (n^3 - 6n^2 + 11n - 6)}
\end{aligned}$$

input

```
int(((e + f*x)^(n - 4)*(c + d*x))/(a + b*x)^n,x)
```

output

```
(x*(e + f*x)^(n - 4)*(2*a^3*c*f^3 + 6*b^3*c*e^3 + a^3*c*f^3*n^2 + b^3*c*e^3*n^2 + 4*a^3*d*e*f^2 - 3*a^3*c*f^3*n - 5*b^3*c*e^3*n + 6*a*b^2*c*e^2*f - 6*a^2*b*c*e*f^2 - 12*a^2*b*d*e^2*f - 3*a*b^2*d*e^3*n - 5*a^3*d*e*f^2*n + a*b^2*d*e^3*n^2 + a^3*d*e*f^2*n^2 + a*b^2*c*e^2*f*n + 7*a^2*b*c*e*f^2*n + 8*a^2*b*d*e^2*f*n - a*b^2*c*e^2*f*n^2 - a^2*b*c*e*f^2*n^2 - 2*a^2*b*d*e^2*f*n^2))/((a*f - b*e)^3*(a + b*x)^n*(11*n - 6*n^2 + n^3 - 6)) + (x^2*(e + f*x)^(n - 4)*(3*a^3*d*f^3 + 3*b^3*d*e^3 + a^3*d*f^3*n^2 + b^3*d*e^3*n^2 + 12*b^3*c*e^2*f - 4*a^3*d*f^3*n - 4*b^3*d*e^3*n - 9*a*b^2*d*e^2*f - 9*a^2*b*d*e*f^2 - a^2*b*c*f^3*n - 7*b^3*c*e^2*f*n + a^2*b*c*f^3*n^2 + b^3*c*e^2*f*n^2 + 8*a*b^2*c*e*f^2*n + 4*a*b^2*d*e^2*f*n + 4*a^2*b*d*e*f^2*n - 2*a*b^2*c*e*f^2*n^2 - a*b^2*d*e^2*f*n^2 - a^2*b*d*e*f^2*n^2))/((a*f - b*e)^3*(a + b*x)^n*(11*n - 6*n^2 + n^3 - 6)) + (a*e*(e + f*x)^(n - 4)*(2*a^2*c*f^2 + 6*b^2*c*e^2 + a^2*c*f^2*n^2 + b^2*c*e^2*n^2 - 3*a*b*d*e^2 + a^2*d*e*f - 3*a^2*c*f^2*n - 5*b^2*c*e^2*n - 6*a*b*c*e*f + a*b*d*e^2*n - a^2*d*e*f*n - 2*a*b*c*e*f*n^2 + 8*a*b*c*e*f*n))/((a*f - b*e)^3*(a + b*x)^n*(11*n - 6*n^2 + n^3 - 6)) + (b^2*f^2*x^4*(e + f*x)^(n - 4)*(2*b*c*f - 3*a*d*f + b*d*e + a*d*f*n - b*d*e*n))/((a*f - b*e)^3*(a + b*x)^n*(11*n - 6*n^2 + n^3 - 6)) + (b*f*x^3*(e + f*x)^(n - 4)*(4*b*e + a*f*n - b*e*n)*(2*b*c*f - 3*a*d*f + b*d*e + a*d*f*n - b*d*e*n))/((a*f - b*e)^3*(a + b*x)^n*(11*n - 6*n^2 + n^3 - 6))
```

Reduce [F]

$$\int (a + bx)^{-n} (c + dx) (e + fx)^{-4+n} dx$$

$$= \left(\int \frac{(fx + e)^n}{(bx + a)^n e^4 + 4(bx + a)^n e^3 fx + 6(bx + a)^n e^2 f^2 x^2 + 4(bx + a)^n e f^3 x^3 + (bx + a)^n f^4 x^4} dx \right) c$$

$$+ \left(\int \frac{(fx + e)^n x}{(bx + a)^n e^4 + 4(bx + a)^n e^3 fx + 6(bx + a)^n e^2 f^2 x^2 + 4(bx + a)^n e f^3 x^3 + (bx + a)^n f^4 x^4} dx \right) d$$

input

```
int((d*x+c)*(f*x+e)^(-4+n)/((b*x+a)^n),x)
```

output

```
int((e + f*x)**n/((a + b*x)**n*e**4 + 4*(a + b*x)**n*e**3*f*x + 6*(a + b*x)**n*e**2*f**2*x**2 + 4*(a + b*x)**n*e*f**3*x**3 + (a + b*x)**n*f**4*x**4),x)*c + int(((e + f*x)**n*x)/((a + b*x)**n*e**4 + 4*(a + b*x)**n*e**3*f*x + 6*(a + b*x)**n*e**2*f**2*x**2 + 4*(a + b*x)**n*e*f**3*x**3 + (a + b*x)**n*f**4*x**4),x)*d
```

3.263 $\int (a + bx)^{-n}(c + dx)(e + fx)^{-5+n} dx$

Optimal result	2282
Mathematica [A] (verified)	2283
Rubi [A] (verified)	2283
Maple [B] (verified)	2285
Fricas [B] (verification not implemented)	2286
Sympy [F(-2)]	2287
Maxima [F]	2288
Giac [F]	2288
Mupad [B] (verification not implemented)	2288
Reduce [F]	2289

Optimal result

Integrand size = 24, antiderivative size = 297

$$\begin{aligned} & \int (a + bx)^{-n}(c + dx)(e + fx)^{-5+n} dx \\ &= -\frac{(de - cf)(a + bx)^{1-n}(e + fx)^{-4+n}}{f(be - af)(4 - n)} \\ &+ \frac{(3bcf - adf(4 - n) + bd(e - en))(a + bx)^{1-n}(e + fx)^{-3+n}}{f(be - af)^2(3 - n)(4 - n)} \\ &+ \frac{2b(3bcf - adf(4 - n) + bd(e - en))(a + bx)^{1-n}(e + fx)^{-2+n}}{f(be - af)^3(2 - n)(3 - n)(4 - n)} \\ &+ \frac{2b^2(3bcf - adf(4 - n) + bd(e - en))(a + bx)^{1-n}(e + fx)^{-1+n}}{f(be - af)^4(1 - n)(2 - n)(3 - n)(4 - n)} \end{aligned}$$

output

```

-(-c*f+d*e)*(b*x+a)^(1-n)*(f*x+e)^(-4+n)/f/(-a*f+b*e)/(4-n)+(3*b*c*f-a*d*f
*(4-n)+b*d*(-e*n+e))*(b*x+a)^(1-n)*(f*x+e)^(-3+n)/f/(-a*f+b*e)^2/(3-n)/(4-
n)+2*b*(3*b*c*f-a*d*f*(4-n)+b*d*(-e*n+e))*(b*x+a)^(1-n)*(f*x+e)^(-2+n)/f/(
-a*f+b*e)^3/(2-n)/(3-n)/(4-n)+2*b^2*(3*b*c*f-a*d*f*(4-n)+b*d*(-e*n+e))*(b*
x+a)^(1-n)*(f*x+e)^(-1+n)/f/(-a*f+b*e)^4/(1-n)/(2-n)/(3-n)/(4-n)
    
```

Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.50

$$\int (a + bx)^{-n}(c + dx)(e + fx)^{-5+n} dx$$

$$= \frac{(a + bx)^{1-n}(e + fx)^{-4+n} \left(-de + cf - \frac{(3bcf + adf(-4+n) - bde(-1+n))(e + fx)((be - af)^2(-2+n)(-1+n) - 2b(be - af)(-1+n))}{(be - af)^3(-3+n)(-2+n)(-1+n)} \right)}{f(-be + af)(-4 + n)}$$

input `Integrate[((c + d*x)*(e + f*x)^(-5 + n))/(a + b*x)^n,x]`

output `((a + b*x)^(1 - n)*(e + f*x)^(-4 + n)*(-d*e) + c*f - ((3*b*c*f + a*d*f*(-4 + n) - b*d*e*(-1 + n))*(e + f*x)*((b*e - a*f)^2*(-2 + n)*(-1 + n) - 2*b*(b*e - a*f)*(-1 + n)*(e + f*x) + 2*b^2*(e + f*x)^2))/((b*e - a*f)^3*(-3 + n)*(-2 + n)*(-1 + n)))/(f*(-(b*e) + a*f)*(-4 + n))`

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 241, normalized size of antiderivative = 0.81, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {88, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + dx)(a + bx)^{-n}(e + fx)^{n-5} dx$$

$$\downarrow 88$$

$$\frac{(-adf(4 - n) + 3bcf + bde(1 - n)) \int (a + bx)^{-n}(e + fx)^{n-4} dx}{\frac{f(4 - n)(be - af)}{(a + bx)^{1-n}(de - cf)(e + fx)^{n-4}}}$$

$$\downarrow 55$$

$$\frac{(-adf(4-n) + 3bcf + bde(1-n)) \left(\frac{2b \int (a+bx)^{-n} (e+fx)^{n-3} dx}{(3-n)(be-af)} + \frac{(a+bx)^{1-n} (e+fx)^{n-3}}{(3-n)(be-af)} \right)}{\frac{f(4-n)(be-af)}{(a+bx)^{1-n}(de-cf)(e+fx)^{n-4}} \cdot f(4-n)(be-af)}$$

↓ 55

$$\frac{(-adf(4-n) + 3bcf + bde(1-n)) \left(\frac{2b \left(\frac{b \int (a+bx)^{-n} (e+fx)^{n-2} dx}{(2-n)(be-af)} + \frac{(a+bx)^{1-n} (e+fx)^{n-2}}{(2-n)(be-af)} \right)}{(3-n)(be-af)} + \frac{(a+bx)^{1-n} (e+fx)^{n-3}}{(3-n)(be-af)} \right)}{\frac{f(4-n)(be-af)}{(a+bx)^{1-n}(de-cf)(e+fx)^{n-4}} \cdot f(4-n)(be-af)}$$

↓ 48

$$\frac{\left(\frac{(a+bx)^{1-n} (e+fx)^{n-3}}{(3-n)(be-af)} + \frac{2b \left(\frac{(a+bx)^{1-n} (e+fx)^{n-2}}{(2-n)(be-af)} + \frac{b(a+bx)^{1-n} (e+fx)^{n-1}}{(1-n)(2-n)(be-af)^2} \right)}{(3-n)(be-af)} \right) (-adf(4-n) + 3bcf + bde(1-n))}{\frac{f(4-n)(be-af)}{(a+bx)^{1-n}(de-cf)(e+fx)^{n-4}} \cdot f(4-n)(be-af)}$$

input `Int[((c + d*x)*(e + f*x)^(-5 + n))/(a + b*x)^n,x]`

output `-(((d*e - c*f)*(a + b*x)^(1 - n)*(e + f*x)^(-4 + n))/(f*(b*e - a*f)*(4 - n))) + ((3*b*c*f + b*d*e*(1 - n) - a*d*f*(4 - n))*(((a + b*x)^(1 - n)*(e + f*x)^(-3 + n))/((b*e - a*f)*(3 - n)) + (2*b*((a + b*x)^(1 - n)*(e + f*x)^(-2 + n))/((b*e - a*f)*(2 - n)) + (b*(a + b*x)^(1 - n)*(e + f*x)^(-1 + n))/((b*e - a*f)^2*(1 - n)*(2 - n))))/(b*e - a*f)*(3 - n)))/(f*(b*e - a*f)*(4 - n))`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
 [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
 a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 55 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
 (a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
 implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
 c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
 c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
 lerQ[n, 1])`

rule 88 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
), x] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
 + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
 + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^Simplify[p + 1],
 x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && !RationalQ[p] && SumSimpl
 erQ[p, 1]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1187 vs. 2(297) = 594.

Time = 0.56 (sec) , antiderivative size = 1188, normalized size of antiderivative = 4.00

method	result	size
gospers	Expression too large to display	1188
orering	Expression too large to display	1193
parallelsch	Expression too large to display	3499

input `int((d*x+c)*(f*x+e)^(-5+n)/((b*x+a)^n),x,method=_RETURNVERBOSE)`

output

```
(b*x+a)*(f*x+e)^(-4+n)/((b*x+a)^n)/(a^4*f^4*n^4-4*a^3*b*e*f^3*n^4+6*a^2*b^2*e^2*f^2*n^4-4*a*b^3*e^3*f*n^4+b^4*e^4*n^4-10*a^4*f^4*n^3+40*a^3*b*e*f^3*n^3-60*a^2*b^2*e^2*f^2*n^3+40*a*b^3*e^3*f*n^3-10*b^4*e^4*n^3+35*a^4*f^4*n^2-140*a^3*b*e*f^3*n^2+210*a^2*b^2*e^2*f^2*n^2-140*a*b^3*e^3*f*n^2+35*b^4*e^4*n^2-50*a^4*f^4*n+200*a^3*b*e*f^3*n-300*a^2*b^2*e^2*f^2*n+200*a*b^3*e^3*f*n-50*b^4*e^4*n+24*a^4*f^4-96*a^3*b*e*f^3+144*a^2*b^2*e^2*f^2-96*a*b^3*e^3*f+24*b^4*e^4)*(a^3*d*f^3*n^3*x-3*a^2*b*d*e*f^2*n^3*x+2*a^2*b*d*f^3*n^2*x^2+3*a*b^2*d*e^2*f*n^3*x-4*a*b^2*d*e*f^2*n^2*x^2+2*a*b^2*d*f^3*n*x^3-b^3*d*e^3*n^3*x+2*b^3*d*e^2*f*n^2*x^2-2*b^3*d*e*f^2*n*x^3+a^3*c*f^3*n^3-7*a^3*d*f^3*n^2*x-3*a^2*b*c*e*f^2*n^3+3*a^2*b*c*f^3*n^2*x+22*a^2*b*d*e*f^2*n^2*x-10*a^2*b*d*f^3*n*x^2+3*a*b^2*c*e^2*f*n^3-6*a*b^2*c*e*f^2*n^2*x+6*a*b^2*c*f^3*n*x^2-23*a*b^2*d*e^2*f*n^2*x+20*a*b^2*d*e*f^2*n*x^2-8*a*b^2*d*f^3*x^3-b^3*c*e^3*n^3+3*b^3*c*e^2*f*n^2*x-6*b^3*c*e*f^2*n*x^2+6*b^3*c*f^3*x^3+8*b^3*d*e^3*n^2*x-10*b^3*d*e^2*f*n*x^2+2*b^3*d*e*f^2*x^3-6*a^3*c*f^3*n^2-a^3*d*e*f^2*n^2+14*a^3*d*f^3*n*x+21*a^2*b*c*e*f^2*n^2-9*a^2*b*c*f^3*n*x+2*a^2*b*d*e^2*f*n^2-53*a^2*b*d*e*f^2*n*x+8*a^2*b*d*f^3*x^2-24*a*b^2*c*e^2*f*n^2+30*a*b^2*c*e*f^2*n*x-6*a*b^2*c*f^3*x^2-a*b^2*d*e^3*n^2+58*a*b^2*d*e^2*f*n*x-34*a*b^2*d*e*f^2*x^2+9*b^3*c*e^3*n^2-21*b^3*c*e^2*f*n*x+24*b^3*c*e*f^2*x^2-19*b^3*d*e^3*n*x+8*b^3*d*e^2*f*x^2+11*a^3*c*f^3*n+3*a^3*d*e*f^2*n-8*a^3*d*f^3*x-42*a^2*b*c*e*f^2*n+6*a^2*b*c*f^3*x-10*a^2*b*d*e^2*f*n+34*a^2*b*d...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1740 vs. $2(273) = 546$.

Time = 0.18 (sec) , antiderivative size = 1740, normalized size of antiderivative = 5.86

$$\int (a + bx)^{-n}(c + dx)(e + fx)^{-5+n} dx = \text{Too large to display}$$

input

```
integrate((d*x+c)*(f*x+e)^(-5+n)/((b*x+a)^n),x, algorithm="fricas")
```

output

```

-(6*a^4*c*e*f^3 - 2*(b^4*d*e*f^3 + (3*b^4*c - 4*a*b^3*d)*f^4 - (b^4*d*e*f^
3 - a*b^3*d*f^4)*n)*x^5 - 12*(2*a*b^3*c - a^2*b^2*d)*e^4 + 4*(9*a^2*b^2*c
- 2*a^3*b*d)*e^3*f - 2*(12*a^3*b*c - a^4*d)*e^2*f^2 - 2*(5*b^4*d*e^2*f^2 +
5*(3*b^4*c - 4*a*b^3*d)*e*f^3 + (b^4*d*e^2*f^2 - 2*a*b^3*d*e*f^3 + a^2*b^
2*d*f^4)*n^2 - (6*b^4*d*e^2*f^2 + (3*b^4*c - 10*a*b^3*d)*e*f^3 - (3*a*b^3*
c - 4*a^2*b^2*d)*f^4)*n)*x^4 + (a*b^3*c*e^4 - 3*a^2*b^2*c*e^3*f + 3*a^3*b*
c*e^2*f^2 - a^4*c*e*f^3)*n^3 - (20*b^4*d*e^3*f + 20*(3*b^4*c - 4*a*b^3*d)*
e^2*f^2 - (b^4*d*e^3*f - 3*a*b^3*d*e^2*f^2 + 3*a^2*b^2*d*e*f^3 - a^3*b*d*f
^4)*n^3 + (10*b^4*d*e^3*f + (3*b^4*c - 25*a*b^3*d)*e^2*f^2 - 2*(3*a*b^3*c
- 10*a^2*b^2*d)*e*f^3 + (3*a^2*b^2*c - 5*a^3*b*d)*f^4)*n^2 - (29*b^4*d*e^3
*f + 3*(9*b^4*c - 22*a*b^3*d)*e^2*f^2 - (30*a*b^3*c - 41*a^2*b^2*d)*e*f^3
+ (3*a^2*b^2*c - 4*a^3*b*d)*f^4)*n)*x^3 + (6*a^4*c*e*f^3 - (9*a*b^3*c - a^
2*b^2*d)*e^4 + 2*(12*a^2*b^2*c - a^3*b*d)*e^3*f - (21*a^3*b*c - a^4*d)*e^2
*f^2)*n^2 - (12*b^4*d*e^4 - 48*a^2*b^2*d*e^2*f^2 + 32*a^3*b*d*e*f^3 - 8*a^
4*d*f^4 + 12*(5*b^4*c - 4*a*b^3*d)*e^3*f - (b^4*d*e^4 - 3*a*b^3*c*e^2*f^2
+ (b^4*c - 2*a*b^3*d)*e^3*f + (3*a^2*b^2*c + 2*a^3*b*d)*e*f^3 - (a^3*b*c +
a^4*d)*f^4)*n^3 + (8*b^4*d*e^4 + 2*(6*b^4*c - 7*a*b^3*d)*e^3*f - 3*(9*a*b
^3*c + a^2*b^2*d)*e^2*f^2 + 2*(9*a^2*b^2*c + 8*a^3*b*d)*e*f^3 - (3*a^3*b*c
+ 7*a^4*d)*f^4)*n^2 - (19*b^4*d*e^4 + (47*b^4*c - 36*a*b^3*d)*e^3*f - 15*
(4*a*b^3*c + a^2*b^2*d)*e^2*f^2 + (15*a^2*b^2*c + 46*a^3*b*d)*e*f^3 - 2...

```

Sympy [F(-2)]

Exception generated.

$$\int (a + bx)^{-n}(c + dx)(e + fx)^{-5+n} dx = \text{Exception raised: HeuristicGCDFailed}$$

input

```
integrate((d*x+c)*(f*x+e)**(-5+n)/((b*x+a)**n),x)
```

output

```
Exception raised: HeuristicGCDFailed >> no luck
```

Maxima [F]

$$\int (a + bx)^{-n}(c + dx)(e + fx)^{-5+n} dx = \int \frac{(dx + c)(fx + e)^{n-5}}{(bx + a)^n} dx$$

input `integrate((d*x+c)*(f*x+e)^(-5+n)/((b*x+a)^n),x, algorithm="maxima")`

output `integrate((d*x + c)*(f*x + e)^(n - 5)/(b*x + a)^n, x)`

Giac [F]

$$\int (a + bx)^{-n}(c + dx)(e + fx)^{-5+n} dx = \int \frac{(dx + c)(fx + e)^{n-5}}{(bx + a)^n} dx$$

input `integrate((d*x+c)*(f*x+e)^(-5+n)/((b*x+a)^n),x, algorithm="giac")`

output `integrate((d*x + c)*(f*x + e)^(n - 5)/(b*x + a)^n, x)`

Mupad [B] (verification not implemented)

Time = 2.39 (sec) , antiderivative size = 1659, normalized size of antiderivative = 5.59

$$\int (a + bx)^{-n}(c + dx)(e + fx)^{-5+n} dx = \text{Too large to display}$$

input `int(((e + f*x)^(n - 5)*(c + d*x))/(a + b*x)^n,x)`

output

```
(2*b^3*f^3*x^5*(e + f*x)^(n - 5)*(3*b*c*f - 4*a*d*f + b*d*e + a*d*f*n - b*
d*e*n))/((a*f - b*e)^4*(a + b*x)^n*(35*n^2 - 50*n - 10*n^3 + n^4 + 24)) -
(x*(e + f*x)^(n - 5)*(6*a^4*c*f^4 - 24*b^4*c*e^4 + 6*a^4*c*f^4*n^2 - 9*b^4
*c*e^4*n^2 - a^4*c*f^4*n^3 + b^4*c*e^4*n^3 + 10*a^4*d*e*f^3 - 11*a^4*c*f^4
*n + 26*b^4*c*e^4*n - 24*a*b^3*c*e^3*f - 24*a^3*b*c*e*f^3 + 12*a*b^3*d*e^4
*n - 17*a^4*d*e*f^3*n + 60*a^2*b^2*d*e^3*f - 40*a^3*b*d*e^2*f^2 - 7*a*b^3*
d*e^4*n^2 + a*b^3*d*e^4*n^3 + 8*a^4*d*e*f^3*n^2 - a^4*d*e*f^3*n^3 + 36*a^2
*b^2*c*e^2*f^2 - 45*a^2*b^2*c*e^2*f^2*n + 22*a^2*b^2*d*e^3*f*n^2 - 23*a^3*
b*d*e^2*f^2*n^2 - 3*a^2*b^2*d*e^3*f*n^3 + 3*a^3*b*d*e^2*f^2*n^3 - 10*a*b^3
*c*e^3*f*n + 40*a^3*b*c*e*f^3*n + 9*a^2*b^2*c*e^2*f^2*n^2 + 12*a*b^3*c*e^3
*f*n^2 - 18*a^3*b*c*e*f^3*n^2 - 2*a*b^3*c*e^3*f*n^3 + 2*a^3*b*c*e*f^3*n^3
- 55*a^2*b^2*d*e^3*f*n + 60*a^3*b*d*e^2*f^2*n))/((a*f - b*e)^4*(a + b*x)^n
*(35*n^2 - 50*n - 10*n^3 + n^4 + 24)) - (x^2*(e + f*x)^(n - 5)*(8*a^4*d*f^
4 - 12*b^4*d*e^4 + 7*a^4*d*f^4*n^2 - 8*b^4*d*e^4*n^2 - a^4*d*f^4*n^3 + b^4
*d*e^4*n^3 - 60*b^4*c*e^3*f - 14*a^4*d*f^4*n + 19*b^4*d*e^4*n + 48*a*b^3*d
*e^3*f - 32*a^3*b*d*e*f^3 - 2*a^3*b*c*f^4*n + 47*b^4*c*e^3*f*n + 3*a^3*b*c
*f^4*n^2 - a^3*b*c*f^4*n^3 - 12*b^4*c*e^3*f*n^2 + b^4*c*e^3*f*n^3 + 48*a^2
*b^2*d*e^2*f^2 + 27*a*b^3*c*e^2*f^2*n^2 - 18*a^2*b^2*c*e*f^3*n^2 - 3*a*b^3
*c*e^2*f^2*n^3 + 3*a^2*b^2*c*e*f^3*n^3 - 15*a^2*b^2*d*e^2*f^2*n - 36*a*b^3
*d*e^3*f*n + 46*a^3*b*d*e*f^3*n + 3*a^2*b^2*d*e^2*f^2*n^2 - 60*a*b^3*c*...
```

Reduce [F]

$$\int (a + bx)^{-n} (c + dx) (e + fx)^{-5+n} dx$$

$$= \left(\int \frac{(fx + e)^n}{(bx + a)^n e^5 + 5(bx + a)^n e^4 fx + 10(bx + a)^n e^3 f^2 x^2 + 10(bx + a)^n e^2 f^3 x^3 + 5(bx + a)^n e f^4 x^4 +} \right.$$

$$\left. + \left(\int \frac{(fx + e)^n x}{(bx + a)^n e^5 + 5(bx + a)^n e^4 fx + 10(bx + a)^n e^3 f^2 x^2 + 10(bx + a)^n e^2 f^3 x^3 + 5(bx + a)^n e f^4 x^4} \right) \right)$$

input

```
int((d*x+c)*(f*x+e)^(-5+n)/((b*x+a)^n),x)
```

output

```
int((e + f*x)**n/((a + b*x)**n*e**5 + 5*(a + b*x)**n*e**4*f*x + 10*(a + b*x)**n*e**3*f**2*x**2 + 10*(a + b*x)**n*e**2*f**3*x**3 + 5*(a + b*x)**n*e*f**4*x**4 + (a + b*x)**n*f**5*x**5),x)*c + int(((e + f*x)**n*x)/((a + b*x)**n*e**5 + 5*(a + b*x)**n*e**4*f*x + 10*(a + b*x)**n*e**3*f**2*x**2 + 10*(a + b*x)**n*e**2*f**3*x**3 + 5*(a + b*x)**n*e*f**4*x**4 + (a + b*x)**n*f**5*x**5),x)*d
```

3.264 $\int (a + bx)^{1-n}(c + dx)^{-2+n}(-ad + bc(3 + 2n) + 2bd(1 + n)x) dx$

Optimal result	2291
Mathematica [A] (verified)	2291
Rubi [A] (verified)	2292
Maple [F]	2293
Fricas [F]	2294
Sympy [F(-2)]	2294
Maxima [F]	2294
Giac [F]	2295
Mupad [F(-1)]	2295
Reduce [F]	2296

Optimal result

Integrand size = 42, antiderivative size = 83

$$\int (a + bx)^{1-n}(c + dx)^{-2+n}(-ad + bc(3 + 2n) + 2bd(1 + n)x) dx$$

$$= 2(1 + n)(a + bx)^{2-n}(c + dx)^{-1+n}$$

$$- \frac{(1 - 2n^2)(a + bx)^{2-n}(c + dx)^{-1+n} \operatorname{Hypergeometric2F1}\left(1, 1, n, \frac{b(c+dx)}{bc-ad}\right)}{1 - n}$$

output

```
2*(1+n)*(b*x+a)^(2-n)*(d*x+c)^(-1+n)-(-2*n^2+1)*(b*x+a)^(2-n)*(d*x+c)^(-1+n)*hypergeom([1, 1], [n], b*(d*x+c)/(-a*d+b*c))/(1-n)
```

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.28

$$\int (a + bx)^{1-n}(c + dx)^{-2+n}(-ad + bc(3 + 2n) + 2bd(1 + n)x) dx =$$

$$\frac{(a + bx)^{-n}(c + dx)^n \left(\frac{(a+bx)^2}{c+dx} + \frac{b(bc-ad)(-1+2n^2) \left(\frac{d(a+bx)}{-bc+ad} \right)^n \operatorname{Hypergeometric2F1}\left(-1+n, n, 1+n, \frac{b(c+dx)}{bc-ad}\right)}{d^2 n} \right)}{-1 + n}$$

input

```
Integrate[(a + b*x)^(1 - n)*(c + d*x)^(-2 + n)*(-(a*d) + b*c*(3 + 2*n) + 2
*b*d*(1 + n)*x), x]
```

output

```
-(((c + d*x)^n*((a + b*x)^2/(c + d*x) + (b*(b*c - a*d)*(-1 + 2*n^2)*((d*(a
+ b*x))/(-b*c) + a*d))^n*Hypergeometric2F1[-1 + n, n, 1 + n, (b*(c + d*x
))/b*c - a*d]))/(d^2*n))/((-1 + n)*(a + b*x)^n))
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.46, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$, Rules used = {88, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + bx)^{1-n} (c + dx)^{n-2} (-ad + bc(2n + 3) + 2bd(n + 1)x) dx \\
 & \quad \downarrow 88 \\
 & \frac{b(1 - 2n^2) \int (a + bx)^{1-n} (c + dx)^{n-1} dx}{1 - n} + \frac{(a + bx)^{2-n} (c + dx)^{n-1}}{1 - n} \\
 & \quad \downarrow 80 \\
 & \frac{(a + bx)^{2-n} (c + dx)^{n-1}}{1 - n} - \frac{b(1 - 2n^2) (bc - ad) (a + bx)^{-n} \left(-\frac{d(a+bx)}{bc-ad}\right)^n \int (c + dx)^{n-1} \left(-\frac{bxd}{bc-ad} - \frac{ad}{bc-ad}\right)^{1-n} dx}{d(1 - n)} \\
 & \quad \downarrow 79 \\
 & \frac{(a + bx)^{2-n} (c + dx)^{n-1}}{1 - n} - \frac{b(1 - 2n^2) (bc - ad) (a + bx)^{-n} (c + dx)^n \left(-\frac{d(a+bx)}{bc-ad}\right)^n \text{Hypergeometric2F1}\left(n - 1, n, n + 1, \frac{b(c+dx)}{bc-ad}\right)}{d^2(1 - n)n}
 \end{aligned}$$

input

```
Int[(a + b*x)^(1 - n)*(c + d*x)^(-2 + n)*(-(a*d) + b*c*(3 + 2*n) + 2*b*d*(
1 + n)*x), x]
```

output
$$\frac{((a + bx)^{2-n}(c + dx)^{-1+n})/(1-n) - (b(bc - ad)(1 - 2n^2)) * (-((d(a + bx))/(b*c - a*d)))^n * (c + dx)^n * \text{Hypergeometric2F1}[-1 + n, n, 1 + n, (b(c + dx))/(b*c - a*d)]}{d^2(1-n)n(a + bx)^n}$$

Defintions of rubi rules used

rule 79
$$\text{Int}[(a + b*x)^m * (c + d*x)^n, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{m+1} / (b*(m+1)*(b/(b*c - a*d))^n) * \text{Hypergeometric2F1}[-n, m+1, m+2, (-d)*(a + b*x)/(b*c - a*d)], x] /;$$
 FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

rule 80
$$\text{Int}[(a + b*x)^m * (c + d*x)^n, x_Symbol] \rightarrow \text{Simp}[(c + d*x)^{\text{FracPart}[n]} / ((b/(b*c - a*d))^{\text{IntPart}[n]} * (b*((c + d*x)/(b*c - a*d)))^{\text{FracPart}[n]}) \text{Int}[(a + b*x)^m * \text{Simp}[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n, x], x] /;$$
 FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])

rule 88
$$\text{Int}[(a + b*x)^m * (c + d*x)^n * (e + f*x)^p, x_Symbol] \rightarrow \text{Simp}[-(b*e - a*f) * (c + d*x)^{n+1} * (e + f*x)^{p+1} / (f*(p+1)*(c*f - d*e)), x] - \text{Simp}[(a*d*f*(n+p+2) - b*(d*e*(n+1) + c*f*(p+1))] / (f*(p+1)*(c*f - d*e)) \text{Int}[(c + d*x)^n * (e + f*x)^{\text{Simplify}[p+1]}, x], x] /;$$
 FreeQ[{a, b, c, d, e, f, n, p}, x] && !RationalQ[p] && SumSimplerQ[p, 1]

Maple [F]

$$\int (bx + a)^{-n+1} (xd + c)^{n-2} (-ad + bc(2n + 3) + 2bd(1 + n)x) dx$$

input
$$\text{int}((b*x+a)^{-n+1}*(d*x+c)^{n-2}*(-a*d+b*c*(2*n+3)+2*b*d*(1+n)*x), x)$$

output
$$\text{int}((b*x+a)^{-n+1}*(d*x+c)^{n-2}*(-a*d+b*c*(2*n+3)+2*b*d*(1+n)*x), x)$$

Fricas [F]

$$\int (a + bx)^{1-n} (c + dx)^{-2+n} (-ad + bc(3 + 2n) + 2bd(1 + n)x) dx$$

$$= \int (2bd(n + 1)x + bc(2n + 3) - ad)(bx + a)^{-n+1} (dx + c)^{n-2} dx$$

input `integrate((b*x+a)^(1-n)*(d*x+c)^(-2+n)*(-a*d+b*c*(3+2*n)+2*b*d*(1+n)*x),x,
algorithm="fricas")`

output `integral((2*b*c*n + 3*b*c - a*d + 2*(b*d*n + b*d)*x)*(b*x + a)^(-n + 1)*(d
*x + c)^(n - 2), x)`

Sympy [F(-2)]

Exception generated.

$$\int (a + bx)^{1-n} (c + dx)^{-2+n} (-ad + bc(3 + 2n) + 2bd(1 + n)x) dx$$

$$= \text{Exception raised: HeuristicGCDFailed}$$

input `integrate((b*x+a)**(1-n)*(d*x+c)**(-2+n)*(-a*d+b*c*(3+2*n)+2*b*d*(1+n)*x),
x)`

output `Exception raised: HeuristicGCDFailed >> no luck`

Maxima [F]

$$\int (a + bx)^{1-n} (c + dx)^{-2+n} (-ad + bc(3 + 2n) + 2bd(1 + n)x) dx$$

$$= \int (2bd(n + 1)x + bc(2n + 3) - ad)(bx + a)^{-n+1} (dx + c)^{n-2} dx$$

input `integrate((b*x+a)^(1-n)*(d*x+c)^(-2+n)*(-a*d+b*c*(3+2*n)+2*b*d*(1+n)*x),x,
algorithm="maxima")`

output `integrate((2*b*d*(n + 1)*x + b*c*(2*n + 3) - a*d)*(b*x + a)^(-n + 1)*(d*x
+ c)^(n - 2), x)`

Giac [F]

$$\begin{aligned} & \int (a + bx)^{1-n} (c + dx)^{-2+n} (-ad + bc(3 + 2n) + 2bd(1 + n)x) dx \\ &= \int (2bd(n + 1)x + bc(2n + 3) - ad)(bx + a)^{-n+1} (dx + c)^{n-2} dx \end{aligned}$$

input `integrate((b*x+a)^(1-n)*(d*x+c)^(-2+n)*(-a*d+b*c*(3+2*n)+2*b*d*(1+n)*x),x,
algorithm="giac")`

output `integrate((2*b*d*(n + 1)*x + b*c*(2*n + 3) - a*d)*(b*x + a)^(-n + 1)*(d*x
+ c)^(n - 2), x)`

Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (a + bx)^{1-n} (c + dx)^{-2+n} (-ad + bc(3 + 2n) + 2bd(1 + n)x) dx \\ &= \int (a + bx)^{1-n} (c + dx)^{n-2} (bc(2n + 3) - ad + 2bdx(n + 1)) dx \end{aligned}$$

input `int((a + b*x)^(1 - n)*(c + d*x)^(n - 2)*(b*c*(2*n + 3) - a*d + 2*b*d*x*(n
+ 1)),x)`

output `int((a + b*x)^(1 - n)*(c + d*x)^(n - 2)*(b*c*(2*n + 3) - a*d + 2*b*d*x*(n
+ 1)), x)`

Reduce [F]

$$\begin{aligned}
& \int (a + bx)^{1-n} (c + dx)^{-2+n} (-ad + bc(3 + 2n) + 2bd(1 + n)x) dx \\
&= - \left(\int \frac{(dx + c)^n}{(bx + a)^n c^2 + 2(bx + a)^n cdx + (bx + a)^n d^2 x^2} dx \right) a^2 d \\
&+ 2 \left(\int \frac{(dx + c)^n}{(bx + a)^n c^2 + 2(bx + a)^n cdx + (bx + a)^n d^2 x^2} dx \right) abc n \\
&+ 3 \left(\int \frac{(dx + c)^n}{(bx + a)^n c^2 + 2(bx + a)^n cdx + (bx + a)^n d^2 x^2} dx \right) abc \\
&+ 2 \left(\int \frac{(dx + c)^n x^2}{(bx + a)^n c^2 + 2(bx + a)^n cdx + (bx + a)^n d^2 x^2} dx \right) b^2 dn \\
&+ 2 \left(\int \frac{(dx + c)^n x^2}{(bx + a)^n c^2 + 2(bx + a)^n cdx + (bx + a)^n d^2 x^2} dx \right) b^2 d \\
&+ 2 \left(\int \frac{(dx + c)^n x}{(bx + a)^n c^2 + 2(bx + a)^n cdx + (bx + a)^n d^2 x^2} dx \right) abd n \\
&+ \left(\int \frac{(dx + c)^n x}{(bx + a)^n c^2 + 2(bx + a)^n cdx + (bx + a)^n d^2 x^2} dx \right) abd \\
&+ 2 \left(\int \frac{(dx + c)^n x}{(bx + a)^n c^2 + 2(bx + a)^n cdx + (bx + a)^n d^2 x^2} dx \right) b^2 cn \\
&+ 3 \left(\int \frac{(dx + c)^n x}{(bx + a)^n c^2 + 2(bx + a)^n cdx + (bx + a)^n d^2 x^2} dx \right) b^2 c
\end{aligned}$$

input `int((b*x+a)^(1-n)*(d*x+c)^(-2+n)*(-a*d+b*c*(3+2*n)+2*b*d*(1+n)*x),x)`

output

```

- int((c + d*x)**n/((a + b*x)**n*c**2 + 2*(a + b*x)**n*c*d*x + (a + b*x)*
*n*d**2*x**2),x)*a**2*d + 2*int((c + d*x)**n/((a + b*x)**n*c**2 + 2*(a + b
*x)**n*c*d*x + (a + b*x)**n*d**2*x**2),x)*a*b*c*n + 3*int((c + d*x)**n/((a
+ b*x)**n*c**2 + 2*(a + b*x)**n*c*d*x + (a + b*x)**n*d**2*x**2),x)*a*b*c
+ 2*int(((c + d*x)**n*x**2)/((a + b*x)**n*c**2 + 2*(a + b*x)**n*c*d*x + (a
+ b*x)**n*d**2*x**2),x)*b**2*d*n + 2*int(((c + d*x)**n*x**2)/((a + b*x)**
n*c**2 + 2*(a + b*x)**n*c*d*x + (a + b*x)**n*d**2*x**2),x)*b**2*d + 2*int(
((c + d*x)**n*x)/((a + b*x)**n*c**2 + 2*(a + b*x)**n*c*d*x + (a + b*x)**n*
d**2*x**2),x)*a*b*d*n + int(((c + d*x)**n*x)/((a + b*x)**n*c**2 + 2*(a + b
*x)**n*c*d*x + (a + b*x)**n*d**2*x**2),x)*a*b*d + 2*int(((c + d*x)**n*x)/
(a + b*x)**n*c**2 + 2*(a + b*x)**n*c*d*x + (a + b*x)**n*d**2*x**2),x)*b**2
*c*n + 3*int(((c + d*x)**n*x)/((a + b*x)**n*c**2 + 2*(a + b*x)**n*c*d*x +
(a + b*x)**n*d**2*x**2),x)*b**2*c

```

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions 2298
4.2 Links to plain text integration problems used in this report for each CAS . 2316

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
      ]
    ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "
    ,

```



```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]==Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]==Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]==Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]==Plus || Head[expn]==Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```

```

    Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 5]],
    If [AppellFunctionQ [Head [expn]],
        Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 6]],
        If [Head [expn] === RootSum,
            Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 7]],
            If [Head [expn] === Integrate || Head [expn] === Int,
                Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 8]],
                9]]]]]]]]]]]

```

```
ElementaryFunctionQ [func_] :=
```

```

    MemberQ [{
        Exp, Log,
        Sin, Cos, Tan, Cot, Sec, Csc,
        ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
        Sinh, Cosh, Tanh, Coth, Sech, Csch,
        ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
    }, func]

```

```
SpecialFunctionQ [func_] :=
```

```

    MemberQ [{
        Erf, Erfc, Erfi,
        FresnelS, FresnelC,
        ExpIntegralE, ExpIntegralEi, LogIntegral,
        SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
        Gamma, LogGamma, PolyGamma,
        Zeta, PolyLog, ProductLog,
        EllipticF, EllipticE, EllipticPi
    }, func]

```

```
HypergeometricFunctionQ [func_] :=
```

```

    MemberQ [{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

```

```
AppellFunctionQ [func_] :=
```

```

    MemberQ [{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

#   antiderivative
#   "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
            fi;
        fi;
    else #ExpnType(result) > ExpnType(optimal)
        if debug then
            print("ExpnType(result) > ExpnType(optimal)");
        fi;
        return "C",cat("Result contains higher order function than in optimal. Order ",
                        convert(ExpnType_result,string)," vs. order ",
                        convert(ExpnType_optimal,string),".");
    fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
    9
  end if
end proc
```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```



```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```


4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file