

Computer Algebra Independent Integration Tests

Summer 2024

1-Algebraic-functions/1.1-Binomial/1.1.2-Quadratic-
binomial/38-1.1.2.11

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 5:54pm

Contents

1	Introduction	4
1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Time and leaf size Performance	10
1.4	Performance based on number of rules Rubi used	12
1.5	Performance based on number of steps Rubi used	13
1.6	Solved integrals histogram based on leaf size of result	14
1.7	Solved integrals histogram based on CPU time used	15
1.8	Leaf size vs. CPU time used	16
1.9	list of integrals with no known antiderivative	17
1.10	List of integrals solved by CAS but has no known antiderivative	17
1.11	list of integrals solved by CAS but failed verification	17
1.12	Timing	18
1.13	Verification	18
1.14	Important notes about some of the results	19
1.15	Current tree layout of integration tests	22
1.16	Design of the test system	23
2	detailed summary tables of results	24
2.1	List of integrals sorted by grade for each CAS	25
2.2	Detailed conclusion table per each integral for all CAS systems	29
2.3	Detailed conclusion table specific for Rubi results	42
3	Listing of integrals	44
3.1	$\int \frac{A+Bx^2}{(a+bx^2)^{3/2}(c+dx^2)(e+fx^2)} dx$	46
3.2	$\int \frac{A+Bx^2}{(a+bx^2)^{3/2}(c+dx^2)(e+fx^2)^2} dx$	53
3.3	$\int \frac{A+Bx^2}{(a+bx^2)^{3/2}(c+dx^2)(e+fx^2)^3} dx$	61
3.4	$\int \frac{A+Bx^2}{(a+bx^2)^{5/2}(c+dx^2)(e+fx^2)} dx$	70
3.5	$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)(e+fx^2)^2} dx$	77

3.6	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}(c+dx^2)(e+fx^2)^2} dx$	84
3.7	$\int \frac{A+Bx^2+Cx^4}{(a+bx^2)^{3/2}(c+dx^2)(e+fx^2)} dx$	92
3.8	$\int \frac{\sqrt{a+bx^2}(c+dx^2)^{3/2}(A+Bx^2+Cx^4)}{e+fx^2} dx$	99
3.9	$\int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{e+fx^2} dx$	107
3.10	$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{\sqrt{c+dx^2}(e+fx^2)} dx$	116
3.11	$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{3/2}(e+fx^2)} dx$	124
3.12	$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{5/2}(e+fx^2)} dx$	132
3.13	$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{7/2}(e+fx^2)} dx$	140
3.14	$\int \frac{A+Bx^2}{(a+bx^2)\sqrt{c+dx^2}\sqrt{e+fx^2}} dx$	148
3.15	$\int \frac{A+Bx^2}{(a+bx^2)\sqrt{c-dx^2}\sqrt{e+fx^2}} dx$	154
3.16	$\int \frac{A+Bx^2}{(a+bx^2)\sqrt{c+dx^2}\sqrt{e-fx^2}} dx$	160
3.17	$\int \frac{A+Bx^2}{(a+bx^2)\sqrt{c-dx^2}\sqrt{e-fx^2}} dx$	166
3.18	$\int \frac{A+Bx^2+Cx^4}{(a+bx^2)\sqrt{c+dx^2}\sqrt{e+fx^2}} dx$	172
3.19	$\int \frac{A+Bx^2+Cx^4}{(a+bx^2)\sqrt{c-dx^2}\sqrt{e+fx^2}} dx$	178
3.20	$\int \frac{A+Bx^2+Cx^4}{(a+bx^2)\sqrt{c+dx^2}\sqrt{e-fx^2}} dx$	184
3.21	$\int \frac{A+Bx^2+Cx^4}{(a+bx^2)\sqrt{c-dx^2}\sqrt{e-fx^2}} dx$	190
3.22	$\int \frac{A+Bx^2}{\sqrt{a+bx^2}(c+dx^2)^{3/2}(e+fx^2)} dx$	196
3.23	$\int \frac{A+Bx^2}{\sqrt{a+bx^2}(c-dx^2)^{3/2}(e+fx^2)} dx$	202
3.24	$\int \frac{A+Bx^2}{\sqrt{a-bx^2}(c+dx^2)^{3/2}(e+fx^2)} dx$	209
3.25	$\int \frac{A+Bx^2}{\sqrt{a-bx^2}(c-dx^2)^{3/2}(e+fx^2)} dx$	216
3.26	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}(c+dx^2)^{3/2}(e+fx^2)} dx$	223
3.27	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}(c-dx^2)^{3/2}(e+fx^2)} dx$	231
3.28	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}(c+dx^2)^{3/2}(e+fx^2)} dx$	239
3.29	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}(c-dx^2)^{3/2}(e+fx^2)} dx$	247
3.30	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e+fx^2)^2} dx$	255
3.31	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e+fx^2)^3} dx$	264
3.32	$\int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{(e+fx^2)^{3/2}} dx$	272
3.33	$\int \frac{A+Bx^2}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e+fx^2)^{3/2}} dx$	278
3.34	$\int \frac{A+Bx^2}{\sqrt{a-bx^2}\sqrt{c+dx^2}(e+fx^2)^{3/2}} dx$	283
3.35	$\int \frac{A+Bx^2}{\sqrt{a+bx^2}\sqrt{c-dx^2}(e+fx^2)^{3/2}} dx$	288

3.36	$\int \frac{A+Bx^2}{\sqrt{a-bx^2}\sqrt{c-dx^2}(e+fx^2)^{3/2}} dx$	293
3.37	$\int \frac{A+Bx^2}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e-fx^2)^{3/2}} dx$	298
3.38	$\int \frac{A+Bx^2}{\sqrt{a-bx^2}\sqrt{c+dx^2}(e-fx^2)^{3/2}} dx$	303
3.39	$\int \frac{A+Bx^2}{\sqrt{a+bx^2}\sqrt{c-dx^2}(e-fx^2)^{3/2}} dx$	308
3.40	$\int \frac{A+Bx^2}{\sqrt{a-bx^2}\sqrt{c-dx^2}(e-fx^2)^{3/2}} dx$	313
3.41	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e+fx^2)^{3/2}} dx$	318
3.42	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}\sqrt{c+dx^2}(e+fx^2)^{3/2}} dx$	324
3.43	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c-dx^2}(e+fx^2)^{3/2}} dx$	330
3.44	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}\sqrt{c-dx^2}(e+fx^2)^{3/2}} dx$	336
3.45	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e-fx^2)^{3/2}} dx$	342
3.46	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}\sqrt{c+dx^2}(e-fx^2)^{3/2}} dx$	348
3.47	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c-dx^2}(e-fx^2)^{3/2}} dx$	354
3.48	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}\sqrt{c-dx^2}(e-fx^2)^{3/2}} dx$	360
4	Appendix	366
4.1	Listing of Grading functions	366
4.2	Links to plain text integration problems used in this report for each CAS384	

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Time and leaf size Performance	10
1.4	Performance based on number of rules Rubi used	12
1.5	Performance based on number of steps Rubi used	13
1.6	Solved integrals histogram based on leaf size of result	14
1.7	Solved integrals histogram based on CPU time used	15
1.8	Leaf size vs. CPU time used	16
1.9	list of integrals with no known antiderivative	17
1.10	List of integrals solved by CAS but has no known antiderivative	17
1.11	list of integrals solved by CAS but failed verification	17
1.12	Timing	18
1.13	Verification	18
1.14	Important notes about some of the results	19
1.15	Current tree layout of integration tests	22
1.16	Design of the test system	23

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [48]. This is test number [38].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Mathematica	64.58 (31)	35.42 (17)
Maple	64.58 (31)	35.42 (17)
Rubi	62.50 (30)	37.50 (18)
Reduce	14.58 (7)	85.42 (41)
Giac	12.50 (6)	87.50 (42)
Fricas	0.00 (0)	100.00 (48)
Mupad	0.00 (0)	100.00 (48)
Maxima	0.00 (0)	100.00 (48)
Sympy	0.00 (0)	100.00 (48)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	54.167	8.333	0.000	37.500
Maple	39.583	25.000	0.000	35.417
Mathematica	10.417	0.000	54.167	35.417
Giac	4.167	8.333	0.000	87.500
Fricas	0.000	0.000	0.000	100.000
Mupad	0.000	0.000	0.000	100.000
Maxima	0.000	0.000	0.000	100.000
Reduce	0.000	14.583	0.000	85.417
Sympy	0.000	0.000	0.000	100.000

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Mathematica	17	100.00	0.00	0.00
Maple	17	100.00	0.00	0.00
Rubi	18	100.00	0.00	0.00
Reduce	41	100.00	0.00	0.00
Giac	42	97.62	0.00	2.38
Fricas	48	27.08	72.92	0.00
Mupad	48	0.00	100.00	0.00
Maxima	48	100.00	0.00	0.00
Sympy	48	79.17	20.83	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Rubi	1.51
Reduce	1.86
Giac	2.40
Maple	6.81
Mathematica	9.41
Sympy	-nan(ind)
Maxima	-nan(ind)
Mupad	-nan(ind)
Fricas	-nan(ind)

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Rubi	595.67	1.43	554.50	1.27
Mathematica	799.97	1.45	337.00	0.85
Giac	1104.33	3.06	731.50	2.32
Maple	1419.81	2.35	536.00	1.20
Reduce	5008.86	14.51	4615.00	14.89
Sympy	-nan(ind)	-nan(ind)	nan	nan
Maxima	-nan(ind)	-nan(ind)	nan	nan
Mupad	-nan(ind)	-nan(ind)	nan	nan
Fricas	-nan(ind)	-nan(ind)	nan	nan

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {2, 3}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer. the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	25
2.2	Detailed conclusion table per each integral for all CAS systems	29
2.3	Detailed conclusion table specific for Rubi results	42

2.1 List of integrals sorted by grade for each CAS

Rubi	25
Mma	25
Maple	26
Fricas	26
Maxima	26
Giac	27
Mupad	27
Sympy	27
Reduce	28

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 26, 30 }
}

B grade { 25, 27, 28, 29 }

C grade { }

F normal fail { 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 4, 5, 6, 7 }

B grade { }

C grade { 2, 3, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31 }

F normal fail { 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 14, 15, 16, 17, 18, 19, 20, 21, 22 }

B grade { 11, 12, 13, 23, 24, 25, 26, 27, 28, 29, 30, 31 }

C grade { }

F normal fail { 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48 }

F(-1) timeout fail { }

F(-2) exception fail { }

Fricas

A grade { }

B grade { }

C grade { }

F normal fail { 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 46, 47, 48 }

F(-1) timeout fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 43, 44, 45 }

F(-2) exception fail { }

Maxima

A grade { }

B grade { }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48 }

F(-1) timeout fail { }

F(-2) exception fail { }

Giac

A grade { 1, 7 }

B grade { 2, 3, 4, 6 }

C grade { }

F normal fail { 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48 }

F(-1) timedout fail { }

F(-2) exception fail { 5 }

Mupad

A grade { }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48 }

F(-2) exception fail { }

Sympy

A grade { }

B grade { }

C grade { }

F normal fail { 8, 9, 10, 11, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48 }

F(-1) timedout fail { 1, 2, 3, 4, 5, 6, 7, 12, 13, 31 }

F(-2) exception fail { }

Reduce

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7 }

C grade { }

F normal fail { 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F(-1)	F(-1)	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	183	229	216	187	0	0	0	277	508	0
N.S.	1	1.25	1.18	1.02	0.00	0.00	0.00	1.51	2.78	0.00
time (sec)	N/A	1.061	1.991	1.461	0.000	0.000	0.000	0.705	0.482	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F(-1)	F(-1)	B	B	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	356	431	1481	421	0	0	0	785	3033	0
N.S.	1	1.21	4.16	1.18	0.00	0.00	0.00	2.21	8.52	0.00
time (sec)	N/A	1.260	18.033	1.716	0.000	0.000	0.000	2.140	1.305	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F(-1)	F(-1)	B	B	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	706	736	2913	953	0	0	0	2528	12786	0
N.S.	1	1.04	4.13	1.35	0.00	0.00	0.00	3.58	18.11	0.00
time (sec)	N/A	1.810	20.692	3.352	0.000	0.000	0.000	8.728	5.652	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F(-1)	F(-1)	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	288	352	337	291	0	0	0	2061	2231	0
N.S.	1	1.22	1.17	1.01	0.00	0.00	0.00	7.16	7.75	0.00
time (sec)	N/A	1.300	3.577	1.473	0.000	0.000	0.000	1.214	1.322	0.000

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F(-1)	F(-1)	F(-2)	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	310	422	343	348	0	0	0	0	4615	0
N.S.	1	1.36	1.11	1.12	0.00	0.00	0.00	0.00	14.89	0.00
time (sec)	N/A	1.170	2.295	1.565	0.000	0.000	0.000	0.000	1.518	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F(-1)	F(-1)	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	278	316	295	302	0	0	0	678	6592	0
N.S.	1	1.14	1.06	1.09	0.00	0.00	0.00	2.44	23.71	0.00
time (sec)	N/A	1.125	2.465	1.521	0.000	0.000	0.000	0.859	1.388	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F(-1)	F(-1)	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	205	288	238	208	0	0	0	297	5297	0
N.S.	1	1.40	1.16	1.01	0.00	0.00	0.00	1.45	25.84	0.00
time (sec)	N/A	1.049	1.549	1.361	0.000	0.000	0.000	0.747	1.376	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	1070	1321	6720	1360	0	0	0	0	42	0
N.S.	1	1.23	6.28	1.27	0.00	0.00	0.00	0.00	0.04	0.00
time (sec)	N/A	2.450	13.161	19.544	0.000	0.000	0.000	0.000	200.015	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	666	1029	541	802	0	0	0	0	42	0
N.S.	1	1.55	0.81	1.20	0.00	0.00	0.00	0.00	0.06	0.00
time (sec)	N/A	1.766	7.823	9.103	0.000	0.000	0.000	0.000	200.020	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	443	623	383	506	0	0	0	0	867	0
N.S.	1	1.41	0.86	1.14	0.00	0.00	0.00	0.00	1.96	0.00
time (sec)	N/A	1.265	5.507	8.606	0.000	0.000	0.000	0.000	7.958	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	481	550	368	1300	0	0	0	0	0	0
N.S.	1	1.14	0.77	2.70	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.267	6.687	6.558	0.000	0.000	0.000	0.000	39.043	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	F(-1)	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	637	979	3341	6013	0	0	0	0	0	0
N.S.	1	1.54	5.24	9.44	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.930	10.948	8.853	0.000	0.000	0.000	0.000	93.766	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	F(-1)	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	1093	1396	1031	13344	0	0	0	0	0	0
N.S.	1	1.28	0.94	12.21	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.745	12.758	11.495	0.000	0.000	0.000	0.000	152.510	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	226	203	143	191	0	0	0	0	43	0
N.S.	1	0.90	0.63	0.85	0.00	0.00	0.00	0.00	0.19	0.00
time (sec)	N/A	0.990	8.576	5.780	0.000	0.000	0.000	0.000	0.163	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	203	203	151	192	0	0	0	0	46	0
N.S.	1	1.00	0.74	0.95	0.00	0.00	0.00	0.00	0.23	0.00
time (sec)	N/A	1.037	8.552	6.018	0.000	0.000	0.000	0.000	0.164	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	203	203	147	192	0	0	0	0	46	0
N.S.	1	1.00	0.72	0.95	0.00	0.00	0.00	0.00	0.23	0.00
time (sec)	N/A	1.049	8.566	5.906	0.000	0.000	0.000	0.000	0.164	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	205	205	151	191	0	0	0	0	47	0
N.S.	1	1.00	0.74	0.93	0.00	0.00	0.00	0.00	0.23	0.00
time (sec)	N/A	1.110	8.520	5.935	0.000	0.000	0.000	0.000	0.157	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	371	342	200	328	0	0	0	0	238	0
N.S.	1	0.92	0.54	0.88	0.00	0.00	0.00	0.00	0.64	0.00
time (sec)	N/A	1.103	7.309	5.918	0.000	0.000	0.000	0.000	1.800	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	323	409	210	328	0	0	0	0	253	0
N.S.	1	1.27	0.65	1.02	0.00	0.00	0.00	0.00	0.78	0.00
time (sec)	N/A	1.252	7.295	5.763	0.000	0.000	0.000	0.000	1.786	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	323	409	206	328	0	0	0	0	253	0
N.S.	1	1.27	0.64	1.02	0.00	0.00	0.00	0.00	0.78	0.00
time (sec)	N/A	1.237	7.332	5.839	0.000	0.000	0.000	0.000	1.758	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	326	413	210	323	0	0	0	0	256	0
N.S.	1	1.27	0.64	0.99	0.00	0.00	0.00	0.00	0.79	0.00
time (sec)	N/A	1.302	7.342	5.780	0.000	0.000	0.000	0.000	1.750	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	377	572	240	537	0	0	0	0	67	0
N.S.	1	1.52	0.64	1.42	0.00	0.00	0.00	0.00	0.18	0.00
time (sec)	N/A	1.471	11.123	8.571	0.000	0.000	0.000	0.000	8.398	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	390	750	240	752	0	0	0	0	68	0
N.S.	1	1.92	0.62	1.93	0.00	0.00	0.00	0.00	0.17	0.00
time (sec)	N/A	1.845	11.113	8.437	0.000	0.000	0.000	0.000	8.088	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	294	559	250	536	0	0	0	0	272	0
N.S.	1	1.90	0.85	1.82	0.00	0.00	0.00	0.00	0.93	0.00
time (sec)	N/A	1.523	11.305	8.680	0.000	0.000	0.000	0.000	11.148	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	C	B	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	294	673	249	762	0	0	0	0	274	0
N.S.	1	2.29	0.85	2.59	0.00	0.00	0.00	0.00	0.93	0.00
time (sec)	N/A	1.789	11.185	8.652	0.000	0.000	0.000	0.000	10.245	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	415	805	352	1042	0	0	0	0	394	0
N.S.	1	1.94	0.85	2.51	0.00	0.00	0.00	0.00	0.95	0.00
time (sec)	N/A	1.720	10.886	8.391	0.000	0.000	0.000	0.000	18.991	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	C	B	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	426	956	350	1127	0	0	0	0	397	0
N.S.	1	2.24	0.82	2.65	0.00	0.00	0.00	0.00	0.93	0.00
time (sec)	N/A	2.057	10.946	8.454	0.000	0.000	0.000	0.000	16.913	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	C	B	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	416	844	362	1043	0	0	0	0	409	0
N.S.	1	2.03	0.87	2.51	0.00	0.00	0.00	0.00	0.98	0.00
time (sec)	N/A	1.871	13.345	8.464	0.000	0.000	0.000	0.000	17.830	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	C	B	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	418	875	361	1134	0	0	0	0	412	0
N.S.	1	2.09	0.86	2.71	0.00	0.00	0.00	0.00	0.99	0.00
time (sec)	N/A	2.013	11.109	8.434	0.000	0.000	0.000	0.000	15.644	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	545	777	2038	2573	0	0	0	0	394	0
N.S.	1	1.43	3.74	4.72	0.00	0.00	0.00	0.00	0.72	0.00
time (sec)	N/A	1.714	14.672	8.472	0.000	0.000	0.000	0.000	19.078	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	C	B	F	F(-1)	F(-1)	F	F	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	967	0	732	6400	0	0	0	0	550	0
N.S.	1	0.00	0.76	6.62	0.00	0.00	0.00	0.00	0.57	0.00
time (sec)	N/A	0.000	15.175	10.911	0.000	0.000	0.000	0.000	60.354	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	F	F	F	F(-1)	F	F	F	F(-1)
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	813	0	0	0	0	0	0	0	42	0
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.05	0.00
time (sec)	N/A	0.000	0.000	0.000	0.000	0.000	0.000	0.000	200.024	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	F	F	F	F	F	F	F	F(-1)
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	332	0	0	0	0	0	0	0	75	0
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.23	0.00
time (sec)	N/A	0.000	0.000	0.000	0.000	0.000	0.000	0.000	15.541	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	F	F	F	F	F	F	F	F(-1)
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	332	0	0	0	0	0	0	0	288	0
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.87	0.00
time (sec)	N/A	0.000	0.000	0.000	0.000	0.000	0.000	0.000	19.616	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	F	F	F	F	F	F	F	F(-1)
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	333	0	0	0	0	0	0	0	78	0
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.23	0.00
time (sec)	N/A	0.000	0.000	0.000	0.000	0.000	0.000	0.000	14.651	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	F	F	F	F	F	F	F	F(-1)
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	329	0	0	0	0	0	0	0	290	0
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.88	0.00
time (sec)	N/A	0.000	0.000	0.000	0.000	0.000	0.000	0.000	18.640	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	F	F	F	F	F	F	F	F(-1)
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	319	0	0	0	0	0	0	0	76	0
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.24	0.00
time (sec)	N/A	0.000	0.000	0.000	0.000	0.000	0.000	0.000	14.175	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	F	F	F	F	F	F	F	F(-1)
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	339	0	0	0	0	0	0	0	290	0
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.86	0.00
time (sec)	N/A	0.000	0.000	0.000	0.000	0.000	0.000	0.000	18.230	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	F	F	F	F	F	F	F	F(-1)
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	340	0	0	0	0	0	0	0	79	0
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.23	0.00
time (sec)	N/A	0.000	0.000	0.000	0.000	0.000	0.000	0.000	14.111	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	F	F	F	F	F	F	F	F(-1)
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	344	0	0	0	0	0	0	0	292	0
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.85	0.00
time (sec)	N/A	0.000	0.000	0.000	0.000	0.000	0.000	0.000	18.085	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	F	F	F	F	F	F	F	F(-1)
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	527	0	0	0	0	0	0	0	418	0
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.79	0.00
time (sec)	N/A	0.000	0.000	0.000	0.000	0.000	0.000	0.000	41.264	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	F	F	F	F	F	F	F	F(-1)
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	525	0	0	0	0	0	0	0	433	0
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.82	0.00
time (sec)	N/A	0.000	0.000	0.000	0.000	0.000	0.000	0.000	39.285	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	F	F	F	F(-1)	F	F	F	F(-1)
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	533	0	0	0	0	0	0	0	433	0
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.81	0.00
time (sec)	N/A	0.000	0.000	0.000	0.000	0.000	0.000	0.000	38.337	0.000

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	F	F	F	F(-1)	F	F	F	F(-1)
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	521	0	0	0	0	0	0	0	436	0
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.84	0.00
time (sec)	N/A	0.000	0.000	0.000	0.000	0.000	0.000	0.000	35.919	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	F	F	F	F(-1)	F	F	F	F(-1)
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	496	0	0	0	0	0	0	0	421	0
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.85	0.00
time (sec)	N/A	0.000	0.000	0.000	0.000	0.000	0.000	0.000	37.486	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	F	F	F	F	F	F	F	F(-1)
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	538	0	0	0	0	0	0	0	436	0
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.81	0.00
time (sec)	N/A	0.000	0.000	0.000	0.000	0.000	0.000	0.000	35.527	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	F	F	F	F	F	F	F	F(-1)
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	545	0	0	0	0	0	0	0	436	0
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.80	0.00
time (sec)	N/A	0.000	0.000	0.000	0.000	0.000	0.000	0.000	34.581	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	F	F	F	F	F	F	F	F(-1)
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	546	0	0	0	0	0	0	0	439	0
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.80	0.00
time (sec)	N/A	0.000	0.000	0.000	0.000	0.000	0.000	0.000	34.558	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [1] had the largest ratio of [.540541000000000008e-1]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	2	2	1.25	37	0.054
2	A	2	2	1.21	37	0.054
3	A	2	2	1.04	37	0.054
4	A	2	2	1.22	37	0.054
5	A	2	2	1.36	42	0.048
6	A	2	2	1.14	42	0.048
7	A	2	2	1.40	42	0.048
8	A	2	2	1.23	44	0.045
9	A	2	2	1.55	44	0.045
10	A	2	2	1.41	44	0.045
11	A	2	2	1.14	44	0.045
12	A	2	2	1.54	44	0.045
13	A	2	2	1.28	44	0.045
14	A	2	2	0.90	39	0.051
15	A	2	2	1.00	40	0.050
16	A	2	2	1.00	40	0.050
17	A	2	2	1.00	41	0.049
18	A	2	2	0.92	44	0.045
19	A	2	2	1.27	45	0.044
20	A	2	2	1.27	45	0.044
21	A	2	2	1.27	46	0.043

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	2	2	1.52	39	0.051
23	A	2	2	1.92	40	0.050
24	A	2	2	1.90	40	0.050
25	B	2	2	2.29	41	0.049
26	A	2	2	1.94	44	0.045
27	B	2	2	2.24	45	0.044
28	B	2	2	2.03	45	0.044
29	B	2	2	2.09	46	0.043
30	A	2	2	1.43	44	0.045
31	F	0	0	N/A	0.000	N/A
32	F	0	0	N/A	0.000	N/A
33	F	0	0	N/A	0.000	N/A
34	F	0	0	N/A	0.000	N/A
35	F	0	0	N/A	0.000	N/A
36	F	0	0	N/A	0.000	N/A
37	F	0	0	N/A	0.000	N/A
38	F	0	0	N/A	0.000	N/A
39	F	0	0	N/A	0.000	N/A
40	F	0	0	N/A	0.000	N/A
41	F	0	0	N/A	0.000	N/A
42	F	0	0	N/A	0.000	N/A
43	F	0	0	N/A	0.000	N/A
44	F	0	0	N/A	0.000	N/A
45	F	0	0	N/A	0.000	N/A
46	F	0	0	N/A	0.000	N/A
47	F	0	0	N/A	0.000	N/A
48	F	0	0	N/A	0.000	N/A

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \frac{A+Bx^2}{(a+bx^2)^{3/2}(c+dx^2)(e+fx^2)} dx$	46
3.2	$\int \frac{A+Bx^2}{(a+bx^2)^{3/2}(c+dx^2)(e+fx^2)^2} dx$	53
3.3	$\int \frac{A+Bx^2}{(a+bx^2)^{3/2}(c+dx^2)(e+fx^2)^3} dx$	61
3.4	$\int \frac{A+Bx^2}{(a+bx^2)^{5/2}(c+dx^2)(e+fx^2)} dx$	70
3.5	$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)(e+fx^2)^2} dx$	77
3.6	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}(c+dx^2)(e+fx^2)^2} dx$	84
3.7	$\int \frac{A+Bx^2+Cx^4}{(a+bx^2)^{3/2}(c+dx^2)(e+fx^2)} dx$	92
3.8	$\int \frac{\sqrt{a+bx^2}(c+dx^2)^{3/2}(A+Bx^2+Cx^4)}{e+fx^2} dx$	99
3.9	$\int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{e+fx^2} dx$	107
3.10	$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{\sqrt{c+dx^2}(e+fx^2)} dx$	116
3.11	$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{3/2}(e+fx^2)} dx$	124
3.12	$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{5/2}(e+fx^2)} dx$	132
3.13	$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{7/2}(e+fx^2)} dx$	140
3.14	$\int \frac{A+Bx^2}{(a+bx^2)\sqrt{c+dx^2}\sqrt{e+fx^2}} dx$	148
3.15	$\int \frac{A+Bx^2}{(a+bx^2)\sqrt{c-dx^2}\sqrt{e+fx^2}} dx$	154
3.16	$\int \frac{A+Bx^2}{(a+bx^2)\sqrt{c+dx^2}\sqrt{e-fx^2}} dx$	160
3.17	$\int \frac{A+Bx^2}{(a+bx^2)\sqrt{c-dx^2}\sqrt{e-fx^2}} dx$	166
3.18	$\int \frac{A+Bx^2+Cx^4}{(a+bx^2)\sqrt{c+dx^2}\sqrt{e+fx^2}} dx$	172
3.19	$\int \frac{A+Bx^2+Cx^4}{(a+bx^2)\sqrt{c-dx^2}\sqrt{e+fx^2}} dx$	178
3.20	$\int \frac{A+Bx^2+Cx^4}{(a+bx^2)\sqrt{c+dx^2}\sqrt{e-fx^2}} dx$	184

3.21	$\int \frac{A+Bx^2+Cx^4}{(a+bx^2)\sqrt{c-dx^2}\sqrt{e-fx^2}} dx$	190
3.22	$\int \frac{A+Bx^2}{\sqrt{a+bx^2}(c+dx^2)^{3/2}(e+fx^2)} dx$	196
3.23	$\int \frac{A+Bx^2}{\sqrt{a+bx^2}(c-dx^2)^{3/2}(e+fx^2)} dx$	202
3.24	$\int \frac{A+Bx^2}{\sqrt{a-bx^2}(c+dx^2)^{3/2}(e+fx^2)} dx$	209
3.25	$\int \frac{A+Bx^2}{\sqrt{a-bx^2}(c-dx^2)^{3/2}(e+fx^2)} dx$	216
3.26	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}(c+dx^2)^{3/2}(e+fx^2)} dx$	223
3.27	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}(c-dx^2)^{3/2}(e+fx^2)} dx$	231
3.28	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}(c+dx^2)^{3/2}(e+fx^2)} dx$	239
3.29	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}(c-dx^2)^{3/2}(e+fx^2)} dx$	247
3.30	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e+fx^2)^2} dx$	255
3.31	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e+fx^2)^3} dx$	264
3.32	$\int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{(e+fx^2)^{3/2}} dx$	272
3.33	$\int \frac{A+Bx^2}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e+fx^2)^{3/2}} dx$	278
3.34	$\int \frac{A+Bx^2}{\sqrt{a-bx^2}\sqrt{c+dx^2}(e+fx^2)^{3/2}} dx$	283
3.35	$\int \frac{A+Bx^2}{\sqrt{a+bx^2}\sqrt{c-dx^2}(e+fx^2)^{3/2}} dx$	288
3.36	$\int \frac{A+Bx^2}{\sqrt{a-bx^2}\sqrt{c-dx^2}(e+fx^2)^{3/2}} dx$	293
3.37	$\int \frac{A+Bx^2}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e-fx^2)^{3/2}} dx$	298
3.38	$\int \frac{A+Bx^2}{\sqrt{a-bx^2}\sqrt{c+dx^2}(e-fx^2)^{3/2}} dx$	303
3.39	$\int \frac{A+Bx^2}{\sqrt{a+bx^2}\sqrt{c-dx^2}(e-fx^2)^{3/2}} dx$	308
3.40	$\int \frac{A+Bx^2}{\sqrt{a-bx^2}\sqrt{c-dx^2}(e-fx^2)^{3/2}} dx$	313
3.41	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e+fx^2)^{3/2}} dx$	318
3.42	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}\sqrt{c+dx^2}(e+fx^2)^{3/2}} dx$	324
3.43	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c-dx^2}(e+fx^2)^{3/2}} dx$	330
3.44	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}\sqrt{c-dx^2}(e+fx^2)^{3/2}} dx$	336
3.45	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e-fx^2)^{3/2}} dx$	342
3.46	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}\sqrt{c+dx^2}(e-fx^2)^{3/2}} dx$	348
3.47	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c-dx^2}(e-fx^2)^{3/2}} dx$	354
3.48	$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}\sqrt{c-dx^2}(e-fx^2)^{3/2}} dx$	360

$$3.1 \quad \int \frac{A+Bx^2}{(a+bx^2)^{3/2}(c+dx^2)(e+fx^2)} dx$$

Optimal result	46
Mathematica [A] (verified)	47
Rubi [A] (verified)	47
Maple [A] (verified)	48
Fricas [F(-1)]	49
Sympy [F(-1)]	49
Maxima [F]	50
Giac [A] (verification not implemented)	50
Mupad [F(-1)]	51
Reduce [B] (verification not implemented)	51

Optimal result

Integrand size = 37, antiderivative size = 183

$$\int \frac{A+Bx^2}{(a+bx^2)^{3/2}(c+dx^2)(e+fx^2)} dx = \frac{b(Ab-aB)x}{a(bc-ad)(be-af)\sqrt{a+bx^2}} + \frac{d(Bc-Ad)\operatorname{arctanh}\left(\frac{\sqrt{bc-ad}x}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}(bc-ad)^{3/2}(de-cf)} - \frac{f(Be-Af)\operatorname{arctanh}\left(\frac{\sqrt{be-af}x}{\sqrt{e}\sqrt{a+bx^2}}\right)}{\sqrt{e}(be-af)^{3/2}(de-cf)}$$

output

```
b*(A*b-B*a)*x/a/(-a*d+b*c)/(-a*f+b*e)/(b*x^2+a)^(1/2)+d*(-A*d+B*c)*arctanh
((-a*d+b*c)^(1/2)*x/c^(1/2)/(b*x^2+a)^(1/2))/c^(1/2)/(-a*d+b*c)^(3/2)/(-c*
f+d*e)-f*(-A*f+B*e)*arctanh((-a*f+b*e)^(1/2)*x/e^(1/2)/(b*x^2+a)^(1/2))/e^
(1/2)/(-a*f+b*e)^(3/2)/(-c*f+d*e)
```

Mathematica [A] (verified)

Time = 1.99 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.18

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)} dx = \frac{b(Ab - aB)x}{a(-bc + ad)(-be + af)\sqrt{a + bx^2}}$$

$$+ \frac{d(Bc - Ad) \arctan\left(\frac{-dx\sqrt{a+bx^2} + \sqrt{b}(c+dx^2)}{\sqrt{c}\sqrt{-bc+ad}}\right)}{\sqrt{c}(-bc + ad)^{3/2}(de - cf)}$$

$$+ \frac{f(Be - Af) \arctan\left(\frac{-fx\sqrt{a+bx^2} + \sqrt{b}(e+fx^2)}{\sqrt{e}\sqrt{-be+af}}\right)}{\sqrt{e}(-be + af)^{3/2}(-de + cf)}$$

input

```
Integrate[(A + B*x^2)/((a + b*x^2)^(3/2)*(c + d*x^2)*(e + f*x^2)),x]
```

output

```
(b*(A*b - a*B)*x)/(a*(-(b*c) + a*d)*(-(b*e) + a*f)*Sqrt[a + b*x^2]) + (d*(B*c - A*d)*ArcTan[(-(d*x*Sqrt[a + b*x^2]) + Sqrt[b]*(c + d*x^2))/(Sqrt[c]*Sqrt[-(b*c) + a*d])]/(Sqrt[c]*(-(b*c) + a*d)^(3/2)*(d*e - c*f)) + (f*(B*e - A*f)*ArcTan[(-(f*x*Sqrt[a + b*x^2]) + Sqrt[b]*(e + f*x^2))/(Sqrt[e]*Sqrt[-(b*e) + a*f])]/(Sqrt[e]*(-(b*e) + a*f)^(3/2)*(-(d*e) + c*f))
```

Rubi [A] (verified)

Time = 1.06 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.25, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.054$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)} dx$$

$$\downarrow \text{7276}$$

$$\int \left(\frac{Be - Af}{(a + bx^2)^{3/2} (e + fx^2) (de - cf)} - \frac{Bc - Ad}{(a + bx^2)^{3/2} (c + dx^2) (de - cf)} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{d(Bc - Ad)\operatorname{arctanh}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}(bc - ad)^{3/2}(de - cf)} - \frac{f(Be - Af)\operatorname{arctanh}\left(\frac{x\sqrt{be-af}}{\sqrt{e}\sqrt{a+bx^2}}\right)}{\sqrt{e}(be - af)^{3/2}(de - cf)} - \frac{bx(Bc - Ad)}{a\sqrt{a + bx^2}(bc - ad)(de - cf)} + \frac{bx(Be - Af)}{a\sqrt{a + bx^2}(be - af)(de - cf)}$$

input `Int[(A + B*x^2)/((a + b*x^2)^(3/2)*(c + d*x^2)*(e + f*x^2)),x]`

output `-((b*(B*c - A*d)*x)/(a*(b*c - a*d)*(d*e - c*f)*Sqrt[a + b*x^2])) + (b*(B*e - A*f)*x)/(a*(b*e - a*f)*(d*e - c*f)*Sqrt[a + b*x^2]) + (d*(B*c - A*d)*ArcTanh[(Sqrt[b*c - a*d]*x)/(Sqrt[c]*Sqrt[a + b*x^2])])/(Sqrt[c]*(b*c - a*d)^(3/2)*(d*e - c*f)) - (f*(B*e - A*f)*ArcTanh[(Sqrt[b*e - a*f]*x)/(Sqrt[e]*Sqrt[a + b*x^2])])/(Sqrt[e]*(b*e - a*f)^(3/2)*(d*e - c*f))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 1.46 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.02

method	result	size
pseudoelliptic	$\frac{(Ad - Bc)ad \arctan\left(\frac{c\sqrt{bx^2+a}}{x\sqrt{(ad-bc)c}}\right)}{(cf-de)(ad-bc)\sqrt{(ad-bc)c}} + \frac{b(Ab - Ba)x}{(af-be)(ad-bc)\sqrt{bx^2+a}} - \frac{(Af - Be)af \arctan\left(\frac{e\sqrt{bx^2+a}}{x\sqrt{(af-be)e}}\right)}{(cf-de)(af-be)\sqrt{(af-be)e}}$	187
default	Expression too large to display	1640

input `int((B*x^2+A)/(b*x^2+a)^(3/2)/(d*x^2+c)/(f*x^2+e),x,method=_RETURNVERBOSE)`

output

```
((A*d-B*c)*a*d/(c*f-d*e)/(a*d-b*c)/((a*d-b*c)*c)^(1/2)*arctan(c*(b*x^2+a)^(1/2)/x/((a*d-b*c)*c)^(1/2))+b*(A*b-B*a)/(a*f-b*e)/(a*d-b*c)*x/(b*x^2+a)^(1/2)-(A*f-B*e)*a*f/(c*f-d*e)/(a*f-b*e)/((a*f-b*e)*e)^(1/2)*arctan(e*(b*x^2+a)^(1/2)/x/((a*f-b*e)*e)^(1/2)))/a
```

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)} dx = \text{Timed out}$$

input

```
integrate((B*x^2+A)/(b*x^2+a)^(3/2)/(d*x^2+c)/(f*x^2+e),x, algorithm="fricas")
```

output

Timed out

Sympy [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)} dx = \text{Timed out}$$

input

```
integrate((B*x**2+A)/(b*x**2+a)**(3/2)/(d*x**2+c)/(f*x**2+e),x)
```

output

Timed out

Maxima [F]

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)} dx = \int \frac{Bx^2 + A}{(bx^2 + a)^{3/2} (dx^2 + c)(fx^2 + e)} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)^(3/2)/(d*x^2+c)/(f*x^2+e),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/((b*x^2 + a)^(3/2)*(d*x^2 + c)*(f*x^2 + e)), x)`

Giac [A] (verification not implemented)

Time = 0.70 (sec) , antiderivative size = 277, normalized size of antiderivative = 1.51

$$\begin{aligned} & \int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)} dx = \\ & \frac{(Bab - Ab^2)x}{(ab^2ce - a^2bde - a^2bcf + a^3df)\sqrt{bx^2 + a}} \\ & - \frac{(B\sqrt{bcd} - A\sqrt{bd^2}) \arctan\left(\frac{(\sqrt{bx} - \sqrt{bx^2+a})^2 d + 2bc - ad}{2\sqrt{-b^2c^2 + abcd}}\right)}{\sqrt{-b^2c^2 + abcd}(bcde - ad^2e - bc^2f + acdf)} \\ & + \frac{(B\sqrt{bef} - A\sqrt{bf^2}) \arctan\left(\frac{(\sqrt{bx} - \sqrt{bx^2+a})^2 f + 2be - af}{2\sqrt{-b^2e^2 + abef}}\right)}{\sqrt{-b^2e^2 + abef}(bde^2 - bcef - adef + acf^2)} \end{aligned}$$

input `integrate((B*x^2+A)/(b*x^2+a)^(3/2)/(d*x^2+c)/(f*x^2+e),x, algorithm="giac")`

output

```

-(B*a*b - A*b^2)*x/((a*b^2*c*e - a^2*b*d*e - a^2*b*c*f + a^3*d*f)*sqrt(b*x
^2 + a)) - (B*sqrt(b)*c*d - A*sqrt(b)*d^2)*arctan(1/2*((sqrt(b)*x - sqrt(b
*x^2 + a))^2*d + 2*b*c - a*d)/sqrt(-b^2*c^2 + a*b*c*d))/(sqrt(-b^2*c^2 + a
*b*c*d)*(b*c*d*e - a*d^2*e - b*c^2*f + a*c*d*f)) + (B*sqrt(b)*e*f - A*sqrt
(b)*f^2)*arctan(1/2*((sqrt(b)*x - sqrt(b*x^2 + a))^2*f + 2*b*e - a*f)/sqrt
(-b^2*e^2 + a*b*e*f))/(sqrt(-b^2*e^2 + a*b*e*f)*(b*d*e^2 - b*c*e*f - a*d*e
*f + a*c*f^2))

```

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)} dx = \int \frac{Bx^2 + A}{(bx^2 + a)^{3/2} (dx^2 + c) (fx^2 + e)} dx$$

input

```
int((A + B*x^2)/((a + b*x^2)^(3/2)*(c + d*x^2)*(e + f*x^2)),x)
```

output

```
int((A + B*x^2)/((a + b*x^2)^(3/2)*(c + d*x^2)*(e + f*x^2)), x)
```

Reduce [B] (verification not implemented)

Time = 0.48 (sec) , antiderivative size = 508, normalized size of antiderivative = 2.78

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)} dx = \frac{\sqrt{c} \sqrt{ad - bc} \operatorname{atan}\left(\frac{\sqrt{ad - bc} - \sqrt{d} \sqrt{bx^2 + a} - \sqrt{d} \sqrt{bx}}{\sqrt{c} \sqrt{b}}\right) adef - \sqrt{c} \sqrt{ad - bc}}{\dots}$$

input

```
int((B*x^2+A)/(b*x^2+a)^(3/2)/(d*x^2+c)/(f*x^2+e),x)
```

output

```
(sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2)
- sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a*d*e**f - sqrt(c)*sqrt(a*d - b*c)*
atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sq
rt(c)*sqrt(b)))*b*d*e**2 + sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) +
sqrt(d)*sqrt(a + b*x**2) + sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a*d*e**f -
sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) + sqrt(d)*sqrt(a + b*x**2)
+ sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*b*d*e**2 - sqrt(e)*sqrt(a*f - b*e)
*atan((sqrt(a*f - b*e) - sqrt(f)*sqrt(a + b*x**2) - sqrt(f)*sqrt(b)*x)/(sq
rt(e)*sqrt(b)))*a*c*d*f + sqrt(e)*sqrt(a*f - b*e)*atan((sqrt(a*f - b*e) -
sqrt(f)*sqrt(a + b*x**2) - sqrt(f)*sqrt(b)*x)/(sqrt(e)*sqrt(b)))*b*c**2*f
- sqrt(e)*sqrt(a*f - b*e)*atan((sqrt(a*f - b*e) + sqrt(f)*sqrt(a + b*x**2)
+ sqrt(f)*sqrt(b)*x)/(sqrt(e)*sqrt(b)))*a*c*d*f + sqrt(e)*sqrt(a*f - b*e)
*atan((sqrt(a*f - b*e) + sqrt(f)*sqrt(a + b*x**2) + sqrt(f)*sqrt(b)*x)/(sq
rt(e)*sqrt(b)))*b*c**2*f)/(c*e*(a**2*c*d*f**2 - a**2*d**2*e*f - a*b*c**2*f
**2 + a*b*d**2*e**2 + b**2*c**2*e*f - b**2*c*d*e**2))
```

$$3.2 \quad \int \frac{A+Bx^2}{(a+bx^2)^{3/2}(c+dx^2)(e+fx^2)^2} dx$$

Optimal result	53
Mathematica [C] (warning: unable to verify)	54
Rubi [A] (verified)	55
Maple [A] (verified)	56
Fricas [F(-1)]	57
Sympy [F(-1)]	57
Maxima [F]	57
Giac [B] (verification not implemented)	58
Mupad [F(-1)]	59
Reduce [B] (verification not implemented)	60

Optimal result

Integrand size = 37, antiderivative size = 356

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)^2} dx =$$

$$\frac{b(aBe(2bde - 3bcf + adf) + A(abc f^2 - a^2 d f^2 - 2b^2 e(de - cf))) x}{2a(bc - ad)e(be - af)^2(de - cf)\sqrt{a + bx^2}}$$

$$- \frac{f(Be - Af)x}{2e(be - af)(de - cf)\sqrt{a + bx^2}(e + fx^2)} + \frac{d^2(Bc - Ad)\operatorname{arctanh}\left(\frac{\sqrt{bc - ad}x}{\sqrt{c}\sqrt{a + bx^2}}\right)}{\sqrt{c}(bc - ad)^{3/2}(de - cf)^2}$$

$$+ \frac{f(2be(Af(3de - 2cf) - Be(2de - cf)) - af(Af(3de - cf) - Be(de + cf)))\operatorname{arctanh}\left(\frac{\sqrt{be - af}x}{\sqrt{e}\sqrt{a + bx^2}}\right)}{2e^{3/2}(be - af)^{5/2}(de - cf)^2}$$

output

```
-1/2*b*(a*B*e*(a*d*f-3*b*c*f+2*b*d*e)+A*(a*b*c*f^2-a^2*d*f^2-2*b^2*e*(-c*f+d*e)))*x/a/(-a*d+b*c)/e/(-a*f+b*e)^2/(-c*f+d*e)/(b*x^2+a)^(1/2)-1/2*f*(-A*f+B*e)*x/e/(-a*f+b*e)/(-c*f+d*e)/(b*x^2+a)^(1/2)/(f*x^2+e)+d^2*(-A*d+B*c)*arctanh((-a*d+b*c)^(1/2)*x/c^(1/2)/(b*x^2+a)^(1/2))/c^(1/2)/(-a*d+b*c)^(3/2)/(-c*f+d*e)^2+1/2*f*(2*b*e*(A*f*(-2*c*f+3*d*e)-B*e*(-c*f+2*d*e))-a*f*(A*f*(-c*f+3*d*e)-B*e*(c*f+d*e)))*arctanh((-a*f+b*e)^(1/2)*x/e^(1/2)/(b*x^2+a)^(1/2))/e^(3/2)/(-a*f+b*e)^(5/2)/(-c*f+d*e)^2
```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 18.03 (sec) , antiderivative size = 1481, normalized size of antiderivative = 4.16

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)^2} dx = \text{Too large to display}$$

input `Integrate[(A + B*x^2)/((a + b*x^2)^(3/2)*(c + d*x^2)*(e + f*x^2)^2),x]`

output

```
(d*(-(B*c) + A*d)*x*(-15*c*sqrt(((b*c - a*d)*x^2)/(c*(a + b*x^2))) - 10*d*x^2*sqrt(((b*c - a*d)*x^2)/(c*(a + b*x^2))) + 15*c*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]] + 10*d*x^2*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]] + 2*c*(((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(5/2)*Hypergeometric2F1[2, 5/2, 7/2, ((b*c - a*d)*x^2)/(c*(a + b*x^2))] + 2*d*x^2*(((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(5/2)*Hypergeometric2F1[2, 5/2, 7/2, ((b*c - a*d)*x^2)/(c*(a + b*x^2))])/((5*a*c^2*(d*e - c*f)^2*(((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(3/2)*sqrt[a + b*x^2]*(1 + (b*x^2)/a)) + ((B*c - A*d)*f*x*(-15*e*sqrt(((b*e - a*f)*x^2)/(e*(a + b*x^2))) - 10*f*x^2*sqrt(((b*e - a*f)*x^2)/(e*(a + b*x^2))) + 15*e*ArcTanh[Sqrt[((b*e - a*f)*x^2)/(e*(a + b*x^2))]] + 10*f*x^2*ArcTanh[Sqrt[((b*e - a*f)*x^2)/(e*(a + b*x^2))]] + 2*e*(((b*e - a*f)*x^2)/(e*(a + b*x^2)))^(5/2)*Hypergeometric2F1[2, 5/2, 7/2, ((b*e - a*f)*x^2)/(e*(a + b*x^2))] + 2*f*x^2*(((b*e - a*f)*x^2)/(e*(a + b*x^2)))^(5/2)*Hypergeometric2F1[2, 5/2, 7/2, ((b*e - a*f)*x^2)/(e*(a + b*x^2))])/((5*a*e^2*(-(d*e) + c*f)^2*(((b*e - a*f)*x^2)/(e*(a + b*x^2)))^(3/2)*sqrt[a + b*x^2]*(1 + (b*x^2)/a)) + ((B*e - A*f)*x*(-2625*sqrt(((b*e - a*f)*x^2)/(e*(a + b*x^2))) - (5250*f*x^2*sqrt(((b*e - a*f)*x^2)/(e*(a + b*x^2))))/e - (2310*f^2*x^4*sqrt(((b*e - a*f)*x^2)/(e*(a + b*x^2))))/e^2 + 70*(((b*e - a*f)*x^2)/(e*(a + b*x^2)))^(3/2) + (560*f*x^2*(((b*e - a*f)*x^2)/(e*(a + b*x^2)))^(3/2))/e + (280*f^2*x^4*(((b*e - a*f)*x^2)/(e*(a + b*x^2)))^(3/2))...
```

Rubi [A] (verified)

Time = 1.26 (sec) , antiderivative size = 431, normalized size of antiderivative = 1.21, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.054$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)^2} dx$$

↓ 7276

$$\int \left(\frac{d(Ad - Bc)}{(a + bx^2)^{3/2} (c + dx^2) (de - cf)^2} + \frac{f(Bc - Ad)}{(a + bx^2)^{3/2} (e + fx^2) (cf - de)^2} + \frac{Be - Af}{(a + bx^2)^{3/2} (e + fx^2)^2 (de - cf)} \right) dx$$

↓ 2009

$$\frac{d^2(Bc - Ad)\operatorname{arctanh}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}(bc - ad)^{3/2}(de - cf)^2} - \frac{f(4be - af)(Be - Af)\operatorname{arctanh}\left(\frac{x\sqrt{be-af}}{\sqrt{e}\sqrt{a+bx^2}}\right)}{2e^{3/2}(be - af)^{5/2}(de - cf)} -$$

$$\frac{f^2(Bc - Ad)\operatorname{arctanh}\left(\frac{x\sqrt{be-af}}{\sqrt{e}\sqrt{a+bx^2}}\right)}{\sqrt{e}(be - af)^{3/2}(de - cf)^2} - \frac{bdx(Bc - Ad)}{a\sqrt{a + bx^2}(bc - ad)(de - cf)^2} +$$

$$\frac{bx(af + 2be)(Be - Af)}{2ae\sqrt{a + bx^2}(be - af)^2(de - cf)} + \frac{bf x(Bc - Ad)}{a\sqrt{a + bx^2}(be - af)(de - cf)^2} -$$

$$\frac{fx(Be - Af)}{2e\sqrt{a + bx^2}(e + fx^2)(be - af)(de - cf)}$$

input

```
Int[(A + B*x^2)/((a + b*x^2)^(3/2)*(c + d*x^2)*(e + f*x^2)^2),x]
```

output

```
-((b*d*(B*c - A*d)*x)/(a*(b*c - a*d)*(d*e - c*f)^2*Sqrt[a + b*x^2])) + (b*(B*c - A*d)*f*x)/(a*(b*e - a*f)*(d*e - c*f)^2*Sqrt[a + b*x^2]) + (b*(2*b*e + a*f)*(B*e - A*f)*x)/(2*a*e*(b*e - a*f)^2*(d*e - c*f)*Sqrt[a + b*x^2]) - (f*(B*e - A*f)*x)/(2*e*(b*e - a*f)*(d*e - c*f)*Sqrt[a + b*x^2]*(e + f*x^2)) + (d^2*(B*c - A*d)*ArcTanh[(Sqrt[b*c - a*d]*x)/(Sqrt[c]*Sqrt[a + b*x^2])])/(Sqrt[c]*(b*c - a*d)^(3/2)*(d*e - c*f)^2) - ((B*c - A*d)*f^2*ArcTanh[(Sqrt[b*e - a*f]*x)/(Sqrt[e]*Sqrt[a + b*x^2])])/(Sqrt[e]*(b*e - a*f)^(3/2)*(d*e - c*f)^2) - (f*(4*b*e - a*f)*(B*e - A*f)*ArcTanh[(Sqrt[b*e - a*f]*x)/(Sqrt[e]*Sqrt[a + b*x^2])])/(2*e^(3/2)*(b*e - a*f)^(5/2)*(d*e - c*f))
```


Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 7276 Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xprand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Maple [A] (verified)

Time = 1.72 (sec) , antiderivative size = 421, normalized size of antiderivative = 1.18

method	result
pseudoelliptic	$\frac{-a\sqrt{(ad-bc)c(ad-bc)}\left(\left(Af^2+Be\right)c-3Adef+Bde^2\right)fa-4be\left(f\left(Af-\frac{Be}{2}\right)c-\frac{3Adef}{2}+Bde^2\right)\sqrt{bx^2+a}f\left(fx^2+e\right)}{\dots}$
default	Expression too large to display

```
input int((B*x^2+A)/(b*x^2+a)^(3/2)/(d*x^2+c)/(f*x^2+e)^2,x,method=_RETURNVERBOS
E)
```

```
output 1/2/(b*x^2+a)^(1/2)/((a*d-b*c)*c)^(1/2)/((a*f-b*e)*e)^(1/2)*(-a*((a*d-b*c)
*c)^(1/2)*(a*d-b*c)*((A*f^2+B*e*f)*c-3*A*d*e*f+B*d*e^2)*f*a-4*b*e*(f*(A*f
-1/2*B*e)*c-3/2*A*d*e*f+B*d*e^2))*(b*x^2+a)^(1/2)*f*(f*x^2+e)*arctan(e*(b*
x^2+a)^(1/2)/x/((a*f-b*e)*e)^(1/2))+((a*f-b*e)*e)^(1/2)*(-2*(b*x^2+a)^(1/2)
)*a*d^2*e*(f*x^2+e)*(a*f-b*e)^2*(A*d-B*c)*arctan(c*(b*x^2+a)^(1/2)/x/((a*d
-b*c)*c)^(1/2))+x*(d*f^2*(A*f-B*e)*a^3-b*f^2*(-d*x^2+c)*(A*f-B*e)*a^2-(f*(
A*f^2*x^2-3*B*e*f*x^2-2*B*e^2)*c+2*B*e^2*d*(f*x^2+e))*b^2*a-2*A*b^3*e*(f*x
^2+e)*(c*f-d*e))*((a*d-b*c)*c)^(1/2)*(c*f-d*e))/(c*f-d*e)^2/(a*d-b*c)/e/(
f*x^2+e)/(a*f-b*e)^2/a
```

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)^2} dx = \text{Timed out}$$

input `integrate((B*x^2+A)/(b*x^2+a)^(3/2)/(d*x^2+c)/(f*x^2+e)^2,x, algorithm="fricas")`

output `Timed out`

Sympy [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)^2} dx = \text{Timed out}$$

input `integrate((B*x**2+A)/(b*x**2+a)**(3/2)/(d*x**2+c)/(f*x**2+e)**2,x)`

output `Timed out`

Maxima [F]

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)^2} dx = \int \frac{Bx^2 + A}{(bx^2 + a)^{\frac{3}{2}} (dx^2 + c) (fx^2 + e)^2} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)^(3/2)/(d*x^2+c)/(f*x^2+e)^2,x, algorithm="maxima")`

output `integrate((B*x^2 + A)/((b*x^2 + a)^(3/2)*(d*x^2 + c)*(f*x^2 + e)^2), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 785 vs. $2(326) = 652$.

Time = 2.14 (sec) , antiderivative size = 785, normalized size of antiderivative = 2.21

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)^2} dx =$$

$$\frac{(Bab^2 - Ab^3)x}{(ab^3ce^2 - a^2b^2de^2 - 2a^2b^2cef + 2a^3bdef + a^3bcf^2 - a^4df^2)\sqrt{bx^2 + a}}$$

$$- \frac{(B\sqrt{bcd^2} - A\sqrt{bd^3}) \arctan\left(\frac{(\sqrt{bx} - \sqrt{bx^2 + a})^2 d + 2bc - ad}{2\sqrt{-b^2c^2 + abcd}}\right)}{(bcd^2e^2 - ad^3e^2 - 2bc^2def + 2acd^2ef + bc^3f^2 - ac^2df^2)\sqrt{-b^2c^2 + abcd}}$$

$$+ \frac{(4Bb^{\frac{3}{2}}de^3f - 2Bb^{\frac{3}{2}}ce^2f^2 - Ba\sqrt{bde^2}f^2 - 6Ab^{\frac{3}{2}}de^2f^2 - Ba\sqrt{bce}f^3 + 4Ab^{\frac{3}{2}}cef^3 + 3Aa\sqrt{bde}f^3 - Aa\sqrt{bce}f^3)}{2(b^2d^2e^5 - 2b^2cde^4f - 2abd^2e^4f + b^2c^2e^3f^2 + 4abcde^3f^2 + a^2d^2e^3f^2 - 2abc^2e^2f^3 - 2a^2cde^2f^3)}$$

$$+ \frac{2(\sqrt{bx} - \sqrt{bx^2 + a})^2 Bb^{\frac{3}{2}}e^2f - (\sqrt{bx} - \sqrt{bx^2 + a})^2 Ba\sqrt{bde}f^2 - 2(\sqrt{bx} - \sqrt{bx^2 + a})^2 Ab^{\frac{3}{2}}ef^2 + (\sqrt{bx} - \sqrt{bx^2 + a})^2 Aa\sqrt{bde}f^3}{(b^2de^4 - b^2ce^3f - 2abde^3f + 2abce^2f^2 + a^2de^2f^2 - a^2cef^3)\left((\sqrt{bx} - \sqrt{bx^2 + a})^4 f + 4(\sqrt{bx} - \sqrt{bx^2 + a})^3\right)}$$

input

```
integrate((B*x^2+A)/(b*x^2+a)^(3/2)/(d*x^2+c)/(f*x^2+e)^2,x, algorithm="giac")
```

output

```

-(B*a*b^2 - A*b^3)*x/((a*b^3*c*e^2 - a^2*b^2*d*e^2 - 2*a^2*b^2*c*e*f + 2*a^3*b*d*e*f + a^3*b*c*f^2 - a^4*d*f^2)*sqrt(b*x^2 + a)) - (B*sqrt(b)*c*d^2 - A*sqrt(b)*d^3)*arctan(1/2*((sqrt(b)*x - sqrt(b*x^2 + a))^2*d + 2*b*c - a*d)/sqrt(-b^2*c^2 + a*b*c*d))/((b*c*d^2*e^2 - a*d^3*e^2 - 2*b*c^2*d*e*f + 2*a*c*d^2*e*f + b*c^3*f^2 - a*c^2*d*f^2)*sqrt(-b^2*c^2 + a*b*c*d)) + 1/2*(4*B*b^(3/2)*d*e^3*f - 2*B*b^(3/2)*c*e^2*f^2 - B*a*sqrt(b)*d*e^2*f^2 - 6*A*b^(3/2)*d*e^2*f^2 - B*a*sqrt(b)*c*e*f^3 + 4*A*b^(3/2)*c*e*f^3 + 3*A*a*sqrt(b)*d*e*f^3 - A*a*sqrt(b)*c*f^4)*arctan(1/2*((sqrt(b)*x - sqrt(b*x^2 + a))^2*f + 2*b*e - a*f)/sqrt(-b^2*e^2 + a*b*e*f))/((b^2*d^2*e^5 - 2*b^2*c*d*e^4*f - 2*a*b*d^2*e^4*f + b^2*c^2*e^3*f^2 + 4*a*b*c*d*e^3*f^2 + a^2*d^2*e^3*f^2 - 2*a*b*c^2*e^2*f^3 - 2*a^2*c*d*e^2*f^3 + a^2*c^2*e*f^4)*sqrt(-b^2*e^2 + a*b*e*f)) + (2*(sqrt(b)*x - sqrt(b*x^2 + a))^2*B*b^(3/2)*e^2*f - (sqrt(b)*x - sqrt(b*x^2 + a))^2*B*a*sqrt(b)*e*f^2 - 2*(sqrt(b)*x - sqrt(b*x^2 + a))^2*A*b^(3/2)*e*f^2 + (sqrt(b)*x - sqrt(b*x^2 + a))^2*A*a*sqrt(b)*f^3 + B*a^2*sqrt(b)*e*f^2 - A*a^2*sqrt(b)*f^3)/((b^2*d*e^4 - b^2*c*e^3*f - 2*a*b*d*e^3*f + 2*a*b*c*e^2*f^2 + a^2*d*e^2*f^2 - a^2*c*e*f^3)*((sqrt(b)*x - sqrt(b*x^2 + a))^4*f + 4*(sqrt(b)*x - sqrt(b*x^2 + a))^2*b*e - 2*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a*f + a^2*f))

```

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)^2} dx = \int \frac{Bx^2 + A}{(bx^2 + a)^{3/2} (dx^2 + c) (fx^2 + e)^2} dx$$

input

```
int((A + B*x^2)/((a + b*x^2)^(3/2)*(c + d*x^2)*(e + f*x^2)^2),x)
```

output

```
int((A + B*x^2)/((a + b*x^2)^(3/2)*(c + d*x^2)*(e + f*x^2)^2), x)
```

Reduce [B] (verification not implemented)

Time = 1.31 (sec) , antiderivative size = 3033, normalized size of antiderivative = 8.52

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)^2} dx = \text{Too large to display}$$

input `int((B*x^2+A)/(b*x^2+a)^(3/2)/(d*x^2+c)/(f*x^2+e)^2,x)`

output `(- 2*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**2*d**2*e**3*f**2 - 2*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**2*d**2*e**2*f**3*x**2 + 4*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a*b*d**2*e**4*f + 4*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a*b*d**2*e**3*f**2*x**2 - 2*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*b**2*d**2*e**5 - 2*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*b**2*d**2*e**4*f*x**2 - 2*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) + sqrt(d)*sqrt(a + b*x**2) + sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**2*d**2*e**3*f**2 - 2*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) + sqrt(d)*sqrt(a + b*x**2) + sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**2*d**2*e**2*f**3*x**2 + 4*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) + sqrt(d)*sqrt(a + b*x**2) + sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a*b*d**2*e**4*f + 4*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) + sqrt(d)*sqrt(a + b*x**2) + sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a*b*d**2*e**3*f**2*x**2 - 2*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) + sqrt(d)*sqrt(a + b*x**2) + sqrt(d)*sqrt(b)*x)/...`

$$3.3 \quad \int \frac{A+Bx^2}{(a+bx^2)^{3/2}(c+dx^2)(e+fx^2)^3} dx$$

Optimal result	61
Mathematica [C] (warning: unable to verify)	62
Rubi [A] (verified)	63
Maple [A] (verified)	65
Fricas [F(-1)]	66
Sympy [F(-1)]	67
Maxima [F]	67
Giac [B] (verification not implemented)	67
Mupad [F(-1)]	68
Reduce [B] (verification not implemented)	69

Optimal result

Integrand size = 37, antiderivative size = 706

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)^3} dx =$$

$$\frac{b(A(2ab^2cef^2(7de - 5cf) + a^3df^3(7de - 3cf) - 8b^3e^2(de - cf)^2 - a^2bf^2(14d^2e^2 - 3cdef - 3c^2f^2)) - a^2d^3e^2(7de - 5cf) + a^2d^3f^3(7de - 3cf) - 8b^3e^2(de - cf)^2 - a^2bf^2(14d^2e^2 - 3cdef - 3c^2f^2)) - a^2d^3e^2(7de - 5cf) + a^2d^3f^3(7de - 3cf) - 8b^3e^2(de - cf)^2 - a^2bf^2(14d^2e^2 - 3cdef - 3c^2f^2))}{8a(bc - ad)e^2(be - af)^3(de - cf)}$$

$$- \frac{f(Be - Af)x}{4e(be - af)(de - cf)\sqrt{a + bx^2}(e + fx^2)^2}$$

$$+ \frac{f(4be(Af(3de - 2cf) - Be(2de - cf)) - af(Af(7de - 3cf) - Be(3de + cf)))x}{8e^2(be - af)^2(de - cf)^2\sqrt{a + bx^2}(e + fx^2)}$$

$$+ \frac{d^3(Bc - Ad)\operatorname{arctanh}\left(\frac{\sqrt{bc - ad}x}{\sqrt{c}\sqrt{a + bx^2}}\right)}{\sqrt{c}(bc - ad)^{3/2}(de - cf)^3}$$

$$+ \frac{f(4abef(Be(3d^2e^2 + 3cdef - 2c^2f^2) - Af(12d^2e^2 - 11cdef + 3c^2f^2)) - a^2f^2(Be(3d^2e^2 + 6cdef - c^2f^2) - Af(12d^2e^2 - 11cdef + 3c^2f^2))}{8e^{5/2}(bc - ad)^{3/2}(de - cf)^3}$$

output

```

-1/8*b*(A*(2*a*b^2*c*e*f^2*(-5*c*f+7*d*e)+a^3*d*f^3*(-3*c*f+7*d*e)-8*b^3*e
^2*(-c*f+d*e)^2-a^2*b*f^2*(-3*c^2*f^2-3*c*d*e*f+14*d^2*e^2))-a*B*e*(a^2*d*
f^2*(c*f+3*d*e)-a*b*f*(c^2*f^2-3*c*d*e*f+10*d^2*e^2)-2*b^2*e*(7*c^2*f^2-13
*c*d*e*f+4*d^2*e^2))*x/a/(-a*d+b*c)/e^2/(-a*f+b*e)^3/(-c*f+d*e)^2/(b*x^2+
a)^(1/2)-1/4*f*(-A*f+B*e)*x/e/(-a*f+b*e)/(-c*f+d*e)/(b*x^2+a)^(1/2)/(f*x^2
+e)^2+1/8*f*(4*b*e*(A*f*(-2*c*f+3*d*e)-B*e*(-c*f+2*d*e))-a*f*(A*f*(-3*c*f+
7*d*e)-B*e*(c*f+3*d*e))*x/e^2/(-a*f+b*e)^2/(-c*f+d*e)^2/(b*x^2+a)^(1/2)/((
f*x^2+e)+d^3*(-A*d+B*c)*arctanh((-a*d+b*c)^(1/2)*x/c^(1/2)/(b*x^2+a)^(1/2)
)/c^(1/2)/(-a*d+b*c)^(3/2)/(-c*f+d*e)^3+1/8*f*(4*a*b*e*f*(B*e*(-2*c^2*f^2+
3*c*d*e*f+3*d^2*e^2)-A*f*(3*c^2*f^2-11*c*d*e*f+12*d^2*e^2))-a^2*f^2*(B*e*(
-c^2*f^2+6*c*d*e*f+3*d^2*e^2)-A*f*(3*c^2*f^2-10*c*d*e*f+15*d^2*e^2))-8*b^2
*e^2*(B*e*(c^2*f^2-3*c*d*e*f+3*d^2*e^2)-A*f*(3*c^2*f^2-8*c*d*e*f+6*d^2*e^2
))*arctanh((-a*f+b*e)^(1/2)*x/e^(1/2)/(b*x^2+a)^(1/2))/e^(5/2)/(-a*f+b*e)
^(7/2)/(-c*f+d*e)^3

```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 20.69 (sec) , antiderivative size = 2913, normalized size of antiderivative = 4.13

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)^3} dx = \text{Result too large to show}$$

input

```
Integrate[(A + B*x^2)/((a + b*x^2)^(3/2)*(c + d*x^2)*(e + f*x^2)^3),x]
```

output

```
(d^2*(-(B*c) + A*d)*x*(-15*c*Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))] - 10*
d*x^2*Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))] + 15*c*ArcTanh[Sqrt[((b*c -
a*d)*x^2)/(c*(a + b*x^2))] + 10*d*x^2*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(
a + b*x^2))] + 2*c*(((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(5/2)*Hypergeometr
ic2F1[2, 5/2, 7/2, ((b*c - a*d)*x^2)/(c*(a + b*x^2))] + 2*d*x^2*(((b*c - a
*d)*x^2)/(c*(a + b*x^2)))^(5/2)*Hypergeometric2F1[2, 5/2, 7/2, ((b*c - a*d
)*x^2)/(c*(a + b*x^2))])/((5*a*c^2*(d*e - c*f)^3*(((b*c - a*d)*x^2)/(c*(a
+ b*x^2)))^(3/2)*Sqrt[a + b*x^2]*(1 + (b*x^2)/a)) + (d*(-(B*c) + A*d)*f*x*
(-15*e*Sqrt[((b*e - a*f)*x^2)/(e*(a + b*x^2))] - 10*f*x^2*Sqrt[((b*e - a*f
)*x^2)/(e*(a + b*x^2))] + 15*e*ArcTanh[Sqrt[((b*e - a*f)*x^2)/(e*(a + b*x^
2))] + 10*f*x^2*ArcTanh[Sqrt[((b*e - a*f)*x^2)/(e*(a + b*x^2))] + 2*e*((
(b*e - a*f)*x^2)/(e*(a + b*x^2)))^(5/2)*Hypergeometric2F1[2, 5/2, 7/2, ((b
*e - a*f)*x^2)/(e*(a + b*x^2))] + 2*f*x^2*(((b*e - a*f)*x^2)/(e*(a + b*x^2
)))^(5/2)*Hypergeometric2F1[2, 5/2, 7/2, ((b*e - a*f)*x^2)/(e*(a + b*x^2))
])/((5*a*e^2*(-(d*e) + c*f)^3*(((b*e - a*f)*x^2)/(e*(a + b*x^2)))^(3/2)*Sq
rt[a + b*x^2]*(1 + (b*x^2)/a)) + ((B*c - A*d)*f*x*(-2625*Sqrt[((b*e - a*f)
*x^2)/(e*(a + b*x^2))] - (5250*f*x^2*Sqrt[((b*e - a*f)*x^2)/(e*(a + b*x^2
))])/e - (2310*f^2*x^4*Sqrt[((b*e - a*f)*x^2)/(e*(a + b*x^2))])/e^2 + 70*((
(b*e - a*f)*x^2)/(e*(a + b*x^2)))^(3/2) + (560*f*x^2*(((b*e - a*f)*x^2)/(e
*(a + b*x^2)))^(3/2))/e + (280*f^2*x^4*(((b*e - a*f)*x^2)/(e*(a + b*x^2)...
```

Rubi [A] (verified)

Time = 1.81 (sec) , antiderivative size = 736, normalized size of antiderivative = 1.04, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.054$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)^3} dx$$

↓ 7276

$$\int \left(\frac{d^2(Ad - Bc)}{(a + bx^2)^{3/2} (c + dx^2) (de - cf)^3} + \frac{df(Ad - Bc)}{(a + bx^2)^{3/2} (e + fx^2) (cf - de)^3} + \frac{f(Bc - Ad)}{(a + bx^2)^{3/2} (e + fx^2)^2 (cf - de)^2} \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{3f(a^2f^2 - 4abef + 8b^2e^2)(Be - Af)\operatorname{arctanh}\left(\frac{x\sqrt{be-af}}{\sqrt{e}\sqrt{a+bx^2}}\right)}{8e^{5/2}(be-af)^{7/2}(de-cf)} + \\
& \frac{d^3(Bc - Ad)\operatorname{arctanh}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}(bc-ad)^{3/2}(de-cf)^3} - \frac{f^2(4be-af)(Bc-Ad)\operatorname{arctanh}\left(\frac{x\sqrt{be-af}}{\sqrt{e}\sqrt{a+bx^2}}\right)}{2e^{3/2}(be-af)^{5/2}(de-cf)^2} - \\
& \frac{df^2(Bc-Ad)\operatorname{arctanh}\left(\frac{x\sqrt{be-af}}{\sqrt{e}\sqrt{a+bx^2}}\right)}{\sqrt{e}(be-af)^{3/2}(de-cf)^3} - \frac{bd^2x(Bc-Ad)}{a\sqrt{a+bx^2}(bc-ad)(de-cf)^3} + \\
& \frac{fx\sqrt{a+bx^2}(4be-af)(3af+2be)(Be-Af)}{8ae^2(e+fx^2)(be-af)^3(de-cf)} - \frac{f^2x(Bc-Ad)}{2e\sqrt{a+bx^2}(e+fx^2)(be-af)(de-cf)^2} + \\
& \frac{bdfx(Bc-Ad)}{a\sqrt{a+bx^2}(be-af)(de-cf)^3} + \frac{2ae\sqrt{a+bx^2}(be-af)^2(de-cf)^2}{bfx(af+2be)(Bc-Ad)} + \\
& \frac{bx(af+4be)(Be-Af)}{4ae\sqrt{a+bx^2}(e+fx^2)(be-af)^2(de-cf)} - \frac{fx(Be-Af)}{4e\sqrt{a+bx^2}(e+fx^2)^2(be-af)(de-cf)}
\end{aligned}$$

input

```
Int[(A + B*x^2)/((a + b*x^2)^(3/2)*(c + d*x^2)*(e + f*x^2)^3),x]
```

output

```

-((b*d^2*(B*c - A*d)*x)/(a*(b*c - a*d)*(d*e - c*f)^3*Sqrt[a + b*x^2])) + (
b*d*(B*c - A*d)*f*x)/(a*(b*e - a*f)*(d*e - c*f)^3*Sqrt[a + b*x^2]) + (b*(B
*c - A*d)*f*(2*b*e + a*f)*x)/(2*a*e*(b*e - a*f)^2*(d*e - c*f)^2*Sqrt[a + b
*x^2]) - (f*(B*e - A*f)*x)/(4*e*(b*e - a*f)*(d*e - c*f)*Sqrt[a + b*x^2]*(e
+ f*x^2)^2) - ((B*c - A*d)*f^2*x)/(2*e*(b*e - a*f)*(d*e - c*f)^2*Sqrt[a +
b*x^2]*(e + f*x^2)) + (b*(4*b*e + a*f)*(B*e - A*f)*x)/(4*a*e*(b*e - a*f)^
2*(d*e - c*f)*Sqrt[a + b*x^2]*(e + f*x^2)) + (f*(4*b*e - a*f)*(2*b*e + 3*a
*f)*(B*e - A*f)*x*Sqrt[a + b*x^2])/(8*a*e^2*(b*e - a*f)^3*(d*e - c*f)*(e +
f*x^2)) + (d^3*(B*c - A*d)*ArcTanh[(Sqrt[b*c - a*d]*x)/(Sqrt[c]*Sqrt[a +
b*x^2])])/(Sqrt[c]*(b*c - a*d)^(3/2)*(d*e - c*f)^3) - (d*(B*c - A*d)*f^2*A
rcTanh[(Sqrt[b*e - a*f]*x)/(Sqrt[e]*Sqrt[a + b*x^2])])/(Sqrt[e]*(b*e - a*f
)^(3/2)*(d*e - c*f)^3) - ((B*c - A*d)*f^2*(4*b*e - a*f)*ArcTanh[(Sqrt[b*e
- a*f]*x)/(Sqrt[e]*Sqrt[a + b*x^2])])/(2*e^(3/2)*(b*e - a*f)^(5/2)*(d*e -
c*f)^2) - (3*f*(B*e - A*f)*(8*b^2*e^2 - 4*a*b*e*f + a^2*f^2)*ArcTanh[(Sqrt
[b*e - a*f]*x)/(Sqrt[e]*Sqrt[a + b*x^2])])/(8*e^(5/2)*(b*e - a*f)^(7/2)*(d
*e - c*f))

```

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
x
pand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]`

Maple [A] (verified)

Time = 3.35 (sec) , antiderivative size = 953, normalized size of antiderivative = 1.35

method	result	size
pseudoelliptic	Expression too large to display	953
default	Expression too large to display	5470

input `int((B*x^2+A)/(b*x^2+a)^(3/2)/(d*x^2+c)/(f*x^2+e)^3,x,method=_RETURNVERBOS
E)`

output

```

5/8/(b*x^2+a)^(1/2)/((a*d-b*c)*c)^(1/2)/((a*f-b*e)*e)^(1/2)*(-3/5*(f^2*(f^
2*(A*f+1/3*B*e)*c^2-10/3*(A*f+3/5*B*e)*f*e*d*c+5*A*d^2*e^2*f-B*d^2*e^3)*a^
2-4*b*(f^2*(A*f+2/3*B*e)*c^2+d*(-11/3*e*A*f^2-B*e^2*f)*c+4*A*d^2*e^2*f-B*d
^2*e^3)*f*e*a+8*b^2*(f^2*(A*f-1/3*B*e)*c^2-8/3*f*e*d*(A*f-3/8*B*e)*c+2*A*d
^2*e^2*f-B*d^2*e^3)*e^2)*a*((a*d-b*c)*c)^(1/2)*(a*d-b*c)*(b*x^2+a)^(1/2)*f
*(f*x^2+e)^2*arctan(e*(b*x^2+a)^(1/2)/x/((a*f-b*e)*e)^(1/2))+((a*f-b*e)*e)
^(1/2)*(8/5*(b*x^2+a)^(1/2)*a*d^3*e^2*(f*x^2+e)^2*(a*f-b*e)^3*(A*d-B*c)*ar
ctan(c*(b*x^2+a)^(1/2)/x/((a*d-b*c)*c)^(1/2))+x*((f*(-1/5*B*e^2+f*(1/5*x^2
*B+A)*e+3/5*f^2*x^2*A)*c-9/5*(-5/9*B*e^2+f*(-1/3*x^2*B+A)*e+7/9*f^2*x^2*A)
*e*d)*f^3*d*a^4-(f^2*(-1/5*B*e^2+f*(1/5*x^2*B+A)*e+3/5*f^2*x^2*A)*c^2+3/5*
(-B*e^3+f*(-2/3*x^2*B+A)*e^2-2/3*x^2*(1/2*x^2*B+A)*f^2*e-A*f^3*x^4)*f*d*c-
16/5*(-3/4*B*e^3+f*(-5/16*x^2*B+A)*e^2+5/16*x^2*(3/5*x^2*B+A)*f^2*e-7/16*A
*f^3*x^4)*e*d^2)*b*f^2*a^3+12/5*b^2*((-2/3*B*e^3+f*(-5/12*x^2*B+A)*e^2+5/1
2*x^2*f^2*(-1/5*x^2*B+A)*e-1/4*A*f^3*x^4)*f*c^2-4/3*e*d*(-3/4*B*e^3+f*(-13
/16*x^2*B+A)*e^2+17/16*x^2*f^2*(-3/17*x^2*B+A)*e+3/16*A*f^3*x^4)*c+4/3*x^2
*e^2*d^2*(-3/4*B*e^2+f*(-5/8*x^2*B+A)*e+7/8*f^2*x^2*A))*f^2*a^2+12/5*b^3*(
(-2/3*B*e^3-2*B*e^2*f*x^2+f^2*x^2*(-7/6*x^2*B+A)*e+5/6*A*f^3*x^4)*f^2*c^2-
4/3*f*e*d*(-B*e^3-11/4*B*e^2*f*x^2+f^2*x^2*(-13/8*x^2*B+A)*e+7/8*A*f^3*x^4
)*c-2/3*B*d^2*e^3*(f*x^2+e)^2)*e*a+8/5*A*b^4*e^2*(f*x^2+e)^2*(c*f-d*e)^2)*
((a*d-b*c)*c)^(1/2)*(c*f-d*e)))/(c*f-d*e)^3/(a*f-b*e)^3/(a*d-b*c)/e^2/(...

```

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)^3} dx = \text{Timed out}$$

input

```

integrate((B*x^2+A)/(b*x^2+a)^(3/2)/(d*x^2+c)/(f*x^2+e)^3,x, algorithm="fr
icas")

```

output

Timed out

Sympy [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)^3} dx = \text{Timed out}$$

input `integrate((B*x**2+A)/(b*x**2+a)**(3/2)/(d*x**2+c)/(f*x**2+e)**3,x)`

output `Timed out`

Maxima [F]

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)^3} dx = \int \frac{Bx^2 + A}{(bx^2 + a)^{\frac{3}{2}} (dx^2 + c) (fx^2 + e)^3} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)^(3/2)/(d*x^2+c)/(f*x^2+e)^3,x, algorithm="maxima")`

output `integrate((B*x^2 + A)/((b*x^2 + a)^(3/2)*(d*x^2 + c)*(f*x^2 + e)^3), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2528 vs. $2(671) = 1342$.

Time = 8.73 (sec) , antiderivative size = 2528, normalized size of antiderivative = 3.58

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)^3} dx = \text{Too large to display}$$

input `integrate((B*x^2+A)/(b*x^2+a)^(3/2)/(d*x^2+c)/(f*x^2+e)^3,x, algorithm="giac")`

output

```

-(B*a*b^3 - A*b^4)*x/((a*b^4*c*e^3 - a^2*b^3*d*e^3 - 3*a^2*b^3*c*e^2*f + 3
*a^3*b^2*d*e^2*f + 3*a^3*b^2*c*e*f^2 - 3*a^4*b*d*e*f^2 - a^4*b*c*f^3 + a^5
*d*f^3)*sqrt(b*x^2 + a)) - (B*sqrt(b)*c*d^3 - A*sqrt(b)*d^4)*arctan(1/2*((
sqrt(b)*x - sqrt(b*x^2 + a))^2*d + 2*b*c - a*d)/sqrt(-b^2*c^2 + a*b*c*d))/
((b*c*d^3*e^3 - a*d^4*e^3 - 3*b*c^2*d^2*e^2*f + 3*a*c*d^3*e^2*f + 3*b*c^3*
d*e*f^2 - 3*a*c^2*d^2*e*f^2 - b*c^4*f^3 + a*c^3*d*f^3)*sqrt(-b^2*c^2 + a*b
*c*d)) + 1/8*(24*B*b^(5/2)*d^2*e^5*f - 24*B*b^(5/2)*c*d*e^4*f^2 - 12*B*a*b
^(3/2)*d^2*e^4*f^2 - 48*A*b^(5/2)*d^2*e^4*f^2 + 8*B*b^(5/2)*c^2*e^3*f^3 -
12*B*a*b^(3/2)*c*d*e^3*f^3 + 64*A*b^(5/2)*c*d*e^3*f^3 + 3*B*a^2*sqrt(b)*d^
2*e^3*f^3 + 48*A*a*b^(3/2)*d^2*e^3*f^3 + 8*B*a*b^(3/2)*c^2*e^2*f^4 - 24*A*
b^(5/2)*c^2*e^2*f^4 + 6*B*a^2*sqrt(b)*c*d*e^2*f^4 - 44*A*a*b^(3/2)*c*d*e^2
*f^4 - 15*A*a^2*sqrt(b)*d^2*e^2*f^4 - B*a^2*sqrt(b)*c^2*e*f^5 + 12*A*a*b^(
3/2)*c^2*e*f^5 + 10*A*a^2*sqrt(b)*c*d*e*f^5 - 3*A*a^2*sqrt(b)*c^2*f^6)*arc
tan(1/2*((sqrt(b)*x - sqrt(b*x^2 + a))^2*f + 2*b*e - a*f)/sqrt(-b^2*e^2 +
a*b*e*f))/((b^3*d^3*e^8 - 3*b^3*c*d^2*e^7*f - 3*a*b^2*d^3*e^7*f + 3*b^3*c^
2*d*e^6*f^2 + 9*a*b^2*c*d^2*e^6*f^2 + 3*a^2*b*d^3*e^6*f^2 - b^3*c^3*e^5*f^
3 - 9*a*b^2*c^2*d*e^5*f^3 - 9*a^2*b*c*d^2*e^5*f^3 - a^3*d^3*e^5*f^3 + 3*a*
b^2*c^3*e^4*f^4 + 9*a^2*b*c^2*d*e^4*f^4 + 3*a^3*c*d^2*e^4*f^4 - 3*a^2*b*c^
3*e^3*f^5 - 3*a^3*c^2*d*e^3*f^5 + a^3*c^3*e^2*f^6)*sqrt(-b^2*e^2 + a*b*e*f
)) + 1/4*(16*(sqrt(b)*x - sqrt(b*x^2 + a))^6*B*b^(5/2)*d*e^4*f^2 - 8*(s...

```

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)^3} dx = \int \frac{Bx^2 + A}{(bx^2 + a)^{3/2} (dx^2 + c) (fx^2 + e)^3} dx$$

input

```
int((A + B*x^2)/((a + b*x^2)^(3/2)*(c + d*x^2)*(e + f*x^2)^3),x)
```

output

```
int((A + B*x^2)/((a + b*x^2)^(3/2)*(c + d*x^2)*(e + f*x^2)^3), x)
```

Reduce [B] (verification not implemented)

Time = 5.65 (sec) , antiderivative size = 12786, normalized size of antiderivative = 18.11

$$\int \frac{A + Bx^2}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)^3} dx = \text{Too large to display}$$

input `int((B*x^2+A)/(b*x^2+a)^(3/2)/(d*x^2+c)/(f*x^2+e)^3,x)`

output

```
(16*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**4*d**3*e**5*f**4 + 32*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**4*d**3*e**4*f**5*x**2 + 16*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**4*d**3*e**3*f**6*x**4 - 80*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**3*b*d**3*e**6*f**3 - 160*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**3*b*d**3*e**5*f**4*x**2 - 80*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**3*b*d**3*e**4*f**5*x**4 + 144*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**2*b**2*d**3*e**7*f**2 + 288*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**2*b**2*d**3*e**6*f**3*x**2 + 144*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**2*b**2*d**3*e**5*f**4*x**4 - 112*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a*b**3*d**3*e**8*f - 224*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*...
```

$$3.4 \quad \int \frac{A+Bx^2}{(a+bx^2)^{5/2}(c+dx^2)(e+fx^2)} dx$$

Optimal result	70
Mathematica [A] (verified)	71
Rubi [A] (verified)	71
Maple [A] (verified)	73
Fricas [F(-1)]	73
Sympy [F(-1)]	74
Maxima [F]	74
Giac [B] (verification not implemented)	74
Mupad [F(-1)]	75
Reduce [B] (verification not implemented)	76

Optimal result

Integrand size = 37, antiderivative size = 288

$$\int \frac{A + Bx^2}{(a + bx^2)^{5/2} (c + dx^2) (e + fx^2)} dx = \frac{b(Ab - aB)x}{3a(bc - ad)(be - af)(a + bx^2)^{3/2}} + \frac{b(Ab(2b^2ce + 8a^2df - 5ab(de + cf)) + aB(b^2ce - 5a^2df + 2ab(de + cf)))x}{3a^2(bc - ad)^2(be - af)^2\sqrt{a + bx^2}} - \frac{d^2(Bc - Ad)\operatorname{arctanh}\left(\frac{\sqrt{bc-ad}x}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}(bc - ad)^{5/2}(de - cf)} + \frac{f^2(Be - Af)\operatorname{arctanh}\left(\frac{\sqrt{be-af}x}{\sqrt{e}\sqrt{a+bx^2}}\right)}{\sqrt{e}(be - af)^{5/2}(de - cf)}$$

output

```
1/3*b*(A*b-B*a)*x/a/(-a*d+b*c)/(-a*f+b*e)/(b*x^2+a)^(3/2)+1/3*b*(A*b*(2*b^2*c*e+8*a^2*d*f-5*a*b*(c*f+d*e))+a*B*(b^2*c*e-5*a^2*d*f+2*a*b*(c*f+d*e)))*x/a^2/(-a*d+b*c)^2/(-a*f+b*e)^2/(b*x^2+a)^(1/2)-d^2*(-A*d+B*c)*arctanh((-a*d+b*c)^(1/2)*x/c^(1/2)/(b*x^2+a)^(1/2))/c^(1/2)/(-a*d+b*c)^(5/2)/(-c*f+d*e)+f^2*(-A*f+B*e)*arctanh((-a*f+b*e)^(1/2)*x/e^(1/2)/(b*x^2+a)^(1/2))/e^(1/2)/(-a*f+b*e)^(5/2)/(-c*f+d*e)
```

Mathematica [A] (verified)

Time = 3.58 (sec) , antiderivative size = 337, normalized size of antiderivative = 1.17

$$\int \frac{A + Bx^2}{(a + bx^2)^{5/2} (c + dx^2) (e + fx^2)} dx = \frac{bx(-6a^4Bdf + 2Ab^4cex^2 + ab^3(Bcex^2 + A(3ce - 5dex^2 - 5cfx^2))}{3a^2(b^2c^2 - 5c^2fx^2)} + \frac{d^2(Bc - Ad) \arctan\left(\frac{-dx\sqrt{a+bx^2} + \sqrt{b}(c+dx^2)}{\sqrt{c}\sqrt{-bc+ad}}\right)}{\sqrt{c}(-bc + ad)^{5/2}(de - cf)} + \frac{f^2(-Be + Af) \arctan\left(\frac{-fx\sqrt{a+bx^2} + \sqrt{b}(e+fx^2)}{\sqrt{e}\sqrt{-be+af}}\right)}{\sqrt{e}(-be + af)^{5/2}(de - cf)}$$

input

```
Integrate[(A + B*x^2)/((a + b*x^2)^(5/2)*(c + d*x^2)*(e + f*x^2)),x]
```

output

```
(b*x*(-6*a^4*B*d*f + 2*A*b^4*c*e*x^2 + a*b^3*(B*c*e*x^2 + A*(3*c*e - 5*d*e*x^2 - 5*c*f*x^2)) + a^3*b*(9*A*d*f + B*(3*d*e + 3*c*f - 5*d*f*x^2)) + 2*a^2*b^2*(B*(d*e + c*f)*x^2 + A*(-3*d*e - 3*c*f + 4*d*f*x^2)))/(3*a^2*(b*c - a*d)^2*(b*e - a*f)^2*(a + b*x^2)^(3/2)) + (d^2*(B*c - A*d)*ArcTan[(-(d*x*Sqrt[a + b*x^2]) + Sqrt[b]*(c + d*x^2))/(Sqrt[c]*Sqrt[-(b*c) + a*d])]/(Sqrt[c]*(-(b*c) + a*d)^(5/2)*(d*e - c*f)) + (f^2*(-(B*e) + A*f)*ArcTan[(-(f*x*Sqrt[a + b*x^2]) + Sqrt[b]*(e + f*x^2))/(Sqrt[e]*Sqrt[-(b*e) + a*f])]/(Sqrt[e]*(-(b*e) + a*f)^(5/2)*(d*e - c*f)))/
```

Rubi [A] (verified)

Time = 1.30 (sec) , antiderivative size = 352, normalized size of antiderivative = 1.22, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.054$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{(a + bx^2)^{5/2} (c + dx^2) (e + fx^2)} dx$$

↓ 7276

$$\int \left(\frac{Be - Af}{(a + bx^2)^{5/2} (e + fx^2) (de - cf)} - \frac{Bc - Ad}{(a + bx^2)^{5/2} (c + dx^2) (de - cf)} \right) dx$$

↓ 2009

$$\begin{aligned} & -\frac{bx(2bc - 5ad)(Bc - Ad)}{3a^2\sqrt{a + bx^2}(bc - ad)^2(de - cf)} + \frac{bx(2be - 5af)(Be - Af)}{3a^2\sqrt{a + bx^2}(be - af)^2(de - cf)} - \\ & \frac{d^2(Bc - Ad)\operatorname{arctanh}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}(bc - ad)^{5/2}(de - cf)} + \frac{f^2(Be - Af)\operatorname{arctanh}\left(\frac{x\sqrt{be-af}}{\sqrt{e}\sqrt{a+bx^2}}\right)}{\sqrt{e}(be - af)^{5/2}(de - cf)} - \\ & \frac{bx(Bc - Ad)}{3a(a + bx^2)^{3/2}(bc - ad)(de - cf)} + \frac{bx(Be - Af)}{3a(a + bx^2)^{3/2}(be - af)(de - cf)} \end{aligned}$$

input `Int[(A + B*x^2)/((a + b*x^2)^(5/2)*(c + d*x^2)*(e + f*x^2)),x]`

output `-1/3*(b*(B*c - A*d)*x)/(a*(b*c - a*d)*(d*e - c*f)*(a + b*x^2)^(3/2)) + (b*(B*e - A*f)*x)/(3*a*(b*e - a*f)*(d*e - c*f)*(a + b*x^2)^(3/2)) - (b*(2*b*c - 5*a*d)*(B*c - A*d)*x)/(3*a^2*(b*c - a*d)^2*(d*e - c*f)*Sqrt[a + b*x^2]) + (b*(2*b*e - 5*a*f)*(B*e - A*f)*x)/(3*a^2*(b*e - a*f)^2*(d*e - c*f)*Sqrt[a + b*x^2]) - (d^2*(B*c - A*d)*ArcTanh[(Sqrt[b*c - a*d]*x)/(Sqrt[c]*Sqrt[a + b*x^2])])/(Sqrt[c]*(b*c - a*d)^(5/2)*(d*e - c*f)) + (f^2*(B*e - A*f)*ArcTanh[(Sqrt[b*e - a*f]*x)/(Sqrt[e]*Sqrt[a + b*x^2])])/(Sqrt[e]*(b*e - a*f)^(5/2)*(d*e - c*f))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 1.47 (sec) , antiderivative size = 291, normalized size of antiderivative = 1.01

method	result
pseudoelliptic	$\frac{(Ad-Bc)a^2 d^2 \arctan\left(\frac{c\sqrt{bx^2+a}}{x\sqrt{(ad-bc)c}}\right)}{(cf-de)(ad-bc)^2\sqrt{(ad-bc)c}} - \frac{(Af-Be)a^2 f^2 \arctan\left(\frac{e\sqrt{bx^2+a}}{x\sqrt{(af-be)e}}\right)}{(cf-de)(af-be)^2\sqrt{(af-be)e}} + \frac{b(3Aa^2 bdf - 2Aa b^2 cf - 2Aa b^2 de + Ab^3 ce - 2Ba^3 df + B^2 a^2 ce)}{(af-be)^2(ad-bc)^2\sqrt{bx^2+a}}$
default	Expression too large to display

input `int((B*x^2+A)/(b*x^2+a)^(5/2)/(d*x^2+c)/(f*x^2+e),x,method=_RETURNVERBOSE)`

output
$$\frac{((A*d-B*c)*a^2*d^2/(c*f-d*e)/(a*d-b*c)^2/((a*d-b*c)*c)^(1/2)*\arctan(c*(b*x^2+a)^(1/2)/x/((a*d-b*c)*c)^(1/2)-(A*f-B*e)*a^2*f^2/(c*f-d*e)/(a*f-b*e)^2/((a*f-b*e)*e)^(1/2)*\arctan(e*(b*x^2+a)^(1/2)/x/((a*f-b*e)*e)^(1/2))+b*(3*A*a^2*b*d*f-2*A*a*b^2*c*f-2*A*a*b^2*d*e+A*b^3*c*e-2*B*a^3*d*f+B*a^2*b*c*f+B*a^2*b*d*e)/(a*f-b*e)^2/(a*d-b*c)^2*x/(b*x^2+a)^(1/2)-1/3*b^2*(A*b-B*a)/(a*f-b*e)/(a*d-b*c)*x^3/(b*x^2+a)^(3/2))/a^2}$$

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2)^{5/2} (c + dx^2) (e + fx^2)} dx = \text{Timed out}$$

input `integrate((B*x^2+A)/(b*x^2+a)^(5/2)/(d*x^2+c)/(f*x^2+e),x, algorithm="fricas")`

output `Timed out`

Sympy [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2)^{5/2} (c + dx^2) (e + fx^2)} dx = \text{Timed out}$$

input `integrate((B*x**2+A)/(b*x**2+a)**(5/2)/(d*x**2+c)/(f*x**2+e),x)`

output `Timed out`

Maxima [F]

$$\int \frac{A + Bx^2}{(a + bx^2)^{5/2} (c + dx^2) (e + fx^2)} dx = \int \frac{Bx^2 + A}{(bx^2 + a)^{5/2} (dx^2 + c)(fx^2 + e)} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)^(5/2)/(d*x^2+c)/(f*x^2+e),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/((b*x^2 + a)^(5/2)*(d*x^2 + c)*(f*x^2 + e)), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2061 vs. 2(260) = 520.

Time = 1.21 (sec) , antiderivative size = 2061, normalized size of antiderivative = 7.16

$$\int \frac{A + Bx^2}{(a + bx^2)^{5/2} (c + dx^2) (e + fx^2)} dx = \text{Too large to display}$$

input `integrate((B*x^2+A)/(b*x^2+a)^(5/2)/(d*x^2+c)/(f*x^2+e),x, algorithm="giac")`

output

```

1/3*((B*a*b^9*c^3*e^3 + 2*A*b^10*c^3*e^3 - 9*A*a*b^9*c^2*d*e^3 - 3*B*a^3*b
^7*c*d^2*e^3 + 12*A*a^2*b^8*c*d^2*e^3 + 2*B*a^4*b^6*d^3*e^3 - 5*A*a^3*b^7*
d^3*e^3 - 9*A*a*b^9*c^3*e^2*f - 9*B*a^3*b^7*c^2*d*e^2*f + 36*A*a^2*b^8*c^2
*d*e^2*f + 18*B*a^4*b^6*c*d^2*e^2*f - 45*A*a^3*b^7*c*d^2*e^2*f - 9*B*a^5*b
^5*d^3*e^2*f + 18*A*a^4*b^6*d^3*e^2*f - 3*B*a^3*b^7*c^3*e*f^2 + 12*A*a^2*b
^8*c^3*e*f^2 + 18*B*a^4*b^6*c^2*d*e*f^2 - 45*A*a^3*b^7*c^2*d*e*f^2 - 27*B*
a^5*b^5*c*d^2*e*f^2 + 54*A*a^4*b^6*c*d^2*e*f^2 + 12*B*a^6*b^4*d^3*e*f^2 -
21*A*a^5*b^5*d^3*e*f^2 + 2*B*a^4*b^6*c^3*f^3 - 5*A*a^3*b^7*c^3*f^3 - 9*B*a
^5*b^5*c^2*d*f^3 + 18*A*a^4*b^6*c^2*d*f^3 + 12*B*a^6*b^4*c*d^2*f^3 - 21*A*
a^5*b^5*c*d^2*f^3 - 5*B*a^7*b^3*d^3*f^3 + 8*A*a^6*b^4*d^3*f^3)*x^2/(a^2*b^
9*c^4*e^4 - 4*a^3*b^8*c^3*d*e^4 + 6*a^4*b^7*c^2*d^2*e^4 - 4*a^5*b^6*c*d^3*
e^4 + a^6*b^5*d^4*e^4 - 4*a^3*b^8*c^4*e^3*f + 16*a^4*b^7*c^3*d*e^3*f - 24*
a^5*b^6*c^2*d^2*e^3*f + 16*a^6*b^5*c*d^3*e^3*f - 4*a^7*b^4*d^4*e^3*f + 6*a
^4*b^7*c^4*e^2*f^2 - 24*a^5*b^6*c^3*d*e^2*f^2 + 36*a^6*b^5*c^2*d^2*e^2*f^2
- 24*a^7*b^4*c*d^3*e^2*f^2 + 6*a^8*b^3*d^4*e^2*f^2 - 4*a^5*b^6*c^4*e*f^3
+ 16*a^6*b^5*c^3*d*e*f^3 - 24*a^7*b^4*c^2*d^2*e*f^3 + 16*a^8*b^3*c*d^3*e*f
^3 - 4*a^9*b^2*d^4*e*f^3 + a^6*b^5*c^4*f^4 - 4*a^7*b^4*c^3*d*f^4 + 6*a^8*b
^3*c^2*d^2*f^4 - 4*a^9*b^2*c*d^3*f^4 + a^10*b*d^4*f^4) + 3*(A*a*b^9*c^3*e^
3 + B*a^3*b^7*c^2*d*e^3 - 4*A*a^2*b^8*c^2*d*e^3 - 2*B*a^4*b^6*c*d^2*e^3 +
5*A*a^3*b^7*c*d^2*e^3 + B*a^5*b^5*d^3*e^3 - 2*A*a^4*b^6*d^3*e^3 + B*a^3...

```

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2)^{5/2} (c + dx^2) (e + fx^2)} dx = \int \frac{Bx^2 + A}{(bx^2 + a)^{5/2} (dx^2 + c) (fx^2 + e)} dx$$

input

```
int((A + B*x^2)/((a + b*x^2)^(5/2)*(c + d*x^2)*(e + f*x^2)),x)
```

output

```
int((A + B*x^2)/((a + b*x^2)^(5/2)*(c + d*x^2)*(e + f*x^2)), x)
```

Reduce [B] (verification not implemented)

Time = 1.32 (sec) , antiderivative size = 2231, normalized size of antiderivative = 7.75

$$\int \frac{A + Bx^2}{(a + bx^2)^{5/2} (c + dx^2) (e + fx^2)} dx = \text{Too large to display}$$

input `int((B*x^2+A)/(b*x^2+a)^(5/2)/(d*x^2+c)/(f*x^2+e),x)`

output

```
(sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2)
- sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**4*d**2*e*f**2 - 2*sqrt(c)*sqrt(
a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt
(b)*x)/(sqrt(c)*sqrt(b)))*a**3*b*d**2*e**2*f + sqrt(c)*sqrt(a*d - b*c)*ata
n((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)
)*sqrt(b)))*a**3*b*d**2*e*f**2*x**2 + sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a
*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)
))*a**2*b**2*d**2*e**3 - 2*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) -
sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**2*b**
2*d**2*e**2*f*x**2 + sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(
d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a*b**3*d**2*e*
*3*x**2 + sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) + sqrt(d)*sqrt(a +
b*x**2) + sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**4*d**2*e*f**2 - 2*sqrt
(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) + sqrt(d)*sqrt(a + b*x**2) + sqr
t(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**3*b*d**2*e**2*f + sqrt(c)*sqrt(a*d -
b*c)*atan((sqrt(a*d - b*c) + sqrt(d)*sqrt(a + b*x**2) + sqrt(d)*sqrt(b)*x
)/(sqrt(c)*sqrt(b)))*a**3*b*d**2*e*f**2*x**2 + sqrt(c)*sqrt(a*d - b*c)*ata
n((sqrt(a*d - b*c) + sqrt(d)*sqrt(a + b*x**2) + sqrt(d)*sqrt(b)*x)/(sqrt(c)
)*sqrt(b)))*a**2*b**2*d**2*e**3 - 2*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d
- b*c) + sqrt(d)*sqrt(a + b*x**2) + sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b...
```

$$3.5 \quad \int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)(e+fx^2)^2} dx$$

Optimal result	77
Mathematica [A] (verified)	78
Rubi [A] (verified)	78
Maple [A] (verified)	80
Fricas [F(-1)]	81
Sympy [F(-1)]	81
Maxima [F]	81
Giac [F(-2)]	82
Mupad [F(-1)]	82
Reduce [B] (verification not implemented)	82

Optimal result

Integrand size = 42, antiderivative size = 310

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)(e+fx^2)^2} dx = -\frac{(Ce^2 - Bef + Af^2) x\sqrt{a+bx^2}}{2ef(de - cf)(e + fx^2)} + \frac{\sqrt{b}C \operatorname{arctanh}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{df^2} - \frac{\sqrt{bc-ad}(c^2C - Bcd + Ad^2) \operatorname{arctanh}\left(\frac{\sqrt{bc-ad}x}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{cd}(de - cf)^2} - \frac{(2be^2((Bc - Ad)f^2 + Ce(de - 2cf)) - af(Ce^2(de - 3cf) - f(Af(3de - cf) - Be(de + cf)))) \operatorname{arctanh}\left(\frac{\sqrt{a+bx^2}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{2e^{3/2}f^2\sqrt{be - af}(de - cf)^2}$$

output

```
-1/2*(A*f^2-B*e*f+C*e^2)*x*(b*x^2+a)^(1/2)/e/f/(-c*f+d*e)/(f*x^2+e)+b^(1/2)*C*arctanh(b^(1/2)*x/(b*x^2+a)^(1/2))/d/f^2-(-a*d+b*c)^(1/2)*(A*d^2-B*c*d+C*c^2)*arctanh((-a*d+b*c)^(1/2)*x/c^(1/2)/(b*x^2+a)^(1/2))/c^(1/2)/d/(-c*f+d*e)^2-1/2*(2*b*e^2*((-A*d+B*c)*f^2+C*e*(-2*c*f+d*e))-a*f*(C*e^2*(-3*c*f+d*e)-f*(A*f*(-c*f+3*d*e)-B*e*(c*f+d*e))))*arctanh((-a*f+b*e)^(1/2)*x/e^(1/2)/(b*x^2+a)^(1/2))/e^(3/2)/f^2/(-a*f+b*e)^(1/2)/(-c*f+d*e)^2
```

Mathematica [A] (verified)

Time = 2.29 (sec) , antiderivative size = 343, normalized size of antiderivative = 1.11

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)(e+fx^2)^2} dx = \frac{1}{2} \left(-\frac{(Ce^2 + f(-Be + Af))x\sqrt{a+bx^2}}{ef(de-cf)(e+fx^2)} \right. \\ \left. - \frac{2\sqrt{-bc+ad}(c^2C - Bcd + Ad^2) \arctan\left(\frac{-dx\sqrt{a+bx^2} + \sqrt{b}(c+dx^2)}{\sqrt{c}\sqrt{-bc+ad}}\right)}{\sqrt{cd}(de-cf)^2} \right. \\ \left. + \frac{(2be^2((Bc - Ad)f^2 + Ce(de - 2cf)) - af(Ce^2(de - 3cf) + f(Af(-3de + cf) + Be(de + cf)))) \arctan\left(\frac{-\sqrt{bx} + \sqrt{a+bx^2}}{\sqrt{c}\sqrt{-bc+ad}}\right)}{e^{3/2}f^2\sqrt{-be+af}(de-cf)^2} \right. \\ \left. - \frac{2\sqrt{b}C \log\left(-\sqrt{bx} + \sqrt{a+bx^2}\right)}{df^2} \right)$$

input

```
Integrate[(Sqrt[a + b*x^2]*(A + B*x^2 + C*x^4))/((c + d*x^2)*(e + f*x^2)^2),x]
```

output

```
(-(((C*e^2 + f*(-(B*e) + A*f))*x*Sqrt[a + b*x^2])/(e*f*(d*e - c*f)*(e + f*x^2))) - (2*Sqrt[-(b*c) + a*d]*(c^2*C - B*c*d + A*d^2)*ArcTan[(-(d*x*Sqrt[a + b*x^2]) + Sqrt[b]*(c + d*x^2))/(Sqrt[c]*Sqrt[-(b*c) + a*d])])/(Sqrt[c]*d*(d*e - c*f)^2) + (((2*b*e^2*((B*c - A*d)*f^2 + C*e*(d*e - 2*c*f)) - a*f*(C*e^2*(d*e - 3*c*f) + f*(A*f*(-3*d*e + c*f) + B*e*(d*e + c*f))))*ArcTan[(-(f*x*Sqrt[a + b*x^2]) + Sqrt[b]*(e + f*x^2))/(Sqrt[e]*Sqrt[-(b*e) + a*f])])/(e^(3/2)*f^2*Sqrt[-(b*e) + a*f]*(d*e - c*f)^2) - (2*Sqrt[b]*C*Log[-(Sqrt[b]*x) + Sqrt[a + b*x^2])]/(d*f^2))/2
```

Rubi [A] (verified)

Time = 1.17 (sec) , antiderivative size = 422, normalized size of antiderivative = 1.36, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)(e+fx^2)^2} dx$$

↓ 7276

$$\int \left(\frac{\sqrt{a+bx^2}(Ad^2 - Bcd + c^2C)}{(c+dx^2)(de-cf)^2} + \frac{\sqrt{a+bx^2}(Af^2 - Bef + Ce^2)}{f(e+fx^2)^2(cf-de)} + \frac{\sqrt{a+bx^2}(f^2(Bc - Ad) + Ce(de - 2cf))}{f(e+fx^2)(de-cf)^2} \right)$$

↓ 2009

$$\begin{aligned} & \frac{\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right) (Ad^2 - Bcd + c^2C)}{d(de - cf)^2} - \\ & \frac{\sqrt{bc - ad}(Ad^2 - Bcd + c^2C) \operatorname{arctanh}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{cd}(de - cf)^2} - \\ & \frac{a(Af^2 - Bef + Ce^2) \operatorname{arctanh}\left(\frac{x\sqrt{be-af}}{\sqrt{e}\sqrt{a+bx^2}}\right)}{2e^{3/2}f\sqrt{be - af}(de - cf)} + \\ & \frac{\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right) (f^2(Bc - Ad) + Ce(de - 2cf))}{f^2(de - cf)^2} - \\ & \frac{\sqrt{be - af} \operatorname{arctanh}\left(\frac{x\sqrt{be-af}}{\sqrt{e}\sqrt{a+bx^2}}\right) (f^2(Bc - Ad) + Ce(de - 2cf))}{\sqrt{e}f^2(de - cf)^2} - \\ & \frac{x\sqrt{a+bx^2}(Af^2 - Bef + Ce^2)}{2ef(e+fx^2)(de - cf)} \end{aligned}$$

input `Int[(Sqrt[a + b*x^2]*(A + B*x^2 + C*x^4))/((c + d*x^2)*(e + f*x^2)^2),x]`

output `-1/2*((C*e^2 - B*e*f + A*f^2)*x*Sqrt[a + b*x^2])/(e*f*(d*e - c*f)*(e + f*x^2)) + (Sqrt[b]*(c^2*C - B*c*d + A*d^2)*ArcTanh[(Sqrt[b]*x)/Sqrt[a + b*x^2]])/(d*(d*e - c*f)^2) + (Sqrt[b]*((B*c - A*d)*f^2 + C*e*(d*e - 2*c*f))*ArcTanh[(Sqrt[b]*x)/Sqrt[a + b*x^2]])/(f^2*(d*e - c*f)^2) - (Sqrt[b*c - a*d]*(c^2*C - B*c*d + A*d^2)*ArcTanh[(Sqrt[b*c - a*d]*x)/(Sqrt[c]*Sqrt[a + b*x^2])])/(Sqrt[c]*d*(d*e - c*f)^2) - (a*(C*e^2 - B*e*f + A*f^2)*ArcTanh[(Sqrt[b*e - a*f]*x)/(Sqrt[e]*Sqrt[a + b*x^2])])/(2*e^(3/2)*f*Sqrt[b*e - a*f]*(d*e - c*f)) - (Sqrt[b*e - a*f]*((B*c - A*d)*f^2 + C*e*(d*e - 2*c*f))*ArcTanh[(Sqrt[b*e - a*f]*x)/(Sqrt[e]*Sqrt[a + b*x^2])])/(Sqrt[e]*f^2*(d*e - c*f)^2)`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
x
pand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]`

Maple [A] (verified)

Time = 1.56 (sec) , antiderivative size = 348, normalized size of antiderivative = 1.12

method	result
pseudoelliptic	$\frac{-\sqrt{(ad-bc)c}(fx^2+e)\left(-2Cbd e^4+Cf(ad+4bc)e^3+2f^2\left(-Bb-\frac{3Ca}{2}\right)c+d\left(Ab+\frac{Ba}{2}\right)\right)e^2-3\left(Ad-\frac{Bc}{3}\right)f^3ae+Aac f^4}{d}$
default	Expression too large to display

input `int((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)/(f*x^2+e)^2,x,method=_RETURN
VERBOSE)`

output
$$\frac{1/2}{(a*d-b*c)*c}^{1/2} / ((a*f-b*e)*e)^{1/2} * (-((a*d-b*c)*c)^{1/2} * (f*x^2+e) * (-2*C*b*d*e^4+C*f*(a*d+4*b*c)*e^3+2*f^2*((-B*b-3/2*C*a)*c+d*(A*b+1/2*B*a)) * e^2-3*(A*d-1/3*B*c)*f^3*a*e+A*a*c*f^4) * d * \arctan(e*(b*x^2+a)^{1/2}/x / ((a*f-b*e)*e)^{1/2}) + (-2*e*f^2*(f*x^2+e)*(a*d-b*c)*(A*d^2-B*c*d+C*c^2) * \arctan(c*(b*x^2+a)^{1/2}/x / ((a*d-b*c)*c)^{1/2}) + ((a*d-b*c)*c)^{1/2} * (2*b^{1/2}*C * e*(f*x^2+e)*(c*f-d*e) * \operatorname{arctanh}((b*x^2+a)^{1/2}/x/b^{1/2}) + x*(b*x^2+a)^{1/2}) * d * f * (A*f^2-B*e*f+C*e^2) * (c*f-d*e) * ((a*f-b*e)*e)^{1/2} / d / (c*f-d*e)^2 / e / (f*x^2+e) / f^2$$

Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)(e+fx^2)^2} dx = \text{Timed out}$$

input `integrate((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)/(f*x^2+e)^2,x, algorithm="fricas")`

output Timed out

Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)(e+fx^2)^2} dx = \text{Timed out}$$

input `integrate((b*x**2+a)**(1/2)*(C*x**4+B*x**2+A)/(d*x**2+c)/(f*x**2+e)**2,x)`

output Timed out

Maxima [F]

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)(e+fx^2)^2} dx = \int \frac{(Cx^4+Bx^2+A)\sqrt{bx^2+a}}{(dx^2+c)(fx^2+e)^2} dx$$

input `integrate((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)/(f*x^2+e)^2,x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)*sqrt(b*x^2 + a)/((d*x^2 + c)*(f*x^2 + e)^2), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)(e+fx^2)^2} dx = \text{Exception raised: TypeError}$$

input `integrate((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)/(f*x^2+e)^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx);OUTPUT:index.cc index_m i_lex_is_greater E rror: Bad Argument Value`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)(e+fx^2)^2} dx = \int \frac{\sqrt{bx^2+a}(Cx^4+Bx^2+A)}{(dx^2+c)(fx^2+e)^2} dx$$

input `int(((a + b*x^2)^(1/2)*(A + B*x^2 + C*x^4))/((c + d*x^2)*(e + f*x^2)^2),x)`

output `int(((a + b*x^2)^(1/2)*(A + B*x^2 + C*x^4))/((c + d*x^2)*(e + f*x^2)^2), x)`

Reduce [B] (verification not implemented)

Time = 1.52 (sec) , antiderivative size = 4615, normalized size of antiderivative = 14.89

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)(e+fx^2)^2} dx = \text{Too large to display}$$

input `int((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)/(f*x^2+e)^2,x)`

output

```
( - 2*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x
**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**2*d**2*e**3*f**3 - 2*sqrt(
c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt
(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**2*d**2*e**2*f**4*x**2 + 2*sqrt(c)*sqr
t(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sq
rt(b)*x)/(sqrt(c)*sqrt(b)))*a*b*c*d*e**3*f**3 + 2*sqrt(c)*sqrt(a*d - b*c)*
atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqr
t(c)*sqrt(b)))*a*b*c*d*e**2*f**4*x**2 + 2*sqrt(c)*sqrt(a*d - b*c)*atan((sq
rt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqr
t(b)))*a*b*d**2*e**4*f**2 + 2*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c
) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a*b*d
**2*e**3*f**3*x**2 - 2*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqr
t(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a*c**3*e**3*
f**3 - 2*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a +
b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a*c**3*e**2*f**4*x**2 - 2*
sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) -
sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*b**2*c*d*e**4*f**2 - 2*sqrt(c)*sqrt
(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqr
t(b)*x)/(sqrt(c)*sqrt(b)))*b**2*c*d*e**3*f**3*x**2 + 2*sqrt(c)*sqrt(a*d -
b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)...
```

3.6 $\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}(c+dx^2)(e+fx^2)^2} dx$

Optimal result	84
Mathematica [A] (verified)	85
Rubi [A] (verified)	85
Maple [A] (verified)	87
Fricas [F(-1)]	87
Sympy [F(-1)]	88
Maxima [F]	88
Giac [B] (verification not implemented)	88
Mupad [F(-1)]	90
Reduce [B] (verification not implemented)	90

Optimal result

Integrand size = 42, antiderivative size = 278

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}(c + dx^2)(e + fx^2)^2} dx$$

$$= \frac{(Ce^2 - Bef + Af^2)x\sqrt{a + bx^2}}{2e(be - af)(de - cf)(e + fx^2)} + \frac{(c^2C - Bcd + Ad^2) \operatorname{arctanh}\left(\frac{\sqrt{bc - ad}x}{\sqrt{c}\sqrt{a + bx^2}}\right)}{\sqrt{c}\sqrt{bc - ad}(de - cf)^2}$$

$$+ \frac{(2be(de(Be - 2Af) - c(Ce^2 - Af^2)) - a(Ce^2(de - 3cf) - f(Af(3de - cf) - Be(de + cf)))) \operatorname{arctan}}{2e^{3/2}(be - af)^{3/2}(de - cf)^2}$$

output

```
1/2*(A*f^2-B*e*f+C*e^2)*x*(b*x^2+a)^(1/2)/e/(-a*f+b*e)/(-c*f+d*e)/(f*x^2+e
)+(A*d^2-B*c*d+C*c^2)*arctanh((-a*d+b*c)^(1/2)*x/c^(1/2)/(b*x^2+a)^(1/2))/
c^(1/2)/(-a*d+b*c)^(1/2)/(-c*f+d*e)^2+1/2*(2*b*e*(d*e*(-2*A*f+B*e)-c*(-A*f
^2+C*e^2))-a*(C*e^2*(-3*c*f+d*e)-f*(A*f*(-c*f+3*d*e)-B*e*(c*f+d*e))))*arct
anh((-a*f+b*e)^(1/2)*x/e^(1/2)/(b*x^2+a)^(1/2))/e^(3/2)/(-a*f+b*e)^(3/2)/(
-c*f+d*e)^2
```

Mathematica [A] (verified)

Time = 2.46 (sec) , antiderivative size = 295, normalized size of antiderivative = 1.06

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c + dx^2) (e + fx^2)^2} dx$$

$$= \frac{(de - cf)(Ce^2 + f(-Be + Af))x\sqrt{a + bx^2}}{e(be - af)(e + fx^2)} - \frac{2(c^2C - Bcd + Ad^2) \arctan\left(\frac{-dx\sqrt{a + bx^2} + \sqrt{b}(c + dx^2)}{\sqrt{c}\sqrt{-bc + ad}}\right)}{\sqrt{c}\sqrt{-bc + ad}} - \frac{(-2be(de(Be - 2Af) + c(-Ce^2 + Af^2)))}{2(de - cf)^2}$$

input

```
Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*(c + d*x^2)*(e + f*x^2)^2), x]
```

output

```
((d*e - c*f)*(C*e^2 + f*(-(B*e) + A*f))*x*Sqrt[a + b*x^2])/(e*(b*e - a*f)*(e + f*x^2)) - (2*(c^2*C - B*c*d + A*d^2)*ArcTan[(-(d*x*Sqrt[a + b*x^2]) + Sqrt[b]*(c + d*x^2))/(Sqrt[c]*Sqrt[-(b*c) + a*d])]/(Sqrt[c]*Sqrt[-(b*c) + a*d]) - ((-2*b*e*(d*e*(B*e - 2*A*f) + c*(-(C*e^2) + A*f^2)) + a*(C*e^2*(d*e - 3*c*f) + f*(A*f*(-3*d*e + c*f) + B*e*(d*e + c*f))))*ArcTan[(-(f*x*Sqrt[a + b*x^2]) + Sqrt[b]*(e + f*x^2))/(Sqrt[e]*Sqrt[-(b*e) + a*f])]/(e^(3/2)*(-(b*e) + a*f)^(3/2)))/(2*(d*e - c*f)^2)
```

Rubi [A] (verified)

Time = 1.12 (sec) , antiderivative size = 316, normalized size of antiderivative = 1.14, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c + dx^2) (e + fx^2)^2} dx$$

↓ 7276

$$\int \left(\frac{Ad^2 - Bcd + c^2C}{\sqrt{a + bx^2} (c + dx^2) (de - cf)^2} + \frac{Af^2 - Bef + Ce^2}{f\sqrt{a + bx^2} (e + fx^2)^2 (cf - de)} + \frac{f^2(Bc - Ad) + Ce(de - 2cf)}{f\sqrt{a + bx^2} (e + fx^2) (de - cf)^2} \right) dx$$

↓ 2009

$$\frac{(Ad^2 - Bcd + c^2C) \operatorname{arctanh}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}\sqrt{bc-ad}(de-cf)^2} - \frac{(2be-af)(Af^2 - Bef + Ce^2) \operatorname{arctanh}\left(\frac{x\sqrt{be-af}}{\sqrt{e}\sqrt{a+bx^2}}\right)}{2e^{3/2}f(be-af)^{3/2}(de-cf)} + \frac{\operatorname{arctanh}\left(\frac{x\sqrt{be-af}}{\sqrt{e}\sqrt{a+bx^2}}\right)(f^2(Bc-Ad) + Ce(de-2cf))}{\sqrt{e}f\sqrt{be-af}(de-cf)^2} + \frac{x\sqrt{a+bx^2}(Af^2 - Bef + Ce^2)}{2e(e+fx^2)(be-af)(de-cf)}$$

input `Int[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*(c + d*x^2)*(e + f*x^2)^2),x]`

output `((C*e^2 - B*e*f + A*f^2)*x*Sqrt[a + b*x^2])/(2*e*(b*e - a*f)*(d*e - c*f)*(e + f*x^2)) + ((c^2*C - B*c*d + A*d^2)*ArcTanh[(Sqrt[b*c - a*d]*x)/(Sqrt[c]*Sqrt[a + b*x^2])])/(Sqrt[c]*Sqrt[b*c - a*d]*(d*e - c*f)^2) - ((2*b*e - a*f)*(C*e^2 - B*e*f + A*f^2)*ArcTanh[(Sqrt[b*e - a*f]*x)/(Sqrt[e]*Sqrt[a + b*x^2])])/(2*e^(3/2)*f*(b*e - a*f)^(3/2)*(d*e - c*f)) + ((B*c - A*d)*f^2 + C*e*(d*e - 2*c*f))*ArcTanh[(Sqrt[b*e - a*f]*x)/(Sqrt[e]*Sqrt[a + b*x^2])]/(Sqrt[e]*f*Sqrt[b*e - a*f]*(d*e - c*f)^2)`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 1.52 (sec) , antiderivative size = 302, normalized size of antiderivative = 1.09

method	result
pseudoelliptic	$\frac{-(f x^2+e) \sqrt{(a d-b c) c} \left((-2 B b d+C a d+2 C b c) e^3+4 f \left(-\frac{3 C a c}{4}+d \left(A b+\frac{B a}{4} \right) \right) e^2-3 f^2 \left(\frac{(2 A b-B a) c}{3}+A a d \right) e+A a c f^3 \right) \arctan \left(\frac{e \sqrt{(b x^2+a)} / x}{\sqrt{(a f-b e) e}} \right)}{2 \sqrt{(a d-b c) c}}$
default	Expression too large to display

input `int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)/(f*x^2+e)^2,x,method=_RETURN
VERBOSE)`

output `1/2/((a*d-b*c)*c)^(1/2)*(-(f*x^2+e)*((a*d-b*c)*c)^(1/2)*((-2*B*b*d+C*a*d+2
*C*b*c)*e^3+4*f*(-3/4*C*a*c+d*(A*b+1/4*B*a))*e^2-3*f^2*(1/3*(2*A*b-B*a)*c+
A*a*d)*e+A*a*c*f^3)*arctan(e*(b*x^2+a)^(1/2)/x/((a*f-b*e)*e)^(1/2))+(-2*e*
(f*x^2+e)*(a*f-b*e)*(A*d^2-B*c*d+C*c^2)*arctan(c*(b*x^2+a)^(1/2)/x/((a*d-b
*c)*c)^(1/2))+x*((a*d-b*c)*c)^(1/2)*(A*f^2-B*e*f+C*e^2)*(c*f-d*e)*(b*x^2+a
)^(1/2))*((a*f-b*e)*e)^(1/2))/((a*f-b*e)*e)^(1/2)/(c*f-d*e)^2/(a*f-b*e)/e/
(f*x^2+e)`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c + dx^2) (e + fx^2)^2} dx = \text{Timed out}$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)/(f*x^2+e)^2,x, algorit
hm="fricas")`

output `Timed out`

Sympy [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c + dx^2) (e + fx^2)^2} dx = \text{Timed out}$$

input `integrate((C*x**4+B*x**2+A)/(b*x**2+a)**(1/2)/(d*x**2+c)/(f*x**2+e)**2,x)`

output `Timed out`

Maxima [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c + dx^2) (e + fx^2)^2} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}(dx^2 + c)(fx^2 + e)^2} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)/(f*x^2+e)^2,x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(b*x^2 + a)*(d*x^2 + c)*(f*x^2 + e)^2), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 678 vs. $2(251) = 502$.

Time = 0.86 (sec) , antiderivative size = 678, normalized size of antiderivative = 2.44

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c + dx^2) (e + fx^2)^2} dx$$

$$= - \frac{\left(C\sqrt{bc^2} - B\sqrt{bcd} + A\sqrt{bd^2} \right) \arctan \left(\frac{(\sqrt{bx - \sqrt{bx^2 + a}})^2 d + 2bc - ad}{2\sqrt{-b^2c^2 + abcd}} \right)}{\sqrt{-b^2c^2 + abcd}(d^2e^2 - 2cdef + c^2f^2)}$$

$$+ \frac{\left(2Cb^{\frac{3}{2}}ce^3 + Ca\sqrt{bde^3} - 2Bb^{\frac{3}{2}}de^3 - 3Ca\sqrt{bce^2f} + Ba\sqrt{bde^2f} + 4Ab^{\frac{3}{2}}de^2f + Ba\sqrt{bce^2f} - 2Ab^{\frac{3}{2}}ce^2 \right)}{2(bd^2e^4 - 2bcde^3f - ad^2e^3f + bc^2e^2f^2 + 2acde^2f^2 - ac^2e^2f^2)}$$

$$+ \frac{2(\sqrt{bx} - \sqrt{bx^2 + a})^2 Cb^{\frac{3}{2}}e^3 - (\sqrt{bx} - \sqrt{bx^2 + a})^2 Ca\sqrt{be^2f} - 2(\sqrt{bx} - \sqrt{bx^2 + a})^2 Bb^{\frac{3}{2}}e^2f + (\sqrt{bx} - \sqrt{bx^2 + a})^2 Aa\sqrt{bf^3}}{(bde^3f - bce^2f^2 - ade^2f^2 + acef^3) \left((\sqrt{bx} - \sqrt{bx^2 + a})^2 \right)}$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)/(f*x^2+e)^2,x, algorithm="giac")`

output `-(C*sqrt(b)*c^2 - B*sqrt(b)*c*d + A*sqrt(b)*d^2)*arctan(1/2*((sqrt(b)*x - sqrt(b*x^2 + a))^2*d + 2*b*c - a*d)/sqrt(-b^2*c^2 + a*b*c*d))/(sqrt(-b^2*c^2 + a*b*c*d)*(d^2*e^2 - 2*c*d*e*f + c^2*f^2)) + 1/2*(2*C*b^(3/2)*c*e^3 + C*a*sqrt(b)*d*e^3 - 2*B*b^(3/2)*d*e^3 - 3*C*a*sqrt(b)*c*e^2*f + B*a*sqrt(b)*d*e^2*f + 4*A*b^(3/2)*d*e^2*f + B*a*sqrt(b)*c*e*f^2 - 2*A*b^(3/2)*c*e*f^2 - 3*A*a*sqrt(b)*d*e*f^2 + A*a*sqrt(b)*c*f^3)*arctan(1/2*((sqrt(b)*x - sqrt(b*x^2 + a))^2*f + 2*b*e - a*f)/sqrt(-b^2*e^2 + a*b*e*f))/((b*d^2*e^4 - 2*b*c*d*e^3*f - a*d^2*e^3*f + b*c^2*e^2*f^2 + 2*a*c*d*e^2*f^2 - a*c^2*e*f^3)*sqrt(-b^2*e^2 + a*b*e*f)) + (2*(sqrt(b)*x - sqrt(b*x^2 + a))^2*C*b^(3/2)*e^3 - (sqrt(b)*x - sqrt(b*x^2 + a))^2*C*a*sqrt(b)*e^2*f - 2*(sqrt(b)*x - sqrt(b*x^2 + a))^2*B*b^(3/2)*e^2*f + (sqrt(b)*x - sqrt(b*x^2 + a))^2*B*a*sqrt(b)*e*f^2 + 2*(sqrt(b)*x - sqrt(b*x^2 + a))^2*A*b^(3/2)*e*f^2 - (sqrt(b)*x - sqrt(b*x^2 + a))^2*A*a*sqrt(b)*f^3 + C*a^2*sqrt(b)*e^2*f - B*a^2*sqrt(b)*e*f^2 + A*a^2*sqrt(b)*f^3)/((b*d*e^3*f - b*c*e^2*f^2 - a*d*e^2*f^2 + a*c*e*f^3)*((sqrt(b)*x - sqrt(b*x^2 + a))^4*f + 4*(sqrt(b)*x - sqrt(b*x^2 + a))^2*b*e - 2*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a*f + a^2*f))`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c + dx^2) (e + fx^2)^2} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a} (dx^2 + c) (fx^2 + e)^2} dx$$

input `int((A + B*x^2 + C*x^4)/((a + b*x^2)^(1/2)*(c + d*x^2)*(e + f*x^2)^2),x)`

output `int((A + B*x^2 + C*x^4)/((a + b*x^2)^(1/2)*(c + d*x^2)*(e + f*x^2)^2), x)`

Reduce [B] (verification not implemented)

Time = 1.39 (sec) , antiderivative size = 6592, normalized size of antiderivative = 23.71

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c + dx^2) (e + fx^2)^2} dx = \text{Too large to display}$$

input `int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)/(f*x^2+e)^2,x)`

output

```
( - 2*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x
**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**3*d**2*e**3*f**2 - 2*sqrt(
c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt
(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**3*d**2*e**2*f**3*x**2 + 2*sqrt(c)*sqr
t(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sq
rt(b)*x)/(sqrt(c)*sqrt(b)))*a**2*b*c*d*e**3*f**2 + 2*sqrt(c)*sqrt(a*d - b*
c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(
sqrt(c)*sqrt(b)))*a**2*b*c*d*e**2*f**3*x**2 + 4*sqrt(c)*sqrt(a*d - b*c)*at
an((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(
c)*sqrt(b)))*a**2*b*d**2*e**4*f + 4*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d
- b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b))
)*a**2*b*d**2*e**3*f**2*x**2 - 2*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b
*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**
2*c**3*e**3*f**2 - 2*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(
d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**2*c**3*e**2
*f**3*x**2 - 4*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqr
t(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a*b**2*c*d*e**4*f -
4*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2)
- sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a*b**2*c*d*e**3*f**2*x**2 - 2*sqr
t(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - ...
```

3.7
$$\int \frac{A+Bx^2+Cx^4}{(a+bx^2)^{3/2}(c+dx^2)(e+fx^2)} dx$$

Optimal result	92
Mathematica [A] (verified)	93
Rubi [A] (verified)	93
Maple [A] (verified)	95
Fricas [F(-1)]	95
Sympy [F(-1)]	96
Maxima [F]	96
Giac [A] (verification not implemented)	96
Mupad [F(-1)]	97
Reduce [B] (verification not implemented)	97

Optimal result

Integrand size = 42, antiderivative size = 205

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)} dx = \frac{(Ab^2 - a(bB - aC)) x}{a(bc - ad)(be - af)\sqrt{a + bx^2}}$$

$$- \frac{(c^2C - Bcd + Ad^2) \operatorname{arctanh}\left(\frac{\sqrt{bc-ad}x}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}(bc - ad)^{3/2}(de - cf)}$$

$$+ \frac{(Ce^2 - Bef + Af^2) \operatorname{arctanh}\left(\frac{\sqrt{be-af}x}{\sqrt{e}\sqrt{a+bx^2}}\right)}{\sqrt{e}(be - af)^{3/2}(de - cf)}$$

output

```
(A*b^2-a*(B*b-C*a))*x/a/(-a*d+b*c)/(-a*f+b*e)/(b*x^2+a)^(1/2)-(A*d^2-B*c*d
+C*c^2)*arctanh((-a*d+b*c)^(1/2)*x/c^(1/2)/(b*x^2+a)^(1/2))/c^(1/2)/(-a*d+
b*c)^(3/2)/(-c*f+d*e)+(A*f^2-B*e*f+C*e^2)*arctanh((-a*f+b*e)^(1/2)*x/e^(1/
2)/(b*x^2+a)^(1/2))/e^(1/2)/(-a*f+b*e)^(3/2)/(-c*f+d*e)
```

Mathematica [A] (verified)

Time = 1.55 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.16

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)} dx = \frac{(Ab^2 + a(-bB + aC))x}{a(-bc + ad)(-be + af)\sqrt{a + bx^2}} - \frac{(c^2C - Bcd + Ad^2) \arctan\left(\frac{-dx\sqrt{a+bx^2} + \sqrt{b}(c+dx^2)}{\sqrt{c}\sqrt{-bc+ad}}\right)}{\sqrt{c}(-bc + ad)^{3/2}(de - cf)} + \frac{(Ce^2 + f(-Be + Af)) \arctan\left(\frac{-fx\sqrt{a+bx^2} + \sqrt{b}(e+fx^2)}{\sqrt{e}\sqrt{-be+af}}\right)}{\sqrt{e}(-be + af)^{3/2}(de - cf)}$$

input

```
Integrate[(A + B*x^2 + C*x^4)/((a + b*x^2)^(3/2)*(c + d*x^2)*(e + f*x^2)), x]
```

output

```
((A*b^2 + a*(-b*B) + a*C)*x)/(a*(-b*c) + a*d)*(-b*e) + a*f)*Sqrt[a + b*x^2]) - ((c^2*C - B*c*d + A*d^2)*ArcTan[(-d*x*Sqrt[a + b*x^2]) + Sqrt[b]*(c + d*x^2)]/(Sqrt[c]*Sqrt[-(b*c) + a*d]))/(Sqrt[c]*(-b*c) + a*d)^(3/2)*(d*e - c*f)) + ((C*e^2 + f*(-B*e) + A*f)*ArcTan[(-f*x*Sqrt[a + b*x^2]) + Sqrt[b]*(e + f*x^2)]/(Sqrt[e]*Sqrt[-(b*e) + a*f]))/(Sqrt[e]*(-b*e) + a*f)^(3/2)*(d*e - c*f))
```

Rubi [A] (verified)

Time = 1.05 (sec) , antiderivative size = 288, normalized size of antiderivative = 1.40, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)} dx$$

↓ 7276

$$\int \left(\frac{Ad^2 - Bcd + c^2C}{d(a + bx^2)^{3/2}(c + dx^2)(de - cf)} + \frac{Af^2 - Bef + Ce^2}{f(a + bx^2)^{3/2}(e + fx^2)(cf - de)} + \frac{C}{df(a + bx^2)^{3/2}} \right) dx$$

↓ 2009

$$-\frac{(Ad^2 - Bcd + c^2C) \operatorname{arctanh}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}(bc - ad)^{3/2}(de - cf)} + \frac{(Af^2 - Bef + Ce^2) \operatorname{arctanh}\left(\frac{x\sqrt{be-af}}{\sqrt{e}\sqrt{a+bx^2}}\right)}{\sqrt{e}(be - af)^{3/2}(de - cf)} +$$

$$\frac{bx(Ad^2 - Bcd + c^2C)}{ad\sqrt{a + bx^2}(bc - ad)(de - cf)} - \frac{bx(Af^2 - Bef + Ce^2)}{af\sqrt{a + bx^2}(be - af)(de - cf)} + \frac{Cx}{adf\sqrt{a + bx^2}}$$

input `Int[(A + B*x^2 + C*x^4)/((a + b*x^2)^(3/2)*(c + d*x^2)*(e + f*x^2)),x]`

output `(C*x)/(a*d*f*Sqrt[a + b*x^2]) + (b*(c^2*C - B*c*d + A*d^2)*x)/(a*d*(b*c - a*d)*(d*e - c*f)*Sqrt[a + b*x^2]) - (b*(C*e^2 - B*e*f + A*f^2)*x)/(a*f*(b*e - a*f)*(d*e - c*f)*Sqrt[a + b*x^2]) - ((c^2*C - B*c*d + A*d^2)*ArcTanh[(Sqrt[b*c - a*d]*x)/(Sqrt[c]*Sqrt[a + b*x^2])])/(Sqrt[c]*(b*c - a*d)^(3/2)*(d*e - c*f)) + ((C*e^2 - B*e*f + A*f^2)*ArcTanh[(Sqrt[b*e - a*f]*x)/(Sqrt[e]*Sqrt[a + b*x^2])])/(Sqrt[e]*(b*e - a*f)^(3/2)*(d*e - c*f))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 1.36 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.01

method	result	size
pseudoelliptic	$\frac{(d^2 A - cdB + C c^2) a \arctan\left(\frac{c\sqrt{bx^2+a}}{x\sqrt{(ad-bc)c}}\right)}{(cf-de)(ad-bc)\sqrt{(ad-bc)c}} + \frac{(b^2 A - abB + a^2 C) x}{(af-be)(ad-bc)\sqrt{bx^2+a}} - \frac{(A f^2 - B e f + C e^2) a \arctan\left(\frac{e\sqrt{bx^2+a}}{x\sqrt{(af-be)e}}\right)}{(cf-de)(af-be)\sqrt{(af-be)e}}$	208
default	Expression too large to display	1689

input `int((C*x^4+B*x^2+A)/(b*x^2+a)^(3/2)/(d*x^2+c)/(f*x^2+e),x,method=_RETURNVE
RBOSE)`

output `((A*d^2-B*c*d+C*c^2)*a/(c*f-d*e)/(a*d-b*c)/((a*d-b*c)*c)^(1/2)*arctan(c*(b
*x^2+a)^(1/2)/x/((a*d-b*c)*c)^(1/2))+ (A*b^2-B*a*b+C*a^2)/(a*f-b*e)/(a*d-b*
c)*x/(b*x^2+a)^(1/2)- (A*f^2-B*e*f+C*e^2)*a/(c*f-d*e)/(a*f-b*e)/((a*f-b*e)*
e)^(1/2)*arctan(e*(b*x^2+a)^(1/2)/x/((a*f-b*e)*e)^(1/2))/a`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)} dx = \text{Timed out}$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(3/2)/(d*x^2+c)/(f*x^2+e),x, algorithm
="fricas")`

output `Timed out`

Sympy [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)} dx = \text{Timed out}$$

input `integrate((C*x**4+B*x**2+A)/(b*x**2+a)**(3/2)/(d*x**2+c)/(f*x**2+e),x)`

output `Timed out`

Maxima [F]

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)} dx = \int \frac{Cx^4 + Bx^2 + A}{(bx^2 + a)^{\frac{3}{2}} (dx^2 + c)(fx^2 + e)} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(3/2)/(d*x^2+c)/(f*x^2+e),x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)/((b*x^2 + a)^(3/2)*(d*x^2 + c)*(f*x^2 + e)), x)`

Giac [A] (verification not implemented)

Time = 0.75 (sec) , antiderivative size = 297, normalized size of antiderivative = 1.45

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)} dx = \frac{(Ca^2 - Bab + Ab^2)x}{(ab^2ce - a^2bde - a^2bcf + a^3df)\sqrt{bx^2 + a}}$$

$$+ \frac{(C\sqrt{bc^2} - B\sqrt{bcd} + A\sqrt{bd^2}) \arctan\left(\frac{(\sqrt{bx} - \sqrt{bx^2 + a})^2 d + 2bc - ad}{2\sqrt{-b^2c^2 + abcd}}\right)}{\sqrt{-b^2c^2 + abcd}(bcde - ad^2e - bc^2f + acdf)}$$

$$- \frac{(C\sqrt{be^2} - B\sqrt{bef} + A\sqrt{bf^2}) \arctan\left(\frac{(\sqrt{bx} - \sqrt{bx^2 + a})^2 f + 2be - af}{2\sqrt{-b^2e^2 + abef}}\right)}{\sqrt{-b^2e^2 + abef}(bde^2 - bcef - adef + acf^2)}$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(3/2)/(d*x^2+c)/(f*x^2+e),x, algorithm="giac")`

output `(C*a^2 - B*a*b + A*b^2)*x/((a*b^2*c*e - a^2*b*d*e - a^2*b*c*f + a^3*d*f)*sqrt(b*x^2 + a)) + (C*sqrt(b)*c^2 - B*sqrt(b)*c*d + A*sqrt(b)*d^2)*arctan(1/2*((sqrt(b)*x - sqrt(b*x^2 + a))^2*d + 2*b*c - a*d)/sqrt(-b^2*c^2 + a*b*c*d))/(sqrt(-b^2*c^2 + a*b*c*d)*(b*c*d*e - a*d^2*e - b*c^2*f + a*c*d*f)) - (C*sqrt(b)*e^2 - B*sqrt(b)*e*f + A*sqrt(b)*f^2)*arctan(1/2*((sqrt(b)*x - sqrt(b*x^2 + a))^2*f + 2*b*e - a*f)/sqrt(-b^2*e^2 + a*b*e*f))/(sqrt(-b^2*e^2 + a*b*e*f)*(b*d*e^2 - b*c*e*f - a*d*e*f + a*c*f^2))`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)} dx = \int \frac{Cx^4 + Bx^2 + A}{(bx^2 + a)^{3/2} (dx^2 + c) (fx^2 + e)} dx$$

input `int((A + B*x^2 + C*x^4)/((a + b*x^2)^(3/2)*(c + d*x^2)*(e + f*x^2)),x)`

output `int((A + B*x^2 + C*x^4)/((a + b*x^2)^(3/2)*(c + d*x^2)*(e + f*x^2)), x)`

Reduce [B] (verification not implemented)

Time = 1.38 (sec) , antiderivative size = 5297, normalized size of antiderivative = 25.84

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)^{3/2} (c + dx^2) (e + fx^2)} dx = \text{Too large to display}$$

input `int((C*x^4+B*x^2+A)/(b*x^2+a)^(3/2)/(d*x^2+c)/(f*x^2+e),x)`

output

```

(sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2)
- sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**4*b*d**2*e*f**2 - sqrt(c)*sqrt(
a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt
(b)*x)/(sqrt(c)*sqrt(b)))*a**3*b**2*c*d*e*f**2 - 2*sqrt(c)*sqrt(a*d - b*c)
*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sq
rt(c)*sqrt(b)))*a**3*b**2*d**2*e**2*f + sqrt(c)*sqrt(a*d - b*c)*atan((sqrt
(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(
b)))*a**3*b**2*d**2*e*f**2*x**2 + sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d -
b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a
**3*b*c**3*e*f**2 + 2*sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt
(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**2*b**3*c*d
*e**2*f - sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a +
b*x**2) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**2*b**3*c*d*e*f**2*x**2
+ sqrt(c)*sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2)
) - sqrt(d)*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**2*b**3*d**2*e**3 - 2*sqrt(c)*
sqrt(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)
*sqrt(b)*x)/(sqrt(c)*sqrt(b)))*a**2*b**3*d**2*e**2*f*x**2 - 2*sqrt(c)*sqrt
(a*d - b*c)*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqr
t(b)*x)/(sqrt(c)*sqrt(b)))*a**2*b**2*c**3*e**2*f + sqrt(c)*sqrt(a*d - b*c)
*atan((sqrt(a*d - b*c) - sqrt(d)*sqrt(a + b*x**2) - sqrt(d)*sqrt(b)*x)/...

```

$$3.8 \quad \int \frac{\sqrt{a+bx^2}(c+dx^2)^{3/2}(A+Bx^2+Cx^4)}{e+fx^2} dx$$

Optimal result	99
Mathematica [C] (verified)	100
Rubi [A] (verified)	101
Maple [A] (verified)	103
Fricas [F(-1)]	104
Sympy [F]	105
Maxima [F]	105
Giac [F]	105
Mupad [F(-1)]	106
Reduce [F]	106

Optimal result

Integrand size = 44, antiderivative size = 1070

$$\int \frac{\sqrt{a+bx^2}(c+dx^2)^{3/2}(A+Bx^2+Cx^4)}{e+fx^2} dx = \text{Too large to display}$$

output

```

1/105*(8*a^3*C*d^3*f^3+a^2*b*d^2*f^2*(-14*B*d*f-19*C*c*f+14*C*d*e)-a*b^2*d
*f*(7*d*f*(-5*A*d*f-7*B*c*f+5*B*d*e)-C*(9*c^2*f^2-49*c*d*e*f+35*d^2*e^2))-
b^3*(C*(6*c^3*f^3+21*c^2*d*e*f^2-140*c*d^2*e^2*f+105*d^3*e^3)+7*d*f*(5*A*d
*f*(-4*c*f+3*d*e)-B*(3*c^2*f^2-20*c*d*e*f+15*d^2*e^2))))*x*(d*x^2+c)^(1/2)
/b^2/d^2/f^4/(b*x^2+a)^(1/2)-1/105*(4*a^2*C*d^2*f^2+a*b*d*f*(-7*B*d*f-9*C*
*c*f+7*C*d*e)+b^2*(7*d*f*(-5*A*d*f-6*B*c*f+5*B*d*e)-C*(3*c^2*f^2-42*c*d*e*f
+35*d^2*e^2))))*x*(b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)/b^2/d/f^3+1/35*(a*C*d*f-b
*(-7*B*d*f-8*C*c*f+7*C*d*e))*x^3*(b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)/b/f^2+1/7
*C*d*x^5*(b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)/f-1/105*a^(1/2)*(8*a^3*C*d^3*f^3+
a^2*b*d^2*f^2*(-14*B*d*f-19*C*c*f+14*C*d*e)-a*b^2*d*f*(7*d*f*(-5*A*d*f-7*B
*c*f+5*B*d*e)-C*(9*c^2*f^2-49*c*d*e*f+35*d^2*e^2))-b^3*(C*(6*c^3*f^3+21*c^
2*d*e*f^2-140*c*d^2*e^2*f+105*d^3*e^3)+7*d*f*(5*A*d*f*(-4*c*f+3*d*e)-B*(3*
c^2*f^2-20*c*d*e*f+15*d^2*e^2))))*(d*x^2+c)^(1/2)*EllipticE(b^(1/2)*x/a^(1
/2)/(1+b*x^2/a)^(1/2),(1-a*d/b/c)^(1/2))/b^(5/2)/d^2/f^4/(b*x^2+a)^(1/2)/(
a*(d*x^2+c)/c/(b*x^2+a)^(1/2)+1/105*a^(3/2)*(4*a^2*c*C*d^2*f^3+a*b*c*d*f^
2*(-7*B*d*f-9*C*c*f+7*C*d*e)-b^2*(C*(3*c^3*f^3+63*c^2*d*e*f^2-175*c*d^2*e^
2*f+105*d^3*e^3)+7*d*f*(5*A*d*f*(-5*c*f+3*d*e)-B*(9*c^2*f^2-25*c*d*e*f+15*
d^2*e^2))))*(d*x^2+c)^(1/2)*InverseJacobiAM(arctan(b^(1/2)*x/a^(1/2)),(1-a
*d/b/c)^(1/2))/b^(5/2)/c/d/f^4/(b*x^2+a)^(1/2)/(a*(d*x^2+c)/c/(b*x^2+a)^(
1/2)+a^(3/2)*(-c*f+d*e)^2*(A*f^2-B*e*f+C*e^2)*(d*x^2+c)^(1/2)*EllipticP...

```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 13.16 (sec) , antiderivative size = 6720, normalized size of antiderivative = 6.28

$$\int \frac{\sqrt{a+bx^2}(c+dx^2)^{3/2}(A+Bx^2+Cx^4)}{e+fx^2} dx = \text{Result too large to show}$$

input

```
Integrate[(Sqrt[a + b*x^2]*(c + d*x^2)^(3/2)*(A + B*x^2 + C*x^4))/(e + f*x
^2),x]
```

output

Result too large to show

Rubi [A] (verified)

Time = 2.45 (sec) , antiderivative size = 1321, normalized size of antiderivative = 1.23, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.045$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a+bx^2}(c+dx^2)^{3/2}(A+Bx^2+Cx^4)}{e+fx^2} dx$$

↓ 7276

$$\int \left(\frac{\sqrt{a+bx^2}(c+dx^2)^{3/2}(Af^2 - Bef + Ce^2)}{f^2(e+fx^2)} - \frac{\sqrt{a+bx^2}(c+dx^2)^{3/2}(Ce - Bf)}{f^2} + \frac{Cx^2\sqrt{a+bx^2}(c+dx^2)^3}{f} \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{C\sqrt{bx^2+a}(dx^2+c)^{3/2}x^3}{7f} + \frac{C(3bc+ad)\sqrt{bx^2+a}\sqrt{dx^2+cx}^3}{35bf} - \\
& \frac{d(Ce-Bf)(bx^2+a)^{3/2}\sqrt{dx^2+cx}}{5bf^2} - \frac{2(3bc-ad)(Ce-Bf)\sqrt{bx^2+a}\sqrt{dx^2+cx}}{15bf^2} + \\
& \frac{d(Ce^2-Bfe+Af^2)\sqrt{bx^2+a}\sqrt{dx^2+cx}}{3f^3} + \frac{C\left(-\frac{4da^2}{b}+9ca+\frac{3bc^2}{d}\right)\sqrt{bx^2+a}\sqrt{dx^2+cx}}{105bf} - \\
& \frac{(3b^2c^2+7abdc-2a^2d^2)(Ce-Bf)\sqrt{bx^2+ax}}{15b^2f^2\sqrt{dx^2+c}} - \\
& \frac{d(3bde-4bcf-adf)(Ce^2-Bfe+Af^2)\sqrt{bx^2+ax}}{3bf^4\sqrt{dx^2+c}} - \\
& \frac{C(2bc-ad)(3b^2c^2-3abdc+8a^2d^2)\sqrt{bx^2+ax}}{105b^3df\sqrt{dx^2+c}} + \\
& \frac{\sqrt{c}(3b^2c^2+7abdc-2a^2d^2)(Ce-Bf)\sqrt{bx^2+ax}E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\left|1-\frac{bc}{ad}\right.\right)}{15b^2\sqrt{d}f^2\sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}}\sqrt{dx^2+c}} + \\
& \frac{\sqrt{c}\sqrt{d}(3bde-4bcf-adf)(Ce^2-Bfe+Af^2)\sqrt{bx^2+ax}E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\left|1-\frac{bc}{ad}\right.\right)}{3bf^4\sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}}\sqrt{dx^2+c}} + \\
& \frac{\sqrt{c}C(2bc-ad)(3b^2c^2-3abdc+8a^2d^2)\sqrt{bx^2+ax}E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\left|1-\frac{bc}{ad}\right.\right)}{105b^3d^{3/2}f\sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}}\sqrt{dx^2+c}} - \\
& \frac{c^{3/2}(9bc-ad)(Ce-Bf)\sqrt{bx^2+ax}\text{EllipticF}\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right),1-\frac{bc}{ad}\right)}{15b\sqrt{d}f^2\sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}}\sqrt{dx^2+c}} - \\
& \frac{\sqrt{c}\sqrt{d}(3de-5cf)(Ce^2-Bfe+Af^2)\sqrt{bx^2+ax}\text{EllipticF}\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right),1-\frac{bc}{ad}\right)}{3f^4\sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}}\sqrt{dx^2+c}} - \\
& \frac{c^{3/2}C(3b^2c^2+9abdc-4a^2d^2)\sqrt{bx^2+ax}\text{EllipticF}\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right),1-\frac{bc}{ad}\right)}{105b^2d^{3/2}f\sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}}\sqrt{dx^2+c}} + \\
& \frac{a^{3/2}(de-cf)^2(Ce^2-Bfe+Af^2)\sqrt{dx^2+c}\text{EllipticPi}\left(1-\frac{af}{be},\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right),1-\frac{ad}{bc}\right)}{\sqrt{bce}f^4\sqrt{bx^2+a}\sqrt{\frac{a(dx^2+c)}{c(bx^2+a)}}}
\end{aligned}$$

input `Int[(Sqrt[a + b*x^2]*(c + d*x^2)^(3/2)*(A + B*x^2 + C*x^4))/(e + f*x^2),x]`

output

```

-1/105*(C*(2*b*c - a*d)*(3*b^2*c^2 - 3*a*b*c*d + 8*a^2*d^2)*x*Sqrt[a + b*x
^2])/(b^3*d*f*Sqrt[c + d*x^2]) - ((3*b^2*c^2 + 7*a*b*c*d - 2*a^2*d^2)*(C*e
- B*f)*x*Sqrt[a + b*x^2])/(15*b^2*f^2*Sqrt[c + d*x^2]) - (d*(3*b*d*e - 4*
b*c*f - a*d*f)*(C*e^2 - B*e*f + A*f^2)*x*Sqrt[a + b*x^2])/(3*b*f^4*Sqrt[c
+ d*x^2]) + (C*(9*a*c + (3*b*c^2)/d - (4*a^2*d)/b)*x*Sqrt[a + b*x^2]*Sqrt[
c + d*x^2])/(105*b*f) - (2*(3*b*c - a*d)*(C*e - B*f)*x*Sqrt[a + b*x^2]*Sqr
t[c + d*x^2])/(15*b*f^2) + (d*(C*e^2 - B*e*f + A*f^2)*x*Sqrt[a + b*x^2]*Sq
rt[c + d*x^2])/(3*f^3) + (C*(3*b*c + a*d)*x^3*Sqrt[a + b*x^2]*Sqrt[c + d*x
^2])/(35*b*f) - (d*(C*e - B*f)*x*(a + b*x^2)^(3/2)*Sqrt[c + d*x^2])/(5*b*f
^2) + (C*x^3*Sqrt[a + b*x^2]*(c + d*x^2)^(3/2))/(7*f) + (Sqrt[c]*C*(2*b*c
- a*d)*(3*b^2*c^2 - 3*a*b*c*d + 8*a^2*d^2)*Sqrt[a + b*x^2]*EllipticE[ArcTan
[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)])/(105*b^3*d^(3/2)*f*Sqrt[(c*(a +
b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) + (Sqrt[c]*(3*b^2*c^2 + 7*a*b*c*
d - 2*a^2*d^2)*(C*e - B*f)*Sqrt[a + b*x^2]*EllipticE[ArcTan[(Sqrt[d]*x)/Sq
rt[c]], 1 - (b*c)/(a*d)])/(15*b^2*Sqrt[d]*f^2*Sqrt[(c*(a + b*x^2))/(a*(c +
d*x^2))]*Sqrt[c + d*x^2]) + (Sqrt[c]*Sqrt[d]*(3*b*d*e - 4*b*c*f - a*d*f)*
(C*e^2 - B*e*f + A*f^2)*Sqrt[a + b*x^2]*EllipticE[ArcTan[(Sqrt[d]*x)/Sqrt[
c]], 1 - (b*c)/(a*d)])/(3*b*f^4*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt
[c + d*x^2]) - (c^(3/2)*C*(3*b^2*c^2 + 9*a*b*c*d - 4*a^2*d^2)*Sqrt[a + b*x
^2]*EllipticF[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)])/(105*b^2*d...

```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 7276

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Maple [A] (verified)

Time = 19.54 (sec) , antiderivative size = 1360, normalized size of antiderivative = 1.27

method	result	size
risch	Expression too large to display	1360
default	Expression too large to display	5349
elliptic	Expression too large to display	6489

input `int((b*x^2+a)^(1/2)*(d*x^2+c)^(3/2)*(C*x^4+B*x^2+A)/(f*x^2+e),x,method=_RE
TURNVERBOSE)`

output `1/105*x/d*(15*C*b^2*d^2*f^2*x^4+21*B*b^2*d^2*f^2*x^2+3*C*a*b*d^2*f^2*x^2+
4*C*b^2*c*d*f^2*x^2-21*C*b^2*d^2*e*f*x^2+35*A*b^2*d^2*f^2+7*B*a*b*d^2*f^2+
42*B*b^2*c*d*f^2-35*B*b^2*d^2*e*f-4*C*a^2*d^2*f^2+9*C*a*b*c*d*f^2-7*C*a*b*
d^2*e*f+3*C*b^2*c^2*f^2-42*C*b^2*c*d*e*f+35*C*b^2*d^2*e^2)*(b*x^2+a)^(1/2)
*(d*x^2+c)^(1/2)/b^2/f^3+1/105/f^3/d/b^2*((175*A*a*b^2*c*d^2*f^4-105*A*a*b
^2*d^3*e*f^3+105*A*b^3*c^2*d*f^4-210*A*b^3*c*d^2*e*f^3+105*A*b^3*d^3*e^2*f
^2-7*B*a^2*b*c*d^2*f^4+63*B*a*b^2*c^2*d*f^4-175*B*a*b^2*c*d^2*e*f^3+105*B*
a*b^2*d^3*e^2*f^2-105*B*b^3*c^2*d*e*f^3+210*B*b^3*c*d^2*e^2*f^2-105*B*b^3*
d^3*e^3*f+4*C*a^3*c*d^2*f^4-9*C*a^2*b*c^2*d*f^4+7*C*a^2*b*c*d^2*e*f^3-3*C*
a*b^2*c^3*f^4-63*C*a*b^2*c^2*d*e*f^3+175*C*a*b^2*c*d^2*e^2*f^2-105*C*a*b^2
*d^3*e^3*f+105*C*b^3*c^2*d*e^2*f^2-210*C*b^3*c*d^2*e^3*f+105*C*b^3*d^3*e^4
)/f^2/(-b/a)^(1/2)*(1+b*x^2/a)^(1/2)*(1+d*x^2/c)^(1/2)/(b*d*x^4+a*d*x^2+b*
c*x^2+a*c)^(1/2)*EllipticF(x*(-b/a)^(1/2),(-1+(a*d+b*c)/c/b)^(1/2))-1/f*(3
5*A*a*b^2*d^3*f^3+140*A*b^3*c*d^2*f^3-105*A*b^3*d^3*e*f^2-14*B*a^2*b*d^3*f
^3+49*B*a*b^2*c*d^2*f^3-35*B*a*b^2*d^3*e*f^2+21*B*b^3*c^2*d*f^3-140*B*b^3*
c*d^2*e*f^2+105*B*b^3*d^3*e^2*f+8*C*a^3*d^3*f^3-19*C*a^2*b*c*d^2*f^3+14*C*
a^2*b*d^3*e*f^2+9*C*a*b^2*c^2*d*f^3-49*C*a*b^2*c*d^2*e*f^2+35*C*a*b^2*d^3*
e^2*f-6*C*b^3*c^3*f^3-21*C*b^3*c^2*d*e*f^2+140*C*b^3*c*d^2*e^2*f-105*C*b^3
*d^3*e^3)*c/(-b/a)^(1/2)*(1+b*x^2/a)^(1/2)*(1+d*x^2/c)^(1/2)/(b*d*x^4+a*d*
x^2+b*c*x^2+a*c)^(1/2)/d*(EllipticF(x*(-b/a)^(1/2),(-1+(a*d+b*c)/c/b)^(...`

Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{a+bx^2}(c+dx^2)^{3/2}(A+Bx^2+Cx^4)}{e+fx^2} dx = \text{Timed out}$$

input `integrate((b*x^2+a)^(1/2)*(d*x^2+c)^(3/2)*(C*x^4+B*x^2+A)/(f*x^2+e),x,alg
orithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{\sqrt{a+bx^2}(c+dx^2)^{3/2}(A+Bx^2+Cx^4)}{e+fx^2} dx = \int \frac{\sqrt{a+bx^2}(c+dx^2)^{3/2}(A+Bx^2+Cx^4)}{e+fx^2} dx$$

input `integrate((b*x**2+a)**(1/2)*(d*x**2+c)**(3/2)*(C*x**4+B*x**2+A)/(f*x**2+e),x)`

output `Integral(sqrt(a + b*x**2)*(c + d*x**2)**(3/2)*(A + B*x**2 + C*x**4)/(e + f*x**2), x)`

Maxima [F]

$$\int \frac{\sqrt{a+bx^2}(c+dx^2)^{3/2}(A+Bx^2+Cx^4)}{e+fx^2} dx = \int \frac{(Cx^4+Bx^2+A)\sqrt{bx^2+a}(dx^2+c)^{3/2}}{fx^2+e} dx$$

input `integrate((b*x^2+a)^(1/2)*(d*x^2+c)^(3/2)*(C*x^4+B*x^2+A)/(f*x^2+e),x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)*sqrt(b*x^2 + a)*(d*x^2 + c)^(3/2)/(f*x^2 + e), x)`

Giac [F]

$$\int \frac{\sqrt{a+bx^2}(c+dx^2)^{3/2}(A+Bx^2+Cx^4)}{e+fx^2} dx = \int \frac{(Cx^4+Bx^2+A)\sqrt{bx^2+a}(dx^2+c)^{3/2}}{fx^2+e} dx$$

input `integrate((b*x^2+a)^(1/2)*(d*x^2+c)^(3/2)*(C*x^4+B*x^2+A)/(f*x^2+e),x, algorithm="giac")`

output

```
integrate((C*x^4 + B*x^2 + A)*sqrt(b*x^2 + a)*(d*x^2 + c)^(3/2)/(f*x^2 + e), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + bx^2}(c + dx^2)^{3/2}(A + Bx^2 + Cx^4)}{e + fx^2} dx = \int \frac{\sqrt{bx^2 + a}(dx^2 + c)^{3/2}(Cx^4 + Bx^2 + A)}{fx^2 + e} dx$$

input

```
int(((a + b*x^2)^(1/2)*(c + d*x^2)^(3/2)*(A + B*x^2 + C*x^4))/(e + f*x^2), x)
```

output

```
int(((a + b*x^2)^(1/2)*(c + d*x^2)^(3/2)*(A + B*x^2 + C*x^4))/(e + f*x^2), x)
```

Reduce [F]

$$\int \frac{\sqrt{a + bx^2}(c + dx^2)^{3/2}(A + Bx^2 + Cx^4)}{e + fx^2} dx = \int \frac{\sqrt{bx^2 + a}(dx^2 + c)^{\frac{3}{2}}(Cx^4 + Bx^2 + A)}{fx^2 + e} dx$$

input

```
int((b*x^2+a)^(1/2)*(d*x^2+c)^(3/2)*(C*x^4+B*x^2+A)/(f*x^2+e), x)
```

output

```
int((b*x^2+a)^(1/2)*(d*x^2+c)^(3/2)*(C*x^4+B*x^2+A)/(f*x^2+e), x)
```

$$3.9 \quad \int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{e+fx^2} dx$$

Optimal result	107
Mathematica [C] (verified)	108
Rubi [A] (verified)	109
Maple [A] (verified)	112
Fricas [F(-1)]	113
Sympy [F]	113
Maxima [F]	113
Giac [F]	114
Mupad [F(-1)]	114
Reduce [F]	114

Optimal result

Integrand size = 44, antiderivative size = 666

$$\int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{e+fx^2} dx =$$

$$\frac{\left(2a^2Cdf + b^2\left(5cCe + 15Bde - \frac{15Cde^2}{f} - 5Bcf + \frac{2c^2Cf}{d} - 15Adf\right) + ab(5Cde - 2cCf - 5Bdf)\right) x\sqrt{c+dx^2}}{15bdf^2\sqrt{a+bx^2}}$$

$$+ \frac{(aCdf - b(5Cde + 2cCf - 5Bdf))x\sqrt{a+bx^2}\sqrt{c+dx^2}}{15bdf^2} + \frac{Cx\sqrt{a+bx^2}(c+dx^2)^{3/2}}{5df}$$

$$+ \frac{\sqrt{a}(2a^2Cd^2f^2 + abdf(5Cde - 2cCf - 5Bdf) + b^2(5df(3Bde - Bcf - 3Adf) - C(15d^2e^2 - 5cdef - 15b^3/2d^2f^3\sqrt{a+bx^2}\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}))}{15b^3/2d^2f^3\sqrt{a+bx^2}\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}}$$

$$- \frac{a^{3/2}(acCdf^2 + 5bdf(3Bde - 2Bcf - 3Adf) - bC(15d^2e^2 - 10cdef - c^2f^2))\sqrt{c+dx^2}\text{EllipticF}\left(\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right), 1 - \frac{ad}{bc}\right)}{15b^3/2cdf^3\sqrt{a+bx^2}\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}}$$

$$- \frac{a^{3/2}(de - cf)(Ce^2 - Bef + Af^2)\sqrt{c+dx^2}\text{EllipticPi}\left(1 - \frac{af}{be}, \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right), 1 - \frac{ad}{bc}\right)}{\sqrt{bce}f^3\sqrt{a+bx^2}\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}}$$

output

```

-1/15*(2*a^2*C*d*f+b^2*(5*c*C*e+15*B*d*e-15*C*d*e^2/f-5*B*c*f+2*c^2*C*f/d-
15*A*d*f)+a*b*(-5*B*d*f-2*C*c*f+5*C*d*e))*x*(d*x^2+c)^(1/2)/b/d/f^2/(b*x^2
+a)^(1/2)+1/15*(a*C*d*f-b*(-5*B*d*f+2*C*c*f+5*C*d*e))*x*(b*x^2+a)^(1/2)*(d
*x^2+c)^(1/2)/b/d/f^2+1/5*C*x*(b*x^2+a)^(1/2)*(d*x^2+c)^(3/2)/d/f+1/15*a^(
1/2)*(2*a^2*C*d^2*f^2+a*b*d*f*(-5*B*d*f-2*C*c*f+5*C*d*e)+b^2*(5*d*f*(-3*A*
d*f-B*c*f+3*B*d*e)-C*(-2*c^2*f^2-5*c*d*e*f+15*d^2*e^2)))*(d*x^2+c)^(1/2)*E
llipticE(b^(1/2)*x/a^(1/2)/(1+b*x^2/a)^(1/2),(1-a*d/b/c)^(1/2))/b^(3/2)/d^
2/f^3/(b*x^2+a)^(1/2)/(a*(d*x^2+c)/c/(b*x^2+a))^(1/2)-1/15*a^(3/2)*(a*c*C*
d*f^2+5*b*d*f*(-3*A*d*f-2*B*c*f+3*B*d*e)-b*C*(-c^2*f^2-10*c*d*e*f+15*d^2*e
^2))*(d*x^2+c)^(1/2)*InverseJacobiAM(arctan(b^(1/2)*x/a^(1/2)),(1-a*d/b/c)
^(1/2))/b^(3/2)/c/d/f^3/(b*x^2+a)^(1/2)/(a*(d*x^2+c)/c/(b*x^2+a))^(1/2)-a^
(3/2)*(-c*f+d*e)*(A*f^2-B*e*f+C*e^2)*(d*x^2+c)^(1/2)*EllipticPi(b^(1/2)*x/
a^(1/2)/(1+b*x^2/a)^(1/2),1-a*f/b/e,(1-a*d/b/c)^(1/2))/b^(1/2)/c/e/f^3/(b*
x^2+a)^(1/2)/(a*(d*x^2+c)/c/(b*x^2+a))^(1/2)

```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 7.82 (sec) , antiderivative size = 541, normalized size of antiderivative = 0.81

$$\int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{e+fx^2} dx$$

$$= \frac{icf(2a^2Cd^2f^2+abdf(5Cde-2cCf-5Bdf))+b^2(-5df(-3Bde+Bcf+3Adf))+C(-15d^2e^2+5cde}}{e^2+2efx^2+fx^4}$$

input

```

Integrate[(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(A + B*x^2 + C*x^4))/(e + f*x^2
),x]

```

output

```
(I*c*e*f*(2*a^2*C*d^2*f^2 + a*b*d*f*(5*C*d*e - 2*c*C*f - 5*B*d*f) + b^2*(-
5*d*f*(-3*B*d*e + B*c*f + 3*A*d*f) + C*(-15*d^2*e^2 + 5*c*d*e*f + 2*c^2*f^
2)))*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]
*x], (a*d)/(b*c)] - I*e*(a^2*c*C*d^2*f^3 + a*b*d*f*(5*d*f*(-3*B*d*e + B*c*
f + 3*A*d*f) + C*(15*d^2*e^2 - 5*c*d*e*f - 3*c^2*f^2)) + b^2*(-5*d*f*(-3*B
*d^2*e^2 + 3*A*d^2*e*f + B*c^2*f^2) + C*(-15*d^3*e^3 + 5*c^2*d*e*f^2 + 2*c
^3*f^3))*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt
[b/a]*x], (a*d)/(b*c)] + d*(Sqrt[b/a]*e*f^2*x*(a + b*x^2)*(c + d*x^2)*(5*b
*B*d*f + a*C*d*f + b*C*(-5*d*e + c*f + 3*d*f*x^2)) - (15*I)*b*d*(-(b*e) +
a*f)*(-(d*e) + c*f)*(C*e^2 + f*(-(B*e) + A*f))*Sqrt[1 + (b*x^2)/a]*Sqrt[1
+ (d*x^2)/c]*EllipticPi[(a*f)/(b*e), I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)])
)/(15*b*Sqrt[b/a]*d^2*e*f^4*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])
```

Rubi [A] (verified)

Time = 1.77 (sec) , antiderivative size = 1029, normalized size of antiderivative = 1.55, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.045$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{e+fx^2} dx$$

↓ 7276

$$\int \left(\frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(Af^2 - Bef + Ce^2)}{f^2(e+fx^2)} - \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(Ce - Bf)}{f^2} + \frac{Cx^2\sqrt{a+bx^2}\sqrt{c+dx^2}}{f} \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{C\sqrt{bx^2+a}\sqrt{dx^2+cx^3}}{5f} - \frac{(Ce-Bf)\sqrt{bx^2+a}\sqrt{dx^2+cx}}{3f^2} + \\
& \frac{C(bc+ad)\sqrt{bx^2+a}\sqrt{dx^2+cx}}{15bdf} - \frac{(bc+ad)(Ce-Bf)\sqrt{bx^2+ax}}{3bf^2\sqrt{dx^2+c}} + \\
& \frac{d(Ce^2-Bfe+Af^2)\sqrt{bx^2+ax}}{f^3\sqrt{dx^2+c}} - \frac{2C(b^2c^2-abdc+a^2d^2)\sqrt{bx^2+ax}}{15b^2df\sqrt{dx^2+c}} + \\
& \frac{\sqrt{c}(bc+ad)(Ce-Bf)\sqrt{bx^2+a}E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\middle|1-\frac{bc}{ad}\right)}{3b\sqrt{d}f^2\sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}}\sqrt{dx^2+c}} - \\
& \frac{\sqrt{c}\sqrt{d}(Ce^2-Bfe+Af^2)\sqrt{bx^2+a}E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\middle|1-\frac{bc}{ad}\right)}{f^3\sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}}\sqrt{dx^2+c}} + \\
& \frac{2\sqrt{c}C(b^2c^2-abdc+a^2d^2)\sqrt{bx^2+a}E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\middle|1-\frac{bc}{ad}\right)}{15b^2d^{3/2}f\sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}}\sqrt{dx^2+c}} - \\
& \frac{2c^{3/2}(Ce-Bf)\sqrt{bx^2+a}\operatorname{EllipticF}\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right),1-\frac{bc}{ad}\right)}{3\sqrt{d}f^2\sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}}\sqrt{dx^2+c}} + \\
& \frac{bc^{3/2}(Ce^2-Bfe+Af^2)\sqrt{bx^2+a}\operatorname{EllipticF}\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right),1-\frac{bc}{ad}\right)}{a\sqrt{d}f^3\sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}}\sqrt{dx^2+c}} - \\
& \frac{c^{3/2}C(bc+ad)\sqrt{bx^2+a}\operatorname{EllipticF}\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right),1-\frac{bc}{ad}\right)}{15bd^{3/2}f\sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}}\sqrt{dx^2+c}} - \\
& \frac{c^{3/2}(be-af)(Ce^2-Bfe+Af^2)\sqrt{bx^2+a}\operatorname{EllipticPi}\left(1-\frac{cf}{de},\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right),1-\frac{bc}{ad}\right)}{a\sqrt{de}f^3\sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}}\sqrt{dx^2+c}}
\end{aligned}$$

input `Int[(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(A + B*x^2 + C*x^4))/(e + f*x^2),x]`

output

```
(-2*C*(b^2*c^2 - a*b*c*d + a^2*d^2)*x*Sqrt[a + b*x^2])/(15*b^2*d*f*Sqrt[c + d*x^2]) - ((b*c + a*d)*(C*e - B*f)*x*Sqrt[a + b*x^2])/(3*b*f^2*Sqrt[c + d*x^2]) + (d*(C*e^2 - B*e*f + A*f^2)*x*Sqrt[a + b*x^2])/(f^3*Sqrt[c + d*x^2]) + (C*(b*c + a*d)*x*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])/(15*b*d*f) - ((C*e - B*f)*x*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])/(3*f^2) + (C*x^3*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])/(5*f) + (2*Sqrt[c]*C*(b^2*c^2 - a*b*c*d + a^2*d^2)*Sqrt[a + b*x^2]*EllipticE[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(15*b^2*d^(3/2)*f*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) + (Sqrt[c]*(b*c + a*d)*(C*e - B*f)*Sqrt[a + b*x^2]*EllipticE[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(3*b*Sqrt[d]*f^2*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) - (Sqrt[c]*Sqrt[d]*(C*e^2 - B*e*f + A*f^2)*Sqrt[a + b*x^2]*EllipticE[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(f^3*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) - (c^(3/2)*C*(b*c + a*d)*Sqrt[a + b*x^2]*EllipticF[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(15*b*d^(3/2)*f*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) - (2*c^(3/2)*(C*e - B*f)*Sqrt[a + b*x^2]*EllipticF[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(3*Sqrt[d]*f^2*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) + (b*c^(3/2)*(C*e^2 - B*e*f + A*f^2)*Sqrt[a + b*x^2]*EllipticF[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(a*Sqrt[d]*f^3*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) - (c^(3/2)*(b*e - ...
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 7276

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE  
x  
pand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ  
[n, 0]
```


Maple [A] (verified)

Time = 9.10 (sec) , antiderivative size = 802, normalized size of antiderivative = 1.20

method	result
risch	$\frac{x(3C x^2 bdf + 5bBdf + aCdf + Cbfc - 5Cbde)\sqrt{bx^2+a}\sqrt{x^2d+c}}{15bd f^2} + \left(\frac{(15Aab d^2 f^3 + 15A b^2 cd f^3 - 15A b^2 d^2 e f^2 + 10Babcd f^3 - 15Bab d^2 e f^2}{15bd f^2} \right)$
default	Expression too large to display
elliptic	Expression too large to display

input

```
int((b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*(C*x^4+B*x^2+A)/(f*x^2+e),x,method=_RE
TURNVERBOSE)
```

output

```
1/15*x*(3*C*b*d*f*x^2+5*B*b*d*f+C*a*d*f+C*b*c*f-5*C*b*d*e)*(b*x^2+a)^(1/2)
*(d*x^2+c)^(1/2)/b/d/f^2+1/15/d/b/f^2*((15*A*a*b*d^2*f^3+15*A*b^2*c*d*f^3-
15*A*b^2*d^2*e*f^2+10*B*a*b*c*d*f^3-15*B*a*b*d^2*e*f^2-15*B*b^2*c*d*e*f^2+
15*B*b^2*d^2*e^2*f-C*a^2*c*d*f^3-C*a*b*c^2*f^3-10*C*a*b*c*d*e*f^2+15*C*a*b
*d^2*e^2*f+15*C*b^2*c*d*e^2*f-15*C*b^2*d^2*e^3)/f^2/(-b/a)^(1/2)*(1+b*x^2/
a)^(1/2)*(1+d*x^2/c)^(1/2)/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)^(1/2)*EllipticF(x
*(-b/a)^(1/2),(-1+(a*d+b*c)/c/b)^(1/2))-1/f*(15*A*b^2*d^2*f^2+5*B*a*b*d^2*
f^2+5*B*b^2*c*d*f^2-15*B*b^2*d^2*e*f-2*C*a^2*d^2*f^2+2*C*a*b*c*d*f^2-5*C*a
*b*d^2*e*f-2*C*b^2*c^2*f^2-5*C*b^2*c*d*e*f+15*C*b^2*d^2*e^2)*c/(-b/a)^(1/2)
*(1+b*x^2/a)^(1/2)*(1+d*x^2/c)^(1/2)/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)^(1/2)/
d*(EllipticF(x*(-b/a)^(1/2),(-1+(a*d+b*c)/c/b)^(1/2))-EllipticE(x*(-b/a)^(
1/2),(-1+(a*d+b*c)/c/b)^(1/2)))+15*(A*a*c*f^4-A*a*d*e*f^3-A*b*c*e*f^3+A*b*
d*e^2*f^2-B*a*c*e*f^3+B*a*d*e^2*f^2+B*b*c*e^2*f^2-B*b*d*e^3*f+C*a*c*e^2*f^
2-C*a*d*e^3*f-C*b*c*e^3*f+C*b*d*e^4)*b*d/f^2/e/(-b/a)^(1/2)*(1+b*x^2/a)^(1
/2)*(1+d*x^2/c)^(1/2)/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)^(1/2)*EllipticPi(x*(-b
/a)^(1/2),a*f/b/e,(-1/c*d)^(1/2)/(-b/a)^(1/2))*((b*x^2+a)*(d*x^2+c))^(1/2)
)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)
```

Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + bx^2} \sqrt{c + dx^2} (A + Bx^2 + Cx^4)}{e + fx^2} dx = \text{Timed out}$$

input `integrate((b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*(C*x^4+B*x^2+A)/(f*x^2+e),x, alg
orithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{\sqrt{a + bx^2} \sqrt{c + dx^2} (A + Bx^2 + Cx^4)}{e + fx^2} dx = \int \frac{\sqrt{a + bx^2} \sqrt{c + dx^2} (A + Bx^2 + Cx^4)}{e + fx^2} dx$$

input `integrate((b*x**2+a)**(1/2)*(d*x**2+c)**(1/2)*(C*x**4+B*x**2+A)/(f*x**2+e)
,x)`

output `Integral(sqrt(a + b*x**2)*sqrt(c + d*x**2)*(A + B*x**2 + C*x**4)/(e + f*x*
*2), x)`

Maxima [F]

$$\int \frac{\sqrt{a + bx^2} \sqrt{c + dx^2} (A + Bx^2 + Cx^4)}{e + fx^2} dx = \int \frac{(Cx^4 + Bx^2 + A) \sqrt{bx^2 + a} \sqrt{dx^2 + c}}{fx^2 + e} dx$$

input `integrate((b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*(C*x^4+B*x^2+A)/(f*x^2+e),x, alg
orithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)*sqrt(b*x^2 + a)*sqrt(d*x^2 + c)/(f*x^2 + e),
x)`

Giac [F]

$$\int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{e+fx^2} dx = \int \frac{(Cx^4+Bx^2+A)\sqrt{bx^2+a}\sqrt{dx^2+c}}{fx^2+e} dx$$

input `integrate((b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*(C*x^4+B*x^2+A)/(f*x^2+e),x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)*sqrt(b*x^2 + a)*sqrt(d*x^2 + c)/(f*x^2 + e), x)`

Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{e+fx^2} dx \\ &= \int \frac{\sqrt{bx^2+a}\sqrt{dx^2+c}(Cx^4+Bx^2+A)}{fx^2+e} dx \end{aligned}$$

input `int(((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(A + B*x^2 + C*x^4))/(e + f*x^2), x)`

output `int(((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(A + B*x^2 + C*x^4))/(e + f*x^2), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{e+fx^2} dx \\ &= \int \frac{\sqrt{bx^2+a}\sqrt{dx^2+c}(Cx^4+Bx^2+A)}{fx^2+e} dx \end{aligned}$$

input `int((b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*(C*x^4+B*x^2+A)/(f*x^2+e),x)`

output `int((b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*(C*x^4+B*x^2+A)/(f*x^2+e),x)`

$$3.10 \quad \int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{\sqrt{c+dx^2}(e+fx^2)} dx$$

Optimal result	116
Mathematica [C] (verified)	117
Rubi [A] (verified)	118
Maple [A] (verified)	119
Fricas [F(-1)]	120
Sympy [F]	120
Maxima [F]	121
Giac [F]	121
Mupad [F(-1)]	122
Reduce [F]	122

Optimal result

Integrand size = 44, antiderivative size = 443

$$\begin{aligned}
& \int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{\sqrt{c+dx^2}(e+fx^2)} dx \\
&= \frac{(aCdf - b(3Cde + 2cCf - 3Bdf))x\sqrt{c+dx^2}}{3d^2 f^2 \sqrt{a+bx^2}} + \frac{Cx\sqrt{a+bx^2}\sqrt{c+dx^2}}{3df} \\
&\quad - \frac{\sqrt{a}(aCdf - b(3Cde + 2cCf - 3Bdf))\sqrt{c+dx^2} E\left(\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) \mid 1 - \frac{ad}{bc}\right)}{3\sqrt{bd^2} f^2 \sqrt{a+bx^2} \sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}} \\
&\quad - \frac{a^{3/2}(3Cde + cCf - 3Bdf)\sqrt{c+dx^2} \operatorname{EllipticF}\left(\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right), 1 - \frac{ad}{bc}\right)}{3\sqrt{bcd} f^2 \sqrt{a+bx^2} \sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}} \\
&\quad + \frac{a^{3/2}(Ce^2 - Bef + Af^2)\sqrt{c+dx^2} \operatorname{EllipticPi}\left(1 - \frac{af}{be}, \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right), 1 - \frac{ad}{bc}\right)}{\sqrt{bce} f^2 \sqrt{a+bx^2} \sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}}
\end{aligned}$$

output

```

1/3*(a*C*d*f-b*(-3*B*d*f+2*C*c*f+3*C*d*e))*x*(d*x^2+c)^(1/2)/d^2/f^2/(b*x^
2+a)^(1/2)+1/3*C*x*(b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)/d/f-1/3*a^(1/2)*(a*C*d*
f-b*(-3*B*d*f+2*C*c*f+3*C*d*e))*(d*x^2+c)^(1/2)*EllipticE(b^(1/2)*x/a^(1/2
)/(1+b*x^2/a)^(1/2),(1-a*d/b/c)^(1/2))/b^(1/2)/d^2/f^2/(b*x^2+a)^(1/2)/(a*
(d*x^2+c)/c/(b*x^2+a)^(1/2)-1/3*a^(3/2)*(-3*B*d*f+C*c*f+3*C*d*e)*(d*x^2+c
)^(1/2)*InverseJacobiAM(arctan(b^(1/2)*x/a^(1/2)),(1-a*d/b/c)^(1/2))/b^(1/
2)/c/d/f^2/(b*x^2+a)^(1/2)/(a*(d*x^2+c)/c/(b*x^2+a))^(1/2)+a^(3/2)*(A*f^2-
B*e*f+C*e^2)*(d*x^2+c)^(1/2)*EllipticPi(b^(1/2)*x/a^(1/2)/(1+b*x^2/a)^(1/2
),1-a*f/b/e,(1-a*d/b/c)^(1/2))/b^(1/2)/c/e/f^2/(b*x^2+a)^(1/2)/(a*(d*x^2+c
)/c/(b*x^2+a))^(1/2)

```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 5.51 (sec) , antiderivative size = 383, normalized size of antiderivative = 0.86

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{\sqrt{c+dx^2}(e+fx^2)} dx$$

$$= \frac{-icf(aCdf + b(-3Cde - 2cCf + 3Bdf))\sqrt{1 + \frac{bx^2}{a}}\sqrt{1 + \frac{dx^2}{c}}E\left(i\operatorname{arcsinh}\left(\sqrt{\frac{b}{a}}x\right)\middle|\frac{ad}{bc}\right) + ie(adf(3Cde +$$

input

```

Integrate[(Sqrt[a + b*x^2]*(A + B*x^2 + C*x^4))/(Sqrt[c + d*x^2]*(e + f*x^
2)),x]

```

output

```

((-I)*c*e*f*(a*C*d*f + b*(-3*C*d*e - 2*c*C*f + 3*B*d*f))*Sqrt[1 + (b*x^2)/
a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] + I*
e*(a*d*f*(3*C*d*e + 2*c*C*f - 3*B*d*f) - b*(C*(3*d^2*e^2 + 3*c*d*e*f + 2*c
^2*f^2) + 3*d*f*(A*d*f - B*(d*e + c*f))))*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*
x^2)/c]*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] + d*(Sqrt[b/a]*C*e*
f^2*x*(a + b*x^2)*(c + d*x^2) - (3*I)*d*(-(b*e) + a*f)*(C*e^2 + f*(-(B*e)
+ A*f))*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticPi[(a*f)/(b*e), I*
ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)))/(3*Sqrt[b/a]*d^2*e*f^3*Sqrt[a + b*x^2
]*Sqrt[c + d*x^2])

```

Rubi [A] (verified)

Time = 1.27 (sec) , antiderivative size = 623, normalized size of antiderivative = 1.41, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.045$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{\sqrt{c+dx^2}(e+fx^2)} dx$$

↓ 7276

$$\int \left(\frac{\sqrt{a+bx^2}(Af^2 - Bef + Ce^2)}{f^2\sqrt{c+dx^2}(e+fx^2)} - \frac{\sqrt{a+bx^2}(Ce - Bf)}{f^2\sqrt{c+dx^2}} + \frac{Cx^2\sqrt{a+bx^2}}{f\sqrt{c+dx^2}} \right) dx$$

↓ 2009

$$\frac{a^{3/2}\sqrt{c+dx^2}(Af^2 - Bef + Ce^2) \text{EllipticPi}\left(1 - \frac{af}{be}, \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right), 1 - \frac{ad}{bc}\right) - \sqrt{bce}f^2\sqrt{a+bx^2}\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}}{\sqrt{c}\sqrt{a+bx^2}(Ce - Bf) \text{EllipticF}\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right) + \frac{\sqrt{d}f^2\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}{\sqrt{c}\sqrt{a+bx^2}(Ce - Bf)E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right) - \frac{\sqrt{d}f^2\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}{c^{3/2}C\sqrt{a+bx^2} \text{EllipticF}\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right) + \frac{3d^{3/2}f\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}{\sqrt{c}C\sqrt{a+bx^2}(2bc - ad)E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right) - \frac{x\sqrt{a+bx^2}(Ce - Bf)}{f^2\sqrt{c+dx^2}} + \frac{3bd^{3/2}f\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}{3df} - \frac{Cx\sqrt{a+bx^2}\sqrt{c+dx^2}}{3bdf\sqrt{c+dx^2}} - \frac{Cx\sqrt{a+bx^2}(2bc - ad)}{3bdf\sqrt{c+dx^2}}$$

input

```
Int[(Sqrt[a + b*x^2]*(A + B*x^2 + C*x^4))/(Sqrt[c + d*x^2]*(e + f*x^2)),x]
```

output

```
-1/3*(C*(2*b*c - a*d)*x*Sqrt[a + b*x^2])/(b*d*f*Sqrt[c + d*x^2]) - ((C*e - B*f)*x*Sqrt[a + b*x^2])/(f^2*Sqrt[c + d*x^2]) + (C*x*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])/(3*d*f) + (Sqrt[c]*C*(2*b*c - a*d)*Sqrt[a + b*x^2]*EllipticE[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(3*b*d^(3/2)*f*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) + (Sqrt[c]*(C*e - B*f)*Sqrt[a + b*x^2]*EllipticE[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(Sqrt[d]*f^2*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) - (c^(3/2)*C*Sqrt[a + b*x^2]*EllipticF[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(3*d^(3/2)*f*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) - (Sqrt[c]*(C*e - B*f)*Sqrt[a + b*x^2]*EllipticF[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(Sqrt[d]*f^2*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) + (a^(3/2)*(C*e^2 - B*e*f + A*f^2)*Sqrt[c + d*x^2]*EllipticPi[1 - (a*f)/(b*e), ArcTan[(Sqrt[b]*x)/Sqrt[a]], 1 - (a*d)/(b*c)]/(Sqrt[b]*c*e*f^2*Sqrt[a + b*x^2]*Sqrt[(a*(c + d*x^2))/(c*(a + b*x^2))])
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 7276

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]
```

Maple [A] (verified)

Time = 8.61 (sec) , antiderivative size = 506, normalized size of antiderivative = 1.14

method	result
risch	$\frac{Cx\sqrt{bx^2+a}\sqrt{x^2d+c}}{3df} + \frac{\left((3Abdf^2+3Badf^2-3Bbdef-Cacf^2-3aCdef+3e^2Cbd)\sqrt{1+\frac{bx^2}{a}}\sqrt{1+\frac{dx^2}{c}} \operatorname{EllipticF}\left(x\sqrt{-\frac{b}{a}},\sqrt{-1+\frac{ad+bc}{cb}}\right) \right)}{f^2\sqrt{-\frac{b}{a}}\sqrt{bdx^4+adx^2+x^2bc+ac}}$
default	Expression too large to display
elliptic	Expression too large to display

input `int((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)^(1/2)/(f*x^2+e),x,method=_RE
TURNVERBOSE)`

output `1/3*C*x*(b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)/d/f+1/3/d/f*((3*A*b*d*f^2+3*B*a*d*
f^2-3*B*b*d*e*f-C*a*c*f^2-3*C*a*d*e*f+3*C*b*d*e^2)/f^2/(-b/a)^(1/2)*(1+b*x
^2/a)^(1/2)*(1+d*x^2/c)^(1/2)/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)^(1/2)*Elliptic
F(x*(-b/a)^(1/2),(-1+(a*d+b*c)/c/b)^(1/2))-1/f*(3*B*b*d*f+C*a*d*f-2*C*b*c*
f-3*C*b*d*e)*c/(-b/a)^(1/2)*(1+b*x^2/a)^(1/2)*(1+d*x^2/c)^(1/2)/(b*d*x^4+a
*d*x^2+b*c*x^2+a*c)^(1/2)/d*(EllipticF(x*(-b/a)^(1/2),(-1+(a*d+b*c)/c/b)^(
1/2))-EllipticE(x*(-b/a)^(1/2),(-1+(a*d+b*c)/c/b)^(1/2)))+3*(A*a*f^3-A*b*e
*f^2-B*a*e*f^2+B*b*e^2*f+C*a*e^2*f-C*b*e^3)*d/f^2/e/(-b/a)^(1/2)*(1+b*x^2/
a)^(1/2)*(1+d*x^2/c)^(1/2)/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)^(1/2)*EllipticPi(
x*(-b/a)^(1/2),a*f/b/e,(-1/c*d)^(1/2)/(-b/a)^(1/2))*((b*x^2+a)*(d*x^2+c)
^(1/2)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2))`

Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{\sqrt{c+dx^2}(e+fx^2)} dx = \text{Timed out}$$

input `integrate((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)^(1/2)/(f*x^2+e),x,alg
orithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{\sqrt{c+dx^2}(e+fx^2)} dx = \int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{\sqrt{c+dx^2}(e+fx^2)} dx$$

input `integrate((b*x**2+a)**(1/2)*(C*x**4+B*x**2+A)/(d*x**2+c)**(1/2)/(f*x**2+e)
,x)`

output `Integral(sqrt(a + b*x**2)*(A + B*x**2 + C*x**4)/(sqrt(c + d*x**2)*(e + f*x**2)), x)`

Maxima [F]

$$\int \frac{\sqrt{a + bx^2}(A + Bx^2 + Cx^4)}{\sqrt{c + dx^2}(e + fx^2)} dx = \int \frac{(Cx^4 + Bx^2 + A)\sqrt{bx^2 + a}}{\sqrt{dx^2 + c}(fx^2 + e)} dx$$

input `integrate((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)^(1/2)/(f*x^2+e),x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)*sqrt(b*x^2 + a)/(sqrt(d*x^2 + c)*(f*x^2 + e)), x)`

Giac [F]

$$\int \frac{\sqrt{a + bx^2}(A + Bx^2 + Cx^4)}{\sqrt{c + dx^2}(e + fx^2)} dx = \int \frac{(Cx^4 + Bx^2 + A)\sqrt{bx^2 + a}}{\sqrt{dx^2 + c}(fx^2 + e)} dx$$

input `integrate((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)^(1/2)/(f*x^2+e),x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)*sqrt(b*x^2 + a)/(sqrt(d*x^2 + c)*(f*x^2 + e)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{\sqrt{c+dx^2}(e+fx^2)} dx = \int \frac{\sqrt{bx^2+a}(Cx^4+Bx^2+A)}{\sqrt{dx^2+c}(fx^2+e)} dx$$

input

```
int(((a + b*x^2)^(1/2)*(A + B*x^2 + C*x^4))/((c + d*x^2)^(1/2)*(e + f*x^2)),x)
```

output

```
int(((a + b*x^2)^(1/2)*(A + B*x^2 + C*x^4))/((c + d*x^2)^(1/2)*(e + f*x^2)), x)
```

Reduce [F]

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{\sqrt{c+dx^2}(e+fx^2)} dx$$

$$= \frac{\sqrt{dx^2+c}\sqrt{bx^2+a}cx + \left(\int \frac{\sqrt{dx^2+c}\sqrt{bx^2+a}x^4}{bdfx^6+adf x^4+bcf x^4+bde x^4+acf x^2+ade x^2+bce x^2+ace} dx \right) acdf + 3 \left(\int \frac{\sqrt{dx^2+c}\sqrt{bx^2+a}x^4}{bdf x^6+adf x^4+bcf x^4+bde x^4+acf x^2+ade x^2+bce x^2+ace} dx \right)}{\dots}$$

input

```
int((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)^(1/2)/(f*x^2+e),x)
```

output

```
(sqrt(c + d*x**2)*sqrt(a + b*x**2)*c*x + int((sqrt(c + d*x**2)*sqrt(a + b*
x**2)*x**4)/(a*c*e + a*c*f*x**2 + a*d*e*x**2 + a*d*f*x**4 + b*c*e*x**2 + b
*c*f*x**4 + b*d*e*x**4 + b*d*f*x**6),x)*a*c*d*f + 3*int((sqrt(c + d*x**2)*
sqrt(a + b*x**2)*x**4)/(a*c*e + a*c*f*x**2 + a*d*e*x**2 + a*d*f*x**4 + b*c
*e*x**2 + b*c*f*x**4 + b*d*e*x**4 + b*d*f*x**6),x)*b**2*d*f - 2*int((sqrt(
c + d*x**2)*sqrt(a + b*x**2)*x**4)/(a*c*e + a*c*f*x**2 + a*d*e*x**2 + a*d*
f*x**4 + b*c*e*x**2 + b*c*f*x**4 + b*d*e*x**4 + b*d*f*x**6),x)*b*c**2*f -
3*int((sqrt(c + d*x**2)*sqrt(a + b*x**2)*x**4)/(a*c*e + a*c*f*x**2 + a*d*e
*x**2 + a*d*f*x**4 + b*c*e*x**2 + b*c*f*x**4 + b*d*e*x**4 + b*d*f*x**6),x)
*b*c*d*e + 6*int((sqrt(c + d*x**2)*sqrt(a + b*x**2)*x**2)/(a*c*e + a*c*f*x
**2 + a*d*e*x**2 + a*d*f*x**4 + b*c*e*x**2 + b*c*f*x**4 + b*d*e*x**4 + b*d
*f*x**6),x)*a*b*d*f - int((sqrt(c + d*x**2)*sqrt(a + b*x**2)*x**2)/(a*c*e
+ a*c*f*x**2 + a*d*e*x**2 + a*d*f*x**4 + b*c*e*x**2 + b*c*f*x**4 + b*d*e*x
**4 + b*d*f*x**6),x)*a*c**2*f - 2*int((sqrt(c + d*x**2)*sqrt(a + b*x**2)*x
**2)/(a*c*e + a*c*f*x**2 + a*d*e*x**2 + a*d*f*x**4 + b*c*e*x**2 + b*c*f*x*
**4 + b*d*e*x**4 + b*d*f*x**6),x)*a*c*d*e - 2*int((sqrt(c + d*x**2)*sqrt(a
+ b*x**2)*x**2)/(a*c*e + a*c*f*x**2 + a*d*e*x**2 + a*d*f*x**4 + b*c*e*x**2
+ b*c*f*x**4 + b*d*e*x**4 + b*d*f*x**6),x)*b*c**2*e + 3*int((sqrt(c + d*x
**2)*sqrt(a + b*x**2))/(a*c*e + a*c*f*x**2 + a*d*e*x**2 + a*d*f*x**4 + b*c
*e*x**2 + b*c*f*x**4 + b*d*e*x**4 + b*d*f*x**6),x)*a**2*d*f - int((sqrt...
```

$$3.11 \quad \int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{3/2}(e+fx^2)} dx$$

Optimal result	124
Mathematica [C] (verified)	125
Rubi [A] (verified)	126
Maple [B] (verified)	127
Fricas [F(-1)]	128
Sympy [F]	129
Maxima [F]	129
Giac [F]	129
Mupad [F(-1)]	130
Reduce [F]	130

Optimal result

Integrand size = 44, antiderivative size = 481

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{3/2}(e+fx^2)} dx = \frac{(c^2C - Bcd + Ad^2) x\sqrt{a+bx^2}}{cd(de - cf)\sqrt{c+dx^2}} - \frac{b(2c^2Cf + Ad^2f - cd(Ce + Bf)) x\sqrt{c+dx^2}}{cd^2f(de - cf)\sqrt{a+bx^2}} + \frac{\sqrt{a}\sqrt{b}(2c^2Cf + Ad^2f - cd(Ce + Bf))\sqrt{c+dx^2}E\left(\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) \mid 1 - \frac{ad}{bc}\right)}{cd^2f(de - cf)\sqrt{a+bx^2}\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}} + \frac{a^{3/2}C\sqrt{c+dx^2}\text{EllipticF}\left(\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right), 1 - \frac{ad}{bc}\right)}{\sqrt{bcd}f\sqrt{a+bx^2}\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}} - \frac{a^{3/2}(Ce^2 - Bef + Af^2)\sqrt{c+dx^2}\text{EllipticPi}\left(1 - \frac{af}{be}, \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right), 1 - \frac{ad}{bc}\right)}{\sqrt{bcef}(de - cf)\sqrt{a+bx^2}\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}}$$

output

```
(A*d^2-B*c*d+C*c^2)*x*(b*x^2+a)^(1/2)/c/d/(-c*f+d*e)/(d*x^2+c)^(1/2)-b*(2*c^2*C*f+A*d^2*f-c*d*(B*f+C*e))*x*(d*x^2+c)^(1/2)/c/d^2/f/(-c*f+d*e)/(b*x^2+a)^(1/2)+a^(1/2)*b^(1/2)*(2*c^2*C*f+A*d^2*f-c*d*(B*f+C*e))*(d*x^2+c)^(1/2)*EllipticE(b^(1/2)*x/a^(1/2)/(1+b*x^2/a)^(1/2),(1-a*d/b/c)^(1/2))/c/d^2/f/(-c*f+d*e)/(b*x^2+a)^(1/2)/(a*(d*x^2+c)/c/(b*x^2+a))^(1/2)+a^(3/2)*C*(d*x^2+c)^(1/2)*InverseJacobiAM(arctan(b^(1/2)*x/a^(1/2)),(1-a*d/b/c)^(1/2))/b^(1/2)/c/d/f/(b*x^2+a)^(1/2)/(a*(d*x^2+c)/c/(b*x^2+a))^(1/2)-a^(3/2)*(A*f^2-B*e*f+C*e^2)*(d*x^2+c)^(1/2)*EllipticPi(b^(1/2)*x/a^(1/2)/(1+b*x^2/a)^(1/2),1-a*f/b/e,(1-a*d/b/c)^(1/2))/b^(1/2)/c/e/f/(-c*f+d*e)/(b*x^2+a)^(1/2)/(a*(d*x^2+c)/c/(b*x^2+a))^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.69 (sec) , antiderivative size = 368, normalized size of antiderivative = 0.77

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{3/2}(e+fx^2)} dx = \frac{-ibcef(2c^2Cf+Ad^2f-cd(Ce+Bf))\sqrt{1+\frac{bx^2}{a}}\sqrt{1+\frac{dx^2}{c}}E\left(\operatorname{arcsinh}\left(\frac{bx}{\sqrt{a}}\right)\right)}{(c+dx^2)^{3/2}(e+fx^2)}$$

input

```
Integrate[(Sqrt[a + b*x^2]*(A + B*x^2 + C*x^4))/((c + d*x^2)^(3/2)*(e + f*x^2)),x]
```

output

```
((-I)*b*c*e*f*(2*c^2*C*f + A*d^2*f - c*d*(C*e + B*f))*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - I*c*e*(-(d*e) + c*f)*(b*B*d*f + a*C*d*f - b*C*(d*e + 2*c*f))*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - d*(Sqrt[b/a]*(c^2*C - B*c*d + A*d^2)*e*f^2*x*(a + b*x^2) + I*c*d*(-(b*e) + a*f)*(C*e^2 + f*(-(B*e) + A*f))*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticPi[(a*f)/(b*e), I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)))/(Sqrt[b/a]*c*d^2*e*f^2*(-(d*e) + c*f)*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])
```

Rubi [A] (verified)

Time = 1.27 (sec) , antiderivative size = 550, normalized size of antiderivative = 1.14, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.045$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{3/2}(e+fx^2)} dx$$

↓ 7276

$$\int \left(\frac{\sqrt{a+bx^2}(Af^2 - Bef + Ce^2)}{f^2(c+dx^2)^{3/2}(e+fx^2)} - \frac{\sqrt{a+bx^2}(Ce - Bf)}{f^2(c+dx^2)^{3/2}} + \frac{Cx^2\sqrt{a+bx^2}}{f(c+dx^2)^{3/2}} \right) dx$$

↓ 2009

$$\frac{a^{3/2}\sqrt{c+dx^2}(Af^2 - Bef + Ce^2) \text{EllipticPi}\left(1 - \frac{af}{be}, \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right), 1 - \frac{ad}{bc}\right) + \sqrt{bc}ef\sqrt{a+bx^2}(de - cf)\sqrt{\frac{c(a+bx^2)}{c(a+bx^2)}}}{\sqrt{d}\sqrt{a+bx^2}(Af^2 - Bef + Ce^2) E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right) - \sqrt{cf^2}\sqrt{c+dx^2}(de - cf)\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{\sqrt{a+bx^2}(Ce - Bf)E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{c}\sqrt{d}f^2\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{\sqrt{c}C\sqrt{a+bx^2} \text{EllipticF}\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right)}{d^{3/2}f\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} - \frac{2\sqrt{c}C\sqrt{a+bx^2}E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{d^{3/2}f\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{Cx\sqrt{a+bx^2}}{df\sqrt{c+dx^2}}$$

input

```
Int[(Sqrt[a + b*x^2]*(A + B*x^2 + C*x^4))/((c + d*x^2)^(3/2)*(e + f*x^2)), x]
```

output

```
(C*x*Sqrt[a + b*x^2])/(d*f*Sqrt[c + d*x^2]) - (2*Sqrt[c]*C*Sqrt[a + b*x^2]
*EllipticE[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(d^(3/2)*f*Sqrt[
(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) - ((C*e - B*f)*Sqrt[a +
b*x^2]*EllipticE[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(Sqrt[c]*S
qrt[d]*f^2*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) + (Sqrt[
d]*(C*e^2 - B*e*f + A*f^2)*Sqrt[a + b*x^2]*EllipticE[ArcTan[(Sqrt[d]*x)/Sq
rt[c]], 1 - (b*c)/(a*d)]/(Sqrt[c]*f^2*(d*e - c*f)*Sqrt[(c*(a + b*x^2))/(a
*(c + d*x^2))]*Sqrt[c + d*x^2]) + (Sqrt[c]*C*Sqrt[a + b*x^2]*EllipticF[Arc
Tan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(d^(3/2)*f*Sqrt[(c*(a + b*x^2)
)/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) - (a^(3/2)*(C*e^2 - B*e*f + A*f^2)*Sqr
t[c + d*x^2]*EllipticPi[1 - (a*f)/(b*e), ArcTan[(Sqrt[b]*x)/Sqrt[a]], 1 -
(a*d)/(b*c)]/(Sqrt[b]*c*e*f*(d*e - c*f)*Sqrt[a + b*x^2]*Sqrt[(a*(c + d*x^
2))/(c*(a + b*x^2))])
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 7276

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1299 vs. $2(459) = 918$.

Time = 6.56 (sec) , antiderivative size = 1300, normalized size of antiderivative = 2.70

method	result	size
default	Expression too large to display	1300
elliptic	Expression too large to display	1542

input

```
int((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)^(3/2)/(f*x^2+e),x,method=_RE
TURNVERBOSE)
```



```

output (-A*(-b/a)^(1/2)*b*d^3*e*f^2*x^3+B*(-b/a)^(1/2)*b*c*d^2*e*f^2*x^3-C*(-b/a)
^(1/2)*b*c^2*d*e*f^2*x^3+A*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*Ellipti
cE(x*(-b/a)^(1/2),(a*d/b/c)^(1/2))*b*c*d^2*e*f^2+A*((b*x^2+a)/a)^(1/2)*((d
*x^2+c)/c)^(1/2)*EllipticPi(x*(-b/a)^(1/2),a*f/b/e,(-1/c*d)^(1/2)/(-b/a)^(
1/2))*a*c*d^2*f^3-A*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticPi(x*(
-b/a)^(1/2),a*f/b/e,(-1/c*d)^(1/2)/(-b/a)^(1/2))*b*c*d^2*e*f^2-B*((b*x^2+a
)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE(x*(-b/a)^(1/2),(a*d/b/c)^(1/2))*b
*c^2*d*e*f^2-B*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticPi(x*(-b/a)
^(1/2),a*f/b/e,(-1/c*d)^(1/2)/(-b/a)^(1/2))*a*c*d^2*e*f^2+B*((b*x^2+a)/a)
^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticPi(x*(-b/a)^(1/2),a*f/b/e,(-1/c*d)^(1/2)
/(-b/a)^(1/2))*b*c*d^2*e^2*f+B*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*Ell
ipticF(x*(-b/a)^(1/2),(a*d/b/c)^(1/2))*b*c^2*d*e*f^2-B*((b*x^2+a)/a)^(1/2)
*((d*x^2+c)/c)^(1/2)*EllipticF(x*(-b/a)^(1/2),(a*d/b/c)^(1/2))*b*c*d^2*e^2
*f+2*C*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE(x*(-b/a)^(1/2),(a
*d/b/c)^(1/2))*b*c^3*e*f^2-C*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*Ellip
ticE(x*(-b/a)^(1/2),(a*d/b/c)^(1/2))*b*c^2*d*e^2*f+C*((b*x^2+a)/a)^(1/2)*((
d*x^2+c)/c)^(1/2)*EllipticPi(x*(-b/a)^(1/2),a*f/b/e,(-1/c*d)^(1/2)/(-b/a)
^(1/2))*a*c*d^2*e^2*f-C*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticPi
(x*(-b/a)^(1/2),a*f/b/e,(-1/c*d)^(1/2)/(-b/a)^(1/2))*b*c*d^2*e^3+C*((b*x^2
+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF(x*(-b/a)^(1/2),(a*d/b/c)^(1/...
    
```

Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{3/2}(e+fx^2)} dx = \text{Timed out}$$

```

input integrate((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)^(3/2)/(f*x^2+e),x, alg
orithm="fricas")
    
```

output Timed out

Sympy [F]

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{3/2}(e+fx^2)} dx = \int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{\frac{3}{2}}(e+fx^2)} dx$$

input `integrate((b*x**2+a)**(1/2)*(C*x**4+B*x**2+A)/(d*x**2+c)**(3/2)/(f*x**2+e),x)`

output `Integral(sqrt(a + b*x**2)*(A + B*x**2 + C*x**4)/((c + d*x**2)**(3/2)*(e + f*x**2)), x)`

Maxima [F]

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{3/2}(e+fx^2)} dx = \int \frac{(Cx^4+Bx^2+A)\sqrt{bx^2+a}}{(dx^2+c)^{\frac{3}{2}}(fx^2+e)} dx$$

input `integrate((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)*sqrt(b*x^2 + a)/((d*x^2 + c)^(3/2)*(f*x^2 + e)), x)`

Giac [F]

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{3/2}(e+fx^2)} dx = \int \frac{(Cx^4+Bx^2+A)\sqrt{bx^2+a}}{(dx^2+c)^{\frac{3}{2}}(fx^2+e)} dx$$

input `integrate((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm="giac")`

output

```
integrate((C*x^4 + B*x^2 + A)*sqrt(b*x^2 + a)/((d*x^2 + c)^(3/2)*(f*x^2 + e)), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + bx^2}(A + Bx^2 + Cx^4)}{(c + dx^2)^{3/2}(e + fx^2)} dx = \int \frac{\sqrt{bx^2 + a}(Cx^4 + Bx^2 + A)}{(dx^2 + c)^{3/2}(fx^2 + e)} dx$$

input

```
int(((a + b*x^2)^(1/2)*(A + B*x^2 + C*x^4))/((c + d*x^2)^(3/2)*(e + f*x^2)), x)
```

output

```
int(((a + b*x^2)^(1/2)*(A + B*x^2 + C*x^4))/((c + d*x^2)^(3/2)*(e + f*x^2)), x)
```

Reduce [F]

$$\int \frac{\sqrt{a + bx^2}(A + Bx^2 + Cx^4)}{(c + dx^2)^{3/2}(e + fx^2)} dx = \text{too large to display}$$

input

```
int((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)^(3/2)/(f*x^2+e), x)
```

output

```
(sqrt(c + d*x**2)*sqrt(a + b*x**2)*a*c*x + sqrt(c + d*x**2)*sqrt(a + b*x**
2)*b**2*x - 2*int((sqrt(c + d*x**2)*sqrt(a + b*x**2)*x**6)/(2*a*c**3*e*f +
2*a*c**3*f**2*x**2 + a*c**2*d*e**2 + 5*a*c**2*d*e*f*x**2 + 4*a*c**2*d*f**
2*x**4 + 2*a*c*d**2*e**2*x**2 + 4*a*c*d**2*e*f*x**4 + 2*a*c*d**2*f**2*x**6
+ a*d**3*e**2*x**4 + a*d**3*e*f*x**6 + 2*b*c**3*e*f*x**2 + 2*b*c**3*f**2*
x**4 + b*c**2*d*e**2*x**2 + 5*b*c**2*d*e*f*x**4 + 4*b*c**2*d*f**2*x**6 + 2
*b*c*d**2*e**2*x**4 + 4*b*c*d**2*e*f*x**6 + 2*b*c*d**2*f**2*x**8 + b*d**3*
e**2*x**6 + b*d**3*e*f*x**8),x)*a*b*c**3*d*f**2 - int((sqrt(c + d*x**2)*sq
rt(a + b*x**2)*x**6)/(2*a*c**3*e*f + 2*a*c**3*f**2*x**2 + a*c**2*d*e**2 +
5*a*c**2*d*e*f*x**2 + 4*a*c**2*d*f**2*x**4 + 2*a*c*d**2*e**2*x**2 + 4*a*c*
d**2*e*f*x**4 + 2*a*c*d**2*f**2*x**6 + a*d**3*e**2*x**4 + a*d**3*e*f*x**6
+ 2*b*c**3*e*f*x**2 + 2*b*c**3*f**2*x**4 + b*c**2*d*e**2*x**2 + 5*b*c**2*d
*e*f*x**4 + 4*b*c**2*d*f**2*x**6 + 2*b*c*d**2*e**2*x**4 + 4*b*c*d**2*e*f*x
**6 + 2*b*c*d**2*f**2*x**8 + b*d**3*e**2*x**6 + b*d**3*e*f*x**8),x)*a*b*c*
*2*d**2*e*f - 2*int((sqrt(c + d*x**2)*sqrt(a + b*x**2)*x**6)/(2*a*c**3*e*f
+ 2*a*c**3*f**2*x**2 + a*c**2*d*e**2 + 5*a*c**2*d*e*f*x**2 + 4*a*c**2*d*f
**2*x**4 + 2*a*c*d**2*e**2*x**2 + 4*a*c*d**2*e*f*x**4 + 2*a*c*d**2*f**2*x*
*6 + a*d**3*e**2*x**4 + a*d**3*e*f*x**6 + 2*b*c**3*e*f*x**2 + 2*b*c**3*f**
2*x**4 + b*c**2*d*e**2*x**2 + 5*b*c**2*d*e*f*x**4 + 4*b*c**2*d*f**2*x**6 +
2*b*c*d**2*e**2*x**4 + 4*b*c*d**2*e*f*x**6 + 2*b*c*d**2*f**2*x**8 + b...
```

$$3.12 \quad \int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{5/2}(e+fx^2)} dx$$

Optimal result	132
Mathematica [C] (verified)	133
Rubi [A] (verified)	134
Maple [B] (verified)	136
Fricas [F(-1)]	137
Sympy [F(-1)]	137
Maxima [F]	138
Giac [F]	138
Mupad [F(-1)]	138
Reduce [F]	139

Optimal result

Integrand size = 44, antiderivative size = 637

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{5/2}(e+fx^2)} dx = \frac{(c^2C - Bcd + Ad^2)x\sqrt{a+bx^2}}{3cd(de - cf)(c+dx^2)^{3/2}}$$

$$\frac{(ad(2Ad^3e + c^3Cf + cd^2(Be - 5Af)) - c^2(4Cde - 2Bdf)) - bc(Ad^3e + 2c^3Cf + 2cd^2(Be - 2Af)) - c^2(3b^2c^3d(Ce^2 - f(Be - Af)) + 3a^2cd^3(Ce^2 - f(Be - Af)) - ab(Ad^4e^2 + c^4Cf^2 - cd^3e(Be + 2Af)) - c^2(3c^{3/2}d^{3/2}(bc - ad)(de - cf)^2\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2})}{3a\sqrt{cd}^{3/2}(bc - ad)(de - cf)^3\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}}$$

$$+ \frac{c^{3/2}(be - af)(Ce^2 - Bef + Af^2)\sqrt{a+bx^2}\text{EllipticPi}\left(1 - \frac{cf}{de}, \arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right)}{a\sqrt{de}(de - cf)^3\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}}$$

output

```

1/3*(A*d^2-B*c*d+C*c^2)*x*(b*x^2+a)^(1/2)/c/d/(-c*f+d*e)/(d*x^2+c)^(3/2)-1
/3*(a*d*(2*A*d^3*e+c^3*C*f+c*d^2*(-5*A*f+B*e)-c^2*(-2*B*d*f+4*C*d*e))-b*c*
(A*d^3*e+2*c^3*C*f+2*c*d^2*(-2*A*f+B*e)-c^2*(-B*d*f+5*C*d*e)))*(b*x^2+a)^(
1/2)*EllipticE(d^(1/2)*x/c^(1/2)/(1+d*x^2/c)^(1/2),(1-b*c/a/d)^(1/2))/c^(3
/2)/d^(3/2)/(-a*d+b*c)/(-c*f+d*e)^2/(c*(b*x^2+a)/a/(d*x^2+c))^(1/2)/(d*x^2
+c)^(1/2)-1/3*(3*b^2*c^3*d*(C*e^2-f*(-A*f+B*e))+3*a^2*c*d^3*(C*e^2-f*(-A*f
+B*e))-a*b*(A*d^4*e^2+c^4*C*f^2-c*d^3*e*(2*A*f+B*e)-c^3*d*f*(B*f+2*C*e)+c^
2*d^2*(7*A*f^2-4*B*e*f+7*C*e^2)))*(b*x^2+a)^(1/2)*InverseJacobiAM(arctan(d
^(1/2)*x/c^(1/2),(1-b*c/a/d)^(1/2))/a/c^(1/2)/d^(3/2)/(-a*d+b*c)/(-c*f+d*
e)^3/(c*(b*x^2+a)/a/(d*x^2+c))^(1/2)/(d*x^2+c)^(1/2)+c^(3/2)*(-a*f+b*e)*(A
*f^2-B*e*f+C*e^2)*(b*x^2+a)^(1/2)*EllipticPi(d^(1/2)*x/c^(1/2)/(1+d*x^2/c)
^(1/2),1-c*f/d/e,(1-b*c/a/d)^(1/2))/a/d^(1/2)/e/(-c*f+d*e)^3/(c*(b*x^2+a)/
a/(d*x^2+c))^(1/2)/(d*x^2+c)^(1/2)

```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 10.95 (sec) , antiderivative size = 3341, normalized size of antiderivative = 5.24

$$\int \frac{\sqrt{a + bx^2}(A + Bx^2 + Cx^4)}{(c + dx^2)^{5/2}(e + fx^2)} dx = \text{Result too large to show}$$

input

```

Integrate[(Sqrt[a + b*x^2]*(A + B*x^2 + C*x^4))/((c + d*x^2)^(5/2)*(e + f*
x^2)),x]

```

output

```
(4*a*b*Sqrt[b/a]*c^4*C*d^2*e^2*f*x - a*b*Sqrt[b/a]*B*c^3*d^3*e^2*f*x - 3*a^2*Sqrt[b/a]*c^3*C*d^3*e^2*f*x - 2*a*A*b*Sqrt[b/a]*c^2*d^4*e^2*f*x + 3*a^2*A*Sqrt[b/a]*c*d^5*e^2*f*x - a*b*Sqrt[b/a]*c^5*C*d*e*f^2*x - 2*a*b*Sqrt[b/a]*B*c^4*d^2*e*f^2*x + 5*a*A*b*Sqrt[b/a]*c^3*d^3*e*f^2*x + 3*a^2*Sqrt[b/a]*B*c^3*d^3*e*f^2*x - 6*a^2*A*Sqrt[b/a]*c^2*d^4*e*f^2*x + 4*a*b*(b/a)^(3/2)*c^4*C*d^2*e^2*f*x^3 - a*b*(b/a)^(3/2)*B*c^3*d^3*e^2*f*x^3 + 2*a*b*Sqrt[b/a]*c^3*C*d^3*e^2*f*x^3 - 2*a*A*b*(b/a)^(3/2)*c^2*d^4*e^2*f*x^3 - 2*a*b*Sqrt[b/a]*B*c^2*d^4*e^2*f*x^3 - 4*a^2*Sqrt[b/a]*c^2*C*d^4*e^2*f*x^3 + 2*a*A*b*Sqrt[b/a]*c*d^5*e^2*f*x^3 + a^2*Sqrt[b/a]*B*c*d^5*e^2*f*x^3 + 2*a^2*A*Sqrt[b/a]*d^6*e^2*f*x^3 - a*b*(b/a)^(3/2)*c^5*C*d*e*f^2*x^3 - 2*a*b*(b/a)^(3/2)*B*c^4*d^2*e*f^2*x^3 - 2*a*b*Sqrt[b/a]*c^4*C*d^2*e*f^2*x^3 + 5*a*A*b*(b/a)^(3/2)*c^3*d^3*e*f^2*x^3 + 2*a*b*Sqrt[b/a]*B*c^3*d^3*e*f^2*x^3 + a^2*Sqrt[b/a]*c^3*C*d^3*e*f^2*x^3 - 2*a*A*b*Sqrt[b/a]*c^2*d^4*e*f^2*x^3 + 2*a^2*Sqrt[b/a]*B*c^2*d^4*e*f^2*x^3 - 5*a^2*A*Sqrt[b/a]*c*d^5*e*f^2*x^3 + 5*a*b*(b/a)^(3/2)*c^3*C*d^3*e^2*f*x^5 - 2*a*b*(b/a)^(3/2)*B*c^2*d^4*e^2*f*x^5 - 4*a*b*Sqrt[b/a]*c^2*C*d^4*e^2*f*x^5 - a*A*b*(b/a)^(3/2)*c*d^5*e^2*f*x^5 + a*b*Sqrt[b/a]*B*c*d^5*e^2*f*x^5 + 2*a*A*b*Sqrt[b/a]*d^6*e^2*f*x^5 - 2*a*b*(b/a)^(3/2)*c^4*C*d^2*e*f^2*x^5 - a*b*(b/a)^(3/2)*B*c^3*d^3*e*f^2*x^5 + a*b*Sqrt[b/a]*c^3*C*d^3*e*f^2*x^5 + 4*a*A*b*(b/a)^(3/2)*c^2*d^4*e*f^2*x^5 + 2*a*b*Sqrt[b/a]*B*c^2*d^4*e*f^2*x^5 - 5*a*A*b*Sqrt[b/a]*c*d^5*e*f^2*x^5 ...
```

Rubi [A] (verified)

Time = 1.93 (sec) , antiderivative size = 979, normalized size of antiderivative = 1.54, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.045$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{5/2}(e+fx^2)} dx$$

$$\downarrow 7276$$

$$\int \left(\frac{\sqrt{a+bx^2}(Af^2 - Bef + Ce^2)}{f^2(c+dx^2)^{5/2}(e+fx^2)} - \frac{\sqrt{a+bx^2}(Ce - Bf)}{f^2(c+dx^2)^{5/2}} + \frac{Cx^2\sqrt{a+bx^2}}{f(c+dx^2)^{5/2}} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
& \frac{(Ce^2 - Bfe + Af^2) \sqrt{dx^2 + c} \operatorname{EllipticPi} \left(1 - \frac{af}{be}, \arctan \left(\frac{\sqrt{bx}}{\sqrt{a}} \right), 1 - \frac{ad}{bc} \right) a^{3/2}}{\sqrt{bce}(de - cf)^2 \sqrt{bx^2 + a} \sqrt{\frac{a(dx^2+c)}{c(bx^2+a)}}} \\
& \frac{(bc - 2ad)(Ce - Bf) \sqrt{bx^2 + a} E \left(\arctan \left(\frac{\sqrt{dx}}{\sqrt{c}} \right) \mid 1 - \frac{bc}{ad} \right)}{3c^{3/2} \sqrt{d}(bc - ad) f^2 \sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}} \sqrt{dx^2 + c}} \\
& \frac{\sqrt{d}(Ce^2 - Bfe + Af^2) (ad(2de - 5cf) - bc(de - 4cf)) \sqrt{bx^2 + a} E \left(\arctan \left(\frac{\sqrt{dx}}{\sqrt{c}} \right) \mid 1 - \frac{bc}{ad} \right)}{3c^{3/2}(bc - ad) f^2 (de - cf)^2 \sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}} \sqrt{dx^2 + c}} + \\
& \frac{C(2bc - ad) \sqrt{bx^2 + a} E \left(\arctan \left(\frac{\sqrt{dx}}{\sqrt{c}} \right) \mid 1 - \frac{bc}{ad} \right)}{3\sqrt{cd}^{3/2}(bc - ad) f \sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}} \sqrt{dx^2 + c}} \\
& \frac{b(Ce - Bf) \sqrt{bx^2 + a} \operatorname{EllipticF} \left(\arctan \left(\frac{\sqrt{dx}}{\sqrt{c}} \right), 1 - \frac{bc}{ad} \right)}{3\sqrt{c} \sqrt{d}(bc - ad) f^2 \sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}} \sqrt{dx^2 + c}} + \\
& \frac{b\sqrt{d}(Ce^2 - Bfe + Af^2) \sqrt{bx^2 + a} \operatorname{EllipticF} \left(\arctan \left(\frac{\sqrt{dx}}{\sqrt{c}} \right), 1 - \frac{bc}{ad} \right)}{3\sqrt{c}(bc - ad) f^2 (de - cf) \sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}} \sqrt{dx^2 + c}} \\
& \frac{b\sqrt{c} C \sqrt{bx^2 + a} \operatorname{EllipticF} \left(\arctan \left(\frac{\sqrt{dx}}{\sqrt{c}} \right), 1 - \frac{bc}{ad} \right)}{3d^{3/2}(bc - ad) f \sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}} \sqrt{dx^2 + c}} - \frac{(Ce - Bf) x \sqrt{bx^2 + a}}{3cf^2 (dx^2 + c)^{3/2}} + \\
& \frac{d(Ce^2 - Bfe + Af^2) x \sqrt{bx^2 + a}}{3cf^2 (de - cf) (dx^2 + c)^{3/2}} - \frac{Cx \sqrt{bx^2 + a}}{3df (dx^2 + c)^{3/2}}
\end{aligned}$$

input

```
Int[(Sqrt[a + b*x^2]*(A + B*x^2 + C*x^4))/((c + d*x^2)^(5/2)*(e + f*x^2)),
x]
```


output

```

-1/3*(C*x*Sqrt[a + b*x^2])/(d*f*(c + d*x^2)^(3/2)) - ((C*e - B*f)*x*Sqrt[a
+ b*x^2])/(3*c*f^2*(c + d*x^2)^(3/2)) + (d*(C*e^2 - B*e*f + A*f^2)*x*Sqrt
[a + b*x^2])/(3*c*f^2*(d*e - c*f)*(c + d*x^2)^(3/2)) + (C*(2*b*c - a*d)*Sq
rt[a + b*x^2]*EllipticE[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)])/(3*
Sqrt[c]*d^(3/2)*(b*c - a*d)*f*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c
+ d*x^2]) - ((b*c - 2*a*d)*(C*e - B*f)*Sqrt[a + b*x^2]*EllipticE[ArcTan[(
Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)])/(3*c^(3/2)*Sqrt[d]*(b*c - a*d)*f^2*
Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) - (Sqrt[d]*(C*e^2 -
B*e*f + A*f^2)*(a*d*(2*d*e - 5*c*f) - b*c*(d*e - 4*c*f))*Sqrt[a + b*x^2]*
EllipticE[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)])/(3*c^(3/2)*(b*c -
a*d)*f^2*(d*e - c*f)^2*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x
^2]) - (b*Sqrt[c]*C*Sqrt[a + b*x^2]*EllipticF[ArcTan[(Sqrt[d]*x)/Sqrt[c]],
1 - (b*c)/(a*d)])/(3*d^(3/2)*(b*c - a*d)*f*Sqrt[(c*(a + b*x^2))/(a*(c + d
*x^2))]*Sqrt[c + d*x^2]) - (b*(C*e - B*f)*Sqrt[a + b*x^2]*EllipticF[ArcTan
[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)])/(3*Sqrt[c]*Sqrt[d]*(b*c - a*d)*f^
2*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) + (b*Sqrt[d]*(C*e
^2 - B*e*f + A*f^2)*Sqrt[a + b*x^2]*EllipticF[ArcTan[(Sqrt[d]*x)/Sqrt[c]],
1 - (b*c)/(a*d)])/(3*Sqrt[c]*(b*c - a*d)*f^2*(d*e - c*f)*Sqrt[(c*(a + b*x
^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) + (a^(3/2)*(C*e^2 - B*e*f + A*f^2)*
Sqrt[c + d*x^2]*EllipticPi[1 - (a*f)/(b*e), ArcTan[(Sqrt[b]*x)/Sqrt[a]]...

```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 7276

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 6012 vs. $2(613) = 1226$.

Time = 8.85 (sec) , antiderivative size = 6013, normalized size of antiderivative = 9.44

method	result	size
elliptic	Expression too large to display	6013
default	Expression too large to display	6326

input `int((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)^(5/2)/(f*x^2+e),x,method=_RE
TURNVERBOSE)`

output `result too large to display`

Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{5/2}(e+fx^2)} dx = \text{Timed out}$$

input `integrate((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)^(5/2)/(f*x^2+e),x,alg
orithm="fricas")`

output `Timed out`

Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{5/2}(e+fx^2)} dx = \text{Timed out}$$

input `integrate((b*x**2+a)**(1/2)*(C*x**4+B*x**2+A)/(d*x**2+c)**(5/2)/(f*x**2+e)
,x)`

output `Timed out`

Maxima [F]

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{5/2}(e+fx^2)} dx = \int \frac{(Cx^4+Bx^2+A)\sqrt{bx^2+a}}{(dx^2+c)^{5/2}(fx^2+e)} dx$$

input `integrate((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)^(5/2)/(f*x^2+e),x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)*sqrt(b*x^2 + a)/((d*x^2 + c)^(5/2)*(f*x^2 + e)), x)`

Giac [F]

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{5/2}(e+fx^2)} dx = \int \frac{(Cx^4+Bx^2+A)\sqrt{bx^2+a}}{(dx^2+c)^{5/2}(fx^2+e)} dx$$

input `integrate((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)^(5/2)/(f*x^2+e),x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)*sqrt(b*x^2 + a)/((d*x^2 + c)^(5/2)*(f*x^2 + e)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{5/2}(e+fx^2)} dx = \int \frac{\sqrt{bx^2+a}(Cx^4+Bx^2+A)}{(dx^2+c)^{5/2}(fx^2+e)} dx$$

input `int(((a + b*x^2)^(1/2)*(A + B*x^2 + C*x^4))/((c + d*x^2)^(5/2)*(e + f*x^2)),x)`

output

```
int(((a + b*x^2)^(1/2)*(A + B*x^2 + C*x^4))/((c + d*x^2)^(5/2)*(e + f*x^2)
), x)
```

Reduce [F]

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{5/2}(e+fx^2)} dx = \text{too large to display}$$

input

```
int((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)^(5/2)/(f*x^2+e),x)
```

output

```
( - sqrt(c + d*x**2)*sqrt(a + b*x**2)*a*c*x - sqrt(c + d*x**2)*sqrt(a + b*
x**2)*b**2*x + 2*int((sqrt(c + d*x**2)*sqrt(a + b*x**2)*x**6)/(2*a**2*c**3
*d*e*f + 2*a**2*c**3*d*f**2*x**2 + 6*a**2*c**2*d**2*e*f*x**2 + 6*a**2*c**2
*d**2*f**2*x**4 + 6*a**2*c*d**3*e*f*x**4 + 6*a**2*c*d**3*f**2*x**6 + 2*a**
2*d**4*e*f*x**6 + 2*a**2*d**4*f**2*x**8 - 2*a*b*c**4*e*f - 2*a*b*c**4*f**2
*x**2 + a*b*c**3*d*e**2 - 3*a*b*c**3*d*e*f*x**2 - 4*a*b*c**3*d*f**2*x**4 +
3*a*b*c**2*d**2*e**2*x**2 + 3*a*b*c**2*d**2*e*f*x**4 + 3*a*b*c*d**3*e**2*
x**4 + 7*a*b*c*d**3*e*f*x**6 + 4*a*b*c*d**3*f**2*x**8 + a*b*d**4*e**2*x**6
+ 3*a*b*d**4*e*f*x**8 + 2*a*b*d**4*f**2*x**10 - 2*b**2*c**4*e*f*x**2 - 2*
b**2*c**4*f**2*x**4 + b**2*c**3*d*e**2*x**2 - 5*b**2*c**3*d*e*f*x**4 - 6*b
**2*c**3*d*f**2*x**6 + 3*b**2*c**2*d**2*e**2*x**4 - 3*b**2*c**2*d**2*e*f*x
**6 - 6*b**2*c**2*d**2*f**2*x**8 + 3*b**2*c*d**3*e**2*x**6 + b**2*c*d**3*e
*f*x**8 - 2*b**2*c*d**3*f**2*x**10 + b**2*d**4*e**2*x**8 + b**2*d**4*e*f*x
**10),x)*a**2*b*c**3*d**2*f**2 + 4*int((sqrt(c + d*x**2)*sqrt(a + b*x**2)*
x**6)/(2*a**2*c**3*d*e*f + 2*a**2*c**3*d*f**2*x**2 + 6*a**2*c**2*d**2*e*f*
x**2 + 6*a**2*c**2*d**2*f**2*x**4 + 6*a**2*c*d**3*e*f*x**4 + 6*a**2*c*d**3
*f**2*x**6 + 2*a**2*d**4*e*f*x**6 + 2*a**2*d**4*f**2*x**8 - 2*a*b*c**4*e*f
- 2*a*b*c**4*f**2*x**2 + a*b*c**3*d*e**2 - 3*a*b*c**3*d*e*f*x**2 - 4*a*b*
c**3*d*f**2*x**4 + 3*a*b*c**2*d**2*e**2*x**2 + 3*a*b*c**2*d**2*e*f*x**4 +
3*a*b*c*d**3*e**2*x**4 + 7*a*b*c*d**3*e*f*x**6 + 4*a*b*c*d**3*f**2*x**8...
```

3.13
$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{7/2}(e+fx^2)} dx$$

Optimal result	140
Mathematica [C] (verified)	141
Rubi [A] (verified)	142
Maple [B] (verified)	144
Fricas [F(-1)]	145
Sympy [F(-1)]	145
Maxima [F]	146
Giac [F]	146
Mupad [F(-1)]	146
Reduce [F]	147

Optimal result

Integrand size = 44, antiderivative size = 1093

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{7/2}(e+fx^2)} dx = \text{Too large to display}$$

output

```

1/5*(A*d^2-B*c*d+C*c^2)*x*(b*x^2+a)^(1/2)/c/d/(-c*f+d*e)/(d*x^2+c)^(5/2)-1
/15*(a*d*(4*A*d^3*e+c^3*C*f+c*d^2*(-9*A*f+B*e)-c^2*(-4*B*d*f+6*C*d*e))-b*c
*(3*A*d^3*e+2*c^3*C*f+2*c*d^2*(-4*A*f+B*e)-c^2*(-3*B*d*f+7*C*d*e)))*x*(b*x
^2+a)^(1/2)/c^2/d/(-a*d+b*c)/(-c*f+d*e)^2/(d*x^2+c)^(3/2)+1/15*(a^2*d^2*(8
*A*d^4*e^2-2*c^4*C*f^2+2*c*d^3*e*(-13*A*f+B*e)+2*c^3*d*f*(-4*B*f+7*C*e)+3*
c^2*d^2*(11*A*f^2-3*B*e*f+C*e^2))+b^2*c^2*(3*A*d^4*e^2-2*c^4*C*f^2+c*d^3*e
*(-11*A*f+2*B*e)+3*c^3*d*f*(-B*f+3*C*e)+c^2*d^2*(23*A*f^2-14*B*e*f+8*C*e^2
))-a*b*c*d*(13*A*d^4*e^2-2*c^4*C*f^2+c*d^3*e*(-41*A*f+2*B*e)+c^3*d*f*(-13*
B*f+19*C*e)+c^2*d^2*(58*A*f^2-19*B*e*f+13*C*e^2)))*(b*x^2+a)^(1/2)*Ellipti
cE(d^(1/2)*x/c^(1/2)/(1+d*x^2/c)^(1/2),(1-b*c/a/d)^(1/2))/c^(5/2)/d^(3/2)/
(-a*d+b*c)^2/(-c*f+d*e)^3/(c*(b*x^2+a)/a/(d*x^2+c))^(1/2)/(d*x^2+c)^(1/2)+
1/15*(15*b^3*c^5*d*f*(C*e^2-f*(-A*f+B*e))-15*a^3*c^2*d^4*f*(C*e^2-f*(-A*f+
B*e))-a^2*b*d*(4*A*d^5*e^3+c^5*C*f^3+c*d^4*e^2*(-17*A*f+B*e)-4*c^4*d*f^2*(
-B*f+2*C*e)-2*c^2*d^3*e*(-11*A*f^2-B*e*f+3*C*e^2)-2*c^3*d^2*f*(27*A*f^2-19
*B*e*f+16*C*e^2))+a*b^2*c*(6*A*d^5*e^3-c^5*C*f^3-c*d^4*e^2*(23*A*f+B*e)-2*
c^4*d*f^2*(-3*B*f+C*e)-2*c^3*d^2*f*(19*C*e^2-4*f*(-7*A*f+4*B*e))-4*c^2*d^3
*e*(C*e^2-f*(7*A*f+2*B*e))))*(b*x^2+a)^(1/2)*InverseJacobiAM(arctan(d^(1/2)
)*x/c^(1/2),(1-b*c/a/d)^(1/2))/a/c^(3/2)/d^(3/2)/(-a*d+b*c)^2/(-c*f+d*e)^
4/(c*(b*x^2+a)/a/(d*x^2+c))^(1/2)/(d*x^2+c)^(1/2)-c^(3/2)*f*(-a*f+b*e)*(A*
f^2-B*e*f+C*e^2)*(b*x^2+a)^(1/2)*EllipticPi(d^(1/2)*x/c^(1/2)/(1+d*x^2/...

```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 12.76 (sec) , antiderivative size = 1031, normalized size of antiderivative = 0.94

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{7/2}(e+fx^2)} dx = \frac{\sqrt{\frac{b}{a}} \operatorname{dex}(a+bx^2) \left(3c^2(bc-ad)^2(c^2C-Bcd+Ad^2)(de-cf)^2 - c(bc-ad)^2 \right)}{(c+dx^2)^{7/2}(e+fx^2)}$$

input

```

Integrate[(Sqrt[a + b*x^2]*(A + B*x^2 + C*x^4))/((c + d*x^2)^(7/2)*(e + f*
x^2)), x]

```

output

```
(Sqrt[b/a]*d*e*x*(a + b*x^2)*(3*c^2*(b*c - a*d)^2*(c^2*C - B*c*d + A*d^2)*
(d*e - c*f)^2 - c*(b*c - a*d)*(d*e - c*f)*(-(b*c*(3*A*d^3*e + 2*c^3*C*f +
2*c*d^2*(B*e - 4*A*f) + c^2*(-7*C*d*e + 3*B*d*f))) + a*d*(4*A*d^3*e + c^3*
C*f + c*d^2*(B*e - 9*A*f) + c^2*(-6*C*d*e + 4*B*d*f)))*(c + d*x^2) + (a*b*
c*d*(-13*A*d^4*e^2 + 2*c^4*C*f^2 + c*d^3*e*(-2*B*e + 41*A*f) + c^3*d*f*(-1
9*C*e + 13*B*f) + c^2*d^2*(-13*C*e^2 + 19*B*e*f - 58*A*f^2)) + a^2*d^2*(8*
A*d^4*e^2 - 2*c^4*C*f^2 + 2*c*d^3*e*(B*e - 13*A*f) + 2*c^3*d*f*(7*C*e - 4*
B*f) + 3*c^2*d^2*(C*e^2 - 3*B*e*f + 11*A*f^2)) + b^2*c^2*(3*A*d^4*e^2 - 2*
c^4*C*f^2 + c*d^3*e*(2*B*e - 11*A*f) - 3*c^3*d*f*(-3*C*e + B*f) + c^2*d^2*
(8*C*e^2 - 14*B*e*f + 23*A*f^2)))*(c + d*x^2)^2) + I*c*Sqrt[1 + (b*x^2)/a]
*(c + d*x^2)^2*Sqrt[1 + (d*x^2)/c]*(b*e*(a*b*c*d*(-13*A*d^4*e^2 + 2*c^4*C*
f^2 + c*d^3*e*(-2*B*e + 41*A*f) + c^3*d*f*(-19*C*e + 13*B*f) + c^2*d^2*(-1
3*C*e^2 + 19*B*e*f - 58*A*f^2)) + a^2*d^2*(8*A*d^4*e^2 - 2*c^4*C*f^2 + 2*c
*d^3*e*(B*e - 13*A*f) + 2*c^3*d*f*(7*C*e - 4*B*f) + 3*c^2*d^2*(C*e^2 - 3*B
*e*f + 11*A*f^2)) + b^2*c^2*(3*A*d^4*e^2 - 2*c^4*C*f^2 + c*d^3*e*(2*B*e -
11*A*f) - 3*c^3*d*f*(-3*C*e + B*f) + c^2*d^2*(8*C*e^2 - 14*B*e*f + 23*A*f^
2)))*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] + (b*c - a*d)*(b*e*(-(
d*e) + c*f)*(b*c*(3*A*d^3*e + 2*c^3*C*f + 2*c*d^2*(B*e - 4*A*f) + c^2*(-7*
C*d*e + 3*B*d*f)) - a*d*(4*A*d^3*e + c^3*C*f + c*d^2*(B*e - 9*A*f) + c^2*(-
6*C*d*e + 4*B*d*f)))*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - ...
```

Rubi [A] (verified)

Time = 2.74 (sec) , antiderivative size = 1396, normalized size of antiderivative = 1.28, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.045$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{7/2}(e+fx^2)} dx$$

$$\downarrow 7276$$

$$\int \left(\frac{\sqrt{a+bx^2}(Af^2 - Bef + Ce^2)}{f^2(c+dx^2)^{7/2}(e+fx^2)} - \frac{\sqrt{a+bx^2}(Ce - Bf)}{f^2(c+dx^2)^{7/2}} + \frac{Cx^2\sqrt{a+bx^2}}{f(c+dx^2)^{7/2}} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
& \frac{f(Ce^2 - Bfe + Af^2) \sqrt{dx^2 + c} \operatorname{EllipticPi}\left(1 - \frac{af}{be}, \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right), 1 - \frac{ad}{bc}\right) a^{3/2}}{\sqrt{bce}(de - cf)^3 \sqrt{bx^2 + a} \sqrt{\frac{a(dx^2+c)}{c(bx^2+a)}}} \\
& \frac{(3b^2c^2 - 13abdc + 8a^2d^2)(Ce - Bf) \sqrt{bx^2 + a} E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{15c^{5/2} \sqrt{d}(bc - ad)^2 f^2 \sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}} \sqrt{dx^2 + c}} + \\
& \frac{\sqrt{d}(Ce^2 - Bfe + Af^2) \sqrt{bx^2 + a} E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{c}(de - cf)^3 \sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}} \sqrt{dx^2 + c}} \\
& \frac{\sqrt{d}(Ce^2 - Bfe + Af^2) (-b^2(3de - 8cf)c^2 + abd(13de - 28cf)c - 2a^2d^2(4de - 9cf)) \sqrt{bx^2 + a} E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\right)}{15c^{5/2}(bc - ad)^2 f^2 (de - cf)^2 \sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}} \sqrt{dx^2 + c}} \\
& \frac{2C(b^2c^2 - abdc + a^2d^2) \sqrt{bx^2 + a} E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{15c^{3/2} d^{3/2} (bc - ad)^2 f \sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}} \sqrt{dx^2 + c}} \\
& \frac{2b(3bc - 2ad)(Ce - Bf) \sqrt{bx^2 + a} \operatorname{EllipticF}\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right)}{15c^{3/2} \sqrt{d}(bc - ad)^2 f^2 \sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}} \sqrt{dx^2 + c}} + \\
& \frac{b\sqrt{d}(Ce^2 - Bfe + Af^2) (bc(6de - 11cf) - ad(4de - 9cf)) \sqrt{bx^2 + a} \operatorname{EllipticF}\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right)}{15c^{3/2}(bc - ad)^2 f^2 (de - cf)^2 \sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}} \sqrt{dx^2 + c}} \\
& \frac{bC(bc + ad) \sqrt{bx^2 + a} \operatorname{EllipticF}\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right)}{15\sqrt{cd}^{3/2}(bc - ad)^2 f \sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}} \sqrt{dx^2 + c}} - \frac{(3bc - 4ad)(Ce - Bf)x \sqrt{bx^2 + a}}{15c^2(bc - ad) f^2 (dx^2 + c)^{3/2}} \\
& \frac{d(Ce^2 - Bfe + Af^2) (ad(4de - 9cf) - bc(3de - 8cf))x \sqrt{bx^2 + a}}{15c^2(bc - ad) f^2 (de - cf)^2 (dx^2 + c)^{3/2}} + \\
& \frac{C(2bc - ad)x \sqrt{bx^2 + a}}{15cd(bc - ad) f (dx^2 + c)^{3/2}} - \frac{(Ce - Bf)x \sqrt{bx^2 + a}}{5cf^2 (dx^2 + c)^{5/2}} + \frac{d(Ce^2 - Bfe + Af^2) x \sqrt{bx^2 + a}}{5cf^2 (de - cf) (dx^2 + c)^{5/2}} - \\
& \frac{Cx \sqrt{bx^2 + a}}{5df (dx^2 + c)^{5/2}}
\end{aligned}$$

input

```
Int[(Sqrt[a + b*x^2]*(A + B*x^2 + C*x^4))/((c + d*x^2)^(7/2)*(e + f*x^2)),
x]
```


output

```

-1/5*(C*x*Sqrt[a + b*x^2])/(d*f*(c + d*x^2)^(5/2)) - ((C*e - B*f)*x*Sqrt[a
+ b*x^2])/(5*c*f^2*(c + d*x^2)^(5/2)) + (d*(C*e^2 - B*e*f + A*f^2)*x*Sqrt
[a + b*x^2])/(5*c*f^2*(d*e - c*f)*(c + d*x^2)^(5/2)) + (C*(2*b*c - a*d)*x*
Sqrt[a + b*x^2])/(15*c*d*(b*c - a*d)*f*(c + d*x^2)^(3/2)) - ((3*b*c - 4*a*
d)*(C*e - B*f)*x*Sqrt[a + b*x^2])/(15*c^2*(b*c - a*d)*f^2*(c + d*x^2)^(3/2
)) - (d*(C*e^2 - B*e*f + A*f^2)*(a*d*(4*d*e - 9*c*f) - b*c*(3*d*e - 8*c*f)
)*x*Sqrt[a + b*x^2])/(15*c^2*(b*c - a*d)*f^2*(d*e - c*f)^2*(c + d*x^2)^(3/
2)) + (2*C*(b^2*c^2 - a*b*c*d + a^2*d^2)*Sqrt[a + b*x^2]*EllipticE[ArcTan[
(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(15*c^(3/2)*d^(3/2)*(b*c - a*d)^2*
f*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) - ((3*b^2*c^2 - 1
3*a*b*c*d + 8*a^2*d^2)*(C*e - B*f)*Sqrt[a + b*x^2]*EllipticE[ArcTan[(Sqrt[
d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(15*c^(5/2)*Sqrt[d]*(b*c - a*d)^2*f^2*Sq
rt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) + (Sqrt[d]*(C*e^2 - B
*e*f + A*f^2)*Sqrt[a + b*x^2]*EllipticE[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (
b*c)/(a*d)]/(Sqrt[c]*(d*e - c*f)^3*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*
Sqrt[c + d*x^2]) - (Sqrt[d]*(C*e^2 - B*e*f + A*f^2)*(a*b*c*d*(13*d*e - 28*
c*f) - 2*a^2*d^2*(4*d*e - 9*c*f) - b^2*c^2*(3*d*e - 8*c*f))*Sqrt[a + b*x^2
]*EllipticE[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(15*c^(5/2)*(b*
c - a*d)^2*f^2*(d*e - c*f)^2*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c
+ d*x^2]) - (b*C*(b*c + a*d)*Sqrt[a + b*x^2]*EllipticF[ArcTan[(Sqrt[d]*...

```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 7276

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 13343 vs. $2(1063) = 2126$.

Time = 11.50 (sec) , antiderivative size = 13344, normalized size of antiderivative = 12.21

method	result	size
elliptic	Expression too large to display	13344
default	Expression too large to display	18396

input `int((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)^(7/2)/(f*x^2+e),x,method=_RE
TURNVERBOSE)`

output `result too large to display`

Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{7/2}(e+fx^2)} dx = \text{Timed out}$$

input `integrate((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)^(7/2)/(f*x^2+e),x,alg
orithm="fricas")`

output `Timed out`

Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{7/2}(e+fx^2)} dx = \text{Timed out}$$

input `integrate((b*x**2+a)**(1/2)*(C*x**4+B*x**2+A)/(d*x**2+c)**(7/2)/(f*x**2+e)
,x)`

output `Timed out`

Maxima [F]

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{7/2}(e+fx^2)} dx = \int \frac{(Cx^4+Bx^2+A)\sqrt{bx^2+a}}{(dx^2+c)^{7/2}(fx^2+e)} dx$$

input `integrate((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)^(7/2)/(f*x^2+e),x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)*sqrt(b*x^2 + a)/((d*x^2 + c)^(7/2)*(f*x^2 + e)), x)`

Giac [F]

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{7/2}(e+fx^2)} dx = \int \frac{(Cx^4+Bx^2+A)\sqrt{bx^2+a}}{(dx^2+c)^{7/2}(fx^2+e)} dx$$

input `integrate((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)^(7/2)/(f*x^2+e),x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)*sqrt(b*x^2 + a)/((d*x^2 + c)^(7/2)*(f*x^2 + e)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{7/2}(e+fx^2)} dx = \int \frac{\sqrt{bx^2+a}(Cx^4+Bx^2+A)}{(dx^2+c)^{7/2}(fx^2+e)} dx$$

input `int(((a + b*x^2)^(1/2)*(A + B*x^2 + C*x^4))/((c + d*x^2)^(7/2)*(e + f*x^2)),x)`

output

```
int(((a + b*x^2)^(1/2)*(A + B*x^2 + C*x^4))/((c + d*x^2)^(7/2)*(e + f*x^2)
), x)
```

Reduce [F]

$$\int \frac{\sqrt{a+bx^2}(A+Bx^2+Cx^4)}{(c+dx^2)^{7/2}(e+fx^2)} dx = \text{too large to display}$$

input

```
int((b*x^2+a)^(1/2)*(C*x^4+B*x^2+A)/(d*x^2+c)^(7/2)/(f*x^2+e),x)
```

output

```
( - sqrt(c + d*x**2)*sqrt(a + b*x**2)*a*c*x - sqrt(c + d*x**2)*sqrt(a + b*
x**2)*b**2*x + 4*int((sqrt(c + d*x**2)*sqrt(a + b*x**2)*x**6)/(4*a**2*c**4
*d*e*f + 4*a**2*c**4*d*f**2*x**2 + 16*a**2*c**3*d**2*e*f*x**2 + 16*a**2*c*
*3*d**2*f**2*x**4 + 24*a**2*c**2*d**3*e*f*x**4 + 24*a**2*c**2*d**3*f**2*x*
*6 + 16*a**2*c*d**4*e*f*x**6 + 16*a**2*c*d**4*f**2*x**8 + 4*a**2*d**5*e*f*
x**8 + 4*a**2*d**5*f**2*x**10 - 2*a*b*c**5*e*f - 2*a*b*c**5*f**2*x**2 + 3*
a*b*c**4*d*e**2 - a*b*c**4*d*e*f*x**2 - 4*a*b*c**4*d*f**2*x**4 + 12*a*b*c*
*3*d**2*e**2*x**2 + 16*a*b*c**3*d**2*e*f*x**4 + 4*a*b*c**3*d**2*f**2*x**6
+ 18*a*b*c**2*d**3*e**2*x**4 + 34*a*b*c**2*d**3*e*f*x**6 + 16*a*b*c**2*d**
3*f**2*x**8 + 12*a*b*c*d**4*e**2*x**6 + 26*a*b*c*d**4*e*f*x**8 + 14*a*b*c*
d**4*f**2*x**10 + 3*a*b*d**5*e**2*x**8 + 7*a*b*d**5*e*f*x**10 + 4*a*b*d**5
*f**2*x**12 - 2*b**2*c**5*e*f*x**2 - 2*b**2*c**5*f**2*x**4 + 3*b**2*c**4*d
*e**2*x**2 - 5*b**2*c**4*d*e*f*x**4 - 8*b**2*c**4*d*f**2*x**6 + 12*b**2*c*
*3*d**2*e**2*x**4 - 12*b**2*c**3*d**2*f**2*x**8 + 18*b**2*c**2*d**3*e**2*x
**6 + 10*b**2*c**2*d**3*e*f*x**8 - 8*b**2*c**2*d**3*f**2*x**10 + 12*b**2*c
*d**4*e**2*x**8 + 10*b**2*c*d**4*e*f*x**10 - 2*b**2*c*d**4*f**2*x**12 + 3*
b**2*d**5*e**2*x**10 + 3*b**2*d**5*e*f*x**12),x)*a**2*b*c**4*d**2*f**2 + 1
2*int((sqrt(c + d*x**2)*sqrt(a + b*x**2)*x**6)/(4*a**2*c**4*d*e*f + 4*a**2
*c**4*d*f**2*x**2 + 16*a**2*c**3*d**2*e*f*x**2 + 16*a**2*c**3*d**2*f**2*x*
*4 + 24*a**2*c**2*d**3*e*f*x**4 + 24*a**2*c**2*d**3*f**2*x**6 + 16*a**2...
```

3.14
$$\int \frac{A+Bx^2}{(a+bx^2)\sqrt{c+dx^2}\sqrt{e+fx^2}} dx$$

Optimal result	148
Mathematica [C] (verified)	149
Rubi [A] (verified)	149
Maple [A] (verified)	150
Fricas [F(-1)]	151
Sympy [F]	151
Maxima [F]	152
Giac [F]	152
Mupad [F(-1)]	152
Reduce [F]	153

Optimal result

Integrand size = 39, antiderivative size = 226

$$\int \frac{A+Bx^2}{(a+bx^2)\sqrt{c+dx^2}\sqrt{e+fx^2}} dx$$

$$= \frac{\sqrt{c}(Be-Af)\sqrt{e+fx^2} \operatorname{EllipticF}\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1-\frac{cf}{de}\right)}{\sqrt{de}(be-af)\sqrt{c+dx^2}\sqrt{\frac{c(e+fx^2)}{e(c+dx^2)}}}$$

$$+ \frac{(Ab-aB)e^{3/2}\sqrt{c+dx^2} \operatorname{EllipticPi}\left(1-\frac{be}{af}, \arctan\left(\frac{\sqrt{fx}}{\sqrt{e}}\right), 1-\frac{de}{cf}\right)}{ac\sqrt{f}(be-af)\sqrt{\frac{e(c+dx^2)}{e+fx^2}}\sqrt{e+fx^2}}$$

output

```
c^(1/2)*(-A*f+B*e)*(f*x^2+e)^(1/2)*InverseJacobiAM(arctan(d^(1/2)*x/c^(1/2)), (1-c*f/d/e)^(1/2))/d^(1/2)/e/(-a*f+b*e)/(d*x^2+c)^(1/2)/(c*(f*x^2+e)/e/(d*x^2+c))^(1/2)+(A*b-B*a)*e^(3/2)*(d*x^2+c)^(1/2)*EllipticPi(f^(1/2)*x/e^(1/2)/(1+f*x^2/e)^(1/2), 1-b*e/a/f, (1-d*e/c/f)^(1/2))/a/c/f^(1/2)/(-a*f+b*e)/(e*(d*x^2+c)/c/(f*x^2+e))^(1/2)/(f*x^2+e)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 8.58 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.63

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e + fx^2}} dx = \frac{i\sqrt{1 + \frac{dx^2}{c}}\sqrt{1 + \frac{fx^2}{e}} \left(aB \operatorname{EllipticF} \left(\operatorname{arcsinh} \left(\sqrt{\frac{d}{c}}x \right), \frac{cf}{de} \right) + (Ab - aB) \operatorname{EllipticPi} \left(\frac{bc}{ad}, \operatorname{arcsinh} \left(\sqrt{\frac{d}{c}}x \right) \right) \right)}{ab\sqrt{\frac{d}{c}}\sqrt{c + dx^2}\sqrt{e + fx^2}}$$

input

```
Integrate[(A + B*x^2)/((a + b*x^2)*Sqrt[c + d*x^2]*Sqrt[e + f*x^2]),x]
```

output

```
((-I)*Sqrt[1 + (d*x^2)/c]*Sqrt[1 + (f*x^2)/e]*(a*B*EllipticF[I*ArcSinh[Sqrt[d/c]*x], (c*f)/(d*e)] + (A*b - a*B)*EllipticPi[(b*c)/(a*d), I*ArcSinh[Sqrt[d/c]*x], (c*f)/(d*e)))/(a*b*Sqrt[d/c]*Sqrt[c + d*x^2]*Sqrt[e + f*x^2])
```

Rubi [A] (verified)

Time = 0.99 (sec) , antiderivative size = 203, normalized size of antiderivative = 0.90, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.051$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e + fx^2}} dx$$

↓ 7276

$$\int \left(\frac{Ab - aB}{b(a + bx^2)\sqrt{c + dx^2}\sqrt{e + fx^2}} + \frac{B}{b\sqrt{c + dx^2}\sqrt{e + fx^2}} \right) dx$$

↓ 2009

$$\frac{\sqrt{-c}\sqrt{\frac{dx^2}{c} + 1}\sqrt{\frac{fx^2}{e} + 1}(Ab - aB) \operatorname{EllipticPi}\left(\frac{bc}{ad}, \arcsin\left(\frac{\sqrt{dx}}{\sqrt{-c}}\right), \frac{cf}{de}\right) + B\sqrt{e}\sqrt{c + dx^2} \operatorname{EllipticF}\left(\arctan\left(\frac{\sqrt{fx}}{\sqrt{e}}\right), 1 - \frac{de}{cf}\right)}{ab\sqrt{d}\sqrt{c + dx^2}\sqrt{e + fx^2} + bc\sqrt{f}\sqrt{e + fx^2}\sqrt{\frac{e(c+dx^2)}{c(e+fx^2)}}}$$

input `Int[(A + B*x^2)/((a + b*x^2)*Sqrt[c + d*x^2]*Sqrt[e + f*x^2]),x]`

output `(B*Sqrt[e]*Sqrt[c + d*x^2]*EllipticF[ArcTan[(Sqrt[f]*x)/Sqrt[e]], 1 - (d*e)/(c*f)]/(b*c*Sqrt[f]*Sqrt[(e*(c + d*x^2))/(c*(e + f*x^2))]*Sqrt[e + f*x^2]) + ((A*b - a*B)*Sqrt[-c]*Sqrt[1 + (d*x^2)/c]*Sqrt[1 + (f*x^2)/e]*EllipticPi[(b*c)/(a*d), ArcSin[(Sqrt[d]*x)/Sqrt[-c]], (c*f)/(d*e)]/(a*b*Sqrt[d]*Sqrt[c + d*x^2]*Sqrt[e + f*x^2]))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 5.78 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.85

method	result
default	$\frac{\left(A \operatorname{EllipticPi}\left(x\sqrt{-\frac{d}{c}}, \frac{bc}{ad}, \sqrt{\frac{-f}{e}}\right) b + B \operatorname{EllipticF}\left(x\sqrt{-\frac{d}{c}}, \sqrt{\frac{cf}{de}}\right) a - B \operatorname{EllipticPi}\left(x\sqrt{-\frac{d}{c}}, \frac{bc}{ad}, \sqrt{\frac{-f}{e}}\right) a\right) \sqrt{\frac{fx^2+e}{e}} \sqrt{\frac{x^2d+c}{c}} \sqrt{x^2d+c}}{ba\sqrt{-\frac{d}{c}}(dfx^4+cfx^2+dex^2+ce)}$
elliptic	$\frac{\sqrt{(x^2d+c)(fx^2+e)} \left(\frac{B\sqrt{1+\frac{dx^2}{c}}\sqrt{1+\frac{fx^2}{e}} \operatorname{EllipticF}\left(x\sqrt{-\frac{d}{c}}, \sqrt{-1+\frac{cf+de}{ed}}\right)}{b\sqrt{-\frac{d}{c}}\sqrt{dfx^4+cfx^2+dex^2+ce}} + \frac{\sqrt{1+\frac{dx^2}{c}}\sqrt{1+\frac{fx^2}{e}} \operatorname{EllipticPi}\left(x\sqrt{-\frac{d}{c}}, \frac{bc}{ad}, \sqrt{\frac{-f}{e}}\right) A}{a\sqrt{-\frac{d}{c}}\sqrt{dfx^4+cfx^2+dex^2+ce}} - \frac{A}{\sqrt{x^2d+c}\sqrt{fx^2+e}} \right)}{\sqrt{x^2d+c}\sqrt{fx^2+e}}$

input `int((B*x^2+A)/(b*x^2+a)/(d*x^2+c)^(1/2)/(f*x^2+e)^(1/2),x,method=_RETURNVE
RBOSE)`

output `(A*EllipticPi(x*(-1/c*d)^(1/2),b*c/a/d,(-f/e)^(1/2)/(-1/c*d)^(1/2))*b+B*El
lipticF(x*(-1/c*d)^(1/2),(c*f/d/e)^(1/2))*a-B*EllipticPi(x*(-1/c*d)^(1/2),
b*c/a/d,(-f/e)^(1/2)/(-1/c*d)^(1/2))*a)*((f*x^2+e)/e)^(1/2)*((d*x^2+c)/c)
^(1/2)/b*(d*x^2+c)^(1/2)*(f*x^2+e)^(1/2)/a/(-1/c*d)^(1/2)/(d*f*x^4+c*f*x^2+
d*e*x^2+c*e)`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e + fx^2}} dx = \text{Timed out}$$

input `integrate((B*x^2+A)/(b*x^2+a)/(d*x^2+c)^(1/2)/(f*x^2+e)^(1/2),x, algorithm
="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e + fx^2}} dx = \int \frac{A + Bx^2}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e + fx^2}} dx$$

input `integrate((B*x**2+A)/(b*x**2+a)/(d*x**2+c)**(1/2)/(f*x**2+e)**(1/2),x)`

output `Integral((A + B*x**2)/((a + b*x**2)*sqrt(c + d*x**2)*sqrt(e + f*x**2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e + fx^2}} dx = \int \frac{Bx^2 + A}{(bx^2 + a)\sqrt{dx^2 + c}\sqrt{fx^2 + e}} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)/(d*x^2+c)^(1/2)/(f*x^2+e)^(1/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/((b*x^2 + a)*sqrt(d*x^2 + c)*sqrt(f*x^2 + e)), x)`

Giac [F]

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e + fx^2}} dx = \int \frac{Bx^2 + A}{(bx^2 + a)\sqrt{dx^2 + c}\sqrt{fx^2 + e}} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)/(d*x^2+c)^(1/2)/(f*x^2+e)^(1/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/((b*x^2 + a)*sqrt(d*x^2 + c)*sqrt(f*x^2 + e)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e + fx^2}} dx = \int \frac{Bx^2 + A}{(bx^2 + a)\sqrt{dx^2 + c}\sqrt{fx^2 + e}} dx$$

input `int((A + B*x^2)/((a + b*x^2)*(c + d*x^2)^(1/2)*(e + f*x^2)^(1/2)),x)`

output `int((A + B*x^2)/((a + b*x^2)*(c + d*x^2)^(1/2)*(e + f*x^2)^(1/2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e + fx^2}} dx = \int \frac{\sqrt{fx^2 + e}\sqrt{dx^2 + c}}{dfx^4 + cfx^2 + dex^2 + ce} dx$$

input `int((B*x^2+A)/(b*x^2+a)/(d*x^2+c)^(1/2)/(f*x^2+e)^(1/2),x)`

output `int((sqrt(e + f*x**2)*sqrt(c + d*x**2))/(c*e + c*f*x**2 + d*e*x**2 + d*f*x**4),x)`

3.15
$$\int \frac{A+Bx^2}{(a+bx^2)\sqrt{c-dx^2}\sqrt{e+fx^2}} dx$$

Optimal result	154
Mathematica [C] (verified)	155
Rubi [A] (verified)	155
Maple [A] (verified)	156
Fricas [F(-1)]	157
Sympy [F]	157
Maxima [F]	158
Giac [F]	158
Mupad [F(-1)]	158
Reduce [F]	159

Optimal result

Integrand size = 40, antiderivative size = 203

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} dx$$

$$= \frac{B\sqrt{c}\sqrt{1 - \frac{dx^2}{c}}\sqrt{1 + \frac{fx^2}{e}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), -\frac{cf}{de}\right)}{b\sqrt{d}\sqrt{c - dx^2}\sqrt{e + fx^2}}$$

$$+ \frac{(Ab - aB)\sqrt{c}\sqrt{1 - \frac{dx^2}{c}}\sqrt{1 + \frac{fx^2}{e}} \operatorname{EllipticPi}\left(-\frac{bc}{ad}, \arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), -\frac{cf}{de}\right)}{ab\sqrt{d}\sqrt{c - dx^2}\sqrt{e + fx^2}}$$

output

```
B*c^(1/2)*(1-d*x^2/c)^(1/2)*(1+f*x^2/e)^(1/2)*EllipticF(d^(1/2)*x/c^(1/2),
(-c*f/d/e)^(1/2))/b/d^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(1/2)+(A*b-B*a)*c^(
1/2)*(1-d*x^2/c)^(1/2)*(1+f*x^2/e)^(1/2)*EllipticPi(d^(1/2)*x/c^(1/2),-b*c
/a/d,(-c*f/d/e)^(1/2))/a/b/d^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 8.55 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.74

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} dx = \frac{i\sqrt{1 - \frac{dx^2}{c}}\sqrt{1 + \frac{fx^2}{e}} \left(aB \operatorname{EllipticF} \left(\operatorname{arcsinh} \left(\sqrt{-\frac{d}{c}}x \right), -\frac{cf}{de} \right) + (Ab - aB) \operatorname{EllipticPi} \left(-\frac{bc}{ad}, \operatorname{arcsinh} \left(\sqrt{-\frac{d}{c}}x \right) \right) \right)}{ab\sqrt{-\frac{d}{c}}\sqrt{c - dx^2}\sqrt{e + fx^2}}$$

input

```
Integrate[(A + B*x^2)/((a + b*x^2)*Sqrt[c - d*x^2]*Sqrt[e + f*x^2]),x]
```

output

```
((-I)*Sqrt[1 - (d*x^2)/c]*Sqrt[1 + (f*x^2)/e]*(a*B*EllipticF[I*ArcSinh[Sqrt[-(d/c)]*x], -((c*f)/(d*e))] + (A*b - a*B)*EllipticPi[-((b*c)/(a*d)), I*ArcSinh[Sqrt[-(d/c)]*x], -((c*f)/(d*e)))]/(a*b*Sqrt[-(d/c)]*Sqrt[c - d*x^2]*Sqrt[e + f*x^2])
```

Rubi [A] (verified)

Time = 1.04 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} dx$$

↓ 7276

$$\int \left(\frac{Ab - aB}{b(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} + \frac{B}{b\sqrt{c - dx^2}\sqrt{e + fx^2}} \right) dx$$

↓ 2009

$$\frac{\sqrt{c}\sqrt{1-\frac{dx^2}{c}}\sqrt{\frac{fx^2}{e}+1}(Ab-aB)\text{EllipticPi}\left(-\frac{bc}{ad}, \arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), -\frac{cf}{de}\right)}{ab\sqrt{d}\sqrt{c-dx^2}\sqrt{e+fx^2}} + \frac{B\sqrt{c}\sqrt{1-\frac{dx^2}{c}}\sqrt{\frac{fx^2}{e}+1}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), -\frac{cf}{de}\right)}{b\sqrt{d}\sqrt{c-dx^2}\sqrt{e+fx^2}}$$

input `Int[(A + B*x^2)/((a + b*x^2)*Sqrt[c - d*x^2]*Sqrt[e + f*x^2]),x]`

output `(B*Sqrt[c]*Sqrt[1 - (d*x^2)/c]*Sqrt[1 + (f*x^2)/e]*EllipticF[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -(c*f)/(d*e))]/(b*Sqrt[d]*Sqrt[c - d*x^2]*Sqrt[e + f*x^2]) + ((A*b - a*B)*Sqrt[c]*Sqrt[1 - (d*x^2)/c]*Sqrt[1 + (f*x^2)/e]*EllipticPi[-((b*c)/(a*d)), ArcSin[(Sqrt[d]*x)/Sqrt[c]], -(c*f)/(d*e))]/(a*b*Sqrt[d]*Sqrt[c - d*x^2]*Sqrt[e + f*x^2])`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 6.02 (sec) , antiderivative size = 192, normalized size of antiderivative = 0.95

method	result
default	$\frac{\left(A\text{EllipticPi}\left(x\sqrt{\frac{d}{c}}, -\frac{bc}{ad}, \frac{\sqrt{-\frac{f}{e}}}{\sqrt{\frac{d}{c}}}\right)b + B\text{EllipticF}\left(x\sqrt{\frac{d}{c}}, \sqrt{-\frac{cf}{de}}\right)a - B\text{EllipticPi}\left(x\sqrt{\frac{d}{c}}, -\frac{bc}{ad}, \frac{\sqrt{-\frac{f}{e}}}{\sqrt{\frac{d}{c}}}\right)a\right)\sqrt{\frac{fx^2+e}{e}}\sqrt{\frac{-x^2d+c}{c}}\sqrt{-x^2d+c}}{ba\sqrt{\frac{d}{c}}(-dfx^4+cfx^2-dex^2+ce)}$
elliptic	$\frac{\sqrt{(-x^2d+c)(fx^2+e)}\left(\frac{B\sqrt{1-\frac{dx^2}{c}}\sqrt{1+\frac{fx^2}{e}}\text{EllipticF}\left(x\sqrt{\frac{d}{c}}, \sqrt{-1-\frac{cf-de}{ed}}\right)}{b\sqrt{\frac{d}{c}}\sqrt{-dfx^4+cfx^2-dex^2+ce}} + \frac{\sqrt{1-\frac{dx^2}{c}}\sqrt{1+\frac{fx^2}{e}}\text{EllipticPi}\left(x\sqrt{\frac{d}{c}}, -\frac{bc}{ad}, \frac{\sqrt{-\frac{f}{e}}}{\sqrt{\frac{d}{c}}}\right)A}{a\sqrt{\frac{d}{c}}\sqrt{-dfx^4+cfx^2-dex^2+ce}}\right)}{\sqrt{-x^2d+c}\sqrt{fx^2+e}}$

input `int((B*x^2+A)/(b*x^2+a)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(1/2),x,method=_RETURNV
ERBOSE)`

output `(A*EllipticPi(x*(1/c*d)^(1/2),-b*c/a/d,(-f/e)^(1/2)/(1/c*d)^(1/2))*b+B*EllipticF(x*(1/c*d)^(1/2),(-c*f/d/e)^(1/2))*a-B*EllipticPi(x*(1/c*d)^(1/2),-b*c/a/d,(-f/e)^(1/2)/(1/c*d)^(1/2))*a)/b*((f*x^2+e)/e)^(1/2)*((-d*x^2+c)/c)^(1/2)*(-d*x^2+c)^(1/2)*(f*x^2+e)^(1/2)/a/(1/c*d)^(1/2)/(-d*f*x^4+c*f*x^2-d*e*x^2+c*e)`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} dx = \text{Timed out}$$

input `integrate((B*x^2+A)/(b*x^2+a)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(1/2),x,algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} dx = \int \frac{A + Bx^2}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} dx$$

input `integrate((B*x**2+A)/(b*x**2+a)/(-d*x**2+c)**(1/2)/(f*x**2+e)**(1/2),x)`

output `Integral((A + B*x**2)/((a + b*x**2)*sqrt(c - d*x**2)*sqrt(e + f*x**2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} dx = \int \frac{Bx^2 + A}{(bx^2 + a)\sqrt{-dx^2 + c}\sqrt{fx^2 + e}} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(1/2),x, algorithm m="maxima")`

output `integrate((B*x^2 + A)/((b*x^2 + a)*sqrt(-d*x^2 + c)*sqrt(f*x^2 + e)), x)`

Giac [F]

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} dx = \int \frac{Bx^2 + A}{(bx^2 + a)\sqrt{-dx^2 + c}\sqrt{fx^2 + e}} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(1/2),x, algorithm m="giac")`

output `integrate((B*x^2 + A)/((b*x^2 + a)*sqrt(-d*x^2 + c)*sqrt(f*x^2 + e)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} dx = \int \frac{Bx^2 + A}{(bx^2 + a)\sqrt{c - dx^2}\sqrt{fx^2 + e}} dx$$

input `int((A + B*x^2)/((a + b*x^2)*(c - d*x^2)^(1/2)*(e + f*x^2)^(1/2)),x)`

output `int((A + B*x^2)/((a + b*x^2)*(c - d*x^2)^(1/2)*(e + f*x^2)^(1/2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} dx = \int \frac{\sqrt{fx^2 + e}\sqrt{-dx^2 + c}}{-dfx^4 + cfx^2 - dex^2 + ce} dx$$

input `int((B*x^2+A)/(b*x^2+a)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(1/2),x)`

output `int((sqrt(e + f*x**2)*sqrt(c - d*x**2))/(c*e + c*f*x**2 - d*e*x**2 - d*f*x**4),x)`

3.16
$$\int \frac{A+Bx^2}{(a+bx^2)\sqrt{c+dx^2}\sqrt{e-fx^2}} dx$$

Optimal result	160
Mathematica [C] (verified)	161
Rubi [A] (verified)	161
Maple [A] (verified)	162
Fricas [F(-1)]	163
Sympy [F]	163
Maxima [F]	164
Giac [F]	164
Mupad [F(-1)]	164
Reduce [F]	165

Optimal result

Integrand size = 40, antiderivative size = 203

$$\int \frac{A+Bx^2}{(a+bx^2)\sqrt{c+dx^2}\sqrt{e-fx^2}} dx$$

$$= \frac{B\sqrt{e}\sqrt{1+\frac{dx^2}{c}}\sqrt{1-\frac{fx^2}{e}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{fx}}{\sqrt{e}}\right), -\frac{de}{cf}\right)}{b\sqrt{f}\sqrt{c+dx^2}\sqrt{e-fx^2}}$$

$$+ \frac{(Ab-aB)\sqrt{e}\sqrt{1+\frac{dx^2}{c}}\sqrt{1-\frac{fx^2}{e}} \operatorname{EllipticPi}\left(-\frac{be}{af}, \arcsin\left(\frac{\sqrt{fx}}{\sqrt{e}}\right), -\frac{de}{cf}\right)}{ab\sqrt{f}\sqrt{c+dx^2}\sqrt{e-fx^2}}$$

output

```
B*e^(1/2)*(1+d*x^2/c)^(1/2)*(1-f*x^2/e)^(1/2)*EllipticF(f^(1/2)*x/e^(1/2),
(-d*e/c/f)^(1/2))/b/f^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2)+(A*b-B*a)*e^(
1/2)*(1+d*x^2/c)^(1/2)*(1-f*x^2/e)^(1/2)*EllipticPi(f^(1/2)*x/e^(1/2),-b*e
/a/f,(-d*e/c/f)^(1/2))/a/b/f^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 8.57 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.72

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e - fx^2}} dx = \frac{i\sqrt{1 + \frac{dx^2}{c}}\sqrt{1 - \frac{fx^2}{e}} \left(aB \operatorname{EllipticF} \left(\operatorname{arcsinh} \left(\sqrt{\frac{d}{c}} x \right), -\frac{cf}{de} \right) + (Ab - aB) \operatorname{EllipticPi} \left(\frac{bc}{ad}, \operatorname{arcsinh} \left(\sqrt{\frac{d}{c}} x \right) \right) \right)}{ab\sqrt{\frac{d}{c}}\sqrt{c + dx^2}\sqrt{e - fx^2}}$$

input `Integrate[(A + B*x^2)/((a + b*x^2)*Sqrt[c + d*x^2]*Sqrt[e - f*x^2]),x]`

output `((-I)*Sqrt[1 + (d*x^2)/c]*Sqrt[1 - (f*x^2)/e]*(a*B*EllipticF[I*ArcSinh[Sqrt[d/c]*x], -((c*f)/(d*e))] + (A*b - a*B)*EllipticPi[(b*c)/(a*d), I*ArcSinh[Sqrt[d/c]*x], -((c*f)/(d*e))])/(a*b*Sqrt[d/c]*Sqrt[c + d*x^2]*Sqrt[e - f*x^2])`

Rubi [A] (verified)

Time = 1.05 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e - fx^2}} dx$$

↓ 7276

$$\int \left(\frac{Ab - aB}{b(a + bx^2)\sqrt{c + dx^2}\sqrt{e - fx^2}} + \frac{B}{b\sqrt{c + dx^2}\sqrt{e - fx^2}} \right) dx$$

↓ 2009

$$\frac{\sqrt{e}\sqrt{\frac{dx^2}{c} + 1}\sqrt{1 - \frac{fx^2}{e}}(Ab - aB) \operatorname{EllipticPi}\left(-\frac{be}{af}, \arcsin\left(\frac{\sqrt{fx}}{\sqrt{e}}\right), -\frac{de}{cf}\right) + B\sqrt{e}\sqrt{\frac{dx^2}{c} + 1}\sqrt{1 - \frac{fx^2}{e}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{fx}}{\sqrt{e}}\right), -\frac{de}{cf}\right)}{ab\sqrt{f}\sqrt{c + dx^2}\sqrt{e - fx^2} + b\sqrt{f}\sqrt{c + dx^2}\sqrt{e - fx^2}}$$

input `Int[(A + B*x^2)/((a + b*x^2)*Sqrt[c + d*x^2]*Sqrt[e - f*x^2]),x]`

output `(B*Sqrt[e]*Sqrt[1 + (d*x^2)/c]*Sqrt[1 - (f*x^2)/e]*EllipticF[ArcSin[(Sqrt[f]*x)/Sqrt[e]], -(d*e)/(c*f))]/(b*Sqrt[f]*Sqrt[c + d*x^2]*Sqrt[e - f*x^2]) + ((A*b - a*B)*Sqrt[e]*Sqrt[1 + (d*x^2)/c]*Sqrt[1 - (f*x^2)/e]*EllipticPi[-(b*e)/(a*f), ArcSin[(Sqrt[f]*x)/Sqrt[e]], -(d*e)/(c*f)])/ (a*b*Sqrt[f]*Sqrt[c + d*x^2]*Sqrt[e - f*x^2])`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 5.91 (sec) , antiderivative size = 192, normalized size of antiderivative = 0.95

method	result
default	$\frac{\left(A \operatorname{EllipticPi}\left(x\sqrt{\frac{f}{e}}, -\frac{be}{af}, \sqrt{\frac{-d}{c}}\right) b + B \operatorname{EllipticF}\left(x\sqrt{\frac{f}{e}}, \sqrt{-\frac{de}{cf}}\right) a - B \operatorname{EllipticPi}\left(x\sqrt{\frac{f}{e}}, -\frac{be}{af}, \sqrt{\frac{-d}{c}}\right) a\right) \sqrt{\frac{x^2 d + c}{c}} \sqrt{\frac{-f x^2 + e}{e}} \sqrt{x^2 d + c}}{ba\sqrt{\frac{f}{e}}(-df x^4 - cf x^2 + de x^2 + ce)}$
elliptic	$\frac{\sqrt{(x^2 d + c)(-f x^2 + e)} \left(\frac{B \sqrt{1 - \frac{f x^2}{e}} \sqrt{1 + \frac{d x^2}{c}} \operatorname{EllipticF}\left(x\sqrt{\frac{f}{e}}, \sqrt{-1 - \frac{-cf + de}{cf}}\right)}{b\sqrt{\frac{f}{e}} \sqrt{-df x^4 - cf x^2 + de x^2 + ce}} + \frac{\sqrt{1 - \frac{f x^2}{e}} \sqrt{1 + \frac{d x^2}{c}} \operatorname{EllipticPi}\left(x\sqrt{\frac{f}{e}}, -\frac{be}{af}, \sqrt{\frac{-d}{c}}\right) A}{a\sqrt{\frac{f}{e}} \sqrt{-df x^4 - cf x^2 + de x^2 + ce}} \right)}{\sqrt{x^2 d + c} \sqrt{-f x^2 + e}}$

input `int((B*x^2+A)/(b*x^2+a)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2),x,method=_RETURNV
ERBOSE)`

output `(A*EllipticPi(x*(f/e)^(1/2),-b*e/a/f,(-1/c*d)^(1/2)/(f/e)^(1/2))*b+B*Ellip
ticF(x*(f/e)^(1/2),(-d*e/c/f)^(1/2))*a-B*EllipticPi(x*(f/e)^(1/2),-b*e/a/f
,(-1/c*d)^(1/2)/(f/e)^(1/2))*a)/b*((d*x^2+c)/c)^(1/2)*((-f*x^2+e)/e)^(1/2)
*(d*x^2+c)^(1/2)*(-f*x^2+e)^(1/2)/a/(f/e)^(1/2)/(-d*f*x^4-c*f*x^2+d*e*x^2+
c*e)`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e - fx^2}} dx = \text{Timed out}$$

input `integrate((B*x^2+A)/(b*x^2+a)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2),x,algorith
m="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e - fx^2}} dx = \int \frac{A + Bx^2}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e - fx^2}} dx$$

input `integrate((B*x**2+A)/(b*x**2+a)/(d*x**2+c)**(1/2)/(-f*x**2+e)**(1/2),x)`

output `Integral((A + B*x**2)/((a + b*x**2)*sqrt(c + d*x**2)*sqrt(e - f*x**2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e - fx^2}} dx = \int \frac{Bx^2 + A}{(bx^2 + a)\sqrt{dx^2 + c}\sqrt{-fx^2 + e}} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2),x, algorithm m="maxima")`

output `integrate((B*x^2 + A)/((b*x^2 + a)*sqrt(d*x^2 + c)*sqrt(-f*x^2 + e)), x)`

Giac [F]

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e - fx^2}} dx = \int \frac{Bx^2 + A}{(bx^2 + a)\sqrt{dx^2 + c}\sqrt{-fx^2 + e}} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2),x, algorithm m="giac")`

output `integrate((B*x^2 + A)/((b*x^2 + a)*sqrt(d*x^2 + c)*sqrt(-f*x^2 + e)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e - fx^2}} dx = \int \frac{Bx^2 + A}{(bx^2 + a)\sqrt{dx^2 + c}\sqrt{e - fx^2}} dx$$

input `int((A + B*x^2)/((a + b*x^2)*(c + d*x^2)^(1/2)*(e - f*x^2)^(1/2)),x)`

output `int((A + B*x^2)/((a + b*x^2)*(c + d*x^2)^(1/2)*(e - f*x^2)^(1/2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e - fx^2}} dx = \int \frac{\sqrt{-fx^2 + e}\sqrt{dx^2 + c}}{-dfx^4 - cfx^2 + dex^2 + ce} dx$$

input `int((B*x^2+A)/(b*x^2+a)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2),x)`

output `int((sqrt(e - f*x**2)*sqrt(c + d*x**2))/(c*e - c*f*x**2 + d*e*x**2 - d*f*x**4),x)`

3.17
$$\int \frac{A+Bx^2}{(a+bx^2)\sqrt{c-dx^2}\sqrt{e-fx^2}} dx$$

Optimal result	166
Mathematica [C] (verified)	167
Rubi [A] (verified)	167
Maple [A] (verified)	168
Fricas [F(-1)]	169
Sympy [F]	169
Maxima [F]	170
Giac [F]	170
Mupad [F(-1)]	170
Reduce [F]	171

Optimal result

Integrand size = 41, antiderivative size = 205

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx$$

$$= \frac{B\sqrt{c}\sqrt{1 - \frac{dx^2}{c}}\sqrt{1 - \frac{fx^2}{e}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), \frac{cf}{de}\right)}{b\sqrt{d}\sqrt{c - dx^2}\sqrt{e - fx^2}}$$

$$+ \frac{(Ab - aB)\sqrt{c}\sqrt{1 - \frac{dx^2}{c}}\sqrt{1 - \frac{fx^2}{e}} \operatorname{EllipticPi}\left(-\frac{bc}{ad}, \arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), \frac{cf}{de}\right)}{ab\sqrt{d}\sqrt{c - dx^2}\sqrt{e - fx^2}}$$

output

```
B*c^(1/2)*(1-d*x^2/c)^(1/2)*(1-f*x^2/e)^(1/2)*EllipticF(d^(1/2)*x/c^(1/2),
(c*f/d/e)^(1/2))/b/d^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2)+(A*b-B*a)*c^(
1/2)*(1-d*x^2/c)^(1/2)*(1-f*x^2/e)^(1/2)*EllipticPi(d^(1/2)*x/c^(1/2),-b*c
/a/d,(c*f/d/e)^(1/2))/a/b/d^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 8.52 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.74

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx = \frac{i\sqrt{1 - \frac{dx^2}{c}}\sqrt{1 - \frac{fx^2}{e}} \left(aB \operatorname{EllipticF} \left(\operatorname{arcsinh} \left(\sqrt{-\frac{d}{c}}x \right), \frac{cf}{de} \right) + (Ab - aB) \operatorname{EllipticPi} \left(-\frac{bc}{ad}, \operatorname{arcsinh} \left(\sqrt{-\frac{d}{c}}x \right) \right) \right)}{ab\sqrt{-\frac{d}{c}}\sqrt{c - dx^2}\sqrt{e - fx^2}}$$

input

```
Integrate[(A + B*x^2)/((a + b*x^2)*Sqrt[c - d*x^2]*Sqrt[e - f*x^2]),x]
```

output

```
((-I)*Sqrt[1 - (d*x^2)/c]*Sqrt[1 - (f*x^2)/e]*(a*B*EllipticF[I*ArcSinh[Sqrt[-(d/c)]*x], (c*f)/(d*e)] + (A*b - a*B)*EllipticPi[-((b*c)/(a*d)), I*ArcSinh[Sqrt[-(d/c)]*x], (c*f)/(d*e)]))/(a*b*Sqrt[-(d/c)]*Sqrt[c - d*x^2]*Sqrt[e - f*x^2])
```

Rubi [A] (verified)

Time = 1.11 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.049$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx$$

↓ 7276

$$\int \left(\frac{Ab - aB}{b(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} + \frac{B}{b\sqrt{c - dx^2}\sqrt{e - fx^2}} \right) dx$$

↓ 2009

$$\frac{\sqrt{c}\sqrt{1-\frac{dx^2}{c}}\sqrt{1-\frac{fx^2}{e}}(Ab-aB)\text{EllipticPi}\left(-\frac{bc}{ad},\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right),\frac{cf}{de}\right)}{ab\sqrt{d}\sqrt{c-dx^2}\sqrt{e-fx^2}} + \frac{B\sqrt{e}\sqrt{1-\frac{dx^2}{c}}\sqrt{1-\frac{fx^2}{e}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{fx}}{\sqrt{e}}\right),\frac{de}{cf}\right)}{b\sqrt{f}\sqrt{c-dx^2}\sqrt{e-fx^2}}$$

input

```
Int[(A + B*x^2)/((a + b*x^2)*Sqrt[c - d*x^2]*Sqrt[e - f*x^2]),x]
```

output

```
(B*Sqrt[e]*Sqrt[1 - (d*x^2)/c]*Sqrt[1 - (f*x^2)/e]*EllipticF[ArcSin[(Sqrt[f]*x)/Sqrt[e]], (d*e)/(c*f)])/(b*Sqrt[f]*Sqrt[c - d*x^2]*Sqrt[e - f*x^2]) + ((A*b - a*B)*Sqrt[c]*Sqrt[1 - (d*x^2)/c]*Sqrt[1 - (f*x^2)/e]*EllipticPi[-((b*c)/(a*d)), ArcSin[(Sqrt[d]*x)/Sqrt[c]], (c*f)/(d*e)])/(a*b*Sqrt[d]*Sqrt[c - d*x^2]*Sqrt[e - f*x^2])
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 7276

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]
```

Maple [A] (verified)

Time = 5.94 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.93

method	result
default	$\frac{\left(A\text{EllipticPi}\left(x\sqrt{\frac{f}{e}},-\frac{be}{af},\sqrt{\frac{d}{c}}\right)b+B\text{EllipticF}\left(x\sqrt{\frac{f}{e}},\sqrt{\frac{de}{cf}}\right)a-B\text{EllipticPi}\left(x\sqrt{\frac{f}{e}},-\frac{be}{af},\sqrt{\frac{d}{c}}\right)a\right)\sqrt{\frac{-x^2d+c}{c}}\sqrt{\frac{-fx^2+e}{e}}\sqrt{-x^2d+c}}{ba\sqrt{\frac{f}{e}}(dfx^4-cfx^2-dex^2+ce)}$
elliptic	$\frac{\sqrt{(-x^2d+c)(-fx^2+e)}}{\sqrt{-x^2d+c}\sqrt{-fx^2+e}}\left(\frac{B\sqrt{1-\frac{fx^2}{e}}\sqrt{1-\frac{dx^2}{c}}\text{EllipticF}\left(x\sqrt{\frac{f}{e}},\sqrt{-1-\frac{-cf-de}{cf}}\right)}{b\sqrt{\frac{f}{e}}\sqrt{dfx^4-cfx^2-dex^2+ce}}+\frac{\sqrt{1-\frac{fx^2}{e}}\sqrt{1-\frac{dx^2}{c}}\text{EllipticPi}\left(x\sqrt{\frac{f}{e}},-\frac{be}{af},\sqrt{\frac{d}{c}}\right)A}{a\sqrt{\frac{f}{e}}\sqrt{dfx^4-cfx^2-dex^2+ce}}\right)$

input `int((B*x^2+A)/(b*x^2+a)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2),x,method=_RETURN
VERBOSE)`

output `(A*EllipticPi(x*(f/e)^(1/2),-b*e/a/f,(1/c*d)^(1/2)/(f/e)^(1/2))*b+B*EllipticF(x*(f/e)^(1/2),(d*e/c/f)^(1/2))*a-B*EllipticPi(x*(f/e)^(1/2),-b*e/a/f,(1/c*d)^(1/2)/(f/e)^(1/2))*a)*((-d*x^2+c)/c)^(1/2)*((-f*x^2+e)/e)^(1/2)/b*(-d*x^2+c)^(1/2)*(-f*x^2+e)^(1/2)/a/(f/e)^(1/2)/(d*f*x^4-c*f*x^2-d*e*x^2+c*e)`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx = \text{Timed out}$$

input `integrate((B*x^2+A)/(b*x^2+a)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2),x, algorithm="fricas")`

output Timed out

Sympy [F]

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx = \int \frac{A + Bx^2}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx$$

input `integrate((B*x**2+A)/(b*x**2+a)/(-d*x**2+c)**(1/2)/(-f*x**2+e)**(1/2),x)`

output `Integral((A + B*x**2)/((a + b*x**2)*sqrt(c - d*x**2)*sqrt(e - f*x**2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx = \int \frac{Bx^2 + A}{(bx^2 + a)\sqrt{-dx^2 + c}\sqrt{-fx^2 + e}} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/((b*x^2 + a)*sqrt(-d*x^2 + c)*sqrt(-f*x^2 + e)), x)`

Giac [F]

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx = \int \frac{Bx^2 + A}{(bx^2 + a)\sqrt{-dx^2 + c}\sqrt{-fx^2 + e}} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/((b*x^2 + a)*sqrt(-d*x^2 + c)*sqrt(-f*x^2 + e)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx = \int \frac{Bx^2 + A}{(bx^2 + a)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx$$

input `int((A + B*x^2)/((a + b*x^2)*(c - d*x^2)^(1/2)*(e - f*x^2)^(1/2)),x)`

output `int((A + B*x^2)/((a + b*x^2)*(c - d*x^2)^(1/2)*(e - f*x^2)^(1/2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx = \int \frac{\sqrt{-fx^2 + e}\sqrt{-dx^2 + c}}{dfx^4 - cfx^2 - dex^2 + ce} dx$$

input `int((B*x^2+A)/(b*x^2+a)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2),x)`

output `int((sqrt(e - f*x**2)*sqrt(c - d*x**2))/(c*e - c*f*x**2 - d*e*x**2 + d*f*x**4),x)`

3.18 $\int \frac{A+Bx^2+Cx^4}{(a+bx^2)\sqrt{c+dx^2}\sqrt{e+fx^2}} dx$

Optimal result	172
Mathematica [C] (verified)	173
Rubi [A] (verified)	173
Maple [A] (verified)	175
Fricas [F(-1)]	175
Sympy [F]	176
Maxima [F]	176
Giac [F]	176
Mupad [F(-1)]	177
Reduce [F]	177

Optimal result

Integrand size = 44, antiderivative size = 371

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e + fx^2}} dx$$

$$= \frac{Cx\sqrt{e + fx^2}}{bf\sqrt{c + dx^2}} - \frac{\sqrt{c}C\sqrt{e + fx^2}E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \mid 1 - \frac{cf}{de}\right)}{b\sqrt{d}f\sqrt{c + dx^2}\sqrt{\frac{c(e+fx^2)}{e(c+dx^2)}}}$$

$$+ \frac{\sqrt{c}(bBc - acC - Abd)\sqrt{e + fx^2} \text{EllipticF}\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1 - \frac{cf}{de}\right)}{b\sqrt{d}(bc - ad)e\sqrt{c + dx^2}\sqrt{\frac{c(e+fx^2)}{e(c+dx^2)}}}$$

$$+ \frac{c^{3/2}(Ab^2 - a(bB - aC))\sqrt{e + fx^2} \text{EllipticPi}\left(1 - \frac{bc}{ad}, \arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1 - \frac{cf}{de}\right)}{ab\sqrt{d}(bc - ad)e\sqrt{c + dx^2}\sqrt{\frac{c(e+fx^2)}{e(c+dx^2)}}}$$

output

```
C*x*(f*x^2+e)^(1/2)/b/f/(d*x^2+c)^(1/2)-c^(1/2)*C*(f*x^2+e)^(1/2)*Elliptic
E(d^(1/2)*x/c^(1/2)/(1+d*x^2/c)^(1/2),(1-c*f/d/e)^(1/2))/b/d^(1/2)/f/(d*x^
2+c)^(1/2)/(c*(f*x^2+e)/e/(d*x^2+c)^(1/2)+c^(1/2)*(-A*b*d+B*b*c-C*a*c)*(f
*x^2+e)^(1/2)*InverseJacobiAM(arctan(d^(1/2)*x/c^(1/2)),(1-c*f/d/e)^(1/2))
/b/d^(1/2)/(-a*d+b*c)/e/(d*x^2+c)^(1/2)/(c*(f*x^2+e)/e/(d*x^2+c)^(1/2)+c^
(3/2)*(A*b^2-a*(B*b-C*a))*(f*x^2+e)^(1/2)*EllipticPi(d^(1/2)*x/c^(1/2)/(1+
d*x^2/c)^(1/2),1-b*c/a/d,(1-c*f/d/e)^(1/2))/a/b/d^(1/2)/(-a*d+b*c)/e/(d*x^
2+c)^(1/2)/(c*(f*x^2+e)/e/(d*x^2+c)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 7.31 (sec) , antiderivative size = 200, normalized size of antiderivative = 0.54

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e + fx^2}} dx = \frac{i\sqrt{1 + \frac{dx^2}{c}}\sqrt{1 + \frac{fx^2}{e}}\left(abCeE\left(\operatorname{arcsinh}\left(\sqrt{\frac{d}{c}}x\right)\middle|\frac{cf}{de}\right) - a(bCe - bBf + aCf)\operatorname{EllipticF}\left(\operatorname{arcsinh}\left(\sqrt{\frac{d}{c}}x\right)\right)\right)}{ab^2\sqrt{\frac{d}{c}}f\sqrt{c + dx^2}\sqrt{e + fx^2}}$$

input

```
Integrate[(A + B*x^2 + C*x^4)/((a + b*x^2)*Sqrt[c + d*x^2]*Sqrt[e + f*x^2]),x]
```

output

```
((-I)*Sqrt[1 + (d*x^2)/c]*Sqrt[1 + (f*x^2)/e]*(a*b*C*e*EllipticE[I*ArcSinh[Sqrt[d/c]*x], (c*f)/(d*e)] - a*(b*C*e - b*B*f + a*C*f)*EllipticF[I*ArcSinh[Sqrt[d/c]*x], (c*f)/(d*e)] + (A*b^2 + a*(-(b*B) + a*C))*f*EllipticPi[(b*c)/(a*d), I*ArcSinh[Sqrt[d/c]*x], (c*f)/(d*e)))/(a*b^2*Sqrt[d/c]*f*Sqrt[c + d*x^2]*Sqrt[e + f*x^2])
```

Rubi [A] (verified)

Time = 1.10 (sec) , antiderivative size = 342, normalized size of antiderivative = 0.92, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.045$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e + fx^2}} dx$$

↓ 7276

$$\int \left(\frac{a^2C - abB + Ab^2}{b^2(a + bx^2)\sqrt{c + dx^2}\sqrt{e + fx^2}} + \frac{bB - aC}{b^2\sqrt{c + dx^2}\sqrt{e + fx^2}} + \frac{Cx^2}{b\sqrt{c + dx^2}\sqrt{e + fx^2}} \right) dx$$

↓ 2009

$$\frac{\sqrt{-c}\sqrt{\frac{dx^2}{c}+1}\sqrt{\frac{fx^2}{e}+1}(Ab^2-a(bB-aC))\text{EllipticPi}\left(\frac{bc}{ad}, \arcsin\left(\frac{\sqrt{dx}}{\sqrt{-c}}\right), \frac{cf}{de}\right)}{ab^2\sqrt{d}\sqrt{c+dx^2}\sqrt{e+fx^2}} + \frac{\sqrt{e}\sqrt{c+dx^2}(bB-aC)\text{EllipticF}\left(\arctan\left(\frac{\sqrt{fx}}{\sqrt{e}}\right), 1-\frac{de}{cf}\right)}{b^2c\sqrt{f}\sqrt{e+fx^2}\sqrt{\frac{e(c+dx^2)}{c(e+fx^2)}}} - \frac{C\sqrt{e}\sqrt{c+dx^2}E\left(\arctan\left(\frac{\sqrt{fx}}{\sqrt{e}}\right) \middle| 1-\frac{de}{cf}\right)}{bd\sqrt{f}\sqrt{e+fx^2}\sqrt{\frac{e(c+dx^2)}{c(e+fx^2)}}} + \frac{Cx\sqrt{c+dx^2}}{bd\sqrt{e+fx^2}}$$

input `Int[(A + B*x^2 + C*x^4)/((a + b*x^2)*Sqrt[c + d*x^2]*Sqrt[e + f*x^2]),x]`

output `(C*x*Sqrt[c + d*x^2])/(b*d*Sqrt[e + f*x^2]) - (C*Sqrt[e]*Sqrt[c + d*x^2]*EllipticE[ArcTan[(Sqrt[f]*x)/Sqrt[e]], 1 - (d*e)/(c*f)])/(b*d*Sqrt[f]*Sqrt[(e*(c + d*x^2))/(c*(e + f*x^2))]*Sqrt[e + f*x^2]) + ((b*B - a*C)*Sqrt[e]*Sqrt[c + d*x^2]*EllipticF[ArcTan[(Sqrt[f]*x)/Sqrt[e]], 1 - (d*e)/(c*f)])/(b^2*c*Sqrt[f]*Sqrt[(e*(c + d*x^2))/(c*(e + f*x^2))]*Sqrt[e + f*x^2]) + (Sqrt[-c]*(A*b^2 - a*(b*B - a*C))*Sqrt[1 + (d*x^2)/c]*Sqrt[1 + (f*x^2)/e]*EllipticPi[(b*c)/(a*d), ArcSin[(Sqrt[d]*x)/Sqrt[-c]], (c*f)/(d*e)]/(a*b^2*Sqrt[d]*Sqrt[c + d*x^2]*Sqrt[e + f*x^2])`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 5.92 (sec) , antiderivative size = 328, normalized size of antiderivative = 0.88

method	result
default	$\left(A \operatorname{EllipticPi}\left(x\sqrt{-\frac{d}{c}}, \frac{bc}{ad}, \sqrt{\frac{-f}{e}}\right) b^2 f + B \operatorname{EllipticF}\left(x\sqrt{-\frac{d}{c}}, \sqrt{\frac{cf}{de}}\right) abf - B \operatorname{EllipticPi}\left(x\sqrt{-\frac{d}{c}}, \frac{bc}{ad}, \sqrt{\frac{-f}{e}}\right) abf - C \operatorname{EllipticF}\left(x\sqrt{-\frac{d}{c}}, \sqrt{\frac{cf}{de}}\right) abf \right) \sqrt{(x^2 d + c)(f x^2 + e)}$
elliptic	$\frac{B \sqrt{1 + \frac{d x^2}{c}} \sqrt{1 + \frac{f x^2}{e}} \operatorname{EllipticF}\left(x\sqrt{-\frac{d}{c}}, \sqrt{-1 + \frac{cf + de}{ed}}\right) - \sqrt{1 + \frac{d x^2}{c}} \sqrt{1 + \frac{f x^2}{e}} \operatorname{EllipticF}\left(x\sqrt{-\frac{d}{c}}, \sqrt{-1 + \frac{cf + de}{ed}}\right) C a}{b \sqrt{-\frac{d}{c}} \sqrt{d f x^4 + c f x^2 + d e x^2 + c e}}$

input `int((C*x^4+B*x^2+A)/(b*x^2+a)/(d*x^2+c)^(1/2)/(f*x^2+e)^(1/2),x,method=_RETURNVERBOSE)`

output `(A*EllipticPi(x*(-1/c*d)^(1/2),b*c/a/d,(-f/e)^(1/2)/(-1/c*d)^(1/2))*b^2*f+B*EllipticF(x*(-1/c*d)^(1/2),(c*f/d/e)^(1/2))*a*b*f-B*EllipticPi(x*(-1/c*d)^(1/2),b*c/a/d,(-f/e)^(1/2)/(-1/c*d)^(1/2))*a*b*f-C*EllipticF(x*(-1/c*d)^(1/2),(c*f/d/e)^(1/2))*a^2*f-C*EllipticF(x*(-1/c*d)^(1/2),(c*f/d/e)^(1/2))*a*b*e+C*EllipticE(x*(-1/c*d)^(1/2),(c*f/d/e)^(1/2))*a*b*e+C*EllipticPi(x*(-1/c*d)^(1/2),b*c/a/d,(-f/e)^(1/2)/(-1/c*d)^(1/2))*a^2*f*((f*x^2+e)/e)^(1/2)*((d*x^2+c)/c)^(1/2)*(d*x^2+c)^(1/2)*(f*x^2+e)^(1/2)/f/a/(-1/c*d)^(1/2)/b^2/(d*f*x^4+c*f*x^2+d*e*x^2+c*e)`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2) \sqrt{c + dx^2} \sqrt{e + fx^2}} dx = \text{Timed out}$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)/(d*x^2+c)^(1/2)/(f*x^2+e)^(1/2),x,algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e + fx^2}} dx = \int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e + fx^2}} dx$$

input `integrate((C*x**4+B*x**2+A)/(b*x**2+a)/(d*x**2+c)**(1/2)/(f*x**2+e)**(1/2),x)`

output `Integral((A + B*x**2 + C*x**4)/((a + b*x**2)*sqrt(c + d*x**2)*sqrt(e + f*x**2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e + fx^2}} dx = \int \frac{Cx^4 + Bx^2 + A}{(bx^2 + a)\sqrt{dx^2 + c}\sqrt{fx^2 + e}} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)/(d*x^2+c)^(1/2)/(f*x^2+e)^(1/2),x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)/((b*x^2 + a)*sqrt(d*x^2 + c)*sqrt(f*x^2 + e)), x)`

Giac [F]

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e + fx^2}} dx = \int \frac{Cx^4 + Bx^2 + A}{(bx^2 + a)\sqrt{dx^2 + c}\sqrt{fx^2 + e}} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)/(d*x^2+c)^(1/2)/(f*x^2+e)^(1/2),x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)/((b*x^2 + a)*sqrt(d*x^2 + c)*sqrt(f*x^2 + e)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e + fx^2}} dx = \int \frac{Cx^4 + Bx^2 + A}{(bx^2 + a)\sqrt{dx^2 + c}\sqrt{fx^2 + e}} dx$$

input

```
int((A + B*x^2 + C*x^4)/((a + b*x^2)*(c + d*x^2)^(1/2)*(e + f*x^2)^(1/2)),
x)
```

output

```
int((A + B*x^2 + C*x^4)/((a + b*x^2)*(c + d*x^2)^(1/2)*(e + f*x^2)^(1/2)),
x)
```

Reduce [F]

$$\begin{aligned} & \int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e + fx^2}} dx \\ &= \left(\int \frac{\sqrt{fx^2 + e}\sqrt{dx^2 + c}x^4}{bdfx^6 + adfx^4 + bcfx^4 + bde x^4 + acf x^2 + ade x^2 + bce x^2 + ace} dx \right) c \\ &+ \left(\int \frac{\sqrt{fx^2 + e}\sqrt{dx^2 + c}x^2}{bdfx^6 + adfx^4 + bcfx^4 + bde x^4 + acf x^2 + ade x^2 + bce x^2 + ace} dx \right) b \\ &+ \left(\int \frac{\sqrt{fx^2 + e}\sqrt{dx^2 + c}}{bdfx^6 + adfx^4 + bcfx^4 + bde x^4 + acf x^2 + ade x^2 + bce x^2 + ace} dx \right) a \end{aligned}$$

input

```
int((C*x^4+B*x^2+A)/(b*x^2+a)/(d*x^2+c)^(1/2)/(f*x^2+e)^(1/2),x)
```

output

```
int((sqrt(e + f*x**2)*sqrt(c + d*x**2)*x**4)/(a*c*e + a*c*f*x**2 + a*d*e*x
**2 + a*d*f*x**4 + b*c*e*x**2 + b*c*f*x**4 + b*d*e*x**4 + b*d*f*x**6),x)*c
+ int((sqrt(e + f*x**2)*sqrt(c + d*x**2)*x**2)/(a*c*e + a*c*f*x**2 + a*d*
e*x**2 + a*d*f*x**4 + b*c*e*x**2 + b*c*f*x**4 + b*d*e*x**4 + b*d*f*x**6),x
)*b + int((sqrt(e + f*x**2)*sqrt(c + d*x**2))/(a*c*e + a*c*f*x**2 + a*d*e*
x**2 + a*d*f*x**4 + b*c*e*x**2 + b*c*f*x**4 + b*d*e*x**4 + b*d*f*x**6),x)*
a
```

3.19 $\int \frac{A+Bx^2+Cx^4}{(a+bx^2)\sqrt{c-dx^2}\sqrt{e+fx^2}} dx$

Optimal result	178
Mathematica [C] (verified)	179
Rubi [A] (verified)	179
Maple [A] (verified)	181
Fricas [F(-1)]	181
Sympy [F]	182
Maxima [F]	182
Giac [F]	182
Mupad [F(-1)]	183
Reduce [F]	183

Optimal result

Integrand size = 45, antiderivative size = 323

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} dx$$

$$= \frac{\sqrt{c}C\sqrt{1 - \frac{dx^2}{c}}\sqrt{e + fx^2}E\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \mid -\frac{cf}{de}\right)}{b\sqrt{d}f\sqrt{c - dx^2}\sqrt{1 + \frac{fx^2}{e}}}$$

$$- \frac{\sqrt{c}(bCe - bBf + aCf)\sqrt{1 - \frac{dx^2}{c}}\sqrt{1 + \frac{fx^2}{e}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), -\frac{cf}{de}\right)}{b^2\sqrt{d}f\sqrt{c - dx^2}\sqrt{e + fx^2}}$$

$$+ \frac{\sqrt{c}(Ab^2 - a(bB - aC))\sqrt{1 - \frac{dx^2}{c}}\sqrt{1 + \frac{fx^2}{e}}\text{EllipticPi}\left(-\frac{bc}{ad}, \arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), -\frac{cf}{de}\right)}{ab^2\sqrt{d}\sqrt{c - dx^2}\sqrt{e + fx^2}}$$

output

```
c^(1/2)*C*(1-d*x^2/c)^(1/2)*(f*x^2+e)^(1/2)*EllipticE(d^(1/2)*x/c^(1/2),(-
c*f/d/e)^(1/2))/b/d^(1/2)/f/(-d*x^2+c)^(1/2)/(1+f*x^2/e)^(1/2)-c^(1/2)*(-B
*b*f+C*a*f+C*b*e)*(1-d*x^2/c)^(1/2)*(1+f*x^2/e)^(1/2)*EllipticF(d^(1/2)*x/
c^(1/2),(-c*f/d/e)^(1/2))/b^2/d^(1/2)/f/(-d*x^2+c)^(1/2)/(f*x^2+e)^(1/2)+c
^(1/2)*(A*b^2-a*(B*b-C*a))*(1-d*x^2/c)^(1/2)*(1+f*x^2/e)^(1/2)*EllipticPi(
d^(1/2)*x/c^(1/2),-b*c/a/d,(-c*f/d/e)^(1/2))/a/b^2/d^(1/2)/(-d*x^2+c)^(1/2
)/(f*x^2+e)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 7.29 (sec) , antiderivative size = 210, normalized size of antiderivative = 0.65

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} dx = \frac{i\sqrt{1 - \frac{dx^2}{c}}\sqrt{1 + \frac{fx^2}{e}}\left(abCeE\left(\operatorname{arcsinh}\left(\sqrt{-\frac{d}{c}}x\right)\middle| -\frac{cf}{de}\right) - a(bCe - bBf + aCf)\operatorname{EllipticF}\left(\operatorname{arcsinh}\left(\sqrt{-\frac{d}{c}}x\right)\right)\right)}{ab^2\sqrt{-\frac{d}{c}}f\sqrt{c - dx^2}\sqrt{e + fx^2}}$$

input

```
Integrate[(A + B*x^2 + C*x^4)/((a + b*x^2)*Sqrt[c - d*x^2]*Sqrt[e + f*x^2]),x]
```

output

```
((-I)*Sqrt[1 - (d*x^2)/c]*Sqrt[1 + (f*x^2)/e]*(a*b*C*e*EllipticE[I*ArcSinh[Sqrt[-(d/c)]*x], -((c*f)/(d*e))] - a*(b*C*e - b*B*f + a*C*f)*EllipticF[I*ArcSinh[Sqrt[-(d/c)]*x], -((c*f)/(d*e))] + (A*b^2 + a*(-(b*B) + a*C))*f*EllipticPi[-((b*c)/(a*d)), I*ArcSinh[Sqrt[-(d/c)]*x], -((c*f)/(d*e))]))/(a*b^2*Sqrt[-(d/c)]*f*Sqrt[c - d*x^2]*Sqrt[e + f*x^2])
```

Rubi [A] (verified)

Time = 1.25 (sec) , antiderivative size = 409, normalized size of antiderivative = 1.27, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.044$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} dx$$

↓ 7276

$$\int \left(\frac{a^2C - abB + Ab^2}{b^2(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} + \frac{bB - aC}{b^2\sqrt{c - dx^2}\sqrt{e + fx^2}} + \frac{Cx^2}{b\sqrt{c - dx^2}\sqrt{e + fx^2}} \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{\sqrt{c}\sqrt{1-\frac{dx^2}{c}}\sqrt{\frac{fx^2}{e}+1}(Ab^2-a(bB-aC))\text{EllipticPi}\left(-\frac{bc}{ad}, \arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), -\frac{cf}{de}\right)}{ab^2\sqrt{d}\sqrt{c-dx^2}\sqrt{e+fx^2}} + \\
& \frac{\sqrt{c}\sqrt{1-\frac{dx^2}{c}}\sqrt{\frac{fx^2}{e}+1}(bB-aC)\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), -\frac{cf}{de}\right)}{b^2\sqrt{d}\sqrt{c-dx^2}\sqrt{e+fx^2}} - \\
& \frac{\sqrt{c}Ce\sqrt{1-\frac{dx^2}{c}}\sqrt{\frac{fx^2}{e}+1}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), -\frac{cf}{de}\right)}{b\sqrt{d}f\sqrt{c-dx^2}\sqrt{e+fx^2}} + \\
& \frac{\sqrt{c}C\sqrt{1-\frac{dx^2}{c}}\sqrt{e+fx^2}E\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \middle| -\frac{cf}{de}\right)}{b\sqrt{d}f\sqrt{c-dx^2}\sqrt{\frac{fx^2}{e}+1}}
\end{aligned}$$

input `Int[(A + B*x^2 + C*x^4)/((a + b*x^2)*Sqrt[c - d*x^2]*Sqrt[e + f*x^2]),x]`

output `(Sqrt[c]*C*Sqrt[1 - (d*x^2)/c]*Sqrt[e + f*x^2]*EllipticE[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -((c*f)/(d*e))]/(b*Sqrt[d]*f*Sqrt[c - d*x^2]*Sqrt[1 + (f*x^2)/e]) + (Sqrt[c]*(b*B - a*C)*Sqrt[1 - (d*x^2)/c]*Sqrt[1 + (f*x^2)/e]*EllipticF[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -((c*f)/(d*e))]/(b^2*Sqrt[d]*Sqrt[c - d*x^2]*Sqrt[e + f*x^2]) - (Sqrt[c]*C*e*Sqrt[1 - (d*x^2)/c]*Sqrt[1 + (f*x^2)/e]*EllipticF[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -((c*f)/(d*e))]/(b*Sqrt[d]*f*Sqrt[c - d*x^2]*Sqrt[e + f*x^2]) + (Sqrt[c]*(A*b^2 - a*(b*B - a*C))*Sqrt[1 - (d*x^2)/c]*Sqrt[1 + (f*x^2)/e]*EllipticPi[-((b*c)/(a*d)), ArcSin[(Sqrt[d]*x)/Sqrt[c]], -((c*f)/(d*e))]/(a*b^2*Sqrt[d]*Sqrt[c - d*x^2]*Sqrt[e + f*x^2]))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 5.76 (sec) , antiderivative size = 328, normalized size of antiderivative = 1.02

method	result
default	$\left(A \operatorname{EllipticPi}\left(x\sqrt{\frac{d}{c}}, -\frac{bc}{ad}, \sqrt{\frac{-f}{e}}\right) b^2 f + B \operatorname{EllipticF}\left(x\sqrt{\frac{d}{c}}, \sqrt{-\frac{cf}{de}}\right) abf - B \operatorname{EllipticPi}\left(x\sqrt{\frac{d}{c}}, -\frac{bc}{ad}, \sqrt{\frac{-f}{e}}\right) abf - C \operatorname{EllipticF}\left(x\sqrt{\frac{d}{c}}, \sqrt{-\frac{cf}{de}}\right) abf \right) f a$
elliptic	$\sqrt{(-x^2 d + c)(f x^2 + e)} \left(\frac{B \sqrt{1 - \frac{d x^2}{c}} \sqrt{1 + \frac{f x^2}{e}} \operatorname{EllipticF}\left(x\sqrt{\frac{d}{c}}, \sqrt{-1 - \frac{cf - de}{ed}}\right)}{b \sqrt{\frac{d}{c}} \sqrt{-df x^4 + cf x^2 - de x^2 + ce}} - \frac{\sqrt{1 - \frac{d x^2}{c}} \sqrt{1 + \frac{f x^2}{e}} \operatorname{EllipticF}\left(x\sqrt{\frac{d}{c}}, \sqrt{-1 - \frac{cf - de}{ed}}\right) C a}{b^2 \sqrt{\frac{d}{c}} \sqrt{-df x^4 + cf x^2 - de x^2 + ce}} \right) C a$

input `int((C*x^4+B*x^2+A)/(b*x^2+a)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(1/2),x,method=_RETURVERBOSE)`

output `(A*EllipticPi(x*(1/c*d)^(1/2),-b*c/a/d,(-f/e)^(1/2)/(1/c*d)^(1/2))*b^2*f+B*EllipticF(x*(1/c*d)^(1/2),(-c*f/d/e)^(1/2))*a*b*f-B*EllipticPi(x*(1/c*d)^(1/2),-b*c/a/d,(-f/e)^(1/2)/(1/c*d)^(1/2))*a*b*f-C*EllipticF(x*(1/c*d)^(1/2),(-c*f/d/e)^(1/2))*a^2*f-C*EllipticF(x*(1/c*d)^(1/2),(-c*f/d/e)^(1/2))*a*b*e+C*EllipticE(x*(1/c*d)^(1/2),(-c*f/d/e)^(1/2))*a*b*e+C*EllipticPi(x*(1/c*d)^(1/2),-b*c/a/d,(-f/e)^(1/2)/(1/c*d)^(1/2))*a^2*f)*((f*x^2+e)/e)^(1/2))*((-d*x^2+c)/c)^(1/2)*(-d*x^2+c)^(1/2)*(f*x^2+e)^(1/2)/f/a/(1/c*d)^(1/2)/b^2/(-d*f*x^4+c*f*x^2-d*e*x^2+c*e)`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2) \sqrt{c - dx^2} \sqrt{e + fx^2}} dx = \text{Timed out}$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(1/2),x,algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} dx = \int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} dx$$

input `integrate((C*x**4+B*x**2+A)/(b*x**2+a)/(-d*x**2+c)**(1/2)/(f*x**2+e)**(1/2),x)`

output `Integral((A + B*x**2 + C*x**4)/((a + b*x**2)*sqrt(c - d*x**2)*sqrt(e + f*x**2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} dx = \int \frac{Cx^4 + Bx^2 + A}{(bx^2 + a)\sqrt{-dx^2 + c}\sqrt{fx^2 + e}} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(1/2),x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)/((b*x^2 + a)*sqrt(-d*x^2 + c)*sqrt(f*x^2 + e)), x)`

Giac [F]

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} dx = \int \frac{Cx^4 + Bx^2 + A}{(bx^2 + a)\sqrt{-dx^2 + c}\sqrt{fx^2 + e}} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(1/2),x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)/((b*x^2 + a)*sqrt(-d*x^2 + c)*sqrt(f*x^2 + e)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} dx = \int \frac{Cx^4 + Bx^2 + A}{(bx^2 + a)\sqrt{c - dx^2}\sqrt{fx^2 + e}} dx$$

input

```
int((A + B*x^2 + C*x^4)/((a + b*x^2)*(c - d*x^2)^(1/2)*(e + f*x^2)^(1/2)),
x)
```

output

```
int((A + B*x^2 + C*x^4)/((a + b*x^2)*(c - d*x^2)^(1/2)*(e + f*x^2)^(1/2)),
x)
```

Reduce [F]

$$\begin{aligned} & \int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e + fx^2}} dx \\ &= \left(\int \frac{\sqrt{fx^2 + e}\sqrt{-dx^2 + cx^4}}{-bdfx^6 - adfx^4 + bcfx^4 - bdex^4 + acfx^2 - adex^2 + bce x^2 + ace} dx \right) c \\ &+ \left(\int \frac{\sqrt{fx^2 + e}\sqrt{-dx^2 + cx^2}}{-bdfx^6 - adfx^4 + bcfx^4 - bdex^4 + acfx^2 - adex^2 + bce x^2 + ace} dx \right) b \\ &+ \left(\int \frac{\sqrt{fx^2 + e}\sqrt{-dx^2 + c}}{-bdfx^6 - adfx^4 + bcfx^4 - bdex^4 + acfx^2 - adex^2 + bce x^2 + ace} dx \right) a \end{aligned}$$

input

```
int((C*x^4+B*x^2+A)/(b*x^2+a)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(1/2),x)
```

output

```
int((sqrt(e + f*x**2)*sqrt(c - d*x**2)*x**4)/(a*c*e + a*c*f*x**2 - a*d*e*x
**2 - a*d*f*x**4 + b*c*e*x**2 + b*c*f*x**4 - b*d*e*x**4 - b*d*f*x**6),x)*c
+ int((sqrt(e + f*x**2)*sqrt(c - d*x**2)*x**2)/(a*c*e + a*c*f*x**2 - a*d*
e*x**2 - a*d*f*x**4 + b*c*e*x**2 + b*c*f*x**4 - b*d*e*x**4 - b*d*f*x**6),x
)*b + int((sqrt(e + f*x**2)*sqrt(c - d*x**2))/(a*c*e + a*c*f*x**2 - a*d*e*
x**2 - a*d*f*x**4 + b*c*e*x**2 + b*c*f*x**4 - b*d*e*x**4 - b*d*f*x**6),x)*
a
```


3.20 $\int \frac{A+Bx^2+Cx^4}{(a+bx^2)\sqrt{c+dx^2}\sqrt{e-fx^2}} dx$

Optimal result	184
Mathematica [C] (verified)	185
Rubi [A] (verified)	185
Maple [A] (verified)	187
Fricas [F(-1)]	187
Sympy [F]	188
Maxima [F]	188
Giac [F]	188
Mupad [F(-1)]	189
Reduce [F]	189

Optimal result

Integrand size = 45, antiderivative size = 323

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e - fx^2}} dx$$

$$= \frac{C\sqrt{e}\sqrt{c + dx^2}\sqrt{1 - \frac{fx^2}{e}} E\left(\arcsin\left(\frac{\sqrt{fx}}{\sqrt{e}}\right) \mid -\frac{de}{cf}\right)}{bd\sqrt{f}\sqrt{1 + \frac{dx^2}{c}}\sqrt{e - fx^2}}$$

$$- \frac{(bcC - bBd + aCd)\sqrt{e}\sqrt{1 + \frac{dx^2}{c}}\sqrt{1 - \frac{fx^2}{e}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{fx}}{\sqrt{e}}\right), -\frac{de}{cf}\right)}{b^2d\sqrt{f}\sqrt{c + dx^2}\sqrt{e - fx^2}}$$

$$+ \frac{(Ab^2 - a(bB - aC))\sqrt{e}\sqrt{1 + \frac{dx^2}{c}}\sqrt{1 - \frac{fx^2}{e}} \text{EllipticPi}\left(-\frac{be}{af}, \arcsin\left(\frac{\sqrt{fx}}{\sqrt{e}}\right), -\frac{de}{cf}\right)}{ab^2\sqrt{f}\sqrt{c + dx^2}\sqrt{e - fx^2}}$$

output

```
C*e^(1/2)*(d*x^2+c)^(1/2)*(1-f*x^2/e)^(1/2)*EllipticE(f^(1/2)*x/e^(1/2),(-
d*e/c/f)^(1/2))/b/d/f^(1/2)/(1+d*x^2/c)^(1/2)/(-f*x^2+e)^(1/2)-(-B*b*d+C*a
*d+C*b*c)*e^(1/2)*(1+d*x^2/c)^(1/2)*(1-f*x^2/e)^(1/2)*EllipticF(f^(1/2)*x/
e^(1/2),(-d*e/c/f)^(1/2))/b^2/d/f^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2)+(
A*b^2-a*(B*b-C*a))*e^(1/2)*(1+d*x^2/c)^(1/2)*(1-f*x^2/e)^(1/2)*EllipticPi(
f^(1/2)*x/e^(1/2),-b*e/a/f,(-d*e/c/f)^(1/2))/a/b^2/f^(1/2)/(d*x^2+c)^(1/2)
/(-f*x^2+e)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 7.33 (sec) , antiderivative size = 206, normalized size of antiderivative = 0.64

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e - fx^2}} dx$$

$$= \frac{i\sqrt{1 + \frac{dx^2}{c}}\sqrt{1 - \frac{fx^2}{e}}\left(abCeE\left(\operatorname{arcsinh}\left(\sqrt{\frac{d}{c}}x\right)\middle| -\frac{cf}{de}\right) + a(aCf - b(Ce + Bf))\operatorname{EllipticF}\left(\operatorname{arcsinh}\left(\sqrt{\frac{d}{c}}x\right)\right)\right)}{ab^2\sqrt{\frac{d}{c}}f\sqrt{c + dx^2}\sqrt{e - fx^2}}$$

input

```
Integrate[(A + B*x^2 + C*x^4)/((a + b*x^2)*Sqrt[c + d*x^2]*Sqrt[e - f*x^2]),x]
```

output

```
(I*Sqrt[1 + (d*x^2)/c]*Sqrt[1 - (f*x^2)/e]*(a*b*C*e*EllipticE[I*ArcSinh[Sqrt[d/c]*x], -((c*f)/(d*e))] + a*(a*C*f - b*(C*e + B*f))*EllipticF[I*ArcSinh[Sqrt[d/c]*x], -((c*f)/(d*e))] - (A*b^2 + a*(-(b*B) + a*C))*f*EllipticPi[(b*c)/(a*d), I*ArcSinh[Sqrt[d/c]*x], -((c*f)/(d*e))])/((a*b^2*Sqrt[d/c]*f*Sqrt[c + d*x^2]*Sqrt[e - f*x^2]))
```

Rubi [A] (verified)

Time = 1.24 (sec) , antiderivative size = 409, normalized size of antiderivative = 1.27, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.044$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e - fx^2}} dx$$

$$\downarrow \text{7276}$$

$$\int \left(\frac{a^2C - abB + Ab^2}{b^2(a + bx^2)\sqrt{c + dx^2}\sqrt{e - fx^2}} + \frac{bB - aC}{b^2\sqrt{c + dx^2}\sqrt{e - fx^2}} + \frac{Cx^2}{b\sqrt{c + dx^2}\sqrt{e - fx^2}} \right) dx$$

$$\downarrow \text{2009}$$

$$\begin{aligned}
& \frac{\sqrt{e}\sqrt{\frac{dx^2}{c} + 1}\sqrt{1 - \frac{fx^2}{e}}(Ab^2 - a(bB - aC)) \operatorname{EllipticPi}\left(-\frac{be}{af}, \arcsin\left(\frac{\sqrt{fx}}{\sqrt{e}}\right), -\frac{de}{cf}\right)}{ab^2\sqrt{f}\sqrt{c + dx^2}\sqrt{e - fx^2}} + \\
& \frac{\sqrt{e}\sqrt{\frac{dx^2}{c} + 1}\sqrt{1 - \frac{fx^2}{e}}(bB - aC) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{fx}}{\sqrt{e}}\right), -\frac{de}{cf}\right)}{b^2\sqrt{f}\sqrt{c + dx^2}\sqrt{e - fx^2}} - \\
& \frac{cC\sqrt{e}\sqrt{\frac{dx^2}{c} + 1}\sqrt{1 - \frac{fx^2}{e}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{fx}}{\sqrt{e}}\right), -\frac{de}{cf}\right)}{bd\sqrt{f}\sqrt{c + dx^2}\sqrt{e - fx^2}} + \\
& \frac{C\sqrt{e}\sqrt{c + dx^2}\sqrt{1 - \frac{fx^2}{e}} E\left(\arcsin\left(\frac{\sqrt{fx}}{\sqrt{e}}\right) \middle| -\frac{de}{cf}\right)}{bd\sqrt{f}\sqrt{\frac{dx^2}{c} + 1}\sqrt{e - fx^2}}
\end{aligned}$$

input `Int[(A + B*x^2 + C*x^4)/((a + b*x^2)*Sqrt[c + d*x^2]*Sqrt[e - f*x^2]),x]`

output `(C*Sqrt[e]*Sqrt[c + d*x^2]*Sqrt[1 - (f*x^2)/e]*EllipticE[ArcSin[(Sqrt[f]*x)/Sqrt[e]], -((d*e)/(c*f))]/(b*d*Sqrt[f]*Sqrt[1 + (d*x^2)/c]*Sqrt[e - f*x^2]) + ((b*B - a*C)*Sqrt[e]*Sqrt[1 + (d*x^2)/c]*Sqrt[1 - (f*x^2)/e]*EllipticF[ArcSin[(Sqrt[f]*x)/Sqrt[e]], -((d*e)/(c*f))]/(b^2*Sqrt[f]*Sqrt[c + d*x^2]*Sqrt[e - f*x^2]) - (c*C*Sqrt[e]*Sqrt[1 + (d*x^2)/c]*Sqrt[1 - (f*x^2)/e]*EllipticF[ArcSin[(Sqrt[f]*x)/Sqrt[e]], -((d*e)/(c*f))]/(b*d*Sqrt[f]*Sqrt[c + d*x^2]*Sqrt[e - f*x^2]) + ((A*b^2 - a*(b*B - a*C))*Sqrt[e]*Sqrt[1 + (d*x^2)/c]*Sqrt[1 - (f*x^2)/e]*EllipticPi[-((b*e)/(a*f)), ArcSin[(Sqrt[f]*x)/Sqrt[e]], -((d*e)/(c*f))]/(a*b^2*Sqrt[f]*Sqrt[c + d*x^2]*Sqrt[e - f*x^2]))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 5.84 (sec) , antiderivative size = 328, normalized size of antiderivative = 1.02

method	result
default	$\left(A \operatorname{EllipticPi}\left(x\sqrt{\frac{f}{e}}, -\frac{be}{af}, \sqrt{\frac{-d}{f/e}}\right) b^2 d + B \operatorname{EllipticF}\left(x\sqrt{\frac{f}{e}}, \sqrt{-\frac{de}{cf}}\right) abd - B \operatorname{EllipticPi}\left(x\sqrt{\frac{f}{e}}, -\frac{be}{af}, \sqrt{\frac{-d}{f/e}}\right) abd - C \operatorname{EllipticF}\left(x\sqrt{\frac{f}{e}}, \sqrt{-\frac{de}{cf}}\right) abd \right) dx$
elliptic	$\frac{\sqrt{(x^2 d + c)(-f x^2 + e)} \left(\frac{B \sqrt{1 - \frac{f x^2}{e}} \sqrt{1 + \frac{d x^2}{c}} \operatorname{EllipticF}\left(x\sqrt{\frac{f}{e}}, \sqrt{-1 - \frac{-cf + de}{cf}}\right)}{b \sqrt{\frac{f}{e}} \sqrt{-df x^4 - cf x^2 + de x^2 + ce}} - \frac{\sqrt{1 - \frac{f x^2}{e}} \sqrt{1 + \frac{d x^2}{c}} \operatorname{EllipticF}\left(x\sqrt{\frac{f}{e}}, \sqrt{-1 - \frac{-cf + de}{cf}}\right) C a}{b^2 \sqrt{\frac{f}{e}} \sqrt{-df x^4 - cf x^2 + de x^2 + ce}} \right)}{\sqrt{(x^2 d + c)(-f x^2 + e)}}$

input `int((C*x^4+B*x^2+A)/(b*x^2+a)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2),x,method=_RETURNVERBOSE)`

output `(A*EllipticPi(x*(f/e)^(1/2),-b*e/a/f,(-1/c*d)^(1/2)/(f/e)^(1/2))*b^2*d+B*EllipticF(x*(f/e)^(1/2),(-d*e/c/f)^(1/2))*a*b*d-B*EllipticPi(x*(f/e)^(1/2),-b*e/a/f,(-1/c*d)^(1/2)/(f/e)^(1/2))*a*b*d-C*EllipticF(x*(f/e)^(1/2),(-d*e/c/f)^(1/2))*a^2*d-C*EllipticF(x*(f/e)^(1/2),(-d*e/c/f)^(1/2))*a*b*c+C*EllipticE(x*(f/e)^(1/2),(-d*e/c/f)^(1/2))*a*b*c+C*EllipticPi(x*(f/e)^(1/2),-b*e/a/f,(-1/c*d)^(1/2)/(f/e)^(1/2))*a^2*d)*((d*x^2+c)/c)^(1/2)*((-f*x^2+e)/e)^(1/2)*(d*x^2+c)^(1/2)*(-f*x^2+e)^(1/2)/d/a/(f/e)^(1/2)/b^2/(-d*f*x^4-c*f*x^2+d*e*x^2+c*e)`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2) \sqrt{c + dx^2} \sqrt{e - fx^2}} dx = \text{Timed out}$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2),x,algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e - fx^2}} dx = \int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e - fx^2}} dx$$

input `integrate((C*x**4+B*x**2+A)/(b*x**2+a)/(d*x**2+c)**(1/2)/(-f*x**2+e)**(1/2),x)`

output `Integral((A + B*x**2 + C*x**4)/((a + b*x**2)*sqrt(c + d*x**2)*sqrt(e - f*x**2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e - fx^2}} dx = \int \frac{Cx^4 + Bx^2 + A}{(bx^2 + a)\sqrt{dx^2 + c}\sqrt{-fx^2 + e}} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2),x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)/((b*x^2 + a)*sqrt(d*x^2 + c)*sqrt(-f*x^2 + e)), x)`

Giac [F]

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e - fx^2}} dx = \int \frac{Cx^4 + Bx^2 + A}{(bx^2 + a)\sqrt{dx^2 + c}\sqrt{-fx^2 + e}} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2),x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)/((b*x^2 + a)*sqrt(d*x^2 + c)*sqrt(-f*x^2 + e)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e - fx^2}} dx = \int \frac{Cx^4 + Bx^2 + A}{(bx^2 + a)\sqrt{dx^2 + c}\sqrt{e - fx^2}} dx$$

input

```
int((A + B*x^2 + C*x^4)/((a + b*x^2)*(c + d*x^2)^(1/2)*(e - f*x^2)^(1/2)),
x)
```

output

```
int((A + B*x^2 + C*x^4)/((a + b*x^2)*(c + d*x^2)^(1/2)*(e - f*x^2)^(1/2)),
x)
```

Reduce [F]

$$\begin{aligned} & \int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c + dx^2}\sqrt{e - fx^2}} dx \\ &= \left(\int \frac{\sqrt{-fx^2 + e}\sqrt{dx^2 + c}x^4}{-bdfx^6 - adfx^4 - bcfx^4 + bdex^4 - acfx^2 + adex^2 + bce x^2 + ace} dx \right) c \\ &+ \left(\int \frac{\sqrt{-fx^2 + e}\sqrt{dx^2 + c}x^2}{-bdfx^6 - adfx^4 - bcfx^4 + bdex^4 - acfx^2 + adex^2 + bce x^2 + ace} dx \right) b \\ &+ \left(\int \frac{\sqrt{-fx^2 + e}\sqrt{dx^2 + c}}{-bdfx^6 - adfx^4 - bcfx^4 + bdex^4 - acfx^2 + adex^2 + bce x^2 + ace} dx \right) a \end{aligned}$$

input

```
int((C*x^4+B*x^2+A)/(b*x^2+a)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2),x)
```

output

```
int((sqrt(e - f*x**2)*sqrt(c + d*x**2)*x**4)/(a*c*e - a*c*f*x**2 + a*d*e*x
**2 - a*d*f*x**4 + b*c*e*x**2 - b*c*f*x**4 + b*d*e*x**4 - b*d*f*x**6),x)*c
+ int((sqrt(e - f*x**2)*sqrt(c + d*x**2)*x**2)/(a*c*e - a*c*f*x**2 + a*d*
e*x**2 - a*d*f*x**4 + b*c*e*x**2 - b*c*f*x**4 + b*d*e*x**4 - b*d*f*x**6),x
)*b + int((sqrt(e - f*x**2)*sqrt(c + d*x**2))/(a*c*e - a*c*f*x**2 + a*d*e*
x**2 - a*d*f*x**4 + b*c*e*x**2 - b*c*f*x**4 + b*d*e*x**4 - b*d*f*x**6),x)*
a
```

3.21 $\int \frac{A+Bx^2+Cx^4}{(a+bx^2)\sqrt{c-dx^2}\sqrt{e-fx^2}} dx$

Optimal result	190
Mathematica [C] (verified)	191
Rubi [A] (verified)	191
Maple [A] (verified)	193
Fricas [F(-1)]	193
Sympy [F]	194
Maxima [F]	194
Giac [F]	194
Mupad [F(-1)]	195
Reduce [F]	195

Optimal result

Integrand size = 46, antiderivative size = 326

$$\int \frac{A+Bx^2+Cx^4}{(a+bx^2)\sqrt{c-dx^2}\sqrt{e-fx^2}} dx$$

$$= -\frac{\sqrt{c}C\sqrt{1-\frac{dx^2}{c}}\sqrt{e-fx^2}E\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\middle|\frac{cf}{de}\right)}{b\sqrt{d}f\sqrt{c-dx^2}\sqrt{1-\frac{fx^2}{e}}}$$

$$+ \frac{\sqrt{c}(bCe+bBf-aCf)\sqrt{1-\frac{dx^2}{c}}\sqrt{1-\frac{fx^2}{e}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right),\frac{cf}{de}\right)}{b^2\sqrt{d}f\sqrt{c-dx^2}\sqrt{e-fx^2}}$$

$$+ \frac{\sqrt{c}(Ab^2-a(bB-aC))\sqrt{1-\frac{dx^2}{c}}\sqrt{1-\frac{fx^2}{e}}\text{EllipticPi}\left(-\frac{bc}{ad},\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right),\frac{cf}{de}\right)}{ab^2\sqrt{d}\sqrt{c-dx^2}\sqrt{e-fx^2}}$$

output

```
-c^(1/2)*C*(1-d*x^2/c)^(1/2)*(-f*x^2+e)^(1/2)*EllipticE(d^(1/2)*x/c^(1/2),
(c*f/d/e)^(1/2))/b/d^(1/2)/f/(-d*x^2+c)^(1/2)/(1-f*x^2/e)^(1/2)+c^(1/2)*(B
*b*f-C*a*f+C*b*e)*(1-d*x^2/c)^(1/2)*(1-f*x^2/e)^(1/2)*EllipticF(d^(1/2)*x/
c^(1/2),(c*f/d/e)^(1/2))/b^2/d^(1/2)/f/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2)+c
^(1/2)*(A*b^2-a*(B*b-C*a))*(1-d*x^2/c)^(1/2)*(1-f*x^2/e)^(1/2)*EllipticPi(
d^(1/2)*x/c^(1/2),-b*c/a/d,(c*f/d/e)^(1/2))/a/b^2/d^(1/2)/(-d*x^2+c)^(1/2)
/(-f*x^2+e)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 7.34 (sec) , antiderivative size = 210, normalized size of antiderivative = 0.64

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx$$

$$= \frac{i\sqrt{1 - \frac{dx^2}{c}}\sqrt{1 - \frac{fx^2}{e}}\left(abCeE\left(\operatorname{arcsinh}\left(\sqrt{-\frac{d}{c}}x\right)\middle|\frac{cf}{de}\right) + a(aCf - b(Ce + Bf))\operatorname{EllipticF}\left(\operatorname{arcsinh}\left(\sqrt{-\frac{d}{c}}x\right)\middle|\frac{cf}{de}\right)\right)}{ab^2\sqrt{-\frac{d}{c}}f\sqrt{c - dx^2}\sqrt{e - fx^2}}$$

input

```
Integrate[(A + B*x^2 + C*x^4)/((a + b*x^2)*Sqrt[c - d*x^2]*Sqrt[e - f*x^2]),x]
```

output

```
(I*Sqrt[1 - (d*x^2)/c]*Sqrt[1 - (f*x^2)/e]*(a*b*C*e*EllipticE[I*ArcSinh[Sqrt[-(d/c)]*x], (c*f)/(d*e)] + a*(a*C*f - b*(C*e + B*f))*EllipticF[I*ArcSinh[Sqrt[-(d/c)]*x], (c*f)/(d*e)] - (A*b^2 + a*(-(b*B) + a*C))*f*EllipticPi[-((b*c)/(a*d)), I*ArcSinh[Sqrt[-(d/c)]*x], (c*f)/(d*e)))/(a*b^2*Sqrt[-(d/c)]*f*Sqrt[c - d*x^2]*Sqrt[e - f*x^2])
```

Rubi [A] (verified)

Time = 1.30 (sec) , antiderivative size = 413, normalized size of antiderivative = 1.27, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx$$

$$\downarrow \text{7276}$$

$$\int \left(\frac{a^2C - abB + Ab^2}{b^2(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} + \frac{bB - aC}{b^2\sqrt{c - dx^2}\sqrt{e - fx^2}} + \frac{Cx^2}{b\sqrt{c - dx^2}\sqrt{e - fx^2}} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{\sqrt{c}\sqrt{1-\frac{dx^2}{c}}\sqrt{1-\frac{fx^2}{e}}(Ab^2-a(bB-aC))\text{EllipticPi}\left(-\frac{bc}{ad}, \arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), \frac{cf}{de}\right)}{ab^2\sqrt{d}\sqrt{c-dx^2}\sqrt{e-fx^2}} +$$

$$\frac{\sqrt{e}\sqrt{1-\frac{dx^2}{c}}\sqrt{1-\frac{fx^2}{e}}(bB-aC)\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{fx}}{\sqrt{e}}\right), \frac{de}{cf}\right)}{b^2\sqrt{f}\sqrt{c-dx^2}\sqrt{e-fx^2}} +$$

$$\frac{cC\sqrt{e}\sqrt{1-\frac{dx^2}{c}}\sqrt{1-\frac{fx^2}{e}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{fx}}{\sqrt{e}}\right), \frac{de}{cf}\right)}{bd\sqrt{f}\sqrt{c-dx^2}\sqrt{e-fx^2}} -$$

$$\frac{C\sqrt{e}\sqrt{c-dx^2}\sqrt{1-\frac{fx^2}{e}}E\left(\arcsin\left(\frac{\sqrt{fx}}{\sqrt{e}}\right)\middle|\frac{de}{cf}\right)}{bd\sqrt{f}\sqrt{1-\frac{dx^2}{c}}\sqrt{e-fx^2}}$$

input `Int[(A + B*x^2 + C*x^4)/((a + b*x^2)*Sqrt[c - d*x^2]*Sqrt[e - f*x^2]),x]`

output `-((C*Sqrt[e]*Sqrt[c - d*x^2]*Sqrt[1 - (f*x^2)/e]*EllipticE[ArcSin[(Sqrt[f]*x)/Sqrt[e]], (d*e)/(c*f)])/(b*d*Sqrt[f]*Sqrt[1 - (d*x^2)/c]*Sqrt[e - f*x^2])) + ((b*B - a*C)*Sqrt[e]*Sqrt[1 - (d*x^2)/c]*Sqrt[1 - (f*x^2)/e]*EllipticF[ArcSin[(Sqrt[f]*x)/Sqrt[e]], (d*e)/(c*f)])/(b^2*Sqrt[f]*Sqrt[c - d*x^2]*Sqrt[e - f*x^2]) + (c*C*Sqrt[e]*Sqrt[1 - (d*x^2)/c]*Sqrt[1 - (f*x^2)/e]*EllipticF[ArcSin[(Sqrt[f]*x)/Sqrt[e]], (d*e)/(c*f)])/(b*d*Sqrt[f]*Sqrt[c - d*x^2]*Sqrt[e - f*x^2]) + (Sqrt[c]*(A*b^2 - a*(b*B - a*C))*Sqrt[1 - (d*x^2)/c]*Sqrt[1 - (f*x^2)/e]*EllipticPi[-((b*c)/(a*d)), ArcSin[(Sqrt[d]*x)/Sqrt[c]], (c*f)/(d*e)])/(a*b^2*Sqrt[d]*Sqrt[c - d*x^2]*Sqrt[e - f*x^2])`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 5.78 (sec) , antiderivative size = 323, normalized size of antiderivative = 0.99

method	result
default	$\left(A \operatorname{EllipticPi}\left(x\sqrt{\frac{f}{e}}, -\frac{be}{af}, \sqrt{\frac{d}{cf}}\right) b^2 d + B \operatorname{EllipticF}\left(x\sqrt{\frac{f}{e}}, \sqrt{\frac{de}{cf}}\right) abd - B \operatorname{EllipticPi}\left(x\sqrt{\frac{f}{e}}, -\frac{be}{af}, \sqrt{\frac{d}{cf}}\right) abd - C \operatorname{EllipticF}\left(x\sqrt{\frac{f}{e}}, \sqrt{\frac{de}{cf}}\right) da\sqrt{\frac{d}{cf}} \right)$
elliptic	$\sqrt{(-x^2 d + c)(-f x^2 + e)} \left(\frac{B \sqrt{1 - \frac{f x^2}{e}} \sqrt{1 - \frac{d x^2}{c}} \operatorname{EllipticF}\left(x\sqrt{\frac{f}{e}}, \sqrt{-1 - \frac{-cf - de}{cf}}\right)}{b \sqrt{\frac{f}{e}} \sqrt{df x^4 - cf x^2 - de x^2 + ce}} - \frac{\sqrt{1 - \frac{f x^2}{e}} \sqrt{1 - \frac{d x^2}{c}} \operatorname{EllipticF}\left(x\sqrt{\frac{f}{e}}, \sqrt{-1 - \frac{-cf - de}{cf}}\right)}{b^2 \sqrt{\frac{f}{e}} \sqrt{df x^4 - cf x^2 - de x^2 + ce}} \right) C$

input `int((C*x^4+B*x^2+A)/(b*x^2+a)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2),x,method=_RETURNVERBOSE)`

output `(A*EllipticPi(x*(f/e)^(1/2),-b*e/a/f,(1/c*d)^(1/2)/(f/e)^(1/2))*b^2*d+B*EllipticF(x*(f/e)^(1/2),(d*e/c/f)^(1/2))*a*b*d-B*EllipticPi(x*(f/e)^(1/2),-b*e/a/f,(1/c*d)^(1/2)/(f/e)^(1/2))*a*b*d-C*EllipticF(x*(f/e)^(1/2),(d*e/c/f)^(1/2))*a^2*d+C*EllipticF(x*(f/e)^(1/2),(d*e/c/f)^(1/2))*a*b*c-C*EllipticE(x*(f/e)^(1/2),(d*e/c/f)^(1/2))*a*b*c+C*EllipticPi(x*(f/e)^(1/2),-b*e/a/f,(1/c*d)^(1/2)/(f/e)^(1/2))*a^2*d*((-d*x^2+c)/c)^(1/2)*((-f*x^2+e)/e)^(1/2)*((-d*x^2+c)^(1/2)*(-f*x^2+e)^(1/2)/d/a/(f/e)^(1/2)/b^2/(d*f*x^4-c*f*x^2-d*e*x^2+c*e)`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx = \text{Timed out}$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2),x,algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx = \int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx$$

input `integrate((C*x**4+B*x**2+A)/(b*x**2+a)/(-d*x**2+c)**(1/2)/(-f*x**2+e)**(1/2),x)`

output `Integral((A + B*x**2 + C*x**4)/((a + b*x**2)*sqrt(c - d*x**2)*sqrt(e - f*x**2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx = \int \frac{Cx^4 + Bx^2 + A}{(bx^2 + a)\sqrt{-dx^2 + c}\sqrt{-fx^2 + e}} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2),x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)/((b*x^2 + a)*sqrt(-d*x^2 + c)*sqrt(-f*x^2 + e)), x)`

Giac [F]

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx = \int \frac{Cx^4 + Bx^2 + A}{(bx^2 + a)\sqrt{-dx^2 + c}\sqrt{-fx^2 + e}} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2),x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)/((b*x^2 + a)*sqrt(-d*x^2 + c)*sqrt(-f*x^2 + e)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx = \int \frac{Cx^4 + Bx^2 + A}{(bx^2 + a)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx$$

input

```
int((A + B*x^2 + C*x^4)/((a + b*x^2)*(c - d*x^2)^(1/2)*(e - f*x^2)^(1/2)),
x)
```

output

```
int((A + B*x^2 + C*x^4)/((a + b*x^2)*(c - d*x^2)^(1/2)*(e - f*x^2)^(1/2)),
x)
```

Reduce [F]

$$\begin{aligned} & \int \frac{A + Bx^2 + Cx^4}{(a + bx^2)\sqrt{c - dx^2}\sqrt{e - fx^2}} dx \\ &= \left(\int \frac{\sqrt{-fx^2 + e}\sqrt{-dx^2 + cx^4}}{bdfx^6 + adfx^4 - bcfx^4 - bde x^4 - acfx^2 - ade x^2 + bce x^2 + ace} dx \right) c \\ &+ \left(\int \frac{\sqrt{-fx^2 + e}\sqrt{-dx^2 + cx^2}}{bdfx^6 + adfx^4 - bcfx^4 - bde x^4 - acfx^2 - ade x^2 + bce x^2 + ace} dx \right) b \\ &+ \left(\int \frac{\sqrt{-fx^2 + e}\sqrt{-dx^2 + c}}{bdfx^6 + adfx^4 - bcfx^4 - bde x^4 - acfx^2 - ade x^2 + bce x^2 + ace} dx \right) a \end{aligned}$$

input

```
int((C*x^4+B*x^2+A)/(b*x^2+a)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(1/2),x)
```

output

```
int((sqrt(e - f*x**2)*sqrt(c - d*x**2)*x**4)/(a*c*e - a*c*f*x**2 - a*d*e*x
**2 + a*d*f*x**4 + b*c*e*x**2 - b*c*f*x**4 - b*d*e*x**4 + b*d*f*x**6),x)*c
+ int((sqrt(e - f*x**2)*sqrt(c - d*x**2)*x**2)/(a*c*e - a*c*f*x**2 - a*d*
e*x**2 + a*d*f*x**4 + b*c*e*x**2 - b*c*f*x**4 - b*d*e*x**4 + b*d*f*x**6),x
)*b + int((sqrt(e - f*x**2)*sqrt(c - d*x**2))/(a*c*e - a*c*f*x**2 - a*d*e*
x**2 + a*d*f*x**4 + b*c*e*x**2 - b*c*f*x**4 - b*d*e*x**4 + b*d*f*x**6),x)*
a
```

$$3.22 \quad \int \frac{A+Bx^2}{\sqrt{a+bx^2}(c+dx^2)^{3/2}(e+fx^2)} dx$$

Optimal result	196
Mathematica [C] (verified)	197
Rubi [A] (verified)	197
Maple [A] (verified)	199
Fricas [F(-1)]	199
Sympy [F]	200
Maxima [F]	200
Giac [F]	200
Mupad [F(-1)]	201
Reduce [F]	201

Optimal result

Integrand size = 39, antiderivative size = 377

$$\int \frac{A+Bx^2}{\sqrt{a+bx^2}(c+dx^2)^{3/2}(e+fx^2)} dx = \frac{\sqrt{d}(Bc-Ad)\sqrt{a+bx^2}E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\mid 1-\frac{bc}{ad}\right)}{\sqrt{c}(bc-ad)(de-cf)\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}}$$

$$\frac{\sqrt{c}(aBd^2e-bBc^2f-Ad(bde-2bcf+adf))\sqrt{a+bx^2}\text{EllipticF}\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1-\frac{bc}{ad}\right)}{a\sqrt{d}(bc-ad)(de-cf)^2\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}}$$

$$-\frac{c^{3/2}f(Be-Af)\sqrt{a+bx^2}\text{EllipticPi}\left(1-\frac{cf}{de}, \arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1-\frac{bc}{ad}\right)}{a\sqrt{de}(de-cf)^2\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}}$$

output

```
d^(1/2)*(-A*d+B*c)*(b*x^2+a)^(1/2)*EllipticE(d^(1/2)*x/c^(1/2)/(1+d*x^2/c)
^(1/2), (1-b*c/a/d)^(1/2))/c^(1/2)/(-a*d+b*c)/(-c*f+d*e)/(c*(b*x^2+a)/a/(d*
x^2+c))^(1/2)/(d*x^2+c)^(1/2)-c^(1/2)*(a*B*d^2*e-b*B*c^2*f-A*d*(a*d*f-2*b*
c*f+b*d*e))*(b*x^2+a)^(1/2)*InverseJacobiAM(arctan(d^(1/2)*x/c^(1/2)), (1-b
*c/a/d)^(1/2))/a/d^(1/2)/(-a*d+b*c)/(-c*f+d*e)^2/(c*(b*x^2+a)/a/(d*x^2+c))
^(1/2)/(d*x^2+c)^(1/2)-c^(3/2)*f*(-A*f+B*e)*(b*x^2+a)^(1/2)*EllipticPi(d^(
1/2)*x/c^(1/2)/(1+d*x^2/c)^(1/2), 1-c*f/d/e, (1-b*c/a/d)^(1/2))/a/d^(1/2)/e/
(-c*f+d*e)^2/(c*(b*x^2+a)/a/(d*x^2+c))^(1/2)/(d*x^2+c)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 11.12 (sec) , antiderivative size = 240, normalized size of antiderivative = 0.64

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \frac{\sqrt{\frac{b}{a}} d (Bc - Ad) e x (a + bx^2) + ibc (Bc - Ad) e \sqrt{1 + \frac{bx^2}{a}} \sqrt{1 + \frac{dx^2}{c}} E\left(\frac{b}{a}\right) + I \sqrt{\frac{b}{a}} d (Bc - Ad) e \sqrt{1 + \frac{bx^2}{a}} \sqrt{1 + \frac{dx^2}{c}} \operatorname{EllipticE}\left[\operatorname{ArcSinh}\left[\sqrt{\frac{b}{a}} x\right], \frac{a*d}{b*c}\right] + I c * (-b*c) + a*d * (B*e - A*f) * \sqrt{1 + \frac{bx^2}{a}} * \sqrt{1 + \frac{dx^2}{c}} * \operatorname{EllipticPi}\left[\frac{a*f}{b*e}, \operatorname{ArcSinh}\left[\sqrt{\frac{b}{a}} x\right], \frac{a*d}{b*c}\right]}{\sqrt{\frac{b}{a}} d (Bc - Ad) e x (a + bx^2) + ibc (Bc - Ad) e \sqrt{1 + \frac{bx^2}{a}} \sqrt{1 + \frac{dx^2}{c}} E\left(\frac{b}{a}\right) + I \sqrt{\frac{b}{a}} d (Bc - Ad) e \sqrt{1 + \frac{bx^2}{a}} \sqrt{1 + \frac{dx^2}{c}} \operatorname{EllipticE}\left[\operatorname{ArcSinh}\left[\sqrt{\frac{b}{a}} x\right], \frac{a*d}{b*c}\right] + I c * (-b*c) + a*d * (B*e - A*f) * \sqrt{1 + \frac{bx^2}{a}} * \sqrt{1 + \frac{dx^2}{c}} * \operatorname{EllipticPi}\left[\frac{a*f}{b*e}, \operatorname{ArcSinh}\left[\sqrt{\frac{b}{a}} x\right], \frac{a*d}{b*c}\right]}$$

input `Integrate[(A + B*x^2)/(Sqrt[a + b*x^2]*(c + d*x^2)^(3/2)*(e + f*x^2)),x]`

output `(Sqrt[b/a]*d*(B*c - A*d)*e*x*(a + b*x^2) + I*b*c*(B*c - A*d)*e*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] + I*c*(-(b*c) + a*d)*(B*e - A*f)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticPi[(a*f)/(b*e), I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)]/(Sqrt[b/a]*c*(-(b*c) + a*d)*e*(-(d*e) + c*f)*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])`

Rubi [A] (verified)

Time = 1.47 (sec) , antiderivative size = 572, normalized size of antiderivative = 1.52, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.051$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx$$

↓ 7276

$$\int \left(\frac{Af - Be}{f\sqrt{a + bx^2} (c + dx^2)^{3/2} (e + fx^2)} + \frac{B}{f\sqrt{a + bx^2} (c + dx^2)^{3/2}} \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{c^{3/2} f \sqrt{a + bx^2} (Be - Af) \operatorname{EllipticPi} \left(1 - \frac{cf}{de}, \arctan \left(\frac{\sqrt{dx}}{\sqrt{c}} \right), 1 - \frac{bc}{ad} \right)}{a \sqrt{de} \sqrt{c + dx^2} (de - cf)^2 \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \\
& \frac{d^{3/2} \sqrt{a + bx^2} (Be - Af) E \left(\arctan \left(\frac{\sqrt{dx}}{\sqrt{c}} \right) \middle| 1 - \frac{bc}{ad} \right)}{\sqrt{cf} \sqrt{c + dx^2} (bc - ad) (de - cf) \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} - \\
& \frac{\sqrt{c} \sqrt{d} \sqrt{a + bx^2} (Be - Af) (adf - 2bcf + bde) \operatorname{EllipticF} \left(\arctan \left(\frac{\sqrt{dx}}{\sqrt{c}} \right), 1 - \frac{bc}{ad} \right)}{af \sqrt{c + dx^2} (bc - ad) (de - cf)^2 \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \\
& \frac{bB \sqrt{c} \sqrt{a + bx^2} \operatorname{EllipticF} \left(\arctan \left(\frac{\sqrt{dx}}{\sqrt{c}} \right), 1 - \frac{bc}{ad} \right)}{a \sqrt{d} f \sqrt{c + dx^2} (bc - ad) \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} - \frac{B \sqrt{d} \sqrt{a + bx^2} E \left(\arctan \left(\frac{\sqrt{dx}}{\sqrt{c}} \right) \middle| 1 - \frac{bc}{ad} \right)}{\sqrt{cf} \sqrt{c + dx^2} (bc - ad) \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}
\end{aligned}$$

input `Int[(A + B*x^2)/(Sqrt[a + b*x^2]*(c + d*x^2)^(3/2)*(e + f*x^2)),x]`

output `-((B*Sqrt[d]*Sqrt[a + b*x^2]*EllipticE[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)])/(Sqrt[c]*(b*c - a*d)*f*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2])) + (d^(3/2)*(B*e - A*f)*Sqrt[a + b*x^2]*EllipticE[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)])/(Sqrt[c]*(b*c - a*d)*f*(d*e - c*f)*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) + (b*B*Sqrt[c]*Sqrt[a + b*x^2]*EllipticF[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)])/(a*Sqrt[d]*(b*c - a*d)*f*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) - (Sqrt[c]*Sqrt[d]*(B*e - A*f)*(b*d*e - 2*b*c*f + a*d*f)*Sqrt[a + b*x^2]*EllipticF[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)])/(a*(b*c - a*d)*f*(d*e - c*f)^2*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) - (c^(3/2)*f*(B*e - A*f)*Sqrt[a + b*x^2]*EllipticPi[1 - (c*f)/(d*e), ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)])/(a*Sqrt[d]*e*(d*e - c*f)^2*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2])`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 8.57 (sec) , antiderivative size = 537, normalized size of antiderivative = 1.42

method	result
default	$\left(-A\sqrt{-\frac{b}{a}}bd^2ex^3+B\sqrt{-\frac{b}{a}}bcde x^3+A\sqrt{\frac{bx^2+a}{a}}\sqrt{\frac{x^2d+c}{c}}\operatorname{EllipticE}\left(x\sqrt{-\frac{b}{a}},\sqrt{\frac{ad}{bc}}\right)bcde+A\sqrt{\frac{bx^2+a}{a}}\sqrt{\frac{x^2d+c}{c}}\operatorname{EllipticPi}\left(x\sqrt{-\frac{b}{a}},\sqrt{\frac{ad}{bc}}\right)\right)$
elliptic	Expression too large to display

input `int((B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(3/2)/(f*x^2+e),x,method=_RETURNVE
RBOSE)`

output
$$\begin{aligned} &(-A*(-b/a)^{(1/2)}*b*d^2*e*x^3+B*(-b/a)^{(1/2)}*b*c*d*e*x^3+A*((b*x^2+a)/a)^{(1/2)} \\ &/2)*((d*x^2+c)/c)^{(1/2)}*\operatorname{EllipticE}(x*(-b/a)^{(1/2)},(a*d/b/c)^{(1/2)})*b*c*d*e+ \\ &A*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)}*\operatorname{EllipticPi}(x*(-b/a)^{(1/2)},a*f/b/ \\ &e,(-1/c*d)^{(1/2)}*(-b/a)^{(1/2)})*a*c*d*f-A*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c) \\ &^{(1/2)}*\operatorname{EllipticPi}(x*(-b/a)^{(1/2)},a*f/b/e,(-1/c*d)^{(1/2)}*(-b/a)^{(1/2)})*b*c^ \\ &2*f-B*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)}*\operatorname{EllipticE}(x*(-b/a)^{(1/2)},(a* \\ &d/b/c)^{(1/2)})*b*c^2*e-B*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)}*\operatorname{EllipticPi} \\ &(x*(-b/a)^{(1/2)},a*f/b/e,(-1/c*d)^{(1/2)}*(-b/a)^{(1/2)})*a*c*d*e+B*((b*x^2+a)/ \\ &a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)}*\operatorname{EllipticPi}(x*(-b/a)^{(1/2)},a*f/b/e,(-1/c*d)^{(1/2)} \\ &/2)*b*c^2*e-A*(-b/a)^{(1/2)}*a*d^2*e*x+B*(-b/a)^{(1/2)}*a*c*d*e* \\ &x*(d*x^2+c)^{(1/2)}*(b*x^2+a)^{(1/2)}/c/(a*d-b*c)/e/(-b/a)^{(1/2)}/(c*f-d*e)/(b \\ &*d*x^4+a*d*x^2+b*c*x^2+a*c) \end{aligned}$$

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \text{Timed out}$$

input `integrate((B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm
="fricas")`

output Timed out

Sympy [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \int \frac{A + Bx^2}{\sqrt{a + bx^2} (c + dx^2)^{\frac{3}{2}} (e + fx^2)} dx$$

input `integrate((B*x**2+A)/(b*x**2+a)**(1/2)/(d*x**2+c)**(3/2)/(f*x**2+e),x)`

output `Integral((A + B*x**2)/(sqrt(a + b*x**2)*(c + d*x**2)**(3/2)*(e + f*x**2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a} (dx^2 + c)^{\frac{3}{2}} (fx^2 + e)} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/(sqrt(b*x^2 + a)*(d*x^2 + c)^(3/2)*(f*x^2 + e)), x)`

Giac [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a} (dx^2 + c)^{\frac{3}{2}} (fx^2 + e)} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm="giac")`

output `integrate((B*x^2 + A)/(sqrt(b*x^2 + a)*(d*x^2 + c)^(3/2)*(f*x^2 + e)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a} (dx^2 + c)^{3/2} (fx^2 + e)} dx$$

input `int((A + B*x^2)/((a + b*x^2)^(1/2)*(c + d*x^2)^(3/2)*(e + f*x^2)),x)`

output `int((A + B*x^2)/((a + b*x^2)^(1/2)*(c + d*x^2)^(3/2)*(e + f*x^2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \int \frac{\sqrt{dx^2 + c} \sqrt{bx^2 + a}}{d^2 f x^6 + 2cdf x^4 + d^2 e x^4 + c^2 f x^2 + 2cde x^2 + c^2 e} dx$$

input `int((B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(3/2)/(f*x^2+e),x)`

output `int((sqrt(c + d*x**2)*sqrt(a + b*x**2))/(c**2*e + c**2*f*x**2 + 2*c*d*e*x**2 + 2*c*d*f*x**4 + d**2*e*x**4 + d**2*f*x**6),x)`

3.23 $\int \frac{A+Bx^2}{\sqrt{a+bx^2}(c-dx^2)^{3/2}(e+fx^2)} dx$

Optimal result	202
Mathematica [C] (verified)	203
Rubi [A] (verified)	203
Maple [B] (verified)	205
Fricas [F(-1)]	206
Sympy [F]	206
Maxima [F]	207
Giac [F]	207
Mupad [F(-1)]	208
Reduce [F]	208

Optimal result

Integrand size = 40, antiderivative size = 390

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \frac{d(Bc + Ad)x\sqrt{a + bx^2}}{c(bc + ad)(de + cf)\sqrt{c - dx^2}} - \frac{a\sqrt{d}(Bc + Ad)\sqrt{1 + \frac{bx^2}{a}}\sqrt{c - dx^2}E\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \mid -\frac{bc}{ad}\right)}{c^{3/2}(bc + ad)(de + cf)\sqrt{a + bx^2}\sqrt{1 - \frac{dx^2}{c}}} + \frac{(Bc + Ad)\sqrt{1 + \frac{bx^2}{a}}\sqrt{c - dx^2}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), -\frac{bc}{ad}\right)}{c^{3/2}\sqrt{d}(de + cf)\sqrt{a + bx^2}\sqrt{1 - \frac{dx^2}{c}}} - \frac{\sqrt{c}(Be - Af)\sqrt{1 + \frac{bx^2}{a}}\sqrt{1 - \frac{dx^2}{c}}\text{EllipticPi}\left(-\frac{cf}{de}, \arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), -\frac{bc}{ad}\right)}{\sqrt{de}(de + cf)\sqrt{a + bx^2}\sqrt{c - dx^2}}$$

output

```
d*(A*d+B*c)*x*(b*x^2+a)^(1/2)/c/(a*d+b*c)/(c*f+d*e)/(-d*x^2+c)^(1/2)-a*d^(1/2)*(A*d+B*c)*(1+b*x^2/a)^(1/2)*(-d*x^2+c)^(1/2)*EllipticE(d^(1/2)*x/c^(1/2),(-b*c/a/d)^(1/2))/c^(3/2)/(a*d+b*c)/(c*f+d*e)/(b*x^2+a)^(1/2)/(1-d*x^2/c)^(1/2)+(A*d+B*c)*(1+b*x^2/a)^(1/2)*(-d*x^2+c)^(1/2)*EllipticF(d^(1/2)*x/c^(1/2),(-b*c/a/d)^(1/2))/c^(3/2)/d^(1/2)/(c*f+d*e)/(b*x^2+a)^(1/2)/(1-d*x^2/c)^(1/2)-c^(1/2)*(-A*f+B*e)*(1+b*x^2/a)^(1/2)*(1-d*x^2/c)^(1/2)*EllipticPi(d^(1/2)*x/c^(1/2),-c*f/d/e,(-b*c/a/d)^(1/2))/d^(1/2)/e/(c*f+d*e)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 11.11 (sec) , antiderivative size = 240, normalized size of antiderivative = 0.62

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \frac{\sqrt{\frac{b}{a}} d (Bc + Ad) e x (a + bx^2) - i b c (Bc + Ad) e \sqrt{1 + \frac{bx^2}{a}} \sqrt{1 - \frac{dx^2}{c}}}{\dots}$$

input `Integrate[(A + B*x^2)/(Sqrt[a + b*x^2]*(c - d*x^2)^(3/2)*(e + f*x^2)),x]`

output `(Sqrt[b/a]*d*(B*c + A*d)*e*x*(a + b*x^2) - I*b*c*(B*c + A*d)*e*Sqrt[1 + (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], -((a*d)/(b*c))] + I*c*(b*c + a*d)*(B*e - A*f)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticPi[(a*f)/(b*e), I*ArcSinh[Sqrt[b/a]*x], -((a*d)/(b*c))]/(Sqrt[b/a]*c*(b*c + a*d)*e*(d*e + c*f)*Sqrt[a + b*x^2]*Sqrt[c - d*x^2])`

Rubi [A] (verified)

Time = 1.85 (sec) , antiderivative size = 750, normalized size of antiderivative = 1.92, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx$$

↓ 7276

$$\int \left(\frac{Af - Be}{f\sqrt{a + bx^2} (c - dx^2)^{3/2} (e + fx^2)} + \frac{B}{f\sqrt{a + bx^2} (c - dx^2)^{3/2}} \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{d^{3/2}\sqrt{a+bx^2}\sqrt{1-\frac{dx^2}{c}}(Be-Af)E\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\middle|-\frac{bc}{ad}\right)}{\sqrt{cf}\sqrt{\frac{bx^2}{a}+1}\sqrt{c-dx^2}(ad+bc)(cf+de)} - \\
& \frac{\sqrt{d}\sqrt{\frac{bx^2}{a}+1}\sqrt{1-\frac{dx^2}{c}}(Be-Af)(2cf+de)\operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right),-\frac{bc}{ad}\right)}{\sqrt{cf}\sqrt{a+bx^2}\sqrt{c-dx^2}(cf+de)^2} + \\
& \frac{\sqrt{c}\sqrt{d}\sqrt{\frac{bx^2}{a}+1}\sqrt{1-\frac{dx^2}{c}}(Be-Af)\operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right),-\frac{bc}{ad}\right)}{\sqrt{a+bx^2}\sqrt{c-dx^2}(cf+de)^2} - \\
& \frac{\sqrt{c}\sqrt{\frac{bx^2}{a}+1}\sqrt{1-\frac{dx^2}{c}}(Be-Af)\operatorname{EllipticPi}\left(-\frac{cf}{de},\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right),-\frac{bc}{ad}\right)}{\sqrt{de}\sqrt{a+bx^2}\sqrt{c-dx^2}(cf+de)} - \\
& \frac{d^2x\sqrt{a+bx^2}(Be-Af)}{cf\sqrt{c-dx^2}(ad+bc)(cf+de)} + \frac{B\sqrt{\frac{bx^2}{a}+1}\sqrt{1-\frac{dx^2}{c}}\operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right),-\frac{bc}{ad}\right)}{\sqrt{c}\sqrt{d}\sqrt{a+bx^2}\sqrt{c-dx^2}} - \\
& \frac{B\sqrt{d}\sqrt{a+bx^2}\sqrt{1-\frac{dx^2}{c}}E\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\middle|-\frac{bc}{ad}\right)}{\sqrt{c}\sqrt{\frac{bx^2}{a}+1}\sqrt{c-dx^2}(ad+bc)} + \frac{Bdx\sqrt{a+bx^2}}{cf\sqrt{c-dx^2}(ad+bc)}
\end{aligned}$$

input `Int[(A + B*x^2)/(Sqrt[a + b*x^2]*(c - d*x^2)^(3/2)*(e + f*x^2)),x]`

output `(B*d*x*Sqrt[a + b*x^2])/(c*(b*c + a*d)*f*Sqrt[c - d*x^2]) - (d^2*(B*e - A*f)*x*Sqrt[a + b*x^2])/(c*(b*c + a*d)*f*(d*e + c*f)*Sqrt[c - d*x^2]) - (B*Sqrt[d]*Sqrt[a + b*x^2]*Sqrt[1 - (d*x^2)/c]*EllipticE[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -((b*c)/(a*d))])/(Sqrt[c]*(b*c + a*d)*f*Sqrt[1 + (b*x^2)/a]*Sqrt[c - d*x^2]) + (d^(3/2)*(B*e - A*f)*Sqrt[a + b*x^2]*Sqrt[1 - (d*x^2)/c]*EllipticE[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -((b*c)/(a*d))])/(Sqrt[c]*(b*c + a*d)*f*(d*e + c*f)*Sqrt[1 + (b*x^2)/a]*Sqrt[c - d*x^2]) + (B*Sqrt[1 + (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticF[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -((b*c)/(a*d))])/(Sqrt[c]*Sqrt[d]*f*Sqrt[a + b*x^2]*Sqrt[c - d*x^2]) + (Sqrt[c]*Sqrt[d]*(B*e - A*f)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticF[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -((b*c)/(a*d))])/(d*e + c*f)^2*Sqrt[a + b*x^2]*Sqrt[c - d*x^2]) - (Sqrt[d]*(B*e - A*f)*(d*e + 2*c*f)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticF[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -((b*c)/(a*d))])/(Sqrt[c]*f*(d*e + c*f)^2*Sqrt[a + b*x^2]*Sqrt[c - d*x^2]) - (Sqrt[c]*(B*e - A*f)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticPi[-((c*f)/(d*e)), ArcSin[(Sqrt[d]*x)/Sqrt[c]], -((b*c)/(a*d))])/(Sqrt[d]*e*(d*e + c*f)*Sqrt[a + b*x^2]*Sqrt[c - d*x^2])`

Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 7276 Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] :> With[{v = RationalFunctionE
xprand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 751 vs. 2(341) = 682.

Time = 8.44 (sec) , antiderivative size = 752, normalized size of antiderivative = 1.93

method	result
elliptic	$\sqrt{(-x^2d+c)(bx^2+a)} \left(-\frac{(-bdx^2-ad)x(Ad+Bc)}{c(ad+bc)(cf+de)\sqrt{(x^2-\frac{c}{d})(-bdx^2-ad)}} + \frac{\sqrt{1-\frac{dx^2}{c}}\sqrt{1+\frac{bx^2}{a}} \operatorname{EllipticF}\left(x\sqrt{\frac{d}{c}}, \sqrt{-1-\frac{-ad+bc}{ad}}\right)Ad}{\sqrt{\frac{d}{c}}\sqrt{-bdx^4-adx^2+x^2bc+ac}c(cf+de)} + \frac{\sqrt{1-\frac{dx^2}{c}}}{\sqrt{\frac{d}{c}}}$
default	$\left(A\sqrt{\frac{d}{c}}bd^2ex^3 + B\sqrt{\frac{d}{c}}bcde x^3 + A\sqrt{\frac{-x^2d+c}{c}}\sqrt{\frac{bx^2+a}{a}} \operatorname{EllipticF}\left(x\sqrt{\frac{d}{c}}, \sqrt{-\frac{bc}{ad}}\right)ad^2e + A\sqrt{\frac{-x^2d+c}{c}}\sqrt{\frac{bx^2+a}{a}} \operatorname{EllipticF}\left(x\sqrt{\frac{d}{c}}, \sqrt{\frac{d}{c}}\right) \right)$

```
input int((B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(3/2)/(f*x^2+e),x,method=_RETURNV
ERBOSE)
```

output

```
((-d*x^2+c)*(b*x^2+a))^(1/2)/(-d*x^2+c)^(1/2)/(b*x^2+a)^(1/2)*(-(-b*d*x^2-
a*d)/c/(a*d+b*c)*x*(A*d+B*c)/(c*f+d*e)/((x^2-c/d)*(-b*d*x^2-a*d))^(1/2)+1/
(1/c*d)^(1/2)*(1-d*x^2/c)^(1/2)*(1+b*x^2/a)^(1/2)/(-b*d*x^4-a*d*x^2+b*c*x^
2+a*c)^(1/2)*EllipticF(x*(1/c*d)^(1/2),(-1-(-a*d+b*c)/a/d)^(1/2))/c/(c*f+d
*e)*A*d+1/(1/c*d)^(1/2)*(1-d*x^2/c)^(1/2)*(1+b*x^2/a)^(1/2)/(-b*d*x^4-a*d*
x^2+b*c*x^2+a*c)^(1/2)*EllipticF(x*(1/c*d)^(1/2),(-1-(-a*d+b*c)/a/d)^(1/2)
)/(c*f+d*e)*B-d^2/c/(a*d+b*c)/(c*f+d*e)*a/(1/c*d)^(1/2)*(1-d*x^2/c)^(1/2)*
(1+b*x^2/a)^(1/2)/(-b*d*x^4-a*d*x^2+b*c*x^2+a*c)^(1/2)*EllipticE(x*(1/c*d)
^(1/2),(-1-(-a*d+b*c)/a/d)^(1/2))*A-d/(a*d+b*c)/(c*f+d*e)*a/(1/c*d)^(1/2)*
(1-d*x^2/c)^(1/2)*(1+b*x^2/a)^(1/2)/(-b*d*x^4-a*d*x^2+b*c*x^2+a*c)^(1/2)*E
llipticE(x*(1/c*d)^(1/2),(-1-(-a*d+b*c)/a/d)^(1/2))*B+1/(c*f+d*e)/e*f/(1/c
*d)^(1/2)*(1-d*x^2/c)^(1/2)*(1+b*x^2/a)^(1/2)/(-b*d*x^4-a*d*x^2+b*c*x^2+a*
c)^(1/2)*EllipticPi(x*(1/c*d)^(1/2),-c*f/d/e,(-b/a)^(1/2)/(1/c*d)^(1/2))*A
-1/(c*f+d*e)/(1/c*d)^(1/2)*(1-d*x^2/c)^(1/2)*(1+b*x^2/a)^(1/2)/(-b*d*x^4-a
*d*x^2+b*c*x^2+a*c)^(1/2)*EllipticPi(x*(1/c*d)^(1/2),-c*f/d/e,(-b/a)^(1/2)
/(1/c*d)^(1/2))*B)
```

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \text{Timed out}$$

input

```
integrate((B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm
m="fricas")
```

output

Timed out

Sympy [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \int \frac{A + Bx^2}{\sqrt{a + bx^2} (c - dx^2)^{\frac{3}{2}} (e + fx^2)} dx$$

input

```
integrate((B*x**2+A)/(b*x**2+a)**(1/2)/(-d*x**2+c)**(3/2)/(f*x**2+e),x)
```

output `Integral((A + B*x**2)/(sqrt(a + b*x**2)*(c - d*x**2)**(3/2)*(e + f*x**2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a} (-dx^2 + c)^{\frac{3}{2}} (fx^2 + e)} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm m="maxima")`

output `integrate((B*x^2 + A)/(sqrt(b*x^2 + a)*(-d*x^2 + c)^(3/2)*(f*x^2 + e)), x)`

Giac [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a} (-dx^2 + c)^{\frac{3}{2}} (fx^2 + e)} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm m="giac")`

output `integrate((B*x^2 + A)/(sqrt(b*x^2 + a)*(-d*x^2 + c)^(3/2)*(f*x^2 + e)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a} (c - dx^2)^{3/2} (fx^2 + e)} dx$$

input `int((A + B*x^2)/((a + b*x^2)^(1/2)*(c - d*x^2)^(3/2)*(e + f*x^2)),x)`

output `int((A + B*x^2)/((a + b*x^2)^(1/2)*(c - d*x^2)^(3/2)*(e + f*x^2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \int \frac{\sqrt{-dx^2 + c} \sqrt{bx^2 + a}}{d^2 f x^6 - 2cdf x^4 + d^2 e x^4 + c^2 f x^2 - 2cde x^2 + c^2 e} dx$$

input `int((B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(3/2)/(f*x^2+e),x)`

output `int((sqrt(c - d*x**2)*sqrt(a + b*x**2))/(c**2*e + c**2*f*x**2 - 2*c*d*e*x**2 - 2*c*d*f*x**4 + d**2*e*x**4 + d**2*f*x**6),x)`

$$3.24 \quad \int \frac{A+Bx^2}{\sqrt{a-bx^2}(c+dx^2)^{3/2}(e+fx^2)} dx$$

Optimal result	209
Mathematica [C] (verified)	210
Rubi [A] (verified)	210
Maple [B] (verified)	212
Fricas [F(-1)]	213
Sympy [F]	213
Maxima [F]	213
Giac [F]	214
Mupad [F(-1)]	214
Reduce [F]	214

Optimal result

Integrand size = 40, antiderivative size = 294

$$\int \frac{A+Bx^2}{\sqrt{a-bx^2}(c+dx^2)^{3/2}(e+fx^2)} dx = -\frac{d(Bc-Ad)x\sqrt{a-bx^2}}{c(bc+ad)(de-cf)\sqrt{c+dx^2}} - \frac{\sqrt{a}\sqrt{b}(Bc-Ad)\sqrt{1-\frac{bx^2}{a}}\sqrt{c+dx^2}E\left(\arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|-\frac{ad}{bc}\right)}{c(bc+ad)(de-cf)\sqrt{a-bx^2}\sqrt{1+\frac{dx^2}{c}}} + \frac{\sqrt{a}(Be-Af)\sqrt{1-\frac{bx^2}{a}}\sqrt{1+\frac{dx^2}{c}}\text{EllipticPi}\left(-\frac{af}{be}, \arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right), -\frac{ad}{bc}\right)}{\sqrt{be}(de-cf)\sqrt{a-bx^2}\sqrt{c+dx^2}}$$

output

```
-d*(-A*d+B*c)*x*(-b*x^2+a)^(1/2)/c/(a*d+b*c)/(-c*f+d*e)/(d*x^2+c)^(1/2)-a^(1/2)*b^(1/2)*(-A*d+B*c)*(1-b*x^2/a)^(1/2)*(d*x^2+c)^(1/2)*EllipticE(b^(1/2)*x/a^(1/2),(-a*d/b/c)^(1/2))/c/(a*d+b*c)/(-c*f+d*e)/(-b*x^2+a)^(1/2)/(1+d*x^2/c)^(1/2)+a^(1/2)*(-A*f+B*e)*(1-b*x^2/a)^(1/2)*(1+d*x^2/c)^(1/2)*EllipticPi(b^(1/2)*x/a^(1/2),-a*f/b/e,(-a*d/b/c)^(1/2))/b^(1/2)/e/(-c*f+d*e)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 11.31 (sec) , antiderivative size = 250, normalized size of antiderivative = 0.85

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \frac{-\sqrt{-\frac{b}{a}} d(-Bc + Ad)ex(a - bx^2) - ibc(Bc - Ad)e\sqrt{1 - \frac{bx^2}{a}}\sqrt{1 - \frac{bx^2}{a}}}{\dots}$$

input `Integrate[(A + B*x^2)/(Sqrt[a - b*x^2]*(c + d*x^2)^(3/2)*(e + f*x^2)),x]`

output `(-(Sqrt[-(b/a)]*d*(-(B*c) + A*d)*e*x*(a - b*x^2)) - I*b*c*(B*c - A*d)*e*Sqrt[1 - (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[-(b/a)]*x], -(a*d)/(b*c)]) + I*c*(b*c + a*d)*(B*e - A*f)*Sqrt[1 - (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticPi[-((a*f)/(b*e)), I*ArcSinh[Sqrt[-(b/a)]*x], -(a*d)/(b*c))]/(Sqrt[-(b/a)]*c*(b*c + a*d)*e*(-(d*e) + c*f)*Sqrt[a - b*x^2]*Sqrt[c + d*x^2])`

Rubi [A] (verified)

Time = 1.52 (sec) , antiderivative size = 559, normalized size of antiderivative = 1.90, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx$$

↓ 7276

$$\int \left(\frac{Af - Be}{f\sqrt{a - bx^2} (c + dx^2)^{3/2} (e + fx^2)} + \frac{B}{f\sqrt{a - bx^2} (c + dx^2)^{3/2}} \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{\sqrt{ad}\sqrt{1-\frac{bx^2}{a}}\sqrt{\frac{dx^2}{c}+1}(Be-Af)\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right),-\frac{ad}{bc}\right)}{\sqrt{b}\sqrt{a-bx^2}\sqrt{c+dx^2}(de-cf)^2} - \\
& \frac{\sqrt{a}\sqrt{bd}\sqrt{1-\frac{bx^2}{a}}\sqrt{c+dx^2}(Be-Af)E\left(\arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|-\frac{ad}{bc}\right)}{cf\sqrt{a-bx^2}\sqrt{\frac{dx^2}{c}+1}(ad+bc)(de-cf)} - \\
& \frac{c^{3/2}f\sqrt{a-bx^2}(Be-Af)\text{EllipticPi}\left(1-\frac{cf}{de},\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right),\frac{bc}{ad}+1\right)}{a\sqrt{de}\sqrt{c+dx^2}(de-cf)^2\sqrt{\frac{c(a-bx^2)}{a(c+dx^2)}}} - \\
& \frac{d^2x\sqrt{a-bx^2}(Be-Af)}{cf\sqrt{c+dx^2}(ad+bc)(de-cf)} + \frac{\sqrt{a}\sqrt{b}B\sqrt{1-\frac{bx^2}{a}}\sqrt{c+dx^2}E\left(\arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|-\frac{ad}{bc}\right)}{cf\sqrt{a-bx^2}\sqrt{\frac{dx^2}{c}+1}(ad+bc)} + \\
& \frac{Bdx\sqrt{a-bx^2}}{cf\sqrt{c+dx^2}(ad+bc)}
\end{aligned}$$

input

```
Int[(A + B*x^2)/(Sqrt[a - b*x^2]*(c + d*x^2)^(3/2)*(e + f*x^2)),x]
```

output

```
(B*d*x*Sqrt[a - b*x^2])/(c*(b*c + a*d)*f*Sqrt[c + d*x^2]) - (d^2*(B*e - A*f)*x*Sqrt[a - b*x^2])/(c*(b*c + a*d)*f*(d*e - c*f)*Sqrt[c + d*x^2]) + (Sqrt[a]*Sqrt[b]*B*Sqrt[1 - (b*x^2)/a]*Sqrt[c + d*x^2]*EllipticE[ArcSin[(Sqrt[b]*x)/Sqrt[a]], -((a*d)/(b*c))])/(c*(b*c + a*d)*f*Sqrt[a - b*x^2]*Sqrt[1 + (d*x^2)/c]) - (Sqrt[a]*Sqrt[b]*d*(B*e - A*f)*Sqrt[1 - (b*x^2)/a]*Sqrt[c + d*x^2]*EllipticE[ArcSin[(Sqrt[b]*x)/Sqrt[a]], -((a*d)/(b*c))])/(c*(b*c + a*d)*f*(d*e - c*f)*Sqrt[a - b*x^2]*Sqrt[1 + (d*x^2)/c]) + (Sqrt[a]*d*(B*e - A*f)*Sqrt[1 - (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[ArcSin[(Sqrt[b]*x)/Sqrt[a]], -((a*d)/(b*c))])/(Sqrt[b]*(d*e - c*f)^2*Sqrt[a - b*x^2]*Sqrt[c + d*x^2]) - (c^(3/2)*f*(B*e - A*f)*Sqrt[a - b*x^2]*EllipticPi[1 - (c*f)/(d*e), ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 + (b*c)/(a*d)])/(a*Sqrt[d]*e*(d*e - c*f)^2*Sqrt[(c*(a - b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2])
```

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xexpand[u/(a + b*xn), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 535 vs. 2(260) = 520.

Time = 8.68 (sec) , antiderivative size = 536, normalized size of antiderivative = 1.82

method	result
default	$\left(A\sqrt{\frac{b}{a}} b d^2 e x^3 - B\sqrt{\frac{b}{a}} bcde x^3 - A\sqrt{\frac{-b x^2 + a}{a}} \sqrt{\frac{x^2 d + c}{c}} \operatorname{EllipticE}\left(x\sqrt{\frac{b}{a}}, \sqrt{-\frac{ad}{bc}}\right) bcde + A\sqrt{\frac{-b x^2 + a}{a}} \sqrt{\frac{x^2 d + c}{c}} \operatorname{EllipticPi}\left(x\sqrt{\frac{b}{a}}, -\right.$
elliptic	Expression too large to display

input `int((B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(3/2)/(f*x^2+e),x,method=_RETURNV
ERBOSE)`

output `(A*(b/a)^(1/2)*b*d^2*e*x^3-B*(b/a)^(1/2)*b*c*d*e*x^3-A*((-b*x^2+a)/a)^(1/2
)*((d*x^2+c)/c)^(1/2)*EllipticE(x*(b/a)^(1/2),(-a*d/b/c)^(1/2))*b*c*d*e+A*
((-b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticPi(x*(b/a)^(1/2),-a*f/b/e
,(-1/c*d)^(1/2)/(b/a)^(1/2))*a*c*d*f+A*((-b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(
1/2)*EllipticPi(x*(b/a)^(1/2),-a*f/b/e,(-1/c*d)^(1/2)/(b/a)^(1/2))*b*c^2*
f+B*((-b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE(x*(b/a)^(1/2),(-a*d
/b/c)^(1/2))*b*c^2*e-B*((-b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticPi
(x*(b/a)^(1/2),-a*f/b/e,(-1/c*d)^(1/2)/(b/a)^(1/2))*a*c*d*e-B*((-b*x^2+a)/
a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticPi(x*(b/a)^(1/2),-a*f/b/e,(-1/c*d)^(1
/2)/(b/a)^(1/2))*b*c^2*e-A*(b/a)^(1/2)*a*d^2*e*x+B*(b/a)^(1/2)*a*c*d*e*x)*
(d*x^2+c)^(1/2)*(-b*x^2+a)^(1/2)/c/(a*d+b*c)/e/(b/a)^(1/2)/(c*f-d*e)/(-b*d
*x^4+a*d*x^2-b*c*x^2+a*c)`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \text{Timed out}$$

input `integrate((B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm m="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \int \frac{A + Bx^2}{\sqrt{a - bx^2} (c + dx^2)^{\frac{3}{2}} (e + fx^2)} dx$$

input `integrate((B*x**2+A)/(-b*x**2+a)**(1/2)/(d*x**2+c)**(3/2)/(f*x**2+e),x)`

output `Integral((A + B*x**2)/(sqrt(a - b*x**2)*(c + d*x**2)**(3/2)*(e + f*x**2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Bx^2 + A}{\sqrt{-bx^2 + a} (dx^2 + c)^{\frac{3}{2}} (fx^2 + e)} dx$$

input `integrate((B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm m="maxima")`

output `integrate((B*x^2 + A)/(sqrt(-b*x^2 + a)*(d*x^2 + c)^(3/2)*(f*x^2 + e)), x)`

Giac [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Bx^2 + A}{\sqrt{-bx^2 + a} (dx^2 + c)^{3/2} (fx^2 + e)} dx$$

input `integrate((B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm m="giac")`

output `integrate((B*x^2 + A)/(sqrt(-b*x^2 + a)*(d*x^2 + c)^(3/2)*(f*x^2 + e)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Bx^2 + A}{\sqrt{a - bx^2} (dx^2 + c)^{3/2} (fx^2 + e)} dx$$

input `int((A + B*x^2)/((a - b*x^2)^(1/2)*(c + d*x^2)^(3/2)*(e + f*x^2)),x)`

output `int((A + B*x^2)/((a - b*x^2)^(1/2)*(c + d*x^2)^(3/2)*(e + f*x^2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \left(\int \frac{\sqrt{dx^2 + c}}{-bd^2fx^8 + ad^2fx^6 - 2bcdfx^6 - bd^2ex^6 + 2acdfx^4 + ad^2ex^4} dx \right) + \left(\int \frac{\sqrt{dx^2 + c} \sqrt{-bx^2 + a}}{-bd^2fx^8 + ad^2fx^6 - 2bcdfx^6 - bd^2ex^6 + 2acdfx^4 + ad^2ex^4 - bc^2fx^4 - 2bcdex^4 + ac^2fx^2 + 2ac^2e} dx \right)$$

input `int((B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(3/2)/(f*x^2+e),x)`

output

```
int((sqrt(c + d*x**2)*sqrt(a - b*x**2)*x**2)/(a*c**2*e + a*c**2*f*x**2 + 2
*a*c*d*e*x**2 + 2*a*c*d*f*x**4 + a*d**2*e*x**4 + a*d**2*f*x**6 - b*c**2*e*
x**2 - b*c**2*f*x**4 - 2*b*c*d*e*x**4 - 2*b*c*d*f*x**6 - b*d**2*e*x**6 - b
*d**2*f*x**8),x)*b + int((sqrt(c + d*x**2)*sqrt(a - b*x**2))/(a*c**2*e + a
*c**2*f*x**2 + 2*a*c*d*e*x**2 + 2*a*c*d*f*x**4 + a*d**2*e*x**4 + a*d**2*f*
x**6 - b*c**2*e*x**2 - b*c**2*f*x**4 - 2*b*c*d*e*x**4 - 2*b*c*d*f*x**6 - b
*d**2*e*x**6 - b*d**2*f*x**8),x)*a
```


3.25
$$\int \frac{A+Bx^2}{\sqrt{a-bx^2}(c-dx^2)^{3/2}(e+fx^2)} dx$$

Optimal result	216
Mathematica [C] (verified)	217
Rubi [B] (verified)	217
Maple [B] (verified)	219
Fricas [F(-1)]	220
Sympy [F]	221
Maxima [F]	221
Giac [F]	221
Mupad [F(-1)]	222
Reduce [F]	222

Optimal result

Integrand size = 41, antiderivative size = 294

$$\int \frac{A+Bx^2}{\sqrt{a-bx^2}(c-dx^2)^{3/2}(e+fx^2)} dx = -\frac{d(Bc+Ad)x\sqrt{a-bx^2}}{c(bc-ad)(de+cf)\sqrt{c-dx^2}} + \frac{\sqrt{a}\sqrt{b}(Bc+Ad)\sqrt{1-\frac{bx^2}{a}}\sqrt{c-dx^2}E\left(\arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|\frac{ad}{bc}\right)}{c(bc-ad)(de+cf)\sqrt{a-bx^2}\sqrt{1-\frac{dx^2}{c}}} - \frac{\sqrt{a}(Be-Af)\sqrt{1-\frac{bx^2}{a}}\sqrt{1-\frac{dx^2}{c}}\text{EllipticPi}\left(-\frac{af}{be},\arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right),\frac{ad}{bc}\right)}{\sqrt{be}(de+cf)\sqrt{a-bx^2}\sqrt{c-dx^2}}$$

output

```
-d*(A*d+B*c)*x*(-b*x^2+a)^(1/2)/c/(-a*d+b*c)/(c*f+d*e)/(-d*x^2+c)^(1/2)+a^(1/2)*b^(1/2)*(A*d+B*c)*(1-b*x^2/a)^(1/2)*(-d*x^2+c)^(1/2)*EllipticE(b^(1/2)*x/a^(1/2),(a*d/b/c)^(1/2))/c/(-a*d+b*c)/(c*f+d*e)/(-b*x^2+a)^(1/2)/(1-d*x^2/c)^(1/2)-a^(1/2)*(-A*f+B*e)*(1-b*x^2/a)^(1/2)*(1-d*x^2/c)^(1/2)*EllipticPi(b^(1/2)*x/a^(1/2),-a*f/b/e,(a*d/b/c)^(1/2))/b^(1/2)/e/(c*f+d*e)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 11.19 (sec) , antiderivative size = 249, normalized size of antiderivative = 0.85

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \frac{\sqrt{-\frac{b}{a}} d (Bc + Ad) e x (a - bx^2) + i b c (Bc + Ad) e \sqrt{1 - \frac{bx^2}{a}} \sqrt{1 - \frac{dx^2}{c}}}{\dots}$$

input `Integrate[(A + B*x^2)/(Sqrt[a - b*x^2]*(c - d*x^2)^(3/2)*(e + f*x^2)),x]`

output `(Sqrt[-(b/a)]*d*(B*c + A*d)*e*x*(a - b*x^2) + I*b*c*(B*c + A*d)*e*Sqrt[1 - (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[-(b/a)]*x], (a*d)/(b*c)] + I*c*(-(b*c) + a*d)*(B*e - A*f)*Sqrt[1 - (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticPi[-((a*f)/(b*e)), I*ArcSinh[Sqrt[-(b/a)]*x], (a*d)/(b*c)]/(Sqrt[-(b/a)]*c*(-(b*c) + a*d)*e*(d*e + c*f)*Sqrt[a - b*x^2]*Sqrt[c - d*x^2])`

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 673 vs. $2(294) = 588$.

Time = 1.79 (sec) , antiderivative size = 673, normalized size of antiderivative = 2.29, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.049$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx$$

↓ 7276

$$\int \left(\frac{Af - Be}{f\sqrt{a - bx^2} (c - dx^2)^{3/2} (e + fx^2)} + \frac{B}{f\sqrt{a - bx^2} (c - dx^2)^{3/2}} \right) dx$$

↓ 2009

$$\begin{aligned}
& - \frac{d^{3/2} \sqrt{a - bx^2} \sqrt{1 - \frac{dx^2}{c}} (Be - Af) E\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \middle| \frac{bc}{ad}\right)}{\sqrt{cf} \sqrt{1 - \frac{bx^2}{a}} \sqrt{c - dx^2} (bc - ad) (cf + de)} \\
& \frac{\sqrt{d} \sqrt{1 - \frac{bx^2}{a}} \sqrt{1 - \frac{dx^2}{c}} (Be - Af) (2cf + de) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), \frac{bc}{ad}\right)}{\sqrt{cf} \sqrt{a - bx^2} \sqrt{c - dx^2} (cf + de)^2} + \\
& \frac{\sqrt{c} \sqrt{d} \sqrt{1 - \frac{bx^2}{a}} \sqrt{1 - \frac{dx^2}{c}} (Be - Af) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), \frac{bc}{ad}\right)}{\sqrt{a - bx^2} \sqrt{c - dx^2} (cf + de)^2} - \\
& \frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{1 - \frac{dx^2}{c}} (Be - Af) \operatorname{EllipticPi}\left(-\frac{af}{be}, \arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right), \frac{ad}{bc}\right)}{\sqrt{be} \sqrt{a - bx^2} \sqrt{c - dx^2} (cf + de)} + \\
& \frac{d^2 x \sqrt{a - bx^2} (Be - Af)}{cf \sqrt{c - dx^2} (bc - ad) (cf + de)} + \frac{\sqrt{a} \sqrt{b} B \sqrt{1 - \frac{bx^2}{a}} \sqrt{c - dx^2} E\left(\arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) \middle| \frac{ad}{bc}\right)}{cf \sqrt{a - bx^2} \sqrt{1 - \frac{dx^2}{c}} (bc - ad)} - \\
& \frac{B dx \sqrt{a - bx^2}}{cf \sqrt{c - dx^2} (bc - ad)}
\end{aligned}$$

input `Int[(A + B*x^2)/(Sqrt[a - b*x^2]*(c - d*x^2)^(3/2)*(e + f*x^2)),x]`

output `-((B*d*x*Sqrt[a - b*x^2])/(c*(b*c - a*d)*f*Sqrt[c - d*x^2])) + (d^2*(B*e - A*f)*x*Sqrt[a - b*x^2])/(c*(b*c - a*d)*f*(d*e + c*f)*Sqrt[c - d*x^2]) + (Sqrt[a]*Sqrt[b]*B*Sqrt[1 - (b*x^2)/a]*Sqrt[c - d*x^2]*EllipticE[ArcSin[(Sqrt[b]*x)/Sqrt[a]], (a*d)/(b*c)])/(c*(b*c - a*d)*f*Sqrt[a - b*x^2]*Sqrt[1 - (d*x^2)/c]) - (d^(3/2)*(B*e - A*f)*Sqrt[a - b*x^2]*Sqrt[1 - (d*x^2)/c]*EllipticE[ArcSin[(Sqrt[d]*x)/Sqrt[c]], (b*c)/(a*d)])/(Sqrt[c]*(b*c - a*d)*f*(d*e + c*f)*Sqrt[1 - (b*x^2)/a]*Sqrt[c - d*x^2]) + (Sqrt[c]*Sqrt[d]*(B*e - A*f)*Sqrt[1 - (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticF[ArcSin[(Sqrt[d]*x)/Sqrt[c]], (b*c)/(a*d)])/(Sqrt[c]) - ((d*e + c*f)^2*Sqrt[a - b*x^2]*Sqrt[c - d*x^2]) - (Sqrt[d]*(B*e - A*f)*(d*e + 2*c*f)*Sqrt[1 - (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticF[ArcSin[(Sqrt[d]*x)/Sqrt[c]], (b*c)/(a*d)])/(Sqrt[c]*f*(d*e + c*f)^2*Sqrt[a - b*x^2]*Sqrt[c - d*x^2]) - (Sqrt[a]*(B*e - A*f)*Sqrt[1 - (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticPi[-((a*f)/(b*e)), ArcSin[(Sqrt[b]*x)/Sqrt[a]], (a*d)/(b*c)])/(Sqrt[b]*e*(d*e + c*f)*Sqrt[a - b*x^2]*Sqrt[c - d*x^2])`

Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 7276 Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xexpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 761 vs. 2(260) = 520.

Time = 8.65 (sec) , antiderivative size = 762, normalized size of antiderivative = 2.59

method	result
default	$\left(-A\sqrt{\frac{d}{c}}bd^2ex^3 - B\sqrt{\frac{d}{c}}bcde x^3 - A \operatorname{EllipticE}\left(x\sqrt{\frac{d}{c}}, \sqrt{\frac{bc}{ad}}\right)\sqrt{\frac{-x^2d+c}{c}}\sqrt{\frac{-bx^2+a}{a}}ad^2e + A \operatorname{EllipticPi}\left(x\sqrt{\frac{d}{c}}, -\frac{cf}{de}, \sqrt{\frac{b}{a}}\right)\sqrt{\frac{-x^2d+c}{c}}\right)$
elliptic	$\sqrt{(-x^2d+c)(-bx^2+a)}\left(-\frac{(bdx^2-ad)x(Ad+Bc)}{c(ad-bc)(cf+de)\sqrt{(x^2-\frac{c}{d})(bdx^2-ad)}} + \frac{\sqrt{1-\frac{dx^2}{c}}\sqrt{1-\frac{bx^2}{a}}\operatorname{EllipticF}\left(x\sqrt{\frac{d}{c}}, \sqrt{-1-\frac{-ad-bc}{ad}}\right)Ad}{\sqrt{\frac{d}{c}}\sqrt{bdx^4-adx^2-x^2bc+ac}c(cf+de)} + \frac{\sqrt{1-\frac{dx^2}{c}}}{\sqrt{1-\frac{dx^2}{c}}}\right)$

```
input int((B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(3/2)/(f*x^2+e), x, method=_RETURN
VERBOSE)
```

output

```
(-A*(1/c*d)^(1/2)*b*d^2*e*x^3-B*(1/c*d)^(1/2)*b*c*d*e*x^3-A*EllipticE(x*(1/c*d)^(1/2),(b*c/a/d)^(1/2))*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*a*d^2*e+A*EllipticPi(x*(1/c*d)^(1/2),-c*f/d/e,(b/a)^(1/2)/(1/c*d)^(1/2))*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*a*c*d*f-A*EllipticPi(x*(1/c*d)^(1/2),-c*f/d/e,(b/a)^(1/2)/(1/c*d)^(1/2))*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*b*c^2*f+A*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*EllipticF(x*(1/c*d)^(1/2),(b*c/a/d)^(1/2))*a*d^2*e-A*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*EllipticF(x*(1/c*d)^(1/2),(b*c/a/d)^(1/2))*b*c*d*e-B*EllipticE(x*(1/c*d)^(1/2),(b*c/a/d)^(1/2))*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*a*c*d*e-B*EllipticPi(x*(1/c*d)^(1/2),-c*f/d/e,(b/a)^(1/2)/(1/c*d)^(1/2))*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*a*c*d*e+B*EllipticPi(x*(1/c*d)^(1/2),-c*f/d/e,(b/a)^(1/2)/(1/c*d)^(1/2))*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*b*c^2*e+B*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*EllipticF(x*(1/c*d)^(1/2),(b*c/a/d)^(1/2))*a*c*d*e-B*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*EllipticF(x*(1/c*d)^(1/2),(b*c/a/d)^(1/2))*b*c^2*e+A*(1/c*d)^(1/2)*a*d^2*e*x+B*(1/c*d)^(1/2)*a*c*d*e*x*(-d*x^2+c)^(1/2)*(-b*x^2+a)^(1/2)/c/(a*d-b*c)/e/(1/c*d)^(1/2)/(c*f+d*e)/(b*d*x^4-a*d*x^2-b*c*x^2+a*c)
```

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \text{Timed out}$$

input

```
integrate((B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm="fricas")
```

output

Timed out

Sympy [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \int \frac{A + Bx^2}{\sqrt{a - bx^2} (c - dx^2)^{\frac{3}{2}} (e + fx^2)} dx$$

input `integrate((B*x**2+A)/(-b*x**2+a)**(1/2)/(-d*x**2+c)**(3/2)/(f*x**2+e),x)`

output `Integral((A + B*x**2)/(sqrt(a - b*x**2)*(c - d*x**2)**(3/2)*(e + f*x**2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Bx^2 + A}{\sqrt{-bx^2 + a} (-dx^2 + c)^{\frac{3}{2}} (fx^2 + e)} dx$$

input `integrate((B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/(sqrt(-b*x^2 + a)*(-d*x^2 + c)^(3/2)*(f*x^2 + e)), x)`

Giac [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Bx^2 + A}{\sqrt{-bx^2 + a} (-dx^2 + c)^{\frac{3}{2}} (fx^2 + e)} dx$$

input `integrate((B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm="giac")`

output `integrate((B*x^2 + A)/(sqrt(-b*x^2 + a)*(-d*x^2 + c)^(3/2)*(f*x^2 + e)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Bx^2 + A}{\sqrt{a - bx^2} (c - dx^2)^{3/2} (fx^2 + e)} dx$$

input `int((A + B*x^2)/((a - b*x^2)^(1/2)*(c - d*x^2)^(3/2)*(e + f*x^2)),x)`

output `int((A + B*x^2)/((a - b*x^2)^(1/2)*(c - d*x^2)^(3/2)*(e + f*x^2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \left(\int \frac{\sqrt{-dx^2 + c}}{-bd^2fx^8 + ad^2fx^6 + 2bcdfx^6 - bd^2ex^6 - 2acdfx^4 + ad^2e} \right. \\ \left. + \left(\int \frac{\sqrt{-dx^2 + c} \sqrt{-bx^2 + a}}{-bd^2fx^8 + ad^2fx^6 + 2bcdfx^6 - bd^2ex^6 - 2acdfx^4 + ad^2ex^4 - bc^2fx^4 + 2bcde x^4 + ac^2fx^2 - 2} \right) \right)$$

input `int((B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(3/2)/(f*x^2+e),x)`

output `int((sqrt(c - d*x**2)*sqrt(a - b*x**2)*x**2)/(a*c**2*e + a*c**2*f*x**2 - 2*a*c*d*e*x**2 - 2*a*c*d*f*x**4 + a*d**2*e*x**4 + a*d**2*f*x**6 - b*c**2*e*x**2 - b*c**2*f*x**4 + 2*b*c*d*e*x**4 + 2*b*c*d*f*x**6 - b*d**2*e*x**6 - b*d**2*f*x**8),x)*b + int((sqrt(c - d*x**2)*sqrt(a - b*x**2))/(a*c**2*e + a*c**2*f*x**2 - 2*a*c*d*e*x**2 - 2*a*c*d*f*x**4 + a*d**2*e*x**4 + a*d**2*f*x**6 - b*c**2*e*x**2 - b*c**2*f*x**4 + 2*b*c*d*e*x**4 + 2*b*c*d*f*x**6 - b*d**2*e*x**6 - b*d**2*f*x**8),x)*a`

3.26
$$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}(c+dx^2)^{3/2}(e+fx^2)} dx$$

Optimal result	223
Mathematica [C] (verified)	224
Rubi [A] (verified)	225
Maple [B] (verified)	227
Fricas [F(-1)]	228
Sympy [F]	228
Maxima [F]	228
Giac [F]	229
Mupad [F(-1)]	229
Reduce [F]	229

Optimal result

Integrand size = 44, antiderivative size = 415

$$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}(c+dx^2)^{3/2}(e+fx^2)} dx =$$

$$\frac{(c^2C - Bcd + Ad^2) \sqrt{a+bx^2} E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \mid 1 - \frac{bc}{ad}\right)}{\sqrt{c}\sqrt{d}(bc - ad)(de - cf) \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}} \sqrt{c+dx^2}}$$

$$+ \frac{\sqrt{c}(a(2cCde - c^2Cf - d^2(Be - Af)) + b(Ad^2e - 2Acdf - c^2(Ce - Bf))) \sqrt{a+bx^2} \text{EllipticF}\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right)}{a\sqrt{d}(bc - ad)(de - cf)^2 \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}} \sqrt{c+dx^2}}$$

$$+ \frac{c^{3/2}(Ce^2 - Bef + Af^2) \sqrt{a+bx^2} \text{EllipticPi}\left(1 - \frac{cf}{de}, \arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right)}{a\sqrt{de}(de - cf)^2 \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}} \sqrt{c+dx^2}}$$

output

```

-(A*d^2-B*c*d+C*c^2)*(b*x^2+a)^(1/2)*EllipticE(d^(1/2)*x/c^(1/2)/(1+d*x^2/
c)^(1/2),(1-b*c/a/d)^(1/2))/c^(1/2)/d^(1/2)/(-a*d+b*c)/(-c*f+d*e)/(c*(b*x^
2+a)/a/(d*x^2+c))^(1/2)/(d*x^2+c)^(1/2)+c^(1/2)*(a*(2*c*C*d*e-c^2*C*f-d^2*
(-A*f+B*e))+b*(A*d^2*e-2*A*c*d*f-c^2*(-B*f+C*e)))*(b*x^2+a)^(1/2)*InverseJ
acobiAM(arctan(d^(1/2)*x/c^(1/2)),(1-b*c/a/d)^(1/2))/a/d^(1/2)/(-a*d+b*c)/
(-c*f+d*e)^2/(c*(b*x^2+a)/a/(d*x^2+c))^(1/2)/(d*x^2+c)^(1/2)+c^(3/2)*(A*f^
2-B*e*f+C*e^2)*(b*x^2+a)^(1/2)*EllipticPi(d^(1/2)*x/c^(1/2)/(1+d*x^2/c)^(1
/2),1-c*f/d/e,(1-b*c/a/d)^(1/2))/a/d^(1/2)/e/(-c*f+d*e)^2/(c*(b*x^2+a)/a/(
d*x^2+c))^(1/2)/(d*x^2+c)^(1/2)

```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 10.89 (sec) , antiderivative size = 352, normalized size of antiderivative = 0.85

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \frac{-ibc(c^2C - Bcd + Ad^2) ef \sqrt{1 + \frac{bx^2}{a}} \sqrt{1 + \frac{dx^2}{c}} E\left(i \operatorname{arcsinh}\left(\sqrt{\frac{b}{a}} x\right)\right)}{\sqrt{a + bx^2} (c + dx^2)^{3/2} (e + fx^2)}$$

input

```

Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*(c + d*x^2)^(3/2)*(e + f*x^
2)),x]

```

output

```

((-I)*b*c*(c^2*C - B*c*d + A*d^2)*e*f*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)
/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - I*c*C*(-(b*c) + a*d)*
e*(-(d*e) + c*f)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSi
nh[Sqrt[b/a]*x], (a*d)/(b*c)] - d*(Sqrt[b/a]*(c^2*C - B*c*d + A*d^2)*e*f*x
*(a + b*x^2) + I*c*(-(b*c) + a*d)*(C*e^2 + f*(-(B*e) + A*f))*Sqrt[1 + (b*x
^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticPi[(a*f)/(b*e), I*ArcSinh[Sqrt[b/a]*x],
(a*d)/(b*c)))/(Sqrt[b/a]*c*d*(-(b*c) + a*d)*e*f*(-(d*e) + c*f)*Sqrt[a +
b*x^2]*Sqrt[c + d*x^2])

```

Rubi [A] (verified)

Time = 1.72 (sec) , antiderivative size = 805, normalized size of antiderivative = 1.94, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.045$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx$$

↓ 7276

$$\int \left(\frac{Af^2 - Bef + Ce^2}{f^2 \sqrt{a + bx^2} (c + dx^2)^{3/2} (e + fx^2)} - \frac{Ce - Bf}{f^2 \sqrt{a + bx^2} (c + dx^2)^{3/2}} + \frac{Cx^2}{f \sqrt{a + bx^2} (c + dx^2)^{3/2}} \right) dx$$

↓ 2009

$$\frac{(Ce^2 - Bfe + Af^2) \sqrt{bx^2 + a} \text{EllipticPi} \left(1 - \frac{cf}{de}, \arctan \left(\frac{\sqrt{dx}}{\sqrt{c}} \right), 1 - \frac{bc}{ad} \right) c^{3/2} +}{a\sqrt{de}(de - cf)^2 \sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}} \sqrt{dx^2 + c} + \frac{C\sqrt{bx^2 + a} E \left(\arctan \left(\frac{\sqrt{dx}}{\sqrt{c}} \right) \mid 1 - \frac{bc}{ad} \right) \sqrt{c}}{\sqrt{d}(bc - ad) f \sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}} \sqrt{dx^2 + c}} - \frac{b(Ce - Bf) \sqrt{bx^2 + a} \text{EllipticF} \left(\arctan \left(\frac{\sqrt{dx}}{\sqrt{c}} \right), 1 - \frac{bc}{ad} \right) \sqrt{c}}{a\sqrt{d}(bc - ad) f^2 \sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}} \sqrt{dx^2 + c}} + \frac{\sqrt{d}(bde - 2bcf + adf) (Ce^2 - Bfe + Af^2) \sqrt{bx^2 + a} \text{EllipticF} \left(\arctan \left(\frac{\sqrt{dx}}{\sqrt{c}} \right), 1 - \frac{bc}{ad} \right) \sqrt{c}}{a(bc - ad) f^2 (de - cf)^2 \sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}} \sqrt{dx^2 + c} + \frac{C\sqrt{bx^2 + a} \text{EllipticF} \left(\arctan \left(\frac{\sqrt{dx}}{\sqrt{c}} \right), 1 - \frac{bc}{ad} \right) \sqrt{c}}{\sqrt{d}(bc - ad) f \sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}} \sqrt{dx^2 + c}} + \frac{\sqrt{d}(Ce - Bf) \sqrt{bx^2 + a} E \left(\arctan \left(\frac{\sqrt{dx}}{\sqrt{c}} \right) \mid 1 - \frac{bc}{ad} \right)}{(bc - ad) f^2 \sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}} \sqrt{dx^2 + c} \sqrt{c}} - \frac{d^{3/2} (Ce^2 - Bfe + Af^2) \sqrt{bx^2 + a} E \left(\arctan \left(\frac{\sqrt{dx}}{\sqrt{c}} \right) \mid 1 - \frac{bc}{ad} \right)}{(bc - ad) f^2 (de - cf) \sqrt{\frac{c(bx^2+a)}{a(dx^2+c)}} \sqrt{dx^2 + c} \sqrt{c}}$$

input `Int[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*(c + d*x^2)^(3/2)*(e + f*x^2)),x]`

output `(Sqrt[c]*C*Sqrt[a + b*x^2]*EllipticE[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(Sqrt[d]*(b*c - a*d)*f*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) + (Sqrt[d]*(C*e - B*f)*Sqrt[a + b*x^2]*EllipticE[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(Sqrt[c]*(b*c - a*d)*f^2*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) - (d^(3/2)*(C*e^2 - B*e*f + A*f^2)*Sqrt[a + b*x^2]*EllipticE[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(Sqrt[c]*(b*c - a*d)*f^2*(d*e - c*f)*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) - (Sqrt[c]*C*Sqrt[a + b*x^2]*EllipticF[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(Sqrt[d]*(b*c - a*d)*f*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) - (b*Sqrt[c]*(C*e - B*f)*Sqrt[a + b*x^2]*EllipticF[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(a*Sqrt[d]*(b*c - a*d)*f^2*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) + (Sqrt[c]*Sqrt[d]*(b*d*e - 2*b*c*f + a*d*f)*(C*e^2 - B*e*f + A*f^2)*Sqrt[a + b*x^2]*EllipticF[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(a*(b*c - a*d)*f^2*(d*e - c*f)^2*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) + (c^(3/2)*(C*e^2 - B*e*f + A*f^2)*Sqrt[a + b*x^2]*EllipticPi[1 - (c*f)/(d*e), ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)]/(a*Sqrt[d]*e*(d*e - c*f)^2*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2])`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1041 vs. $2(401) = 802$.

Time = 8.39 (sec) , antiderivative size = 1042, normalized size of antiderivative = 2.51

method	result	size
default	Expression too large to display	1042
elliptic	Expression too large to display	1833

input

```
int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(3/2)/(f*x^2+e),x,method=_RE
TURNVERBOSE)
```

output

```
(-A*(-b/a)^(1/2)*b*d^3*e*f*x^3+B*(-b/a)^(1/2)*b*c*d^2*e*f*x^3-C*(-b/a)^(1/2)
)*b*c^2*d*e*f*x^3+A*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE(x*(
-b/a)^(1/2),(a*d/b/c)^(1/2))*b*c*d^2*e*f+A*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/
c)^(1/2)*EllipticPi(x*(-b/a)^(1/2),a*f/b/e,(-1/c*d)^(1/2)/(-b/a)^(1/2))*a*
c*d^2*f^2-A*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticPi(x*(-b/a)^(1
/2),a*f/b/e,(-1/c*d)^(1/2)/(-b/a)^(1/2))*b*c^2*d*f^2-B*((b*x^2+a)/a)^(1/2)
)*((d*x^2+c)/c)^(1/2)*EllipticE(x*(-b/a)^(1/2),(a*d/b/c)^(1/2))*b*c^2*d*e*f
-B*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticPi(x*(-b/a)^(1/2),a*f/b
/e,(-1/c*d)^(1/2)/(-b/a)^(1/2))*a*c*d^2*e*f+B*((b*x^2+a)/a)^(1/2)*((d*x^2+
c)/c)^(1/2)*EllipticPi(x*(-b/a)^(1/2),a*f/b/e,(-1/c*d)^(1/2)/(-b/a)^(1/2))
)*b*c^2*d*e*f+C*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE(x*(-b/a)^(
1/2),(a*d/b/c)^(1/2))*b*c^3*e*f+C*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)
)*EllipticPi(x*(-b/a)^(1/2),a*f/b/e,(-1/c*d)^(1/2)/(-b/a)^(1/2))*a*c*d^2*e^
2-C*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticPi(x*(-b/a)^(1/2),a*f/
b/e,(-1/c*d)^(1/2)/(-b/a)^(1/2))*b*c^2*d*e^2+C*((b*x^2+a)/a)^(1/2)*((d*x^2
+c)/c)^(1/2)*EllipticF(x*(-b/a)^(1/2),(a*d/b/c)^(1/2))*a*c^2*d*e*f-C*((b*x
^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF(x*(-b/a)^(1/2),(a*d/b/c)^(1/2)
))*a*c*d^2*e^2-C*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF(x*(-b/a)
)^(1/2),(a*d/b/c)^(1/2))*b*c^3*e*f+C*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/
2)*EllipticF(x*(-b/a)^(1/2),(a*d/b/c)^(1/2))*b*c^2*d*e^2-A*(-b/a)^(1/2)...
```

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \text{Timed out}$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm="fricas")`

output Timed out

Sympy [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c + dx^2)^{\frac{3}{2}} (e + fx^2)} dx$$

input `integrate((C*x**4+B*x**2+A)/(b*x**2+a)**(1/2)/(d*x**2+c)**(3/2)/(f*x**2+e),x)`

output `Integral((A + B*x**2 + C*x**4)/(sqrt(a + b*x**2)*(c + d*x**2)**(3/2)*(e + f*x**2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a} (dx^2 + c)^{\frac{3}{2}} (fx^2 + e)} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(b*x^2 + a)*(d*x^2 + c)^(3/2)*(f*x^2 + e)), x)`

Giac [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a} (dx^2 + c)^{3/2} (fx^2 + e)} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(b*x^2 + a)*(d*x^2 + c)^(3/2)*(f*x^2 + e)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a} (dx^2 + c)^{3/2} (fx^2 + e)} dx$$

input `int((A + B*x^2 + C*x^4)/((a + b*x^2)^(1/2)*(c + d*x^2)^(3/2)*(e + f*x^2)), x)`

output `int((A + B*x^2 + C*x^4)/((a + b*x^2)^(1/2)*(c + d*x^2)^(3/2)*(e + f*x^2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \left(\int \frac{\sqrt{dx^2 + c}}{bd^2fx^8 + ad^2fx^6 + 2bcdfx^6 + bd^2ex^6 + 2acdfx^4 + ad^2ex^4} \right) + \left(\int \frac{\sqrt{dx^2 + c}\sqrt{bx^2 + a}}{bd^2fx^8 + ad^2fx^6 + 2bcdfx^6 + bd^2ex^6 + 2acdfx^4 + ad^2ex^4 + bc^2fx^4 + 2bcdex^4 + ac^2fx^2 + 2ac} \right) + \left(\int \frac{\sqrt{dx^2 + c}\sqrt{bx^2 + a}}{bd^2fx^8 + ad^2fx^6 + 2bcdfx^6 + bd^2ex^6 + 2acdfx^4 + ad^2ex^4 + bc^2fx^4 + 2bcdex^4 + ac^2fx^2 + 2ac} \right)$$

input `int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(3/2)/(f*x^2+e),x)`

output `int((sqrt(c + d*x**2)*sqrt(a + b*x**2)*x**4)/(a*c**2*e + a*c**2*f*x**2 + 2*a*c*d*e*x**2 + 2*a*c*d*f*x**4 + a*d**2*e*x**4 + a*d**2*f*x**6 + b*c**2*e*x**2 + b*c**2*f*x**4 + 2*b*c*d*e*x**4 + 2*b*c*d*f*x**6 + b*d**2*e*x**6 + b*d**2*f*x**8),x)*c + int((sqrt(c + d*x**2)*sqrt(a + b*x**2)*x**2)/(a*c**2*e + a*c**2*f*x**2 + 2*a*c*d*e*x**2 + 2*a*c*d*f*x**4 + a*d**2*e*x**4 + a*d**2*f*x**6 + b*c**2*e*x**2 + b*c**2*f*x**4 + 2*b*c*d*e*x**4 + 2*b*c*d*f*x**6 + b*d**2*e*x**6 + b*d**2*f*x**8),x)*b + int((sqrt(c + d*x**2)*sqrt(a + b*x**2))/(a*c**2*e + a*c**2*f*x**2 + 2*a*c*d*e*x**2 + 2*a*c*d*f*x**4 + a*d**2*e*x**4 + a*d**2*f*x**6 + b*c**2*e*x**2 + b*c**2*f*x**4 + 2*b*c*d*e*x**4 + 2*b*c*d*f*x**6 + b*d**2*e*x**6 + b*d**2*f*x**8),x)*a`

$$3.27 \quad \int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}(c-dx^2)^{3/2}(e+fx^2)} dx$$

Optimal result	231
Mathematica [C] (verified)	232
Rubi [B] (verified)	232
Maple [B] (verified)	234
Fricas [F(-1)]	235
Sympy [F]	236
Maxima [F]	236
Giac [F]	237
Mupad [F(-1)]	237
Reduce [F]	237

Optimal result

Integrand size = 45, antiderivative size = 426

$$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}(c-dx^2)^{3/2}(e+fx^2)} dx = \frac{(c^2C+Bcd+Ad^2)x\sqrt{a+bx^2}}{c(bc+ad)(de+cf)\sqrt{c-dx^2}} - \frac{(c^2C+Bcd+Ad^2)\sqrt{a+bx^2}\sqrt{1-\frac{dx^2}{c}}E\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\middle|-\frac{bc}{ad}\right)}{\sqrt{c}\sqrt{d}(bc+ad)(de+cf)\sqrt{1+\frac{bx^2}{a}}\sqrt{c-dx^2}} - \frac{(cCe-Bcf-Adf)\sqrt{1+\frac{bx^2}{a}}\sqrt{1-\frac{dx^2}{c}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right),-\frac{bc}{ad}\right)}{\sqrt{c}\sqrt{d}f(de+cf)\sqrt{a+bx^2}\sqrt{c-dx^2}} + \frac{\sqrt{c}(Ce^2-Bef+Af^2)\sqrt{1+\frac{bx^2}{a}}\sqrt{1-\frac{dx^2}{c}}\text{EllipticPi}\left(-\frac{cf}{de},\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right),-\frac{bc}{ad}\right)}{\sqrt{def}(de+cf)\sqrt{a+bx^2}\sqrt{c-dx^2}}$$

output

```
(A*d^2+B*c*d+C*c^2)*x*(b*x^2+a)^(1/2)/c/(a*d+b*c)/(c*f+d*e)/(-d*x^2+c)^(1/2)-(A*d^2+B*c*d+C*c^2)*(b*x^2+a)^(1/2)*(1-d*x^2/c)^(1/2)*EllipticE(d^(1/2)*x/c^(1/2),(-b*c/a/d)^(1/2))/c^(1/2)/d^(1/2)/(a*d+b*c)/(c*f+d*e)/(1+b*x^2/a)^(1/2)/(-d*x^2+c)^(1/2)-(-A*d*f-B*c*f+C*c*e)*(1+b*x^2/a)^(1/2)*(1-d*x^2/c)^(1/2)*EllipticF(d^(1/2)*x/c^(1/2),(-b*c/a/d)^(1/2))/c^(1/2)/d^(1/2)/f/(c*f+d*e)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)+c^(1/2)*(A*f^2-B*e*f+C*e^2)*(1+b*x^2/a)^(1/2)*(1-d*x^2/c)^(1/2)*EllipticPi(d^(1/2)*x/c^(1/2),-c*f/d/e,(-b*c/a/d)^(1/2))/d^(1/2)/e/f/(c*f+d*e)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)
```


Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 10.95 (sec) , antiderivative size = 350, normalized size of antiderivative = 0.82

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \frac{\sqrt{\frac{b}{a}} d (c^2 C + Bcd + Ad^2) \operatorname{erf} x(a + bx^2) - ibc(c^2 C + Bcd + Ad^2) e}{\dots}$$

input

```
Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*(c - d*x^2)^(3/2)*(e + f*x^2)),x]
```

output

```
(Sqrt[b/a]*d*(c^2*C + B*c*d + A*d^2)*e*f*x*(a + b*x^2) - I*b*c*(c^2*C + B*c*d + A*d^2)*e*f*Sqrt[1 + (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], -((a*d)/(b*c))] + I*c*C*(b*c + a*d)*e*(d*e + c*f)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt[b/a]*x], -((a*d)/(b*c))] - I*c*d*(b*c + a*d)*(C*e^2 + f*(-(B*e) + A*f))*Sqrt[1 + (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticPi[(a*f)/(b*e), I*ArcSinh[Sqrt[b/a]*x], -((a*d)/(b*c))]/(Sqrt[b/a]*c*d*(b*c + a*d)*e*f*(d*e + c*f)*Sqrt[a + b*x^2]*Sqrt[c - d*x^2])
```

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 956 vs. 2(426) = 852.

Time = 2.06 (sec) , antiderivative size = 956, normalized size of antiderivative = 2.24, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.044$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx$$

↓ 7276

$$\int \left(\frac{Af^2 - Bef + Ce^2}{f^2\sqrt{a+bx^2}(c-dx^2)^{3/2}(e+fx^2)} - \frac{Ce - Bf}{f^2\sqrt{a+bx^2}(c-dx^2)^{3/2}} + \frac{Cx^2}{f\sqrt{a+bx^2}(c-dx^2)^{3/2}} \right) dx$$

↓ 2009

$$\frac{(Ce^2 - Bfe + Af^2) x\sqrt{bx^2 + ad^2}}{c(bc + ad)f^2(de + cf)\sqrt{c - dx^2}} - \frac{(Ce^2 - Bfe + Af^2) \sqrt{bx^2 + a}\sqrt{1 - \frac{dx^2}{c}} E\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \mid -\frac{bc}{ad}\right) d^{3/2}}{\sqrt{c}(bc + ad)f^2(de + cf)\sqrt{\frac{bx^2}{a} + 1}\sqrt{c - dx^2}}$$

$$+ \frac{(Ce - Bf)x\sqrt{bx^2 + ad}}{c(bc + ad)f^2\sqrt{c - dx^2}} + \frac{(Ce - Bf)\sqrt{bx^2 + a}\sqrt{1 - \frac{dx^2}{c}} E\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \mid -\frac{bc}{ad}\right) \sqrt{d}}{\sqrt{c}(bc + ad)f^2\sqrt{\frac{bx^2}{a} + 1}\sqrt{c - dx^2}}$$

$$- \frac{(de + 2cf)(Ce^2 - Bfe + Af^2) \sqrt{\frac{bx^2}{a} + 1}\sqrt{1 - \frac{dx^2}{c}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), -\frac{bc}{ad}\right) \sqrt{d}}{\sqrt{c}f^2(de + cf)^2\sqrt{bx^2 + a}\sqrt{c - dx^2}}$$

$$+ \frac{\sqrt{c}(Ce^2 - Bfe + Af^2) \sqrt{\frac{bx^2}{a} + 1}\sqrt{1 - \frac{dx^2}{c}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), -\frac{bc}{ad}\right) \sqrt{d}}{f(de + cf)^2\sqrt{bx^2 + a}\sqrt{c - dx^2}}$$

$$- \frac{Cx\sqrt{bx^2 + a}}{(bc + ad)f\sqrt{c - dx^2}} - \frac{\sqrt{c}C\sqrt{bx^2 + a}\sqrt{1 - \frac{dx^2}{c}} E\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \mid -\frac{bc}{ad}\right)}{(bc + ad)f\sqrt{\frac{bx^2}{a} + 1}\sqrt{c - dx^2}\sqrt{d}}$$

$$+ \frac{(Ce - Bf)\sqrt{\frac{bx^2}{a} + 1}\sqrt{1 - \frac{dx^2}{c}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), -\frac{bc}{ad}\right)}{\sqrt{c}f^2\sqrt{bx^2 + a}\sqrt{c - dx^2}\sqrt{d}}$$

$$+ \frac{\sqrt{c}(Ce^2 - Bfe + Af^2) \sqrt{\frac{bx^2}{a} + 1}\sqrt{1 - \frac{dx^2}{c}} \text{EllipticPi}\left(-\frac{cf}{de}, \arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), -\frac{bc}{ad}\right)}{ef(de + cf)\sqrt{bx^2 + a}\sqrt{c - dx^2}\sqrt{d}}$$

input

```
Int[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*(c - d*x^2)^(3/2)*(e + f*x^2)),x]
```

output

```
(C*x*Sqrt[a + b*x^2])/((b*c + a*d)*f*Sqrt[c - d*x^2]) - (d*(C*e - B*f)*x*S
qrt[a + b*x^2])/(c*(b*c + a*d)*f^2*Sqrt[c - d*x^2]) + (d^2*(C*e^2 - B*e*f
+ A*f^2)*x*Sqrt[a + b*x^2])/(c*(b*c + a*d)*f^2*(d*e + c*f)*Sqrt[c - d*x^2]
) - (Sqrt[c]*C*Sqrt[a + b*x^2]*Sqrt[1 - (d*x^2)/c]*EllipticE[ArcSin[(Sqrt[
d]*x)/Sqrt[c]], -((b*c)/(a*d))])/(Sqrt[d]*(b*c + a*d)*f*Sqrt[1 + (b*x^2)/a
]*Sqrt[c - d*x^2]) + (Sqrt[d]*(C*e - B*f)*Sqrt[a + b*x^2]*Sqrt[1 - (d*x^2)
/c]*EllipticE[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -((b*c)/(a*d))])/(Sqrt[c]*(b*c
+ a*d)*f^2*Sqrt[1 + (b*x^2)/a]*Sqrt[c - d*x^2]) - (d^(3/2)*(C*e^2 - B*e*f
+ A*f^2)*Sqrt[a + b*x^2]*Sqrt[1 - (d*x^2)/c]*EllipticE[ArcSin[(Sqrt[d]*x)/
Sqrt[c]], -((b*c)/(a*d))])/(Sqrt[c]*(b*c + a*d)*f^2*(d*e + c*f)*Sqrt[1 + (
b*x^2)/a]*Sqrt[c - d*x^2]) - ((C*e - B*f)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 - (d*
x^2)/c]*EllipticF[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -((b*c)/(a*d))])/(Sqrt[c]*S
qrt[d]*f^2*Sqrt[a + b*x^2]*Sqrt[c - d*x^2]) - (Sqrt[c]*Sqrt[d]*(C*e^2 - B*
e*f + A*f^2)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticF[ArcSin[(Sqr
t[d]*x)/Sqrt[c]], -((b*c)/(a*d))])/(f*(d*e + c*f)^2*Sqrt[a + b*x^2]*Sqrt[c
- d*x^2]) + (Sqrt[d]*(d*e + 2*c*f)*(C*e^2 - B*e*f + A*f^2)*Sqrt[1 + (b*x^
2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticF[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -((b*c)/(
a*d))])/(Sqrt[c]*f^2*(d*e + c*f)^2*Sqrt[a + b*x^2]*Sqrt[c - d*x^2]) + (Sqr
t[c]*(C*e^2 - B*e*f + A*f^2)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*Ellip
ticPi[-((c*f)/(d*e)), ArcSin[(Sqrt[d]*x)/Sqrt[c]], -((b*c)/(a*d))])/(Sq...
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 7276

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1126 vs. $2(377) = 754$.

Time = 8.45 (sec) , antiderivative size = 1127, normalized size of antiderivative = 2.65

method	result	size
default	Expression too large to display	1127
elliptic	Expression too large to display	1183

input `int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(3/2)/(f*x^2+e),x,method=_RETURVERBOSE)`

output
$$\begin{aligned} & (A*(1/c*d)^{(1/2)}*b*d^2*e*f*x^3+B*(1/c*d)^{(1/2)}*b*c*d*e*f*x^3+C*(1/c*d)^{(1/2)} \\ & *b*c^2*e*f*x^3+A*((-d*x^2+c)/c)^{(1/2)}*((b*x^2+a)/a)^{(1/2)}*EllipticF(x*(1/c*d)^{(1/2)}, \\ & (-b*c/a/d)^{(1/2)})*a*d^2*e*f+A*((-d*x^2+c)/c)^{(1/2)}*((b*x^2+a)/a)^{(1/2)}*EllipticF(x*(1/c*d)^{(1/2)}, \\ & (-b*c/a/d)^{(1/2)})*b*c*d*e*f-A*((-d*x^2+c)/c)^{(1/2)}*((b*x^2+a)/a)^{(1/2)}*EllipticE(x*(1/c*d)^{(1/2)}, \\ & (-b*c/a/d)^{(1/2)})*a*d^2*e*f+A*((-d*x^2+c)/c)^{(1/2)}*((b*x^2+a)/a)^{(1/2)}*EllipticPi(x*(1/c*d)^{(1/2)}, \\ & -c*f/d/e, (-b/a)^{(1/2)}/(1/c*d)^{(1/2)})*a*c*d*f^2+A*((-d*x^2+c)/c)^{(1/2)}*((b*x^2+a)/a)^{(1/2)}*EllipticPi(x*(1/c*d)^{(1/2)}, \\ & -c*f/d/e, (-b/a)^{(1/2)}/(1/c*d)^{(1/2)})*b*c^2*f^2+B*((-d*x^2+c)/c)^{(1/2)}*((b*x^2+a)/a)^{(1/2)}*EllipticF(x*(1/c*d)^{(1/2)}, \\ & (-b*c/a/d)^{(1/2)})*a*c*d*e*f+B*((-d*x^2+c)/c)^{(1/2)}*((b*x^2+a)/a)^{(1/2)}*EllipticF(x*(1/c*d)^{(1/2)}, \\ & (-b*c/a/d)^{(1/2)})*b*c^2*e*f-B*((-d*x^2+c)/c)^{(1/2)}*((b*x^2+a)/a)^{(1/2)}*EllipticE(x*(1/c*d)^{(1/2)}, \\ & (-b*c/a/d)^{(1/2)})*a*c*d*e*f-B*((-d*x^2+c)/c)^{(1/2)}*((b*x^2+a)/a)^{(1/2)}*EllipticPi(x*(1/c*d)^{(1/2)}, \\ & -c*f/d/e, (-b/a)^{(1/2)}/(1/c*d)^{(1/2)})*a*c*d*e*f-B*((-d*x^2+c)/c)^{(1/2)}*((b*x^2+a)/a)^{(1/2)}*EllipticPi(x*(1/c*d)^{(1/2)}, \\ & -c*f/d/e, (-b/a)^{(1/2)}/(1/c*d)^{(1/2)})*b*c^2*e*f-C*((-d*x^2+c)/c)^{(1/2)}*((b*x^2+a)/a)^{(1/2)}* \\ & EllipticF(x*(1/c*d)^{(1/2)}, (-b*c/a/d)^{(1/2)})*a*c*d*e^2-C*((-d*x^2+c)/c)^{(1/2)}*((b*x^2+a)/a)^{(1/2)}*EllipticF(x*(1/c*d)^{(1/2)}, \\ & (-b*c/a/d)^{(1/2)})*b*c^2*e^2-C*((-d*x^2+c)/c)^{(1/2)}*((b*x^2+a)/a)^{(1/2)}*EllipticE(x*(1/c*d)^{(1/2)}, \\ & (-b*c/a/d)^{(1/2)})*a*c^2*e*f+C*((-d*x^2+c)/c)^{(1/2)}*((b*x^2+a)/a)^{(1/2)}*El... \end{aligned}$$

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \text{Timed out}$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(3/2)/(f*x^2+e),x,algorithm="fricas")`

output Timed out

Sympy [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c - dx^2)^{\frac{3}{2}} (e + fx^2)} dx$$

input `integrate((C*x**4+B*x**2+A)/(b*x**2+a)**(1/2)/(-d*x**2+c)**(3/2)/(f*x**2+e),x)`

output `Integral((A + B*x**2 + C*x**4)/(sqrt(a + b*x**2)*(c - d*x**2)**(3/2)*(e + f*x**2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a} (-dx^2 + c)^{\frac{3}{2}} (fx^2 + e)} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(b*x^2 + a)*(-d*x^2 + c)^(3/2)*(f*x^2 + e)), x)`

Giac [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a} (-dx^2 + c)^{\frac{3}{2}} (fx^2 + e)} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(b*x^2 + a)*(-d*x^2 + c)^(3/2)*(f*x^2 + e)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a} (c - dx^2)^{3/2} (fx^2 + e)} dx$$

input `int((A + B*x^2 + C*x^4)/((a + b*x^2)^(1/2)*(c - d*x^2)^(3/2)*(e + f*x^2)), x)`

output `int((A + B*x^2 + C*x^4)/((a + b*x^2)^(1/2)*(c - d*x^2)^(3/2)*(e + f*x^2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \left(\int \frac{\sqrt{-dx^2 + c}}{b d^2 f x^8 + a d^2 f x^6 - 2 b c d f x^6 + b d^2 e x^6 - 2 a c d f x^4 + a d^2 e x^4} dx \right) + \left(\int \frac{\sqrt{-dx^2 + c} \sqrt{bx^2 + a}}{b d^2 f x^8 + a d^2 f x^6 - 2 b c d f x^6 + b d^2 e x^6 - 2 a c d f x^4 + a d^2 e x^4 + b c^2 f x^4 - 2 b c d e x^4 + a c^2 f x^2 - 2 a c d e} dx \right) + \left(\int \frac{\sqrt{-dx^2 + c} \sqrt{bx^2 + a}}{b d^2 f x^8 + a d^2 f x^6 - 2 b c d f x^6 + b d^2 e x^6 - 2 a c d f x^4 + a d^2 e x^4 + b c^2 f x^4 - 2 b c d e x^4 + a c^2 f x^2 - 2 a c d e} dx \right)$$

input `int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(3/2)/(f*x^2+e),x)`

output `int((sqrt(c - d*x**2)*sqrt(a + b*x**2)*x**4)/(a*c**2*e + a*c**2*f*x**2 - 2*a*c*d*e*x**2 - 2*a*c*d*f*x**4 + a*d**2*e*x**4 + a*d**2*f*x**6 + b*c**2*e*x**2 + b*c**2*f*x**4 - 2*b*c*d*e*x**4 - 2*b*c*d*f*x**6 + b*d**2*e*x**6 + b*d**2*f*x**8),x)*c + int((sqrt(c - d*x**2)*sqrt(a + b*x**2)*x**2)/(a*c**2*e + a*c**2*f*x**2 - 2*a*c*d*e*x**2 - 2*a*c*d*f*x**4 + a*d**2*e*x**4 + a*d**2*f*x**6 + b*c**2*e*x**2 + b*c**2*f*x**4 - 2*b*c*d*e*x**4 - 2*b*c*d*f*x**6 + b*d**2*e*x**6 + b*d**2*f*x**8),x)*b + int((sqrt(c - d*x**2)*sqrt(a + b*x**2))/(a*c**2*e + a*c**2*f*x**2 - 2*a*c*d*e*x**2 - 2*a*c*d*f*x**4 + a*d**2*e*x**4 + a*d**2*f*x**6 + b*c**2*e*x**2 + b*c**2*f*x**4 - 2*b*c*d*e*x**4 - 2*b*c*d*f*x**6 + b*d**2*e*x**6 + b*d**2*f*x**8),x)*a`

3.28
$$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}(c+dx^2)^{3/2}(e+fx^2)} dx$$

Optimal result	239
Mathematica [C] (verified)	240
Rubi [B] (verified)	240
Maple [B] (verified)	242
Fricas [F(-1)]	243
Sympy [F]	244
Maxima [F]	244
Giac [F]	244
Mupad [F(-1)]	245
Reduce [F]	245

Optimal result

Integrand size = 45, antiderivative size = 416

$$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}(c+dx^2)^{3/2}(e+fx^2)} dx = \frac{(c^2C - Bcd + Ad^2)x\sqrt{a-bx^2}}{c(bc+ad)(de-cf)\sqrt{c+dx^2}} + \frac{\sqrt{a}\sqrt{b}(c^2C - Bcd + Ad^2)\sqrt{1-\frac{bx^2}{a}}\sqrt{c+dx^2}E\left(\arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|-\frac{ad}{bc}\right)}{cd(bc+ad)(de-cf)\sqrt{a-bx^2}\sqrt{1+\frac{dx^2}{c}}} + \frac{\sqrt{a}C\sqrt{1-\frac{bx^2}{a}}\sqrt{1+\frac{dx^2}{c}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right),-\frac{ad}{bc}\right)}{\sqrt{bdf}\sqrt{a-bx^2}\sqrt{c+dx^2}} - \frac{\sqrt{a}(Ce^2 - Bef + Af^2)\sqrt{1-\frac{bx^2}{a}}\sqrt{1+\frac{dx^2}{c}}\text{EllipticPi}\left(-\frac{af}{be},\arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right),-\frac{ad}{bc}\right)}{\sqrt{bef}(de-cf)\sqrt{a-bx^2}\sqrt{c+dx^2}}$$

output

```
(A*d^2-B*c*d+C*c^2)*x*(-b*x^2+a)^(1/2)/c/(a*d+b*c)/(-c*f+d*e)/(d*x^2+c)^(1/2)+a^(1/2)*b^(1/2)*(A*d^2-B*c*d+C*c^2)*(1-b*x^2/a)^(1/2)*(d*x^2+c)^(1/2)*EllipticE(b^(1/2)*x/a^(1/2),(-a*d/b/c)^(1/2))/c/d/(a*d+b*c)/(-c*f+d*e)/(-b*x^2+a)^(1/2)/(1+d*x^2/c)^(1/2)+a^(1/2)*C*(1-b*x^2/a)^(1/2)*(1+d*x^2/c)^(1/2)*EllipticF(b^(1/2)*x/a^(1/2),(-a*d/b/c)^(1/2))/b^(1/2)/d/f/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)-a^(1/2)*(A*f^2-B*e*f+C*e^2)*(1-b*x^2/a)^(1/2)*(1+d*x^2/c)^(1/2)*EllipticPi(b^(1/2)*x/a^(1/2),-a*f/b/e,(-a*d/b/c)^(1/2))/b^(1/2)/e/f/(-c*f+d*e)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)
```


Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 13.35 (sec) , antiderivative size = 362, normalized size of antiderivative = 0.87

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \frac{-\sqrt{-\frac{b}{a}}d(c^2C - Bcd + Ad^2)efx(a - bx^2) + ibc(c^2C - Bcd + Ad^2)}{\dots}$$

input

```
Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a - b*x^2]*(c + d*x^2)^(3/2)*(e + f*x^2)),x]
```

output

```
(- (Sqrt[-(b/a)]*d*(c^2*C - B*c*d + A*d^2)*e*f*x*(a - b*x^2)) + I*b*c*(c^2*C - B*c*d + A*d^2)*e*f*Sqrt[1 - (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[-(b/a)]*x], -((a*d)/(b*c))] - I*c*C*(b*c + a*d)*e*(-(d*e) + c*f)*Sqrt[1 - (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt[-(b/a)]*x], -((a*d)/(b*c))] - I*c*d*(b*c + a*d)*(C*e^2 + f*(-(B*e) + A*f))*Sqrt[1 - (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticPi[-((a*f)/(b*e)), I*ArcSinh[Sqrt[-(b/a)]*x], -((a*d)/(b*c))])/(Sqrt[-(b/a)]*c*d*(b*c + a*d)*e*f*(-(d*e) + c*f)*Sqrt[a - b*x^2]*Sqrt[c + d*x^2])
```

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 844 vs. 2(416) = 832.

Time = 1.87 (sec) , antiderivative size = 844, normalized size of antiderivative = 2.03, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.044$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx$$

↓ 7276

$$\int \left(\frac{Af^2 - Bef + Ce^2}{f^2\sqrt{a - bx^2}(c + dx^2)^{3/2}(e + fx^2)} - \frac{Ce - Bf}{f^2\sqrt{a - bx^2}(c + dx^2)^{3/2}} + \frac{Cx^2}{f\sqrt{a - bx^2}(c + dx^2)^{3/2}} \right) dx$$

↓ 2009

$$\frac{\frac{(Ce^2 - Bfe + Af^2)x\sqrt{a - bx^2}d^2}{c(bc + ad)f^2(de - cf)\sqrt{dx^2 + c}} + \frac{\sqrt{a}\sqrt{b}(Ce^2 - Bfe + Af^2)\sqrt{1 - \frac{bx^2}{a}}\sqrt{dx^2 + c}E\left(\arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)d}{c(bc + ad)f^2(de - cf)\sqrt{a - bx^2}\sqrt{\frac{dx^2}{c} + 1}}}{\frac{\sqrt{a}(Ce^2 - Bfe + Af^2)\sqrt{1 - \frac{bx^2}{a}}\sqrt{\frac{dx^2}{c} + 1}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right), -\frac{ad}{bc}\right)d}{\sqrt{b}f(de - cf)^2\sqrt{a - bx^2}\sqrt{dx^2 + c}}} - \frac{(Ce - Bf)x\sqrt{a - bx^2}d}{c(bc + ad)f^2\sqrt{dx^2 + c}} - \frac{\sqrt{a}\sqrt{b}(Ce - Bf)\sqrt{1 - \frac{bx^2}{a}}\sqrt{dx^2 + c}E\left(\arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{c(bc + ad)f^2\sqrt{a - bx^2}\sqrt{\frac{dx^2}{c} + 1}} + \frac{\frac{Cx\sqrt{a - bx^2}}{(bc + ad)f\sqrt{dx^2 + c}}}{c^{3/2}(Ce^2 - Bfe + Af^2)\sqrt{a - bx^2}\text{EllipticPi}\left(1 - \frac{cf}{de}, \arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), \frac{bc}{ad} + 1\right)} - \frac{\frac{ae(de - cf)^2\sqrt{\frac{c(a - bx^2)}{a(dx^2 + c)}}\sqrt{dx^2 + c}\sqrt{d}}{\sqrt{a}\sqrt{b}C\sqrt{1 - \frac{bx^2}{a}}\sqrt{dx^2 + c}E\left(\arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)} + \frac{(bc + ad)f\sqrt{a - bx^2}\sqrt{\frac{dx^2}{c} + 1}d}{\sqrt{a}C\sqrt{1 - \frac{bx^2}{a}}\sqrt{\frac{dx^2}{c} + 1}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right), -\frac{ad}{bc}\right)}{\sqrt{b}f\sqrt{a - bx^2}\sqrt{dx^2 + cd}}$$

input `Int[(A + B*x^2 + C*x^4)/(Sqrt[a - b*x^2]*(c + d*x^2)^(3/2)*(e + f*x^2)),x]`

output

```

-((C*x*Sqrt[a - b*x^2])/((b*c + a*d)*f*Sqrt[c + d*x^2])) - (d*(C*e - B*f)*
x*Sqrt[a - b*x^2])/(c*(b*c + a*d)*f^2*Sqrt[c + d*x^2]) + (d^2*(C*e^2 - B*e
*f + A*f^2)*x*Sqrt[a - b*x^2])/(c*(b*c + a*d)*f^2*(d*e - c*f)*Sqrt[c + d*x
^2]) - (Sqrt[a]*Sqrt[b]*C*Sqrt[1 - (b*x^2)/a]*Sqrt[c + d*x^2]*EllipticE[Ar
cSin[(Sqrt[b]*x)/Sqrt[a]], -((a*d)/(b*c))])/(d*(b*c + a*d)*f*Sqrt[a - b*x^
2]*Sqrt[1 + (d*x^2)/c]) - (Sqrt[a]*Sqrt[b]*(C*e - B*f)*Sqrt[1 - (b*x^2)/a]
*Sqrt[c + d*x^2]*EllipticE[ArcSin[(Sqrt[b]*x)/Sqrt[a]], -((a*d)/(b*c))])/(
c*(b*c + a*d)*f^2*Sqrt[a - b*x^2]*Sqrt[1 + (d*x^2)/c]) + (Sqrt[a]*Sqrt[b]*
d*(C*e^2 - B*e*f + A*f^2)*Sqrt[1 - (b*x^2)/a]*Sqrt[c + d*x^2]*EllipticE[Ar
cSin[(Sqrt[b]*x)/Sqrt[a]], -((a*d)/(b*c))])/(c*(b*c + a*d)*f^2*(d*e - c*f)
*Sqrt[a - b*x^2]*Sqrt[1 + (d*x^2)/c]) + (Sqrt[a]*C*Sqrt[1 - (b*x^2)/a]*Sqr
t[1 + (d*x^2)/c]*EllipticF[ArcSin[(Sqrt[b]*x)/Sqrt[a]], -((a*d)/(b*c))])/(
Sqrt[b]*d*f*Sqrt[a - b*x^2]*Sqrt[c + d*x^2]) - (Sqrt[a]*d*(C*e^2 - B*e*f +
A*f^2)*Sqrt[1 - (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[ArcSin[(Sqrt[b]*
x)/Sqrt[a]], -((a*d)/(b*c))])/(Sqrt[b]*f*(d*e - c*f)^2*Sqrt[a - b*x^2]*Sqr
t[c + d*x^2]) + (c^(3/2)*(C*e^2 - B*e*f + A*f^2)*Sqrt[a - b*x^2]*EllipticP
i[1 - (c*f)/(d*e), ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 + (b*c)/(a*d)]/(a*Sqrt[
d]*e*(d*e - c*f)^2*Sqrt[(c*(a - b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2])

```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 7276

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1042 vs. $2(367) = 734$.

Time = 8.46 (sec) , antiderivative size = 1043, normalized size of antiderivative = 2.51

method	result	size
default	Expression too large to display	1043
elliptic	Expression too large to display	1869

input `int((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(3/2)/(f*x^2+e),x,method=_RETURNNVERBOSE)`

output `(A*(b/a)^(1/2)*b*d^3*e*f*x^3-B*(b/a)^(1/2)*b*c*d^2*e*f*x^3+C*(b/a)^(1/2)*b*c^2*d*e*f*x^3-A*((-b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE(x*(b/a)^(1/2),(-a*d/b/c)^(1/2))*b*c*d^2*e*f+A*((-b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticPi(x*(b/a)^(1/2),-a*f/b/e,(-1/c*d)^(1/2)/(b/a)^(1/2))*a*c*d^2*f^2+A*((-b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticPi(x*(b/a)^(1/2),-a*f/b/e,(-1/c*d)^(1/2)/(b/a)^(1/2))*b*c^2*d*f^2+B*((-b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE(x*(b/a)^(1/2),(-a*d/b/c)^(1/2))*b*c^2*d*e*f-B*((-b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticPi(x*(b/a)^(1/2),-a*f/b/e,(-1/c*d)^(1/2)/(b/a)^(1/2))*a*c*d^2*e*f-B*((-b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticPi(x*(b/a)^(1/2),-a*f/b/e,(-1/c*d)^(1/2)/(b/a)^(1/2))*b*c^2*d*e*f+C*((-b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF(x*(b/a)^(1/2),(-a*d/b/c)^(1/2))*a*c^2*d*e*f-C*((-b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF(x*(b/a)^(1/2),(-a*d/b/c)^(1/2))*a*c*d^2*e^2+C*((-b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF(x*(b/a)^(1/2),(-a*d/b/c)^(1/2))*b*c^3*e*f-C*((-b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF(x*(b/a)^(1/2),(-a*d/b/c)^(1/2))*b*c^2*d*e^2-C*((-b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE(x*(b/a)^(1/2),(-a*d/b/c)^(1/2))*b*c^3*e*f+C*((-b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticPi(x*(b/a)^(1/2),-a*f/b/e,(-1/c*d)^(1/2)/(b/a)^(1/2))*a*c*d^2*e^2+C*((-b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticPi(x*(b/a)^(1/2),-a*f/b/e,(-1/c*d)^(1/2)/(b/a)^(1/2))*b*c^2*d*e^2-A*(b/a)^(1/...`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \text{Timed out}$$

input `integrate((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(3/2)/(f*x^2+e),x,algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2} (c + dx^2)^{\frac{3}{2}} (e + fx^2)} dx$$

input `integrate((C*x**4+B*x**2+A)/(-b*x**2+a)**(1/2)/(d*x**2+c)**(3/2)/(f*x**2+e),x)`

output `Integral((A + B*x**2 + C*x**4)/(sqrt(a - b*x**2)*(c + d*x**2)**(3/2)*(e + f*x**2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{-bx^2 + a}(dx^2 + c)^{\frac{3}{2}}(fx^2 + e)} dx$$

input `integrate((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(-b*x^2 + a)*(d*x^2 + c)^(3/2)*(f*x^2 + e)), x)`

Giac [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{-bx^2 + a}(dx^2 + c)^{\frac{3}{2}}(fx^2 + e)} dx$$

input `integrate((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(-b*x^2 + a)*(d*x^2 + c)^(3/2)*(f*x^2 + e)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{a - bx^2} (dx^2 + c)^{3/2} (fx^2 + e)} dx$$

input `int((A + B*x^2 + C*x^4)/((a - b*x^2)^(1/2)*(c + d*x^2)^(3/2)*(e + f*x^2)), x)`

output `int((A + B*x^2 + C*x^4)/((a - b*x^2)^(1/2)*(c + d*x^2)^(3/2)*(e + f*x^2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2} (c + dx^2)^{3/2} (e + fx^2)} dx = \left(\int \frac{\sqrt{dx^2 + c}}{-bd^2fx^8 + ad^2fx^6 - 2bcdfx^6 - bd^2ex^6 + 2acdfx^4 + ad^2ex^4 - bc^2fx^4 - 2bcdex^4 + ac^2fx^2 + 2acde} dx \right) + \left(\int \frac{\sqrt{dx^2 + c} \sqrt{-bx^2 + a}}{-bd^2fx^8 + ad^2fx^6 - 2bcdfx^6 - bd^2ex^6 + 2acdfx^4 + ad^2ex^4 - bc^2fx^4 - 2bcdex^4 + ac^2fx^2 + 2acde} dx \right) + \left(\int \frac{\sqrt{dx^2 + c} \sqrt{-bx^2 + a}}{-bd^2fx^8 + ad^2fx^6 - 2bcdfx^6 - bd^2ex^6 + 2acdfx^4 + ad^2ex^4 - bc^2fx^4 - 2bcdex^4 + ac^2fx^2 + 2acde} dx \right)$$

input `int((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(3/2)/(f*x^2+e), x)`

output

```

int((sqrt(c + d*x**2)*sqrt(a - b*x**2)*x**4)/(a*c**2*e + a*c**2*f*x**2 + 2
*a*c*d*e*x**2 + 2*a*c*d*f*x**4 + a*d**2*e*x**4 + a*d**2*f*x**6 - b*c**2*e
*x**2 - b*c**2*f*x**4 - 2*b*c*d*e*x**4 - 2*b*c*d*f*x**6 - b*d**2*e*x**6 - b
*d**2*f*x**8),x)*c + int((sqrt(c + d*x**2)*sqrt(a - b*x**2)*x**2)/(a*c**2*
e + a*c**2*f*x**2 + 2*a*c*d*e*x**2 + 2*a*c*d*f*x**4 + a*d**2*e*x**4 + a*d*
*2*f*x**6 - b*c**2*e*x**2 - b*c**2*f*x**4 - 2*b*c*d*e*x**4 - 2*b*c*d*f*x**
6 - b*d**2*e*x**6 - b*d**2*f*x**8),x)*b + int((sqrt(c + d*x**2)*sqrt(a - b
*x**2))/(a*c**2*e + a*c**2*f*x**2 + 2*a*c*d*e*x**2 + 2*a*c*d*f*x**4 + a*d*
*2*e*x**4 + a*d**2*f*x**6 - b*c**2*e*x**2 - b*c**2*f*x**4 - 2*b*c*d*e*x**4
- 2*b*c*d*f*x**6 - b*d**2*e*x**6 - b*d**2*f*x**8),x)*a

```

3.29
$$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}(c-dx^2)^{3/2}(e+fx^2)} dx$$

Optimal result	247
Mathematica [C] (verified)	248
Rubi [B] (verified)	248
Maple [B] (verified)	250
Fricas [F(-1)]	251
Sympy [F]	252
Maxima [F]	252
Giac [F]	252
Mupad [F(-1)]	253
Reduce [F]	253

Optimal result

Integrand size = 46, antiderivative size = 418

$$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}(c-dx^2)^{3/2}(e+fx^2)} dx = -\frac{(c^2C+Bcd+Ad^2)x\sqrt{a-bx^2}}{c(bc-ad)(de+cf)\sqrt{c-dx^2}}$$

$$+\frac{\sqrt{a}\sqrt{b}(c^2C+Bcd+Ad^2)\sqrt{1-\frac{bx^2}{a}}\sqrt{c-dx^2}E\left(\arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|\frac{ad}{bc}\right)}{cd(bc-ad)(de+cf)\sqrt{a-bx^2}\sqrt{1-\frac{dx^2}{c}}}$$

$$-\frac{\sqrt{a}C\sqrt{1-\frac{bx^2}{a}}\sqrt{1-\frac{dx^2}{c}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right),\frac{ad}{bc}\right)}{\sqrt{bdf}\sqrt{a-bx^2}\sqrt{c-dx^2}}$$

$$+\frac{\sqrt{a}(Ce^2-Bef+Af^2)\sqrt{1-\frac{bx^2}{a}}\sqrt{1-\frac{dx^2}{c}}\text{EllipticPi}\left(-\frac{af}{be},\arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right),\frac{ad}{bc}\right)}{\sqrt{bef}(de+cf)\sqrt{a-bx^2}\sqrt{c-dx^2}}$$

output

```
-(A*d^2+B*c*d+C*c^2)*x*(-b*x^2+a)^(1/2)/c/(-a*d+b*c)/(c*f+d*e)/(-d*x^2+c)^(1/2)+a^(1/2)*b^(1/2)*(A*d^2+B*c*d+C*c^2)*(1-b*x^2/a)^(1/2)*(-d*x^2+c)^(1/2)*EllipticE(b^(1/2)*x/a^(1/2),(a*d/b/c)^(1/2))/c/d/(-a*d+b*c)/(c*f+d*e)/(-b*x^2+a)^(1/2)/(1-d*x^2/c)^(1/2)-a^(1/2)*C*(1-b*x^2/a)^(1/2)*(1-d*x^2/c)^(1/2)*EllipticF(b^(1/2)*x/a^(1/2),(a*d/b/c)^(1/2))/b^(1/2)/d/f/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)+a^(1/2)*(A*f^2-B*e*f+C*e^2)*(1-b*x^2/a)^(1/2)*(1-d*x^2/c)^(1/2)*EllipticPi(b^(1/2)*x/a^(1/2),-a*f/b/e,(a*d/b/c)^(1/2))/b^(1/2)/e/f/(c*f+d*e)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)
```


Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 11.11 (sec) , antiderivative size = 361, normalized size of antiderivative = 0.86

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \frac{\sqrt{-\frac{b}{a}} d (c^2 C + Bcd + Ad^2) \operatorname{efx}(a - bx^2) + ibc(c^2 C + Bcd + Ad^2)}{\dots}$$

input

```
Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a - b*x^2]*(c - d*x^2)^(3/2)*(e + f*x^2)),x]
```

output

```
(Sqrt[-(b/a)]*d*(c^2*C + B*c*d + A*d^2)*e*f*x*(a - b*x^2) + I*b*c*(c^2*C + B*c*d + A*d^2)*e*f*Sqrt[1 - (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[-(b/a)]*x], (a*d)/(b*c)] + I*c*C*(-(b*c) + a*d)*e*(d*e + c*f)*Sqrt[1 - (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt[-(b/a)]*x], (a*d)/(b*c)] - I*c*d*(-(b*c) + a*d)*(C*e^2 + f*(-(B*e) + A*f))*Sqrt[1 - (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticPi[-((a*f)/(b*e)), I*ArcSinh[Sqrt[-(b/a)]*x], (a*d)/(b*c)])/(Sqrt[-(b/a)]*c*d*(-(b*c) + a*d)*e*f*(d*e + c*f)*Sqrt[a - b*x^2]*Sqrt[c - d*x^2])
```

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 875 vs. $2(418) = 836$.

Time = 2.01 (sec) , antiderivative size = 875, normalized size of antiderivative = 2.09, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx$$

↓ 7276

$$\begin{aligned}
& \int \left(\frac{Af^2 - Bef + Ce^2}{f^2\sqrt{a - bx^2}(c - dx^2)^{3/2}(e + fx^2)} - \frac{Ce - Bf}{f^2\sqrt{a - bx^2}(c - dx^2)^{3/2}} + \frac{Cx^2}{f\sqrt{a - bx^2}(c - dx^2)^{3/2}} \right) dx \\
& \quad \downarrow \text{2009} \\
& \quad - \frac{(Ce^2 - Bfe + Af^2)x\sqrt{a - bx^2}d^2}{c(bc - ad)f^2(de + cf)\sqrt{c - dx^2}} + \\
& \quad \frac{(Ce^2 - Bfe + Af^2)\sqrt{a - bx^2}\sqrt{1 - \frac{dx^2}{c}}E\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \middle| \frac{bc}{ad}\right)d^{3/2}}{\sqrt{c}(bc - ad)f^2(de + cf)\sqrt{1 - \frac{bx^2}{a}}\sqrt{c - dx^2}} + \frac{(Ce - Bf)x\sqrt{a - bx^2}d}{c(bc - ad)f^2\sqrt{c - dx^2}} + \\
& \quad \frac{(de + 2cf)(Ce^2 - Bfe + Af^2)\sqrt{1 - \frac{bx^2}{a}}\sqrt{1 - \frac{dx^2}{c}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), \frac{bc}{ad}\right)\sqrt{d}}{\sqrt{cf^2}(de + cf)^2\sqrt{a - bx^2}\sqrt{c - dx^2}} - \\
& \quad \frac{\sqrt{c}(Ce^2 - Bfe + Af^2)\sqrt{1 - \frac{bx^2}{a}}\sqrt{1 - \frac{dx^2}{c}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), \frac{bc}{ad}\right)\sqrt{d}}{f(de + cf)^2\sqrt{a - bx^2}\sqrt{c - dx^2}} - \\
& \quad \frac{\sqrt{a}\sqrt{b}(Ce - Bf)\sqrt{1 - \frac{bx^2}{a}}\sqrt{c - dx^2}E\left(\arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) \middle| \frac{ad}{bc}\right)}{c(bc - ad)f^2\sqrt{a - bx^2}\sqrt{1 - \frac{dx^2}{c}}} + \\
& \quad \frac{\sqrt{a}(Ce^2 - Bfe + Af^2)\sqrt{1 - \frac{bx^2}{a}}\sqrt{1 - \frac{dx^2}{c}}\text{EllipticPi}\left(-\frac{af}{be}, \arcsin\left(\frac{\sqrt{bx}}{\sqrt{a}}\right), \frac{ad}{bc}\right)}{\sqrt{bef}(de + cf)\sqrt{a - bx^2}\sqrt{c - dx^2}} - \\
& \quad \frac{Cx\sqrt{a - bx^2}}{(bc - ad)f\sqrt{c - dx^2}} + \frac{\sqrt{c}C\sqrt{a - bx^2}\sqrt{1 - \frac{dx^2}{c}}E\left(\arcsin\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \middle| \frac{bc}{ad}\right)}{(bc - ad)f\sqrt{1 - \frac{bx^2}{a}}\sqrt{c - dx^2}\sqrt{d}}
\end{aligned}$$

input

```
Int[(A + B*x^2 + C*x^4)/(Sqrt[a - b*x^2]*(c - d*x^2)^(3/2)*(e + f*x^2)),x]
```

output

```

-((C*x*Sqrt[a - b*x^2])/((b*c - a*d)*f*Sqrt[c - d*x^2])) + (d*(C*e - B*f)*
x*Sqrt[a - b*x^2])/(c*(b*c - a*d)*f^2*Sqrt[c - d*x^2]) - (d^2*(C*e^2 - B*e
*f + A*f^2)*x*Sqrt[a - b*x^2])/(c*(b*c - a*d)*f^2*(d*e + c*f)*Sqrt[c - d*x
^2]) - (Sqrt[a]*Sqrt[b]*(C*e - B*f)*Sqrt[1 - (b*x^2)/a]*Sqrt[c - d*x^2]*El
lipticE[ArcSin[(Sqrt[b]*x)/Sqrt[a]], (a*d)/(b*c)]/(c*(b*c - a*d)*f^2*Sqrt
[a - b*x^2]*Sqrt[1 - (d*x^2)/c]) + (Sqrt[c]*C*Sqrt[a - b*x^2]*Sqrt[1 - (d*
x^2)/c]*EllipticE[ArcSin[(Sqrt[d]*x)/Sqrt[c]], (b*c)/(a*d)]/(Sqrt[d]*(b*c
- a*d)*f*Sqrt[1 - (b*x^2)/a]*Sqrt[c - d*x^2]) + (d^(3/2)*(C*e^2 - B*e*f +
A*f^2)*Sqrt[a - b*x^2]*Sqrt[1 - (d*x^2)/c]*EllipticE[ArcSin[(Sqrt[d]*x)/S
qrt[c]], (b*c)/(a*d)]/(Sqrt[c]*(b*c - a*d)*f^2*(d*e + c*f)*Sqrt[1 - (b*x^
2)/a]*Sqrt[c - d*x^2]) - (Sqrt[c]*Sqrt[d]*(C*e^2 - B*e*f + A*f^2)*Sqrt[1 -
(b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticF[ArcSin[(Sqrt[d]*x)/Sqrt[c]], (b*
c)/(a*d)]/(f*(d*e + c*f)^2*Sqrt[a - b*x^2]*Sqrt[c - d*x^2]) + (Sqrt[d]*(d
*e + 2*c*f)*(C*e^2 - B*e*f + A*f^2)*Sqrt[1 - (b*x^2)/a]*Sqrt[1 - (d*x^2)/c
]*EllipticF[ArcSin[(Sqrt[d]*x)/Sqrt[c]], (b*c)/(a*d)]/(Sqrt[c]*f^2*(d*e +
c*f)^2*Sqrt[a - b*x^2]*Sqrt[c - d*x^2]) + (Sqrt[a]*(C*e^2 - B*e*f + A*f^2
)*Sqrt[1 - (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticPi[-((a*f)/(b*e)), ArcSi
n[(Sqrt[b]*x)/Sqrt[a]], (a*d)/(b*c)]/(Sqrt[b]*e*f*(d*e + c*f)*Sqrt[a - b*
x^2]*Sqrt[c - d*x^2])

```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 7276

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1133 vs. $2(369) = 738$.

Time = 8.43 (sec) , antiderivative size = 1134, normalized size of antiderivative = 2.71

method	result	size
default	Expression too large to display	1134
elliptic	Expression too large to display	1201

input `int((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(3/2)/(f*x^2+e),x,method=_RETURNVERBOSE)`

output `(-A*(1/c*d)^(1/2)*b*d^2*e*f*x^3-B*(1/c*d)^(1/2)*b*c*d*e*f*x^3-C*(1/c*d)^(1/2)*b*c^2*e*f*x^3+A*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*EllipticF(x*(1/c*d)^(1/2),(b*c/a/d)^(1/2))*a*d^2*e*f-A*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*EllipticF(x*(1/c*d)^(1/2),(b*c/a/d)^(1/2))*b*c*d*e*f-A*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*EllipticE(x*(1/c*d)^(1/2),(b*c/a/d)^(1/2))*a*d^2*e*f+A*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*EllipticPi(x*(1/c*d)^(1/2),(b*c/a/d)^(1/2),-c*f/d/e,(b/a)^(1/2)/(1/c*d)^(1/2))*a*c*d*f^2-A*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*EllipticPi(x*(1/c*d)^(1/2),-c*f/d/e,(b/a)^(1/2)/(1/c*d)^(1/2))*b*c^2*f^2+B*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*EllipticF(x*(1/c*d)^(1/2),(b*c/a/d)^(1/2))*a*c*d*e*f-B*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*EllipticF(x*(1/c*d)^(1/2),(b*c/a/d)^(1/2))*b*c^2*e*f-B*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*EllipticE(x*(1/c*d)^(1/2),(b*c/a/d)^(1/2))*a*c*d*e*f-B*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*EllipticPi(x*(1/c*d)^(1/2),-c*f/d/e,(b/a)^(1/2)/(1/c*d)^(1/2))*a*c*d*e*f+B*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*EllipticPi(x*(1/c*d)^(1/2),-c*f/d/e,(b/a)^(1/2)/(1/c*d)^(1/2))*b*c^2*e*f-C*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*EllipticF(x*(1/c*d)^(1/2),(b*c/a/d)^(1/2))*a*c*d*e^2+C*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*EllipticF(x*(1/c*d)^(1/2),(b*c/a/d)^(1/2))*b*c^2*e^2-C*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*EllipticE(x*(1/c*d)^(1/2),(b*c/a/d)^(1/2))*a*c^2*e*f+C*((-d*x^2+c)/c)^(1/2)*((-b*x^2+a)/a)^(1/2)*...`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \text{Timed out}$$

input `integrate((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(3/2)/(f*x^2+e),x,algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2} (c - dx^2)^{\frac{3}{2}} (e + fx^2)} dx$$

input `integrate((C*x**4+B*x**2+A)/(-b*x**2+a)**(1/2)/(-d*x**2+c)**(3/2)/(f*x**2+e),x)`

output `Integral((A + B*x**2 + C*x**4)/(sqrt(a - b*x**2)*(c - d*x**2)**(3/2)*(e + f*x**2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{-bx^2 + a} (-dx^2 + c)^{\frac{3}{2}} (fx^2 + e)} dx$$

input `integrate((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(-b*x^2 + a)*(-d*x^2 + c)^(3/2)*(f*x^2 + e)), x)`

Giac [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{-bx^2 + a} (-dx^2 + c)^{\frac{3}{2}} (fx^2 + e)} dx$$

input `integrate((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(3/2)/(f*x^2+e),x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(-b*x^2 + a)*(-d*x^2 + c)^(3/2)*(f*x^2 + e)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{a - bx^2} (c - dx^2)^{3/2} (fx^2 + e)} dx$$

input `int((A + B*x^2 + C*x^4)/((a - b*x^2)^(1/2)*(c - d*x^2)^(3/2)*(e + f*x^2)), x)`

output `int((A + B*x^2 + C*x^4)/((a - b*x^2)^(1/2)*(c - d*x^2)^(3/2)*(e + f*x^2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2} (c - dx^2)^{3/2} (e + fx^2)} dx = \left(\int \frac{\sqrt{-dx^2 + c}}{-bd^2fx^8 + ad^2fx^6 + 2bcdfx^6 - bd^2ex^6 - 2acdfx^4 + ad^2ex^4 - bc^2fx^4 + 2bcdex^4 + ac^2fx^2 - 2bcde} \right) \\ + \left(\int \frac{\sqrt{-dx^2 + c}\sqrt{-bx^2 + a}x^2}{-bd^2fx^8 + ad^2fx^6 + 2bcdfx^6 - bd^2ex^6 - 2acdfx^4 + ad^2ex^4 - bc^2fx^4 + 2bcdex^4 + ac^2fx^2 - 2bcde} \right) \\ + \left(\int \frac{\sqrt{-dx^2 + c}\sqrt{-bx^2 + a}}{-bd^2fx^8 + ad^2fx^6 + 2bcdfx^6 - bd^2ex^6 - 2acdfx^4 + ad^2ex^4 - bc^2fx^4 + 2bcdex^4 + ac^2fx^2 - 2bcde} \right)$$

input `int((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(3/2)/(f*x^2+e), x)`

output

```
int((sqrt(c - d*x**2)*sqrt(a - b*x**2)*x**4)/(a*c**2*e + a*c**2*f*x**2 - 2
*a*c*d*e*x**2 - 2*a*c*d*f*x**4 + a*d**2*e*x**4 + a*d**2*f*x**6 - b*c**2*e
*x**2 - b*c**2*f*x**4 + 2*b*c*d*e*x**4 + 2*b*c*d*f*x**6 - b*d**2*e*x**6 - b
*d**2*f*x**8),x)*c + int((sqrt(c - d*x**2)*sqrt(a - b*x**2)*x**2)/(a*c**2*
e + a*c**2*f*x**2 - 2*a*c*d*e*x**2 - 2*a*c*d*f*x**4 + a*d**2*e*x**4 + a*d*
*2*f*x**6 - b*c**2*e*x**2 - b*c**2*f*x**4 + 2*b*c*d*e*x**4 + 2*b*c*d*f*x**
6 - b*d**2*e*x**6 - b*d**2*f*x**8),x)*b + int((sqrt(c - d*x**2)*sqrt(a - b
*x**2))/(a*c**2*e + a*c**2*f*x**2 - 2*a*c*d*e*x**2 - 2*a*c*d*f*x**4 + a*d*
*2*e*x**4 + a*d**2*f*x**6 - b*c**2*e*x**2 - b*c**2*f*x**4 + 2*b*c*d*e*x**4
+ 2*b*c*d*f*x**6 - b*d**2*e*x**6 - b*d**2*f*x**8),x)*a
```

3.30
$$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e+fx^2)^2} dx$$

Optimal result	255
Mathematica [C] (verified)	256
Rubi [A] (verified)	257
Maple [B] (verified)	259
Fricas [F(-1)]	260
Sympy [F]	261
Maxima [F]	261
Giac [F]	261
Mupad [F(-1)]	262
Reduce [F]	262

Optimal result

Integrand size = 44, antiderivative size = 545

$$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e+fx^2)^2} dx = -\frac{(Ce^2 - Bef + Af^2)x\sqrt{a+bx^2}}{2ef(be - af)\sqrt{c+dx^2}(e+fx^2)}$$

$$+ \frac{\sqrt{c}\sqrt{d}(Ce^2 - Bef + Af^2)\sqrt{a+bx^2}E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \mid 1 - \frac{bc}{ad}\right)}{2ef(be - af)(de - cf)\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}}$$

$$+ \frac{\sqrt{c}(2c^2Cef + 2Ad^2ef - cd(Ce^2 + f(Be + Af)))\sqrt{a+bx^2}\text{EllipticF}\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right)}{2a\sqrt{def}(de - cf)^2\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}}$$

$$c^{3/2}(af(Ce^2(2de - 3cf) + f^2(Bce - 2Ade + Acf)) - b(Ce^3(de - 2cf) + ef(Bde^2 - Af(3de - 2cf))))$$

$$2a\sqrt{de^2f}(be - af)(de - cf)^2\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}$$

output

```
-1/2*(A*f^2-B*e*f+C*e^2)*x*(b*x^2+a)^(1/2)/e/f/(-a*f+b*e)/(d*x^2+c)^(1/2)/
(f*x^2+e)+1/2*c^(1/2)*d^(1/2)*(A*f^2-B*e*f+C*e^2)*(b*x^2+a)^(1/2)*Elliptic
E(d^(1/2)*x/c^(1/2)/(1+d*x^2/c)^(1/2),(1-b*c/a/d)^(1/2))/e/f/(-a*f+b*e)/(-
c*f+d*e)/(c*(b*x^2+a)/a/(d*x^2+c))^(1/2)/(d*x^2+c)^(1/2)+1/2*c^(1/2)*(2*c^
2*C*e*f+2*A*d^2*e*f-c*d*(C*e^2+f*(A*f+B*e)))*(b*x^2+a)^(1/2)*InverseJacobi
AM(arctan(d^(1/2)*x/c^(1/2)),(1-b*c/a/d)^(1/2))/a/d^(1/2)/e/f/(-c*f+d*e)^2
/(c*(b*x^2+a)/a/(d*x^2+c))^(1/2)/(d*x^2+c)^(1/2)-1/2*c^(3/2)*(a*f*(C*e^2*(
-3*c*f+2*d*e)+f^2*(A*c*f-2*A*d*e+B*c*e))-b*(C*e^3*(-2*c*f+d*e)+e*f*(B*d*e^
2-A*f*(-2*c*f+3*d*e)))*(b*x^2+a)^(1/2)*EllipticPi(d^(1/2)*x/c^(1/2)/(1+d*
x^2/c)^(1/2),1-c*f/d/e,(1-b*c/a/d)^(1/2))/a/d^(1/2)/e^2/f/(-a*f+b*e)/(-c*f
+d*e)^2/(c*(b*x^2+a)/a/(d*x^2+c))^(1/2)/(d*x^2+c)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 14.67 (sec) , antiderivative size = 2038, normalized size of antiderivative = 3.74

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^2} dx = \text{Result too large to show}$$

input

```
Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(e + f*x^2)
^2),x]
```

output

```
(a*Sqrt[b/a]*c*C*e^3*f^2*x - a*Sqrt[b/a]*B*c*e^2*f^3*x + a*A*Sqrt[b/a]*c*e
*f^4*x + b*Sqrt[b/a]*c*C*e^3*f^2*x^3 + a*Sqrt[b/a]*C*d*e^3*f^2*x^3 - b*Sqr
t[b/a]*B*c*e^2*f^3*x^3 - a*Sqrt[b/a]*B*d*e^2*f^3*x^3 + A*b*Sqrt[b/a]*c*e*f
^4*x^3 + a*A*Sqrt[b/a]*d*e*f^4*x^3 + b*Sqrt[b/a]*C*d*e^3*f^2*x^5 - b*Sqrt[
b/a]*B*d*e^2*f^3*x^5 + A*b*Sqrt[b/a]*d*e*f^4*x^5 + I*b*c*e*f*(C*e^2 + f*(-
(B*e) + A*f))*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*(e + f*x^2)*Elliptic
E[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - I*e*(-(d*e) + c*f)*(-(b*C*e^2) -
b*B*e*f + 2*a*C*e*f + A*b*f^2)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*(e
+ f*x^2)*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] + I*b*C*d*e^5*Sqrt
[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticPi[(a*f)/(b*e), I*ArcSinh[Sqrt
[b/a]*x], (a*d)/(b*c)] - (2*I)*b*c*C*e^4*f*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d
*x^2)/c]*EllipticPi[(a*f)/(b*e), I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] + I*
b*B*d*e^4*f*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticPi[(a*f)/(b*e)
, I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - (2*I)*a*C*d*e^4*f*Sqrt[1 + (b*x^2
)/a]*Sqrt[1 + (d*x^2)/c]*EllipticPi[(a*f)/(b*e), I*ArcSinh[Sqrt[b/a]*x], (
a*d)/(b*c)] + (3*I)*a*c*C*e^3*f^2*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*
EllipticPi[(a*f)/(b*e), I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - (3*I)*A*b*d
*e^3*f^2*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticPi[(a*f)/(b*e), I
*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] + (2*I)*A*b*c*e^2*f^3*Sqrt[1 + (b*x^2
)/a]*Sqrt[1 + (d*x^2)/c]*EllipticPi[(a*f)/(b*e), I*ArcSinh[Sqrt[b/a]*x],...
```

Rubi [A] (verified)

Time = 1.71 (sec) , antiderivative size = 777, normalized size of antiderivative = 1.43, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.045$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^2} dx$$

↓ 7293

$$\int \left(\frac{Af^2 - Bef + Ce^2}{f^2\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^2} + \frac{Bf - 2Ce}{f^2\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)} + \frac{C}{f^2\sqrt{a + bx^2}\sqrt{c + dx^2}} \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{\sqrt{-a}\sqrt{\frac{bx^2}{a}+1}\sqrt{\frac{dx^2}{c}+1}(Af^2 - Bef + Ce^2)(be(3de - 2cf) - af(2de - cf)) \operatorname{EllipticPi}\left(\frac{af}{be}, \arcsin\left(\frac{\sqrt{bx}}{\sqrt{-a}}\right), \frac{ad}{bc}\right)}{2\sqrt{be^2f^2}\sqrt{a+bx^2}\sqrt{c+dx^2}(be-af)(de-cf)} \\
& + \frac{b\sqrt{c}\sqrt{d}\sqrt{a+bx^2}(Af^2 - Bef + Ce^2) \operatorname{EllipticF}\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right)}{2af^2\sqrt{c+dx^2}(be-af)(de-cf)\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} \\
& - \frac{\sqrt{c}\sqrt{d}\sqrt{a+bx^2}(Af^2 - Bef + Ce^2) E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{2ef\sqrt{c+dx^2}(be-af)(de-cf)\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} \\
& + \frac{dx\sqrt{a+bx^2}(Af^2 - Bef + Ce^2)}{2ef\sqrt{c+dx^2}(be-af)(de-cf)} + \frac{x\sqrt{a+bx^2}\sqrt{c+dx^2}(Af^2 - Bef + Ce^2)}{2e(e+fx^2)(be-af)(de-cf)} \\
& + \frac{\sqrt{-a}\sqrt{\frac{bx^2}{a}+1}\sqrt{\frac{dx^2}{c}+1}(2Ce - Bf) \operatorname{EllipticPi}\left(\frac{af}{be}, \arcsin\left(\frac{\sqrt{bx}}{\sqrt{-a}}\right), \frac{ad}{bc}\right)}{\sqrt{be^2f^2}\sqrt{a+bx^2}\sqrt{c+dx^2}} \\
& + \frac{\sqrt{c}C\sqrt{a+bx^2} \operatorname{EllipticF}\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right)}{a\sqrt{d}f^2\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}
\end{aligned}$$

input

```
Int[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(e + f*x^2)^2),x]
```

output

```
-1/2*(d*(C*e^2 - B*e*f + A*f^2)*x*Sqrt[a + b*x^2])/(e*f*(b*e - a*f)*(d*e - c*f)*Sqrt[c + d*x^2]) + ((C*e^2 - B*e*f + A*f^2)*x*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])/(2*e*(b*e - a*f)*(d*e - c*f)*(e + f*x^2)) + (Sqrt[c]*Sqrt[d]*(C*e^2 - B*e*f + A*f^2)*Sqrt[a + b*x^2]*EllipticE[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)])/(2*e*f*(b*e - a*f)*(d*e - c*f)*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) + (Sqrt[c]*C*Sqrt[a + b*x^2]*EllipticF[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)])/(a*Sqrt[d]*f^2*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) - (b*Sqrt[c]*Sqrt[d]*(C*e^2 - B*e*f + A*f^2)*Sqrt[a + b*x^2]*EllipticF[ArcTan[(Sqrt[d]*x)/Sqrt[c]], 1 - (b*c)/(a*d)])/(2*a*f^2*(b*e - a*f)*(d*e - c*f)*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*Sqrt[c + d*x^2]) - (Sqrt[-a]*(2*C*e - B*f)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticPi[(a*f)/(b*e), ArcSin[(Sqrt[b]*x)/Sqrt[-a]], (a*d)/(b*c)])/(Sqrt[b]*e*f^2*Sqrt[a + b*x^2]*Sqrt[c + d*x^2]) + (Sqrt[-a]*(C*e^2 - B*e*f + A*f^2)*(b*e*(3*d*e - 2*c*f) - a*f*(2*d*e - c*f))*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticPi[(a*f)/(b*e), ArcSin[(Sqrt[b]*x)/Sqrt[-a]], (a*d)/(b*c)])/(2*Sqrt[b]*e^2*f^2*(b*e - a*f)*(d*e - c*f)*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])
```

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2572 vs. $2(519) = 1038$.

Time = 8.47 (sec) , antiderivative size = 2573, normalized size of antiderivative = 4.72

method	result	size
elliptic	Expression too large to display	2573
default	Expression too large to display	3121

input `int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^2,x,method=_
RETURNVERBOSE)`

output

```

((b*x^2+a)*(d*x^2+c)^(1/2)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)*(1/2/e/(a*c*f^
2-a*d*e*f-b*c*e*f+b*d*e^2))*x*(A*f^2-B*e*f+C*e^2)*(b*d*x^4+a*d*x^2+b*c*x^2+
a*c)^(1/2)/(f*x^2+e)+1/2/(a*c*f^2-a*d*e*f-b*c*e*f+b*d*e^2)*f/e/(-b/a)^(1/2
)*(1+b*x^2/a)^(1/2)*(1+d*x^2/c)^(1/2)/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)^(1/2)*
EllipticPi(x*(-b/a)^(1/2),a*f/b/e,(-1/c*d)^(1/2)/(-b/a)^(1/2))*B*a*c-1/2/(
a*c*f^2-a*d*e*f-b*c*e*f+b*d*e^2)*e/f/(-b/a)^(1/2)*(1+b*x^2/a)^(1/2)*(1+d*x
^2/c)^(1/2)/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)^(1/2)*EllipticPi(x*(-b/a)^(1/2),
a*f/b/e,(-1/c*d)^(1/2)/(-b/a)^(1/2))*B*b*d+1/(a*c*f^2-a*d*e*f-b*c*e*f+b*d*
e^2)*e/f/(-b/a)^(1/2)*(1+b*x^2/a)^(1/2)*(1+d*x^2/c)^(1/2)/(b*d*x^4+a*d*x^2
+b*c*x^2+a*c)^(1/2)*EllipticPi(x*(-b/a)^(1/2),a*f/b/e,(-1/c*d)^(1/2)/(-b/a
)^(1/2))*C*a*d+1/(a*c*f^2-a*d*e*f-b*c*e*f+b*d*e^2)*e/f/(-b/a)^(1/2)*(1+b*x
^2/a)^(1/2)*(1+d*x^2/c)^(1/2)/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)^(1/2)*Elliptic
Pi(x*(-b/a)^(1/2),a*f/b/e,(-1/c*d)^(1/2)/(-b/a)^(1/2))*C*b*c-1/2/(a*c*f^2-
a*d*e*f-b*c*e*f+b*d*e^2)*e^2/f^2/(-b/a)^(1/2)*(1+b*x^2/a)^(1/2)*(1+d*x^2/c
)^(1/2)/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)^(1/2)*EllipticPi(x*(-b/a)^(1/2),a*f/
b/e,(-1/c*d)^(1/2)/(-b/a)^(1/2))*C*b*d+1/2/(-b/a)^(1/2)*(1+b*x^2/a)^(1/2)*
(1+d*x^2/c)^(1/2)/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)^(1/2)*EllipticF(x*(-b/a)^(
1/2),(-1+(a*d+b*c)/c/b)^(1/2))*d*b/f/(a*c*f^2-a*d*e*f-b*c*e*f+b*d*e^2)*B*e
-1/2/(-b/a)^(1/2)*(1+b*x^2/a)^(1/2)*(1+d*x^2/c)^(1/2)/(b*d*x^4+a*d*x^2+b*c
*x^2+a*c)^(1/2)*EllipticF(x*(-b/a)^(1/2),(-1+(a*d+b*c)/c/b)^(1/2))*d*b/...

```

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^2} dx = \text{Timed out}$$

input

```

integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^2,x, a
lgorithm="fricas")

```

output

Timed out

Sympy [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^2} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^2} dx$$

input `integrate((C*x**4+B*x**2+A)/(b*x**2+a)**(1/2)/(d*x**2+c)**(1/2)/(f*x**2+e)**2,x)`

output `Integral((A + B*x**2 + C*x**4)/(sqrt(a + b*x**2)*sqrt(c + d*x**2)*(e + f*x**2)**2), x)`

Maxima [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^2} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(fx^2 + e)^2} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^2,x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)*(f*x^2 + e)^2), x)`

Giac [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^2} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(fx^2 + e)^2} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^2,x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)*(f*x^2 + e)^2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^2} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(fx^2 + e)^2} dx$$

input

```
int((A + B*x^2 + C*x^4)/((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(e + f*x^2)^2),x)
```

output

```
int((A + B*x^2 + C*x^4)/((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(e + f*x^2)^2), x)
```

Reduce [F]

$$\begin{aligned} & \int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^2} dx \\ &= \left(\int \frac{\sqrt{dx^2 + c}\sqrt{bx^2 + a}x^4}{bd f^2 x^8 + ad f^2 x^6 + bc f^2 x^6 + 2bdef x^6 + ac f^2 x^4 + 2adef x^4 + 2bcef x^4 + bde^2 x^4 + 2acef x^2 + ae^2} dx \right) \\ &+ \left(\int \frac{\sqrt{dx^2 + c}\sqrt{bx^2 + a}x^2}{bd f^2 x^8 + ad f^2 x^6 + bc f^2 x^6 + 2bdef x^6 + ac f^2 x^4 + 2adef x^4 + 2bcef x^4 + bde^2 x^4 + 2acef x^2 + ae^2} dx \right) \\ &+ \left(\int \frac{\sqrt{dx^2 + c}\sqrt{bx^2 + a}}{bd f^2 x^8 + ad f^2 x^6 + bc f^2 x^6 + 2bdef x^6 + ac f^2 x^4 + 2adef x^4 + 2bcef x^4 + bde^2 x^4 + 2acef x^2 + ae^2} dx \right) \end{aligned}$$

input

```
int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^2,x)
```

output

```

int((sqrt(c + d*x**2)*sqrt(a + b*x**2)*x**4)/(a*c**2 + 2*a*c*e*f*x**2 +
a*c*f**2*x**4 + a*d*e**2*x**2 + 2*a*d*e*f*x**4 + a*d*f**2*x**6 + b*c*e**2*
x**2 + 2*b*c*e*f*x**4 + b*c*f**2*x**6 + b*d*e**2*x**4 + 2*b*d*e*f*x**6 + b
*d*f**2*x**8),x)*c + int((sqrt(c + d*x**2)*sqrt(a + b*x**2)*x**2)/(a*c**
2 + 2*a*c*e*f*x**2 + a*c*f**2*x**4 + a*d*e**2*x**2 + 2*a*d*e*f*x**4 + a*d*
f**2*x**6 + b*c*e**2*x**2 + 2*b*c*e*f*x**4 + b*c*f**2*x**6 + b*d*e**2*x**4
+ 2*b*d*e*f*x**6 + b*d*f**2*x**8),x)*b + int((sqrt(c + d*x**2)*sqrt(a + b
*x**2))/(a*c**2 + 2*a*c*e*f*x**2 + a*c*f**2*x**4 + a*d*e**2*x**2 + 2*a*d
*e*f*x**4 + a*d*f**2*x**6 + b*c*e**2*x**2 + 2*b*c*e*f*x**4 + b*c*f**2*x**6
+ b*d*e**2*x**4 + 2*b*d*e*f*x**6 + b*d*f**2*x**8),x)*a

```


output

```

1/4*(A*f^2-B*e*f+C*e^2)*x*(b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)/e/(-a*f+b*e)/(-c
*f+d*e)/(f*x^2+e)^2-1/8*(b*(C*e^3*(2*c*f+d*e)+e*f*(3*A*f*(-2*c*f+3*d*e)-B*
e*(-2*c*f+5*d*e)))+a*f*(C*e^2*(-5*c*f+2*d*e)-f*(3*A*f*(-c*f+2*d*e)-B*e*(c*
f+2*d*e))))*x*(b*x^2+a)^(1/2)/e^2/f/(-a*f+b*e)^2/(-c*f+d*e)/(d*x^2+c)^(1/2
)/(f*x^2+e)+1/8*c^(1/2)*d^(1/2)*(b*(C*e^3*(2*c*f+d*e)+e*f*(3*A*f*(-2*c*f+3
*d*e)-B*e*(-2*c*f+5*d*e)))+a*f*(C*e^2*(-5*c*f+2*d*e)-f*(3*A*f*(-c*f+2*d*e)
-B*e*(c*f+2*d*e))))*(b*x^2+a)^(1/2)*EllipticE(d^(1/2)*x/c^(1/2)/(1+d*x^2/c
)^(1/2),(1-b*c/a/d)^(1/2))/e^2/f/(-a*f+b*e)^2/(-c*f+d*e)^2/(c*(b*x^2+a)/a/
(d*x^2+c))^(1/2)/(d*x^2+c)^(1/2)-1/8*c^(1/2)*d^(1/2)*(a*f^2*(8*A*d^2*e^2-4
*c*d*e*(2*A*f+B*e)+c^2*(3*A*f^2+B*e*f+3*C*e^2))-b*e*(8*A*d^2*e^2*f-c*d*e*(
9*A*f^2+3*B*e*f+C*e^2)+4*c^2*(A*f^3+C*e^2*f)))*(b*x^2+a)^(1/2)*InverseJaco
biAM(arctan(d^(1/2)*x/c^(1/2)),(1-b*c/a/d)^(1/2))/a/e^2/f/(-a*f+b*e)/(-c*f
+d*e)^3/(c*(b*x^2+a)/a/(d*x^2+c))^(1/2)/(d*x^2+c)^(1/2)-1/8*c^(3/2)*(a^2*f
^4*(8*A*d^2*e^2-4*c*d*e*(2*A*f+B*e)+c^2*(3*A*f^2+B*e*f+3*C*e^2))+2*a*b*e*f
*(C*d*e^3*(-5*c*f+2*d*e)+f^2*(B*c*e*(-2*c*f+5*d*e)-A*(4*c^2*f^2-11*c*d*e*f
+10*d^2*e^2)))-b^2*(C*d*e^5*(-4*c*f+d*e)+e^2*f*(3*B*d^2*e^3-A*f*(8*c^2*f^2
-20*c*d*e*f+15*d^2*e^2)))*(b*x^2+a)^(1/2)*EllipticPi(d^(1/2)*x/c^(1/2)/(1
+d*x^2/c)^(1/2),1-c*f/d/e,(1-b*c/a/d)^(1/2))/a/d^(1/2)/e^3/f/(-a*f+b*e)^2/
(-c*f+d*e)^3/(c*(b*x^2+a)/a/(d*x^2+c))^(1/2)/(d*x^2+c)^(1/2)

```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 15.17 (sec) , antiderivative size = 732, normalized size of antiderivative = 0.76

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^3} dx$$

$$= \frac{\sqrt{\frac{b}{a}} e f^2 x (a + bx^2) (c + dx^2) (2e(be - af)(de - cf) (Ce^2 + f(-Be + Af)) + (af(Ce^2(2de - 5cf) + f(3$$

input

```

Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(e + f*x^2)
^3),x]

```

output

```
(Sqrt[b/a]*e*f^2*x*(a + b*x^2)*(c + d*x^2)*(2*e*(b*e - a*f)*(d*e - c*f)*(C
*e^2 + f*(-(B*e) + A*f)) + (a*f*(C*e^2*(2*d*e - 5*c*f) + f*(3*A*f*(-2*d*e
+ c*f) + B*e*(2*d*e + c*f))) + b*(C*e^3*(d*e + 2*c*f) + e*f*(3*A*f*(3*d*e
- 2*c*f) + B*e*(-5*d*e + 2*c*f))))*(e + f*x^2)) - I*Sqrt[1 + (b*x^2)/a]*Sq
rt[1 + (d*x^2)/c]*(e + f*x^2)^2*(-(b*c*e*f*(a*f*(C*e^2*(2*d*e - 5*c*f) + f
*(3*A*f*(-2*d*e + c*f) + B*e*(2*d*e + c*f))) + b*(C*e^3*(d*e + 2*c*f) + e
*f*(3*A*f*(3*d*e - 2*c*f) + B*e*(-5*d*e + 2*c*f))))*EllipticE[I*ArcSinh[Sqr
t[b/a]*x], (a*d)/(b*c)]) + b*e*(d*e - c*f)*(-(a*f*(C*e^2*(4*d*e - 5*c*f) +
f^2*(B*c*e - 4*A*d*e + 3*A*c*f))) + b*(C*e^3*(d*e - 2*c*f) + e*f*(B*e*(3*
d*e - 2*c*f) + A*f*(-7*d*e + 6*c*f))))*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (
a*d)/(b*c)] + (a^2*f^4*(8*A*d^2*e^2 - 4*c*d*e*(B*e + 2*A*f) + c^2*(3*C*e^2
+ B*e*f + 3*A*f^2)) + 2*a*b*e*f*(C*d*e^3*(2*d*e - 5*c*f) + f^2*(B*c*e*(5*
d*e - 2*c*f) + A*(-10*d^2*e^2 + 11*c*d*e*f - 4*c^2*f^2))) + b^2*(C*d*e^5*(
-(d*e) + 4*c*f) + e^2*f*(-3*B*d^2*e^3 + A*f*(15*d^2*e^2 - 20*c*d*e*f + 8*c
^2*f^2))))*EllipticPi[(a*f)/(b*e), I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)])))/
(8*Sqrt[b/a]*e^3*f^2*(b*e - a*f)^2*(d*e - c*f)^2*Sqrt[a + b*x^2]*Sqrt[c +
d*x^2]*(e + f*x^2)^2)
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^3} dx$$

↓ 7293

$$\int \left(\frac{Af^2 - Bef + Ce^2}{f^2\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^3} + \frac{Bf - 2Ce}{f^2\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^2} + \frac{C}{f^2\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)} \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{d(2Ce - Bf)\sqrt{bx^2 + ax}}{2ef(be - af)(de - cf)\sqrt{dx^2 + c}} - \frac{(2Ce - Bf)\sqrt{bx^2 + a}\sqrt{dx^2 + cx}}{2e(be - af)(de - cf)(fx^2 + e)} - \\
& \frac{\sqrt{c}\sqrt{d}(2Ce - Bf)\sqrt{bx^2 + a}E\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \mid 1 - \frac{bc}{ad}\right)}{2ef(be - af)(de - cf)\sqrt{\frac{c(bx^2 + a)}{a(dx^2 + c)}}\sqrt{dx^2 + c}} + \\
& \frac{b\sqrt{c}\sqrt{d}(2Ce - Bf)\sqrt{bx^2 + a}\operatorname{EllipticF}\left(\arctan\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right)}{2af^2(be - af)(de - cf)\sqrt{\frac{c(bx^2 + a)}{a(dx^2 + c)}}\sqrt{dx^2 + c}} + \\
& \frac{3\sqrt{-a}(Ce^2 - Bfe + Af^2)\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{dx^2}{c} + 1}\operatorname{EllipticPi}\left(\frac{af}{be}, \arcsin\left(\frac{\sqrt{bx}}{\sqrt{-a}}\right), \frac{ad}{bc}\right)}{8\sqrt{be^3}f^2\sqrt{bx^2 + a}\sqrt{dx^2 + c}} - \\
& \frac{\sqrt{-a}(2Ce - Bf)(be(3de - 2cf) - af(2de - cf))\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{dx^2}{c} + 1}\operatorname{EllipticPi}\left(\frac{af}{be}, \arcsin\left(\frac{\sqrt{bx}}{\sqrt{-a}}\right), \frac{ad}{bc}\right)}{2\sqrt{be^2}f^2(be - af)(de - cf)\sqrt{bx^2 + a}\sqrt{dx^2 + c}} + \\
& \frac{\sqrt{-a}C\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{dx^2}{c} + 1}\operatorname{EllipticPi}\left(\frac{af}{be}, \arcsin\left(\frac{\sqrt{bx}}{\sqrt{-a}}\right), \frac{ad}{bc}\right)}{\sqrt{bef^2}\sqrt{bx^2 + a}\sqrt{dx^2 + c}} - \\
& \frac{(Ce^2 - Bfe + Af^2)\int\frac{1}{(\sqrt{-e}\sqrt{f} - fx)^3\sqrt{bx^2 + a}\sqrt{dx^2 + c}}dx}{8(-e)^{3/2}\sqrt{f}} - \\
& \frac{\frac{3}{16}\left(\frac{C}{f} - \frac{Be - Af}{e^2}\right)\int\frac{1}{(\sqrt{-e}\sqrt{f} - fx)^2\sqrt{bx^2 + a}\sqrt{dx^2 + c}}dx}{(Ce^2 - Bfe + Af^2)\int\frac{1}{(fx + \sqrt{-e}\sqrt{f})^3\sqrt{bx^2 + a}\sqrt{dx^2 + c}}dx} - \\
& \frac{\frac{3}{16}\left(\frac{C}{f} - \frac{Be - Af}{e^2}\right)\int\frac{1}{(fx + \sqrt{-e}\sqrt{f})^2\sqrt{bx^2 + a}\sqrt{dx^2 + c}}dx}{8(-e)^{3/2}\sqrt{f}}
\end{aligned}$$

input `Int[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(e + f*x^2)^3),x]`

output `$Aborted`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 6399 vs. $2(935) = 1870$.

Time = 10.91 (sec) , antiderivative size = 6400, normalized size of antiderivative = 6.62

method	result	size
elliptic	Expression too large to display	6400
default	Expression too large to display	11835

input

```
int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^3,x,method=_
RETURNVERBOSE)
```

output

```
result too large to display
```

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^3} dx = \text{Timed out}$$

input

```
integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^3,x, a
lgorithm="fricas")
```

output

```
Timed out
```

Sympy [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^3} dx = \text{Timed out}$$

input

```
integrate((C*x**4+B*x**2+A)/(b*x**2+a)**(1/2)/(d*x**2+c)**(1/2)/(f*x**2+e)
**3,x)
```

output Timed out

Maxima [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^3} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(fx^2 + e)^3} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^3,x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)*(f*x^2 + e)^3), x)`

Giac [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^3} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(fx^2 + e)^3} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^3,x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)*(f*x^2 + e)^3), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^3} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(fx^2 + e)^3} dx$$

input

```
int((A + B*x^2 + C*x^4)/((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(e + f*x^2)^3),x)
```

output

```
int((A + B*x^2 + C*x^4)/((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(e + f*x^2)^3), x)
```

Reduce [F]

$$\begin{aligned} & \int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^3} dx \\ &= \left(\int \frac{\sqrt{dx^2 + c}\sqrt{bx^2 + a}}{bd f^3 x^{10} + ad f^3 x^8 + bc f^3 x^8 + 3bde f^2 x^8 + ac f^3 x^6 + 3ade f^2 x^6 + 3bce f^2 x^6 + 3bd e^2 f x^6 + 3ace f^2 x^6} dx \right. \\ & \quad + \left(\int \frac{\sqrt{dx^2 + c}\sqrt{bx^2 + a}}{bd f^3 x^{10} + ad f^3 x^8 + bc f^3 x^8 + 3bde f^2 x^8 + ac f^3 x^6 + 3ade f^2 x^6 + 3bce f^2 x^6 + 3bd e^2 f x^6 + 3ace f^2 x^6} dx \right. \\ & \quad \left. + \left(\int \frac{\sqrt{dx^2 + c}\sqrt{bx^2 + a}}{bd f^3 x^{10} + ad f^3 x^8 + bc f^3 x^8 + 3bde f^2 x^8 + ac f^3 x^6 + 3ade f^2 x^6 + 3bce f^2 x^6 + 3bd e^2 f x^6 + 3ace f^2 x^6} dx \right) \right) \end{aligned}$$

input

```
int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^3,x)
```

output

```

int((sqrt(c + d*x**2)*sqrt(a + b*x**2)*x**4)/(a*c**3 + 3*a*c**2*f*x**2
+ 3*a*c*e*f**2*x**4 + a*c*f**3*x**6 + a*d*e**3*x**2 + 3*a*d*e**2*f*x**4 +
3*a*d*e*f**2*x**6 + a*d*f**3*x**8 + b*c*e**3*x**2 + 3*b*c*e**2*f*x**4 + 3
*b*c*e*f**2*x**6 + b*c*f**3*x**8 + b*d*e**3*x**4 + 3*b*d*e**2*f*x**6 + 3*b
*d*e*f**2*x**8 + b*d*f**3*x**10),x)*c + int((sqrt(c + d*x**2)*sqrt(a + b*x
**2)*x**2)/(a*c**3 + 3*a*c**2*f*x**2 + 3*a*c*e*f**2*x**4 + a*c*f**3*x*
*6 + a*d*e**3*x**2 + 3*a*d*e**2*f*x**4 + 3*a*d*e*f**2*x**6 + a*d*f**3*x**8
+ b*c*e**3*x**2 + 3*b*c*e**2*f*x**4 + 3*b*c*e*f**2*x**6 + b*c*f**3*x**8 +
b*d*e**3*x**4 + 3*b*d*e**2*f*x**6 + 3*b*d*e*f**2*x**8 + b*d*f**3*x**10),x
)*b + int((sqrt(c + d*x**2)*sqrt(a + b*x**2))/(a*c**3 + 3*a*c**2*f*x**
2 + 3*a*c*e*f**2*x**4 + a*c*f**3*x**6 + a*d*e**3*x**2 + 3*a*d*e**2*f*x**4
+ 3*a*d*e*f**2*x**6 + a*d*f**3*x**8 + b*c*e**3*x**2 + 3*b*c*e**2*f*x**4 +
3*b*c*e*f**2*x**6 + b*c*f**3*x**8 + b*d*e**3*x**4 + 3*b*d*e**2*f*x**6 + 3*
b*d*e*f**2*x**8 + b*d*f**3*x**10),x)*a

```


3.32
$$\int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{(e+fx^2)^{3/2}} dx$$

Optimal result	272
Mathematica [F]	273
Rubi [F]	274
Maple [F]	275
Fricas [F(-1)]	275
Sympy [F]	276
Maxima [F]	276
Giac [F]	276
Mupad [F(-1)]	277
Reduce [F]	277

Optimal result

Integrand size = 46, antiderivative size = 813

$$\int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{(e+fx^2)^{3/2}} dx = \frac{(aCdf - b(5Cde - cCf - 4Bdf))x\sqrt{a+bx^2}\sqrt{c+dx^2}}{8bdf^2\sqrt{e+fx^2}} + \frac{Cx^3\sqrt{a+bx^2}\sqrt{c+dx^2}}{4f\sqrt{e+fx^2}} - \frac{c\sqrt{-be+af}(aCdef + 4bdf(3Be - 2Af) - bCe(15de - cf))\sqrt{a+bx^2}\sqrt{\frac{e(c+dx^2)}{c(e+fx^2)}}E\left(\arcsin\left(\frac{\sqrt{-be+afx}}{\sqrt{a}\sqrt{e+fx^2}}\right)\right)}{8\sqrt{abdef^3}\sqrt{c+dx^2}\sqrt{\frac{e(a+bx^2)}{a(e+fx^2)}}} + \frac{(a^2Cdef^2 + abf(Ce(6de - 13cf) - 4Bf(de - 2cf)) - b^2(3Ce^2(5de - 7cf) - 4f(Be(3de - 4cf) - 2Af(3de - 4cf))))\sqrt{a+bx^2}\sqrt{c+dx^2}}{8\sqrt{ab}f^4\sqrt{-be+af}\sqrt{c+dx^2}\sqrt{\frac{e(a+bx^2)}{a(e+fx^2)}}} + \frac{e(a^2Cd^2f^2 + 2abdf(3Cde - cCf - 2Bdf) + b^2(4df(3Bde - Bcf - 2Adf) - C(15d^2e^2 - 6cdef - c^2f^2))\sqrt{a+bx^2}\sqrt{c+dx^2}}{8\sqrt{abdf^4}\sqrt{-be+af}\sqrt{c+dx^2}\sqrt{\frac{e(a+bx^2)}{a(e+fx^2)}}}$$

output

```

1/8*(a*C*d*f-b*(-4*B*d*f-C*c*f+5*C*d*e))*x*(b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)
/b/d/f^2/(f*x^2+e)^(1/2)+1/4*C*x^3*(b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)/f/(f*x^
2+e)^(1/2)-1/8*c*(a*f-b*e)^(1/2)*(a*C*d*e*f+4*b*d*f*(-2*A*f+3*B*e)-b*C*e*(
-c*f+15*d*e))*(b*x^2+a)^(1/2)*(e*(d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticE((a
*f-b*e)^(1/2)*x/a^(1/2)/(f*x^2+e)^(1/2),(a*(-c*f+d*e)/c/(-a*f+b*e))^(1/2))
/a^(1/2)/b/d/e/f^3/(d*x^2+c)^(1/2)/(e*(b*x^2+a)/a/(f*x^2+e))^(1/2)+1/8*(a^
2*C*d*e*f^2+a*b*f*(C*e*(-13*c*f+6*d*e)-4*B*f*(-2*c*f+d*e))-b^2*(3*C*e^2*(-
7*c*f+5*d*e)-4*f*(B*e*(-4*c*f+3*d*e)-2*A*f*(-c*f+d*e))))*(b*x^2+a)^(1/2)*(
e*(d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticF((a*f-b*e)^(1/2)*x/a^(1/2)/(f*x^2+
e)^(1/2),(a*(-c*f+d*e)/c/(-a*f+b*e))^(1/2))/a^(1/2)/b/f^4/(a*f-b*e)^(1/2)/
(d*x^2+c)^(1/2)/(e*(b*x^2+a)/a/(f*x^2+e))^(1/2)-1/8*e*(a^2*C*d^2*f^2+2*a*b
*d*f*(-2*B*d*f-C*c*f+3*C*d*e)+b^2*(4*d*f*(-2*A*d*f-B*c*f+3*B*d*e)-C*(-c^2*
f^2-6*c*d*e*f+15*d^2*e^2)))*(b*x^2+a)^(1/2)*(e*(d*x^2+c)/c/(f*x^2+e))^(1/2
)*EllipticPi((a*f-b*e)^(1/2)*x/a^(1/2)/(f*x^2+e)^(1/2),-a*f/(-a*f+b*e),(a
(-c*f+d*e)/c/(-a*f+b*e))^(1/2))/a^(1/2)/b/d/f^4/(a*f-b*e)^(1/2)/(d*x^2+c)^(
1/2)/(e*(b*x^2+a)/a/(f*x^2+e))^(1/2)

```

Mathematica [F]

$$\int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{(e+fx^2)^{3/2}} dx = \int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{(e+fx^2)^{3/2}} dx$$

input

```

Integrate[(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(A + B*x^2 + C*x^4))/(e + f*x^2
)^(3/2), x]

```

output

```

Integrate[(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(A + B*x^2 + C*x^4))/(e + f*x^2
)^(3/2), x]

```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{(e+fx^2)^{3/2}} dx$$

↓ 7293

$$\int \left(\frac{A\sqrt{a+bx^2}\sqrt{c+dx^2}}{(e+fx^2)^{3/2}} + \frac{Bx^2\sqrt{a+bx^2}\sqrt{c+dx^2}}{(e+fx^2)^{3/2}} + \frac{Cx^4\sqrt{a+bx^2}\sqrt{c+dx^2}}{(e+fx^2)^{3/2}} \right) dx$$

↓ 2009

$$\frac{B \int \frac{x^2\sqrt{bx^2+a}\sqrt{dx^2+c}}{(fx^2+e)^{3/2}} dx + C \int \frac{x^4\sqrt{bx^2+a}\sqrt{dx^2+c}}{(fx^2+e)^{3/2}} dx + \frac{A\sqrt{c}\sqrt{a+bx^2}\sqrt{de-cf}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}} \operatorname{EllipticPi}\left(\frac{de}{de-cf}, \arcsin\left(\frac{\sqrt{de-cf}x}{\sqrt{e}\sqrt{dx^2+c}}\right), -\frac{(bc-ad)e}{a(de-cf)}\right)}{af\sqrt{e+fx^2}\sqrt{de-cf}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} - \frac{Ac^{3/2}\sqrt{a+bx^2}(be-af) \operatorname{EllipticF}\left(\arctan\left(\frac{\sqrt{de-cf}x}{\sqrt{c}\sqrt{fx^2+e}}\right), -\frac{(bc-ad)e}{a(de-cf)}\right)}{aef\sqrt{c+dx^2}\sqrt{de-cf}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{A\sqrt{c}\sqrt{a+bx^2}\sqrt{de-cf}E\left(\arctan\left(\frac{\sqrt{de-cf}x}{\sqrt{c}\sqrt{fx^2+e}}\right) \middle| -\frac{(bc-ad)e}{a(de-cf)}\right)}{ef\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} - \frac{Ax\sqrt{a+bx^2}(de-cf)}{ef\sqrt{c+dx^2}\sqrt{e+fx^2}}$$

input

```
Int[(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(A + B*x^2 + C*x^4))/(e + f*x^2)^(3/2), x]
```

output

```
$Aborted
```

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]`

Maple [F]

$$\int \frac{\sqrt{bx^2+a}\sqrt{x^2d+c}(Cx^4+x^2B+A)}{(fx^2+e)^{\frac{3}{2}}} dx$$

input `int((b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*(C*x^4+B*x^2+A)/(f*x^2+e)^(3/2),x)`

output `int((b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*(C*x^4+B*x^2+A)/(f*x^2+e)^(3/2),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{(e+fx^2)^{3/2}} dx = \text{Timed out}$$

input `integrate((b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*(C*x^4+B*x^2+A)/(f*x^2+e)^(3/2),
x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{(e+fx^2)^{3/2}} dx = \int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{(e+fx^2)^{\frac{3}{2}}} dx$$

input `integrate((b*x**2+a)**(1/2)*(d*x**2+c)**(1/2)*(C*x**4+B*x**2+A)/(f*x**2+e)**(3/2),x)`

output `Integral(sqrt(a + b*x**2)*sqrt(c + d*x**2)*(A + B*x**2 + C*x**4)/(e + f*x**2)**(3/2), x)`

Maxima [F]

$$\int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{(e+fx^2)^{3/2}} dx = \int \frac{(Cx^4+Bx^2+A)\sqrt{bx^2+a}\sqrt{dx^2+c}}{(fx^2+e)^{\frac{3}{2}}} dx$$

input `integrate((b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*(C*x^4+B*x^2+A)/(f*x^2+e)^(3/2), x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)*sqrt(b*x^2 + a)*sqrt(d*x^2 + c)/(f*x^2 + e)^(3/2), x)`

Giac [F]

$$\int \frac{\sqrt{a+bx^2}\sqrt{c+dx^2}(A+Bx^2+Cx^4)}{(e+fx^2)^{3/2}} dx = \int \frac{(Cx^4+Bx^2+A)\sqrt{bx^2+a}\sqrt{dx^2+c}}{(fx^2+e)^{\frac{3}{2}}} dx$$

input `integrate((b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*(C*x^4+B*x^2+A)/(f*x^2+e)^(3/2), x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)*sqrt(b*x^2 + a)*sqrt(d*x^2 + c)/(f*x^2 + e)^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + bx^2} \sqrt{c + dx^2} (A + Bx^2 + Cx^4)}{(e + fx^2)^{3/2}} dx = \int \frac{\sqrt{bx^2 + a} \sqrt{dx^2 + c} (Cx^4 + Bx^2 + A)}{(fx^2 + e)^{3/2}} dx$$

input `int(((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(A + B*x^2 + C*x^4))/(e + f*x^2)^(3/2), x)`

output `int(((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(A + B*x^2 + C*x^4))/(e + f*x^2)^(3/2), x)`

Reduce [F]

$$\int \frac{\sqrt{a + bx^2} \sqrt{c + dx^2} (A + Bx^2 + Cx^4)}{(e + fx^2)^{3/2}} dx = \int \frac{\sqrt{bx^2 + a} \sqrt{dx^2 + c} (Cx^4 + Bx^2 + A)}{(fx^2 + e)^{\frac{3}{2}}} dx$$

input `int((b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*(C*x^4+B*x^2+A)/(f*x^2+e)^(3/2), x)`

output `int((b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*(C*x^4+B*x^2+A)/(f*x^2+e)^(3/2), x)`

3.33
$$\int \frac{A+Bx^2}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e+fx^2)^{3/2}} dx$$

Optimal result	278
Mathematica [F]	279
Rubi [F]	279
Maple [F]	280
Fricas [F]	280
Sympy [F]	281
Maxima [F]	281
Giac [F]	281
Mupad [F(-1)]	282
Reduce [F]	282

Optimal result

Integrand size = 41, antiderivative size = 332

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \frac{c(Be - Af)\sqrt{a + bx^2}\sqrt{\frac{e(c+dx^2)}{c(e+fx^2)}}E\left(\arcsin\left(\frac{\sqrt{-be+afx}}{\sqrt{a}\sqrt{e+fx^2}}\right) \mid \frac{a(de-cf)}{c(be-af)}\right)}{\sqrt{ae}\sqrt{-be + af}(de - cf)\sqrt{c + dx^2}\sqrt{\frac{e(a+bx^2)}{a(e+fx^2)}}} - \frac{(Bc - Ad)\sqrt{a + bx^2}\sqrt{\frac{e(c+dx^2)}{c(e+fx^2)}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{-be+afx}}{\sqrt{a}\sqrt{e+fx^2}}\right), \frac{a(de-cf)}{c(be-af)}\right)}{\sqrt{a}\sqrt{-be + af}(de - cf)\sqrt{c + dx^2}\sqrt{\frac{e(a+bx^2)}{a(e+fx^2)}}}$$

output

```
c*(-A*f+B*e)*(b*x^2+a)^(1/2)*(e*(d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticE((a*f-b*e)^(1/2)*x/a^(1/2)/(f*x^2+e)^(1/2),(a*(-c*f+d*e)/c/(-a*f+b*e))^(1/2))/a^(1/2)/e/(a*f-b*e)^(1/2)/(-c*f+d*e)/(d*x^2+c)^(1/2)/(e*(b*x^2+a)/a/(f*x^2+e))^(1/2)-(-A*d+B*c)*(b*x^2+a)^(1/2)*(e*(d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticF((a*f-b*e)^(1/2)*x/a^(1/2)/(f*x^2+e)^(1/2),(a*(-c*f+d*e)/c/(-a*f+b*e))^(1/2))/a^(1/2)/(a*f-b*e)^(1/2)/(-c*f+d*e)/(d*x^2+c)^(1/2)/(e*(b*x^2+a)/a/(f*x^2+e))^(1/2)
```

Mathematica [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx$$

input

```
Integrate[(A + B*x^2)/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(e + f*x^2)^(3/2)), x]
```

output

```
Integrate[(A + B*x^2)/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(e + f*x^2)^(3/2)), x]
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx$$

↓ 7293

$$\int \left(\frac{A}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} + \frac{Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} \right) dx$$

↓ 2009

$$A \int \frac{1}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(fx^2 + e)^{3/2}} dx + B \int \frac{x^2}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(fx^2 + e)^{3/2}} dx$$

input

```
Int[(A + B*x^2)/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(e + f*x^2)^(3/2)), x]
```

output

```
$Aborted
```


Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]`

Maple [F]

$$\int \frac{x^2 B + A}{\sqrt{b x^2 + a} \sqrt{x^2 d + c} (f x^2 + e)^{\frac{3}{2}}} dx$$

input `int((B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)`

output `int((B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)`

Fricas [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} \sqrt{c + dx^2} (e + fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a} \sqrt{dx^2 + c} (fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, alg
orithm="fricas")`

output `integral((B*x^2 + A)*sqrt(b*x^2 + a)*sqrt(d*x^2 + c)*sqrt(f*x^2 + e)/(b*d*
f^2*x^8 + (2*b*d*e*f + (b*c + a*d)*f^2)*x^6 + (b*d*e^2 + a*c*f^2 + 2*(b*c
+ a*d)*e*f)*x^4 + a*c*e^2 + (2*a*c*e*f + (b*c + a*d)*e^2)*x^2), x)`

Sympy [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{\frac{3}{2}}} dx$$

input `integrate((B*x**2+A)/(b*x**2+a)**(1/2)/(d*x**2+c)**(1/2)/(f*x**2+e)**(3/2),x)`

output `Integral((A + B*x**2)/(sqrt(a + b*x**2)*sqrt(c + d*x**2)*(e + f*x**2)**(3/2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)*(f*x^2 + e)^(3/2)), x)`

Giac [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)*(f*x^2 + e)^(3/2)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(fx^2 + e)^{3/2}} dx$$

input `int((A + B*x^2)/((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(e + f*x^2)^(3/2)), x)`

output `int((A + B*x^2)/((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(e + f*x^2)^(3/2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{\sqrt{fx^2 + e}\sqrt{dx^2 + c}\sqrt{bx^2 + a}}{df^2x^6 + cf^2x^4 + 2defx^4 + 2cef x^2 + de^2x^2 + ce^2} dx$$

input `int((B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2), x)`

output `int((sqrt(e + f*x**2)*sqrt(c + d*x**2)*sqrt(a + b*x**2))/(c*e**2 + 2*c*e*f*x**2 + c*f**2*x**4 + d*e**2*x**2 + 2*d*e*f*x**4 + d*f**2*x**6), x)`

$$3.34 \quad \int \frac{A+Bx^2}{\sqrt{a-bx^2}\sqrt{c+dx^2}(e+fx^2)^{3/2}} dx$$

Optimal result	283
Mathematica [F]	284
Rubi [F]	284
Maple [F]	285
Fricas [F]	285
Sympy [F]	286
Maxima [F]	286
Giac [F]	286
Mupad [F(-1)]	287
Reduce [F]	287

Optimal result

Integrand size = 42, antiderivative size = 332

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \frac{c(Be - Af)\sqrt{a - bx^2}\sqrt{\frac{e(c+dx^2)}{c(e+fx^2)}}E\left(\arcsin\left(\frac{\sqrt{be+afx}}{\sqrt{a}\sqrt{e+fx^2}}\right) \mid -\frac{a(de-cf)}{c(be+af)}\right)}{\sqrt{ae}\sqrt{be + af}(de - cf)\sqrt{c + dx^2}\sqrt{\frac{e(a-bx^2)}{a(e+fx^2)}}} - \frac{(Bc - Ad)\sqrt{a - bx^2}\sqrt{\frac{e(c+dx^2)}{c(e+fx^2)}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{be+afx}}{\sqrt{a}\sqrt{e+fx^2}}\right), -\frac{a(de-cf)}{c(be+af)}\right)}{\sqrt{a}\sqrt{be + af}(de - cf)\sqrt{c + dx^2}\sqrt{\frac{e(a-bx^2)}{a(e+fx^2)}}}$$

output

```
c*(-A*f+B*e)*(-b*x^2+a)^(1/2)*(e*(d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticE((a*f+b*e)^(1/2)*x/a^(1/2)/(f*x^2+e)^(1/2),(-a*(-c*f+d*e)/c/(a*f+b*e))^(1/2))/a^(1/2)/e/(a*f+b*e)^(1/2)/(-c*f+d*e)/(d*x^2+c)^(1/2)/(e*(-b*x^2+a)/a/(f*x^2+e))^(1/2)-(-A*d+B*c)*(-b*x^2+a)^(1/2)*(e*(d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticF((a*f+b*e)^(1/2)*x/a^(1/2)/(f*x^2+e)^(1/2),(-a*(-c*f+d*e)/c/(a*f+b*e))^(1/2))/a^(1/2)/(a*f+b*e)^(1/2)/(-c*f+d*e)/(d*x^2+c)^(1/2)/(e*(-b*x^2+a)/a/(f*x^2+e))^(1/2)
```

Mathematica [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx$$

input

```
Integrate[(A + B*x^2)/(Sqrt[a - b*x^2]*Sqrt[c + d*x^2]*(e + f*x^2)^(3/2)), x]
```

output

```
Integrate[(A + B*x^2)/(Sqrt[a - b*x^2]*Sqrt[c + d*x^2]*(e + f*x^2)^(3/2)), x]
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx$$

↓ 7293

$$\int \left(\frac{A}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} + \frac{Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} \right) dx$$

↓ 2009

$$A \int \frac{1}{\sqrt{a - bx^2}\sqrt{dx^2 + c}(fx^2 + e)^{3/2}} dx + B \int \frac{x^2}{\sqrt{a - bx^2}\sqrt{dx^2 + c}(fx^2 + e)^{3/2}} dx$$

input

```
Int[(A + B*x^2)/(Sqrt[a - b*x^2]*Sqrt[c + d*x^2]*(e + f*x^2)^(3/2)), x]
```

output

```
$Aborted
```

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]`

Maple [F]

$$\int \frac{x^2 B + A}{\sqrt{-bx^2 + a} \sqrt{x^2 d + c} (fx^2 + e)^{\frac{3}{2}}} dx$$

input `int((B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)`

output `int((B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)`

Fricas [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2} \sqrt{c + dx^2} (e + fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{-bx^2 + a} \sqrt{dx^2 + c} (fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, al
gorithm="fricas")`

output `integral(-(B*x^2 + A)*sqrt(-b*x^2 + a)*sqrt(d*x^2 + c)*sqrt(f*x^2 + e)/(b*
d*f^2*x^8 + (2*b*d*e*f + (b*c - a*d)*f^2)*x^6 + (b*d*e^2 - a*c*f^2 + 2*(b*
c - a*d)*e*f)*x^4 - a*c*e^2 - (2*a*c*e*f - (b*c - a*d)*e^2)*x^2), x)`

Sympy [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{\frac{3}{2}}} dx$$

input `integrate((B*x**2+A)/(-b*x**2+a)**(1/2)/(d*x**2+c)**(1/2)/(f*x**2+e)**(3/2),x)`

output `Integral((A + B*x**2)/(sqrt(a - b*x**2)*sqrt(c + d*x**2)*(e + f*x**2)**(3/2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{-bx^2 + a}\sqrt{dx^2 + c}(fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/(sqrt(-b*x^2 + a)*sqrt(d*x^2 + c)*(f*x^2 + e)^(3/2)), x)`

Giac [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{-bx^2 + a}\sqrt{dx^2 + c}(fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/(sqrt(-b*x^2 + a)*sqrt(d*x^2 + c)*(f*x^2 + e)^(3/2)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{a - bx^2}\sqrt{dx^2 + c}(fx^2 + e)^{3/2}} dx$$

input `int((A + B*x^2)/((a - b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(e + f*x^2)^(3/2)), x)`

output `int((A + B*x^2)/((a - b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(e + f*x^2)^(3/2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \left(\int \frac{\sqrt{fx^2 + e}\sqrt{dx^2 + c}}{-bd f^2 x^8 + ad f^2 x^6 - bc f^2 x^6 - 2bdef x^6 + ac f^2 x^4 + 2adef x^4 + 2acef x^2 + a} dx \right) + \left(\int \frac{\sqrt{fx^2 + e}\sqrt{dx^2 + c}\sqrt{-bx^2 + a}}{-bd f^2 x^8 + ad f^2 x^6 - bc f^2 x^6 - 2bdef x^6 + ac f^2 x^4 + 2adef x^4 - 2bcef x^4 - bde^2 x^4 + 2acef x^2 + a} dx \right)$$

input `int((B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2), x)`

output `int((sqrt(e + f*x**2)*sqrt(c + d*x**2)*sqrt(a - b*x**2)*x**2)/(a*c*e**2 + 2*a*c*e*f*x**2 + a*c*f**2*x**4 + a*d*e**2*x**2 + 2*a*d*e*f*x**4 + a*d*f**2*x**6 - b*c*e**2*x**2 - 2*b*c*e*f*x**4 - b*c*f**2*x**6 - b*d*e**2*x**4 - 2*b*d*e*f*x**6 - b*d*f**2*x**8), x)*b + int((sqrt(e + f*x**2)*sqrt(c + d*x**2)*sqrt(a - b*x**2))/(a*c*e**2 + 2*a*c*e*f*x**2 + a*c*f**2*x**4 + a*d*e**2*x**2 + 2*a*d*e*f*x**4 + a*d*f**2*x**6 - b*c*e**2*x**2 - 2*b*c*e*f*x**4 - b*c*f**2*x**6 - b*d*e**2*x**4 - 2*b*d*e*f*x**6 - b*d*f**2*x**8), x)*a`

3.35 $\int \frac{A+Bx^2}{\sqrt{a+bx^2}\sqrt{c-dx^2}(e+fx^2)^{3/2}} dx$

Optimal result	288
Mathematica [F]	289
Rubi [F]	289
Maple [F]	290
Fricas [F]	290
Sympy [F]	291
Maxima [F]	291
Giac [F]	291
Mupad [F(-1)]	292
Reduce [F]	292

Optimal result

Integrand size = 42, antiderivative size = 333

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \frac{\sqrt{c}(Be - Af)\sqrt{a + bx^2}\sqrt{\frac{e(c-dx^2)}{c(e+fx^2)}}E\left(\arcsin\left(\frac{\sqrt{de+cf}x}{\sqrt{c}\sqrt{e+fx^2}}\right)\right) - \frac{c(be-af)}{a(de+cf)}}{e(be - af)\sqrt{de + cf}\sqrt{c - dx^2}\sqrt{\frac{e(a+bx^2)}{a(e+fx^2)}}} + \frac{(Ab - aB)\sqrt{c}\sqrt{a + bx^2}\sqrt{\frac{e(c-dx^2)}{c(e+fx^2)}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{de+cf}x}{\sqrt{c}\sqrt{e+fx^2}}\right), -\frac{c(be-af)}{a(de+cf)}\right)}{a(be - af)\sqrt{de + cf}\sqrt{c - dx^2}\sqrt{\frac{e(a+bx^2)}{a(e+fx^2)}}}$$

output

```
c^(1/2)*(-A*f+B*e)*(b*x^2+a)^(1/2)*(e*(-d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticE((c*f+d*e)^(1/2)*x/c^(1/2)/(f*x^2+e)^(1/2),(-c*(-a*f+b*e)/a/(c*f+d*e))^(1/2))/e/(-a*f+b*e)/(c*f+d*e)^(1/2)/(-d*x^2+c)^(1/2)/(e*(b*x^2+a)/a/(f*x^2+e))^(1/2)+(A*b-B*a)*c^(1/2)*(b*x^2+a)^(1/2)*(e*(-d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticF((c*f+d*e)^(1/2)*x/c^(1/2)/(f*x^2+e)^(1/2),(-c*(-a*f+b*e)/a/(c*f+d*e))^(1/2))/a/(-a*f+b*e)/(c*f+d*e)^(1/2)/(-d*x^2+c)^(1/2)/(e*(b*x^2+a)/a/(f*x^2+e))^(1/2)
```

Mathematica [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx$$

input

```
Integrate[(A + B*x^2)/(Sqrt[a + b*x^2]*Sqrt[c - d*x^2]*(e + f*x^2)^(3/2)),
x]
```

output

```
Integrate[(A + B*x^2)/(Sqrt[a + b*x^2]*Sqrt[c - d*x^2]*(e + f*x^2)^(3/2)),
x]
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx$$

↓ 7293

$$\int \left(\frac{A}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} + \frac{Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} \right) dx$$

↓ 2009

$$A \int \frac{1}{\sqrt{bx^2 + a}\sqrt{c - dx^2}(fx^2 + e)^{3/2}} dx + B \int \frac{x^2}{\sqrt{bx^2 + a}\sqrt{c - dx^2}(fx^2 + e)^{3/2}} dx$$

input

```
Int[(A + B*x^2)/(Sqrt[a + b*x^2]*Sqrt[c - d*x^2]*(e + f*x^2)^(3/2)),x]
```

output

```
$Aborted
```

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

Maple [F]

$$\int \frac{x^2 B + A}{\sqrt{b x^2 + a} \sqrt{-x^2 d + c} (f x^2 + e)^{\frac{3}{2}}} dx$$

input `int((B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)`

output `int((B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)`

Fricas [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} \sqrt{c - dx^2} (e + fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a} \sqrt{-dx^2 + c} (fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, algorithm="fricas")`

output `integral(-(B*x^2 + A)*sqrt(b*x^2 + a)*sqrt(-d*x^2 + c)*sqrt(f*x^2 + e)/(b*d*f^2*x^8 + (2*b*d*e*f - (b*c - a*d)*f^2)*x^6 + (b*d*e^2 - a*c*f^2 - 2*(b*c - a*d)*e*f)*x^4 - a*c*e^2 - (2*a*c*e*f + (b*c - a*d)*e^2)*x^2), x)`

Sympy [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{\frac{3}{2}}} dx$$

input `integrate((B*x**2+A)/(b*x**2+a)**(1/2)/(-d*x**2+c)**(1/2)/(f*x**2+e)**(3/2),x)`

output `Integral((A + B*x**2)/(sqrt(a + b*x**2)*sqrt(c - d*x**2)*(e + f*x**2)**(3/2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{-dx^2 + c}(fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/(sqrt(b*x^2 + a)*sqrt(-d*x^2 + c)*(f*x^2 + e)^(3/2)), x)`

Giac [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{-dx^2 + c}(fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/(sqrt(b*x^2 + a)*sqrt(-d*x^2 + c)*(f*x^2 + e)^(3/2)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{c - dx^2}(fx^2 + e)^{3/2}} dx$$

input `int((A + B*x^2)/((a + b*x^2)^(1/2)*(c - d*x^2)^(1/2)*(e + f*x^2)^(3/2)),x)`

output `int((A + B*x^2)/((a + b*x^2)^(1/2)*(c - d*x^2)^(1/2)*(e + f*x^2)^(3/2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{\sqrt{fx^2 + e}\sqrt{-dx^2 + c}\sqrt{bx^2 + a}}{-df^2x^6 + cf^2x^4 - 2defx^4 + 2cef x^2 - de^2x^2 + ce^2} dx$$

input `int((B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)`

output `int((sqrt(e + f*x**2)*sqrt(c - d*x**2)*sqrt(a + b*x**2))/(c*e**2 + 2*c*e*f*x**2 + c*f**2*x**4 - d*e**2*x**2 - 2*d*e*f*x**4 - d*f**2*x**6),x)`

3.36
$$\int \frac{A+Bx^2}{\sqrt{a-bx^2}\sqrt{c-dx^2}(e+fx^2)^{3/2}} dx$$

Optimal result	293
Mathematica [F]	294
Rubi [F]	294
Maple [F]	295
Fricas [F]	295
Sympy [F]	296
Maxima [F]	296
Giac [F]	296
Mupad [F(-1)]	297
Reduce [F]	297

Optimal result

Integrand size = 43, antiderivative size = 329

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx =$$

$$-\frac{c(Be - Af)\sqrt{a - bx^2}\sqrt{\frac{e(c-dx^2)}{c(e+fx^2)}}E\left(\arcsin\left(\frac{\sqrt{be+afx}}{\sqrt{a}\sqrt{e+fx^2}}\right)\middle|\frac{a(de+cf)}{c(be+af)}\right)}{\sqrt{ae}\sqrt{be + af}(de + cf)\sqrt{c - dx^2}\sqrt{\frac{e(a-bx^2)}{a(e+fx^2)}}}$$

$$+ \frac{(Bc + Ad)\sqrt{a - bx^2}\sqrt{\frac{e(c-dx^2)}{c(e+fx^2)}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{be+afx}}{\sqrt{a}\sqrt{e+fx^2}}\right), \frac{a(de+cf)}{c(be+af)}\right)}{\sqrt{ae}\sqrt{be + af}(de + cf)\sqrt{c - dx^2}\sqrt{\frac{e(a-bx^2)}{a(e+fx^2)}}}$$

output

```
-c*(-A*f+B*e)*(-b*x^2+a)^(1/2)*(e*(-d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticE(
(a*f+b*e)^(1/2)*x/a^(1/2)/(f*x^2+e)^(1/2),(a*(c*f+d*e)/c/(a*f+b*e))^(1/2))
/a^(1/2)/e/(a*f+b*e)^(1/2)/(c*f+d*e)/(-d*x^2+c)^(1/2)/(e*(-b*x^2+a)/a/(f*x
^2+e))^(1/2)+(A*d+B*c)*(-b*x^2+a)^(1/2)*(e*(-d*x^2+c)/c/(f*x^2+e))^(1/2)*E
llipticF((a*f+b*e)^(1/2)*x/a^(1/2)/(f*x^2+e)^(1/2),(a*(c*f+d*e)/c/(a*f+b*
e))^(1/2))/a^(1/2)/(a*f+b*e)^(1/2)/(c*f+d*e)/(-d*x^2+c)^(1/2)/(e*(-b*x^2+a)
/a/(f*x^2+e))^(1/2)
```

Mathematica [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx$$

input

```
Integrate[(A + B*x^2)/(Sqrt[a - b*x^2]*Sqrt[c - d*x^2]*(e + f*x^2)^(3/2)),
x]
```

output

```
Integrate[(A + B*x^2)/(Sqrt[a - b*x^2]*Sqrt[c - d*x^2]*(e + f*x^2)^(3/2)),
x]
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx$$

↓ 7293

$$\int \left(\frac{A}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} + \frac{Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} \right) dx$$

↓ 2009

$$A \int \frac{1}{\sqrt{a - bx^2}\sqrt{c - dx^2}(fx^2 + e)^{3/2}} dx + B \int \frac{x^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(fx^2 + e)^{3/2}} dx$$

input

```
Int[(A + B*x^2)/(Sqrt[a - b*x^2]*Sqrt[c - d*x^2]*(e + f*x^2)^(3/2)),x]
```

output

```
$Aborted
```

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]`

Maple [F]

$$\int \frac{x^2 B + A}{\sqrt{-bx^2 + a} \sqrt{-x^2 d + c} (fx^2 + e)^{\frac{3}{2}}} dx$$

input `int((B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)`

output `int((B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)`

Fricas [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2} \sqrt{c - dx^2} (e + fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{-bx^2 + a} \sqrt{-dx^2 + c} (fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, a
lgorithm="fricas")`

output `integral((B*x^2 + A)*sqrt(-b*x^2 + a)*sqrt(-d*x^2 + c)*sqrt(f*x^2 + e)/(b*
d*f^2*x^8 + (2*b*d*e*f - (b*c + a*d)*f^2)*x^6 + (b*d*e^2 + a*c*f^2 - 2*(b*
c + a*d)*e*f)*x^4 + a*c*e^2 + (2*a*c*e*f - (b*c + a*d)*e^2)*x^2), x)`

Sympy [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{\frac{3}{2}}} dx$$

input `integrate((B*x**2+A)/(-b*x**2+a)**(1/2)/(-d*x**2+c)**(1/2)/(f*x**2+e)**(3/2),x)`

output `Integral((A + B*x**2)/(sqrt(a - b*x**2)*sqrt(c - d*x**2)*(e + f*x**2)**(3/2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{-bx^2 + a}\sqrt{-dx^2 + c}(fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/(sqrt(-b*x^2 + a)*sqrt(-d*x^2 + c)*(f*x^2 + e)^(3/2)), x)`

Giac [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{-bx^2 + a}\sqrt{-dx^2 + c}(fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/(sqrt(-b*x^2 + a)*sqrt(-d*x^2 + c)*(f*x^2 + e)^(3/2)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{a - bx^2}\sqrt{c - dx^2}(fx^2 + e)^{3/2}} dx$$

input `int((A + B*x^2)/((a - b*x^2)^(1/2)*(c - d*x^2)^(1/2)*(e + f*x^2)^(3/2)),x)`

output `int((A + B*x^2)/((a - b*x^2)^(1/2)*(c - d*x^2)^(1/2)*(e + f*x^2)^(3/2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \left(\int \frac{\sqrt{fx^2 + e}\sqrt{-dx^2}}{bd f^2 x^8 - ad f^2 x^6 - bc f^2 x^6 + 2bdef x^6 + ac f^2 x^4 - 2adef x^4} dx \right) + \left(\int \frac{\sqrt{fx^2 + e}\sqrt{-dx^2 + c}\sqrt{-bx^2 + a}}{bd f^2 x^8 - ad f^2 x^6 - bc f^2 x^6 + 2bdef x^6 + ac f^2 x^4 - 2adef x^4 - 2bcef x^4 + bde^2 x^4 + 2acef x^2 - ad} dx \right)$$

input `int((B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)`

output `int((sqrt(e + f*x**2)*sqrt(c - d*x**2)*sqrt(a - b*x**2)*x**2)/(a*c*e**2 + 2*a*c*e*f*x**2 + a*c*f**2*x**4 - a*d*e**2*x**2 - 2*a*d*e*f*x**4 - a*d*f**2*x**6 - b*c*e**2*x**2 - 2*b*c*e*f*x**4 - b*c*f**2*x**6 + b*d*e**2*x**4 + 2*b*d*e*f*x**6 + b*d*f**2*x**8),x)*b + int((sqrt(e + f*x**2)*sqrt(c - d*x**2)*sqrt(a - b*x**2))/(a*c*e**2 + 2*a*c*e*f*x**2 + a*c*f**2*x**4 - a*d*e**2*x**2 - 2*a*d*e*f*x**4 - a*d*f**2*x**6 - b*c*e**2*x**2 - 2*b*c*e*f*x**4 - b*c*f**2*x**6 + b*d*e**2*x**4 + 2*b*d*e*f*x**6 + b*d*f**2*x**8),x)*a`

$$3.37 \quad \int \frac{A+Bx^2}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e-fx^2)^{3/2}} dx$$

Optimal result	298
Mathematica [F]	299
Rubi [F]	299
Maple [F]	300
Fricas [F]	300
Sympy [F]	301
Maxima [F]	301
Giac [F]	301
Mupad [F(-1)]	302
Reduce [F]	302

Optimal result

Integrand size = 42, antiderivative size = 319

$$\int \frac{A+Bx^2}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e-fx^2)^{3/2}} dx = \frac{(Be+Af)x\sqrt{c+dx^2}}{e(de+cf)\sqrt{a+bx^2}\sqrt{e-fx^2}} - \frac{\sqrt{a}(Be+Af)\sqrt{c+dx^2}E\left(\arctan\left(\frac{\sqrt{be+afx}}{\sqrt{a}\sqrt{e-fx^2}}\right) \middle| \frac{(bc-ad)e}{c(be+af)}\right)}{e\sqrt{be+af}(de+cf)\sqrt{a+bx^2}\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}} + \frac{\sqrt{a}A\sqrt{c+dx^2}\text{EllipticF}\left(\arctan\left(\frac{\sqrt{be+afx}}{\sqrt{a}\sqrt{e-fx^2}}\right), \frac{(bc-ad)e}{c(be+af)}\right)}{ce\sqrt{be+af}\sqrt{a+bx^2}\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}}$$

output

```
(A*f+B*e)*x*(d*x^2+c)^(1/2)/e/(c*f+d*e)/(b*x^2+a)^(1/2)/(-f*x^2+e)^(1/2)-a
^(1/2)*(A*f+B*e)*(d*x^2+c)^(1/2)*EllipticE((a*f+b*e)^(1/2)*x/a^(1/2)/(-f*x
^2+e)^(1/2)/(1+(a*f+b*e)*x^2/a/(-f*x^2+e))^(1/2),((-a*d+b*c)*e/c/(a*f+b*e)
)^(1/2))/e/(a*f+b*e)^(1/2)/(c*f+d*e)/(b*x^2+a)^(1/2)/(a*(d*x^2+c)/c/(b*x^2
+a))^(1/2)+a^(1/2)*A*(d*x^2+c)^(1/2)*InverseJacobiAM(arctan((a*f+b*e)^(1/2)
)*x/a^(1/2)/(-f*x^2+e)^(1/2),((-a*d+b*c)*e/c/(a*f+b*e))^(1/2))/c/e/(a*f+b
*e)^(1/2)/(b*x^2+a)^(1/2)/(a*(d*x^2+c)/c/(b*x^2+a))^(1/2)
```

Mathematica [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx$$

input

```
Integrate[(A + B*x^2)/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(e - f*x^2)^(3/2)),
x]
```

output

```
Integrate[(A + B*x^2)/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(e - f*x^2)^(3/2)),
x]
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx$$

↓ 7293

$$\int \left(\frac{A}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} + \frac{Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} \right) dx$$

↓ 2009

$$A \int \frac{1}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(e - fx^2)^{3/2}} dx + B \int \frac{x^2}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(e - fx^2)^{3/2}} dx$$

input

```
Int[(A + B*x^2)/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(e - f*x^2)^(3/2)),x]
```

output

```
$Aborted
```

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]`

Maple [F]

$$\int \frac{x^2 B + A}{\sqrt{b x^2 + a} \sqrt{x^2 d + c} (-f x^2 + e)^{\frac{3}{2}}} dx$$

input `int((B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

output `int((B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

Fricas [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} \sqrt{c + dx^2} (e - fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a} \sqrt{dx^2 + c} (-fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, al
gorithm="fricas")`

output `integral((B*x^2 + A)*sqrt(b*x^2 + a)*sqrt(d*x^2 + c)*sqrt(-f*x^2 + e)/(b*d
*f^2*x^8 - (2*b*d*e*f - (b*c + a*d)*f^2)*x^6 + (b*d*e^2 + a*c*f^2 - 2*(b*c
+ a*d)*e*f)*x^4 + a*c*e^2 - (2*a*c*e*f - (b*c + a*d)*e^2)*x^2), x)`

Sympy [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{\frac{3}{2}}} dx$$

input `integrate((B*x**2+A)/(b*x**2+a)**(1/2)/(d*x**2+c)**(1/2)/(-f*x**2+e)**(3/2),x)`

output `Integral((A + B*x**2)/(sqrt(a + b*x**2)*sqrt(c + d*x**2)*(e - f*x**2)**(3/2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(-fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)*(-f*x^2 + e)^(3/2)), x)`

Giac [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(-fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)*(-f*x^2 + e)^(3/2)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(e - fx^2)^{3/2}} dx$$

input `int((A + B*x^2)/((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(e - f*x^2)^(3/2)),x)`

output `int((A + B*x^2)/((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(e - f*x^2)^(3/2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{\sqrt{-fx^2 + e}\sqrt{dx^2 + c}\sqrt{bx^2 + a}}{df^2x^6 + cf^2x^4 - 2defx^4 - 2cef x^2 + de^2x^2 + ce^2} dx$$

input `int((B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

output `int((sqrt(e - f*x**2)*sqrt(c + d*x**2)*sqrt(a + b*x**2))/(c*e**2 - 2*c*e*f*x**2 + c*f**2*x**4 + d*e**2*x**2 - 2*d*e*f*x**4 + d*f**2*x**6),x)`

3.38
$$\int \frac{A+Bx^2}{\sqrt{a-bx^2}\sqrt{c+dx^2}(e-fx^2)^{3/2}} dx$$

Optimal result	303
Mathematica [F]	304
Rubi [F]	304
Maple [F]	305
Fricas [F]	305
Sympy [F]	306
Maxima [F]	306
Giac [F]	306
Mupad [F(-1)]	307
Reduce [F]	307

Optimal result

Integrand size = 43, antiderivative size = 339

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \frac{c(Be + Af)\sqrt{a - bx^2}\sqrt{\frac{e(c+dx^2)}{c(e-fx^2)}}E\left(\arcsin\left(\frac{\sqrt{be-afx}}{\sqrt{a}\sqrt{e-fx^2}}\right) \mid -\frac{a(de+cf)}{c(be-af)}\right)}{\sqrt{ae}\sqrt{be - af}(de + cf)\sqrt{c + dx^2}\sqrt{\frac{e(a-bx^2)}{a(e-fx^2)}}} - \frac{(Bc - Ad)\sqrt{a - bx^2}\sqrt{\frac{e(c+dx^2)}{c(e-fx^2)}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{be-afx}}{\sqrt{a}\sqrt{e-fx^2}}\right), -\frac{a(de+cf)}{c(be-af)}\right)}{\sqrt{a}\sqrt{be - af}(de + cf)\sqrt{c + dx^2}\sqrt{\frac{e(a-bx^2)}{a(e-fx^2)}}}$$

output

```
c*(A*f+B*e)*(-b*x^2+a)^(1/2)*(e*(d*x^2+c)/c/(-f*x^2+e))^(1/2)*EllipticE((-a*f+b*e)^(1/2)*x/a^(1/2)/(-f*x^2+e)^(1/2),(-a*(c*f+d*e)/c/(-a*f+b*e))^(1/2))/a^(1/2)/e/(-a*f+b*e)^(1/2)/(c*f+d*e)/(d*x^2+c)^(1/2)/(e*(-b*x^2+a)/a/(-f*x^2+e))^(1/2)-(-A*d+B*c)*(-b*x^2+a)^(1/2)*(e*(d*x^2+c)/c/(-f*x^2+e))^(1/2)*EllipticF((-a*f+b*e)^(1/2)*x/a^(1/2)/(-f*x^2+e)^(1/2),(-a*(c*f+d*e)/c/(-a*f+b*e))^(1/2))/a^(1/2)/(-a*f+b*e)^(1/2)/(c*f+d*e)/(d*x^2+c)^(1/2)/(e*(-b*x^2+a)/a/(-f*x^2+e))^(1/2)
```


Mathematica [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx$$

input

```
Integrate[(A + B*x^2)/(Sqrt[a - b*x^2]*Sqrt[c + d*x^2]*(e - f*x^2)^(3/2)),
x]
```

output

```
Integrate[(A + B*x^2)/(Sqrt[a - b*x^2]*Sqrt[c + d*x^2]*(e - f*x^2)^(3/2)),
x]
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx$$

↓ 7293

$$\int \left(\frac{A}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} + \frac{Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} \right) dx$$

↓ 2009

$$A \int \frac{1}{\sqrt{a - bx^2}\sqrt{dx^2 + c}(e - fx^2)^{3/2}} dx + B \int \frac{x^2}{\sqrt{a - bx^2}\sqrt{dx^2 + c}(e - fx^2)^{3/2}} dx$$

input

```
Int[(A + B*x^2)/(Sqrt[a - b*x^2]*Sqrt[c + d*x^2]*(e - f*x^2)^(3/2)),x]
```

output

```
$Aborted
```

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]`

Maple [F]

$$\int \frac{x^2 B + A}{\sqrt{-bx^2 + a} \sqrt{x^2 d + c} (-fx^2 + e)^{\frac{3}{2}}} dx$$

input `int((B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

output `int((B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

Fricas [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2} \sqrt{c + dx^2} (e - fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{-bx^2 + a} \sqrt{dx^2 + c} (-fx^2 + e)^{3/2}} dx$$

input `integrate((B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, a
lgorithm="fricas")`

output `integral(-(B*x^2 + A)*sqrt(-b*x^2 + a)*sqrt(d*x^2 + c)*sqrt(-f*x^2 + e)/(b
*d*f^2*x^8 - (2*b*d*e*f - (b*c - a*d)*f^2)*x^6 + (b*d*e^2 - a*c*f^2 - 2*(b
*c - a*d)*e*f)*x^4 - a*c*e^2 + (2*a*c*e*f + (b*c - a*d)*e^2)*x^2), x)`

Sympy [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{\frac{3}{2}}} dx$$

input `integrate((B*x**2+A)/(-b*x**2+a)**(1/2)/(d*x**2+c)**(1/2)/(-f*x**2+e)**(3/2),x)`

output `Integral((A + B*x**2)/(sqrt(a - b*x**2)*sqrt(c + d*x**2)*(e - f*x**2)**(3/2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{-bx^2 + a}\sqrt{dx^2 + c}(-fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/(sqrt(-b*x^2 + a)*sqrt(d*x^2 + c)*(-f*x^2 + e)^(3/2)), x)`

Giac [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{-bx^2 + a}\sqrt{dx^2 + c}(-fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/(sqrt(-b*x^2 + a)*sqrt(d*x^2 + c)*(-f*x^2 + e)^(3/2)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{a - bx^2}\sqrt{dx^2 + c}(e - fx^2)^{3/2}} dx$$

input `int((A + B*x^2)/((a - b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(e - f*x^2)^(3/2)),x)`

output `int((A + B*x^2)/((a - b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(e - f*x^2)^(3/2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \left(\int \frac{\sqrt{-fx^2 + e}\sqrt{dx^2 + c}}{-bd f^2 x^8 + ad f^2 x^6 - bc f^2 x^6 + 2bdef x^6 + ac f^2 x^4 - 2adef x^4 - 2acef x^2 + a} \right) + \left(\int \frac{\sqrt{-fx^2 + e}\sqrt{dx^2 + c}\sqrt{-bx^2 + a}}{-bd f^2 x^8 + ad f^2 x^6 - bc f^2 x^6 + 2bdef x^6 + ac f^2 x^4 - 2adef x^4 + 2bcef x^4 - bde^2 x^4 - 2acef x^2 + a} \right)$$

input `int((B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

output `int((sqrt(e - f*x**2)*sqrt(c + d*x**2)*sqrt(a - b*x**2)*x**2)/(a*c*e**2 - 2*a*c*e*f*x**2 + a*c*f**2*x**4 + a*d*e**2*x**2 - 2*a*d*e*f*x**4 + a*d*f**2*x**6 - b*c*e**2*x**2 + 2*b*c*e*f*x**4 - b*c*f**2*x**6 - b*d*e**2*x**4 + 2*b*d*e*f*x**6 - b*d*f**2*x**8),x)*b + int((sqrt(e - f*x**2)*sqrt(c + d*x**2)*sqrt(a - b*x**2))/(a*c*e**2 - 2*a*c*e*f*x**2 + a*c*f**2*x**4 + a*d*e**2*x**2 - 2*a*d*e*f*x**4 + a*d*f**2*x**6 - b*c*e**2*x**2 + 2*b*c*e*f*x**4 - b*c*f**2*x**6 - b*d*e**2*x**4 + 2*b*d*e*f*x**6 - b*d*f**2*x**8),x)*a`

3.39
$$\int \frac{A+Bx^2}{\sqrt{a+bx^2}\sqrt{c-dx^2}(e-fx^2)^{3/2}} dx$$

Optimal result	308
Mathematica [F]	309
Rubi [F]	309
Maple [F]	310
Fricas [F]	310
Sympy [F]	311
Maxima [F]	311
Giac [F]	311
Mupad [F(-1)]	312
Reduce [F]	312

Optimal result

Integrand size = 43, antiderivative size = 340

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \frac{\sqrt{c}(Be + Af)\sqrt{a + bx^2}\sqrt{\frac{e(c-dx^2)}{c(e-fx^2)}}E\left(\arcsin\left(\frac{\sqrt{de-cfx}}{\sqrt{c}\sqrt{e-fx^2}}\right) \mid -\frac{c(be+af)}{a(de-cf)}\right)}{e(be + af)\sqrt{de - cf}\sqrt{c - dx^2}\sqrt{\frac{e(a+bx^2)}{a(e-fx^2)}}} + \frac{(Ab - aB)\sqrt{c}\sqrt{a + bx^2}\sqrt{\frac{e(c-dx^2)}{c(e-fx^2)}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{de-cfx}}{\sqrt{c}\sqrt{e-fx^2}}\right), -\frac{c(be+af)}{a(de-cf)}\right)}{a(be + af)\sqrt{de - cf}\sqrt{c - dx^2}\sqrt{\frac{e(a+bx^2)}{a(e-fx^2)}}}$$

output

```
c^(1/2)*(A*f+B*e)*(b*x^2+a)^(1/2)*(e*(-d*x^2+c)/c/(-f*x^2+e))^(1/2)*EllipticE((-c*f+d*e)^(1/2)*x/c^(1/2)/(-f*x^2+e)^(1/2),(-c*(a*f+b*e)/a/(-c*f+d*e))^(1/2))/e/(a*f+b*e)/(-c*f+d*e)^(1/2)/(-d*x^2+c)^(1/2)/(e*(b*x^2+a)/a/(-f*x^2+e))^(1/2)+(A*b-B*a)*c^(1/2)*(b*x^2+a)^(1/2)*(e*(-d*x^2+c)/c/(-f*x^2+e))^(1/2)*EllipticF((-c*f+d*e)^(1/2)*x/c^(1/2)/(-f*x^2+e)^(1/2),(-c*(a*f+b*e)/a/(-c*f+d*e))^(1/2))/a/(a*f+b*e)/(-c*f+d*e)^(1/2)/(-d*x^2+c)^(1/2)/(e*(b*x^2+a)/a/(-f*x^2+e))^(1/2)
```

Mathematica [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx$$

input

```
Integrate[(A + B*x^2)/(Sqrt[a + b*x^2]*Sqrt[c - d*x^2]*(e - f*x^2)^(3/2)),
x]
```

output

```
Integrate[(A + B*x^2)/(Sqrt[a + b*x^2]*Sqrt[c - d*x^2]*(e - f*x^2)^(3/2)),
x]
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx$$

↓ 7293

$$\int \left(\frac{A}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} + \frac{Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} \right) dx$$

↓ 2009

$$A \int \frac{1}{\sqrt{bx^2 + a}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx + B \int \frac{x^2}{\sqrt{bx^2 + a}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx$$

input

```
Int[(A + B*x^2)/(Sqrt[a + b*x^2]*Sqrt[c - d*x^2]*(e - f*x^2)^(3/2)),x]
```

output

```
$Aborted
```

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]`
`]`

Maple [F]

$$\int \frac{x^2 B + A}{\sqrt{b x^2 + a} \sqrt{-x^2 d + c} (-f x^2 + e)^{\frac{3}{2}}} dx$$

input `int((B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

output `int((B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

Fricas [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2} \sqrt{c - dx^2} (e - fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a} \sqrt{-dx^2 + c} (-fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, a`
`lgorithm="fricas")`

output `integral(-(B*x^2 + A)*sqrt(b*x^2 + a)*sqrt(-d*x^2 + c)*sqrt(-f*x^2 + e)/(b`
`*d*f^2*x^8 - (2*b*d*e*f + (b*c - a*d)*f^2)*x^6 + (b*d*e^2 - a*c*f^2 + 2*(b`
`*c - a*d)*e*f)*x^4 - a*c*e^2 + (2*a*c*e*f - (b*c - a*d)*e^2)*x^2), x)`

Sympy [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{\frac{3}{2}}} dx$$

input `integrate((B*x**2+A)/(b*x**2+a)**(1/2)/(-d*x**2+c)**(1/2)/(-f*x**2+e)**(3/2),x)`

output `Integral((A + B*x**2)/(sqrt(a + b*x**2)*sqrt(c - d*x**2)*(e - f*x**2)**(3/2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{-dx^2 + c}(-fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/(sqrt(b*x^2 + a)*sqrt(-d*x^2 + c)*(-f*x^2 + e)^(3/2)), x)`

Giac [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{-dx^2 + c}(-fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/(sqrt(b*x^2 + a)*sqrt(-d*x^2 + c)*(-f*x^2 + e)^(3/2)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx$$

input `int((A + B*x^2)/((a + b*x^2)^(1/2)*(c - d*x^2)^(1/2)*(e - f*x^2)^(3/2)),x)`

output `int((A + B*x^2)/((a + b*x^2)^(1/2)*(c - d*x^2)^(1/2)*(e - f*x^2)^(3/2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{\sqrt{-fx^2 + e}\sqrt{-dx^2 + c}\sqrt{bx^2 + a}}{-df^2x^6 + cf^2x^4 + 2defx^4 - 2cef x^2 - de^2x^2 + ce^2} dx$$

input `int((B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

output `int((sqrt(e - f*x**2)*sqrt(c - d*x**2)*sqrt(a + b*x**2))/(c*e**2 - 2*c*e*f*x**2 + c*f**2*x**4 - d*e**2*x**2 + 2*d*e*f*x**4 - d*f**2*x**6),x)`

$$3.40 \quad \int \frac{A+Bx^2}{\sqrt{a-bx^2}\sqrt{c-dx^2}(e-fx^2)^{3/2}} dx$$

Optimal result	313
Mathematica [F]	314
Rubi [F]	314
Maple [F]	315
Fricas [F]	315
Sympy [F]	316
Maxima [F]	316
Giac [F]	316
Mupad [F(-1)]	317
Reduce [F]	317

Optimal result

Integrand size = 44, antiderivative size = 344

$$\int \frac{A+Bx^2}{\sqrt{a-bx^2}\sqrt{c-dx^2}(e-fx^2)^{3/2}} dx =$$

$$-\frac{c(Be+Af)\sqrt{a-bx^2}\sqrt{\frac{e(c-dx^2)}{c(e-fx^2)}}E\left(\arcsin\left(\frac{\sqrt{be-afx}}{\sqrt{a}\sqrt{e-fx^2}}\right)\middle|\frac{a(de-cf)}{c(be-af)}\right)}{\sqrt{ae}\sqrt{be-af}(de-cf)\sqrt{c-dx^2}\sqrt{\frac{e(a-bx^2)}{a(e-fx^2)}}}$$

$$+\frac{(Bc+Ad)\sqrt{a-bx^2}\sqrt{\frac{e(c-dx^2)}{c(e-fx^2)}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{be-afx}}{\sqrt{a}\sqrt{e-fx^2}}\right),\frac{a(de-cf)}{c(be-af)}\right)}{\sqrt{ae}\sqrt{be-af}(de-cf)\sqrt{c-dx^2}\sqrt{\frac{e(a-bx^2)}{a(e-fx^2)}}}$$

output

```
-c*(A*f+B*e)*(-b*x^2+a)^(1/2)*(e*(-d*x^2+c)/c/(-f*x^2+e))^(1/2)*EllipticE(
(-a*f+b*e)^(1/2)*x/a^(1/2)/(-f*x^2+e)^(1/2),(a*(-c*f+d*e)/c/(-a*f+b*e))^(1
/2))/a^(1/2)/e/(-a*f+b*e)^(1/2)/(-c*f+d*e)/(-d*x^2+c)^(1/2)/(e*(-b*x^2+a)
a/(-f*x^2+e))^(1/2)+(A*d+B*c)*(-b*x^2+a)^(1/2)*(e*(-d*x^2+c)/c/(-f*x^2+e)
)^(1/2)*EllipticF((-a*f+b*e)^(1/2)*x/a^(1/2)/(-f*x^2+e)^(1/2),(a*(-c*f+d*e)
/c/(-a*f+b*e))^(1/2))/a^(1/2)/(-a*f+b*e)^(1/2)/(-c*f+d*e)/(-d*x^2+c)^(1/2)
/(e*(-b*x^2+a)/a/(-f*x^2+e))^(1/2)
```

Mathematica [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx$$

input

```
Integrate[(A + B*x^2)/(Sqrt[a - b*x^2]*Sqrt[c - d*x^2]*(e - f*x^2)^(3/2)), x]
```

output

```
Integrate[(A + B*x^2)/(Sqrt[a - b*x^2]*Sqrt[c - d*x^2]*(e - f*x^2)^(3/2)), x]
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx$$

↓ 7293

$$\int \left(\frac{A}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} + \frac{Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} \right) dx$$

↓ 2009

$$A \int \frac{1}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx + B \int \frac{x^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx$$

input

```
Int[(A + B*x^2)/(Sqrt[a - b*x^2]*Sqrt[c - d*x^2]*(e - f*x^2)^(3/2)), x]
```

output

```
$Aborted
```

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]`

Maple [F]

$$\int \frac{x^2 B + A}{\sqrt{-b x^2 + a} \sqrt{-x^2 d + c} (-f x^2 + e)^{\frac{3}{2}}} dx$$

input `int((B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

output `int((B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

Fricas [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2} \sqrt{c - dx^2} (e - fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{-bx^2 + a} \sqrt{-dx^2 + c} (-fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x,
algorithm="fricas")`

output `integral((B*x^2 + A)*sqrt(-b*x^2 + a)*sqrt(-d*x^2 + c)*sqrt(-f*x^2 + e)/(b
*d*f^2*x^8 - (2*b*d*e*f + (b*c + a*d)*f^2)*x^6 + (b*d*e^2 + a*c*f^2 + 2*(b
*c + a*d)*e*f)*x^4 + a*c*e^2 - (2*a*c*e*f + (b*c + a*d)*e^2)*x^2), x)`

Sympy [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{\frac{3}{2}}} dx$$

input `integrate((B*x**2+A)/(-b*x**2+a)**(1/2)/(-d*x**2+c)**(1/2)/(-f*x**2+e)**(3/2),x)`

output `Integral((A + B*x**2)/(sqrt(a - b*x**2)*sqrt(c - d*x**2)*(e - f*x**2)**(3/2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{-bx^2 + a}\sqrt{-dx^2 + c}(-fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/(sqrt(-b*x^2 + a)*sqrt(-d*x^2 + c)*(-f*x^2 + e)^(3/2)), x)`

Giac [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{-bx^2 + a}\sqrt{-dx^2 + c}(-fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, algorithm="giac")`

output

```
integrate((B*x^2 + A)/(sqrt(-b*x^2 + a)*sqrt(-d*x^2 + c)*(-f*x^2 + e)^(3/2)), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Bx^2 + A}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx$$

input

```
int((A + B*x^2)/((a - b*x^2)^(1/2)*(c - d*x^2)^(1/2)*(e - f*x^2)^(3/2)),x)
```

output

```
int((A + B*x^2)/((a - b*x^2)^(1/2)*(c - d*x^2)^(1/2)*(e - f*x^2)^(3/2)), x)
```

Reduce [F]

$$\int \frac{A + Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \left(\int \frac{\sqrt{-fx^2 + e}\sqrt{-dx^2 + c}}{bd f^2 x^8 - ad f^2 x^6 - bc f^2 x^6 - 2bdef x^6 + ac f^2 x^4 + 2adef x^4} dx \right) + \left(\int \frac{\sqrt{-fx^2 + e}\sqrt{-dx^2 + c}\sqrt{-bx^2 + a}}{bd f^2 x^8 - ad f^2 x^6 - bc f^2 x^6 - 2bdef x^6 + ac f^2 x^4 + 2adef x^4 + 2bcef x^4 + bd e^2 x^4 - 2acef x^2 - ad e^2} dx \right)$$

input

```
int((B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)
```

output

```
int((sqrt(e - f*x**2)*sqrt(c - d*x**2)*sqrt(a - b*x**2)*x**2)/(a*c*e**2 - 2*a*c*e*f*x**2 + a*c*f**2*x**4 - a*d*e**2*x**2 + 2*a*d*e*f*x**4 - a*d*f**2*x**6 - b*c*e**2*x**2 + 2*b*c*e*f*x**4 - b*c*f**2*x**6 + b*d*e**2*x**4 - 2*b*d*e*f*x**6 + b*d*f**2*x**8),x)*b + int((sqrt(e - f*x**2)*sqrt(c - d*x**2)*sqrt(a - b*x**2))/(a*c*e**2 - 2*a*c*e*f*x**2 + a*c*f**2*x**4 - a*d*e**2*x**2 + 2*a*d*e*f*x**4 - a*d*f**2*x**6 - b*c*e**2*x**2 + 2*b*c*e*f*x**4 - b*c*f**2*x**6 + b*d*e**2*x**4 - 2*b*d*e*f*x**6 + b*d*f**2*x**8),x)*a
```

$$3.41 \quad \int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e+fx^2)^{3/2}} dx$$

Optimal result	318
Mathematica [F]	319
Rubi [F]	319
Maple [F]	320
Fricas [F]	321
Sympy [F]	321
Maxima [F]	321
Giac [F]	322
Mupad [F(-1)]	322
Reduce [F]	323

Optimal result

Integrand size = 46, antiderivative size = 527

$$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e+fx^2)^{3/2}} dx =$$

$$\frac{c(Ce^2 - Bef + Af^2) \sqrt{a+bx^2} \sqrt{\frac{e(c+dx^2)}{c(e+fx^2)}} E\left(\arcsin\left(\frac{\sqrt{-be+afx}}{\sqrt{a}\sqrt{e+fx^2}}\right) \middle| \frac{a(de-cf)}{c(be-af)}\right)}{\sqrt{aef}\sqrt{-be+af}(de-cf)\sqrt{c+dx^2}\sqrt{\frac{e(a+bx^2)}{a(e+fx^2)}}}$$

$$- \frac{((Bc - Ad)f^2 + Ce(de - 2cf)) \sqrt{a+bx^2} \sqrt{\frac{e(c+dx^2)}{c(e+fx^2)}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{-be+afx}}{\sqrt{a}\sqrt{e+fx^2}}\right), \frac{a(de-cf)}{c(be-af)}\right)}{\sqrt{a}f^2\sqrt{-be+af}(de-cf)\sqrt{c+dx^2}\sqrt{\frac{e(a+bx^2)}{a(e+fx^2)}}}$$

$$+ \frac{Ce\sqrt{a+bx^2} \sqrt{\frac{e(c+dx^2)}{c(e+fx^2)}} \text{EllipticPi}\left(-\frac{af}{be-af}, \arcsin\left(\frac{\sqrt{-be+afx}}{\sqrt{a}\sqrt{e+fx^2}}\right), \frac{a(de-cf)}{c(be-af)}\right)}{\sqrt{a}f^2\sqrt{-be+af}\sqrt{c+dx^2}\sqrt{\frac{e(a+bx^2)}{a(e+fx^2)}}}$$

output

```
-c*(A*f^2-B*e*f+C*e^2)*(b*x^2+a)^(1/2)*(e*(d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticE((a*f-b*e)^(1/2)*x/a^(1/2)/(f*x^2+e)^(1/2),(a*(-c*f+d*e)/c/(-a*f+b*e))^(1/2))/a^(1/2)/e/f/(a*f-b*e)^(1/2)/(-c*f+d*e)/(d*x^2+c)^(1/2)/(e*(b*x^2+a)/a/(f*x^2+e))^(1/2)-((-A*d+B*c)*f^2+C*e*(-2*c*f+d*e))*(b*x^2+a)^(1/2)*(e*(d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticF((a*f-b*e)^(1/2)*x/a^(1/2)/(f*x^2+e)^(1/2),(a*(-c*f+d*e)/c/(-a*f+b*e))^(1/2))/a^(1/2)/f^2/(a*f-b*e)^(1/2)/(-c*f+d*e)/(d*x^2+c)^(1/2)/(e*(b*x^2+a)/a/(f*x^2+e))^(1/2)+C*e*(b*x^2+a)^(1/2)*(e*(d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticPi((a*f-b*e)^(1/2)*x/a^(1/2)/(f*x^2+e)^(1/2),-a*f/(-a*f+b*e),(a*(-c*f+d*e)/c/(-a*f+b*e))^(1/2))/a^(1/2)/f^2/(a*f-b*e)^(1/2)/(d*x^2+c)^(1/2)/(e*(b*x^2+a)/a/(f*x^2+e))^(1/2)
```

Mathematica [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx$$

input

```
Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(e + f*x^2)^(3/2)),x]
```

output

```
Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(e + f*x^2)^(3/2)), x]
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx$$

↓ 7293

$$\int \left(\frac{A}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} + \frac{Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} + \frac{Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} \right) dx$$

$$\begin{array}{c}
 \downarrow \text{2009} \\
 A \int \frac{1}{\sqrt{bx^2+a}\sqrt{dx^2+c}(fx^2+e)^{3/2}} dx + B \int \frac{x^2}{\sqrt{bx^2+a}\sqrt{dx^2+c}(fx^2+e)^{3/2}} dx + \\
 C \int \frac{x^4}{\sqrt{bx^2+a}\sqrt{dx^2+c}(fx^2+e)^{3/2}} dx
 \end{array}$$

input `Int[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(e + f*x^2)^(3/2)),x]`

output `$Aborted`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

Maple [F]

$$\int \frac{C x^4 + x^2 B + A}{\sqrt{b x^2 + a} \sqrt{x^2 d + c} (f x^2 + e)^{\frac{3}{2}}} dx$$

input `int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)`

output `int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)`

Fricas [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(fx^2 + e)^{3/2}} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2), x, algorithm="fricas")`

output `integral((C*x^4 + B*x^2 + A)*sqrt(b*x^2 + a)*sqrt(d*x^2 + c)*sqrt(f*x^2 + e)/(b*d*f^2*x^8 + (2*b*d*e*f + (b*c + a*d)*f^2)*x^6 + (b*d*e^2 + a*c*f^2 + 2*(b*c + a*d)*e*f)*x^4 + a*c*e^2 + (2*a*c*e*f + (b*c + a*d)*e^2)*x^2), x)`

Sympy [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx$$

input `integrate((C*x**4+B*x**2+A)/(b*x**2+a)**(1/2)/(d*x**2+c)**(1/2)/(f*x**2+e)**(3/2), x)`

output `Integral((A + B*x**2 + C*x**4)/(sqrt(a + b*x**2)*sqrt(c + d*x**2)*(e + f*x**2)**(3/2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(fx^2 + e)^{3/2}} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2), x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)*(f*x^2 + e)^(3/2)), x)`

Giac [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(fx^2 + e)^{3/2}} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2), x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)*(f*x^2 + e)^(3/2)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(fx^2 + e)^{3/2}} dx$$

input `int((A + B*x^2 + C*x^4)/((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(e + f*x^2)^(3/2)),x)`

output `int((A + B*x^2 + C*x^4)/((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(e + f*x^2)^(3/2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \left(\int \frac{\sqrt{fx^2 + e}\sqrt{dx^2}}{bd f^2x^8 + ad f^2x^6 + bc f^2x^6 + 2bdef x^6 + ac f^2x^4 + 2adef x^4} \right.$$

$$+ \left(\int \frac{\sqrt{fx^2 + e}\sqrt{dx^2 + c}\sqrt{bx^2 + a}x^2}{bd f^2x^8 + ad f^2x^6 + bc f^2x^6 + 2bdef x^6 + ac f^2x^4 + 2adef x^4 + 2bcef x^4 + bd e^2x^4 + 2acef x^2 + ad} \right.$$

$$+ \left(\int \frac{\sqrt{fx^2 + e}\sqrt{dx^2 + c}\sqrt{bx^2 + a}}{bd f^2x^8 + ad f^2x^6 + bc f^2x^6 + 2bdef x^6 + ac f^2x^4 + 2adef x^4 + 2bcef x^4 + bd e^2x^4 + 2acef x^2 + ad} \right)$$

input

```
int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)
```

output

```
int((sqrt(e + f*x**2)*sqrt(c + d*x**2)*sqrt(a + b*x**2)*x**4)/(a*c*e**2 +
2*a*c*e*f*x**2 + a*c*f**2*x**4 + a*d*e**2*x**2 + 2*a*d*e*f*x**4 + a*d*f**2
*x**6 + b*c*e**2*x**2 + 2*b*c*e*f*x**4 + b*c*f**2*x**6 + b*d*e**2*x**4 + 2
*b*d*e*f*x**6 + b*d*f**2*x**8),x)*c + int((sqrt(e + f*x**2)*sqrt(c + d*x**
2)*sqrt(a + b*x**2)*x**2)/(a*c*e**2 + 2*a*c*e*f*x**2 + a*c*f**2*x**4 + a*d
*e**2*x**2 + 2*a*d*e*f*x**4 + a*d*f**2*x**6 + b*c*e**2*x**2 + 2*b*c*e*f*x*
*4 + b*c*f**2*x**6 + b*d*e**2*x**4 + 2*b*d*e*f*x**6 + b*d*f**2*x**8),x)*b
+ int((sqrt(e + f*x**2)*sqrt(c + d*x**2)*sqrt(a + b*x**2))/(a*c*e**2 + 2*a
*c*e*f*x**2 + a*c*f**2*x**4 + a*d*e**2*x**2 + 2*a*d*e*f*x**4 + a*d*f**2*x*
*6 + b*c*e**2*x**2 + 2*b*c*e*f*x**4 + b*c*f**2*x**6 + b*d*e**2*x**4 + 2*b*
d*e*f*x**6 + b*d*f**2*x**8),x)*a
```

$$3.42 \quad \int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}\sqrt{c+dx^2}(e+fx^2)^{3/2}} dx$$

Optimal result	324
Mathematica [F]	325
Rubi [F]	325
Maple [F]	326
Fricas [F]	327
Sympy [F]	327
Maxima [F]	327
Giac [F]	328
Mupad [F(-1)]	328
Reduce [F]	329

Optimal result

Integrand size = 47, antiderivative size = 525

$$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}\sqrt{c+dx^2}(e+fx^2)^{3/2}} dx =$$

$$\frac{c(Ce^2 - Bef + Af^2) \sqrt{a-bx^2} \sqrt{\frac{e(c+dx^2)}{c(e+fx^2)}} E\left(\arcsin\left(\frac{\sqrt{be+afx}}{\sqrt{a}\sqrt{e+fx^2}}\right) \mid -\frac{a(de-cf)}{c(be+af)}\right)}{\sqrt{aef}\sqrt{be+af}(de-cf)\sqrt{c+dx^2}\sqrt{\frac{e(a-bx^2)}{a(e+fx^2)}}}$$

$$- \frac{((Bc - Ad)f^2 + Ce(de - 2cf)) \sqrt{a-bx^2} \sqrt{\frac{e(c+dx^2)}{c(e+fx^2)}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{be+afx}}{\sqrt{a}\sqrt{e+fx^2}}\right), -\frac{a(de-cf)}{c(be+af)}\right)}{\sqrt{a}f^2\sqrt{be+af}(de-cf)\sqrt{c+dx^2}\sqrt{\frac{e(a-bx^2)}{a(e+fx^2)}}}$$

$$+ \frac{Ce\sqrt{a-bx^2} \sqrt{\frac{e(c+dx^2)}{c(e+fx^2)}} \text{EllipticPi}\left(\frac{af}{be+af}, \arcsin\left(\frac{\sqrt{be+afx}}{\sqrt{a}\sqrt{e+fx^2}}\right), -\frac{a(de-cf)}{c(be+af)}\right)}{\sqrt{a}f^2\sqrt{be+af}\sqrt{c+dx^2}\sqrt{\frac{e(a-bx^2)}{a(e+fx^2)}}}$$

output

```
-c*(A*f^2-B*e*f+C*e^2)*(-b*x^2+a)^(1/2)*(e*(d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticE((a*f+b*e)^(1/2)*x/a^(1/2)/(f*x^2+e)^(1/2),(-a*(-c*f+d*e)/c/(a*f+b*e))^(1/2))/a^(1/2)/e/f/(a*f+b*e)^(1/2)/(-c*f+d*e)/(d*x^2+c)^(1/2)/(e*(-b*x^2+a)/a/(f*x^2+e))^(1/2)-((-A*d+B*c)*f^2+C*e*(-2*c*f+d*e))*(-b*x^2+a)^(1/2)*(e*(d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticF((a*f+b*e)^(1/2)*x/a^(1/2)/(f*x^2+e)^(1/2),(-a*(-c*f+d*e)/c/(a*f+b*e))^(1/2))/a^(1/2)/f^2/(a*f+b*e)^(1/2)/(-c*f+d*e)/(d*x^2+c)^(1/2)/(e*(-b*x^2+a)/a/(f*x^2+e))^(1/2)+C*e*(-b*x^2+a)^(1/2)*(e*(d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticPi((a*f+b*e)^(1/2)*x/a^(1/2)/(f*x^2+e)^(1/2),a*f/(a*f+b*e),(-a*(-c*f+d*e)/c/(a*f+b*e))^(1/2))/a^(1/2)/f^2/(a*f+b*e)^(1/2)/(d*x^2+c)^(1/2)/(e*(-b*x^2+a)/a/(f*x^2+e))^(1/2)
```

Mathematica [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx$$

input

```
Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a - b*x^2]*Sqrt[c + d*x^2]*(e + f*x^2)^(3/2)),x]
```

output

```
Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a - b*x^2]*Sqrt[c + d*x^2]*(e + f*x^2)^(3/2)), x]
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx$$

↓ 7293

$$\int \left(\frac{A}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} + \frac{Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} + \frac{Cx^4}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} \right) dx$$

$$\begin{array}{c}
 \downarrow \text{2009} \\
 A \int \frac{1}{\sqrt{a - bx^2} \sqrt{dx^2 + c} (fx^2 + e)^{3/2}} dx + B \int \frac{x^2}{\sqrt{a - bx^2} \sqrt{dx^2 + c} (fx^2 + e)^{3/2}} dx + \\
 C \int \frac{x^4}{\sqrt{a - bx^2} \sqrt{dx^2 + c} (fx^2 + e)^{3/2}} dx
 \end{array}$$

input `Int[(A + B*x^2 + C*x^4)/(Sqrt[a - b*x^2]*Sqrt[c + d*x^2]*(e + f*x^2)^(3/2)),x]`

output `$Aborted`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

Maple [F]

$$\int \frac{C x^4 + x^2 B + A}{\sqrt{-b x^2 + a} \sqrt{x^2 d + c} (f x^2 + e)^{\frac{3}{2}}} dx$$

input `int((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)`

output `int((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)`

Fricas [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{-bx^2 + a}\sqrt{dx^2 + c}(fx^2 + e)^{3/2}} dx$$

input `integrate((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, algorithm="fricas")`

output `integral(-(C*x^4 + B*x^2 + A)*sqrt(-b*x^2 + a)*sqrt(d*x^2 + c)*sqrt(f*x^2 + e)/(b*d*f^2*x^8 + (2*b*d*e*f + (b*c - a*d)*f^2)*x^6 + (b*d*e^2 - a*c*f^2 + 2*(b*c - a*d)*e*f)*x^4 - a*c*e^2 - (2*a*c*e*f - (b*c - a*d)*e^2)*x^2), x)`

Sympy [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx$$

input `integrate((C*x**4+B*x**2+A)/(-b*x**2+a)**(1/2)/(d*x**2+c)**(1/2)/(f*x**2+e)**(3/2),x)`

output `Integral((A + B*x**2 + C*x**4)/(sqrt(a - b*x**2)*sqrt(c + d*x**2)*(e + f*x**2)**(3/2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{-bx^2 + a}\sqrt{dx^2 + c}(fx^2 + e)^{3/2}} dx$$

input `integrate((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, algorithm="maxima")`

output

```
integrate((C*x^4 + B*x^2 + A)/(sqrt(-b*x^2 + a)*sqrt(d*x^2 + c)*(f*x^2 + e)^(3/2)), x)
```

Giac [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{-bx^2 + a}\sqrt{dx^2 + c}(fx^2 + e)^{3/2}} dx$$

input

```
integrate((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, algorithm="giac")
```

output

```
integrate((C*x^4 + B*x^2 + A)/(sqrt(-b*x^2 + a)*sqrt(d*x^2 + c)*(f*x^2 + e)^(3/2)), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{a - bx^2}\sqrt{dx^2 + c}(fx^2 + e)^{3/2}} dx$$

input

```
int((A + B*x^2 + C*x^4)/((a - b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(e + f*x^2)^(3/2)),x)
```

output

```
int((A + B*x^2 + C*x^4)/((a - b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(e + f*x^2)^(3/2)), x)
```

Reduce [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2} \sqrt{c + dx^2} (e + fx^2)^{3/2}} dx = \left(\int \frac{\sqrt{fx^2 + e} \sqrt{dx^2 + c}}{-bd f^2 x^8 + ad f^2 x^6 - bc f^2 x^6 - 2bdef x^6 + ac f^2 x^4 + 2adef x^4} \right.$$

$$+ \left(\int \frac{\sqrt{fx^2 + e} \sqrt{dx^2 + c} \sqrt{-bx^2 + a} x^2}{-bd f^2 x^8 + ad f^2 x^6 - bc f^2 x^6 - 2bdef x^6 + ac f^2 x^4 + 2adef x^4 - 2bcef x^4 - bde^2 x^4 + 2acef x^2 + a} \right.$$

$$+ \left(\int \frac{\sqrt{fx^2 + e} \sqrt{dx^2 + c} \sqrt{-bx^2 + a}}{-bd f^2 x^8 + ad f^2 x^6 - bc f^2 x^6 - 2bdef x^6 + ac f^2 x^4 + 2adef x^4 - 2bcef x^4 - bde^2 x^4 + 2acef x^2 + a} \right)$$

input

```
int((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)
```

output

```
int((sqrt(e + f*x**2)*sqrt(c + d*x**2)*sqrt(a - b*x**2)*x**4)/(a*c*e**2 +
2*a*c*e*f*x**2 + a*c*f**2*x**4 + a*d*e**2*x**2 + 2*a*d*e*f*x**4 + a*d*f**2
*x**6 - b*c*e**2*x**2 - 2*b*c*e*f*x**4 - b*c*f**2*x**6 - b*d*e**2*x**4 - 2
*b*d*e*f*x**6 - b*d*f**2*x**8),x)*c + int((sqrt(e + f*x**2)*sqrt(c + d*x**
2)*sqrt(a - b*x**2)*x**2)/(a*c*e**2 + 2*a*c*e*f*x**2 + a*c*f**2*x**4 + a*d
*e**2*x**2 + 2*a*d*e*f*x**4 + a*d*f**2*x**6 - b*c*e**2*x**2 - 2*b*c*e*f*x**
4 - b*c*f**2*x**6 - b*d*e**2*x**4 - 2*b*d*e*f*x**6 - b*d*f**2*x**8),x)*b
+ int((sqrt(e + f*x**2)*sqrt(c + d*x**2)*sqrt(a - b*x**2))/(a*c*e**2 + 2*a
*c*e*f*x**2 + a*c*f**2*x**4 + a*d*e**2*x**2 + 2*a*d*e*f*x**4 + a*d*f**2*x**
*6 - b*c*e**2*x**2 - 2*b*c*e*f*x**4 - b*c*f**2*x**6 - b*d*e**2*x**4 - 2*b*
d*e*f*x**6 - b*d*f**2*x**8),x)*a
```

3.43
$$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c-dx^2}(e+fx^2)^{3/2}} dx$$

Optimal result	330
Mathematica [F]	331
Rubi [F]	331
Maple [F]	332
Fricas [F(-1)]	333
Sympy [F]	333
Maxima [F]	333
Giac [F]	334
Mupad [F(-1)]	334
Reduce [F]	334

Optimal result

Integrand size = 47, antiderivative size = 533

$$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c-dx^2}(e+fx^2)^{3/2}} dx =$$

$$\frac{\sqrt{c}(Ce^2 - Bef + Af^2) \sqrt{a+bx^2} \sqrt{\frac{e(c-dx^2)}{c(e+fx^2)}} E\left(\arcsin\left(\frac{\sqrt{de+cf}x}{\sqrt{c}\sqrt{e+fx^2}}\right) \mid -\frac{c(be-af)}{a(de+cf)}\right) - ef(be-af)\sqrt{de+cf}\sqrt{c-dx^2}\sqrt{\frac{e(a+bx^2)}{a(e+fx^2)}}}{af^2(be-af)\sqrt{de+cf}\sqrt{c-dx^2}\sqrt{\frac{e(a+bx^2)}{a(e+fx^2)}}}$$

$$+ \frac{\sqrt{c}(af(2Ce - Bf) - b(Ce^2 - Af^2)) \sqrt{a+bx^2} \sqrt{\frac{e(c-dx^2)}{c(e+fx^2)}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{de+cf}x}{\sqrt{c}\sqrt{e+fx^2}}\right), -\frac{c(be-af)}{a(de+cf)}\right)}{af^2(be-af)\sqrt{de+cf}\sqrt{c-dx^2}\sqrt{\frac{e(a+bx^2)}{a(e+fx^2)}}}$$

$$+ \frac{\sqrt{c}Ce\sqrt{a+bx^2}\sqrt{\frac{e(c-dx^2)}{c(e+fx^2)}} \text{EllipticPi}\left(\frac{cf}{de+cf}, \arcsin\left(\frac{\sqrt{de+cf}x}{\sqrt{c}\sqrt{e+fx^2}}\right), -\frac{c(be-af)}{a(de+cf)}\right)}{af^2\sqrt{de+cf}\sqrt{c-dx^2}\sqrt{\frac{e(a+bx^2)}{a(e+fx^2)}}}$$

output

```

-c^(1/2)*(A*f^2-B*e*f+C*e^2)*(b*x^2+a)^(1/2)*(e*(-d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticE((c*f+d*e)^(1/2)*x/c^(1/2)/(f*x^2+e)^(1/2),(-c*(-a*f+b*e)/a/(c*f+d*e))^(1/2))/e/f/(-a*f+b*e)/(c*f+d*e)^(1/2)/(-d*x^2+c)^(1/2)/(e*(b*x^2+a)/a/(f*x^2+e))^(1/2)+c^(1/2)*(a*f*(-B*f+2*C*e)-b*(-A*f^2+C*e^2))*(b*x^2+a)^(1/2)*(e*(-d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticF((c*f+d*e)^(1/2)*x/c^(1/2)/(f*x^2+e)^(1/2),(-c*(-a*f+b*e)/a/(c*f+d*e))^(1/2))/a/f^2/(-a*f+b*e)/(c*f+d*e)^(1/2)/(-d*x^2+c)^(1/2)/(e*(b*x^2+a)/a/(f*x^2+e))^(1/2)+c^(1/2)*C*e*(b*x^2+a)^(1/2)*(e*(-d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticPi((c*f+d*e)^(1/2)*x/c^(1/2)/(f*x^2+e)^(1/2),c*f/(c*f+d*e),(-c*(-a*f+b*e)/a/(c*f+d*e))^(1/2))/a/f^2/(c*f+d*e)^(1/2)/(-d*x^2+c)^(1/2)/(e*(b*x^2+a)/a/(f*x^2+e))^(1/2)

```

Mathematica [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx$$

input

```

Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*Sqrt[c - d*x^2]*(e + f*x^2)^(3/2)),x]

```

output

```

Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*Sqrt[c - d*x^2]*(e + f*x^2)^(3/2)), x]

```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx$$

↓ 7293

$$\int \left(\frac{A}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} + \frac{Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} + \frac{Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} \right) dx$$

$$\begin{array}{c}
 \downarrow \text{2009} \\
 A \int \frac{1}{\sqrt{bx^2 + a}\sqrt{c - dx^2} (fx^2 + e)^{3/2}} dx + B \int \frac{x^2}{\sqrt{bx^2 + a}\sqrt{c - dx^2} (fx^2 + e)^{3/2}} dx + \\
 C \int \frac{x^4}{\sqrt{bx^2 + a}\sqrt{c - dx^2} (fx^2 + e)^{3/2}} dx
 \end{array}$$

input `Int[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*Sqrt[c - d*x^2]*(e + f*x^2)^(3/2)),x]`

output `$Aborted`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

Maple [F]

$$\int \frac{C x^4 + x^2 B + A}{\sqrt{b x^2 + a} \sqrt{-x^2 d + c} (f x^2 + e)^{\frac{3}{2}}} dx$$

input `int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)`

output `int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \text{Timed out}$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, algorithm="fricas")`

output Timed out

Sympy [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{\frac{3}{2}}} dx$$

input `integrate((C*x**4+B*x**2+A)/(b*x**2+a)**(1/2)/(-d*x**2+c)**(1/2)/(f*x**2+e)**(3/2),x)`

output `Integral((A + B*x**2 + C*x**4)/(sqrt(a + b*x**2)*sqrt(c - d*x**2)*(e + f*x**2)**(3/2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{-dx^2 + c}(fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(b*x^2 + a)*sqrt(-d*x^2 + c)*(f*x^2 + e)^(3/2)), x)`

Giac [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{-dx^2 + c}(fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(b*x^2 + a)*sqrt(-d*x^2 + c)*(f*x^2 + e)^(3/2)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{c - dx^2}(fx^2 + e)^{3/2}} dx$$

input `int((A + B*x^2 + C*x^4)/((a + b*x^2)^(1/2)*(c - d*x^2)^(1/2)*(e + f*x^2)^(3/2)),x)`

output `int((A + B*x^2 + C*x^4)/((a + b*x^2)^(1/2)*(c - d*x^2)^(1/2)*(e + f*x^2)^(3/2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \left(\int \frac{\sqrt{fx^2 + e}\sqrt{-dx^2 + c}\sqrt{bx^2 + a}}{-bd f^2 x^8 - ad f^2 x^6 + bc f^2 x^6 - 2bdef x^6 + ac f^2 x^4 - 2adef x^4} \right.$$

$$+ \left(\int \frac{\sqrt{fx^2 + e}\sqrt{-dx^2 + c}\sqrt{bx^2 + a}}{-bd f^2 x^8 - ad f^2 x^6 + bc f^2 x^6 - 2bdef x^6 + ac f^2 x^4 - 2adef x^4 + 2bcef x^4 - bde^2 x^4 + 2acef x^2 - c} \right.$$

$$+ \left(\int \frac{\sqrt{fx^2 + e}\sqrt{-dx^2 + c}\sqrt{bx^2 + a}}{-bd f^2 x^8 - ad f^2 x^6 + bc f^2 x^6 - 2bdef x^6 + ac f^2 x^4 - 2adef x^4 + 2bcef x^4 - bde^2 x^4 + 2acef x^2 - c} \right.$$

input `int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)`

output `int((sqrt(e + f*x**2)*sqrt(c - d*x**2)*sqrt(a + b*x**2)*x**4)/(a*c*e**2 + 2*a*c*e*f*x**2 + a*c*f**2*x**4 - a*d*e**2*x**2 - 2*a*d*e*f*x**4 - a*d*f**2*x**6 + b*c*e**2*x**2 + 2*b*c*e*f*x**4 + b*c*f**2*x**6 - b*d*e**2*x**4 - 2*b*d*e*f*x**6 - b*d*f**2*x**8),x)*c + int((sqrt(e + f*x**2)*sqrt(c - d*x**2)*sqrt(a + b*x**2)*x**2)/(a*c*e**2 + 2*a*c*e*f*x**2 + a*c*f**2*x**4 - a*d*e**2*x**2 - 2*a*d*e*f*x**4 - a*d*f**2*x**6 + b*c*e**2*x**2 + 2*b*c*e*f*x**4 + b*c*f**2*x**6 - b*d*e**2*x**4 - 2*b*d*e*f*x**6 - b*d*f**2*x**8),x)*b + int((sqrt(e + f*x**2)*sqrt(c - d*x**2)*sqrt(a + b*x**2))/(a*c*e**2 + 2*a*c*e*f*x**2 + a*c*f**2*x**4 - a*d*e**2*x**2 - 2*a*d*e*f*x**4 - a*d*f**2*x**6 + b*c*e**2*x**2 + 2*b*c*e*f*x**4 + b*c*f**2*x**6 - b*d*e**2*x**4 - 2*b*d*e*f*x**6 - b*d*f**2*x**8),x)*a`

$$3.44 \quad \int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}\sqrt{c-dx^2}(e+fx^2)^{3/2}} dx$$

Optimal result	336
Mathematica [F]	337
Rubi [F]	337
Maple [F]	338
Fricas [F(-1)]	338
Sympy [F]	339
Maxima [F]	339
Giac [F]	339
Mupad [F(-1)]	340
Reduce [F]	340

Optimal result

Integrand size = 48, antiderivative size = 521

$$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}\sqrt{c-dx^2}(e+fx^2)^{3/2}} dx = \frac{c(Ce^2 - Bef + Af^2) \sqrt{a-bx^2} \sqrt{\frac{e(c-dx^2)}{c(e+fx^2)}} E\left(\arcsin\left(\frac{\sqrt{be+afx}}{\sqrt{a}\sqrt{e+fx^2}}\right)\right)}{\sqrt{aef}\sqrt{be+af}(de+cf)\sqrt{c-dx^2} \sqrt{\frac{e(a-bx^2)}{a(e+fx^2)}}} + \frac{((Bc + Ad)f^2 - Ce(de + 2cf)) \sqrt{a-bx^2} \sqrt{\frac{e(c-dx^2)}{c(e+fx^2)}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{be+afx}}{\sqrt{a}\sqrt{e+fx^2}}\right), \frac{a(de+cf)}{c(be+af)}\right)}{\sqrt{af^2}\sqrt{be+af}(de+cf)\sqrt{c-dx^2} \sqrt{\frac{e(a-bx^2)}{a(e+fx^2)}}} + \frac{Ce\sqrt{a-bx^2} \sqrt{\frac{e(c-dx^2)}{c(e+fx^2)}} \text{EllipticPi}\left(\frac{af}{be+af}, \arcsin\left(\frac{\sqrt{be+afx}}{\sqrt{a}\sqrt{e+fx^2}}\right), \frac{a(de+cf)}{c(be+af)}\right)}{\sqrt{af^2}\sqrt{be+af}\sqrt{c-dx^2} \sqrt{\frac{e(a-bx^2)}{a(e+fx^2)}}}$$

output

```
c*(A*f^2-B*e*f+C*e^2)*(-b*x^2+a)^(1/2)*(e*(-d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticE((a*f+b*e)^(1/2)*x/a^(1/2)/(f*x^2+e)^(1/2), (a*(c*f+d*e)/c/(a*f+b*e))^(1/2))/a^(1/2)/e/f/(a*f+b*e)^(1/2)/(c*f+d*e)/(-d*x^2+c)^(1/2)/(e*(-b*x^2+a)/a/(f*x^2+e))^(1/2)+((A*d+B*c)*f^2-C*e*(2*c*f+d*e))*(-b*x^2+a)^(1/2)*(e*(-d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticF((a*f+b*e)^(1/2)*x/a^(1/2)/(f*x^2+e)^(1/2), (a*(c*f+d*e)/c/(a*f+b*e))^(1/2))/a^(1/2)/f^2/(a*f+b*e)^(1/2)/(c*f+d*e)/(-d*x^2+c)^(1/2)/(e*(-b*x^2+a)/a/(f*x^2+e))^(1/2)+C*e*(-b*x^2+a)^(1/2)*(e*(-d*x^2+c)/c/(f*x^2+e))^(1/2)*EllipticPi((a*f+b*e)^(1/2)*x/a^(1/2)/(f*x^2+e)^(1/2), a*f/(a*f+b*e), (a*(c*f+d*e)/c/(a*f+b*e))^(1/2))/a^(1/2)/f^2/(a*f+b*e)^(1/2)/(-d*x^2+c)^(1/2)/(e*(-b*x^2+a)/a/(f*x^2+e))^(1/2)
```

Mathematica [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx$$

input

```
Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a - b*x^2]*Sqrt[c - d*x^2]*(e + f*x^2)^(3/2)),x]
```

output

```
Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a - b*x^2]*Sqrt[c - d*x^2]*(e + f*x^2)^(3/2)), x]
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx$$

↓ 7293

$$\int \left(\frac{A}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} + \frac{Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} + \frac{Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} \right) dx$$

↓ 2009

$$A \int \frac{1}{\sqrt{a - bx^2}\sqrt{c - dx^2}(fx^2 + e)^{3/2}} dx + B \int \frac{x^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(fx^2 + e)^{3/2}} dx + C \int \frac{x^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(fx^2 + e)^{3/2}} dx$$

input

```
Int[(A + B*x^2 + C*x^4)/(Sqrt[a - b*x^2]*Sqrt[c - d*x^2]*(e + f*x^2)^(3/2)),x]
```

output

```
$Aborted
```

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

Maple [F]

$$\int \frac{Cx^4 + x^2B + A}{\sqrt{-bx^2 + a}\sqrt{-x^2d + c}(fx^2 + e)^{\frac{3}{2}}} dx$$

input `int((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)`

output `int((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \text{Timed out}$$

input `integrate((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{\frac{3}{2}}} dx$$

input `integrate((C*x**4+B*x**2+A)/(-b*x**2+a)**(1/2)/(-d*x**2+c)**(1/2)/(f*x**2+e)**(3/2),x)`

output `Integral((A + B*x**2 + C*x**4)/(sqrt(a - b*x**2)*sqrt(c - d*x**2)*(e + f*x**2)**(3/2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{-bx^2 + a}\sqrt{-dx^2 + c}(fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(-b*x^2 + a)*sqrt(-d*x^2 + c)*(f*x^2 + e)^(3/2)), x)`

Giac [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{-bx^2 + a}\sqrt{-dx^2 + c}(fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2),x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(-b*x^2 + a)*sqrt(-d*x^2 + c)*(f*x^2 + e)^(3/2)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{a - bx^2}\sqrt{c - dx^2}(fx^2 + e)^{3/2}} dx$$

input `int((A + B*x^2 + C*x^4)/((a - b*x^2)^(1/2)*(c - d*x^2)^(1/2)*(e + f*x^2)^(3/2)), x)`

output `int((A + B*x^2 + C*x^4)/((a - b*x^2)^(1/2)*(c - d*x^2)^(1/2)*(e + f*x^2)^(3/2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e + fx^2)^{3/2}} dx = \left(\int \frac{\sqrt{fx^2 + e}\sqrt{-dx^2}}{bd f^2 x^8 - ad f^2 x^6 - bc f^2 x^6 + 2bdef x^6 + ac f^2 x^4 - 2adef x^4} \right. \\ \left. + \left(\int \frac{\sqrt{fx^2 + e}\sqrt{-dx^2 + c}\sqrt{-bx^2 + a}}{bd f^2 x^8 - ad f^2 x^6 - bc f^2 x^6 + 2bdef x^6 + ac f^2 x^4 - 2adef x^4 - 2bcef x^4 + bde^2 x^4 + 2acef x^2 - ad} \right) \right. \\ \left. + \left(\int \frac{\sqrt{fx^2 + e}\sqrt{-dx^2 + c}\sqrt{-bx^2 + a}}{bd f^2 x^8 - ad f^2 x^6 - bc f^2 x^6 + 2bdef x^6 + ac f^2 x^4 - 2adef x^4 - 2bcef x^4 + bde^2 x^4 + 2acef x^2 - ad} \right) \right)$$

input `int((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(f*x^2+e)^(3/2), x)`

output

```

int((sqrt(e + f*x**2)*sqrt(c - d*x**2)*sqrt(a - b*x**2)*x**4)/(a*c*e**2 +
2*a*c*e*f*x**2 + a*c*f**2*x**4 - a*d*e**2*x**2 - 2*a*d*e*f*x**4 - a*d*f**2
*x**6 - b*c*e**2*x**2 - 2*b*c*e*f*x**4 - b*c*f**2*x**6 + b*d*e**2*x**4 + 2
*b*d*e*f*x**6 + b*d*f**2*x**8),x)*c + int((sqrt(e + f*x**2)*sqrt(c - d*x**
2)*sqrt(a - b*x**2)*x**2)/(a*c*e**2 + 2*a*c*e*f*x**2 + a*c*f**2*x**4 - a*d
*e**2*x**2 - 2*a*d*e*f*x**4 - a*d*f**2*x**6 - b*c*e**2*x**2 - 2*b*c*e*f*x
**4 - b*c*f**2*x**6 + b*d*e**2*x**4 + 2*b*d*e*f*x**6 + b*d*f**2*x**8),x)*b
+ int((sqrt(e + f*x**2)*sqrt(c - d*x**2)*sqrt(a - b*x**2))/(a*c*e**2 + 2*a
*c*e*f*x**2 + a*c*f**2*x**4 - a*d*e**2*x**2 - 2*a*d*e*f*x**4 - a*d*f**2*x
**6 - b*c*e**2*x**2 - 2*b*c*e*f*x**4 - b*c*f**2*x**6 + b*d*e**2*x**4 + 2*b
*d*e*f*x**6 + b*d*f**2*x**8),x)*a

```

3.45
$$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e-fx^2)^{3/2}} dx$$

Optimal result	342
Mathematica [F]	343
Rubi [F]	343
Maple [F]	344
Fricas [F(-1)]	345
Sympy [F]	345
Maxima [F]	345
Giac [F]	346
Mupad [F(-1)]	346
Reduce [F]	346

Optimal result

Integrand size = 47, antiderivative size = 496

$$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c+dx^2}(e-fx^2)^{3/2}} dx = \frac{(Ce^2+f(Be+Af))x\sqrt{c+dx^2}}{ef(de+cf)\sqrt{a+bx^2}\sqrt{e-fx^2}}$$

$$- \frac{\sqrt{a}(Ce^2+f(Be+Af))\sqrt{c+dx^2}E\left(\arctan\left(\frac{\sqrt{be+afx}}{\sqrt{a}\sqrt{e-fx^2}}\right) \mid \frac{(bc-ad)e}{c(be+af)}\right)}{ef\sqrt{be+af}(de+cf)\sqrt{a+bx^2}\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}}$$

$$+ \frac{\sqrt{a}(aCe+Abf)\sqrt{c+dx^2}\text{EllipticF}\left(\arctan\left(\frac{\sqrt{be+afx}}{\sqrt{a}\sqrt{e-fx^2}}\right), \frac{(bc-ad)e}{c(be+af)}\right)}{bcef\sqrt{be+af}\sqrt{a+bx^2}\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}}$$

$$- \frac{a^{3/2}C\sqrt{c+dx^2}\text{EllipticPi}\left(\frac{be}{be+af}, \arctan\left(\frac{\sqrt{be+afx}}{\sqrt{a}\sqrt{e-fx^2}}\right), \frac{(bc-ad)e}{c(be+af)}\right)}{bcf\sqrt{be+af}\sqrt{a+bx^2}\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}}$$

output

```
(C*e^2+f*(A*f+B*e))*x*(d*x^2+c)^(1/2)/e/f/(c*f+d*e)/(b*x^2+a)^(1/2)/(-f*x^2+e)^(1/2)-a^(1/2)*(C*e^2+f*(A*f+B*e))*(d*x^2+c)^(1/2)*EllipticE((a*f+b*e)^(1/2)*x/a^(1/2)/(-f*x^2+e)^(1/2)/(1+(a*f+b*e)*x^2/a/(-f*x^2+e)^(1/2)),((-a*d+b*c)*e/c/(a*f+b*e))^(1/2))/e/f/(a*f+b*e)^(1/2)/(c*f+d*e)/(b*x^2+a)^(1/2)/(a*(d*x^2+c)/c/(b*x^2+a))^(1/2)+a^(1/2)*(A*b*f+C*a*e)*(d*x^2+c)^(1/2)*InverseJacobiAM(arctan((a*f+b*e)^(1/2)*x/a^(1/2)/(-f*x^2+e)^(1/2)),((-a*d+b*c)*e/c/(a*f+b*e))^(1/2))/b/c/e/f/(a*f+b*e)^(1/2)/(b*x^2+a)^(1/2)/(a*(d*x^2+c)/c/(b*x^2+a))^(1/2)-a^(3/2)*C*(d*x^2+c)^(1/2)*EllipticPi((a*f+b*e)^(1/2)*x/a^(1/2)/(-f*x^2+e)^(1/2)/(1+(a*f+b*e)*x^2/a/(-f*x^2+e)^(1/2)),b*e/(a*f+b*e)),((-a*d+b*c)*e/c/(a*f+b*e))^(1/2))/b/c/f/(a*f+b*e)^(1/2)/(b*x^2+a)^(1/2)/(a*(d*x^2+c)/c/(b*x^2+a))^(1/2)
```

Mathematica [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx$$

input

```
Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(e - f*x^2)^(3/2)), x]
```

output

```
Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(e - f*x^2)^(3/2)), x]
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx$$

↓ 7293

$$\int \left(\frac{A}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} + \frac{Bx^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} + \frac{Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} \right) dx$$

$$\begin{array}{c}
 \downarrow \text{2009} \\
 A \int \frac{1}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(e - fx^2)^{3/2}} dx + B \int \frac{x^2}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(e - fx^2)^{3/2}} dx + \\
 C \int \frac{x^4}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(e - fx^2)^{3/2}} dx
 \end{array}$$

input `Int[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]*(e - f*x^2)^(3/2)),x]`

output `$Aborted`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

Maple [F]

$$\int \frac{C x^4 + x^2 B + A}{\sqrt{b x^2 + a} \sqrt{x^2 d + c} (-f x^2 + e)^{\frac{3}{2}}} dx$$

input `int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

output `int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \text{Timed out}$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, algorithm="fricas")`

output Timed out

Sympy [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{\frac{3}{2}}} dx$$

input `integrate((C*x**4+B*x**2+A)/(b*x**2+a)**(1/2)/(d*x**2+c)**(1/2)/(-f*x**2+e)**(3/2),x)`

output `Integral((A + B*x**2 + C*x**4)/(sqrt(a + b*x**2)*sqrt(c + d*x**2)*(e - f*x**2)**(3/2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(-fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)*(-f*x^2 + e)^(3/2)), x)`

Giac [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(-fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)*(-f*x^2 + e)^(3/2)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{dx^2 + c}(e - fx^2)^{3/2}} dx$$

input `int((A + B*x^2 + C*x^4)/((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(e - f*x^2)^(3/2)),x)`

output `int((A + B*x^2 + C*x^4)/((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(e - f*x^2)^(3/2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \left(\int \frac{\sqrt{-fx^2 + e}\sqrt{dx^2 + c}}{bd f^2 x^8 + ad f^2 x^6 + bc f^2 x^6 - 2bdef x^6 + ac f^2 x^4 - 2adef x^4} \right. \\ + \left(\int \frac{\sqrt{-fx^2 + e}\sqrt{dx^2 + c}\sqrt{bx^2 + a}}{bd f^2 x^8 + ad f^2 x^6 + bc f^2 x^6 - 2bdef x^6 + ac f^2 x^4 - 2adef x^4 - 2bcef x^4 + bde^2 x^4 - 2acef x^2 + ad} \right) \\ + \left. \left(\int \frac{\sqrt{-fx^2 + e}\sqrt{dx^2 + c}\sqrt{bx^2 + a}}{bd f^2 x^8 + ad f^2 x^6 + bc f^2 x^6 - 2bdef x^6 + ac f^2 x^4 - 2adef x^4 - 2bcef x^4 + bde^2 x^4 - 2acef x^2 + ad} \right) \right)$$

input `int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

output `int((sqrt(e - f*x**2)*sqrt(c + d*x**2)*sqrt(a + b*x**2)*x**4)/(a*c*e**2 - 2*a*c*e*f*x**2 + a*c*f**2*x**4 + a*d*e**2*x**2 - 2*a*d*e*f*x**4 + a*d*f**2*x**6 + b*c*e**2*x**2 - 2*b*c*e*f*x**4 + b*c*f**2*x**6 + b*d*e**2*x**4 - 2*b*d*e*f*x**6 + b*d*f**2*x**8),x)*c + int((sqrt(e - f*x**2)*sqrt(c + d*x**2)*sqrt(a + b*x**2)*x**2)/(a*c*e**2 - 2*a*c*e*f*x**2 + a*c*f**2*x**4 + a*d*e**2*x**2 - 2*a*d*e*f*x**4 + a*d*f**2*x**6 + b*c*e**2*x**2 - 2*b*c*e*f*x**4 + b*c*f**2*x**6 + b*d*e**2*x**4 - 2*b*d*e*f*x**6 + b*d*f**2*x**8),x)*b + int((sqrt(e - f*x**2)*sqrt(c + d*x**2)*sqrt(a + b*x**2))/(a*c*e**2 - 2*a*c*e*f*x**2 + a*c*f**2*x**4 + a*d*e**2*x**2 - 2*a*d*e*f*x**4 + a*d*f**2*x**6 + b*c*e**2*x**2 - 2*b*c*e*f*x**4 + b*c*f**2*x**6 + b*d*e**2*x**4 - 2*b*d*e*f*x**6 + b*d*f**2*x**8),x)*a`

3.46
$$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}\sqrt{c+dx^2}(e-fx^2)^{3/2}} dx$$

Optimal result	348
Mathematica [F]	349
Rubi [F]	349
Maple [F]	350
Fricas [F]	351
Sympy [F]	351
Maxima [F]	351
Giac [F]	352
Mupad [F(-1)]	352
Reduce [F]	353

Optimal result

Integrand size = 48, antiderivative size = 538

$$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}\sqrt{c+dx^2}(e-fx^2)^{3/2}} dx = \frac{c(Ce^2+f(Be+Af))\sqrt{a-bx^2}\sqrt{\frac{e(c+dx^2)}{c(e-fx^2)}}E\left(\arcsin\left(\frac{\sqrt{be-afx}}{\sqrt{a}\sqrt{e-fx^2}}\right)\right)}{\sqrt{aef}\sqrt{be-af}(de+cf)\sqrt{c+dx^2}\sqrt{\frac{e(a-bx^2)}{a(e-fx^2)}}} - \frac{((Bc-Ad)f^2+Ce(de+2cf))\sqrt{a-bx^2}\sqrt{\frac{e(c+dx^2)}{c(e-fx^2)}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{be-afx}}{\sqrt{a}\sqrt{e-fx^2}}\right),-\frac{a(de+cf)}{c(be-af)}\right)}{\sqrt{a}f^2\sqrt{be-af}(de+cf)\sqrt{c+dx^2}\sqrt{\frac{e(a-bx^2)}{a(e-fx^2)}}} + \frac{Ce\sqrt{a-bx^2}\sqrt{\frac{e(c+dx^2)}{c(e-fx^2)}}\text{EllipticPi}\left(-\frac{af}{be-af},\arcsin\left(\frac{\sqrt{be-afx}}{\sqrt{a}\sqrt{e-fx^2}}\right),-\frac{a(de+cf)}{c(be-af)}\right)}{\sqrt{a}f^2\sqrt{be-af}\sqrt{c+dx^2}\sqrt{\frac{e(a-bx^2)}{a(e-fx^2)}}}$$

output

```
c*(C*e^2+f*(A*f+B*e))*(-b*x^2+a)^(1/2)*(e*(d*x^2+c)/c/(-f*x^2+e))^(1/2)*EllipticE((-a*f+b*e)^(1/2)*x/a^(1/2)/(-f*x^2+e)^(1/2),(-a*(c*f+d*e)/c/(-a*f+b*e))^(1/2))/a^(1/2)/e/f/(-a*f+b*e)^(1/2)/(c*f+d*e)/(d*x^2+c)^(1/2)/(e*(-b*x^2+a)/a/(-f*x^2+e))^(1/2)-((-A*d+B*c)*f^2+C*e*(2*c*f+d*e))*(-b*x^2+a)^(1/2)*(e*(d*x^2+c)/c/(-f*x^2+e))^(1/2)*EllipticF((-a*f+b*e)^(1/2)*x/a^(1/2)/(-f*x^2+e)^(1/2),(-a*(c*f+d*e)/c/(-a*f+b*e))^(1/2))/a^(1/2)/f^2/(-a*f+b*e)^(1/2)/(c*f+d*e)/(d*x^2+c)^(1/2)/(e*(-b*x^2+a)/a/(-f*x^2+e))^(1/2)+C*e*(-b*x^2+a)^(1/2)*(e*(d*x^2+c)/c/(-f*x^2+e))^(1/2)*EllipticPi((-a*f+b*e)^(1/2)*x/a^(1/2)/(-f*x^2+e)^(1/2),-a*f/(-a*f+b*e),(-a*(c*f+d*e)/c/(-a*f+b*e))^(1/2))/a^(1/2)/f^2/(-a*f+b*e)^(1/2)/(d*x^2+c)^(1/2)/(e*(-b*x^2+a)/a/(-f*x^2+e))^(1/2)
```

Mathematica [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx$$

input

```
Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a - b*x^2]*Sqrt[c + d*x^2]*(e - f*x^2)^(3/2)), x]
```

output

```
Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a - b*x^2]*Sqrt[c + d*x^2]*(e - f*x^2)^(3/2)), x]
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx$$

↓ 7293

$$\int \left(\frac{A}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} + \frac{Bx^2}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} + \frac{Cx^4}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} \right) dx$$

$$\begin{array}{c}
 \downarrow \text{2009} \\
 A \int \frac{1}{\sqrt{a - bx^2} \sqrt{dx^2 + c} (e - fx^2)^{3/2}} dx + B \int \frac{x^2}{\sqrt{a - bx^2} \sqrt{dx^2 + c} (e - fx^2)^{3/2}} dx + \\
 C \int \frac{x^4}{\sqrt{a - bx^2} \sqrt{dx^2 + c} (e - fx^2)^{3/2}} dx
 \end{array}$$

input `Int[(A + B*x^2 + C*x^4)/(Sqrt[a - b*x^2]*Sqrt[c + d*x^2]*(e - f*x^2)^(3/2)),x]`

output `$Aborted`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

Maple [F]

$$\int \frac{C x^4 + x^2 B + A}{\sqrt{-b x^2 + a} \sqrt{x^2 d + c} (-f x^2 + e)^{\frac{3}{2}}} dx$$

input `int((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

output `int((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

Fricas [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{-bx^2 + a}\sqrt{dx^2 + c}(-fx^2 + e)^{3/2}} dx$$

input `integrate((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, algorithm="fricas")`

output `integral(-(C*x^4 + B*x^2 + A)*sqrt(-b*x^2 + a)*sqrt(d*x^2 + c)*sqrt(-f*x^2 + e)/(b*d*f^2*x^8 - (2*b*d*e*f - (b*c - a*d)*f^2)*x^6 + (b*d*e^2 - a*c*f^2 - 2*(b*c - a*d)*e*f)*x^4 - a*c*e^2 + (2*a*c*e*f + (b*c - a*d)*e^2)*x^2), x)`

Sympy [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx$$

input `integrate((C*x**4+B*x**2+A)/(-b*x**2+a)**(1/2)/(d*x**2+c)**(1/2)/(-f*x**2+e)**(3/2),x)`

output `Integral((A + B*x**2 + C*x**4)/(sqrt(a - b*x**2)*sqrt(c + d*x**2)*(e - f*x**2)**(3/2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{-bx^2 + a}\sqrt{dx^2 + c}(-fx^2 + e)^{3/2}} dx$$

input `integrate((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(-b*x^2 + a)*sqrt(d*x^2 + c)*(-f*x^2 + e)^(3/2)), x)`

Giac [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{-bx^2 + a}\sqrt{dx^2 + c}(-fx^2 + e)^{3/2}} dx$$

input `integrate((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(-b*x^2 + a)*sqrt(d*x^2 + c)*(-f*x^2 + e)^(3/2)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c + dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{a - bx^2}\sqrt{dx^2 + c}(e - fx^2)^{3/2}} dx$$

input `int((A + B*x^2 + C*x^4)/((a - b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(e - f*x^2)^(3/2)),x)`

output `int((A + B*x^2 + C*x^4)/((a - b*x^2)^(1/2)*(c + d*x^2)^(1/2)*(e - f*x^2)^(3/2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2} \sqrt{c + dx^2} (e - fx^2)^{3/2}} dx = \left(\int \frac{\sqrt{-fx^2 + e} \sqrt{dx^2 + c}}{-bd f^2 x^8 + ad f^2 x^6 - bc f^2 x^6 + 2bdef x^6 + ac f^2 x^4 - 2adef x^4 - 2acef x^2 + a^2} dx \right.$$

$$+ \left(\int \frac{\sqrt{-fx^2 + e} \sqrt{dx^2 + c} \sqrt{-bx^2 + a} x^2}{-bd f^2 x^8 + ad f^2 x^6 - bc f^2 x^6 + 2bdef x^6 + ac f^2 x^4 - 2adef x^4 + 2bcef x^4 - bd e^2 x^4 - 2acef x^2 + a^2} dx \right.$$

$$+ \left(\int \frac{\sqrt{-fx^2 + e} \sqrt{dx^2 + c} \sqrt{-bx^2 + a} x^4}{-bd f^2 x^8 + ad f^2 x^6 - bc f^2 x^6 + 2bdef x^6 + ac f^2 x^4 - 2adef x^4 + 2bcef x^4 - bd e^2 x^4 - 2acef x^2 + a^2} dx \right)$$

input `int((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

output

```
int((sqrt(e - f*x**2)*sqrt(c + d*x**2)*sqrt(a - b*x**2)*x**4)/(a*c*e**2 -
2*a*c*e*f*x**2 + a*c*f**2*x**4 + a*d*e**2*x**2 - 2*a*d*e*f*x**4 + a*d*f**2
*x**6 - b*c*e**2*x**2 + 2*b*c*e*f*x**4 - b*c*f**2*x**6 - b*d*e**2*x**4 + 2
*b*d*e*f*x**6 - b*d*f**2*x**8),x)*c + int((sqrt(e - f*x**2)*sqrt(c + d*x**
2)*sqrt(a - b*x**2)*x**2)/(a*c*e**2 - 2*a*c*e*f*x**2 + a*c*f**2*x**4 + a*d
*e**2*x**2 - 2*a*d*e*f*x**4 + a*d*f**2*x**6 - b*c*e**2*x**2 + 2*b*c*e*f*x**
4 - b*c*f**2*x**6 - b*d*e**2*x**4 + 2*b*d*e*f*x**6 - b*d*f**2*x**8),x)*b
+ int((sqrt(e - f*x**2)*sqrt(c + d*x**2)*sqrt(a - b*x**2))/(a*c*e**2 - 2*a
*c*e*f*x**2 + a*c*f**2*x**4 + a*d*e**2*x**2 - 2*a*d*e*f*x**4 + a*d*f**2*x**
6 - b*c*e**2*x**2 + 2*b*c*e*f*x**4 - b*c*f**2*x**6 - b*d*e**2*x**4 + 2*b*
d*e*f*x**6 - b*d*f**2*x**8),x)*a
```

$$3.47 \quad \int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c-dx^2}(e-fx^2)^{3/2}} dx$$

Optimal result	354
Mathematica [F]	355
Rubi [F]	355
Maple [F]	356
Fricas [F]	357
Sympy [F]	357
Maxima [F]	357
Giac [F]	358
Mupad [F(-1)]	358
Reduce [F]	359

Optimal result

Integrand size = 48, antiderivative size = 545

$$\int \frac{A+Bx^2+Cx^4}{\sqrt{a+bx^2}\sqrt{c-dx^2}(e-fx^2)^{3/2}} dx = \frac{\sqrt{c}(Ce^2+f(Be+Af))\sqrt{a+bx^2}\sqrt{\frac{e(c-dx^2)}{c(e-fx^2)}}E\left(\arcsin\left(\frac{\sqrt{de-cfx}}{\sqrt{c}\sqrt{e-fx^2}}\right)\right)}{ef(be+af)\sqrt{de-cf}\sqrt{c-dx^2}\sqrt{\frac{e(a+bx^2)}{a(e-fx^2)}}}$$

$$- \frac{\sqrt{c}(af(2Ce+Bf)+b(Ce^2-Af^2))\sqrt{a+bx^2}\sqrt{\frac{e(c-dx^2)}{c(e-fx^2)}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{de-cfx}}{\sqrt{c}\sqrt{e-fx^2}}\right), -\frac{c(be+af)}{a(de-cf)}\right)}{af^2(be+af)\sqrt{de-cf}\sqrt{c-dx^2}\sqrt{\frac{e(a+bx^2)}{a(e-fx^2)}}}$$

$$+ \frac{\sqrt{c}Ce\sqrt{a+bx^2}\sqrt{\frac{e(c-dx^2)}{c(e-fx^2)}}\text{EllipticPi}\left(-\frac{cf}{de-cf}, \arcsin\left(\frac{\sqrt{de-cfx}}{\sqrt{c}\sqrt{e-fx^2}}\right), -\frac{c(be+af)}{a(de-cf)}\right)}{af^2\sqrt{de-cf}\sqrt{c-dx^2}\sqrt{\frac{e(a+bx^2)}{a(e-fx^2)}}}$$

output

```

c^(1/2)*(C*e^2+f*(A*f+B*e))*(b*x^2+a)^(1/2)*(e*(-d*x^2+c)/c/(-f*x^2+e))^(1/2)*EllipticE((-c*f+d*e)^(1/2)*x/c^(1/2)/(-f*x^2+e)^(1/2),(-c*(a*f+b*e)/a/(-c*f+d*e))^(1/2))/e/f/(a*f+b*e)/(-c*f+d*e)^(1/2)/(-d*x^2+c)^(1/2)/(e*(b*x^2+a)/a/(-f*x^2+e))^(1/2)-c^(1/2)*(a*f*(B*f+2*C*e)+b*(-A*f^2+C*e^2))*(b*x^2+a)^(1/2)*(e*(-d*x^2+c)/c/(-f*x^2+e))^(1/2)*EllipticF((-c*f+d*e)^(1/2)*x/c^(1/2)/(-f*x^2+e)^(1/2),(-c*(a*f+b*e)/a/(-c*f+d*e))^(1/2))/a/f^2/(a*f+b*e)/(-c*f+d*e)^(1/2)/(-d*x^2+c)^(1/2)/(e*(b*x^2+a)/a/(-f*x^2+e))^(1/2)+c^(1/2)*C*e*(b*x^2+a)^(1/2)*(e*(-d*x^2+c)/c/(-f*x^2+e))^(1/2)*EllipticPi((-c*f+d*e)^(1/2)*x/c^(1/2)/(-f*x^2+e)^(1/2),-c*f/(-c*f+d*e),(-c*(a*f+b*e)/a/(-c*f+d*e))^(1/2))/a/f^2/(-c*f+d*e)^(1/2)/(-d*x^2+c)^(1/2)/(e*(b*x^2+a)/a/(-f*x^2+e))^(1/2)

```

Mathematica [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx$$

input

```

Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*Sqrt[c - d*x^2]*(e - f*x^2)^(3/2)), x]

```

output

```

Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*Sqrt[c - d*x^2]*(e - f*x^2)^(3/2)), x]

```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx$$

↓ 7293

$$\int \left(\frac{A}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} + \frac{Bx^2}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} + \frac{Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} \right) dx$$

$$\begin{array}{c}
 \downarrow \text{2009} \\
 A \int \frac{1}{\sqrt{bx^2 + a}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx + B \int \frac{x^2}{\sqrt{bx^2 + a}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx + \\
 C \int \frac{x^4}{\sqrt{bx^2 + a}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx
 \end{array}$$

input `Int[(A + B*x^2 + C*x^4)/(Sqrt[a + b*x^2]*Sqrt[c - d*x^2]*(e - f*x^2)^(3/2)),x]`

output `$Aborted`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

Maple [F]

$$\int \frac{C x^4 + x^2 B + A}{\sqrt{b x^2 + a} \sqrt{-x^2 d + c} (-f x^2 + e)^{\frac{3}{2}}} dx$$

input `int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

output `int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

Fricas [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{-dx^2 + c}(-fx^2 + e)^{3/2}} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, algorithm="fricas")`

output `integral(-(C*x^4 + B*x^2 + A)*sqrt(b*x^2 + a)*sqrt(-d*x^2 + c)*sqrt(-f*x^2 + e)/(b*d*f^2*x^8 - (2*b*d*e*f + (b*c - a*d)*f^2)*x^6 + (b*d*e^2 - a*c*f^2 + 2*(b*c - a*d)*e*f)*x^4 - a*c*e^2 + (2*a*c*e*f - (b*c - a*d)*e^2)*x^2), x)`

Sympy [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx$$

input `integrate((C*x**4+B*x**2+A)/(b*x**2+a)**(1/2)/(-d*x**2+c)**(1/2)/(-f*x**2+e)**(3/2),x)`

output `Integral((A + B*x**2 + C*x**4)/(sqrt(a + b*x**2)*sqrt(c - d*x**2)*(e - f*x**2)**(3/2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{-dx^2 + c}(-fx^2 + e)^{3/2}} dx$$

input `integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, algorithm="maxima")`

output

```
integrate((C*x^4 + B*x^2 + A)/(sqrt(b*x^2 + a)*sqrt(-d*x^2 + c)*(-f*x^2 + e)^(3/2)), x)
```

Giac [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{-dx^2 + c}(-fx^2 + e)^{3/2}} dx$$

input

```
integrate((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, algorithm="giac")
```

output

```
integrate((C*x^4 + B*x^2 + A)/(sqrt(b*x^2 + a)*sqrt(-d*x^2 + c)*(-f*x^2 + e)^(3/2)), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{bx^2 + a}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx$$

input

```
int((A + B*x^2 + C*x^4)/((a + b*x^2)^(1/2)*(c - d*x^2)^(1/2)*(e - f*x^2)^(3/2)),x)
```

output

```
int((A + B*x^2 + C*x^4)/((a + b*x^2)^(1/2)*(c - d*x^2)^(1/2)*(e - f*x^2)^(3/2)), x)
```

Reduce [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a + bx^2} \sqrt{c - dx^2} (e - fx^2)^{3/2}} dx = \left(\int \frac{\sqrt{-fx^2 + e} \sqrt{-d}}{-bd f^2 x^8 - ad f^2 x^6 + bc f^2 x^6 + 2bdef x^6 + ac f^2 x^4 + 2adef x^4 - 2bcef x^4 - bd e^2 x^4 - 2acef x^2 - a^2 e^2} \right)$$

$$+ \left(\int \frac{\sqrt{-fx^2 + e} \sqrt{-dx^2 + c} \sqrt{bx^2 + a} x^2}{-bd f^2 x^8 - ad f^2 x^6 + bc f^2 x^6 + 2bdef x^6 + ac f^2 x^4 + 2adef x^4 - 2bcef x^4 - bd e^2 x^4 - 2acef x^2 - a^2 e^2} \right)$$

$$+ \left(\int \frac{\sqrt{-fx^2 + e} \sqrt{-dx^2 + c} \sqrt{bx^2 + a}}{-bd f^2 x^8 - ad f^2 x^6 + bc f^2 x^6 + 2bdef x^6 + ac f^2 x^4 + 2adef x^4 - 2bcef x^4 - bd e^2 x^4 - 2acef x^2 - a^2 e^2} \right)$$

input `int((C*x^4+B*x^2+A)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

output

```
int((sqrt(e - f*x**2)*sqrt(c - d*x**2)*sqrt(a + b*x**2)*x**4)/(a*c*e**2 -
2*a*c*e*f*x**2 + a*c*f**2*x**4 - a*d*e**2*x**2 + 2*a*d*e*f*x**4 - a*d*f**2
*x**6 + b*c*e**2*x**2 - 2*b*c*e*f*x**4 + b*c*f**2*x**6 - b*d*e**2*x**4 + 2
*b*d*e*f*x**6 - b*d*f**2*x**8),x)*c + int((sqrt(e - f*x**2)*sqrt(c - d*x**
2)*sqrt(a + b*x**2)*x**2)/(a*c*e**2 - 2*a*c*e*f*x**2 + a*c*f**2*x**4 - a*d
*e**2*x**2 + 2*a*d*e*f*x**4 - a*d*f**2*x**6 + b*c*e**2*x**2 - 2*b*c*e*f*x
**4 + b*c*f**2*x**6 - b*d*e**2*x**4 + 2*b*d*e*f*x**6 - b*d*f**2*x**8),x)*b
+ int((sqrt(e - f*x**2)*sqrt(c - d*x**2)*sqrt(a + b*x**2))/(a*c*e**2 - 2*a
*c*e*f*x**2 + a*c*f**2*x**4 - a*d*e**2*x**2 + 2*a*d*e*f*x**4 - a*d*f**2*x
**6 + b*c*e**2*x**2 - 2*b*c*e*f*x**4 + b*c*f**2*x**6 - b*d*e**2*x**4 + 2*b
*d*e*f*x**6 - b*d*f**2*x**8),x)*a
```


$$3.48 \quad \int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}\sqrt{c-dx^2}(e-fx^2)^{3/2}} dx$$

Optimal result	360
Mathematica [F]	361
Rubi [F]	361
Maple [F]	362
Fricas [F]	363
Sympy [F]	363
Maxima [F]	363
Giac [F]	364
Mupad [F(-1)]	364
Reduce [F]	365

Optimal result

Integrand size = 49, antiderivative size = 546

$$\int \frac{A+Bx^2+Cx^4}{\sqrt{a-bx^2}\sqrt{c-dx^2}(e-fx^2)^{3/2}} dx =$$

$$\frac{c(Ce^2+f(Be+Af))\sqrt{a-bx^2}\sqrt{\frac{e(c-dx^2)}{c(e-fx^2)}}E\left(\arcsin\left(\frac{\sqrt{be-afx}}{\sqrt{a}\sqrt{e-fx^2}}\right)\mid\frac{a(de-cf)}{c(be-af)}\right)}{\sqrt{aef}\sqrt{be-af}(de-cf)\sqrt{c-dx^2}\sqrt{\frac{e(a-bx^2)}{a(e-fx^2)}}}$$

$$+\frac{((Bc+Ad)f^2-Ce(de-2cf))\sqrt{a-bx^2}\sqrt{\frac{e(c-dx^2)}{c(e-fx^2)}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{be-afx}}{\sqrt{a}\sqrt{e-fx^2}}\right),\frac{a(de-cf)}{c(be-af)}\right)}{\sqrt{af^2}\sqrt{be-af}(de-cf)\sqrt{c-dx^2}\sqrt{\frac{e(a-bx^2)}{a(e-fx^2)}}}$$

$$+\frac{Ce\sqrt{a-bx^2}\sqrt{\frac{e(c-dx^2)}{c(e-fx^2)}}\text{EllipticPi}\left(-\frac{af}{be-af},\arcsin\left(\frac{\sqrt{be-afx}}{\sqrt{a}\sqrt{e-fx^2}}\right),\frac{a(de-cf)}{c(be-af)}\right)}{\sqrt{af^2}\sqrt{be-af}\sqrt{c-dx^2}\sqrt{\frac{e(a-bx^2)}{a(e-fx^2)}}}$$

output

```
-c*(C*e^2+f*(A*f+B*e))*(-b*x^2+a)^(1/2)*(e*(-d*x^2+c)/c/(-f*x^2+e))^(1/2)*
EllipticE((-a*f+b*e)^(1/2)*x/a^(1/2)/(-f*x^2+e)^(1/2),(a*(-c*f+d*e)/c/(-a*
f+b*e))^(1/2))/a^(1/2)/e/f/(-a*f+b*e)^(1/2)/(-c*f+d*e)/(-d*x^2+c)^(1/2)/(e
*(-b*x^2+a)/a/(-f*x^2+e))^(1/2)+((A*d+B*c)*f^2-C*e*(-2*c*f+d*e))*(-b*x^2+a
)^(1/2)*(e*(-d*x^2+c)/c/(-f*x^2+e))^(1/2)*EllipticF((-a*f+b*e)^(1/2)*x/a^(
1/2)/(-f*x^2+e)^(1/2),(a*(-c*f+d*e)/c/(-a*f+b*e))^(1/2))/a^(1/2)/f^2/(-a*f
+b*e)^(1/2)/(-c*f+d*e)/(-d*x^2+c)^(1/2)/(e*(-b*x^2+a)/a/(-f*x^2+e))^(1/2)+
C*e*(-b*x^2+a)^(1/2)*(e*(-d*x^2+c)/c/(-f*x^2+e))^(1/2)*EllipticPi((-a*f+b*
e)^(1/2)*x/a^(1/2)/(-f*x^2+e)^(1/2),-a*f/(-a*f+b*e),(a*(-c*f+d*e)/c/(-a*f+
b*e))^(1/2))/a^(1/2)/f^2/(-a*f+b*e)^(1/2)/(-d*x^2+c)^(1/2)/(e*(-b*x^2+a)/
a/(-f*x^2+e))^(1/2)
```

Mathematica [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx$$

input

```
Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a - b*x^2]*Sqrt[c - d*x^2]*(e - f*x^2)
^(3/2)),x]
```

output

```
Integrate[(A + B*x^2 + C*x^4)/(Sqrt[a - b*x^2]*Sqrt[c - d*x^2]*(e - f*x^2)
^(3/2)), x]
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx$$

↓ 7293

$$\int \left(\frac{A}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} + \frac{Bx^2}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} + \frac{Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} \right) dx$$

$$\begin{array}{c}
 \downarrow \text{2009} \\
 A \int \frac{1}{\sqrt{a-bx^2}\sqrt{c-dx^2}(e-fx^2)^{3/2}} dx + B \int \frac{x^2}{\sqrt{a-bx^2}\sqrt{c-dx^2}(e-fx^2)^{3/2}} dx + \\
 C \int \frac{x^4}{\sqrt{a-bx^2}\sqrt{c-dx^2}(e-fx^2)^{3/2}} dx
 \end{array}$$

input `Int[(A + B*x^2 + C*x^4)/(Sqrt[a - b*x^2]*Sqrt[c - d*x^2]*(e - f*x^2)^(3/2)),x]`

output `$Aborted`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

Maple [F]

$$\int \frac{C x^4 + x^2 B + A}{\sqrt{-b x^2 + a} \sqrt{-x^2 d + c} (-f x^2 + e)^{\frac{3}{2}}} dx$$

input `int((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

output `int((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)`

Fricas [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{-bx^2 + a}\sqrt{-dx^2 + c}(-fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, algorithm="fricas")`

output `integral((C*x^4 + B*x^2 + A)*sqrt(-b*x^2 + a)*sqrt(-d*x^2 + c)*sqrt(-f*x^2 + e)/(b*d*f^2*x^8 - (2*b*d*e*f + (b*c + a*d)*f^2)*x^6 + (b*d*e^2 + a*c*f^2 + 2*(b*c + a*d)*e*f)*x^4 + a*c*e^2 - (2*a*c*e*f + (b*c + a*d)*e^2)*x^2), x)`

Sympy [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{\frac{3}{2}}} dx$$

input `integrate((C*x**4+B*x**2+A)/(-b*x**2+a)**(1/2)/(-d*x**2+c)**(1/2)/(-f*x**2+e)**(3/2),x)`

output `Integral((A + B*x**2 + C*x**4)/(sqrt(a - b*x**2)*sqrt(c - d*x**2)*(e - f*x**2)**(3/2)), x)`

Maxima [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{-bx^2 + a}\sqrt{-dx^2 + c}(-fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, algorithm="maxima")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(-b*x^2 + a)*sqrt(-d*x^2 + c)*(-f*x^2 + e)^(3/2)), x)`

Giac [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{-bx^2 + a}\sqrt{-dx^2 + c}(-fx^2 + e)^{\frac{3}{2}}} dx$$

input `integrate((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x, algorithm="giac")`

output `integrate((C*x^4 + B*x^2 + A)/(sqrt(-b*x^2 + a)*sqrt(-d*x^2 + c)*(-f*x^2 + e)^(3/2)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \int \frac{Cx^4 + Bx^2 + A}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx$$

input `int((A + B*x^2 + C*x^4)/((a - b*x^2)^(1/2)*(c - d*x^2)^(1/2)*(e - f*x^2)^(3/2)),x)`

output `int((A + B*x^2 + C*x^4)/((a - b*x^2)^(1/2)*(c - d*x^2)^(1/2)*(e - f*x^2)^(3/2)), x)`

Reduce [F]

$$\int \frac{A + Bx^2 + Cx^4}{\sqrt{a - bx^2}\sqrt{c - dx^2}(e - fx^2)^{3/2}} dx = \left(\int \frac{\sqrt{-fx^2 + e}\sqrt{-dx^2 + c}\sqrt{-bx^2 + a}}{bd f^2 x^8 - ad f^2 x^6 - bc f^2 x^6 - 2bdef x^6 + ac f^2 x^4 + 2adef x^4} \right.$$

$$+ \left(\int \frac{\sqrt{-fx^2 + e}\sqrt{-dx^2 + c}\sqrt{-bx^2 + a} x^2}{bd f^2 x^8 - ad f^2 x^6 - bc f^2 x^6 - 2bdef x^6 + ac f^2 x^4 + 2adef x^4 + 2bcef x^4 + bd e^2 x^4 - 2acef x^2 - ad} \right.$$

$$+ \left(\int \frac{\sqrt{-fx^2 + e}\sqrt{-dx^2 + c}\sqrt{-bx^2 + a}}{bd f^2 x^8 - ad f^2 x^6 - bc f^2 x^6 - 2bdef x^6 + ac f^2 x^4 + 2adef x^4 + 2bcef x^4 + bd e^2 x^4 - 2acef x^2 - ad} \right)$$

input

```
int((C*x^4+B*x^2+A)/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)/(-f*x^2+e)^(3/2),x)
```

output

```
int((sqrt(e - f*x**2)*sqrt(c - d*x**2)*sqrt(a - b*x**2)*x**4)/(a*c*e**2 -
2*a*c*e*f*x**2 + a*c*f**2*x**4 - a*d*e**2*x**2 + 2*a*d*e*f*x**4 - a*d*f**2
*x**6 - b*c*e**2*x**2 + 2*b*c*e*f*x**4 - b*c*f**2*x**6 + b*d*e**2*x**4 - 2
*b*d*e*f*x**6 + b*d*f**2*x**8),x)*c + int((sqrt(e - f*x**2)*sqrt(c - d*x**
2)*sqrt(a - b*x**2)*x**2)/(a*c*e**2 - 2*a*c*e*f*x**2 + a*c*f**2*x**4 - a*d
*e**2*x**2 + 2*a*d*e*f*x**4 - a*d*f**2*x**6 - b*c*e**2*x**2 + 2*b*c*e*f*x
**4 - b*c*f**2*x**6 + b*d*e**2*x**4 - 2*b*d*e*f*x**6 + b*d*f**2*x**8),x)*b
+ int((sqrt(e - f*x**2)*sqrt(c - d*x**2)*sqrt(a - b*x**2))/(a*c*e**2 - 2*a
*c*e*f*x**2 + a*c*f**2*x**4 - a*d*e**2*x**2 + 2*a*d*e*f*x**4 - a*d*f**2*x
**6 - b*c*e**2*x**2 + 2*b*c*e*f*x**4 - b*c*f**2*x**6 + b*d*e**2*x**4 - 2*b*
d*e*f*x**6 + b*d*f**2*x**8),x)*a
```

CHAPTER 4

APPENDIX

4.1	Listing of Grading functions	366
4.2	Links to plain text integration problems used in this report for each CAS .	384

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
  ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "
  ,

```



```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]===Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]===Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]===Plus || Head[expn]===Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```

```
Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
If[AppellFunctionQ[Head[expn]],
Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
If[Head[expn]===RootSum,
Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
If[Head[expn]===Integrate || Head[expn]===Int,
Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
9]]]]]]]]]]
```

```
ElementaryFunctionQ[func_] :=
MemberQ[{
Exp, Log,
Sin, Cos, Tan, Cot, Sec, Csc,
ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
Sinh, Cosh, Tanh, Coth, Sech, Csch,
ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
}, func]
```

```
SpecialFunctionQ[func_] :=
MemberQ[{
Erf, Erfc, Erfi,
FresnelS, FresnelC,
ExpIntegralE, ExpIntegralEi, LogIntegral,
SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
Gamma, LogGamma, PolyGamma,
Zeta, PolyLog, ProductLog,
EllipticF, EllipticE, EllipticPi
}, func]
```

```
HypergeometricFunctionQ[func_] :=
MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]
```

```
AppellFunctionQ[func_] :=
MemberQ[{AppellF1}, func]
```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

#   antiderivative
#   "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co

        fi;
    fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                    convert(ExpnType_result,string)," vs. order ",
                    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```



```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>Enter expnType, expn=", expn)
        print (">>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```


4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file