

Computer Algebra Independent Integration Tests

Summer 2024

1-Algebraic-functions/1.1-Binomial/1.1.3-General-
binomial/1.1.3.1/40-1.1.3.1-a

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 6:01pm

Contents

1	Introduction	4
1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Time and leaf size Performance	10
1.4	Performance based on number of rules Rubi used	12
1.5	Performance based on number of steps Rubi used	13
1.6	Solved integrals histogram based on leaf size of result	14
1.7	Solved integrals histogram based on CPU time used	15
1.8	Leaf size vs. CPU time used	16
1.9	list of integrals with no known antiderivative	17
1.10	List of integrals solved by CAS but has no known antiderivative	17
1.11	list of integrals solved by CAS but failed verification	17
1.12	Timing	18
1.13	Verification	18
1.14	Important notes about some of the results	19
1.15	Current tree layout of integration tests	22
1.16	Design of the test system	23
2	detailed summary tables of results	24
2.1	List of integrals sorted by grade for each CAS	25
2.2	Detailed conclusion table per each integral for all CAS systems	29
2.3	Detailed conclusion table specific for Rubi results	40
3	Listing of integrals	42
3.1	$\int (a + bx^3)^5 dx$	44
3.2	$\int (a + bx^3)^4 dx$	49
3.3	$\int (a + bx^3)^3 dx$	54
3.4	$\int (a + bx^3)^2 dx$	59
3.5	$\int (a + bx^3) dx$	64
3.6	$\int \frac{1}{a+bx^3} dx$	69
3.7	$\int \frac{1}{(a+bx^3)^2} dx$	77

3.8	$\int \frac{1}{(a+bx^3)^3} dx$	87
3.9	$\int \frac{1}{1+a+bx^3} dx$	99
3.10	$\int \frac{1}{1+a-bx^3} dx$	108
3.11	$\int \frac{1}{-1+a+bx^3} dx$	117
3.12	$\int \frac{1}{-1+a-bx^3} dx$	125
3.13	$\int (a+bx^3)^{5/2} dx$	134
3.14	$\int (a+bx^3)^{3/2} dx$	141
3.15	$\int \sqrt{a+bx^3} dx$	148
3.16	$\int \frac{1}{\sqrt{a+bx^3}} dx$	154
3.17	$\int \frac{1}{(a+bx^3)^{3/2}} dx$	160
3.18	$\int \frac{1}{(a+bx^3)^{5/2}} dx$	167
3.19	$\int \frac{1}{(a+bx^3)^{7/2}} dx$	174
3.20	$\int \frac{1}{\sqrt{1+x^3}} dx$	181
3.21	$\int \frac{1}{\sqrt{1-x^3}} dx$	186
3.22	$\int \frac{1}{\sqrt{-1+x^3}} dx$	192
3.23	$\int \frac{1}{\sqrt{-1-x^3}} dx$	198
3.24	$\int (a+bx^3)^{5/3} dx$	204
3.25	$\int (a+bx^3)^{2/3} dx$	211
3.26	$\int \frac{1}{\sqrt[3]{a+bx^3}} dx$	218
3.27	$\int \frac{1}{(a+bx^3)^{4/3}} dx$	224
3.28	$\int \frac{1}{(a+bx^3)^{7/3}} dx$	229
3.29	$\int \frac{1}{(a+bx^3)^{10/3}} dx$	234
3.30	$\int \frac{1}{(a+bx^3)^{13/3}} dx$	240
3.31	$\int (a+bx^3)^{4/3} dx$	246
3.32	$\int \sqrt[3]{a+bx^3} dx$	252
3.33	$\int \frac{1}{(a+bx^3)^{2/3}} dx$	257
3.34	$\int \frac{1}{(a+bx^3)^{5/3}} dx$	262
3.35	$\int \frac{1}{\sqrt[3]{a-bx^3}} dx$	267
3.36	$\int \frac{1}{\sqrt[3]{2+x^3}} dx$	273
3.37	$\int (a+bx^3)^{3/4} dx$	278
3.38	$\int \sqrt[4]{a+bx^3} dx$	283
3.39	$\int \frac{1}{\sqrt[4]{a+bx^3}} dx$	288
3.40	$\int \frac{1}{(a+bx^3)^{3/4}} dx$	293
3.41	$\int \frac{1}{(a+bx^3)^{5/4}} dx$	298

4 Appendix	303
4.1 Listing of Grading functions	303
4.2 Links to plain text integration problems used in this report for each CAS321	

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Time and leaf size Performance	10
1.4	Performance based on number of rules Rubi used	12
1.5	Performance based on number of steps Rubi used	13
1.6	Solved integrals histogram based on leaf size of result	14
1.7	Solved integrals histogram based on CPU time used	15
1.8	Leaf size vs. CPU time used	16
1.9	list of integrals with no known antiderivative	17
1.10	List of integrals solved by CAS but has no known antiderivative	17
1.11	list of integrals solved by CAS but failed verification	17
1.12	Timing	18
1.13	Verification	18
1.14	Important notes about some of the results	19
1.15	Current tree layout of integration tests	22
1.16	Design of the test system	23

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [41]. This is test number [40].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (41)	0.00 (0)
Mathematica	100.00 (41)	0.00 (0)
Mupad	100.00 (41)	0.00 (0)
Sympy	100.00 (41)	0.00 (0)
Maple	78.05 (32)	21.95 (9)
Fricas	78.05 (32)	21.95 (9)
Maxima	46.34 (19)	53.66 (22)
Giac	29.27 (12)	70.73 (29)
Reduce	29.27 (12)	70.73 (29)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Fricas	58.537	19.512	0.000	21.951
Sympy	56.098	9.756	34.146	0.000
Maple	48.780	0.000	29.268	21.951
Maxima	43.902	2.439	0.000	53.659
Mathematica	36.585	0.000	63.415	0.000
Giac	29.268	0.000	0.000	70.732
Mupad	0.000	100.000	0.000	0.000
Reduce	0.000	29.268	0.000	70.732

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Mupad	0	0.00	0.00	0.00
Sympy	0	0.00	0.00	0.00
Fricas	9	100.00	0.00	0.00
Maple	9	100.00	0.00	0.00
Maxima	22	90.91	0.00	9.09
Giac	29	100.00	0.00	0.00
Reduce	29	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.07
Fricas	0.07
Mupad	0.11
Giac	0.12
Rubi	0.20
Reduce	0.23
Sympy	0.44
Maple	0.71
Mathematica	7.48

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mupad	65.61	0.75	37.00	0.80
Maxima	78.79	1.03	69.00	0.88
Reduce	83.50	0.90	72.00	0.93
Sympy	85.88	1.27	36.00	0.51
Giac	90.67	0.94	119.50	0.93
Maple	103.22	0.77	40.00	0.87
Rubi	105.27	1.01	92.00	1.00
Fricas	165.25	1.62	63.50	1.03
Mathematica	1107.10	18.42	110.00	1.00

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram shows that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {25, 27, 28, 29, 30, 31, 32, 33, 34, 37, 38, 39, 40, 41}

Maple {22}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand,the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	25
2.2	Detailed conclusion table per each integral for all CAS systems	29
2.3	Detailed conclusion table specific for Rubi results	40

2.1 List of integrals sorted by grade for each CAS

Rubi	25
Mma	25
Maple	26
Fricas	26
Maxima	26
Giac	27
Mupad	27
Sympy	27
Reduce	28

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 26, 35, 36 }

B grade { }

C grade { 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 27, 28, 29, 30, 31, 32, 33, 34, 37, 38, 39, 40, 41 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 5, 13, 14, 15, 16, 17, 18, 19, 24, 25, 26, 27, 28, 29, 30, 35 }

B grade { }

C grade { 6, 7, 8, 9, 10, 11, 12, 20, 21, 22, 23, 36 }

F normal fail { 31, 32, 33, 34, 37, 38, 39, 40, 41 }

F(-1) timeout fail { }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 3, 4, 5, 6, 7, 11, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 27, 28, 29, 30, 36 }

B grade { 8, 9, 10, 12, 24, 25, 26, 35 }

C grade { }

F normal fail { 31, 32, 33, 34, 37, 38, 39, 40, 41 }

F(-1) timeout fail { }

F(-2) exception fail { }

Maxima

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 25, 26, 27, 28, 29, 30, 35, 36 }

B grade { 24 }

C grade { }

F normal fail { 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 31, 32, 33, 34, 37, 38, 39, 40, 41 }

F(-1) timeout fail { }

F(-2) exception fail { 11, 12 }

Giac

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 }

B grade { }

C grade { }

F normal fail { 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41 }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Sympy

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23 }

B grade { 27, 28, 29, 30 }

C grade { 24, 25, 26, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Reduce

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 }

C grade { }

F normal fail { 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33,
34, 35, 36, 37, 38, 39, 40, 41 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	61	54	53	53	60	53	57	53
N.S.	1	1.00	1.00	0.89	0.87	0.87	0.98	0.87	0.93	0.87
time (sec)	N/A	0.350	0.038	0.345	0.035	0.070	0.023	0.117	0.210	0.132

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	48	48	43	42	42	46	42	46	42
N.S.	1	1.00	1.00	0.90	0.88	0.88	0.96	0.88	0.96	0.88
time (sec)	N/A	0.169	0.001	0.355	0.026	0.062	0.020	0.122	0.234	0.015

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	38	33	32	32	36	32	35	32
N.S.	1	1.00	1.00	0.87	0.84	0.84	0.95	0.84	0.92	0.84
time (sec)	N/A	0.161	0.001	0.339	0.025	0.057	0.019	0.128	0.236	0.026

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	25	22	21	21	20	21	24	21
N.S.	1	1.00	1.00	0.88	0.84	0.84	0.80	0.84	0.96	0.84
time (sec)	N/A	0.149	0.001	0.348	0.029	0.061	0.021	0.111	0.230	0.037

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	12	11	10	10	8	10	12	10
N.S.	1	1.00	1.00	0.92	0.83	0.83	0.67	0.83	1.00	0.83
time (sec)	N/A	0.136	0.000	0.037	0.025	0.059	0.016	0.119	0.205	0.034

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	111	89	27	98	299	20	112	67	99
N.S.	1	0.97	0.77	0.23	0.85	2.60	0.17	0.97	0.58	0.86
time (sec)	N/A	0.370	0.035	0.464	0.110	0.083	0.093	0.124	0.264	0.259

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	134	136	118	46	122	389	39	127	171	128
N.S.	1	1.01	0.88	0.34	0.91	2.90	0.29	0.95	1.28	0.96
time (sec)	N/A	0.269	0.040	0.430	0.107	0.081	0.128	0.120	0.213	0.147

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	151	161	135	57	145	499	63	137	278	142
N.S.	1	1.07	0.89	0.38	0.96	3.30	0.42	0.91	1.84	0.94
time (sec)	N/A	0.295	0.036	0.457	0.129	0.081	0.199	0.128	0.230	0.148

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	125	123	101	28	114	446	32	143	79	137
N.S.	1	0.98	0.81	0.22	0.91	3.57	0.26	1.14	0.63	1.10
time (sec)	N/A	0.275	0.020	0.461	0.109	0.088	0.130	0.125	0.242	0.208

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	124	122	124	30	113	467	34	131	77	165
N.S.	1	0.98	1.00	0.24	0.91	3.77	0.27	1.06	0.62	1.33
time (sec)	N/A	0.256	0.047	0.439	0.110	0.096	0.149	0.119	0.241	0.296

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F(-2)	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	139	134	101	28	0	446	32	142	79	137
N.S.	1	0.96	0.73	0.20	0.00	3.21	0.23	1.02	0.57	0.99
time (sec)	N/A	0.288	0.022	0.463	0.000	0.081	0.138	0.119	0.252	0.221

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F(-2)	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	138	135	124	30	0	467	34	138	77	165
N.S.	1	0.98	0.90	0.22	0.00	3.38	0.25	1.00	0.56	1.20
time (sec)	N/A	0.282	0.030	0.439	0.000	0.083	0.156	0.125	0.203	0.278

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	265	271	49	320	0	58	37	0	71	37
N.S.	1	1.02	0.18	1.21	0.00	0.22	0.14	0.00	0.27	0.14
time (sec)	N/A	0.290	4.394	2.528	0.000	0.073	0.543	0.000	0.261	0.142

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	246	249	47	309	0	47	37	0	52	37
N.S.	1	1.01	0.19	1.26	0.00	0.19	0.15	0.00	0.21	0.15
time (sec)	N/A	0.243	3.817	0.500	0.000	0.070	0.667	0.000	0.265	0.077

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	227	227	46	297	0	33	37	0	35	37
N.S.	1	1.00	0.20	1.31	0.00	0.15	0.16	0.00	0.15	0.16
time (sec)	N/A	0.228	3.015	0.472	0.000	0.071	0.461	0.000	0.234	0.072

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	207	207	46	283	0	14	36	0	20	37
N.S.	1	1.00	0.22	1.37	0.00	0.07	0.17	0.00	0.10	0.18
time (sec)	N/A	0.213	10.014	0.469	0.000	0.068	0.424	0.000	0.233	0.076

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	232	232	55	306	0	51	36	0	31	37
N.S.	1	1.00	0.24	1.32	0.00	0.22	0.16	0.00	0.13	0.16
time (sec)	N/A	0.230	3.328	0.440	0.000	0.068	0.487	0.000	0.218	0.110

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	251	259	72	335	0	88	36	0	42	37
N.S.	1	1.03	0.29	1.33	0.00	0.35	0.14	0.00	0.17	0.15
time (sec)	N/A	0.257	4.155	0.435	0.000	0.070	0.517	0.000	0.224	0.102

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	270	286	85	364	0	121	36	0	53	37
N.S.	1	1.06	0.31	1.35	0.00	0.45	0.13	0.00	0.20	0.14
time (sec)	N/A	0.283	4.700	0.451	0.000	0.076	0.661	0.000	0.219	0.131

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	103	103	17	14	0	6	27	0	16	155
N.S.	1	1.00	0.17	0.14	0.00	0.06	0.26	0.00	0.16	1.50
time (sec)	N/A	0.178	10.018	0.908	0.000	0.064	0.342	0.000	0.240	0.155

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	115	15	12	0	6	29	0	20	172
N.S.	1	1.00	0.13	0.10	0.00	0.05	0.25	0.00	0.17	1.50
time (sec)	N/A	0.181	10.018	0.825	0.000	0.087	0.366	0.000	0.257	0.143

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	A	F	F	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	120	120	35	30	0	6	26	0	16	156
N.S.	1	1.00	0.29	0.25	0.00	0.05	0.22	0.00	0.13	1.30
time (sec)	N/A	0.193	10.020	0.903	0.000	0.070	0.366	0.000	0.243	0.064

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	112	112	37	16	0	6	31	0	19	172
N.S.	1	1.00	0.33	0.14	0.00	0.05	0.28	0.00	0.17	1.54
time (sec)	N/A	0.182	10.020	0.680	0.000	0.077	0.344	0.000	0.249	0.026

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	B	B	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	112	114	272	144	179	399	37	0	45	37
N.S.	1	1.02	2.43	1.29	1.60	3.56	0.33	0.00	0.40	0.33
time (sec)	N/A	0.188	0.171	2.274	0.123	0.091	0.968	0.000	0.219	0.077

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	C	F	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	92	203	118	134	362	37	0	27	37
N.S.	1	1.01	2.23	1.30	1.47	3.98	0.41	0.00	0.30	0.41
time (sec)	N/A	0.171	0.087	0.641	0.116	0.078	0.656	0.000	0.240	0.067

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	110	99	100	330	36	0	11	37
N.S.	1	1.00	1.57	1.41	1.43	4.71	0.51	0.00	0.16	0.53
time (sec)	N/A	0.156	0.041	0.498	0.112	0.078	0.558	0.000	0.219	0.083

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	F	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	524	15	14	23	29	0	30	14
N.S.	1	1.00	32.75	0.94	0.88	1.44	1.81	0.00	1.88	0.88
time (sec)	N/A	0.132	3.924	0.489	0.027	0.074	0.376	0.000	0.244	0.071

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	F	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	5268	26	31	47	126	0	50	28
N.S.	1	1.00	135.08	0.67	0.79	1.21	3.23	0.00	1.28	0.72
time (sec)	N/A	0.150	16.910	0.479	0.031	0.068	0.594	0.000	0.243	0.065

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	F	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	66	11702	37	50	69	515	0	70	44
N.S.	1	1.14	201.76	0.64	0.86	1.19	8.88	0.00	1.21	0.76
time (sec)	N/A	0.170	55.029	0.484	0.026	0.070	0.992	0.000	0.207	0.078

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	F	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	93	21879	48	67	91	1550	0	90	61
N.S.	1	1.21	284.14	0.62	0.87	1.18	20.13	0.00	1.17	0.79
time (sec)	N/A	0.190	144.932	0.496	0.039	0.075	1.675	0.000	0.266	0.085

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	C	F	F	B
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	47	47	272	0	0	0	37	0	45	37
N.S.	1	1.00	5.79	0.00	0.00	0.00	0.79	0.00	0.96	0.79
time (sec)	N/A	0.157	0.211	0.000	0.000	0.000	0.544	0.000	0.258	0.082

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	C	F	F	B
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	203	0	0	0	37	0	27	37
N.S.	1	1.00	4.41	0.00	0.00	0.00	0.80	0.00	0.59	0.80
time (sec)	N/A	0.159	0.134	0.000	0.000	0.000	0.465	0.000	0.237	0.062

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	C	F	F	B
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	183	0	0	0	36	0	11	37
N.S.	1	1.00	3.98	0.00	0.00	0.00	0.78	0.00	0.24	0.80
time (sec)	N/A	0.157	0.159	0.000	0.000	0.000	0.448	0.000	0.261	0.078

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	C	F	F	B
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	2028	0	0	0	36	0	30	37
N.S.	1	1.00	41.39	0.00	0.00	0.00	0.73	0.00	0.61	0.76
time (sec)	N/A	0.153	10.785	0.000	0.000	0.000	0.662	0.000	0.210	0.113

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	116	103	103	284	37	0	12	38
N.S.	1	1.00	1.61	1.43	1.43	3.94	0.51	0.00	0.17	0.53
time (sec)	N/A	0.156	0.051	1.858	0.107	0.074	0.517	0.000	0.242	0.113

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	78	18	69	76	34	0	9	16
N.S.	1	1.00	1.70	0.39	1.50	1.65	0.74	0.00	0.20	0.35
time (sec)	N/A	0.147	0.023	2.256	0.107	0.076	0.431	0.000	0.231	0.091

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	C	F	F	B
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	203	0	0	0	37	0	27	37
N.S.	1	1.00	4.41	0.00	0.00	0.00	0.80	0.00	0.59	0.80
time (sec)	N/A	0.153	0.125	0.000	0.000	0.000	0.638	0.000	0.252	0.096

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	C	F	F	B
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	203	0	0	0	37	0	27	37
N.S.	1	1.00	4.41	0.00	0.00	0.00	0.80	0.00	0.59	0.80
time (sec)	N/A	0.154	0.126	0.000	0.000	0.000	0.480	0.000	0.258	0.079

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	C	F	F	B
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	203	0	0	0	36	0	11	37
N.S.	1	1.00	4.41	0.00	0.00	0.00	0.78	0.00	0.24	0.80
time (sec)	N/A	0.158	0.098	0.000	0.000	0.000	0.432	0.000	0.231	0.093

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	C	F	F	B
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	201	0	0	0	36	0	11	37
N.S.	1	1.00	4.37	0.00	0.00	0.00	0.78	0.00	0.24	0.80
time (sec)	N/A	0.154	0.109	0.000	0.000	0.000	0.445	0.000	0.274	0.093

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	C	F	F	B
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	263	0	0	0	36	0	30	37
N.S.	1	1.00	5.37	0.00	0.00	0.00	0.73	0.00	0.61	0.76
time (sec)	N/A	0.156	0.150	0.000	0.000	0.000	0.693	0.000	0.237	0.114

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [8] had the largest ratio of [1.11111000000000004]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	2	2	1.00	9	0.222
2	A	2	2	1.00	9	0.222
3	A	2	2	1.00	9	0.222
4	A	2	2	1.00	9	0.222
5	A	1	1	1.00	7	0.143
6	A	9	8	0.97	9	0.889
7	A	10	9	1.01	9	1.000
8	A	11	10	1.07	9	1.111
9	A	9	8	0.98	10	0.800
10	A	8	7	0.98	11	0.636
11	A	9	8	0.96	10	0.800
12	A	10	9	0.98	11	0.818
13	A	4	4	1.02	11	0.364
14	A	3	3	1.01	11	0.273
15	A	2	2	1.00	11	0.182
16	A	1	1	1.00	11	0.091
17	A	2	2	1.00	11	0.182
18	A	3	3	1.03	11	0.273
19	A	4	4	1.06	11	0.364
20	A	1	1	1.00	9	0.111
21	A	1	1	1.00	11	0.091

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	1	1	1.00	9	0.111
23	A	1	1	1.00	11	0.091
24	A	3	3	1.02	11	0.273
25	A	2	2	1.01	11	0.182
26	A	1	1	1.00	11	0.091
27	A	1	1	1.00	11	0.091
28	A	2	2	1.00	11	0.182
29	A	3	3	1.14	11	0.273
30	A	4	4	1.21	11	0.364
31	A	2	2	1.00	11	0.182
32	A	2	2	1.00	11	0.182
33	A	2	2	1.00	11	0.182
34	A	2	2	1.00	11	0.182
35	A	1	1	1.00	12	0.083
36	A	1	1	1.00	9	0.111
37	A	2	2	1.00	11	0.182
38	A	2	2	1.00	11	0.182
39	A	2	2	1.00	11	0.182
40	A	2	2	1.00	11	0.182
41	A	2	2	1.00	11	0.182

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int (a + bx^3)^5 dx$	44
3.2	$\int (a + bx^3)^4 dx$	49
3.3	$\int (a + bx^3)^3 dx$	54
3.4	$\int (a + bx^3)^2 dx$	59
3.5	$\int (a + bx^3) dx$	64
3.6	$\int \frac{1}{a+bx^3} dx$	69
3.7	$\int \frac{1}{(a+bx^3)^2} dx$	77
3.8	$\int \frac{1}{(a+bx^3)^3} dx$	87
3.9	$\int \frac{1}{1+a+bx^3} dx$	99
3.10	$\int \frac{1}{1+a-bx^3} dx$	108
3.11	$\int \frac{1}{-1+a+bx^3} dx$	117
3.12	$\int \frac{1}{-1+a-bx^3} dx$	125
3.13	$\int (a + bx^3)^{5/2} dx$	134
3.14	$\int (a + bx^3)^{3/2} dx$	141
3.15	$\int \sqrt{a + bx^3} dx$	148
3.16	$\int \frac{1}{\sqrt{a+bx^3}} dx$	154
3.17	$\int \frac{1}{(a+bx^3)^{3/2}} dx$	160
3.18	$\int \frac{1}{(a+bx^3)^{5/2}} dx$	167
3.19	$\int \frac{1}{(a+bx^3)^{7/2}} dx$	174
3.20	$\int \frac{1}{\sqrt{1+x^3}} dx$	181
3.21	$\int \frac{1}{\sqrt{1-x^3}} dx$	186
3.22	$\int \frac{1}{\sqrt{-1+x^3}} dx$	192
3.23	$\int \frac{1}{\sqrt{-1-x^3}} dx$	198
3.24	$\int (a + bx^3)^{5/3} dx$	204
3.25	$\int (a + bx^3)^{2/3} dx$	211

3.26	$\int \frac{1}{\sqrt[3]{a+bx^3}} dx$	218
3.27	$\int \frac{1}{(a+bx^3)^{4/3}} dx$	224
3.28	$\int \frac{1}{(a+bx^3)^{7/3}} dx$	229
3.29	$\int \frac{1}{(a+bx^3)^{10/3}} dx$	234
3.30	$\int \frac{1}{(a+bx^3)^{13/3}} dx$	240
3.31	$\int (a+bx^3)^{4/3} dx$	246
3.32	$\int \sqrt[3]{a+bx^3} dx$	252
3.33	$\int \frac{1}{(a+bx^3)^{2/3}} dx$	257
3.34	$\int \frac{1}{(a+bx^3)^{5/3}} dx$	262
3.35	$\int \frac{1}{\sqrt[3]{a-bx^3}} dx$	267
3.36	$\int \frac{1}{\sqrt[3]{2+x^3}} dx$	273
3.37	$\int (a+bx^3)^{3/4} dx$	278
3.38	$\int \sqrt[4]{a+bx^3} dx$	283
3.39	$\int \frac{1}{\sqrt[4]{a+bx^3}} dx$	288
3.40	$\int \frac{1}{(a+bx^3)^{3/4}} dx$	293
3.41	$\int \frac{1}{(a+bx^3)^{5/4}} dx$	298

3.1 $\int (a + bx^3)^5 dx$

Optimal result	44
Mathematica [A] (verified)	44
Rubi [A] (verified)	45
Maple [A] (verified)	46
Fricas [A] (verification not implemented)	46
Sympy [A] (verification not implemented)	47
Maxima [A] (verification not implemented)	47
Giac [A] (verification not implemented)	47
Mupad [B] (verification not implemented)	48
Reduce [B] (verification not implemented)	48

Optimal result

Integrand size = 9, antiderivative size = 61

$$\int (a + bx^3)^5 dx = a^5x + \frac{5}{4}a^4bx^4 + \frac{10}{7}a^3b^2x^7 + a^2b^3x^{10} + \frac{5}{13}ab^4x^{13} + \frac{b^5x^{16}}{16}$$

output

```
a^5*x+5/4*a^4*b*x^4+10/7*a^3*b^2*x^7+a^2*b^3*x^10+5/13*a*b^4*x^13+1/16*b^5*x^16
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00

$$\int (a + bx^3)^5 dx = a^5x + \frac{5}{4}a^4bx^4 + \frac{10}{7}a^3b^2x^7 + a^2b^3x^{10} + \frac{5}{13}ab^4x^{13} + \frac{b^5x^{16}}{16}$$

input

```
Integrate[(a + b*x^3)^5,x]
```

output

```
a^5*x + (5*a^4*b*x^4)/4 + (10*a^3*b^2*x^7)/7 + a^2*b^3*x^10 + (5*a*b^4*x^13)/13 + (b^5*x^16)/16
```

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {747, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx^3)^5 dx$$

$$\downarrow 747$$

$$\int (a^5 + 5a^4bx^3 + 10a^3b^2x^6 + 10a^2b^3x^9 + 5ab^4x^{12} + b^5x^{15}) dx$$

$$\downarrow 2009$$

$$a^5x + \frac{5}{4}a^4bx^4 + \frac{10}{7}a^3b^2x^7 + a^2b^3x^{10} + \frac{5}{13}ab^4x^{13} + \frac{b^5x^{16}}{16}$$

input

```
Int[(a + b*x^3)^5, x]
```

output

```
a^5*x + (5*a^4*b*x^4)/4 + (10*a^3*b^2*x^7)/7 + a^2*b^3*x^10 + (5*a*b^4*x^13)/13 + (b^5*x^16)/16
```

Defintions of rubi rules used

rule 747

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*x^n)^p, x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && IGtQ[p, 0]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.89

method	result	size
gosper	$a^5x + \frac{5}{4}a^4bx^4 + \frac{10}{7}a^3b^2x^7 + a^2b^3x^{10} + \frac{5}{13}ab^4x^{13} + \frac{1}{16}b^5x^{16}$	54
default	$a^5x + \frac{5}{4}a^4bx^4 + \frac{10}{7}a^3b^2x^7 + a^2b^3x^{10} + \frac{5}{13}ab^4x^{13} + \frac{1}{16}b^5x^{16}$	54
norman	$a^5x + \frac{5}{4}a^4bx^4 + \frac{10}{7}a^3b^2x^7 + a^2b^3x^{10} + \frac{5}{13}ab^4x^{13} + \frac{1}{16}b^5x^{16}$	54
risch	$a^5x + \frac{5}{4}a^4bx^4 + \frac{10}{7}a^3b^2x^7 + a^2b^3x^{10} + \frac{5}{13}ab^4x^{13} + \frac{1}{16}b^5x^{16}$	54
parallelrisch	$a^5x + \frac{5}{4}a^4bx^4 + \frac{10}{7}a^3b^2x^7 + a^2b^3x^{10} + \frac{5}{13}ab^4x^{13} + \frac{1}{16}b^5x^{16}$	54
orering	$\frac{x(91b^5x^{15} + 560ab^4x^{12} + 1456a^2b^3x^9 + 2080a^3b^2x^6 + 1820a^4bx^3 + 1456a^5)}{1456}$	58

input `int((b*x^3+a)^5,x,method=_RETURNVERBOSE)`output `a^5*x+5/4*a^4*b*x^4+10/7*a^3*b^2*x^7+a^2*b^3*x^10+5/13*a*b^4*x^13+1/16*b^5*x^16`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.87

$$\int (a + bx^3)^5 dx = \frac{1}{16} b^5 x^{16} + \frac{5}{13} ab^4 x^{13} + a^2 b^3 x^{10} + \frac{10}{7} a^3 b^2 x^7 + \frac{5}{4} a^4 b x^4 + a^5 x$$

input `integrate((b*x^3+a)^5,x, algorithm="fricas")`output `1/16*b^5*x^16 + 5/13*a*b^4*x^13 + a^2*b^3*x^10 + 10/7*a^3*b^2*x^7 + 5/4*a^4*b*x^4 + a^5*x`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.98

$$\int (a + bx^3)^5 dx = a^5x + \frac{5a^4bx^4}{4} + \frac{10a^3b^2x^7}{7} + a^2b^3x^{10} + \frac{5ab^4x^{13}}{13} + \frac{b^5x^{16}}{16}$$

input `integrate((b*x**3+a)**5,x)`output `a**5*x + 5*a**4*b*x**4/4 + 10*a**3*b**2*x**7/7 + a**2*b**3*x**10 + 5*a*b**4*x**13/13 + b**5*x**16/16`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.87

$$\int (a + bx^3)^5 dx = \frac{1}{16} b^5x^{16} + \frac{5}{13} ab^4x^{13} + a^2b^3x^{10} + \frac{10}{7} a^3b^2x^7 + \frac{5}{4} a^4bx^4 + a^5x$$

input `integrate((b*x^3+a)^5,x, algorithm="maxima")`output `1/16*b^5*x^16 + 5/13*a*b^4*x^13 + a^2*b^3*x^10 + 10/7*a^3*b^2*x^7 + 5/4*a^4*b*x^4 + a^5*x`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.87

$$\int (a + bx^3)^5 dx = \frac{1}{16} b^5x^{16} + \frac{5}{13} ab^4x^{13} + a^2b^3x^{10} + \frac{10}{7} a^3b^2x^7 + \frac{5}{4} a^4bx^4 + a^5x$$

input `integrate((b*x^3+a)^5,x, algorithm="giac")`output `1/16*b^5*x^16 + 5/13*a*b^4*x^13 + a^2*b^3*x^10 + 10/7*a^3*b^2*x^7 + 5/4*a^4*b*x^4 + a^5*x`

Mupad [B] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.87

$$\int (a + bx^3)^5 dx = a^5 x + \frac{5a^4 b x^4}{4} + \frac{10a^3 b^2 x^7}{7} + a^2 b^3 x^{10} + \frac{5a b^4 x^{13}}{13} + \frac{b^5 x^{16}}{16}$$

input `int((a + b*x^3)^5,x)`output `a^5*x + (b^5*x^16)/16 + (5*a^4*b*x^4)/4 + (5*a*b^4*x^13)/13 + (10*a^3*b^2*x^7)/7 + a^2*b^3*x^10`**Reduce [B] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.93

$$\int (a + bx^3)^5 dx = \frac{x(91b^5x^{15} + 560ab^4x^{12} + 1456a^2b^3x^9 + 2080a^3b^2x^6 + 1820a^4bx^3 + 1456a^5)}{1456}$$

input `int((b*x^3+a)^5,x)`output `(x*(1456*a**5 + 1820*a**4*b*x**3 + 2080*a**3*b**2*x**6 + 1456*a**2*b**3*x**9 + 560*a*b**4*x**12 + 91*b**5*x**15))/1456`

3.2 $\int (a + bx^3)^4 dx$

Optimal result	49
Mathematica [A] (verified)	49
Rubi [A] (verified)	50
Maple [A] (verified)	51
Fricas [A] (verification not implemented)	51
Sympy [A] (verification not implemented)	52
Maxima [A] (verification not implemented)	52
Giac [A] (verification not implemented)	52
Mupad [B] (verification not implemented)	53
Reduce [B] (verification not implemented)	53

Optimal result

Integrand size = 9, antiderivative size = 48

$$\int (a + bx^3)^4 dx = a^4x + a^3bx^4 + \frac{6}{7}a^2b^2x^7 + \frac{2}{5}ab^3x^{10} + \frac{b^4x^{13}}{13}$$

output

```
a^4*x+a^3*b*x^4+6/7*a^2*b^2*x^7+2/5*a*b^3*x^10+1/13*b^4*x^13
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00

$$\int (a + bx^3)^4 dx = a^4x + a^3bx^4 + \frac{6}{7}a^2b^2x^7 + \frac{2}{5}ab^3x^{10} + \frac{b^4x^{13}}{13}$$

input

```
Integrate[(a + b*x^3)^4,x]
```

output

```
a^4*x + a^3*b*x^4 + (6*a^2*b^2*x^7)/7 + (2*a*b^3*x^10)/5 + (b^4*x^13)/13
```

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {747, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx^3)^4 dx$$

$$\downarrow 747$$

$$\int (a^4 + 4a^3bx^3 + 6a^2b^2x^6 + 4ab^3x^9 + b^4x^{12}) dx$$

$$\downarrow 2009$$

$$a^4x + a^3bx^4 + \frac{6}{7}a^2b^2x^7 + \frac{2}{5}ab^3x^{10} + \frac{b^4x^{13}}{13}$$

input

```
Int[(a + b*x^3)^4,x]
```

output

```
a^4*x + a^3*b*x^4 + (6*a^2*b^2*x^7)/7 + (2*a*b^3*x^10)/5 + (b^4*x^13)/13
```

Defintions of rubi rules used

rule 747

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Int[ExpandIntegrand[(a + b*x^n)^p, x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && IGtQ[p, 0]
```

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.90

method	result	size
gospers	$a^4x + a^3bx^4 + \frac{6}{7}a^2b^2x^7 + \frac{2}{5}ab^3x^{10} + \frac{1}{13}b^4x^{13}$	43
default	$a^4x + a^3bx^4 + \frac{6}{7}a^2b^2x^7 + \frac{2}{5}ab^3x^{10} + \frac{1}{13}b^4x^{13}$	43
norman	$a^4x + a^3bx^4 + \frac{6}{7}a^2b^2x^7 + \frac{2}{5}ab^3x^{10} + \frac{1}{13}b^4x^{13}$	43
risch	$a^4x + a^3bx^4 + \frac{6}{7}a^2b^2x^7 + \frac{2}{5}ab^3x^{10} + \frac{1}{13}b^4x^{13}$	43
parallelrisch	$a^4x + a^3bx^4 + \frac{6}{7}a^2b^2x^7 + \frac{2}{5}ab^3x^{10} + \frac{1}{13}b^4x^{13}$	43
orering	$\frac{x(35b^4x^{12} + 182ab^3x^9 + 390a^2b^2x^6 + 455a^3bx^3 + 455a^4)}{455}$	47

input `int((b*x^3+a)^4,x,method=_RETURNVERBOSE)`output `a^4*x+a^3*b*x^4+6/7*a^2*b^2*x^7+2/5*a*b^3*x^10+1/13*b^4*x^13`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.88

$$\int (a + bx^3)^4 dx = \frac{1}{13} b^4 x^{13} + \frac{2}{5} ab^3 x^{10} + \frac{6}{7} a^2 b^2 x^7 + a^3 b x^4 + a^4 x$$

input `integrate((b*x^3+a)^4,x, algorithm="fricas")`output `1/13*b^4*x^13 + 2/5*a*b^3*x^10 + 6/7*a^2*b^2*x^7 + a^3*b*x^4 + a^4*x`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.96

$$\int (a + bx^3)^4 dx = a^4x + a^3bx^4 + \frac{6a^2b^2x^7}{7} + \frac{2ab^3x^{10}}{5} + \frac{b^4x^{13}}{13}$$

input `integrate((b*x**3+a)**4,x)`output `a**4*x + a**3*b*x**4 + 6*a**2*b**2*x**7/7 + 2*a*b**3*x**10/5 + b**4*x**13/13`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.88

$$\int (a + bx^3)^4 dx = \frac{1}{13} b^4 x^{13} + \frac{2}{5} ab^3 x^{10} + \frac{6}{7} a^2 b^2 x^7 + a^3 b x^4 + a^4 x$$

input `integrate((b*x^3+a)^4,x, algorithm="maxima")`output `1/13*b^4*x^13 + 2/5*a*b^3*x^10 + 6/7*a^2*b^2*x^7 + a^3*b*x^4 + a^4*x`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.88

$$\int (a + bx^3)^4 dx = \frac{1}{13} b^4 x^{13} + \frac{2}{5} ab^3 x^{10} + \frac{6}{7} a^2 b^2 x^7 + a^3 b x^4 + a^4 x$$

input `integrate((b*x^3+a)^4,x, algorithm="giac")`output `1/13*b^4*x^13 + 2/5*a*b^3*x^10 + 6/7*a^2*b^2*x^7 + a^3*b*x^4 + a^4*x`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.88

$$\int (a + bx^3)^4 dx = a^4 x + a^3 b x^4 + \frac{6 a^2 b^2 x^7}{7} + \frac{2 a b^3 x^{10}}{5} + \frac{b^4 x^{13}}{13}$$

input `int((a + b*x^3)^4,x)`output `a^4*x + (b^4*x^13)/13 + a^3*b*x^4 + (2*a*b^3*x^10)/5 + (6*a^2*b^2*x^7)/7`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.96

$$\int (a + bx^3)^4 dx = \frac{x(35b^4x^{12} + 182ab^3x^9 + 390a^2b^2x^6 + 455a^3bx^3 + 455a^4)}{455}$$

input `int((b*x^3+a)^4,x)`output `(x*(455*a**4 + 455*a**3*b*x**3 + 390*a**2*b**2*x**6 + 182*a*b**3*x**9 + 35*b**4*x**12))/455`

3.3 $\int (a + bx^3)^3 dx$

Optimal result	54
Mathematica [A] (verified)	54
Rubi [A] (verified)	55
Maple [A] (verified)	56
Fricas [A] (verification not implemented)	56
Sympy [A] (verification not implemented)	57
Maxima [A] (verification not implemented)	57
Giac [A] (verification not implemented)	57
Mupad [B] (verification not implemented)	58
Reduce [B] (verification not implemented)	58

Optimal result

Integrand size = 9, antiderivative size = 38

$$\int (a + bx^3)^3 dx = a^3x + \frac{3}{4}a^2bx^4 + \frac{3}{7}ab^2x^7 + \frac{b^3x^{10}}{10}$$

output

```
a^3*x+3/4*a^2*b*x^4+3/7*a*b^2*x^7+1/10*b^3*x^10
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int (a + bx^3)^3 dx = a^3x + \frac{3}{4}a^2bx^4 + \frac{3}{7}ab^2x^7 + \frac{b^3x^{10}}{10}$$

input

```
Integrate[(a + b*x^3)^3,x]
```

output

```
a^3*x + (3*a^2*b*x^4)/4 + (3*a*b^2*x^7)/7 + (b^3*x^10)/10
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {747, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx^3)^3 dx$$

$$\downarrow 747$$

$$\int (a^3 + 3a^2bx^3 + 3ab^2x^6 + b^3x^9) dx$$

$$\downarrow 2009$$

$$a^3x + \frac{3}{4}a^2bx^4 + \frac{3}{7}ab^2x^7 + \frac{b^3x^{10}}{10}$$

input `Int[(a + b*x^3)^3,x]`

output `a^3*x + (3*a^2*b*x^4)/4 + (3*a*b^2*x^7)/7 + (b^3*x^10)/10`

Defintions of rubi rules used

rule 747 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Int[ExpandIntegrand[(a + b*x^n)^p, x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.87

method	result	size
gospers	$a^3x + \frac{3}{4}a^2bx^4 + \frac{3}{7}ab^2x^7 + \frac{1}{10}b^3x^{10}$	33
default	$a^3x + \frac{3}{4}a^2bx^4 + \frac{3}{7}ab^2x^7 + \frac{1}{10}b^3x^{10}$	33
norman	$a^3x + \frac{3}{4}a^2bx^4 + \frac{3}{7}ab^2x^7 + \frac{1}{10}b^3x^{10}$	33
risch	$a^3x + \frac{3}{4}a^2bx^4 + \frac{3}{7}ab^2x^7 + \frac{1}{10}b^3x^{10}$	33
parallelrisch	$a^3x + \frac{3}{4}a^2bx^4 + \frac{3}{7}ab^2x^7 + \frac{1}{10}b^3x^{10}$	33
orering	$\frac{x(14b^3x^9 + 60ab^2x^6 + 105a^2bx^3 + 140a^3)}{140}$	36

input `int((b*x^3+a)^3,x,method=_RETURNVERBOSE)`output `a^3*x+3/4*a^2*b*x^4+3/7*a*b^2*x^7+1/10*b^3*x^10`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int (a + bx^3)^3 dx = \frac{1}{10}b^3x^{10} + \frac{3}{7}ab^2x^7 + \frac{3}{4}a^2bx^4 + a^3x$$

input `integrate((b*x^3+a)^3,x, algorithm="fricas")`output `1/10*b^3*x^10 + 3/7*a*b^2*x^7 + 3/4*a^2*b*x^4 + a^3*x`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.95

$$\int (a + bx^3)^3 dx = a^3x + \frac{3a^2bx^4}{4} + \frac{3ab^2x^7}{7} + \frac{b^3x^{10}}{10}$$

input `integrate((b*x**3+a)**3,x)`output `a**3*x + 3*a**2*b*x**4/4 + 3*a*b**2*x**7/7 + b**3*x**10/10`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int (a + bx^3)^3 dx = \frac{1}{10} b^3x^{10} + \frac{3}{7} ab^2x^7 + \frac{3}{4} a^2bx^4 + a^3x$$

input `integrate((b*x^3+a)^3,x, algorithm="maxima")`output `1/10*b^3*x^10 + 3/7*a*b^2*x^7 + 3/4*a^2*b*x^4 + a^3*x`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int (a + bx^3)^3 dx = \frac{1}{10} b^3x^{10} + \frac{3}{7} ab^2x^7 + \frac{3}{4} a^2bx^4 + a^3x$$

input `integrate((b*x^3+a)^3,x, algorithm="giac")`output `1/10*b^3*x^10 + 3/7*a*b^2*x^7 + 3/4*a^2*b*x^4 + a^3*x`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int (a + bx^3)^3 dx = a^3 x + \frac{3a^2 b x^4}{4} + \frac{3a b^2 x^7}{7} + \frac{b^3 x^{10}}{10}$$

input `int((a + b*x^3)^3,x)`output `a^3*x + (b^3*x^10)/10 + (3*a^2*b*x^4)/4 + (3*a*b^2*x^7)/7`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.92

$$\int (a + bx^3)^3 dx = \frac{x(14b^3x^9 + 60ab^2x^6 + 105a^2bx^3 + 140a^3)}{140}$$

input `int((b*x^3+a)^3,x)`output `(x*(140*a**3 + 105*a**2*b*x**3 + 60*a*b**2*x**6 + 14*b**3*x**9))/140`

3.4 $\int (a + bx^3)^2 dx$

Optimal result	59
Mathematica [A] (verified)	59
Rubi [A] (verified)	60
Maple [A] (verified)	61
Fricas [A] (verification not implemented)	61
Sympy [A] (verification not implemented)	62
Maxima [A] (verification not implemented)	62
Giac [A] (verification not implemented)	62
Mupad [B] (verification not implemented)	63
Reduce [B] (verification not implemented)	63

Optimal result

Integrand size = 9, antiderivative size = 25

$$\int (a + bx^3)^2 dx = a^2x + \frac{1}{2}abx^4 + \frac{b^2x^7}{7}$$

output

```
a^2*x+1/2*a*b*x^4+1/7*b^2*x^7
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int (a + bx^3)^2 dx = a^2x + \frac{1}{2}abx^4 + \frac{b^2x^7}{7}$$

input

```
Integrate[(a + b*x^3)^2,x]
```

output

```
a^2*x + (a*b*x^4)/2 + (b^2*x^7)/7
```

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {747, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx^3)^2 dx$$

$$\downarrow 747$$

$$\int (a^2 + 2abx^3 + b^2x^6) dx$$

$$\downarrow 2009$$

$$a^2x + \frac{1}{2}abx^4 + \frac{b^2x^7}{7}$$

input `Int[(a + b*x^3)^2,x]`

output `a^2*x + (a*b*x^4)/2 + (b^2*x^7)/7`

Defintions of rubi rules used

rule 747 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*x^n)^p, x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

method	result	size
gospers	$a^2x + \frac{1}{2}abx^4 + \frac{1}{7}b^2x^7$	22
default	$a^2x + \frac{1}{2}abx^4 + \frac{1}{7}b^2x^7$	22
norman	$a^2x + \frac{1}{2}abx^4 + \frac{1}{7}b^2x^7$	22
risch	$a^2x + \frac{1}{2}abx^4 + \frac{1}{7}b^2x^7$	22
parallelrisch	$a^2x + \frac{1}{2}abx^4 + \frac{1}{7}b^2x^7$	22
orering	$\frac{x(2b^2x^6+7abx^3+14a^2)}{14}$	25

input `int((b*x^3+a)^2,x,method=_RETURNVERBOSE)`

output `a^2*x+1/2*a*b*x^4+1/7*b^2*x^7`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int (a + bx^3)^2 dx = \frac{1}{7}b^2x^7 + \frac{1}{2}abx^4 + a^2x$$

input `integrate((b*x^3+a)^2,x, algorithm="fricas")`

output `1/7*b^2*x^7 + 1/2*a*b*x^4 + a^2*x`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int (a + bx^3)^2 dx = a^2x + \frac{abx^4}{2} + \frac{b^2x^7}{7}$$

input `integrate((b*x**3+a)**2,x)`output `a**2*x + a*b*x**4/2 + b**2*x**7/7`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int (a + bx^3)^2 dx = \frac{1}{7}b^2x^7 + \frac{1}{2}abx^4 + a^2x$$

input `integrate((b*x^3+a)^2,x, algorithm="maxima")`output `1/7*b^2*x^7 + 1/2*a*b*x^4 + a^2*x`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int (a + bx^3)^2 dx = \frac{1}{7}b^2x^7 + \frac{1}{2}abx^4 + a^2x$$

input `integrate((b*x^3+a)^2,x, algorithm="giac")`output `1/7*b^2*x^7 + 1/2*a*b*x^4 + a^2*x`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int (a + bx^3)^2 dx = a^2 x + \frac{abx^4}{2} + \frac{b^2 x^7}{7}$$

input `int((a + b*x^3)^2,x)`

output `a^2*x + (b^2*x^7)/7 + (a*b*x^4)/2`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int (a + bx^3)^2 dx = \frac{x(2b^2x^6 + 7abx^3 + 14a^2)}{14}$$

input `int((b*x^3+a)^2,x)`

output `(x*(14*a**2 + 7*a*b*x**3 + 2*b**2*x**6))/14`

3.5 $\int (a + bx^3) dx$

Optimal result	64
Mathematica [A] (verified)	64
Rubi [A] (verified)	65
Maple [A] (verified)	66
Fricas [A] (verification not implemented)	66
Sympy [A] (verification not implemented)	67
Maxima [A] (verification not implemented)	67
Giac [A] (verification not implemented)	67
Mupad [B] (verification not implemented)	68
Reduce [B] (verification not implemented)	68

Optimal result

Integrand size = 7, antiderivative size = 12

$$\int (a + bx^3) dx = ax + \frac{bx^4}{4}$$

output `a*x+1/4*b*x^4`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int (a + bx^3) dx = ax + \frac{bx^4}{4}$$

input `Integrate[a + b*x^3,x]`

output `a*x + (b*x^4)/4`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx^3) dx$$

↓ 2009

$$ax + \frac{bx^4}{4}$$

input `Int[a + b*x^3,x]`

output `a*x + (b*x^4)/4`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

method	result	size
gospers	$ax + \frac{1}{4}bx^4$	11
default	$ax + \frac{1}{4}bx^4$	11
norman	$ax + \frac{1}{4}bx^4$	11
risch	$ax + \frac{1}{4}bx^4$	11
parallelrisch	$ax + \frac{1}{4}bx^4$	11
parts	$ax + \frac{1}{4}bx^4$	11
orering	$\frac{x(bx^3+4a)}{4}$	13

input `int(b*x^3+a,x,method=_RETURNVERBOSE)`output `a*x+1/4*b*x^4`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int (a + bx^3) dx = \frac{1}{4}bx^4 + ax$$

input `integrate(b*x^3+a,x, algorithm="fricas")`output `1/4*b*x^4 + a*x`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int (a + bx^3) dx = ax + \frac{bx^4}{4}$$

input `integrate(b*x**3+a,x)`

output `a*x + b*x**4/4`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int (a + bx^3) dx = \frac{1}{4}bx^4 + ax$$

input `integrate(b*x^3+a,x, algorithm="maxima")`

output `1/4*b*x^4 + a*x`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int (a + bx^3) dx = \frac{1}{4}bx^4 + ax$$

input `integrate(b*x^3+a,x, algorithm="giac")`

output `1/4*b*x^4 + a*x`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int (a + bx^3) dx = \frac{bx^4}{4} + ax$$

input `int(a + b*x^3,x)`

output `a*x + (b*x^4)/4`

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int (a + bx^3) dx = \frac{x(bx^3 + 4a)}{4}$$

input `int(b*x^3+a,x)`

output `(x*(4*a + b*x**3))/4`

3.6 $\int \frac{1}{a+bx^3} dx$

Optimal result	69
Mathematica [A] (verified)	69
Rubi [A] (verified)	70
Maple [C] (verified)	73
Fricas [A] (verification not implemented)	73
Sympy [A] (verification not implemented)	74
Maxima [A] (verification not implemented)	74
Giac [A] (verification not implemented)	75
Mupad [B] (verification not implemented)	75
Reduce [B] (verification not implemented)	76

Optimal result

Integrand size = 9, antiderivative size = 115

$$\int \frac{1}{a+bx^3} dx = -\frac{\arctan\left(\frac{\sqrt[3]{a}-2\sqrt[3]{bx}}{\sqrt{3}\sqrt[3]{a}}\right)}{\sqrt{3}a^{2/3}\sqrt[3]{b}} + \frac{\log\left(\sqrt[3]{a} + \sqrt[3]{bx}\right)}{3a^{2/3}\sqrt[3]{b}} - \frac{\log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx} + b^{2/3}x^2\right)}{6a^{2/3}\sqrt[3]{b}}$$

```
output -1/3*arctan(1/3*(a^(1/3)-2*b^(1/3)*x)*3^(1/2)/a^(1/3))*3^(1/2)/a^(2/3)/b^(1/3)+1/3*ln(a^(1/3)+b^(1/3)*x)/a^(2/3)/b^(1/3)-1/6*ln(a^(2/3)-a^(1/3)*b^(1/3)*x+b^(2/3)*x^2)/a^(2/3)/b^(1/3)
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.77

$$\int \frac{1}{a+bx^3} dx = \frac{2\sqrt{3} \arctan\left(\frac{1-2\sqrt[3]{bx}}{\sqrt{3}\sqrt[3]{a}}\right) - 2\log\left(\sqrt[3]{a} + \sqrt[3]{bx}\right) + \log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx} + b^{2/3}x^2\right)}{6a^{2/3}\sqrt[3]{b}}$$

input `Integrate[(a + b*x^3)^(-1),x]`

output
$$-1/6*(2*\text{Sqrt}[3]*\text{ArcTan}[(1 - (2*b^{(1/3)}*x)/a^{(1/3)})/\text{Sqrt}[3]] - 2*\text{Log}[a^{(1/3)} + b^{(1/3)}*x] + \text{Log}[a^{(2/3)} - a^{(1/3)}*b^{(1/3)}*x + b^{(2/3)}*x^2])/(a^{(2/3)}*b^{(1/3)})$$

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.97, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.889$, Rules used = {750, 16, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{a + bx^3} dx \\ & \quad \downarrow 750 \\ & \frac{\int \frac{2\sqrt[3]{a} - \sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{3a^{2/3}} + \frac{\int \frac{1}{\sqrt[3]{bx} + \sqrt[3]{a}}} dx}{3a^{2/3}} \\ & \quad \downarrow 16 \\ & \frac{\int \frac{2\sqrt[3]{a} - \sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \\ & \quad \downarrow 1142 \\ & \frac{\frac{3}{2}\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \frac{\int -\frac{\sqrt[3]{b}(\sqrt[3]{a} - 2\sqrt[3]{bx})}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{2\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \\ & \quad \downarrow 25 \end{aligned}$$

$$\begin{aligned}
& \frac{\frac{3}{2}\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx + \frac{\int \frac{\sqrt[3]{b}(\sqrt[3]{a}-2\sqrt[3]{bx})}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{2\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \\
& \quad \downarrow 27 \\
& \frac{\frac{3}{2}\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx + \frac{1}{2} \int \frac{\sqrt[3]{a}-2\sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \\
& \quad \downarrow 1082 \\
& \frac{\frac{1}{2} \int \frac{\sqrt[3]{a}-2\sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx + \frac{3 \int \frac{1}{\left(1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}\right)^2} d\left(1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}\right)}{-\left(1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}\right)^{-3}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \\
& \quad \downarrow 217 \\
& \frac{\frac{1}{2} \int \frac{\sqrt[3]{a}-2\sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \\
& \quad \downarrow 1103 \\
& \frac{\frac{\log(a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx+b^{2/3}x^2})}{2\sqrt[3]{b}} - \frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}}
\end{aligned}$$

input

```
Int[(a + b*x^3)^(-1), x]
```

output

```
Log[a^(1/3) + b^(1/3)*x]/(3*a^(2/3)*b^(1/3)) + (-((Sqrt[3]*ArcTan[(1 - (2*
b^(1/3)*x)/a^(1/3)]/Sqrt[3])/b^(1/3)) - Log[a^(2/3) - a^(1/3)*b^(1/3)*x +
b^(2/3)*x^2]/(2*b^(1/3)))/(3*a^(2/3))
```


Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 25 $\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[Fx, x], x]$
- rule 27 $\text{Int}[(a_)*(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_)*(Gx_)] /; \text{FreeQ}[b, x]$
- rule 217 $\text{Int}[(a_)+(b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-(\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])]$
- rule 750 $\text{Int}[(a_)+(b_)*(x_)^3)^{-1}, x_Symbol] \rightarrow \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[(2*\text{Rt}[a, 3] - \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$
- rule 1082 $\text{Int}[(a_)+(b_)*(x_)+(c_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S\text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$
- rule 1142 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \text{ Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \text{ Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.46 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.23

method	result	size
risch	$\frac{\sum_{R=\text{RootOf}(bZ^3+a)} \frac{\ln(x-R)}{R^2}}{3b}$	27
default	$\frac{\ln\left(x+\left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3b\left(\frac{a}{b}\right)^{\frac{2}{3}}} - \frac{\ln\left(x^2-\left(\frac{a}{b}\right)^{\frac{1}{3}}x+\left(\frac{a}{b}\right)^{\frac{2}{3}}\right)}{6b\left(\frac{a}{b}\right)^{\frac{2}{3}}} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{-2x}{\left(\frac{a}{b}\right)^{\frac{1}{3}}}-1\right)}{3}\right)}{3b\left(\frac{a}{b}\right)^{\frac{2}{3}}}$	91

input `int(1/(b*x^3+a),x,method=_RETURNVERBOSE)`

output `1/3/b*sum(1/_R^2*ln(x-_R),_R=RootOf(_Z^3*b+a))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 299, normalized size of antiderivative = 2.60

$$\int \frac{1}{a + bx^3} dx$$

$$= \frac{3 \sqrt{\frac{1}{3}} ab \sqrt{-\frac{(a^2b)^{\frac{1}{3}}}{b}} \log\left(\frac{2abx^3 - 3(a^2b)^{\frac{1}{3}}ax - a^2 + 3\sqrt{\frac{1}{3}}\left(2abx^2 + (a^2b)^{\frac{2}{3}}x - (a^2b)^{\frac{1}{3}}a\right)\sqrt{-\frac{(a^2b)^{\frac{1}{3}}}{b}}}{bx^3 + a}\right) - (a^2b)^{\frac{2}{3}} \log\left(abx^2 - \dots\right)}{6a^2b}$$

input `integrate(1/(b*x^3+a),x, algorithm="fricas")`

output

```
[1/6*(3*sqrt(1/3)*a*b*sqrt(-(a^2*b)^(1/3)/b)*log((2*a*b*x^3 - 3*(a^2*b)^(1/3)*a*x - a^2 + 3*sqrt(1/3)*(2*a*b*x^2 + (a^2*b)^(2/3)*x - (a^2*b)^(1/3)*a)*sqrt(-(a^2*b)^(1/3)/b))/(b*x^3 + a) - (a^2*b)^(2/3)*log(a*b*x^2 - (a^2*b)^(2/3)*x + (a^2*b)^(1/3)*a) + 2*(a^2*b)^(2/3)*log(a*b*x + (a^2*b)^(2/3)))/(a^2*b), 1/6*(6*sqrt(1/3)*a*b*sqrt((a^2*b)^(1/3)/b)*arctan(sqrt(1/3)*(2*(a^2*b)^(2/3)*x - (a^2*b)^(1/3)*a)*sqrt((a^2*b)^(1/3)/b)/a^2 - (a^2*b)^(2/3)*log(a*b*x^2 - (a^2*b)^(2/3)*x + (a^2*b)^(1/3)*a) + 2*(a^2*b)^(2/3)*log(a*b*x + (a^2*b)^(2/3)))/(a^2*b)]
```

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.17

$$\int \frac{1}{a + bx^3} dx = \text{RootSum}(27t^3a^2b - 1, (t \mapsto t \log(3ta + x)))$$

input

```
integrate(1/(b*x**3+a),x)
```

output

```
RootSum(27*_t**3*a**2*b - 1, Lambda(_t, _t*log(3*_t*a + x)))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.85

$$\int \frac{1}{a + bx^3} dx = \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(2x - \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{3b\left(\frac{a}{b}\right)^{\frac{2}{3}}} - \frac{\log\left(x^2 - x\left(\frac{a}{b}\right)^{\frac{1}{3}} + \left(\frac{a}{b}\right)^{\frac{2}{3}}\right)}{6b\left(\frac{a}{b}\right)^{\frac{2}{3}}} + \frac{\log\left(x + \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3b\left(\frac{a}{b}\right)^{\frac{2}{3}}}$$

input

```
integrate(1/(b*x^3+a),x, algorithm="maxima")
```

output

```
1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - (a/b)^(1/3))/(a/b)^(1/3))/(b*(a/b)^(2/3)) - 1/6*log(x^2 - x*(a/b)^(1/3) + (a/b)^(2/3))/(b*(a/b)^(2/3)) + 1/3*log(x + (a/b)^(1/3))/(b*(a/b)^(2/3))
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.97

$$\int \frac{1}{a + bx^3} dx = -\frac{\left(-\frac{a}{b}\right)^{\frac{1}{3}} \log\left(\left|x - \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right|\right)}{3a} + \frac{\sqrt{3}(-ab^2)^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}\left(2x + \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(-\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{3ab}$$

$$+ \frac{(-ab^2)^{\frac{1}{3}} \log\left(x^2 + x\left(-\frac{a}{b}\right)^{\frac{1}{3}} + \left(-\frac{a}{b}\right)^{\frac{2}{3}}\right)}{6ab}$$

input `integrate(1/(b*x^3+a),x, algorithm="giac")`output `-1/3*(-a/b)^(1/3)*log(abs(x - (-a/b)^(1/3)))/a + 1/3*sqrt(3)*(-a*b^2)^(1/3)*arctan(1/3*sqrt(3)*(2*x + (-a/b)^(1/3))/(-a/b)^(1/3))/(a*b) + 1/6*(-a*b^2)^(1/3)*log(x^2 + x*(-a/b)^(1/3) + (-a/b)^(2/3))/(a*b)`**Mupad [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.86

$$\int \frac{1}{a + bx^3} dx = \frac{\ln\left(b^{1/3}x + a^{1/3}\right)}{3a^{2/3}b^{1/3}} + \frac{\ln\left(3b^2x + \frac{3a^{1/3}b^{5/3}(-1+\sqrt{3}1i)}{2}\right)(-1+\sqrt{3}1i)}{6a^{2/3}b^{1/3}}$$

$$- \frac{\ln\left(3b^2x - \frac{3a^{1/3}b^{5/3}(1+\sqrt{3}1i)}{2}\right)(1+\sqrt{3}1i)}{6a^{2/3}b^{1/3}}$$

input `int(1/(a + b*x^3),x)`output `log(b^(1/3)*x + a^(1/3))/(3*a^(2/3)*b^(1/3)) + (log(3*b^2*x + (3*a^(1/3)*b^(5/3)*(3^(1/2)*1i - 1))/2)*(3^(1/2)*1i - 1))/(6*a^(2/3)*b^(1/3)) - (log(3*b^2*x - (3*a^(1/3)*b^(5/3)*(3^(1/2)*1i + 1))/2)*(3^(1/2)*1i + 1))/(6*a^(2/3)*b^(1/3))`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.58

$$\int \frac{1}{a + bx^3} dx = \frac{-2\sqrt{3} \operatorname{atan}\left(\frac{a^{\frac{1}{3}} - 2b^{\frac{1}{3}}x}{a^{\frac{1}{3}}\sqrt{3}}\right) - \log\left(a^{\frac{2}{3}} - b^{\frac{1}{3}}a^{\frac{1}{3}}x + b^{\frac{2}{3}}x^2\right) + 2\log\left(a^{\frac{1}{3}} + b^{\frac{1}{3}}x\right)}{6a^{\frac{2}{3}}b^{\frac{1}{3}}}$$

input `int(1/(b*x^3+a),x)`

output `(a**(1/3)*(-2*sqrt(3)*atan((a**(1/3)-2*b**(1/3)*x)/(a**(1/3)*sqrt(3))) - log(a**(2/3)-b**(1/3)*a**(1/3)*x+b**(2/3)*x**2)+2*log(a**(1/3)+b**(1/3)*x))/(6*b**(1/3)*a)`

3.7 $\int \frac{1}{(a+bx^3)^2} dx$

Optimal result	77
Mathematica [A] (verified)	78
Rubi [A] (verified)	78
Maple [C] (verified)	82
Fricas [A] (verification not implemented)	83
Sympy [A] (verification not implemented)	83
Maxima [A] (verification not implemented)	84
Giac [A] (verification not implemented)	84
Mupad [B] (verification not implemented)	85
Reduce [B] (verification not implemented)	85

Optimal result

Integrand size = 9, antiderivative size = 134

$$\int \frac{1}{(a+bx^3)^2} dx = \frac{x}{3a(a+bx^3)} - \frac{2 \arctan\left(\frac{\sqrt[3]{a}-2\sqrt[3]{b}x}{\sqrt{3}\sqrt[3]{a}}\right)}{3\sqrt{3}a^{5/3}\sqrt[3]{b}} + \frac{2 \log\left(\sqrt[3]{a} + \sqrt[3]{b}x\right)}{9a^{5/3}\sqrt[3]{b}} - \frac{\log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{b}x + b^{2/3}x^2\right)}{9a^{5/3}\sqrt[3]{b}}$$

output

```
1/3*x/a/(b*x^3+a)-2/9*arctan(1/3*(a^(1/3)-2*b^(1/3)*x)*3^(1/2)/a^(1/3))*3^(1/2)/a^(5/3)/b^(1/3)+2/9*ln(a^(1/3)+b^(1/3)*x)/a^(5/3)/b^(1/3)-1/9*ln(a^(2/3)-a^(1/3)*b^(1/3)*x+b^(2/3)*x^2)/a^(5/3)/b^(1/3)
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.88

$$\int \frac{1}{(a + bx^3)^2} dx$$

$$= \frac{\frac{3a^{2/3}x}{a+bx^3} - \frac{2\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{b}x}{\sqrt[3]{a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}} + \frac{2 \log\left(\sqrt[3]{a} + \sqrt[3]{b}x\right)}{\sqrt[3]{b}} - \frac{\log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{b}x + b^{2/3}x^2\right)}{\sqrt[3]{b}}}{9a^{5/3}}$$

input

```
Integrate[(a + b*x^3)^(-2), x]
```

output

```
((3*a^(2/3)*x)/(a + b*x^3) - (2*Sqrt[3]*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3))/Sqrt[3]])/b^(1/3) + (2*Log[a^(1/3) + b^(1/3)*x])/b^(1/3) - Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/b^(1/3))/(9*a^(5/3))
```

Rubi [A] (verified)Time = 0.27 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.01, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {749, 750, 16, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + bx^3)^2} dx$$

$$\downarrow 749$$

$$\frac{2 \int \frac{1}{bx^3+a} dx}{3a} + \frac{x}{3a(a + bx^3)}$$

$$\downarrow 750$$

$$2 \left(\frac{\int \frac{{}_2\sqrt[3]{a} - \sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{3a^{2/3}} + \frac{\int \frac{1}{\sqrt[3]{bx} + \sqrt[3]{a}} dx}{3a^{2/3}} \right) + \frac{x}{3a(a+bx^3)}$$

↓ 16

$$2 \left(\frac{\int \frac{{}_2\sqrt[3]{a} - \sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \right) + \frac{x}{3a(a+bx^3)}$$

↓ 1142

$$2 \left(\frac{\frac{{}_2\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \frac{\int \frac{\sqrt[3]{b}(\sqrt[3]{a} - 2\sqrt[3]{bx})}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{2\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \right) + \frac{x}{3a(a+bx^3)}$$

↓ 25

$$2 \left(\frac{\frac{{}_2\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx + \frac{\int \frac{\sqrt[3]{b}(\sqrt[3]{a} - 2\sqrt[3]{bx})}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{2\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \right) + \frac{x}{3a(a+bx^3)}$$

↓ 27

$$2 \left(\frac{\frac{{}_2\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx + \frac{1}{2} \int \frac{\sqrt[3]{a} - 2\sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \right) + \frac{x}{3a(a+bx^3)}$$

↓ 1082

$$2 \left(\frac{\int \frac{\sqrt[3]{a-2\sqrt[3]{bx}}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx + \frac{\int \frac{1}{\left(1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}\right)^2} dx - \frac{\int \frac{1}{\left(1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}\right)^{-3}} dx}{\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \right) + \frac{x}{3a(a+bx^3)}$$

217

$$2 \left(\frac{\int \frac{\sqrt[3]{a-2\sqrt[3]{bx}}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \right) + \frac{x}{3a(a+bx^3)}$$

1103

$$2 \left(\frac{\frac{\log(a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx+b^{2/3}x^2})}{2\sqrt[3]{b}} - \frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \right) + \frac{x}{3a(a+bx^3)}$$

input `Int[(a + b*x^3)^(-2),x]`

output `x/(3*a*(a + b*x^3)) + (2*(Log[a^(1/3) + b^(1/3)*x]/(3*a^(2/3)*b^(1/3)) + (-((Sqrt[3]*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3)]/Sqrt[3])]/b^(1/3)) - Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/(2*b^(1/3)))/(3*a^(2/3)))/(3*a)`

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}\{a, b, c\}, x]$
- rule 25 $\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[Fx, x], x]$
- rule 27 $\text{Int}[(a_)*(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \&\& \text{!MatchQ}[Fx, (b_)*(Gx_)] /; \text{FreeQ}[b, x]$
- rule 217 $\text{Int}[(a_)+(b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-(\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}\{a, b\}, x \&\& \text{PosQ}[a/b] \&\& (\text{LtQ}[a, 0] \parallel \text{LtQ}[b, 0])]$
- rule 749 $\text{Int}[(a_)+(b_)*(x_)^{(n)})^{(p)}, x_Symbol] \rightarrow \text{Simp}[(-x)*((a + b*x^n)^{(p+1)}/(a*n*(p+1))), x] + \text{Simp}[(n*(p+1)+1)/(a*n*(p+1)) \text{ Int}[(a + b*x^n)^{(p+1)}, x], x] /; \text{FreeQ}\{a, b\}, x \&\& \text{IGtQ}[n, 0] \&\& \text{LtQ}[p, -1] \&\& (\text{IntegerQ}[2*p] \parallel \text{Denominator}[p + 1/n] < \text{Denominator}[p])]$
- rule 750 $\text{Int}[(a_)+(b_)*(x_)^3)^{-1}, x_Symbol] \rightarrow \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[(2*\text{Rt}[a, 3] - \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}\{a, b\}, x]$
- rule 1082 $\text{Int}[(a_)+(b_)*(x_)+(c_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \&\& (\text{EqQ}[q^2, 1] \parallel \text{!RationalQ}[b^2 - 4*a*c]) /; \text{FreeQ}\{a, b, c\}, x]$
- rule 1103 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}\{a, b, c, d, e\}, x \&\& \text{EqQ}[2*c*d - b*e, 0]$

rule 1142

```
Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c)
Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.43 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.34

method	result	size
risch	$\frac{x}{3a(bx^3+a)} + \frac{2 \left(\sum_{R=\text{RootOf}(b-Z^3+a)} \frac{\ln(x-R)}{-R^2} \right)}{9ab}$	46
default	$\frac{x}{3a(bx^3+a)} + \frac{\frac{2 \ln\left(x + \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{9b\left(\frac{a}{b}\right)^{\frac{2}{3}}}}{a} - \frac{\ln\left(x^2 - \left(\frac{a}{b}\right)^{\frac{1}{3}}x + \left(\frac{a}{b}\right)^{\frac{2}{3}}\right)}{9b\left(\frac{a}{b}\right)^{\frac{2}{3}}} + \frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2x}{\left(\frac{a}{b}\right)^{\frac{1}{3}}}-1\right)}{3}\right)}{9b\left(\frac{a}{b}\right)^{\frac{2}{3}}}$	112

input

```
int(1/(b*x^3+a)^2,x,method=_RETURNVERBOSE)
```

output

```
1/3*x/a/(b*x^3+a)+2/9/a/b*sum(1/_R^2*ln(x-_R),_R=RootOf(_Z^3*b+a))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 389, normalized size of antiderivative = 2.90

$$\int \frac{1}{(a + bx^3)^2} dx$$

$$= \frac{3a^2bx + 3\sqrt{\frac{1}{3}}(ab^2x^3 + a^2b)\sqrt{-\frac{(a^2b)^{\frac{1}{3}}}{b}} \log\left(\frac{2abx^3 - 3(a^2b)^{\frac{1}{3}}ax - a^2 + 3\sqrt{\frac{1}{3}}(2abx^2 + (a^2b)^{\frac{2}{3}}x - (a^2b)^{\frac{1}{3}}a)\sqrt{-\frac{(a^2b)^{\frac{1}{3}}}{b}}}{bx^3 + a}\right)}{9(a^3b^2x^3 + a^4)}$$

input `integrate(1/(b*x^3+a)^2,x, algorithm="fricas")`

output

```
[1/9*(3*a^2*b*x + 3*sqrt(1/3)*(a*b^2*x^3 + a^2*b)*sqrt(-(a^2*b)^(1/3)/b)*log((2*a*b*x^3 - 3*(a^2*b)^(1/3)*a*x - a^2 + 3*sqrt(1/3)*(2*a*b*x^2 + (a^2*b)^(2/3)*x - (a^2*b)^(1/3)*a)*sqrt(-(a^2*b)^(1/3)/b))/(b*x^3 + a) - (b*x^3 + a)*(a^2*b)^(2/3)*log(a*b*x^2 - (a^2*b)^(2/3)*x + (a^2*b)^(1/3)*a) + 2*(b*x^3 + a)*(a^2*b)^(2/3)*log(a*b*x + (a^2*b)^(2/3)))/(a^3*b^2*x^3 + a^4*b), 1/9*(3*a^2*b*x + 6*sqrt(1/3)*(a*b^2*x^3 + a^2*b)*sqrt((a^2*b)^(1/3)/b)*arctan(sqrt(1/3)*(2*(a^2*b)^(2/3)*x - (a^2*b)^(1/3)*a)*sqrt((a^2*b)^(1/3)/b)/a^2) - (b*x^3 + a)*(a^2*b)^(2/3)*log(a*b*x^2 - (a^2*b)^(2/3)*x + (a^2*b)^(1/3)*a) + 2*(b*x^3 + a)*(a^2*b)^(2/3)*log(a*b*x + (a^2*b)^(2/3)))/(a^3*b^2*x^3 + a^4*b)]
```

Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.29

$$\int \frac{1}{(a + bx^3)^2} dx = \frac{x}{3a^2 + 3abx^3} + \text{RootSum}\left(729t^3a^5b - 8, \left(t \mapsto t \log\left(\frac{9ta^2}{2} + x\right)\right)\right)$$

input `integrate(1/(b*x**3+a)**2,x)`

output `x/(3*a**2 + 3*a*b*x**3) + RootSum(729*_t**3*a**5*b - 8, Lambda(_t, _t*log(9*_t*a**2/2 + x)))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.91

$$\int \frac{1}{(a + bx^3)^2} dx = \frac{x}{3(abx^3 + a^2)} + \frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(2x - \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{9ab\left(\frac{a}{b}\right)^{\frac{2}{3}}} - \frac{\log\left(x^2 - x\left(\frac{a}{b}\right)^{\frac{1}{3}} + \left(\frac{a}{b}\right)^{\frac{2}{3}}\right)}{9ab\left(\frac{a}{b}\right)^{\frac{2}{3}}} + \frac{2\log\left(x + \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{9ab\left(\frac{a}{b}\right)^{\frac{2}{3}}}$$

input `integrate(1/(b*x^3+a)^2,x, algorithm="maxima")`

output `1/3*x/(a*b*x^3 + a^2) + 2/9*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - (a/b)^(1/3))/(a/b)^(1/3))/(a*b*(a/b)^(2/3)) - 1/9*log(x^2 - x*(a/b)^(1/3) + (a/b)^(2/3))/(a*b*(a/b)^(2/3)) + 2/9*log(x + (a/b)^(1/3))/(a*b*(a/b)^(2/3))`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.95

$$\int \frac{1}{(a + bx^3)^2} dx = -\frac{2\left(-\frac{a}{b}\right)^{\frac{1}{3}} \log\left(\left|x - \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right|\right)}{9a^2} + \frac{x}{3(bx^3 + a)a} + \frac{2\sqrt{3}(-ab^2)^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}\left(2x + \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(-\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{9a^2b} + \frac{\left(-ab^2\right)^{\frac{1}{3}} \log\left(x^2 + x\left(-\frac{a}{b}\right)^{\frac{1}{3}} + \left(-\frac{a}{b}\right)^{\frac{2}{3}}\right)}{9a^2b}$$

input `integrate(1/(b*x^3+a)^2,x, algorithm="giac")`

output

```
-2/9*(-a/b)^(1/3)*log(abs(x - (-a/b)^(1/3)))/a^2 + 1/3*x/((b*x^3 + a)*a) +
2/9*sqrt(3)*(-a*b^2)^(1/3)*arctan(1/3*sqrt(3)*(2*x + (-a/b)^(1/3))/(-a/b)
^(1/3))/(a^2*b) + 1/9*(-a*b^2)^(1/3)*log(x^2 + x*(-a/b)^(1/3) + (-a/b)^(2/
3))/(a^2*b)
```

Mupad [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.96

$$\int \frac{1}{(a + bx^3)^2} dx = \frac{x}{3a(bx^3 + a)} + \frac{2 \ln \left(\frac{2b^{5/3}}{a^{2/3}} + \frac{2b^2x}{a} \right)}{9a^{5/3}b^{1/3}}$$

$$+ \frac{\ln \left(\frac{2b^2x}{a} + \frac{b^{5/3}(-1 + \sqrt{3}i)}{a^{2/3}} \right) (-1 + \sqrt{3}i)}{9a^{5/3}b^{1/3}}$$

$$- \frac{\ln \left(\frac{2b^2x}{a} - \frac{b^{5/3}(1 + \sqrt{3}i)}{a^{2/3}} \right) (1 + \sqrt{3}i)}{9a^{5/3}b^{1/3}}$$

input

```
int(1/(a + b*x^3)^2,x)
```

output

```
x/(3*a*(a + b*x^3)) + (2*log((2*b^(5/3))/a^(2/3) + (2*b^2*x)/a))/(9*a^(5/3)
)*b^(1/3)) + (log((2*b^2*x)/a + (b^(5/3)*(3^(1/2)*1i - 1))/a^(2/3))*(3^(1/
2)*1i - 1))/(9*a^(5/3)*b^(1/3)) - (log((2*b^2*x)/a - (b^(5/3)*(3^(1/2)*1i
+ 1))/a^(2/3))*(3^(1/2)*1i + 1))/(9*a^(5/3)*b^(1/3))
```

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.28

$$\int \frac{1}{(a + bx^3)^2} dx$$

$$= \frac{-2a^{4/3}\sqrt{3} \operatorname{atan}\left(\frac{a^{1/3}-2b^{1/3}x}{a^{1/3}\sqrt{3}}\right) - 2a^{1/3}\sqrt{3} \operatorname{atan}\left(\frac{a^{1/3}-2b^{1/3}x}{a^{1/3}\sqrt{3}}\right) b x^3 - a^{4/3} \log\left(a^{2/3} - b^{1/3}a^{1/3}x + b^{2/3}x^2\right) - a^{1/3} \log\left(a^{2/3} - b^{1/3}x\right)}{9b^{1/3}a^2(bx^3 + a)}$$

input

```
int(1/(b*x^3+a)^2,x)
```

output

```
( - 2*a**(1/3)*sqrt(3)*atan((a**(1/3) - 2*b**(1/3)*x)/(a**(1/3)*sqrt(3)))*
a - 2*a**(1/3)*sqrt(3)*atan((a**(1/3) - 2*b**(1/3)*x)/(a**(1/3)*sqrt(3)))*
b*x**3 - a**(1/3)*log(a**(2/3) - b**(1/3)*a**(1/3)*x + b**(2/3)*x**2)*a -
a**(1/3)*log(a**(2/3) - b**(1/3)*a**(1/3)*x + b**(2/3)*x**2)*b*x**3 + 2*a*
*(1/3)*log(a**(1/3) + b**(1/3)*x)*a + 2*a**(1/3)*log(a**(1/3) + b**(1/3)*x
)*b*x**3 + 3*b**(1/3)*a*x)/(9*b**(1/3)*a**2*(a + b*x**3))
```

3.8 $\int \frac{1}{(a+bx^3)^3} dx$

Optimal result	87
Mathematica [A] (verified)	88
Rubi [A] (verified)	88
Maple [C] (verified)	94
Fricas [B] (verification not implemented)	94
Sympy [A] (verification not implemented)	95
Maxima [A] (verification not implemented)	96
Giac [A] (verification not implemented)	96
Mupad [B] (verification not implemented)	97
Reduce [B] (verification not implemented)	98

Optimal result

Integrand size = 9, antiderivative size = 151

$$\int \frac{1}{(a+bx^3)^3} dx = \frac{x}{6a(a+bx^3)^2} + \frac{5x}{18a^2(a+bx^3)} - \frac{5 \arctan\left(\frac{\sqrt[3]{a}-2\sqrt[3]{bx}}{\sqrt{3}\sqrt[3]{a}}\right)}{9\sqrt{3}a^{8/3}\sqrt[3]{b}}$$

$$+ \frac{5 \log\left(\sqrt[3]{a} + \sqrt[3]{bx}\right)}{27a^{8/3}\sqrt[3]{b}} - \frac{5 \log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx} + b^{2/3}x^2\right)}{54a^{8/3}\sqrt[3]{b}}$$

output

```
1/6*x/a/(b*x^3+a)^2+5/18*x/a^2/(b*x^3+a)-5/27*arctan(1/3*(a^(1/3)-2*b^(1/3)
)*x)*3^(1/2)/a^(1/3))*3^(1/2)/a^(8/3)/b^(1/3)+5/27*ln(a^(1/3)+b^(1/3)*x)/a
^(8/3)/b^(1/3)-5/54*ln(a^(2/3)-a^(1/3)*b^(1/3)*x+b^(2/3)*x^2)/a^(8/3)/b^(1
/3)
```


Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.89

$$\int \frac{1}{(a + bx^3)^3} dx$$

$$= \frac{\frac{9a^{5/3}x}{(a+bx^3)^2} + \frac{15a^{2/3}x}{a+bx^3} - \frac{10\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{b}x}{\sqrt[3]{a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}} + \frac{10 \log\left(\sqrt[3]{a} + \sqrt[3]{bx}\right)}{\sqrt[3]{b}} - \frac{5 \log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx} + b^{2/3}x^2\right)}{\sqrt[3]{b}}}{54a^{8/3}}$$

input

```
Integrate[(a + b*x^3)^(-3), x]
```

output

```
((9*a^(5/3)*x)/(a + b*x^3)^2 + (15*a^(2/3)*x)/(a + b*x^3) - (10*sqrt(3)*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3))/sqrt(3)]/b^(1/3) + (10*Log[a^(1/3) + b^(1/3)*x])/b^(1/3) - (5*Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2])/b^(1/3))/(54*a^(8/3))
```

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.07, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 1.111$, Rules used = {749, 749, 750, 16, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + bx^3)^3} dx$$

$$\downarrow 749$$

$$\frac{5 \int \frac{1}{(bx^3+a)^2} dx}{6a} + \frac{x}{6a(a + bx^3)^2}$$

$$\downarrow 749$$

$$\frac{5 \left(\frac{2 \int \frac{1}{bx^3+a} dx}{3a} + \frac{x}{3a(a+bx^3)} \right)}{6a} + \frac{x}{6a(a+bx^3)^2}$$

↓ 750

$$\frac{5 \left(\frac{2 \left(\frac{\int \frac{{}_2\sqrt[3]{a}-\sqrt[3]{b}x}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{b}x+a^{2/3}} dx}{3a^{2/3}} + \frac{\int \frac{1}{\sqrt[3]{b}x+\sqrt[3]{a}} dx}{3a^{2/3}} \right)}{3a} + \frac{x}{3a(a+bx^3)} \right)}{6a} + \frac{x}{6a(a+bx^3)^2}$$

↓ 16

$$\frac{5 \left(\frac{2 \left(\frac{\int \frac{{}_2\sqrt[3]{a}-\sqrt[3]{b}x}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{b}x+a^{2/3}} dx}{3a^{2/3}} + \frac{\log\left(\sqrt[3]{a}+\sqrt[3]{b}x\right)}{3a^{2/3}\sqrt[3]{b}} \right)}{3a} + \frac{x}{3a(a+bx^3)} \right)}{6a} + \frac{x}{6a(a+bx^3)^2}$$

↓ 1142

$$\frac{5 \left(\frac{2 \left(\frac{{}_2\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{b}x+a^{2/3}} dx - \frac{\sqrt[3]{b} \left(\sqrt[3]{a}-2\sqrt[3]{b}x \right)}{2\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log\left(\sqrt[3]{a}+\sqrt[3]{b}x\right)}{3a^{2/3}\sqrt[3]{b}} \right)}{3a} + \frac{x}{3a(a+bx^3)} \right)}{6a} + \frac{x}{6a(a+bx^3)^2}$$

$$\begin{array}{c}
 \downarrow 25 \\
 \left(\left(\frac{\frac{\frac{3}{2} \sqrt[3]{a} \int \frac{1}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b} \sqrt[3]{bx+a^{2/3}}} dx + \frac{\int \frac{\sqrt[3]{b} (\sqrt[3]{a-2} \sqrt[3]{b_x})}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b} \sqrt[3]{bx+a^{2/3}}} dx}{2 \sqrt[3]{b}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{b_x})}{3a^{2/3} \sqrt[3]{b}}}{3a^{2/3}} \right) \right) + \frac{x}{3a(a+bx^3)} \\
 \left. \vphantom{\frac{\frac{3}{2} \sqrt[3]{a} \int \frac{1}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b} \sqrt[3]{bx+a^{2/3}}} dx + \frac{\int \frac{\sqrt[3]{b} (\sqrt[3]{a-2} \sqrt[3]{b_x})}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b} \sqrt[3]{bx+a^{2/3}}} dx}{2 \sqrt[3]{b}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{b_x})}{3a^{2/3} \sqrt[3]{b}}}{3a^{2/3}}} \right) + \frac{x}{3a(a+bx^3)}
 \end{array}$$

$$\frac{6a}{x} \frac{1}{6a(a+bx^3)^2}$$

$$\begin{array}{c}
 \downarrow 27 \\
 \left(\left(\frac{\frac{\frac{3}{2} \sqrt[3]{a} \int \frac{1}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b} \sqrt[3]{bx+a^{2/3}}} dx + \frac{1}{2} \int \frac{\sqrt[3]{a-2} \sqrt[3]{b_x}}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b} \sqrt[3]{bx+a^{2/3}}} dx}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{b_x})}{3a^{2/3} \sqrt[3]{b}}}{3a} \right) \right) + \frac{x}{3a(a+bx^3)} \\
 \left. \vphantom{\frac{\frac{3}{2} \sqrt[3]{a} \int \frac{1}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b} \sqrt[3]{bx+a^{2/3}}} dx + \frac{1}{2} \int \frac{\sqrt[3]{a-2} \sqrt[3]{b_x}}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b} \sqrt[3]{bx+a^{2/3}}} dx}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{b_x})}{3a^{2/3} \sqrt[3]{b}}}{3a}} \right) + \frac{x}{3a(a+bx^3)}
 \end{array}$$

$$\frac{6a}{x} \frac{1}{6a(a+bx^3)^2}$$

\downarrow 1082

$$\left(\frac{2}{5} \left(\frac{\int \frac{\sqrt[3]{a-2\sqrt[3]{b}x}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{b}x + a^{2/3}} dx + \frac{\int \frac{1 - 2\frac{\sqrt[3]{b}x}{\sqrt[3]{a}}}{\left(1 - 2\frac{\sqrt[3]{b}x}{\sqrt[3]{a}}\right)^2} d\left(1 - 2\frac{\sqrt[3]{b}x}{\sqrt[3]{a}}\right)}{\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log\left(\sqrt[3]{a} + \sqrt[3]{b}x\right)}{3a^{2/3}\sqrt[3]{b}} \right) + \frac{x}{3a(a+bx^3)} \right) +$$

$$\frac{6a}{6a(a+bx^3)^2}$$

↓ 217

$$\left(\frac{2}{5} \left(\frac{\int \frac{\sqrt[3]{a-2\sqrt[3]{b}x}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{b}x + a^{2/3}} dx - \frac{\sqrt{3} \arctan\left(\frac{1 - 2\frac{\sqrt[3]{b}x}{\sqrt[3]{a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log\left(\sqrt[3]{a} + \sqrt[3]{b}x\right)}{3a^{2/3}\sqrt[3]{b}} \right) + \frac{x}{3a(a+bx^3)} \right) +$$

$$\frac{6a}{6a(a+bx^3)^2}$$

↓ 1103

$$\frac{\left(\frac{2 \left(\frac{\log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{b}x + b^{2/3}x^2\right)}{2\sqrt[3]{b}} - \frac{\sqrt{3} \arctan\left(\frac{1 - 2\sqrt[3]{b}x}{\sqrt[3]{a}}\right)}{\sqrt{3}} \right)}{3a^{2/3}} + \frac{\log\left(\sqrt[3]{a} + \sqrt[3]{b}x\right)}{3a^{2/3}\sqrt[3]{b}} \right)}{3a} + \frac{x}{3a(a+bx^3)} \right) + \frac{6a}{6a(a+bx^3)^2}$$

input

```
Int[(a + b*x^3)^(-3), x]
```

output

```
x/(6*a*(a + b*x^3)^2) + (5*(x/(3*a*(a + b*x^3))) + (2*(Log[a^(1/3) + b^(1/3)*x]/(3*a^(2/3)*b^(1/3)) + (-((Sqrt[3]*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3)])/Sqrt[3])/b^(1/3)) - Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/(2*b^(1/3)))/(3*a^(2/3))))/(3*a))/(6*a)
```

Defintions of rubi rules used

rule 16

```
Int[(c_)/((a_) + (b_)*(x_)), x_Symbol] :> Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]
```

rule 25

```
Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]
```

rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$

rule 217 $\text{Int}[(a_*) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1} * \text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 749 $\text{Int}[(a_*) + (b_*)(x_)^{(n_*)})^{(p_*)}, x_Symbol] \rightarrow \text{Simp}[(-x)*((a + b*x^n)^{(p+1})/(a*n*(p+1))), x] + \text{Simp}[(n*(p+1) + 1)/(a*n*(p+1)) \text{ Int}[(a + b*x^n)^{(p+1)}, x], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ (\text{IntegerQ}[2*p] \ || \ \text{Denominator}[p + 1/n] < \text{Denominator}[p])$

rule 750 $\text{Int}[(a_*) + (b_*)(x_)^3)^{-1}, x_Symbol] \rightarrow \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[(2*\text{Rt}[a, 3] - \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$

rule 1082 $\text{Int}[(a_*) + (b_*)(x_) + (c_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1103 $\text{Int}[(d_*) + (e_*)(x_)/((a_*) + (b_*)(x_) + (c_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$

rule 1142 $\text{Int}[(d_*) + (e_*)(x_)/((a_*) + (b_*)(x_) + (c_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \text{ Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \text{ Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.46 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.38

method	result	size
risch	$\frac{5bx^4 + 4x}{18a^2 + 9a} + \frac{5 \left(\sum_{-R=\text{RootOf}(bZ^3+a)} \frac{\ln(x-R)}{-R^2} \right)}{27a^2b}$ $+ \frac{5 \left(\frac{2 \ln \left(x + \left(\frac{a}{b} \right)^{\frac{1}{3}} \right) - \ln \left(x^2 - \left(\frac{a}{b} \right)^{\frac{1}{3}} x + \left(\frac{a}{b} \right)^{\frac{2}{3}} \right)}{9b \left(\frac{a}{b} \right)^{\frac{2}{3}}} + \frac{2\sqrt{3} \arctan \left(\frac{\sqrt{3} \left(\frac{2x}{\left(\frac{a}{b} \right)^{\frac{1}{3}} - 1 \right)} \right)}{9b \left(\frac{a}{b} \right)^{\frac{2}{3}}} \right)}{9b \left(\frac{a}{b} \right)^{\frac{2}{3}}}$	57
default	$\frac{x}{6a(bx^3+a)^2} + \frac{5x}{18a(bx^3+a)} + \frac{6a}{a}$	133

input `int(1/(b*x^3+a)^3,x,method=_RETURNVERBOSE)`

output `(5/18*b/a^2*x^4+4/9*x/a)/(b*x^3+a)^2+5/27/a^2/b*sum(1/_R^2*ln(x-_R),_R=RootOf(_Z^3*b+a))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 229 vs. 2(110) = 220.

Time = 0.08 (sec) , antiderivative size = 499, normalized size of antiderivative = 3.30

$$\int \frac{1}{(a + bx^3)^3} dx$$

$$= \frac{15a^2b^2x^4 + 24a^3bx + 15\sqrt{\frac{1}{3}}(ab^3x^6 + 2a^2b^2x^3 + a^3b)\sqrt{-\frac{(a^2b)^{\frac{1}{3}}}{b}} \log\left(\frac{2abx^3 - 3(a^2b)^{\frac{1}{3}}ax - a^2 + 3\sqrt{\frac{1}{3}}(2abx^2 + (a^2b)^{\frac{1}{3}}ax - a^2)}{bx^3 + a}\right)}{15a^2b^2x^4 + 24a^3bx + 15\sqrt{\frac{1}{3}}(ab^3x^6 + 2a^2b^2x^3 + a^3b)\sqrt{-\frac{(a^2b)^{\frac{1}{3}}}{b}} \log\left(\frac{2abx^3 - 3(a^2b)^{\frac{1}{3}}ax - a^2 + 3\sqrt{\frac{1}{3}}(2abx^2 + (a^2b)^{\frac{1}{3}}ax - a^2)}{bx^3 + a}\right)}$$

input `integrate(1/(b*x^3+a)^3,x, algorithm="fricas")`

output `[1/54*(15*a^2*b^2*x^4 + 24*a^3*b*x + 15*sqrt(1/3)*(a*b^3*x^6 + 2*a^2*b^2*x^3 + a^3*b)*sqrt(-(a^2*b)^(1/3)/b)*log((2*a*b*x^3 - 3*(a^2*b)^(1/3)*a*x - a^2 + 3*sqrt(1/3)*(2*a*b*x^2 + (a^2*b)^(2/3)*x - (a^2*b)^(1/3)*a)*sqrt(-(a^2*b)^(1/3)/b))/(b*x^3 + a)) - 5*(b^2*x^6 + 2*a*b*x^3 + a^2)*(a^2*b)^(2/3)*log(a*b*x^2 - (a^2*b)^(2/3)*x + (a^2*b)^(1/3)*a) + 10*(b^2*x^6 + 2*a*b*x^3 + a^2)*(a^2*b)^(2/3)*log(a*b*x + (a^2*b)^(2/3)))/(a^4*b^3*x^6 + 2*a^5*b^2*x^3 + a^6*b), 1/54*(15*a^2*b^2*x^4 + 24*a^3*b*x + 30*sqrt(1/3)*(a*b^3*x^6 + 2*a^2*b^2*x^3 + a^3*b)*sqrt((a^2*b)^(1/3)/b)*arctan(sqrt(1/3)*(2*(a^2*b)^(2/3)*x - (a^2*b)^(1/3)*a)*sqrt((a^2*b)^(1/3)/b)/a^2) - 5*(b^2*x^6 + 2*a*b*x^3 + a^2)*(a^2*b)^(2/3)*log(a*b*x^2 - (a^2*b)^(2/3)*x + (a^2*b)^(1/3)*a) + 10*(b^2*x^6 + 2*a*b*x^3 + a^2)*(a^2*b)^(2/3)*log(a*b*x + (a^2*b)^(2/3)))/(a^4*b^3*x^6 + 2*a^5*b^2*x^3 + a^6*b)]`

Sympy [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.42

$$\int \frac{1}{(a + bx^3)^3} dx = \frac{8ax + 5bx^4}{18a^4 + 36a^3bx^3 + 18a^2b^2x^6} + \text{RootSum}\left(19683t^3a^8b - 125, \left(t \mapsto t \log\left(\frac{27ta^3}{5} + x\right)\right)\right)$$

input `integrate(1/(b*x**3+a)**3,x)`

output $(8ax + 5bx^4)/(18a^4 + 36a^3bx^3 + 18a^2b^2x^6) + \text{RootSum}(19683_t^3a^8b - 125, \text{Lambda}(t, _t \log(27_t a^3/5 + x)))$

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.96

$$\int \frac{1}{(a + bx^3)^3} dx = \frac{5bx^4 + 8ax}{18(a^2b^2x^6 + 2a^3bx^3 + a^4)} + \frac{5\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(2x - \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{27a^2b\left(\frac{a}{b}\right)^{\frac{2}{3}}} - \frac{5 \log\left(x^2 - x\left(\frac{a}{b}\right)^{\frac{1}{3}} + \left(\frac{a}{b}\right)^{\frac{2}{3}}\right)}{54a^2b\left(\frac{a}{b}\right)^{\frac{2}{3}}} + \frac{5 \log\left(x + \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{27a^2b\left(\frac{a}{b}\right)^{\frac{2}{3}}}$$

input `integrate(1/(b*x^3+a)^3,x, algorithm="maxima")`

output $1/18*(5bx^4 + 8ax)/(a^2b^2x^6 + 2a^3bx^3 + a^4) + 5/27*\text{sqrt}(3)*\text{arctan}(1/3*\text{sqrt}(3)*(2x - (a/b)^{(1/3)})/(a/b)^{(1/3)})/(a^2b*(a/b)^{(2/3)}) - 5/54*\log(x^2 - x*(a/b)^{(1/3)} + (a/b)^{(2/3)})/(a^2b*(a/b)^{(2/3)}) + 5/27*\log(x + (a/b)^{(1/3)})/(a^2b*(a/b)^{(2/3)})$

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.91

$$\int \frac{1}{(a + bx^3)^3} dx = -\frac{5\left(-\frac{a}{b}\right)^{\frac{1}{3}} \log\left(\left|x - \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right|\right)}{27a^3} + \frac{5\sqrt{3}(-ab^2)^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}\left(2x + \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(-\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{27a^3b} + \frac{5(-ab^2)^{\frac{1}{3}} \log\left(x^2 + x\left(-\frac{a}{b}\right)^{\frac{1}{3}} + \left(-\frac{a}{b}\right)^{\frac{2}{3}}\right)}{54a^3b} + \frac{5bx^4 + 8ax}{18(bx^3 + a)^2a^2}$$

input `integrate(1/(b*x^3+a)^3,x, algorithm="giac")`

output `-5/27*(-a/b)^(1/3)*log(abs(x - (-a/b)^(1/3)))/a^3 + 5/27*sqrt(3)*(-a*b^2)^(1/3)*arctan(1/3*sqrt(3)*(2*x + (-a/b)^(1/3))/(-a/b)^(1/3))/(a^3*b) + 5/54*(-a*b^2)^(1/3)*log(x^2 + x*(-a/b)^(1/3) + (-a/b)^(2/3))/(a^3*b) + 1/18*(5*b*x^4 + 8*a*x)/((b*x^3 + a)^2*a^2)`

Mupad [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.94

$$\int \frac{1}{(a + bx^3)^3} dx = \frac{\frac{4x}{9a} + \frac{5bx^4}{18a^2}}{a^2 + 2abx^3 + b^2x^6} + \frac{5 \ln(b^{1/3}x + a^{1/3})}{27a^{8/3}b^{1/3}}$$

$$+ \frac{\ln\left(\frac{5b^2x}{3a^2} + \frac{b^{5/3}(-5+\sqrt{3}5i)}{6a^{5/3}}\right) (-5 + \sqrt{3}5i)}{54a^{8/3}b^{1/3}}$$

$$- \frac{\ln\left(\frac{5b^2x}{3a^2} - \frac{b^{5/3}(5+\sqrt{3}5i)}{6a^{5/3}}\right) (5 + \sqrt{3}5i)}{54a^{8/3}b^{1/3}}$$

input `int(1/(a + b*x^3)^3,x)`

output `((4*x)/(9*a) + (5*b*x^4)/(18*a^2))/(a^2 + b^2*x^6 + 2*a*b*x^3) + (5*log(b^(1/3)*x + a^(1/3)))/(27*a^(8/3)*b^(1/3)) + (log((5*b^2*x)/(3*a^2) + (b^(5/3)*(3^(1/2)*5i - 5))/(6*a^(5/3))))*(3^(1/2)*5i - 5)/(54*a^(8/3)*b^(1/3)) - (log((5*b^2*x)/(3*a^2) - (b^(5/3)*(3^(1/2)*5i + 5))/(6*a^(5/3))))*(3^(1/2)*5i + 5)/(54*a^(8/3)*b^(1/3))`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 278, normalized size of antiderivative = 1.84

$$\int \frac{1}{(a + bx^3)^3} dx$$

$$= \frac{-10a^{\frac{7}{3}}\sqrt{3} \operatorname{atan}\left(\frac{a^{\frac{1}{3}} - 2b^{\frac{1}{3}}x}{a^{\frac{1}{3}}\sqrt{3}}\right) - 20a^{\frac{4}{3}}\sqrt{3} \operatorname{atan}\left(\frac{a^{\frac{1}{3}} - 2b^{\frac{1}{3}}x}{a^{\frac{1}{3}}\sqrt{3}}\right) b x^3 - 10a^{\frac{1}{3}}\sqrt{3} \operatorname{atan}\left(\frac{a^{\frac{1}{3}} - 2b^{\frac{1}{3}}x}{a^{\frac{1}{3}}\sqrt{3}}\right) b^2 x^6 - 5a^{\frac{7}{3}} \log\left(a^{\frac{2}{3}} - b^{\frac{2}{3}}x^2\right) a^{\frac{1}{3}} x^2 - 10a^{\frac{4}{3}} \log\left(a^{\frac{2}{3}} - b^{\frac{2}{3}}x^2\right) b x^3 - 5a^{\frac{1}{3}} \log\left(a^{\frac{2}{3}} - b^{\frac{2}{3}}x^2\right) b^2 x^6 + 10a^{\frac{1}{3}} \log\left(a^{\frac{1}{3}} + b^{\frac{1}{3}}x\right) a^{\frac{2}{3}} x^2 + 20a^{\frac{1}{3}} \log\left(a^{\frac{1}{3}} + b^{\frac{1}{3}}x\right) a^{\frac{1}{3}} x b x^3 + 10a^{\frac{1}{3}} \log\left(a^{\frac{1}{3}} + b^{\frac{1}{3}}x\right) b^2 x^6 + 24b^{\frac{1}{3}} a^{\frac{2}{3}} x + 15b^{\frac{1}{3}} a^{\frac{1}{3}} b^2 x^4}{54b^{\frac{1}{3}}(a^{\frac{2}{3}} + 2abx^3 + b^2x^6)}$$

input `int(1/(b*x^3+a)^3,x)`output

```
( - 10*a**(1/3)*sqrt(3)*atan((a**(1/3) - 2*b**(1/3)*x)/(a**(1/3)*sqrt(3)))*a**2 - 20*a**(1/3)*sqrt(3)*atan((a**(1/3) - 2*b**(1/3)*x)/(a**(1/3)*sqrt(3)))*a*b*x**3 - 10*a**(1/3)*sqrt(3)*atan((a**(1/3) - 2*b**(1/3)*x)/(a**(1/3)*sqrt(3)))*b**2*x**6 - 5*a**(1/3)*log(a**(2/3) - b**(1/3)*a**(1/3)*x + b**(2/3)*x**2)*a**2 - 10*a**(1/3)*log(a**(2/3) - b**(1/3)*a**(1/3)*x + b**(2/3)*x**2)*a*b*x**3 - 5*a**(1/3)*log(a**(2/3) - b**(1/3)*a**(1/3)*x + b**(2/3)*x**2)*b**2*x**6 + 10*a**(1/3)*log(a**(1/3) + b**(1/3)*x)*a**2 + 20*a*(1/3)*log(a**(1/3) + b**(1/3)*x)*a*b*x**3 + 10*a**(1/3)*log(a**(1/3) + b*(1/3)*x)*b**2*x**6 + 24*b**(1/3)*a**2*x + 15*b**(1/3)*a*b*x**4)/(54*b**(1/3)*a**3*(a**2 + 2*a*b*x**3 + b**2*x**6))
```

3.9 $\int \frac{1}{1+a+bx^3} dx$

Optimal result	99
Mathematica [A] (verified)	100
Rubi [A] (verified)	100
Maple [C] (verified)	103
Fricas [B] (verification not implemented)	104
Sympy [A] (verification not implemented)	104
Maxima [A] (verification not implemented)	105
Giac [A] (verification not implemented)	105
Mupad [B] (verification not implemented)	106
Reduce [B] (verification not implemented)	107

Optimal result

Integrand size = 10, antiderivative size = 125

$$\int \frac{1}{1+a+bx^3} dx = -\frac{\arctan\left(\frac{1-\frac{2\sqrt[3]{bx}}{\sqrt[3]{1+a}}}{\sqrt{3}}\right)}{\sqrt{3}(1+a)^{2/3}\sqrt[3]{b}} + \frac{\log\left(\sqrt[3]{1+a} + \sqrt[3]{bx}\right)}{3(1+a)^{2/3}\sqrt[3]{b}} - \frac{\log\left((1+a)^{2/3} - \sqrt[3]{1+a}\sqrt[3]{bx} + b^{2/3}x^2\right)}{6(1+a)^{2/3}\sqrt[3]{b}}$$

output

```
-1/3*arctan(1/3*(1-2*b^(1/3)*x/(1+a)^(1/3))*3^(1/2))*3^(1/2)/(1+a)^(2/3)/b
^(1/3)+1/3*ln((1+a)^(1/3)+b^(1/3)*x)/(1+a)^(2/3)/b^(1/3)-1/6*ln((1+a)^(2/3)
)-(1+a)^(1/3)*b^(1/3)*x+b^(2/3)*x^2)/(1+a)^(2/3)/b^(1/3)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.81

$$\int \frac{1}{1+a+bx^3} dx$$

$$= \frac{2\sqrt{3} \arctan\left(\frac{-1 + \frac{2\sqrt[3]{bx}}{\sqrt[3]{1+a}}}{\sqrt{3}}\right) + 2 \log\left(\sqrt[3]{1+a} + \sqrt[3]{bx}\right) - \log\left((1+a)^{2/3} - \sqrt[3]{1+a}\sqrt[3]{bx} + b^{2/3}x^2\right)}{6(1+a)^{2/3}\sqrt[3]{b}}$$

input `Integrate[(1 + a + b*x^3)^(-1), x]`

output `(2*sqrt(3)*ArcTan[(-1 + (2*b^(1/3)*x)/(1 + a)^(1/3))/sqrt(3)] + 2*Log[(1 + a)^(1/3) + b^(1/3)*x] - Log[(1 + a)^(2/3) - (1 + a)^(1/3)*b^(1/3)*x + b^(2/3)*x^2])/(6*(1 + a)^(2/3)*b^(1/3))`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.98, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {750, 16, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{a+bx^3+1} dx$$

$$\downarrow 750$$

$$\int \frac{2\sqrt[3]{a+1}-\sqrt[3]{bx}}{b^{2/3}x^2-\sqrt[3]{a+1}\sqrt[3]{bx+(a+1)^{2/3}}} dx + \int \frac{1}{\sqrt[3]{bx+\sqrt[3]{a+1}}}$$

$$\frac{dx}{3(a+1)^{2/3}} + \frac{dx}{3(a+1)^{2/3}}$$

$$\downarrow 16$$

$$\begin{aligned}
& \frac{\int \frac{2\sqrt[3]{a+1}-\sqrt[3]{bx}}{b^{2/3}x^2-\sqrt[3]{a+1}\sqrt[3]{bx+(a+1)^{2/3}}} dx}{3(a+1)^{2/3}} + \frac{\log\left(\sqrt[3]{a+1} + \sqrt[3]{bx}\right)}{3(a+1)^{2/3}\sqrt[3]{b}} \\
& \quad \downarrow 1142 \\
& \frac{\frac{3}{2}\sqrt[3]{a+1} \int \frac{1}{b^{2/3}x^2-\sqrt[3]{a+1}\sqrt[3]{bx+(a+1)^{2/3}}} dx - \frac{\int \frac{\sqrt[3]{b}\left(\sqrt[3]{a+1}-2\sqrt[3]{bx}\right)}{b^{2/3}x^2-\sqrt[3]{a+1}\sqrt[3]{bx+(a+1)^{2/3}}} dx}{2\sqrt[3]{b}}}{3(a+1)^{2/3}} + \\
& \quad \frac{\log\left(\sqrt[3]{a+1} + \sqrt[3]{bx}\right)}{3(a+1)^{2/3}\sqrt[3]{b}} \\
& \quad \downarrow 25 \\
& \frac{\frac{3}{2}\sqrt[3]{a+1} \int \frac{1}{b^{2/3}x^2-\sqrt[3]{a+1}\sqrt[3]{bx+(a+1)^{2/3}}} dx + \frac{\int \frac{\sqrt[3]{b}\left(\sqrt[3]{a+1}-2\sqrt[3]{bx}\right)}{b^{2/3}x^2-\sqrt[3]{a+1}\sqrt[3]{bx+(a+1)^{2/3}}} dx}{2\sqrt[3]{b}}}{3(a+1)^{2/3}} + \\
& \quad \frac{\log\left(\sqrt[3]{a+1} + \sqrt[3]{bx}\right)}{3(a+1)^{2/3}\sqrt[3]{b}} \\
& \quad \downarrow 27 \\
& \frac{\frac{3}{2}\sqrt[3]{a+1} \int \frac{1}{b^{2/3}x^2-\sqrt[3]{a+1}\sqrt[3]{bx+(a+1)^{2/3}}} dx + \frac{1}{2} \int \frac{\sqrt[3]{a+1}-2\sqrt[3]{bx}}{b^{2/3}x^2-\sqrt[3]{a+1}\sqrt[3]{bx+(a+1)^{2/3}}} dx}{3(a+1)^{2/3}} + \\
& \quad \frac{\log\left(\sqrt[3]{a+1} + \sqrt[3]{bx}\right)}{3(a+1)^{2/3}\sqrt[3]{b}} \\
& \quad \downarrow 1082 \\
& \frac{\frac{1}{2} \int \frac{\sqrt[3]{a+1}-2\sqrt[3]{bx}}{b^{2/3}x^2-\sqrt[3]{a+1}\sqrt[3]{bx+(a+1)^{2/3}}} dx + \frac{3 \int \frac{1}{\left(1-\frac{2\sqrt[3]{bx}}{\sqrt[3]{a+1}}\right)^2} d\left(1-\frac{2\sqrt[3]{bx}}{\sqrt[3]{a+1}}\right)}{\left(1-\frac{2\sqrt[3]{bx}}{\sqrt[3]{a+1}}\right)^{-3}}}{3(a+1)^{2/3}} + \\
& \quad \frac{\log\left(\sqrt[3]{a+1} + \sqrt[3]{bx}\right)}{3(a+1)^{2/3}\sqrt[3]{b}} \\
& \quad \downarrow 217
\end{aligned}$$

$$\frac{\frac{1}{2} \int \frac{\sqrt[3]{a+1} - 2\sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a+1}\sqrt[3]{bx} + (a+1)^{2/3}} dx - \frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a+1}}}{\sqrt{3}}\right)}{\sqrt[3]{b}}}{3(a+1)^{2/3}} + \frac{\log\left(\sqrt[3]{a+1} + \sqrt[3]{bx}\right)}{3(a+1)^{2/3}\sqrt[3]{b}}$$

↓ 1103

$$\frac{\frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a+1}}}{\sqrt{3}}\right)}{\sqrt[3]{b}} - \frac{\log\left(-\sqrt[3]{a+1}\sqrt[3]{bx} + (a+1)^{2/3} + b^{2/3}x^2\right)}{2\sqrt[3]{b}}}{3(a+1)^{2/3}} + \frac{\log\left(\sqrt[3]{a+1} + \sqrt[3]{bx}\right)}{3(a+1)^{2/3}\sqrt[3]{b}}$$

input `Int[(1 + a + b*x^3)^(-1), x]`

output `Log[(1 + a)^(1/3) + b^(1/3)*x]/(3*(1 + a)^(2/3)*b^(1/3)) + (-((Sqrt[3]*ArcTan[(1 - (2*b^(1/3)*x)/(1 + a)^(1/3)]/Sqrt[3])/b^(1/3)) - Log[(1 + a)^(2/3) - (1 + a)^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/(2*b^(1/3)))/(3*(1 + a)^(2/3))`

Defintions of rubi rules used

rule 16 `Int[(c_)/((a_) + (b_)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

```
rule 750 Int[((a_) + (b_)*(x_)^3)^(-1), x_Symbol] := Simp[1/(3*Rt[a, 3]^2) Int[1/
(Rt[a, 3] + Rt[b, 3]*x), x], x] + Simp[1/(3*Rt[a, 3]^2) Int[(2*Rt[a, 3] -
Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /;
FreeQ[{a, b}, x]
```

```
rule 1082 Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

```
rule 1103 Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

```
rule 1142 Int(((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c)
Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.46 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.22

method	result	size
risch	$\frac{\sum_{-R=\text{RootOf}(bZ^3+a+1)} \frac{\ln(x-R)}{-R^2}}{3b}$	28
default	$\frac{\ln\left(x + \left(\frac{1+a}{b}\right)^{\frac{1}{3}}\right)}{3b\left(\frac{1+a}{b}\right)^{\frac{2}{3}}} - \frac{\ln\left(x^2 - \left(\frac{1+a}{b}\right)^{\frac{1}{3}}x + \left(\frac{1+a}{b}\right)^{\frac{2}{3}}\right)}{6b\left(\frac{1+a}{b}\right)^{\frac{2}{3}}} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2x}{\left(\frac{1+a}{b}\right)^{\frac{1}{3}}}-1\right)}{3}\right)}{3b\left(\frac{1+a}{b}\right)^{\frac{2}{3}}}$	105

```
input int(1/(b*x^3+a+1),x,method=_RETURNVERBOSE)
```


output `1/3/b*sum(1/_R^2*ln(x-_R),_R=RootOf(_Z^3*b+a+1))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 198 vs. 2(90) = 180.

Time = 0.09 (sec) , antiderivative size = 446, normalized size of antiderivative = 3.57

$$\int \frac{1}{1+a+bx^3} dx = \text{Too large to display}$$

input `integrate(1/(b*x^3+a+1),x, algorithm="fricas")`

output `[1/6*(3*sqrt(1/3)*(a + 1)*b*sqrt(-((a^2 + 2*a + 1)*b)^(1/3)/b)*log((2*(a + 1)*b*x^3 - 3*((a^2 + 2*a + 1)*b)^(1/3)*(a + 1)*x - a^2 + 3*sqrt(1/3)*(2*(a + 1)*b*x^2 + ((a^2 + 2*a + 1)*b)^(2/3)*x - ((a^2 + 2*a + 1)*b)^(1/3)*(a + 1))*sqrt(-((a^2 + 2*a + 1)*b)^(1/3)/b) - 2*a - 1)/(b*x^3 + a + 1)) - ((a^2 + 2*a + 1)*b)^(2/3)*log((a + 1)*b*x^2 - ((a^2 + 2*a + 1)*b)^(2/3)*x + ((a^2 + 2*a + 1)*b)^(1/3)*(a + 1)) + 2*((a^2 + 2*a + 1)*b)^(2/3)*log((a + 1)*b*x + ((a^2 + 2*a + 1)*b)^(2/3)))/((a^2 + 2*a + 1)*b), 1/6*(6*sqrt(1/3)*(a + 1)*b*sqrt(((a^2 + 2*a + 1)*b)^(1/3)/b)*arctan(sqrt(1/3)*(2*((a^2 + 2*a + 1)*b)^(2/3)*x - ((a^2 + 2*a + 1)*b)^(1/3)*(a + 1))*sqrt(((a^2 + 2*a + 1)*b)^(1/3)/b)/(a^2 + 2*a + 1)) - ((a^2 + 2*a + 1)*b)^(2/3)*log((a + 1)*b*x^2 - ((a^2 + 2*a + 1)*b)^(2/3)*x + ((a^2 + 2*a + 1)*b)^(1/3)*(a + 1)) + 2*((a^2 + 2*a + 1)*b)^(2/3)*log((a + 1)*b*x + ((a^2 + 2*a + 1)*b)^(2/3)))/((a^2 + 2*a + 1)*b)]`

Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.26

$$\int \frac{1}{1+a+bx^3} dx = \text{RootSum}(t^3 \cdot (27a^2b + 54ab + 27b) - 1, (t \mapsto t \log(3ta + 3t + x)))$$

input `integrate(1/(b*x**3+a+1),x)`

output `RootSum(_t**3*(27*a**2*b + 54*a*b + 27*b) - 1, Lambda(_t, _t*log(3*_t*a + 3*_t + x)))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.91

$$\int \frac{1}{1+a+bx^3} dx = \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(2x - \left(\frac{a+1}{b}\right)^{\frac{1}{3}}\right)}{3\left(\frac{a+1}{b}\right)^{\frac{1}{3}}}\right)}{3b\left(\frac{a+1}{b}\right)^{\frac{2}{3}}} - \frac{\log\left(x^2 - x\left(\frac{a+1}{b}\right)^{\frac{1}{3}} + \left(\frac{a+1}{b}\right)^{\frac{2}{3}}\right)}{6b\left(\frac{a+1}{b}\right)^{\frac{2}{3}}} + \frac{\log\left(x + \left(\frac{a+1}{b}\right)^{\frac{1}{3}}\right)}{3b\left(\frac{a+1}{b}\right)^{\frac{2}{3}}}$$

input `integrate(1/(b*x^3+a+1),x, algorithm="maxima")`

output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - ((a + 1)/b)^(1/3))/((a + 1)/b)^(1/3))/(b*((a + 1)/b)^(2/3)) - 1/6*log(x^2 - x*((a + 1)/b)^(1/3) + ((a + 1)/b)^(2/3))/(b*((a + 1)/b)^(2/3)) + 1/3*log(x + ((a + 1)/b)^(1/3))/(b*((a + 1)/b)^(2/3))`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.14

$$\int \frac{1}{1+a+bx^3} dx = \frac{(-ab^2 - b^2)^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}\left(2x + \left(-\frac{a+1}{b}\right)^{\frac{1}{3}}\right)}{3\left(-\frac{a+1}{b}\right)^{\frac{1}{3}}}\right)}{\sqrt{3}ab + \sqrt{3}b} + \frac{(-ab^2 - b^2)^{\frac{1}{3}} \log\left(x^2 + x\left(-\frac{a+1}{b}\right)^{\frac{1}{3}} + \left(-\frac{a+1}{b}\right)^{\frac{2}{3}}\right)}{6(ab + b)} - \frac{\left(-\frac{a+1}{b}\right)^{\frac{1}{3}} \log\left(\left|x - \left(-\frac{a+1}{b}\right)^{\frac{1}{3}}\right|\right)}{3(a + 1)}$$

input `integrate(1/(b*x^3+a+1),x, algorithm="giac")`

output $(-a*b^2 - b^2)^{1/3}*\arctan(1/3*\sqrt{3}*(2*x + (-a + 1)/b)^{1/3})/(-a + 1)/b)^{1/3})/(\sqrt{3}*a*b + \sqrt{3}*b) + 1/6*(-a*b^2 - b^2)^{1/3}*\log(x^2 + x*(-a + 1)/b)^{1/3} + (-a + 1)/b)^{2/3})/(a*b + b) - 1/3*(-a + 1)/b)^{1/3}*\log(\text{abs}(x - (-a + 1)/b)^{1/3})/(a + 1)$

Mupad [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.10

$$\int \frac{1}{1+a+bx^3} dx = \frac{\ln\left(a + b^{1/3} x (a+1)^{2/3} + 1\right)}{3 b^{1/3} (a+1)^{2/3}} + \frac{\ln\left(3 b^2 x + \frac{(9 a b^2 + 9 b^2) (-1 + \sqrt{3} i i)}{6 b^{1/3} (a+1)^{2/3}}\right) (-1 + \sqrt{3} i i)}{6 b^{1/3} (a+1)^{2/3}} - \frac{\ln\left(3 b^2 x - \frac{(9 a b^2 + 9 b^2) (1 + \sqrt{3} i i)}{6 b^{1/3} (a+1)^{2/3}}\right) (1 + \sqrt{3} i i)}{6 b^{1/3} (a+1)^{2/3}}$$

input `int(1/(a + b*x^3 + 1),x)`

output $\log(a + b^{1/3}*x*(a + 1)^{2/3} + 1)/(3*b^{1/3}*(a + 1)^{2/3}) + (\log(3*b^2*x + ((9*a*b^2 + 9*b^2)*(3^{1/2}*1i - 1))/(6*b^{1/3}*(a + 1)^{2/3}))*3^{1/2}*1i - 1)/(6*b^{1/3}*(a + 1)^{2/3}) - (\log(3*b^2*x - ((9*a*b^2 + 9*b^2)*(3^{1/2}*1i + 1))/(6*b^{1/3}*(a + 1)^{2/3}))*3^{1/2}*1i + 1)/(6*b^{1/3}*(a + 1)^{2/3})$

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.63

$$\int \frac{1}{1+a+bx^3} dx$$

$$= \frac{-2\sqrt{3} \operatorname{atan}\left(\frac{(a+1)^{\frac{1}{3}} - 2b^{\frac{1}{3}}x}{(a+1)^{\frac{1}{3}}\sqrt{3}}\right) - \log\left((a+1)^{\frac{2}{3}} - b^{\frac{1}{3}}(a+1)^{\frac{1}{3}}x + b^{\frac{2}{3}}x^2\right) + 2\log\left((a+1)^{\frac{1}{3}} + b^{\frac{1}{3}}x\right)}{6(a+1)^{\frac{2}{3}}b^{\frac{1}{3}}}$$

input `int(1/(b*x^3+a+1),x)`output `((a + 1)**(1/3)*(- 2*sqrt(3)*atan(((a + 1)**(1/3) - 2*b**(1/3)*x)/((a + 1)**(1/3)*sqrt(3)))) - log((a + 1)**(2/3) - b**(1/3)*(a + 1)**(1/3)*x + b**(2/3)*x**2) + 2*log((a + 1)**(1/3) + b**(1/3)*x))/(6*b**(1/3)*(a + 1))`

3.10 $\int \frac{1}{1+a-bx^3} dx$

Optimal result	108
Mathematica [A] (verified)	109
Rubi [A] (verified)	109
Maple [C] (verified)	112
Fricas [B] (verification not implemented)	113
Sympy [A] (verification not implemented)	113
Maxima [A] (verification not implemented)	114
Giac [A] (verification not implemented)	114
Mupad [B] (verification not implemented)	115
Reduce [B] (verification not implemented)	115

Optimal result

Integrand size = 11, antiderivative size = 124

$$\int \frac{1}{1+a-bx^3} dx = \frac{\arctan\left(\frac{1+\frac{2\sqrt[3]{bx}}{\sqrt[3]{1+a}}}{\sqrt{3}}\right)}{\sqrt{3}(1+a)^{2/3}\sqrt[3]{b}} - \frac{\log\left(\sqrt[3]{1+a}-\sqrt[3]{bx}\right)}{3(1+a)^{2/3}\sqrt[3]{b}} + \frac{\log\left((1+a)^{2/3}+\sqrt[3]{1+a}\sqrt[3]{bx}+b^{2/3}x^2\right)}{6(1+a)^{2/3}\sqrt[3]{b}}$$

output

```
1/3*arctan(1/3*(1+2*b^(1/3)*x/(1+a)^(1/3))*3^(1/2))*3^(1/2)/(1+a)^(2/3)/b^(1/3)-1/3*ln((1+a)^(1/3)-b^(1/3)*x)/(1+a)^(2/3)/b^(1/3)+1/6*ln((1+a)^(2/3)+(1+a)^(1/3)*b^(1/3)*x+b^(2/3)*x^2)/(1+a)^(2/3)/b^(1/3)
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+a-bx^3} dx$$

$$= \frac{(-1)^{2/3} \left(-2\sqrt{3} \arctan \left(\frac{-1 + \sqrt[3]{-1} \sqrt[3]{bx}}{\sqrt[3]{1+a}} \right) - 2 \log \left(\sqrt[3]{1+a} + \sqrt[3]{-1} \sqrt[3]{bx} \right) + \log \left((1+a)^{2/3} - \sqrt[3]{-1} \sqrt[3]{1+a} \right) \right)}{6(1+a)^{2/3} \sqrt[3]{b}}$$

input `Integrate[(1 + a - b*x^3)^(-1),x]`

output `((-1)^(2/3)*(-2*Sqrt[3]*ArcTan[(-1 + (2*(-1)^(1/3)*b^(1/3)*x)/(1 + a)^(1/3)])/Sqrt[3]] - 2*Log[(1 + a)^(1/3) + (-1)^(1/3)*b^(1/3)*x] + Log[(1 + a)^(2/3) - (-1)^(1/3)*(1 + a)^(1/3)*b^(1/3)*x + (-1)^(2/3)*b^(2/3)*x^2])/(6*(1 + a)^(2/3)*b^(1/3))`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.98, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {750, 16, 1142, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{a - bx^3 + 1} dx$$

$$\downarrow 750$$

$$\frac{\int \frac{\sqrt[3]{bx+2}\sqrt[3]{a+1}}{b^{2/3}x^2 + \sqrt[3]{a+1}\sqrt[3]{bx+(a+1)^{2/3}}} dx}{3(a+1)^{2/3}} + \frac{\int \frac{1}{\sqrt[3]{a+1} - \sqrt[3]{bx}} dx}{3(a+1)^{2/3}}$$

$$\downarrow 16$$

$$\begin{aligned}
& \frac{\int \frac{\sqrt[3]{bx+2}\sqrt[3]{a+1}}{b^{2/3}x^2+\sqrt[3]{a+1}\sqrt[3]{bx+(a+1)^{2/3}}} dx}{3(a+1)^{2/3}} - \frac{\log\left(\sqrt[3]{a+1}-\sqrt[3]{bx}\right)}{3(a+1)^{2/3}\sqrt[3]{b}} \\
& \quad \downarrow 1142 \\
& \frac{\frac{3}{2}\sqrt[3]{a+1} \int \frac{1}{b^{2/3}x^2+\sqrt[3]{a+1}\sqrt[3]{bx+(a+1)^{2/3}}} dx + \frac{\int \frac{\sqrt[3]{b}\left(2\sqrt[3]{bx}+\sqrt[3]{a+1}\right)}{b^{2/3}x^2+\sqrt[3]{a+1}\sqrt[3]{bx+(a+1)^{2/3}}} dx}{2\sqrt[3]{b}}}{3(a+1)^{2/3}} - \\
& \quad \frac{\log\left(\sqrt[3]{a+1}-\sqrt[3]{bx}\right)}{3(a+1)^{2/3}\sqrt[3]{b}} \\
& \quad \downarrow 27 \\
& \frac{\frac{3}{2}\sqrt[3]{a+1} \int \frac{1}{b^{2/3}x^2+\sqrt[3]{a+1}\sqrt[3]{bx+(a+1)^{2/3}}} dx + \frac{1}{2} \int \frac{2\sqrt[3]{bx}+\sqrt[3]{a+1}}{b^{2/3}x^2+\sqrt[3]{a+1}\sqrt[3]{bx+(a+1)^{2/3}}} dx}{3(a+1)^{2/3}} - \\
& \quad \frac{\log\left(\sqrt[3]{a+1}-\sqrt[3]{bx}\right)}{3(a+1)^{2/3}\sqrt[3]{b}} \\
& \quad \downarrow 1082 \\
& \frac{\frac{1}{2} \int \frac{2\sqrt[3]{bx}+\sqrt[3]{a+1}}{b^{2/3}x^2+\sqrt[3]{a+1}\sqrt[3]{bx+(a+1)^{2/3}}} dx - \frac{3 \int \frac{1}{\left(\frac{2\sqrt[3]{bx}}{\sqrt[3]{a+1}}+1\right)^2} d\left(\frac{2\sqrt[3]{bx}}{\sqrt[3]{a+1}}+1\right)}{\left(\frac{2\sqrt[3]{bx}}{\sqrt[3]{a+1}}+1\right)^{-3}}}{\sqrt[3]{b}}}{3(a+1)^{2/3}} - \\
& \quad \frac{\log\left(\sqrt[3]{a+1}-\sqrt[3]{bx}\right)}{3(a+1)^{2/3}\sqrt[3]{b}} \\
& \quad \downarrow 217 \\
& \frac{\frac{1}{2} \int \frac{2\sqrt[3]{bx}+\sqrt[3]{a+1}}{b^{2/3}x^2+\sqrt[3]{a+1}\sqrt[3]{bx+(a+1)^{2/3}}} dx + \frac{\sqrt{3} \arctan\left(\frac{\frac{2\sqrt[3]{bx}}{\sqrt[3]{a+1}}+1}{\sqrt{3}}\right)}{\sqrt[3]{b}}}{3(a+1)^{2/3}} - \frac{\log\left(\sqrt[3]{a+1}-\sqrt[3]{bx}\right)}{3(a+1)^{2/3}\sqrt[3]{b}} \\
& \quad \downarrow 1103
\end{aligned}$$

$$\frac{\sqrt{3} \arctan\left(\frac{\frac{2\sqrt[3]{bx}+1}{\sqrt[3]{a+1}}}{\sqrt{3}}\right)}{\sqrt[3]{b}} + \frac{\log\left(\frac{\sqrt[3]{a+1}\sqrt[3]{bx}+(a+1)^{2/3}+b^{2/3}x^2}{2\sqrt[3]{b}}\right)}{3(a+1)^{2/3}} - \frac{\log\left(\sqrt[3]{a+1}-\sqrt[3]{bx}\right)}{3(a+1)^{2/3}\sqrt[3]{b}}$$

input `Int[(1 + a - b*x^3)^(-1), x]`

output `-1/3*Log[(1 + a)^(1/3) - b^(1/3)*x]/((1 + a)^(2/3)*b^(1/3)) + ((Sqrt[3]*ArcTan[(1 + (2*b^(1/3)*x)/(1 + a)^(1/3))/Sqrt[3]])/b^(1/3) + Log[(1 + a)^(2/3) + (1 + a)^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/(2*b^(1/3)))/(3*(1 + a)^(2/3))`

Defintions of rubi rules used

rule 16 `Int[(c_)/((a_) + (b_)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 750 `Int[((a_) + (b_)*(x_)^3)^(-1), x_Symbol] := Simp[1/(3*Rt[a, 3]^2) Int[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + Simp[1/(3*Rt[a, 3]^2) Int[(2*Rt[a, 3] - Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /; FreeQ[{a, b}, x]`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

rule 1142

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c)
Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.44 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.24

method	result	size
risch	$-\frac{\sum_{R=\text{RootOf}(bZ^3-a-1)} \frac{\ln(x-R)}{-R^2}}{3b}$	30
default	$-\frac{\ln\left(x - \left(\frac{1+a}{b}\right)^{\frac{1}{3}}\right)}{3b\left(\frac{1+a}{b}\right)^{\frac{2}{3}}} + \frac{\ln\left(x^2 + \left(\frac{1+a}{b}\right)^{\frac{1}{3}}x + \left(\frac{1+a}{b}\right)^{\frac{2}{3}}\right)}{6b\left(\frac{1+a}{b}\right)^{\frac{2}{3}}} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2x}{\left(\frac{1+a}{b}\right)^{\frac{1}{3}}+1\right)}\right)}{3b\left(\frac{1+a}{b}\right)^{\frac{2}{3}}}$	106

input

```
int(1/(-b*x^3+a+1),x,method=_RETURNVERBOSE)
```

output

```
-1/3/b*sum(1/_R^2*ln(x-_R),_R=RootOf(_Z^3*b-a-1))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 210 vs. $2(90) = 180$.

Time = 0.10 (sec) , antiderivative size = 467, normalized size of antiderivative = 3.77

$$\int \frac{1}{1+a-bx^3} dx = \text{Too large to display}$$

input `integrate(1/(-b*x^3+a+1),x, algorithm="fricas")`

output `[1/6*(3*sqrt(1/3)*(a + 1)*b*sqrt((-a^2 + 2*a + 1)*b)^(1/3)/b)*log((2*(a + 1)*b*x^3 + 3*(-a^2 + 2*a + 1)*b)^(1/3)*(a + 1)*x + a^2 + 3*sqrt(1/3)*(2*(a + 1)*b*x^2 - (-a^2 + 2*a + 1)*b)^(2/3)*x + (-a^2 + 2*a + 1)*b)^(1/3)*(a + 1))*sqrt((-a^2 + 2*a + 1)*b)^(1/3)/b + 2*a + 1)/(b*x^3 - a - 1)) + (-a^2 + 2*a + 1)*b)^(2/3)*log((a + 1)*b*x^2 + (-a^2 + 2*a + 1)*b)^(2/3)*x - (-a^2 + 2*a + 1)*b)^(1/3)*(a + 1)) - 2*(-a^2 + 2*a + 1)*b)^(2/3)*log((a + 1)*b*x - (-a^2 + 2*a + 1)*b)^(2/3)))/((a^2 + 2*a + 1)*b), 1/6*(6*sqrt(1/3)*(a + 1)*b*sqrt(-(-a^2 + 2*a + 1)*b)^(1/3)/b)*arctan(sqrt(1/3)*(2*(-a^2 + 2*a + 1)*b)^(2/3)*x - (-a^2 + 2*a + 1)*b)^(1/3)*(a + 1))*sqrt(-(-a^2 + 2*a + 1)*b)^(1/3)/b)/(a^2 + 2*a + 1)) + (-a^2 + 2*a + 1)*b)^(2/3)*log((a + 1)*b*x^2 + (-a^2 + 2*a + 1)*b)^(2/3)*x - (-a^2 + 2*a + 1)*b)^(1/3)*(a + 1)) - 2*(-a^2 + 2*a + 1)*b)^(2/3)*log((a + 1)*b*x - (-a^2 + 2*a + 1)*b)^(2/3)))/((a^2 + 2*a + 1)*b)]`

Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.27

$$\int \frac{1}{1+a-bx^3} dx = -\text{RootSum}(t^3 \cdot (27a^2b + 54ab + 27b) - 1, (t \mapsto t \log(-3ta - 3t + x)))$$

input `integrate(1/(-b*x**3+a+1),x)`

output `-RootSum(_t**3*(27*a**2*b + 54*a*b + 27*b) - 1, Lambda(_t, _t*log(-3*_t*a - 3*_t + x)))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.91

$$\int \frac{1}{1+a-bx^3} dx = \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(2x+\left(\frac{a+1}{b}\right)^{\frac{1}{3}}\right)}{3\left(\frac{a+1}{b}\right)^{\frac{1}{3}}}\right)}{3b\left(\frac{a+1}{b}\right)^{\frac{2}{3}}} + \frac{\log\left(x^2+x\left(\frac{a+1}{b}\right)^{\frac{1}{3}}+\left(\frac{a+1}{b}\right)^{\frac{2}{3}}\right)}{6b\left(\frac{a+1}{b}\right)^{\frac{2}{3}}} - \frac{\log\left(x-\left(\frac{a+1}{b}\right)^{\frac{1}{3}}\right)}{3b\left(\frac{a+1}{b}\right)^{\frac{2}{3}}}$$

input `integrate(1/(-b*x^3+a+1),x, algorithm="maxima")`

output

```
1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + ((a + 1)/b)^(1/3))/((a + 1)/b)^(1/3))
)/(b*((a + 1)/b)^(2/3)) + 1/6*log(x^2 + x*((a + 1)/b)^(1/3) + ((a + 1)/b)^(2/3))
)/(b*((a + 1)/b)^(2/3)) - 1/3*log(x - ((a + 1)/b)^(1/3))/(b*((a + 1)/
b)^(2/3))
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.06

$$\int \frac{1}{1+a-bx^3} dx = \frac{(ab^2+b^2)^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}\left(2x+\left(\frac{a+1}{b}\right)^{\frac{1}{3}}\right)}{3\left(\frac{a+1}{b}\right)^{\frac{1}{3}}}\right)}{\sqrt{3}ab + \sqrt{3}b} + \frac{(ab^2+b^2)^{\frac{1}{3}} \log\left(x^2+x\left(\frac{a+1}{b}\right)^{\frac{1}{3}}+\left(\frac{a+1}{b}\right)^{\frac{2}{3}}\right)}{6(ab+b)} - \frac{\left(\frac{a+1}{b}\right)^{\frac{1}{3}} \log\left(\left|x-\left(\frac{a+1}{b}\right)^{\frac{1}{3}}\right|\right)}{3(a+1)}$$

input `integrate(1/(-b*x^3+a+1),x, algorithm="giac")`

output

```
(a*b^2 + b^2)^(1/3)*arctan(1/3*sqrt(3)*(2*x + ((a + 1)/b)^(1/3))/((a + 1)/
b)^(1/3))/(sqrt(3)*a*b + sqrt(3)*b) + 1/6*(a*b^2 + b^2)^(1/3)*log(x^2 + x*
((a + 1)/b)^(1/3) + ((a + 1)/b)^(2/3))/(a*b + b) - 1/3*((a + 1)/b)^(1/3)*l
og(abs(x - ((a + 1)/b)^(1/3)))/(a + 1)
```

Mupad [B] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.33

$$\int \frac{1}{1 + a - bx^3} dx = \frac{\ln \left(3b^2 x + \frac{9ab^2 + 9b^2}{3(-b)^{1/3}(a+1)^{2/3}} \right)}{3(-b)^{1/3}(a+1)^{2/3}} + \frac{\ln \left(3b^2 x + \frac{(9ab^2 + 9b^2)(-1 + \sqrt{3}i)}{6(-b)^{1/3}(a+1)^{2/3}} \right) (-1 + \sqrt{3}i)}{6(-b)^{1/3}(a+1)^{2/3}} - \frac{\ln \left(3b^2 x - \frac{(9ab^2 + 9b^2)(1 + \sqrt{3}i)}{6(-b)^{1/3}(a+1)^{2/3}} \right) (1 + \sqrt{3}i)}{6(-b)^{1/3}(a+1)^{2/3}}$$

input

```
int(1/(a - b*x^3 + 1),x)
```

output

```
log(3*b^2*x + (9*a*b^2 + 9*b^2)/(3*(-b)^(1/3)*(a + 1)^(2/3)))/(3*(-b)^(1/3)
)*(a + 1)^(2/3)) + (log(3*b^2*x + ((9*a*b^2 + 9*b^2)*(3^(1/2)*1i - 1))/(6*
(-b)^(1/3)*(a + 1)^(2/3)))*(3^(1/2)*1i - 1))/(6*(-b)^(1/3)*(a + 1)^(2/3))
- (log(3*b^2*x - ((9*a*b^2 + 9*b^2)*(3^(1/2)*1i + 1))/(6*(-b)^(1/3)*(a + 1
)^(2/3)))*(3^(1/2)*1i + 1))/(6*(-b)^(1/3)*(a + 1)^(2/3))
```

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.62

$$\int \frac{1}{1 + a - bx^3} dx = \frac{2\sqrt{3} \operatorname{atan} \left(\frac{(a+1)^{\frac{1}{3}} + 2b^{\frac{1}{3}}x}{(a+1)^{\frac{1}{3}}\sqrt{3}} \right) + \log \left((a+1)^{\frac{2}{3}} + b^{\frac{1}{3}}(a+1)^{\frac{1}{3}}x + b^{\frac{2}{3}}x^2 \right) - 2 \log \left((a+1)^{\frac{1}{3}} - b^{\frac{1}{3}}x \right)}{6(a+1)^{\frac{2}{3}}b^{\frac{1}{3}}}$$

input `int(1/(-b*x^3+a+1),x)`

output
$$\frac{((a + 1)^{1/3} * (2 * \sqrt{3}) * \operatorname{atan}((a + 1)^{1/3} + 2 * b^{1/3} * x) / ((a + 1)^{1/3} * \sqrt{3})) + \log((a + 1)^{2/3} + b^{1/3} * (a + 1)^{1/3} * x + b^{2/3} * x^2) - 2 * \log((a + 1)^{1/3} - b^{1/3} * x))}{6 * b^{1/3} * (a + 1)}$$

3.11 $\int \frac{1}{-1+a+bx^3} dx$

Optimal result	117
Mathematica [A] (verified)	118
Rubi [A] (verified)	118
Maple [C] (verified)	121
Fricas [A] (verification not implemented)	122
Sympy [A] (verification not implemented)	122
Maxima [F(-2)]	123
Giac [A] (verification not implemented)	123
Mupad [B] (verification not implemented)	124
Reduce [B] (verification not implemented)	124

Optimal result

Integrand size = 10, antiderivative size = 139

$$\int \frac{1}{-1+a+bx^3} dx = -\frac{\arctan\left(\frac{1+\frac{2\sqrt[3]{b}x}{\sqrt[3]{1-a}}}{\sqrt{3}}\right)}{\sqrt{3}(1-a)^{2/3}\sqrt[3]{b}} + \frac{\log\left(\sqrt[3]{1-a}-\sqrt[3]{bx}\right)}{3(1-a)^{2/3}\sqrt[3]{b}} - \frac{\log\left((1-a)^{2/3}+\sqrt[3]{1-a}\sqrt[3]{bx}+b^{2/3}x^2\right)}{6(1-a)^{2/3}\sqrt[3]{b}}$$

output

```
-1/3*arctan(1/3*(1+2*b^(1/3)*x/(1-a)^(1/3))*3^(1/2))*3^(1/2)/(1-a)^(2/3)/b
^(1/3)+1/3*ln((1-a)^(1/3)-b^(1/3)*x)/(1-a)^(2/3)/b^(1/3)-1/6*ln((1-a)^(2/3)
)+(1-a)^(1/3)*b^(1/3)*x+b^(2/3)*x^2)/(1-a)^(2/3)/b^(1/3)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.73

$$\int \frac{1}{-1 + a + bx^3} dx$$

$$= \frac{2\sqrt{3} \arctan\left(\frac{-1 + \frac{2\sqrt[3]{bx}}{\sqrt[3]{-1+a}}}{\sqrt[3]{-1+a}}\right) + 2 \log\left(\sqrt[3]{-1+a} + \sqrt[3]{bx}\right) - \log\left((-1+a)^{2/3} - \sqrt[3]{-1+a}\sqrt[3]{bx} + b^{2/3}x^2\right)}{6(-1+a)^{2/3}\sqrt[3]{b}}$$

input `Integrate[(-1 + a + b*x^3)^(-1),x]`

output `(2*sqrt[3]*ArcTan[(-1 + (2*b^(1/3)*x)/(-1 + a)^(1/3))/sqrt[3]] + 2*Log[(-1 + a)^(1/3) + b^(1/3)*x] - Log[(-1 + a)^(2/3) - (-1 + a)^(1/3)*b^(1/3)*x + b^(2/3)*x^2])/(6*(-1 + a)^(2/3)*b^(1/3))`

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.96, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {750, 16, 25, 1142, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{a + bx^3 - 1} dx$$

$$\downarrow 750$$

$$\int -\frac{\sqrt[3]{bx+2}\sqrt[3]{1-a}}{b^{2/3}x^2 + \sqrt[3]{1-a}\sqrt[3]{bx+(1-a)^{2/3}}} dx + \int \frac{1}{\sqrt[3]{bx}-\sqrt[3]{1-a}} dx$$

$$\downarrow 16$$

$$\begin{aligned}
& \frac{\int -\frac{\sqrt[3]{bx+2}\sqrt[3]{1-a}}{b^{2/3}x^2+\sqrt[3]{1-a}\sqrt[3]{bx+(1-a)^{2/3}}}dx + \frac{\log\left(\sqrt[3]{1-a}-\sqrt[3]{bx}\right)}{3(1-a)^{2/3}\sqrt[3]{b}}}{3(1-a)^{2/3}} \\
& \quad \downarrow 25 \\
& \frac{\log\left(\sqrt[3]{1-a}-\sqrt[3]{bx}\right)}{3(1-a)^{2/3}\sqrt[3]{b}} - \frac{\int \frac{\sqrt[3]{bx+2}\sqrt[3]{1-a}}{b^{2/3}x^2+\sqrt[3]{1-a}\sqrt[3]{bx+(1-a)^{2/3}}}dx}{3(1-a)^{2/3}} \\
& \quad \downarrow 1142 \\
& \frac{\log\left(\sqrt[3]{1-a}-\sqrt[3]{bx}\right)}{3(1-a)^{2/3}\sqrt[3]{b}} - \\
& \frac{\frac{3}{2}\sqrt[3]{1-a} \int \frac{1}{b^{2/3}x^2+\sqrt[3]{1-a}\sqrt[3]{bx+(1-a)^{2/3}}}dx + \frac{\int \frac{\sqrt[3]{b}\left(2\sqrt[3]{bx}+\sqrt[3]{1-a}\right)}{b^{2/3}x^2+\sqrt[3]{1-a}\sqrt[3]{bx+(1-a)^{2/3}}}dx}{2\sqrt[3]{b}}}{3(1-a)^{2/3}} \\
& \quad \downarrow 27 \\
& \frac{\log\left(\sqrt[3]{1-a}-\sqrt[3]{bx}\right)}{3(1-a)^{2/3}\sqrt[3]{b}} - \\
& \frac{\frac{3}{2}\sqrt[3]{1-a} \int \frac{1}{b^{2/3}x^2+\sqrt[3]{1-a}\sqrt[3]{bx+(1-a)^{2/3}}}dx + \frac{1}{2} \int \frac{2\sqrt[3]{bx}+\sqrt[3]{1-a}}{b^{2/3}x^2+\sqrt[3]{1-a}\sqrt[3]{bx+(1-a)^{2/3}}}dx}{3(1-a)^{2/3}} \\
& \quad \downarrow 1082 \\
& \frac{\log\left(\sqrt[3]{1-a}-\sqrt[3]{bx}\right)}{3(1-a)^{2/3}\sqrt[3]{b}} - \\
& \frac{\frac{1}{2} \int \frac{2\sqrt[3]{bx}+\sqrt[3]{1-a}}{b^{2/3}x^2+\sqrt[3]{1-a}\sqrt[3]{bx+(1-a)^{2/3}}}dx - \frac{3 \int \frac{1}{\left(\frac{2\sqrt[3]{bx}}{\sqrt[3]{1-a}}+1\right)^2}d\left(\frac{2\sqrt[3]{bx}}{\sqrt[3]{1-a}}+1\right)}{\left(\frac{2\sqrt[3]{bx}}{\sqrt[3]{1-a}}+1\right)^{-3}}}{\sqrt[3]{b}}}{3(1-a)^{2/3}} \\
& \quad \downarrow 217 \\
& \frac{\log\left(\sqrt[3]{1-a}-\sqrt[3]{bx}\right)}{3(1-a)^{2/3}\sqrt[3]{b}} - \frac{\frac{1}{2} \int \frac{2\sqrt[3]{bx}+\sqrt[3]{1-a}}{b^{2/3}x^2+\sqrt[3]{1-a}\sqrt[3]{bx+(1-a)^{2/3}}}dx + \frac{\sqrt{3} \arctan\left(\frac{\frac{2\sqrt[3]{bx}}{\sqrt[3]{1-a}}+1}{\sqrt{3}}\right)}{\sqrt[3]{b}}}{3(1-a)^{2/3}}
\end{aligned}$$

$$\frac{\log\left(\sqrt[3]{1-a}-\sqrt[3]{bx}\right)}{3(1-a)^{2/3}\sqrt[3]{b}} - \frac{\sqrt{3} \arctan\left(\frac{\frac{2\sqrt[3]{bx}+1}{\sqrt[3]{1-a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}} + \frac{\log\left(\sqrt[3]{1-a}\sqrt[3]{bx}+(1-a)^{2/3}+b^{2/3}x^2\right)}{2\sqrt[3]{b}}}{3(1-a)^{2/3}}$$

input `Int[(-1 + a + b*x^3)^(-1), x]`

output `Log[(1 - a)^(1/3) - b^(1/3)*x]/(3*(1 - a)^(2/3)*b^(1/3)) - ((Sqrt[3]*ArcTan[(1 + (2*b^(1/3)*x)/(1 - a)^(1/3))/Sqrt[3]])/b^(1/3) + Log[(1 - a)^(2/3) + (1 - a)^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/(2*b^(1/3)))/(3*(1 - a)^(2/3))`

Defintions of rubi rules used

rule 16 `Int[(c_)/((a_) + (b_)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 750 `Int[((a_) + (b_)*(x_)^3)^(-1), x_Symbol] := Simp[1/(3*Rt[a, 3]^2) Int[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + Simp[1/(3*Rt[a, 3]^2) Int[(2*Rt[a, 3] - Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /; FreeQ[{a, b}, x]`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1142 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.46 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.20

method	result	size
risch	$\frac{\sum_{-R=\text{RootOf}(bZ^3+a-1)} \frac{\ln(x-R)}{-R^2}}{3b}$	28
default	$\frac{\ln\left(x + \left(\frac{a-1}{b}\right)^{\frac{1}{3}}\right)}{3b\left(\frac{a-1}{b}\right)^{\frac{2}{3}}} - \frac{\ln\left(x^2 - \left(\frac{a-1}{b}\right)^{\frac{1}{3}}x + \left(\frac{a-1}{b}\right)^{\frac{2}{3}}\right)}{6b\left(\frac{a-1}{b}\right)^{\frac{2}{3}}} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2x}{\left(\frac{a-1}{b}\right)^{\frac{1}{3}}}-1\right)}{\frac{\left(\frac{a-1}{b}\right)^{\frac{1}{3}}}{3}}\right)}{3b\left(\frac{a-1}{b}\right)^{\frac{2}{3}}}$	105

input `int(1/(b*x^3+a-1),x,method=_RETURNVERBOSE)`

output `1/3/b*sum(1/_R^2*ln(x-_R),_R=RootOf(_Z^3*b+a-1))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 446, normalized size of antiderivative = 3.21

$$\int \frac{1}{-1 + a + bx^3} dx = \text{Too large to display}$$

input `integrate(1/(b*x^3+a-1),x, algorithm="fricas")`

output `[1/6*(3*sqrt(1/3)*(a - 1)*b*sqrt(-((a^2 - 2*a + 1)*b)^(1/3)/b)*log((2*(a - 1)*b*x^3 - 3*((a^2 - 2*a + 1)*b)^(1/3)*(a - 1)*x - a^2 + 3*sqrt(1/3)*(2*(a - 1)*b*x^2 + ((a^2 - 2*a + 1)*b)^(2/3)*x - ((a^2 - 2*a + 1)*b)^(1/3)*(a - 1))*sqrt(-((a^2 - 2*a + 1)*b)^(1/3)/b + 2*a - 1)/(b*x^3 + a - 1)) - ((a^2 - 2*a + 1)*b)^(2/3)*log((a - 1)*b*x^2 - ((a^2 - 2*a + 1)*b)^(2/3)*x + ((a^2 - 2*a + 1)*b)^(1/3)*(a - 1)) + 2*((a^2 - 2*a + 1)*b)^(2/3)*log((a - 1)*b*x + ((a^2 - 2*a + 1)*b)^(2/3)))/((a^2 - 2*a + 1)*b), 1/6*(6*sqrt(1/3)*(a - 1)*b*sqrt(((a^2 - 2*a + 1)*b)^(1/3)/b)*arctan(sqrt(1/3)*(2*((a^2 - 2*a + 1)*b)^(2/3)*x - ((a^2 - 2*a + 1)*b)^(1/3)*(a - 1))*sqrt(((a^2 - 2*a + 1)*b)^(1/3)/b)/(a^2 - 2*a + 1)) - ((a^2 - 2*a + 1)*b)^(2/3)*log((a - 1)*b*x^2 - ((a^2 - 2*a + 1)*b)^(2/3)*x + ((a^2 - 2*a + 1)*b)^(1/3)*(a - 1)) + 2*((a^2 - 2*a + 1)*b)^(2/3)*log((a - 1)*b*x + ((a^2 - 2*a + 1)*b)^(2/3)))/((a^2 - 2*a + 1)*b)]`

Sympy [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.23

$$\int \frac{1}{-1 + a + bx^3} dx$$

$$= \text{RootSum}(t^3 \cdot (27a^2b - 54ab + 27b) - 1, (t \mapsto t \log(3ta - 3t + x)))$$

input `integrate(1/(b*x**3+a-1),x)`

output `RootSum(_t**3*(27*a**2*b - 54*a*b + 27*b) - 1, Lambda(_t, _t*log(3*_t*a - 3*_t + x)))`

Maxima [F(-2)]

Exception generated.

$$\int \frac{1}{-1 + a + bx^3} dx = \text{Exception raised: ValueError}$$

input `integrate(1/(b*x^3+a-1),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a-1.0>0)', see `assume?` for more details)

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.02

$$\int \frac{1}{-1 + a + bx^3} dx = \frac{(-ab^2 + b^2)^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}\left(2x + \left(-\frac{a-1}{b}\right)^{\frac{1}{3}}\right)}{3\left(-\frac{a-1}{b}\right)^{\frac{1}{3}}}\right)}{\sqrt{3}ab - \sqrt{3}b} + \frac{(-ab^2 + b^2)^{\frac{1}{3}} \log\left(x^2 + x\left(-\frac{a-1}{b}\right)^{\frac{1}{3}} + \left(-\frac{a-1}{b}\right)^{\frac{2}{3}}\right)}{6(ab - b)} - \frac{\left(-\frac{a-1}{b}\right)^{\frac{1}{3}} \log\left(\left|x - \left(-\frac{a-1}{b}\right)^{\frac{1}{3}}\right|\right)}{3(a - 1)}$$

input `integrate(1/(b*x^3+a-1),x, algorithm="giac")`

output `(-a*b^2 + b^2)^(1/3)*arctan(1/3*sqrt(3)*(2*x + (-a - 1)/b)^(1/3))/(-a - 1)/b)^(1/3))/(sqrt(3)*a*b - sqrt(3)*b) + 1/6*(-a*b^2 + b^2)^(1/3)*log(x^2 + x*(-a - 1)/b)^(1/3) + (-a - 1)/b)^(2/3))/(a*b - b) - 1/3*(-a - 1)/b)^(1/3)*log(abs(x - (-a - 1)/b)^(1/3)))/(a - 1)`

Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.99

$$\int \frac{1}{-1 + a + bx^3} dx = \frac{\ln\left(a + b^{1/3} x (a - 1)^{2/3} - 1\right)}{3 b^{1/3} (a - 1)^{2/3}} + \frac{\ln\left(3 b^2 x + \frac{(9 a b^2 - 9 b^2)(-1 + \sqrt{3} i)}{6 b^{1/3} (a - 1)^{2/3}}\right) (-1 + \sqrt{3} i)}{6 b^{1/3} (a - 1)^{2/3}} - \frac{\ln\left(3 b^2 x - \frac{(9 a b^2 - 9 b^2)(1 + \sqrt{3} i)}{6 b^{1/3} (a - 1)^{2/3}}\right) (1 + \sqrt{3} i)}{6 b^{1/3} (a - 1)^{2/3}}$$

input `int(1/(a + b*x^3 - 1),x)`output `log(a + b^(1/3)*x*(a - 1)^(2/3) - 1)/(3*b^(1/3)*(a - 1)^(2/3)) + (log(3*b^2*x + ((9*a*b^2 - 9*b^2)*(3^(1/2)*1i - 1))/(6*b^(1/3)*(a - 1)^(2/3)))*(3^(1/2)*1i - 1))/(6*b^(1/3)*(a - 1)^(2/3)) - (log(3*b^2*x - ((9*a*b^2 - 9*b^2)*(3^(1/2)*1i + 1))/(6*b^(1/3)*(a - 1)^(2/3)))*(3^(1/2)*1i + 1))/(6*b^(1/3)*(a - 1)^(2/3))`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.57

$$\int \frac{1}{-1 + a + bx^3} dx = \frac{-2\sqrt{3} \operatorname{atan}\left(\frac{(a-1)^{1/3} - 2b^{1/3}x}{(a-1)^{1/3}\sqrt{3}}\right) - \log\left((a-1)^{2/3} - b^{1/3}(a-1)^{1/3}x + b^{2/3}x^2\right) + 2\log\left((a-1)^{1/3} + b^{1/3}x\right)}{6(a-1)^{2/3}b^{1/3}}$$

input `int(1/(b*x^3+a-1),x)`output `((a - 1)**(1/3)*(- 2*sqrt(3)*atan(((a - 1)**(1/3) - 2*b**(1/3)*x)/((a - 1)**(1/3)*sqrt(3)))) - log(((a - 1)**(2/3) - b**(1/3)*(a - 1)**(1/3)*x + b**(2/3)*x**2) + 2*log((a - 1)**(1/3) + b**(1/3)*x))/(6*b**(1/3)*(a - 1))`

3.12 $\int \frac{1}{-1+a-bx^3} dx$

Optimal result	125
Mathematica [A] (verified)	126
Rubi [A] (verified)	126
Maple [C] (verified)	130
Fricas [B] (verification not implemented)	130
Sympy [A] (verification not implemented)	131
Maxima [F(-2)]	131
Giac [A] (verification not implemented)	132
Mupad [B] (verification not implemented)	133
Reduce [B] (verification not implemented)	133

Optimal result

Integrand size = 11, antiderivative size = 138

$$\int \frac{1}{-1+a-bx^3} dx = \frac{\arctan\left(\frac{1-\frac{2\sqrt[3]{bx}}{\sqrt[3]{1-a}}}{\sqrt{3}}\right)}{\sqrt{3}(1-a)^{2/3}\sqrt[3]{b}} - \frac{\log\left(\sqrt[3]{1-a} + \sqrt[3]{bx}\right)}{3(1-a)^{2/3}\sqrt[3]{b}} + \frac{\log\left((1-a)^{2/3} - \sqrt[3]{1-a}\sqrt[3]{bx} + b^{2/3}x^2\right)}{6(1-a)^{2/3}\sqrt[3]{b}}$$

output

```
1/3*arctan(1/3*(1-2*b^(1/3)*x/(1-a)^(1/3))*3^(1/2))*3^(1/2)/(1-a)^(2/3)/b^(1/3)-1/3*ln((1-a)^(1/3)+b^(1/3)*x)/(1-a)^(2/3)/b^(1/3)+1/6*ln((1-a)^(2/3)-(1-a)^(1/3)*b^(1/3)*x+b^(2/3)*x^2)/(1-a)^(2/3)/b^(1/3)
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.90

$$\int \frac{1}{-1 + a - bx^3} dx$$

$$= \frac{(-1)^{2/3} \left(-2\sqrt{3} \arctan \left(\frac{-1 + \sqrt[3]{-1} \sqrt[3]{bx}}{\sqrt{3}} \right) - 2 \log \left(\sqrt[3]{-1 + a} + \sqrt[3]{-1} \sqrt[3]{bx} \right) + \log \left((-1 + a)^{2/3} - \sqrt[3]{-1} \right) \right)}{6(-1 + a)^{2/3} \sqrt[3]{b}}$$

input `Integrate[(-1 + a - b*x^3)^(-1),x]`

output

```
((-1)^(2/3)*(-2*Sqrt[3]*ArcTan[(-1 + (2*(-1)^(1/3)*b^(1/3)*x)/(-1 + a)^(1/3)])/Sqrt[3]] - 2*Log[(-1 + a)^(1/3) + (-1)^(1/3)*b^(1/3)*x] + Log[(-1 + a)^(2/3) - (-1)^(1/3)*(-1 + a)^(1/3)*b^(1/3)*x + (-1)^(2/3)*b^(2/3)*x^2])/(6*(-1 + a)^(2/3)*b^(1/3))
```

Rubi [A] (verified)Time = 0.28 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.98, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.818$, Rules used = {750, 16, 25, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{a - bx^3 - 1} dx$$

$$\downarrow 750$$

$$\frac{\int -\frac{2\sqrt[3]{1-a} - \sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{1-a}\sqrt[3]{bx} + (1-a)^{2/3}} dx}{3(1-a)^{2/3}} + \frac{\int \frac{1}{-\sqrt[3]{bx} - \sqrt[3]{1-a}} dx}{3(1-a)^{2/3}}$$

$$\downarrow 16$$

$$\frac{\int -\frac{2\sqrt[3]{1-a}-\sqrt[3]{bx}}{b^{2/3}x^2-\sqrt[3]{1-a}\sqrt[3]{bx+(1-a)^{2/3}}}dx}{3(1-a)^{2/3}} - \frac{\log\left(\sqrt[3]{1-a} + \sqrt[3]{bx}\right)}{3(1-a)^{2/3}\sqrt[3]{b}}$$

↓ 25

$$\frac{\int \frac{2\sqrt[3]{1-a}-\sqrt[3]{bx}}{b^{2/3}x^2-\sqrt[3]{1-a}\sqrt[3]{bx+(1-a)^{2/3}}}dx}{3(1-a)^{2/3}} - \frac{\log\left(\sqrt[3]{1-a} + \sqrt[3]{bx}\right)}{3(1-a)^{2/3}\sqrt[3]{b}}$$

↓ 1142

$$\frac{\frac{3}{2}\sqrt[3]{1-a} \int \frac{1}{b^{2/3}x^2-\sqrt[3]{1-a}\sqrt[3]{bx+(1-a)^{2/3}}}dx - \frac{\int -\frac{\sqrt[3]{b}\left(\sqrt[3]{1-a}-2\sqrt[3]{bx}\right)}{b^{2/3}x^2-\sqrt[3]{1-a}\sqrt[3]{bx+(1-a)^{2/3}}}dx}{2\sqrt[3]{b}}}{\frac{3(1-a)^{2/3} \log\left(\sqrt[3]{1-a} + \sqrt[3]{bx}\right)}{3(1-a)^{2/3}\sqrt[3]{b}}}$$

↓ 25

$$\frac{\frac{3}{2}\sqrt[3]{1-a} \int \frac{1}{b^{2/3}x^2-\sqrt[3]{1-a}\sqrt[3]{bx+(1-a)^{2/3}}}dx + \frac{\int \frac{\sqrt[3]{b}\left(\sqrt[3]{1-a}-2\sqrt[3]{bx}\right)}{b^{2/3}x^2-\sqrt[3]{1-a}\sqrt[3]{bx+(1-a)^{2/3}}}dx}{2\sqrt[3]{b}}}{\frac{3(1-a)^{2/3} \log\left(\sqrt[3]{1-a} + \sqrt[3]{bx}\right)}{3(1-a)^{2/3}\sqrt[3]{b}}}$$

↓ 27

$$\frac{\frac{3}{2}\sqrt[3]{1-a} \int \frac{1}{b^{2/3}x^2-\sqrt[3]{1-a}\sqrt[3]{bx+(1-a)^{2/3}}}dx + \frac{1}{2} \int \frac{\sqrt[3]{1-a}-2\sqrt[3]{bx}}{b^{2/3}x^2-\sqrt[3]{1-a}\sqrt[3]{bx+(1-a)^{2/3}}}dx}{\frac{3(1-a)^{2/3} \log\left(\sqrt[3]{1-a} + \sqrt[3]{bx}\right)}{3(1-a)^{2/3}\sqrt[3]{b}}}$$

↓ 1082

$$\begin{aligned}
& \frac{\frac{1}{2} \int \frac{\sqrt[3]{1-a} - 2\sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{1-a}\sqrt[3]{bx} + (1-a)^{2/3}} dx + \frac{3 \int \frac{1}{\left(1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{1-a}}\right)^2} d\left(1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{1-a}}\right)}{-3\sqrt[3]{1-a}}}{\frac{3(1-a)^{2/3}}{\log\left(\sqrt[3]{1-a} + \sqrt[3]{bx}\right)}} - \frac{\sqrt[3]{b}}{3(1-a)^{2/3}\sqrt[3]{b}} \\
& \quad \downarrow \text{217} \\
& \frac{\frac{1}{2} \int \frac{\sqrt[3]{1-a} - 2\sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{1-a}\sqrt[3]{bx} + (1-a)^{2/3}} dx - \frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{1-a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}}}{\frac{3(1-a)^{2/3}}{\log\left(\sqrt[3]{1-a} + \sqrt[3]{bx}\right)}} - \frac{\log\left(\sqrt[3]{1-a} + \sqrt[3]{bx}\right)}{3(1-a)^{2/3}\sqrt[3]{b}} \\
& \quad \downarrow \text{1103} \\
& \frac{\frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{1-a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}} - \frac{\log\left(-\sqrt[3]{1-a}\sqrt[3]{bx} + (1-a)^{2/3} + b^{2/3}x^2\right)}{2\sqrt[3]{b}}}{\frac{3(1-a)^{2/3}}{\log\left(\sqrt[3]{1-a} + \sqrt[3]{bx}\right)}} - \frac{\log\left(\sqrt[3]{1-a} + \sqrt[3]{bx}\right)}{3(1-a)^{2/3}\sqrt[3]{b}}
\end{aligned}$$

input `Int[(-1 + a - b*x^3)^(-1), x]`

output `-1/3*Log[(1 - a)^(1/3) + b^(1/3)*x]/((1 - a)^(2/3)*b^(1/3)) - (-((Sqrt[3]*ArcTan[(1 - (2*b^(1/3)*x)/(1 - a)^(1/3)]/Sqrt[3])/b^(1/3)) - Log[(1 - a)^(2/3) - (1 - a)^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/(2*b^(1/3)))/(3*(1 - a)^(2/3)))`

Defintions of rubi rules used

rule 16 `Int[(c.)/((a.) + (b.)*(x.)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 750 `Int[((a_) + (b_)*(x_)^3)^(-1), x_Symbol] := Simp[1/(3*Rt[a, 3]^2) Int[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + Simp[1/(3*Rt[a, 3]^2) Int[(2*Rt[a, 3] - Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /; FreeQ[{a, b}, x]`
- rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1142 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.44 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.22

method	result	size
risch	$-\frac{\sum_{R=\text{RootOf}(bZ^3-a+1)} \frac{\ln(x-R)}{R^2}}{3b}$	30
default	$-\frac{\ln\left(x-\left(\frac{a-1}{b}\right)^{\frac{1}{3}}\right)}{3b\left(\frac{a-1}{b}\right)^{\frac{2}{3}}} + \frac{\ln\left(x^2+\left(\frac{a-1}{b}\right)^{\frac{1}{3}}x+\left(\frac{a-1}{b}\right)^{\frac{2}{3}}\right)}{6b\left(\frac{a-1}{b}\right)^{\frac{2}{3}}} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2x}{\left(\frac{a-1}{b}\right)^{\frac{1}{3}}}+1\right)}{\left(\frac{a-1}{b}\right)^{\frac{1}{3}}}\right)}{3b\left(\frac{a-1}{b}\right)^{\frac{2}{3}}}$	106

input `int(1/(-b*x^3+a-1),x,method=_RETURNVERBOSE)`

output `-1/3/b*sum(1/_R^2*ln(x-_R),_R=RootOf(_Z^3*b-a+1))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 210 vs. 2(104) = 208.

Time = 0.08 (sec) , antiderivative size = 467, normalized size of antiderivative = 3.38

$$\int \frac{1}{-1+a-bx^3} dx = \text{Too large to display}$$

input `integrate(1/(-b*x^3+a-1),x, algorithm="fricas")`

output

```
[1/6*(3*sqrt(1/3)*(a - 1)*b*sqrt((-a^2 - 2*a + 1)*b)^(1/3)/b)*log((2*(a - 1)*b*x^3 + 3*(-a^2 - 2*a + 1)*b)^(1/3)*(a - 1)*x + a^2 + 3*sqrt(1/3)*(2*(a - 1)*b*x^2 - (-a^2 - 2*a + 1)*b)^(2/3)*x + (-a^2 - 2*a + 1)*b)^(1/3)*(a - 1))*sqrt((-a^2 - 2*a + 1)*b)^(1/3)/b - 2*a + 1)/(b*x^3 - a + 1)) + (-a^2 - 2*a + 1)*b)^(2/3)*log((a - 1)*b*x^2 + (-a^2 - 2*a + 1)*b)^(2/3)*x - (-a^2 - 2*a + 1)*b)^(1/3)*(a - 1)) - 2*(-a^2 - 2*a + 1)*b)^(2/3)*log((a - 1)*b*x - (-a^2 - 2*a + 1)*b)^(2/3)))/((a^2 - 2*a + 1)*b), 1/6*(6*sqrt(1/3)*(a - 1)*b*sqrt(-(-a^2 - 2*a + 1)*b)^(1/3)/b)*arctan(sqrt(1/3)*(2*(-a^2 - 2*a + 1)*b)^(2/3)*x - (-a^2 - 2*a + 1)*b)^(1/3)*(a - 1))*sqrt(-(-a^2 - 2*a + 1)*b)^(1/3)/b)/(a^2 - 2*a + 1)) + (-a^2 - 2*a + 1)*b)^(2/3)*log((a - 1)*b*x^2 + (-a^2 - 2*a + 1)*b)^(2/3)*x - (-a^2 - 2*a + 1)*b)^(1/3)*(a - 1)) - 2*(-a^2 - 2*a + 1)*b)^(2/3)*log((a - 1)*b*x - (-a^2 - 2*a + 1)*b)^(2/3)))/((a^2 - 2*a + 1)*b)]
```

Sympy [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.25

$$\int \frac{1}{-1 + a - bx^3} dx$$

$$= -\text{RootSum}(t^3 \cdot (27a^2b - 54ab + 27b) - 1, (t \mapsto t \log(-3ta + 3t + x)))$$

input

```
integrate(1/(-b*x**3+a-1),x)
```

output

```
-RootSum(_t**3*(27*a**2*b - 54*a*b + 27*b) - 1, Lambda(_t, _t*log(-3*_t*a + 3*_t + x)))
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{1}{-1 + a - bx^3} dx = \text{Exception raised: ValueError}$$

input

```
integrate(1/(-b*x^3+a-1),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a-1.0>0)', see `assume?` for mor
e details)
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.00

$$\int \frac{1}{-1 + a - bx^3} dx = \frac{(ab^2 - b^2)^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}\left(2x + \left(\frac{a-1}{b}\right)^{\frac{1}{3}}\right)}{3\left(\frac{a-1}{b}\right)^{\frac{1}{3}}}\right)}{\sqrt{3}ab - \sqrt{3}b} + \frac{(ab^2 - b^2)^{\frac{1}{3}} \log\left(x^2 + x\left(\frac{a-1}{b}\right)^{\frac{1}{3}} + \left(\frac{a-1}{b}\right)^{\frac{2}{3}}\right)}{6(ab - b)} - \frac{\left(\frac{a-1}{b}\right)^{\frac{1}{3}} \log\left(\left|x - \left(\frac{a-1}{b}\right)^{\frac{1}{3}}\right|\right)}{3(a - 1)}$$

input

```
integrate(1/(-b*x^3+a-1),x, algorithm="giac")
```

output

```
(a*b^2 - b^2)^(1/3)*arctan(1/3*sqrt(3)*(2*x + ((a - 1)/b)^(1/3))/((a - 1)/
b)^(1/3))/(sqrt(3)*a*b - sqrt(3)*b) + 1/6*(a*b^2 - b^2)^(1/3)*log(x^2 + x*
((a - 1)/b)^(1/3) + ((a - 1)/b)^(2/3))/(a*b - b) - 1/3*((a - 1)/b)^(1/3)*l
og(abs(x - ((a - 1)/b)^(1/3)))/(a - 1)
```

Mupad [B] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.20

$$\int \frac{1}{-1 + a - bx^3} dx = \frac{\ln\left(3b^2x + \frac{9ab^2 - 9b^2}{3(-b)^{1/3}(a-1)^{2/3}}\right)}{3(-b)^{1/3}(a-1)^{2/3}} + \frac{\ln\left(3b^2x + \frac{(9ab^2 - 9b^2)(-1 + \sqrt{3}i)}{6(-b)^{1/3}(a-1)^{2/3}}\right)(-1 + \sqrt{3}i)}{6(-b)^{1/3}(a-1)^{2/3}} - \frac{\ln\left(3b^2x - \frac{(9ab^2 - 9b^2)(1 + \sqrt{3}i)}{6(-b)^{1/3}(a-1)^{2/3}}\right)(1 + \sqrt{3}i)}{6(-b)^{1/3}(a-1)^{2/3}}$$

input `int(-1/(b*x^3 - a + 1),x)`output `log(3*b^2*x + (9*a*b^2 - 9*b^2)/(3*(-b)^(1/3)*(a - 1)^(2/3)))/(3*(-b)^(1/3)*(a - 1)^(2/3)) + (log(3*b^2*x + ((9*a*b^2 - 9*b^2)*(3^(1/2)*1i - 1))/(6*(-b)^(1/3)*(a - 1)^(2/3)))*(3^(1/2)*1i - 1))/(6*(-b)^(1/3)*(a - 1)^(2/3)) - (log(3*b^2*x - ((9*a*b^2 - 9*b^2)*(3^(1/2)*1i + 1))/(6*(-b)^(1/3)*(a - 1)^(2/3)))*(3^(1/2)*1i + 1))/(6*(-b)^(1/3)*(a - 1)^(2/3))`**Reduce [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.56

$$\int \frac{1}{-1 + a - bx^3} dx = \frac{2\sqrt{3} \operatorname{atan}\left(\frac{(a-1)^{\frac{1}{3}} + 2b^{\frac{1}{3}}x}{(a-1)^{\frac{1}{3}}\sqrt{3}}\right) + \log\left((a-1)^{\frac{2}{3}} + b^{\frac{1}{3}}(a-1)^{\frac{1}{3}}x + b^{\frac{2}{3}}x^2\right) - 2\log\left((a-1)^{\frac{1}{3}} - b^{\frac{1}{3}}x\right)}{6(a-1)^{\frac{2}{3}}b^{\frac{1}{3}}}$$

input `int(1/(-b*x^3+a-1),x)`output `((a - 1)**(1/3)*(2*sqrt(3)*atan(((a - 1)**(1/3) + 2*b**(1/3)*x)/((a - 1)**(1/3)*sqrt(3)))) + log((a - 1)**(2/3) + b**(1/3)*(a - 1)**(1/3)*x + b**(2/3)*x**2) - 2*log((a - 1)**(1/3) - b**(1/3)*x))/(6*b**(1/3)*(a - 1))`

3.13 $\int (a + bx^3)^{5/2} dx$

Optimal result	134
Mathematica [C] (verified)	135
Rubi [A] (verified)	135
Maple [A] (verified)	137
Fricas [A] (verification not implemented)	138
Sympy [A] (verification not implemented)	138
Maxima [F]	139
Giac [F]	139
Mupad [B] (verification not implemented)	139
Reduce [F]	140

Optimal result

Integrand size = 11, antiderivative size = 265

$$\int (a + bx^3)^{5/2} dx = \frac{54}{187}a^2x\sqrt{a + bx^3} + \frac{30}{187}ax(a + bx^3)^{3/2} + \frac{2}{17}x(a + bx^3)^{5/2} + \frac{54 \cdot 3^{3/4} \sqrt{2 + \sqrt{3}} a^3 (\sqrt[3]{a} + \sqrt[3]{bx}) \sqrt{\frac{a^{2/3} - \sqrt[3]{a} \sqrt[3]{bx} + b^{2/3} x^2}{((1 + \sqrt{3}) \sqrt[3]{a} + \sqrt[3]{bx})^2}} \operatorname{EllipticF}\left(\arcsin\left(\frac{(1 - \sqrt{3}) \sqrt[3]{a} + \sqrt[3]{bx}}{(1 + \sqrt{3}) \sqrt[3]{a} + \sqrt[3]{bx}}\right), -7 - 4\sqrt{3}\right)}{187 \sqrt[3]{b} \sqrt{\frac{\sqrt[3]{a} (\sqrt[3]{a} + \sqrt[3]{bx})}{((1 + \sqrt{3}) \sqrt[3]{a} + \sqrt[3]{bx})^2}} \sqrt{a + bx^3}}$$

```
output 54/187*a^2*x*(b*x^3+a)^(1/2)+30/187*a*x*(b*x^3+a)^(3/2)+2/17*x*(b*x^3+a)^(5/2)+54/187*3^(3/4)*(1/2*6^(1/2)+1/2*2^(1/2))*a^3*(a^(1/3)+b^(1/3)*x)*((a^(2/3)-a^(1/3)*b^(1/3)*x+b^(2/3)*x^2)/((1+3^(1/2))*a^(1/3)+b^(1/3)*x)^2)^(1/2)*EllipticF(((1-3^(1/2))*a^(1/3)+b^(1/3)*x)/((1+3^(1/2))*a^(1/3)+b^(1/3)*x),I*3^(1/2)+2*I)/b^(1/3)/(a^(1/3)*(a^(1/3)+b^(1/3)*x)/((1+3^(1/2))*a^(1/3)+b^(1/3)*x)^2)^(1/2)/(b*x^3+a)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 4.39 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.18

$$\int (a + bx^3)^{5/2} dx = \frac{a^2 x \sqrt{a + bx^3} \operatorname{Hypergeometric2F1}\left(-\frac{5}{2}, \frac{1}{3}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{\sqrt{1 + \frac{bx^3}{a}}}$$

input `Integrate[(a + b*x^3)^(5/2), x]`

output `(a^2*x*Sqrt[a + b*x^3]*Hypergeometric2F1[-5/2, 1/3, 4/3, -((b*x^3)/a)]/Sqrt[1 + (b*x^3)/a])`

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 271, normalized size of antiderivative = 1.02, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {748, 748, 748, 759}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + bx^3)^{5/2} dx \\ & \quad \downarrow 748 \\ & \frac{15}{17}a \int (bx^3 + a)^{3/2} dx + \frac{2}{17}x(a + bx^3)^{5/2} \\ & \quad \downarrow 748 \\ & \frac{15}{17}a \left(\frac{9}{11}a \int \sqrt{bx^3 + a} dx + \frac{2}{11}x(a + bx^3)^{3/2} \right) + \frac{2}{17}x(a + bx^3)^{5/2} \\ & \quad \downarrow 748 \\ & \frac{15}{17}a \left(\frac{9}{11}a \left(\frac{3}{5}a \int \frac{1}{\sqrt{bx^3 + a}} dx + \frac{2}{5}x\sqrt{a + bx^3} \right) + \frac{2}{11}x(a + bx^3)^{3/2} \right) + \frac{2}{17}x(a + bx^3)^{5/2} \end{aligned}$$

$$\begin{array}{c} \downarrow 759 \\ \frac{15}{17}a \left(\frac{9}{11}a \left(\frac{2 \cdot 3^{3/4} \sqrt{2 + \sqrt{3}} a (\sqrt[3]{a} + \sqrt[3]{bx}) \sqrt{\frac{a^{2/3} - \sqrt[3]{a} \sqrt[3]{bx} + b^{2/3} x^2}{((1 + \sqrt{3}) \sqrt[3]{a} + \sqrt[3]{bx})^2}} \operatorname{EllipticF} \left(\arcsin \left(\frac{\sqrt[3]{bx} + (1 - \sqrt{3}) \sqrt[3]{a}}{\sqrt[3]{bx} + (1 + \sqrt{3}) \sqrt[3]{a}} \right), -7 - \dots \right)}{5 \sqrt[3]{b} \sqrt{\frac{\sqrt[3]{a} (\sqrt[3]{a} + \sqrt[3]{bx})}{((1 + \sqrt{3}) \sqrt[3]{a} + \sqrt[3]{bx})^2}} \sqrt{a + bx^3}} \right) \right) \\ \frac{2}{17}x(a + bx^3)^{5/2} \end{array}$$

input `Int[(a + b*x^3)^(5/2), x]`

output `(2*x*(a + b*x^3)^(5/2))/17 + (15*a*((2*x*(a + b*x^3)^(3/2))/11 + (9*a*((2*x*Sqrt[a + b*x^3])/5 + (2*3^(3/4)*Sqrt[2 + Sqrt[3]]*a*(a^(1/3) + b^(1/3)*x)*Sqrt[(a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/((1 + Sqrt[3])*a^(1/3) + b^(1/3)*x)^2)*EllipticF[ArcSin[((1 - Sqrt[3])*a^(1/3) + b^(1/3)*x)/((1 + Sqrt[3])*a^(1/3) + b^(1/3)*x)], -7 - 4*Sqrt[3]])/(5*b^(1/3)*Sqrt[(a^(1/3) * (a^(1/3) + b^(1/3)*x))/((1 + Sqrt[3])*a^(1/3) + b^(1/3)*x)^2]*Sqrt[a + b*x^3])))/11))/17`

Defintions of rubi rules used

rule 748 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p + 1)), x] + Simp[a*n*(p/(n*p + 1)) Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || LtQ[Denominator[p + 1/n], Denominator[p]])`

rule 759 `Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[2*Sqrt[2 + Sqrt[3]]*(s + r*x)*(Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 + Sqrt[3])*s + r*x)^2]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[s*(s + r*x)/((1 + Sqrt[3])*s + r*x)^2]))*EllipticF[ArcSin[((1 - Sqrt[3])*s + r*x)/((1 + Sqrt[3])*s + r*x)], -7 - 4*Sqrt[3]], x] /; FreeQ[{a, b}, x] & & PosQ[a]`

Maple [A] (verified)

Time = 2.53 (sec) , antiderivative size = 320, normalized size of antiderivative = 1.21

method	result
risch	$\frac{2x(11b^2x^6+37abx^3+53a^2)\sqrt{bx^3+a}}{187} - \frac{54ia^3\sqrt{3}(-ab^2)^{\frac{1}{3}}\sqrt{\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b}-\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}{(-ab^2)^{\frac{1}{3}}}}{\sqrt{\frac{x-\frac{(-ab^2)^{\frac{1}{3}}}{b}}{-\frac{3(-ab^2)^{\frac{1}{3}}}{2b}+\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}}}}$
default	$\frac{2b^2x^7\sqrt{bx^3+a}}{17} + \frac{74abx^4\sqrt{bx^3+a}}{187} + \frac{106a^2x\sqrt{bx^3+a}}{187} - \frac{54ia^3\sqrt{3}(-ab^2)^{\frac{1}{3}}\sqrt{\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b}-\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}{(-ab^2)^{\frac{1}{3}}}}{\sqrt{\frac{x-\frac{(-ab^2)^{\frac{1}{3}}}{b}}{-\frac{3(-ab^2)^{\frac{1}{3}}}{2b}+\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}}}}$
elliptic	$\frac{2b^2x^7\sqrt{bx^3+a}}{17} + \frac{74abx^4\sqrt{bx^3+a}}{187} + \frac{106a^2x\sqrt{bx^3+a}}{187} - \frac{54ia^3\sqrt{3}(-ab^2)^{\frac{1}{3}}\sqrt{\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b}-\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}{(-ab^2)^{\frac{1}{3}}}}{\sqrt{\frac{x-\frac{(-ab^2)^{\frac{1}{3}}}{b}}{-\frac{3(-ab^2)^{\frac{1}{3}}}{2b}+\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}}}}$

```
input int((b*x^3+a)^(5/2), x, method=_RETURNVERBOSE)
```

```
output 2/187*x*(11*b^2*x^6+37*a*b*x^3+53*a^2)*(b*x^3+a)^(1/2)-54/187*I*a^3*3^(1/2)/b*(-a*b^2)^(1/3)*(I*(x+1/2/b*(-a*b^2)^(1/3)-1/2*I*3^(1/2)/b*(-a*b^2)^(1/3))*3^(1/2)*b/(-a*b^2)^(1/3))^(1/2)*((x-1/b*(-a*b^2)^(1/3))/(-3/2/b*(-a*b^2)^(1/3)+1/2*I*3^(1/2)/b*(-a*b^2)^(1/3)))^(1/2)*(-I*(x+1/2/b*(-a*b^2)^(1/3)+1/2*I*3^(1/2)/b*(-a*b^2)^(1/3))*3^(1/2)*b/(-a*b^2)^(1/3))^(1/2)/(b*x^3+a)^(1/2)*EllipticF(1/3*3^(1/2)*(I*(x+1/2/b*(-a*b^2)^(1/3)-1/2*I*3^(1/2)/b*(-a*b^2)^(1/3))*3^(1/2)*b/(-a*b^2)^(1/3))^(1/2), (I*3^(1/2)/b*(-a*b^2)^(1/3)/(-3/2/b*(-a*b^2)^(1/3)+1/2*I*3^(1/2)/b*(-a*b^2)^(1/3)))^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.22

$$\int (a + bx^3)^{5/2} dx = \frac{2 \left(81 a^3 \sqrt{b} \text{weierstrassPInverse}(0, -\frac{4a}{b}, x) + (11 b^3 x^7 + 37 ab^2 x^4 + 53 a^2 bx) \sqrt{bx^3 + a} \right)}{187 b}$$

input `integrate((b*x^3+a)^(5/2),x, algorithm="fricas")`output `2/187*(81*a^3*sqrt(b)*weierstrassPInverse(0, -4*a/b, x) + (11*b^3*x^7 + 37*a*b^2*x^4 + 53*a^2*b*x)*sqrt(b*x^3 + a))/b`**Sympy [A] (verification not implemented)**

Time = 0.54 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.14

$$\int (a + bx^3)^{5/2} dx = \frac{a^{5/2} x \Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\begin{matrix} -\frac{5}{2}, \frac{1}{3} \\ \frac{4}{3} \end{matrix} \middle| \frac{bx^3 e^{i\pi}}{a} \right)}{3 \Gamma\left(\frac{4}{3}\right)}$$

input `integrate((b*x**3+a)**(5/2),x)`output `a**(5/2)*x*gamma(1/3)*hyper((-5/2, 1/3), (4/3,), b*x**3*exp_polar(I*pi)/a)/(3*gamma(4/3))`

Maxima [F]

$$\int (a + bx^3)^{5/2} dx = \int (bx^3 + a)^{5/2} dx$$

input `integrate((b*x^3+a)^(5/2),x, algorithm="maxima")`

output `integrate((b*x^3 + a)^(5/2), x)`

Giac [F]

$$\int (a + bx^3)^{5/2} dx = \int (bx^3 + a)^{5/2} dx$$

input `integrate((b*x^3+a)^(5/2),x, algorithm="giac")`

output `integrate((b*x^3 + a)^(5/2), x)`

Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.14

$$\int (a + bx^3)^{5/2} dx = \frac{x (bx^3 + a)^{5/2} {}_2F_1\left(-\frac{5}{2}, \frac{1}{3}; \frac{4}{3}; -\frac{bx^3}{a}\right)}{\left(\frac{bx^3}{a} + 1\right)^{5/2}}$$

input `int((a + b*x^3)^(5/2),x)`

output `(x*(a + b*x^3)^(5/2)*hypergeom([-5/2, 1/3], 4/3, -(b*x^3)/a))/((b*x^3)/a + 1)^(5/2)`

Reduce [F]

$$\int (a + bx^3)^{5/2} dx = \frac{106\sqrt{bx^3+a}a^2x}{187} + \frac{74\sqrt{bx^3+a}abx^4}{187} + \frac{2\sqrt{bx^3+a}b^2x^7}{17} + \frac{81\left(\int \frac{\sqrt{bx^3+a}}{bx^3+a} dx\right)a^3}{187}$$

input `int((b*x^3+a)^(5/2),x)`

output `(106*sqrt(a + b*x**3)*a**2*x + 74*sqrt(a + b*x**3)*a*b*x**4 + 22*sqrt(a + b*x**3)*b**2*x**7 + 81*int(sqrt(a + b*x**3)/(a + b*x**3),x)*a**3)/187`

3.14 $\int (a + bx^3)^{3/2} dx$

Optimal result	141
Mathematica [C] (verified)	142
Rubi [A] (verified)	142
Maple [A] (verified)	144
Fricas [A] (verification not implemented)	145
Sympy [A] (verification not implemented)	145
Maxima [F]	146
Giac [F]	146
Mupad [B] (verification not implemented)	146
Reduce [F]	147

Optimal result

Integrand size = 11, antiderivative size = 246

$$\int (a + bx^3)^{3/2} dx = \frac{18}{55}ax\sqrt{a + bx^3} + \frac{2}{11}x(a + bx^3)^{3/2} + \frac{18 \cdot 3^{3/4} \sqrt{2 + \sqrt{3}} a^2 (\sqrt[3]{a} + \sqrt[3]{bx}) \sqrt{\frac{a^{2/3} - \sqrt[3]{a} \sqrt[3]{bx} + b^{2/3} x^2}{((1 + \sqrt{3}) \sqrt[3]{a} + \sqrt[3]{bx})^2}} \operatorname{EllipticF}\left(\arcsin\left(\frac{(1 - \sqrt{3}) \sqrt[3]{a} + \sqrt[3]{bx}}{(1 + \sqrt{3}) \sqrt[3]{a} + \sqrt[3]{bx}}\right), -7 - 4\sqrt{3}\right)}{55 \sqrt[3]{b} \sqrt{\frac{\sqrt[3]{a} (\sqrt[3]{a} + \sqrt[3]{bx})}{((1 + \sqrt{3}) \sqrt[3]{a} + \sqrt[3]{bx})^2}} \sqrt{a + bx^3}}$$

```
output 18/55*a*x*(b*x^3+a)^(1/2)+2/11*x*(b*x^3+a)^(3/2)+18/55*3^(3/4)*(1/2*6^(1/2)
)+1/2*2^(1/2))*a^2*(a^(1/3)+b^(1/3)*x)*((a^(2/3)-a^(1/3)*b^(1/3)*x+b^(2/3)
)*x^2)/((1+3^(1/2))*a^(1/3)+b^(1/3)*x)^2)^(1/2)*EllipticF(((1-3^(1/2))*a^(1
/3)+b^(1/3)*x)/((1+3^(1/2))*a^(1/3)+b^(1/3)*x),I*3^(1/2)+2*I)/b^(1/3)/(a^(
1/3)*(a^(1/3)+b^(1/3)*x)/((1+3^(1/2))*a^(1/3)+b^(1/3)*x)^2)^(1/2)/(b*x^3+a
)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 3.82 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.19

$$\int (a + bx^3)^{3/2} dx = \frac{ax\sqrt{a + bx^3} \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, \frac{1}{3}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{\sqrt{1 + \frac{bx^3}{a}}}$$

input `Integrate[(a + b*x^3)^(3/2),x]`

output `(a*x*Sqrt[a + b*x^3]*Hypergeometric2F1[-3/2, 1/3, 4/3, -(b*x^3)/a])/Sqrt[1 + (b*x^3)/a]`

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.01, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {748, 748, 759}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + bx^3)^{3/2} dx \\ & \quad \downarrow \text{748} \\ & \frac{9}{11}a \int \sqrt{bx^3 + a} dx + \frac{2}{11}x(a + bx^3)^{3/2} \\ & \quad \downarrow \text{748} \\ & \frac{9}{11}a \left(\frac{3}{5}a \int \frac{1}{\sqrt{bx^3 + a}} dx + \frac{2}{5}x\sqrt{a + bx^3} \right) + \frac{2}{11}x(a + bx^3)^{3/2} \\ & \quad \downarrow \text{759} \end{aligned}$$

$$\frac{9}{11} a \left(\frac{2 \cdot 3^{3/4} \sqrt{2 + \sqrt{3}} a (\sqrt[3]{a} + \sqrt[3]{bx}) \sqrt{\frac{a^{2/3} - \sqrt[3]{a} \sqrt[3]{bx} + b^{2/3} x^2}{((1 + \sqrt{3}) \sqrt[3]{a} + \sqrt[3]{bx})^2}} \operatorname{EllipticF} \left(\arcsin \left(\frac{\sqrt[3]{bx} + (1 - \sqrt{3}) \sqrt[3]{a}}{\sqrt[3]{bx} + (1 + \sqrt{3}) \sqrt[3]{a}} \right), -7 - 4\sqrt{3} \right)}{5 \sqrt[3]{b} \sqrt{\frac{\sqrt[3]{a} (\sqrt[3]{a} + \sqrt[3]{bx})}{((1 + \sqrt{3}) \sqrt[3]{a} + \sqrt[3]{bx})^2}} \sqrt{a + bx^3}} \right. \\ \left. \frac{2}{11} x (a + bx^3)^{3/2} \right)$$

input `Int[(a + b*x^3)^(3/2), x]`

output `(2*x*(a + b*x^3)^(3/2))/11 + (9*a*((2*x*Sqrt[a + b*x^3])/5 + (2*3^(3/4)*Sqrt[2 + Sqrt[3]]*a*(a^(1/3) + b^(1/3)*x)*Sqrt[(a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2])/((1 + Sqrt[3])*a^(1/3) + b^(1/3)*x)^2)*EllipticF[ArcSin[((1 - Sqrt[3])*a^(1/3) + b^(1/3)*x)/((1 + Sqrt[3])*a^(1/3) + b^(1/3)*x)], -7 - 4*Sqrt[3]])/(5*b^(1/3)*Sqrt[(a^(1/3)*(a^(1/3) + b^(1/3)*x))/((1 + Sqrt[3])*a^(1/3) + b^(1/3)*x)^2]*Sqrt[a + b*x^3]))/11`

Defintions of rubi rules used

rule 748 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p + 1)), x] + Simp[a*n*(p/(n*p + 1)) Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || LtQ[Denominator[p + 1/n], Denominator[p]])`

rule 759 `Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[2*Sqrt[2 + Sqrt[3]]*(s + r*x)*(Sqrt[(s^2 - r*s*x + r^2*x^2])/((1 + Sqrt[3])*s + r*x)^2]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[s*((s + r*x)/((1 + Sqrt[3])*s + r*x)^2])]*EllipticF[ArcSin[((1 - Sqrt[3])*s + r*x)/((1 + Sqrt[3])*s + r*x)], -7 - 4*Sqrt[3]], x] /; FreeQ[{a, b}, x] && PosQ[a]`

Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 309, normalized size of antiderivative = 1.26

method	result
risch	$\frac{2x(5bx^3+14a)\sqrt{bx^3+a}}{55} - \frac{18ia^2\sqrt{3}(-ab^2)^{\frac{1}{3}}}{(-ab^2)^{\frac{1}{3}}} \sqrt{\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b}-\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}{(-ab^2)^{\frac{1}{3}}}} \sqrt{\frac{x-\frac{(-ab^2)^{\frac{1}{3}}}{b}}{-\frac{3(-ab^2)^{\frac{1}{3}}}{2b}+\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}}} \sqrt{-\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b}-\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}{(-ab^2)^{\frac{1}{3}}}}$
default	$\frac{2bx^4\sqrt{bx^3+a}}{11} + \frac{28ax\sqrt{bx^3+a}}{55} - \frac{18ia^2\sqrt{3}(-ab^2)^{\frac{1}{3}}}{(-ab^2)^{\frac{1}{3}}} \sqrt{\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b}-\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}{(-ab^2)^{\frac{1}{3}}}} \sqrt{\frac{x-\frac{(-ab^2)^{\frac{1}{3}}}{b}}{-\frac{3(-ab^2)^{\frac{1}{3}}}{2b}+\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}}} \sqrt{-\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b}-\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}{(-ab^2)^{\frac{1}{3}}}}$
elliptic	$\frac{2bx^4\sqrt{bx^3+a}}{11} + \frac{28ax\sqrt{bx^3+a}}{55} - \frac{18ia^2\sqrt{3}(-ab^2)^{\frac{1}{3}}}{(-ab^2)^{\frac{1}{3}}} \sqrt{\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b}-\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}{(-ab^2)^{\frac{1}{3}}}} \sqrt{\frac{x-\frac{(-ab^2)^{\frac{1}{3}}}{b}}{-\frac{3(-ab^2)^{\frac{1}{3}}}{2b}+\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}}} \sqrt{-\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b}-\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}{(-ab^2)^{\frac{1}{3}}}}$

input `int((b*x^3+a)^(3/2), x, method=_RETURNVERBOSE)`

output `2/55*x*(5*b*x^3+14*a)*(b*x^3+a)^(1/2)-18/55*I*a^2*3^(1/2)/b*(-a*b^2)^(1/3)*((I*(x+1/2/b*(-a*b^2)^(1/3))-1/2*I*3^(1/2)/b*(-a*b^2)^(1/3))*3^(1/2)*b/(-a*b^2)^(1/3))^(1/2)*((x-1/b*(-a*b^2)^(1/3))/(-3/2/b*(-a*b^2)^(1/3)+1/2*I*3^(1/2)/b*(-a*b^2)^(1/3)))^(1/2)*(-I*(x+1/2/b*(-a*b^2)^(1/3)+1/2*I*3^(1/2)/b*(-a*b^2)^(1/3))*3^(1/2)*b/(-a*b^2)^(1/3))^(1/2)/(b*x^3+a)^(1/2)*EllipticF(1/3*3^(1/2)*(I*(x+1/2/b*(-a*b^2)^(1/3))-1/2*I*3^(1/2)/b*(-a*b^2)^(1/3))*3^(1/2)*b/(-a*b^2)^(1/3))^(1/2), (I*3^(1/2)/b*(-a*b^2)^(1/3))/(-3/2/b*(-a*b^2)^(1/3)+1/2*I*3^(1/2)/b*(-a*b^2)^(1/3))^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.19

$$\int (a + bx^3)^{3/2} dx = \frac{2 \left(27 a^2 \sqrt{b} \operatorname{weierstrassPInverse} \left(0, -\frac{4a}{b}, x \right) + (5 b^2 x^4 + 14 a b x) \sqrt{b x^3 + a} \right)}{55 b}$$

input `integrate((b*x^3+a)^(3/2),x, algorithm="fricas")`

output `2/55*(27*a^2*sqrt(b)*weierstrassPInverse(0, -4*a/b, x) + (5*b^2*x^4 + 14*a*b*x)*sqrt(b*x^3 + a))/b`

Sympy [A] (verification not implemented)

Time = 0.67 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.15

$$\int (a + bx^3)^{3/2} dx = \frac{a^{3/2} x \Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\begin{matrix} -\frac{3}{2}, \frac{1}{3} \\ \frac{4}{3} \end{matrix} \middle| \frac{bx^3 e^{i\pi}}{a} \right)}{3 \Gamma\left(\frac{4}{3}\right)}$$

input `integrate((b*x**3+a)**(3/2),x)`

output `a**(3/2)*x*gamma(1/3)*hyper((-3/2, 1/3), (4/3,), b*x**3*exp_polar(I*pi)/a)/(3*gamma(4/3))`

Maxima [F]

$$\int (a + bx^3)^{3/2} dx = \int (bx^3 + a)^{\frac{3}{2}} dx$$

input `integrate((b*x^3+a)^(3/2),x, algorithm="maxima")`

output `integrate((b*x^3 + a)^(3/2), x)`

Giac [F]

$$\int (a + bx^3)^{3/2} dx = \int (bx^3 + a)^{\frac{3}{2}} dx$$

input `integrate((b*x^3+a)^(3/2),x, algorithm="giac")`

output `integrate((b*x^3 + a)^(3/2), x)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.15

$$\int (a + bx^3)^{3/2} dx = \frac{x (bx^3 + a)^{3/2} {}_2F_1\left(-\frac{3}{2}, \frac{1}{3}; \frac{4}{3}; -\frac{bx^3}{a}\right)}{\left(\frac{bx^3}{a} + 1\right)^{3/2}}$$

input `int((a + b*x^3)^(3/2),x)`

output `(x*(a + b*x^3)^(3/2)*hypergeom([-3/2, 1/3], 4/3, -(b*x^3)/a))/((b*x^3)/a + 1)^(3/2)`

Reduce [F]

$$\int (a + bx^3)^{3/2} dx = \frac{28\sqrt{bx^3 + a}ax}{55} + \frac{2\sqrt{bx^3 + a}bx^4}{11} + \frac{27\left(\int \frac{\sqrt{bx^3 + a}}{bx^3 + a} dx\right) a^2}{55}$$

input `int((b*x^3+a)^(3/2),x)`

output `(28*sqrt(a + b*x**3)*a*x + 10*sqrt(a + b*x**3)*b*x**4 + 27*int(sqrt(a + b*x**3)/(a + b*x**3),x)*a**2)/55`

3.15 $\int \sqrt{a + bx^3} dx$

Optimal result	148
Mathematica [C] (verified)	149
Rubi [A] (verified)	149
Maple [A] (verified)	151
Fricas [A] (verification not implemented)	152
Sympy [A] (verification not implemented)	152
Maxima [F]	152
Giac [F]	153
Mupad [B] (verification not implemented)	153
Reduce [F]	153

Optimal result

Integrand size = 11, antiderivative size = 227

$$\int \sqrt{a + bx^3} dx = \frac{2}{5}x\sqrt{a + bx^3} + \frac{2 \cdot 3^{3/4} \sqrt{2 + \sqrt{3}} a \left(\sqrt[3]{a} + \sqrt[3]{bx} \right) \sqrt{\frac{a^{2/3} - \sqrt[3]{a} \sqrt[3]{bx} + b^{2/3} x^2}{\left((1 + \sqrt{3}) \sqrt[3]{a} + \sqrt[3]{bx} \right)^2}} \operatorname{EllipticF} \left(\arcsin \left(\frac{(1 - \sqrt{3}) \sqrt[3]{a} + \sqrt[3]{bx}}{(1 + \sqrt{3}) \sqrt[3]{a} + \sqrt[3]{bx}} \right), -7 - 4\sqrt{3} \right)}{5 \sqrt[3]{b} \sqrt{\frac{\sqrt[3]{a} \left(\sqrt[3]{a} + \sqrt[3]{bx} \right)}{\left((1 + \sqrt{3}) \sqrt[3]{a} + \sqrt[3]{bx} \right)^2}} \sqrt{a + bx^3}}$$

output

```
2/5*x*(b*x^3+a)^(1/2)+2/5*3^(3/4)*(1/2*6^(1/2)+1/2*2^(1/2))*a*(a^(1/3)+b^(1/3)*x)*((a^(2/3)-a^(1/3)*b^(1/3)*x+b^(2/3)*x^2)/((1+3^(1/2))*a^(1/3)+b^(1/3)*x)^2)^(1/2)*EllipticF(((1-3^(1/2))*a^(1/3)+b^(1/3)*x)/((1+3^(1/2))*a^(1/3)+b^(1/3)*x),I*3^(1/2)+2*I)/b^(1/3)/(a^(1/3)*(a^(1/3)+b^(1/3)*x)/((1+3^(1/2))*a^(1/3)+b^(1/3)*x)^2)^(1/2)/(b*x^3+a)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 3.01 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.20

$$\int \sqrt{a + bx^3} dx = \frac{x\sqrt{a + bx^3} \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, \frac{1}{3}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{\sqrt{1 + \frac{bx^3}{a}}}$$

input `Integrate[Sqrt[a + b*x^3],x]`

output `(x*Sqrt[a + b*x^3]*Hypergeometric2F1[-1/2, 1/3, 4/3, -((b*x^3)/a)]/Sqrt[1 + (b*x^3)/a]`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {748, 759}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \sqrt{a + bx^3} dx \\ \downarrow 748 \\ \frac{3}{5}a \int \frac{1}{\sqrt{bx^3 + a}} dx + \frac{2}{5}x\sqrt{a + bx^3} \\ \downarrow 759 \end{array}$$

$$\frac{2 \cdot 3^{3/4} \sqrt{2 + \sqrt{3}} a \left(\sqrt[3]{a} + \sqrt[3]{bx} \right) \sqrt{\frac{a^{2/3} - \sqrt[3]{a} \sqrt[3]{bx} + b^{2/3} x^2}{\left((1 + \sqrt{3}) \sqrt[3]{a} + \sqrt[3]{bx} \right)^2}} \operatorname{EllipticF} \left(\arcsin \left(\frac{\sqrt[3]{bx} + (1 - \sqrt{3}) \sqrt[3]{a}}{\sqrt[3]{bx} + (1 + \sqrt{3}) \sqrt[3]{a}} \right), -7 - 4\sqrt{3} \right)}{5 \sqrt[3]{b} \sqrt{\frac{\sqrt[3]{a} \left(\sqrt[3]{a} + \sqrt[3]{bx} \right)}{\left((1 + \sqrt{3}) \sqrt[3]{a} + \sqrt[3]{bx} \right)^2}} \sqrt{a + bx^3} + \frac{2}{5} x \sqrt{a + bx^3}}$$

input `Int[Sqrt[a + b*x^3], x]`

output `(2*x*Sqrt[a + b*x^3])/5 + (2*3^(3/4)*Sqrt[2 + Sqrt[3]]*a*(a^(1/3) + b^(1/3)*x)*Sqrt[(a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/((1 + Sqrt[3])*a^(1/3) + b^(1/3)*x)^2]*EllipticF[ArcSin[((1 - Sqrt[3])*a^(1/3) + b^(1/3)*x)/((1 + Sqrt[3])*a^(1/3) + b^(1/3)*x)], -7 - 4*Sqrt[3]]/(5*b^(1/3)*Sqrt[(a^(1/3)*(a^(1/3) + b^(1/3)*x))/((1 + Sqrt[3])*a^(1/3) + b^(1/3)*x)^2]*Sqrt[a + b*x^3])`

Defintions of rubi rules used

rule 748 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p + 1)), x] + Simp[a*n*(p/(n*p + 1)) Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || LtQ[Denominator[p] + 1/n], Denominator[p])`

rule 759 `Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[2*Sqrt[2 + Sqrt[3]]*(s + r*x)*(Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 + Sqrt[3])*s + r*x)^2]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[s*((s + r*x)/((1 + Sqrt[3])*s + r*x)^2]))*EllipticF[ArcSin[((1 - Sqrt[3])*s + r*x)/((1 + Sqrt[3])*s + r*x)], -7 - 4*Sqrt[3]], x] /; FreeQ[{a, b}, x] && PosQ[a]`

Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 297, normalized size of antiderivative = 1.31

method	result
default	$\frac{2x\sqrt{bx^3+a}}{5} - \frac{2ia\sqrt{3}(-ab^2)^{\frac{1}{3}} \sqrt{\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b} - \frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}{(-ab^2)^{\frac{1}{3}}}}{\sqrt{-\frac{x-\frac{(-ab^2)^{\frac{1}{3}}}{b}}{\frac{3(-ab^2)^{\frac{1}{3}}}{2b} + \frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}}}} \sqrt{-\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b} + \frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)}{(-ab^2)^{\frac{1}{3}}}}{5b\sqrt{bx^3+a}}$
risch	$\frac{2x\sqrt{bx^3+a}}{5} - \frac{2ia\sqrt{3}(-ab^2)^{\frac{1}{3}} \sqrt{\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b} - \frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}{(-ab^2)^{\frac{1}{3}}}}{\sqrt{-\frac{x-\frac{(-ab^2)^{\frac{1}{3}}}{b}}{\frac{3(-ab^2)^{\frac{1}{3}}}{2b} + \frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}}}} \sqrt{-\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b} + \frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)}{(-ab^2)^{\frac{1}{3}}}}{5b\sqrt{bx^3+a}}$
elliptic	$\frac{2x\sqrt{bx^3+a}}{5} - \frac{2ia\sqrt{3}(-ab^2)^{\frac{1}{3}} \sqrt{\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b} - \frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}{(-ab^2)^{\frac{1}{3}}}}{\sqrt{-\frac{x-\frac{(-ab^2)^{\frac{1}{3}}}{b}}{\frac{3(-ab^2)^{\frac{1}{3}}}{2b} + \frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}}}} \sqrt{-\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b} + \frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)}{(-ab^2)^{\frac{1}{3}}}}{5b\sqrt{bx^3+a}}$

input `int((b*x^3+a)^(1/2), x, method=_RETURNVERBOSE)`

output `2/5*x*(b*x^3+a)^(1/2)-2/5*I*a*3^(1/2)/b*(-a*b^2)^(1/3)*(I*(x+1/2/b*(-a*b^2)^(1/3))-1/2*I*3^(1/2)/b*(-a*b^2)^(1/3))*3^(1/2)*b/(-a*b^2)^(1/3))^((x-1/b*(-a*b^2)^(1/3))/(-3/2/b*(-a*b^2)^(1/3)+1/2*I*3^(1/2)/b*(-a*b^2)^(1/3)))^(1/2)*(-I*(x+1/2/b*(-a*b^2)^(1/3)+1/2*I*3^(1/2)/b*(-a*b^2)^(1/3))*3^(1/2)*b/(-a*b^2)^(1/3))^((x+1/2/b*(-a*b^2)^(1/3)-1/2*I*3^(1/2)/b*(-a*b^2)^(1/3))*3^(1/2)*b/(-a*b^2)^(1/3))^((1/2), (I*3^(1/2)/b*(-a*b^2)^(1/3))/(-3/2/b*(-a*b^2)^(1/3)+1/2*I*3^(1/2)/b*(-a*b^2)^(1/3)))^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.15

$$\int \sqrt{a + bx^3} dx = \frac{2 \left(\sqrt{bx^3 + abx} + 3a\sqrt{b} \operatorname{weierstrassPInverse}(0, -\frac{4a}{b}, x) \right)}{5b}$$

input `integrate((b*x^3+a)^(1/2),x, algorithm="fricas")`output `2/5*(sqrt(b*x^3 + a)*b*x + 3*a*sqrt(b)*weierstrassPInverse(0, -4*a/b, x))/b`**Sympy [A] (verification not implemented)**

Time = 0.46 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.16

$$\int \sqrt{a + bx^3} dx = \frac{\sqrt{ax}\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\begin{matrix} -\frac{1}{2}, \frac{1}{3} \\ \frac{4}{3} \end{matrix} \middle| \frac{bx^3 e^{i\pi}}{a} \right)}{3\Gamma\left(\frac{4}{3}\right)}$$

input `integrate((b*x**3+a)**(1/2),x)`output `sqrt(a)*x*gamma(1/3)*hyper((-1/2, 1/3), (4/3,), b*x**3*exp_polar(I*pi)/a)/(3*gamma(4/3))`**Maxima [F]**

$$\int \sqrt{a + bx^3} dx = \int \sqrt{bx^3 + a} dx$$

input `integrate((b*x^3+a)^(1/2),x, algorithm="maxima")`output `integrate(sqrt(b*x^3 + a), x)`

Giac [F]

$$\int \sqrt{a + bx^3} dx = \int \sqrt{bx^3 + a} dx$$

input `integrate((b*x^3+a)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*x^3 + a), x)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.16

$$\int \sqrt{a + bx^3} dx = \frac{x \sqrt{bx^3 + a} {}_2F_1\left(-\frac{1}{2}, \frac{1}{3}; \frac{4}{3}; -\frac{bx^3}{a}\right)}{\sqrt{\frac{bx^3}{a} + 1}}$$

input `int((a + b*x^3)^(1/2),x)`

output `(x*(a + b*x^3)^(1/2)*hypergeom([-1/2, 1/3], 4/3, -(b*x^3)/a))/((b*x^3)/a + 1)^(1/2)`

Reduce [F]

$$\int \sqrt{a + bx^3} dx = \frac{2\sqrt{bx^3 + a}x}{5} + \frac{3\left(\int \frac{\sqrt{bx^3+a}}{bx^3+a} dx\right)a}{5}$$

input `int((b*x^3+a)^(1/2),x)`

output `(2*sqrt(a + b*x**3)*x + 3*int(sqrt(a + b*x**3)/(a + b*x**3),x)*a)/5`

3.16 $\int \frac{1}{\sqrt{a+bx^3}} dx$

Optimal result	154
Mathematica [C] (verified)	155
Rubi [A] (verified)	155
Maple [A] (verified)	156
Fricas [A] (verification not implemented)	157
Sympy [A] (verification not implemented)	157
Maxima [F]	158
Giac [F]	158
Mupad [B] (verification not implemented)	158
Reduce [F]	159

Optimal result

Integrand size = 11, antiderivative size = 207

$$\int \frac{1}{\sqrt{a+bx^3}} dx$$

$$= \frac{2\sqrt{2+\sqrt{3}}(\sqrt[3]{a}+\sqrt[3]{bx})\sqrt{\frac{a^{2/3}-\sqrt[3]{a}\sqrt[3]{bx}+b^{2/3}x^2}{((1+\sqrt{3})\sqrt[3]{a}+\sqrt[3]{bx})^2}}\text{EllipticF}\left(\arcsin\left(\frac{(1-\sqrt{3})\sqrt[3]{a}+\sqrt[3]{bx}}{(1+\sqrt{3})\sqrt[3]{a}+\sqrt[3]{bx}}\right),-7-4\sqrt{3}\right)}{\sqrt[4]{3}\sqrt[3]{b}\sqrt{\frac{\sqrt[3]{a}(\sqrt[3]{a}+\sqrt[3]{bx})}{((1+\sqrt{3})\sqrt[3]{a}+\sqrt[3]{bx})^2}}\sqrt{a+bx^3}}$$

output

```
2/3*(1/2*6^(1/2)+1/2*2^(1/2))*(a^(1/3)+b^(1/3)*x)*((a^(2/3)-a^(1/3)*b^(1/3)
)*x+b^(2/3)*x^2)/((1+3^(1/2))*a^(1/3)+b^(1/3)*x)^2)^(1/2)*EllipticF(((1-3^(
1/2))*a^(1/3)+b^(1/3)*x)/((1+3^(1/2))*a^(1/3)+b^(1/3)*x),I*3^(1/2)+2*I)*3
^(3/4)/b^(1/3)/(a^(1/3)*(a^(1/3)+b^(1/3)*x)/((1+3^(1/2))*a^(1/3)+b^(1/3)*x
)^2)^(1/2)/(b*x^3+a)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.01 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.22

$$\int \frac{1}{\sqrt{a + bx^3}} dx = \frac{x\sqrt{1 + \frac{bx^3}{a}} \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{\sqrt{a + bx^3}}$$

input `Integrate[1/Sqrt[a + b*x^3],x]`

output `(x*Sqrt[1 + (b*x^3)/a]*Hypergeometric2F1[1/3, 1/2, 4/3, -((b*x^3)/a)]/Sqrt[a + b*x^3]`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {759}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{a + bx^3}} dx$$

↓ 759

$$\frac{2\sqrt{2 + \sqrt{3}}(\sqrt[3]{a} + \sqrt[3]{bx}) \sqrt{\frac{a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx} + b^{2/3}x^2}{((1 + \sqrt{3})\sqrt[3]{a} + \sqrt[3]{bx})^2}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt[3]{bx} + (1 - \sqrt{3})\sqrt[3]{a}}{\sqrt[3]{bx} + (1 + \sqrt{3})\sqrt[3]{a}}\right), -7 - 4\sqrt{3}\right)}{\sqrt[4]{3}\sqrt[3]{b} \sqrt{\frac{\sqrt[3]{a}(\sqrt[3]{a} + \sqrt[3]{bx})}{((1 + \sqrt{3})\sqrt[3]{a} + \sqrt[3]{bx})^2}} \sqrt{a + bx^3}}$$

input `Int[1/Sqrt[a + b*x^3],x]`

output

```
(2*Sqrt[2 + Sqrt[3]]*(a^(1/3) + b^(1/3)*x)*Sqrt[(a^(2/3) - a^(1/3)*b^(1/3)
*x + b^(2/3)*x^2)/((1 + Sqrt[3])*a^(1/3) + b^(1/3)*x)^2]*EllipticF[ArcSin[
((1 - Sqrt[3])*a^(1/3) + b^(1/3)*x)/((1 + Sqrt[3])*a^(1/3) + b^(1/3)*x)],
-7 - 4*Sqrt[3]]/(3^(1/4)*b^(1/3)*Sqrt[(a^(1/3)*(a^(1/3) + b^(1/3)*x))/((1
+ Sqrt[3])*a^(1/3) + b^(1/3)*x)^2]*Sqrt[a + b*x^3])
```

Defintions of rubi rules used

rule 759

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] :> With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[2*Sqrt[2 + Sqrt[3]]*(s + r*x)*(Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 + Sqrt[3])*s + r*x)^2]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[s*
((s + r*x)/((1 + Sqrt[3])*s + r*x)^2)))*EllipticF[ArcSin[((1 - Sqrt[3])*s
+ r*x)/((1 + Sqrt[3])*s + r*x)], -7 - 4*Sqrt[3]], x] /; FreeQ[{a, b}, x] &
& PosQ[a]
```

Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.37

method	result
default	$2i\sqrt{3}(-ab^2)^{\frac{1}{3}} \sqrt{\frac{i\left(x + \frac{(-ab^2)^{\frac{1}{3}}}{2b} - \frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}{(-ab^2)^{\frac{1}{3}}}} \sqrt{\frac{x - \frac{(-ab^2)^{\frac{1}{3}}}{b}}{-\frac{3(-ab^2)^{\frac{1}{3}}}{2b} + \frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}}} \sqrt{\frac{i\left(x + \frac{(-ab^2)^{\frac{1}{3}}}{2b} + \frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}{(-ab^2)^{\frac{1}{3}}}}$ <hr/> $3b\sqrt{bx^3+a}$
elliptic	$2i\sqrt{3}(-ab^2)^{\frac{1}{3}} \sqrt{\frac{i\left(x + \frac{(-ab^2)^{\frac{1}{3}}}{2b} - \frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}{(-ab^2)^{\frac{1}{3}}}} \sqrt{\frac{x - \frac{(-ab^2)^{\frac{1}{3}}}{b}}{-\frac{3(-ab^2)^{\frac{1}{3}}}{2b} + \frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}}} \sqrt{\frac{i\left(x + \frac{(-ab^2)^{\frac{1}{3}}}{2b} + \frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}{(-ab^2)^{\frac{1}{3}}}}$ <hr/> $3b\sqrt{bx^3+a}$

input

```
int(1/(b*x^3+a)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
-2/3*I*3^(1/2)/b*(-a*b^2)^(1/3)*(I*(x+1/2/b*(-a*b^2)^(1/3)-1/2*I*3^(1/2)/b
*(-a*b^2)^(1/3))*3^(1/2)*b/(-a*b^2)^(1/3))^(1/2)*((x-1/b*(-a*b^2)^(1/3))/(
-3/2/b*(-a*b^2)^(1/3)+1/2*I*3^(1/2)/b*(-a*b^2)^(1/3)))^(1/2)*(-I*(x+1/2/b*
(-a*b^2)^(1/3)+1/2*I*3^(1/2)/b*(-a*b^2)^(1/3))*3^(1/2)*b/(-a*b^2)^(1/3))^(
1/2)/(b*x^3+a)^(1/2)*EllipticF(1/3*3^(1/2)*(I*(x+1/2/b*(-a*b^2)^(1/3)-1/2*
I*3^(1/2)/b*(-a*b^2)^(1/3))*3^(1/2)*b/(-a*b^2)^(1/3))^(1/2),(I*3^(1/2)/b*(
-a*b^2)^(1/3)/(-3/2/b*(-a*b^2)^(1/3)+1/2*I*3^(1/2)/b*(-a*b^2)^(1/3)))^(1/2
))
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.07

$$\int \frac{1}{\sqrt{a+bx^3}} dx = \frac{2 \operatorname{weierstrassPInverse}\left(0, -\frac{4a}{b}, x\right)}{\sqrt{b}}$$

input

```
integrate(1/(b*x^3+a)^(1/2),x, algorithm="fricas")
```

output

```
2*weierstrassPInverse(0, -4*a/b, x)/sqrt(b)
```

Sympy [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.17

$$\int \frac{1}{\sqrt{a+bx^3}} dx = \frac{x\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{1}{2} \middle| \frac{bx^3 e^{i\pi}}{a}\right)}{3\sqrt{a}\Gamma\left(\frac{4}{3}\right)}$$

input

```
integrate(1/(b*x**3+a)**(1/2),x)
```

output

```
x*gamma(1/3)*hyper((1/3, 1/2), (4/3,), b*x**3*exp_polar(I*pi)/a)/(3*sqrt(a)
)*gamma(4/3))
```

Maxima [F]

$$\int \frac{1}{\sqrt{a + bx^3}} dx = \int \frac{1}{\sqrt{bx^3 + a}} dx$$

input `integrate(1/(b*x^3+a)^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(b*x^3 + a), x)`

Giac [F]

$$\int \frac{1}{\sqrt{a + bx^3}} dx = \int \frac{1}{\sqrt{bx^3 + a}} dx$$

input `integrate(1/(b*x^3+a)^(1/2),x, algorithm="giac")`

output `integrate(1/sqrt(b*x^3 + a), x)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.18

$$\int \frac{1}{\sqrt{a + bx^3}} dx = \frac{x \sqrt{\frac{bx^3}{a} + 1} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; -\frac{bx^3}{a}\right)}{\sqrt{bx^3 + a}}$$

input `int(1/(a + b*x^3)^(1/2),x)`

output `(x*((b*x^3)/a + 1)^(1/2)*hypergeom([1/3, 1/2], 4/3, -(b*x^3)/a))/(a + b*x^3)^(1/2)`

Reduce [F]

$$\int \frac{1}{\sqrt{a+bx^3}} dx = \int \frac{\sqrt{bx^3+a}}{bx^3+a} dx$$

input `int(1/(b*x^3+a)^(1/2),x)`

output `int(sqrt(a + b*x**3)/(a + b*x**3),x)`

3.17 $\int \frac{1}{(a+bx^3)^{3/2}} dx$

Optimal result	160
Mathematica [C] (verified)	161
Rubi [A] (verified)	161
Maple [A] (verified)	163
Fricas [A] (verification not implemented)	164
Sympy [A] (verification not implemented)	164
Maxima [F]	164
Giac [F]	165
Mupad [B] (verification not implemented)	165
Reduce [F]	165

Optimal result

Integrand size = 11, antiderivative size = 232

$$\int \frac{1}{(a+bx^3)^{3/2}} dx = \frac{2x}{3a\sqrt{a+bx^3}} + \frac{2\sqrt{2+\sqrt{3}}(\sqrt[3]{a} + \sqrt[3]{bx}) \sqrt{\frac{a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx} + b^{2/3}x^2}{((1+\sqrt{3})\sqrt[3]{a} + \sqrt[3]{bx})^2}} \operatorname{EllipticF}\left(\arcsin\left(\frac{(1-\sqrt{3})\sqrt[3]{a} + \sqrt[3]{bx}}{(1+\sqrt{3})\sqrt[3]{a} + \sqrt[3]{bx}}\right), -7 - 4\sqrt{3}\right)}{3\sqrt[4]{3}a\sqrt[3]{b} \sqrt{\frac{\sqrt[3]{a}(\sqrt[3]{a} + \sqrt[3]{bx})}{((1+\sqrt{3})\sqrt[3]{a} + \sqrt[3]{bx})^2}} \sqrt{a+bx^3}}$$

output

```
2/3*x/a/(b*x^3+a)^(1/2)+2/9*(1/2*6^(1/2)+1/2*2^(1/2))*(a^(1/3)+b^(1/3)*x)*
((a^(2/3)-a^(1/3)*b^(1/3)*x+b^(2/3)*x^2)/((1+3^(1/2))*a^(1/3)+b^(1/3)*x)^(2
)^(1/2)*EllipticF(((1-3^(1/2))*a^(1/3)+b^(1/3)*x)/((1+3^(1/2))*a^(1/3)+b^(
1/3)*x),I*3^(1/2)+2*I)*3^(3/4)/a/b^(1/3)/(a^(1/3)*(a^(1/3)+b^(1/3)*x)/((1+
3^(1/2))*a^(1/3)+b^(1/3)*x)^(1/2)/(b*x^3+a)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 3.33 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.24

$$\int \frac{1}{(a + bx^3)^{3/2}} dx = \frac{x \left(2 + \sqrt{1 + \frac{bx^3}{a}} \operatorname{Hypergeometric2F1} \left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, -\frac{bx^3}{a} \right) \right)}{3a\sqrt{a + bx^3}}$$

input `Integrate[(a + b*x^3)^(-3/2),x]`

output `(x*(2 + Sqrt[1 + (b*x^3)/a]*Hypergeometric2F1[1/3, 1/2, 4/3, -((b*x^3)/a)])/(3*a*Sqrt[a + b*x^3])`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 232, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {749, 759}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{1}{(a + bx^3)^{3/2}} dx \\ \downarrow 749 \\ \frac{\int \frac{1}{\sqrt{bx^3+a}} dx}{3a} + \frac{2x}{3a\sqrt{a + bx^3}} \\ \downarrow 759 \end{array}$$

$$\frac{2\sqrt{2+\sqrt{3}}\left(\sqrt[3]{a}+\sqrt[3]{bx}\right)\sqrt{\frac{a^{2/3}-\sqrt[3]{a}\sqrt[3]{bx}+b^{2/3}x^2}{\left((1+\sqrt{3})\sqrt[3]{a}+\sqrt[3]{bx}\right)^2}}\operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt[3]{bx}+(1-\sqrt{3})\sqrt[3]{a}}{\sqrt[3]{bx}+(1+\sqrt{3})\sqrt[3]{a}}\right),-7-4\sqrt{3}\right)}{3^4\sqrt{3}a\sqrt[3]{b}\sqrt{\frac{\sqrt[3]{a}\left(\sqrt[3]{a}+\sqrt[3]{bx}\right)}{\left((1+\sqrt{3})\sqrt[3]{a}+\sqrt[3]{bx}\right)^2}}\sqrt{a+bx^3}}+\frac{2x}{3a\sqrt{a+bx^3}}$$

input `Int[(a + b*x^3)^(-3/2),x]`

output `(2*x)/(3*a*Sqrt[a + b*x^3]) + (2*Sqrt[2 + Sqrt[3]]*(a^(1/3) + b^(1/3)*x)*Sqrt[(a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/((1 + Sqrt[3])*a^(1/3) + b^(1/3)*x)^2)*EllipticF[ArcSin[((1 - Sqrt[3])*a^(1/3) + b^(1/3)*x)/((1 + Sqrt[3])*a^(1/3) + b^(1/3)*x)], -7 - 4*Sqrt[3]]/(3*3^(1/4)*a*b^(1/3)*Sqrt[(a^(1/3)*(a^(1/3) + b^(1/3)*x))/((1 + Sqrt[3])*a^(1/3) + b^(1/3)*x)^2])*Sqrt[a + b*x^3])`

Defintions of rubi rules used

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

rule 759 `Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[2*Sqrt[2 + Sqrt[3]]*(s + r*x)*(Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 + Sqrt[3])*s + r*x)^2]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[s*((s + r*x)/((1 + Sqrt[3])*s + r*x)^2])))*EllipticF[ArcSin[((1 - Sqrt[3])*s + r*x)/((1 + Sqrt[3])*s + r*x)], -7 - 4*Sqrt[3]], x] /; FreeQ[{a, b}, x] && PosQ[a]`

Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 306, normalized size of antiderivative = 1.32

method	result
default	$\frac{2i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{3a\sqrt{(x^3+\frac{a}{b})b}} - \frac{\sqrt{\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b}-\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}{(-ab^2)^{\frac{1}{3}}}}}{\sqrt{-\frac{x-\frac{(-ab^2)^{\frac{1}{3}}}{b}}{3\frac{(-ab^2)^{\frac{1}{3}}}{2b}+\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}}}} \sqrt{\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b}+\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)}{(-ab^2)^{\frac{1}{3}}}}$ $\frac{2x}{3a\sqrt{(x^3+\frac{a}{b})b}} - \frac{9ab\sqrt{bx}}{9ab\sqrt{bx}}$
elliptic	$\frac{2i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{3a\sqrt{(x^3+\frac{a}{b})b}} - \frac{\sqrt{\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b}-\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}{(-ab^2)^{\frac{1}{3}}}}}{\sqrt{-\frac{x-\frac{(-ab^2)^{\frac{1}{3}}}{b}}{3\frac{(-ab^2)^{\frac{1}{3}}}{2b}+\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}}}} \sqrt{\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b}+\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)}{(-ab^2)^{\frac{1}{3}}}}$ $\frac{2x}{3a\sqrt{(x^3+\frac{a}{b})b}} - \frac{9ab\sqrt{bx}}{9ab\sqrt{bx}}$

```
input int(1/(b*x^3+a)^(3/2), x, method=_RETURNVERBOSE)
```

```
output 2/3*x/a/((x^3+a/b)*b)^(1/2)-2/9*I/a*3^(1/2)/b*(-a*b^2)^(1/3)*(I*(x+1/2/b*(-a*b^2)^(1/3)-1/2*I*3^(1/2)/b*(-a*b^2)^(1/3))*3^(1/2)*b/(-a*b^2)^(1/3))^(1/2)*((x-1/b*(-a*b^2)^(1/3))/(-3/2/b*(-a*b^2)^(1/3)+1/2*I*3^(1/2)/b*(-a*b^2)^(1/3)))^(1/2)*(-I*(x+1/2/b*(-a*b^2)^(1/3)+1/2*I*3^(1/2)/b*(-a*b^2)^(1/3))*3^(1/2)*b/(-a*b^2)^(1/3))^(1/2)/(b*x^3+a)^(1/2)*EllipticF(1/3*3^(1/2)*(I*(x+1/2/b*(-a*b^2)^(1/3)-1/2*I*3^(1/2)/b*(-a*b^2)^(1/3))*3^(1/2)*b/(-a*b^2)^(1/3))^(1/2), (I*3^(1/2)/b*(-a*b^2)^(1/3))/(-3/2/b*(-a*b^2)^(1/3)+1/2*I*3^(1/2)/b*(-a*b^2)^(1/3)))^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.22

$$\int \frac{1}{(a + bx^3)^{3/2}} dx = \frac{2 \left(\sqrt{bx^3 + abx} + (bx^3 + a)\sqrt{b} \operatorname{weierstrassPInverse}(0, -\frac{4a}{b}, x) \right)}{3(ab^2x^3 + a^2b)}$$

input `integrate(1/(b*x^3+a)^(3/2),x, algorithm="fricas")`

output `2/3*(sqrt(b*x^3 + a)*b*x + (b*x^3 + a)*sqrt(b)*weierstrassPInverse(0, -4*a/b, x))/(a*b^2*x^3 + a^2*b)`

Sympy [A] (verification not implemented)

Time = 0.49 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.16

$$\int \frac{1}{(a + bx^3)^{3/2}} dx = \frac{x\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{3}{2} \middle| \frac{4}{3}, \frac{bx^3 e^{i\pi}}{a}\right)}{3a^{\frac{3}{2}}\Gamma\left(\frac{4}{3}\right)}$$

input `integrate(1/(b*x**3+a)**(3/2),x)`

output `x*gamma(1/3)*hyper((1/3, 3/2), (4/3,), b*x**3*exp_polar(I*pi)/a)/(3*a**(3/2)*gamma(4/3))`

Maxima [F]

$$\int \frac{1}{(a + bx^3)^{3/2}} dx = \int \frac{1}{(bx^3 + a)^{\frac{3}{2}}} dx$$

input `integrate(1/(b*x^3+a)^(3/2),x, algorithm="maxima")`

output `integrate((b*x^3 + a)^(-3/2), x)`

Giac [F]

$$\int \frac{1}{(a + bx^3)^{3/2}} dx = \int \frac{1}{(bx^3 + a)^{\frac{3}{2}}} dx$$

input `integrate(1/(b*x^3+a)^(3/2),x, algorithm="giac")`

output `integrate((b*x^3 + a)^(-3/2), x)`

Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.16

$$\int \frac{1}{(a + bx^3)^{3/2}} dx = \frac{x \left(\frac{bx^3}{a} + 1 \right)^{3/2} {}_2F_1 \left(\frac{1}{3}, \frac{3}{2}; \frac{4}{3}; -\frac{bx^3}{a} \right)}{(bx^3 + a)^{3/2}}$$

input `int(1/(a + b*x^3)^(3/2),x)`

output `(x*((b*x^3)/a + 1)^(3/2)*hypergeom([1/3, 3/2], 4/3, -(b*x^3)/a))/(a + b*x^3)^(3/2)`

Reduce [F]

$$\int \frac{1}{(a + bx^3)^{3/2}} dx = \int \frac{\sqrt{bx^3 + a}}{b^2x^6 + 2abx^3 + a^2} dx$$

input `int(1/(b*x^3+a)^(3/2),x)`

output `int(sqrt(a + b*x**3)/(a**2 + 2*a*b*x**3 + b**2*x**6),x)`

3.18 $\int \frac{1}{(a+bx^3)^{5/2}} dx$

Optimal result	167
Mathematica [C] (verified)	168
Rubi [A] (verified)	168
Maple [A] (verified)	170
Fricas [A] (verification not implemented)	171
Sympy [A] (verification not implemented)	171
Maxima [F]	172
Giac [F]	172
Mupad [B] (verification not implemented)	172
Reduce [F]	173

Optimal result

Integrand size = 11, antiderivative size = 251

$$\int \frac{1}{(a+bx^3)^{5/2}} dx = \frac{2x}{9a(a+bx^3)^{3/2}} + \frac{14x}{27a^2\sqrt{a+bx^3}}$$

$$+ \frac{14\sqrt{2+\sqrt{3}}(\sqrt[3]{a} + \sqrt[3]{bx}) \sqrt{\frac{a^{2/3}-\sqrt[3]{a}\sqrt[3]{bx}+b^{2/3}x^2}{((1+\sqrt{3})\sqrt[3]{a}+\sqrt[3]{bx})^2}} \operatorname{EllipticF}\left(\arcsin\left(\frac{(1-\sqrt{3})\sqrt[3]{a}+\sqrt[3]{bx}}{(1+\sqrt{3})\sqrt[3]{a}+\sqrt[3]{bx}}\right), -7-4\sqrt{3}\right)}{27\sqrt[4]{3}a^2\sqrt[3]{b} \sqrt{\frac{\sqrt[3]{a}(\sqrt[3]{a}+\sqrt[3]{bx})}{((1+\sqrt{3})\sqrt[3]{a}+\sqrt[3]{bx})^2}} \sqrt{a+bx^3}}$$

output

```
2/9*x/a/(b*x^3+a)^(3/2)+14/27*x/a^2/(b*x^3+a)^(1/2)+14/81*(1/2*6^(1/2)+1/2
*2^(1/2))*(a^(1/3)+b^(1/3)*x)*((a^(2/3)-a^(1/3)*b^(1/3)*x+b^(2/3)*x^2)/((1
+3^(1/2))*a^(1/3)+b^(1/3)*x)^2)^(1/2)*EllipticF(((1-3^(1/2))*a^(1/3)+b^(1/
3)*x)/((1+3^(1/2))*a^(1/3)+b^(1/3)*x),I*3^(1/2)+2*I)*3^(3/4)/a^2/b^(1/3)/(
a^(1/3)*(a^(1/3)+b^(1/3)*x)/((1+3^(1/2))*a^(1/3)+b^(1/3)*x)^2)^(1/2)/(b*x^
3+a)^(1/2)
```


Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 4.16 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.29

$$\int \frac{1}{(a + bx^3)^{5/2}} dx = \frac{20ax + 14bx^4 + 7x(a + bx^3) \sqrt{1 + \frac{bx^3}{a}} \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{27a^2 (a + bx^3)^{3/2}}$$

input

```
Integrate[(a + b*x^3)^(-5/2), x]
```

output

```
(20*a*x + 14*b*x^4 + 7*x*(a + b*x^3)*Sqrt[1 + (b*x^3)/a]*Hypergeometric2F1
[1/3, 1/2, 4/3, -((b*x^3)/a)]/(27*a^2*(a + b*x^3)^(3/2))
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 259, normalized size of antiderivative = 1.03, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {749, 749, 759}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(a + bx^3)^{5/2}} dx \\ & \quad \downarrow 749 \\ & \frac{7 \int \frac{1}{(bx^3+a)^{3/2}} dx}{9a} + \frac{2x}{9a (a + bx^3)^{3/2}} \\ & \quad \downarrow 749 \\ & \frac{7 \left(\frac{\int \frac{1}{\sqrt{bx^3+a}} dx}{3a} + \frac{2x}{3a\sqrt{a+bx^3}} \right)}{9a} + \frac{2x}{9a (a + bx^3)^{3/2}} \\ & \quad \downarrow 759 \end{aligned}$$

$$7 \left(\frac{2\sqrt{2+\sqrt{3}} \left(\sqrt[3]{a} + \sqrt[3]{bx} \right) \sqrt{\frac{a^{2/3} - \sqrt[3]{a} \sqrt[3]{bx} + b^{2/3} x^2}{\left((1+\sqrt{3}) \sqrt[3]{a} + \sqrt[3]{bx} \right)^2}} \operatorname{EllipticF} \left(\arcsin \left(\frac{\sqrt[3]{bx} + (1-\sqrt{3}) \sqrt[3]{a}}{\sqrt[3]{bx} + (1+\sqrt{3}) \sqrt[3]{a}} \right), -7-4\sqrt{3} \right)}{3^4 \sqrt[3]{3a} \sqrt[3]{b} \sqrt{\frac{\sqrt[3]{a} \left(\sqrt[3]{a} + \sqrt[3]{bx} \right)}{\left((1+\sqrt{3}) \sqrt[3]{a} + \sqrt[3]{bx} \right)^2}} \sqrt{a+bx^3}} + \frac{2x}{3a\sqrt{a+bx^3}} \right) + \frac{9a}{2x} \frac{1}{9a(a+bx^3)^{3/2}}$$

input `Int[(a + b*x^3)^(-5/2), x]`

output `(2*x)/(9*a*(a + b*x^3)^(3/2)) + (7*((2*x)/(3*a*Sqrt[a + b*x^3]) + (2*Sqrt[2 + Sqrt[3]]*(a^(1/3) + b^(1/3)*x)*Sqrt[(a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/((1 + Sqrt[3])*a^(1/3) + b^(1/3)*x)^2)*EllipticF[ArcSin[((1 - Sqrt[3])*a^(1/3) + b^(1/3)*x)/((1 + Sqrt[3])*a^(1/3) + b^(1/3)*x)], -7 - 4*Sqrt[3]])/(3*3^(1/4)*a*b^(1/3)*Sqrt[(a^(1/3)*(a^(1/3) + b^(1/3)*x)]/((1 + Sqrt[3])*a^(1/3) + b^(1/3)*x)^2)*Sqrt[a + b*x^3]))/(9*a)`

Defintions of rubi rules used

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

rule 759 `Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[2*Sqrt[2 + Sqrt[3]]*(s + r*x)*(Sqrt[(s^2 - r*s*x + r^2*x^2)]/((1 + Sqrt[3])*s + r*x)^2)/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[s*((s + r*x)/((1 + Sqrt[3])*s + r*x)^2])]*EllipticF[ArcSin[((1 - Sqrt[3])*s + r*x)/((1 + Sqrt[3])*s + r*x)], -7 - 4*Sqrt[3]], x] /; FreeQ[{a, b}, x] && PosQ[a]`

Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 335, normalized size of antiderivative = 1.33

method	result
default	$\frac{2x\sqrt{bx^3+a}}{9ab^2(x^3+\frac{a}{b})^2} + \frac{14x}{27a^2\sqrt{(x^3+\frac{a}{b})b}} - \frac{14i\sqrt{3}(-ab^2)^{\frac{1}{3}}\sqrt{\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b}-\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}{(-ab^2)^{\frac{1}{3}}}}}{\sqrt{\frac{x-\frac{(-ab^2)^{\frac{1}{3}}}{b}}{3\frac{(-ab^2)^{\frac{1}{3}}}{2b}+\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}}}}$
elliptic	$\frac{2x\sqrt{bx^3+a}}{9ab^2(x^3+\frac{a}{b})^2} + \frac{14x}{27a^2\sqrt{(x^3+\frac{a}{b})b}} - \frac{14i\sqrt{3}(-ab^2)^{\frac{1}{3}}\sqrt{\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b}-\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}{(-ab^2)^{\frac{1}{3}}}}}{\sqrt{\frac{x-\frac{(-ab^2)^{\frac{1}{3}}}{b}}{3\frac{(-ab^2)^{\frac{1}{3}}}{2b}+\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}}}}$

```
input int(1/(b*x^3+a)^(5/2), x, method=_RETURNVERBOSE)
```

```
output 2/9*x/a/b^2*(b*x^3+a)^(1/2)/(x^3+a/b)^2+14/27*x/a^2/((x^3+a/b)*b)^(1/2)-14/81*I/a^2*3^(1/2)/b*(-a*b^2)^(1/3)*(I*(x+1/2/b*(-a*b^2)^(1/3)-1/2*I*3^(1/2)/b*(-a*b^2)^(1/3))*3^(1/2)*b/(-a*b^2)^(1/3))^(1/2)*((x-1/b*(-a*b^2)^(1/3))/(-3/2/b*(-a*b^2)^(1/3)+1/2*I*3^(1/2)/b*(-a*b^2)^(1/3)))^(1/2)*(-I*(x+1/2/b*(-a*b^2)^(1/3)+1/2*I*3^(1/2)/b*(-a*b^2)^(1/3))*3^(1/2)*b/(-a*b^2)^(1/3))^(1/2)/(b*x^3+a)^(1/2)*EllipticF(1/3*3^(1/2)*(I*(x+1/2/b*(-a*b^2)^(1/3)-1/2*I*3^(1/2)/b*(-a*b^2)^(1/3))*3^(1/2)*b/(-a*b^2)^(1/3))^(1/2), (I*3^(1/2)/b*(-a*b^2)^(1/3)/(-3/2/b*(-a*b^2)^(1/3)+1/2*I*3^(1/2)/b*(-a*b^2)^(1/3)))^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.35

$$\int \frac{1}{(a + bx^3)^{5/2}} dx = \frac{2 \left(7(b^2x^6 + 2abx^3 + a^2)\sqrt{b}\text{weierstrassPInverse}(0, -\frac{4a}{b}, x) + (7b^2x^4 + 10abx)\sqrt{bx^3} \right)}{27(a^2b^3x^6 + 2a^3b^2x^3 + a^4b)}$$

input `integrate(1/(b*x^3+a)^(5/2),x, algorithm="fricas")`

output `2/27*(7*(b^2*x^6 + 2*a*b*x^3 + a^2)*sqrt(b)*weierstrassPInverse(0, -4*a/b, x) + (7*b^2*x^4 + 10*a*b*x)*sqrt(b*x^3 + a))/(a^2*b^3*x^6 + 2*a^3*b^2*x^3 + a^4*b)`

Sympy [A] (verification not implemented)

Time = 0.52 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.14

$$\int \frac{1}{(a + bx^3)^{5/2}} dx = \frac{x\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{5}{2} \middle| \frac{4}{3}, \frac{bx^3 e^{i\pi}}{a}\right)}{3a^{5/2}\Gamma\left(\frac{4}{3}\right)}$$

input `integrate(1/(b*x**3+a)**(5/2),x)`

output `x*gamma(1/3)*hyper((1/3, 5/2), (4/3,), b*x**3*exp_polar(I*pi)/a)/(3*a**(5/2)*gamma(4/3))`

Maxima [F]

$$\int \frac{1}{(a + bx^3)^{5/2}} dx = \int \frac{1}{(bx^3 + a)^{5/2}} dx$$

input `integrate(1/(b*x^3+a)^(5/2),x, algorithm="maxima")`

output `integrate((b*x^3 + a)^(-5/2), x)`

Giac [F]

$$\int \frac{1}{(a + bx^3)^{5/2}} dx = \int \frac{1}{(bx^3 + a)^{5/2}} dx$$

input `integrate(1/(b*x^3+a)^(5/2),x, algorithm="giac")`

output `integrate((b*x^3 + a)^(-5/2), x)`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.15

$$\int \frac{1}{(a + bx^3)^{5/2}} dx = \frac{x \left(\frac{bx^3}{a} + 1 \right)^{5/2} {}_2F_1 \left(\frac{1}{3}, \frac{5}{2}; \frac{4}{3}; -\frac{bx^3}{a} \right)}{(bx^3 + a)^{5/2}}$$

input `int(1/(a + b*x^3)^(5/2),x)`

output `(x*((b*x^3)/a + 1)^(5/2)*hypergeom([1/3, 5/2], 4/3, -(b*x^3)/a))/(a + b*x^3)^(5/2)`

Reduce [F]

$$\int \frac{1}{(a + bx^3)^{5/2}} dx = \int \frac{\sqrt{bx^3 + a}}{b^3x^9 + 3ab^2x^6 + 3a^2bx^3 + a^3} dx$$

input `int(1/(b*x^3+a)^(5/2),x)`

output `int(sqrt(a + b*x**3)/(a**3 + 3*a**2*b*x**3 + 3*a*b**2*x**6 + b**3*x**9),x)`

3.19 $\int \frac{1}{(a+bx^3)^{7/2}} dx$

Optimal result	174
Mathematica [C] (verified)	175
Rubi [A] (verified)	175
Maple [A] (verified)	177
Fricas [A] (verification not implemented)	178
Sympy [A] (verification not implemented)	178
Maxima [F]	179
Giac [F]	179
Mupad [B] (verification not implemented)	179
Reduce [F]	180

Optimal result

Integrand size = 11, antiderivative size = 270

$$\int \frac{1}{(a+bx^3)^{7/2}} dx = \frac{2x}{15a(a+bx^3)^{5/2}} + \frac{26x}{135a^2(a+bx^3)^{3/2}} + \frac{182x}{405a^3\sqrt{a+bx^3}}$$

$$+ \frac{182\sqrt{2+\sqrt{3}}(\sqrt[3]{a}+\sqrt[3]{bx})\sqrt{\frac{a^{2/3}-\sqrt[3]{a}\sqrt[3]{bx}+b^{2/3}x^2}{((1+\sqrt{3})\sqrt[3]{a}+\sqrt[3]{bx})^2}} \operatorname{EllipticF}\left(\arcsin\left(\frac{(1-\sqrt{3})\sqrt[3]{a}+\sqrt[3]{bx}}{(1+\sqrt{3})\sqrt[3]{a}+\sqrt[3]{bx}}\right), -7-4\sqrt{3}\right)}{405\sqrt[4]{3}a^3\sqrt[3]{b}\sqrt{\frac{\sqrt[3]{a}(\sqrt[3]{a}+\sqrt[3]{bx})}{((1+\sqrt{3})\sqrt[3]{a}+\sqrt[3]{bx})^2}}\sqrt{a+bx^3}}$$

output

```
2/15*x/a/(b*x^3+a)^(5/2)+26/135*x/a^2/(b*x^3+a)^(3/2)+182/405*x/a^3/(b*x^3+a)^(1/2)+182/1215*(1/2*6^(1/2)+1/2*2^(1/2))*(a^(1/3)+b^(1/3)*x)*((a^(2/3)-a^(1/3)*b^(1/3)*x+b^(2/3)*x^2)/((1+3^(1/2))*a^(1/3)+b^(1/3)*x)^2)^(1/2)*EllipticF(((1-3^(1/2))*a^(1/3)+b^(1/3)*x)/((1+3^(1/2))*a^(1/3)+b^(1/3)*x),I*3^(1/2)+2*I)*3^(3/4)/a^3/b^(1/3)/(a^(1/3)*(a^(1/3)+b^(1/3)*x)/((1+3^(1/2))*a^(1/3)+b^(1/3)*x)^2)^(1/2)/(b*x^3+a)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 4.70 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.31

$$\int \frac{1}{(a + bx^3)^{7/2}} dx = \frac{314a^2x + 442abx^4 + 182b^2x^7 + 91x(a + bx^3)^2 \sqrt{1 + \frac{bx^3}{a}} \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \frac{bx^3}{a}\right)}{405a^3 (a + bx^3)^{5/2}}$$

input

```
Integrate[(a + b*x^3)^(-7/2), x]
```

output

```
(314*a^2*x + 442*a*b*x^4 + 182*b^2*x^7 + 91*x*(a + b*x^3)^2*sqrt[1 + (b*x^3)/a]*Hypergeometric2F1[1/3, 1/2, 4/3, -(b*x^3)/a])/(405*a^3*(a + b*x^3)^(5/2))
```

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 286, normalized size of antiderivative = 1.06, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {749, 749, 749, 759}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(a + bx^3)^{7/2}} dx \\ & \quad \downarrow 749 \\ & \frac{13 \int \frac{1}{(bx^3+a)^{5/2}} dx}{15a} + \frac{2x}{15a (a + bx^3)^{5/2}} \\ & \quad \downarrow 749 \\ & \frac{13 \left(\frac{7 \int \frac{1}{(bx^3+a)^{3/2}} dx}{9a} + \frac{2x}{9a(a+bx^3)^{3/2}} \right)}{15a} + \frac{2x}{15a (a + bx^3)^{5/2}} \end{aligned}$$

$$\begin{aligned}
 & \downarrow 749 \\
 & \frac{13 \left(\frac{7 \left(\frac{\int \frac{1}{\sqrt{bx^3+a}} dx}{3a} + \frac{2x}{3a\sqrt{a+bx^3}} \right)}{9a} + \frac{2x}{9a(a+bx^3)^{3/2}} \right)}{15a} + \frac{2x}{15a(a+bx^3)^{5/2}} \\
 & \downarrow 759 \\
 & \frac{13 \left(\frac{7 \left(\frac{2\sqrt{2+\sqrt{3}} \left(\sqrt[3]{a} + \sqrt[3]{bx} \right) \sqrt{\frac{a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx} + b^{2/3}x^2}{(1+\sqrt{3})\sqrt[3]{a} + \sqrt[3]{bx}}}}^2 \text{EllipticF} \left(\arcsin \left(\frac{\sqrt[3]{bx} + (1-\sqrt{3})\sqrt[3]{a}}{\sqrt[3]{bx} + (1+\sqrt{3})\sqrt[3]{a}} \right), -7-4\sqrt{3} \right)}{3a\sqrt{a+bx^3}} \right)}{3\sqrt[4]{3a}\sqrt[3]{b} \sqrt{\frac{\sqrt[3]{a} \left(\sqrt[3]{a} + \sqrt[3]{bx} \right)}{(1+\sqrt{3})\sqrt[3]{a} + \sqrt[3]{bx}}}}^2 \sqrt{a+bx^3}} \right)}{9a} + \frac{2x}{9a(a+bx^3)^{3/2}} \right)}{15a} + \frac{2x}{15a(a+bx^3)^{5/2}}
 \end{aligned}$$

input `Int[(a + b*x^3)^(-7/2),x]`

output `(2*x)/(15*a*(a + b*x^3)^(5/2)) + (13*((2*x)/(9*a*(a + b*x^3)^(3/2)) + (7*((2*x)/(3*a*Sqrt[a + b*x^3]) + (2*Sqrt[2 + Sqrt[3]]*(a^(1/3) + b^(1/3)*x)*Sqrt[(a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/((1 + Sqrt[3])*a^(1/3) + b^(1/3)*x)^2)*EllipticF[ArcSin[((1 - Sqrt[3])*a^(1/3) + b^(1/3)*x)/((1 + Sqrt[3])*a^(1/3) + b^(1/3)*x)], -7 - 4*Sqrt[3]])/(3*3^(1/4)*a*b^(1/3)*Sqrt[(a^(1/3)*(a^(1/3) + b^(1/3)*x))/((1 + Sqrt[3])*a^(1/3) + b^(1/3)*x)^2]*Sqrt[a + b*x^3])))/(9*a)))/(15*a)`

Defintions of rubi rules used

```
rule 749 Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])
```

```
rule 759 Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[2*Sqrt[2 + Sqrt[3]]*(s + r*x)*(Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 + Sqrt[3])*s + r*x)^2]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[s*((s + r*x)/((1 + Sqrt[3])*s + r*x)^2]))*EllipticF[ArcSin[((1 - Sqrt[3])*s + r*x)/((1 + Sqrt[3])*s + r*x)], -7 - 4*Sqrt[3]], x] /; FreeQ[{a, b}, x] && PosQ[a]
```

Maple [A] (verified)

Time = 0.45 (sec) , antiderivative size = 364, normalized size of antiderivative = 1.35

method	result
default	$\frac{2x\sqrt{bx^3+a}}{15ab^3(x^3+\frac{a}{b})^3} + \frac{26x\sqrt{bx^3+a}}{135a^2b^2(x^3+\frac{a}{b})^2} + \frac{182x}{405a^3\sqrt{(x^3+\frac{a}{b})b}} - \frac{182i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{\sqrt{\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b}-\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}}{\sqrt{(-ab^2)^{\frac{1}{3}}}}}\sqrt{\frac{3(-ab^2)^{\frac{1}{3}}}{(-ab^2)^{\frac{1}{3}}}}$
elliptic	$\frac{2x\sqrt{bx^3+a}}{15ab^3(x^3+\frac{a}{b})^3} + \frac{26x\sqrt{bx^3+a}}{135a^2b^2(x^3+\frac{a}{b})^2} + \frac{182x}{405a^3\sqrt{(x^3+\frac{a}{b})b}} - \frac{182i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{\sqrt{\frac{i\left(x+\frac{(-ab^2)^{\frac{1}{3}}}{2b}-\frac{i\sqrt{3}(-ab^2)^{\frac{1}{3}}}{2b}\right)\sqrt{3}b}}{\sqrt{(-ab^2)^{\frac{1}{3}}}}}\sqrt{\frac{3(-ab^2)^{\frac{1}{3}}}{(-ab^2)^{\frac{1}{3}}}}$

```
input int(1/(b*x^3+a)^(7/2), x, method=_RETURNVERBOSE)
```

output

```
2/15*x/a/b^3*(b*x^3+a)^(1/2)/(x^3+a/b)^3+26/135*x/a^2/b^2*(b*x^3+a)^(1/2)/
(x^3+a/b)^2+182/405*x/a^3/((x^3+a/b)*b)^(1/2)-182/1215*I/a^3*3^(1/2)/b*(-a
*b^2)^(1/3)*(I*(x+1/2/b*(-a*b^2)^(1/3)-1/2*I*3^(1/2)/b*(-a*b^2)^(1/3))*3^(
1/2)*b/(-a*b^2)^(1/3))^(1/2)*((x-1/b*(-a*b^2)^(1/3))/(-3/2/b*(-a*b^2)^(1/3
)+1/2*I*3^(1/2)/b*(-a*b^2)^(1/3)))^(1/2)*(-I*(x+1/2/b*(-a*b^2)^(1/3)+1/2*I
*3^(1/2)/b*(-a*b^2)^(1/3))*3^(1/2)*b/(-a*b^2)^(1/3))^(1/2)/(b*x^3+a)^(1/2)
*EllipticF(1/3*3^(1/2)*(I*(x+1/2/b*(-a*b^2)^(1/3)-1/2*I*3^(1/2)/b*(-a*b^2)
^(1/3))*3^(1/2)*b/(-a*b^2)^(1/3))^(1/2),(I*3^(1/2)/b*(-a*b^2)^(1/3)/(-3/2/
b*(-a*b^2)^(1/3)+1/2*I*3^(1/2)/b*(-a*b^2)^(1/3)))^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.45

$$\int \frac{1}{(a + bx^3)^{7/2}} dx = \frac{2 \left(91 (b^3 x^9 + 3 ab^2 x^6 + 3 a^2 b x^3 + a^3) \sqrt{b} \text{weierstrassPInverse}(0, -\frac{4a}{b}, x) + (91 b^3 x^7 + 405 (a^3 b^4 x^9 + 3 a^4 b^3 x^6 + 3 a^5 b^2 x^3 + a^6 b)) \right)}{405 (a^3 b^4 x^9 + 3 a^4 b^3 x^6 + 3 a^5 b^2 x^3 + a^6 b)}$$

input

```
integrate(1/(b*x^3+a)^(7/2),x, algorithm="fricas")
```

output

```
2/405*(91*(b^3*x^9 + 3*a*b^2*x^6 + 3*a^2*b*x^3 + a^3)*sqrt(b)*weierstrassP
Inverse(0, -4*a/b, x) + (91*b^3*x^7 + 221*a*b^2*x^4 + 157*a^2*b*x)*sqrt(b*
x^3 + a))/(a^3*b^4*x^9 + 3*a^4*b^3*x^6 + 3*a^5*b^2*x^3 + a^6*b)
```

Sympy [A] (verification not implemented)

Time = 0.66 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.13

$$\int \frac{1}{(a + bx^3)^{7/2}} dx = \frac{x \Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{7}{2} \middle| \frac{bx^3 e^{i\pi}}{a}\right)}{3a^{\frac{7}{2}} \Gamma\left(\frac{4}{3}\right)}$$

input

```
integrate(1/(b*x**3+a)**(7/2),x)
```

output `x*gamma(1/3)*hyper((1/3, 7/2), (4/3,), b*x**3*exp_polar(I*pi)/a)/(3*a**(7/2)*gamma(4/3))`

Maxima [F]

$$\int \frac{1}{(a + bx^3)^{7/2}} dx = \int \frac{1}{(bx^3 + a)^{7/2}} dx$$

input `integrate(1/(b*x^3+a)^(7/2),x, algorithm="maxima")`

output `integrate((b*x^3 + a)^(-7/2), x)`

Giac [F]

$$\int \frac{1}{(a + bx^3)^{7/2}} dx = \int \frac{1}{(bx^3 + a)^{7/2}} dx$$

input `integrate(1/(b*x^3+a)^(7/2),x, algorithm="giac")`

output `integrate((b*x^3 + a)^(-7/2), x)`

Mupad [B] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.14

$$\int \frac{1}{(a + bx^3)^{7/2}} dx = \frac{x \left(\frac{bx^3}{a} + 1 \right)^{7/2} {}_2F_1 \left(\frac{1}{3}, \frac{7}{2}; \frac{4}{3}; -\frac{bx^3}{a} \right)}{(bx^3 + a)^{7/2}}$$

input `int(1/(a + b*x^3)^(7/2),x)`

output `(x*((b*x^3)/a + 1)^(7/2)*hypergeom([1/3, 7/2], 4/3, -(b*x^3)/a))/(a + b*x^3)^(7/2)`

Reduce [F]

$$\int \frac{1}{(a + bx^3)^{7/2}} dx = \int \frac{\sqrt{bx^3 + a}}{b^4x^{12} + 4ab^3x^9 + 6a^2b^2x^6 + 4a^3bx^3 + a^4} dx$$

input `int(1/(b*x^3+a)^(7/2),x)`

output `int(sqrt(a + b*x**3)/(a**4 + 4*a**3*b*x**3 + 6*a**2*b**2*x**6 + 4*a*b**3*x**9 + b**4*x**12),x)`

3.20 $\int \frac{1}{\sqrt{1+x^3}} dx$

Optimal result	181
Mathematica [C] (verified)	181
Rubi [A] (verified)	182
Maple [C] (verified)	183
Fricas [A] (verification not implemented)	183
Sympy [A] (verification not implemented)	184
Maxima [F]	184
Giac [F]	184
Mupad [B] (verification not implemented)	185
Reduce [F]	185

Optimal result

Integrand size = 9, antiderivative size = 103

$$\int \frac{1}{\sqrt{1+x^3}} dx = \frac{2\sqrt{2+\sqrt{3}}(1+x)\sqrt{\frac{1-x+x^2}{(1+\sqrt{3}+x)^2}} \text{EllipticF}\left(\arcsin\left(\frac{1-\sqrt{3}+x}{1+\sqrt{3}+x}\right), -7-4\sqrt{3}\right)}{\sqrt[4]{3}\sqrt{\frac{1+x}{(1+\sqrt{3}+x)^2}}\sqrt{1+x^3}}$$

output

```
2/3*(1/2*6^(1/2)+1/2*2^(1/2))*(1+x)*((x^2-x+1)/(1+x+3^(1/2))^2)^(1/2)*EllipticF((1+x-3^(1/2))/(1+x+3^(1/2)),I*3^(1/2)+2*I)*3^(3/4)/((1+x)/(1+x+3^(1/2))^2)^(1/2)/(x^3+1)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.17

$$\int \frac{1}{\sqrt{1+x^3}} dx = x \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, -x^3\right)$$

input

```
Integrate[1/Sqrt[1 + x^3],x]
```

output `x*Hypergeometric2F1[1/3, 1/2, 4/3, -x^3]`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {759}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x^3+1}} dx$$

↓ 759

$$\frac{2\sqrt{2+\sqrt{3}}(x+1)\sqrt{\frac{x^2-x+1}{(x+\sqrt{3}+1)^2}} \operatorname{EllipticF}\left(\arcsin\left(\frac{x-\sqrt{3}+1}{x+\sqrt{3}+1}\right), -7-4\sqrt{3}\right)}{\sqrt[4]{3}\sqrt{\frac{x+1}{(x+\sqrt{3}+1)^2}}\sqrt{x^3+1}}$$

input `Int[1/Sqrt[1 + x^3], x]`

output `(2*Sqrt[2 + Sqrt[3]]*(1 + x)*Sqrt[(1 - x + x^2)/(1 + Sqrt[3] + x)^2]*EllipticF[ArcSin[(1 - Sqrt[3] + x)/(1 + Sqrt[3] + x)], -7 - 4*Sqrt[3]])/(3^(1/4)*Sqrt[(1 + x)/(1 + Sqrt[3] + x)^2]*Sqrt[1 + x^3])`

Defintions of rubi rules used

rule 759 `Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[2*Sqrt[2 + Sqrt[3]]*(s + r*x)*(Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 + Sqrt[3])*s + r*x)^2]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[s*((s + r*x)/((1 + Sqrt[3])*s + r*x)^2]))*EllipticF[ArcSin[((1 - Sqrt[3])*s + r*x)/((1 + Sqrt[3])*s + r*x)], -7 - 4*Sqrt[3]], x]] /; FreeQ[{a, b}, x] && PosQ[a]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4.

Time = 0.91 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.14

method	result	size
meijerg	$x \operatorname{hypergeom}\left(\left[\frac{1}{3}, \frac{1}{2}\right], \left[\frac{4}{3}\right], -x^3\right)$	14
default	$\frac{2\left(\frac{3}{2} - \frac{i\sqrt{3}}{2}\right) \sqrt{\frac{1+x}{\frac{3}{2} - \frac{i\sqrt{3}}{2}}} \sqrt{\frac{x - \frac{1}{2} - \frac{i\sqrt{3}}{2}}{-\frac{3}{2} - \frac{i\sqrt{3}}{2}}} \sqrt{\frac{x - \frac{1}{2} + \frac{i\sqrt{3}}{2}}{-\frac{3}{2} + \frac{i\sqrt{3}}{2}}} \operatorname{EllipticF}\left(\sqrt{\frac{1+x}{\frac{3}{2} - \frac{i\sqrt{3}}{2}}}, \sqrt{\frac{-\frac{3}{2} + \frac{i\sqrt{3}}{2}}{-\frac{3}{2} - \frac{i\sqrt{3}}{2}}}\right)}{\sqrt{x^3+1}}$	116
elliptic	$\frac{2\left(\frac{3}{2} - \frac{i\sqrt{3}}{2}\right) \sqrt{\frac{1+x}{\frac{3}{2} - \frac{i\sqrt{3}}{2}}} \sqrt{\frac{x - \frac{1}{2} - \frac{i\sqrt{3}}{2}}{-\frac{3}{2} - \frac{i\sqrt{3}}{2}}} \sqrt{\frac{x - \frac{1}{2} + \frac{i\sqrt{3}}{2}}{-\frac{3}{2} + \frac{i\sqrt{3}}{2}}} \operatorname{EllipticF}\left(\sqrt{\frac{1+x}{\frac{3}{2} - \frac{i\sqrt{3}}{2}}}, \sqrt{\frac{-\frac{3}{2} + \frac{i\sqrt{3}}{2}}{-\frac{3}{2} - \frac{i\sqrt{3}}{2}}}\right)}{\sqrt{x^3+1}}$	116

input `int(1/(x^3+1)^(1/2), x, method=_RETURNVERBOSE)`

output `x*hypergeom([1/3, 1/2], [4/3], -x^3)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.06

$$\int \frac{1}{\sqrt{1+x^3}} dx = 2 \operatorname{weierstrassPInverse}(0, -4, x)$$

input `integrate(1/(x^3+1)^(1/2), x, algorithm="fricas")`

output `2*weierstrassPInverse(0, -4, x)`

Sympy [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.26

$$\int \frac{1}{\sqrt{1+x^3}} dx = \frac{x\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{1}{2} \middle| \frac{4}{3} \middle| x^3 e^{i\pi}\right)}{3\Gamma\left(\frac{4}{3}\right)}$$

input `integrate(1/(x**3+1)**(1/2),x)`output `x*gamma(1/3)*hyper((1/3, 1/2), (4/3,), x**3*exp_polar(I*pi))/(3*gamma(4/3))`**Maxima [F]**

$$\int \frac{1}{\sqrt{1+x^3}} dx = \int \frac{1}{\sqrt{x^3+1}} dx$$

input `integrate(1/(x^3+1)^(1/2),x, algorithm="maxima")`output `integrate(1/sqrt(x^3 + 1), x)`**Giac [F]**

$$\int \frac{1}{\sqrt{1+x^3}} dx = \int \frac{1}{\sqrt{x^3+1}} dx$$

input `integrate(1/(x^3+1)^(1/2),x, algorithm="giac")`output `integrate(1/sqrt(x^3 + 1), x)`

Mupad [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.50

$$\int \frac{1}{\sqrt{1+x^3}} dx$$

$$= \frac{(3 + \sqrt{3} 1i) \sqrt{\frac{x - \frac{1}{2} + \frac{\sqrt{3} 1i}{2}}{-\frac{3}{2} + \frac{\sqrt{3} 1i}{2}}} \sqrt{\frac{x+1}{\frac{3}{2} + \frac{\sqrt{3} 1i}{2}}} \sqrt{\frac{\frac{1}{2} - x + \frac{\sqrt{3} 1i}{2}}{\frac{3}{2} + \frac{\sqrt{3} 1i}{2}}} F\left(\operatorname{asin}\left(\sqrt{\frac{x+1}{\frac{3}{2} + \frac{\sqrt{3} 1i}{2}}}\right) \middle| -\frac{\frac{3}{2} + \frac{\sqrt{3} 1i}{2}}{-\frac{3}{2} + \frac{\sqrt{3} 1i}{2}}\right)}{\sqrt{x^3 + \left(-\left(-\frac{1}{2} + \frac{\sqrt{3} 1i}{2}\right)\left(\frac{1}{2} + \frac{\sqrt{3} 1i}{2}\right) - 1\right) x - \left(-\frac{1}{2} + \frac{\sqrt{3} 1i}{2}\right)\left(\frac{1}{2} + \frac{\sqrt{3} 1i}{2}\right)}}$$

input `int(1/(x^3 + 1)^(1/2), x)`output `((3^(1/2)*1i + 3)*((x + (3^(1/2)*1i)/2 - 1/2)/((3^(1/2)*1i)/2 - 3/2))^(1/2) * ((x + 1)/((3^(1/2)*1i)/2 + 3/2))^(1/2) * ((3^(1/2)*1i)/2 - x + 1/2)/((3^(1/2)*1i)/2 + 3/2)^(1/2) * ellipticF(asin(((x + 1)/((3^(1/2)*1i)/2 + 3/2))^(1/2)), -((3^(1/2)*1i)/2 + 3/2)/((3^(1/2)*1i)/2 - 3/2))/(x^3 - x*((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2) + 1) - ((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2))^(1/2)`**Reduce [F]**

$$\int \frac{1}{\sqrt{1+x^3}} dx = \int \frac{\sqrt{x^3+1}}{x^3+1} dx$$

input `int(1/(x^3+1)^(1/2), x)`output `int(sqrt(x**3 + 1)/(x**3 + 1), x)`

3.21 $\int \frac{1}{\sqrt{1-x^3}} dx$

Optimal result	186
Mathematica [C] (verified)	186
Rubi [A] (verified)	187
Maple [C] (verified)	188
Fricas [A] (verification not implemented)	189
Sympy [A] (verification not implemented)	189
Maxima [F]	189
Giac [F]	190
Mupad [B] (verification not implemented)	190
Reduce [F]	191

Optimal result

Integrand size = 11, antiderivative size = 115

$$\int \frac{1}{\sqrt{1-x^3}} dx = \frac{2\sqrt{2+\sqrt{3}}(1-x)\sqrt{\frac{1+x+x^2}{(1+\sqrt{3}-x)^2}} \text{EllipticF}\left(\arcsin\left(\frac{1-\sqrt{3}-x}{1+\sqrt{3}-x}\right), -7-4\sqrt{3}\right)}{\sqrt[4]{3}\sqrt{\frac{1-x}{(1+\sqrt{3}-x)^2}}\sqrt{1-x^3}}$$

output

```
-2/3*(1/2*6^(1/2)+1/2*2^(1/2))*(1-x)*((x^2+x+1)/(1+3^(1/2)-x)^2)^(1/2)*EllipticF((1-3^(1/2)-x)/(1+3^(1/2)-x), I*3^(1/2)+2*I)*3^(3/4)/((1-x)/(1+3^(1/2)-x)^2)^(1/2)/(-x^3+1)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.13

$$\int \frac{1}{\sqrt{1-x^3}} dx = x \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, x^3\right)$$

input `Integrate[1/Sqrt[1 - x^3],x]`

output `x*Hypergeometric2F1[1/3, 1/2, 4/3, x^3]`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {759}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{1-x^3}} dx$$

↓ 759

$$\frac{2\sqrt{2+\sqrt{3}}(1-x)\sqrt{\frac{x^2+x+1}{(-x+\sqrt{3}+1)^2}} \operatorname{EllipticF}\left(\arcsin\left(\frac{-x-\sqrt{3}+1}{-x+\sqrt{3}+1}\right), -7-4\sqrt{3}\right)}{\sqrt[4]{3}\sqrt{\frac{1-x}{(-x+\sqrt{3}+1)^2}}\sqrt{1-x^3}}$$

input `Int[1/Sqrt[1 - x^3],x]`

output `(-2*Sqrt[2 + Sqrt[3]]*(1 - x)*Sqrt[(1 + x + x^2)/(1 + Sqrt[3] - x)^2]*EllipticF[ArcSin[(1 - Sqrt[3] - x)/(1 + Sqrt[3] - x)], -7 - 4*Sqrt[3]])/(3^(1/4)*Sqrt[(1 - x)/(1 + Sqrt[3] - x)^2]*Sqrt[1 - x^3])`

Defintions of rubi rules used

rule 759

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[2*Sqrt[2 + Sqrt[3]]*(s + r*x)*(Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 + Sqrt[3])*s + r*x)^2]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[s*
((s + r*x)/((1 + Sqrt[3])*s + r*x)^2)))*EllipticF[ArcSin[((1 - Sqrt[3])*s
+ r*x)/((1 + Sqrt[3])*s + r*x)], -7 - 4*Sqrt[3]], x] /; FreeQ[{a, b}, x] &
& PosQ[a]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4.

Time = 0.82 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.10

method	result	size
meijerg	$x \operatorname{hypergeom}\left(\left[\frac{1}{3}, \frac{1}{2}\right], \left[\frac{4}{3}\right], x^3\right)$	12
default	$\frac{2i\sqrt{3} \sqrt{i\left(x+\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)}\sqrt{3} \sqrt{\frac{-1+x}{-\frac{3}{2}+\frac{i\sqrt{3}}{2}}} \sqrt{-i\left(x+\frac{1}{2}+\frac{i\sqrt{3}}{2}\right)}\sqrt{3} \operatorname{EllipticF}\left(\frac{\sqrt{3} \sqrt{i\left(x+\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)}\sqrt{3}}{3}, \sqrt{\frac{i\sqrt{3}}{-\frac{3}{2}+\frac{i\sqrt{3}}{2}}}\right)}{3\sqrt{-x^3+1}}$	107
elliptic	$\frac{2i\sqrt{3} \sqrt{i\left(x+\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)}\sqrt{3} \sqrt{\frac{-1+x}{-\frac{3}{2}+\frac{i\sqrt{3}}{2}}} \sqrt{-i\left(x+\frac{1}{2}+\frac{i\sqrt{3}}{2}\right)}\sqrt{3} \operatorname{EllipticF}\left(\frac{\sqrt{3} \sqrt{i\left(x+\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)}\sqrt{3}}{3}, \sqrt{\frac{i\sqrt{3}}{-\frac{3}{2}+\frac{i\sqrt{3}}{2}}}\right)}{3\sqrt{-x^3+1}}$	107

input `int(1/(-x^3+1)^(1/2), x, method=_RETURNVERBOSE)`

output `x*hypergeom([1/3, 1/2], [4/3], x^3)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.05

$$\int \frac{1}{\sqrt{1-x^3}} dx = -2i \operatorname{weierstrassPInverse}(0, 4, x)$$

input `integrate(1/(-x^3+1)^(1/2),x, algorithm="fricas")`output `-2*I*weierstrassPInverse(0, 4, x)`**Sympy [A] (verification not implemented)**

Time = 0.37 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.25

$$\int \frac{1}{\sqrt{1-x^3}} dx = \frac{x\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{1}{2} \middle| \frac{4}{3} \middle| x^3 e^{2i\pi}\right)}{3\Gamma\left(\frac{4}{3}\right)}$$

input `integrate(1/(-x**3+1)**(1/2),x)`output `x*gamma(1/3)*hyper((1/3, 1/2), (4/3,), x**3*exp_polar(2*I*pi))/(3*gamma(4/3))`**Maxima [F]**

$$\int \frac{1}{\sqrt{1-x^3}} dx = \int \frac{1}{\sqrt{-x^3+1}} dx$$

input `integrate(1/(-x^3+1)^(1/2),x, algorithm="maxima")`output `integrate(1/sqrt(-x^3 + 1), x)`

Giac [F]

$$\int \frac{1}{\sqrt{1-x^3}} dx = \int \frac{1}{\sqrt{-x^3+1}} dx$$

input `integrate(1/(-x^3+1)^(1/2),x, algorithm="giac")`

output `integrate(1/sqrt(-x^3 + 1), x)`

Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.50

$$\int \frac{1}{\sqrt{1-x^3}} dx = \frac{2 \left(\frac{3}{2} + \frac{\sqrt{3}1i}{2} \right) \sqrt{x^3-1} \sqrt{-\frac{x+\frac{1}{2}-\frac{\sqrt{3}1i}{2}}{-\frac{3}{2}+\frac{\sqrt{3}1i}{2}}} \sqrt{\frac{x+\frac{1}{2}+\frac{\sqrt{3}1i}{2}}{\frac{3}{2}+\frac{\sqrt{3}1i}{2}}} \sqrt{-\frac{x-1}{\frac{3}{2}+\frac{\sqrt{3}1i}{2}}} F\left(\operatorname{asin}\left(\sqrt{-\frac{x-1}{\frac{3}{2}+\frac{\sqrt{3}1i}{2}}}\right) \middle| -\frac{\frac{3}{2}+\frac{\sqrt{3}1i}{2}}{-\frac{3}{2}+\frac{\sqrt{3}1i}{2}}\right)}{\sqrt{1-x^3} \sqrt{x^3 + \left(-\left(-\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right) \left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right) - 1\right) x + \left(-\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right) \left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)}}$$

input `int(1/(1 - x^3)^(1/2),x)`

output `-(2*((3^(1/2)*1i)/2 + 3/2)*(x^3 - 1)^(1/2)*(-(x - (3^(1/2)*1i)/2 + 1/2)/((3^(1/2)*1i)/2 - 3/2))^(1/2)*((x + (3^(1/2)*1i)/2 + 1/2)/((3^(1/2)*1i)/2 + 3/2))^(1/2)*(-(x - 1)/((3^(1/2)*1i)/2 + 3/2))^(1/2)*ellipticF(asin((-x - 1)/((3^(1/2)*1i)/2 + 3/2))^(1/2), -((3^(1/2)*1i)/2 + 3/2)/((3^(1/2)*1i)/2 - 3/2))/((1 - x^3)^(1/2)*(((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2) - x*((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2) + x^3)^(1/2))`

Reduce [F]

$$\int \frac{1}{\sqrt{1-x^3}} dx = - \left(\int \frac{\sqrt{-x^3+1}}{x^3-1} dx \right)$$

input `int(1/(-x3+1)(1/2),x)`

output `- int(sqrt(- x**3 + 1)/(x**3 - 1),x)`

3.22 $\int \frac{1}{\sqrt{-1+x^3}} dx$

Optimal result	192
Mathematica [C] (verified)	192
Rubi [A] (verified)	193
Maple [C] (warning: unable to verify)	194
Fricas [A] (verification not implemented)	195
Sympy [A] (verification not implemented)	195
Maxima [F]	195
Giac [F]	196
Mupad [B] (verification not implemented)	196
Reduce [F]	197

Optimal result

Integrand size = 9, antiderivative size = 120

$$\int \frac{1}{\sqrt{-1+x^3}} dx = \frac{2\sqrt{2-\sqrt{3}}(1-x)\sqrt{\frac{1+x+x^2}{(1-\sqrt{3}-x)^2}} \operatorname{EllipticF}\left(\arcsin\left(\frac{1+\sqrt{3}-x}{1-\sqrt{3}-x}\right), -7+4\sqrt{3}\right)}{\sqrt[4]{3}\sqrt{-\frac{1-x}{(1-\sqrt{3}-x)^2}}\sqrt{-1+x^3}}$$

```
output -2/3*(1/2*6^(1/2)-1/2*2^(1/2))*(1-x)*((x^2+x+1)/(1-3^(1/2)-x)^2)^(1/2)*EllipticF((1+3^(1/2)-x)/(1-3^(1/2)-x),2*I-I*3^(1/2))*3^(3/4)/(-(1-x)/(1-3^(1/2)-x)^2)^(1/2)/(x^3-1)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.02 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.29

$$\int \frac{1}{\sqrt{-1+x^3}} dx = \frac{x\sqrt{1-x^3} \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, x^3\right)}{\sqrt{-1+x^3}}$$

input `Integrate[1/Sqrt[-1 + x^3],x]`

output `(x*Sqrt[1 - x^3]*Hypergeometric2F1[1/3, 1/2, 4/3, x^3])/Sqrt[-1 + x^3]`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {760}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x^3 - 1}} dx$$

↓ 760

$$\frac{2\sqrt{2 - \sqrt{3}}(1 - x) \sqrt{\frac{x^2 + x + 1}{(-x - \sqrt{3} + 1)^2}} \text{EllipticF}\left(\arcsin\left(\frac{-x + \sqrt{3} + 1}{-x - \sqrt{3} + 1}\right), -7 + 4\sqrt{3}\right)}{\sqrt[4]{3} \sqrt{-\frac{1 - x}{(-x - \sqrt{3} + 1)^2}} \sqrt{x^3 - 1}}$$

input `Int[1/Sqrt[-1 + x^3],x]`

output `(-2*Sqrt[2 - Sqrt[3]]*(1 - x)*Sqrt[(1 + x + x^2)/(1 - Sqrt[3] - x)^2]*EllipticF[ArcSin[(1 + Sqrt[3] - x)/(1 - Sqrt[3] - x)], -7 + 4*Sqrt[3]])/(3^(1/4)*Sqrt[-((1 - x)/(1 - Sqrt[3] - x)^2)]*Sqrt[-1 + x^3])`

Defintions of rubi rules used

rule 760

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] :> With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[2*Sqrt[2 - Sqrt[3]]*(s + r*x)*(Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[(-
s)*((s + r*x)/((1 - Sqrt[3])*s + r*x)^2)))*EllipticF[ArcSin[((1 + Sqrt[3])
*s + r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]], x] /; FreeQ[{a, b}, x
] && NegQ[a]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.90 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.25

method	result	size
meijerg	$\frac{\sqrt{-\text{signum}(x^3-1)} x \text{ hypergeom}\left(\left[\frac{1}{3}, \frac{1}{2}\right], \left[\frac{4}{3}\right], x^3\right)}{\sqrt{\text{signum}(x^3-1)}}$	30
default	$\frac{2\left(-\frac{3}{2} - \frac{i\sqrt{3}}{2}\right) \sqrt{\frac{-1+x}{-\frac{3}{2} - \frac{i\sqrt{3}}{2}}} \sqrt{\frac{x+\frac{1}{2} - \frac{i\sqrt{3}}{2}}{\frac{3}{2} - \frac{i\sqrt{3}}{2}}} \sqrt{\frac{x+\frac{1}{2} + \frac{i\sqrt{3}}{2}}{\frac{3}{2} + \frac{i\sqrt{3}}{2}}} \text{EllipticF}\left(\sqrt{\frac{-1+x}{-\frac{3}{2} - \frac{i\sqrt{3}}{2}}}, \sqrt{\frac{\frac{3}{2} + \frac{i\sqrt{3}}{2}}{\frac{3}{2} - \frac{i\sqrt{3}}{2}}}\right)}{\sqrt{x^3-1}}$	116
elliptic	$\frac{2\left(-\frac{3}{2} - \frac{i\sqrt{3}}{2}\right) \sqrt{\frac{-1+x}{-\frac{3}{2} - \frac{i\sqrt{3}}{2}}} \sqrt{\frac{x+\frac{1}{2} - \frac{i\sqrt{3}}{2}}{\frac{3}{2} - \frac{i\sqrt{3}}{2}}} \sqrt{\frac{x+\frac{1}{2} + \frac{i\sqrt{3}}{2}}{\frac{3}{2} + \frac{i\sqrt{3}}{2}}} \text{EllipticF}\left(\sqrt{\frac{-1+x}{-\frac{3}{2} - \frac{i\sqrt{3}}{2}}}, \sqrt{\frac{\frac{3}{2} + \frac{i\sqrt{3}}{2}}{\frac{3}{2} - \frac{i\sqrt{3}}{2}}}\right)}{\sqrt{x^3-1}}$	116

input

```
int(1/(x^3-1)^(1/2), x, method=_RETURNVERBOSE)
```

output

```
1/signum(x^3-1)^(1/2)*(-signum(x^3-1))^(1/2)*x*hypergeom([1/3,1/2],[4/3],x^3)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.05

$$\int \frac{1}{\sqrt{-1+x^3}} dx = 2 \operatorname{weierstrassPInverse}(0, 4, x)$$

input `integrate(1/(x^3-1)^(1/2),x, algorithm="fricas")`output `2*weierstrassPInverse(0, 4, x)`**Sympy [A] (verification not implemented)**

Time = 0.37 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.22

$$\int \frac{1}{\sqrt{-1+x^3}} dx = -\frac{ix\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{1}{2} \middle| \frac{4}{3} \middle| x^3\right)}{3\Gamma\left(\frac{4}{3}\right)}$$

input `integrate(1/(x**3-1)**(1/2),x)`output `-I*x*gamma(1/3)*hyper((1/3, 1/2), (4/3,), x**3)/(3*gamma(4/3))`**Maxima [F]**

$$\int \frac{1}{\sqrt{-1+x^3}} dx = \int \frac{1}{\sqrt{x^3-1}} dx$$

input `integrate(1/(x^3-1)^(1/2),x, algorithm="maxima")`output `integrate(1/sqrt(x^3 - 1), x)`

Giac [F]

$$\int \frac{1}{\sqrt{-1+x^3}} dx = \int \frac{1}{\sqrt{x^3-1}} dx$$

input `integrate(1/(x^3-1)^(1/2),x, algorithm="giac")`

output `integrate(1/sqrt(x^3 - 1), x)`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.30

$$\int \frac{1}{\sqrt{-1+x^3}} dx = \frac{(3 + \sqrt{3} i) \sqrt{-\frac{x+\frac{1}{2}-\frac{\sqrt{3}i}{2}}{-\frac{3}{2}+\frac{\sqrt{3}i}{2}}} \sqrt{\frac{x+\frac{1}{2}+\frac{\sqrt{3}i}{2}}{\frac{3}{2}+\frac{\sqrt{3}i}{2}}} \sqrt{-\frac{x-1}{\frac{3}{2}+\frac{\sqrt{3}i}{2}}} F\left(\operatorname{asin}\left(\sqrt{-\frac{x-1}{\frac{3}{2}+\frac{\sqrt{3}i}{2}}}\right) \middle| -\frac{\frac{3}{2}+\frac{\sqrt{3}i}{2}}{-\frac{3}{2}+\frac{\sqrt{3}i}{2}}\right)}{\sqrt{x^3 + \left(-\left(-\frac{1}{2} + \frac{\sqrt{3}i}{2}\right) \left(\frac{1}{2} + \frac{\sqrt{3}i}{2}\right) - 1\right) x + \left(-\frac{1}{2} + \frac{\sqrt{3}i}{2}\right) \left(\frac{1}{2} + \frac{\sqrt{3}i}{2}\right)}}$$

input `int(1/(x^3 - 1)^(1/2),x)`

output `-((3^(1/2)*1i + 3)*(-(x - (3^(1/2)*1i)/2 + 1/2)/((3^(1/2)*1i)/2 - 3/2))^(1/2)*((x + (3^(1/2)*1i)/2 + 1/2)/((3^(1/2)*1i)/2 + 3/2))^(1/2)*(-(x - 1)/((3^(1/2)*1i)/2 + 3/2))^(1/2)*ellipticF(asin(-(x - 1)/((3^(1/2)*1i)/2 + 3/2))^(1/2), -(3^(1/2)*1i)/2 + 3/2)/((3^(1/2)*1i)/2 - 3/2))/(((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2) - x*(((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2) + 1) + x^3)^(1/2)`

Reduce [F]

$$\int \frac{1}{\sqrt{-1+x^3}} dx = \int \frac{\sqrt{x^3-1}}{x^3-1} dx$$

input `int(1/(x^3-1)^(1/2),x)`

output `int(sqrt(x**3 - 1)/(x**3 - 1),x)`

3.23 $\int \frac{1}{\sqrt{-1-x^3}} dx$

Optimal result	198
Mathematica [C] (verified)	198
Rubi [A] (verified)	199
Maple [C] (verified)	200
Fricas [A] (verification not implemented)	201
Sympy [A] (verification not implemented)	201
Maxima [F]	201
Giac [F]	202
Mupad [B] (verification not implemented)	202
Reduce [F]	203

Optimal result

Integrand size = 11, antiderivative size = 112

$$\int \frac{1}{\sqrt{-1-x^3}} dx = \frac{2\sqrt{2-\sqrt{3}}(1+x) \sqrt{\frac{1-x+x^2}{(1-\sqrt{3}+x)^2}} \text{EllipticF}\left(\arcsin\left(\frac{1+\sqrt{3}+x}{1-\sqrt{3}+x}\right), -7+4\sqrt{3}\right)}{\sqrt[4]{3} \sqrt{-\frac{1+x}{(1-\sqrt{3}+x)^2}} \sqrt{-1-x^3}}$$

output `2/3*(1/2*6^(1/2)-1/2*2^(1/2))*(1+x)*((x^2-x+1)/(1+x-3^(1/2)))^(1/2)*EllipticF((1+x+3^(1/2))/(1+x-3^(1/2)),2*I-I*3^(1/2))*3^(3/4)/(-(1+x)/(1+x-3^(1/2)))^(1/2)/(-x^3-1)^(1/2)`

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.02 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.33

$$\int \frac{1}{\sqrt{-1-x^3}} dx = \frac{x\sqrt{1+x^3} \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, -x^3\right)}{\sqrt{-1-x^3}}$$

input `Integrate[1/Sqrt[-1 - x^3],x]`

output `(x*Sqrt[1 + x^3]*Hypergeometric2F1[1/3, 1/2, 4/3, -x^3])/Sqrt[-1 - x^3]`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {760}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{-x^3 - 1}} dx$$

↓ 760

$$\frac{2\sqrt{2 - \sqrt{3}}(x + 1) \sqrt{\frac{x^2 - x + 1}{(x - \sqrt{3} + 1)^2}} \text{EllipticF}\left(\arcsin\left(\frac{x + \sqrt{3} + 1}{x - \sqrt{3} + 1}\right), -7 + 4\sqrt{3}\right)}{\sqrt[4]{3} \sqrt{-\frac{x + 1}{(x - \sqrt{3} + 1)^2}} \sqrt{-x^3 - 1}}$$

input `Int[1/Sqrt[-1 - x^3],x]`

output `(2*Sqrt[2 - Sqrt[3]]*(1 + x)*Sqrt[(1 - x + x^2)/(1 - Sqrt[3] + x)^2]*EllipticF[ArcSin[(1 + Sqrt[3] + x)/(1 - Sqrt[3] + x)], -7 + 4*Sqrt[3]])/(3^(1/4)*Sqrt[-((1 + x)/(1 - Sqrt[3] + x)^2)]*Sqrt[-1 - x^3])`

Defintions of rubi rules used

rule 760

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] :> With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[2*Sqrt[2 - Sqrt[3]]*(s + r*x)*(Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[(-
s)*((s + r*x)/((1 - Sqrt[3])*s + r*x)^2)])]*EllipticF[ArcSin[((1 + Sqrt[3])
*s + r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]], x]] /; FreeQ[{a, b}, x
] && NegQ[a]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4.

Time = 0.68 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.14

method	result	size
meijerg	$-ix \operatorname{hypergeom}\left(\left[\frac{1}{3}, \frac{1}{2}\right], \left[\frac{4}{3}\right], -x^3\right)$	16
default	$\frac{2i\sqrt{3} \sqrt{i\left(x-\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{3}} \sqrt{\frac{1+x}{\frac{3}{2}+\frac{i\sqrt{3}}{2}}} \sqrt{-i\left(x-\frac{1}{2}+\frac{i\sqrt{3}}{2}\right)\sqrt{3}} \operatorname{EllipticF}\left(\frac{\sqrt{3} \sqrt{i\left(x-\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{3}}}{3}, \sqrt{\frac{i\sqrt{3}}{\frac{3}{2}+\frac{i\sqrt{3}}{2}}}\right)}{3\sqrt{-x^3-1}}$	107
elliptic	$\frac{2i\sqrt{3} \sqrt{i\left(x-\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{3}} \sqrt{\frac{1+x}{\frac{3}{2}+\frac{i\sqrt{3}}{2}}} \sqrt{-i\left(x-\frac{1}{2}+\frac{i\sqrt{3}}{2}\right)\sqrt{3}} \operatorname{EllipticF}\left(\frac{\sqrt{3} \sqrt{i\left(x-\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{3}}}{3}, \sqrt{\frac{i\sqrt{3}}{\frac{3}{2}+\frac{i\sqrt{3}}{2}}}\right)}{3\sqrt{-x^3-1}}$	107

input

```
int(1/(-x^3-1)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
-I*x*hypergeom([1/3,1/2],[4/3],-x^3)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.05

$$\int \frac{1}{\sqrt{-1-x^3}} dx = -2i \operatorname{weierstrassPInverse}(0, -4, x)$$

input `integrate(1/(-x^3-1)^(1/2),x, algorithm="fricas")`output `-2*I*weierstrassPInverse(0, -4, x)`**Sympy [A] (verification not implemented)**

Time = 0.34 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.28

$$\int \frac{1}{\sqrt{-1-x^3}} dx = -\frac{ix\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{1}{2} \middle| \frac{4}{3} \middle| x^3 e^{i\pi}\right)}{3\Gamma\left(\frac{4}{3}\right)}$$

input `integrate(1/(-x**3-1)**(1/2),x)`output `-I*x*gamma(1/3)*hyper((1/3, 1/2), (4/3,), x**3*exp_polar(I*pi))/(3*gamma(4/3))`**Maxima [F]**

$$\int \frac{1}{\sqrt{-1-x^3}} dx = \int \frac{1}{\sqrt{-x^3-1}} dx$$

input `integrate(1/(-x^3-1)^(1/2),x, algorithm="maxima")`output `integrate(1/sqrt(-x^3 - 1), x)`

Giac [F]

$$\int \frac{1}{\sqrt{-1-x^3}} dx = \int \frac{1}{\sqrt{-x^3-1}} dx$$

input `integrate(1/(-x^3-1)^(1/2),x, algorithm="giac")`

output `integrate(1/sqrt(-x^3 - 1), x)`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.54

$$\int \frac{1}{\sqrt{-1-x^3}} dx$$

$$= \frac{2 \left(\frac{3}{2} + \frac{\sqrt{3}1i}{2} \right) \sqrt{x^3+1} \sqrt{\frac{x-\frac{1}{2}+\frac{\sqrt{3}1i}{2}}{-\frac{3}{2}+\frac{\sqrt{3}1i}{2}}} \sqrt{\frac{x+1}{\frac{3}{2}+\frac{\sqrt{3}1i}{2}}} \sqrt{\frac{\frac{1}{2}-x+\frac{\sqrt{3}1i}{2}}{\frac{3}{2}+\frac{\sqrt{3}1i}{2}}} F\left(\operatorname{asin}\left(\sqrt{\frac{x+1}{\frac{3}{2}+\frac{\sqrt{3}1i}{2}}}\right) \middle| -\frac{\frac{3}{2}+\frac{\sqrt{3}1i}{2}}{-\frac{3}{2}+\frac{\sqrt{3}1i}{2}}\right)}{\sqrt{-x^3-1} \sqrt{x^3 + \left(-\left(-\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right) \left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right) - 1\right) x - \left(-\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right) \left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)}$$

input `int(1/(- x^3 - 1)^(1/2),x)`

output `(2*((3^(1/2)*1i)/2 + 3/2)*(x^3 + 1)^(1/2)*((x + (3^(1/2)*1i)/2 - 1/2)/((3^(1/2)*1i)/2 - 3/2))^(1/2)*((x + 1)/((3^(1/2)*1i)/2 + 3/2))^(1/2)*(((3^(1/2)*1i)/2 - x + 1/2)/((3^(1/2)*1i)/2 + 3/2))^(1/2)*ellipticF(asin(((x + 1)/((3^(1/2)*1i)/2 + 3/2))^(1/2)), -((3^(1/2)*1i)/2 + 3/2)/((3^(1/2)*1i)/2 - 3/2))/((- x^3 - 1)^(1/2)*(x^3 - x*((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2) + 1) - ((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2))^(1/2)`

Reduce [F]

$$\int \frac{1}{\sqrt{-1-x^3}} dx = - \left(\int \frac{\sqrt{x^3+1}}{x^3+1} dx \right) i$$

input `int(1/(-x^3-1)^(1/2),x)`

output `- int(sqrt(x**3 + 1)/(x**3 + 1),x)*i`

3.24 $\int (a + bx^3)^{5/3} dx$

Optimal result	204
Mathematica [C] (verified)	204
Rubi [A] (verified)	205
Maple [A] (verified)	207
Fricas [B] (verification not implemented)	207
Sympy [C] (verification not implemented)	208
Maxima [B] (verification not implemented)	209
Giac [F]	209
Mupad [B] (verification not implemented)	210
Reduce [F]	210

Optimal result

Integrand size = 11, antiderivative size = 112

$$\int (a + bx^3)^{5/3} dx = \frac{5}{18}ax(a + bx^3)^{2/3} + \frac{1}{6}x(a + bx^3)^{5/3} + \frac{5a^2 \arctan\left(\frac{1 + \frac{2\sqrt[3]{b}x}{\sqrt[3]{a + bx^3}}}{\sqrt{3}}\right)}{9\sqrt{3}\sqrt[3]{b}} - \frac{5a^2 \log\left(-\sqrt[3]{b}x + \sqrt[3]{a + bx^3}\right)}{18\sqrt[3]{b}}$$

output

```
5/18*a*x*(b*x^3+a)^(2/3)+1/6*x*(b*x^3+a)^(5/3)+5/27*a^2*arctan(1/3*(1+2*b^(1/3)*x/(b*x^3+a)^(1/3))*3^(1/2))*3^(1/2)/b^(1/3)-5/18*a^2*ln(-b^(1/3)*x+(b*x^3+a)^(1/3))/b^(1/3)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 0.17 (sec) , antiderivative size = 272, normalized size of antiderivative = 2.43

$$\int (a + bx^3)^{5/3} dx = \frac{3 \left(\frac{(-1)^{2/3} \sqrt[3]{a}}{\sqrt[3]{b}} + x \right) (a + bx^3)^{5/3} \operatorname{AppellF1} \left(\frac{8}{3}, -\frac{5}{3}, -\frac{5}{3}, \frac{11}{3}, -\frac{(-1)^{2/3} \sqrt[3]{a} + x}{\sqrt[3]{b}}, -\frac{(-1)^{2/3} \sqrt[3]{a}}{\sqrt[3]{b}} \right)}{8 \left(1 + \frac{(-1)^{2/3} \sqrt[3]{a} + x}{\sqrt[3]{b}} \right)^{5/3} \left(1 + \frac{(-1)^{2/3} \sqrt[3]{a} + x}{\sqrt[3]{b}} \right)^{5/3}}$$

input

```
Integrate[(a + b*x^3)^(5/3), x]
```

output

```
(3*(((−1)^(2/3)*a^(1/3))/b^(1/3) + x)*(a + b*x^3)^(5/3)*AppellF1[8/3, −5/3, −5/3, 11/3, −(((−1)^(2/3)*a^(1/3))/b^(1/3) + x)/((−1)^(1/3)*a^(1/3))/b^(1/3) − ((−1)^(2/3)*a^(1/3))/b^(1/3)], −(((−1)^(2/3)*a^(1/3))/b^(1/3) + x)/(a^(1/3)/b^(1/3) − ((−1)^(2/3)*a^(1/3))/b^(1/3)))]/(8*(1 + (((−1)^(2/3)*a^(1/3))/b^(1/3) + x)/(a^(1/3)/b^(1/3) − ((−1)^(2/3)*a^(1/3))/b^(1/3)))^(5/3)*(1 + (((−1)^(2/3)*a^(1/3))/b^(1/3) + x)/((−1)^(1/3)*a^(1/3))/b^(1/3) − ((−1)^(2/3)*a^(1/3))/b^(1/3)))^(5/3))
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.02, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {748, 748, 769}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx^3)^{5/3} dx$$

↓ 748

$$\frac{5}{6}a \int (bx^3 + a)^{2/3} dx + \frac{1}{6}x(a + bx^3)^{5/3}$$

$$\begin{array}{c}
 \downarrow 748 \\
 \frac{5}{6}a \left(\frac{2}{3}a \int \frac{1}{\sqrt[3]{bx^3+a}} dx + \frac{1}{3}x(a+bx^3)^{2/3} \right) + \frac{1}{6}x(a+bx^3)^{5/3} \\
 \downarrow 769 \\
 \frac{5}{6}a \left(\frac{2}{3}a \left(\frac{\arctan\left(\frac{\frac{2\sqrt[3]{b}x+1}{\sqrt[3]{a+bx^3}}}{\sqrt{3}}\right)}{\sqrt{3}\sqrt[3]{b}} - \frac{\log\left(\frac{\sqrt[3]{a+bx^3}-\sqrt[3]{bx}}{2\sqrt[3]{b}}\right)}{2\sqrt[3]{b}} \right) + \frac{1}{3}x(a+bx^3)^{2/3} \right) + \frac{1}{6}x(a+bx^3)^{5/3}
 \end{array}$$

input `Int[(a + b*x^3)^(5/3), x]`

output `(x*(a + b*x^3)^(5/3))/6 + (5*a*((x*(a + b*x^3)^(2/3))/3 + (2*a*(ArcTan[(1 + (2*b^(1/3)*x)/(a + b*x^3)^(1/3)]/Sqrt[3])/(Sqrt[3]*b^(1/3)) - Log[-(b^(1/3)*x) + (a + b*x^3)^(1/3)]/(2*b^(1/3)))]/3))/6`

Defintions of rubi rules used

rule 748 `Int[((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p + 1)), x] + Simp[a*n*(p/(n*p + 1)) Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || LtQ[Denominator[p + 1/n], Denominator[p]])`

rule 769 `Int[((a_) + (b_)*(x_)^3)^(-1/3), x_Symbol] := Simp[ArcTan[(1 + 2*Rt[b, 3]*(x/(a + b*x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*Rt[b, 3]), x] - Simp[Log[(a + b*x^3)^(1/3) - Rt[b, 3]*x]/(2*Rt[b, 3]), x] /; FreeQ[{a, b}, x]`

Maple [A] (verified)

Time = 2.27 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.29

method	result
pseudoelliptic	$\frac{9(bx^3+a)^{\frac{2}{3}}b^{\frac{4}{3}}x^4+24ax(bx^3+a)^{\frac{2}{3}}b^{\frac{1}{3}}-10\sqrt{3}\arctan\left(\frac{\sqrt{3}\left(b^{\frac{1}{3}}x+2(bx^3+a)^{\frac{1}{3}}\right)}{3b^{\frac{1}{3}}x}\right)a^2-10\ln\left(\frac{-b^{\frac{1}{3}}x+(bx^3+a)^{\frac{1}{3}}}{x}\right)a^2+5\ln\left(\frac{-b^{\frac{1}{3}}x+(bx^3+a)^{\frac{1}{3}}}{x}\right)a^2}{54b^{\frac{1}{3}}}$

input `int((b*x^3+a)^(5/3),x,method=_RETURNVERBOSE)`

output `1/54*(9*(b*x^3+a)^(2/3)*b^(4/3)*x^4+24*a*x*(b*x^3+a)^(2/3)*b^(1/3)-10*3^(1/2)*arctan(1/3*3^(1/2)*(b^(1/3)*x+2*(b*x^3+a)^(1/3))/b^(1/3)/x)*a^2-10*ln((-b^(1/3)*x+(b*x^3+a)^(1/3))/x)*a^2+5*ln((b^(2/3)*x^2+b^(1/3)*(b*x^3+a)^(1/3)*x+(b*x^3+a)^(2/3))/x^2)*a^2)/b^(1/3)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 175 vs. 2(85) = 170.

Time = 0.09 (sec) , antiderivative size = 399, normalized size of antiderivative = 3.56

$$\int (a + bx^3)^{5/3} dx = \frac{15\sqrt{\frac{1}{3}}a^2b\sqrt{\frac{(-b)^{\frac{1}{3}}}{b}}\log\left(3bx^3 - 3(bx^3 + a)^{\frac{1}{3}}(-b)^{\frac{2}{3}}x^2 - 3\sqrt{\frac{1}{3}}\left((-b)^{\frac{1}{3}}bx^3 - (bx^3 + a)^{\frac{1}{3}}bx^2 + (-b)^{\frac{2}{3}}x\right)\right)}{30\sqrt{\frac{1}{3}}a^2b\sqrt{-\frac{(-b)^{\frac{1}{3}}}{b}}\arctan\left(-\frac{\sqrt{\frac{1}{3}}\left((-b)^{\frac{1}{3}}x - 2(bx^3 + a)^{\frac{1}{3}}\right)\sqrt{-\frac{(-b)^{\frac{1}{3}}}{b}}}{x}\right) + 10a^2(-b)^{\frac{2}{3}}\log\left(\frac{(-b)^{\frac{1}{3}}x + (bx^3 + a)^{\frac{1}{3}}}{x}\right) - 5a^2}{54b}$$

input `integrate((b*x^3+a)^(5/3),x, algorithm="fricas")`

output `[1/54*(15*sqrt(1/3)*a^2*b*sqrt((-b)^(1/3)/b)*log(3*b*x^3 - 3*(b*x^3 + a)^(1/3)*(-b)^(2/3)*x^2 - 3*sqrt(1/3)*((-b)^(1/3)*b*x^3 - (b*x^3 + a)^(1/3)*b*x^2 + 2*(b*x^3 + a)^(2/3)*(-b)^(2/3)*x)*sqrt((-b)^(1/3)/b) + 2*a) - 10*a^2*(-b)^(2/3)*log(((b)^(1/3)*x + (b*x^3 + a)^(1/3))/x) + 5*a^2*(-b)^(2/3)*log(((b)^(2/3)*x^2 - (b*x^3 + a)^(1/3)*(-b)^(1/3)*x + (b*x^3 + a)^(2/3))/x^2) + 3*(3*b^2*x^4 + 8*a*b*x)*(b*x^3 + a)^(2/3))/b, -1/54*(30*sqrt(1/3)*a^2*b*sqrt(-(-b)^(1/3)/b)*arctan(-sqrt(1/3)*((-b)^(1/3)*x - 2*(b*x^3 + a)^(1/3))*sqrt(-(-b)^(1/3)/b)/x) + 10*a^2*(-b)^(2/3)*log(((b)^(1/3)*x + (b*x^3 + a)^(1/3))/x) - 5*a^2*(-b)^(2/3)*log(((b)^(2/3)*x^2 - (b*x^3 + a)^(1/3)*(-b)^(1/3)*x + (b*x^3 + a)^(2/3))/x^2) - 3*(3*b^2*x^4 + 8*a*b*x)*(b*x^3 + a)^(2/3))/b]`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.97 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.33

$$\int (a + bx^3)^{5/3} dx = \frac{a^{5/3} x \Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\begin{matrix} -\frac{5}{3}, \frac{1}{3} \\ \frac{4}{3} \end{matrix} \middle| \frac{bx^3 e^{i\pi}}{a} \right)}{3\Gamma\left(\frac{4}{3}\right)}$$

input `integrate((b*x**3+a)**(5/3),x)`

output `a**(5/3)*x*gamma(1/3)*hyper((-5/3, 1/3), (4/3,), b*x**3*exp_polar(I*pi)/a)/(3*gamma(4/3))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 179 vs. $2(85) = 170$.

Time = 0.12 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.60

$$\int (a + bx^3)^{5/3} dx = -\frac{5\sqrt{3}a^2 \arctan\left(\frac{\sqrt{3}\left(b^{1/3} + \frac{2(bx^3+a)^{1/3}}{x}\right)}{3b^{1/3}}\right)}{27b^{1/3}} + \frac{5a^2 \log\left(b^{2/3} + \frac{(bx^3+a)^{1/3}b^{1/3}}{x} + \frac{(bx^3+a)^{2/3}}{x^2}\right)}{54b^{1/3}} - \frac{5a^2 \log\left(-b^{1/3} + \frac{(bx^3+a)^{1/3}}{x}\right)}{27b^{1/3}} - \frac{\frac{5(bx^3+a)^{2/3}a^2b}{x^2} - \frac{8(bx^3+a)^{5/3}a^2}{x^5}}{18\left(b^2 - \frac{2(bx^3+a)b}{x^3} + \frac{(bx^3+a)^2}{x^6}\right)}$$

input `integrate((b*x^3+a)^(5/3),x, algorithm="maxima")`

output `-5/27*sqrt(3)*a^2*arctan(1/3*sqrt(3)*(b^(1/3) + 2*(b*x^3 + a)^(1/3)/x)/b^(1/3))/b^(1/3) + 5/54*a^2*log(b^(2/3) + (b*x^3 + a)^(1/3)*b^(1/3)/x + (b*x^3 + a)^(2/3)/x^2)/b^(1/3) - 5/27*a^2*log(-b^(1/3) + (b*x^3 + a)^(1/3)/x)/b^(1/3) - 1/18*(5*(b*x^3 + a)^(2/3)*a^2*b/x^2 - 8*(b*x^3 + a)^(5/3)*a^2/x^5)/(b^2 - 2*(b*x^3 + a)*b/x^3 + (b*x^3 + a)^2/x^6)`

Giac [F]

$$\int (a + bx^3)^{5/3} dx = \int (bx^3 + a)^{5/3} dx$$

input `integrate((b*x^3+a)^(5/3),x, algorithm="giac")`

output `integrate((b*x^3 + a)^(5/3), x)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.33

$$\int (a + bx^3)^{5/3} dx = \frac{x(bx^3 + a)^{5/3} {}_2F_1\left(-\frac{5}{3}, \frac{1}{3}; \frac{4}{3}; -\frac{bx^3}{a}\right)}{\left(\frac{bx^3}{a} + 1\right)^{5/3}}$$

input `int((a + b*x^3)^(5/3),x)`output `(x*(a + b*x^3)^(5/3)*hypergeom([-5/3, 1/3], 4/3, -(b*x^3)/a))/((b*x^3)/a + 1)^(5/3)`**Reduce [F]**

$$\int (a + bx^3)^{5/3} dx = \frac{4(bx^3 + a)^{2/3} ax}{9} + \frac{(bx^3 + a)^{2/3} bx^4}{6} + \frac{5\left(\int \frac{1}{(bx^3+a)^{1/3}} dx\right) a^2}{9}$$

input `int((b*x^3+a)^(5/3),x)`output `(8*(a + b*x**3)**(2/3)*a*x + 3*(a + b*x**3)**(2/3)*b*x**4 + 10*int((a + b*x**3)**(2/3)/(a + b*x**3),x)*a**2)/18`

3.25 $\int (a + bx^3)^{2/3} dx$

Optimal result	211
Mathematica [C] (warning: unable to verify)	211
Rubi [A] (verified)	212
Maple [A] (verified)	213
Fricas [B] (verification not implemented)	214
Sympy [C] (verification not implemented)	215
Maxima [A] (verification not implemented)	216
Giac [F]	216
Mupad [B] (verification not implemented)	217
Reduce [F]	217

Optimal result

Integrand size = 11, antiderivative size = 91

$$\int (a + bx^3)^{2/3} dx = \frac{1}{3}x(a + bx^3)^{2/3} + \frac{2a \arctan\left(\frac{1 + \frac{2\sqrt[3]{b}x}{\sqrt[3]{a + bx^3}}}{\sqrt{3}}\right) - \frac{a \log\left(-\sqrt[3]{b}x + \sqrt[3]{a + bx^3}\right)}{3\sqrt[3]{b}}}{3\sqrt{3}\sqrt[3]{b}}$$

output

```
1/3*x*(b*x^3+a)^(2/3)+2/9*a*arctan(1/3*(1+2*b^(1/3)*x/(b*x^3+a)^(1/3))*3^(1/2))*3^(1/2)/b^(1/3)-1/3*a*ln(-b^(1/3)*x+(b*x^3+a)^(1/3))/b^(1/3)
```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 0.09 (sec) , antiderivative size = 203, normalized size of antiderivative = 2.23

$$\int (a + bx^3)^{2/3} dx = \frac{3 \left((-1)^{2/3} \sqrt[3]{a} + \sqrt[3]{bx} \right) (a + bx^3)^{2/3} \operatorname{AppellF1} \left(\frac{5}{3}, -\frac{2}{3}, -\frac{2}{3}, \frac{8}{3}, -\frac{(-1)^{2/3} \left((-1)^{2/3} \sqrt[3]{a} + \sqrt[3]{bx} \right)}{(1 + \sqrt[3]{-1}) \sqrt[3]{a}}, \frac{i + \sqrt{3}}{3} \right)}{5 \cdot 2^{2/3} \sqrt[3]{b} \left(\frac{\sqrt[3]{a} + (-1)^{2/3} \sqrt[3]{bx}}{(1 + \sqrt[3]{-1}) \sqrt[3]{a}} \right)^{2/3} \left(\frac{i \left(1 + \frac{\sqrt[3]{bx}}{\sqrt[3]{a}} \right)}{3i + \sqrt{3}} \right)^{2/3}}$$

input `Integrate[(a + b*x^3)^(2/3),x]`

output `(3*((-1)^(2/3)*a^(1/3) + b^(1/3)*x)*(a + b*x^3)^(2/3)*AppellF1[5/3, -2/3, -2/3, 8/3, -(((-1)^(2/3)*((-1)^(2/3)*a^(1/3) + b^(1/3)*x))/((1 + (-1)^(1/3))*a^(1/3))], (I + Sqrt[3] - ((2*I)*b^(1/3)*x)/a^(1/3))/(3*I + Sqrt[3]))/(5*2^(2/3)*b^(1/3)*((a^(1/3) + (-1)^(2/3)*b^(1/3)*x)/((1 + (-1)^(1/3))*a^(1/3)))^(2/3)*((I*(1 + (b^(1/3)*x)/a^(1/3)))/(3*I + Sqrt[3]))^(2/3)`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.01, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {748, 769}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx^3)^{2/3} dx$$

↓ 748

$$\frac{2}{3}a \int \frac{1}{\sqrt[3]{bx^3 + a}} dx + \frac{1}{3}x(a + bx^3)^{2/3}$$

↓ 769

$$\frac{2}{3}a \left(\frac{\arctan\left(\frac{\frac{2\sqrt[3]{bx}}{\sqrt[3]{a+bx^3}}+1}{\sqrt{3}}\right)}{\sqrt{3}\sqrt[3]{b}} - \frac{\log\left(\frac{\sqrt[3]{a+bx^3}-\sqrt[3]{bx}}{2\sqrt[3]{b}}\right)}{2\sqrt[3]{b}} \right) + \frac{1}{3}x(a+bx^3)^{2/3}$$

input `Int[(a + b*x^3)^(2/3), x]`

output `(x*(a + b*x^3)^(2/3))/3 + (2*a*(ArcTan[(1 + (2*b^(1/3)*x)/(a + b*x^3)^(1/3))]/Sqrt[3])/(Sqrt[3]*b^(1/3)) - Log[-(b^(1/3)*x) + (a + b*x^3)^(1/3)]/(2*b^(1/3)))/3`

Defintions of rubi rules used

rule 748 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p + 1)), x] + Simp[a*n*(p/(n*p + 1)) Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || LtQ[Denominator[p + 1/n], Denominator[p]])`

rule 769 `Int[((a_) + (b_.)*(x_)^3)^(-1/3), x_Symbol] := Simp[ArcTan[(1 + 2*Rt[b, 3]*(x/(a + b*x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*Rt[b, 3]), x] - Simp[Log[(a + b*x^3)^(1/3) - Rt[b, 3]*x]/(2*Rt[b, 3]), x] /; FreeQ[{a, b}, x]`

Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.30

method	result
pseudoelliptic	$\frac{2 \left(\sqrt{3} \arctan\left(\frac{\sqrt{3} \left(b^{\frac{1}{3}} x + 2(b x^3 + a)^{\frac{1}{3}} \right)}{3 b^{\frac{1}{3}} x}\right) a^{-\frac{3(b x^3 + a)^{\frac{2}{3}} x b^{\frac{1}{3}}}{2}} + \ln\left(\frac{-b^{\frac{1}{3}} x + (b x^3 + a)^{\frac{1}{3}}}{x}\right) a^{-\frac{\ln\left(\frac{b^{\frac{2}{3}} x^2 + b^{\frac{1}{3}}(b x^3 + a)^{\frac{1}{3}} x + (b x^3 + a)^{\frac{1}{3}}}{x^2}}{2}} \right)}{9 b^{\frac{1}{3}}}$

input `int((b*x^3+a)^(2/3),x,method=_RETURNVERBOSE)`

output `-2/9*(3^(1/2)*arctan(1/3*3^(1/2)*(b^(1/3)*x+2*(b*x^3+a)^(1/3))/b^(1/3)/x)*
a-3/2*(b*x^3+a)^(2/3)*x*b^(1/3)+ln((-b^(1/3)*x+(b*x^3+a)^(1/3))/x)*a-1/2*ln((b^(2/3)*x^2+b^(1/3)*(b*x^3+a)^(1/3)*x+(b*x^3+a)^(2/3))/x^2)*a)/b^(1/3)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 157 vs. 2(68) = 136.

Time = 0.08 (sec) , antiderivative size = 362, normalized size of antiderivative = 3.98

$$\int (a + bx^3)^{2/3} dx = \frac{3 \sqrt{\frac{1}{3}} ab \sqrt{\frac{(-b)^{\frac{1}{3}}}{b}} \log \left(3bx^3 - 3(bx^3 + a)^{\frac{1}{3}}(-b)^{\frac{2}{3}}x^2 - 3 \sqrt{\frac{1}{3}} \left((-b)^{\frac{1}{3}}bx^3 - (bx^3 + a)^{\frac{1}{3}}bx^2 + 2a(-b)^{\frac{1}{3}} \right) \right)}{9b} + \frac{6 \sqrt{\frac{1}{3}} ab \sqrt{-\frac{(-b)^{\frac{1}{3}}}{b}} \arctan \left(-\frac{\sqrt{\frac{1}{3}} \left((-b)^{\frac{1}{3}}x - 2(bx^3 + a)^{\frac{1}{3}} \right) \sqrt{-\frac{(-b)^{\frac{1}{3}}}{b}}}{x} \right) - 3(bx^3 + a)^{\frac{2}{3}}bx + 2a(-b)^{\frac{2}{3}} \log \left(\frac{(-b)^{\frac{1}{3}}x + (bx^3 + a)^{\frac{1}{3}}}{x} \right)}{9b}$$

input `integrate((b*x^3+a)^(2/3),x, algorithm="fricas")`

output

```
[1/9*(3*sqrt(1/3)*a*b*sqrt((-b)^(1/3)/b)*log(3*b*x^3 - 3*(b*x^3 + a)^(1/3)
*(-b)^(2/3)*x^2 - 3*sqrt(1/3)*((-b)^(1/3)*b*x^3 - (b*x^3 + a)^(1/3)*b*x^2
+ 2*(b*x^3 + a)^(2/3)*(-b)^(2/3)*x)*sqrt((-b)^(1/3)/b) + 2*a) + 3*(b*x^3 +
a)^(2/3)*b*x - 2*a*(-b)^(2/3)*log(((b)^(1/3)*x + (b*x^3 + a)^(1/3))/x) +
a*(-b)^(2/3)*log(((b)^(2/3)*x^2 - (b*x^3 + a)^(1/3)*(-b)^(1/3)*x + (b*x^
3 + a)^(2/3))/x^2))/b, -1/9*(6*sqrt(1/3)*a*b*sqrt(-(-b)^(1/3)/b)*arctan(-s
qrt(1/3)*((-b)^(1/3)*x - 2*(b*x^3 + a)^(1/3))*sqrt(-(-b)^(1/3)/b)/x) - 3*(
b*x^3 + a)^(2/3)*b*x + 2*a*(-b)^(2/3)*log(((b)^(1/3)*x + (b*x^3 + a)^(1/3)
))/x) - a*(-b)^(2/3)*log(((b)^(2/3)*x^2 - (b*x^3 + a)^(1/3)*(-b)^(1/3)*x
+ (b*x^3 + a)^(2/3))/x^2))/b]
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.66 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.41

$$\int (a + bx^3)^{2/3} dx = \frac{a^{2/3} x \Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{1}{3} \\ \frac{4}{3} \end{matrix} \middle| \frac{bx^3 e^{i\pi}}{a} \right)}{3\Gamma\left(\frac{4}{3}\right)}$$

input

```
integrate((b*x**3+a)**(2/3),x)
```

output

```
a**(2/3)*x*gamma(1/3)*hyper((-2/3, 1/3), (4/3,), b*x**3*exp_polar(I*pi)/a)
/(3*gamma(4/3))
```


Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.47

$$\int (a + bx^3)^{2/3} dx = -\frac{2\sqrt{3}a \arctan\left(\frac{\sqrt{3}\left(b^{1/3} + \frac{2(bx^3+a)^{1/3}}{x}\right)}{3b^{1/3}}\right)}{9b^{1/3}} + \frac{a \log\left(b^{2/3} + \frac{(bx^3+a)^{1/3}b^{1/3}}{x} + \frac{(bx^3+a)^{2/3}}{x^2}\right)}{9b^{1/3}} - \frac{2a \log\left(-b^{1/3} + \frac{(bx^3+a)^{1/3}}{x}\right)}{9b^{1/3}} - \frac{(bx^3+a)^{2/3}a}{3\left(b - \frac{bx^3+a}{x^3}\right)x^2}$$

input `integrate((b*x^3+a)^(2/3),x, algorithm="maxima")`output `-2/9*sqrt(3)*a*arctan(1/3*sqrt(3)*(b^(1/3) + 2*(b*x^3 + a)^(1/3)/x)/b^(1/3))/
b^(1/3) + 1/9*a*log(b^(2/3) + (b*x^3 + a)^(1/3)*b^(1/3)/x + (b*x^3 + a)^(2/3)/x^2)/b^(1/3) - 2/9*a*log(-b^(1/3) + (b*x^3 + a)^(1/3)/x)/b^(1/3) - 1/3*(b*x^3 + a)^(2/3)*a/((b - (b*x^3 + a)/x^3)*x^2)`**Giac [F]**

$$\int (a + bx^3)^{2/3} dx = \int (bx^3 + a)^{2/3} dx$$

input `integrate((b*x^3+a)^(2/3),x, algorithm="giac")`output `integrate((b*x^3 + a)^(2/3), x)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.41

$$\int (a + bx^3)^{2/3} dx = \frac{x (bx^3 + a)^{2/3} {}_2F_1\left(-\frac{2}{3}, \frac{1}{3}; \frac{4}{3}; -\frac{bx^3}{a}\right)}{\left(\frac{bx^3}{a} + 1\right)^{2/3}}$$

input `int((a + b*x^3)^(2/3),x)`output `(x*(a + b*x^3)^(2/3)*hypergeom([-2/3, 1/3], 4/3, -(b*x^3)/a))/((b*x^3)/a + 1)^(2/3)`**Reduce [F]**

$$\int (a + bx^3)^{2/3} dx = \frac{(bx^3 + a)^{2/3} x}{3} + \frac{2 \left(\int \frac{1}{(bx^3 + a)^{1/3}} dx \right) a}{3}$$

input `int((b*x^3+a)^(2/3),x)`output `((a + b*x**3)**(2/3)*x + 2*int((a + b*x**3)**(2/3)/(a + b*x**3),x)*a)/3`

3.26 $\int \frac{1}{\sqrt[3]{a + bx^3}} dx$

Optimal result	218
Mathematica [A] (verified)	218
Rubi [A] (verified)	219
Maple [A] (verified)	220
Fricas [B] (verification not implemented)	220
Sympy [C] (verification not implemented)	221
Maxima [A] (verification not implemented)	222
Giac [F]	222
Mupad [B] (verification not implemented)	223
Reduce [F]	223

Optimal result

Integrand size = 11, antiderivative size = 70

$$\int \frac{1}{\sqrt[3]{a + bx^3}} dx = \frac{\arctan\left(\frac{1 + \frac{2\sqrt[3]{bx}}{\sqrt[3]{a + bx^3}}}{\sqrt{3}}\right)}{\sqrt{3}\sqrt[3]{b}} - \frac{\log\left(-\sqrt[3]{bx} + \sqrt[3]{a + bx^3}\right)}{2\sqrt[3]{b}}$$

output

```
1/3*arctan(1/3*(1+2*b^(1/3)*x/(b*x^3+a)^(1/3))*3^(1/2))*3^(1/2)/b^(1/3)-1/2*ln(-b^(1/3)*x+(b*x^3+a)^(1/3))/b^(1/3)
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.57

$$\int \frac{1}{\sqrt[3]{a + bx^3}} dx = \frac{2\sqrt{3} \arctan\left(\frac{1 + \frac{2\sqrt[3]{bx}}{\sqrt[3]{a + bx^3}}}{\sqrt{3}}\right) - 2 \log\left(1 - \frac{\sqrt[3]{bx}}{\sqrt[3]{a + bx^3}}\right) + \log\left(1 + \frac{b^{2/3}x^2}{(a + bx^3)^{2/3}} + \frac{\sqrt[3]{bx}}{\sqrt[3]{a + bx^3}}\right)}{6\sqrt[3]{b}}$$

input `Integrate[(a + b*x^3)^(-1/3),x]`

output `(2*sqrt(3)*ArcTan[(1 + (2*b^(1/3)*x)/(a + b*x^3)^(1/3))/sqrt(3)] - 2*Log[1 - (b^(1/3)*x)/(a + b*x^3)^(1/3)] + Log[1 + (b^(2/3)*x^2)/(a + b*x^3)^(2/3) + (b^(1/3)*x)/(a + b*x^3)^(1/3)])/(6*b^(1/3))`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {769}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt[3]{a + bx^3}} dx$$

↓ 769

$$\frac{\arctan\left(\frac{\frac{2\sqrt[3]{bx}}{\sqrt[3]{a + bx^3}} + 1}{\sqrt{3}}\right)}{\sqrt{3}\sqrt[3]{b}} - \frac{\log\left(\sqrt[3]{a + bx^3} - \sqrt[3]{bx}\right)}{2\sqrt[3]{b}}$$

input `Int[(a + b*x^3)^(-1/3),x]`

output `ArcTan[(1 + (2*b^(1/3)*x)/(a + b*x^3)^(1/3))/sqrt(3)]/(sqrt(3)*b^(1/3)) - Log[-(b^(1/3)*x) + (a + b*x^3)^(1/3)]/(2*b^(1/3))`

Defintions of rubi rules used

rule 769

```
Int[((a_) + (b_.)*(x_)^3)^(-1/3), x_Symbol] :> Simp[ArcTan[(1 + 2*Rt[b, 3]*
(x/(a + b*x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*Rt[b, 3]), x] - Simp[Log[(a + b*x^
3)^(1/3) - Rt[b, 3]*x]/(2*Rt[b, 3]), x] /; FreeQ[{a, b}, x]
```

Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.41

method	result	size
pseudoelliptic	$\frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(b^{\frac{1}{3}}x + 2(bx^3+a)^{\frac{1}{3}}\right)}{3b^{\frac{1}{3}}x}\right) + \ln\left(\frac{-b^{\frac{1}{3}}x + (bx^3+a)^{\frac{1}{3}}}{x}\right) - \frac{\ln\left(\frac{b^{\frac{2}{3}}x^2 + b^{\frac{1}{3}}(bx^3+a)^{\frac{1}{3}}x + (bx^3+a)^{\frac{2}{3}}}{x^2}\right)}{2}}{3b^{\frac{1}{3}}}$	99

input

```
int(1/(b*x^3+a)^(1/3),x,method=_RETURNVERBOSE)
```

output

```
-1/3*(3^(1/2)*arctan(1/3*3^(1/2)*(b^(1/3)*x+2*(b*x^3+a)^(1/3))/b^(1/3)/x)+
ln((-b^(1/3)*x+(b*x^3+a)^(1/3))/x)-1/2*ln((b^(2/3)*x^2+b^(1/3)*(b*x^3+a)^(
1/3)*x+(b*x^3+a)^(2/3))/x^2))/b^(1/3)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 141 vs. 2(54) = 108.

Time = 0.08 (sec) , antiderivative size = 330, normalized size of antiderivative = 4.71

$$\int \frac{1}{\sqrt[3]{a+bx^3}} dx$$

$$= \frac{3 \sqrt{\frac{1}{3}} b \sqrt{\frac{(-b)^{\frac{1}{3}}}{b}} \log \left(3bx^3 - 3(bx^3+a)^{\frac{1}{3}}(-b)^{\frac{2}{3}}x^2 - 3\sqrt{\frac{1}{3}} \left((-b)^{\frac{1}{3}}bx^3 - (bx^3+a)^{\frac{1}{3}}bx^2 + 2(bx^3+a)^{\frac{2}{3}}(-b)^{\frac{1}{3}} \right) \right)}{6} - \frac{6 \sqrt{\frac{1}{3}} b \sqrt{-\frac{(-b)^{\frac{1}{3}}}{b}} \arctan \left(-\frac{\sqrt{\frac{1}{3}} \left((-b)^{\frac{1}{3}}x - 2(bx^3+a)^{\frac{1}{3}} \right) \sqrt{-\frac{(-b)^{\frac{1}{3}}}{b}}}{x} \right) + 2(-b)^{\frac{2}{3}} \log \left(\frac{(-b)^{\frac{1}{3}}x + (bx^3+a)^{\frac{1}{3}}}{x} \right) - (-b)^{\frac{2}{3}}}{6b}$$

input `integrate(1/(b*x^3+a)^(1/3),x, algorithm="fricas")`

output `[1/6*(3*sqrt(1/3)*b*sqrt((-b)^(1/3)/b)*log(3*b*x^3 - 3*(b*x^3 + a)^(1/3)*(-b)^(2/3)*x^2 - 3*sqrt(1/3)*((-b)^(1/3)*b*x^3 - (b*x^3 + a)^(1/3)*b*x^2 + 2*(b*x^3 + a)^(2/3)*(-b)^(2/3)*x)*sqrt((-b)^(1/3)/b) + 2*a - 2*(-b)^(2/3)*log(((b)^(1/3)*x + (b*x^3 + a)^(1/3))/x) + (-b)^(2/3)*log(((b)^(2/3)*x^2 - (b*x^3 + a)^(1/3)*(-b)^(1/3)*x + (b*x^3 + a)^(2/3))/x^2))/b, -1/6*(6*sqrt(1/3)*b*sqrt(-(-b)^(1/3)/b)*arctan(-sqrt(1/3)*((-b)^(1/3)*x - 2*(b*x^3 + a)^(1/3))*sqrt(-(-b)^(1/3)/b)/x) + 2*(-b)^(2/3)*log(((b)^(1/3)*x + (b*x^3 + a)^(1/3))/x) - (-b)^(2/3)*log(((b)^(2/3)*x^2 - (b*x^3 + a)^(1/3)*(-b)^(1/3)*x + (b*x^3 + a)^(2/3))/x^2))/b]`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.56 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.51

$$\int \frac{1}{\sqrt[3]{a+bx^3}} dx = \frac{x\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{1}{3} \middle| \frac{bx^3 e^{i\pi}}{a}\right)}{3\sqrt[3]{a}\Gamma\left(\frac{4}{3}\right)}$$

input `integrate(1/(b*x**3+a)**(1/3),x)`

output `x*gamma(1/3)*hyper((1/3, 1/3), (4/3,), b*x**3*exp_polar(I*pi)/a)/(3*a**(1/3)*gamma(4/3))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.43

$$\int \frac{1}{\sqrt[3]{a+bx^3}} dx = -\frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(b^{\frac{1}{3}} + \frac{2(bx^3+a)^{\frac{1}{3}}}{x}\right)}{3b^{\frac{1}{3}}}\right)}{3b^{\frac{1}{3}}} + \frac{\log\left(b^{\frac{2}{3}} + \frac{(bx^3+a)^{\frac{1}{3}}b^{\frac{1}{3}}}{x} + \frac{(bx^3+a)^{\frac{2}{3}}}{x^2}\right)}{6b^{\frac{1}{3}}} - \frac{\log\left(-b^{\frac{1}{3}} + \frac{(bx^3+a)^{\frac{1}{3}}}{x}\right)}{3b^{\frac{1}{3}}}$$

input `integrate(1/(b*x^3+a)^(1/3),x, algorithm="maxima")`

output `-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(b^(1/3) + 2*(b*x^3 + a)^(1/3)/x)/b^(1/3)) /b^(1/3) + 1/6*log(b^(2/3) + (b*x^3 + a)^(1/3)*b^(1/3)/x + (b*x^3 + a)^(2/3)/x^2)/b^(1/3) - 1/3*log(-b^(1/3) + (b*x^3 + a)^(1/3)/x)/b^(1/3)`

Giac [F]

$$\int \frac{1}{\sqrt[3]{a+bx^3}} dx = \int \frac{1}{(bx^3+a)^{\frac{1}{3}}} dx$$

input `integrate(1/(b*x^3+a)^(1/3),x, algorithm="giac")`

output `integrate((b*x^3 + a)^(-1/3), x)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.53

$$\int \frac{1}{\sqrt[3]{a + bx^3}} dx = \frac{x \left(\frac{bx^3}{a} + 1 \right)^{1/3} {}_2F_1 \left(\frac{1}{3}, \frac{1}{3}; \frac{4}{3}; -\frac{bx^3}{a} \right)}{(bx^3 + a)^{1/3}}$$

input `int(1/(a + b*x^3)^(1/3),x)`output `(x*((b*x^3)/a + 1)^(1/3)*hypergeom([1/3, 1/3], 4/3, -(b*x^3)/a))/(a + b*x^3)^(1/3)`**Reduce [F]**

$$\int \frac{1}{\sqrt[3]{a + bx^3}} dx = \int \frac{1}{(bx^3 + a)^{1/3}} dx$$

input `int(1/(b*x^3+a)^(1/3),x)`output `int(1/(a + b*x**3)**(1/3),x)`

3.27 $\int \frac{1}{(a+bx^3)^{4/3}} dx$

Optimal result	224
Mathematica [C] (warning: unable to verify)	224
Rubi [A] (verified)	225
Maple [A] (verified)	226
Fricas [A] (verification not implemented)	226
Sympy [B] (verification not implemented)	227
Maxima [A] (verification not implemented)	227
Giac [F]	227
Mupad [B] (verification not implemented)	228
Reduce [F]	228

Optimal result

Integrand size = 11, antiderivative size = 16

$$\int \frac{1}{(a + bx^3)^{4/3}} dx = \frac{x}{a\sqrt[3]{a + bx^3}}$$

output

```
x/a/(b*x^3+a)^(1/3)
```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 2 in optimal.

Time = 3.92 (sec) , antiderivative size = 524, normalized size of antiderivative = 32.75

$$\int \frac{1}{(a + bx^3)^{4/3}} dx = \frac{i\sqrt{\frac{\pi}{3}} \left(\frac{(-1)^{2/3} \sqrt[3]{a}}{\sqrt[3]{b}} + x \right) \left(1 + \frac{\sqrt[3]{bx}}{\sqrt[3]{a}} \right) \text{Gamma} \left(\frac{1}{3} \right) \left(4 \left(2(-3i + \sqrt{3}) a - (9i + \sqrt{3}) a^{2/3} \right)}{\dots} \right)}{\dots}$$

input

```
Integrate[(a + b*x^3)^(-4/3), x]
```

output

```
((I/20)*Sqrt[Pi/3]*((( -1)^(2/3)*a^(1/3))/b^(1/3) + x)*(1 + (b^(1/3)*x)/a^(1/3))*Gamma[1/3]*(4*(2*(-3*I + Sqrt[3])*a - (9*I + Sqrt[3])*a^(2/3)*b^(1/3)*x + 6*(-I + Sqrt[3])*a^(1/3)*b^(2/3)*x^2 + 6*Sqrt[3]*b*x^3)*Hypergeometric2F1[1, 4/3, 8/3, (2*Sqrt[3]*a^(1/3) - (3*I + Sqrt[3])*b^(1/3)*x)/((3*I + Sqrt[3])*a^(1/3) - 2*Sqrt[3]*b^(1/3)*x)] - 3*(a^(1/3) + b^(1/3)*x)*((2*I + 4*Sqrt[3])*a^(2/3) + (-7*I + 5*Sqrt[3])*a^(1/3)*b^(1/3)*x - 3*(3*I + Sqrt[3])*b^(2/3)*x^2)*Hypergeometric2F1[2, 7/3, 11/3, (2*Sqrt[3]*a^(1/3) - (3*I + Sqrt[3])*b^(1/3)*x)/((3*I + Sqrt[3])*a^(1/3) - 2*Sqrt[3]*b^(1/3)*x)] - 3*(a^(1/3) + b^(1/3)*x)^2*(2*Sqrt[3]*a^(1/3) - (3*I + Sqrt[3])*b^(1/3)*x)*HypergeometricPFQ[{2, 2, 7/3}, {1, 11/3}, (2*Sqrt[3]*a^(1/3) - (3*I + Sqrt[3])*b^(1/3)*x)/((3*I + Sqrt[3])*a^(1/3) + (-1)^(2/3)*b^(1/3)*x)]))/(2^(1/3)*(3*I + Sqrt[3])*a^(2/3)*(a^(1/3) + (-1)^(2/3)*b^(1/3)*x)*(a + b*x^3)^(4/3)*Gamma[2/3]*Gamma[7/6])
```

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {746}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + bx^3)^{4/3}} dx$$

$$\downarrow 746$$

$$\frac{x}{a^3 \sqrt[3]{a + bx^3}}$$

input

```
Int[(a + b*x^3)^(-4/3), x]
```

output

```
x/(a*(a + b*x^3)^(1/3))
```

Defintions of rubi rules used

rule 746 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[x*((a + b*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]`

Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result	size
gosper	$\frac{x}{a(bx^3+a)^{\frac{1}{3}}}$	15
trager	$\frac{x}{a(bx^3+a)^{\frac{1}{3}}}$	15
pseudoelliptic	$\frac{x}{a(bx^3+a)^{\frac{1}{3}}}$	15
orering	$\frac{x}{a(bx^3+a)^{\frac{1}{3}}}$	15

input `int(1/(b*x^3+a)^(4/3),x,method=_RETURNVERBOSE)`

output `x/a/(b*x^3+a)^(1/3)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.44

$$\int \frac{1}{(a + bx^3)^{4/3}} dx = \frac{(bx^3 + a)^{\frac{2}{3}} x}{abx^3 + a^2}$$

input `integrate(1/(b*x^3+a)^(4/3),x, algorithm="fricas")`

output `(b*x^3 + a)^(2/3)*x/(a*b*x^3 + a^2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. $2(12) = 24$.

Time = 0.38 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.81

$$\int \frac{1}{(a + bx^3)^{4/3}} dx = \frac{x\Gamma(\frac{1}{3})}{3a^{\frac{4}{3}}\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma(\frac{4}{3})}$$

input `integrate(1/(b*x**3+a)**(4/3),x)`

output `x*gamma(1/3)/(3*a**(4/3)*(1 + b*x**3/a)**(1/3)*gamma(4/3))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1}{(a + bx^3)^{4/3}} dx = \frac{x}{(bx^3 + a)^{\frac{1}{3}}a}$$

input `integrate(1/(b*x^3+a)^(4/3),x, algorithm="maxima")`

output `x/((b*x^3 + a)^(1/3)*a)`

Giac [F]

$$\int \frac{1}{(a + bx^3)^{4/3}} dx = \int \frac{1}{(bx^3 + a)^{\frac{4}{3}}} dx$$

input `integrate(1/(b*x^3+a)^(4/3),x, algorithm="giac")`

output `integrate((b*x^3 + a)^(-4/3), x)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1}{(a + bx^3)^{4/3}} dx = \frac{x}{a(bx^3 + a)^{1/3}}$$

input `int(1/(a + b*x^3)^(4/3),x)`output `x/(a*(a + b*x^3)^(1/3))`**Reduce [F]**

$$\int \frac{1}{(a + bx^3)^{4/3}} dx = \int \frac{1}{(bx^3 + a)^{\frac{1}{3}} a + (bx^3 + a)^{\frac{1}{3}} bx^3} dx$$

input `int(1/(b*x^3+a)^(4/3),x)`output `int(1/((a + b*x**3)**(1/3)*a + (a + b*x**3)**(1/3)*b*x**3),x)`

3.28 $\int \frac{1}{(a+bx^3)^{7/3}} dx$

Optimal result	229
Mathematica [C] (warning: unable to verify)	229
Rubi [A] (verified)	230
Maple [A] (verified)	231
Fricas [A] (verification not implemented)	231
Sympy [B] (verification not implemented)	232
Maxima [A] (verification not implemented)	232
Giac [F]	233
Mupad [B] (verification not implemented)	233
Reduce [F]	233

Optimal result

Integrand size = 11, antiderivative size = 39

$$\int \frac{1}{(a + bx^3)^{7/3}} dx = \frac{x}{4a(a + bx^3)^{4/3}} + \frac{3x}{4a^2\sqrt[3]{a + bx^3}}$$

output `1/4*x/a/(b*x^3+a)^(4/3)+3/4*x/a^2/(b*x^3+a)^(1/3)`

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 2 in optimal.

Time = 16.91 (sec) , antiderivative size = 5268, normalized size of antiderivative = 135.08

$$\int \frac{1}{(a + bx^3)^{7/3}} dx = \text{Result too large to show}$$

input `Integrate[(a + b*x^3)^(-7/3),x]`

output `Result too large to show`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {749, 746}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + bx^3)^{7/3}} dx$$

$$\downarrow 749$$

$$\frac{3 \int \frac{1}{(bx^3+a)^{4/3}} dx}{4a} + \frac{x}{4a (a + bx^3)^{4/3}}$$

$$\downarrow 746$$

$$\frac{3x}{4a^2 \sqrt[3]{a + bx^3}} + \frac{x}{4a (a + bx^3)^{4/3}}$$

input `Int[(a + b*x^3)^(-7/3), x]`

output `x/(4*a*(a + b*x^3)^(4/3)) + (3*x)/(4*a^2*(a + b*x^3)^(1/3))`

Defintions of rubi rules used

rule 746 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]`

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.67

method	result	size
gospers	$\frac{x(3bx^3+4a)}{4(bx^3+a)^{\frac{4}{3}}a^2}$	26
trager	$\frac{x(3bx^3+4a)}{4(bx^3+a)^{\frac{4}{3}}a^2}$	26
pseudoelliptic	$\frac{x(3bx^3+4a)}{4(bx^3+a)^{\frac{4}{3}}a^2}$	26
orering	$\frac{x(3bx^3+4a)}{4(bx^3+a)^{\frac{4}{3}}a^2}$	26

input `int(1/(b*x^3+a)^(7/3),x,method=_RETURNVERBOSE)`

output `1/4*x*(3*b*x^3+4*a)/(b*x^3+a)^(4/3)/a^2`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.21

$$\int \frac{1}{(a+bx^3)^{7/3}} dx = \frac{(3bx^4+4ax)(bx^3+a)^{\frac{2}{3}}}{4(a^2b^2x^6+2a^3bx^3+a^4)}$$

input `integrate(1/(b*x^3+a)^(7/3),x, algorithm="fricas")`

output `1/4*(3*b*x^4 + 4*a*x)*(b*x^3 + a)^(2/3)/(a^2*b^2*x^6 + 2*a^3*b*x^3 + a^4)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 126 vs. $2(32) = 64$.

Time = 0.59 (sec) , antiderivative size = 126, normalized size of antiderivative = 3.23

$$\int \frac{1}{(a + bx^3)^{7/3}} dx = \frac{4ax\Gamma(\frac{1}{3})}{9a^{\frac{10}{3}}\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma(\frac{7}{3}) + 9a^{\frac{7}{3}}bx^3\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma(\frac{7}{3})} + \frac{3bx^4\Gamma(\frac{1}{3})}{9a^{\frac{10}{3}}\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma(\frac{7}{3}) + 9a^{\frac{7}{3}}bx^3\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma(\frac{7}{3})}$$

input `integrate(1/(b*x**3+a)**(7/3),x)`

output `4*a*x*gamma(1/3)/(9*a**(10/3)*(1 + b*x**3/a)**(1/3)*gamma(7/3) + 9*a**(7/3)*b*x**3*(1 + b*x**3/a)**(1/3)*gamma(7/3)) + 3*b*x**4*gamma(1/3)/(9*a**(10/3)*(1 + b*x**3/a)**(1/3)*gamma(7/3) + 9*a**(7/3)*b*x**3*(1 + b*x**3/a)**(1/3)*gamma(7/3))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.79

$$\int \frac{1}{(a + bx^3)^{7/3}} dx = -\frac{\left(b - \frac{4(bx^3+a)}{x^3}\right)x^4}{4(bx^3 + a)^{\frac{4}{3}}a^2}$$

input `integrate(1/(b*x^3+a)^(7/3),x, algorithm="maxima")`

output `-1/4*(b - 4*(b*x^3 + a)/x^3)*x^4/((b*x^3 + a)^(4/3)*a^2)`

Giac [F]

$$\int \frac{1}{(a + bx^3)^{7/3}} dx = \int \frac{1}{(bx^3 + a)^{7/3}} dx$$

input `integrate(1/(b*x^3+a)^(7/3),x, algorithm="giac")`

output `integrate((b*x^3 + a)^(-7/3), x)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.72

$$\int \frac{1}{(a + bx^3)^{7/3}} dx = \frac{3x(bx^3 + a) + ax}{4a^2(bx^3 + a)^{4/3}}$$

input `int(1/(a + b*x^3)^(7/3),x)`

output `(3*x*(a + b*x^3) + a*x)/(4*a^2*(a + b*x^3)^(4/3))`

Reduce [F]

$$\int \frac{1}{(a + bx^3)^{7/3}} dx = \int \frac{1}{(bx^3 + a)^{1/3} a^2 + 2(bx^3 + a)^{1/3} abx^3 + (bx^3 + a)^{1/3} b^2x^6} dx$$

input `int(1/(b*x^3+a)^(7/3),x)`

output `int(1/((a + b*x**3)**(1/3)*a**2 + 2*(a + b*x**3)**(1/3)*a*b*x**3 + (a + b*x**3)**(1/3)*b**2*x**6),x)`

3.29 $\int \frac{1}{(a+bx^3)^{10/3}} dx$

Optimal result	234
Mathematica [C] (warning: unable to verify)	234
Rubi [A] (verified)	235
Maple [A] (verified)	236
Fricas [A] (verification not implemented)	236
Sympy [B] (verification not implemented)	237
Maxima [A] (verification not implemented)	238
Giac [F]	238
Mupad [B] (verification not implemented)	239
Reduce [F]	239

Optimal result

Integrand size = 11, antiderivative size = 58

$$\int \frac{1}{(a+bx^3)^{10/3}} dx = \frac{x}{7a(a+bx^3)^{7/3}} + \frac{3x}{14a^2(a+bx^3)^{4/3}} + \frac{9x}{14a^3\sqrt[3]{a+bx^3}}$$

output

```
1/7*x/a/(b*x^3+a)^(7/3)+3/14*x/a^2/(b*x^3+a)^(4/3)+9/14*x/a^3/(b*x^3+a)^(1/3)
```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 2 in optimal.

Time = 55.03 (sec) , antiderivative size = 11702, normalized size of antiderivative = 201.76

$$\int \frac{1}{(a+bx^3)^{10/3}} dx = \text{Result too large to show}$$

input

```
Integrate[(a + b*x^3)^(-10/3),x]
```

output

```
Result too large to show
```

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.14, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {749, 749, 746}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + bx^3)^{10/3}} dx$$

$$\downarrow 749$$

$$\frac{6 \int \frac{1}{(bx^3+a)^{7/3}} dx}{7a} + \frac{x}{7a(a+bx^3)^{7/3}}$$

$$\downarrow 749$$

$$\frac{6 \left(\frac{3 \int \frac{1}{(bx^3+a)^{4/3}} dx}{4a} + \frac{x}{4a(a+bx^3)^{4/3}} \right)}{7a} + \frac{x}{7a(a+bx^3)^{7/3}}$$

$$\downarrow 746$$

$$\frac{6 \left(\frac{3x}{4a^2 \sqrt[3]{a+bx^3}} + \frac{x}{4a(a+bx^3)^{4/3}} \right)}{7a} + \frac{x}{7a(a+bx^3)^{7/3}}$$

input `Int[(a + b*x^3)^(-10/3),x]`

output `x/(7*a*(a + b*x^3)^(7/3)) + (6*(x/(4*a*(a + b*x^3)^(4/3)) + (3*x)/(4*a^2*(a + b*x^3)^(1/3))))/(7*a)`

Defintions of rubi rules used

rule 746 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]`

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.64

method	result	size
gosper	$\frac{x(9b^2x^6+21abx^3+14a^2)}{14(bx^3+a)^{\frac{7}{3}}a^3}$	37
trager	$\frac{x(9b^2x^6+21abx^3+14a^2)}{14(bx^3+a)^{\frac{7}{3}}a^3}$	37
pseudoelliptic	$\frac{x(9b^2x^6+21abx^3+14a^2)}{14(bx^3+a)^{\frac{7}{3}}a^3}$	37
orering	$\frac{x(9b^2x^6+21abx^3+14a^2)}{14(bx^3+a)^{\frac{7}{3}}a^3}$	37

input `int(1/(b*x^3+a)^(10/3),x,method=_RETURNVERBOSE)`

output `1/14*x*(9*b^2*x^6+21*a*b*x^3+14*a^2)/(b*x^3+a)^(7/3)/a^3`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.19

$$\int \frac{1}{(a + bx^3)^{10/3}} dx = \frac{(9b^2x^7 + 21abx^4 + 14a^2x)(bx^3 + a)^{\frac{2}{3}}}{14(a^3b^3x^9 + 3a^4b^2x^6 + 3a^5bx^3 + a^6)}$$

input `integrate(1/(b*x^3+a)^(10/3),x, algorithm="fricas")`

output $\frac{1}{14} \cdot (9b^2x^7 + 21a \cdot bx^4 + 14a^2x) \cdot (bx^3 + a)^{2/3} / (a^3b^3x^9 + 3a^4b^2x^6 + 3a^5bx^3 + a^6)$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 515 vs. $2(51) = 102$.

Time = 0.99 (sec) , antiderivative size = 515, normalized size of antiderivative = 8.88

$$\int \frac{1}{(a + bx^3)^{10/3}} dx = \frac{28a^5x\Gamma\left(\frac{1}{3}\right)}{27a^{\frac{25}{3}}\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma\left(\frac{10}{3}\right) + 81a^{\frac{22}{3}}bx^3\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma\left(\frac{10}{3}\right) + 81a^{\frac{19}{3}}b^2x^6\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma\left(\frac{10}{3}\right) + 27a^{\frac{16}{3}}b^3x^9\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma\left(\frac{10}{3}\right) + 18a^2b^3x^{10}\Gamma\left(\frac{1}{3}\right)} + \frac{70a^4bx^4\Gamma\left(\frac{1}{3}\right)}{27a^{\frac{25}{3}}\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma\left(\frac{10}{3}\right) + 81a^{\frac{22}{3}}bx^3\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma\left(\frac{10}{3}\right) + 81a^{\frac{19}{3}}b^2x^6\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma\left(\frac{10}{3}\right) + 27a^{\frac{16}{3}}b^3x^9\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma\left(\frac{10}{3}\right) + 18a^2b^3x^{10}\Gamma\left(\frac{1}{3}\right)} + \frac{60a^3b^2x^7\Gamma\left(\frac{1}{3}\right)}{27a^{\frac{25}{3}}\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma\left(\frac{10}{3}\right) + 81a^{\frac{22}{3}}bx^3\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma\left(\frac{10}{3}\right) + 81a^{\frac{19}{3}}b^2x^6\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma\left(\frac{10}{3}\right) + 27a^{\frac{16}{3}}b^3x^9\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma\left(\frac{10}{3}\right) + 18a^2b^3x^{10}\Gamma\left(\frac{1}{3}\right)} + \frac{18a^2b^3x^{10}\Gamma\left(\frac{1}{3}\right)}{27a^{\frac{25}{3}}\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma\left(\frac{10}{3}\right) + 81a^{\frac{22}{3}}bx^3\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma\left(\frac{10}{3}\right) + 81a^{\frac{19}{3}}b^2x^6\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma\left(\frac{10}{3}\right) + 27a^{\frac{16}{3}}b^3x^9\sqrt[3]{1 + \frac{bx^3}{a}}\Gamma\left(\frac{10}{3}\right) + 18a^2b^3x^{10}\Gamma\left(\frac{1}{3}\right)}$$

input `integrate(1/(b*x**3+a)**(10/3),x)`

output

```

28*a**5*x*gamma(1/3)/(27*a**(25/3)*(1 + b*x**3/a)**(1/3)*gamma(10/3) + 81*
a**(22/3)*b*x**3*(1 + b*x**3/a)**(1/3)*gamma(10/3) + 81*a**(19/3)*b**2*x**
6*(1 + b*x**3/a)**(1/3)*gamma(10/3) + 27*a**(16/3)*b**3*x**9*(1 + b*x**3/a
)**(1/3)*gamma(10/3)) + 70*a**4*b*x**4*gamma(1/3)/(27*a**(25/3)*(1 + b*x**
3/a)**(1/3)*gamma(10/3) + 81*a**(22/3)*b*x**3*(1 + b*x**3/a)**(1/3)*gamma(
10/3) + 81*a**(19/3)*b**2*x**6*(1 + b*x**3/a)**(1/3)*gamma(10/3) + 27*a**(
16/3)*b**3*x**9*(1 + b*x**3/a)**(1/3)*gamma(10/3)) + 60*a**3*b**2*x**7*gam
ma(1/3)/(27*a**(25/3)*(1 + b*x**3/a)**(1/3)*gamma(10/3) + 81*a**(22/3)*b*x
**3*(1 + b*x**3/a)**(1/3)*gamma(10/3) + 81*a**(19/3)*b**2*x**6*(1 + b*x**3
/a)**(1/3)*gamma(10/3) + 27*a**(16/3)*b**3*x**9*(1 + b*x**3/a)**(1/3)*gamm
a(10/3)) + 18*a**2*b**3*x**10*gamma(1/3)/(27*a**(25/3)*(1 + b*x**3/a)**(1/
3)*gamma(10/3) + 81*a**(22/3)*b*x**3*(1 + b*x**3/a)**(1/3)*gamma(10/3) + 8
1*a**(19/3)*b**2*x**6*(1 + b*x**3/a)**(1/3)*gamma(10/3) + 27*a**(16/3)*b**
3*x**9*(1 + b*x**3/a)**(1/3)*gamma(10/3))

```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.86

$$\int \frac{1}{(a + bx^3)^{10/3}} dx = \frac{\left(2b^2 - \frac{7(bx^3+a)b}{x^3} + \frac{14(bx^3+a)^2}{x^6}\right)x^7}{14(bx^3 + a)^{\frac{7}{3}}a^3}$$

input

```
integrate(1/(b*x^3+a)^(10/3),x, algorithm="maxima")
```

output

```
1/14*(2*b^2 - 7*(b*x^3 + a)*b/x^3 + 14*(b*x^3 + a)^2/x^6)*x^7/((b*x^3 + a)
^(7/3)*a^3)
```

Giac [F]

$$\int \frac{1}{(a + bx^3)^{10/3}} dx = \int \frac{1}{(bx^3 + a)^{\frac{10}{3}}} dx$$

input

```
integrate(1/(b*x^3+a)^(10/3),x, algorithm="giac")
```

output `integrate((b*x^3 + a)^(-10/3), x)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.76

$$\int \frac{1}{(a + bx^3)^{10/3}} dx = \frac{9x(bx^3 + a)^2 + 2a^2x + 3ax(bx^3 + a)}{14a^3(bx^3 + a)^{7/3}}$$

input `int(1/(a + b*x^3)^(10/3),x)`

output `(9*x*(a + b*x^3)^2 + 2*a^2*x + 3*a*x*(a + b*x^3))/(14*a^3*(a + b*x^3)^(7/3))`

Reduce [F]

$$\int \frac{1}{(a + bx^3)^{10/3}} dx = \int \frac{1}{(bx^3 + a)^{\frac{1}{3}} a^3 + 3(bx^3 + a)^{\frac{1}{3}} a^2 bx^3 + 3(bx^3 + a)^{\frac{1}{3}} a b^2 x^6 + (bx^3 + a)^{\frac{1}{3}} b^3 x^9} dx$$

input `int(1/(b*x^3+a)^(10/3),x)`

output `int(1/((a + b*x**3)**(1/3)*a**3 + 3*(a + b*x**3)**(1/3)*a**2*b*x**3 + 3*(a + b*x**3)**(1/3)*a*b**2*x**6 + (a + b*x**3)**(1/3)*b**3*x**9),x)`

3.30 $\int \frac{1}{(a+bx^3)^{13/3}} dx$

Optimal result	240
Mathematica [C] (warning: unable to verify)	240
Rubi [A] (verified)	241
Maple [A] (verified)	242
Fricas [A] (verification not implemented)	243
Sympy [B] (verification not implemented)	243
Maxima [A] (verification not implemented)	244
Giac [F]	245
Mupad [B] (verification not implemented)	245
Reduce [F]	245

Optimal result

Integrand size = 11, antiderivative size = 77

$$\int \frac{1}{(a+bx^3)^{13/3}} dx = \frac{x}{10a(a+bx^3)^{10/3}} + \frac{9x}{70a^2(a+bx^3)^{7/3}} + \frac{27x}{140a^3(a+bx^3)^{4/3}} + \frac{81x}{140a^4\sqrt[3]{a+bx^3}}$$

output

$1/10*x/a/(b*x^3+a)^{(10/3)}+9/70*x/a^2/(b*x^3+a)^{(7/3)}+27/140*x/a^3/(b*x^3+a)^{(4/3)}+81/140*x/a^4/(b*x^3+a)^{(1/3)}$

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 2 in optimal.

Time = 144.93 (sec) , antiderivative size = 21879, normalized size of antiderivative = 284.14

$$\int \frac{1}{(a+bx^3)^{13/3}} dx = \text{Result too large to show}$$

input

`Integrate[(a + b*x^3)^(-13/3),x]`

output

Result too large to show

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.21, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {749, 749, 749, 746}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a + bx^3)^{13/3}} dx \\
 & \quad \downarrow \text{749} \\
 & \frac{9 \int \frac{1}{(bx^3+a)^{10/3}} dx}{10a} + \frac{x}{10a(a + bx^3)^{10/3}} \\
 & \quad \downarrow \text{749} \\
 & \frac{9 \left(\frac{6 \int \frac{1}{(bx^3+a)^{7/3}} dx}{7a} + \frac{x}{7a(a+bx^3)^{7/3}} \right)}{10a} + \frac{x}{10a(a + bx^3)^{10/3}} \\
 & \quad \downarrow \text{749} \\
 & \frac{9 \left(\frac{6 \left(\frac{3 \int \frac{1}{(bx^3+a)^{4/3}} dx}{4a} + \frac{x}{4a(a+bx^3)^{4/3}} \right)}{7a} + \frac{x}{7a(a+bx^3)^{7/3}} \right)}{10a} + \frac{x}{10a(a + bx^3)^{10/3}} \\
 & \quad \downarrow \text{746} \\
 & \frac{9 \left(\frac{6 \left(\frac{3x}{4a^2 \sqrt[3]{a + bx^3}} + \frac{x}{4a(a+bx^3)^{4/3}} \right)}{7a} + \frac{x}{7a(a+bx^3)^{7/3}} \right)}{10a} + \frac{x}{10a(a + bx^3)^{10/3}}
 \end{aligned}$$

input `Int[(a + b*x^3)^(-13/3),x]`

output `x/(10*a*(a + b*x^3)^(10/3)) + (9*(x/(7*a*(a + b*x^3)^(7/3)) + (6*(x/(4*a*(a + b*x^3)^(4/3)) + (3*x)/(4*a^2*(a + b*x^3)^(1/3))))/(7*a)))/(10*a)`

Defintions of rubi rules used

rule 746 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]`

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.62

method	result	size
gospers	$\frac{x(81b^3x^9+270ab^2x^6+315a^2bx^3+140a^3)}{140(bx^3+a)^{\frac{10}{3}}a^4}$	48
trager	$\frac{x(81b^3x^9+270ab^2x^6+315a^2bx^3+140a^3)}{140(bx^3+a)^{\frac{10}{3}}a^4}$	48
pseudoelliptic	$\frac{x(81b^3x^9+270ab^2x^6+315a^2bx^3+140a^3)}{140(bx^3+a)^{\frac{10}{3}}a^4}$	48
orering	$\frac{x(81b^3x^9+270ab^2x^6+315a^2bx^3+140a^3)}{140(bx^3+a)^{\frac{10}{3}}a^4}$	48

input `int(1/(b*x^3+a)^(13/3),x,method=_RETURNVERBOSE)`

output `1/140*x*(81*b^3*x^9+270*a*b^2*x^6+315*a^2*b*x^3+140*a^3)/(b*x^3+a)^(10/3)/a^4`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.18

$$\int \frac{1}{(a + bx^3)^{13/3}} dx = \frac{(81 b^3 x^{10} + 270 a b^2 x^7 + 315 a^2 b x^4 + 140 a^3 x)(bx^3 + a)^{2/3}}{140 (a^4 b^4 x^{12} + 4 a^5 b^3 x^9 + 6 a^6 b^2 x^6 + 4 a^7 b x^3 + a^8)}$$

input `integrate(1/(b*x^3+a)^(13/3),x, algorithm="fricas")`

output `1/140*(81*b^3*x^10 + 270*a*b^2*x^7 + 315*a^2*b*x^4 + 140*a^3*x)*(b*x^3 + a)^(2/3)/(a^4*b^4*x^12 + 4*a^5*b^3*x^9 + 6*a^6*b^2*x^6 + 4*a^7*b*x^3 + a^8)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1550 vs. $2(70) = 140$.

Time = 1.68 (sec) , antiderivative size = 1550, normalized size of antiderivative = 20.13

$$\int \frac{1}{(a + bx^3)^{13/3}} dx = \text{Too large to display}$$

input `integrate(1/(b*x**3+a)**(13/3),x)`

output

```

280*a**14*x*gamma(1/3)/(81*a**(55/3)*(1 + b*x**3/a)**(1/3)*gamma(13/3) + 4
86*a**(52/3)*b*x**3*(1 + b*x**3/a)**(1/3)*gamma(13/3) + 1215*a**(49/3)*b**
2*x**6*(1 + b*x**3/a)**(1/3)*gamma(13/3) + 1620*a**(46/3)*b**3*x**9*(1 + b
*x**3/a)**(1/3)*gamma(13/3) + 1215*a**(43/3)*b**4*x**12*(1 + b*x**3/a)**(1
/3)*gamma(13/3) + 486*a**(40/3)*b**5*x**15*(1 + b*x**3/a)**(1/3)*gamma(13/
3) + 81*a**(37/3)*b**6*x**18*(1 + b*x**3/a)**(1/3)*gamma(13/3)) + 1470*a**
13*b*x**4*gamma(1/3)/(81*a**(55/3)*(1 + b*x**3/a)**(1/3)*gamma(13/3) + 486
*a**(52/3)*b*x**3*(1 + b*x**3/a)**(1/3)*gamma(13/3) + 1215*a**(49/3)*b**2*
x**6*(1 + b*x**3/a)**(1/3)*gamma(13/3) + 1620*a**(46/3)*b**3*x**9*(1 + b*x
**3/a)**(1/3)*gamma(13/3) + 1215*a**(43/3)*b**4*x**12*(1 + b*x**3/a)**(1/3
)*gamma(13/3) + 486*a**(40/3)*b**5*x**15*(1 + b*x**3/a)**(1/3)*gamma(13/3)
+ 81*a**(37/3)*b**6*x**18*(1 + b*x**3/a)**(1/3)*gamma(13/3)) + 3270*a**12
*b**2*x**7*gamma(1/3)/(81*a**(55/3)*(1 + b*x**3/a)**(1/3)*gamma(13/3) + 48
6*a**(52/3)*b*x**3*(1 + b*x**3/a)**(1/3)*gamma(13/3) + 1215*a**(49/3)*b**2
*x**6*(1 + b*x**3/a)**(1/3)*gamma(13/3) + 1620*a**(46/3)*b**3*x**9*(1 + b*
x**3/a)**(1/3)*gamma(13/3) + 1215*a**(43/3)*b**4*x**12*(1 + b*x**3/a)**(1/
3)*gamma(13/3) + 486*a**(40/3)*b**5*x**15*(1 + b*x**3/a)**(1/3)*gamma(13/3
) + 81*a**(37/3)*b**6*x**18*(1 + b*x**3/a)**(1/3)*gamma(13/3)) + 3952*a**1
1*b**3*x**10*gamma(1/3)/(81*a**(55/3)*(1 + b*x**3/a)**(1/3)*gamma(13/3) +
486*a**(52/3)*b*x**3*(1 + b*x**3/a)**(1/3)*gamma(13/3) + 1215*a**(49/3)...

```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.87

$$\int \frac{1}{(a + bx^3)^{13/3}} dx = -\frac{\left(14b^3 - \frac{60(bx^3+a)b^2}{x^3} + \frac{105(bx^3+a)^2b}{x^6} - \frac{140(bx^3+a)^3}{x^9}\right)x^{10}}{140(bx^3+a)^{\frac{10}{3}}a^4}$$

input

```
integrate(1/(b*x^3+a)^(13/3),x, algorithm="maxima")
```

output

```
-1/140*(14*b^3 - 60*(b*x^3 + a)*b^2/x^3 + 105*(b*x^3 + a)^2*b/x^6 - 140*(b
*x^3 + a)^3/x^9)*x^10/((b*x^3 + a)^(10/3)*a^4)
```

Giac [F]

$$\int \frac{1}{(a + bx^3)^{13/3}} dx = \int \frac{1}{(bx^3 + a)^{13/3}} dx$$

input `integrate(1/(b*x^3+a)^(13/3),x, algorithm="giac")`

output `integrate((b*x^3 + a)^(-13/3), x)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.79

$$\int \frac{1}{(a + bx^3)^{13/3}} dx = \frac{81x}{140a^4(bx^3 + a)^{1/3}} + \frac{27x}{140a^3(bx^3 + a)^{4/3}} + \frac{9x}{70a^2(bx^3 + a)^{7/3}} + \frac{x}{10a(bx^3 + a)^{10/3}}$$

input `int(1/(a + b*x^3)^(13/3),x)`

output `(81*x)/(140*a^4*(a + b*x^3)^(1/3)) + (27*x)/(140*a^3*(a + b*x^3)^(4/3)) + (9*x)/(70*a^2*(a + b*x^3)^(7/3)) + x/(10*a*(a + b*x^3)^(10/3))`

Reduce [F]

$$\int \frac{1}{(a + bx^3)^{13/3}} dx = \int \frac{1}{(bx^3 + a)^{1/3} a^4 + 4(bx^3 + a)^{1/3} a^3 b x^3 + 6(bx^3 + a)^{1/3} a^2 b^2 x^6 + 4(bx^3 + a)^{1/3} a b^3 x^9 + (bx^3 + a)^{4/3}}$$

input `int(1/(b*x^3+a)^(13/3),x)`

output `int(1/((a + b*x**3)**(1/3)*a**4 + 4*(a + b*x**3)**(1/3)*a**3*b*x**3 + 6*(a + b*x**3)**(1/3)*a**2*b**2*x**6 + 4*(a + b*x**3)**(1/3)*a*b**3*x**9 + (a + b*x**3)**(1/3)*b**4*x**12),x)`

3.31 $\int (a + bx^3)^{4/3} dx$

Optimal result	246
Mathematica [C] (warning: unable to verify)	247
Rubi [A] (verified)	247
Maple [F]	249
Fricas [F]	249
Sympy [C] (verification not implemented)	249
Maxima [F]	250
Giac [F]	250
Mupad [B] (verification not implemented)	250
Reduce [F]	251

Optimal result

Integrand size = 11, antiderivative size = 47

$$\int (a + bx^3)^{4/3} dx = \frac{ax\sqrt[3]{a + bx^3} \operatorname{Hypergeometric2F1}\left(-\frac{4}{3}, \frac{1}{3}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{\sqrt[3]{1 + \frac{bx^3}{a}}}$$

output `a*x*(b*x^3+a)^(1/3)*hypergeom([-4/3, 1/3], [4/3], -b*x^3/a)/(1+b*x^3/a)^(1/3)`

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 0.21 (sec) , antiderivative size = 272, normalized size of antiderivative = 5.79

$$\int (a + bx^3)^{4/3} dx = \frac{3 \left(\frac{(-1)^{2/3} \sqrt[3]{a}}{\sqrt[3]{b}} + x \right) (a + bx^3)^{4/3} \operatorname{AppellF1} \left(\frac{7}{3}, -\frac{4}{3}, -\frac{4}{3}, \frac{10}{3}, -\frac{\frac{(-1)^{2/3} \sqrt[3]{a}}{\sqrt[3]{b}} + x}{\sqrt[3]{-1} \sqrt[3]{a}} - \frac{(-1)^{2/3} \sqrt[3]{a}}{\sqrt[3]{b}}, -\frac{\frac{(-1)^{2/3} \sqrt[3]{a}}{\sqrt[3]{b}}}{\sqrt[3]{b}} \right)}{7 \left(1 + \frac{\frac{(-1)^{2/3} \sqrt[3]{a}}{\sqrt[3]{b}} + x}{\sqrt[3]{a}} - \frac{(-1)^{2/3} \sqrt[3]{a}}{\sqrt[3]{b}} \right)^{4/3} \left(1 + \frac{\frac{(-1)^{2/3} \sqrt[3]{a}}{\sqrt[3]{b}} + x}{\sqrt[3]{-1} \sqrt[3]{a}} - \frac{(-1)^{2/3} \sqrt[3]{a}}{\sqrt[3]{b}} \right)^{4/3}}$$

input

```
Integrate[(a + b*x^3)^(4/3), x]
```

output

```
(3*(((−1)^(2/3)*a^(1/3))/b^(1/3) + x)*(a + b*x^3)^(4/3)*AppellF1[7/3, −4/3, −4/3, 10/3, −(((−1)^(2/3)*a^(1/3))/b^(1/3) + x)/((−1)^(1/3)*a^(1/3))/b^(1/3) − ((−1)^(2/3)*a^(1/3))/b^(1/3)), −(((−1)^(2/3)*a^(1/3))/b^(1/3) + x)/(a^(1/3)/b^(1/3) − ((−1)^(2/3)*a^(1/3))/b^(1/3))]/(7*(1 + (((−1)^(2/3)*a^(1/3))/b^(1/3) + x)/(a^(1/3)/b^(1/3) − ((−1)^(2/3)*a^(1/3))/b^(1/3)))^(4/3)*(1 + (((−1)^(2/3)*a^(1/3))/b^(1/3) + x)/((−1)^(1/3)*a^(1/3))/b^(1/3) − ((−1)^(2/3)*a^(1/3))/b^(1/3)))^(4/3))
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {779, 778}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx^3)^{4/3} dx$$

$$\begin{array}{c}
 \downarrow 779 \\
 \frac{a \sqrt[3]{a + bx^3} \int \left(\frac{bx^3}{a} + 1 \right)^{4/3} dx}{\sqrt[3]{\frac{bx^3}{a} + 1}} \\
 \downarrow 778 \\
 \frac{ax \sqrt[3]{a + bx^3} \operatorname{Hypergeometric2F1} \left(-\frac{4}{3}, \frac{1}{3}, \frac{4}{3}, -\frac{bx^3}{a} \right)}{\sqrt[3]{\frac{bx^3}{a} + 1}}
 \end{array}$$

input `Int[(a + b*x^3)^(4/3), x]`

output `(a*x*(a + b*x^3)^(1/3)*Hypergeometric2F1[-4/3, 1/3, 4/3, -(b*x^3)/a])/(1 + (b*x^3)/a)^(1/3)`

Defintions of rubi rules used

rule 778 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F1[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] || GtQ[a, 0])`

rule 779 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p]) Int[(1 + b*(x^n/a))^p, x], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && !(IntegerQ[p] || GtQ[a, 0])`

Maple [F]

$$\int (bx^3 + a)^{\frac{4}{3}} dx$$

input `int((b*x^3+a)^(4/3),x)`

output `int((b*x^3+a)^(4/3),x)`

Fricas [F]

$$\int (a + bx^3)^{4/3} dx = \int (bx^3 + a)^{\frac{4}{3}} dx$$

input `integrate((b*x^3+a)^(4/3),x, algorithm="fricas")`

output `integral((b*x^3 + a)^(4/3), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.54 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.79

$$\int (a + bx^3)^{4/3} dx = \frac{a^{\frac{4}{3}} x \Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\begin{matrix} -\frac{4}{3}, \frac{1}{3} \\ \frac{4}{3} \end{matrix} \middle| \frac{bx^3 e^{i\pi}}{a}\right)}{3\Gamma\left(\frac{4}{3}\right)}$$

input `integrate((b*x**3+a)**(4/3),x)`

output `a**(4/3)*x*gamma(1/3)*hyper((-4/3, 1/3), (4/3,), b*x**3*exp_polar(I*pi)/a)/(3*gamma(4/3))`

Maxima [F]

$$\int (a + bx^3)^{4/3} dx = \int (bx^3 + a)^{\frac{4}{3}} dx$$

input `integrate((b*x^3+a)^(4/3),x, algorithm="maxima")`

output `integrate((b*x^3 + a)^(4/3), x)`

Giac [F]

$$\int (a + bx^3)^{4/3} dx = \int (bx^3 + a)^{\frac{4}{3}} dx$$

input `integrate((b*x^3+a)^(4/3),x, algorithm="giac")`

output `integrate((b*x^3 + a)^(4/3), x)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.79

$$\int (a + bx^3)^{4/3} dx = \frac{x (bx^3 + a)^{4/3} {}_2F_1\left(-\frac{4}{3}, \frac{1}{3}; \frac{4}{3}; -\frac{bx^3}{a}\right)}{\left(\frac{bx^3}{a} + 1\right)^{4/3}}$$

input `int((a + b*x^3)^(4/3),x)`

output `(x*(a + b*x^3)^(4/3)*hypergeom([-4/3, 1/3], 4/3, -(b*x^3)/a))/((b*x^3)/a + 1)^(4/3)`

Reduce [F]

$$\int (a + bx^3)^{4/3} dx = \frac{3(bx^3 + a)^{1/3} ax}{5} + \frac{(bx^3 + a)^{1/3} bx^4}{5} + \frac{2 \left(\int \frac{1}{(bx^3+a)^{2/3}} dx \right) a^2}{5}$$

input `int((b*x^3+a)^(4/3),x)`

output `(3*(a + b*x**3)**(1/3)*a*x + (a + b*x**3)**(1/3)*b*x**4 + 2*int((a + b*x**3)**(1/3)/(a + b*x**3),x)*a**2)/5`

3.32 $\int \sqrt[3]{a + bx^3} dx$

Optimal result	252
Mathematica [C] (warning: unable to verify)	252
Rubi [A] (verified)	253
Maple [F]	254
Fricas [F]	254
Sympy [C] (verification not implemented)	255
Maxima [F]	255
Giac [F]	255
Mupad [B] (verification not implemented)	256
Reduce [F]	256

Optimal result

Integrand size = 11, antiderivative size = 46

$$\int \sqrt[3]{a + bx^3} dx = \frac{x \sqrt[3]{a + bx^3} \operatorname{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{1}{3}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{\sqrt[3]{1 + \frac{bx^3}{a}}}$$

output `x*(b*x^3+a)^(1/3)*hypergeom([-1/3, 1/3], [4/3], -b*x^3/a)/(1+b*x^3/a)^(1/3)`

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 0.13 (sec) , antiderivative size = 203, normalized size of antiderivative = 4.41

$$\int \sqrt[3]{a + bx^3} dx = \frac{3 \left((-1)^{2/3} \sqrt[3]{a} + \sqrt[3]{bx} \right) \sqrt[3]{a + bx^3} \operatorname{AppellF1}\left(\frac{4}{3}, -\frac{1}{3}, -\frac{1}{3}, \frac{7}{3}, -\frac{(-1)^{2/3} \left((-1)^{2/3} \sqrt[3]{a} + \sqrt[3]{bx} \right)}{(1 + \sqrt[3]{-1}) \sqrt[3]{a}}, \frac{i + \sqrt{3} - 2i \sqrt[3]{\frac{bx}{a}}}{3i + \sqrt{3}}\right)}{4 \sqrt[3]{2} \sqrt[3]{b} \sqrt[3]{\frac{\sqrt[3]{a} + (-1)^{2/3} \sqrt[3]{bx}}{(1 + \sqrt[3]{-1}) \sqrt[3]{a}}}} \sqrt[3]{\frac{i \left(1 + \frac{\sqrt[3]{bx}}{\sqrt[3]{a}} \right)}{3i + \sqrt{3}}}$$

input `Integrate[(a + b*x^3)^(1/3), x]`

output $(3*((-1)^{(2/3)}*a^{(1/3)} + b^{(1/3)}*x)*(a + b*x^3)^{(1/3)}*AppellF1[4/3, -1/3, -1/3, 7/3, -(((-1)^{(2/3)}*((-1)^{(2/3)}*a^{(1/3)} + b^{(1/3)}*x))/((1 + (-1)^{(1/3)})) * a^{(1/3)})), (I + Sqrt[3] - ((2*I)*b^{(1/3)}*x)/a^{(1/3)})/(3*I + Sqrt[3]))/(4*2^{(1/3)}*b^{(1/3)}*((a^{(1/3)} + (-1)^{(2/3)}*b^{(1/3)}*x)/((1 + (-1)^{(1/3)})*a^{(1/3)}))^{(1/3)}*((I*(1 + (b^{(1/3)}*x)/a^{(1/3)}))/(3*I + Sqrt[3]))^{(1/3)})$

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {779, 778}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt[3]{a + bx^3} dx$$

$$\downarrow \text{779}$$

$$\frac{\sqrt[3]{a + bx^3} \int \sqrt[3]{\frac{bx^3}{a} + 1} dx}{\sqrt[3]{\frac{bx^3}{a} + 1}}$$

$$\downarrow \text{778}$$

$$\frac{x \sqrt[3]{a + bx^3} \text{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{1}{3}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{\sqrt[3]{\frac{bx^3}{a} + 1}}$$

input `Int[(a + b*x^3)^(1/3), x]`

output $(x*(a + b*x^3)^{(1/3)}*Hypergeometric2F1[-1/3, 1/3, 4/3, -(b*x^3)/a])/(1 + (b*x^3)/a)^{(1/3)}$

Defintions of rubi rules used

rule 778 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F1[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] || GtQ[a, 0])`

rule 779 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p]) Int[(1 + b*(x^n/a))^p, x], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && !(IntegerQ[p] || GtQ[a, 0])`

Maple [F]

$$\int (bx^3 + a)^{\frac{1}{3}} dx$$

input `int((b*x^3+a)^(1/3),x)`

output `int((b*x^3+a)^(1/3),x)`

Fricas [F]

$$\int \sqrt[3]{a + bx^3} dx = \int (bx^3 + a)^{\frac{1}{3}} dx$$

input `integrate((b*x^3+a)^(1/3),x, algorithm="fricas")`

output `integral((b*x^3 + a)^(1/3), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.47 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.80

$$\int \sqrt[3]{a + bx^3} dx = \frac{\sqrt[3]{ax}\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\begin{matrix} -\frac{1}{3}, \frac{1}{3} \\ \frac{4}{3} \end{matrix} \middle| \frac{bx^3 e^{i\pi}}{a} \right)}{3\Gamma\left(\frac{4}{3}\right)}$$

input `integrate((b*x**3+a)**(1/3),x)`

output `a**(1/3)*x*gamma(1/3)*hyper((-1/3, 1/3), (4/3,), b*x**3*exp_polar(I*pi)/a)/(3*gamma(4/3))`

Maxima [F]

$$\int \sqrt[3]{a + bx^3} dx = \int (bx^3 + a)^{\frac{1}{3}} dx$$

input `integrate((b*x^3+a)^(1/3),x, algorithm="maxima")`

output `integrate((b*x^3 + a)^(1/3), x)`

Giac [F]

$$\int \sqrt[3]{a + bx^3} dx = \int (bx^3 + a)^{\frac{1}{3}} dx$$

input `integrate((b*x^3+a)^(1/3),x, algorithm="giac")`

output `integrate((b*x^3 + a)^(1/3), x)`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.80

$$\int \sqrt[3]{a + bx^3} dx = \frac{x (bx^3 + a)^{1/3} {}_2F_1\left(-\frac{1}{3}, \frac{1}{3}; \frac{4}{3}; -\frac{bx^3}{a}\right)}{\left(\frac{bx^3}{a} + 1\right)^{1/3}}$$

input `int((a + b*x^3)^(1/3),x)`output `(x*(a + b*x^3)^(1/3)*hypergeom([-1/3, 1/3], 4/3, -(b*x^3)/a))/((b*x^3)/a + 1)^(1/3)`**Reduce [F]**

$$\int \sqrt[3]{a + bx^3} dx = \frac{(bx^3 + a)^{\frac{1}{3}} x}{2} + \frac{\left(\int \frac{1}{(bx^3+a)^{\frac{2}{3}}} dx\right) a}{2}$$

input `int((b*x^3+a)^(1/3),x)`output `((a + b*x**3)**(1/3)*x + int((a + b*x**3)**(1/3)/(a + b*x**3),x)*a)/2`

3.33 $\int \frac{1}{(a+bx^3)^{2/3}} dx$

Optimal result	257
Mathematica [C] (warning: unable to verify)	257
Rubi [A] (verified)	258
Maple [F]	259
Fricas [F]	259
Sympy [C] (verification not implemented)	260
Maxima [F]	260
Giac [F]	260
Mupad [B] (verification not implemented)	261
Reduce [F]	261

Optimal result

Integrand size = 11, antiderivative size = 46

$$\int \frac{1}{(a + bx^3)^{2/3}} dx = \frac{x \left(1 + \frac{bx^3}{a}\right)^{2/3} \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{(a + bx^3)^{2/3}}$$

output `x*(1+b*x^3/a)^(2/3)*hypergeom([1/3, 2/3], [4/3], -b*x^3/a)/(b*x^3+a)^(2/3)`

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.16 (sec) , antiderivative size = 183, normalized size of antiderivative = 3.98

$$\int \frac{1}{(a + bx^3)^{2/3}} dx = \frac{3\sqrt[3]{2} \left((-1)^{2/3} \sqrt[3]{a} + \sqrt[3]{bx}\right) \left(\frac{\sqrt[3]{a+(-1)^{2/3}\sqrt[3]{bx}}}{(1+\sqrt[3]{-1})\sqrt[3]{a}}\right)^{2/3} \sqrt[3]{\frac{i \left(1 + \frac{\sqrt[3]{bx}}{\sqrt[3]{a}}\right)}{3i + \sqrt{3}} \text{Hypergeometric2F1}}{\sqrt[3]{b} (a + bx^3)^{2/3}}}$$

input `Integrate[(a + b*x^3)^(-2/3), x]`

output

$$\frac{(3 \cdot 2^{1/3} \cdot (-1)^{2/3} \cdot a^{1/3} + b^{1/3} \cdot x) \cdot ((a^{1/3} + (-1)^{2/3} \cdot b^{1/3}) \cdot x) / ((1 + (-1)^{1/3}) \cdot a^{1/3})^{2/3} \cdot ((I \cdot (1 + (b^{1/3} \cdot x) / a^{1/3})) / (3 \cdot I + \text{Sqrt}[3]))^{1/3} \cdot \text{Hypergeometric2F1}[1/3, 2/3, 4/3, ((2 \cdot I) \cdot \text{Sqrt}[3] \cdot a^{1/3} + (3 - I \cdot \text{Sqrt}[3]) \cdot b^{1/3} \cdot x) / (2 \cdot (1 + (-1)^{1/3}) \cdot (a^{1/3} + b^{1/3} \cdot x)))] / (b^{1/3} \cdot (a + b \cdot x^3)^{2/3})$$
Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {779, 778}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + bx^3)^{2/3}} dx$$

$$\downarrow 779$$

$$\frac{\left(\frac{bx^3}{a} + 1\right)^{2/3} \int \frac{1}{\left(\frac{bx^3}{a} + 1\right)^{2/3}} dx}{(a + bx^3)^{2/3}}$$

$$\downarrow 778$$

$$\frac{x \left(\frac{bx^3}{a} + 1\right)^{2/3} \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{(a + bx^3)^{2/3}}$$

input

$$\text{Int}[(a + b \cdot x^3)^{-2/3}, x]$$

output

$$(x \cdot (1 + (b \cdot x^3) / a)^{2/3} \cdot \text{Hypergeometric2F1}[1/3, 2/3, 4/3, -((b \cdot x^3) / a)]) / (a + b \cdot x^3)^{2/3}$$

Defintions of rubi rules used

rule 778 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F1[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] || GtQ[a, 0])`

rule 779 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p]) Int[(1 + b*(x^n/a))^p, x], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && !(IntegerQ[p] || GtQ[a, 0])`

Maple [F]

$$\int \frac{1}{(bx^3 + a)^{\frac{2}{3}}} dx$$

input `int(1/(b*x^3+a)^(2/3),x)`

output `int(1/(b*x^3+a)^(2/3),x)`

Fricas [F]

$$\int \frac{1}{(a + bx^3)^{2/3}} dx = \int \frac{1}{(bx^3 + a)^{\frac{2}{3}}} dx$$

input `integrate(1/(b*x^3+a)^(2/3),x, algorithm="fricas")`

output `integral((b*x^3 + a)^(-2/3), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.45 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.78

$$\int \frac{1}{(a + bx^3)^{2/3}} dx = \frac{x\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{2}{3} \middle| \frac{bx^3 e^{i\pi}}{a}\right)}{3a^{2/3}\Gamma\left(\frac{4}{3}\right)}$$

input `integrate(1/(b*x**3+a)**(2/3),x)`

output `x*gamma(1/3)*hyper((1/3, 2/3), (4/3,), b*x**3*exp_polar(I*pi)/a)/(3*a**(2/3)*gamma(4/3))`

Maxima [F]

$$\int \frac{1}{(a + bx^3)^{2/3}} dx = \int \frac{1}{(bx^3 + a)^{2/3}} dx$$

input `integrate(1/(b*x^3+a)^(2/3),x, algorithm="maxima")`

output `integrate((b*x^3 + a)^(-2/3), x)`

Giac [F]

$$\int \frac{1}{(a + bx^3)^{2/3}} dx = \int \frac{1}{(bx^3 + a)^{2/3}} dx$$

input `integrate(1/(b*x^3+a)^(2/3),x, algorithm="giac")`

output `integrate((b*x^3 + a)^(-2/3), x)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.80

$$\int \frac{1}{(a + bx^3)^{2/3}} dx = \frac{x \left(\frac{bx^3}{a} + 1 \right)^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{2}{3}; \frac{4}{3}; -\frac{bx^3}{a}\right)}{(bx^3 + a)^{2/3}}$$

input `int(1/(a + b*x^3)^(2/3),x)`output `(x*((b*x^3)/a + 1)^(2/3)*hypergeom([1/3, 2/3], 4/3, -(b*x^3)/a))/(a + b*x^3)^(2/3)`**Reduce [F]**

$$\int \frac{1}{(a + bx^3)^{2/3}} dx = \int \frac{1}{(bx^3 + a)^{2/3}} dx$$

input `int(1/(b*x^3+a)^(2/3),x)`output `int(1/(a + b*x**3)**(2/3),x)`

3.34 $\int \frac{1}{(a+bx^3)^{5/3}} dx$

Optimal result	262
Mathematica [C] (warning: unable to verify)	262
Rubi [A] (verified)	263
Maple [F]	264
Fricas [F]	265
Sympy [C] (verification not implemented)	265
Maxima [F]	265
Giac [F]	266
Mupad [B] (verification not implemented)	266
Reduce [F]	266

Optimal result

Integrand size = 11, antiderivative size = 49

$$\int \frac{1}{(a+bx^3)^{5/3}} dx = \frac{x \left(1 + \frac{bx^3}{a}\right)^{2/3} \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{5}{3}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{a (a+bx^3)^{2/3}}$$

output `x*(1+b*x^3/a)^(2/3)*hypergeom([1/3, 5/3], [4/3], -b*x^3/a)/a/(b*x^3+a)^(2/3)`

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 10.79 (sec) , antiderivative size = 2028, normalized size of antiderivative = 41.39

$$\int \frac{1}{(a+bx^3)^{5/3}} dx = \text{Result too large to show}$$

input `Integrate[(a + b*x^3)^(-5/3), x]`

output

```
(9*(-1)^(2/3)*((-1)^(1/3)*a^(1/3) - b^(1/3)*x)*((-1)^(2/3)*a^(1/3) + b^(1/3)
(3)*x)*Gamma[2/3]*(-52*a^(4/3)*Hypergeometric2F1[1, 5/3, 10/3, (2*Sqrt[3]*a
^(1/3) - (3*I + Sqrt[3])*b^(1/3)*x)/((3*I + Sqrt[3])*a^(1/3) - 2*Sqrt[3]*b
^(1/3)*x)] - (76*I)*Sqrt[3]*a^(4/3)*Hypergeometric2F1[1, 5/3, 10/3, (2*Sqr
t[3]*a^(1/3) - (3*I + Sqrt[3])*b^(1/3)*x)/((3*I + Sqrt[3])*a^(1/3) - 2*Sqr
t[3]*b^(1/3)*x)] - 40*a*b^(1/3)*x*Hypergeometric2F1[1, 5/3, 10/3, (2*Sqrt[
3]*a^(1/3) - (3*I + Sqrt[3])*b^(1/3)*x)/((3*I + Sqrt[3])*a^(1/3) - 2*Sqrt[
3]*b^(1/3)*x)] - (48*I)*Sqrt[3]*a*b^(1/3)*x*Hypergeometric2F1[1, 5/3, 10/3
, (2*Sqrt[3]*a^(1/3) - (3*I + Sqrt[3])*b^(1/3)*x)/((3*I + Sqrt[3])*a^(1/3)
- 2*Sqrt[3]*b^(1/3)*x)] - 72*a^(2/3)*b^(2/3)*x^2*Hypergeometric2F1[1, 5/3
, 10/3, (2*Sqrt[3]*a^(1/3) - (3*I + Sqrt[3])*b^(1/3)*x)/((3*I + Sqrt[3])*a
^(1/3) - 2*Sqrt[3]*b^(1/3)*x)] - (24*I)*Sqrt[3]*a^(2/3)*b^(2/3)*x^2*Hyperg
eometric2F1[1, 5/3, 10/3, (2*Sqrt[3]*a^(1/3) - (3*I + Sqrt[3])*b^(1/3)*x)/
((3*I + Sqrt[3])*a^(1/3) - 2*Sqrt[3]*b^(1/3)*x)] - 36*a^(1/3)*b*x^3*Hyperg
eometric2F1[1, 5/3, 10/3, (2*Sqrt[3]*a^(1/3) - (3*I + Sqrt[3])*b^(1/3)*x)/
((3*I + Sqrt[3])*a^(1/3) - 2*Sqrt[3]*b^(1/3)*x)] - (108*I)*Sqrt[3]*a^(1/3)
*b*x^3*Hypergeometric2F1[1, 5/3, 10/3, (2*Sqrt[3]*a^(1/3) - (3*I + Sqrt[3]
)*b^(1/3)*x)/((3*I + Sqrt[3])*a^(1/3) - 2*Sqrt[3]*b^(1/3)*x)] - (72*I)*Sqr
t[3]*b^(4/3)*x^4*Hypergeometric2F1[1, 5/3, 10/3, (2*Sqrt[3]*a^(1/3) - (3*I
+ Sqrt[3])*b^(1/3)*x)/((3*I + Sqrt[3])*a^(1/3) - 2*Sqrt[3]*b^(1/3)*x)]...
```

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {779, 778}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + bx^3)^{5/3}} dx$$

$$\downarrow 779$$

$$\frac{\left(\frac{bx^3}{a} + 1\right)^{2/3} \int \frac{1}{\left(\frac{bx^3}{a} + 1\right)^{5/3}} dx}{a(a + bx^3)^{2/3}}$$

$$\downarrow 778$$

$$\frac{x \left(\frac{bx^3}{a} + 1 \right)^{2/3} \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{5}{3}, \frac{4}{3}, -\frac{bx^3}{a} \right)}{a (a + bx^3)^{2/3}}$$

input `Int[(a + b*x^3)^(-5/3), x]`

output `(x*(1 + (b*x^3)/a)^(2/3)*Hypergeometric2F1[1/3, 5/3, 4/3, -(b*x^3)/a])/ (a*(a + b*x^3)^(2/3))`

Defintions of rubi rules used

rule 778 `Int[((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F1[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] || GtQ[a, 0])`

rule 779 `Int[((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p]) Int[(1 + b*(x^n/a))^p, x], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && !(IntegerQ[p] || GtQ[a, 0])`

Maple [F]

$$\int \frac{1}{(bx^3 + a)^{5/3}} dx$$

input `int(1/(b*x^3+a)^(5/3), x)`

output `int(1/(b*x^3+a)^(5/3), x)`

Fricas [F]

$$\int \frac{1}{(a + bx^3)^{5/3}} dx = \int \frac{1}{(bx^3 + a)^{5/3}} dx$$

input `integrate(1/(b*x^3+a)^(5/3),x, algorithm="fricas")`

output `integral((b*x^3 + a)^(1/3)/(b^2*x^6 + 2*a*b*x^3 + a^2), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.66 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.73

$$\int \frac{1}{(a + bx^3)^{5/3}} dx = \frac{x\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{5}{3} \middle| \frac{bx^3 e^{i\pi}}{a}\right)}{3a^{5/3}\Gamma\left(\frac{4}{3}\right)}$$

input `integrate(1/(b*x**3+a)**(5/3),x)`

output `x*gamma(1/3)*hyper((1/3, 5/3), (4/3,), b*x**3*exp_polar(I*pi)/a)/(3*a**(5/3)*gamma(4/3))`

Maxima [F]

$$\int \frac{1}{(a + bx^3)^{5/3}} dx = \int \frac{1}{(bx^3 + a)^{5/3}} dx$$

input `integrate(1/(b*x^3+a)^(5/3),x, algorithm="maxima")`

output `integrate((b*x^3 + a)^(-5/3), x)`

Giac [F]

$$\int \frac{1}{(a + bx^3)^{5/3}} dx = \int \frac{1}{(bx^3 + a)^{5/3}} dx$$

input `integrate(1/(b*x^3+a)^(5/3),x, algorithm="giac")`

output `integrate((b*x^3 + a)^(-5/3), x)`

Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.76

$$\int \frac{1}{(a + bx^3)^{5/3}} dx = \frac{x \left(\frac{bx^3}{a} + 1 \right)^{5/3} {}_2F_1 \left(\frac{1}{3}, \frac{5}{3}; \frac{4}{3}; -\frac{bx^3}{a} \right)}{(bx^3 + a)^{5/3}}$$

input `int(1/(a + b*x^3)^(5/3),x)`

output `(x*((b*x^3)/a + 1)^(5/3)*hypergeom([1/3, 5/3], 4/3, -(b*x^3)/a))/(a + b*x^3)^(5/3)`

Reduce [F]

$$\int \frac{1}{(a + bx^3)^{5/3}} dx = \int \frac{1}{(bx^3 + a)^{2/3} a + (bx^3 + a)^{2/3} bx^3} dx$$

input `int(1/(b*x^3+a)^(5/3),x)`

output `int(1/((a + b*x**3)**(2/3)*a + (a + b*x**3)**(2/3)*b*x**3),x)`

3.35 $\int \frac{1}{\sqrt[3]{a - bx^3}} dx$

Optimal result	267
Mathematica [A] (verified)	267
Rubi [A] (verified)	268
Maple [A] (verified)	269
Fricas [B] (verification not implemented)	269
Sympy [C] (verification not implemented)	270
Maxima [A] (verification not implemented)	271
Giac [F]	271
Mupad [B] (verification not implemented)	272
Reduce [F]	272

Optimal result

Integrand size = 12, antiderivative size = 72

$$\int \frac{1}{\sqrt[3]{a - bx^3}} dx = -\frac{\arctan\left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a - bx^3}}}{\sqrt{3}}\right)}{\sqrt{3}\sqrt[3]{b}} + \frac{\log\left(\sqrt[3]{bx} + \sqrt[3]{a - bx^3}\right)}{2\sqrt[3]{b}}$$

output

```
-1/3*arctan(1/3*(1-2*b^(1/3)*x/(-b*x^3+a)^(1/3))*3^(1/2))*3^(1/2)/b^(1/3)+
1/2*ln(b^(1/3)*x+(-b*x^3+a)^(1/3))/b^(1/3)
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.61

$$\int \frac{1}{\sqrt[3]{a - bx^3}} dx = \frac{2\sqrt{3} \arctan\left(\frac{-1 + \frac{2\sqrt[3]{bx}}{\sqrt[3]{a - bx^3}}}{\sqrt{3}}\right) - \log\left(1 + \frac{b^{2/3}x^2}{(a - bx^3)^{2/3}} - \frac{\sqrt[3]{bx}}{\sqrt[3]{a - bx^3}}\right) + 2\log\left(1 + \frac{\sqrt[3]{bx}}{\sqrt[3]{a - bx^3}}\right)}{6\sqrt[3]{b}}$$

input `Integrate[(a - b*x^3)^(-1/3),x]`

output `(2*Sqrt[3]*ArcTan[(-1 + (2*b^(1/3)*x)/(a - b*x^3)^(1/3))/Sqrt[3]] - Log[1 + (b^(2/3)*x^2)/(a - b*x^3)^(2/3) - (b^(1/3)*x)/(a - b*x^3)^(1/3)] + 2*Log[1 + (b^(1/3)*x)/(a - b*x^3)^(1/3)])/(6*b^(1/3))`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {769}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt[3]{a - bx^3}} dx$$

↓ 769

$$\frac{\log\left(\sqrt[3]{a - bx^3} + \sqrt[3]{bx}\right)}{2\sqrt[3]{b}} - \frac{\arctan\left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a - bx^3}}}{\sqrt{3}}\right)}{\sqrt{3}\sqrt[3]{b}}$$

input `Int[(a - b*x^3)^(-1/3),x]`

output `-(ArcTan[(1 - (2*b^(1/3)*x)/(a - b*x^3)^(1/3))/Sqrt[3]]/(Sqrt[3]*b^(1/3))) + Log[b^(1/3)*x + (a - b*x^3)^(1/3)]/(2*b^(1/3))`

Defintions of rubi rules used

rule 769

```
Int[((a_) + (b_.)*(x_)^3)^(-1/3), x_Symbol] :> Simp[ArcTan[(1 + 2*Rt[b, 3]*
(x/(a + b*x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*Rt[b, 3]), x] - Simp[Log[(a + b*x^
3)^(1/3) - Rt[b, 3]*x]/(2*Rt[b, 3]), x] /; FreeQ[{a, b}, x]
```

Maple [A] (verified)

Time = 1.86 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.43

method	result	size
pseudoelliptic	$\frac{\sqrt{3} \arctan\left(\frac{\sqrt{3} \left(b^{\frac{1}{3}} x - 2(-bx^3+a)^{\frac{1}{3}}\right)}{3b^{\frac{1}{3}} x}\right) + \ln\left(\frac{b^{\frac{1}{3}} x + (-bx^3+a)^{\frac{1}{3}}}{x}\right) - \frac{\ln\left(\frac{b^{\frac{2}{3}} x^2 - b^{\frac{1}{3}}(-bx^3+a)^{\frac{1}{3}} x + (-bx^3+a)^{\frac{2}{3}}}{x^2}\right)}{2}}{3b^{\frac{1}{3}}}$	103

input

```
int(1/(-b*x^3+a)^(1/3), x, method=_RETURNVERBOSE)
```

output

```
1/3/b^(1/3)*(3^(1/2)*arctan(1/3*3^(1/2)*(b^(1/3)*x-2*(-b*x^3+a)^(1/3))/b^(
1/3)/x)+ln((b^(1/3)*x+(-b*x^3+a)^(1/3))/x)-1/2*ln((b^(2/3)*x^2-b^(1/3)*(-b
*x^3+a)^(1/3)*x+(-b*x^3+a)^(2/3))/x^2))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 113 vs. 2(55) = 110.

Time = 0.07 (sec) , antiderivative size = 284, normalized size of antiderivative = 3.94

$$\int \frac{1}{\sqrt[3]{a - bx^3}} dx$$

$$= \left[\frac{3 \sqrt{\frac{1}{3}b} \sqrt{-\frac{1}{b^{\frac{2}{3}}}} \log\left(-3bx^3 - 3(-bx^3+a)^{\frac{1}{3}}b^{\frac{2}{3}}x^2 + 3\sqrt{\frac{1}{3}}\left(b^{\frac{4}{3}}x^3 - (-bx^3+a)^{\frac{1}{3}}bx^2 - 2(-bx^3+a)^{\frac{2}{3}}b^{\frac{2}{3}}x\right)\right)}{6b} \right]$$

input `integrate(1/(-b*x^3+a)^(1/3),x, algorithm="fricas")`

output `[1/6*(3*sqrt(1/3)*b*sqrt(-1/b^(2/3))*log(-3*b*x^3 - 3*(-b*x^3 + a)^(1/3)*b^(2/3)*x^2 + 3*sqrt(1/3)*(b^(4/3)*x^3 - (-b*x^3 + a)^(1/3)*b*x^2 - 2*(-b*x^3 + a)^(2/3)*b^(2/3)*x)*sqrt(-1/b^(2/3)) + 2*a) + 2*b^(2/3)*log((b^(1/3)*x + (-b*x^3 + a)^(1/3))/x) - b^(2/3)*log((b^(2/3)*x^2 - (-b*x^3 + a)^(1/3)*b^(1/3)*x + (-b*x^3 + a)^(2/3))/x^2))/b, 1/6*(6*sqrt(1/3)*b^(2/3)*arctan(sqrt(1/3)*(b^(1/3)*x - 2*(-b*x^3 + a)^(1/3))/(b^(1/3)*x)) + 2*b^(2/3)*log((b^(1/3)*x + (-b*x^3 + a)^(1/3))/x) - b^(2/3)*log((b^(2/3)*x^2 - (-b*x^3 + a)^(1/3)*b^(1/3)*x + (-b*x^3 + a)^(2/3))/x^2))/b]`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.52 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.51

$$\int \frac{1}{\sqrt[3]{a - bx^3}} dx = \frac{x\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{1}{3} \middle| \frac{bx^3 e^{2i\pi}}{a}\right)}{3\sqrt[3]{a}\Gamma\left(\frac{4}{3}\right)}$$

input `integrate(1/(-b*x**3+a)**(1/3),x)`

output `x*gamma(1/3)*hyper((1/3, 1/3), (4/3,), b*x**3*exp_polar(2*I*pi)/a)/(3*a**
1/3)*gamma(4/3)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.43

$$\int \frac{1}{\sqrt[3]{a-bx^3}} dx = -\frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(b^{\frac{1}{3}} - \frac{2(-bx^3+a)^{\frac{1}{3}}}{x}\right)}{3b^{\frac{1}{3}}}\right)}{3b^{\frac{1}{3}}} - \frac{\log\left(b^{\frac{2}{3}} - \frac{(-bx^3+a)^{\frac{1}{3}}b^{\frac{1}{3}}}{x} + \frac{(-bx^3+a)^{\frac{2}{3}}}{x^2}\right)}{6b^{\frac{1}{3}}} + \frac{\log\left(b^{\frac{1}{3}} + \frac{(-bx^3+a)^{\frac{1}{3}}}{x}\right)}{3b^{\frac{1}{3}}}$$

input `integrate(1/(-b*x^3+a)^(1/3),x, algorithm="maxima")`output `-1/3*sqrt(3)*arctan(-1/3*sqrt(3)*(b^(1/3) - 2*(-b*x^3 + a)^(1/3)/x)/b^(1/3)))/b^(1/3) - 1/6*log(b^(2/3) - (-b*x^3 + a)^(1/3)*b^(1/3)/x + (-b*x^3 + a)^(2/3)/x^2)/b^(1/3) + 1/3*log(b^(1/3) + (-b*x^3 + a)^(1/3)/x)/b^(1/3)`**Giac [F]**

$$\int \frac{1}{\sqrt[3]{a-bx^3}} dx = \int \frac{1}{(-bx^3 + a)^{\frac{1}{3}}} dx$$

input `integrate(1/(-b*x^3+a)^(1/3),x, algorithm="giac")`output `integrate((-b*x^3 + a)^(-1/3), x)`

Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.53

$$\int \frac{1}{\sqrt[3]{a - bx^3}} dx = \frac{x \left(1 - \frac{bx^3}{a}\right)^{1/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{3}; \frac{4}{3}; \frac{bx^3}{a}\right)}{(a - bx^3)^{1/3}}$$

input `int(1/(a - b*x^3)^(1/3),x)`output `(x*(1 - (b*x^3)/a)^(1/3)*hypergeom([1/3, 1/3], 4/3, (b*x^3)/a))/(a - b*x^3)^(1/3)`**Reduce [F]**

$$\int \frac{1}{\sqrt[3]{a - bx^3}} dx = \int \frac{1}{(-bx^3 + a)^{1/3}} dx$$

input `int(1/(-b*x^3+a)^(1/3),x)`output `int(1/(a - b*x**3)**(1/3),x)`

3.36 $\int \frac{1}{\sqrt[3]{2+x^3}} dx$

Optimal result	273
Mathematica [A] (verified)	273
Rubi [A] (verified)	274
Maple [C] (verified)	275
Fricas [A] (verification not implemented)	275
Sympy [C] (verification not implemented)	276
Maxima [A] (verification not implemented)	276
Giac [F]	277
Mupad [B] (verification not implemented)	277
Reduce [F]	277

Optimal result

Integrand size = 9, antiderivative size = 46

$$\int \frac{1}{\sqrt[3]{2+x^3}} dx = \frac{\arctan\left(\frac{1+\frac{2x}{\sqrt[3]{2+x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{2} \log\left(-x + \sqrt[3]{2+x^3}\right)$$

output

`1/3*arctan(1/3*(1+2*x/(x^3+2)^(1/3))*3^(1/2))*3^(1/2)-1/2*ln(-x+(x^3+2)^(1/3))`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.70

$$\int \frac{1}{\sqrt[3]{2+x^3}} dx = \frac{\arctan\left(\frac{1+\frac{2x}{\sqrt[3]{2+x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{3} \log\left(1 - \frac{x}{\sqrt[3]{2+x^3}}\right) + \frac{1}{6} \log\left(1 + \frac{x^2}{(2+x^3)^{2/3}} + \frac{x}{\sqrt[3]{2+x^3}}\right)$$

input

`Integrate[(2 + x^3)^(-1/3), x]`

output `ArcTan[(1 + (2*x)/(2 + x^3)^(1/3))/Sqrt[3]]/Sqrt[3] - Log[1 - x/(2 + x^3)^(1/3)]/3 + Log[1 + x^2/(2 + x^3)^(2/3) + x/(2 + x^3)^(1/3)]/6`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {769}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt[3]{x^3 + 2}} dx$$

↓ 769

$$\frac{\arctan\left(\frac{\frac{2x}{\sqrt[3]{x^3 + 2}} + 1}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{2} \log\left(\sqrt[3]{x^3 + 2} - x\right)$$

input `Int[(2 + x^3)^(-1/3), x]`

output `ArcTan[(1 + (2*x)/(2 + x^3)^(1/3))/Sqrt[3]]/Sqrt[3] - Log[-x + (2 + x^3)^(1/3)]/2`

Defintions of rubi rules used

rule 769 `Int[((a_) + (b_.)*(x_)^3)^(-1/3), x_Symbol] :=> Simp[ArcTan[(1 + 2*Rt[b, 3]*(x/(a + b*x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*Rt[b, 3]), x] - Simp[Log[(a + b*x^3)^(1/3) - Rt[b, 3]*x]/(2*Rt[b, 3]), x] /; FreeQ[{a, b}, x]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3.

Time = 2.26 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.39

method	result
meijerg	$\frac{2^{\frac{2}{3}} x \operatorname{hypergeom}\left(\left[\frac{1}{3}, \frac{1}{3}\right], \left[\frac{4}{3}\right], -\frac{x^3}{2}\right)}{2}$
pseudoelliptic	$-\frac{\ln\left(\frac{-x+(x^3+2)^{\frac{1}{3}}}{x}\right)}{3} + \frac{\ln\left(\frac{(x^3+2)^{\frac{2}{3}}+(x^3+2)^{\frac{1}{3}}x+x^2}{x^2}\right)}{6} - \frac{\sqrt{3} \arctan\left(\frac{\left(x+2(x^3+2)^{\frac{1}{3}}\right)\sqrt{3}}{3x}\right)}{3}$
trager	$\frac{\operatorname{RootOf}\left(-Z^2-Z+1\right) \ln\left(\operatorname{RootOf}\left(-Z^2-Z+1\right)^2 x^3-3 \operatorname{RootOf}\left(-Z^2-Z+1\right)\left(x^3+2\right)^{\frac{2}{3}} x-3 \operatorname{RootOf}\left(-Z^2-Z+1\right)\right)}{3}$

input `int(1/(x^3+2)^(1/3), x, method=_RETURNVERBOSE)`

output `1/2*2^(2/3)*x*hypergeom([1/3,1/3],[4/3],-1/2*x^3)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.65

$$\int \frac{1}{\sqrt[3]{2+x^3}} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{\sqrt{3}x + 2\sqrt{3}(x^3+2)^{\frac{1}{3}}}{3x}\right) - \frac{1}{3} \log\left(-\frac{x - (x^3+2)^{\frac{1}{3}}}{x}\right) + \frac{1}{6} \log\left(\frac{x^2 + (x^3+2)^{\frac{1}{3}}x + (x^3+2)^{\frac{2}{3}}}{x^2}\right)$$

input `integrate(1/(x^3+2)^(1/3), x, algorithm="fricas")`

output `-1/3*sqrt(3)*arctan(1/3*(sqrt(3)*x + 2*sqrt(3)*(x^3 + 2)^(1/3))/x) - 1/3*log(-(x - (x^3 + 2)^(1/3))/x) + 1/6*log((x^2 + (x^3 + 2)^(1/3)*x + (x^3 + 2)^(2/3))/x^2)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.43 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.74

$$\int \frac{1}{\sqrt[3]{2+x^3}} dx = \frac{2^{\frac{2}{3}} x \Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{1}{3} \middle| \frac{x^3 e^{i\pi}}{2}\right)}{6 \Gamma\left(\frac{4}{3}\right)}$$

input `integrate(1/(x**3+2)**(1/3),x)`

output `2**(2/3)*x*gamma(1/3)*hyper((1/3, 1/3), (4/3,), x**3*exp_polar(I*pi)/2)/(6*gamma(4/3))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.50

$$\int \frac{1}{\sqrt[3]{2+x^3}} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3} \left(\frac{2(x^3+2)^{\frac{1}{3}}}{x} + 1\right)\right) + \frac{1}{6} \log\left(\frac{(x^3+2)^{\frac{1}{3}}}{x} + \frac{(x^3+2)^{\frac{2}{3}}}{x^2} + 1\right) - \frac{1}{3} \log\left(\frac{(x^3+2)^{\frac{1}{3}}}{x} - 1\right)$$

input `integrate(1/(x^3+2)^(1/3),x,algorithm="maxima")`

output `-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*(x^3 + 2)^(1/3)/x + 1)) + 1/6*log((x^3 + 2)^(1/3)/x + (x^3 + 2)^(2/3)/x^2 + 1) - 1/3*log((x^3 + 2)^(1/3)/x - 1)`

Giac [F]

$$\int \frac{1}{\sqrt[3]{2+x^3}} dx = \int \frac{1}{(x^3+2)^{\frac{1}{3}}} dx$$

input `integrate(1/(x^3+2)^(1/3),x, algorithm="giac")`

output `integrate((x^3 + 2)^(-1/3), x)`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.35

$$\int \frac{1}{\sqrt[3]{2+x^3}} dx = \frac{2^{2/3} x {}_2F_1\left(\frac{1}{3}, \frac{1}{3}; \frac{4}{3}; -\frac{x^3}{2}\right)}{2}$$

input `int(1/(x^3 + 2)^(1/3),x)`

output `(2^(2/3)*x*hypergeom([1/3, 1/3], 4/3, -x^3/2))/2`

Reduce [F]

$$\int \frac{1}{\sqrt[3]{2+x^3}} dx = \int \frac{1}{(x^3+2)^{\frac{1}{3}}} dx$$

input `int(1/(x^3+2)^(1/3),x)`

output `int(1/(x**3 + 2)**(1/3),x)`

3.37 $\int (a + bx^3)^{3/4} dx$

Optimal result	278
Mathematica [C] (warning: unable to verify)	278
Rubi [A] (verified)	279
Maple [F]	280
Fricas [F]	280
Sympy [C] (verification not implemented)	281
Maxima [F]	281
Giac [F]	281
Mupad [B] (verification not implemented)	282
Reduce [F]	282

Optimal result

Integrand size = 11, antiderivative size = 46

$$\int (a + bx^3)^{3/4} dx = \frac{x(a + bx^3)^{3/4} \operatorname{Hypergeometric2F1}\left(-\frac{3}{4}, \frac{1}{3}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{\left(1 + \frac{bx^3}{a}\right)^{3/4}}$$

output

`x*(b*x^3+a)^(3/4)*hypergeom([-3/4, 1/3], [4/3], -b*x^3/a)/(1+b*x^3/a)^(3/4)`

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 0.12 (sec) , antiderivative size = 203, normalized size of antiderivative = 4.41

$$\int (a$$

$$+bx^3)^{3/4} dx = \frac{2^4 \sqrt{2} \left((-1)^{2/3} \sqrt[3]{a} + \sqrt[3]{bx} \right) (a + bx^3)^{3/4} \operatorname{AppellF1} \left(\frac{7}{4}, -\frac{3}{4}, -\frac{3}{4}, \frac{11}{4}, -\frac{(-1)^{2/3} \left((-1)^{2/3} \sqrt[3]{a} + \sqrt[3]{bx} \right)}{(1 + \sqrt[3]{-1}) \sqrt[3]{a}} \right)}{7 \sqrt[3]{b} \left(\frac{\sqrt[3]{a} + (-1)^{2/3} \sqrt[3]{bx}}{(1 + \sqrt[3]{-1}) \sqrt[3]{a}} \right)^{3/4} \left(\frac{i \left(1 + \frac{\sqrt[3]{bx}}{\sqrt[3]{a}} \right)}{3i + \sqrt{3}} \right)^{3/4}}$$

input `Integrate[(a + b*x^3)^(3/4), x]`

output `(2*2^(1/4)*((-1)^(2/3)*a^(1/3) + b^(1/3)*x)*(a + b*x^3)^(3/4)*AppellF1[7/4, -3/4, -3/4, 11/4, -(((1)^(2/3)*((-1)^(2/3)*a^(1/3) + b^(1/3)*x))/((1 + (-1)^(1/3))*a^(1/3))], (I + Sqrt[3] - ((2*I)*b^(1/3)*x)/a^(1/3))/(3*I + Sqrt[3])]/(7*b^(1/3)*((a^(1/3) + (-1)^(2/3)*b^(1/3)*x)/((1 + (-1)^(1/3))*a^(1/3)))^(3/4)*((I*(1 + (b^(1/3)*x)/a^(1/3)))/(3*I + Sqrt[3]))^(3/4)`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {779, 778}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx^3)^{3/4} dx$$

$$\downarrow 779$$

$$\frac{(a + bx^3)^{3/4} \int \left(\frac{bx^3}{a} + 1\right)^{3/4} dx}{\left(\frac{bx^3}{a} + 1\right)^{3/4}}$$

$$\downarrow 778$$

$$\frac{x(a + bx^3)^{3/4} \text{Hypergeometric2F1}\left(-\frac{3}{4}, \frac{1}{3}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{\left(\frac{bx^3}{a} + 1\right)^{3/4}}$$

input `Int[(a + b*x^3)^(3/4), x]`

output `(x*(a + b*x^3)^(3/4)*Hypergeometric2F1[-3/4, 1/3, 4/3, -(b*x^3)/a])/((1 + (b*x^3)/a)^(3/4))`

Defintions of rubi rules used

rule 778 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F1[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] || GtQ[a, 0])`

rule 779 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p]) Int[(1 + b*(x^n/a))^p, x], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && !(IntegerQ[p] || GtQ[a, 0])`

Maple [F]

$$\int (bx^3 + a)^{\frac{3}{4}} dx$$

input `int((b*x^3+a)^(3/4),x)`

output `int((b*x^3+a)^(3/4),x)`

Fricas [F]

$$\int (a + bx^3)^{3/4} dx = \int (bx^3 + a)^{\frac{3}{4}} dx$$

input `integrate((b*x^3+a)^(3/4),x, algorithm="fricas")`

output `integral((b*x^3 + a)^(3/4), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.64 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.80

$$\int (a + bx^3)^{3/4} dx = \frac{a^{3/4} x \Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\begin{matrix} -\frac{3}{4}, \frac{1}{3} \\ \frac{4}{3} \end{matrix} \middle| \frac{bx^3 e^{i\pi}}{a}\right)}{3\Gamma\left(\frac{4}{3}\right)}$$

input `integrate((b*x**3+a)**(3/4),x)`

output `a**(3/4)*x*gamma(1/3)*hyper((-3/4, 1/3), (4/3,), b*x**3*exp_polar(I*pi)/a)/(3*gamma(4/3))`

Maxima [F]

$$\int (a + bx^3)^{3/4} dx = \int (bx^3 + a)^{3/4} dx$$

input `integrate((b*x^3+a)^(3/4),x, algorithm="maxima")`

output `integrate((b*x^3 + a)^(3/4), x)`

Giac [F]

$$\int (a + bx^3)^{3/4} dx = \int (bx^3 + a)^{3/4} dx$$

input `integrate((b*x^3+a)^(3/4),x, algorithm="giac")`

output `integrate((b*x^3 + a)^(3/4), x)`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.80

$$\int (a + bx^3)^{3/4} dx = \frac{x (bx^3 + a)^{3/4} {}_2F_1\left(-\frac{3}{4}, \frac{1}{3}; \frac{4}{3}; -\frac{bx^3}{a}\right)}{\left(\frac{bx^3}{a} + 1\right)^{3/4}}$$

input `int((a + b*x^3)^(3/4),x)`output `(x*(a + b*x^3)^(3/4)*hypergeom([-3/4, 1/3], 4/3, -(b*x^3)/a))/((b*x^3)/a + 1)^(3/4)`**Reduce [F]**

$$\int (a + bx^3)^{3/4} dx = \frac{4(bx^3 + a)^{3/4} x}{13} + \frac{9 \left(\int \frac{1}{(bx^3 + a)^{1/4}} dx \right) a}{13}$$

input `int((b*x^3+a)^(3/4),x)`output `(4*(a + b*x**3)**(3/4)*x + 9*int((a + b*x**3)**(3/4)/(a + b*x**3),x)*a)/13`

3.38 $\int \sqrt[4]{a + bx^3} dx$

Optimal result	283
Mathematica [C] (warning: unable to verify)	283
Rubi [A] (verified)	284
Maple [F]	285
Fricas [F]	285
Sympy [C] (verification not implemented)	286
Maxima [F]	286
Giac [F]	286
Mupad [B] (verification not implemented)	287
Reduce [F]	287

Optimal result

Integrand size = 11, antiderivative size = 46

$$\int \sqrt[4]{a + bx^3} dx = \frac{x \sqrt[4]{a + bx^3} \operatorname{Hypergeometric2F1}\left(-\frac{1}{4}, \frac{1}{3}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{\sqrt[4]{1 + \frac{bx^3}{a}}}$$

```
output x*(b*x^3+a)^(1/4)*hypergeom([-1/4, 1/3], [4/3], -b*x^3/a)/(1+b*x^3/a)^(1/4)
```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 0.13 (sec) , antiderivative size = 203, normalized size of antiderivative = 4.41

$$\int \sqrt[4]{a + bx^3} dx$$

$$= \frac{2 \cdot 2^{3/4} \left((-1)^{2/3} \sqrt[3]{a} + \sqrt[3]{bx} \right) \sqrt[4]{a + bx^3} \operatorname{AppellF1} \left(\frac{5}{4}, -\frac{1}{4}, -\frac{1}{4}, \frac{9}{4}, -\frac{(-1)^{2/3} \left((-1)^{2/3} \sqrt[3]{a} + \sqrt[3]{bx} \right)}{(1 + \sqrt[3]{-1}) \sqrt[3]{a}}, \frac{i + \sqrt{3} - \frac{2i \sqrt[3]{bx}}{\sqrt[3]{a}}}{3i + \sqrt{3}} \right)}{5 \sqrt[3]{b} \sqrt[4]{\frac{\sqrt[3]{a} + (-1)^{2/3} \sqrt[3]{bx}}{(1 + \sqrt[3]{-1}) \sqrt[3]{a}}} \sqrt[4]{\frac{i \left(1 + \frac{\sqrt[3]{bx}}{\sqrt[3]{a}} \right)}{3i + \sqrt{3}}}}$$

input `Integrate[(a + b*x^3)^(1/4), x]`

output $(2^{3/4} * (-1)^{2/3} * a^{1/3} + b^{1/3} * x) * (a + b * x^3)^{1/4} * \text{AppellF1}[5/4, -1/4, -1/4, 9/4, -(((-1)^{2/3} * ((-1)^{2/3} * a^{1/3} + b^{1/3} * x)) / ((1 + (-1)^{1/3}) * a^{1/3}))], (I + \text{Sqrt}[3] - ((2 * I) * b^{1/3} * x) / a^{1/3}) / (3 * I + \text{Sqrt}[3])] / (5 * b^{1/3} * ((a^{1/3} + (-1)^{2/3} * b^{1/3} * x) / ((1 + (-1)^{1/3}) * a^{1/3}))^{1/4} * ((I * (1 + (b^{1/3} * x) / a^{1/3})) / (3 * I + \text{Sqrt}[3]))^{1/4})$

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {779, 778}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt[4]{a + bx^3} dx$$

$$\downarrow 779$$

$$\frac{\sqrt[4]{a + bx^3} \int \sqrt[4]{\frac{bx^3}{a} + 1} dx}{\sqrt[4]{\frac{bx^3}{a} + 1}}$$

$$\downarrow 778$$

$$\frac{x \sqrt[4]{a + bx^3} \text{Hypergeometric2F1}\left(-\frac{1}{4}, \frac{1}{3}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{\sqrt[4]{\frac{bx^3}{a} + 1}}$$

input `Int[(a + b*x^3)^(1/4), x]`

output $(x * (a + b * x^3)^{1/4} * \text{Hypergeometric2F1}[-1/4, 1/3, 4/3, -(b * x^3) / a]) / (1 + (b * x^3) / a)^{1/4}$

Defintions of rubi rules used

rule 778 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F1[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] || GtQ[a, 0])`

rule 779 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p]) Int[(1 + b*(x^n/a))^p, x], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && !(IntegerQ[p] || GtQ[a, 0])`

Maple [F]

$$\int (bx^3 + a)^{\frac{1}{4}} dx$$

input `int((b*x^3+a)^(1/4),x)`

output `int((b*x^3+a)^(1/4),x)`

Fricas [F]

$$\int \sqrt[4]{a + bx^3} dx = \int (bx^3 + a)^{\frac{1}{4}} dx$$

input `integrate((b*x^3+a)^(1/4),x, algorithm="fricas")`

output `integral((b*x^3 + a)^(1/4), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.48 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.80

$$\int \sqrt[4]{a + bx^3} dx = \frac{\sqrt[4]{ax}\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\begin{matrix} -\frac{1}{4}, \frac{1}{3} \\ \frac{4}{3} \end{matrix} \middle| \frac{bx^3 e^{i\pi}}{a}\right)}{3\Gamma\left(\frac{4}{3}\right)}$$

input `integrate((b*x**3+a)**(1/4),x)`

output `a**(1/4)*x*gamma(1/3)*hyper((-1/4, 1/3), (4/3,), b*x**3*exp_polar(I*pi)/a)/(3*gamma(4/3))`

Maxima [F]

$$\int \sqrt[4]{a + bx^3} dx = \int (bx^3 + a)^{\frac{1}{4}} dx$$

input `integrate((b*x^3+a)^(1/4),x, algorithm="maxima")`

output `integrate((b*x^3 + a)^(1/4), x)`

Giac [F]

$$\int \sqrt[4]{a + bx^3} dx = \int (bx^3 + a)^{\frac{1}{4}} dx$$

input `integrate((b*x^3+a)^(1/4),x, algorithm="giac")`

output `integrate((b*x^3 + a)^(1/4), x)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.80

$$\int \sqrt[4]{a + bx^3} dx = \frac{x (bx^3 + a)^{1/4} {}_2F_1\left(-\frac{1}{4}, \frac{1}{3}; \frac{4}{3}; -\frac{bx^3}{a}\right)}{\left(\frac{bx^3}{a} + 1\right)^{1/4}}$$

input `int((a + b*x^3)^(1/4),x)`output `(x*(a + b*x^3)^(1/4)*hypergeom([-1/4, 1/3], 4/3, -(b*x^3)/a))/((b*x^3)/a + 1)^(1/4)`**Reduce [F]**

$$\int \sqrt[4]{a + bx^3} dx = \frac{4(bx^3 + a)^{\frac{1}{4}} x}{7} + \frac{3 \left(\int \frac{1}{(bx^3+a)^{\frac{3}{4}}} dx \right) a}{7}$$

input `int((b*x^3+a)^(1/4),x)`output `(4*(a + b*x**3)**(1/4)*x + 3*int((a + b*x**3)**(1/4)/(a + b*x**3),x)*a)/7`

3.39 $\int \frac{1}{\sqrt[4]{a + bx^3}} dx$

Optimal result	288
Mathematica [C] (warning: unable to verify)	288
Rubi [A] (verified)	289
Maple [F]	290
Fricas [F]	290
Sympy [C] (verification not implemented)	291
Maxima [F]	291
Giac [F]	291
Mupad [B] (verification not implemented)	292
Reduce [F]	292

Optimal result

Integrand size = 11, antiderivative size = 46

$$\int \frac{1}{\sqrt[4]{a + bx^3}} dx = \frac{x \sqrt[4]{1 + \frac{bx^3}{a}} \operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1}{3}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{\sqrt[4]{a + bx^3}}$$

output `x*(1+b*x^3/a)^(1/4)*hypergeom([1/4, 1/3], [4/3], -b*x^3/a)/(b*x^3+a)^(1/4)`

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 0.10 (sec) , antiderivative size = 203, normalized size of antiderivative = 4.41

$$\int \frac{1}{\sqrt[4]{a + bx^3}} dx = \frac{4\sqrt[4]{2} \left((-1)^{2/3} \sqrt[3]{a} + \sqrt[3]{bx} \right) \sqrt[4]{\frac{\sqrt[3]{a} + (-1)^{2/3} \sqrt[3]{bx}}{(1 + \sqrt[3]{-1}) \sqrt[3]{a}}} \sqrt[4]{\frac{i \left(1 + \frac{\sqrt[3]{bx}}{\sqrt[3]{a}} \right)}{3i + \sqrt{3}}} \operatorname{AppellF1}\left(\frac{3}{4}, \frac{1}{4}, \frac{1}{4}, \frac{7}{4}, -\frac{(-1)^{2/3} \left((-1)^{2/3} \sqrt[3]{a} + \sqrt[3]{bx} \right)}{(1 + \sqrt[3]{-1}) \sqrt[3]{a}}\right)}{3\sqrt[3]{b} \sqrt[4]{a + bx^3}}$$

input `Integrate[(a + b*x^3)^(-1/4), x]`

output
$$\frac{(4 \cdot 2^{1/4} \cdot (-1)^{2/3} \cdot a^{1/3} + b^{1/3} \cdot x) \cdot ((a^{1/3} + (-1)^{2/3} \cdot b^{1/3}) \cdot x) / ((1 + (-1)^{1/3}) \cdot a^{1/3})^{1/4} \cdot ((I \cdot (1 + (b^{1/3} \cdot x) / a^{1/3})) / (3 \cdot I + \text{Sqrt}[3]))^{1/4} \cdot \text{AppellF1}[3/4, 1/4, 1/4, 7/4, -(((-1)^{2/3} \cdot (-1)^{2/3} \cdot a^{1/3} + b^{1/3} \cdot x) / ((1 + (-1)^{1/3}) \cdot a^{1/3}))], (I + \text{Sqrt}[3] - ((2 \cdot I) \cdot b^{1/3} \cdot x) / a^{1/3}) / (3 \cdot I + \text{Sqrt}[3])]}{3 \cdot b^{1/3} \cdot (a + b \cdot x^3)^{1/4}}$$

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {779, 778}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt[4]{a + bx^3}} dx$$

↓ 779

$$\frac{\sqrt[4]{\frac{bx^3}{a}} + 1 \int \frac{1}{\sqrt[4]{\frac{bx^3}{a}} + 1} dx}{\sqrt[4]{a + bx^3}}$$

↓ 778

$$\frac{x \sqrt[4]{\frac{bx^3}{a}} + 1 \text{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1}{3}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{\sqrt[4]{a + bx^3}}$$

input `Int[(a + b*x^3)^(-1/4), x]`

output
$$(x \cdot (1 + (b \cdot x^3) / a)^{1/4} \cdot \text{Hypergeometric2F1}[1/4, 1/3, 4/3, -((b \cdot x^3) / a)]) / (a + b \cdot x^3)^{1/4}$$

Defintions of rubi rules used

rule 778 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F1[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] || GtQ[a, 0])`

rule 779 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p]) Int[(1 + b*(x^n/a))^p, x], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && !(IntegerQ[p] || GtQ[a, 0])`

Maple [F]

$$\int \frac{1}{(bx^3 + a)^{\frac{1}{4}}} dx$$

input `int(1/(b*x^3+a)^(1/4),x)`

output `int(1/(b*x^3+a)^(1/4),x)`

Fricas [F]

$$\int \frac{1}{\sqrt[4]{a + bx^3}} dx = \int \frac{1}{(bx^3 + a)^{\frac{1}{4}}} dx$$

input `integrate(1/(b*x^3+a)^(1/4),x, algorithm="fricas")`

output `integral((b*x^3 + a)^(-1/4), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.43 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.78

$$\int \frac{1}{\sqrt[4]{a+bx^3}} dx = \frac{x\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{3} \middle| \frac{bx^3 e^{i\pi}}{a}\right)}{3\sqrt[4]{a}\Gamma\left(\frac{4}{3}\right)}$$

input `integrate(1/(b*x**3+a)**(1/4),x)`

output `x*gamma(1/3)*hyper((1/4, 1/3), (4/3,), b*x**3*exp_polar(I*pi)/a)/(3*a**(1/4)*gamma(4/3))`

Maxima [F]

$$\int \frac{1}{\sqrt[4]{a+bx^3}} dx = \int \frac{1}{(bx^3+a)^{\frac{1}{4}}} dx$$

input `integrate(1/(b*x^3+a)^(1/4),x, algorithm="maxima")`

output `integrate((b*x^3 + a)^(-1/4), x)`

Giac [F]

$$\int \frac{1}{\sqrt[4]{a+bx^3}} dx = \int \frac{1}{(bx^3+a)^{\frac{1}{4}}} dx$$

input `integrate(1/(b*x^3+a)^(1/4),x, algorithm="giac")`

output `integrate((b*x^3 + a)^(-1/4), x)`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.80

$$\int \frac{1}{\sqrt[4]{a + bx^3}} dx = \frac{x \left(\frac{bx^3}{a} + 1 \right)^{1/4} {}_2F_1 \left(\frac{1}{4}, \frac{1}{3}; \frac{4}{3}; -\frac{bx^3}{a} \right)}{(bx^3 + a)^{1/4}}$$

input `int(1/(a + b*x^3)^(1/4),x)`output `(x*((b*x^3)/a + 1)^(1/4)*hypergeom([1/4, 1/3], 4/3, -(b*x^3)/a))/(a + b*x^3)^(1/4)`**Reduce [F]**

$$\int \frac{1}{\sqrt[4]{a + bx^3}} dx = \int \frac{1}{(bx^3 + a)^{1/4}} dx$$

input `int(1/(b*x^3+a)^(1/4),x)`output `int(1/(a + b*x**3)**(1/4),x)`

3.40 $\int \frac{1}{(a+bx^3)^{3/4}} dx$

Optimal result	293
Mathematica [C] (warning: unable to verify)	293
Rubi [A] (verified)	294
Maple [F]	295
Fricas [F]	295
Sympy [C] (verification not implemented)	296
Maxima [F]	296
Giac [F]	296
Mupad [B] (verification not implemented)	297
Reduce [F]	297

Optimal result

Integrand size = 11, antiderivative size = 46

$$\int \frac{1}{(a + bx^3)^{3/4}} dx = \frac{x \left(1 + \frac{bx^3}{a}\right)^{3/4} \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{3}{4}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{(a + bx^3)^{3/4}}$$

output `x*(1+b*x^3/a)^(3/4)*hypergeom([1/3, 3/4], [4/3], -b*x^3/a)/(b*x^3+a)^(3/4)`

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 0.11 (sec) , antiderivative size = 201, normalized size of antiderivative = 4.37

$$\int \frac{1}{(a + bx^3)^{3/4}} dx = \frac{4 \cdot 2^{3/4} \left((-1)^{2/3} \sqrt[3]{a} + \sqrt[3]{bx}\right) \left(\frac{\sqrt[3]{a} + (-1)^{2/3} \sqrt[3]{bx}}{(1 + \sqrt[3]{-1}) \sqrt[3]{a}}\right)^{3/4} \left(\frac{i \left(1 + \frac{\sqrt[3]{bx}}{\sqrt[3]{a}}\right)}{3i + \sqrt{3}}\right)^{3/4} \text{AppellF1}\left(\frac{1}{4}, \frac{3}{4}, \frac{3}{4}, \frac{3}{4}, \frac{\sqrt[3]{bx}}{\sqrt[3]{a}}\right)}{\sqrt[3]{b} (a + bx^3)^{3/4}}$$

input `Integrate[(a + b*x^3)^(-3/4), x]`

output

$$\frac{(4 \cdot 2^{3/4} \cdot (-1)^{2/3} \cdot a^{1/3} + b^{1/3} \cdot x) \cdot ((a^{1/3} + (-1)^{2/3} \cdot b^{1/3}) \cdot x) / ((1 + (-1)^{1/3}) \cdot a^{1/3})^{3/4} \cdot ((I \cdot (1 + (b^{1/3} \cdot x) / a^{1/3})) / (3 \cdot I + \text{Sqrt}[3]))^{3/4} \cdot \text{AppellF1}[1/4, 3/4, 3/4, 5/4, -(((-1)^{2/3} \cdot (-1)^{2/3} \cdot a^{1/3} + b^{1/3} \cdot x) / ((1 + (-1)^{1/3}) \cdot a^{1/3}))], (I + \text{Sqrt}[3] - ((2 \cdot I) \cdot b^{1/3} \cdot x) / a^{1/3}) / (3 \cdot I + \text{Sqrt}[3])]}{(b^{1/3} \cdot (a + b \cdot x^3)^{3/4})}$$
Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {779, 778}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + bx^3)^{3/4}} dx$$

$$\downarrow 779$$

$$\frac{\left(\frac{bx^3}{a} + 1\right)^{3/4} \int \frac{1}{\left(\frac{bx^3}{a} + 1\right)^{3/4}} dx}{(a + bx^3)^{3/4}}$$

$$\downarrow 778$$

$$\frac{x \left(\frac{bx^3}{a} + 1\right)^{3/4} \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{3}{4}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{(a + bx^3)^{3/4}}$$

input

$$\text{Int}[(a + b \cdot x^3)^{-3/4}, x]$$

output

$$\frac{(x \cdot (1 + (b \cdot x^3) / a)^{3/4} \cdot \text{Hypergeometric2F1}[1/3, 3/4, 4/3, -((b \cdot x^3) / a)])}{(a + b \cdot x^3)^{3/4}}$$

Defintions of rubi rules used

rule 778 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F1[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] || GtQ[a, 0])`

rule 779 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p]) Int[(1 + b*(x^n/a))^p, x], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && !(IntegerQ[p] || GtQ[a, 0])`

Maple [F]

$$\int \frac{1}{(bx^3 + a)^{\frac{3}{4}}} dx$$

input `int(1/(b*x^3+a)^(3/4),x)`

output `int(1/(b*x^3+a)^(3/4),x)`

Fricas [F]

$$\int \frac{1}{(a + bx^3)^{3/4}} dx = \int \frac{1}{(bx^3 + a)^{\frac{3}{4}}} dx$$

input `integrate(1/(b*x^3+a)^(3/4),x, algorithm="fricas")`

output `integral((b*x^3 + a)^(-3/4), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.44 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.78

$$\int \frac{1}{(a + bx^3)^{3/4}} dx = \frac{x\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{3}{4} \middle| \frac{bx^3 e^{i\pi}}{a}\right)}{3a^{3/4}\Gamma\left(\frac{4}{3}\right)}$$

input `integrate(1/(b*x**3+a)**(3/4),x)`

output `x*gamma(1/3)*hyper((1/3, 3/4), (4/3,), b*x**3*exp_polar(I*pi)/a)/(3*a**(3/4)*gamma(4/3))`

Maxima [F]

$$\int \frac{1}{(a + bx^3)^{3/4}} dx = \int \frac{1}{(bx^3 + a)^{3/4}} dx$$

input `integrate(1/(b*x^3+a)^(3/4),x, algorithm="maxima")`

output `integrate((b*x^3 + a)^(-3/4), x)`

Giac [F]

$$\int \frac{1}{(a + bx^3)^{3/4}} dx = \int \frac{1}{(bx^3 + a)^{3/4}} dx$$

input `integrate(1/(b*x^3+a)^(3/4),x, algorithm="giac")`

output `integrate((b*x^3 + a)^(-3/4), x)`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.80

$$\int \frac{1}{(a + bx^3)^{3/4}} dx = \frac{x \left(\frac{bx^3}{a} + 1 \right)^{3/4} {}_2F_1 \left(\frac{1}{3}, \frac{3}{4}; \frac{4}{3}; -\frac{bx^3}{a} \right)}{(bx^3 + a)^{3/4}}$$

input `int(1/(a + b*x^3)^(3/4),x)`output `(x*((b*x^3)/a + 1)^(3/4)*hypergeom([1/3, 3/4], 4/3, -(b*x^3)/a))/(a + b*x^3)^(3/4)`**Reduce [F]**

$$\int \frac{1}{(a + bx^3)^{3/4}} dx = \int \frac{1}{(bx^3 + a)^{3/4}} dx$$

input `int(1/(b*x^3+a)^(3/4),x)`output `int(1/(a + b*x**3)**(3/4),x)`

3.41 $\int \frac{1}{(a+bx^3)^{5/4}} dx$

Optimal result	298
Mathematica [C] (warning: unable to verify)	298
Rubi [A] (verified)	299
Maple [F]	300
Fricas [F]	300
Sympy [C] (verification not implemented)	301
Maxima [F]	301
Giac [F]	301
Mupad [B] (verification not implemented)	302
Reduce [F]	302

Optimal result

Integrand size = 11, antiderivative size = 49

$$\int \frac{1}{(a+bx^3)^{5/4}} dx = \frac{x \sqrt[4]{1 + \frac{bx^3}{a}} \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{5}{4}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{a \sqrt[4]{a+bx^3}}$$

output

```
x*(1+b*x^3/a)^(1/4)*hypergeom([1/3, 5/4], [4/3], -b*x^3/a)/a/(b*x^3+a)^(1/4)
```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 0.15 (sec) , antiderivative size = 263, normalized size of antiderivative = 5.37

$$\int \frac{1}{(a+bx^3)^{5/4}} dx = \frac{4(-1)^{2/3} \sqrt[4]{2} \left(\sqrt[3]{-1} \sqrt[3]{a} - \sqrt[3]{bx}\right) \left((-1)^{2/3} \sqrt[3]{a} + \sqrt[3]{bx}\right) \sqrt[4]{\frac{\sqrt[3]{a} + (-1)^{2/3} \sqrt[3]{bx}}{(1 + \sqrt[3]{-1}) \sqrt[3]{a}}} \sqrt{\frac{i \left(1 + \sqrt[3]{-1}\right)}{3i + \sqrt[3]{-1}}}}{3a^{2/3} \sqrt[3]{b} \left(a^{2/3} - \sqrt[3]{a} \sqrt[3]{bx} + b^{2/3}\right)}$$

input

```
Integrate[(a + b*x^3)^(-5/4), x]
```

output

```
(4*(-1)^(2/3)*2^(1/4)*((-1)^(1/3)*a^(1/3) - b^(1/3)*x)*((-1)^(2/3)*a^(1/3)
+ b^(1/3)*x)*((a^(1/3) + (-1)^(2/3)*b^(1/3)*x)/((1 + (-1)^(1/3))*a^(1/3))
)^(1/4)*((I*(1 + (b^(1/3)*x)/a^(1/3)))/(3*I + Sqrt[3]))^(1/4)*AppellF1[-1/
4, 5/4, 5/4, 3/4, -(((1)^(2/3)*((-1)^(2/3)*a^(1/3) + b^(1/3)*x))/((1 + (-
1)^(1/3))*a^(1/3))], (I + Sqrt[3] - ((2*I)*b^(1/3)*x)/a^(1/3))/(3*I + Sqrt
[3])]/(3*a^(2/3)*b^(1/3)*(a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2)*(a +
b*x^3)^(1/4))
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {779, 778}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + bx^3)^{5/4}} dx$$

$$\downarrow 779$$

$$\frac{\sqrt[4]{\frac{bx^3}{a}} + 1 \int \frac{1}{\left(\frac{bx^3}{a} + 1\right)^{5/4}} dx}{a^4 \sqrt[4]{a + bx^3}}$$

$$\downarrow 778$$

$$\frac{x \sqrt[4]{\frac{bx^3}{a}} + 1 \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{5}{4}, \frac{4}{3}, -\frac{bx^3}{a}\right)}{a^4 \sqrt[4]{a + bx^3}}$$

input

```
Int[(a + b*x^3)^(-5/4),x]
```

output

```
(x*(1 + (b*x^3)/a)^(1/4)*Hypergeometric2F1[1/3, 5/4, 4/3, -(b*x^3)/a])/
(a*(a + b*x^3)^(1/4))
```

Defintions of rubi rules used

rule 778 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F1[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] || GtQ[a, 0])`

rule 779 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p]) Int[(1 + b*(x^n/a))^p, x], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && !(IntegerQ[p] || GtQ[a, 0])`

Maple [F]

$$\int \frac{1}{(bx^3 + a)^{\frac{5}{4}}} dx$$

input `int(1/(b*x^3+a)^(5/4),x)`

output `int(1/(b*x^3+a)^(5/4),x)`

Fricas [F]

$$\int \frac{1}{(a + bx^3)^{5/4}} dx = \int \frac{1}{(bx^3 + a)^{5/4}} dx$$

input `integrate(1/(b*x^3+a)^(5/4),x, algorithm="fricas")`

output `integral((b*x^3 + a)^(3/4)/(b^2*x^6 + 2*a*b*x^3 + a^2), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.69 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.73

$$\int \frac{1}{(a + bx^3)^{5/4}} dx = \frac{x\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{5}{4} \middle| \frac{4}{3} \middle| \frac{bx^3 e^{i\pi}}{a}\right)}{3a^{5/4}\Gamma\left(\frac{4}{3}\right)}$$

input `integrate(1/(b*x**3+a)**(5/4),x)`

output `x*gamma(1/3)*hyper((1/3, 5/4), (4/3,), b*x**3*exp_polar(I*pi)/a)/(3*a**(5/4)*gamma(4/3))`

Maxima [F]

$$\int \frac{1}{(a + bx^3)^{5/4}} dx = \int \frac{1}{(bx^3 + a)^{5/4}} dx$$

input `integrate(1/(b*x^3+a)^(5/4),x, algorithm="maxima")`

output `integrate((b*x^3 + a)^(-5/4), x)`

Giac [F]

$$\int \frac{1}{(a + bx^3)^{5/4}} dx = \int \frac{1}{(bx^3 + a)^{5/4}} dx$$

input `integrate(1/(b*x^3+a)^(5/4),x, algorithm="giac")`

output `integrate((b*x^3 + a)^(-5/4), x)`

Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.76

$$\int \frac{1}{(a + bx^3)^{5/4}} dx = \frac{x \left(\frac{bx^3}{a} + 1 \right)^{5/4} {}_2F_1 \left(\frac{1}{3}, \frac{5}{4}; \frac{4}{3}; -\frac{bx^3}{a} \right)}{(bx^3 + a)^{5/4}}$$

input `int(1/(a + b*x^3)^(5/4),x)`output `(x*((b*x^3)/a + 1)^(5/4)*hypergeom([1/3, 5/4], 4/3, -(b*x^3)/a))/(a + b*x^3)^(5/4)`**Reduce [F]**

$$\int \frac{1}{(a + bx^3)^{5/4}} dx = \int \frac{1}{(bx^3 + a)^{1/4} a + (bx^3 + a)^{1/4} bx^3} dx$$

input `int(1/(b*x^3+a)^(5/4),x)`output `int(1/((a + b*x**3)**(1/4)*a + (a + b*x**3)**(1/4)*b*x**3),x)`

CHAPTER 4

APPENDIX

4.1	Listing of Grading functions	303
4.2	Links to plain text integration problems used in this report for each CAS .	321

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```



```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
  ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "
  ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]==Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]==Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]==Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]==Plus || Head[expn]==Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```

```

    Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
    If[AppellFunctionQ[Head[expn]],
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
    If[Head[expn]===RootSum,
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
    If[Head[expn]===Integrate || Head[expn]===Int,
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
    9]]]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

```

```

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

```

```

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

```

```

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

#   antiderivative
#   "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
            fi;
        fi;
    else #ExpnType(result) > ExpnType(optimal)
        if debug then
            print("ExpnType(result) > ExpnType(optimal)");
        fi;
        return "C",cat("Result contains higher order function than in optimal. Order ",
                        convert(ExpnType_result,string)," vs. order ",
                        convert(ExpnType_optimal,string),".");
    fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```



```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```



```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file