

Computer Algebra Independent Integration Tests

Summer 2024

1-Algebraic-functions/1.1-Binomial/1.1.3-General-
binomial/1.1.3.1/41-1.1.3.1-b

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 6:06pm

Contents

1	Introduction	4
1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Time and leaf size Performance	10
1.4	Performance based on number of rules Rubi used	12
1.5	Performance based on number of steps Rubi used	13
1.6	Solved integrals histogram based on leaf size of result	14
1.7	Solved integrals histogram based on CPU time used	15
1.8	Leaf size vs. CPU time used	16
1.9	list of integrals with no known antiderivative	17
1.10	List of integrals solved by CAS but has no known antiderivative	17
1.11	list of integrals solved by CAS but failed verification	17
1.12	Timing	18
1.13	Verification	18
1.14	Important notes about some of the results	19
1.15	Current tree layout of integration tests	22
1.16	Design of the test system	23
2	detailed summary tables of results	24
2.1	List of integrals sorted by grade for each CAS	25
2.2	Detailed conclusion table per each integral for all CAS systems	29
2.3	Detailed conclusion table specific for Rubi results	43
3	Listing of integrals	45
3.1	$\int (a + bx^4)^4 dx$	47
3.2	$\int (a + bx^4)^3 dx$	52
3.3	$\int (a + bx^4)^2 dx$	57
3.4	$\int (a + bx^4) dx$	62
3.5	$\int \frac{1}{a+cx^4} dx$	67
3.6	$\int \frac{1}{(a+cx^4)^2} dx$	76

3.7	$\int \frac{1}{(a+bx^4)^3} dx$	86
3.8	$\int \frac{1}{1+a+(-1+a)x^4} dx$	98
3.9	$\int (a-bx^4)^{3/2} dx$	106
3.10	$\int \sqrt{a-bx^4} dx$	112
3.11	$\int \frac{1}{\sqrt{a-bx^4}} dx$	118
3.12	$\int \frac{1}{(a-bx^4)^{3/2}} dx$	123
3.13	$\int \frac{1}{(a-bx^4)^{5/2}} dx$	129
3.14	$\int (a+bx^4)^{3/2} dx$	135
3.15	$\int \sqrt{a+bx^4} dx$	141
3.16	$\int \frac{1}{\sqrt{a+bx^4}} dx$	147
3.17	$\int \frac{1}{(a+bx^4)^{3/2}} dx$	152
3.18	$\int \frac{1}{(a+bx^4)^{5/2}} dx$	158
3.19	$\int (1-x^4)^{5/2} dx$	164
3.20	$\int (1-x^4)^{3/2} dx$	169
3.21	$\int \sqrt{1-x^4} dx$	174
3.22	$\int \frac{1}{\sqrt{1-x^4}} dx$	179
3.23	$\int \frac{1}{(1-x^4)^{3/2}} dx$	184
3.24	$\int \frac{1}{(1-x^4)^{5/2}} dx$	189
3.25	$\int (1+x^4)^{5/2} dx$	194
3.26	$\int (1+x^4)^{3/2} dx$	200
3.27	$\int \sqrt{1+x^4} dx$	206
3.28	$\int \frac{1}{\sqrt{1+x^4}} dx$	211
3.29	$\int \frac{1}{(1+x^4)^{3/2}} dx$	216
3.30	$\int \frac{1}{(1+x^4)^{5/2}} dx$	221
3.31	$\int (a+bx^4)^{7/4} dx$	227
3.32	$\int (a+bx^4)^{3/4} dx$	234
3.33	$\int \frac{1}{\sqrt[4]{a+bx^4}} dx$	241
3.34	$\int \frac{1}{(a+bx^4)^{5/4}} dx$	247
3.35	$\int \frac{1}{(a+bx^4)^{9/4}} dx$	252
3.36	$\int \frac{1}{(a+bx^4)^{13/4}} dx$	257
3.37	$\int \frac{1}{(a+bx^4)^{17/4}} dx$	263
3.38	$\int (a+bx^4)^{9/4} dx$	269
3.39	$\int (a+bx^4)^{5/4} dx$	276
3.40	$\int \sqrt[4]{a+bx^4} dx$	282

3.41	$\int \frac{1}{(a+bx^4)^{3/4}} dx$	288
3.42	$\int \frac{1}{(a+bx^4)^{7/4}} dx$	293
3.43	$\int \frac{1}{(a+bx^4)^{11/4}} dx$	299
3.44	$\int \frac{1}{(a+bx^4)^{15/4}} dx$	305
3.45	$\int (a - bx^4)^{3/4} dx$	312
3.46	$\int \frac{1}{\sqrt[4]{a - bx^4}} dx$	322
3.47	$\int \frac{1}{(a-bx^4)^{5/4}} dx$	331
3.48	$\int \frac{1}{(a-bx^4)^{9/4}} dx$	336
3.49	$\int \frac{1}{(a-bx^4)^{13/4}} dx$	341
3.50	$\int \sqrt[4]{a - bx^4} dx$	347
3.51	$\int \frac{1}{(a-bx^4)^{3/4}} dx$	353
3.52	$\int \frac{1}{(a-bx^4)^{7/4}} dx$	358
4	Appendix	364
4.1	Listing of Grading functions	364
4.2	Links to plain text integration problems used in this report for each CAS382	

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Time and leaf size Performance	10
1.4	Performance based on number of rules Rubi used	12
1.5	Performance based on number of steps Rubi used	13
1.6	Solved integrals histogram based on leaf size of result	14
1.7	Solved integrals histogram based on CPU time used	15
1.8	Leaf size vs. CPU time used	16
1.9	list of integrals with no known antiderivative	17
1.10	List of integrals solved by CAS but has no known antiderivative	17
1.11	list of integrals solved by CAS but failed verification	17
1.12	Timing	18
1.13	Verification	18
1.14	Important notes about some of the results	19
1.15	Current tree layout of integration tests	22
1.16	Design of the test system	23

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [52]. This is test number [41].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (52)	0.00 (0)
Mathematica	100.00 (52)	0.00 (0)
Mupad	100.00 (52)	0.00 (0)
Sympy	100.00 (52)	0.00 (0)
Maple	80.77 (42)	19.23 (10)
Fricas	80.77 (42)	19.23 (10)
Maxima	38.46 (20)	61.54 (32)
Giac	15.38 (8)	84.62 (44)
Reduce	15.38 (8)	84.62 (44)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Maple	51.923	0.000	28.846	19.231
Fricas	51.923	0.000	28.846	19.231
Mathematica	42.308	0.000	57.692	0.000
Maxima	36.538	1.923	0.000	61.538
Sympy	34.615	9.615	55.769	0.000
Giac	11.538	3.846	0.000	84.615
Mupad	0.000	100.000	0.000	0.000
Reduce	0.000	15.385	0.000	84.615

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Mupad	0	0.00	0.00	0.00
Sympy	0	0.00	0.00	0.00
Fricas	10	100.00	0.00	0.00
Maple	10	100.00	0.00	0.00
Maxima	32	100.00	0.00	0.00
Giac	44	100.00	0.00	0.00
Reduce	44	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.07
Fricas	0.08
Mupad	0.08
Giac	0.12
Rubi	0.23
Reduce	0.24
Maple	0.58
Sympy	0.59
Mathematica	4.01

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mupad	40.88	0.62	37.00	0.42
Maple	50.21	0.74	37.00	0.79
Mathematica	60.54	0.82	48.50	0.75
Fricas	71.79	0.99	47.50	0.84
Rubi	85.25	1.08	77.00	1.00
Maxima	98.10	1.17	61.00	0.89
Sympy	99.48	1.77	36.50	0.48
Giac	118.75	1.30	111.00	1.03
Reduce	146.00	1.42	79.00	0.98

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	25
2.2	Detailed conclusion table per each integral for all CAS systems	29
2.3	Detailed conclusion table specific for Rubi results	43

2.1 List of integrals sorted by grade for each CAS

Rubi	25
Mma	25
Maple	26
Fricas	26
Maxima	26
Giac	27
Mupad	27
Sympy	27
Reduce	28

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 21, 22, 31, 32, 33, 34, 35, 36, 37, 45, 46, 47, 48, 49 }

B grade { }

C grade { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 23, 24, 25, 26, 27, 28, 29, 30, 38, 39, 40, 41, 42, 43, 44, 50, 51, 52 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 9, 10, 11, 12, 13, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 45, 46, 47, 48, 49 }

B grade { }

C grade { 5, 6, 7, 8, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 }

F normal fail { 38, 39, 40, 41, 42, 43, 44, 50, 51, 52 }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 3, 4, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 34, 35, 36, 37, 47, 48, 49 }

B grade { }

C grade { 5, 6, 7, 8, 25, 26, 27, 28, 29, 30, 31, 32, 33, 45, 46 }

F normal fail { 38, 39, 40, 41, 42, 43, 44, 50, 51, 52 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maxima

A grade { 1, 2, 3, 4, 5, 6, 7, 31, 32, 33, 34, 35, 36, 37, 45, 46, 47, 48, 49 }

B grade { 8 }

C grade { }

F normal fail { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 38, 39, 40, 41, 42, 43, 44, 50, 51, 52 }

F(-1) timedout fail { }

F(-2) exception fail { }

Giac

A grade { 1, 2, 3, 4, 6, 7 }

B grade { 5, 8 }

C grade { }

F normal fail { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Sympy

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 19, 20, 21, 23, 24 }

B grade { 22, 34, 35, 36, 37 }

C grade { 14, 15, 16, 17, 18, 25, 26, 27, 28, 29, 30, 31, 32, 33, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Reduce

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8 }

C grade { }

F normal fail { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29,
30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	51	44	43	43	49	43	46	43
N.S.	1	1.00	1.00	0.86	0.84	0.84	0.96	0.84	0.90	0.84
time (sec)	N/A	0.174	0.002	0.357	0.027	0.057	0.023	0.117	0.273	0.014

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	38	33	32	32	34	32	35	32
N.S.	1	1.00	1.00	0.87	0.84	0.84	0.89	0.84	0.92	0.84
time (sec)	N/A	0.163	0.002	0.348	0.030	0.069	0.021	0.121	0.243	0.023

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	25	22	21	21	22	21	24	21
N.S.	1	1.00	1.00	0.88	0.84	0.84	0.88	0.84	0.96	0.84
time (sec)	N/A	0.152	0.002	0.337	0.033	0.066	0.020	0.121	0.221	0.017

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	12	11	10	10	8	10	12	10
N.S.	1	1.00	1.00	0.92	0.83	0.83	0.67	0.83	1.00	0.83
time (sec)	N/A	0.134	0.000	0.038	0.034	0.058	0.022	0.119	0.230	0.011

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	C	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	134	200	134	27	169	112	20	179	112	33
N.S.	1	1.49	1.00	0.20	1.26	0.84	0.15	1.34	0.84	0.25
time (sec)	N/A	0.369	0.036	0.481	0.112	0.082	0.082	0.116	0.232	0.085

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	C	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	151	225	183	46	189	183	39	194	305	58
N.S.	1	1.49	1.21	0.30	1.25	1.21	0.26	1.28	2.02	0.38
time (sec)	N/A	0.389	0.078	0.444	0.110	0.077	0.155	0.128	0.249	0.056

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	C	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	168	250	200	57	212	255	63	204	474	80
N.S.	1	1.49	1.19	0.34	1.26	1.52	0.38	1.21	2.82	0.48
time (sec)	N/A	0.403	0.067	0.461	0.127	0.070	0.289	0.128	0.241	0.094

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	C	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	83	160	37	318	196	32	267	160	543
N.S.	1	1.00	1.93	0.45	3.83	2.36	0.39	3.22	1.93	6.54
time (sec)	N/A	0.205	0.058	0.500	0.118	0.080	0.206	0.122	0.249	0.141

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	97	48	92	0	52	39	0	56	38
N.S.	1	1.05	0.52	1.00	0.00	0.57	0.42	0.00	0.61	0.41
time (sec)	N/A	0.197	5.310	0.546	0.000	0.086	0.535	0.000	0.267	0.120

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	88	80	0	42	39	0	38	38
N.S.	1	1.00	1.19	1.08	0.00	0.57	0.53	0.00	0.51	0.51
time (sec)	N/A	0.172	4.220	0.536	0.000	0.070	0.490	0.000	0.239	0.071

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	72	64	0	26	37	0	22	38
N.S.	1	1.00	1.36	1.21	0.00	0.49	0.70	0.00	0.42	0.72
time (sec)	N/A	0.158	10.045	0.457	0.000	0.075	0.475	0.000	0.254	0.085

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	56	90	0	66	37	0	32	38
N.S.	1	1.00	0.73	1.17	0.00	0.86	0.48	0.00	0.42	0.49
time (sec)	N/A	0.175	5.856	0.472	0.000	0.077	0.537	0.000	0.238	0.118

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	105	74	121	0	100	37	0	44	38
N.S.	1	1.08	0.76	1.25	0.00	1.03	0.38	0.00	0.45	0.39
time (sec)	N/A	0.197	10.030	0.477	0.000	0.075	0.609	0.000	0.235	0.129

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	122	127	47	96	0	51	37	0	52	37
N.S.	1	1.04	0.39	0.79	0.00	0.42	0.30	0.00	0.43	0.30
time (sec)	N/A	0.195	5.413	0.553	0.000	0.076	0.520	0.000	0.227	0.091

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	105	105	89	85	0	41	37	0	35	37
N.S.	1	1.00	0.85	0.81	0.00	0.39	0.35	0.00	0.33	0.35
time (sec)	N/A	0.176	4.043	0.530	0.000	0.084	0.473	0.000	0.225	0.064

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	74	70	0	29	36	0	20	37
N.S.	1	1.00	0.84	0.80	0.00	0.33	0.41	0.00	0.23	0.42
time (sec)	N/A	0.163	10.049	0.464	0.000	0.075	0.479	0.000	0.238	0.086

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	108	108	55	94	0	64	36	0	31	37
N.S.	1	1.00	0.51	0.87	0.00	0.59	0.33	0.00	0.29	0.34
time (sec)	N/A	0.182	5.683	0.464	0.000	0.084	0.526	0.000	0.232	0.108

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	127	135	72	123	0	101	36	0	42	37
N.S.	1	1.06	0.57	0.97	0.00	0.80	0.28	0.00	0.33	0.29
time (sec)	N/A	0.198	10.026	0.471	0.000	0.087	0.703	0.000	0.273	0.122

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	67	15	12	0	34	31	0	58	10
N.S.	1	1.18	0.26	0.21	0.00	0.60	0.54	0.00	1.02	0.18
time (sec)	N/A	0.174	4.712	0.509	0.000	0.072	0.508	0.000	0.234	0.085

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	46	15	12	0	27	31	0	45	10
N.S.	1	1.12	0.37	0.29	0.00	0.66	0.76	0.00	1.10	0.24
time (sec)	N/A	0.153	4.043	0.497	0.000	0.072	0.453	0.000	0.253	0.065

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	39	12	0	21	31	0	32	10
N.S.	1	1.00	1.56	0.48	0.00	0.84	1.24	0.00	1.28	0.40
time (sec)	N/A	0.141	3.169	0.477	0.000	0.068	0.419	0.000	0.214	0.063

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	4	4	4	12	0	4	29	0	20	10
N.S.	1	1.00	1.00	3.00	0.00	1.00	7.25	0.00	5.00	2.50
time (sec)	N/A	0.120	10.025	0.451	0.000	0.076	0.393	0.000	0.230	0.067

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	30	12	0	32	29	0	23	10
N.S.	1	1.00	1.20	0.48	0.00	1.28	1.16	0.00	0.92	0.40
time (sec)	N/A	0.141	3.485	0.497	0.000	0.070	0.452	0.000	0.257	0.073

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	46	51	12	0	51	29	0	30	10
N.S.	1	1.12	1.24	0.29	0.00	1.24	0.71	0.00	0.73	0.24
time (sec)	N/A	0.153	4.458	0.506	0.000	0.078	0.476	0.000	0.257	0.075

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	C	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	96	17	14	0	39	29	0	50	12
N.S.	1	1.12	0.20	0.16	0.00	0.45	0.34	0.00	0.58	0.14
time (sec)	N/A	0.186	4.333	0.481	0.000	0.085	0.524	0.000	0.244	0.043

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	C	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	77	17	14	0	32	29	0	39	12
N.S.	1	1.07	0.24	0.19	0.00	0.44	0.40	0.00	0.54	0.17
time (sec)	N/A	0.170	3.753	0.459	0.000	0.079	0.476	0.000	0.258	0.029

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	C	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	48	14	0	26	29	0	28	12
N.S.	1	1.00	0.83	0.24	0.00	0.45	0.50	0.00	0.48	0.21
time (sec)	N/A	0.156	2.970	0.444	0.000	0.077	0.402	0.000	0.228	0.026

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	C	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	21	14	0	13	27	0	16	12
N.S.	1	1.00	0.49	0.33	0.00	0.30	0.63	0.00	0.37	0.28
time (sec)	N/A	0.144	10.026	0.429	0.000	0.083	0.351	0.000	0.228	0.069

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	C	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	30	14	0	38	27	0	21	12
N.S.	1	1.00	0.52	0.24	0.00	0.66	0.47	0.00	0.36	0.21
time (sec)	N/A	0.158	3.493	0.460	0.000	0.083	0.428	0.000	0.212	0.070

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	C	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	77	49	14	0	58	27	0	26	12
N.S.	1	1.07	0.68	0.19	0.00	0.81	0.38	0.00	0.36	0.17
time (sec)	N/A	0.168	4.261	0.476	0.000	0.084	0.494	0.000	0.250	0.076

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	96	101	89	102	143	198	37	0	45	37
N.S.	1	1.05	0.93	1.06	1.49	2.06	0.39	0.00	0.47	0.39
time (sec)	N/A	0.205	0.393	1.256	0.110	0.086	1.235	0.000	0.240	0.080

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	79	75	80	102	188	37	0	27	37
N.S.	1	1.05	1.00	1.07	1.36	2.51	0.49	0.00	0.36	0.49
time (sec)	N/A	0.177	0.262	0.637	0.117	0.090	0.731	0.000	0.243	0.061

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	57	76	61	68	106	36	0	11	37
N.S.	1	1.00	1.33	1.07	1.19	1.86	0.63	0.00	0.19	0.65
time (sec)	N/A	0.161	0.028	0.563	0.116	0.075	0.573	0.000	0.200	0.080

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	15	14	23	29	0	30	14
N.S.	1	1.00	1.00	0.94	0.88	1.44	1.81	0.00	1.88	0.88
time (sec)	N/A	0.130	0.200	0.486	0.035	0.074	0.404	0.000	0.244	0.060

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	29	26	31	47	126	0	50	28
N.S.	1	1.00	0.74	0.67	0.79	1.21	3.23	0.00	1.28	0.72
time (sec)	N/A	0.143	0.279	0.490	0.045	0.074	0.673	0.000	0.233	0.069

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	66	40	37	50	69	515	0	70	44
N.S.	1	1.14	0.69	0.64	0.86	1.19	8.88	0.00	1.21	0.76
time (sec)	N/A	0.166	0.374	0.492	0.028	0.073	1.450	0.000	0.222	0.077

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	93	51	48	67	91	1550	0	90	61
N.S.	1	1.21	0.66	0.62	0.87	1.18	20.13	0.00	1.17	0.79
time (sec)	N/A	0.291	0.704	0.537	0.026	0.070	2.772	0.000	0.219	0.084

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	C	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	116	124	49	0	0	0	37	0	65	37
N.S.	1	1.07	0.42	0.00	0.00	0.00	0.32	0.00	0.56	0.32
time (sec)	N/A	0.393	8.582	0.000	0.000	0.000	1.004	0.000	0.272	0.102

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	C	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	97	102	47	0	0	0	37	0	45	37
N.S.	1	1.05	0.48	0.00	0.00	0.00	0.38	0.00	0.46	0.38
time (sec)	N/A	0.346	7.599	0.000	0.000	0.000	0.636	0.000	0.209	0.091

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	C	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	80	80	46	0	0	0	37	0	27	37
N.S.	1	1.00	0.58	0.00	0.00	0.00	0.46	0.00	0.34	0.46
time (sec)	N/A	0.322	6.306	0.000	0.000	0.000	0.475	0.000	0.210	0.082

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	C	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	46	0	0	0	36	0	11	37
N.S.	1	1.00	0.75	0.00	0.00	0.00	0.59	0.00	0.18	0.61
time (sec)	N/A	0.294	6.126	0.000	0.000	0.000	0.484	0.000	0.282	0.089

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	C	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	83	83	56	0	0	0	36	0	30	37
N.S.	1	1.00	0.67	0.00	0.00	0.00	0.43	0.00	0.36	0.45
time (sec)	N/A	0.325	7.441	0.000	0.000	0.000	0.610	0.000	0.233	0.104

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	C	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	102	110	72	0	0	0	36	0	50	37
N.S.	1	1.08	0.71	0.00	0.00	0.00	0.35	0.00	0.49	0.36
time (sec)	N/A	0.363	9.477	0.000	0.000	0.000	1.000	0.000	0.226	0.121

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	C	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	121	137	85	0	0	0	36	0	70	37
N.S.	1	1.13	0.70	0.00	0.00	0.00	0.30	0.00	0.58	0.31
time (sec)	N/A	0.413	10.040	0.000	0.000	0.000	1.721	0.000	0.213	0.116

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	168	237	145	173	213	201	39	0	29	38
N.S.	1	1.41	0.86	1.03	1.27	1.20	0.23	0.00	0.17	0.23
time (sec)	N/A	0.580	0.456	3.975	0.119	0.082	0.754	0.000	0.222	0.081

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	148	214	173	138	176	146	37	0	12	38
N.S.	1	1.45	1.17	0.93	1.19	0.99	0.25	0.00	0.08	0.26
time (sec)	N/A	0.515	0.068	0.779	0.110	0.081	0.578	0.000	0.200	0.085

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	16	15	27	73	0	33	15
N.S.	1	1.00	1.00	0.94	0.88	1.59	4.29	0.00	1.94	0.88
time (sec)	N/A	0.215	0.229	0.513	0.027	0.069	0.458	0.000	0.215	0.056

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	30	27	34	48	274	0	53	30
N.S.	1	1.00	0.73	0.66	0.83	1.17	6.68	0.00	1.29	0.73
time (sec)	N/A	0.237	0.326	0.490	0.027	0.075	0.713	0.000	0.264	0.064

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	69	41	38	55	72	1066	0	75	47
N.S.	1	1.13	0.67	0.62	0.90	1.18	17.48	0.00	1.23	0.77
time (sec)	N/A	0.276	0.439	0.623	0.047	0.076	1.397	0.000	0.222	0.077

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	C	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	83	83	47	0	0	0	39	0	29	38
N.S.	1	1.00	0.57	0.00	0.00	0.00	0.47	0.00	0.35	0.46
time (sec)	N/A	0.347	6.122	0.000	0.000	0.000	0.498	0.000	0.210	0.067

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	C	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	47	0	0	0	37	0	12	38
N.S.	1	1.00	0.75	0.00	0.00	0.00	0.59	0.00	0.19	0.60
time (sec)	N/A	0.295	6.120	0.000	0.000	0.000	0.510	0.000	0.198	0.082

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	C	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	86	86	57	0	0	0	37	0	33	38
N.S.	1	1.00	0.66	0.00	0.00	0.00	0.43	0.00	0.38	0.44
time (sec)	N/A	0.324	7.513	0.000	0.000	0.000	0.602	0.000	0.251	0.103

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [7] had the largest ratio of [1.11111000000000004]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	2	2	1.00	9	0.222
2	A	2	2	1.00	9	0.222
3	A	2	2	1.00	9	0.222
4	A	1	1	1.00	7	0.143
5	A	9	8	1.49	9	0.889
6	A	10	9	1.49	9	1.000
7	A	11	10	1.49	9	1.111
8	A	3	3	1.00	12	0.250
9	A	4	4	1.05	12	0.333
10	A	3	3	1.00	12	0.250
11	A	2	2	1.00	12	0.167
12	A	3	3	1.00	12	0.250
13	A	4	4	1.08	12	0.333
14	A	3	3	1.04	11	0.273
15	A	2	2	1.00	11	0.182
16	A	1	1	1.00	11	0.091
17	A	2	2	1.00	11	0.182
18	A	3	3	1.06	11	0.273
19	A	4	4	1.18	11	0.364
20	A	3	3	1.12	11	0.273
21	A	2	2	1.00	11	0.182

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	1	1	1.00	11	0.091
23	A	2	2	1.00	11	0.182
24	A	3	3	1.12	11	0.273
25	A	4	4	1.12	9	0.444
26	A	3	3	1.07	9	0.333
27	A	2	2	1.00	9	0.222
28	A	1	1	1.00	9	0.111
29	A	2	2	1.00	9	0.222
30	A	3	3	1.07	9	0.333
31	A	7	6	1.05	11	0.545
32	A	6	5	1.05	11	0.455
33	A	5	4	1.00	11	0.364
34	A	1	1	1.00	11	0.091
35	A	2	2	1.00	11	0.182
36	A	3	3	1.14	11	0.273
37	A	4	4	1.21	11	0.364
38	A	8	7	1.07	11	0.636
39	A	7	6	1.05	11	0.545
40	A	6	5	1.00	11	0.455
41	A	5	4	1.00	11	0.364
42	A	6	5	1.00	11	0.455
43	A	7	6	1.08	11	0.545
44	A	8	7	1.13	11	0.636
45	A	11	10	1.41	12	0.833
46	A	10	9	1.45	12	0.750
47	A	1	1	1.00	12	0.083
48	A	2	2	1.00	12	0.167
49	A	3	3	1.13	12	0.250
50	A	6	5	1.00	12	0.417
51	A	5	4	1.00	12	0.333
52	A	6	5	1.00	12	0.417

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int (a + bx^4)^4 dx$	47
3.2	$\int (a + bx^4)^3 dx$	52
3.3	$\int (a + bx^4)^2 dx$	57
3.4	$\int (a + bx^4) dx$	62
3.5	$\int \frac{1}{a+cx^4} dx$	67
3.6	$\int \frac{1}{(a+cx^4)^2} dx$	76
3.7	$\int \frac{1}{(a+cx^4)^3} dx$	86
3.8	$\int \frac{1}{1+a+(-1+a)x^4} dx$	98
3.9	$\int (a - bx^4)^{3/2} dx$	106
3.10	$\int \sqrt{a - bx^4} dx$	112
3.11	$\int \frac{1}{\sqrt{a-bx^4}} dx$	118
3.12	$\int \frac{1}{(a-bx^4)^{3/2}} dx$	123
3.13	$\int \frac{1}{(a-bx^4)^{5/2}} dx$	129
3.14	$\int (a + bx^4)^{3/2} dx$	135
3.15	$\int \sqrt{a + bx^4} dx$	141
3.16	$\int \frac{1}{\sqrt{a+bx^4}} dx$	147
3.17	$\int \frac{1}{(a+bx^4)^{3/2}} dx$	152
3.18	$\int \frac{1}{(a+bx^4)^{5/2}} dx$	158
3.19	$\int (1 - x^4)^{5/2} dx$	164
3.20	$\int (1 - x^4)^{3/2} dx$	169
3.21	$\int \sqrt{1 - x^4} dx$	174
3.22	$\int \frac{1}{\sqrt{1-x^4}} dx$	179
3.23	$\int \frac{1}{(1-x^4)^{3/2}} dx$	184
3.24	$\int \frac{1}{(1-x^4)^{5/2}} dx$	189

3.25	$\int (1 + x^4)^{5/2} dx$	194
3.26	$\int (1 + x^4)^{3/2} dx$	200
3.27	$\int \sqrt{1 + x^4} dx$	206
3.28	$\int \frac{1}{\sqrt{1+x^4}} dx$	211
3.29	$\int \frac{1}{(1+x^4)^{3/2}} dx$	216
3.30	$\int \frac{1}{(1+x^4)^{5/2}} dx$	221
3.31	$\int (a + bx^4)^{7/4} dx$	227
3.32	$\int (a + bx^4)^{3/4} dx$	234
3.33	$\int \frac{1}{\sqrt[4]{a + bx^4}} dx$	241
3.34	$\int \frac{1}{(a+bx^4)^{5/4}} dx$	247
3.35	$\int \frac{1}{(a+bx^4)^{9/4}} dx$	252
3.36	$\int \frac{1}{(a+bx^4)^{13/4}} dx$	257
3.37	$\int \frac{1}{(a+bx^4)^{17/4}} dx$	263
3.38	$\int (a + bx^4)^{9/4} dx$	269
3.39	$\int (a + bx^4)^{5/4} dx$	276
3.40	$\int \sqrt[4]{a + bx^4} dx$	282
3.41	$\int \frac{1}{(a+bx^4)^{3/4}} dx$	288
3.42	$\int \frac{1}{(a+bx^4)^{7/4}} dx$	293
3.43	$\int \frac{1}{(a+bx^4)^{11/4}} dx$	299
3.44	$\int \frac{1}{(a+bx^4)^{15/4}} dx$	305
3.45	$\int (a - bx^4)^{3/4} dx$	312
3.46	$\int \frac{1}{\sqrt[4]{a - bx^4}} dx$	322
3.47	$\int \frac{1}{(a-bx^4)^{5/4}} dx$	331
3.48	$\int \frac{1}{(a-bx^4)^{9/4}} dx$	336
3.49	$\int \frac{1}{(a-bx^4)^{13/4}} dx$	341
3.50	$\int \sqrt[4]{a - bx^4} dx$	347
3.51	$\int \frac{1}{(a-bx^4)^{3/4}} dx$	353
3.52	$\int \frac{1}{(a-bx^4)^{7/4}} dx$	358

3.1 $\int (a + bx^4)^4 dx$

Optimal result	47
Mathematica [A] (verified)	47
Rubi [A] (verified)	48
Maple [A] (verified)	49
Fricas [A] (verification not implemented)	49
Sympy [A] (verification not implemented)	50
Maxima [A] (verification not implemented)	50
Giac [A] (verification not implemented)	50
Mupad [B] (verification not implemented)	51
Reduce [B] (verification not implemented)	51

Optimal result

Integrand size = 9, antiderivative size = 51

$$\int (a + bx^4)^4 dx = a^4x + \frac{4}{5}a^3bx^5 + \frac{2}{3}a^2b^2x^9 + \frac{4}{13}ab^3x^{13} + \frac{b^4x^{17}}{17}$$

output

```
a^4*x+4/5*a^3*b*x^5+2/3*a^2*b^2*x^9+4/13*a*b^3*x^13+1/17*b^4*x^17
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00

$$\int (a + bx^4)^4 dx = a^4x + \frac{4}{5}a^3bx^5 + \frac{2}{3}a^2b^2x^9 + \frac{4}{13}ab^3x^{13} + \frac{b^4x^{17}}{17}$$

input

```
Integrate[(a + b*x^4)^4,x]
```

output

```
a^4*x + (4*a^3*b*x^5)/5 + (2*a^2*b^2*x^9)/3 + (4*a*b^3*x^13)/13 + (b^4*x^17)/17
```


Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {747, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx^4)^4 dx$$

↓ 747

$$\int (a^4 + 4a^3bx^4 + 6a^2b^2x^8 + 4ab^3x^{12} + b^4x^{16}) dx$$

↓ 2009

$$a^4x + \frac{4}{5}a^3bx^5 + \frac{2}{3}a^2b^2x^9 + \frac{4}{13}ab^3x^{13} + \frac{b^4x^{17}}{17}$$

input `Int[(a + b*x^4)^4, x]`

output `a^4*x + (4*a^3*b*x^5)/5 + (2*a^2*b^2*x^9)/3 + (4*a*b^3*x^13)/13 + (b^4*x^17)/17`

Defintions of rubi rules used

rule 747 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*x^n)^p, x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.86

method	result	size
gospers	$a^4x + \frac{4}{5}a^3bx^5 + \frac{2}{3}a^2b^2x^9 + \frac{4}{13}ab^3x^{13} + \frac{1}{17}b^4x^{17}$	44
default	$a^4x + \frac{4}{5}a^3bx^5 + \frac{2}{3}a^2b^2x^9 + \frac{4}{13}ab^3x^{13} + \frac{1}{17}b^4x^{17}$	44
norman	$a^4x + \frac{4}{5}a^3bx^5 + \frac{2}{3}a^2b^2x^9 + \frac{4}{13}ab^3x^{13} + \frac{1}{17}b^4x^{17}$	44
risch	$a^4x + \frac{4}{5}a^3bx^5 + \frac{2}{3}a^2b^2x^9 + \frac{4}{13}ab^3x^{13} + \frac{1}{17}b^4x^{17}$	44
parallelrisch	$a^4x + \frac{4}{5}a^3bx^5 + \frac{2}{3}a^2b^2x^9 + \frac{4}{13}ab^3x^{13} + \frac{1}{17}b^4x^{17}$	44
orering	$\frac{x(195b^4x^{16} + 1020ab^3x^{12} + 2210a^2b^2x^8 + 2652a^3bx^4 + 3315a^4)}{3315}$	47

input `int((b*x^4+a)^4,x,method=_RETURNVERBOSE)`output `a^4*x+4/5*a^3*b*x^5+2/3*a^2*b^2*x^9+4/13*a*b^3*x^13+1/17*b^4*x^17`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.84

$$\int (a + bx^4)^4 dx = \frac{1}{17}b^4x^{17} + \frac{4}{13}ab^3x^{13} + \frac{2}{3}a^2b^2x^9 + \frac{4}{5}a^3bx^5 + a^4x$$

input `integrate((b*x^4+a)^4,x, algorithm="fricas")`output `1/17*b^4*x^17 + 4/13*a*b^3*x^13 + 2/3*a^2*b^2*x^9 + 4/5*a^3*b*x^5 + a^4*x`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.96

$$\int (a + bx^4)^4 dx = a^4x + \frac{4a^3bx^5}{5} + \frac{2a^2b^2x^9}{3} + \frac{4ab^3x^{13}}{13} + \frac{b^4x^{17}}{17}$$

input `integrate((b*x**4+a)**4,x)`output `a**4*x + 4*a**3*b*x**5/5 + 2*a**2*b**2*x**9/3 + 4*a*b**3*x**13/13 + b**4*x**17/17`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.84

$$\int (a + bx^4)^4 dx = \frac{1}{17} b^4 x^{17} + \frac{4}{13} ab^3 x^{13} + \frac{2}{3} a^2 b^2 x^9 + \frac{4}{5} a^3 b x^5 + a^4 x$$

input `integrate((b*x^4+a)^4,x, algorithm="maxima")`output `1/17*b^4*x^17 + 4/13*a*b^3*x^13 + 2/3*a^2*b^2*x^9 + 4/5*a^3*b*x^5 + a^4*x`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.84

$$\int (a + bx^4)^4 dx = \frac{1}{17} b^4 x^{17} + \frac{4}{13} ab^3 x^{13} + \frac{2}{3} a^2 b^2 x^9 + \frac{4}{5} a^3 b x^5 + a^4 x$$

input `integrate((b*x^4+a)^4,x, algorithm="giac")`output `1/17*b^4*x^17 + 4/13*a*b^3*x^13 + 2/3*a^2*b^2*x^9 + 4/5*a^3*b*x^5 + a^4*x`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.84

$$\int (a + bx^4)^4 dx = a^4 x + \frac{4a^3 b x^5}{5} + \frac{2a^2 b^2 x^9}{3} + \frac{4a b^3 x^{13}}{13} + \frac{b^4 x^{17}}{17}$$

input `int((a + b*x^4)^4,x)`output `a^4*x + (b^4*x^17)/17 + (4*a^3*b*x^5)/5 + (4*a*b^3*x^13)/13 + (2*a^2*b^2*x^9)/3`**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.90

$$\int (a + bx^4)^4 dx = \frac{x(195b^4x^{16} + 1020ab^3x^{12} + 2210a^2b^2x^8 + 2652a^3bx^4 + 3315a^4)}{3315}$$

input `int((b*x^4+a)^4,x)`output `(x*(3315*a**4 + 2652*a**3*b*x**4 + 2210*a**2*b**2*x**8 + 1020*a*b**3*x**12 + 195*b**4*x**16))/3315`

3.2 $\int (a + bx^4)^3 dx$

Optimal result	52
Mathematica [A] (verified)	52
Rubi [A] (verified)	53
Maple [A] (verified)	54
Fricas [A] (verification not implemented)	54
Sympy [A] (verification not implemented)	55
Maxima [A] (verification not implemented)	55
Giac [A] (verification not implemented)	55
Mupad [B] (verification not implemented)	56
Reduce [B] (verification not implemented)	56

Optimal result

Integrand size = 9, antiderivative size = 38

$$\int (a + bx^4)^3 dx = a^3x + \frac{3}{5}a^2bx^5 + \frac{1}{3}ab^2x^9 + \frac{b^3x^{13}}{13}$$

output

```
a^3*x+3/5*a^2*b*x^5+1/3*a*b^2*x^9+1/13*b^3*x^13
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int (a + bx^4)^3 dx = a^3x + \frac{3}{5}a^2bx^5 + \frac{1}{3}ab^2x^9 + \frac{b^3x^{13}}{13}$$

input

```
Integrate[(a + b*x^4)^3,x]
```

output

```
a^3*x + (3*a^2*b*x^5)/5 + (a*b^2*x^9)/3 + (b^3*x^13)/13
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {747, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx^4)^3 dx$$

$$\downarrow 747$$

$$\int (a^3 + 3a^2bx^4 + 3ab^2x^8 + b^3x^{12}) dx$$

$$\downarrow 2009$$

$$a^3x + \frac{3}{5}a^2bx^5 + \frac{1}{3}ab^2x^9 + \frac{b^3x^{13}}{13}$$

input `Int[(a + b*x^4)^3,x]`

output `a^3*x + (3*a^2*b*x^5)/5 + (a*b^2*x^9)/3 + (b^3*x^13)/13`

Defintions of rubi rules used

rule 747 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*x^n)^p, x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.87

method	result	size
gosper	$a^3x + \frac{3}{5}a^2bx^5 + \frac{1}{3}ab^2x^9 + \frac{1}{13}b^3x^{13}$	33
default	$a^3x + \frac{3}{5}a^2bx^5 + \frac{1}{3}ab^2x^9 + \frac{1}{13}b^3x^{13}$	33
norman	$a^3x + \frac{3}{5}a^2bx^5 + \frac{1}{3}ab^2x^9 + \frac{1}{13}b^3x^{13}$	33
risch	$a^3x + \frac{3}{5}a^2bx^5 + \frac{1}{3}ab^2x^9 + \frac{1}{13}b^3x^{13}$	33
parallelrisch	$a^3x + \frac{3}{5}a^2bx^5 + \frac{1}{3}ab^2x^9 + \frac{1}{13}b^3x^{13}$	33
orering	$\frac{x(15b^3x^{12}+65a^2bx^8+117a^2bx^4+195a^3)}{195}$	36

input `int((b*x^4+a)^3,x,method=_RETURNVERBOSE)`output `a^3*x+3/5*a^2*b*x^5+1/3*a*b^2*x^9+1/13*b^3*x^13`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int (a + bx^4)^3 dx = \frac{1}{13}b^3x^{13} + \frac{1}{3}ab^2x^9 + \frac{3}{5}a^2bx^5 + a^3x$$

input `integrate((b*x^4+a)^3,x, algorithm="fricas")`output `1/13*b^3*x^13 + 1/3*a*b^2*x^9 + 3/5*a^2*b*x^5 + a^3*x`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.89

$$\int (a + bx^4)^3 dx = a^3x + \frac{3a^2bx^5}{5} + \frac{ab^2x^9}{3} + \frac{b^3x^{13}}{13}$$

input `integrate((b*x**4+a)**3,x)`output `a**3*x + 3*a**2*b*x**5/5 + a*b**2*x**9/3 + b**3*x**13/13`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int (a + bx^4)^3 dx = \frac{1}{13}b^3x^{13} + \frac{1}{3}ab^2x^9 + \frac{3}{5}a^2bx^5 + a^3x$$

input `integrate((b*x^4+a)^3,x, algorithm="maxima")`output `1/13*b^3*x^13 + 1/3*a*b^2*x^9 + 3/5*a^2*b*x^5 + a^3*x`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int (a + bx^4)^3 dx = \frac{1}{13}b^3x^{13} + \frac{1}{3}ab^2x^9 + \frac{3}{5}a^2bx^5 + a^3x$$

input `integrate((b*x^4+a)^3,x, algorithm="giac")`output `1/13*b^3*x^13 + 1/3*a*b^2*x^9 + 3/5*a^2*b*x^5 + a^3*x`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int (a + bx^4)^3 dx = a^3 x + \frac{3a^2 b x^5}{5} + \frac{a b^2 x^9}{3} + \frac{b^3 x^{13}}{13}$$

input `int((a + b*x^4)^3,x)`output `a^3*x + (b^3*x^13)/13 + (3*a^2*b*x^5)/5 + (a*b^2*x^9)/3`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.92

$$\int (a + bx^4)^3 dx = \frac{x(15b^3x^{12} + 65ab^2x^8 + 117a^2bx^4 + 195a^3)}{195}$$

input `int((b*x^4+a)^3,x)`output `(x*(195*a**3 + 117*a**2*b*x**4 + 65*a*b**2*x**8 + 15*b**3*x**12))/195`

3.3 $\int (a + bx^4)^2 dx$

Optimal result	57
Mathematica [A] (verified)	57
Rubi [A] (verified)	58
Maple [A] (verified)	59
Fricas [A] (verification not implemented)	59
Sympy [A] (verification not implemented)	60
Maxima [A] (verification not implemented)	60
Giac [A] (verification not implemented)	60
Mupad [B] (verification not implemented)	61
Reduce [B] (verification not implemented)	61

Optimal result

Integrand size = 9, antiderivative size = 25

$$\int (a + bx^4)^2 dx = a^2x + \frac{2}{5}abx^5 + \frac{b^2x^9}{9}$$

output

```
a^2*x+2/5*a*b*x^5+1/9*b^2*x^9
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int (a + bx^4)^2 dx = a^2x + \frac{2}{5}abx^5 + \frac{b^2x^9}{9}$$

input

```
Integrate[(a + b*x^4)^2,x]
```

output

```
a^2*x + (2*a*b*x^5)/5 + (b^2*x^9)/9
```

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {747, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx^4)^2 dx$$

$$\downarrow 747$$

$$\int (a^2 + 2abx^4 + b^2x^8) dx$$

$$\downarrow 2009$$

$$a^2x + \frac{2}{5}abx^5 + \frac{b^2x^9}{9}$$

input `Int[(a + b*x^4)^2,x]`

output `a^2*x + (2*a*b*x^5)/5 + (b^2*x^9)/9`

Defintions of rubi rules used

rule 747 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*x^n)^p, x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

method	result	size
gospers	$a^2x + \frac{2}{5}abx^5 + \frac{1}{9}b^2x^9$	22
default	$a^2x + \frac{2}{5}abx^5 + \frac{1}{9}b^2x^9$	22
norman	$a^2x + \frac{2}{5}abx^5 + \frac{1}{9}b^2x^9$	22
risch	$a^2x + \frac{2}{5}abx^5 + \frac{1}{9}b^2x^9$	22
parallelrisch	$a^2x + \frac{2}{5}abx^5 + \frac{1}{9}b^2x^9$	22
orering	$\frac{x(5b^2x^8+18abx^4+45a^2)}{45}$	25

input `int((b*x^4+a)^2,x,method=_RETURNVERBOSE)`output `a^2*x+2/5*a*b*x^5+1/9*b^2*x^9`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int (a + bx^4)^2 dx = \frac{1}{9}b^2x^9 + \frac{2}{5}abx^5 + a^2x$$

input `integrate((b*x^4+a)^2,x, algorithm="fricas")`output `1/9*b^2*x^9 + 2/5*a*b*x^5 + a^2*x`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int (a + bx^4)^2 dx = a^2x + \frac{2abx^5}{5} + \frac{b^2x^9}{9}$$

input `integrate((b*x**4+a)**2,x)`output `a**2*x + 2*a*b*x**5/5 + b**2*x**9/9`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int (a + bx^4)^2 dx = \frac{1}{9}b^2x^9 + \frac{2}{5}abx^5 + a^2x$$

input `integrate((b*x^4+a)^2,x, algorithm="maxima")`output `1/9*b^2*x^9 + 2/5*a*b*x^5 + a^2*x`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int (a + bx^4)^2 dx = \frac{1}{9}b^2x^9 + \frac{2}{5}abx^5 + a^2x$$

input `integrate((b*x^4+a)^2,x, algorithm="giac")`output `1/9*b^2*x^9 + 2/5*a*b*x^5 + a^2*x`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int (a + bx^4)^2 dx = a^2 x + \frac{2abx^5}{5} + \frac{b^2 x^9}{9}$$

input `int((a + b*x^4)^2,x)`

output `a^2*x + (b^2*x^9)/9 + (2*a*b*x^5)/5`

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int (a + bx^4)^2 dx = \frac{x(5b^2x^8 + 18abx^4 + 45a^2)}{45}$$

input `int((b*x^4+a)^2,x)`

output `(x*(45*a**2 + 18*a*b*x**4 + 5*b**2*x**8))/45`

3.4 $\int (a + bx^4) dx$

Optimal result	62
Mathematica [A] (verified)	62
Rubi [A] (verified)	63
Maple [A] (verified)	64
Fricas [A] (verification not implemented)	64
Sympy [A] (verification not implemented)	65
Maxima [A] (verification not implemented)	65
Giac [A] (verification not implemented)	65
Mupad [B] (verification not implemented)	66
Reduce [B] (verification not implemented)	66

Optimal result

Integrand size = 7, antiderivative size = 12

$$\int (a + bx^4) dx = ax + \frac{bx^5}{5}$$

output `a*x+1/5*b*x^5`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int (a + bx^4) dx = ax + \frac{bx^5}{5}$$

input `Integrate[a + b*x^4,x]`

output `a*x + (b*x^5)/5`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx^4) dx$$

↓ 2009

$$ax + \frac{bx^5}{5}$$

input `Int[a + b*x^4,x]`

output `a*x + (b*x^5)/5`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

method	result	size
gospers	$ax + \frac{1}{5}bx^5$	11
default	$ax + \frac{1}{5}bx^5$	11
norman	$ax + \frac{1}{5}bx^5$	11
risch	$ax + \frac{1}{5}bx^5$	11
parallelrisch	$ax + \frac{1}{5}bx^5$	11
parts	$ax + \frac{1}{5}bx^5$	11
orering	$\frac{x(bx^4+5a)}{5}$	13

input `int(b*x^4+a,x,method=_RETURNVERBOSE)`

output `a*x+1/5*b*x^5`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int (a + bx^4) dx = \frac{1}{5}bx^5 + ax$$

input `integrate(b*x^4+a,x, algorithm="fricas")`

output `1/5*b*x^5 + a*x`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int (a + bx^4) dx = ax + \frac{bx^5}{5}$$

input `integrate(b*x**4+a,x)`

output `a*x + b*x**5/5`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int (a + bx^4) dx = \frac{1}{5} bx^5 + ax$$

input `integrate(b*x^4+a,x, algorithm="maxima")`

output `1/5*b*x^5 + a*x`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int (a + bx^4) dx = \frac{1}{5} bx^5 + ax$$

input `integrate(b*x^4+a,x, algorithm="giac")`

output `1/5*b*x^5 + a*x`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int (a + bx^4) dx = \frac{bx^5}{5} + ax$$

input `int(a + b*x^4,x)`

output `a*x + (b*x^5)/5`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int (a + bx^4) dx = \frac{x(bx^4 + 5a)}{5}$$

input `int(b*x^4+a,x)`

output `(x*(5*a + b*x**4))/5`

3.5 $\int \frac{1}{a+cx^4} dx$

Optimal result	67
Mathematica [A] (verified)	67
Rubi [A] (verified)	68
Maple [C] (verified)	71
Fricas [C] (verification not implemented)	71
Sympy [A] (verification not implemented)	72
Maxima [A] (verification not implemented)	73
Giac [B] (verification not implemented)	73
Mupad [B] (verification not implemented)	74
Reduce [B] (verification not implemented)	74

Optimal result

Integrand size = 9, antiderivative size = 134

$$\int \frac{1}{a+cx^4} dx = -\frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{cx}}{\sqrt[4]{a}}\right)}{2\sqrt{2}a^{3/4}\sqrt[4]{c}} + \frac{\arctan\left(1 + \frac{\sqrt{2}\sqrt[4]{cx}}{\sqrt[4]{a}}\right)}{2\sqrt{2}a^{3/4}\sqrt[4]{c}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{a}\sqrt[4]{cx}}{\sqrt{a+\sqrt{cx^2}}}\right)}{2\sqrt{2}a^{3/4}\sqrt[4]{c}}$$

```
output 1/4*arctan(-1+2^(1/2)*c^(1/4)*x/a^(1/4))*2^(1/2)/a^(3/4)/c^(1/4)+1/4*arctan(1+2^(1/2)*c^(1/4)*x/a^(1/4))*2^(1/2)/a^(3/4)/c^(1/4)+1/4*arctanh(2^(1/2)*a^(1/4)*c^(1/4)*x/(a^(1/2)+c^(1/2)*x^2))*2^(1/2)/a^(3/4)/c^(1/4)
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.00

$$\int \frac{1}{a+cx^4} dx = \frac{-2 \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{cx}}{\sqrt[4]{a}}\right) + 2 \arctan\left(1 + \frac{\sqrt{2}\sqrt[4]{cx}}{\sqrt[4]{a}}\right) - \log(\sqrt{a} - \sqrt{2}\sqrt[4]{a}\sqrt[4]{cx} + \sqrt{cx^2}) + \log(\sqrt{a} + \sqrt{2}\sqrt[4]{a}\sqrt[4]{cx} + \sqrt{cx^2})}{4\sqrt{2}a^{3/4}\sqrt[4]{c}}$$

```
input Integrate[(a + c*x^4)^(-1),x]
```

output

$$(-2*\text{ArcTan}[1 - (\text{Sqrt}[2]*c^{(1/4)}*x)/a^{(1/4)}] + 2*\text{ArcTan}[1 + (\text{Sqrt}[2]*c^{(1/4)}*x)/a^{(1/4)}] - \text{Log}[\text{Sqrt}[a] - \text{Sqrt}[2]*a^{(1/4)}*c^{(1/4)}*x + \text{Sqrt}[c]*x^2] + \text{Log}[\text{Sqrt}[a] + \text{Sqrt}[2]*a^{(1/4)}*c^{(1/4)}*x + \text{Sqrt}[c]*x^2])/(4*\text{Sqrt}[2]*a^{(3/4)}*c^{(1/4)})$$
Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.49, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.889$, Rules used = {755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{a + cx^4} dx \\
 & \quad \downarrow 755 \\
 & \frac{\int \frac{\sqrt{a}-\sqrt{cx^2}}{cx^4+a} dx}{2\sqrt{a}} + \frac{\int \frac{\sqrt{cx^2}+\sqrt{a}}{cx^4+a} dx}{2\sqrt{a}} \\
 & \quad \downarrow 1476 \\
 & \frac{\int \frac{1}{x^2 - \frac{\sqrt{2}\sqrt[4]{a}x + \sqrt{c}}{\sqrt{c}}} dx}{2\sqrt{a}} + \frac{\int \frac{1}{x^2 + \frac{\sqrt{2}\sqrt[4]{a}x + \sqrt{c}}{\sqrt{c}}} dx}{2\sqrt{a}} + \frac{\int \frac{\sqrt{a}-\sqrt{cx^2}}{cx^4+a} dx}{2\sqrt{a}} \\
 & \quad \downarrow 1082 \\
 & \frac{\int \frac{\sqrt{a}-\sqrt{cx^2}}{cx^4+a} dx}{2\sqrt{a}} + \frac{\int \frac{1}{\left(1 - \frac{\sqrt{2}\sqrt[4]{c}x}{\sqrt{a}}\right)^2 - 1} d\left(1 - \frac{\sqrt{2}\sqrt[4]{c}x}{\sqrt{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} - \frac{\int \frac{1}{\left(\frac{\sqrt{2}\sqrt[4]{c}x}{\sqrt{a}} + 1\right)^2 - 1} d\left(\frac{\sqrt{2}\sqrt[4]{c}x}{\sqrt{a}} + 1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} \\
 & \quad \downarrow 217 \\
 & \frac{\int \frac{\sqrt{a}-\sqrt{cx^2}}{cx^4+a} dx}{2\sqrt{a}} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{c}x}{\sqrt{a}} + 1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} - \frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{c}x}{\sqrt{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} \\
 & \quad \downarrow 1479
 \end{aligned}$$

$$\begin{aligned}
& \frac{\int -\frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{c}x}{\sqrt[4]{c}\left(x^2-\frac{\sqrt{2}\sqrt[4]{a}x+\sqrt{a}}{\sqrt[4]{c}}\right)}dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} - \frac{\int -\frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{c}x+\sqrt[4]{a}\right)}{\sqrt[4]{c}\left(x^2+\frac{\sqrt{2}\sqrt[4]{a}x+\sqrt{a}}{\sqrt[4]{c}}\right)}dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} + \\
& \frac{2\sqrt{a}}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{c}x}{\sqrt[4]{a}}+1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{c}x}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} \\
& \qquad \qquad \qquad \downarrow 25 \\
& \frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{c}x}{\sqrt[4]{c}\left(x^2-\frac{\sqrt{2}\sqrt[4]{a}x+\sqrt{a}}{\sqrt[4]{c}}\right)}dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} + \frac{\int \frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{c}x+\sqrt[4]{a}\right)}{\sqrt[4]{c}\left(x^2+\frac{\sqrt{2}\sqrt[4]{a}x+\sqrt{a}}{\sqrt[4]{c}}\right)}dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{c}x}{\sqrt[4]{a}}+1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{c}x}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} \\
& \qquad \qquad \qquad \downarrow 27 \\
& \frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{c}x}{x^2-\frac{\sqrt{2}\sqrt[4]{a}x+\sqrt{a}}{\sqrt[4]{c}}}dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} + \frac{\int -\frac{\sqrt{2}\sqrt[4]{c}x+\sqrt[4]{a}}{x^2+\frac{\sqrt{2}\sqrt[4]{a}x+\sqrt{a}}{\sqrt[4]{c}}}dx}{2\sqrt[4]{a}\sqrt[4]{c}} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{c}x}{\sqrt[4]{a}}+1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{c}x}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} \\
& \qquad \qquad \qquad \downarrow 1103 \\
& \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{c}x}{\sqrt[4]{a}}+1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{c}x}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} + \\
& \frac{\log\left(\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}x+\sqrt{a}+\sqrt{cx^2}\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} - \frac{\log\left(-\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}x+\sqrt{a}+\sqrt{cx^2}\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} \\
& \qquad \qquad \qquad \downarrow \\
& \frac{2\sqrt{a}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}}
\end{aligned}$$

input `Int[(a + c*x^4)^(-1),x]`

output `(-(ArcTan[1 - (Sqrt[2]*c^(1/4)*x)/a^(1/4)]/(Sqrt[2]*a^(1/4)*c^(1/4))) + ArcTan[1 + (Sqrt[2]*c^(1/4)*x)/a^(1/4)]/(Sqrt[2]*a^(1/4)*c^(1/4)))/(2*Sqrt[a]) + (-1/2*Log[Sqrt[a] - Sqrt[2]*a^(1/4)*c^(1/4)*x + Sqrt[c]*x^2]/(Sqrt[2]*a^(1/4)*c^(1/4)) + Log[Sqrt[a] + Sqrt[2]*a^(1/4)*c^(1/4)*x + Sqrt[c]*x^2]/(2*Sqrt[2]*a^(1/4)*c^(1/4)))/(2*Sqrt[a])`

Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.48 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.20

method	result	size
risch	$\frac{\sum_{-R=\text{RootOf}(cZ^4+a)} \frac{\ln(x-R)}{-R^3}}{4c}$	27
default	$\frac{\left(\frac{a}{c}\right)^{\frac{1}{4}}\sqrt{2} \left(\ln\left(\frac{x^2+\left(\frac{a}{c}\right)^{\frac{1}{4}}x\sqrt{2}+\sqrt{\frac{a}{c}}}{x^2-\left(\frac{a}{c}\right)^{\frac{1}{4}}x\sqrt{2}+\sqrt{\frac{a}{c}}}\right) + 2\arctan\left(\frac{\sqrt{2}x}{\left(\frac{a}{c}\right)^{\frac{1}{4}}}\right) + 2\arctan\left(\frac{\sqrt{2}x}{\left(\frac{a}{c}\right)^{\frac{1}{4}}}-1\right) \right)}{8a}$	102

input `int(1/(c*x^4+a),x,method=_RETURNVERBOSE)`

output `1/4/c*sum(1/_R^3*ln(x-_R),_R=RootOf(_Z^4*c+a))`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.84

$$\begin{aligned} \int \frac{1}{a + cx^4} dx &= \frac{1}{4} \left(-\frac{1}{a^3c} \right)^{\frac{1}{4}} \log \left(a \left(-\frac{1}{a^3c} \right)^{\frac{1}{4}} + x \right) \\ &+ \frac{1}{4} i \left(-\frac{1}{a^3c} \right)^{\frac{1}{4}} \log \left(i a \left(-\frac{1}{a^3c} \right)^{\frac{1}{4}} + x \right) \\ &- \frac{1}{4} i \left(-\frac{1}{a^3c} \right)^{\frac{1}{4}} \log \left(-i a \left(-\frac{1}{a^3c} \right)^{\frac{1}{4}} + x \right) \\ &- \frac{1}{4} \left(-\frac{1}{a^3c} \right)^{\frac{1}{4}} \log \left(-a \left(-\frac{1}{a^3c} \right)^{\frac{1}{4}} + x \right) \end{aligned}$$

input `integrate(1/(c*x^4+a),x, algorithm="fricas")`

output `1/4*(-1/(a^3*c))^(1/4)*log(a*(-1/(a^3*c))^(1/4) + x) + 1/4*I*(-1/(a^3*c))^(1/4)*log(I*a*(-1/(a^3*c))^(1/4) + x) - 1/4*I*(-1/(a^3*c))^(1/4)*log(-I*a*(-1/(a^3*c))^(1/4) + x) - 1/4*(-1/(a^3*c))^(1/4)*log(-a*(-1/(a^3*c))^(1/4) + x)`

Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.15

$$\int \frac{1}{a + cx^4} dx = \text{RootSum}(256t^4a^3c + 1, (t \mapsto t \log(4ta + x)))$$

input `integrate(1/(c*x**4+a),x)`

output `RootSum(256*_t**4*a**3*c + 1, Lambda(_t, _t*log(4*_t*a + x)))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.26

$$\int \frac{1}{a + cx^4} dx = \frac{\sqrt{2} \arctan\left(\frac{\sqrt{2}(2\sqrt{cx} + \sqrt{2}a^{\frac{1}{4}}c^{\frac{1}{4}})}{2\sqrt{a}\sqrt{c}}\right)}{4\sqrt{a}\sqrt{\sqrt{a}\sqrt{c}}} + \frac{\sqrt{2} \arctan\left(\frac{\sqrt{2}(2\sqrt{cx} - \sqrt{2}a^{\frac{1}{4}}c^{\frac{1}{4}})}{2\sqrt{a}\sqrt{c}}\right)}{4\sqrt{a}\sqrt{\sqrt{a}\sqrt{c}}}$$

$$+ \frac{\sqrt{2} \log\left(\sqrt{cx^2} + \sqrt{2}a^{\frac{1}{4}}c^{\frac{1}{4}}x + \sqrt{a}\right)}{8a^{\frac{3}{4}}c^{\frac{1}{4}}} - \frac{\sqrt{2} \log\left(\sqrt{cx^2} - \sqrt{2}a^{\frac{1}{4}}c^{\frac{1}{4}}x + \sqrt{a}\right)}{8a^{\frac{3}{4}}c^{\frac{1}{4}}}$$

input `integrate(1/(c*x^4+a),x, algorithm="maxima")`

output `1/4*sqrt(2)*arctan(1/2*sqrt(2)*(2*sqrt(c)*x + sqrt(2)*a^(1/4)*c^(1/4))/sqrt(a)*sqrt(c))/sqrt(a)*sqrt(sqrt(a)*sqrt(c)) + 1/4*sqrt(2)*arctan(1/2*sqrt(2)*(2*sqrt(c)*x - sqrt(2)*a^(1/4)*c^(1/4))/sqrt(a)*sqrt(c))/sqrt(a)*sqrt(sqrt(a)*sqrt(c)) + 1/8*sqrt(2)*log(sqrt(c)*x^2 + sqrt(2)*a^(1/4)*c^(1/4)*x + sqrt(a))/(a^(3/4)*c^(1/4)) - 1/8*sqrt(2)*log(sqrt(c)*x^2 - sqrt(2)*a^(1/4)*c^(1/4)*x + sqrt(a))/(a^(3/4)*c^(1/4))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 179 vs. 2(87) = 174.

Time = 0.12 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.34

$$\int \frac{1}{a + cx^4} dx = \frac{\sqrt{2}(ac^3)^{\frac{1}{4}} \arctan\left(\frac{\sqrt{2}\left(2x + \sqrt{2}\left(\frac{a}{c}\right)^{\frac{1}{4}}\right)}{2\left(\frac{a}{c}\right)^{\frac{1}{4}}}\right)}{4ac}$$

$$+ \frac{\sqrt{2}(ac^3)^{\frac{1}{4}} \arctan\left(\frac{\sqrt{2}\left(2x - \sqrt{2}\left(\frac{a}{c}\right)^{\frac{1}{4}}\right)}{2\left(\frac{a}{c}\right)^{\frac{1}{4}}}\right)}{4ac}$$

$$+ \frac{\sqrt{2}(ac^3)^{\frac{1}{4}} \log\left(x^2 + \sqrt{2}x\left(\frac{a}{c}\right)^{\frac{1}{4}} + \sqrt{\frac{a}{c}}\right)}{8ac}$$

$$- \frac{\sqrt{2}(ac^3)^{\frac{1}{4}} \log\left(x^2 - \sqrt{2}x\left(\frac{a}{c}\right)^{\frac{1}{4}} + \sqrt{\frac{a}{c}}\right)}{8ac}$$

input `integrate(1/(c*x^4+a),x, algorithm="giac")`

output `1/4*sqrt(2)*(a*c^3)^(1/4)*arctan(1/2*sqrt(2)*(2*x + sqrt(2)*(a/c)^(1/4))/(a/c)^(1/4))/(a*c) + 1/4*sqrt(2)*(a*c^3)^(1/4)*arctan(1/2*sqrt(2)*(2*x - sqrt(2)*(a/c)^(1/4))/(a/c)^(1/4))/(a*c) + 1/8*sqrt(2)*(a*c^3)^(1/4)*log(x^2 + sqrt(2)*x*(a/c)^(1/4) + sqrt(a/c))/(a*c) - 1/8*sqrt(2)*(a*c^3)^(1/4)*log(x^2 - sqrt(2)*x*(a/c)^(1/4) + sqrt(a/c))/(a*c)`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.25

$$\int \frac{1}{a + cx^4} dx = -\frac{\operatorname{atan}\left(\frac{c^{1/4}x}{(-a)^{1/4}}\right) + \operatorname{atanh}\left(\frac{c^{1/4}x}{(-a)^{1/4}}\right)}{2(-a)^{3/4}c^{1/4}}$$

input `int(1/(a + c*x^4),x)`

output `-(atan((c^(1/4)*x)/(-a)^(1/4)) + atanh((c^(1/4)*x)/(-a)^(1/4)))/(2*(-a)^(3/4)*c^(1/4))`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.84

$$\int \frac{1}{a + cx^4} dx = \frac{\sqrt{2} \left(-2 \operatorname{atan}\left(\frac{c^{1/4}a^{1/4}\sqrt{2}-2\sqrt{cx}}{c^{1/4}a^{1/4}\sqrt{2}}\right) + 2 \operatorname{atan}\left(\frac{c^{1/4}a^{1/4}\sqrt{2}+2\sqrt{cx}}{c^{1/4}a^{1/4}\sqrt{2}}\right) - \log\left(-c^{1/4}a^{1/4}\sqrt{2}x + \sqrt{a} + \sqrt{cx^2}\right) + \log\left(c^{1/4}a^{1/4}\sqrt{2}x + \sqrt{a} + \sqrt{cx^2}\right) \right)}{8c^{1/4}a^{3/4}}$$

input `int(1/(c*x^4+a),x)`

output

```
(c**(3/4)*a**(1/4)*sqrt(2)*( - 2*atan((c**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(c)*x)/(c**(1/4)*a**(1/4)*sqrt(2))) + 2*atan((c**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(c)*x)/(c**(1/4)*a**(1/4)*sqrt(2))) - log( - c**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(c)*x**2) + log(c**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(c)*x**2)))/(8*a*c)
```

3.6 $\int \frac{1}{(a+cx^4)^2} dx$

Optimal result	76
Mathematica [A] (verified)	77
Rubi [A] (verified)	77
Maple [C] (verified)	82
Fricas [C] (verification not implemented)	82
Sympy [A] (verification not implemented)	83
Maxima [A] (verification not implemented)	83
Giac [A] (verification not implemented)	84
Mupad [B] (verification not implemented)	84
Reduce [B] (verification not implemented)	85

Optimal result

Integrand size = 9, antiderivative size = 151

$$\int \frac{1}{(a+cx^4)^2} dx = \frac{x}{4a(a+cx^4)} - \frac{3 \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{cx}}{\sqrt[4]{a}}\right)}{8\sqrt{2}a^{7/4}\sqrt[4]{c}} + \frac{3 \arctan\left(1 + \frac{\sqrt{2}\sqrt[4]{cx}}{\sqrt[4]{a}}\right)}{8\sqrt{2}a^{7/4}\sqrt[4]{c}} + \frac{3 \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{a}\sqrt[4]{cx}}{\sqrt{a+\sqrt{cx^2}}}\right)}{8\sqrt{2}a^{7/4}\sqrt[4]{c}}$$

output

```
1/4*x/a/(c*x^4+a)+3/16*arctan(-1+2^(1/2)*c^(1/4)*x/a^(1/4))*2^(1/2)/a^(7/4)
)/c^(1/4)+3/16*arctan(1+2^(1/2)*c^(1/4)*x/a^(1/4))*2^(1/2)/a^(7/4)/c^(1/4)
+3/16*arctanh(2^(1/2)*a^(1/4)*c^(1/4)*x/(a^(1/2)+c^(1/2)*x^2))*2^(1/2)/a^(
7/4)/c^(1/4)
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.21

$$\int \frac{1}{(a + cx^4)^2} dx$$

$$= \frac{\frac{8a^{3/4}x}{a+cx^4} - \frac{6\sqrt{2} \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{c}x}{\sqrt[4]{a}}\right)}{\sqrt[4]{c}} + \frac{6\sqrt{2} \arctan\left(1 + \frac{\sqrt{2}\sqrt[4]{c}x}{\sqrt[4]{a}}\right)}{\sqrt[4]{c}} - \frac{3\sqrt{2} \log\left(\sqrt{a} - \sqrt{2}\sqrt[4]{a}\sqrt[4]{cx + \sqrt{cx^2}}\right)}{\sqrt[4]{c}} + \frac{3\sqrt{2} \log\left(\sqrt{a} + \sqrt{2}\sqrt[4]{a}\sqrt[4]{cx + \sqrt{cx^2}}\right)}{\sqrt[4]{c}}}{32a^{7/4}}$$

input `Integrate[(a + c*x^4)^(-2), x]`

output $((8*a^{(3/4)*x})/(a + c*x^4) - (6*\text{Sqrt}[2]*\text{ArcTan}[1 - (\text{Sqrt}[2]*c^{(1/4)*x})/a^{(1/4)}])/c^{(1/4)} + (6*\text{Sqrt}[2]*\text{ArcTan}[1 + (\text{Sqrt}[2]*c^{(1/4)*x})/a^{(1/4)}])/c^{(1/4)} - (3*\text{Sqrt}[2]*\text{Log}[\text{Sqrt}[a] - \text{Sqrt}[2]*a^{(1/4)*c^{(1/4)*x} + \text{Sqrt}[c]*x^2])/c^{(1/4)} + (3*\text{Sqrt}[2]*\text{Log}[\text{Sqrt}[a] + \text{Sqrt}[2]*a^{(1/4)*c^{(1/4)*x} + \text{Sqrt}[c]*x^2])/c^{(1/4)})/(32*a^{(7/4)})$

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.49, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {749, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + cx^4)^2} dx$$

$$\downarrow 749$$

$$\frac{3 \int \frac{1}{cx^4 + a} dx}{4a} + \frac{x}{4a(a + cx^4)}$$

$$\downarrow 755$$

$$\begin{aligned}
 & \frac{3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{cx^2}}{cx^4+a} dx}{2\sqrt{a}} + \frac{\int \frac{\sqrt{cx^2}+\sqrt{a}}{cx^4+a} dx}{2\sqrt{a}} \right)}{4a} + \frac{x}{4a(a+cx^4)} \\
 & \quad \downarrow 1476 \\
 & \frac{3 \left(\frac{\int \frac{1}{x^2 - \frac{\sqrt{2}\sqrt[4]{a}x + \sqrt{a}}{\sqrt{c}}} dx}{2\sqrt{c}} + \frac{\int \frac{1}{x^2 + \frac{\sqrt{2}\sqrt[4]{a}x + \sqrt{a}}{\sqrt{c}}} dx}{2\sqrt{c}} + \frac{\int \frac{\sqrt{a}-\sqrt{cx^2}}{cx^4+a} dx}{2\sqrt{a}} \right)}{4a} + \frac{x}{4a(a+cx^4)} \\
 & \quad \downarrow 1082 \\
 & \frac{3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{cx^2}}{cx^4+a} dx}{2\sqrt{a}} + \frac{\int \frac{1}{\left(1 - \frac{\sqrt{2}\sqrt[4]{cx}}{\sqrt{a}}\right)^2 - 1} d\left(1 - \frac{\sqrt{2}\sqrt[4]{cx}}{\sqrt{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} - \frac{\int \frac{1}{\left(\frac{\sqrt{2}\sqrt[4]{cx}}{\sqrt{a}} + 1\right)^2 - 1} d\left(\frac{\sqrt{2}\sqrt[4]{cx}}{\sqrt{a}} + 1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} \right)}{2\sqrt{a}} \right)}{4a} + \frac{x}{4a(a+cx^4)} \\
 & \quad \downarrow 217 \\
 & \frac{3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{cx^2}}{cx^4+a} dx}{2\sqrt{a}} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{cx}}{\sqrt{a}} + 1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} - \frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{cx}}{\sqrt{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} \right)}{4a} + \frac{x}{4a(a+cx^4)} \\
 & \quad \downarrow 1479
 \end{aligned}$$

$$3 \left(\frac{\int -\frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{c} x}{\sqrt[4]{c} \left(x^2 - \frac{\sqrt{2} \sqrt[4]{a} x + \sqrt{a}}{\sqrt[4]{c}} \right)} dx}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} - \frac{\int -\frac{\sqrt{2} (\sqrt{2} \sqrt[4]{c} x + \sqrt[4]{a})}{\sqrt[4]{c} \left(x^2 + \frac{\sqrt{2} \sqrt[4]{a} x + \sqrt{a}}{\sqrt[4]{c}} \right)} dx}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} + \frac{\arctan \left(\frac{\sqrt{2} \sqrt[4]{c} x + 1}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[4]{c} x}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} \right) +$$

$$\frac{4a}{x} \\ 4a(a + cx^4)$$

25

$$3 \left(\frac{\int \frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{c} x}{\sqrt[4]{c} \left(x^2 - \frac{\sqrt{2} \sqrt[4]{a} x + \sqrt{a}}{\sqrt[4]{c}} \right)} dx}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} + \frac{\int \frac{\sqrt{2} (\sqrt{2} \sqrt[4]{c} x + \sqrt[4]{a})}{\sqrt[4]{c} \left(x^2 + \frac{\sqrt{2} \sqrt[4]{a} x + \sqrt{a}}{\sqrt[4]{c}} \right)} dx}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} + \frac{\arctan \left(\frac{\sqrt{2} \sqrt[4]{c} x + 1}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[4]{c} x}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} \right) +$$

$$\frac{4a}{x} \\ 4a(a + cx^4)$$

27

$$3 \left(\frac{\int \frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{c} x}{x^2 - \frac{\sqrt{2} \sqrt[4]{a} x + \sqrt{a}}{\sqrt[4]{c}}} dx}{2\sqrt{2} \sqrt[4]{a} \sqrt{c}} + \frac{\int \frac{\sqrt{2} \sqrt[4]{c} x + \sqrt[4]{a}}{x^2 + \frac{\sqrt{2} \sqrt[4]{a} x + \sqrt{a}}{\sqrt[4]{c}}} dx}{2 \sqrt[4]{a} \sqrt{c}} + \frac{\arctan \left(\frac{\sqrt{2} \sqrt[4]{c} x + 1}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[4]{c} x}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} \right) +$$

$$\frac{4a}{x} \\ 4a(a + cx^4)$$

1103

$$3 \left(\frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{cx}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} - \frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{cx}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}}}{2\sqrt{a}} + \frac{\frac{\log\left(\sqrt{2}\sqrt[4]{a}\sqrt[4]{cx} + \sqrt{a} + \sqrt{cx^2}\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}} - \frac{\log\left(-\sqrt{2}\sqrt[4]{a}\sqrt[4]{cx} + \sqrt{a} + \sqrt{cx^2}\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}}}{2\sqrt{a}} \right) + \frac{4a}{4a(a+cx^4)x}$$

input `Int[(a + c*x^4)^(-2),x]`

output `x/(4*a*(a + c*x^4)) + (3*((-(ArcTan[1 - (Sqrt[2]*c^(1/4)*x)/a^(1/4)]/(Sqrt[2]*a^(1/4)*c^(1/4))) + ArcTan[1 + (Sqrt[2]*c^(1/4)*x)/a^(1/4)]/(Sqrt[2]*a^(1/4)*c^(1/4)))/(2*Sqrt[a]) + (-1/2*Log[Sqrt[a] - Sqrt[2]*a^(1/4)*c^(1/4)*x + Sqrt[c]*x^2]/(Sqrt[2]*a^(1/4)*c^(1/4)) + Log[Sqrt[a] + Sqrt[2]*a^(1/4)*c^(1/4)*x + Sqrt[c]*x^2]/(2*Sqrt[2]*a^(1/4)*c^(1/4)))/(2*Sqrt[a]))/(4*a)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.44 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.30

method	result	size
risch	$\frac{x}{4a(cx^4+a)} + \frac{3 \left(\sum_{R=\text{RootOf}(cZ^4+a)} \frac{\ln(x-R)}{-R^3} \right)}{16ac}$	46
default	$\frac{x}{4a(cx^4+a)} + \frac{3 \left(\frac{a}{c} \right)^{\frac{1}{4}} \sqrt{2} \left(\ln \left(\frac{x^2 + \left(\frac{a}{c} \right)^{\frac{1}{4}} x \sqrt{2} + \sqrt{\frac{a}{c}}}{x^2 - \left(\frac{a}{c} \right)^{\frac{1}{4}} x \sqrt{2} + \sqrt{\frac{a}{c}}} \right) + 2 \arctan \left(\frac{\sqrt{2} x}{\left(\frac{a}{c} \right)^{\frac{1}{4}}} + 1 \right) + 2 \arctan \left(\frac{\sqrt{2} x}{\left(\frac{a}{c} \right)^{\frac{1}{4}}} - 1 \right) \right)}{32a^2}$	118

input `int(1/(c*x^4+a)^2,x,method=_RETURNVERBOSE)`

output `1/4*x/a/(c*x^4+a)+3/16/a/c*sum(1/_R^3*ln(x-_R),_R=RootOf(_Z^4*c+a))`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.21

$$\int \frac{1}{(a+cx^4)^2} dx$$

$$= \frac{3(acx^4+a^2)\left(-\frac{1}{a^7c}\right)^{\frac{1}{4}} \log\left(a^2\left(-\frac{1}{a^7c}\right)^{\frac{1}{4}}+x\right) - 3(-iacx^4-ia^2)\left(-\frac{1}{a^7c}\right)^{\frac{1}{4}} \log\left(ia^2\left(-\frac{1}{a^7c}\right)^{\frac{1}{4}}+x\right) - 3(iacx^4+ia^2)\left(-\frac{1}{a^7c}\right)^{\frac{1}{4}} \log\left(-ia^2\left(-\frac{1}{a^7c}\right)^{\frac{1}{4}}+x\right) - 3(-iacx^4-ia^2)\left(-\frac{1}{a^7c}\right)^{\frac{1}{4}} \log\left(-ia^2\left(-\frac{1}{a^7c}\right)^{\frac{1}{4}}+x\right) + 4x}{16(acx^4+a^2)}$$

input `integrate(1/(c*x^4+a)^2,x, algorithm="fricas")`

output `1/16*(3*(a*c*x^4 + a^2)*(-1/(a^7*c))^(1/4)*log(a^2*(-1/(a^7*c))^(1/4) + x) - 3*(-I*a*c*x^4 - I*a^2)*(-1/(a^7*c))^(1/4)*log(I*a^2*(-1/(a^7*c))^(1/4) + x) - 3*(I*a*c*x^4 + I*a^2)*(-1/(a^7*c))^(1/4)*log(-I*a^2*(-1/(a^7*c))^(1/4) + x) - 3*(a*c*x^4 + a^2)*(-1/(a^7*c))^(1/4)*log(-a^2*(-1/(a^7*c))^(1/4) + x) + 4*x)/(a*c*x^4 + a^2)`

Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.26

$$\int \frac{1}{(a + cx^4)^2} dx = \frac{x}{4a^2 + 4acx^4} + \text{RootSum} \left(65536t^4 a^7 c + 81, \left(t \mapsto t \log \left(\frac{16ta^2}{3} + x \right) \right) \right)$$

input `integrate(1/(c*x**4+a)**2,x)`output `x/(4*a**2 + 4*a*c*x**4) + RootSum(65536*_t**4*a**7*c + 81, Lambda(_t, _t*log(16*_t*a**2/3 + x)))`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.25

$$\int \frac{1}{(a + cx^4)^2} dx = \frac{x}{4(acx^4 + a^2)} + \frac{3}{32a} \left(\frac{2\sqrt{2} \arctan \left(\frac{\sqrt{2} (2\sqrt{cx} + \sqrt{2a} \frac{1}{4} c \frac{1}{4})}{2\sqrt{a}\sqrt{c}} \right)}{\sqrt{a}\sqrt{a}\sqrt{c}} + \frac{2\sqrt{2} \arctan \left(\frac{\sqrt{2} (2\sqrt{cx} - \sqrt{2a} \frac{1}{4} c \frac{1}{4})}{2\sqrt{a}\sqrt{c}} \right)}{\sqrt{a}\sqrt{a}\sqrt{c}} + \frac{\sqrt{2} \log(\sqrt{cx^2 + \sqrt{2a} \frac{1}{4} c \frac{1}{4} x + \sqrt{a}})}{a^{\frac{3}{4}} c^{\frac{1}{4}}} - \frac{\sqrt{2} \log(\sqrt{cx^2 - \sqrt{2a} \frac{1}{4} c \frac{1}{4} x + \sqrt{a}})}{a^{\frac{3}{4}} c^{\frac{1}{4}}} \right)$$

input `integrate(1/(c*x^4+a)^2,x, algorithm="maxima")`output `1/4*x/(a*c*x^4 + a^2) + 3/32*(2*sqrt(2)*arctan(1/2*sqrt(2)*(2*sqrt(c)*x + sqrt(2)*a^(1/4)*c^(1/4))/sqrt(sqrt(a)*sqrt(c)))/(sqrt(a)*sqrt(sqrt(a)*sqrt(c))) + 2*sqrt(2)*arctan(1/2*sqrt(2)*(2*sqrt(c)*x - sqrt(2)*a^(1/4)*c^(1/4))/sqrt(sqrt(a)*sqrt(c)))/(sqrt(a)*sqrt(sqrt(a)*sqrt(c))) + sqrt(2)*log(sqrt(c)*x^2 + sqrt(2)*a^(1/4)*c^(1/4)*x + sqrt(a))/(a^(3/4)*c^(1/4)) - sqrt(2)*log(sqrt(c)*x^2 - sqrt(2)*a^(1/4)*c^(1/4)*x + sqrt(a))/(a^(3/4)*c^(1/4))/a`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.28

$$\int \frac{1}{(a + cx^4)^2} dx = \frac{x}{4(cx^4 + a)a} + \frac{3\sqrt{2}(ac^3)^{\frac{1}{4}} \arctan\left(\frac{\sqrt{2}\left(2x + \sqrt{2}\left(\frac{a}{c}\right)^{\frac{1}{4}}\right)}{2\left(\frac{a}{c}\right)^{\frac{1}{4}}}\right)}{16a^2c}$$

$$+ \frac{3\sqrt{2}(ac^3)^{\frac{1}{4}} \arctan\left(\frac{\sqrt{2}\left(2x - \sqrt{2}\left(\frac{a}{c}\right)^{\frac{1}{4}}\right)}{2\left(\frac{a}{c}\right)^{\frac{1}{4}}}\right)}{16a^2c}$$

$$+ \frac{3\sqrt{2}(ac^3)^{\frac{1}{4}} \log\left(x^2 + \sqrt{2}x\left(\frac{a}{c}\right)^{\frac{1}{4}} + \sqrt{\frac{a}{c}}\right)}{32a^2c}$$

$$- \frac{3\sqrt{2}(ac^3)^{\frac{1}{4}} \log\left(x^2 - \sqrt{2}x\left(\frac{a}{c}\right)^{\frac{1}{4}} + \sqrt{\frac{a}{c}}\right)}{32a^2c}$$

input `integrate(1/(c*x^4+a)^2,x, algorithm="giac")`output `1/4*x/((c*x^4 + a)*a) + 3/16*sqrt(2)*(a*c^3)^(1/4)*arctan(1/2*sqrt(2)*(2*x + sqrt(2)*(a/c)^(1/4))/(a/c)^(1/4))/(a^2*c) + 3/16*sqrt(2)*(a*c^3)^(1/4)*arctan(1/2*sqrt(2)*(2*x - sqrt(2)*(a/c)^(1/4))/(a/c)^(1/4))/(a^2*c) + 3/32*sqrt(2)*(a*c^3)^(1/4)*log(x^2 + sqrt(2)*x*(a/c)^(1/4) + sqrt(a/c))/(a^2*c) - 3/32*sqrt(2)*(a*c^3)^(1/4)*log(x^2 - sqrt(2)*x*(a/c)^(1/4) + sqrt(a/c))/(a^2*c)`**Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.38

$$\int \frac{1}{(a + cx^4)^2} dx = \frac{x}{4a(cx^4 + a)} + \frac{3 \operatorname{atan}\left(\frac{c^{1/4}x}{(-a)^{1/4}}\right)}{8(-a)^{7/4}c^{1/4}} + \frac{3 \operatorname{atanh}\left(\frac{c^{1/4}x}{(-a)^{1/4}}\right)}{8(-a)^{7/4}c^{1/4}}$$

input `int(1/(a + c*x^4)^2,x)`

output

$$\frac{x}{4a(a + cx^4)} + \frac{3 \operatorname{atan}\left(\frac{c^{1/4}x}{(-a)^{1/4}}\right)}{8(-a)^{7/4}c^{1/4}} + \frac{3 \operatorname{atanh}\left(\frac{c^{1/4}x}{(-a)^{1/4}}\right)}{8(-a)^{7/4}c^{1/4}}$$

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 305, normalized size of antiderivative = 2.02

$$\int \frac{1}{(a + cx^4)^2} dx$$

$$= \frac{-6c^{\frac{3}{4}}a^{\frac{5}{4}}\sqrt{2} \operatorname{atan}\left(\frac{c^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}-2\sqrt{c}x}{c^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) - 6c^{\frac{7}{4}}a^{\frac{1}{4}}\sqrt{2} \operatorname{atan}\left(\frac{c^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}-2\sqrt{c}x}{c^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) x^4 + 6c^{\frac{3}{4}}a^{\frac{5}{4}}\sqrt{2} \operatorname{atan}\left(\frac{c^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}+2\sqrt{c}x}{c^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) + \dots}{\dots}$$

input

$$\operatorname{int}(1/(c*x^4+a)^2,x)$$

output

$$\begin{aligned} & \left(-6c^{3/4}a^{5/4}\sqrt{2}\operatorname{atan}\left(\frac{c^{1/4}a^{1/4}\sqrt{2}-2\sqrt{c}x}{c^{1/4}a^{1/4}\sqrt{2}}\right) - 6c^{7/4}a^{1/4}\sqrt{2}\operatorname{atan}\left(\frac{c^{1/4}a^{1/4}\sqrt{2}-2\sqrt{c}x}{c^{1/4}a^{1/4}\sqrt{2}}\right) \right) a - 6c^{3/4}a^{5/4}\sqrt{2}\operatorname{atan}\left(\frac{c^{1/4}a^{1/4}\sqrt{2}-2\sqrt{c}x}{c^{1/4}a^{1/4}\sqrt{2}}\right) c x^4 \\ & + 6c^{3/4}a^{5/4}\sqrt{2}\operatorname{atan}\left(\frac{c^{1/4}a^{1/4}\sqrt{2}+2\sqrt{c}x}{c^{1/4}a^{1/4}\sqrt{2}}\right) a + 6c^{7/4}a^{1/4}\sqrt{2}\operatorname{atan}\left(\frac{c^{1/4}a^{1/4}\sqrt{2}+2\sqrt{c}x}{c^{1/4}a^{1/4}\sqrt{2}}\right) c x^4 \\ & - 3c^{3/4}a^{5/4}\sqrt{2}\log(-c^{1/4}a^{1/4}\sqrt{2}x + \sqrt{a} + \sqrt{c}x^2) a - 3c^{7/4}a^{1/4}\sqrt{2}\log(-c^{1/4}a^{1/4}\sqrt{2}x + \sqrt{a} + \sqrt{c}x^2) c x^4 + 3c^{3/4}a^{5/4}\sqrt{2}\log(c^{1/4}a^{1/4}\sqrt{2}x + \sqrt{a} + \sqrt{c}x^2) a \\ & + 3c^{7/4}a^{1/4}\sqrt{2}\log(c^{1/4}a^{1/4}\sqrt{2}x + \sqrt{a} + \sqrt{c}x^2) c x^4 + 8a^2c^2x/(32a^2c(a + cx^4)) \end{aligned}$$

3.7 $\int \frac{1}{(a+cx^4)^3} dx$

Optimal result	86
Mathematica [A] (verified)	87
Rubi [A] (verified)	87
Maple [C] (verified)	93
Fricas [C] (verification not implemented)	94
Sympy [A] (verification not implemented)	94
Maxima [A] (verification not implemented)	95
Giac [A] (verification not implemented)	95
Mupad [B] (verification not implemented)	96
Reduce [B] (verification not implemented)	96

Optimal result

Integrand size = 9, antiderivative size = 168

$$\int \frac{1}{(a+cx^4)^3} dx = \frac{x}{8a(a+cx^4)^2} + \frac{7x}{32a^2(a+cx^4)} - \frac{21 \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{cx}}{\sqrt[4]{a}}\right)}{64\sqrt{2}a^{11/4}\sqrt[4]{c}} + \frac{21 \arctan\left(1 + \frac{\sqrt{2}\sqrt[4]{cx}}{\sqrt[4]{a}}\right)}{64\sqrt{2}a^{11/4}\sqrt[4]{c}} + \frac{21 \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{a}\sqrt[4]{cx}}{\sqrt{a+\sqrt{cx^2}}}\right)}{64\sqrt{2}a^{11/4}\sqrt[4]{c}}$$

output

```
1/8*x/a/(c*x^4+a)^2+7/32*x/a^2/(c*x^4+a)+21/128*arctan(-1+2^(1/2)*c^(1/4)*
x/a^(1/4))*2^(1/2)/a^(11/4)/c^(1/4)+21/128*arctan(1+2^(1/2)*c^(1/4)*x/a^(1
/4))*2^(1/2)/a^(11/4)/c^(1/4)+21/128*arctanh(2^(1/2)*a^(1/4)*c^(1/4)*x/(a^
(1/2)+c^(1/2)*x^2))*2^(1/2)/a^(11/4)/c^(1/4)
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.19

$$\int \frac{1}{(a + cx^4)^3} dx$$

$$= \frac{\frac{32a^{7/4}x}{(a+cx^4)^2} + \frac{56a^{3/4}x}{a+cx^4} - \frac{42\sqrt{2} \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{cx}}{\sqrt[4]{a}}\right)}{\sqrt[4]{c}} + \frac{42\sqrt{2} \arctan\left(1 + \frac{\sqrt{2}\sqrt[4]{cx}}{\sqrt[4]{a}}\right)}{\sqrt[4]{c}} - \frac{21\sqrt{2} \log\left(\sqrt{a} - \sqrt{2}\sqrt[4]{a}\sqrt[4]{cx} + \sqrt{cx^2}\right)}{\sqrt[4]{c}} + \frac{21\sqrt{2} \log\left(\sqrt{a} + \sqrt{2}\sqrt[4]{a}\sqrt[4]{cx} + \sqrt{cx^2}\right)}{\sqrt[4]{c}}}{256a^{11/4}}$$

input `Integrate[(a + c*x^4)^(-3),x]`

output `((32*a^(7/4)*x)/(a + c*x^4)^2 + (56*a^(3/4)*x)/(a + c*x^4) - (42*sqrt(2)*ArcTan[1 - (sqrt(2)*c^(1/4)*x)/a^(1/4)]/c^(1/4) + (42*sqrt(2)*ArcTan[1 + (sqrt(2)*c^(1/4)*x)/a^(1/4)]/c^(1/4) - (21*sqrt(2)*Log[sqrt(a) - sqrt(2)*a^(1/4)*c^(1/4)*x + sqrt(c)*x^2])/c^(1/4) + (21*sqrt(2)*Log[sqrt(a) + sqrt(2)*a^(1/4)*c^(1/4)*x + sqrt(c)*x^2])/c^(1/4))/(256*a^(11/4))`

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.49, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 1.111$, Rules used = {749, 749, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + cx^4)^3} dx$$

$$\downarrow 749$$

$$\frac{7 \int \frac{1}{(cx^4+a)^2} dx}{8a} + \frac{x}{8a(a + cx^4)^2}$$

$$\downarrow 749$$

$$7 \left(\frac{3 \int \frac{1}{cx^4+a} dx}{4a} + \frac{x}{4a(a+cx^4)} \right) + \frac{x}{8a(a+cx^4)^2}$$

↓ 755

$$7 \left(\frac{3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{cx^2}}{cx^4+a} dx}{2\sqrt{a}} + \frac{\int \frac{\sqrt{cx^2}+\sqrt{a}}{cx^4+a} dx}{2\sqrt{a}} \right)}{4a} + \frac{x}{4a(a+cx^4)} \right) + \frac{x}{8a(a+cx^4)^2}$$

↓ 1476

$$7 \left(\frac{3 \left(\frac{\int \frac{1}{x^2 - \sqrt{2} \sqrt[4]{a}x + \sqrt{a}} dx}{\frac{\sqrt{c}}{2\sqrt{a}}} + \frac{\int \frac{1}{x^2 + \sqrt{2} \sqrt[4]{a}x + \sqrt{a}} dx}{\frac{\sqrt{c}}{2\sqrt{a}}} + \frac{\int \frac{\sqrt{a}-\sqrt{cx^2}}{cx^4+a} dx}{2\sqrt{a}} \right)}{4a} + \frac{x}{4a(a+cx^4)} \right) + \frac{x}{8a(a+cx^4)^2}$$

↓ 1082

$$\left(\frac{3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{cx^2}}{cx^4+a} dx}{2\sqrt{a}} + \frac{\frac{\int \frac{1}{\left(1-\frac{\sqrt{2}\sqrt[4]{cx}\right)^2} d\left(1-\frac{\sqrt{2}\sqrt[4]{cx}\right)}{\sqrt[4]{a}}\right)^{-1}}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}}}{2\sqrt{a}} - \frac{\frac{\int \frac{1}{\left(\frac{\sqrt{2}\sqrt[4]{cx}+1\right)^2} d\left(\frac{\sqrt{2}\sqrt[4]{cx}+1\right)}{\sqrt[4]{a}}\right)^{-1}}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}}}{2\sqrt{a}} \right)}{4a} + \frac{x}{4a(ax^4)} \right) +$$

$$\frac{8a}{x} \frac{x}{8a(a+cx^4)^2}$$

217

$$\left(\frac{3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{cx^2}}{cx^4+a} dx}{2\sqrt{a}} + \frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{cx}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}}}{2\sqrt{a}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{cx}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{c}}}{2\sqrt{a}} \right)}{4a} + \frac{x}{4a(ax^4)} \right) + \frac{x}{8a(a+cx^4)^2}$$

1479

$$\left(\frac{3}{7} \left(\frac{\int -\frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{c} x}{\sqrt[4]{c} \left(x^2 - \frac{\sqrt{2} \sqrt[4]{a} x + \sqrt{a}}{\sqrt[4]{c}} \right)} dx}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} - \frac{\int -\frac{\sqrt{2} (\sqrt{2} \sqrt[4]{c} x + \sqrt[4]{a})}{\sqrt[4]{c} \left(x^2 + \frac{\sqrt{2} \sqrt[4]{a} x + \sqrt{a}}{\sqrt[4]{c}} \right)} dx}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} + \frac{\arctan \left(\frac{\sqrt{2} \sqrt[4]{c} x + 1}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[4]{c} x}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} \right) + \frac{x}{4a(a+cx^4)} \right)$$

$$\frac{x}{8a(a+cx^4)^2}$$

↓ 25

$$\left(\frac{3}{7} \left(\frac{\int \frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{c} x}{\sqrt[4]{c} \left(x^2 - \frac{\sqrt{2} \sqrt[4]{a} x + \sqrt{a}}{\sqrt[4]{c}} \right)} dx}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} + \frac{\int \frac{\sqrt{2} (\sqrt{2} \sqrt[4]{c} x + \sqrt[4]{a})}{\sqrt[4]{c} \left(x^2 + \frac{\sqrt{2} \sqrt[4]{a} x + \sqrt{a}}{\sqrt[4]{c}} \right)} dx}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} + \frac{\arctan \left(\frac{\sqrt{2} \sqrt[4]{c} x + 1}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[4]{c} x}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} \right) + \frac{x}{4a(a+cx^4)} \right)$$

$$\frac{x}{8a(a+cx^4)^2}$$

↓ 27

$$\left(\frac{3 \left(\frac{\int \frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{c} x}{x^2 - \sqrt{2} \sqrt[4]{a} x + \sqrt{a}} dx}{2\sqrt{2} \sqrt[4]{a} \sqrt{c}} + \frac{\int \frac{\sqrt{2} \sqrt[4]{c} x + \sqrt{a}}{x^2 + \sqrt{2} \sqrt[4]{a} x + \sqrt{a}} dx}{2 \sqrt[4]{a} \sqrt{c}} + \frac{\arctan\left(\frac{\sqrt{2} \sqrt[4]{c} x + 1}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} - \frac{\arctan\left(1 - \frac{\sqrt{2} \sqrt[4]{c} x}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} \right)}{4a} + \frac{x}{4a(a+cx^4)} \right) + \frac{8a}{x(a+cx^4)^2}$$

1103

$$\left(\frac{3 \left(\frac{\arctan\left(\frac{\sqrt{2} \sqrt[4]{c} x + 1}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} - \frac{\arctan\left(1 - \frac{\sqrt{2} \sqrt[4]{c} x}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} + \frac{\log\left(\sqrt{2} \sqrt[4]{a} \sqrt[4]{c} x + \sqrt{a} + \sqrt{cx^2}\right)}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} - \frac{\log\left(-\sqrt{2} \sqrt[4]{a} \sqrt[4]{c} x + \sqrt{a} + \sqrt{cx^2}\right)}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{c}} \right)}{4a} + \frac{x}{4a(a+cx^4)} \right) + \frac{x}{8a(a+cx^4)^2}$$

input `Int[(a + c*x^4)^(-3), x]`

output

$$\frac{x/(8*a*(a + c*x^4)^2) + (7*(x/(4*a*(a + c*x^4)) + (3*((-(ArcTan[1 - (Sqrt[2]*c^(1/4)*x)/a^(1/4)]/(Sqrt[2]*a^(1/4)*c^(1/4))) + ArcTan[1 + (Sqrt[2]*c^(1/4)*x)/a^(1/4)]/(Sqrt[2]*a^(1/4)*c^(1/4)))/(2*Sqrt[a]) + (-1/2*Log[Sqrt[a] - Sqrt[2]*a^(1/4)*c^(1/4)*x + Sqrt[c]*x^2]/(Sqrt[2]*a^(1/4)*c^(1/4)) + Log[Sqrt[a] + Sqrt[2]*a^(1/4)*c^(1/4)*x + Sqrt[c]*x^2]/(2*Sqrt[2]*a^(1/4)*c^(1/4)))/(2*Sqrt[a])))/(4*a)))/(8*a)}$$

Defintions of rubi rules used

rule 25

$$\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$$

rule 27

$$\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_) \text{ ; FreeQ}[\text{b}, \text{x}]$$

rule 217

$$\text{Int}[(\text{a}_) + (\text{b}_)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(-\text{Rt}[-\text{a}, 2]*\text{Rt}[-\text{b}, 2])^{-1})*\text{ArcTan}[\text{Rt}[-\text{b}, 2]*(x/\text{Rt}[-\text{a}, 2])], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ \text{PosQ}[\text{a}/\text{b}] \ \&\& \ (\text{LtQ}[\text{a}, 0] \ || \ \text{LtQ}[\text{b}, 0])$$

rule 749

$$\text{Int}[(\text{a}_) + (\text{b}_)*(x_)^{(\text{n}_)})^{(\text{p}_)}, \text{x_Symbol}] \rightarrow \text{Simp}[(-x)*(\text{a} + \text{b}*x^{\text{n}})^{(\text{p} + 1)}/(\text{a}*n*(\text{p} + 1))), \text{x}] + \text{Simp}[(n*(\text{p} + 1) + 1)/(\text{a}*n*(\text{p} + 1)) \quad \text{Int}[(\text{a} + \text{b}*x^{\text{n}})^{(\text{p} + 1)}, \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ \text{IGtQ}[\text{n}, 0] \ \&\& \ \text{LtQ}[\text{p}, -1] \ \&\& \ (\text{IntegerQ}[2*\text{p}] \ || \ \text{Denominator}[\text{p} + 1/\text{n}] < \text{Denominator}[\text{p}])$$

rule 755

$$\text{Int}[(\text{a}_) + (\text{b}_)*(x_)^4)^{-1}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{r} = \text{Numerator}[\text{Rt}[\text{a}/\text{b}, 2]], \text{s} = \text{Denominator}[\text{Rt}[\text{a}/\text{b}, 2]]\}, \text{Simp}[1/(2*\text{r}) \quad \text{Int}[(\text{r} - \text{s}*x^2)/(\text{a} + \text{b}*x^4), \text{x}], \text{x}] + \text{Simp}[1/(2*\text{r}) \quad \text{Int}[(\text{r} + \text{s}*x^2)/(\text{a} + \text{b}*x^4), \text{x}], \text{x}]] \text{ ; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ (\text{GtQ}[\text{a}/\text{b}, 0] \ || \ (\text{PosQ}[\text{a}/\text{b}] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, \text{a}]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, \text{b}]])$$

rule 1082

$$\text{Int}[(\text{a}_) + (\text{b}_)*(x_) + (\text{c}_)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{q} = 1 - 4*\text{Simplify}[\text{a}*(\text{c}/\text{b}^2)]\}, \text{Simp}[-2/\text{b} \quad \text{Subst}[\text{Int}[1/(\text{q} - x^2), \text{x}], \text{x}, 1 + 2*\text{c}*(x/\text{b})], \text{x}] \text{ ; RationalQ}[\text{q}] \ \&\& \ (\text{EqQ}[\text{q}^2, 1] \ || \ \text{!RationalQ}[\text{b}^2 - 4*\text{a}*\text{c}]) \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}\}, \text{x}]$$

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[2*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.46 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.34

method	result	size
risch	$\frac{\frac{7cx^5 + 11x}{32a^2} + \frac{32a}{(cx^4 + a)^2} + \frac{21 \left(\sum_{-R=\text{RootOf}(cZ^4+a)} \frac{\ln(x-R)}{-R^3} \right)}{128a^2c}$	57
default	$\frac{x}{8a(cx^4+a)^2} + \frac{32a \left(\frac{7x}{(cx^4+a)} + \frac{21 \left(\frac{a}{c} \right)^{\frac{1}{4}} \sqrt{2} \left(\ln \left(\frac{x^2 + \left(\frac{a}{c} \right)^{\frac{1}{4}} x \sqrt{2} + \sqrt{\frac{a}{c}} \right)}{x^2 - \left(\frac{a}{c} \right)^{\frac{1}{4}} x \sqrt{2} + \sqrt{\frac{a}{c}}} \right) + 2 \arctan \left(\frac{\sqrt{2}x}{\left(\frac{a}{c} \right)^{\frac{1}{4}} + 1} \right) + 2 \arctan \left(\frac{\sqrt{2}x}{\left(\frac{a}{c} \right)^{\frac{1}{4}} - 1} \right) \right)}{256a^2}$	139

input `int(1/(c*x^4+a)^3,x,method=_RETURNVERBOSE)`

output `(7/32/a^2*c*x^5+11/32*x/a)/(c*x^4+a)^2+21/128/a^2/c*sum(1/_R^3*ln(x-_R),_R=RootOf(_Z^4*c+a))`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.07 (sec) , antiderivative size = 255, normalized size of antiderivative = 1.52

$$\int \frac{1}{(a + cx^4)^3} dx$$

$$= \frac{28cx^5 + 21(a^2c^2x^8 + 2a^3cx^4 + a^4)\left(-\frac{1}{a^{11}c}\right)^{\frac{1}{4}} \log\left(a^3\left(-\frac{1}{a^{11}c}\right)^{\frac{1}{4}} + x\right) - 21(-ia^2c^2x^8 - 2ia^3cx^4 - ia^4)\left(-\frac{1}{a^{11}c}\right)^{\frac{1}{4}} \log\left(-ia^3\left(-\frac{1}{a^{11}c}\right)^{\frac{1}{4}} + x\right)}{32a^4 + 64a^3cx^4 + 32a^2c^2x^8}$$

input `integrate(1/(c*x^4+a)^3,x, algorithm="fricas")`

output $\frac{1}{128}*(28*c*x^5 + 21*(a^2*c^2*x^8 + 2*a^3*c*x^4 + a^4)*(-1/(a^{11}*c))^{(1/4)} * \log(a^3*(-1/(a^{11}*c))^{(1/4)} + x) - 21*(-I*a^2*c^2*x^8 - 2*I*a^3*c*x^4 - I*a^4)*(-1/(a^{11}*c))^{(1/4)} * \log(I*a^3*(-1/(a^{11}*c))^{(1/4)} + x) - 21*(I*a^2*c^2*x^8 + 2*I*a^3*c*x^4 + I*a^4)*(-1/(a^{11}*c))^{(1/4)} * \log(-I*a^3*(-1/(a^{11}*c))^{(1/4)} + x) - 21*(a^2*c^2*x^8 + 2*a^3*c*x^4 + a^4)*(-1/(a^{11}*c))^{(1/4)} * \log(-a^3*(-1/(a^{11}*c))^{(1/4)} + x) + 44*a*x)/(a^2*c^2*x^8 + 2*a^3*c*x^4 + a^4)$

Sympy [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.38

$$\int \frac{1}{(a + cx^4)^3} dx$$

$$= \frac{11ax + 7cx^5}{32a^4 + 64a^3cx^4 + 32a^2c^2x^8} + \text{RootSum}\left(268435456t^4a^{11}c + 194481, \left(t \mapsto t \log\left(\frac{128ta^3}{21} + x\right)\right)\right)$$

input `integrate(1/(c*x**4+a)**3,x)`

output $(11*a*x + 7*c*x**5)/(32*a**4 + 64*a**3*c*x**4 + 32*a**2*c**2*x**8) + \text{RootSum}(268435456*_t**4*a**11*c + 194481, \text{Lambda}(_t, _t*\log(128*_t*a**3/21 + x)))$

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.26

$$\int \frac{1}{(a + cx^4)^3} dx = \frac{7cx^5 + 11ax}{32(a^2c^2x^8 + 2a^3cx^4 + a^4)}$$

$$+ \frac{21 \left(\frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(2\sqrt{cx} + \sqrt{2}a^{\frac{1}{4}}c^{\frac{1}{4}})}{2\sqrt{a}\sqrt{c}}\right)}{\sqrt{a}\sqrt{a}\sqrt{c}} + \frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(2\sqrt{cx} - \sqrt{2}a^{\frac{1}{4}}c^{\frac{1}{4}})}{2\sqrt{a}\sqrt{c}}\right)}{\sqrt{a}\sqrt{a}\sqrt{c}} + \frac{\sqrt{2} \log(\sqrt{cx^2 + \sqrt{2}a^{\frac{1}{4}}c^{\frac{1}{4}}x + \sqrt{a}})}{a^{\frac{3}{4}}c^{\frac{1}{4}}} - \frac{\sqrt{2} \log(\sqrt{cx^2 - \sqrt{2}a^{\frac{1}{4}}c^{\frac{1}{4}}x + \sqrt{a}})}{a^{\frac{3}{4}}c^{\frac{1}{4}}} \right)}{256a^2}$$

input `integrate(1/(c*x^4+a)^3,x, algorithm="maxima")`output

```
1/32*(7*c*x^5 + 11*a*x)/(a^2*c^2*x^8 + 2*a^3*c*x^4 + a^4) + 21/256*(2*sqrt(2)*arctan(1/2*sqrt(2)*(2*sqrt(c)*x + sqrt(2)*a^(1/4)*c^(1/4))/sqrt(sqrt(a)*sqrt(c)))/sqrt(a)*sqrt(sqrt(a)*sqrt(c)) + 2*sqrt(2)*arctan(1/2*sqrt(2)*(2*sqrt(c)*x - sqrt(2)*a^(1/4)*c^(1/4))/sqrt(sqrt(a)*sqrt(c)))/sqrt(a)*sqrt(sqrt(a)*sqrt(c)) + sqrt(2)*log(sqrt(c)*x^2 + sqrt(2)*a^(1/4)*c^(1/4)*x + sqrt(a))/(a^(3/4)*c^(1/4)) - sqrt(2)*log(sqrt(c)*x^2 - sqrt(2)*a^(1/4)*c^(1/4)*x + sqrt(a))/(a^(3/4)*c^(1/4))/a^2
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.21

$$\int \frac{1}{(a + cx^4)^3} dx = \frac{21\sqrt{2}(ac^3)^{\frac{1}{4}} \arctan\left(\frac{\sqrt{2}(2x + \sqrt{2}(\frac{a}{c})^{\frac{1}{4}})}{2(\frac{a}{c})^{\frac{1}{4}}}\right)}{128a^3c}$$

$$+ \frac{21\sqrt{2}(ac^3)^{\frac{1}{4}} \arctan\left(\frac{\sqrt{2}(2x - \sqrt{2}(\frac{a}{c})^{\frac{1}{4}})}{2(\frac{a}{c})^{\frac{1}{4}}}\right)}{128a^3c}$$

$$+ \frac{21\sqrt{2}(ac^3)^{\frac{1}{4}} \log\left(x^2 + \sqrt{2}x(\frac{a}{c})^{\frac{1}{4}} + \sqrt{\frac{a}{c}}\right)}{256a^3c}$$

$$- \frac{21\sqrt{2}(ac^3)^{\frac{1}{4}} \log\left(x^2 - \sqrt{2}x(\frac{a}{c})^{\frac{1}{4}} + \sqrt{\frac{a}{c}}\right)}{256a^3c} + \frac{7cx^5 + 11ax}{32(cx^4 + a)^2a^2}$$

input `integrate(1/(c*x^4+a)^3,x, algorithm="giac")`

output
$$\frac{21\sqrt{2}\sqrt{c^3}^{1/4}\arctan\left(\frac{1}{2}\sqrt{2}\left(2x + \sqrt{2}\sqrt{c}^{1/4}\right)\sqrt{c}^{1/4}\right)}{c^{3/4}} + \frac{21\sqrt{2}\sqrt{c^3}^{1/4}\arctan\left(\frac{1}{2}\sqrt{2}\left(2x - \sqrt{2}\sqrt{c}^{1/4}\right)\sqrt{c}^{1/4}\right)}{c^{3/4}} + \frac{21\sqrt{2}\sqrt{c^3}^{1/4}\log\left(x^2 + \sqrt{2}\sqrt{c}^{1/4}x + \sqrt{c}\right)}{c^{3/4}} - \frac{21\sqrt{2}\sqrt{c^3}^{1/4}\log\left(x^2 - \sqrt{2}\sqrt{c}^{1/4}x + \sqrt{c}\right)}{c^{3/4}} + \frac{1}{32} \frac{(7cx^5 + 11ax)}{(cx^4 + a)^2a^2}$$

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.48

$$\int \frac{1}{(a + cx^4)^3} dx = \frac{\frac{11x}{32a} + \frac{7cx^5}{32a^2}}{a^2 + 2acx^4 + c^2x^8} - \frac{21 \operatorname{atan}\left(\frac{c^{1/4}x}{(-a)^{1/4}}\right)}{64(-a)^{11/4}c^{1/4}} - \frac{21 \operatorname{atanh}\left(\frac{c^{1/4}x}{(-a)^{1/4}}\right)}{64(-a)^{11/4}c^{1/4}}$$

input `int(1/(a + c*x^4)^3,x)`

output
$$\left(\frac{11x}{32a} + \frac{7cx^5}{32a^2}\right)/(a^2 + c^2x^8 + 2acx^4) - \frac{21\operatorname{atan}\left(\frac{c^{1/4}x}{(-a)^{1/4}}\right)}{64(-a)^{11/4}c^{1/4}} - \frac{21\operatorname{atanh}\left(\frac{c^{1/4}x}{(-a)^{1/4}}\right)}{64(-a)^{11/4}c^{1/4}}$$

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 474, normalized size of antiderivative = 2.82

$$\int \frac{1}{(a + cx^4)^3} dx = \text{Too large to display}$$

input `int(1/(c*x^4+a)^3,x)`

output

```
( - 42*c**(3/4)*a**(1/4)*sqrt(2)*atan((c**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(c)*x)/(c**(1/4)*a**(1/4)*sqrt(2)))*a**2 - 84*c**(3/4)*a**(1/4)*sqrt(2)*atan((c**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(c)*x)/(c**(1/4)*a**(1/4)*sqrt(2)))*a*c*x**4 - 42*c**(3/4)*a**(1/4)*sqrt(2)*atan((c**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(c)*x)/(c**(1/4)*a**(1/4)*sqrt(2)))*c**2*x**8 + 42*c**(3/4)*a**(1/4)*sqrt(2)*atan((c**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(c)*x)/(c**(1/4)*a**(1/4)*sqrt(2)))*a**2 + 84*c**(3/4)*a**(1/4)*sqrt(2)*atan((c**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(c)*x)/(c**(1/4)*a**(1/4)*sqrt(2)))*a*c*x**4 + 42*c**(3/4)*a**(1/4)*sqrt(2)*atan((c**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(c)*x)/(c**(1/4)*a**(1/4)*sqrt(2)))*c**2*x**8 - 21*c**(3/4)*a**(1/4)*sqrt(2)*log(-c**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(c)*x**2)*a**2 - 42*c**(3/4)*a**(1/4)*sqrt(2)*log(-c**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(c)*x**2)*a*c*x**4 - 21*c**(3/4)*a**(1/4)*sqrt(2)*log(-c**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(c)*x**2)*c**2*x**8 + 21*c**(3/4)*a**(1/4)*sqrt(2)*log(c**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(c)*x**2)*a**2 + 42*c**(3/4)*a**(1/4)*sqrt(2)*log(c**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(c)*x**2)*a*c*x**4 + 21*c**(3/4)*a**(1/4)*sqrt(2)*log(c**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(c)*x**2)*c**2*x**8 + 88*a**2*c*x + 56*a*c**2*x**5)/(256*a**3*c*(a**2 + 2*a*c*x**4 + c**2*x**8))
```

3.8 $\int \frac{1}{1+a+(-1+a)x^4} dx$

Optimal result	98
Mathematica [A] (verified)	98
Rubi [A] (verified)	99
Maple [C] (verified)	100
Fricas [C] (verification not implemented)	101
Sympy [A] (verification not implemented)	101
Maxima [B] (verification not implemented)	102
Giac [B] (verification not implemented)	103
Mupad [B] (verification not implemented)	104
Reduce [B] (verification not implemented)	105

Optimal result

Integrand size = 12, antiderivative size = 83

$$\int \frac{1}{1+a+(-1+a)x^4} dx = \frac{\arctan\left(\frac{\sqrt[4]{1-ax}}{\sqrt[4]{1+a}}\right)}{2\sqrt{1+a}\sqrt[4]{1-a^2}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{1-ax}}{\sqrt[4]{1+a}}\right)}{2\sqrt{1+a}\sqrt[4]{1-a^2}}$$

output

```
1/2*arctan((1-a)^(1/4)*x/(1+a)^(1/4))/(1+a)^(1/2)/(-a^2+1)^(1/4)+1/2*arctanh((1-a)^(1/4)*x/(1+a)^(1/4))/(1+a)^(1/2)/(-a^2+1)^(1/4)
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.93

$$\int \frac{1}{1+a+(-1+a)x^4} dx = \frac{-2 \arctan\left(1 - \sqrt{2}\sqrt[4]{\frac{-1+a}{1+a}}x\right) + 2 \arctan\left(1 + \sqrt{2}\sqrt[4]{\frac{-1+a}{1+a}}x\right) - \log\left(\sqrt{1+a} - \sqrt{2}\sqrt[4]{-1+a}\sqrt[4]{1+a}\right)}{4\sqrt{2}\sqrt[4]{-1+a}(1+a)^{3/4}}$$

input

```
Integrate[(1 + a + (-1 + a)*x^4)^(-1), x]
```

output

```
(-2*ArcTan[1 - Sqrt[2]*((-1 + a)/(1 + a))^(1/4)*x] + 2*ArcTan[1 + Sqrt[2]*
((-1 + a)/(1 + a))^(1/4)*x] - Log[Sqrt[1 + a] - Sqrt[2]*(-1 + a)^(1/4)*(1
+ a)^(1/4)*x + Sqrt[-1 + a]*x^2] + Log[Sqrt[1 + a] + Sqrt[2]*(-1 + a)^(1/4
)*(1 + a)^(1/4)*x + Sqrt[-1 + a]*x^2])/(4*Sqrt[2]*(-1 + a)^(1/4)*(1 + a)^(
3/4))
```

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {756, 218, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a-1)x^4 + a + 1} dx$$

$$\downarrow \text{756}$$

$$\frac{\int \frac{1}{\sqrt{a+1}-\sqrt{1-ax^2}} dx}{2\sqrt{a+1}} + \frac{\int \frac{1}{\sqrt{1-ax^2}+\sqrt{a+1}} dx}{2\sqrt{a+1}}$$

$$\downarrow \text{218}$$

$$\frac{\int \frac{1}{\sqrt{a+1}-\sqrt{1-ax^2}} dx}{2\sqrt{a+1}} + \frac{\arctan\left(\frac{\sqrt[4]{1-ax}}{\sqrt[4]{a+1}}\right)}{2\sqrt{a+1}\sqrt[4]{1-a^2}}$$

$$\downarrow \text{221}$$

$$\frac{\arctan\left(\frac{\sqrt[4]{1-ax}}{\sqrt[4]{a+1}}\right)}{2\sqrt{a+1}\sqrt[4]{1-a^2}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{1-ax}}{\sqrt[4]{a+1}}\right)}{2\sqrt{a+1}\sqrt[4]{1-a^2}}$$

input

```
Int[(1 + a + (-1 + a)*x^4)^(-1), x]
```

output

```
ArcTan[((1 - a)^(1/4)*x)/(1 + a)^(1/4)]/(2*Sqrt[1 + a]*(1 - a^2)^(1/4)) +
ArcTanh[((1 - a)^(1/4)*x)/(1 + a)^(1/4)]/(2*Sqrt[1 + a]*(1 - a^2)^(1/4))
```

Definitions of rubi rules used

rule 218 $\text{Int}[(a_+ + (b_+)(x_+)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a) * \text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$

rule 221 $\text{Int}[(a_+ + (b_+)(x_+)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a) * \text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$

rule 756 $\text{Int}[(a_+ + (b_+)(x_+)^4)^{-1}, x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[r/(2*a) \ \text{Int}[1/(r - s*x^2), x], x] + \text{Simp}[r/(2*a) \ \text{Int}[1/(r + s*x^2), x], x]] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ !\text{GtQ}[a/b, 0]$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.50 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.45

method	result	size
risch	$\frac{\left(\sum_{-R=\text{RootOf}((a-1)Z^4+1+a)} \frac{\ln(x-R)}{R^3 a - R^3} \right)}{4}$	37
default	$\frac{\left(\frac{1+a}{a-1} \right)^{\frac{1}{4}} \sqrt{2} \left(\ln \left(\frac{x^2 + \left(\frac{1+a}{a-1} \right)^{\frac{1}{4}} x \sqrt{2} + \sqrt{\frac{1+a}{a-1}}}{x^2 - \left(\frac{1+a}{a-1} \right)^{\frac{1}{4}} x \sqrt{2} + \sqrt{\frac{1+a}{a-1}}} \right) + 2 \arctan \left(\frac{\sqrt{2} x}{\left(\frac{1+a}{a-1} \right)^{\frac{1}{4}}} + 1 \right) + 2 \arctan \left(\frac{\sqrt{2} x}{\left(\frac{1+a}{a-1} \right)^{\frac{1}{4}}} - 1 \right) \right)}{8+8a}$	132

input `int(1/(1+a+(a-1)*x^4), x, method=_RETURNVERBOSE)`

output `1/4*sum(1/(_R^3*a-_R^3)*ln(x-_R), _R=RootOf((a-1)*_Z^4+1+a))`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 196, normalized size of antiderivative = 2.36

$$\int \frac{1}{1+a+(-1+a)x^4} dx$$

$$= \frac{1}{4} \left(-\frac{1}{a^4+2a^3-2a-1} \right)^{\frac{1}{4}} \log \left((a+1) \left(-\frac{1}{a^4+2a^3-2a-1} \right)^{\frac{1}{4}} + x \right)$$

$$- \frac{1}{4} \left(-\frac{1}{a^4+2a^3-2a-1} \right)^{\frac{1}{4}} \log \left(-(a+1) \left(-\frac{1}{a^4+2a^3-2a-1} \right)^{\frac{1}{4}} + x \right)$$

$$- \frac{1}{4} i \left(-\frac{1}{a^4+2a^3-2a-1} \right)^{\frac{1}{4}} \log \left(-(ia+i) \left(-\frac{1}{a^4+2a^3-2a-1} \right)^{\frac{1}{4}} + x \right)$$

$$+ \frac{1}{4} i \left(-\frac{1}{a^4+2a^3-2a-1} \right)^{\frac{1}{4}} \log \left(-(-ia-i) \left(-\frac{1}{a^4+2a^3-2a-1} \right)^{\frac{1}{4}} + x \right)$$

input `integrate(1/(1+a+(-1+a)*x^4),x, algorithm="fricas")`

output `1/4*(-1/(a^4 + 2*a^3 - 2*a - 1))^(1/4)*log((a + 1)*(-1/(a^4 + 2*a^3 - 2*a - 1))^(1/4) + x) - 1/4*(-1/(a^4 + 2*a^3 - 2*a - 1))^(1/4)*log(-(a + 1)*(-1/(a^4 + 2*a^3 - 2*a - 1))^(1/4) + x) - 1/4*I*(-1/(a^4 + 2*a^3 - 2*a - 1))^(1/4)*log(-(I*a + I)*(-1/(a^4 + 2*a^3 - 2*a - 1))^(1/4) + x) + 1/4*I*(-1/(a^4 + 2*a^3 - 2*a - 1))^(1/4)*log(-(-I*a - I)*(-1/(a^4 + 2*a^3 - 2*a - 1))^(1/4) + x)`

Sympy [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.39

$$\int \frac{1}{1+a+(-1+a)x^4} dx$$

$$= \text{RootSum}(t^4 \cdot (256a^4 + 512a^3 - 512a - 256) + 1, (t \mapsto t \log(4ta + 4t + x)))$$

input `integrate(1/(1+a+(-1+a)*x**4),x)`

output `RootSum(_t**4*(256*a**4 + 512*a**3 - 512*a - 256) + 1, Lambda(_t, _t*log(4*_t*a + 4*_t + x)))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 318 vs. $2(63) = 126$.

Time = 0.12 (sec) , antiderivative size = 318, normalized size of antiderivative = 3.83

$$\int \frac{1}{1+a+(-1+a)x^4} dx = \frac{\sqrt{2} \log\left(\sqrt{a-1}x^2 + \sqrt{2}(a+1)^{\frac{1}{4}}(a-1)^{\frac{1}{4}}x + \sqrt{a+1}\right)}{8(a+1)^{\frac{3}{4}}(a-1)^{\frac{1}{4}}} - \frac{\sqrt{2} \log\left(\sqrt{a-1}x^2 - \sqrt{2}(a+1)^{\frac{1}{4}}(a-1)^{\frac{1}{4}}x + \sqrt{a+1}\right)}{8(a+1)^{\frac{3}{4}}(a-1)^{\frac{1}{4}}} + \frac{\sqrt{2} \log\left(\frac{2\sqrt{a-1}x - \sqrt{2}\sqrt{-\sqrt{a+1}\sqrt{a-1}} + \sqrt{2}(a+1)^{\frac{1}{4}}(a-1)^{\frac{1}{4}}}{2\sqrt{a-1}x + \sqrt{2}\sqrt{-\sqrt{a+1}\sqrt{a-1}} + \sqrt{2}(a+1)^{\frac{1}{4}}(a-1)^{\frac{1}{4}}}\right)}{8\sqrt{-\sqrt{a+1}\sqrt{a-1}}\sqrt{a+1}} + \frac{\sqrt{2} \log\left(\frac{2\sqrt{a-1}x - \sqrt{2}\sqrt{-\sqrt{a+1}\sqrt{a-1}} - \sqrt{2}(a+1)^{\frac{1}{4}}(a-1)^{\frac{1}{4}}}{2\sqrt{a-1}x + \sqrt{2}\sqrt{-\sqrt{a+1}\sqrt{a-1}} - \sqrt{2}(a+1)^{\frac{1}{4}}(a-1)^{\frac{1}{4}}}\right)}{8\sqrt{-\sqrt{a+1}\sqrt{a-1}}\sqrt{a+1}}$$

input `integrate(1/(1+a+(-1+a)*x^4),x, algorithm="maxima")`

output `1/8*sqrt(2)*log(sqrt(a - 1)*x^2 + sqrt(2)*(a + 1)^(1/4)*(a - 1)^(1/4)*x + sqrt(a + 1))/((a + 1)^(3/4)*(a - 1)^(1/4)) - 1/8*sqrt(2)*log(sqrt(a - 1)*x^2 - sqrt(2)*(a + 1)^(1/4)*(a - 1)^(1/4)*x + sqrt(a + 1))/((a + 1)^(3/4)*(a - 1)^(1/4)) + 1/8*sqrt(2)*log((2*sqrt(a - 1)*x - sqrt(2)*sqrt(-sqrt(a + 1)*sqrt(a - 1)) + sqrt(2)*(a + 1)^(1/4)*(a - 1)^(1/4))/(2*sqrt(a - 1)*x + sqrt(2)*sqrt(-sqrt(a + 1)*sqrt(a - 1)) + sqrt(2)*(a + 1)^(1/4)*(a - 1)^(1/4)))/(sqrt(-sqrt(a + 1)*sqrt(a - 1))*sqrt(a + 1)) + 1/8*sqrt(2)*log((2*sqrt(a - 1)*x - sqrt(2)*sqrt(-sqrt(a + 1)*sqrt(a - 1)) - sqrt(2)*(a + 1)^(1/4)*(a - 1)^(1/4))/(2*sqrt(a - 1)*x + sqrt(2)*sqrt(-sqrt(a + 1)*sqrt(a - 1)) - sqrt(2)*(a + 1)^(1/4)*(a - 1)^(1/4)))/(sqrt(-sqrt(a + 1)*sqrt(a - 1))*sqrt(a + 1))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 267 vs. $2(63) = 126$.

Time = 0.12 (sec) , antiderivative size = 267, normalized size of antiderivative = 3.22

$$\int \frac{1}{1+a+(-1+a)x^4} dx = \frac{(a^4 - 2a^3 + 2a - 1)^{\frac{1}{4}} \arctan\left(\frac{\sqrt{2}\left(2x + \sqrt{2}\left(\frac{a+1}{a-1}\right)^{\frac{1}{4}}\right)}{2\left(\frac{a+1}{a-1}\right)^{\frac{1}{4}}}\right)}{2(\sqrt{2}a^2 - \sqrt{2})} + \frac{(a^4 - 2a^3 + 2a - 1)^{\frac{1}{4}} \arctan\left(\frac{\sqrt{2}\left(2x - \sqrt{2}\left(\frac{a+1}{a-1}\right)^{\frac{1}{4}}\right)}{2\left(\frac{a+1}{a-1}\right)^{\frac{1}{4}}}\right)}{2(\sqrt{2}a^2 - \sqrt{2})} + \frac{(a^4 - 2a^3 + 2a - 1)^{\frac{1}{4}} \log\left(x^2 + \sqrt{2}x\left(\frac{a+1}{a-1}\right)^{\frac{1}{4}} + \sqrt{\frac{a+1}{a-1}}\right)}{4(\sqrt{2}a^2 - \sqrt{2})} - \frac{(a^4 - 2a^3 + 2a - 1)^{\frac{1}{4}} \log\left(x^2 - \sqrt{2}x\left(\frac{a+1}{a-1}\right)^{\frac{1}{4}} + \sqrt{\frac{a+1}{a-1}}\right)}{4(\sqrt{2}a^2 - \sqrt{2})}$$

input `integrate(1/(1+a+(-1+a)*x^4),x, algorithm="giac")`

output

```
1/2*(a^4 - 2*a^3 + 2*a - 1)^(1/4)*arctan(1/2*sqrt(2)*(2*x + sqrt(2)*((a +
1)/(a - 1))^(1/4))/((a + 1)/(a - 1))^(1/4))/(sqrt(2)*a^2 - sqrt(2)) + 1/2*
(a^4 - 2*a^3 + 2*a - 1)^(1/4)*arctan(1/2*sqrt(2)*(2*x - sqrt(2)*((a + 1)/(
a - 1))^(1/4))/((a + 1)/(a - 1))^(1/4))/(sqrt(2)*a^2 - sqrt(2)) + 1/4*(a^4
- 2*a^3 + 2*a - 1)^(1/4)*log(x^2 + sqrt(2)*x*((a + 1)/(a - 1))^(1/4) + sq
rt((a + 1)/(a - 1)))/(sqrt(2)*a^2 - sqrt(2)) - 1/4*(a^4 - 2*a^3 + 2*a - 1)
^(1/4)*log(x^2 - sqrt(2)*x*((a + 1)/(a - 1))^(1/4) + sqrt((a + 1)/(a - 1)
))/(sqrt(2)*a^2 - sqrt(2))
```


Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 543, normalized size of antiderivative = 6.54

$$\int \frac{1}{1+a+(-1+a)x^4} dx$$

$$= \frac{\operatorname{atan}\left(\frac{\left(\frac{x(4a^3-12a^2+12a-4)}{4} - \frac{4a^4-8a^3+8a-4}{4(1-a)^{1/4}(a+1)^{3/4}}\right) \operatorname{li}\left(\frac{x(4a^3-12a^2+12a-4)}{4} + \frac{4a^4-8a^3+8a-4}{4(1-a)^{1/4}(a+1)^{3/4}}\right)}{\frac{(1-a)^{1/4}(a+1)^{3/4}}{x(4a^3-12a^2+12a-4)} - \frac{4a^4-8a^3+8a-4}{4(1-a)^{1/4}(a+1)^{3/4}}}, \frac{(1-a)^{1/4}(a+1)^{3/4}}{x(4a^3-12a^2+12a-4)} + \frac{4a^4-8a^3+8a-4}{4(1-a)^{1/4}(a+1)^{3/4}}}\right) \operatorname{li}\left(\frac{x(4a^3-12a^2+12a-4)}{4} - \frac{4a^4-8a^3+8a-4}{4(1-a)^{1/4}(a+1)^{3/4}}\right) \operatorname{li}\left(\frac{x(4a^3-12a^2+12a-4)}{4} + \frac{4a^4-8a^3+8a-4}{4(1-a)^{1/4}(a+1)^{3/4}}\right)}{2(1-a)^{1/4}(a+1)^{3/4}} + \frac{\operatorname{atan}\left(\frac{\frac{x(4a^3-12a^2+12a-4)}{4} - \frac{4a^4-8a^3+8a-4}{4(1-a)^{1/4}(a+1)^{3/4}} \operatorname{li}\left(\frac{x(4a^3-12a^2+12a-4)}{4} + \frac{4a^4-8a^3+8a-4}{4(1-a)^{1/4}(a+1)^{3/4}}\right)}{(1-a)^{1/4}(a+1)^{3/4}} + \frac{\frac{x(4a^3-12a^2+12a-4)}{4} + \frac{4a^4-8a^3+8a-4}{4(1-a)^{1/4}(a+1)^{3/4}} \operatorname{li}\left(\frac{x(4a^3-12a^2+12a-4)}{4} - \frac{4a^4-8a^3+8a-4}{4(1-a)^{1/4}(a+1)^{3/4}}\right)}{(1-a)^{1/4}(a+1)^{3/4}}}{\frac{(1-a)^{1/4}(a+1)^{3/4}}{x(4a^3-12a^2+12a-4)} - \frac{4a^4-8a^3+8a-4}{4(1-a)^{1/4}(a+1)^{3/4}}}, \frac{(1-a)^{1/4}(a+1)^{3/4}}{x(4a^3-12a^2+12a-4)} + \frac{4a^4-8a^3+8a-4}{4(1-a)^{1/4}(a+1)^{3/4}}}\right) \operatorname{li}\left(\frac{x(4a^3-12a^2+12a-4)}{4} - \frac{4a^4-8a^3+8a-4}{4(1-a)^{1/4}(a+1)^{3/4}}\right) \operatorname{li}\left(\frac{x(4a^3-12a^2+12a-4)}{4} + \frac{4a^4-8a^3+8a-4}{4(1-a)^{1/4}(a+1)^{3/4}}\right)}{2(1-a)^{1/4}(a+1)^{3/4}}$$

input `int(1/(a + x^4*(a - 1) + 1),x)`

output

```
(atan((((x*(12*a - 12*a^2 + 4*a^3 - 4))/4 - (8*a - 8*a^3 + 4*a^4 - 4)/(4*(1 - a)^(1/4)*(a + 1)^(3/4))))*li)/((1 - a)^(1/4)*(a + 1)^(3/4)) + (((x*(12*a - 12*a^2 + 4*a^3 - 4))/4 + (8*a - 8*a^3 + 4*a^4 - 4)/(4*(1 - a)^(1/4)*(a + 1)^(3/4))))*li)/((1 - a)^(1/4)*(a + 1)^(3/4)))/(((x*(12*a - 12*a^2 + 4*a^3 - 4))/4 - (8*a - 8*a^3 + 4*a^4 - 4)/(4*(1 - a)^(1/4)*(a + 1)^(3/4)))/((1 - a)^(1/4)*(a + 1)^(3/4)) - ((x*(12*a - 12*a^2 + 4*a^3 - 4))/4 + (8*a - 8*a^3 + 4*a^4 - 4)/(4*(1 - a)^(1/4)*(a + 1)^(3/4)))/((1 - a)^(1/4)*(a + 1)^(3/4))))*li)/(2*(1 - a)^(1/4)*(a + 1)^(3/4)) + atan((((x*(12*a - 12*a^2 + 4*a^3 - 4))/4 - ((8*a - 8*a^3 + 4*a^4 - 4)*li)/(4*(1 - a)^(1/4)*(a + 1)^(3/4)))/((1 - a)^(1/4)*(a + 1)^(3/4)) + ((x*(12*a - 12*a^2 + 4*a^3 - 4))/4 + ((8*a - 8*a^3 + 4*a^4 - 4)*li)/(4*(1 - a)^(1/4)*(a + 1)^(3/4)))/((1 - a)^(1/4)*(a + 1)^(3/4)))/(((x*(12*a - 12*a^2 + 4*a^3 - 4))/4 - ((8*a - 8*a^3 + 4*a^4 - 4)*li)/(4*(1 - a)^(1/4)*(a + 1)^(3/4)))*li)/((1 - a)^(1/4)*(a + 1)^(3/4)) - (((x*(12*a - 12*a^2 + 4*a^3 - 4))/4 + ((8*a - 8*a^3 + 4*a^4 - 4)*li)/(4*(1 - a)^(1/4)*(a + 1)^(3/4))))*li)/((1 - a)^(1/4)*(a + 1)^(3/4))))/((1 - a)^(1/4)*(a + 1)^(3/4))
```

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.93

$$\int \frac{1}{1+a+(-1+a)x^4} dx$$

$$= \frac{(a+1)^{\frac{1}{4}}(a-1)^{\frac{3}{4}}\sqrt{2} \left(-2\operatorname{atan}\left(\frac{(a+1)^{\frac{1}{4}}(a-1)^{\frac{1}{4}}\sqrt{2}-2\sqrt{a-1}x}{(a+1)^{\frac{1}{4}}(a-1)^{\frac{1}{4}}\sqrt{2}}\right) + 2\operatorname{atan}\left(\frac{(a+1)^{\frac{1}{4}}(a-1)^{\frac{1}{4}}\sqrt{2}+2\sqrt{a-1}x}{(a+1)^{\frac{1}{4}}(a-1)^{\frac{1}{4}}\sqrt{2}}\right) - \log\left(-\frac{(a+1)^{\frac{1}{4}}(a-1)^{\frac{1}{4}}\sqrt{2}x + \sqrt{a-1}x^2 + \sqrt{a+1}}{(a+1)^{\frac{1}{4}}(a-1)^{\frac{1}{4}}\sqrt{2}}\right) + \log\left(\frac{(a+1)^{\frac{1}{4}}(a-1)^{\frac{1}{4}}\sqrt{2}x + \sqrt{a-1}x^2 + \sqrt{a+1}}{(a+1)^{\frac{1}{4}}(a-1)^{\frac{1}{4}}\sqrt{2}}\right) \right)}{8a^2 - 1}$$

input `int(1/(1+a+(-1+a)*x^4),x)`output `((a + 1)**(1/4)*(a - 1)**(3/4)*sqrt(2)*(- 2*atan(((a + 1)**(1/4)*(a - 1)**(1/4)*sqrt(2) - 2*sqrt(a - 1)*x)/((a + 1)**(1/4)*(a - 1)**(1/4)*sqrt(2))) + 2*atan(((a + 1)**(1/4)*(a - 1)**(1/4)*sqrt(2) + 2*sqrt(a - 1)*x)/((a + 1)**(1/4)*(a - 1)**(1/4)*sqrt(2))) - log(- (a + 1)**(1/4)*(a - 1)**(1/4)*sqrt(2)*x + sqrt(a - 1)*x**2 + sqrt(a + 1)) + log((a + 1)**(1/4)*(a - 1)**(1/4)*sqrt(2)*x + sqrt(a - 1)*x**2 + sqrt(a + 1)))/(8*(a**2 - 1))`

3.9 $\int (a - bx^4)^{3/2} dx$

Optimal result	106
Mathematica [C] (verified)	106
Rubi [A] (verified)	107
Maple [A] (verified)	108
Fricas [A] (verification not implemented)	109
Sympy [A] (verification not implemented)	109
Maxima [F]	110
Giac [F]	110
Mupad [B] (verification not implemented)	110
Reduce [F]	111

Optimal result

Integrand size = 12, antiderivative size = 92

$$\int (a - bx^4)^{3/2} dx = \frac{2}{7}ax\sqrt{a - bx^4} + \frac{1}{7}x(a - bx^4)^{3/2} + \frac{4a^{9/4}\sqrt{1 - \frac{bx^4}{a}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt[4]{bx^4}}{\sqrt[4]{a}}\right), -1\right)}{7\sqrt[4]{b}\sqrt{a - bx^4}}$$

output

$2/7*a*x*(-b*x^4+a)^{(1/2)}+1/7*x*(-b*x^4+a)^{(3/2)}+4/7*a^{(9/4)}*(1-b*x^4/a)^{(1/2)}*\operatorname{EllipticF}(b^{(1/4)}*x/a^{(1/4)}, I)/b^{(1/4)}/(-b*x^4+a)^{(1/2)}$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 5.31 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.52

$$\int (a - bx^4)^{3/2} dx = \frac{ax\sqrt{a - bx^4} \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, \frac{1}{4}, \frac{5}{4}, \frac{bx^4}{a}\right)}{\sqrt{1 - \frac{bx^4}{a}}}$$

input

`Integrate[(a - b*x^4)^(3/2), x]`

output $(a*x*\text{Sqrt}[a - b*x^4]*\text{Hypergeometric2F1}[-3/2, 1/4, 5/4, (b*x^4)/a])/\text{Sqrt}[1 - (b*x^4)/a]$

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.05, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {748, 748, 765, 762}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a - bx^4)^{3/2} dx$$

$$\downarrow 748$$

$$\frac{6}{7}a \int \sqrt{a - bx^4} dx + \frac{1}{7}x(a - bx^4)^{3/2}$$

$$\downarrow 748$$

$$\frac{6}{7}a \left(\frac{2}{3}a \int \frac{1}{\sqrt{a - bx^4}} dx + \frac{1}{3}x\sqrt{a - bx^4} \right) + \frac{1}{7}x(a - bx^4)^{3/2}$$

$$\downarrow 765$$

$$\frac{6}{7}a \left(\frac{2a\sqrt{1 - \frac{bx^4}{a}} \int \frac{1}{\sqrt{1 - \frac{bx^4}{a}}} dx}{3\sqrt{a - bx^4}} + \frac{1}{3}x\sqrt{a - bx^4} \right) + \frac{1}{7}x(a - bx^4)^{3/2}$$

$$\downarrow 762$$

$$\frac{6}{7}a \left(\frac{2a^{5/4}\sqrt{1 - \frac{bx^4}{a}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt[4]{bx^4}}{\sqrt[4]{a}}\right), -1\right)}{3\sqrt[4]{b}\sqrt{a - bx^4}} + \frac{1}{3}x\sqrt{a - bx^4} \right) + \frac{1}{7}x(a - bx^4)^{3/2}$$

input $\text{Int}[(a - b*x^4)^(3/2), x]$

output

$$\frac{(x*(a - b*x^4)^{(3/2)})/7 + (6*a*((x*\text{Sqrt}[a - b*x^4])/3 + (2*a^{(5/4)}*\text{Sqrt}[1 - (b*x^4)/a]*\text{EllipticF}[\text{ArcSin}[(b^{(1/4)}*x)/a^{(1/4)}], -1])/(3*b^{(1/4)}*\text{Sqrt}[a - b*x^4])))/7$$
Defintions of rubi rules used

rule 748

$$\text{Int}[(a_ + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Simp}[x*((a + b*x^n)^p/(n*p + 1)), x] + \text{Simp}[a*n*(p/(n*p + 1)) \text{Int}[(a + b*x^n)^{(p - 1)}, x], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ (\text{IntegerQ}[2*p] \ || \ \text{LtQ}[\text{Denominator}[p + 1/n], \text{Denominator}[p]])$$

rule 762

$$\text{Int}[1/\text{Sqrt}[(a_ + (b_)*(x_)^4], x_Symbol] \rightarrow \text{Simp}[(1/(\text{Sqrt}[a]*\text{Rt}[-b/a, 4]))*\text{EllipticF}[\text{ArcSin}[\text{Rt}[-b/a, 4]*x], -1], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[b/a] \ \&\& \ \text{GtQ}[a, 0]$$

rule 765

$$\text{Int}[1/\text{Sqrt}[(a_ + (b_)*(x_)^4], x_Symbol] \rightarrow \text{Simp}[\text{Sqrt}[1 + b*(x^4/a)]/\text{Sqrt}[a + b*x^4] \text{Int}[1/\text{Sqrt}[1 + b*(x^4/a)], x], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[b/a] \ \&\& \ !\text{GtQ}[a, 0]$$
Maple [A] (verified)

Time = 0.55 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.00

method	result	size
risch	$\frac{x(-bx^4+3a)\sqrt{-bx^4+a}}{7} + \frac{4a^2\sqrt{1-\frac{\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1+\frac{\sqrt{b}x^2}{\sqrt{a}}}\text{EllipticF}\left(x\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}, i\right)}{7\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}\sqrt{-bx^4+a}}$	92
default	$-\frac{bx^5\sqrt{-bx^4+a}}{7} + \frac{3ax\sqrt{-bx^4+a}}{7} + \frac{4a^2\sqrt{1-\frac{\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1+\frac{\sqrt{b}x^2}{\sqrt{a}}}\text{EllipticF}\left(x\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}, i\right)}{7\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}\sqrt{-bx^4+a}}$	99
elliptic	$-\frac{bx^5\sqrt{-bx^4+a}}{7} + \frac{3ax\sqrt{-bx^4+a}}{7} + \frac{4a^2\sqrt{1-\frac{\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1+\frac{\sqrt{b}x^2}{\sqrt{a}}}\text{EllipticF}\left(x\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}, i\right)}{7\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}\sqrt{-bx^4+a}}$	99

input

$$\text{int}((-b*x^4+a)^{(3/2)}, x, \text{method}=_RETURNVERBOSE)$$

output

```
1/7*x*(-b*x^4+3*a)*(-b*x^4+a)^(1/2)+4/7*a^2/(1/a^(1/2)*b^(1/2))^(1/2)*(1-b
^(1/2)*x^2/a^(1/2))^(1/2)*(1+b^(1/2)*x^2/a^(1/2))^(1/2)/(-b*x^4+a)^(1/2)*E
llipticF(x*(1/a^(1/2)*b^(1/2))^(1/2),I)
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.57

$$\int (a - bx^4)^{3/2} dx = \frac{4}{7} a \sqrt{-b} \left(\frac{a}{b}\right)^{\frac{3}{4}} F(\arcsin\left(\frac{\left(\frac{a}{b}\right)^{\frac{1}{4}}}{x}\right) | -1) - \frac{1}{7} (bx^5 - 3ax) \sqrt{-bx^4 + a}$$

input

```
integrate((-b*x^4+a)^(3/2),x, algorithm="fricas")
```

output

```
4/7*a*sqrt(-b)*(a/b)^(3/4)*elliptic_f(arcsin((a/b)^(1/4)/x), -1) - 1/7*(b*
x^5 - 3*a*x)*sqrt(-b*x^4 + a)
```

Sympy [A] (verification not implemented)

Time = 0.54 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.42

$$\int (a - bx^4)^{3/2} dx = \frac{a^{\frac{3}{2}} x \Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\begin{matrix} -\frac{3}{2}, \frac{1}{4} \\ \frac{5}{4} \end{matrix} \middle| \frac{bx^4 e^{2i\pi}}{a}\right)}{4\Gamma\left(\frac{5}{4}\right)}$$

input

```
integrate((-b*x**4+a)**(3/2),x)
```

output

```
a**(3/2)*x*gamma(1/4)*hyper((-3/2, 1/4), (5/4,), b*x**4*exp_polar(2*I*pi)/
a)/(4*gamma(5/4))
```

Maxima [F]

$$\int (a - bx^4)^{3/2} dx = \int (-bx^4 + a)^{\frac{3}{2}} dx$$

input `integrate((-b*x^4+a)^(3/2),x, algorithm="maxima")`

output `integrate((-b*x^4 + a)^(3/2), x)`

Giac [F]

$$\int (a - bx^4)^{3/2} dx = \int (-bx^4 + a)^{\frac{3}{2}} dx$$

input `integrate((-b*x^4+a)^(3/2),x, algorithm="giac")`

output `integrate((-b*x^4 + a)^(3/2), x)`

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.41

$$\int (a - bx^4)^{3/2} dx = \frac{x(a - bx^4)^{3/2} {}_2F_1\left(-\frac{3}{2}, \frac{1}{4}; \frac{5}{4}, \frac{bx^4}{a}\right)}{\left(1 - \frac{bx^4}{a}\right)^{3/2}}$$

input `int((a - b*x^4)^(3/2),x)`

output `(x*(a - b*x^4)^(3/2)*hypergeom([-3/2, 1/4], 5/4, (b*x^4)/a))/(1 - (b*x^4)/a)^(3/2)`

Reduce [F]

$$\int (a - bx^4)^{3/2} dx = \frac{3\sqrt{-bx^4 + a} ax}{7} - \frac{\sqrt{-bx^4 + a} bx^5}{7} + \frac{4\left(\int \frac{\sqrt{-bx^4 + a}}{-bx^4 + a} dx\right) a^2}{7}$$

input `int((-b*x^4+a)^(3/2),x)`

output `(3*sqrt(a - b*x**4)*a*x - sqrt(a - b*x**4)*b*x**5 + 4*int(sqrt(a - b*x**4)/(a - b*x**4),x)*a**2)/7`

3.10 $\int \sqrt{a - bx^4} dx$

Optimal result	112
Mathematica [C] (verified)	112
Rubi [A] (verified)	113
Maple [A] (verified)	114
Fricas [A] (verification not implemented)	115
Sympy [A] (verification not implemented)	115
Maxima [F]	115
Giac [F]	116
Mupad [B] (verification not implemented)	116
Reduce [F]	116

Optimal result

Integrand size = 12, antiderivative size = 74

$$\int \sqrt{a - bx^4} dx = \frac{1}{3}x\sqrt{a - bx^4} + \frac{2a^{5/4}\sqrt{1 - \frac{bx^4}{a}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a}}\right), -1\right)}{3\sqrt[4]{b}\sqrt{a - bx^4}}$$

output

```
1/3*x*(-b*x^4+a)^(1/2)+2/3*a^(5/4)*(1-b*x^4/a)^(1/2)*EllipticF(b^(1/4)*x/a^(1/4),I)/b^(1/4)/(-b*x^4+a)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.22 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.19

$$\int \sqrt{a - bx^4} dx = \frac{x(a - bx^4) - \frac{2ia\sqrt{1 - \frac{bx^4}{a}} \operatorname{EllipticF}\left(i\operatorname{arcsinh}\left(\sqrt{-\frac{\sqrt{b}}{\sqrt{a}}}x\right), -1\right)}{\sqrt{-\frac{\sqrt{b}}{\sqrt{a}}}}}{3\sqrt{a - bx^4}}$$

input

```
Integrate[Sqrt[a - b*x^4], x]
```

output

```
(x*(a - b*x^4) - ((2*I)*a*Sqrt[1 - (b*x^4)/a]*EllipticF[I*ArcSinh[Sqrt[-(Sqrt[b]/Sqrt[a])]*x], -1])/Sqrt[-(Sqrt[b]/Sqrt[a])]/(3*Sqrt[a - b*x^4])
```

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {748, 765, 762}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{a - bx^4} dx$$

$$\downarrow 748$$

$$\frac{2}{3}a \int \frac{1}{\sqrt{a - bx^4}} dx + \frac{1}{3}x\sqrt{a - bx^4}$$

$$\downarrow 765$$

$$\frac{2a\sqrt{1 - \frac{bx^4}{a}} \int \frac{1}{\sqrt{1 - \frac{bx^4}{a}}} dx}{3\sqrt{a - bx^4}} + \frac{1}{3}x\sqrt{a - bx^4}$$

$$\downarrow 762$$

$$\frac{2a^{5/4}\sqrt{1 - \frac{bx^4}{a}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt[4]{bx^4}}{\sqrt[4]{a}}\right), -1\right)}{3\sqrt[4]{b}\sqrt{a - bx^4}} + \frac{1}{3}x\sqrt{a - bx^4}$$

input

```
Int[Sqrt[a - b*x^4], x]
```

output

```
(x*Sqrt[a - b*x^4])/3 + (2*a^(5/4)*Sqrt[1 - (b*x^4)/a]*EllipticF[ArcSin[(b^(1/4)*x)/a^(1/4)], -1])/ (3*b^(1/4)*Sqrt[a - b*x^4])
```

Definitions of rubi rules used

rule 748 $\text{Int}[(a_+ + (b_+)(x_+)^{n_+})^{p_+}, x_Symbol] \rightarrow \text{Simp}[x((a + b*x^n)^p/(n*p + 1)), x] + \text{Simp}[a*n*(p/(n*p + 1)) \text{Int}[(a + b*x^n)^{p-1}, x], x] /;$ FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || LtQ[Denominator[p + 1/n], Denominator[p]])

rule 762 $\text{Int}[1/\text{Sqrt}[(a_+ + (b_+)(x_+)^4], x_Symbol] \rightarrow \text{Simp}[(1/(\text{Sqrt}[a]*\text{Rt}[-b/a, 4]))*\text{EllipticF}[\text{ArcSin}[\text{Rt}[-b/a, 4]*x], -1], x] /;$ FreeQ[{a, b}, x] && NegQ[b/a] && GtQ[a, 0]

rule 765 $\text{Int}[1/\text{Sqrt}[(a_+ + (b_+)(x_+)^4], x_Symbol] \rightarrow \text{Simp}[\text{Sqrt}[1 + b*(x^4/a)]/\text{Sqrt}[a + b*x^4] \text{Int}[1/\text{Sqrt}[1 + b*(x^4/a)], x], x] /;$ FreeQ[{a, b}, x] && NegQ[b/a] && !GtQ[a, 0]

Maple [A] (verified)

Time = 0.54 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.08

method	result	size
default	$\frac{x\sqrt{-bx^4+a}}{3} + \frac{2a\sqrt{1-\frac{\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1+\frac{\sqrt{b}x^2}{\sqrt{a}}}\text{EllipticF}\left(x\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}, i\right)}{3\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}\sqrt{-bx^4+a}}$	80
risch	$\frac{x\sqrt{-bx^4+a}}{3} + \frac{2a\sqrt{1-\frac{\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1+\frac{\sqrt{b}x^2}{\sqrt{a}}}\text{EllipticF}\left(x\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}, i\right)}{3\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}\sqrt{-bx^4+a}}$	80
elliptic	$\frac{x\sqrt{-bx^4+a}}{3} + \frac{2a\sqrt{1-\frac{\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1+\frac{\sqrt{b}x^2}{\sqrt{a}}}\text{EllipticF}\left(x\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}, i\right)}{3\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}\sqrt{-bx^4+a}}$	80

input $\text{int}((-b*x^4+a)^{(1/2)}, x, \text{method}=_RETURNVERBOSE)$

output $\frac{1}{3}x*(-b*x^4+a)^{(1/2)} + \frac{2}{3}a/(1/a^{(1/2)}*b^{(1/2)})^{(1/2)}*(1-b^{(1/2)}*x^2/a^{(1/2)})^{(1/2)}*(1+b^{(1/2)}*x^2/a^{(1/2)})^{(1/2)}/(-b*x^4+a)^{(1/2)}*\text{EllipticF}(x*(1/a^{(1/2)}*b^{(1/2)})^{(1/2)}, I)$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.57

$$\int \sqrt{a - bx^4} dx = \frac{2}{3} \sqrt{-b} \left(\frac{a}{b}\right)^{\frac{3}{4}} F(\arcsin\left(\frac{\left(\frac{a}{b}\right)^{\frac{1}{4}}}{x}\right) | -1) + \frac{1}{3} \sqrt{-bx^4 + ax}$$

input `integrate((-b*x^4+a)^(1/2),x, algorithm="fricas")`output `2/3*sqrt(-b)*(a/b)^(3/4)*elliptic_f(arcsin((a/b)^(1/4)/x), -1) + 1/3*sqrt(-b*x^4 + a)*x`**Sympy [A] (verification not implemented)**

Time = 0.49 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.53

$$\int \sqrt{a - bx^4} dx = \frac{\sqrt{ax} \Gamma\left(\frac{1}{4}\right) {}_2F_1\left(-\frac{1}{2}, \frac{1}{4} \mid \frac{bx^4 e^{2i\pi}}{a}\right)}{4\Gamma\left(\frac{5}{4}\right)}$$

input `integrate((-b*x**4+a)**(1/2),x)`output `sqrt(a)*x*gamma(1/4)*hyper((-1/2, 1/4), (5/4,), b*x**4*exp_polar(2*I*pi)/a)/(4*gamma(5/4))`**Maxima [F]**

$$\int \sqrt{a - bx^4} dx = \int \sqrt{-bx^4 + a} dx$$

input `integrate((-b*x^4+a)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(-b*x^4 + a), x)`

Giac [F]

$$\int \sqrt{a - bx^4} dx = \int \sqrt{-bx^4 + a} dx$$

input `integrate((-b*x^4+a)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(-b*x^4 + a), x)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.51

$$\int \sqrt{a - bx^4} dx = \frac{x \sqrt{a - bx^4} {}_2F_1\left(-\frac{1}{2}, \frac{1}{4}; \frac{5}{4}; \frac{bx^4}{a}\right)}{\sqrt{1 - \frac{bx^4}{a}}}$$

input `int((a - b*x^4)^(1/2),x)`

output `(x*(a - b*x^4)^(1/2)*hypergeom([-1/2, 1/4], 5/4, (b*x^4)/a))/(1 - (b*x^4)/a)^(1/2)`

Reduce [F]

$$\int \sqrt{a - bx^4} dx = \frac{\sqrt{-bx^4 + a} x}{3} + \frac{2\left(\int \frac{\sqrt{-bx^4+a}}{-bx^4+a} dx\right) a}{3}$$

input `int((-b*x^4+a)^(1/2),x)`

output $(\sqrt{a - b*x**4}*x + 2*\text{int}(\sqrt{a - b*x**4}/(a - b*x**4),x)*a)/3$

3.11 $\int \frac{1}{\sqrt{a-bx^4}} dx$

Optimal result	118
Mathematica [C] (verified)	118
Rubi [A] (verified)	119
Maple [A] (verified)	120
Fricas [A] (verification not implemented)	120
Sympy [A] (verification not implemented)	121
Maxima [F]	121
Giac [F]	122
Mupad [B] (verification not implemented)	122
Reduce [F]	122

Optimal result

Integrand size = 12, antiderivative size = 53

$$\int \frac{1}{\sqrt{a-bx^4}} dx = \frac{\sqrt[4]{a}\sqrt{1-\frac{bx^4}{a}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt[4]{b}x}{\sqrt[4]{a}}\right), -1\right)}{\sqrt[4]{b}\sqrt{a-bx^4}}$$

output `a^(1/4)*(1-b*x^4/a)^(1/2)*EllipticF(b^(1/4)*x/a^(1/4),I)/b^(1/4)/(-b*x^4+a)^(1/2)`

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 10.04 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.36

$$\int \frac{1}{\sqrt{a-bx^4}} dx = -\frac{i\sqrt{1-\frac{bx^4}{a}} \operatorname{EllipticF}\left(i\operatorname{arcsinh}\left(\sqrt{-\frac{\sqrt{b}}{\sqrt{a}}}x\right), -1\right)}{\sqrt{-\frac{\sqrt{b}}{\sqrt{a}}}\sqrt{a-bx^4}}$$

input `Integrate[1/Sqrt[a - b*x^4],x]`

output $((-I)*\text{Sqrt}[1 - (b*x^4)/a]*\text{EllipticF}[I*\text{ArcSinh}[\text{Sqrt}[-(\text{Sqrt}[b]/\text{Sqrt}[a])]]*x, -1])/(\text{Sqrt}[-(\text{Sqrt}[b]/\text{Sqrt}[a])]*\text{Sqrt}[a - b*x^4])$

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {765, 762}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{a - bx^4}} dx$$

$$\downarrow 765$$

$$\frac{\sqrt{1 - \frac{bx^4}{a}} \int \frac{1}{\sqrt{1 - \frac{bx^4}{a}}} dx}{\sqrt{a - bx^4}}$$

$$\downarrow 762$$

$$\frac{\sqrt[4]{a} \sqrt{1 - \frac{bx^4}{a}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a}}\right), -1\right)}{\sqrt[4]{b} \sqrt{a - bx^4}}$$

input $\text{Int}[1/\text{Sqrt}[a - b*x^4], x]$

output $(a^{(1/4)}*\text{Sqrt}[1 - (b*x^4)/a]*\text{EllipticF}[\text{ArcSin}[(b^{(1/4)}*x)/a^{(1/4)}], -1])/ (b^{(1/4)}*\text{Sqrt}[a - b*x^4])$

Defintions of rubi rules used

rule 762 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := Simp[(1/(Sqrt[a]*Rt[-b/a, 4])
)*EllipticF[ArcSin[Rt[-b/a, 4]*x], -1], x] /; FreeQ[{a, b}, x] && NegQ[b/a]
&& GtQ[a, 0]`

rule 765 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := Simp[Sqrt[1 + b*(x^4/a)]/Sqrt
[a + b*x^4] Int[1/Sqrt[1 + b*(x^4/a)], x], x] /; FreeQ[{a, b}, x] && NegQ
[b/a] && !GtQ[a, 0]`

Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.21

method	result	size
default	$\frac{\sqrt{1 - \frac{\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1 + \frac{\sqrt{b}x^2}{\sqrt{a}}}\operatorname{EllipticF}\left(x\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}, i\right)}{\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}\sqrt{-bx^4+a}}$	64
elliptic	$\frac{\sqrt{1 - \frac{\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1 + \frac{\sqrt{b}x^2}{\sqrt{a}}}\operatorname{EllipticF}\left(x\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}, i\right)}{\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}\sqrt{-bx^4+a}}$	64

input `int(1/(-b*x^4+a)^(1/2),x,method=_RETURNVERBOSE)`

output `1/(1/a^(1/2)*b^(1/2))^(1/2)*(1-b^(1/2)*x^2/a^(1/2))^(1/2)*(1+b^(1/2)*x^2/a
^(1/2))^(1/2)/(-b*x^4+a)^(1/2)*EllipticF(x*(1/a^(1/2)*b^(1/2))^(1/2),I)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.49

$$\int \frac{1}{\sqrt{a - bx^4}} dx = \frac{\sqrt{a}\left(\frac{b}{a}\right)^{\frac{3}{4}} F\left(\arcsin\left(x\left(\frac{b}{a}\right)^{\frac{1}{4}}\right) \mid -1\right)}{b}$$

input `integrate(1/(-b*x^4+a)^(1/2),x, algorithm="fricas")`

output `sqrt(a)*(b/a)^(3/4)*elliptic_f(arcsin(x*(b/a)^(1/4)), -1)/b`

Sympy [A] (verification not implemented)

Time = 0.47 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.70

$$\int \frac{1}{\sqrt{a - bx^4}} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2} \middle| \frac{bx^4 e^{2i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{5}{4}\right)}$$

input `integrate(1/(-b*x**4+a)**(1/2),x)`

output `x*gamma(1/4)*hyper((1/4, 1/2), (5/4,), b*x**4*exp_polar(2*I*pi)/a)/(4*sqrt(a)*gamma(5/4))`

Maxima [F]

$$\int \frac{1}{\sqrt{a - bx^4}} dx = \int \frac{1}{\sqrt{-bx^4 + a}} dx$$

input `integrate(1/(-b*x^4+a)^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(-b*x^4 + a), x)`

Giac [F]

$$\int \frac{1}{\sqrt{a - bx^4}} dx = \int \frac{1}{\sqrt{-bx^4 + a}} dx$$

input `integrate(1/(-b*x^4+a)^(1/2),x, algorithm="giac")`

output `integrate(1/sqrt(-b*x^4 + a), x)`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.72

$$\int \frac{1}{\sqrt{a - bx^4}} dx = \frac{x \sqrt{1 - \frac{bx^4}{a}} {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}, \frac{bx^4}{a}\right)}{\sqrt{a - bx^4}}$$

input `int(1/(a - b*x^4)^(1/2),x)`

output `(x*(1 - (b*x^4)/a)^(1/2)*hypergeom([1/4, 1/2], 5/4, (b*x^4)/a))/(a - b*x^4)^(1/2)`

Reduce [F]

$$\int \frac{1}{\sqrt{a - bx^4}} dx = \int \frac{\sqrt{-bx^4 + a}}{-bx^4 + a} dx$$

input `int(1/(-b*x^4+a)^(1/2),x)`

output `int(sqrt(a - b*x**4)/(a - b*x**4),x)`

3.12 $\int \frac{1}{(a-bx^4)^{3/2}} dx$

Optimal result	123
Mathematica [C] (verified)	123
Rubi [A] (verified)	124
Maple [A] (verified)	125
Fricas [A] (verification not implemented)	126
Sympy [A] (verification not implemented)	126
Maxima [F]	126
Giac [F]	127
Mupad [B] (verification not implemented)	127
Reduce [F]	127

Optimal result

Integrand size = 12, antiderivative size = 77

$$\int \frac{1}{(a-bx^4)^{3/2}} dx = \frac{x}{2a\sqrt{a-bx^4}} + \frac{\sqrt{1-\frac{bx^4}{a}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt[4]{bx^4}}{\sqrt[4]{a}}\right), -1\right)}{2a^{3/4}\sqrt[4]{b}\sqrt{a-bx^4}}$$

output `1/2*x/a/(-b*x^4+a)^(1/2)+1/2*(1-b*x^4/a)^(1/2)*EllipticF(b^(1/4)*x/a^(1/4),1)/a^(3/4)/b^(1/4)/(-b*x^4+a)^(1/2)`

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 5.86 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.73

$$\int \frac{1}{(a-bx^4)^{3/2}} dx = \frac{x + x\sqrt{1-\frac{bx^4}{a}} \operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, \frac{bx^4}{a}\right)}{2a\sqrt{a-bx^4}}$$

input `Integrate[(a - b*x^4)^(-3/2),x]`

output $(x + x\sqrt{1 - (bx^4)/a})\text{Hypergeometric2F1}[1/4, 1/2, 5/4, (bx^4)/a]/(2a\sqrt{a - bx^4})$

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {749, 765, 762}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(a - bx^4)^{3/2}} dx \\ & \quad \downarrow 749 \\ & \frac{\int \frac{1}{\sqrt{a - bx^4}} dx}{2a} + \frac{x}{2a\sqrt{a - bx^4}} \\ & \quad \downarrow 765 \\ & \frac{\sqrt{1 - \frac{bx^4}{a}} \int \frac{1}{\sqrt{1 - \frac{bx^4}{a}}} dx}{2a\sqrt{a - bx^4}} + \frac{x}{2a\sqrt{a - bx^4}} \\ & \quad \downarrow 762 \\ & \frac{\sqrt{1 - \frac{bx^4}{a}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt[4]{bx^4}}{\sqrt[4]{a}}\right), -1\right)}{2a^{3/4}\sqrt[4]{b}\sqrt{a - bx^4}} + \frac{x}{2a\sqrt{a - bx^4}} \end{aligned}$$

input $\text{Int}[(a - bx^4)^{-3/2}, x]$

output $x/(2a\sqrt{a - bx^4}) + (\sqrt{1 - (bx^4)/a})\text{EllipticF}[\text{ArcSin}[(b^{1/4})x/a^{1/4}], -1]/(2a^{3/4}b^{1/4}\sqrt{a - bx^4})$

Definitions of rubi rules used

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

rule 762 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := Simp[(1/(Sqrt[a]*Rt[-b/a, 4]))*EllipticF[ArcSin[Rt[-b/a, 4]*x], -1], x] /; FreeQ[{a, b}, x] && NegQ[b/a] && GtQ[a, 0]`

rule 765 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := Simp[Sqrt[1 + b*(x^4/a)]/Sqrt[a + b*x^4] Int[1/Sqrt[1 + b*(x^4/a)], x], x] /; FreeQ[{a, b}, x] && NegQ[b/a] && !GtQ[a, 0]`

Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.17

method	result	size
default	$\frac{x}{2a\sqrt{-(x^4-\frac{a}{b})b}} + \frac{\sqrt{1-\frac{\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1+\frac{\sqrt{b}x^2}{\sqrt{a}}}\text{EllipticF}\left(x\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}, i\right)}{2a\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}\sqrt{-bx^4+a}}$	90
elliptic	$\frac{x}{2a\sqrt{-(x^4-\frac{a}{b})b}} + \frac{\sqrt{1-\frac{\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1+\frac{\sqrt{b}x^2}{\sqrt{a}}}\text{EllipticF}\left(x\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}, i\right)}{2a\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}\sqrt{-bx^4+a}}$	90

input `int(1/(-b*x^4+a)^(3/2), x, method=_RETURNVERBOSE)`

output `1/2*x/a/(-(x^4-a/b)*b)^(1/2)+1/2/a/(1/a^(1/2)*b^(1/2))^(1/2)*(1-b^(1/2)*x^2/a^(1/2))^(1/2)*(1+b^(1/2)*x^2/a^(1/2))^(1/2)/(-b*x^4+a)^(1/2)*EllipticF(x*(1/a^(1/2)*b^(1/2))^(1/2), I)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.86

$$\int \frac{1}{(a - bx^4)^{3/2}} dx = \frac{(bx^4 - a)\sqrt{a}\left(\frac{b}{a}\right)^{3/4} F(\arcsin\left(x\left(\frac{b}{a}\right)^{1/4}\right) | -1) - \sqrt{-bx^4 + abx}}{2(ab^2x^4 - a^2b)}$$

input `integrate(1/(-b*x^4+a)^(3/2),x, algorithm="fricas")`output `1/2*((b*x^4 - a)*sqrt(a)*(b/a)^(3/4)*elliptic_f(arcsin(x*(b/a)^(1/4)), -1) - sqrt(-b*x^4 + a)*b*x)/(a*b^2*x^4 - a^2*b)`**Sympy [A] (verification not implemented)**

Time = 0.54 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.48

$$\int \frac{1}{(a - bx^4)^{3/2}} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{3}{2} \middle| \frac{bx^4 e^{2i\pi}}{a}\right)}{4a^{3/2}\Gamma\left(\frac{5}{4}\right)}$$

input `integrate(1/(-b*x**4+a)**(3/2),x)`output `x*gamma(1/4)*hyper((1/4, 3/2), (5/4,), b*x**4*exp_polar(2*I*pi)/a)/(4*a**(3/2)*gamma(5/4))`**Maxima [F]**

$$\int \frac{1}{(a - bx^4)^{3/2}} dx = \int \frac{1}{(-bx^4 + a)^{3/2}} dx$$

input `integrate(1/(-b*x^4+a)^(3/2),x, algorithm="maxima")`

output `integrate((-b*x^4 + a)^(-3/2), x)`

Giac [F]

$$\int \frac{1}{(a - bx^4)^{3/2}} dx = \int \frac{1}{(-bx^4 + a)^{\frac{3}{2}}} dx$$

input `integrate(1/(-b*x^4+a)^(3/2),x, algorithm="giac")`

output `integrate((-b*x^4 + a)^(-3/2), x)`

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.49

$$\int \frac{1}{(a - bx^4)^{3/2}} dx = \frac{x \left(1 - \frac{bx^4}{a}\right)^{3/2} {}_2F_1\left(\frac{1}{4}, \frac{3}{2}; \frac{5}{4}; \frac{bx^4}{a}\right)}{(a - bx^4)^{3/2}}$$

input `int(1/(a - b*x^4)^(3/2),x)`

output `(x*(1 - (b*x^4)/a)^(3/2)*hypergeom([1/4, 3/2], 5/4, (b*x^4)/a))/(a - b*x^4)^(3/2)`

Reduce [F]

$$\int \frac{1}{(a - bx^4)^{3/2}} dx = \int \frac{\sqrt{-bx^4 + a}}{b^2x^8 - 2abx^4 + a^2} dx$$

input `int(1/(-b*x^4+a)^(3/2),x)`

output `int(sqrt(a - b*x**4)/(a**2 - 2*a*b*x**4 + b**2*x**8),x)`

3.13 $\int \frac{1}{(a-bx^4)^{5/2}} dx$

Optimal result	129
Mathematica [C] (verified)	129
Rubi [A] (verified)	130
Maple [A] (verified)	131
Fricas [A] (verification not implemented)	132
Sympy [A] (verification not implemented)	132
Maxima [F]	133
Giac [F]	133
Mupad [B] (verification not implemented)	133
Reduce [F]	134

Optimal result

Integrand size = 12, antiderivative size = 97

$$\int \frac{1}{(a-bx^4)^{5/2}} dx = \frac{x}{6a(a-bx^4)^{3/2}} + \frac{5x}{12a^2\sqrt{a-bx^4}} + \frac{5\sqrt{1-\frac{bx^4}{a}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt[4]{bx^4}}{\sqrt[4]{a}}\right), -1\right)}{12a^{7/4}\sqrt[4]{b}\sqrt{a-bx^4}}$$

output

```
1/6*x/a/(-b*x^4+a)^(3/2)+5/12*x/a^2/(-b*x^4+a)^(1/2)+5/12*(1-b*x^4/a)^(1/2)*EllipticF(b^(1/4)*x/a^(1/4),I)/a^(7/4)/b^(1/4)/(-b*x^4+a)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.03 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.76

$$\int \frac{1}{(a-bx^4)^{5/2}} dx = \frac{7ax - 5bx^5 + 5x(a-bx^4)\sqrt{1-\frac{bx^4}{a}} \operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, \frac{bx^4}{a}\right)}{12a^2(a-bx^4)^{3/2}}$$

input

```
Integrate[(a - b*x^4)^(-5/2), x]
```

output

```
(7*a*x - 5*b*x^5 + 5*x*(a - b*x^4)*Sqrt[1 - (b*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, (b*x^4)/a])/(12*a^2*(a - b*x^4)^(3/2))
```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.08, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {749, 749, 765, 762}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a - bx^4)^{5/2}} dx \\
 & \quad \downarrow \text{749} \\
 & \frac{5 \int \frac{1}{(a - bx^4)^{3/2}} dx}{6a} + \frac{x}{6a(a - bx^4)^{3/2}} \\
 & \quad \downarrow \text{749} \\
 & \frac{5 \left(\frac{\int \frac{1}{\sqrt{a - bx^4}} dx}{2a} + \frac{x}{2a\sqrt{a - bx^4}} \right)}{6a} + \frac{x}{6a(a - bx^4)^{3/2}} \\
 & \quad \downarrow \text{765} \\
 & \frac{5 \left(\frac{\sqrt{1 - \frac{bx^4}{a}} \int \frac{1}{\sqrt{1 - \frac{bx^4}{a}}} dx}{2a\sqrt{a - bx^4}} + \frac{x}{2a\sqrt{a - bx^4}} \right)}{6a} + \frac{x}{6a(a - bx^4)^{3/2}} \\
 & \quad \downarrow \text{762} \\
 & \frac{5 \left(\frac{\sqrt{1 - \frac{bx^4}{a}} \operatorname{EllipticF} \left(\arcsin \left(\frac{\sqrt[4]{bx^4}}{\sqrt[4]{a}} \right), -1 \right)}{2a^{3/4} \sqrt[4]{b} \sqrt{a - bx^4}} + \frac{x}{2a\sqrt{a - bx^4}} \right)}{6a} + \frac{x}{6a(a - bx^4)^{3/2}}
 \end{aligned}$$

input

```
Int[(a - b*x^4)^(-5/2), x]
```

output
$$\frac{x/(6*a*(a - b*x^4)^{(3/2)}) + (5*(x/(2*a*\text{Sqrt}[a - b*x^4]) + (\text{Sqrt}[1 - (b*x^4)/a]*\text{EllipticF}[\text{ArcSin}[(b^{(1/4)}*x)/a^{(1/4)}], -1]))/(2*a^{(3/4)}*b^{(1/4)}*\text{Sqrt}[a - b*x^4])))/(6*a)}$$

Defintions of rubi rules used

rule 749
$$\text{Int}[(a_ + (b_)*(x_)^{(n)})^{(p)}, x_Symbol] \rightarrow \text{Simp}[(-x)*((a + b*x^n)^{(p + 1))/(a*n*(p + 1))], x] + \text{Simp}[(n*(p + 1) + 1)/(a*n*(p + 1)) \text{Int}[(a + b*x^n)^{(p + 1)}, x], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ (\text{IntegerQ}[2*p] \ || \ \text{Denominator}[p + 1/n] < \text{Denominator}[p])$$

rule 762
$$\text{Int}[1/\text{Sqrt}[(a_ + (b_)*(x_)^4], x_Symbol] \rightarrow \text{Simp}[(1/(\text{Sqrt}[a]*\text{Rt}[-b/a, 4]))*\text{EllipticF}[\text{ArcSin}[\text{Rt}[-b/a, 4]*x], -1], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[b/a] \ \&\& \ \text{GtQ}[a, 0]$$

rule 765
$$\text{Int}[1/\text{Sqrt}[(a_ + (b_)*(x_)^4], x_Symbol] \rightarrow \text{Simp}[\text{Sqrt}[1 + b*(x^4/a)]/\text{Sqrt}[a + b*x^4] \text{Int}[1/\text{Sqrt}[1 + b*(x^4/a)], x], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[b/a] \ \&\& \ !\text{GtQ}[a, 0]$$

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.25

method	result	size
default	$\frac{x\sqrt{-bx^4+a}}{6ab^2(x^4-\frac{a}{b})^2} + \frac{5x}{12a^2\sqrt{-(x^4-\frac{a}{b})b}} + \frac{5\sqrt{1-\frac{\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1+\frac{\sqrt{b}x^2}{\sqrt{a}}}\text{EllipticF}\left(x\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}, i\right)}{12a^2\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}\sqrt{-bx^4+a}}$	121
elliptic	$\frac{x\sqrt{-bx^4+a}}{6ab^2(x^4-\frac{a}{b})^2} + \frac{5x}{12a^2\sqrt{-(x^4-\frac{a}{b})b}} + \frac{5\sqrt{1-\frac{\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1+\frac{\sqrt{b}x^2}{\sqrt{a}}}\text{EllipticF}\left(x\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}, i\right)}{12a^2\sqrt{\frac{\sqrt{b}}{\sqrt{a}}}\sqrt{-bx^4+a}}$	121

input
$$\text{int}(1/(-b*x^4+a)^{(5/2)}, x, \text{method}=_RETURNVERBOSE)$$

output $\frac{1}{6}x/a/b^2*(-b*x^4+a)^{(1/2)}/(x^4-a/b)^2+5/12*x/a^2/(-(x^4-a/b)*b)^{(1/2)}+5/12/a^2/(1/a^{(1/2)}*b^{(1/2)})^{(1/2)}*(1-b^{(1/2)}*x^2/a^{(1/2)})^{(1/2)}*(1+b^{(1/2)}*x^2/a^{(1/2)})^{(1/2)}/(-b*x^4+a)^{(1/2)}*EllipticF(x*(1/a^{(1/2)}*b^{(1/2)})^{(1/2)},I)$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.03

$$\int \frac{1}{(a - bx^4)^{5/2}} dx = \frac{5(b^2x^8 - 2abx^4 + a^2)\sqrt{a}\left(\frac{b}{a}\right)^{3/4} F(\arcsin\left(x\left(\frac{b}{a}\right)^{1/4}\right) | -1) - (5b^2x^5 - 7abx)\sqrt{-bx^4 + a}}{12(a^2b^3x^8 - 2a^3b^2x^4 + a^4b)}$$

input `integrate(1/(-b*x^4+a)^(5/2),x, algorithm="fricas")`

output $\frac{1}{12}*(5*(b^2*x^8 - 2*a*b*x^4 + a^2)*sqrt(a)*(b/a)^{(3/4)}*elliptic_f(arcsin(x*(b/a)^{(1/4)}), -1) - (5*b^2*x^5 - 7*a*b*x)*sqrt(-b*x^4 + a))/(a^2*b^3*x^8 - 2*a^3*b^2*x^4 + a^4*b)$

Sympy [A] (verification not implemented)

Time = 0.61 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.38

$$\int \frac{1}{(a - bx^4)^{5/2}} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{5}{2} \middle| \frac{bx^4 e^{2i\pi}}{a}\right)}{4a^{5/2}\Gamma\left(\frac{5}{4}\right)}$$

input `integrate(1/(-b*x**4+a)**(5/2),x)`

output $x*\gamma(1/4)*hyper((1/4, 5/2), (5/4,), b*x**4*exp_polar(2*I*pi)/a)/(4*a**5/2)*\gamma(5/4)$

Maxima [F]

$$\int \frac{1}{(a - bx^4)^{5/2}} dx = \int \frac{1}{(-bx^4 + a)^{\frac{5}{2}}} dx$$

input `integrate(1/(-b*x^4+a)^(5/2),x, algorithm="maxima")`

output `integrate((-b*x^4 + a)^(-5/2), x)`

Giac [F]

$$\int \frac{1}{(a - bx^4)^{5/2}} dx = \int \frac{1}{(-bx^4 + a)^{\frac{5}{2}}} dx$$

input `integrate(1/(-b*x^4+a)^(5/2),x, algorithm="giac")`

output `integrate((-b*x^4 + a)^(-5/2), x)`

Mupad [B] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.39

$$\int \frac{1}{(a - bx^4)^{5/2}} dx = \frac{x \left(1 - \frac{bx^4}{a}\right)^{5/2} {}_2F_1\left(\frac{1}{4}, \frac{5}{2}, \frac{5}{4}, \frac{bx^4}{a}\right)}{(a - bx^4)^{5/2}}$$

input `int(1/(a - b*x^4)^(5/2),x)`

output `(x*(1 - (b*x^4)/a)^(5/2)*hypergeom([1/4, 5/2], 5/4, (b*x^4)/a))/(a - b*x^4)^(5/2)`

Reduce [F]

$$\int \frac{1}{(a - bx^4)^{5/2}} dx = \int \frac{\sqrt{-bx^4 + a}}{-b^3x^{12} + 3ab^2x^8 - 3a^2bx^4 + a^3} dx$$

input `int(1/(-b*x^4+a)^(5/2),x)`

output `int(sqrt(a - b*x**4)/(a**3 - 3*a**2*b*x**4 + 3*a*b**2*x**8 - b**3*x**12),x)`

3.14 $\int (a + bx^4)^{3/2} dx$

Optimal result	135
Mathematica [C] (verified)	135
Rubi [A] (verified)	136
Maple [C] (verified)	137
Fricas [A] (verification not implemented)	138
Sympy [C] (verification not implemented)	138
Maxima [F]	139
Giac [F]	139
Mupad [B] (verification not implemented)	139
Reduce [F]	140

Optimal result

Integrand size = 11, antiderivative size = 122

$$\int (a + bx^4)^{3/2} dx = \frac{2}{7}ax\sqrt{a + bx^4} + \frac{1}{7}x(a + bx^4)^{3/2} + \frac{2a^{7/4}(\sqrt{a} + \sqrt{bx^2})\sqrt{\frac{a+bx^4}{(\sqrt{a}+\sqrt{bx^2})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{7\sqrt[4]{b}\sqrt{a + bx^4}}$$

output

```
2/7*a*x*(b*x^4+a)^(1/2)+1/7*x*(b*x^4+a)^(3/2)+2/7*a^(7/4)*(a^(1/2)+b^(1/2)*x^2)*((b*x^4+a)/(a^(1/2)+b^(1/2)*x^2)^2)^(1/2)*InverseJacobiAM(2*arctan(b^(1/4)*x/a^(1/4)),1/2*2^(1/2))/b^(1/4)/(b*x^4+a)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 5.41 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.39

$$\int (a + bx^4)^{3/2} dx = \frac{ax\sqrt{a + bx^4}\text{Hypergeometric2F1}\left(-\frac{3}{2}, \frac{1}{4}, \frac{5}{4}, -\frac{bx^4}{a}\right)}{\sqrt{1 + \frac{bx^4}{a}}}$$

input `Integrate[(a + b*x^4)^(3/2),x]`

output `(a*x*Sqrt[a + b*x^4]*Hypergeometric2F1[-3/2, 1/4, 5/4, -(b*x^4)/a])/Sqrt[1 + (b*x^4)/a]`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.04, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {748, 748, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + bx^4)^{3/2} dx \\
 & \quad \downarrow 748 \\
 & \frac{6}{7}a \int \sqrt{bx^4 + a} dx + \frac{1}{7}x(a + bx^4)^{3/2} \\
 & \quad \downarrow 748 \\
 & \frac{6}{7}a \left(\frac{2}{3}a \int \frac{1}{\sqrt{bx^4 + a}} dx + \frac{1}{3}x\sqrt{a + bx^4} \right) + \frac{1}{7}x(a + bx^4)^{3/2} \\
 & \quad \downarrow 761 \\
 & \frac{6}{7}a \left(\frac{a^{3/4}(\sqrt{a} + \sqrt{bx^2}) \sqrt{\frac{a+bx^4}{(\sqrt{a}+\sqrt{bx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{3\sqrt[4]{b}\sqrt{a + bx^4}} + \frac{1}{3}x\sqrt{a + bx^4} \right) + \\
 & \quad \frac{1}{7}x(a + bx^4)^{3/2}
 \end{aligned}$$

input `Int[(a + b*x^4)^(3/2),x]`

output

$$\frac{(x*(a + b*x^4)^{(3/2)})/7 + (6*a*((x*\text{Sqrt}[a + b*x^4])/3 + (a^{(3/4)}*(\text{Sqrt}[a + \text{Sqrt}[b]*x^2)*\text{Sqrt}[(a + b*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[b]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(b^{(1/4)}*x)/a^{(1/4)}], 1/2]))/(3*b^{(1/4)}*\text{Sqrt}[a + b*x^4]))/7$$

Defintions of rubi rules used

rule 748

$$\text{Int}[(a_+ + (b_+)*(x_+)^{(n_+))^{(p_+)}, x_Symbol] \rightarrow \text{Simp}[x*((a + b*x^n)^p/(n*p + 1)), x] + \text{Simp}[a*n*(p/(n*p + 1)) \text{Int}[(a + b*x^n)^{(p-1)}, x], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{IGtQ}[n, 0] \&\& \text{GtQ}[p, 0] \&\& (\text{IntegerQ}[2*p] \parallel \text{LtQ}[\text{Denominator}[p + 1/n], \text{Denominator}[p]])$$

rule 761

$$\text{Int}[1/\text{Sqrt}[(a_+ + (b_+)*(x_+)^4], x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b/a, 4]\}, \text{Simp}[(1 + q^2*x^2)*(\text{Sqrt}[(a + b*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*\text{Sqrt}[a + b*x^4]))*\text{EllipticF}[2*\text{ArcTan}[q*x], 1/2], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[b/a]$$

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.55 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.79

method	result	size
risch	$\frac{x(bx^4+3a)\sqrt{bx^4+a}}{7} + \frac{4a^2\sqrt{1-\frac{i\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{b}x^2}{\sqrt{a}}}\text{EllipticF}\left(x\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}, i\right)}{7\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}\sqrt{bx^4+a}}$	96
default	$\frac{bx^5\sqrt{bx^4+a}}{7} + \frac{3ax\sqrt{bx^4+a}}{7} + \frac{4a^2\sqrt{1-\frac{i\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{b}x^2}{\sqrt{a}}}\text{EllipticF}\left(x\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}, i\right)}{7\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}\sqrt{bx^4+a}}$	103
elliptic	$\frac{bx^5\sqrt{bx^4+a}}{7} + \frac{3ax\sqrt{bx^4+a}}{7} + \frac{4a^2\sqrt{1-\frac{i\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{b}x^2}{\sqrt{a}}}\text{EllipticF}\left(x\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}, i\right)}{7\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}\sqrt{bx^4+a}}$	103

input

$$\text{int}((b*x^4+a)^{(3/2)}, x, \text{method}=_RETURNVERBOSE)$$

output `1/7*x*(b*x^4+3*a)*(b*x^4+a)^(1/2)+4/7*a^2/(I/a^(1/2)*b^(1/2))^(1/2)*(1-I*b^(1/2)*x^2/a^(1/2))^(1/2)*(1+I*b^(1/2)*x^2/a^(1/2))^(1/2)/(b*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*b^(1/2))^(1/2),I)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.42

$$\int (a + bx^4)^{3/2} dx = \frac{4}{7} a\sqrt{b} \left(-\frac{a}{b}\right)^{\frac{3}{4}} F(\arcsin\left(\frac{\left(-\frac{a}{b}\right)^{\frac{1}{4}}}{x}\right) | -1) + \frac{1}{7} (bx^5 + 3ax)\sqrt{bx^4 + a}$$

input `integrate((b*x^4+a)^(3/2),x, algorithm="fricas")`

output `4/7*a*sqrt(b)*(-a/b)^(3/4)*elliptic_f(arcsin((-a/b)^(1/4)/x), -1) + 1/7*(b*x^5 + 3*a*x)*sqrt(b*x^4 + a)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.52 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.30

$$\int (a + bx^4)^{3/2} dx = \frac{a^{\frac{3}{2}} x \Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\begin{matrix} -\frac{3}{2}, \frac{1}{4} \\ \frac{5}{4} \end{matrix} \middle| \frac{bx^4 e^{i\pi}}{a} \right)}{4\Gamma\left(\frac{5}{4}\right)}$$

input `integrate((b*x**4+a)**(3/2),x)`

output `a**(3/2)*x*gamma(1/4)*hyper((-3/2, 1/4), (5/4,), b*x**4*exp_polar(I*pi)/a)/(4*gamma(5/4))`

Maxima [F]

$$\int (a + bx^4)^{3/2} dx = \int (bx^4 + a)^{\frac{3}{2}} dx$$

input `integrate((b*x^4+a)^(3/2),x, algorithm="maxima")`

output `integrate((b*x^4 + a)^(3/2), x)`

Giac [F]

$$\int (a + bx^4)^{3/2} dx = \int (bx^4 + a)^{\frac{3}{2}} dx$$

input `integrate((b*x^4+a)^(3/2),x, algorithm="giac")`

output `integrate((b*x^4 + a)^(3/2), x)`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.30

$$\int (a + bx^4)^{3/2} dx = \frac{x (bx^4 + a)^{3/2} {}_2F_1\left(-\frac{3}{2}, \frac{1}{4}; \frac{5}{4}; -\frac{bx^4}{a}\right)}{\left(\frac{bx^4}{a} + 1\right)^{3/2}}$$

input `int((a + b*x^4)^(3/2),x)`

output `(x*(a + b*x^4)^(3/2)*hypergeom([-3/2, 1/4], 5/4, -(b*x^4)/a))/((b*x^4)/a + 1)^(3/2)`

Reduce [F]

$$\int (a + bx^4)^{3/2} dx = \frac{3\sqrt{bx^4 + a} ax}{7} + \frac{\sqrt{bx^4 + a} bx^5}{7} + \frac{4\left(\int \frac{\sqrt{bx^4 + a}}{bx^4 + a} dx\right) a^2}{7}$$

input `int((b*x^4+a)^(3/2),x)`

output `(3*sqrt(a + b*x**4)*a*x + sqrt(a + b*x**4)*b*x**5 + 4*int(sqrt(a + b*x**4)/(a + b*x**4),x)*a**2)/7`

3.15 $\int \sqrt{a + bx^4} dx$

Optimal result	141
Mathematica [C] (verified)	141
Rubi [A] (verified)	142
Maple [C] (verified)	143
Fricas [A] (verification not implemented)	144
Sympy [C] (verification not implemented)	144
Maxima [F]	145
Giac [F]	145
Mupad [B] (verification not implemented)	145
Reduce [F]	146

Optimal result

Integrand size = 11, antiderivative size = 105

$$\int \sqrt{a + bx^4} dx = \frac{1}{3}x\sqrt{a + bx^4} + \frac{a^{3/4}(\sqrt{a} + \sqrt{bx^2}) \sqrt{\frac{a+bx^4}{(\sqrt{a}+\sqrt{bx^2})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{3\sqrt[4]{b}\sqrt{a + bx^4}}$$

output

```
1/3*x*(b*x^4+a)^(1/2)+1/3*a^(3/4)*(a^(1/2)+b^(1/2)*x^2)*((b*x^4+a)/(a^(1/2)+b^(1/2)*x^2)^2)^(1/2)*InverseJacobiAM(2*arctan(b^(1/4)*x/a^(1/4)),1/2*2^(1/2))/b^(1/4)/(b*x^4+a)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.04 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.85

$$\int \sqrt{a + bx^4} dx = \frac{x(a + bx^4) - \frac{2ia\sqrt{1+\frac{bx^4}{a}} \operatorname{EllipticF}\left(i\operatorname{arcsinh}\left(\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}x\right), -1\right)}{\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}}}{3\sqrt{a + bx^4}}$$

input `Integrate[Sqrt[a + b*x^4], x]`

output `(x*(a + b*x^4) - ((2*I)*a*Sqrt[1 + (b*x^4)/a]*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[b])/Sqrt[a]]*x], -1])/Sqrt[(I*Sqrt[b])/Sqrt[a]]/(3*Sqrt[a + b*x^4])`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {748, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{a + bx^4} dx$$

$$\downarrow 748$$

$$\frac{2}{3}a \int \frac{1}{\sqrt{bx^4 + a}} dx + \frac{1}{3}x\sqrt{a + bx^4}$$

$$\downarrow 761$$

$$\frac{a^{3/4}(\sqrt{a} + \sqrt{bx^2}) \sqrt{\frac{a+bx^4}{(\sqrt{a}+\sqrt{bx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{3\sqrt[4]{b}\sqrt{a + bx^4}} + \frac{1}{3}x\sqrt{a + bx^4}$$

input `Int[Sqrt[a + b*x^4], x]`

output `(x*Sqrt[a + b*x^4])/3 + (a^(3/4)*(Sqrt[a] + Sqrt[b]*x^2)*Sqrt[(a + b*x^4)/(Sqrt[a] + Sqrt[b]*x^2)^2]*EllipticF[2*ArcTan[(b^(1/4)*x)/a^(1/4)], 1/2])/(3*b^(1/4)*Sqrt[a + b*x^4])`

Definitions of rubi rules used

rule 748

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p
+ 1)), x] + Simp[a*n*(p/(n*p + 1)) Int[(a + b*x^n)^(p - 1), x], x] /; Fre
eQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || LtQ[Denominat
or[p + 1/n], Denominator[p]])
```

rule 761

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(
1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*
EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]
```

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.53 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.81

method	result	size
default	$\frac{x\sqrt{bx^4+a}}{3} + \frac{2a\sqrt{1-\frac{i\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{b}x^2}{\sqrt{a}}}\operatorname{EllipticF}\left(x\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}, i\right)}{3\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}\sqrt{bx^4+a}}$	85
risch	$\frac{x\sqrt{bx^4+a}}{3} + \frac{2a\sqrt{1-\frac{i\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{b}x^2}{\sqrt{a}}}\operatorname{EllipticF}\left(x\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}, i\right)}{3\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}\sqrt{bx^4+a}}$	85
elliptic	$\frac{x\sqrt{bx^4+a}}{3} + \frac{2a\sqrt{1-\frac{i\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{b}x^2}{\sqrt{a}}}\operatorname{EllipticF}\left(x\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}, i\right)}{3\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}\sqrt{bx^4+a}}$	85

input `int((b*x^4+a)^(1/2), x, method=_RETURNVERBOSE)`

output

```
1/3*x*(b*x^4+a)^(1/2)+2/3*a/(I/a^(1/2)*b^(1/2))^(1/2)*(1-I*b^(1/2)*x^2/a^(
1/2))^(1/2)*(1+I*b^(1/2)*x^2/a^(1/2))^(1/2)/(b*x^4+a)^(1/2)*EllipticF(x*(I
/a^(1/2)*b^(1/2))^(1/2), I)
```


Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.39

$$\int \sqrt{a + bx^4} dx = \frac{2}{3} \sqrt{b} \left(-\frac{a}{b}\right)^{\frac{3}{4}} F(\arcsin\left(\frac{\left(-\frac{a}{b}\right)^{\frac{1}{4}}}{x}\right) | -1) + \frac{1}{3} \sqrt{bx^4 + ax}$$

input `integrate((b*x^4+a)^(1/2),x, algorithm="fricas")`

output `2/3*sqrt(b)*(-a/b)^(3/4)*elliptic_f(arcsin((-a/b)^(1/4)/x), -1) + 1/3*sqrt(b*x^4 + a)*x`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.47 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.35

$$\int \sqrt{a + bx^4} dx = \frac{\sqrt{ax} \Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\begin{matrix} -\frac{1}{2}, \frac{1}{4} \\ \frac{5}{4} \end{matrix} \middle| \frac{bx^4 e^{i\pi}}{a}\right)}{4\Gamma\left(\frac{5}{4}\right)}$$

input `integrate((b*x**4+a)**(1/2),x)`

output `sqrt(a)*x*gamma(1/4)*hyper((-1/2, 1/4), (5/4,), b*x**4*exp_polar(I*pi)/a)/(4*gamma(5/4))`

Maxima [F]

$$\int \sqrt{a + bx^4} dx = \int \sqrt{bx^4 + a} dx$$

input `integrate((b*x^4+a)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(b*x^4 + a), x)`

Giac [F]

$$\int \sqrt{a + bx^4} dx = \int \sqrt{bx^4 + a} dx$$

input `integrate((b*x^4+a)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*x^4 + a), x)`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.35

$$\int \sqrt{a + bx^4} dx = \frac{x \sqrt{bx^4 + a} {}_2F_1\left(-\frac{1}{2}, \frac{1}{4}; \frac{5}{4}; -\frac{bx^4}{a}\right)}{\sqrt{\frac{bx^4}{a} + 1}}$$

input `int((a + b*x^4)^(1/2),x)`

output `(x*(a + b*x^4)^(1/2)*hypergeom([-1/2, 1/4], 5/4, -(b*x^4)/a))/((b*x^4)/a + 1)^(1/2)`

Reduce [F]

$$\int \sqrt{a + bx^4} dx = \frac{\sqrt{bx^4 + a} x}{3} + \frac{2 \left(\int \frac{\sqrt{bx^4 + a}}{bx^4 + a} dx \right) a}{3}$$

input `int((b*x^4+a)^(1/2),x)`

output `(sqrt(a + b*x**4)*x + 2*int(sqrt(a + b*x**4)/(a + b*x**4),x)*a)/3`

3.16 $\int \frac{1}{\sqrt{a+bx^4}} dx$

Optimal result	147
Mathematica [C] (verified)	147
Rubi [A] (verified)	148
Maple [C] (verified)	149
Fricas [A] (verification not implemented)	149
Sympy [C] (verification not implemented)	150
Maxima [F]	150
Giac [F]	150
Mupad [B] (verification not implemented)	151
Reduce [F]	151

Optimal result

Integrand size = 11, antiderivative size = 88

$$\int \frac{1}{\sqrt{a+bx^4}} dx = \frac{(\sqrt{a} + \sqrt{bx^2}) \sqrt{\frac{a+bx^4}{(\sqrt{a}+\sqrt{bx^2})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{b}x}{\sqrt{a}}\right), \frac{1}{2}\right)}{2\sqrt[4]{a}\sqrt[4]{b}\sqrt{a+bx^4}}$$

output

```
1/2*(a^(1/2)+b^(1/2)*x^2)*((b*x^4+a)/(a^(1/2)+b^(1/2)*x^2)^2)^(1/2)*Invers
eJacobiAM(2*arctan(b^(1/4)*x/a^(1/4)),1/2*2^(1/2))/a^(1/4)/b^(1/4)/(b*x^4+
a)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 10.05 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.84

$$\int \frac{1}{\sqrt{a+bx^4}} dx = -\frac{i\sqrt{1+\frac{bx^4}{a}} \operatorname{EllipticF}\left(i \operatorname{arcsinh}\left(\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}x\right), -1\right)}{\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}\sqrt{a+bx^4}}$$

input

```
Integrate[1/Sqrt[a + b*x^4], x]
```

output $((-I)*\text{Sqrt}[1 + (b*x^4)/a]*\text{EllipticF}[I*\text{ArcSinh}[\text{Sqrt}[(I*\text{Sqrt}[b])/ \text{Sqrt}[a]]*x, -1])/(\text{Sqrt}[(I*\text{Sqrt}[b])/ \text{Sqrt}[a]]*\text{Sqrt}[a + b*x^4])$

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{a + bx^4}} dx$$

↓ 761

$$\frac{(\sqrt{a} + \sqrt{bx^2}) \sqrt{\frac{a+bx^4}{(\sqrt{a}+\sqrt{bx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2\sqrt[4]{a}\sqrt[4]{b}\sqrt{a + bx^4}}$$

input `Int[1/Sqrt[a + b*x^4], x]`

output $((\text{Sqrt}[a] + \text{Sqrt}[b]*x^2)*\text{Sqrt}[(a + b*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[b]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(b^{1/4})*x]/a^{1/4}], 1/2))/(2*a^{1/4}*b^{1/4}*\text{Sqrt}[a + b*x^4])$

Defintions of rubi rules used

rule 761 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] :> With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2]]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.46 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.80

method	result	size
default	$\frac{\sqrt{1 - \frac{i\sqrt{b}x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{b}x^2}{\sqrt{a}}} \operatorname{EllipticF}\left(x\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}, i\right)}{\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}} \sqrt{bx^4 + a}}$	70
elliptic	$\frac{\sqrt{1 - \frac{i\sqrt{b}x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{b}x^2}{\sqrt{a}}} \operatorname{EllipticF}\left(x\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}, i\right)}{\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}} \sqrt{bx^4 + a}}$	70

input `int(1/(b*x^4+a)^(1/2), x, method=_RETURNVERBOSE)`

output `1/(I/a^(1/2)*b^(1/2))^(1/2)*(1-I*b^(1/2)*x^2/a^(1/2))^(1/2)*(1+I*b^(1/2)*x^2/a^(1/2))^(1/2)/(b*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*b^(1/2))^(1/2), I)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.33

$$\int \frac{1}{\sqrt{a + bx^4}} dx = -\frac{\sqrt{a}\left(-\frac{b}{a}\right)^{\frac{3}{4}} F\left(\arcsin\left(x\left(-\frac{b}{a}\right)^{\frac{1}{4}}\right) \mid -1\right)}{b}$$

input `integrate(1/(b*x^4+a)^(1/2), x, algorithm="fricas")`

output `-sqrt(a)*(-b/a)^(3/4)*elliptic_f(arcsin(x*(-b/a)^(1/4)), -1)/b`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.48 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.41

$$\int \frac{1}{\sqrt{a+bx^4}} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2} \middle| \frac{bx^4 e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{5}{4}\right)}$$

input `integrate(1/(b*x**4+a)**(1/2),x)`

output `x*gamma(1/4)*hyper((1/4, 1/2), (5/4,), b*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(5/4))`

Maxima [F]

$$\int \frac{1}{\sqrt{a+bx^4}} dx = \int \frac{1}{\sqrt{bx^4+a}} dx$$

input `integrate(1/(b*x^4+a)^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(b*x^4 + a), x)`

Giac [F]

$$\int \frac{1}{\sqrt{a+bx^4}} dx = \int \frac{1}{\sqrt{bx^4+a}} dx$$

input `integrate(1/(b*x^4+a)^(1/2),x, algorithm="giac")`

output `integrate(1/sqrt(b*x^4 + a), x)`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.42

$$\int \frac{1}{\sqrt{a + bx^4}} dx = \frac{x \sqrt{\frac{bx^4}{a} + 1} {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; -\frac{bx^4}{a}\right)}{\sqrt{bx^4 + a}}$$

input `int(1/(a + b*x^4)^(1/2),x)`output `(x*((b*x^4)/a + 1)^(1/2)*hypergeom([1/4, 1/2], 5/4, -(b*x^4)/a))/(a + b*x^4)^(1/2)`**Reduce [F]**

$$\int \frac{1}{\sqrt{a + bx^4}} dx = \int \frac{\sqrt{bx^4 + a}}{bx^4 + a} dx$$

input `int(1/(b*x^4+a)^(1/2),x)`output `int(sqrt(a + b*x**4)/(a + b*x**4),x)`

$$3.17 \quad \int \frac{1}{(a+bx^4)^{3/2}} dx$$

Optimal result	152
Mathematica [C] (verified)	152
Rubi [A] (verified)	153
Maple [C] (verified)	154
Fricas [A] (verification not implemented)	155
Sympy [C] (verification not implemented)	155
Maxima [F]	156
Giac [F]	156
Mupad [B] (verification not implemented)	156
Reduce [F]	157

Optimal result

Integrand size = 11, antiderivative size = 108

$$\int \frac{1}{(a+bx^4)^{3/2}} dx = \frac{x}{2a\sqrt{a+bx^4}} + \frac{(\sqrt{a} + \sqrt{bx^2}) \sqrt{\frac{a+bx^4}{(\sqrt{a} + \sqrt{bx^2})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{4a^{5/4} \sqrt[4]{b} \sqrt{a+bx^4}}$$

output

```
1/2*x/a/(b*x^4+a)^(1/2)+1/4*(a^(1/2)+b^(1/2)*x^2)*((b*x^4+a)/(a^(1/2)+b^(1/2)*x^2)^2)^(1/2)*InverseJacobiAM(2*arctan(b^(1/4)*x/a^(1/4)),1/2*2^(1/2))/a^(5/4)/b^(1/4)/(b*x^4+a)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 5.68 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.51

$$\int \frac{1}{(a+bx^4)^{3/2}} dx = \frac{x + x \sqrt{1 + \frac{bx^4}{a}} \operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, -\frac{bx^4}{a}\right)}{2a\sqrt{a+bx^4}}$$

input `Integrate[(a + b*x^4)^(-3/2),x]`

output `(x + x*Sqrt[1 + (b*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, -((b*x^4)/a)]) / (2*a*Sqrt[a + b*x^4])`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {749, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + bx^4)^{3/2}} dx$$

$$\downarrow 749$$

$$\frac{\int \frac{1}{\sqrt{bx^4+a}} dx}{2a} + \frac{x}{2a\sqrt{a + bx^4}}$$

$$\downarrow 761$$

$$\frac{(\sqrt{a} + \sqrt{bx^2}) \sqrt{\frac{a+bx^4}{(\sqrt{a}+\sqrt{bx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{4a^{5/4}\sqrt[4]{b}\sqrt{a + bx^4}} + \frac{x}{2a\sqrt{a + bx^4}}$$

input `Int[(a + b*x^4)^(-3/2),x]`

output `x/(2*a*Sqrt[a + b*x^4]) + ((Sqrt[a] + Sqrt[b]*x^2)*Sqrt[(a + b*x^4)/(Sqrt[a] + Sqrt[b]*x^2)^2]*EllipticF[2*ArcTan[(b^(1/4)*x)/a^(1/4)], 1/2])/(4*a^(5/4)*b^(1/4)*Sqrt[a + b*x^4])`

Definitions of rubi rules used

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

rule 761 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.46 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.87

method	result	size
default	$\frac{x}{2a\sqrt{(x^4 + \frac{a}{b})b}} + \frac{\sqrt{1 - \frac{i\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{b}x^2}{\sqrt{a}}}\operatorname{EllipticF}\left(x\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}, i\right)}{2a\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}\sqrt{bx^4 + a}}$	94
elliptic	$\frac{x}{2a\sqrt{(x^4 + \frac{a}{b})b}} + \frac{\sqrt{1 - \frac{i\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{b}x^2}{\sqrt{a}}}\operatorname{EllipticF}\left(x\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}, i\right)}{2a\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}\sqrt{bx^4 + a}}$	94

input `int(1/(b*x^4+a)^(3/2),x,method=_RETURNVERBOSE)`

output `1/2*x/a/((x^4+a/b)*b)^(1/2)+1/2/a/(I/a^(1/2)*b^(1/2))^(1/2)*(1-I*b^(1/2)*x^2/a^(1/2))^(1/2)*(1+I*b^(1/2)*x^2/a^(1/2))^(1/2)/(b*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*b^(1/2))^(1/2),I)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.59

$$\int \frac{1}{(a + bx^4)^{3/2}} dx = -\frac{(bx^4 + a)\sqrt{a}\left(-\frac{b}{a}\right)^{\frac{3}{4}} F(\arcsin\left(x\left(-\frac{b}{a}\right)^{\frac{1}{4}}\right) | -1) - \sqrt{bx^4 + abx}}{2(ab^2x^4 + a^2b)}$$

input `integrate(1/(b*x^4+a)^(3/2),x, algorithm="fricas")`

output `-1/2*((b*x^4 + a)*sqrt(a)*(-b/a)^(3/4)*elliptic_f(arcsin(x*(-b/a)^(1/4)), -1) - sqrt(b*x^4 + a)*b*x)/(a*b^2*x^4 + a^2*b)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.53 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.33

$$\int \frac{1}{(a + bx^4)^{3/2}} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{3}{2} \middle| \frac{bx^4 e^{i\pi}}{a}\right)}{4a^{\frac{3}{2}}\Gamma\left(\frac{5}{4}\right)}$$

input `integrate(1/(b*x**4+a)**(3/2),x)`

output `x*gamma(1/4)*hyper((1/4, 3/2), (5/4,), b*x**4*exp_polar(I*pi)/a)/(4*a**(3/2)*gamma(5/4))`

Maxima [F]

$$\int \frac{1}{(a + bx^4)^{3/2}} dx = \int \frac{1}{(bx^4 + a)^{\frac{3}{2}}} dx$$

input `integrate(1/(b*x^4+a)^(3/2),x, algorithm="maxima")`

output `integrate((b*x^4 + a)^(-3/2), x)`

Giac [F]

$$\int \frac{1}{(a + bx^4)^{3/2}} dx = \int \frac{1}{(bx^4 + a)^{\frac{3}{2}}} dx$$

input `integrate(1/(b*x^4+a)^(3/2),x, algorithm="giac")`

output `integrate((b*x^4 + a)^(-3/2), x)`

Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.34

$$\int \frac{1}{(a + bx^4)^{3/2}} dx = \frac{x \left(\frac{bx^4}{a} + 1 \right)^{3/2} {}_2F_1 \left(\frac{1}{4}, \frac{3}{2}; \frac{5}{4}; -\frac{bx^4}{a} \right)}{(bx^4 + a)^{3/2}}$$

input `int(1/(a + b*x^4)^(3/2),x)`

output `(x*((b*x^4)/a + 1)^(3/2)*hypergeom([1/4, 3/2], 5/4, -(b*x^4)/a))/(a + b*x^4)^(3/2)`

Reduce [F]

$$\int \frac{1}{(a + bx^4)^{3/2}} dx = \int \frac{\sqrt{bx^4 + a}}{b^2x^8 + 2abx^4 + a^2} dx$$

input `int(1/(b*x^4+a)^(3/2),x)`

output `int(sqrt(a + b*x**4)/(a**2 + 2*a*b*x**4 + b**2*x**8),x)`

3.18 $\int \frac{1}{(a+bx^4)^{5/2}} dx$

Optimal result	158
Mathematica [C] (verified)	158
Rubi [A] (verified)	159
Maple [C] (verified)	160
Fricas [A] (verification not implemented)	161
Sympy [C] (verification not implemented)	161
Maxima [F]	162
Giac [F]	162
Mupad [B] (verification not implemented)	162
Reduce [F]	163

Optimal result

Integrand size = 11, antiderivative size = 127

$$\int \frac{1}{(a+bx^4)^{5/2}} dx = \frac{x}{6a(a+bx^4)^{3/2}} + \frac{5x}{12a^2\sqrt{a+bx^4}} + \frac{5(\sqrt{a} + \sqrt{bx^2}) \sqrt{\frac{a+bx^4}{(\sqrt{a}+\sqrt{bx^2})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{24a^{9/4}\sqrt[4]{b}\sqrt{a+bx^4}}$$

output

```
1/6*x/a/(b*x^4+a)^(3/2)+5/12*x/a^2/(b*x^4+a)^(1/2)+5/24*(a^(1/2)+b^(1/2)*x^2)*((b*x^4+a)/(a^(1/2)+b^(1/2)*x^2)^(1/2)*InverseJacobiAM(2*arctan(b^(1/4)*x/a^(1/4)),1/2*2^(1/2))/a^(9/4)/b^(1/4)/(b*x^4+a)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.03 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.57

$$\int \frac{1}{(a+bx^4)^{5/2}} dx = \frac{7ax + 5bx^5 + 5x(a+bx^4) \sqrt{1 + \frac{bx^4}{a}} \operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, -\frac{bx^4}{a}\right)}{12a^2(a+bx^4)^{3/2}}$$

input `Integrate[(a + b*x^4)^(-5/2),x]`

output `(7*a*x + 5*b*x^5 + 5*x*(a + b*x^4)*Sqrt[1 + (b*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, -((b*x^4)/a)]/(12*a^2*(a + b*x^4)^(3/2))`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.06, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {749, 749, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a + bx^4)^{5/2}} dx \\
 & \quad \downarrow 749 \\
 & \frac{5 \int \frac{1}{(bx^4+a)^{3/2}} dx}{6a} + \frac{x}{6a(a + bx^4)^{3/2}} \\
 & \quad \downarrow 749 \\
 & \frac{5 \left(\frac{\int \frac{1}{\sqrt{bx^4+a}} dx}{2a} + \frac{x}{2a\sqrt{a+bx^4}} \right)}{6a} + \frac{x}{6a(a + bx^4)^{3/2}} \\
 & \quad \downarrow 761 \\
 & \frac{5 \left(\frac{(\sqrt{a} + \sqrt{bx^2}) \sqrt{\frac{a+bx^4}{(\sqrt{a} + \sqrt{bx^2})^2}} \operatorname{EllipticF} \left(2 \arctan \left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a}} \right), \frac{1}{2} \right)}{4a^{5/4} \sqrt[4]{b} \sqrt{a+bx^4}} + \frac{x}{2a\sqrt{a+bx^4}} \right)}{6a} + \frac{x}{6a(a + bx^4)^{3/2}}
 \end{aligned}$$

input `Int[(a + b*x^4)^(-5/2),x]`

output
$$\frac{x/(6*a*(a + b*x^4)^{(3/2)}) + (5*(x/(2*a*Sqrt[a + b*x^4]) + ((Sqrt[a] + Sqrt[b]*x^2)*Sqrt[(a + b*x^4)/(Sqrt[a] + Sqrt[b]*x^2)^2]*EllipticF[2*ArcTan[(b^{(1/4)}*x)/a^{(1/4)}], 1/2)]/(4*a^{(5/4)}*b^{(1/4)}*Sqrt[a + b*x^4])))/(6*a)}$$

Defintions of rubi rules used

rule 749
$$\text{Int}[(a + b*x^n)^p, x_Symbol] \rightarrow \text{Simp}[(-x)*(a + b*x^n)^{p+1}/(a*n*(p+1)), x] + \text{Simp}[(n*(p+1) + 1)/(a*n*(p+1)) \text{Int}[(a + b*x^n)^{p+1}, x], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{IGtQ}\{n, 0\} \ \&\& \ \text{LtQ}\{p, -1\} \ \&\& \ (\text{IntegerQ}\{2*p\} \ || \ \text{Denominator}\{p + 1/n\} < \text{Denominator}\{p\})$$

rule 761
$$\text{Int}[1/\text{Sqrt}[(a + b*x^4), x_Symbol] \rightarrow \text{With}\{q = \text{Rt}\{b/a, 4\}\}, \text{Simp}[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))* \text{EllipticF}[2*ArcTan[q*x], 1/2], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}\{b/a\}$$

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.47 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.97

method	result	size
default	$\frac{x\sqrt{bx^4+a}}{6ab^2(x^4+\frac{a}{b})^2} + \frac{5x}{12a^2\sqrt{(x^4+\frac{a}{b})b}} + \frac{5\sqrt{1-\frac{i\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{b}x^2}{\sqrt{a}}}\text{EllipticF}\left(x\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}, i\right)}{12a^2\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}\sqrt{bx^4+a}}$	123
elliptic	$\frac{x\sqrt{bx^4+a}}{6ab^2(x^4+\frac{a}{b})^2} + \frac{5x}{12a^2\sqrt{(x^4+\frac{a}{b})b}} + \frac{5\sqrt{1-\frac{i\sqrt{b}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{b}x^2}{\sqrt{a}}}\text{EllipticF}\left(x\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}, i\right)}{12a^2\sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}\sqrt{bx^4+a}}$	123

input `int(1/(b*x^4+a)^(5/2), x, method=_RETURNVERBOSE)`

output
$$\frac{1}{6}x/a/b^2*(b*x^4+a)^{(1/2)}/(x^4+a/b)^2+5/12*x/a^2/((x^4+a/b)*b)^{(1/2)}+5/12/a^2/(I/a^{(1/2)}*b^{(1/2)})^{(1/2)}*(1-I*b^{(1/2)}*x^2/a^{(1/2)})^{(1/2)}*(1+I*b^{(1/2)}*x^2/a^{(1/2)})^{(1/2)}/(b*x^4+a)^{(1/2)}*\text{EllipticF}(x*(I/a^{(1/2)}*b^{(1/2)})^{(1/2)}, I)$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.80

$$\int \frac{1}{(a + bx^4)^{5/2}} dx = \frac{5(b^2x^8 + 2abx^4 + a^2)\sqrt{a}\left(-\frac{b}{a}\right)^{\frac{3}{4}} F(\arcsin\left(x\left(-\frac{b}{a}\right)^{\frac{1}{4}}\right) \mid -1) - (5b^2x^5 + 7abx)\sqrt{bx^4 + a}}{12(a^2b^3x^8 + 2a^3b^2x^4 + a^4b)}$$

input `integrate(1/(b*x^4+a)^(5/2),x, algorithm="fricas")`

output `-1/12*(5*(b^2*x^8 + 2*a*b*x^4 + a^2)*sqrt(a)*(-b/a)^(3/4)*elliptic_f(arcsin(x*(-b/a)^(1/4)), -1) - (5*b^2*x^5 + 7*a*b*x)*sqrt(b*x^4 + a))/(a^2*b^3*x^8 + 2*a^3*b^2*x^4 + a^4*b)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.70 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.28

$$\int \frac{1}{(a + bx^4)^{5/2}} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{5}{2} \mid \frac{5}{4}, \frac{bx^4 e^{i\pi}}{a}\right)}{4a^{\frac{5}{2}}\Gamma\left(\frac{5}{4}\right)}$$

input `integrate(1/(b*x**4+a)**(5/2),x)`

output `x*gamma(1/4)*hyper((1/4, 5/2), (5/4,), b*x**4*exp_polar(I*pi)/a)/(4*a**(5/2)*gamma(5/4))`

Maxima [F]

$$\int \frac{1}{(a + bx^4)^{5/2}} dx = \int \frac{1}{(bx^4 + a)^{5/2}} dx$$

input `integrate(1/(b*x^4+a)^(5/2),x, algorithm="maxima")`

output `integrate((b*x^4 + a)^(-5/2), x)`

Giac [F]

$$\int \frac{1}{(a + bx^4)^{5/2}} dx = \int \frac{1}{(bx^4 + a)^{5/2}} dx$$

input `integrate(1/(b*x^4+a)^(5/2),x, algorithm="giac")`

output `integrate((b*x^4 + a)^(-5/2), x)`

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.29

$$\int \frac{1}{(a + bx^4)^{5/2}} dx = \frac{x \left(\frac{bx^4}{a} + 1 \right)^{5/2} {}_2F_1 \left(\frac{1}{4}, \frac{5}{2}; \frac{5}{4}; -\frac{bx^4}{a} \right)}{(bx^4 + a)^{5/2}}$$

input `int(1/(a + b*x^4)^(5/2),x)`

output `(x*((b*x^4)/a + 1)^(5/2)*hypergeom([1/4, 5/2], 5/4, -(b*x^4)/a))/(a + b*x^4)^(5/2)`

Reduce [F]

$$\int \frac{1}{(a + bx^4)^{5/2}} dx = \int \frac{\sqrt{bx^4 + a}}{b^3x^{12} + 3ab^2x^8 + 3a^2bx^4 + a^3} dx$$

input `int(1/(b*x^4+a)^(5/2),x)`

output `int(sqrt(a + b*x**4)/(a**3 + 3*a**2*b*x**4 + 3*a*b**2*x**8 + b**3*x**12),x)`

3.19 $\int (1 - x^4)^{5/2} dx$

Optimal result	164
Mathematica [C] (verified)	164
Rubi [A] (verified)	165
Maple [C] (verified)	166
Fricas [A] (verification not implemented)	167
Sympy [A] (verification not implemented)	167
Maxima [F]	167
Giac [F]	168
Mupad [B] (verification not implemented)	168
Reduce [F]	168

Optimal result

Integrand size = 11, antiderivative size = 57

$$\int (1 - x^4)^{5/2} dx = \frac{20}{77}x\sqrt{1 - x^4} + \frac{10}{77}x(1 - x^4)^{3/2} + \frac{1}{11}x(1 - x^4)^{5/2} + \frac{40}{77}\text{EllipticF}(\arcsin(x), -1)$$

output

```
20/77*x*(-x^4+1)^(1/2)+10/77*x*(-x^4+1)^(3/2)+1/11*x*(-x^4+1)^(5/2)+40/77*
EllipticF(x,I)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 4.71 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.26

$$\int (1 - x^4)^{5/2} dx = x \text{Hypergeometric2F1} \left(-\frac{5}{2}, \frac{1}{4}, \frac{5}{4}, x^4 \right)$$

input

```
Integrate[(1 - x^4)^(5/2),x]
```

output

```
x*Hypergeometric2F1[-5/2, 1/4, 5/4, x^4]
```

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.18, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {748, 748, 748, 762}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (1 - x^4)^{5/2} dx \\
 & \quad \downarrow 748 \\
 & \frac{10}{11} \int (1 - x^4)^{3/2} dx + \frac{1}{11} x(1 - x^4)^{5/2} \\
 & \quad \downarrow 748 \\
 & \frac{10}{11} \left(\frac{6}{7} \int \sqrt{1 - x^4} dx + \frac{1}{7} x(1 - x^4)^{3/2} \right) + \frac{1}{11} x(1 - x^4)^{5/2} \\
 & \quad \downarrow 748 \\
 & \frac{10}{11} \left(\frac{6}{7} \left(\frac{2}{3} \int \frac{1}{\sqrt{1 - x^4}} dx + \frac{1}{3} \sqrt{1 - x^4} x \right) + \frac{1}{7} x(1 - x^4)^{3/2} \right) + \frac{1}{11} x(1 - x^4)^{5/2} \\
 & \quad \downarrow 762 \\
 & \frac{10}{11} \left(\frac{6}{7} \left(\frac{2}{3} \text{EllipticF}(\arcsin(x), -1) + \frac{1}{3} \sqrt{1 - x^4} x \right) + \frac{1}{7} x(1 - x^4)^{3/2} \right) + \frac{1}{11} x(1 - x^4)^{5/2}
 \end{aligned}$$

input `Int[(1 - x^4)^(5/2), x]`

output `(x*(1 - x^4)^(5/2))/11 + (10*((x*(1 - x^4)^(3/2))/7 + (6*((x*Sqrt[1 - x^4])/3 + (2*EllipticF[ArcSin[x], -1])/3))/7))/11`

Definitions of rubi rules used

rule 748 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p + 1)), x] + Simp[a*n*(p/(n*p + 1)) Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || LtQ[Denominator[p + 1/n], Denominator[p]])`

rule 762 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := Simp[(1/(Sqrt[a]*Rt[-b/a, 4]))*EllipticF[ArcSin[Rt[-b/a, 4]*x], -1], x] /; FreeQ[{a, b}, x] && NegQ[b/a] && GtQ[a, 0]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4.

Time = 0.51 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.21

method	result	size
meijerg	$x \operatorname{hypergeom}\left(\left[-\frac{5}{2}, \frac{1}{4}\right], \left[\frac{5}{4}\right], x^4\right)$	12
risch	$-\frac{x(7x^8-24x^4+37)(x^4-1)}{77\sqrt{-x^4+1}} + \frac{40\sqrt{-x^2+1}\sqrt{x^2+1}\operatorname{EllipticF}(x,i)}{77\sqrt{-x^4+1}}$	62
default	$\frac{x^9\sqrt{-x^4+1}}{11} - \frac{24x^5\sqrt{-x^4+1}}{77} + \frac{37x\sqrt{-x^4+1}}{77} + \frac{40\sqrt{-x^2+1}\sqrt{x^2+1}\operatorname{EllipticF}(x,i)}{77\sqrt{-x^4+1}}$	73
elliptic	$\frac{x^9\sqrt{-x^4+1}}{11} - \frac{24x^5\sqrt{-x^4+1}}{77} + \frac{37x\sqrt{-x^4+1}}{77} + \frac{40\sqrt{-x^2+1}\sqrt{x^2+1}\operatorname{EllipticF}(x,i)}{77\sqrt{-x^4+1}}$	73

input `int((-x^4+1)^(5/2),x,method=_RETURNVERBOSE)`

output `x*hypergeom([-5/2,1/4],[5/4],x^4)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.60

$$\int (1 - x^4)^{5/2} dx = \frac{1}{77} (7x^9 - 24x^5 + 37x) \sqrt{-x^4 + 1} + \frac{40}{77} i F(\arcsin\left(\frac{1}{x}\right) | -1)$$

input `integrate((-x^4+1)^(5/2),x, algorithm="fricas")`output `1/77*(7*x^9 - 24*x^5 + 37*x)*sqrt(-x^4 + 1) + 40/77*I*elliptic_f(arcsin(1/x), -1)`**Sympy [A] (verification not implemented)**

Time = 0.51 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.54

$$\int (1 - x^4)^{5/2} dx = \frac{x \Gamma\left(\frac{1}{4}\right) {}_2F_1\left(-\frac{5}{2}, \frac{1}{4} \middle| \frac{5}{4}, x^4 e^{2i\pi}\right)}{4 \Gamma\left(\frac{5}{4}\right)}$$

input `integrate((-x**4+1)**(5/2),x)`output `x*gamma(1/4)*hyper((-5/2, 1/4), (5/4,), x**4*exp_polar(2*I*pi))/(4*gamma(5/4))`**Maxima [F]**

$$\int (1 - x^4)^{5/2} dx = \int (-x^4 + 1)^{\frac{5}{2}} dx$$

input `integrate((-x^4+1)^(5/2),x, algorithm="maxima")`output `integrate((-x^4 + 1)^(5/2), x)`

Giac [F]

$$\int (1 - x^4)^{5/2} dx = \int (-x^4 + 1)^{\frac{5}{2}} dx$$

input `integrate((-x^4+1)^(5/2),x, algorithm="giac")`

output `integrate((-x^4 + 1)^(5/2), x)`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.18

$$\int (1 - x^4)^{5/2} dx = x {}_2F_1\left(-\frac{5}{2}, \frac{1}{4}; \frac{5}{4}; x^4\right)$$

input `int((1 - x^4)^(5/2),x)`

output `x*hypergeom([-5/2, 1/4], 5/4, x^4)`

Reduce [F]

$$\int (1 - x^4)^{5/2} dx = \frac{\sqrt{-x^4 + 1} x^9}{11} - \frac{24\sqrt{-x^4 + 1} x^5}{77} + \frac{37\sqrt{-x^4 + 1} x}{77} - \frac{40\left(\int \frac{\sqrt{-x^4 + 1}}{x^4 - 1} dx\right)}{77}$$

input `int((-x^4+1)^(5/2),x)`

output `(7*sqrt(-x**4 + 1)*x**9 - 24*sqrt(-x**4 + 1)*x**5 + 37*sqrt(-x**4 + 1)*x - 40*int(sqrt(-x**4 + 1)/(x**4 - 1),x))/77`

3.20 $\int (1 - x^4)^{3/2} dx$

Optimal result	169
Mathematica [C] (verified)	169
Rubi [A] (verified)	170
Maple [C] (verified)	171
Fricas [A] (verification not implemented)	171
Sympy [A] (verification not implemented)	172
Maxima [F]	172
Giac [F]	172
Mupad [B] (verification not implemented)	173
Reduce [F]	173

Optimal result

Integrand size = 11, antiderivative size = 41

$$\int (1 - x^4)^{3/2} dx = \frac{2}{7}x\sqrt{1 - x^4} + \frac{1}{7}x(1 - x^4)^{3/2} + \frac{4}{7}\text{EllipticF}(\arcsin(x), -1)$$

output

```
2/7*x*(-x^4+1)^(1/2)+1/7*x*(-x^4+1)^(3/2)+4/7*EllipticF(x,I)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 4.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.37

$$\int (1 - x^4)^{3/2} dx = x \text{Hypergeometric2F1} \left(-\frac{3}{2}, \frac{1}{4}, \frac{5}{4}, x^4 \right)$$

input

```
Integrate[(1 - x^4)^(3/2),x]
```

output

```
x*Hypergeometric2F1[-3/2, 1/4, 5/4, x^4]
```

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.12, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {748, 748, 762}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (1 - x^4)^{3/2} dx \\ & \quad \downarrow 748 \\ & \frac{6}{7} \int \sqrt{1 - x^4} dx + \frac{1}{7} x (1 - x^4)^{3/2} \\ & \quad \downarrow 748 \\ & \frac{6}{7} \left(\frac{2}{3} \int \frac{1}{\sqrt{1 - x^4}} dx + \frac{1}{3} \sqrt{1 - x^4} x \right) + \frac{1}{7} x (1 - x^4)^{3/2} \\ & \quad \downarrow 762 \\ & \frac{6}{7} \left(\frac{2}{3} \text{EllipticF}(\arcsin(x), -1) + \frac{1}{3} \sqrt{1 - x^4} x \right) + \frac{1}{7} x (1 - x^4)^{3/2} \end{aligned}$$

input `Int[(1 - x^4)^(3/2), x]`

output `(x*(1 - x^4)^(3/2))/7 + (6*((x*Sqrt[1 - x^4])/3 + (2*EllipticF[ArcSin[x], -1])/3))/7`

Defintions of rubi rules used

rule 748

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p + 1)), x] + Simp[a*n*(p/(n*p + 1)) Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || LtQ[Denominator[p + 1/n], Denominator[p]])
```

rule 762

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := Simp[(1/(Sqrt[a]*Rt[-b/a, 4])
)*EllipticF[ArcSin[Rt[-b/a, 4]*x], -1], x] /; FreeQ[{a, b}, x] && NegQ[b/a]
&& GtQ[a, 0]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4.

Time = 0.50 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.29

method	result	size
meijerg	$x \operatorname{hypergeom}\left(\left[-\frac{3}{2}, \frac{1}{4}\right], \left[\frac{5}{4}\right], x^4\right)$	12
risch	$\frac{x(x^4-3)(x^4-1)}{7\sqrt{-x^4+1}} + \frac{4\sqrt{-x^2+1}\sqrt{x^2+1}\operatorname{EllipticF}(x,i)}{7\sqrt{-x^4+1}}$	55
default	$-\frac{x^5\sqrt{-x^4+1}}{7} + \frac{3x\sqrt{-x^4+1}}{7} + \frac{4\sqrt{-x^2+1}\sqrt{x^2+1}\operatorname{EllipticF}(x,i)}{7\sqrt{-x^4+1}}$	59
elliptic	$-\frac{x^5\sqrt{-x^4+1}}{7} + \frac{3x\sqrt{-x^4+1}}{7} + \frac{4\sqrt{-x^2+1}\sqrt{x^2+1}\operatorname{EllipticF}(x,i)}{7\sqrt{-x^4+1}}$	59

input `int((-x^4+1)^(3/2),x,method=_RETURNVERBOSE)`

output `x*hypergeom([-3/2,1/4],[5/4],x^4)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.66

$$\int (1-x^4)^{3/2} dx = -\frac{1}{7}(x^5-3x)\sqrt{-x^4+1} + \frac{4}{7}i F\left(\arcsin\left(\frac{1}{x}\right) \mid -1\right)$$

input `integrate((-x^4+1)^(3/2),x, algorithm="fricas")`

output `-1/7*(x^5 - 3*x)*sqrt(-x^4 + 1) + 4/7*I*elliptic_f(arcsin(1/x), -1)`

Sympy [A] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.76

$$\int (1 - x^4)^{3/2} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\begin{matrix} -\frac{3}{2}, \frac{1}{4} \\ \frac{5}{4} \end{matrix} \middle| x^4 e^{2i\pi}\right)}{4\Gamma\left(\frac{5}{4}\right)}$$

input `integrate((-x**4+1)**(3/2),x)`output `x*gamma(1/4)*hyper((-3/2, 1/4), (5/4,), x**4*exp_polar(2*I*pi))/(4*gamma(5/4))`**Maxima [F]**

$$\int (1 - x^4)^{3/2} dx = \int (-x^4 + 1)^{\frac{3}{2}} dx$$

input `integrate((-x^4+1)^(3/2),x, algorithm="maxima")`output `integrate((-x^4 + 1)^(3/2), x)`**Giac [F]**

$$\int (1 - x^4)^{3/2} dx = \int (-x^4 + 1)^{\frac{3}{2}} dx$$

input `integrate((-x^4+1)^(3/2),x, algorithm="giac")`output `integrate((-x^4 + 1)^(3/2), x)`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.24

$$\int (1 - x^4)^{3/2} dx = x {}_2F_1\left(-\frac{3}{2}, \frac{1}{4}; \frac{5}{4}; x^4\right)$$

input `int((1 - x^4)^(3/2),x)`output `x*hypergeom([-3/2, 1/4], 5/4, x^4)`**Reduce [F]**

$$\int (1 - x^4)^{3/2} dx = -\frac{\sqrt{-x^4 + 1} x^5}{7} + \frac{3\sqrt{-x^4 + 1} x}{7} - \frac{4\left(\int \frac{\sqrt{-x^4 + 1}}{x^4 - 1} dx\right)}{7}$$

input `int((-x^4+1)^(3/2),x)`output `(- sqrt(- x**4 + 1)*x**5 + 3*sqrt(- x**4 + 1)*x - 4*int(sqrt(- x**4 + 1)/(x**4 - 1),x))/7`

3.21 $\int \sqrt{1-x^4} dx$

Optimal result	174
Mathematica [A] (verified)	174
Rubi [A] (verified)	175
Maple [C] (verified)	176
Fricas [A] (verification not implemented)	176
Sympy [A] (verification not implemented)	177
Maxima [F]	177
Giac [F]	177
Mupad [B] (verification not implemented)	178
Reduce [F]	178

Optimal result

Integrand size = 11, antiderivative size = 25

$$\int \sqrt{1-x^4} dx = \frac{1}{3}x\sqrt{1-x^4} + \frac{2}{3}\text{EllipticF}(\arcsin(x), -1)$$

output `1/3*x*(-x^4+1)^(1/2)+2/3*EllipticF(x,I)`

Mathematica [A] (verified)

Time = 3.17 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.56

$$\int \sqrt{1-x^4} dx = \frac{x-x^5+2\sqrt{1-x^4}\text{EllipticF}(\arcsin(x), -1)}{3\sqrt{1-x^4}}$$

input `Integrate[Sqrt[1-x^4],x]`

output `(x-x^5+2*Sqrt[1-x^4]*EllipticF[ArcSin[x],-1])/(3*Sqrt[1-x^4])`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {748, 762}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{1-x^4} dx$$

$$\downarrow 748$$

$$\frac{2}{3} \int \frac{1}{\sqrt{1-x^4}} dx + \frac{1}{3} \sqrt{1-x^4} x$$

$$\downarrow 762$$

$$\frac{2}{3} \text{EllipticF}(\arcsin(x), -1) + \frac{1}{3} \sqrt{1-x^4} x$$

input `Int[Sqrt[1 - x^4], x]`

output `(x*Sqrt[1 - x^4])/3 + (2*EllipticF[ArcSin[x], -1])/3`

Defintions of rubi rules used

rule 748 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p + 1)), x] + Simp[a*n*(p/(n*p + 1)) Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || LtQ[Denominator[p + 1/n], Denominator[p]])`

rule 762 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := Simp[(1/(Sqrt[a]*Rt[-b/a, 4]))*EllipticF[ArcSin[Rt[-b/a, 4]*x], -1], x] /; FreeQ[{a, b}, x] && NegQ[b/a] && GtQ[a, 0]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4.

Time = 0.48 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.48

method	result	size
meijerg	$x \operatorname{hypergeom}\left(\left[-\frac{1}{2}, \frac{1}{4}\right], \left[\frac{5}{4}\right], x^4\right)$	12
default	$\frac{x\sqrt{-x^4+1}}{3} + \frac{2\sqrt{-x^2+1}\sqrt{x^2+1}\operatorname{EllipticF}(x,i)}{3\sqrt{-x^4+1}}$	45
elliptic	$\frac{x\sqrt{-x^4+1}}{3} + \frac{2\sqrt{-x^2+1}\sqrt{x^2+1}\operatorname{EllipticF}(x,i)}{3\sqrt{-x^4+1}}$	45
risch	$-\frac{x(x^4-1)}{3\sqrt{-x^4+1}} + \frac{2\sqrt{-x^2+1}\sqrt{x^2+1}\operatorname{EllipticF}(x,i)}{3\sqrt{-x^4+1}}$	50

input `int((-x^4+1)^(1/2),x,method=_RETURNVERBOSE)`

output `x*hypergeom([-1/2,1/4],[5/4],x^4)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int \sqrt{1-x^4} dx = \frac{1}{3} \sqrt{-x^4+1} x + \frac{2}{3} i F\left(\arcsin\left(\frac{1}{x}\right) \mid -1\right)$$

input `integrate((-x^4+1)^(1/2),x, algorithm="fricas")`

output `1/3*sqrt(-x^4 + 1)*x + 2/3*I*elliptic_f(arcsin(1/x), -1)`

Sympy [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.24

$$\int \sqrt{1-x^4} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\begin{matrix} -\frac{1}{2}, \frac{1}{4} \\ \frac{5}{4} \end{matrix} \middle| x^4 e^{2i\pi}\right)}{4\Gamma\left(\frac{5}{4}\right)}$$

input `integrate((-x**4+1)**(1/2),x)`output `x*gamma(1/4)*hyper((-1/2, 1/4), (5/4,), x**4*exp_polar(2*I*pi))/(4*gamma(5/4))`**Maxima [F]**

$$\int \sqrt{1-x^4} dx = \int \sqrt{-x^4+1} dx$$

input `integrate((-x^4+1)^(1/2),x, algorithm="maxima")`output `integrate(sqrt(-x^4 + 1), x)`**Giac [F]**

$$\int \sqrt{1-x^4} dx = \int \sqrt{-x^4+1} dx$$

input `integrate((-x^4+1)^(1/2),x, algorithm="giac")`output `integrate(sqrt(-x^4 + 1), x)`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.40

$$\int \sqrt{1-x^4} dx = x {}_2F_1\left(-\frac{1}{2}, \frac{1}{4}; \frac{5}{4}; x^4\right)$$

input `int((1 - x^4)^(1/2),x)`output `x*hypergeom([-1/2, 1/4], 5/4, x^4)`**Reduce [F]**

$$\int \sqrt{1-x^4} dx = \frac{\sqrt{-x^4+1}x}{3} - \frac{2\left(\int \frac{\sqrt{-x^4+1}}{x^4-1} dx\right)}{3}$$

input `int((-x^4+1)^(1/2),x)`output `(sqrt(-x**4+1)*x - 2*int(sqrt(-x**4+1)/(x**4-1),x))/3`

3.22 $\int \frac{1}{\sqrt{1-x^4}} dx$

Optimal result	179
Mathematica [A] (verified)	179
Rubi [A] (verified)	180
Maple [C] (verified)	180
Fricas [A] (verification not implemented)	181
Sympy [B] (verification not implemented)	181
Maxima [F]	182
Giac [F]	182
Mupad [B] (verification not implemented)	182
Reduce [F]	183

Optimal result

Integrand size = 11, antiderivative size = 4

$$\int \frac{1}{\sqrt{1-x^4}} dx = \text{EllipticF}(\arcsin(x), -1)$$

output `EllipticF(x,I)`

Mathematica [A] (verified)

Time = 10.03 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1-x^4}} dx = \text{EllipticF}(\arcsin(x), -1)$$

input `Integrate[1/Sqrt[1 - x^4],x]`

output `EllipticF[ArcSin[x], -1]`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {762}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{1-x^4}} dx$$

↓ 762

$$\text{EllipticF}(\arcsin(x), -1)$$

input `Int[1/Sqrt[1 - x^4], x]`

output `EllipticF[ArcSin[x], -1]`

Defintions of rubi rules used

rule 762 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] :> Simp[(1/(Sqrt[a]*Rt[-b/a, 4])
) * EllipticF[ArcSin[Rt[-b/a, 4]*x], -1], x] /; FreeQ[{a, b}, x] && NegQ[b/a]
&& GtQ[a, 0]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4.

Time = 0.45 (sec) , antiderivative size = 12, normalized size of antiderivative = 3.00

method	result	size
meijerg	$x \text{ hypergeom} \left(\left[\frac{1}{4}, \frac{1}{2} \right], \left[\frac{5}{4} \right], x^4 \right)$	12
default	$\frac{\sqrt{-x^2+1} \sqrt{x^2+1} \text{EllipticF}(x, i)}{\sqrt{-x^4+1}}$	31
elliptic	$\frac{\sqrt{-x^2+1} \sqrt{x^2+1} \text{EllipticF}(x, i)}{\sqrt{-x^4+1}}$	31

input `int(1/(-x^4+1)^(1/2),x,method=_RETURNVERBOSE)`

output `x*hypergeom([1/4,1/2],[5/4],x^4)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1-x^4}} dx = F(\arcsin(x) | -1)$$

input `integrate(1/(-x^4+1)^(1/2),x, algorithm="fricas")`

output `elliptic_f(arcsin(x), -1)`

Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 29 vs. $2(2) = 4$.

Time = 0.39 (sec) , antiderivative size = 29, normalized size of antiderivative = 7.25

$$\int \frac{1}{\sqrt{1-x^4}} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2} \middle| \frac{5}{4} \middle| x^4 e^{2i\pi}\right)}{4\Gamma\left(\frac{5}{4}\right)}$$

input `integrate(1/(-x**4+1)**(1/2),x)`

output `x*gamma(1/4)*hyper((1/4, 1/2), (5/4,), x**4*exp_polar(2*I*pi))/(4*gamma(5/4))`

Maxima [F]

$$\int \frac{1}{\sqrt{1-x^4}} dx = \int \frac{1}{\sqrt{-x^4+1}} dx$$

input `integrate(1/(-x^4+1)^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(-x^4 + 1), x)`

Giac [F]

$$\int \frac{1}{\sqrt{1-x^4}} dx = \int \frac{1}{\sqrt{-x^4+1}} dx$$

input `integrate(1/(-x^4+1)^(1/2),x, algorithm="giac")`

output `integrate(1/sqrt(-x^4 + 1), x)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 2.50

$$\int \frac{1}{\sqrt{1-x^4}} dx = x {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; x^4\right)$$

input `int(1/(1 - x^4)^(1/2),x)`

output `x*hypergeom([1/4, 1/2], 5/4, x^4)`

Reduce [F]

$$\int \frac{1}{\sqrt{1-x^4}} dx = - \left(\int \frac{\sqrt{-x^4+1}}{x^4-1} dx \right)$$

input `int(1/(-x^4+1)^(1/2),x)`

output `- int(sqrt(-x**4+1)/(x**4-1),x)`

3.23 $\int \frac{1}{(1-x^4)^{3/2}} dx$

Optimal result	184
Mathematica [C] (verified)	184
Rubi [A] (verified)	185
Maple [C] (verified)	186
Fricas [A] (verification not implemented)	186
Sympy [A] (verification not implemented)	187
Maxima [F]	187
Giac [F]	187
Mupad [B] (verification not implemented)	188
Reduce [F]	188

Optimal result

Integrand size = 11, antiderivative size = 25

$$\int \frac{1}{(1-x^4)^{3/2}} dx = \frac{x}{2\sqrt{1-x^4}} + \frac{1}{2} \text{EllipticF}(\arcsin(x), -1)$$

output `1/2*x/(-x^4+1)^(1/2)+1/2*EllipticF(x,I)`

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 3.49 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.20

$$\int \frac{1}{(1-x^4)^{3/2}} dx = \frac{1}{2}x \left(\frac{1}{\sqrt{1-x^4}} + \text{Hypergeometric2F1} \left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, x^4 \right) \right)$$

input `Integrate[(1 - x^4)^(-3/2),x]`

output `(x*(1/Sqrt[1 - x^4] + Hypergeometric2F1[1/4, 1/2, 5/4, x^4]))/2`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {749, 762}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(1-x^4)^{3/2}} dx$$

$$\downarrow \text{749}$$

$$\frac{1}{2} \int \frac{1}{\sqrt{1-x^4}} dx + \frac{x}{2\sqrt{1-x^4}}$$

$$\downarrow \text{762}$$

$$\frac{1}{2} \text{EllipticF}(\arcsin(x), -1) + \frac{x}{2\sqrt{1-x^4}}$$

input `Int[(1 - x^4)^(-3/2), x]`

output `x/(2*sqrt[1 - x^4]) + EllipticF[ArcSin[x], -1]/2`

Defintions of rubi rules used

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

rule 762 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := Simp[(1/(Sqrt[a]*Rt[-b/a, 4]))*EllipticF[ArcSin[Rt[-b/a, 4]*x], -1], x] /; FreeQ[{a, b}, x] && NegQ[b/a] && GtQ[a, 0]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4.

Time = 0.50 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.48

method	result	size
meijerg	$x \operatorname{hypergeom}\left(\left[\frac{1}{4}, \frac{3}{2}\right], \left[\frac{5}{4}\right], x^4\right)$	12
default	$\frac{x}{2\sqrt{-x^4+1}} + \frac{\sqrt{-x^2+1}\sqrt{x^2+1}\operatorname{EllipticF}(x,i)}{2\sqrt{-x^4+1}}$	45
risch	$\frac{x}{2\sqrt{-x^4+1}} + \frac{\sqrt{-x^2+1}\sqrt{x^2+1}\operatorname{EllipticF}(x,i)}{2\sqrt{-x^4+1}}$	45
elliptic	$\frac{x}{2\sqrt{-x^4+1}} + \frac{\sqrt{-x^2+1}\sqrt{x^2+1}\operatorname{EllipticF}(x,i)}{2\sqrt{-x^4+1}}$	45

input `int(1/(-x^4+1)^(3/2),x,method=_RETURNVERBOSE)`

output `x*hypergeom([1/4,3/2],[5/4],x^4)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.28

$$\int \frac{1}{(1-x^4)^{3/2}} dx = \frac{(x^4-1)F(\arcsin(x) | -1) - \sqrt{-x^4+1}x}{2(x^4-1)}$$

input `integrate(1/(-x^4+1)^(3/2),x, algorithm="fricas")`

output `1/2*((x^4 - 1)*elliptic_f(arcsin(x), -1) - sqrt(-x^4 + 1)*x)/(x^4 - 1)`

Sympy [A] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.16

$$\int \frac{1}{(1-x^4)^{3/2}} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{3}{2} \middle| \frac{5}{4}, x^4 e^{2i\pi}\right)}{4\Gamma\left(\frac{5}{4}\right)}$$

input `integrate(1/(-x**4+1)**(3/2),x)`output `x*gamma(1/4)*hyper((1/4, 3/2), (5/4,), x**4*exp_polar(2*I*pi))/(4*gamma(5/4))`**Maxima [F]**

$$\int \frac{1}{(1-x^4)^{3/2}} dx = \int \frac{1}{(-x^4+1)^{\frac{3}{2}}} dx$$

input `integrate(1/(-x^4+1)^(3/2),x, algorithm="maxima")`output `integrate((-x^4 + 1)^(-3/2), x)`**Giac [F]**

$$\int \frac{1}{(1-x^4)^{3/2}} dx = \int \frac{1}{(-x^4+1)^{\frac{3}{2}}} dx$$

input `integrate(1/(-x^4+1)^(3/2),x, algorithm="giac")`output `integrate((-x^4 + 1)^(-3/2), x)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.40

$$\int \frac{1}{(1-x^4)^{3/2}} dx = x {}_2F_1\left(\frac{1}{4}, \frac{3}{2}; \frac{5}{4}; x^4\right)$$

input `int(1/(1 - x^4)^(3/2),x)`output `x*hypergeom([1/4, 3/2], 5/4, x^4)`**Reduce [F]**

$$\int \frac{1}{(1-x^4)^{3/2}} dx = \int \frac{\sqrt{-x^4+1}}{x^8-2x^4+1} dx$$

input `int(1/(-x^4+1)^(3/2),x)`output `int(sqrt(-x**4+1)/(x**8-2*x**4+1),x)`

3.24 $\int \frac{1}{(1-x^4)^{5/2}} dx$

Optimal result	189
Mathematica [C] (verified)	189
Rubi [A] (verified)	190
Maple [C] (verified)	191
Fricas [A] (verification not implemented)	191
Sympy [A] (verification not implemented)	192
Maxima [F]	192
Giac [F]	192
Mupad [B] (verification not implemented)	193
Reduce [F]	193

Optimal result

Integrand size = 11, antiderivative size = 41

$$\int \frac{1}{(1-x^4)^{5/2}} dx = \frac{x}{6(1-x^4)^{3/2}} + \frac{5x}{12\sqrt{1-x^4}} + \frac{5}{12} \text{EllipticF}(\arcsin(x), -1)$$

output `1/6*x/(-x^4+1)^(3/2)+5/12*x/(-x^4+1)^(1/2)+5/12*EllipticF(x,I)`

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 4.46 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.24

$$\int \frac{1}{(1-x^4)^{5/2}} dx = \frac{x}{6(1-x^4)^{3/2}} + \frac{5x}{12\sqrt{1-x^4}} + \frac{5}{12} x \text{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, x^4\right)$$

input `Integrate[(1 - x^4)^(-5/2), x]`

output `x/(6*(1 - x^4)^(3/2)) + (5*x)/(12*Sqrt[1 - x^4]) + (5*x*Hypergeometric2F1[1/4, 1/2, 5/4, x^4])/12`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.12, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {749, 749, 762}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(1-x^4)^{5/2}} dx$$

↓ 749

$$\frac{5}{6} \int \frac{1}{(1-x^4)^{3/2}} dx + \frac{x}{6(1-x^4)^{3/2}}$$

↓ 749

$$\frac{5}{6} \left(\frac{1}{2} \int \frac{1}{\sqrt{1-x^4}} dx + \frac{x}{2\sqrt{1-x^4}} \right) + \frac{x}{6(1-x^4)^{3/2}}$$

↓ 762

$$\frac{5}{6} \left(\frac{1}{2} \text{EllipticF}(\arcsin(x), -1) + \frac{x}{2\sqrt{1-x^4}} \right) + \frac{x}{6(1-x^4)^{3/2}}$$

input `Int[(1 - x^4)^(-5/2), x]`

output `x/(6*(1 - x^4)^(3/2)) + (5*(x/(2*sqrt[1 - x^4]) + EllipticF[ArcSin[x], -1]/2))/6`

Defintions of rubi rules used

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

rule 762

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := Simp[(1/(Sqrt[a]*Rt[-b/a, 4])
)*EllipticF[ArcSin[Rt[-b/a, 4]*x], -1], x] /; FreeQ[{a, b}, x] && NegQ[b/a]
&& GtQ[a, 0]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4.

Time = 0.51 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.29

method	result	size
meijerg	$x \operatorname{hypergeom}\left(\left[\frac{1}{4}, \frac{5}{2}\right], \left[\frac{5}{4}\right], x^4\right)$	12
risch	$\frac{x(5x^4-7)}{12(x^4-1)\sqrt{-x^4+1}} + \frac{5\sqrt{-x^2+1}\sqrt{x^2+1}\operatorname{EllipticF}(x,i)}{12\sqrt{-x^4+1}}$	59
default	$\frac{x\sqrt{-x^4+1}}{6(x^4-1)^2} + \frac{5x}{12\sqrt{-x^4+1}} + \frac{5\sqrt{-x^2+1}\sqrt{x^2+1}\operatorname{EllipticF}(x,i)}{12\sqrt{-x^4+1}}$	64
elliptic	$\frac{x\sqrt{-x^4+1}}{6(x^4-1)^2} + \frac{5x}{12\sqrt{-x^4+1}} + \frac{5\sqrt{-x^2+1}\sqrt{x^2+1}\operatorname{EllipticF}(x,i)}{12\sqrt{-x^4+1}}$	64

input

```
int(1/(-x^4+1)^(5/2), x, method=_RETURNVERBOSE)
```

output

```
x*hypergeom([1/4, 5/2], [5/4], x^4)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.24

$$\int \frac{1}{(1-x^4)^{5/2}} dx = \frac{5(x^8 - 2x^4 + 1)F(\arcsin(x) | -1) - (5x^5 - 7x)\sqrt{-x^4 + 1}}{12(x^8 - 2x^4 + 1)}$$

input

```
integrate(1/(-x^4+1)^(5/2), x, algorithm="fricas")
```

output

```
1/12*(5*(x^8 - 2*x^4 + 1)*elliptic_f(arcsin(x), -1) - (5*x^5 - 7*x)*sqrt(-
x^4 + 1))/(x^8 - 2*x^4 + 1)
```


Sympy [A] (verification not implemented)

Time = 0.48 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.71

$$\int \frac{1}{(1-x^4)^{5/2}} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{5}{2} \middle| \frac{5}{4} \right) x^4 e^{2i\pi}}{4\Gamma\left(\frac{5}{4}\right)}$$

input `integrate(1/(-x**4+1)**(5/2),x)`output `x*gamma(1/4)*hyper((1/4, 5/2), (5/4,), x**4*exp_polar(2*I*pi))/(4*gamma(5/4))`**Maxima [F]**

$$\int \frac{1}{(1-x^4)^{5/2}} dx = \int \frac{1}{(-x^4+1)^{\frac{5}{2}}} dx$$

input `integrate(1/(-x^4+1)^(5/2),x, algorithm="maxima")`output `integrate((-x^4 + 1)^(-5/2), x)`**Giac [F]**

$$\int \frac{1}{(1-x^4)^{5/2}} dx = \int \frac{1}{(-x^4+1)^{\frac{5}{2}}} dx$$

input `integrate(1/(-x^4+1)^(5/2),x, algorithm="giac")`output `integrate((-x^4 + 1)^(-5/2), x)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.24

$$\int \frac{1}{(1-x^4)^{5/2}} dx = x {}_2F_1\left(\frac{1}{4}, \frac{5}{2}; \frac{5}{4}; x^4\right)$$

input `int(1/(1 - x^4)^(5/2),x)`output `x*hypergeom([1/4, 5/2], 5/4, x^4)`**Reduce [F]**

$$\int \frac{1}{(1-x^4)^{5/2}} dx = -\left(\int \frac{\sqrt{-x^4+1}}{x^{12}-3x^8+3x^4-1} dx\right)$$

input `int(1/(-x^4+1)^(5/2),x)`output `- int(sqrt(- x**4 + 1)/(x**12 - 3*x**8 + 3*x**4 - 1),x)`

3.25 $\int (1 + x^4)^{5/2} dx$

Optimal result	194
Mathematica [C] (verified)	194
Rubi [A] (verified)	195
Maple [A] (verified)	196
Fricas [C] (verification not implemented)	197
Sympy [C] (verification not implemented)	197
Maxima [F]	198
Giac [F]	198
Mupad [B] (verification not implemented)	198
Reduce [F]	199

Optimal result

Integrand size = 9, antiderivative size = 86

$$\int (1 + x^4)^{5/2} dx = \frac{20}{77}x\sqrt{1 + x^4} + \frac{10}{77}x(1 + x^4)^{3/2} + \frac{1}{11}x(1 + x^4)^{5/2} + \frac{20(1 + x^2) \sqrt{\frac{1+x^4}{(1+x^2)^2}} \operatorname{EllipticF}\left(2 \arctan(x), \frac{1}{2}\right)}{77\sqrt{1 + x^4}}$$

output

```
20/77*x*(x^4+1)^(1/2)+10/77*x*(x^4+1)^(3/2)+1/11*x*(x^4+1)^(5/2)+20/77*(x^2+1)*((x^4+1)/(x^2+1)^2)^(1/2)*InverseJacobiAM(2*arctan(x),1/2*2^(1/2))/(x^4+1)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 4.33 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.20

$$\int (1 + x^4)^{5/2} dx = x \operatorname{Hypergeometric2F1}\left(-\frac{5}{2}, \frac{1}{4}, \frac{5}{4}, -x^4\right)$$

input

```
Integrate[(1 + x^4)^(5/2), x]
```

output `x*Hypergeometric2F1[-5/2, 1/4, 5/4, -x^4]`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.12, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {748, 748, 748, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (x^4 + 1)^{5/2} dx \\
 & \quad \downarrow 748 \\
 & \frac{10}{11} \int (x^4 + 1)^{3/2} dx + \frac{1}{11} x (x^4 + 1)^{5/2} \\
 & \quad \downarrow 748 \\
 & \frac{10}{11} \left(\frac{6}{7} \int \sqrt{x^4 + 1} dx + \frac{1}{7} x (x^4 + 1)^{3/2} \right) + \frac{1}{11} x (x^4 + 1)^{5/2} \\
 & \quad \downarrow 748 \\
 & \frac{10}{11} \left(\frac{6}{7} \left(\frac{2}{3} \int \frac{1}{\sqrt{x^4 + 1}} dx + \frac{1}{3} \sqrt{x^4 + 1} x \right) + \frac{1}{7} x (x^4 + 1)^{3/2} \right) + \frac{1}{11} x (x^4 + 1)^{5/2} \\
 & \quad \downarrow 761 \\
 & \frac{10}{11} \left(\frac{6}{7} \left(\frac{(x^2 + 1) \sqrt{\frac{x^4 + 1}{(x^2 + 1)^2}} \operatorname{EllipticF}\left(2 \arctan(x), \frac{1}{2}\right)}{3\sqrt{x^4 + 1}} + \frac{1}{3} \sqrt{x^4 + 1} x \right) + \frac{1}{7} x (x^4 + 1)^{3/2} \right) + \\
 & \quad \frac{1}{11} x (x^4 + 1)^{5/2}
 \end{aligned}$$

input `Int[(1 + x^4)^(5/2), x]`

output `(x*(1 + x^4)^(5/2))/11 + (10*((x*(1 + x^4)^(3/2))/7 + (6*((x*Sqrt[1 + x^4])/3 + ((1 + x^2)*Sqrt[(1 + x^4)/(1 + x^2)^2]*EllipticF[2*ArcTan[x], 1/2])/(3*Sqrt[1 + x^4])))/7))/11`

Definitions of rubi rules used

rule 748

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p
+ 1)), x] + Simp[a*n*(p/(n*p + 1)) Int[(a + b*x^n)^(p - 1), x], x] /; Fre
eQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || LtQ[Denominat
or[p + 1/n], Denominator[p]])
```

rule 761

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(
1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*
EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]
```

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.16

method	result	size
meijerg	$x \operatorname{hypergeom}\left(\left[-\frac{5}{2}, \frac{1}{4}\right], \left[\frac{5}{4}\right], -x^4\right)$	14
risch	$\frac{x(7x^8+24x^4+37)\sqrt{x^4+1}}{77} + \frac{40\sqrt{-ix^2+1}\sqrt{ix^2+1}\operatorname{EllipticF}\left(x\left(\frac{\sqrt{2}+i\sqrt{2}}{2}\right), i\right)}{77\left(\frac{\sqrt{2}+i\sqrt{2}}{2}\right)\sqrt{x^4+1}}$	84
default	$\frac{x^9\sqrt{x^4+1}}{11} + \frac{24x^5\sqrt{x^4+1}}{77} + \frac{37x\sqrt{x^4+1}}{77} + \frac{40\sqrt{-ix^2+1}\sqrt{ix^2+1}\operatorname{EllipticF}\left(x\left(\frac{\sqrt{2}+i\sqrt{2}}{2}\right), i\right)}{77\left(\frac{\sqrt{2}+i\sqrt{2}}{2}\right)\sqrt{x^4+1}}$	96
elliptic	$\frac{x^9\sqrt{x^4+1}}{11} + \frac{24x^5\sqrt{x^4+1}}{77} + \frac{37x\sqrt{x^4+1}}{77} + \frac{40\sqrt{-ix^2+1}\sqrt{ix^2+1}\operatorname{EllipticF}\left(x\left(\frac{\sqrt{2}+i\sqrt{2}}{2}\right), i\right)}{77\left(\frac{\sqrt{2}+i\sqrt{2}}{2}\right)\sqrt{x^4+1}}$	96

input

```
int((x^4+1)^(5/2), x, method=_RETURNVERBOSE)
```

output

```
x*hypergeom([-5/2, 1/4], [5/4], -x^4)
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.45

$$\int (1 + x^4)^{5/2} dx = \frac{1}{77} (7x^9 + 24x^5 + 37x)\sqrt{x^4 + 1} + \frac{40}{77}i\sqrt{i}F(\arcsin\left(\frac{\sqrt{i}}{x}\right) | -1)$$

input `integrate((x^4+1)^(5/2),x, algorithm="fricas")`

output `1/77*(7*x^9 + 24*x^5 + 37*x)*sqrt(x^4 + 1) + 40/77*I*sqrt(I)*elliptic_f(arcsin(sqrt(I)/x), -1)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.52 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.34

$$\int (1 + x^4)^{5/2} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(-\frac{5}{2}, \frac{1}{4} \middle| \frac{5}{4}, x^4 e^{i\pi}\right)}{4\Gamma\left(\frac{5}{4}\right)}$$

input `integrate((x**4+1)**(5/2),x)`

output `x*gamma(1/4)*hyper((-5/2, 1/4), (5/4,), x**4*exp_polar(I*pi))/(4*gamma(5/4))`

Maxima [F]

$$\int (1 + x^4)^{5/2} dx = \int (x^4 + 1)^{\frac{5}{2}} dx$$

input `integrate((x^4+1)^(5/2),x, algorithm="maxima")`

output `integrate((x^4 + 1)^(5/2), x)`

Giac [F]

$$\int (1 + x^4)^{5/2} dx = \int (x^4 + 1)^{\frac{5}{2}} dx$$

input `integrate((x^4+1)^(5/2),x, algorithm="giac")`

output `integrate((x^4 + 1)^(5/2), x)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.14

$$\int (1 + x^4)^{5/2} dx = x {}_2F_1\left(-\frac{5}{2}, \frac{1}{4}; \frac{5}{4}; -x^4\right)$$

input `int((x^4 + 1)^(5/2),x)`

output `x*hypergeom([-5/2, 1/4], 5/4, -x^4)`

Reduce [F]

$$\int (1+x^4)^{5/2} dx = \frac{\sqrt{x^4+1}x^9}{11} + \frac{24\sqrt{x^4+1}x^5}{77} + \frac{37\sqrt{x^4+1}x}{77} + \frac{40\left(\int \frac{\sqrt{x^4+1}}{x^4+1} dx\right)}{77}$$

input `int((x^4+1)^(5/2),x)`

output `(7*sqrt(x**4 + 1)*x**9 + 24*sqrt(x**4 + 1)*x**5 + 37*sqrt(x**4 + 1)*x + 40
*int(sqrt(x**4 + 1)/(x**4 + 1),x))/77`

3.26 $\int (1 + x^4)^{3/2} dx$

Optimal result	200
Mathematica [C] (verified)	200
Rubi [A] (verified)	201
Maple [A] (verified)	202
Fricas [C] (verification not implemented)	203
Sympy [C] (verification not implemented)	203
Maxima [F]	204
Giac [F]	204
Mupad [B] (verification not implemented)	204
Reduce [F]	205

Optimal result

Integrand size = 9, antiderivative size = 72

$$\int (1 + x^4)^{3/2} dx = \frac{2}{7}x\sqrt{1 + x^4} + \frac{1}{7}x(1 + x^4)^{3/2} + \frac{2(1 + x^2)\sqrt{\frac{1+x^4}{(1+x^2)^2}} \operatorname{EllipticF}\left(2 \arctan(x), \frac{1}{2}\right)}{7\sqrt{1 + x^4}}$$

output

```
2/7*x*(x^4+1)^(1/2)+1/7*x*(x^4+1)^(3/2)+2/7*(x^2+1)*((x^4+1)/(x^2+1)^2)^(1/2)*InverseJacobiAM(2*arctan(x),1/2*2^(1/2))/(x^4+1)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 3.75 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.24

$$\int (1 + x^4)^{3/2} dx = x \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, \frac{1}{4}, \frac{5}{4}, -x^4\right)$$

input

```
Integrate[(1 + x^4)^(3/2), x]
```

output `x*Hypergeometric2F1[-3/2, 1/4, 5/4, -x^4]`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.07, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {748, 748, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (x^4 + 1)^{3/2} dx \\
 & \quad \downarrow 748 \\
 & \frac{6}{7} \int \sqrt{x^4 + 1} dx + \frac{1}{7} x (x^4 + 1)^{3/2} \\
 & \quad \downarrow 748 \\
 & \frac{6}{7} \left(\frac{2}{3} \int \frac{1}{\sqrt{x^4 + 1}} dx + \frac{1}{3} \sqrt{x^4 + 1} x \right) + \frac{1}{7} x (x^4 + 1)^{3/2} \\
 & \quad \downarrow 761 \\
 & \frac{6}{7} \left(\frac{(x^2 + 1) \sqrt{\frac{x^4 + 1}{(x^2 + 1)^2}} \operatorname{EllipticF}\left(2 \arctan(x), \frac{1}{2}\right)}{3 \sqrt{x^4 + 1}} + \frac{1}{3} \sqrt{x^4 + 1} x \right) + \frac{1}{7} x (x^4 + 1)^{3/2}
 \end{aligned}$$

input `Int[(1 + x^4)^(3/2), x]`

output `(x*(1 + x^4)^(3/2))/7 + (6*((x*Sqrt[1 + x^4])/3 + ((1 + x^2)*Sqrt[(1 + x^4)/(1 + x^2)^2]*EllipticF[2*ArcTan[x], 1/2])/(3*Sqrt[1 + x^4]))) / 7`

Definitions of rubi rules used

rule 748

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p
+ 1)), x] + Simp[a*n*(p/(n*p + 1)) Int[(a + b*x^n)^(p - 1), x], x] /; Fre
eQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || LtQ[Denominat
or[p + 1/n], Denominator[p]])
```

rule 761

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(
1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*
EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]
```

Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.19

method	result	size
meijerg	$x \operatorname{hypergeom}\left(\left[-\frac{3}{2}, \frac{1}{4}\right], \left[\frac{5}{4}\right], -x^4\right)$	14
risch	$\frac{x(x^4+3)\sqrt{x^4+1}}{7} + \frac{4\sqrt{-ix^2+1}\sqrt{ix^2+1}\operatorname{EllipticF}\left(x\left(\frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right), i\right)}{7\left(\frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right)\sqrt{x^4+1}}$	77
default	$\frac{x^5\sqrt{x^4+1}}{7} + \frac{3x\sqrt{x^4+1}}{7} + \frac{4\sqrt{-ix^2+1}\sqrt{ix^2+1}\operatorname{EllipticF}\left(x\left(\frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right), i\right)}{7\left(\frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right)\sqrt{x^4+1}}$	84
elliptic	$\frac{x^5\sqrt{x^4+1}}{7} + \frac{3x\sqrt{x^4+1}}{7} + \frac{4\sqrt{-ix^2+1}\sqrt{ix^2+1}\operatorname{EllipticF}\left(x\left(\frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right), i\right)}{7\left(\frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right)\sqrt{x^4+1}}$	84

input

```
int((x^4+1)^(3/2), x, method=_RETURNVERBOSE)
```

output

```
x*hypergeom([-3/2, 1/4], [5/4], -x^4)
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.44

$$\int (1+x^4)^{3/2} dx = \frac{1}{7}(x^5+3x)\sqrt{x^4+1} + \frac{4}{7}i\sqrt{i}F\left(\arcsin\left(\frac{\sqrt{i}}{x}\right) \mid -1\right)$$

input `integrate((x^4+1)^(3/2),x, algorithm="fricas")`

output `1/7*(x^5 + 3*x)*sqrt(x^4 + 1) + 4/7*I*sqrt(I)*elliptic_f(arcsin(sqrt(I)/x), -1)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.48 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.40

$$\int (1+x^4)^{3/2} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(-\frac{3}{2}, \frac{1}{4} \mid \frac{5}{4} \mid x^4 e^{i\pi}\right)}{4\Gamma\left(\frac{5}{4}\right)}$$

input `integrate((x**4+1)**(3/2),x)`

output `x*gamma(1/4)*hyper((-3/2, 1/4), (5/4,), x**4*exp_polar(I*pi))/(4*gamma(5/4))`

Maxima [F]

$$\int (1 + x^4)^{3/2} dx = \int (x^4 + 1)^{\frac{3}{2}} dx$$

input `integrate((x^4+1)^(3/2),x, algorithm="maxima")`

output `integrate((x^4 + 1)^(3/2), x)`

Giac [F]

$$\int (1 + x^4)^{3/2} dx = \int (x^4 + 1)^{\frac{3}{2}} dx$$

input `integrate((x^4+1)^(3/2),x, algorithm="giac")`

output `integrate((x^4 + 1)^(3/2), x)`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.17

$$\int (1 + x^4)^{3/2} dx = x {}_2F_1\left(-\frac{3}{2}, \frac{1}{4}; \frac{5}{4}; -x^4\right)$$

input `int((x^4 + 1)^(3/2),x)`

output `x*hypergeom([-3/2, 1/4], 5/4, -x^4)`

Reduce [F]

$$\int (1 + x^4)^{3/2} dx = \frac{\sqrt{x^4 + 1} x^5}{7} + \frac{3\sqrt{x^4 + 1} x}{7} + \frac{4 \left(\int \frac{\sqrt{x^4 + 1}}{x^4 + 1} dx \right)}{7}$$

input `int((x^4+1)^(3/2),x)`

output `(sqrt(x**4 + 1)*x**5 + 3*sqrt(x**4 + 1)*x + 4*int(sqrt(x**4 + 1)/(x**4 + 1),x))/7`

3.27 $\int \sqrt{1+x^4} dx$

Optimal result	206
Mathematica [C] (verified)	206
Rubi [A] (verified)	207
Maple [A] (verified)	208
Fricas [C] (verification not implemented)	208
Sympy [C] (verification not implemented)	209
Maxima [F]	209
Giac [F]	209
Mupad [B] (verification not implemented)	210
Reduce [F]	210

Optimal result

Integrand size = 9, antiderivative size = 58

$$\int \sqrt{1+x^4} dx = \frac{1}{3}x\sqrt{1+x^4} + \frac{(1+x^2)\sqrt{\frac{1+x^4}{(1+x^2)^2}} \operatorname{EllipticF}\left(2\arctan(x), \frac{1}{2}\right)}{3\sqrt{1+x^4}}$$

output

```
1/3*x*(x^4+1)^(1/2)+1/3*(x^2+1)*((x^4+1)/(x^2+1)^2)^(1/2)*InverseJacobiAM(
2*arctan(x),1/2*2^(1/2))/(x^4+1)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.97 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.83

$$\int \sqrt{1+x^4} dx = \frac{x+x^5-2\sqrt[4]{-1}\sqrt{1+x^4}\operatorname{EllipticF}\left(i\operatorname{arcsinh}(\sqrt[4]{-1}x), -1\right)}{3\sqrt{1+x^4}}$$

input

```
Integrate[Sqrt[1+x^4],x]
```

output

```
(x+x^5-2*(-1)^(1/4)*Sqrt[1+x^4]*EllipticF[I*ArcSinh[(-1)^(1/4)*x],-
1])/(3*Sqrt[1+x^4])
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {748, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{x^4 + 1} dx$$

$$\downarrow 748$$

$$\frac{2}{3} \int \frac{1}{\sqrt{x^4 + 1}} dx + \frac{1}{3} \sqrt{x^4 + 1} x$$

$$\downarrow 761$$

$$\frac{(x^2 + 1) \sqrt{\frac{x^4 + 1}{(x^2 + 1)^2}} \text{EllipticF}\left(2 \arctan(x), \frac{1}{2}\right)}{3\sqrt{x^4 + 1}} + \frac{1}{3} \sqrt{x^4 + 1} x$$

input `Int[Sqrt[1 + x^4], x]`

output `(x*Sqrt[1 + x^4])/3 + ((1 + x^2)*Sqrt[(1 + x^4)/(1 + x^2)^2]*EllipticF[2*ArcTan[x], 1/2])/(3*Sqrt[1 + x^4])`

Defintions of rubi rules used

rule 748 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p + 1)), x] + Simp[a*n*(p/(n*p + 1)) Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || LtQ[Denominator[p + 1/n], Denominator[p]])`

rule 761 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]`

Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.24

method	result	size
meijerg	$x \operatorname{hypergeom}\left(\left[-\frac{1}{2}, \frac{1}{4}\right], \left[\frac{5}{4}\right], -x^4\right)$	14
default	$\frac{x\sqrt{x^4+1}}{3} + \frac{2\sqrt{-ix^2+1}\sqrt{ix^2+1}\operatorname{EllipticF}\left(x\left(\frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right), i\right)}{3\left(\frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right)\sqrt{x^4+1}}$	72
risch	$\frac{x\sqrt{x^4+1}}{3} + \frac{2\sqrt{-ix^2+1}\sqrt{ix^2+1}\operatorname{EllipticF}\left(x\left(\frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right), i\right)}{3\left(\frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right)\sqrt{x^4+1}}$	72
elliptic	$\frac{x\sqrt{x^4+1}}{3} + \frac{2\sqrt{-ix^2+1}\sqrt{ix^2+1}\operatorname{EllipticF}\left(x\left(\frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right), i\right)}{3\left(\frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right)\sqrt{x^4+1}}$	72

input `int((x^4+1)^(1/2),x,method=_RETURNVERBOSE)`

output `x*hypergeom([-1/2,1/4],[5/4],-x^4)`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.45

$$\int \sqrt{1+x^4} dx = \frac{1}{3} \sqrt{x^4+1}x + \frac{2}{3}i \sqrt{i} F\left(\arcsin\left(\frac{\sqrt{i}}{x}\right) \mid -1\right)$$

input `integrate((x^4+1)^(1/2),x, algorithm="fricas")`

output `1/3*sqrt(x^4 + 1)*x + 2/3*I*sqrt(I)*elliptic_f(arcsin(sqrt(I)/x), -1)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.40 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.50

$$\int \sqrt{1+x^4} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\begin{matrix} -\frac{1}{2}, \frac{1}{4} \\ \frac{5}{4} \end{matrix} \middle| x^4 e^{i\pi}\right)}{4\Gamma\left(\frac{5}{4}\right)}$$

input `integrate((x**4+1)**(1/2),x)`

output `x*gamma(1/4)*hyper((-1/2, 1/4), (5/4,), x**4*exp_polar(I*pi))/(4*gamma(5/4))`

Maxima [F]

$$\int \sqrt{1+x^4} dx = \int \sqrt{x^4+1} dx$$

input `integrate((x^4+1)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(x^4 + 1), x)`

Giac [F]

$$\int \sqrt{1+x^4} dx = \int \sqrt{x^4+1} dx$$

input `integrate((x^4+1)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(x^4 + 1), x)`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.21

$$\int \sqrt{1+x^4} dx = x {}_2F_1\left(-\frac{1}{2}, \frac{1}{4}; \frac{5}{4}; -x^4\right)$$

input `int((x^4 + 1)^(1/2),x)`output `x*hypergeom([-1/2, 1/4], 5/4, -x^4)`**Reduce [F]**

$$\int \sqrt{1+x^4} dx = \frac{\sqrt{x^4+1}x}{3} + \frac{2\left(\int \frac{\sqrt{x^4+1}}{x^4+1} dx\right)}{3}$$

input `int((x^4+1)^(1/2),x)`output `(sqrt(x**4 + 1)*x + 2*int(sqrt(x**4 + 1)/(x**4 + 1),x))/3`

3.28 $\int \frac{1}{\sqrt{1+x^4}} dx$

Optimal result	211
Mathematica [C] (verified)	211
Rubi [A] (verified)	212
Maple [A] (verified)	212
Fricas [C] (verification not implemented)	213
Sympy [C] (verification not implemented)	213
Maxima [F]	214
Giac [F]	214
Mupad [B] (verification not implemented)	215
Reduce [F]	215

Optimal result

Integrand size = 9, antiderivative size = 43

$$\int \frac{1}{\sqrt{1+x^4}} dx = \frac{(1+x^2) \sqrt{\frac{1+x^4}{(1+x^2)^2}} \text{EllipticF}\left(2 \arctan(x), \frac{1}{2}\right)}{2\sqrt{1+x^4}}$$

output `1/2*(x^2+1)*((x^4+1)/(x^2+1)^2)^(1/2)*InverseJacobiAM(2*arctan(x),1/2*2^(1/2))/(x^4+1)^(1/2)`

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 10.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.49

$$\int \frac{1}{\sqrt{1+x^4}} dx = -\sqrt[4]{-1} \text{EllipticF}\left(i \operatorname{arcsinh}(\sqrt[4]{-1}x), -1\right)$$

input `Integrate[1/Sqrt[1 + x^4],x]`

output `-((-1)^(1/4)*EllipticF[I*ArcSinh[(-1)^(1/4)*x], -1])`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x^4 + 1}} dx$$

↓ 761

$$\frac{(x^2 + 1) \sqrt{\frac{x^4 + 1}{(x^2 + 1)^2}} \text{EllipticF}\left(2 \arctan(x), \frac{1}{2}\right)}{2\sqrt{x^4 + 1}}$$

input `Int[1/Sqrt[1 + x^4], x]`

output `((1 + x^2)*Sqrt[(1 + x^4)/(1 + x^2)^2]*EllipticF[2*ArcTan[x], 1/2])/(2*Sqrt[1 + x^4])`

Defintions of rubi rules used

rule 761 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] :> With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, b}, x] && PosQ[b/a]`

Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.33

method	result	size
meijerg	$x \operatorname{hypergeom}\left(\left[\frac{1}{4}, \frac{1}{2}\right], \left[\frac{5}{4}\right], -x^4\right)$	14
default	$\frac{\sqrt{-ix^2+1}\sqrt{ix^2+1}\operatorname{EllipticF}\left(x\left(\frac{\sqrt{2}+i\sqrt{2}}{2}\right), i\right)}{\left(\frac{\sqrt{2}+i\sqrt{2}}{2}\right)\sqrt{x^4+1}}$	60
elliptic	$\frac{\sqrt{-ix^2+1}\sqrt{ix^2+1}\operatorname{EllipticF}\left(x\left(\frac{\sqrt{2}+i\sqrt{2}}{2}\right), i\right)}{\left(\frac{\sqrt{2}+i\sqrt{2}}{2}\right)\sqrt{x^4+1}}$	60

input `int(1/(x^4+1)^(1/2), x, method=_RETURNVERBOSE)`

output `x*hypergeom([1/4, 1/2], [5/4], -x^4)`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.30

$$\int \frac{1}{\sqrt{1+x^4}} dx = -i \sqrt{i} F(\arcsin(\sqrt{ix}) \mid -1)$$

input `integrate(1/(x^4+1)^(1/2), x, algorithm="fricas")`

output `-I*sqrt(I)*elliptic_f(arcsin(sqrt(I)*x), -1)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.35 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.63

$$\int \frac{1}{\sqrt{1+x^4}} dx = \frac{x \Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2} \mid \frac{5}{4} \mid x^4 e^{i\pi}\right)}{4 \Gamma\left(\frac{5}{4}\right)}$$

input `integrate(1/(x**4+1)**(1/2),x)`

output `x*gamma(1/4)*hyper((1/4, 1/2), (5/4,), x**4*exp_polar(I*pi))/(4*gamma(5/4))`

Maxima [F]

$$\int \frac{1}{\sqrt{1+x^4}} dx = \int \frac{1}{\sqrt{x^4+1}} dx$$

input `integrate(1/(x^4+1)^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(x^4 + 1), x)`

Giac [F]

$$\int \frac{1}{\sqrt{1+x^4}} dx = \int \frac{1}{\sqrt{x^4+1}} dx$$

input `integrate(1/(x^4+1)^(1/2),x, algorithm="giac")`

output `integrate(1/sqrt(x^4 + 1), x)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.28

$$\int \frac{1}{\sqrt{1+x^4}} dx = x {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; -x^4\right)$$

input `int(1/(x^4 + 1)^(1/2),x)`

output `x*hypergeom([1/4, 1/2], 5/4, -x^4)`

Reduce [F]

$$\int \frac{1}{\sqrt{1+x^4}} dx = \int \frac{\sqrt{x^4+1}}{x^4+1} dx$$

input `int(1/(x^4+1)^(1/2),x)`

output `int(sqrt(x**4 + 1)/(x**4 + 1),x)`

3.29 $\int \frac{1}{(1+x^4)^{3/2}} dx$

Optimal result	216
Mathematica [C] (verified)	216
Rubi [A] (verified)	217
Maple [A] (verified)	218
Fricas [C] (verification not implemented)	218
Sympy [C] (verification not implemented)	219
Maxima [F]	219
Giac [F]	219
Mupad [B] (verification not implemented)	220
Reduce [F]	220

Optimal result

Integrand size = 9, antiderivative size = 58

$$\int \frac{1}{(1+x^4)^{3/2}} dx = \frac{x}{2\sqrt{1+x^4}} + \frac{(1+x^2)\sqrt{\frac{1+x^4}{(1+x^2)^2}} \operatorname{EllipticF}\left(2 \arctan(x), \frac{1}{2}\right)}{4\sqrt{1+x^4}}$$

output

```
1/2*x/(x^4+1)^(1/2)+1/4*(x^2+1)*((x^4+1)/(x^2+1)^2)^(1/2)*InverseJacobiAM(
2*arctan(x),1/2*2^(1/2))/(x^4+1)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 3.49 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.52

$$\int \frac{1}{(1+x^4)^{3/2}} dx = \frac{1}{2}x \left(\frac{1}{\sqrt{1+x^4}} + \operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, -x^4\right) \right)$$

input

```
Integrate[(1 + x^4)^(-3/2), x]
```

output

```
(x*(1/Sqrt[1 + x^4] + Hypergeometric2F1[1/4, 1/2, 5/4, -x^4]))/2
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {749, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(x^4 + 1)^{3/2}} dx$$

↓ 749

$$\frac{1}{2} \int \frac{1}{\sqrt{x^4 + 1}} dx + \frac{x}{2\sqrt{x^4 + 1}}$$

↓ 761

$$\frac{(x^2 + 1) \sqrt{\frac{x^4 + 1}{(x^2 + 1)^2}} \text{EllipticF}\left(2 \arctan(x), \frac{1}{2}\right)}{4\sqrt{x^4 + 1}} + \frac{x}{2\sqrt{x^4 + 1}}$$

input `Int[(1 + x^4)^(-3/2), x]`

output `x/(2*Sqrt[1 + x^4]) + ((1 + x^2)*Sqrt[(1 + x^4)/(1 + x^2)^2]*EllipticF[2*ArcTan[x], 1/2])/(4*Sqrt[1 + x^4])`

Defintions of rubi rules used

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

rule 761 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]`

Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.24

method	result	size
meijerg	x hypergeom $\left(\left[\frac{1}{4}, \frac{3}{2}\right], \left[\frac{5}{4}\right], -x^4\right)$	14
default	$\frac{x}{2\sqrt{x^4+1}} + \frac{\sqrt{-ix^2+1}\sqrt{ix^2+1}\operatorname{EllipticF}\left(x\left(\frac{\sqrt{2}+i\sqrt{2}}{2}\right), i\right)}{2\left(\frac{\sqrt{2}+i\sqrt{2}}{2}\right)\sqrt{x^4+1}}$	72
risch	$\frac{x}{2\sqrt{x^4+1}} + \frac{\sqrt{-ix^2+1}\sqrt{ix^2+1}\operatorname{EllipticF}\left(x\left(\frac{\sqrt{2}+i\sqrt{2}}{2}\right), i\right)}{2\left(\frac{\sqrt{2}+i\sqrt{2}}{2}\right)\sqrt{x^4+1}}$	72
elliptic	$\frac{x}{2\sqrt{x^4+1}} + \frac{\sqrt{-ix^2+1}\sqrt{ix^2+1}\operatorname{EllipticF}\left(x\left(\frac{\sqrt{2}+i\sqrt{2}}{2}\right), i\right)}{2\left(\frac{\sqrt{2}+i\sqrt{2}}{2}\right)\sqrt{x^4+1}}$	72

input `int(1/(x^4+1)^(3/2), x, method=_RETURNVERBOSE)`

output `x*hypergeom([1/4, 3/2], [5/4], -x^4)`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.66

$$\int \frac{1}{(1+x^4)^{3/2}} dx = \frac{\sqrt{i}(-ix^4 - i)F(\arcsin(\sqrt{i}x) | -1) + \sqrt{x^4+1}x}{2(x^4+1)}$$

input `integrate(1/(x^4+1)^(3/2), x, algorithm="fricas")`

output `1/2*(sqrt(I)*(-I*x^4 - I)*elliptic_f(arcsin(sqrt(I)*x), -1) + sqrt(x^4 + 1)*x)/(x^4 + 1)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.43 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.47

$$\int \frac{1}{(1+x^4)^{3/2}} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{3}{2} \middle| \frac{5}{4} \middle| x^4 e^{i\pi}\right)}{4\Gamma\left(\frac{5}{4}\right)}$$

input `integrate(1/(x**4+1)**(3/2),x)`

output `x*gamma(1/4)*hyper((1/4, 3/2), (5/4,), x**4*exp_polar(I*pi))/(4*gamma(5/4))`

Maxima [F]

$$\int \frac{1}{(1+x^4)^{3/2}} dx = \int \frac{1}{(x^4+1)^{\frac{3}{2}}} dx$$

input `integrate(1/(x^4+1)^(3/2),x, algorithm="maxima")`

output `integrate((x^4 + 1)^(-3/2), x)`

Giac [F]

$$\int \frac{1}{(1+x^4)^{3/2}} dx = \int \frac{1}{(x^4+1)^{\frac{3}{2}}} dx$$

input `integrate(1/(x^4+1)^(3/2),x, algorithm="giac")`

output `integrate((x^4 + 1)^(-3/2), x)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.21

$$\int \frac{1}{(1+x^4)^{3/2}} dx = x {}_2F_1\left(\frac{1}{4}, \frac{3}{2}; \frac{5}{4}; -x^4\right)$$

input `int(1/(x^4 + 1)^(3/2),x)`output `x*hypergeom([1/4, 3/2], 5/4, -x^4)`**Reduce [F]**

$$\int \frac{1}{(1+x^4)^{3/2}} dx = \int \frac{\sqrt{x^4+1}}{x^8+2x^4+1} dx$$

input `int(1/(x^4+1)^(3/2),x)`output `int(sqrt(x**4 + 1)/(x**8 + 2*x**4 + 1),x)`

3.30 $\int \frac{1}{(1+x^4)^{5/2}} dx$

Optimal result	221
Mathematica [C] (verified)	221
Rubi [A] (verified)	222
Maple [A] (verified)	223
Fricas [C] (verification not implemented)	224
Sympy [C] (verification not implemented)	224
Maxima [F]	225
Giac [F]	225
Mupad [B] (verification not implemented)	225
Reduce [F]	226

Optimal result

Integrand size = 9, antiderivative size = 72

$$\int \frac{1}{(1+x^4)^{5/2}} dx = \frac{x}{6(1+x^4)^{3/2}} + \frac{5x}{12\sqrt{1+x^4}} + \frac{5(1+x^2)\sqrt{\frac{1+x^4}{(1+x^2)^2}} \operatorname{EllipticF}\left(2\arctan(x), \frac{1}{2}\right)}{24\sqrt{1+x^4}}$$

output

$1/6*x/(x^4+1)^{(3/2)}+5/12*x/(x^4+1)^{(1/2)}+5/24*(x^2+1)*((x^4+1)/(x^2+1)^2)^{(1/2)}*\operatorname{InverseJacobiAM}(2*\arctan(x), 1/2*2^{(1/2)})/(x^4+1)^{(1/2)}$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 4.26 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.68

$$\int \frac{1}{(1+x^4)^{5/2}} dx = \frac{x}{6(1+x^4)^{3/2}} + \frac{5x}{12\sqrt{1+x^4}} + \frac{5}{12}x \operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, -x^4\right)$$

input

`Integrate[(1 + x^4)^(-5/2), x]`

output

$$\frac{x}{6(1+x^4)^{3/2}} + \frac{5x}{12\sqrt{1+x^4}} + \frac{5x \operatorname{Hypergeometric2F1}\left[\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, -x^4\right]}{12}$$
Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.07, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {749, 749, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(x^4+1)^{5/2}} dx \\ & \quad \downarrow 749 \\ & \frac{5}{6} \int \frac{1}{(x^4+1)^{3/2}} dx + \frac{x}{6(x^4+1)^{3/2}} \\ & \quad \downarrow 749 \\ & \frac{5}{6} \left(\frac{1}{2} \int \frac{1}{\sqrt{x^4+1}} dx + \frac{x}{2\sqrt{x^4+1}} \right) + \frac{x}{6(x^4+1)^{3/2}} \\ & \quad \downarrow 761 \\ & \frac{5}{6} \left(\frac{(x^2+1) \sqrt{\frac{x^4+1}{(x^2+1)^2}} \operatorname{EllipticF}\left(2 \arctan(x), \frac{1}{2}\right)}{4\sqrt{x^4+1}} + \frac{x}{2\sqrt{x^4+1}} \right) + \frac{x}{6(x^4+1)^{3/2}} \end{aligned}$$

input

$$\operatorname{Int}[(1+x^4)^{-5/2}, x]$$

output

$$\frac{x}{6(1+x^4)^{3/2}} + \frac{5(x/(2\sqrt{1+x^4}) + ((1+x^2)\sqrt{(1+x^4)})/(1+x^2)^2) \operatorname{EllipticF}[2 \operatorname{ArcTan}[x], 1/2]}{4\sqrt{1+x^4}})/6$$

Definitions of rubi rules used

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

rule 761 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]`

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.19

method	result	size
meijerg	$x \text{ hypergeom} \left(\left[\frac{1}{4}, \frac{5}{2} \right], \left[\frac{5}{4} \right], -x^4 \right)$	14
risch	$\frac{x(5x^4+7)}{12(x^4+1)^{\frac{3}{2}}} + \frac{5\sqrt{-ix^2+1}\sqrt{ix^2+1}\text{EllipticF}\left(x\left(\frac{\sqrt{2}}{2}+\frac{i\sqrt{2}}{2}\right),i\right)}{12\left(\frac{\sqrt{2}}{2}+\frac{i\sqrt{2}}{2}\right)\sqrt{x^4+1}}$	79
default	$\frac{x}{6(x^4+1)^{\frac{3}{2}}} + \frac{5x}{12\sqrt{x^4+1}} + \frac{5\sqrt{-ix^2+1}\sqrt{ix^2+1}\text{EllipticF}\left(x\left(\frac{\sqrt{2}}{2}+\frac{i\sqrt{2}}{2}\right),i\right)}{12\left(\frac{\sqrt{2}}{2}+\frac{i\sqrt{2}}{2}\right)\sqrt{x^4+1}}$	82
elliptic	$\frac{x}{6(x^4+1)^{\frac{3}{2}}} + \frac{5x}{12\sqrt{x^4+1}} + \frac{5\sqrt{-ix^2+1}\sqrt{ix^2+1}\text{EllipticF}\left(x\left(\frac{\sqrt{2}}{2}+\frac{i\sqrt{2}}{2}\right),i\right)}{12\left(\frac{\sqrt{2}}{2}+\frac{i\sqrt{2}}{2}\right)\sqrt{x^4+1}}$	82

input `int(1/(x^4+1)^(5/2),x,method=_RETURNVERBOSE)`

output `x*hypergeom([1/4,5/2],[5/4],-x^4)`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.81

$$\int \frac{1}{(1+x^4)^{5/2}} dx = -\frac{5\sqrt{i}(ix^8 + 2ix^4 + i)F(\arcsin(\sqrt{i}x) | -1) - (5x^5 + 7x)\sqrt{x^4 + 1}}{12(x^8 + 2x^4 + 1)}$$

input `integrate(1/(x^4+1)^(5/2),x, algorithm="fricas")`

output `-1/12*(5*sqrt(I)*(I*x^8 + 2*I*x^4 + I)*elliptic_f(arcsin(sqrt(I)*x), -1) - (5*x^5 + 7*x)*sqrt(x^4 + 1))/(x^8 + 2*x^4 + 1)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.49 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.38

$$\int \frac{1}{(1+x^4)^{5/2}} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{5}{2} \middle| \frac{5}{4} \right) x^4 e^{i\pi}}{4\Gamma\left(\frac{5}{4}\right)}$$

input `integrate(1/(x**4+1)**(5/2),x)`

output `x*gamma(1/4)*hyper((1/4, 5/2), (5/4,), x**4*exp_polar(I*pi))/(4*gamma(5/4))`

Maxima [F]

$$\int \frac{1}{(1+x^4)^{5/2}} dx = \int \frac{1}{(x^4+1)^{\frac{5}{2}}} dx$$

input `integrate(1/(x^4+1)^(5/2),x, algorithm="maxima")`

output `integrate((x^4 + 1)^(-5/2), x)`

Giac [F]

$$\int \frac{1}{(1+x^4)^{5/2}} dx = \int \frac{1}{(x^4+1)^{\frac{5}{2}}} dx$$

input `integrate(1/(x^4+1)^(5/2),x, algorithm="giac")`

output `integrate((x^4 + 1)^(-5/2), x)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.17

$$\int \frac{1}{(1+x^4)^{5/2}} dx = x {}_2F_1\left(\frac{1}{4}, \frac{5}{2}; \frac{5}{4}; -x^4\right)$$

input `int(1/(x^4 + 1)^(5/2),x)`

output `x*hypergeom([1/4, 5/2], 5/4, -x^4)`

Reduce [F]

$$\int \frac{1}{(1+x^4)^{5/2}} dx = \int \frac{\sqrt{x^4+1}}{x^{12}+3x^8+3x^4+1} dx$$

input `int(1/(x^4+1)^(5/2),x)`

output `int(sqrt(x**4 + 1)/(x**12 + 3*x**8 + 3*x**4 + 1),x)`

3.31 $\int (a + bx^4)^{7/4} dx$

Optimal result	227
Mathematica [A] (verified)	227
Rubi [A] (verified)	228
Maple [A] (verified)	230
Fricas [C] (verification not implemented)	231
Sympy [C] (verification not implemented)	232
Maxima [A] (verification not implemented)	232
Giac [F]	233
Mupad [B] (verification not implemented)	233
Reduce [F]	233

Optimal result

Integrand size = 11, antiderivative size = 96

$$\int (a + bx^4)^{7/4} dx = \frac{7}{32}ax(a + bx^4)^{3/4} + \frac{1}{8}x(a + bx^4)^{7/4} + \frac{21a^2 \arctan\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a + bx^4}}\right)}{64\sqrt[4]{b}} + \frac{21a^2 \operatorname{arctanh}\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a + bx^4}}\right)}{64\sqrt[4]{b}}$$

output

```
7/32*a*x*(b*x^4+a)^(3/4)+1/8*x*(b*x^4+a)^(7/4)+21/64*a^2*arctan(b^(1/4)*x/
(b*x^4+a)^(1/4))/b^(1/4)+21/64*a^2*arctanh(b^(1/4)*x/(b*x^4+a)^(1/4))/b^(1/4)
```

Mathematica [A] (verified)

Time = 0.39 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.93

$$\int (a + bx^4)^{7/4} dx = \frac{1}{32}x(a + bx^4)^{3/4} (11a + 4bx^4) + \frac{21a^2 \arctan\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a + bx^4}}\right)}{64\sqrt[4]{b}} + \frac{21a^2 \operatorname{arctanh}\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a + bx^4}}\right)}{64\sqrt[4]{b}}$$

input `Integrate[(a + b*x^4)^(7/4), x]`

output $(x*(a + b*x^4)^{(3/4)}*(11*a + 4*b*x^4))/32 + (21*a^2*ArcTan[(b^{(1/4)}*x)/(a + b*x^4)^{(1/4)}])/(64*b^{(1/4)}) + (21*a^2*ArcTanh[(b^{(1/4)}*x)/(a + b*x^4)^{(1/4)}])/(64*b^{(1/4)})$

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.05, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {748, 748, 770, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + bx^4)^{7/4} dx \\
 & \quad \downarrow 748 \\
 & \frac{7}{8}a \int (bx^4 + a)^{3/4} dx + \frac{1}{8}x(a + bx^4)^{7/4} \\
 & \quad \downarrow 748 \\
 & \frac{7}{8}a \left(\frac{3}{4}a \int \frac{1}{\sqrt[4]{bx^4 + a}} dx + \frac{1}{4}x(a + bx^4)^{3/4} \right) + \frac{1}{8}x(a + bx^4)^{7/4} \\
 & \quad \downarrow 770 \\
 & \frac{7}{8}a \left(\frac{3}{4}a \int \frac{1}{1 - \frac{bx^4}{bx^4 + a}} d \frac{x}{\sqrt[4]{bx^4 + a}} + \frac{1}{4}x(a + bx^4)^{3/4} \right) + \frac{1}{8}x(a + bx^4)^{7/4} \\
 & \quad \downarrow 756 \\
 & \frac{7}{8}a \left(\frac{3}{4}a \left(\frac{1}{2} \int \frac{1}{1 - \frac{\sqrt{bx^2}}{\sqrt{bx^4 + a}}} d \frac{x}{\sqrt[4]{bx^4 + a}} + \frac{1}{2} \int \frac{1}{\frac{\sqrt{bx^2}}{\sqrt{bx^4 + a}} + 1} d \frac{x}{\sqrt[4]{bx^4 + a}} \right) + \frac{1}{4}x(a + bx^4)^{3/4} \right) + \\
 & \quad \frac{1}{8}x(a + bx^4)^{7/4} \\
 & \quad \downarrow 216
 \end{aligned}$$

$$\frac{7}{8}a \left(\frac{3}{4}a \left(\frac{1}{2} \int \frac{1}{1 - \frac{\sqrt{bx^2}}{\sqrt{bx^4+a}}} d \frac{x}{\sqrt{bx^4+a}} + \frac{\arctan\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a+bx^4}}\right)}{2\sqrt[4]{b}} \right) + \frac{1}{4}x(a+bx^4)^{3/4} \right) + \frac{1}{8}x(a+bx^4)^{7/4}$$

↓ 219

$$\frac{7}{8}a \left(\frac{3}{4}a \left(\frac{\arctan\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a+bx^4}}\right)}{2\sqrt[4]{b}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a+bx^4}}\right)}{2\sqrt[4]{b}} \right) + \frac{1}{4}x(a+bx^4)^{3/4} \right) + \frac{1}{8}x(a+bx^4)^{7/4}$$

input `Int[(a + b*x^4)^(7/4), x]`

output `(x*(a + b*x^4)^(7/4))/8 + (7*a*((x*(a + b*x^4)^(3/4))/4 + (3*a*(ArcTan[(b^(1/4)*x)/(a + b*x^4)^(1/4)]/(2*b^(1/4)) + ArcTanh[(b^(1/4)*x)/(a + b*x^4)^(1/4)]/(2*b^(1/4)))/4))/8`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 748 `Int[((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p + 1)), x] + Simp[a*n*(p/(n*p + 1)) Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || LtQ[Denominator[p + 1/n], Denominator[p]])`

rule 756 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 770 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^(p + 1/n) Subst[Int[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/n]`

Maple [A] (verified)

Time = 1.26 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.06

method	result	size
pseudoelliptic	$\frac{16(bx^4+a)^{\frac{3}{4}}b^{\frac{5}{4}}x^5+44ax(bx^4+a)^{\frac{3}{4}}b^{\frac{1}{4}}-42\arctan\left(\frac{(bx^4+a)^{\frac{1}{4}}}{b^{\frac{1}{4}}x}\right)a^2+21\ln\left(\frac{b^{\frac{1}{4}}x+(bx^4+a)^{\frac{1}{4}}}{-b^{\frac{1}{4}}x+(bx^4+a)^{\frac{1}{4}}}\right)a^2}{128b^{\frac{1}{4}}}$	102

input `int((b*x^4+a)^(7/4),x,method=_RETURNVERBOSE)`

output `1/128*(16*(b*x^4+a)^(3/4)*b^(5/4)*x^5+44*a*x*(b*x^4+a)^(3/4)*b^(1/4)-42*arctan(1/b^(1/4)/x*(b*x^4+a)^(1/4))*a^2+21*ln((b^(1/4)*x+(b*x^4+a)^(1/4))/(-b^(1/4)*x+(b*x^4+a)^(1/4)))*a^2)/b^(1/4)`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 198, normalized size of antiderivative = 2.06

$$\begin{aligned} \int (a + bx^4)^{7/4} dx &= \frac{1}{32} (4bx^5 + 11ax)(bx^4 + a)^{3/4} \\ &+ \frac{21}{128} \left(\frac{a^8}{b}\right)^{1/4} \log\left(\frac{9261 \left((bx^4 + a)^{1/4} a^6 + \left(\frac{a^8}{b}\right)^{3/4} bx\right)}{x}\right) \\ &- \frac{21}{128} i \left(\frac{a^8}{b}\right)^{1/4} \log\left(\frac{9261 \left((bx^4 + a)^{1/4} a^6 + i \left(\frac{a^8}{b}\right)^{3/4} bx\right)}{x}\right) \\ &+ \frac{21}{128} i \left(\frac{a^8}{b}\right)^{1/4} \log\left(\frac{9261 \left((bx^4 + a)^{1/4} a^6 - i \left(\frac{a^8}{b}\right)^{3/4} bx\right)}{x}\right) \\ &- \frac{21}{128} \left(\frac{a^8}{b}\right)^{1/4} \log\left(\frac{9261 \left((bx^4 + a)^{1/4} a^6 - \left(\frac{a^8}{b}\right)^{3/4} bx\right)}{x}\right) \end{aligned}$$

input `integrate((b*x^4+a)^(7/4),x, algorithm="fricas")`

output `1/32*(4*b*x^5 + 11*a*x)*(b*x^4 + a)^(3/4) + 21/128*(a^8/b)^(1/4)*log(9261*
((b*x^4 + a)^(1/4)*a^6 + (a^8/b)^(3/4)*b*x)/x) - 21/128*I*(a^8/b)^(1/4)*lo
g(9261*((b*x^4 + a)^(1/4)*a^6 + I*(a^8/b)^(3/4)*b*x)/x) + 21/128*I*(a^8/b)
^(1/4)*log(9261*((b*x^4 + a)^(1/4)*a^6 - I*(a^8/b)^(3/4)*b*x)/x) - 21/128*
(a^8/b)^(1/4)*log(9261*((b*x^4 + a)^(1/4)*a^6 - (a^8/b)^(3/4)*b*x)/x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.24 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.39

$$\int (a + bx^4)^{7/4} dx = \frac{a^{7/4} x \Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\begin{matrix} -\frac{7}{4}, \frac{1}{4} \\ \frac{5}{4} \end{matrix} \middle| \frac{bx^4 e^{i\pi}}{a} \right)}{4\Gamma\left(\frac{5}{4}\right)}$$

input `integrate((b*x**4+a)**(7/4),x)`

output `a**(7/4)*x*gamma(1/4)*hyper((-7/4, 1/4), (5/4,), b*x**4*exp_polar(I*pi)/a)/(4*gamma(5/4))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.49

$$\int (a + bx^4)^{7/4} dx = -\frac{21}{128} a^2 \left(\frac{2 \arctan\left(\frac{(bx^4+a)^{1/4}}{b^{1/4}x}\right)}{b^{1/4}} + \frac{\log\left(-\frac{b^{1/4} - (bx^4+a)^{1/4}}{b^{1/4} + (bx^4+a)^{1/4}}\right)}{b^{1/4}} \right) - \frac{\frac{7(bx^4+a)^{3/4}a^2b}{x^3} - \frac{11(bx^4+a)^{7/4}a^2}{x^7}}{32\left(b^2 - \frac{2(bx^4+a)b}{x^4} + \frac{(bx^4+a)^2}{x^8}\right)}$$

input `integrate((b*x^4+a)^(7/4),x, algorithm="maxima")`

output `-21/128*a^2*(2*arctan((b*x^4 + a)^(1/4)/(b^(1/4)*x))/b^(1/4) + log(-(b^(1/4) - (b*x^4 + a)^(1/4)/x)/(b^(1/4) + (b*x^4 + a)^(1/4)/x))/b^(1/4)) - 1/32*(7*(b*x^4 + a)^(3/4)*a^2*b/x^3 - 11*(b*x^4 + a)^(7/4)*a^2/x^7)/(b^2 - 2*(b*x^4 + a)*b/x^4 + (b*x^4 + a)^2/x^8)`

Giac [F]

$$\int (a + bx^4)^{7/4} dx = \int (bx^4 + a)^{7/4} dx$$

input `integrate((b*x^4+a)^(7/4),x, algorithm="giac")`

output `integrate((b*x^4 + a)^(7/4), x)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.39

$$\int (a + bx^4)^{7/4} dx = \frac{x (bx^4 + a)^{7/4} {}_2F_1\left(-\frac{7}{4}, \frac{1}{4}; \frac{5}{4}; -\frac{bx^4}{a}\right)}{\left(\frac{bx^4}{a} + 1\right)^{7/4}}$$

input `int((a + b*x^4)^(7/4),x)`

output `(x*(a + b*x^4)^(7/4)*hypergeom([-7/4, 1/4], 5/4, -(b*x^4)/a))/((b*x^4)/a + 1)^(7/4)`

Reduce [F]

$$\int (a + bx^4)^{7/4} dx = \frac{11(bx^4 + a)^{3/4} ax}{32} + \frac{(bx^4 + a)^{3/4} bx^5}{8} + \frac{21\left(\int \frac{1}{(bx^4+a)^{1/4}} dx\right) a^2}{32}$$

input `int((b*x^4+a)^(7/4),x)`

output `(11*(a + b*x**4)**(3/4)*a*x + 4*(a + b*x**4)**(3/4)*b*x**5 + 21*int((a + b*x**4)**(3/4)/(a + b*x**4),x)*a**2)/32`

3.32 $\int (a + bx^4)^{3/4} dx$

Optimal result	234
Mathematica [A] (verified)	234
Rubi [A] (verified)	235
Maple [A] (verified)	237
Fricas [C] (verification not implemented)	237
Sympy [C] (verification not implemented)	238
Maxima [A] (verification not implemented)	238
Giac [F]	239
Mupad [B] (verification not implemented)	239
Reduce [F]	240

Optimal result

Integrand size = 11, antiderivative size = 75

$$\int (a + bx^4)^{3/4} dx = \frac{1}{4}x(a + bx^4)^{3/4} + \frac{3a \arctan\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a + bx^4}}\right)}{8\sqrt[4]{b}} + \frac{3a \operatorname{arctanh}\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a + bx^4}}\right)}{8\sqrt[4]{b}}$$

```
output 1/4*x*(b*x^4+a)^(3/4)+3/8*a*arctan(b^(1/4)*x/(b*x^4+a)^(1/4))/b^(1/4)+3/8*
a*arctanh(b^(1/4)*x/(b*x^4+a)^(1/4))/b^(1/4)
```

Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.00

$$\int (a + bx^4)^{3/4} dx = \frac{1}{4}x(a + bx^4)^{3/4} + \frac{3a \arctan\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a + bx^4}}\right)}{8\sqrt[4]{b}} + \frac{3a \operatorname{arctanh}\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a + bx^4}}\right)}{8\sqrt[4]{b}}$$

```
input Integrate[(a + b*x^4)^(3/4),x]
```

```
output (x*(a + b*x^4)^(3/4))/4 + (3*a*ArcTan[(b^(1/4)*x)/(a + b*x^4)^(1/4)])/(8*b
^(1/4)) + (3*a*ArcTanh[(b^(1/4)*x)/(a + b*x^4)^(1/4)])/(8*b^(1/4))
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.05, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {748, 770, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + bx^4)^{3/4} dx \\
 & \quad \downarrow 748 \\
 & \frac{3}{4}a \int \frac{1}{\sqrt[4]{bx^4 + a}} dx + \frac{1}{4}x(a + bx^4)^{3/4} \\
 & \quad \downarrow 770 \\
 & \frac{3}{4}a \int \frac{1}{1 - \frac{bx^4}{bx^4 + a}} d \frac{x}{\sqrt[4]{bx^4 + a}} + \frac{1}{4}x(a + bx^4)^{3/4} \\
 & \quad \downarrow 756 \\
 & \frac{3}{4}a \left(\frac{1}{2} \int \frac{1}{1 - \frac{\sqrt{bx^2}}{\sqrt{bx^4 + a}}} d \frac{x}{\sqrt[4]{bx^4 + a}} + \frac{1}{2} \int \frac{1}{\frac{\sqrt{bx^2}}{\sqrt{bx^4 + a}} + 1} d \frac{x}{\sqrt[4]{bx^4 + a}} \right) + \frac{1}{4}x(a + bx^4)^{3/4} \\
 & \quad \downarrow 216 \\
 & \frac{3}{4}a \left(\frac{1}{2} \int \frac{1}{1 - \frac{\sqrt{bx^2}}{\sqrt{bx^4 + a}}} d \frac{x}{\sqrt[4]{bx^4 + a}} + \frac{\arctan\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a + bx^4}}\right)}{2\sqrt[4]{b}} \right) + \frac{1}{4}x(a + bx^4)^{3/4} \\
 & \quad \downarrow 219 \\
 & \frac{3}{4}a \left(\frac{\arctan\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a + bx^4}}\right)}{2\sqrt[4]{b}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a + bx^4}}\right)}{2\sqrt[4]{b}} \right) + \frac{1}{4}x(a + bx^4)^{3/4}
 \end{aligned}$$

input `Int[(a + b*x^4)^(3/4), x]`

output $(x*(a + b*x^4)^{(3/4)}/4 + (3*a*(\text{ArcTan}[(b^{(1/4)}*x)/(a + b*x^4)^{(1/4)}]/(2*b^{(1/4)}) + \text{ArcTanh}[(b^{(1/4)}*x)/(a + b*x^4)^{(1/4)}]/(2*b^{(1/4)})))/4$

Defintions of rubi rules used

rule 216 $\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

rule 219 $\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 748 $\text{Int}[(a_ + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Simp}[x*((a + b*x^n)^p/(n*p + 1)), x] + \text{Simp}[a*n*(p/(n*p + 1)) \ \text{Int}[(a + b*x^n)^{(p - 1)}, x], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ (\text{IntegerQ}[2*p] \ || \ \text{LtQ}[\text{Denominator}[p + 1/n], \text{Denominator}[p]])$

rule 756 $\text{Int}[(a_ + (b_)*(x_)^4)^{-1}, x_Symbol] \rightarrow \text{With}\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[r/(2*a) \ \text{Int}[1/(r - s*x^2), x], x] + \text{Simp}[r/(2*a) \ \text{Int}[1/(r + s*x^2), x], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ !\text{GtQ}[a/b, 0]$

rule 770 $\text{Int}[(a_ + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Simp}[a^{(p + 1/n)} \ \text{Subst}[\text{Int}[1/(1 - b*x^n)^{(p + 1/n + 1)}, x], x, x/(a + b*x^n)^{(1/n)}], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[-1, p, 0] \ \&\& \ \text{NeQ}[p, -2^{(-1)}] \ \&\& \ \text{IntegerQ}[p + 1/n]$

Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.07

method	result	size
pseudoelliptic	$\frac{4(bx^4+a)^{\frac{3}{4}}x b^{\frac{1}{4}} - 6 \arctan\left(\frac{(bx^4+a)^{\frac{1}{4}}}{b^{\frac{1}{4}}x}\right) a + 3 \ln\left(\frac{b^{\frac{1}{4}}x + (bx^4+a)^{\frac{1}{4}}}{-b^{\frac{1}{4}}x + (bx^4+a)^{\frac{1}{4}}}\right) a}{16b^{\frac{1}{4}}}$	80

input `int((b*x^4+a)^(3/4),x,method=_RETURNVERBOSE)`output `1/16*(4*(b*x^4+a)^(3/4)*x*b^(1/4)-6*arctan(1/b^(1/4)/x*(b*x^4+a)^(1/4))*a+3*ln((b^(1/4)*x+(b*x^4+a)^(1/4))/(-b^(1/4)*x+(b*x^4+a)^(1/4)))*a)/b^(1/4)`**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 188, normalized size of antiderivative = 2.51

$$\int (a + bx^4)^{3/4} dx = \frac{1}{4} (bx^4 + a)^{\frac{3}{4}} x$$

$$+ \frac{3}{16} \left(\frac{a^4}{b}\right)^{\frac{1}{4}} \log\left(\frac{27 \left((bx^4 + a)^{\frac{1}{4}} a^3 + \left(\frac{a^4}{b}\right)^{\frac{3}{4}} bx\right)}{x}\right)$$

$$- \frac{3}{16} i \left(\frac{a^4}{b}\right)^{\frac{1}{4}} \log\left(\frac{27 \left((bx^4 + a)^{\frac{1}{4}} a^3 + i \left(\frac{a^4}{b}\right)^{\frac{3}{4}} bx\right)}{x}\right)$$

$$+ \frac{3}{16} i \left(\frac{a^4}{b}\right)^{\frac{1}{4}} \log\left(\frac{27 \left((bx^4 + a)^{\frac{1}{4}} a^3 - i \left(\frac{a^4}{b}\right)^{\frac{3}{4}} bx\right)}{x}\right)$$

$$- \frac{3}{16} \left(\frac{a^4}{b}\right)^{\frac{1}{4}} \log\left(\frac{27 \left((bx^4 + a)^{\frac{1}{4}} a^3 - \left(\frac{a^4}{b}\right)^{\frac{3}{4}} bx\right)}{x}\right)$$

input `integrate((b*x^4+a)^(3/4),x, algorithm="fricas")`

output `1/4*(b*x^4 + a)^(3/4)*x + 3/16*(a^4/b)^(1/4)*log(27*((b*x^4 + a)^(1/4)*a^3 + (a^4/b)^(3/4)*b*x)/x) - 3/16*I*(a^4/b)^(1/4)*log(27*((b*x^4 + a)^(1/4)*a^3 + I*(a^4/b)^(3/4)*b*x)/x) + 3/16*I*(a^4/b)^(1/4)*log(27*((b*x^4 + a)^(1/4)*a^3 - I*(a^4/b)^(3/4)*b*x)/x) - 3/16*(a^4/b)^(1/4)*log(27*((b*x^4 + a)^(1/4)*a^3 - (a^4/b)^(3/4)*b*x)/x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.73 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.49

$$\int (a + bx^4)^{3/4} dx = \frac{a^{3/4} x \Gamma\left(\frac{1}{4}\right) {}_2F_1\left(-\frac{3}{4}, \frac{1}{4} \middle| \frac{bx^4 e^{i\pi}}{a}\right)}{4\Gamma\left(\frac{5}{4}\right)}$$

input `integrate((b*x**4+a)**(3/4),x)`

output `a**(3/4)*x*gamma(1/4)*hyper((-3/4, 1/4), (5/4,), b*x**4*exp_polar(I*pi)/a)/(4*gamma(5/4))`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.36

$$\int (a + bx^4)^{3/4} dx = -\frac{3}{16} a \left(\frac{2 \arctan\left(\frac{(bx^4+a)^{1/4}}{b^{1/4}x}\right)}{b^{1/4}} + \frac{\log\left(-\frac{b^{1/4} - \frac{(bx^4+a)^{1/4}}{x}}{b^{1/4} + \frac{(bx^4+a)^{1/4}}{x}}\right)}{b^{1/4}} \right) - \frac{(bx^4 + a)^{3/4} a}{4 \left(b - \frac{bx^4+a}{x^4}\right) x^3}$$

input `integrate((b*x^4+a)^(3/4),x, algorithm="maxima")`

output
$$-\frac{3}{16}a \cdot (2 \arctan((b \cdot x^4 + a)^{1/4} / (b^{1/4} \cdot x)) / b^{1/4} + \log(-(b^{1/4} - (b \cdot x^4 + a)^{1/4} / x) / (b^{1/4} + (b \cdot x^4 + a)^{1/4} / x)) / b^{1/4}) - \frac{1}{4} \cdot (b \cdot x^4 + a)^{3/4} \cdot a / ((b - (b \cdot x^4 + a) / x^4) \cdot x^3)$$

Giac [F]

$$\int (a + bx^4)^{3/4} dx = \int (bx^4 + a)^{3/4} dx$$

input `integrate((b*x^4+a)^(3/4),x, algorithm="giac")`

output `integrate((b*x^4 + a)^(3/4), x)`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.49

$$\int (a + bx^4)^{3/4} dx = \frac{x (bx^4 + a)^{3/4} {}_2F_1\left(-\frac{3}{4}, \frac{1}{4}; \frac{5}{4}; -\frac{bx^4}{a}\right)}{\left(\frac{bx^4}{a} + 1\right)^{3/4}}$$

input `int((a + b*x^4)^(3/4),x)`

output
$$(x \cdot (a + b \cdot x^4)^{3/4} \cdot \text{hypergeom}([-3/4, 1/4], 5/4, -(b \cdot x^4) / a)) / ((b \cdot x^4) / a + 1)^{3/4}$$

Reduce [F]

$$\int (a + bx^4)^{3/4} dx = \frac{(bx^4 + a)^{3/4} x}{4} + \frac{3 \left(\int \frac{1}{(bx^4 + a)^{1/4}} dx \right) a}{4}$$

input `int((b*x^4+a)^(3/4),x)`

output `((a + b*x**4)**(3/4)*x + 3*int((a + b*x**4)**(3/4)/(a + b*x**4),x)*a)/4`

3.33 $\int \frac{1}{\sqrt[4]{a + bx^4}} dx$

Optimal result	241
Mathematica [A] (verified)	241
Rubi [A] (verified)	242
Maple [A] (verified)	243
Fricas [C] (verification not implemented)	244
Sympy [C] (verification not implemented)	244
Maxima [A] (verification not implemented)	245
Giac [F]	245
Mupad [B] (verification not implemented)	246
Reduce [F]	246

Optimal result

Integrand size = 11, antiderivative size = 57

$$\int \frac{1}{\sqrt[4]{a + bx^4}} dx = \frac{\arctan\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a + bx^4}}\right)}{2\sqrt[4]{b}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a + bx^4}}\right)}{2\sqrt[4]{b}}$$

output `1/2*arctan(b^(1/4)*x/(b*x^4+a)^(1/4))/b^(1/4)+1/2*arctanh(b^(1/4)*x/(b*x^4+a)^(1/4))/b^(1/4)`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.33

$$\begin{aligned} & \int \frac{1}{\sqrt[4]{a + bx^4}} dx \\ &= \frac{2 \arctan\left(\frac{\sqrt[4]{bx}}{\sqrt[4]{a + bx^4}}\right) - \log\left(1 - \frac{\sqrt[4]{bx}}{\sqrt[4]{a + bx^4}}\right) + \log\left(1 + \frac{\sqrt[4]{bx}}{\sqrt[4]{a + bx^4}}\right)}{4\sqrt[4]{b}} \end{aligned}$$

input `Integrate[(a + b*x^4)^(-1/4),x]`

output

```
(2*ArcTan[(b^(1/4)*x)/(a + b*x^4)^(1/4)] - Log[1 - (b^(1/4)*x)/(a + b*x^4)^(1/4)] + Log[1 + (b^(1/4)*x)/(a + b*x^4)^(1/4)])/(4*b^(1/4))
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {770, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt[4]{a + bx^4}} dx$$

$$\downarrow 770$$

$$\int \frac{1}{1 - \frac{bx^4}{a+bx^4}} d \frac{x}{\sqrt[4]{a + bx^4}}$$

$$\downarrow 756$$

$$\frac{1}{2} \int \frac{1}{1 - \frac{\sqrt{bx^2}}{\sqrt{bx^4+a}}} d \frac{x}{\sqrt[4]{bx^4 + a}} + \frac{1}{2} \int \frac{1}{\frac{\sqrt{bx^2}}{\sqrt{bx^4+a}} + 1} d \frac{x}{\sqrt[4]{bx^4 + a}}$$

$$\downarrow 216$$

$$\frac{1}{2} \int \frac{1}{1 - \frac{\sqrt{bx^2}}{\sqrt{bx^4+a}}} d \frac{x}{\sqrt[4]{bx^4 + a}} + \frac{\arctan\left(\frac{\sqrt[4]{b}x}{\sqrt[4]{a + bx^4}}\right)}{2\sqrt[4]{b}}$$

$$\downarrow 219$$

$$\frac{\arctan\left(\frac{\sqrt[4]{b}x}{\sqrt[4]{a + bx^4}}\right)}{2\sqrt[4]{b}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{b}x}{\sqrt[4]{a + bx^4}}\right)}{2\sqrt[4]{b}}$$

input

```
Int[(a + b*x^4)^(-1/4), x]
```

output $\text{ArcTan}[(b^{1/4}x)/(a + b x^4)^{1/4}]/(2b^{1/4}) + \text{ArcTanh}[(b^{1/4}x)/(a + b x^4)^{1/4}]/(2b^{1/4})$

Defintions of rubi rules used

rule 216 $\text{Int}[(a_ + (b_)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

rule 219 $\text{Int}[(a_ + (b_)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 756 $\text{Int}[(a_ + (b_)(x_)^4)^{-1}, x_Symbol] \rightarrow \text{With}\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[r/(2*a) \ \text{Int}[1/(r - s*x^2), x], x] + \text{Simp}[r/(2*a) \ \text{Int}[1/(r + s*x^2), x], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ !\text{GtQ}[a/b, 0]$

rule 770 $\text{Int}[(a_ + (b_)(x_)^{n_})^{p_}, x_Symbol] \rightarrow \text{Simp}[a^{(p + 1/n)} \ \text{Subst}[\text{Int}[1/(1 - b*x^n)^{(p + 1/n + 1)}, x], x, x/(a + b*x^n)^{1/n}], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[-1, p, 0] \ \&\& \ \text{NeQ}[p, -2^{(-1)}] \ \&\& \ \text{IntegerQ}[p + 1/n]$

Maple [A] (verified)

Time = 0.56 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.07

method	result	size
pseudoelliptic	$\frac{-2 \arctan\left(\frac{(bx^4+a)^{\frac{1}{4}}}{b^{\frac{1}{4}}x}\right) + \ln\left(\frac{b^{\frac{1}{4}}x + (bx^4+a)^{\frac{1}{4}}}{-b^{\frac{1}{4}}x + (bx^4+a)^{\frac{1}{4}}}\right)}{4b^{\frac{1}{4}}}$	61

input $\text{int}(1/(b*x^4+a)^{1/4}, x, \text{method}=_RETURNVERBOSE)$

output $\frac{1}{4} \cdot (-2 \cdot \arctan(1/b^{1/4}/x \cdot (b \cdot x^4 + a)^{1/4}) + \ln((b^{1/4} \cdot x + (b \cdot x^4 + a)^{1/4}) / (-b^{1/4} \cdot x + (b \cdot x^4 + a)^{1/4}))) / b^{1/4}$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.07 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.86

$$\int \frac{1}{\sqrt[4]{a + bx^4}} dx = \frac{\log\left(\frac{b^{1/4}x + (bx^4 + a)^{1/4}}{x}\right)}{4b^{1/4}} - \frac{\log\left(-\frac{b^{1/4}x - (bx^4 + a)^{1/4}}{x}\right)}{4b^{1/4}} - \frac{i \log\left(\frac{i b^{1/4}x + (bx^4 + a)^{1/4}}{x}\right)}{4b^{1/4}} + \frac{i \log\left(\frac{-i b^{1/4}x + (bx^4 + a)^{1/4}}{x}\right)}{4b^{1/4}}$$

input `integrate(1/(b*x^4+a)^(1/4),x, algorithm="fricas")`

output $\frac{1}{4} \cdot \log((b^{1/4} \cdot x + (b \cdot x^4 + a)^{1/4})/x) / b^{1/4} - \frac{1}{4} \cdot \log(-(b^{1/4} \cdot x - (b \cdot x^4 + a)^{1/4})/x) / b^{1/4} - \frac{1}{4} \cdot I \cdot \log((I \cdot b^{1/4} \cdot x + (b \cdot x^4 + a)^{1/4})/x) / b^{1/4} + \frac{1}{4} \cdot I \cdot \log((-I \cdot b^{1/4} \cdot x + (b \cdot x^4 + a)^{1/4})/x) / b^{1/4}$

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.57 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.63

$$\int \frac{1}{\sqrt[4]{a + bx^4}} dx = \frac{x \Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{4} \middle| \frac{bx^4 e^{i\pi}}{a}\right)}{4 \sqrt[4]{a} \Gamma\left(\frac{5}{4}\right)}$$

input `integrate(1/(b*x**4+a)**(1/4),x)`

output `x*gamma(1/4)*hyper((1/4, 1/4), (5/4,), b*x**4*exp_polar(I*pi)/a)/(4*a**(1/4)*gamma(5/4))`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.19

$$\int \frac{1}{\sqrt[4]{a+bx^4}} dx = -\frac{\arctan\left(\frac{(bx^4+a)^{\frac{1}{4}}}{b^{\frac{1}{4}}x}\right)}{2b^{\frac{1}{4}}} - \frac{\log\left(\frac{-b^{\frac{1}{4}} - \frac{(bx^4+a)^{\frac{1}{4}}}{x}}{b^{\frac{1}{4}} + \frac{(bx^4+a)^{\frac{1}{4}}}{x}}\right)}{4b^{\frac{1}{4}}}$$

input `integrate(1/(b*x^4+a)^(1/4),x, algorithm="maxima")`

output `-1/2*arctan((b*x^4 + a)^(1/4)/(b^(1/4)*x))/b^(1/4) - 1/4*log(-(b^(1/4) - (b*x^4 + a)^(1/4)/x)/(b^(1/4) + (b*x^4 + a)^(1/4)/x))/b^(1/4)`

Giac [F]

$$\int \frac{1}{\sqrt[4]{a+bx^4}} dx = \int \frac{1}{(bx^4+a)^{\frac{1}{4}}} dx$$

input `integrate(1/(b*x^4+a)^(1/4),x, algorithm="giac")`

output `integrate((b*x^4 + a)^(-1/4), x)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.65

$$\int \frac{1}{\sqrt[4]{a + bx^4}} dx = \frac{x \left(\frac{bx^4}{a} + 1 \right)^{1/4} {}_2F_1 \left(\frac{1}{4}, \frac{1}{4}; \frac{5}{4}; -\frac{bx^4}{a} \right)}{(bx^4 + a)^{1/4}}$$

input `int(1/(a + b*x^4)^(1/4),x)`output `(x*((b*x^4)/a + 1)^(1/4)*hypergeom([1/4, 1/4], 5/4, -(b*x^4)/a))/(a + b*x^4)^(1/4)`**Reduce [F]**

$$\int \frac{1}{\sqrt[4]{a + bx^4}} dx = \int \frac{1}{(bx^4 + a)^{1/4}} dx$$

input `int(1/(b*x^4+a)^(1/4),x)`output `int(1/(a + b*x**4)**(1/4),x)`

3.34 $\int \frac{1}{(a+bx^4)^{5/4}} dx$

Optimal result	247
Mathematica [A] (verified)	247
Rubi [A] (verified)	248
Maple [A] (verified)	248
Fricas [A] (verification not implemented)	249
Sympy [B] (verification not implemented)	249
Maxima [A] (verification not implemented)	250
Giac [F]	250
Mupad [B] (verification not implemented)	250
Reduce [F]	251

Optimal result

Integrand size = 11, antiderivative size = 16

$$\int \frac{1}{(a+bx^4)^{5/4}} dx = \frac{x}{a\sqrt[4]{a+bx^4}}$$

output `x/a/(b*x^4+a)^(1/4)`

Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a+bx^4)^{5/4}} dx = \frac{x}{a\sqrt[4]{a+bx^4}}$$

input `Integrate[(a + b*x^4)^(-5/4),x]`

output `x/(a*(a + b*x^4)^(1/4))`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {746}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + bx^4)^{5/4}} dx$$

\downarrow 746
 $\frac{x}{a\sqrt[4]{a + bx^4}}$

input `Int[(a + b*x^4)^(-5/4),x]`

output `x/(a*(a + b*x^4)^(1/4))`

Defintions of rubi rules used

rule 746 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[x*((a + b*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]`

Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result	size
gosper	$\frac{x}{a(bx^4+a)^{\frac{1}{4}}}$	15
trager	$\frac{x}{a(bx^4+a)^{\frac{1}{4}}}$	15
pseudoelliptic	$\frac{x}{a(bx^4+a)^{\frac{1}{4}}}$	15
orering	$\frac{x}{a(bx^4+a)^{\frac{1}{4}}}$	15

input `int(1/(b*x^4+a)^(5/4),x,method=_RETURNVERBOSE)`

output `x/a/(b*x^4+a)^(1/4)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.44

$$\int \frac{1}{(a + bx^4)^{5/4}} dx = \frac{(bx^4 + a)^{\frac{3}{4}} x}{abx^4 + a^2}$$

input `integrate(1/(b*x^4+a)^(5/4),x, algorithm="fricas")`

output `(b*x^4 + a)^(3/4)*x/(a*b*x^4 + a^2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. 2(12) = 24.

Time = 0.40 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.81

$$\int \frac{1}{(a + bx^4)^{5/4}} dx = \frac{x\Gamma(\frac{1}{4})}{4a^{\frac{5}{4}}\sqrt[4]{1 + \frac{bx^4}{a}}\Gamma(\frac{5}{4})}$$

input `integrate(1/(b*x**4+a)**(5/4),x)`

output `x*gamma(1/4)/(4*a**(5/4)*(1 + b*x**4/a)**(1/4)*gamma(5/4))`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1}{(a + bx^4)^{5/4}} dx = \frac{x}{(bx^4 + a)^{1/4} a}$$

input `integrate(1/(b*x^4+a)^(5/4),x, algorithm="maxima")`output `x/((b*x^4 + a)^(1/4)*a)`**Giac [F]**

$$\int \frac{1}{(a + bx^4)^{5/4}} dx = \int \frac{1}{(bx^4 + a)^{5/4}} dx$$

input `integrate(1/(b*x^4+a)^(5/4),x, algorithm="giac")`output `integrate((b*x^4 + a)^(-5/4), x)`**Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1}{(a + bx^4)^{5/4}} dx = \frac{x}{a (bx^4 + a)^{1/4}}$$

input `int(1/(a + b*x^4)^(5/4),x)`output `x/(a*(a + b*x^4)^(1/4))`

Reduce [F]

$$\int \frac{1}{(a + bx^4)^{5/4}} dx = \int \frac{1}{(bx^4 + a)^{1/4} a + (bx^4 + a)^{1/4} bx^4} dx$$

input `int(1/(b*x^4+a)^(5/4),x)`

output `int(1/((a + b*x**4)**(1/4)*a + (a + b*x**4)**(1/4)*b*x**4),x)`

3.35 $\int \frac{1}{(a+bx^4)^{9/4}} dx$

Optimal result	252
Mathematica [A] (verified)	252
Rubi [A] (verified)	253
Maple [A] (verified)	254
Fricas [A] (verification not implemented)	254
Sympy [B] (verification not implemented)	255
Maxima [A] (verification not implemented)	255
Giac [F]	256
Mupad [B] (verification not implemented)	256
Reduce [F]	256

Optimal result

Integrand size = 11, antiderivative size = 39

$$\int \frac{1}{(a + bx^4)^{9/4}} dx = \frac{x}{5a(a + bx^4)^{5/4}} + \frac{4x}{5a^2 \sqrt[4]{a + bx^4}}$$

output `1/5*x/a/(b*x^4+a)^(5/4)+4/5*x/a^2/(b*x^4+a)^(1/4)`

Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.74

$$\int \frac{1}{(a + bx^4)^{9/4}} dx = \frac{5ax + 4bx^5}{5a^2(a + bx^4)^{5/4}}$$

input `Integrate[(a + b*x^4)^(-9/4),x]`

output `(5*a*x + 4*b*x^5)/(5*a^2*(a + b*x^4)^(5/4))`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {749, 746}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + bx^4)^{9/4}} dx$$

$$\downarrow 749$$

$$\frac{4 \int \frac{1}{(bx^4+a)^{5/4}} dx}{5a} + \frac{x}{5a(a + bx^4)^{5/4}}$$

$$\downarrow 746$$

$$\frac{4x}{5a^2 \sqrt[4]{a + bx^4}} + \frac{x}{5a(a + bx^4)^{5/4}}$$

input `Int[(a + b*x^4)^(-9/4), x]`

output `x/(5*a*(a + b*x^4)^(5/4)) + (4*x)/(5*a^2*(a + b*x^4)^(1/4))`

Defintions of rubi rules used

rule 746 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]`

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.67

method	result	size
gospers	$\frac{x(4bx^4+5a)}{5(bx^4+a)^{\frac{5}{4}}a^2}$	26
trager	$\frac{x(4bx^4+5a)}{5(bx^4+a)^{\frac{5}{4}}a^2}$	26
pseudoelliptic	$\frac{x(4bx^4+5a)}{5(bx^4+a)^{\frac{5}{4}}a^2}$	26
orering	$\frac{x(4bx^4+5a)}{5(bx^4+a)^{\frac{5}{4}}a^2}$	26

input `int(1/(b*x^4+a)^(9/4),x,method=_RETURNVERBOSE)`

output `1/5*x*(4*b*x^4+5*a)/(b*x^4+a)^(5/4)/a^2`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.21

$$\int \frac{1}{(a+bx^4)^{9/4}} dx = \frac{(4bx^5+5ax)(bx^4+a)^{3/4}}{5(a^2b^2x^8+2a^3bx^4+a^4)}$$

input `integrate(1/(b*x^4+a)^(9/4),x, algorithm="fricas")`

output `1/5*(4*b*x^5+5*a*x)*(b*x^4+a)^(3/4)/(a^2*b^2*x^8+2*a^3*b*x^4+a^4)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 126 vs. $2(32) = 64$.

Time = 0.67 (sec) , antiderivative size = 126, normalized size of antiderivative = 3.23

$$\int \frac{1}{(a + bx^4)^{9/4}} dx = \frac{5ax\Gamma\left(\frac{1}{4}\right)}{16a^{13/4}\sqrt[4]{1 + \frac{bx^4}{a}}\Gamma\left(\frac{9}{4}\right) + 16a^{9/4}bx^4\sqrt[4]{1 + \frac{bx^4}{a}}\Gamma\left(\frac{9}{4}\right)} + \frac{4bx^5\Gamma\left(\frac{1}{4}\right)}{16a^{13/4}\sqrt[4]{1 + \frac{bx^4}{a}}\Gamma\left(\frac{9}{4}\right) + 16a^{9/4}bx^4\sqrt[4]{1 + \frac{bx^4}{a}}\Gamma\left(\frac{9}{4}\right)}$$

input `integrate(1/(b*x**4+a)**(9/4),x)`

output `5*a*x*gamma(1/4)/(16*a**(13/4)*(1 + b*x**4/a)**(1/4)*gamma(9/4) + 16*a**(9/4)*b*x**4*(1 + b*x**4/a)**(1/4)*gamma(9/4)) + 4*b*x**5*gamma(1/4)/(16*a**(13/4)*(1 + b*x**4/a)**(1/4)*gamma(9/4) + 16*a**(9/4)*b*x**4*(1 + b*x**4/a)**(1/4)*gamma(9/4))`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.79

$$\int \frac{1}{(a + bx^4)^{9/4}} dx = -\frac{\left(b - \frac{5(bx^4+a)}{x^4}\right)x^5}{5(bx^4 + a)^{5/4}a^2}$$

input `integrate(1/(b*x^4+a)^(9/4),x, algorithm="maxima")`

output `-1/5*(b - 5*(b*x^4 + a)/x^4)*x^5/((b*x^4 + a)^(5/4)*a^2)`

Giac [F]

$$\int \frac{1}{(a + bx^4)^{9/4}} dx = \int \frac{1}{(bx^4 + a)^{9/4}} dx$$

input `integrate(1/(b*x^4+a)^(9/4),x, algorithm="giac")`

output `integrate((b*x^4 + a)^(-9/4), x)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.72

$$\int \frac{1}{(a + bx^4)^{9/4}} dx = \frac{4x(bx^4 + a) + ax}{5a^2(bx^4 + a)^{5/4}}$$

input `int(1/(a + b*x^4)^(9/4),x)`

output `(4*x*(a + b*x^4) + a*x)/(5*a^2*(a + b*x^4)^(5/4))`

Reduce [F]

$$\int \frac{1}{(a + bx^4)^{9/4}} dx = \int \frac{1}{(bx^4 + a)^{1/4} a^2 + 2(bx^4 + a)^{1/4} abx^4 + (bx^4 + a)^{1/4} b^2x^8} dx$$

input `int(1/(b*x^4+a)^(9/4),x)`

output `int(1/((a + b*x**4)**(1/4)*a**2 + 2*(a + b*x**4)**(1/4)*a*b*x**4 + (a + b*x**4)**(1/4)*b**2*x**8),x)`

3.36 $\int \frac{1}{(a+bx^4)^{13/4}} dx$

Optimal result	257
Mathematica [A] (verified)	257
Rubi [A] (verified)	258
Maple [A] (verified)	259
Fricas [A] (verification not implemented)	259
Sympy [B] (verification not implemented)	260
Maxima [A] (verification not implemented)	261
Giac [F]	261
Mupad [B] (verification not implemented)	262
Reduce [F]	262

Optimal result

Integrand size = 11, antiderivative size = 58

$$\int \frac{1}{(a+bx^4)^{13/4}} dx = \frac{x}{9a(a+bx^4)^{9/4}} + \frac{8x}{45a^2(a+bx^4)^{5/4}} + \frac{32x}{45a^3\sqrt[4]{a+bx^4}}$$

output $1/9*x/a/(b*x^4+a)^{(9/4)}+8/45*x/a^2/(b*x^4+a)^{(5/4)}+32/45*x/a^3/(b*x^4+a)^{(1/4)}$

Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.69

$$\int \frac{1}{(a+bx^4)^{13/4}} dx = \frac{45a^2x + 72abx^5 + 32b^2x^9}{45a^3(a+bx^4)^{9/4}}$$

input `Integrate[(a + b*x^4)^(-13/4), x]`

output $(45*a^2*x + 72*a*b*x^5 + 32*b^2*x^9)/(45*a^3*(a + b*x^4)^{(9/4)})$

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.14, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {749, 749, 746}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + bx^4)^{13/4}} dx$$

$$\downarrow 749$$

$$\frac{8 \int \frac{1}{(bx^4+a)^{9/4}} dx}{9a} + \frac{x}{9a (a + bx^4)^{9/4}}$$

$$\downarrow 749$$

$$\frac{8 \left(\frac{4 \int \frac{1}{(bx^4+a)^{5/4}} dx}{5a} + \frac{x}{5a(a+bx^4)^{5/4}} \right)}{9a} + \frac{x}{9a (a + bx^4)^{9/4}}$$

$$\downarrow 746$$

$$\frac{8 \left(\frac{4x}{5a^2 \sqrt[4]{a + bx^4}} + \frac{x}{5a(a+bx^4)^{5/4}} \right)}{9a} + \frac{x}{9a (a + bx^4)^{9/4}}$$

input `Int[(a + b*x^4)^(-13/4),x]`

output `x/(9*a*(a + b*x^4)^(9/4)) + (8*(x/(5*a*(a + b*x^4)^(5/4)) + (4*x)/(5*a^2*(a + b*x^4)^(1/4))))/(9*a)`

Defintions of rubi rules used

rule 746 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[x*((a + b*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]`

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.64

method	result	size
gospers	$\frac{x(32b^2x^8+72abx^4+45a^2)}{45(bx^4+a)^{\frac{9}{4}}a^3}$	37
trager	$\frac{x(32b^2x^8+72abx^4+45a^2)}{45(bx^4+a)^{\frac{9}{4}}a^3}$	37
pseudoelliptic	$\frac{x(32b^2x^8+72abx^4+45a^2)}{45(bx^4+a)^{\frac{9}{4}}a^3}$	37
orering	$\frac{x(32b^2x^8+72abx^4+45a^2)}{45(bx^4+a)^{\frac{9}{4}}a^3}$	37

input `int(1/(b*x^4+a)^(13/4), x, method=_RETURNVERBOSE)`

output `1/45*x*(32*b^2*x^8+72*a*b*x^4+45*a^2)/(b*x^4+a)^(9/4)/a^3`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.19

$$\int \frac{1}{(a + bx^4)^{13/4}} dx = \frac{(32b^2x^9 + 72abx^5 + 45a^2x)(bx^4 + a)^{\frac{3}{4}}}{45(a^3b^3x^{12} + 3a^4b^2x^8 + 3a^5bx^4 + a^6)}$$

input `integrate(1/(b*x^4+a)^(13/4),x, algorithm="fricas")`

output $\frac{1}{45} \cdot (32 \cdot b^2 \cdot x^9 + 72 \cdot a \cdot b \cdot x^5 + 45 \cdot a^2 \cdot x) \cdot (b \cdot x^4 + a)^{3/4} / (a^3 \cdot b^3 \cdot x^{12} + 3 \cdot a^4 \cdot b^2 \cdot x^8 + 3 \cdot a^5 \cdot b \cdot x^4 + a^6)$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 515 vs. $2(51) = 102$.

Time = 1.45 (sec) , antiderivative size = 515, normalized size of antiderivative = 8.88

$$\int \frac{1}{(a + bx^4)^{13/4}} dx = \frac{45a^5 x \Gamma\left(\frac{1}{4}\right)}{64a^{\frac{33}{4}} \sqrt[4]{1 + \frac{bx^4}{a}} \Gamma\left(\frac{13}{4}\right) + 192a^{\frac{29}{4}} bx^4 \sqrt[4]{1 + \frac{bx^4}{a}} \Gamma\left(\frac{13}{4}\right) + 192a^{\frac{25}{4}} b^2 x^8 \sqrt[4]{1 + \frac{bx^4}{a}} \Gamma\left(\frac{13}{4}\right) + 117a^4 bx^5 \Gamma\left(\frac{1}{4}\right)} + \frac{64a^{\frac{33}{4}} \sqrt[4]{1 + \frac{bx^4}{a}} \Gamma\left(\frac{13}{4}\right) + 192a^{\frac{29}{4}} bx^4 \sqrt[4]{1 + \frac{bx^4}{a}} \Gamma\left(\frac{13}{4}\right) + 192a^{\frac{25}{4}} b^2 x^8 \sqrt[4]{1 + \frac{bx^4}{a}} \Gamma\left(\frac{13}{4}\right) + 64a^{\frac{21}{4}} b^3 x^{12} \sqrt[4]{1 + \frac{bx^4}{a}} \Gamma\left(\frac{13}{4}\right)}{104a^3 b^2 x^9 \Gamma\left(\frac{1}{4}\right)} + \frac{64a^{\frac{33}{4}} \sqrt[4]{1 + \frac{bx^4}{a}} \Gamma\left(\frac{13}{4}\right) + 192a^{\frac{29}{4}} bx^4 \sqrt[4]{1 + \frac{bx^4}{a}} \Gamma\left(\frac{13}{4}\right) + 192a^{\frac{25}{4}} b^2 x^8 \sqrt[4]{1 + \frac{bx^4}{a}} \Gamma\left(\frac{13}{4}\right) + 64a^{\frac{21}{4}} b^3 x^{12} \sqrt[4]{1 + \frac{bx^4}{a}} \Gamma\left(\frac{13}{4}\right)}{32a^2 b^3 x^{13} \Gamma\left(\frac{1}{4}\right)} + \frac{64a^{\frac{33}{4}} \sqrt[4]{1 + \frac{bx^4}{a}} \Gamma\left(\frac{13}{4}\right) + 192a^{\frac{29}{4}} bx^4 \sqrt[4]{1 + \frac{bx^4}{a}} \Gamma\left(\frac{13}{4}\right) + 192a^{\frac{25}{4}} b^2 x^8 \sqrt[4]{1 + \frac{bx^4}{a}} \Gamma\left(\frac{13}{4}\right) + 64a^{\frac{21}{4}} b^3 x^{12} \sqrt[4]{1 + \frac{bx^4}{a}} \Gamma\left(\frac{13}{4}\right)}{\dots}$$

input `integrate(1/(b*x**4+a)**(13/4),x)`

output

```
45*a**5*x*gamma(1/4)/(64*a**(33/4)*(1 + b*x**4/a)**(1/4)*gamma(13/4) + 192
*a**(29/4)*b*x**4*(1 + b*x**4/a)**(1/4)*gamma(13/4) + 192*a**(25/4)*b**2*x
**8*(1 + b*x**4/a)**(1/4)*gamma(13/4) + 64*a**(21/4)*b**3*x**12*(1 + b*x**
4/a)**(1/4)*gamma(13/4)) + 117*a**4*b*x**5*gamma(1/4)/(64*a**(33/4)*(1 + b
*x**4/a)**(1/4)*gamma(13/4) + 192*a**(29/4)*b*x**4*(1 + b*x**4/a)**(1/4)*g
amma(13/4) + 192*a**(25/4)*b**2*x**8*(1 + b*x**4/a)**(1/4)*gamma(13/4) + 6
4*a**(21/4)*b**3*x**12*(1 + b*x**4/a)**(1/4)*gamma(13/4)) + 104*a**3*b**2*
x**9*gamma(1/4)/(64*a**(33/4)*(1 + b*x**4/a)**(1/4)*gamma(13/4) + 192*a**(
29/4)*b*x**4*(1 + b*x**4/a)**(1/4)*gamma(13/4) + 192*a**(25/4)*b**2*x**8*(
1 + b*x**4/a)**(1/4)*gamma(13/4) + 64*a**(21/4)*b**3*x**12*(1 + b*x**4/a)*
*(1/4)*gamma(13/4)) + 32*a**2*b**3*x**13*gamma(1/4)/(64*a**(33/4)*(1 + b*x
**4/a)**(1/4)*gamma(13/4) + 192*a**(29/4)*b*x**4*(1 + b*x**4/a)**(1/4)*gam
ma(13/4) + 192*a**(25/4)*b**2*x**8*(1 + b*x**4/a)**(1/4)*gamma(13/4) + 64*
a**(21/4)*b**3*x**12*(1 + b*x**4/a)**(1/4)*gamma(13/4))
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.86

$$\int \frac{1}{(a + bx^4)^{13/4}} dx = \frac{\left(5b^2 - \frac{18(bx^4+a)b}{x^4} + \frac{45(bx^4+a)^2}{x^8}\right)x^9}{45(bx^4 + a)^{9/4}a^3}$$

input

```
integrate(1/(b*x^4+a)^(13/4),x, algorithm="maxima")
```

output

```
1/45*(5*b^2 - 18*(b*x^4 + a)*b/x^4 + 45*(b*x^4 + a)^2/x^8)*x^9/((b*x^4 + a
)^(9/4)*a^3)
```

Giac [F]

$$\int \frac{1}{(a + bx^4)^{13/4}} dx = \int \frac{1}{(bx^4 + a)^{\frac{13}{4}}} dx$$

input

```
integrate(1/(b*x^4+a)^(13/4),x, algorithm="giac")
```

output `integrate((b*x^4 + a)^(-13/4), x)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.76

$$\int \frac{1}{(a + bx^4)^{13/4}} dx = \frac{32x(bx^4 + a)^2 + 5a^2x + 8ax(bx^4 + a)}{45a^3(bx^4 + a)^{9/4}}$$

input `int(1/(a + b*x^4)^(13/4),x)`

output `(32*x*(a + b*x^4)^2 + 5*a^2*x + 8*a*x*(a + b*x^4))/(45*a^3*(a + b*x^4)^(9/4))`

Reduce [F]

$$\int \frac{1}{(a + bx^4)^{13/4}} dx = \int \frac{1}{(bx^4 + a)^{\frac{1}{4}} a^3 + 3(bx^4 + a)^{\frac{1}{4}} a^2 bx^4 + 3(bx^4 + a)^{\frac{1}{4}} a b^2 x^8 + (bx^4 + a)^{\frac{1}{4}} b^3 x^{12}} dx$$

input `int(1/(b*x^4+a)^(13/4),x)`

output `int(1/((a + b*x**4)**(1/4)*a**3 + 3*(a + b*x**4)**(1/4)*a**2*b*x**4 + 3*(a + b*x**4)**(1/4)*a*b**2*x**8 + (a + b*x**4)**(1/4)*b**3*x**12),x)`

$$3.37 \quad \int \frac{1}{(a+bx^4)^{17/4}} dx$$

Optimal result	263
Mathematica [A] (verified)	263
Rubi [A] (verified)	264
Maple [A] (verified)	265
Fricas [A] (verification not implemented)	266
Sympy [B] (verification not implemented)	266
Maxima [A] (verification not implemented)	267
Giac [F]	268
Mupad [B] (verification not implemented)	268
Reduce [F]	268

Optimal result

Integrand size = 11, antiderivative size = 77

$$\int \frac{1}{(a+bx^4)^{17/4}} dx = \frac{x}{13a(a+bx^4)^{13/4}} + \frac{4x}{39a^2(a+bx^4)^{9/4}} + \frac{32x}{195a^3(a+bx^4)^{5/4}} + \frac{128x}{195a^4\sqrt[4]{a+bx^4}}$$

output

```
1/13*x/a/(b*x^4+a)^(13/4)+4/39*x/a^2/(b*x^4+a)^(9/4)+32/195*x/a^3/(b*x^4+a)^(5/4)+128/195*x/a^4/(b*x^4+a)^(1/4)
```

Mathematica [A] (verified)

Time = 0.70 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.66

$$\int \frac{1}{(a+bx^4)^{17/4}} dx = \frac{195a^3x + 468a^2bx^5 + 416ab^2x^9 + 128b^3x^{13}}{195a^4(a+bx^4)^{13/4}}$$

input

```
Integrate[(a + b*x^4)^(-17/4), x]
```


output

$$(195*a^3*x + 468*a^2*b*x^5 + 416*a*b^2*x^9 + 128*b^3*x^{13})/(195*a^4*(a + b*x^4)^{(13/4)})$$
Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.21, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {749, 749, 749, 746}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + bx^4)^{17/4}} dx$$

$$\downarrow 749$$

$$\frac{12 \int \frac{1}{(bx^4+a)^{13/4}} dx}{13a} + \frac{x}{13a(a+bx^4)^{13/4}}$$

$$\downarrow 749$$

$$\frac{12 \left(\frac{8 \int \frac{1}{(bx^4+a)^{9/4}} dx}{9a} + \frac{x}{9a(a+bx^4)^{9/4}} \right)}{13a} + \frac{x}{13a(a+bx^4)^{13/4}}$$

$$\downarrow 749$$

$$\frac{12 \left(\frac{8 \left(\frac{4 \int \frac{1}{(bx^4+a)^{5/4}} dx}{5a} + \frac{x}{5a(a+bx^4)^{5/4}} \right)}{9a} + \frac{x}{9a(a+bx^4)^{9/4}} \right)}{13a} + \frac{x}{13a(a+bx^4)^{13/4}}$$

$$\downarrow 746$$

$$\frac{12 \left(\frac{8 \left(\frac{4x}{5a^2 \sqrt[4]{a+bx^4}} + \frac{x}{5a(a+bx^4)^{5/4}} \right)}{9a} + \frac{x}{9a(a+bx^4)^{9/4}} \right)}{13a} + \frac{x}{13a(a+bx^4)^{13/4}}$$

input `Int[(a + b*x^4)^(-17/4),x]`

output
$$\frac{x/(13*a*(a + b*x^4)^{(13/4)}) + (12*(x/(9*a*(a + b*x^4)^{(9/4)}) + (8*(x/(5*a*(a + b*x^4)^{(5/4)}) + (4*x)/(5*a^2*(a + b*x^4)^{(1/4)))/(9*a)))/(13*a)}{1}$$

Defintions of rubi rules used

rule 746 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]`

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

Maple [A] (verified)

Time = 0.54 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.62

method	result	size
gospers	$\frac{x(128b^3x^{12}+416ab^2x^8+468a^2bx^4+195a^3)}{195(bx^4+a)^{\frac{13}{4}}a^4}$	48
trager	$\frac{x(128b^3x^{12}+416ab^2x^8+468a^2bx^4+195a^3)}{195(bx^4+a)^{\frac{13}{4}}a^4}$	48
pseudoelliptic	$\frac{x(128b^3x^{12}+416ab^2x^8+468a^2bx^4+195a^3)}{195(bx^4+a)^{\frac{13}{4}}a^4}$	48
orering	$\frac{x(128b^3x^{12}+416ab^2x^8+468a^2bx^4+195a^3)}{195(bx^4+a)^{\frac{13}{4}}a^4}$	48

input `int(1/(b*x^4+a)^(17/4),x,method=_RETURNVERBOSE)`

output
$$\frac{1/195*x*(128*b^3*x^{12}+416*a*b^2*x^8+468*a^2*b*x^4+195*a^3)/(b*x^4+a)^{(13/4)}}{a^4}$$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.18

$$\int \frac{1}{(a + bx^4)^{17/4}} dx = \frac{(128 b^3 x^{13} + 416 ab^2 x^9 + 468 a^2 b x^5 + 195 a^3 x)(bx^4 + a)^{3/4}}{195 (a^4 b^4 x^{16} + 4 a^5 b^3 x^{12} + 6 a^6 b^2 x^8 + 4 a^7 b x^4 + a^8)}$$

input `integrate(1/(b*x^4+a)^(17/4),x, algorithm="fricas")`

output `1/195*(128*b^3*x^13 + 416*a*b^2*x^9 + 468*a^2*b*x^5 + 195*a^3*x)*(b*x^4 + a)^(3/4)/(a^4*b^4*x^16 + 4*a^5*b^3*x^12 + 6*a^6*b^2*x^8 + 4*a^7*b*x^4 + a^8)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1550 vs. 2(70) = 140.

Time = 2.77 (sec) , antiderivative size = 1550, normalized size of antiderivative = 20.13

$$\int \frac{1}{(a + bx^4)^{17/4}} dx = \text{Too large to display}$$

input `integrate(1/(b*x**4+a)**(17/4),x)`

output

```

585*a**14*x*gamma(1/4)/(256*a**(73/4)*(1 + b*x**4/a)**(1/4)*gamma(17/4) +
1536*a**(69/4)*b*x**4*(1 + b*x**4/a)**(1/4)*gamma(17/4) + 3840*a**(65/4)*b
**2*x**8*(1 + b*x**4/a)**(1/4)*gamma(17/4) + 5120*a**(61/4)*b**3*x**12*(1
+ b*x**4/a)**(1/4)*gamma(17/4) + 3840*a**(57/4)*b**4*x**16*(1 + b*x**4/a)*
*(1/4)*gamma(17/4) + 1536*a**(53/4)*b**5*x**20*(1 + b*x**4/a)**(1/4)*gamma
(17/4) + 256*a**(49/4)*b**6*x**24*(1 + b*x**4/a)**(1/4)*gamma(17/4)) + 315
9*a**13*b*x**5*gamma(1/4)/(256*a**(73/4)*(1 + b*x**4/a)**(1/4)*gamma(17/4)
+ 1536*a**(69/4)*b*x**4*(1 + b*x**4/a)**(1/4)*gamma(17/4) + 3840*a**(65/4)
)*b**2*x**8*(1 + b*x**4/a)**(1/4)*gamma(17/4) + 5120*a**(61/4)*b**3*x**12*
(1 + b*x**4/a)**(1/4)*gamma(17/4) + 3840*a**(57/4)*b**4*x**16*(1 + b*x**4/
a)**(1/4)*gamma(17/4) + 1536*a**(53/4)*b**5*x**20*(1 + b*x**4/a)**(1/4)*ga
mma(17/4) + 256*a**(49/4)*b**6*x**24*(1 + b*x**4/a)**(1/4)*gamma(17/4)) +
7215*a**12*b**2*x**9*gamma(1/4)/(256*a**(73/4)*(1 + b*x**4/a)**(1/4)*gamma
(17/4) + 1536*a**(69/4)*b*x**4*(1 + b*x**4/a)**(1/4)*gamma(17/4) + 3840*a*
*(65/4)*b**2*x**8*(1 + b*x**4/a)**(1/4)*gamma(17/4) + 5120*a**(61/4)*b**3*
x**12*(1 + b*x**4/a)**(1/4)*gamma(17/4) + 3840*a**(57/4)*b**4*x**16*(1 + b
*x**4/a)**(1/4)*gamma(17/4) + 1536*a**(53/4)*b**5*x**20*(1 + b*x**4/a)**(1
/4)*gamma(17/4) + 256*a**(49/4)*b**6*x**24*(1 + b*x**4/a)**(1/4)*gamma(17/
4)) + 8925*a**11*b**3*x**13*gamma(1/4)/(256*a**(73/4)*(1 + b*x**4/a)**(1/4)
)*gamma(17/4) + 1536*a**(69/4)*b*x**4*(1 + b*x**4/a)**(1/4)*gamma(17/4)...

```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.87

$$\int \frac{1}{(a + bx^4)^{17/4}} dx = -\frac{\left(15b^3 - \frac{65(bx^4+a)b^2}{x^4} + \frac{117(bx^4+a)^2b}{x^8} - \frac{195(bx^4+a)^3}{x^{12}}\right)x^{13}}{195(bx^4+a)^{\frac{13}{4}}a^4}$$

input

```
integrate(1/(b*x^4+a)^(17/4),x, algorithm="maxima")
```

output

```
-1/195*(15*b^3 - 65*(b*x^4 + a)*b^2/x^4 + 117*(b*x^4 + a)^2*b/x^8 - 195*(b
*x^4 + a)^3/x^12)*x^13/((b*x^4 + a)^(13/4)*a^4)
```

Giac [F]

$$\int \frac{1}{(a + bx^4)^{17/4}} dx = \int \frac{1}{(bx^4 + a)^{17/4}} dx$$

input `integrate(1/(b*x^4+a)^(17/4),x, algorithm="giac")`

output `integrate((b*x^4 + a)^(-17/4), x)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.79

$$\int \frac{1}{(a + bx^4)^{17/4}} dx = \frac{128x}{195a^4(bx^4 + a)^{1/4}} + \frac{32x}{195a^3(bx^4 + a)^{5/4}} + \frac{4x}{39a^2(bx^4 + a)^{9/4}} + \frac{x}{13a(bx^4 + a)^{13/4}}$$

input `int(1/(a + b*x^4)^(17/4),x)`

output `(128*x)/(195*a^4*(a + b*x^4)^(1/4)) + (32*x)/(195*a^3*(a + b*x^4)^(5/4)) + (4*x)/(39*a^2*(a + b*x^4)^(9/4)) + x/(13*a*(a + b*x^4)^(13/4))`

Reduce [F]

$$\int \frac{1}{(a + bx^4)^{17/4}} dx = \int \frac{1}{(bx^4 + a)^{1/4} a^4 + 4(bx^4 + a)^{1/4} a^3 bx^4 + 6(bx^4 + a)^{1/4} a^2 b^2 x^8 + 4(bx^4 + a)^{1/4} a b^3 x^{12}}$$

input `int(1/(b*x^4+a)^(17/4),x)`

output `int(1/((a + b*x**4)**(1/4)*a**4 + 4*(a + b*x**4)**(1/4)*a**3*b*x**4 + 6*(a + b*x**4)**(1/4)*a**2*b**2*x**8 + 4*(a + b*x**4)**(1/4)*a*b**3*x**12 + (a + b*x**4)**(1/4)*b**4*x**16),x)`

3.38 $\int (a + bx^4)^{9/4} dx$

Optimal result	269
Mathematica [C] (verified)	269
Rubi [A] (verified)	270
Maple [F]	272
Fricas [F]	273
Sympy [C] (verification not implemented)	273
Maxima [F]	273
Giac [F]	274
Mupad [B] (verification not implemented)	274
Reduce [F]	274

Optimal result

Integrand size = 11, antiderivative size = 116

$$\int (a + bx^4)^{9/4} dx = \frac{3}{8}a^2x^4\sqrt{a + bx^4} + \frac{3}{20}ax(a + bx^4)^{5/4} + \frac{1}{10}x(a + bx^4)^{9/4} - \frac{3a^{5/2}\sqrt{b}\left(1 + \frac{a}{bx^4}\right)^{3/4}x^3 \operatorname{EllipticF}\left(\frac{1}{2}\cot^{-1}\left(\frac{\sqrt{bx^2}}{\sqrt{a}}\right), 2\right)}{8(a + bx^4)^{3/4}}$$

output $\frac{3}{8}a^2x^4\sqrt{a + bx^4} + \frac{3}{20}ax(a + bx^4)^{5/4} + \frac{1}{10}x(a + bx^4)^{9/4} - \frac{3a^{5/2}\sqrt{b}\left(1 + \frac{a}{bx^4}\right)^{3/4}x^3 \operatorname{InverseJacobiAM}\left(\frac{1}{2}\operatorname{arccot}\left(\frac{\sqrt{bx^2}}{\sqrt{a}}\right), 2\right)}{8(a + bx^4)^{3/4}}$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 8.58 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.42

$$\int (a + bx^4)^{9/4} dx = \frac{a^2x^4\sqrt{a + bx^4} \operatorname{Hypergeometric2F1}\left(-\frac{9}{4}, \frac{1}{4}, \frac{5}{4}, -\frac{bx^4}{a}\right)}{\sqrt[4]{1 + \frac{bx^4}{a}}}$$

input `Integrate[(a + b*x^4)^(9/4), x]`

output $(a^2 x (a + b x^4)^{1/4} \text{Hypergeometric2F1}[-9/4, 1/4, 5/4, -((b x^4)/a)]) / (1 + (b x^4)/a)^{1/4}$

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.07, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {748, 748, 748, 768, 858, 807, 229}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + bx^4)^{9/4} dx \\
 & \quad \downarrow 748 \\
 & \frac{9}{10}a \int (bx^4 + a)^{5/4} dx + \frac{1}{10}x(a + bx^4)^{9/4} \\
 & \quad \downarrow 748 \\
 & \frac{9}{10}a \left(\frac{5}{6}a \int \sqrt[4]{bx^4 + a} dx + \frac{1}{6}x(a + bx^4)^{5/4} \right) + \frac{1}{10}x(a + bx^4)^{9/4} \\
 & \quad \downarrow 748 \\
 & \frac{9}{10}a \left(\frac{5}{6}a \left(\frac{1}{2}a \int \frac{1}{(bx^4 + a)^{3/4}} dx + \frac{1}{2}x^4 \sqrt[4]{a + bx^4} \right) + \frac{1}{6}x(a + bx^4)^{5/4} \right) + \frac{1}{10}x(a + bx^4)^{9/4} \\
 & \quad \downarrow 768 \\
 & \frac{9}{10}a \left(\frac{5}{6}a \left(\frac{ax^3 \left(\frac{a}{bx^4} + 1 \right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^4} + 1 \right)^{3/4} x^3} dx}{2(a + bx^4)^{3/4}} + \frac{1}{2}x^4 \sqrt[4]{a + bx^4} \right) + \frac{1}{6}x(a + bx^4)^{5/4} \right) + \\
 & \quad \frac{1}{10}x(a + bx^4)^{9/4} \\
 & \quad \downarrow 858
 \end{aligned}$$

$$\frac{9}{10}a \left(\frac{5}{6}a \left(\frac{1}{2}x^4 \sqrt{a+bx^4} - \frac{ax^3 \left(\frac{a}{bx^4} + 1\right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^4} + 1\right)^{3/4} x} d\frac{1}{x}}{2(a+bx^4)^{3/4}} \right) + \frac{1}{6}x(a+bx^4)^{5/4} \right) + \frac{1}{10}x(a+bx^4)^{9/4}$$

↓ 807

$$\frac{9}{10}a \left(\frac{5}{6}a \left(\frac{1}{2}x^4 \sqrt{a+bx^4} - \frac{ax^3 \left(\frac{a}{bx^2} + 1\right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^2} + 1\right)^{3/4} d\frac{1}{x^2}}}{4(a+bx^4)^{3/4}} \right) + \frac{1}{6}x(a+bx^4)^{5/4} \right) + \frac{1}{10}x(a+bx^4)^{9/4}$$

↓ 229

$$\frac{9}{10}a \left(\frac{5}{6}a \left(\frac{1}{2}x^4 \sqrt{a+bx^4} - \frac{\sqrt{a}\sqrt{bx^3} \left(\frac{a}{bx^4} + 1\right)^{3/4} \text{EllipticF}\left(\frac{1}{2} \arctan\left(\frac{\sqrt{a}}{\sqrt{bx^2}}\right), 2\right)}{2(a+bx^4)^{3/4}} \right) + \frac{1}{6}x(a+bx^4)^{5/4} \right) + \frac{1}{10}x(a+bx^4)^{9/4}$$

input `Int[(a + b*x^4)^(9/4), x]`

output `(x*(a + b*x^4)^(9/4))/10 + (9*a*((x*(a + b*x^4)^(5/4))/6 + (5*a*((x*(a + b*x^4)^(1/4))/2 - (Sqrt[a]*Sqrt[b]*(1 + a/(b*x^4))^(3/4)*x^3*EllipticF[ArcTan[Sqrt[a]/(Sqrt[b]*x^2)]/2, 2)]/(2*(a + b*x^4)^(3/4)))/6))/10`

Definitions of rubi rules used

rule 229 $\text{Int}[(a_ + (b_.)*(x_)^2)^{-3/4}, x_Symbol] \rightarrow \text{Simp}[(2/(a^{3/4})*\text{Rt}[b/a, 2]) * \text{EllipticF}[(1/2)*\text{ArcTan}[\text{Rt}[b/a, 2]*x], 2], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{GtQ}[a, 0] \ \&\& \ \text{PosQ}[b/a]$

rule 748 $\text{Int}[(a_ + (b_.)*(x_)^{n_})^{p_}, x_Symbol] \rightarrow \text{Simp}[x*((a + b*x^n)^p/(n*p + 1)), x] + \text{Simp}[a*n*(p/(n*p + 1)) \ \text{Int}[(a + b*x^n)^{p-1}, x], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ (\text{IntegerQ}[2*p] \ || \ \text{LtQ}[\text{Denominator}[p + 1/n], \text{Denominator}[p]])$

rule 768 $\text{Int}[(a_ + (b_.)*(x_)^4)^{-3/4}, x_Symbol] \rightarrow \text{Simp}[x^3*((1 + a/(b*x^4))^{3/4})/(a + b*x^4)^{3/4}) \ \text{Int}[1/(x^3*(1 + a/(b*x^4))^{3/4}), x], x] /; \text{FreeQ}[\{a, b\}, x]$

rule 807 $\text{Int}[(x_)^{m_}*(a_ + (b_.)*(x_)^{n_})^{p_}, x_Symbol] \rightarrow \text{With}[\{k = \text{GCD}[m + 1, n]\}, \text{Simp}[1/k \ \text{Subst}[\text{Int}[x^{(m+1)/k-1}*(a + b*x^{n/k})^p, x], x, x^k], x] /; k \neq 1] /; \text{FreeQ}[\{a, b, p\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IntegerQ}[m]$

rule 858 $\text{Int}[(x_)^{m_}*(a_ + (b_.)*(x_)^{n_})^{p_}, x_Symbol] \rightarrow -\text{Subst}[\text{Int}[(a + b/x^n)^p/x^{m+2}, x], x, 1/x] /; \text{FreeQ}[\{a, b, p\}, x] \ \&\& \ \text{ILtQ}[n, 0] \ \&\& \ \text{IntegerQ}[m]$

Maple [F]

$$\int (bx^4 + a)^{\frac{9}{4}} dx$$

input $\text{int}((b*x^4+a)^{9/4}, x)$

output $\text{int}((b*x^4+a)^{9/4}, x)$

Fricas [F]

$$\int (a + bx^4)^{9/4} dx = \int (bx^4 + a)^{9/4} dx$$

input `integrate((b*x^4+a)^(9/4),x, algorithm="fricas")`

output `integral((b^2*x^8 + 2*a*b*x^4 + a^2)*(b*x^4 + a)^(1/4), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.00 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.32

$$\int (a + bx^4)^{9/4} dx = \frac{a^{9/4} x \Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\begin{matrix} -\frac{9}{4}, \frac{1}{4} \\ \frac{5}{4} \end{matrix} \middle| \frac{bx^4 e^{i\pi}}{a}\right)}{4\Gamma\left(\frac{5}{4}\right)}$$

input `integrate((b*x**4+a)**(9/4),x)`

output `a**(9/4)*x*gamma(1/4)*hyper((-9/4, 1/4), (5/4,), b*x**4*exp_polar(I*pi)/a)/(4*gamma(5/4))`

Maxima [F]

$$\int (a + bx^4)^{9/4} dx = \int (bx^4 + a)^{9/4} dx$$

input `integrate((b*x^4+a)^(9/4),x, algorithm="maxima")`

output `integrate((b*x^4 + a)^(9/4), x)`

Giac [F]

$$\int (a + bx^4)^{9/4} dx = \int (bx^4 + a)^{9/4} dx$$

input `integrate((b*x^4+a)^(9/4),x, algorithm="giac")`

output `integrate((b*x^4 + a)^(9/4), x)`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.32

$$\int (a + bx^4)^{9/4} dx = \frac{x (bx^4 + a)^{9/4} {}_2F_1\left(-\frac{9}{4}, \frac{1}{4}; \frac{5}{4}; -\frac{bx^4}{a}\right)}{\left(\frac{bx^4}{a} + 1\right)^{9/4}}$$

input `int((a + b*x^4)^(9/4),x)`

output `(x*(a + b*x^4)^(9/4)*hypergeom([-9/4, 1/4], 5/4, -(b*x^4)/a))/((b*x^4)/a + 1)^(9/4)`

Reduce [F]

$$\begin{aligned} \int (a + bx^4)^{9/4} dx &= \frac{5(bx^4 + a)^{1/4} a^2 x}{8} + \frac{7(bx^4 + a)^{1/4} ab x^5}{20} \\ &+ \frac{(bx^4 + a)^{1/4} b^2 x^9}{10} + \frac{3 \left(\int \frac{1}{(bx^4 + a)^{3/4}} dx \right) a^3}{8} \end{aligned}$$

input `int((b*x^4+a)^(9/4),x)`

output

```
(25*(a + b*x**4)**(1/4)*a**2*x + 14*(a + b*x**4)**(1/4)*a*b*x**5 + 4*(a +
b*x**4)**(1/4)*b**2*x**9 + 15*int((a + b*x**4)**(1/4)/(a + b*x**4),x)*a**3
)/40
```

3.39 $\int (a + bx^4)^{5/4} dx$

Optimal result	276
Mathematica [C] (verified)	276
Rubi [A] (verified)	277
Maple [F]	279
Fricas [F]	279
Sympy [C] (verification not implemented)	279
Maxima [F]	280
Giac [F]	280
Mupad [B] (verification not implemented)	280
Reduce [F]	281

Optimal result

Integrand size = 11, antiderivative size = 97

$$\int (a + bx^4)^{5/4} dx = \frac{5}{12}ax\sqrt[4]{a + bx^4} + \frac{1}{6}x(a + bx^4)^{5/4} - \frac{5a^{3/2}\sqrt{b}\left(1 + \frac{a}{bx^4}\right)^{3/4} x^3 \operatorname{EllipticF}\left(\frac{1}{2} \cot^{-1}\left(\frac{\sqrt{bx^2}}{\sqrt{a}}\right), 2\right)}{12(a + bx^4)^{3/4}}$$

output `5/12*a*x*(b*x^4+a)^(1/4)+1/6*x*(b*x^4+a)^(5/4)-5/12*a^(3/2)*b^(1/2)*(1+a/b/x^4)^(3/4)*x^3*InverseJacobiAM(1/2*arccot(b^(1/2)*x^2/a^(1/2)),2^(1/2))/(b*x^4+a)^(3/4)`

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.60 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.48

$$\int (a + bx^4)^{5/4} dx = \frac{ax\sqrt[4]{a + bx^4} \operatorname{Hypergeometric2F1}\left(-\frac{5}{4}, \frac{1}{4}, \frac{5}{4}, -\frac{bx^4}{a}\right)}{\sqrt[4]{1 + \frac{bx^4}{a}}}$$

input `Integrate[(a + b*x^4)^(5/4), x]`

output `(a*x*(a + b*x^4)^(1/4)*Hypergeometric2F1[-5/4, 1/4, 5/4, -((b*x^4)/a)]/(1 + (b*x^4)/a)^(1/4)`

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.05, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {748, 748, 768, 858, 807, 229}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + bx^4)^{5/4} dx \\
 & \quad \downarrow 748 \\
 & \frac{5}{6}a \int \sqrt[4]{bx^4 + a} dx + \frac{1}{6}x(a + bx^4)^{5/4} \\
 & \quad \downarrow 748 \\
 & \frac{5}{6}a \left(\frac{1}{2}a \int \frac{1}{(bx^4 + a)^{3/4}} dx + \frac{1}{2}x \sqrt[4]{a + bx^4} \right) + \frac{1}{6}x(a + bx^4)^{5/4} \\
 & \quad \downarrow 768 \\
 & \frac{5}{6}a \left(\frac{ax^3 \left(\frac{a}{bx^4} + 1\right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^4} + 1\right)^{3/4} x^3} dx}{2(a + bx^4)^{3/4}} + \frac{1}{2}x \sqrt[4]{a + bx^4} \right) + \frac{1}{6}x(a + bx^4)^{5/4} \\
 & \quad \downarrow 858 \\
 & \frac{5}{6}a \left(\frac{1}{2}x \sqrt[4]{a + bx^4} - \frac{ax^3 \left(\frac{a}{bx^4} + 1\right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^4} + 1\right)^{3/4} x} d\frac{1}{x}}{2(a + bx^4)^{3/4}} \right) + \frac{1}{6}x(a + bx^4)^{5/4} \\
 & \quad \downarrow 807
 \end{aligned}$$

$$\frac{5}{6}a \left(\frac{1}{2}x^4\sqrt{a+bx^4} - \frac{ax^3\left(\frac{a}{bx^4}+1\right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^2}+1\right)^{3/4}} d\frac{1}{x^2}}{4(a+bx^4)^{3/4}} \right) + \frac{1}{6}x(a+bx^4)^{5/4}$$

↓ 229

$$\frac{5}{6}a \left(\frac{1}{2}x^4\sqrt{a+bx^4} - \frac{\sqrt{a}\sqrt{bx^3\left(\frac{a}{bx^4}+1\right)^{3/4}} \operatorname{EllipticF}\left(\frac{1}{2}\arctan\left(\frac{\sqrt{a}}{\sqrt{bx^2}}\right), 2\right)}{2(a+bx^4)^{3/4}} \right) + \frac{1}{6}x(a+bx^4)^{5/4}$$

input `Int[(a + b*x^4)^(5/4), x]`

output `(x*(a + b*x^4)^(5/4))/6 + (5*a*((x*(a + b*x^4)^(1/4))/2 - (Sqrt[a]*Sqrt[b]*(1 + a/(b*x^4))^(3/4)*x^3*EllipticF[ArcTan[Sqrt[a]/(Sqrt[b]*x^2)]/2, 2)]/(2*(a + b*x^4)^(3/4)))/6`

Defintions of rubi rules used

rule 229 `Int[((a_) + (b_.)*(x_)^2)^(-3/4), x_Symbol] := Simp[(2/(a^(3/4)*Rt[b/a, 2])*)*EllipticF[(1/2)*ArcTan[Rt[b/a, 2]*x], 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]`

rule 748 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p + 1)), x] + Simp[a*n*(p/(n*p + 1)) Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || LtQ[Denominator[p + 1/n], Denominator[p]])`

rule 768 `Int[((a_) + (b_.)*(x_)^4)^(-3/4), x_Symbol] := Simp[x^3*((1 + a/(b*x^4))^(3/4))/(a + b*x^4)^(3/4) Int[1/(x^3*(1 + a/(b*x^4))^(3/4)), x], x] /; FreeQ[{a, b}, x]`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

rule 858

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a +
b/x^n)^p/x^(m + 2), x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0] && IntegerQ[m]
```

Maple [F]

$$\int (bx^4 + a)^{\frac{5}{4}} dx$$

input

```
int((b*x^4+a)^(5/4),x)
```

output

```
int((b*x^4+a)^(5/4),x)
```

Fricas [F]

$$\int (a + bx^4)^{5/4} dx = \int (bx^4 + a)^{5/4} dx$$

input

```
integrate((b*x^4+a)^(5/4),x, algorithm="fricas")
```

output

```
integral((b*x^4 + a)^(5/4), x)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.64 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.38

$$\int (a + bx^4)^{5/4} dx = \frac{a^{\frac{5}{4}} x \Gamma\left(\frac{1}{4}\right) {}_2F_1\left(-\frac{5}{4}, \frac{1}{4} \middle| \frac{bx^4 e^{i\pi}}{a}\right)}{4\Gamma\left(\frac{5}{4}\right)}$$

input

```
integrate((b*x**4+a)**(5/4),x)
```


output `a**(5/4)*x*gamma(1/4)*hyper((-5/4, 1/4), (5/4,), b*x**4*exp_polar(I*pi)/a)/(4*gamma(5/4))`

Maxima [F]

$$\int (a + bx^4)^{5/4} dx = \int (bx^4 + a)^{5/4} dx$$

input `integrate((b*x^4+a)^(5/4),x, algorithm="maxima")`

output `integrate((b*x^4 + a)^(5/4), x)`

Giac [F]

$$\int (a + bx^4)^{5/4} dx = \int (bx^4 + a)^{5/4} dx$$

input `integrate((b*x^4+a)^(5/4),x, algorithm="giac")`

output `integrate((b*x^4 + a)^(5/4), x)`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.38

$$\int (a + bx^4)^{5/4} dx = \frac{x (bx^4 + a)^{5/4} {}_2F_1\left(-\frac{5}{4}, \frac{1}{4}; \frac{5}{4}; -\frac{bx^4}{a}\right)}{\left(\frac{bx^4}{a} + 1\right)^{5/4}}$$

input `int((a + b*x^4)^(5/4),x)`

output $(x*(a + b*x^4)^{(5/4)}*\text{hypergeom}([-5/4, 1/4], 5/4, -(b*x^4)/a))/((b*x^4)/a + 1)^{(5/4)}$

Reduce [F]

$$\int (a + bx^4)^{5/4} dx = \frac{7(bx^4 + a)^{1/4} ax}{12} + \frac{(bx^4 + a)^{1/4} bx^5}{6} + \frac{5 \left(\int \frac{1}{(bx^4+a)^{3/4}} dx \right) a^2}{12}$$

input $\text{int}((b*x^4+a)^{(5/4)}, x)$

output $(7*(a + b*x**4)**(1/4)*a*x + 2*(a + b*x**4)**(1/4)*b*x**5 + 5*\text{int}((a + b*x**4)**(1/4)/(a + b*x**4), x)*a**2)/12$

3.40 $\int \sqrt[4]{a + bx^4} dx$

Optimal result	282
Mathematica [C] (verified)	282
Rubi [A] (verified)	283
Maple [F]	285
Fricas [F]	285
Sympy [C] (verification not implemented)	285
Maxima [F]	286
Giac [F]	286
Mupad [B] (verification not implemented)	286
Reduce [F]	287

Optimal result

Integrand size = 11, antiderivative size = 80

$$\int \sqrt[4]{a + bx^4} dx = \frac{1}{2}x\sqrt[4]{a + bx^4} - \frac{\sqrt{a}\sqrt{b}\left(1 + \frac{a}{bx^4}\right)^{3/4} x^3 \operatorname{EllipticF}\left(\frac{1}{2} \cot^{-1}\left(\frac{\sqrt{bx^2}}{\sqrt{a}}\right), 2\right)}{2(a + bx^4)^{3/4}}$$

output

`1/2*x*(b*x^4+a)^(1/4)-1/2*a^(1/2)*b^(1/2)*(1+a/b/x^4)^(3/4)*x^3*InverseJacobiAM(1/2*arccot(b^(1/2)*x^2/a^(1/2)),2^(1/2))/(b*x^4+a)^(3/4)`

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.31 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.58

$$\int \sqrt[4]{a + bx^4} dx = \frac{x\sqrt[4]{a + bx^4} \operatorname{Hypergeometric2F1}\left(-\frac{1}{4}, \frac{1}{4}, \frac{5}{4}, -\frac{bx^4}{a}\right)}{\sqrt[4]{1 + \frac{bx^4}{a}}}$$

input

`Integrate[(a + b*x^4)^(1/4),x]`

output

```
(x*(a + b*x^4)^(1/4)*Hypergeometric2F1[-1/4, 1/4, 5/4, -((b*x^4)/a)]/(1 +
(b*x^4)/a)^(1/4)
```

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {748, 768, 858, 807, 229}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt[4]{a + bx^4} dx \\
 & \quad \downarrow 748 \\
 & \frac{1}{2}a \int \frac{1}{(bx^4 + a)^{3/4}} dx + \frac{1}{2}x \sqrt[4]{a + bx^4} \\
 & \quad \downarrow 768 \\
 & \frac{ax^3 \left(\frac{a}{bx^4} + 1\right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^4} + 1\right)^{3/4} x^3} dx}{2(a + bx^4)^{3/4}} + \frac{1}{2}x \sqrt[4]{a + bx^4} \\
 & \quad \downarrow 858 \\
 & \frac{1}{2}x \sqrt[4]{a + bx^4} - \frac{ax^3 \left(\frac{a}{bx^4} + 1\right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^4} + 1\right)^{3/4} x} d\frac{1}{x}}{2(a + bx^4)^{3/4}} \\
 & \quad \downarrow 807 \\
 & \frac{1}{2}x \sqrt[4]{a + bx^4} - \frac{ax^3 \left(\frac{a}{bx^4} + 1\right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^2} + 1\right)^{3/4} d\frac{1}{x^2}}}{4(a + bx^4)^{3/4}} \\
 & \quad \downarrow 229 \\
 & \frac{1}{2}x \sqrt[4]{a + bx^4} - \frac{\sqrt{a}\sqrt{bx^3} \left(\frac{a}{bx^4} + 1\right)^{3/4} \text{EllipticF}\left(\frac{1}{2} \arctan\left(\frac{\sqrt{a}}{\sqrt{bx^2}}\right), 2\right)}{2(a + bx^4)^{3/4}}
 \end{aligned}$$

input `Int[(a + b*x^4)^(1/4),x]`

output `(x*(a + b*x^4)^(1/4))/2 - (Sqrt[a]*Sqrt[b]*(1 + a/(b*x^4))^(3/4)*x^3*EllipticF[ArcTan[Sqrt[a]/(Sqrt[b]*x^2)]/2, 2])/(2*(a + b*x^4)^(3/4))`

Defintions of rubi rules used

rule 229 `Int[((a_) + (b_.)*(x_)^2)^(-3/4), x_Symbol] := Simp[(2/(a^(3/4)*Rt[b/a, 2]))*EllipticF[(1/2)*ArcTan[Rt[b/a, 2]*x], 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]`

rule 748 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p + 1)), x] + Simp[a*n*(p/(n*p + 1)) Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || LtQ[Denominator[p + 1/n], Denominator[p]])`

rule 768 `Int[((a_) + (b_.)*(x_)^4)^(-3/4), x_Symbol] := Simp[x^3*((1 + a/(b*x^4))^(3/4)/(a + b*x^4)^(3/4)) Int[1/(x^3*(1 + a/(b*x^4))^(3/4)), x], x] /; FreeQ[{a, b}, x]`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

rule 858 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^(m + 2), x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0] && IntegerQ[m]`

Maple [F]

$$\int (bx^4 + a)^{\frac{1}{4}} dx$$

input `int((b*x^4+a)^(1/4),x)`

output `int((b*x^4+a)^(1/4),x)`

Fricas [F]

$$\int \sqrt[4]{a + bx^4} dx = \int (bx^4 + a)^{\frac{1}{4}} dx$$

input `integrate((b*x^4+a)^(1/4),x, algorithm="fricas")`

output `integral((b*x^4 + a)^(1/4), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.48 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.46

$$\int \sqrt[4]{a + bx^4} dx = \frac{\sqrt[4]{ax}\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\begin{matrix} -\frac{1}{4}, \frac{1}{4} \\ \frac{5}{4} \end{matrix} \middle| \frac{bx^4 e^{i\pi}}{a}\right)}{4\Gamma\left(\frac{5}{4}\right)}$$

input `integrate((b*x**4+a)**(1/4),x)`

output `a**(1/4)*x*gamma(1/4)*hyper((-1/4, 1/4), (5/4,), b*x**4*exp_polar(I*pi)/a)/(4*gamma(5/4))`

Maxima [F]

$$\int \sqrt[4]{a + bx^4} dx = \int (bx^4 + a)^{\frac{1}{4}} dx$$

input `integrate((b*x^4+a)^(1/4),x, algorithm="maxima")`

output `integrate((b*x^4 + a)^(1/4), x)`

Giac [F]

$$\int \sqrt[4]{a + bx^4} dx = \int (bx^4 + a)^{\frac{1}{4}} dx$$

input `integrate((b*x^4+a)^(1/4),x, algorithm="giac")`

output `integrate((b*x^4 + a)^(1/4), x)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.46

$$\int \sqrt[4]{a + bx^4} dx = \frac{x (bx^4 + a)^{1/4} {}_2F_1\left(-\frac{1}{4}, \frac{1}{4}; \frac{5}{4}; -\frac{bx^4}{a}\right)}{\left(\frac{bx^4}{a} + 1\right)^{1/4}}$$

input `int((a + b*x^4)^(1/4),x)`

output `(x*(a + b*x^4)^(1/4)*hypergeom([-1/4, 1/4], 5/4, -(b*x^4)/a))/((b*x^4)/a + 1)^(1/4)`

Reduce [F]

$$\int \sqrt[4]{a + bx^4} dx = \frac{(bx^4 + a)^{\frac{1}{4}} x}{2} + \frac{\left(\int \frac{1}{(bx^4 + a)^{\frac{3}{4}}} dx \right) a}{2}$$

input `int((b*x^4+a)^(1/4),x)`

output `((a + b*x**4)**(1/4)*x + int((a + b*x**4)**(1/4)/(a + b*x**4),x)*a)/2`

3.41 $\int \frac{1}{(a+bx^4)^{3/4}} dx$

Optimal result	288
Mathematica [C] (verified)	288
Rubi [A] (verified)	289
Maple [F]	290
Fricas [F]	291
Sympy [C] (verification not implemented)	291
Maxima [F]	291
Giac [F]	292
Mupad [B] (verification not implemented)	292
Reduce [F]	292

Optimal result

Integrand size = 11, antiderivative size = 61

$$\int \frac{1}{(a + bx^4)^{3/4}} dx = -\frac{\sqrt{b}\left(1 + \frac{a}{bx^4}\right)^{3/4} x^3 \operatorname{EllipticF}\left(\frac{1}{2} \cot^{-1}\left(\frac{\sqrt{bx^2}}{\sqrt{a}}\right), 2\right)}{\sqrt{a}(a + bx^4)^{3/4}}$$

output `-b^(1/2)*(1+a/b/x^4)^(3/4)*x^3*InverseJacobiAM(1/2*arccot(b^(1/2)*x^2/a^(1/2)),2^(1/2))/a^(1/2)/(b*x^4+a)^(3/4)`

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.13 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.75

$$\int \frac{1}{(a + bx^4)^{3/4}} dx = \frac{x\left(1 + \frac{bx^4}{a}\right)^{3/4} \operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{3}{4}, \frac{5}{4}, -\frac{bx^4}{a}\right)}{(a + bx^4)^{3/4}}$$

input `Integrate[(a + b*x^4)^(-3/4),x]`

output $(x*(1 + (b*x^4)/a)^{(3/4)}*\text{Hypergeometric2F1}[1/4, 3/4, 5/4, -((b*x^4)/a)])/(a + b*x^4)^{(3/4)}$

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {768, 858, 807, 229}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a + bx^4)^{3/4}} dx \\
 & \quad \downarrow \text{768} \\
 & \frac{x^3 \left(\frac{a}{bx^4} + 1\right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^4} + 1\right)^{3/4} x^3} dx}{(a + bx^4)^{3/4}} \\
 & \quad \downarrow \text{858} \\
 & \frac{x^3 \left(\frac{a}{bx^4} + 1\right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^4} + 1\right)^{3/4} x} d\frac{1}{x}}{(a + bx^4)^{3/4}} \\
 & \quad \downarrow \text{807} \\
 & \frac{x^3 \left(\frac{a}{bx^4} + 1\right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^2} + 1\right)^{3/4} d\frac{1}{x^2}}}{2(a + bx^4)^{3/4}} \\
 & \quad \downarrow \text{229} \\
 & \frac{\sqrt{b}x^3 \left(\frac{a}{bx^4} + 1\right)^{3/4} \text{EllipticF}\left(\frac{1}{2} \arctan\left(\frac{\sqrt{a}}{\sqrt{bx^2}}\right), 2\right)}{\sqrt{a}(a + bx^4)^{3/4}}
 \end{aligned}$$

input $\text{Int}[(a + b*x^4)^{-3/4}, x]$

output $-\left(\frac{\sqrt{b}(1 + a/(bx^4))^{3/4}x^3 \operatorname{EllipticF}[\operatorname{ArcTan}[\sqrt{a}/(\sqrt{b}x^2)]/2, 2]}{\sqrt{a}(a + bx^4)^{3/4}}\right)$

Defintions of rubi rules used

rule 229 $\operatorname{Int}[(a_ + (b_ \cdot)(x_)^2)^{-3/4}, x_Symbol] \rightarrow \operatorname{Simp}[(2/(a^{3/4} \operatorname{Rt}[b/a, 2]) \cdot \operatorname{EllipticF}[(1/2) \operatorname{ArcTan}[\operatorname{Rt}[b/a, 2]x], 2], x] /; \operatorname{FreeQ}\{a, b, x\} \ \&\& \ \operatorname{GtQ}[a, 0] \ \&\& \ \operatorname{PosQ}[b/a]$

rule 768 $\operatorname{Int}[(a_ + (b_ \cdot)(x_)^4)^{-3/4}, x_Symbol] \rightarrow \operatorname{Simp}[x^3 \cdot ((1 + a/(bx^4))^{3/4}) / (a + bx^4)^{3/4}] \operatorname{Int}[1/(x^3(1 + a/(bx^4))^{3/4}), x], x] /; \operatorname{FreeQ}\{a, b, x\}$

rule 807 $\operatorname{Int}[(x_)^{(m_ \cdot)} \cdot ((a_ + (b_ \cdot)(x_)^{(n_)})^{(p_)}), x_Symbol] \rightarrow \operatorname{With}\{k = \operatorname{GCD}[m + 1, n]\}, \operatorname{Simp}[1/k \operatorname{Subst}[\operatorname{Int}[x^{(m+1)/k - 1} (a + bx^{n/k})^p, x], x, x^k], x] /; k \neq 1] /; \operatorname{FreeQ}\{a, b, p, x\} \ \&\& \ \operatorname{IGtQ}[n, 0] \ \&\& \ \operatorname{IntegerQ}[m]$

rule 858 $\operatorname{Int}[(x_)^{(m_ \cdot)} \cdot ((a_ + (b_ \cdot)(x_)^{(n_)})^{(p_)}), x_Symbol] \rightarrow -\operatorname{Subst}[\operatorname{Int}[(a + b/x^n)^p / x^{m+2}, x], x, 1/x] /; \operatorname{FreeQ}\{a, b, p, x\} \ \&\& \ \operatorname{ILtQ}[n, 0] \ \&\& \ \operatorname{IntegerQ}[m]$

Maple [F]

$$\int \frac{1}{(bx^4 + a)^{3/4}} dx$$

input $\operatorname{int}(1/(bx^4+a)^{3/4}, x)$

output $\operatorname{int}(1/(bx^4+a)^{3/4}, x)$

Fricas [F]

$$\int \frac{1}{(a + bx^4)^{3/4}} dx = \int \frac{1}{(bx^4 + a)^{3/4}} dx$$

input `integrate(1/(b*x^4+a)^(3/4),x, algorithm="fricas")`

output `integral((b*x^4 + a)^(-3/4), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.48 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.59

$$\int \frac{1}{(a + bx^4)^{3/4}} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{3}{4} \middle| \frac{bx^4 e^{i\pi}}{a}\right)}{4a^{3/4}\Gamma\left(\frac{5}{4}\right)}$$

input `integrate(1/(b*x**4+a)**(3/4),x)`

output `x*gamma(1/4)*hyper((1/4, 3/4), (5/4,), b*x**4*exp_polar(I*pi)/a)/(4*a**(3/4)*gamma(5/4))`

Maxima [F]

$$\int \frac{1}{(a + bx^4)^{3/4}} dx = \int \frac{1}{(bx^4 + a)^{3/4}} dx$$

input `integrate(1/(b*x^4+a)^(3/4),x, algorithm="maxima")`

output `integrate((b*x^4 + a)^(-3/4), x)`

Giac [F]

$$\int \frac{1}{(a + bx^4)^{3/4}} dx = \int \frac{1}{(bx^4 + a)^{3/4}} dx$$

input `integrate(1/(b*x^4+a)^(3/4),x, algorithm="giac")`

output `integrate((b*x^4 + a)^(-3/4), x)`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.61

$$\int \frac{1}{(a + bx^4)^{3/4}} dx = \frac{x \left(\frac{bx^4}{a} + 1 \right)^{3/4} {}_2F_1 \left(\frac{1}{4}, \frac{3}{4}; \frac{5}{4}; -\frac{bx^4}{a} \right)}{(bx^4 + a)^{3/4}}$$

input `int(1/(a + b*x^4)^(3/4),x)`

output `(x*((b*x^4)/a + 1)^(3/4)*hypergeom([1/4, 3/4], 5/4, -(b*x^4)/a))/(a + b*x^4)^(3/4)`

Reduce [F]

$$\int \frac{1}{(a + bx^4)^{3/4}} dx = \int \frac{1}{(bx^4 + a)^{3/4}} dx$$

input `int(1/(b*x^4+a)^(3/4),x)`

output `int(1/(a + b*x**4)**(3/4),x)`

3.42 $\int \frac{1}{(a+bx^4)^{7/4}} dx$

Optimal result	293
Mathematica [C] (verified)	293
Rubi [A] (verified)	294
Maple [F]	296
Fricas [F]	296
Sympy [C] (verification not implemented)	296
Maxima [F]	297
Giac [F]	297
Mupad [B] (verification not implemented)	297
Reduce [F]	298

Optimal result

Integrand size = 11, antiderivative size = 83

$$\int \frac{1}{(a + bx^4)^{7/4}} dx = \frac{x}{3a(a + bx^4)^{3/4}} - \frac{2\sqrt{b}(1 + \frac{a}{bx^4})^{3/4} x^3 \text{EllipticF}\left(\frac{1}{2} \cot^{-1}\left(\frac{\sqrt{bx^2}}{\sqrt{a}}\right), 2\right)}{3a^{3/2}(a + bx^4)^{3/4}}$$

output `1/3*x/a/(b*x^4+a)^(3/4)-2/3*b^(1/2)*(1+a/b/x^4)^(3/4)*x^3*InverseJacobiAM(1/2*arccot(b^(1/2)*x^2/a^(1/2)),2^(1/2))/a^(3/2)/(b*x^4+a)^(3/4)`

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.44 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.67

$$\int \frac{1}{(a + bx^4)^{7/4}} dx = \frac{x + 2x\left(1 + \frac{bx^4}{a}\right)^{3/4} \text{Hypergeometric2F1}\left(\frac{1}{4}, \frac{3}{4}, \frac{5}{4}, -\frac{bx^4}{a}\right)}{3a(a + bx^4)^{3/4}}$$

input `Integrate[(a + b*x^4)^(-7/4),x]`

output

$$\frac{(x + 2*x*(1 + (b*x^4)/a)^{(3/4)}*Hypergeometric2F1[1/4, 3/4, 5/4, -((b*x^4)/a)])/(3*a*(a + b*x^4)^{(3/4)})$$
Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {749, 768, 858, 807, 229}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(a + bx^4)^{7/4}} dx \\ & \quad \downarrow \text{749} \\ & \frac{2 \int \frac{1}{(bx^4+a)^{3/4}} dx}{3a} + \frac{x}{3a(a + bx^4)^{3/4}} \\ & \quad \downarrow \text{768} \\ & \frac{2x^3 \left(\frac{a}{bx^4} + 1\right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^4} + 1\right)^{3/4} x^3} dx}{3a(a + bx^4)^{3/4}} + \frac{x}{3a(a + bx^4)^{3/4}} \\ & \quad \downarrow \text{858} \\ & \frac{x}{3a(a + bx^4)^{3/4}} - \frac{2x^3 \left(\frac{a}{bx^4} + 1\right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^4} + 1\right)^{3/4} x} d\frac{1}{x}}{3a(a + bx^4)^{3/4}} \\ & \quad \downarrow \text{807} \\ & \frac{x}{3a(a + bx^4)^{3/4}} - \frac{x^3 \left(\frac{a}{bx^4} + 1\right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^2} + 1\right)^{3/4} d\frac{1}{x^2}}}{3a(a + bx^4)^{3/4}} \\ & \quad \downarrow \text{229} \\ & \frac{x}{3a(a + bx^4)^{3/4}} - \frac{2\sqrt{b}x^3 \left(\frac{a}{bx^4} + 1\right)^{3/4} \text{EllipticF}\left(\frac{1}{2} \arctan\left(\frac{\sqrt{a}}{\sqrt{bx^2}}\right), 2\right)}{3a^{3/2}(a + bx^4)^{3/4}} \end{aligned}$$

input `Int[(a + b*x^4)^(-7/4),x]`

output `x/(3*a*(a + b*x^4)^(3/4)) - (2*Sqrt[b]*(1 + a/(b*x^4))^(3/4)*x^3*EllipticF[ArcTan[Sqrt[a]/(Sqrt[b]*x^2)]/2, 2])/(3*a^(3/2)*(a + b*x^4)^(3/4))`

Defintions of rubi rules used

rule 229 `Int[((a_) + (b_.)*(x_)^2)^(-3/4), x_Symbol] := Simp[(2/(a^(3/4)*Rt[b/a, 2]))*EllipticF[(1/2)*ArcTan[Rt[b/a, 2]*x], 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]`

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

rule 768 `Int[((a_) + (b_.)*(x_)^4)^(-3/4), x_Symbol] := Simp[x^3*((1 + a/(b*x^4))^(3/4)/(a + b*x^4)^(3/4)) Int[1/(x^3*(1 + a/(b*x^4))^(3/4)), x], x] /; FreeQ[{a, b}, x]`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

rule 858 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^(m + 2), x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0] && IntegerQ[m]`

Maple [F]

$$\int \frac{1}{(bx^4 + a)^{7/4}} dx$$

input `int(1/(b*x^4+a)^(7/4),x)`

output `int(1/(b*x^4+a)^(7/4),x)`

Fricas [F]

$$\int \frac{1}{(a + bx^4)^{7/4}} dx = \int \frac{1}{(bx^4 + a)^{7/4}} dx$$

input `integrate(1/(b*x^4+a)^(7/4),x, algorithm="fricas")`

output `integral((b*x^4 + a)^(1/4)/(b^2*x^8 + 2*a*b*x^4 + a^2), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.61 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.43

$$\int \frac{1}{(a + bx^4)^{7/4}} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{7}{4} \middle| \frac{bx^4 e^{i\pi}}{a}\right)}{4a^{7/4}\Gamma\left(\frac{5}{4}\right)}$$

input `integrate(1/(b*x**4+a)**(7/4),x)`

output `x*gamma(1/4)*hyper((1/4, 7/4), (5/4,), b*x**4*exp_polar(I*pi)/a)/(4*a**(7/4)*gamma(5/4))`

Maxima [F]

$$\int \frac{1}{(a + bx^4)^{7/4}} dx = \int \frac{1}{(bx^4 + a)^{7/4}} dx$$

input `integrate(1/(b*x^4+a)^(7/4),x, algorithm="maxima")`

output `integrate((b*x^4 + a)^(-7/4), x)`

Giac [F]

$$\int \frac{1}{(a + bx^4)^{7/4}} dx = \int \frac{1}{(bx^4 + a)^{7/4}} dx$$

input `integrate(1/(b*x^4+a)^(7/4),x, algorithm="giac")`

output `integrate((b*x^4 + a)^(-7/4), x)`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.45

$$\int \frac{1}{(a + bx^4)^{7/4}} dx = \frac{x \left(\frac{bx^4}{a} + 1 \right)^{7/4} {}_2F_1 \left(\frac{1}{4}, \frac{7}{4}; \frac{5}{4}; -\frac{bx^4}{a} \right)}{(bx^4 + a)^{7/4}}$$

input `int(1/(a + b*x^4)^(7/4),x)`

output `(x*((b*x^4)/a + 1)^(7/4)*hypergeom([1/4, 7/4], 5/4, -(b*x^4)/a))/(a + b*x^4)^(7/4)`

Reduce [F]

$$\int \frac{1}{(a + bx^4)^{7/4}} dx = \int \frac{1}{(bx^4 + a)^{3/4} a + (bx^4 + a)^{3/4} bx^4} dx$$

input `int(1/(b*x^4+a)^(7/4),x)`

output `int(1/((a + b*x**4)**(3/4)*a + (a + b*x**4)**(3/4)*b*x**4),x)`

3.43 $\int \frac{1}{(a+bx^4)^{11/4}} dx$

Optimal result	299
Mathematica [C] (verified)	299
Rubi [A] (verified)	300
Maple [F]	302
Fricas [F]	302
Sympy [C] (verification not implemented)	303
Maxima [F]	303
Giac [F]	303
Mupad [B] (verification not implemented)	304
Reduce [F]	304

Optimal result

Integrand size = 11, antiderivative size = 102

$$\int \frac{1}{(a+bx^4)^{11/4}} dx = \frac{x}{7a(a+bx^4)^{7/4}} + \frac{2x}{7a^2(a+bx^4)^{3/4}} - \frac{4\sqrt{b}\left(1+\frac{a}{bx^4}\right)^{3/4} x^3 \operatorname{EllipticF}\left(\frac{1}{2} \cot^{-1}\left(\frac{\sqrt{bx^2}}{\sqrt{a}}\right), 2\right)}{7a^{5/2}(a+bx^4)^{3/4}}$$

output

```
1/7*x/a/(b*x^4+a)^(7/4)+2/7*x/a^2/(b*x^4+a)^(3/4)-4/7*b^(1/2)*(1+a/b/x^4)^(3/4)*x^3*InverseJacobiAM(1/2*arccot(b^(1/2)*x^2/a^(1/2)),2^(1/2))/a^(5/2)/(b*x^4+a)^(3/4)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 9.48 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.71

$$\int \frac{1}{(a+bx^4)^{11/4}} dx = \frac{3ax + 2bx^5 + 4x(a+bx^4) \left(1 + \frac{bx^4}{a}\right)^{3/4} \operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{3}{4}, \frac{5}{4}, -\frac{bx^4}{a}\right)}{7a^2(a+bx^4)^{7/4}}$$

input `Integrate[(a + b*x^4)^(-11/4),x]`

output `(3*a*x + 2*b*x^5 + 4*x*(a + b*x^4)*(1 + (b*x^4)/a)^(3/4)*Hypergeometric2F1[1/4, 3/4, 5/4, -((b*x^4)/a)]/(7*a^2*(a + b*x^4)^(7/4))`

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.08, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {749, 749, 768, 858, 807, 229}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a + bx^4)^{11/4}} dx \\
 & \quad \downarrow 749 \\
 & \frac{6 \int \frac{1}{(bx^4+a)^{7/4}} dx}{7a} + \frac{x}{7a(a + bx^4)^{7/4}} \\
 & \quad \downarrow 749 \\
 & \frac{6 \left(\frac{2 \int \frac{1}{(bx^4+a)^{3/4}} dx}{3a} + \frac{x}{3a(a+bx^4)^{3/4}} \right)}{7a} + \frac{x}{7a(a + bx^4)^{7/4}} \\
 & \quad \downarrow 768 \\
 & \frac{6 \left(\frac{2x^3 \left(\frac{a}{bx^4} + 1 \right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^4} + 1 \right)^{3/4} x^3} dx}{3a(a+bx^4)^{3/4}} + \frac{x}{3a(a+bx^4)^{3/4}} \right)}{7a} + \frac{x}{7a(a + bx^4)^{7/4}} \\
 & \quad \downarrow 858
 \end{aligned}$$

$$\begin{aligned}
& \frac{6 \left(\frac{x}{3a(ax^4)^{3/4}} - \frac{2x^3 \left(\frac{a}{bx^4} + 1\right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^4} + 1\right)^{3/4} x} d\frac{1}{x}}{3a(ax^4)^{3/4}} \right)}{7a} + \frac{x}{7a(a+bx^4)^{7/4}} \\
& \quad \downarrow \text{807} \\
& \frac{6 \left(\frac{x}{3a(ax^4)^{3/4}} - \frac{x^3 \left(\frac{a}{bx^4} + 1\right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^2} + 1\right)^{3/4} d\frac{1}{x^2}}}{3a(ax^4)^{3/4}} \right)}{7a} + \frac{x}{7a(a+bx^4)^{7/4}} \\
& \quad \downarrow \text{229} \\
& \frac{6 \left(\frac{x}{3a(ax^4)^{3/4}} - \frac{2\sqrt{b}x^3 \left(\frac{a}{bx^4} + 1\right)^{3/4} \text{EllipticF}\left(\frac{1}{2} \arctan\left(\frac{\sqrt{a}}{\sqrt{bx^2}}\right), 2\right)}{3a^{3/2}(a+bx^4)^{3/4}} \right)}{7a} + \frac{x}{7a(a+bx^4)^{7/4}}
\end{aligned}$$

input `Int[(a + b*x^4)^(-11/4), x]`

output `x/(7*a*(a + b*x^4)^(7/4)) + (6*(x/(3*a*(a + b*x^4)^(3/4)) - (2*sqrt[b]*(1 + a/(b*x^4))^(3/4)*x^3*EllipticF[ArcTan[Sqrt[a]/(Sqrt[b]*x^2)]/2, 2)]/(3*a^(3/2)*(a + b*x^4)^(3/4)))/(7*a)`

Defintions of rubi rules used

rule 229 `Int[((a_) + (b_.)*(x_)^2)^(-3/4), x_Symbol] := Simp[(2/(a^(3/4)*Rt[b/a, 2]) * EllipticF[(1/2)*ArcTan[Rt[b/a, 2]*x], 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]`

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

rule 768 `Int[((a_) + (b_.)*(x_)^4)^(-3/4), x_Symbol] := Simp[x^3*((1 + a/(b*x^4))^(3/4))/(a + b*x^4)^(3/4) Int[1/(x^3*(1 + a/(b*x^4))^(3/4)), x], x] /; FreeQ[{a, b}, x]`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

rule 858 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^(m + 2), x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0] && IntegerQ[m]`

Maple [F]

$$\int \frac{1}{(bx^4 + a)^{\frac{11}{4}}} dx$$

input `int(1/(b*x^4+a)^(11/4),x)`

output `int(1/(b*x^4+a)^(11/4),x)`

Fricas [F]

$$\int \frac{1}{(a + bx^4)^{11/4}} dx = \int \frac{1}{(bx^4 + a)^{\frac{11}{4}}} dx$$

input `integrate(1/(b*x^4+a)^(11/4),x, algorithm="fricas")`

output `integral((b*x^4 + a)^(1/4)/(b^3*x^12 + 3*a*b^2*x^8 + 3*a^2*b*x^4 + a^3), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.00 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.35

$$\int \frac{1}{(a + bx^4)^{11/4}} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{11}{4} \middle| \frac{bx^4 e^{i\pi}}{a}\right)}{4a^{\frac{11}{4}} \Gamma\left(\frac{5}{4}\right)}$$

input `integrate(1/(b*x**4+a)**(11/4),x)`

output `x*gamma(1/4)*hyper((1/4, 11/4), (5/4,), b*x**4*exp_polar(I*pi)/a)/(4*a**(11/4)*gamma(5/4))`

Maxima [F]

$$\int \frac{1}{(a + bx^4)^{11/4}} dx = \int \frac{1}{(bx^4 + a)^{\frac{11}{4}}} dx$$

input `integrate(1/(b*x^4+a)^(11/4),x, algorithm="maxima")`

output `integrate((b*x^4 + a)^(-11/4), x)`

Giac [F]

$$\int \frac{1}{(a + bx^4)^{11/4}} dx = \int \frac{1}{(bx^4 + a)^{\frac{11}{4}}} dx$$

input `integrate(1/(b*x^4+a)^(11/4),x, algorithm="giac")`

output `integrate((b*x^4 + a)^(-11/4), x)`

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.36

$$\int \frac{1}{(a + bx^4)^{11/4}} dx = \frac{x \left(\frac{bx^4}{a} + 1 \right)^{11/4} {}_2F_1\left(\frac{1}{4}, \frac{11}{4}; \frac{5}{4}; -\frac{bx^4}{a}\right)}{(bx^4 + a)^{11/4}}$$

input `int(1/(a + b*x^4)^(11/4),x)`output `(x*((b*x^4)/a + 1)^(11/4)*hypergeom([1/4, 11/4], 5/4, -(b*x^4)/a))/(a + b*x^4)^(11/4)`**Reduce [F]**

$$\int \frac{1}{(a + bx^4)^{11/4}} dx = \int \frac{1}{(bx^4 + a)^{3/4} a^2 + 2(bx^4 + a)^{3/4} abx^4 + (bx^4 + a)^{3/4} b^2x^8} dx$$

input `int(1/(b*x^4+a)^(11/4),x)`output `int(1/((a + b*x**4)**(3/4)*a**2 + 2*(a + b*x**4)**(3/4)*a*b*x**4 + (a + b*x**4)**(3/4)*b**2*x**8),x)`

3.44 $\int \frac{1}{(a+bx^4)^{15/4}} dx$

Optimal result	305
Mathematica [C] (verified)	305
Rubi [A] (verified)	306
Maple [F]	309
Fricas [F]	309
Sympy [C] (verification not implemented)	309
Maxima [F]	310
Giac [F]	310
Mupad [B] (verification not implemented)	310
Reduce [F]	311

Optimal result

Integrand size = 11, antiderivative size = 121

$$\int \frac{1}{(a+bx^4)^{15/4}} dx = \frac{x}{11a(a+bx^4)^{11/4}} + \frac{10x}{77a^2(a+bx^4)^{7/4}} + \frac{20x}{77a^3(a+bx^4)^{3/4}} - \frac{40\sqrt{b}\left(1+\frac{a}{bx^4}\right)^{3/4}x^3 \operatorname{EllipticF}\left(\frac{1}{2}\cot^{-1}\left(\frac{\sqrt{bx^2}}{\sqrt{a}}\right), 2\right)}{77a^{7/2}(a+bx^4)^{3/4}}$$

output

```
1/11*x/a/(b*x^4+a)^(11/4)+10/77*x/a^2/(b*x^4+a)^(7/4)+20/77*x/a^3/(b*x^4+a)^(3/4)-40/77*b^(1/2)*(1+a/b/x^4)^(3/4)*x^3*InverseJacobiAM(1/2*arccot(b^(1/2)*x^2/a^(1/2)),2^(1/2))/a^(7/2)/(b*x^4+a)^(3/4)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.04 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.70

$$\int \frac{1}{(a+bx^4)^{15/4}} dx = \frac{37a^2x + 50abx^5 + 20b^2x^9 + 40x(a+bx^4)^2 \left(1 + \frac{bx^4}{a}\right)^{3/4} \operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{3}{4}, \frac{5}{4}\right)}{77a^3(a+bx^4)^{11/4}}$$

input `Integrate[(a + b*x^4)^(-15/4),x]`

output `(37*a^2*x + 50*a*b*x^5 + 20*b^2*x^9 + 40*x*(a + b*x^4)^2*(1 + (b*x^4)/a)^(3/4)*Hypergeometric2F1[1/4, 3/4, 5/4, -((b*x^4)/a)]/(77*a^3*(a + b*x^4)^(11/4))`

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.13, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {749, 749, 749, 768, 858, 807, 229}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a + bx^4)^{15/4}} dx \\
 & \quad \downarrow 749 \\
 & \frac{10 \int \frac{1}{(bx^4+a)^{11/4}} dx}{11a} + \frac{x}{11a (a + bx^4)^{11/4}} \\
 & \quad \downarrow 749 \\
 & \frac{10 \left(\frac{6 \int \frac{1}{(bx^4+a)^{7/4}} dx}{7a} + \frac{x}{7a(a+bx^4)^{7/4}} \right)}{11a} + \frac{x}{11a (a + bx^4)^{11/4}} \\
 & \quad \downarrow 749 \\
 & \frac{10 \left(\frac{6 \left(\frac{2 \int \frac{1}{(bx^4+a)^{3/4}} dx}{3a} + \frac{x}{3a(a+bx^4)^{3/4}} \right)}{7a} + \frac{x}{7a(a+bx^4)^{7/4}} \right)}{11a} + \frac{x}{11a (a + bx^4)^{11/4}} \\
 & \quad \downarrow 768
 \end{aligned}$$

$$10 \left(\frac{6 \left(\frac{2x^3 \left(\frac{a}{bx^4} + 1 \right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^4} + 1 \right)^{3/4} x^3} dx}{3a(a+bx^4)^{3/4}} + \frac{x}{3a(a+bx^4)^{3/4}} \right)}{7a} + \frac{x}{7a(a+bx^4)^{7/4}} \right) + \frac{x}{11a(a+bx^4)^{11/4}}$$

858

$$10 \left(\frac{6 \left(\frac{x}{3a(a+bx^4)^{3/4}} - \frac{2x^3 \left(\frac{a}{bx^4} + 1 \right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^4} + 1 \right)^{3/4} x} d\frac{1}{x}}{3a(a+bx^4)^{3/4}} \right)}{7a} + \frac{x}{7a(a+bx^4)^{7/4}} \right) + \frac{x}{11a(a+bx^4)^{11/4}}$$

807

$$10 \left(\frac{6 \left(\frac{x}{3a(a+bx^4)^{3/4}} - \frac{x^3 \left(\frac{a}{bx^4} + 1 \right)^{3/4} \int \frac{1}{\left(\frac{a}{bx^2} + 1 \right)^{3/4} d\frac{1}{x^2}}}{3a(a+bx^4)^{3/4}} \right)}{7a} + \frac{x}{7a(a+bx^4)^{7/4}} \right) + \frac{x}{11a(a+bx^4)^{11/4}}$$

229

$$10 \left(\frac{6 \left(\frac{x}{3a(a+bx^4)^{3/4}} - \frac{2\sqrt{b}x^3 \left(\frac{a}{bx^4} + 1 \right)^{3/4} \operatorname{EllipticF}\left(\frac{1}{2} \arctan\left(\frac{\sqrt{a}}{\sqrt{bx^2}}\right), 2\right)}{3a^{3/2}(a+bx^4)^{3/4}} \right)}{7a} + \frac{x}{7a(a+bx^4)^{7/4}} \right) + \frac{11a}{x} \frac{x}{11a(a+bx^4)^{11/4}}$$

input `Int[(a + b*x^4)^(-15/4),x]`

output `x/(11*a*(a + b*x^4)^(11/4)) + (10*(x/(7*a*(a + b*x^4)^(7/4)) + (6*(x/(3*a*(a + b*x^4)^(3/4)) - (2*Sqrt[b]*(1 + a/(b*x^4))^(3/4)*x^3*EllipticF[ArcTan[Sqrt[a]/(Sqrt[b]*x^2)]/2, 2)]/(3*a^(3/2)*(a + b*x^4)^(3/4))))/(7*a)))/(11*a)`

Defintions of rubi rules used

rule 229 `Int[((a_) + (b_.)*(x_)^2)^(-3/4), x_Symbol] := Simp[(2/(a^(3/4)*Rt[b/a, 2]))*EllipticF[(1/2)*ArcTan[Rt[b/a, 2]*x], 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]`

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

rule 768 `Int[((a_) + (b_.)*(x_)^4)^(-3/4), x_Symbol] := Simp[x^3*((1 + a/(b*x^4))^(3/4)/(a + b*x^4)^(3/4)) Int[1/(x^3*(1 + a/(b*x^4))^(3/4)), x], x] /; FreeQ[{a, b}, x]`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

rule 858 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^(m + 2), x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0] && IntegerQ[m]`

Maple [F]

$$\int \frac{1}{(bx^4 + a)^{\frac{15}{4}}} dx$$

input `int(1/(b*x^4+a)^(15/4),x)`

output `int(1/(b*x^4+a)^(15/4),x)`

Fricas [F]

$$\int \frac{1}{(a + bx^4)^{15/4}} dx = \int \frac{1}{(bx^4 + a)^{\frac{15}{4}}} dx$$

input `integrate(1/(b*x^4+a)^(15/4),x, algorithm="fricas")`

output `integral((b*x^4 + a)^(1/4)/(b^4*x^16 + 4*a*b^3*x^12 + 6*a^2*b^2*x^8 + 4*a^3*b*x^4 + a^4), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.72 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.30

$$\int \frac{1}{(a + bx^4)^{15/4}} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{15}{4} \middle| \frac{bx^4 e^{i\pi}}{a}\right)}{4a^{\frac{15}{4}} \Gamma\left(\frac{5}{4}\right)}$$

input `integrate(1/(b*x**4+a)**(15/4),x)`

output `x*gamma(1/4)*hyper((1/4, 15/4), (5/4,), b*x**4*exp_polar(I*pi)/a)/(4*a**(15/4)*gamma(5/4))`

Maxima [F]

$$\int \frac{1}{(a + bx^4)^{15/4}} dx = \int \frac{1}{(bx^4 + a)^{15/4}} dx$$

input `integrate(1/(b*x^4+a)^(15/4),x, algorithm="maxima")`

output `integrate((b*x^4 + a)^(-15/4), x)`

Giac [F]

$$\int \frac{1}{(a + bx^4)^{15/4}} dx = \int \frac{1}{(bx^4 + a)^{15/4}} dx$$

input `integrate(1/(b*x^4+a)^(15/4),x, algorithm="giac")`

output `integrate((b*x^4 + a)^(-15/4), x)`

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.31

$$\int \frac{1}{(a + bx^4)^{15/4}} dx = \frac{x \left(\frac{bx^4}{a} + 1 \right)^{15/4} {}_2F_1 \left(\frac{1}{4}, \frac{15}{4}, \frac{5}{4}, -\frac{bx^4}{a} \right)}{(bx^4 + a)^{15/4}}$$

input `int(1/(a + b*x^4)^(15/4),x)`

output `(x*((b*x^4)/a + 1)^(15/4)*hypergeom([1/4, 15/4], 5/4, -(b*x^4)/a))/(a + b*x^4)^(15/4)`

Reduce [F]

$$\int \frac{1}{(a + bx^4)^{15/4}} dx = \int \frac{1}{(bx^4 + a)^{3/4} a^3 + 3(bx^4 + a)^{3/4} a^2 bx^4 + 3(bx^4 + a)^{3/4} a b^2 x^8 + (bx^4 + a)^{3/4} b^3 x^{12}} dx$$

input `int(1/(b*x^4+a)^(15/4),x)`

output `int(1/((a + b*x**4)**(3/4)*a**3 + 3*(a + b*x**4)**(3/4)*a**2*b*x**4 + 3*(a + b*x**4)**(3/4)*a*b**2*x**8 + (a + b*x**4)**(3/4)*b**3*x**12),x)`

3.45 $\int (a - bx^4)^{3/4} dx$

Optimal result	312
Mathematica [A] (verified)	313
Rubi [A] (verified)	313
Maple [A] (verified)	317
Fricas [C] (verification not implemented)	318
Sympy [C] (verification not implemented)	319
Maxima [A] (verification not implemented)	319
Giac [F]	320
Mupad [B] (verification not implemented)	320
Reduce [F]	321

Optimal result

Integrand size = 12, antiderivative size = 168

$$\int (a - bx^4)^{3/4} dx = \frac{1}{4}x(a - bx^4)^{3/4} - \frac{3a \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a - bx^4}}\right)}{8\sqrt{2}\sqrt[4]{b}} + \frac{3a \arctan\left(1 + \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a - bx^4}}\right)}{8\sqrt{2}\sqrt[4]{b}} + \frac{3a \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a - bx^4}\left(1 + \frac{\sqrt{bx^2}}{\sqrt{a - bx^4}}\right)}\right)}{8\sqrt{2}\sqrt[4]{b}}$$

output

```
1/4*x*(-b*x^4+a)^(3/4)+3/16*a*arctan(-1+2^(1/2)*b^(1/4)*x/(-b*x^4+a)^(1/4))
)*2^(1/2)/b^(1/4)+3/16*a*arctan(1+2^(1/2)*b^(1/4)*x/(-b*x^4+a)^(1/4))*2^(1
/2)/b^(1/4)+3/16*a*arctanh(2^(1/2)*b^(1/4)*x/(-b*x^4+a)^(1/4)/(1+b^(1/2)*x
^2/(-b*x^4+a)^(1/2)))*2^(1/2)/b^(1/4)
```

Mathematica [A] (verified)

Time = 0.46 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.86

$$\int (a - bx^4)^{3/4} dx = \frac{4\sqrt[4]{b}x(a - bx^4)^{3/4} + 3\sqrt{2}a \arctan\left(\frac{\sqrt{2}\sqrt[4]{b}x\sqrt[4]{a - bx^4}}{-\sqrt{bx^2 + \sqrt{a - bx^4}}}\right) + 3\sqrt{2}a \operatorname{arctanh}\left(\frac{\sqrt{bx^2 + \sqrt{a - bx^4}}}{\sqrt{2}\sqrt[4]{b}x\sqrt[4]{a - bx^4}}\right)}{16\sqrt[4]{b}}$$

input `Integrate[(a - b*x^4)^(3/4), x]`

output `(4*b^(1/4)*x*(a - b*x^4)^(3/4) + 3*Sqrt[2]*a*ArcTan[(Sqrt[2]*b^(1/4)*x*(a - b*x^4)^(1/4))/(-Sqrt[b]*x^2 + Sqrt[a - b*x^4])] + 3*Sqrt[2]*a*ArcTanh[(Sqrt[b]*x^2 + Sqrt[a - b*x^4])/(Sqrt[2]*b^(1/4)*x*(a - b*x^4)^(1/4))]/(16*b^(1/4))`

Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.41, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {748, 770, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a - bx^4)^{3/4} dx \\ & \quad \downarrow 748 \\ & \frac{3}{4}a \int \frac{1}{\sqrt[4]{a - bx^4}} dx + \frac{1}{4}x(a - bx^4)^{3/4} \\ & \quad \downarrow 770 \\ & \frac{3}{4}a \int \frac{1}{\frac{bx^4}{a - bx^4} + 1} d\frac{x}{\sqrt[4]{a - bx^4}} + \frac{1}{4}x(a - bx^4)^{3/4} \\ & \quad \downarrow 755 \end{aligned}$$

$$\frac{3}{4}a \left(\frac{1}{2} \int \frac{1 - \frac{\sqrt{bx^2}}{\sqrt{a-bx^4}}}{\frac{bx^4}{a-bx^4} + 1} d \frac{x}{\sqrt[4]{a-bx^4}} + \frac{1}{2} \int \frac{\frac{\sqrt{bx^2}}{\sqrt{a-bx^4}} + 1}{\frac{bx^4}{a-bx^4} + 1} d \frac{x}{\sqrt[4]{a-bx^4}} \right) + \frac{1}{4}x(a-bx^4)^{3/4}$$

↓ 1476

$$\frac{3}{4}a \left(\frac{1}{2} \int \frac{1 - \frac{\sqrt{bx^2}}{\sqrt{a-bx^4}}}{\frac{bx^4}{a-bx^4} + 1} d \frac{x}{\sqrt[4]{a-bx^4}} + \frac{1}{2} \left(\frac{\int \frac{\frac{x^2}{\sqrt{a-bx^4}} - \frac{1}{\sqrt[4]{b} \sqrt[4]{a-bx^4}} + \frac{1}{\sqrt{b}}}{\sqrt[4]{a-bx^4} + \frac{1}{\sqrt{b}}} d \frac{x}{\sqrt[4]{a-bx^4}}}{2\sqrt{b}} + \frac{\int \frac{\frac{x^2}{\sqrt{a-bx^4}} + \frac{1}{\sqrt[4]{b} \sqrt[4]{a-bx^4}} + \frac{1}{\sqrt{b}}}{\sqrt[4]{a-bx^4} + \frac{1}{\sqrt{b}}} d \frac{x}{\sqrt[4]{a-bx^4}}}{2\sqrt{b}} \right) \right) + \frac{1}{4}x(a-bx^4)^{3/4}$$

↓ 1082

$$\frac{3}{4}a \left(\frac{1}{2} \left(\frac{\int \frac{1}{-\frac{x^2}{\sqrt{a-bx^4}} - 1} d \left(1 - \frac{\sqrt[4]{2} \sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} \right)}{\sqrt{2} \sqrt[4]{b}} - \frac{\int \frac{1}{-\frac{x^2}{\sqrt{a-bx^4}} - 1} d \left(\frac{\sqrt[4]{2} \sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} + 1 \right)}{\sqrt{2} \sqrt[4]{b}} \right) + \frac{1}{2} \int \frac{1 - \frac{\sqrt{bx^2}}{\sqrt{a-bx^4}}}{\frac{bx^4}{a-bx^4} + 1} d \frac{x}{\sqrt[4]{a-bx^4}} \right) + \frac{1}{4}x(a-bx^4)^{3/4}$$

↓ 217

$$\frac{3}{4}a \left(\frac{1}{2} \int \frac{1 - \frac{\sqrt{bx^2}}{\sqrt{a-bx^4}}}{\frac{bx^4}{a-bx^4} + 1} d \frac{x}{\sqrt[4]{a-bx^4}} + \frac{1}{2} \left(\frac{\arctan \left(\frac{\sqrt[4]{2} \sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} + 1 \right)}{\sqrt{2} \sqrt[4]{b}} - \frac{\arctan \left(1 - \frac{\sqrt[4]{2} \sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} \right)}{\sqrt{2} \sqrt[4]{b}} \right) \right) + \frac{1}{4}x(a-bx^4)^{3/4}$$

↓ 1479

$$\frac{3}{4}a \left(\frac{1}{2} \left(\frac{\int \frac{\sqrt{2} - \frac{2 \sqrt[4]{bx}}{\sqrt[4]{a-bx^4}}}{\sqrt[4]{b} \left(\frac{x^2}{\sqrt{a-bx^4}} - \frac{1}{\sqrt[4]{b} \sqrt[4]{a-bx^4}} + \frac{1}{\sqrt{b}} \right)} d \frac{x}{\sqrt[4]{a-bx^4}}}{2\sqrt{2} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{2} \left(\frac{\sqrt[4]{2} \sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} + 1 \right)}{\sqrt[4]{b} \left(\frac{x^2}{\sqrt{a-bx^4}} + \frac{1}{\sqrt[4]{b} \sqrt[4]{a-bx^4}} + \frac{1}{\sqrt{b}} \right)} d \frac{x}{\sqrt[4]{a-bx^4}}}{2\sqrt{2} \sqrt[4]{b}} \right) \right) + \frac{1}{4}x(a-bx^4)^{3/4}$$

↓ 25

$$\frac{3}{4}a \left(\frac{1}{2} \left(\frac{\int \frac{\sqrt{2} - \frac{2\sqrt[4]{b}x}{\sqrt{a-bx^4}}}{\sqrt[4]{b} \left(\frac{x^2}{\sqrt{a-bx^4}} - \frac{\sqrt{2}x}{\sqrt[4]{b}\sqrt{a-bx^4}} + \frac{1}{\sqrt{b}} \right)} d\sqrt[4]{a-bx^4}}{2\sqrt{2}\sqrt[4]{b}} + \frac{\int \frac{\sqrt{2} \left(\frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt{a-bx^4}} + 1 \right)}{\sqrt[4]{b} \left(\frac{x^2}{\sqrt{a-bx^4}} + \frac{\sqrt{2}x}{\sqrt[4]{b}\sqrt{a-bx^4}} + \frac{1}{\sqrt{b}} \right)} d\sqrt[4]{a-bx^4}}{2\sqrt{2}\sqrt[4]{b}} \right) + \frac{1}{2} \right)$$

$$\frac{1}{4}x(a-bx^4)^{3/4}$$

↓ 27

$$\frac{3}{4}a \left(\frac{1}{2} \left(\frac{\int \frac{\sqrt{2} - \frac{2\sqrt[4]{b}x}{\sqrt{a-bx^4}}}{\sqrt[4]{b} \left(\frac{x^2}{\sqrt{a-bx^4}} - \frac{\sqrt{2}x}{\sqrt[4]{b}\sqrt{a-bx^4}} + \frac{1}{\sqrt{b}} \right)} d\sqrt[4]{a-bx^4}}{2\sqrt{2}\sqrt[4]{b}} + \frac{\int \frac{\frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt{a-bx^4}} + 1}{\sqrt[4]{b} \left(\frac{x^2}{\sqrt{a-bx^4}} + \frac{\sqrt{2}x}{\sqrt[4]{b}\sqrt{a-bx^4}} + \frac{1}{\sqrt{b}} \right)} d\sqrt[4]{a-bx^4}}{2\sqrt{b}} \right) + \frac{1}{2} \left(\arctan \left(\frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt{a-bx^4}} + 1 \right) - \arctan \left(1 - \frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt{a-bx^4}} \right) \right) \right)$$

$$\frac{1}{4}x(a-bx^4)^{3/4}$$

↓ 1103

$$\frac{3}{4}a \left(\frac{1}{2} \left(\frac{\arctan \left(\frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt{a-bx^4}} + 1 \right)}{\sqrt{2}\sqrt[4]{b}} - \frac{\arctan \left(1 - \frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt{a-bx^4}} \right)}{\sqrt{2}\sqrt[4]{b}} \right) + \frac{1}{2} \left(\frac{\log \left(\frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt{a-bx^4}} + \frac{\sqrt{bx^2}}{\sqrt{a-bx^4}} + 1 \right)}{2\sqrt{2}\sqrt[4]{b}} - \frac{\log \left(\frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt{a-bx^4}} - \frac{\sqrt{bx^2}}{\sqrt{a-bx^4}} + 1 \right)}{2\sqrt{2}\sqrt[4]{b}} \right) \right)$$

$$\frac{1}{4}x(a-bx^4)^{3/4}$$

input `Int[(a - b*x^4)^(3/4),x]`

output

$$\begin{aligned} & (x*(a - b*x^4)^{(3/4)}/4 + (3*a*((-ArcTan[1 - (Sqrt[2]*b^{(1/4)}*x)/(a - b*x \\ & ^4)^{(1/4)]/(Sqrt[2]*b^{(1/4)})}) + ArcTan[1 + (Sqrt[2]*b^{(1/4)}*x)/(a - b*x^4) \\ & ^{(1/4)]/(Sqrt[2]*b^{(1/4)})})/2 + (-1/2*Log[1 + (Sqrt[b]*x^2)/Sqrt[a - b*x^4] \\ & - (Sqrt[2]*b^{(1/4)}*x)/(a - b*x^4)^{(1/4)]/(Sqrt[2]*b^{(1/4)}) + Log[1 + (Sqr \\ & t[b]*x^2)/Sqrt[a - b*x^4] + (Sqrt[2]*b^{(1/4)}*x)/(a - b*x^4)^{(1/4)]/(2*Sqrt \\ & [2]*b^{(1/4)})/2))/4 \end{aligned}$$

Defintions of rubi rules used

rule 25

$$\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$$

rule 27

$$\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F_x, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ !\text{Ma} \\ \text{tchQ}[F_x, (b_)*(G_x)] \text{ ; FreeQ}[b, x]$$

rule 217

$$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-(\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1}) \\ * \text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] \text{ ; FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \& \\ \& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$$

rule 748

$$\text{Int}[(a_ + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Simp}[x*((a + b*x^n)^p/(n*p \\ + 1)), x] + \text{Simp}[a*n*(p/(n*p + 1)) \quad \text{Int}[(a + b*x^n)^{(p - 1)}, x], x] \text{ ; Fre} \\ \text{eQ}[\{a, b\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ (\text{IntegerQ}[2*p] \ || \ \text{LtQ}[\text{Denominat} \\ \text{or}[p + 1/n], \text{Denominator}[p]])$$

rule 755

$$\text{Int}[(a_ + (b_)*(x_)^4)^{-1}, x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2] \\], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2*r) \quad \text{Int}[(r - s*x^2)/(a + b*x^4) \\ , x], x] + \text{Simp}[1/(2*r) \quad \text{Int}[(r + s*x^2)/(a + b*x^4), x], x]] \text{ ; FreeQ}[\{a, \\ b\}, x] \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \& \\ \& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$$

rule 770

$$\text{Int}[(a_ + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Simp}[a^{(p + 1/n)} \quad \text{Subst}[\text{In} \\ \text{t}[1/(1 - b*x^n)^{(p + 1/n + 1)}, x], x, x/(a + b*x^n)^{(1/n)}, x] \text{ ; FreeQ}[\{a, \\ b\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[-1, p, 0] \ \&\& \ \text{NeQ}[p, -2^{(-1)}] \ \&\& \ \text{IntegerQ}[p + 1 \\ /n]$$

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

Maple [A] (verified)

Time = 3.98 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.03

method	result
pseudoelliptic	$\frac{3 \left(\arctan \left(\frac{\sqrt{2}(-bx^4+a)^{\frac{1}{4}} - b^{\frac{1}{4}}x}{b^{\frac{1}{4}}x} \right) \sqrt{2}a + \arctan \left(\frac{\sqrt{2}(-bx^4+a)^{\frac{1}{4}} + b^{\frac{1}{4}}x}{b^{\frac{1}{4}}x} \right) \sqrt{2}a + \frac{\ln \left(\frac{-b^{\frac{1}{4}}(-bx^4+a)^{\frac{1}{4}} \sqrt{2}x + \sqrt{b}x^2 + \sqrt{-bx^4+a}}{b^{\frac{1}{4}}(-bx^4+a)^{\frac{1}{4}} \sqrt{2}x + \sqrt{b}x^2 + \sqrt{-bx^4+a}} \right)}{2}}{16b^{\frac{1}{4}}}$

input `int((-b*x^4+a)^(3/4), x, method=_RETURNVERBOSE)`

output

```
-3/16*(arctan((2^(1/2)*(-b*x^4+a)^(1/4)-b^(1/4)*x)/b^(1/4)/x)*2^(1/2)*a+arctan((2^(1/2)*(-b*x^4+a)^(1/4)+b^(1/4)*x)/b^(1/4)/x)*2^(1/2)*a+1/2*ln((-b^(1/4)*(-b*x^4+a)^(1/4)*2^(1/2)*x+b^(1/2)*x^2+(-b*x^4+a)^(1/2))/b^(1/4)*(-b*x^4+a)^(1/4)*2^(1/2)*x+b^(1/2)*x^2+(-b*x^4+a)^(1/2)))*2^(1/2)*a-4/3*(-b*x^4+a)^(3/4)*x*b^(1/4)/b^(1/4)
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.20

$$\int (a - bx^4)^{3/4} dx = \frac{1}{4} (-bx^4 + a)^{3/4} x - \frac{3}{16} \left(-\frac{a^4}{b} \right)^{1/4} \log \left(\frac{27 \left((-bx^4 + a)^{1/4} a^3 + \left(-\frac{a^4}{b} \right)^{3/4} bx \right)}{x} \right) + \frac{3}{16} i \left(-\frac{a^4}{b} \right)^{1/4} \log \left(\frac{27 \left((-bx^4 + a)^{1/4} a^3 + i \left(-\frac{a^4}{b} \right)^{3/4} bx \right)}{x} \right) - \frac{3}{16} i \left(-\frac{a^4}{b} \right)^{1/4} \log \left(\frac{27 \left((-bx^4 + a)^{1/4} a^3 - i \left(-\frac{a^4}{b} \right)^{3/4} bx \right)}{x} \right) + \frac{3}{16} \left(-\frac{a^4}{b} \right)^{1/4} \log \left(\frac{27 \left((-bx^4 + a)^{1/4} a^3 - \left(-\frac{a^4}{b} \right)^{3/4} bx \right)}{x} \right)$$

input

```
integrate((-b*x^4+a)^(3/4),x, algorithm="fricas")
```

output

```
1/4*(-b*x^4 + a)^(3/4)*x - 3/16*(-a^4/b)^(1/4)*log(27*((-b*x^4 + a)^(1/4)*a^3 + (-a^4/b)^(3/4)*b*x)/x) + 3/16*I*(-a^4/b)^(1/4)*log(27*((-b*x^4 + a)^(1/4)*a^3 + I*(-a^4/b)^(3/4)*b*x)/x) - 3/16*I*(-a^4/b)^(1/4)*log(27*((-b*x^4 + a)^(1/4)*a^3 - I*(-a^4/b)^(3/4)*b*x)/x) + 3/16*(-a^4/b)^(1/4)*log(27*((-b*x^4 + a)^(1/4)*a^3 - (-a^4/b)^(3/4)*b*x)/x)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.75 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.23

$$\int (a - bx^4)^{3/4} dx = \frac{a^{3/4} x \Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\begin{matrix} -\frac{3}{4}, \frac{1}{4} \\ \frac{5}{4} \end{matrix} \middle| \frac{bx^4 e^{2i\pi}}{a}\right)}{4\Gamma\left(\frac{5}{4}\right)}$$

input `integrate((-b*x**4+a)**(3/4),x)`

output `a**(3/4)*x*gamma(1/4)*hyper((-3/4, 1/4), (5/4,), b*x**4*exp_polar(2*I*pi)/a)/(4*gamma(5/4))`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.27

$$\int (a - bx^4)^{3/4} dx =$$

$$-\frac{3}{32} \left(\frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}b^{1/4} + \frac{2(-bx^4+a)^{1/4}}{x}\right)}{2b^{1/4}}\right)}{b^{1/4}} + \frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}b^{1/4} - \frac{2(-bx^4+a)^{1/4}}{x}\right)}{2b^{1/4}}\right)}{b^{1/4}} - \sqrt{2} \log\left(\sqrt{b} + \right.$$

$$\left. + \frac{(-bx^4+a)^{3/4} a}{4\left(b - \frac{bx^4-a}{x^4}\right)x^3} \right)$$

input `integrate((-b*x^4+a)^(3/4),x, algorithm="maxima")`

output

```
-3/32*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*b^(1/4) + 2*(-b*x^4 + a)^(1/4)
)/x)/b^(1/4))/b^(1/4) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*b^(1/4) - 2
*(-b*x^4 + a)^(1/4)/x)/b^(1/4))/b^(1/4) - sqrt(2)*log(sqrt(b) + sqrt(2)*(-
b*x^4 + a)^(1/4)*b^(1/4)/x + sqrt(-b*x^4 + a)/x^2)/b^(1/4) + sqrt(2)*log(s
qrt(b) - sqrt(2)*(-b*x^4 + a)^(1/4)*b^(1/4)/x + sqrt(-b*x^4 + a)/x^2)/b^(1
/4))*a + 1/4*(-b*x^4 + a)^(3/4)*a/((b - (b*x^4 - a)/x^4)*x^3)
```

Giac [F]

$$\int (a - bx^4)^{3/4} dx = \int (-bx^4 + a)^{3/4} dx$$

input

```
integrate((-b*x^4+a)^(3/4),x, algorithm="giac")
```

output

```
integrate((-b*x^4 + a)^(3/4), x)
```

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.23

$$\int (a - bx^4)^{3/4} dx = \frac{x(a - bx^4)^{3/4} {}_2F_1\left(-\frac{3}{4}, \frac{1}{4}; \frac{5}{4}; \frac{bx^4}{a}\right)}{\left(1 - \frac{bx^4}{a}\right)^{3/4}}$$

input

```
int((a - b*x^4)^(3/4),x)
```

output

```
(x*(a - b*x^4)^(3/4)*hypergeom([-3/4, 1/4], 5/4, (b*x^4)/a))/(1 - (b*x^4)/
a)^(3/4)
```

Reduce [F]

$$\int (a - bx^4)^{3/4} dx = \frac{(-bx^4 + a)^{3/4} x}{4} + \frac{3 \left(\int \frac{1}{(-bx^4 + a)^{1/4}} dx \right) a}{4}$$

input `int((-b*x^4+a)^(3/4),x)`

output `((a - b*x**4)**(3/4)*x + 3*int((a - b*x**4)**(3/4)/(a - b*x**4),x)*a)/4`

3.46 $\int \frac{1}{\sqrt[4]{a - bx^4}} dx$

Optimal result	322
Mathematica [A] (verified)	323
Rubi [A] (verified)	323
Maple [A] (verified)	327
Fricas [C] (verification not implemented)	328
Sympy [C] (verification not implemented)	328
Maxima [A] (verification not implemented)	329
Giac [F]	330
Mupad [B] (verification not implemented)	330
Reduce [F]	330

Optimal result

Integrand size = 12, antiderivative size = 148

$$\int \frac{1}{\sqrt[4]{a - bx^4}} dx = -\frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a - bx^4}}\right)}{2\sqrt{2}\sqrt[4]{b}} + \frac{\arctan\left(1 + \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a - bx^4}}\right)}{2\sqrt{2}\sqrt[4]{b}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a - bx^4}\left(1 + \frac{\sqrt{bx^2}}{\sqrt[4]{a - bx^4}}\right)}\right)}{2\sqrt{2}\sqrt[4]{b}}$$

output

```
1/4*arctan(-1+2^(1/2)*b^(1/4)*x/(-b*x^4+a)^(1/4))*2^(1/2)/b^(1/4)+1/4*arctan(1+2^(1/2)*b^(1/4)*x/(-b*x^4+a)^(1/4))*2^(1/2)/b^(1/4)+1/4*arctanh(2^(1/2)*b^(1/4)*x/(-b*x^4+a)^(1/4)/(1+b^(1/2)*x^2/(-b*x^4+a)^(1/2)))*2^(1/2)/b^(1/4)
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.17

$$\int \frac{1}{\sqrt[4]{a-bx^4}} dx$$

$$= \frac{-2 \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a-bx^4}}\right) + 2 \arctan\left(1 + \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a-bx^4}}\right) - \log\left(1 + \frac{\sqrt{bx^2}}{\sqrt{a-bx^4}} - \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a-bx^4}}\right) + \log\left(1 + \frac{\sqrt{bx^2}}{\sqrt{a-bx^4}} + \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a-bx^4}}\right)}{4\sqrt{2}\sqrt[4]{b}}$$

input `Integrate[(a - b*x^4)^(-1/4), x]`output `(-2*ArcTan[1 - (Sqrt[2]*b^(1/4)*x)/(a - b*x^4)^(1/4)] + 2*ArcTan[1 + (Sqrt[2]*b^(1/4)*x)/(a - b*x^4)^(1/4)] - Log[1 + (Sqrt[b]*x^2)/Sqrt[a - b*x^4] - (Sqrt[2]*b^(1/4)*x)/(a - b*x^4)^(1/4)] + Log[1 + (Sqrt[b]*x^2)/Sqrt[a - b*x^4] + (Sqrt[2]*b^(1/4)*x)/(a - b*x^4)^(1/4)])/(4*Sqrt[2]*b^(1/4))`**Rubi [A] (verified)**Time = 0.51 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.45, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {770, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt[4]{a-bx^4}} dx$$

$$\downarrow 770$$

$$\int \frac{1}{\frac{bx^4}{a-bx^4} + 1} d \frac{x}{\sqrt[4]{a-bx^4}}$$

$$\downarrow 755$$

$$\frac{1}{2} \int \frac{1 - \frac{\sqrt{bx^2}}{\sqrt{a-bx^4}}}{\frac{bx^4}{a-bx^4} + 1} d \frac{x}{\sqrt[4]{a-bx^4}} + \frac{1}{2} \int \frac{\frac{\sqrt{bx^2}}{\sqrt{a-bx^4}} + 1}{\frac{bx^4}{a-bx^4} + 1} d \frac{x}{\sqrt[4]{a-bx^4}}$$

$$\begin{aligned} & \downarrow 1476 \\ & \frac{1}{2} \int \frac{1 - \frac{\sqrt{bx^2}}{\sqrt{a-bx^4}}}{\frac{bx^4}{a-bx^4} + 1} d \frac{x}{\sqrt[4]{a-bx^4}} + \\ & \frac{1}{2} \left(\frac{\int \frac{x^2}{\sqrt{a-bx^4}} - \frac{1}{\sqrt[4]{b} \sqrt[4]{a-bx^4}} + \frac{1}{\sqrt{b}} d \frac{x}{\sqrt[4]{a-bx^4}}}{2\sqrt{b}} + \frac{\int \frac{x^2}{\sqrt{a-bx^4}} + \frac{1}{\sqrt[4]{b} \sqrt[4]{a-bx^4}} + \frac{1}{\sqrt{b}} d \frac{x}{\sqrt[4]{a-bx^4}}}{2\sqrt{b}} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 1082 \\ & \frac{1}{2} \left(\frac{\int \frac{1}{-\frac{x^2}{\sqrt{a-bx^4}} - 1} d \left(1 - \frac{\sqrt{2} \sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} \right)}{\sqrt{2} \sqrt[4]{b}} - \frac{\int \frac{1}{-\frac{x^2}{\sqrt{a-bx^4}} - 1} d \left(\frac{\sqrt{2} \sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} + 1 \right)}{\sqrt{2} \sqrt[4]{b}} \right) + \end{aligned}$$

$$\frac{1}{2} \int \frac{1 - \frac{\sqrt{bx^2}}{\sqrt{a-bx^4}}}{\frac{bx^4}{a-bx^4} + 1} d \frac{x}{\sqrt[4]{a-bx^4}}$$

$$\begin{aligned} & \downarrow 217 \\ & \frac{1}{2} \int \frac{1 - \frac{\sqrt{bx^2}}{\sqrt{a-bx^4}}}{\frac{bx^4}{a-bx^4} + 1} d \frac{x}{\sqrt[4]{a-bx^4}} + \frac{1}{2} \left(\frac{\arctan \left(\frac{\sqrt{2} \sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} + 1 \right)}{\sqrt{2} \sqrt[4]{b}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} \right)}{\sqrt{2} \sqrt[4]{b}} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 1479 \\ & \frac{1}{2} \left(\frac{\int -\frac{\sqrt{2} - \frac{2 \sqrt[4]{bx}}{\sqrt[4]{a-bx^4}}}{\sqrt[4]{b} \left(\frac{x^2}{\sqrt{a-bx^4}} - \frac{\sqrt{2x}}{\sqrt[4]{b} \sqrt[4]{a-bx^4}} + \frac{1}{\sqrt{b}} \right)} d \frac{x}{\sqrt[4]{a-bx^4}}}{2\sqrt{2} \sqrt[4]{b}} - \frac{\int -\frac{\sqrt{2} \left(\frac{\sqrt{2} \sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} + 1 \right)}{\sqrt[4]{b} \left(\frac{x^2}{\sqrt{a-bx^4}} + \frac{\sqrt{2x}}{\sqrt[4]{b} \sqrt[4]{a-bx^4}} + \frac{1}{\sqrt{b}} \right)} d \frac{x}{\sqrt[4]{a-bx^4}}}{2\sqrt{2} \sqrt[4]{b}} \right) + \\ & \frac{1}{2} \left(\frac{\arctan \left(\frac{\sqrt{2} \sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} + 1 \right)}{\sqrt{2} \sqrt[4]{b}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} \right)}{\sqrt{2} \sqrt[4]{b}} \right) \end{aligned}$$

25

$$\begin{aligned}
& \frac{1}{2} \left(\frac{\int \frac{\sqrt{2} - \frac{2\sqrt[4]{bx}}{\sqrt[4]{a-bx^4}}}{\sqrt[4]{b} \left(\frac{x^2}{\sqrt{a-bx^4}} - \frac{\sqrt{2x}}{\sqrt[4]{b} \sqrt[4]{a-bx^4}} + \frac{1}{\sqrt[4]{b}} \right)} d \frac{x}{\sqrt[4]{a-bx^4}}}{2\sqrt{2}\sqrt[4]{b}} + \frac{\int \frac{\sqrt{2} \left(\frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} + 1 \right)}{\sqrt[4]{b} \left(\frac{x^2}{\sqrt{a-bx^4}} + \frac{\sqrt{2x}}{\sqrt[4]{b} \sqrt[4]{a-bx^4}} + \frac{1}{\sqrt[4]{b}} \right)} d \frac{x}{\sqrt[4]{a-bx^4}}}{2\sqrt{2}\sqrt[4]{b}} \right) + \\
& \frac{1}{2} \left(\frac{\arctan \left(\frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} + 1 \right)}{\sqrt{2}\sqrt[4]{b}} - \frac{\arctan \left(1 - \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} \right)}{\sqrt{2}\sqrt[4]{b}} \right) \\
& \quad \downarrow 27 \\
& \frac{1}{2} \left(\frac{\int \frac{\sqrt{2} - \frac{2\sqrt[4]{bx}}{\sqrt[4]{a-bx^4}}}{\frac{x^2}{\sqrt{a-bx^4}} - \frac{\sqrt{2x}}{\sqrt[4]{b} \sqrt[4]{a-bx^4}} + \frac{1}{\sqrt[4]{b}}} d \frac{x}{\sqrt[4]{a-bx^4}}}{2\sqrt{2}\sqrt[4]{b}} + \frac{\int \frac{\frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} + 1}{\frac{x^2}{\sqrt{a-bx^4}} + \frac{\sqrt{2x}}{\sqrt[4]{b} \sqrt[4]{a-bx^4}} + \frac{1}{\sqrt[4]{b}}} d \frac{x}{\sqrt[4]{a-bx^4}}}{2\sqrt[4]{b}} \right) + \\
& \frac{1}{2} \left(\frac{\arctan \left(\frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} + 1 \right)}{\sqrt{2}\sqrt[4]{b}} - \frac{\arctan \left(1 - \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} \right)}{\sqrt{2}\sqrt[4]{b}} \right) \\
& \quad \downarrow 1103 \\
& \frac{1}{2} \left(\frac{\arctan \left(\frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} + 1 \right)}{\sqrt{2}\sqrt[4]{b}} - \frac{\arctan \left(1 - \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} \right)}{\sqrt{2}\sqrt[4]{b}} \right) + \\
& \frac{1}{2} \left(\frac{\log \left(\frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} + \frac{\sqrt{bx^2}}{\sqrt{a-bx^4}} + 1 \right)}{2\sqrt{2}\sqrt[4]{b}} - \frac{\log \left(-\frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a-bx^4}} + \frac{\sqrt{bx^2}}{\sqrt{a-bx^4}} + 1 \right)}{2\sqrt{2}\sqrt[4]{b}} \right)
\end{aligned}$$

input

Int[(a - b*x^4)^(-1/4), x]

output

$$\begin{aligned} & (-\text{ArcTan}[1 - (\text{Sqrt}[2]*b^{(1/4)*x})/(a - b*x^4)^{(1/4)}]/(\text{Sqrt}[2]*b^{(1/4)}) + \\ & \text{ArcTan}[1 + (\text{Sqrt}[2]*b^{(1/4)*x})/(a - b*x^4)^{(1/4)}]/(\text{Sqrt}[2]*b^{(1/4)}))/2 + (\\ & -1/2*\text{Log}[1 + (\text{Sqrt}[b]*x^2)/\text{Sqrt}[a - b*x^4] - (\text{Sqrt}[2]*b^{(1/4)*x})/(a - b*x^4)^{(1/4)}]/(\text{Sqrt}[2]*b^{(1/4)}) + \text{Log}[1 + (\text{Sqrt}[b]*x^2)/\text{Sqrt}[a - b*x^4] + (\text{Sqrt}[2]*b^{(1/4)*x})/(a - b*x^4)^{(1/4)}]/(2*\text{Sqrt}[2]*b^{(1/4)})/2 \end{aligned}$$
Defintions of rubi rules used

rule 25

$$\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \text{ :> } \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$$

rule 27

$$\text{Int}[(a_)*(\text{Fx}_), \text{x_Symbol}] \text{ :> } \text{Simp}[a \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[a, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (b_)*(Gx_)] \text{ ; FreeQ}[b, \text{x}]$$

rule 217

$$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, \text{x_Symbol}] \text{ :> } \text{Simp}[(-(\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], \text{x}] \text{ ; FreeQ}[\{a, b\}, \text{x}] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$$

rule 755

$$\text{Int}[(a_ + (b_)*(x_)^4)^{-1}, \text{x_Symbol}] \text{ :> } \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2*r) \quad \text{Int}[(r - s*x^2)/(a + b*x^4), \text{x}], \text{x}] + \text{Simp}[1/(2*r) \quad \text{Int}[(r + s*x^2)/(a + b*x^4), \text{x}], \text{x}]] \text{ ; FreeQ}[\{a, b\}, \text{x}] \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$$

rule 770

$$\text{Int}[(a_ + (b_)*(x_)^{(n_)})^{(p_)}, \text{x_Symbol}] \text{ :> } \text{Simp}[a^{(p + 1/n)} \quad \text{Subst}[\text{Int}[1/(1 - b*x^n)^{(p + 1/n + 1)}, \text{x}], \text{x}, x/(a + b*x^n)^{(1/n)}], \text{x}] \text{ ; FreeQ}[\{a, b\}, \text{x}] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[-1, p, 0] \ \&\& \ \text{NeQ}[p, -2^{(-1)}] \ \&\& \ \text{IntegerQ}[p + 1/n]$$

rule 1082

$$\text{Int}[(a_ + (b_)*(x_) + (c_)*(x_)^2)^{-1}, \text{x_Symbol}] \text{ :> } \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \quad \text{Subst}[\text{Int}[1/(q - x^2), \text{x}], \text{x}, 1 + 2*c*(x/b)], \text{x}] \text{ ; RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ \text{!RationalQ}[b^2 - 4*a*c])] \text{ ; FreeQ}[\{a, b, c\}, \text{x}]$$

rule 1103 $\text{Int}[\frac{(d_.) + (e_.)x}{(a_.) + (b_.)x + (c_.)x^2}, x_Symbol] \rightarrow \text{Simp}[d \cdot \text{Log}[\text{RemoveContent}[a + bx + cx^2, x]]/b, x] /; \text{FreeQ}\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2cd - b^2e, 0]$

rule 1476 $\text{Int}[\frac{(d_.) + (e_.)x^2}{(a_.) + (c_.)x^4}, x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[2(d/e), 2]\}, \text{Simp}[e/(2c) \text{Int}[1/\text{Simp}[d/e + qx + x^2, x], x], x] + \text{Simp}[e/(2c) \text{Int}[1/\text{Simp}[d/e - qx + x^2, x], x], x]] /; \text{FreeQ}\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c^2d - a^2e, 0] \ \&\& \ \text{PosQ}[d^2e]$

rule 1479 $\text{Int}[\frac{(d_.) + (e_.)x^2}{(a_.) + (c_.)x^4}, x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[-2(d/e), 2]\}, \text{Simp}[e/(2cq) \text{Int}[(q - 2x)/\text{Simp}[d/e + qx - x^2, x], x], x] + \text{Simp}[e/(2cq) \text{Int}[(q + 2x)/\text{Simp}[d/e - qx - x^2, x], x], x]] /; \text{FreeQ}\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c^2d - a^2e, 0] \ \&\& \ \text{NegQ}[d^2e]$

Maple [A] (verified)

Time = 0.78 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.93

method	result
pseudoelliptic	$-\frac{\sqrt{2} \left(\ln \left(\frac{-b^{\frac{1}{4}}(-bx^4+a)^{\frac{1}{4}}\sqrt{2x+\sqrt{bx^2+\sqrt{-bx^4+a}}}}{b^{\frac{1}{4}}(-bx^4+a)^{\frac{1}{4}}\sqrt{2x+\sqrt{bx^2+\sqrt{-bx^4+a}}}} \right) + 2 \arctan \left(\frac{\sqrt{2}(-bx^4+a)^{\frac{1}{4}}}{b^{\frac{1}{4}}x} + 1 \right) - 2 \arctan \left(-\frac{\sqrt{2}(-bx^4+a)^{\frac{1}{4}}}{b^{\frac{1}{4}}x} + 1 \right) \right)}{8b^{\frac{1}{4}}}$

input `int(1/(-b*x^4+a)^(1/4),x,method=_RETURNVERBOSE)`

output
$$-1/8/b^{1/4} \cdot 2^{1/2} \cdot (\ln((-b^{1/4} \cdot (-bx^4+a)^{1/4} \cdot 2^{1/2} \cdot x + b^{1/2} \cdot x^2 + (-bx^4+a)^{1/2})) / (b^{1/4} \cdot (-bx^4+a)^{1/4} \cdot 2^{1/2} \cdot x + b^{1/2} \cdot x^2 + (-bx^4+a)^{1/2})) + 2 \cdot \arctan(2^{1/2} / b^{1/4} \cdot (-bx^4+a)^{1/4} / x + 1) - 2 \cdot \arctan(-2^{1/2} / b^{1/4} \cdot (-bx^4+a)^{1/4} / x + 1))$$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.99

$$\int \frac{1}{\sqrt[4]{a-bx^4}} dx = -\frac{1}{4} \left(-\frac{1}{b}\right)^{\frac{1}{4}} \log\left(\frac{bx\left(-\frac{1}{b}\right)^{\frac{3}{4}} + (-bx^4 + a)^{\frac{1}{4}}}{x}\right) \\ + \frac{1}{4} \left(-\frac{1}{b}\right)^{\frac{1}{4}} \log\left(-\frac{bx\left(-\frac{1}{b}\right)^{\frac{3}{4}} - (-bx^4 + a)^{\frac{1}{4}}}{x}\right) \\ + \frac{1}{4} i \left(-\frac{1}{b}\right)^{\frac{1}{4}} \log\left(\frac{ibx\left(-\frac{1}{b}\right)^{\frac{3}{4}} + (-bx^4 + a)^{\frac{1}{4}}}{x}\right) \\ - \frac{1}{4} i \left(-\frac{1}{b}\right)^{\frac{1}{4}} \log\left(\frac{-ibx\left(-\frac{1}{b}\right)^{\frac{3}{4}} + (-bx^4 + a)^{\frac{1}{4}}}{x}\right)$$

input `integrate(1/(-b*x^4+a)^(1/4),x, algorithm="fricas")`

output `-1/4*(-1/b)^(1/4)*log((b*x*(-1/b)^(3/4) + (-b*x^4 + a)^(1/4))/x) + 1/4*(-1/b)^(1/4)*log(-b*x*(-1/b)^(3/4) - (-b*x^4 + a)^(1/4))/x + 1/4*I*(-1/b)^(1/4)*log((I*b*x*(-1/b)^(3/4) + (-b*x^4 + a)^(1/4))/x) - 1/4*I*(-1/b)^(1/4)*log((-I*b*x*(-1/b)^(3/4) + (-b*x^4 + a)^(1/4))/x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.58 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.25

$$\int \frac{1}{\sqrt[4]{a-bx^4}} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{4} \middle| \frac{5}{4} \middle| \frac{bx^4 e^{2i\pi}}{a}\right)}{4\sqrt[4]{a}\Gamma\left(\frac{5}{4}\right)}$$

input `integrate(1/(-b*x**4+a)**(1/4),x)`

output `x*gamma(1/4)*hyper((1/4, 1/4), (5/4,), b*x**4*exp_polar(2*I*pi)/a)/(4*a**(1/4)*gamma(5/4))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.19

$$\int \frac{1}{\sqrt[4]{a-bx^4}} dx = -\frac{\sqrt{2} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}b^{\frac{1}{4}} + \frac{2(-bx^4+a)^{\frac{1}{4}}}{x}\right)}{2b^{\frac{1}{4}}}\right)}{4b^{\frac{1}{4}}} - \frac{\sqrt{2} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}b^{\frac{1}{4}} - \frac{2(-bx^4+a)^{\frac{1}{4}}}{x}\right)}{2b^{\frac{1}{4}}}\right)}{4b^{\frac{1}{4}}} + \frac{\sqrt{2} \log\left(\sqrt{b} + \frac{\sqrt{2}(-bx^4+a)^{\frac{1}{4}}b^{\frac{1}{4}}}{x} + \frac{\sqrt{-bx^4+a}}{x^2}\right)}{8b^{\frac{1}{4}}} - \frac{\sqrt{2} \log\left(\sqrt{b} - \frac{\sqrt{2}(-bx^4+a)^{\frac{1}{4}}b^{\frac{1}{4}}}{x} + \frac{\sqrt{-bx^4+a}}{x^2}\right)}{8b^{\frac{1}{4}}}$$

input `integrate(1/(-b*x^4+a)^(1/4),x, algorithm="maxima")`

output `-1/4*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*b^(1/4) + 2*(-b*x^4 + a)^(1/4)/x)/b^(1/4))/b^(1/4) - 1/4*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*b^(1/4) - 2*(-b*x^4 + a)^(1/4)/x)/b^(1/4))/b^(1/4) + 1/8*sqrt(2)*log(sqrt(b) + sqrt(2)*(-b*x^4 + a)^(1/4)*b^(1/4)/x + sqrt(-b*x^4 + a)/x^2)/b^(1/4) - 1/8*sqrt(2)*log(sqrt(b) - sqrt(2)*(-b*x^4 + a)^(1/4)*b^(1/4)/x + sqrt(-b*x^4 + a)/x^2)/b^(1/4)`

Giac [F]

$$\int \frac{1}{\sqrt[4]{a - bx^4}} dx = \int \frac{1}{(-bx^4 + a)^{\frac{1}{4}}} dx$$

input `integrate(1/(-b*x^4+a)^(1/4),x, algorithm="giac")`

output `integrate((-b*x^4 + a)^(-1/4), x)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.26

$$\int \frac{1}{\sqrt[4]{a - bx^4}} dx = \frac{x \left(1 - \frac{bx^4}{a}\right)^{1/4} {}_2F_1\left(\frac{1}{4}, \frac{1}{4}, \frac{5}{4}, \frac{bx^4}{a}\right)}{(a - bx^4)^{1/4}}$$

input `int(1/(a - b*x^4)^(1/4),x)`

output `(x*(1 - (b*x^4)/a)^(1/4)*hypergeom([1/4, 1/4], 5/4, (b*x^4)/a))/(a - b*x^4)^(1/4)`

Reduce [F]

$$\int \frac{1}{\sqrt[4]{a - bx^4}} dx = \int \frac{1}{(-bx^4 + a)^{\frac{1}{4}}} dx$$

input `int(1/(-b*x^4+a)^(1/4),x)`

output `int(1/(a - b*x**4)**(1/4),x)`

$$3.47 \quad \int \frac{1}{(a-bx^4)^{5/4}} dx$$

Optimal result	331
Mathematica [A] (verified)	331
Rubi [A] (verified)	332
Maple [A] (verified)	332
Fricas [A] (verification not implemented)	333
Sympy [C] (verification not implemented)	333
Maxima [A] (verification not implemented)	334
Giac [F]	334
Mupad [B] (verification not implemented)	334
Reduce [F]	335

Optimal result

Integrand size = 12, antiderivative size = 17

$$\int \frac{1}{(a-bx^4)^{5/4}} dx = \frac{x}{a\sqrt[4]{a-bx^4}}$$

output `x/a/(-b*x^4+a)^(1/4)`

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a-bx^4)^{5/4}} dx = \frac{x}{a\sqrt[4]{a-bx^4}}$$

input `Integrate[(a - b*x^4)^(-5/4),x]`

output `x/(a*(a - b*x^4)^(1/4))`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {746}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a - bx^4)^{5/4}} dx$$

$$\downarrow \text{746}$$

$$\frac{x}{a \sqrt[4]{a - bx^4}}$$

input `Int[(a - b*x^4)^(-5/4), x]`

output `x/(a*(a - b*x^4)^(1/4))`

Defintions of rubi rules used

rule 746 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[x*((a + b*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]`

Maple [A] (verified)

Time = 0.51 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

method	result	size
gosper	$\frac{x}{a(-bx^4+a)^{\frac{1}{4}}}$	16
trager	$\frac{x}{a(-bx^4+a)^{\frac{1}{4}}}$	16
pseudoelliptic	$\frac{x}{a(-bx^4+a)^{\frac{1}{4}}}$	16
orering	$\frac{x}{a(-bx^4+a)^{\frac{1}{4}}}$	16

input `int(1/(-b*x^4+a)^(5/4),x,method=_RETURNVERBOSE)`

output `x/a/(-b*x^4+a)^(1/4)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \frac{1}{(a - bx^4)^{5/4}} dx = -\frac{(-bx^4 + a)^{3/4} x}{abx^4 - a^2}$$

input `integrate(1/(-b*x^4+a)^(5/4),x, algorithm="fricas")`

output `-(-b*x^4 + a)^(3/4)*x/(a*b*x^4 - a^2)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.46 (sec) , antiderivative size = 73, normalized size of antiderivative = 4.29

$$\int \frac{1}{(a - bx^4)^{5/4}} dx = \begin{cases} \frac{xe^{-\frac{i\pi}{4}}\Gamma(\frac{1}{4})}{4a^{\frac{5}{4}}\sqrt[4]{-1 + \frac{bx^4}{a}}\Gamma(\frac{5}{4})} & \text{for } \left|\frac{bx^4}{a}\right| > 1 \\ \frac{x\Gamma(\frac{1}{4})}{4a^{\frac{5}{4}}\sqrt[4]{1 - \frac{bx^4}{a}}\Gamma(\frac{5}{4})} & \text{otherwise} \end{cases}$$

input `integrate(1/(-b*x**4+a)**(5/4),x)`

output `Piecewise((x*exp(-I*pi/4)*gamma(1/4)/(4*a**(5/4)*(-1 + b*x**4/a)**(1/4)*gamma(5/4)), Abs(b*x**4/a) > 1), (x*gamma(1/4)/(4*a**(5/4)*(1 - b*x**4/a)**(1/4)*gamma(5/4)), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{1}{(a - bx^4)^{5/4}} dx = \frac{x}{(-bx^4 + a)^{1/4} a}$$

input `integrate(1/(-b*x^4+a)^(5/4),x, algorithm="maxima")`output `x/((-b*x^4 + a)^(1/4)*a)`**Giac [F]**

$$\int \frac{1}{(a - bx^4)^{5/4}} dx = \int \frac{1}{(-bx^4 + a)^{5/4}} dx$$

input `integrate(1/(-b*x^4+a)^(5/4),x, algorithm="giac")`output `integrate((-b*x^4 + a)^(-5/4), x)`**Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{1}{(a - bx^4)^{5/4}} dx = \frac{x}{a(a - bx^4)^{1/4}}$$

input `int(1/(a - b*x^4)^(5/4),x)`output `x/(a*(a - b*x^4)^(1/4))`

Reduce [F]

$$\int \frac{1}{(a - bx^4)^{5/4}} dx = \int \frac{1}{(-bx^4 + a)^{1/4} a - (-bx^4 + a)^{1/4} bx^4} dx$$

input `int(1/(-b*x^4+a)^(5/4),x)`

output `int(1/((a - b*x**4)**(1/4)*a - (a - b*x**4)**(1/4)*b*x**4),x)`

3.48 $\int \frac{1}{(a-bx^4)^{9/4}} dx$

Optimal result	336
Mathematica [A] (verified)	336
Rubi [A] (verified)	337
Maple [A] (verified)	338
Fricas [A] (verification not implemented)	338
Sympy [C] (verification not implemented)	339
Maxima [A] (verification not implemented)	339
Giac [F]	340
Mupad [B] (verification not implemented)	340
Reduce [F]	340

Optimal result

Integrand size = 12, antiderivative size = 41

$$\int \frac{1}{(a-bx^4)^{9/4}} dx = \frac{x}{5a(a-bx^4)^{5/4}} + \frac{4x}{5a^2\sqrt[4]{a-bx^4}}$$

output

```
1/5*x/a/(-b*x^4+a)^(5/4)+4/5*x/a^2/(-b*x^4+a)^(1/4)
```

Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.73

$$\int \frac{1}{(a-bx^4)^{9/4}} dx = \frac{5ax - 4bx^5}{5a^2(a-bx^4)^{5/4}}$$

input

```
Integrate[(a - b*x^4)^(-9/4),x]
```

output

```
(5*a*x - 4*b*x^5)/(5*a^2*(a - b*x^4)^(5/4))
```

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {749, 746}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a - bx^4)^{9/4}} dx$$

$$\downarrow 749$$

$$\frac{4 \int \frac{1}{(a - bx^4)^{5/4}} dx}{5a} + \frac{x}{5a(a - bx^4)^{5/4}}$$

$$\downarrow 746$$

$$\frac{4x}{5a^2 \sqrt[4]{a - bx^4}} + \frac{x}{5a(a - bx^4)^{5/4}}$$

input `Int[(a - b*x^4)^(-9/4), x]`

output `x/(5*a*(a - b*x^4)^(5/4)) + (4*x)/(5*a^2*(a - b*x^4)^(1/4))`

Defintions of rubi rules used

rule 746 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]`

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.66

method	result	size
gospers	$\frac{x(-4bx^4+5a)}{5(-bx^4+a)^{\frac{5}{4}}a^2}$	27
trager	$\frac{x(-4bx^4+5a)}{5(-bx^4+a)^{\frac{5}{4}}a^2}$	27
pseudoelliptic	$\frac{x(-4bx^4+5a)}{5(-bx^4+a)^{\frac{5}{4}}a^2}$	27
orering	$\frac{x(-4bx^4+5a)}{5(-bx^4+a)^{\frac{5}{4}}a^2}$	27

input `int(1/(-b*x^4+a)^(9/4),x,method=_RETURNVERBOSE)`output `1/5*x*(-4*b*x^4+5*a)/(-b*x^4+a)^(5/4)/a^2`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.17

$$\int \frac{1}{(a-bx^4)^{9/4}} dx = -\frac{(4bx^5-5ax)(-bx^4+a)^{3/4}}{5(a^2b^2x^8-2a^3bx^4+a^4)}$$

input `integrate(1/(-b*x^4+a)^(9/4),x, algorithm="fricas")`output `-1/5*(4*b*x^5 - 5*a*x)*(-b*x^4 + a)^(3/4)/(a^2*b^2*x^8 - 2*a^3*b*x^4 + a^4)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.71 (sec) , antiderivative size = 274, normalized size of antiderivative = 6.68

$$\int \frac{1}{(a - bx^4)^{9/4}} dx = \begin{cases} -\frac{5axe^{-\frac{i\pi}{4}}\Gamma(\frac{1}{4})}{-16a^{\frac{13}{4}}\sqrt[4]{-1 + \frac{bx^4}{a}}\Gamma(\frac{9}{4}) + 16a^{\frac{9}{4}}bx^4\sqrt[4]{-1 + \frac{bx^4}{a}}\Gamma(\frac{9}{4})} + \frac{4bx^5e^{-\frac{i\pi}{4}}\Gamma(\frac{1}{4})}{-16a^{\frac{13}{4}}\sqrt[4]{-1 + \frac{bx^4}{a}}\Gamma(\frac{9}{4}) + 16a^{\frac{9}{4}}bx^4\sqrt[4]{-1 + \frac{bx^4}{a}}\Gamma(\frac{9}{4})} \\ -\frac{5ax\Gamma(\frac{1}{4})}{-16a^{\frac{13}{4}}\sqrt[4]{1 - \frac{bx^4}{a}}\Gamma(\frac{9}{4}) + 16a^{\frac{9}{4}}bx^4\sqrt[4]{1 - \frac{bx^4}{a}}\Gamma(\frac{9}{4})} + \frac{4bx^5\Gamma(\frac{1}{4})}{-16a^{\frac{13}{4}}\sqrt[4]{1 - \frac{bx^4}{a}}\Gamma(\frac{9}{4}) + 16a^{\frac{9}{4}}bx^4\sqrt[4]{1 - \frac{bx^4}{a}}\Gamma(\frac{9}{4})} \end{cases}$$

input `integrate(1/(-b*x**4+a)**(9/4),x)`

output `Piecewise((-5*a*x*exp(-I*pi/4)*gamma(1/4)/(-16*a**(13/4)*(-1 + b*x**4/a)**(1/4)*gamma(9/4) + 16*a**(9/4)*b*x**4*(-1 + b*x**4/a)**(1/4)*gamma(9/4)) + 4*b*x**5*exp(-I*pi/4)*gamma(1/4)/(-16*a**(13/4)*(-1 + b*x**4/a)**(1/4)*gamma(9/4) + 16*a**(9/4)*b*x**4*(-1 + b*x**4/a)**(1/4)*gamma(9/4)), Abs(b*x**4/a) > 1), (-5*a*x*gamma(1/4)/(-16*a**(13/4)*(1 - b*x**4/a)**(1/4)*gamma(9/4) + 16*a**(9/4)*b*x**4*(1 - b*x**4/a)**(1/4)*gamma(9/4)) + 4*b*x**5*gamma(1/4)/(-16*a**(13/4)*(1 - b*x**4/a)**(1/4)*gamma(9/4) + 16*a**(9/4)*b*x**4*(1 - b*x**4/a)**(1/4)*gamma(9/4)), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.83

$$\int \frac{1}{(a - bx^4)^{9/4}} dx = \frac{\left(b - \frac{5(bx^4 - a)}{x^4}\right)x^5}{5(-bx^4 + a)^{\frac{5}{4}}a^2}$$

input `integrate(1/(-b*x^4+a)^(9/4),x, algorithm="maxima")`

output `1/5*(b - 5*(b*x^4 - a)/x^4)*x^5/((-b*x^4 + a)^(5/4)*a^2)`

Giac [F]

$$\int \frac{1}{(a - bx^4)^{9/4}} dx = \int \frac{1}{(-bx^4 + a)^{9/4}} dx$$

input `integrate(1/(-b*x^4+a)^(9/4),x, algorithm="giac")`

output `integrate((-b*x^4 + a)^(-9/4), x)`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.73

$$\int \frac{1}{(a - bx^4)^{9/4}} dx = \frac{4x(a - bx^4) + ax}{5a^2(a - bx^4)^{5/4}}$$

input `int(1/(a - b*x^4)^(9/4),x)`

output `(4*x*(a - b*x^4) + a*x)/(5*a^2*(a - b*x^4)^(5/4))`

Reduce [F]

$$\int \frac{1}{(a - bx^4)^{9/4}} dx = \int \frac{1}{(-bx^4 + a)^{1/4} a^2 - 2(-bx^4 + a)^{1/4} abx^4 + (-bx^4 + a)^{1/4} b^2x^8} dx$$

input `int(1/(-b*x^4+a)^(9/4),x)`

output `int(1/((a - b*x**4)**(1/4)*a**2 - 2*(a - b*x**4)**(1/4)*a*b*x**4 + (a - b*x**4)**(1/4)*b**2*x**8),x)`

3.49 $\int \frac{1}{(a-bx^4)^{13/4}} dx$

Optimal result	341
Mathematica [A] (verified)	341
Rubi [A] (verified)	342
Maple [A] (verified)	343
Fricas [A] (verification not implemented)	343
Sympy [C] (verification not implemented)	344
Maxima [A] (verification not implemented)	345
Giac [F]	345
Mupad [B] (verification not implemented)	345
Reduce [F]	346

Optimal result

Integrand size = 12, antiderivative size = 61

$$\int \frac{1}{(a-bx^4)^{13/4}} dx = \frac{x}{9a(a-bx^4)^{9/4}} + \frac{8x}{45a^2(a-bx^4)^{5/4}} + \frac{32x}{45a^3\sqrt[4]{a-bx^4}}$$

output `1/9*x/a/(-b*x^4+a)^(9/4)+8/45*x/a^2/(-b*x^4+a)^(5/4)+32/45*x/a^3/(-b*x^4+a)^(1/4)`

Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.67

$$\int \frac{1}{(a-bx^4)^{13/4}} dx = \frac{45a^2x - 72abx^5 + 32b^2x^9}{45a^3(a-bx^4)^{9/4}}$$

input `Integrate[(a - b*x^4)^(-13/4),x]`

output `(45*a^2*x - 72*a*b*x^5 + 32*b^2*x^9)/(45*a^3*(a - b*x^4)^(9/4))`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.13, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {749, 749, 746}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a - bx^4)^{13/4}} dx \\
 & \quad \downarrow \text{749} \\
 & \frac{8 \int \frac{1}{(a - bx^4)^{9/4}} dx}{9a} + \frac{x}{9a(a - bx^4)^{9/4}} \\
 & \quad \downarrow \text{749} \\
 & \frac{8 \left(\frac{4 \int \frac{1}{(a - bx^4)^{5/4}} dx}{5a} + \frac{x}{5a(a - bx^4)^{5/4}} \right)}{9a} + \frac{x}{9a(a - bx^4)^{9/4}} \\
 & \quad \downarrow \text{746} \\
 & \frac{8 \left(\frac{4x}{5a^2 \sqrt[4]{a - bx^4}} + \frac{x}{5a(a - bx^4)^{5/4}} \right)}{9a} + \frac{x}{9a(a - bx^4)^{9/4}}
 \end{aligned}$$

input `Int[(a - b*x^4)^(-13/4),x]`

output `x/(9*a*(a - b*x^4)^(9/4)) + (8*(x/(5*a*(a - b*x^4)^(5/4)) + (4*x)/(5*a^2*(a - b*x^4)^(1/4))))/(9*a)`

Defintions of rubi rules used

rule 746 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[x*((a + b*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]`

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

Maple [A] (verified)

Time = 0.62 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.62

method	result	size
gospers	$\frac{x(32b^2x^8 - 72abx^4 + 45a^2)}{45(-bx^4 + a)^{\frac{9}{4}}a^3}$	38
trager	$\frac{x(32b^2x^8 - 72abx^4 + 45a^2)}{45(-bx^4 + a)^{\frac{9}{4}}a^3}$	38
pseudoelliptic	$\frac{x(32b^2x^8 - 72abx^4 + 45a^2)}{45(-bx^4 + a)^{\frac{9}{4}}a^3}$	38
orering	$\frac{x(32b^2x^8 - 72abx^4 + 45a^2)}{45(-bx^4 + a)^{\frac{9}{4}}a^3}$	38

input `int(1/(-b*x^4+a)^(13/4),x,method=_RETURNVERBOSE)`

output `1/45*x*(32*b^2*x^8-72*a*b*x^4+45*a^2)/(-b*x^4+a)^(9/4)/a^3`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.18

$$\int \frac{1}{(a - bx^4)^{13/4}} dx = -\frac{(32b^2x^9 - 72abx^5 + 45a^2x)(-bx^4 + a)^{\frac{3}{4}}}{45(a^3b^3x^{12} - 3a^4b^2x^8 + 3a^5bx^4 - a^6)}$$

input `integrate(1/(-b*x^4+a)^(13/4),x, algorithm="fricas")`

output
$$-1/45*(32*b^2*x^9 - 72*a*b*x^5 + 45*a^2*x)*(-b*x^4 + a)^{3/4}/(a^3*b^3*x^{12} - 3*a^4*b^2*x^8 + 3*a^5*b*x^4 - a^6)$$

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.40 (sec) , antiderivative size = 1066, normalized size of antiderivative = 17.48

$$\int \frac{1}{(a - bx^4)^{13/4}} dx = \text{Too large to display}$$

input `integrate(1/(-b*x**4+a)**(13/4),x)`

output `Piecewise((-45*a**5*x*exp(-I*pi/4)*gamma(1/4)/(-64*a**(33/4)*(-1 + b*x**4/a)**(1/4)*gamma(13/4) + 192*a**(29/4)*b*x**4*(-1 + b*x**4/a)**(1/4)*gamma(13/4) - 192*a**(25/4)*b**2*x**8*(-1 + b*x**4/a)**(1/4)*gamma(13/4) + 64*a**(21/4)*b**3*x**12*(-1 + b*x**4/a)**(1/4)*gamma(13/4)) + 117*a**4*b*x**5*exp(-I*pi/4)*gamma(1/4)/(-64*a**(33/4)*(-1 + b*x**4/a)**(1/4)*gamma(13/4) + 192*a**(29/4)*b*x**4*(-1 + b*x**4/a)**(1/4)*gamma(13/4) - 192*a**(25/4)*b**2*x**8*(-1 + b*x**4/a)**(1/4)*gamma(13/4) + 64*a**(21/4)*b**3*x**12*(-1 + b*x**4/a)**(1/4)*gamma(13/4)) - 104*a**3*b**2*x**9*exp(-I*pi/4)*gamma(1/4)/(-64*a**(33/4)*(-1 + b*x**4/a)**(1/4)*gamma(13/4) + 192*a**(29/4)*b*x**4*(-1 + b*x**4/a)**(1/4)*gamma(13/4) - 192*a**(25/4)*b**2*x**8*(-1 + b*x**4/a)**(1/4)*gamma(13/4) + 64*a**(21/4)*b**3*x**12*(-1 + b*x**4/a)**(1/4)*gamma(13/4)) + 32*a**2*b**3*x**13*exp(-I*pi/4)*gamma(1/4)/(-64*a**(33/4)*(-1 + b*x**4/a)**(1/4)*gamma(13/4) + 192*a**(29/4)*b*x**4*(-1 + b*x**4/a)**(1/4)*gamma(13/4) - 192*a**(25/4)*b**2*x**8*(-1 + b*x**4/a)**(1/4)*gamma(13/4) + 64*a**(21/4)*b**3*x**12*(-1 + b*x**4/a)**(1/4)*gamma(13/4)), Abs(b*x**4/a) > 1), (-45*a**5*x*gamma(1/4)/(-64*a**(33/4)*(1 - b*x**4/a)**(1/4)*gamma(13/4) + 192*a**(29/4)*b*x**4*(1 - b*x**4/a)**(1/4)*gamma(13/4) - 192*a**(25/4)*b**2*x**8*(1 - b*x**4/a)**(1/4)*gamma(13/4) + 64*a**(21/4)*b**3*x**12*(1 - b*x**4/a)**(1/4)*gamma(13/4)) + 117*a**4*b*x**5*gamma(1/4)/(-64*a**(33/4)*(1 - b*x**4/a)**(1/4)*gamma(13/4) + 192*a**(29/4)*b*x**4*(1 - b*x**4/a)**(1/4)*gamma(13/4) - 192*a**(25/4)*b**2*x**8*(1 - b*x**4/a)**(1/4)*gamma(13/4) + 64*a**(21/4)*b**3*x**12*(1 - b*x**4/a)**(1/4)*gamma(13/4)), Abs(b*x**4/a) < 1))`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.90

$$\int \frac{1}{(a - bx^4)^{13/4}} dx = \frac{\left(5b^2 - \frac{18(bx^4 - a)b}{x^4} + \frac{45(bx^4 - a)^2}{x^8}\right)x^9}{45(-bx^4 + a)^{9/4}a^3}$$

input `integrate(1/(-b*x^4+a)^(13/4),x, algorithm="maxima")`output `1/45*(5*b^2 - 18*(b*x^4 - a)*b/x^4 + 45*(b*x^4 - a)^2/x^8)*x^9/((-b*x^4 + a)^(9/4)*a^3)`**Giac [F]**

$$\int \frac{1}{(a - bx^4)^{13/4}} dx = \int \frac{1}{(-bx^4 + a)^{13/4}} dx$$

input `integrate(1/(-b*x^4+a)^(13/4),x, algorithm="giac")`output `integrate((-b*x^4 + a)^(-13/4), x)`**Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.77

$$\int \frac{1}{(a - bx^4)^{13/4}} dx = \frac{32x(a - bx^4)^2 + 5a^2x + 8ax(a - bx^4)}{45a^3(a - bx^4)^{9/4}}$$

input `int(1/(a - b*x^4)^(13/4),x)`output `(32*x*(a - b*x^4)^2 + 5*a^2*x + 8*a*x*(a - b*x^4))/(45*a^3*(a - b*x^4)^(9/4))`

Reduce [F]

$$\int \frac{1}{(a - bx^4)^{13/4}} dx = \int \frac{1}{(-bx^4 + a)^{1/4} a^3 - 3(-bx^4 + a)^{1/4} a^2 bx^4 + 3(-bx^4 + a)^{1/4} a b^2 x^8 - (-bx^4 + a)^{1/4} b^3 x^{12}} dx$$

input `int(1/(-b*x^4+a)^(13/4),x)`

output `int(1/((a - b*x**4)**(1/4)*a**3 - 3*(a - b*x**4)**(1/4)*a**2*b*x**4 + 3*(a - b*x**4)**(1/4)*a*b**2*x**8 - (a - b*x**4)**(1/4)*b**3*x**12),x)`

3.50 $\int \sqrt[4]{a - bx^4} dx$

Optimal result	347
Mathematica [C] (verified)	347
Rubi [A] (verified)	348
Maple [F]	350
Fricas [F]	350
Sympy [C] (verification not implemented)	350
Maxima [F]	351
Giac [F]	351
Mupad [B] (verification not implemented)	351
Reduce [F]	352

Optimal result

Integrand size = 12, antiderivative size = 83

$$\int \sqrt[4]{a - bx^4} dx = \frac{1}{2}x\sqrt[4]{a - bx^4} - \frac{\sqrt{a}\sqrt{b}\left(1 - \frac{a}{bx^4}\right)^{3/4} x^3 \operatorname{EllipticF}\left(\frac{1}{2} \operatorname{csc}^{-1}\left(\frac{\sqrt{bx^2}}{\sqrt{a}}\right), 2\right)}{2(a - bx^4)^{3/4}}$$

output

```
1/2*x*(-b*x^4+a)^(1/4)-1/2*a^(1/2)*b^(1/2)*(1-a/b/x^4)^(3/4)*x^3*InverseJacobiAM(1/2*arccsc(b^(1/2)*x^2/a^(1/2)),2^(1/2))/(-b*x^4+a)^(3/4)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.12 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.57

$$\int \sqrt[4]{a - bx^4} dx = \frac{x\sqrt[4]{a - bx^4} \operatorname{Hypergeometric2F1}\left(-\frac{1}{4}, \frac{1}{4}, \frac{5}{4}, \frac{bx^4}{a}\right)}{\sqrt[4]{1 - \frac{bx^4}{a}}}$$

input

```
Integrate[(a - b*x^4)^(1/4),x]
```

output

$$(x*(a - b*x^4)^{(1/4)}*Hypergeometric2F1[-1/4, 1/4, 5/4, (b*x^4)/a])/(1 - (b*x^4)/a)^{(1/4)}$$
Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {748, 768, 858, 807, 230}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt[4]{a - bx^4} dx$$

$$\downarrow 748$$

$$\frac{1}{2}a \int \frac{1}{(a - bx^4)^{3/4}} dx + \frac{1}{2}x \sqrt[4]{a - bx^4}$$

$$\downarrow 768$$

$$\frac{ax^3(1 - \frac{a}{bx^4})^{3/4} \int \frac{1}{(1 - \frac{a}{bx^4})^{3/4} x^3} dx}{2(a - bx^4)^{3/4}} + \frac{1}{2}x \sqrt[4]{a - bx^4}$$

$$\downarrow 858$$

$$\frac{1}{2}x \sqrt[4]{a - bx^4} - \frac{ax^3(1 - \frac{a}{bx^4})^{3/4} \int \frac{1}{(1 - \frac{a}{bx^4})^{3/4} x} d\frac{1}{x}}{2(a - bx^4)^{3/4}}$$

$$\downarrow 807$$

$$\frac{1}{2}x \sqrt[4]{a - bx^4} - \frac{ax^3(1 - \frac{a}{bx^4})^{3/4} \int \frac{1}{(1 - \frac{a}{bx^2})^{3/4}} d\frac{1}{x^2}}{4(a - bx^4)^{3/4}}$$

$$\downarrow 230$$

$$\frac{1}{2}x \sqrt[4]{a - bx^4} - \frac{\sqrt{a}\sqrt{bx^3}(1 - \frac{a}{bx^4})^{3/4} \text{EllipticF}\left(\frac{1}{2} \arcsin\left(\frac{\sqrt{a}}{\sqrt{bx^2}}\right), 2\right)}{2(a - bx^4)^{3/4}}$$

input `Int[(a - b*x^4)^(1/4),x]`

output `(x*(a - b*x^4)^(1/4))/2 - (Sqrt[a]*Sqrt[b]*(1 - a/(b*x^4))^(3/4)*x^3*EllipticF[ArcSin[Sqrt[a]/(Sqrt[b]*x^2)]/2, 2])/(2*(a - b*x^4)^(3/4))`

Defintions of rubi rules used

rule 230 `Int[((a_) + (b_.)*(x_)^2)^(-3/4), x_Symbol] := Simp[(2/(a^(3/4)*Rt[-b/a, 2]))*EllipticF[(1/2)*ArcSin[Rt[-b/a, 2]*x], 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b/a]`

rule 748 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p + 1)), x] + Simp[a*n*(p/(n*p + 1)) Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || LtQ[Denominator[p + 1/n], Denominator[p]])`

rule 768 `Int[((a_) + (b_.)*(x_)^4)^(-3/4), x_Symbol] := Simp[x^3*((1 + a/(b*x^4))^(3/4)/(a + b*x^4)^(3/4)) Int[1/(x^3*(1 + a/(b*x^4))^(3/4)), x], x] /; FreeQ[{a, b}, x]`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

rule 858 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^(m + 2), x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0] && IntegerQ[m]`

Maple [F]

$$\int (-bx^4 + a)^{\frac{1}{4}} dx$$

input `int((-b*x^4+a)^(1/4),x)`

output `int((-b*x^4+a)^(1/4),x)`

Fricas [F]

$$\int \sqrt[4]{a - bx^4} dx = \int (-bx^4 + a)^{\frac{1}{4}} dx$$

input `integrate((-b*x^4+a)^(1/4),x, algorithm="fricas")`

output `integral((-b*x^4 + a)^(1/4), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.50 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.47

$$\int \sqrt[4]{a - bx^4} dx = \frac{\sqrt[4]{ax}\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\begin{matrix} -\frac{1}{4}, \frac{1}{4} \\ \frac{5}{4} \end{matrix} \middle| \frac{bx^4 e^{2i\pi}}{a}\right)}{4\Gamma\left(\frac{5}{4}\right)}$$

input `integrate((-b*x**4+a)**(1/4),x)`

output `a**(1/4)*x*gamma(1/4)*hyper((-1/4, 1/4), (5/4,), b*x**4*exp_polar(2*I*pi)/a)/(4*gamma(5/4))`

Maxima [F]

$$\int \sqrt[4]{a - bx^4} dx = \int (-bx^4 + a)^{\frac{1}{4}} dx$$

input `integrate((-b*x^4+a)^(1/4),x, algorithm="maxima")`

output `integrate((-b*x^4 + a)^(1/4), x)`

Giac [F]

$$\int \sqrt[4]{a - bx^4} dx = \int (-bx^4 + a)^{\frac{1}{4}} dx$$

input `integrate((-b*x^4+a)^(1/4),x, algorithm="giac")`

output `integrate((-b*x^4 + a)^(1/4), x)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.46

$$\int \sqrt[4]{a - bx^4} dx = \frac{x (a - bx^4)^{1/4} {}_2F_1\left(-\frac{1}{4}, \frac{1}{4}; \frac{5}{4}, \frac{bx^4}{a}\right)}{\left(1 - \frac{bx^4}{a}\right)^{1/4}}$$

input `int((a - b*x^4)^(1/4),x)`

output `(x*(a - b*x^4)^(1/4)*hypergeom([-1/4, 1/4], 5/4, (b*x^4)/a))/(1 - (b*x^4)/a)^(1/4)`

Reduce [F]

$$\int \sqrt[4]{a - bx^4} dx = \frac{(-bx^4 + a)^{\frac{1}{4}} x}{2} + \frac{\left(\int \frac{1}{(-bx^4 + a)^{\frac{3}{4}}} dx \right) a}{2}$$

input `int((-b*x^4+a)^(1/4),x)`

output `((a - b*x**4)**(1/4)*x + int((a - b*x**4)**(1/4)/(a - b*x**4),x)*a)/2`

3.51 $\int \frac{1}{(a-bx^4)^{3/4}} dx$

Optimal result	353
Mathematica [C] (verified)	353
Rubi [A] (verified)	354
Maple [F]	355
Fricas [F]	356
Sympy [C] (verification not implemented)	356
Maxima [F]	356
Giac [F]	357
Mupad [B] (verification not implemented)	357
Reduce [F]	357

Optimal result

Integrand size = 12, antiderivative size = 63

$$\int \frac{1}{(a-bx^4)^{3/4}} dx = -\frac{\sqrt{b}\left(1-\frac{a}{bx^4}\right)^{3/4} x^3 \operatorname{EllipticF}\left(\frac{1}{2} \operatorname{csc}^{-1}\left(\frac{\sqrt{bx^2}}{\sqrt{a}}\right), 2\right)}{\sqrt{a}(a-bx^4)^{3/4}}$$

output `-b^(1/2)*(1-a/b/x^4)^(3/4)*x^3*InverseJacobiAM(1/2*arccsc(b^(1/2)*x^2/a^(1/2)),2^(1/2))/a^(1/2)/(-b*x^4+a)^(3/4)`

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.12 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.75

$$\int \frac{1}{(a-bx^4)^{3/4}} dx = \frac{x\left(1-\frac{bx^4}{a}\right)^{3/4} \operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{3}{4}, \frac{5}{4}, \frac{bx^4}{a}\right)}{(a-bx^4)^{3/4}}$$

input `Integrate[(a - b*x^4)^(-3/4), x]`

output

```
(x*(1 - (b*x^4)/a)^(3/4)*Hypergeometric2F1[1/4, 3/4, 5/4, (b*x^4)/a])/(a - b*x^4)^(3/4)
```

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {768, 858, 807, 230}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a - bx^4)^{3/4}} dx \\
 & \quad \downarrow \text{768} \\
 & \frac{x^3 \left(1 - \frac{a}{bx^4}\right)^{3/4} \int \frac{1}{\left(1 - \frac{a}{bx^4}\right)^{3/4} x^3} dx}{(a - bx^4)^{3/4}} \\
 & \quad \downarrow \text{858} \\
 & \frac{x^3 \left(1 - \frac{a}{bx^4}\right)^{3/4} \int \frac{1}{\left(1 - \frac{a}{bx^4}\right)^{3/4} x} d\frac{1}{x}}{(a - bx^4)^{3/4}} \\
 & \quad \downarrow \text{807} \\
 & \frac{x^3 \left(1 - \frac{a}{bx^4}\right)^{3/4} \int \frac{1}{\left(1 - \frac{a}{bx^2}\right)^{3/4} d\frac{1}{x^2}}}{2(a - bx^4)^{3/4}} \\
 & \quad \downarrow \text{230} \\
 & \frac{\sqrt{b}x^3 \left(1 - \frac{a}{bx^4}\right)^{3/4} \text{EllipticF}\left(\frac{1}{2} \arcsin\left(\frac{\sqrt{a}}{\sqrt{bx^2}}\right), 2\right)}{\sqrt{a}(a - bx^4)^{3/4}}
 \end{aligned}$$

input

```
Int[(a - b*x^4)^(-3/4), x]
```

output $-\left(\frac{\sqrt{b}(1 - a/(bx^4))^{3/4}x^3 \operatorname{EllipticF}[\operatorname{ArcSin}[\sqrt{a}/(\sqrt{b}x^2)]/2, 2]}{\sqrt{a}(a - bx^4)^{3/4}}\right)$

Defintions of rubi rules used

rule 230 $\operatorname{Int}[(a_ + (b_ \cdot)(x_)^2)^{-3/4}, x_Symbol] \rightarrow \operatorname{Simp}[(2/(a^{3/4} \operatorname{Rt}[-b/a, 2])) \cdot \operatorname{EllipticF}[(1/2) \cdot \operatorname{ArcSin}[\operatorname{Rt}[-b/a, 2] \cdot x], 2], x] \;/; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{GtQ}[a, 0] \ \&\& \operatorname{NegQ}[b/a]$

rule 768 $\operatorname{Int}[(a_ + (b_ \cdot)(x_)^4)^{-3/4}, x_Symbol] \rightarrow \operatorname{Simp}[x^3 \cdot ((1 + a/(bx^4))^{3/4}) / (a + bx^4)^{3/4}] \operatorname{Int}[1/(x^3(1 + a/(bx^4))^{3/4}), x], x] \;/; \operatorname{FreeQ}\{a, b\}, x]$

rule 807 $\operatorname{Int}[(x_)^{(m_ \cdot)} \cdot ((a_ + (b_ \cdot)(x_)^{(n_)})^{(p_)}), x_Symbol] \rightarrow \operatorname{With}\{k = \operatorname{GCD}[m + 1, n]\}, \operatorname{Simp}[1/k \operatorname{Subst}[\operatorname{Int}[x^{(m+1)/k - 1} \cdot (a + bx^{n/k})^p, x], x, x^k], x] \;/; k \neq 1] \;/; \operatorname{FreeQ}\{a, b, p\}, x \ \&\& \operatorname{IGtQ}[n, 0] \ \&\& \operatorname{IntegerQ}[m]$

rule 858 $\operatorname{Int}[(x_)^{(m_ \cdot)} \cdot ((a_ + (b_ \cdot)(x_)^{(n_)})^{(p_)}), x_Symbol] \rightarrow -\operatorname{Subst}[\operatorname{Int}[(a + b/x^n)^p / x^{(m+2)}, x], x, 1/x] \;/; \operatorname{FreeQ}\{a, b, p\}, x \ \&\& \operatorname{ILtQ}[n, 0] \ \&\& \operatorname{IntegerQ}[m]$

Maple [F]

$$\int \frac{1}{(-bx^4 + a)^{\frac{3}{4}}} dx$$

input $\operatorname{int}(1/(-bx^4+a)^{3/4}, x)$

output $\operatorname{int}(1/(-bx^4+a)^{3/4}, x)$

Fricas [F]

$$\int \frac{1}{(a - bx^4)^{3/4}} dx = \int \frac{1}{(-bx^4 + a)^{3/4}} dx$$

input `integrate(1/(-b*x^4+a)^(3/4),x, algorithm="fricas")`

output `integral(-(-b*x^4 + a)^(1/4)/(b*x^4 - a), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.51 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.59

$$\int \frac{1}{(a - bx^4)^{3/4}} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{3}{4} \middle| \frac{bx^4 e^{2i\pi}}{a}\right)}{4a^{3/4}\Gamma\left(\frac{5}{4}\right)}$$

input `integrate(1/(-b*x**4+a)**(3/4),x)`

output `x*gamma(1/4)*hyper((1/4, 3/4), (5/4,), b*x**4*exp_polar(2*I*pi)/a)/(4*a**(3/4)*gamma(5/4))`

Maxima [F]

$$\int \frac{1}{(a - bx^4)^{3/4}} dx = \int \frac{1}{(-bx^4 + a)^{3/4}} dx$$

input `integrate(1/(-b*x^4+a)^(3/4),x, algorithm="maxima")`

output `integrate((-b*x^4 + a)^(-3/4), x)`

Giac [F]

$$\int \frac{1}{(a - bx^4)^{3/4}} dx = \int \frac{1}{(-bx^4 + a)^{3/4}} dx$$

input `integrate(1/(-b*x^4+a)^(3/4),x, algorithm="giac")`

output `integrate((-b*x^4 + a)^(-3/4), x)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.60

$$\int \frac{1}{(a - bx^4)^{3/4}} dx = \frac{x \left(1 - \frac{bx^4}{a}\right)^{3/4} {}_2F_1\left(\frac{1}{4}, \frac{3}{4}; \frac{5}{4}; \frac{bx^4}{a}\right)}{(a - bx^4)^{3/4}}$$

input `int(1/(a - b*x^4)^(3/4),x)`

output `(x*(1 - (b*x^4)/a)^(3/4)*hypergeom([1/4, 3/4], 5/4, (b*x^4)/a))/(a - b*x^4)^(3/4)`

Reduce [F]

$$\int \frac{1}{(a - bx^4)^{3/4}} dx = \int \frac{1}{(-bx^4 + a)^{3/4}} dx$$

input `int(1/(-b*x^4+a)^(3/4),x)`

output `int(1/(a - b*x**4)**(3/4),x)`

3.52 $\int \frac{1}{(a-bx^4)^{7/4}} dx$

Optimal result	358
Mathematica [C] (verified)	358
Rubi [A] (verified)	359
Maple [F]	361
Fricas [F]	361
Sympy [C] (verification not implemented)	361
Maxima [F]	362
Giac [F]	362
Mupad [B] (verification not implemented)	362
Reduce [F]	363

Optimal result

Integrand size = 12, antiderivative size = 86

$$\int \frac{1}{(a-bx^4)^{7/4}} dx = \frac{x}{3a(a-bx^4)^{3/4}} - \frac{2\sqrt{b}\left(1-\frac{a}{bx^4}\right)^{3/4} x^3 \operatorname{EllipticF}\left(\frac{1}{2} \operatorname{csc}^{-1}\left(\frac{\sqrt{bx^2}}{\sqrt{a}}\right), 2\right)}{3a^{3/2}(a-bx^4)^{3/4}}$$

output `1/3*x/a/(-b*x^4+a)^(3/4)-2/3*b^(1/2)*(1-a/b/x^4)^(3/4)*x^3*InverseJacobiAM(1/2*arccsc(b^(1/2)*x^2/a^(1/2)),2^(1/2))/a^(3/2)/(-b*x^4+a)^(3/4)`

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.51 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.66

$$\int \frac{1}{(a-bx^4)^{7/4}} dx = \frac{x + 2x\left(1-\frac{bx^4}{a}\right)^{3/4} \operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{3}{4}, \frac{5}{4}, \frac{bx^4}{a}\right)}{3a(a-bx^4)^{3/4}}$$

input `Integrate[(a - b*x^4)^(-7/4), x]`

output

```
(x + 2*x*(1 - (b*x^4)/a)^(3/4)*Hypergeometric2F1[1/4, 3/4, 5/4, (b*x^4)/a]
)/(3*a*(a - b*x^4)^(3/4))
```

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {749, 768, 858, 807, 230}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a - bx^4)^{7/4}} dx \\
 & \quad \downarrow \text{749} \\
 & \frac{2 \int \frac{1}{(a - bx^4)^{3/4}} dx}{3a} + \frac{x}{3a(a - bx^4)^{3/4}} \\
 & \quad \downarrow \text{768} \\
 & \frac{2x^3 \left(1 - \frac{a}{bx^4}\right)^{3/4} \int \frac{1}{\left(1 - \frac{a}{bx^4}\right)^{3/4} x^3} dx}{3a(a - bx^4)^{3/4}} + \frac{x}{3a(a - bx^4)^{3/4}} \\
 & \quad \downarrow \text{858} \\
 & \frac{x}{3a(a - bx^4)^{3/4}} - \frac{2x^3 \left(1 - \frac{a}{bx^4}\right)^{3/4} \int \frac{1}{\left(1 - \frac{a}{bx^4}\right)^{3/4} x} d\frac{1}{x}}{3a(a - bx^4)^{3/4}} \\
 & \quad \downarrow \text{807} \\
 & \frac{x}{3a(a - bx^4)^{3/4}} - \frac{x^3 \left(1 - \frac{a}{bx^4}\right)^{3/4} \int \frac{1}{\left(1 - \frac{a}{bx^4}\right)^{3/4} x^2} d\frac{1}{x^2}}{3a(a - bx^4)^{3/4}} \\
 & \quad \downarrow \text{230} \\
 & \frac{x}{3a(a - bx^4)^{3/4}} - \frac{2\sqrt{b}x^3 \left(1 - \frac{a}{bx^4}\right)^{3/4} \text{EllipticF}\left(\frac{1}{2} \arcsin\left(\frac{\sqrt{a}}{\sqrt{bx^2}}\right), 2\right)}{3a^{3/2}(a - bx^4)^{3/4}}
 \end{aligned}$$

input `Int[(a - b*x^4)^(-7/4),x]`

output `x/(3*a*(a - b*x^4)^(3/4)) - (2*Sqrt[b]*(1 - a/(b*x^4))^(3/4)*x^3*EllipticF[ArcSin[Sqrt[a]/(Sqrt[b]*x^2)]/2, 2])/(3*a^(3/2)*(a - b*x^4)^(3/4))`

Defintions of rubi rules used

rule 230 `Int[((a_) + (b_.)*(x_)^2)^(-3/4), x_Symbol] := Simp[(2/(a^(3/4)*Rt[-b/a, 2]))*EllipticF[(1/2)*ArcSin[Rt[-b/a, 2]*x], 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b/a]`

rule 749 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || Denominator[p + 1/n] < Denominator[p])`

rule 768 `Int[((a_) + (b_.)*(x_)^4)^(-3/4), x_Symbol] := Simp[x^3*((1 + a/(b*x^4))^(3/4)/(a + b*x^4)^(3/4)) Int[1/(x^3*(1 + a/(b*x^4))^(3/4)), x], x] /; FreeQ[{a, b}, x]`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

rule 858 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^(m + 2), x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0] && IntegerQ[m]`

Maple [F]

$$\int \frac{1}{(-bx^4 + a)^{7/4}} dx$$

input `int(1/(-b*x^4+a)^(7/4),x)`

output `int(1/(-b*x^4+a)^(7/4),x)`

Fricas [F]

$$\int \frac{1}{(a - bx^4)^{7/4}} dx = \int \frac{1}{(-bx^4 + a)^{7/4}} dx$$

input `integrate(1/(-b*x^4+a)^(7/4),x, algorithm="fricas")`

output `integral((-b*x^4 + a)^(1/4)/(b^2*x^8 - 2*a*b*x^4 + a^2), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.60 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.43

$$\int \frac{1}{(a - bx^4)^{7/4}} dx = \frac{x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{7}{4} \middle| \frac{bx^4 e^{2i\pi}}{a}\right)}{4a^{7/4}\Gamma\left(\frac{5}{4}\right)}$$

input `integrate(1/(-b*x**4+a)**(7/4),x)`

output `x*gamma(1/4)*hyper((1/4, 7/4), (5/4,), b*x**4*exp_polar(2*I*pi)/a)/(4*a**
7/4)*gamma(5/4)`

Maxima [F]

$$\int \frac{1}{(a - bx^4)^{7/4}} dx = \int \frac{1}{(-bx^4 + a)^{7/4}} dx$$

input `integrate(1/(-b*x^4+a)^(7/4),x, algorithm="maxima")`

output `integrate((-b*x^4 + a)^(-7/4), x)`

Giac [F]

$$\int \frac{1}{(a - bx^4)^{7/4}} dx = \int \frac{1}{(-bx^4 + a)^{7/4}} dx$$

input `integrate(1/(-b*x^4+a)^(7/4),x, algorithm="giac")`

output `integrate((-b*x^4 + a)^(-7/4), x)`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.44

$$\int \frac{1}{(a - bx^4)^{7/4}} dx = \frac{x \left(1 - \frac{bx^4}{a}\right)^{7/4} {}_2F_1\left(\frac{1}{4}, \frac{7}{4}, \frac{5}{4}, \frac{bx^4}{a}\right)}{(a - bx^4)^{7/4}}$$

input `int(1/(a - b*x^4)^(7/4),x)`

output `(x*(1 - (b*x^4)/a)^(7/4)*hypergeom([1/4, 7/4], 5/4, (b*x^4)/a))/(a - b*x^4)^(7/4)`

Reduce [F]

$$\int \frac{1}{(a - bx^4)^{7/4}} dx = \int \frac{1}{(-bx^4 + a)^{3/4} a - (-bx^4 + a)^{3/4} bx^4} dx$$

input `int(1/(-b*x^4+a)^(7/4),x)`

output `int(1/((a - b*x**4)**(3/4)*a - (a - b*x**4)**(3/4)*b*x**4),x)`

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions	364
4.2 Links to plain text integration problems used in this report for each CAS .	382

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
          finalresult={"C","Result contains complex when optimal does not."}
        ]
      ,(*ELSE*)(*result does not contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
          ]
        ]
      ,(*ELSE*)(*expnResult>expnOptimal*)
        If[FreeQ[result,Integrate] && FreeQ[result,Int],
          finalresult={"C","Result contains higher order function than in optimal. Order "
          ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]==Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]==Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]==Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]==Plus || Head[expn]==Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```


Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

#   antiderivative
#   "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

        end if
    else #result contains complex but optimal is not
        if debug then
            print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
    fi;
else # result do not contain complex
    # this assumes optimal do not as well. No check is needed here.
    if debug then
        print("result do not contain complex, this assumes optimal do not as well
    fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co

        fi;
    fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                    convert(ExpnType_result,string)," vs. order ",
                    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```



```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file