

Computer Algebra Independent Integration Tests

Summer 2024

1-Algebraic-functions/1.1-Binomial/1.1.3-General-
binomial/1.1.3.2/49-1.1.3.2-g

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 9:07pm

Contents

1	Introduction	5
1.1	Listing of CAS systems tested	6
1.2	Results	7
1.3	Time and leaf size Performance	11
1.4	Performance based on number of rules Rubi used	13
1.5	Performance based on number of steps Rubi used	14
1.6	Solved integrals histogram based on leaf size of result	15
1.7	Solved integrals histogram based on CPU time used	16
1.8	Leaf size vs. CPU time used	17
1.9	list of integrals with no known antiderivative	18
1.10	List of integrals solved by CAS but has no known antiderivative	18
1.11	list of integrals solved by CAS but failed verification	18
1.12	Timing	19
1.13	Verification	19
1.14	Important notes about some of the results	20
1.15	Current tree layout of integration tests	23
1.16	Design of the test system	24
2	detailed summary tables of results	25
2.1	List of integrals sorted by grade for each CAS	26
2.2	Detailed conclusion table per each integral for all CAS systems	30
2.3	Detailed conclusion table specific for Rubi results	48
3	Listing of integrals	51
3.1	$\int x^5 (c(a + bx^2)^2)^{3/2} dx$	55
3.2	$\int x^3 (c(a + bx^2)^2)^{3/2} dx$	61
3.3	$\int x (c(a + bx^2)^2)^{3/2} dx$	67
3.4	$\int \frac{(c(a+bx^2)^2)^{3/2}}{x} dx$	72

3.5	$\int \frac{(c+bx^2)^2)^{3/2}}{x^3} dx$	78
3.6	$\int \frac{(c+bx^2)^2)^{3/2}}{x^5} dx$	85
3.7	$\int x^4 (c+bx^2)^2)^{3/2} dx$	92
3.8	$\int x^2 (c+bx^2)^2)^{3/2} dx$	98
3.9	$\int (c+bx^2)^2)^{3/2} dx$	104
3.10	$\int \frac{(c+bx^2)^2)^{3/2}}{x^2} dx$	109
3.11	$\int \frac{(c+bx^2)^2)^{3/2}}{x^4} dx$	115
3.12	$\int x^3 (c+bx^2)^3)^{3/2} dx$	121
3.13	$\int x (c+bx^2)^3)^{3/2} dx$	127
3.14	$\int \frac{(c+bx^2)^3)^{3/2}}{x} dx$	133
3.15	$\int \frac{(c+bx^2)^3)^{3/2}}{x^3} dx$	141
3.16	$\int x^2 (c+bx^2)^3)^{3/2} dx$	150
3.17	$\int (c+bx^2)^3)^{3/2} dx$	158
3.18	$\int \frac{(c+bx^2)^3)^{3/2}}{x^2} dx$	165
3.19	$\int \frac{(c+bx^2)^3)^{3/2}}{x^4} dx$	173
3.20	$\int x^3 \left(\frac{c}{a+bx^2}\right)^{3/2} dx$	181
3.21	$\int x \left(\frac{c}{a+bx^2}\right)^{3/2} dx$	187
3.22	$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x} dx$	192
3.23	$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^3} dx$	198
3.24	$\int x^4 \left(\frac{c}{a+bx^2}\right)^{3/2} dx$	205
3.25	$\int x^2 \left(\frac{c}{a+bx^2}\right)^{3/2} dx$	211
3.26	$\int \left(\frac{c}{a+bx^2}\right)^{3/2} dx$	217
3.27	$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^2} dx$	222
3.28	$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^4} dx$	228
3.29	$\int x^5 \left(\frac{c}{(a+bx^2)^2}\right)^{3/2} dx$	234
3.30	$\int x^3 \left(\frac{c}{(a+bx^2)^2}\right)^{3/2} dx$	240

3.31	$\int x \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx$	245
3.32	$\int \frac{\left(\frac{c}{(a+bx^2)^2} \right)^{3/2}}{x} dx$	250
3.33	$\int \frac{\left(\frac{c}{(a+bx^2)^2} \right)^{3/2}}{x^3} dx$	256
3.34	$\int x^4 \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx$	263
3.35	$\int x^2 \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx$	269
3.36	$\int \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx$	275
3.37	$\int \frac{\left(\frac{c}{(a+bx^2)^2} \right)^{3/2}}{x^2} dx$	281
3.38	$\int \frac{\left(\frac{c}{(a+bx^2)^2} \right)^{3/2}}{x^4} dx$	288
3.39	$\int x^7 (c\sqrt{a+bx^2})^{3/2} dx$	296
3.40	$\int x^5 (c\sqrt{a+bx^2})^{3/2} dx$	303
3.41	$\int x^3 (c\sqrt{a+bx^2})^{3/2} dx$	309
3.42	$\int x (c\sqrt{a+bx^2})^{3/2} dx$	315
3.43	$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x} dx$	320
3.44	$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^3} dx$	328
3.45	$\int x^2 (c\sqrt{a+bx^2})^{3/2} dx$	336
3.46	$\int (c\sqrt{a+bx^2})^{3/2} dx$	342
3.47	$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^2} dx$	348
3.48	$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^4} dx$	354
3.49	$\int x^7 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx$	361
3.50	$\int x^5 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx$	367
3.51	$\int x^3 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx$	373
3.52	$\int x \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx$	379
3.53	$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2}}{x} dx$	384
3.54	$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2}}{x^3} dx$	391

3.55	$\int x^4 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx$	398
3.56	$\int x^2 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx$	404
3.57	$\int \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx$	409
3.58	$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2}}{x^2} dx$	414
3.59	$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2}}{x^4} dx$	420
3.60	$\int (dx)^m \left(c(a+bx^2)^3 \right)^{3/2} dx$	426
3.61	$\int (dx)^m \left(c(a+bx^2)^2 \right)^{3/2} dx$	431
3.62	$\int (dx)^m \left(c(a+bx^2) \right)^{3/2} dx$	438
3.63	$\int (dx)^m \left(\frac{c}{a+bx^2} \right)^{3/2} dx$	443
3.64	$\int (dx)^m \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx$	448
3.65	$\int (dx)^m \left(c(a+bx^n)^3 \right)^{3/2} dx$	453
3.66	$\int (dx)^m \left(c(a+bx^n)^2 \right)^{3/2} dx$	459
3.67	$\int (dx)^m \left(c(a+bx^n) \right)^{3/2} dx$	466
3.68	$\int (dx)^m \left(\frac{c}{a+bx^n} \right)^{3/2} dx$	472
3.69	$\int (dx)^m \left(\frac{c}{(a+bx^n)^2} \right)^{3/2} dx$	477
4	Appendix	482
4.1	Listing of Grading functions	482
4.2	Links to plain text integration problems used in this report for each CAS500	

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	6
1.2	Results	7
1.3	Time and leaf size Performance	11
1.4	Performance based on number of rules Rubi used	13
1.5	Performance based on number of steps Rubi used	14
1.6	Solved integrals histogram based on leaf size of result	15
1.7	Solved integrals histogram based on CPU time used	16
1.8	Leaf size vs. CPU time used	17
1.9	list of integrals with no known antiderivative	18
1.10	List of integrals solved by CAS but has no known antiderivative	18
1.11	list of integrals solved by CAS but failed verification	18
1.12	Timing	19
1.13	Verification	19
1.14	Important notes about some of the results	20
1.15	Current tree layout of integration tests	23
1.16	Design of the test system	24

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [69]. This is test number [49].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (69)	0.00 (0)
Mathematica	100.00 (69)	0.00 (0)
Giac	75.36 (52)	24.64 (17)
Reduce	75.36 (52)	24.64 (17)
Fricas	69.57 (48)	30.43 (21)
Maple	68.12 (47)	31.88 (22)
Maxima	62.32 (43)	37.68 (26)
Mupad	28.99 (20)	71.01 (49)
Sympy	15.94 (11)	84.06 (58)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Mathematica	86.957	0.000	13.043	0.000
Giac	69.565	5.797	0.000	24.638
Maple	66.667	0.000	1.449	31.884
Fricas	66.667	2.899	0.000	30.435
Maxima	59.420	2.899	0.000	37.681
Sympy	13.043	2.899	0.000	84.058
Mupad	0.000	28.986	0.000	71.014
Reduce	0.000	75.362	0.000	24.638

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Giac	17	88.24	0.00	11.76
Reduce	17	100.00	0.00	0.00
Fricas	21	66.67	19.05	14.29
Maple	22	100.00	0.00	0.00
Maxima	26	100.00	0.00	0.00
Mupad	49	0.00	100.00	0.00
Sympy	58	79.31	18.97	1.72

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.06
Fricas	0.09
Giac	0.13
Reduce	0.21
Rubi	0.38
Mupad	0.54
Maple	0.62
Mathematica	1.19
Sympy	3.92

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mupad	61.10	0.94	46.50	0.87
Mathematica	76.04	0.73	66.00	0.67
Maple	76.23	0.69	60.00	0.72
Maxima	85.93	0.87	70.00	0.73
Sympy	87.27	1.44	87.00	1.35
Rubi	90.25	0.87	83.00	0.90
Reduce	102.60	1.01	108.00	1.01
Giac	149.71	1.07	72.50	0.79
Fricas	151.81	1.25	75.50	0.98

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram shows that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand,the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	26
2.2	Detailed conclusion table per each integral for all CAS systems	30
2.3	Detailed conclusion table specific for Rubi results	48

2.1 List of integrals sorted by grade for each CAS

Rubi	26
Mma	26
Maple	27
Fricas	27
Maxima	27
Giac	28
Mupad	28
Sympy	28
Reduce	29

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 49, 50, 51, 52, 53, 54, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69 }

B grade { }

C grade { 45, 46, 47, 48, 55, 56, 57, 58, 59 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 38, 39, 40, 41, 42, 49, 50, 51, 52, 61 }

B grade { }

C grade { 37 }

F normal fail { 43, 44, 45, 46, 47, 48, 53, 54, 55, 56, 57, 58, 59, 60, 62, 63, 64, 65, 66, 67, 68, 69 }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 49, 50, 51, 52, 61, 66 }

B grade { 3, 13 }

C grade { }

F normal fail { 45, 46, 47, 48, 55, 56, 57, 58, 59, 60, 62, 63, 64, 69 }

F(-1) timedout fail { 43, 44, 53, 54 }

F(-2) exception fail { 65, 67, 68 }

Maxima

A grade { 1, 2, 4, 5, 6, 7, 8, 9, 10, 11, 12, 20, 21, 22, 23, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 49, 50, 51, 52, 53, 54, 61, 66 }

B grade { 3, 13 }

C grade { }

F normal fail { 14, 15, 16, 17, 18, 19, 24, 25, 26, 45, 46, 47, 48, 55, 56, 57, 58, 59, 60, 62, 63, 64, 65, 67, 68, 69 }

F(-1) timedout fail { }

F(-2) exception fail { }

Giac

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 49, 50, 51, 52, 54 }

B grade { 28, 53, 61, 66 }

C grade { }

F normal fail { 45, 46, 47, 48, 55, 56, 57, 60, 62, 63, 64, 65, 67, 68, 69 }

F(-1) timedout fail { }

F(-2) exception fail { 58, 59 }

Mupad

A grade { }

B grade { 2, 3, 12, 13, 20, 21, 26, 27, 28, 30, 31, 39, 40, 41, 42, 49, 50, 51, 52, 61 }

C grade { }

F normal fail { }

F(-1) timedout fail { 1, 4, 5, 6, 7, 8, 9, 10, 11, 14, 15, 16, 17, 18, 19, 22, 23, 24, 25, 29, 32, 33, 34, 35, 36, 37, 38, 43, 44, 45, 46, 47, 48, 53, 54, 55, 56, 57, 58, 59, 60, 62, 63, 64, 65, 66, 67, 68, 69 }

F(-2) exception fail { }

Sympy

A grade { 20, 39, 40, 41, 42, 49, 50, 51, 52 }

B grade { 21, 26 }

C grade { }

F normal fail { 3, 4, 5, 6, 8, 9, 10, 11, 13, 14, 15, 16, 17, 18, 19, 22, 23, 24, 25, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 43, 44, 45, 46, 47, 48, 53, 54, 55, 56, 57, 58, 59, 63, 68 }

F(-1) timedout fail { 1, 2, 7, 12, 61, 62, 64, 65, 66, 67, 69 }

F(-2) exception fail { 60 }

Reduce

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26,
27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 49, 50, 51, 52, 53, 54, 61,
66 }

C grade { }

F normal fail { 45, 46, 47, 48, 55, 56, 57, 58, 59, 60, 62, 63, 64, 65, 67, 68, 69 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	143	72	63	60	136	74	0	72	40	0
N.S.	1	0.50	0.44	0.42	0.95	0.52	0.00	0.50	0.28	0.00
time (sec)	N/A	0.391	0.805	0.479	0.037	0.082	0.000	0.120	0.184	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	63	63	60	98	74	0	48	40	50
N.S.	1	0.95	0.95	0.91	1.48	1.12	0.00	0.73	0.61	0.76
time (sec)	N/A	0.375	0.777	0.357	0.036	0.072	0.000	0.124	0.179	0.607

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	29	29	26	60	73	0	25	39	40
N.S.	1	0.91	0.91	0.81	1.88	2.28	0.00	0.78	1.22	1.25
time (sec)	N/A	0.273	0.010	0.362	0.040	0.076	0.000	0.116	0.236	0.642

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	139	68	234	59	171	73	0	73	39	0
N.S.	1	0.49	1.68	0.42	1.23	0.53	0.00	0.53	0.28	0.00
time (sec)	N/A	0.372	0.498	0.261	0.038	0.074	0.000	0.118	0.190	0.000

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	140	67	237	61	176	76	0	91	41	0
N.S.	1	0.48	1.69	0.44	1.26	0.54	0.00	0.65	0.29	0.00
time (sec)	N/A	0.363	0.678	0.309	0.042	0.079	0.000	0.129	0.222	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	140	66	237	62	251	77	0	92	42	0
N.S.	1	0.47	1.69	0.44	1.79	0.55	0.00	0.66	0.30	0.00
time (sec)	N/A	0.378	0.658	0.207	0.042	0.079	0.000	0.127	0.180	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	143	69	63	60	47	74	0	72	40	0
N.S.	1	0.48	0.44	0.42	0.33	0.52	0.00	0.50	0.28	0.00
time (sec)	N/A	0.372	0.077	1.168	0.038	0.072	0.000	0.117	0.201	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	143	69	63	60	47	74	0	72	40	0
N.S.	1	0.48	0.44	0.42	0.33	0.52	0.00	0.50	0.28	0.00
time (sec)	N/A	0.370	0.079	0.904	0.035	0.075	0.000	0.127	0.206	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	135	65	61	58	43	72	0	46	38	0
N.S.	1	0.48	0.45	0.43	0.32	0.53	0.00	0.34	0.28	0.00
time (sec)	N/A	0.334	0.071	0.559	0.036	0.072	0.000	0.117	0.189	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	134	60	62	60	48	72	0	69	39	0
N.S.	1	0.45	0.46	0.45	0.36	0.54	0.00	0.51	0.29	0.00
time (sec)	N/A	0.360	0.086	0.747	0.036	0.071	0.000	0.123	0.207	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	137	63	61	58	48	74	0	71	39	0
N.S.	1	0.46	0.45	0.42	0.35	0.54	0.00	0.52	0.28	0.00
time (sec)	N/A	0.360	0.083	1.023	0.046	0.072	0.000	0.116	0.224	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	86	39	36	86	101	0	55	80	74
N.S.	1	1.30	0.59	0.55	1.30	1.53	0.00	0.83	1.21	1.12
time (sec)	N/A	0.409	0.042	2.418	0.044	0.098	0.000	0.131	0.209	0.607

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	29	29	26	70	87	0	28	67	62
N.S.	1	0.91	0.91	0.81	2.19	2.72	0.00	0.88	2.09	1.94
time (sec)	N/A	0.271	0.010	1.685	0.045	0.089	0.000	0.127	0.203	0.507

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	192	143	111	221	0	382	0	185	146	0
N.S.	1	0.74	0.58	1.15	0.00	1.99	0.00	0.96	0.76	0.00
time (sec)	N/A	0.425	0.107	1.626	0.000	0.095	0.000	0.126	0.237	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	200	153	118	223	0	402	0	196	157	0
N.S.	1	0.76	0.59	1.12	0.00	2.01	0.00	0.98	0.78	0.00
time (sec)	N/A	0.451	0.116	2.470	0.000	0.105	0.000	0.134	0.259	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	253	215	142	155	0	433	0	177	140	0
N.S.	1	0.85	0.56	0.61	0.00	1.71	0.00	0.70	0.55	0.00
time (sec)	N/A	0.550	0.602	2.154	0.000	0.130	0.000	0.145	0.231	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	207	177	123	138	0	402	0	153	121	0
N.S.	1	0.86	0.59	0.67	0.00	1.94	0.00	0.74	0.58	0.00
time (sec)	N/A	0.419	0.141	1.618	0.000	0.109	0.000	0.148	0.199	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	234	179	121	141	0	396	0	185	126	0
N.S.	1	0.76	0.52	0.60	0.00	1.69	0.00	0.79	0.54	0.00
time (sec)	N/A	0.468	0.139	2.414	0.000	0.099	0.000	0.150	0.243	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	236	183	123	143	0	410	0	262	132	0
N.S.	1	0.78	0.52	0.61	0.00	1.74	0.00	1.11	0.56	0.00
time (sec)	N/A	0.494	0.149	3.188	0.000	0.098	0.000	0.141	0.233	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	82	29	28	36	28	66	50	33	42
N.S.	1	1.67	0.59	0.57	0.73	0.57	1.35	1.02	0.67	0.86
time (sec)	N/A	0.401	0.027	0.208	0.029	0.076	0.441	0.120	0.187	0.513

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	21	20	19	19	42	28	25	19
N.S.	1	1.00	1.00	0.95	0.90	0.90	2.00	1.33	1.19	0.90
time (sec)	N/A	0.270	0.004	0.350	0.027	0.077	0.257	0.119	0.202	0.452

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	71	71	59	64	80	138	0	59	139	0
N.S.	1	1.00	0.83	0.90	1.13	1.94	0.00	0.83	1.96	0.00
time (sec)	N/A	0.366	0.042	0.134	0.133	0.086	0.000	0.137	0.193	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	104	99	78	79	121	175	0	103	175	0
N.S.	1	0.95	0.75	0.76	1.16	1.68	0.00	0.99	1.68	0.00
time (sec)	N/A	0.395	0.081	0.189	0.105	0.094	0.000	0.124	0.205	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	115	85	70	0	164	0	93	117	0
N.S.	1	1.02	0.75	0.62	0.00	1.45	0.00	0.82	1.04	0.00
time (sec)	N/A	0.386	0.183	0.169	0.000	0.085	0.000	0.136	0.187	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	80	73	60	0	141	0	71	91	0
N.S.	1	1.04	0.95	0.78	0.00	1.83	0.00	0.92	1.18	0.00
time (sec)	N/A	0.350	0.094	0.113	0.000	0.105	0.000	0.138	0.249	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	21	20	0	19	46	28	42	19
N.S.	1	1.00	1.00	0.95	0.00	0.90	2.19	1.33	2.00	0.90
time (sec)	N/A	0.264	0.030	0.105	0.000	0.091	0.294	0.128	0.205	0.431

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	75	32	37	46	32	0	81	59	54
N.S.	1	1.56	0.67	0.77	0.96	0.67	0.00	1.69	1.23	1.12
time (sec)	N/A	0.347	0.048	0.159	0.036	0.089	0.000	0.140	0.185	0.491

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	105	45	48	61	45	0	178	84	43
N.S.	1	1.30	0.56	0.59	0.75	0.56	0.00	2.20	1.04	0.53
time (sec)	N/A	0.387	0.062	0.185	0.037	0.082	0.000	0.134	0.184	0.547

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	74	61	77	64	88	0	71	84	0
N.S.	1	0.78	0.64	0.81	0.67	0.93	0.00	0.75	0.88	0.00
time (sec)	N/A	0.418	1.027	0.271	0.037	0.073	0.000	0.131	0.187	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	37	34	43	52	0	33	31	36
N.S.	1	1.00	1.06	0.97	1.23	1.49	0.00	0.94	0.89	1.03
time (sec)	N/A	0.301	1.013	0.141	0.038	0.075	0.000	0.117	0.224	0.478

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	29	29	26	29	40	0	25	28	28
N.S.	1	0.91	0.91	0.81	0.91	1.25	0.00	0.78	0.88	0.88
time (sec)	N/A	0.278	0.005	0.348	0.036	0.074	0.000	0.123	0.189	0.463

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	78	74	102	72	113	0	71	112	0
N.S.	1	0.63	0.60	0.83	0.59	0.92	0.00	0.58	0.91	0.00
time (sec)	N/A	0.414	1.030	0.169	0.039	0.085	0.000	0.126	0.209	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	159	92	96	122	92	141	0	94	136	0
N.S.	1	0.58	0.60	0.77	0.58	0.89	0.00	0.59	0.86	0.00
time (sec)	N/A	0.451	1.033	0.183	0.044	0.084	0.000	0.124	0.204	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	110	96	84	99	68	285	0	74	116	0
N.S.	1	0.87	0.76	0.90	0.62	2.59	0.00	0.67	1.05	0.00
time (sec)	N/A	0.387	1.035	0.145	0.121	0.086	0.000	0.128	0.199	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	111	94	81	101	71	282	0	77	113	0
N.S.	1	0.85	0.73	0.91	0.64	2.54	0.00	0.69	1.02	0.00
time (sec)	N/A	0.379	1.034	0.146	0.122	0.082	0.000	0.126	0.197	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	108	92	83	99	67	282	0	57	116	0
N.S.	1	0.85	0.77	0.92	0.62	2.61	0.00	0.53	1.07	0.00
time (sec)	N/A	0.339	1.033	0.148	0.119	0.089	0.000	0.123	0.203	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	141	112	92	114	83	296	0	69	124	0
N.S.	1	0.79	0.65	0.81	0.59	2.10	0.00	0.49	0.88	0.00
time (sec)	N/A	0.399	1.045	0.247	0.115	0.090	0.000	0.128	0.227	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	179	129	105	141	101	334	0	84	140	0
N.S.	1	0.72	0.59	0.79	0.56	1.87	0.00	0.47	0.78	0.00
time (sec)	N/A	0.435	1.045	0.200	0.123	0.087	0.000	0.122	0.233	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	138	132	63	58	85	75	144	137	192	109
N.S.	1	0.96	0.46	0.42	0.62	0.54	1.04	0.99	1.39	0.79
time (sec)	N/A	0.475	0.053	0.150	0.042	0.098	10.963	0.123	0.191	0.564

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	102	108	52	47	64	63	116	109	160	88
N.S.	1	1.06	0.51	0.46	0.63	0.62	1.14	1.07	1.57	0.86
time (sec)	N/A	0.437	0.035	0.106	0.040	0.093	6.098	0.126	0.192	0.527

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	84	41	36	43	51	87	81	128	67
N.S.	1	1.27	0.62	0.55	0.65	0.77	1.32	1.23	1.94	1.02
time (sec)	N/A	0.390	0.035	0.102	0.034	0.094	2.904	0.121	0.216	0.509

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	31	31	26	25	37	58	17	93	28
N.S.	1	0.86	0.86	0.72	0.69	1.03	1.61	0.47	2.58	0.78
time (sec)	N/A	0.273	0.006	0.339	0.032	0.098	1.396	0.121	0.194	0.487

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	A	F(-1)	F	A	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	117	95	96	0	118	0	0	190	125	0
N.S.	1	0.81	0.82	0.00	1.01	0.00	0.00	1.62	1.07	0.00
time (sec)	N/A	0.395	0.071	0.000	0.116	0.000	0.000	0.125	0.203	0.000

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	A	F(-1)	F	A	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	133	100	107	0	138	0	0	209	144	0
N.S.	1	0.75	0.80	0.00	1.04	0.00	0.00	1.57	1.08	0.00
time (sec)	N/A	0.407	0.112	0.000	0.112	0.000	0.000	0.127	0.200	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	152	139	68	0	0	0	0	0	53	0
N.S.	1	0.91	0.45	0.00	0.00	0.00	0.00	0.00	0.35	0.00
time (sec)	N/A	0.415	6.977	0.000	0.000	0.000	0.000	0.000	0.273	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	119	105	52	0	0	0	0	0	32	0
N.S.	1	0.88	0.44	0.00	0.00	0.00	0.00	0.00	0.27	0.00
time (sec)	N/A	0.344	6.703	0.000	0.000	0.000	0.000	0.000	0.212	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	115	109	55	0	0	0	0	0	48	0
N.S.	1	0.95	0.48	0.00	0.00	0.00	0.00	0.00	0.42	0.00
time (sec)	N/A	0.363	7.209	0.000	0.000	0.000	0.000	0.000	0.205	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	154	139	57	0	0	0	0	0	51	0
N.S.	1	0.90	0.37	0.00	0.00	0.00	0.00	0.00	0.33	0.00
time (sec)	N/A	0.401	10.013	0.000	0.000	0.000	0.000	0.000	0.232	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	136	130	59	58	76	63	144	70	149	91
N.S.	1	0.96	0.43	0.43	0.56	0.46	1.06	0.51	1.10	0.67
time (sec)	N/A	0.480	0.050	0.145	0.034	0.138	10.732	0.129	0.199	0.546

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	106	48	47	59	51	116	56	117	70
N.S.	1	1.06	0.48	0.47	0.59	0.51	1.16	0.56	1.17	0.70
time (sec)	N/A	0.430	0.041	0.103	0.039	0.135	5.766	0.128	0.206	0.546

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	64	82	36	36	42	38	87	40	84	38
N.S.	1	1.28	0.56	0.56	0.66	0.59	1.36	0.62	1.31	0.59
time (sec)	N/A	0.408	0.034	0.103	0.036	0.156	2.812	0.123	0.223	0.536

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	29	25	26	25	28	54	27	59	28
N.S.	1	0.85	0.74	0.76	0.74	0.82	1.59	0.79	1.74	0.82
time (sec)	N/A	0.284	0.009	0.341	0.034	0.115	1.467	0.131	0.219	0.519

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	A	F(-1)	F	B	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	97	84	72	0	110	0	0	200	104	0
N.S.	1	0.87	0.74	0.00	1.13	0.00	0.00	2.06	1.07	0.00
time (sec)	N/A	0.395	0.060	0.000	0.114	0.000	0.000	0.134	0.196	0.000

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	A	F(-1)	F	A	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	143	104	107	0	157	0	0	222	144	0
N.S.	1	0.73	0.75	0.00	1.10	0.00	0.00	1.55	1.01	0.00
time (sec)	N/A	0.412	0.137	0.000	0.117	0.000	0.000	0.131	0.204	0.000

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	132	123	83	0	0	0	0	0	52	0
N.S.	1	0.93	0.63	0.00	0.00	0.00	0.00	0.00	0.39	0.00
time (sec)	N/A	0.416	6.374	0.000	0.000	0.000	0.000	0.000	0.206	0.000

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	97	89	68	0	0	0	0	0	34	0
N.S.	1	0.92	0.70	0.00	0.00	0.00	0.00	0.00	0.35	0.00
time (sec)	N/A	0.377	5.908	0.000	0.000	0.000	0.000	0.000	0.261	0.000

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	62	62	52	0	0	0	0	0	15	0
N.S.	1	1.00	0.84	0.00	0.00	0.00	0.00	0.00	0.24	0.00
time (sec)	N/A	0.291	5.588	0.000	0.000	0.000	0.000	0.000	0.195	0.000

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	95	83	55	0	0	0	0	0	29	0
N.S.	1	0.87	0.58	0.00	0.00	0.00	0.00	0.00	0.31	0.00
time (sec)	N/A	0.364	6.176	0.000	0.000	0.000	0.000	0.000	0.197	0.000

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	134	113	57	0	0	0	0	0	29	0
N.S.	1	0.84	0.43	0.00	0.00	0.00	0.00	0.00	0.22	0.00
time (sec)	N/A	0.401	10.018	0.000	0.000	0.000	0.000	0.000	0.206	0.000

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	82	0	0	0	0	0	118	0
N.S.	1	1.00	1.08	0.00	0.00	0.00	0.00	0.00	1.55	0.00
time (sec)	N/A	0.372	0.406	0.000	0.000	0.000	0.000	0.000	1.486	0.000

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	181	107	133	201	131	235	0	387	182	236
N.S.	1	0.59	0.73	1.11	0.72	1.30	0.00	2.14	1.01	1.30
time (sec)	N/A	0.485	0.102	0.144	0.039	0.116	0.000	0.130	0.235	0.758

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	68	0	0	0	0	0	43	0
N.S.	1	1.00	0.93	0.00	0.00	0.00	0.00	0.00	0.59	0.00
time (sec)	N/A	0.352	0.208	0.000	0.000	0.000	0.000	0.000	0.275	0.000

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	72	0	0	0	0	0	39	0
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	0.51	0.00
time (sec)	N/A	0.340	0.216	0.000	0.000	0.000	0.000	0.000	0.215	0.000

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	63	0	0	0	0	0	44	0
N.S.	1	1.00	0.94	0.00	0.00	0.00	0.00	0.00	0.66	0.00
time (sec)	N/A	0.349	0.039	0.000	0.000	0.000	0.000	0.000	0.194	0.000

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F(-2)	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	82	0	0	0	0	0	0	0
N.S.	1	1.00	1.06	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.380	0.080	0.000	0.000	0.000	0.000	0.000	0.331	0.000

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	A	A	F(-1)	B	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	284	126	92	0	288	546	0	2722	483	0
N.S.	1	0.44	0.32	0.00	1.01	1.92	0.00	9.58	1.70	0.00
time (sec)	N/A	0.516	0.089	0.000	0.046	0.117	0.000	0.241	0.193	0.000

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F(-2)	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	69	0	0	0	0	0	731	0
N.S.	1	1.00	0.93	0.00	0.00	0.00	0.00	0.00	9.88	0.00
time (sec)	N/A	0.393	0.024	0.000	0.000	0.000	0.000	0.000	0.225	0.000

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F(-2)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	72	0	0	0	0	0	43	0
N.S.	1	1.00	0.94	0.00	0.00	0.00	0.00	0.00	0.56	0.00
time (sec)	N/A	0.393	0.033	0.000	0.000	0.000	0.000	0.000	0.238	0.000

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	71	71	66	0	0	0	0	0	48	0
N.S.	1	1.00	0.93	0.00	0.00	0.00	0.00	0.00	0.68	0.00
time (sec)	N/A	0.366	0.025	0.000	0.000	0.000	0.000	0.000	0.216	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [14] had the largest ratio of [.4736839999999999994]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	6	5	0.50	19	0.263
2	A	6	5	0.95	19	0.263
3	A	4	3	0.91	17	0.176
4	A	6	5	0.49	19	0.263
5	A	6	5	0.48	19	0.263
6	A	6	5	0.47	19	0.263
7	A	4	4	0.48	19	0.211
8	A	4	4	0.48	19	0.211
9	A	3	3	0.48	15	0.200
10	A	4	4	0.45	19	0.211
11	A	4	4	0.46	19	0.211
12	A	5	4	1.30	19	0.211
13	A	4	3	0.91	17	0.176
14	A	10	9	0.74	19	0.474
15	A	10	9	0.76	19	0.474
16	A	8	8	0.85	19	0.421
17	A	7	7	0.86	15	0.467
18	A	7	7	0.76	19	0.368
19	A	7	7	0.78	19	0.368
20	A	5	4	1.67	19	0.211
21	A	3	2	1.00	17	0.118

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	6	5	1.00	19	0.263
23	A	7	6	0.95	19	0.316
24	A	4	4	1.02	19	0.211
25	A	3	3	1.04	19	0.158
26	A	2	2	1.00	15	0.133
27	A	3	3	1.56	19	0.158
28	A	4	4	1.30	19	0.211
29	A	6	5	0.78	19	0.263
30	A	3	3	1.00	19	0.158
31	A	4	3	0.91	17	0.176
32	A	6	5	0.63	19	0.263
33	A	6	5	0.58	19	0.263
34	A	5	5	0.87	19	0.263
35	A	5	5	0.85	19	0.263
36	A	4	4	0.85	15	0.267
37	A	6	6	0.79	19	0.316
38	A	7	7	0.72	19	0.368
39	A	5	4	0.96	21	0.190
40	A	5	4	1.06	21	0.190
41	A	5	4	1.27	21	0.190
42	A	4	3	0.86	19	0.158
43	A	10	9	0.81	21	0.429
44	A	10	9	0.75	21	0.429
45	A	5	5	0.91	21	0.238
46	A	4	4	0.88	17	0.235
47	A	4	4	0.95	21	0.190
48	A	5	5	0.90	21	0.238
49	A	5	4	0.96	21	0.190
50	A	5	4	1.06	21	0.190
51	A	5	4	1.28	21	0.190
52	A	4	3	0.85	19	0.158
53	A	7	6	0.87	21	0.286

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
54	A	8	7	0.73	21	0.333
55	A	4	4	0.93	21	0.190
56	A	3	3	0.92	21	0.143
57	A	2	2	1.00	17	0.118
58	A	3	3	0.87	21	0.143
59	A	4	4	0.84	21	0.190
60	A	2	2	1.00	21	0.095
61	A	4	4	0.59	21	0.190
62	A	3	3	1.00	19	0.158
63	A	2	2	1.00	21	0.095
64	A	3	3	1.00	21	0.143
65	A	2	2	1.00	21	0.095
66	A	3	3	0.44	21	0.143
67	A	3	3	1.00	19	0.158
68	A	2	2	1.00	21	0.095
69	A	2	2	1.00	21	0.095

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int x^5 (c(a + bx^2)^2)^{3/2} dx \dots\dots\dots$	55
3.2	$\int x^3 (c(a + bx^2)^2)^{3/2} dx \dots\dots\dots$	61
3.3	$\int x (c(a + bx^2)^2)^{3/2} dx \dots\dots\dots$	67
3.4	$\int \frac{(c(a+bx^2)^2)^{3/2}}{x} dx \dots\dots\dots$	72
3.5	$\int \frac{(c(a+bx^2)^2)^{3/2}}{x^3} dx \dots\dots\dots$	78
3.6	$\int \frac{(c(a+bx^2)^2)^{3/2}}{x^5} dx \dots\dots\dots$	85
3.7	$\int x^4 (c(a + bx^2)^2)^{3/2} dx \dots\dots\dots$	92
3.8	$\int x^2 (c(a + bx^2)^2)^{3/2} dx \dots\dots\dots$	98
3.9	$\int (c(a + bx^2)^2)^{3/2} dx \dots\dots\dots$	104
3.10	$\int \frac{(c(a+bx^2)^2)^{3/2}}{x^2} dx \dots\dots\dots$	109
3.11	$\int \frac{(c(a+bx^2)^2)^{3/2}}{x^4} dx \dots\dots\dots$	115
3.12	$\int x^3 (c(a + bx^2)^3)^{3/2} dx \dots\dots\dots$	121
3.13	$\int x (c(a + bx^2)^3)^{3/2} dx \dots\dots\dots$	127
3.14	$\int \frac{(c(a+bx^2)^3)^{3/2}}{x} dx \dots\dots\dots$	133
3.15	$\int \frac{(c(a+bx^2)^3)^{3/2}}{x^3} dx \dots\dots\dots$	141
3.16	$\int x^2 (c(a + bx^2)^3)^{3/2} dx \dots\dots\dots$	150
3.17	$\int (c(a + bx^2)^3)^{3/2} dx \dots\dots\dots$	158

3.18	$\int \frac{(c(ax^2+b)^3)^{3/2}}{x^2} dx$	165
3.19	$\int \frac{(c(ax^2+b)^3)^{3/2}}{x^4} dx$	173
3.20	$\int x^3 \left(\frac{c}{a+bx^2}\right)^{3/2} dx$	181
3.21	$\int x \left(\frac{c}{a+bx^2}\right)^{3/2} dx$	187
3.22	$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x} dx$	192
3.23	$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^3} dx$	198
3.24	$\int x^4 \left(\frac{c}{a+bx^2}\right)^{3/2} dx$	205
3.25	$\int x^2 \left(\frac{c}{a+bx^2}\right)^{3/2} dx$	211
3.26	$\int \left(\frac{c}{a+bx^2}\right)^{3/2} dx$	217
3.27	$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^2} dx$	222
3.28	$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^4} dx$	228
3.29	$\int x^5 \left(\frac{c}{(a+bx^2)^2}\right)^{3/2} dx$	234
3.30	$\int x^3 \left(\frac{c}{(a+bx^2)^2}\right)^{3/2} dx$	240
3.31	$\int x \left(\frac{c}{(a+bx^2)^2}\right)^{3/2} dx$	245
3.32	$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x} dx$	250
3.33	$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^3} dx$	256
3.34	$\int x^4 \left(\frac{c}{(a+bx^2)^2}\right)^{3/2} dx$	263
3.35	$\int x^2 \left(\frac{c}{(a+bx^2)^2}\right)^{3/2} dx$	269
3.36	$\int \left(\frac{c}{(a+bx^2)^2}\right)^{3/2} dx$	275
3.37	$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^2} dx$	281
3.38	$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^4} dx$	288
3.39	$\int x^7 (c\sqrt{a+bx^2})^{3/2} dx$	296
3.40	$\int x^5 (c\sqrt{a+bx^2})^{3/2} dx$	303
3.41	$\int x^3 (c\sqrt{a+bx^2})^{3/2} dx$	309
3.42	$\int x (c\sqrt{a+bx^2})^{3/2} dx$	315

3.43	$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x} dx$	320
3.44	$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^3} dx$	328
3.45	$\int x^2 (c\sqrt{a+bx^2})^{3/2} dx$	336
3.46	$\int (c\sqrt{a+bx^2})^{3/2} dx$	342
3.47	$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^2} dx$	348
3.48	$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^4} dx$	354
3.49	$\int x^7 \left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} dx$	361
3.50	$\int x^5 \left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} dx$	367
3.51	$\int x^3 \left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} dx$	373
3.52	$\int x \left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} dx$	379
3.53	$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x} dx$	384
3.54	$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^3} dx$	391
3.55	$\int x^4 \left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} dx$	398
3.56	$\int x^2 \left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} dx$	404
3.57	$\int \left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} dx$	409
3.58	$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^2} dx$	414
3.59	$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^4} dx$	420
3.60	$\int (dx)^m (c(a+bx^2)^3)^{3/2} dx$	426
3.61	$\int (dx)^m (c(a+bx^2)^2)^{3/2} dx$	431
3.62	$\int (dx)^m (c(a+bx^2))^{3/2} dx$	438
3.63	$\int (dx)^m \left(\frac{c}{a+bx^2}\right)^{3/2} dx$	443
3.64	$\int (dx)^m \left(\frac{c}{(a+bx^2)^2}\right)^{3/2} dx$	448
3.65	$\int (dx)^m (c(a+bx^n)^3)^{3/2} dx$	453
3.66	$\int (dx)^m (c(a+bx^n)^2)^{3/2} dx$	459
3.67	$\int (dx)^m (c(a+bx^n))^{3/2} dx$	466
3.68	$\int (dx)^m \left(\frac{c}{a+bx^n}\right)^{3/2} dx$	472

3.69 $\int (dx)^m \left(\frac{c}{(a+bx^n)^2} \right)^{3/2} dx \dots\dots\dots 477$

3.1 $\int x^5 \left(c(a + bx^2)^2 \right)^{3/2} dx$

Optimal result	55
Mathematica [A] (verified)	55
Rubi [A] (verified)	56
Maple [A] (verified)	58
Fricas [A] (verification not implemented)	58
Sympy [F(-1)]	59
Maxima [A] (verification not implemented)	59
Giac [A] (verification not implemented)	60
Mupad [F(-1)]	60
Reduce [B] (verification not implemented)	60

Optimal result

Integrand size = 19, antiderivative size = 143

$$\int x^5 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{a^3 cx^6 \sqrt{c(a + bx^2)^2}}{6(a + bx^2)} + \frac{3a^2 bcx^8 \sqrt{c(a + bx^2)^2}}{8(a + bx^2)} + \frac{3ab^2 cx^{10} \sqrt{c(a + bx^2)^2}}{10(a + bx^2)} + \frac{b^3 cx^{12} \sqrt{c(a + bx^2)^2}}{12(a + bx^2)}$$

output

```
a^3*c*x^6*(c*(b*x^2+a)^2)^(1/2)/(6*b*x^2+6*a)+3*a^2*b*c*x^8*(c*(b*x^2+a)^2)^(1/2)/(8*b*x^2+8*a)+3*a*b^2*c*x^10*(c*(b*x^2+a)^2)^(1/2)/(10*b*x^2+10*a)+b^3*c*x^12*(c*(b*x^2+a)^2)^(1/2)/(12*b*x^2+12*a)
```

Mathematica [A] (verified)

Time = 0.80 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.44

$$\int x^5 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{x^6 \left(c(a + bx^2)^2 \right)^{3/2} (20a^3 + 45a^2bx^2 + 36ab^2x^4 + 10b^3x^6)}{120(a + bx^2)^3}$$

input

```
Integrate[x^5*(c*(a + b*x^2)^2)^(3/2),x]
```


output

$$\frac{(x^6*(c*(a + b*x^2)^2)^{(3/2)}*(20*a^3 + 45*a^2*b*x^2 + 36*a*b^2*x^4 + 10*b^3*x^6))/(120*(a + b*x^2)^3)}$$

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.50, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {2045, 27, 243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^5 (c(a + bx^2)^2)^{3/2} dx \\ & \quad \downarrow \text{2045} \\ & \frac{a^3 c \sqrt{c(a + bx^2)^2} \int \frac{x^5 (bx^2 + a)^3}{a^3} dx}{a + bx^2} \\ & \quad \downarrow \text{27} \\ & \frac{c \sqrt{c(a + bx^2)^2} \int x^5 (bx^2 + a)^3 dx}{a + bx^2} \\ & \quad \downarrow \text{243} \\ & \frac{c \sqrt{c(a + bx^2)^2} \int x^4 (bx^2 + a)^3 dx^2}{2(a + bx^2)} \\ & \quad \downarrow \text{49} \\ & \frac{c \sqrt{c(a + bx^2)^2} \int (b^3 x^{10} + 3ab^2 x^8 + 3a^2 b x^6 + a^3 x^4) dx^2}{2(a + bx^2)} \\ & \quad \downarrow \text{2009} \\ & \frac{c \left(\frac{a^3 x^6}{3} + \frac{3}{4} a^2 b x^8 + \frac{3}{5} a b^2 x^{10} + \frac{b^3 x^{12}}{6} \right) \sqrt{c(a + bx^2)^2}}{2(a + bx^2)} \end{aligned}$$

input

$$\text{Int}[x^5*(c*(a + b*x^2)^2)^{(3/2)},x]$$

output
$$\frac{(c\sqrt{c(a + bx^2)^2}((a^3x^6)/3 + (3a^2bx^8)/4 + (3ab^2x^{10})/5 + (b^3x^{12})/6))/(2(a + bx^2))$$

Defintions of rubi rules used

rule 27
$$\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] \text{ ; FreeQ}[b, x]$$

rule 49
$$\text{Int}[(a_*) + (b_*)(x_)^{(m_*)}((c_*) + (d_*)(x_)^{(n_*)}), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + bx)^m(c + dx)^n, x], x] \text{ ; FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{IGtQ}[m + n + 2, 0]$$

rule 243
$$\text{Int}[(x_)^{(m_*)}((a_*) + (b_*)(x_)^2)^{(p_*)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{(m-1)/2}(a + bx)^p, x], x, x^2], x] \text{ ; FreeQ}\{a, b, m, p\}, x \ \&\& \ \text{IntegerQ}[(m-1)/2]$$

rule 2009
$$\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ ; SumQ}[u]$$

rule 2045
$$\text{Int}[(u_)*((c_*)((a_*) + (b_*)(x_)^{(n_*)})^{(q_*)})^{(p_*)}, x_Symbol] \rightarrow \text{Simp}[\text{Simp}[(c*(a + bx^n)^q)^p/(1 + b*(x^n/a))^{(p*q)} \text{ Int}[u*(1 + b*(x^n/a))^{(p*q)}, x], x] \text{ ; FreeQ}\{a, b, c, n, p, q\}, x \ \&\& \ !\text{GeQ}[a, 0]$$

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.42

method	result	size
gospers	$\frac{x^6(10b^3x^6+36ab^2x^4+45a^2bx^2+20a^3)(c(bx^2+a)^2)^{\frac{3}{2}}}{120(bx^2+a)^3}$	60
default	$\frac{x^6(10b^3x^6+36ab^2x^4+45a^2bx^2+20a^3)(c(bx^2+a)^2)^{\frac{3}{2}}}{120(bx^2+a)^3}$	60
orering	$\frac{x^6(10b^3x^6+36ab^2x^4+45a^2bx^2+20a^3)(c(bx^2+a)^2)^{\frac{3}{2}}}{120(bx^2+a)^3}$	60
pseudoelliptic	$\frac{x^6(10b^3x^6+36ab^2x^4+45a^2bx^2+20a^3)c\sqrt{c(bx^2+a)^2}}{120bx^2+120a}$	63
trager	$\frac{cx^6(10b^3x^6+36ab^2x^4+45a^2bx^2+20a^3)\sqrt{b^2cx^4+2abcx^2+a^2c}}{120bx^2+120a}$	72
risch	$\frac{a^3cx^6\sqrt{c(bx^2+a)^2}}{6bx^2+6a} + \frac{3c\sqrt{c(bx^2+a)^2}a^2bx^8}{8(bx^2+a)} + \frac{3c\sqrt{c(bx^2+a)^2}ab^2x^{10}}{10(bx^2+a)} + \frac{b^3cx^{12}\sqrt{c(bx^2+a)^2}}{12bx^2+12a}$	128

input `int(x^5*(c*(b*x^2+a)^2)^(3/2),x,method=_RETURNVERBOSE)`

output `1/120*x^6*(10*b^3*x^6+36*a*b^2*x^4+45*a^2*b*x^2+20*a^3)*(c*(b*x^2+a)^2)^(3/2)/(b*x^2+a)^3`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.52

$$\int x^5 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{(10b^3cx^{12} + 36ab^2cx^{10} + 45a^2bcx^8 + 20a^3cx^6)\sqrt{b^2cx^4 + 2abcx^2 + a^2c}}{120(bx^2 + a)}$$

input `integrate(x^5*(c*(b*x^2+a)^2)^(3/2),x, algorithm="fricas")`

output `1/120*(10*b^3*c*x^12 + 36*a*b^2*c*x^10 + 45*a^2*b*c*x^8 + 20*a^3*c*x^6)*sqrt(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)/(b*x^2 + a)`

Sympy [F(-1)]

Timed out.

$$\int x^5 (c(a + bx^2)^2)^{3/2} dx = \text{Timed out}$$

input `integrate(x**5*(c*(b*x**2+a)**2)**(3/2),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.95

$$\begin{aligned} \int x^5 (c(a + bx^2)^2)^{3/2} dx &= \frac{(b^2cx^4 + 2abcx^2 + a^2c)^{3/2} a^2 x^2}{8b^2} \\ &+ \frac{(b^2cx^4 + 2abcx^2 + a^2c)^{3/2} a^3}{8b^3} + \frac{(b^2cx^4 + 2abcx^2 + a^2c)^{5/2} x^2}{12b^2c} \\ &- \frac{7(b^2cx^4 + 2abcx^2 + a^2c)^{5/2} a}{60b^3c} \end{aligned}$$

input `integrate(x^5*(c*(b*x^2+a)^2)^(3/2),x, algorithm="maxima")`

output `1/8*(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)^(3/2)*a^2*x^2/b^2 + 1/8*(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)^(3/2)*a^3/b^3 + 1/12*(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)^(5/2)*x^2/(b^2*c) - 7/60*(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)^(5/2)*a/(b^3*c)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.50

$$\int x^5 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{1}{120} \left(10b^3x^{12} \operatorname{sgn}(bx^2 + a) + 36ab^2x^{10} \operatorname{sgn}(bx^2 + a) + 45a^2bx^8 \operatorname{sgn}(bx^2 + a) + 20a^3x^6 \operatorname{sgn}(bx^2 + a) \right) c^{3/2}$$

input `integrate(x^5*(c*(b*x^2+a)^2)^(3/2),x, algorithm="giac")`

output `1/120*(10*b^3*x^12*sgn(b*x^2 + a) + 36*a*b^2*x^10*sgn(b*x^2 + a) + 45*a^2*b*x^8*sgn(b*x^2 + a) + 20*a^3*x^6*sgn(b*x^2 + a))*c^(3/2)`

Mupad [F(-1)]

Timed out.

$$\int x^5 \left(c(a + bx^2)^2 \right)^{3/2} dx = \int x^5 \left(c(bx^2 + a)^2 \right)^{3/2} dx$$

input `int(x^5*(c*(a + b*x^2)^2)^(3/2),x)`

output `int(x^5*(c*(a + b*x^2)^2)^(3/2), x)`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.28

$$\int x^5 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{\sqrt{c}cx^6(10b^3x^6 + 36ab^2x^4 + 45a^2bx^2 + 20a^3)}{120}$$

input `int(x^5*(c*(b*x^2+a)^2)^(3/2),x)`

output `(sqrt(c)*c*x**6*(20*a**3 + 45*a**2*b*x**2 + 36*a*b**2*x**4 + 10*b**3*x**6)/120)`

3.2 $\int x^3 \left(c(a + bx^2)^2 \right)^{3/2} dx$

Optimal result	61
Mathematica [A] (verified)	61
Rubi [A] (verified)	62
Maple [A] (verified)	63
Fricas [A] (verification not implemented)	64
Sympy [F(-1)]	65
Maxima [A] (verification not implemented)	65
Giac [A] (verification not implemented)	65
Mupad [B] (verification not implemented)	66
Reduce [B] (verification not implemented)	66

Optimal result

Integrand size = 19, antiderivative size = 66

$$\int x^3 \left(c(a + bx^2)^2 \right)^{3/2} dx = -\frac{ac(a + bx^2)^3 \sqrt{c(a + bx^2)^2}}{8b^2} + \frac{c(a + bx^2)^4 \sqrt{c(a + bx^2)^2}}{10b^2}$$

output

```
-1/8*a*c*(b*x^2+a)^3*(c*(b*x^2+a)^2)^(1/2)/b^2+1/10*c*(b*x^2+a)^4*(c*(b*x^2+a)^2)^(1/2)/b^2
```

Mathematica [A] (verified)

Time = 0.78 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.95

$$\int x^3 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{x^4 \left(c(a + bx^2)^2 \right)^{3/2} (10a^3 + 20a^2bx^2 + 15ab^2x^4 + 4b^3x^6)}{40(a + bx^2)^3}$$

input

```
Integrate[x^3*(c*(a + b*x^2)^2)^(3/2),x]
```

output

```
(x^4*(c*(a + b*x^2)^2)^(3/2)*(10*a^3 + 20*a^2*b*x^2 + 15*a*b^2*x^4 + 4*b^3*x^6))/(40*(a + b*x^2)^3)
```

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.95, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {2045, 27, 243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 (c(a + bx^2)^2)^{3/2} dx \\
 & \quad \downarrow \text{2045} \\
 & \frac{a^3 c \sqrt{c(a + bx^2)^2} \int \frac{x^3 (bx^2 + a)^3}{a^3} dx}{a + bx^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{c \sqrt{c(a + bx^2)^2} \int x^3 (bx^2 + a)^3 dx}{a + bx^2} \\
 & \quad \downarrow \text{243} \\
 & \frac{c \sqrt{c(a + bx^2)^2} \int x^2 (bx^2 + a)^3 dx^2}{2(a + bx^2)} \\
 & \quad \downarrow \text{49} \\
 & \frac{c \sqrt{c(a + bx^2)^2} \int \left(\frac{(bx^2 + a)^4}{b} - \frac{a(bx^2 + a)^3}{b} \right) dx^2}{2(a + bx^2)} \\
 & \quad \downarrow \text{2009} \\
 & \frac{c \left(\frac{(a + bx^2)^5}{5b^2} - \frac{a(a + bx^2)^4}{4b^2} \right) \sqrt{c(a + bx^2)^2}}{2(a + bx^2)}
 \end{aligned}$$

input

$$\text{Int}[x^3(c*(a + b*x^2)^2)^(3/2), x]$$

output

$$\frac{(c*\text{Sqrt}[c*(a + b*x^2)^2]*(-1/4*(a*(a + b*x^2)^4)/b^2 + (a + b*x^2)^5/(5*b^2)))/(2*(a + b*x^2))$$

Defintions of rubi rules used

rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$

rule 49 $\text{Int}[(a_.) + (b_.)(x_)^{(m_.)}((c_.) + (d_.)(x_)^{(n_.)}), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{IGtQ}[m + n + 2, 0]$

rule 243 $\text{Int}[(x_)^{(m_.)}((a_) + (b_.)(x_)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{((m - 1)/2)*(a + b*x)^p}, x], x, x^2], x] /; \text{FreeQ}\{a, b, m, p\}, x \ \&\& \ \text{IntegerQ}[(m - 1)/2]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 2045 $\text{Int}[(u_.)*((c_.)*((a_) + (b_.)(x_)^{(n_.)})^{(q_.)})^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[\text{Simp}[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^{(p*q)} \text{ Int}[u*(1 + b*(x^n/a))^{(p*q)}, x], x] /; \text{FreeQ}\{a, b, c, n, p, q\}, x \ \&\& \ !\text{GeQ}[a, 0]$

Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.91

method	result	size
gospers	$\frac{x^4(4b^3x^6+15ab^2x^4+20a^2bx^2+10a^3)(c(bx^2+a)^2)^{\frac{3}{2}}}{40(bx^2+a)^3}$	60
default	$\frac{x^4(4b^3x^6+15ab^2x^4+20a^2bx^2+10a^3)(c(bx^2+a)^2)^{\frac{3}{2}}}{40(bx^2+a)^3}$	60
orering	$\frac{x^4(4b^3x^6+15ab^2x^4+20a^2bx^2+10a^3)(c(bx^2+a)^2)^{\frac{3}{2}}}{40(bx^2+a)^3}$	60
pseudoelliptic	$\frac{x^4(4b^3x^6+15ab^2x^4+20a^2bx^2+10a^3)c\sqrt{c(bx^2+a)^2}}{40bx^2+40a}$	63
trager	$\frac{cx^4(4b^3x^6+15ab^2x^4+20a^2bx^2+10a^3)\sqrt{b^2cx^4+2abcx^2+a^2c}}{40bx^2+40a}$	72
risch	$\frac{c\sqrt{c(bx^2+a)^2}b^3x^{10}}{10bx^2+10a} + \frac{3c\sqrt{c(bx^2+a)^2}ab^2x^8}{8(bx^2+a)} + \frac{c\sqrt{c(bx^2+a)^2}a^2bx^6}{2bx^2+2a} + \frac{c\sqrt{c(bx^2+a)^2}a^3x^4}{4bx^2+4a}$	128

input `int(x^3*(c*(b*x^2+a)^2)^(3/2),x,method=_RETURNVERBOSE)`

output $\frac{1}{40}x^4(4b^3x^6+15ab^2x^4+20a^2bx^2+10a^3)(c(bx^2+a)^2)^{3/2}/(bx^2+a)^3$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.12

$$\int x^3 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{(4b^3cx^{10} + 15ab^2cx^8 + 20a^2bcx^6 + 10a^3cx^4)\sqrt{b^2cx^4 + 2abcx^2 + a^2c}}{40(bx^2 + a)}$$

input `integrate(x^3*(c*(b*x^2+a)^2)^(3/2),x, algorithm="fricas")`

output $\frac{1}{40}(4b^3c*x^{10} + 15*a*b^2*c*x^8 + 20*a^2*b*c*x^6 + 10*a^3*c*x^4)*\text{sqrt}(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)/(b*x^2 + a)$

Sympy [F(-1)]

Timed out.

$$\int x^3 (c(a + bx^2)^2)^{3/2} dx = \text{Timed out}$$

input `integrate(x**3*(c*(b*x**2+a)**2)**(3/2),x)`output `Timed out`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.48

$$\int x^3 (c(a + bx^2)^2)^{3/2} dx = -\frac{(b^2cx^4 + 2abcx^2 + a^2c)^{\frac{3}{2}}ax^2}{8b} - \frac{(b^2cx^4 + 2abcx^2 + a^2c)^{\frac{3}{2}}a^2}{8b^2} + \frac{(b^2cx^4 + 2abcx^2 + a^2c)^{\frac{5}{2}}}{10b^2c}$$

input `integrate(x^3*(c*(b*x^2+a)^2)^(3/2),x, algorithm="maxima")`output `-1/8*(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)^(3/2)*a*x^2/b - 1/8*(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)^(3/2)*a^2/b^2 + 1/10*(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)^(5/2)/(b^2*c)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.73

$$\int x^3 (c(a + bx^2)^2)^{3/2} dx = \frac{1}{40} (4b^3x^{10} + 15ab^2x^8 + 20a^2bx^6 + 10a^3x^4)c^{\frac{3}{2}}\text{sgn}(bx^2 + a)$$

input `integrate(x^3*(c*(b*x^2+a)^2)^(3/2),x, algorithm="giac")`

output $1/40*(4*b^3*x^{10} + 15*a*b^2*x^8 + 20*a^2*b*x^6 + 10*a^3*x^4)*c^{(3/2)}*sgn(b*x^2 + a)$

Mupad [B] (verification not implemented)

Time = 0.61 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.76

$$\int x^3 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{(-a^2 + 3abx^2 + 4b^2x^4)(ca^2 + 2cabx^2 + cb^2x^4)^{3/2}}{40b^2}$$

input $\text{int}(x^3*(c*(a + b*x^2)^2)^{(3/2)},x)$

output $((4*b^2*x^4 - a^2 + 3*a*b*x^2)*(a^2*c + b^2*c*x^4 + 2*a*b*c*x^2)^{(3/2)})/(40*b^2)$

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.61

$$\int x^3 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{\sqrt{c} c x^4 (4b^3 x^6 + 15a b^2 x^4 + 20a^2 b x^2 + 10a^3)}{40}$$

input $\text{int}(x^3*(c*(b*x^2+a)^2)^{(3/2)},x)$

output $(\text{sqrt}(c)*c*x**4*(10*a**3 + 20*a**2*b*x**2 + 15*a*b**2*x**4 + 4*b**3*x**6))/40$

3.3 $\int x \left(c(a + bx^2)^2 \right)^{3/2} dx$

Optimal result	67
Mathematica [A] (verified)	67
Rubi [A] (verified)	68
Maple [A] (verified)	69
Fricas [B] (verification not implemented)	69
Sympy [F]	70
Maxima [B] (verification not implemented)	70
Giac [A] (verification not implemented)	71
Mupad [B] (verification not implemented)	71
Reduce [B] (verification not implemented)	71

Optimal result

Integrand size = 17, antiderivative size = 32

$$\int x \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{c(a + bx^2)^3 \sqrt{c(a + bx^2)^2}}{8b}$$

output $1/8*c*(b*x^2+a)^3*(c*(b*x^2+a)^2)^{(1/2)}/b$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.91

$$\int x \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{(a + bx^2) \left(c(a + bx^2)^2 \right)^{3/2}}{8b}$$

input `Integrate[x*(c*(a + b*x^2)^2)^(3/2),x]`

output $((a + b*x^2)*(c*(a + b*x^2)^2)^{(3/2)})/(8*b)$

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.91, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2024, 20, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \left(c(a + bx^2)^2 \right)^{3/2} dx \\
 & \quad \downarrow \text{2024} \\
 & \frac{\int \left(c(bx^2 + a)^2 \right)^{3/2} d(bx^2 + a)}{2b} \\
 & \quad \downarrow \text{20} \\
 & \frac{\left(c(a + bx^2)^2 \right)^{3/2} \int (bx^2 + a)^3 d(bx^2 + a)}{2b(a + bx^2)^3} \\
 & \quad \downarrow \text{15} \\
 & \frac{(a + bx^2) \left(c(a + bx^2)^2 \right)^{3/2}}{8b}
 \end{aligned}$$

input `Int[x*(c*(a + b*x^2)^2)^(3/2),x]`

output `((a + b*x^2)*(c*(a + b*x^2)^2)^(3/2))/(8*b)`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 20 `Int[((a_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a*x^n)^p/x^(n*p) Int[x^(n*p), x], x] /; FreeQ[{a, n, p}, x] && !IntegerQ[p]`

rule 2024

```
Int[((a_.) + (b_.)*(Pq_)^(n_.))^(p_.)*(Qr_), x_Symbol] := With[{q = Expon[P
q, x], r = Expon[Qr, x]}, Simp[Coeff[Qr, x, r]/(q*Coeff[Pq, x, q]) Subst[
Int[(a + b*x^n)^p, x], x, Pq], x] /; EqQ[r, q - 1] && EqQ[Coeff[Qr, x, r]*D
[Pq, x], q*Coeff[Pq, x, q]*Qr]] /; FreeQ[{a, b, n, p}, x] && PolyQ[Pq, x] &
& PolyQ[Qr, x]
```

Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.81

method	result	size
default	$\frac{(c(bx^2+a)^2)^{\frac{3}{2}}(bx^2+a)}{8b}$	26
risch	$\frac{c(bx^2+a)^3\sqrt{c(bx^2+a)^2}}{8b}$	29
gospers	$\frac{x^2(b^3x^6+4ab^2x^4+6a^2bx^2+4a^3)(c(bx^2+a)^2)^{\frac{3}{2}}}{8(bx^2+a)^3}$	59
orering	$\frac{x^2(b^3x^6+4ab^2x^4+6a^2bx^2+4a^3)(c(bx^2+a)^2)^{\frac{3}{2}}}{8(bx^2+a)^3}$	59
pseudoelliptic	$\frac{x^2(b^3x^6+4ab^2x^4+6a^2bx^2+4a^3)c\sqrt{c(bx^2+a)^2}}{8bx^2+8a}$	62
trager	$\frac{cx^2(b^3x^6+4ab^2x^4+6a^2bx^2+4a^3)\sqrt{b^2cx^4+2abcx^2+a^2c}}{8bx^2+8a}$	71

input

```
int(x*(c*(b*x^2+a)^2)^(3/2),x,method=_RETURNVERBOSE)
```

output

```
1/8*(c*(b*x^2+a)^2)^(3/2)*(b*x^2+a)/b
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 73 vs. 2(28) = 56.

Time = 0.08 (sec) , antiderivative size = 73, normalized size of antiderivative = 2.28

$$\int x(c(a+bx^2)^2)^{3/2} dx = \frac{(b^3cx^8 + 4ab^2cx^6 + 6a^2bcx^4 + 4a^3cx^2)\sqrt{b^2cx^4 + 2abcx^2 + a^2c}}{8(bx^2 + a)}$$

input `integrate(x*(c*(b*x^2+a)^2)^(3/2),x, algorithm="fricas")`

output `1/8*(b^3*c*x^8 + 4*a*b^2*c*x^6 + 6*a^2*b*c*x^4 + 4*a^3*c*x^2)*sqrt(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)/(b*x^2 + a)`

Sympy [F]

$$\int x \left(c(a + bx^2)^2 \right)^{3/2} dx = \int x \left(c(a + bx^2)^2 \right)^{\frac{3}{2}} dx$$

input `integrate(x*(c*(b*x**2+a)**2)**(3/2),x)`

output `Integral(x*(c*(a + b*x**2)**2)**(3/2), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 60 vs. 2(28) = 56.

Time = 0.04 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.88

$$\int x \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{1}{8} (b^2cx^4 + 2abcx^2 + a^2c)^{\frac{3}{2}}x^2 + \frac{(b^2cx^4 + 2abcx^2 + a^2c)^{\frac{3}{2}}a}{8b}$$

input `integrate(x*(c*(b*x^2+a)^2)^(3/2),x, algorithm="maxima")`

output `1/8*(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)^(3/2)*x^2 + 1/8*(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)^(3/2)*a/b`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.78

$$\int x \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{(bx^2 + a)^4 c^{3/2} \operatorname{sgn}(bx^2 + a)}{8b}$$

input `integrate(x*(c*(b*x^2+a)^2)^(3/2),x, algorithm="giac")`

output `1/8*(b*x^2 + a)^4*c^(3/2)*sgn(b*x^2 + a)/b`

Mupad [B] (verification not implemented)

Time = 0.64 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.25

$$\int x \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{(b^2 x^2 + a b) (c a^2 + 2 c a b x^2 + c b^2 x^4)^{3/2}}{8 b^2}$$

input `int(x*(c*(a + b*x^2)^2)^(3/2),x)`

output `((a*b + b^2*x^2)*(a^2*c + b^2*c*x^4 + 2*a*b*c*x^2)^(3/2))/(8*b^2)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.22

$$\int x \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{\sqrt{c} c x^2 (b^3 x^6 + 4a b^2 x^4 + 6a^2 b x^2 + 4a^3)}{8}$$

input `int(x*(c*(b*x^2+a)^2)^(3/2),x)`

output `(sqrt(c)*c*x**2*(4*a**3 + 6*a**2*b*x**2 + 4*a*b**2*x**4 + b**3*x**6))/8`

3.4
$$\int \frac{\left(c(a+bx^2)^2\right)^{3/2}}{x} dx$$

Optimal result	72
Mathematica [A] (verified)	72
Rubi [A] (verified)	73
Maple [A] (verified)	75
Fricas [A] (verification not implemented)	75
Sympy [F]	76
Maxima [A] (verification not implemented)	76
Giac [A] (verification not implemented)	77
Mupad [F(-1)]	77
Reduce [B] (verification not implemented)	77

Optimal result

Integrand size = 19, antiderivative size = 139

$$\int \frac{\left(c(a+bx^2)^2\right)^{3/2}}{x} dx = \frac{3a^2bcx^2\sqrt{c(a+bx^2)^2}}{2(a+bx^2)} + \frac{3ab^2cx^4\sqrt{c(a+bx^2)^2}}{4(a+bx^2)} + \frac{b^3cx^6\sqrt{c(a+bx^2)^2}}{6(a+bx^2)} + \frac{a^3c\sqrt{c(a+bx^2)^2}\log(x)}{a+bx^2}$$

output `3*a^2*b*c*x^2*(c*(b*x^2+a)^2)^(1/2)/(2*b*x^2+2*a)+3*a*b^2*c*x^4*(c*(b*x^2+a)^2)^(1/2)/(4*b*x^2+4*a)+b^3*c*x^6*(c*(b*x^2+a)^2)^(1/2)/(6*b*x^2+6*a)+a^3*c*(c*(b*x^2+a)^2)^(1/2)*ln(x)/(b*x^2+a)`

Mathematica [A] (verified)

Time = 0.50 (sec) , antiderivative size = 234, normalized size of antiderivative = 1.68

$$\int \frac{\left(c(a+bx^2)^2\right)^{3/2}}{x} dx = \frac{c\left(b\left(2b^2x^4\left(-\sqrt{b^2cx^2} + \sqrt{c(a+bx^2)^2}\right) + abx^2\left(-9\sqrt{b^2cx^2} + 7\sqrt{c(a+bx^2)^2}\right)\right)}{x}$$

input `Integrate[(c*(a + b*x^2)^2)^(3/2)/x,x]`

output

```
(c*(b*(2*b^2*x^4*(-(Sqrt[b^2*c]*x^2) + Sqrt[c*(a + b*x^2)^2]) + a*b*x^2*(-
9*Sqrt[b^2*c]*x^2 + 7*Sqrt[c*(a + b*x^2)^2]) + a^2*(-18*Sqrt[b^2*c]*x^2 +
11*Sqrt[c*(a + b*x^2)^2])) + 12*a^3*b*Sqrt[c]*ArcTanh[(Sqrt[b^2*c]*x^2 - S
qrt[c*(a + b*x^2)^2])/(a*Sqrt[c])] - 6*a^3*Sqrt[b^2*c]*Log[x^2*(a*b*c + b^
2*c*x^2 - Sqrt[b^2*c]*Sqrt[c*(a + b*x^2)^2])])/(24*b)
```

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.49, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {2045, 27, 243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c(a + bx^2)^2)^{3/2}}{x} dx \\
 & \quad \downarrow \text{2045} \\
 & \frac{a^3 c \sqrt{c(a + bx^2)^2} \int \frac{(bx^2+a)^3}{a^3 x} dx}{a + bx^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{c \sqrt{c(a + bx^2)^2} \int \frac{(bx^2+a)^3}{x} dx}{a + bx^2} \\
 & \quad \downarrow \text{243} \\
 & \frac{c \sqrt{c(a + bx^2)^2} \int \frac{(bx^2+a)^3}{x^2} dx^2}{2(a + bx^2)} \\
 & \quad \downarrow \text{49} \\
 & \frac{c \sqrt{c(a + bx^2)^2} \int (b^3 x^4 + 3ab^2 x^2 + 3a^2 b + \frac{a^3}{x^2}) dx^2}{2(a + bx^2)} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$\frac{c\sqrt{c(a+bx^2)^2}\left(a^3\log(x^2)+3a^2bx^2+\frac{3}{2}ab^2x^4+\frac{b^3x^6}{3}\right)}{2(a+bx^2)}$$

input `Int[(c*(a + b*x^2)^2)^(3/2)/x,x]`

output `(c*Sqrt[c*(a + b*x^2)^2]*(3*a^2*b*x^2 + (3*a*b^2*x^4)/2 + (b^3*x^6)/3 + a^3*Log[x^2]))/(2*(a + b*x^2))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 49 `Int[((a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 243 `Int[(x_.)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_.))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.42

method	result	size
default	$\frac{(c(bx^2+a)^2)^{\frac{3}{2}}(2b^3x^6+9ab^2x^4+18a^2bx^2+12a^3\ln(x))}{12(bx^2+a)^3}$	59
pseudoelliptic	$\frac{c\sqrt{c(bx^2+a)^2}(2b^3x^6+9ab^2x^4+6a^3\ln(x^2)+18a^2bx^2)}{12bx^2+12a}$	64
risch	$\frac{c\sqrt{c(bx^2+a)^2}b(\frac{1}{6}b^2x^6+\frac{3}{4}abx^4+\frac{3}{2}a^2x^2)}{bx^2+a} + \frac{a^3c\sqrt{c(bx^2+a)^2}\ln(x)}{bx^2+a}$	80

input `int((c*(b*x^2+a)^2)^(3/2)/x,x,method=_RETURNVERBOSE)`

output `1/12*(c*(b*x^2+a)^2)^(3/2)*(2*b^3*x^6+9*a*b^2*x^4+18*a^2*b*x^2+12*a^3*ln(x))/ (b*x^2+a)^3`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.53

$$\int \frac{(c(a+bx^2)^2)^{3/2}}{x} dx = \frac{(2b^3cx^6 + 9ab^2cx^4 + 18a^2bcx^2 + 12a^3c \log(x))\sqrt{b^2cx^4 + 2abcx^2 + a^2c}}{12(bx^2 + a)}$$

input `integrate((c*(b*x^2+a)^2)^(3/2)/x,x, algorithm="fricas")`

output `1/12*(2*b^3*c*x^6 + 9*a*b^2*c*x^4 + 18*a^2*b*c*x^2 + 12*a^3*c*log(x))*sqrt (b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)/(b*x^2 + a)`

Sympy [F]

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x} dx = \int \frac{(c(a + bx^2)^2)^{\frac{3}{2}}}{x} dx$$

input `integrate((c*(b*x**2+a)**2)**(3/2)/x,x)`

output `Integral((c*(a + b*x**2)**2)**(3/2)/x, x)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.23

$$\begin{aligned} \int \frac{(c(a + bx^2)^2)^{3/2}}{x} dx &= \frac{1}{2} (-1)^{2b^2cx^2+2abc} a^3 c^{\frac{3}{2}} \log(2b^2cx^2 + 2abc) \\ &- \frac{1}{2} (-1)^{2abcx^2+2a^2c} a^3 c^{\frac{3}{2}} \log\left(2abc + \frac{2a^2c}{x^2}\right) + \frac{1}{4} \sqrt{b^2cx^4 + 2abcx^2 + a^2cabx^2} \\ &+ \frac{3}{4} \sqrt{b^2cx^4 + 2abcx^2 + a^2ca^2c} + \frac{1}{6} (b^2cx^4 + 2abcx^2 + a^2c)^{\frac{3}{2}} \end{aligned}$$

input `integrate((c*(b*x^2+a)^2)^(3/2)/x,x, algorithm="maxima")`

output `1/2*(-1)^(2*b^2*c*x^2 + 2*a*b*c)*a^3*c^(3/2)*log(2*b^2*c*x^2 + 2*a*b*c) -
1/2*(-1)^(2*a*b*c*x^2 + 2*a^2*c)*a^3*c^(3/2)*log(2*a*b*c + 2*a^2*c/x^2) +
1/4*sqrt(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)*a*b*c*x^2 + 3/4*sqrt(b^2*c*x^4 +
2*a*b*c*x^2 + a^2*c)*a^2*c + 1/6*(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)^(3/2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.53

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x} dx = \frac{1}{12} (2b^3x^6 \operatorname{sgn}(bx^2 + a) + 9ab^2x^4 \operatorname{sgn}(bx^2 + a) + 18a^2bx^2 \operatorname{sgn}(bx^2 + a) + 6a^3 \log(x)) c^{3/2}$$

input `integrate((c*(b*x^2+a)^2)^(3/2)/x,x, algorithm="giac")`

output `1/12*(2*b^3*x^6*sgn(b*x^2 + a) + 9*a*b^2*x^4*sgn(b*x^2 + a) + 18*a^2*b*x^2*sgn(b*x^2 + a) + 6*a^3*log(x^2)*sgn(b*x^2 + a))*c^(3/2)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x} dx = \int \frac{(c(bx^2 + a)^2)^{3/2}}{x} dx$$

input `int((c*(a + b*x^2)^2)^(3/2)/x,x)`

output `int((c*(a + b*x^2)^2)^(3/2)/x, x)`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.28

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x} dx = \frac{\sqrt{c}c(12 \log(x) a^3 + 18a^2bx^2 + 9ab^2x^4 + 2b^3x^6)}{12}$$

input `int((c*(b*x^2+a)^2)^(3/2)/x,x)`

output `(sqrt(c)*c*(12*log(x)*a**3 + 18*a**2*b*x**2 + 9*a*b**2*x**4 + 2*b**3*x**6))/12`

3.5
$$\int \frac{(c(a+bx^2)^2)^{3/2}}{x^3} dx$$

Optimal result	78
Mathematica [A] (verified)	79
Rubi [A] (verified)	79
Maple [A] (verified)	81
Fricas [A] (verification not implemented)	82
Sympy [F]	82
Maxima [A] (verification not implemented)	82
Giac [A] (verification not implemented)	83
Mupad [F(-1)]	83
Reduce [B] (verification not implemented)	84

Optimal result

Integrand size = 19, antiderivative size = 140

$$\int \frac{(c(a+bx^2)^2)^{3/2}}{x^3} dx = -\frac{a^3c\sqrt{c(a+bx^2)^2}}{2x^2(a+bx^2)} + \frac{3ab^2cx^2\sqrt{c(a+bx^2)^2}}{2(a+bx^2)} + \frac{b^3cx^4\sqrt{c(a+bx^2)^2}}{4(a+bx^2)} + \frac{3a^2bc\sqrt{c(a+bx^2)^2}\log(x)}{a+bx^2}$$

output

```
-1/2*a^3*c*(c*(b*x^2+a)^2)^(1/2)/x^2/(b*x^2+a)+3*a*b^2*c*x^2*(c*(b*x^2+a)^2)^(1/2)/(2*b*x^2+2*a)+b^3*c*x^4*(c*(b*x^2+a)^2)^(1/2)/(4*b*x^2+4*a)+3*a^2*c*b*c*(c*(b*x^2+a)^2)^(1/2)*ln(x)/(b*x^2+a)
```

Mathematica [A] (verified)

Time = 0.68 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.69

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^3} dx = \frac{1}{16}c \left(\frac{(8a^3 + 21a^2bx^2 - 24ab^2x^4 - 4b^3x^6) \left(abc + b^2cx^2 - \sqrt{b^2c}\sqrt{c(a + bx^2)^2} \right)}{x^2 \left(a\sqrt{b^2c} + b\sqrt{b^2cx^2} - b\sqrt{c(a + bx^2)^2} \right)} \right. \\ \left. + 24a^2b\sqrt{c}\operatorname{arctanh} \left(\frac{\sqrt{b^2cx^2} - \sqrt{c(a + bx^2)^2}}{a\sqrt{c}} \right) \right. \\ \left. - 12a^2\sqrt{b^2c} \log \left(x^2 \left(abc + b^2cx^2 - \sqrt{b^2c}\sqrt{c(a + bx^2)^2} \right) \right) \right)$$

input `Integrate[(c*(a + b*x^2)^2)^(3/2)/x^3,x]`

output `(c*(((8*a^3 + 21*a^2*b*x^2 - 24*a*b^2*x^4 - 4*b^3*x^6)*(a*b*c + b^2*c*x^2 - Sqrt[b^2*c]*Sqrt[c*(a + b*x^2)^2]))/(x^2*(a*Sqrt[b^2*c] + b*Sqrt[b^2*c]*x^2 - b*Sqrt[c*(a + b*x^2)^2])) + 24*a^2*b*Sqrt[c]*ArcTanh[(Sqrt[b^2*c]*x^2 - Sqrt[c*(a + b*x^2)^2])/(a*Sqrt[c])] - 12*a^2*Sqrt[b^2*c]*Log[x^2*(a*b*c + b^2*c*x^2 - Sqrt[b^2*c]*Sqrt[c*(a + b*x^2)^2])]))/16`

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.48, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {2045, 27, 243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^3} dx$$

↓ 2045

$$\begin{aligned}
& \frac{a^3 c \sqrt{c(a+bx^2)^2} \int \frac{(bx^2+a)^3}{a^3 x^3} dx}{a+bx^2} \\
& \quad \downarrow \text{27} \\
& \frac{c \sqrt{c(a+bx^2)^2} \int \frac{(bx^2+a)^3}{x^3} dx}{a+bx^2} \\
& \quad \downarrow \text{243} \\
& \frac{c \sqrt{c(a+bx^2)^2} \int \frac{(bx^2+a)^3}{x^4} dx^2}{2(a+bx^2)} \\
& \quad \downarrow \text{49} \\
& \frac{c \sqrt{c(a+bx^2)^2} \int \left(\frac{a^3}{x^4} + \frac{3ba^2}{x^2} + 3b^2a + b^3x^2 \right) dx^2}{2(a+bx^2)} \\
& \quad \downarrow \text{2009} \\
& \frac{c \sqrt{c(a+bx^2)^2} \left(-\frac{a^3}{x^2} + 3a^2b \log(x^2) + 3ab^2x^2 + \frac{b^3x^4}{2} \right)}{2(a+bx^2)}
\end{aligned}$$

input `Int[(c*(a + b*x^2)^2)^(3/2)/x^3,x]`

output `(c*Sqrt[c*(a + b*x^2)^2]*(-(a^3/x^2) + 3*a*b^2*x^2 + (b^3*x^4)/2 + 3*a^2*b*Log[x^2]))/(2*(a + b*x^2))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_.))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.44

method	result	size
default	$\frac{(c(bx^2+a)^2)^{\frac{3}{2}}(b^3x^6+6ab^2x^4+12a^2b\ln(x)x^2-2a^3)}{4x^2(bx^2+a)^3}$	61
pseudoelliptic	$-\frac{c\left(-\frac{b^3x^6}{2}-3ab^2x^4-3a^2b\ln(x^2)x^2+a^3\right)\sqrt{c(bx^2+a)^2}}{2(bx^2+a)x^2}$	63
risch	$\frac{c\sqrt{c(bx^2+a)^2}b(bx^2+3a)^2}{4bx^2+4a} - \frac{a^3c\sqrt{c(bx^2+a)^2}}{2x^2(bx^2+a)} + \frac{3a^2bc\sqrt{c(bx^2+a)^2}\ln(x)}{bx^2+a}$	101

input `int((c*(b*x^2+a)^2)^(3/2)/x^3,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{4}*(c*(b*x^2+a)^2)^(3/2)*(b^3*x^6+6*a*b^2*x^4+12*a^2*b*\ln(x)*x^2-2*a^3)/x^2/(b*x^2+a)^3$$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.54

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^3} dx = \frac{(b^3cx^6 + 6ab^2cx^4 + 12a^2bcx^2 \log(x) - 2a^3c)\sqrt{b^2cx^4 + 2abcx^2 + a^2c}}{4(bx^4 + ax^2)}$$

input `integrate((c*(b*x^2+a)^2)^(3/2)/x^3,x, algorithm="fricas")`output `1/4*(b^3*c*x^6 + 6*a*b^2*c*x^4 + 12*a^2*b*c*x^2*log(x) - 2*a^3*c)*sqrt(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)/(b*x^4 + a*x^2)`**Sympy [F]**

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^3} dx = \int \frac{(c(a + bx^2)^2)^{\frac{3}{2}}}{x^3} dx$$

input `integrate((c*(b*x**2+a)**2)**(3/2)/x**3,x)`output `Integral((c*(a + b*x**2)**2)**(3/2)/x**3, x)`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.26

$$\begin{aligned} \int \frac{(c(a + bx^2)^2)^{3/2}}{x^3} dx &= \frac{3}{2} (-1)^{2b^2cx^2+2abc} a^2bc^{\frac{3}{2}} \log(2b^2cx^2 + 2abc) \\ &- \frac{3}{2} (-1)^{2abcx^2+2a^2c} a^2bc^{\frac{3}{2}} \log\left(2abc + \frac{2a^2c}{x^2}\right) + \frac{3}{4} \sqrt{b^2cx^4 + 2abcx^2 + a^2cb^2cx^2} \\ &+ \frac{9}{4} \sqrt{b^2cx^4 + 2abcx^2 + a^2c} abc - \frac{(b^2cx^4 + 2abcx^2 + a^2c)^{\frac{3}{2}}}{2x^2} \end{aligned}$$

input `integrate((c*(b*x^2+a)^2)^(3/2)/x^3,x, algorithm="maxima")`

output
$$\begin{aligned} & 3/2*(-1)^(2*b^2*c*x^2 + 2*a*b*c)*a^2*b*c^(3/2)*\log(2*b^2*c*x^2 + 2*a*b*c) \\ & - 3/2*(-1)^(2*a*b*c*x^2 + 2*a^2*c)*a^2*b*c^(3/2)*\log(2*a*b*c + 2*a^2*c/x^2) \\ & + 3/4*\sqrt{b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c}*b^2*c*x^2 + 9/4*\sqrt{b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c} \\ & *a*b*c - 1/2*(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)^(3/2)/x^2 \end{aligned}$$

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.65

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^3} dx = \frac{1}{4} \left(b^3 x^4 \operatorname{sgn}(bx^2 + a) + 6 ab^2 x^2 \operatorname{sgn}(bx^2 + a) + 6 a^2 b \log(x^2) \operatorname{sgn}(bx^2 + a) - \frac{2}{3} \right)$$

input `integrate((c*(b*x^2+a)^2)^(3/2)/x^3,x, algorithm="giac")`

output
$$\begin{aligned} & 1/4*(b^3*x^4*\operatorname{sgn}(b*x^2 + a) + 6*a*b^2*x^2*\operatorname{sgn}(b*x^2 + a) + 6*a^2*b*\log(x^2) \\ &)*\operatorname{sgn}(b*x^2 + a) - 2*(3*a^2*b*x^2*\operatorname{sgn}(b*x^2 + a) + a^3*\operatorname{sgn}(b*x^2 + a))/x^2 \\ &)*c^(3/2) \end{aligned}$$

Mupad [F(-1)]

Timed out.

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^3} dx = \int \frac{(c(bx^2 + a)^2)^{3/2}}{x^3} dx$$

input `int((c*(a + b*x^2)^2)^(3/2)/x^3,x)`

output `int((c*(a + b*x^2)^2)^(3/2)/x^3, x)`

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.29

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^3} dx = \frac{\sqrt{c}c(12 \log(x) a^2 b x^2 - 2a^3 + 6a b^2 x^4 + b^3 x^6)}{4x^2}$$

input `int((c*(b*x^2+a)^2)^(3/2)/x^3,x)`output `(sqrt(c)*c*(12*log(x)*a**2*b*x**2 - 2*a**3 + 6*a*b**2*x**4 + b**3*x**6))/(4*x**2)`

3.6 $\int \frac{(c(a+bx^2)^2)^{3/2}}{x^5} dx$

Optimal result	85
Mathematica [A] (verified)	86
Rubi [A] (verified)	86
Maple [A] (verified)	88
Fricas [A] (verification not implemented)	89
Sympy [F]	89
Maxima [A] (verification not implemented)	89
Giac [A] (verification not implemented)	90
Mupad [F(-1)]	90
Reduce [B] (verification not implemented)	91

Optimal result

Integrand size = 19, antiderivative size = 140

$$\int \frac{(c(a+bx^2)^2)^{3/2}}{x^5} dx = -\frac{a^3c\sqrt{c(a+bx^2)^2}}{4x^4(a+bx^2)} - \frac{3a^2bc\sqrt{c(a+bx^2)^2}}{2x^2(a+bx^2)} + \frac{b^3cx^2\sqrt{c(a+bx^2)^2}}{2(a+bx^2)} + \frac{3ab^2c\sqrt{c(a+bx^2)^2} \log(x)}{a+bx^2}$$

output

```
-1/4*a^3*c*(c*(b*x^2+a)^2)^(1/2)/x^4/(b*x^2+a)-3/2*a^2*b*c*(c*(b*x^2+a)^2)^(1/2)/x^2/(b*x^2+a)+b^3*c*x^2*(c*(b*x^2+a)^2)^(1/2)/(2*b*x^2+2*a)+3*a*b^2*c*(c*(b*x^2+a)^2)^(1/2)*ln(x)/(b*x^2+a)
```

Mathematica [A] (verified)

Time = 0.66 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.69

$$\int \frac{(c(a+bx^2)^2)^{3/2}}{x^5} dx = \frac{1}{4}c \left(-\frac{c(-a^3 - 6a^2bx^2 + ab^2x^4 + 2b^3x^6) \left(a\sqrt{b^2c} + b\sqrt{b^2cx^2} - b\sqrt{c(a+bx^2)^2} \right)}{x^4 \left(abc + b^2cx^2 - \sqrt{b^2c}\sqrt{c(a+bx^2)^2} \right)} \right. \\ \left. + 6ab^2\sqrt{c} \operatorname{arctanh} \left(\frac{\sqrt{b^2cx^2} - \sqrt{c(a+bx^2)^2}}{a\sqrt{c}} \right) \right. \\ \left. - 3ab\sqrt{b^2c} \log \left(x^2 \left(abc + b^2cx^2 - \sqrt{b^2c}\sqrt{c(a+bx^2)^2} \right) \right) \right)$$

input `Integrate[(c*(a + b*x^2)^2)^(3/2)/x^5,x]`

output `(c*(-((c*(-a^3 - 6*a^2*b*x^2 + a*b^2*x^4 + 2*b^3*x^6)*(a*Sqrt[b^2*c] + b*Sqrt[b^2*c]*x^2 - b*Sqrt[c*(a + b*x^2)^2]))/(x^4*(a*b*c + b^2*c*x^2 - Sqrt[b^2*c]*Sqrt[c*(a + b*x^2)^2]))) + 6*a*b^2*Sqrt[c]*ArcTanh[(Sqrt[b^2*c]*x^2 - Sqrt[c*(a + b*x^2)^2])/(a*Sqrt[c])] - 3*a*b*Sqrt[b^2*c]*Log[x^2*(a*b*c + b^2*c*x^2 - Sqrt[b^2*c]*Sqrt[c*(a + b*x^2)^2])])/4`

Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.47, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {2045, 27, 243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c(a+bx^2)^2)^{3/2}}{x^5} dx$$

↓ 2045

$$\begin{aligned}
& \frac{a^3 c \sqrt{c(a+bx^2)^2} \int \frac{(bx^2+a)^3}{a^3 x^5} dx}{a+bx^2} \\
& \quad \downarrow \text{27} \\
& \frac{c \sqrt{c(a+bx^2)^2} \int \frac{(bx^2+a)^3}{x^5} dx}{a+bx^2} \\
& \quad \downarrow \text{243} \\
& \frac{c \sqrt{c(a+bx^2)^2} \int \frac{(bx^2+a)^3}{x^6} dx^2}{2(a+bx^2)} \\
& \quad \downarrow \text{49} \\
& \frac{c \sqrt{c(a+bx^2)^2} \int \left(\frac{a^3}{x^6} + \frac{3ba^2}{x^4} + \frac{3b^2a}{x^2} + b^3 \right) dx^2}{2(a+bx^2)} \\
& \quad \downarrow \text{2009} \\
& \frac{c \sqrt{c(a+bx^2)^2} \left(-\frac{a^3}{2x^4} - \frac{3a^2b}{x^2} + 3ab^2 \log(x^2) + b^3 x^2 \right)}{2(a+bx^2)}
\end{aligned}$$

input `Int[(c*(a + b*x^2)^2)^(3/2)/x^5,x]`

output `(c*Sqrt[c*(a + b*x^2)^2]*(-1/2*a^3/x^4 - (3*a^2*b)/x^2 + b^3*x^2 + 3*a*b^2*Log[x^2]))/(2*(a + b*x^2))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_.))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.44

method	result	size
default	$\frac{(c(bx^2+a)^2)^{\frac{3}{2}}(2b^3x^6+12ab^2\ln(x)x^4-6a^2bx^2-a^3)}{4x^4(bx^2+a)^3}$	62
pseudoelliptic	$-\frac{c\sqrt{c(bx^2+a)^2}(-6ab^2\ln(x)x^4-2b^3x^6+6a^2bx^2+a^3)}{4(bx^2+a)x^4}$	63
risch	$\frac{b^3cx^2\sqrt{c(bx^2+a)^2}}{2bx^2+2a} + \frac{c\sqrt{c(bx^2+a)^2}(-\frac{3}{2}a^2bx^2-\frac{1}{4}a^3)}{(bx^2+a)x^4} + \frac{3ab^2c\sqrt{c(bx^2+a)^2}\ln(x)}{bx^2+a}$	106

input `int((c*(b*x^2+a)^2)^(3/2)/x^5,x,method=_RETURNVERBOSE)`

output `1/4*(c*(b*x^2+a)^2)^(3/2)*(2*b^3*x^6+12*a*b^2*ln(x)*x^4-6*a^2*b*x^2-a^3)/x^4/(b*x^2+a)^3`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.55

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^5} dx = \frac{(2b^3cx^6 + 12ab^2cx^4 \log(x) - 6a^2bcx^2 - a^3c)\sqrt{b^2cx^4 + 2abcx^2 + a^2c}}{4(bx^6 + ax^4)}$$

input `integrate((c*(b*x^2+a)^2)^(3/2)/x^5,x, algorithm="fricas")`output `1/4*(2*b^3*c*x^6 + 12*a*b^2*c*x^4*log(x) - 6*a^2*b*c*x^2 - a^3*c)*sqrt(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)/(b*x^6 + a*x^4)`**Sympy [F]**

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^5} dx = \int \frac{(c(a + bx^2)^2)^{\frac{3}{2}}}{x^5} dx$$

input `integrate((c*(b*x**2+a)**2)**(3/2)/x**5,x)`output `Integral((c*(a + b*x**2)**2)**(3/2)/x**5, x)`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.79

$$\begin{aligned} \int \frac{(c(a + bx^2)^2)^{3/2}}{x^5} dx &= \frac{3}{2} (-1)^{2b^2cx^2+2abc} ab^2c^{\frac{3}{2}} \log(2b^2cx^2 + 2abc) \\ &- \frac{3}{2} (-1)^{2abcx^2+2a^2c} ab^2c^{\frac{3}{2}} \log\left(2abc + \frac{2a^2c}{x^2}\right) + \frac{3\sqrt{b^2cx^4 + 2abcx^2 + a^2c}b^3cx^2}{4a} \\ &+ \frac{9}{4} \sqrt{b^2cx^4 + 2abcx^2 + a^2c}b^2c + \frac{(b^2cx^4 + 2abcx^2 + a^2c)^{\frac{3}{2}}b^2}{4a^2} \\ &- \frac{(b^2cx^4 + 2abcx^2 + a^2c)^{\frac{3}{2}}b}{4ax^2} - \frac{(b^2cx^4 + 2abcx^2 + a^2c)^{\frac{5}{2}}}{4a^2cx^4} \end{aligned}$$

input `integrate((c*(b*x^2+a)^2)^(3/2)/x^5,x, algorithm="maxima")`

output
$$\begin{aligned} & 3/2*(-1)^(2*b^2*c*x^2 + 2*a*b*c)*a*b^2*c^(3/2)*\log(2*b^2*c*x^2 + 2*a*b*c) \\ & - 3/2*(-1)^(2*a*b*c*x^2 + 2*a^2*c)*a*b^2*c^(3/2)*\log(2*a*b*c + 2*a^2*c/x^2) \\ & + 3/4*\sqrt{b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c}*b^3*c*x^2/a + 9/4*\sqrt{b^2*c} \\ & *x^4 + 2*a*b*c*x^2 + a^2*c)*b^2*c + 1/4*(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)^(3/2)*b^2/a^2 \\ & - 1/4*(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)^(3/2)*b/(a*x^2) - 1/4*(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)^(5/2)/(a^2*c*x^4) \end{aligned}$$

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.66

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^5} dx = \frac{1}{4} \left(2b^3x^2 \operatorname{sgn}(bx^2 + a) + 6ab^2 \log(x^2) \operatorname{sgn}(bx^2 + a) - \frac{9ab^2x^4 \operatorname{sgn}(bx^2 + a) + 6a^3 \operatorname{sgn}(bx^2 + a)}{x^4} \right) * c^{3/2}$$

input `integrate((c*(b*x^2+a)^2)^(3/2)/x^5,x, algorithm="giac")`

output
$$\frac{1}{4}*(2*b^3*x^2*\operatorname{sgn}(b*x^2 + a) + 6*a*b^2*\log(x^2)*\operatorname{sgn}(b*x^2 + a) - (9*a*b^2*x^4*\operatorname{sgn}(b*x^2 + a) + 6*a^2*b*x^2*\operatorname{sgn}(b*x^2 + a) + a^3*\operatorname{sgn}(b*x^2 + a))/x^4)*c^(3/2)$$

Mupad [F(-1)]

Timed out.

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^5} dx = \int \frac{(c(bx^2 + a)^2)^{3/2}}{x^5} dx$$

input `int((c*(a + b*x^2)^2)^(3/2)/x^5,x)`

output `int((c*(a + b*x^2)^2)^(3/2)/x^5, x)`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.30

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^5} dx = \frac{\sqrt{c}c(12 \log(x) a b^2 x^4 - a^3 - 6a^2 b x^2 + 2b^3 x^6)}{4x^4}$$

input `int((c*(b*x^2+a)^2)^(3/2)/x^5,x)`

output `(sqrt(c)*c*(12*log(x)*a*b**2*x**4 - a**3 - 6*a**2*b*x**2 + 2*b**3*x**6))/(4*x**4)`

3.7 $\int x^4 \left(c(a + bx^2)^2 \right)^{3/2} dx$

Optimal result	92
Mathematica [A] (verified)	92
Rubi [A] (verified)	93
Maple [A] (verified)	94
Fricas [A] (verification not implemented)	95
Sympy [F(-1)]	95
Maxima [A] (verification not implemented)	96
Giac [A] (verification not implemented)	96
Mupad [F(-1)]	96
Reduce [B] (verification not implemented)	97

Optimal result

Integrand size = 19, antiderivative size = 143

$$\int x^4 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{a^3 cx^5 \sqrt{c(a + bx^2)^2}}{5(a + bx^2)} + \frac{3a^2 bcx^7 \sqrt{c(a + bx^2)^2}}{7(a + bx^2)} + \frac{ab^2 cx^9 \sqrt{c(a + bx^2)^2}}{3(a + bx^2)} + \frac{b^3 cx^{11} \sqrt{c(a + bx^2)^2}}{11(a + bx^2)}$$

output

```
a^3*c*x^5*(c*(b*x^2+a)^2)^(1/2)/(5*b*x^2+5*a)+3*a^2*b*c*x^7*(c*(b*x^2+a)^2)^(1/2)/(7*b*x^2+7*a)+a*b^2*c*x^9*(c*(b*x^2+a)^2)^(1/2)/(3*b*x^2+3*a)+b^3*c*x^11*(c*(b*x^2+a)^2)^(1/2)/(11*b*x^2+11*a)
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.44

$$\int x^4 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{x^5 \left(c(a + bx^2)^2 \right)^{3/2} (231a^3 + 495a^2bx^2 + 385ab^2x^4 + 105b^3x^6)}{1155(a + bx^2)^3}$$

input

```
Integrate[x^4*(c*(a + b*x^2)^2)^(3/2),x]
```

output $(x^5*(c*(a + b*x^2)^2)^{(3/2)*(231*a^3 + 495*a^2*b*x^2 + 385*a*b^2*x^4 + 105*b^3*x^6)))/(1155*(a + b*x^2)^3)$

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.48, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {2045, 27, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^4 (c(a + bx^2)^2)^{3/2} dx \\ & \quad \downarrow 2045 \\ & \frac{a^3 c \sqrt{c(a + bx^2)^2} \int \frac{x^4 (bx^2 + a)^3}{a^3} dx}{a + bx^2} \\ & \quad \downarrow 27 \\ & \frac{c \sqrt{c(a + bx^2)^2} \int x^4 (bx^2 + a)^3 dx}{a + bx^2} \\ & \quad \downarrow 244 \\ & \frac{c \sqrt{c(a + bx^2)^2} \int (b^3 x^{10} + 3ab^2 x^8 + 3a^2 b x^6 + a^3 x^4) dx}{a + bx^2} \\ & \quad \downarrow 2009 \\ & \frac{c \left(\frac{a^3 x^5}{5} + \frac{3}{7} a^2 b x^7 + \frac{1}{3} a b^2 x^9 + \frac{b^3 x^{11}}{11} \right) \sqrt{c(a + bx^2)^2}}{a + bx^2} \end{aligned}$$

input $\text{Int}[x^4*(c*(a + b*x^2)^2)^{(3/2)}, x]$

output $(c*\text{Sqrt}[c*(a + b*x^2)^2]*((a^3*x^5)/5 + (3*a^2*b*x^7)/7 + (a*b^2*x^9)/3 + (b^3*x^11)/11))/(a + b*x^2)$

Definitions of rubi rules used

rule 27 `Int[(a_)*(F x_), x_Symbol] := Simp[a Int[F x, x], x] /; FreeQ[a, x] && !MatchQ[F x, (b_)*(G x_) /; FreeQ[b, x]]`

rule 244 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2045 `Int[(u_)*((c_)*((a_) + (b_)*(x_)^(n_))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 1.17 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.42

method	result	size
gospers	$\frac{x^5(105b^3x^6+385ab^2x^4+495a^2bx^2+231a^3)(c(bx^2+a)^2)^{\frac{3}{2}}}{1155(bx^2+a)^3}$	60
default	$\frac{x^5(105b^3x^6+385ab^2x^4+495a^2bx^2+231a^3)(c(bx^2+a)^2)^{\frac{3}{2}}}{1155(bx^2+a)^3}$	60
orering	$\frac{x^5(105b^3x^6+385ab^2x^4+495a^2bx^2+231a^3)(c(bx^2+a)^2)^{\frac{3}{2}}}{1155(bx^2+a)^3}$	60
trager	$\frac{cx^5(105b^3x^6+385ab^2x^4+495a^2bx^2+231a^3)\sqrt{b^2cx^4+2abcx^2+a^2c}}{1155bx^2+1155a}$	72
risch	$\frac{a^3cx^5\sqrt{c(bx^2+a)^2}}{5bx^2+5a} + \frac{3c\sqrt{c(bx^2+a)^2}a^2bx^7}{7(bx^2+a)} + \frac{ab^2cx^9\sqrt{c(bx^2+a)^2}}{3bx^2+3a} + \frac{b^3cx^{11}\sqrt{c(bx^2+a)^2}}{11bx^2+11a}$	128

input `int(x^4*(c*(b*x^2+a)^2)^(3/2),x,method=_RETURNVERBOSE)`

output $1/1155*x^5*(105*b^3*x^6+385*a*b^2*x^4+495*a^2*b*x^2+231*a^3)*(c*(b*x^2+a)^2)^{(3/2)}/(b*x^2+a)^3$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.52

$$\int x^4 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{(105 b^3 cx^{11} + 385 ab^2 cx^9 + 495 a^2 bcx^7 + 231 a^3 cx^5) \sqrt{b^2 cx^4 + 2 abcx^2 + a^2 c}}{1155 (bx^2 + a)}$$

input `integrate(x^4*(c*(b*x^2+a)^2)^(3/2),x, algorithm="fricas")`

output $1/1155*(105*b^3*c*x^{11} + 385*a*b^2*c*x^9 + 495*a^2*b*c*x^7 + 231*a^3*c*x^5)*\text{sqrt}(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)/(b*x^2 + a)$

Sympy [F(-1)]

Timed out.

$$\int x^4 \left(c(a + bx^2)^2 \right)^{3/2} dx = \text{Timed out}$$

input `integrate(x**4*(c*(b*x**2+a)**2)**(3/2),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.33

$$\int x^4 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{1}{11} b^3 c^{3/2} x^{11} + \frac{1}{3} ab^2 c^{3/2} x^9 + \frac{3}{7} a^2 b c^{3/2} x^7 + \frac{1}{5} a^3 c^{3/2} x^5$$

input `integrate(x^4*(c*(b*x^2+a)^2)^(3/2),x, algorithm="maxima")`output `1/11*b^3*c^(3/2)*x^11 + 1/3*a*b^2*c^(3/2)*x^9 + 3/7*a^2*b*c^(3/2)*x^7 + 1/5*a^3*c^(3/2)*x^5`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.50

$$\int x^4 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{1}{1155} (105 b^3 x^{11} \operatorname{sgn}(bx^2 + a) + 385 ab^2 x^9 \operatorname{sgn}(bx^2 + a) + 495 a^2 b x^7 \operatorname{sgn}(bx^2 + a) + 231 a^3 \operatorname{sgn}(bx^2 + a)) c^{3/2}$$

input `integrate(x^4*(c*(b*x^2+a)^2)^(3/2),x, algorithm="giac")`output `1/1155*(105*b^3*x^11*sgn(b*x^2 + a) + 385*a*b^2*x^9*sgn(b*x^2 + a) + 495*a^2*b*x^7*sgn(b*x^2 + a) + 231*a^3*x^5*sgn(b*x^2 + a))*c^(3/2)`**Mupad [F(-1)]**

Timed out.

$$\int x^4 \left(c(a + bx^2)^2 \right)^{3/2} dx = \int x^4 \left(c(bx^2 + a)^2 \right)^{3/2} dx$$

input `int(x^4*(c*(a + b*x^2)^2)^(3/2),x)`output `int(x^4*(c*(a + b*x^2)^2)^(3/2), x)`

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.28

$$\int x^4 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{\sqrt{c} c x^5 (105b^3 x^6 + 385a b^2 x^4 + 495a^2 b x^2 + 231a^3)}{1155}$$

input `int(x^4*(c*(b*x^2+a)^2)^(3/2),x)`

output `(sqrt(c)*c*x**5*(231*a**3 + 495*a**2*b*x**2 + 385*a*b**2*x**4 + 105*b**3*x**6))/1155`

3.8 $\int x^2 \left(c(a + bx^2)^2 \right)^{3/2} dx$

Optimal result	98
Mathematica [A] (verified)	98
Rubi [A] (verified)	99
Maple [A] (verified)	100
Fricas [A] (verification not implemented)	101
Sympy [F]	101
Maxima [A] (verification not implemented)	102
Giac [A] (verification not implemented)	102
Mupad [F(-1)]	102
Reduce [B] (verification not implemented)	103

Optimal result

Integrand size = 19, antiderivative size = 143

$$\int x^2 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{a^3 cx^3 \sqrt{c(a + bx^2)^2}}{3(a + bx^2)} + \frac{3a^2 bcx^5 \sqrt{c(a + bx^2)^2}}{5(a + bx^2)} + \frac{3ab^2 cx^7 \sqrt{c(a + bx^2)^2}}{7(a + bx^2)} + \frac{b^3 cx^9 \sqrt{c(a + bx^2)^2}}{9(a + bx^2)}$$

output

$$a^3 c x^3 (c (b x^2 + a)^2)^{1/2} / (3 b x^2 + 3 a) + 3 a^2 b c x^5 (c (b x^2 + a)^2)^{1/2} / (5 b x^2 + 5 a) + 3 a b^2 c x^7 (c (b x^2 + a)^2)^{1/2} / (7 b x^2 + 7 a) + b^3 c x^9 (c (b x^2 + a)^2)^{1/2} / (9 b x^2 + 9 a)$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.44

$$\int x^2 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{x^3 \left(c(a + bx^2)^2 \right)^{3/2} (105a^3 + 189a^2bx^2 + 135ab^2x^4 + 35b^3x^6)}{315(a + bx^2)^3}$$

input

$$\text{Integrate}[x^2 (c (a + b x^2)^2)^{3/2}, x]$$

output

$$(x^3*(c*(a + b*x^2)^2)^(3/2)*(105*a^3 + 189*a^2*b*x^2 + 135*a*b^2*x^4 + 35*b^3*x^6))/(315*(a + b*x^2)^3)$$

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.48, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {2045, 27, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^2 (c(a + bx^2)^2)^{3/2} dx \\ & \quad \downarrow \text{2045} \\ & \frac{a^3 c \sqrt{c(a + bx^2)^2} \int \frac{x^2 (bx^2 + a)^3}{a^3} dx}{a + bx^2} \\ & \quad \downarrow \text{27} \\ & \frac{c \sqrt{c(a + bx^2)^2} \int x^2 (bx^2 + a)^3 dx}{a + bx^2} \\ & \quad \downarrow \text{244} \\ & \frac{c \sqrt{c(a + bx^2)^2} \int (b^3 x^8 + 3ab^2 x^6 + 3a^2 b x^4 + a^3 x^2) dx}{a + bx^2} \\ & \quad \downarrow \text{2009} \\ & \frac{c \left(\frac{a^3 x^3}{3} + \frac{3}{5} a^2 b x^5 + \frac{3}{7} a b^2 x^7 + \frac{b^3 x^9}{9} \right) \sqrt{c(a + bx^2)^2}}{a + bx^2} \end{aligned}$$

input

$$\text{Int}[x^2*(c*(a + b*x^2)^2)^(3/2), x]$$

output

$$(c*\text{Sqrt}[c*(a + b*x^2)^2]*((a^3*x^3)/3 + (3*a^2*b*x^5)/5 + (3*a*b^2*x^7)/7 + (b^3*x^9)/9))/(a + b*x^2)$$

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 244 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2045 `Int[(u_)*((c_)*((a_) + (b_)*(x_)^(n_))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 0.90 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.42

method	result	size
gospers	$\frac{x^3(35b^3x^6+135ab^2x^4+189a^2bx^2+105a^3)(cx^2+a)^2}{315(bx^2+a)^3}$	60
default	$\frac{x^3(35b^3x^6+135ab^2x^4+189a^2bx^2+105a^3)(cx^2+a)^2}{315(bx^2+a)^3}$	60
orering	$\frac{x^3(35b^3x^6+135ab^2x^4+189a^2bx^2+105a^3)(cx^2+a)^2}{315(bx^2+a)^3}$	60
trager	$\frac{cx^3(35b^3x^6+135ab^2x^4+189a^2bx^2+105a^3)\sqrt{b^2cx^4+2abcx^2+a^2c}}{315bx^2+315a}$	72
risch	$\frac{a^3cx^3\sqrt{c(bx^2+a)^2}}{3bx^2+3a} + \frac{3c\sqrt{c(bx^2+a)^2}a^2bx^5}{5(bx^2+a)} + \frac{3c\sqrt{c(bx^2+a)^2}ab^2x^7}{7(bx^2+a)} + \frac{b^3cx^9\sqrt{c(bx^2+a)^2}}{9bx^2+9a}$	128

input `int(x^2*(c*(b*x^2+a)^2)^(3/2),x,method=_RETURNVERBOSE)`

output

```
1/315*x^3*(35*b^3*x^6+135*a*b^2*x^4+189*a^2*b*x^2+105*a^3)*(c*(b*x^2+a)^2)^(3/2)/(b*x^2+a)^3
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.52

$$\int x^2 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{(35b^3cx^9 + 135ab^2cx^7 + 189a^2bcx^5 + 105a^3cx^3)\sqrt{b^2cx^4 + 2abcx^2 + a^2c}}{315(bx^2 + a)}$$

input

```
integrate(x^2*(c*(b*x^2+a)^2)^(3/2),x, algorithm="fricas")
```

output

```
1/315*(35*b^3*c*x^9 + 135*a*b^2*c*x^7 + 189*a^2*b*c*x^5 + 105*a^3*c*x^3)*s
qrt(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)/(b*x^2 + a)
```

Sympy [F]

$$\int x^2 \left(c(a + bx^2)^2 \right)^{3/2} dx = \int x^2 \left(c(a + bx^2)^2 \right)^{\frac{3}{2}} dx$$

input

```
integrate(x**2*(c*(b*x**2+a)**2)**(3/2),x)
```

output

```
Integral(x**2*(c*(a + b*x**2)**2)**(3/2), x)
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.33

$$\int x^2 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{1}{9} b^3 c^{3/2} x^9 + \frac{3}{7} ab^2 c^{3/2} x^7 + \frac{3}{5} a^2 b c^{3/2} x^5 + \frac{1}{3} a^3 c^{3/2} x^3$$

input `integrate(x^2*(c*(b*x^2+a)^2)^(3/2),x, algorithm="maxima")`output `1/9*b^3*c^(3/2)*x^9 + 3/7*a*b^2*c^(3/2)*x^7 + 3/5*a^2*b*c^(3/2)*x^5 + 1/3*a^3*c^(3/2)*x^3`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.50

$$\int x^2 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{1}{315} (35 b^3 x^9 \operatorname{sgn}(bx^2 + a) + 135 ab^2 x^7 \operatorname{sgn}(bx^2 + a) + 189 a^2 b x^5 \operatorname{sgn}(bx^2 + a) + 105 a^3 x^3 \operatorname{sgn}(bx^2 + a)) c^{3/2}$$

input `integrate(x^2*(c*(b*x^2+a)^2)^(3/2),x, algorithm="giac")`output `1/315*(35*b^3*x^9*sgn(b*x^2 + a) + 135*a*b^2*x^7*sgn(b*x^2 + a) + 189*a^2*b*x^5*sgn(b*x^2 + a) + 105*a^3*x^3*sgn(b*x^2 + a))*c^(3/2)`**Mupad [F(-1)]**

Timed out.

$$\int x^2 \left(c(a + bx^2)^2 \right)^{3/2} dx = \int x^2 \left(c(bx^2 + a)^2 \right)^{3/2} dx$$

input `int(x^2*(c*(a + b*x^2)^2)^(3/2),x)`output `int(x^2*(c*(a + b*x^2)^2)^(3/2), x)`

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.28

$$\int x^2 \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{\sqrt{c} c x^3 (35b^3 x^6 + 135a b^2 x^4 + 189a^2 b x^2 + 105a^3)}{315}$$

input `int(x^2*(c*(b*x^2+a)^2)^(3/2),x)`

output `(sqrt(c)*c*x**3*(105*a**3 + 189*a**2*b*x**2 + 135*a*b**2*x**4 + 35*b**3*x**6))/315`

3.9 $\int \left(c(a + bx^2)^2\right)^{3/2} dx$

Optimal result	104
Mathematica [A] (verified)	104
Rubi [A] (verified)	105
Maple [A] (verified)	106
Fricas [A] (verification not implemented)	107
Sympy [F]	107
Maxima [A] (verification not implemented)	107
Giac [A] (verification not implemented)	108
Mupad [F(-1)]	108
Reduce [B] (verification not implemented)	108

Optimal result

Integrand size = 15, antiderivative size = 135

$$\int \left(c(a + bx^2)^2\right)^{3/2} dx = \frac{a^3 cx \sqrt{c(a + bx^2)^2}}{a + bx^2} + \frac{a^2 b c x^3 \sqrt{c(a + bx^2)^2}}{a + bx^2} + \frac{3 a b^2 c x^5 \sqrt{c(a + bx^2)^2}}{5(a + bx^2)} + \frac{b^3 c x^7 \sqrt{c(a + bx^2)^2}}{7(a + bx^2)}$$

output

```
a^3*c*x*(c*(b*x^2+a)^2)^(1/2)/(b*x^2+a)+a^2*b*c*x^3*(c*(b*x^2+a)^2)^(1/2)/(b*x^2+a)+3*a*b^2*c*x^5*(c*(b*x^2+a)^2)^(1/2)/(5*b*x^2+5*a)+b^3*c*x^7*(c*(b*x^2+a)^2)^(1/2)/(7*b*x^2+7*a)
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.45

$$\int \left(c(a + bx^2)^2\right)^{3/2} dx = \frac{x \left(c(a + bx^2)^2\right)^{3/2} (35a^3 + 35a^2bx^2 + 21ab^2x^4 + 5b^3x^6)}{35(a + bx^2)^3}$$

input

```
Integrate[(c*(a + b*x^2)^2)^(3/2),x]
```

output

$$\frac{(x*(c*(a + b*x^2)^2)^{(3/2)*(35*a^3 + 35*a^2*b*x^2 + 21*a*b^2*x^4 + 5*b^3*x^6))}{(35*(a + b*x^2)^3)}$$

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.48, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2045, 210, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (c(a + bx^2)^2)^{3/2} dx \\ & \quad \downarrow 2045 \\ & \frac{a^3 c \sqrt{c(a + bx^2)^2} \int \left(\frac{bx^2}{a} + 1\right)^3 dx}{a + bx^2} \\ & \quad \downarrow 210 \\ & \frac{a^3 c \sqrt{c(a + bx^2)^2} \int \left(\frac{b^3 x^6}{a^3} + \frac{3b^2 x^4}{a^2} + \frac{3bx^2}{a} + 1\right) dx}{a + bx^2} \\ & \quad \downarrow 2009 \\ & \frac{a^3 c \left(\frac{b^3 x^7}{7a^3} + \frac{3b^2 x^5}{5a^2} + \frac{bx^3}{a} + x\right) \sqrt{c(a + bx^2)^2}}{a + bx^2} \end{aligned}$$

input

$$\text{Int}[(c*(a + b*x^2)^2)^{(3/2),x}]$$

output

$$\frac{(a^3*c*\text{Sqrt}[c*(a + b*x^2)^2]*(x + (b*x^3)/a + (3*b^2*x^5)/(5*a^2) + (b^3*x^7)/(7*a^3)))/(a + b*x^2)}$$

Defintions of rubi rules used

rule 210 $\text{Int}[(a_+) + (b_+)(x_+)^2]^{(p_+)} , x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x^2)^p, x], x] /;$ $\text{FreeQ}\{a, b\}, x] \ \&\& \ \text{IGtQ}[p, 0]$

rule 2009 $\text{Int}[u_ , x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /;$ $\text{SumQ}[u]$

rule 2045 $\text{Int}[(u_+)((c_+)((a_+ + (b_+)(x_+)^{n_+}))^{(q_+)})^{(p_+)} , x_Symbol] \rightarrow \text{Simp}[\text{Simp}[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^{(p*q)}] \text{Int}[u*(1 + b*(x^n/a))^{(p*q)}, x], x] /;$ $\text{FreeQ}\{a, b, c, n, p, q\}, x] \ \&\& \ !\text{GeQ}[a, 0]$

Maple [A] (verified)

Time = 0.56 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.43

method	result	size
gospers	$\frac{x(5b^3x^6+21ab^2x^4+35a^2bx^2+35a^3)(c(bx^2+a)^2)^{\frac{3}{2}}}{35(bx^2+a)^3}$	58
default	$\frac{x(5b^3x^6+21ab^2x^4+35a^2bx^2+35a^3)(c(bx^2+a)^2)^{\frac{3}{2}}}{35(bx^2+a)^3}$	58
orering	$\frac{x(5b^3x^6+21ab^2x^4+35a^2bx^2+35a^3)(c(bx^2+a)^2)^{\frac{3}{2}}}{35(bx^2+a)^3}$	58
trager	$\frac{cx(5b^3x^6+21ab^2x^4+35a^2bx^2+35a^3)\sqrt{b^2cx^4+2abcx^2+a^2c}}{35bx^2+35a}$	70
risch	$\frac{a^3cx\sqrt{c(bx^2+a)^2}}{bx^2+a} + \frac{a^2bcx^3\sqrt{c(bx^2+a)^2}}{bx^2+a} + \frac{3c\sqrt{c(bx^2+a)^2}ab^2x^5}{5(bx^2+a)} + \frac{b^3cx^7\sqrt{c(bx^2+a)^2}}{7bx^2+7a}$	124

input $\text{int}((c*(b*x^2+a)^2)^{(3/2)}, x, \text{method}=_RETURNVERBOSE)$

output $1/35*x*(5*b^3*x^6+21*a*b^2*x^4+35*a^2*b*x^2+35*a^3)*(c*(b*x^2+a)^2)^{(3/2)}/(b*x^2+a)^3$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.53

$$\int \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{(5b^3cx^7 + 21ab^2cx^5 + 35a^2bcx^3 + 35a^3cx)\sqrt{b^2cx^4 + 2abcx^2 + a^2c}}{35(bx^2 + a)}$$

input `integrate((c*(b*x^2+a)^2)^(3/2),x, algorithm="fricas")`output `1/35*(5*b^3*c*x^7 + 21*a*b^2*c*x^5 + 35*a^2*b*c*x^3 + 35*a^3*c*x)*sqrt(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)/(b*x^2 + a)`**Sympy [F]**

$$\int \left(c(a + bx^2)^2 \right)^{3/2} dx = \int \left(c(a + bx^2)^2 \right)^{\frac{3}{2}} dx$$

input `integrate((c*(b*x**2+a)**2)**(3/2),x)`output `Integral((c*(a + b*x**2)**2)**(3/2), x)`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.32

$$\int \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{1}{7} b^3 c^{\frac{3}{2}} x^7 + \frac{3}{5} ab^2 c^{\frac{3}{2}} x^5 + a^2 bc^{\frac{3}{2}} x^3 + a^3 c^{\frac{3}{2}} x$$

input `integrate((c*(b*x^2+a)^2)^(3/2),x, algorithm="maxima")`output `1/7*b^3*c^(3/2)*x^7 + 3/5*a*b^2*c^(3/2)*x^5 + a^2*b*c^(3/2)*x^3 + a^3*c^(3/2)*x`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.34

$$\int \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{1}{35} (5b^3x^7 + 21ab^2x^5 + 35a^2bx^3 + 35a^3x) c^{3/2} \operatorname{sgn}(bx^2 + a)$$

input `integrate((c*(b*x^2+a)^2)^(3/2),x, algorithm="giac")`

output `1/35*(5*b^3*x^7 + 21*a*b^2*x^5 + 35*a^2*b*x^3 + 35*a^3*x)*c^(3/2)*sgn(b*x^2 + a)`

Mupad [F(-1)]

Timed out.

$$\int \left(c(a + bx^2)^2 \right)^{3/2} dx = \int \left(c(bx^2 + a)^2 \right)^{3/2} dx$$

input `int((c*(a + b*x^2)^2)^(3/2),x)`

output `int((c*(a + b*x^2)^2)^(3/2), x)`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.28

$$\int \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{\sqrt{c}cx(5b^3x^6 + 21ab^2x^4 + 35a^2bx^2 + 35a^3)}{35}$$

input `int((c*(b*x^2+a)^2)^(3/2),x)`

output `(sqrt(c)*c*x*(35*a**3 + 35*a**2*b*x**2 + 21*a*b**2*x**4 + 5*b**3*x**6))/35`

3.10
$$\int \frac{\left(c(a+bx^2)^2\right)^{3/2}}{x^2} dx$$

Optimal result	109
Mathematica [A] (verified)	109
Rubi [A] (verified)	110
Maple [A] (verified)	111
Fricas [A] (verification not implemented)	112
Sympy [F]	112
Maxima [A] (verification not implemented)	113
Giac [A] (verification not implemented)	113
Mupad [F(-1)]	113
Reduce [B] (verification not implemented)	114

Optimal result

Integrand size = 19, antiderivative size = 134

$$\int \frac{\left(c(a+bx^2)^2\right)^{3/2}}{x^2} dx = -\frac{a^3c\sqrt{c(a+bx^2)^2}}{x(a+bx^2)} + \frac{3a^2bcx\sqrt{c(a+bx^2)^2}}{a+bx^2} + \frac{ab^2cx^3\sqrt{c(a+bx^2)^2}}{a+bx^2} + \frac{b^3cx^5\sqrt{c(a+bx^2)^2}}{5(a+bx^2)}$$

output

```
-a^3*c*(c*(b*x^2+a)^2)^(1/2)/x/(b*x^2+a)+3*a^2*b*c*x*(c*(b*x^2+a)^2)^(1/2)
/(b*x^2+a)+a*b^2*c*x^3*(c*(b*x^2+a)^2)^(1/2)/(b*x^2+a)+b^3*c*x^5*(c*(b*x^2
+a)^2)^(1/2)/(5*b*x^2+5*a)
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.46

$$\int \frac{\left(c(a+bx^2)^2\right)^{3/2}}{x^2} dx = \frac{\left(c(a+bx^2)^2\right)^{3/2} (-5a^3 + 15a^2bx^2 + 5ab^2x^4 + b^3x^6)}{5x(a+bx^2)^3}$$

input

```
Integrate[(c*(a + b*x^2)^2)^(3/2)/x^2,x]
```

output $((c*(a + b*x^2)^2)^{(3/2)*(-5*a^3 + 15*a^2*b*x^2 + 5*a*b^2*x^4 + b^3*x^6)})/(5*x*(a + b*x^2)^3)$

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.45, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {2045, 27, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c(a + bx^2)^2)^{3/2}}{x^2} dx \\ & \quad \downarrow \text{2045} \\ & \frac{a^3 c \sqrt{c(a + bx^2)^2} \int \frac{(bx^2 + a)^3}{a^3 x^2} dx}{a + bx^2} \\ & \quad \downarrow \text{27} \\ & \frac{c \sqrt{c(a + bx^2)^2} \int \frac{(bx^2 + a)^3}{x^2} dx}{a + bx^2} \\ & \quad \downarrow \text{244} \\ & \frac{c \sqrt{c(a + bx^2)^2} \int (b^3 x^4 + 3ab^2 x^2 + 3a^2 b + \frac{a^3}{x^2}) dx}{a + bx^2} \\ & \quad \downarrow \text{2009} \\ & \frac{c \left(-\frac{a^3}{x} + 3a^2 b x + ab^2 x^3 + \frac{b^3 x^5}{5} \right) \sqrt{c(a + bx^2)^2}}{a + bx^2} \end{aligned}$$

input $\text{Int}[(c*(a + b*x^2)^2)^{(3/2)}/x^2, x]$

output $(c*\text{Sqrt}[c*(a + b*x^2)^2]*(-a^3/x) + 3*a^2*b*x + a*b^2*x^3 + (b^3*x^5)/5)/(a + b*x^2)$

Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 244 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2045 `Int[(u_)*((c_)*((a_) + (b_)*(x_)^(n_))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 0.75 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.45

method	result	size
gospers	$-\frac{(-b^3x^6 - 5ab^2x^4 - 15a^2bx^2 + 5a^3)(c(bx^2 + a)^2)^{3/2}}{5x(bx^2 + a)^3}$	60
default	$-\frac{(-b^3x^6 - 5ab^2x^4 - 15a^2bx^2 + 5a^3)(c(bx^2 + a)^2)^{3/2}}{5x(bx^2 + a)^3}$	60
orering	$-\frac{(-b^3x^6 - 5ab^2x^4 - 15a^2bx^2 + 5a^3)(c(bx^2 + a)^2)^{3/2}}{5x(bx^2 + a)^3}$	60
risch	$\frac{c\sqrt{c(bx^2 + a)^2} b(\frac{1}{5}b^2x^5 + abx^3 + 3a^2x)}{bx^2 + a} - \frac{a^3c\sqrt{c(bx^2 + a)^2}}{x(bx^2 + a)}$	79
trager	$\frac{c(b^3x^5 + b^3x^4 + 5ab^2x^3 + b^3x^3 + 5a^2bx^2 + b^3x^2 + 15a^2bx + 5ab^2x + b^3x + 5a^3)(-1+x)\sqrt{b^2cx^4 + 2abcx^2 + a^2c}}{5x(bx^2 + a)}$	114

input `int((c*(b*x^2+a)^2)^(3/2)/x^2,x,method=_RETURNVERBOSE)`

output

```
-1/5*(-b^3*x^6-5*a*b^2*x^4-15*a^2*b*x^2+5*a^3)*(c*(b*x^2+a)^2)^(3/2)/x/(b*x^2+a)^3
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.54

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^2} dx = \frac{(b^3cx^6 + 5ab^2cx^4 + 15a^2bcx^2 - 5a^3c)\sqrt{b^2cx^4 + 2abcx^2 + a^2c}}{5(bx^3 + ax)}$$

input

```
integrate((c*(b*x^2+a)^2)^(3/2)/x^2,x, algorithm="fricas")
```

output

```
1/5*(b^3*c*x^6 + 5*a*b^2*c*x^4 + 15*a^2*b*c*x^2 - 5*a^3*c)*sqrt(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)/(b*x^3 + a*x)
```

Sympy [F]

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^2} dx = \int \frac{(c(a + bx^2)^2)^{\frac{3}{2}}}{x^2} dx$$

input

```
integrate((c*(b*x**2+a)**2)**(3/2)/x**2,x)
```

output

```
Integral((c*(a + b*x**2)**2)**(3/2)/x**2, x)
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.36

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^2} dx = \frac{b^3 c^{3/2} x^6 + 5 ab^2 c^{3/2} x^4 + 15 a^2 b c^{3/2} x^2 - 5 a^3 c^{3/2}}{5x}$$

input `integrate((c*(b*x^2+a)^2)^(3/2)/x^2,x, algorithm="maxima")`output `1/5*(b^3*c^(3/2)*x^6 + 5*a*b^2*c^(3/2)*x^4 + 15*a^2*b*c^(3/2)*x^2 - 5*a^3*c^(3/2))/x`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.51

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^2} dx = \frac{1}{5} \left(b^3 x^5 \operatorname{sgn}(bx^2 + a) + 5 ab^2 x^3 \operatorname{sgn}(bx^2 + a) + 15 a^2 b x \operatorname{sgn}(bx^2 + a) - \frac{5 a^3 \operatorname{sgn}(bx^2 + a)}{x} \right)$$

input `integrate((c*(b*x^2+a)^2)^(3/2)/x^2,x, algorithm="giac")`output `1/5*(b^3*x^5*sgn(b*x^2 + a) + 5*a*b^2*x^3*sgn(b*x^2 + a) + 15*a^2*b*x*sgn(b*x^2 + a) - 5*a^3*sgn(b*x^2 + a)/x)*c^(3/2)`**Mupad [F(-1)]**

Timed out.

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^2} dx = \int \frac{(c(bx^2 + a)^2)^{3/2}}{x^2} dx$$

input `int((c*(a + b*x^2)^2)^(3/2)/x^2,x)`output `int((c*(a + b*x^2)^2)^(3/2)/x^2, x)`

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.29

$$\int \frac{\left(c(a + bx^2)^2\right)^{3/2}}{x^2} dx = \frac{\sqrt{c}(b^3x^6 + 5ab^2x^4 + 15a^2bx^2 - 5a^3)}{5x}$$

input `int((c*(b*x^2+a)^2)^(3/2)/x^2,x)`

output `(sqrt(c)*c*(- 5*a**3 + 15*a**2*b*x**2 + 5*a*b**2*x**4 + b**3*x**6))/(5*x)`

$$3.11 \quad \int \frac{(c(a+bx^2)^2)^{3/2}}{x^4} dx$$

Optimal result	115
Mathematica [A] (verified)	115
Rubi [A] (verified)	116
Maple [A] (verified)	117
Fricas [A] (verification not implemented)	118
Sympy [F]	118
Maxima [A] (verification not implemented)	119
Giac [A] (verification not implemented)	119
Mupad [F(-1)]	119
Reduce [B] (verification not implemented)	120

Optimal result

Integrand size = 19, antiderivative size = 137

$$\int \frac{(c(a+bx^2)^2)^{3/2}}{x^4} dx = -\frac{a^3 c \sqrt{c(a+bx^2)^2}}{3x^3(a+bx^2)} - \frac{3a^2 bc \sqrt{c(a+bx^2)^2}}{x(a+bx^2)} + \frac{3ab^2 cx \sqrt{c(a+bx^2)^2}}{a+bx^2} + \frac{b^3 cx^3 \sqrt{c(a+bx^2)^2}}{3(a+bx^2)}$$

output

```
-1/3*a^3*c*(c*(b*x^2+a)^2)^(1/2)/x^3/(b*x^2+a)-3*a^2*b*c*(c*(b*x^2+a)^2)^(1/2)/x/(b*x^2+a)+3*a*b^2*c*x*(c*(b*x^2+a)^2)^(1/2)/(b*x^2+a)+b^3*c*x^3*(c*(b*x^2+a)^2)^(1/2)/(3*b*x^2+3*a)
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.45

$$\int \frac{(c(a+bx^2)^2)^{3/2}}{x^4} dx = -\frac{(c(a+bx^2)^2)^{3/2} (a^3 + 9a^2bx^2 - 9ab^2x^4 - b^3x^6)}{3x^3(a+bx^2)^3}$$

input

```
Integrate[(c*(a + b*x^2)^2)^(3/2)/x^4,x]
```

output
$$\frac{-1/3*((c*(a + b*x^2)^2)^{(3/2)*(a^3 + 9*a^2*b*x^2 - 9*a*b^2*x^4 - b^3*x^6))}{(x^3*(a + b*x^2)^3)}$$

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.46, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {2045, 27, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c(a + bx^2)^2)^{3/2}}{x^4} dx \\ & \quad \downarrow \text{2045} \\ & \frac{a^3 c \sqrt{c(a + bx^2)^2} \int \frac{(bx^2 + a)^3}{a^3 x^4} dx}{a + bx^2} \\ & \quad \downarrow \text{27} \\ & \frac{c \sqrt{c(a + bx^2)^2} \int \frac{(bx^2 + a)^3}{x^4} dx}{a + bx^2} \\ & \quad \downarrow \text{244} \\ & \frac{c \sqrt{c(a + bx^2)^2} \int \left(\frac{a^3}{x^4} + \frac{3ba^2}{x^2} + 3b^2 a + b^3 x^2 \right) dx}{a + bx^2} \\ & \quad \downarrow \text{2009} \\ & \frac{c \left(-\frac{a^3}{3x^3} - \frac{3a^2 b}{x} + 3ab^2 x + \frac{b^3 x^3}{3} \right) \sqrt{c(a + bx^2)^2}}{a + bx^2} \end{aligned}$$

input
$$\text{Int}[(c*(a + b*x^2)^2)^{(3/2)}/x^4, x]$$

output
$$(c*\text{Sqrt}[c*(a + b*x^2)^2]*(-1/3*a^3/x^3 - (3*a^2*b)/x + 3*a*b^2*x + (b^3*x^3)/3))/(a + b*x^2)$$

Definitions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 244 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2045 `Int[(u_)*((c_)*((a_) + (b_)*(x_)^(n_))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 1.02 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.42

method	result	size
gospers	$-\frac{(-b^3x^6-9ab^2x^4+9a^2bx^2+a^3)(c(bx^2+a)^2)^{\frac{3}{2}}}{3x^3(bx^2+a)^3}$	58
default	$-\frac{(-b^3x^6-9ab^2x^4+9a^2bx^2+a^3)(c(bx^2+a)^2)^{\frac{3}{2}}}{3x^3(bx^2+a)^3}$	58
orering	$-\frac{(-b^3x^6-9ab^2x^4+9a^2bx^2+a^3)(c(bx^2+a)^2)^{\frac{3}{2}}}{3x^3(bx^2+a)^3}$	58
risch	$\frac{c\sqrt{c(bx^2+a)^2}b^2(\frac{1}{3}bx^3+3ax)}{bx^2+a} + \frac{c\sqrt{c(bx^2+a)^2}(-3a^2bx^2-\frac{1}{3}a^3)}{(bx^2+a)^3}$	82
trager	$\frac{c(b^3x^5+b^3x^4+9ab^2x^3+b^3x^3+a^3x^2+9a^2bx^2+a^3x+a^3)(-1+x)\sqrt{b^2cx^4+2abcx^2+a^2c}}{3x^3(bx^2+a)}$	98

input `int((c*(b*x^2+a)^2)^(3/2)/x^4,x,method=_RETURNVERBOSE)`

output

```
-1/3*(-b^3*x^6-9*a*b^2*x^4+9*a^2*b*x^2+a^3)*(c*(b*x^2+a)^2)^(3/2)/x^3/(b*x^2+a)^3
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.54

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^4} dx = \frac{(b^3cx^6 + 9ab^2cx^4 - 9a^2bcx^2 - a^3c)\sqrt{b^2cx^4 + 2abcx^2 + a^2c}}{3(bx^5 + ax^3)}$$

input

```
integrate((c*(b*x^2+a)^2)^(3/2)/x^4,x, algorithm="fricas")
```

output

```
1/3*(b^3*c*x^6 + 9*a*b^2*c*x^4 - 9*a^2*b*c*x^2 - a^3*c)*sqrt(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)/(b*x^5 + a*x^3)
```

Sympy [F]

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^4} dx = \int \frac{(c(a + bx^2)^2)^{\frac{3}{2}}}{x^4} dx$$

input

```
integrate((c*(b*x**2+a)**2)**(3/2)/x**4,x)
```

output

```
Integral((c*(a + b*x**2)**2)**(3/2)/x**4, x)
```

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.35

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^4} dx = \frac{b^3 c^{3/2} x^6 + 9 ab^2 c^{3/2} x^4 - 9 a^2 b c^{3/2} x^2 - a^3 c^{3/2}}{3 x^3}$$

input `integrate((c*(b*x^2+a)^2)^(3/2)/x^4,x, algorithm="maxima")`output `1/3*(b^3*c^(3/2)*x^6 + 9*a*b^2*c^(3/2)*x^4 - 9*a^2*b*c^(3/2)*x^2 - a^3*c^(3/2))/x^3`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.52

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^4} dx = \frac{1}{3} \left(b^3 x^3 \operatorname{sgn}(bx^2 + a) + 9 ab^2 x \operatorname{sgn}(bx^2 + a) - \frac{9 a^2 b x^2 \operatorname{sgn}(bx^2 + a) + a^3 \operatorname{sgn}(bx^2 + a)}{x^3} \right) c^{3/2}$$

input `integrate((c*(b*x^2+a)^2)^(3/2)/x^4,x, algorithm="giac")`output `1/3*(b^3*x^3*sgn(b*x^2 + a) + 9*a*b^2*x*sgn(b*x^2 + a) - (9*a^2*b*x^2*sgn(b*x^2 + a) + a^3*sgn(b*x^2 + a))/x^3)*c^(3/2)`**Mupad [F(-1)]**

Timed out.

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^4} dx = \int \frac{(c(bx^2 + a)^2)^{3/2}}{x^4} dx$$

input `int((c*(a + b*x^2)^2)^(3/2)/x^4,x)`output `int((c*(a + b*x^2)^2)^(3/2)/x^4, x)`

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.28

$$\int \frac{(c(a + bx^2)^2)^{3/2}}{x^4} dx = \frac{\sqrt{c}(b^3x^6 + 9ab^2x^4 - 9a^2bx^2 - a^3)}{3x^3}$$

input `int((c*(b*x^2+a)^2)^(3/2)/x^4,x)`

output `(sqrt(c)*c*(- a**3 - 9*a**2*b*x**2 + 9*a*b**2*x**4 + b**3*x**6))/(3*x**3)`

3.12 $\int x^3 \left(c(a + bx^2)^3 \right)^{3/2} dx$

Optimal result	121
Mathematica [A] (verified)	121
Rubi [A] (verified)	122
Maple [A] (verified)	123
Fricas [A] (verification not implemented)	124
Sympy [F(-1)]	124
Maxima [A] (verification not implemented)	125
Giac [A] (verification not implemented)	125
Mupad [B] (verification not implemented)	126
Reduce [B] (verification not implemented)	126

Optimal result

Integrand size = 19, antiderivative size = 66

$$\int x^3 \left(c(a + bx^2)^3 \right)^{3/2} dx = -\frac{ac(a + bx^2)^4 \sqrt{c(a + bx^2)^3}}{11b^2} + \frac{c(a + bx^2)^5 \sqrt{c(a + bx^2)^3}}{13b^2}$$

output

```
-1/11*a*c*(b*x^2+a)^4*(c*(b*x^2+a)^3)^(1/2)/b^2+1/13*c*(b*x^2+a)^5*(c*(b*x^2+a)^3)^(1/2)/b^2
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.59

$$\int x^3 \left(c(a + bx^2)^3 \right)^{3/2} dx = \frac{(a + bx^2) \left(c(a + bx^2)^3 \right)^{3/2} (-2a + 11bx^2)}{143b^2}$$

input

```
Integrate[x^3*(c*(a + b*x^2)^3)^(3/2),x]
```

output

```
((a + b*x^2)*(c*(a + b*x^2)^3)^(3/2)*(-2*a + 11*b*x^2))/(143*b^2)
```

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.30, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {2045, 243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 (c(a + bx^2)^3)^{3/2} dx \\
 & \quad \downarrow \text{2045} \\
 & \frac{a^3 c \sqrt{c(a + bx^2)^3} \int x^3 \left(\frac{bx^2}{a} + 1\right)^{9/2} dx}{\left(\frac{bx^2}{a} + 1\right)^{3/2}} \\
 & \quad \downarrow \text{243} \\
 & \frac{a^3 c \sqrt{c(a + bx^2)^3} \int x^2 \left(\frac{bx^2}{a} + 1\right)^{9/2} dx^2}{2 \left(\frac{bx^2}{a} + 1\right)^{3/2}} \\
 & \quad \downarrow \text{53} \\
 & \frac{a^3 c \sqrt{c(a + bx^2)^3} \int \left(\frac{a\left(\frac{bx^2}{a} + 1\right)^{11/2}}{b} - \frac{a\left(\frac{bx^2}{a} + 1\right)^{9/2}}{b}\right) dx^2}{2 \left(\frac{bx^2}{a} + 1\right)^{3/2}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a^3 c \left(\frac{2a^2 \left(\frac{bx^2}{a} + 1\right)^{13/2}}{13b^2} - \frac{2a^2 \left(\frac{bx^2}{a} + 1\right)^{11/2}}{11b^2}\right) \sqrt{c(a + bx^2)^3}}{2 \left(\frac{bx^2}{a} + 1\right)^{3/2}}
 \end{aligned}$$

input `Int[x^3*(c*(a + b*x^2)^3)^(3/2),x]`

output `(a^3*c*Sqrt[c*(a + b*x^2)^3]*((-2*a^2*(1 + (b*x^2)/a)^(11/2))/(11*b^2) + (2*a^2*(1 + (b*x^2)/a)^(13/2))/(13*b^2)))/(2*(1 + (b*x^2)/a)^(3/2))`

Definitions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 2.42 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.55

method	result	size
gospers	$-\frac{(bx^2+a)(-11bx^2+2a)(c(bx^2+a)^3)^{\frac{3}{2}}}{143b^2}$	36
orering	$-\frac{(bx^2+a)(-11bx^2+2a)(c(bx^2+a)^3)^{\frac{3}{2}}}{143b^2}$	36
default	$-\frac{(-11bx^2+2a)(x^2bc+ac)^{\frac{5}{2}}(c(bx^2+a)^3)^{\frac{3}{2}}}{143b^2c(bx^2+a)c^{\frac{3}{2}}}$	55
risch	$-\frac{c\sqrt{c(bx^2+a)^3}(-11b^6x^{12}-53ab^5x^{10}-100a^2b^4x^8-90a^3b^3x^6-35a^4b^2x^4-a^5bx^2+2a^6)}{143(bx^2+a)b^2}$	94
trager	$-\frac{c(-11b^5x^{10}-42ab^4x^8-58a^2b^3x^6-32a^3b^2x^4-3a^4bx^2+2a^5)\sqrt{b^3cx^6+3ab^2cx^4+3a^2bcx^2+a^3c}}{143b^2}$	97

input `int(x^3*(c*(b*x^2+a)^3)^(3/2),x,method=_RETURNVERBOSE)`

output $-1/143*(b*x^2+a)*(-11*b*x^2+2*a)*(c*(b*x^2+a)^3)^{(3/2)}/b^2$

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.53

$$\int x^3 \left(c(a + bx^2)^3 \right)^{3/2} dx = \frac{(11 b^5 c x^{10} + 42 a b^4 c x^8 + 58 a^2 b^3 c x^6 + 32 a^3 b^2 c x^4 + 3 a^4 b c x^2 - 2 a^5 c) \sqrt{b^3 c x^6 + 3 a b^2 c x^4 + 3 a^2 b c x^2 + a^3 c}}{143 b^2}$$

input `integrate(x^3*(c*(b*x^2+a)^3)^(3/2),x, algorithm="fricas")`

output $1/143*(11*b^5*c*x^10 + 42*a*b^4*c*x^8 + 58*a^2*b^3*c*x^6 + 32*a^3*b^2*c*x^4 + 3*a^4*b*c*x^2 - 2*a^5*c)*\text{sqrt}(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c)/b^2$

Sympy [F(-1)]

Timed out.

$$\int x^3 \left(c(a + bx^2)^3 \right)^{3/2} dx = \text{Timed out}$$

input `integrate(x**3*(c*(b*x**2+a)**3)**(3/2),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.30

$$\int x^3 \left(c(a + bx^2)^3 \right)^{3/2} dx = \frac{\left(11 b^5 c^{3/2} x^{10} + 42 a b^4 c^{3/2} x^8 + 58 a^2 b^3 c^{3/2} x^6 + 32 a^3 b^2 c^{3/2} x^4 + 3 a^4 b c^{3/2} x^2 - 2 a^5 c^{3/2} \right) (bx^2 + a)^{3/2}}{143 b^2}$$

input `integrate(x^3*(c*(b*x^2+a)^3)^(3/2),x, algorithm="maxima")`output `1/143*(11*b^5*c^(3/2)*x^10 + 42*a*b^4*c^(3/2)*x^8 + 58*a^2*b^3*c^(3/2)*x^6 + 32*a^3*b^2*c^(3/2)*x^4 + 3*a^4*b*c^(3/2)*x^2 - 2*a^5*c^(3/2))*(b*x^2 + a)^(3/2)/b^2`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.83

$$\int x^3 \left(c(a + bx^2)^3 \right)^{3/2} dx = -\frac{13 (bcx^2 + ac)^{\frac{11}{2}} a c \operatorname{sgn}(bx^2 + a) - 11 (bcx^2 + ac)^{\frac{13}{2}} \operatorname{sgn}(bx^2 + a)}{143 b^2 c^5}$$

input `integrate(x^3*(c*(b*x^2+a)^3)^(3/2),x, algorithm="giac")`output `-1/143*(13*(b*c*x^2 + a*c)^(11/2)*a*c*sgn(b*x^2 + a) - 11*(b*c*x^2 + a*c)^(13/2)*sgn(b*x^2 + a))/(b^2*c^5)`

Mupad [B] (verification not implemented)

Time = 0.61 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.12

$$\int x^3 \left(c(a + bx^2)^3 \right)^{3/2} dx = \sqrt{c(bx^2 + a)^3} \left(\frac{32a^3cx^4}{143} - \frac{2a^5c}{143b^2} + \frac{b^3cx^{10}}{13} + \frac{3a^4cx^2}{143b} + \frac{58a^2bcx^6}{143} + \frac{42ab^2cx^8}{143} \right)$$

input `int(x^3*(c*(a + b*x^2)^3)^(3/2),x)`output `(c*(a + b*x^2)^3)^(1/2)*((32*a^3*c*x^4)/143 - (2*a^5*c)/(143*b^2) + (b^3*c*x^10)/13 + (3*a^4*c*x^2)/(143*b) + (58*a^2*b*c*x^6)/143 + (42*a*b^2*c*x^8)/143)`**Reduce [B] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.21

$$\int x^3 \left(c(a + bx^2)^3 \right)^{3/2} dx = \frac{\sqrt{c} \sqrt{bx^2 + a} c (11b^6x^{12} + 53ab^5x^{10} + 100a^2b^4x^8 + 90a^3b^3x^6 + 35a^4b^2x^4 + a^5bx^2 - 2a^6)}{143b^2}$$

input `int(x^3*(c*(b*x^2+a)^3)^(3/2),x)`output `(sqrt(c)*sqrt(a + b*x**2)*c*(- 2*a**6 + a**5*b*x**2 + 35*a**4*b**2*x**4 + 90*a**3*b**3*x**6 + 100*a**2*b**4*x**8 + 53*a*b**5*x**10 + 11*b**6*x**12))/(143*b**2)`

3.13 $\int x \left(c(a + bx^2)^3 \right)^{3/2} dx$

Optimal result	127
Mathematica [A] (verified)	127
Rubi [A] (verified)	128
Maple [A] (verified)	129
Fricas [B] (verification not implemented)	129
Sympy [F]	130
Maxima [B] (verification not implemented)	130
Giac [A] (verification not implemented)	131
Mupad [B] (verification not implemented)	131
Reduce [B] (verification not implemented)	131

Optimal result

Integrand size = 17, antiderivative size = 32

$$\int x \left(c(a + bx^2)^3 \right)^{3/2} dx = \frac{c(a + bx^2)^4 \sqrt{c(a + bx^2)^3}}{11b}$$

output `1/11*c*(b*x^2+a)^4*(c*(b*x^2+a)^3)^(1/2)/b`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.91

$$\int x \left(c(a + bx^2)^3 \right)^{3/2} dx = \frac{(a + bx^2) \left(c(a + bx^2)^3 \right)^{3/2}}{11b}$$

input `Integrate[x*(c*(a + b*x^2)^3)^(3/2),x]`

output `((a + b*x^2)*(c*(a + b*x^2)^3)^(3/2))/(11*b)`

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.91, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2024, 20, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int x \left(c(a + bx^2)^3 \right)^{3/2} dx \\
 \downarrow \text{2024} \\
 \frac{\int \left(c(bx^2 + a)^3 \right)^{3/2} d(bx^2 + a)}{2b} \\
 \downarrow \text{20} \\
 \frac{\left(c(a + bx^2)^3 \right)^{3/2} \int (bx^2 + a)^{9/2} d(bx^2 + a)}{2b(a + bx^2)^{9/2}} \\
 \downarrow \text{15} \\
 \frac{(a + bx^2) \left(c(a + bx^2)^3 \right)^{3/2}}{11b}
 \end{array}$$

input `Int[x*(c*(a + b*x^2)^3)^(3/2),x]`

output `((a + b*x^2)*(c*(a + b*x^2)^3)^(3/2))/(11*b)`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 20 `Int[((a_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a*x^n)^p/x^(n*p) Int[x^(n*p), x], x] /; FreeQ[{a, n, p}, x] && !IntegerQ[p]`

rule 2024

```
Int[((a_.) + (b_.)*(Pq_)^(n_.))^(p_.)*(Qr_), x_Symbol] := With[{q = Expon[P
q, x], r = Expon[Qr, x]}, Simp[Coeff[Qr, x, r]/(q*Coeff[Pq, x, q]) Subst[
Int[(a + b*x^n)^p, x], x, Pq], x] /; EqQ[r, q - 1] && EqQ[Coeff[Qr, x, r]*D
[Pq, x], q*Coeff[Pq, x, q]*Qr]] /; FreeQ[{a, b, n, p}, x] && PolyQ[Pq, x] &
& PolyQ[Qr, x]
```

Maple [A] (verified)

Time = 1.68 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.81

method	result
gospers	$\frac{(bx^2+a)(c(bx^2+a)^3)^{\frac{3}{2}}}{11b}$
orering	$\frac{(bx^2+a)(c(bx^2+a)^3)^{\frac{3}{2}}}{11b}$
risch	$\frac{c\sqrt{c(bx^2+a)^3(b^5x^{10}+5ab^4x^8+10a^2b^3x^6+10a^3b^2x^4+5a^4bx^2+a^5)}}{11(bx^2+a)b}$
trager	$\frac{c(b^4x^8+4ab^3x^6+6a^2b^2x^4+4a^3bx^2+a^4)\sqrt{b^3cx^6+3ab^2cx^4+3a^2bcx^2+a^3c}}{11b}$
default	$-\frac{(c(bx^2+a)^3)^{\frac{3}{2}}\left(-5x^6(x^2bc+ac)^{\frac{5}{2}}b^3-15(x^2bc+ac)^{\frac{5}{2}}ab^2x^4-15(x^2bc+ac)^{\frac{5}{2}}a^2bx^2+6(x^2bc+ac)^{\frac{5}{2}}a^3-11a^3((bx^2+a)c)^{\frac{5}{2}}\right)}{55b(bx^2+a)^3((bx^2+a)c)^{\frac{3}{2}}c}$

input

```
int(x*(c*(b*x^2+a)^3)^(3/2),x,method=_RETURNVERBOSE)
```

output

```
1/11*(b*x^2+a)/b*(c*(b*x^2+a)^3)^(3/2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 87 vs. 2(28) = 56.

Time = 0.09 (sec) , antiderivative size = 87, normalized size of antiderivative = 2.72

$$\int x(c(a + bx^2)^3)^{3/2} dx = \frac{(b^4cx^8 + 4ab^3cx^6 + 6a^2b^2cx^4 + 4a^3bcx^2 + a^4c)\sqrt{b^3cx^6 + 3ab^2cx^4 + 3a^2bcx^2 + a^3c}}{11b}$$

input `integrate(x*(c*(b*x^2+a)^3)^(3/2),x, algorithm="fricas")`

output `1/11*(b^4*c*x^8 + 4*a*b^3*c*x^6 + 6*a^2*b^2*c*x^4 + 4*a^3*b*c*x^2 + a^4*c)*sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c)/b`

Sympy [F]

$$\int x \left(c(a + bx^2)^3 \right)^{3/2} dx = \int x \left(c(a + bx^2)^3 \right)^{\frac{3}{2}} dx$$

input `integrate(x*(c*(b*x**2+a)**3)**(3/2),x)`

output `Integral(x*(c*(a + b*x**2)**3)**(3/2), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 70 vs. 2(28) = 56.

Time = 0.04 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.19

$$\int x \left(c(a + bx^2)^3 \right)^{3/2} dx = \frac{\left(b^4 c^{\frac{3}{2}} x^8 + 4 a b^3 c^{\frac{3}{2}} x^6 + 6 a^2 b^2 c^{\frac{3}{2}} x^4 + 4 a^3 b c^{\frac{3}{2}} x^2 + a^4 c^{\frac{3}{2}} \right) (bx^2 + a)^{\frac{3}{2}}}{11 b}$$

input `integrate(x*(c*(b*x^2+a)^3)^(3/2),x, algorithm="maxima")`

output `1/11*(b^4*c^(3/2)*x^8 + 4*a*b^3*c^(3/2)*x^6 + 6*a^2*b^2*c^(3/2)*x^4 + 4*a^3*b*c^(3/2)*x^2 + a^4*c^(3/2))*(b*x^2 + a)^(3/2)/b`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.88

$$\int x \left(c(a + bx^2)^3 \right)^{3/2} dx = \frac{(bcx^2 + ac)^{\frac{11}{2}} \operatorname{sgn}(bx^2 + a)}{11bc^4}$$

input `integrate(x*(c*(b*x^2+a)^3)^(3/2),x, algorithm="giac")`output `1/11*(b*c*x^2 + a*c)^(11/2)*sgn(b*x^2 + a)/(b*c^4)`**Mupad [B] (verification not implemented)**

Time = 0.51 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.94

$$\int x \left(c(a + bx^2)^3 \right)^{3/2} dx = \sqrt{c(bx^2 + a)^3} \left(\frac{a^4 c}{11b} + \frac{4a^3 c x^2}{11} + \frac{b^3 c x^8}{11} + \frac{6a^2 b c x^4}{11} + \frac{4ab^2 c x^6}{11} \right)$$

input `int(x*(c*(a + b*x^2)^3)^(3/2),x)`output `(c*(a + b*x^2)^3)^(1/2)*((a^4*c)/(11*b) + (4*a^3*c*x^2)/11 + (b^3*c*x^8)/11 + (6*a^2*b*c*x^4)/11 + (4*a*b^2*c*x^6)/11)`**Reduce [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 67, normalized size of antiderivative = 2.09

$$\int x \left(c(a + bx^2)^3 \right)^{3/2} dx = \frac{\sqrt{c} \sqrt{bx^2 + a} c (b^5 x^{10} + 5ab^4 x^8 + 10a^2 b^3 x^6 + 10a^3 b^2 x^4 + 5a^4 b x^2 + a^5)}{11b}$$

input `int(x*(c*(b*x^2+a)^3)^(3/2),x)`

output $(\sqrt{c})\sqrt{a + b*x**2}*c*(a**5 + 5*a**4*b*x**2 + 10*a**3*b**2*x**4 + 10*a**2*b**3*x**6 + 5*a*b**4*x**8 + b**5*x**10))/(11*b)$

$$3.14 \quad \int \frac{\left(c(a+bx^2)^3\right)^{3/2}}{x} dx$$

Optimal result	133
Mathematica [A] (verified)	134
Rubi [A] (verified)	134
Maple [A] (verified)	137
Fricas [A] (verification not implemented)	137
Sympy [F]	138
Maxima [F]	138
Giac [A] (verification not implemented)	139
Mupad [F(-1)]	139
Reduce [B] (verification not implemented)	139

Optimal result

Integrand size = 19, antiderivative size = 192

$$\int \frac{\left(c(a+bx^2)^3\right)^{3/2}}{x} dx = \frac{1}{3}a^3c\sqrt{c(a+bx^2)^3} + \frac{a^4c\sqrt{c(a+bx^2)^3}}{a+bx^2} + \frac{1}{5}a^2c(a+bx^2)\sqrt{c(a+bx^2)^3} + \frac{1}{7}ac(a+bx^2)^2\sqrt{c(a+bx^2)^3} + \frac{1}{9}c(a+bx^2)^3\sqrt{c(a+bx^2)^3} - \frac{a^3c\sqrt{c(a+bx^2)^3}\operatorname{arctanh}\left(\sqrt{1+\frac{bx^2}{a}}\right)}{\left(1+\frac{bx^2}{a}\right)^{3/2}}$$

output

```
1/3*a^3*c*(c*(b*x^2+a)^3)^(1/2)+a^4*c*(c*(b*x^2+a)^3)^(1/2)/(b*x^2+a)+1/5*
a^2*c*(b*x^2+a)*(c*(b*x^2+a)^3)^(1/2)+1/7*a*c*(b*x^2+a)^2*(c*(b*x^2+a)^3)^(
1/2)+1/9*c*(b*x^2+a)^3*(c*(b*x^2+a)^3)^(1/2)-a^3*c*(c*(b*x^2+a)^3)^(1/2)*
arctanh((1+b*x^2/a)^(1/2))/(1+b*x^2/a)^(3/2)
```

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.58

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x} dx = \frac{(c(a + bx^2)^3)^{3/2} (\sqrt{a + bx^2}(563a^4 + 506a^3bx^2 + 408a^2b^2x^4 + 185ab^3x^6 + 35b^4x^8) - 315a^{9/2} \operatorname{ArcTanh}[\frac{\sqrt{a + bx^2}}{\sqrt{a}}])}{315(a + bx^2)^{9/2}}$$

input `Integrate[(c*(a + b*x^2)^3)^(3/2)/x,x]`

output `((c*(a + b*x^2)^3)^(3/2)*(Sqrt[a + b*x^2]*(563*a^4 + 506*a^3*b*x^2 + 408*a^2*b^2*x^4 + 185*a*b^3*x^6 + 35*b^4*x^8) - 315*a^(9/2)*ArcTanh[Sqrt[a + b*x^2]/Sqrt[a]])/(315*(a + b*x^2)^(9/2))`

Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.74, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.474$, Rules used = {2045, 243, 60, 60, 60, 60, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c(a + bx^2)^3)^{3/2}}{x} dx \\ & \quad \downarrow \text{2045} \\ & \frac{a^3 c \sqrt{c(a + bx^2)^3} \int \frac{(\frac{bx^2}{a} + 1)^{9/2}}{x} dx}{\left(\frac{bx^2}{a} + 1\right)^{3/2}} \\ & \quad \downarrow \text{243} \\ & \frac{a^3 c \sqrt{c(a + bx^2)^3} \int \frac{(\frac{bx^2}{a} + 1)^{9/2}}{x^2} dx^2}{2 \left(\frac{bx^2}{a} + 1\right)^{3/2}} \end{aligned}$$

$$\begin{aligned} & \downarrow 60 \\ & \frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\int \frac{\left(\frac{bx^2}{a}+1\right)^{7/2}}{x^2} dx^2 + \frac{2}{9} \left(\frac{bx^2}{a}+1\right)^{9/2} \right)}{2 \left(\frac{bx^2}{a}+1\right)^{3/2}} \\ & \downarrow 60 \\ & \frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\int \frac{\left(\frac{bx^2}{a}+1\right)^{5/2}}{x^2} dx^2 + \frac{2}{9} \left(\frac{bx^2}{a}+1\right)^{9/2} + \frac{2}{7} \left(\frac{bx^2}{a}+1\right)^{7/2} \right)}{2 \left(\frac{bx^2}{a}+1\right)^{3/2}} \\ & \downarrow 60 \\ & \frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\int \frac{\left(\frac{bx^2}{a}+1\right)^{3/2}}{x^2} dx^2 + \frac{2}{9} \left(\frac{bx^2}{a}+1\right)^{9/2} + \frac{2}{7} \left(\frac{bx^2}{a}+1\right)^{7/2} + \frac{2}{5} \left(\frac{bx^2}{a}+1\right)^{5/2} \right)}{2 \left(\frac{bx^2}{a}+1\right)^{3/2}} \\ & \downarrow 60 \\ & \frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\int \frac{\sqrt{\frac{bx^2}{a}+1}}{x^2} dx^2 + \frac{2}{9} \left(\frac{bx^2}{a}+1\right)^{9/2} + \frac{2}{7} \left(\frac{bx^2}{a}+1\right)^{7/2} + \frac{2}{5} \left(\frac{bx^2}{a}+1\right)^{5/2} + \frac{2}{3} \left(\frac{bx^2}{a}+1\right)^{3/2} \right)}{2 \left(\frac{bx^2}{a}+1\right)^{3/2}} \\ & \downarrow 60 \\ & \frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\int \frac{1}{x^2 \sqrt{\frac{bx^2}{a}+1}} dx^2 + \frac{2}{9} \left(\frac{bx^2}{a}+1\right)^{9/2} + \frac{2}{7} \left(\frac{bx^2}{a}+1\right)^{7/2} + \frac{2}{5} \left(\frac{bx^2}{a}+1\right)^{5/2} + \frac{2}{3} \left(\frac{bx^2}{a}+1\right)^{3/2} + 2 \sqrt{\frac{bx^2}{a}+1} \right)}{2 \left(\frac{bx^2}{a}+1\right)^{3/2}} \\ & \downarrow 73 \\ & \frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\frac{2a \int \frac{1}{\frac{ax^4}{b} - \frac{a}{b}} d\sqrt{\frac{bx^2}{a}+1}}{b} + \frac{2}{9} \left(\frac{bx^2}{a}+1\right)^{9/2} + \frac{2}{7} \left(\frac{bx^2}{a}+1\right)^{7/2} + \frac{2}{5} \left(\frac{bx^2}{a}+1\right)^{5/2} + \frac{2}{3} \left(\frac{bx^2}{a}+1\right)^{3/2} + 2 \sqrt{\frac{bx^2}{a}+1} \right)}{2 \left(\frac{bx^2}{a}+1\right)^{3/2}} \\ & \downarrow 221 \end{aligned}$$

$$\frac{a^3 c \left(-2 \operatorname{arctanh} \left(\sqrt{\frac{bx^2}{a} + 1} \right) + \frac{2}{9} \left(\frac{bx^2}{a} + 1 \right)^{9/2} + \frac{2}{7} \left(\frac{bx^2}{a} + 1 \right)^{7/2} + \frac{2}{5} \left(\frac{bx^2}{a} + 1 \right)^{5/2} + \frac{2}{3} \left(\frac{bx^2}{a} + 1 \right)^{3/2} + 2 \sqrt{\frac{bx^2}{a} + 1} \right)}{2 \left(\frac{bx^2}{a} + 1 \right)^{3/2}}$$

input `Int[(c*(a + b*x^2)^3)^(3/2)/x,x]`

output `(a^3*c*Sqrt[c*(a + b*x^2)^3]*(2*Sqrt[1 + (b*x^2)/a] + (2*(1 + (b*x^2)/a)^(3/2))/3 + (2*(1 + (b*x^2)/a)^(5/2))/5 + (2*(1 + (b*x^2)/a)^(7/2))/7 + (2*(1 + (b*x^2)/a)^(9/2))/9 - 2*ArcTanh[Sqrt[1 + (b*x^2)/a]])/(2*(1 + (b*x^2)/a)^(3/2))`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2045

```
Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_.))^(p_), x_Symbol] := Simp[Simp[Si
mp[(c*(a + b*x^n)^q]^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q)
, x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]
```

Maple [A] (verified)

Time = 1.63 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.15

method	result
default	$\frac{(c(bx^2+a)^3)^{\frac{3}{2}} \left(35\sqrt{ac} (x^2bc+ac)^{\frac{5}{2}} b^2 x^4 + 115\sqrt{ac} (x^2bc+ac)^{\frac{5}{2}} ab x^2 - 315 \ln\left(\frac{2ac+2\sqrt{ac}\sqrt{x^2bc+ac}}{x}\right) a^5 c^3 + 189a^2 ((bx^2+a)c)^{\frac{5}{2}} \sqrt{ac} \right)}{315(bx^2+a)^3 (bx^2+a)c^{\frac{3}{2}} \sqrt{ac}}$

input

```
int((c*(b*x^2+a)^3)^(3/2)/x,x,method=_RETURNVERBOSE)
```

output

```
1/315*(c*(b*x^2+a)^3)^(3/2)*(35*(a*c)^(1/2)*(b*c*x^2+a*c)^(5/2)*b^2*x^4+11
5*(a*c)^(1/2)*(b*c*x^2+a*c)^(5/2)*a*b*x^2-315*ln(2*((a*c)^(1/2)*(b*c*x^2+a
*c)^(1/2)+a*c)/x)*a^5*c^3+189*a^2*((b*x^2+a)*c)^(5/2)*(a*c)^(1/2)-46*(a*c)
^(1/2)*(b*c*x^2+a*c)^(5/2)*a^2+105*(a*c)^(1/2)*(b*c*x^2+a*c)^(3/2)*a^3*c+3
15*(a*c)^(1/2)*(b*c*x^2+a*c)^(1/2)*a^4*c^2/(b*x^2+a)^3/((b*x^2+a)*c)^(3/2
)/(a*c)^(1/2)/c
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 382, normalized size of antiderivative = 1.99

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x} dx = \left[\frac{315 (a^4 bcx^2 + a^5 c) \sqrt{ac} \log\left(-\frac{b^2 cx^4 + 3 abcx^2 + 2 a^2 c - 2 \sqrt{b^3 cx^6 + 3 ab^2 cx^4 + 3 a^2 bcx^2 + a^3 c \sqrt{ac}}{bx^4 + ax^2}\right)}{bx^4 + ax^2} \right]$$

input

```
integrate((c*(b*x^2+a)^3)^(3/2)/x,x, algorithm="fricas")
```

output

```
[1/630*(315*(a^4*b*c*x^2 + a^5*c)*sqrt(a*c)*log(-(b^2*c*x^4 + 3*a*b*c*x^2 + 2*a^2*c - 2*sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c))*sqrt(a*c))/(b*x^4 + a*x^2)) + 2*(35*b^4*c*x^8 + 185*a*b^3*c*x^6 + 408*a^2*b^2*c*x^4 + 506*a^3*b*c*x^2 + 563*a^4*c)*sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c))/(b*x^2 + a), 1/315*(315*(a^4*b*c*x^2 + a^5*c)*sqrt(-a*c)*arctan(sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c))*sqrt(-a*c)/(a*b*c*x^2 + a^2*c)) + (35*b^4*c*x^8 + 185*a*b^3*c*x^6 + 408*a^2*b^2*c*x^4 + 506*a^3*b*c*x^2 + 563*a^4*c)*sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c))/(b*x^2 + a)]
```

Sympy [F]

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x} dx = \int \frac{(c(a + bx^2)^3)^{\frac{3}{2}}}{x} dx$$

input

```
integrate((c*(b*x**2+a)**3)**(3/2)/x,x)
```

output

```
Integral((c*(a + b*x**2)**3)**(3/2)/x, x)
```

Maxima [F]

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x} dx = \int \frac{((bx^2 + a)^3 c)^{\frac{3}{2}}}{x} dx$$

input

```
integrate((c*(b*x^2+a)^3)^(3/2)/x,x, algorithm="maxima")
```

output

```
integrate(((b*x^2 + a)^3*c)^(3/2)/x, x)
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 185, normalized size of antiderivative = 0.96

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x} dx = \frac{1}{315} \left(\frac{315 a^5 \arctan\left(\frac{\sqrt{bcx^2+ac}}{\sqrt{-ac}}\right) \operatorname{sgn}(bx^2 + a)}{\sqrt{-ac}} + \frac{315 \sqrt{bcx^2 + ac} a^4 c^{44} \operatorname{sgn}(bx^2 + a)}{\dots} \right)$$

input `integrate((c*(b*x^2+a)^3)^(3/2)/x,x, algorithm="giac")`output `1/315*(315*a^5*arctan(sqrt(b*c*x^2 + a*c)/sqrt(-a*c))*sgn(b*x^2 + a)/sqrt(-a*c) + (315*sqrt(b*c*x^2 + a*c)*a^4*c^44*sgn(b*x^2 + a) + 105*(b*c*x^2 + a*c)^(3/2)*a^3*c^43*sgn(b*x^2 + a) + 63*(b*c*x^2 + a*c)^(5/2)*a^2*c^42*sgn(b*x^2 + a) + 45*(b*c*x^2 + a*c)^(7/2)*a*c^41*sgn(b*x^2 + a) + 35*(b*c*x^2 + a*c)^(9/2)*c^40*sgn(b*x^2 + a))/c^45)*c^2`**Mupad [F(-1)]**

Timed out.

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x} dx = \int \frac{(c(bx^2 + a)^3)^{3/2}}{x} dx$$

input `int((c*(a + b*x^2)^3)^(3/2)/x,x)`output `int((c*(a + b*x^2)^3)^(3/2)/x, x)`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.76

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x} dx = \frac{\sqrt{c} c \left(563 \sqrt{bx^2 + a} a^4 + 506 \sqrt{bx^2 + a} a^3 b x^2 + 408 \sqrt{bx^2 + a} a^2 b^2 x^4 + 185 \sqrt{bx^2 + a} b^3 x^6 \right)}{\dots}$$

input `int((c*(b*x^2+a)^3)^(3/2)/x,x)`

output

```
(sqrt(c)*c*(563*sqrt(a + b*x**2)*a**4 + 506*sqrt(a + b*x**2)*a**3*b*x**2 +
408*sqrt(a + b*x**2)*a**2*b**2*x**4 + 185*sqrt(a + b*x**2)*a*b**3*x**6 +
35*sqrt(a + b*x**2)*b**4*x**8 + 315*sqrt(a)*log((sqrt(a + b*x**2) - sqrt(a)
) + sqrt(b)*x)/sqrt(a))*a**4 - 315*sqrt(a)*log((sqrt(a + b*x**2) + sqrt(a)
+ sqrt(b)*x)/sqrt(a))*a**4)/315
```

3.15
$$\int \frac{(c(a+bx^2)^3)^{3/2}}{x^3} dx$$

Optimal result	141
Mathematica [A] (verified)	142
Rubi [A] (verified)	142
Maple [A] (verified)	146
Fricas [A] (verification not implemented)	146
Sympy [F]	147
Maxima [F]	147
Giac [A] (verification not implemented)	148
Mupad [F(-1)]	148
Reduce [B] (verification not implemented)	148

Optimal result

Integrand size = 19, antiderivative size = 200

$$\int \frac{(c(a+bx^2)^3)^{3/2}}{x^3} dx = a^2bc\sqrt{c(a+bx^2)^3} + \frac{4a^3bc\sqrt{c(a+bx^2)^3}}{a+bx^2} - \frac{a^4c\sqrt{c(a+bx^2)^3}}{2x^2(a+bx^2)} + \frac{2}{5}abc(a+bx^2)\sqrt{c(a+bx^2)^3} + \frac{1}{7}bc(a+bx^2)^2\sqrt{c(a+bx^2)^3} - \frac{9a^2bc\sqrt{c(a+bx^2)^3}\operatorname{arctanh}\left(\sqrt{1+\frac{bx^2}{a}}\right)}{2\left(1+\frac{bx^2}{a}\right)^{3/2}}$$

output

```
a^2*b*c*(c*(b*x^2+a)^3)^(1/2)+4*a^3*b*c*(c*(b*x^2+a)^3)^(1/2)/(b*x^2+a)-1/2*a^4*c*(c*(b*x^2+a)^3)^(1/2)/x^2/(b*x^2+a)+2/5*a*b*c*(b*x^2+a)*(c*(b*x^2+a)^3)^(1/2)+1/7*b*c*(b*x^2+a)^2*(c*(b*x^2+a)^3)^(1/2)-9/2*a^2*b*c*(c*(b*x^2+a)^3)^(1/2)*arctanh((1+b*x^2/a)^(1/2))/(1+b*x^2/a)^(3/2)
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.59

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^3} dx = \frac{(c(a + bx^2)^3)^{3/2} \left(\sqrt{a + bx^2} (35a^4 - 388a^3bx^2 - 156a^2b^2x^4 - 58ab^3x^6 - 10b^4x^8) + 315a^{7/2}bx^2 \operatorname{arctanh}\left(\frac{\sqrt{a + bx^2}}{\sqrt{a}}\right) \right)}{70x^2 (a + bx^2)^{9/2}}$$

input `Integrate[(c*(a + b*x^2)^3)^(3/2)/x^3,x]`

output `-1/70*((c*(a + b*x^2)^3)^(3/2)*(Sqrt[a + b*x^2]*(35*a^4 - 388*a^3*b*x^2 - 156*a^2*b^2*x^4 - 58*a*b^3*x^6 - 10*b^4*x^8) + 315*a^(7/2)*b*x^2*ArcTanh[Sqrt[a + b*x^2]/Sqrt[a]]))/(x^2*(a + b*x^2)^(9/2))`

Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.76, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.474$, Rules used = {2045, 243, 51, 60, 60, 60, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^3} dx$$

↓ 2045

$$\frac{a^3 c \sqrt{c(a + bx^2)^3} \int \frac{\left(\frac{bx^2}{a} + 1\right)^{9/2}}{x^3} dx}{\left(\frac{bx^2}{a} + 1\right)^{3/2}}$$

↓ 243

$$\begin{aligned}
& \frac{a^3 c \sqrt{c(a+bx^2)^3} \int \frac{\left(\frac{bx^2}{a}+1\right)^{9/2}}{x^4} dx^2}{2\left(\frac{bx^2}{a}+1\right)^{3/2}} \\
& \quad \downarrow 51 \\
& \frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\frac{9b \int \frac{\left(\frac{bx^2}{a}+1\right)^{7/2}}{x^2} dx^2}{2a} - \frac{\left(\frac{bx^2}{a}+1\right)^{9/2}}{x^2} \right)}{2\left(\frac{bx^2}{a}+1\right)^{3/2}} \\
& \quad \downarrow 60 \\
& \frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\frac{9b \left(\int \frac{\left(\frac{bx^2}{a}+1\right)^{5/2}}{x^2} dx^2 + \frac{2}{7} \left(\frac{bx^2}{a}+1\right)^{7/2} \right)}{2a} - \frac{\left(\frac{bx^2}{a}+1\right)^{9/2}}{x^2} \right)}{2\left(\frac{bx^2}{a}+1\right)^{3/2}} \\
& \quad \downarrow 60 \\
& \frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\frac{9b \left(\int \frac{\left(\frac{bx^2}{a}+1\right)^{3/2}}{x^2} dx^2 + \frac{2}{7} \left(\frac{bx^2}{a}+1\right)^{7/2} + \frac{2}{5} \left(\frac{bx^2}{a}+1\right)^{5/2} \right)}{2a} - \frac{\left(\frac{bx^2}{a}+1\right)^{9/2}}{x^2} \right)}{2\left(\frac{bx^2}{a}+1\right)^{3/2}} \\
& \quad \downarrow 60 \\
& \frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\frac{9b \left(\int \frac{\sqrt{\frac{bx^2}{a}+1}}{x^2} dx^2 + \frac{2}{7} \left(\frac{bx^2}{a}+1\right)^{7/2} + \frac{2}{5} \left(\frac{bx^2}{a}+1\right)^{5/2} + \frac{2}{3} \left(\frac{bx^2}{a}+1\right)^{3/2} \right)}{2a} - \frac{\left(\frac{bx^2}{a}+1\right)^{9/2}}{x^2} \right)}{2\left(\frac{bx^2}{a}+1\right)^{3/2}} \\
& \quad \downarrow 60
\end{aligned}$$

$$a^3 c \sqrt{c(a + bx^2)^3} \left(\frac{9b \left(\int \frac{1}{x^2 \sqrt{\frac{bx^2}{a} + 1}} dx^2 + \frac{2}{7} \left(\frac{bx^2}{a} + 1 \right)^{7/2} + \frac{2}{5} \left(\frac{bx^2}{a} + 1 \right)^{5/2} + \frac{2}{3} \left(\frac{bx^2}{a} + 1 \right)^{3/2} + 2\sqrt{\frac{bx^2}{a} + 1} \right)}{2a} - \frac{\left(\frac{bx^2}{a} + 1 \right)^{9/2}}{x^2} \right)$$

$$2 \left(\frac{bx^2}{a} + 1 \right)^{3/2}$$

↓ 73

$$a^3 c \sqrt{c(a + bx^2)^3} \left(\frac{9b \left(\frac{2a \int \frac{1}{\frac{ax^4}{b} - \frac{a}{b}} d\sqrt{\frac{bx^2}{a} + 1}}{\frac{ax^4}{b} - \frac{a}{b}} + \frac{2}{7} \left(\frac{bx^2}{a} + 1 \right)^{7/2} + \frac{2}{5} \left(\frac{bx^2}{a} + 1 \right)^{5/2} + \frac{2}{3} \left(\frac{bx^2}{a} + 1 \right)^{3/2} + 2\sqrt{\frac{bx^2}{a} + 1} \right)}{2a} - \frac{\left(\frac{bx^2}{a} + 1 \right)^{9/2}}{x^2} \right)$$

$$2 \left(\frac{bx^2}{a} + 1 \right)^{3/2}$$

↓ 221

$$a^3 c \left(\frac{9b \left(-2 \operatorname{arctanh} \left(\sqrt{\frac{bx^2}{a} + 1} \right) + \frac{2}{7} \left(\frac{bx^2}{a} + 1 \right)^{7/2} + \frac{2}{5} \left(\frac{bx^2}{a} + 1 \right)^{5/2} + \frac{2}{3} \left(\frac{bx^2}{a} + 1 \right)^{3/2} + 2\sqrt{\frac{bx^2}{a} + 1} \right)}{2a} - \frac{\left(\frac{bx^2}{a} + 1 \right)^{9/2}}{x^2} \right) \sqrt{c(a + bx^2)^3}$$

$$2 \left(\frac{bx^2}{a} + 1 \right)^{3/2}$$

input `Int[(c*(a + b*x^2)^3)^(3/2)/x^3,x]`

output `(a^3*c*Sqrt[c*(a + b*x^2)^3]*(-(1 + (b*x^2)/a)^(9/2)/x^2) + (9*b*(2*Sqrt[1 + (b*x^2)/a] + (2*(1 + (b*x^2)/a)^(3/2))/3 + (2*(1 + (b*x^2)/a)^(5/2))/5 + (2*(1 + (b*x^2)/a)^(7/2))/7 - 2*ArcTanh[Sqrt[1 + (b*x^2)/a]]))/(2*a)))/(2*(1 + (b*x^2)/a)^(3/2))`

Definitions of rubi rules used

rule 51 $\text{Int}[(a_.) + (b_.)(x_)^{(m_.)}((c_.) + (d_.)(x_)^{(n_.)}), x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}((c + d*x)^n/(b*(m + 1))), x] - \text{Simp}[d*(n/(b*(m + 1))), x] - \text{Simp}[(a + b*x)^{(m + 1)}(c + d*x)^{(n - 1)}, x], x] /;$ FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]

rule 60 $\text{Int}[(a_.) + (b_.)(x_)^{(m_.)}((c_.) + (d_.)(x_)^{(n_.)}), x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}((c + d*x)^n/(b*(m + n + 1))), x] + \text{Simp}[n*((b*c - a*d)/(b*(m + n + 1))) \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /;$ FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

rule 73 $\text{Int}[(a_.) + (b_.)(x_)^{(m_.)}((c_.) + (d_.)(x_)^{(n_.)}), x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \text{Subst}[\text{Int}[x^{(p*(m + 1) - 1)}*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^{(1/p)}, x]] /;$ FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

rule 221 $\text{Int}[(a_.) + (b_.)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /;$ FreeQ[{a, b}, x] && NegQ[a/b]

rule 243 $\text{Int}[(x_)^{(m_.)}((a_.) + (b_.)(x_)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{Subst}[\text{Int}[x^{((m - 1)/2)*(a + b*x)^p}, x], x, x^2], x] /;$ FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]

rule 2045 $\text{Int}[(u_.)*((c_.)*((a_.) + (b_.)(x_)^{(n_.)})^{(q_.)})^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[\text{Simp}[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^{(p*q)} \text{Int}[u*(1 + b*(x^n/a))^{(p*q)}, x], x] /;$ FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]

Maple [A] (verified)

Time = 2.47 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.12

method	result
risch	$-\frac{a^4 c \sqrt{c(bx^2+a)^3}}{2x^2(bx^2+a)} + \frac{\left(-\frac{9b a^4 \ln\left(\frac{2ac+2\sqrt{ac}\sqrt{x^2bc+ac}}{x}\right)}{2\sqrt{ac}} + \frac{b^4 x^6 \sqrt{x^2bc+ac}}{7c} + \frac{29b^3 a x^4 \sqrt{x^2bc+ac}}{35c} + \frac{78b^2 a^2 x^2 \sqrt{x^2bc+ac}}{35c} - \frac{156b a^3 \sqrt{x^2bc+ac}}{35c} \right)}{(bx^2+a)^2}$
default	$\frac{(c(bx^2+a)^3)^{\frac{3}{2}} \left(10\sqrt{ac}(x^2bc+ac)^{\frac{5}{2}} b^2 x^4 - 315 \ln\left(\frac{2ac+2\sqrt{ac}\sqrt{x^2bc+ac}}{x}\right) a^4 b c^3 x^2 + 42ab((bx^2+a)c)^{\frac{5}{2}} x^2 \sqrt{ac} - 4\sqrt{ac}(x^2bc+ac)^{\frac{5}{2}} \right)}{70(bx^2+a)^3 ((bx^2+a)c)^{\frac{3}{2}} c x^2 \sqrt{ac}}$

```
input int((c*(b*x^2+a)^3)^(3/2)/x^3,x,method=_RETURNVERBOSE)
```

```
output -1/2*a^4*c*(c*(b*x^2+a)^3)^(1/2)/x^2/(b*x^2+a)+(-9/2*b*a^4/(a*c)^(1/2)*ln(
(2*a*c+2*(a*c)^(1/2)*(b*c*x^2+a*c)^(1/2))/x)+1/7*b^4*x^6/c*(b*c*x^2+a*c)^(
1/2)+29/35*b^3*a*x^4/c*(b*c*x^2+a*c)^(1/2)+78/35*b^2*a^2*x^2/c*(b*c*x^2+a*
c)^(1/2)-156/35*b*a^3/c*(b*c*x^2+a*c)^(1/2)+10*b*a^3/c*((b*x^2+a)*c)^(1/2)
)*c/(b*x^2+a)^2*(c*(b*x^2+a)^3)^(1/2)*((b*x^2+a)*c)^(1/2)
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 402, normalized size of antiderivative = 2.01

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^3} dx = \left[\frac{315 (a^3 b^2 c x^4 + a^4 b c x^2) \sqrt{ac} \log\left(-\frac{b^2 c x^4 + 3 a b c x^2 + 2 a^2 c - 2 \sqrt{b^3 c x^6 + 3 a b^2 c x^4 + 3 a^2 b c x^2 + a^3 c} \sqrt{bx^2+ax^2}}{bx^4+ax^2}\right)}{\dots} \right]$$

```
input integrate((c*(b*x^2+a)^3)^(3/2)/x^3,x, algorithm="fricas")
```

output

```
[1/140*(315*(a^3*b^2*c*x^4 + a^4*b*c*x^2)*sqrt(a*c)*log(-(b^2*c*x^4 + 3*a*
b*c*x^2 + 2*a^2*c - 2*sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3
*c)*sqrt(a*c))/(b*x^4 + a*x^2)) + 2*(10*b^4*c*x^8 + 58*a*b^3*c*x^6 + 156*a
^2*b^2*c*x^4 + 388*a^3*b*c*x^2 - 35*a^4*c)*sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4
+ 3*a^2*b*c*x^2 + a^3*c))/(b*x^4 + a*x^2), 1/70*(315*(a^3*b^2*c*x^4 + a^4*
b*c*x^2)*sqrt(-a*c)*arctan(sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2
+ a^3*c)*sqrt(-a*c)/(a*b*c*x^2 + a^2*c)) + (10*b^4*c*x^8 + 58*a*b^3*c*x^6
+ 156*a^2*b^2*c*x^4 + 388*a^3*b*c*x^2 - 35*a^4*c)*sqrt(b^3*c*x^6 + 3*a*b^2
*c*x^4 + 3*a^2*b*c*x^2 + a^3*c))/(b*x^4 + a*x^2)]
```

Sympy [F]

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^3} dx = \int \frac{(c(a + bx^2)^3)^{\frac{3}{2}}}{x^3} dx$$

input

```
integrate((c*(b*x**2+a)**3)**(3/2)/x**3,x)
```

output

```
Integral((c*(a + b*x**2)**3)**(3/2)/x**3, x)
```

Maxima [F]

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^3} dx = \int \frac{((bx^2 + a)^3 c)^{\frac{3}{2}}}{x^3} dx$$

input

```
integrate((c*(b*x^2+a)^3)^(3/2)/x^3,x, algorithm="maxima")
```

output

```
integrate(((b*x^2 + a)^3*c)^(3/2)/x^3, x)
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.98

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^3} dx = \frac{1}{70} \left(\frac{315 a^4 \arctan\left(\frac{\sqrt{bcx^2+ac}}{\sqrt{-ac}}\right) \operatorname{sgn}(bx^2 + a)}{\sqrt{-acc}} - \frac{35 \sqrt{bcx^2 + ac} a^4 \operatorname{sgn}(bx^2 + a)}{bc^2 x^2} + \dots \right)$$

input `integrate((c*(b*x^2+a)^3)^(3/2)/x^3,x, algorithm="giac")`

output `1/70*(315*a^4*arctan(sqrt(b*c*x^2 + a*c)/sqrt(-a*c))*sgn(b*x^2 + a)/(sqrt(-a*c)*c) - 35*sqrt(b*c*x^2 + a*c)*a^4*sgn(b*x^2 + a)/(b*c^2*x^2) + 2*(140*sqrt(b*c*x^2 + a*c)*a^3*c^33*sgn(b*x^2 + a) + 35*(b*c*x^2 + a*c)^(3/2)*a^2*c^32*sgn(b*x^2 + a) + 14*(b*c*x^2 + a*c)^(5/2)*a*c^31*sgn(b*x^2 + a) + 5*(b*c*x^2 + a*c)^(7/2)*c^30*sgn(b*x^2 + a))/c^35*b*c^3`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^3} dx = \int \frac{(c(bx^2 + a)^3)^{3/2}}{x^3} dx$$

input `int((c*(a + b*x^2)^3)^(3/2)/x^3,x)`

output `int((c*(a + b*x^2)^3)^(3/2)/x^3, x)`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.78

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^3} dx = \frac{\sqrt{c} c \left(-35 \sqrt{bx^2 + a} a^4 + 388 \sqrt{bx^2 + a} a^3 b x^2 + 156 \sqrt{bx^2 + a} a^2 b^2 x^4 + 58 \sqrt{bx^2 + a} a b^3 x^6 + 14 b^4 x^8 \right)}{70 x^7}$$

input `int((c*(b*x^2+a)^3)^(3/2)/x^3,x)`

output

```
(sqrt(c)*c*( - 35*sqrt(a + b*x**2)*a**4 + 388*sqrt(a + b*x**2)*a**3*b*x**2
+ 156*sqrt(a + b*x**2)*a**2*b**2*x**4 + 58*sqrt(a + b*x**2)*a*b**3*x**6 +
10*sqrt(a + b*x**2)*b**4*x**8 + 315*sqrt(a)*log((sqrt(a + b*x**2) - sqrt(
a) + sqrt(b)*x)/sqrt(a))*a**3*b*x**2 - 315*sqrt(a)*log((sqrt(a + b*x**2) +
sqrt(a) + sqrt(b)*x)/sqrt(a))*a**3*b*x**2))/(70*x**2)
```

3.16 $\int x^2 \left(c(a + bx^2)^3 \right)^{3/2} dx$

Optimal result	150
Mathematica [A] (verified)	151
Rubi [A] (verified)	151
Maple [A] (verified)	154
Fricas [A] (verification not implemented)	154
Sympy [F]	155
Maxima [F]	155
Giac [A] (verification not implemented)	156
Mupad [F(-1)]	156
Reduce [B] (verification not implemented)	157

Optimal result

Integrand size = 19, antiderivative size = 253

$$\begin{aligned} \int x^2 \left(c(a + bx^2)^3 \right)^{3/2} dx &= \frac{7}{128} a^3 c x^3 \sqrt{c(a + bx^2)^3} \\ &+ \frac{21a^5 c x \sqrt{c(a + bx^2)^3}}{1024b(a + bx^2)} + \frac{21a^4 c x^3 \sqrt{c(a + bx^2)^3}}{512(a + bx^2)} \\ &+ \frac{21}{320} a^2 c x^3 (a + bx^2) \sqrt{c(a + bx^2)^3} + \frac{3}{40} a c x^3 (a + bx^2)^2 \sqrt{c(a + bx^2)^3} \\ &+ \frac{1}{12} c x^3 (a + bx^2)^3 \sqrt{c(a + bx^2)^3} - \frac{21a^{9/2} c \sqrt{c(a + bx^2)^3} \operatorname{arcsinh}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{1024b^{3/2} \left(1 + \frac{bx^2}{a}\right)^{3/2}} \end{aligned}$$

output $\frac{7}{128}a^3c*x^3*(c*(b*x^2+a)^3)^{(1/2)}+21/1024*a^5*c*x*(c*(b*x^2+a)^3)^{(1/2)}/b/(b*x^2+a)+21*a^4*c*x^3*(c*(b*x^2+a)^3)^{(1/2)}/(512*b*x^2+512*a)+21/320*a^2*c*x^3*(b*x^2+a)*(c*(b*x^2+a)^3)^{(1/2)}+3/40*a*c*x^3*(b*x^2+a)^2*(c*(b*x^2+a)^3)^{(1/2)}+1/12*c*x^3*(b*x^2+a)^3*(c*(b*x^2+a)^3)^{(1/2)}-21/1024*a^{(9/2)}*c*(c*(b*x^2+a)^3)^{(1/2)}*\operatorname{arcsinh}(b^{(1/2)}*x/a^{(1/2)})/b^{(3/2)}/(1+b*x^2/a)^{(3/2)}$

Mathematica [A] (verified)

Time = 0.60 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.56

$$\int x^2 \left(c(a + bx^2)^3 \right)^{3/2} dx = \frac{\left(c(a + bx^2)^3 \right)^{3/2} \left(\sqrt{bx} \sqrt{a + bx^2} (315a^5 + 4910a^4bx^2 + 11432a^3b^2x^4 + 12144a^2b^3x^6 + 6272ab^4x^8 + 1280b^5x^{10}) + 630a^6 \operatorname{ArcTanh} \left[\frac{\sqrt{bx}}{\sqrt{a + bx^2}} \right] \right)}{15360b^{3/2} (a + bx^2)^{9/2}}$$

input `Integrate[x^2*(c*(a + b*x^2)^3)^(3/2),x]`

output `((c*(a + b*x^2)^3)^(3/2)*(Sqrt[b]*x*Sqrt[a + b*x^2]*(315*a^5 + 4910*a^4*b*x^2 + 11432*a^3*b^2*x^4 + 12144*a^2*b^3*x^6 + 6272*a*b^4*x^8 + 1280*b^5*x^10) + 630*a^6*ArcTanh[(Sqrt[b]*x)/(Sqrt[a] - Sqrt[a + b*x^2])]))/(15360*b^(3/2)*(a + b*x^2)^(9/2))`

Rubi [A] (verified)

Time = 0.55 (sec) , antiderivative size = 215, normalized size of antiderivative = 0.85, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$, Rules used = {2045, 248, 248, 248, 248, 248, 262, 222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^2 \left(c(a + bx^2)^3 \right)^{3/2} dx \\ & \quad \downarrow \text{2045} \\ & \frac{a^3 c \sqrt{c(a + bx^2)^3} \int x^2 \left(\frac{bx^2}{a} + 1 \right)^{9/2} dx}{\left(\frac{bx^2}{a} + 1 \right)^{3/2}} \\ & \quad \downarrow \text{248} \end{aligned}$$

$$\frac{a^3 c \sqrt{c(a+bx^2)}^3 \left(\frac{3}{4} \int x^2 \left(\frac{bx^2}{a} + 1 \right)^{7/2} dx + \frac{1}{12} x^3 \left(\frac{bx^2}{a} + 1 \right)^{9/2} \right)}{\left(\frac{bx^2}{a} + 1 \right)^{3/2}}$$

↓ 248

$$\frac{a^3 c \sqrt{c(a+bx^2)}^3 \left(\frac{3}{4} \left(\frac{7}{10} \int x^2 \left(\frac{bx^2}{a} + 1 \right)^{5/2} dx + \frac{1}{10} x^3 \left(\frac{bx^2}{a} + 1 \right)^{7/2} \right) + \frac{1}{12} x^3 \left(\frac{bx^2}{a} + 1 \right)^{9/2} \right)}{\left(\frac{bx^2}{a} + 1 \right)^{3/2}}$$

↓ 248

$$\frac{a^3 c \sqrt{c(a+bx^2)}^3 \left(\frac{3}{4} \left(\frac{7}{10} \left(\frac{5}{8} \int x^2 \left(\frac{bx^2}{a} + 1 \right)^{3/2} dx + \frac{1}{8} x^3 \left(\frac{bx^2}{a} + 1 \right)^{5/2} \right) + \frac{1}{10} x^3 \left(\frac{bx^2}{a} + 1 \right)^{7/2} \right) + \frac{1}{12} x^3 \left(\frac{bx^2}{a} + 1 \right)^{9/2} \right)}{\left(\frac{bx^2}{a} + 1 \right)^{3/2}}$$

↓ 248

$$\frac{a^3 c \sqrt{c(a+bx^2)}^3 \left(\frac{3}{4} \left(\frac{7}{10} \left(\frac{5}{8} \left(\frac{1}{2} \int x^2 \sqrt{\frac{bx^2}{a} + 1} dx + \frac{1}{6} x^3 \left(\frac{bx^2}{a} + 1 \right)^{3/2} \right) + \frac{1}{8} x^3 \left(\frac{bx^2}{a} + 1 \right)^{5/2} \right) + \frac{1}{10} x^3 \left(\frac{bx^2}{a} + 1 \right)^{7/2} \right) \right)}{\left(\frac{bx^2}{a} + 1 \right)^{3/2}}$$

↓ 248

$$\frac{a^3 c \sqrt{c(a+bx^2)}^3 \left(\frac{3}{4} \left(\frac{7}{10} \left(\frac{5}{8} \left(\frac{1}{2} \left(\frac{1}{4} \int \frac{x^2}{\sqrt{\frac{bx^2}{a} + 1}} dx + \frac{1}{4} x^3 \sqrt{\frac{bx^2}{a} + 1} \right) + \frac{1}{6} x^3 \left(\frac{bx^2}{a} + 1 \right)^{3/2} \right) + \frac{1}{8} x^3 \left(\frac{bx^2}{a} + 1 \right)^{5/2} \right) \right) \right)}{\left(\frac{bx^2}{a} + 1 \right)^{3/2}}$$

↓ 262

$$\frac{a^3 c \sqrt{c(a+bx^2)}^3 \left(\frac{3}{4} \left(\frac{7}{10} \left(\frac{5}{8} \left(\frac{1}{2} \left(\frac{1}{4} \left(\frac{ax \sqrt{\frac{bx^2}{a} + 1}}{2b} - \frac{a \int \frac{1}{\sqrt{\frac{bx^2}{a} + 1}} dx}{2b} \right) + \frac{1}{4} x^3 \sqrt{\frac{bx^2}{a} + 1} \right) + \frac{1}{6} x^3 \left(\frac{bx^2}{a} + 1 \right)^{3/2} \right) \right) \right) + \frac{1}{8} x^3 \left(\frac{bx^2}{a} + 1 \right)^{5/2} \right)}{\left(\frac{bx^2}{a} + 1 \right)^{3/2}}$$

↓ 222

$$\frac{a^3 c \left(\frac{3}{4} \left(\frac{7}{10} \left(\frac{5}{8} \left(\frac{1}{2} \left(\frac{1}{4} \left(\frac{ax \sqrt{\frac{bx^2}{a} + 1}}{2b} - \frac{a^{3/2} \operatorname{arcsinh}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{2b^{3/2}} \right) + \frac{1}{4} x^3 \sqrt{\frac{bx^2}{a} + 1} \right) + \frac{1}{6} x^3 \left(\frac{bx^2}{a} + 1\right)^{3/2} \right) + \frac{1}{8} x^3 \left(\frac{bx^2}{a} + 1\right)^{5/2} \right) \right) \right) \right) \right) \right) \right) \left(\frac{bx^2}{a} + 1\right)^{3/2}}$$

input `Int[x^2*(c*(a + b*x^2)^3)^(3/2),x]`

output `(a^3*c*Sqrt[c*(a + b*x^2)^3]*((x^3*(1 + (b*x^2)/a)^(9/2))/12 + (3*((x^3*(1 + (b*x^2)/a)^(7/2))/10 + (7*((x^3*(1 + (b*x^2)/a)^(5/2))/8 + (5*((x^3*(1 + (b*x^2)/a)^(3/2))/6 + ((x^3*Sqrt[1 + (b*x^2)/a])/4 + ((a*x*Sqrt[1 + (b*x^2)/a])/(2*b) - (a^(3/2)*ArcSinh[(Sqrt[b]*x)/Sqrt[a]])/(2*b^(3/2)))/4)/2))/8))/10))/4)/(1 + (b*x^2)/a)^(3/2)`

Defintions of rubi rules used

rule 222 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

rule 248 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^p/(c*(m + 2*p + 1))), x] + Simp[2*a*(p/(m + 2*p + 1)) Int[(c*x)^m*(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c, m}, x] && GtQ[p, 0] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 262 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2045 `Int[(u_)*((c_)*((a_) + (b_)*(x_)^(n_))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 2.15 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.61

method	result
risch	$\frac{x(1280b^5x^{10}+6272ab^4x^8+12144a^2b^3x^6+11432a^3b^2x^4+4910a^4bx^2+315a^5)c\sqrt{c(bx^2+a)^3}}{15360b(bx^2+a)} - \frac{21a^6 \ln\left(\frac{bcx}{\sqrt{bc}} + \sqrt{x^2bc+ac}\right)c\sqrt{c(bx^2+a)^3}}{1024b\sqrt{bc}(bx^2+a)}$
default	$\frac{(c(bx^2+a)^3)^{\frac{3}{2}} \left(1280x^7(x^2bc+ac)^{\frac{5}{2}}b^3\sqrt{bc}+3712\sqrt{bc}(x^2bc+ac)^{\frac{5}{2}}ab^2x^5+3440\sqrt{bc}(x^2bc+ac)^{\frac{5}{2}}a^2bx^3+840\sqrt{bc}(x^2bc+ac)^{\frac{5}{2}}a^3x\right)}{15360b(bx^2+a)^3((bx^2+a)c)^{\frac{3}{2}}c\sqrt{bc}}$

input `int(x^2*(c*(b*x^2+a)^3)^(3/2),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{15360}xx(1280b^5x^{10}+6272a^2b^4x^8+12144a^2b^3x^6+11432a^3b^2x^4+4910a^4bx^2+315a^5)/b/(bx^2+a)*c*(c*(bx^2+a)^3)^{(1/2)}-21/1024/b*a^6*\ln(b*c*x/(b*c)^{(1/2)}+(b*c*x^2+a*c)^{(1/2)})/(b*c)^{(1/2)}*c/(b*x^2+a)^2*(c*(b*x^2+a)^3)^{(1/2)}*((b*x^2+a)*c)^{(1/2)}$$

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 433, normalized size of antiderivative = 1.71

$$\int x^2 \left(c(a + bx^2)^3 \right)^{3/2} dx = \left[\frac{315 (a^6bcx^2 + a^7c) \sqrt{\frac{c}{b}} \log \left(-\frac{2b^2cx^4 + 3abcx^2 + a^2c - 2\sqrt{b^3cx^6 + 3ab^2cx^4 + 3a^2bcx^2 + a^3cbx} \sqrt{\frac{c}{b}}}{bx^2+a} \right) + 2(1280x^7(x^2bc+ac)^{\frac{5}{2}}b^3\sqrt{bc}+3712\sqrt{bc}(x^2bc+ac)^{\frac{5}{2}}ab^2x^5+3440\sqrt{bc}(x^2bc+ac)^{\frac{5}{2}}a^2bx^3+840\sqrt{bc}(x^2bc+ac)^{\frac{5}{2}}a^3x)}{15360b(bx^2+a)^3((bx^2+a)c)^{\frac{3}{2}}c\sqrt{bc}} \right]$$

input `integrate(x^2*(c*(b*x^2+a)^3)^(3/2),x, algorithm="fricas")`

output

```
[1/30720*(315*(a^6*b*c*x^2 + a^7*c)*sqrt(c/b)*log(-(2*b^2*c*x^4 + 3*a*b*c*x^2 + a^2*c - 2*sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c))*x*sqrt(c/b))/(b*x^2 + a)) + 2*(1280*b^5*c*x^11 + 6272*a*b^4*c*x^9 + 12144*a^2*b^3*c*x^7 + 11432*a^3*b^2*c*x^5 + 4910*a^4*b*c*x^3 + 315*a^5*c*x)*sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c))/(b^2*x^2 + a*b), 1/15360*(315*(a^6*b*c*x^2 + a^7*c)*sqrt(-c/b)*arctan(sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c))*b*x*sqrt(-c/b)/(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)) + (1280*b^5*c*x^11 + 6272*a*b^4*c*x^9 + 12144*a^2*b^3*c*x^7 + 11432*a^3*b^2*c*x^5 + 4910*a^4*b*c*x^3 + 315*a^5*c*x)*sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c))/(b^2*x^2 + a*b)]
```

Sympy [F]

$$\int x^2 \left(c(a + bx^2)^3 \right)^{3/2} dx = \int x^2 \left(c(a + bx^2)^3 \right)^{\frac{3}{2}} dx$$

input

```
integrate(x**2*(c*(b*x**2+a)**3)**(3/2),x)
```

output

```
Integral(x**2*(c*(a + b*x**2)**3)**(3/2), x)
```

Maxima [F]

$$\int x^2 \left(c(a + bx^2)^3 \right)^{3/2} dx = \int \left((bx^2 + a)^3 c \right)^{\frac{3}{2}} x^2 dx$$

input

```
integrate(x^2*(c*(b*x^2+a)^3)^(3/2),x, algorithm="maxima")
```

output

```
integrate(((b*x^2 + a)^3*c)^(3/2)*x^2, x)
```

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.70

$$\int x^2 \left(c(a + bx^2)^3 \right)^{3/2} dx = \frac{1}{15360} \left(\frac{315 a^6 c \log \left(\left| -\sqrt{bc}x + \sqrt{bcx^2 + ac} \right| \right) \operatorname{sgn}(bx^2 + a)}{\sqrt{bc}b} + \left(\frac{315 a^5 \operatorname{sgn}(bx^2 + a)}{b} + 2 \right) \right)$$

input `integrate(x^2*(c*(b*x^2+a)^3)^(3/2),x, algorithm="giac")`

output `1/15360*(315*a^6*c*log(abs(-sqrt(b*c)*x + sqrt(b*c*x^2 + a*c)))*sgn(b*x^2 + a)/(sqrt(b*c)*b) + (315*a^5*sgn(b*x^2 + a)/b + 2*(2455*a^4*sgn(b*x^2 + a) + 4*(1429*a^3*b*sgn(b*x^2 + a) + 2*(759*a^2*b^2*sgn(b*x^2 + a) + 8*(10*b^4*x^2*sgn(b*x^2 + a) + 49*a*b^3*sgn(b*x^2 + a))*x^2)*x^2)*sqrt(b*c*x^2 + a*c)*x)*c`

Mupad [F(-1)]

Timed out.

$$\int x^2 \left(c(a + bx^2)^3 \right)^{3/2} dx = \int x^2 \left(c(bx^2 + a)^3 \right)^{3/2} dx$$

input `int(x^2*(c*(a + b*x^2)^3)^(3/2),x)`

output `int(x^2*(c*(a + b*x^2)^3)^(3/2), x)`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.55

$$\int x^2 \left(c(a + bx^2)^3 \right)^{3/2} dx = \frac{\sqrt{c} c \left(315\sqrt{bx^2 + a} a^5 bx + 4910\sqrt{bx^2 + a} a^4 b^2 x^3 + 11432\sqrt{bx^2 + a} a^3 b^3 x^5 + 12144\sqrt{bx^2 + a} a^2 b^4 x^7 + 6272\sqrt{bx^2 + a} a b^5 x^9 + 1280\sqrt{bx^2 + a} b^6 x^{11} - 315\sqrt{b} \log\left(\frac{\sqrt{bx^2 + a} + \sqrt{b}x}{\sqrt{a}}\right) a^6 \right)}{15360 b^2}$$

input

```
int(x^2*(c*(b*x^2+a)^3)^(3/2),x)
```

output

```
(sqrt(c)*c*(315*sqrt(a + b*x**2)*a**5*b*x + 4910*sqrt(a + b*x**2)*a**4*b**2*x**3 + 11432*sqrt(a + b*x**2)*a**3*b**3*x**5 + 12144*sqrt(a + b*x**2)*a**2*b**4*x**7 + 6272*sqrt(a + b*x**2)*a*b**5*x**9 + 1280*sqrt(a + b*x**2)*b**6*x**11 - 315*sqrt(b)*log((sqrt(a + b*x**2) + sqrt(b)*x)/sqrt(a))*a**6)/(15360*b**2)
```

$$3.17 \quad \int \left(c(a + bx^2)^3 \right)^{3/2} dx$$

Optimal result	158
Mathematica [A] (verified)	159
Rubi [A] (verified)	159
Maple [A] (verified)	161
Fricas [A] (verification not implemented)	162
Sympy [F]	162
Maxima [F]	163
Giac [A] (verification not implemented)	163
Mupad [F(-1)]	163
Reduce [B] (verification not implemented)	164

Optimal result

Integrand size = 15, antiderivative size = 207

$$\begin{aligned} \int \left(c(a + bx^2)^3 \right)^{3/2} dx &= \frac{21}{128} a^3 cx \sqrt{c(a + bx^2)^3} + \frac{63a^4 cx \sqrt{c(a + bx^2)^3}}{256(a + bx^2)} \\ &+ \frac{21}{160} a^2 cx (a + bx^2) \sqrt{c(a + bx^2)^3} + \frac{9}{80} acx (a + bx^2)^2 \sqrt{c(a + bx^2)^3} \\ &+ \frac{1}{10} cx (a + bx^2)^3 \sqrt{c(a + bx^2)^3} + \frac{63a^{7/2} c \sqrt{c(a + bx^2)^3} \operatorname{arcsinh}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{256\sqrt{b} \left(1 + \frac{bx^2}{a}\right)^{3/2}} \end{aligned}$$

output

```
21/128*a^3*c*x*(c*(b*x^2+a)^3)^(1/2)+63*a^4*c*x*(c*(b*x^2+a)^3)^(1/2)/(256
*b*x^2+256*a)+21/160*a^2*c*x*(b*x^2+a)*(c*(b*x^2+a)^3)^(1/2)+9/80*a*c*x*(b
*x^2+a)^2*(c*(b*x^2+a)^3)^(1/2)+1/10*c*x*(b*x^2+a)^3*(c*(b*x^2+a)^3)^(1/2)
+63/256*a^(7/2)*c*(c*(b*x^2+a)^3)^(1/2)*arcsinh(b^(1/2)*x/a^(1/2))/b^(1/2)
/(1+b*x^2/a)^(3/2)
```

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.59

$$\int (c(a + bx^2)^3)^{3/2} dx = \frac{(c(a + bx^2)^3)^{3/2} (\sqrt{bx}\sqrt{a + bx^2}(965a^4 + 1490a^3bx^2 + 1368a^2b^2x^4 + 656ab^3x^6 + 128b^4x^8) - 315a^5 \operatorname{Log}[-(\operatorname{Sqrt}[b]x) + \operatorname{Sqrt}[a + bx^2]])}{1280\sqrt{b}(a + bx^2)^{9/2}}$$

input `Integrate[(c*(a + b*x^2)^3)^(3/2), x]`

output `((c*(a + b*x^2)^3)^(3/2)*(Sqrt[b]*x*Sqrt[a + b*x^2]*(965*a^4 + 1490*a^3*b*x^2 + 1368*a^2*b^2*x^4 + 656*a*b^3*x^6 + 128*b^4*x^8) - 315*a^5*Log[-(Sqrt[b]*x) + Sqrt[a + b*x^2]]))/(1280*Sqrt[b]*(a + b*x^2)^(9/2))`

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.86, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {2045, 211, 211, 211, 211, 211, 222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (c(a + bx^2)^3)^{3/2} dx \\ & \quad \downarrow \text{2045} \\ & \frac{a^3 c \sqrt{c(a + bx^2)^3} \int \left(\frac{bx^2}{a} + 1\right)^{9/2} dx}{\left(\frac{bx^2}{a} + 1\right)^{3/2}} \\ & \quad \downarrow \text{211} \\ & \frac{a^3 c \sqrt{c(a + bx^2)^3} \left(\frac{9}{10} \int \left(\frac{bx^2}{a} + 1\right)^{7/2} dx + \frac{1}{10} x \left(\frac{bx^2}{a} + 1\right)^{9/2}\right)}{\left(\frac{bx^2}{a} + 1\right)^{3/2}} \end{aligned}$$

$$\begin{aligned} & \downarrow 211 \\ & \frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\frac{9}{10} \left(\frac{7}{8} \int \left(\frac{bx^2}{a} + 1 \right)^{5/2} dx + \frac{1}{8} x \left(\frac{bx^2}{a} + 1 \right)^{7/2} \right) + \frac{1}{10} x \left(\frac{bx^2}{a} + 1 \right)^{9/2} \right)}{\left(\frac{bx^2}{a} + 1 \right)^{3/2}} \\ & \downarrow 211 \\ & \frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\frac{9}{10} \left(\frac{7}{8} \left(\frac{5}{6} \int \left(\frac{bx^2}{a} + 1 \right)^{3/2} dx + \frac{1}{6} x \left(\frac{bx^2}{a} + 1 \right)^{5/2} \right) + \frac{1}{8} x \left(\frac{bx^2}{a} + 1 \right)^{7/2} \right) + \frac{1}{10} x \left(\frac{bx^2}{a} + 1 \right)^{9/2} \right)}{\left(\frac{bx^2}{a} + 1 \right)^{3/2}} \\ & \downarrow 211 \\ & \frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\frac{9}{10} \left(\frac{7}{8} \left(\frac{5}{6} \left(\frac{3}{4} \int \sqrt{\frac{bx^2}{a} + 1} dx + \frac{1}{4} x \left(\frac{bx^2}{a} + 1 \right)^{3/2} \right) + \frac{1}{6} x \left(\frac{bx^2}{a} + 1 \right)^{5/2} \right) + \frac{1}{8} x \left(\frac{bx^2}{a} + 1 \right)^{7/2} \right) + \frac{1}{10} x \left(\frac{bx^2}{a} + 1 \right)^{9/2} \right)}{\left(\frac{bx^2}{a} + 1 \right)^{3/2}} \\ & \downarrow 211 \\ & \frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\frac{9}{10} \left(\frac{7}{8} \left(\frac{5}{6} \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{1}{\sqrt{\frac{bx^2}{a} + 1}} dx + \frac{1}{2} x \sqrt{\frac{bx^2}{a} + 1} \right) + \frac{1}{4} x \left(\frac{bx^2}{a} + 1 \right)^{3/2} \right) + \frac{1}{6} x \left(\frac{bx^2}{a} + 1 \right)^{5/2} \right) + \frac{1}{8} x \left(\frac{bx^2}{a} + 1 \right)^{7/2} \right) \right)}{\left(\frac{bx^2}{a} + 1 \right)^{3/2}} \\ & \downarrow 222 \\ & \frac{a^3 c \left(\frac{9}{10} \left(\frac{7}{8} \left(\frac{5}{6} \left(\frac{3}{4} \left(\frac{\sqrt{a} \operatorname{arcsinh} \left(\frac{\sqrt{bx}}{\sqrt{a}} \right)}{2\sqrt{b}} + \frac{1}{2} x \sqrt{\frac{bx^2}{a} + 1} \right) + \frac{1}{4} x \left(\frac{bx^2}{a} + 1 \right)^{3/2} \right) + \frac{1}{6} x \left(\frac{bx^2}{a} + 1 \right)^{5/2} \right) + \frac{1}{8} x \left(\frac{bx^2}{a} + 1 \right)^{7/2} \right) \right)}{\left(\frac{bx^2}{a} + 1 \right)^{3/2}} \end{aligned}$$

input `Int[(c*(a + b*x^2)^3)^(3/2),x]`

output `(a^3*c*Sqrt[c*(a + b*x^2)^3]*((x*(1 + (b*x^2)/a)^(9/2))/10 + (9*((x*(1 + (b*x^2)/a)^(7/2))/8 + (7*((x*(1 + (b*x^2)/a)^(5/2))/6 + (5*((x*(1 + (b*x^2)/a)^(3/2))/4 + (3*((x*Sqrt[1 + (b*x^2)/a])/2 + (Sqrt[a]*ArcSinh[(Sqrt[b]*x)/Sqrt[a]])/(2*Sqrt[b])))/4))/6))/8))/10)/(1 + (b*x^2)/a)^(3/2)`

Definitions of rubi rules used

rule 211

$$\text{Int}[(a_+) + (b_+)(x_+)^2]^{(p_+)}, x_Symbol] \rightarrow \text{Simp}[x*((a + b*x^2)^p/(2*p + 1)), x] + \text{Simp}[2*a*(p/(2*p + 1)) \text{Int}[(a + b*x^2)^{(p - 1)}, x], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{GtQ}[p, 0] \&\& (\text{IntegerQ}[4*p] \parallel \text{IntegerQ}[6*p])$$

rule 222

$$\text{Int}[1/\text{Sqrt}[(a_+) + (b_+)(x_+)^2], x_Symbol] \rightarrow \text{Simp}[\text{ArcSinh}[\text{Rt}[b, 2]*(x/\text{Sqrt}[a])]/\text{Rt}[b, 2], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{GtQ}[a, 0] \&\& \text{PosQ}[b]$$

rule 2045

$$\text{Int}[(u_+)((c_+)((a_+) + (b_+)(x_+)^{n_+})^{(q_+)})^{(p_+)}, x_Symbol] \rightarrow \text{Simp}[\text{Simp}[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^{(p*q)}] \text{Int}[u*(1 + b*(x^n/a))^{(p*q)}, x], x] /; \text{FreeQ}[\{a, b, c, n, p, q\}, x] \&\& !\text{GeQ}[a, 0]$$

Maple [A] (verified)

Time = 1.62 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.67

method	result
risch	$\frac{x(128b^4x^8+656ab^3x^6+1368a^2b^2x^4+1490a^3bx^2+965a^4)c\sqrt{c(bx^2+a)^3}}{1280bx^2+1280a} + \frac{63a^5 \ln\left(\frac{bcx}{\sqrt{bc}} + \sqrt{x^2bc+ac}\right)c\sqrt{c(bx^2+a)^3}\sqrt{(bx^2+a)c}}{256\sqrt{bc}(bx^2+a)^2}$
default	$\frac{(c(bx^2+a)^3)^{\frac{3}{2}} \left(128x^5(x^2bc+ac)^{\frac{5}{2}}b^2\sqrt{bc}+400(x^2bc+ac)^{\frac{5}{2}}\sqrt{bc}abx^3+440(x^2bc+ac)^{\frac{5}{2}}\sqrt{bc}a^2x+210(x^2bc+ac)^{\frac{3}{2}}\sqrt{bc}a^3cx+315\sqrt{bc}a^4\right)}{1280(bx^2+a)^3((bx^2+a)c)^{\frac{3}{2}}c\sqrt{bc}}$

input

$$\text{int}((c*(b*x^2+a)^3)^{(3/2)}, x, \text{method}=_RETURNVERBOSE)$$

output

$$\frac{1}{1280}x*(128*b^4*x^8+656*a*b^3*x^6+1368*a^2*b^2*x^4+1490*a^3*b*x^2+965*a^4)/(b*x^2+a)*c*(c*(b*x^2+a)^3)^{(1/2)}+63/256*a^5*\ln(b*c*x/(b*c)^{(1/2)}+(b*c*x^2+a*c)^{(1/2)})/(b*c)^{(1/2)}*c/(b*x^2+a)^2*(c*(b*x^2+a)^3)^{(1/2)}*((b*x^2+a)*c)^{(1/2)}$$

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 402, normalized size of antiderivative = 1.94

$$\int (c(a + bx^2)^3)^{3/2} dx = \frac{315(a^5bcx^2 + a^6c)\sqrt{\frac{c}{b}} \log\left(-\frac{2b^2cx^4 + 3abcx^2 + a^2c + 2\sqrt{b^3cx^6 + 3ab^2cx^4 + 3a^2bcx^2 + a^3cbx}\sqrt{\frac{c}{b}}}{bx^2 + a}\right) + 2(128b^4cx^9 + 656ab^3cx^7 + 1368a^2b^2cx^5 + 1490a^3b^2cx^3 + 965a^4cx)\sqrt{b^3cx^6 + 3ab^2cx^4 + 3a^2bcx^2 + a^3cbx}\sqrt{\frac{c}{b}}}{1280(bx^2 + a)} - \frac{315(a^5bcx^2 + a^6c)\sqrt{-\frac{c}{b}} \arctan\left(\frac{\sqrt{b^3cx^6 + 3ab^2cx^4 + 3a^2bcx^2 + a^3cbx}\sqrt{-\frac{c}{b}}}{b^2cx^4 + 2abcx^2 + a^2c}\right) - (128b^4cx^9 + 656ab^3cx^7 + 1368a^2b^2cx^5 + 1490a^3b^2cx^3 + 965a^4cx)\sqrt{b^3cx^6 + 3ab^2cx^4 + 3a^2bcx^2 + a^3cbx}\sqrt{-\frac{c}{b}}}{1280(bx^2 + a)}$$

input `integrate((c*(b*x^2+a)^3)^(3/2),x, algorithm="fricas")`output `[1/2560*(315*(a^5*b*c*x^2 + a^6*c)*sqrt(c/b)*log(-(2*b^2*c*x^4 + 3*a*b*c*x^2 + a^2*c + 2*sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c))*b*x*sqrt(c/b))/(b*x^2 + a)) + 2*(128*b^4*c*x^9 + 656*a*b^3*c*x^7 + 1368*a^2*b^2*c*x^5 + 1490*a^3*b^2*c*x^3 + 965*a^4*c*x)*sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c))/(b*x^2 + a), -1/1280*(315*(a^5*b*c*x^2 + a^6*c)*sqrt(-c/b)*arctan(sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c))*b*x*sqrt(-c/b)/(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)) - (128*b^4*c*x^9 + 656*a*b^3*c*x^7 + 1368*a^2*b^2*c*x^5 + 1490*a^3*b^2*c*x^3 + 965*a^4*c*x)*sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c))/(b*x^2 + a)]`**Sympy [F]**

$$\int (c(a + bx^2)^3)^{3/2} dx = \int (c(a + bx^2)^3)^{\frac{3}{2}} dx$$

input `integrate((c*(b*x**2+a)**3)**(3/2),x)`output `Integral((c*(a + b*x**2)**3)**(3/2), x)`

Maxima [F]

$$\int \left(c(a + bx^2)^3 \right)^{3/2} dx = \int \left((bx^2 + a)^3 c \right)^{3/2} dx$$

input `integrate((c*(b*x^2+a)^3)^(3/2),x, algorithm="maxima")`

output `integrate(((b*x^2 + a)^3*c)^(3/2), x)`

Giac [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.74

$$\int \left(c(a + bx^2)^3 \right)^{3/2} dx =$$

$$-\frac{1}{1280} \left(\frac{315 a^5 c \log \left(\left| -\sqrt{bc} x + \sqrt{bcx^2 + ac} \right| \right) \operatorname{sgn}(bx^2 + a)}{\sqrt{bc}} - (965 a^4 \operatorname{sgn}(bx^2 + a) + 2 (745 a^3 b \operatorname{sgn}(bx^2 + a) + 4 (171 a^2 b^2 \operatorname{sgn}(bx^2 + a) + 2 (8 b^4 x^2 \operatorname{sgn}(bx^2 + a) + 41 a b^3 \operatorname{sgn}(bx^2 + a)) x^2) x^2) x^2) \sqrt{bc} x^2 + a c) x \right) c$$

input `integrate((c*(b*x^2+a)^3)^(3/2),x, algorithm="giac")`

output `-1/1280*(315*a^5*c*log(abs(-sqrt(b*c)*x + sqrt(b*c*x^2 + a*c)))*sgn(b*x^2 + a)/sqrt(b*c) - (965*a^4*sgn(b*x^2 + a) + 2*(745*a^3*b*sgn(b*x^2 + a) + 4*(171*a^2*b^2*sgn(b*x^2 + a) + 2*(8*b^4*x^2*sgn(b*x^2 + a) + 41*a*b^3*sgn(b*x^2 + a))*x^2)*x^2)*sqrt(b*c*x^2 + a*c)*x)*c`

Mupad [F(-1)]

Timed out.

$$\int \left(c(a + bx^2)^3 \right)^{3/2} dx = \int \left(c(bx^2 + a)^3 \right)^{3/2} dx$$

input `int((c*(a + b*x^2)^3)^(3/2),x)`

output `int((c*(a + b*x^2)^3)^(3/2), x)`

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.58

$$\int (c(a + bx^2)^3)^{3/2} dx = \frac{\sqrt{c}c(965\sqrt{bx^2+a}a^4bx + 1490\sqrt{bx^2+a}a^3b^2x^3 + 1368\sqrt{bx^2+a}a^2b^3x^5 + 656\sqrt{bx^2+a}ab^4x^7 + 128\sqrt{bx^2+a}b^5x^9 + 315\sqrt{b}\log((\sqrt{a+bx^2}) + \sqrt{b}x/\sqrt{a})a^{5/2})}{1280b}$$

input `int((c*(b*x^2+a)^3)^(3/2),x)`

output `(sqrt(c)*c*(965*sqrt(a + b*x**2)*a**4*b*x + 1490*sqrt(a + b*x**2)*a**3*b**2*x**3 + 1368*sqrt(a + b*x**2)*a**2*b**3*x**5 + 656*sqrt(a + b*x**2)*a*b**4*x**7 + 128*sqrt(a + b*x**2)*b**5*x**9 + 315*sqrt(b)*log((sqrt(a + b*x**2) + sqrt(b)*x)/sqrt(a))*a**5))/(1280*b)`

$$3.18 \quad \int \frac{(c(a+bx^2)^3)^{3/2}}{x^2} dx$$

Optimal result	165
Mathematica [A] (verified)	166
Rubi [A] (verified)	166
Maple [A] (verified)	169
Fricas [A] (verification not implemented)	169
Sympy [F]	170
Maxima [F]	170
Giac [A] (verification not implemented)	171
Mupad [F(-1)]	171
Reduce [B] (verification not implemented)	172

Optimal result

Integrand size = 19, antiderivative size = 234

$$\begin{aligned} \int \frac{(c(a+bx^2)^3)^{3/2}}{x^2} dx &= -\frac{a^4 c \sqrt{c(a+bx^2)^3}}{x(a+bx^2)} + \frac{325a^3 b c x \sqrt{c(a+bx^2)^3}}{128(a+bx^2)} \\ &+ \frac{105a^2 b^2 c x^3 \sqrt{c(a+bx^2)^3}}{64(a+bx^2)} + \frac{11ab^3 c x^5 \sqrt{c(a+bx^2)^3}}{16(a+bx^2)} \\ &+ \frac{b^4 c x^7 \sqrt{c(a+bx^2)^3}}{8(a+bx^2)} + \frac{315a^{5/2} \sqrt{bc} \sqrt{c(a+bx^2)^3} \operatorname{arcsinh}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{128\left(1+\frac{bx^2}{a}\right)^{3/2}} \end{aligned}$$

output

```
-a^4*c*(c*(b*x^2+a)^3)^(1/2)/x/(b*x^2+a)+325*a^3*b*c*x*(c*(b*x^2+a)^3)^(1/2)/(128*b*x^2+128*a)+105*a^2*b^2*c*x^3*(c*(b*x^2+a)^3)^(1/2)/(64*b*x^2+64*a)+11*a*b^3*c*x^5*(c*(b*x^2+a)^3)^(1/2)/(16*b*x^2+16*a)+b^4*c*x^7*(c*(b*x^2+a)^3)^(1/2)/(8*b*x^2+8*a)+315/128*a^(5/2)*b^(1/2)*c*(c*(b*x^2+a)^3)^(1/2)*arcsinh(b^(1/2)*x/a^(1/2))/(1+b*x^2/a)^(3/2)
```

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.52

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^2} dx = \frac{(c(a + bx^2)^3)^{3/2} \left(\sqrt{a + bx^2} (128a^4 - 325a^3bx^2 - 210a^2b^2x^4 - 88ab^3x^6 - 16b^4x^8) + 315a^4\sqrt{bx} \log(-\sqrt{bx} + \sqrt{a + bx^2}) \right)}{128x(a + bx^2)^{9/2}}$$

input `Integrate[(c*(a + b*x^2)^3)^(3/2)/x^2,x]`

output `-1/128*((c*(a + b*x^2)^3)^(3/2)*(Sqrt[a + b*x^2]*(128*a^4 - 325*a^3*b*x^2 - 210*a^2*b^2*x^4 - 88*a*b^3*x^6 - 16*b^4*x^8) + 315*a^4*Sqrt[b]*x*Log[-(Sqrt[b]*x) + Sqrt[a + b*x^2]]))/(x*(a + b*x^2)^(9/2))`

Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 179, normalized size of antiderivative = 0.76, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {2045, 247, 211, 211, 211, 211, 222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^2} dx$$

↓ 2045

$$\frac{a^3c\sqrt{c(a + bx^2)^3} \int \frac{\left(\frac{bx^2}{a} + 1\right)^{9/2}}{x^2} dx}{\left(\frac{bx^2}{a} + 1\right)^{3/2}}$$

↓ 247

$$\frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\frac{9b \int \left(\frac{bx^2}{a} + 1\right)^{7/2} dx}{a} - \frac{\left(\frac{bx^2}{a} + 1\right)^{9/2}}{x} \right)}{\left(\frac{bx^2}{a} + 1\right)^{3/2}}$$

↓ 211

$$\frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\frac{9b \left(\frac{7}{8} \int \left(\frac{bx^2}{a} + 1\right)^{5/2} dx + \frac{1}{8} x \left(\frac{bx^2}{a} + 1\right)^{7/2} \right)}{a} - \frac{\left(\frac{bx^2}{a} + 1\right)^{9/2}}{x} \right)}{\left(\frac{bx^2}{a} + 1\right)^{3/2}}$$

↓ 211

$$\frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\frac{9b \left(\frac{7}{8} \left(\frac{5}{6} \int \left(\frac{bx^2}{a} + 1\right)^{3/2} dx + \frac{1}{6} x \left(\frac{bx^2}{a} + 1\right)^{5/2} \right) + \frac{1}{8} x \left(\frac{bx^2}{a} + 1\right)^{7/2} \right)}{a} - \frac{\left(\frac{bx^2}{a} + 1\right)^{9/2}}{x} \right)}{\left(\frac{bx^2}{a} + 1\right)^{3/2}}$$

↓ 211

$$\frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\frac{9b \left(\frac{7}{8} \left(\frac{5}{6} \left(\frac{3}{4} \int \sqrt{\frac{bx^2}{a} + 1} dx + \frac{1}{4} x \left(\frac{bx^2}{a} + 1\right)^{3/2} \right) + \frac{1}{6} x \left(\frac{bx^2}{a} + 1\right)^{5/2} \right) + \frac{1}{8} x \left(\frac{bx^2}{a} + 1\right)^{7/2} \right)}{a} - \frac{\left(\frac{bx^2}{a} + 1\right)^{9/2}}{x} \right)}{\left(\frac{bx^2}{a} + 1\right)^{3/2}}$$

↓ 211

$$\frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\frac{9b \left(\frac{7}{8} \left(\frac{5}{6} \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{1}{\sqrt{\frac{bx^2}{a} + 1}} dx + \frac{1}{2} x \sqrt{\frac{bx^2}{a} + 1} \right) + \frac{1}{4} x \left(\frac{bx^2}{a} + 1\right)^{3/2} \right) + \frac{1}{6} x \left(\frac{bx^2}{a} + 1\right)^{5/2} \right) + \frac{1}{8} x \left(\frac{bx^2}{a} + 1\right)^{7/2} \right)}{a} - \frac{\left(\frac{bx^2}{a} + 1\right)^{9/2}}{x} \right)}{\left(\frac{bx^2}{a} + 1\right)^{3/2}}$$

↓ 222

$$\frac{a^3 c \left(\frac{9b \left(\frac{7}{8} \left(\frac{5}{6} \left(\frac{3}{4} \left(\frac{\sqrt{a} \operatorname{arcsinh} \left(\frac{\sqrt{bx^2}}{\sqrt{a}} \right)}{2\sqrt{b}} + \frac{1}{2} x \sqrt{\frac{bx^2}{a} + 1} \right) + \frac{1}{4} x \left(\frac{bx^2}{a} + 1\right)^{3/2} \right) + \frac{1}{6} x \left(\frac{bx^2}{a} + 1\right)^{5/2} \right) + \frac{1}{8} x \left(\frac{bx^2}{a} + 1\right)^{7/2} \right)}{a} - \frac{\left(\frac{bx^2}{a} + 1\right)^{9/2}}{x} \right) \sqrt{c}}{\left(\frac{bx^2}{a} + 1\right)^{3/2}}$$

input `Int[(c*(a + b*x^2)^3)^(3/2)/x^2,x]`

output `(a^3*c*Sqrt[c*(a + b*x^2)^3]*(-((1 + (b*x^2)/a)^(9/2)/x) + (9*b*((x*(1 + (b*x^2)/a)^(7/2))/8 + (7*((x*(1 + (b*x^2)/a)^(5/2))/6 + (5*((x*(1 + (b*x^2)/a)^(3/2))/4 + (3*((x*Sqrt[1 + (b*x^2)/a])/2 + (Sqrt[a]*ArcSinh[(Sqrt[b]*x)/Sqrt[a]])/(2*Sqrt[b])))/4))/6))/8))/a)/(1 + (b*x^2)/a)^(3/2)`

Defintions of rubi rules used

rule 211 `Int[((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[x*((a + b*x^2)^p/(2*p + 1)), x] + Simp[2*a*(p/(2*p + 1)) Int[(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 222 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

rule 247 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^p/(c*(m + 1))), x] - Simp[2*b*(p/(c^2*(m + 1))) Int[(c*x)^(m + 2)*(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && GtQ[p, 0] && LtQ[m, -1] && !ILtQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2045 `Int[(u_)*((c_)*((a_) + (b_)*(x_)^(n_))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 2.41 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.60

method	result
risch	$-\frac{(-16b^4x^8-88ab^3x^6-210a^2b^2x^4-325a^3bx^2+128a^4)c\sqrt{c(bx^2+a)^3}}{128(bx^2+a)x} + \frac{315a^4b\ln\left(\frac{bcx}{\sqrt{bc}}+\sqrt{x^2bc+ac}\right)c\sqrt{c(bx^2+a)^3}\sqrt{(bx^2+a)}}{128\sqrt{bc}(bx^2+a)^2}$
default	$-\frac{(c(bx^2+a)^3)^{\frac{3}{2}}\left(-16(x^2bc+ac)^{\frac{5}{2}}\sqrt{bc}b^2x^4-56(x^2bc+ac)^{\frac{5}{2}}\sqrt{bc}abx^2-210(x^2bc+ac)^{\frac{3}{2}}\sqrt{bc}a^2bcx^2-315\sqrt{x^2bc+ac}\sqrt{bc}a^3bc^2x\right)}{128(bx^2+a)^3((bx^2+a)c)^{\frac{3}{2}}cx\sqrt{bc}}$

input `int((c*(b*x^2+a)^3)^(3/2)/x^2,x,method=_RETURNVERBOSE)`

output `-1/128/(b*x^2+a)*(-16*b^4*x^8-88*a*b^3*x^6-210*a^2*b^2*x^4-325*a^3*b*x^2+128*a^4)/x*c*(c*(b*x^2+a)^3)^(1/2)+315/128*a^4*b*ln(b*c*x/(b*c)^(1/2)+(b*c*x^2+a*c)^(1/2))/(b*c)^(1/2)*c/(b*x^2+a)^2*(c*(b*x^2+a)^3)^(1/2)*((b*x^2+a)*c)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 396, normalized size of antiderivative = 1.69

$$\int \frac{(c(a+bx^2)^3)^{3/2}}{x^2} dx = \frac{315(a^4bcx^3+a^5cx)\sqrt{bc}\log\left(\frac{-2b^2cx^4+3abcx^2+a^2c+2\sqrt{b^3cx^6+3ab^2cx^4+3a^2bcx^2+a^3c}\sqrt{bcx}}{bx^2+a}\right) - 315(a^4bcx^3+a^5cx)\sqrt{-bc}\arctan\left(\frac{\sqrt{b^3cx^6+3ab^2cx^4+3a^2bcx^2+a^3c}\sqrt{-bcx}}{b^2cx^4+2abcx^2+a^2c}\right) - (16b^4cx^8+88ab^3cx^6+210a^2b^2cx^4)}{128(bx^3+ax)}$$

input `integrate((c*(b*x^2+a)^3)^(3/2)/x^2,x, algorithm="fricas")`

output

```
[1/256*(315*(a^4*b*c*x^3 + a^5*c*x)*sqrt(b*c)*log(-(2*b^2*c*x^4 + 3*a*b*c*x^2 + a^2*c + 2*sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c))*sqrt(b*c)*x)/(b*x^2 + a)) + 2*(16*b^4*c*x^8 + 88*a*b^3*c*x^6 + 210*a^2*b^2*c*x^4 + 325*a^3*b*c*x^2 - 128*a^4*c)*sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c))/(b*x^3 + a*x), -1/128*(315*(a^4*b*c*x^3 + a^5*c*x)*sqrt(-b*c)*arctan(sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c))*sqrt(-b*c)*x/(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)) - (16*b^4*c*x^8 + 88*a*b^3*c*x^6 + 210*a^2*b^2*c*x^4 + 325*a^3*b*c*x^2 - 128*a^4*c)*sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c))/(b*x^3 + a*x)]
```

Sympy [F]

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^2} dx = \int \frac{(c(a + bx^2)^3)^{\frac{3}{2}}}{x^2} dx$$

input

```
integrate((c*(b*x**2+a)**3)**(3/2)/x**2,x)
```

output

```
Integral((c*(a + b*x**2)**3)**(3/2)/x**2, x)
```

Maxima [F]

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^2} dx = \int \frac{((bx^2 + a)^3 c)^{\frac{3}{2}}}{x^2} dx$$

input

```
integrate((c*(b*x^2+a)^3)^(3/2)/x^2,x, algorithm="maxima")
```

output

```
integrate(((b*x^2 + a)^3*c)^(3/2)/x^2, x)
```

Giac [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 185, normalized size of antiderivative = 0.79

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^2} dx = \frac{1}{256} \left(\frac{512 \sqrt{bca^5} \operatorname{sgn}(bx^2 + a)}{(\sqrt{bcx} - \sqrt{bcx^2 + ac})^2 - ac} - 315 \sqrt{bca^4} \log \left((\sqrt{bcx} - \sqrt{bcx^2 + ac})^2 \right) \right)$$

input `integrate((c*(b*x^2+a)^3)^(3/2)/x^2,x, algorithm="giac")`

output `1/256*(512*sqrt(b*c)*a^5*c*sgn(b*x^2 + a)/((sqrt(b*c)*x - sqrt(b*c*x^2 + a*c))^2 - a*c) - 315*sqrt(b*c)*a^4*log((sqrt(b*c)*x - sqrt(b*c*x^2 + a*c))^2)*sgn(b*x^2 + a) + 2*(325*a^3*b*sgn(b*x^2 + a) + 2*(105*a^2*b^2*sgn(b*x^2 + a) + 4*(2*b^4*x^2*sgn(b*x^2 + a) + 11*a*b^3*sgn(b*x^2 + a))*x^2)*x^2)*sqrt(b*c*x^2 + a*c)*x)*c`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^2} dx = \int \frac{(c(bx^2 + a)^3)^{3/2}}{x^2} dx$$

input `int((c*(a + b*x^2)^3)^(3/2)/x^2,x)`

output `int((c*(a + b*x^2)^3)^(3/2)/x^2, x)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.54

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^2} dx = \frac{\sqrt{c}c(-128\sqrt{bx^2 + a}a^4 + 325\sqrt{bx^2 + a}a^3bx^2 + 210\sqrt{bx^2 + a}a^2b^2x^4 + 88\sqrt{bx^2 + a}ab^3x^6 + 16\sqrt{bx^2 + a}b^4x^8 + 315\sqrt{b}\log((\sqrt{a + bx^2}) + \sqrt{(b)x}/\sqrt{a})a^4x - 189\sqrt{b}a^4x)/(128x)}{128}$$

input `int((c*(b*x^2+a)^3)^(3/2)/x^2,x)`output `(sqrt(c)*c*(- 128*sqrt(a + b*x**2)*a**4 + 325*sqrt(a + b*x**2)*a**3*b*x**2 + 210*sqrt(a + b*x**2)*a**2*b**2*x**4 + 88*sqrt(a + b*x**2)*a*b**3*x**6 + 16*sqrt(a + b*x**2)*b**4*x**8 + 315*sqrt(b)*log((sqrt(a + b*x**2) + sqrt(b)*x)/sqrt(a))*a**4*x - 189*sqrt(b)*a**4*x)/(128*x)`

$$3.19 \quad \int \frac{(c(a+bx^2)^3)^{3/2}}{x^4} dx$$

Optimal result	173
Mathematica [A] (verified)	174
Rubi [A] (verified)	174
Maple [A] (verified)	177
Fricas [A] (verification not implemented)	178
Sympy [F]	178
Maxima [F]	179
Giac [A] (verification not implemented)	179
Mupad [F(-1)]	180
Reduce [B] (verification not implemented)	180

Optimal result

Integrand size = 19, antiderivative size = 236

$$\begin{aligned} \int \frac{(c(a+bx^2)^3)^{3/2}}{x^4} dx = & -\frac{a^4 c \sqrt{c(a+bx^2)^3}}{3x^3(a+bx^2)} - \frac{13a^3 b c \sqrt{c(a+bx^2)^3}}{3x(a+bx^2)} \\ & + \frac{55a^2 b^2 c x \sqrt{c(a+bx^2)^3}}{16(a+bx^2)} + \frac{25ab^3 c x^3 \sqrt{c(a+bx^2)^3}}{24(a+bx^2)} \\ & + \frac{b^4 c x^5 \sqrt{c(a+bx^2)^3}}{6(a+bx^2)} + \frac{105a^{3/2} b^{3/2} c \sqrt{c(a+bx^2)^3} \operatorname{arcsinh}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{16\left(1+\frac{bx^2}{a}\right)^{3/2}} \end{aligned}$$

output

```
-1/3*a^4*c*(c*(b*x^2+a)^3)^(1/2)/x^3/(b*x^2+a)-13/3*a^3*b*c*(c*(b*x^2+a)^3)^(1/2)/x/(b*x^2+a)+55*a^2*b^2*c*x*(c*(b*x^2+a)^3)^(1/2)/(16*b*x^2+16*a)+25*a*b^3*c*x^3*(c*(b*x^2+a)^3)^(1/2)/(24*b*x^2+24*a)+b^4*c*x^5*(c*(b*x^2+a)^3)^(1/2)/(6*b*x^2+6*a)+105/16*a^(3/2)*b^(3/2)*c*(c*(b*x^2+a)^3)^(1/2)*arc sinh(b^(1/2)*x/a^(1/2))/(1+b*x^2/a)^(3/2)
```

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.52

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^4} dx = \frac{(c(a + bx^2)^3)^{3/2} \left(\sqrt{a + bx^2} (16a^4 + 208a^3bx^2 - 165a^2b^2x^4 - 50ab^3x^6 - 8b^4x^8) + 315a^3b^{3/2}x^3 \log(-\sqrt{bx} + \sqrt{a + bx^2}) \right)}{48x^3 (a + bx^2)^{9/2}}$$

input `Integrate[(c*(a + b*x^2)^3)^(3/2)/x^4,x]`

output `-1/48*((c*(a + b*x^2)^3)^(3/2)*(Sqrt[a + b*x^2]*(16*a^4 + 208*a^3*b*x^2 - 165*a^2*b^2*x^4 - 50*a*b^3*x^6 - 8*b^4*x^8) + 315*a^3*b^(3/2)*x^3*Log[-(Sqrt[b]*x) + Sqrt[a + b*x^2]]))/(x^3*(a + b*x^2)^(9/2))`

Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 183, normalized size of antiderivative = 0.78, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {2045, 247, 247, 211, 211, 211, 222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^4} dx$$

↓ 2045

$$\frac{a^3 c \sqrt{c(a + bx^2)^3} \int \frac{\left(\frac{bx^2}{a} + 1\right)^{9/2}}{x^4} dx}{\left(\frac{bx^2}{a} + 1\right)^{3/2}}$$

↓ 247

$$\frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\frac{3b \int \frac{\left(\frac{bx^2}{a} + 1\right)^{7/2}}{x^2} dx}{a} - \frac{\left(\frac{bx^2}{a} + 1\right)^{9/2}}{3x^3} \right)}{\left(\frac{bx^2}{a} + 1\right)^{3/2}}$$

↓ 247

$$\frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\frac{3b \left(\frac{7b \int \left(\frac{bx^2}{a} + 1\right)^{5/2} dx}{a} - \frac{\left(\frac{bx^2}{a} + 1\right)^{7/2}}{x} \right)}{a} - \frac{\left(\frac{bx^2}{a} + 1\right)^{9/2}}{3x^3} \right)}{\left(\frac{bx^2}{a} + 1\right)^{3/2}}$$

↓ 211

$$\frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\frac{3b \left(\frac{7b \left(\frac{5}{6} \int \left(\frac{bx^2}{a} + 1\right)^{3/2} dx + \frac{1}{6} x \left(\frac{bx^2}{a} + 1\right)^{5/2} \right)}{a} - \frac{\left(\frac{bx^2}{a} + 1\right)^{7/2}}{x} \right)}{a} - \frac{\left(\frac{bx^2}{a} + 1\right)^{9/2}}{3x^3} \right)}{\left(\frac{bx^2}{a} + 1\right)^{3/2}}$$

↓ 211

$$\frac{a^3 c \sqrt{c(a+bx^2)^3} \left(\frac{3b \left(\frac{7b \left(\frac{5}{6} \left(\frac{3}{4} \int \sqrt{\frac{bx^2}{a} + 1} dx + \frac{1}{4} x \left(\frac{bx^2}{a} + 1\right)^{3/2} \right) + \frac{1}{6} x \left(\frac{bx^2}{a} + 1\right)^{5/2} \right)}{a} - \frac{\left(\frac{bx^2}{a} + 1\right)^{7/2}}{x} \right)}{a} - \frac{\left(\frac{bx^2}{a} + 1\right)^{9/2}}{3x^3} \right)}{\left(\frac{bx^2}{a} + 1\right)^{3/2}}$$

↓ 211

$$a^3 c \sqrt{c(a + bx^2)^3} \left(\frac{3b \left(\frac{7b \left(\frac{5}{6} \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{1}{\sqrt{\frac{bx^2}{a} + 1}} dx + \frac{1}{2} x \sqrt{\frac{bx^2}{a} + 1} \right) + \frac{1}{4} x \left(\frac{bx^2}{a} + 1 \right)^{3/2} \right) + \frac{1}{6} x \left(\frac{bx^2}{a} + 1 \right)^{5/2} \right) \right) \left(\frac{bx^2}{a} + 1 \right)^{7/2}}{a} - \frac{\left(\frac{bx^2}{a} + 1 \right)^{9/2}}{3x^3} \right)}{a} - \frac{\left(\frac{bx^2}{a} + 1 \right)^{9/2}}{3x^3} \right)$$

$\left(\frac{bx^2}{a} + 1 \right)^{3/2}$
 \downarrow 222

$$a^3 c \left(\frac{3b \left(\frac{7b \left(\frac{5}{6} \left(\frac{3}{4} \left(\frac{\sqrt{a} \operatorname{arcsinh} \left(\frac{\sqrt{bx}}{\sqrt{a}} \right)}{2\sqrt{b}} + \frac{1}{2} x \sqrt{\frac{bx^2}{a} + 1} \right) + \frac{1}{4} x \left(\frac{bx^2}{a} + 1 \right)^{3/2} \right) + \frac{1}{6} x \left(\frac{bx^2}{a} + 1 \right)^{5/2} \right) \right) \left(\frac{bx^2}{a} + 1 \right)^{7/2}}{a} - \frac{\left(\frac{bx^2}{a} + 1 \right)^{9/2}}{3x^3} \right) \sqrt{c(a + bx^2)^3}}{a} - \frac{\left(\frac{bx^2}{a} + 1 \right)^{9/2}}{3x^3} \right) \sqrt{c(a + bx^2)^3}$$

input `Int[(c*(a + b*x^2)^3)^(3/2)/x^4,x]`

output `(a^3*c*Sqrt[c*(a + b*x^2)^3]*(-1/3*(1 + (b*x^2)/a)^(9/2)/x^3 + (3*b*(-((1 + (b*x^2)/a)^(7/2)/x) + (7*b*((x*(1 + (b*x^2)/a)^(5/2))/6 + (5*((x*(1 + (b*x^2)/a)^(3/2))/4 + (3*((x*Sqrt[1 + (b*x^2)/a]))/2 + (Sqrt[a]*ArcSinh[(Sqrt[b]*x)/Sqrt[a]])/(2*Sqrt[b])))/4))/6)/a))/a)/(1 + (b*x^2)/a)^(3/2)`

Definitions of rubi rules used

rule 211 $\text{Int}[(a_+) + (b_+)(x_+)^2]^{(p_+)}, x_Symbol] \rightarrow \text{Simp}[x*((a + b*x^2)^p/(2*p + 1)), x] + \text{Simp}[2*a*(p/(2*p + 1)) \text{Int}[(a + b*x^2)^{(p - 1)}, x], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{GtQ}[p, 0] \&\& (\text{IntegerQ}[4*p] \parallel \text{IntegerQ}[6*p])$

rule 222 $\text{Int}[1/\text{Sqrt}[(a_+) + (b_+)(x_+)^2], x_Symbol] \rightarrow \text{Simp}[\text{ArcSinh}[\text{Rt}[b, 2]*(x/\text{Sqrt}[a])]/\text{Rt}[b, 2], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{GtQ}[a, 0] \&\& \text{PosQ}[b]$

rule 247 $\text{Int}[(c_+)(x_+)^{(m_+)}*((a_+) + (b_+)(x_+)^2)^{(p_+)}, x_Symbol] \rightarrow \text{Simp}[(c*x)^{(m + 1)}*((a + b*x^2)^p/(c*(m + 1))), x] - \text{Simp}[2*b*(p/(c^2*(m + 1))) \text{Int}[(c*x)^{(m + 2)}*(a + b*x^2)^{(p - 1)}, x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&\& \text{GtQ}[p, 0] \&\& \text{LtQ}[m, -1] \&\& !\text{LtQ}[(m + 2*p + 3)/2, 0] \&\& \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 2045 $\text{Int}[(u_+)*((c_+)*((a_+) + (b_+)(x_+)^{(n_+)})^{(q_+)})^{(p_+)}, x_Symbol] \rightarrow \text{Simp}[\text{Simp}[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^{(p*q)} \text{Int}[u*(1 + b*(x^n/a))^{(p*q)}, x], x] /; \text{FreeQ}[\{a, b, c, n, p, q\}, x] \&\& !\text{GeQ}[a, 0]$

Maple [A] (verified)

Time = 3.19 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.61

method	result
risch	$-\frac{(-8b^4x^8 - 50ab^3x^6 - 165a^2b^2x^4 + 208a^3bx^2 + 16a^4)c\sqrt{c(bx^2+a)^3}}{48(bx^2+a)x^3} + \frac{105a^3b^2 \ln\left(\frac{bcx}{\sqrt{bc}} + \sqrt{x^2bc+ac}\right)c\sqrt{c(bx^2+a)^3} \sqrt{(bx^2+a)c}}{16\sqrt{bc}(bx^2+a)^2}$
default	$-\frac{(c(bx^2+a)^3)^{\frac{3}{2}} \left(-8(x^2bc+ac)^{\frac{5}{2}} \sqrt{bc} b^2 x^4 - 210(x^2bc+ac)^{\frac{3}{2}} \sqrt{bc} a b^2 c x^4 - 315 \sqrt{x^2bc+ac} \sqrt{bc} a^2 b^2 c^2 x^4 - 315 \ln\left(\frac{bcx + \sqrt{x^2bc+ac}}{\sqrt{bc}}\right) \right)}{48(bx^2+a)^3 ((bx^2+a)c)^{\frac{3}{2}} x^3 c \sqrt{bc}}$

input $\text{int}((c*(b*x^2+a)^3)^{(3/2)}/x^4, x, \text{method}=_RETURNVERBOSE)$

output

```
-1/48/(b*x^2+a)*(-8*b^4*x^8-50*a*b^3*x^6-165*a^2*b^2*x^4+208*a^3*b*x^2+16*a^4)/x^3*c*(c*(b*x^2+a)^3)^(1/2)+105/16*a^3*b^2*ln(b*c*x/(b*c)^(1/2)+(b*c*x^2+a*c)^(1/2))/(b*c)^(1/2)*c/(b*x^2+a)^2*(c*(b*x^2+a)^3)^(1/2)*((b*x^2+a)*c)^(1/2)
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 410, normalized size of antiderivative = 1.74

$$\int \frac{(c(a+bx^2))^3}{x^4} dx = \frac{\left[\frac{315(a^3b^2cx^5 + a^4bcx^3)\sqrt{bc} \log\left(-\frac{2b^2cx^4+3abcx^2+a^2c+2\sqrt{b^3cx^6+3ab^2cx^4+3a^2bcx^2+a^3c}\sqrt{b}}{bx^2+a}\right)}{48(bx^5+ax^3)} + \frac{315(a^3b^2cx^5 + a^4bcx^3)\sqrt{-bc} \arctan\left(\frac{\sqrt{b^3cx^6+3ab^2cx^4+3a^2bcx^2+a^3c}\sqrt{-bc}}{b^2cx^4+2abcx^2+a^2c}\right) - (8b^4cx^8 + 50ab^3cx^6 + 165a^2b^2cx^4 - 208a^3b^2cx^2 - 16a^4c)\sqrt{b^3cx^6+3ab^2cx^4+3a^2bcx^2+a^3c}}{48(bx^5+ax^3)} \right]}{48(bx^5+ax^3)}$$

input

```
integrate((c*(b*x^2+a)^3)^(3/2)/x^4,x, algorithm="fricas")
```

output

```
[1/96*(315*(a^3*b^2*c*x^5 + a^4*b*c*x^3)*sqrt(b*c)*log(-(2*b^2*c*x^4 + 3*a*b*c*x^2 + a^2*c + 2*sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c)*sqrt(b*c)*x)/(b*x^2 + a)) + 2*(8*b^4*c*x^8 + 50*a*b^3*c*x^6 + 165*a^2*b^2*c*x^4 - 208*a^3*b^2*c*x^2 - 16*a^4*c)*sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c))/(b*x^5 + a*x^3), -1/48*(315*(a^3*b^2*c*x^5 + a^4*b*c*x^3)*sqrt(-b*c)*arctan(sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c)*sqrt(-b*c)*x/(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)) - (8*b^4*c*x^8 + 50*a*b^3*c*x^6 + 165*a^2*b^2*c*x^4 - 208*a^3*b^2*c*x^2 - 16*a^4*c)*sqrt(b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c))/(b*x^5 + a*x^3)]
```

Sympy [F]

$$\int \frac{(c(a+bx^2))^3}{x^4} dx = \int \frac{(c(a+bx^2))^3}{x^4} dx$$

input

```
integrate((c*(b*x**2+a)**3)**(3/2)/x**4,x)
```

output `Integral((c*(a + b*x**2)**3)**(3/2)/x**4, x)`

Maxima [F]

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^4} dx = \int \frac{((bx^2 + a)^3 c)^{3/2}}{x^4} dx$$

input `integrate((c*(b*x^2+a)^3)^(3/2)/x^4,x, algorithm="maxima")`

output `integrate(((b*x^2 + a)^3*c)^(3/2)/x^4, x)`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 262, normalized size of antiderivative = 1.11

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^4} dx =$$

$$-\frac{1}{96} \left(315 \sqrt{bca}^3 b \log \left(\left(\sqrt{bcx} - \sqrt{bcx^2 + ac} \right)^2 \right) \operatorname{sgn}(bx^2 + a) - 2 (165 a^2 b^2 \operatorname{sgn}(bx^2 + a) + 2 (4 b^4 x^2 \operatorname{sgn}(bx^2 + a) + 25 a b^3 \operatorname{sgn}(bx^2 + a)) x^2) \sqrt{b c x^2 + a c} x - 64 (15 \sqrt{b c} (\sqrt{b c} x - \sqrt{b c x^2 + a c})^4 a^4 b c \operatorname{sgn}(b x^2 + a) - 24 \sqrt{b c} (\sqrt{b c} x - \sqrt{b c x^2 + a c})^2 a^5 b c^2 \operatorname{sgn}(b x^2 + a) + 13 \sqrt{b c} a^6 b c^3 \operatorname{sgn}(b x^2 + a)) / ((\sqrt{b c} x - \sqrt{b c x^2 + a c})^2 - a c)^3 \right) c$$

input `integrate((c*(b*x^2+a)^3)^(3/2)/x^4,x, algorithm="giac")`

output `-1/96*(315*sqrt(b*c)*a^3*b*log((sqrt(b*c)*x - sqrt(b*c*x^2 + a*c))^2)*sgn(b*x^2 + a) - 2*(165*a^2*b^2*sgn(b*x^2 + a) + 2*(4*b^4*x^2*sgn(b*x^2 + a) + 25*a*b^3*sgn(b*x^2 + a))*x^2)*sqrt(b*c*x^2 + a*c)*x - 64*(15*sqrt(b*c)*(sqrt(b*c)*x - sqrt(b*c*x^2 + a*c))^4*a^4*b*c*sgn(b*x^2 + a) - 24*sqrt(b*c)*(sqrt(b*c)*x - sqrt(b*c*x^2 + a*c))^2*a^5*b*c^2*sgn(b*x^2 + a) + 13*sqrt(b*c)*a^6*b*c^3*sgn(b*x^2 + a))/((sqrt(b*c)*x - sqrt(b*c*x^2 + a*c))^2 - a*c)^3)*c`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^4} dx = \int \frac{(c(bx^2 + a)^3)^{3/2}}{x^4} dx$$

input `int((c*(a + b*x^2)^3)^(3/2)/x^4,x)`output `int((c*(a + b*x^2)^3)^(3/2)/x^4, x)`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.56

$$\int \frac{(c(a + bx^2)^3)^{3/2}}{x^4} dx = \frac{\sqrt{c}c(-128\sqrt{bx^2 + a}a^4 - 1664\sqrt{bx^2 + a}a^3bx^2 + 1320\sqrt{bx^2 + a}a^2b^2x^4 + 400\sqrt{bx^2 + a}ab^3x^6 + 64\sqrt{bx^2 + a}b^4x^8 + 2520\sqrt{b}\log(\sqrt{a + bx^2} + \sqrt{b}x)/\sqrt{a})a^3bx^3 + 567\sqrt{b}a^3bx^3)/(384x^3)}$$

input `int((c*(b*x^2+a)^3)^(3/2)/x^4,x)`output `(sqrt(c)*c*(- 128*sqrt(a + b*x**2)*a**4 - 1664*sqrt(a + b*x**2)*a**3*b*x**2 + 1320*sqrt(a + b*x**2)*a**2*b**2*x**4 + 400*sqrt(a + b*x**2)*a*b**3*x**6 + 64*sqrt(a + b*x**2)*b**4*x**8 + 2520*sqrt(b)*log((sqrt(a + b*x**2) + sqrt(b)*x)/sqrt(a))*a**3*b*x**3 + 567*sqrt(b)*a**3*b*x**3)/(384*x**3)`

$$3.20 \quad \int x^3 \left(\frac{c}{a+bx^2} \right)^{3/2} dx$$

Optimal result	181
Mathematica [A] (verified)	181
Rubi [A] (verified)	182
Maple [A] (verified)	183
Fricas [A] (verification not implemented)	184
Sympy [A] (verification not implemented)	184
Maxima [A] (verification not implemented)	185
Giac [A] (verification not implemented)	185
Mupad [B] (verification not implemented)	185
Reduce [B] (verification not implemented)	186

Optimal result

Integrand size = 19, antiderivative size = 49

$$\int x^3 \left(\frac{c}{a+bx^2} \right)^{3/2} dx = \frac{ac\sqrt{\frac{c}{a+bx^2}}}{b^2} + \frac{c\sqrt{\frac{c}{a+bx^2}}(a+bx^2)}{b^2}$$

output

```
a*c*(c/(b*x^2+a))^(1/2)/b^2+c*(c/(b*x^2+a))^(1/2)*(b*x^2+a)/b^2
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.59

$$\int x^3 \left(\frac{c}{a+bx^2} \right)^{3/2} dx = \frac{c\sqrt{\frac{c}{a+bx^2}}(2a+bx^2)}{b^2}$$

input

```
Integrate[x^3*(c/(a + b*x^2))^(3/2),x]
```

output

```
(c*Sqrt[c/(a + b*x^2)]*(2*a + b*x^2))/b^2
```

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.67, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {2045, 243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \left(\frac{c}{a + bx^2} \right)^{3/2} dx \\
 & \quad \downarrow \text{2045} \\
 & \frac{c \sqrt{\frac{bx^2}{a} + 1} \sqrt{\frac{c}{a + bx^2}} \int \frac{x^3}{\left(\frac{bx^2}{a} + 1\right)^{3/2}} dx}{a} \\
 & \quad \downarrow \text{243} \\
 & \frac{c \sqrt{\frac{bx^2}{a} + 1} \sqrt{\frac{c}{a + bx^2}} \int \frac{x^2}{\left(\frac{bx^2}{a} + 1\right)^{3/2}} dx^2}{2a} \\
 & \quad \downarrow \text{53} \\
 & \frac{c \sqrt{\frac{bx^2}{a} + 1} \sqrt{\frac{c}{a + bx^2}} \int \left(\frac{a}{b \sqrt{\frac{bx^2}{a} + 1}} - \frac{a}{b \left(\frac{bx^2}{a} + 1\right)^{3/2}} \right) dx^2}{2a} \\
 & \quad \downarrow \text{2009} \\
 & \frac{c \sqrt{\frac{bx^2}{a} + 1} \left(\frac{2a^2 \sqrt{\frac{bx^2}{a} + 1}}{b^2} + \frac{2a^2}{b^2 \sqrt{\frac{bx^2}{a} + 1}} \right) \sqrt{\frac{c}{a + bx^2}}}{2a}
 \end{aligned}$$

input `Int[x^3*(c/(a + b*x^2))^(3/2),x]`

output `(c*Sqrt[c/(a + b*x^2)]*Sqrt[1 + (b*x^2)/a]*((2*a^2)/(b^2*Sqrt[1 + (b*x^2)/a]) + (2*a^2*Sqrt[1 + (b*x^2)/a])/b^2))/(2*a)`

Definitions of rubi rules used

rule 53 $\text{Int}[(a_.) + (b_.)(x_)^{(m_.)}*((c_.) + (d_.)(x_)^{(n_.)}), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \text{FreeQ}\{a, b, c, d, n\}, x] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ (\ !\text{IntegerQ}[n] \ || \ (\text{EqQ}[c, 0] \ \&\& \ \text{LeQ}[7*m + 4*n + 4, 0]) \ || \ \text{LtQ}[9*m + 5*(n + 1), 0] \ || \ \text{GtQ}[m + n + 2, 0])$

rule 243 $\text{Int}[(x_)^{(m_.)}*((a_) + (b_.)(x_)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[1/2 \ \text{Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] /; \text{FreeQ}\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 2045 $\text{Int}[(u_.)*((c_.)*((a_) + (b_.)(x_)^{(n_.)})^{(q_.)})^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[\text{Simp}[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^{(p*q)} \ \text{Int}[u*(1 + b*(x^n/a))^{(p*q)}, x], x] /; \text{FreeQ}\{a, b, c, n, p, q\}, x] \ \&\& \ !\text{GeQ}[a, 0]$

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.57

method	result	size
trager	$\frac{c(bx^2+2a)\sqrt{\frac{c}{bx^2+a}}}{b^2}$	28
gospers	$\frac{(bx^2+a)(bx^2+2a)\left(\frac{c}{bx^2+a}\right)^{\frac{3}{2}}}{b^2}$	34
default	$\frac{(bx^2+a)(bx^2+2a)\left(\frac{c}{bx^2+a}\right)^{\frac{3}{2}}}{b^2}$	34
orering	$\frac{(bx^2+a)(bx^2+2a)\left(\frac{c}{bx^2+a}\right)^{\frac{3}{2}}}{b^2}$	34
risch	$\frac{ac\sqrt{\frac{c}{bx^2+a}}}{b^2} + \frac{c\sqrt{\frac{c}{bx^2+a}}(bx^2+a)}{b^2}$	46

input $\text{int}(x^3*(c/(b*x^2+a))^{(3/2)}, x, \text{method}=_RETURNVERBOSE)$

output `c*(b*x^2+2*a)/b^2*(c/(b*x^2+a))^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.57

$$\int x^3 \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \frac{(bcx^2 + 2ac) \sqrt{\frac{c}{bx^2+a}}}{b^2}$$

input `integrate(x^3*(c/(b*x^2+a))^(3/2),x, algorithm="fricas")`

output `(b*c*x^2 + 2*a*c)*sqrt(c/(b*x^2 + a))/b^2`

Sympy [A] (verification not implemented)

Time = 0.44 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.35

$$\int x^3 \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \begin{cases} \frac{2a^2 \left(\frac{c}{a+bx^2} \right)^{3/2}}{b^2} + \frac{3ax^2 \left(\frac{c}{a+bx^2} \right)^{3/2}}{b} + x^4 \left(\frac{c}{a+bx^2} \right)^{3/2} & \text{for } b \neq 0 \\ \frac{x^4 \left(\frac{c}{a} \right)^{3/2}}{4} & \text{otherwise} \end{cases}$$

input `integrate(x**3*(c/(b*x**2+a))**(3/2),x)`

output `Piecewise((2*a**2*(c/(a + b*x**2))**(3/2)/b**2 + 3*a*x**2*(c/(a + b*x**2))
(3/2)/b + x4*(c/(a + b*x**2))**(3/2), Ne(b, 0)), (x**4*(c/a)**(3/2)/4,
True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.73

$$\int x^3 \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \frac{\left(a \sqrt{\frac{c}{bx^2+a}} + \frac{c}{\sqrt{\frac{c}{bx^2+a}}} \right) c}{b^2}$$

input `integrate(x^3*(c/(b*x^2+a))^(3/2),x, algorithm="maxima")`output `(a*sqrt(c/(b*x^2 + a)) + c/sqrt(c/(b*x^2 + a)))*c/b^2`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.02

$$\int x^3 \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \frac{\left(\frac{ac^2}{\sqrt{bcx^2+acb}} + \frac{\sqrt{bcx^2+acc}}{b} \right) \text{sgn}(bx^2 + a)}{b}$$

input `integrate(x^3*(c/(b*x^2+a))^(3/2),x, algorithm="giac")`output `(a*c^2/(sqrt(b*c*x^2 + a*c)*b) + sqrt(b*c*x^2 + a*c)*c/b)*sgn(b*x^2 + a)/b`**Mupad [B] (verification not implemented)**

Time = 0.51 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.86

$$\int x^3 \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \frac{cx^2 \sqrt{\frac{c}{bx^2+a}}}{b} + \frac{2ac \sqrt{\frac{c}{bx^2+a}}}{b^2}$$

input `int(x^3*(c/(a + b*x^2))^(3/2),x)`output `(c*x^2*(c/(a + b*x^2))^(1/2))/b + (2*a*c*(c/(a + b*x^2))^(1/2))/b^2`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.67

$$\int x^3 \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \frac{\sqrt{c} \sqrt{bx^2 + a} c (bx^2 + 2a)}{b^2 (bx^2 + a)}$$

input `int(x^3*(c/(b*x^2+a))^(3/2),x)`

output `(sqrt(c)*sqrt(a + b*x**2)*c*(2*a + b*x**2))/(b**2*(a + b*x**2))`

$$3.21 \quad \int x \left(\frac{c}{a+bx^2} \right)^{3/2} dx$$

Optimal result	187
Mathematica [A] (verified)	187
Rubi [A] (verified)	188
Maple [A] (verified)	189
Fricas [A] (verification not implemented)	189
Sympy [B] (verification not implemented)	190
Maxima [A] (verification not implemented)	190
Giac [A] (verification not implemented)	190
Mupad [B] (verification not implemented)	191
Reduce [B] (verification not implemented)	191

Optimal result

Integrand size = 17, antiderivative size = 21

$$\int x \left(\frac{c}{a+bx^2} \right)^{3/2} dx = -\frac{c\sqrt{\frac{c}{a+bx^2}}}{b}$$

output `-c*(c/(b*x^2+a))^(1/2)/b`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int x \left(\frac{c}{a+bx^2} \right)^{3/2} dx = -\frac{c\sqrt{\frac{c}{a+bx^2}}}{b}$$

input `Integrate[x*(c/(a + b*x^2))^(3/2),x]`

output `-((c*Sqrt[c/(a + b*x^2)])/b)`

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2024, 19}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \left(\frac{c}{a + bx^2} \right)^{3/2} dx$$

$$\downarrow \text{2024}$$

$$\int \left(\frac{c}{bx^2 + a} \right)^{3/2} d(bx^2 + a)$$

$$\frac{2b}{2b}$$

$$\downarrow \text{19}$$

$$-\frac{c\sqrt{\frac{c}{a+bx^2}}}{b}$$

input `Int[x*(c/(a + b*x^2))^(3/2),x]`

output `-((c*Sqrt[c/(a + b*x^2)])/b)`

Defintions of rubi rules used

rule 19 `Int[((a_.)/(x_))^(p_), x_Symbol] := Simp[(-a)*((a/x)^(p - 1)/(p - 1)), x] / ; FreeQ[{a, p}, x] && !IntegerQ[p]`

rule 2024 `Int[((a_.) + (b_.)*(Pq_)^(n_.))^(p_.)*(Qr_), x_Symbol] := With[{q = Expon[Pq, x], r = Expon[Qr, x]}, Simp[Coeff[Qr, x, r]/(q*Coeff[Pq, x, q]) Subst[Int[(a + b*x^n)^p, x], x, Pq], x] /; EqQ[r, q - 1] && EqQ[Coeff[Qr, x, r]*D[Pq, x], q*Coeff[Pq, x, q]*Qr] /; FreeQ[{a, b, n, p}, x] && PolyQ[Pq, x] & & PolyQ[Qr, x]`

Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.95

method	result	size
trager	$-\frac{c\sqrt{\frac{c}{bx^2+a}}}{b}$	20
gospers	$-\frac{(bx^2+a)\left(\frac{c}{bx^2+a}\right)^{3/2}}{b}$	26
derivativdivides	$-\frac{(bx^2+a)\left(\frac{c}{bx^2+a}\right)^{3/2}}{b}$	26
default	$-\frac{(bx^2+a)\left(\frac{c}{bx^2+a}\right)^{3/2}}{b}$	26
orering	$-\frac{(bx^2+a)\left(\frac{c}{bx^2+a}\right)^{3/2}}{b}$	26

input `int(x*(c/(b*x^2+a))^(3/2),x,method=_RETURNVERBOSE)`

output `-c*(c/(b*x^2+a))^(1/2)/b`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int x \left(\frac{c}{a + bx^2} \right)^{3/2} dx = -\frac{c\sqrt{\frac{c}{bx^2+a}}}{b}$$

input `integrate(x*(c/(b*x^2+a))^(3/2),x, algorithm="fricas")`

output `-c*sqrt(c/(b*x^2 + a))/b`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 42 vs. 2(15) = 30.

Time = 0.26 (sec) , antiderivative size = 42, normalized size of antiderivative = 2.00

$$\int x \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \begin{cases} -\frac{a \left(\frac{c}{a + bx^2} \right)^{3/2}}{b} - x^2 \left(\frac{c}{a + bx^2} \right)^{3/2} & \text{for } b \neq 0 \\ \frac{x^2 \left(\frac{c}{a} \right)^{3/2}}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*(c/(b*x**2+a))**(3/2),x)`

output `Piecewise((-a*(c/(a + b*x**2))**(3/2)/b - x**2*(c/(a + b*x**2))**(3/2), Ne(b, 0)), (x**2*(c/a)**(3/2)/2, True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int x \left(\frac{c}{a + bx^2} \right)^{3/2} dx = -\frac{c \sqrt{\frac{c}{bx^2 + a}}}{b}$$

input `integrate(x*(c/(b*x^2+a))^(3/2),x, algorithm="maxima")`

output `-c*sqrt(c/(b*x^2 + a))/b`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.33

$$\int x \left(\frac{c}{a + bx^2} \right)^{3/2} dx = -\frac{c^2 \operatorname{sgn}(bx^2 + a)}{\sqrt{bcx^2 + acb}}$$

input `integrate(x*(c/(b*x^2+a))^(3/2),x, algorithm="giac")`

output $-c^2 \operatorname{sgn}(bx^2 + a) / (\sqrt{bcx^2 + ac}) * b$

Mupad [B] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int x \left(\frac{c}{a + bx^2} \right)^{3/2} dx = -\frac{c \sqrt{\frac{c}{bx^2 + a}}}{b}$$

input $\operatorname{int}(x*(c/(a + b*x^2))^(3/2), x)$

output $-(c*(c/(a + b*x^2))^(1/2))/b$

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.19

$$\int x \left(\frac{c}{a + bx^2} \right)^{3/2} dx = -\frac{\sqrt{c} \sqrt{bx^2 + a} c}{b(bx^2 + a)}$$

input $\operatorname{int}(x*(c/(b*x^2+a))^(3/2), x)$

output $(- \operatorname{sqrt}(c) * \operatorname{sqrt}(a + b*x**2) * c) / (b * (a + b*x**2))$

3.22 $\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x} dx$

Optimal result	192
Mathematica [A] (verified)	192
Rubi [A] (verified)	193
Maple [A] (verified)	195
Fricas [A] (verification not implemented)	195
Sympy [F]	196
Maxima [A] (verification not implemented)	196
Giac [A] (verification not implemented)	196
Mupad [F(-1)]	197
Reduce [B] (verification not implemented)	197

Optimal result

Integrand size = 19, antiderivative size = 71

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x} dx = \frac{c\sqrt{\frac{c}{a+bx^2}}}{a} - \frac{c\sqrt{\frac{c}{a+bx^2}}\sqrt{1+\frac{bx^2}{a}}\operatorname{arctanh}\left(\sqrt{1+\frac{bx^2}{a}}\right)}{a}$$

output

```
c*(c/(b*x^2+a))^(1/2)/a-c*(c/(b*x^2+a))^(1/2)*(1+b*x^2/a)^(1/2)*arctanh((1+b*x^2/a)^(1/2))/a
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.83

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x} dx = \frac{c\sqrt{\frac{c}{a+bx^2}}\left(\sqrt{a}-\sqrt{a+bx^2}\operatorname{arctanh}\left(\frac{\sqrt{a+bx^2}}{\sqrt{a}}\right)\right)}{a^{3/2}}$$

input

```
Integrate[(c/(a + b*x^2))^(3/2)/x,x]
```

output

```
(c*Sqrt[c/(a + b*x^2)]*(Sqrt[a] - Sqrt[a + b*x^2]*ArcTanh[Sqrt[a + b*x^2]/Sqrt[a]]))/a^(3/2)
```

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {2045, 243, 61, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x} dx \\
 & \quad \downarrow \text{2045} \\
 & \frac{c\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{c}{a+bx^2}} \int \frac{1}{x\left(\frac{bx^2}{a} + 1\right)^{3/2}} dx}{a} \\
 & \quad \downarrow \text{243} \\
 & \frac{c\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{c}{a+bx^2}} \int \frac{1}{x^2\left(\frac{bx^2}{a} + 1\right)^{3/2}} dx^2}{2a} \\
 & \quad \downarrow \text{61} \\
 & \frac{c\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{c}{a+bx^2}} \left(\int \frac{1}{x^2\sqrt{\frac{bx^2}{a} + 1}} dx^2 + \frac{2}{\sqrt{\frac{bx^2}{a} + 1}} \right)}{2a} \\
 & \quad \downarrow \text{73} \\
 & \frac{c\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{c}{a+bx^2}} \left(\frac{2a \int \frac{1}{\frac{ax^4}{b} - \frac{a}{b}} d\sqrt{\frac{bx^2}{a} + 1}}{\frac{ax^4}{b} - \frac{a}{b}} + \frac{2}{\sqrt{\frac{bx^2}{a} + 1}} \right)}{2a} \\
 & \quad \downarrow \text{221} \\
 & \frac{c\sqrt{\frac{bx^2}{a} + 1} \left(\frac{2}{\sqrt{\frac{bx^2}{a} + 1}} - 2\operatorname{arctanh}\left(\sqrt{\frac{bx^2}{a} + 1}\right) \right) \sqrt{\frac{c}{a+bx^2}}}{2a}
 \end{aligned}$$

input

```
Int[(c/(a + b*x^2))^(3/2)/x,x]
```

output $(c\sqrt{c/(a + b*x^2)}*\sqrt{1 + (b*x^2)/a}*(2/\sqrt{1 + (b*x^2)/a} - 2*\text{ArcTanh}[\sqrt{1 + (b*x^2)/a}]])/ (2*a)$

Defintions of rubi rules used

rule 61 $\text{Int}[(a_.) + (b_.)*(x_)^(m_)*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] \rightarrow \text{Simp}[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - \text{Simp}[d*((m + n + 2)/((b*c - a*d)*(m + 1))) \text{Int}[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /;$ $\text{FreeQ}\{a, b, c, d, n\}, x\} \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{!(LtQ}[n, -1] \ \&\& \ (\text{EqQ}[a, 0] \ || \ (\text{NeQ}[c, 0] \ \&\& \ \text{LtQ}[m - n, 0] \ \&\& \ \text{IntegerQ}[n])) \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 73 $\text{Int}[(a_.) + (b_.)*(x_)^(m_)*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] \rightarrow \text{With}\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \ \text{Subst}[\text{Int}[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /;$ $\text{FreeQ}\{a, b, c, d\}, x\} \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 221 $\text{Int}[(a_) + (b_.)*(x_)^2)^(-1), x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /;$ $\text{FreeQ}\{a, b\}, x\} \ \&\& \ \text{NegQ}[a/b]$

rule 243 $\text{Int}[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] \rightarrow \text{Simp}[1/2 \ \text{Subst}[\text{Int}[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /;$ $\text{FreeQ}\{a, b, m, p\}, x\} \ \&\& \ \text{IntegerQ}[(m - 1)/2]$

rule 2045 $\text{Int}[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_))^(p_), x_Symbol] \rightarrow \text{Simp}[\text{Simp}[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] \ \text{Int}[u*(1 + b*(x^n/a))^(p*q), x], x] /;$ $\text{FreeQ}\{a, b, c, n, p, q\}, x\} \ \&\& \ \text{!GeQ}[a, 0]$

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.90

method	result	size
default	$-\frac{\left(\frac{c}{bx^2+a}\right)^{\frac{3}{2}}(bx^2+a)\left(\ln\left(\frac{2a+2\sqrt{a}\sqrt{bx^2+a}}{x}\right)a\sqrt{bx^2+a}-a^{\frac{3}{2}}\right)}{a^{\frac{5}{2}}}$	64

input `int((c/(b*x^2+a))^(3/2)/x,x,method=_RETURNVERBOSE)`output `-(c/(b*x^2+a))^(3/2)*(b*x^2+a)*(ln(2*(a^(1/2))*(b*x^2+a)^(1/2)+a)/x)*a*(b*x^2+a)^(1/2)-a^(3/2))/a^(5/2)`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.94

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x} dx = \left[\frac{c\sqrt{\frac{c}{a}} \log\left(-\frac{bcx^2+2ac-2(abx^2+a^2)\sqrt{\frac{c}{bx^2+a}}\sqrt{\frac{c}{a}}}{x^2}\right) + 2c\sqrt{\frac{c}{bx^2+a}}}{2a}, \frac{c\sqrt{-\frac{c}{a}} \arctan\left(\frac{a\sqrt{\frac{c}{bx^2+a}}\sqrt{-\frac{c}{a}}}{c}\right)}{a} \right]$$

input `integrate((c/(b*x^2+a))^(3/2)/x,x, algorithm="fricas")`output `[1/2*(c*sqrt(c/a)*log(-(b*c*x^2 + 2*a*c - 2*(a*b*x^2 + a^2)*sqrt(c/(b*x^2 + a))*sqrt(c/a))/x^2) + 2*c*sqrt(c/(b*x^2 + a)))/a, (c*sqrt(-c/a)*arctan(a*sqrt(c/(b*x^2 + a))*sqrt(-c/a)/c) + c*sqrt(c/(b*x^2 + a)))/a]`

Sympy [F]

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x} dx = \int \frac{\left(\frac{c}{a+bx^2}\right)^{\frac{3}{2}}}{x} dx$$

input `integrate((c/(b*x**2+a))**(3/2)/x,x)`

output `Integral((c/(a + b*x**2))**(3/2)/x, x)`

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.13

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x} dx = \frac{1}{2} c \left(\frac{c \log \left(\frac{a \sqrt{\frac{c}{bx^2+a}} - \sqrt{ac}}{a \sqrt{\frac{c}{bx^2+a}} + \sqrt{ac}} \right)}{\sqrt{aca}} + \frac{2 \sqrt{\frac{c}{bx^2+a}}}{a} \right)$$

input `integrate((c/(b*x^2+a))^(3/2)/x,x, algorithm="maxima")`

output `1/2*c*(c*log((a*sqrt(c/(b*x^2 + a)) - sqrt(a*c))/(a*sqrt(c/(b*x^2 + a)) + sqrt(a*c)))/(sqrt(a*c)*a) + 2*sqrt(c/(b*x^2 + a))/a)`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.83

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x} dx = c \left(\frac{c \arctan \left(\frac{\sqrt{bcx^2+ac}}{\sqrt{-ac}} \right)}{\sqrt{-aca}} + \frac{c}{\sqrt{bcx^2 + aca}} \right) \operatorname{sgn}(bx^2 + a)$$

input `integrate((c/(b*x^2+a))^(3/2)/x,x, algorithm="giac")`

output

```
c*(c*arctan(sqrt(b*c*x^2 + a*c)/sqrt(-a*c))/(sqrt(-a*c)*a) + c/(sqrt(b*c*x^2 + a*c)*a))*sgn(b*x^2 + a)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x} dx = \int \frac{\left(\frac{c}{bx^2+a}\right)^{3/2}}{x} dx$$

input

```
int((c/(a + b*x^2))^(3/2)/x,x)
```

output

```
int((c/(a + b*x^2))^(3/2)/x, x)
```

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.96

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x} dx = \frac{\sqrt{c}c\left(\sqrt{bx^2+a}a + \sqrt{a}\log\left(\frac{\sqrt{bx^2+a}-\sqrt{a}+\sqrt{bx}}{\sqrt{a}}\right)a + \sqrt{a}\log\left(\frac{\sqrt{bx^2+a}-\sqrt{a}+\sqrt{bx}}{\sqrt{a}}\right)bx^2 - \sqrt{a}\log\left(\frac{\sqrt{bx^2+a}+\sqrt{a}+\sqrt{bx}}{\sqrt{a}}\right)a - \sqrt{a}\log\left(\frac{\sqrt{bx^2+a}+\sqrt{a}+\sqrt{bx}}{\sqrt{a}}\right)bx^2\right)}{a^2(bx^2+a)}$$

input

```
int((c/(b*x^2+a))^(3/2)/x,x)
```

output

```
(sqrt(c)*c*(sqrt(a + b*x**2)*a + sqrt(a)*log((sqrt(a + b*x**2) - sqrt(a) + sqrt(b)*x)/sqrt(a))*a + sqrt(a)*log((sqrt(a + b*x**2) - sqrt(a) + sqrt(b)*x)/sqrt(a))*b*x**2 - sqrt(a)*log((sqrt(a + b*x**2) + sqrt(a) + sqrt(b)*x)/sqrt(a))*a - sqrt(a)*log((sqrt(a + b*x**2) + sqrt(a) + sqrt(b)*x)/sqrt(a))*b*x**2)/(a**2*(a + b*x**2))
```

3.23 $\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^3} dx$

Optimal result	198
Mathematica [A] (verified)	198
Rubi [A] (verified)	199
Maple [A] (verified)	201
Fricas [A] (verification not implemented)	202
Sympy [F]	202
Maxima [A] (verification not implemented)	203
Giac [A] (verification not implemented)	203
Mupad [F(-1)]	204
Reduce [B] (verification not implemented)	204

Optimal result

Integrand size = 19, antiderivative size = 104

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^3} dx = -\frac{3bc\sqrt{\frac{c}{a+bx^2}}}{2a^2} - \frac{c\sqrt{\frac{c}{a+bx^2}}}{2ax^2} + \frac{3bc\sqrt{\frac{c}{a+bx^2}}\sqrt{1+\frac{bx^2}{a}}\operatorname{arctanh}\left(\sqrt{1+\frac{bx^2}{a}}\right)}{2a^2}$$

output

$$-3/2*b*c*(c/(b*x^2+a))^(1/2)/a^2-1/2*c*(c/(b*x^2+a))^(1/2)/a/x^2+3/2*b*c*(c/(b*x^2+a))^(1/2)*(1+b*x^2/a)^(1/2)*\operatorname{arctanh}((1+b*x^2/a)^(1/2))/a^2$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.75

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^3} dx = -\frac{c\sqrt{\frac{c}{a+bx^2}}\left(\sqrt{a}(a+3bx^2) - 3bx^2\sqrt{a+bx^2}\operatorname{arctanh}\left(\frac{\sqrt{a+bx^2}}{\sqrt{a}}\right)\right)}{2a^{5/2}x^2}$$

input

```
Integrate[(c/(a + b*x^2))^(3/2)/x^3,x]
```

output

$$-1/2*(c*\operatorname{Sqrt}[c/(a + b*x^2)]*(\operatorname{Sqrt}[a]*(a + 3*b*x^2) - 3*b*x^2*\operatorname{Sqrt}[a + b*x^2])*ArcTanh[\operatorname{Sqrt}[a + b*x^2]/\operatorname{Sqrt}[a]])/(a^(5/2)*x^2)$$

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.95, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {2045, 243, 52, 61, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^3} dx \\
 & \quad \downarrow \text{2045} \\
 & \frac{c\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{c}{a+bx^2}} \int \frac{1}{x^3\left(\frac{bx^2}{a} + 1\right)^{3/2}} dx}{a} \\
 & \quad \downarrow \text{243} \\
 & \frac{c\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{c}{a+bx^2}} \int \frac{1}{x^4\left(\frac{bx^2}{a} + 1\right)^{3/2}} dx^2}{2a} \\
 & \quad \downarrow \text{52} \\
 & \frac{c\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{c}{a+bx^2}} \left(-\frac{3b \int \frac{1}{x^2\left(\frac{bx^2}{a} + 1\right)^{3/2}} dx^2}{2a} - \frac{1}{x^2\sqrt{\frac{bx^2}{a} + 1}} \right)}{2a} \\
 & \quad \downarrow \text{61} \\
 & \frac{c\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{c}{a+bx^2}} \left(-\frac{3b \left(\int \frac{1}{x^2\sqrt{\frac{bx^2}{a} + 1}} dx^2 + \frac{2}{\sqrt{\frac{bx^2}{a} + 1}} \right)}{2a} - \frac{1}{x^2\sqrt{\frac{bx^2}{a} + 1}} \right)}{2a} \\
 & \quad \downarrow \text{73}
 \end{aligned}$$

$$\frac{c\sqrt{\frac{bx^2}{a}+1}\sqrt{\frac{c}{a+bx^2}}}{2a} \left(-\frac{3b\left(\frac{2a\int\frac{1}{ax^4-\frac{a}{b}}dx\sqrt{\frac{bx^2}{a}+1}}{b}+\frac{2}{\sqrt{\frac{bx^2}{a}+1}}\right)}{2a}-\frac{1}{x^2\sqrt{\frac{bx^2}{a}+1}} \right)$$

221

$$\frac{c\sqrt{\frac{bx^2}{a}+1}}{2a} \left(-\frac{3b\left(\frac{2}{\sqrt{\frac{bx^2}{a}+1}}-2\operatorname{arctanh}\left(\sqrt{\frac{bx^2}{a}+1}\right)\right)}{2a}-\frac{1}{x^2\sqrt{\frac{bx^2}{a}+1}} \right) \sqrt{\frac{c}{a+bx^2}}$$

input `Int[(c/(a + b*x^2))^(3/2)/x^3,x]`

output `(c*Sqrt[c/(a + b*x^2)]*Sqrt[1 + (b*x^2)/a]*(-(1/(x^2*Sqrt[1 + (b*x^2)/a])) - (3*b*(2/Sqrt[1 + (b*x^2)/a] - 2*ArcTanh[Sqrt[1 + (b*x^2)/a]]))/(2*a)))/(2*a)`

Defintions of rubi rules used

rule 52 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 61 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
 /Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[In
 t[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && I
 ntegerQ[(m - 1)/2]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_.))^(p_), x_Symbol] := Simp[Si
 mp[(c*(a + b*x^n)^q]^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q)
 , x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.76

method	result
default	$-\frac{\left(\frac{c}{bx^2+a}\right)^{\frac{3}{2}}(bx^2+a)\left(3a^{\frac{3}{2}}bx^2-3\ln\left(\frac{2a+2\sqrt{a}\sqrt{bx^2+a}}{x}\right)\sqrt{bx^2+a}abx^2+a^{\frac{5}{2}}\right)}{2a^{\frac{7}{2}}x^2}$
risch	$-\frac{(bx^2+a)c\sqrt{\frac{c}{bx^2+a}}}{2a^2x^2} - \frac{b\left(-\frac{3\ln\left(\frac{2ac+2\sqrt{ac}\sqrt{x^2bc+ac}}{\sqrt{ac}}\right)}{\sqrt{ac}} + \frac{\sqrt{bc\left(x-\frac{\sqrt{-ab}}{b}\right)^2+2c\sqrt{-ab}\left(x-\frac{\sqrt{-ab}}{b}\right)}}{c\sqrt{-ab}\left(x-\frac{\sqrt{-ab}}{b}\right)} - \frac{\sqrt{bc\left(x+\frac{\sqrt{-ab}}{b}\right)^2-2c\sqrt{-ab}\left(x+\frac{\sqrt{-ab}}{b}\right)}}{c\sqrt{-ab}\left(x+\frac{\sqrt{-ab}}{b}\right)}\right)}{2a^2}$

input `int((c/(b*x^2+a))^(3/2)/x^3,x,method=_RETURNVERBOSE)`

output `-1/2*(c/(b*x^2+a))^(3/2)*(b*x^2+a)*(3*a^(3/2)*b*x^2-3*ln(2*(a^(1/2)*(b*x^2
 +a)^(1/2)+a)/x)*(b*x^2+a)^(1/2)*a*b*x^2+a^(5/2))/a^(7/2)/x^2`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.68

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^3} dx = \left[\frac{3bcx^2 \sqrt{\frac{c}{a}} \log\left(-\frac{bcx^2+2ac+2(abx^2+a^2)\sqrt{\frac{c}{bx^2+a}}\sqrt{\frac{c}{a}}}{x^2}\right) - 2(3bcx^2+ac)\sqrt{\frac{c}{bx^2+a}}}{4a^2x^2}, \right. \\ \left. - \frac{3bcx^2 \sqrt{-\frac{c}{a}} \arctan\left(\frac{a\sqrt{\frac{c}{bx^2+a}}\sqrt{-\frac{c}{a}}}{c}\right) + (3bcx^2+ac)\sqrt{\frac{c}{bx^2+a}}}{2a^2x^2} \right]$$

input `integrate((c/(b*x^2+a))^(3/2)/x^3,x, algorithm="fricas")`output `[1/4*(3*b*c*x^2*sqrt(c/a)*log(-(b*c*x^2 + 2*a*c + 2*(a*b*x^2 + a^2)*sqrt(c/(b*x^2 + a))*sqrt(c/a))/x^2) - 2*(3*b*c*x^2 + a*c)*sqrt(c/(b*x^2 + a)))/(a^2*x^2), -1/2*(3*b*c*x^2*sqrt(-c/a)*arctan(a*sqrt(c/(b*x^2 + a))*sqrt(-c/a)/c) + (3*b*c*x^2 + a*c)*sqrt(c/(b*x^2 + a)))/(a^2*x^2)]`**Sympy [F]**

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^3} dx = \int \frac{\left(\frac{c}{a+bx^2}\right)^{\frac{3}{2}}}{x^3} dx$$

input `integrate((c/(b*x**2+a))**(3/2)/x**3,x)`output `Integral((c/(a + b*x**2))**(3/2)/x**3, x)`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.16

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^3} dx = -\frac{1}{4} bc \left(\frac{2c\sqrt{\frac{c}{bx^2+a}}}{a^2c - \frac{a^3c}{bx^2+a}} + \frac{3c \log\left(\frac{a\sqrt{\frac{c}{bx^2+a}} - \sqrt{ac}}{a\sqrt{\frac{c}{bx^2+a}} + \sqrt{ac}}\right)}{\sqrt{aca^2}} + \frac{4\sqrt{\frac{c}{bx^2+a}}}{a^2} \right)$$

input `integrate((c/(b*x^2+a))^(3/2)/x^3,x, algorithm="maxima")`

output `-1/4*b*c*(2*c*sqrt(c/(b*x^2 + a))/(a^2*c - a^3*c/(b*x^2 + a)) + 3*c*log((a*sqrt(c/(b*x^2 + a)) - sqrt(a*c))/(a*sqrt(c/(b*x^2 + a)) + sqrt(a*c)))/(sqrt(a*c)*a^2) + 4*sqrt(c/(b*x^2 + a))/a^2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.99

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^3} dx = -\frac{1}{2} c \left(\frac{3bc \arctan\left(\frac{\sqrt{bcx^2+ac}}{\sqrt{-ac}}\right)}{\sqrt{-aca^2}} + \frac{2abc^2 - 3(bc^2 + ac)bc}{\left(\sqrt{bcx^2 + ac}ac - (bcx^2 + ac)^{\frac{3}{2}}\right)a^2} \right) \operatorname{sgn}(bx^2 + a)$$

input `integrate((c/(b*x^2+a))^(3/2)/x^3,x, algorithm="giac")`

output `-1/2*c*(3*b*c*arctan(sqrt(b*c*x^2 + a*c)/sqrt(-a*c))/(sqrt(-a*c)*a^2) + (2*a*b*c^2 - 3*(b*c*x^2 + a*c)*b*c)/((sqrt(b*c*x^2 + a*c)*a*c - (b*c*x^2 + a*c)^(3/2))*a^2))*sgn(b*x^2 + a)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^3} dx = \int \frac{\left(\frac{c}{bx^2+a}\right)^{3/2}}{x^3} dx$$

input `int((c/(a + b*x^2))^(3/2)/x^3,x)`output `int((c/(a + b*x^2))^(3/2)/x^3, x)`**Reduce [B] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.68

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^3} dx = \frac{\sqrt{c}c\left(-\sqrt{bx^2+a}a^2 - 3\sqrt{bx^2+a}abx^2 - 3\sqrt{a}\log\left(\frac{\sqrt{bx^2+a}-\sqrt{a}+\sqrt{bx}}{\sqrt{a}}\right)abx^2 - 3\sqrt{a}\log\left(\frac{\sqrt{bx^2+a}+\sqrt{a}+\sqrt{bx}}{\sqrt{a}}\right)abx^2\right)}{2a^3x^2}$$

input `int((c/(b*x^2+a))^(3/2)/x^3,x)`output `(sqrt(c)*c*(- sqrt(a + b*x**2)*a**2 - 3*sqrt(a + b*x**2)*a*b*x**2 - 3*sqrt(a)*log((sqrt(a + b*x**2) - sqrt(a) + sqrt(b)*x)/sqrt(a))*a*b*x**2 - 3*sqrt(a)*log((sqrt(a + b*x**2) - sqrt(a) + sqrt(b)*x)/sqrt(a))*b**2*x**4 + 3*sqrt(a)*log((sqrt(a + b*x**2) + sqrt(a) + sqrt(b)*x)/sqrt(a))*a*b*x**2 + 3*sqrt(a)*log((sqrt(a + b*x**2) + sqrt(a) + sqrt(b)*x)/sqrt(a))*b**2*x**4))/(2*a**3*x**2*(a + b*x**2))`

3.24 $\int x^4 \left(\frac{c}{a+bx^2}\right)^{3/2} dx$

Optimal result	205
Mathematica [A] (verified)	205
Rubi [A] (verified)	206
Maple [A] (verified)	208
Fricas [A] (verification not implemented)	208
Sympy [F]	209
Maxima [F]	209
Giac [A] (verification not implemented)	209
Mupad [F(-1)]	210
Reduce [B] (verification not implemented)	210

Optimal result

Integrand size = 19, antiderivative size = 113

$$\int x^4 \left(\frac{c}{a+bx^2}\right)^{3/2} dx = -\frac{cx^3 \sqrt{\frac{c}{a+bx^2}}}{b} + \frac{3cx \sqrt{\frac{c}{a+bx^2}}(a+bx^2)}{2b^2} - \frac{3a^{3/2}c \sqrt{\frac{c}{a+bx^2}} \sqrt{1 + \frac{bx^2}{a}} \operatorname{arcsinh}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{2b^{5/2}}$$

output

```
-c*x^3*(c/(b*x^2+a))^(1/2)/b+3/2*c*x*(c/(b*x^2+a))^(1/2)*(b*x^2+a)/b^2-3/2
*a^(3/2)*c*(c/(b*x^2+a))^(1/2)*(1+b*x^2/a)^(1/2)*arcsinh(b^(1/2)*x/a^(1/2)
)/b^(5/2)
```

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.75

$$\int x^4 \left(\frac{c}{a+bx^2}\right)^{3/2} dx = \frac{c \sqrt{\frac{c}{a+bx^2}} \left(\sqrt{bx}(3a+bx^2) + 6a \sqrt{a+bx^2} \operatorname{arctanh}\left(\frac{\sqrt{bx}}{\sqrt{a}-\sqrt{a+bx^2}}\right) \right)}{2b^{5/2}}$$

input

```
Integrate[x^4*(c/(a + b*x^2))^(3/2),x]
```

output

```
(c*Sqrt[c/(a + b*x^2)]*(Sqrt[b]*x*(3*a + b*x^2) + 6*a*Sqrt[a + b*x^2]*ArcTanh[(Sqrt[b]*x)/(Sqrt[a] - Sqrt[a + b*x^2])]))/(2*b^(5/2))
```

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.02, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {2045, 252, 262, 222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^4 \left(\frac{c}{a + bx^2} \right)^{3/2} dx \\
 & \quad \downarrow \text{2045} \\
 & \frac{c \sqrt{\frac{bx^2}{a} + 1} \sqrt{\frac{c}{a + bx^2}} \int \frac{x^4}{\left(\frac{bx^2}{a} + 1\right)^{3/2}} dx}{a} \\
 & \quad \downarrow \text{252} \\
 & \frac{c \sqrt{\frac{bx^2}{a} + 1} \sqrt{\frac{c}{a + bx^2}} \left(\frac{3a \int \frac{x^2}{\sqrt{\frac{bx^2}{a} + 1}} dx}{b} - \frac{ax^3}{b \sqrt{\frac{bx^2}{a} + 1}} \right)}{a} \\
 & \quad \downarrow \text{262} \\
 & \frac{c \sqrt{\frac{bx^2}{a} + 1} \sqrt{\frac{c}{a + bx^2}} \left(\frac{3a \left(\frac{ax \sqrt{\frac{bx^2}{a} + 1}}{2b} - \frac{a \int \frac{1}{\sqrt{\frac{bx^2}{a} + 1}} dx}{2b} \right)}{b} - \frac{ax^3}{b \sqrt{\frac{bx^2}{a} + 1}} \right)}{a} \\
 & \quad \downarrow \text{222}
 \end{aligned}$$

$$\frac{c\sqrt{\frac{bx^2}{a} + 1} \left(\frac{3a \left(\frac{ax\sqrt{\frac{bx^2}{a} + 1}}{2b} - \frac{a^{3/2} \operatorname{arcsinh}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{2b^{3/2}} \right)}{b} - \frac{ax^3}{b\sqrt{\frac{bx^2}{a} + 1}} \right) \sqrt{\frac{c}{a+bx^2}}}{a}$$

input `Int[x^4*(c/(a + b*x^2))^(3/2),x]`

output `(c*Sqrt[c/(a + b*x^2)]*Sqrt[1 + (b*x^2)/a]*(-(a*x^3)/(b*Sqrt[1 + (b*x^2)/a])) + (3*a*((a*x*Sqrt[1 + (b*x^2)/a])/(2*b) - (a^(3/2)*ArcSinh[(Sqrt[b]*x)/Sqrt[a]])/(2*b^(3/2))))/b)/a`

Defintions of rubi rules used

rule 222 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

rule 252 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(2*b*(p + 1))), x] - Simp[c^2*(m - 1)/(2*b*(p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && LtQ[p, -1] && GtQ[m, 1] && !ILtQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 262 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.)))^(q_)]^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.62

method	result
default	$\frac{\left(\frac{c}{bx^2+a}\right)^{\frac{3}{2}}(bx^2+a)\left(b^{\frac{5}{2}}x^3+3b^{\frac{3}{2}}ax-3\ln(\sqrt{b}x+\sqrt{bx^2+a})\sqrt{bx^2+a}ab\right)}{2b^{\frac{7}{2}}}$
risch	$\frac{cx\sqrt{\frac{c}{bx^2+a}}(bx^2+a)}{2b^2} - \frac{a\left(-\frac{\sqrt{bc}\left(x+\frac{\sqrt{-ab}}{b}\right)^2-2c\sqrt{-ab}\left(x+\frac{\sqrt{-ab}}{b}\right)}{bc\left(x+\frac{\sqrt{-ab}}{b}\right)} + \frac{3\ln\left(\frac{bcx}{\sqrt{bc}}+\sqrt{x^2bc+ac}\right)}{\sqrt{bc}} - \frac{\sqrt{bc}\left(x-\frac{\sqrt{-ab}}{b}\right)^2+2c\sqrt{-ab}\left(x-\frac{\sqrt{-ab}}{b}\right)}{bc\left(x-\frac{\sqrt{-ab}}{b}\right)}\right)}{2b^2}$

input `int(x^4*(c/(b*x^2+a))^(3/2),x,method=_RETURNVERBOSE)`output
$$\frac{1}{2}\left(\frac{c}{bx^2+a}\right)^{\frac{3}{2}}(bx^2+a)\left(b^{\frac{5}{2}}x^3+3b^{\frac{3}{2}}ax-3\ln\left(b^{\frac{1}{2}}x+\sqrt{bx^2+a}\right)\sqrt{bx^2+a}ab\right)/b^{\frac{7}{2}}$$
Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.45

$$\int x^4 \left(\frac{c}{a+bx^2}\right)^{3/2} dx = \left[\frac{3ac\sqrt{\frac{c}{b}} \log\left(-2bcx^2 - ac + 2(b^2x^3 + abx)\sqrt{\frac{c}{bx^2+a}}\sqrt{\frac{c}{b}}\right) + 2(bc x^3 + 3acx)\sqrt{\frac{c}{bx^2+a}}}{4b^2} \right]$$

input `integrate(x^4*(c/(b*x^2+a))^(3/2),x, algorithm="fricas")`output
$$\left[\frac{1}{4}\left(3ac\sqrt{\frac{c}{b}}\log\left(-2bcx^2 - ac + 2(b^2x^3 + abx)\sqrt{\frac{c}{bx^2+a}}\sqrt{\frac{c}{b}}\right) + 2(bc x^3 + 3acx)\sqrt{\frac{c}{bx^2+a}}\right)/b^2, \frac{1}{2}\left(3ac\sqrt{-\frac{c}{b}}\arctan\left(bx\sqrt{\frac{c}{bx^2+a}}\sqrt{-\frac{c}{b}}\right) + (bcx^3 + 3acx)\sqrt{\frac{c}{bx^2+a}}\right)/b^2 \right]$$

Sympy [F]

$$\int x^4 \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \int x^4 \left(\frac{c}{a + bx^2} \right)^{\frac{3}{2}} dx$$

input `integrate(x**4*(c/(b*x**2+a))**(3/2), x)`

output `Integral(x**4*(c/(a + b*x**2))**(3/2), x)`

Maxima [F]

$$\int x^4 \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \int x^4 \left(\frac{c}{bx^2 + a} \right)^{\frac{3}{2}} dx$$

input `integrate(x^4*(c/(b*x^2+a))^(3/2), x, algorithm="maxima")`

output `integrate(x^4*(c/(b*x^2 + a))^(3/2), x)`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.82

$$\int x^4 \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \frac{1}{2} \left(\frac{\left(\frac{cx^2 \operatorname{sgn}(bx^2+a)}{b} + \frac{3ac \operatorname{sgn}(bx^2+a)}{b^2} \right) x}{\sqrt{bcx^2 + ac}} + \frac{3ac \log \left(\left| -\sqrt{bc}x + \sqrt{bcx^2 + ac} \right| \right) \operatorname{sgn}(bx^2)}{\sqrt{bcb^2}} \right)$$

input `integrate(x^4*(c/(b*x^2+a))^(3/2), x, algorithm="giac")`

output `1/2*((c*x^2*sgn(b*x^2 + a)/b + 3*a*c*sgn(b*x^2 + a)/b^2)*x/sqrt(b*c*x^2 + a*c) + 3*a*c*log(abs(-sqrt(b*c)*x + sqrt(b*c*x^2 + a*c)))*sgn(b*x^2 + a)/(sqrt(b*c)*b^2))*c`

Mupad [F(-1)]

Timed out.

$$\int x^4 \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \int x^4 \left(\frac{c}{bx^2 + a} \right)^{3/2} dx$$

input `int(x^4*(c/(a + b*x^2))^(3/2),x)`output `int(x^4*(c/(a + b*x^2))^(3/2), x)`**Reduce [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.04

$$\int x^4 \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \frac{\sqrt{c} c \left(12\sqrt{bx^2 + a} abx + 4\sqrt{bx^2 + a} b^2 x^3 - 12\sqrt{b} \log\left(\frac{\sqrt{bx^2 + a} + \sqrt{bx}}{\sqrt{a}}\right) a^2 - 12\sqrt{b} \log\left(\frac{\sqrt{bx^2 + a} - \sqrt{bx}}{\sqrt{a}}\right) a^2 \right)}{8b^3 (bx^2 + a)}$$

input `int(x^4*(c/(b*x^2+a))^(3/2),x)`output `(sqrt(c)*c*(12*sqrt(a + b*x**2)*a*b*x + 4*sqrt(a + b*x**2)*b**2*x**3 - 12*sqrt(b)*log((sqrt(a + b*x**2) + sqrt(b)*x)/sqrt(a))*a**2 - 12*sqrt(b)*log((sqrt(a + b*x**2) - sqrt(b)*x)/sqrt(a))*a*b*x**2 + 9*sqrt(b)*a**2 + 9*sqrt(b)*a*b*x**2))/(8*b**3*(a + b*x**2))`

3.25 $\int x^2 \left(\frac{c}{a+bx^2} \right)^{3/2} dx$

Optimal result	211
Mathematica [A] (verified)	211
Rubi [A] (verified)	212
Maple [A] (verified)	213
Fricas [A] (verification not implemented)	214
Sympy [F]	214
Maxima [F]	215
Giac [A] (verification not implemented)	215
Mupad [F(-1)]	215
Reduce [B] (verification not implemented)	216

Optimal result

Integrand size = 19, antiderivative size = 77

$$\int x^2 \left(\frac{c}{a+bx^2} \right)^{3/2} dx = -\frac{cx\sqrt{\frac{c}{a+bx^2}}}{b} + \frac{\sqrt{ac}\sqrt{\frac{c}{a+bx^2}}\sqrt{1+\frac{bx^2}{a}}\operatorname{arcsinh}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{b^{3/2}}$$

output

```
-c*x*(c/(b*x^2+a))^(1/2)/b+a^(1/2)*c*(c/(b*x^2+a))^(1/2)*(1+b*x^2/a)^(1/2)
*arcsinh(b^(1/2)*x/a^(1/2))/b^(3/2)
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.95

$$\int x^2 \left(\frac{c}{a+bx^2} \right)^{3/2} dx = -\frac{c\sqrt{\frac{c}{a+bx^2}}\left(\sqrt{bx} + 2\sqrt{a+bx^2}\operatorname{arctanh}\left(\frac{\sqrt{bx}}{\sqrt{a}-\sqrt{a+bx^2}}\right)\right)}{b^{3/2}}$$

input

```
Integrate[x^2*(c/(a + b*x^2))^(3/2),x]
```

output

```
-((c*Sqrt[c/(a + b*x^2)]*(Sqrt[b]*x + 2*Sqrt[a + b*x^2]*ArcTanh[(Sqrt[b]*x)
]/(Sqrt[a] - Sqrt[a + b*x^2])))/b^(3/2))
```

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.04, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2045, 252, 222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int x^2 \left(\frac{c}{a + bx^2} \right)^{3/2} dx \\
 \downarrow \text{2045} \\
 \frac{c \sqrt{\frac{bx^2}{a} + 1} \sqrt{\frac{c}{a + bx^2}} \int \frac{x^2}{\left(\frac{bx^2}{a} + 1 \right)^{3/2}} dx}{a} \\
 \downarrow \text{252} \\
 \frac{c \sqrt{\frac{bx^2}{a} + 1} \sqrt{\frac{c}{a + bx^2}} \left(a \int \frac{1}{\sqrt{\frac{bx^2}{a} + 1}} dx - \frac{ax}{b \sqrt{\frac{bx^2}{a} + 1}} \right)}{a} \\
 \downarrow \text{222} \\
 \frac{c \sqrt{\frac{bx^2}{a} + 1} \left(\frac{a^{3/2} \operatorname{arcsinh} \left(\frac{\sqrt{bx}}{\sqrt{a}} \right)}{b^{3/2}} - \frac{ax}{b \sqrt{\frac{bx^2}{a} + 1}} \right) \sqrt{\frac{c}{a + bx^2}}}{a}
 \end{array}$$

input `Int [x^2*(c/(a + b*x^2))^(3/2),x]`

output `(c*Sqrt[c/(a + b*x^2)]*Sqrt[1 + (b*x^2)/a]*(-(a*x)/(b*Sqrt[1 + (b*x^2)/a]) + (a^(3/2)*ArcSinh[(Sqrt[b]*x)/Sqrt[a]])/b^(3/2)))/a`

Definitions of rubi rules used

rule 222 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

rule 252 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(2*b*(p + 1))), x] - Simp[c^2*(m - 1)/(2*b*(p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && LtQ[p, -1] && GtQ[m, 1] && !ILtQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.78

method	result	size
default	$-\left(\frac{c}{bx^2+a}\right)^{\frac{3}{2}}(bx^2+a)\left(\frac{3}{2}x - \ln\left(\sqrt{bx} + \sqrt{bx^2+a}\right)b\sqrt{bx^2+a}\right)$	60

input `int(x^2*(c/(b*x^2+a))^(3/2),x,method=_RETURNVERBOSE)`

output `-(c/(b*x^2+a))^(3/2)*(b*x^2+a)*(b^(3/2)*x-ln(b^(1/2)*x+(b*x^2+a)^(1/2))*b*(b*x^2+a)^(1/2))/b^(5/2)`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.83

$$\int x^2 \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \left[-\frac{2cx\sqrt{\frac{c}{bx^2+a}} - c\sqrt{\frac{c}{b}} \log\left(-2bcx^2 - ac - 2(b^2x^3 + abx)\sqrt{\frac{c}{bx^2+a}}\sqrt{\frac{c}{b}}\right)}{2b}, \right. \\ \left. -\frac{cx\sqrt{\frac{c}{bx^2+a}} + c\sqrt{-\frac{c}{b}} \arctan\left(\frac{bx\sqrt{\frac{c}{bx^2+a}}\sqrt{-\frac{c}{b}}}{c}\right)}{b} \right]$$

input `integrate(x^2*(c/(b*x^2+a))^(3/2),x, algorithm="fricas")`

output `[-1/2*(2*c*x*sqrt(c/(b*x^2 + a)) - c*sqrt(c/b)*log(-2*b*c*x^2 - a*c - 2*(b^2*x^3 + a*b*x)*sqrt(c/(b*x^2 + a))*sqrt(c/b)))/b, -(c*x*sqrt(c/(b*x^2 + a)) + c*sqrt(-c/b)*arctan(b*x*sqrt(c/(b*x^2 + a))*sqrt(-c/b)/c))/b]`

Sympy [F]

$$\int x^2 \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \int x^2 \left(\frac{c}{a + bx^2} \right)^{\frac{3}{2}} dx$$

input `integrate(x**2*(c/(b*x**2+a))**(3/2),x)`

output `Integral(x**2*(c/(a + b*x**2))**(3/2), x)`

Maxima [F]

$$\int x^2 \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \int x^2 \left(\frac{c}{bx^2 + a} \right)^{3/2} dx$$

input `integrate(x^2*(c/(b*x^2+a))^(3/2),x, algorithm="maxima")`

output `integrate(x^2*(c/(b*x^2 + a))^(3/2), x)`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.92

$$\int x^2 \left(\frac{c}{a + bx^2} \right)^{3/2} dx = - \left(\frac{cx \operatorname{sgn}(bx^2 + a)}{\sqrt{bcx^2 + ac}} + \frac{c \log \left(\left| -\sqrt{bc}x + \sqrt{bcx^2 + ac} \right| \operatorname{sgn}(bx^2 + a) \right)}{\sqrt{bc}b} \right) c$$

input `integrate(x^2*(c/(b*x^2+a))^(3/2),x, algorithm="giac")`

output `-(c*x*sgn(b*x^2 + a)/(sqrt(b*c*x^2 + a*c)*b) + c*log(abs(-sqrt(b*c)*x + sqrt(b*c*x^2 + a*c)))*sgn(b*x^2 + a)/(sqrt(b*c)*b))*c`

Mupad [F(-1)]

Timed out.

$$\int x^2 \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \int x^2 \left(\frac{c}{bx^2 + a} \right)^{3/2} dx$$

input `int(x^2*(c/(a + b*x^2))^(3/2),x)`

output `int(x^2*(c/(a + b*x^2))^(3/2), x)`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.18

$$\int x^2 \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \frac{\sqrt{c} c \left(-\sqrt{bx^2 + a} bx + \sqrt{b} \log \left(\frac{\sqrt{bx^2 + a} + \sqrt{bx}}{\sqrt{a}} \right) a + \sqrt{b} \log \left(\frac{\sqrt{bx^2 + a} + \sqrt{bx}}{\sqrt{a}} \right) bx^2 - \sqrt{b} \right)}{b^2 (bx^2 + a)}$$

input `int(x^2*(c/(b*x^2+a))^(3/2),x)`output `(sqrt(c)*c*(- sqrt(a + b*x**2)*b*x + sqrt(b)*log((sqrt(a + b*x**2) + sqrt(b)*x)/sqrt(a))*a + sqrt(b)*log((sqrt(a + b*x**2) + sqrt(b)*x)/sqrt(a))*b*x**2 - sqrt(b)*a - sqrt(b)*b*x**2)/(b**2*(a + b*x**2))`

$$3.26 \quad \int \left(\frac{c}{a+bx^2} \right)^{3/2} dx$$

Optimal result	217
Mathematica [A] (verified)	217
Rubi [A] (verified)	218
Maple [A] (verified)	219
Fricas [A] (verification not implemented)	219
Sympy [B] (verification not implemented)	220
Maxima [F]	220
Giac [A] (verification not implemented)	220
Mupad [B] (verification not implemented)	221
Reduce [B] (verification not implemented)	221

Optimal result

Integrand size = 15, antiderivative size = 21

$$\int \left(\frac{c}{a+bx^2} \right)^{3/2} dx = \frac{cx \sqrt{\frac{c}{a+bx^2}}}{a}$$

output

```
c*x*(c/(b*x^2+a))^(1/2)/a
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \left(\frac{c}{a+bx^2} \right)^{3/2} dx = \frac{cx \sqrt{\frac{c}{a+bx^2}}}{a}$$

input

```
Integrate[(c/(a + b*x^2))^(3/2),x]
```

output

```
(c*x*Sqrt[c/(a + b*x^2)])/a
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2045, 208}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\frac{c}{a + bx^2} \right)^{3/2} dx$$

$$\downarrow \text{2045}$$

$$\frac{c\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{c}{a+bx^2}} \int \frac{1}{\left(\frac{bx^2}{a} + 1\right)^{3/2}} dx}{a}$$

$$\downarrow \text{208}$$

$$\frac{cx\sqrt{\frac{c}{a+bx^2}}}{a}$$

input `Int[(c/(a + b*x^2))^(3/2),x]`

output `(c*x*Sqrt[c/(a + b*x^2)])/a`

Defintions of rubi rules used

rule 208

```
Int[((a_) + (b_.)*(x_)^2)^(-3/2), x_Symbol] := Simp[x/(a*Sqrt[a + b*x^2]),
x] /; FreeQ[{a, b}, x]
```

rule 2045

```
Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_.))^(p_), x_Symbol] := Simp[Si
mp[(c*(a + b*x^n)^q]^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q)
, x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]
```

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.95

method	result	size
trager	$\frac{cx\sqrt{\frac{c}{bx^2+a}}}{a}$	20
gospers	$\frac{(bx^2+a)x\left(\frac{c}{bx^2+a}\right)^{\frac{3}{2}}}{a}$	26
default	$\frac{(bx^2+a)x\left(\frac{c}{bx^2+a}\right)^{\frac{3}{2}}}{a}$	26
orering	$\frac{(bx^2+a)x\left(\frac{c}{bx^2+a}\right)^{\frac{3}{2}}}{a}$	26

input `int((c/(b*x^2+a))^(3/2),x,method=_RETURNVERBOSE)`

output `c*x*(c/(b*x^2+a))^(1/2)/a`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \left(\frac{c}{a+bx^2}\right)^{3/2} dx = \frac{cx\sqrt{\frac{c}{bx^2+a}}}{a}$$

input `integrate((c/(b*x^2+a))^(3/2),x, algorithm="fricas")`

output `c*x*sqrt(c/(b*x^2 + a))/a`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 46 vs. $2(15) = 30$.

Time = 0.29 (sec) , antiderivative size = 46, normalized size of antiderivative = 2.19

$$\int \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \begin{cases} x \left(\frac{c}{a + bx^2} \right)^{3/2} + \frac{bx^3 \left(\frac{c}{a + bx^2} \right)^{3/2}}{a} & \text{for } a \neq 0 \\ -\frac{x \left(\frac{c}{bx^2} \right)^{3/2}}{2} & \text{otherwise} \end{cases}$$

input `integrate((c/(b*x**2+a))**(3/2),x)`

output `Piecewise((x*(c/(a + b*x**2))**(3/2) + b*x**3*(c/(a + b*x**2))**(3/2)/a, Ne(a, 0)), (-x*(c/(b*x**2))**(3/2)/2, True))`

Maxima [F]

$$\int \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \int \left(\frac{c}{bx^2 + a} \right)^{3/2} dx$$

input `integrate((c/(b*x^2+a))^(3/2),x, algorithm="maxima")`

output `integrate((c/(b*x^2 + a))^(3/2), x)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.33

$$\int \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \frac{c^2 x \operatorname{sgn}(bx^2 + a)}{\sqrt{bcx^2 + aca}}$$

input `integrate((c/(b*x^2+a))^(3/2),x, algorithm="giac")`

output `c^2*x*sgn(b*x^2 + a)/(sqrt(b*c*x^2 + a*c)*a)`

Mupad [B] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \frac{cx \sqrt{\frac{c}{bx^2+a}}}{a}$$

input `int((c/(a + b*x^2))^(3/2),x)`

output `(c*x*(c/(a + b*x^2))^(1/2))/a`

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 42, normalized size of antiderivative = 2.00

$$\int \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \frac{\sqrt{c}c(\sqrt{bx^2+a}bx + \sqrt{b}a + \sqrt{b}bx^2)}{ab(bx^2+a)}$$

input `int((c/(b*x^2+a))^(3/2),x)`

output `(sqrt(c)*c*(sqrt(a + b*x**2)*b*x + sqrt(b)*a + sqrt(b)*b*x**2))/(a*b*(a + b*x**2))`

$$3.27 \quad \int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^2} dx$$

Optimal result	222
Mathematica [A] (verified)	222
Rubi [A] (verified)	223
Maple [A] (verified)	224
Fricas [A] (verification not implemented)	225
Sympy [F]	225
Maxima [A] (verification not implemented)	225
Giac [A] (verification not implemented)	226
Mupad [B] (verification not implemented)	226
Reduce [B] (verification not implemented)	226

Optimal result

Integrand size = 19, antiderivative size = 48

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^2} dx = -\frac{c\sqrt{\frac{c}{a+bx^2}}}{ax} - \frac{2bcx\sqrt{\frac{c}{a+bx^2}}}{a^2}$$

output `-c*(c/(b*x^2+a))^(1/2)/a/x-2*b*c*x*(c/(b*x^2+a))^(1/2)/a^2`

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.67

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^2} dx = -\frac{c\sqrt{\frac{c}{a+bx^2}}(a+2bx^2)}{a^2x}$$

input `Integrate[(c/(a + b*x^2))^(3/2)/x^2,x]`

output `-((c*Sqrt[c/(a + b*x^2)]*(a + 2*b*x^2))/(a^2*x))`

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.56, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2045, 245, 208}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^2} dx \\
 \downarrow \text{2045} \\
 \frac{c\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{c}{a+bx^2}}}{a} \int \frac{1}{x^2\left(\frac{bx^2}{a} + 1\right)^{3/2}} dx \\
 \downarrow \text{245} \\
 \frac{c\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{c}{a+bx^2}}}{a} \left(-\frac{2b \int \frac{1}{\left(\frac{bx^2}{a} + 1\right)^{3/2}} dx}{a} - \frac{1}{x\sqrt{\frac{bx^2}{a} + 1}} \right) \\
 \downarrow \text{208} \\
 \frac{c\sqrt{\frac{bx^2}{a} + 1} \left(-\frac{2bx}{a\sqrt{\frac{bx^2}{a} + 1}} - \frac{1}{x\sqrt{\frac{bx^2}{a} + 1}} \right) \sqrt{\frac{c}{a+bx^2}}}{a}
 \end{array}$$

input `Int[(c/(a + b*x^2))^(3/2)/x^2,x]`

output `(c*Sqrt[c/(a + b*x^2)]*Sqrt[1 + (b*x^2)/a]*(-(1/(x*Sqrt[1 + (b*x^2)/a])) - (2*b*x)/(a*Sqrt[1 + (b*x^2)/a]))) / a`

Definitions of rubi rules used

rule 208 $\text{Int}[(a_ + (b_ \cdot)(x_)^2)^{-3/2}, x_Symbol] \rightarrow \text{Simp}[x/(a\sqrt{a + b \cdot x^2}), x] \text{ /; FreeQ}\{a, b\}, x]$

rule 245 $\text{Int}[(x_)^{(m_)} \cdot ((a_ + (b_ \cdot)(x_)^2)^{(p_)}), x_Symbol] \rightarrow \text{Simp}[x^{(m + 1)} \cdot ((a + b \cdot x^2)^{(p + 1)} / (a \cdot (m + 1))), x] - \text{Simp}[b \cdot ((m + 2) \cdot (p + 1) + 1) / (a \cdot (m + 1)) \cdot \text{Int}[x^{(m + 2)} \cdot (a + b \cdot x^2)^p, x], x] \text{ /; FreeQ}\{a, b, m, p\}, x] \&\& \text{ILtQ}[\text{Simplify}[(m + 1)/2 + p + 1], 0] \&\& \text{NeQ}[m, -1]$

rule 2045 $\text{Int}[(u_) \cdot ((c_) \cdot ((a_ + (b_ \cdot)(x_)^{(n_)})^{(q_)})^{(p_)}), x_Symbol] \rightarrow \text{Simp}[\text{Simp}[(c \cdot (a + b \cdot x^n)^q)^p / (1 + b \cdot (x^n/a)^{(p \cdot q))} \cdot \text{Int}[u \cdot (1 + b \cdot (x^n/a)^{(p \cdot q))}, x], x] \text{ /; FreeQ}\{a, b, c, n, p, q\}, x] \&\& !\text{GeQ}[a, 0]$

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.77

method	result	size
gospers	$-\frac{(bx^2+a)(2bx^2+a)\left(\frac{c}{bx^2+a}\right)^{\frac{3}{2}}}{a^2x}$	37
default	$-\frac{(bx^2+a)(2bx^2+a)\left(\frac{c}{bx^2+a}\right)^{\frac{3}{2}}}{a^2x}$	37
orering	$-\frac{(bx^2+a)(2bx^2+a)\left(\frac{c}{bx^2+a}\right)^{\frac{3}{2}}}{a^2x}$	37
trager	$-\frac{(ac+bc)(2bx^2+a)\sqrt{\frac{c}{bx^2+a}}}{a^2x(a+b)}$	42
risch	$-\frac{(bx^2+a)c\sqrt{\frac{c}{bx^2+a}}}{a^2x} - \frac{bcx\sqrt{\frac{c}{bx^2+a}}}{a^2}$	52

input $\text{int}((c/(b \cdot x^2+a))^{3/2}/x^2,x,\text{method}=_RETURNVERBOSE)$

output $-(b \cdot x^2+a) \cdot (2 \cdot b \cdot x^2+a) \cdot (c/(b \cdot x^2+a))^{3/2}/a^2/x$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.67

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^2} dx = -\frac{(2bcx^2 + ac)\sqrt{\frac{c}{bx^2+a}}}{a^2x}$$

input `integrate((c/(b*x^2+a))^(3/2)/x^2,x, algorithm="fricas")`output `-(2*b*c*x^2 + a*c)*sqrt(c/(b*x^2 + a))/(a^2*x)`**Sympy [F]**

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^2} dx = \int \frac{\left(\frac{c}{a+bx^2}\right)^{\frac{3}{2}}}{x^2} dx$$

input `integrate((c/(b*x**2+a))**(3/2)/x**2,x)`output `Integral((c/(a + b*x**2))**(3/2)/x**2, x)`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.96

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^2} dx = -\frac{2b^2c^{\frac{3}{2}}x^4 + 3abc^{\frac{3}{2}}x^2 + a^2c^{\frac{3}{2}}}{(bx^2 + a)^{\frac{3}{2}}a^2x}$$

input `integrate((c/(b*x^2+a))^(3/2)/x^2,x, algorithm="maxima")`output `-(2*b^2*c^(3/2)*x^4 + 3*a*b*c^(3/2)*x^2 + a^2*c^(3/2))/((b*x^2 + a)^(3/2)*a^2*x)`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.69

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^2} dx = - \left(\frac{bcx \operatorname{sgn}(bx^2 + a)}{\sqrt{bcx^2 + aca^2}} - \frac{2\sqrt{bcc} \operatorname{sgn}(bx^2 + a)}{\left(\left(\sqrt{bcx} - \sqrt{bcx^2 + ac}\right)^2 - ac\right)a} \right) c$$

input `integrate((c/(b*x^2+a))^(3/2)/x^2,x, algorithm="giac")`output `-(b*c*x*sgn(b*x^2 + a)/(sqrt(b*c*x^2 + a*c)*a^2) - 2*sqrt(b*c)*c*sgn(b*x^2 + a)/(((sqrt(b*c)*x - sqrt(b*c*x^2 + a*c))^2 - a*c)*a)*c`**Mupad [B] (verification not implemented)**

Time = 0.49 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.12

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^2} dx = - \frac{\left(\frac{bc}{a} + \frac{2b^2cx^2}{a^2}\right) \sqrt{\frac{c}{bx^2+a}} \left(\frac{a}{b} + x^2\right)}{bx^3 + ax}$$

input `int((c/(a + b*x^2))^(3/2)/x^2,x)`output `-(((b*c)/a + (2*b^2*c*x^2)/a^2)*(c/(a + b*x^2))^(1/2)*(a/b + x^2))/(a*x + b*x^3)`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.23

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^2} dx = \frac{\sqrt{c}c\left(-\sqrt{bx^2+a}a - 2\sqrt{bx^2+a}bx^2 - 2\sqrt{b}ax - 2\sqrt{b}bx^3\right)}{a^2x(bx^2+a)}$$

input `int((c/(b*x^2+a))^(3/2)/x^2,x)`

output
$$\frac{(\sqrt{c})c(-\sqrt{a+bx^2})a - 2\sqrt{a+bx^2}bx^2 - 2\sqrt{b}ax - 2\sqrt{b}bx^3}{a^2x(a+bx^2)}$$

3.28 $\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^4} dx$

Optimal result	228
Mathematica [A] (verified)	228
Rubi [A] (verified)	229
Maple [A] (verified)	230
Fricas [A] (verification not implemented)	231
Sympy [F]	231
Maxima [A] (verification not implemented)	231
Giac [B] (verification not implemented)	232
Mupad [B] (verification not implemented)	232
Reduce [B] (verification not implemented)	233

Optimal result

Integrand size = 19, antiderivative size = 81

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^4} dx = -\frac{c\sqrt{\frac{c}{a+bx^2}}}{3ax^3} + \frac{4bc\sqrt{\frac{c}{a+bx^2}}}{3a^2x} + \frac{8b^2cx\sqrt{\frac{c}{a+bx^2}}}{3a^3}$$

output

```
-1/3*c*(c/(b*x^2+a))^(1/2)/a/x^3+4/3*b*c*(c/(b*x^2+a))^(1/2)/a^2/x+8/3*b^2*c*x*(c/(b*x^2+a))^(1/2)/a^3
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.56

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^4} dx = -\frac{c\sqrt{\frac{c}{a+bx^2}}(a^2 - 4abx^2 - 8b^2x^4)}{3a^3x^3}$$

input

```
Integrate[(c/(a + b*x^2))^(3/2)/x^4,x]
```

output

```
-1/3*(c*Sqrt[c/(a + b*x^2)]*(a^2 - 4*a*b*x^2 - 8*b^2*x^4))/(a^3*x^3)
```

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.30, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {2045, 245, 245, 208}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^4} dx \\
 \downarrow \text{2045} \\
 \frac{c\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{c}{a+bx^2}} \int \frac{1}{x^4\left(\frac{bx^2}{a} + 1\right)^{3/2}} dx}{a} \\
 \downarrow \text{245} \\
 \frac{c\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{c}{a+bx^2}} \left(-\frac{4b \int \frac{1}{x^2\left(\frac{bx^2}{a} + 1\right)^{3/2}} dx}{3a} - \frac{1}{3x^3\sqrt{\frac{bx^2}{a} + 1}} \right)}{a} \\
 \downarrow \text{245} \\
 \frac{c\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{c}{a+bx^2}} \left(-\frac{4b \left(-\frac{2b \int \frac{1}{\left(\frac{bx^2}{a} + 1\right)^{3/2}} dx}{a} - \frac{1}{x\sqrt{\frac{bx^2}{a} + 1}} \right)}{3a} - \frac{1}{3x^3\sqrt{\frac{bx^2}{a} + 1}} \right)}{a} \\
 \downarrow \text{208} \\
 \frac{c\sqrt{\frac{bx^2}{a} + 1} \left(-\frac{4b \left(-\frac{2bx}{a\sqrt{\frac{bx^2}{a} + 1}} - \frac{1}{x\sqrt{\frac{bx^2}{a} + 1}} \right)}{3a} - \frac{1}{3x^3\sqrt{\frac{bx^2}{a} + 1}} \right) \sqrt{\frac{c}{a+bx^2}}}{a}
 \end{array}$$

input

```
Int[(c/(a + b*x^2))^(3/2)/x^4,x]
```

output
$$\frac{(c\sqrt{c/(a + bx^2)}\sqrt{1 + (bx^2)/a}*(-1/3*1/(x^3\sqrt{1 + (bx^2)/a}) - (4*b*(-1/(x*\sqrt{1 + (bx^2)/a})) - (2*b*x)/(a*\sqrt{1 + (bx^2)/a}))/ (3*a)))/a$$

Defintions of rubi rules used

rule 208
$$\text{Int}[(a_ + (b_)*(x_)^2)^{-3/2}, x_Symbol] \rightarrow \text{Simp}[x/(a*\sqrt{a + b*x^2}), x] \text{ /; FreeQ}\{a, b\}, x]$$

rule 245
$$\text{Int}[(x_)^{(m_)}*((a_ + (b_)*(x_)^2)^{(p_)}), x_Symbol] \rightarrow \text{Simp}[x^{(m + 1)}*((a + b*x^2)^{(p + 1)/(a*(m + 1))}), x] - \text{Simp}[b*(m + 2*(p + 1) + 1)/(a*(m + 1)) \text{ Int}[x^{(m + 2)}*(a + b*x^2)^p, x], x] \text{ /; FreeQ}\{a, b, m, p\}, x] \&\& \text{ILtQ}[\text{Simplify}[(m + 1)/2 + p + 1], 0] \&\& \text{NeQ}[m, -1]$$

rule 2045
$$\text{Int}[(u_)*((c_)*((a_ + (b_)*(x_)^{(n_)}))^{(q_)})^{(p_)}), x_Symbol] \rightarrow \text{Simp}[\text{Simp}[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^{(p*q)} \text{ Int}[u*(1 + b*(x^n/a))^{(p*q)}, x], x] \text{ /; FreeQ}\{a, b, c, n, p, q\}, x] \&\& !\text{GeQ}[a, 0]$$

Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.59

method	result	size
gospers	$-\frac{(bx^2+a)(-8b^2x^4-4abx^2+a^2)\left(\frac{c}{bx^2+a}\right)^{\frac{3}{2}}}{3a^3x^3}$	48
default	$-\frac{(bx^2+a)(-8b^2x^4-4abx^2+a^2)\left(\frac{c}{bx^2+a}\right)^{\frac{3}{2}}}{3a^3x^3}$	48
orering	$-\frac{(bx^2+a)(-8b^2x^4-4abx^2+a^2)\left(\frac{c}{bx^2+a}\right)^{\frac{3}{2}}}{3a^3x^3}$	48
trager	$-\frac{(ac+bc)(-8b^2x^4-4abx^2+a^2)\sqrt{\frac{c}{bx^2+a}}}{3(a+b)a^3x^3}$	53
risch	$-\frac{(bx^2+a)(-5bx^2+a)c\sqrt{\frac{c}{bx^2+a}}}{3a^3x^3} + \frac{b^2cx\sqrt{\frac{c}{bx^2+a}}}{a^3}$	61

input
$$\text{int}((c/(b*x^2+a))^{3/2}/x^4,x,\text{method}=_RETURNVERBOSE)$$

output $-1/3*(b*x^2+a)*(-8*b^2*x^4-4*a*b*x^2+a^2)*(c/(b*x^2+a))^{3/2}/a^3/x^3$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.56

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^4} dx = \frac{(8b^2cx^4 + 4abcx^2 - a^2c)\sqrt{\frac{c}{bx^2+a}}}{3a^3x^3}$$

input `integrate((c/(b*x^2+a))^(3/2)/x^4,x, algorithm="fricas")`

output $1/3*(8*b^2*c*x^4 + 4*a*b*c*x^2 - a^2*c)*\text{sqrt}(c/(b*x^2 + a))/(a^3*x^3)$

Sympy [F]

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^4} dx = \int \frac{\left(\frac{c}{a+bx^2}\right)^{\frac{3}{2}}}{x^4} dx$$

input `integrate((c/(b*x**2+a))**(3/2)/x**4,x)`

output `Integral((c/(a + b*x**2))**(3/2)/x**4, x)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.75

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^4} dx = \frac{8b^3c^{\frac{3}{2}}x^6 + 12ab^2c^{\frac{3}{2}}x^4 + 3a^2bc^{\frac{3}{2}}x^2 - a^3c^{\frac{3}{2}}}{3(bx^2 + a)^{\frac{3}{2}}a^3x^3}$$

input `integrate((c/(b*x^2+a))^(3/2)/x^4,x, algorithm="maxima")`

output

$$\frac{1}{3} \frac{(8b^3c^{3/2}x^6 + 12a^2b^2c^{3/2}x^4 + 3a^3b^2c^{3/2}x^2 - a^3c^{3/2})}{(bx^2 + a)^{3/2}a^3x^3}$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 178 vs. $2(69) = 138$.

Time = 0.13 (sec) , antiderivative size = 178, normalized size of antiderivative = 2.20

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^4} dx = \frac{1}{3} \left(\frac{3b^2cx \operatorname{sgn}(bx^2 + a)}{\sqrt{bcx^2 + aca^3}} - \frac{2 \left(3\sqrt{bc}(\sqrt{bcx} - \sqrt{bcx^2 + ac})^4 bcs \operatorname{sgn}(bx^2 + a) - 12\sqrt{bc}(\sqrt{bcx} - \sqrt{bcx^2 + ac})^3 \right)}{\left((\sqrt{bcx} - \sqrt{bcx^2 + ac})^2 - a^3 \right)} \right)$$

input

```
integrate((c/(b*x^2+a))^(3/2)/x^4,x, algorithm="giac")
```

output

$$\frac{1}{3} \frac{(3b^2cx \operatorname{sgn}(bx^2 + a) / (\sqrt{bcx^2 + aca^3}) - 2(3\sqrt{bc}(\sqrt{bcx} - \sqrt{bcx^2 + ac})^4 bcs \operatorname{sgn}(bx^2 + a) - 12\sqrt{bc}(\sqrt{bcx} - \sqrt{bcx^2 + ac})^3) / ((\sqrt{bcx} - \sqrt{bcx^2 + ac})^2 - a^3))}{a^3}$$

Mupad [B] (verification not implemented)

Time = 0.55 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.53

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^4} dx = \frac{c \sqrt{\frac{c}{bx^2+a}} (-a^2 + 4abx^2 + 8b^2x^4)}{3a^3x^3}$$

input

```
int((c/(a + b*x^2))^(3/2)/x^4,x)
```

output

$$(c*(c/(a + b*x^2))^(1/2)*(8*b^2*x^4 - a^2 + 4*a*b*x^2))/(3*a^3*x^3)$$

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.04

$$\int \frac{\left(\frac{c}{a+bx^2}\right)^{3/2}}{x^4} dx = \frac{\sqrt{c}c\left(-\sqrt{bx^2+a}a^2 + 4\sqrt{bx^2+a}abx^2 + 8\sqrt{bx^2+a}b^2x^4 - 8\sqrt{b}abx^3 - 8\sqrt{b}b^2x^5\right)}{3a^3x^3(bx^2+a)}$$

input `int((c/(b*x^2+a))^(3/2)/x^4,x)`

output `(sqrt(c)*c*(-sqrt(a+b*x**2)*a**2+4*sqrt(a+b*x**2)*a*b*x**2+8*sqrt(a+b*x**2)*b**2*x**4-8*sqrt(b)*a*b*x**3-8*sqrt(b)*b**2*x**5))/(3*a**3*x**3*(a+b*x**2))`

$$3.29 \quad \int x^5 \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx$$

Optimal result	234
Mathematica [A] (verified)	234
Rubi [A] (verified)	235
Maple [A] (verified)	236
Fricas [A] (verification not implemented)	237
Sympy [F]	237
Maxima [A] (verification not implemented)	238
Giac [A] (verification not implemented)	238
Mupad [F(-1)]	238
Reduce [B] (verification not implemented)	239

Optimal result

Integrand size = 19, antiderivative size = 95

$$\int x^5 \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx = \frac{ac \sqrt{\frac{c}{(a+bx^2)^2}}}{b^3} - \frac{a^2 c \sqrt{\frac{c}{(a+bx^2)^2}}}{4b^3 (a+bx^2)} + \frac{c \sqrt{\frac{c}{(a+bx^2)^2}} (a+bx^2) \log(a+bx^2)}{2b^3}$$

output

```
a*c*(c/(b*x^2+a)^2)^(1/2)/b^3-1/4*a^2*c*(c/(b*x^2+a)^2)^(1/2)/b^3/(b*x^2+a)
)+1/2*c*(c/(b*x^2+a)^2)^(1/2)*(b*x^2+a)*ln(b*x^2+a)/b^3
```

Mathematica [A] (verified)

Time = 1.03 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.64

$$\int x^5 \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx = \frac{\left(\frac{c}{(a+bx^2)^2} \right)^{3/2} (a+bx^2) \left(a(3a+4bx^2) + 2(a+bx^2)^2 \log(a+bx^2) \right)}{4b^3}$$

input

```
Integrate[x^5*(c/(a + b*x^2)^2)^(3/2),x]
```

output

$$\left(\frac{c}{(a + bx^2)^2} \right)^{3/2} (a + bx^2) (a(3a + 4bx^2) + 2(a + bx^2)^2 \operatorname{Log}[a + bx^2]) / (4b^3)$$
Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.78, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {2045, 27, 243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^5 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx \\ & \quad \downarrow \text{2045} \\ & \frac{c(a + bx^2) \sqrt{\frac{c}{(a + bx^2)^2}} \int \frac{a^3 x^5}{(bx^2 + a)^3} dx}{a^3} \\ & \quad \downarrow \text{27} \\ & c(a + bx^2) \sqrt{\frac{c}{(a + bx^2)^2}} \int \frac{x^5}{(bx^2 + a)^3} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{2} c(a + bx^2) \sqrt{\frac{c}{(a + bx^2)^2}} \int \frac{x^4}{(bx^2 + a)^3} dx \\ & \quad \downarrow \text{49} \\ & \frac{1}{2} c(a + bx^2) \sqrt{\frac{c}{(a + bx^2)^2}} \int \left(\frac{a^2}{b^2 (bx^2 + a)^3} - \frac{2a}{b^2 (bx^2 + a)^2} + \frac{1}{b^2 (bx^2 + a)} \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2} c(a + bx^2) \sqrt{\frac{c}{(a + bx^2)^2}} \left(-\frac{a^2}{2b^3 (a + bx^2)^2} + \frac{2a}{b^3 (a + bx^2)} + \frac{\log(a + bx^2)}{b^3} \right) \end{aligned}$$

input

$$\operatorname{Int}[x^5 (c/(a + bx^2)^2)^{3/2}, x]$$

output
$$\frac{(c\sqrt{c/(a + bx^2)^2}*(a + bx^2)*(-1/2*a^2/(b^3*(a + bx^2)^2) + (2*a)/(b^3*(a + bx^2))) + \text{Log}[a + bx^2]/b^3)/2}$$

Defintions of rubi rules used

rule 27
$$\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_)*(Gx_)] \text{ ; FreeQ}[b, x]$$

rule 49
$$\text{Int}[(a_.) + (b_.)*(x_)^(m_.)*((c_.) + (d_.)*(x_)^(n_.), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + bx)^m*(c + dx)^n, x], x] \text{ ; FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{IGtQ}[m + n + 2, 0]$$

rule 243
$$\text{Int}[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{(m-1)/2}*(a + bx)^p, x], x, x^2], x] \text{ ; FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$$

rule 2009
$$\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ ; SumQ}[u]$$

rule 2045
$$\text{Int}[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_))^(p_), x_Symbol] \rightarrow \text{Simp}[\text{Simp}[(c*(a + bx^n)^q)^p/(1 + b*(x^n/a))^(p*q)] \text{ Int}[u*(1 + b*(x^n/a))^(p*q), x], x] \text{ ; FreeQ}[\{a, b, c, n, p, q\}, x] \ \&\& \ !\text{GeQ}[a, 0]$$

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.81

method	result	size
risch	$\frac{c\sqrt{\frac{c}{(bx^2+a)^2}}\left(\frac{ax^2}{b^2} + \frac{3a^2}{4b^3}\right)}{bx^2+a} + \frac{c\sqrt{\frac{c}{(bx^2+a)^2}}(bx^2+a)\ln(bx^2+a)}{2b^3}$	77
default	$\frac{(2\ln(bx^2+a)b^2x^4 + 4\ln(bx^2+a)abx^2 + 4abx^2 + 2\ln(bx^2+a)a^2 + 3a^2)(bx^2+a)\left(\frac{c}{(bx^2+a)^2}\right)^{\frac{3}{2}}}{4b^3}$	83

input `int(x^5*(c/(b*x^2+a)^2)^(3/2),x,method=_RETURNVERBOSE)`

output `c*(c/(b*x^2+a)^2)^(1/2)/(b*x^2+a)*(a*x^2/b^2+3/4*a^2/b^3)+1/2*c*(c/(b*x^2+a)^2)^(1/2)*(b*x^2+a)*ln(b*x^2+a)/b^3`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.93

$$\int x^5 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \frac{(4abcx^2 + 3a^2c + 2(b^2cx^4 + 2abcx^2 + a^2c) \log(bx^2 + a)) \sqrt{\frac{c}{b^2x^4 + 2abx^2 + a^2}}}{4(b^4x^2 + ab^3)}$$

input `integrate(x^5*(c/(b*x^2+a)^2)^(3/2),x, algorithm="fricas")`

output `1/4*(4*a*b*c*x^2 + 3*a^2*c + 2*(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)*log(b*x^2 + a))*sqrt(c/(b^2*x^4 + 2*a*b*x^2 + a^2))/(b^4*x^2 + a*b^3)`

Sympy [F]

$$\int x^5 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \int x^5 \left(\frac{c}{a^2 + 2abx^2 + b^2x^4} \right)^{\frac{3}{2}} dx$$

input `integrate(x**5*(c/(b*x**2+a)**2)**(3/2),x)`

output `Integral(x**5*(c/(a**2 + 2*a*b*x**2 + b**2*x**4))**(3/2), x)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.67

$$\int x^5 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \frac{4abc^{\frac{3}{2}}x^2 + 3a^2c^{\frac{3}{2}}}{4(b^5x^4 + 2ab^4x^2 + a^2b^3)} + \frac{c^{\frac{3}{2}} \log(bx^2 + a)}{2b^3}$$

input `integrate(x^5*(c/(b*x^2+a)^2)^(3/2),x, algorithm="maxima")`output `1/4*(4*a*b*c^(3/2)*x^2 + 3*a^2*c^(3/2))/(b^5*x^4 + 2*a*b^4*x^2 + a^2*b^3) + 1/2*c^(3/2)*log(b*x^2 + a)/b^3`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.75

$$\int x^5 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \frac{1}{4} c^{\frac{3}{2}} \left(\frac{2 \log(|bx^2 + a|) \operatorname{sgn}(bx^2 + a)}{b^3} - \frac{3bx^4 \operatorname{sgn}(bx^2 + a) + 2ax^2 \operatorname{sgn}(bx^2 + a)}{(bx^2 + a)^2 b^2} \right)$$

input `integrate(x^5*(c/(b*x^2+a)^2)^(3/2),x, algorithm="giac")`output `1/4*c^(3/2)*(2*log(abs(b*x^2 + a))*sgn(b*x^2 + a)/b^3 - (3*b*x^4*sgn(b*x^2 + a) + 2*a*x^2*sgn(b*x^2 + a))/((b*x^2 + a)^2*b^2))`**Mupad [F(-1)]**

Timed out.

$$\int x^5 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \int x^5 \left(\frac{c}{(bx^2 + a)^2} \right)^{3/2} dx$$

input `int(x^5*(c/(a + b*x^2)^2)^(3/2),x)`output `int(x^5*(c/(a + b*x^2)^2)^(3/2), x)`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.88

$$\int x^5 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \frac{\sqrt{c} c (2 \log(bx^2 + a) a^2 + 4 \log(bx^2 + a) abx^2 + 2 \log(bx^2 + a) b^2 x^4 + a^2 - 2b^2 x^4)}{4b^3 (b^2 x^4 + 2abx^2 + a^2)}$$

input `int(x^5*(c/(b*x^2+a)^2)^(3/2),x)`

output `(sqrt(c)*c*(2*log(a + b*x**2)*a**2 + 4*log(a + b*x**2)*a*b*x**2 + 2*log(a + b*x**2)*b**2*x**4 + a**2 - 2*b**2*x**4))/(4*b**3*(a**2 + 2*a*b*x**2 + b**2*x**4))`

$$3.30 \quad \int x^3 \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx$$

Optimal result	240
Mathematica [A] (verified)	240
Rubi [A] (verified)	241
Maple [A] (verified)	242
Fricas [A] (verification not implemented)	243
Sympy [F]	243
Maxima [A] (verification not implemented)	243
Giac [A] (verification not implemented)	244
Mupad [B] (verification not implemented)	244
Reduce [B] (verification not implemented)	244

Optimal result

Integrand size = 19, antiderivative size = 35

$$\int x^3 \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx = \frac{cx^4 \sqrt{\frac{c}{(a+bx^2)^2}}}{4a(a+bx^2)}$$

output `1/4*c*x^4*(c/(b*x^2+a)^2)^(1/2)/a/(b*x^2+a)`

Mathematica [A] (verified)

Time = 1.01 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.06

$$\int x^3 \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx = -\frac{\left(\frac{c}{(a+bx^2)^2} \right)^{3/2} (a+bx^2)(a+2bx^2)}{4b^2}$$

input `Integrate[x^3*(c/(a + b*x^2)^2)^(3/2),x]`

output `-1/4*((c/(a + b*x^2)^2)^(3/2)*(a + b*x^2)*(a + 2*b*x^2))/b^2`

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2045, 27, 242}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx$$

$$\downarrow \text{2045}$$

$$\frac{c(a + bx^2) \sqrt{\frac{c}{(a+bx^2)^2}} \int \frac{a^3 x^3}{(bx^2+a)^3} dx}{a^3}$$

$$\downarrow \text{27}$$

$$c(a + bx^2) \sqrt{\frac{c}{(a + bx^2)^2}} \int \frac{x^3}{(bx^2 + a)^3} dx$$

$$\downarrow \text{242}$$

$$\frac{cx^4 \sqrt{\frac{c}{(a+bx^2)^2}}}{4a(a + bx^2)}$$

input `Int[x^3*(c/(a + b*x^2)^2)^(3/2),x]`

output `(c*x^4*Sqrt[c/(a + b*x^2)^2])/(4*a*(a + b*x^2))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] :> Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 242 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, p}, x] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.97

method	result	size
gospers	$-\frac{(bx^2+a)(2bx^2+a)\left(\frac{c}{(bx^2+a)^2}\right)^{\frac{3}{2}}}{4b^2}$	34
default	$-\frac{(bx^2+a)(2bx^2+a)\left(\frac{c}{(bx^2+a)^2}\right)^{\frac{3}{2}}}{4b^2}$	34
orering	$-\frac{(bx^2+a)(2bx^2+a)\left(\frac{c}{(bx^2+a)^2}\right)^{\frac{3}{2}}}{4b^2}$	34
risch	$\frac{c\sqrt{\frac{c}{(bx^2+a)^2}}\left(-\frac{x^2}{2b}-\frac{a}{4b^2}\right)}{bx^2+a}$	40
trager	$\frac{cx^4\sqrt{\frac{c}{b^2x^4+2abx^2+a^2}}}{4a(bx^2+a)}$	43

input `int(x^3*(c/(b*x^2+a)^2)^(3/2),x,method=_RETURNVERBOSE)`

output `-1/4*(b*x^2+a)*(2*b*x^2+a)*(c/(b*x^2+a)^2)^(3/2)/b^2`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.49

$$\int x^3 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = -\frac{(2bcx^2 + ac) \sqrt{\frac{c}{b^2x^4 + 2abx^2 + a^2}}}{4(b^3x^2 + ab^2)}$$

input `integrate(x^3*(c/(b*x^2+a)^2)^(3/2),x, algorithm="fricas")`output `-1/4*(2*b*c*x^2 + a*c)*sqrt(c/(b^2*x^4 + 2*a*b*x^2 + a^2))/(b^3*x^2 + a*b^2)`**Sympy [F]**

$$\int x^3 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \int x^3 \left(\frac{c}{a^2 + 2abx^2 + b^2x^4} \right)^{\frac{3}{2}} dx$$

input `integrate(x**3*(c/(b*x**2+a)**2)**(3/2),x)`output `Integral(x**3*(c/(a**2 + 2*a*b*x**2 + b**2*x**4))**(3/2), x)`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.23

$$\int x^3 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = -\frac{2bc^{\frac{3}{2}}x^2 + ac^{\frac{3}{2}}}{4(b^4x^4 + 2ab^3x^2 + a^2b^2)}$$

input `integrate(x^3*(c/(b*x^2+a)^2)^(3/2),x, algorithm="maxima")`output `-1/4*(2*b*c^(3/2)*x^2 + a*c^(3/2))/(b^4*x^4 + 2*a*b^3*x^2 + a^2*b^2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int x^3 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = -\frac{(2bx^2 + a)c^{3/2} \operatorname{sgn}(bx^2 + a)}{4(bx^2 + a)^2 b^2}$$

input `integrate(x^3*(c/(b*x^2+a)^2)^(3/2),x, algorithm="giac")`output `-1/4*(2*b*x^2 + a)*c^(3/2)*sgn(b*x^2 + a)/((b*x^2 + a)^2*b^2)`**Mupad [B] (verification not implemented)**

Time = 0.48 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.03

$$\int x^3 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = -\frac{c(2bx^2 + a) \sqrt{\frac{c}{(bx^2+a)^2}}}{4b^2(bx^2 + a)}$$

input `int(x^3*(c/(a + b*x^2)^2)^(3/2),x)`output `-(c*(a + 2*b*x^2)*(c/(a + b*x^2)^2)^(1/2))/(4*b^2*(a + b*x^2))`**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.89

$$\int x^3 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \frac{\sqrt{c} c x^4}{4a(b^2 x^4 + 2abx^2 + a^2)}$$

input `int(x^3*(c/(b*x^2+a)^2)^(3/2),x)`output `(sqrt(c)*c*x**4)/(4*a*(a**2 + 2*a*b*x**2 + b**2*x**4))`

$$3.31 \quad \int x \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx$$

Optimal result	245
Mathematica [A] (verified)	245
Rubi [A] (verified)	246
Maple [A] (verified)	247
Fricas [A] (verification not implemented)	248
Sympy [F]	248
Maxima [A] (verification not implemented)	248
Giac [A] (verification not implemented)	249
Mupad [B] (verification not implemented)	249
Reduce [B] (verification not implemented)	249

Optimal result

Integrand size = 17, antiderivative size = 32

$$\int x \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx = -\frac{c\sqrt{\frac{c}{(a+bx^2)^2}}}{4b(a+bx^2)}$$

output `-1/4*c*(c/(b*x^2+a)^2)^(1/2)/b/(b*x^2+a)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.91

$$\int x \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx = -\frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2} (a+bx^2)}{4b}$$

input `Integrate[x*(c/(a + b*x^2)^2)^(3/2), x]`

output `-1/4*((c/(a + b*x^2)^2)^(3/2)*(a + b*x^2))/b`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.91, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2024, 20, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int x \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx \\
 \downarrow \text{2024} \\
 \frac{\int \left(\frac{c}{(bx^2+a)^2} \right)^{3/2} d(bx^2 + a)}{2b} \\
 \downarrow \text{20} \\
 \frac{(a + bx^2)^3 \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} \int \frac{1}{(bx^2+a)^3} d(bx^2 + a)}{2b} \\
 \downarrow \text{15} \\
 \frac{(a + bx^2) \left(\frac{c}{(a+bx^2)^2} \right)^{3/2}}{4b}
 \end{array}$$

input `Int[x*(c/(a + b*x^2)^2)^(3/2),x]`

output `-1/4*((c/(a + b*x^2)^2)^(3/2)*(a + b*x^2))/b`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 20 `Int[((a_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a*x^n)^p/x^(n*p) Int[x^(n*p), x], x] /; FreeQ[{a, n, p}, x] && !IntegerQ[p]`

rule 2024

```
Int[((a_.) + (b_.)*(Pq_)^(n_.))^(p_.)*(Qr_), x_Symbol] := With[{q = Expon[P
q, x], r = Expon[Qr, x]}, Simp[Coeff[Qr, x, r]/(q*Coeff[Pq, x, q]) Subst[
Int[(a + b*x^n)^p, x], x, Pq], x] /; EqQ[r, q - 1] && EqQ[Coeff[Qr, x, r]*D
[Pq, x], q*Coeff[Pq, x, q]*Qr]] /; FreeQ[{a, b, n, p}, x] && PolyQ[Pq, x] &
& PolyQ[Qr, x]
```

Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.81

method	result	size
gospers	$-\frac{(bx^2+a)\left(\frac{c}{(bx^2+a)^2}\right)^{\frac{3}{2}}}{4b}$	26
derivativedivides	$-\frac{(bx^2+a)\left(\frac{c}{(bx^2+a)^2}\right)^{\frac{3}{2}}}{4b}$	26
default	$-\frac{(bx^2+a)\left(\frac{c}{(bx^2+a)^2}\right)^{\frac{3}{2}}}{4b}$	26
oring	$-\frac{(bx^2+a)\left(\frac{c}{(bx^2+a)^2}\right)^{\frac{3}{2}}}{4b}$	26
risch	$-\frac{c\sqrt{\frac{c}{(bx^2+a)^2}}}{4b(bx^2+a)}$	29
trager	$\frac{c(bx^2+2a)x^2\sqrt{\frac{c}{b^2x^4+2abx^2+a^2}}}{4a^2(bx^2+a)}$	52

input `int(x*(c/(b*x^2+a)^2)^(3/2),x,method=_RETURNVERBOSE)`

output `-1/4*(b*x^2+a)/b*(c/(b*x^2+a)^2)^(3/2)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.25

$$\int x \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = -\frac{c \sqrt{b^2 x^4 + 2abx^2 + a^2}}{4(b^2 x^2 + ab)}$$

input `integrate(x*(c/(b*x^2+a)^2)^(3/2),x, algorithm="fricas")`output `-1/4*c*sqrt(c/(b^2*x^4 + 2*a*b*x^2 + a^2))/(b^2*x^2 + a*b)`**Sympy [F]**

$$\int x \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \int x \left(\frac{c}{a^2 + 2abx^2 + b^2x^4} \right)^{\frac{3}{2}} dx$$

input `integrate(x*(c/(b*x**2+a)**2)**(3/2),x)`output `Integral(x*(c/(a**2 + 2*a*b*x**2 + b**2*x**4))**(3/2), x)`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.91

$$\int x \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = -\frac{c^{\frac{3}{2}}}{4(b^3 x^4 + 2ab^2 x^2 + a^2 b)}$$

input `integrate(x*(c/(b*x^2+a)^2)^(3/2),x, algorithm="maxima")`output `-1/4*c^(3/2)/(b^3*x^4 + 2*a*b^2*x^2 + a^2*b)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.78

$$\int x \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = -\frac{c^{3/2} \operatorname{sgn}(bx^2 + a)}{4(bx^2 + a)^2 b}$$

input `integrate(x*(c/(b*x^2+a)^2)^(3/2),x, algorithm="giac")`output `-1/4*c^(3/2)*sgn(b*x^2 + a)/((b*x^2 + a)^2*b)`**Mupad [B] (verification not implemented)**

Time = 0.46 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.88

$$\int x \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = -\frac{c \sqrt{\frac{c}{(bx^2+a)^2}}}{4b(bx^2 + a)}$$

input `int(x*(c/(a + b*x^2)^2)^(3/2),x)`output `-(c*(c/(a + b*x^2)^2)^(1/2))/(4*b*(a + b*x^2))`**Reduce [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.88

$$\int x \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = -\frac{\sqrt{c} c}{4b(b^2x^4 + 2abx^2 + a^2)}$$

input `int(x*(c/(b*x^2+a)^2)^(3/2),x)`output `(- sqrt(c)*c)/(4*b*(a**2 + 2*a*b*x**2 + b**2*x**4))`

3.32
$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x} dx$$

Optimal result	250
Mathematica [A] (verified)	251
Rubi [A] (verified)	251
Maple [A] (verified)	253
Fricas [A] (verification not implemented)	253
Sympy [F]	254
Maxima [A] (verification not implemented)	254
Giac [A] (verification not implemented)	254
Mupad [F(-1)]	255
Reduce [B] (verification not implemented)	255

Optimal result

Integrand size = 19, antiderivative size = 123

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x} dx = \frac{c\sqrt{\frac{c}{(a+bx^2)^2}}}{2a^2} + \frac{c\sqrt{\frac{c}{(a+bx^2)^2}}}{4a(a+bx^2)} + \frac{c\sqrt{\frac{c}{(a+bx^2)^2}}(a+bx^2)\log(x)}{a^3} - \frac{c\sqrt{\frac{c}{(a+bx^2)^2}}(a+bx^2)\log(a+bx^2)}{2a^3}$$

output

```
1/2*c*(c/(b*x^2+a)^2)^(1/2)/a^2+1/4*c*(c/(b*x^2+a)^2)^(1/2)/a/(b*x^2+a)+c*(c/(b*x^2+a)^2)^(1/2)*(b*x^2+a)*ln(x)/a^3-1/2*c*(c/(b*x^2+a)^2)^(1/2)*(b*x^2+a)*ln(b*x^2+a)/a^3
```

Mathematica [A] (verified)

Time = 1.03 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.60

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x} dx = \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2} (a+bx^2) \left(a(3a+2bx^2) + 4(a+bx^2)^2 \log(x) - 2(a+bx^2)^2 \log(a+bx^2)\right)}{4a^3}$$

input `Integrate[(c/(a + b*x^2)^2)^(3/2)/x,x]`

output `((c/(a + b*x^2)^2)^(3/2)*(a + b*x^2)*(a*(3*a + 2*b*x^2) + 4*(a + b*x^2)^2*Log[x] - 2*(a + b*x^2)^2*Log[a + b*x^2]))/(4*a^3)`

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.63, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {2045, 27, 243, 54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x} dx \\ & \quad \downarrow \text{2045} \\ & \frac{c(a+bx^2) \sqrt{\frac{c}{(a+bx^2)^2}} \int \frac{a^3}{x(bx^2+a)^3} dx}{a^3} \\ & \quad \downarrow \text{27} \\ & c(a+bx^2) \sqrt{\frac{c}{(a+bx^2)^2}} \int \frac{1}{x(bx^2+a)^3} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{2}c(a+bx^2) \sqrt{\frac{c}{(a+bx^2)^2}} \int \frac{1}{x^2(bx^2+a)^3} dx^2 \\ & \quad \downarrow \text{54} \end{aligned}$$

$$\frac{1}{2}c(a+bx^2)\sqrt{\frac{c}{(a+bx^2)^2}}\int\left(-\frac{b}{a^3(bx^2+a)}-\frac{b}{a^2(bx^2+a)^2}-\frac{b}{a(bx^2+a)^3}+\frac{1}{a^3x^2}\right)dx^2$$

↓ 2009

$$\frac{1}{2}c(a+bx^2)\sqrt{\frac{c}{(a+bx^2)^2}}\left(-\frac{\log(a+bx^2)}{a^3}+\frac{\log(x^2)}{a^3}+\frac{1}{a^2(a+bx^2)}+\frac{1}{2a(a+bx^2)^2}\right)$$

input `Int[(c/(a + b*x^2)^2)^(3/2)/x,x]`

output `(c*Sqrt[c/(a + b*x^2)^2]*(a + b*x^2)*(1/(2*a*(a + b*x^2)^2) + 1/(a^2*(a + b*x^2))) + Log[x^2]/a^3 - Log[a + b*x^2]/a^3))/2`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 54 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && !LtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 243 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2045 `Int[(u_)*((c_)*((a_) + (b_)*(x_)^(n_))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.83

method	result
risch	$\frac{c \sqrt{\frac{c}{(bx^2+a)^2}} \left(\frac{bx^2}{2a^2} + \frac{3}{4a}\right)}{bx^2+a} + \frac{c \sqrt{\frac{c}{(bx^2+a)^2}} (bx^2+a) \ln(x)}{a^3} - \frac{c \sqrt{\frac{c}{(bx^2+a)^2}} (bx^2+a) \ln(bx^2+a)}{2a^3}$
default	$-\frac{(2 \ln(bx^2+a) b^2 x^4 - 4 \ln(x) b^2 x^4 + 4 \ln(bx^2+a) abx^2 - 8 \ln(x) abx^2 - 2abx^2 + 2 \ln(bx^2+a) a^2 - 4a^2 \ln(x) - 3a^2) (bx^2+a) \left(\frac{c}{(bx^2+a)}\right)}{4a^3}$

input `int((c/(b*x^2+a)^2)^(3/2)/x,x,method=_RETURNVERBOSE)`output
$$\frac{c/(bx^2+a) * (c/(bx^2+a)^2)^{(1/2)} * (1/2 * b/a^2 * x^2 + 3/4/a) + c * (c/(bx^2+a)^2)^{(1/2)} * (bx^2+a) * \ln(x)/a^3 - 1/2 * c * (c/(bx^2+a)^2)^{(1/2)} * (bx^2+a) * \ln(bx^2+a)}{a^3}$$
Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.92

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x} dx = \frac{(2abcx^2 + 3a^2c - 2(b^2cx^4 + 2abcx^2 + a^2c) \log(bx^2 + a) + 4(b^2cx^4 + 2abcx^2 + a^2c) \log(x)) \sqrt{c/(b^2x^4 + 2abx^2 + a^2)}}{4(a^3bx^2 + a^4)}$$

input `integrate((c/(b*x^2+a)^2)^(3/2)/x,x, algorithm="fricas")`output
$$\frac{1/4 * (2*a*b*c*x^2 + 3*a^2*c - 2*(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c) * \log(b*x^2 + a) + 4*(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c) * \log(x)) * \sqrt{c/(b^2*x^4 + 2*a*b*x^2 + a^2)}}{(a^3*b*x^2 + a^4)}$$

Sympy [F]

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x} dx = \int \frac{\left(\frac{c}{a^2+2abx^2+b^2x^4}\right)^{3/2}}{x} dx$$

input `integrate((c/(b*x**2+a)**2)**(3/2)/x,x)`

output `Integral((c/(a**2 + 2*a*b*x**2 + b**2*x**4))**(3/2)/x, x)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.59

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x} dx = \frac{2bc^{\frac{3}{2}}x^2 + 3ac^{\frac{3}{2}}}{4(a^2b^2x^4 + 2a^3bx^2 + a^4)} - \frac{c^{\frac{3}{2}} \log(bx^2 + a)}{2a^3} + \frac{c^{\frac{3}{2}} \log(x^2)}{2a^3}$$

input `integrate((c/(b*x^2+a)^2)^(3/2)/x,x, algorithm="maxima")`

output `1/4*(2*b*c^(3/2)*x^2 + 3*a*c^(3/2))/(a^2*b^2*x^4 + 2*a^3*b*x^2 + a^4) - 1/2*c^(3/2)*log(b*x^2 + a)/a^3 + 1/2*c^(3/2)*log(x^2)/a^3`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.58

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x} dx = \frac{1}{4} c^{\frac{3}{2}} \left(\frac{2 \log(x^2)}{a^3} - \frac{2 \log(|bx^2 + a|)}{a^3} + \frac{3b^2x^4 + 8abx^2 + 6a^2}{(bx^2 + a)^2 a^3} \right) \operatorname{sgn}(bx^2 + a)$$

input `integrate((c/(b*x^2+a)^2)^(3/2)/x,x, algorithm="giac")`

output $\frac{1}{4}c^{3/2}(2\log(x^2)/a^3 - 2\log(\text{abs}(b*x^2 + a))/a^3 + (3*b^2*x^4 + 8*a*b*x^2 + 6*a^2)/((b*x^2 + a)^2*a^3))*\text{sgn}(b*x^2 + a)$

Mupad [F(-1)]

Timed out.

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x} dx = \int \frac{\left(\frac{c}{(bx^2+a)^2}\right)^{3/2}}{x} dx$$

input `int((c/(a + b*x^2)^2)^(3/2)/x,x)`

output `int((c/(a + b*x^2)^2)^(3/2)/x, x)`

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.91

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x} dx = \frac{\sqrt{c}c(-2\log(bx^2 + a)a^2 - 4\log(bx^2 + a)abx^2 - 2\log(bx^2 + a)b^2x^4 + 4\log(x)a^2 + 4\log(x)abx^2 + 4\log(x)b^2x^4 + 2a^2 - b^2x^4)}{4a^3(b^2x^4 + 2abx^2 + a^2)}$$

input `int((c/(b*x^2+a)^2)^(3/2)/x,x)`

output `(sqrt(c)*c*(- 2*log(a + b*x**2)*a**2 - 4*log(a + b*x**2)*a*b*x**2 - 2*log(a + b*x**2)*b**2*x**4 + 4*log(x)*a**2 + 8*log(x)*a*b*x**2 + 4*log(x)*b**2*x**4 + 2*a**2 - b**2*x**4))/(4*a**3*(a**2 + 2*a*b*x**2 + b**2*x**4))`

3.33
$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^3} dx$$

Optimal result	256
Mathematica [A] (verified)	257
Rubi [A] (verified)	257
Maple [A] (verified)	259
Fricas [A] (verification not implemented)	259
Sympy [F]	260
Maxima [A] (verification not implemented)	260
Giac [A] (verification not implemented)	261
Mupad [F(-1)]	261
Reduce [B] (verification not implemented)	262

Optimal result

Integrand size = 19, antiderivative size = 159

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^3} dx = -\frac{bc\sqrt{\frac{c}{(a+bx^2)^2}}}{a^3} - \frac{bc\sqrt{\frac{c}{(a+bx^2)^2}}}{4a^2(a+bx^2)} - \frac{c\sqrt{\frac{c}{(a+bx^2)^2}}(a+bx^2)}{2a^3x^2}$$

$$- \frac{3bc\sqrt{\frac{c}{(a+bx^2)^2}}(a+bx^2)\log(x)}{a^4} + \frac{3bc\sqrt{\frac{c}{(a+bx^2)^2}}(a+bx^2)\log(a+bx^2)}{2a^4}$$

output

```
-b*c*(c/(b*x^2+a)^2)^(1/2)/a^3-1/4*b*c*(c/(b*x^2+a)^2)^(1/2)/a^2/(b*x^2+a)
-1/2*c*(c/(b*x^2+a)^2)^(1/2)*(b*x^2+a)/a^3/x^2-3*b*c*(c/(b*x^2+a)^2)^(1/2)
*(b*x^2+a)*ln(x)/a^4+3/2*b*c*(c/(b*x^2+a)^2)^(1/2)*(b*x^2+a)*ln(b*x^2+a)/a
^4
```

Mathematica [A] (verified)

Time = 1.03 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.60

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^3} dx = \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2} (a+bx^2) \left(a(2a^2+9abx^2+6b^2x^4) + 12bx^2(a+bx^2)^2 \log(x) - 6bx^2(a+bx^2)^2 \log(a+bx^2)\right)}{4a^4x^2}$$

input `Integrate[(c/(a + b*x^2)^2)^(3/2)/x^3,x]`

output `-1/4*((c/(a + b*x^2)^2)^(3/2)*(a + b*x^2)*(a*(2*a^2 + 9*a*b*x^2 + 6*b^2*x^4) + 12*b*x^2*(a + b*x^2)^2*Log[x] - 6*b*x^2*(a + b*x^2)^2*Log[a + b*x^2]))/(a^4*x^2)`

Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.58, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {2045, 27, 243, 54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^3} dx \\ & \quad \downarrow \text{2045} \\ & \frac{c(a+bx^2) \sqrt{\frac{c}{(a+bx^2)^2}} \int \frac{a^3}{x^3(bx^2+a)^3} dx}{a^3} \\ & \quad \downarrow \text{27} \\ & c(a+bx^2) \sqrt{\frac{c}{(a+bx^2)^2}} \int \frac{1}{x^3(bx^2+a)^3} dx \\ & \quad \downarrow \text{243} \end{aligned}$$

$$\frac{1}{2}c(a + bx^2) \sqrt{\frac{c}{(a + bx^2)^2}} \int \frac{1}{x^4 (bx^2 + a)^3} dx^2$$

↓ 54

$$\frac{1}{2}c(a + bx^2) \sqrt{\frac{c}{(a + bx^2)^2}} \int \left(\frac{3b^2}{a^4 (bx^2 + a)} + \frac{2b^2}{a^3 (bx^2 + a)^2} + \frac{b^2}{a^2 (bx^2 + a)^3} - \frac{3b}{a^4 x^2} + \frac{1}{a^3 x^4} \right) dx^2$$

↓ 2009

$$\frac{1}{2}c(a + bx^2) \sqrt{\frac{c}{(a + bx^2)^2}} \left(-\frac{3b \log(x^2)}{a^4} + \frac{3b \log(a + bx^2)}{a^4} - \frac{2b}{a^3 (a + bx^2)} - \frac{1}{a^3 x^2} - \frac{b}{2a^2 (a + bx^2)^2} \right)$$

input `Int[(c/(a + b*x^2)^2)^(3/2)/x^3,x]`

output `(c*Sqrt[c/(a + b*x^2)^2]*(a + b*x^2)*(-(1/(a^3*x^2)) - b/(2*a^2*(a + b*x^2)^2) - (2*b)/(a^3*(a + b*x^2)) - (3*b*Log[x^2])/a^4 + (3*b*Log[a + b*x^2])/a^4))/2`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 54 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && !LtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2045

```
Int[(u_)*((c_)*((a_) + (b_)*(x_)^(n_))^(q_))^(p_), x_Symbol] := Simp[Simp[
mp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q)
, x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]
```

Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.77

method	result
risch	$\frac{c \sqrt{\frac{c}{(bx^2+a)^2}} \left(-\frac{3b^2x^4}{2a^3} - \frac{9bx^2}{4a^2} - \frac{1}{2a} \right)}{(bx^2+a)x^2} - \frac{3bc \sqrt{\frac{c}{(bx^2+a)^2}} (bx^2+a) \ln(x)}{a^4} + \frac{3c(bx^2+a) \sqrt{\frac{c}{(bx^2+a)^2}} b \ln(-bx^2-a)}{2a^4}$
default	$-\frac{(12 \ln(x)b^3x^6 - 6 \ln(bx^2+a)b^3x^6 + 24ab^2 \ln(x)x^4 - 12 \ln(bx^2+a)ab^2x^4 + 6ab^2x^4 + 12a^2b \ln(x)x^2 - 6 \ln(bx^2+a)a^2bx^2 + 9a^2bx^2 + a^2b \ln(-bx^2-a))}{4a^4x^2}$

input

```
int((c/(b*x^2+a)^2)^(3/2)/x^3,x,method=_RETURNVERBOSE)
```

output

```
c/(b*x^2+a)*(c/(b*x^2+a)^2)^(1/2)*(-3/2*b^2/a^3*x^4-9/4*b/a^2*x^2-1/2/a)/x
^2-3*b*c*(c/(b*x^2+a)^2)^(1/2)*(b*x^2+a)*ln(x)/a^4+3/2*c*(b*x^2+a)*(c/(b*x
^2+a)^2)^(1/2)/a^4*b*ln(-b*x^2-a)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.89

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^3} dx =$$

$$\frac{(6ab^2cx^4 + 9a^2bcx^2 + 2a^3c - 6(b^3cx^6 + 2ab^2cx^4 + a^2bcx^2) \log(bx^2 + a) + 12(b^3cx^6 + 2ab^2cx^4 + a^2bcx^2) \log(-bx^2 - a))}{4(a^4bx^4 + a^5x^2)}$$

input

```
integrate((c/(b*x^2+a)^2)^(3/2)/x^3,x, algorithm="fricas")
```

output

```
-1/4*(6*a*b^2*c*x^4 + 9*a^2*b*c*x^2 + 2*a^3*c - 6*(b^3*c*x^6 + 2*a*b^2*c*x^4 + a^2*b*c*x^2)*log(b*x^2 + a) + 12*(b^3*c*x^6 + 2*a*b^2*c*x^4 + a^2*b*c*x^2)*log(x))*sqrt(c/(b^2*x^4 + 2*a*b*x^2 + a^2))/(a^4*b*x^4 + a^5*x^2)
```

Sympy [F]

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^3} dx = \int \frac{\left(\frac{c}{a^2+2abx^2+b^2x^4}\right)^{3/2}}{x^3} dx$$

input

```
integrate((c/(b*x**2+a)**2)**(3/2)/x**3,x)
```

output

```
Integral((c/(a**2 + 2*a*b*x**2 + b**2*x**4))**(3/2)/x**3, x)
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.58

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^3} dx = -\frac{6b^2c^{\frac{3}{2}}x^4 + 9abc^{\frac{3}{2}}x^2 + 2a^2c^{\frac{3}{2}}}{4(a^3b^2x^6 + 2a^4bx^4 + a^5x^2)} + \frac{3bc^{\frac{3}{2}}\log(bx^2 + a)}{2a^4} - \frac{3bc^{\frac{3}{2}}\log(x^2)}{2a^4}$$

input

```
integrate((c/(b*x^2+a)^2)^(3/2)/x^3,x, algorithm="maxima")
```

output

```
-1/4*(6*b^2*c^(3/2)*x^4 + 9*a*b*c^(3/2)*x^2 + 2*a^2*c^(3/2))/(a^3*b^2*x^6 + 2*a^4*b*x^4 + a^5*x^2) + 3/2*b*c^(3/2)*log(b*x^2 + a)/a^4 - 3/2*b*c^(3/2)*log(x^2)/a^4
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.59

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^3} dx = -\frac{1}{4} c^{\frac{3}{2}} \left(\frac{6b \log(x^2)}{a^4} - \frac{6b \log(|bx^2 + a|)}{a^4} + \frac{9b^3x^4 + 22ab^2x^2 + 14a^2b}{(bx^2 + a)^2 a^4} - \frac{2(3bx^2 - a)}{a^4 x^2} \right) \operatorname{sgn}(bx^2 + a)$$

input `integrate((c/(b*x^2+a)^2)^(3/2)/x^3,x, algorithm="giac")`

output `-1/4*c^(3/2)*(6*b*log(x^2)/a^4 - 6*b*log(abs(b*x^2 + a))/a^4 + (9*b^3*x^4 + 22*a*b^2*x^2 + 14*a^2*b)/((b*x^2 + a)^2*a^4) - 2*(3*b*x^2 - a)/(a^4*x^2))*sgn(b*x^2 + a)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^3} dx = \int \frac{\left(\frac{c}{(bx^2+a)^2}\right)^{3/2}}{x^3} dx$$

input `int((c/(a + b*x^2)^2)^(3/2)/x^3,x)`

output `int((c/(a + b*x^2)^2)^(3/2)/x^3, x)`

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.86

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^3} dx = \frac{\sqrt{c}c(6\log(bx^2+a)a^2bx^2 + 12\log(bx^2+a)ab^2x^4 + 6\log(bx^2+a)b^3x^6 - 12\log(x))}{4a^4x^2(b^2x^4 + 2abx^2 - a^2)}$$

input `int((c/(b*x^2+a)^2)^(3/2)/x^3,x)`output `(sqrt(c)*c*(6*log(a + b*x**2)*a**2*b*x**2 + 12*log(a + b*x**2)*a*b**2*x**4 + 6*log(a + b*x**2)*b**3*x**6 - 12*log(x)*a**2*b*x**2 - 24*log(x)*a*b**2*x**4 - 12*log(x)*b**3*x**6 - 2*a**3 - 6*a**2*b*x**2 + 3*b**3*x**6))/(4*a**4*x**2*(a**2 + 2*a*b*x**2 + b**2*x**4))`

3.34
$$\int x^4 \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx$$

Optimal result	263
Mathematica [A] (verified)	263
Rubi [A] (verified)	264
Maple [A] (verified)	266
Fricas [A] (verification not implemented)	266
Sympy [F]	267
Maxima [A] (verification not implemented)	267
Giac [A] (verification not implemented)	267
Mupad [F(-1)]	268
Reduce [B] (verification not implemented)	268

Optimal result

Integrand size = 19, antiderivative size = 110

$$\int x^4 \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx = -\frac{3cx\sqrt{\frac{c}{(a+bx^2)^2}}}{8b^2} - \frac{cx^3\sqrt{\frac{c}{(a+bx^2)^2}}}{4b(a+bx^2)} + \frac{3c\sqrt{\frac{c}{(a+bx^2)^2}}(a+bx^2)\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{8\sqrt{ab^5/2}}$$

output

```
-3/8*c*x*(c/(b*x^2+a)^2)^(1/2)/b^2-1/4*c*x^3*(c/(b*x^2+a)^2)^(1/2)/b/(b*x^2+a)+3/8*c*(c/(b*x^2+a)^2)^(1/2)*(b*x^2+a)*arctan(b^(1/2)*x/a^(1/2))/a^(1/2)/b^(5/2)
```

Mathematica [A] (verified)

Time = 1.04 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.76

$$\int x^4 \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx = \frac{\left(\frac{c}{(a+bx^2)^2} \right)^{3/2} (a+bx^2) \left(-\sqrt{a}\sqrt{bx}(3a+5bx^2) + 3(a+bx^2)^2 \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) \right)}{8\sqrt{ab^5/2}}$$

input `Integrate[x^4*(c/(a + b*x^2)^2)^(3/2),x]`

output `((c/(a + b*x^2)^2)^(3/2)*(a + b*x^2)*(-(Sqrt[a]*Sqrt[b]*x*(3*a + 5*b*x^2)) + 3*(a + b*x^2)^2*ArcTan[(Sqrt[b]*x)/Sqrt[a]]))/(8*Sqrt[a]*b^(5/2))`

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.87, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {2045, 27, 252, 252, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^4 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx \\
 & \quad \downarrow \text{2045} \\
 & \frac{c(a + bx^2) \sqrt{\frac{c}{(a+bx^2)^2}} \int \frac{a^3 x^4}{(bx^2+a)^3} dx}{a^3} \\
 & \quad \downarrow \text{27} \\
 & c(a + bx^2) \sqrt{\frac{c}{(a + bx^2)^2}} \int \frac{x^4}{(bx^2 + a)^3} dx \\
 & \quad \downarrow \text{252} \\
 & c(a + bx^2) \sqrt{\frac{c}{(a + bx^2)^2}} \left(\frac{3 \int \frac{x^2}{(bx^2+a)^2} dx}{4b} - \frac{x^3}{4b(a + bx^2)^2} \right) \\
 & \quad \downarrow \text{252} \\
 & c(a + bx^2) \sqrt{\frac{c}{(a + bx^2)^2}} \left(\frac{3 \left(\frac{\int \frac{1}{bx^2+a} dx}{2b} - \frac{x}{2b(a+bx^2)} \right)}{4b} - \frac{x^3}{4b(a + bx^2)^2} \right) \\
 & \quad \downarrow \text{218}
 \end{aligned}$$

$$c(a + bx^2) \left(\frac{3 \left(\frac{\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{2\sqrt{ab^{3/2}}} - \frac{x}{2b(a+bx^2)} \right)}{4b} - \frac{x^3}{4b(a+bx^2)^2} \right) \sqrt{\frac{c}{(a+bx^2)^2}}$$

input `Int[x^4*(c/(a + b*x^2)^2)^(3/2),x]`

output `c*Sqrt[c/(a + b*x^2)^2]*(a + b*x^2)*(-1/4*x^3/(b*(a + b*x^2)^2) + (3*(-1/2*x/(b*(a + b*x^2)) + ArcTan[(Sqrt[b]*x)/Sqrt[a]]/(2*Sqrt[a]*b^(3/2))))/(4*b))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 252 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(2*b*(p + 1))), x] - Simp[c^2*((m - 1)/(2*b*(p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && LtQ[p, -1] && GtQ[m, 1] && !ILtQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_.))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^(p/(1 + b*(x^n/a))^(p*q)) Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.90

method	result	size
default	$-\frac{\left(-3 \arctan\left(\frac{bx}{\sqrt{ab}}\right) b^2 x^4 + 5\sqrt{ab} b x^3 - 6 \arctan\left(\frac{bx}{\sqrt{ab}}\right) a b x^2 + 3\sqrt{ab} a x - 3 \arctan\left(\frac{bx}{\sqrt{ab}}\right) a^2\right) (b x^2 + a) \left(\frac{c}{(b x^2 + a)^2}\right)^{3/2}}{8 b^2 \sqrt{ab}}$	99
risch	$\frac{c \sqrt{\frac{c}{(b x^2 + a)^2}} \left(-\frac{5 x^3}{8 b} - \frac{3 a x}{8 b^2}\right)}{b x^2 + a} - \frac{3 c (b x^2 + a) \sqrt{\frac{c}{(b x^2 + a)^2}} \ln(b x + \sqrt{-ab})}{16 \sqrt{-ab} b^2} + \frac{3 c (b x^2 + a) \sqrt{\frac{c}{(b x^2 + a)^2}} \ln(-b x + \sqrt{-ab})}{16 \sqrt{-ab} b^2}$	129

input `int(x^4*(c/(b*x^2+a)^2)^(3/2),x,method=_RETURNVERBOSE)`

output
$$-1/8*(-3*\arctan(b*x/(a*b)^(1/2))*b^2*x^4+5*(a*b)^(1/2)*b*x^3-6*\arctan(b*x/(a*b)^(1/2))*a*b*x^2+3*(a*b)^(1/2)*a*x-3*\arctan(b*x/(a*b)^(1/2))*a^2)*(b*x^2+a)*(c/(b*x^2+a)^2)^(3/2)/b^2/(a*b)^(1/2)$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 285, normalized size of antiderivative = 2.59

$$\int x^4 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \left[\frac{3 (bcx^2 + ac) \sqrt{-\frac{c}{ab}} \log \left(\frac{bcx^2 - ac + 2 (ab^2 x^3 + a^2 bx) \sqrt{\frac{c}{b^2 x^4 + 2 abx^2 + a^2}} \sqrt{-\frac{c}{ab}}}{bx^2 + a} \right)}{16 (b^3 x^2 + ab^2)} - 2 (5 b c x^3 + a^2 b^2 x) \sqrt{c/(b^2 x^4 + 2 a b x^2 + a^2)} \sqrt{-c/(a b)}}{16 (b^3 x^2 + ab^2)} \right]$$

input `integrate(x^4*(c/(b*x^2+a)^2)^(3/2),x, algorithm="fricas")`

output
$$[1/16*(3*(b*c*x^2 + a*c)*\sqrt{-c/(a*b)}*\log((b*c*x^2 - a*c + 2*(a*b^2*x^3 + a^2*b*x)*\sqrt{c/(b^2*x^4 + 2*a*b*x^2 + a^2)}*\sqrt{-c/(a*b)}))/(b*x^2 + a) - 2*(5*b*c*x^3 + 3*a*c*x)*\sqrt{c/(b^2*x^4 + 2*a*b*x^2 + a^2)}}/(b^3*x^2 + a*b^2), 1/8*(3*(b*c*x^2 + a*c)*\sqrt{c/(a*b)}*\arctan((b^2*x^3 + a*b*x)*\sqrt{c/(b^2*x^4 + 2*a*b*x^2 + a^2)}*\sqrt{c/(a*b)})/c - (5*b*c*x^3 + 3*a*c*x)*\sqrt{c/(b^2*x^4 + 2*a*b*x^2 + a^2)}}/(b^3*x^2 + a*b^2)]$$

Sympy [F]

$$\int x^4 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \int x^4 \left(\frac{c}{a^2 + 2abx^2 + b^2x^4} \right)^{3/2} dx$$

input `integrate(x**4*(c/(b*x**2+a)**2)**(3/2),x)`

output `Integral(x**4*(c/(a**2 + 2*a*b*x**2 + b**2*x**4))**(3/2), x)`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.62

$$\int x^4 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = -\frac{5bc^{\frac{3}{2}}x^3 + 3ac^{\frac{3}{2}}x}{8(b^4x^4 + 2ab^3x^2 + a^2b^2)} + \frac{3c^{\frac{3}{2}} \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{8\sqrt{abb^2}}$$

input `integrate(x^4*(c/(b*x^2+a)^2)^(3/2),x, algorithm="maxima")`

output `-1/8*(5*b*c^(3/2)*x^3 + 3*a*c^(3/2)*x)/(b^4*x^4 + 2*a*b^3*x^2 + a^2*b^2) + 3/8*c^(3/2)*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*b^2)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.67

$$\int x^4 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \frac{1}{8} c^{\frac{3}{2}} \left(\frac{3 \arctan\left(\frac{bx}{\sqrt{ab}}\right) \operatorname{sgn}(bx^2 + a)}{\sqrt{abb^2}} - \frac{5bx^3 \operatorname{sgn}(bx^2 + a) + 3ax \operatorname{sgn}(bx^2 + a)}{(bx^2 + a)^2 b^2} \right)$$

input `integrate(x^4*(c/(b*x^2+a)^2)^(3/2),x, algorithm="giac")`

output `1/8*c^(3/2)*(3*arctan(b*x/sqrt(a*b))*sgn(b*x^2 + a)/(sqrt(a*b)*b^2) - (5*b*x^3*sgn(b*x^2 + a) + 3*a*x*sgn(b*x^2 + a))/((b*x^2 + a)^2*b^2)`

Mupad [F(-1)]

Timed out.

$$\int x^4 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \int x^4 \left(\frac{c}{(bx^2 + a)^2} \right)^{3/2} dx$$

input `int(x^4*(c/(a + b*x^2)^2)^(3/2), x)`output `int(x^4*(c/(a + b*x^2)^2)^(3/2), x)`**Reduce [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.05

$$\int x^4 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \frac{\sqrt{c} c \left(3\sqrt{b} \sqrt{a} \operatorname{atan} \left(\frac{bx}{\sqrt{b}\sqrt{a}} \right) a^2 + 6\sqrt{b} \sqrt{a} \operatorname{atan} \left(\frac{bx}{\sqrt{b}\sqrt{a}} \right) abx^2 + 3\sqrt{b} \sqrt{a} \operatorname{atan} \left(\frac{bx}{\sqrt{b}\sqrt{a}} \right) \right)}{8ab^3(b^2x^4 + 2abx^2 + a^2)}$$

input `int(x^4*(c/(b*x^2+a)^2)^(3/2), x)`output `(sqrt(c)*c*(3*sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*sqrt(a)))*a**2 + 6*sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*sqrt(a)))*a*b*x**2 + 3*sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*sqrt(a)))*b**2*x**4 - 3*a**2*b*x - 5*a*b**2*x**3)/(8*a*b**3*(a**2 + 2*a*b*x**2 + b**2*x**4))`

3.35
$$\int x^2 \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx$$

Optimal result	269
Mathematica [A] (verified)	269
Rubi [A] (verified)	270
Maple [A] (verified)	272
Fricas [A] (verification not implemented)	272
Sympy [F]	273
Maxima [A] (verification not implemented)	273
Giac [A] (verification not implemented)	273
Mupad [F(-1)]	274
Reduce [B] (verification not implemented)	274

Optimal result

Integrand size = 19, antiderivative size = 111

$$\int x^2 \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx = \frac{cx \sqrt{\frac{c}{(a+bx^2)^2}}}{8ab} - \frac{cx \sqrt{\frac{c}{(a+bx^2)^2}}}{4b(a+bx^2)} + \frac{c \sqrt{\frac{c}{(a+bx^2)^2}} (a+bx^2) \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{8a^{3/2}b^{3/2}}$$

output

```
1/8*c*x*(c/(b*x^2+a)^2)^(1/2)/a/b-1/4*c*x*(c/(b*x^2+a)^2)^(1/2)/b/(b*x^2+a)
)+1/8*c*(c/(b*x^2+a)^2)^(1/2)*(b*x^2+a)*arctan(b^(1/2)*x/a^(1/2))/a^(3/2)/
b^(3/2)
```

Mathematica [A] (verified)

Time = 1.03 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.73

$$\int x^2 \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx = \frac{\left(\frac{c}{(a+bx^2)^2} \right)^{3/2} (a+bx^2) \left(\sqrt{a}\sqrt{bx}(-a+bx^2) + (a+bx^2)^2 \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) \right)}{8a^{3/2}b^{3/2}}$$

input `Integrate[x^2*(c/(a + b*x^2)^2)^(3/2),x]`

output `((c/(a + b*x^2)^2)^(3/2)*(a + b*x^2)*(Sqrt[a]*Sqrt[b]*x*(-a + b*x^2) + (a + b*x^2)^2*ArcTan[(Sqrt[b]*x)/Sqrt[a]]))/(8*a^(3/2)*b^(3/2))`

Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.85, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {2045, 27, 252, 215, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx \\
 & \quad \downarrow \text{2045} \\
 & \frac{c(a + bx^2) \sqrt{\frac{c}{(a+bx^2)^2}} \int \frac{a^3 x^2}{(bx^2+a)^3} dx}{a^3} \\
 & \quad \downarrow \text{27} \\
 & c(a + bx^2) \sqrt{\frac{c}{(a + bx^2)^2}} \int \frac{x^2}{(bx^2 + a)^3} dx \\
 & \quad \downarrow \text{252} \\
 & c(a + bx^2) \sqrt{\frac{c}{(a + bx^2)^2}} \left(\frac{\int \frac{1}{(bx^2+a)^2} dx}{4b} - \frac{x}{4b(a + bx^2)^2} \right) \\
 & \quad \downarrow \text{215} \\
 & c(a + bx^2) \sqrt{\frac{c}{(a + bx^2)^2}} \left(\frac{\int \frac{1}{bx^2+a} dx}{2a} + \frac{x}{2a(a+bx^2)} - \frac{x}{4b(a + bx^2)^2} \right) \\
 & \quad \downarrow \text{218}
 \end{aligned}$$

$$c(a + bx^2) \left(\frac{\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{2a^{3/2}\sqrt{b}} + \frac{x}{2a(a+bx^2)} - \frac{x}{4b(a+bx^2)^2} \right) \sqrt{\frac{c}{(a+bx^2)^2}}$$

input `Int[x^2*(c/(a + b*x^2)^2)^(3/2),x]`

output `c*Sqrt[c/(a + b*x^2)^2]*(a + b*x^2)*(-1/4*x/(b*(a + b*x^2)^2) + (x/(2*a*(a + b*x^2)) + ArcTan[(Sqrt[b]*x)/Sqrt[a]]/(2*a^(3/2)*Sqrt[b]))/(4*b)`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 215 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && LtQ[p, -1] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 252 `Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(2*b*(p + 1))), x] - Simp[c^2*(m - 1)/(2*b*(p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && LtQ[p, -1] && GtQ[m, 1] && !ILtQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.)))^(q_)^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.91

method	result	size
default	$-\frac{\left(-\arctan\left(\frac{bx}{\sqrt{ab}}\right)b^2x^4-\sqrt{ab}bx^3-2\arctan\left(\frac{bx}{\sqrt{ab}}\right)abx^2+\sqrt{ab}ax-\arctan\left(\frac{bx}{\sqrt{ab}}\right)a^2\right)(bx^2+a)\left(\frac{c}{(bx^2+a)^2}\right)^{\frac{3}{2}}}{8ab\sqrt{ab}}$	101
risch	$\frac{c\sqrt{\frac{c}{(bx^2+a)^2}}\left(\frac{x^3}{8a}-\frac{x}{8b}\right)}{bx^2+a}-\frac{c(bx^2+a)\sqrt{\frac{c}{(bx^2+a)^2}}\ln(bx+\sqrt{-ab})}{16\sqrt{-ab}ba}+\frac{c(bx^2+a)\sqrt{\frac{c}{(bx^2+a)^2}}\ln(-bx+\sqrt{-ab})}{16\sqrt{-ab}ba}$	134

input `int(x^2*(c/(b*x^2+a)^2)^(3/2),x,method=_RETURNVERBOSE)`

output
$$-1/8*(-\arctan(b*x/(a*b)^(1/2))*b^2*x^4-(a*b)^(1/2)*b*x^3-2*\arctan(b*x/(a*b)^(1/2))*a*b*x^2+(a*b)^(1/2)*a*x-\arctan(b*x/(a*b)^(1/2))*a^2)*(b*x^2+a)*(c/(b*x^2+a)^2)^(3/2)/a/b/(a*b)^(1/2)$$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 282, normalized size of antiderivative = 2.54

$$\int x^2 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \left[\frac{(bcx^2 + ac) \sqrt{-\frac{c}{ab}} \log \left(\frac{bcx^2 - ac + 2(ab^2x^3 + a^2bx) \sqrt{\frac{c}{b^2x^4 + 2abx^2 + a^2}} \sqrt{-\frac{c}{ab}}}{bx^2 + a} \right) + 2(bc x^3 - a^2 b)}{16(ab^2x^2 + a^2b)} \right]$$

input `integrate(x^2*(c/(b*x^2+a)^2)^(3/2),x, algorithm="fricas")`

output
$$[1/16*((b*c*x^2 + a*c)*\sqrt{-c/(a*b)})*\log((b*c*x^2 - a*c + 2*(a*b^2*x^3 + a^2*b*x)*\sqrt{c/(b^2*x^4 + 2*a*b*x^2 + a^2)}*\sqrt{-c/(a*b)})/(b*x^2 + a)) + 2*(b*c*x^3 - a*c*x)*\sqrt{c/(b^2*x^4 + 2*a*b*x^2 + a^2)})/(a*b^2*x^2 + a^2*b), 1/8*((b*c*x^2 + a*c)*\sqrt{c/(a*b)})*\arctan((b^2*x^3 + a*b*x)*\sqrt{c/(b^2*x^4 + 2*a*b*x^2 + a^2)}*\sqrt{c/(a*b)})/c + (b*c*x^3 - a*c*x)*\sqrt{c/(b^2*x^4 + 2*a*b*x^2 + a^2)})/(a*b^2*x^2 + a^2*b)]$$

Sympy [F]

$$\int x^2 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \int x^2 \left(\frac{c}{a^2 + 2abx^2 + b^2x^4} \right)^{3/2} dx$$

input `integrate(x**2*(c/(b*x**2+a)**2)**(3/2),x)`

output `Integral(x**2*(c/(a**2 + 2*a*b*x**2 + b**2*x**4))**(3/2), x)`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.64

$$\int x^2 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \frac{bc^{\frac{3}{2}}x^3 - ac^{\frac{3}{2}}x}{8(ab^3x^4 + 2a^2b^2x^2 + a^3b)} + \frac{c^{\frac{3}{2}} \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{8\sqrt{abab}}$$

input `integrate(x^2*(c/(b*x^2+a)^2)^(3/2),x, algorithm="maxima")`

output `1/8*(b*c^(3/2)*x^3 - a*c^(3/2)*x)/(a*b^3*x^4 + 2*a^2*b^2*x^2 + a^3*b) + 1/8*c^(3/2)*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*a*b)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.69

$$\int x^2 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \frac{1}{8} c^{\frac{3}{2}} \left(\frac{\arctan\left(\frac{bx}{\sqrt{ab}}\right) \operatorname{sgn}(bx^2 + a)}{\sqrt{abab}} + \frac{bx^3 \operatorname{sgn}(bx^2 + a) - ax \operatorname{sgn}(bx^2 + a)}{(bx^2 + a)^2 ab} \right)$$

input `integrate(x^2*(c/(b*x^2+a)^2)^(3/2),x, algorithm="giac")`

output `1/8*c^(3/2)*(arctan(b*x/sqrt(a*b))*sgn(b*x^2 + a)/(sqrt(a*b)*a*b) + (b*x^3*sgn(b*x^2 + a) - a*x*sgn(b*x^2 + a))/((b*x^2 + a)^2*a*b))`

Mupad [F(-1)]

Timed out.

$$\int x^2 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \int x^2 \left(\frac{c}{(bx^2 + a)^2} \right)^{3/2} dx$$

input `int(x^2*(c/(a + b*x^2)^2)^(3/2), x)`output `int(x^2*(c/(a + b*x^2)^2)^(3/2), x)`**Reduce [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.02

$$\int x^2 \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \frac{\sqrt{c} c \left(\sqrt{b} \sqrt{a} \operatorname{atan} \left(\frac{bx}{\sqrt{b} \sqrt{a}} \right) a^2 + 2\sqrt{b} \sqrt{a} \operatorname{atan} \left(\frac{bx}{\sqrt{b} \sqrt{a}} \right) abx^2 + \sqrt{b} \sqrt{a} \operatorname{atan} \left(\frac{bx}{\sqrt{b} \sqrt{a}} \right) \right)}{8a^2b^2 (b^2x^4 + 2abx^2 + a^2)}$$

input `int(x^2*(c/(b*x^2+a)^2)^(3/2), x)`output `(sqrt(c)*c*(sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*sqrt(a)))*a**2 + 2*sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*sqrt(a)))*a*b*x**2 + sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*sqrt(a)))*b**2*x**4 - a**2*b*x + a*b**2*x**3))/(8*a**2*b**2*(a**2 + 2*a*b*x**2 + b**2*x**4))`

3.36
$$\int \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx$$

Optimal result	275
Mathematica [A] (verified)	275
Rubi [A] (verified)	276
Maple [A] (verified)	277
Fricas [A] (verification not implemented)	278
Sympy [F]	278
Maxima [A] (verification not implemented)	279
Giac [A] (verification not implemented)	279
Mupad [F(-1)]	279
Reduce [B] (verification not implemented)	280

Optimal result

Integrand size = 15, antiderivative size = 108

$$\int \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx = \frac{3cx\sqrt{\frac{c}{(a+bx^2)^2}}}{8a^2} + \frac{cx\sqrt{\frac{c}{(a+bx^2)^2}}}{4a(a+bx^2)} + \frac{3c\sqrt{\frac{c}{(a+bx^2)^2}}(a+bx^2)\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{8a^{5/2}\sqrt{b}}$$

output

```
3/8*c*x*(c/(b*x^2+a)^2)^(1/2)/a^2+1/4*c*x*(c/(b*x^2+a)^2)^(1/2)/a/(b*x^2+a)
)+3/8*c*(c/(b*x^2+a)^2)^(1/2)*(b*x^2+a)*arctan(b^(1/2)*x/a^(1/2))/a^(5/2)/
b^(1/2)
```

Mathematica [A] (verified)

Time = 1.03 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.77

$$\int \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx = \frac{\left(\frac{c}{(a+bx^2)^2} \right)^{3/2} (a+bx^2) \left(\sqrt{a}\sqrt{bx}(5a+3bx^2) + 3(a+bx^2)^2 \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) \right)}{8a^{5/2}\sqrt{b}}$$

input `Integrate[(c/(a + b*x^2)^2)^(3/2),x]`

output `((c/(a + b*x^2)^2)^(3/2)*(a + b*x^2)*(Sqrt[a]*Sqrt[b]*x*(5*a + 3*b*x^2) + 3*(a + b*x^2)^2*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(8*a^(5/2)*Sqrt[b])`

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.85, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {2045, 215, 215, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx \\
 & \quad \downarrow \text{2045} \\
 & \frac{c(a + bx^2) \sqrt{\frac{c}{(a+bx^2)^2}} \int \frac{1}{\left(\frac{bx^2}{a} + 1\right)^3} dx}{a^3} \\
 & \quad \downarrow \text{215} \\
 & \frac{c(a + bx^2) \sqrt{\frac{c}{(a+bx^2)^2}} \left(\frac{3}{4} \int \frac{1}{\left(\frac{bx^2}{a} + 1\right)^2} dx + \frac{a^2 x}{4(a+bx^2)^2} \right)}{a^3} \\
 & \quad \downarrow \text{215} \\
 & \frac{c(a + bx^2) \sqrt{\frac{c}{(a+bx^2)^2}} \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{1}{\frac{bx^2}{a} + 1} dx + \frac{ax}{2(a+bx^2)} \right) + \frac{a^2 x}{4(a+bx^2)^2} \right)}{a^3} \\
 & \quad \downarrow \text{216} \\
 & \frac{c(a + bx^2) \left(\frac{a^2 x}{4(a+bx^2)^2} + \frac{3}{4} \left(\frac{\sqrt{a} \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{2\sqrt{b}} + \frac{ax}{2(a+bx^2)} \right) \right) \sqrt{\frac{c}{(a+bx^2)^2}}}{a^3}
 \end{aligned}$$

input `Int[(c/(a + b*x^2)^2)^(3/2),x]`

output
$$\frac{(c\sqrt{c/(a + b*x^2)^2}*(a + b*x^2)*((a^2*x)/(4*(a + b*x^2)^2) + (3*((a*x)/(2*(a + b*x^2)) + (\sqrt{a}*\text{ArcTan}[(\sqrt{b}*x)/\sqrt{a}])/(2*\sqrt{b}))))/4)}{a^3}$$

Defintions of rubi rules used

rule 215 `Int[((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && LtQ[p, -1] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 2045 `Int[(u_)*((c_)*((a_) + (b_)*(x_)^(n_))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a)^(p*q)) Int[u*(1 + b*(x^n/a)^(p*q)), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.92

method	result	size
default	$\frac{\left(3 \arctan\left(\frac{bx}{\sqrt{ab}}\right) b^2 x^4 + 3\sqrt{ab} b x^3 + 6 \arctan\left(\frac{bx}{\sqrt{ab}}\right) ab x^2 + 5\sqrt{ab} ax + 3 \arctan\left(\frac{bx}{\sqrt{ab}}\right) a^2\right) (bx^2+a) \left(\frac{c}{(bx^2+a)^2}\right)^{\frac{3}{2}}}{8\sqrt{ab} a^2}$	99
risch	$\frac{c \sqrt{\frac{c}{(bx^2+a)^2}} \left(\frac{3bx^3}{8a^2} + \frac{5x}{8a}\right)}{bx^2+a} - \frac{3c(bx^2+a) \sqrt{\frac{c}{(bx^2+a)^2}} \ln(bx+\sqrt{-ab})}{16\sqrt{-ab} a^2} + \frac{3c(bx^2+a) \sqrt{\frac{c}{(bx^2+a)^2}} \ln(-bx+\sqrt{-ab})}{16\sqrt{-ab} a^2}$	129

input `int((c/(b*x^2+a)^2)^(3/2),x,method=_RETURNVERBOSE)`

output

```
1/8*(3*arctan(b*x/(a*b)^(1/2))*b^2*x^4+3*(a*b)^(1/2)*b*x^3+6*arctan(b*x/(a
*b)^(1/2))*a*b*x^2+5*(a*b)^(1/2)*a*x+3*arctan(b*x/(a*b)^(1/2))*a^2)*(b*x^2
+a)*(c/(b*x^2+a)^2)^(3/2)/(a*b)^(1/2)/a^2
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 282, normalized size of antiderivative = 2.61

$$\int \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \frac{3(bc x^2 + ac) \sqrt{-\frac{c}{ab}} \log \left(\frac{bc x^2 - ac + 2(ab^2 x^3 + a^2 bx) \sqrt{\frac{c}{b^2 x^4 + 2abx^2 + a^2}} \sqrt{-\frac{c}{ab}}}{bx^2 + a} \right) + 2(3bcx^3 + a^2 b^2 x^2 + a^3)}{16(a^2 bx^2 + a^3)}$$

input

```
integrate((c/(b*x^2+a)^2)^(3/2),x, algorithm="fricas")
```

output

```
[1/16*(3*(b*c*x^2 + a*c)*sqrt(-c/(a*b))*log((b*c*x^2 - a*c + 2*(a*b^2*x^3
+ a^2*b*x)*sqrt(c/(b^2*x^4 + 2*a*b*x^2 + a^2))*sqrt(-c/(a*b)))/(b*x^2 + a)
) + 2*(3*b*c*x^3 + 5*a*c*x)*sqrt(c/(b^2*x^4 + 2*a*b*x^2 + a^2)))/(a^2*b*x^
2 + a^3), 1/8*(3*(b*c*x^2 + a*c)*sqrt(c/(a*b))*arctan((b^2*x^3 + a*b*x)*sq
rt(c/(b^2*x^4 + 2*a*b*x^2 + a^2))*sqrt(c/(a*b)))/c + (3*b*c*x^3 + 5*a*c*x)
*sqrt(c/(b^2*x^4 + 2*a*b*x^2 + a^2)))/(a^2*b*x^2 + a^3)]
```

Sympy [F]

$$\int \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \int \left(\frac{c}{(a + bx^2)^2} \right)^{\frac{3}{2}} dx$$

input

```
integrate((c/(b*x**2+a)**2)**(3/2),x)
```

output

```
Integral((c/(a + b*x**2)**2)**(3/2), x)
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.62

$$\int \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \frac{3bc^{3/2}x^3 + 5ac^{3/2}x}{8(a^2b^2x^4 + 2a^3bx^2 + a^4)} + \frac{3c^{3/2} \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{8\sqrt{aba^2}}$$

input `integrate((c/(b*x^2+a)^2)^(3/2),x, algorithm="maxima")`output `1/8*(3*b*c^(3/2)*x^3 + 5*a*c^(3/2)*x)/(a^2*b^2*x^4 + 2*a^3*b*x^2 + a^4) + 3/8*c^(3/2)*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*a^2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.53

$$\int \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \frac{1}{8} c^{3/2} \left(\frac{3 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{aba^2}} + \frac{3bx^3 + 5ax}{(bx^2 + a)^2 a^2} \right) \operatorname{sgn}(bx^2 + a)$$

input `integrate((c/(b*x^2+a)^2)^(3/2),x, algorithm="giac")`output `1/8*c^(3/2)*(3*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*a^2) + (3*b*x^3 + 5*a*x)/((b*x^2 + a)^2*a^2))*sgn(b*x^2 + a)`**Mupad [F(-1)]**

Timed out.

$$\int \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \int \left(\frac{c}{(bx^2 + a)^2} \right)^{3/2} dx$$

input `int((c/(a + b*x^2)^2)^(3/2),x)`output `int((c/(a + b*x^2)^2)^(3/2), x)`

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.07

$$\int \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \frac{\sqrt{c} c \left(3\sqrt{b} \sqrt{a} \operatorname{atan} \left(\frac{bx}{\sqrt{b}\sqrt{a}} \right) a^2 + 6\sqrt{b} \sqrt{a} \operatorname{atan} \left(\frac{bx}{\sqrt{b}\sqrt{a}} \right) abx^2 + 3\sqrt{b} \sqrt{a} \operatorname{atan} \left(\frac{bx}{\sqrt{b}\sqrt{a}} \right) \right)}{8a^3b(b^2x^4 + 2abx^2 + a^2)}$$

input `int((c/(b*x^2+a)^2)^(3/2),x)`output `(sqrt(c)*c*(3*sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*sqrt(a)))*a**2 + 6*sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*sqrt(a)))*a*b*x**2 + 3*sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*sqrt(a)))*b**2*x**4 + 5*a**2*b*x + 3*a*b**2*x**3))/(8*a**3*b*(a**2 + 2*a*b*x**2 + b**2*x**4))`

3.37
$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^2} dx$$

Optimal result	281
Mathematica [A] (verified)	282
Rubi [A] (verified)	282
Maple [C] (verified)	284
Fricas [A] (verification not implemented)	285
Sympy [F]	286
Maxima [A] (verification not implemented)	286
Giac [A] (verification not implemented)	286
Mupad [F(-1)]	287
Reduce [B] (verification not implemented)	287

Optimal result

Integrand size = 19, antiderivative size = 141

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^2} dx = -\frac{7bcx\sqrt{\frac{c}{(a+bx^2)^2}}}{8a^3} - \frac{bcx\sqrt{\frac{c}{(a+bx^2)^2}}}{4a^2(a+bx^2)}$$

$$- \frac{c\sqrt{\frac{c}{(a+bx^2)^2}}(a+bx^2)}{a^3x} - \frac{15\sqrt{bc}\sqrt{\frac{c}{(a+bx^2)^2}}(a+bx^2)\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{8a^{7/2}}$$

output

```
-7/8*b*c*x*(c/(b*x^2+a)^2)^(1/2)/a^3-1/4*b*c*x*(c/(b*x^2+a)^2)^(1/2)/a^2/(
b*x^2+a)-c*(c/(b*x^2+a)^2)^(1/2)*(b*x^2+a)/a^3/x-15/8*b^(1/2)*c*(c/(b*x^2+
a)^2)^(1/2)*(b*x^2+a)*arctan(b^(1/2)*x/a^(1/2))/a^(7/2)
```

Mathematica [A] (verified)

Time = 1.05 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.65

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^2} dx = \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2} (a+bx^2) \left(\sqrt{a}(8a^2+25abx^2+15b^2x^4) + 15\sqrt{b}x(a+bx^2)^2 \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\right)}{8a^{7/2}x}$$

input

```
Integrate[(c/(a + b*x^2)^2)^(3/2)/x^2,x]
```

output

```
-1/8*((c/(a + b*x^2)^2)^(3/2)*(a + b*x^2)*(Sqrt[a]*(8*a^2 + 25*a*b*x^2 + 15*b^2*x^4) + 15*Sqrt[b]*x*(a + b*x^2)^2*ArcTan[(Sqrt[b]*x)/Sqrt[a]]))/(a^(7/2)*x)
```

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.79, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {2045, 27, 253, 253, 264, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^2} dx \\ & \quad \downarrow \text{2045} \\ & \frac{c(a+bx^2) \sqrt{\frac{c}{(a+bx^2)^2}} \int \frac{a^3}{x^2(bx^2+a)^3} dx}{a^3} \\ & \quad \downarrow \text{27} \\ & c(a+bx^2) \sqrt{\frac{c}{(a+bx^2)^2}} \int \frac{1}{x^2(bx^2+a)^3} dx \\ & \quad \downarrow \text{253} \end{aligned}$$

$$\begin{aligned}
& c(a+bx^2) \sqrt{\frac{c}{(a+bx^2)^2}} \left(\frac{5 \int \frac{1}{x^2(bx^2+a)^2} dx}{4a} + \frac{1}{4ax(a+bx^2)^2} \right) \\
& \quad \downarrow \text{253} \\
& c(a+bx^2) \sqrt{\frac{c}{(a+bx^2)^2}} \left(\frac{5 \left(\frac{3 \int \frac{1}{x^2(bx^2+a)} dx}{2a} + \frac{1}{2ax(a+bx^2)} \right)}{4a} + \frac{1}{4ax(a+bx^2)^2} \right) \\
& \quad \downarrow \text{264} \\
& c(a+bx^2) \sqrt{\frac{c}{(a+bx^2)^2}} \left(\frac{5 \left(\frac{3 \left(-\frac{b \int \frac{1}{bx^2+a} dx}{a} - \frac{1}{ax} \right)}{2a} + \frac{1}{2ax(a+bx^2)} \right)}{4a} + \frac{1}{4ax(a+bx^2)^2} \right) \\
& \quad \downarrow \text{218} \\
& c(a+bx^2) \left(\frac{5 \left(\frac{3 \left(-\frac{\sqrt{b} \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{a^{3/2}} - \frac{1}{ax} \right)}{2a} + \frac{1}{2ax(a+bx^2)} \right)}{4a} + \frac{1}{4ax(a+bx^2)^2} \right) \sqrt{\frac{c}{(a+bx^2)^2}}
\end{aligned}$$

input `Int[(c/(a + b*x^2)^2)^(3/2)/x^2,x]`

output `c*Sqrt[c/(a + b*x^2)^2]*(a + b*x^2)*(1/(4*a*x*(a + b*x^2)^2) + (5*(1/(2*a*x*(a + b*x^2)) + (3*(-(1/(a*x)) - (Sqrt[b]*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/a^(3/2)))/(2*a)))/(4*a))`

Defintions of rubi rules used

rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$

rule 218 $\text{Int}[((a_) + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$

rule 253 $\text{Int}[((c_)*(x_))^{(m_)}*((a_) + (b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[(-(c*x)^{(m+1))*((a + b*x^2)^{(p+1})/(2*a*c*(p+1))), x] + \text{Simp}[(m + 2*p + 3)/(2*a*(p+1)) \text{ Int}[(c*x)^m*(a + b*x^2)^{(p+1)}, x], x] /; \text{FreeQ}[\{a, b, c, m\}, x] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 264 $\text{Int}[((c_)*(x_))^{(m_)}*((a_) + (b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[(c*x)^{(m+1))*((a + b*x^2)^{(p+1})/(a*c*(m+1))), x] - \text{Simp}[b*(m + 2*p + 3)/(a*c^{2*(m+1)}) \text{ Int}[(c*x)^{(m+2)}*(a + b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 2045 $\text{Int}[(u_)*((c_)*((a_) + (b_)*(x_)^{(n_)}))^{(q_)}]^{(p_)}, x_Symbol] \rightarrow \text{Simp}[\text{Simp}[(c*(a + b*x^n)^q]^p/(1 + b*(x^n/a))^{(p*q)} \text{ Int}[u*(1 + b*(x^n/a))^{(p*q)}, x], x] /; \text{FreeQ}[\{a, b, c, n, p, q\}, x] \ \&\& \ !\text{GeQ}[a, 0]$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.25 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.81

method	result
risch	$\frac{c \sqrt{\frac{c}{(bx^2+a)^2}} \left(-\frac{15b^2x^4}{8a^3} - \frac{25bx^2}{8a^2} - \frac{1}{a} \right)}{(bx^2+a)x} + \frac{15c(bx^2+a) \sqrt{\frac{c}{(bx^2+a)^2}} \left(\sum_{-R=\text{RootOf}(a^7-Z^2+b)} -R \ln((3a^7-R^2+2b)x+a^4-R) \right)}{16}$
default	$\frac{\left(15 \arctan\left(\frac{bx}{\sqrt{ab}}\right) b^3 x^5 + 30 \arctan\left(\frac{bx}{\sqrt{ab}}\right) a b^2 x^3 + 15 \sqrt{ab} b^2 x^4 + 15 \arctan\left(\frac{bx}{\sqrt{ab}}\right) a^2 b x + 25 \sqrt{ab} a b x^2 + 8 \sqrt{ab} a^2 \right) (bx^2+a) \left(\frac{c}{(bx^2+a)^2} \right)}{8x\sqrt{ab}a^3}$

input `int((c/(b*x^2+a)^2)^(3/2)/x^2,x,method=_RETURNVERBOSE)`

output `c/(b*x^2+a)*(c/(b*x^2+a)^(1/2))*(-15/8*b^2/a^3*x^4-25/8*b/a^2*x^2-1/a)/x+15/16*c*(b*x^2+a)*(c/(b*x^2+a)^(1/2))*sum(_R*ln((3*_R^2*a^7+2*b)*x+a^4*_R),_R=RootOf(_Z^2*a^7+b))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 296, normalized size of antiderivative = 2.10

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^2} dx = \left[\frac{15 (bcx^3 + acx) \sqrt{-\frac{bc}{a}} \log\left(\frac{bcx^2 - ac - 2(abx^3 + a^2x) \sqrt{-\frac{bc}{a}} \sqrt{\frac{c}{b^2x^4 + 2abx^2 + a^2}}}{bx^2 + a}\right) - 2(15b^2cx^4 + 15bcx^3 + acx) \sqrt{\frac{bc}{a}} \arctan\left(\frac{(bx^3 + ax) \sqrt{\frac{bc}{a}} \sqrt{\frac{c}{b^2x^4 + 2abx^2 + a^2}}}{c}\right) + (15b^2cx^4 + 25abcx^2 + 8a^2c) \sqrt{\frac{c}{b^2x^4 + 2abx^2 + a^2}}}{16(a^3bx^3 + a^4x)} \right]$$

input `integrate((c/(b*x^2+a)^2)^(3/2)/x^2,x, algorithm="fricas")`

output `[1/16*(15*(b*c*x^3 + a*c*x)*sqrt(-b*c/a)*log((b*c*x^2 - a*c - 2*(a*b*x^3 + a^2*x)*sqrt(-b*c/a)*sqrt(c/(b^2*x^4 + 2*a*b*x^2 + a^2)))/(b*x^2 + a)) - 2*(15*b^2*c*x^4 + 25*a*b*c*x^2 + 8*a^2*c)*sqrt(c/(b^2*x^4 + 2*a*b*x^2 + a^2)))/(a^3*b*x^3 + a^4*x), -1/8*(15*(b*c*x^3 + a*c*x)*sqrt(b*c/a)*arctan((b*x^3 + a*x)*sqrt(b*c/a)*sqrt(c/(b^2*x^4 + 2*a*b*x^2 + a^2)))/c) + (15*b^2*c*x^4 + 25*a*b*c*x^2 + 8*a^2*c)*sqrt(c/(b^2*x^4 + 2*a*b*x^2 + a^2)))/(a^3*b*x^3 + a^4*x)]`

Sympy [F]

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^2} dx = \int \frac{\left(\frac{c}{a^2+2abx^2+b^2x^4}\right)^{3/2}}{x^2} dx$$

input `integrate((c/(b*x**2+a)**2)**(3/2)/x**2,x)`

output `Integral((c/(a**2 + 2*a*b*x**2 + b**2*x**4))**(3/2)/x**2, x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.59

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^2} dx = -\frac{15b^2c^{3/2}x^4 + 25abc^{3/2}x^2 + 8a^2c^{3/2}}{8(a^3b^2x^5 + 2a^4bx^3 + a^5x)} - \frac{15bc^{3/2} \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{8\sqrt{aba^3}}$$

input `integrate((c/(b*x^2+a)^2)^(3/2)/x^2,x, algorithm="maxima")`

output `-1/8*(15*b^2*c^(3/2)*x^4 + 25*a*b*c^(3/2)*x^2 + 8*a^2*c^(3/2))/(a^3*b^2*x^5 + 2*a^4*b*x^3 + a^5*x) - 15/8*b*c^(3/2)*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*a^3)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.49

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^2} dx = -\frac{1}{8}c^{3/2} \left(\frac{15b \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{aba^3}} + \frac{7b^2x^3 + 9abx}{(bx^2 + a)^2a^3} + \frac{8}{a^3x} \right) \operatorname{sgn}(bx^2 + a)$$

input `integrate((c/(b*x^2+a)^2)^(3/2)/x^2,x, algorithm="giac")`

output

$$-1/8*c^{(3/2)}*(15*b*\arctan(b*x/\sqrt{a*b})/(\sqrt{a*b})*a^3) + (7*b^2*x^3 + 9*a*b*x)/((b*x^2 + a)^2*a^3) + 8/(a^3*x))*\operatorname{sgn}(b*x^2 + a)$$

Mupad [F(-1)]

Timed out.

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^2} dx = \int \frac{\left(\frac{c}{(bx^2+a)^2}\right)^{3/2}}{x^2} dx$$

input

$$\operatorname{int}((c/(a + b*x^2)^2)^{(3/2)}/x^2,x)$$

output

$$\operatorname{int}((c/(a + b*x^2)^2)^{(3/2)}/x^2, x)$$

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.88

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^2} dx = \frac{\sqrt{c}c\left(-15\sqrt{b}\sqrt{a}\operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right)a^2x - 30\sqrt{b}\sqrt{a}\operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right)abx^3 - 15\sqrt{b}\sqrt{a}\operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right)a^2x^5 - 8a^3 - 25a^2bx^2 - 15abx^4\right)}{8a^4x(b^2x^4 + 2abx^2 + a^2)}$$

input

$$\operatorname{int}((c/(b*x^2+a)^2)^{(3/2)}/x^2,x)$$

output

$$\left(\sqrt{c}c\left(-15\sqrt{b}\sqrt{a}\operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right)a^2x - 30\sqrt{b}\sqrt{a}\operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right)abx^3 - 15\sqrt{b}\sqrt{a}\operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right)a^2x^5 - 8a^3 - 25a^2bx^2 - 15abx^4\right)\right)/\left(8a^4x\left(a^2 + 2abx^2 + b^2x^4\right)\right)$$

3.38
$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^4} dx$$

Optimal result	288
Mathematica [A] (verified)	289
Rubi [A] (verified)	289
Maple [A] (verified)	292
Fricas [A] (verification not implemented)	293
Sympy [F]	293
Maxima [A] (verification not implemented)	294
Giac [A] (verification not implemented)	294
Mupad [F(-1)]	295
Reduce [B] (verification not implemented)	295

Optimal result

Integrand size = 19, antiderivative size = 179

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^4} dx = \frac{11b^2cx\sqrt{\frac{c}{(a+bx^2)^2}}}{8a^4} + \frac{b^2cx\sqrt{\frac{c}{(a+bx^2)^2}}}{4a^3(a+bx^2)} - \frac{c\sqrt{\frac{c}{(a+bx^2)^2}}(a+bx^2)}{3a^3x^3} + \frac{3bc\sqrt{\frac{c}{(a+bx^2)^2}}(a+bx^2)}{a^4x} + \frac{35b^{3/2}c\sqrt{\frac{c}{(a+bx^2)^2}}(a+bx^2)\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{8a^{9/2}}$$

output

```
11/8*b^2*c*x*(c/(b*x^2+a)^2)^(1/2)/a^4+1/4*b^2*c*x*(c/(b*x^2+a)^2)^(1/2)/a^3/(b*x^2+a)-1/3*c*(c/(b*x^2+a)^2)^(1/2)*(b*x^2+a)/a^3/x^3+3*b*c*(c/(b*x^2+a)^2)^(1/2)*(b*x^2+a)/a^4/x+35/8*b^(3/2)*c*(c/(b*x^2+a)^2)^(1/2)*(b*x^2+a)*arctan(b^(1/2)*x/a^(1/2))/a^(9/2)
```

Mathematica [A] (verified)

Time = 1.04 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.59

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^4} dx = \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2} (a+bx^2) \left(\sqrt{a}(8a^3 - 56a^2bx^2 - 175ab^2x^4 - 105b^3x^6) - 105b^{3/2}x^3(a+bx^2)^2 \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\right)}{24a^{9/2}x^3}$$

input

```
Integrate[(c/(a + b*x^2)^2)^(3/2)/x^4,x]
```

output

```
-1/24*((c/(a + b*x^2)^2)^(3/2)*(a + b*x^2)*(Sqrt[a]*(8*a^3 - 56*a^2*b*x^2 - 175*a*b^2*x^4 - 105*b^3*x^6) - 105*b^(3/2)*x^3*(a + b*x^2)^2*ArcTan[(Sqrt[b]*x)/Sqrt[a]]))/(a^(9/2)*x^3)
```

Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.72, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {2045, 27, 253, 253, 264, 264, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^4} dx \\ & \quad \downarrow \text{2045} \\ & \frac{c(a+bx^2) \sqrt{\frac{c}{(a+bx^2)^2}} \int \frac{a^3}{x^4(bx^2+a)^3} dx}{a^3} \\ & \quad \downarrow \text{27} \\ & c(a+bx^2) \sqrt{\frac{c}{(a+bx^2)^2}} \int \frac{1}{x^4(bx^2+a)^3} dx \\ & \quad \downarrow \text{253} \end{aligned}$$

$$c(a + bx^2) \sqrt{\frac{c}{(a + bx^2)^2}} \left(\frac{7 \int \frac{1}{x^4(bx^2+a)^2} dx}{4a} + \frac{1}{4ax^3(a + bx^2)^2} \right)$$

↓ 253

$$c(a + bx^2) \sqrt{\frac{c}{(a + bx^2)^2}} \left(\frac{7 \left(\frac{5 \int \frac{1}{x^4(bx^2+a)} dx}{2a} + \frac{1}{2ax^3(a+bx^2)} \right)}{4a} + \frac{1}{4ax^3(a + bx^2)^2} \right)$$

↓ 264

$$c(a + bx^2) \sqrt{\frac{c}{(a + bx^2)^2}} \left(\frac{7 \left(\frac{5 \left(-\frac{b \int \frac{1}{x^2(bx^2+a)} dx}{a} - \frac{1}{3ax^3} \right)}{2a} + \frac{1}{2ax^3(a+bx^2)} \right)}{4a} + \frac{1}{4ax^3(a + bx^2)^2} \right)$$

↓ 264

$$c(a + bx^2) \sqrt{\frac{c}{(a + bx^2)^2}} \left(\frac{7 \left(\frac{5 \left(-\frac{b \left(-\frac{b \int \frac{1}{bx^2+a} dx}{a} - \frac{1}{ax} \right)}{a} - \frac{1}{3ax^3} \right)}{2a} + \frac{1}{2ax^3(a+bx^2)} \right)}{4a} + \frac{1}{4ax^3(a + bx^2)^2} \right)$$

↓ 218

$$c(a + bx^2) \left(\frac{7 \left(\frac{5 \left(b \left(-\frac{\sqrt{b} \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) - \frac{1}{ax}}{a^{3/2}} \right) - \frac{1}{3ax^3} \right)}{a} \right)}{2a} + \frac{1}{2ax^3(a+bx^2)} \right)}{4a} + \frac{1}{4ax^3(a+bx^2)^2} \sqrt{\frac{c}{(a+bx^2)^2}} \right)$$

input `Int[(c/(a + b*x^2)^2)^(3/2)/x^4,x]`

output `c*Sqrt[c/(a + b*x^2)^2]*(a + b*x^2)*(1/(4*a*x^3*(a + b*x^2)^2) + (7*(1/(2*a*x^3*(a + b*x^2)) + (5*(-1/3*1/(a*x^3) - (b*(-1/(a*x)) - (Sqrt[b]*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/a^(3/2)))/a))/(2*a)))/(4*a))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 253 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-(c*x)^(m + 1))*((a + b*x^2)^(p + 1)/(2*a*c*(p + 1))), x] + Simp[(m + 2*p + 3)/(2*a*(p + 1)) Int[(c*x)^m*(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, m}, x] && LtQ[p, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 264 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*(m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_.))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.79

method	result
default	$\frac{\left(-105 \arctan\left(\frac{bx}{\sqrt{ab}}\right) b^4 x^7 - 105 \sqrt{ab} b^3 x^6 - 210 \arctan\left(\frac{bx}{\sqrt{ab}}\right) a b^3 x^5 - 175 \sqrt{ab} a b^2 x^4 - 105 \arctan\left(\frac{bx}{\sqrt{ab}}\right) a^2 b^2 x^3 - 56 \sqrt{ab} a^2 b x^2 + 8 a^3 x\right)}{24 x^3 \sqrt{ab} a^4}$
risch	$\frac{c \sqrt{\frac{c}{(b x^2+a)^2}} \left(\frac{35 b^3 x^6}{8 a^4} + \frac{175 b^2 x^4}{24 a^3} + \frac{7 b x^2}{3 a^2} - \frac{1}{3 a}\right)}{(b x^2+a) x^3} + \frac{35 c (b x^2+a) \sqrt{\frac{c}{(b x^2+a)^2}} \sqrt{-a b} b \ln(-b x-\sqrt{-a b})}{16 a^5} - \frac{35 c (b x^2+a) \sqrt{\frac{c}{(b x^2+a)^2}}}{16 a^5}$

input `int((c/(b*x^2+a)^2)^(3/2)/x^4,x,method=_RETURNVERBOSE)`

output `-1/24*(-105*arctan(b*x/(a*b)^(1/2))*b^4*x^7-105*(a*b)^(1/2)*b^3*x^6-210*arctan(b*x/(a*b)^(1/2))*a*b^3*x^5-175*(a*b)^(1/2)*a*b^2*x^4-105*arctan(b*x/(a*b)^(1/2))*a^2*b^2*x^3-56*(a*b)^(1/2)*a^2*b*x^2+8*(a*b)^(1/2)*a^3)*(b*x^2+a)*(c/(b*x^2+a)^2)^(3/2)/x^3/(a*b)^(1/2)/a^4`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 334, normalized size of antiderivative = 1.87

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^4} dx = \frac{105(b^2cx^5 + abcx^3)\sqrt{-\frac{bc}{a}} \log\left(\frac{bcx^2 - ac + 2(abx^3 + a^2x)\sqrt{-\frac{bc}{a}}\sqrt{\frac{c}{b^2x^4 + 2abx^2 + a^2}}}{bx^2 + a}\right) + 2(105b^3c}{48(a^4bx^5 + a^5x^3)}$$

input `integrate((c/(b*x^2+a)^2)^(3/2)/x^4,x, algorithm="fricas")`

output `[1/48*(105*(b^2*c*x^5 + a*b*c*x^3)*sqrt(-b*c/a)*log((b*c*x^2 - a*c + 2*(a*b*x^3 + a^2*x)*sqrt(-b*c/a)*sqrt(c/(b^2*x^4 + 2*a*b*x^2 + a^2)))/(b*x^2 + a)) + 2*(105*b^3*c*x^6 + 175*a*b^2*c*x^4 + 56*a^2*b*c*x^2 - 8*a^3*c)*sqrt(c/(b^2*x^4 + 2*a*b*x^2 + a^2)))/(a^4*b*x^5 + a^5*x^3), 1/24*(105*(b^2*c*x^5 + a*b*c*x^3)*sqrt(b*c/a)*arctan((b*x^3 + a*x)*sqrt(b*c/a)*sqrt(c/(b^2*x^4 + 2*a*b*x^2 + a^2)))/c) + (105*b^3*c*x^6 + 175*a*b^2*c*x^4 + 56*a^2*b*c*x^2 - 8*a^3*c)*sqrt(c/(b^2*x^4 + 2*a*b*x^2 + a^2)))/(a^4*b*x^5 + a^5*x^3)]`

Sympy [F]

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^4} dx = \int \frac{\left(\frac{c}{a^2+2abx^2+b^2x^4}\right)^{\frac{3}{2}}}{x^4} dx$$

input `integrate((c/(b*x**2+a)**2)**(3/2)/x**4,x)`

output `Integral((c/(a**2 + 2*a*b*x**2 + b**2*x**4))**(3/2)/x**4, x)`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.56

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^4} dx = \frac{105 b^3 c^{\frac{3}{2}} x^6 + 175 ab^2 c^{\frac{3}{2}} x^4 + 56 a^2 b c^{\frac{3}{2}} x^2 - 8 a^3 c^{\frac{3}{2}}}{24 (a^4 b^2 x^7 + 2 a^5 b x^5 + a^6 x^3)} + \frac{35 b^2 c^{\frac{3}{2}} \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{8 \sqrt{aba^4}}$$

input `integrate((c/(b*x^2+a)^2)^(3/2)/x^4,x, algorithm="maxima")`output `1/24*(105*b^3*c^(3/2)*x^6 + 175*a*b^2*c^(3/2)*x^4 + 56*a^2*b*c^(3/2)*x^2 - 8*a^3*c^(3/2))/(a^4*b^2*x^7 + 2*a^5*b*x^5 + a^6*x^3) + 35/8*b^2*c^(3/2)*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*a^4)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.47

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^4} dx = \frac{1}{24} c^{\frac{3}{2}} \left(\frac{105 b^2 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{aba^4}} + \frac{3(11 b^3 x^3 + 13 ab^2 x)}{(bx^2 + a)^2 a^4} + \frac{8(9 bx^2 - a)}{a^4 x^3} \right) \operatorname{sgn}(bx^2 + a)$$

input `integrate((c/(b*x^2+a)^2)^(3/2)/x^4,x, algorithm="giac")`output `1/24*c^(3/2)*(105*b^2*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*a^4) + 3*(11*b^3*x^3 + 13*a*b^2*x)/((b*x^2 + a)^2*a^4) + 8*(9*b*x^2 - a)/(a^4*x^3))*sgn(b*x^2 + a)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^4} dx = \int \frac{\left(\frac{c}{(bx^2+a)^2}\right)^{3/2}}{x^4} dx$$

input `int((c/(a + b*x^2)^2)^(3/2)/x^4,x)`output `int((c/(a + b*x^2)^2)^(3/2)/x^4, x)`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.78

$$\int \frac{\left(\frac{c}{(a+bx^2)^2}\right)^{3/2}}{x^4} dx = \frac{\sqrt{c} c \left(105\sqrt{b} \sqrt{a} \operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right) a^2 b x^3 + 210\sqrt{b} \sqrt{a} \operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right) a b^2 x^5 + 105\sqrt{b} \sqrt{a} \operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right) a^2 b^2 x^7 - 8a^4 + 56a^3 b x^2 + 175a^2 b^2 x^4 + 105a b^3 x^6\right)}{24a^5 x^3 (b^2 x^4 + 2ab x^2 + a^2)}$$

input `int((c/(b*x^2+a)^2)^(3/2)/x^4,x)`output `(sqrt(c)*c*(105*sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*sqrt(a)))*a**2*b*x**3 + 210*sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*sqrt(a)))*a*b**2*x**5 + 105*sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*sqrt(a)))*b**3*x**7 - 8*a**4 + 56*a**3*b*x**2 + 175*a**2*b**2*x**4 + 105*a*b**3*x**6)/(24*a**5*x**3*(a**2 + 2*a*b*x**2 + b**2*x**4))`

3.39 $\int x^7 \left(c\sqrt{a + bx^2} \right)^{3/2} dx$

Optimal result	296
Mathematica [A] (verified)	297
Rubi [A] (verified)	297
Maple [A] (verified)	299
Fricas [A] (verification not implemented)	299
Sympy [A] (verification not implemented)	299
Maxima [A] (verification not implemented)	300
Giac [A] (verification not implemented)	300
Mupad [B] (verification not implemented)	301
Reduce [B] (verification not implemented)	301

Optimal result

Integrand size = 21, antiderivative size = 138

$$\int x^7 \left(c\sqrt{a + bx^2} \right)^{3/2} dx = -\frac{2a^3 (c\sqrt{a + bx^2})^{3/2} (a + bx^2)}{7b^4} + \frac{6a^2 (c\sqrt{a + bx^2})^{3/2} (a + bx^2)^2}{11b^4} - \frac{2a (c\sqrt{a + bx^2})^{3/2} (a + bx^2)^3}{5b^4} + \frac{2 (c\sqrt{a + bx^2})^{3/2} (a + bx^2)^4}{19b^4}$$

output -2/7*a^3*(c*(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)/b^4+6/11*a^2*(c*(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)^2/b^4-2/5*a*(c*(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)^3/b^4+2/19*(c*(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)^4/b^4

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.46

$$\int x^7 \left(c\sqrt{a+bx^2} \right)^{3/2} dx = \frac{2(c\sqrt{a+bx^2})^{3/2} (a+bx^2) (-128a^3 + 224a^2bx^2 - 308ab^2x^4 + 385b^3x^6)}{7315b^4}$$

input `Integrate[x^7*(c*Sqrt[a + b*x^2])^(3/2),x]`

output `(2*(c*Sqrt[a + b*x^2])^(3/2)*(a + b*x^2)*(-128*a^3 + 224*a^2*b*x^2 - 308*a*b^2*x^4 + 385*b^3*x^6))/(7315*b^4)`

Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.96, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2045, 243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^7 \left(c\sqrt{a+bx^2} \right)^{3/2} dx \\ & \quad \downarrow \text{2045} \\ & \frac{\left(c\sqrt{a+bx^2} \right)^{3/2} \int x^7 \left(\frac{bx^2}{a} + 1 \right)^{3/4} dx}{\left(\frac{bx^2}{a} + 1 \right)^{3/4}} \\ & \quad \downarrow \text{243} \\ & \frac{\left(c\sqrt{a+bx^2} \right)^{3/2} \int x^6 \left(\frac{bx^2}{a} + 1 \right)^{3/4} dx^2}{2 \left(\frac{bx^2}{a} + 1 \right)^{3/4}} \\ & \quad \downarrow \text{53} \end{aligned}$$

$$\frac{(c\sqrt{a+bx^2})^{3/2} \int \left(\frac{a^3 \left(\frac{bx^2}{a} + 1\right)^{15/4}}{b^3} - \frac{3a^3 \left(\frac{bx^2}{a} + 1\right)^{11/4}}{b^3} + \frac{3a^3 \left(\frac{bx^2}{a} + 1\right)^{7/4}}{b^3} - \frac{a^3 \left(\frac{bx^2}{a} + 1\right)^{3/4}}{b^3} \right) dx^2}{2 \left(\frac{bx^2}{a} + 1\right)^{3/4}}$$

↓ 2009

$$\frac{\left(\frac{4a^4 \left(\frac{bx^2}{a} + 1\right)^{19/4}}{19b^4} - \frac{4a^4 \left(\frac{bx^2}{a} + 1\right)^{15/4}}{5b^4} + \frac{12a^4 \left(\frac{bx^2}{a} + 1\right)^{11/4}}{11b^4} - \frac{4a^4 \left(\frac{bx^2}{a} + 1\right)^{7/4}}{7b^4} \right) (c\sqrt{a+bx^2})^{3/2}}{2 \left(\frac{bx^2}{a} + 1\right)^{3/4}}$$

input `Int[x^7*(c*Sqrt[a + b*x^2])^(3/2),x]`

output `((c*Sqrt[a + b*x^2])^(3/2)*((-4*a^4*(1 + (b*x^2)/a)^(7/4))/(7*b^4) + (12*a^4*(1 + (b*x^2)/a)^(11/4))/(11*b^4) - (4*a^4*(1 + (b*x^2)/a)^(15/4))/(5*b^4) + (4*a^4*(1 + (b*x^2)/a)^(19/4))/(19*b^4)))/(2*(1 + (b*x^2)/a)^(3/4))`

Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_.))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.42

method	result	size
gospers	$-\frac{2(bx^2+a)(-385b^3x^6+308ab^2x^4-224a^2bx^2+128a^3)(c\sqrt{bx^2+a})^{\frac{3}{2}}}{7315b^4}$	58
orering	$-\frac{2(bx^2+a)(-385b^3x^6+308ab^2x^4-224a^2bx^2+128a^3)(c\sqrt{bx^2+a})^{\frac{3}{2}}}{7315b^4}$	58

input `int(x^7*(c*(b*x^2+a)^(1/2))^(3/2),x,method=_RETURNVERBOSE)`output
$$-2/7315*(b*x^2+a)*(-385*b^3*x^6+308*a*b^2*x^4-224*a^2*b*x^2+128*a^3)*(c*(b*x^2+a)^(1/2))^(3/2)/b^4$$
Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.54

$$\int x^7 (c\sqrt{a+bx^2})^{3/2} dx = \frac{2(385b^4cx^8 + 77ab^3cx^6 - 84a^2b^2cx^4 + 96a^3bcx^2 - 128a^4c)\sqrt{bx^2+a}\sqrt{\sqrt{bx^2+a}}}{7315b^4}$$

input `integrate(x^7*(c*(b*x^2+a)^(1/2))^(3/2),x, algorithm="fricas")`output
$$2/7315*(385*b^4*c*x^8 + 77*a*b^3*c*x^6 - 84*a^2*b^2*c*x^4 + 96*a^3*b*c*x^2 - 128*a^4*c)*\text{sqrt}(b*x^2 + a)*\text{sqrt}(\text{sqrt}(b*x^2 + a)*c)/b^4$$
Sympy [A] (verification not implemented)

Time = 10.96 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.04

$$\int x^7 (c\sqrt{a+bx^2})^{3/2} dx = \left\{ \begin{array}{l} -\frac{256a^4(c\sqrt{a+bx^2})^{\frac{3}{2}}}{7315b^4} + \frac{192a^3x^2(c\sqrt{a+bx^2})^{\frac{3}{2}}}{7315b^3} - \frac{24a^2x^4(c\sqrt{a+bx^2})^{\frac{3}{2}}}{1045b^2} + \frac{2ax^6(c\sqrt{a+bx^2})^{\frac{3}{2}}}{95b} + \frac{x^8(\sqrt{ac})^{\frac{3}{2}}}{8} \end{array} \right.$$

input `integrate(x**7*(c*(b*x**2+a)**(1/2))**(3/2),x)`

output `Piecewise((-256*a**4*(c*sqrt(a + b*x**2))**(3/2)/(7315*b**4) + 192*a**3*x*
2(c*sqrt(a + b*x**2))**(3/2)/(7315*b**3) - 24*a**2*x**4*(c*sqrt(a + b*x*
*2))**(3/2)/(1045*b**2) + 2*a*x**6*(c*sqrt(a + b*x**2))**(3/2)/(95*b) + 2*
x**8*(c*sqrt(a + b*x**2))**(3/2)/19, Ne(b, 0)), (x**8*(sqrt(a)*c)**(3/2)/8
, True))`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.62

$$\int x^7 \left(c\sqrt{a + bx^2} \right)^{3/2} dx = \frac{2 \left(1045 (\sqrt{bx^2 + ac})^{7/2} a^3 c^6 - 1995 (\sqrt{bx^2 + ac})^{11/2} a^2 c^4 + 1463 (\sqrt{bx^2 + ac})^{15/2} ac^2 - 385 (\sqrt{bx^2 + ac})^{19/2} \right)}{7315 b^4 c^8}$$

input `integrate(x^7*(c*(b*x^2+a)^(1/2))^(3/2),x, algorithm="maxima")`

output `-2/7315*(1045*(sqrt(b*x^2 + a)*c)^(7/2)*a^3*c^6 - 1995*(sqrt(b*x^2 + a)*c)
^(11/2)*a^2*c^4 + 1463*(sqrt(b*x^2 + a)*c)^(15/2)*a*c^2 - 385*(sqrt(b*x^2
+ a)*c)^(19/2))/(b^4*c^8)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.99

$$\int x^7 \left(c\sqrt{a + bx^2} \right)^{3/2} dx = \frac{2 c^{3/2} \left(\frac{19 \left(77 (bx^2+a)^{15/4} - 315 (bx^2+a)^{11/4} a + 495 (bx^2+a)^{7/4} a^2 - 385 (bx^2+a)^{3/4} a^3 \right) a}{b^3} + \frac{1155 (bx^2+a)^{19/4}}{21945 b} \right)}{21945 b}$$

input `integrate(x^7*(c*(b*x^2+a)^(1/2))^(3/2),x, algorithm="giac")`

output

```
2/21945*c^(3/2)*(19*(77*(b*x^2 + a)^(15/4) - 315*(b*x^2 + a)^(11/4)*a + 49
5*(b*x^2 + a)^(7/4)*a^2 - 385*(b*x^2 + a)^(3/4)*a^3)*a/b^3 + (1155*(b*x^2
+ a)^(19/4) - 5852*(b*x^2 + a)^(15/4)*a + 11970*(b*x^2 + a)^(11/4)*a^2 - 1
2540*(b*x^2 + a)^(7/4)*a^3 + 7315*(b*x^2 + a)^(3/4)*a^4)/b^3)/b
```

Mupad [B] (verification not implemented)

Time = 0.56 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.79

$$\int x^7 (c\sqrt{a+bx^2})^{3/2} dx = \sqrt{c\sqrt{bx^2+a}} \left(\frac{2cx^8\sqrt{bx^2+a}}{19} - \frac{256a^4c\sqrt{bx^2+a}}{7315b^4} \right) + \frac{2acx^6\sqrt{bx^2+a}}{95b} - \frac{24a^2cx^4\sqrt{bx^2+a}}{1045b^2} + \frac{192a^3cx^2\sqrt{bx^2+a}}{7315b^3}$$

input

```
int(x^7*(c*(a + b*x^2)^(1/2))^(3/2),x)
```

output

```
(c*(a + b*x^2)^(1/2))^(1/2)*((2*c*x^8*(a + b*x^2)^(1/2))/19 - (256*a^4*c*(
a + b*x^2)^(1/2))/(7315*b^4) + (2*a*c*x^6*(a + b*x^2)^(1/2))/(95*b) - (24*
a^2*c*x^4*(a + b*x^2)^(1/2))/(1045*b^2) + (192*a^3*c*x^2*(a + b*x^2)^(1/2)
)/(7315*b^3))
```

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.39

$$\int x^7 (c\sqrt{a+bx^2})^{3/2} dx = \frac{2\sqrt{c}\sqrt{\sqrt{b}\sqrt{bx^2+a}x+a+bx^2}\sqrt{\sqrt{bx^2+a}+\sqrt{b}x}c(128\sqrt{b}\sqrt{bx^2+a}a^4x -$$

input

```
int(x^7*(c*(b*x^2+a)^(1/2))^(3/2),x)
```

output

```
(2*sqrt(c)*sqrt(sqrt(b)*sqrt(a + b*x**2))*x + a + b*x**2)*sqrt(sqrt(a + b*x**2) + sqrt(b)*x)*c*(128*sqrt(b)*sqrt(a + b*x**2)*a**4*x - 96*sqrt(b)*sqrt(a + b*x**2)*a**3*b*x**3 + 84*sqrt(b)*sqrt(a + b*x**2)*a**2*b**2*x**5 - 77*sqrt(b)*sqrt(a + b*x**2)*a*b**3*x**7 - 385*sqrt(b)*sqrt(a + b*x**2)*b**4*x**9 - 128*a**5 - 32*a**4*b*x**2 + 12*a**3*b**2*x**4 - 7*a**2*b**3*x**6 + 462*a*b**4*x**8 + 385*b**5*x**10))/(7315*a*b**4)
```

3.40 $\int x^5 \left(c\sqrt{a + bx^2} \right)^{3/2} dx$

Optimal result	303
Mathematica [A] (verified)	303
Rubi [A] (verified)	304
Maple [A] (verified)	305
Fricas [A] (verification not implemented)	306
Sympy [A] (verification not implemented)	306
Maxima [A] (verification not implemented)	307
Giac [A] (verification not implemented)	307
Mupad [B] (verification not implemented)	307
Reduce [B] (verification not implemented)	308

Optimal result

Integrand size = 21, antiderivative size = 102

$$\int x^5 \left(c\sqrt{a + bx^2} \right)^{3/2} dx = \frac{2a^2 (c\sqrt{a + bx^2})^{3/2} (a + bx^2)}{7b^3} - \frac{4a (c\sqrt{a + bx^2})^{3/2} (a + bx^2)^2}{11b^3} + \frac{2 (c\sqrt{a + bx^2})^{3/2} (a + bx^2)^3}{15b^3}$$

```
output 2/7*a^2*(c*(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)/b^3-4/11*a*(c*(b*x^2+a)^(1/2))
^(3/2)*(b*x^2+a)^2/b^3+2/15*(c*(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)^3/b^3
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.51

$$\int x^5 \left(c\sqrt{a + bx^2} \right)^{3/2} dx = \frac{2 (c\sqrt{a + bx^2})^{3/2} (a + bx^2) (32a^2 - 56abx^2 + 77b^2x^4)}{1155b^3}$$

```
input Integrate[x^5*(c*Sqrt[a + b*x^2])^(3/2),x]
```


output

```
(2*(c*Sqrt[a + b*x^2])^(3/2)*(a + b*x^2)*(32*a^2 - 56*a*b*x^2 + 77*b^2*x^4
))/(1155*b^3)
```

Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.06, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2045, 243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^5 (c\sqrt{a + bx^2})^{3/2} dx \\
 & \quad \downarrow \text{2045} \\
 & \frac{(c\sqrt{a + bx^2})^{3/2} \int x^5 \left(\frac{bx^2}{a} + 1\right)^{3/4} dx}{\left(\frac{bx^2}{a} + 1\right)^{3/4}} \\
 & \quad \downarrow \text{243} \\
 & \frac{(c\sqrt{a + bx^2})^{3/2} \int x^4 \left(\frac{bx^2}{a} + 1\right)^{3/4} dx^2}{2 \left(\frac{bx^2}{a} + 1\right)^{3/4}} \\
 & \quad \downarrow \text{53} \\
 & \frac{(c\sqrt{a + bx^2})^{3/2} \int \left(\frac{a^2 \left(\frac{bx^2}{a} + 1\right)^{11/4}}{b^2} - \frac{2a^2 \left(\frac{bx^2}{a} + 1\right)^{7/4}}{b^2} + \frac{a^2 \left(\frac{bx^2}{a} + 1\right)^{3/4}}{b^2} \right) dx^2}{2 \left(\frac{bx^2}{a} + 1\right)^{3/4}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{\left(\frac{4a^3 \left(\frac{bx^2}{a} + 1\right)^{15/4}}{15b^3} - \frac{8a^3 \left(\frac{bx^2}{a} + 1\right)^{11/4}}{11b^3} + \frac{4a^3 \left(\frac{bx^2}{a} + 1\right)^{7/4}}{7b^3} \right) (c\sqrt{a + bx^2})^{3/2}}{2 \left(\frac{bx^2}{a} + 1\right)^{3/4}}
 \end{aligned}$$

input `Int[x^5*(c*Sqrt[a + b*x^2])^(3/2),x]`

output `((c*Sqrt[a + b*x^2])^(3/2)*((4*a^3*(1 + (b*x^2)/a)^(7/4))/(7*b^3) - (8*a^3*(1 + (b*x^2)/a)^(11/4))/(11*b^3) + (4*a^3*(1 + (b*x^2)/a)^(15/4))/(15*b^3)))/(2*(1 + (b*x^2)/a)^(3/4))`

Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a)^(p*q)) Int[u*(1 + b*(x^n/a)^(p*q)), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.46

method	result	size
gospers	$\frac{2(bx^2+a)(77b^2x^4-56abx^2+32a^2)(c\sqrt{bx^2+a})^{\frac{3}{2}}}{1155b^3}$	47
orering	$\frac{2(bx^2+a)(77b^2x^4-56abx^2+32a^2)(c\sqrt{bx^2+a})^{\frac{3}{2}}}{1155b^3}$	47

input `int(x^5*(c*(b*x^2+a)^(1/2))^(3/2),x,method=_RETURNVERBOSE)`

output `2/1155*(b*x^2+a)*(77*b^2*x^4-56*a*b*x^2+32*a^2)*(c*(b*x^2+a)^(1/2))^(3/2)/b^3`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.62

$$\int x^5 \left(c\sqrt{a+bx^2} \right)^{3/2} dx = \frac{2(77b^3cx^6 + 21ab^2cx^4 - 24a^2bcx^2 + 32a^3c)\sqrt{bx^2+a}\sqrt{\sqrt{bx^2+a}c}}{1155b^3}$$

input `integrate(x^5*(c*(b*x^2+a)^(1/2))^(3/2),x, algorithm="fricas")`

output `2/1155*(77*b^3*c*x^6 + 21*a*b^2*c*x^4 - 24*a^2*b*c*x^2 + 32*a^3*c)*sqrt(b*x^2 + a)*sqrt(sqrt(b*x^2 + a)*c)/b^3`

Sympy [A] (verification not implemented)

Time = 6.10 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.14

$$\int x^5 \left(c\sqrt{a+bx^2} \right)^{3/2} dx = \begin{cases} \frac{64a^3 \left(c\sqrt{a+bx^2} \right)^{3/2}}{1155b^3} - \frac{16a^2x^2 \left(c\sqrt{a+bx^2} \right)^{3/2}}{385b^2} + \frac{2ax^4 \left(c\sqrt{a+bx^2} \right)^{3/2}}{55b} + \frac{2x^6 \left(c\sqrt{a+bx^2} \right)^{3/2}}{15} & \text{for } b \neq 0 \\ \frac{x^6 (\sqrt{ac})^{3/2}}{6} & \text{otherwise} \end{cases}$$

input `integrate(x**5*(c*(b*x**2+a)**(1/2))**(3/2),x)`

output `Piecewise((64*a**3*(c*sqrt(a + b*x**2))**(3/2)/(1155*b**3) - 16*a**2*x**2*(c*sqrt(a + b*x**2))**(3/2)/(385*b**2) + 2*a*x**4*(c*sqrt(a + b*x**2))**(3/2)/(55*b) + 2*x**6*(c*sqrt(a + b*x**2))**(3/2)/15, Ne(b, 0)), (x**6*(sqrt(a)*c)**(3/2)/6, True))`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.63

$$\int x^5 \left(c\sqrt{a+bx^2} \right)^{3/2} dx = \frac{2 \left(165 (\sqrt{bx^2+ac})^{7/2} a^2 c^4 - 210 (\sqrt{bx^2+ac})^{11/2} ac^2 + 77 (\sqrt{bx^2+ac})^{15/2} \right)}{1155 b^3 c^6}$$

input `integrate(x^5*(c*(b*x^2+a)^(1/2))^(3/2),x, algorithm="maxima")`output `2/1155*(165*(sqrt(b*x^2 + a)*c)^(7/2)*a^2*c^4 - 210*(sqrt(b*x^2 + a)*c)^(11/2)*a*c^2 + 77*(sqrt(b*x^2 + a)*c)^(15/2))/(b^3*c^6)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.07

$$\int x^5 \left(c\sqrt{a+bx^2} \right)^{3/2} dx = \frac{2 c^{3/2} \left(\frac{5 \left(21 (bx^2+a)^{11/4} - 66 (bx^2+a)^{7/4} a + 77 (bx^2+a)^{3/4} a^2 \right) a}{b^2} + \frac{77 (bx^2+a)^{15/4} - 315 (bx^2+a)^{11/4} a + 495 (bx^2+a)^{7/4} a^2}{b^2} \right)}{1155 b}$$

input `integrate(x^5*(c*(b*x^2+a)^(1/2))^(3/2),x, algorithm="giac")`output `2/1155*c^(3/2)*(5*(21*(b*x^2 + a)^(11/4) - 66*(b*x^2 + a)^(7/4)*a + 77*(b*x^2 + a)^(3/4)*a^2)*a/b^2 + (77*(b*x^2 + a)^(15/4) - 315*(b*x^2 + a)^(11/4)*a + 495*(b*x^2 + a)^(7/4)*a^2 - 385*(b*x^2 + a)^(3/4)*a^3)/b^2)/b`**Mupad [B] (verification not implemented)**

Time = 0.53 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.86

$$\int x^5 \left(c\sqrt{a+bx^2} \right)^{3/2} dx = \sqrt{c\sqrt{bx^2+a}} \left(\frac{2cx^6\sqrt{bx^2+a}}{15} + \frac{64a^3c\sqrt{bx^2+a}}{1155b^3} + \frac{2acx^4\sqrt{bx^2+a}}{55b} - \frac{16a^2cx^2\sqrt{bx^2+a}}{385b^2} \right)$$

input `int(x^5*(c*(a + b*x^2)^(1/2))^(3/2),x)`

output `(c*(a + b*x^2)^(1/2))^(1/2)*((2*c*x^6*(a + b*x^2)^(1/2))/15 + (64*a^3*c*(a + b*x^2)^(1/2))/(1155*b^3) + (2*a*c*x^4*(a + b*x^2)^(1/2))/(55*b) - (16*a^2*c*x^2*(a + b*x^2)^(1/2))/(385*b^2))`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.57

$$\int x^5 \left(c\sqrt{a + bx^2} \right)^{3/2} dx = \frac{2\sqrt{c} \sqrt{\sqrt{b} \sqrt{bx^2 + a} x + a + bx^2} \sqrt{\sqrt{bx^2 + a} + \sqrt{b} x} c \left(-32\sqrt{b} \sqrt{bx^2 + a} a^3 x - \dots \right)}{\dots}$$

input `int(x^5*(c*(b*x^2+a)^(1/2))^(3/2),x)`

output `(2*sqrt(c)*sqrt(sqrt(b)*sqrt(a + b*x**2)*x + a + b*x**2)*sqrt(sqrt(a + b*x**2) + sqrt(b)*x)*c*(- 32*sqrt(b)*sqrt(a + b*x**2)*a**3*x + 24*sqrt(b)*sqrt(a + b*x**2)*a**2*b*x**3 - 21*sqrt(b)*sqrt(a + b*x**2)*a*b**2*x**5 - 77*sqrt(b)*sqrt(a + b*x**2)*b**3*x**7 + 32*a**4 + 8*a**3*b*x**2 - 3*a**2*b**2*x**4 + 98*a*b**3*x**6 + 77*b**4*x**8))/(1155*a*b**3)`

3.41 $\int x^3 \left(c\sqrt{a + bx^2} \right)^{3/2} dx$

Optimal result	309
Mathematica [A] (verified)	309
Rubi [A] (verified)	310
Maple [A] (verified)	311
Fricas [A] (verification not implemented)	312
Sympy [A] (verification not implemented)	312
Maxima [A] (verification not implemented)	312
Giac [A] (verification not implemented)	313
Mupad [B] (verification not implemented)	313
Reduce [B] (verification not implemented)	314

Optimal result

Integrand size = 21, antiderivative size = 66

$$\int x^3 \left(c\sqrt{a + bx^2} \right)^{3/2} dx = -\frac{2a(c\sqrt{a + bx^2})^{3/2} (a + bx^2)}{7b^2} + \frac{2(c\sqrt{a + bx^2})^{3/2} (a + bx^2)^2}{11b^2}$$

output

$$-2/7*a*(c*(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)/b^2+2/11*(c*(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)^2/b^2$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.62

$$\int x^3 \left(c\sqrt{a + bx^2} \right)^{3/2} dx = \frac{2(c\sqrt{a + bx^2})^{3/2} (a + bx^2) (-4a + 7bx^2)}{77b^2}$$

input

```
Integrate[x^3*(c*Sqrt[a + b*x^2])^(3/2),x]
```

output

$$(2*(c*Sqrt[a + b*x^2])^(3/2)*(a + b*x^2)*(-4*a + 7*b*x^2))/(77*b^2)$$

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.27, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2045, 243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 (c\sqrt{a+bx^2})^{3/2} dx \\
 & \quad \downarrow \text{2045} \\
 & \frac{(c\sqrt{a+bx^2})^{3/2} \int x^3 \left(\frac{bx^2}{a} + 1\right)^{3/4} dx}{\left(\frac{bx^2}{a} + 1\right)^{3/4}} \\
 & \quad \downarrow \text{243} \\
 & \frac{(c\sqrt{a+bx^2})^{3/2} \int x^2 \left(\frac{bx^2}{a} + 1\right)^{3/4} dx^2}{2 \left(\frac{bx^2}{a} + 1\right)^{3/4}} \\
 & \quad \downarrow \text{53} \\
 & \frac{(c\sqrt{a+bx^2})^{3/2} \int \left(\frac{a \left(\frac{bx^2}{a} + 1\right)^{7/4}}{b} - \frac{a \left(\frac{bx^2}{a} + 1\right)^{3/4}}{b} \right) dx^2}{2 \left(\frac{bx^2}{a} + 1\right)^{3/4}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{\left(\frac{4a^2 \left(\frac{bx^2}{a} + 1\right)^{11/4}}{11b^2} - \frac{4a^2 \left(\frac{bx^2}{a} + 1\right)^{7/4}}{7b^2} \right) (c\sqrt{a+bx^2})^{3/2}}{2 \left(\frac{bx^2}{a} + 1\right)^{3/4}}
 \end{aligned}$$

input `Int[x^3*(c*Sqrt[a + b*x^2])^(3/2),x]`

output `((c*Sqrt[a + b*x^2])^(3/2)*((-4*a^2*(1 + (b*x^2)/a)^(7/4))/(7*b^2) + (4*a^2*(1 + (b*x^2)/a)^(11/4))/(11*b^2)))/(2*(1 + (b*x^2)/a)^(3/4))`

Defintions of rubi rules used

rule 53 $\text{Int}[(a_.) + (b_.)(x_)^{(m_.)}((c_.) + (d_.)(x_)^{(n_.)}), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \text{FreeQ}\{a, b, c, d, n\}, x] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ (\ !\text{IntegerQ}[n] \ || \ (\text{EqQ}[c, 0] \ \&\& \ \text{LeQ}[7*m + 4*n + 4, 0]) \ || \ \text{LtQ}[9*m + 5*(n + 1), 0] \ || \ \text{GtQ}[m + n + 2, 0])$

rule 243 $\text{Int}[(x_)^{(m_.)}((a_) + (b_.)(x_)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[1/2 \ \text{Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] /; \text{FreeQ}\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 2045 $\text{Int}[(u_.)*((c_.)*((a_) + (b_.)(x_)^{(n_.)})^{(q_.)})^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[\text{Simp}[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^{(p*q)}] \ \text{Int}[u*(1 + b*(x^n/a))^{(p*q)}, x], x] /; \text{FreeQ}\{a, b, c, n, p, q\}, x] \ \&\& \ !\text{GeQ}[a, 0]$

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.55

method	result	size
gospers	$-\frac{2(bx^2+a)(-7bx^2+4a)(c\sqrt{bx^2+a})^{\frac{3}{2}}}{77b^2}$	36
orering	$-\frac{2(bx^2+a)(-7bx^2+4a)(c\sqrt{bx^2+a})^{\frac{3}{2}}}{77b^2}$	36

input $\text{int}(x^3*(c*(b*x^2+a)^{(1/2}))^{(3/2)}, x, \text{method}=_RETURNVERBOSE)$

output $-2/77*(b*x^2+a)*(-7*b*x^2+4*a)*(c*(b*x^2+a)^{(1/2}))^{(3/2)}/b^2$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.77

$$\int x^3 \left(c\sqrt{a+bx^2} \right)^{3/2} dx = \frac{2(7b^2cx^4 + 3abcx^2 - 4a^2c)\sqrt{bx^2+a}\sqrt{\sqrt{bx^2+ac}}}{77b^2}$$

input `integrate(x^3*(c*(b*x^2+a)^(1/2))^(3/2),x, algorithm="fricas")`output `2/77*(7*b^2*c*x^4 + 3*a*b*c*x^2 - 4*a^2*c)*sqrt(b*x^2 + a)*sqrt(sqrt(b*x^2 + a)*c)/b^2`**Sympy [A] (verification not implemented)**

Time = 2.90 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.32

$$\int x^3 \left(c\sqrt{a+bx^2} \right)^{3/2} dx = \begin{cases} -\frac{8a^2(c\sqrt{a+bx^2})^{3/2}}{77b^2} + \frac{6ax^2(c\sqrt{a+bx^2})^{3/2}}{77b} + \frac{2x^4(c\sqrt{a+bx^2})^{3/2}}{11} & \text{for } b \neq 0 \\ \frac{x^4(\sqrt{ac})^{3/2}}{4} & \text{otherwise} \end{cases}$$

input `integrate(x**3*(c*(b*x**2+a)**(1/2))**(3/2),x)`output `Piecewise((-8*a**2*(c*sqrt(a + b*x**2))**(3/2)/(77*b**2) + 6*a*x**2*(c*sqrt(a + b*x**2))**(3/2)/(77*b) + 2*x**4*(c*sqrt(a + b*x**2))**(3/2)/11, Ne(b, 0)), (x**4*(sqrt(a)*c)**(3/2)/4, True))`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.65

$$\int x^3 \left(c\sqrt{a+bx^2} \right)^{3/2} dx = -\frac{2 \left(11 (\sqrt{bx^2+ac})^{7/2} ac^2 - 7 (\sqrt{bx^2+ac})^{11/2} \right)}{77b^2c^4}$$

input `integrate(x^3*(c*(b*x^2+a)^(1/2))^(3/2),x, algorithm="maxima")`

output

$$\frac{-2/77*(11*(\sqrt{b*x^2 + a})*c)^{(7/2)}*a*c^2 - 7*(\sqrt{b*x^2 + a})*c^{(11/2)}}{(b^2*c^4)}$$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.23

$$\int x^3 \left(c\sqrt{a + bx^2} \right)^{3/2} dx = \frac{2 \left(\frac{11 \left(3 (bx^2+a)^{7/4} - 7 (bx^2+a)^{3/4} a \right) a}{b} + \frac{21 (bx^2+a)^{11/4} - 66 (bx^2+a)^{7/4} a + 77 (bx^2+a)^{3/4} a^2}{b} \right) c^{3/2}}{231 b}$$

input

```
integrate(x^3*(c*(b*x^2+a)^(1/2))^(3/2),x, algorithm="giac")
```

output

$$\frac{2/231*(11*(3*(b*x^2 + a)^{(7/4)} - 7*(b*x^2 + a)^{(3/4)}*a)*a/b + (21*(b*x^2 + a)^{(11/4)} - 66*(b*x^2 + a)^{(7/4)}*a + 77*(b*x^2 + a)^{(3/4)}*a^2)/b)*c^{(3/2)}}{b}$$

Mupad [B] (verification not implemented)

Time = 0.51 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.02

$$\int x^3 \left(c\sqrt{a + bx^2} \right)^{3/2} dx = \sqrt{c\sqrt{bx^2 + a}} \left(\frac{2cx^4\sqrt{bx^2 + a}}{11} - \frac{8a^2c\sqrt{bx^2 + a}}{77b^2} + \frac{6acx^2\sqrt{bx^2 + a}}{77b} \right)$$

input

```
int(x^3*(c*(a + b*x^2)^(1/2))^(3/2),x)
```

output

$$(c*(a + b*x^2)^(1/2))^(1/2)*((2*c*x^4*(a + b*x^2)^(1/2))/11 - (8*a^2*c*(a + b*x^2)^(1/2))/(77*b^2) + (6*a*c*x^2*(a + b*x^2)^(1/2))/(77*b))$$

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.94

$$\int x^3 \left(c\sqrt{a+bx^2} \right)^{3/2} dx = \frac{2\sqrt{c} \sqrt{\sqrt{b}\sqrt{bx^2+a}x+a+bx^2} \sqrt{\sqrt{bx^2+a}+\sqrt{b}x} c \left(4\sqrt{b}\sqrt{bx^2+a}a^2x - 3\sqrt{b}\sqrt{bx^2+a}a^2x - 3\sqrt{b}\sqrt{bx^2+a}a^2x - 3\sqrt{b}\sqrt{bx^2+a}a^2x \right)}{77a}$$

input `int(x^3*(c*(b*x^2+a)^(1/2))^(3/2),x)`output `(2*sqrt(c)*sqrt(sqrt(b)*sqrt(a + b*x**2)*x + a + b*x**2)*sqrt(sqrt(a + b*x**2) + sqrt(b)*x)*c*(4*sqrt(b)*sqrt(a + b*x**2)*a**2*x - 3*sqrt(b)*sqrt(a + b*x**2)*a*b*x**3 - 7*sqrt(b)*sqrt(a + b*x**2)*b**2*x**5 - 4*a**3 - a**2*b*x**2 + 10*a*b**2*x**4 + 7*b**3*x**6))/(77*a*b**2)`

$$3.42 \quad \int x \left(c\sqrt{a + bx^2} \right)^{3/2} dx$$

Optimal result	315
Mathematica [A] (verified)	315
Rubi [A] (verified)	316
Maple [A] (verified)	317
Fricas [A] (verification not implemented)	317
Sympy [A] (verification not implemented)	318
Maxima [A] (verification not implemented)	318
Giac [A] (verification not implemented)	318
Mupad [B] (verification not implemented)	319
Reduce [B] (verification not implemented)	319

Optimal result

Integrand size = 19, antiderivative size = 36

$$\int x \left(c\sqrt{a + bx^2} \right)^{3/2} dx = \frac{2c\sqrt{c\sqrt{a + bx^2}}(a + bx^2)^{3/2}}{7b}$$

output `2/7*c*(c*(b*x^2+a)^(1/2))^(1/2)*(b*x^2+a)^(3/2)/b`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.86

$$\int x \left(c\sqrt{a + bx^2} \right)^{3/2} dx = \frac{2(c\sqrt{a + bx^2})^{3/2} (a + bx^2)}{7b}$$

input `Integrate[x*(c*Sqrt[a + b*x^2])^(3/2),x]`

output `(2*(c*Sqrt[a + b*x^2])^(3/2)*(a + b*x^2))/(7*b)`

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.86, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2024, 20, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int x \left(c\sqrt{a + bx^2} \right)^{3/2} dx \\
 \downarrow \text{2024} \\
 \frac{\int \left(c\sqrt{bx^2 + a} \right)^{3/2} d(bx^2 + a)}{2b} \\
 \downarrow \text{20} \\
 \frac{\left(c\sqrt{a + bx^2} \right)^{3/2} \int (bx^2 + a)^{3/4} d(bx^2 + a)}{2b(a + bx^2)^{3/4}} \\
 \downarrow \text{15} \\
 \frac{2(a + bx^2) \left(c\sqrt{a + bx^2} \right)^{3/2}}{7b}
 \end{array}$$

input `Int[x*(c*Sqrt[a + b*x^2])^(3/2),x]`

output `(2*(c*Sqrt[a + b*x^2])^(3/2)*(a + b*x^2))/(7*b)`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 20 `Int[((a_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a*x^n)^p/x^(n*p) Int[x^(n*p), x], x] /; FreeQ[{a, n, p}, x] && !IntegerQ[p]`

rule 2024

```
Int[((a_.) + (b_.)*(Pq_)^(n_.))^(p_.)*(Qr_), x_Symbol] := With[{q = Expon[P
q, x], r = Expon[Qr, x]}, Simp[Coeff[Qr, x, r]/(q*Coeff[Pq, x, q]) Subst[
Int[(a + b*x^n)^p, x], x, Pq], x] /; EqQ[r, q - 1] && EqQ[Coeff[Qr, x, r]*D
[Pq, x], q*Coeff[Pq, x, q]*Qr] /; FreeQ[{a, b, n, p}, x] && PolyQ[Pq, x] &
& PolyQ[Qr, x]
```

Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.72

method	result	size
gospers	$\frac{2(bx^2+a)(c\sqrt{bx^2+a})^{\frac{3}{2}}}{7b}$	26
derivativedivides	$\frac{2(bx^2+a)(c\sqrt{bx^2+a})^{\frac{3}{2}}}{7b}$	26
default	$\frac{2(bx^2+a)(c\sqrt{bx^2+a})^{\frac{3}{2}}}{7b}$	26
orering	$\frac{2(bx^2+a)(c\sqrt{bx^2+a})^{\frac{3}{2}}}{7b}$	26

input

```
int(x*(c*(b*x^2+a)^(1/2))^(3/2),x,method=_RETURNVERBOSE)
```

output

```
2/7*(b*x^2+a)*(c*(b*x^2+a)^(1/2))^(3/2)/b
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.03

$$\int x \left(c\sqrt{a + bx^2} \right)^{3/2} dx = \frac{2(bc x^2 + ac)\sqrt{bx^2 + a}\sqrt{\sqrt{bx^2 + ac}}}{7b}$$

input

```
integrate(x*(c*(b*x^2+a)^(1/2))^(3/2),x, algorithm="fricas")
```

output

```
2/7*(b*c*x^2 + a*c)*sqrt(b*x^2 + a)*sqrt(sqrt(b*x^2 + a)*c)/b
```

Sympy [A] (verification not implemented)

Time = 1.40 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.61

$$\int x \left(c\sqrt{a+bx^2} \right)^{3/2} dx = \begin{cases} \frac{2a(c\sqrt{a+bx^2})^{3/2}}{7b} + \frac{2x^2(c\sqrt{a+bx^2})^{3/2}}{7} & \text{for } b \neq 0 \\ \frac{x^2(\sqrt{ac})^{3/2}}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*(c*(b*x**2+a)**(1/2))**(3/2),x)`output `Piecewise((2*a*(c*sqrt(a + b*x**2))**(3/2)/(7*b) + 2*x**2*(c*sqrt(a + b*x**2))**(3/2)/7, Ne(b, 0)), (x**2*(sqrt(a)*c)**(3/2)/2, True))`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.69

$$\int x \left(c\sqrt{a+bx^2} \right)^{3/2} dx = \frac{2(bx^2 + a)(\sqrt{bx^2 + ac})^{3/2}}{7b}$$

input `integrate(x*(c*(b*x^2+a)^(1/2))^(3/2),x, algorithm="maxima")`output `2/7*(b*x^2 + a)*(sqrt(b*x^2 + a)*c)^(3/2)/b`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.47

$$\int x \left(c\sqrt{a+bx^2} \right)^{3/2} dx = \frac{2(bx^2 + a)^{7/4} c^{3/2}}{7b}$$

input `integrate(x*(c*(b*x^2+a)^(1/2))^(3/2),x, algorithm="giac")`output `2/7*(b*x^2 + a)^(7/4)*c^(3/2)/b`

Mupad [B] (verification not implemented)

Time = 0.49 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.78

$$\int x \left(c\sqrt{a + bx^2} \right)^{3/2} dx = \frac{2c(bx^2 + a)^{3/2} \sqrt{c\sqrt{bx^2 + a}}}{7b}$$

input `int(x*(c*(a + b*x^2)^(1/2))^(3/2),x)`output `(2*c*(a + b*x^2)^(3/2)*(c*(a + b*x^2)^(1/2))^(1/2))/(7*b)`**Reduce [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 93, normalized size of antiderivative = 2.58

$$\int x \left(c\sqrt{a + bx^2} \right)^{3/2} dx = \frac{2\sqrt{c} \sqrt{\sqrt{b}\sqrt{bx^2 + a}x + a + bx^2} \sqrt{\sqrt{bx^2 + a} + \sqrt{b}x} c \left(-\sqrt{b}\sqrt{bx^2 + a}ax - \sqrt{b}\sqrt{bx^2 + a} \right)}{7ab}$$

input `int(x*(c*(b*x^2+a)^(1/2))^(3/2),x)`output `(2*sqrt(c)*sqrt(sqrt(b)*sqrt(a + b*x**2)*x + a + b*x**2)*sqrt(sqrt(a + b*x**2) + sqrt(b)*x)*c*(- sqrt(b)*sqrt(a + b*x**2)*a*x - sqrt(b)*sqrt(a + b*x**2)*b*x**3 + a**2 + 2*a*b*x**2 + b**2*x**4))/(7*a*b)`

3.43 $\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x} dx$

Optimal result	320
Mathematica [A] (verified)	321
Rubi [A] (verified)	321
Maple [F]	324
Fricas [F(-1)]	325
Sympy [F]	325
Maxima [A] (verification not implemented)	325
Giac [A] (verification not implemented)	326
Mupad [F(-1)]	326
Reduce [B] (verification not implemented)	327

Optimal result

Integrand size = 21, antiderivative size = 117

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x} dx = \frac{2}{3}(c\sqrt{a+bx^2})^{3/2} + \frac{(c\sqrt{a+bx^2})^{3/2} \arctan\left(\sqrt[4]{1+\frac{bx^2}{a}}\right)}{\left(1+\frac{bx^2}{a}\right)^{3/4}} - \frac{(c\sqrt{a+bx^2})^{3/2} \operatorname{arctanh}\left(\sqrt[4]{1+\frac{bx^2}{a}}\right)}{\left(1+\frac{bx^2}{a}\right)^{3/4}}$$

output

```
2/3*(c*(b*x^2+a)^(1/2))^(3/2)+(c*(b*x^2+a)^(1/2))^(3/2)*arctan((1+b*x^2/a)^(1/4))/(1+b*x^2/a)^(3/4)-(c*(b*x^2+a)^(1/2))^(3/2)*arctanh((1+b*x^2/a)^(1/4))/(1+b*x^2/a)^(3/4)
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.82

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x} dx = \frac{(c\sqrt{a+bx^2})^{3/2} \left(2(a+bx^2)^{3/4} + 3a^{3/4} \arctan\left(\frac{\sqrt[4]{a+bx^2}}{\sqrt[4]{a}}\right) - 3a^{3/4} \operatorname{arctanh}\left(\frac{\sqrt[4]{a+bx^2}}{\sqrt[4]{a}}\right) \right)}{3(a+bx^2)^{3/4}}$$

input `Integrate[(c*Sqrt[a + b*x^2])^(3/2)/x,x]`

output `((c*Sqrt[a + b*x^2])^(3/2)*(2*(a + b*x^2)^(3/4) + 3*a^(3/4)*ArcTan[(a + b*x^2)^(1/4)/a^(1/4)] - 3*a^(3/4)*ArcTanh[(a + b*x^2)^(1/4)/a^(1/4)]))/(3*(a + b*x^2)^(3/4))`

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.81, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {2045, 243, 60, 73, 25, 27, 827, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c\sqrt{a+bx^2})^{3/2}}{x} dx \\ & \quad \downarrow \text{2045} \\ & \frac{(c\sqrt{a+bx^2})^{3/2} \int \frac{(\frac{bx^2}{a}+1)^{3/4}}{x} dx}{(\frac{bx^2}{a}+1)^{3/4}} \\ & \quad \downarrow \text{243} \\ & \frac{(c\sqrt{a+bx^2})^{3/2} \int \frac{(\frac{bx^2}{a}+1)^{3/4}}{x^2} dx^2}{2(\frac{bx^2}{a}+1)^{3/4}} \end{aligned}$$

$$\begin{aligned} & \downarrow 60 \\ & \frac{(c\sqrt{a+bx^2})^{3/2} \left(\int \frac{1}{x^2 \sqrt[4]{\frac{bx^2}{a} + 1}} dx^2 + \frac{4}{3} \left(\frac{bx^2}{a} + 1 \right)^{3/4} \right)}{2 \left(\frac{bx^2}{a} + 1 \right)^{3/4}} \\ & \downarrow 73 \\ & \frac{(c\sqrt{a+bx^2})^{3/2} \left(\frac{4a \int -\frac{bx^4}{a(1-x^8)} d \sqrt[4]{\frac{bx^2}{a} + 1}}{b} + \frac{4}{3} \left(\frac{bx^2}{a} + 1 \right)^{3/4} \right)}{2 \left(\frac{bx^2}{a} + 1 \right)^{3/4}} \\ & \downarrow 25 \\ & \frac{(c\sqrt{a+bx^2})^{3/2} \left(\frac{4}{3} \left(\frac{bx^2}{a} + 1 \right)^{3/4} - \frac{4a \int \frac{bx^4}{a(1-x^8)} d \sqrt[4]{\frac{bx^2}{a} + 1}}{b} \right)}{2 \left(\frac{bx^2}{a} + 1 \right)^{3/4}} \\ & \downarrow 27 \\ & \frac{(c\sqrt{a+bx^2})^{3/2} \left(\frac{4}{3} \left(\frac{bx^2}{a} + 1 \right)^{3/4} - 4 \int \frac{x^4}{1-x^8} d \sqrt[4]{\frac{bx^2}{a} + 1} \right)}{2 \left(\frac{bx^2}{a} + 1 \right)^{3/4}} \\ & \downarrow 827 \\ & \frac{(c\sqrt{a+bx^2})^{3/2} \left(\frac{4}{3} \left(\frac{bx^2}{a} + 1 \right)^{3/4} - 4 \left(\frac{1}{2} \int \frac{1}{1-x^4} d \sqrt[4]{\frac{bx^2}{a} + 1} - \frac{1}{2} \int \frac{1}{x^4+1} d \sqrt[4]{\frac{bx^2}{a} + 1} \right) \right)}{2 \left(\frac{bx^2}{a} + 1 \right)^{3/4}} \\ & \downarrow 216 \\ & \frac{(c\sqrt{a+bx^2})^{3/2} \left(\frac{4}{3} \left(\frac{bx^2}{a} + 1 \right)^{3/4} - 4 \left(\frac{1}{2} \int \frac{1}{1-x^4} d \sqrt[4]{\frac{bx^2}{a} + 1} - \frac{1}{2} \arctan \left(\sqrt[4]{\frac{bx^2}{a} + 1} \right) \right) \right)}{2 \left(\frac{bx^2}{a} + 1 \right)^{3/4}} \end{aligned}$$

$$\frac{(c\sqrt{a+bx^2})^{3/2} \left(\frac{4}{3} \left(\frac{bx^2}{a} + 1 \right)^{3/4} - 4 \left(\frac{1}{2} \operatorname{arctanh} \left(\sqrt[4]{\frac{bx^2}{a} + 1} \right) - \frac{1}{2} \arctan \left(\sqrt[4]{\frac{bx^2}{a} + 1} \right) \right) \right)}{2 \left(\frac{bx^2}{a} + 1 \right)^{3/4}}$$

input `Int[(c*Sqrt[a + b*x^2])^(3/2)/x,x]`

output `((c*Sqrt[a + b*x^2])^(3/2)*((4*(1 + (b*x^2)/a)^(3/4))/3 - 4*(-1/2*ArcTan[(1 + (b*x^2)/a)^(1/4)] + ArcTanh[(1 + (b*x^2)/a)^(1/4)]/2)))/(2*(1 + (b*x^2)/a)^(3/4))`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 827 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[s/(2*b) Int[1/(r + s*x^2), x], x] - Simp[s/(2*b) Int[1/(r - s*x^2), x], x] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q) Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [F]

$$\int \frac{(c\sqrt{bx^2+a})^{\frac{3}{2}}}{x} dx$$

input `int((c*(b*x^2+a)^(1/2))^(3/2)/x,x)`

output `int((c*(b*x^2+a)^(1/2))^(3/2)/x,x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x} dx = \text{Timed out}$$

input `integrate((c*(b*x^2+a)^(1/2))^(3/2)/x,x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x} dx = \int \frac{(c\sqrt{a+bx^2})^{\frac{3}{2}}}{x} dx$$

input `integrate((c*(b*x**2+a)**(1/2))**(3/2)/x,x)`

output `Integral((c*sqrt(a + b*x**2))**(3/2)/x, x)`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.01

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x} dx = \frac{3ac^4 \left(\frac{2 \arctan\left(\frac{\sqrt{\sqrt{bx^2+ac}}}{(ac^2)^{\frac{1}{4}}}\right)}{(ac^2)^{\frac{1}{4}}} + \frac{\log\left(\frac{\sqrt{\sqrt{bx^2+ac}} - (ac^2)^{\frac{1}{4}}}{\sqrt{\sqrt{bx^2+ac}} + (ac^2)^{\frac{1}{4}}}\right)}{(ac^2)^{\frac{1}{4}}}\right) + 4(\sqrt{bx^2+ac})^{\frac{3}{2}}c^2}{6c^2}$$

input `integrate((c*(b*x^2+a)^(1/2))^(3/2)/x,x, algorithm="maxima")`

output

```
1/6*(3*a*c^4*(2*arctan(sqrt(sqrt(b*x^2 + a)*c)/(a*c^2)^(1/4))/(a*c^2)^(1/4)
) + log((sqrt(sqrt(b*x^2 + a)*c) - (a*c^2)^(1/4))/(sqrt(sqrt(b*x^2 + a)*c)
+ (a*c^2)^(1/4)))/(a*c^2)^(1/4)) + 4*(sqrt(b*x^2 + a)*c)^(3/2)*c^2/c^2
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.62

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x} dx = -\frac{1}{12} \left(6\sqrt{2}(-a)^{\frac{3}{4}} \arctan \left(\frac{\sqrt{2}(\sqrt{2}(-a)^{\frac{1}{4}} + 2(bx^2 + a)^{\frac{1}{4}})}{2(-a)^{\frac{1}{4}}} \right) + 6\sqrt{2}(-a)^{\frac{3}{4}} \arctan \left(-\frac{\sqrt{2}(\sqrt{2}(-a)^{\frac{1}{4}} - 2(bx^2 + a)^{\frac{1}{4}})}{2(-a)^{\frac{1}{4}}} \right) \right)$$

input

```
integrate((c*(b*x^2+a)^(1/2))^(3/2)/x,x, algorithm="giac")
```

output

```
-1/12*(6*sqrt(2)*(-a)^(3/4)*arctan(1/2*sqrt(2)*(sqrt(2)*(-a)^(1/4) + 2*(b*
x^2 + a)^(1/4))/(-a)^(1/4)) + 6*sqrt(2)*(-a)^(3/4)*arctan(-1/2*sqrt(2)*(sq
rt(2)*(-a)^(1/4) - 2*(b*x^2 + a)^(1/4))/(-a)^(1/4)) - 3*sqrt(2)*(-a)^(3/4)
*log(sqrt(2)*(b*x^2 + a)^(1/4)*(-a)^(1/4) + sqrt(b*x^2 + a) + sqrt(-a)) +
3*sqrt(2)*(-a)^(3/4)*log(-sqrt(2)*(b*x^2 + a)^(1/4)*(-a)^(1/4) + sqrt(b*x^
2 + a) + sqrt(-a)) - 8*(b*x^2 + a)^(3/4)*c^(3/2)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x} dx = \int \frac{(c\sqrt{bx^2+a})^{3/2}}{x} dx$$

input

```
int((c*(a + b*x^2)^(1/2))^(3/2)/x,x)
```

output

```
int((c*(a + b*x^2)^(1/2))^(3/2)/x, x)
```

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.07

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x} dx = \frac{\sqrt{c}c \left(-3a^{3/4} \operatorname{atan} \left(\frac{a^{5/4}(bx^2+a)^{3/4} - a^{7/4}(bx^2+a)^{1/4} - a^{3/4}(bx^2+a)^{1/4}bx^2}{2abx^2+2a^2} \right) + 4(bx^2+a)^{3/4} - 3a^{3/4} \log \right)}{6}$$

input `int((c*(b*x^2+a)^(1/2))^(3/2)/x,x)`output `(sqrt(c)*c*(- 3*a**(3/4)*atan((a**(1/4)*(a + b*x**2)**(3/4)*a - a**(3/4)*(a + b*x**2)**(1/4)*a - a**(3/4)*(a + b*x**2)**(1/4)*b*x**2)/(2*a**2 + 2*a*b*x**2)) + 4*(a + b*x**2)**(3/4) - 3*a**(3/4)*log((a + b*x**2)**(1/4) + a**(1/4)) + 3*a**(3/4)*log((a + b*x**2)**(1/4) - a**(1/4))))/6`

3.44 $\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^3} dx$

Optimal result	328
Mathematica [A] (verified)	329
Rubi [A] (verified)	329
Maple [F]	333
Fricas [F(-1)]	333
Sympy [F]	333
Maxima [A] (verification not implemented)	334
Giac [A] (verification not implemented)	334
Mupad [F(-1)]	335
Reduce [B] (verification not implemented)	335

Optimal result

Integrand size = 21, antiderivative size = 133

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^3} dx = -\frac{(c\sqrt{a+bx^2})^{3/2}}{2x^2} + \frac{3b(c\sqrt{a+bx^2})^{3/2} \arctan\left(\sqrt[4]{1+\frac{bx^2}{a}}\right)}{4a\left(1+\frac{bx^2}{a}\right)^{3/4}} - \frac{3b(c\sqrt{a+bx^2})^{3/2} \operatorname{arctanh}\left(\sqrt[4]{1+\frac{bx^2}{a}}\right)}{4a\left(1+\frac{bx^2}{a}\right)^{3/4}}$$

output `-1/2*(c*(b*x^2+a)^(1/2))^(3/2)/x^2+3/4*b*(c*(b*x^2+a)^(1/2))^(3/2)*arctan((1+b*x^2/a)^(1/4))/a/(1+b*x^2/a)^(3/4)-3/4*b*(c*(b*x^2+a)^(1/2))^(3/2)*arctanh((1+b*x^2/a)^(1/4))/a/(1+b*x^2/a)^(3/4)`

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.80

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^3} dx = \frac{(c\sqrt{a+bx^2})^{3/2} \left(2\sqrt[4]{a}(a+bx^2)^{3/4} - 3bx^2 \arctan\left(\frac{\sqrt[4]{a+bx^2}}{\sqrt[4]{a}}\right) + 3bx^2 \operatorname{arctanh}\left(\frac{\sqrt[4]{a+bx^2}}{\sqrt[4]{a}}\right) \right)}{4\sqrt[4]{a}x^2(a+bx^2)^{3/4}}$$

input `Integrate[(c*Sqrt[a + b*x^2])^(3/2)/x^3,x]`

output `-1/4*((c*Sqrt[a + b*x^2])^(3/2)*(2*a^(1/4)*(a + b*x^2)^(3/4) - 3*b*x^2*ArcTan[(a + b*x^2)^(1/4)/a^(1/4)] + 3*b*x^2*ArcTanh[(a + b*x^2)^(1/4)/a^(1/4)]))/(a^(1/4)*x^2*(a + b*x^2)^(3/4))`

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.75, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {2045, 243, 51, 73, 25, 27, 827, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^3} dx$$

↓ 2045

$$\frac{(c\sqrt{a+bx^2})^{3/2} \int \frac{(\frac{bx^2}{a}+1)^{3/4}}{x^3} dx}{(\frac{bx^2}{a}+1)^{3/4}}$$

↓ 243

$$\begin{aligned}
& \frac{(c\sqrt{a+bx^2})^{3/2} \int \frac{(\frac{bx^2}{a}+1)^{3/4}}{x^4} dx^2}{2\left(\frac{bx^2}{a}+1\right)^{3/4}} \\
& \quad \downarrow \mathbf{51} \\
& \frac{(c\sqrt{a+bx^2})^{3/2} \left(\frac{3b \int \frac{1}{x^2 \sqrt[4]{\frac{bx^2}{a}+1}} dx^2}{4a} - \frac{(\frac{bx^2}{a}+1)^{3/4}}{x^2} \right)}{2\left(\frac{bx^2}{a}+1\right)^{3/4}} \\
& \quad \downarrow \mathbf{73} \\
& \frac{(c\sqrt{a+bx^2})^{3/2} \left(3 \int -\frac{bx^4}{a(1-x^8)} d\sqrt[4]{\frac{bx^2}{a}+1} - \frac{(\frac{bx^2}{a}+1)^{3/4}}{x^2} \right)}{2\left(\frac{bx^2}{a}+1\right)^{3/4}} \\
& \quad \downarrow \mathbf{25} \\
& \frac{(c\sqrt{a+bx^2})^{3/2} \left(-3 \int \frac{bx^4}{a(1-x^8)} d\sqrt[4]{\frac{bx^2}{a}+1} - \frac{(\frac{bx^2}{a}+1)^{3/4}}{x^2} \right)}{2\left(\frac{bx^2}{a}+1\right)^{3/4}} \\
& \quad \downarrow \mathbf{27} \\
& \frac{(c\sqrt{a+bx^2})^{3/2} \left(-\frac{3b \int \frac{x^4}{1-x^8} d\sqrt[4]{\frac{bx^2}{a}+1}}{a} - \frac{(\frac{bx^2}{a}+1)^{3/4}}{x^2} \right)}{2\left(\frac{bx^2}{a}+1\right)^{3/4}} \\
& \quad \downarrow \mathbf{827} \\
& \frac{(c\sqrt{a+bx^2})^{3/2} \left(-\frac{3b \left(\frac{1}{2} \int \frac{1}{1-x^4} d\sqrt[4]{\frac{bx^2}{a}+1} + 1 - \frac{1}{2} \int \frac{1}{x^4+1} d\sqrt[4]{\frac{bx^2}{a}+1} \right)}{a} - \frac{(\frac{bx^2}{a}+1)^{3/4}}{x^2} \right)}{2\left(\frac{bx^2}{a}+1\right)^{3/4}} \\
& \quad \downarrow \mathbf{216}
\end{aligned}$$

$$\frac{(c\sqrt{a+bx^2})^{3/2} \left(-\frac{3b \left(\frac{1}{2} \int \frac{1}{1-x^4} d\sqrt[4]{\frac{bx^2}{a} + 1} - \frac{1}{2} \arctan \left(\sqrt[4]{\frac{bx^2}{a} + 1} \right) \right)}{a} - \frac{\left(\frac{bx^2}{a} + 1\right)^{3/4}}{x^2} \right)}{2 \left(\frac{bx^2}{a} + 1\right)^{3/4}}$$

↓ 219

$$\frac{(c\sqrt{a+bx^2})^{3/2} \left(-\frac{3b \left(\frac{1}{2} \operatorname{arctanh} \left(\sqrt[4]{\frac{bx^2}{a} + 1} \right) - \frac{1}{2} \arctan \left(\sqrt[4]{\frac{bx^2}{a} + 1} \right) \right)}{a} - \frac{\left(\frac{bx^2}{a} + 1\right)^{3/4}}{x^2} \right)}{2 \left(\frac{bx^2}{a} + 1\right)^{3/4}}$$

input `Int[(c*Sqrt[a + b*x^2])^(3/2)/x^3,x]`

output `((c*Sqrt[a + b*x^2])^(3/2)*(-(1 + (b*x^2)/a)^(3/4)/x^2) - (3*b*(-1/2*ArcTan[(1 + (b*x^2)/a)^(1/4)] + ArcTanh[(1 + (b*x^2)/a)^(1/4)]/2))/a)/(2*(1 + (b*x^2)/a)^(3/4))`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 51 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))] Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]`

- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`
- rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
 rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
 , 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
 ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
 Q[a, 0] || LtQ[b, 0])`
- rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int
 t[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && I
 ntegerQ[(m - 1)/2]`
- rule 827 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b,
 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[s/(2*b) Int[1/(r + s*x^2), x],
 x] - Simp[s/(2*b) Int[1/(r - s*x^2), x], x] /; FreeQ[{a, b}, x] && !GtQ
 [a/b, 0]`
- rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_))^(p_), x_Symbol] := Simp[Si
 mp[(c*(a + b*x^n)^q]^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q)
 , x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [F]

$$\int \frac{(c\sqrt{bx^2+a})^{\frac{3}{2}}}{x^3} dx$$

input `int((c*(b*x^2+a)^(1/2))^(3/2)/x^3,x)`

output `int((c*(b*x^2+a)^(1/2))^(3/2)/x^3,x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^3} dx = \text{Timed out}$$

input `integrate((c*(b*x^2+a)^(1/2))^(3/2)/x^3,x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^3} dx = \int \frac{(c\sqrt{a+bx^2})^{\frac{3}{2}}}{x^3} dx$$

input `integrate((c*(b*x**2+a)**(1/2))**(3/2)/x**3,x)`

output `Integral((c*sqrt(a + b*x**2))**(3/2)/x**3, x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.04

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^3} dx = \frac{\left(3c^4 \left(\frac{2 \arctan\left(\frac{\sqrt{\sqrt{bx^2+ac}}}{(ac^2)^{1/4}}\right)}{(ac^2)^{1/4}} + \frac{\log\left(\frac{\sqrt{\sqrt{bx^2+ac}} - (ac^2)^{1/4}}{\sqrt{\sqrt{bx^2+ac}} + (ac^2)^{1/4}}\right)}{(ac^2)^{1/4}} \right) - \frac{4(\sqrt{bx^2+ac})^{3/2}c^4}{(bx^2+a)c^2-ac^2} \right) b}{8c^2}$$

input `integrate((c*(b*x^2+a)^(1/2))^(3/2)/x^3,x, algorithm="maxima")`

output `1/8*(3*c^4*(2*arctan(sqrt(sqrt(b*x^2 + a)*c)/(a*c^2)^(1/4))/(a*c^2)^(1/4) + log((sqrt(sqrt(b*x^2 + a)*c) - (a*c^2)^(1/4))/(sqrt(sqrt(b*x^2 + a)*c) + (a*c^2)^(1/4)))/(a*c^2)^(1/4)) - 4*(sqrt(b*x^2 + a)*c)^(3/2)*c^4/((b*x^2 + a)*c^2 - a*c^2))*b/c^2`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.57

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^3} dx = -\frac{1}{16} \left(\frac{6\sqrt{2}(-a)^{3/4} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}(-a)^{1/4} + 2(bx^2+a)^{1/4}\right)}{2(-a)^{1/4}}\right)}{a} + \frac{6\sqrt{2}(-a)^{3/4} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}(-a)^{1/4} - 2(bx^2+a)^{1/4}\right)}{2(-a)^{1/4}}\right)}{a} \right) + \dots$$

input `integrate((c*(b*x^2+a)^(1/2))^(3/2)/x^3,x, algorithm="giac")`

output `-1/16*(6*sqrt(2)*(-a)^(3/4)*arctan(1/2*sqrt(2)*(sqrt(2)*(-a)^(1/4) + 2*(b*x^2 + a)^(1/4))/(-a)^(1/4))/a + 6*sqrt(2)*(-a)^(3/4)*arctan(-1/2*sqrt(2)*(sqrt(2)*(-a)^(1/4) - 2*(b*x^2 + a)^(1/4))/(-a)^(1/4))/a - 3*sqrt(2)*(-a)^(3/4)*log(sqrt(2)*(b*x^2 + a)^(1/4)*(-a)^(1/4) + sqrt(b*x^2 + a) + sqrt(-a))/a + 3*sqrt(2)*(-a)^(3/4)*log(-sqrt(2)*(b*x^2 + a)^(1/4)*(-a)^(1/4) + sqrt(b*x^2 + a) + sqrt(-a))/a + 8*(b*x^2 + a)^(3/4)/(b*x^2))*b*c^(3/2)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^3} dx = \int \frac{(c\sqrt{bx^2+a})^{3/2}}{x^3} dx$$

input `int((c*(a + b*x^2)^(1/2))^(3/2)/x^3,x)`output `int((c*(a + b*x^2)^(1/2))^(3/2)/x^3, x)`**Reduce [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.08

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^3} dx = \frac{\sqrt{c}c \left(-3a^{3/4} \operatorname{atan} \left(\frac{a^{5/4}(bx^2+a)^{3/4} - a^{7/4}(bx^2+a)^{1/4} - a^{3/4}(bx^2+a)^{1/4}bx^2}{2abx^2+2a^2} \right) bx^2 - 4(bx^2+a)^{3/4}a - 3 \right)}{8ax^2}$$

input `int((c*(b*x^2+a)^(1/2))^(3/2)/x^3,x)`output `(sqrt(c)*c*(- 3*a**(3/4)*atan((a**(1/4)*(a + b*x**2)**(3/4)*a - a**(3/4)*(a + b*x**2)**(1/4)*a - a**(3/4)*(a + b*x**2)**(1/4)*b*x**2)/(2*a**2 + 2*a*b*x**2))*b*x**2 - 4*(a + b*x**2)**(3/4)*a - 3*a**(3/4)*log((a + b*x**2)**(1/4) + a**(1/4))*b*x**2 + 3*a**(3/4)*log((a + b*x**2)**(1/4) - a**(1/4))*b*x**2))/(8*a*x**2)`

3.45 $\int x^2 \left(c\sqrt{a + bx^2} \right)^{3/2} dx$

Optimal result	336
Mathematica [C] (verified)	336
Rubi [A] (verified)	337
Maple [F]	339
Fricas [F]	340
Sympy [F]	340
Maxima [F]	340
Giac [F]	341
Mupad [F(-1)]	341
Reduce [F]	341

Optimal result

Integrand size = 21, antiderivative size = 152

$$\int x^2 \left(c\sqrt{a + bx^2} \right)^{3/2} dx = \frac{2ax \left(c\sqrt{a + bx^2} \right)^{3/2}}{15b} + \frac{2}{9} x^3 \left(c\sqrt{a + bx^2} \right)^{3/2} - \frac{4a^2 x \left(c\sqrt{a + bx^2} \right)^{3/2}}{15b(a + bx^2)} + \frac{4a^{3/2} \left(c\sqrt{a + bx^2} \right)^{3/2} E\left(\frac{1}{2} \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) \middle| 2\right)}{15b^{3/2} \left(1 + \frac{bx^2}{a}\right)^{3/4}}$$

output

```
2/15*a*x*(c*(b*x^2+a)^(1/2))^(3/2)/b+2/9*x^3*(c*(b*x^2+a)^(1/2))^(3/2)-4/15*a^2*x*(c*(b*x^2+a)^(1/2))^(3/2)/b/(b*x^2+a)+4/15*a^(3/2)*(c*(b*x^2+a)^(1/2))^(3/2)*EllipticE(sin(1/2*arctan(b^(1/2)*x/a^(1/2))),2^(1/2))/b^(3/2)/(1+b*x^2/a)^(3/4)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.98 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.45

$$\int x^2 \left(c\sqrt{a + bx^2} \right)^{3/2} dx = \frac{2x \left(c\sqrt{a + bx^2} \right)^{3/2} \left(a + bx^2 - \frac{{}_2F_1\left(-\frac{3}{4}, \frac{1}{2}, \frac{3}{2}, -\frac{bx^2}{a}\right)}{\left(1 + \frac{bx^2}{a}\right)^{3/4}} \right)}{9b}$$

input `Integrate[x^2*(c*Sqrt[a + b*x^2])^(3/2),x]`

output `(2*x*(c*Sqrt[a + b*x^2])^(3/2)*(a + b*x^2 - (a*Hypergeometric2F1[-3/4, 1/2, 3/2, -(b*x^2)/a]))/(1 + (b*x^2)/a)^(3/4))/(9*b)`

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.91, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {2045, 248, 262, 225, 212}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 (c\sqrt{a+bx^2})^{3/2} dx \\
 & \quad \downarrow \text{2045} \\
 & \frac{(c\sqrt{a+bx^2})^{3/2} \int x^2 \left(\frac{bx^2}{a} + 1\right)^{3/4} dx}{\left(\frac{bx^2}{a} + 1\right)^{3/4}} \\
 & \quad \downarrow \text{248} \\
 & \frac{(c\sqrt{a+bx^2})^{3/2} \left(\frac{1}{3} \int \frac{x^2}{\sqrt[4]{\frac{bx^2}{a} + 1}} dx + \frac{2}{9} x^3 \left(\frac{bx^2}{a} + 1\right)^{3/4} \right)}{\left(\frac{bx^2}{a} + 1\right)^{3/4}} \\
 & \quad \downarrow \text{262} \\
 & \frac{(c\sqrt{a+bx^2})^{3/2} \left(\frac{1}{3} \left(\frac{2ax \left(\frac{bx^2}{a} + 1\right)^{3/4}}{5b} - \frac{2a \int \frac{1}{\sqrt[4]{\frac{bx^2}{a} + 1}} dx}{5b} \right) + \frac{2}{9} x^3 \left(\frac{bx^2}{a} + 1\right)^{3/4} \right)}{\left(\frac{bx^2}{a} + 1\right)^{3/4}} \\
 & \quad \downarrow \text{225}
 \end{aligned}$$

$$\frac{\left((c\sqrt{a+bx^2})^{3/2} \left(\frac{1}{3} \frac{2ax\left(\frac{bx^2}{a}+1\right)^{3/4}}{5b} - \frac{2a \left(\frac{2x}{\sqrt[4]{\frac{bx^2}{a}+1}} - \int \frac{1}{\left(\frac{bx^2}{a}+1\right)^{5/4}} dx \right)}{5b} \right) + \frac{2}{9}x^3\left(\frac{bx^2}{a}+1\right)^{3/4} \right)}{\left(\frac{bx^2}{a}+1\right)^{3/4}}$$

↓ 212

$$\frac{\left((c\sqrt{a+bx^2})^{3/2} \left(\frac{1}{3} \frac{2ax\left(\frac{bx^2}{a}+1\right)^{3/4}}{5b} - \frac{2a \left(\frac{2x}{\sqrt[4]{\frac{bx^2}{a}+1}} - \frac{2\sqrt{a}E\left(\frac{1}{2}\arctan\left(\frac{\sqrt{bx^2}}{\sqrt{a}}\right)\middle|2\right)}{\sqrt{b}} \right)}{5b} \right) + \frac{2}{9}x^3\left(\frac{bx^2}{a}+1\right)^{3/4} \right)}{\left(\frac{bx^2}{a}+1\right)^{3/4}}$$

input `Int[x^2*(c*Sqrt[a + b*x^2])^(3/2),x]`

output `((c*Sqrt[a + b*x^2])^(3/2))*((2*x^3*(1 + (b*x^2)/a)^(3/4))/9 + ((2*a*x*(1 + (b*x^2)/a)^(3/4))/(5*b) - (2*a*((2*x)/(1 + (b*x^2)/a)^(1/4) - (2*Sqrt[a]*EllipticE[ArcTan[(Sqrt[b]*x)/Sqrt[a]]/2, 2])/Sqrt[b]))/(5*b))/3)/(1 + (b*x^2)/a)^(3/4)`

Defintions of rubi rules used

rule 212 `Int[((a_) + (b_.)*(x_)^2)^(-5/4), x_Symbol] := Simp[(2/(a^(5/4)*Rt[b/a, 2]))*EllipticE[(1/2)*ArcTan[Rt[b/a, 2]*x], 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]`

rule 225 `Int[((a_) + (b_.)*(x_)^2)^(-1/4), x_Symbol] := Simp[2*(x/(a + b*x^2)^(1/4)), x] - Simp[a Int[1/(a + b*x^2)^(5/4), x], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]`

rule 248 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^p/(c*(m + 2*p + 1))), x] + Simp[2*a*(p/(m + 2*p + 1)) Int[(c*x)^m*(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c, m}, x] && GtQ[p, 0] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 262 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_.))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a)^(p*q)) Int[u*(1 + b*(x^n/a)^(p*q)), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [F]

$$\int x^2 \left(c\sqrt{bx^2 + a} \right)^{\frac{3}{2}} dx$$

input `int(x^2*(c*(b*x^2+a)^(1/2))^(3/2),x)`

output `int(x^2*(c*(b*x^2+a)^(1/2))^(3/2),x)`

Fricas [F]

$$\int x^2 (c\sqrt{a+bx^2})^{3/2} dx = \int (\sqrt{bx^2+ac})^{\frac{3}{2}} x^2 dx$$

input `integrate(x^2*(c*(b*x^2+a)^(1/2))^(3/2),x, algorithm="fricas")`

output `integral(sqrt(b*x^2 + a)*sqrt(sqrt(b*x^2 + a)*c)*c*x^2, x)`

Sympy [F]

$$\int x^2 (c\sqrt{a+bx^2})^{3/2} dx = \int x^2 (c\sqrt{a+bx^2})^{\frac{3}{2}} dx$$

input `integrate(x**2*(c*(b*x**2+a)**(1/2))**(3/2),x)`

output `Integral(x**2*(c*sqrt(a + b*x**2))**(3/2), x)`

Maxima [F]

$$\int x^2 (c\sqrt{a+bx^2})^{3/2} dx = \int (\sqrt{bx^2+ac})^{\frac{3}{2}} x^2 dx$$

input `integrate(x^2*(c*(b*x^2+a)^(1/2))^(3/2),x, algorithm="maxima")`

output `integrate((sqrt(b*x^2 + a)*c)^(3/2)*x^2, x)`

Giac [F]

$$\int x^2 (c\sqrt{a+bx^2})^{3/2} dx = \int (\sqrt{bx^2+ac})^{3/2} x^2 dx$$

input `integrate(x^2*(c*(b*x^2+a)^(1/2))^(3/2),x, algorithm="giac")`

output `integrate((sqrt(b*x^2 + a)*c)^(3/2)*x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 (c\sqrt{a+bx^2})^{3/2} dx = \int x^2 (c\sqrt{bx^2+a})^{3/2} dx$$

input `int(x^2*(c*(a + b*x^2)^(1/2))^(3/2),x)`

output `int(x^2*(c*(a + b*x^2)^(1/2))^(3/2), x)`

Reduce [F]

$$\int x^2 (c\sqrt{a+bx^2})^{3/2} dx = \frac{2\sqrt{c}c \left(3(bx^2+a)^{3/4}ax + 5(bx^2+a)^{3/4}bx^3 - 3 \left(\int \frac{1}{(bx^2+a)^{1/4}} dx \right) a^2 \right)}{45b}$$

input `int(x^2*(c*(b*x^2+a)^(1/2))^(3/2),x)`

output `(2*sqrt(c)*c*(3*(a + b*x**2)**(3/4)*a*x + 5*(a + b*x**2)**(3/4)*b*x**3 - 3*int((a + b*x**2)**(3/4)/(a + b*x**2),x)*a**2))/(45*b)`

3.46 $\int \left(c\sqrt{a + bx^2} \right)^{3/2} dx$

Optimal result	342
Mathematica [C] (verified)	342
Rubi [A] (verified)	343
Maple [F]	345
Fricas [F]	345
Sympy [F]	345
Maxima [F]	346
Giac [F]	346
Mupad [F(-1)]	346
Reduce [F]	347

Optimal result

Integrand size = 17, antiderivative size = 119

$$\int \left(c\sqrt{a + bx^2} \right)^{3/2} dx = \frac{2}{5}x \left(c\sqrt{a + bx^2} \right)^{3/2} + \frac{6ax \left(c\sqrt{a + bx^2} \right)^{3/2}}{5(a + bx^2)} - \frac{6\sqrt{a} \left(c\sqrt{a + bx^2} \right)^{3/2} E\left(\frac{1}{2} \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}} \right) \middle| 2 \right)}{5\sqrt{b} \left(1 + \frac{bx^2}{a} \right)^{3/4}}$$

output

```
2/5*x*(c*(b*x^2+a)^(1/2))^(3/2)+6*a*x*(c*(b*x^2+a)^(1/2))^(3/2)/(5*b*x^2+5
*a)-6/5*a^(1/2)*(c*(b*x^2+a)^(1/2))^(3/2)*EllipticE(sin(1/2*arctan(b^(1/2)
*x/a^(1/2))),2^(1/2))/b^(1/2)/(1+b*x^2/a)^(3/4)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.70 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.44

$$\int \left(c\sqrt{a + bx^2} \right)^{3/2} dx = \frac{x \left(c\sqrt{a + bx^2} \right)^{3/2} \text{Hypergeometric2F1}\left(-\frac{3}{4}, \frac{1}{2}, \frac{3}{2}, -\frac{bx^2}{a} \right)}{\left(1 + \frac{bx^2}{a} \right)^{3/4}}$$

input `Integrate[(c*Sqrt[a + b*x^2])^(3/2),x]`

output `(x*(c*Sqrt[a + b*x^2])^(3/2)*Hypergeometric2F1[-3/4, 1/2, 3/2, -((b*x^2)/a)])/(1 + (b*x^2)/a)^(3/4)`

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.88, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {2045, 211, 225, 212}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c\sqrt{a+bx^2})^{3/2} dx \\
 & \quad \downarrow \text{2045} \\
 & \frac{(c\sqrt{a+bx^2})^{3/2} \int \left(\frac{bx^2}{a} + 1\right)^{3/4} dx}{\left(\frac{bx^2}{a} + 1\right)^{3/4}} \\
 & \quad \downarrow \text{211} \\
 & \frac{(c\sqrt{a+bx^2})^{3/2} \left(\frac{3}{5} \int \frac{1}{\sqrt[4]{\frac{bx^2}{a} + 1}} dx + \frac{2}{5} x \left(\frac{bx^2}{a} + 1\right)^{3/4} \right)}{\left(\frac{bx^2}{a} + 1\right)^{3/4}} \\
 & \quad \downarrow \text{225} \\
 & \frac{(c\sqrt{a+bx^2})^{3/2} \left(\frac{3}{5} \left(\frac{2x}{\sqrt[4]{\frac{bx^2}{a} + 1}} - \int \frac{1}{\left(\frac{bx^2}{a} + 1\right)^{5/4}} dx \right) + \frac{2}{5} x \left(\frac{bx^2}{a} + 1\right)^{3/4} \right)}{\left(\frac{bx^2}{a} + 1\right)^{3/4}} \\
 & \quad \downarrow \text{212}
 \end{aligned}$$

$$\frac{(c\sqrt{a+bx^2})^{3/2} \left(\frac{3}{5} \left(\frac{2x}{\sqrt[4]{\frac{bx^2}{a}+1}} - \frac{2\sqrt{a}E\left(\frac{1}{2}\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|2\right)}{\sqrt{b}} \right) + \frac{2}{5}x\left(\frac{bx^2}{a}+1\right)^{3/4} \right)}{\left(\frac{bx^2}{a}+1\right)^{3/4}}$$

input `Int[(c*Sqrt[a + b*x^2])^(3/2),x]`

output `((c*Sqrt[a + b*x^2])^(3/2)*((2*x*(1 + (b*x^2)/a)^(3/4))/5 + (3*((2*x)/(1 + (b*x^2)/a)^(1/4) - (2*Sqrt[a]*EllipticE[ArcTan[(Sqrt[b]*x)/Sqrt[a]]/2, 2])/Sqrt[b]))/5)/(1 + (b*x^2)/a)^(3/4)`

Defintions of rubi rules used

rule 211 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[x*((a + b*x^2)^p/(2*p + 1)), x] + Simp[2*a*(p/(2*p + 1)) Int[(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 212 `Int[((a_) + (b_.)*(x_)^2)^(-5/4), x_Symbol] := Simp[(2/(a^(5/4)*Rt[b/a, 2])*EllipticE[(1/2)*ArcTan[Rt[b/a, 2]*x], 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]`

rule 225 `Int[((a_) + (b_.)*(x_)^2)^(-1/4), x_Symbol] := Simp[2*(x/(a + b*x^2)^(1/4)), x] - Simp[a Int[1/(a + b*x^2)^(5/4), x], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_.))^p, x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a)^(p*q)) Int[u*(1 + b*(x^n/a)^(p*q)), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [F]

$$\int \left(c\sqrt{bx^2 + a} \right)^{\frac{3}{2}} dx$$

input `int((c*(b*x^2+a)^(1/2))^(3/2),x)`

output `int((c*(b*x^2+a)^(1/2))^(3/2),x)`

Fricas [F]

$$\int \left(c\sqrt{a + bx^2} \right)^{3/2} dx = \int \left(\sqrt{bx^2 + ac} \right)^{\frac{3}{2}} dx$$

input `integrate((c*(b*x^2+a)^(1/2))^(3/2),x, algorithm="fricas")`

output `integral(sqrt(b*x^2 + a)*sqrt(sqrt(b*x^2 + a)*c)*c, x)`

Sympy [F]

$$\int \left(c\sqrt{a + bx^2} \right)^{3/2} dx = \int \left(c\sqrt{a + bx^2} \right)^{\frac{3}{2}} dx$$

input `integrate((c*(b*x**2+a)**(1/2))**(3/2),x)`

output `Integral((c*sqrt(a + b*x**2))**(3/2), x)`

Maxima [F]

$$\int (c\sqrt{a+bx^2})^{3/2} dx = \int (\sqrt{bx^2+ac})^{3/2} dx$$

input `integrate((c*(b*x^2+a)^(1/2))^(3/2),x, algorithm="maxima")`

output `integrate((sqrt(b*x^2 + a)*c)^(3/2), x)`

Giac [F]

$$\int (c\sqrt{a+bx^2})^{3/2} dx = \int (\sqrt{bx^2+ac})^{3/2} dx$$

input `integrate((c*(b*x^2+a)^(1/2))^(3/2),x, algorithm="giac")`

output `integrate((sqrt(b*x^2 + a)*c)^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int (c\sqrt{a+bx^2})^{3/2} dx = \int (c\sqrt{bx^2+a})^{3/2} dx$$

input `int((c*(a + b*x^2)^(1/2))^(3/2),x)`

output `int((c*(a + b*x^2)^(1/2))^(3/2), x)`

Reduce [F]

$$\int \left(c\sqrt{a+bx^2} \right)^{3/2} dx = \frac{\sqrt{c}c \left(2(bx^2+a)^{3/4}x + 3 \left(\int \frac{1}{(bx^2+a)^{1/4}} dx \right) a \right)}{5}$$

input `int((c*(b*x^2+a)^(1/2))^(3/2),x)`

output `(sqrt(c)*c*(2*(a + b*x**2)**(3/4)*x + 3*int((a + b*x**2)**(3/4)/(a + b*x**2),x)*a))/5`

3.47 $\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^2} dx$

Optimal result	348
Mathematica [C] (verified)	348
Rubi [A] (verified)	349
Maple [F]	351
Fricas [F]	351
Sympy [F]	352
Maxima [F]	352
Giac [F]	352
Mupad [F(-1)]	353
Reduce [F]	353

Optimal result

Integrand size = 21, antiderivative size = 115

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^2} dx = -\frac{(c\sqrt{a+bx^2})^{3/2}}{x} + \frac{3bx(c\sqrt{a+bx^2})^{3/2}}{a+bx^2} - \frac{3\sqrt{b}(c\sqrt{a+bx^2})^{3/2} E\left(\frac{1}{2} \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) \middle| 2\right)}{\sqrt{a} \left(1 + \frac{bx^2}{a}\right)^{3/4}}$$

output

```
-(c*(b*x^2+a)^(1/2))^(3/2)/x+3*b*x*(c*(b*x^2+a)^(1/2))^(3/2)/(b*x^2+a)-3*b
^(1/2)*(c*(b*x^2+a)^(1/2))^(3/2)*EllipticE(sin(1/2*arctan(b^(1/2)*x/a^(1/2)
)),2^(1/2))/a^(1/2)/(1+b*x^2/a)^(3/4)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.21 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.48

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^2} dx = -\frac{(c\sqrt{a+bx^2})^{3/2} \text{Hypergeometric2F1}\left(-\frac{3}{4}, -\frac{1}{2}, \frac{1}{2}, -\frac{bx^2}{a}\right)}{x \left(1 + \frac{bx^2}{a}\right)^{3/4}}$$

input `Integrate[(c*Sqrt[a + b*x^2])^(3/2)/x^2,x]`

output `-(((c*Sqrt[a + b*x^2])^(3/2)*Hypergeometric2F1[-3/4, -1/2, 1/2, -((b*x^2)/a)])/(x*(1 + (b*x^2)/a)^(3/4)))`

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.95, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2045, 247, 225, 212}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^2} dx \\
 & \quad \downarrow \text{2045} \\
 & \frac{(c\sqrt{a+bx^2})^{3/2} \int \frac{(\frac{bx^2}{a}+1)^{3/4}}{x^2} dx}{(\frac{bx^2}{a}+1)^{3/4}} \\
 & \quad \downarrow \text{247} \\
 & \frac{(c\sqrt{a+bx^2})^{3/2} \left(\frac{3b \int \frac{1}{\sqrt[4]{\frac{bx^2}{a}+1}} dx}{2a} - \frac{(\frac{bx^2}{a}+1)^{3/4}}{x} \right)}{(\frac{bx^2}{a}+1)^{3/4}} \\
 & \quad \downarrow \text{225}
 \end{aligned}$$

$$\frac{\left(c\sqrt{a+bx^2} \right)^{3/2} \left(\frac{3b \left(\frac{2x}{\sqrt[4]{\frac{bx^2}{a}+1}} - \int \frac{1}{\left(\frac{bx^2}{a}+1\right)^{5/4}} dx \right)}{2a} - \frac{\left(\frac{bx^2}{a}+1\right)^{3/4}}{x} \right)}{\left(\frac{bx^2}{a}+1\right)^{3/4}}$$

↓ 212

$$\frac{\left(c\sqrt{a+bx^2} \right)^{3/2} \left(\frac{3b \left(\frac{2x}{\sqrt[4]{\frac{bx^2}{a}+1}} - \frac{2\sqrt{a}E\left(\frac{1}{2}\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|2\right)}{\sqrt{b}} \right)}{2a} - \frac{\left(\frac{bx^2}{a}+1\right)^{3/4}}{x} \right)}{\left(\frac{bx^2}{a}+1\right)^{3/4}}$$

input `Int[(c*Sqrt[a + b*x^2])^(3/2)/x^2,x]`

output `((c*Sqrt[a + b*x^2])^(3/2))*(-(1 + (b*x^2)/a)^(3/4)/x) + (3*b*((2*x)/(1 + (b*x^2)/a)^(1/4) - (2*Sqrt[a]*EllipticE[ArcTan[(Sqrt[b]*x)/Sqrt[a]]/2, 2])/Sqrt[b]))/(2*a))/(1 + (b*x^2)/a)^(3/4)`

Defintions of rubi rules used

rule 212 `Int[((a_) + (b_.)*(x_)^2)^(-5/4), x_Symbol] := Simp[(2/(a^(5/4)*Rt[b/a, 2]))*EllipticE[(1/2)*ArcTan[Rt[b/a, 2]*x], 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]`

rule 225 `Int[((a_) + (b_.)*(x_)^2)^(-1/4), x_Symbol] := Simp[2*(x/(a + b*x^2)^(1/4)), x] - Simp[a Int[1/(a + b*x^2)^(5/4), x], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]`

rule 247

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(
(m + 1)*((a + b*x^2)^p/(c*(m + 1))), x] - Simp[2*b*(p/(c^2*(m + 1))) Int[
(c*x)^(m + 2)*(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && GtQ[p,
0] && LtQ[m, -1] && !LtQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2,
m, p, x]
```

rule 2045

```
Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_))^(p_), x_Symbol] := Simp[Si
mp[(c*(a + b*x^n)^q]^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q)
, x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]
```

Maple [F]

$$\int \frac{(c\sqrt{bx^2+a})^{\frac{3}{2}}}{x^2} dx$$

input

```
int((c*(b*x^2+a)^(1/2))^(3/2)/x^2,x)
```

output

```
int((c*(b*x^2+a)^(1/2))^(3/2)/x^2,x)
```

Fricas [F]

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^2} dx = \int \frac{(\sqrt{bx^2+ac})^{\frac{3}{2}}}{x^2} dx$$

input

```
integrate((c*(b*x^2+a)^(1/2))^(3/2)/x^2,x, algorithm="fricas")
```

output

```
integral(sqrt(b*x^2 + a)*sqrt(sqrt(b*x^2 + a)*c)*c/x^2, x)
```


Sympy [F]

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^2} dx = \int \frac{(c\sqrt{a+bx^2})^{\frac{3}{2}}}{x^2} dx$$

input `integrate((c*(b*x**2+a)**(1/2))**(3/2)/x**2,x)`

output `Integral((c*sqrt(a + b*x**2))**(3/2)/x**2, x)`

Maxima [F]

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^2} dx = \int \frac{(\sqrt{bx^2+ac})^{\frac{3}{2}}}{x^2} dx$$

input `integrate((c*(b*x^2+a)^(1/2))^(3/2)/x^2,x, algorithm="maxima")`

output `integrate((sqrt(b*x^2 + a)*c)^(3/2)/x^2, x)`

Giac [F]

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^2} dx = \int \frac{(\sqrt{bx^2+ac})^{\frac{3}{2}}}{x^2} dx$$

input `integrate((c*(b*x^2+a)^(1/2))^(3/2)/x^2,x, algorithm="giac")`

output `integrate((sqrt(b*x^2 + a)*c)^(3/2)/x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^2} dx = \int \frac{(c\sqrt{bx^2+a})^{3/2}}{x^2} dx$$

input `int((c*(a + b*x^2)^(1/2))^(3/2)/x^2,x)`output `int((c*(a + b*x^2)^(1/2))^(3/2)/x^2, x)`**Reduce [F]**

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^2} dx = \frac{\sqrt{c}c\left(2(bx^2+a)^{3/4} + 3\left(\int \frac{(bx^2+a)^{3/4}}{bx^4+ax^2} dx\right)ax\right)}{x}$$

input `int((c*(b*x^2+a)^(1/2))^(3/2)/x^2,x)`output `(sqrt(c)*c*(2*(a + b*x**2)**(3/4) + 3*int((a + b*x**2)**(3/4)/(a*x**2 + b*x**4),x)*a*x))/x`

3.48
$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^4} dx$$

Optimal result	354
Mathematica [C] (verified)	355
Rubi [A] (verified)	355
Maple [F]	358
Fricas [F]	358
Sympy [F]	359
Maxima [F]	359
Giac [F]	359
Mupad [F(-1)]	360
Reduce [F]	360

Optimal result

Integrand size = 21, antiderivative size = 154

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^4} dx = -\frac{(c\sqrt{a+bx^2})^{3/2}}{3x^3} - \frac{b(c\sqrt{a+bx^2})^{3/2}}{2ax} + \frac{b^2x(c\sqrt{a+bx^2})^{3/2}}{2a(a+bx^2)} - \frac{b^{3/2}(c\sqrt{a+bx^2})^{3/2} E\left(\frac{1}{2} \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) \middle| 2\right)}{2a^{3/2} \left(1 + \frac{bx^2}{a}\right)^{3/4}}$$

output

```
-1/3*(c*(b*x^2+a)^(1/2))^(3/2)/x^3-1/2*b*(c*(b*x^2+a)^(1/2))^(3/2)/a/x+1/2
*b^2*x*(c*(b*x^2+a)^(1/2))^(3/2)/a/(b*x^2+a)-1/2*b^(3/2)*(c*(b*x^2+a)^(1/2))
)^(3/2)*EllipticE(sin(1/2*arctan(b^(1/2)*x/a^(1/2))),2^(1/2))/a^(3/2)/(1+
b*x^2/a)^(3/4)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.01 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.37

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^4} dx = -\frac{(c\sqrt{a+bx^2})^{3/2} \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, -\frac{3}{4}, -\frac{1}{2}, -\frac{bx^2}{a}\right)}{3x^3 \left(1 + \frac{bx^2}{a}\right)^{3/4}}$$

input `Integrate[(c*Sqrt[a + b*x^2])^(3/2)/x^4,x]`

output `-1/3*((c*Sqrt[a + b*x^2])^(3/2)*Hypergeometric2F1[-3/2, -3/4, -1/2, -(b*x^2/a)])/(x^3*(1 + (b*x^2)/a)^(3/4))`

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.90, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {2045, 247, 264, 225, 212}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^4} dx \\ & \quad \downarrow \text{2045} \\ & \frac{(c\sqrt{a+bx^2})^{3/2} \int \frac{(\frac{bx^2}{a}+1)^{3/4}}{x^4} dx}{\left(\frac{bx^2}{a}+1\right)^{3/4}} \\ & \quad \downarrow \text{247} \end{aligned}$$

$$\begin{aligned}
 & \frac{(c\sqrt{a+bx^2})^{3/2} \left(\frac{b \int \frac{1}{x^2 \sqrt[4]{\frac{bx^2}{a} + 1}} dx}{2a} - \frac{(\frac{bx^2}{a} + 1)^{3/4}}{3x^3} \right)}{(\frac{bx^2}{a} + 1)^{3/4}} \\
 & \quad \downarrow \text{264} \\
 & \frac{(c\sqrt{a+bx^2})^{3/2} \left(\frac{b \left(\frac{b \int \frac{1}{x^2 \sqrt[4]{\frac{bx^2}{a} + 1}} dx}{2a} - \frac{(\frac{bx^2}{a} + 1)^{3/4}}{x} \right)}{2a} - \frac{(\frac{bx^2}{a} + 1)^{3/4}}{3x^3} \right)}{(\frac{bx^2}{a} + 1)^{3/4}} \\
 & \quad \downarrow \text{225} \\
 & \frac{(c\sqrt{a+bx^2})^{3/2} \left(\frac{b \left(\frac{b \left(\frac{2x}{\sqrt[4]{\frac{bx^2}{a} + 1}} - \int \frac{1}{(\frac{bx^2}{a} + 1)^{5/4}} dx \right)}{2a} - \frac{(\frac{bx^2}{a} + 1)^{3/4}}{x} \right)}{2a} - \frac{(\frac{bx^2}{a} + 1)^{3/4}}{3x^3} \right)}{(\frac{bx^2}{a} + 1)^{3/4}} \\
 & \quad \downarrow \text{212}
 \end{aligned}$$

$$\left(c\sqrt{a+bx^2} \right)^{3/2} \left(\frac{b \left(\frac{2x}{\sqrt[4]{\frac{bx^2}{a}+1}} - \frac{2\sqrt{a}E\left(\frac{1}{2}\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|2\right)}{\sqrt{b}} \right)}{2a} - \frac{\left(\frac{bx^2}{a}+1\right)^{3/4}}{x} \right)}{2a} - \frac{\left(\frac{bx^2}{a}+1\right)^{3/4}}{3x^3} \right)$$

$$\left(\frac{bx^2}{a} + 1\right)^{3/4}$$

input `Int[(c*Sqrt[a + b*x^2])^(3/2)/x^4,x]`

output `((c*Sqrt[a + b*x^2])^(3/2)*(-1/3*(1 + (b*x^2)/a)^(3/4)/x^3 + (b*(-((1 + (b*x^2)/a)^(3/4)/x) + (b*((2*x)/(1 + (b*x^2)/a)^(1/4) - (2*Sqrt[a]*EllipticE[ArcTan[(Sqrt[b]*x)/Sqrt[a]]/2, 2)]/Sqrt[b]))/(2*a)))/(2*a)))/(1 + (b*x^2)/a)^(3/4)`

Defintions of rubi rules used

rule 212 `Int[((a_) + (b_.)*(x_)^2)^(-5/4), x_Symbol] := Simp[(2/(a^(5/4)*Rt[b/a, 2]))*EllipticE[(1/2)*ArcTan[Rt[b/a, 2]*x], 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]`

rule 225 `Int[((a_) + (b_.)*(x_)^2)^(-1/4), x_Symbol] := Simp[2*(x/(a + b*x^2)^(1/4)), x] - Simp[a Int[1/(a + b*x^2)^(5/4), x], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]`

rule 247 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^p/(c*(m + 1))), x] - Simp[2*b*(p/(c^2*(m + 1))) Int[(c*x)^(m + 2)*(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && GtQ[p, 0] && LtQ[m, -1] && !LtQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 264 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*(m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [F]

$$\int \frac{(c\sqrt{bx^2+a})^{\frac{3}{2}}}{x^4} dx$$

input `int((c*(b*x^2+a)^(1/2))^(3/2)/x^4,x)`

output `int((c*(b*x^2+a)^(1/2))^(3/2)/x^4,x)`

Fricas [F]

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^4} dx = \int \frac{(\sqrt{bx^2+ac})^{\frac{3}{2}}}{x^4} dx$$

input `integrate((c*(b*x^2+a)^(1/2))^(3/2)/x^4,x, algorithm="fricas")`

output `integral(sqrt(b*x^2 + a)*sqrt(sqrt(b*x^2 + a)*c)*c/x^4, x)`

Sympy [F]

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^4} dx = \int \frac{(c\sqrt{a+bx^2})^{\frac{3}{2}}}{x^4} dx$$

input `integrate((c*(b*x**2+a)**(1/2))**(3/2)/x**4,x)`

output `Integral((c*sqrt(a + b*x**2))**(3/2)/x**4, x)`

Maxima [F]

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^4} dx = \int \frac{(\sqrt{bx^2+ac})^{\frac{3}{2}}}{x^4} dx$$

input `integrate((c*(b*x^2+a)^(1/2))^(3/2)/x^4,x, algorithm="maxima")`

output `integrate((sqrt(b*x^2 + a)*c)^(3/2)/x^4, x)`

Giac [F]

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^4} dx = \int \frac{(\sqrt{bx^2+ac})^{\frac{3}{2}}}{x^4} dx$$

input `integrate((c*(b*x^2+a)^(1/2))^(3/2)/x^4,x, algorithm="giac")`

output `integrate((sqrt(b*x^2 + a)*c)^(3/2)/x^4, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^4} dx = \int \frac{(c\sqrt{bx^2+a})^{3/2}}{x^4} dx$$

input `int((c*(a + b*x^2)^(1/2))^(3/2)/x^4, x)`output `int((c*(a + b*x^2)^(1/2))^(3/2)/x^4, x)`**Reduce [F]**

$$\int \frac{(c\sqrt{a+bx^2})^{3/2}}{x^4} dx = \frac{\sqrt{c}c \left(-2(bx^2+a)^{3/4} - 3 \left(\int \frac{(bx^2+a)^{3/4}}{bx^6+ax^4} dx \right) ax^3 \right)}{3x^3}$$

input `int((c*(b*x^2+a)^(1/2))^(3/2)/x^4, x)`output `(sqrt(c)*c*(- 2*(a + b*x**2)**(3/4) - 3*int((a + b*x**2)**(3/4)/(a*x**4 + b*x**6), x)*a*x**3))/(3*x**3)`

3.49 $\int x^7 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx$

Optimal result	361
Mathematica [A] (verified)	361
Rubi [A] (verified)	362
Maple [A] (verified)	363
Fricas [A] (verification not implemented)	364
Sympy [A] (verification not implemented)	364
Maxima [A] (verification not implemented)	365
Giac [A] (verification not implemented)	365
Mupad [B] (verification not implemented)	365
Reduce [B] (verification not implemented)	366

Optimal result

Integrand size = 21, antiderivative size = 136

$$\int x^7 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = -\frac{2a^3 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} (a+bx^2)}{b^4} + \frac{6a^2 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} (a+bx^2)^2}{5b^4} - \frac{2a \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} (a+bx^2)^3}{3b^4} + \frac{2 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} (a+bx^2)^4}{13b^4}$$

output `-2*a^3*(c/(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)/b^4+6/5*a^2*(c/(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)^2/b^4-2/3*a*(c/(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)^3/b^4+2/13*(c/(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)^4/b^4`

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.43

$$\int x^7 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \frac{2c^2(-128a^3 + 32a^2bx^2 - 20ab^2x^4 + 15b^3x^6)}{195b^4 \sqrt{\frac{c}{\sqrt{a+bx^2}}}}$$

input `Integrate[x^7*(c/Sqrt[a + b*x^2])^(3/2),x]`

output

```
(2*c^2*(-128*a^3 + 32*a^2*b*x^2 - 20*a*b^2*x^4 + 15*b^3*x^6))/(195*b^4*Sqr
t[c/Sqrt[a + b*x^2]])
```

Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.96, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2045, 243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^7 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx$$

$$\downarrow \text{2045}$$

$$\left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} \int \frac{x^7}{\left(\frac{bx^2}{a} + 1 \right)^{3/4}} dx$$

$$\downarrow \text{243}$$

$$\frac{1}{2} \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} \int \frac{x^6}{\left(\frac{bx^2}{a} + 1 \right)^{3/4}} dx^2$$

$$\downarrow \text{53}$$

$$\frac{1}{2} \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} \int \left(\frac{\left(\frac{bx^2}{a} + 1 \right)^{9/4} a^3}{b^3} - \frac{3 \left(\frac{bx^2}{a} + 1 \right)^{5/4} a^3}{b^3} + \frac{3 \sqrt[4]{\frac{bx^2}{a} + 1} a^3}{b^3} - \frac{a^3}{b^3 \left(\frac{bx^2}{a} + 1 \right)^{3/4}} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{1}{2} \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} \left(\frac{4a^4 \left(\frac{bx^2}{a} + 1 \right)^{13/4}}{13b^4} - \frac{4a^4 \left(\frac{bx^2}{a} + 1 \right)^{9/4}}{3b^4} + \frac{12a^4 \left(\frac{bx^2}{a} + 1 \right)^{5/4}}{5b^4} - \frac{4a^4 \sqrt[4]{\frac{bx^2}{a} + 1}}{b^4} \right) \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2}$$

input `Int[x^7*(c/Sqrt[a + b*x^2])^(3/2),x]`

output
$$\left(\frac{c}{\sqrt{a + bx^2}}\right)^{3/2} \left(1 + \frac{bx^2}{a}\right)^{3/4} \left(\frac{-4a^4(1 + \frac{bx^2}{a})^{1/4}}{b^4} + \frac{12a^4(1 + \frac{bx^2}{a})^{5/4}}{(5b^4)} - \frac{4a^4(1 + \frac{bx^2}{a})^{9/4}}{(3b^4)} + \frac{4a^4(1 + \frac{bx^2}{a})^{13/4}}{(13b^4)}\right) / 2$$

Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.43

method	result	size
gospers	$\frac{2(bx^2+a)(-15b^3x^6+20ab^2x^4-32a^2bx^2+128a^3)\left(\frac{c}{\sqrt{bx^2+a}}\right)^{3/2}}{195b^4}$	58
orering	$\frac{2(bx^2+a)(-15b^3x^6+20ab^2x^4-32a^2bx^2+128a^3)\left(\frac{c}{\sqrt{bx^2+a}}\right)^{3/2}}{195b^4}$	58

input `int(x^7*(c/(b*x^2+a)^(1/2))^(3/2),x,method=_RETURNVERBOSE)`

output
$$-2/195*(b*x^2+a)*(-15*b^3*x^6+20*a*b^2*x^4-32*a^2*b*x^2+128*a^3)*(c/(b*x^2+a)^(1/2))^(3/2)/b^4$$

Fricas [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.46

$$\int x^7 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \frac{2(15b^3cx^6 - 20ab^2cx^4 + 32a^2bcx^2 - 128a^3c)\sqrt{bx^2+a}\sqrt{\frac{c}{\sqrt{bx^2+a}}}}{195b^4}$$

input `integrate(x^7*(c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="fricas")`

output
$$2/195*(15*b^3*c*x^6 - 20*a*b^2*c*x^4 + 32*a^2*b*c*x^2 - 128*a^3*c)*\text{sqrt}(b*x^2 + a)*\text{sqrt}(c/\text{sqrt}(b*x^2 + a))/b^4$$

Sympy [A] (verification not implemented)

Time = 10.73 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.06

$$\int x^7 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \begin{cases} -\frac{256a^4 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2}}{195b^4} - \frac{64a^3x^2 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2}}{65b^3} + \frac{8a^2x^4 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2}}{65b^2} - \frac{2ax^6 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2}}{39b} + \frac{2x^8}{8} \\ \frac{x^8 \left(\frac{c}{\sqrt{a}} \right)^{3/2}}{8} \end{cases}$$

input `integrate(x**7*(c/(b*x**2+a)**(1/2))**(3/2),x)`

output `Piecewise((-256*a**4*(c/sqrt(a + b*x**2))**(3/2)/(195*b**4) - 64*a**3*x**2*(c/sqrt(a + b*x**2))**(3/2)/(65*b**3) + 8*a**2*x**4*(c/sqrt(a + b*x**2))**(3/2)/(65*b**2) - 2*a*x**6*(c/sqrt(a + b*x**2))**(3/2)/(39*b) + 2*x**8*(c/sqrt(a + b*x**2))**(3/2)/13, Ne(b, 0)), (x**8*(c/sqrt(a))**(3/2)/8, True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.56

$$\int x^7 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = -\frac{2 \left(\frac{65ac^6}{bx^2+a} - \frac{117a^2c^6}{(bx^2+a)^2} + \frac{195a^3c^6}{(bx^2+a)^3} - 15c^6 \right) c^2}{195b^4 \left(\frac{c}{\sqrt{bx^2+a}} \right)^{\frac{13}{2}}}$$

input `integrate(x^7*(c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="maxima")`output
$$-2/195*(65*a*c^6/(b*x^2 + a) - 117*a^2*c^6/(b*x^2 + a)^2 + 195*a^3*c^6/(b*x^2 + a)^3 - 15*c^6)*c^2/(b^4*(c/\text{sqrt}(b*x^2 + a))^(13/2))$$
Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.51

$$\int x^7 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \frac{2 \left(15 (bx^2 + a)^{\frac{13}{4}} - 65 (bx^2 + a)^{\frac{9}{4}} a + 117 (bx^2 + a)^{\frac{5}{4}} a^2 - 195 (bx^2 + a)^{\frac{1}{4}} a^3 \right) c^{\frac{3}{2}}}{195 b^4 \sqrt{\text{sgn}(bx^2 + a)}}$$

input `integrate(x^7*(c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="giac")`output
$$2/195*(15*(b*x^2 + a)^(13/4) - 65*(b*x^2 + a)^(9/4)*a + 117*(b*x^2 + a)^(5/4)*a^2 - 195*(b*x^2 + a)^(1/4)*a^3)*c^(3/2)/(b^4*\text{sqrt}(\text{sgn}(b*x^2 + a)))$$
Mupad [B] (verification not implemented)

Time = 0.55 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.67

$$\int x^7 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \sqrt{\frac{c}{\sqrt{bx^2+a}}} \left(\frac{2cx^6\sqrt{bx^2+a}}{13b} - \frac{256a^3c\sqrt{bx^2+a}}{195b^4} - \frac{8acx^4\sqrt{bx^2+a}}{39b^2} + \frac{64a^2cx^2\sqrt{bx^2+a}}{195b^3} \right)$$

input `int(x^7*(c/(a + b*x^2)^(1/2))^(3/2),x)`

output `(c/(a + b*x^2)^(1/2))^(1/2)*((2*c*x^6*(a + b*x^2)^(1/2))/(13*b) - (256*a^3*c*(a + b*x^2)^(1/2))/(195*b^4) - (8*a*c*x^4*(a + b*x^2)^(1/2))/(39*b^2) + (64*a^2*c*x^2*(a + b*x^2)^(1/2))/(195*b^3))`

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.10

$$\int x^7 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \frac{2\sqrt{c} \sqrt{\sqrt{b} \sqrt{bx^2 + a}} x + a + bx^2 \sqrt{\sqrt{bx^2 + a} + \sqrt{b}} x c \left(-128\sqrt{bx^2 + a} a^3 + 32\sqrt{bx^2 + a} a^2 + 32\sqrt{bx^2 + a} a - 128\sqrt{bx^2 + a} \right)}{195 a^4 b^3}$$

input `int(x^7*(c/(b*x^2+a)^(1/2))^(3/2),x)`

output `(2*sqrt(c)*sqrt(sqrt(b)*sqrt(a + b*x**2)*x + a + b*x**2)*sqrt(sqrt(a + b*x**2) + sqrt(b)*x)*c*(- 128*sqrt(a + b*x**2)*a**3 + 32*sqrt(a + b*x**2)*a**2*b*x**2 - 20*sqrt(a + b*x**2)*a*b**2*x**4 + 15*sqrt(a + b*x**2)*b**3*x**6 + 128*sqrt(b)*a**3*x - 32*sqrt(b)*a**2*b*x**3 + 20*sqrt(b)*a*b**2*x**5 - 15*sqrt(b)*b**3*x**7))/(195*a*b**4)`

3.50 $\int x^5 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx$

Optimal result	367
Mathematica [A] (verified)	367
Rubi [A] (verified)	368
Maple [A] (verified)	369
Fricas [A] (verification not implemented)	370
Sympy [A] (verification not implemented)	370
Maxima [A] (verification not implemented)	371
Giac [A] (verification not implemented)	371
Mupad [B] (verification not implemented)	371
Reduce [B] (verification not implemented)	372

Optimal result

Integrand size = 21, antiderivative size = 100

$$\int x^5 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \frac{2a^2 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} (a+bx^2)}{b^3} - \frac{4a \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} (a+bx^2)^2}{5b^3} + \frac{2 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} (a+bx^2)^3}{9b^3}$$

output `2*a^2*(c/(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)/b^3-4/5*a*(c/(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)^2/b^3+2/9*(c/(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)^3/b^3`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.48

$$\int x^5 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \frac{2c^2(32a^2 - 8abx^2 + 5b^2x^4)}{45b^3 \sqrt{\frac{c}{a+bx^2}}}$$

input `Integrate[x^5*(c/Sqrt[a + b*x^2])^(3/2),x]`

output $(2*c^2*(32*a^2 - 8*a*b*x^2 + 5*b^2*x^4))/(45*b^3*sqrt[c/Sqrt[a + b*x^2]])$

Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.06, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2045, 243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^5 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx$$

↓ 2045

$$\left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} \int \frac{x^5}{\left(\frac{bx^2}{a} + 1 \right)^{3/4}} dx$$

↓ 243

$$\frac{1}{2} \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} \int \frac{x^4}{\left(\frac{bx^2}{a} + 1 \right)^{3/4}} dx^2$$

↓ 53

$$\frac{1}{2} \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} \int \left(\frac{\left(\frac{bx^2}{a} + 1 \right)^{5/4} a^2}{b^2} - \frac{2 \sqrt[4]{\frac{bx^2}{a} + 1} a^2}{b^2} + \frac{a^2}{b^2 \left(\frac{bx^2}{a} + 1 \right)^{3/4}} \right) dx^2$$

↓ 2009

$$\frac{1}{2} \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} \left(\frac{4a^3 \left(\frac{bx^2}{a} + 1 \right)^{9/4}}{9b^3} - \frac{8a^3 \left(\frac{bx^2}{a} + 1 \right)^{5/4}}{5b^3} + \frac{4a^3 \sqrt[4]{\frac{bx^2}{a} + 1}}{b^3} \right)$$

input $\text{Int}[x^5*(c/Sqrt[a + b*x^2])^(3/2), x]$

output

$$\left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} (1 + (bx^2)/a)^{3/4} \left(\frac{4a^3(1 + (bx^2)/a)^{1/4}}{b^3} - \frac{8a^3(1 + (bx^2)/a)^{5/4}}{(5b^3)} + \frac{4a^3(1 + (bx^2)/a)^{9/4}}{(9b^3)} \right) / 2$$
Defintions of rubi rules used

rule 53

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && IGtQ[m, 0] && ( !IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0])
|| LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

rule 243

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int
[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && I
ntegerQ[(m - 1)/2]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2045

```
Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_))^(p_), x_Symbol] := Simp[Simp
[(c*(a + b*x^n)^q]^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q)
, x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]
```

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.47

method	result	size
gospers	$\frac{2(bx^2+a)(5b^2x^4-8abx^2+32a^2)\left(\frac{c}{\sqrt{bx^2+a}}\right)^{\frac{3}{2}}}{45b^3}$	47
orering	$\frac{2(bx^2+a)(5b^2x^4-8abx^2+32a^2)\left(\frac{c}{\sqrt{bx^2+a}}\right)^{\frac{3}{2}}}{45b^3}$	47

input

```
int(x^5*(c/(b*x^2+a)^(1/2))^(3/2),x,method=_RETURNVERBOSE)
```

output $2/45*(b*x^2+a)*(5*b^2*x^4-8*a*b*x^2+32*a^2)*(c/(b*x^2+a)^{(1/2)})^{(3/2)}/b^3$

Fricas [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.51

$$\int x^5 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \frac{2(5b^2cx^4 - 8abcx^2 + 32a^2c)\sqrt{bx^2+a}\sqrt{\frac{c}{\sqrt{bx^2+a}}}}{45b^3}$$

input `integrate(x^5*(c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="fricas")`

output $2/45*(5*b^2*c*x^4 - 8*a*b*c*x^2 + 32*a^2*c)*\text{sqrt}(b*x^2 + a)*\text{sqrt}(c/\text{sqrt}(b*x^2 + a))/b^3$

Sympy [A] (verification not implemented)

Time = 5.77 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.16

$$\int x^5 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \begin{cases} \frac{64a^3 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2}}{45b^3} + \frac{16a^2x^2 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2}}{15b^2} - \frac{2ax^4 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2}}{15b} + \frac{2x^6 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2}}{9} & \text{for } b \neq 0 \\ \frac{x^6 \left(\frac{c}{\sqrt{a}} \right)^{3/2}}{6} & \text{otherwise} \end{cases}$$

input `integrate(x**5*(c/(b*x**2+a)**(1/2))**(3/2),x)`

output `Piecewise((64*a**3*(c/sqrt(a + b*x**2))**(3/2)/(45*b**3) + 16*a**2*x**2*(c/sqrt(a + b*x**2))**(3/2)/(15*b**2) - 2*a*x**4*(c/sqrt(a + b*x**2))**(3/2)/(15*b) + 2*x**6*(c/sqrt(a + b*x**2))**(3/2)/9, Ne(b, 0)), (x**6*(c/sqrt(a))**(3/2)/6, True))`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.59

$$\int x^5 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = -\frac{2 \left(\frac{18ac^4}{bx^2+a} - \frac{45a^2c^4}{(bx^2+a)^2} - 5c^4 \right) c^2}{45b^3 \left(\frac{c}{\sqrt{bx^2+a}} \right)^{\frac{9}{2}}}$$

input `integrate(x^5*(c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="maxima")`output `-2/45*(18*a*c^4/(b*x^2 + a) - 45*a^2*c^4/(b*x^2 + a)^2 - 5*c^4)*c^2/(b^3*(c/sqrt(b*x^2 + a))^(9/2))`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.56

$$\int x^5 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \frac{2 \left(5(bx^2+a)^{\frac{9}{4}} - 18(bx^2+a)^{\frac{5}{4}}a + 45(bx^2+a)^{\frac{1}{4}}a^2 \right) c^{\frac{3}{2}}}{45b^3 \sqrt{\text{sgn}(bx^2+a)}}$$

input `integrate(x^5*(c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="giac")`output `2/45*(5*(b*x^2 + a)^(9/4) - 18*(b*x^2 + a)^(5/4)*a + 45*(b*x^2 + a)^(1/4)*a^2)*c^(3/2)/(b^3*sqrt(sgn(b*x^2 + a)))`**Mupad [B] (verification not implemented)**

Time = 0.55 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.70

$$\int x^5 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \sqrt{\frac{c}{\sqrt{bx^2+a}}} \left(\frac{2cx^4\sqrt{bx^2+a}}{9b} + \frac{64a^2c\sqrt{bx^2+a}}{45b^3} - \frac{16acx^2\sqrt{bx^2+a}}{45b^2} \right)$$

input `int(x^5*(c/(a + b*x^2)^(1/2))^(3/2),x)`

output `(c/(a + b*x^2)^(1/2))^(1/2)*((2*c*x^4*(a + b*x^2)^(1/2))/(9*b) + (64*a^2*c*(a + b*x^2)^(1/2))/(45*b^3) - (16*a*c*x^2*(a + b*x^2)^(1/2))/(45*b^2))`

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.17

$$\int x^5 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \frac{2\sqrt{c} \sqrt{\sqrt{b} \sqrt{bx^2 + a}} x + a + bx^2 \sqrt{\sqrt{bx^2 + a} + \sqrt{b}} c (32\sqrt{bx^2 + a} a^2 - 8\sqrt{b} a^2 + 5\sqrt{a + bx^2} b^2 x^4 - 32\sqrt{b} a^2 x + 8\sqrt{b} a b x^3 - 5\sqrt{b} b^2 x^5)}{45 a b^3}$$

input `int(x^5*(c/(b*x^2+a)^(1/2))^(3/2),x)`

output `(2*sqrt(c)*sqrt(sqrt(b)*sqrt(a + b*x**2)*x + a + b*x**2)*sqrt(sqrt(a + b*x**2) + sqrt(b)*x)*c*(32*sqrt(a + b*x**2)*a**2 - 8*sqrt(a + b*x**2)*a*b*x**2 + 5*sqrt(a + b*x**2)*b**2*x**4 - 32*sqrt(b)*a**2*x + 8*sqrt(b)*a*b*x**3 - 5*sqrt(b)*b**2*x**5))/(45*a*b**3)`

3.51 $\int x^3 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx$

Optimal result	373
Mathematica [A] (verified)	373
Rubi [A] (verified)	374
Maple [A] (verified)	375
Fricas [A] (verification not implemented)	376
Sympy [A] (verification not implemented)	376
Maxima [A] (verification not implemented)	376
Giac [A] (verification not implemented)	377
Mupad [B] (verification not implemented)	377
Reduce [B] (verification not implemented)	378

Optimal result

Integrand size = 21, antiderivative size = 64

$$\int x^3 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = -\frac{2a \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} (a+bx^2)}{b^2} + \frac{2 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} (a+bx^2)^2}{5b^2}$$

output `-2*a*(c/(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)/b^2+2/5*(c/(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)^2/b^2`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.56

$$\int x^3 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \frac{2c^2(-4a+bx^2)}{5b^2 \sqrt{\frac{c}{\sqrt{a+bx^2}}}}$$

input `Integrate[x^3*(c/Sqrt[a + b*x^2])^(3/2),x]`

output `(2*c^2*(-4*a + b*x^2))/(5*b^2*Sqrt[c/Sqrt[a + b*x^2]])`

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.28, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2045, 243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx \\
 & \quad \downarrow \text{2045} \\
 & \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} \int \frac{x^3}{\left(\frac{bx^2}{a} + 1 \right)^{3/4}} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{2} \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} \int \frac{x^2}{\left(\frac{bx^2}{a} + 1 \right)^{3/4}} dx^2 \\
 & \quad \downarrow \text{53} \\
 & \frac{1}{2} \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} \int \left(\frac{a \sqrt[4]{\frac{bx^2}{a} + 1}}{b} - \frac{a}{b \left(\frac{bx^2}{a} + 1 \right)^{3/4}} \right) dx^2 \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{2} \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} \left(\frac{4a^2 \left(\frac{bx^2}{a} + 1 \right)^{5/4}}{5b^2} - \frac{4a^2 \sqrt[4]{\frac{bx^2}{a} + 1}}{b^2} \right)
 \end{aligned}$$

input `Int[x^3*(c/Sqrt[a + b*x^2])^(3/2),x]`

output `((c/Sqrt[a + b*x^2])^(3/2)*(1 + (b*x^2)/a)^(3/4)*((-4*a^2*(1 + (b*x^2)/a)^(1/4))/b^2 + (4*a^2*(1 + (b*x^2)/a)^(5/4))/(5*b^2)))/2`

Definitions of rubi rules used

rule 53 $\text{Int}[(a_.) + (b_.)(x_)^{(m_.)}*((c_.) + (d_.)(x_)^{(n_.)}), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \text{FreeQ}\{a, b, c, d, n\}, x] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ (\ !\text{IntegerQ}[n] \ || \ (\text{EqQ}[c, 0] \ \&\& \ \text{LeQ}[7*m + 4*n + 4, 0]) \ || \ \text{LtQ}[9*m + 5*(n + 1), 0] \ || \ \text{GtQ}[m + n + 2, 0])$

rule 243 $\text{Int}[(x_)^{(m_.)}*((a_) + (b_.)(x_)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[1/2 \ \text{Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] /; \text{FreeQ}\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 2045 $\text{Int}[(u_.)*((c_.)*((a_) + (b_.)(x_)^{(n_.)})^{(q_.)})^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[\text{Simp}[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^{(p*q)}] \ \text{Int}[u*(1 + b*(x^n/a))^{(p*q)}, x], x] /; \text{FreeQ}\{a, b, c, n, p, q\}, x] \ \&\& \ !\text{GeQ}[a, 0]$

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.56

method	result	size
gospers	$-\frac{2(bx^2+a)(-bx^2+4a)\left(\frac{c}{\sqrt{bx^2+a}}\right)^{3/2}}{5b^2}$	36
orering	$-\frac{2(bx^2+a)(-bx^2+4a)\left(\frac{c}{\sqrt{bx^2+a}}\right)^{3/2}}{5b^2}$	36

input $\text{int}(x^3*(c/(b*x^2+a)^{(1/2}))^{(3/2)}, x, \text{method}=_RETURNVERBOSE)$

output $-2/5*(b*x^2+a)*(-b*x^2+4*a)*(c/(b*x^2+a)^{(1/2}))^{(3/2)}/b^2$

Fricas [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.59

$$\int x^3 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \frac{2(bc x^2 - 4ac) \sqrt{bx^2+a} \sqrt{\frac{c}{\sqrt{bx^2+a}}}}{5b^2}$$

input `integrate(x^3*(c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="fricas")`output `2/5*(b*c*x^2 - 4*a*c)*sqrt(b*x^2 + a)*sqrt(c/sqrt(b*x^2 + a))/b^2`**Sympy [A] (verification not implemented)**

Time = 2.81 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.36

$$\int x^3 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \begin{cases} -\frac{8a^2 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2}}{5b^2} - \frac{6ax^2 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2}}{5b} + \frac{2x^4 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2}}{5} & \text{for } b \neq 0 \\ \frac{x^4 \left(\frac{c}{\sqrt{a}} \right)^{3/2}}{4} & \text{otherwise} \end{cases}$$

input `integrate(x**3*(c/(b*x**2+a)**(1/2))**(3/2),x)`output `Piecewise((-8*a**2*(c/sqrt(a + b*x**2))**(3/2)/(5*b**2) - 6*a*x**2*(c/sqrt(a + b*x**2))**(3/2)/(5*b) + 2*x**4*(c/sqrt(a + b*x**2))**(3/2)/5, Ne(b, 0)), (x**4*(c/sqrt(a))**(3/2)/4, True))`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.66

$$\int x^3 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = -\frac{2 \left(\frac{5ac^2}{bx^2+a} - c^2 \right) c^2}{5b^2 \left(\frac{c}{\sqrt{bx^2+a}} \right)^{5/2}}$$

input `integrate(x^3*(c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="maxima")`

output `-2/5*(5*a*c^2/(b*x^2 + a) - c^2)*c^2/(b^2*(c/sqrt(b*x^2 + a))^(5/2))`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.62

$$\int x^3 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \frac{2 \left((bx^2 + a)^{5/4} - 5 (bx^2 + a)^{1/4} a \right) c^{3/2}}{5 b^2 \sqrt{\operatorname{sgn}(bx^2 + a)}}$$

input `integrate(x^3*(c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="giac")`

output `2/5*((b*x^2 + a)^(5/4) - 5*(b*x^2 + a)^(1/4)*a)*c^(3/2)/(b^2*sqrt(sgn(b*x^2 + a)))`

Mupad [B] (verification not implemented)

Time = 0.54 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.59

$$\int x^3 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = -\frac{2 c \sqrt{bx^2 + a} \sqrt{\frac{c}{\sqrt{bx^2 + a}}} (4a - bx^2)}{5 b^2}$$

input `int(x^3*(c/(a + b*x^2)^(1/2))^(3/2),x)`

output `-(2*c*(a + b*x^2)^(1/2)*(c/(a + b*x^2)^(1/2))^(1/2)*(4*a - b*x^2))/(5*b^2)`

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.31

$$\int x^3 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \frac{2\sqrt{c} \sqrt{\sqrt{b}\sqrt{bx^2+a}x+a+bx^2} \sqrt{\sqrt{bx^2+a}+\sqrt{b}x} c (-4\sqrt{bx^2+a}a+\sqrt{bx^2+a}a+\sqrt{bx^2+a}a)}{5ab^2}$$

input `int(x^3*(c/(b*x^2+a)^(1/2))^(3/2),x)`output `(2*sqrt(c)*sqrt(sqrt(b)*sqrt(a + b*x**2)*x + a + b*x**2)*sqrt(sqrt(a + b*x**2) + sqrt(b)*x)*c*(- 4*sqrt(a + b*x**2)*a + sqrt(a + b*x**2)*b*x**2 + 4*sqrt(b)*a*x - sqrt(b)*b*x**3))/(5*a*b**2)`

$$3.52 \quad \int x \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx$$

Optimal result	379
Mathematica [A] (verified)	379
Rubi [A] (verified)	380
Maple [A] (verified)	381
Fricas [A] (verification not implemented)	381
Sympy [A] (verification not implemented)	382
Maxima [A] (verification not implemented)	382
Giac [A] (verification not implemented)	383
Mupad [B] (verification not implemented)	383
Reduce [B] (verification not implemented)	383

Optimal result

Integrand size = 19, antiderivative size = 34

$$\int x \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \frac{2c \sqrt{\frac{c}{\sqrt{a+bx^2}}} \sqrt{a+bx^2}}{b}$$

output `2*c*(c/(b*x^2+a)^(1/2))^(1/2)*(b*x^2+a)^(1/2)/b`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.74

$$\int x \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \frac{2c^2}{b \sqrt{\frac{c}{\sqrt{a+bx^2}}}}$$

input `Integrate[x*(c/Sqrt[a + b*x^2])^(3/2),x]`

output `(2*c^2)/(b*Sqrt[c/Sqrt[a + b*x^2]])`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.85, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2024, 20, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int x \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx \\
 \downarrow 2024 \\
 \frac{\int \left(\frac{c}{\sqrt{bx^2+a}} \right)^{3/2} d(bx^2+a)}{2b} \\
 \downarrow 20 \\
 \frac{(a+bx^2)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} \int \frac{1}{(bx^2+a)^{3/4}} d(bx^2+a)}{2b} \\
 \downarrow 15 \\
 \frac{2(a+bx^2) \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2}}{b}
 \end{array}$$

input `Int[x*(c/Sqrt[a + b*x^2])^(3/2),x]`

output `(2*(c/Sqrt[a + b*x^2])^(3/2)*(a + b*x^2))/b`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 20 `Int[((a_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a*x^n)^p/x^(n*p) Int[x^(n*p), x], x] /; FreeQ[{a, n, p}, x] && !IntegerQ[p]`

rule 2024

```
Int[((a_.) + (b_.)*(Pq_)^(n_.))^(p_.)*(Qr_), x_Symbol] := With[{q = Expon[P
q, x], r = Expon[Qr, x]}, Simp[Coeff[Qr, x, r]/(q*Coeff[Pq, x, q]) Subst[
Int[(a + b*x^n)^p, x], x, Pq], x] /; EqQ[r, q - 1] && EqQ[Coeff[Qr, x, r]*D
[Pq, x], q*Coeff[Pq, x, q]*Qr]] /; FreeQ[{a, b, n, p}, x] && PolyQ[Pq, x] &
& PolyQ[Qr, x]
```

Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.76

method	result	size
gospers	$\frac{2(bx^2+a)\left(\frac{c}{\sqrt{bx^2+a}}\right)^{3/2}}{b}$	26
derivativedivides	$\frac{2(bx^2+a)\left(\frac{c}{\sqrt{bx^2+a}}\right)^{3/2}}{b}$	26
default	$\frac{2(bx^2+a)\left(\frac{c}{\sqrt{bx^2+a}}\right)^{3/2}}{b}$	26
orering	$\frac{2(bx^2+a)\left(\frac{c}{\sqrt{bx^2+a}}\right)^{3/2}}{b}$	26

input

```
int(x*(c/(b*x^2+a)^(1/2))^(3/2),x,method=_RETURNVERBOSE)
```

output

```
2*(b*x^2+a)*(c/(b*x^2+a)^(1/2))^(3/2)/b
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.82

$$\int x \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \frac{2\sqrt{bx^2+ac}\sqrt{\frac{c}{\sqrt{bx^2+a}}}}{b}$$

input

```
integrate(x*(c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="fricas")
```

output `2*sqrt(b*x^2 + a)*c*sqrt(c/sqrt(b*x^2 + a))/b`

Sympy [A] (verification not implemented)

Time = 1.47 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.59

$$\int x \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \begin{cases} \frac{2a \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2}}{b} + 2x^2 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} & \text{for } b \neq 0 \\ \frac{x^2 \left(\frac{c}{\sqrt{a}} \right)^{3/2}}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*(c/(b*x**2+a)**(1/2))**(3/2),x)`

output `Piecewise((2*a*(c/sqrt(a + b*x**2))**(3/2)/b + 2*x**2*(c/sqrt(a + b*x**2))**(3/2), Ne(b, 0)), (x**2*(c/sqrt(a))**(3/2)/2, True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.74

$$\int x \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \frac{2(bx^2 + a) \left(\frac{c}{\sqrt{bx^2+a}} \right)^{3/2}}{b}$$

input `integrate(x*(c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="maxima")`

output `2*(b*x^2 + a)*(c/sqrt(b*x^2 + a))^(3/2)/b`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.79

$$\int x \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \frac{2 (bx^2 + a)^{1/4} c^{3/2}}{b \sqrt{\operatorname{sgn}(bx^2 + a)}}$$

input `integrate(x*(c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="giac")`output `2*(b*x^2 + a)^(1/4)*c^(3/2)/(b*sqrt(sgn(b*x^2 + a)))`**Mupad [B] (verification not implemented)**

Time = 0.52 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.82

$$\int x \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \frac{2c \sqrt{bx^2 + a} \sqrt{\frac{c}{\sqrt{bx^2 + a}}}}{b}$$

input `int(x*(c/(a + b*x^2)^(1/2))^(3/2),x)`output `(2*c*(a + b*x^2)^(1/2)*(c/(a + b*x^2)^(1/2))^(1/2))/b`**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.74

$$\int x \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \frac{2\sqrt{c} \sqrt{\sqrt{b} \sqrt{bx^2 + a} x + a + bx^2} \sqrt{\sqrt{bx^2 + a} + \sqrt{b} x} c (\sqrt{bx^2 + a} - \sqrt{b} x)}{ab}$$

input `int(x*(c/(b*x^2+a)^(1/2))^(3/2),x)`output `(2*sqrt(c)*sqrt(sqrt(b)*sqrt(a + b*x**2)*x + a + b*x**2)*sqrt(sqrt(a + b*x**2) + sqrt(b)*x)*c*(sqrt(a + b*x**2) - sqrt(b)*x)/(a*b)`

3.53
$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x} dx$$

Optimal result	384
Mathematica [A] (verified)	384
Rubi [A] (verified)	385
Maple [F]	387
Fricas [F(-1)]	387
Sympy [F]	388
Maxima [A] (verification not implemented)	388
Giac [B] (verification not implemented)	388
Mupad [F(-1)]	389
Reduce [B] (verification not implemented)	390

Optimal result

Integrand size = 21, antiderivative size = 97

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x} dx = -\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} \left(1+\frac{bx^2}{a}\right)^{3/4} \arctan\left(\sqrt[4]{1+\frac{bx^2}{a}}\right) - \left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} \left(1+\frac{bx^2}{a}\right)^{3/4} \operatorname{arctanh}\left(\sqrt[4]{1+\frac{bx^2}{a}}\right)$$

```
output - (c/(b*x^2+a)^(1/2))^(3/2)*(1+b*x^2/a)^(3/4)*arctan((1+b*x^2/a)^(1/4)) - (c/(b*x^2+a)^(1/2))^(3/2)*(1+b*x^2/a)^(3/4)*arctanh((1+b*x^2/a)^(1/4))
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.74

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x} dx = \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} (a+bx^2)^{3/4} \left(\arctan\left(\frac{\sqrt[4]{a+bx^2}}{\sqrt[4]{a}}\right) + \operatorname{arctanh}\left(\frac{\sqrt[4]{a+bx^2}}{\sqrt[4]{a}}\right)\right)}{a^{3/4}}$$

input `Integrate[(c/Sqrt[a + b*x^2])^(3/2)/x,x]`

output `-(((c/Sqrt[a + b*x^2])^(3/2)*(a + b*x^2)^(3/4)*(ArcTan[(a + b*x^2)^(1/4)/a^(1/4)] + ArcTanh[(a + b*x^2)^(1/4)/a^(1/4)]))/a^(3/4))`

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.87, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2045, 243, 73, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x} dx \\
 & \quad \downarrow \text{2045} \\
 & \left(\frac{bx^2}{a} + 1\right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} \int \frac{1}{x \left(\frac{bx^2}{a} + 1\right)^{3/4}} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{2} \left(\frac{bx^2}{a} + 1\right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} \int \frac{1}{x^2 \left(\frac{bx^2}{a} + 1\right)^{3/4}} dx^2 \\
 & \quad \downarrow \text{73} \\
 & \frac{2a \left(\frac{bx^2}{a} + 1\right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} \int \frac{1}{\frac{ax^8}{b} - \frac{a}{b}} d^4 \sqrt{\frac{bx^2}{a} + 1}}{b} \\
 & \quad \downarrow \text{756} \\
 & \frac{2a \left(\frac{bx^2}{a} + 1\right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} \left(-\frac{b \int \frac{1}{1-x^4} d^4 \sqrt{\frac{bx^2}{a} + 1}}{2a} - \frac{b \int \frac{1}{x^4+1} d^4 \sqrt{\frac{bx^2}{a} + 1}}{2a} \right)}{b}
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow 216 \\
 \frac{2a \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} \left(-\frac{b \int \frac{1}{1-x^4} dx \sqrt[4]{\frac{bx^2}{a} + 1}}{2a} - \frac{b \arctan \left(\sqrt[4]{\frac{bx^2}{a} + 1} \right)}{2a} \right)}{b} \\
 \downarrow 219 \\
 \frac{2a \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} \left(-\frac{b \arctan \left(\sqrt[4]{\frac{bx^2}{a} + 1} \right)}{2a} - \frac{b \operatorname{arctanh} \left(\sqrt[4]{\frac{bx^2}{a} + 1} \right)}{2a} \right)}{b}
 \end{array}$$

input `Int[(c/Sqrt[a + b*x^2])^(3/2)/x,x]`

output `(2*a*(c/Sqrt[a + b*x^2])^(3/2)*(1 + (b*x^2)/a)^(3/4)*(-1/2*(b*ArcTan[(1 + (b*x^2)/a)^(1/4)])/a - (b*ArcTanh[(1 + (b*x^2)/a)^(1/4)]/(2*a)))/b`

Defintions of rubi rules used

rule 73 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 756 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^q)^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q) Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [F]

$$\int \frac{\left(\frac{c}{\sqrt{bx^2+a}}\right)^{\frac{3}{2}}}{x} dx$$

input `int((c/(b*x^2+a)^(1/2))^(3/2)/x,x)`

output `int((c/(b*x^2+a)^(1/2))^(3/2)/x,x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x} dx = \text{Timed out}$$

input `integrate((c/(b*x^2+a)^(1/2))^(3/2)/x,x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x} dx = \int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x} dx$$

input `integrate((c/(b*x**2+a)**(1/2))**(3/2)/x,x)`

output `Integral((c/sqrt(a + b*x**2))**(3/2)/x, x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.13

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x} dx = \frac{1}{2} c^2 \left(\frac{2 \arctan\left(\frac{\sqrt{a}\sqrt{\frac{c}{\sqrt{bx^2+a}}}}{\sqrt{\sqrt{ac}}}\right)}{\sqrt{\sqrt{ac}}\sqrt{a}} + \frac{\log\left(\frac{\sqrt{a}\sqrt{\frac{c}{\sqrt{bx^2+a}}} - \sqrt{\sqrt{ac}}}{\sqrt{a}\sqrt{\frac{c}{\sqrt{bx^2+a}}} + \sqrt{\sqrt{ac}}}\right)}{\sqrt{\sqrt{ac}}\sqrt{a}} \right)$$

input `integrate((c/(b*x^2+a)^(1/2))^(3/2)/x,x, algorithm="maxima")`

output `1/2*c^2*(2*arctan(sqrt(a)*sqrt(c/sqrt(b*x^2 + a))/sqrt(sqrt(a)*c))/(sqrt(sqrt(a)*c)*sqrt(a)) + log((sqrt(a)*sqrt(c/sqrt(b*x^2 + a)) - sqrt(sqrt(a)*c))/(sqrt(a)*sqrt(c/sqrt(b*x^2 + a)) + sqrt(sqrt(a)*c)))/(sqrt(sqrt(a)*c)*sqrt(a))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 200 vs. 2(81) = 162.

Time = 0.13 (sec) , antiderivative size = 200, normalized size of antiderivative = 2.06

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x} dx =$$

$$c^{\frac{3}{2}} \left(\frac{2\sqrt{2}(-a)^{\frac{1}{4}} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}(-a)^{\frac{1}{4}}+2(bx^2+a)^{\frac{1}{4}}\right)}{2(-a)^{\frac{1}{4}}}\right)}{a} + \frac{2\sqrt{2}(-a)^{\frac{1}{4}} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}(-a)^{\frac{1}{4}}-2(bx^2+a)^{\frac{1}{4}}\right)}{2(-a)^{\frac{1}{4}}}\right)}{a} + \frac{\sqrt{2}(-a)^{\frac{1}{4}} \log\left(\sqrt{2}(bx^2+a)\right)}{4\sqrt{\operatorname{sgn}(bx^2+a)}}$$

input `integrate((c/(b*x^2+a)^(1/2))^(3/2)/x,x, algorithm="giac")`

output

```
-1/4*c^(3/2)*(2*sqrt(2)*(-a)^(1/4)*arctan(1/2*sqrt(2)*(sqrt(2)*(-a)^(1/4)
+ 2*(b*x^2 + a)^(1/4))/(-a)^(1/4))/a + 2*sqrt(2)*(-a)^(1/4)*arctan(-1/2*sq
rt(2)*(sqrt(2)*(-a)^(1/4) - 2*(b*x^2 + a)^(1/4))/(-a)^(1/4))/a + sqrt(2)*(-
-a)^(1/4)*log(sqrt(2)*(b*x^2 + a)^(1/4)*(-a)^(1/4) + sqrt(b*x^2 + a) + sqr
t(-a))/a - sqrt(2)*(-a)^(1/4)*log(-sqrt(2)*(b*x^2 + a)^(1/4)*(-a)^(1/4) +
sqrt(b*x^2 + a) + sqrt(-a))/a)/sqrt(sgn(b*x^2 + a))
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x} dx = \int \frac{\left(\frac{c}{\sqrt{bx^2+a}}\right)^{3/2}}{x} dx$$

input `int((c/(a + b*x^2)^(1/2))^(3/2)/x,x)`

output

```
int((c/(a + b*x^2)^(1/2))^(3/2)/x, x)
```

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.07

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x} dx = \frac{\sqrt{c} c \left(\operatorname{atan}\left(\frac{a^{5/4}(bx^2+a)^{3/4} - a^{7/4}(bx^2+a)^{1/4} - a^{3/4}(bx^2+a)^{1/4}bx^2}{2abx^2+2a^2}\right) - \log\left((bx^2+a)^{1/4} + a^{1/4}\right) + \log\left(\left(\frac{a+bx^2}{a}\right)^{1/4} - a^{1/4}\right) \right)}{2a^{3/4}}$$

input `int((c/(b*x^2+a)^(1/2))^(3/2)/x,x)`

output

```
(sqrt(c)*a**(1/4)*c*(atan((a**(1/4)*(a + b*x**2)**(3/4)*a - a**(3/4)*(a +
b*x**2)**(1/4)*a - a**(3/4)*(a + b*x**2)**(1/4)*b*x**2)/(2*a**2 + 2*a*b*x*
*2)) - log((a + b*x**2)**(1/4) + a**(1/4)) + log((a + b*x**2)**(1/4) - a**
(1/4))))/(2*a)
```

3.54 $\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^3} dx$

Optimal result	391
Mathematica [A] (verified)	392
Rubi [A] (verified)	392
Maple [F]	395
Fricas [F(-1)]	395
Sympy [F]	395
Maxima [A] (verification not implemented)	396
Giac [A] (verification not implemented)	396
Mupad [F(-1)]	397
Reduce [B] (verification not implemented)	397

Optimal result

Integrand size = 21, antiderivative size = 143

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^3} dx = -\frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} (a + bx^2)}{2ax^2} + \frac{3b\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} \left(1 + \frac{bx^2}{a}\right)^{3/4} \arctan\left(\sqrt[4]{1 + \frac{bx^2}{a}}\right)}{4a} + \frac{3b\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} \left(1 + \frac{bx^2}{a}\right)^{3/4} \operatorname{arctanh}\left(\sqrt[4]{1 + \frac{bx^2}{a}}\right)}{4a}$$

output

```
-1/2*(c/(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)/a/x^2+3/4*b*(c/(b*x^2+a)^(1/2))^(3/2)*(1+b*x^2/a)^(3/4)*arctan((1+b*x^2/a)^(1/4))/a+3/4*b*(c/(b*x^2+a)^(1/2))^(3/2)*(1+b*x^2/a)^(3/4)*arctanh((1+b*x^2/a)^(1/4))/a
```


Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.75

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^3} dx = \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} (a+bx^2)^{3/4} \left(-2a^{3/4}\sqrt[4]{a+bx^2} + 3bx^2 \arctan\left(\frac{\sqrt[4]{a+bx^2}}{\sqrt[4]{a}}\right) + 3bx^2 \arctan\right)}{4a^{7/4}x^2}$$

input `Integrate[(c/Sqrt[a + b*x^2])^(3/2)/x^3,x]`

output `((c/Sqrt[a + b*x^2])^(3/2)*(a + b*x^2)^(3/4)*(-2*a^(3/4)*(a + b*x^2)^(1/4) + 3*b*x^2*ArcTan[(a + b*x^2)^(1/4)/a^(1/4)] + 3*b*x^2*ArcTanh[(a + b*x^2)^(1/4)/a^(1/4)]))/(4*a^(7/4)*x^2)`

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.73, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2045, 243, 52, 73, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^3} dx \\ & \quad \downarrow \text{2045} \\ & \left(\frac{bx^2}{a} + 1\right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} \int \frac{1}{x^3 \left(\frac{bx^2}{a} + 1\right)^{3/4}} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{2} \left(\frac{bx^2}{a} + 1\right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} \int \frac{1}{x^4 \left(\frac{bx^2}{a} + 1\right)^{3/4}} dx^2 \\ & \quad \downarrow \text{52} \end{aligned}$$

$$\frac{1}{2} \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} \left(-\frac{3b \int \frac{1}{x^2 \left(\frac{bx^2}{a} + 1 \right)^{3/4}} dx^2}{4a} - \frac{\sqrt[4]{\frac{bx^2}{a} + 1}}{x^2} \right)$$

↓ 73

$$\frac{1}{2} \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} \left(-3 \int \frac{1}{\frac{ax^8}{b} - \frac{a}{b}} d\sqrt[4]{\frac{bx^2}{a} + 1} - \frac{\sqrt[4]{\frac{bx^2}{a} + 1}}{x^2} \right)$$

↓ 756

$$\frac{1}{2} \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} \left(-3 \left(-\frac{b \int \frac{1}{1-x^4} d\sqrt[4]{\frac{bx^2}{a} + 1}}{2a} - \frac{b \int \frac{1}{x^4+1} d\sqrt[4]{\frac{bx^2}{a} + 1}}{2a} \right) - \frac{\sqrt[4]{\frac{bx^2}{a} + 1}}{x^2} \right)$$

↓ 216

$$\frac{1}{2} \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} \left(-3 \left(-\frac{b \int \frac{1}{1-x^4} d\sqrt[4]{\frac{bx^2}{a} + 1}}{2a} - \frac{b \arctan \left(\sqrt[4]{\frac{bx^2}{a} + 1} \right)}{2a} \right) - \frac{\sqrt[4]{\frac{bx^2}{a} + 1}}{x^2} \right)$$

↓ 219

$$\frac{1}{2} \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} \left(-3 \left(-\frac{b \arctan \left(\sqrt[4]{\frac{bx^2}{a} + 1} \right)}{2a} - \frac{b \operatorname{arctanh} \left(\sqrt[4]{\frac{bx^2}{a} + 1} \right)}{2a} \right) - \frac{\sqrt[4]{\frac{bx^2}{a} + 1}}{x^2} \right)$$

input `Int[(c/Sqrt[a + b*x^2])^(3/2)/x^3,x]`

output `((c/Sqrt[a + b*x^2])^(3/2)*(1 + (b*x^2)/a)^(3/4)*(-(1 + (b*x^2)/a)^(1/4)/x^2 - 3*(-1/2*(b*ArcTan[(1 + (b*x^2)/a)^(1/4)]/a - (b*ArcTanh[(1 + (b*x^2)/a)^(1/4)]/(2*a)))))/2`

Definitions of rubi rules used

- rule 52 $\text{Int}[(a_.) + (b_.)(x_)^{(m_)}((c_.) + (d_.)(x_)^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}((c + d*x)^{(n + 1)} / ((b*c - a*d)*(m + 1))), x] - \text{Simp}[d*((m + n + 2) / ((b*c - a*d)*(m + 1))) \text{Int}[(a + b*x)^{(m + 1)}(c + d*x)^n, x], x] /; \text{FreeQ}\{a, b, c, d, n\}, x \ \&\& \ \text{ILtQ}[m, -1] \ \&\& \ \text{FractionQ}[n] \ \&\& \ \text{LtQ}[n, 0]$
- rule 73 $\text{Int}[(a_.) + (b_.)(x_)^{(m_)}((c_.) + (d_.)(x_)^{(n_)}, x_Symbol] \rightarrow \text{With}\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \ \text{Subst}[\text{Int}[x^{(p*(m + 1) - 1)}(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^{(1/p)}, x]] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$
- rule 216 $\text{Int}[(a_) + (b_.)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))* \text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$
- rule 219 $\text{Int}[(a_) + (b_.)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))* \text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 243 $\text{Int}[(x_)^{(m_.)}((a_) + (b_.)(x_)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[1/2 \ \text{Subst}[\text{Int}[x^{((m - 1)/2)*(a + b*x)^p}, x], x, x^2], x] /; \text{FreeQ}\{a, b, m, p\}, x \ \&\& \ \text{IntegerQ}[(m - 1)/2]$
- rule 756 $\text{Int}[(a_) + (b_.)(x_)^4)^{-1}, x_Symbol] \rightarrow \text{With}\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[r/(2*a) \ \text{Int}[1/(r - s*x^2), x], x] + \text{Simp}[r/(2*a) \ \text{Int}[1/(r + s*x^2), x], x]] /; \text{FreeQ}\{a, b\}, x \ \&\& \ !\text{GtQ}[a/b, 0]$
- rule 2045 $\text{Int}[(u_.)*((c_.)*((a_) + (b_.)(x_)^{(n_.)})^{(q_.)})^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[\text{Simp}[(c*(a + b*x^n)^q)^p / (1 + b*(x^n/a))^{(p*q)} \ \text{Int}[u*(1 + b*(x^n/a))^{(p*q)}, x], x] /; \text{FreeQ}\{a, b, c, n, p, q\}, x \ \&\& \ !\text{GeQ}[a, 0]$

Maple [F]

$$\int \frac{\left(\frac{c}{\sqrt{bx^2+a}}\right)^{\frac{3}{2}}}{x^3} dx$$

input `int((c/(b*x^2+a)^(1/2))^(3/2)/x^3,x)`

output `int((c/(b*x^2+a)^(1/2))^(3/2)/x^3,x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^3} dx = \text{Timed out}$$

input `integrate((c/(b*x^2+a)^(1/2))^(3/2)/x^3,x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^3} dx = \int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{\frac{3}{2}}}{x^3} dx$$

input `integrate((c/(b*x**2+a)**(1/2))**(3/2)/x**3,x)`

output `Integral((c/sqrt(a + b*x**2))**(3/2)/x**3, x)`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.10

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^3} dx = -\frac{1}{8}bc^2 \left(\frac{4\left(\frac{c}{\sqrt{bx^2+a}}\right)^{3/2}}{ac^2 - \frac{a^2c^2}{bx^2+a}} + \frac{3 \left(\frac{2 \arctan\left(\frac{\sqrt{a}\sqrt{\frac{c}{bx^2+a}}}{\sqrt{ac}}\right)}{\sqrt{ac}\sqrt{a}} + \frac{\log\left(\frac{\sqrt{a}\sqrt{\frac{c}{bx^2+a}} - \sqrt{ac}}{\sqrt{a}\sqrt{\frac{c}{bx^2+a}} + \sqrt{ac}}\right)}{\sqrt{ac}\sqrt{a}} \right)}{a} \right)$$

input `integrate((c/(b*x^2+a)^(1/2))^(3/2)/x^3,x, algorithm="maxima")`output `-1/8*b*c^2*(4*(c/sqrt(b*x^2 + a))^(3/2)/(a*c^2 - a^2*c^2/(b*x^2 + a)) + 3*(2*arctan(sqrt(a)*sqrt(c/sqrt(b*x^2 + a))/sqrt(sqrt(a)*c))/(sqrt(sqrt(a)*c)*sqrt(a)) + log((sqrt(a)*sqrt(c/sqrt(b*x^2 + a)) - sqrt(sqrt(a)*c))/(sqrt(a)*sqrt(c/sqrt(b*x^2 + a)) + sqrt(sqrt(a)*c)))/(sqrt(sqrt(a)*c)*sqrt(a)))/a)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.55

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^3} dx = bc^{\frac{3}{2}} \left(\frac{6\sqrt{2}(-a)^{\frac{1}{4}} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}(-a)^{\frac{1}{4}}+2\left(bx^2+a\right)^{\frac{1}{4}}\right)}{2(-a)^{\frac{1}{4}}}\right)}{a^2} + \frac{6\sqrt{2}(-a)^{\frac{1}{4}} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}(-a)^{\frac{1}{4}}-2\left(bx^2+a\right)^{\frac{1}{4}}\right)}{2(-a)^{\frac{1}{4}}}\right)}{a^2} \right)$$

input `integrate((c/(b*x^2+a)^(1/2))^(3/2)/x^3,x, algorithm="giac")`

output

```
1/16*b*c^(3/2)*(6*sqrt(2)*(-a)^(1/4)*arctan(1/2*sqrt(2)*(sqrt(2)*(-a)^(1/4)
)+ 2*(b*x^2 + a)^(1/4))/(-a)^(1/4))/a^2 + 6*sqrt(2)*(-a)^(1/4)*arctan(-1/
2*sqrt(2)*(sqrt(2)*(-a)^(1/4) - 2*(b*x^2 + a)^(1/4))/(-a)^(1/4))/a^2 + 3*s
qrt(2)*(-a)^(1/4)*log(sqrt(2)*(b*x^2 + a)^(1/4)*(-a)^(1/4) + sqrt(b*x^2 +
a) + sqrt(-a))/a^2 - 3*sqrt(2)*(-a)^(1/4)*log(-sqrt(2)*(b*x^2 + a)^(1/4)*(-
a)^(1/4) + sqrt(b*x^2 + a) + sqrt(-a))/a^2 - 8*(b*x^2 + a)^(1/4)/(a*b*x^2
))/sqrt(sgn(b*x^2 + a))
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^3} dx = \int \frac{\left(\frac{c}{\sqrt{bx^2+a}}\right)^{3/2}}{x^3} dx$$

input

```
int((c/(a + b*x^2)^(1/2))^(3/2)/x^3,x)
```

output

```
int((c/(a + b*x^2)^(1/2))^(3/2)/x^3, x)
```

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.01

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^3} dx = \frac{\sqrt{c} c \left(-3a^{1/4} \operatorname{atan} \left(\frac{a^{5/4} (bx^2+a)^{3/4} - a^{7/4} (bx^2+a)^{1/4} - a^{3/4} (bx^2+a)^{1/4} bx^2}{2abx^2+2a^2} \right) bx^2 - 4(bx^2 + a)^{1/4} a + 3a^{1/4}}{8a^2x^2} \right)}{8a^2x^2}$$

input

```
int((c/(b*x^2+a)^(1/2))^(3/2)/x^3,x)
```

output

```
(sqrt(c)*c*( - 3*a**(1/4)*atan((a**(1/4)*(a + b*x**2)**(3/4)*a - a**(3/4)*
(a + b*x**2)**(1/4)*a - a**(3/4)*(a + b*x**2)**(1/4)*b*x**2)/(2*a**2 + 2*a
*b*x**2))*b*x**2 - 4*(a + b*x**2)**(1/4)*a + 3*a**(1/4)*log((a + b*x**2)**
(1/4) + a**(1/4))*b*x**2 - 3*a**(1/4)*log((a + b*x**2)**(1/4) - a**(1/4))*
b*x**2))/(8*a**2*x**2)
```

3.55 $\int x^4 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx$

Optimal result	398
Mathematica [C] (verified)	398
Rubi [A] (verified)	399
Maple [F]	401
Fricas [F]	401
Sympy [F]	401
Maxima [F]	402
Giac [F]	402
Mupad [F(-1)]	402
Reduce [F]	403

Optimal result

Integrand size = 21, antiderivative size = 132

$$\int x^4 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = -\frac{4ax \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} (a+bx^2)}{7b^2} + \frac{2x^3 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} (a+bx^2)}{7b} + \frac{8a^{5/2} \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} \left(1 + \frac{bx^2}{a} \right)^{3/4} \text{EllipticF} \left(\frac{1}{2} \arctan \left(\frac{\sqrt{bx}}{\sqrt{a}} \right), 2 \right)}{7b^{5/2}}$$

output `-4/7*a*x*(c/(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)/b^2+2/7*x^3*(c/(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)/b+8/7*a^(5/2)*(c/(b*x^2+a)^(1/2))^(3/2)*(1+b*x^2/a)^(3/4)*InverseJacobiAM(1/2*arctan(b^(1/2)*x/a^(1/2)),2^(1/2))/b^(5/2)`

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.37 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.63

$$\int x^4 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \frac{2x \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} \left(-2a^2 - abx^2 + b^2x^4 + 2a^2 \left(1 + \frac{bx^2}{a} \right)^{3/4} \text{Hypergeometric2F1} \left(\right)}{7b^2}$$

input `Integrate[x^4*(c/Sqrt[a + b*x^2])^(3/2),x]`

output $(2*x*(c/Sqrt[a + b*x^2])^(3/2)*(-2*a^2 - a*b*x^2 + b^2*x^4 + 2*a^2*(1 + (b*x^2)/a)^(3/4)*Hypergeometric2F1[1/2, 3/4, 3/2, -((b*x^2)/a)]))/(7*b^2)$

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.93, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2045, 262, 262, 229}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^4 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx \\
 & \quad \downarrow \text{2045} \\
 & \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} \int \frac{x^4}{\left(\frac{bx^2}{a} + 1 \right)^{3/4}} dx \\
 & \quad \downarrow \text{262} \\
 & \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} \left(\frac{2ax^3 \sqrt{\frac{bx^2}{a} + 1}}{7b} - \frac{6a \int \frac{x^2}{\left(\frac{bx^2}{a} + 1 \right)^{3/4}} dx}{7b} \right) \\
 & \quad \downarrow \text{262} \\
 & \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} \left(\frac{2ax^3 \sqrt{\frac{bx^2}{a} + 1}}{7b} - \frac{6a \left(\frac{2ax^4 \sqrt{\frac{bx^2}{a} + 1}}{3b} - \frac{2a \int \frac{1}{\left(\frac{bx^2}{a} + 1 \right)^{3/4}} dx}{3b} \right)}{7b} \right) \\
 & \quad \downarrow \text{229}
 \end{aligned}$$

$$\left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} \left(\frac{2ax^3 \sqrt[4]{\frac{bx^2}{a} + 1}}{7b} - \frac{6a \left(\frac{2ax \sqrt[4]{\frac{bx^2}{a} + 1}}{3b} - \frac{4a^{3/2} \operatorname{EllipticF}\left(\frac{1}{2} \arctan\left(\frac{\sqrt{bx^2}}{\sqrt{a}}\right), 2\right)}{3b^{3/2}} \right)}{7b} \right)$$

input `Int[x^4*(c/Sqrt[a + b*x^2])^(3/2),x]`

output `(c/Sqrt[a + b*x^2])^(3/2)*(1 + (b*x^2)/a)^(3/4)*((2*a*x^3*(1 + (b*x^2)/a)^(1/4))/(7*b) - (6*a*((2*a*x*(1 + (b*x^2)/a)^(1/4))/(3*b) - (4*a^(3/2)*EllipticF[ArcTan[(Sqrt[b]*x)/Sqrt[a]]/2, 2)]/(3*b^(3/2))))/(7*b)`

Defintions of rubi rules used

rule 229 `Int[((a_) + (b_)*(x_)^2)^(-3/4), x_Symbol] := Simp[(2/(a^(3/4)*Rt[b/a, 2]) * EllipticF[(1/2)*ArcTan[Rt[b/a, 2]*x], 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]`

rule 262 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*(m - 1)/(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2045 `Int[(u_)*((c_)*((a_) + (b_)*(x_)^(n_))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [F]

$$\int x^4 \left(\frac{c}{\sqrt{bx^2 + a}} \right)^{\frac{3}{2}} dx$$

input `int(x^4*(c/(b*x^2+a)^(1/2))^(3/2),x)`

output `int(x^4*(c/(b*x^2+a)^(1/2))^(3/2),x)`

Fricas [F]

$$\int x^4 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \int x^4 \left(\frac{c}{\sqrt{bx^2 + a}} \right)^{\frac{3}{2}} dx$$

input `integrate(x^4*(c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="fricas")`

output `integral(c*x^4*sqrt(c/sqrt(b*x^2 + a))/sqrt(b*x^2 + a), x)`

Sympy [F]

$$\int x^4 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \int x^4 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{\frac{3}{2}} dx$$

input `integrate(x**4*(c/(b*x**2+a)**(1/2))**(3/2),x)`

output `Integral(x**4*(c/sqrt(a + b*x**2))**(3/2), x)`

Maxima [F]

$$\int x^4 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \int x^4 \left(\frac{c}{\sqrt{bx^2 + a}} \right)^{3/2} dx$$

input `integrate(x^4*(c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="maxima")`

output `integrate(x^4*(c/sqrt(b*x^2 + a))^(3/2), x)`

Giac [F]

$$\int x^4 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \int x^4 \left(\frac{c}{\sqrt{bx^2 + a}} \right)^{3/2} dx$$

input `integrate(x^4*(c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="giac")`

output `integrate(x^4*(c/sqrt(b*x^2 + a))^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int x^4 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \int x^4 \left(\frac{c}{\sqrt{bx^2 + a}} \right)^{3/2} dx$$

input `int(x^4*(c/(a + b*x^2)^(1/2))^(3/2),x)`

output `int(x^4*(c/(a + b*x^2)^(1/2))^(3/2), x)`

Reduce [F]

$$\int x^4 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \frac{2\sqrt{c}c \left(-2(bx^2+a)^{\frac{1}{4}}ax + (bx^2+a)^{\frac{1}{4}}bx^3 + 2 \left(\int \frac{1}{(bx^2+a)^{\frac{3}{4}}} dx \right) a^2 \right)}{7b^2}$$

input `int(x^4*(c/(b*x^2+a)^(1/2))^(3/2),x)`

output `(2*sqrt(c)*c*(- 2*(a + b*x**2)**(1/4)*a*x + (a + b*x**2)**(1/4)*b*x**3 + 2*int((a + b*x**2)**(1/4)/(a + b*x**2),x)*a**2))/(7*b**2)`

3.56 $\int x^2 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx$

Optimal result	404
Mathematica [C] (verified)	404
Rubi [A] (verified)	405
Maple [F]	406
Fricas [F]	407
Sympy [F]	407
Maxima [F]	407
Giac [F]	408
Mupad [F(-1)]	408
Reduce [F]	408

Optimal result

Integrand size = 21, antiderivative size = 97

$$\int x^2 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \frac{2x \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} (a+bx^2)}{3b} - \frac{4a^{3/2} \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} \left(1 + \frac{bx^2}{a} \right)^{3/4} \text{EllipticF} \left(\frac{1}{2} \arctan \left(\frac{\sqrt{bx}}{\sqrt{a}} \right), 2 \right)}{3b^{3/2}}$$

output `2/3*x*(c/(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)/b-4/3*a^(3/2)*(c/(b*x^2+a)^(1/2))^(3/2)*(1+b*x^2/a)^(3/4)*InverseJacobiAM(1/2*arctan(b^(1/2)*x/a^(1/2)),2^(1/2))/b^(3/2)`

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 5.91 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.70

$$\int x^2 \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \frac{2x \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} \left(a+bx^2 - a \left(1 + \frac{bx^2}{a} \right)^{3/4} \text{Hypergeometric2F1} \left(\frac{1}{2}, \frac{3}{4}, \frac{3}{2}, -\frac{bx^2}{a} \right) \right)}{3b}$$

input `Integrate[x^2*(c/Sqrt[a + b*x^2])^(3/2),x]`

output `(2*x*(c/Sqrt[a + b*x^2])^(3/2)*(a + b*x^2 - a*(1 + (b*x^2)/a)^(3/4)*Hypergeometric2F1[1/2, 3/4, 3/2, -((b*x^2)/a)])/(3*b)`

Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.92, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2045, 262, 229}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx \\
 & \quad \downarrow \text{2045} \\
 & \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} \int \frac{x^2}{\left(\frac{bx^2}{a} + 1 \right)^{3/4}} dx \\
 & \quad \downarrow \text{262} \\
 & \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} \left(\frac{2ax \sqrt{\frac{bx^2}{a} + 1}}{3b} - \frac{2a \int \frac{1}{\left(\frac{bx^2}{a} + 1 \right)^{3/4}} dx}{3b} \right) \\
 & \quad \downarrow \text{229} \\
 & \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} \left(\frac{2ax \sqrt{\frac{bx^2}{a} + 1}}{3b} - \frac{4a^{3/2} \text{EllipticF} \left(\frac{1}{2} \arctan \left(\frac{\sqrt{bx}}{\sqrt{a}} \right), 2 \right)}{3b^{3/2}} \right)
 \end{aligned}$$

input `Int[x^2*(c/Sqrt[a + b*x^2])^(3/2),x]`

output

```
(c/Sqrt[a + b*x^2])^(3/2)*(1 + (b*x^2)/a)^(3/4)*((2*a*x*(1 + (b*x^2)/a)^(1/4))/(3*b) - (4*a^(3/2)*EllipticF[ArcTan[(Sqrt[b]*x)/Sqrt[a]]/2, 2])/(3*b^(3/2)))
```

Defintions of rubi rules used

rule 229

```
Int[((a_) + (b_.)*(x_)^2)^(-3/4), x_Symbol] := Simp[(2/(a^(3/4)*Rt[b/a, 2]) * EllipticF[(1/2)*ArcTan[Rt[b/a, 2]*x], 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]
```

rule 262

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]
```

rule 2045

```
Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]
```

Maple [F]

$$\int x^2 \left(\frac{c}{\sqrt{bx^2 + a}} \right)^{\frac{3}{2}} dx$$

input

```
int(x^2*(c/(b*x^2+a)^(1/2))^(3/2),x)
```

output

```
int(x^2*(c/(b*x^2+a)^(1/2))^(3/2),x)
```

Fricas [F]

$$\int x^2 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \int x^2 \left(\frac{c}{\sqrt{bx^2 + a}} \right)^{\frac{3}{2}} dx$$

input `integrate(x^2*(c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="fricas")`

output `integral(c*x^2*sqrt(c/sqrt(b*x^2 + a))/sqrt(b*x^2 + a), x)`

Sympy [F]

$$\int x^2 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \int x^2 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{\frac{3}{2}} dx$$

input `integrate(x**2*(c/(b*x**2+a)**(1/2))**(3/2),x)`

output `Integral(x**2*(c/sqrt(a + b*x**2))**(3/2), x)`

Maxima [F]

$$\int x^2 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \int x^2 \left(\frac{c}{\sqrt{bx^2 + a}} \right)^{\frac{3}{2}} dx$$

input `integrate(x^2*(c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="maxima")`

output `integrate(x^2*(c/sqrt(b*x^2 + a))^(3/2), x)`

Giac [F]

$$\int x^2 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \int x^2 \left(\frac{c}{\sqrt{bx^2 + a}} \right)^{3/2} dx$$

input `integrate(x^2*(c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="giac")`

output `integrate(x^2*(c/sqrt(b*x^2 + a))^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \int x^2 \left(\frac{c}{\sqrt{bx^2 + a}} \right)^{3/2} dx$$

input `int(x^2*(c/(a + b*x^2)^(1/2))^(3/2),x)`

output `int(x^2*(c/(a + b*x^2)^(1/2))^(3/2), x)`

Reduce [F]

$$\int x^2 \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \frac{2\sqrt{c}c \left((bx^2 + a)^{1/4} x - \left(\int \frac{1}{(bx^2+a)^{3/4}} dx \right) a \right)}{3b}$$

input `int(x^2*(c/(b*x^2+a)^(1/2))^(3/2),x)`

output `(2*sqrt(c)*c*((a + b*x**2)**(1/4)*x - int((a + b*x**2)**(1/4)/(a + b*x**2),x)*a))/(3*b)`

$$3.57 \quad \int \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx$$

Optimal result	409
Mathematica [C] (verified)	409
Rubi [A] (verified)	410
Maple [F]	411
Fricas [F]	411
Sympy [F]	412
Maxima [F]	412
Giac [F]	412
Mupad [F(-1)]	413
Reduce [F]	413

Optimal result

Integrand size = 17, antiderivative size = 62

$$\int \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \frac{2\sqrt{a} \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} \left(1 + \frac{bx^2}{a} \right)^{3/4} \text{EllipticF} \left(\frac{1}{2} \arctan \left(\frac{\sqrt{bx}}{\sqrt{a}} \right), 2 \right)}{\sqrt{b}}$$

output

```
2*a^(1/2)*(c/(b*x^2+a)^(1/2))^(3/2)*(1+b*x^2/a)^(3/4)*InverseJacobiAM(1/2*
arctan(b^(1/2)*x/a^(1/2)),2^(1/2))/b^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 5.59 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.84

$$\int \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = x \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} \left(1 + \frac{bx^2}{a} \right)^{3/4} \text{Hypergeometric2F1} \left(\frac{1}{2}, \frac{3}{4}, \frac{3}{2}, -\frac{bx^2}{a} \right)$$

input

```
Integrate[(c/Sqrt[a + b*x^2])^(3/2),x]
```

output

```
x*(c/Sqrt[a + b*x^2])^(3/2)*(1 + (b*x^2)/a)^(3/4)*Hypergeometric2F1[1/2, 3/4, 3/2, -(b*x^2)/a]
```

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2045, 229}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx$$

$$\downarrow \text{2045}$$

$$\left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} \int \frac{1}{\left(\frac{bx^2}{a} + 1 \right)^{3/4}} dx$$

$$\downarrow \text{229}$$

$$\frac{2\sqrt{a} \left(\frac{bx^2}{a} + 1 \right)^{3/4} \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} \text{EllipticF} \left(\frac{1}{2} \arctan \left(\frac{\sqrt{bx}}{\sqrt{a}} \right), 2 \right)}{\sqrt{b}}$$

input

```
Int[(c/Sqrt[a + b*x^2])^(3/2),x]
```

output

```
(2*Sqrt[a]*(c/Sqrt[a + b*x^2])^(3/2)*(1 + (b*x^2)/a)^(3/4)*EllipticF[ArcTan[(Sqrt[b]*x)/Sqrt[a]]/2, 2])/Sqrt[b]
```

Defintions of rubi rules used

rule 229 `Int[((a_) + (b_)*(x_)^2)^(-3/4), x_Symbol] := Simp[(2/(a^(3/4)*Rt[b/a, 2])
)*EllipticF[(1/2)*ArcTan[Rt[b/a, 2]*x], 2], x] /; FreeQ[{a, b}, x] && GtQ[a
, 0] && PosQ[b/a]`

rule 2045 `Int[(u_)*((c_)*((a_) + (b_)*(x_)^(n_))^(q_))^(p_), x_Symbol] := Simp[Simp[Si
mp[(c*(a + b*x^n)^q]^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q)
, x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [F]

$$\int \left(\frac{c}{\sqrt{bx^2 + a}} \right)^{\frac{3}{2}} dx$$

input `int((c/(b*x^2+a)^(1/2))^(3/2),x)`

output `int((c/(b*x^2+a)^(1/2))^(3/2),x)`

Fricas [F]

$$\int \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \int \left(\frac{c}{\sqrt{bx^2 + a}} \right)^{\frac{3}{2}} dx$$

input `integrate((c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="fricas")`

output `integral(c*sqrt(c/sqrt(b*x^2 + a))/sqrt(b*x^2 + a), x)`

Sympy [F]

$$\int \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \int \left(\frac{c}{\sqrt{a + bx^2}} \right)^{\frac{3}{2}} dx$$

input `integrate((c/(b*x**2+a)**(1/2))**(3/2),x)`

output `Integral((c/sqrt(a + b*x**2))**(3/2), x)`

Maxima [F]

$$\int \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \int \left(\frac{c}{\sqrt{bx^2 + a}} \right)^{\frac{3}{2}} dx$$

input `integrate((c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="maxima")`

output `integrate((c/sqrt(b*x^2 + a))^(3/2), x)`

Giac [F]

$$\int \left(\frac{c}{\sqrt{a + bx^2}} \right)^{3/2} dx = \int \left(\frac{c}{\sqrt{bx^2 + a}} \right)^{\frac{3}{2}} dx$$

input `integrate((c/(b*x^2+a)^(1/2))^(3/2),x, algorithm="giac")`

output `integrate((c/sqrt(b*x^2 + a))^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \int \left(\frac{c}{\sqrt{bx^2+a}} \right)^{3/2} dx$$

input `int((c/(a + b*x^2)^(1/2))^(3/2),x)`output `int((c/(a + b*x^2)^(1/2))^(3/2), x)`**Reduce [F]**

$$\int \left(\frac{c}{\sqrt{a+bx^2}} \right)^{3/2} dx = \sqrt{c} \left(\int \frac{1}{(bx^2+a)^{3/4}} dx \right) c$$

input `int((c/(b*x^2+a)^(1/2))^(3/2),x)`output `sqrt(c)*int((a + b*x**2)**(1/4)/(a + b*x**2),x)*c`

3.58
$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^2} dx$$

Optimal result	414
Mathematica [C] (verified)	414
Rubi [A] (verified)	415
Maple [F]	416
Fricas [F]	417
Sympy [F]	417
Maxima [F]	417
Giac [F(-2)]	418
Mupad [F(-1)]	418
Reduce [F]	418

Optimal result

Integrand size = 21, antiderivative size = 95

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^2} dx = -\frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} (a + bx^2)}{ax} - \frac{\sqrt{b}\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} \left(1 + \frac{bx^2}{a}\right)^{3/4} \text{EllipticF}\left(\frac{1}{2} \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right), 2\right)}{\sqrt{a}}$$

output

```
-(c/(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)/a/x-b^(1/2)*(c/(b*x^2+a)^(1/2))^(3/2)
*(1+b*x^2/a)^(3/4)*InverseJacobiAM(1/2*arctan(b^(1/2)*x/a^(1/2)),2^(1/2))/
a^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.18 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.58

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^2} dx = -\frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} \left(1 + \frac{bx^2}{a}\right)^{3/4} \text{Hypergeometric2F1}\left(-\frac{1}{2}, \frac{3}{4}, \frac{1}{2}, -\frac{bx^2}{a}\right)}{x}$$

input `Integrate[(c/Sqrt[a + b*x^2])^(3/2)/x^2,x]`

output `-(((c/Sqrt[a + b*x^2])^(3/2)*(1 + (b*x^2)/a)^(3/4)*Hypergeometric2F1[-1/2, 3/4, 1/2, -(b*x^2)/a]))/x`

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.87, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2045, 264, 229}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^2} dx \\
 & \quad \downarrow \text{2045} \\
 & \left(\frac{bx^2}{a} + 1\right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} \int \frac{1}{x^2 \left(\frac{bx^2}{a} + 1\right)^{3/4}} dx \\
 & \quad \downarrow \text{264} \\
 & \left(\frac{bx^2}{a} + 1\right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} \left(-\frac{b \int \frac{1}{\left(\frac{bx^2}{a} + 1\right)^{3/4}} dx}{2a} - \frac{\sqrt[4]{\frac{bx^2}{a} + 1}}{x} \right) \\
 & \quad \downarrow \text{229} \\
 & \left(\frac{bx^2}{a} + 1\right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} \left(-\frac{\sqrt{b} \operatorname{EllipticF}\left(\frac{1}{2} \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right), 2\right)}{\sqrt{a}} - \frac{\sqrt[4]{\frac{bx^2}{a} + 1}}{x} \right)
 \end{aligned}$$

input `Int[(c/Sqrt[a + b*x^2])^(3/2)/x^2,x]`

output $(c/\sqrt{a + b*x^2})^{3/2}*(1 + (b*x^2)/a)^{3/4}*(-(1 + (b*x^2)/a)^{1/4}/x) - (\sqrt{b}*\text{EllipticF}[\text{ArcTan}[(\sqrt{b}*x)/\sqrt{a}]/2, 2])/\sqrt{a}$

Defintions of rubi rules used

rule 229 $\text{Int}[(a_ + (b_)*(x_)^2)^{-3/4}, x_Symbol] \rightarrow \text{Simp}[(2/(a^{3/4}*\text{Rt}[b/a, 2]))*\text{EllipticF}[(1/2)*\text{ArcTan}[\text{Rt}[b/a, 2]*x], 2], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{GtQ}[a, 0] \ \&\& \ \text{PosQ}[b/a]$

rule 264 $\text{Int}[(c_)*(x_)^m*((a_ + (b_)*(x_)^2)^{p_}), x_Symbol] \rightarrow \text{Simp}[(c*x)^{(m+1)}*((a + b*x^2)^{(p+1})/(a*c*(m+1))), x] - \text{Simp}[b*((m+2*p+3)/(a*c^2*(m+1)) \ \text{Int}[(c*x)^{(m+2)}*(a + b*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, p\}, x] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 2045 $\text{Int}[(u_)*((c_)*((a_ + (b_)*(x_)^n)^{q_}))^{p_}], x_Symbol] \rightarrow \text{Simp}[\text{Simp}[c*(a + b*x^n)^q]^p/(1 + b*(x^n/a))^{p*q}] \ \text{Int}[u*(1 + b*(x^n/a))^{p*q}, x], x] /; \text{FreeQ}\{a, b, c, n, p, q\}, x] \ \&\& \ !\text{GeQ}[a, 0]$

Maple [F]

$$\int \frac{\left(\frac{c}{\sqrt{bx^2+a}}\right)^{\frac{3}{2}}}{x^2} dx$$

input $\text{int}((c/(b*x^2+a)^{(1/2}))^{3/2}/x^2,x)$

output $\text{int}((c/(b*x^2+a)^{(1/2}))^{3/2}/x^2,x)$

Fricas [F]

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^2} dx = \int \frac{\left(\frac{c}{\sqrt{bx^2+a}}\right)^{3/2}}{x^2} dx$$

input `integrate((c/(b*x^2+a)^(1/2))^(3/2)/x^2,x, algorithm="fricas")`

output `integral(sqrt(b*x^2 + a)*c*sqrt(c/sqrt(b*x^2 + a))/(b*x^4 + a*x^2), x)`

Sympy [F]

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^2} dx = \int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^2} dx$$

input `integrate((c/(b*x**2+a)**(1/2))**(3/2)/x**2,x)`

output `Integral((c/sqrt(a + b*x**2))**(3/2)/x**2, x)`

Maxima [F]

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^2} dx = \int \frac{\left(\frac{c}{\sqrt{bx^2+a}}\right)^{3/2}}{x^2} dx$$

input `integrate((c/(b*x^2+a)^(1/2))^(3/2)/x^2,x, algorithm="maxima")`

output `integrate((c/sqrt(b*x^2 + a))^(3/2)/x^2, x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^2} dx = \text{Exception raised: TypeError}$$

input `integrate((c/(b*x^2+a)^(1/2))^(3/2)/x^2,x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to ro
unding error%%{1,[0,1,0,0]%%} / %%{1,[0,0,1,2]%%} Error: Bad Argument
Value

Mupad [F(-1)]

Timed out.

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^2} dx = \int \frac{\left(\frac{c}{\sqrt{bx^2+a}}\right)^{3/2}}{x^2} dx$$

input `int((c/(a + b*x^2)^(1/2))^(3/2)/x^2,x)`

output `int((c/(a + b*x^2)^(1/2))^(3/2)/x^2, x)`

Reduce [F]

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^2} dx = \sqrt{c} \left(\int \frac{(bx^2 + a)^{\frac{1}{4}}}{bx^4 + ax^2} dx \right) c$$

input `int((c/(b*x^2+a)^(1/2))^(3/2)/x^2,x)`

output `sqrt(c)*int((a + b*x**2)**(1/4)/(a*x**2 + b*x**4),x)*c`

3.59
$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^4} dx$$

Optimal result	420
Mathematica [C] (verified)	420
Rubi [A] (verified)	421
Maple [F]	423
Fricas [F]	423
Sympy [F]	423
Maxima [F]	424
Giac [F(-2)]	424
Mupad [F(-1)]	424
Reduce [F]	425

Optimal result

Integrand size = 21, antiderivative size = 134

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^4} dx = -\frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} (a + bx^2)}{3ax^3} + \frac{5b\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} (a + bx^2)}{6a^2x} + \frac{5b^{3/2}\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} \left(1 + \frac{bx^2}{a}\right)^{3/4} \text{EllipticF}\left(\frac{1}{2} \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right), 2\right)}{6a^{3/2}}$$

output

```
-1/3*(c/(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)/a/x^3+5/6*b*(c/(b*x^2+a)^(1/2))^(3/2)*(b*x^2+a)/a^2/x+5/6*b^(3/2)*(c/(b*x^2+a)^(1/2))^(3/2)*(1+b*x^2/a)^(3/4)*InverseJacobiAM(1/2*arctan(b^(1/2)*x/a^(1/2)),2^(1/2))/a^(3/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.02 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.43

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^4} dx = -\frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} \left(1 + \frac{bx^2}{a}\right)^{3/4} \text{Hypergeometric2F1}\left(-\frac{3}{2}, \frac{3}{4}, -\frac{1}{2}, -\frac{bx^2}{a}\right)}{3x^3}$$

input `Integrate[(c/Sqrt[a + b*x^2])^(3/2)/x^4,x]`

output `-1/3*((c/Sqrt[a + b*x^2])^(3/2)*(1 + (b*x^2)/a)^(3/4)*Hypergeometric2F1[-3/2, 3/4, -1/2, -((b*x^2)/a)])/x^3`

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.84, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2045, 264, 264, 229}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^4} dx \\
 & \quad \downarrow \text{2045} \\
 & \left(\frac{bx^2}{a} + 1\right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} \int \frac{1}{x^4 \left(\frac{bx^2}{a} + 1\right)^{3/4}} dx \\
 & \quad \downarrow \text{264} \\
 & \left(\frac{bx^2}{a} + 1\right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} \left(-\frac{5b \int \frac{1}{x^2 \left(\frac{bx^2}{a} + 1\right)^{3/4}} dx}{6a} - \frac{\sqrt[4]{\frac{bx^2}{a} + 1}}{3x^3} \right) \\
 & \quad \downarrow \text{264} \\
 & \left(\frac{bx^2}{a} + 1\right)^{3/4} \left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2} \left(-\frac{5b \left(-\frac{b \int \frac{1}{\left(\frac{bx^2}{a} + 1\right)^{3/4}} dx}{2a} - \frac{\sqrt[4]{\frac{bx^2}{a} + 1}}{x} \right)}{6a} - \frac{\sqrt[4]{\frac{bx^2}{a} + 1}}{3x^3} \right) \\
 & \quad \downarrow \text{229}
 \end{aligned}$$

$$\left(\frac{bx^2}{a} + 1\right)^{3/4} \left(\frac{c}{\sqrt{a + bx^2}}\right)^{3/2} \left(-\frac{5b \left(\frac{\sqrt{b} \operatorname{EllipticF}\left(\frac{1}{2} \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right), 2\right)}{\sqrt{a}} - \frac{\sqrt[4]{\frac{bx^2}{a} + 1}}{x} \right)}{6a} - \frac{\sqrt[4]{\frac{bx^2}{a} + 1}}{3x^3} \right)$$

input `Int[(c/Sqrt[a + b*x^2])^(3/2)/x^4,x]`

output `(c/Sqrt[a + b*x^2])^(3/2)*(1 + (b*x^2)/a)^(3/4)*(-1/3*(1 + (b*x^2)/a)^(1/4)/x^3 - (5*b*(-((1 + (b*x^2)/a)^(1/4)/x) - (Sqrt[b]*EllipticF[ArcTan[(Sqrt[b]*x)/Sqrt[a]], 2])/Sqrt[a]))/(6*a))`

Defintions of rubi rules used

rule 229 `Int[((a_) + (b_.)*(x_)^2)^(-3/4), x_Symbol] := Simp[(2/(a^(3/4)*Rt[b/a, 2]) * EllipticF[(1/2)*ArcTan[Rt[b/a, 2]*x], 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]`

rule 264 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*(m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [F]

$$\int \frac{\left(\frac{c}{\sqrt{bx^2+a}}\right)^{\frac{3}{2}}}{x^4} dx$$

input `int((c/(b*x^2+a)^(1/2))^(3/2)/x^4,x)`

output `int((c/(b*x^2+a)^(1/2))^(3/2)/x^4,x)`

Fricas [F]

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^4} dx = \int \frac{\left(\frac{c}{\sqrt{bx^2+a}}\right)^{\frac{3}{2}}}{x^4} dx$$

input `integrate((c/(b*x^2+a)^(1/2))^(3/2)/x^4,x, algorithm="fricas")`

output `integral(sqrt(b*x^2 + a)*c*sqrt(c/sqrt(b*x^2 + a))/(b*x^6 + a*x^4), x)`

Sympy [F]

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^4} dx = \int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{\frac{3}{2}}}{x^4} dx$$

input `integrate((c/(b*x**2+a)**(1/2))**(3/2)/x**4,x)`

output `Integral((c/sqrt(a + b*x**2))**(3/2)/x**4, x)`

Maxima [F]

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^4} dx = \int \frac{\left(\frac{c}{\sqrt{bx^2+a}}\right)^{3/2}}{x^4} dx$$

input `integrate((c/(b*x^2+a)^(1/2))^(3/2)/x^4,x, algorithm="maxima")`

output `integrate((c/sqrt(b*x^2 + a))^(3/2)/x^4, x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^4} dx = \text{Exception raised: TypeError}$$

input `integrate((c/(b*x^2+a)^(1/2))^(3/2)/x^4,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to ro
unding error%%{1,[0,1,1,2,0,0]%%}+%%{-1,[0,1,0,0,1,0]%%} / %%{1,[0,0,
2,0,0,2]%`

Mupad [F(-1)]

Timed out.

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^4} dx = \int \frac{\left(\frac{c}{\sqrt{bx^2+a}}\right)^{3/2}}{x^4} dx$$

input `int((c/(a + b*x^2)^(1/2))^(3/2)/x^4,x)`

output `int((c/(a + b*x^2)^(1/2))^(3/2)/x^4, x)`

Reduce [F]

$$\int \frac{\left(\frac{c}{\sqrt{a+bx^2}}\right)^{3/2}}{x^4} dx = \sqrt{c} \left(\int \frac{(bx^2+a)^{1/4}}{bx^6+ax^4} dx \right) c$$

input `int((c/(b*x^2+a)^(1/2))^(3/2)/x^4,x)`

output `sqrt(c)*int((a + b*x**2)**(1/4)/(a*x**4 + b*x**6),x)*c`

3.60 $\int (dx)^m \left(c(a + bx^2)^3 \right)^{3/2} dx$

Optimal result	426
Mathematica [A] (verified)	426
Rubi [A] (verified)	427
Maple [F]	428
Fricas [F]	428
Sympy [F(-2)]	429
Maxima [F]	429
Giac [F]	429
Mupad [F(-1)]	430
Reduce [F]	430

Optimal result

Integrand size = 21, antiderivative size = 76

$$\int (dx)^m \left(c(a + bx^2)^3 \right)^{3/2} dx = \frac{a^3 c (dx)^{1+m} \sqrt{c(a + bx^2)^3} \operatorname{Hypergeometric2F1} \left(-\frac{9}{2}, \frac{1+m}{2}, \frac{3+m}{2}, -\frac{bx^2}{a} \right)}{d(1+m) \left(1 + \frac{bx^2}{a} \right)^{3/2}}$$

output

```
a^3*c*(d*x)^(1+m)*(c*(b*x^2+a)^3)^(1/2)*hypergeom([-9/2, 1/2+1/2*m], [3/2+1/2*m], -b*x^2/a)/d/(1+m)/(1+b*x^2/a)^(3/2)
```

Mathematica [A] (verified)

Time = 0.41 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.08

$$\int (dx)^m \left(c(a + bx^2)^3 \right)^{3/2} dx = \frac{a^4 x (dx)^m \left(c(a + bx^2)^3 \right)^{3/2} \operatorname{Hypergeometric2F1} \left(-\frac{9}{2}, \frac{1+m}{2}, 1 + \frac{1+m}{2}, -\frac{bx^2}{a} \right)}{(1+m) (a + bx^2)^4 \sqrt{1 + \frac{bx^2}{a}}}$$

input `Integrate[(d*x)^m*(c*(a + b*x^2)^3)^(3/2),x]`

output `(a^4*x*(d*x)^m*(c*(a + b*x^2)^3)^(3/2)*Hypergeometric2F1[-9/2, (1 + m)/2, 1 + (1 + m)/2, -((b*x^2)/a)]/((1 + m)*(a + b*x^2)^4*Sqrt[1 + (b*x^2)/a])`

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2045, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m \left(c(a + bx^2)^3 \right)^{3/2} dx$$

$$\downarrow \text{2045}$$

$$\frac{a^3 c \sqrt{c(a + bx^2)^3} \int (dx)^m \left(\frac{bx^2}{a} + 1 \right)^{9/2} dx}{\left(\frac{bx^2}{a} + 1 \right)^{3/2}}$$

$$\downarrow \text{278}$$

$$\frac{a^3 c (dx)^{m+1} \sqrt{c(a + bx^2)^3} \text{Hypergeometric2F1} \left(-\frac{9}{2}, \frac{m+1}{2}, \frac{m+3}{2}, -\frac{bx^2}{a} \right)}{d(m+1) \left(\frac{bx^2}{a} + 1 \right)^{3/2}}$$

input `Int[(d*x)^m*(c*(a + b*x^2)^3)^(3/2),x]`

output `(a^3*c*(d*x)^(1 + m)*Sqrt[c*(a + b*x^2)^3]*Hypergeometric2F1[-9/2, (1 + m)/2, (3 + m)/2, -((b*x^2)/a)]/(d*(1 + m)*(1 + (b*x^2)/a)^(3/2))`

Defintions of rubi rules used

rule 278

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])
```

rule 2045

```
Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]
```

Maple [F]

$$\int (xd)^m (c(bx^2 + a)^3)^{\frac{3}{2}} dx$$

input

```
int((x*d)^m*(c*(b*x^2+a)^3)^(3/2),x)
```

output

```
int((x*d)^m*(c*(b*x^2+a)^3)^(3/2),x)
```

Fricas [F]

$$\int (dx)^m (c(a + bx^2)^3)^{3/2} dx = \int ((bx^2 + a)^3 c)^{\frac{3}{2}} (dx)^m dx$$

input

```
integrate((d*x)^m*(c*(b*x^2+a)^3)^(3/2),x, algorithm="fricas")
```

output

```
integral((b^3*c*x^6 + 3*a*b^2*c*x^4 + 3*a^2*b*c*x^2 + a^3*c)^(3/2)*(d*x)^m, x)
```

Sympy [F(-2)]

Exception generated.

$$\int (dx)^m \left(c(a + bx^2)^3 \right)^{3/2} dx = \text{Exception raised: HeuristicGCDFailed}$$

input `integrate((d*x)**m*(c*(b*x**2+a)**3)**(3/2),x)`

output `Exception raised: HeuristicGCDFailed >> no luck`

Maxima [F]

$$\int (dx)^m \left(c(a + bx^2)^3 \right)^{3/2} dx = \int \left((bx^2 + a)^3 c \right)^{\frac{3}{2}} (dx)^m dx$$

input `integrate((d*x)^m*(c*(b*x^2+a)^3)^(3/2),x, algorithm="maxima")`

output `integrate(((b*x^2 + a)^3*c)^(3/2)*(d*x)^m, x)`

Giac [F]

$$\int (dx)^m \left(c(a + bx^2)^3 \right)^{3/2} dx = \int \left((bx^2 + a)^3 c \right)^{\frac{3}{2}} (dx)^m dx$$

input `integrate((d*x)^m*(c*(b*x^2+a)^3)^(3/2),x, algorithm="giac")`

output `integrate(((b*x^2 + a)^3*c)^(3/2)*(d*x)^m, x)`

Mupad [F(-1)]

Timed out.

$$\int (dx)^m (c(a + bx^2)^3)^{3/2} dx = \int (dx)^m (c(bx^2 + a)^3)^{3/2} dx$$

input `int((d*x)^m*(c*(a + b*x^2)^3)^(3/2), x)`

output `int((d*x)^m*(c*(a + b*x^2)^3)^(3/2), x)`

Reduce [F]

$$\begin{aligned} \int (dx)^m (c(a + bx^2)^3)^{3/2} dx &= d^m \sqrt{c} c \left(\left(\int x^m \sqrt{bx^2 + a} x^8 dx \right) b^4 \right. \\ &+ 4 \left(\int x^m \sqrt{bx^2 + a} x^6 dx \right) a b^3 + 6 \left(\int x^m \sqrt{bx^2 + a} x^4 dx \right) a^2 b^2 \\ &\left. + 4 \left(\int x^m \sqrt{bx^2 + a} x^2 dx \right) a^3 b + \left(\int x^m \sqrt{bx^2 + a} dx \right) a^4 \right) \end{aligned}$$

input `int((d*x)^m*(c*(b*x^2+a)^3)^(3/2), x)`

output `d**m*sqrt(c)*c*(int(x**m*sqrt(a + b*x**2)*x**8,x)*b**4 + 4*int(x**m*sqrt(a + b*x**2)*x**6,x)*a*b**3 + 6*int(x**m*sqrt(a + b*x**2)*x**4,x)*a**2*b**2 + 4*int(x**m*sqrt(a + b*x**2)*x**2,x)*a**3*b + int(x**m*sqrt(a + b*x**2),x)*a**4)`

3.61 $\int (dx)^m \left(c(a + bx^2)^2 \right)^{3/2} dx$

Optimal result	431
Mathematica [A] (verified)	432
Rubi [A] (verified)	432
Maple [A] (verified)	434
Fricas [A] (verification not implemented)	434
Sympy [F(-1)]	435
Maxima [A] (verification not implemented)	435
Giac [B] (verification not implemented)	435
Mupad [B] (verification not implemented)	436
Reduce [B] (verification not implemented)	437

Optimal result

Integrand size = 21, antiderivative size = 181

$$\int (dx)^m \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{a^3 c (dx)^{1+m} \sqrt{c(a + bx^2)^2}}{d(1 + m)(a + bx^2)} + \frac{3a^2 b c (dx)^{3+m} \sqrt{c(a + bx^2)^2}}{d^3(3 + m)(a + bx^2)} + \frac{3ab^2 c (dx)^{5+m} \sqrt{c(a + bx^2)^2}}{d^5(5 + m)(a + bx^2)} + \frac{b^3 c (dx)^{7+m} \sqrt{c(a + bx^2)^2}}{d^7(7 + m)(a + bx^2)}$$

output

```
a^3*c*(d*x)^(1+m)*(c*(b*x^2+a)^2)^(1/2)/d/(1+m)/(b*x^2+a)+3*a^2*b*c*(d*x)^(3+m)*(c*(b*x^2+a)^2)^(1/2)/d^3/(3+m)/(b*x^2+a)+3*a*b^2*c*(d*x)^(5+m)*(c*(b*x^2+a)^2)^(1/2)/d^5/(5+m)/(b*x^2+a)+b^3*c*(d*x)^(7+m)*(c*(b*x^2+a)^2)^(1/2)/d^7/(7+m)/(b*x^2+a)
```


Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.73

$$\int (dx)^m \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{x(dx)^m \left(c(a + bx^2)^2 \right)^{3/2} \left(a^3(105 + 71m + 15m^2 + m^3) + 3a^2b(35 + 47m + 13m^2 + m^3) + 3ab^2(21 + 31m + 11m^2 + m^3) + b^3(15 + 23m + 9m^2 + m^3)x^6 \right)}{(1+m)(3+m)(5+m)(7+m)}$$

input `Integrate[(d*x)^m*(c*(a + b*x^2)^2)^(3/2),x]`output `(x*(d*x)^m*(c*(a + b*x^2)^2)^(3/2)*(a^3*(105 + 71*m + 15*m^2 + m^3) + 3*a^2*b*(35 + 47*m + 13*m^2 + m^3)*x^2 + 3*a*b^2*(21 + 31*m + 11*m^2 + m^3)*x^4 + b^3*(15 + 23*m + 9*m^2 + m^3)*x^6))/((1 + m)*(3 + m)*(5 + m)*(7 + m)*(a + b*x^2)^3)`**Rubi [A] (verified)**Time = 0.49 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.59, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2045, 27, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (dx)^m \left(c(a + bx^2)^2 \right)^{3/2} dx \\ & \quad \downarrow \text{2045} \\ & \frac{a^3 c \sqrt{c(a + bx^2)^2} \int \frac{(dx)^m (bx^2 + a)^3}{a^3} dx}{a + bx^2} \\ & \quad \downarrow \text{27} \\ & \frac{c \sqrt{c(a + bx^2)^2} \int (dx)^m (bx^2 + a)^3 dx}{a + bx^2} \\ & \quad \downarrow \text{244} \end{aligned}$$

$$\frac{c\sqrt{c(a+bx^2)^2} \int \left(a^3(dx)^m + \frac{3a^2b(dx)^{m+2}}{d^2} + \frac{3ab^2(dx)^{m+4}}{d^4} + \frac{b^3(dx)^{m+6}}{d^6} \right) dx}{a+bx^2}$$

↓ 2009

$$\frac{c\sqrt{c(a+bx^2)^2} \left(\frac{a^3(dx)^{m+1}}{d(m+1)} + \frac{3a^2b(dx)^{m+3}}{d^3(m+3)} + \frac{3ab^2(dx)^{m+5}}{d^5(m+5)} + \frac{b^3(dx)^{m+7}}{d^7(m+7)} \right)}{a+bx^2}$$

input `Int[(d*x)^m*(c*(a + b*x^2)^2)^(3/2),x]`

output `(c*Sqrt[c*(a + b*x^2)^2]*((a^3*(d*x)^(1 + m))/(d*(1 + m)) + (3*a^2*b*(d*x)^(3 + m))/(d^3*(3 + m)) + (3*a*b^2*(d*x)^(5 + m))/(d^5*(5 + m)) + (b^3*(d*x)^(7 + m))/(d^7*(7 + m)))/(a + b*x^2)`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 244 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2045 `Int[(u_)*((c_)*((a_) + (b_)*(x_)^(n_))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.11

method	result
gospers	$\frac{x(b^3m^3x^6+9b^3m^2x^6+3ab^2m^3x^4+23mx^6b^3+33ab^2m^2x^4+15b^3x^6+3a^2bm^3x^2+93mx^4ab^2+39a^2bm^2x^2+63ab^2x^4+a^3m^3+141a^2m^2x^2+15a^3m^2x^2+105a^2m^2x^2+71a^3m+105a^3)(b^2x^2+a)^{3/2}}{(7+m)(5+m)(3+m)(1+m)(b^2x^2+a)^3}$
orering	$\frac{x(b^3m^3x^6+9b^3m^2x^6+3ab^2m^3x^4+23mx^6b^3+33ab^2m^2x^4+15b^3x^6+3a^2bm^3x^2+93mx^4ab^2+39a^2bm^2x^2+63ab^2x^4+a^3m^3+141a^2m^2x^2+15a^3m^2x^2+105a^2m^2x^2+71a^3m+105a^3)(b^2x^2+a)^{3/2}}{(7+m)(5+m)(3+m)(1+m)(b^2x^2+a)^3}$
risch	$\frac{c\sqrt{c(b^2x^2+a)^2}(b^3m^3x^6+9b^3m^2x^6+3ab^2m^3x^4+23mx^6b^3+33ab^2m^2x^4+15b^3x^6+3a^2bm^3x^2+93mx^4ab^2+39a^2bm^2x^2+63ab^2x^4+a^3m^3+141a^2m^2x^2+15a^3m^2x^2+105a^2m^2x^2+71a^3m+105a^3)(b^2x^2+a)^{3/2}}{(b^2x^2+a)(7+m)(5+m)(3+m)(1+m)}$

input `int((x*d)^m*(c*(b*x^2+a)^2)^(3/2),x,method=_RETURNVERBOSE)`

output
$$x*(b^3m^3x^6+9b^3m^2x^6+3a*b^2m^3x^4+23*b^3m*x^6+33*a*b^2m^2x^4+15*b^3x^6+3a^2*b*m^3x^2+93*a*b^2m^2x^2+63*a*b^2x^4+a^3m^3+141*a^2*b*m*x^2+15*a^3m^2+105*a^2*b*x^2+71*a^3m+105*a^3)*(x*d)^m*(c*(b*x^2+a)^2)^(3/2)/(7+m)/(5+m)/(3+m)/(1+m)/(b*x^2+a)^3$$

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.30

$$\int (dx)^m \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{((b^3cm^3 + 9b^3cm^2 + 23b^3cm + 15b^3c)x^7 + 3(ab^2cm^3 + 11ab^2cm^2 + 31ab^2cm + 21ab^2c)am^4 + 16am^3 + 86am^2 -$$

input `integrate((d*x)^m*(c*(b*x^2+a)^2)^(3/2),x, algorithm="fricas")`

output
$$((b^3*c*m^3 + 9*b^3*c*m^2 + 23*b^3*c*m + 15*b^3*c)*x^7 + 3*(a*b^2*c*m^3 + 11*a*b^2*c*m^2 + 31*a*b^2*c*m + 21*a*b^2*c)*x^5 + 3*(a^2*b*c*m^3 + 13*a^2*b*c*m^2 + 47*a^2*b*c*m + 35*a^2*b*c)*x^3 + (a^3*c*m^3 + 15*a^3*c*m^2 + 71*a^3*c*m + 105*a^3*c)*x)*sqrt(b^2*c*x^4 + 2*a*b*c*x^2 + a^2*c)*(d*x)^m/(a*m^4 + 16*a*m^3 + 86*a*m^2 + (b*m^4 + 16*b*m^3 + 86*b*m^2 + 176*b*m + 105*b)*x^2 + 176*a*m + 105*a)$$

Sympy [F(-1)]

Timed out.

$$\int (dx)^m \left(c(a + bx^2)^2 \right)^{3/2} dx = \text{Timed out}$$

input `integrate((d*x)**m*(c*(b*x**2+a)**2)**(3/2),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.72

$$\int (dx)^m \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{\left((m^3 + 9m^2 + 23m + 15)b^3c^{\frac{3}{2}}d^m x^7 + 3(m^3 + 11m^2 + 31m + 21)ab^2c^{\frac{3}{2}}d^m x^5 + 3(m^3 + 17m^2 + 47m + 35)a^2b^2c^{\frac{3}{2}}d^m x^3 + (m^3 + 15m^2 + 71m + 105)a^3c^{\frac{3}{2}}d^m x \right)}{m^4 + 16m^3 + 86m^2 + 176m + 105}$$

input `integrate((d*x)^m*(c*(b*x^2+a)^2)^(3/2),x, algorithm="maxima")`

output `((m^3 + 9*m^2 + 23*m + 15)*b^3*c^(3/2)*d^m*x^7 + 3*(m^3 + 11*m^2 + 31*m + 21)*a*b^2*c^(3/2)*d^m*x^5 + 3*(m^3 + 13*m^2 + 47*m + 35)*a^2*b*c^(3/2)*d^m*x^3 + (m^3 + 15*m^2 + 71*m + 105)*a^3*c^(3/2)*d^m*x)/(m^4 + 16*m^3 + 86*m^2 + 176*m + 105)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 387 vs. 2(173) = 346.

Time = 0.13 (sec) , antiderivative size = 387, normalized size of antiderivative = 2.14

$$\int (dx)^m \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{\left((dx)^m b^3 m^3 x^7 \operatorname{sgn}(bx^2 + a) + 9(dx)^m b^3 m^2 x^7 \operatorname{sgn}(bx^2 + a) + 3(dx)^m ab^2 m^3 x^5 \operatorname{sgn}(bx^2 + a) + 3(dx)^m a^2 b^2 m^3 x^3 \operatorname{sgn}(bx^2 + a) + 3(dx)^m a^3 m \operatorname{sgn}(bx^2 + a) \right)}{m^4 + 16m^3 + 86m^2 + 176m + 105}$$

input `integrate((d*x)^m*(c*(b*x^2+a)^2)^(3/2),x, algorithm="giac")`

output
$$\begin{aligned} & ((d*x)^m*b^3*m^3*x^7*sgn(b*x^2 + a) + 9*(d*x)^m*b^3*m^2*x^7*sgn(b*x^2 + a) \\ & + 3*(d*x)^m*a*b^2*m^3*x^5*sgn(b*x^2 + a) + 23*(d*x)^m*b^3*m*x^7*sgn(b*x^2 \\ & + a) + 33*(d*x)^m*a*b^2*m^2*x^5*sgn(b*x^2 + a) + 15*(d*x)^m*b^3*x^7*sgn(b \\ & *x^2 + a) + 3*(d*x)^m*a^2*b*m^3*x^3*sgn(b*x^2 + a) + 93*(d*x)^m*a*b^2*m*x^ \\ & 5*sgn(b*x^2 + a) + 39*(d*x)^m*a^2*b*m^2*x^3*sgn(b*x^2 + a) + 63*(d*x)^m*a* \\ & b^2*x^5*sgn(b*x^2 + a) + (d*x)^m*a^3*m^3*x*sgn(b*x^2 + a) + 141*(d*x)^m*a^ \\ & 2*b*m*x^3*sgn(b*x^2 + a) + 15*(d*x)^m*a^3*m^2*x*sgn(b*x^2 + a) + 105*(d*x) \\ & ^m*a^2*b*x^3*sgn(b*x^2 + a) + 71*(d*x)^m*a^3*m*x*sgn(b*x^2 + a) + 105*(d*x) \\ & ^m*a^3*x*sgn(b*x^2 + a)) * c^(3/2) / (m^4 + 16*m^3 + 86*m^2 + 176*m + 105) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.76 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.30

$$\int (dx)^m \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{(dx)^m \left(\frac{3a^2 cx^3 \sqrt{c(bx^2+a)^2} (m^3+13m^2+47m+35)}{m^4+16m^3+86m^2+176m+105} + \frac{b^2 cx^7 \sqrt{c(bx^2+a)^2} (m^3+9m^2+23m+15)}{m^4+16m^3+86m^2+176m+105} + \frac{3abcx^5 \sqrt{c(bx^2+a)^2}}{m^4} \right)}{\frac{a}{b} + x^2}$$

input `int((d*x)^m*(c*(a + b*x^2)^2)^(3/2),x)`

output
$$\begin{aligned} & ((d*x)^m*((3*a^2*c*x^3*(c*(a + b*x^2)^2)^(1/2)*(47*m + 13*m^2 + m^3 + 35)) \\ & / (176*m + 86*m^2 + 16*m^3 + m^4 + 105) + (b^2*c*x^7*(c*(a + b*x^2)^2)^(1/2) \\ &)*(23*m + 9*m^2 + m^3 + 15))/(176*m + 86*m^2 + 16*m^3 + m^4 + 105) + (3*a* \\ & b*c*x^5*(c*(a + b*x^2)^2)^(1/2)*(31*m + 11*m^2 + m^3 + 21))/(176*m + 86*m^ \\ & 2 + 16*m^3 + m^4 + 105) + (a^3*c*x*(c*(a + b*x^2)^2)^(1/2)*(71*m + 15*m^2 \\ & + m^3 + 105))/(b*(176*m + 86*m^2 + 16*m^3 + m^4 + 105))))/(a/b + x^2) \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.01

$$\int (dx)^m \left(c(a + bx^2)^2 \right)^{3/2} dx = \frac{x^m d^m \sqrt{c} c x (b^3 m^3 x^6 + 9b^3 m^2 x^6 + 3a b^2 m^3 x^4 + 23b^3 m x^6 + 33a b^2 m^2 x^4 + 15b^3 x^6 + 3a^2 b m^3 x^6 + 16m^4 + 105m^3 + 86m^2 + 176m + 105)}{m^4 + 16m^3 + 86m^2 + 176m + 105}$$

input `int((d*x)^m*(c*(b*x^2+a)^2)^(3/2),x)`output `(x**m*d**m*sqrt(c)*c*x*(a**3*m**3 + 15*a**3*m**2 + 71*a**3*m + 105*a**3 + 3*a**2*b*m**3*x**2 + 39*a**2*b*m**2*x**2 + 141*a**2*b*m*x**2 + 105*a**2*b*x**2 + 3*a*b**2*m**3*x**4 + 33*a*b**2*m**2*x**4 + 93*a*b**2*m*x**4 + 63*a*b**2*x**4 + b**3*m**3*x**6 + 9*b**3*m**2*x**6 + 23*b**3*m*x**6 + 15*b**3*x**6))/(m**4 + 16*m**3 + 86*m**2 + 176*m + 105)`

3.62 $\int (dx)^m (c(a + bx^2))^{3/2} dx$

Optimal result	438
Mathematica [A] (verified)	438
Rubi [A] (verified)	439
Maple [F]	440
Fricas [F]	440
Sympy [F(-1)]	441
Maxima [F]	441
Giac [F]	441
Mupad [F(-1)]	442
Reduce [F]	442

Optimal result

Integrand size = 19, antiderivative size = 73

$$\int (dx)^m (c(a + bx^2))^{3/2} dx = \frac{ac(dx)^{1+m} \sqrt{ac + bcx^2} \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, \frac{1+m}{2}, \frac{3+m}{2}, -\frac{bx^2}{a}\right)}{d(1+m)\sqrt{1 + \frac{bx^2}{a}}}$$

output

```
a*c*(d*x)^(1+m)*(b*c*x^2+a*c)^(1/2)*hypergeom([-3/2, 1/2+1/2*m], [3/2+1/2*m], -b*x^2/a)/d/(1+m)/(1+b*x^2/a)^(1/2)
```

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.93

$$\int (dx)^m (c(a + bx^2))^{3/2} dx = \frac{acx(dx)^m \sqrt{c(a + bx^2)} \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, \frac{1+m}{2}, \frac{3+m}{2}, -\frac{bx^2}{a}\right)}{(1+m)\sqrt{1 + \frac{bx^2}{a}}}$$

input

```
Integrate[(d*x)^m*(c*(a + b*x^2))^(3/2),x]
```

output

$$(a*c*x*(d*x)^m*sqrt[c*(a + b*x^2)]*Hypergeometric2F1[-3/2, (1 + m)/2, (3 + m)/2, -((b*x^2)/a)]/((1 + m)*sqrt[1 + (b*x^2)/a])$$
Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2073, 279, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (dx)^m (c(a + bx^2))^{3/2} dx \\ & \quad \downarrow \text{2073} \\ & \int (dx)^m (ac + bcx^2)^{3/2} dx \\ & \quad \downarrow \text{279} \\ & \frac{ac\sqrt{ac + bcx^2} \int (dx)^m \left(\frac{bx^2}{a} + 1\right)^{3/2} dx}{\sqrt{\frac{bx^2}{a} + 1}} \\ & \quad \downarrow \text{278} \\ & \frac{ac(dx)^{m+1}\sqrt{ac + bcx^2} \text{Hypergeometric2F1}\left(-\frac{3}{2}, \frac{m+1}{2}, \frac{m+3}{2}, -\frac{bx^2}{a}\right)}{d(m+1)\sqrt{\frac{bx^2}{a} + 1}} \end{aligned}$$

input

$$\text{Int}[(d*x)^m*(c*(a + b*x^2))^(3/2), x]$$

output

$$(a*c*(d*x)^(1 + m)*sqrt[a*c + b*c*x^2]*Hypergeometric2F1[-3/2, (1 + m)/2, (3 + m)/2, -((b*x^2)/a)]/(d*(1 + m)*sqrt[1 + (b*x^2)/a])$$

Defintions of rubi rules used

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 279 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(c*x)^m*(1 + b*(x^2/a))^p, x], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && !(ILtQ[p, 0] || GtQ[a, 0])`

rule 2073 `Int[(u_)^(p_.)*((c_.)*(x_))^(m_.), x_Symbol] := Int[(c*x)^m*ExpandToSum[u, x]^p, x] /; FreeQ[{c, m, p}, x] && BinomialQ[u, x] && !BinomialMatchQ[u, x]`

Maple [F]

$$\int (xd)^m ((bx^2 + a)c)^{\frac{3}{2}} dx$$

input `int((x*d)^m*((b*x^2+a)*c)^(3/2),x)`

output `int((x*d)^m*((b*x^2+a)*c)^(3/2),x)`

Fricas [F]

$$\int (dx)^m (c(a + bx^2))^{3/2} dx = \int ((bx^2 + a)c)^{\frac{3}{2}} (dx)^m dx$$

input `integrate((d*x)^m*(c*(b*x^2+a))^(3/2),x, algorithm="fricas")`

output `integral((b*c*x^2 + a*c)^(3/2)*(d*x)^m, x)`

Sympy [F(-1)]

Timed out.

$$\int (dx)^m (c(a + bx^2))^{3/2} dx = \text{Timed out}$$

input `integrate((d*x)**m*(c*(b*x**2+a))**(3/2),x)`

output `Timed out`

Maxima [F]

$$\int (dx)^m (c(a + bx^2))^{3/2} dx = \int ((bx^2 + a)c)^{\frac{3}{2}} (dx)^m dx$$

input `integrate((d*x)^m*(c*(b*x^2+a))^(3/2),x, algorithm="maxima")`

output `integrate(((b*x^2 + a)*c)^(3/2)*(d*x)^m, x)`

Giac [F]

$$\int (dx)^m (c(a + bx^2))^{3/2} dx = \int ((bx^2 + a)c)^{\frac{3}{2}} (dx)^m dx$$

input `integrate((d*x)^m*(c*(b*x^2+a))^(3/2),x, algorithm="giac")`

output `integrate(((b*x^2 + a)*c)^(3/2)*(d*x)^m, x)`

Mupad [F(-1)]

Timed out.

$$\int (dx)^m (c(a + bx^2))^{3/2} dx = \int (c(bx^2 + a))^{3/2} (dx)^m dx$$

input `int((c*(a + b*x^2))^(3/2)*(d*x)^m,x)`output `int((c*(a + b*x^2))^(3/2)*(d*x)^m, x)`**Reduce [F]**

$$\int (dx)^m (c(a + bx^2))^{3/2} dx = d^m \sqrt{c} c \left(\left(\int x^m \sqrt{bx^2 + a} x^2 dx \right) b \right. \\ \left. + \left(\int x^m \sqrt{bx^2 + a} dx \right) a \right)$$

input `int((d*x)^m*(c*(b*x^2+a))^(3/2),x)`output `d**m*sqrt(c)*c*(int(x**m*sqrt(a + b*x**2)*x**2,x)*b + int(x**m*sqrt(a + b*x**2),x)*a)`

$$3.63 \quad \int (dx)^m \left(\frac{c}{a+bx^2} \right)^{3/2} dx$$

Optimal result	443
Mathematica [A] (verified)	443
Rubi [A] (verified)	444
Maple [F]	445
Fricas [F]	445
Sympy [F]	445
Maxima [F]	446
Giac [F]	446
Mupad [F(-1)]	446
Reduce [F]	447

Optimal result

Integrand size = 21, antiderivative size = 76

$$\int (dx)^m \left(\frac{c}{a+bx^2} \right)^{3/2} dx = \frac{c(dx)^{1+m} \sqrt{\frac{c}{a+bx^2}} \sqrt{1 + \frac{bx^2}{a}} \operatorname{Hypergeometric2F1} \left(\frac{3}{2}, \frac{1+m}{2}, \frac{3+m}{2}, -\frac{bx^2}{a} \right)}{ad(1+m)}$$

output

```
c*(d*x)^(1+m)*(c/(b*x^2+a))^(1/2)*(1+b*x^2/a)^(1/2)*hypergeom([3/2, 1/2+1/2*m], [3/2+1/2*m], -b*x^2/a)/a/d/(1+m)
```

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.95

$$\int (dx)^m \left(\frac{c}{a+bx^2} \right)^{3/2} dx = \frac{cx(dx)^m \sqrt{\frac{c}{a+bx^2}} \sqrt{1 + \frac{bx^2}{a}} \operatorname{Hypergeometric2F1} \left(\frac{3}{2}, \frac{1+m}{2}, \frac{3+m}{2}, -\frac{bx^2}{a} \right)}{a(1+m)}$$

input

```
Integrate[(d*x)^m*(c/(a + b*x^2))^(3/2),x]
```

output

```
(c*x*(d*x)^m*Sqrt[c/(a + b*x^2)]*Sqrt[1 + (b*x^2)/a]*Hypergeometric2F1[3/2, (1 + m)/2, (3 + m)/2, -((b*x^2)/a)]/(a*(1 + m))
```

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2045, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m \left(\frac{c}{a + bx^2} \right)^{3/2} dx$$

$$\downarrow \text{2045}$$

$$\frac{c \sqrt{\frac{bx^2}{a} + 1} \sqrt{\frac{c}{a + bx^2}} \int \frac{(dx)^m}{\left(\frac{bx^2}{a} + 1\right)^{3/2}} dx}{a}$$

$$\downarrow \text{278}$$

$$\frac{c \sqrt{\frac{bx^2}{a} + 1} (dx)^{m+1} \sqrt{\frac{c}{a + bx^2}} \text{Hypergeometric2F1} \left(\frac{3}{2}, \frac{m+1}{2}, \frac{m+3}{2}, -\frac{bx^2}{a} \right)}{ad(m+1)}$$

input `Int[(d*x)^m*(c/(a + b*x^2))^(3/2),x]`

output `(c*(d*x)^(1 + m)*Sqrt[c/(a + b*x^2)]*Sqrt[1 + (b*x^2)/a]*Hypergeometric2F1[3/2, (1 + m)/2, (3 + m)/2, -(b*x^2)/a])/(a*d*(1 + m))`

Defintions of rubi rules used

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [F]

$$\int (xd)^m \left(\frac{c}{bx^2 + a} \right)^{\frac{3}{2}} dx$$

input `int((x*d)^m*(c/(b*x^2+a))^(3/2),x)`

output `int((x*d)^m*(c/(b*x^2+a))^(3/2),x)`

Fricas [F]

$$\int (dx)^m \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \int (dx)^m \left(\frac{c}{bx^2 + a} \right)^{\frac{3}{2}} dx$$

input `integrate((d*x)^m*(c/(b*x^2+a))^(3/2),x, algorithm="fricas")`

output `integral((d*x)^m*c*sqrt(c/(b*x^2 + a))/(b*x^2 + a), x)`

Sympy [F]

$$\int (dx)^m \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \int \left(\frac{c}{a + bx^2} \right)^{\frac{3}{2}} (dx)^m dx$$

input `integrate((d*x)**m*(c/(b*x**2+a))**(3/2),x)`

output `Integral((c/(a + b*x**2))**(3/2)*(d*x)**m, x)`

Maxima [F]

$$\int (dx)^m \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \int (dx)^m \left(\frac{c}{bx^2 + a} \right)^{\frac{3}{2}} dx$$

input `integrate((d*x)^m*(c/(b*x^2+a))^(3/2),x, algorithm="maxima")`

output `integrate((d*x)^m*(c/(b*x^2 + a))^(3/2), x)`

Giac [F]

$$\int (dx)^m \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \int (dx)^m \left(\frac{c}{bx^2 + a} \right)^{\frac{3}{2}} dx$$

input `integrate((d*x)^m*(c/(b*x^2+a))^(3/2),x, algorithm="giac")`

output `integrate((d*x)^m*(c/(b*x^2 + a))^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int (dx)^m \left(\frac{c}{a + bx^2} \right)^{3/2} dx = \int (dx)^m \left(\frac{c}{bx^2 + a} \right)^{3/2} dx$$

input `int((d*x)^m*(c/(a + b*x^2))^(3/2),x)`

output `int((d*x)^m*(c/(a + b*x^2))^(3/2), x)`

Reduce [F]

$$\int (dx)^m \left(\frac{c}{a + bx^2} \right)^{3/2} dx = d^m \sqrt{c} \left(\int \frac{x^m}{\sqrt{bx^2 + a} a + \sqrt{bx^2 + a} bx^2} dx \right) c$$

input `int((d*x)^m*(c/(b*x^2+a))^(3/2),x)`

output `d**m*sqrt(c)*int(x**m/(sqrt(a + b*x**2)*a + sqrt(a + b*x**2)*b*x**2),x)*c`

3.64 $\int (dx)^m \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx$

Optimal result	448
Mathematica [A] (verified)	448
Rubi [A] (verified)	449
Maple [F]	450
Fricas [F]	450
Sympy [F(-1)]	451
Maxima [F]	451
Giac [F]	451
Mupad [F(-1)]	452
Reduce [F]	452

Optimal result

Integrand size = 21, antiderivative size = 67

$$\int (dx)^m \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx = \frac{c(dx)^{1+m} \sqrt{\frac{c}{(a+bx^2)^2}} (a+bx^2) \text{Hypergeometric2F1} \left(3, \frac{1+m}{2}, \frac{3+m}{2}, -\frac{bx^2}{a} \right)}{a^3 d(1+m)}$$

output

```
c*(d*x)^(1+m)*(c/(b*x^2+a)^2)^(1/2)*(b*x^2+a)*hypergeom([3, 1/2+1/2*m], [3/2+1/2*m], -b*x^2/a)/a^3/d/(1+m)
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.94

$$\int (dx)^m \left(\frac{c}{(a+bx^2)^2} \right)^{3/2} dx = \frac{cx(dx)^m \sqrt{\frac{c}{(a+bx^2)^2}} (a+bx^2) \text{Hypergeometric2F1} \left(3, \frac{1+m}{2}, \frac{3+m}{2}, -\frac{bx^2}{a} \right)}{a^3(1+m)}$$

input

```
Integrate[(d*x)^m*(c/(a + b*x^2)^2)^(3/2),x]
```

output

```
(c*x*(d*x)^m*Sqrt[c/(a + b*x^2)^2]*(a + b*x^2)*Hypergeometric2F1[3, (1 + m)/2, (3 + m)/2, -((b*x^2)/a)]/(a^3*(1 + m))
```

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2045, 27, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx$$

$$\downarrow \text{2045}$$

$$\frac{c(a + bx^2) \sqrt{\frac{c}{(a+bx^2)^2}} \int \frac{a^3(dx)^m}{(bx^2+a)^3} dx}{a^3}$$

$$\downarrow \text{27}$$

$$c(a + bx^2) \sqrt{\frac{c}{(a + bx^2)^2}} \int \frac{(dx)^m}{(bx^2 + a)^3} dx$$

$$\downarrow \text{278}$$

$$\frac{c(a + bx^2) (dx)^{m+1} \sqrt{\frac{c}{(a+bx^2)^2}} \text{Hypergeometric2F1} \left(3, \frac{m+1}{2}, \frac{m+3}{2}, -\frac{bx^2}{a} \right)}{a^3 d(m+1)}$$

input

```
Int[(d*x)^m*(c/(a + b*x^2)^2)^(3/2), x]
```

output

```
(c*(d*x)^(1 + m)*Sqrt[c/(a + b*x^2)^2]*(a + b*x^2)*Hypergeometric2F1[3, (1 + m)/2, (3 + m)/2, -((b*x^2)/a)]/(a^3*d*(1 + m))
```

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 278 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1))]*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 2045 `Int[(u_)*((c_)*((a_) + (b_)*(x_)^(n_))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [F]

$$\int (xd)^m \left(\frac{c}{(bx^2 + a)^2} \right)^{\frac{3}{2}} dx$$

input `int((x*d)^m*(c/(b*x^2+a)^2)^(3/2),x)`

output `int((x*d)^m*(c/(b*x^2+a)^2)^(3/2),x)`

Fricas [F]

$$\int (dx)^m \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \int (dx)^m \left(\frac{c}{(bx^2 + a)^2} \right)^{\frac{3}{2}} dx$$

input `integrate((d*x)^m*(c/(b*x^2+a)^2)^(3/2),x, algorithm="fricas")`

output `integral((d*x)^m*c*sqrt(c/(b^2*x^4 + 2*a*b*x^2 + a^2))/(b^2*x^4 + 2*a*b*x^2 + a^2), x)`

Sympy [F(-1)]

Timed out.

$$\int (dx)^m \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \text{Timed out}$$

input `integrate((d*x)**m*(c/(b*x**2+a)**2)**(3/2),x)`

output `Timed out`

Maxima [F]

$$\int (dx)^m \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \int (dx)^m \left(\frac{c}{(bx^2 + a)^2} \right)^{\frac{3}{2}} dx$$

input `integrate((d*x)^m*(c/(b*x^2+a)^2)^(3/2),x, algorithm="maxima")`

output `integrate((d*x)^m*(c/(b*x^2 + a)^2)^(3/2), x)`

Giac [F]

$$\int (dx)^m \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \int (dx)^m \left(\frac{c}{(bx^2 + a)^2} \right)^{\frac{3}{2}} dx$$

input `integrate((d*x)^m*(c/(b*x^2+a)^2)^(3/2),x, algorithm="giac")`

output `integrate((d*x)^m*(c/(b*x^2 + a)^2)^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int (dx)^m \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = \int (dx)^m \left(\frac{c}{(bx^2 + a)^2} \right)^{3/2} dx$$

input `int((d*x)^m*(c/(a + b*x^2)^2)^(3/2), x)`output `int((d*x)^m*(c/(a + b*x^2)^2)^(3/2), x)`**Reduce [F]**

$$\int (dx)^m \left(\frac{c}{(a + bx^2)^2} \right)^{3/2} dx = d^m \sqrt{c} \left(\int \frac{x^m}{b^3 x^6 + 3a b^2 x^4 + 3a^2 b x^2 + a^3} dx \right) c$$

input `int((d*x)^m*(c/(b*x^2+a)^2)^(3/2), x)`output `d**m*sqrt(c)*int(x**m/(a**3 + 3*a**2*b*x**2 + 3*a*b**2*x**4 + b**3*x**6), x)*c`

3.65 $\int (dx)^m \left(c(a + bx^n)^3 \right)^{3/2} dx$

Optimal result	453
Mathematica [A] (verified)	453
Rubi [A] (verified)	454
Maple [F]	455
Fricas [F(-2)]	455
Sympy [F(-1)]	456
Maxima [F]	456
Giac [F]	456
Mupad [F(-1)]	457
Reduce [F]	457

Optimal result

Integrand size = 21, antiderivative size = 77

$$\int (dx)^m (c(a + bx^n)^3)^{3/2} dx = \frac{a^3 c (dx)^{1+m} \sqrt{c(a + bx^n)^3} \operatorname{Hypergeometric2F1}\left(-\frac{9}{2}, \frac{1+m}{n}, \frac{1+m+n}{n}, -\frac{bx^n}{a}\right)}{d(1+m) \left(1 + \frac{bx^n}{a}\right)^{3/2}}$$

output

```
a^3*c*(d*x)^(1+m)*(c*(a+b*x^n)^3)^(1/2)*hypergeom([-9/2, (1+m)/n], [(1+m+n)/n], -b*x^n/a)/d/(1+m)/(1+b*x^n/a)^(3/2)
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.06

$$\int (dx)^m (c(a + bx^n)^3)^{3/2} dx = \frac{a^4 x (dx)^m (c(a + bx^n)^3)^{3/2} \operatorname{Hypergeometric2F1}\left(-\frac{9}{2}, \frac{1+m}{n}, 1 + \frac{1+m}{n}, -\frac{bx^n}{a}\right)}{(1+m) (a + bx^n)^4 \sqrt{1 + \frac{bx^n}{a}}}$$

input `Integrate[(d*x)^m*(c*(a + b*x^n)^3)^(3/2),x]`

output `(a^4*x*(d*x)^m*(c*(a + b*x^n)^3)^(3/2)*Hypergeometric2F1[-9/2, (1 + m)/n, 1 + (1 + m)/n, -((b*x^n)/a)]/((1 + m)*(a + b*x^n)^4*Sqrt[1 + (b*x^n)/a])`

Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2045, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m \left(c(a + bx^n)^3 \right)^{3/2} dx$$

$$\downarrow \text{2045}$$

$$\frac{a^3 c \sqrt{c(a + bx^n)^3} \int (dx)^m \left(\frac{bx^n}{a} + 1 \right)^{9/2} dx}{\left(\frac{bx^n}{a} + 1 \right)^{3/2}}$$

$$\downarrow \text{888}$$

$$\frac{a^3 c (dx)^{m+1} \sqrt{c(a + bx^n)^3} \text{Hypergeometric2F1} \left(-\frac{9}{2}, \frac{m+1}{n}, \frac{m+n+1}{n}, -\frac{bx^n}{a} \right)}{d(m+1) \left(\frac{bx^n}{a} + 1 \right)^{3/2}}$$

input `Int[(d*x)^m*(c*(a + b*x^n)^3)^(3/2),x]`

output `(a^3*c*(d*x)^(1 + m)*Sqrt[c*(a + b*x^n)^3]*Hypergeometric2F1[-9/2, (1 + m)/n, (1 + m + n)/n, -((b*x^n)/a)]/(d*(1 + m)*(1 + (b*x^n)/a)^(3/2))`

Definitions of rubi rules used

rule 888 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p * ((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [F]

$$\int (xd)^m (c(a + bx^n)^3)^{\frac{3}{2}} dx$$

input `int((x*d)^m*(c*(a+b*x^n)^3)^(3/2),x)`

output `int((x*d)^m*(c*(a+b*x^n)^3)^(3/2),x)`

Fricas [F(-2)]

Exception generated.

$$\int (dx)^m (c(a + bx^n)^3)^{3/2} dx = \text{Exception raised: TypeError}$$

input `integrate((d*x)^m*(c*(a+b*x^n)^3)^(3/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)`

Sympy [F(-1)]

Timed out.

$$\int (dx)^m (c(a + bx^n)^3)^{3/2} dx = \text{Timed out}$$

input `integrate((d*x)**m*(c*(a+b*x**n)**3)**(3/2),x)`

output `Timed out`

Maxima [F]

$$\int (dx)^m (c(a + bx^n)^3)^{3/2} dx = \int ((bx^n + a)^3 c)^{\frac{3}{2}} (dx)^m dx$$

input `integrate((d*x)^m*(c*(a+b*x^n)^3)^(3/2),x, algorithm="maxima")`

output `integrate(((b*x^n + a)^3*c)^(3/2)*(d*x)^m, x)`

Giac [F]

$$\int (dx)^m (c(a + bx^n)^3)^{3/2} dx = \int ((bx^n + a)^3 c)^{\frac{3}{2}} (dx)^m dx$$

input `integrate((d*x)^m*(c*(a+b*x^n)^3)^(3/2),x, algorithm="giac")`

output `integrate(((b*x^n + a)^3*c)^(3/2)*(d*x)^m, x)`

Mupad [F(-1)]

Timed out.

$$\int (dx)^m (c(a + bx^n)^3)^{3/2} dx = \int (dx)^m (c(a + bx^n)^3)^{3/2} dx$$

input `int((d*x)^m*(c*(a + b*x^n)^3)^(3/2), x)`output `int((d*x)^m*(c*(a + b*x^n)^3)^(3/2), x)`**Reduce [F]**

$$\int (dx)^m (c(a + bx^n)^3)^{3/2} dx = \text{too large to display}$$

input `int((d*x)^m*(c*(a+b*x^n)^3)^(3/2), x)`

output

```
(d**m*sqrt(c)*c*(32*x**(m + 4*n)*sqrt(x**n*b + a)*b**4*m**4*x + 256*x**(m
+ 4*n)*sqrt(x**n*b + a)*b**4*m**3*n*x + 128*x**(m + 4*n)*sqrt(x**n*b + a)*
b**4*m**3*x + 688*x**(m + 4*n)*sqrt(x**n*b + a)*b**4*m**2*n**2*x + 768*x**
(m + 4*n)*sqrt(x**n*b + a)*b**4*m**2*n*x + 192*x**(m + 4*n)*sqrt(x**n*b +
a)*b**4*m**2*x + 704*x**(m + 4*n)*sqrt(x**n*b + a)*b**4*m*n**3*x + 1376*x*
*(m + 4*n)*sqrt(x**n*b + a)*b**4*m*n**2*x + 768*x**(m + 4*n)*sqrt(x**n*b +
a)*b**4*m*n*x + 128*x**(m + 4*n)*sqrt(x**n*b + a)*b**4*m*x + 210*x**(m +
4*n)*sqrt(x**n*b + a)*b**4*n**4*x + 704*x**(m + 4*n)*sqrt(x**n*b + a)*b**4
*n**3*x + 688*x**(m + 4*n)*sqrt(x**n*b + a)*b**4*n**2*x + 256*x**(m + 4*n)
*sqrt(x**n*b + a)*b**4*n*x + 32*x**(m + 4*n)*sqrt(x**n*b + a)*b**4*x + 128
*x**(m + 3*n)*sqrt(x**n*b + a)*a*b**3*m**4*x + 1168*x**(m + 3*n)*sqrt(x**n
*b + a)*a*b**3*m**3*n*x + 512*x**(m + 3*n)*sqrt(x**n*b + a)*a*b**3*m**3*x
+ 3400*x**(m + 3*n)*sqrt(x**n*b + a)*a*b**3*m**2*n**2*x + 3504*x**(m + 3*n
)*sqrt(x**n*b + a)*a*b**3*m**2*n*x + 768*x**(m + 3*n)*sqrt(x**n*b + a)*a*b
**3*m**2*x + 3644*x**(m + 3*n)*sqrt(x**n*b + a)*a*b**3*m*n**3*x + 6800*x**
(m + 3*n)*sqrt(x**n*b + a)*a*b**3*m*n**2*x + 3504*x**(m + 3*n)*sqrt(x**n*b
+ a)*a*b**3*m*n*x + 512*x**(m + 3*n)*sqrt(x**n*b + a)*a*b**3*m*x + 1110*x
**(m + 3*n)*sqrt(x**n*b + a)*a*b**3*n**4*x + 3644*x**(m + 3*n)*sqrt(x**n*b
+ a)*a*b**3*n**3*x + 3400*x**(m + 3*n)*sqrt(x**n*b + a)*a*b**3*n**2*x + 1
168*x**(m + 3*n)*sqrt(x**n*b + a)*a*b**3*n*x + 128*x**(m + 3*n)*sqrt(x**...
```

3.66 $\int (dx)^m \left(c(a + bx^n)^2 \right)^{3/2} dx$

Optimal result	459
Mathematica [A] (verified)	460
Rubi [A] (verified)	460
Maple [F]	461
Fricas [A] (verification not implemented)	462
Sympy [F(-1)]	462
Maxima [A] (verification not implemented)	463
Giac [B] (verification not implemented)	463
Mupad [F(-1)]	464
Reduce [B] (verification not implemented)	465

Optimal result

Integrand size = 21, antiderivative size = 284

$$\int (dx)^m (c(a + bx^n)^2)^{3/2} dx = \frac{a^3 c^2 (dx)^{1+m} \sqrt{a^2 c + 2abcx^n + b^2 cx^{2n}}}{d(1+m)(ac + bcx^n)} + \frac{3a^2 b^2 c^2 x^n (dx)^{1+m} \sqrt{a^2 c + 2abcx^n + b^2 cx^{2n}}}{d(1+m+n)(abc + b^2 cx^n)} + \frac{3ab^3 c^2 x^{2n} (dx)^{1+m} \sqrt{a^2 c + 2abcx^n + b^2 cx^{2n}}}{d(1+m+2n)(abc + b^2 cx^n)} + \frac{b^4 c^2 x^{3n} (dx)^{1+m} \sqrt{a^2 c + 2abcx^n + b^2 cx^{2n}}}{d(1+m+3n)(abc + b^2 cx^n)}$$

output

```
a^3*c^2*(d*x)^(1+m)*(a^2*c+2*a*b*c*x^n+b^2*c*x^(2*n))^(1/2)/d/(1+m)/(a*c+b*c*x^n)+3*a^2*b^2*c^2*x^n*(d*x)^(1+m)*(a^2*c+2*a*b*c*x^n+b^2*c*x^(2*n))^(1/2)/d/(1+m+n)/(a*b*c+b^2*c*x^n)+3*a*b^3*c^2*x^(2*n)*(d*x)^(1+m)*(a^2*c+2*a*b*c*x^n+b^2*c*x^(2*n))^(1/2)/d/(1+m+2*n)/(a*b*c+b^2*c*x^n)+b^4*c^2*x^(3*n)*(d*x)^(1+m)*(a^2*c+2*a*b*c*x^n+b^2*c*x^(2*n))^(1/2)/d/(1+m+3*n)/(a*b*c+b^2*c*x^n)
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.32

$$\int (dx)^m (c(a + bx^n)^2)^{3/2} dx = \frac{x(dx)^m (c(a + bx^n)^2)^{3/2} \left(\frac{a^3}{1+m} + \frac{3a^2bx^n}{1+m+n} + \frac{3ab^2x^{2n}}{1+m+2n} + \frac{b^3x^{3n}}{1+m+3n} \right)}{(a + bx^n)^3}$$

input `Integrate[(d*x)^m*(c*(a + b*x^n)^2)^(3/2),x]`output `(x*(d*x)^m*(c*(a + b*x^n)^2)^(3/2)*(a^3/(1 + m) + (3*a^2*b*x^n)/(1 + m + n) + (3*a*b^2*x^(2*n))/(1 + m + 2*n) + (b^3*x^(3*n))/(1 + m + 3*n)))/(a + b*x^n)^3`**Rubi [A] (verified)**Time = 0.52 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.44, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2045, 802, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (dx)^m (c(a + bx^n)^2)^{3/2} dx \\ & \quad \downarrow \text{2045} \\ & \frac{a^2c\sqrt{c(a + bx^n)^2} \int (dx)^m \left(\frac{bx^n}{a} + 1\right)^3 dx}{\frac{bx^n}{a} + 1} \\ & \quad \downarrow \text{802} \\ & \frac{a^2c\sqrt{c(a + bx^n)^2} \int \left(\frac{3bx^n(dx)^m}{a} + \frac{3b^2x^{2n}(dx)^m}{a^2} + \frac{b^3x^{3n}(dx)^m}{a^3} + (dx)^m\right) dx}{\frac{bx^n}{a} + 1} \\ & \quad \downarrow \text{2009} \end{aligned}$$

$$\frac{a^2 c \sqrt{c(a + bx^n)^2} \left(\frac{b^3 x^{3n+1} (dx)^m}{a^3 (m+3n+1)} + \frac{3b^2 x^{2n+1} (dx)^m}{a^2 (m+2n+1)} + \frac{3bx^{n+1} (dx)^m}{a(m+n+1)} + \frac{(dx)^{m+1}}{d(m+1)} \right)}{\frac{bx^n}{a} + 1}$$

input `Int[(d*x)^m*(c*(a + b*x^n)^2)^(3/2),x]`

output `(a^2*c*Sqrt[c*(a + b*x^n)^2]*((3*b*x^(1 + n)*(d*x)^m)/(a*(1 + m + n)) + (3*b^2*x^(1 + 2*n)*(d*x)^m)/(a^2*(1 + m + 2*n)) + (b^3*x^(1 + 3*n)*(d*x)^m)/(a^3*(1 + m + 3*n)) + (d*x)^(1 + m)/(d*(1 + m))))/(1 + (b*x^n)/a)`

Defintions of rubi rules used

rule 802 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] := Int[Exp andIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_))^(q_))^(p_), x_Symbol] := Simp[Simp[(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple **[F]**

$$\int (xd)^m (c(a + bx^n)^2)^{\frac{3}{2}} dx$$

input `int((x*d)^m*(c*(a+b*x^n)^2)^(3/2),x)`

output `int((x*d)^m*(c*(a+b*x^n)^2)^(3/2),x)`

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 546, normalized size of antiderivative = 1.92

$$\int (dx)^m (c(a + bx^n)^2)^{3/2} dx = \frac{\sqrt{b^2cx^{2n} + 2abcx^n + a^2c}((b^3cm^3 + 3b^3cm^2 + 3b^3cm + b^3c + 2(b^3cm + b^3c)n^2 + 3(b^3cm^2 + 3b^3cm + b^3c)n + b^3c)n + a^2c)}{\dots}$$

input `integrate((d*x)^m*(c*(a+b*x^n)^2)^(3/2),x, algorithm="fricas")`

output

```
sqrt(b^2*c*x^(2*n) + 2*a*b*c*x^n + a^2*c)*((b^3*c*m^3 + 3*b^3*c*m^2 + 3*b^3*c*m + b^3*c + 2*(b^3*c*m + b^3*c)*n^2 + 3*(b^3*c*m^2 + 2*b^3*c*m + b^3*c)*n)*x*x^(3*n)*e^(m*log(d) + m*log(x)) + 3*(a*b^2*c*m^3 + 3*a*b^2*c*m^2 + 3*a*b^2*c*m + a*b^2*c + 3*(a*b^2*c*m + a*b^2*c)*n^2 + 4*(a*b^2*c*m^2 + 2*a*b^2*c*m + a*b^2*c)*n)*x*x^(2*n)*e^(m*log(d) + m*log(x)) + 3*(a^2*b*c*m^3 + 3*a^2*b*c*m^2 + 3*a^2*b*c*m + a^2*b*c + 6*(a^2*b*c*m + a^2*b*c)*n^2 + 5*(a^2*b*c*m^2 + 2*a^2*b*c*m + a^2*b*c)*n)*x*x^n*e^(m*log(d) + m*log(x)) + (a^3*c*m^3 + 6*a^3*c*n^3 + 3*a^3*c*m^2 + 3*a^3*c*m + a^3*c + 11*(a^3*c*m + a^3*c)*n^2 + 6*(a^3*c*m^2 + 2*a^3*c*m + a^3*c)*n)*x*e^(m*log(d) + m*log(x)))/(a*m^4 + 4*a*m^3 + 6*(a*m + a)*n^3 + 6*a*m^2 + 11*(a*m^2 + 2*a*m + a)*n^2 + 4*a*m + 6*(a*m^3 + 3*a*m^2 + 3*a*m + a)*n + (b*m^4 + 4*b*m^3 + 6*(b*m + b)*n^3 + 6*b*m^2 + 11*(b*m^2 + 2*b*m + b)*n^2 + 4*b*m + 6*(b*m^3 + 3*b*m^2 + 3*b*m + b)*n + b)*x^n + a)
```

Sympy [F(-1)]

Timed out.

$$\int (dx)^m (c(a + bx^n)^2)^{3/2} dx = \text{Timed out}$$

input `integrate((d*x)**m*(c*(a+b*x**n)**2)**(3/2),x)`

output

Timed out

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 288, normalized size of antiderivative = 1.01

$$\int (dx)^m (c(a + bx^n)^2)^{3/2} dx = \frac{(m^3 + 3m^2(2n + 1) + 6n^3 + (11n^2 + 12n + 3)m + 11n^2 + 6n + 1)a^3 c^{3/2} d^m x x^m + (m^3 + 3m^2(2n + 1) + 6n^3 + (11n^2 + 12n + 3)m + 11n^2 + 6n + 1)a^3 c^{3/2} d^m x x^m + (m^3 + 3m^2(2n + 1) + 6n^3 + (11n^2 + 12n + 3)m + 11n^2 + 6n + 1)a^3 c^{3/2} d^m x x^m + \dots}{(m^4 + 2m^3(3n + 2) + (11n^2 + 18n + 6)m^2 + 6n^3 + 2(3n^3 + 11n^2 + 9n + 2)m + 11n^2 + 6n + 1)}$$

input `integrate((d*x)^m*(c*(a+b*x^n)^2)^(3/2),x, algorithm="maxima")`

output `((m^3 + 3*m^2*(2*n + 1) + 6*n^3 + (11*n^2 + 12*n + 3)*m + 11*n^2 + 6*n + 1)*a^3*c^(3/2)*d^m*x*x^m + (m^3 + 3*m^2*(n + 1) + (2*n^2 + 6*n + 3)*m + 2*n^2 + 3*n + 1)*b^3*c^(3/2)*d^m*x*e^(m*log(x) + 3*n*log(x)) + 3*(m^3 + m^2*(4*n + 3) + (3*n^2 + 8*n + 3)*m + 3*n^2 + 4*n + 1)*a*b^2*c^(3/2)*d^m*x*e^(m*log(x) + 2*n*log(x)) + 3*(m^3 + m^2*(5*n + 3) + (6*n^2 + 10*n + 3)*m + 6*n^2 + 5*n + 1)*a^2*b*c^(3/2)*d^m*x*e^(m*log(x) + n*log(x)))/(m^4 + 2*m^3*(3*n + 2) + (11*n^2 + 18*n + 6)*m^2 + 6*n^3 + 2*(3*n^3 + 11*n^2 + 9*n + 2)*m + 11*n^2 + 6*n + 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2722 vs. 2(276) = 552.

Time = 0.24 (sec) , antiderivative size = 2722, normalized size of antiderivative = 9.58

$$\int (dx)^m (c(a + bx^n)^2)^{3/2} dx = \text{Too large to display}$$

input `integrate((d*x)^m*(c*(a+b*x^n)^2)^(3/2),x, algorithm="giac")`

output

```
(b^3*m^3*x*x^(3*n)*e^(m*log(d) + m*log(x))*sgn(b*x^n + a) + 3*b^3*m^2*n*x*
x^(3*n)*e^(m*log(d) + m*log(x))*sgn(b*x^n + a) + 2*b^3*m*n^2*x*x^(3*n)*e^(
m*log(d) + m*log(x))*sgn(b*x^n + a) + 3*a*b^2*m^3*x*x^(2*n)*e^(m*log(d) +
m*log(x))*sgn(b*x^n + a) + b^3*m^3*x*x^(2*n)*e^(m*log(d) + m*log(x))*sgn(b
*x^n + a) + 12*a*b^2*m^2*n*x*x^(2*n)*e^(m*log(d) + m*log(x))*sgn(b*x^n + a
) + 3*b^3*m^2*n*x*x^(2*n)*e^(m*log(d) + m*log(x))*sgn(b*x^n + a) + 9*a*b^2
*m*n^2*x*x^(2*n)*e^(m*log(d) + m*log(x))*sgn(b*x^n + a) + 2*b^3*m*n^2*x*x^(
2*n)*e^(m*log(d) + m*log(x))*sgn(b*x^n + a) + 3*a^2*b*m^3*x*x^n*e^(m*log(
d) + m*log(x))*sgn(b*x^n + a) + 3*a*b^2*m^3*x*x^n*e^(m*log(d) + m*log(x))*
sgn(b*x^n + a) + b^3*m^3*x*x^n*e^(m*log(d) + m*log(x))*sgn(b*x^n + a) + 15
*a^2*b*m^2*n*x*x^n*e^(m*log(d) + m*log(x))*sgn(b*x^n + a) + 12*a*b^2*m^2*n
*x*x^n*e^(m*log(d) + m*log(x))*sgn(b*x^n + a) + 3*b^3*m^2*n*x*x^n*e^(m*log
(d) + m*log(x))*sgn(b*x^n + a) + 18*a^2*b*m*n^2*x*x^n*e^(m*log(d) + m*log(
x))*sgn(b*x^n + a) + 9*a*b^2*m*n^2*x*x^n*e^(m*log(d) + m*log(x))*sgn(b*x^n
+ a) + 2*b^3*m*n^2*x*x^n*e^(m*log(d) + m*log(x))*sgn(b*x^n + a) + a^3*m^3
*x*e^(m*log(d) + m*log(x))*sgn(b*x^n + a) + 3*a^2*b*m^3*x*e^(m*log(d) + m*
log(x))*sgn(b*x^n + a) + 3*a*b^2*m^3*x*e^(m*log(d) + m*log(x))*sgn(b*x^n +
a) + b^3*m^3*x*e^(m*log(d) + m*log(x))*sgn(b*x^n + a) + 6*a^3*m^2*n*x*e^(
m*log(d) + m*log(x))*sgn(b*x^n + a) + 15*a^2*b*m^2*n*x*e^(m*log(d) + m*log
(x))*sgn(b*x^n + a) + 12*a*b^2*m^2*n*x*e^(m*log(d) + m*log(x))*sgn(b*x^...
```

Mupad [F(-1)]

Timed out.

$$\int (dx)^m (c(a + bx^n)^2)^{3/2} dx = \int (dx)^m (c(a + bx^n)^2)^{3/2} dx$$

input

```
int((d*x)^m*(c*(a + b*x^n)^2)^(3/2), x)
```

output

```
int((d*x)^m*(c*(a + b*x^n)^2)^(3/2), x)
```

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 483, normalized size of antiderivative = 1.70

$$\int (dx)^m (c(a + bx^n)^2)^{3/2} dx = \frac{x^m d^m \sqrt{c} cx(a^3 + 2x^{3n}b^3n^2 + 3x^{3n}b^3n + 3x^{2n}ab^2 + 3x^na^2b + 9x^{2n}ab^2n^2 + 12x^{2n}ab^2n + 1$$

input `int((d*x)^m*(c*(a+b*x^n)^2)^(3/2),x)`

output

```
(x**m*d**m*sqrt(c)*c*x*(x**(3*n)*b**3*m**3 + 3*x**(3*n)*b**3*m**2*n + 3*x**
*(3*n)*b**3*m**2 + 2*x**(3*n)*b**3*m*n**2 + 6*x**(3*n)*b**3*m*n + 3*x**(3*
n)*b**3*m + 2*x**(3*n)*b**3*n**2 + 3*x**(3*n)*b**3*n + x**(3*n)*b**3 + 3*x
**(2*n)*a*b**2*m**3 + 12*x**(2*n)*a*b**2*m**2*n + 9*x**(2*n)*a*b**2*m**2 +
9*x**(2*n)*a*b**2*m*n**2 + 24*x**(2*n)*a*b**2*m*n + 9*x**(2*n)*a*b**2*m +
9*x**(2*n)*a*b**2*n**2 + 12*x**(2*n)*a*b**2*n + 3*x**(2*n)*a*b**2 + 3*x**
n*a**2*b*m**3 + 15*x**n*a**2*b*m**2*n + 9*x**n*a**2*b*m**2 + 18*x**n*a**2*
b*m*n**2 + 30*x**n*a**2*b*m*n + 9*x**n*a**2*b*m + 18*x**n*a**2*b*n**2 + 15
*x**n*a**2*b*n + 3*x**n*a**2*b + a**3*m**3 + 6*a**3*m**2*n + 3*a**3*m**2 +
11*a**3*m*n**2 + 12*a**3*m*n + 3*a**3*m + 6*a**3*n**3 + 11*a**3*n**2 + 6*
a**3*n + a**3))/(m**4 + 6*m**3*n + 4*m**3 + 11*m**2*n**2 + 18*m**2*n + 6*m
**2 + 6*m*n**3 + 22*m*n**2 + 18*m*n + 4*m + 6*n**3 + 11*n**2 + 6*n + 1)
```

3.67 $\int (dx)^m (c(a + bx^n))^{3/2} dx$

Optimal result	466
Mathematica [A] (verified)	466
Rubi [A] (verified)	467
Maple [F]	468
Fricas [F(-2)]	468
Sympy [F(-1)]	469
Maxima [F]	469
Giac [F]	469
Mupad [F(-1)]	470
Reduce [F]	470

Optimal result

Integrand size = 19, antiderivative size = 74

$$\int (dx)^m (c(a + bx^n))^{3/2} dx = \frac{ac(dx)^{1+m} \sqrt{ac + bcx^n} \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, \frac{1+m}{n}, \frac{1+m+n}{n}, -\frac{bx^n}{a}\right)}{d(1+m) \sqrt{1 + \frac{bx^n}{a}}}$$

output

```
a*c*(d*x)^(1+m)*(a*c+b*c*x^n)^(1/2)*hypergeom([-3/2, (1+m)/n], [(1+m+n)/n], -b*x^n/a)/d/(1+m)/(1+b*x^n/a)^(1/2)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.93

$$\int (dx)^m (c(a + bx^n))^{3/2} dx = \frac{acx(dx)^m \sqrt{c(a + bx^n)} \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, \frac{1+m}{n}, \frac{1+m+n}{n}, -\frac{bx^n}{a}\right)}{(1+m) \sqrt{1 + \frac{bx^n}{a}}}$$

input

```
Integrate[(d*x)^m*(c*(a + b*x^n))^(3/2),x]
```

output

```
(a*c*x*(d*x)^m*Sqrt[c*(a + b*x^n)]*Hypergeometric2F1[-3/2, (1 + m)/n, (1 + m + n)/n, -((b*x^n)/a)]/((1 + m)*Sqrt[1 + (b*x^n)/a])
```

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2073, 889, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (dx)^m (c(a + bx^n))^{3/2} dx \\
 & \quad \downarrow \text{2073} \\
 & \int (dx)^m (ac + bcx^n)^{3/2} dx \\
 & \quad \downarrow \text{889} \\
 & \frac{ac\sqrt{ac + bcx^n} \int (dx)^m \left(\frac{bx^n}{a} + 1\right)^{3/2} dx}{\sqrt{\frac{bx^n}{a} + 1}} \\
 & \quad \downarrow \text{888} \\
 & \frac{ac(dx)^{m+1}\sqrt{ac + bcx^n} \text{Hypergeometric2F1}\left(-\frac{3}{2}, \frac{m+1}{n}, \frac{m+n+1}{n}, -\frac{bx^n}{a}\right)}{d(m+1)\sqrt{\frac{bx^n}{a} + 1}}
 \end{aligned}$$

input

```
Int[(d*x)^m*(c*(a + b*x^n))^(3/2),x]
```

output

```
(a*c*(d*x)^(1 + m)*Sqrt[a*c + b*c*x^n]*Hypergeometric2F1[-3/2, (1 + m)/n, (1 + m + n)/n, -((b*x^n)/a)]/(d*(1 + m)*Sqrt[1 + (b*x^n)/a])
```

Definitions of rubi rules used

rule 888 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p * ((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 889 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p]) Int[(c*x)^m*(1 + b*(x^n/a))^p, x], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && !(ILtQ[p, 0] || GtQ[a, 0])`

rule 2073 `Int[(u_)^(p_.)*((c_.)*(x_))^(m_.), x_Symbol] := Int[(c*x)^m*ExpandToSum[u, x]^p, x] /; FreeQ[{c, m, p}, x] && BinomialQ[u, x] && !BinomialMatchQ[u, x]`

Maple [F]

$$\int (xd)^m (c(a + bx^n))^{\frac{3}{2}} dx$$

input `int((x*d)^m*(c*(a+b*x^n))^(3/2),x)`

output `int((x*d)^m*(c*(a+b*x^n))^(3/2),x)`

Fricas [F(-2)]

Exception generated.

$$\int (dx)^m (c(a + bx^n))^{3/2} dx = \text{Exception raised: TypeError}$$

input `integrate((d*x)^m*(c*(a+b*x^n))^(3/2),x, algorithm="fricas")`

output Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)

Sympy [F(-1)]

Timed out.

$$\int (dx)^m (c(a + bx^n))^{3/2} dx = \text{Timed out}$$

input `integrate((d*x)**m*(c*(a+b*x**n))**(3/2),x)`

output Timed out

Maxima [F]

$$\int (dx)^m (c(a + bx^n))^{3/2} dx = \int ((bx^n + a)c)^{\frac{3}{2}} (dx)^m dx$$

input `integrate((d*x)^m*(c*(a+b*x^n))^(3/2),x, algorithm="maxima")`

output `integrate(((b*x^n + a)*c)^(3/2)*(d*x)^m, x)`

Giac [F]

$$\int (dx)^m (c(a + bx^n))^{3/2} dx = \int ((bx^n + a)c)^{\frac{3}{2}} (dx)^m dx$$

input `integrate((d*x)^m*(c*(a+b*x^n))^(3/2),x, algorithm="giac")`

output `integrate(((b*x^n + a)*c)^(3/2)*(d*x)^m, x)`

Mupad [F(-1)]

Timed out.

$$\int (dx)^m (c(a + bx^n))^{3/2} dx = \int (dx)^m (c(a + bx^n))^{3/2} dx$$

input `int((d*x)^m*(c*(a + b*x^n))^(3/2),x)`output `int((d*x)^m*(c*(a + b*x^n))^(3/2), x)`**Reduce [F]**

$$\int (dx)^m (c(a + bx^n))^{3/2} dx = \frac{d^m \sqrt{c} c \left(4x^{m+n} \sqrt{x^n b + a} b m x + 2x^{m+n} \sqrt{x^n b + a} b n x + 4x^{m+n} \sqrt{x^n b + a} b x + 4x^m \sqrt{x^n b + a} \right)}{...}$$

input `int((d*x)^m*(c*(a+b*x^n))^(3/2),x)`

output

```
(d**m*sqrt(c)*c*(4*x**(m + n)*sqrt(x**n*b + a)*b*m*x + 2*x**(m + n)*sqrt(x
**n*b + a)*b*n*x + 4*x**(m + n)*sqrt(x**n*b + a)*b*x + 4*x**m*sqrt(x**n*b
+ a)*a*m*x + 8*x**m*sqrt(x**n*b + a)*a*n*x + 4*x**m*sqrt(x**n*b + a)*a*x +
12*int((x**m*sqrt(x**n*b + a))/(4*x**n*b*m**2 + 8*x**n*b*m*n + 8*x**n*b*m
+ 3*x**n*b*n**2 + 8*x**n*b*n + 4*x**n*b + 4*a*m**2 + 8*a*m*n + 8*a*m + 3*
a*n**2 + 8*a*n + 4*a),x)*a**2*m**2*n**2 + 24*int((x**m*sqrt(x**n*b + a))/(
4*x**n*b*m**2 + 8*x**n*b*m*n + 8*x**n*b*m + 3*x**n*b*n**2 + 8*x**n*b*n + 4
*x**n*b + 4*a*m**2 + 8*a*m*n + 8*a*m + 3*a*n**2 + 8*a*n + 4*a),x)*a**2*m*n
**3 + 24*int((x**m*sqrt(x**n*b + a))/(4*x**n*b*m**2 + 8*x**n*b*m*n + 8*x**
n*b*m + 3*x**n*b*n**2 + 8*x**n*b*n + 4*x**n*b + 4*a*m**2 + 8*a*m*n + 8*a*m
+ 3*a*n**2 + 8*a*n + 4*a),x)*a**2*m*n**2 + 9*int((x**m*sqrt(x**n*b + a))/
(4*x**n*b*m**2 + 8*x**n*b*m*n + 8*x**n*b*m + 3*x**n*b*n**2 + 8*x**n*b*n +
4*x**n*b + 4*a*m**2 + 8*a*m*n + 8*a*m + 3*a*n**2 + 8*a*n + 4*a),x)*a**2*n*
*4 + 24*int((x**m*sqrt(x**n*b + a))/(4*x**n*b*m**2 + 8*x**n*b*m*n + 8*x**n
*b*m + 3*x**n*b*n**2 + 8*x**n*b*n + 4*x**n*b + 4*a*m**2 + 8*a*m*n + 8*a*m
+ 3*a*n**2 + 8*a*n + 4*a),x)*a**2*n**3 + 12*int((x**m*sqrt(x**n*b + a))/(4
*x**n*b*m**2 + 8*x**n*b*m*n + 8*x**n*b*m + 3*x**n*b*n**2 + 8*x**n*b*n + 4*
x**n*b + 4*a*m**2 + 8*a*m*n + 8*a*m + 3*a*n**2 + 8*a*n + 4*a),x)*a**2*n**2
))/(4*m**2 + 8*m*n + 8*m + 3*n**2 + 8*n + 4)
```


3.68 $\int (dx)^m \left(\frac{c}{a+bx^n}\right)^{3/2} dx$

Optimal result	472
Mathematica [A] (verified)	472
Rubi [A] (verified)	473
Maple [F]	474
Fricas [F(-2)]	474
Sympy [F]	474
Maxima [F]	475
Giac [F]	475
Mupad [F(-1)]	475
Reduce [F]	476

Optimal result

Integrand size = 21, antiderivative size = 77

$$\int (dx)^m \left(\frac{c}{a+bx^n}\right)^{3/2} dx = \frac{c(dx)^{1+m} \sqrt{\frac{c}{a+bx^n}} \sqrt{1 + \frac{bx^n}{a}} \operatorname{Hypergeometric2F1}\left(\frac{3}{2}, \frac{1+m}{n}, \frac{1+m+n}{n}, -\frac{bx^n}{a}\right)}{ad(1+m)}$$

output `c*(d*x)^(1+m)*(c/(a+b*x^n))^(1/2)*(1+b*x^n/a)^(1/2)*hypergeom([3/2, (1+m)/n], [(1+m+n)/n], -b*x^n/a)/a/d/(1+m)`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.94

$$\int (dx)^m \left(\frac{c}{a+bx^n}\right)^{3/2} dx = \frac{cx(dx)^m \sqrt{\frac{c}{a+bx^n}} \sqrt{1 + \frac{bx^n}{a}} \operatorname{Hypergeometric2F1}\left(\frac{3}{2}, \frac{1+m}{n}, \frac{1+m+n}{n}, -\frac{bx^n}{a}\right)}{a+am}$$

input `Integrate[(d*x)^m*(c/(a + b*x^n))^(3/2),x]`

output `(c*x*(d*x)^m*sqrt[c/(a + b*x^n)]*sqrt[1 + (b*x^n)/a]*Hypergeometric2F1[3/2, (1 + m)/n, (1 + m + n)/n, -((b*x^n)/a)]/(a + a*m)`

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2045, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m \left(\frac{c}{a + bx^n} \right)^{3/2} dx$$

↓ 2045

$$\frac{c \sqrt{\frac{bx^n}{a} + 1} \sqrt{\frac{c}{a + bx^n}} \int \frac{(dx)^m}{\left(\frac{bx^n}{a} + 1\right)^{3/2}} dx}{a}$$

↓ 888

$$\frac{c(dx)^{m+1} \sqrt{\frac{bx^n}{a} + 1} \sqrt{\frac{c}{a + bx^n}} \text{Hypergeometric2F1}\left(\frac{3}{2}, \frac{m+1}{n}, \frac{m+n+1}{n}, -\frac{bx^n}{a}\right)}{ad(m+1)}$$

input `Int[(d*x)^m*(c/(a + b*x^n))^(3/2),x]`

output `(c*(d*x)^(1 + m)*Sqrt[c/(a + b*x^n)]*Sqrt[1 + (b*x^n)/a]*Hypergeometric2F1[3/2, (1 + m)/n, (1 + m + n)/n, -(b*x^n)/a])/(a*d*(1 + m))`

Defintions of rubi rules used

rule 888 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p *((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 2045 `Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_.))^(q_))^(p_), x_Symbol] :> Simp[Simp[(c*(a + b*x^n)^q]^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]`

Maple [F]

$$\int (xd)^m \left(\frac{c}{a + bx^n} \right)^{\frac{3}{2}} dx$$

input `int((x*d)^m*(c/(a+b*x^n))^(3/2),x)`

output `int((x*d)^m*(c/(a+b*x^n))^(3/2),x)`

Fricas [F(-2)]

Exception generated.

$$\int (dx)^m \left(\frac{c}{a + bx^n} \right)^{3/2} dx = \text{Exception raised: TypeError}$$

input `integrate((d*x)^m*(c/(a+b*x^n))^(3/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

Sympy [F]

$$\int (dx)^m \left(\frac{c}{a + bx^n} \right)^{3/2} dx = \int \left(\frac{c}{a + bx^n} \right)^{\frac{3}{2}} (dx)^m dx$$

input `integrate((d*x)**m*(c/(a+b*x**n))**(3/2),x)`

output `Integral((c/(a + b*x**n))**(3/2)*(d*x)**m, x)`

Maxima [F]

$$\int (dx)^m \left(\frac{c}{a + bx^n} \right)^{3/2} dx = \int (dx)^m \left(\frac{c}{bx^n + a} \right)^{3/2} dx$$

input `integrate((d*x)^m*(c/(a+b*x^n))^(3/2),x, algorithm="maxima")`

output `integrate((d*x)^m*(c/(b*x^n + a))^(3/2), x)`

Giac [F]

$$\int (dx)^m \left(\frac{c}{a + bx^n} \right)^{3/2} dx = \int (dx)^m \left(\frac{c}{bx^n + a} \right)^{3/2} dx$$

input `integrate((d*x)^m*(c/(a+b*x^n))^(3/2),x, algorithm="giac")`

output `integrate((d*x)^m*(c/(b*x^n + a))^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int (dx)^m \left(\frac{c}{a + bx^n} \right)^{3/2} dx = \int (dx)^m \left(\frac{c}{a + bx^n} \right)^{3/2} dx$$

input `int((d*x)^m*(c/(a + b*x^n))^(3/2),x)`

output `int((d*x)^m*(c/(a + b*x^n))^(3/2), x)`

Reduce [F]

$$\int (dx)^m \left(\frac{c}{a + bx^n} \right)^{3/2} dx = d^m \sqrt{c} \left(\int \frac{x^m \sqrt{x^n b + a}}{x^{2n} b^2 + 2x^n a b + a^2} dx \right) c$$

input `int((d*x)^m*(c/(a+b*x^n))^(3/2),x)`

output `d**m*sqrt(c)*int((x**m*sqrt(x**n*b + a))/(x**(2*n)*b**2 + 2*x**n*a*b + a**2),x)*c`

3.69 $\int (dx)^m \left(\frac{c}{(a+bx^n)^2} \right)^{3/2} dx$

Optimal result	477
Mathematica [A] (verified)	477
Rubi [A] (verified)	478
Maple [F]	479
Fricas [F]	479
Sympy [F(-1)]	479
Maxima [F]	480
Giac [F]	480
Mupad [F(-1)]	480
Reduce [F]	481

Optimal result

Integrand size = 21, antiderivative size = 71

$$\int (dx)^m \left(\frac{c}{(a+bx^n)^2} \right)^{3/2} dx = \frac{c(dx)^{1+m} \sqrt{\frac{c}{(a+bx^n)^2}} \left(1 + \frac{bx^n}{a}\right) \text{Hypergeometric2F1}\left(3, \frac{1+m}{n}, \frac{1+m+n}{n}, -\frac{bx^n}{a}\right)}{a^2 d(1+m)}$$

output `c*(d*x)^(1+m)*(c/(a+b*x^n)^2)^(1/2)*(1+b*x^n/a)*hypergeom([3, (1+m)/n], [(1+m+n)/n], -b*x^n/a)/a^2/d/(1+m)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.93

$$\int (dx)^m \left(\frac{c}{(a+bx^n)^2} \right)^{3/2} dx = \frac{x(dx)^m \left(\frac{c}{(a+bx^n)^2} \right)^{3/2} (a+bx^n)^3 \text{Hypergeometric2F1}\left(3, \frac{1+m}{n}, 1 + \frac{1+m}{n}, -\frac{bx^n}{a}\right)}{a^3(1+m)}$$

input `Integrate[(d*x)^m*(c/(a + b*x^n)^2)^(3/2),x]`

output `(x*(d*x)^m*(c/(a + b*x^n)^2)^(3/2)*(a + b*x^n)^3*Hypergeometric2F1[3, (1 + m)/n, 1 + (1 + m)/n, -((b*x^n)/a)]/(a^3*(1 + m))`

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2045, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (dx)^m \left(\frac{c}{(a + bx^n)^2} \right)^{3/2} dx$$

↓ 2045

$$\frac{c \left(\frac{bx^n}{a} + 1 \right) \sqrt{\frac{c}{(a+bx^n)^2}} \int \frac{(dx)^m}{\left(\frac{bx^n}{a} + 1 \right)^3} dx}{a^2}$$

↓ 888

$$\frac{c(dx)^{m+1} \left(\frac{bx^n}{a} + 1 \right) \sqrt{\frac{c}{(a+bx^n)^2}} \text{Hypergeometric2F1} \left(3, \frac{m+1}{n}, \frac{m+n+1}{n}, -\frac{bx^n}{a} \right)}{a^2 d(m+1)}$$

input `Int[(d*x)^m*(c/(a + b*x^n)^2)^(3/2),x]`

output `(c*(d*x)^(1 + m)*Sqrt[c/(a + b*x^n)^2]*(1 + (b*x^n)/a)*Hypergeometric2F1[3, (1 + m)/n, (1 + m + n)/n, -(b*x^n)/a]]/(a^2*d*(1 + m))`

Defintions of rubi rules used

rule 888

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p
*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1
, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILt
Q[p, 0] || GtQ[a, 0])
```

rule 2045

```
Int[(u_.)*((c_.)*((a_) + (b_.)*(x_)^(n_))^(q_))^(p_), x_Symbol] :> Simp[Simp[
(c*(a + b*x^n)^q)^p/(1 + b*(x^n/a))^(p*q)] Int[u*(1 + b*(x^n/a))^(p*q)
, x], x] /; FreeQ[{a, b, c, n, p, q}, x] && !GeQ[a, 0]
```

Maple [F]

$$\int (xd)^m \left(\frac{c}{(a + bx^n)^2} \right)^{\frac{3}{2}} dx$$

input `int((x*d)^m*(c/(a+b*x^n)^2)^(3/2),x)`

output `int((x*d)^m*(c/(a+b*x^n)^2)^(3/2),x)`

Fricas [F]

$$\int (dx)^m \left(\frac{c}{(a + bx^n)^2} \right)^{3/2} dx = \int (dx)^m \left(\frac{c}{(bx^n + a)^2} \right)^{\frac{3}{2}} dx$$

input `integrate((d*x)^m*(c/(a+b*x^n)^2)^(3/2),x, algorithm="fricas")`

output `integral((d*x)^m*c*sqrt(c/(b^2*x^(2*n) + 2*a*b*x^n + a^2))/(b^2*x^(2*n) + 2*a*b*x^n + a^2), x)`

Sympy [F(-1)]

Timed out.

$$\int (dx)^m \left(\frac{c}{(a + bx^n)^2} \right)^{3/2} dx = \text{Timed out}$$

input `integrate((d*x)**m*(c/(a+b*x**n)**2)**(3/2),x)`

output `Timed out`

Maxima [F]

$$\int (dx)^m \left(\frac{c}{(a+bx^n)^2} \right)^{3/2} dx = \int (dx)^m \left(\frac{c}{(bx^n+a)^2} \right)^{3/2} dx$$

input `integrate((d*x)^m*(c/(a+b*x^n)^2)^(3/2),x, algorithm="maxima")`

output `(m^2 - m*(3*n - 2) + 2*n^2 - 3*n + 1)*c^(3/2)*d^m*integrate(1/2*x^m/(a^2*b*n^2*x^n + a^3*n^2), x) - 1/2*(a*c^(3/2)*d^m*(m - 3*n + 1)*x*x^m + b*c^(3/2)*d^m*(m - 2*n + 1)*x*e^(m*log(x) + n*log(x)))/(a^2*b^2*n^2*x^(2*n) + 2*a^3*b*n^2*x^n + a^4*n^2)`

Giac [F]

$$\int (dx)^m \left(\frac{c}{(a+bx^n)^2} \right)^{3/2} dx = \int (dx)^m \left(\frac{c}{(bx^n+a)^2} \right)^{3/2} dx$$

input `integrate((d*x)^m*(c/(a+b*x^n)^2)^(3/2),x, algorithm="giac")`

output `integrate((d*x)^m*(c/(b*x^n + a)^2)^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int (dx)^m \left(\frac{c}{(a+bx^n)^2} \right)^{3/2} dx = \int (dx)^m \left(\frac{c}{(a+bx^n)^2} \right)^{3/2} dx$$

input `int((d*x)^m*(c/(a + b*x^n)^2)^(3/2), x)`

output `int((d*x)^m*(c/(a + b*x^n)^2)^(3/2), x)`

Reduce [F]

$$\int (dx)^m \left(\frac{c}{(a + bx^n)^2} \right)^{3/2} dx = d^m \sqrt{c} \left(\int \frac{x^m}{x^{3n}b^3 + 3x^{2n}ab^2 + 3x^na^2b + a^3} dx \right) c$$

input `int((d*x)^m*(c/(a+b*x^n)^2)^(3/2),x)`

output `d**m*sqrt(c)*int(x**m/(x**(3*n)*b**3 + 3*x**(2*n)*a*b**2 + 3*x**n*a**2*b + a**3),x)*c`

CHAPTER 4

APPENDIX

4.1	Listing of Grading functions	482
4.2	Links to plain text integration problems used in this report for each CAS .	500

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
  ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "
  ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType, expn]],
  If[Head[expn]===Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]===Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
        1,
        Max[ExpnType[expn[[1]], 2]],
      Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]],
    If[Head[expn]===Plus || Head[expn]===Times,
      Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3, ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
    If[HypergeometricFunctionQ[Head[expn]],

```

```

    Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 5]],
    If [AppellFunctionQ [Head [expn]],
        Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 6]],
        If [Head [expn] === RootSum,
            Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 7]],
            If [Head [expn] === Integrate || Head [expn] === Int,
                Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 8]],
                9]]]]]]]]]]]]

```

```

ElementaryFunctionQ [func_] :=
  MemberQ [{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

```

```

SpecialFunctionQ [func_] :=
  MemberQ [{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

```

```

HypergeometricFunctionQ [func_] :=
  MemberQ [{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

```

```

AppellFunctionQ [func_] :=
  MemberQ [{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue
fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

#   antiderivative
#   "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```



```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
            fi;
        fi;
    else #ExpnType(result) > ExpnType(optimal)
        if debug then
            print("ExpnType(result) > ExpnType(optimal)");
        fi;
        return "C",cat("Result contains higher order function than in optimal. Order ",
                        convert(ExpnType_result,string)," vs. order ",
                        convert(ExpnType_optimal,string),".");
    fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#      Albert Rich to use with Sagemath. This is used to
#      grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#      'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#      issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```



```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file