

Computer Algebra Independent Integration Tests

Summer 2024

1-Algebraic-functions/1.2-Trinomial/1.2.3-General-trinomial/127-1.2.3.3-a

Nasser M. Abbasi

May 18, 2024 Compiled on May 18, 2024 at 9:37pm

Contents

1	Introduction	4
1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Time and leaf size Performance	10
1.4	Performance based on number of rules Rubi used	12
1.5	Performance based on number of steps Rubi used	13
1.6	Solved integrals histogram based on leaf size of result	14
1.7	Solved integrals histogram based on CPU time used	15
1.8	Leaf size vs. CPU time used	16
1.9	list of integrals with no known antiderivative	17
1.10	List of integrals solved by CAS but has no known antiderivative	17
1.11	list of integrals solved by CAS but failed verification	17
1.12	Timing	18
1.13	Verification	18
1.14	Important notes about some of the results	19
1.15	Current tree layout of integration tests	22
1.16	Design of the test system	23
2	detailed summary tables of results	24
2.1	List of integrals sorted by grade for each CAS	25
2.2	Detailed conclusion table per each integral for all CAS systems	29
2.3	Detailed conclusion table specific for Rubi results	44
3	Listing of integrals	47
3.1	$\int \frac{d+ex^3}{a+cx^6} dx$	49
3.2	$\int \frac{d+ex^3}{a-cx^6} dx$	61
3.3	$\int \frac{(d+ex^3)^{3/2}}{a+cx^6} dx$	74
3.4	$\int \frac{\sqrt{d+ex^3}}{a+cx^6} dx$	80
3.5	$\int \frac{1}{\sqrt{d+ex^3}(a+cx^6)} dx$	87
3.6	$\int \frac{1}{(d+ex^3)^{3/2}(a+cx^6)} dx$	94

3.7	$\int \frac{\sqrt{1-x^6}}{\sqrt{1-x^3}} dx$	101
3.8	$\int (d+ex^3)(a-cx^6)^p dx$	107
3.9	$\int (d+ex^3)(a+cx^6)^p dx$	113
3.10	$\int (d+ex^3)^q (a+cx^6)^p dx$	119
3.11	$\int (1-x^3)^p (1-x^6)^p dx$	124
3.12	$\int (d+ex^3)^q (d^2-e^2x^6)^p dx$	129
3.13	$\int (1-x^3)^{-p} (1-x^6)^p dx$	134
3.14	$\int \frac{d+ex^4}{a-cx^8} dx$	139
3.15	$\int \frac{d+ex^4}{a+cx^8} dx$	152
3.16	$\int \frac{d+ex^4}{\sqrt{1-\frac{e^2x^8}{d^2}}} dx$	166
3.17	$\int \frac{d+ex^4}{\sqrt{d^2-e^2x^8}} dx$	172
3.18	$\int \frac{d+ex^4}{\sqrt{a-cx^8}} dx$	177
3.19	$\int \frac{d+ex^4}{\sqrt{a+cx^8}} dx$	182
3.20	$\int (d+ex^4)(a-cx^8)^p dx$	188
3.21	$\int (d+ex^4)(a+cx^8)^p dx$	194
3.22	$\int (d+ex^4)^q (a+cx^8)^p dx$	200
3.23	$\int (1-x^4)^p (1-x^8)^p dx$	205
3.24	$\int (d+ex^4)^q (d^2-e^2x^8)^p dx$	210
3.25	$\int (1-x^4)^{-p} (1-x^8)^p dx$	215
3.26	$\int \frac{(d+ex^n)^3}{a+cx^{2n}} dx$	220
3.27	$\int \frac{(d+ex^n)^2}{a+cx^{2n}} dx$	226
3.28	$\int \frac{d+ex^n}{a+cx^{2n}} dx$	231
3.29	$\int \frac{1}{(d+ex^n)(a+cx^{2n})} dx$	236
3.30	$\int \frac{1}{(d+ex^n)^2(a+cx^{2n})} dx$	241
3.31	$\int \frac{d+ex^n}{a-cx^{2n}} dx$	246
3.32	$\int \frac{(d+ex^n)^3}{(a+cx^{2n})^2} dx$	251
3.33	$\int \frac{(d+ex^n)^2}{(a+cx^{2n})^2} dx$	257
3.34	$\int \frac{d+ex^n}{(a+cx^{2n})^2} dx$	263
3.35	$\int \frac{1}{(d+ex^n)(a+cx^{2n})^2} dx$	269
3.36	$\int \frac{1}{(d+ex^n)^2(a+cx^{2n})^2} dx$	275
3.37	$\int \frac{(d+ex^n)^3}{(a+cx^{2n})^3} dx$	281
3.38	$\int \frac{(d+ex^n)^2}{(a+cx^{2n})^3} dx$	287
3.39	$\int \frac{d+ex^n}{(a+cx^{2n})^3} dx$	293
3.40	$\int \frac{1}{(d+ex^n)(a+cx^{2n})^3} dx$	299
3.41	$\int \frac{1}{(d+ex^n)^2(a+cx^{2n})^3} dx$	306

3.42	$\int \frac{(d+ex^n)^{3/2}}{a+cx^{2n}} dx$	313
3.43	$\int \frac{\sqrt{d+ex^n}}{a+cx^{2n}} dx$	318
3.44	$\int \frac{1}{\sqrt{d+ex^n}(a+cx^{2n})} dx$	323
3.45	$\int \frac{1}{(d+ex^n)^{3/2}(a+cx^{2n})} dx$	328
3.46	$\int \frac{1}{(d+ex^n)\sqrt{a+cx^{2n}}} dx$	334
3.47	$\int (d+ex^n)^3 (a+cx^{2n})^p dx$	339
3.48	$\int (d+ex^n)^2 (a+cx^{2n})^p dx$	345
3.49	$\int (d+ex^n) (a+cx^{2n})^p dx$	351
3.50	$\int \frac{(a+cx^{2n})^p}{d+ex^n} dx$	357
3.51	$\int \frac{(a+cx^{2n})^p}{(d+ex^n)^2} dx$	362
3.52	$\int \frac{(a+cx^{2n})^p}{(d+ex^n)^3} dx$	367
3.53	$\int (d+ex^n)^q (a+cx^{2n})^p dx$	372
3.54	$\int (d+ex^n)^q (cd^2 - ce^2 x^{2n})^p dx$	377
3.55	$\int (2-ex^n)^p (2+ex^n)^{p+q} dx$	382
3.56	$\int (2+ex^n)^q (4-e^2 x^{2n})^p dx$	387
4	Appendix	392
4.1	Listing of Grading functions	392
4.2	Links to plain text integration problems used in this report for each CAS410	

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Time and leaf size Performance	10
1.4	Performance based on number of rules Rubi used	12
1.5	Performance based on number of steps Rubi used	13
1.6	Solved integrals histogram based on leaf size of result	14
1.7	Solved integrals histogram based on CPU time used	15
1.8	Leaf size vs. CPU time used	16
1.9	list of integrals with no known antiderivative	17
1.10	List of integrals solved by CAS but has no known antiderivative	17
1.11	list of integrals solved by CAS but failed verification	17
1.12	Timing	18
1.13	Verification	18
1.14	Important notes about some of the results	19
1.15	Current tree layout of integration tests	22
1.16	Design of the test system	23

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [56]. This is test number [127].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (56)	0.00 (0)
Mathematica	67.86 (38)	32.14 (18)
Sympy	28.57 (16)	71.43 (40)
Maple	23.21 (13)	76.79 (43)
Fricas	14.29 (8)	85.71 (48)
Mupad	12.50 (7)	87.50 (49)
Giac	12.50 (7)	87.50 (49)
Reduce	12.50 (7)	87.50 (49)
Maxima	8.93 (5)	91.07 (51)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	85.714	3.571	5.357	5.357
Mathematica	53.571	1.786	7.143	37.500
Giac	5.357	1.786	0.000	92.857
Maxima	3.571	0.000	0.000	96.429
Sympy	3.571	0.000	25.000	71.429
Maple	1.786	1.786	14.286	82.143
Fricas	1.786	7.143	0.000	91.071
Mupad	0.000	7.143	0.000	92.857
Reduce	0.000	7.143	0.000	92.857

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sageMath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	18	100.00	0.00	0.00
Sympy	40	37.50	55.00	7.50
Maple	43	100.00	0.00	0.00
Fricas	48	91.67	8.33	0.00
Mupad	49	0.00	100.00	0.00
Giac	49	95.92	0.00	4.08
Reduce	49	100.00	0.00	0.00
Maxima	51	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.09
Fricas	0.22
Giac	0.24
Rubi	0.39
Maple	0.46
Reduce	1.61
Mathematica	3.02
Mupad	9.92
Sympy	44.73

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Maxima	132.00	1.04	23.00	1.10
Rubi	202.59	1.17	143.00	1.00
Sympy	204.88	1.69	113.00	0.80
Giac	267.00	1.22	283.00	1.10
Maple	269.38	2.11	36.00	1.00
Mathematica	483.29	3.20	133.50	0.86
Reduce	761.43	25.09	820.00	1.79
Mupad	1091.00	3.61	1293.00	4.00
Fricas	1105.00	3.65	827.00	2.93

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.





Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the precentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.



1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{10, 22, 53}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {6, 16, 17, 18, 19}

Mathematica {4, 5, 6, 55}

Maple {3, 4, 5, 6}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each `integrate` call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int', int(expr,x)),output='realtime'
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A `time limit` of 3 CPU minutes was used for each integral. If the `integrate` command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError.`

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    """
    Return the tree size of this expression.
    """

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	25
2.2	Detailed conclusion table per each integral for all CAS systems	29
2.3	Detailed conclusion table specific for Rubi results	44

2.1 List of integrals sorted by grade for each CAS

Rubi	25
Mma	25
Maple	26
Fricas	26
Maxima	26
Giac	27
Mupad	27
Sympy	27
Reduce	28

Rubi

A grade { 1, 2, 3, 4, 5, 7, 8, 9, 11, 12, 13, 14, 15, 20, 21, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 54, 55, 56 }

B grade { 6, 18 }

C grade { 16, 17, 19 }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 8, 9, 14, 15, 16, 17, 18, 20, 21, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 47, 48, 49 }

B grade { 55 }

C grade { 4, 5, 6, 19 }

F normal fail { 3, 7, 11, 12, 13, 23, 24, 25, 42, 43, 44, 45, 46, 50, 51, 52, 54, 56 }

F(-1) timeout fail { }

F(-2) exception fail { }

Maple**A grade** { 16 }**B grade** { 7 }**C grade** { 1, 2, 3, 4, 5, 6, 14, 15 }**F normal fail** { 8, 9, 11, 12, 13, 17, 18, 19, 20, 21, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 54, 55, 56 }**F(-1) timeout fail** { }**F(-2) exception fail** { }**Fricas****A grade** { 7 }**B grade** { 1, 2, 14, 15 }**C grade** { }**F normal fail** { 8, 9, 11, 12, 13, 16, 17, 18, 19, 20, 21, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 54, 55, 56 }**F(-1) timeout fail** { 3, 4, 5, 6 }**F(-2) exception fail** { }**Maxima****A grade** { 1, 2 }**B grade** { }**C grade** { }**F normal fail** { 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 54, 55, 56 }**F(-1) timeout fail** { }**F(-2) exception fail** { }

Giac

A grade { 1, 2, 15 }

B grade { 14 }

C grade { }

F normal fail { 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 16, 17, 18, 19, 20, 21, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 49, 50, 51, 52, 54, 55, 56 }

F(-1) timeout fail { }

F(-2) exception fail { 47, 48 }

Mupad

A grade { }

B grade { 1, 2, 14, 15 }

C grade { }

F normal fail { }

F(-1) timeout fail { 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 16, 17, 18, 19, 20, 21, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 54, 55, 56 }

F(-2) exception fail { }

Sympy

A grade { 1, 2 }

B grade { }

C grade { 8, 9, 16, 17, 18, 19, 20, 21, 26, 27, 28, 31, 34, 49 }

F normal fail { 3, 4, 5, 6, 7, 11, 23, 32, 33, 42, 43, 44, 45, 46, 55 }

F(-1) timeout fail { 10, 12, 13, 14, 15, 22, 24, 25, 35, 36, 37, 38, 39, 40, 41, 47, 48, 51, 52, 53, 54, 56 }

F(-2) exception fail { 29, 30, 50 }

Reduce

A grade { }

B grade { 1, 2, 14, 15 }

C grade { }

F normal fail { 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 16, 17, 18, 19, 20, 21, 23, 24, 25, 26, 27, 28, 29,
30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 54, 55,
56 }

F(-1) timeout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	305	352	334	34	282	1631	165	283	276	1331
N.S.	1	1.15	1.10	0.11	0.92	5.35	0.54	0.93	0.90	4.36
time (sec)	N/A	0.649	0.109	0.148	0.115	0.144	1.918	0.129	0.253	0.946

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	323	279	337	36	313	1613	168	303	328	1293
N.S.	1	0.86	1.04	0.11	0.97	4.99	0.52	0.94	1.02	4.00
time (sec)	N/A	0.456	0.117	0.139	0.122	0.148	1.853	0.143	0.250	11.900

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	C	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	N/A	No	TBD	TBD	TBD	TBD	TBD	TBD
size	142	142	0	902	0	0	0	0	48	0
N.S.	1	1.00	0.00	6.35	0.00	0.00	0.00	0.00	0.34	0.00
time (sec)	N/A	0.312	0.000	0.722	0.000	0.000	0.000	0.000	0.734	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	140	140	5647	605	0	0	0	0	20	0
N.S.	1	1.00	40.34	4.32	0.00	0.00	0.00	0.00	0.14	0.00
time (sec)	N/A	0.291	21.193	0.346	0.000	0.000	0.000	0.000	0.295	0.000

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	140	140	932	595	0	0	0	0	35	0
N.S.	1	1.00	6.66	4.25	0.00	0.00	0.00	0.00	0.25	0.00
time (sec)	N/A	0.295	19.887	0.322	0.000	0.000	0.000	0.000	0.246	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	C	C	F	F(-1)	F	F	F	F(-1)
verified	N/A	No	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	146	438	5979	937	0	0	0	0	59	0
N.S.	1	3.00	40.95	6.42	0.00	0.00	0.00	0.00	0.40	0.00
time (sec)	N/A	0.977	21.281	0.365	0.000	0.000	0.000	0.000	0.337	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	B	F	A	F	F	F	F(-1)
verified	N/A	Yes	N/A	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	120	120	0	221	0	41	0	0	28	0
N.S.	1	1.00	0.00	1.84	0.00	0.34	0.00	0.00	0.23	0.00
time (sec)	N/A	0.218	0.000	0.267	0.000	0.070	0.000	0.000	0.202	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	75	0	0	0	78	0	547	0
N.S.	1	1.00	0.77	0.00	0.00	0.00	0.80	0.00	5.58	0.00
time (sec)	N/A	0.252	0.572	0.000	0.000	0.000	39.398	0.000	0.311	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	96	96	75	0	0	0	75	0	535	0
N.S.	1	1.00	0.78	0.00	0.00	0.00	0.78	0.00	5.57	0.00
time (sec)	N/A	0.240	0.567	0.000	0.000	0.000	39.303	0.000	0.320	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	21	19	21	21	0	21	1249	21
N.S.	1	1.00	1.11	1.00	1.11	1.11	0.00	1.11	65.74	1.11
time (sec)	N/A	0.159	0.208	0.066	0.069	0.177	0.000	0.306	0.855	10.794

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F(-1)						
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	0	0	0	0	0	0	199	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	8.65	0.00
time (sec)	N/A	0.168	0.000	0.000	0.000	0.000	0.000	0.000	0.205	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	0	0	0	0	0	0	659	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	7.16	0.00
time (sec)	N/A	0.261	0.000	0.000	0.000	0.000	0.000	0.000	0.263	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	0	0	0	0	0	0	23	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	1.35	0.00
time (sec)	N/A	0.157	0.000	0.000	0.000	0.000	0.000	0.000	0.201	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	263	293	425	36	0	2741	0	625	375	2438
N.S.	1	1.11	1.62	0.14	0.00	10.42	0.00	2.38	1.43	9.27
time (sec)	N/A	0.531	0.201	0.082	0.000	0.387	0.000	0.170	0.205	11.464

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	578	849	534	34	0	2749	0	593	1033	2510
N.S.	1	1.47	0.92	0.06	0.00	4.76	0.00	1.03	1.79	4.34
time (sec)	N/A	1.501	0.690	0.089	0.000	0.368	0.000	0.173	0.221	11.778

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	F	F	C	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	52	85	52	43	0	0	78	0	28	0
N.S.	1	1.63	1.00	0.83	0.00	0.00	1.50	0.00	0.54	0.00
time (sec)	N/A	0.263	10.034	3.301	0.000	0.000	1.051	0.000	0.216	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	F	F	F	C	F	F	F(-1)
verified	N/A	No	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	118	84	87	0	0	0	78	0	26	0
N.S.	1	0.71	0.74	0.00	0.00	0.00	0.66	0.00	0.22	0.00
time (sec)	N/A	0.252	10.056	0.000	0.000	0.000	1.081	0.000	0.206	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	A	F	F	F	C	F	F	F(-1)
verified	N/A	No	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	102	482	77	0	0	0	82	0	52	0
N.S.	1	4.73	0.75	0.00	0.00	0.00	0.80	0.00	0.51	0.00
time (sec)	N/A	0.617	10.053	0.000	0.000	0.000	0.958	0.000	0.274	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	C	F	F	F	C	F	F	F(-1)
verified	N/A	No	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	905	433	77	0	0	0	78	0	48	0
N.S.	1	0.48	0.09	0.00	0.00	0.00	0.09	0.00	0.05	0.00
time (sec)	N/A	0.493	10.055	0.000	0.000	0.000	1.004	0.000	0.384	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	75	0	0	0	78	0	547	0
N.S.	1	1.00	0.77	0.00	0.00	0.00	0.80	0.00	5.58	0.00
time (sec)	N/A	0.233	0.569	0.000	0.000	0.000	150.529	0.000	0.338	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	96	96	75	0	0	0	75	0	535	0
N.S.	1	1.00	0.78	0.00	0.00	0.00	0.78	0.00	5.57	0.00
time (sec)	N/A	0.227	0.572	0.000	0.000	0.000	154.484	0.000	0.437	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	21	19	21	21	0	21	1249	21
N.S.	1	1.00	1.11	1.00	1.11	1.11	0.00	1.11	65.74	1.11
time (sec)	N/A	0.157	0.205	0.071	0.082	0.392	0.000	0.607	9.236	11.175

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	0	0	0	0	0	0	199	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	8.65	0.00
time (sec)	N/A	0.163	0.000	0.000	0.000	0.000	0.000	0.000	0.249	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	0	0	0	0	0	0	659	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	7.16	0.00
time (sec)	N/A	0.262	0.000	0.000	0.000	0.000	0.000	0.000	0.297	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	0	0	0	0	0	0	23	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	1.35	0.00
time (sec)	N/A	0.155	0.000	0.000	0.000	0.000	0.000	0.000	0.227	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	141	141	127	0	0	0	466	0	207	0
N.S.	1	1.00	0.90	0.00	0.00	0.00	3.30	0.00	1.47	0.00
time (sec)	N/A	0.327	0.739	0.000	0.000	0.000	4.490	0.000	0.238	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	107	107	107	0	0	0	296	0	62	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	2.77	0.00	0.58	0.00
time (sec)	N/A	0.281	0.304	0.000	0.000	0.000	3.099	0.000	0.232	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	83	83	83	0	0	0	209	0	35	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	2.52	0.00	0.42	0.00
time (sec)	N/A	0.214	0.092	0.000	0.000	0.000	2.367	0.000	0.241	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	152	152	131	0	0	0	0	0	30	0
N.S.	1	1.00	0.86	0.00	0.00	0.00	0.00	0.00	0.20	0.00
time (sec)	N/A	0.305	0.232	0.000	0.000	0.000	0.000	0.000	0.238	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	224	205	186	0	0	0	0	0	58	0
N.S.	1	0.92	0.83	0.00	0.00	0.00	0.00	0.00	0.26	0.00
time (sec)	N/A	0.394	0.361	0.000	0.000	0.000	0.000	0.000	0.250	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	81	0	0	0	214	0	41	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	2.64	0.00	0.51	0.00
time (sec)	N/A	0.220	0.116	0.000	0.000	0.000	2.397	0.000	0.268	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	210	288	188	0	0	0	0	0	1562	0
N.S.	1	1.37	0.90	0.00	0.00	0.00	0.00	0.00	7.44	0.00
time (sec)	N/A	0.462	0.503	0.000	0.000	0.000	0.000	0.000	0.227	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	166	203	136	0	0	0	0	0	351	0
N.S.	1	1.22	0.82	0.00	0.00	0.00	0.00	0.00	2.11	0.00
time (sec)	N/A	0.373	0.288	0.000	0.000	0.000	0.000	0.000	0.212	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	134	133	83	0	0	0	994	0	61	0
N.S.	1	0.99	0.62	0.00	0.00	0.00	7.42	0.00	0.46	0.00
time (sec)	N/A	0.281	0.096	0.000	0.000	0.000	173.594	0.000	0.197	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	249	333	227	0	0	0	0	0	58	0
N.S.	1	1.34	0.91	0.00	0.00	0.00	0.00	0.00	0.23	0.00
time (sec)	N/A	0.489	0.367	0.000	0.000	0.000	0.000	0.000	0.215	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	347	410	298	0	0	0	0	0	105	0
N.S.	1	1.18	0.86	0.00	0.00	0.00	0.00	0.00	0.30	0.00
time (sec)	N/A	0.632	0.652	0.000	0.000	0.000	0.000	0.000	0.210	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	259	424	188	0	0	0	0	0	0	0
N.S.	1	1.64	0.73	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.613	0.896	0.000	0.000	0.000	0.000	0.000	0.234	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	221	272	136	0	0	0	0	0	710	0
N.S.	1	1.23	0.62	0.00	0.00	0.00	0.00	0.00	3.21	0.00
time (sec)	N/A	0.456	0.530	0.000	0.000	0.000	0.000	0.000	0.222	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	184	194	83	0	0	0	0	0	87	0
N.S.	1	1.05	0.45	0.00	0.00	0.00	0.00	0.00	0.47	0.00
time (sec)	N/A	0.367	0.112	0.000	0.000	0.000	0.000	0.000	0.222	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	462	582	346	0	0	0	0	0	86	0
N.S.	1	1.26	0.75	0.00	0.00	0.00	0.00	0.00	0.19	0.00
time (sec)	N/A	0.753	0.574	0.000	0.000	0.000	0.000	0.000	0.213	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	660	701	426	0	0	0	0	0	152	0
N.S.	1	1.06	0.65	0.00	0.00	0.00	0.00	0.00	0.23	0.00
time (sec)	N/A	0.998	1.017	0.000	0.000	0.000	0.000	0.000	0.226	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	146	146	0	0	0	0	0	0	52	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.36	0.00
time (sec)	N/A	0.339	0.000	0.000	0.000	0.000	0.000	0.000	0.257	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	144	144	0	0	0	0	0	0	22	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.15	0.00
time (sec)	N/A	0.317	0.000	0.000	0.000	0.000	0.000	0.000	0.203	0.000

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	144	144	0	0	0	0	0	0	39	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.27	0.00
time (sec)	N/A	0.317	0.000	0.000	0.000	0.000	0.000	0.000	0.195	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	150	260	0	0	0	0	0	0	67	0
N.S.	1	1.73	0.00	0.00	0.00	0.00	0.00	0.00	0.45	0.00
time (sec)	N/A	0.976	0.000	0.000	0.000	0.000	0.000	0.000	0.309	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	171	171	0	0	0	0	0	0	41	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.24	0.00
time (sec)	N/A	0.362	0.000	0.000	0.000	0.000	0.000	0.000	0.194	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	263	299	213	0	0	0	0	0	0	0
N.S.	1	1.14	0.81	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.416	0.350	0.000	0.000	0.000	0.000	0.000	0.342	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	201	217	171	0	0	0	0	0	0	0
N.S.	1	1.08	0.85	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.342	0.209	0.000	0.000	0.000	0.000	0.000	0.239	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	135	135	110	0	0	0	144	0	0	0
N.S.	1	1.00	0.81	0.00	0.00	0.00	1.07	0.00	0.00	0.00
time (sec)	N/A	0.268	0.117	0.000	0.000	0.000	138.224	0.000	0.218	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	167	167	0	0	0	0	0	0	23	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.14	0.00
time (sec)	N/A	0.346	0.000	0.000	0.000	0.000	0.000	0.000	0.205	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	261	261	0	0	0	0	0	0	36	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.14	0.00
time (sec)	N/A	0.473	0.000	0.000	0.000	0.000	0.000	0.000	0.216	0.000

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	357	357	0	0	0	0	0	0	49	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.14	0.00
time (sec)	N/A	0.608	0.000	0.000	0.000	0.000	0.000	0.000	0.223	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	23	21	23	23	0	23	820	23
N.S.	1	1.00	1.10	1.00	1.10	1.10	0.00	1.10	39.05	1.10
time (sec)	N/A	0.162	0.383	0.093	0.084	0.081	0.000	0.181	0.262	11.377

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	99	99	0	0	0	0	0	0	31	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.31	0.00
time (sec)	N/A	0.294	0.000	0.000	0.000	0.000	0.000	0.000	0.201	0.000

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	44	44	199	0	0	0	0	0	24	0
N.S.	1	1.00	4.52	0.00	0.00	0.00	0.00	0.00	0.55	0.00
time (sec)	N/A	0.176	0.398	0.000	0.000	0.000	0.000	0.000	0.216	0.000

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	44	44	0	0	0	0	0	0	26	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.59	0.00
time (sec)	N/A	0.185	0.000	0.000	0.000	0.000	0.000	0.000	0.216	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [14] had the largest ratio of [.72222200000000031]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	12	11	1.15	17	0.647
2	A	11	10	0.86	18	0.556
3	A	4	4	1.00	21	0.190
4	A	4	4	1.00	21	0.190
5	A	4	4	1.00	21	0.190
6	B	5	5	3.00	21	0.238
7	A	3	3	1.00	23	0.130
8	A	2	2	1.00	18	0.111
9	A	2	2	1.00	17	0.118
10	N/A	1	0	1.00	19	0.000
11	A	2	2	1.00	19	0.105
12	A	4	4	1.00	24	0.167
13	A	2	2	1.00	21	0.095
14	A	14	13	1.11	18	0.722
15	A	12	11	1.47	17	0.647
16	C	4	4	1.63	25	0.160
17	C	4	4	0.71	24	0.167
18	B	2	2	4.73	20	0.100
19	C	2	2	0.48	19	0.105
20	A	2	2	1.00	18	0.111
21	A	2	2	1.00	17	0.118

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	N/A	1	0	1.00	19	0.000
23	A	2	2	1.00	19	0.105
24	A	4	4	1.00	24	0.167
25	A	2	2	1.00	21	0.095
26	A	2	2	1.00	21	0.095
27	A	2	2	1.00	21	0.095
28	A	3	3	1.00	19	0.158
29	A	2	2	1.00	21	0.095
30	A	2	2	0.92	21	0.095
31	A	3	3	1.00	20	0.150
32	A	2	2	1.37	21	0.095
33	A	2	2	1.22	21	0.095
34	A	4	4	0.99	19	0.211
35	A	2	2	1.34	21	0.095
36	A	2	2	1.18	21	0.095
37	A	2	2	1.64	21	0.095
38	A	2	2	1.23	21	0.095
39	A	5	5	1.05	19	0.263
40	A	2	2	1.26	21	0.095
41	A	2	2	1.06	21	0.095
42	A	3	3	1.00	23	0.130
43	A	3	3	1.00	23	0.130
44	A	3	3	1.00	23	0.130
45	A	5	5	1.73	23	0.217
46	A	2	2	1.00	23	0.087
47	A	2	2	1.14	21	0.095
48	A	2	2	1.08	21	0.095
49	A	2	2	1.00	19	0.105
50	A	2	2	1.00	21	0.095
51	A	2	2	1.00	21	0.095
52	A	2	2	1.00	21	0.095
53	N/A	1	0	1.00	21	0.000

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
54	A	4	4	1.00	29	0.138
55	A	1	1	1.00	22	0.045
56	A	2	2	1.00	24	0.083

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \frac{d+ex^3}{a+cx^6} dx$	49
3.2	$\int \frac{d+ex^3}{a-cx^6} dx$	61
3.3	$\int \frac{(d+ex^3)^{3/2}}{a+cx^6} dx$	74
3.4	$\int \frac{\sqrt{d+ex^3}}{a+cx^6} dx$	80
3.5	$\int \frac{1}{\sqrt{d+ex^3}(a+cx^6)} dx$	87
3.6	$\int \frac{1}{(d+ex^3)^{3/2}(a+cx^6)} dx$	94
3.7	$\int \frac{\sqrt{1-x^6}}{\sqrt{1-x^3}} dx$	101
3.8	$\int (d+ex^3)(a-cx^6)^p dx$	107
3.9	$\int (d+ex^3)(a+cx^6)^p dx$	113
3.10	$\int (d+ex^3)^q (a+cx^6)^p dx$	119
3.11	$\int (1-x^3)^p (1-x^6)^p dx$	124
3.12	$\int (d+ex^3)^q (d^2-e^2x^6)^p dx$	129
3.13	$\int (1-x^3)^{-p} (1-x^6)^p dx$	134
3.14	$\int \frac{d+ex^4}{a-cx^8} dx$	139
3.15	$\int \frac{d+ex^4}{a+cx^8} dx$	152
3.16	$\int \frac{d+ex^4}{\sqrt{1-\frac{e^2x^8}{d^2}}} dx$	166
3.17	$\int \frac{d+ex^4}{\sqrt{d^2-e^2x^8}} dx$	172
3.18	$\int \frac{d+ex^4}{\sqrt{a-cx^8}} dx$	177
3.19	$\int \frac{d+ex^4}{\sqrt{a+cx^8}} dx$	182
3.20	$\int (d+ex^4)(a-cx^8)^p dx$	188
3.21	$\int (d+ex^4)(a+cx^8)^p dx$	194
3.22	$\int (d+ex^4)^q (a+cx^8)^p dx$	200
3.23	$\int (1-x^4)^p (1-x^8)^p dx$	205
3.24	$\int (d+ex^4)^q (d^2-e^2x^8)^p dx$	210
3.25	$\int (1-x^4)^{-p} (1-x^8)^p dx$	215

3.26	$\int \frac{(d+ex^n)^3}{a+cx^{2n}} dx$	220
3.27	$\int \frac{(d+ex^n)^2}{a+cx^{2n}} dx$	226
3.28	$\int \frac{d+ex^n}{a+cx^{2n}} dx$	231
3.29	$\int \frac{1}{(d+ex^n)(a+cx^{2n})} dx$	236
3.30	$\int \frac{1}{(d+ex^n)^2(a+cx^{2n})} dx$	241
3.31	$\int \frac{d+ex^n}{a-cx^{2n}} dx$	246
3.32	$\int \frac{(d+ex^n)^3}{(a+cx^{2n})^2} dx$	251
3.33	$\int \frac{(d+ex^n)^2}{(a+cx^{2n})^2} dx$	257
3.34	$\int \frac{d+ex^n}{(a+cx^{2n})^2} dx$	263
3.35	$\int \frac{1}{(d+ex^n)(a+cx^{2n})^2} dx$	269
3.36	$\int \frac{1}{(d+ex^n)^2(a+cx^{2n})^2} dx$	275
3.37	$\int \frac{(d+ex^n)^3}{(a+cx^{2n})^3} dx$	281
3.38	$\int \frac{(d+ex^n)^2}{(a+cx^{2n})^3} dx$	287
3.39	$\int \frac{d+ex^n}{(a+cx^{2n})^3} dx$	293
3.40	$\int \frac{1}{(d+ex^n)(a+cx^{2n})^3} dx$	299
3.41	$\int \frac{1}{(d+ex^n)^2(a+cx^{2n})^3} dx$	306
3.42	$\int \frac{(d+ex^n)^{3/2}}{a+cx^{2n}} dx$	313
3.43	$\int \frac{\sqrt{d+ex^n}}{a+cx^{2n}} dx$	318
3.44	$\int \frac{1}{\sqrt{d+ex^n}(a+cx^{2n})} dx$	323
3.45	$\int \frac{1}{(d+ex^n)^{3/2}(a+cx^{2n})} dx$	328
3.46	$\int \frac{1}{(d+ex^n)\sqrt{a+cx^{2n}}} dx$	334
3.47	$\int (d+ex^n)^3 (a+cx^{2n})^p dx$	339
3.48	$\int (d+ex^n)^2 (a+cx^{2n})^p dx$	345
3.49	$\int (d+ex^n) (a+cx^{2n})^p dx$	351
3.50	$\int \frac{(a+cx^{2n})^p}{d+ex^n} dx$	357
3.51	$\int \frac{(a+cx^{2n})^p}{(d+ex^n)^2} dx$	362
3.52	$\int \frac{(a+cx^{2n})^p}{(d+ex^n)^3} dx$	367
3.53	$\int (d+ex^n)^q (a+cx^{2n})^p dx$	372
3.54	$\int (d+ex^n)^q (cd^2 - ce^2 x^{2n})^p dx$	377
3.55	$\int (2-ex^n)^p (2+ex^n)^{p+q} dx$	382
3.56	$\int (2+ex^n)^q (4-e^2 x^{2n})^p dx$	387

3.1 $\int \frac{d+ex^3}{a+cx^6} dx$

Optimal result	49
Mathematica [A] (verified)	50
Rubi [A] (verified)	50
Maple [C] (verified)	55
Fricas [B] (verification not implemented)	56
Sympy [A] (verification not implemented)	57
Maxima [A] (verification not implemented)	57
Giac [A] (verification not implemented)	58
Mupad [B] (verification not implemented)	59
Reduce [B] (verification not implemented)	60

Optimal result

Integrand size = 17, antiderivative size = 305

$$\begin{aligned} \int \frac{d+ex^3}{a+cx^6} dx = & \frac{d \arctan\left(\frac{\sqrt[6]{cx}}{\sqrt[6]{a}}\right)}{3a^{5/6}\sqrt[6]{c}} - \frac{(\sqrt{cd} + \sqrt{3}\sqrt{ae}) \arctan\left(\sqrt{3} - \frac{2\sqrt[6]{cx}}{\sqrt[6]{a}}\right)}{6a^{5/6}c^{2/3}} \\ & + \frac{(\sqrt{cd} - \sqrt{3}\sqrt{ae}) \arctan\left(\sqrt{3} + \frac{2\sqrt[6]{cx}}{\sqrt[6]{a}}\right)}{6a^{5/6}c^{2/3}} - \frac{e \log(\sqrt[3]{a} + \sqrt[3]{cx^2})}{6\sqrt[3]{ac^{2/3}}} \\ & - \frac{(\sqrt{3}\sqrt{cd} - \sqrt{ae}) \log(\sqrt[3]{a} - \sqrt{3}\sqrt[6]{a}\sqrt[6]{cx} + \sqrt[3]{cx^2})}{12a^{5/6}c^{2/3}} \\ & + \frac{(\sqrt{3}\sqrt{cd} + \sqrt{ae}) \log(\sqrt[3]{a} + \sqrt{3}\sqrt[6]{a}\sqrt[6]{cx} + \sqrt[3]{cx^2})}{12a^{5/6}c^{2/3}} \end{aligned}$$

output

```
1/3*d*arctan(c^(1/6)*x/a^(1/6))/a^(5/6)/c^(1/6)+1/6*(c^(1/2)*d+3^(1/2)*a^(1/2)*e)*arctan(-3^(1/2)+2*c^(1/6)*x/a^(1/6))/a^(5/6)/c^(2/3)+1/6*(c^(1/2)*d-3^(1/2)*a^(1/2)*e)*arctan(3^(1/2)+2*c^(1/6)*x/a^(1/6))/a^(5/6)/c^(2/3)-1/6*e*ln(a^(1/3)+c^(1/3)*x^2)/a^(1/3)/c^(2/3)-1/12*(3^(1/2)*c^(1/2)*d-a^(1/2)*e)*ln(a^(1/3)-3^(1/2)*a^(1/6)*c^(1/6)*x+c^(1/3)*x^2)/a^(5/6)/c^(2/3)+1/12*(3^(1/2)*c^(1/2)*d+a^(1/2)*e)*ln(a^(1/3)+3^(1/2)*a^(1/6)*c^(1/6)*x+c^(1/3)*x^2)/a^(5/6)/c^(2/3)
```

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 334, normalized size of antiderivative = 1.10

$$\int \frac{d + ex^3}{a + cx^6} dx = \frac{d \arctan\left(\frac{\sqrt[6]{cx}}{\sqrt[6]{a}}\right)}{3a^{5/6}\sqrt[6]{c}} + \frac{(\sqrt[6]{a}\sqrt{cd} + \sqrt{3}a^{2/3}e) \arctan\left(\frac{-\sqrt{3}\sqrt[6]{a+2}\sqrt[6]{cx}}{\sqrt[6]{a}}\right)}{6ac^{2/3}} \\ + \frac{(\sqrt[6]{a}\sqrt{cd} - \sqrt{3}a^{2/3}e) \arctan\left(\frac{\sqrt{3}\sqrt[6]{a+2}\sqrt[6]{cx}}{\sqrt[6]{a}}\right)}{6ac^{2/3}} - \frac{e \log(\sqrt[3]{a} + \sqrt[3]{cx^2})}{6\sqrt[3]{ac^{2/3}}} \\ - \frac{(\sqrt{3}\sqrt[6]{a}\sqrt{cd} - a^{2/3}e) \log(\sqrt[3]{a} - \sqrt{3}\sqrt[6]{a}\sqrt[6]{cx} + \sqrt[3]{cx^2})}{12ac^{2/3}} \\ - \frac{(-\sqrt{3}\sqrt[6]{a}\sqrt{cd} - a^{2/3}e) \log(\sqrt[3]{a} + \sqrt{3}\sqrt[6]{a}\sqrt[6]{cx} + \sqrt[3]{cx^2})}{12ac^{2/3}}$$

input `Integrate[(d + e*x^3)/(a + c*x^6), x]`

output `(d*ArcTan[(c^(1/6)*x)/a^(1/6)])/(3*a^(5/6)*c^(1/6)) + ((a^(1/6)*Sqrt[c]*d + Sqrt[3]*a^(2/3)*e)*ArcTan[(-(Sqrt[3]*a^(1/6)) + 2*c^(1/6)*x)/a^(1/6)])/(6*a*c^(2/3)) + ((a^(1/6)*Sqrt[c]*d - Sqrt[3]*a^(2/3)*e)*ArcTan[(Sqrt[3]*a^(1/6) + 2*c^(1/6)*x)/a^(1/6)])/(6*a*c^(2/3)) - (e*Log[a^(1/3) + c^(1/3)*x^2])/(6*a^(1/3)*c^(2/3)) - ((Sqrt[3]*a^(1/6)*Sqrt[c]*d - a^(2/3)*e)*Log[a^(1/3) - Sqrt[3]*a^(1/6)*c^(1/6)*x + c^(1/3)*x^2])/(12*a*c^(2/3)) - ((-(Sqrt[3]*a^(1/6)*Sqrt[c]*d) - a^(2/3)*e)*Log[a^(1/3) + Sqrt[3]*a^(1/6)*c^(1/6)*x + c^(1/3)*x^2])/(12*a*c^(2/3))`

Rubi [A] (verified)

Time = 0.65 (sec) , antiderivative size = 352, normalized size of antiderivative = 1.15, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.647$, Rules used = {1746, 27, 452, 218, 240, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{d + ex^3}{a + cx^6} dx$$

$$\begin{aligned}
& \downarrow 1746 \\
& \frac{\int \frac{\frac{2}{3}\sqrt[3]{cd} - \sqrt[3]{a}\left(\frac{\sqrt{3}\sqrt{cd}}{\sqrt{a}} - e\right)x}{\sqrt[3]{a}\left(\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} - \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1\right)} dx}{\frac{3a^{2/3}\sqrt[3]{c}}{3a^{2/3}\sqrt[3]{c}}} + \frac{\int \frac{2\sqrt[3]{cd} + \sqrt[3]{a}\left(\frac{\sqrt{3}\sqrt{cd}}{\sqrt{a}} + e\right)x}{\sqrt[3]{a}\left(\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} + \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1\right)} dx}{\frac{6a^{2/3}\sqrt[3]{c}}{6a^{2/3}\sqrt[3]{c}}} \\
& \downarrow 27 \\
& \frac{\int \frac{\frac{2}{3}\sqrt[3]{cd} - \sqrt[3]{a}\left(\frac{\sqrt{3}\sqrt{cd}}{\sqrt{a}} - e\right)x}{\sqrt[3]{a}\left(\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} - \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1\right)} dx}{\frac{3a^{2/3}\sqrt[3]{c}}{3a^{2/3}\sqrt[3]{c}}} + \frac{\int \frac{2\sqrt[3]{cd} + \sqrt[3]{a}\left(\frac{\sqrt{3}\sqrt{cd}}{\sqrt{a}} + e\right)x}{\sqrt[3]{a}\left(\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} + \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1\right)} dx}{\frac{6a^{2/3}\sqrt[3]{c}}{6a^{2/3}\sqrt[3]{c}}} \\
& \downarrow 452 \\
& \frac{\sqrt{cd} \int \frac{1}{\sqrt[3]{c_x^2} + \sqrt[3]{a}} dx - \sqrt{ae} \int \frac{x}{\sqrt[3]{c_x^2} + \sqrt[3]{a}} dx}{\frac{3a^{2/3}\sqrt[3]{c}}{3a^{2/3}\sqrt[3]{c}}} + \frac{\int \frac{2\sqrt[3]{cd} - \sqrt[3]{a}\left(\frac{\sqrt{3}\sqrt{cd}}{\sqrt{a}} - e\right)x}{\sqrt[3]{a}\left(\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} - \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1\right)} dx}{\frac{6a^{2/3}\sqrt[3]{c}}{6a^{2/3}\sqrt[3]{c}}} + \\
& \quad \frac{\int \frac{2\sqrt[3]{cd} + \sqrt[3]{a}\left(\frac{\sqrt{3}\sqrt{cd}}{\sqrt{a}} + e\right)x}{\sqrt[3]{a}\left(\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} + \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1\right)} dx}{\frac{6a^{2/3}\sqrt[3]{c}}{6a^{2/3}\sqrt[3]{c}}} \\
& \downarrow 218 \\
& \frac{\frac{\sqrt{cd} \arctan\left(\frac{\sqrt[6]{c_x}}{\sqrt[6]{a}}\right)}{\sqrt[6]{a}} - \sqrt{ae} \int \frac{x}{\sqrt[3]{c_x^2} + \sqrt[3]{a}} dx}{\frac{3a^{2/3}\sqrt[3]{c}}{3a^{2/3}\sqrt[3]{c}}} + \frac{\int \frac{2\sqrt[3]{cd} - \sqrt[3]{a}\left(\frac{\sqrt{3}\sqrt{cd}}{\sqrt{a}} - e\right)x}{\sqrt[3]{a}\left(\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} - \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1\right)} dx}{\frac{6a^{2/3}\sqrt[3]{c}}{6a^{2/3}\sqrt[3]{c}}} + \\
& \quad \frac{\int \frac{2\sqrt[3]{cd} + \sqrt[3]{a}\left(\frac{\sqrt{3}\sqrt{cd}}{\sqrt{a}} + e\right)x}{\sqrt[3]{a}\left(\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} + \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1\right)} dx}{\frac{6a^{2/3}\sqrt[3]{c}}{6a^{2/3}\sqrt[3]{c}}} \\
& \downarrow 240 \\
& \frac{\int \frac{2\sqrt[3]{cd} - \sqrt[3]{a}\left(\frac{\sqrt{3}\sqrt{cd}}{\sqrt{a}} - e\right)x}{\sqrt[3]{a}\left(\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} - \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1\right)} dx}{\frac{6a^{2/3}\sqrt[3]{c}}{6a^{2/3}\sqrt[3]{c}}} + \frac{\int \frac{2\sqrt[3]{cd} + \sqrt[3]{a}\left(\frac{\sqrt{3}\sqrt{cd}}{\sqrt{a}} + e\right)x}{\sqrt[3]{a}\left(\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} + \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1\right)} dx}{\frac{6a^{2/3}\sqrt[3]{c}}{6a^{2/3}\sqrt[3]{c}}} + \\
& \quad \frac{\frac{\sqrt{cd} \arctan\left(\frac{\sqrt[6]{c_x}}{\sqrt[6]{a}}\right)}{\sqrt[6]{a}} - \frac{\sqrt{a}e \log\left(\sqrt[3]{a} + \sqrt[3]{c_x^2}\right)}{2\sqrt[3]{c}}}{\frac{3a^{2/3}\sqrt[3]{c}}{3a^{2/3}\sqrt[3]{c}}} \\
& \downarrow 1142
\end{aligned}$$

$$\begin{aligned}
& \frac{\left(\sqrt{3}\sqrt{ae}+\sqrt{cd}\right) \int \frac{1}{\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} - \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1} dx}{2\sqrt[6]{c}} - \frac{a^{2/3} \left(\frac{\sqrt{3}\sqrt{cd}}{\sqrt{a}} - e \right) \int -\frac{\sqrt[6]{c}(\sqrt{3}\sqrt[6]{a} - 2\sqrt[6]{c_x})}{\sqrt[3]{a} \left(\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} - \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1 \right)} dx}{2\sqrt[3]{c}} + \\
& \frac{6a\sqrt[3]{c}}{6a\sqrt[3]{c}} \\
& \frac{\left(\sqrt{cd}-\sqrt{3}\sqrt{ae}\right) \int \frac{1}{\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} + \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1} dx}{2\sqrt[6]{c}} + \frac{\sqrt[6]{a} \left(\sqrt{ae}+\sqrt{3}\sqrt{cd}\right) \int \frac{\sqrt[6]{c}(2\sqrt[6]{c_x} + \sqrt{3}\sqrt[6]{a})}{\sqrt[3]{a} \left(\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} + \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1 \right)} dx}{2\sqrt[3]{c}} + \\
& \frac{6a\sqrt[3]{c}}{6a\sqrt[3]{c}} \\
& \frac{\frac{\sqrt[6]{c}d \arctan\left(\frac{\sqrt[6]{c_x}}{\sqrt[6]{a}}\right)}{\sqrt[6]{a}} - \frac{\sqrt[3]{a}e \log\left(\sqrt[3]{a} + \sqrt[3]{c_x^2}\right)}{2\sqrt[3]{c}}}{3a^{2/3}\sqrt[3]{c}} \\
& \downarrow 25 \\
& \frac{a^{2/3} \left(\frac{\sqrt{3}\sqrt{cd}}{\sqrt{a}} - e \right) \int \frac{\sqrt[6]{c}(\sqrt{3}\sqrt[6]{a} - 2\sqrt[6]{c_x})}{\sqrt[3]{a} \left(\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} - \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1 \right)} dx}{2\sqrt[3]{c}} + \frac{\left(\sqrt{3}\sqrt{ae}+\sqrt{cd}\right) \int \frac{1}{\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} - \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1} dx}{2\sqrt[6]{c}} + \\
& \frac{6a\sqrt[3]{c}}{6a\sqrt[3]{c}} \\
& \frac{\left(\sqrt{cd}-\sqrt{3}\sqrt{ae}\right) \int \frac{1}{\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} + \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1} dx}{2\sqrt[6]{c}} + \frac{\sqrt[6]{a} \left(\sqrt{ae}+\sqrt{3}\sqrt{cd}\right) \int \frac{\sqrt[6]{c}(2\sqrt[6]{c_x} + \sqrt{3}\sqrt[6]{a})}{\sqrt[3]{a} \left(\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} + \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1 \right)} dx}{2\sqrt[3]{c}} + \\
& \frac{6a\sqrt[3]{c}}{6a\sqrt[3]{c}} \\
& \frac{\frac{\sqrt[6]{c}d \arctan\left(\frac{\sqrt[6]{c_x}}{\sqrt[6]{a}}\right)}{\sqrt[6]{a}} - \frac{\sqrt[3]{a}e \log\left(\sqrt[3]{a} + \sqrt[3]{c_x^2}\right)}{2\sqrt[3]{c}}}{3a^{2/3}\sqrt[3]{c}} \\
& \downarrow 27 \\
& \frac{\left(\sqrt{3}\sqrt{ae}+\sqrt{cd}\right) \int \frac{1}{\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} - \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1} dx}{2\sqrt[6]{c}} + \frac{\sqrt[3]{a} \left(\frac{\sqrt{3}\sqrt{cd}}{\sqrt{a}} - e \right) \int \frac{\sqrt{3}\sqrt[6]{a} - 2\sqrt[6]{c_x}}{\sqrt[3]{a} \left(\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} - \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1 \right)} dx}{2\sqrt[3]{c}} + \\
& \frac{6a\sqrt[3]{c}}{6a\sqrt[3]{c}} \\
& \frac{\left(\sqrt{cd}-\sqrt{3}\sqrt{ae}\right) \int \frac{1}{\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} + \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1} dx}{2\sqrt[6]{c}} + \frac{\left(\sqrt{ae}+\sqrt{3}\sqrt{cd}\right) \int \frac{2\sqrt[6]{c_x} + \sqrt{3}\sqrt[6]{a}}{\sqrt[3]{a} \left(\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} + \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1 \right)} dx}{2\sqrt[3]{a}\sqrt[6]{c}} + \\
& \frac{6a\sqrt[3]{c}}{6a\sqrt[3]{c}} \\
& \frac{\frac{\sqrt[6]{c}d \arctan\left(\frac{\sqrt[6]{c_x}}{\sqrt[6]{a}}\right)}{\sqrt[6]{a}} - \frac{\sqrt[3]{a}e \log\left(\sqrt[3]{a} + \sqrt[3]{c_x^2}\right)}{2\sqrt[3]{c}}}{3a^{2/3}\sqrt[3]{c}}
\end{aligned}$$

$$\begin{aligned}
& \downarrow \textcolor{blue}{1082} \\
& \frac{\sqrt[3]{a} \left(\frac{\sqrt{3}\sqrt{cd}}{\sqrt{a}} - e \right) \int \frac{\sqrt{3}\sqrt[6]{a} - 2\sqrt[6]{c_x}}{\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} - \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1} dx}{2\sqrt[6]{c}} + \frac{\sqrt[6]{a} \left(\sqrt{3}\sqrt{ae} + \sqrt{cd} \right) \int \frac{1}{-\left(1 - \frac{2\sqrt[6]{c_x}}{\sqrt[3]\sqrt[6]{a}} \right)^2 - \frac{1}{3}} d \left(1 - \frac{2\sqrt[6]{c_x}}{\sqrt[3]\sqrt[6]{a}} \right)}{\sqrt[3]\sqrt[3]{c}} + \\
& \frac{\left(\sqrt{ae} + \sqrt{3}\sqrt{cd} \right) \int \frac{2\sqrt[6]{c_x} + \sqrt{3}\sqrt[6]{a}}{\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} + \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1} dx}{2\sqrt[6]{a}\sqrt[6]{c}} - \frac{\sqrt[6]{a} \left(\sqrt{cd} - \sqrt{3}\sqrt{ae} \right) \int \frac{1}{-\left(\frac{2\sqrt[6]{c_x}}{\sqrt[3]\sqrt[6]{a}} + 1 \right)^2 - \frac{1}{3}} d \left(\frac{2\sqrt[6]{c_x}}{\sqrt[3]\sqrt[6]{a}} + 1 \right)}{\sqrt[3]\sqrt[3]{c}} + \\
& \frac{\sqrt[6]{c} \arctan \left(\frac{\sqrt[6]{c_x}}{\sqrt[6]{a}} \right) - \frac{\sqrt[3]{ae} \log \left(\sqrt[3]{a} + \sqrt[3]{c_x^2} \right)}{2\sqrt[3]{c}}}{3a^{2/3}\sqrt[3]{c}} \\
& \downarrow \textcolor{blue}{217} \\
& \frac{\sqrt[3]{a} \left(\frac{\sqrt{3}\sqrt{cd}}{\sqrt{a}} - e \right) \int \frac{\sqrt{3}\sqrt[6]{a} - 2\sqrt[6]{c_x}}{\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} - \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1} dx}{2\sqrt[6]{c}} - \frac{\sqrt[6]{a} \arctan \left(\sqrt{3} \left(1 - \frac{2\sqrt[6]{c_x}}{\sqrt[3]\sqrt[6]{a}} \right) \right) (\sqrt{3}\sqrt{ae} + \sqrt{cd})}{\sqrt[3]{c}} + \\
& \frac{\left(\sqrt{ae} + \sqrt{3}\sqrt{cd} \right) \int \frac{2\sqrt[6]{c_x} + \sqrt{3}\sqrt[6]{a}}{\frac{\sqrt[3]{c_x^2}}{\sqrt[3]{a}} + \frac{\sqrt{3}\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1} dx}{2\sqrt[6]{a}\sqrt[6]{c}} + \frac{\sqrt[6]{a} \arctan \left(\sqrt{3} \left(\frac{2\sqrt[6]{c_x}}{\sqrt[3]\sqrt[6]{a}} + 1 \right) \right) (\sqrt{cd} - \sqrt{3}\sqrt{ae})}{\sqrt[3]{c}} + \\
& \frac{\sqrt[6]{c} \arctan \left(\frac{\sqrt[6]{c_x}}{\sqrt[6]{a}} \right) - \frac{\sqrt[3]{ae} \log \left(\sqrt[3]{a} + \sqrt[3]{c_x^2} \right)}{2\sqrt[3]{c}}}{3a^{2/3}\sqrt[3]{c}} \\
& \downarrow \textcolor{blue}{1103} \\
& \frac{\sqrt[6]{c} \arctan \left(\frac{\sqrt[6]{c_x}}{\sqrt[6]{a}} \right) - \frac{\sqrt[3]{ae} \log \left(\sqrt[3]{a} + \sqrt[3]{c_x^2} \right)}{2\sqrt[3]{c}}}{3a^{2/3}\sqrt[3]{c}} + \\
& - \frac{a^{2/3} \left(\frac{\sqrt{3}\sqrt{cd}}{\sqrt{a}} - e \right) \log \left(-\sqrt{3}\sqrt[6]{a} \sqrt[6]{c_x} + \sqrt[3]{a} + \sqrt[3]{c_x^2} \right)}{2\sqrt[3]{c}} - \frac{\sqrt[6]{a} \arctan \left(\sqrt{3} \left(1 - \frac{2\sqrt[6]{c_x}}{\sqrt[3]\sqrt[6]{a}} \right) \right) (\sqrt{3}\sqrt{ae} + \sqrt{cd})}{\sqrt[3]{c}} + \\
& \frac{\sqrt[6]{a} \arctan \left(\sqrt{3} \left(\frac{2\sqrt[6]{c_x}}{\sqrt[3]\sqrt[6]{a}} + 1 \right) \right) (\sqrt{cd} - \sqrt{3}\sqrt{ae})}{\sqrt[3]{c}} + \frac{\sqrt[6]{a} (\sqrt{ae} + \sqrt{3}\sqrt{cd}) \log \left(\sqrt{3}\sqrt[6]{a} \sqrt[6]{c_x} + \sqrt[3]{a} + \sqrt[3]{c_x^2} \right)}{2\sqrt[3]{c}} \\
& \frac{6a\sqrt[3]{c}}{6a\sqrt[3]{c}}
\end{aligned}$$

input $\text{Int}[(d + e*x^3)/(a + c*x^6), x]$

output $((c^{(1/6)}*d*\text{ArcTan}[(c^{(1/6)}*x)/a^{(1/6)}])/a^{(1/6)} - (a^{(1/3)}*e*\text{Log}[a^{(1/3)} + c^{(1/3)}*x^2])/(2*c^{(1/3)})) / (3*a^{(2/3)}*c^{(1/3)}) + ((-((a^{(1/6)}*(\text{Sqrt}[c]*d + \text{Sqrt}[3]*\text{Sqrt}[a]*e)*\text{ArcTan}[\text{Sqrt}[3]*(1 - (2*c^{(1/6)}*x)/(\text{Sqrt}[3]*a^{(1/6)}))]) / c^{(1/3)}) - (a^{(2/3)}*((\text{Sqrt}[3]*\text{Sqrt}[c]*d)/\text{Sqrt}[a] - e)*\text{Log}[a^{(1/3)} - \text{Sqrt}[3]*a^{(1/6)}*c^{(1/6)}*x + c^{(1/3)}*x^2]) / (2*c^{(1/3)})) / (6*a*c^{(1/3)}) + ((a^{(1/6)}*(\text{Sqrt}[c]*d - \text{Sqrt}[3]*\text{Sqrt}[a]*e)*\text{ArcTan}[\text{Sqrt}[3]*(1 + (2*c^{(1/6)}*x)/(\text{Sqrt}[3]*a^{(1/6)}))]) / c^{(1/3)} + (a^{(1/6)}*(\text{Sqrt}[3]*\text{Sqrt}[c]*d + \text{Sqrt}[a]*e)*\text{Log}[a^{(1/3)} + \text{Sqrt}[3]*a^{(1/6)}*c^{(1/6)}*x + c^{(1/3)}*x^2]) / (2*c^{(1/3)})) / (6*a*c^{(1/3)})$

Definitions of rubi rules used

rule 25 $\text{Int}[-(F_x), x_{\text{Symbol}}] \Rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$

rule 27 $\text{Int}[(a_)*(F_x), x_{\text{Symbol}}] \Rightarrow \text{Simp}[a \quad \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \&& \text{!MatchQ}[F_x, (b_)*(G_x) /; \text{FreeQ}[b, x]]$

rule 217 $\text{Int}[((a_) + (b_)*(x_)^2)^{-1}, x_{\text{Symbol}}] \Rightarrow \text{Simp}[(-(Rt[-a, 2]*Rt[-b, 2])^{-1})*\text{ArcTan}[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&& \text{PosQ}[a/b] \& \& (\text{LtQ}[a, 0] \text{ || } \text{LtQ}[b, 0])$

rule 218 $\text{Int}[((a_) + (b_)*(x_)^2)^{-1}, x_{\text{Symbol}}] \Rightarrow \text{Simp}[(Rt[a/b, 2]/a)*\text{ArcTan}[x/Rt[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \&& \text{PosQ}[a/b]$

rule 240 $\text{Int}[(x_)/((a_) + (b_)*(x_)^2), x_{\text{Symbol}}] \Rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x^2, x]]/(2*b), x] /; \text{FreeQ}[\{a, b\}, x]$

rule 452 $\text{Int}[((c_) + (d_)*(x_))/((a_) + (b_)*(x_)^2), x_{\text{Symbol}}] \Rightarrow \text{Simp}[c \quad \text{Int}[1/(a + b*x^2), x], x] + \text{Simp}[d \quad \text{Int}[x/(a + b*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&& \text{NeQ}[b*c^2 + a*d^2, 0]$

rule 1082 $\text{Int}[(a_ + b_*)*(x_ + c_*)^2, x] \rightarrow \text{With}[q = 1 - 4S \text{implify}[a*(c/b^2)], \text{Simp}[-2/b \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \&& (\text{EqQ}[q^2, 1] \text{||} \text{!RationalQ}[b^2 - 4*a*c]) /; \text{FreeQ}[\{a, b, c\}, x]]$

rule 1103 $\text{Int}[(d_ + e_*)*(x_)/((a_ + b_*)*(x_ + c_*)^2), x] \rightarrow S \text{imp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{EqQ}[2*c*d - b*e, 0]$

rule 1142 $\text{Int}[(d_ + e_*)*(x_)/((a_ + b_*)*(x_ + c_*)^2), x] \rightarrow S \text{imp}[(2*c*d - b*e)/(2*c) \text{Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \text{Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]]$

rule 1746 $\text{Int}[(d_ + e_*)*(x_)^3/((a_ + c_*)*(x_)^6), x] \rightarrow \text{With}[q = \text{Rt}[c/a, 6], \text{Simp}[1/(3*a*q^2) \text{Int}[(q^2*d - e*x)/(1 + q^2*x^2), x], x] + (\text{Simp}[1/(6*a*q^2) \text{Int}[(2*q^2*d - (\text{Sqrt}[3]*q^3*d - e)*x)/(1 - \text{Sqrt}[3]*q*x + q^2*x^2), x], x] + \text{Simp}[1/(6*a*q^2) \text{Int}[(2*q^2*d + (\text{Sqrt}[3]*q^3*d + e)*x)/(1 + \text{Sqrt}[3]*q*x + q^2*x^2), x], x])] /; \text{FreeQ}[\{a, c, d, e\}, x] \&& \text{NeQ}[c*d^2 + a*e^2, 0] \&& \text{PosQ}[c/a]]$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.15 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.11

method	result
risch	$\frac{\sum_{R=\text{RootOf}(-Z^6 c+a)} \frac{(-R^{3/e+d}) \ln(x-R)}{-R^5}}{6c}$
default	$\frac{\ln\left(x^2-\sqrt{3}\left(\frac{a}{c}\right)^{\frac{1}{6}}x+\left(\frac{a}{c}\right)^{\frac{1}{3}}\right)\left(\frac{a}{c}\right)^{\frac{2}{3}}e}{12a} - \frac{\ln\left(x^2-\sqrt{3}\left(\frac{a}{c}\right)^{\frac{1}{6}}x+\left(\frac{a}{c}\right)^{\frac{1}{3}}\right)\sqrt{3}\left(\frac{a}{c}\right)^{\frac{1}{6}}d}{12a} + \frac{\left(\frac{a}{c}\right)^{\frac{2}{3}}\arctan\left(\frac{2x}{\left(\frac{a}{c}\right)^{\frac{1}{6}}}-\sqrt{3}\right)\sqrt{3}e}{6a} + \dots$

input `int((e*x^3+d)/(c*x^6+a),x,method=_RETURNVERBOSE)`

output $\frac{1}{6}c \operatorname{sum}((R^3 e + d) / R^5 \ln(x - R), R = \operatorname{RootOf}(Z^6 c + a))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1631 vs. $2(207) = 414$.

Time = 0.14 (sec) , antiderivative size = 1631, normalized size of antiderivative = 5.35

$$\int \frac{d + ex^3}{a + cx^6} dx = \text{Too large to display}$$

input `integrate((e*x^3+d)/(c*x^6+a),x, algorithm="fricas")`

output
$$\begin{aligned} & -\frac{1}{12}(\sqrt{-3} + 1)*((a^2*c^2*\sqrt{-(c^2*d^6 - 6*a*c*d^4*e^2 + 9*a^2*d^2*e^4)/(a^5*c^3)} + 3*c*d^2*e - a*e^3)/(a^2*c^2))^{(1/3)}*\log(-(c^2*d^5 - 2*a*c*d^3*e^2 - 3*a^2*d^2*e^4)*x + 1/2*(a*c^2*d^4 - 3*a^2*c*d^2*e^2 + \sqrt{-3}*(a*c^2*d^4 - 3*a^2*c*d^2*e^2) + (\sqrt{-3}*a^4*c^2*e + a^4*c^2*e)*\sqrt{-(c^2*d^6 - 6*a*c*d^4*e^2 + 9*a^2*d^2*e^4)/(a^5*c^3)}))*((a^2*c^2*\sqrt{-(c^2*d^6 - 6*a*c*d^4*e^2 + 9*a^2*d^2*e^4)/(a^5*c^3)}) + 3*c*d^2*e - a*e^3)/(a^2*c^2))^{(1/3)} + 1/12*(\sqrt{-3} - 1)*((a^2*c^2*\sqrt{-(c^2*d^6 - 6*a*c*d^4*e^2 + 9*a^2*d^2*e^4)/(a^5*c^3)}) + 3*c*d^2*e - a*e^3)/(a^2*c^2))^{(1/3)}*\log(-(c^2*d^5 - 2*a*c*d^3*e^2 - 3*a^2*d^2*e^4)*x + 1/2*(a*c^2*d^4 - 3*a^2*c*d^2*e^2 - \sqrt{-3}*(a*c^2*d^4 - 3*a^2*c*d^2*e^2) - (\sqrt{-3}*a^4*c^2*e - a^4*c^2*e)*\sqrt{-(c^2*d^6 - 6*a*c*d^4*e^2 + 9*a^2*d^2*e^4)/(a^5*c^3)}))*((a^2*c^2*\sqrt{-(c^2*d^6 - 6*a*c*d^4*e^2 + 9*a^2*d^2*e^4)/(a^5*c^3)}) + 3*c*d^2*e - a*e^3)/(a^2*c^2))^{(1/3)} - 1/12*(\sqrt{-3} + 1)*(-(\sqrt{-(c^2*d^6 - 6*a*c*d^4*e^2 + 9*a^2*d^2*e^4)/(a^5*c^3)}) - 3*c*d^2*e + a*e^3)/(a^2*c^2))^{(1/3)}*\log(-(c^2*d^5 - 2*a*c*d^3*e^2 - 3*a^2*d^2*e^4)*x + 1/2*(a*c^2*d^4 - 3*a^2*c*d^2*e^2 + \sqrt{-3}*(a*c^2*d^4 - 3*a^2*c*d^2*e^2) - (\sqrt{-3}*a^4*c^2*e + a^4*c^2*e)*\sqrt{-(c^2*d^6 - 6*a*c*d^4*e^2 + 9*a^2*d^2*e^4)/(a^5*c^3)}))*(-(\sqrt{-(c^2*d^6 - 6*a*c*d^4*e^2 + 9*a^2*d^2*e^4)/(a^5*c^3)}) - 3*c*d^2*e + a*e^3)/(a^2*c^2))^{(1/3)} + 1/12*(\sqrt{-3} - 1)*(-(\sqrt{-(c^2*d^6 - 6*a*c*d^4*e^2 + 9*a^2*d^2*e^4)/(a^5*c^3)}) - 3*c*d^2*e + a*e^3...)$$

Sympy [A] (verification not implemented)

Time = 1.92 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.54

$$\int \frac{d + ex^3}{a + cx^6} dx = \text{RootSum}\left(46656t^6a^5c^4 + t^3 \cdot (432a^4c^2e^3 - 1296a^3c^3d^2e) + a^3e^6 + 3a^2cd^2e^4 + 3ac^2d^4e^2 + c^3d^6, \left(t \mapsto t\right)\right)$$

input `integrate((e*x**3+d)/(c*x**6+a),x)`

output `RootSum(46656*_t**6*a**5*c**4 + _t**3*(432*a**4*c**2*e**3 - 1296*a**3*c**3*d**2*e) + a**3*e**6 + 3*a**2*c*d**2*e**4 + 3*a*c**2*d**4*e**2 + c**3*d**6, Lambda(_t, _t*log(x + (-1296*_t**4*a**4*c**2*e - 6*_t*a**3*e**4 + 36*_t*a**2*c*d**2*e**2 - 6*_t*a*c**2*d**4)/(3*a**2*d*e**4 + 2*a*c*d**3*e**2 - c**2*d**5))))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 282, normalized size of antiderivative = 0.92

$$\begin{aligned} \int \frac{d + ex^3}{a + cx^6} dx &= -\frac{e \log\left(c^{\frac{1}{3}}x^2 + a^{\frac{1}{3}}\right)}{6a^{\frac{1}{3}}c^{\frac{2}{3}}} + \frac{d \arctan\left(\frac{c^{\frac{1}{3}}x}{\sqrt{a^{\frac{1}{3}}c^{\frac{1}{3}}}}\right)}{3a^{\frac{2}{3}}\sqrt{a^{\frac{1}{3}}c^{\frac{1}{3}}}} \\ &\quad + \frac{\left(\sqrt{3}a^{\frac{1}{6}}\sqrt{cd} + a^{\frac{2}{3}}e\right) \log\left(c^{\frac{1}{3}}x^2 + \sqrt{3}a^{\frac{1}{6}}c^{\frac{1}{6}}x + a^{\frac{1}{3}}\right)}{12ac^{\frac{2}{3}}} \\ &\quad - \frac{\left(\sqrt{3}a^{\frac{1}{6}}\sqrt{cd} - a^{\frac{2}{3}}e\right) \log\left(c^{\frac{1}{3}}x^2 - \sqrt{3}a^{\frac{1}{6}}c^{\frac{1}{6}}x + a^{\frac{1}{3}}\right)}{12ac^{\frac{2}{3}}} \\ &\quad - \frac{\left(\sqrt{3}a^{\frac{5}{6}}c^{\frac{1}{6}}e - a^{\frac{1}{3}}c^{\frac{2}{3}}d\right) \arctan\left(\frac{2c^{\frac{1}{3}}x + \sqrt{3}a^{\frac{1}{6}}c^{\frac{1}{6}}}{\sqrt{a^{\frac{1}{3}}c^{\frac{1}{3}}}}\right)}{6ac^{\frac{2}{3}}\sqrt{a^{\frac{1}{3}}c^{\frac{1}{3}}}} \\ &\quad + \frac{\left(\sqrt{3}a^{\frac{5}{6}}c^{\frac{1}{6}}e + a^{\frac{1}{3}}c^{\frac{2}{3}}d\right) \arctan\left(\frac{2c^{\frac{1}{3}}x - \sqrt{3}a^{\frac{1}{6}}c^{\frac{1}{6}}}{\sqrt{a^{\frac{1}{3}}c^{\frac{1}{3}}}}\right)}{6ac^{\frac{2}{3}}\sqrt{a^{\frac{1}{3}}c^{\frac{1}{3}}}} \end{aligned}$$

input `integrate((e*x^3+d)/(c*x^6+a),x, algorithm="maxima")`

output

```

-1/6*e*log(c^(1/3)*x^2 + a^(1/3))/(a^(1/3)*c^(2/3)) + 1/3*d*arctan(c^(1/3)
*x/sqrt(a^(1/3)*c^(1/3)))/(a^(2/3)*sqrt(a^(1/3)*c^(1/3))) + 1/12*(sqrt(3)*
a^(1/6)*sqrt(c)*d + a^(2/3)*e)*log(c^(1/3)*x^2 + sqrt(3)*a^(1/6)*c^(1/6)*x
+ a^(1/3))/(a*c^(2/3)) - 1/12*(sqrt(3)*a^(1/6)*sqrt(c)*d - a^(2/3)*e)*log
(c^(1/3)*x^2 - sqrt(3)*a^(1/6)*c^(1/6)*x + a^(1/3))/(a*c^(2/3)) - 1/6*(sqr
t(3)*a^(5/6)*c^(1/6)*e - a^(1/3)*c^(2/3)*d)*arctan((2*c^(1/3)*x + sqrt(3)*
a^(1/6)*c^(1/6))/sqrt(a^(1/3)*c^(1/3)))/(a*c^(2/3)*sqrt(a^(1/3)*c^(1/3)))
+ 1/6*(sqrt(3)*a^(5/6)*c^(1/6)*e + a^(1/3)*c^(2/3)*d)*arctan((2*c^(1/3)*x
- sqrt(3)*a^(1/6)*c^(1/6))/sqrt(a^(1/3)*c^(1/3)))/(a*c^(2/3)*sqrt(a^(1/3)*
c^(1/3)))

```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 283, normalized size of antiderivative = 0.93

$$\begin{aligned}
\int \frac{d + ex^3}{a + cx^6} dx = & -\frac{e|c| \log \left(x^2 + \left(\frac{a}{c}\right)^{\frac{1}{3}} \right)}{6 (ac^5)^{\frac{1}{3}}} + \frac{(ac^5)^{\frac{1}{6}} d \arctan \left(\frac{x}{\left(\frac{a}{c}\right)^{\frac{1}{6}}} \right)}{3 ac} \\
& + \frac{\left((ac^5)^{\frac{1}{6}} c^3 d - \sqrt{3}(ac^5)^{\frac{2}{3}} e \right) \arctan \left(\frac{2x + \sqrt{3}\left(\frac{a}{c}\right)^{\frac{1}{6}}}{\left(\frac{a}{c}\right)^{\frac{1}{6}}} \right)}{6 ac^4} \\
& + \frac{\left((ac^5)^{\frac{1}{6}} c^3 d + \sqrt{3}(ac^5)^{\frac{2}{3}} e \right) \arctan \left(\frac{2x - \sqrt{3}\left(\frac{a}{c}\right)^{\frac{1}{6}}}{\left(\frac{a}{c}\right)^{\frac{1}{6}}} \right)}{6 ac^4} \\
& + \frac{\left(\sqrt{3}(ac^5)^{\frac{1}{6}} c^3 d + (ac^5)^{\frac{2}{3}} e \right) \log \left(x^2 + \sqrt{3}x\left(\frac{a}{c}\right)^{\frac{1}{6}} + \left(\frac{a}{c}\right)^{\frac{1}{3}} \right)}{12 ac^4} \\
& - \frac{\left(\sqrt{3}(ac^5)^{\frac{1}{6}} c^3 d - (ac^5)^{\frac{2}{3}} e \right) \log \left(x^2 - \sqrt{3}x\left(\frac{a}{c}\right)^{\frac{1}{6}} + \left(\frac{a}{c}\right)^{\frac{1}{3}} \right)}{12 ac^4}
\end{aligned}$$

input `integrate((e*x^3+d)/(c*x^6+a),x, algorithm="giac")`

output

$$\begin{aligned} & -\frac{1}{6}e \cdot \text{abs}(c) \cdot \log(x^2 + (a/c)^{(1/3)}) / (a*c^5)^{(1/3)} + \frac{1}{3}*(a*c^5)^{(1/6)} * d * a \\ & \text{rctan}(x/(a/c)^{(1/6)}) / (a*c) + \frac{1}{6}*((a*c^5)^{(1/6)} * c^3 * d - \sqrt{3} * (a*c^5)^{(2/3)} * e) * \arctan((2*x + \sqrt{3} * (a/c)^{(1/6)}) / (a/c)^{(1/6)}) / (a*c^4) + \frac{1}{6}*((a*c^5)^{(1/6)} * c^3 * d + \sqrt{3} * (a*c^5)^{(2/3)} * e) * \arctan((2*x - \sqrt{3} * (a/c)^{(1/6)}) / (a/c)^{(1/6)}) / (a*c^4) + \frac{1}{12} * (\sqrt{3} * (a*c^5)^{(1/6)} * c^3 * d + (a*c^5)^{(2/3)} * e) * \log(x^2 + \sqrt{3} * x * (a/c)^{(1/6)} + (a/c)^{(1/3)}) / (a*c^4) - \frac{1}{12} * (\sqrt{3} * (a*c^5)^{(1/6)} * c^3 * d - (a*c^5)^{(2/3)} * e) * \log(x^2 - \sqrt{3} * x * (a/c)^{(1/6)} + (a/c)^{(1/3)}) / (a*c^4) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.95 (sec), antiderivative size = 1331, normalized size of antiderivative = 4.36

$$\int \frac{d + ex^3}{a + cx^6} dx = \text{Too large to display}$$

input

```
int((d + e*x^3)/(a + c*x^6),x)
```

output

$$\begin{aligned} & \log(a^3 * c^3 * (-a^4 * c^2 * e^3 + c * d^3 * (-a^5 * c^5)^{(1/2)} - 3 * a^3 * c^3 * d^2 * e - 3 * a * d * e^2 * (-a^5 * c^5)^{(1/2)}) / (a^5 * c^4))^{(1/3)} + e * x * (-a^5 * c^5)^{(1/2)} + a^2 * c^3 * d * x) * (-a^4 * c^2 * e^3 + c * d^3 * (-a^5 * c^5)^{(1/2)} - 3 * a^3 * c^3 * d^2 * e - 3 * a * d * e^2 * (-a^5 * c^5)^{(1/2)}) / (216 * a^5 * c^4))^{(1/3)} + \log(a^3 * c^3 * (-a^4 * c^2 * e^3 - c * d^3 * (-a^5 * c^5)^{(1/2)} - 3 * a^3 * c^3 * d^2 * e + 3 * a * d * e^2 * (-a^5 * c^5)^{(1/2)}) / (a^5 * c^4))^{(1/3)} - e * x * (-a^5 * c^5)^{(1/2)} + a^2 * c^3 * d * x) * (-a^4 * c^2 * e^3 - c * d^3 * (-a^5 * c^5)^{(1/2)} - 3 * a^3 * c^3 * d^2 * e + 3 * a * d * e^2 * (-a^5 * c^5)^{(1/2)}) / (216 * a^5 * c^4))^{(1/3)} - \log(a^3 * c^3 * (-a^4 * c^2 * e^3 + c * d^3 * (-a^5 * c^5)^{(1/2)} - 3 * a^3 * c^3 * d^2 * e - 3 * a * d * e^2 * (-a^5 * c^5)^{(1/2)}) / (a^5 * c^4))^{(1/3)} - 2 * e * x * (-a^5 * c^5)^{(1/2)} + 3^{(1/2)} * a^3 * c^3 * (-a^4 * c^2 * e^3 + c * d^3 * (-a^5 * c^5)^{(1/2)} - 3 * a^3 * c^3 * d^2 * e - 3 * a * d * e^2 * (-a^5 * c^5)^{(1/2)}) / (a^5 * c^4))^{(1/3)} * 1i - 2 * a^2 * c^3 * d * x) * ((3^{(1/2)} * 1i) / 2 + 1/2) * (-a^4 * c^2 * e^3 + c * d^3 * (-a^5 * c^5)^{(1/2)} - 3 * a^3 * c^3 * d^2 * e - 3 * a * d * e^2 * (-a^5 * c^5)^{(1/2)}) / (216 * a^5 * c^4))^{(1/3)} + \log(e * x * (-a^5 * c^5)^{(1/2)} - (a^3 * c^3 * (-a^4 * c^2 * e^3 + c * d^3 * (-a^5 * c^5)^{(1/2)} - 3 * a^3 * c^3 * d^2 * e - 3 * a * d * e^2 * (-a^5 * c^5)^{(1/2)}) / (a^5 * c^4))^{(1/3)}) / 2 + (3^{(1/2)} * a^3 * c^3 * (-a^4 * c^2 * e^3 + c * d^3 * (-a^5 * c^5)^{(1/2)} - 3 * a^3 * c^3 * d^2 * e - 3 * a * d * e^2 * (-a^5 * c^5)^{(1/2)}) / (a^5 * c^4))^{(1/3)} * 1i) / 2 + a^2 * c^3 * d * x) * ((3^{(1/2)} * 1i) / 2 - 1/2) * (-a^4 * c^2 * e^3 + c * d^3 * (-a^5 * c^5)^{(1/2)} - 3 * a^3 * c^3 * d^2 * e - 3 * a * d * e^2 * (-a^5 * c^5)^{(1/2)}) / (216 * a^5 * c^4))^{(1/3)} + \log(a^3 * c^3 * (-a^4 * c^2 * e^3 - c * d^3 * (-a^5 * c^5)^{(1/2)} - 3 * a^3 * c^3 * d^2 * e + 3 * a * d * e^2 * (-a^5 * c^5)^{(1/2)}) / (a^5 * c^4))^{(1/3)} \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 276, normalized size of antiderivative = 0.90

$$\int \frac{d + ex^3}{a + cx^6} dx \\ = \frac{-2\sqrt{c}\sqrt{a}\operatorname{atan}\left(\frac{c^{\frac{1}{6}}a^{\frac{1}{6}}\sqrt{3}-2c^{\frac{1}{3}}x}{c^{\frac{1}{6}}a^{\frac{1}{6}}}\right)d - 2\sqrt{3}\operatorname{atan}\left(\frac{c^{\frac{1}{6}}a^{\frac{1}{6}}\sqrt{3}-2c^{\frac{1}{3}}x}{c^{\frac{1}{6}}a^{\frac{1}{6}}}\right)ae + 2\sqrt{c}\sqrt{a}\operatorname{atan}\left(\frac{c^{\frac{1}{6}}a^{\frac{1}{6}}\sqrt{3}+2c^{\frac{1}{3}}x}{c^{\frac{1}{6}}a^{\frac{1}{6}}}\right)d - 2\sqrt{3}a^{\frac{1}{3}}e^{\frac{3}{2}}\operatorname{atan}\left(\frac{c^{\frac{1}{6}}a^{\frac{1}{6}}\sqrt{3}+2c^{\frac{1}{3}}x}{c^{\frac{1}{6}}a^{\frac{1}{6}}}\right)}{c^{\frac{1}{6}}a^{\frac{1}{6}}}$$

input `int((e*x^3+d)/(c*x^6+a),x)`

output

```
( - 2*sqrt(c)*sqrt(a)*atan((c**(1/6)*a**^(1/6)*sqrt(3) - 2*c**^(1/3)*x)/(c**^(1/6)*a**^(1/6)))*d - 2*sqrt(3)*atan((c**^(1/6)*a**^(1/6)*sqrt(3) - 2*c**^(1/3)*x)/(c**^(1/6)*a**^(1/6)))*a*e + 2*sqrt(c)*sqrt(a)*atan((c**^(1/6)*a**^(1/6)*sqrt(3) + 2*c**^(1/3)*x)/(c**^(1/6)*a**^(1/6)))*d - 2*sqrt(3)*atan((c**^(1/6)*a**^(1/6)*sqrt(3) + 2*c**^(1/3)*x)/(c**^(1/6)*a**^(1/6)))*a*e + 4*sqrt(c)*sqrt(a)*atan((c**^(1/3)*x)/(c**^(1/6)*a**^(1/6)))*d - sqrt(c)*sqrt(a)*sqrt(3)*log( - c**^(1/6)*a**^(1/6)*sqrt(3)*x + a**^(1/3) + c**^(1/3)*x**2)*d + sqrt(c)*sqrt(a)*sqrt(3)*log(c**^(1/6)*a**^(1/6)*sqrt(3)*x + a**^(1/3) + c**^(1/3)*x**2)*d - 2*log(a**^(1/3) + c**^(1/3)*x**2)*a*e + log( - c**^(1/6)*a**^(1/6)*sqrt(3)*x + a**^(1/3) + c**^(1/3)*x**2)*a*e + log(c**^(1/6)*a**^(1/6)*sqrt(3)*x + a**^(1/3) + c**^(1/3)*x**2)*a*e)/(12*c**^(2/3)*a**^(1/3)*a)
```

3.2 $\int \frac{d+ex^3}{a-cx^6} dx$

Optimal result	61
Mathematica [A] (verified)	62
Rubi [A] (verified)	62
Maple [C] (verified)	68
Fricas [B] (verification not implemented)	68
Sympy [A] (verification not implemented)	69
Maxima [A] (verification not implemented)	70
Giac [A] (verification not implemented)	71
Mupad [B] (verification not implemented)	72
Reduce [B] (verification not implemented)	73

Optimal result

Integrand size = 18, antiderivative size = 323

$$\begin{aligned} \int \frac{d+ex^3}{a-cx^6} dx = & -\frac{\left(d - \frac{\sqrt{ae}}{\sqrt{c}}\right) \arctan\left(\frac{\sqrt[6]{a}-\sqrt[6]{cx}}{\sqrt{3}\sqrt[6]{a}}\right)}{2\sqrt{3}a^{5/6}\sqrt[6]{c}} + \frac{(\sqrt{cd} + \sqrt{ae}) \arctan\left(\frac{\sqrt[6]{a}+\sqrt[6]{cx}}{\sqrt{3}\sqrt[6]{a}}\right)}{2\sqrt{3}a^{5/6}c^{2/3}} \\ & - \frac{(\sqrt{cd} + \sqrt{ae}) \log(\sqrt[6]{a} - \sqrt[6]{cx})}{6a^{5/6}c^{2/3}} + \frac{\left(d - \frac{\sqrt{ae}}{\sqrt{c}}\right) \log(\sqrt[6]{a} + \sqrt[6]{cx})}{6a^{5/6}\sqrt[6]{c}} \\ & - \frac{\left(d - \frac{\sqrt{ae}}{\sqrt{c}}\right) \log(\sqrt[3]{a} - \sqrt[6]{a}\sqrt{cx} + \sqrt[3]{cx^2})}{12a^{5/6}\sqrt[6]{c}} \\ & + \frac{(\sqrt{cd} + \sqrt{ae}) \log(\sqrt[3]{a} + \sqrt[6]{a}\sqrt{cx} + \sqrt[3]{cx^2})}{12a^{5/6}c^{2/3}} \end{aligned}$$

output

```
-1/6*(d-a^(1/2)*e/c^(1/2))*arctan(1/3*(a^(1/6)-2*c^(1/6)*x)*3^(1/2)/a^(1/6)
)*3^(1/2)/a^(5/6)/c^(1/6)+1/6*(c^(1/2)*d+a^(1/2)*e)*arctan(1/3*(a^(1/6)+2
*c^(1/6)*x)*3^(1/2)/a^(1/6))*3^(1/2)/a^(5/6)/c^(2/3)-1/6*(c^(1/2)*d+a^(1/2
)*e)*ln(a^(1/6)-c^(1/6)*x)/a^(5/6)/c^(2/3)+1/6*(d-a^(1/2)*e/c^(1/2))*ln(a^
(1/6)+c^(1/6)*x)/a^(5/6)/c^(1/6)-1/12*(d-a^(1/2)*e/c^(1/2))*ln(a^(1/3)-a^
(1/6)*c^(1/6)*x+c^(1/3)*x^2)/a^(5/6)/c^(1/6)+1/12*(c^(1/2)*d+a^(1/2)*e)*ln(
a^(1/3)+a^(1/6)*c^(1/6)*x+c^(1/3)*x^2)/a^(5/6)/c^(2/3)
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 337, normalized size of antiderivative = 1.04

$$\int \frac{d + ex^3}{a - cx^6} dx = -2\sqrt{3}(\sqrt{cd} - \sqrt{ae}) \arctan\left(\frac{1 - \frac{2\sqrt[6]{c_x}}{\sqrt{a}}}{\sqrt{3}}\right) + 2\sqrt{3}(\sqrt{cd} + \sqrt{ae}) \arctan\left(\frac{1 + \frac{2\sqrt[6]{c_x}}{\sqrt{a}}}{\sqrt{3}}\right) - 2\sqrt{cd} \log\left(\sqrt[6]{a} - \sqrt[6]{cx^6}\right)$$

input `Integrate[(d + e*x^3)/(a - c*x^6), x]`

output
$$\begin{aligned} & (-2\text{Sqrt}[3]*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*\text{ArcTan}[(1 - (2*c^{(1/6)}*x)/a^{(1/6)})/\text{Sqr}\\ & t[3]] + 2\text{Sqrt}[3]*(\text{Sqrt}[c]*d + \text{Sqrt}[a]*e)*\text{ArcTan}[(1 + (2*c^{(1/6)}*x)/a^{(1/6})/\text{Sqr}\\ & t[3]] - 2\text{Sqrt}[c]*d*\text{Log}[a^{(1/6)} - c^{(1/6)*x}] - 2\text{Sqrt}[a]*e*\text{Log}[a^{(1/}\\ & 6)} - c^{(1/6)*x}] + 2\text{Sqrt}[c]*d*\text{Log}[a^{(1/6)} + c^{(1/6)*x}] - 2\text{Sqrt}[a]*e*\text{Log}[a\\ & ^{(1/6)} + c^{(1/6)*x}] - \text{Sqrt}[c]*d*\text{Log}[a^{(1/3)} - a^{(1/6)}*c^{(1/6)*x} + c^{(1/3)*}\\ & x^{(2)}] + \text{Sqrt}[a]*e*\text{Log}[a^{(1/3)} - a^{(1/6)}*c^{(1/6)*x} + c^{(1/3)*x^{(2)}}] + \text{Sqr}\\ & t[c]*d*\text{Log}[a^{(1/3)} + a^{(1/6)}*c^{(1/6)*x} + c^{(1/3)*x^{(2)}}] + \text{Sqr}\\ & t[a]*e*\text{Log}[a^{(1/3)} + a^{(1/6)}*c^{(1/6)*x} + c^{(1/3)*x^{(2)}}])/(12*a^{(5/6)}*c^{(2/3)}) \end{aligned}$$

Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 279, normalized size of antiderivative = 0.86, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {1747, 750, 16, 27, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{d + ex^3}{a - cx^6} dx \\ & \downarrow 1747 \\ & \frac{1}{2}\left(\frac{\sqrt{ae}}{\sqrt{c}} + d\right) \int \frac{1}{a - \sqrt{a}\sqrt{c}x^3} dx + \frac{1}{2}\left(d - \frac{\sqrt{ae}}{\sqrt{c}}\right) \int \frac{1}{\sqrt{a}\sqrt{c}x^3 + a} dx \end{aligned}$$

$$\begin{aligned}
 & \downarrow \text{750} \\
 & \frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \left(\frac{\int \frac{\sqrt[6]{a}(2\sqrt[6]{a} - \sqrt[6]{cx})}{\sqrt[3]{a}\sqrt[3]{cx^2 - \sqrt{a}\sqrt[6]{cx+a^{2/3}}}} dx}{3a^{2/3}} + \frac{\int \frac{1}{\sqrt[6]{a}\sqrt[6]{cx+\sqrt[3]{a}}} dx}{3a^{2/3}} \right) + \\
 & \frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \left(\frac{\int \frac{\sqrt[6]{a}(\sqrt[6]{cx}+2\sqrt[6]{a})}{\sqrt[3]{a}\sqrt[3]{cx^2 + \sqrt{a}\sqrt[6]{cx+a^{2/3}}}} dx}{3a^{2/3}} + \frac{\int \frac{1}{\sqrt[3]{a}-\sqrt[6]{a}\sqrt[6]{cx}} dx}{3a^{2/3}} \right) \\
 & \downarrow \text{16} \\
 & \frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \left(\frac{\int \frac{\sqrt[6]{a}(2\sqrt[6]{a} - \sqrt[6]{cx})}{\sqrt[3]{a}\sqrt[3]{cx^2 - \sqrt{a}\sqrt[6]{cx+a^{2/3}}}} dx}{3a^{2/3}} + \frac{\log(\sqrt[6]{a} + \sqrt[6]{cx})}{3a^{5/6}\sqrt[6]{c}} \right) + \\
 & \frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \left(\frac{\int \frac{\sqrt[6]{a}(\sqrt[6]{cx}+2\sqrt[6]{a})}{\sqrt[3]{a}\sqrt[3]{cx^2 + \sqrt{a}\sqrt[6]{cx+a^{2/3}}}} dx}{3a^{2/3}} - \frac{\log(\sqrt[6]{a} - \sqrt[6]{cx})}{3a^{5/6}\sqrt[6]{c}} \right) \\
 & \downarrow \text{27} \\
 & \frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \left(\frac{\int \frac{2\sqrt[6]{a} - \sqrt[6]{cx}}{\sqrt[3]{a}\sqrt[3]{cx^2 - \sqrt{a}\sqrt[6]{cx+a^{2/3}}}} dx}{3\sqrt{a}} + \frac{\log(\sqrt[6]{a} + \sqrt[6]{cx})}{3a^{5/6}\sqrt[6]{c}} \right) + \\
 & \frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \left(\frac{\int \frac{\sqrt[6]{cx}+2\sqrt[6]{a}}{\sqrt[3]{a}\sqrt[3]{cx^2 + \sqrt{a}\sqrt[6]{cx+a^{2/3}}}} dx}{3\sqrt{a}} - \frac{\log(\sqrt[6]{a} - \sqrt[6]{cx})}{3a^{5/6}\sqrt[6]{c}} \right) \\
 & \downarrow \text{1142}
 \end{aligned}$$

$$\left\{ \begin{array}{l} \frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \left(\frac{\frac{3}{2} \sqrt[6]{a} \int \frac{1}{\sqrt[3]{a} \sqrt[3]{cx^2 - \sqrt{a} \sqrt[6]{cx+a^{2/3}}} dx - \frac{\int -\frac{\sqrt[3]{a} \sqrt[6]{c} (\sqrt[6]{a} - \sqrt[6]{cx})}{2 \sqrt[3]{a} \sqrt[6]{c}} dx}{3 \sqrt{a}} + \frac{\log(\sqrt[6]{a} + \sqrt[6]{cx})}{3a^{5/6} \sqrt[6]{c}} \right) + \\ \frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \left(\frac{\frac{3}{2} \sqrt[6]{a} \int \frac{1}{\sqrt[3]{a} \sqrt[3]{cx^2 + \sqrt{a} \sqrt[6]{cx+a^{2/3}}} dx + \frac{\int \frac{\sqrt[3]{a} \sqrt[6]{c} (\sqrt[6]{a} + \sqrt[6]{cx})}{2 \sqrt[3]{a} \sqrt[6]{c}} dx}{3 \sqrt{a}} - \frac{\log(\sqrt[6]{a} - \sqrt[6]{cx})}{3a^{5/6} \sqrt[6]{c}} \right) \end{array} \right.$$

↓ 25

$$\left\{ \begin{array}{l} \frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \left(\frac{\frac{3}{2} \sqrt[6]{a} \int \frac{1}{\sqrt[3]{a} \sqrt[3]{cx^2 - \sqrt{a} \sqrt[6]{cx+a^{2/3}}} dx + \frac{\int \frac{\sqrt[6]{c} (\sqrt[6]{a} - \sqrt[6]{cx})}{2 \sqrt[3]{a} \sqrt[6]{c}} dx}{3 \sqrt{a}} + \frac{\log(\sqrt[6]{a} + \sqrt[6]{cx})}{3a^{5/6} \sqrt[6]{c}} \right) + \\ \frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \left(\frac{\frac{3}{2} \sqrt[6]{a} \int \frac{1}{\sqrt[3]{a} \sqrt[3]{cx^2 + \sqrt{a} \sqrt[6]{cx+a^{2/3}}} dx + \frac{\int \frac{\sqrt[6]{c} (\sqrt[6]{a} + \sqrt[6]{cx})}{2 \sqrt[3]{a} \sqrt[6]{c}} dx}{3 \sqrt{a}} - \frac{\log(\sqrt[6]{a} - \sqrt[6]{cx})}{3a^{5/6} \sqrt[6]{c}} \right) \end{array} \right.$$

↓ 27

$$\left\{ \begin{array}{l} \frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \left(\frac{\frac{3}{2} \sqrt[6]{a} \int \frac{1}{\sqrt[3]{a} \sqrt[3]{cx^2 - \sqrt{a} \sqrt[6]{cx+a^{2/3}}} dx + \frac{\int \frac{\sqrt[6]{a} - \sqrt[6]{cx}}{2 \sqrt[3]{a}} dx}{3 \sqrt{a}} + \frac{\log(\sqrt[6]{a} + \sqrt[6]{cx})}{3a^{5/6} \sqrt[6]{c}} \right) + \\ \frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \left(\frac{\frac{3}{2} \sqrt[6]{a} \int \frac{1}{\sqrt[3]{a} \sqrt[3]{cx^2 + \sqrt{a} \sqrt[6]{cx+a^{2/3}}} dx + \frac{1}{2} \int \frac{2 \sqrt[6]{cx} + \sqrt[6]{a}}{2 \sqrt[3]{a} \sqrt[3]{cx^2 + \sqrt{a} \sqrt[6]{cx+a^{2/3}}} dx}{3 \sqrt{a}} - \frac{\log(\sqrt[6]{a} - \sqrt[6]{cx})}{3a^{5/6} \sqrt[6]{c}} \right) \end{array} \right.$$

↓ 1082

$$\frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \left\{ \begin{array}{l} \frac{\int \frac{\sqrt[6]{a} - 2\sqrt[6]{c_x}}{2\sqrt[3]{a}} dx + \frac{3 \int \frac{1}{-\left(1 - \frac{2\sqrt[6]{c_x}}{\sqrt[6]{a}}\right)^2} d\left(1 - \frac{2\sqrt[6]{c_x}}{\sqrt[6]{a}}\right)}{-\left(1 - \frac{2\sqrt[6]{c_x}}{\sqrt[6]{a}}\right)^{-3}} + \frac{\log(\sqrt[6]{a} + \sqrt[6]{c_x})}{3a^{5/6}\sqrt[6]{c}}}{3\sqrt{a}} \end{array} \right\} + \\ \frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \left\{ \begin{array}{l} \frac{\frac{1}{2} \int \frac{2\sqrt[6]{c_x} + \sqrt[6]{a}}{\sqrt[3]{a}\sqrt[3]{c_x^2 + \sqrt{a}\sqrt[6]{c_x+a^{2/3}}} dx - \frac{3 \int \frac{1}{-\left(\frac{2\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1\right)^2} d\left(\frac{2\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1\right)}{-\left(\frac{2\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1\right)^{-3}} - \frac{\log(\sqrt[6]{a} - \sqrt[6]{c_x})}{3a^{5/6}\sqrt[6]{c}}}{3\sqrt{a}} \end{array} \right\}$$

↓ 217

$$\frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \left\{ \begin{array}{l} \frac{\int \frac{\sqrt[6]{a} - 2\sqrt[6]{c_x}}{2\sqrt[3]{a}} dx - \frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[6]{c_x}}{\sqrt[6]{a}}}{\sqrt{3}}\right)}{\sqrt[3]{a}\sqrt[6]{c}} + \frac{\log(\sqrt[6]{a} + \sqrt[6]{c_x})}{3a^{5/6}\sqrt[6]{c}}}{3\sqrt{a}} \end{array} \right\} + \\ \frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \left\{ \begin{array}{l} \frac{\frac{1}{2} \int \frac{2\sqrt[6]{c_x} + \sqrt[6]{a}}{\sqrt[3]{a}\sqrt[3]{c_x^2 + \sqrt{a}\sqrt[6]{c_x+a^{2/3}}} dx + \frac{\sqrt{3} \arctan\left(\frac{2\sqrt[6]{c_x}}{\sqrt[6]{a}} + 1\right)}{\sqrt[3]{a}\sqrt[6]{c}} - \frac{\log(\sqrt[6]{a} - \sqrt[6]{c_x})}{3a^{5/6}\sqrt[6]{c}}}{3\sqrt{a}} \end{array} \right\}$$

↓ 1103

$$\frac{1}{2} \left(d - \frac{\sqrt{a}e}{\sqrt{c}} \right) \left(\begin{array}{l} \frac{\log(\sqrt[6]{a} + \sqrt[6]{c}x)}{3a^{5/6}\sqrt[6]{c}} + \frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[6]{c}x}{\sqrt[6]{a}}}{\sqrt{3}}\right)}{3\sqrt{a}} - \frac{\log(-\sqrt[6]{a}\sqrt[6]{c}x + \sqrt[3]{a} + \sqrt[3]{c}x^2)}{2\sqrt[3]{a}\sqrt[6]{c}} \\ \\ \frac{\sqrt{3} \arctan\left(\frac{\frac{2\sqrt[6]{c}x}{\sqrt[6]{a}} + 1}{\sqrt{3}}\right)}{3\sqrt{a}} + \frac{\log(\sqrt[6]{a}\sqrt[6]{c}x + \sqrt[3]{a} + \sqrt[3]{c}x^2)}{2\sqrt[3]{a}\sqrt[6]{c}} - \frac{\log(\sqrt[6]{a} - \sqrt[6]{c}x)}{3a^{5/6}\sqrt[6]{c}} \end{array} \right) +$$

input `Int[(d + e*x^3)/(a - c*x^6), x]`

output `((d - (Sqrt[a]*e)/Sqrt[c])*(Log[a^(1/6) + c^(1/6)*x]/(3*a^(5/6)*c^(1/6)) + (-((Sqrt[3]*ArcTan[(1 - (2*c^(1/6)*x)/a^(1/6))/Sqrt[3]])/(a^(1/3)*c^(1/6))) - Log[a^(1/3) - a^(1/6)*c^(1/6)*x + c^(1/3)*x^2]/(2*a^(1/3)*c^(1/6)))/(3*Sqrt[a])))/2 + ((d + (Sqrt[a]*e)/Sqrt[c])*(-1/3*Log[a^(1/6) - c^(1/6)*x]/(a^(5/6)*c^(1/6)) + ((Sqrt[3]*ArcTan[(1 + (2*c^(1/6)*x)/a^(1/6))/Sqrt[3]])/(a^(1/3)*c^(1/6)) + Log[a^(1/3) + a^(1/6)*c^(1/6)*x + c^(1/3)*x^2]/(2*a^(1/3)*c^(1/6)))/(3*Sqrt[a])))/2`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.*(x_)), x_Symbol] :> Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 27 $\text{Int}[(a_)*(F_{x_}), x_{\text{Symbol}}] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \&& \text{!Ma} \\ \text{tchQ}[F_x, (b_)*(G_x)] /; \text{FreeQ}[b, x]]$

rule 217 $\text{Int}[((a_) + (b_.)*(x_)^2)^{-1}, x_{\text{Symbol}}] \rightarrow \text{Simp}[(-(Rt[-a, 2]*Rt[-b, 2])^{(-1)}) * \text{ArcTan}[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&& \text{PosQ}[a/b] \& \\ \& (\text{LtQ}[a, 0] \mid\mid \text{LtQ}[b, 0])$

rule 750 $\text{Int}[((a_) + (b_.)*(x_)^3)^{-1}, x_{\text{Symbol}}] \rightarrow \text{Simp}[1/(3*Rt[a, 3]^2) \text{ Int}[1/ \\ (Rt[a, 3] + Rt[b, 3]*x), x], x] + \text{Simp}[1/(3*Rt[a, 3]^2) \text{ Int}[(2*Rt[a, 3] - \\ Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /; \\ \text{FreeQ}[\{a, b\}, x]$

rule 1082 $\text{Int}[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^{-1}, x_{\text{Symbol}}] \rightarrow \text{With}[\{q = 1 - 4*S \\ \text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b) \\], x] /; \text{RationalQ}[q] \&& (\text{EqQ}[q^2, 1] \mid\mid \text{!RationalQ}[b^2 - 4*a*c])] /; \text{Fre} \\ \text{eQ}[\{a, b, c\}, x]$

rule 1103 $\text{Int}[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_{\text{Symbol}}] \rightarrow \text{S} \\ \text{imp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{EqQ}[2*c*d - b*e, 0]$

rule 1142 $\text{Int}[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_{\text{Symbol}}] \rightarrow \text{S} \\ \text{imp}[(2*c*d - b*e)/(2*c) \text{ Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \\ \text{Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

rule 1747 $\text{Int}[((d_) + (e_.)*(x_)^{(n_)})/((a_) + (c_.)*(x_)^{(n2_)}), x_{\text{Symbol}}] \rightarrow \text{With}[\{ \\ q = \text{Rt}[-a/c, 2]\}, \text{Simp}[(d + e*q)/2 \text{ Int}[1/(a + c*q*x^n), x], x] + \text{Simp}[(d \\ - e*q)/2 \text{ Int}[1/(a - c*q*x^n), x], x]] /; \text{FreeQ}[\{a, c, d, e, n\}, x] \&& \text{EqQ} \\ [n2, 2*n] \&& \text{NeQ}[c*d^2 + a*e^2, 0] \&& \text{NegQ}[a*c] \&& \text{IntegerQ}[n]$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.14 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.11

method	result
risch	$-\frac{\sum_{\substack{R=\text{RootOf}(_Z^6 c-a)}} \frac{(-R^3 e+d) \ln(x-R)}{-R^5}}{6c}$
default	$\frac{e(\frac{a}{c})^{\frac{2}{3}} \ln\left(x^2+(\frac{a}{c})^{\frac{1}{6}} x+(\frac{a}{c})^{\frac{1}{3}}\right)}{12a} + \frac{e(\frac{a}{c})^{\frac{2}{3}} \sqrt{3} \arctan\left(\frac{2x\sqrt{3}}{3(\frac{a}{c})^{\frac{1}{6}}} + \frac{\sqrt{3}}{3}\right)}{6a} + \frac{d(\frac{a}{c})^{\frac{1}{6}} \ln\left(x^2+(\frac{a}{c})^{\frac{1}{6}} x+(\frac{a}{c})^{\frac{1}{3}}\right)}{12a} + \frac{d(\frac{a}{c})^{\frac{1}{6}} \sqrt{3} \arctan\left(\frac{2x\sqrt{3}}{3(\frac{a}{c})^{\frac{1}{6}}} + \frac{\sqrt{3}}{3}\right)}{6a}$

input `int((e*x^3+d)/(-c*x^6+a),x,method=_RETURNVERBOSE)`

output `-1/6/c*sum((_R^3*e+d)/_R^5*ln(x-_R),_R=RootOf(_Z^6*c-a))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1613 vs. $2(223) = 446$.

Time = 0.15 (sec) , antiderivative size = 1613, normalized size of antiderivative = 4.99

$$\int \frac{d+ex^3}{a-cx^6} dx = \text{Too large to display}$$

input `integrate((e*x^3+d)/(-c*x^6+a),x, algorithm="fricas")`

output

$$\begin{aligned}
 & -\frac{1}{12} \left(\sqrt{-3} + 1 \right) \left(-\left(a^2 c^2 \sqrt{(c^2 d^6 + 6 a c d^4 e^2 + 9 a^2 d^2 e^4) / (a^5 c^3)} + 3 c d^2 e + a e^3 \right) / (a^2 c^2) \right)^{(1/3)} \log \left(-(c^2 d^5 + 2 a c d^3 e^2 - 3 a^2 d e^4) x + \frac{1}{2} (a c^2 d^4 + 3 a^2 c d^2 e^2 + \sqrt{-3}) \left(a c^2 d^4 + 3 a^2 c d^2 e^2 \right) - (\sqrt{-3} a^4 c^2 e + a^4 c^2 e) \sqrt{(c^2 d^6 + 6 a c d^4 e^2 + 9 a^2 d^2 e^4) / (a^5 c^3)} \right) \left(-\left(a^2 c^2 \sqrt{(c^2 d^6 + 6 a c d^4 e^2 + 9 a^2 d^2 e^4) / (a^5 c^3)} + 3 c d^2 e + a e^3 \right) / (a^2 c^2) \right)^{(1/3)} + \frac{1}{12} \left(\sqrt{-3} - 1 \right) \left(-\left(a^2 c^2 \sqrt{(c^2 d^6 + 6 a c d^4 e^2 + 9 a^2 d^2 e^4) / (a^5 c^3)} + 3 c d^2 e + a e^3 \right) / (a^2 c^2) \right)^{(1/3)} \log \left(-(c^2 d^5 + 2 a c d^3 e^2 - 3 a^2 d e^4) x + \frac{1}{2} (a c^2 d^4 + 3 a^2 c d^2 e^2 - \sqrt{-3}) \left(a c^2 d^4 + 3 a^2 c d^2 e^2 \right) + (\sqrt{-3} a^4 c^2 e - a^4 c^2 e) \sqrt{(c^2 d^6 + 6 a c d^4 e^2 + 9 a^2 d^2 e^4) / (a^5 c^3)} \right) \left(-\left(a^2 c^2 \sqrt{(c^2 d^6 + 6 a c d^4 e^2 + 9 a^2 d^2 e^4) / (a^5 c^3)} + 3 c d^2 e + a e^3 \right) / (a^2 c^2) \right)^{(1/3)} - \frac{1}{12} \left(\sqrt{-3} + 1 \right) \left(\left(a^2 c^2 \sqrt{(c^2 d^6 + 6 a c d^4 e^2 + 9 a^2 d^2 e^4) / (a^5 c^3)} + 3 c d^2 e + a e^3 \right) / (a^2 c^2) \right)^{(1/3)} \log \left(-(c^2 d^5 + 2 a c d^3 e^2 - 3 a^2 d e^4) x + \frac{1}{2} (a c^2 d^4 + 3 a^2 c d^2 e^2 + \sqrt{-3}) \left(a c^2 d^4 + 3 a^2 c d^2 e^2 \right) + (\sqrt{-3} a^4 c^2 e + a^4 c^2 e) \sqrt{(c^2 d^6 + 6 a c d^4 e^2 + 9 a^2 d^2 e^4) / (a^5 c^3)} \right) \left(\left(a^2 c^2 \sqrt{(c^2 d^6 + 6 a c d^4 e^2 + 9 a^2 d^2 e^4) / (a^5 c^3)} + 3 c d^2 e + a e^3 \right) / (a^2 c^2) \right)^{(1/3)} + \frac{1}{12} \left(\sqrt{-3} - 1 \right) \left(\left(a^2 c^2 \sqrt{(c^2 d^6 + 6 a c d^4 e^2 + 9 a^2 d^2 e^4) / (a^5 c^3)} + 3 c d^2 e + a e^3 \right) / (a^2 c^2) \right)^{(1/3)} \log \left(-(c^2 d^5 + 2 a c d^3 e^2 - 3 a^2 d e^4) x + \frac{1}{2} (a c^2 d^4 + 3 a^2 c d^2 e^2 - \sqrt{-3}) \left(a c^2 d^4 + 3 a^2 c d^2 e^2 \right) + (\sqrt{-3} a^4 c^2 e - a^4 c^2 e) \sqrt{(c^2 d^6 + 6 a c d^4 e^2 + 9 a^2 d^2 e^4) / (a^5 c^3)} \right) \left(\left(a^2 c^2 \sqrt{(c^2 d^6 + 6 a c d^4 e^2 + 9 a^2 d^2 e^4) / (a^5 c^3)} + 3 c d^2 e + a e^3 \right) / (a^2 c^2) \right)^{(1/3)} - 3 c d^2 e - a e^3 / (a^2 c^2) \dots
 \end{aligned}$$
Sympy [A] (verification not implemented)

Time = 1.85 (sec), antiderivative size = 168, normalized size of antiderivative = 0.52

$$\begin{aligned}
 & \int \frac{d + e x^3}{a - c x^6} dx = \\
 & -\text{RootSum} \left(46656 t^6 a^5 c^4 + t^3 (-432 a^4 c^2 e^3 - 1296 a^3 c^3 d^2 e) + a^3 e^6 - 3 a^2 c d^2 e^4 + 3 a c^2 d^4 e^2 - c^3 d^6, \left(t \mapsto \right. \right.
 \end{aligned}$$

input

```
integrate((e*x**3+d)/(-c*x**6+a),x)
```

output

```

-RootSum(46656*_t**6*a**5*c**4 + _t**3*(-432*a**4*c**2*e**3 - 1296*a**3*c**3*d**2*e) + a**3*e**6 - 3*a**2*c*d**2*e**4 + 3*a*c**2*d**4*e**2 - c**3*d**6, Lambda(_t, _t*log(x + (-1296*_t**4*a**4*c**2*e + 6*_t*a**3*e**4 + 36*_t*a**2*c*d**2*e**2 + 6*_t*a*c**2*d**4)/(3*a**2*d**4 - 2*a*c*d**3*e**2 - c**2*d**5)))

```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 313, normalized size of antiderivative = 0.97

$$\int \frac{d + ex^3}{a - cx^6} dx = \frac{\sqrt{3}(\sqrt{cd} + \sqrt{ae}) \arctan\left(\frac{\sqrt{3}\left(2x + \left(\frac{\sqrt{a}}{\sqrt{c}}\right)^{\frac{1}{3}}\right)}{3\left(\frac{\sqrt{a}}{\sqrt{c}}\right)^{\frac{1}{3}}}\right)}{6\sqrt{ac}\left(\frac{\sqrt{a}}{\sqrt{c}}\right)^{\frac{2}{3}}} \\ + \frac{\sqrt{3}(\sqrt{cd} - \sqrt{ae}) \arctan\left(\frac{\sqrt{3}\left(2x - \left(\frac{\sqrt{a}}{\sqrt{c}}\right)^{\frac{1}{3}}\right)}{3\left(\frac{\sqrt{a}}{\sqrt{c}}\right)^{\frac{1}{3}}}\right)}{6\sqrt{ac}\left(\frac{\sqrt{a}}{\sqrt{c}}\right)^{\frac{2}{3}}} \\ + \frac{(\sqrt{cd} + \sqrt{ae}) \log\left(x^2 + x\left(\frac{\sqrt{a}}{\sqrt{c}}\right)^{\frac{1}{3}} + \left(\frac{\sqrt{a}}{\sqrt{c}}\right)^{\frac{2}{3}}\right)}{12\sqrt{ac}\left(\frac{\sqrt{a}}{\sqrt{c}}\right)^{\frac{2}{3}}} \\ - \frac{(\sqrt{cd} - \sqrt{ae}) \log\left(x^2 - x\left(\frac{\sqrt{a}}{\sqrt{c}}\right)^{\frac{1}{3}} + \left(\frac{\sqrt{a}}{\sqrt{c}}\right)^{\frac{2}{3}}\right)}{12\sqrt{ac}\left(\frac{\sqrt{a}}{\sqrt{c}}\right)^{\frac{2}{3}}} \\ + \frac{(\sqrt{cd} - \sqrt{ae}) \log\left(x + \left(\frac{\sqrt{a}}{\sqrt{c}}\right)^{\frac{1}{3}}\right)}{6\sqrt{ac}\left(\frac{\sqrt{a}}{\sqrt{c}}\right)^{\frac{2}{3}}} - \frac{(\sqrt{cd} + \sqrt{ae}) \log\left(x - \left(\frac{\sqrt{a}}{\sqrt{c}}\right)^{\frac{1}{3}}\right)}{6\sqrt{ac}\left(\frac{\sqrt{a}}{\sqrt{c}}\right)^{\frac{2}{3}}}$$

```
input integrate((e*x^3+d)/(-c*x^6+a),x, algorithm="maxima")
```

output

```
1/6*sqrt(3)*(sqrt(c)*d + sqrt(a)*e)*arctan(1/3*sqrt(3)*(2*x + (sqrt(a)/sqrt(c))^(1/3))/(sqrt(a)/sqrt(c))^(1/3))/(sqrt(a)*c*(sqrt(a)/sqrt(c))^(2/3))
+ 1/6*sqrt(3)*(sqrt(c)*d - sqrt(a)*e)*arctan(1/3*sqrt(3)*(2*x - (sqrt(a)/sqrt(c))^(1/3))/(sqrt(a)/sqrt(c))^(1/3))/(sqrt(a)*c*(sqrt(a)/sqrt(c))^(2/3))
) + 1/12*(sqrt(c)*d + sqrt(a)*e)*log(x^2 + x*(sqrt(a)/sqrt(c))^(1/3) + (sqrt(a)/sqrt(c))^(2/3))/(sqrt(a)*c*(sqrt(a)/sqrt(c))^(2/3)) - 1/12*(sqrt(c)*d - sqrt(a)*e)*log(x^2 - x*(sqrt(a)/sqrt(c))^(1/3) + (sqrt(a)/sqrt(c))^(2/3))/(sqrt(a)*c*(sqrt(a)/sqrt(c))^(2/3)) + 1/6*(sqrt(c)*d - sqrt(a)*e)*log(x + (sqrt(a)/sqrt(c))^(1/3))/(sqrt(a)*c*(sqrt(a)/sqrt(c))^(2/3)) - 1/6*(sqrt(c)*d + sqrt(a)*e)*log(x - (sqrt(a)/sqrt(c))^(1/3))/(sqrt(a)*c*(sqrt(a)/sqrt(c))^(2/3))
```

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 303, normalized size of antiderivative = 0.94

$$\int \frac{d + ex^3}{a - cx^6} dx = \frac{e|c| \log \left(x^2 + \left(-\frac{a}{c} \right)^{\frac{1}{3}} \right)}{6 (-ac^5)^{\frac{1}{3}}} + \frac{(-ac^5)^{\frac{1}{6}} d \arctan \left(\frac{x}{(-\frac{a}{c})^{\frac{1}{6}}} \right)}{3 ac} \\ + \frac{\left((-ac^5)^{\frac{1}{6}} c^3 d - \sqrt{3}(-ac^5)^{\frac{2}{3}} e \right) \arctan \left(\frac{2x + \sqrt{3}(-\frac{a}{c})^{\frac{1}{6}}}{(-\frac{a}{c})^{\frac{1}{6}}} \right)}{6 ac^4} \\ + \frac{\left((-ac^5)^{\frac{1}{6}} c^3 d + \sqrt{3}(-ac^5)^{\frac{2}{3}} e \right) \arctan \left(\frac{2x - \sqrt{3}(-\frac{a}{c})^{\frac{1}{6}}}{(-\frac{a}{c})^{\frac{1}{6}}} \right)}{6 ac^4} \\ + \frac{\left(\sqrt{3}(-ac^5)^{\frac{1}{6}} c^3 d + (-ac^5)^{\frac{2}{3}} e \right) \log \left(x^2 + \sqrt{3}x(-\frac{a}{c})^{\frac{1}{6}} + (-\frac{a}{c})^{\frac{1}{3}} \right)}{12 ac^4} \\ - \frac{\left(\sqrt{3}(-ac^5)^{\frac{1}{6}} c^3 d - (-ac^5)^{\frac{2}{3}} e \right) \log \left(x^2 - \sqrt{3}x(-\frac{a}{c})^{\frac{1}{6}} + (-\frac{a}{c})^{\frac{1}{3}} \right)}{12 ac^4}$$

input `integrate((e*x^3+d)/(-c*x^6+a),x, algorithm="giac")`

output

```
1/6*e*abs(c)*log(x^2 + (-a/c)^(1/3))/(-a*c^5)^(1/3) + 1/3*(-a*c^5)^(1/6)*d
*arctan(x/(-a/c)^(1/6))/(a*c) + 1/6*((-a*c^5)^(1/6)*c^3*d - sqrt(3)*(-a*c^5)^(2/3)*e)*arctan((2*x + sqrt(3)*(-a/c)^(1/6))/(-a/c)^(1/6))/(a*c^4) + 1/
6*((-a*c^5)^(1/6)*c^3*d + sqrt(3)*(-a*c^5)^(2/3)*e)*arctan((2*x - sqrt(3)*(-a/c)^(1/6))/(-a/c)^(1/6))/(a*c^4) + 1/12*(sqrt(3)*(-a*c^5)^(1/6)*c^3*d +
(-a*c^5)^(2/3)*e)*log(x^2 + sqrt(3)*x*(-a/c)^(1/6) + (-a/c)^(1/3))/(a*c^4) - 1/12*(sqrt(3)*(-a*c^5)^(1/6)*c^3*d - (-a*c^5)^(2/3)*e)*log(x^2 - sqrt(3)*x*(-a/c)^(1/6) + (-a/c)^(1/3))/(a*c^4)
```

Mupad [B] (verification not implemented)

Time = 11.90 (sec), antiderivative size = 1293, normalized size of antiderivative = 4.00

$$\int \frac{d + ex^3}{a - cx^6} dx = \text{Too large to display}$$

input

```
int((d + e*x^3)/(a - c*x^6),x)
```

output

```
log(a^3*c^3*(-(a^4*c^2*e^3 + c*d^3*(a^5*c^5)^(1/2) + 3*a^3*c^3*d^2*e + 3*a
*d*e^2*(a^5*c^5)^(1/2))/(a^5*c^4))^(1/3) + e*x*(a^5*c^5)^(1/2) + a^2*c^3*d
*x)*(-(a^4*c^2*e^3 + c*d^3*(a^5*c^5)^(1/2) + 3*a^3*c^3*d^2*e + 3*a*d*e^2*(a^5*c^5)^(1/2))/(216*a^5*c^4))^(1/3) + log(a^3*c^3*(-(a^4*c^2*e^3 - c*d^3*(a^5*c^5)^(1/2) + 3*a^3*c^3*d^2*e - 3*a*d*e^2*(a^5*c^5)^(1/2))/(a^5*c^4))^(1/3) - e*x*(a^5*c^5)^(1/2) + a^2*c^3*d*x)*(-(a^4*c^2*e^3 - c*d^3*(a^5*c^5)^(1/2) + 3*a^3*c^3*d^2*e - 3*a*d*e^2*(a^5*c^5)^(1/2))/(216*a^5*c^4))^(1/3) - log(a^3*c^3*(-(a^4*c^2*e^3 + c*d^3*(a^5*c^5)^(1/2) + 3*a^3*c^3*d^2*e + 3*a*d*e^2*(a^5*c^5)^(1/2))/(a^5*c^4))^(1/3) - 2*e*x*(a^5*c^5)^(1/2) + 3^(1/2)*a^3*c^3*(-(a^4*c^2*e^3 + c*d^3*(a^5*c^5)^(1/2) + 3*a^3*c^3*d^2*e + 3*a*d*e^2*(a^5*c^5)^(1/2))/(a^5*c^4))^(1/3)*1i - 2*a^2*c^3*d*x)*((3^(1/2)*1i)/2 + 1/2)*(-(a^4*c^2*e^3 + c*d^3*(a^5*c^5)^(1/2) + 3*a^3*c^3*d^2*e + 3*a*d*e^2*(a^5*c^5)^(1/2))/(216*a^5*c^4))^(1/3) + log(e*x*(a^5*c^5)^(1/2) - (a^3*c^3*(-(a^4*c^2*e^3 + c*d^3*(a^5*c^5)^(1/2) + 3*a^3*c^3*d^2*e + 3*a*d*e^2*(a^5*c^5)^(1/2))/(a^5*c^4))^(1/3))/2 + (3^(1/2)*a^3*c^3*(-(a^4*c^2*e^3 + c*d^3*(a^5*c^5)^(1/2) + 3*a^3*c^3*d^2*e + 3*a*d*e^2*(a^5*c^5)^(1/2))/(a^5*c^4))^(1/3)*1i)/2 + a^2*c^3*d*x)*((3^(1/2)*1i)/2 - 1/2)*(-(a^4*c^2*e^3 + c*d^3*(a^5*c^5)^(1/2) + 3*a^3*c^3*d^2*e + 3*a*d*e^2*(a^5*c^5)^(1/2))/(216*a^5*c^4))^(1/3) + log(a^3*c^3*(-(a^4*c^2*e^3 - c*d^3*(a^5*c^5)^(1/2) + 3*a^3*c^3*d^2*e - 3*a*d*e^2*(a^5*c^5)^(1/2))/(a^5*c^4))^(1/3) + 2*e*x*(a^5...)
```

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 328, normalized size of antiderivative = 1.02

$$\int \frac{d + ex^3}{a - cx^6} dx \\ = \frac{-2\sqrt{c}\sqrt{a}\sqrt{3}\operatorname{atan}\left(\frac{c^{\frac{1}{6}}a^{\frac{1}{6}}-2c^{\frac{1}{3}}x}{c^{\frac{1}{6}}a^{\frac{1}{6}}\sqrt{3}}\right)d + 2\sqrt{3}\operatorname{atan}\left(\frac{c^{\frac{1}{6}}a^{\frac{1}{6}}-2c^{\frac{1}{3}}x}{c^{\frac{1}{6}}a^{\frac{1}{6}}\sqrt{3}}\right)ae + 2\sqrt{c}\sqrt{a}\sqrt{3}\operatorname{atan}\left(\frac{c^{\frac{1}{6}}a^{\frac{1}{6}}+2c^{\frac{1}{3}}x}{c^{\frac{1}{6}}a^{\frac{1}{6}}\sqrt{3}}\right)d + 2\sqrt{3}a^{\frac{1}{2}}e^{\frac{3}{2}}x^{\frac{3}{2}}}{a^{\frac{7}{6}}}$$

input `int((e*x^3+d)/(-c*x^6+a),x)`

output

```
( - 2*sqrt(c)*sqrt(a)*sqrt(3)*atan((c**1/6*a**1/6) - 2*c**1/3*x)/(c**1/6*a**1/6)*sqrt(3))*d + 2*sqrt(3)*atan((c**1/6*a**1/6) - 2*c**1/3*x)/(c**1/6*a**1/6)*sqrt(3))*a*e + 2*sqrt(c)*sqrt(a)*sqrt(3)*atan((c*(1/6)*a**1/6) + 2*c**1/3*x)/(c**1/6*a**1/6)*sqrt(3))*d + 2*sqrt(3)*atan((c**1/6*a**1/6) + 2*c**1/3*x)/(c**1/6*a**1/6)*sqrt(3))*a*e - sqrt(c)*sqrt(a)*log( - c**1/6*a**1/6*x + a**1/3) + c**1/3*x**2)*d + 2*sqrt(c)*sqrt(a)*log( - c**1/6*a**1/6) - c**1/3*x)*d + sqrt(c)*sqrt(a)*log(c**1/6*a**1/6*x + a**1/3) + c**1/3*x**2)*d - 2*sqrt(c)*sqrt(a)*log(c**1/6*a**1/6) - c**1/3*x)*d + log( - c**1/6*a**1/6*x + a**1/3) + c**1/3*x**2)*a*e - 2*log( - c**1/6*a**1/6) - c**1/3*x)*a*e + log(c**1/6*a**1/6*x + a**1/3) + c**1/3*x**2)*a*e - 2*log(c**1/6*a**1/6) - c**1/3*x)*a*e)/(12*c**2/3*a**1/3*a)
```

$$3.3 \quad \int \frac{(d+ex^3)^{3/2}}{a+cx^6} dx$$

Optimal result	74
Mathematica [F]	74
Rubi [A] (verified)	75
Maple [C] (warning: unable to verify)	76
Fricas [F(-1)]	77
Sympy [F]	78
Maxima [F]	78
Giac [F]	78
Mupad [F(-1)]	79
Reduce [F]	79

Optimal result

Integrand size = 21, antiderivative size = 142

$$\begin{aligned} \int \frac{(d+ex^3)^{3/2}}{a+cx^6} dx &= \frac{dx\sqrt{d+ex^3} \operatorname{AppellF1}\left(\frac{1}{3}, -\frac{3}{2}, 1, \frac{4}{3}, -\frac{ex^3}{d}, -\frac{\sqrt{cx^3}}{\sqrt{-a}}\right)}{2a\sqrt{1+\frac{ex^3}{d}}} \\ &+ \frac{dx\sqrt{d+ex^3} \operatorname{AppellF1}\left(\frac{1}{3}, -\frac{3}{2}, 1, \frac{4}{3}, -\frac{ex^3}{d}, \frac{\sqrt{cx^3}}{\sqrt{-a}}\right)}{2a\sqrt{1+\frac{ex^3}{d}}} \end{aligned}$$

output $\frac{1/2*d*x*(e*x^3+d)^(1/2)*\operatorname{AppellF1}(1/3, 1, -3/2, 4/3, -c^(1/2)*x^3/(-a)^(1/2), -e*x^3/d)/a/(1+e*x^3/d)^(1/2)+1/2*d*x*(e*x^3+d)^(1/2)*\operatorname{AppellF1}(1/3, 1, -3/2, 4/3, c^(1/2)*x^3/(-a)^(1/2), -e*x^3/d)/a/(1+e*x^3/d)^(1/2)}$

Mathematica [F]

$$\int \frac{(d+ex^3)^{3/2}}{a+cx^6} dx = \int \frac{(d+ex^3)^{3/2}}{a+cx^6} dx$$

input `Integrate[(d + e*x^3)^(3/2)/(a + c*x^6), x]`

output Integrate[(d + e*x^3)^(3/2)/(a + c*x^6), x]

Rubi [A] (verified)

Time = 0.31 (sec), antiderivative size = 142, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.190, Rules used = {1759, 27, 937, 936}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d + ex^3)^{3/2}}{a + cx^6} dx \\
 & \quad \downarrow \textcolor{blue}{1759} \\
 & - \frac{\sqrt{c} \int \frac{(ex^3+d)^{3/2}}{\sqrt{c}(\sqrt{-a}-\sqrt{cx^3})} dx}{2\sqrt{-a}} - \frac{\sqrt{c} \int \frac{(ex^3+d)^{3/2}}{\sqrt{c}(\sqrt{cx^3}+\sqrt{-a})} dx}{2\sqrt{-a}} \\
 & \quad \downarrow \textcolor{blue}{27} \\
 & - \frac{\int \frac{(ex^3+d)^{3/2}}{\sqrt{-a}-\sqrt{cx^3}} dx}{2\sqrt{-a}} - \frac{\int \frac{(ex^3+d)^{3/2}}{\sqrt{cx^3}+\sqrt{-a}} dx}{2\sqrt{-a}} \\
 & \quad \downarrow \textcolor{blue}{937} \\
 & - \frac{d\sqrt{d+ex^3} \int \frac{\left(\frac{ex^3}{d}+1\right)^{3/2}}{\sqrt{-a}-\sqrt{cx^3}} dx}{2\sqrt{-a}\sqrt{\frac{ex^3}{d}+1}} - \frac{d\sqrt{d+ex^3} \int \frac{\left(\frac{ex^3}{d}+1\right)^{3/2}}{\sqrt{cx^3}+\sqrt{-a}} dx}{2\sqrt{-a}\sqrt{\frac{ex^3}{d}+1}} \\
 & \quad \downarrow \textcolor{blue}{936} \\
 & \frac{dx\sqrt{d+ex^3} \operatorname{AppellF1}\left(\frac{1}{3}, 1, -\frac{3}{2}, \frac{4}{3}, -\frac{\sqrt{cx^3}}{\sqrt{-a}}, -\frac{ex^3}{d}\right)}{2a\sqrt{\frac{ex^3}{d}+1}} + \\
 & \frac{dx\sqrt{d+ex^3} \operatorname{AppellF1}\left(\frac{1}{3}, 1, -\frac{3}{2}, \frac{4}{3}, \frac{\sqrt{cx^3}}{\sqrt{-a}}, -\frac{ex^3}{d}\right)}{2a\sqrt{\frac{ex^3}{d}+1}}
 \end{aligned}$$

input Int[(d + e*x^3)^(3/2)/(a + c*x^6), x]

output

$$(d*x*sqrt[d + e*x^3]*AppellF1[1/3, 1, -3/2, 4/3, -(sqrt[c]*x^3)/sqrt[-a], -((e*x^3)/d)]/(2*a*sqrt[1 + (e*x^3)/d]) + (d*x*sqrt[d + e*x^3]*AppellF1[1/3, 1, -3/2, 4/3, (sqrt[c]*x^3)/sqrt[-a], -((e*x^3)/d)])/(2*a*sqrt[1 + (e*x^3)/d]))$$

Definitions of rubi rules used

rule 27

$$\text{Int}[(a_)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \& \text{ !MatchQ}[F_x, (b_)*(G_x_) /; \text{FreeQ}[b, x]]$$

rule 936

$$\text{Int}[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] \rightarrow \text{Simp}[a^p*c^q*x*AppellF1[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /; \text{FreeQ}[\{a, b, c, d, n, p, q\}, x] \& \text{NeQ}[b*c - a*d, 0] \& \text{NeQ}[n, -1] \& \text{ && (IntegerQ[p] || GtQ[a, 0]) \& (IntegerQ[q] || GtQ[c, 0])}$$

rule 937

$$\text{Int}[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] \rightarrow \text{Simp}[a^{\text{IntPart}[p]}*((a + b*x^n)^{\text{FracPart}[p]}/(1 + b*(x^n/a))^{\text{FracPart}[p]}) \text{ Int}[(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /; \text{FreeQ}[\{a, b, c, d, n, p, q\}, x] \& \text{NeQ}[b*c - a*d, 0] \& \text{NeQ}[n, -1] \& \text{ ! (IntegerQ[p] || GtQ[a, 0])}$$

rule 1759

$$\text{Int}[((d_) + (e_)*(x_)^(n_))^(q_)/((a_) + (c_)*(x_)^(n2_)), x_Symbol] \rightarrow \text{With}[\{r = \text{Rt}[-a*c, 2]\}, \text{Simp}[-c/(2*r) \text{ Int}[(d + e*x^n)^q/(r - c*x^n), x], x] - \text{Simp}[c/(2*r) \text{ Int}[(d + e*x^n)^q/(r + c*x^n), x], x]] /; \text{FreeQ}[\{a, c, d, e, n, q\}, x] \& \text{EqQ}[n2, 2*n] \& \text{NeQ}[c*d^2 + a*e^2, 0] \& \text{ ! IntegerQ[q]}$$

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 6.

Time = 0.72 (sec) , antiderivative size = 902, normalized size of antiderivative = 6.35

method	result	size
default	Expression too large to display	902
elliptic	Expression too large to display	902

input `int((e*x^3+d)^(3/2)/(c*x^6+a),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & -\frac{2}{3} \cdot \frac{I \cdot e \cdot c \cdot 3^{(1/2)} \cdot (-d \cdot e^2)^{(1/3)} \cdot (I \cdot (x+1/2) \cdot e \cdot (-d \cdot e^2)^{(1/3)} - 1/2 \cdot I \cdot 3^{(1/2)} \\ & \cdot e \cdot (-d \cdot e^2)^{(1/3)}) \cdot 3^{(1/2)} \cdot e \cdot (-d \cdot e^2)^{(1/3)})^{(1/2)} \cdot ((x-1) \cdot e \cdot (-d \cdot e^2)^{(1/3)}) \\ & / (-3/2 \cdot e \cdot (-d \cdot e^2)^{(1/3)} + 1/2 \cdot I \cdot 3^{(1/2)} \cdot e \cdot (-d \cdot e^2)^{(1/3)})^{(1/2)} \cdot (-I \cdot (x+1/2) \\ & \cdot e \cdot (-d \cdot e^2)^{(1/3)} + 1/2 \cdot I \cdot 3^{(1/2)} \cdot e \cdot (-d \cdot e^2)^{(1/3)}) \cdot 3^{(1/2)} \cdot e \cdot (-d \cdot e^2)^{(1/3)}) \\ & ^{(1/2)} / (e \cdot x^3 + d)^{(1/2)} \cdot \text{EllipticF}(1/3 \cdot 3^{(1/2)} \cdot (I \cdot (x+1/2) \cdot e \cdot (-d \cdot e^2)^{(1/3)} - 1/2 \\ & \cdot I \cdot 3^{(1/2)} \cdot e \cdot (-d \cdot e^2)^{(1/3)}) \cdot 3^{(1/2)} \cdot e \cdot (-d \cdot e^2)^{(1/3)})^{(1/2)}, (I \cdot 3^{(1/2)} \cdot e \\ & \cdot (-d \cdot e^2)^{(1/3)} / (-3/2 \cdot e \cdot (-d \cdot e^2)^{(1/3)} + 1/2 \cdot I \cdot 3^{(1/2)} \cdot e \cdot (-d \cdot e^2)^{(1/3)}))^{(1/2)} \\ & - 1/6 \cdot I \cdot c \cdot e^2 \cdot 2^{(1/2)} \cdot \text{sum}((-2 \cdot \text{_alpha}^3 \cdot c \cdot d \cdot e + a \cdot e^2 \cdot c \cdot d^2) / \text{_alpha}^5 / (a \cdot e \\ & ^2 + c \cdot d^2) \cdot (-d \cdot e^2)^{(1/3)} \cdot (1/2 \cdot I \cdot e \cdot (2 \cdot x + 1) \cdot e \cdot (-I \cdot 3^{(1/2)} \cdot (-d \cdot e^2)^{(1/3)} + (-d \cdot \\ & e^2)^{(1/3)}) / (-d \cdot e^2)^{(1/3)})^{(1/2)} \cdot (e \cdot (x-1) \cdot e \cdot (-d \cdot e^2)^{(1/3)}) / (-3 \cdot (-d \cdot e^2)^{(1/3)} + I \cdot 3^{(1/2)} \cdot (-d \cdot e^2)^{(1/3)}) \\ & \cdot (-d \cdot e^2)^{(1/3)}))^{(1/2)} \cdot (-1/2 \cdot I \cdot e \cdot (2 \cdot x + 1) \cdot e \cdot (I \cdot 3^{(1/2)} \cdot (-d \cdot e^2)^{(1/3)} + (-d \cdot e^2)^{(1/3)} + (-d \cdot e^2)^{(1/3)})) / (-d \cdot e^2)^{(1/3)})^{(1/2)} / (e \cdot x^3 + d)^{(1/2)} \cdot (2 \cdot e^2 \cdot (-\text{_alpha}^5 \cdot e + \text{_alpha}^2 \cdot d) - I \cdot (-d \cdot e^2)^{(1/3)} \cdot \text{_alpha}^4 \cdot 3^{(1/2)} \cdot e \cdot e^2 + I \cdot \text{_alpha}^3 \cdot 3^{(1/2)} \cdot (-d \cdot e^2)^{(2/3)} \cdot e + (-d \cdot e^2)^{(1/3)} \cdot \text{_alpha}^4 \cdot e^2 + I \cdot (-d \cdot e^2)^{(1/3)} \cdot d \cdot \text{_alpha}^3 \cdot 3^{(1/2)} \cdot e + \text{_alpha}^3 \cdot (-d \cdot e^2)^{(2/3)} \cdot d) \cdot \text{EllipticPi}(1/3 \cdot 3^{(1/2)} \cdot (I \cdot (x+1/2) \cdot e \cdot (-d \cdot e^2)^{(1/3)} - 1/2 \cdot I \cdot 3^{(1/2)} \cdot e \cdot (-d \cdot e^2)^{(1/3)}) \cdot 3^{(1/2)} \cdot e \cdot (-d \cdot e^2)^{(1/3)})^{(1/2)}, -1/2 \cdot c \cdot e \cdot (-2 \cdot I \cdot 3^{(1/2)} \cdot (-d \cdot e^2)^{(1/3)}) \cdot \text{_alpha}^5 \cdot e^2 + I \cdot 3^{(1/2)} \cdot (-d \cdot e^2)^{(2/3)} \cdot \text{_alpha}^4 \cdot e - I \cdot 3^{(1/2)} \cdot \text{_alpha}^3 \cdot d + 3 \cdot \text{_alpha}^2 \cdot d \cdot e + 3 \cdot (-d \cdot e^2)^{(2/3)} \cdot \text{_alpha}^4 \cdot e - I \cdot 3^{(1/2)} \cdot (-d \cdot e^2)^{(1/3)} \cdot \text{_alpha}^4 \cdot e + (2/3) \cdot \text{_alpha}^4 \cdot d + 3 \cdot \text{_alpha}^3 \cdot d + 3 \cdot \text{_alpha}^2 \cdot d^2 + 3 \cdot (-d \cdot e^2)^{(1/3)} \cdot \text{_alpha}^4 \cdot d + 3 \cdot (-d \cdot e^2)^{(1/3)} \cdot \text{_alpha}^3 \cdot d^2 + 3 \cdot (-d \cdot e^2)^{(1/3)} \cdot \text{_alpha}^2 \cdot d^3 + 3 \cdot (-d \cdot e^2)^{(1/3)} \cdot \text{_alpha}^1 \cdot d^4 + 3 \cdot (-d \cdot e^2)^{(1/3)} \cdot \text{_alpha}^0 \cdot d^5) \dots \end{aligned}$$

Fricas [F(-1)]

Timed out.

$$\int \frac{(d + ex^3)^{3/2}}{a + cx^6} dx = \text{Timed out}$$

input `integrate((e*x^3+d)^(3/2)/(c*x^6+a),x, algorithm="fricas")`

output Timed out

Sympy [F]

$$\int \frac{(d + ex^3)^{3/2}}{a + cx^6} dx = \int \frac{(d + ex^3)^{\frac{3}{2}}}{a + cx^6} dx$$

input `integrate((e*x**3+d)**(3/2)/(c*x**6+a),x)`

output `Integral((d + e*x**3)**(3/2)/(a + c*x**6), x)`

Maxima [F]

$$\int \frac{(d + ex^3)^{3/2}}{a + cx^6} dx = \int \frac{(ex^3 + d)^{\frac{3}{2}}}{cx^6 + a} dx$$

input `integrate((e*x^3+d)^(3/2)/(c*x^6+a),x, algorithm="maxima")`

output `integrate((e*x^3 + d)^(3/2)/(c*x^6 + a), x)`

Giac [F]

$$\int \frac{(d + ex^3)^{3/2}}{a + cx^6} dx = \int \frac{(ex^3 + d)^{\frac{3}{2}}}{cx^6 + a} dx$$

input `integrate((e*x^3+d)^(3/2)/(c*x^6+a),x, algorithm="giac")`

output `integrate((e*x^3 + d)^(3/2)/(c*x^6 + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(d + ex^3)^{3/2}}{a + cx^6} dx = \int \frac{(e x^3 + d)^{3/2}}{c x^6 + a} dx$$

input `int((d + e*x^3)^(3/2)/(a + c*x^6),x)`

output `int((d + e*x^3)^(3/2)/(a + c*x^6), x)`

Reduce [F]

$$\int \frac{(d + ex^3)^{3/2}}{a + cx^6} dx = \left(\int \frac{\sqrt{e x^3 + d}}{c x^6 + a} dx \right) d + \left(\int \frac{\sqrt{e x^3 + d} x^3}{c x^6 + a} dx \right) e$$

input `int((e*x^3+d)^(3/2)/(c*x^6+a),x)`

output `int(sqrt(d + e*x**3)/(a + c*x**6),x)*d + int((sqrt(d + e*x**3)*x**3)/(a + c*x**6),x)*e`

3.4 $\int \frac{\sqrt{d+ex^3}}{a+cx^6} dx$

Optimal result	80
Mathematica [C] (warning: unable to verify)	80
Rubi [A] (verified)	81
Maple [C] (warning: unable to verify)	83
Fricas [F(-1)]	84
Sympy [F]	84
Maxima [F]	85
Giac [F]	85
Mupad [F(-1)]	85
Reduce [F]	86

Optimal result

Integrand size = 21, antiderivative size = 140

$$\int \frac{\sqrt{d+ex^3}}{a+cx^6} dx = \frac{x\sqrt{d+ex^3} \operatorname{AppellF1}\left(\frac{1}{3}, -\frac{1}{2}, 1, \frac{4}{3}, -\frac{ex^3}{d}, -\frac{\sqrt{cx^3}}{\sqrt{-a}}\right)}{2a\sqrt{1+\frac{ex^3}{d}}} \\ + \frac{x\sqrt{d+ex^3} \operatorname{AppellF1}\left(\frac{1}{3}, -\frac{1}{2}, 1, \frac{4}{3}, -\frac{ex^3}{d}, \frac{\sqrt{cx^3}}{\sqrt{-a}}\right)}{2a\sqrt{1+\frac{ex^3}{d}}}$$

output
$$\frac{1/2*x*(e*x^3+d)^(1/2)*\operatorname{AppellF1}(1/3, 1, -1/2, 4/3, -c^(1/2)*x^3/(-a)^(1/2), -e*x^3/d)/a/(1+e*x^3/d)^(1/2)+1/2*x*(e*x^3+d)^(1/2)*\operatorname{AppellF1}(1/3, 1, -1/2, 4/3, c^(1/2)*x^3/(-a)^(1/2), -e*x^3/d)/a/(1+e*x^3/d)^(1/2)}$$

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 21.19 (sec) , antiderivative size = 5647, normalized size of antiderivative = 40.34

$$\int \frac{\sqrt{d+ex^3}}{a+cx^6} dx = \text{Result too large to show}$$

input `Integrate[Sqrt[d + e*x^3]/(a + c*x^6), x]`

output `Result too large to show`

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {1759, 27, 937, 936}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{d+ex^3}}{a+cx^6} dx \\
 & \downarrow \textcolor{blue}{1759} \\
 & -\frac{\sqrt{c} \int \frac{\sqrt{ex^3+d}}{\sqrt{c}(\sqrt{-a}-\sqrt{cx^3})} dx}{2\sqrt{-a}} - \frac{\sqrt{c} \int \frac{\sqrt{ex^3+d}}{\sqrt{c}(\sqrt{cx^3}+\sqrt{-a})} dx}{2\sqrt{-a}} \\
 & \downarrow \textcolor{blue}{27} \\
 & -\frac{\int \frac{\sqrt{ex^3+d}}{\sqrt{-a}-\sqrt{cx^3}} dx}{2\sqrt{-a}} - \frac{\int \frac{\sqrt{ex^3+d}}{\sqrt{cx^3}+\sqrt{-a}} dx}{2\sqrt{-a}} \\
 & \downarrow \textcolor{blue}{937} \\
 & -\frac{\sqrt{d+ex^3} \int \frac{\sqrt{\frac{ex^3}{d}+1}}{\sqrt{-a}-\sqrt{cx^3}} dx}{2\sqrt{-a}\sqrt{\frac{ex^3}{d}+1}} - \frac{\sqrt{d+ex^3} \int \frac{\sqrt{\frac{ex^3}{d}+1}}{\sqrt{cx^3}+\sqrt{-a}} dx}{2\sqrt{-a}\sqrt{\frac{ex^3}{d}+1}} \\
 & \downarrow \textcolor{blue}{936} \\
 & \frac{x\sqrt{d+ex^3} \operatorname{AppellF1}\left(\frac{1}{3}, 1, -\frac{1}{2}, \frac{4}{3}, -\frac{\sqrt{cx^3}}{\sqrt{-a}}, -\frac{ex^3}{d}\right)}{2a\sqrt{\frac{ex^3}{d}+1}} + \\
 & \frac{x\sqrt{d+ex^3} \operatorname{AppellF1}\left(\frac{1}{3}, 1, -\frac{1}{2}, \frac{4}{3}, \frac{\sqrt{cx^3}}{\sqrt{-a}}, -\frac{ex^3}{d}\right)}{2a\sqrt{\frac{ex^3}{d}+1}}
 \end{aligned}$$

input $\text{Int}[\text{Sqrt}[d + e*x^3]/(a + c*x^6), x]$

output $(x*\text{Sqrt}[d + e*x^3]*\text{AppellF1}[1/3, 1, -1/2, 4/3, -((\text{Sqrt}[c]*x^3)/\text{Sqrt}[-a]), -((e*x^3)/d)])/(2*a*\text{Sqrt}[1 + (e*x^3)/d]) + (x*\text{Sqrt}[d + e*x^3]*\text{AppellF1}[1/3, 1, -1/2, 4/3, (\text{Sqrt}[c]*x^3)/\text{Sqrt}[-a], -((e*x^3)/d)])/(2*a*\text{Sqrt}[1 + (e*x^3)/d])$

Definitions of rubi rules used

rule 27 $\text{Int}[(a_)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \&& \text{!MatchQ}[F_x, (b_)*(G_x_)] /; \text{FreeQ}[b, x]]$

rule 936 $\text{Int}[((a_) + (b_.)*(x_)^(n_.))^(p_.)*((c_) + (d_.)*(x_)^(n_.))^(q_), x_Symbol] \rightarrow \text{Simp}[a^p*c^q*x^q*\text{AppellF1}[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /; \text{FreeQ}[\{a, b, c, d, n, p, q\}, x] \&& \text{NeQ}[b*c - a*d, 0] \&& \text{NeQ}[n, -1] \&& (\text{IntegerQ}[p] \text{ || } \text{GtQ}[a, 0]) \&& (\text{IntegerQ}[q] \text{ || } \text{GtQ}[c, 0])]$

rule 937 $\text{Int}[((a_) + (b_.)*(x_)^(n_.))^(p_.)*((c_) + (d_.)*(x_)^(n_.))^(q_), x_Symbol] \rightarrow \text{Simp}[a^p*\text{IntPart}[p]*((a + b*x^n)^p*\text{FracPart}[p]/(1 + b*(x^n/a))^q*\text{FracPart}[p]) \text{ Int}[(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /; \text{FreeQ}[\{a, b, c, d, n, p, q\}, x] \&& \text{NeQ}[b*c - a*d, 0] \&& \text{NeQ}[n, -1] \&& \text{!(IntegerQ}[p] \text{ || } \text{GtQ}[a, 0])]$

rule 1759 $\text{Int}[((d_) + (e_.)*(x_)^(n_.))^(q_)/((a_) + (c_.)*(x_)^(n2_)), x_Symbol] \rightarrow \text{With}[\{r = \text{Rt}[-(a*c), 2]\}, \text{Simp}[-c/(2*r) \text{ Int}[(d + e*x^n)^q/(r - c*x^n), x], x] - \text{Simp}[c/(2*r) \text{ Int}[(d + e*x^n)^q/(r + c*x^n), x], x]] /; \text{FreeQ}[\{a, c, d, e, n, q\}, x] \&& \text{EqQ}[n2, 2*n] \&& \text{NeQ}[c*d^2 + a*e^2, 0] \&& \text{!IntegerQ}[q]]$

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 6.

Time = 0.35 (sec) , antiderivative size = 605, normalized size of antiderivative = 4.32

method	result
default	$\frac{i\sqrt{2} \sum_{\alpha=\text{RootOf}(_Z^6 c+a)} \left(-e^{-\alpha^3-d} (-d e^2)^{\frac{1}{3}} \sqrt{2} \sqrt{\frac{ie \left(2x + \frac{-i\sqrt{3}(-d e^2)^{\frac{1}{3}} + (-d e^2)^{\frac{1}{3}}}{e} \right)}{(-d e^2)^{\frac{1}{3}}}} \sqrt{\frac{e \left(x - \frac{(-d e^2)^{\frac{1}{3}}}{e} \right)}{-3(-d e^2)^{\frac{1}{3}} + i\sqrt{3}(-d e^2)^{\frac{1}{3}}}} \sqrt{-\frac{ie \left(x - \frac{(-d e^2)^{\frac{1}{3}}}{e} \right)}{(-d e^2)^{\frac{1}{3}}}} \right)}{(-e^{-\alpha^3-d} (-d e^2)^{\frac{1}{3}} \sqrt{2})^{\frac{1}{3}}}$
elliptic	$\frac{i\sqrt{2} \sum_{\alpha=\text{RootOf}(_Z^6 c+a)} \left(-e^{-\alpha^3-d} (-d e^2)^{\frac{1}{3}} \sqrt{2} \sqrt{\frac{ie \left(2x + \frac{-i\sqrt{3}(-d e^2)^{\frac{1}{3}} + (-d e^2)^{\frac{1}{3}}}{e} \right)}{(-d e^2)^{\frac{1}{3}}}} \sqrt{\frac{e \left(x - \frac{(-d e^2)^{\frac{1}{3}}}{e} \right)}{-3(-d e^2)^{\frac{1}{3}} + i\sqrt{3}(-d e^2)^{\frac{1}{3}}}} \sqrt{-\frac{ie \left(x - \frac{(-d e^2)^{\frac{1}{3}}}{e} \right)}{(-d e^2)^{\frac{1}{3}}}} \right)}{(-e^{-\alpha^3-d} (-d e^2)^{\frac{1}{3}} \sqrt{2})^{\frac{1}{3}}}$

input `int((e*x^3+d)^(1/2)/(c*x^6+a),x,method=_RETURNVERBOSE)`

output

```

-1/6*I/e^2*2^(1/2)*sum((-_alpha^3*e-d)/_alpha^5/(a*e^2+c*d^2)*(-d*e^2)^(1/3)*(1/2*I*e*(2*x+1/e*(-I*3^(1/2)*(-d*e^2)^(1/3)+(-d*e^2)^(1/3)))/(-d*e^2)^(1/3))^(1/2)*(e*(x-1/e*(-d*e^2)^(1/3))/(-3*(-d*e^2)^(1/3)+I*3^(1/2)*(-d*e^2)^(1/3)))^(1/2)*(-1/2*I*e*(2*x+1/e*(I*3^(1/2)*(-d*e^2)^(1/3)+(-d*e^2)^(1/3)))/(-d*e^2)^(1/3))/(e*x^3+d)^(1/2)*(2*e^2*(-_alpha^5*e+_alpha^2*d)-I*(-d*e^2)^(1/3)*_alpha^4*3^(1/2)*e^2+I*_alpha^3*3^(1/2)*(-d*e^2)^(2/3)*e+(-d*e^2)^(1/3)*_alpha^4*e^2+I*(-d*e^2)^(1/3)*d*_alpha*3^(1/2)*e+_alpha^3*(-d*e^2)^(2/3)*e-I*3^(1/2)*(-d*e^2)^(2/3)*d-(-d*e^2)^(1/3)*d*_alpha*e-(-d*e^2)^(2/3)*d)*EllipticPi(1/3*3^(1/2)*(I*(x+1/2/e*(-d*e^2)^(1/3)-1/2*I*3^(1/2)/e*(-d*e^2)^(1/3))*3^(1/2)*e/(-d*e^2)^(1/3))^(1/2), -1/2*c/e*(-2*I*3^(1/2)*(-d*e^2)^(1/3)*_alpha^5*e^2+I*3^(1/2)*(-d*e^2)^(2/3)*_alpha^4*e-I*3^(1/2)*_alpha^3*d*e^2+2*I*3^(1/2)*(-d*e^2)^(1/3)*_alpha^2*d*e+3*(-d*e^2)^(2/3)*_alpha^4*e-I*3^(1/2)*(-d*e^2)^(2/3)*_alpha*d+3*_alpha^3*d*e^2+I*3^(1/2)*d^2*2*e-3*(-d*e^2)^(2/3)*_alpha*d-3*d^2*e)/(a*e^2+c*d^2), (I*3^(1/2)/e*(-d*e^2)^(1/3)/(-3/2/e*(-d*e^2)^(1/3)+1/2*I*3^(1/2)/e*(-d*e^2)^(1/3)))^(1/2)), _alpha=RootOf(_Z^6*c+a))

```

Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{d + ex^3}}{a + cx^6} dx = \text{Timed out}$$

input `integrate((e*x^3+d)^(1/2)/(c*x^6+a),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{\sqrt{d + ex^3}}{a + cx^6} dx = \int \frac{\sqrt{d + ex^3}}{a + cx^6} dx$$

input `integrate((e*x**3+d)**(1/2)/(c*x**6+a),x)`

output `Integral(sqrt(d + e*x**3)/(a + c*x**6), x)`

Maxima [F]

$$\int \frac{\sqrt{d + ex^3}}{a + cx^6} dx = \int \frac{\sqrt{ex^3 + d}}{cx^6 + a} dx$$

input `integrate((e*x^3+d)^(1/2)/(c*x^6+a),x, algorithm="maxima")`

output `integrate(sqrt(e*x^3 + d)/(c*x^6 + a), x)`

Giac [F]

$$\int \frac{\sqrt{d + ex^3}}{a + cx^6} dx = \int \frac{\sqrt{ex^3 + d}}{cx^6 + a} dx$$

input `integrate((e*x^3+d)^(1/2)/(c*x^6+a),x, algorithm="giac")`

output `integrate(sqrt(e*x^3 + d)/(c*x^6 + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{d + ex^3}}{a + cx^6} dx = \int \frac{\sqrt{ex^3 + d}}{cx^6 + a} dx$$

input `int((d + e*x^3)^(1/2)/(a + c*x^6),x)`

output `int((d + e*x^3)^(1/2)/(a + c*x^6), x)`

Reduce [F]

$$\int \frac{\sqrt{d + ex^3}}{a + cx^6} dx = \int \frac{\sqrt{e x^3 + d}}{c x^6 + a} dx$$

input `int((e*x^3+d)^(1/2)/(c*x^6+a),x)`

output `int(sqrt(d + e*x**3)/(a + c*x**6),x)`

3.5 $\int \frac{1}{\sqrt{d+ex^3}(a+cx^6)} dx$

Optimal result	87
Mathematica [C] (warning: unable to verify)	87
Rubi [A] (verified)	88
Maple [C] (warning: unable to verify)	90
Fricas [F(-1)]	91
Sympy [F]	92
Maxima [F]	92
Giac [F]	92
Mupad [F(-1)]	93
Reduce [F]	93

Optimal result

Integrand size = 21, antiderivative size = 140

$$\begin{aligned} \int \frac{1}{\sqrt{d+ex^3}(a+cx^6)} dx &= \frac{x\sqrt{1+\frac{ex^3}{d}} \operatorname{AppellF1}\left(\frac{1}{3}, \frac{1}{2}, 1, \frac{4}{3}, -\frac{ex^3}{d}, -\frac{\sqrt{cx^3}}{\sqrt{-a}}\right)}{2a\sqrt{d+ex^3}} \\ &+ \frac{x\sqrt{1+\frac{ex^3}{d}} \operatorname{AppellF1}\left(\frac{1}{3}, \frac{1}{2}, 1, \frac{4}{3}, -\frac{ex^3}{d}, \frac{\sqrt{cx^3}}{\sqrt{-a}}\right)}{2a\sqrt{d+ex^3}} \end{aligned}$$

output
$$\frac{1/2*x*(1+e*x^3/d)^(1/2)*\operatorname{AppellF1}(1/3, 1, 1/2, 4/3, -c^(1/2)*x^3/(-a)^(1/2), -e*x^3/d)/a/(e*x^3+d)^(1/2)+1/2*x*(1+e*x^3/d)^(1/2)*\operatorname{AppellF1}(1/3, 1, 1/2, 4/3, c^(1/2)*x^3/(-a)^(1/2), -e*x^3/d)/a/(e*x^3+d)^(1/2)}$$

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 19.89 (sec) , antiderivative size = 932, normalized size of antiderivative = 6.66

$$\int \frac{1}{\sqrt{d+ex^3}(a+cx^6)} dx = \text{Too large to display}$$

input
$$\text{Integrate}[1/(\text{Sqrt}[d + e*x^3]*(a + c*x^6)), x]$$

output

$$(2*(-1)^(2/3)*d^(1/3)*Sqrt[(d^(1/3) + e^(1/3)*x)/((1 + (-1)^(1/3))*d^(1/3))]*Sqrt[1 - (e^(1/3)*x)/d^(1/3) + (e^(2/3)*x^2)/d^(2/3)]*(((-1)^(2/3)*EllipticPi[((-1)^(1/6)*(1 + (-1)^(1/3))*c^(1/6)*d^(1/3))/((-1)^(1/6)*c^(1/6)*d^(1/3) - a^(1/6)*e^(1/3)), ArcSin[Sqrt[(d^(1/3) + (-1)^(2/3)*e^(1/3)*x)/((1 + (-1)^(1/3))*d^(1/3))], (-1)^(1/3)]])/(-2*a^(2/3)*c^(1/6)*d^(1/3) + (-I + Sqrt[3])*a^(5/6)*e^(1/3)) - EllipticPi[((1 + (-1)^(1/3))*c^(1/6)*d^(1/3))/(c^(1/6)*d^(1/3) - I*a^(1/6)*e^(1/3)), ArcSin[Sqrt[(d^(1/3) + (-1)^(2/3))*e^(1/3)*x]/((1 + (-1)^(1/3))*d^(1/3))], (-1)^(1/3)]/((I + Sqrt[3])*a^(2/3)*(I*c^(1/6)*d^(1/3) + a^(1/6)*e^(1/3))) + (3*EllipticPi[(I*Sqrt[3])*c^(1/6)*d^(1/3))/((-1)^(1/3)*c^(1/6)*d^(1/3) - I*a^(1/6)*e^(1/3)), ArcSin[Sqrt[(d^(1/3) + (-1)^(2/3)*e^(1/3)*x)/((1 + (-1)^(1/3))*d^(1/3))], (-1)^(1/3)]/(6*a^(2/3)*c^(1/6)*d^(1/3) - 3*(I + Sqrt[3])*a^(5/6)*e^(1/3)) - EllipticPi[((1 + (-1)^(1/3))*c^(1/6)*d^(1/3))/(c^(1/6)*d^(1/3) + I*a^(1/6)*e^(1/3)), ArcSin[Sqrt[(d^(1/3) + (-1)^(2/3)*e^(1/3)*x)/((1 + (-1)^(1/3))*d^(1/3))], (-1)^(1/3)]/((I + Sqrt[3])*a^(2/3)*((-I)*c^(1/6)*d^(1/3) + a^(1/6)*e^(1/3))) - EllipticPi[(I*Sqrt[3])*c^(1/6)*d^(1/3))/((-1)^(1/3)*c^(1/6)*d^(1/3) + I*a^(1/6)*e^(1/3)), ArcSin[Sqrt[(d^(1/3) + (-1)^(2/3)*e^(1/3)*x)/((1 + (-1)^(1/3))*d^(1/3))], (-1)^(1/3)]/(2*a^(2/3)*c^(1/6)*d^(1/3) + (I + Sqrt[3])*a^(5/6)*e^(1/3)) + ((-1)^(2/3)*EllipticPi[((-1)^(1/6)*(1 + (-1)^(1/3))*c^(1/6)*d^(1/3))/((-1)^(1/6)*c^(1/6)*d^(1/3) + a^(1/6)*e^(1/3)), Ar...]$$

Rubi [A] (verified)

Time = 0.29 (sec), antiderivative size = 140, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.190, Rules used = {1759, 27, 937, 936}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(a + cx^6)\sqrt{d + ex^3}} dx \\ & \downarrow \textcolor{blue}{1759} \\ & -\frac{\sqrt{c} \int \frac{1}{\sqrt{c}(\sqrt{-a} - \sqrt{cx^3})\sqrt{ex^3 + d}} dx}{2\sqrt{-a}} - \frac{\sqrt{c} \int \frac{1}{\sqrt{c}(\sqrt{cx^3} + \sqrt{-a})\sqrt{ex^3 + d}} dx}{2\sqrt{-a}} \\ & \downarrow \textcolor{blue}{27} \end{aligned}$$

$$\begin{aligned}
 & -\frac{\int \frac{1}{(\sqrt{-a}-\sqrt{cx^3})\sqrt{ex^3+d}} dx}{2\sqrt{-a}} - \frac{\int \frac{1}{(\sqrt{cx^3}+\sqrt{-a})\sqrt{ex^3+d}} dx}{2\sqrt{-a}} \\
 & \quad \downarrow 937 \\
 & -\frac{\sqrt{\frac{ex^3}{d}+1} \int \frac{1}{(\sqrt{-a}-\sqrt{cx^3})\sqrt{\frac{ex^3}{d}+1}} dx}{2\sqrt{-a}\sqrt{d+ex^3}} - \frac{\sqrt{\frac{ex^3}{d}+1} \int \frac{1}{(\sqrt{cx^3}+\sqrt{-a})\sqrt{\frac{ex^3}{d}+1}} dx}{2\sqrt{-a}\sqrt{d+ex^3}} \\
 & \quad \downarrow 936 \\
 & \frac{x\sqrt{\frac{ex^3}{d}+1} \operatorname{AppellF1}\left(\frac{1}{3}, 1, \frac{1}{2}, \frac{4}{3}, -\frac{\sqrt{cx^3}}{\sqrt{-a}}, -\frac{ex^3}{d}\right)}{2a\sqrt{d+ex^3}} + \frac{x\sqrt{\frac{ex^3}{d}+1} \operatorname{AppellF1}\left(\frac{1}{3}, 1, \frac{1}{2}, \frac{4}{3}, \frac{\sqrt{cx^3}}{\sqrt{-a}}, -\frac{ex^3}{d}\right)}{2a\sqrt{d+ex^3}}
 \end{aligned}$$

input `Int[1/(Sqrt[d + e*x^3]*(a + c*x^6)), x]`

output `(x*Sqrt[1 + (e*x^3)/d]*AppellF1[1/3, 1, 1/2, 4/3, -((Sqrt[c]*x^3)/Sqrt[-a]) - ((e*x^3)/d)])/(2*a*Sqrt[d + e*x^3]) + (x*Sqrt[1 + (e*x^3)/d]*AppellF1[1/3, 1, 1/2, 4/3, (Sqrt[c]*x^3)/Sqrt[-a], -((e*x^3)/d)])/(2*a*Sqrt[d + e*x^3])`

Definitions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] :> Simplify[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 936 `Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] :> Simplify[a^p*c^q*x^AppellF1[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`

rule 937 `Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] :> Simplify[a^p*IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p]) Int[(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && !(IntegerQ[p] || GtQ[a, 0])`

rule 1759

```
Int[((d_) + (e_)*(x_)^(n_))^(q_)/((a_) + (c_)*(x_)^(n2_)), x_Symbol] :> W
ith[{r = Rt[(-a)*c, 2]}, Simpl[-c/(2*r) Int[(d + e*x^n)^q/(r - c*x^n), x],
x] - Simpl[c/(2*r) Int[(d + e*x^n)^q/(r + c*x^n), x], x]] /; FreeQ[{a, c,
d, e, n, q}, x] && EqQ[n2, 2*n] && NeQ[c*d^2 + a*e^2, 0] && !IntegerQ[q]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 6.

Time = 0.32 (sec) , antiderivative size = 595, normalized size of antiderivative = 4.25

method	result
default	$\frac{i\sqrt{2} \sum_{-\alpha=\text{RootOf}(-Z^6 c+a)} \left(\frac{\left(ie \left(2x + \frac{-i\sqrt{3} (-de^2)^{\frac{1}{3}} + (-de^2)^{\frac{1}{3}}}{e} \right) \right)^{\frac{1}{3}}}{(-de^2)^{\frac{1}{3}}} \sqrt{\frac{e \left(x - \frac{(-de^2)^{\frac{1}{3}}}{e} \right)}{-3(-de^2)^{\frac{1}{3}} + i\sqrt{3} (-de^2)^{\frac{1}{3}}}} \sqrt{-\frac{ie \left(2x + \frac{i\sqrt{3} (-de^2)^{\frac{1}{3}}}{2(-de^2)^{\frac{1}{3}}} \right)}{2(-de^2)^{\frac{1}{3}}}} \right)^{\frac{1}{3}}}{(-de^2)^{\frac{1}{3}} \sqrt{2}}$
elliptic	$\frac{i\sqrt{2} \sum_{-\alpha=\text{RootOf}(-Z^6 c+a)} \left(\frac{\left(ie \left(2x + \frac{-i\sqrt{3} (-de^2)^{\frac{1}{3}} + (-de^2)^{\frac{1}{3}}}{e} \right) \right)^{\frac{1}{3}}}{(-de^2)^{\frac{1}{3}}} \sqrt{\frac{e \left(x - \frac{(-de^2)^{\frac{1}{3}}}{e} \right)}{-3(-de^2)^{\frac{1}{3}} + i\sqrt{3} (-de^2)^{\frac{1}{3}}}} \sqrt{-\frac{ie \left(2x + \frac{i\sqrt{3} (-de^2)^{\frac{1}{3}}}{2(-de^2)^{\frac{1}{3}}} \right)}{2(-de^2)^{\frac{1}{3}}}} \right)^{\frac{1}{3}}}{(-de^2)^{\frac{1}{3}} \sqrt{2}}$

input `int(1/(e*x^3+d)^(1/2)/(c*x^6+a),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & \frac{1}{6} \cdot \frac{I}{e^2} \cdot 2^{\frac{1}{2}} \cdot \sum \left(\frac{1}{\alpha} \cdot e^{2+c \cdot d^2} \cdot (-d \cdot e^2)^{-\frac{1}{3}} \cdot (1/2 \cdot I \cdot e^{(2 \cdot x+1)/e} \cdot (-I \cdot 3^{\frac{1}{2}} \cdot (-d \cdot e^2)^{\frac{1}{3}} + (-d \cdot e^2)^{\frac{1}{3}})) / (-d \cdot e^2)^{\frac{1}{3}} \right)^{\frac{1}{2}} \cdot (e^{(x-1)/e} \cdot (-d \cdot e^2)^{\frac{1}{3}}) / (-3 \cdot (-d \cdot e^2)^{\frac{1}{3}} + I \cdot 3^{\frac{1}{2}} \cdot (-d \cdot e^2)^{\frac{1}{3}}))^{\frac{1}{2}} \cdot (-\frac{1}{2} \cdot I \cdot e^{(2 \cdot x+1)/e} \cdot (I \cdot 3^{\frac{1}{2}} \cdot (-d \cdot e^2)^{\frac{1}{3}} + (-d \cdot e^2)^{\frac{1}{3}})) / (-d \cdot e^2)^{\frac{1}{3}})^{\frac{1}{2}} / (e \cdot x^3 + d)^{\frac{1}{2}} \cdot (2 \cdot e^2 \cdot (-\alpha^5 \cdot e + \alpha^2 \cdot d) - I \cdot (-d \cdot e^2)^{\frac{1}{3}}) \cdot \alpha^4 \cdot 3^{\frac{1}{2}} \cdot e^2 + I \cdot \alpha^3 \cdot 3^{\frac{1}{2}} \cdot (-d \cdot e^2)^{\frac{2}{3}} \cdot e + (-d \cdot e^2)^{\frac{1}{3}} \cdot \alpha^4 \cdot e^2 + I \cdot (-d \cdot e^2)^{\frac{1}{3}} \cdot d \cdot \alpha^3 \cdot 3^{\frac{1}{2}} \cdot e + \alpha^3 \cdot (-d \cdot e^2)^{\frac{2}{3}} \cdot e - I \cdot 3^{\frac{1}{2}} \cdot (-d \cdot e^2)^{\frac{2}{3}} \cdot d - (-d \cdot e^2)^{\frac{1}{3}} \cdot d \cdot \alpha^2 \cdot e - (-d \cdot e^2)^{\frac{2}{3}} \cdot d \cdot E \cdot \text{ellipticPi}(1/3 \cdot 3^{\frac{1}{2}} \cdot (I \cdot (x+1/2) \cdot e \cdot (-d \cdot e^2)^{\frac{1}{3}} - 1/2 \cdot I \cdot 3^{\frac{1}{2}}) / e \cdot (-d \cdot e^2)^{\frac{1}{3}}) \cdot 3^{\frac{1}{2}} \cdot e / (-d \cdot e^2)^{\frac{1}{3}})^{\frac{1}{2}}, -1/2 \cdot c \cdot e \cdot (-2 \cdot I \cdot 3^{\frac{1}{2}} \cdot (-d \cdot e^2)^{\frac{1}{3}}) \cdot \alpha^5 \cdot e^2 + I \cdot 3^{\frac{1}{2}} \cdot (-d \cdot e^2)^{\frac{2}{3}} \cdot \alpha^4 \cdot e - I \cdot 3^{\frac{1}{2}} \cdot (-d \cdot e^2)^{\frac{1}{3}} \cdot \alpha^2 \cdot d + 3 \cdot \alpha^3 \cdot d \cdot e^2 + I \cdot 3^{\frac{1}{2}} \cdot (-d \cdot e^2)^{\frac{2}{3}} \cdot \alpha^2 \cdot d \cdot e + 3 \cdot (-d \cdot e^2)^{\frac{1}{3}} \cdot \alpha^3 \cdot d \cdot e - I \cdot 3^{\frac{1}{2}} \cdot (-d \cdot e^2)^{\frac{2}{3}} \cdot \alpha^4 \cdot e - I \cdot 3^{\frac{1}{2}} \cdot (-d \cdot e^2)^{\frac{1}{3}} \cdot \alpha^2 \cdot d - 3 \cdot d^2 \cdot e) / (a \cdot e^2 + c \cdot d^2), (I \cdot 3^{\frac{1}{2}} / e \cdot (-d \cdot e^2)^{\frac{1}{3}}) / (-3/2 \cdot e \cdot (-d \cdot e^2)^{\frac{1}{3}} + 1/2 \cdot I \cdot 3^{\frac{1}{2}} / e \cdot (-d \cdot e^2)^{\frac{1}{3}})^{\frac{1}{2}}), \alpha = \text{RootOf}(_Z^6 \cdot c + a) \right) \end{aligned}$$

Fricas [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{d + ex^3} (a + cx^6)} dx = \text{Timed out}$$

input `integrate(1/(e*x^3+d)^(1/2)/(c*x^6+a),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{1}{\sqrt{d + ex^3} (a + cx^6)} dx = \int \frac{1}{(a + cx^6) \sqrt{d + ex^3}} dx$$

input `integrate(1/(e*x**3+d)**(1/2)/(c*x**6+a),x)`

output `Integral(1/((a + c*x**6)*sqrt(d + e*x**3)), x)`

Maxima [F]

$$\int \frac{1}{\sqrt{d + ex^3} (a + cx^6)} dx = \int \frac{1}{(cx^6 + a)\sqrt{ex^3 + d}} dx$$

input `integrate(1/(e*x^3+d)^(1/2)/(c*x^6+a),x, algorithm="maxima")`

output `integrate(1/((c*x^6 + a)*sqrt(e*x^3 + d)), x)`

Giac [F]

$$\int \frac{1}{\sqrt{d + ex^3} (a + cx^6)} dx = \int \frac{1}{(cx^6 + a)\sqrt{ex^3 + d}} dx$$

input `integrate(1/(e*x^3+d)^(1/2)/(c*x^6+a),x, algorithm="giac")`

output `integrate(1/((c*x^6 + a)*sqrt(e*x^3 + d)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{d + ex^3} (a + cx^6)} dx = \int \frac{1}{(cx^6 + a) \sqrt{ex^3 + d}} dx$$

input `int(1/((a + c*x^6)*(d + e*x^3)^(1/2)),x)`

output `int(1/((a + c*x^6)*(d + e*x^3)^(1/2)), x)`

Reduce [F]

$$\int \frac{1}{\sqrt{d + ex^3} (a + cx^6)} dx = \int \frac{\sqrt{ex^3 + d}}{ce x^9 + cd x^6 + ae x^3 + ad} dx$$

input `int(1/(e*x^3+d)^(1/2)/(c*x^6+a),x)`

output `int(sqrt(d + e*x**3)/(a*d + a*e*x**3 + c*d*x**6 + c*e*x**9),x)`

3.6 $\int \frac{1}{(d+ex^3)^{3/2}(a+cx^6)} dx$

Optimal result	94
Mathematica [C] (warning: unable to verify)	94
Rubi [B] (warning: unable to verify)	95
Maple [C] (warning: unable to verify)	97
Fricas [F(-1)]	98
Sympy [F]	99
Maxima [F]	99
Giac [F]	99
Mupad [F(-1)]	100
Reduce [F]	100

Optimal result

Integrand size = 21, antiderivative size = 146

$$\int \frac{1}{(d+ex^3)^{3/2}(a+cx^6)} dx = \frac{x\sqrt{1+\frac{ex^3}{d}} \operatorname{AppellF1}\left(\frac{1}{3}, \frac{3}{2}, 1, \frac{4}{3}, -\frac{ex^3}{d}, -\frac{\sqrt{cx^3}}{\sqrt{-a}}\right)}{2ad\sqrt{d+ex^3}} \\ + \frac{x\sqrt{1+\frac{ex^3}{d}} \operatorname{AppellF1}\left(\frac{1}{3}, \frac{3}{2}, 1, \frac{4}{3}, -\frac{ex^3}{d}, \frac{\sqrt{cx^3}}{\sqrt{-a}}\right)}{2ad\sqrt{d+ex^3}}$$

output $1/2*x*(1+e*x^3/d)^(1/2)*\operatorname{AppellF1}(1/3, 1, 3/2, 4/3, -c^(1/2)*x^3/(-a)^(1/2), -e*x^3/d)/a/d/(e*x^3+d)^(1/2)+1/2*x*(1+e*x^3/d)^(1/2)*\operatorname{AppellF1}(1/3, 1, 3/2, 4/3, c^(1/2)*x^3/(-a)^(1/2), -e*x^3/d)/a/d/(e*x^3+d)^(1/2)$

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 21.28 (sec) , antiderivative size = 5979, normalized size of antiderivative = 40.95

$$\int \frac{1}{(d+ex^3)^{3/2}(a+cx^6)} dx = \text{Result too large to show}$$

input `Integrate[1/((d + e*x^3)^(3/2)*(a + c*x^6)),x]`

output `Result too large to show`

Rubi [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 438 vs. $2(146) = 292$.

Time = 0.98 (sec), antiderivative size = 438, normalized size of antiderivative = 3.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {1757, 749, 759, 7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a + cx^6)(d + ex^3)^{3/2}} dx \\
 & \quad \downarrow \textcolor{blue}{1757} \\
 & \frac{e^2 \int \frac{1}{(ex^3+d)^{3/2}} dx}{ae^2 + cd^2} + \frac{c \int \frac{d-ex^3}{\sqrt{ex^3+d(cx^6+a)}} dx}{ae^2 + cd^2} \\
 & \quad \downarrow \textcolor{blue}{749} \\
 & \frac{e^2 \left(\frac{\int \frac{1}{\sqrt{ex^3+d}} dx}{3d} + \frac{2x}{3d\sqrt{d+ex^3}} \right)}{ae^2 + cd^2} + \frac{c \int \frac{d-ex^3}{\sqrt{ex^3+d(cx^6+a)}} dx}{ae^2 + cd^2} \\
 & \quad \downarrow \textcolor{blue}{759} \\
 & \frac{c \int \frac{d-ex^3}{\sqrt{ex^3+d(cx^6+a)}} dx}{ae^2 + cd^2} + \\
 & e^2 \left(\frac{2\sqrt{2+\sqrt{3}} \left(\sqrt[3]{d} + \sqrt[3]{e_x} \right) \sqrt{\frac{d^{2/3} - \sqrt[3]{d} \sqrt[3]{e_x + e^{2/3} x^2}}{\left((1+\sqrt{3}) \sqrt[3]{d} + \sqrt[3]{e_x} \right)^2}} \text{EllipticF} \left(\arcsin \left(\frac{\sqrt[3]{e_x + (1-\sqrt{3}) \sqrt[3]{d}}}{\sqrt[3]{e_x + (1+\sqrt{3}) \sqrt[3]{d}}} \right), -7-4\sqrt{3} \right)}{3\sqrt{d+ex^3}} + \frac{2x}{3d\sqrt{d+ex^3}} \right)
 \end{aligned}$$

$$\begin{aligned}
& \frac{c \int \left(\frac{-\sqrt{cd} - \sqrt{-ae}}{2\sqrt{-a}\sqrt{c}(\sqrt{cx^3 + \sqrt{-a}})\sqrt{ex^3 + d}} - \frac{\sqrt{cd} - \sqrt{-ae}}{2\sqrt{-a}\sqrt{c}(\sqrt{-a} - \sqrt{cx^3})\sqrt{ex^3 + d}} \right) dx}{ae^2 + cd^2} + \\
& e^2 \left(\frac{2\sqrt{2+\sqrt{3}} \left(\sqrt[3]{d} + \sqrt[3]{e}x \right) \sqrt{\frac{d^{2/3} - \sqrt[3]{d} \sqrt[3]{e}x^{2/3}x^2}{\left((1+\sqrt{3}) \sqrt[3]{d} + \sqrt[3]{e}x \right)^2}} \text{EllipticF} \left(\arcsin \left(\frac{\sqrt[3]{e}x + (1-\sqrt{3}) \sqrt[3]{d}}{\sqrt[3]{e}x + (1+\sqrt{3}) \sqrt[3]{d}} \right), -7 - 4\sqrt{3} \right)}{3\sqrt[4]{3}d^3\sqrt{e} \sqrt{\frac{\sqrt[3]{d} \left(\sqrt[3]{d} + \sqrt[3]{e}x \right)}{\left((1+\sqrt{3}) \sqrt[3]{d} + \sqrt[3]{e}x \right)^2} \sqrt{d+ex^3}}} + \frac{2x}{3d\sqrt{d+ex^3}} \right) \\
& \downarrow \text{2009} \\
& c \left(\frac{x\sqrt{\frac{ex^3}{d} + 1} \left(\frac{\sqrt{-ae}}{\sqrt{c}} + d \right) \text{AppellF1} \left(\frac{1}{3}, 1, \frac{1}{2}, \frac{4}{3}, -\frac{\sqrt{cx^3}}{\sqrt{-a}}, -\frac{ex^3}{d} \right)}{2a\sqrt{d+ex^3}} + \frac{x\sqrt{\frac{ex^3}{d} + 1} \left(d - \frac{\sqrt{-ae}}{\sqrt{c}} \right) \text{AppellF1} \left(\frac{1}{3}, 1, \frac{1}{2}, \frac{4}{3}, \frac{\sqrt{cx^3}}{\sqrt{-a}}, -\frac{ex^3}{d} \right)}{2a\sqrt{d+ex^3}} \right) + \\
& e^2 \left(\frac{2\sqrt{2+\sqrt{3}} \left(\sqrt[3]{d} + \sqrt[3]{e}x \right) \sqrt{\frac{d^{2/3} - \sqrt[3]{d} \sqrt[3]{e}x^{2/3}x^2}{\left((1+\sqrt{3}) \sqrt[3]{d} + \sqrt[3]{e}x \right)^2}} \text{EllipticF} \left(\arcsin \left(\frac{\sqrt[3]{e}x + (1-\sqrt{3}) \sqrt[3]{d}}{\sqrt[3]{e}x + (1+\sqrt{3}) \sqrt[3]{d}} \right), -7 - 4\sqrt{3} \right)}{3\sqrt[4]{3}d^3\sqrt{e} \sqrt{\frac{\sqrt[3]{d} \left(\sqrt[3]{d} + \sqrt[3]{e}x \right)}{\left((1+\sqrt{3}) \sqrt[3]{d} + \sqrt[3]{e}x \right)^2} \sqrt{d+ex^3}}} + \frac{2x}{3d\sqrt{d+ex^3}} \right) \\
& ae^2 + cd^2
\end{aligned}$$

input `Int[1/((d + e*x^3)^(3/2)*(a + c*x^6)),x]`

output

```
(c*((d + (Sqrt[-a]*e)/Sqrt[c])*x*Sqrt[1 + (e*x^3)/d]*AppellF1[1/3, 1, 1/2, 4/3, -(Sqrt[c]*x^3)/Sqrt[-a]], -((e*x^3)/d))/(2*a*Sqrt[d + e*x^3]) + (d - (Sqrt[-a]*e)/Sqrt[c])*x*Sqrt[1 + (e*x^3)/d]*AppellF1[1/3, 1, 1/2, 4/3, (Sqrt[c]*x^3)/Sqrt[-a], -((e*x^3)/d))/(2*a*Sqrt[d + e*x^3]))/(c*d^2 + a*e^2) + (e^2*((2*x)/(3*d*Sqrt[d + e*x^3]) + (2*Sqrt[2 + Sqrt[3]]*(d^(1/3) + e^(1/3)*x)*Sqrt[(d^(2/3) - d^(1/3)*e^(1/3)*x + e^(2/3)*x^2])/((1 + Sqrt[3])*d^(1/3) + e^(1/3)*x^2)*EllipticF[ArcSin[((1 - Sqrt[3])*d^(1/3) + e^(1/3)*x)/((1 + Sqrt[3])*d^(1/3) + e^(1/3)*x)], -7 - 4*Sqrt[3]])/(3*3^(1/4)*d *e^(1/3)*Sqrt[(d^(1/3)*(d^(1/3) + e^(1/3)*x))/((1 + Sqrt[3])*d^(1/3) + e^(1/3)*x)^2]*Sqrt[d + e*x^3]))/(c*d^2 + a*e^2)
```

Definitions of rubi rules used

rule 749 $\text{Int}[(a_0 + b_0 \cdot x_0^{n_0})^{p_0}, x_{\text{Symbol}}] \rightarrow \text{Simp}[(-x) \cdot ((a + b \cdot x^n)^{p+1}) / (a \cdot n \cdot (p+1)), x] + \text{Simp}[(n \cdot (p+1) + 1) / (a \cdot n \cdot (p+1)) \cdot \text{Int}[(a + b \cdot x^{n_0})^{p+1}, x], x] /; \text{FreeQ}[\{a, b\}, x] \& \text{IGtQ}[n, 0] \& \text{LtQ}[p, -1] \& (\text{IntegerQ}[2*p] \text{ || } \text{Denominator}[p+1/n] < \text{Denominator}[p])$

rule 759 $\text{Int}[1/\sqrt{(a_0 + b_0 \cdot x_0^3)}, x_{\text{Symbol}}] \rightarrow \text{With}[\{r = \text{Numer}[\text{Rt}[b/a, 3]], s = \text{Denom}[\text{Rt}[b/a, 3]]\}, \text{Simp}[2*\sqrt[2]{2 + \sqrt[3]{3}} \cdot (s + r \cdot x) \cdot (\sqrt{s^2 - r^2} \cdot x + r^2 \cdot x^2) / ((1 + \sqrt[3]{3}) \cdot s + r \cdot x)^2] / (3^{(1/4)} \cdot r \cdot \sqrt{a + b \cdot x^3} \cdot \sqrt{s^2 - r^2 \cdot x^2}) \cdot \text{EllipticF}[\text{ArcSin}[((1 - \sqrt[3]{3}) \cdot s + r \cdot x) / ((1 + \sqrt[3]{3}) \cdot s + r \cdot x)], -7 - 4 \cdot \sqrt[3]{3}], x] /; \text{FreeQ}[\{a, b\}, x] \& \text{PosQ}[a]$

rule 1757 $\text{Int}[(d_0 + e_0 \cdot x_0^{n_0})^{q_0} / ((a_0 + c_0 \cdot x_0^{n_2}))^{q_1}, x_{\text{Symbol}}] \rightarrow \text{Simp}[e^{2/(c \cdot d^2 + a \cdot e^2)} \cdot \text{Int}[(d + e \cdot x^n)^q, x], x] + \text{Simp}[c / (c \cdot d^2 + a \cdot e^2) \cdot \text{Int}[(d + e \cdot x^n)^{q+1} \cdot ((d - e \cdot x^n) / (a + c \cdot x^{(2 \cdot n)})), x], x] /; \text{FreeQ}[\{a, c, d, e, n\}, x] \& \text{EqQ}[n_2, 2 \cdot n] \& \text{NeQ}[c \cdot d^2 + a \cdot e^2, 0] \& \text{!IntegerQ}[q] \& \text{LtQ}[q, -1]$

rule 2009 $\text{Int}[u_0, x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 7276 $\text{Int}[(u_0) / ((a_0 + b_0 \cdot x_0^{n_0})^{q_0}), x_{\text{Symbol}}] \rightarrow \text{With}[\{v = \text{RationalFunctionExpand}[u / (a + b \cdot x^n), x]\}, \text{Int}[v, x] /; \text{SumQ}[v]] /; \text{FreeQ}[\{a, b\}, x] \& \text{IGtQ}[n, 0]$

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 6.

Time = 0.36 (sec) , antiderivative size = 937, normalized size of antiderivative = 6.42

method	result	size
default	Expression too large to display	937
elliptic	Expression too large to display	937

input `int(1/(e*x^3+d)^(3/2)/(c*x^6+a),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & 2/3*e^2*x/d/(a*e^2+c*d^2)/((x^3+d/e)*e)^(1/2)-2/9*I/d*e/(a*e^2+c*d^2)*3^(1/2)*(-d*e^2)^(1/3)*(I*(x+1/2/e*(-d*e^2)^(1/3)-1/2*I*3^(1/2)/e*(-d*e^2)^(1/3))*3^(1/2)*e/(-d*e^2)^(1/3))^(1/2)*((x-1/e*(-d*e^2)^(1/3))/(-3/2/e*(-d*e^2)^(1/3)+1/2*I*3^(1/2)/e*(-d*e^2)^(1/3)))^(1/2)*(-I*(x+1/2/e*(-d*e^2)^(1/3)+1/2*I*3^(1/2)/e*(-d*e^2)^(1/3))*3^(1/2)*e/(-d*e^2)^(1/3))^(1/2)/(e*x^3+d)^(1/2)*\text{EllipticF}(1/3*3^(1/2)*(I*(x+1/2/e*(-d*e^2)^(1/3)-1/2*I*3^(1/2)/e*(-d*e^2)^(1/3))*3^(1/2)*e/(-d*e^2)^(1/3)),(I*3^(1/2)/e*(-d*e^2)^(1/3)/(-3/2/e*(-d*e^2)^(1/3)+1/2*I*3^(1/2)/e*(-d*e^2)^(1/3)))^(1/2))-1/6*I*c/e^2*2^(1/2)*\text{sum}(_{\alpha}^3 e^{-d}/(a*e^2+c*d^2)^2/_{\alpha}^5 (-d*e^2)^(1/3)*(1/2*I*e*(2*x+1/e*(-I*3^(1/2)*(-d*e^2)^(1/3)+(-d*e^2)^(1/3)))/(-d*e^2)^(1/3))^(1/2)*(e*(x-1/e*(-d*e^2)^(1/3))/(-3*(-d*e^2)^(1/3)+I*3^(1/2)*(-d*e^2)^(1/3)))^(1/2)*(-1/2*I*e*(2*x+1/e*(I*3^(1/2)*(-d*e^2)^(1/3)+(-d*e^2)^(1/3)))/(-d*e^2)^(1/3))^(1/2)/(e*x^3+d)^(1/2)*(2*e^2*(-_alpha^5 e + _alpha^2 d) - I*(-d*e^2)^(1/3)*_alpha^4 3^(1/2)*e^2 + I*_alpha^3 3^(1/2)*(-d*e^2)^(2/3)*e + (-d*e^2)^(1/3)*_alpha^4 e^2 + I*(-d*e^2)^(1/3)*d*_alpha^3 3^(1/2)*e + _alpha^3 (-d*e^2)^(2/3)*e - I*3^(1/2)*(-d*e^2)^(2/3)*d - (-d*e^2)^(1/3)*d*_alpha^2 e - (-d*e^2)^(2/3)*d)*\text{EllipticPi}(1/3*3^(1/2)*(I*(x+1/2/e*(-d*e^2)^(1/3)-1/2*I*3^(1/2)/e*(-d*e^2)^(1/3))*3^(1/2)*e/(-d*e^2)^(1/3))^(1/2),-1/2*c/e*(-2*I*3^(1/2)*(-d*e^2)^(1/3)*_alpha^5 e^2 + I*3^(1/2)*(-d*e^2)^(2/3)*_alpha^4 e - I*3^(1/2)*_alpha^3 d + 2*I*3^(1/2)*(-d*e^2)^(1/3)*_alpha^2 d + 3*(-d*e^2)^(2/3)*_alpha^2 d + 2*I*3^(1/2)*(-d*e^2)^(1/3)*_alpha^1 d + 3*(-d*e^2)^(1/3)*_alpha^0 d))/\dots \end{aligned}$$

Fricas [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex^3)^{3/2} (a + cx^6)} dx = \text{Timed out}$$

input `integrate(1/(e*x^3+d)^(3/2)/(c*x^6+a),x, algorithm="fricas")`

output Timed out

Sympy [F]

$$\int \frac{1}{(d + ex^3)^{3/2} (a + cx^6)} dx = \int \frac{1}{(a + cx^6) (d + ex^3)^{\frac{3}{2}}} dx$$

input `integrate(1/(e*x**3+d)**(3/2)/(c*x**6+a),x)`

output `Integral(1/((a + c*x**6)*(d + e*x**3)**(3/2)), x)`

Maxima [F]

$$\int \frac{1}{(d + ex^3)^{3/2} (a + cx^6)} dx = \int \frac{1}{(cx^6 + a)(ex^3 + d)^{\frac{3}{2}}} dx$$

input `integrate(1/(e*x^3+d)^(3/2)/(c*x^6+a),x, algorithm="maxima")`

output `integrate(1/((c*x^6 + a)*(e*x^3 + d)^(3/2)), x)`

Giac [F]

$$\int \frac{1}{(d + ex^3)^{3/2} (a + cx^6)} dx = \int \frac{1}{(cx^6 + a)(ex^3 + d)^{\frac{3}{2}}} dx$$

input `integrate(1/(e*x^3+d)^(3/2)/(c*x^6+a),x, algorithm="giac")`

output `integrate(1/((c*x^6 + a)*(e*x^3 + d)^(3/2)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex^3)^{3/2} (a + cx^6)} dx = \int \frac{1}{(cx^6 + a) (e x^3 + d)^{3/2}} dx$$

input `int(1/((a + c*x^6)*(d + e*x^3)^(3/2)),x)`

output `int(1/((a + c*x^6)*(d + e*x^3)^(3/2)), x)`

Reduce [F]

$$\int \frac{1}{(d + ex^3)^{3/2} (a + cx^6)} dx = \int \frac{\sqrt{e x^3 + d}}{c e^2 x^{12} + 2 c d e x^9 + a e^2 x^6 + c d^2 x^6 + 2 a d e x^3 + a d^2} dx$$

input `int(1/(e*x^3+d)^(3/2)/(c*x^6+a),x)`

output `int(sqrt(d + e*x**3)/(a*d**2 + 2*a*d*e*x**3 + a*e**2*x**6 + c*d**2*x**6 + 2*c*d*e*x**9 + c*e**2*x**12),x)`

3.7 $\int \frac{\sqrt{1-x^6}}{\sqrt{1-x^3}} dx$

Optimal result	101
Mathematica [F]	102
Rubi [A] (verified)	102
Maple [B] (verified)	103
Fricas [A] (verification not implemented)	104
Sympy [F]	105
Maxima [F]	105
Giac [F]	105
Mupad [F(-1)]	106
Reduce [F]	106

Optimal result

Integrand size = 23, antiderivative size = 120

$$\begin{aligned} & \int \frac{\sqrt{1-x^6}}{\sqrt{1-x^3}} dx \\ &= \frac{2}{5}x\sqrt{1+x^3} \\ &+ \frac{2 \cdot 3^{3/4}\sqrt{2+\sqrt{3}}(1+x)\sqrt{\frac{1-x+x^2}{(1+\sqrt{3}+x)^2}} \operatorname{EllipticF}\left(\arcsin\left(\frac{1-\sqrt{3}+x}{1+\sqrt{3}+x}\right), -7-4\sqrt{3}\right)}{5\sqrt{\frac{1+x}{(1+\sqrt{3}+x)^2}}\sqrt{1+x^3}} \end{aligned}$$

output
$$\frac{2/5*x*(x^3+1)^(1/2)+2/5*3^(3/4)*(1/2*6^(1/2)+1/2*2^(1/2))*(1+x)*((x^2-x+1)/(1+x+3^(1/2))^2)^(1/2)*\operatorname{EllipticF}((1+x-3^(1/2))/(1+x+3^(1/2)), I*3^(1/2)+2*I)/((1+x)/(1+x+3^(1/2))^2)^(1/2)/(x^3+1)^(1/2)}$$

Mathematica [F]

$$\int \frac{\sqrt{1-x^6}}{\sqrt{1-x^3}} dx = \int \frac{\sqrt{1-x^6}}{\sqrt{1-x^3}} dx$$

input `Integrate[Sqrt[1 - x^6]/Sqrt[1 - x^3], x]`

output `Integrate[Sqrt[1 - x^6]/Sqrt[1 - x^3], x]`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {1386, 748, 759}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{1-x^6}}{\sqrt{1-x^3}} dx \\
 & \quad \downarrow 1386 \\
 & \int \sqrt{x^3+1} dx \\
 & \quad \downarrow 748 \\
 & \frac{3}{5} \int \frac{1}{\sqrt{x^3+1}} dx + \frac{2}{5} \sqrt{x^3+1} x \\
 & \quad \downarrow 759 \\
 & \frac{2 \cdot 3^{3/4} \sqrt{2+\sqrt{3}} (x+1) \sqrt{\frac{x^2-x+1}{(x+\sqrt{3}+1)^2}} \operatorname{EllipticF}\left(\arcsin\left(\frac{x-\sqrt{3}+1}{x+\sqrt{3}+1}\right), -7-4\sqrt{3}\right)}{5 \sqrt{\frac{x+1}{(x+\sqrt{3}+1)^2}} \sqrt{x^3+1}} + \frac{2}{5} \sqrt{x^3+1} x
 \end{aligned}$$

input `Int[Sqrt[1 - x^6]/Sqrt[1 - x^3], x]`

output

$$(2*x*sqrt[1 + x^3])/5 + (2*3^(3/4)*sqrt[2 + sqrt[3]]*(1 + x)*sqrt[(1 - x + x^2)/(1 + sqrt[3] + x)^2]*EllipticF[ArcSin[(1 - sqrt[3] + x)/(1 + sqrt[3] + x)], -7 - 4*sqrt[3]]]/(5*sqrt[(1 + x)/(1 + sqrt[3] + x)^2]*sqrt[1 + x^3])$$

Definitions of rubi rules used

rule 748

$$\text{Int}[(a_0 + b_0 \cdot (x_0)^{n_0})^{p_0}, x_{\text{Symbol}}] \rightarrow \text{Simp}[x \cdot ((a + b \cdot x^n)^{p/(n*p + 1)}), x] + \text{Simp}[a \cdot n \cdot (p/(n*p + 1)) \cdot \text{Int}[(a + b \cdot x^n)^{p - 1}, x], x] /; \text{FreeQ}[\{a, b\}, x] \&& \text{IGtQ}[n, 0] \&& \text{GtQ}[p, 0] \&& (\text{IntegerQ}[2*p] \text{ || } \text{LtQ}[\text{Denominator}[p + 1/n], \text{Denominator}[p]])$$

rule 759

$$\text{Int}[1/\sqrt{(a_0 + b_0 \cdot (x_0)^3}], x_{\text{Symbol}}] \rightarrow \text{With}[\{r = \text{Numer}[\text{Rt}[b/a, 3]], s = \text{Denom}[\text{Rt}[b/a, 3]]\}, \text{Simp}[2*\sqrt[2 + sqrt[3]]*(s + r*x)*(sqrt[(s^2 - r*s*x + r^2*x^2)/((1 + sqrt[3])*s + r*x)^2]/(3^(1/4)*r*sqrt[a + b*x^3]*sqrt[s*((s + r*x)/((1 + sqrt[3])*s + r*x)^2)]))*\text{EllipticF}[\text{ArcSin}[(1 - sqrt[3])*s + r*x]/((1 + sqrt[3])*s + r*x)], -7 - 4*sqrt[3]], x] /; \text{FreeQ}[\{a, b\}, x] \& \text{PosQ}[a]$$

rule 1386

$$\text{Int}[(u_0 \cdot ((a_0 + c_0 \cdot (x_0)^{n_0})^{p_0} \cdot (d_0 + e_0 \cdot (x_0)^{n_0})^{q_0}))^{p_0}, x_{\text{Symbol}}] \rightarrow \text{Simp}[(-e^2/c)^q \cdot \text{Int}[u \cdot (d - e \cdot x^n)^p, x], x] /; \text{FreeQ}[\{a, c, d, e, n, p, q\}, x] \&& \text{EqQ}[n_0, 2*n] \&& \text{EqQ}[c*d^2 + a*e^2, 0] \&& \text{EqQ}[p + q, 0] \&& \text{GtQ}[d, 0] \&& \text{LtQ}[c, 0] \&& \text{GtQ}[e^2, 0]$$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 220 vs. $2(97) = 194$.

Time = 0.27 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.84

method	result
risch	$-\frac{2x\sqrt{x^3+1}\sqrt{\frac{(-x^6+1)(-x^3+1)}{(x^3-1)^2}(x^3-1)}}{5\sqrt{-x^6+1}\sqrt{-x^3+1}} - \frac{6\left(\frac{3}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{\frac{x+1}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x-\frac{1}{2}-\frac{i\sqrt{3}}{2}}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x-\frac{1}{2}+\frac{i\sqrt{3}}{2}}{-\frac{3}{2}+\frac{i\sqrt{3}}{2}}}\text{EllipticF}\left(\sqrt{\frac{x+1}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}},\sqrt{\frac{-\frac{3}{2}+\frac{i\sqrt{3}}{2}}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\right)}{5\sqrt{x^3+1}\sqrt{-x^6+1}\sqrt{-x^3+1}}$
default	$-\frac{\sqrt{-x^6+1}\left(3i\sqrt{3}\sqrt{-\frac{2(x+1)}{i\sqrt{3}-3}}\sqrt{\frac{i\sqrt{3}-2x+1}{i\sqrt{3}+3}}\sqrt{\frac{i\sqrt{3}+2x-1}{i\sqrt{3}-3}}\text{EllipticF}\left(\sqrt{-\frac{2(x+1)}{i\sqrt{3}-3}},\sqrt{-\frac{i\sqrt{3}-3}{i\sqrt{3}+3}}\right)-9\sqrt{-\frac{2(x+1)}{i\sqrt{3}-3}}\sqrt{\frac{i\sqrt{3}-2x+1}{i\sqrt{3}+3}}\sqrt{\frac{i\sqrt{3}+2x-1}{i\sqrt{3}-3}}}\right)}{5\sqrt{-x^3+1}(x^3+1)}$

input `int((-x^6+1)^(1/2)/(-x^3+1)^(1/2),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -2/5*x*(x^3+1)^(1/2)*((-x^6+1)*(-x^3+1)/(x^3-1)^2)^(1/2)/(-x^6+1)^(1/2)/(- \\ & x^3+1)^(1/2)*(x^3-1)-6/5*(3/2-1/2*I*3^(1/2))*(x+1)/(3/2-1/2*I*3^(1/2)))^(1/2)*((x-1/2-1/2*I*3^(1/2))/(-3/2-1/2*I*3^(1/2)))^(1/2)*((x-1/2+1/2*I*3^(1/2))/(-3/2+1/2*I*3^(1/2)))^(1/2)/(x^3+1)^(1/2)*\text{EllipticF}(((x+1)/(3/2-1/2*I*3^(1/2)))^(1/2),((-3/2+1/2*I*3^(1/2))/(-3/2-1/2*I*3^(1/2)))^(1/2))*((-x^6+1)*(-x^3+1)^(1/2)*(x^3-1)) \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.34

$$\int \frac{\sqrt{1-x^6}}{\sqrt{1-x^3}} dx = -\frac{2(\sqrt{-x^6+1}\sqrt{-x^3+1}x + 3(x^3-1)\text{weierstrassPI}(0, -4, x))}{5(x^3-1)}$$

input `integrate((-x^6+1)^(1/2)/(-x^3+1)^(1/2),x, algorithm="fricas")`

output
$$-2/5*(\sqrt{-x^6+1}*\sqrt{-x^3+1}*x + 3*(x^3-1)*\text{weierstrassPI}(0, -4, x))/(x^3-1)$$

Sympy [F]

$$\int \frac{\sqrt{1-x^6}}{\sqrt{1-x^3}} dx = \int \frac{\sqrt{-(x-1)(x+1)(x^2-x+1)(x^2+x+1)}}{\sqrt{-(x-1)(x^2+x+1)}} dx$$

input `integrate((-x**6+1)**(1/2)/(-x**3+1)**(1/2),x)`

output `Integral(sqrt(-(x - 1)*(x + 1)*(x**2 - x + 1)*(x**2 + x + 1))/sqrt(-(x - 1)*(x**2 + x + 1)), x)`

Maxima [F]

$$\int \frac{\sqrt{1-x^6}}{\sqrt{1-x^3}} dx = \int \frac{\sqrt{-x^6+1}}{\sqrt{-x^3+1}} dx$$

input `integrate((-x^6+1)^(1/2)/(-x^3+1)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(-x^6 + 1)/sqrt(-x^3 + 1), x)`

Giac [F]

$$\int \frac{\sqrt{1-x^6}}{\sqrt{1-x^3}} dx = \int \frac{\sqrt{-x^6+1}}{\sqrt{-x^3+1}} dx$$

input `integrate((-x^6+1)^(1/2)/(-x^3+1)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(-x^6 + 1)/sqrt(-x^3 + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{1-x^6}}{\sqrt{1-x^3}} dx = \int \frac{\sqrt{1-x^6}}{\sqrt{1-x^3}} dx$$

input `int((1 - x^6)^(1/2)/(1 - x^3)^(1/2),x)`

output `int((1 - x^6)^(1/2)/(1 - x^3)^(1/2), x)`

Reduce [F]

$$\int \frac{\sqrt{1-x^6}}{\sqrt{1-x^3}} dx = -\left(\int \frac{\sqrt{-x^3+1} \sqrt{-x^6+1}}{x^3-1} dx \right)$$

input `int((-x^6+1)^(1/2)/(-x^3+1)^(1/2),x)`

output `- int(sqrt(-x**3 + 1)*sqrt(-x**6 + 1))/(x**3 - 1),x)`

3.8 $\int (d + ex^3) (a - cx^6)^p \, dx$

Optimal result	107
Mathematica [A] (verified)	107
Rubi [A] (verified)	108
Maple [F]	109
Fricas [F]	109
Sympy [C] (verification not implemented)	110
Maxima [F]	110
Giac [F]	111
Mupad [F(-1)]	111
Reduce [F]	111

Optimal result

Integrand size = 18, antiderivative size = 98

$$\int (d + ex^3) (a - cx^6)^p \, dx = dx(a - cx^6)^p \left(1 - \frac{cx^6}{a}\right)^{-p} \text{Hypergeometric2F1} \left(\frac{1}{6}, -p, \frac{7}{6}, \frac{cx^6}{a}\right) + \frac{1}{4} ex^4 (a - cx^6)^p \left(1 - \frac{cx^6}{a}\right)^{-p} \text{Hypergeometric2F1} \left(\frac{2}{3}, -p, \frac{5}{3}, \frac{cx^6}{a}\right)$$

output $d*x*(-c*x^6+a)^p*\text{hypergeom}([1/6, -p], [7/6], c*x^6/a)/((1-c*x^6/a)^p)+1/4*e*x^4*(-c*x^6+a)^p*\text{hypergeom}([2/3, -p], [5/3], c*x^6/a)/((1-c*x^6/a)^p)$

Mathematica [A] (verified)

Time = 0.57 (sec), antiderivative size = 75, normalized size of antiderivative = 0.77

$$\int (d + ex^3) (a - cx^6)^p \, dx = \frac{1}{4} x (a - cx^6)^p \left(1 - \frac{cx^6}{a}\right)^{-p} \left(4d \text{Hypergeometric2F1} \left(\frac{1}{6}, -p, \frac{7}{6}, \frac{cx^6}{a}\right) + ex^3 \text{Hypergeometric2F1} \left(\frac{2}{3}, -p, \frac{5}{3}, \frac{cx^6}{a}\right)\right)$$

input $\text{Integrate}[(d + e*x^3)*(a - c*x^6)^p, x]$

output $(x*(a - c*x^6)^p*(4*d*\text{Hypergeometric2F1}[1/6, -p, 7/6, (c*x^6)/a] + e*x^3*\text{Hypergeometric2F1}[2/3, -p, 5/3, (c*x^6)/a]))/(4*(1 - (c*x^6)/a)^p)$

Rubi [A] (verified)

Time = 0.25 (sec), antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.111, Rules used = {1763, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (d + ex^3) (a - cx^6)^p dx \\ & \quad \downarrow 1763 \\ & \int (d(a - cx^6)^p + ex^3(a - cx^6)^p) dx \\ & \quad \downarrow 2009 \\ & dx(a - cx^6)^p \left(1 - \frac{cx^6}{a}\right)^{-p} \text{Hypergeometric2F1}\left(\frac{1}{6}, -p, \frac{7}{6}, \frac{cx^6}{a}\right) + \\ & \quad \frac{1}{4}ex^4(a - cx^6)^p \left(1 - \frac{cx^6}{a}\right)^{-p} \text{Hypergeometric2F1}\left(\frac{2}{3}, -p, \frac{5}{3}, \frac{cx^6}{a}\right) \end{aligned}$$

input $\text{Int}[(d + e*x^3)*(a - c*x^6)^p, x]$

output $(d*x*(a - c*x^6)^p*\text{Hypergeometric2F1}[1/6, -p, 7/6, (c*x^6)/a])/(1 - (c*x^6)/a)^p + (e*x^4*(a - c*x^6)^p*\text{Hypergeometric2F1}[2/3, -p, 5/3, (c*x^6)/a])/ (4*(1 - (c*x^6)/a)^p)$

Definitions of rubi rules used

rule 1763 $\text{Int}[(d_0 + e_0 x^{n_0}) (a_0 + c_0 x^{n_2})^p, x] \rightarrow \text{Int}[\text{ExpandIntegrand}[(d + e x^n) (a + c x^{2n})^p, x], x] /; \text{FreeQ}[\{a, c, d, e, n\}, x] \& \text{EqQ}[n_2, 2n]$

rule 2009 $\text{Int}[u, x] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

Maple [F]

$$\int (e x^3 + d) (-c x^6 + a)^p dx$$

input `int((e*x^3+d)*(-c*x^6+a)^p,x)`

output `int((e*x^3+d)*(-c*x^6+a)^p,x)`

Fricas [F]

$$\int (d + e x^3) (a - c x^6)^p dx = \int (e x^3 + d) (-c x^6 + a)^p dx$$

input `integrate((e*x^3+d)*(-c*x^6+a)^p,x, algorithm="fricas")`

output `integral((e*x^3 + d)*(-c*x^6 + a)^p, x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 39.40 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.80

$$\int (d + ex^3) (a - cx^6)^p dx = \frac{a^p dx \Gamma\left(\frac{1}{6}\right) {}_2F_1\left(\begin{matrix} \frac{1}{6}, -p \\ \frac{7}{6} \end{matrix} \middle| \frac{cx^6 e^{2i\pi}}{a}\right)}{6\Gamma\left(\frac{7}{6}\right)} + \frac{a^p ex^4 \Gamma\left(\frac{2}{3}\right) {}_2F_1\left(\begin{matrix} \frac{2}{3}, -p \\ \frac{5}{3} \end{matrix} \middle| \frac{cx^6 e^{2i\pi}}{a}\right)}{6\Gamma\left(\frac{5}{3}\right)}$$

input `integrate((e*x**3+d)*(-c*x**6+a)**p,x)`

output `a**p*d*x*gamma(1/6)*hyper((1/6, -p), (7/6,), c*x**6*exp_polar(2*I*pi)/a)/(6*gamma(7/6)) + a**p*e*x**4*gamma(2/3)*hyper((2/3, -p), (5/3,), c*x**6*exp_polar(2*I*pi)/a)/(6*gamma(5/3))`

Maxima [F]

$$\int (d + ex^3) (a - cx^6)^p dx = \int (ex^3 + d) (-cx^6 + a)^p dx$$

input `integrate((e*x^3+d)*(-c*x^6+a)^p,x, algorithm="maxima")`

output `integrate((e*x^3 + d)*(-c*x^6 + a)^p, x)`

Giac [F]

$$\int (d + ex^3) (a - cx^6)^p dx = \int (ex^3 + d)(-cx^6 + a)^p dx$$

input `integrate((e*x^3+d)*(-c*x^6+a)^p,x, algorithm="giac")`

output `integrate((e*x^3 + d)*(-c*x^6 + a)^p, x)`

Mupad [F(-1)]

Timed out.

$$\int (d + ex^3) (a - cx^6)^p dx = \int (a - cx^6)^p (e x^3 + d) dx$$

input `int((a - c*x^6)^p*(d + e*x^3),x)`

output `int((a - c*x^6)^p*(d + e*x^3), x)`

Reduce [F]

$$\begin{aligned} & \int (d + ex^3) (a - cx^6)^p dx \\ &= \frac{6(-cx^6 + a)^p dpx + 4(-cx^6 + a)^p dx + 6(-cx^6 + a)^p ep x^4 + (-cx^6 + a)^p e x^4 + 648 \left(\int \frac{(-}{-18cp^2x^6 - 15cp x^6} \right)}{ } \end{aligned}$$

input `int((e*x^3+d)*(-c*x^6+a)^p,x)`

output

```
(6*(a - c*x**6)**p*d*p*x + 4*(a - c*x**6)**p*d*x + 6*(a - c*x**6)**p*e*p*x
**4 + (a - c*x**6)**p*e*x**4 + 648*int((a - c*x**6)**p/(18*a*p**2 + 15*a*p
+ 2*a - 18*c*p**2*x**6 - 15*c*p*x**6 - 2*c*x**6),x)*a*d*p**4 + 972*int((a
- c*x**6)**p/(18*a*p**2 + 15*a*p + 2*a - 18*c*p**2*x**6 - 15*c*p*x**6 - 2
*c*x**6),x)*a*d*p**3 + 432*int((a - c*x**6)**p/(18*a*p**2 + 15*a*p + 2*a -
18*c*p**2*x**6 - 15*c*p*x**6 - 2*c*x**6),x)*a*d*p**2 + 48*int((a - c*x**6
)**p/(18*a*p**2 + 15*a*p + 2*a - 18*c*p**2*x**6 - 15*c*p*x**6 - 2*c*x**6),
x)*a*d*p + 648*int(((a - c*x**6)**p*x**3)/(18*a*p**2 + 15*a*p + 2*a - 18*c
*p**2*x**6 - 15*c*p*x**6 - 2*c*x**6),x)*a*e*p**4 + 648*int(((a - c*x**6)**p*x**3)/(18*a*p**2 + 15*a*p + 2*a - 18*c*p**2*x**6 - 15*c*p*x**6 - 2*c*x**6),
x)*a*e*p**3 + 162*int(((a - c*x**6)**p*x**3)/(18*a*p**2 + 15*a*p + 2*a -
18*c*p**2*x**6 - 15*c*p*x**6 - 2*c*x**6),x)*a*e*p**2 + 12*int(((a - c*x*
6)**p*x**3)/(18*a*p**2 + 15*a*p + 2*a - 18*c*p**2*x**6 - 15*c*p*x**6 - 2*
c*x**6),x)*a*e*p)/(2*(18*p**2 + 15*p + 2))
```

3.9 $\int (d + ex^3) (a + cx^6)^p \, dx$

Optimal result	113
Mathematica [A] (verified)	113
Rubi [A] (verified)	114
Maple [F]	115
Fricas [F]	115
Sympy [C] (verification not implemented)	116
Maxima [F]	116
Giac [F]	117
Mupad [F(-1)]	117
Reduce [F]	117

Optimal result

Integrand size = 17, antiderivative size = 96

$$\int (d + ex^3) (a + cx^6)^p \, dx = dx(a + cx^6)^p \left(1 + \frac{cx^6}{a}\right)^{-p} \text{Hypergeometric2F1} \left(\frac{1}{6}, -p, \frac{7}{6}, -\frac{cx^6}{a}\right) + \frac{1}{4} ex^4 (a + cx^6)^p \left(1 + \frac{cx^6}{a}\right)^{-p} \text{Hypergeometric2F1} \left(\frac{2}{3}, -p, \frac{5}{3}, -\frac{cx^6}{a}\right)$$

output $d*x*(c*x^6+a)^p*\text{hypergeom}([1/6, -p], [7/6], -c*x^6/a)/((1+c*x^6/a)^p)+1/4*e*x^4*(c*x^6+a)^p*\text{hypergeom}([2/3, -p], [5/3], -c*x^6/a)/((1+c*x^6/a)^p)$

Mathematica [A] (verified)

Time = 0.57 (sec), antiderivative size = 75, normalized size of antiderivative = 0.78

$$\int (d + ex^3) (a + cx^6)^p \, dx = \frac{1}{4} x (a + cx^6)^p \left(1 + \frac{cx^6}{a}\right)^{-p} \left(4d \text{Hypergeometric2F1} \left(\frac{1}{6}, -p, \frac{7}{6}, -\frac{cx^6}{a}\right) + ex^3 \text{Hypergeometric2F1} \left(\frac{2}{3}, -p, \frac{5}{3}, -\frac{cx^6}{a}\right)\right)$$

input $\text{Integrate}[(d + e*x^3)*(a + c*x^6)^p, x]$

output $(x*(a + c*x^6)^p*(4*d*\text{Hypergeometric2F1}[1/6, -p, 7/6, -((c*x^6)/a)] + e*x^3*\text{Hypergeometric2F1}[2/3, -p, 5/3, -((c*x^6)/a)]))/(4*(1 + (c*x^6)/a)^p)$

Rubi [A] (verified)

Time = 0.24 (sec), antiderivative size = 96, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {1763, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (d + ex^3) (a + cx^6)^p dx \\ & \quad \downarrow 1763 \\ & \int (d(a + cx^6)^p + ex^3(a + cx^6)^p) dx \\ & \quad \downarrow 2009 \\ & dx(a + cx^6)^p \left(\frac{cx^6}{a} + 1\right)^{-p} \text{Hypergeometric2F1}\left(\frac{1}{6}, -p, \frac{7}{6}, -\frac{cx^6}{a}\right) + \\ & \frac{1}{4}ex^4(a + cx^6)^p \left(\frac{cx^6}{a} + 1\right)^{-p} \text{Hypergeometric2F1}\left(\frac{2}{3}, -p, \frac{5}{3}, -\frac{cx^6}{a}\right) \end{aligned}$$

input $\text{Int}[(d + e*x^3)*(a + c*x^6)^p, x]$

output $(d*x*(a + c*x^6)^p*\text{Hypergeometric2F1}[1/6, -p, 7/6, -((c*x^6)/a)])/(1 + (c*x^6)/a)^p + (e*x^4*(a + c*x^6)^p*\text{Hypergeometric2F1}[2/3, -p, 5/3, -((c*x^6)/a)])/(4*(1 + (c*x^6)/a)^p)$

Definitions of rubi rules used

rule 1763 $\text{Int}[(d_0 + e_0 x^{n_0}) (a_0 + c_0 x^{n_2})^p, x] \rightarrow \text{Int}[\text{ExpandIntegrand}[(d + e x^n) (a + c x^{2n})^p, x], x] /; \text{FreeQ}[\{a, c, d, e, n\}, x] \& \text{EqQ}[n_2, 2n]$

rule 2009 $\text{Int}[u, x] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

Maple [F]

$$\int (e x^3 + d) (c x^6 + a)^p dx$$

input `int((e*x^3+d)*(c*x^6+a)^p,x)`

output `int((e*x^3+d)*(c*x^6+a)^p,x)`

Fricas [F]

$$\int (d + e x^3) (a + c x^6)^p dx = \int (e x^3 + d) (c x^6 + a)^p dx$$

input `integrate((e*x^3+d)*(c*x^6+a)^p,x, algorithm="fricas")`

output `integral((e*x^3 + d)*(c*x^6 + a)^p, x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 39.30 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.78

$$\int (d + ex^3) (a + cx^6)^p dx = \frac{a^p dx \Gamma\left(\frac{1}{6}\right) {}_2F_1\left(\begin{matrix} \frac{1}{6}, -p \\ \frac{7}{6} \end{matrix} \middle| \frac{cx^6 e^{i\pi}}{a}\right)}{6\Gamma\left(\frac{7}{6}\right)} + \frac{a^p ex^4 \Gamma\left(\frac{2}{3}\right) {}_2F_1\left(\begin{matrix} \frac{2}{3}, -p \\ \frac{5}{3} \end{matrix} \middle| \frac{cx^6 e^{i\pi}}{a}\right)}{6\Gamma\left(\frac{5}{3}\right)}$$

input `integrate((e*x**3+d)*(c*x**6+a)**p,x)`

output `a**p*d*x*gamma(1/6)*hyper((1/6, -p), (7/6,), c*x**6*exp_polar(I*pi)/a)/(6*gamma(7/6)) + a**p*e*x**4*gamma(2/3)*hyper((2/3, -p), (5/3,), c*x**6*exp_polar(I*pi)/a)/(6*gamma(5/3))`

Maxima [F]

$$\int (d + ex^3) (a + cx^6)^p dx = \int (ex^3 + d)(cx^6 + a)^p dx$$

input `integrate((e*x^3+d)*(c*x^6+a)^p,x, algorithm="maxima")`

output `integrate((e*x^3 + d)*(c*x^6 + a)^p, x)`

Giac [F]

$$\int (d + ex^3) (a + cx^6)^p \, dx = \int (ex^3 + d)(cx^6 + a)^p \, dx$$

input `integrate((e*x^3+d)*(c*x^6+a)^p,x, algorithm="giac")`

output `integrate((e*x^3 + d)*(c*x^6 + a)^p, x)`

Mupad [F(-1)]

Timed out.

$$\int (d + ex^3) (a + cx^6)^p \, dx = \int (cx^6 + a)^p (e x^3 + d) \, dx$$

input `int((a + c*x^6)^p*(d + e*x^3),x)`

output `int((a + c*x^6)^p*(d + e*x^3), x)`

Reduce [F]

$$\begin{aligned} & \int (d + ex^3) (a + cx^6)^p \, dx \\ &= \frac{6(cx^6 + a)^p \, dpx + 4(cx^6 + a)^p \, dx + 6(cx^6 + a)^p \, ep \, x^4 + (cx^6 + a)^p \, e \, x^4 + 648 \left(\int \frac{(cx^6 + a)^p}{18cp^2x^6 + 15cp \, x^6 + 2cx^6 + 18a} \right)}{18cp^2x^6 + 15cp \, x^6 + 2cx^6 + 18a} \end{aligned}$$

input `int((e*x^3+d)*(c*x^6+a)^p,x)`

output

```
(6*(a + c*x**6)**p*d*p*x + 4*(a + c*x**6)**p*d*x + 6*(a + c*x**6)**p*e*p*x
**4 + (a + c*x**6)**p*e*x**4 + 648*int((a + c*x**6)**p/(18*a*p**2 + 15*a*p
+ 2*a + 18*c*p**2*x**6 + 15*c*p*x**6 + 2*c*x**6),x)*a*d*p**4 + 972*int((a
+ c*x**6)**p/(18*a*p**2 + 15*a*p + 2*a + 18*c*p**2*x**6 + 15*c*p*x**6 + 2
*c*x**6),x)*a*d*p**3 + 432*int((a + c*x**6)**p/(18*a*p**2 + 15*a*p + 2*a +
18*c*p**2*x**6 + 15*c*p*x**6 + 2*c*x**6),x)*a*d*p**2 + 48*int((a + c*x**6
)**p/(18*a*p**2 + 15*a*p + 2*a + 18*c*p**2*x**6 + 15*c*p*x**6 + 2*c*x**6),
x)*a*d*p + 648*int(((a + c*x**6)**p*x**3)/(18*a*p**2 + 15*a*p + 2*a + 18*c
*p**2*x**6 + 15*c*p*x**6 + 2*c*x**6),x)*a*e*p**4 + 648*int(((a + c*x**6)**p*x**3)/(18*a*p**2 + 15*a*p + 2*a + 18*c*p**2*x**6 + 15*c*p*x**6 + 2*c*x**6),
x)*a*e*p**3 + 162*int(((a + c*x**6)**p*x**3)/(18*a*p**2 + 15*a*p + 2*a +
18*c*p**2*x**6 + 15*c*p*x**6 + 2*c*x**6),x)*a*e*p**2 + 12*int(((a + c*x**6)**p*x**3)/(18*a*p**2 + 15*a*p + 2*a + 18*c*p**2*x**6 + 15*c*p*x**6 + 2*c*x**6),
x)*a*e*p)/(2*(18*p**2 + 15*p + 2))
```

3.10 $\int (d + ex^3)^q (a + cx^6)^p \, dx$

Optimal result	119
Mathematica [N/A]	119
Rubi [N/A]	120
Maple [N/A]	120
Fricas [N/A]	121
Sympy [F(-1)]	121
Maxima [N/A]	121
Giac [N/A]	122
Mupad [N/A]	122
Reduce [N/A]	123

Optimal result

Integrand size = 19, antiderivative size = 19

$$\int (d + ex^3)^q (a + cx^6)^p \, dx = \text{Int}\left((d + ex^3)^q (a + cx^6)^p, x\right)$$

output `Defer(Int)((e*x^3+d)^q*(c*x^6+a)^p,x)`

Mathematica [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int (d + ex^3)^q (a + cx^6)^p \, dx = \int (d + ex^3)^q (a + cx^6)^p \, dx$$

input `Integrate[(d + e*x^3)^q*(a + c*x^6)^p,x]`

output `Integrate[(d + e*x^3)^q*(a + c*x^6)^p, x]`

Rubi [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {1770}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + cx^6)^p (d + ex^3)^q dx \xrightarrow{1770} \int (a + cx^6)^p (d + ex^3)^q dx$$

input `Int[(d + e*x^3)^q*(a + c*x^6)^p,x]`

output `$Aborted`

Definitions of rubi rules used

rule 1770 `Int[((d_) + (e_.)*(x_)^(n_))^(q_)*((a_) + (c_.)*(x_)^(n2_))^(p_), x_Symbol] :> Unintegrable[(d + e*x^n)^q*(a + c*x^(2*n))^p, x] /; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n]`

Maple [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int (e x^3 + d)^q (c x^6 + a)^p dx$$

input `int((e*x^3+d)^q*(c*x^6+a)^p,x)`

output `int((e*x^3+d)^q*(c*x^6+a)^p,x)`

Fricas [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int (d + ex^3)^q (a + cx^6)^p dx = \int (cx^6 + a)^p (ex^3 + d)^q dx$$

input `integrate((e*x^3+d)^q*(c*x^6+a)^p,x, algorithm="fricas")`

output `integral((c*x^6 + a)^p*(e*x^3 + d)^q, x)`

Sympy [F(-1)]

Timed out.

$$\int (d + ex^3)^q (a + cx^6)^p dx = \text{Timed out}$$

input `integrate((e*x**3+d)**q*(c*x**6+a)**p,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int (d + ex^3)^q (a + cx^6)^p dx = \int (cx^6 + a)^p (ex^3 + d)^q dx$$

input `integrate((e*x^3+d)^q*(c*x^6+a)^p,x, algorithm="maxima")`

output `integrate((c*x^6 + a)^p*(e*x^3 + d)^q, x)`

Giac [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int (d + ex^3)^q (a + cx^6)^p dx = \int (cx^6 + a)^p (ex^3 + d)^q dx$$

input `integrate((e*x^3+d)^q*(c*x^6+a)^p,x, algorithm="giac")`

output `integrate((c*x^6 + a)^p*(e*x^3 + d)^q, x)`

Mupad [N/A]

Not integrable

Time = 10.79 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int (d + ex^3)^q (a + cx^6)^p dx = \int (cx^6 + a)^p (ex^3 + d)^q dx$$

input `int((a + c*x^6)^p*(d + e*x^3)^q,x)`

output `int((a + c*x^6)^p*(d + e*x^3)^q, x)`

Reduce [N/A]

Not integrable

Time = 0.85 (sec) , antiderivative size = 1249, normalized size of antiderivative = 65.74

$$\int (d + ex^3)^q (a + cx^6)^p dx = \text{Too large to display}$$

input `int((e*x^3+d)^q*(c*x^6+a)^p,x)`

output `((d + e*x**3)**q*(a + c*x**6)**p*x + 18*int(((d + e*x**3)**q*(a + c*x**6)*
*p*x**6)/(6*a*d*p + 3*a*d*q + a*d + 6*a*e*p*x**3 + 3*a*e*q*x**3 + a*e*x**3
+ 6*c*d*p*x**6 + 3*c*d*q*x**6 + c*d*x**6 + 6*c*e*p*x**9 + 3*c*e*q*x**9 +
c*e*x**9),x)*c*d*p*q + 9*int(((d + e*x**3)**q*(a + c*x**6)**p*x**6)/(6*a*d
*p + 3*a*d*q + a*d + 6*a*e*p*x**3 + 3*a*e*q*x**3 + a*e*x**3 + 6*c*d*p*x**6
+ 3*c*d*q*x**6 + c*d*x**6 + 6*c*e*p*x**9 + 3*c*e*q*x**9 + c*e*x**9),x)*c*
d*q**2 + 3*int(((d + e*x**3)**q*(a + c*x**6)**p*x**6)/(6*a*d*p + 3*a*d*q +
a*d + 6*a*e*p*x**3 + 3*a*e*q*x**3 + a*e*x**3 + 6*c*d*p*x**6 + 3*c*d*q*x**
6 + c*d*x**6 + 6*c*e*p*x**9 + 3*c*e*q*x**9 + c*e*x**9),x)*c*d*q + 36*int((
(d + e*x**3)**q*(a + c*x**6)**p*x**3)/(6*a*d*p + 3*a*d*q + a*d + 6*a*e*p*x
3 + 3*a*e*q*x3 + a*e*x**3 + 6*c*d*p*x**6 + 3*c*d*q*x**6 + c*d*x**6 + 6
*c*e*p*x**9 + 3*c*e*q*x**9 + c*e*x**9),x)*a*e*p**2 + 18*int(((d + e*x**3)*
q(a + c*x**6)**p*x**3)/(6*a*d*p + 3*a*d*q + a*d + 6*a*e*p*x**3 + 3*a*e*q
*x**3 + a*e*x**3 + 6*c*d*p*x**6 + 3*c*d*q*x**6 + c*d*x**6 + 6*c*e*p*x**9 +
3*c*e*q*x**9 + c*e*x**9),x)*a*e*p*q + 6*int(((d + e*x**3)**q*(a + c*x**6)
p*x3)/(6*a*d*p + 3*a*d*q + a*d + 6*a*e*p*x**3 + 3*a*e*q*x**3 + a*e*x**
3 + 6*c*d*p*x**6 + 3*c*d*q*x**6 + c*d*x**6 + 6*c*e*p*x**9 + 3*c*e*q*x**9 +
c*e*x**9),x)*a*e*p + 36*int(((d + e*x**3)**q*(a + c*x**6)**p)/(6*a*d*p +
3*a*d*q + a*d + 6*a*e*p*x**3 + 3*a*e*q*x**3 + a*e*x**3 + 6*c*d*p*x**6 + 3*
c*d*q*x**6 + c*d*x**6 + 6*c*e*p*x**9 + 3*c*e*q*x**9 + c*e*x**9),x)*a*d*...`

3.11 $\int (1 - x^3)^p (1 - x^6)^p \, dx$

Optimal result	124
Mathematica [F]	124
Rubi [A] (verified)	125
Maple [F]	126
Fricas [F]	126
Sympy [F]	126
Maxima [F]	127
Giac [F]	127
Mupad [F(-1)]	127
Reduce [F]	128

Optimal result

Integrand size = 19, antiderivative size = 23

$$\int (1 - x^3)^p (1 - x^6)^p \, dx = x \text{AppellF1} \left(\frac{1}{3}, -2p, -p, \frac{4}{3}, x^3, -x^3 \right)$$

output `x*AppellF1(1/3, -2*p, -p, 4/3, x^3, -x^3)`

Mathematica [F]

$$\int (1 - x^3)^p (1 - x^6)^p \, dx = \int (1 - x^3)^p (1 - x^6)^p \, dx$$

input `Integrate[(1 - x^3)^p*(1 - x^6)^p, x]`

output `Integrate[(1 - x^3)^p*(1 - x^6)^p, x]`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {1388, 936}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (1 - x^3)^p (1 - x^6)^p dx \\ & \quad \downarrow \textcolor{blue}{1388} \\ & \int (1 - x^3)^{2p} (x^3 + 1)^p dx \\ & \quad \downarrow \textcolor{blue}{936} \\ & x \operatorname{AppellF1} \left(\frac{1}{3}, -2p, -p, \frac{4}{3}, x^3, -x^3 \right) \end{aligned}$$

input `Int[(1 - x^3)^p*(1 - x^6)^p, x]`

output `x*AppellF1[1/3, -2*p, -p, 4/3, x^3, -x^3]`

Definitions of rubi rules used

rule 936 `Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] :> Simp[a^p*c^q*x*AppellF1[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`

rule 1388 `Int[(u_)*(a_ + (c_)*(x_)^(n2_))^(p_)*(d_ + (e_)*(x_)^(n_))^(q_), x_Symbol] :> Int[u*(d + e*x^n)^(p + q)*(a/d + (c/e)*x^n)^p, x] /; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[c*d^2 + a*e^2, 0] && (IntegerQ[p] || (GtQ[a, 0] && GtQ[d, 0]))`

Maple [F]

$$\int (-x^3 + 1)^p (-x^6 + 1)^p dx$$

input `int((-x^3+1)^p*(-x^6+1)^p,x)`

output `int((-x^3+1)^p*(-x^6+1)^p,x)`

Fricas [F]

$$\int (1 - x^3)^p (1 - x^6)^p dx = \int (-x^6 + 1)^p (-x^3 + 1)^p dx$$

input `integrate((-x^3+1)^p*(-x^6+1)^p,x, algorithm="fricas")`

output `integral((-x^6 + 1)^p*(-x^3 + 1)^p, x)`

Sympy [F]

$$\int (1 - x^3)^p (1 - x^6)^p dx = \int (-(x - 1) (x^2 + x + 1))^p (-(x - 1) (x + 1) (x^2 - x + 1) (x^2 + x + 1))^p dx$$

input `integrate((-x**3+1)**p*(-x**6+1)**p,x)`

output `Integral((-x - 1)*(x**2 + x + 1))**p*(-(x - 1)*(x + 1)*(x**2 - x + 1)*(x*x**2 + x + 1))**p, x)`

Maxima [F]

$$\int (1 - x^3)^p (1 - x^6)^p dx = \int (-x^6 + 1)^p (-x^3 + 1)^p dx$$

input `integrate((-x^3+1)^p*(-x^6+1)^p,x, algorithm="maxima")`

output `integrate((-x^6 + 1)^p*(-x^3 + 1)^p, x)`

Giac [F]

$$\int (1 - x^3)^p (1 - x^6)^p dx = \int (-x^6 + 1)^p (-x^3 + 1)^p dx$$

input `integrate((-x^3+1)^p*(-x^6+1)^p,x, algorithm="giac")`

output `integrate((-x^6 + 1)^p*(-x^3 + 1)^p, x)`

Mupad [F(-1)]

Timed out.

$$\int (1 - x^3)^p (1 - x^6)^p dx = \int (1 - x^3)^p (1 - x^6)^p dx$$

input `int((1 - x^3)^p*(1 - x^6)^p,x)`

output `int((1 - x^3)^p*(1 - x^6)^p, x)`

Reduce [F]

$$\int (1-x^3)^p (1-x^6)^p dx = \frac{(-x^3+1)^p (-x^6+1)^p x - 27 \left(\int \frac{(-x^3+1)^p (-x^6+1)^p x^3}{9px^6+x^6-9p-1} dx \right) p^2 - 3 \left(\int \frac{(-x^3+1)^p (-x^6+1)^p x^3}{9px^6+x^6-9p-1} dx \right) p - 81 \left(\int \frac{(-x^3+1)^p}{9px^6+x^6-9p-1} dx \right)}{9p+1}$$

input `int((-x^3+1)^p*(-x^6+1)^p,x)`

output `((- x**3 + 1)**p*(- x**6 + 1)**p*x - 27*int(((- x**3 + 1)**p*(- x**6 + 1)**p*x**3)/(9*p*x**6 - 9*p + x**6 - 1),x)*p**2 - 3*int(((- x**3 + 1)**p*(- x**6 + 1)**p*x**3)/(9*p*x**6 - 9*p + x**6 - 1),x)*p - 81*int(((- x**3 + 1)**p*(- x**6 + 1)**p)/(9*p*x**6 - 9*p + x**6 - 1),x)*p**2 - 9*int(((- x**3 + 1)**p*(- x**6 + 1)**p)/(9*p*x**6 - 9*p + x**6 - 1),x)*p)/(9*p + 1)`

3.12 $\int (d + ex^3)^q (d^2 - e^2x^6)^p dx$

Optimal result	129
Mathematica [F]	129
Rubi [A] (verified)	130
Maple [F]	131
Fricas [F]	131
Sympy [F(-1)]	132
Maxima [F]	132
Giac [F]	132
Mupad [F(-1)]	133
Reduce [F]	133

Optimal result

Integrand size = 24, antiderivative size = 92

$$\int (d + ex^3)^q (d^2 - e^2x^6)^p dx = x(d + ex^3)^q \left(1 - \frac{ex^3}{d}\right)^{-p} \left(1 + \frac{ex^3}{d}\right)^{-p-q} (d^2 - e^2x^6)^p \text{AppellF1}\left(\frac{1}{3}, -p, -p - q, \frac{4}{3}, \frac{ex^3}{d}, -\frac{ex^3}{d}\right)$$

output $x*(e*x^3+d)^q*(1+e*x^3/d)^{(-p-q)*(-e^2*x^6+d^2)}*AppellF1(1/3,-p,-p-q,4/3,e*x^3/d,-e*x^3/d)/((1-e*x^3/d)^p)$

Mathematica [F]

$$\int (d + ex^3)^q (d^2 - e^2x^6)^p dx = \int (d + ex^3)^q (d^2 - e^2x^6)^p dx$$

input `Integrate[(d + e*x^3)^q*(d^2 - e^2*x^6)^p, x]`

output `Integrate[(d + e*x^3)^q*(d^2 - e^2*x^6)^p, x]`

Rubi [A] (verified)

Time = 0.26 (sec), antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {1396, 937, 937, 936}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (d^2 - e^2 x^6)^p (d + ex^3)^q dx \\
 & \quad \downarrow \textcolor{blue}{1396} \\
 & (d - ex^3)^{-p} (d + ex^3)^{-p} (d^2 - e^2 x^6)^p \int (d - ex^3)^p (ex^3 + d)^{p+q} dx \\
 & \quad \downarrow \textcolor{blue}{937} \\
 & (d + ex^3)^{-p} \left(1 - \frac{ex^3}{d}\right)^{-p} (d^2 - e^2 x^6)^p \int (ex^3 + d)^{p+q} \left(1 - \frac{ex^3}{d}\right)^p dx \\
 & \quad \downarrow \textcolor{blue}{937} \\
 & \left(1 - \frac{ex^3}{d}\right)^{-p} (d^2 - e^2 x^6)^p (d + ex^3)^q \left(\frac{ex^3}{d} + 1\right)^{-p-q} \int \left(1 - \frac{ex^3}{d}\right)^p \left(\frac{ex^3}{d} + 1\right)^{p+q} dx \\
 & \quad \downarrow \textcolor{blue}{936} \\
 & x \left(1 - \frac{ex^3}{d}\right)^{-p} (d^2 - e^2 x^6)^p (d + ex^3)^q \left(\frac{ex^3}{d} + 1\right)^{-p-q} \text{AppellF1} \left(\frac{1}{3}, -p, -p - q, \frac{4}{3}, \frac{ex^3}{d}, -\frac{ex^3}{d}\right)
 \end{aligned}$$

input `Int[(d + e*x^3)^q*(d^2 - e^2*x^6)^p,x]`

output `(x*(d + e*x^3)^q*(1 + (e*x^3)/d)^(-p - q)*(d^2 - e^2*x^6)^p*AppellF1[1/3, -p, -p - q, 4/3, (e*x^3)/d, -((e*x^3)/d)])/(1 - (e*x^3)/d)^p`

Definitions of rubi rules used

rule 936 $\text{Int}[(a_.) + (b_.)*(x_.)^{(n_.)}^{(p_.)}*((c_.) + (d_.)*(x_.)^{(n_.)}^{(q_.)}), x_\text{Symbol}]$
 $\Rightarrow \text{Simp}[a^p c^q x^q \text{AppellF1}[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /; \text{FreeQ}[\{a, b, c, d, n, p, q\}, x] \&& \text{NeQ}[b*c - a*d, 0] \&& \text{NeQ}[n, -1] \&& (\text{IntegerQ}[p] \&& \text{GtQ}[a, 0]) \&& (\text{IntegerQ}[q] \&& \text{GtQ}[c, 0])$

rule 937 $\text{Int}[(a_.) + (b_.)*(x_.)^{(n_.)}^{(p_.)}*((c_.) + (d_.)*(x_.)^{(n_.)}^{(q_.)}), x_\text{Symbol}]$
 $\Rightarrow \text{Simp}[a^{\text{IntPart}[p]}*((a + b*x^n)^{\text{FracPart}[p]} / (1 + b*(x^n/a))^{\text{FracPart}[p]}) \text{Int}[(1 + b*(x^n/a))^{p_*} (c + d*x^n)^q, x], x] /; \text{FreeQ}[\{a, b, c, d, n, p, q\}, x] \&& \text{NeQ}[b*c - a*d, 0] \&& \text{NeQ}[n, -1] \&& !(\text{IntegerQ}[p] \&& \text{GtQ}[a, 0])$

rule 1396 $\text{Int}[(u_.)*((a_.) + (c_.)*(x_.)^{(n2_.)}^{(p_.)}*((d_.) + (e_.)*(x_.)^{(n_.)}^{(q_.)}), x_\text{Symbol}]$
 $\Rightarrow \text{Simp}[(a + c*x^{(2*n)})^{\text{FracPart}[p]} / ((d + e*x^n)^{\text{FracPart}[p]} * (a/d + c*(x^n/e))^{\text{FracPart}[p]}) \text{Int}[u*(d + e*x^n)^{p+q} * (a/d + (c/e)*x^n)^p, x], x] /; \text{FreeQ}[\{a, c, d, e, n, p, q\}, x] \&& \text{EqQ}[n2, 2*n] \&& \text{EqQ}[c*d^2 + a*e^2, 0] \&& !\text{IntegerQ}[p] \&& !(\text{EqQ}[q, 1] \&& \text{EqQ}[n, 2])$

Maple [F]

$$\int (e x^3 + d)^q (-e^2 x^6 + d^2)^p dx$$

input `int((e*x^3+d)^q*(-e^2*x^6+d^2)^p,x)`

output `int((e*x^3+d)^q*(-e^2*x^6+d^2)^p,x)`

Fricas [F]

$$\int (d + e x^3)^q (d^2 - e^2 x^6)^p dx = \int (-e^2 x^6 + d^2)^p (e x^3 + d)^q dx$$

input `integrate((e*x^3+d)^q*(-e^2*x^6+d^2)^p,x, algorithm="fricas")`

output `integral((-e^2*x^6 + d^2)^p*(e*x^3 + d)^q, x)`

Sympy [F(-1)]

Timed out.

$$\int (d + ex^3)^q (d^2 - e^2x^6)^p dx = \text{Timed out}$$

input `integrate((e*x**3+d)**q*(-e**2*x**6+d**2)**p,x)`

output `Timed out`

Maxima [F]

$$\int (d + ex^3)^q (d^2 - e^2x^6)^p dx = \int (-e^2x^6 + d^2)^p (ex^3 + d)^q dx$$

input `integrate((e*x^3+d)^q*(-e^2*x^6+d^2)^p,x, algorithm="maxima")`

output `integrate((-e^2*x^6 + d^2)^p*(e*x^3 + d)^q, x)`

Giac [F]

$$\int (d + ex^3)^q (d^2 - e^2x^6)^p dx = \int (-e^2x^6 + d^2)^p (ex^3 + d)^q dx$$

input `integrate((e*x^3+d)^q*(-e^2*x^6+d^2)^p,x, algorithm="giac")`

output `integrate((-e^2*x^6 + d^2)^p*(e*x^3 + d)^q, x)`

Mupad [F(-1)]

Timed out.

$$\int (d + ex^3)^q (d^2 - e^2 x^6)^p dx = \int (d^2 - e^2 x^6)^p (e x^3 + d)^q dx$$

input `int((d^2 - e^2*x^6)^p*(d + e*x^3)^q, x)`

output `int((d^2 - e^2*x^6)^p*(d + e*x^3)^q, x)`

Reduce [F]

$$\begin{aligned} & \int (d + ex^3)^q (d^2 - e^2 x^6)^p dx \\ &= \frac{(e x^3 + d)^q (-e^2 x^6 + d^2)^p x - 18 \left(\int \frac{(e x^3 + d)^q (-e^2 x^6 + d^2)^p x^3}{-6e^2 p x^6 - 3e^2 q x^6 - e^2 x^6 + 6d^2 p + 3d^2 q + d^2} dx \right) depq - 9 \left(\int \frac{(e x^3 + d)^q (-e^2 x^6 + d^2)^p}{-6e^2 p x^6 - 3e^2 q x^6 - e^2 x^6 + 6d^2 p + 3d^2 q + d^2} dx \right) depq^2}{\dots} \end{aligned}$$

input `int((e*x^3+d)^q*(-e^2*x^6+d^2)^p, x)`

output `((d + e*x**3)**q*(d**2 - e**2*x**6)**p*x - 18*int(((d + e*x**3)**q*(d**2 - e**2*x**6)**p*x**3)/(6*d**2*p + 3*d**2*q + d**2 - 6*e**2*p*x**6 - 3*e**2*p*x**3), x)*d*e*p*q - 9*int(((d + e*x**3)**q*(d**2 - e**2*x**6)**p*x**3)/(6*d**2*p + 3*d**2*q + d**2 - 6*e**2*p*x**6 - 3*e**2*p*x**3), x)*d*e*p*q - 3*int(((d + e*x**3)**q*(d**2 - e**2*x**6)**p*x**3)/(6*d**2*p + 3*d**2*q + d**2 - 6*e**2*p*x**6 - 3*e**2*p*x**3), x)*d*e*p*q - 36*int(((d + e*x**3)**q*(d**2 - e**2*x**6)**p*x)/(6*d**2*p + 3*d**2*q + d**2 - 6*e**2*p*x**6 - 3*e**2*p*x**3), x)*d*e*p*q + 36*int(((d + e*x**3)**q*(d**2 - e**2*x**6)**p*x)/(6*d**2*p + 3*d**2*q + d**2 - 6*e**2*p*x**6 - 3*e**2*p*x**3), x)*d*e*p*q + 6*int(((d + e*x**3)**q*(d**2 - e**2*x**6)**p*x)/(6*d**2*p + 3*d**2*q + d**2 - 6*e**2*p*x**6 - 3*e**2*p*x**3), x)*d*e*p*q + 6*int(((d + e*x**3)**q*(d**2 - e**2*x**6)**p*x)/(6*d**2*p + 3*d**2*q + d**2 - 6*e**2*p*x**6 - 3*e**2*p*x**3), x)*d*e*p*q + 9*int(((d + e*x**3)**q*(d**2 - e**2*x**6)**p*x)/(6*d**2*p + 3*d**2*q + d**2 - 6*e**2*p*x**6 - 3*e**2*p*x**3), x)*d*e*p*q + 3*int(((d + e*x**3)**q*(d**2 - e**2*x**6)**p*x)/(6*d**2*p + 3*d**2*q + d**2 - 6*e**2*p*x**6 - 3*e**2*p*x**3), x)*d*e*p*q)/(6*p + 3*q + 1)`

3.13 $\int (1 - x^3)^{-p} (1 - x^6)^p \, dx$

Optimal result	134
Mathematica [F]	134
Rubi [A] (verified)	135
Maple [F]	136
Fricas [F]	136
Sympy [F(-1)]	136
Maxima [F]	137
Giac [F]	137
Mupad [F(-1)]	137
Reduce [F]	138

Optimal result

Integrand size = 21, antiderivative size = 17

$$\int (1 - x^3)^{-p} (1 - x^6)^p \, dx = x \text{Hypergeometric2F1}\left(\frac{1}{3}, -p, \frac{4}{3}, -x^3\right)$$

output `x*hypergeom([1/3, -p], [4/3], -x^3)`

Mathematica [F]

$$\int (1 - x^3)^{-p} (1 - x^6)^p \, dx = \int (1 - x^3)^{-p} (1 - x^6)^p \, dx$$

input `Integrate[(1 - x^6)^p/(1 - x^3)^p, x]`

output `Integrate[(1 - x^6)^p/(1 - x^3)^p, x]`

Rubi [A] (verified)

Time = 0.16 (sec), antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {1386, 778}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (1 - x^3)^{-p} (1 - x^6)^p dx \\ & \quad \downarrow \textcolor{blue}{1386} \\ & \int (x^3 + 1)^p dx \\ & \quad \downarrow \textcolor{blue}{778} \\ & x \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, -p, \frac{4}{3}, -x^3\right) \end{aligned}$$

input `Int[(1 - x^6)^p/(1 - x^3)^p, x]`

output `x*Hypergeometric2F1[1/3, -p, 4/3, -x^3]`

Definitions of rubi rules used

rule 778 `Int[((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F1[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] || GtQ[a, 0])`

rule 1386 `Int[(u_)*(a_) + (c_)*(x_)^(n2_))^(p_)*((d_) + (e_)*(x_)^(n_))^(q_), x_Symbol] := Simp[(-e^2/c)^q Int[u*(d - e*x^n)^p, x], x] /; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[c*d^2 + a*e^2, 0] && EqQ[p + q, 0] && GtQ[d, 0] && LtQ[c, 0] && GtQ[e^2, 0]`

Maple [F]

$$\int (-x^6 + 1)^p (-x^3 + 1)^{-p} dx$$

input `int((-x^6+1)^p/((-x^3+1)^p),x)`

output `int((-x^6+1)^p/((-x^3+1)^p),x)`

Fricas [F]

$$\int (1 - x^3)^{-p} (1 - x^6)^p dx = \int \frac{(-x^6 + 1)^p}{(-x^3 + 1)^p} dx$$

input `integrate((-x^6+1)^p/((-x^3+1)^p),x, algorithm="fricas")`

output `integral((-x^6 + 1)^p/(-x^3 + 1)^p, x)`

Sympy [F(-1)]

Timed out.

$$\int (1 - x^3)^{-p} (1 - x^6)^p dx = \text{Timed out}$$

input `integrate((-x**6+1)**p/((-x**3+1)**p),x)`

output `Timed out`

Maxima [F]

$$\int (1 - x^3)^{-p} (1 - x^6)^p \, dx = \int \frac{(-x^6 + 1)^p}{(-x^3 + 1)^p} \, dx$$

input `integrate((-x^6+1)^p/((-x^3+1)^p),x, algorithm="maxima")`

output `integrate((-x^6 + 1)^p/(-x^3 + 1)^p, x)`

Giac [F]

$$\int (1 - x^3)^{-p} (1 - x^6)^p \, dx = \int \frac{(-x^6 + 1)^p}{(-x^3 + 1)^p} \, dx$$

input `integrate((-x^6+1)^p/((-x^3+1)^p),x, algorithm="giac")`

output `integrate((-x^6 + 1)^p/(-x^3 + 1)^p, x)`

Mupad [F(-1)]

Timed out.

$$\int (1 - x^3)^{-p} (1 - x^6)^p \, dx = \int \frac{(1 - x^6)^p}{(1 - x^3)^p} \, dx$$

input `int((1 - x^6)^p/(1 - x^3)^p,x)`

output `int((1 - x^6)^p/(1 - x^3)^p, x)`

Reduce [F]

$$\int (1 - x^3)^{-p} (1 - x^6)^p \, dx = \int \frac{(-x^6 + 1)^p}{(-x^3 + 1)^p} dx$$

input `int((-x^6+1)^p/((-x^3+1)^p),x)`

output `int((- x**6 + 1)**p / (- x**3 + 1)**p, x)`

3.14 $\int \frac{d+ex^4}{a-cx^8} dx$

Optimal result	139
Mathematica [A] (verified)	140
Rubi [A] (verified)	141
Maple [C] (verified)	146
Fricas [B] (verification not implemented)	147
Sympy [F(-1)]	148
Maxima [F]	148
Giac [B] (verification not implemented)	148
Mupad [B] (verification not implemented)	149
Reduce [B] (verification not implemented)	150

Optimal result

Integrand size = 18, antiderivative size = 263

$$\begin{aligned} \int \frac{d+ex^4}{a-cx^8} dx = & \frac{(\sqrt{cd} + \sqrt{ae}) \arctan\left(\frac{\sqrt[8]{cx}}{\sqrt[8]{a}}\right)}{4a^{7/8}c^{5/8}} - \frac{\left(d - \frac{\sqrt{ae}}{\sqrt{c}}\right) \arctan\left(1 - \frac{\sqrt{2}\sqrt[8]{cx}}{\sqrt[8]{a}}\right)}{4\sqrt{2}a^{7/8}\sqrt[8]{c}} \\ & + \frac{\left(d - \frac{\sqrt{ae}}{\sqrt{c}}\right) \arctan\left(1 + \frac{\sqrt{2}\sqrt[8]{cx}}{\sqrt[8]{a}}\right)}{4\sqrt{2}a^{7/8}\sqrt[8]{c}} + \frac{(\sqrt{cd} + \sqrt{ae}) \operatorname{arctanh}\left(\frac{\sqrt[8]{cx}}{\sqrt[8]{a}}\right)}{4a^{7/8}c^{5/8}} \\ & + \frac{\left(d - \frac{\sqrt{ae}}{\sqrt{c}}\right) \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[8]{a}\sqrt[8]{cx}}{\sqrt[4]{a}+\sqrt[4]{cx^2}}\right)}{4\sqrt{2}a^{7/8}\sqrt[8]{c}} \end{aligned}$$

output

```
1/4*(c^(1/2)*d+a^(1/2)*e)*arctan(c^(1/8)*x/a^(1/8))/a^(7/8)/c^(5/8)+1/8*(d-a^(1/2)*e/c^(1/2))*arctan(-1+2^(1/2)*c^(1/8)*x/a^(1/8))*2^(1/2)/a^(7/8)/c^(1/8)+1/8*(d-a^(1/2)*e/c^(1/2))*arctan(1+2^(1/2)*c^(1/8)*x/a^(1/8))*2^(1/2)/a^(7/8)/c^(1/8)+1/4*(c^(1/2)*d+a^(1/2)*e)*arctanh(c^(1/8)*x/a^(1/8))/a^(7/8)/c^(5/8)+1/8*(d-a^(1/2)*e/c^(1/2))*arctanh(2^(1/2)*a^(1/8)*c^(1/8)*x/(a^(1/4)+c^(1/4)*x^2))*2^(1/2)/a^(7/8)/c^(1/8)
```

Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 425, normalized size of antiderivative = 1.62

$$\int \frac{d + ex^4}{a - cx^8} dx = \frac{(\sqrt[8]{a}\sqrt{cd} + a^{5/8}e) \arctan\left(\frac{\sqrt[8]{cx}}{\sqrt[8]{a}}\right)}{4ac^{5/8}} \\ - \frac{(-\sqrt[8]{a}\sqrt{cd} + a^{5/8}e) \arctan\left(\frac{-\sqrt{2}\sqrt[8]{a} + 2\sqrt[8]{cx}}{\sqrt{2}\sqrt[8]{a}}\right)}{4\sqrt{2}ac^{5/8}} \\ - \frac{(-\sqrt[8]{a}\sqrt{cd} + a^{5/8}e) \arctan\left(\frac{\sqrt{2}\sqrt[8]{a} + 2\sqrt[8]{cx}}{\sqrt{2}\sqrt[8]{a}}\right)}{4\sqrt{2}ac^{5/8}} \\ - \frac{(\sqrt[8]{a}\sqrt{cd} + a^{5/8}e) \log(\sqrt[8]{a} - \sqrt[8]{cx})}{8ac^{5/8}} \\ - \frac{(-\sqrt[8]{a}\sqrt{cd} - a^{5/8}e) \log(\sqrt[8]{a} + \sqrt[8]{cx})}{8ac^{5/8}} \\ + \frac{(-\sqrt[8]{a}\sqrt{cd} + a^{5/8}e) \log(\sqrt[4]{a} - \sqrt{2}\sqrt[8]{a}\sqrt[8]{cx} + \sqrt[4]{cx^2})}{8\sqrt{2}ac^{5/8}} \\ - \frac{(-\sqrt[8]{a}\sqrt{cd} + a^{5/8}e) \log(\sqrt[4]{a} + \sqrt{2}\sqrt[8]{a}\sqrt[8]{cx} + \sqrt[4]{cx^2})}{8\sqrt{2}ac^{5/8}}$$

input `Integrate[(d + e*x^4)/(a - c*x^8), x]`

output

```
((a^(1/8)*Sqrt[c]*d + a^(5/8)*e)*ArcTan[(c^(1/8)*x)/a^(1/8)])/(4*a*c^(5/8)) - ((-a^(1/8)*Sqrt[c]*d + a^(5/8)*e)*ArcTan[((-Sqrt[2]*a^(1/8)) + 2*c^(1/8)*x)/(Sqrt[2]*a^(1/8))])/(4*Sqrt[2]*a*c^(5/8)) - ((-a^(1/8)*Sqrt[c]*d + a^(5/8)*e)*ArcTan[(Sqrt[2]*a^(1/8) + 2*c^(1/8)*x)/(Sqrt[2]*a^(1/8))])/(4*Sqrt[2]*a*c^(5/8)) - ((a^(1/8)*Sqrt[c]*d + a^(5/8)*e)*Log[a^(1/8) - c^(1/8)*x])/(8*a*c^(5/8)) - ((-a^(1/8)*Sqrt[c]*d - a^(5/8)*e)*Log[a^(1/8) + c^(1/8)*x])/(8*a*c^(5/8)) + ((-a^(1/8)*Sqrt[c]*d + a^(5/8)*e)*Log[a^(1/4) - Sqrt[2]*a^(1/8)*c^(1/8)*x + c^(1/4)*x^2])/(8*Sqrt[2]*a*c^(5/8)) - ((-a^(1/8)*Sqrt[c]*d + a^(5/8)*e)*Log[a^(1/4) + Sqrt[2]*a^(1/8)*c^(1/8)*x + c^(1/4)*x^2])/(8*Sqrt[2]*a*c^(5/8))
```

Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 293, normalized size of antiderivative = 1.11, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.722$, Rules used = {1747, 755, 27, 756, 218, 221, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{d + ex^4}{a - cx^8} dx \\
 & \downarrow \textcolor{blue}{1747} \\
 & \frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \int \frac{1}{a - \sqrt{a}\sqrt{cx^4}} dx + \frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \int \frac{1}{\sqrt{a}\sqrt{cx^4} + a} dx \\
 & \downarrow \textcolor{blue}{755} \\
 & \frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \int \frac{1}{a - \sqrt{a}\sqrt{cx^4}} dx + \frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \left(\frac{\int \frac{\sqrt[4]{a} - \sqrt[4]{cx^2}}{\sqrt{a}(\sqrt{cx^4} + \sqrt{a})} dx}{2\sqrt[4]{a}} + \frac{\int \frac{\sqrt[4]{cx^2} + \sqrt[4]{a}}{\sqrt{a}(\sqrt{cx^4} + \sqrt{a})} dx}{2\sqrt[4]{a}} \right) \\
 & \downarrow \textcolor{blue}{27} \\
 & \frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \left(\frac{\int \frac{\sqrt[4]{a} - \sqrt[4]{cx^2}}{\sqrt{cx^4} + \sqrt{a}} dx}{2a^{3/4}} + \frac{\int \frac{\sqrt[4]{cx^2} + \sqrt[4]{a}}{\sqrt{cx^4} + \sqrt{a}} dx}{2a^{3/4}} \right) + \frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \int \frac{1}{a - \sqrt{a}\sqrt{cx^4}} dx \\
 & \downarrow \textcolor{blue}{756} \\
 & \frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \left(\frac{\int \frac{1}{\sqrt[4]{a} - \sqrt[4]{cx^2}} dx}{2a^{3/4}} + \frac{\int \frac{1}{\sqrt[4]{cx^2} + \sqrt[4]{a}} dx}{2a^{3/4}} \right) + \\
 & \frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \left(\frac{\int \frac{\sqrt[4]{a} - \sqrt[4]{cx^2}}{\sqrt{cx^4} + \sqrt{a}} dx}{2a^{3/4}} + \frac{\int \frac{\sqrt[4]{cx^2} + \sqrt[4]{a}}{\sqrt{cx^4} + \sqrt{a}} dx}{2a^{3/4}} \right) \\
 & \downarrow \textcolor{blue}{218}
 \end{aligned}$$

$$\frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \left(\frac{\int \frac{1}{\sqrt[4]{a - \sqrt[4]{cx^2}}} dx}{2a^{3/4}} + \frac{\arctan \left(\frac{\sqrt[8]{cx}}{\sqrt[8]{a}} \right)}{2a^{7/8} \sqrt[8]{c}} \right) + \\ \frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \left(\frac{\int \frac{\sqrt[4]{a - \sqrt[4]{cx^2}}}{\sqrt{cx^4 + \sqrt{a}}} dx}{2a^{3/4}} + \frac{\int \frac{\sqrt[4]{cx^2 + \sqrt{a}}}{\sqrt{cx^4 + \sqrt{a}}} dx}{2a^{3/4}} \right)$$

↓ 221

$$\frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \left(\frac{\int \frac{\sqrt[4]{a - \sqrt[4]{cx^2}}}{\sqrt{cx^4 + \sqrt{a}}} dx}{2a^{3/4}} + \frac{\int \frac{\sqrt[4]{cx^2 + \sqrt{a}}}{\sqrt{cx^4 + \sqrt{a}}} dx}{2a^{3/4}} \right) + \\ \frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \left(\frac{\arctan \left(\frac{\sqrt[8]{cx}}{\sqrt[8]{a}} \right)}{2a^{7/8} \sqrt[8]{c}} + \frac{\operatorname{arctanh} \left(\frac{\sqrt[8]{cx}}{\sqrt[8]{a}} \right)}{2a^{7/8} \sqrt[8]{c}} \right)$$

↓ 1476

$$\frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \left(\frac{\int \frac{\frac{1}{x^2 - \frac{\sqrt{2}}{8}\sqrt[8]{a}x + \frac{4}{8}\sqrt{a}}}{\sqrt[8]{c}} dx}{2\sqrt[4]{c}} + \frac{\int \frac{\frac{1}{x^2 + \frac{\sqrt{2}}{8}\sqrt[8]{a}x + \frac{4}{8}\sqrt{a}}}{\sqrt[8]{c}} dx}{2\sqrt[4]{c}} + \frac{\int \frac{\sqrt[4]{a - \sqrt[4]{cx^2}}}{\sqrt{cx^4 + \sqrt{a}}} dx}{2a^{3/4}} \right) + \\ \frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \left(\frac{\arctan \left(\frac{\sqrt[8]{cx}}{\sqrt[8]{a}} \right)}{2a^{7/8} \sqrt[8]{c}} + \frac{\operatorname{arctanh} \left(\frac{\sqrt[8]{cx}}{\sqrt[8]{a}} \right)}{2a^{7/8} \sqrt[8]{c}} \right)$$

↓ 1082

$$\frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \left(\frac{\int \frac{\sqrt[4]{a - \sqrt[4]{cx^2}}}{\sqrt{cx^4 + \sqrt{a}}} dx}{2a^{3/4}} + \frac{\frac{\int \frac{1}{\left(1 - \frac{\sqrt{2}}{8}\sqrt[8]{cx} \right)^2} d \left(1 - \frac{\sqrt{2}}{8}\sqrt[8]{cx} \right)}{\sqrt{2}\sqrt[8]{a}\sqrt[8]{c}} - \frac{\int \frac{1}{\left(\frac{\sqrt{2}}{8}\sqrt[8]{cx} + 1 \right)^2} d \left(\frac{\sqrt{2}}{8}\sqrt[8]{cx} + 1 \right)}{\sqrt{2}\sqrt[8]{a}\sqrt[8]{c}}}{2a^{3/4}} \right) + \\ \frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \left(\frac{\arctan \left(\frac{\sqrt[8]{cx}}{\sqrt[8]{a}} \right)}{2a^{7/8} \sqrt[8]{c}} + \frac{\operatorname{arctanh} \left(\frac{\sqrt[8]{cx}}{\sqrt[8]{a}} \right)}{2a^{7/8} \sqrt[8]{c}} \right)$$

$$\begin{aligned}
& \downarrow 217 \\
& \frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \left(\frac{\int \frac{\sqrt[4]{a} - \sqrt[4]{cx^2}}{\sqrt{cx^4 + \sqrt{a}}} dx}{2a^{3/4}} + \frac{\arctan \left(\frac{\sqrt{2} \sqrt[8]{cx}}{\sqrt[8]{a}} + 1 \right)}{\sqrt{2} \sqrt[8]{a} \sqrt[8]{c}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[8]{cx}}{\sqrt[8]{a}} \right)}{\sqrt{2} \sqrt[8]{a} \sqrt[8]{c}} \right) + \\
& \quad \frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \left(\frac{\arctan \left(\frac{\sqrt[8]{cx}}{\sqrt[8]{a}} \right)}{2a^{7/8} \sqrt[8]{c}} + \frac{\operatorname{arctanh} \left(\frac{\sqrt[8]{cx}}{\sqrt[8]{a}} \right)}{2a^{7/8} \sqrt[8]{c}} \right) \\
& \quad \downarrow 1479 \\
& \frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \left(\frac{- \frac{\int - \frac{\sqrt{2} \sqrt[8]{a} - \sqrt[8]{cx}}{\sqrt[8]{c} \left(x^2 - \frac{\sqrt{2} \sqrt[8]{ax}}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}} \right)} dx}{2\sqrt{2} \sqrt[8]{a} \sqrt[8]{c}} - \frac{\int - \frac{\sqrt{2} \left(\sqrt{2} \sqrt[8]{cx} + \sqrt[8]{a} \right)}{\sqrt[8]{c} \left(x^2 + \frac{\sqrt{2} \sqrt[8]{ax}}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}} \right)} dx}{2\sqrt{2} \sqrt[8]{a} \sqrt[8]{c}} + \frac{\arctan \left(\frac{\sqrt{2} \sqrt[8]{cx}}{\sqrt[8]{a}} + 1 \right)}{\sqrt{2} \sqrt[8]{a} \sqrt[8]{c}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[8]{cx}}{\sqrt[8]{a}} \right)}{\sqrt{2} \sqrt[8]{a} \sqrt[8]{c}}}{2a^{3/4}} \right. \\
& \quad \left. \frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \left(\frac{\arctan \left(\frac{\sqrt[8]{cx}}{\sqrt[8]{a}} \right)}{2a^{7/8} \sqrt[8]{c}} + \frac{\operatorname{arctanh} \left(\frac{\sqrt[8]{cx}}{\sqrt[8]{a}} \right)}{2a^{7/8} \sqrt[8]{c}} \right) \right. \\
& \quad \downarrow 25 \\
& \frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \left(\frac{\int \frac{\sqrt{2} \sqrt[8]{a} - \sqrt[8]{cx}}{\sqrt[8]{c} \left(x^2 - \frac{\sqrt{2} \sqrt[8]{ax}}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}} \right)} dx}{2a^{3/4}} + \frac{\int \frac{\sqrt{2} \left(\sqrt{2} \sqrt[8]{cx} + \sqrt[8]{a} \right)}{\sqrt[8]{c} \left(x^2 + \frac{\sqrt{2} \sqrt[8]{ax}}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}} \right)} dx}{2\sqrt{2} \sqrt[8]{a} \sqrt[8]{c}} + \frac{\arctan \left(\frac{\sqrt{2} \sqrt[8]{cx}}{\sqrt[8]{a}} + 1 \right)}{\sqrt{2} \sqrt[8]{a} \sqrt[8]{c}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[8]{cx}}{\sqrt[8]{a}} \right)}{\sqrt{2} \sqrt[8]{a} \sqrt[8]{c}} \right. \\
& \quad \left. \frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \left(\frac{\arctan \left(\frac{\sqrt[8]{cx}}{\sqrt[8]{a}} \right)}{2a^{7/8} \sqrt[8]{c}} + \frac{\operatorname{arctanh} \left(\frac{\sqrt[8]{cx}}{\sqrt[8]{a}} \right)}{2a^{7/8} \sqrt[8]{c}} \right) \right. \\
& \quad \downarrow 27
\end{aligned}$$

$$\begin{aligned}
& \frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \left(\begin{array}{l} \int \frac{\sqrt{2} \sqrt[8]{a} - 2 \sqrt[8]{c_x} dx}{x^2 - \frac{\sqrt{2} \sqrt[8]{a} x + \sqrt[4]{a}}{\sqrt[8]{c} + \sqrt[4]{c}}} + \int \frac{\sqrt{2} \sqrt[8]{c_x} + \sqrt[8]{a} dx}{x^2 + \frac{\sqrt{2} \sqrt[8]{a} x + \sqrt[4]{a}}{\sqrt[8]{c} + \sqrt[4]{c}}} + \frac{\arctan\left(\frac{\sqrt{2} \sqrt[8]{c_x}}{\sqrt[8]{a}} + 1\right)}{\sqrt{2} \sqrt[8]{a} \sqrt[8]{c}} - \frac{\arctan\left(1 - \frac{\sqrt{2} \sqrt[8]{c_x}}{\sqrt[8]{a}}\right)}{\sqrt{2} \sqrt[8]{a} \sqrt[8]{c}} \\ \frac{2 \sqrt{2} \sqrt[8]{a} \sqrt[4]{c}}{2a^{3/4}} + \frac{2 \sqrt[8]{a} \sqrt[4]{c}}{2a^{3/4}} + \end{array} \right) + \\
& \frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \left(\frac{\arctan\left(\frac{\sqrt[8]{c_x}}{\sqrt[8]{a}}\right)}{2a^{7/8} \sqrt[8]{c}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[8]{c_x}}{\sqrt[8]{a}}\right)}{2a^{7/8} \sqrt[8]{c}} \right) \\
& \quad \downarrow \text{1103} \\
& \frac{1}{2} \left(\frac{\sqrt{ae}}{\sqrt{c}} + d \right) \left(\frac{\arctan\left(\frac{\sqrt[8]{c_x}}{\sqrt[8]{a}}\right)}{2a^{7/8} \sqrt[8]{c}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[8]{c_x}}{\sqrt[8]{a}}\right)}{2a^{7/8} \sqrt[8]{c}} \right) + \\
& \frac{1}{2} \left(d - \frac{\sqrt{ae}}{\sqrt{c}} \right) \left(\begin{array}{l} \frac{\arctan\left(\frac{\sqrt{2} \sqrt[8]{c_x}}{\sqrt[8]{a}} + 1\right)}{\sqrt{2} \sqrt[8]{a} \sqrt[8]{c}} - \frac{\arctan\left(1 - \frac{\sqrt{2} \sqrt[8]{c_x}}{\sqrt[8]{a}}\right)}{\sqrt{2} \sqrt[8]{a} \sqrt[8]{c}} + \frac{\log\left(\sqrt{2} \sqrt[8]{a} \sqrt[8]{c_x} + \sqrt[4]{a} + \sqrt[4]{c_x^2}\right)}{2\sqrt{2} \sqrt[8]{a} \sqrt[8]{c}} - \frac{\log\left(-\sqrt{2} \sqrt[8]{a} \sqrt[8]{c_x} + \sqrt[4]{a} + \sqrt[4]{c_x^2}\right)}{2\sqrt{2} \sqrt[8]{a} \sqrt[8]{c}} \\ 2a^{3/4} \end{array} \right)
\end{aligned}$$

input `Int[(d + e*x^4)/(a - c*x^8), x]`

output

```
((d + (Sqrt[a]*e)/Sqrt[c])*(ArcTan[(c^(1/8)*x)/a^(1/8)]/(2*a^(7/8)*c^(1/8)) + ArcTanh[(c^(1/8)*x)/a^(1/8)]/(2*a^(7/8)*c^(1/8)))/2 + ((d - (Sqrt[a]*e)/Sqrt[c])*((-(ArcTan[1 - (Sqrt[2]*c^(1/8)*x)/a^(1/8)])/(Sqrt[2]*a^(1/8)*c^(1/8)) + ArcTan[1 + (Sqrt[2]*c^(1/8)*x)/a^(1/8)]/(Sqrt[2]*a^(1/8)*c^(1/8)))/(2*a^(3/4)) + (-1/2*Log[a^(1/4) - Sqrt[2]*a^(1/8)*c^(1/8)*x + c^(1/4)*x^2]/(Sqrt[2]*a^(1/8)*c^(1/8)) + Log[a^(1/4) + Sqrt[2]*a^(1/8)*c^(1/8)*x + c^(1/4)*x^2]/(2*Sqrt[2]*a^(1/8)*c^(1/8)))/(2*a^(3/4)))/2
```

Definitions of rubi rules used

rule 25 $\text{Int}[-(\text{Fx}__), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$

rule 27 $\text{Int}[(\text{a}__)*(\text{Fx}__), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] /; \text{FreeQ}[\text{a}, \text{x}] \& \text{!Ma}\\ \text{tchQ}[\text{Fx}, (\text{b}__)*(\text{Gx}__)] /; \text{FreeQ}[\text{b}, \text{x}]$

rule 217 $\text{Int}[((\text{a}__) + (\text{b}__.)*(\text{x}__)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(-(\text{Rt}[-\text{a}, 2]*\text{Rt}[-\text{b}, 2])^{(-1)})*\text{ArcTan}[\text{Rt}[-\text{b}, 2]*(\text{x}/\text{Rt}[-\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \& \text{PosQ}[\text{a}/\text{b}] \& (\text{LtQ}[\text{a}, 0] \mid \text{LtQ}[\text{b}, 0])$

rule 218 $\text{Int}[((\text{a}__) + (\text{b}__.)*(\text{x}__)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[\text{a}/\text{b}, 2]/\text{a})*\text{ArcTan}[\text{x}/\text{Rt}[\text{a}/\text{b}, 2]], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \& \text{PosQ}[\text{a}/\text{b}]$

rule 221 $\text{Int}[((\text{a}__) + (\text{b}__.)*(\text{x}__)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-\text{a}/\text{b}, 2]/\text{a})*\text{ArcTanh}[\text{x}/\text{Rt}[-\text{a}/\text{b}, 2]], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \& \text{NegQ}[\text{a}/\text{b}]$

rule 755 $\text{Int}[((\text{a}__) + (\text{b}__.)*(\text{x}__)^4)^{-1}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{r} = \text{Numerator}[\text{Rt}[\text{a}/\text{b}, 2]], \text{s} = \text{Denominator}[\text{Rt}[\text{a}/\text{b}, 2]]\}, \text{Simp}[1/(2*\text{r}) \quad \text{Int}[(\text{r} - \text{s}*\text{x}^2)/(\text{a} + \text{b}*\text{x}^4), \text{x}], \text{x}] + \text{Simp}[1/(2*\text{r}) \quad \text{Int}[(\text{r} + \text{s}*\text{x}^2)/(\text{a} + \text{b}*\text{x}^4), \text{x}], \text{x}]] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \& (\text{GtQ}[\text{a}/\text{b}, 0] \mid (\text{PosQ}[\text{a}/\text{b}] \& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, \text{a}]] \& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, \text{b}]]))$

rule 756 $\text{Int}[((\text{a}__) + (\text{b}__.)*(\text{x}__)^4)^{-1}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{r} = \text{Numerator}[\text{Rt}[-\text{a}/\text{b}, 2]], \text{s} = \text{Denominator}[\text{Rt}[-\text{a}/\text{b}, 2]]\}, \text{Simp}[\text{r}/(2*\text{a}) \quad \text{Int}[1/(\text{r} - \text{s}*\text{x}^2), \text{x}], \text{x}] + \text{Simp}[\text{r}/(2*\text{a}) \quad \text{Int}[1/(\text{r} + \text{s}*\text{x}^2), \text{x}], \text{x}]] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \& \text{!GtQ}[\text{a}/\text{b}, 0]$

rule 1082 $\text{Int}[((\text{a}__) + (\text{b}__.)*(\text{x}__) + (\text{c}__.)*(\text{x}__)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{q} = 1 - 4*\text{S}\\ \text{implify}[\text{a}*(\text{c}/\text{b}^2)]\}, \text{Simp}[-2/\text{b} \quad \text{Subst}[\text{Int}[1/(\text{q} - \text{x}^2), \text{x}], \text{x}, 1 + 2*\text{c}*(\text{x}/\text{b})], \text{x}]] /; \text{RationalQ}[\text{q}] \& (\text{EqQ}[\text{q}^2, 1] \mid \text{!RationalQ}[\text{b}^2 - 4*\text{a}*\text{c}]) /; \text{Fre}\\ \text{eQ}[\{\text{a}, \text{b}, \text{c}\}, \text{x}]$

rule 1103 $\text{Int}[(d_ + e_)*x_)/((a_ + b_)*x_ + c_)*x_^2], x_{\text{Symbol}}] \Rightarrow S \text{imp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{EqQ}[2*c*d - b*e, 0]$

rule 1476 $\text{Int}[(d_ + e_)*x_^2)/((a_ + c_)*x_^4], x_{\text{Symbol}}] \Rightarrow \text{With}[\{q = \text{Rt}[2*(d/e), 2]\}, \text{Simp}[e/(2*c) \text{Int}[1/\text{Simp}[d/e + q*x + x^2, x], x] + \text{Simp}[e/(2*c) \text{Int}[1/\text{Simp}[d/e - q*x + x^2, x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&& \text{EqQ}[c*d^2 - a*e^2, 0] \&& \text{PosQ}[d*e]$

rule 1479 $\text{Int}[(d_ + e_)*x_^2)/((a_ + c_)*x_^4], x_{\text{Symbol}}] \Rightarrow \text{With}[\{q = \text{Rt}[-2*(d/e), 2]\}, \text{Simp}[e/(2*c*q) \text{Int}[(q - 2*x)/\text{Simp}[d/e + q*x - x^2, x], x] + \text{Simp}[e/(2*c*q) \text{Int}[(q + 2*x)/\text{Simp}[d/e - q*x - x^2, x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&& \text{EqQ}[c*d^2 - a*e^2, 0] \&& \text{NegQ}[d*e]$

rule 1747 $\text{Int}[(d_ + e_)*x_^{(n_)})/((a_ + c_)*x_^{(n2_)}), x_{\text{Symbol}}] \Rightarrow \text{With}[\{q = \text{Rt}[-a/c, 2]\}, \text{Simp}[(d + e*q)/2 \text{Int}[1/(a + c*q*x^n), x] + \text{Simp}[(d - e*q)/2 \text{Int}[1/(a - c*q*x^n), x]] /; \text{FreeQ}[\{a, c, d, e, n\}, x] \&& \text{EqQ}[n2, 2*n] \&& \text{NeQ}[c*d^2 + a*e^2, 0] \&& \text{NegQ}[a*c] \&& \text{IntegerQ}[n]$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.08 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.14

method	result	size
default	$-\frac{\sum_{R=\text{RootOf}(c_Z^8-a)} \frac{(-R^4 e+d) \ln(x-R)}{-R^7}}{8c}$	36
risch	$-\frac{\sum_{R=\text{RootOf}(c_Z^8-a)} \frac{(-R^4 e+d) \ln(x-R)}{-R^7}}{8c}$	36

input `int((e*x^4+d)/(-c*x^8+a),x,method=_RETURNVERBOSE)`

output
$$-1/8/c*\sum((_R^4*e+d)/_R^7*\ln(x-_R), _R=\text{RootOf}(_Z^8*c-a))$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2741 vs. $2(176) = 352$.

Time = 0.39 (sec) , antiderivative size = 2741, normalized size of antiderivative = 10.42

$$\int \frac{d + ex^4}{a - cx^8} dx = \text{Too large to display}$$

input `integrate((e*x^4+d)/(-c*x^8+a),x, algorithm="fricas")`

output
$$1/8*\sqrt{-\sqrt{((c^4*d^8 + 12*a*c^3*d^6*e^2 + 38*a^2*c^2*d^4*e^4 + 12*a^3*c^3*d^2*e^6 + a^4*e^8)/(a^7*c^5)) + 4*c*d^3*e + 4*a*d^2*e^3}/(a^3*c^2))*\log(-(c^3*d^6 + 5*a*c^2*d^4*e^2 - 5*a^2*c*d^2*e^4 - a^3*e^6)*x + (a^5*c^3*e*\sqrt{(c^4*d^8 + 12*a*c^3*d^6*e^2 + 38*a^2*c^2*d^4*e^4 + 12*a^3*c*d^2*e^6 + a^4*e^8)/(a^7*c^5)) - a*c^3*d^5 - 6*a^2*c^2*d^3*e^2 - a^3*c*d^4)*\sqrt{-\sqrt{((c^4*d^8 + 12*a*c^3*d^6*e^2 + 38*a^2*c^2*d^4*e^4 + 12*a^3*c*d^2*e^6 + a^4*e^8)/(a^7*c^5)) + 4*c*d^3*e + 4*a*d^2*e^3}/(a^3*c^2))} - 1/8*\sqrt{-\sqrt{((c^4*d^8 + 12*a*c^3*d^6*e^2 + 38*a^2*c^2*d^4*e^4 + 12*a^3*c*d^2*e^6 + a^4*e^8)/(a^7*c^5)) + 4*c*d^3*e + 4*a*d^2*e^3}/(a^3*c^2))*\log(-(c^3*d^6 + 5*a*c^2*d^4*e^2 - 5*a^2*c*d^2*e^4 - a^3*e^6)*x - (a^5*c^3*e*\sqrt{(c^4*d^8 + 12*a*c^3*d^6*e^2 + 38*a^2*c^2*d^4*e^4 + 12*a^3*c*d^2*e^6 + a^4*e^8)/(a^7*c^5)) - a*c^3*d^5 - 6*a^2*c^2*d^3*e^2 - a^3*c*d^4)*\sqrt{-\sqrt{((c^4*d^8 + 12*a*c^3*d^6*e^2 + 38*a^2*c^2*d^4*e^4 + 12*a^3*c*d^2*e^6 + a^4*e^8)/(a^7*c^5)) + 4*c*d^3*e + 4*a*d^2*e^3}/(a^3*c^2))} - 1/8*\sqrt{-\sqrt{(-(a^3*c^2*\sqrt{(c^4*d^8 + 12*a*c^3*d^6*e^2 + 38*a^2*c^2*d^4*e^4 + 12*a^3*c*d^2*e^6 + a^4*e^8)/(a^7*c^5)) - 4*c*d^3*e - 4*a*d^2*e^3}/(a^3*c^2))}\log(-(c^3*d^6 + 5*a*c^2*d^4*e^2 - 5*a^2*c*d^2*e^4 - a^3*e^6)*x + (a^5*c^3*e*\sqrt{(c^4*d^8 + 12*a*c^3*d^6*e^2 + 38*a^2*c^2*d^4*e^4 + 12*a^3*c*d^2*e^6 + a^4*e^8)/(a^7*c^5)) + a*c^3*d^5 + 6*a^2*c^2*d^3*e^2 + a^3*c*d^4)*\sqrt{-\sqrt{(-(a^3*c^2*\sqrt{(c^4*d^8 + 12*a*c^3*d^6*e^2 + 38*a^2*c^2*d^4*e^4 + 12*a^3*c*d^2*e^6 + a^4*e^8)/(a^7*c^5)) + 4*c*d^3*e + 4*a*d^2*e^3}/(a^3*c^2))}}$$

Sympy [F(-1)]

Timed out.

$$\int \frac{d + ex^4}{a - cx^8} dx = \text{Timed out}$$

input `integrate((e*x**4+d)/(-c*x**8+a),x)`

output `Timed out`

Maxima [F]

$$\int \frac{d + ex^4}{a - cx^8} dx = \int -\frac{ex^4 + d}{cx^8 - a} dx$$

input `integrate((e*x^4+d)/(-c*x^8+a),x, algorithm="maxima")`

output `-integrate((e*x^4 + d)/(c*x^8 - a), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 625 vs. $2(176) = 352$.

Time = 0.17 (sec) , antiderivative size = 625, normalized size of antiderivative = 2.38

$$\int \frac{d + ex^4}{a - cx^8} dx = \text{Too large to display}$$

input `integrate((e*x^4+d)/(-c*x^8+a),x, algorithm="giac")`

output

$$\begin{aligned}
 & -\frac{1}{8} \left(e \sqrt{-\sqrt{2} + 2} (-a/c)^{5/8} - d \sqrt{\sqrt{2} + 2} (-a/c)^{1/8} \right) \\
 & \times \arctan((2x + \sqrt{-\sqrt{2} + 2}) (-a/c)^{1/8}) / (\sqrt{\sqrt{2} + 2} (-a/c)^{1/8}) \\
 & / a - \frac{1}{8} \left(e \sqrt{-\sqrt{2} + 2} (-a/c)^{5/8} - d \sqrt{\sqrt{2} + 2} \right. \\
 & \left. (-a/c)^{1/8} \right) \times \arctan((2x - \sqrt{-\sqrt{2} + 2}) (-a/c)^{1/8}) / (\sqrt{\sqrt{2} + 2} \\
 & \times (-a/c)^{1/8}) / a + \frac{1}{8} \left(e \sqrt{\sqrt{2} + 2} (-a/c)^{5/8} + d \sqrt{-\sqrt{2} + 2} \right. \\
 & \left. (-a/c)^{1/8} \right) \times \arctan((2x + \sqrt{\sqrt{2} + 2}) (-a/c)^{1/8}) / (\sqrt{-\sqrt{2} + 2} \\
 & \times (-a/c)^{1/8}) / a + \frac{1}{8} \left(e \sqrt{\sqrt{2} + 2} (-a/c)^{5/8} + d \sqrt{-\sqrt{2} + 2} \right. \\
 & \left. (-a/c)^{1/8} \right) \times \arctan((2x - \sqrt{\sqrt{2} + 2}) (-a/c)^{1/8}) / (\sqrt{-\sqrt{2} + 2} \\
 & \times (-a/c)^{1/8}) / a - \frac{1}{16} \left(e \sqrt{-\sqrt{2} + 2} \right. \\
 & \left. (-a/c)^{5/8} - d \sqrt{\sqrt{2} + 2} (-a/c)^{1/8} \right) \times \log(x^2 + x \sqrt{\sqrt{2} + 2} \\
 & \times (-a/c)^{1/8} + (-a/c)^{1/4}) / a + \frac{1}{16} \left(e \sqrt{-\sqrt{2} + 2} \right. \\
 & \left. (-a/c)^{5/8} - d \sqrt{\sqrt{2} + 2} (-a/c)^{1/8} \right) \times \log(x^2 - x \sqrt{\sqrt{2} + 2} \\
 & \times (-a/c)^{1/8} + (-a/c)^{1/4}) / a + \frac{1}{16} \left(e \sqrt{\sqrt{2} + 2} (-a/c)^{5/8} + d \sqrt{-\sqrt{2} + 2} \right. \\
 & \left. (-a/c)^{1/8} \right) \times \log(x^2 + x \sqrt{-\sqrt{2} + 2} (-a/c)^{1/8} + (-a/c)^{1/4}) / a \\
 & - \frac{1}{16} \left(e \sqrt{\sqrt{2} + 2} (-a/c)^{5/8} + d \sqrt{-\sqrt{2} + 2} \right. \\
 & \left. (-a/c)^{1/8} \right) \times \log(x^2 - x \sqrt{-\sqrt{2} + 2} (-a/c)^{1/8} + (-a/c)^{1/4}) / a
 \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 11.46 (sec), antiderivative size = 2438, normalized size of antiderivative = 9.27

$$\int \frac{d + ex^4}{a - cx^8} dx = \text{Too large to display}$$

input `int((d + e*x^4)/(a - c*x^8),x)`

```

output
(atan((a^3*c^6*x + c^3*d^6*x - a*c^2*d^4*e^2*x - a^2*c*d^2*e^4*x + (2*d*e*x*(a^2*e^4*(a^7*c^5)^(1/2) + c^2*d^4*(a^7*c^5)^(1/2) + 4*a^4*c^4*d^3*e + 4*a^5*c^3*d*e^3 + 6*a*c*d^2*e^2*(a^7*c^5)^(1/2)))/(a*c^3*d^5*((a^2*e^4*(a^7*c^5)^(1/2) + c^2*d^4*(a^7*c^5)^(1/2) + 4*a^4*c^4*d^3*e + 4*a^5*c^3*d*e^3 + 6*a*c*d^2*e^2*(a^7*c^5)^(1/2))/(a^7*c^5))^(1/4) + a^5*c^3*e*((a^2*e^4*(a^7*c^5)^(1/2) + c^2*d^4*(a^7*c^5)^(1/2) + 4*a^4*c^4*d^3*e + 4*a^5*c^3*d*e^3 + 6*a*c*d^2*e^2*(a^7*c^5)^(1/2))/(a^7*c^5))^(5/4) + 2*a^2*c^2*d^3*e^2*((a^2*e^4*(a^7*c^5)^(1/2) + c^2*d^4*(a^7*c^5)^(1/2) + 4*a^4*c^4*d^3*e + 4*a^5*c^3*d*e^3 + 6*a*c*d^2*e^2*(a^7*c^5)^(1/2))/(a^7*c^5))^(1/4) - 3*a^3*c*d*e^4*((a^2*e^4*(a^7*c^5)^(1/2) + c^2*d^4*(a^7*c^5)^(1/2) + 4*a^4*c^4*d^3*e + 4*a^5*c^3*d*e^3 + 6*a*c*d^2*e^2*(a^7*c^5)^(1/2))/(a^7*c^5))^(1/4) - ((a^2*e^4*(a^7*c^5)^(1/2) + c^2*d^4*(a^7*c^5)^(1/2) + 4*a^4*c^4*d^3*e + 4*a^5*c^3*d*e^3 + 6*a*c*d^2*e^2*(a^7*c^5)^(1/2))/(a^7*c^5))^(1/4))/4 - (atan((a*c^2*d^4*e^2*x - c^3*d^6*x - a^3*e^6*x + a^2*c*d^2*e^4*x + (2*d*e*x*(a^2*e^4*(a^7*c^5)^(1/2) + c^2*d^4*(a^7*c^5)^(1/2) - 4*a^4*c^4*d^3*e - 4*a^5*c^3*d*e^3 + 6*a*c*d^2*e^2*(a^7*c^5)^(1/2)))/(a^3*c^2))/((a*c^3*d^5*(-(a^2*e^4*(a^7*c^5)^(1/2) + c^2*d^4*(a^7*c^5)^(1/2) - 4*a^4*c^4*d^3*e - 4*a^5*c^3*d*e^3 + 6*a*c*d^2*e^2*(a^7*c^5)^(1/2))/(a^7*c^5))^(1/4) + a^5*c^3*e*(-(a^2*e^4*(a^7*c^5)^(1/2) + c^2*d^4*(a^7*c^5)^(1/2) - 4*a^4*c^4*d^3*e - 4*a^5*c^3*d*e^3 + 6*a*c*d^2*e^2*(a^7*c^5)^(1/2))/(a^7*c^5))^(5/4) + 2*...

```

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 375, normalized size of antiderivative = 1.43

$$\int \frac{d + ex^4}{a - cx^8} dx = \frac{2\sqrt{a}\sqrt{2}\operatorname{atan}\left(\frac{c^{\frac{1}{8}}a^{\frac{1}{8}}\sqrt{2}-2c^{\frac{1}{4}}x}{c^{\frac{1}{8}}a^{\frac{1}{8}}\sqrt{2}}\right)e - 2\sqrt{c}\sqrt{2}\operatorname{atan}\left(\frac{c^{\frac{1}{8}}a^{\frac{1}{8}}\sqrt{2}-2c^{\frac{1}{4}}x}{c^{\frac{1}{8}}a^{\frac{1}{8}}\sqrt{2}}\right)d - 2\sqrt{a}\sqrt{2}\operatorname{atan}\left(\frac{c^{\frac{1}{8}}a^{\frac{1}{8}}\sqrt{2}+2c^{\frac{1}{4}}x}{c^{\frac{1}{8}}a^{\frac{1}{8}}\sqrt{2}}\right)e + 2\sqrt{c}\sqrt{2}\operatorname{atan}\left(\frac{c^{\frac{1}{8}}a^{\frac{1}{8}}\sqrt{2}+2c^{\frac{1}{4}}x}{c^{\frac{1}{8}}a^{\frac{1}{8}}\sqrt{2}}\right)d}{a^{\frac{1}{8}}(c^{\frac{1}{8}}a^{\frac{1}{8}})^{\frac{3}{2}}}$$

input `int((e*x^4+d)/(-c*x^8+a),x)`

output

```
(c**(3/8)*a**(1/8)*(2*sqrt(a)*sqrt(2)*atan((c**1/8)*a**1/8)*sqrt(2) - 2*c**1/4*x)/(c**1/8)*a**1/8)*sqrt(2))*e - 2*sqrt(c)*sqrt(2)*atan((c**1/8)*a**1/8)*sqrt(2) - 2*c**1/4*x)/(c**1/8)*a**1/8)*sqrt(2))*d - 2*sqrt(a)*sqrt(2)*atan((c**1/8)*a**1/8)*sqrt(2) + 2*c**1/4*x)/(c**1/8)*a**1/8)*sqrt(2))*e + 2*sqrt(c)*sqrt(2)*atan((c**1/8)*a**1/8)*sqrt(2) + 2*c**1/4*x)/(c**1/8)*a**1/8)*sqrt(2))*d + 4*sqrt(a)*atan((c**1/4*x)/(c**1/8)*a**1/8))*e + 4*sqrt(c)*atan((c**1/4*x)/(c**1/8)*a**1/8))*d + sqrt(a)*sqrt(2)*log(-c**1/8)*a**1/8)*sqrt(2)*x + a**1/4) + c**1/4)*x**2)*e - sqrt(a)*sqrt(2)*log(c**1/8)*a**1/8)*sqrt(2)*x + a**1/4) + c**1/4)*x**2)*e + 2*sqrt(a)*log(-c**1/8)*a**1/8 - c**1/4*x)*e - 2*sqrt(a)*log(c**1/8)*a**1/8 - c**1/4*x)*e - sqrt(c)*sqrt(2)*log(-c**1/8)*a**1/8)*sqrt(2)*x + a**1/4) + c**1/4)*x**2)*d + sqrt(c)*sqrt(2)*log(c**1/8)*a**1/8)*x + a**1/4) + c**1/4)*x**2)*d + 2*sqrt(c)*log(-c**1/8)*a**1/8 - c**1/4*x)*d - 2*sqrt(c)*log(c**1/8)*a**1/8 - c**1/4*x)*d))/(16*a*c)
```

3.15 $\int \frac{d+ex^4}{a+cx^8} dx$

Optimal result	152
Mathematica [A] (verified)	153
Rubi [A] (verified)	154
Maple [C] (verified)	160
Fricas [B] (verification not implemented)	161
Sympy [F(-1)]	162
Maxima [F]	162
Giac [A] (verification not implemented)	162
Mupad [B] (verification not implemented)	163
Reduce [B] (verification not implemented)	164

Optimal result

Integrand size = 17, antiderivative size = 578

$$\begin{aligned} \int \frac{d+ex^4}{a+cx^8} dx = & -\frac{\left(d + \sqrt{2}d - \frac{\sqrt{ae}}{\sqrt{c}}\right) \arctan\left(\frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a-2} \sqrt[8]{c_x}}{\sqrt{2+\sqrt{2}} \sqrt[8]{a}}\right)}{4\sqrt{2(2+\sqrt{2})} a^{7/8} \sqrt[8]{c}} \\ & + \frac{\left((1-\sqrt{2})\sqrt{cd} - \sqrt{ae}\right) \arctan\left(\frac{\sqrt{2+\sqrt{2}} \sqrt[8]{a-2} \sqrt[8]{c_x}}{\sqrt{2-\sqrt{2}} \sqrt[8]{a}}\right)}{4\sqrt{2(2-\sqrt{2})} a^{7/8} c^{5/8}} \\ & + \frac{\left(d + \sqrt{2}d - \frac{\sqrt{ae}}{\sqrt{c}}\right) \arctan\left(\frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a+2} \sqrt[8]{c_x}}{\sqrt{2+\sqrt{2}} \sqrt[8]{a}}\right)}{4\sqrt{2(2+\sqrt{2})} a^{7/8} \sqrt[8]{c}} \\ & - \frac{\left((1-\sqrt{2})\sqrt{cd} - \sqrt{ae}\right) \arctan\left(\frac{\sqrt{2+\sqrt{2}} \sqrt[8]{a+2} \sqrt[8]{c_x}}{\sqrt{2-\sqrt{2}} \sqrt[8]{a}}\right)}{4\sqrt{2(2-\sqrt{2})} a^{7/8} c^{5/8}} \\ & - \frac{\left((1-\sqrt{2})\sqrt{cd} - \sqrt{ae}\right) \operatorname{arctanh}\left(\frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a} \sqrt[8]{c_x}}{\sqrt[4]{a} + \sqrt[4]{c_x^2}}\right)}{4\sqrt{2(2-\sqrt{2})} a^{7/8} c^{5/8}} \\ & + \frac{\left(d + \sqrt{2}d - \frac{\sqrt{ae}}{\sqrt{c}}\right) \operatorname{arctanh}\left(\frac{\sqrt{2+\sqrt{2}} \sqrt[8]{a} \sqrt[8]{c_x}}{\sqrt[4]{a} + \sqrt[4]{c_x^2}}\right)}{4\sqrt{2(2+\sqrt{2})} a^{7/8} \sqrt[8]{c}} \end{aligned}$$

output

$$\begin{aligned}
 & -\frac{1}{4} \cdot (d+2^{(1/2)} \cdot d-a^{(1/2)} \cdot e/c^{(1/2)}) \cdot \arctan(((2-2^{(1/2)})^{(1/2)} \cdot a^{(1/8)} - 2 \cdot c^{(1/8)} \cdot x) / (2+2^{(1/2)})^{(1/2)} \cdot a^{(1/8)}) / (4+2 \cdot 2^{(1/2)})^{(1/2)} \cdot a^{(7/8)} / c^{(1/8)} + 1 \\
 & + \frac{4 \cdot ((1-2^{(1/2)}) \cdot c^{(1/2)} \cdot d-a^{(1/2)} \cdot e) \cdot \arctan(((2+2^{(1/2)})^{(1/2)} \cdot a^{(1/8)} - 2 \cdot c^{(1/8)} \cdot x) / (2-2^{(1/2)})^{(1/2)} \cdot a^{(1/8)}) / (4-2 \cdot 2^{(1/2)})^{(1/2)} \cdot a^{(7/8)} / c^{(5/8)} + 1 \\
 & + \frac{4 \cdot (d+2^{(1/2)} \cdot d-a^{(1/2)} \cdot e/c^{(1/2)}) \cdot \arctan(((2-2^{(1/2)})^{(1/2)} \cdot a^{(1/8)} + 2 \cdot c^{(1/8)} \cdot x) / (2+2^{(1/2)})^{(1/2)} \cdot a^{(1/8)}) / (4+2 \cdot 2^{(1/2)})^{(1/2)} \cdot a^{(7/8)} / c^{(1/8)} - 1/4 \\
 & * ((1-2^{(1/2)}) \cdot c^{(1/2)} \cdot d-a^{(1/2)} \cdot e) \cdot \arctan(((2+2^{(1/2)})^{(1/2)} \cdot a^{(1/8)} + 2 \cdot c^{(1/8)} \cdot x) / (2-2^{(1/2)})^{(1/2)} \cdot a^{(1/8)}) / (4-2 \cdot 2^{(1/2)})^{(1/2)} \cdot a^{(7/8)} / c^{(5/8)} - 1/4 \\
 & * ((1-2^{(1/2)}) \cdot c^{(1/2)} \cdot d-a^{(1/2)} \cdot e) \cdot \operatorname{arctanh}((2-2^{(1/2)})^{(1/2)} \cdot a^{(1/8)} \cdot c^{(1/8)} \cdot x / (a^{(1/4)} + c^{(1/4)} \cdot x^2)) / (4-2 \cdot 2^{(1/2)})^{(1/2)} \cdot a^{(7/8)} / c^{(5/8)} + 1/4 \\
 & * (d+2^{(1/2)} \cdot d-a^{(1/2)} \cdot e/c^{(1/2)}) \cdot \operatorname{arctanh}((2+2^{(1/2)})^{(1/2)} \cdot a^{(1/8)} \cdot c^{(1/8)} \cdot x / (a^{(1/4)} + c^{(1/4)} \cdot x^2)) / (4+2 \cdot 2^{(1/2)})^{(1/2)} \cdot a^{(7/8)} / c^{(1/8)}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.69 (sec), antiderivative size = 534, normalized size of antiderivative = 0.92

$$\begin{aligned}
 & \int \frac{d+ex^4}{a+cx^8} dx \\
 & = \frac{-2\sqrt[8]{a} \arctan \left(\cot \left(\frac{\pi}{8} \right) - \frac{\sqrt[8]{c_x \csc \left(\frac{\pi}{8} \right)}}{\sqrt[8]{a}} \right) (\sqrt{a}e \cos \left(\frac{\pi}{8} \right) + \sqrt{c}d \sin \left(\frac{\pi}{8} \right)) + 2\sqrt[8]{a} \arctan \left(\cot \left(\frac{\pi}{8} \right) + \frac{\sqrt[8]{c_x \csc \left(\frac{\pi}{8} \right)}}{\sqrt[8]{a}} \right) (\sqrt{a}e \cos \left(\frac{\pi}{8} \right) - \sqrt{c}d \sin \left(\frac{\pi}{8} \right))}{\sqrt[8]{a}}
 \end{aligned}$$

input

```
Integrate[(d + e*x^4)/(a + c*x^8), x]
```

output

$$\begin{aligned}
 & (-2 \cdot a^{(1/8)} \cdot \text{ArcTan}[\text{Cot}[\text{Pi}/8] - (c^{(1/8)} \cdot x \cdot \text{Csc}[\text{Pi}/8]) / a^{(1/8)}] \cdot (\text{Sqrt}[a] \cdot e \cdot \text{Cos}[\text{Pi}/8] + \text{Sqrt}[c] \cdot d \cdot \text{Sin}[\text{Pi}/8]) + 2 \cdot a^{(1/8)} \cdot \text{ArcTan}[\text{Cot}[\text{Pi}/8] + (c^{(1/8)} \cdot x \cdot \text{Csc}[\text{Pi}/8]) / a^{(1/8)}] \cdot (\text{Sqrt}[a] \cdot e \cdot \text{Cos}[\text{Pi}/8] + \text{Sqrt}[c] \cdot d \cdot \text{Sin}[\text{Pi}/8]) - a^{(1/8)} \cdot \text{Log}[a^{(1/4)} + c^{(1/4)} \cdot x^2 - 2 \cdot a^{(1/8)} \cdot c^{(1/8)} \cdot x \cdot \text{Sin}[\text{Pi}/8]] \cdot (\text{Sqrt}[a] \cdot e \cdot \text{Cos}[\text{Pi}/8] + \text{Sqrt}[c] \cdot d \cdot \text{Sin}[\text{Pi}/8]) + a^{(1/8)} \cdot \text{Log}[a^{(1/4)} + c^{(1/4)} \cdot x^2 + 2 \cdot a^{(1/8)} \cdot c^{(1/8)} \cdot x \cdot \text{Sin}[\text{Pi}/8]] \cdot (\text{Sqrt}[a] \cdot e \cdot \text{Cos}[\text{Pi}/8] + \text{Sqrt}[c] \cdot d \cdot \text{Sin}[\text{Pi}/8]) + a^{(1/8)} \cdot \text{Log}[a^{(1/4)} + c^{(1/4)} \cdot x^2 - 2 \cdot a^{(1/8)} \cdot c^{(1/8)} \cdot x \cdot \text{Cos}[\text{Pi}/8]] \cdot (-(\text{Sqrt}[c] \cdot d \cdot \text{Cos}[\text{Pi}/8]) + \text{Sqrt}[a] \cdot e \cdot \text{Sin}[\text{Pi}/8]) - a^{(1/8)} \cdot \text{Log}[a^{(1/4)} + c^{(1/4)} \cdot x^2 + 2 \cdot a^{(1/8)} \cdot c^{(1/8)} \cdot x \cdot \text{Cos}[\text{Pi}/8]] \cdot (-(\text{Sqrt}[c] \cdot d \cdot \text{Cos}[\text{Pi}/8]) + \text{Sqrt}[a] \cdot e \cdot \text{Sin}[\text{Pi}/8]) + 2 \cdot \text{ArcTan}[(c^{(1/8)} \cdot x \cdot \text{Sec}[\text{Pi}/8]) / a^{(1/8)} - \text{Tan}[\text{Pi}/8]] \cdot (a^{(1/8)} \cdot \text{Sqrt}[c] \cdot d \cdot \text{Cos}[\text{Pi}/8] - a^{(5/8)} \cdot e \cdot \text{Sin}[\text{Pi}/8]) + 2 \cdot \text{ArcTan}[(c^{(1/8)} \cdot x \cdot \text{Sec}[\text{Pi}/8]) / a^{(1/8)} + \text{Tan}[\text{Pi}/8]] \cdot (a^{(1/8)} \cdot \text{Sqrt}[c] \cdot d \cdot \text{Cos}[\text{Pi}/8] - a^{(5/8)} \cdot e \cdot \text{Sin}[\text{Pi}/8])) / (8 \cdot a \cdot c^{(5/8)})
 \end{aligned}$$

Rubi [A] (verified)

Time = 1.50 (sec) , antiderivative size = 849, normalized size of antiderivative = 1.47, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.647$, Rules used = {1745, 27, 1483, 27, 27, 1142, 25, 27, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{d + ex^4}{a + cx^8} dx \\
 & \downarrow \textcolor{blue}{1745} \\
 & \frac{\int \frac{\sqrt{2} \sqrt[4]{ad} - \sqrt[4]{c} \left(d - \frac{\sqrt{ae}}{\sqrt{c}}\right) x^2}{\sqrt[4]{c} \left(x^4 - \frac{\sqrt{2} \sqrt[4]{ax^2}}{\sqrt[4]{c}} + \frac{\sqrt{a}}{\sqrt{c}}\right)} dx}{2\sqrt{2}a^{3/4}\sqrt[4]{c}} + \frac{\int \frac{\sqrt[4]{c} \left(d - \frac{\sqrt{ae}}{\sqrt{c}}\right) x^2 + \sqrt{2} \sqrt[4]{ad}}{\sqrt[4]{c} \left(x^4 + \frac{\sqrt{2} \sqrt[4]{ax^2}}{\sqrt[4]{c}} + \frac{\sqrt{a}}{\sqrt{c}}\right)} dx}{2\sqrt{2}a^{3/4}\sqrt[4]{c}} \\
 & \downarrow \textcolor{blue}{27} \\
 & \frac{\int \frac{\sqrt{2} \sqrt[4]{ad} - \sqrt[4]{c} \left(d - \frac{\sqrt{ae}}{\sqrt{c}}\right) x^2}{x^4 - \frac{\sqrt{2} \sqrt[4]{ax^2}}{\sqrt[4]{c}} + \frac{\sqrt{a}}{\sqrt{c}}} dx}{2\sqrt{2}a^{3/4}\sqrt{c}} + \frac{\int \frac{\sqrt[4]{c} \left(d - \frac{\sqrt{ae}}{\sqrt{c}}\right) x^2 + \sqrt{2} \sqrt[4]{ad}}{x^4 + \frac{\sqrt{2} \sqrt[4]{ax^2}}{\sqrt[4]{c}} + \frac{\sqrt{a}}{\sqrt{c}}} dx}{2\sqrt{2}a^{3/4}\sqrt{c}} \\
 & \downarrow \textcolor{blue}{1483} \\
 & \frac{c^{3/8} \int \frac{\sqrt[4]{a} \left(\sqrt{2(2-\sqrt{2})} \sqrt[8]{a} d + \sqrt[8]{c} \left(-\sqrt{2}d + d - \frac{\sqrt{ae}}{\sqrt{c}}\right) x\right)}{\sqrt[8]{c} \left(x^2 - \frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a} x}{\sqrt[8]{c}} + \frac{\sqrt{a}}{\sqrt[4]{c}}\right)} dx}{2\sqrt{2-\sqrt{2}}a^{3/8}} + \frac{c^{3/8} \int \frac{\sqrt[4]{a} \left(\sqrt{2(2-\sqrt{2})} \sqrt[8]{a} d - \sqrt[8]{c} \left((1-\sqrt{2})d - \frac{\sqrt{ae}}{\sqrt{c}}\right) x\right)}{\sqrt[8]{c} \left(x^2 + \frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a} x}{\sqrt[8]{c}} + \frac{\sqrt{a}}{\sqrt[4]{c}}\right)} dx}{2\sqrt{2-\sqrt{2}}a^{3/8}} + \\
 & \frac{c^{3/8} \int \frac{\sqrt[4]{a} \left(\sqrt{2(2+\sqrt{2})} \sqrt[8]{a} d - \sqrt[8]{c} \left(\sqrt{2}d + d - \frac{\sqrt{ae}}{\sqrt{c}}\right) x\right)}{\sqrt[8]{c} \left(x^2 - \frac{\sqrt{2+\sqrt{2}} \sqrt[8]{a} x}{\sqrt[8]{c}} + \frac{\sqrt{a}}{\sqrt[4]{c}}\right)} dx}{2\sqrt{2+\sqrt{2}}a^{3/8}} + \frac{c^{3/8} \int \frac{\sqrt[4]{a} \left(\sqrt{2(2+\sqrt{2})} \sqrt[8]{a} d + \sqrt[8]{c} \left(\sqrt{2}d + d - \frac{\sqrt{ae}}{\sqrt{c}}\right) x\right)}{\sqrt[8]{c} \left(x^2 + \frac{\sqrt{2+\sqrt{2}} \sqrt[8]{a} x}{\sqrt[8]{c}} + \frac{\sqrt{a}}{\sqrt[4]{c}}\right)} dx}{2\sqrt{2+\sqrt{2}}a^{3/8}} \\
 & \downarrow \textcolor{blue}{27}
 \end{aligned}$$

$$\frac{\sqrt[4]{c} \int \frac{\sqrt{2(2-\sqrt{2})} \sqrt[8]{a} c^{3/8} d - ((1-\sqrt{2}) \sqrt{cd} - \sqrt{ae}) x}{x^2 + \frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a} x}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}}} dx}{2\sqrt{2-\sqrt{2}} \sqrt[8]{a}} + \frac{\sqrt[4]{c} \int \frac{\sqrt{2(2-\sqrt{2})} \sqrt[8]{a} d + \sqrt[8]{c}(-\sqrt{2d}+d-\frac{\sqrt{ae}}{\sqrt{c}}) x}{x^2 - \frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a} x}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}}} dx}{2\sqrt{2-\sqrt{2}} \sqrt[8]{a}} +$$

$$\frac{2\sqrt{2}a^{3/4}\sqrt{c}}{2\sqrt{2}a^{3/4}\sqrt{c}}$$

↓ 27

$$\frac{\int \frac{\sqrt{2(2-\sqrt{2})} \sqrt[8]{a} c^{3/8} d - ((1-\sqrt{2}) \sqrt{cd} - \sqrt{ae}) x}{x^2 + \frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a} x}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}}} dx}{2\sqrt{2-\sqrt{2}} \sqrt[8]{a} \sqrt[8]{c}} + \frac{\sqrt[4]{c} \int \frac{\sqrt{2(2-\sqrt{2})} \sqrt[8]{a} d + \sqrt[8]{c}(-\sqrt{2d}+d-\frac{\sqrt{ae}}{\sqrt{c}}) x}{x^2 - \frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a} x}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}}} dx}{2\sqrt{2-\sqrt{2}} \sqrt[8]{a}} +$$

$$\frac{2\sqrt{2}a^{3/4}\sqrt{c}}{2\sqrt{2}a^{3/4}\sqrt{c}}$$

↓ 1142

$$\frac{\sqrt[4]{c} \left(\frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a} ((1+\sqrt{2}) \sqrt{cd} - \sqrt{ae}) \int \frac{1}{x^2 - \frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a} x}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}}} dx}{2\sqrt{c}} + \frac{1}{2} \sqrt[8]{c} (-\sqrt{2d}+d-\frac{\sqrt{ae}}{\sqrt{c}}) \int -\frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a} -2 \sqrt[8]{c} x}{\sqrt[8]{c} \left(x^2 - \frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a} x}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}} \right)} dx \right)}{2\sqrt{2-\sqrt{2}} \sqrt[8]{a}} +$$

$$\frac{2\sqrt{2}a^{3/4}\sqrt{c}}{2\sqrt{2}a^{3/4}\sqrt{c}}$$

$$\frac{\sqrt[4]{c} \left(-\frac{\sqrt{2+\sqrt{2}} \sqrt[8]{a} ((1-\sqrt{2}) \sqrt{cd} - \sqrt{ae}) \int \frac{1}{x^2 - \frac{\sqrt{2+\sqrt{2}} \sqrt[8]{a} x}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}}} dx}{2\sqrt{c}} - \frac{1}{2} \sqrt[8]{c} (\sqrt{2d}+d-\frac{\sqrt{ae}}{\sqrt{c}}) \int -\frac{\sqrt{2+\sqrt{2}} \sqrt[8]{a} -2 \sqrt[8]{c} x}{\sqrt[8]{c} \left(x^2 - \frac{\sqrt{2+\sqrt{2}} \sqrt[8]{a} x}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}} \right)} dx \right)}{2\sqrt{2+\sqrt{2}} \sqrt[8]{a}} +$$

$$\frac{2\sqrt{2}a^{3/4}\sqrt{c}}{2\sqrt{2}a^{3/4}\sqrt{c}}$$

↓ 25

$$\begin{aligned}
& \frac{\sqrt[4]{c}}{2\sqrt{c}} \left(\frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a} ((1+\sqrt{2})\sqrt{c}d - \sqrt{a}e) \int \frac{1}{x^2 - \frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a}x}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}}} dx}{\sqrt[8]{c} \left(x^2 - \frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a}x}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}} \right)} - \frac{1}{2} \sqrt[8]{c} \left(-\sqrt{2}d + d - \frac{\sqrt{a}e}{\sqrt{c}} \right) \int \frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a} \sqrt[8]{c}x}{\sqrt[8]{c} \left(x^2 - \frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a}x}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}} \right)} dx \right) \right) \\
& + \frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a} ((1-\sqrt{2})\sqrt{c}d - \sqrt{a}e) \int \frac{1}{x^2 - \frac{\sqrt{2+\sqrt{2}} \sqrt[8]{a}x}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}}} dx}{2\sqrt{2-\sqrt{2}} \sqrt[8]{a} \sqrt[8]{c}} \\
& + \frac{\sqrt[4]{c}}{2\sqrt{c}} \left(\frac{\frac{1}{2} \sqrt[8]{c} \left(\sqrt{2}d + d - \frac{\sqrt{a}e}{\sqrt{c}} \right) \int \frac{\sqrt{2+\sqrt{2}} \sqrt[8]{a} \sqrt[8]{c}x}{\sqrt[8]{c} \left(x^2 - \frac{\sqrt{2+\sqrt{2}} \sqrt[8]{a}x}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}} \right)} dx}{2\sqrt{2+\sqrt{2}} \sqrt[8]{a} \sqrt[8]{c}} \right) \right) \\
& + \frac{2\sqrt{2}a^{3/4}\sqrt{c}}{2\sqrt{2}a^{3/4}\sqrt{c}}
\end{aligned}$$

↓ 27

$$\begin{aligned}
& \frac{\sqrt[4]{c}}{2\sqrt{c}} \left(\frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a} ((1+\sqrt{2})\sqrt{c}d - \sqrt{a}e) \int \frac{1}{x^2 - \frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a}x}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}}} dx}{\sqrt[8]{c} \left(x^2 - \frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a}x}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}} \right)} - \frac{1}{2} \left(-\sqrt{2}d + d - \frac{\sqrt{a}e}{\sqrt{c}} \right) \int \frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a} \sqrt[8]{c}x}{\sqrt[8]{c} \left(x^2 - \frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a}x}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}} \right)} dx \right) \right) \\
& + \frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a} ((1-\sqrt{2})\sqrt{c}d - \sqrt{a}e) \int \frac{1}{x^2 - \frac{\sqrt{2+\sqrt{2}} \sqrt[8]{a}x}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}}} dx}{2\sqrt{2-\sqrt{2}} \sqrt[8]{a} \sqrt[8]{c}} \\
& + \frac{\sqrt[4]{c}}{2\sqrt{c}} \left(\frac{\frac{1}{2} \left(\sqrt{2}d + d - \frac{\sqrt{a}e}{\sqrt{c}} \right) \int \frac{\sqrt{2+\sqrt{2}} \sqrt[8]{a} \sqrt[8]{c}x}{\sqrt[8]{c} \left(x^2 - \frac{\sqrt{2+\sqrt{2}} \sqrt[8]{a}x}{\sqrt[8]{c}} + \frac{\sqrt[4]{a}}{\sqrt[4]{c}} \right)} dx}{2\sqrt{2+\sqrt{2}} \sqrt[8]{a} \sqrt[8]{c}} \right) \right) \\
& + \frac{2\sqrt{2}a^{3/4}\sqrt{c}}{2\sqrt{2}a^{3/4}\sqrt{c}}
\end{aligned}$$

↓ 1083

$$\frac{4\sqrt{c}}{2\sqrt{2-\sqrt{2}}\sqrt[8]{a}} \left(\begin{array}{l} \int \frac{\sqrt{2-\sqrt{2}}\sqrt[8]{a}((1+\sqrt{2})\sqrt{cd}-\sqrt{ae})}{-\left(2x-\frac{\sqrt{2-\sqrt{2}}\sqrt[8]{a}}{\sqrt[8]{c}}\right)^2-\frac{(2+\sqrt{2})\sqrt[4]{a}}{\sqrt[4]{c}}} dx \\ -\frac{1}{2}\left(-\sqrt{2}d+d-\frac{\sqrt{ae}}{\sqrt{c}}\right) \int \frac{\sqrt{2-\sqrt{2}}\sqrt[8]{a}-2\sqrt[8]{c_x}}{x^2-\frac{\sqrt{2-\sqrt{2}}\sqrt[8]{a_x}+\frac{4}{\sqrt[4]{c}}}{\sqrt[8]{c}}} dx \end{array} \right)$$

$$\frac{4\sqrt{c}}{2\sqrt{2+\sqrt{2}}\sqrt[8]{a}} \left(\begin{array}{l} \int \frac{\sqrt{2+\sqrt{2}}\sqrt[8]{a}((1-\sqrt{2})\sqrt{cd}-\sqrt{ae})}{-\left(2x-\frac{\sqrt{2+\sqrt{2}}\sqrt[8]{a}}{\sqrt[8]{c}}\right)^2-\frac{(2-\sqrt{2})\sqrt[4]{a}}{\sqrt[4]{c}}} dx \\ +\frac{1}{2}\left(\sqrt{2}d+d-\frac{\sqrt{ae}}{\sqrt{c}}\right) \int \frac{\sqrt{2+\sqrt{2}}\sqrt[8]{a}-2\sqrt[8]{c_x}}{x^2-\frac{\sqrt{2+\sqrt{2}}\sqrt[8]{a_x}+\frac{4}{\sqrt[4]{c}}}{\sqrt[8]{c}}} dx \end{array} \right)$$

↓ 217

$$\frac{4\sqrt{c}}{2\sqrt{2-\sqrt{2}}\sqrt[8]{a}} \left(\begin{array}{l} \frac{\sqrt{\frac{2-\sqrt{2}}{2+\sqrt{2}}}((1+\sqrt{2})\sqrt{cd}-\sqrt{ae}) \arctan \left(\frac{\frac{8}{\sqrt{c}}\left(2x-\frac{\sqrt{2-\sqrt{2}}\sqrt[8]{a}}{\sqrt[8]{c}}\right)}{\sqrt{2+\sqrt{2}}\sqrt[8]{a}} \right)}{c^{3/8}} \\ -\frac{1}{2}\left(-\sqrt{2}d+d-\frac{\sqrt{ae}}{\sqrt{c}}\right) \int \frac{\sqrt{2-\sqrt{2}}\sqrt[8]{a}-2\sqrt[8]{c_x}}{x^2-\frac{\sqrt{2-\sqrt{2}}\sqrt[8]{a_x}+\frac{4}{\sqrt[4]{c}}}{\sqrt[8]{c}}} dx \end{array} \right) + \frac{\sqrt{\frac{2-\sqrt{2}}{2+\sqrt{2}}}((1+\sqrt{2})\sqrt{cd}-\sqrt{ae}) \arctan \left(\frac{\frac{8}{\sqrt{c}}\left(2x-\frac{\sqrt{2+\sqrt{2}}\sqrt[8]{a}}{\sqrt[8]{c}}\right)}{\sqrt{2-\sqrt{2}}\sqrt[8]{a}} \right)}{c^{3/8}}$$

$$\frac{4\sqrt{c}}{2\sqrt{2+\sqrt{2}}\sqrt[8]{a}} \left(\begin{array}{l} \frac{\sqrt{\frac{2+\sqrt{2}}{2-\sqrt{2}}}((1-\sqrt{2})\sqrt{cd}-\sqrt{ae}) \arctan \left(\frac{\frac{8}{\sqrt{c}}\left(2x-\frac{\sqrt{2+\sqrt{2}}\sqrt[8]{a}}{\sqrt[8]{c}}\right)}{\sqrt{2-\sqrt{2}}\sqrt[8]{a}} \right)}{c^{3/8}} \\ +\frac{1}{2}\left(\sqrt{2}d+d-\frac{\sqrt{ae}}{\sqrt{c}}\right) \int \frac{\sqrt{2+\sqrt{2}}\sqrt[8]{a}-2\sqrt[8]{c_x}}{x^2-\frac{\sqrt{2+\sqrt{2}}\sqrt[8]{a_x}+\frac{4}{\sqrt[4]{c}}}{\sqrt[8]{c}}} dx \end{array} \right) + \frac{2\sqrt{2}a^{3/4}\sqrt{c}}{2\sqrt{2}a^{3/4}\sqrt{c}}$$

↓ 1103

$$\begin{aligned}
& \frac{\sqrt[4]{c} \left(\frac{\sqrt[8]{c} \left(2x - \frac{\sqrt{2-\sqrt{2}} \sqrt[8]{a}}{\sqrt[8]{c}} \right)}{\sqrt{2+\sqrt{2}} \sqrt[8]{a}} + \frac{1}{2} \sqrt[8]{c} \left(-\sqrt{2}d + d - \frac{\sqrt{a}e}{\sqrt[8]{c}} \right) \log \left(\sqrt[4]{c}x^2 - \sqrt{2-\sqrt{2}} \sqrt[8]{a} \sqrt[8]{c}x + \sqrt[4]{a} \right) \right)}{2\sqrt{2-\sqrt{2}} \sqrt[8]{a}} + \frac{\sqrt[2]{\frac{2}{2-\sqrt{2}}} \sqrt[4]{c}}{2\sqrt{2}a^{3/4}\sqrt{c}} \\
& - \frac{\sqrt[4]{c} \left(\frac{\sqrt[8]{c} \left(2x - \frac{\sqrt{2+\sqrt{2}} \sqrt[8]{a}}{\sqrt[8]{c}} \right)}{\sqrt{2-\sqrt{2}} \sqrt[8]{a}} - \frac{1}{2} \sqrt[8]{c} \left(\sqrt{2}d + d - \frac{\sqrt{a}e}{\sqrt[8]{c}} \right) \log \left(\sqrt[4]{c}x^2 - \sqrt{2+\sqrt{2}} \sqrt[8]{a} \sqrt[8]{c}x + \sqrt[4]{a} \right) \right)}{2\sqrt{2+\sqrt{2}} \sqrt[8]{a}} + \frac{\sqrt[2]{\frac{2}{2+\sqrt{2}}} \sqrt[4]{c}}{2\sqrt{2}a^{3/4}\sqrt{c}}
\end{aligned}$$

input `Int[(d + e*x^4)/(a + c*x^8), x]`

output

$$\begin{aligned}
 & ((c^{1/4}) * ((\text{Sqrt}[(2 - \text{Sqrt}[2])/(2 + \text{Sqrt}[2])] * ((1 + \text{Sqrt}[2]) * \text{Sqrt}[c] * d - \text{Sqr}\\
 & \text{t}[a] * e) * \text{ArcTan}[(c^{1/8}) * (-((\text{Sqrt}[2 - \text{Sqrt}[2]] * a^{1/8})/c^{1/8}) + 2*x)) / \\
 & (\text{Sqrt}[2 + \text{Sqrt}[2]] * a^{1/8}))]) / c^{3/8} + (c^{1/8}) * (d - \text{Sqrt}[2] * d - (\text{Sqrt}[a] \\
 & * e) / \text{Sqrt}[c]) * \text{Log}[a^{1/4} - \text{Sqrt}[2 - \text{Sqrt}[2]] * a^{1/8} * c^{1/8} * x + c^{1/4} * x \\
 & ^2) / 2)) / (2 * \text{Sqrt}[2 - \text{Sqrt}[2]] * a^{1/8}) + (\text{Sqrt}[(2 - \text{Sqrt}[2]) / (2 + \text{Sqrt}[2])] * \\
 & ((1 + \text{Sqrt}[2]) * \text{Sqrt}[c] * d - \text{Sqrt}[a] * e) * \text{ArcTan}[(c^{1/8}) * ((\text{Sqrt}[2 - \text{Sqrt}[2]] * \\
 & a^{1/8}) / c^{1/8} + 2*x)) / (\text{Sqrt}[2 + \text{Sqrt}[2]] * a^{1/8})] - (((1 - \text{Sqrt}[2]) * \\
 & \text{Sqrt}[c] * d - \text{Sqrt}[a] * e) * \text{Log}[a^{1/4} + \text{Sqrt}[2 - \text{Sqrt}[2]] * a^{1/8} * c^{1/8} * x + \\
 & c^{1/4} * x^2) / 2) / (2 * \text{Sqrt}[2 - \text{Sqrt}[2]] * a^{1/8} * c^{1/8})) / (2 * \text{Sqrt}[2] * a^{3/4} \\
 & * \text{Sqrt}[c]) + ((c^{1/4}) * (-((\text{Sqrt}[(2 + \text{Sqrt}[2]) / (2 - \text{Sqrt}[2])] * ((1 - \text{Sqrt}[2]) * \\
 & \text{Sqrt}[c] * d - \text{Sqrt}[a] * e) * \text{ArcTan}[(c^{1/8}) * (-((\text{Sqrt}[2 + \text{Sqrt}[2]] * a^{1/8}) / c^{1/8}) \\
 & + 2*x)) / (\text{Sqrt}[2 - \text{Sqrt}[2]] * a^{1/8}))]) / c^{3/8}) - (c^{1/8}) * (d + \text{Sqrt}[2] * d - \\
 & (\text{Sqrt}[a] * e) / \text{Sqrt}[c]) * \text{Log}[a^{1/4} - \text{Sqrt}[2 + \text{Sqrt}[2]] * a^{1/8} * c^{1/8} * x \\
 & + c^{1/4} * x^2) / 2) / (2 * \text{Sqrt}[2 + \text{Sqrt}[2]] * a^{1/8}) + (c^{1/4}) * (-((\text{Sqrt}[(2 + \text{Sqrt}[2]) / (2 - \text{Sqrt}[2])] * \\
 & ((1 - \text{Sqrt}[2]) * \text{Sqrt}[c] * d - \text{Sqrt}[a] * e) * \text{ArcTan}[(c^{1/8}) * ((\text{Sqrt}[2 + \text{Sqrt}[2]] * a^{1/8}) / c^{1/8}) \\
 & + 2*x)) / (\text{Sqrt}[2 - \text{Sqrt}[2]] * a^{1/8})) / c^{3/8}) + (c^{1/8}) * (d + \text{Sqrt}[2] * d - \\
 & (\text{Sqrt}[a] * e) / \text{Sqrt}[c]) * \text{Log}[a^{1/4} + \text{Sqrt}[2 + \text{Sqrt}[2]] * a^{1/8} * c^{1/8} * x + \\
 & c^{1/4} * x^2) / 2) / (2 * \text{Sqrt}[2 + \text{Sqrt}[2]] * a^{1/8})) / (2 * \text{Sqrt}[2] * a^{3/4} * \text{Sqrt}[c])
 \end{aligned}$$

Definitions of rubi rules used

rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$

rule 27 $\text{Int}[(\text{a}_.) * (\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] /; \text{FreeQ}[\text{a}, \text{x}] \&& \text{!Ma}\\ \text{tchQ}[\text{Fx}, (\text{b}_.) * (\text{Gx}_)] /; \text{FreeQ}[\text{b}, \text{x}]]$

rule 217 $\text{Int}[(\text{a}_.) + (\text{b}_.) * (\text{x}_.)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(-(\text{Rt}[-\text{a}, 2] * \text{Rt}[-\text{b}, 2]))^{-1} \\
 - 1) * \text{ArcTan}[\text{Rt}[-\text{b}, 2] * (\text{x}/\text{Rt}[-\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&& \text{PosQ}[\text{a}/\text{b}] \& \\
 \& (\text{LtQ}[\text{a}, 0] \mid\mid \text{LtQ}[\text{b}, 0])]$

rule 1083 $\text{Int}[(\text{a}_.) + (\text{b}_.) * (\text{x}_.) + (\text{c}_.) * (\text{x}_.)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[-2 \quad \text{Subst}[\text{I}\\ \text{nt}[1/\text{Simp}[\text{b}^2 - 4*\text{a}*\text{c} - \text{x}^2, \text{x}], \text{x}], \text{x}, \text{b} + 2*\text{c}*\text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}\}, \text{x}]$

rule 1103 $\text{Int}[(d_.) + (e_.)*(x_.) / ((a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2), x_{\text{Symbol}}] \Rightarrow S$
 $\text{imp}[d * \text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{EqQ}[2*c*d - b*e, 0]$

rule 1142 $\text{Int}[(d_.) + (e_.)*(x_.) / ((a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2), x_{\text{Symbol}}] \Rightarrow S$
 $\text{imp}[(2*c*d - b*e)/(2*c) \quad \text{Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c)$
 $\text{Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

rule 1483 $\text{Int}[(d_.) + (e_.)*(x_.)^2 / ((a_.) + (b_.)*(x_.)^2 + (c_.)*(x_.)^4), x_{\text{Symbol}}] :$
 $> \text{With}[\{q = \text{Rt}[a/c, 2]\}, \text{With}[\{r = \text{Rt}[2*q - b/c, 2]\}, \text{Simp}[1/(2*c*q*r) \quad \text{In}$
 $t[(d*r - (d - e*q)*x)/(q - r*x + x^2), x], x] + \text{Simp}[1/(2*c*q*r) \quad \text{Int}[(d*r$
 $+ (d - e*q)*x)/(q + r*x + x^2), x], x]] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{N}$
 $\text{eQ}[b^2 - 4*a*c, 0] \&& \text{NeQ}[c*d^2 - b*d*e + a*e^2, 0] \&& \text{NegQ}[b^2 - 4*a*c]$

rule 1745 $\text{Int}[(d_.) + (e_.)*(x_.)^{(n_.)} / ((a_.) + (c_.)*(x_.)^{(n2_.)}), x_{\text{Symbol}}] \Rightarrow \text{With}[\{$
 $q = \text{Rt}[a/c, 4]\}, \text{Simp}[1/(2*\text{Sqrt}[2]*c*q^3) \quad \text{Int}[(\text{Sqrt}[2]*d*q - (d - e*q^2)*$
 $x^{(n/2)})/(q^2 - \text{Sqrt}[2]*q*x^{(n/2)} + x^n), x], x] + \text{Simp}[1/(2*\text{Sqrt}[2]*c*q^3)$
 $\text{Int}[(\text{Sqrt}[2]*d*q + (d - e*q^2)*x^{(n/2)})/(q^2 + \text{Sqrt}[2]*q*x^{(n/2)} + x^n),$
 $x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \&& \text{EqQ}[n2, 2*n] \&& \text{NeQ}[c*d^2 + a*e^2, 0]$
 $\&& \text{NeQ}[c*d^2 - a*e^2, 0] \&& \text{IGtQ}[n/2, 0] \&& \text{PosQ}[a*c]$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.09 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.06

method	result	size
default	$\frac{\sum_{R=\text{RootOf}(c_Z^8+a)} \frac{(-R^{4/e+d}) \ln(x-R)}{-R^7}}{8c}$	34
risch	$\frac{\sum_{R=\text{RootOf}(c_Z^8+a)} \frac{(-R^{4/e+d}) \ln(x-R)}{-R^7}}{8c}$	34

input `int((e*x^4+d)/(c*x^8+a),x,method=_RETURNVERBOSE)`

output `1/8/c*sum(_R^4*e+d)/_R^7*ln(x-_R),_R=RootOf(_Z^8*c+a))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2749 vs. $2(391) = 782$.

Time = 0.37 (sec) , antiderivative size = 2749, normalized size of antiderivative = 4.76

$$\int \frac{d + ex^4}{a + cx^8} dx = \text{Too large to display}$$

input `integrate((e*x^4+d)/(c*x^8+a),x, algorithm="fricas")`

output `1/8*sqrt(-sqrt(-(a^3*c^2*sqrt(-(c^4*d^8 - 12*a*c^3*d^6*e^2 + 38*a^2*c^2*d^4*e^4 - 12*a^3*c*d^2*e^6 + a^4*e^8)/(a^7*c^5)) + 4*c*d^3*e - 4*a*d*e^3)/(a^3*c^2))*log((c^3*d^6 - 5*a*c^2*d^4*e^2 - 5*a^2*c*d^2*e^4 + a^3*e^6)*x + (a^5*c^3*e*sqrt(-(c^4*d^8 - 12*a*c^3*d^6*e^2 + 38*a^2*c^2*d^4*e^4 - 12*a^3*c*d^2*e^6 + a^4*e^8)/(a^7*c^5)) + a*c^3*d^5 - 6*a^2*c^2*d^3*e^2 + a^3*c*d^4*e^4)*sqrt(-sqrt(-(a^3*c^2*sqrt(-(c^4*d^8 - 12*a*c^3*d^6*e^2 + 38*a^2*c^2*d^4*e^4 - 12*a^3*c*d^2*e^6 + a^4*e^8)/(a^7*c^5)) + 4*c*d^3*e - 4*a*d*e^3)/(a^3*c^2))) - 1/8*sqrt(-sqrt(-(a^3*c^2*sqrt(-(c^4*d^8 - 12*a*c^3*d^6*e^2 + 38*a^2*c^2*d^4*e^4 - 12*a^3*c*d^2*e^6 + a^4*e^8)/(a^7*c^5)) + 4*c*d^3*e - 4*a*d*e^3)/(a^3*c^2))*log((c^3*d^6 - 5*a*c^2*d^4*e^2 - 5*a^2*c*d^2*e^4 + a^3*e^6)*x - (a^5*c^3*e*sqrt(-(c^4*d^8 - 12*a*c^3*d^6*e^2 + 38*a^2*c^2*d^4*e^4 - 12*a^3*c*d^2*e^6 + a^4*e^8)/(a^7*c^5)) + a*c^3*d^5 - 6*a^2*c^2*d^3*e^2 + a^3*c*d^4*e^4)*sqrt(-sqrt(-(a^3*c^2*sqrt(-(c^4*d^8 - 12*a*c^3*d^6*e^2 + 38*a^2*c^2*d^4*e^4 - 12*a^3*c*d^2*e^6 + a^4*e^8)/(a^7*c^5)) + 4*c*d^3*e - 4*a*d*e^3)/(a^3*c^2))) - 1/8*sqrt(-sqrt((a^3*c^2*sqrt(-(c^4*d^8 - 12*a*c^3*d^6*e^2 + 38*a^2*c^2*d^4*e^4 - 12*a^3*c*d^2*e^6 + a^4*e^8)/(a^7*c^5)) + 4*c*d^3*e + 4*a*d*e^3)/(a^3*c^2))*log((c^3*d^6 - 5*a*c^2*d^4*e^2 - 5*a^2*c*d^2*e^4 + a^3*e^6)*x + (a^5*c^3*e*sqrt(-(c^4*d^8 - 12*a*c^3*d^6*e^2 + 38*a^2*c^2*d^4*e^4 - 12*a^3*c*d^2*e^6 + a^4*e^8)/(a^7*c^5)) - a*c^3*d^5 + 6*a^2*c^2*d^3*e^2 - a^3*c*d^4*e^4)*sqrt(-sqrt((a^3*c^2*sqrt(-(c^4*d^8 - ...`

Sympy [F(-1)]

Timed out.

$$\int \frac{d + ex^4}{a + cx^8} dx = \text{Timed out}$$

input `integrate((e*x**4+d)/(c*x**8+a),x)`

output `Timed out`

Maxima [F]

$$\int \frac{d + ex^4}{a + cx^8} dx = \int \frac{ex^4 + d}{cx^8 + a} dx$$

input `integrate((e*x^4+d)/(c*x^8+a),x, algorithm="maxima")`

output `integrate((e*x^4 + d)/(c*x^8 + a), x)`

Giac [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 593, normalized size of antiderivative = 1.03

$$\int \frac{d + ex^4}{a + cx^8} dx = \text{Too large to display}$$

input `integrate((e*x^4+d)/(c*x^8+a),x, algorithm="giac")`

output

$$\begin{aligned}
 & -\frac{1}{8} \left(e \sqrt{-\sqrt{2} + 2} \right) \left(\frac{a}{c} \right)^{5/8} - d \sqrt{\sqrt{2} + 2} \left(\frac{a}{c} \right)^{1/8} \arctan \left(\frac{2x + \sqrt{-\sqrt{2} + 2} \left(\frac{a}{c} \right)^{1/8}}{\sqrt{\sqrt{2} + 2} \left(\frac{a}{c} \right)^{1/8}} \right) / a \\
 & - \frac{1}{8} \left(e \sqrt{-\sqrt{2} + 2} \right) \left(\frac{a}{c} \right)^{5/8} - d \sqrt{\sqrt{2} + 2} \left(\frac{a}{c} \right)^{1/8} \arctan \left(\frac{(2x - \sqrt{-\sqrt{2} + 2} \left(\frac{a}{c} \right)^{1/8})}{\sqrt{\sqrt{2} + 2} \left(\frac{a}{c} \right)^{1/8}} \right) / a \\
 & + \frac{1}{8} \left(e \sqrt{\sqrt{2} + 2} \right) \left(\frac{a}{c} \right)^{5/8} + d \sqrt{-\sqrt{2} + 2} \left(\frac{a}{c} \right)^{1/8} \arctan \left(\frac{(2x + \sqrt{\sqrt{2} + 2} \left(\frac{a}{c} \right)^{1/8})}{\sqrt{-\sqrt{2} + 2} \left(\frac{a}{c} \right)^{1/8}} \right) / a \\
 & + \frac{1}{8} \left(e \sqrt{\sqrt{2} + 2} \right) \left(\frac{a}{c} \right)^{5/8} + d \sqrt{-\sqrt{2} + 2} \left(\frac{a}{c} \right)^{1/8} \arctan \left(\frac{(2x - \sqrt{\sqrt{2} + 2} \left(\frac{a}{c} \right)^{1/8})}{\sqrt{-\sqrt{2} + 2} \left(\frac{a}{c} \right)^{1/8}} \right) / a \\
 & - \frac{1}{16} \left(e \sqrt{-\sqrt{2} + 2} \right) \left(\frac{a}{c} \right)^{5/8} - d \sqrt{\sqrt{2} + 2} \left(\frac{a}{c} \right)^{1/8} \log(x^2 + x \sqrt{\sqrt{2} + 2} \left(\frac{a}{c} \right)^{1/8} + \left(\frac{a}{c} \right)^{1/4}) / a \\
 & + \frac{1}{16} \left(e \sqrt{-\sqrt{2} + 2} \right) \left(\frac{a}{c} \right)^{5/8} - d \sqrt{\sqrt{2} + 2} \left(\frac{a}{c} \right)^{1/8} \log(x^2 - x \sqrt{\sqrt{2} + 2} \left(\frac{a}{c} \right)^{1/8} + \left(\frac{a}{c} \right)^{1/4}) / a \\
 & + \frac{1}{16} \left(e \sqrt{\sqrt{2} + 2} \right) \left(\frac{a}{c} \right)^{5/8} + d \sqrt{-\sqrt{2} + 2} \left(\frac{a}{c} \right)^{1/8} \log(x^2 + x \sqrt{-\sqrt{2} + 2} \left(\frac{a}{c} \right)^{1/8} + \left(\frac{a}{c} \right)^{1/4}) / a \\
 & - \frac{1}{16} \left(e \sqrt{\sqrt{2} + 2} \right) \left(\frac{a}{c} \right)^{5/8} + d \sqrt{-\sqrt{2} + 2} \left(\frac{a}{c} \right)^{1/8} \log(x^2 - x \sqrt{-\sqrt{2} + 2} \left(\frac{a}{c} \right)^{1/8} + \left(\frac{a}{c} \right)^{1/4}) / a
 \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 11.78 (sec) , antiderivative size = 2510, normalized size of antiderivative = 4.34

$$\int \frac{d + ex^4}{a + cx^8} dx = \text{Too large to display}$$

input `int((d + e*x^4)/(a + c*x^8),x)`

output

$$\begin{aligned}
 & (\text{atan}((c^3*d^6*x - a^3*e^6*x + a*c^2*d^4*e^2*x - a^2*c*d^2*e^4*x + (2*d*e*x*(a^2*e^4*(-a^7*c^5)^(1/2) + c^2*d^4*(-a^7*c^5)^(1/2) - 4*a^4*c^4*d^3*e + 4*a^5*c^3*d*e^3 - 6*a*c*d^2*e^2*(-a^7*c^5)^(1/2)))/(a^3*c^2))/(a*c^3*d^5*((a^2*e^4*(-a^7*c^5)^(1/2) + c^2*d^4*(-a^7*c^5)^(1/2) - 4*a^4*c^4*d^3*e + 4*a^5*c^3*d*e^3 - 6*a*c*d^2*e^2*(-a^7*c^5)^(1/2))/(a^7*c^5))^(1/4) + a^5*c^3*e*((a^2*e^4*(-a^7*c^5)^(1/2) + c^2*d^4*(-a^7*c^5)^(1/2) - 4*a^4*c^4*d^3*e + 4*a^5*c^3*d*e^3 - 6*a*c*d^2*e^2*(-a^7*c^5)^(1/2))/(a^7*c^5))^(5/4) - 2*a^2*c^2*d^3*e^2*((a^2*e^4*(-a^7*c^5)^(1/2) + c^2*d^4*(-a^7*c^5)^(1/2) - 4*a^4*c^4*d^3*e + 4*a^5*c^3*d*e^3 - 6*a*c*d^2*e^2*(-a^7*c^5)^(1/2))/(a^7*c^5))^(1/4) - 3*a^3*c*d*e^4*((a^2*e^4*(-a^7*c^5)^(1/2) + c^2*d^4*(-a^7*c^5)^(1/2) - 4*a^4*c^4*d^3*e + 4*a^5*c^3*d*e^3 - 6*a*c*d^2*e^2*(-a^7*c^5)^(1/2))/(a^7*c^5))^(1/4))*((a^2*e^4*(-a^7*c^5)^(1/2) + c^2*d^4*(-a^7*c^5)^(1/2) - 4*a^4*c^4*d^3*e + 4*a^5*c^3*d*e^3 - 6*a*c*d^2*e^2*(-a^7*c^5)^(1/2))/(a^7*c^5))^(1/4))/4 - (\text{atan}((a^3*e^6*x - c^3*d^6*x - a*c^2*d^4*e^2*x + a^2*c*d^2*e^4*x + (2*d*e*x*(a^2*e^4*(-a^7*c^5)^(1/2) + c^2*d^4*(-a^7*c^5)^(1/2) + 4*a^4*c^4*d^3*e - 4*a^5*c^3*d*e^3 - 6*a*c*d^2*e^2*(-a^7*c^5)^(1/2)))/(a^3*c^2))/(a*c^3*d^5*(-(a^2*e^4*(-a^7*c^5)^(1/2) + c^2*d^4*(-a^7*c^5)^(1/2) + 4*a^4*c^4*d^3*e - 4*a^5*c^3*d*e^3 - 6*a*c*d^2*e^2*(-a^7*c^5)^(1/2))/(a^7*c^5))^(1/4) + a^5*c^3*e*(-(a^2*e^4*(-a^7*c^5)^(1/2) + c^2*d^4*(-a^7*c^5)^(1/2) + 4*a^4*c^4*d^3*e - 4*a^5*c^3*d*e^3 - 6*a*c*d^2*e^2*(-a^7*c^5)^(1/2))...
 \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.22 (sec), antiderivative size = 1033, normalized size of antiderivative = 1.79

$$\int \frac{d + ex^4}{a + cx^8} dx = \text{Too large to display}$$

input `int((e*x^4+d)/(c*x^8+a),x)`

output

$$\begin{aligned}
 & (c^{(3/8)} * a^{(1/8)} * (2 * \sqrt{a}) * \sqrt{\sqrt{2} + 2}) * \sqrt{2} * \operatorname{atan}((c^{(1/8)} * a^{(1/8)} * \sqrt{2}) \\
 & - 2 * \sqrt{2} + 2) - 2 * c^{(1/4)} * x) / (c^{(1/8)} * a^{(1/8)} * \sqrt{\sqrt{2} + 2}) * e - 2 * \sqrt{a} * \\
 & \sqrt{2} + 2) - 2 * c^{(1/4)} * x) / (c^{(1/8)} * a^{(1/8)} * \sqrt{\sqrt{2} + 2}) * e - 2 * \sqrt{c} * \\
 & \sqrt{\sqrt{2} + 2} * \operatorname{atan}((c^{(1/8)} * a^{(1/8)} * \sqrt{\sqrt{2} + 2}) * d - 2 * \sqrt{a} * \sqrt{\sqrt{2} \\
 & + 2} * \sqrt{2} * \operatorname{atan}((c^{(1/8)} * a^{(1/8)} * \sqrt{\sqrt{2} + 2}) * e + 2 * \sqrt{a} * \sqrt{\sqrt{2} + 2} * \\
 & \operatorname{atan}((c^{(1/8)} * a^{(1/8)} * \sqrt{\sqrt{2} + 2}) * e + 2 * \sqrt{c} * \sqrt{\sqrt{2} + 2}) * \operatorname{atan}((c^{(1/8)} * a \\
 & * (1/8)) * \sqrt{\sqrt{2} + 2} + 2 * c^{(1/4)} * x) / (c^{(1/8)} * a^{(1/8)} * \sqrt{\sqrt{2} + 2})) * d - 2 * \sqrt{a} * \sqrt{\sqrt{2} \\
 & + 2} * \sqrt{2} * \operatorname{atan}((c^{(1/8)} * a^{(1/8)} * \sqrt{\sqrt{2} + 2}) * e - 2 * \sqrt{a} * \sqrt{\sqrt{2} + 2} * \\
 & \operatorname{atan}((c^{(1/8)} * a^{(1/8)} * \sqrt{\sqrt{2} + 2}) * e - 2 * \sqrt{c} * \sqrt{\sqrt{2} + 2}) * \operatorname{atan}((c^{(1/8)} * a \\
 & * (1/8)) * \sqrt{\sqrt{2} + 2} + 2 * c^{(1/4)} * x) / (c^{(1/8)} * a^{(1/8)} * \sqrt{\sqrt{2} + 2})) * e - 2 * \sqrt{c} * \\
 & \sqrt{\sqrt{2} + 2} * \operatorname{atan}((c^{(1/8)} * a^{(1/8)} * \sqrt{\sqrt{2} + 2}) * e + 2 * \sqrt{a} * \sqrt{\sqrt{2} \\
 & + 2} * \sqrt{2} * \operatorname{atan}((c^{(1/8)} * a^{(1/8)} * \sqrt{\sqrt{2} + 2}) * e + 2 * \sqrt{c} * \sqrt{\sqrt{2} + 2}) * \operatorname{atan}((c^{(1/8)} * a \\
 & * (1/8)) * \sqrt{\sqrt{2} + 2} + 2 * c^{(1/4)} * x) / (c^{(1/8)} * a^{(1/8)} * \sqrt{\sqrt{2} + 2}) * f
 \end{aligned}$$

3.16 $\int \frac{d+ex^4}{\sqrt{1-\frac{e^2x^8}{d^2}}} dx$

Optimal result	166
Mathematica [A] (verified)	166
Rubi [C] (warning: unable to verify)	167
Maple [A] (verified)	169
Fricas [F]	169
Sympy [C] (verification not implemented)	169
Maxima [F]	170
Giac [F]	170
Mupad [F(-1)]	170
Reduce [F]	171

Optimal result

Integrand size = 25, antiderivative size = 52

$$\begin{aligned} \int \frac{d+ex^4}{\sqrt{1-\frac{e^2x^8}{d^2}}} dx &= dx \text{Hypergeometric2F1}\left(\frac{1}{8}, \frac{1}{2}, \frac{9}{8}, \frac{e^2x^8}{d^2}\right) \\ &\quad + \frac{1}{5}ex^5 \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{8}, \frac{13}{8}, \frac{e^2x^8}{d^2}\right) \end{aligned}$$

output $d*x*\text{hypergeom}([1/8, 1/2], [9/8], e^{2*x^8/d^2}) + 1/5*e*x^5*\text{hypergeom}([1/2, 5/8], [13/8], e^{2*x^8/d^2})$

Mathematica [A] (verified)

Time = 10.03 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.00

$$\begin{aligned} \int \frac{d+ex^4}{\sqrt{1-\frac{e^2x^8}{d^2}}} dx &= dx \text{Hypergeometric2F1}\left(\frac{1}{8}, \frac{1}{2}, \frac{9}{8}, \frac{e^2x^8}{d^2}\right) \\ &\quad + \frac{1}{5}ex^5 \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{8}, \frac{13}{8}, \frac{e^2x^8}{d^2}\right) \end{aligned}$$

input $\text{Integrate}[(d + e*x^4)/\text{Sqrt}[1 - (e^2*x^8)/d^2], x]$

output $d*x*\text{Hypergeometric2F1}[1/8, 1/2, 9/8, (e^2*x^8)/d^2] + (e*x^5*\text{Hypergeometric2F1}[1/2, 5/8, 13/8, (e^2*x^8)/d^2])/5$

Rubi [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 0.26 (sec), antiderivative size = 85, normalized size of antiderivative = 1.63, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.160, Rules used = {1396, 937, 937, 936}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{d + ex^4}{\sqrt{1 - \frac{e^2 x^8}{d^2}}} dx \\
 & \quad \downarrow \textcolor{blue}{1396} \\
 & \frac{\sqrt{d + ex^4} \sqrt{\frac{1}{d} - \frac{ex^4}{d^2}} \int \frac{\sqrt{ex^4 + d}}{\sqrt{\frac{1}{d} - \frac{ex^4}{d^2}}} dx}{\sqrt{1 - \frac{e^2 x^8}{d^2}}} \\
 & \quad \downarrow \textcolor{blue}{937} \\
 & \frac{(d + ex^4) \sqrt{\frac{1}{d} - \frac{ex^4}{d^2}} \int \frac{\sqrt{\frac{ex^4}{d} + 1}}{\sqrt{\frac{1}{d} - \frac{ex^4}{d^2}}} dx}{\sqrt{\frac{ex^4}{d} + 1} \sqrt{1 - \frac{e^2 x^8}{d^2}}} \\
 & \quad \downarrow \textcolor{blue}{937} \\
 & \frac{(d + ex^4) \sqrt{1 - \frac{ex^4}{d}} \int \frac{\sqrt{\frac{ex^4}{d} + 1}}{\sqrt{1 - \frac{ex^4}{d}}} dx}{\sqrt{\frac{ex^4}{d} + 1} \sqrt{1 - \frac{e^2 x^8}{d^2}}} \\
 & \quad \downarrow \textcolor{blue}{936}
 \end{aligned}$$

$$\frac{x(d + ex^4) \sqrt{1 - \frac{ex^4}{d}} \text{AppellF1}\left(\frac{1}{4}, \frac{1}{2}, -\frac{1}{2}, \frac{5}{4}, \frac{ex^4}{d}, -\frac{ex^4}{d}\right)}{\sqrt{\frac{ex^4}{d} + 1} \sqrt{1 - \frac{e^2 x^8}{d^2}}}$$

input `Int[(d + e*x^4)/Sqrt[1 - (e^2*x^8)/d^2], x]`

output `(x*(d + e*x^4)*Sqrt[1 - (e*x^4)/d]*AppellF1[1/4, 1/2, -1/2, 5/4, (e*x^4)/d, -((e*x^4)/d)])/(Sqrt[1 + (e*x^4)/d]*Sqrt[1 - (e^2*x^8)/d^2])`

Definitions of rubi rules used

rule 936 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> Simplify[a^p*c^q*x^q*AppellF1[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`

rule 937 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> Simplify[a^p*IntPart[p]*((a + b*x^n)^p*(c + d*x^n)^q, x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && !(IntegerQ[p] || GtQ[a, 0])`

rule 1396 `Int[(u_)*(a_ + c_)*(x_)^(n2_))^(p_)*((d_) + (e_)*(x_)^(n_))^(q_), x_Symbol] :> Simplify[(a + c*x^(2*n))^p*((d + e*x^n)^q*(a/d + c/e*x^n)^p, x) /; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && !(EqQ[q, 1] && EqQ[n, 2])`

Maple [A] (verified)

Time = 3.30 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.83

method	result	size
meijerg	$dx \text{ hypergeom} \left(\left[\frac{1}{8}, \frac{1}{2} \right], \left[\frac{9}{8} \right], \frac{e^2 x^8}{d^2} \right) + \frac{e x^5 \text{ hypergeom} \left(\left[\frac{1}{2}, \frac{5}{8} \right], \left[\frac{13}{8} \right], \frac{e^2 x^8}{d^2} \right)}{5}$	43

input `int((e*x^4+d)/(1-e^2*x^8/d^2)^(1/2),x,method=_RETURNVERBOSE)`

output `d*x*hypergeom([1/8,1/2],[9/8],e^2*x^8/d^2)+1/5*e*x^5*hypergeom([1/2,5/8],[13/8],e^2*x^8/d^2)`

Fricas [F]

$$\int \frac{d + ex^4}{\sqrt{1 - \frac{e^2 x^8}{d^2}}} dx = \int \frac{ex^4 + d}{\sqrt{-\frac{e^2 x^8}{d^2} + 1}} dx$$

input `integrate((e*x^4+d)/(1-e^2*x^8/d^2)^(1/2),x, algorithm="fricas")`

output `integral(-d^2*sqrt(-(e^2*x^8 - d^2)/d^2)/(e*x^4 - d), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.05 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.50

$$\int \frac{d + ex^4}{\sqrt{1 - \frac{e^2 x^8}{d^2}}} dx = \frac{dx \Gamma \left(\frac{1}{8} \right) {}_2F_1 \left(\begin{matrix} \frac{1}{8}, \frac{1}{2} \\ \frac{9}{8} \end{matrix} \middle| \frac{e^2 x^8 e^{2i\pi}}{d^2} \right)}{8 \Gamma \left(\frac{9}{8} \right)} + \frac{ex^5 \Gamma \left(\frac{5}{8} \right) {}_2F_1 \left(\begin{matrix} \frac{1}{2}, \frac{5}{8} \\ \frac{13}{8} \end{matrix} \middle| \frac{e^2 x^8 e^{2i\pi}}{d^2} \right)}{8 \Gamma \left(\frac{13}{8} \right)}$$

input `integrate((e*x**4+d)/(1-e**2*x**8/d**2)**(1/2),x)`

output
$$\frac{d*x*gamma(1/8)*hyper((1/8, 1/2), (9/8,), e^{**2}*x^{**8}*exp_polar(2*I*pi)/d^{**2})}{(8*gamma(9/8)) + e*x^{**5}*gamma(5/8)*hyper((1/2, 5/8), (13/8,), e^{**2}*x^{**8}*exp_polar(2*I*pi)/d^{**2})/(8*gamma(13/8))}$$

Maxima [F]

$$\int \frac{d + ex^4}{\sqrt{1 - \frac{e^2 x^8}{d^2}}} dx = \int \frac{ex^4 + d}{\sqrt{-\frac{e^2 x^8}{d^2} + 1}} dx$$

input `integrate((e*x^4+d)/(1-e^2*x^8/d^2)^(1/2),x, algorithm="maxima")`

output `integrate((e*x^4 + d)/sqrt(-e^2*x^8/d^2 + 1), x)`

Giac [F]

$$\int \frac{d + ex^4}{\sqrt{1 - \frac{e^2 x^8}{d^2}}} dx = \int \frac{ex^4 + d}{\sqrt{-\frac{e^2 x^8}{d^2} + 1}} dx$$

input `integrate((e*x^4+d)/(1-e^2*x^8/d^2)^(1/2),x, algorithm="giac")`

output `integrate((e*x^4 + d)/sqrt(-e^2*x^8/d^2 + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{d + ex^4}{\sqrt{1 - \frac{e^2 x^8}{d^2}}} dx = \int \frac{e x^4 + d}{\sqrt{1 - \frac{e^2 x^8}{d^2}}} dx$$

input `int((d + e*x^4)/(1 - (e^2*x^8)/d^2)^(1/2),x)`

output `int((d + e*x^4)/(1 - (e^2*x^8)/d^2)^(1/2), x)`

Reduce [F]

$$\int \frac{d + ex^4}{\sqrt{1 - \frac{e^2x^8}{d^2}}} dx = \left(\int \frac{\sqrt{-e^2x^8 + d^2}}{-e x^4 + d} dx \right) d$$

input `int((e*x^4+d)/(1-e^2*x^8/d^2)^(1/2),x)`

output `int(sqrt(d**2 - e**2*x**8)/(d - e*x**4),x)*d`

3.17 $\int \frac{d+ex^4}{\sqrt{d^2-e^2x^8}} dx$

Optimal result	172
Mathematica [A] (verified)	172
Rubi [C] (warning: unable to verify)	173
Maple [F]	174
Fricas [F]	175
Sympy [C] (verification not implemented)	175
Maxima [F]	175
Giac [F]	176
Mupad [F(-1)]	176
Reduce [F]	176

Optimal result

Integrand size = 24, antiderivative size = 118

$$\begin{aligned} \int \frac{d + ex^4}{\sqrt{d^2 - e^2x^8}} dx &= \frac{dx \sqrt{1 - \frac{e^2x^8}{d^2}} \text{Hypergeometric2F1} \left(\frac{1}{8}, \frac{1}{2}, \frac{9}{8}, \frac{e^2x^8}{d^2} \right)}{\sqrt{d^2 - e^2x^8}} \\ &+ \frac{ex^5 \sqrt{1 - \frac{e^2x^8}{d^2}} \text{Hypergeometric2F1} \left(\frac{1}{2}, \frac{5}{8}, \frac{13}{8}, \frac{e^2x^8}{d^2} \right)}{5\sqrt{d^2 - e^2x^8}} \end{aligned}$$

output $d*x*(1-e^2*x^8/d^2)^(1/2)*hypergeom([1/8, 1/2], [9/8], e^2*x^8/d^2)/(-e^2*x^8+d^2)^(1/2)+1/5*e*x^5*(1-e^2*x^8/d^2)^(1/2)*hypergeom([1/2, 5/8], [13/8], e^2*x^8/d^2)/(-e^2*x^8+d^2)^(1/2)$

Mathematica [A] (verified)

Time = 10.06 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.74

$$\begin{aligned} &\int \frac{d + ex^4}{\sqrt{d^2 - e^2x^8}} dx \\ &= \frac{\sqrt{1 - \frac{e^2x^8}{d^2}} \left(5dx \text{Hypergeometric2F1} \left(\frac{1}{8}, \frac{1}{2}, \frac{9}{8}, \frac{e^2x^8}{d^2} \right) + ex^5 \text{Hypergeometric2F1} \left(\frac{1}{2}, \frac{5}{8}, \frac{13}{8}, \frac{e^2x^8}{d^2} \right) \right)}{5\sqrt{d^2 - e^2x^8}} \end{aligned}$$

input $\text{Integrate}[(d + e*x^4)/\text{Sqrt}[d^2 - e^2*x^8], x]$

output $(\text{Sqrt}[1 - (e^2*x^8)/d^2]*(5*d*x*\text{Hypergeometric2F1}[1/8, 1/2, 9/8, (e^2*x^8)/d^2] + e*x^5*\text{Hypergeometric2F1}[1/2, 5/8, 13/8, (e^2*x^8)/d^2]))/(5*\text{Sqrt}[d^2 - e^2*x^8])$

Rubi [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 0.25 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.71, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {1396, 937, 937, 936}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{d + ex^4}{\sqrt{d^2 - e^2x^8}} dx \\
 & \quad \downarrow 1396 \\
 & \frac{\sqrt{d - ex^4}\sqrt{d + ex^4} \int \frac{\sqrt{ex^4 + d}}{\sqrt{d - ex^4}} dx}{\sqrt{d^2 - e^2x^8}} \\
 & \quad \downarrow 937 \\
 & \frac{\sqrt{d + ex^4}\sqrt{1 - \frac{ex^4}{d}} \int \frac{\sqrt{ex^4 + d}}{\sqrt{1 - \frac{ex^4}{d}}} dx}{\sqrt{d^2 - e^2x^8}} \\
 & \quad \downarrow 937 \\
 & \frac{(d + ex^4)\sqrt{1 - \frac{ex^4}{d}} \int \frac{\sqrt{\frac{ex^4}{d} + 1}}{\sqrt{1 - \frac{ex^4}{d}}} dx}{\sqrt{\frac{ex^4}{d} + 1}\sqrt{d^2 - e^2x^8}} \\
 & \quad \downarrow 936 \\
 & \frac{x(d + ex^4)\sqrt{1 - \frac{ex^4}{d}} \text{AppellF1}\left(\frac{1}{4}, \frac{1}{2}, -\frac{1}{2}, \frac{5}{4}, \frac{ex^4}{d}, -\frac{ex^4}{d}\right)}{\sqrt{\frac{ex^4}{d} + 1}\sqrt{d^2 - e^2x^8}}
 \end{aligned}$$

input $\text{Int}[(d + e*x^4)/\text{Sqrt}[d^2 - e^2*x^8], x]$

output $(x*(d + e*x^4)*\text{Sqrt}[1 - (e*x^4)/d]*\text{AppellF1}[1/4, 1/2, -1/2, 5/4, (e*x^4)/d, -((e*x^4)/d)])/(\text{Sqrt}[1 + (e*x^4)/d]*\text{Sqrt}[d^2 - e^2*x^8])$

Definitions of rubi rules used

rule 936 $\text{Int}[(a_ + b_)*(x_)^{(n_)}*(p_)*((c_ + d_)*(x_)^{(n_)})^{(q_)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[a^p*c^q*x*\text{AppellF1}[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /; \text{FreeQ}[\{a, b, c, d, n, p, q\}, x] \&& \text{NeQ}[b*c - a*d, 0] \&& \text{NeQ}[n, -1] \&& (\text{IntegerQ}[p] \mid\mid \text{GtQ}[a, 0]) \&& (\text{IntegerQ}[q] \mid\mid \text{GtQ}[c, 0])$

rule 937 $\text{Int}[(a_ + b_)*(x_)^{(n_)}*(p_)*((c_ + d_)*(x_)^{(n_)})^{(q_)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[a^{\text{IntPart}[p]}*((a + b*x^n)^{\text{FracPart}[p]}/(1 + b*(x^n/a))^{\text{FracPart}[p]}) \text{Int}[(1 + b*(x^n/a))^{p*}(c + d*x^n)^q, x], x] /; \text{FreeQ}[\{a, b, c, d, n, p, q\}, x] \&& \text{NeQ}[b*c - a*d, 0] \&& \text{NeQ}[n, -1] \&& !(\text{IntegerQ}[p] \mid\mid \text{GtQ}[a, 0])$

rule 1396 $\text{Int}[(u_)*(a_ + c_)*(x_)^{(n2_)}*(p_)*((d_ + e_)*(x_)^{(n_)})^{(q_)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[(a + c*x^{(2*n)})^{\text{FracPart}[p]}/((d + e*x^n)^{\text{FracPart}[p]}*(a/d + c*(x^n/e))^{\text{FracPart}[p]}) \text{Int}[u*(d + e*x^n)^{(p + q)}*(a/d + (c/e)*x^n)^p, x], x] /; \text{FreeQ}[\{a, c, d, e, n, p, q\}, x] \&& \text{EqQ}[n2, 2*n] \&& \text{EqQ}[c*d^2 + a*e^2, 0] \&& !\text{IntegerQ}[p] \&& !(\text{EqQ}[q, 1] \&& \text{EqQ}[n, 2])$

Maple [F]

$$\int \frac{x^4 e + d}{\sqrt{-e^2 x^8 + d^2}} dx$$

input $\text{int}((e*x^4+d)/(-e^2*x^8+d^2)^{(1/2)}, x)$

output $\text{int}((e*x^4+d)/(-e^2*x^8+d^2)^{(1/2)}, x)$

Fricas [F]

$$\int \frac{d + ex^4}{\sqrt{d^2 - e^2x^8}} dx = \int \frac{ex^4 + d}{\sqrt{-e^2x^8 + d^2}} dx$$

input `integrate((e*x^4+d)/(-e^2*x^8+d^2)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-e^2*x^8 + d^2)/(e*x^4 - d), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.08 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.66

$$\int \frac{d + ex^4}{\sqrt{d^2 - e^2x^8}} dx = \frac{x\Gamma(\frac{1}{8}) {}_2F_1\left(\begin{matrix} \frac{1}{8}, \frac{1}{2} \\ \frac{9}{8} \end{matrix} \middle| \frac{e^2x^8e^{2i\pi}}{d^2}\right)}{8\Gamma(\frac{9}{8})} + \frac{ex^5\Gamma(\frac{5}{8}) {}_2F_1\left(\begin{matrix} \frac{1}{2}, \frac{5}{8} \\ \frac{13}{8} \end{matrix} \middle| \frac{e^2x^8e^{2i\pi}}{d^2}\right)}{8d\Gamma(\frac{13}{8})}$$

input `integrate((e*x**4+d)/(-e**2*x**8+d**2)**(1/2),x)`

output `x*gamma(1/8)*hyper((1/8, 1/2), (9/8,), e**2*x**8*exp_polar(2*I*pi)/d**2)/(8*gamma(9/8)) + e*x**5*gamma(5/8)*hyper((1/2, 5/8), (13/8,), e**2*x**8*exp_polar(2*I*pi)/d**2)/(8*d*gamma(13/8))`

Maxima [F]

$$\int \frac{d + ex^4}{\sqrt{d^2 - e^2x^8}} dx = \int \frac{ex^4 + d}{\sqrt{-e^2x^8 + d^2}} dx$$

input `integrate((e*x^4+d)/(-e^2*x^8+d^2)^(1/2),x, algorithm="maxima")`

output `integrate((e*x^4 + d)/sqrt(-e^2*x^8 + d^2), x)`

Giac [F]

$$\int \frac{d + ex^4}{\sqrt{d^2 - e^2x^8}} dx = \int \frac{ex^4 + d}{\sqrt{-e^2x^8 + d^2}} dx$$

input `integrate((e*x^4+d)/(-e^2*x^8+d^2)^(1/2),x, algorithm="giac")`

output `integrate((e*x^4 + d)/sqrt(-e^2*x^8 + d^2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{d + ex^4}{\sqrt{d^2 - e^2x^8}} dx = \int \frac{e x^4 + d}{\sqrt{d^2 - e^2 x^8}} dx$$

input `int((d + e*x^4)/(d^2 - e^2*x^8)^(1/2),x)`

output `int((d + e*x^4)/(d^2 - e^2*x^8)^(1/2), x)`

Reduce [F]

$$\int \frac{d + ex^4}{\sqrt{d^2 - e^2x^8}} dx = \int \frac{\sqrt{-e^2x^8 + d^2}}{-e x^4 + d} dx$$

input `int((e*x^4+d)/(-e^2*x^8+d^2)^(1/2),x)`

output `int(sqrt(d**2 - e**2*x**8)/(d - e*x**4),x)`

3.18 $\int \frac{d+ex^4}{\sqrt{a-cx^8}} dx$

Optimal result	177
Mathematica [A] (verified)	177
Rubi [B] (warning: unable to verify)	178
Maple [F]	179
Fricas [F]	180
Sympy [C] (verification not implemented)	180
Maxima [F]	180
Giac [F]	181
Mupad [F(-1)]	181
Reduce [F]	181

Optimal result

Integrand size = 20, antiderivative size = 102

$$\int \frac{d+ex^4}{\sqrt{a-cx^8}} dx = \frac{dx \sqrt{1 - \frac{cx^8}{a}} \text{Hypergeometric2F1} \left(\frac{1}{8}, \frac{1}{2}, \frac{9}{8}, \frac{cx^8}{a} \right)}{\sqrt{a-cx^8}} + \frac{ex^5 \sqrt{1 - \frac{cx^8}{a}} \text{Hypergeometric2F1} \left(\frac{1}{2}, \frac{5}{8}, \frac{13}{8}, \frac{cx^8}{a} \right)}{5\sqrt{a-cx^8}}$$

output $d*x*(1-c*x^8/a)^(1/2)*hypergeom([1/8, 1/2], [9/8], c*x^8/a)/(-c*x^8+a)^(1/2) + 1/5*e*x^5*(1-c*x^8/a)^(1/2)*hypergeom([1/2, 5/8], [13/8], c*x^8/a)/(-c*x^8+a)^(1/2)$

Mathematica [A] (verified)

Time = 10.05 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.75

$$\begin{aligned} & \int \frac{d+ex^4}{\sqrt{a-cx^8}} dx \\ &= \frac{\sqrt{1 - \frac{cx^8}{a}} \left(5dx \text{Hypergeometric2F1} \left(\frac{1}{8}, \frac{1}{2}, \frac{9}{8}, \frac{cx^8}{a} \right) + ex^5 \text{Hypergeometric2F1} \left(\frac{1}{2}, \frac{5}{8}, \frac{13}{8}, \frac{cx^8}{a} \right) \right)}{5\sqrt{a-cx^8}} \end{aligned}$$

input $\text{Integrate}[(d + e*x^4)/\text{Sqrt}[a - c*x^8], x]$

output $(\text{Sqrt}[1 - (c*x^8)/a]*(5*d*x*\text{Hypergeometric2F1}[1/8, 1/2, 9/8, (c*x^8)/a] + e*x^5*\text{Hypergeometric2F1}[1/2, 5/8, 13/8, (c*x^8)/a]))/(5*\text{Sqrt}[a - c*x^8])$

Rubi [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 482 vs. $2(102) = 204$.

Time = 0.62 (sec), antiderivative size = 482, normalized size of antiderivative = 4.73, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {1763, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{d + ex^4}{\sqrt{a - cx^8}} dx \\
 & \quad \downarrow 1763 \\
 & \int \left(\frac{d}{\sqrt{a - cx^8}} + \frac{ex^4}{\sqrt{a - cx^8}} \right) dx \\
 & \quad \downarrow 2009 \\
 & \frac{\sqrt[4]{c} dx^3 \sqrt{\frac{\sqrt[4]{-a} \left(\frac{\sqrt[4]{c} x^2}{\sqrt[4]{-a}} + 1\right)^2}{\sqrt[4]{c} x^2}} \sqrt{\frac{a - cx^8}{\sqrt{-a} \sqrt{c} x^4}} \text{EllipticF}\left(\arcsin\left(\frac{1}{2} \sqrt{-\frac{\sqrt[4]{-a} \left(\frac{\sqrt{2} \sqrt{c} x^4}{\sqrt{-a}} - \frac{2 \sqrt[4]{c} x^2}{\sqrt[4]{-a}} + \sqrt{2}\right)}{\sqrt[4]{c} x^2}}}{\sqrt[4]{c} x^2}\right)}, -2(1 - \sqrt{2})\right)}{2 \sqrt{2 + \sqrt{2}} \sqrt[4]{-a} \left(\frac{\sqrt[4]{c} x^2}{\sqrt[4]{-a}} + 1\right) \sqrt{a - cx^8}} \\
 & \quad - \frac{\sqrt[4]{c} dx^3 \sqrt{-\frac{\sqrt[4]{-a} \left(1 - \frac{\sqrt[4]{c} x^2}{\sqrt[4]{-a}}\right)^2}{\sqrt[4]{c} x^2}} \sqrt{\frac{a - cx^8}{\sqrt{-a} \sqrt{c} x^4}} \text{EllipticF}\left(\arcsin\left(\frac{1}{2} \sqrt{\frac{\sqrt[4]{-a} \left(\frac{\sqrt{2} \sqrt{c} x^4}{\sqrt{-a}} + \frac{2 \sqrt[4]{c} x^2}{\sqrt[4]{-a}} + \sqrt{2}\right)}{\sqrt[4]{c} x^2}}}{\sqrt[4]{c} x^2}\right)}, -2(1 - \sqrt{2})\right)}{2 \sqrt{2 + \sqrt{2}} \sqrt[4]{-a} \left(1 - \frac{\sqrt[4]{c} x^2}{\sqrt[4]{-a}}\right) \sqrt{a - cx^8}} \\
 & \quad + \frac{e x^5 \sqrt{1 - \frac{c x^8}{a}} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{8}, \frac{13}{8}, \frac{c x^8}{a}\right)}{5 \sqrt{a - cx^8}}
 \end{aligned}$$

input $\text{Int}[(d + e*x^4)/\sqrt{a - c*x^8}, x]$

output
$$\begin{aligned} & \left(c^{(1/4)} * d * x^3 * \sqrt{((-a)^{(1/4)} * (1 + (c^{(1/4)} * x^2) / (-a)^{(1/4)})^2) / (c^{(1/4)} * x^2)} \right) * \sqrt{(a - c*x^8) / (\sqrt{-a} * \sqrt{c} * x^4)} * \text{EllipticF}[\text{ArcSin}[\sqrt{-(((-a)^{(1/4)} * (\sqrt{2} - (2*c^{(1/4)} * x^2) / (-a)^{(1/4}) + (\sqrt{2} * \sqrt{c} * x^4) / \sqrt{-a})) / (c^{(1/4)} * x^2))} / 2], -2 * (1 - \sqrt{2})] / (2 * \sqrt{2 + \sqrt{2}} * (-a)^{(1/4)} * (1 + (c^{(1/4)} * x^2) / (-a)^{(1/4)}) * \sqrt{a - c*x^8}) - (c^{(1/4)} * d * x^3 * \sqrt{-(((-a)^{(1/4)} * (1 - (c^{(1/4)} * x^2) / (-a)^{(1/4)})^2) / (c^{(1/4)} * x^2)}) * \sqrt{(a - c*x^8) / (\sqrt{-a} * \sqrt{c} * x^4)} * \text{EllipticF}[\text{ArcSin}[\sqrt{((-a)^{(1/4)} * (\sqrt{2} + (2*c^{(1/4)} * x^2) / (-a)^{(1/4}) + (\sqrt{2} * \sqrt{c} * x^4) / \sqrt{-a})) / (c^{(1/4)} * x^2))} / 2], -2 * (1 - \sqrt{2})] / (2 * \sqrt{2 + \sqrt{2}} * (-a)^{(1/4)} * (1 - (c^{(1/4)} * x^2) / (-a)^{(1/4)}) * \sqrt{a - c*x^8}) + (e*x^5 * \sqrt{1 - (c*x^8) / a} * \text{Hypergeometric2F1}[1/2, 5/8, 13/8, (c*x^8) / a]) / (5 * \sqrt{a - c*x^8}) \end{aligned}$$

Definitions of rubi rules used

rule 1763 $\text{Int}[(d_+ + e_*)*(x_-)^(n_-)*((a_- + c_*)*(x_-)^(n2_-))^p, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(d + e*x^n)*(a + c*x^(2*n))^p, x], x] /; \text{FreeQ}[\{a, c, d, e, n\}, x] \&& \text{EqQ}[n2, 2*n]$

rule 2009 $\text{Int}[u_, x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

Maple [F]

$$\int \frac{x^4 e + d}{\sqrt{-c x^8 + a}} dx$$

input $\text{int}((e*x^4+d)/(-c*x^8+a)^{(1/2)}, x)$

output $\text{int}((e*x^4+d)/(-c*x^8+a)^{(1/2)}, x)$

Fricas [F]

$$\int \frac{d + ex^4}{\sqrt{a - cx^8}} dx = \int \frac{ex^4 + d}{\sqrt{-cx^8 + a}} dx$$

input `integrate((e*x^4+d)/(-c*x^8+a)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c*x^8 + a)*(e*x^4 + d)/(c*x^8 - a), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.96 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.80

$$\int \frac{d + ex^4}{\sqrt{a - cx^8}} dx = \frac{dx\Gamma(\frac{1}{8}) {}_2F_1\left(\begin{matrix} \frac{1}{8}, \frac{1}{2} \\ \frac{9}{8} \end{matrix} \middle| \frac{cx^8 e^{2i\pi}}{a}\right)}{8\sqrt{a}\Gamma(\frac{9}{8})} + \frac{ex^5\Gamma(\frac{5}{8}) {}_2F_1\left(\begin{matrix} \frac{1}{2}, \frac{5}{8} \\ \frac{13}{8} \end{matrix} \middle| \frac{cx^8 e^{2i\pi}}{a}\right)}{8\sqrt{a}\Gamma(\frac{13}{8})}$$

input `integrate((e*x**4+d)/(-c*x**8+a)**(1/2),x)`

output `d*x*gamma(1/8)*hyper((1/8, 1/2), (9/8,), c*x**8*exp_polar(2*I*pi)/a)/(8*sqrt(a)*gamma(9/8)) + e*x**5*gamma(5/8)*hyper((1/2, 5/8), (13/8,), c*x**8*exp_polar(2*I*pi)/a)/(8*sqrt(a)*gamma(13/8))`

Maxima [F]

$$\int \frac{d + ex^4}{\sqrt{a - cx^8}} dx = \int \frac{ex^4 + d}{\sqrt{-cx^8 + a}} dx$$

input `integrate((e*x^4+d)/(-c*x^8+a)^(1/2),x, algorithm="maxima")`

output `integrate((e*x^4 + d)/sqrt(-c*x^8 + a), x)`

Giac [F]

$$\int \frac{d + ex^4}{\sqrt{a - cx^8}} dx = \int \frac{ex^4 + d}{\sqrt{-cx^8 + a}} dx$$

input `integrate((e*x^4+d)/(-c*x^8+a)^(1/2),x, algorithm="giac")`

output `integrate((e*x^4 + d)/sqrt(-c*x^8 + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{d + ex^4}{\sqrt{a - cx^8}} dx = \int \frac{e x^4 + d}{\sqrt{a - c x^8}} dx$$

input `int((d + e*x^4)/(a - c*x^8)^(1/2),x)`

output `int((d + e*x^4)/(a - c*x^8)^(1/2), x)`

Reduce [F]

$$\int \frac{d + ex^4}{\sqrt{a - cx^8}} dx = \left(\int \frac{\sqrt{-cx^8 + a}}{-cx^8 + a} dx \right) d + \left(\int \frac{\sqrt{-cx^8 + a} x^4}{-cx^8 + a} dx \right) e$$

input `int((e*x^4+d)/(-c*x^8+a)^(1/2),x)`

output `int(sqrt(a - c*x**8)/(a - c*x**8),x)*d + int((sqrt(a - c*x**8)*x**4)/(a - c*x**8),x)*e`

3.19 $\int \frac{d+ex^4}{\sqrt{a+cx^8}} dx$

Optimal result	182
Mathematica [C] (verified)	183
Rubi [C] (warning: unable to verify)	184
Maple [F]	185
Fricas [F]	186
Sympy [C] (verification not implemented)	186
Maxima [F]	186
Giac [F]	187
Mupad [F(-1)]	187
Reduce [F]	187

Optimal result

Integrand size = 19, antiderivative size = 905

$$\int \frac{d + ex^4}{\sqrt{a + cx^8}} dx = \text{Too large to display}$$

output

$$\begin{aligned} & \frac{1}{4} \cdot (2+2^{(1/2)})^{(1/2)} \cdot (c^{(1/2)} \cdot d + a^{(1/2)} \cdot e) \cdot x^3 \cdot ((-a^{(1/2)})^{(1/2)} \cdot (1+c^{(1/4)}) \\ & \cdot x^2 \cdot (-a^{(1/2)})^{(1/2)} \cdot c^{(1/4)} / x^2)^{(1/2)} \cdot ((c \cdot x^8 + a) / a^{(1/2)} / c^{(1/2)} / x^4)^{(1/2)} \cdot \text{EllipticE}(1/2 \cdot (-(-a^{(1/2)})^{(1/2)} \cdot (2^{(1/2)} - 2 \cdot c^{(1/4)} \cdot x^2 / (-a^{(1/2)})^{(1/2)} - 2 \cdot (2+2 \cdot 2^{(1/2)})^{(1/2)} / (-a^{(1/2)})^{(1/2)} - 1/4 \cdot (2+2^{(1/2)})^{(1/2)} \cdot (c^{(1/2)} \cdot d + a^{(1/2)} \cdot e) \cdot x^3 \cdot (-(-a^{(1/2)})^{(1/2)} \cdot (1 - c^{(1/4)} \cdot x^2 / (-a^{(1/2)})^{(1/2)} \cdot c^{(1/4)} / x^2)^{(1/2)} \cdot ((c \cdot x^8 + a) / a^{(1/2)} / c^{(1/2)} / x^4)^{(1/2)} \cdot \text{EllipticE}(1/2 \cdot ((-a^{(1/2)})^{(1/2)} \cdot (2^{(1/2)} + 2 \cdot c^{(1/4)} \cdot x^2 / (-a^{(1/2)})^{(1/2)} - 2 \cdot (2+2 \cdot 2^{(1/2)})^{(1/2)} / (-a^{(1/2)})^{(1/2)} - 1/4 \cdot (2+2^{(1/2)})^{(1/2)} \cdot (c^{(1/2)} \cdot d - a^{(1/2)} \cdot e) \cdot x^3 \cdot ((a^{(1/4)} + c^{(1/4)}) \cdot x^2 / a^{(1/4)} / c^{(1/4)} / x^2)^{(1/2)} \cdot ((-c \cdot x^8 + a) / a^{(1/2)} / c^{(1/2)} / x^4)^{(1/2)} \cdot \text{EllipticE}(1/2 \cdot (-a^{(1/4)} \cdot (2^{(1/2)} - 2 \cdot c^{(1/4)} \cdot x^2 / a^{(1/4)} + 2^{(1/2)} \cdot c^{(1/2)} \cdot x^4 / a^{(1/2)}) / c^{(1/4)} / (a^{(1/4)} + c^{(1/4)}) \cdot x^2 / (c \cdot x^8 + a)^{(1/2)} - 1/4 \cdot (2+2^{(1/2)})^{(1/2)} \cdot (c^{(1/2)} \cdot d - a^{(1/2)} \cdot e) \cdot x^3 \cdot (-a^{(1/4)} - c^{(1/4)} \cdot x^2 \cdot 2 / a^{(1/4)} / c^{(1/4)} / x^2)^{(1/2)} \cdot ((-c \cdot x^8 + a) / a^{(1/2)} / c^{(1/2)} / x^4)^{(1/2)} \cdot \text{EllipticE}(1/2 \cdot (a^{(1/4)} \cdot (2^{(1/2)} + 2 \cdot c^{(1/4)} \cdot x^2 / a^{(1/4)} + 2^{(1/2)} \cdot c^{(1/2)} \cdot x^4 / a^{(1/2)}) / c^{(1/4)} / x^2)^{(1/2)} \cdot (-2+2 \cdot 2^{(1/2)})^{(1/2)} / c^{(1/4)} / (a^{(1/4)} - c^{(1/4)} \cdot x^2) / (c \cdot x^8 + a)^{(1/2)}} \end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.06 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.09

$$\begin{aligned} & \int \frac{d + ex^4}{\sqrt{a + cx^8}} dx \\ &= \frac{\sqrt{1 + \frac{cx^8}{a}} \left(5dx \text{Hypergeometric2F1} \left(\frac{1}{8}, \frac{1}{2}, \frac{9}{8}, -\frac{cx^8}{a} \right) + ex^5 \text{Hypergeometric2F1} \left(\frac{1}{2}, \frac{5}{8}, \frac{13}{8}, -\frac{cx^8}{a} \right) \right)}{5\sqrt{a + cx^8}} \end{aligned}$$

input

```
Integrate[(d + e*x^4)/Sqrt[a + c*x^8], x]
```

output

$$\begin{aligned} & (\text{Sqrt}[1 + (c \cdot x^8) / a] \cdot (5 \cdot d \cdot x \cdot \text{Hypergeometric2F1}[1/8, 1/2, 9/8, -(c \cdot x^8) / a] \\ & + e \cdot x^5 \cdot \text{Hypergeometric2F1}[1/2, 5/8, 13/8, -(c \cdot x^8) / a])) / (5 \cdot \text{Sqrt}[a + c \cdot x^8]) \end{aligned}$$

Rubi [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.49 (sec) , antiderivative size = 433, normalized size of antiderivative = 0.48, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.105, Rules used = {1763, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{d + ex^4}{\sqrt{a + cx^8}} dx \\
 & \quad \downarrow \textcolor{blue}{1763} \\
 & \int \left(\frac{d}{\sqrt{a + cx^8}} + \frac{ex^4}{\sqrt{a + cx^8}} \right) dx \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & \frac{\sqrt[4]{c} dx^3 \sqrt{\frac{\left(\sqrt[4]{a} + \sqrt[4]{c} x^2\right)^2}{\sqrt[4]{a} \sqrt[4]{c} x^2}} \sqrt{-\frac{a + cx^8}{\sqrt{a} \sqrt{c} x^4}} \operatorname{EllipticF}\left(\arcsin\left(\frac{1}{2} \sqrt{-\frac{\sqrt[4]{a} \left(\frac{\sqrt{2} \sqrt{c} x^4}{\sqrt{a}} - \frac{2 \sqrt[4]{c} x^2}{\sqrt[4]{a}} + \sqrt{2}\right)}{\sqrt[4]{c} x^2}}}{\sqrt[4]{c} x^2}\right), -2(1 - \sqrt{2})\right)}{2 \sqrt{2 + \sqrt{2}} (\sqrt[4]{a} + \sqrt[4]{c} x^2) \sqrt{a + cx^8}} \\
 & - \frac{\sqrt[4]{c} dx^3 \sqrt{-\frac{\left(\sqrt[4]{a} - \sqrt[4]{c} x^2\right)^2}{\sqrt[4]{a} \sqrt[4]{c} x^2}} \sqrt{-\frac{a + cx^8}{\sqrt{a} \sqrt{c} x^4}} \operatorname{EllipticF}\left(\arcsin\left(\frac{1}{2} \sqrt{\frac{\sqrt[4]{a} \left(\frac{\sqrt{2} \sqrt{c} x^4}{\sqrt{a}} + \frac{2 \sqrt[4]{c} x^2}{\sqrt[4]{a}} + \sqrt{2}\right)}{\sqrt[4]{c} x^2}}}{\sqrt[4]{c} x^2}\right), -2(1 - \sqrt{2})\right)}{2 \sqrt{2 + \sqrt{2}} (\sqrt[4]{a} - \sqrt[4]{c} x^2) \sqrt{a + cx^8}} \\
 & + \frac{e x^5 \sqrt{\frac{c x^8}{a} + 1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{8}, \frac{13}{8}, -\frac{c x^8}{a}\right)}{5 \sqrt{a + cx^8}}
 \end{aligned}$$

input `Int[(d + e*x^4)/Sqrt[a + c*x^8],x]`

output

$$\begin{aligned} & \left(c^{(1/4)} d x^3 \sqrt{(a^{(1/4)} + c^{(1/4)} x^2)^2 / (a^{(1/4)} c^{(1/4)} x^2)} \right) \sqrt{-((a + c x^8) / (\sqrt{a} \sqrt{c} x^4))} \operatorname{EllipticF}[\operatorname{ArcSin}[\sqrt{-((a^{(1/4)} (\sqrt{2} - (2 c^{(1/4)} x^2) / a^{(1/4)} + (\sqrt{2} \sqrt{c} x^4) / \sqrt{a})) / (c^{(1/4)} x^2)) / 2}], \\ & -2(1 - \sqrt{2})] / (2 \sqrt{2 + \sqrt{2}}) * (a^{(1/4)} + c^{(1/4)} x^2) \sqrt{a + c x^8} - (c^{(1/4)} d x^3 \sqrt{-((a^{(1/4)} - c^{(1/4)} x^2)^2 / (a^{(1/4)} c^{(1/4)} x^2))} \sqrt{-((a + c x^8) / (\sqrt{a} \sqrt{c} x^4))} \operatorname{EllipticF}[\operatorname{ArcSin}[\sqrt{(a^{(1/4)} (\sqrt{2} + (2 c^{(1/4)} x^2) / a^{(1/4)} + (\sqrt{2} \sqrt{c} x^4) / \sqrt{a})) / (c^{(1/4)} x^2)) / 2], \\ & -2(1 - \sqrt{2})] / (2 \sqrt{2 + \sqrt{2}}) * (a^{(1/4)} - c^{(1/4)} x^2) \sqrt{a + c x^8} + e x^5 \sqrt{1 + (c x^8) / a} \operatorname{Hypergeometric2F1}[1/2, 5/8, 13/8, -((c x^8) / a)] / (5 \sqrt{a + c x^8}) \end{aligned}$$

Definitions of rubi rules used

rule 1763

```
Int[((d_) + (e_)*(x_)^(n_))*((a_) + (c_)*(x_)^(n2_))^(p_), x_Symbol] :> Int[ExpandIntegrand[(d + e*x^n)*(a + c*x^(2*n))^p, x], x] /; FreeQ[{a, c, d, e, n}, x] && EqQ[n2, 2*n]
```

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [F]

$$\int \frac{x^4 e + d}{\sqrt{c x^8 + a}} dx$$

input

```
int((e*x^4+d)/(c*x^8+a)^(1/2),x)
```

output

```
int((e*x^4+d)/(c*x^8+a)^(1/2),x)
```

Fricas [F]

$$\int \frac{d + ex^4}{\sqrt{a + cx^8}} dx = \int \frac{ex^4 + d}{\sqrt{cx^8 + a}} dx$$

input `integrate((e*x^4+d)/(c*x^8+a)^(1/2),x, algorithm="fricas")`

output `integral((e*x^4 + d)/sqrt(c*x^8 + a), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.00 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.09

$$\int \frac{d + ex^4}{\sqrt{a + cx^8}} dx = \frac{dx\Gamma(\frac{1}{8}) {}_2F_1\left(\begin{matrix} \frac{1}{8}, \frac{1}{2} \\ \frac{9}{8} \end{matrix} \middle| \frac{cx^8 e^{i\pi}}{a}\right)}{8\sqrt{a}\Gamma(\frac{9}{8})} + \frac{ex^5\Gamma(\frac{5}{8}) {}_2F_1\left(\begin{matrix} \frac{1}{2}, \frac{5}{8} \\ \frac{13}{8} \end{matrix} \middle| \frac{cx^8 e^{i\pi}}{a}\right)}{8\sqrt{a}\Gamma(\frac{13}{8})}$$

input `integrate((e*x**4+d)/(c*x**8+a)**(1/2),x)`

output `d*x*gamma(1/8)*hyper((1/8, 1/2), (9/8,), c*x**8*exp_polar(I*pi)/a)/(8*sqrt(a)*gamma(9/8)) + e*x**5*gamma(5/8)*hyper((1/2, 5/8), (13/8,), c*x**8*exp_polar(I*pi)/a)/(8*sqrt(a)*gamma(13/8))`

Maxima [F]

$$\int \frac{d + ex^4}{\sqrt{a + cx^8}} dx = \int \frac{ex^4 + d}{\sqrt{cx^8 + a}} dx$$

input `integrate((e*x^4+d)/(c*x^8+a)^(1/2),x, algorithm="maxima")`

output `integrate((e*x^4 + d)/sqrt(c*x^8 + a), x)`

Giac [F]

$$\int \frac{d + ex^4}{\sqrt{a + cx^8}} dx = \int \frac{ex^4 + d}{\sqrt{cx^8 + a}} dx$$

input `integrate((e*x^4+d)/(c*x^8+a)^(1/2),x, algorithm="giac")`

output `integrate((e*x^4 + d)/sqrt(c*x^8 + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{d + ex^4}{\sqrt{a + cx^8}} dx = \int \frac{e x^4 + d}{\sqrt{c x^8 + a}} dx$$

input `int((d + e*x^4)/(a + c*x^8)^(1/2),x)`

output `int((d + e*x^4)/(a + c*x^8)^(1/2), x)`

Reduce [F]

$$\int \frac{d + ex^4}{\sqrt{a + cx^8}} dx = \left(\int \frac{\sqrt{c x^8 + a}}{c x^8 + a} dx \right) d + \left(\int \frac{\sqrt{c x^8 + a} x^4}{c x^8 + a} dx \right) e$$

input `int((e*x^4+d)/(c*x^8+a)^(1/2),x)`

output `int(sqrt(a + c*x**8)/(a + c*x**8),x)*d + int((sqrt(a + c*x**8)*x**4)/(a + c*x**8),x)*e`

3.20 $\int (d + ex^4) (a - cx^8)^p \, dx$

Optimal result	188
Mathematica [A] (verified)	188
Rubi [A] (verified)	189
Maple [F]	190
Fricas [F]	190
Sympy [C] (verification not implemented)	191
Maxima [F]	191
Giac [F]	192
Mupad [F(-1)]	192
Reduce [F]	192

Optimal result

Integrand size = 18, antiderivative size = 98

$$\int (d + ex^4) (a - cx^8)^p \, dx = dx(a - cx^8)^p \left(1 - \frac{cx^8}{a}\right)^{-p} \text{Hypergeometric2F1} \left(\frac{1}{8}, -p, \frac{9}{8}, \frac{cx^8}{a}\right) + \frac{1}{5} ex^5 (a - cx^8)^p \left(1 - \frac{cx^8}{a}\right)^{-p} \text{Hypergeometric2F1} \left(\frac{5}{8}, -p, \frac{13}{8}, \frac{cx^8}{a}\right)$$

output $d*x*(-c*x^8+a)^p*\text{hypergeom}([1/8, -p], [9/8], c*x^8/a)/((1-c*x^8/a)^p)+1/5*e*x^5*(-c*x^8+a)^p*\text{hypergeom}([5/8, -p], [13/8], c*x^8/a)/((1-c*x^8/a)^p)$

Mathematica [A] (verified)

Time = 0.57 (sec), antiderivative size = 75, normalized size of antiderivative = 0.77

$$\int (d + ex^4) (a - cx^8)^p \, dx = \frac{1}{5} x (a - cx^8)^p \left(1 - \frac{cx^8}{a}\right)^{-p} \left(5d \text{Hypergeometric2F1} \left(\frac{1}{8}, -p, \frac{9}{8}, \frac{cx^8}{a}\right) + ex^4 \text{Hypergeometric2F1} \left(\frac{5}{8}, -p, \frac{13}{8}, \frac{cx^8}{a}\right)\right)$$

input $\text{Integrate}[(d + e*x^4)*(a - c*x^8)^p, x]$

output $(x*(a - c*x^8)^p*(5*d*\text{Hypergeometric2F1}[1/8, -p, 9/8, (c*x^8)/a] + e*x^4*\text{Hypergeometric2F1}[5/8, -p, 13/8, (c*x^8)/a]))/(5*(1 - (c*x^8)/a)^p)$

Rubi [A] (verified)

Time = 0.23 (sec), antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.111, Rules used = {1763, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (d + ex^4) (a - cx^8)^p dx \\ & \quad \downarrow 1763 \\ & \int (d(a - cx^8)^p + ex^4(a - cx^8)^p) dx \\ & \quad \downarrow 2009 \\ & dx(a - cx^8)^p \left(1 - \frac{cx^8}{a}\right)^{-p} \text{Hypergeometric2F1}\left(\frac{1}{8}, -p, \frac{9}{8}, \frac{cx^8}{a}\right) + \\ & \frac{1}{5}ex^5(a - cx^8)^p \left(1 - \frac{cx^8}{a}\right)^{-p} \text{Hypergeometric2F1}\left(\frac{5}{8}, -p, \frac{13}{8}, \frac{cx^8}{a}\right) \end{aligned}$$

input $\text{Int}[(d + e*x^4)*(a - c*x^8)^p, x]$

output $(d*x*(a - c*x^8)^p*\text{Hypergeometric2F1}[1/8, -p, 9/8, (c*x^8)/a])/(1 - (c*x^8)/a)^p + (e*x^5*(a - c*x^8)^p*\text{Hypergeometric2F1}[5/8, -p, 13/8, (c*x^8)/a])/(5*(1 - (c*x^8)/a)^p)$

Definitions of rubi rules used

rule 1763 $\text{Int}[(d_0 + e_0 x^{n_0}) (a_0 + c_0 x^{n_2})^p, x] \rightarrow \text{Int}[\text{ExpandIntegrand}[(d + e x^n) (a + c x^{2n})^p, x], x] /; \text{FreeQ}[\{a, c, d, e, n\}, x] \&& \text{EqQ}[n_2, 2n]$

rule 2009 $\text{Int}[u, x] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

Maple [F]

$$\int (x^4 e + d) (-c x^8 + a)^p dx$$

input `int((e*x^4+d)*(-c*x^8+a)^p,x)`

output `int((e*x^4+d)*(-c*x^8+a)^p,x)`

Fricas [F]

$$\int (d + e x^4) (a - c x^8)^p dx = \int (e x^4 + d) (-c x^8 + a)^p dx$$

input `integrate((e*x^4+d)*(-c*x^8+a)^p,x, algorithm="fricas")`

output `integral((e*x^4 + d)*(-c*x^8 + a)^p, x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 150.53 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.80

$$\int (d + ex^4) (a - cx^8)^p dx = \frac{a^p dx \Gamma\left(\frac{1}{8}\right) {}_2F_1\left(\begin{matrix} \frac{1}{8}, -p \\ \frac{9}{8} \end{matrix} \middle| \frac{cx^8 e^{2i\pi}}{a}\right)}{8\Gamma\left(\frac{9}{8}\right)} + \frac{a^p ex^5 \Gamma\left(\frac{5}{8}\right) {}_2F_1\left(\begin{matrix} \frac{5}{8}, -p \\ \frac{13}{8} \end{matrix} \middle| \frac{cx^8 e^{2i\pi}}{a}\right)}{8\Gamma\left(\frac{13}{8}\right)}$$

input `integrate((e*x**4+d)*(-c*x**8+a)**p,x)`

output `a**p*d*x*gamma(1/8)*hyper((1/8, -p), (9/8,), c*x**8*exp_polar(2*I*pi)/a)/(8*gamma(9/8)) + a**p*e*x**5*gamma(5/8)*hyper((5/8, -p), (13/8,), c*x**8*exp_polar(2*I*pi)/a)/(8*gamma(13/8))`

Maxima [F]

$$\int (d + ex^4) (a - cx^8)^p dx = \int (ex^4 + d) (-cx^8 + a)^p dx$$

input `integrate((e*x^4+d)*(-c*x^8+a)^p,x, algorithm="maxima")`

output `integrate((e*x^4 + d)*(-c*x^8 + a)^p, x)`

Giac [F]

$$\int (d + ex^4) (a - cx^8)^p dx = \int (ex^4 + d) (-cx^8 + a)^p dx$$

input `integrate((e*x^4+d)*(-c*x^8+a)^p,x, algorithm="giac")`

output `integrate((e*x^4 + d)*(-c*x^8 + a)^p, x)`

Mupad [F(-1)]

Timed out.

$$\int (d + ex^4) (a - cx^8)^p dx = \int (a - cx^8)^p (e x^4 + d) dx$$

input `int((a - c*x^8)^p*(d + e*x^4),x)`

output `int((a - c*x^8)^p*(d + e*x^4), x)`

Reduce [F]

$$\begin{aligned} & \int (d + ex^4) (a - cx^8)^p dx \\ &= \frac{8(-cx^8 + a)^p dp}{dx} + 5(-cx^8 + a)^p dx + 8(-cx^8 + a)^p ep x^5 + (-cx^8 + a)^p e x^5 + 4096 \left(\int \frac{(-64cp^2 x^8 - 48cp x^4)}{-64cp^2 x^8 - 48cp x^4} dx \right) \end{aligned}$$

input `int((e*x^4+d)*(-c*x^8+a)^p,x)`

output

```
(8*(a - c*x**8)**p*d*p*x + 5*(a - c*x**8)**p*d*x + 8*(a - c*x**8)**p*e*p*x
**5 + (a - c*x**8)**p*e*x**5 + 4096*int((a - c*x**8)**p/(64*a*p**2 + 48*a*
p + 5*a - 64*c*p**2*x**8 - 48*c*p*x**8 - 5*c*x**8),x)*a*d*p**4 + 5632*int(
(a - c*x**8)**p/(64*a*p**2 + 48*a*p + 5*a - 64*c*p**2*x**8 - 48*c*p*x**8 -
5*c*x**8),x)*a*d*p**3 + 2240*int((a - c*x**8)**p/(64*a*p**2 + 48*a*p + 5*
a - 64*c*p**2*x**8 - 48*c*p*x**8 - 5*c*x**8),x)*a*d*p**2 + 200*int((a - c*
x**8)**p/(64*a*p**2 + 48*a*p + 5*a - 64*c*p**2*x**8 - 48*c*p*x**8 - 5*c*x*
8),x)*a*d*p + 4096*int(((a - c*x**8)**p*x**4)/(64*a*p**2 + 48*a*p + 5*a -
64*c*p**2*x**8 - 48*c*p*x**8 - 5*c*x**8),x)*a*e*p**4 + 3584*int(((a - c*x
**8)**p*x**4)/(64*a*p**2 + 48*a*p + 5*a - 64*c*p**2*x**8 - 48*c*p*x**8 - 5
*c*x**8),x)*a*e*p**3 + 704*int(((a - c*x**8)**p*x**4)/(64*a*p**2 + 48*a*p +
5*a - 64*c*p**2*x**8 - 48*c*p*x**8 - 5*c*x**8),x)*a*e*p**2 + 40*int(((a -
c*x**8)**p*x**4)/(64*a*p**2 + 48*a*p + 5*a - 64*c*p**2*x**8 - 48*c*p*x**8 -
5*c*x**8),x)*a*e*p)/(64*p**2 + 48*p + 5)
```

3.21 $\int (d + ex^4) (a + cx^8)^p \, dx$

Optimal result	194
Mathematica [A] (verified)	194
Rubi [A] (verified)	195
Maple [F]	196
Fricas [F]	196
Sympy [C] (verification not implemented)	197
Maxima [F]	197
Giac [F]	198
Mupad [F(-1)]	198
Reduce [F]	198

Optimal result

Integrand size = 17, antiderivative size = 96

$$\int (d + ex^4) (a + cx^8)^p \, dx = dx(a + cx^8)^p \left(1 + \frac{cx^8}{a}\right)^{-p} \text{Hypergeometric2F1} \left(\frac{1}{8}, -p, \frac{9}{8}, -\frac{cx^8}{a}\right) + \frac{1}{5}ex^5(a + cx^8)^p \left(1 + \frac{cx^8}{a}\right)^{-p} \text{Hypergeometric2F1} \left(\frac{5}{8}, -p, \frac{13}{8}, -\frac{cx^8}{a}\right)$$

output $d*x*(c*x^8+a)^p*\text{hypergeom}([1/8, -p], [9/8], -c*x^8/a)/((1+c*x^8/a)^p)+1/5*e*x^5*(c*x^8+a)^p*\text{hypergeom}([5/8, -p], [13/8], -c*x^8/a)/((1+c*x^8/a)^p)$

Mathematica [A] (verified)

Time = 0.57 (sec), antiderivative size = 75, normalized size of antiderivative = 0.78

$$\int (d + ex^4) (a + cx^8)^p \, dx = \frac{1}{5}x(a + cx^8)^p \left(1 + \frac{cx^8}{a}\right)^{-p} \left(5d \text{Hypergeometric2F1} \left(\frac{1}{8}, -p, \frac{9}{8}, -\frac{cx^8}{a}\right) + ex^4 \text{Hypergeometric2F1} \left(\frac{5}{8}, -p, \frac{13}{8}, -\frac{cx^8}{a}\right)\right)$$

input $\text{Integrate}[(d + e*x^4)*(a + c*x^8)^p, x]$

output $(x*(a + c*x^8)^p*(5*d*\text{Hypergeometric2F1}[1/8, -p, 9/8, -((c*x^8)/a)] + e*x^4*\text{Hypergeometric2F1}[5/8, -p, 13/8, -((c*x^8)/a)]))/((5*(1 + (c*x^8)/a)^p)$

Rubi [A] (verified)

Time = 0.23 (sec), antiderivative size = 96, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {1763, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (d + ex^4) (a + cx^8)^p dx \\ & \quad \downarrow 1763 \\ & \int (d(a + cx^8)^p + ex^4(a + cx^8)^p) dx \\ & \quad \downarrow 2009 \\ & dx(a + cx^8)^p \left(\frac{cx^8}{a} + 1 \right)^{-p} \text{Hypergeometric2F1} \left(\frac{1}{8}, -p, \frac{9}{8}, -\frac{cx^8}{a} \right) + \\ & \frac{1}{5}ex^5(a + cx^8)^p \left(\frac{cx^8}{a} + 1 \right)^{-p} \text{Hypergeometric2F1} \left(\frac{5}{8}, -p, \frac{13}{8}, -\frac{cx^8}{a} \right) \end{aligned}$$

input $\text{Int}[(d + e*x^4)*(a + c*x^8)^p, x]$

output $(d*x*(a + c*x^8)^p*\text{Hypergeometric2F1}[1/8, -p, 9/8, -((c*x^8)/a)])/(1 + (c*x^8)/a)^p + (e*x^5*(a + c*x^8)^p*\text{Hypergeometric2F1}[5/8, -p, 13/8, -((c*x^8)/a)])/(5*(1 + (c*x^8)/a)^p)$

Definitions of rubi rules used

rule 1763 $\text{Int}[(d_0 + e_0 x^{n_0}) (a_0 + c_0 x^{n_2})^p, x] \rightarrow \text{Int}[\text{ExpandIntegrand}[(d + e x^n) (a + c x^{2n})^p, x], x] /; \text{FreeQ}[\{a, c, d, e, n\}, x] \& \text{EqQ}[n_2, 2n]$

rule 2009 $\text{Int}[u, x] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

Maple [F]

$$\int (x^4 e + d) (c x^8 + a)^p dx$$

input `int((e*x^4+d)*(c*x^8+a)^p,x)`

output `int((e*x^4+d)*(c*x^8+a)^p,x)`

Fricas [F]

$$\int (d + e x^4) (a + c x^8)^p dx = \int (e x^4 + d) (c x^8 + a)^p dx$$

input `integrate((e*x^4+d)*(c*x^8+a)^p,x, algorithm="fricas")`

output `integral((e*x^4 + d)*(c*x^8 + a)^p, x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 154.48 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.78

$$\int (d + ex^4) (a + cx^8)^p \, dx = \frac{a^p dx \Gamma\left(\frac{1}{8}\right) {}_2F_1\left(\begin{matrix} \frac{1}{8}, -p \\ \frac{9}{8} \end{matrix} \middle| \frac{cx^8 e^{i\pi}}{a}\right)}{8\Gamma\left(\frac{9}{8}\right)} + \frac{a^p ex^5 \Gamma\left(\frac{5}{8}\right) {}_2F_1\left(\begin{matrix} \frac{5}{8}, -p \\ \frac{13}{8} \end{matrix} \middle| \frac{cx^8 e^{i\pi}}{a}\right)}{8\Gamma\left(\frac{13}{8}\right)}$$

input `integrate((e*x**4+d)*(c*x**8+a)**p,x)`

output `a**p*d*x*gamma(1/8)*hyper((1/8, -p), (9/8,), c*x**8*exp_polar(I*pi)/a)/(8*gamma(9/8)) + a**p*e*x**5*gamma(5/8)*hyper((5/8, -p), (13/8,), c*x**8*exp_polar(I*pi)/a)/(8*gamma(13/8))`

Maxima [F]

$$\int (d + ex^4) (a + cx^8)^p \, dx = \int (ex^4 + d)(cx^8 + a)^p \, dx$$

input `integrate((e*x^4+d)*(c*x^8+a)^p,x, algorithm="maxima")`

output `integrate((e*x^4 + d)*(c*x^8 + a)^p, x)`

Giac [F]

$$\int (d + ex^4) (a + cx^8)^p \, dx = \int (ex^4 + d)(cx^8 + a)^p \, dx$$

input `integrate((e*x^4+d)*(c*x^8+a)^p,x, algorithm="giac")`

output `integrate((e*x^4 + d)*(c*x^8 + a)^p, x)`

Mupad [F(-1)]

Timed out.

$$\int (d + ex^4) (a + cx^8)^p \, dx = \int (cx^8 + a)^p (e x^4 + d) \, dx$$

input `int((a + c*x^8)^p*(d + e*x^4),x)`

output `int((a + c*x^8)^p*(d + e*x^4), x)`

Reduce [F]

$$\begin{aligned} & \int (d + ex^4) (a + cx^8)^p \, dx \\ &= \frac{8(cx^8 + a)^p \, dpx + 5(cx^8 + a)^p \, dx + 8(cx^8 + a)^p \, ep \, x^5 + (cx^8 + a)^p \, e \, x^5 + 4096 \left(\int \frac{(cx^8 + a)^p}{64cp^2x^8 + 48cp \, x^8 + 5cx^8 + 64} \, dx \right)}{64cp^2x^8 + 48cp \, x^8 + 5cx^8 + 64} \end{aligned}$$

input `int((e*x^4+d)*(c*x^8+a)^p,x)`

output

```
(8*(a + c*x**8)**p*d*p*x + 5*(a + c*x**8)**p*d*x + 8*(a + c*x**8)**p*e*p*x
**5 + (a + c*x**8)**p*e*x**5 + 4096*int((a + c*x**8)**p/(64*a*p**2 + 48*a*
p + 5*a + 64*c*p**2*x**8 + 48*c*p*x**8 + 5*c*x**8),x)*a*d*p**4 + 5632*int(
(a + c*x**8)**p/(64*a*p**2 + 48*a*p + 5*a + 64*c*p**2*x**8 + 48*c*p*x**8 +
5*c*x**8),x)*a*d*p**3 + 2240*int((a + c*x**8)**p/(64*a*p**2 + 48*a*p + 5*
a + 64*c*p**2*x**8 + 48*c*p*x**8 + 5*c*x**8),x)*a*d*p**2 + 200*int((a + c*
x**8)**p/(64*a*p**2 + 48*a*p + 5*a + 64*c*p**2*x**8 + 48*c*p*x**8 + 5*c*x*
8),x)*a*d*p + 4096*int(((a + c*x**8)**p*x**4)/(64*a*p**2 + 48*a*p + 5*a +
64*c*p**2*x**8 + 48*c*p*x**8 + 5*c*x**8),x)*a*e*p**4 + 3584*int(((a + c*x*
8)**p*x**4)/(64*a*p**2 + 48*a*p + 5*a + 64*c*p**2*x**8 + 48*c*p*x**8 + 5*
c*x**8),x)*a*e*p**3 + 704*int(((a + c*x**8)**p*x**4)/(64*a*p**2 + 48*a*p +
5*a + 64*c*p**2*x**8 + 48*c*p*x**8 + 5*c*x**8),x)*a*e*p**2 + 40*int(((a +
c*x**8)**p*x**4)/(64*a*p**2 + 48*a*p + 5*a + 64*c*p**2*x**8 + 48*c*p*x**8 +
5*c*x**8),x)*a*e*p)/(64*p**2 + 48*p + 5)
```

3.22 $\int (d + ex^4)^q (a + cx^8)^p \, dx$

Optimal result	200
Mathematica [N/A]	200
Rubi [N/A]	201
Maple [N/A]	201
Fricas [N/A]	202
Sympy [F(-1)]	202
Maxima [N/A]	202
Giac [N/A]	203
Mupad [N/A]	203
Reduce [N/A]	204

Optimal result

Integrand size = 19, antiderivative size = 19

$$\int (d + ex^4)^q (a + cx^8)^p \, dx = \text{Int}((d + ex^4)^q (a + cx^8)^p, x)$$

output `Defer(Int)((e*x^4+d)^q*(c*x^8+a)^p,x)`

Mathematica [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int (d + ex^4)^q (a + cx^8)^p \, dx = \int (d + ex^4)^q (a + cx^8)^p \, dx$$

input `Integrate[(d + e*x^4)^q*(a + c*x^8)^p,x]`

output `Integrate[(d + e*x^4)^q*(a + c*x^8)^p, x]`

Rubi [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {1770}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + cx^8)^p (d + ex^4)^q dx \xrightarrow{1770} \int (a + cx^8)^p (d + ex^4)^q dx$$

input `Int[(d + e*x^4)^q*(a + c*x^8)^p,x]`

output `$Aborted`

Definitions of rubi rules used

rule 1770 `Int[((d_) + (e_.)*(x_)^(n_))^(q_)*((a_) + (c_.)*(x_)^(n2_))^(p_), x_Symbol] :> Unintegrable[(d + e*x^n)^q*(a + c*x^(2*n))^p, x] /; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n]`

Maple [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int (x^4e + d)^q (cx^8 + a)^p dx$$

input `int((e*x^4+d)^q*(c*x^8+a)^p,x)`

output `int((e*x^4+d)^q*(c*x^8+a)^p,x)`

Fricas [N/A]

Not integrable

Time = 0.39 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int (d + ex^4)^q (a + cx^8)^p dx = \int (cx^8 + a)^p (ex^4 + d)^q dx$$

input `integrate((e*x^4+d)^q*(c*x^8+a)^p,x, algorithm="fricas")`

output `integral((c*x^8 + a)^p*(e*x^4 + d)^q, x)`

Sympy [F(-1)]

Timed out.

$$\int (d + ex^4)^q (a + cx^8)^p dx = \text{Timed out}$$

input `integrate((e*x**4+d)**q*(c*x**8+a)**p,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int (d + ex^4)^q (a + cx^8)^p dx = \int (cx^8 + a)^p (ex^4 + d)^q dx$$

input `integrate((e*x^4+d)^q*(c*x^8+a)^p,x, algorithm="maxima")`

output `integrate((c*x^8 + a)^p*(e*x^4 + d)^q, x)`

Giac [N/A]

Not integrable

Time = 0.61 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int (d + ex^4)^q (a + cx^8)^p dx = \int (cx^8 + a)^p (ex^4 + d)^q dx$$

input `integrate((e*x^4+d)^q*(c*x^8+a)^p,x, algorithm="giac")`

output `integrate((c*x^8 + a)^p*(e*x^4 + d)^q, x)`

Mupad [N/A]

Not integrable

Time = 11.17 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int (d + ex^4)^q (a + cx^8)^p dx = \int (cx^8 + a)^p (ex^4 + d)^q dx$$

input `int((a + c*x^8)^p*(d + e*x^4)^q,x)`

output `int((a + c*x^8)^p*(d + e*x^4)^q, x)`

Reduce [N/A]

Not integrable

Time = 9.24 (sec) , antiderivative size = 1249, normalized size of antiderivative = 65.74

$$\int (d + ex^4)^q (a + cx^8)^p dx = \text{Too large to display}$$

input `int((e*x^4+d)^q*(c*x^8+a)^p,x)`

output
$$\begin{aligned} & ((d + e*x**4)**q*(a + c*x**8)**p*x + 32*int(((d + e*x**4)**q*(a + c*x**8)* \\ & *p*x**8)/(8*a*d*p + 4*a*d*q + a*d + 8*a*e*p*x**4 + 4*a*e*q*x**4 + a*e*x**4 \\ & + 8*c*d*p*x**8 + 4*c*d*q*x**8 + c*d*x**8 + 8*c*e*p*x**12 + 4*c*e*q*x**12 \\ & + c*e*x**12),x)*c*d*p*q + 16*int(((d + e*x**4)**q*(a + c*x**8)**p*x**8)/(8 \\ & *a*d*p + 4*a*d*q + a*d + 8*a*e*p*x**4 + 4*a*e*q*x**4 + a*e*x**4 + 8*c*d*p* \\ & x**8 + 4*c*d*q*x**8 + c*d*x**8 + 8*c*e*p*x**12 + 4*c*e*q*x**12 + c*e*x**12 \\ &),x)*c*d*q**2 + 4*int(((d + e*x**4)**q*(a + c*x**8)**p*x**4)/(8*a*d*p + 4* \\ & a*d*q + a*d + 8*a*e*p*x**4 + 4*a*e*q*x**4 + a*e*x**4 + 8*c*d*p*x**8 + 4*c* \\ & d*q*x**8 + c*d*x**8 + 8*c*e*p*x**12 + 4*c*e*q*x**12 + c*e*x**12),x)*c*d*q \\ & + 64*int(((d + e*x**4)**q*(a + c*x**8)**p*x**4)/(8*a*d*p + 4*a*d*q + a*d + \\ & 8*a*e*p*x**4 + 4*a*e*q*x**4 + a*e*x**4 + 8*c*d*p*x**8 + 4*c*d*q*x**8 + c* \\ & d*x**8 + 8*c*e*p*x**12 + 4*c*e*q*x**12 + c*e*x**12),x)*a*e*p**2 + 32*int((\\ & (d + e*x**4)**q*(a + c*x**8)**p*x**4)/(8*a*d*p + 4*a*d*q + a*d + 8*a*e*p*x \\ & **4 + 4*a*e*q*x**4 + a*e*x**4 + 8*c*d*p*x**8 + 4*c*d*q*x**8 + c*d*x**8 + 8 \\ & *c*e*p*x**12 + 4*c*e*q*x**12 + c*e*x**12),x)*a*e*p*q + 8*int(((d + e*x**4) \\ & **q*(a + c*x**8)**p*x**4)/(8*a*d*p + 4*a*d*q + a*d + 8*a*e*p*x**4 + 4*a*e* \\ & q*x**4 + a*e*x**4 + 8*c*d*p*x**8 + 4*c*d*q*x**8 + c*d*x**8 + 8*c*e*p*x**12 \\ & + 4*c*e*q*x**12 + c*e*x**12),x)*a*e*p + 64*int(((d + e*x**4)**q*(a + c*x* \\ & *8)**p)/(8*a*d*p + 4*a*d*q + a*d + 8*a*e*p*x**4 + 4*a*e*q*x**4 + a*e*x**4 \\ & + 8*c*d*p*x**8 + 4*c*d*q*x**8 + c*d*x**8 + 8*c*e*p*x**12 + 4*c*e*q*x**1... \end{aligned}$$

3.23 $\int (1 - x^4)^p (1 - x^8)^p \, dx$

Optimal result	205
Mathematica [F]	205
Rubi [A] (verified)	206
Maple [F]	207
Fricas [F]	207
Sympy [F]	207
Maxima [F]	208
Giac [F]	208
Mupad [F(-1)]	208
Reduce [F]	209

Optimal result

Integrand size = 19, antiderivative size = 23

$$\int (1 - x^4)^p (1 - x^8)^p \, dx = x \text{AppellF1} \left(\frac{1}{4}, -2p, -p, \frac{5}{4}, x^4, -x^4 \right)$$

output `x*AppellF1(1/4, -2*p, -p, 5/4, x^4, -x^4)`

Mathematica [F]

$$\int (1 - x^4)^p (1 - x^8)^p \, dx = \int (1 - x^4)^p (1 - x^8)^p \, dx$$

input `Integrate[(1 - x^4)^p*(1 - x^8)^p, x]`

output `Integrate[(1 - x^4)^p*(1 - x^8)^p, x]`

Rubi [A] (verified)

Time = 0.16 (sec), antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {1388, 936}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (1 - x^4)^p (1 - x^8)^p dx \\ & \quad \downarrow \textcolor{blue}{1388} \\ & \int (1 - x^4)^{2p} (x^4 + 1)^p dx \\ & \quad \downarrow \textcolor{blue}{936} \\ & x \operatorname{AppellF1} \left(\frac{1}{4}, -2p, -p, \frac{5}{4}, x^4, -x^4 \right) \end{aligned}$$

input `Int[(1 - x^4)^p*(1 - x^8)^p, x]`

output `x*AppellF1[1/4, -2*p, -p, 5/4, x^4, -x^4]`

Definitions of rubi rules used

rule 936 `Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] :> Simplify[a^p*c^q*x*AppellF1[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`

rule 1388 `Int[(u_)*(a_ + (c_)*(x_)^(n2_))^(p_)*(d_ + (e_)*(x_)^(n_))^(q_), x_Symbol] :> Int[u*(d + e*x^n)^(p + q)*(a/d + (c/e)*x^n)^p, x] /; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[c*d^2 + a*e^2, 0] && (IntegerQ[p] || (GtQ[a, 0] && GtQ[d, 0]))`

Maple [F]

$$\int (-x^4 + 1)^p (-x^8 + 1)^p dx$$

input `int((-x^4+1)^p*(-x^8+1)^p,x)`

output `int((-x^4+1)^p*(-x^8+1)^p,x)`

Fricas [F]

$$\int (1 - x^4)^p (1 - x^8)^p dx = \int (-x^8 + 1)^p (-x^4 + 1)^p dx$$

input `integrate((-x^4+1)^p*(-x^8+1)^p,x, algorithm="fricas")`

output `integral((-x^8 + 1)^p*(-x^4 + 1)^p, x)`

Sympy [F]

$$\int (1 - x^4)^p (1 - x^8)^p dx = \int (-(x - 1)(x + 1)(x^2 + 1))^p (-(x - 1)(x + 1)(x^2 + 1))(x^4 + 1)^p dx$$

input `integrate((-x**4+1)**p*(-x**8+1)**p,x)`

output `Integral((-x - 1)*(x + 1)*(x**2 + 1))**p*(-(x - 1)*(x + 1)*(x**2 + 1)*(x**4 + 1))**p, x)`

Maxima [F]

$$\int (1 - x^4)^p (1 - x^8)^p \, dx = \int (-x^8 + 1)^p (-x^4 + 1)^p \, dx$$

input `integrate((-x^4+1)^p*(-x^8+1)^p,x, algorithm="maxima")`

output `integrate((-x^8 + 1)^p*(-x^4 + 1)^p, x)`

Giac [F]

$$\int (1 - x^4)^p (1 - x^8)^p \, dx = \int (-x^8 + 1)^p (-x^4 + 1)^p \, dx$$

input `integrate((-x^4+1)^p*(-x^8+1)^p,x, algorithm="giac")`

output `integrate((-x^8 + 1)^p*(-x^4 + 1)^p, x)`

Mupad [F(-1)]

Timed out.

$$\int (1 - x^4)^p (1 - x^8)^p \, dx = \int (1 - x^4)^p (1 - x^8)^p \, dx$$

input `int((1 - x^4)^p*(1 - x^8)^p,x)`

output `int((1 - x^4)^p*(1 - x^8)^p, x)`

Reduce [F]

$$\begin{aligned}
 & \int (1-x^4)^p (1-x^8)^p dx \\
 &= \frac{(-x^4+1)^p (-x^8+1)^p x - 48 \left(\int \frac{(-x^4+1)^p (-x^8+1)^p x^4}{12p x^8+x^8-12p-1} dx \right) p^2 - 4 \left(\int \frac{(-x^4+1)^p (-x^8+1)^p x^4}{12p x^8+x^8-12p-1} dx \right) p - 144 \left(\int \frac{(-x^4+1)^p (-x^8+1)^p x^4}{12p x^8+x^8-12p-1} dx \right)}{12p+1}
 \end{aligned}$$

input `int((-x^4+1)^p*(-x^8+1)^p,x)`

output `((- x**4 + 1)**p*(- x**8 + 1)**p*x - 48*int(((- x**4 + 1)**p*(- x**8 + 1)**p*x**4)/(12*p*x**8 - 12*p + x**8 - 1),x)*p**2 - 4*int(((- x**4 + 1)**p*(- x**8 + 1)**p*x**4)/(12*p*x**8 - 12*p + x**8 - 1),x)*p - 144*int(((- x**4 + 1)**p*(- x**8 + 1)**p)/(12*p*x**8 - 12*p + x**8 - 1),x)*p**2 - 12*int(((- x**4 + 1)**p*(- x**8 + 1)**p)/(12*p*x**8 - 12*p + x**8 - 1),x)*p)/(12*p + 1)`

$$\mathbf{3.24} \quad \int (d + ex^4)^q (d^2 - e^2x^8)^p dx$$

Optimal result	210
Mathematica [F]	210
Rubi [A] (verified)	211
Maple [F]	212
Fricas [F]	212
Sympy [F(-1)]	213
Maxima [F]	213
Giac [F]	213
Mupad [F(-1)]	214
Reduce [F]	214

Optimal result

Integrand size = 24, antiderivative size = 92

$$\begin{aligned} \int (d + ex^4)^q (d^2 - e^2x^8)^p dx = & x(d + ex^4)^q \left(1 - \frac{ex^4}{d}\right)^{-p} \left(1 + \frac{ex^4}{d}\right)^{-p-q} (d^2 \\ & - e^2x^8)^p \text{AppellF1} \left(\frac{1}{4}, -p, -p - q, \frac{5}{4}, \frac{ex^4}{d}, -\frac{ex^4}{d}\right) \end{aligned}$$

output $x*(e*x^4+d)^q*(1+e*x^4/d)^{(-p-q)*(-e^2*x^8+d^2)}*p*\text{AppellF1}(1/4,-p,-p-q,5/4,e*x^4/d,-e*x^4/d)/((1-e*x^4/d)^p)$

Mathematica [F]

$$\int (d + ex^4)^q (d^2 - e^2x^8)^p dx = \int (d + ex^4)^q (d^2 - e^2x^8)^p dx$$

input `Integrate[(d + e*x^4)^q*(d^2 - e^2*x^8)^p, x]`

output `Integrate[(d + e*x^4)^q*(d^2 - e^2*x^8)^p, x]`

Rubi [A] (verified)

Time = 0.26 (sec), antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {1396, 937, 937, 936}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (d^2 - e^2 x^8)^p (d + ex^4)^q dx \\
 & \quad \downarrow \textcolor{blue}{1396} \\
 & (d - ex^4)^{-p} (d + ex^4)^{-p} (d^2 - e^2 x^8)^p \int (d - ex^4)^p (ex^4 + d)^{p+q} dx \\
 & \quad \downarrow \textcolor{blue}{937} \\
 & (d + ex^4)^{-p} \left(1 - \frac{ex^4}{d}\right)^{-p} (d^2 - e^2 x^8)^p \int (ex^4 + d)^{p+q} \left(1 - \frac{ex^4}{d}\right)^p dx \\
 & \quad \downarrow \textcolor{blue}{937} \\
 & \left(1 - \frac{ex^4}{d}\right)^{-p} (d^2 - e^2 x^8)^p (d + ex^4)^q \left(\frac{ex^4}{d} + 1\right)^{-p-q} \int \left(1 - \frac{ex^4}{d}\right)^p \left(\frac{ex^4}{d} + 1\right)^{p+q} dx \\
 & \quad \downarrow \textcolor{blue}{936} \\
 & x \left(1 - \frac{ex^4}{d}\right)^{-p} (d^2 - e^2 x^8)^p (d + ex^4)^q \left(\frac{ex^4}{d} + 1\right)^{-p-q} \text{AppellF1} \left(\frac{1}{4}, -p, -p - q, \frac{5}{4}, \frac{ex^4}{d}, -\frac{ex^4}{d}\right)
 \end{aligned}$$

input `Int[(d + e*x^4)^q*(d^2 - e^2*x^8)^p,x]`

output `(x*(d + e*x^4)^q*(1 + (e*x^4)/d)^(-p - q)*(d^2 - e^2*x^8)^p*AppellF1[1/4, -p, -p - q, 5/4, (e*x^4)/d, -((e*x^4)/d)])/(1 - (e*x^4)/d)^p`

Definitions of rubi rules used

rule 936 $\text{Int}[(a_+ + b_-) \cdot (x_-)^{(n_-)} \cdot (p_-) \cdot ((c_+ + d_-) \cdot (x_-)^{(n_-)})^{(q_-)}, x_{\text{Symbol}}]$
 $\Rightarrow \text{Simp}[a^p \cdot c^q \cdot x \cdot \text{AppellF1}[1/n, -p, -q, 1 + 1/n, (-b) \cdot (x^n/a), (-d) \cdot (x^n/c)], x] /; \text{FreeQ}[\{a, b, c, d, n, p, q\}, x] \& \text{NeQ}[b*c - a*d, 0] \& \text{NeQ}[n, -1] \& (\text{IntegerQ}[p] \mid\mid \text{GtQ}[a, 0]) \& (\text{IntegerQ}[q] \mid\mid \text{GtQ}[c, 0])$

rule 937 $\text{Int}[(a_+ + b_-) \cdot (x_-)^{(n_-)} \cdot (p_-) \cdot ((c_+ + d_-) \cdot (x_-)^{(n_-)})^{(q_-)}, x_{\text{Symbol}}]$
 $\Rightarrow \text{Simp}[a^{\text{IntPart}[p]} \cdot ((a + b \cdot x^n)^{\text{FracPart}[p]} / (1 + b \cdot (x^n/a))^{\text{FracPart}[p]}) \cdot \text{Int}[(1 + b \cdot (x^n/a))^{p_*} \cdot (c + d \cdot x^n)^q, x], x] /; \text{FreeQ}[\{a, b, c, d, n, p, q\}, x] \& \text{NeQ}[b*c - a*d, 0] \& \text{NeQ}[n, -1] \& !(\text{IntegerQ}[p] \mid\mid \text{GtQ}[a, 0])$

rule 1396 $\text{Int}[(u_-) \cdot ((a_+ + c_-) \cdot (x_-)^{(n2_-)})^{(p_-)} \cdot ((d_+ + e_-) \cdot (x_-)^{(n_-)})^{(q_-)}, x_{\text{Symbol}}]$
 $\Rightarrow \text{Simp}[(a + c \cdot x^{(2*n)})^{\text{FracPart}[p]} / ((d + e \cdot x^n)^{\text{FracPart}[p]} \cdot (a/d + c \cdot (x^n/e))^{\text{FracPart}[p]}) \cdot \text{Int}[u \cdot (d + e \cdot x^n)^{(p+q)} \cdot (a/d + (c/e) \cdot x^n)^p, x], x] /; \text{FreeQ}[\{a, c, d, e, n, p, q\}, x] \& \text{EqQ}[n2, 2*n] \& \text{EqQ}[c*d^2 + a*e^2, 0] \& !\text{IntegerQ}[p] \& !(\text{EqQ}[q, 1] \& \text{EqQ}[n, 2])$

Maple [F]

$$\int (x^4 e + d)^q (-e^2 x^8 + d^2)^p dx$$

input `int((e*x^4+d)^q*(-e^2*x^8+d^2)^p,x)`

output `int((e*x^4+d)^q*(-e^2*x^8+d^2)^p,x)`

Fricas [F]

$$\int (d + e x^4)^q (d^2 - e^2 x^8)^p dx = \int (-e^2 x^8 + d^2)^p (e x^4 + d)^q dx$$

input `integrate((e*x^4+d)^q*(-e^2*x^8+d^2)^p,x, algorithm="fricas")`

output `integral((-e^2*x^8 + d^2)^p*(e*x^4 + d)^q, x)`

Sympy [F(-1)]

Timed out.

$$\int (d + ex^4)^q (d^2 - e^2x^8)^p dx = \text{Timed out}$$

input `integrate((e*x**4+d)**q*(-e**2*x**8+d**2)**p,x)`

output `Timed out`

Maxima [F]

$$\int (d + ex^4)^q (d^2 - e^2x^8)^p dx = \int (-e^2x^8 + d^2)^p (ex^4 + d)^q dx$$

input `integrate((e*x^4+d)^q*(-e^2*x^8+d^2)^p,x, algorithm="maxima")`

output `integrate((-e^2*x^8 + d^2)^p*(e*x^4 + d)^q, x)`

Giac [F]

$$\int (d + ex^4)^q (d^2 - e^2x^8)^p dx = \int (-e^2x^8 + d^2)^p (ex^4 + d)^q dx$$

input `integrate((e*x^4+d)^q*(-e^2*x^8+d^2)^p,x, algorithm="giac")`

output `integrate((-e^2*x^8 + d^2)^p*(e*x^4 + d)^q, x)`

Mupad [F(-1)]

Timed out.

$$\int (d + ex^4)^q (d^2 - e^2 x^8)^p dx = \int (d^2 - e^2 x^8)^p (e x^4 + d)^q dx$$

input `int((d^2 - e^2*x^8)^p*(d + e*x^4)^q,x)`

output `int((d^2 - e^2*x^8)^p*(d + e*x^4)^q, x)`

Reduce [F]

$$\begin{aligned} & \int (d + ex^4)^q (d^2 - e^2 x^8)^p dx \\ &= \frac{(e x^4 + d)^q (-e^2 x^8 + d^2)^p x - 32 \left(\int \frac{(e x^4 + d)^q (-e^2 x^8 + d^2)^p x^4}{-8e^2 p x^8 - 4e^2 q x^8 - e^2 x^8 + 8d^2 p + 4d^2 q + d^2} dx \right) depq - 16 \left(\int \frac{(e x^4 + d)^q (-e^2 x^8 + d^2)^p}{-8e^2 p x^8 - 4e^2 q x^8 - e^2 x^8} dx \right) depq}{\text{continued fraction}} \end{aligned}$$

input `int((e*x^4+d)^q*(-e^2*x^8+d^2)^p,x)`

output `((d + e*x**4)**q*(d**2 - e**2*x**8)**p*x - 32*int(((d + e*x**4)**q*(d**2 - e**2*x**8)**p*x**4)/(8*d**2*p + 4*d**2*q + d**2 - 8*e**2*p*x**8 - 4*e**2*q*x**8 - e**2*x**8),x)*d*e*p*q - 16*int(((d + e*x**4)**q*(d**2 - e**2*x**8)**p*x**4)/(8*d**2*p + 4*d**2*q + d**2 - 8*e**2*p*x**8 - 4*e**2*q*x**8 - e**2*x**8),x)*d*e*p*q - 4*int(((d + e*x**4)**q*(d**2 - e**2*x**8)**p*x**4)/(8*d**2*p + 4*d**2*q + d**2 - 8*e**2*p*x**8 - 4*e**2*q*x**8 - e**2*x**8),x)*d*e*p*q + 64*int(((d + e*x**4)**q*(d**2 - e**2*x**8)**p*x**4)/(8*d**2*p + 4*d**2*q + d**2 - 8*e**2*p*x**8 - 4*e**2*q*x**8 - e**2*x**8),x)*d*e*p*q + 64*int(((d + e*x**4)**q*(d**2 - e**2*x**8)**p*x**4)/(8*d**2*p + 4*d**2*q + d**2 - 8*e**2*p*x**8 - 4*e**2*q*x**8 - e**2*x**8),x)*d*e*p*q + 8*int(((d + e*x**4)**q*(d**2 - e**2*x**8)**p*x**4)/(8*d**2*p + 4*d**2*q + d**2 - 8*e**2*p*x**8 - 4*e**2*q*x**8 - e**2*x**8),x)*d*e*p*q + 16*int(((d + e*x**4)**q*(d**2 - e**2*x**8)**p*x**4)/(8*d**2*p + 4*d**2*q + d**2 - 8*e**2*p*x**8 - 4*e**2*q*x**8 - e**2*x**8),x)*d*e*p*q + 4*int(((d + e*x**4)**q*(d**2 - e**2*x**8)**p*x**4)/(8*d**2*p + 4*d**2*q + d**2 - 8*e**2*p*x**8 - 4*e**2*q*x**8 - e**2*x**8),x)*d*e*p*q)/((8*p + 4*q + 1))`

3.25 $\int (1 - x^4)^{-p} (1 - x^8)^p \, dx$

Optimal result	215
Mathematica [F]	215
Rubi [A] (verified)	216
Maple [F]	217
Fricas [F]	217
Sympy [F(-1)]	217
Maxima [F]	218
Giac [F]	218
Mupad [F(-1)]	218
Reduce [F]	219

Optimal result

Integrand size = 21, antiderivative size = 17

$$\int (1 - x^4)^{-p} (1 - x^8)^p \, dx = x \text{Hypergeometric2F1}\left(\frac{1}{4}, -p, \frac{5}{4}, -x^4\right)$$

output `x*hypergeom([1/4, -p], [5/4], -x^4)`

Mathematica [F]

$$\int (1 - x^4)^{-p} (1 - x^8)^p \, dx = \int (1 - x^4)^{-p} (1 - x^8)^p \, dx$$

input `Integrate[(1 - x^8)^p/(1 - x^4)^p, x]`

output `Integrate[(1 - x^8)^p/(1 - x^4)^p, x]`

Rubi [A] (verified)

Time = 0.15 (sec), antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {1386, 778}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (1 - x^4)^{-p} (1 - x^8)^p dx \\ & \quad \downarrow \textcolor{blue}{1386} \\ & \int (x^4 + 1)^p dx \\ & \quad \downarrow \textcolor{blue}{778} \\ & x \operatorname{Hypergeometric2F1} \left(\frac{1}{4}, -p, \frac{5}{4}, -x^4 \right) \end{aligned}$$

input `Int[(1 - x^8)^p/(1 - x^4)^p, x]`

output `x*Hypergeometric2F1[1/4, -p, 5/4, -x^4]`

Definitions of rubi rules used

rule 778 `Int[((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F1[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] || GtQ[a, 0])`

rule 1386 `Int[(u_)*(a_) + (c_)*(x_)^(n2_))^(p_)*((d_) + (e_)*(x_)^(n_))^(q_), x_Symbol] := Simp[(-e^2/c)^q Int[u*(d - e*x^n)^p, x], x] /; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[c*d^2 + a*e^2, 0] && EqQ[p + q, 0] && GtQ[d, 0] && LtQ[c, 0] && GtQ[e^2, 0]`

Maple [F]

$$\int (-x^8 + 1)^p (-x^4 + 1)^{-p} dx$$

input `int((-x^8+1)^p/((-x^4+1)^p),x)`

output `int((-x^8+1)^p/((-x^4+1)^p),x)`

Fricas [F]

$$\int (1 - x^4)^{-p} (1 - x^8)^p dx = \int \frac{(-x^8 + 1)^p}{(-x^4 + 1)^p} dx$$

input `integrate((-x^8+1)^p/((-x^4+1)^p),x, algorithm="fricas")`

output `integral((-x^8 + 1)^p/(-x^4 + 1)^p, x)`

Sympy [F(-1)]

Timed out.

$$\int (1 - x^4)^{-p} (1 - x^8)^p dx = \text{Timed out}$$

input `integrate((-x**8+1)**p/((-x**4+1)**p),x)`

output `Timed out`

Maxima [F]

$$\int (1 - x^4)^{-p} (1 - x^8)^p \, dx = \int \frac{(-x^8 + 1)^p}{(-x^4 + 1)^p} \, dx$$

input `integrate((-x^8+1)^p/((-x^4+1)^p),x, algorithm="maxima")`

output `integrate((-x^8 + 1)^p/(-x^4 + 1)^p, x)`

Giac [F]

$$\int (1 - x^4)^{-p} (1 - x^8)^p \, dx = \int \frac{(-x^8 + 1)^p}{(-x^4 + 1)^p} \, dx$$

input `integrate((-x^8+1)^p/((-x^4+1)^p),x, algorithm="giac")`

output `integrate((-x^8 + 1)^p/(-x^4 + 1)^p, x)`

Mupad [F(-1)]

Timed out.

$$\int (1 - x^4)^{-p} (1 - x^8)^p \, dx = \int \frac{(1 - x^8)^p}{(1 - x^4)^p} \, dx$$

input `int((1 - x^8)^p/(1 - x^4)^p,x)`

output `int((1 - x^8)^p/(1 - x^4)^p, x)`

Reduce [F]

$$\int (1 - x^4)^{-p} (1 - x^8)^p \, dx = \int \frac{(-x^8 + 1)^p}{(-x^4 + 1)^p} dx$$

input `int((-x^8+1)^p/((-x^4+1)^p),x)`

output `int((- x**8 + 1)**p / (- x**4 + 1)**p, x)`

3.26 $\int \frac{(d+ex^n)^3}{a+cx^{2n}} dx$

Optimal result	220
Mathematica [A] (verified)	221
Rubi [A] (verified)	221
Maple [F]	222
Fricas [F]	223
Sympy [C] (verification not implemented)	223
Maxima [F]	224
Giac [F]	224
Mupad [F(-1)]	225
Reduce [F]	225

Optimal result

Integrand size = 21, antiderivative size = 141

$$\begin{aligned} \int \frac{(d+ex^n)^3}{a+cx^{2n}} dx &= \frac{3de^2x}{c} + \frac{e^3x^{1+n}}{c(1+n)} \\ &+ \frac{d(cd^2 - 3ae^2)x \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{ac} \\ &+ \frac{e(3cd^2 - ae^2)x^{1+n} \operatorname{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{ac(1+n)} \end{aligned}$$

output $3*d*e^2*x/c + e^3*x^(1+n)/c/(1+n) + d*(-3*a*e^2 + c*d^2)*x*hypergeom([1, 1/2/n], [1+1/2/n], -c*x^(2*n)/a)/a + c + e*(-a*e^2 + 3*c*d^2)*x^(1+n)*hypergeom([1, 1/2*(1+n)/n], [3/2+1/2/n], -c*x^(2*n)/a)/a + c/(1+n)$

Mathematica [A] (verified)

Time = 0.74 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.90

$$\int \frac{(d + ex^n)^3}{a + cx^{2n}} dx \\ = \frac{x \left(d(cd^2 - 3ae^2)(1+n) \text{Hypergeometric2F1} \left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a} \right) + e \left(ae(3d(1+n) + ex^n) + (3cd^2 - 3ae^2)n \right) \right)}{ac(1+n)}$$

input `Integrate[(d + e*x^n)^3/(a + c*x^(2*n)), x]`

output $(x*(d*(c*d^2 - 3*a*e^2)*(1 + n)*\text{Hypergeometric2F1}[1, 1/(2*n), (2 + n^{-1})/2, -((c*x^{(2*n)})/a)] + e*(a*e*(3*d*(1 + n) + e*x^{(2*n)}) + (3*c*d^2 - a*e^2)*x^{(2*n)}*\text{Hypergeometric2F1}[1, (1 + n)/(2*n), (3 + n^{-1})/2, -((c*x^{(2*n)})/a)]))/(a*c*(1 + n))$

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {1755, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^n)^3}{a + cx^{2n}} dx \\ \downarrow \textcolor{blue}{1755} \\ \int \left(\frac{x^n(3cd^2e - ae^3) - 3ade^2 + cd^3}{c(a + cx^{2n})} + \frac{3de^2}{c} + \frac{e^3x^n}{c} \right) dx \\ \downarrow \textcolor{blue}{2009}$$

$$\frac{ex^{n+1}(3cd^2 - ae^2) \text{Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{ac(n+1)} +$$

$$\frac{dx(cd^2 - 3ae^2) \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{ac} + \frac{3de^2x}{c} + \frac{e^3x^{n+1}}{c(n+1)}$$

input `Int[(d + e*x^n)^3/(a + c*x^(2*n)), x]`

output `(3*d*e^2*x)/c + (e^3*x^(1 + n))/(c*(1 + n)) + (d*(c*d^2 - 3*a*e^2)*x*Hypergeometric2F1[1, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)])/(a*c) + (e*(3*c*d^2 - a*e^2)*x^(1 + n)*Hypergeometric2F1[1, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)])/(a*c*(1 + n))`

Definitions of rubi rules used

rule 1755 `Int[((d_) + (e_)*(x_)^(n_))^(q_)/((a_) + (c_)*(x_)^(n2_)), x_Symbol] :> Int[ExpandIntegrand[(d + e*x^n)^q/(a + c*x^(2*n)), x], x] /; FreeQ[{a, c, d, e, n}, x] && EqQ[n2, 2*n] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[q]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [F]

$$\int \frac{(d + e x^n)^3}{a + c x^{2n}} dx$$

input `int((d+e*x^n)^3/(a+c*x^(2*n)),x)`

output `int((d+e*x^n)^3/(a+c*x^(2*n)),x)`

Fricas [F]

$$\int \frac{(d + ex^n)^3}{a + cx^{2n}} dx = \int \frac{(ex^n + d)^3}{cx^{2n} + a} dx$$

input `integrate((d+e*x^n)^3/(a+c*x^(2*n)),x, algorithm="fricas")`

output `integral((e^3*x^(3*n) + 3*d*e^2*x^(2*n) + 3*d^2*e*x^n + d^3)/(c*x^(2*n) + a), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 4.49 (sec) , antiderivative size = 466, normalized size of antiderivative = 3.30

$$\begin{aligned} \int \frac{(d + ex^n)^3}{a + cx^{2n}} dx &= \frac{a^{\frac{1}{2n}} a^{-1 - \frac{1}{2n}} d^3 x \Phi\left(\frac{cx^{2n} e^{i\pi}}{a}, 1, \frac{1}{2n}\right) \Gamma\left(\frac{1}{2n}\right)}{4n^2 \Gamma\left(1 + \frac{1}{2n}\right)} \\ &+ \frac{3a^{-\frac{5}{2} - \frac{1}{2n}} a^{\frac{3}{2} + \frac{1}{2n}} e^3 x^{3n+1} \Phi\left(\frac{cx^{2n} e^{i\pi}}{a}, 1, \frac{3}{2} + \frac{1}{2n}\right) \Gamma\left(\frac{3}{2} + \frac{1}{2n}\right)}{4n \Gamma\left(\frac{5}{2} + \frac{1}{2n}\right)} \\ &+ \frac{a^{-\frac{5}{2} - \frac{1}{2n}} a^{\frac{3}{2} + \frac{1}{2n}} e^3 x^{3n+1} \Phi\left(\frac{cx^{2n} e^{i\pi}}{a}, 1, \frac{3}{2} + \frac{1}{2n}\right) \Gamma\left(\frac{3}{2} + \frac{1}{2n}\right)}{4n^2 \Gamma\left(\frac{5}{2} + \frac{1}{2n}\right)} \\ &+ \frac{3a^{-\frac{3}{2} - \frac{1}{2n}} a^{\frac{1}{2} + \frac{1}{2n}} d^2 e x^{n+1} \Phi\left(\frac{cx^{2n} e^{i\pi}}{a}, 1, \frac{1}{2} + \frac{1}{2n}\right) \Gamma\left(\frac{1}{2} + \frac{1}{2n}\right)}{4n \Gamma\left(\frac{3}{2} + \frac{1}{2n}\right)} \\ &+ \frac{3a^{-\frac{3}{2} - \frac{1}{2n}} a^{\frac{1}{2} + \frac{1}{2n}} d^2 e x^{n+1} \Phi\left(\frac{cx^{2n} e^{i\pi}}{a}, 1, \frac{1}{2} + \frac{1}{2n}\right) \Gamma\left(\frac{1}{2} + \frac{1}{2n}\right)}{4n^2 \Gamma\left(\frac{3}{2} + \frac{1}{2n}\right)} \\ &- \frac{3a^{-\frac{1}{2n}} a^{1 + \frac{1}{2n}} c^{\frac{1}{2n}} c^{-1 - \frac{1}{2n}} d e^2 x \Phi\left(\frac{ax^{-2n} e^{i\pi}}{c}, 1, \frac{e^{i\pi}}{2n}\right) \Gamma\left(\frac{1}{2n}\right)}{4an^2 \Gamma\left(1 + \frac{1}{2n}\right)} \end{aligned}$$

input `integrate((d+e*x**n)**3/(a+c*x**2*n),x)`

output

```
a**(1/(2*n))*a**(-1 - 1/(2*n))*d**3*x*lerchphi(c*x**(2*n)*exp_polar(I*pi)/a, 1, 1/(2*n))*gamma(1/(2*n))/(4*n**2*gamma(1 + 1/(2*n))) + 3*a**(-5/2 - 1/(2*n))*a**((3/2 + 1/(2*n))*e**3*x**((3*n + 1)*lerchphi(c*x**(2*n)*exp_polar(I*pi)/a, 1, 3/2 + 1/(2*n))*gamma(3/2 + 1/(2*n))/(4*n*gamma(5/2 + 1/(2*n))) + a**(-5/2 - 1/(2*n))*a**((3/2 + 1/(2*n))*e**3*x**((3*n + 1)*lerchphi(c*x*(2*n)*exp_polar(I*pi)/a, 1, 3/2 + 1/(2*n))*gamma(3/2 + 1/(2*n))/(4*n**2*gamma(5/2 + 1/(2*n))) + 3*a**(-3/2 - 1/(2*n))*a**((1/2 + 1/(2*n))*d**2*e*x**((n + 1)*lerchphi(c*x**(2*n)*exp_polar(I*pi)/a, 1, 1/2 + 1/(2*n))*gamma(1/2 + 1/(2*n))/(4*n*gamma(3/2 + 1/(2*n))) + 3*a**(-3/2 - 1/(2*n))*a**((1/2 + 1/(2*n))*d**2*e*x**((n + 1)*lerchphi(c*x**((2*n)*exp_polar(I*pi)/a, 1, 1/2 + 1/(2*n))*gamma(1/2 + 1/(2*n))/(4*n**2*gamma(3/2 + 1/(2*n))) - 3*a**((1 + 1/(2*n))*c**((1/(2*n))*c**((-1 - 1/(2*n))*d**2*x*lerchphi(a*exp_polar(I*pi)/(c*x**((2*n)), 1, exp_polar(I*pi)/(2*n))*gamma(1/(2*n))/(4*a*a**((1/(2*n))*n**2*gamma(1 + 1/(2*n))))
```

Maxima [F]

$$\int \frac{(d + ex^n)^3}{a + cx^{2n}} dx = \int \frac{(ex^n + d)^3}{cx^{2n} + a} dx$$

input

```
integrate((d+e*x^n)^3/(a+c*x^(2*n)),x, algorithm="maxima")
```

output

```
(3*d*e^(2*(n + 1))*x + e^(3*x*x^n)/(c*(n + 1)) - integrate(-(c*d^3 - 3*a*d*e^2 + (3*c*d^2*e - a*e^3)*x^n)/(c^2*x^(2*n) + a*c), x)
```

Giac [F]

$$\int \frac{(d + ex^n)^3}{a + cx^{2n}} dx = \int \frac{(ex^n + d)^3}{cx^{2n} + a} dx$$

input

```
integrate((d+e*x^n)^3/(a+c*x^(2*n)),x, algorithm="giac")
```

output

```
integrate((e*x^n + d)^3/(c*x^(2*n) + a), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(d + e x^n)^3}{a + c x^{2n}} dx = \int \frac{(d + e x^n)^3}{a + c x^{2n}} dx$$

input `int((d + e*x^n)^3/(a + c*x^(2*n)),x)`

output `int((d + e*x^n)^3/(a + c*x^(2*n)), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{(d + e x^n)^3}{a + c x^{2n}} dx \\ &= \frac{x^n e^3 x - \left(\int \frac{x^n}{x^{2n} c + a} dx \right) a e^3 n - \left(\int \frac{x^n}{x^{2n} c + a} dx \right) a e^3 + 3 \left(\int \frac{x^n}{x^{2n} c + a} dx \right) c d^2 e n + 3 \left(\int \frac{x^n}{x^{2n} c + a} dx \right) c d^2 e - 3 \left(\int \frac{1}{x^{2n} c + a} dx \right)}{c(n+1)} \end{aligned}$$

input `int((d+e*x^n)^3/(a+c*x^(2*n)),x)`

output `(x**n*e**3*x - int(x**n/(x**(2*n)*c + a),x)*a*e**3*n - int(x**n/(x**(2*n)*c + a),x)*a*e**3 + 3*int(x**n/(x**(2*n)*c + a),x)*c*d**2*e*n + 3*int(x**n/(x**(2*n)*c + a),x)*c*d**2*e - 3*int(1/(x**(2*n)*c + a),x)*a*d*e**2*n - 3*int(1/(x**(2*n)*c + a),x)*a*d*e**2 + int(1/(x**(2*n)*c + a),x)*c*d**3*n + int(1/(x**(2*n)*c + a),x)*c*d**3 + 3*d*e**2*n*x + 3*d*e**2*x)/(c*(n + 1))`

$$3.27 \quad \int \frac{(d+ex^n)^2}{a+cx^{2n}} dx$$

Optimal result	226
Mathematica [A] (verified)	226
Rubi [A] (verified)	227
Maple [F]	228
Fricas [F]	228
Sympy [C] (verification not implemented)	229
Maxima [F]	229
Giac [F]	230
Mupad [F(-1)]	230
Reduce [F]	230

Optimal result

Integrand size = 21, antiderivative size = 107

$$\begin{aligned} \int \frac{(d+ex^n)^2}{a+cx^{2n}} dx = & \frac{e^2 x}{c} + \frac{(cd^2 - ae^2) x \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{ac} \\ & + \frac{2dex^{1+n} \text{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a(1+n)} \end{aligned}$$

output $e^{2*x}/c + (-a*e^{2*c*d^2})*x*hypergeom([1, 1/2/n], [1+1/2/n], -c*x^(2*n)/a)/a/c + 2*d*e*x^(1+n)*hypergeom([1, 1/2*(1+n)/n], [3/2+1/2/n], -c*x^(2*n)/a)/a/(1+n)$

Mathematica [A] (verified)

Time = 0.30 (sec), antiderivative size = 107, normalized size of antiderivative = 1.00

$$\begin{aligned} \int \frac{(d+ex^n)^2}{a+cx^{2n}} dx = & \frac{e^2 x}{c} + \frac{(cd^2 - ae^2) x \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{ac} \\ & + \frac{2dex^{1+n} \text{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a(1+n)} \end{aligned}$$

input `Integrate[(d + e*x^n)^2/(a + c*x^(2*n)), x]`

output $(e^{2x})/c + ((c*d^2 - a*e^2)*x*Hypergeometric2F1[1, 1/(2*n), (2 + n^{-1})/2, -((c*x^{(2*n)})/a)])/(a*c) + (2*d*e*x^{(1 + n)}*Hypergeometric2F1[1, (1 + n)/(2*n), (3 + n^{-1})/2, -((c*x^{(2*n)})/a)])/(a*(1 + n))$

Rubi [A] (verified)

Time = 0.28 (sec), antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {1755, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(d + ex^n)^2}{a + cx^{2n}} dx \\ & \quad \downarrow \textcolor{blue}{1755} \\ & \int \left(\frac{-ae^2 + cd^2 + 2cdex^n}{c(a + cx^{2n})} + \frac{e^2}{c} \right) dx \\ & \quad \downarrow \textcolor{blue}{2009} \\ & \frac{x(cd^2 - ae^2) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{ac} + \\ & \quad \frac{2dex^{n+1} \operatorname{Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a(n+1)} + \frac{e^2 x}{c} \end{aligned}$$

input $\operatorname{Int}[(d + e*x^n)^2/(a + c*x^{(2*n)}), x]$

output $(e^{2x})/c + ((c*d^2 - a*e^2)*x*Hypergeometric2F1[1, 1/(2*n), (2 + n^{-1})/2, -((c*x^{(2*n)})/a)])/(a*c) + (2*d*e*x^{(1 + n)}*Hypergeometric2F1[1, (1 + n)/(2*n), (3 + n^{-1})/2, -((c*x^{(2*n)})/a)])/(a*(1 + n))$

Definitions of rubi rules used

rule 1755 $\text{Int}[(d_+ + e_*)*(x_-)^n]^q / ((a_+ + c_*)*(x_-)^{n2_+}), x_{\text{Symbol}}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(d + e*x^n)^q / (a + c*x^{2n}), x], x] /; \text{FreeQ}[\{a, c, d, e, n\}, x] \&& \text{EqQ}[n2, 2n] \&& \text{NeQ}[c*d^2 + a*e^2, 0] \&& \text{IntegerQ}[q]$

rule 2009 $\text{Int}[u_-, x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

Maple [F]

$$\int \frac{(d + e x^n)^2}{a + c x^{2n}} dx$$

input $\text{int}((d+e*x^n)^2/(a+c*x^(2*n)), x)$

output $\text{int}((d+e*x^n)^2/(a+c*x^(2*n)), x)$

Fricas [F]

$$\int \frac{(d + e x^n)^2}{a + c x^{2n}} dx = \int \frac{(e x^n + d)^2}{c x^{2n} + a} dx$$

input $\text{integrate}((d+e*x^n)^2/(a+c*x^(2*n)), x, \text{algorithm}=\text{"fricas"})$

output $\text{integral}((e^{2*x^(2*n)} + 2*d*e*x^n + d^2)/(c*x^(2*n) + a), x)$

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 3.10 (sec) , antiderivative size = 296, normalized size of antiderivative = 2.77

$$\int \frac{(d + ex^n)^2}{a + cx^{2n}} dx = \frac{a^{\frac{1}{2n}} a^{-1 - \frac{1}{2n}} d^2 x \Phi\left(\frac{cx^{2n} e^{i\pi}}{a}, 1, \frac{1}{2n}\right) \Gamma\left(\frac{1}{2n}\right)}{4n^2 \Gamma\left(1 + \frac{1}{2n}\right)}$$

$$+ \frac{a^{-\frac{3}{2} - \frac{1}{2n}} a^{\frac{1}{2} + \frac{1}{2n}} d e x^{n+1} \Phi\left(\frac{cx^{2n} e^{i\pi}}{a}, 1, \frac{1}{2} + \frac{1}{2n}\right) \Gamma\left(\frac{1}{2} + \frac{1}{2n}\right)}{2n \Gamma\left(\frac{3}{2} + \frac{1}{2n}\right)}$$

$$+ \frac{a^{-\frac{3}{2} - \frac{1}{2n}} a^{\frac{1}{2} + \frac{1}{2n}} d e x^{n+1} \Phi\left(\frac{cx^{2n} e^{i\pi}}{a}, 1, \frac{1}{2} + \frac{1}{2n}\right) \Gamma\left(\frac{1}{2} + \frac{1}{2n}\right)}{2n^2 \Gamma\left(\frac{3}{2} + \frac{1}{2n}\right)}$$

$$- \frac{a^{-\frac{1}{2n}} a^{1 + \frac{1}{2n}} c^{\frac{1}{2n}} c^{-1 - \frac{1}{2n}} e^2 x \Phi\left(\frac{ax^{-2n} e^{i\pi}}{c}, 1, \frac{e^{i\pi}}{2n}\right) \Gamma\left(\frac{1}{2n}\right)}{4an^2 \Gamma\left(1 + \frac{1}{2n}\right)}$$

input `integrate((d+e*x**n)**2/(a+c*x**(2*n)),x)`

output `a**(-(1/(2*n)))*a**(-1 - 1/(2*n))*d**2*x*lerchphi(c*x**2*n)*exp_polar(I*pi)/a, 1, 1/(2*n))*gamma(1/(2*n))/(4*n**2*gamma(1 + 1/(2*n))) + a**(-3/2 - 1/(2*n))*a**1/2 + 1/(2*n))*d*e*x**2*n*lerchphi(c*x**2*n)*exp_polar(I*pi)/a, 1, 1/2 + 1/(2*n))*gamma(1/2 + 1/(2*n))/(2*n*gamma(3/2 + 1/(2*n))) + a**(-3/2 - 1/(2*n))*a**1/2 + 1/(2*n))*d*e*x**2*n*lerchphi(c*x**2*n)*exp_polar(I*pi)/a, 1, 1/2 + 1/(2*n))*gamma(1/2 + 1/(2*n))/(2*n**2*gamma(3/2 + 1/(2*n))) - a**1 + 1/(2*n))*c**1/(2*n))*c**(-1 - 1/(2*n))*e**2*x*lerchphi(a*exp_polar(I*pi)/(c*x**2*n), 1, exp_polar(I*pi)/(2*n))*gamma(1/(2*n))/(4*a*a**1/(2*n))*n**2*gamma(1 + 1/(2*n)))`

Maxima [F]

$$\int \frac{(d + ex^n)^2}{a + cx^{2n}} dx = \int \frac{(ex^n + d)^2}{cx^{2n} + a} dx$$

input `integrate((d+e*x^n)^2/(a+c*x^(2*n)),x, algorithm="maxima")`

output $e^{2x}/c + \text{integrate}((2*c*d*e*x^n + c*d^2 - a*e^2)/(c^2*x^(2*n) + a*c), x)$

Giac [F]

$$\int \frac{(d + ex^n)^2}{a + cx^{2n}} dx = \int \frac{(ex^n + d)^2}{cx^{2n} + a} dx$$

input `integrate((d+e*x^n)^2/(a+c*x^(2*n)),x, algorithm="giac")`

output $\text{integrate}((e*x^n + d)^2/(c*x^(2*n) + a), x)$

Mupad [F(-1)]

Timed out.

$$\int \frac{(d + ex^n)^2}{a + cx^{2n}} dx = \int \frac{(d + e x^n)^2}{a + c x^{2n}} dx$$

input `int((d + e*x^n)^2/(a + c*x^(2*n)),x)`

output $\text{int}((d + e*x^n)^2/(a + c*x^(2*n)), x)$

Reduce [F]

$$\int \frac{(d + ex^n)^2}{a + cx^{2n}} dx = \left(\int \frac{x^{2n}}{x^{2n}c + a} dx \right) e^2 + 2 \left(\int \frac{x^n}{x^{2n}c + a} dx \right) de + \left(\int \frac{1}{x^{2n}c + a} dx \right) d^2$$

input `int((d+e*x^n)^2/(a+c*x^(2*n)),x)`

output $\text{int}(x^{**(2*n)}/(x^{**2*n}*c + a), x)*e**2 + 2*\text{int}(x**n/(x^{**2*n}*c + a), x)*d*e + \text{int}(1/(x^{**2*n}*c + a), x)*d**2$

3.28 $\int \frac{d+ex^n}{a+cx^{2n}} dx$

Optimal result	231
Mathematica [A] (verified)	231
Rubi [A] (verified)	232
Maple [F]	233
Fricas [F]	233
Sympy [C] (verification not implemented)	234
Maxima [F]	234
Giac [F]	235
Mupad [F(-1)]	235
Reduce [F]	235

Optimal result

Integrand size = 19, antiderivative size = 83

$$\int \frac{d+ex^n}{a+cx^{2n}} dx = \frac{dx \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a} + \frac{ex^{1+n} \text{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a(1+n)}$$

output $d*x*\text{hypergeom}([1, 1/2/n], [1+1/2/n], -c*x^(2*n)/a)/a+e*x^(1+n)*\text{hypergeom}([1, 1/2*(1+n)/n], [3/2+1/2/n], -c*x^(2*n)/a)/a/(1+n)$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.00

$$\int \frac{d+ex^n}{a+cx^{2n}} dx = \frac{dx \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a} + \frac{ex^{1+n} \text{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a(1+n)}$$

input `Integrate[(d + e*x^n)/(a + c*x^(2*n)), x]`

output
$$(d*x*Hypergeometric2F1[1, 1/(2*n), (2 + n^{-1})/2, -((c*x^{(2*n)})/a)])/a + (e*x^{(1 + n)}*Hypergeometric2F1[1, (1 + n)/(2*n), (3 + n^{-1})/2, -((c*x^{(2*n)})/a)])/(a*(1 + n))$$

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {1748, 778, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{d + ex^n}{a + cx^{2n}} dx \\ & \quad \downarrow \textcolor{blue}{1748} \\ & d \int \frac{1}{cx^{2n} + a} dx + e \int \frac{x^n}{cx^{2n} + a} dx \\ & \quad \downarrow \textcolor{blue}{778} \\ & e \int \frac{x^n}{cx^{2n} + a} dx + \frac{dx \text{ Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a} \\ & \quad \downarrow \textcolor{blue}{888} \\ & \frac{dx \text{ Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a} + \\ & \frac{ex^{n+1} \text{ Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a(n+1)} \end{aligned}$$

input $\text{Int}[(d + e*x^n)/(a + c*x^{(2*n)}), x]$

output
$$(d*x*Hypergeometric2F1[1, 1/(2*n), (2 + n^{-1})/2, -((c*x^{(2*n)})/a)])/a + (e*x^{(1 + n)}*Hypergeometric2F1[1, (1 + n)/(2*n), (3 + n^{-1})/2, -((c*x^{(2*n)})/a)])/(a*(1 + n))$$

Definitions of rubi rules used

rule 778 $\text{Int}[(a_.) + (b_.)*(x_.)^{(n_.)}^{(p_.)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[a^p * x * \text{Hypergeometric2F1}[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; \text{FreeQ}[\{a, b, n, p\}, x] \&& \text{!IGtQ}[p, 0] \&& \text{!IntegerQ}[1/n] \&& \text{!ILtQ}[\text{Simplify}[1/n + p], 0] \&& (\text{IntegerQ}[p] \mid \text{GtQ}[a, 0])$

rule 888 $\text{Int}[((c_.)*(x_.))^{(m_.)}*((a_.) + (b_.)*(x_.)^{(n_.)}^{(p_.)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[a^p * ((c*x)^(m + 1)/(c*(m + 1))) * \text{Hypergeometric2F1}[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; \text{FreeQ}[\{a, b, c, m, n, p\}, x] \&& \text{!IGtQ}[p, 0] \&& (\text{ILtQ}[p, 0] \mid \text{GtQ}[a, 0])$

rule 1748 $\text{Int}[((d_.) + (e_.)*(x_.)^{(n_.)})/((a_.) + (c_.)*(x_.)^{(n2_.)}), x_{\text{Symbol}}] \rightarrow \text{Simp}[d \text{Int}[1/(a + c*x^(2*n)), x], x] + \text{Simp}[e \text{Int}[x^n/(a + c*x^(2*n)), x], x] /; \text{FreeQ}[\{a, c, d, e, n\}, x] \&& \text{EqQ}[n2, 2*n] \&& \text{NeQ}[c*d^2 + a*e^2, 0] \&& (\text{PosQ}[a*c] \mid \text{!IntegerQ}[n])$

Maple [F]

$$\int \frac{d + e x^n}{a + c x^{2n}} dx$$

input $\text{int}((d+e*x^n)/(a+c*x^(2*n)), x)$

output $\text{int}((d+e*x^n)/(a+c*x^(2*n)), x)$

Fricas [F]

$$\int \frac{d + e x^n}{a + c x^{2n}} dx = \int \frac{e x^n + d}{c x^{2n} + a} dx$$

input $\text{integrate}((d+e*x^n)/(a+c*x^(2*n)), x, \text{algorithm}=\text{"fricas"})$

output `integral((e*x^n + d)/(c*x^(2*n) + a), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 2.37 (sec) , antiderivative size = 209, normalized size of antiderivative = 2.52

$$\int \frac{d + ex^n}{a + cx^{2n}} dx = \frac{a^{\frac{1}{2n}} a^{-1 - \frac{1}{2n}} dx \Phi\left(\frac{cx^{2n} e^{i\pi}}{a}, 1, \frac{1}{2n}\right) \Gamma\left(\frac{1}{2n}\right)}{4n^2 \Gamma\left(1 + \frac{1}{2n}\right)}$$

$$+ \frac{a^{-\frac{3}{2} - \frac{1}{2n}} a^{\frac{1}{2} + \frac{1}{2n}} ex^{n+1} \Phi\left(\frac{cx^{2n} e^{i\pi}}{a}, 1, \frac{1}{2} + \frac{1}{2n}\right) \Gamma\left(\frac{1}{2} + \frac{1}{2n}\right)}{4n \Gamma\left(\frac{3}{2} + \frac{1}{2n}\right)}$$

$$+ \frac{a^{-\frac{3}{2} - \frac{1}{2n}} a^{\frac{1}{2} + \frac{1}{2n}} ex^{n+1} \Phi\left(\frac{cx^{2n} e^{i\pi}}{a}, 1, \frac{1}{2} + \frac{1}{2n}\right) \Gamma\left(\frac{1}{2} + \frac{1}{2n}\right)}{4n^2 \Gamma\left(\frac{3}{2} + \frac{1}{2n}\right)}$$

input `integrate((d+e*x**n)/(a+c*x**(2*n)),x)`

output `a**((1/(2*n))*a**(-1 - 1/(2*n))*d*x*lerchphi(c*x**2*n)*exp_polar(I*pi)/a, 1, 1/(2*n))*gamma(1/(2*n))/(4*n**2*gamma(1 + 1/(2*n))) + a**(-3/2 - 1/(2*n))*a**((1/2 + 1/(2*n))*e*x**2*n)*lerchphi(c*x**2*n)*exp_polar(I*pi)/a, 1, 1/2 + 1/(2*n))*gamma(1/2 + 1/(2*n))/(4*n*gamma(3/2 + 1/(2*n))) + a**(-3/2 - 1/(2*n))*a**((1/2 + 1/(2*n))*e*x**2*n)*lerchphi(c*x**2*n)*exp_polar(I*pi)/a, 1, 1/2 + 1/(2*n))*gamma(1/2 + 1/(2*n))/(4*n**2*gamma(3/2 + 1/(2*n)))`

Maxima [F]

$$\int \frac{d + ex^n}{a + cx^{2n}} dx = \int \frac{ex^n + d}{cx^{2n} + a} dx$$

input `integrate((d+e*x^n)/(a+c*x^(2*n)),x, algorithm="maxima")`

output `integrate((e*x^n + d)/(c*x^(2*n) + a), x)`

Giac [F]

$$\int \frac{d + ex^n}{a + cx^{2n}} dx = \int \frac{ex^n + d}{cx^{2n} + a} dx$$

input `integrate((d+e*x^n)/(a+c*x^(2*n)),x, algorithm="giac")`

output `integrate((e*x^n + d)/(c*x^(2*n) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{d + ex^n}{a + cx^{2n}} dx = \int \frac{d + e x^n}{a + c x^{2n}} dx$$

input `int((d + e*x^n)/(a + c*x^(2*n)),x)`

output `int((d + e*x^n)/(a + c*x^(2*n)), x)`

Reduce [F]

$$\int \frac{d + ex^n}{a + cx^{2n}} dx = \left(\int \frac{x^n}{x^{2n}c + a} dx \right) e + \left(\int \frac{1}{x^{2n}c + a} dx \right) d$$

input `int((d+e*x^n)/(a+c*x^(2*n)),x)`

output `int(x**n/(x**2*n*c + a),x)*e + int(1/(x**2*n*c + a),x)*d`

$$\mathbf{3.29} \quad \int \frac{1}{(d+ex^n)(a+cx^{2n})} dx$$

Optimal result	236
Mathematica [A] (verified)	237
Rubi [A] (verified)	237
Maple [F]	238
Fricas [F]	239
Sympy [F(-2)]	239
Maxima [F]	239
Giac [F]	240
Mupad [F(-1)]	240
Reduce [F]	240

Optimal result

Integrand size = 21, antiderivative size = 152

$$\int \frac{1}{(d+ex^n)(a+cx^{2n})} dx = \frac{cdx \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a(cd^2 + ae^2)} \\ + \frac{e^2 x \text{Hypergeometric2F1}\left(1, \frac{1}{n}, 1 + \frac{1}{n}, -\frac{ex^n}{d}\right)}{d(cd^2 + ae^2)} \\ - \frac{cex^{1+n} \text{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a(cd^2 + ae^2)(1+n)}$$

output

```
c*d*x*hypergeom([1, 1/2/n],[1+1/2/n],-c*x^(2*n)/a)/a/(a*e^2+c*d^2)+e^2*x*h
ypergeom([1, 1/n],[1+1/n],-e*x^n/d)/d/(a*e^2+c*d^2)-c*e*x^(1+n)*hypergeom(
[1, 1/2*(1+n)/n],[3/2+1/2/n],-c*x^(2*n)/a)/a/(a*e^2+c*d^2)/(1+n)
```

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.86

$$\int \frac{1}{(d + ex^n)(a + cx^{2n})} dx \\ = \frac{x \left(cd^2(1+n) \text{Hypergeometric2F1} \left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a} \right) + e \left(ae(1+n) \text{Hypergeometric2F1} \left(1, \frac{1}{n}, 1, -\frac{ex^n}{a} \right) \right) \right)}{ad(cd^2 + ae^2)(1+n)}$$

input `Integrate[1/((d + e*x^n)*(a + c*x^(2*n))),x]`

output
$$(x*(c*d^2*(1+n)*\text{Hypergeometric2F1}[1, 1/(2*n), (2+n^(-1))/2, -(c*x^(2*n))/a]) + e*(a*e*(1+n)*\text{Hypergeometric2F1}[1, n^(-1), 1+n^(-1), -((e*x^n)/d)]) - c*d*x^n*\text{Hypergeometric2F1}[1, (1+n)/(2*n), (3+n^(-1))/2, -((c*x^(2*n))/a)]))/((a*d*(c*d^2 + a*e^2)*(1+n)))$$

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {1755, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + cx^{2n})(d + ex^n)} dx \\ \downarrow 1755 \\ \int \left(\frac{e^2}{(ae^2 + cd^2)(d + ex^n)} - \frac{c(ex^n - d)}{(ae^2 + cd^2)(a + cx^{2n})} \right) dx \\ \downarrow 2009$$

$$\begin{aligned}
 & - \frac{cex^{n+1} \text{Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a(n+1)(ae^2 + cd^2)} + \\
 & \frac{cdx \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a(ae^2 + cd^2)} + \\
 & \frac{e^2 x \text{Hypergeometric2F1}\left(1, \frac{1}{n}, 1 + \frac{1}{n}, -\frac{ex^n}{d}\right)}{d(ae^2 + cd^2)}
 \end{aligned}$$

input `Int[1/((d + e*x^n)*(a + c*x^(2*n))), x]`

output `(c*d*x*Hypergeometric2F1[1, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)])/(a*(c*d^2 + a*e^2)) + (e^2*x*Hypergeometric2F1[1, n^(-1), 1 + n^(-1), -((e*x^n)/d)])/(d*(c*d^2 + a*e^2)) - (c*e*x^(1 + n)*Hypergeometric2F1[1, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)])/(a*(c*d^2 + a*e^2)*(1 + n))`

Defintions of rubi rules used

rule 1755 `Int[((d_) + (e_.)*(x_)^(n_))^q_ / ((a_) + (c_.)*(x_)^(n2_)), x_Symbol] :> Int[ExpandIntegrand[(d + e*x^n)^q/(a + c*x^(2*n)), x], x] /; FreeQ[{a, c, d, e, n}, x] && EqQ[n2, 2*n] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[q]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [F]

$$\int \frac{1}{(d + e x^n) (a + c x^{2n})} dx$$

input `int(1/(d+e*x^n)/(a+c*x^(2*n)),x)`

output `int(1/(d+e*x^n)/(a+c*x^(2*n)),x)`

Fricas [F]

$$\int \frac{1}{(d + ex^n)(a + cx^{2n})} dx = \int \frac{1}{(cx^{2n} + a)(ex^n + d)} dx$$

input `integrate(1/(d+e*x^n)/(a+c*x^(2*n)),x, algorithm="fricas")`

output `integral(1/(a*e*x^n + a*d + (c*e*x^n + c*d)*x^(2*n)), x)`

Sympy [F(-2)]

Exception generated.

$$\int \frac{1}{(d + ex^n)(a + cx^{2n})} dx = \text{Exception raised: HeuristicGCDFailed}$$

input `integrate(1/(d+e*x**n)/(a+c*x**2*n),x)`

output `Exception raised: HeuristicGCDFailed >> no luck`

Maxima [F]

$$\int \frac{1}{(d + ex^n)(a + cx^{2n})} dx = \int \frac{1}{(cx^{2n} + a)(ex^n + d)} dx$$

input `integrate(1/(d+e*x^n)/(a+c*x^(2*n)),x, algorithm="maxima")`

output `integrate(1/((c*x^(2*n) + a)*(e*x^n + d)), x)`

Giac [F]

$$\int \frac{1}{(d + ex^n)(a + cx^{2n})} dx = \int \frac{1}{(cx^{2n} + a)(ex^n + d)} dx$$

input `integrate(1/(d+e*x^n)/(a+c*x^(2*n)),x, algorithm="giac")`

output `integrate(1/((c*x^(2*n) + a)*(e*x^n + d)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex^n)(a + cx^{2n})} dx = \int \frac{1}{(a + cx^{2n})(d + ex^n)} dx$$

input `int(1/((a + c*x^(2*n))*(d + e*x^n)),x)`

output `int(1/((a + c*x^(2*n))*(d + e*x^n)), x)`

Reduce [F]

$$\int \frac{1}{(d + ex^n)(a + cx^{2n})} dx = \int \frac{1}{x^{3n}ce + x^{2n}cd + x^nae + ad} dx$$

input `int(1/(d+e*x^n)/(a+c*x^(2*n)),x)`

output `int(1/(x**3*n*c*e + x**2*n*c*d + x**n*a*e + a*d),x)`

3.30 $\int \frac{1}{(d+ex^n)^2(a+cx^{2n})} dx$

Optimal result	241
Mathematica [A] (verified)	242
Rubi [A] (verified)	242
Maple [F]	243
Fricas [F]	244
Sympy [F(-2)]	244
Maxima [F]	244
Giac [F]	245
Mupad [F(-1)]	245
Reduce [F]	245

Optimal result

Integrand size = 21, antiderivative size = 224

$$\begin{aligned} & \int \frac{1}{(d + ex^n)^2(a + cx^{2n})} dx \\ &= \frac{e^2 x}{d (cd^2 + ae^2) n (d + ex^n)} \\ &+ \frac{c(cd^2 - ae^2) x \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a (cd^2 + ae^2)^2} \\ &- \frac{e^2 (cd^2(1 - 3n) + ae^2(1 - n)) x \text{Hypergeometric2F1}\left(1, \frac{1}{n}, 1 + \frac{1}{n}, -\frac{ex^n}{d}\right)}{d^2 (cd^2 + ae^2)^2 n} \\ &- \frac{2c^2 d e^{1+n} \text{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a (cd^2 + ae^2)^2 (1 + n)} \end{aligned}$$

output

```
e^2*x/d/(a*e^2+c*d^2)/n/(d+e*x^n)+c*(-a*e^2+c*d^2)*x*hypergeom([1, 1/2/n], [1+1/2/n], -c*x^(2*n)/a)/a/(a*e^2+c*d^2)^2-e^2*(c*d^2*(1-3*n)+a*e^2*(1-n))*x*hypergeom([1, 1/n], [1+1/n], -e*x^n/d)/d^2/(a*e^2+c*d^2)^2/n-2*c^2*d*e*x^(1+n)*hypergeom([1, 1/2*(1+n)/n], [3/2+1/2/n], -c*x^(2*n)/a)/a/(a*e^2+c*d^2)^2/(1+n)
```

Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 186, normalized size of antiderivative = 0.83

$$\int \frac{1}{(d + ex^n)^2 (a + cx^{2n})} dx \\ = \frac{x \left(cd^2 (cd^2 - ae^2) (1 + n) \text{Hypergeometric2F1} \left(1, \frac{1}{2n}, \frac{1}{2} \left(2 + \frac{1}{n} \right), -\frac{cx^{2n}}{a} \right) + e \left(2acd^2 e (1 + n) \text{Hypergeometric2F1} \left(1, \frac{1}{2n}, \frac{1}{2} \left(2 + \frac{1}{n} \right), -\frac{ex^n}{a} \right) + a^2 c^2 d^2 (cd^2 - ae^2)^2 (1 + n)^2 \right) \right)}{(a + cx^{2n})^2}$$

input `Integrate[1/((d + e*x^n)^2*(a + c*x^(2*n))),x]`

output
$$(x*(c*d^2*(c*d^2 - a*e^2)*(1 + n)*\text{Hypergeometric2F1}[1, 1/(2*n), (2 + n^{(-1)})/2, -((c*x^(2*n))/a)] + e*(2*a*c*d^2*e*(1 + n)*\text{Hypergeometric2F1}[1, n^{(-1)}, 1 + n^{(-1)}, -((e*x^n)/d)] - 2*c^2*d^3*x^n*\text{Hypergeometric2F1}[1, (1 + n)/(2*n), (3 + n^{(-1)})/2, -((c*x^(2*n))/a)] + a*e*(c*d^2 + a*e^2)*(1 + n)*\text{Hypergeometric2F1}[2, n^{(-1)}, 1 + n^{(-1)}, -((e*x^n)/d)]))/((a*(c*d^3 + a*d*e^2)^2*(1 + n)))$$

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 205, normalized size of antiderivative = 0.92, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {1755, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + cx^{2n}) (d + ex^n)^2} dx \\ \downarrow 1755 \\ \int \left(\frac{2cde^2}{(ae^2 + cd^2)^2 (d + ex^n)} + \frac{e^2}{(ae^2 + cd^2) (d + ex^n)^2} - \frac{c(ae^2 - cd^2 + 2cdex^n)}{(ae^2 + cd^2)^2 (a + cx^{2n})} \right) dx \\ \downarrow 2009$$

$$\begin{aligned}
 & -\frac{2c^2dex^{n+1} \text{Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a(n+1)(ae^2 + cd^2)^2} + \\
 & \frac{cx(cd^2 - ae^2) \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a(ae^2 + cd^2)^2} + \\
 & \frac{2ce^2x \text{Hypergeometric2F1}\left(1, \frac{1}{n}, 1 + \frac{1}{n}, -\frac{ex^n}{d}\right)}{(ae^2 + cd^2)^2} + \frac{e^2x \text{Hypergeometric2F1}\left(2, \frac{1}{n}, 1 + \frac{1}{n}, -\frac{ex^n}{d}\right)}{d^2(ae^2 + cd^2)}
 \end{aligned}$$

input `Int[1/((d + e*x^n)^2*(a + c*x^(2*n))), x]`

output `(c*(c*d^2 - a*e^2)*x*Hypergeometric2F1[1, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)]/(a*(c*d^2 + a*e^2)^2) + (2*c*e^2*x*Hypergeometric2F1[1, n^(-1), 1 + n^(-1), -(e*x^n)/d])/((c*d^2 + a*e^2)^2 - (2*c^2*d*e*x^(1 + n))*Hypergeometric2F1[1, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)])/(a*(c*d^2 + a*e^2)^2*(1 + n)) + (e^2*x*Hypergeometric2F1[2, n^(-1), 1 + n^(-1), -(e*x^n)/d])/((d^2*(c*d^2 + a*e^2)))`

Definitions of rubi rules used

rule 1755 `Int[((d_) + (e_.)*(x_)^(n_))^q_]/((a_) + (c_.)*(x_)^(n2_)), x_Symbol] :> Int[ExpandIntegrand[(d + e*x^n)^q/(a + c*x^(2*n)), x], x] /; FreeQ[{a, c, d, e, n}, x] && EqQ[n2, 2*n] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[q]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [F]

$$\int \frac{1}{(d + e x^n)^2 (a + c x^{2n})} dx$$

input `int(1/(d+e*x^n)^2/(a+c*x^(2*n)), x)`

output `int(1/(d+e*x^n)^2/(a+c*x^(2*n)), x)`

Fricas [F]

$$\int \frac{1}{(d + ex^n)^2 (a + cx^{2n})} dx = \int \frac{1}{(cx^{2n} + a)(ex^n + d)^2} dx$$

input `integrate(1/(d+e*x^n)^2/(a+c*x^(2*n)),x, algorithm="fricas")`

output `integral(1/(a*e^2*x^(2*n) + 2*a*d*e*x^n + a*d^2 + (c*e^2*x^(2*n) + 2*c*d*e*x^n + c*d^2)*x^(2*n)), x)`

Sympy [F(-2)]

Exception generated.

$$\int \frac{1}{(d + ex^n)^2 (a + cx^{2n})} dx = \text{Exception raised: HeuristicGCDFailed}$$

input `integrate(1/(d+e*x**n)**2/(a+c*x**(2*n)),x)`

output `Exception raised: HeuristicGCDFailed >> no luck`

Maxima [F]

$$\int \frac{1}{(d + ex^n)^2 (a + cx^{2n})} dx = \int \frac{1}{(cx^{2n} + a)(ex^n + d)^2} dx$$

input `integrate(1/(d+e*x^n)^2/(a+c*x^(2*n)),x, algorithm="maxima")`

output `e^2*x/(c*d^4*n + a*d^2*e^2*n + (c*d^3*e*n + a*d*e^3*n)*x^n) + (c*d^2*e^2*(3*n - 1) + a*e^4*(n - 1))*integrate(1/(c^2*d^6*n + 2*a*c*d^4*e^2*n + a^2*d^2*e^4*n + (c^2*d^5*e*n + 2*a*c*d^3*e^3*n + a^2*d*e^5*n)*x^n), x) - integrate((2*c^2*d*e*x^n - c^2*d^2 + a*c*e^2)/(a*c^2*d^4 + 2*a^2*c*d^2*e^2 + a^3*e^4 + (c^3*d^4 + 2*a*c^2*d^2*e^2 + a^2*c*e^4)*x^(2*n)), x)`

Giac [F]

$$\int \frac{1}{(d + ex^n)^2 (a + cx^{2n})} dx = \int \frac{1}{(cx^{2n} + a)(ex^n + d)^2} dx$$

input `integrate(1/(d+e*x^n)^2/(a+c*x^(2*n)),x, algorithm="giac")`

output `integrate(1/((c*x^(2*n) + a)*(e*x^n + d)^2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex^n)^2 (a + cx^{2n})} dx = \int \frac{1}{(a + cx^{2n}) (d + ex^n)^2} dx$$

input `int(1/((a + c*x^(2*n))*(d + e*x^n)^2),x)`

output `int(1/((a + c*x^(2*n))*(d + e*x^n)^2), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{1}{(d + ex^n)^2 (a + cx^{2n})} dx \\ &= \int \frac{1}{x^{4n} c e^2 + 2x^{3n} c d e + x^{2n} a e^2 + x^{2n} c d^2 + 2x^n a d e + a d^2} dx \end{aligned}$$

input `int(1/(d+e*x^n)^2/(a+c*x^(2*n)),x)`

output `int(1/(x**4*n*c*e**2 + 2*x**3*n*c*d*e + x**2*n*a*e**2 + x**2*n*c*d**2 + 2*x**n*a*d*e + a*d**2),x)`

3.31 $\int \frac{d+ex^n}{a-cx^{2n}} dx$

Optimal result	246
Mathematica [A] (verified)	246
Rubi [A] (verified)	247
Maple [F]	248
Fricas [F]	248
Sympy [C] (verification not implemented)	249
Maxima [F]	249
Giac [F]	250
Mupad [F(-1)]	250
Reduce [F]	250

Optimal result

Integrand size = 20, antiderivative size = 81

$$\int \frac{d + ex^n}{a - cx^{2n}} dx = \frac{dx \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), \frac{cx^{2n}}{a}\right)}{a} \\ + \frac{ex^{1+n} \text{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3 + \frac{1}{n}), \frac{cx^{2n}}{a}\right)}{a(1+n)}$$

output `d*x*hypergeom([1, 1/2/n], [1+1/2/n], c*x^(2*n)/a)/a+e*x^(1+n)*hypergeom([1, 1/2*(1+n)/n], [3/2+1/2/n], c*x^(2*n)/a)/a/(1+n)`

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.00

$$\int \frac{d + ex^n}{a - cx^{2n}} dx = \frac{dx \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), \frac{cx^{2n}}{a}\right)}{a} \\ + \frac{ex^{1+n} \text{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3 + \frac{1}{n}), \frac{cx^{2n}}{a}\right)}{a(1+n)}$$

input `Integrate[(d + e*x^n)/(a - c*x^(2*n)), x]`

output $(d*x*Hypergeometric2F1[1, 1/(2*n), (2 + n^{-1})/2, (c*x^(2*n))/a])/a + (e*x^(1 + n)*Hypergeometric2F1[1, (1 + n)/(2*n), (3 + n^{-1})/2, (c*x^(2*n))/a])/(a*(1 + n))$

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {1748, 778, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{d + ex^n}{a - cx^{2n}} dx \\
 & \quad \downarrow \textcolor{blue}{1748} \\
 & d \int \frac{1}{a - cx^{2n}} dx + e \int \frac{x^n}{a - cx^{2n}} dx \\
 & \quad \downarrow \textcolor{blue}{778} \\
 & e \int \frac{x^n}{a - cx^{2n}} dx + \frac{dx \text{ Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), \frac{cx^{2n}}{a}\right)}{a} \\
 & \quad \downarrow \textcolor{blue}{888} \\
 & \frac{dx \text{ Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), \frac{cx^{2n}}{a}\right)}{a} + \\
 & \frac{ex^{n+1} \text{ Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3 + \frac{1}{n}), \frac{cx^{2n}}{a}\right)}{a(n+1)}
 \end{aligned}$$

input $\text{Int}[(d + e*x^n)/(a - c*x^(2*n)), x]$

output $(d*x*Hypergeometric2F1[1, 1/(2*n), (2 + n^{-1})/2, (c*x^(2*n))/a])/a + (e*x^(1 + n)*Hypergeometric2F1[1, (1 + n)/(2*n), (3 + n^{-1})/2, (c*x^(2*n))/a])/(a*(1 + n))$

Definitions of rubi rules used

rule 778 $\text{Int}[(a_.) + (b_.)*(x_.)^{(n_.)}^{(p_.)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[a^p * x * \text{Hypergeometric2F1}[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; \text{FreeQ}[\{a, b, n, p\}, x] \&& \text{!IGtQ}[p, 0] \&& \text{!IntegerQ}[1/n] \&& \text{!ILtQ}[\text{Simplify}[1/n + p], 0] \&& (\text{IntegerQ}[p] \mid \text{GtQ}[a, 0])$

rule 888 $\text{Int}[((c_.)*(x_.))^{(m_.)}*((a_.) + (b_.)*(x_.)^{(n_.)}^{(p_.)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[a^p * ((c*x)^(m + 1)/(c*(m + 1))) * \text{Hypergeometric2F1}[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; \text{FreeQ}[\{a, b, c, m, n, p\}, x] \&& \text{!IGtQ}[p, 0] \&& (\text{ILtQ}[p, 0] \mid \text{GtQ}[a, 0])$

rule 1748 $\text{Int}[((d_.) + (e_.)*(x_.)^{(n_.)})/((a_.) + (c_.)*(x_.)^{(n2_.)}), x_{\text{Symbol}}] \rightarrow \text{Simp}[d \text{Int}[1/(a + c*x^(2*n)), x], x] + \text{Simp}[e \text{Int}[x^n/(a + c*x^(2*n)), x], x] /; \text{FreeQ}[\{a, c, d, e, n\}, x] \&& \text{EqQ}[n2, 2*n] \&& \text{NeQ}[c*d^2 + a*e^2, 0] \&& (\text{PosQ}[a*c] \mid \text{!IntegerQ}[n])$

Maple [F]

$$\int \frac{d + e x^n}{a - c x^{2n}} dx$$

input $\text{int}((d+e*x^n)/(a-c*x^(2*n)), x)$

output $\text{int}((d+e*x^n)/(a-c*x^(2*n)), x)$

Fricas [F]

$$\int \frac{d + e x^n}{a - c x^{2n}} dx = \int -\frac{e x^n + d}{c x^{2n} - a} dx$$

input $\text{integrate}((d+e*x^n)/(a-c*x^(2*n)), x, \text{algorithm}=\text{"fricas"})$

output `integral(-(e*x^n + d)/(c*x^(2*n) - a), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 2.40 (sec) , antiderivative size = 214, normalized size of antiderivative = 2.64

$$\int \frac{d + ex^n}{a - cx^{2n}} dx = \frac{a^{\frac{1}{2n}} a^{-1 - \frac{1}{2n}} dx \Phi\left(\frac{cx^{2n} e^{2i\pi}}{a}, 1, \frac{1}{2n}\right) \Gamma\left(\frac{1}{2n}\right)}{4n^2 \Gamma\left(1 + \frac{1}{2n}\right)}$$

$$+ \frac{a^{-\frac{3}{2} - \frac{1}{2n}} a^{\frac{1}{2} + \frac{1}{2n}} ex^{n+1} \Phi\left(\frac{cx^{2n} e^{2i\pi}}{a}, 1, \frac{1}{2} + \frac{1}{2n}\right) \Gamma\left(\frac{1}{2} + \frac{1}{2n}\right)}{4n \Gamma\left(\frac{3}{2} + \frac{1}{2n}\right)}$$

$$+ \frac{a^{-\frac{3}{2} - \frac{1}{2n}} a^{\frac{1}{2} + \frac{1}{2n}} ex^{n+1} \Phi\left(\frac{cx^{2n} e^{2i\pi}}{a}, 1, \frac{1}{2} + \frac{1}{2n}\right) \Gamma\left(\frac{1}{2} + \frac{1}{2n}\right)}{4n^2 \Gamma\left(\frac{3}{2} + \frac{1}{2n}\right)}$$

input `integrate((d+e*x**n)/(a-c*x**(2*n)),x)`

output `a**(1/(2*n))*a**(-1 - 1/(2*n))*d*x*lerchphi(c*x**(2*n)*exp_polar(2*I*pi)/a, 1, 1/(2*n))*gamma(1/(2*n))/(4*n**2*gamma(1 + 1/(2*n))) + a**(-3/2 - 1/(2*n))*a**(-1/2 + 1/(2*n))*e*x***(n + 1)*lerchphi(c*x**(2*n)*exp_polar(2*I*pi)/a, 1, 1/2 + 1/(2*n))*gamma(1/2 + 1/(2*n))/(4*n*gamma(3/2 + 1/(2*n))) + a*(-3/2 - 1/(2*n))*a**(-1/2 + 1/(2*n))*e*x***(n + 1)*lerchphi(c*x**(2*n)*exp_polar(2*I*pi)/a, 1, 1/2 + 1/(2*n))*gamma(1/2 + 1/(2*n))/(4*n**2*gamma(3/2 + 1/(2*n)))`

Maxima [F]

$$\int \frac{d + ex^n}{a - cx^{2n}} dx = \int -\frac{ex^n + d}{cx^{2n} - a} dx$$

input `integrate((d+e*x^n)/(a-c*x^(2*n)),x, algorithm="maxima")`

output `-integrate((e*x^n + d)/(c*x^(2*n) - a), x)`

Giac [F]

$$\int \frac{d + ex^n}{a - cx^{2n}} dx = \int -\frac{ex^n + d}{cx^{2n} - a} dx$$

input `integrate((d+e*x^n)/(a-c*x^(2*n)),x, algorithm="giac")`

output `integrate(-(e*x^n + d)/(c*x^(2*n) - a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{d + ex^n}{a - cx^{2n}} dx = \int \frac{d + e x^n}{a - c x^{2n}} dx$$

input `int((d + e*x^n)/(a - c*x^(2*n)),x)`

output `int((d + e*x^n)/(a - c*x^(2*n)), x)`

Reduce [F]

$$\int \frac{d + ex^n}{a - cx^{2n}} dx = - \left(\int \frac{x^n}{x^{2n}c - a} dx \right) e - \left(\int \frac{1}{x^{2n}c - a} dx \right) d$$

input `int((d+e*x^n)/(a-c*x^(2*n)),x)`

output `- (int(x**n/(x**2*n)*c - a,x)*e + int(1/(x**2*n)*c - a,x)*d)`

3.32 $\int \frac{(d+ex^n)^3}{(a+cx^{2n})^2} dx$

Optimal result	251
Mathematica [A] (verified)	252
Rubi [A] (verified)	252
Maple [F]	254
Fricas [F]	254
Sympy [F]	254
Maxima [F]	255
Giac [F]	255
Mupad [F(-1)]	255
Reduce [F]	256

Optimal result

Integrand size = 21, antiderivative size = 210

$$\begin{aligned} & \int \frac{(d+ex^n)^3}{(a+cx^{2n})^2} dx \\ &= -\frac{3de^2x}{2acn} - \frac{e^3x^{1+n}}{2acn} + \frac{x(d+ex^n)^3}{2an(a+cx^{2n})} \\ &+ \frac{d(3ae^2 - cd^2(1-2n))x \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{2a^2cn} \\ &- \frac{e(3cd^2(1-n) - ae^2(1+n))x^{1+n} \operatorname{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{2a^2cn(1+n)} \end{aligned}$$

output

```
-3/2*d*e^2*x/a/c/n-1/2*e^3*x^(1+n)/a/c/n+1/2*x*(d+e*x^n)^3/a/n/(a+c*x^(2*n))
)+1/2*d*(3*a*e^2-c*d^2*(1-2*n))*x*hypergeom([1, 1/2/n],[1+1/2/n],-c*x^(2*n)/a)/a^2/c/n-1/2*e*(3*c*d^2*(1-n)-a*e^2*(1+n))*x^(1+n)*hypergeom([1, 1/2*(1+n)/n],[3/2+1/2/n],-c*x^(2*n)/a)/a^2/c/n/(1+n)
```

Mathematica [A] (verified)

Time = 0.50 (sec) , antiderivative size = 188, normalized size of antiderivative = 0.90

$$\int \frac{(d + ex^n)^3}{(a + cx^{2n})^2} dx \\ = \frac{x \left(3ade^2 \text{Hypergeometric2F1} \left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a} \right) + \frac{ae^3 x^n \text{Hypergeometric2F1} \left(1, \frac{1+n}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a} \right)}{1+n} \right) + d(c a e^2 x^{2n} \text{Hypergeometric2F1} \left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a} \right) + \frac{ae^3 x^n \text{Hypergeometric2F1} \left(1, \frac{1+n}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a} \right)}{1+n})}{(a + cx^{2n})^2}$$

input `Integrate[(d + e*x^n)^3/(a + c*x^(2*n))^2, x]`

output
$$(x*(3*a*d*e^2*\text{Hypergeometric2F1}[1, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)] + (a*e^3*x^n*\text{Hypergeometric2F1}[1, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)]/(1 + n) + d*(c*d^2 - 3*a*e^2)*\text{Hypergeometric2F1}[2, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)] + (e*(3*c*d^2 - a*e^2)*x^n*\text{Hypergeometric2F1}[2, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)]/(1 + n)))/(a^2*c))$$

Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 288, normalized size of antiderivative = 1.37, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {1767, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^n)^3}{(a + cx^{2n})^2} dx \\ \downarrow 1767 \\ \int \left(\frac{x^n(3cd^2e - ae^3) - 3ade^2 + cd^3}{c(a + cx^{2n})^2} + \frac{e^2(3d + ex^n)}{c(a + cx^{2n})} \right) dx \\ \downarrow 2009$$

$$\begin{aligned}
 & -\frac{e(1-n)x^{n+1}(3cd^2 - ae^2) \text{Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{2a^2cn(n+1)} - \\
 & \frac{d(1-2n)x(cd^2 - 3ae^2) \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{2a^2cn} + \\
 & \frac{x(ex^n(3cd^2 - ae^2) + d(cd^2 - 3ae^2))}{2acn(a + cx^{2n})} + \frac{3de^2x \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{ac} + \\
 & \frac{e^3x^{n+1} \text{Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{ac(n+1)}
 \end{aligned}$$

input `Int[(d + e*x^n)^3/(a + c*x^(2*n))^2, x]`

output `(x*(d*(c*d^2 - 3*a*e^2) + e*(3*c*d^2 - a*e^2)*x^n)/(2*a*c*n*(a + c*x^(2*n))) + (3*d*e^2*x*Hypergeometric2F1[1, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)])/(a*c) - (d*(c*d^2 - 3*a*e^2)*(1 - 2*n)*x*Hypergeometric2F1[1, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)])/(2*a^2*c*n) + (e^3*x^(1 + n)*Hypergeometric2F1[1, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)])/(a*c*(1 + n)) - (e*(3*c*d^2 - a*e^2)*(1 - n)*x^(1 + n)*Hypergeometric2F1[1, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)])/(2*a^2*c*n*(1 + n))`

Definitions of rubi rules used

rule 1767 `Int[((d_) + (e_.)*(x_)^(n_))^(q_)*((a_) + (c_.)*(x_)^(n2_))^(p_), x_Symbol] :> Int[ExpandIntegrand[(d + e*x^n)^q*(a + c*x^(2*n))^p, x], x] /; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && NeQ[c*d^2 + a*e^2, 0] && ((IntegersQ[p, q] && !IntegerQ[n]) || IGtQ[p, 0] || (IGtQ[q, 0] && !IntegerQ[n]))`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [F]

$$\int \frac{(d + e x^n)^3}{(a + c x^{2n})^2} dx$$

input `int((d+e*x^n)^3/(a+c*x^(2*n))^2,x)`

output `int((d+e*x^n)^3/(a+c*x^(2*n))^2,x)`

Fricas [F]

$$\int \frac{(d + e x^n)^3}{(a + c x^{2n})^2} dx = \int \frac{(e x^n + d)^3}{(c x^{2n} + a)^2} dx$$

input `integrate((d+e*x^n)^3/(a+c*x^(2*n))^2,x, algorithm="fricas")`

output `integral((e^3*x^(3*n) + 3*d*e^2*x^(2*n) + 3*d^2*e*x^n + d^3)/(c^2*x^(4*n) + 2*a*c*x^(2*n) + a^2), x)`

Sympy [F]

$$\int \frac{(d + e x^n)^3}{(a + c x^{2n})^2} dx = \int \frac{(d + e x^n)^3}{(a + c x^{2n})^2} dx$$

input `integrate((d+e*x**n)**3/(a+c*x**2*n)**2,x)`

output `Integral((d + e*x**n)**3/(a + c*x**2*n)**2, x)`

Maxima [F]

$$\int \frac{(d + ex^n)^3}{(a + cx^{2n})^2} dx = \int \frac{(ex^n + d)^3}{(cx^{2n} + a)^2} dx$$

input `integrate((d+e*x^n)^3/(a+c*x^(2*n))^2,x, algorithm="maxima")`

output `1/2*((3*c*d^2*e - a*e^3)*x*x^n + (c*d^3 - 3*a*d*e^2)*x)/(a*c^2*n*x^(2*n) + a^2*c*n) + integrate(1/2*(c*d^3*(2*n - 1) + 3*a*d*e^2 + (a*e^3*(n + 1) + 3*c*d^2*e*(n - 1))*x^n)/(a*c^2*n*x^(2*n) + a^2*c*n), x)`

Giac [F]

$$\int \frac{(d + ex^n)^3}{(a + cx^{2n})^2} dx = \int \frac{(ex^n + d)^3}{(cx^{2n} + a)^2} dx$$

input `integrate((d+e*x^n)^3/(a+c*x^(2*n))^2,x, algorithm="giac")`

output `integrate((e*x^n + d)^3/(c*x^(2*n) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(d + ex^n)^3}{(a + cx^{2n})^2} dx = \int \frac{(d + e x^n)^3}{(a + c x^{2n})^2} dx$$

input `int((d + e*x^n)^3/(a + c*x^(2*n))^2,x)`

output `int((d + e*x^n)^3/(a + c*x^(2*n))^2, x)`

Reduce [F]

$$\int \frac{(d + ex^n)^3}{(a + cx^{2n})^2} dx = \text{Too large to display}$$

input int((d+e*x^n)^3/(a+c*x^(2*n))^2,x)

```

output
(x**((2*n)*int(x**((3*n)/(x**((4*n)*c**2*n + x**((4*n)*c**2 + 2*x**((2*n)*a*c*n
+ 2*x**((2*n)*a*c + a**2*n + a**2),x)*a*c*e**3*n**2 + 2*x**((2*n)*int(x**((3
*n)/(x**((4*n)*c**2*n + x**((4*n)*c**2 + 2*x**((2*n)*a*c*n + 2*x**((2*n)*a*c +
a**2*n + a**2),x)*a*c*e**3*n + x**((2*n)*int(x**((3*n)/(x**((4*n)*c**2*n + x
**((4*n)*c**2 + 2*x**((2*n)*a*c*n + 2*x**((2*n)*a*c + a**2*n + a**2),x)*a*c*e
**3 + 3*x**((2*n)*int(x**((3*n)/(x**((4*n)*c**2*n + x**((4*n)*c**2 + 2*x**((2*n
)*a*c*n + 2*x**((2*n)*a*c + a**2*n + a**2),x)*c**2*d**2*e*n**2 - 3*x**((2*n)
*int(x**((3*n)/(x**((4*n)*c**2*n + x**((4*n)*c**2 + 2*x**((2*n)*a*c*n + 2*x**(
2*n)*a*c + a**2*n + a**2),x)*c**2*d**2*e + 3*x**((2*n)*int(x**((2*n)/(x**((4*
n)*c**2*n + x**((4*n)*c**2 + 2*x**((2*n)*a*c*n + 2*x**((2*n)*a*c + a**2*n + a
**2),x)*a*c*d*e**2*n**2 + 6*x**((2*n)*int(x**((2*n)/(x**((4*n)*c**2*n + x**((4
*n)*c**2 + 2*x**((2*n)*a*c*n + 2*x**((2*n)*a*c + a**2*n + a**2),x)*a*c*d*e**2*n
+ 3*x**((2*n)*int(x**((2*n)/(x**((4*n)*c**2*n + x**((4*n)*c**2 + 2*x**((2*n)
)*a*c*n + 2*x**((2*n)*a*c + a**2*n + a**2),x)*a*c*d*e**2 + 2*x**((2*n)*int(x
**((2*n)/(x**((4*n)*c**2*n + x**((4*n)*c**2 + 2*x**((2*n)*a*c*n + 2*x**((2*n)*a
*c + a**2*n + a**2),x)*c**2*d**3*n**3 + 3*x**((2*n)*int(x**((2*n)/(x**((4*n)*
c**2*n + x**((4*n)*c**2 + 2*x**((2*n)*a*c*n + 2*x**((2*n)*a*c + a**2*n + a**2
),x)*c**2*d**3*n**2 - x**((2*n)*int(x**((2*n)/(x**((4*n)*c**2*n + x**((4*n)*c*
2 + 2*x**((2*n)*a*c*n + 2*x**((2*n)*a*c + a**2*n + a**2),x)*c**2*d**3 + 3*x
**n*d**2*e*x + int(x**((3*n)/(x**((4*n)*c**2*n + x**((4*n)*c**2 + 2*x**((2*...

```

3.33 $\int \frac{(d+ex^n)^2}{(a+cx^{2n})^2} dx$

Optimal result	257
Mathematica [A] (verified)	258
Rubi [A] (verified)	258
Maple [F]	259
Fricas [F]	260
Sympy [F]	260
Maxima [F]	260
Giac [F]	261
Mupad [F(-1)]	261
Reduce [F]	261

Optimal result

Integrand size = 21, antiderivative size = 166

$$\begin{aligned} & \int \frac{(d+ex^n)^2}{(a+cx^{2n})^2} dx \\ &= -\frac{e^2 x}{2acn} + \frac{x(d+ex^n)^2}{2an(a+cx^{2n})} \\ &+ \frac{(ae^2 - cd^2(1-2n))x \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{2a^2 cn} \\ &- \frac{de(1-n)x^{1+n} \operatorname{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a^2 n(1+n)} \end{aligned}$$

output

```
-1/2*e^2*x/a/c/n+1/2*x*(d+e*x^n)^2/a/n/(a+c*x^(2*n))+1/2*(a*e^2-c*d^2*(1-2*n))*x*hypergeom([1, 1/2/n],[1+1/2/n],-c*x^(2*n)/a)/a^2/c/n-d*e*(1-n)*x^(1+n)*hypergeom([1, 1/2*(1+n)/n],[3/2+1/2/n],-c*x^(2*n)/a)/a^2/n/(1+n)
```

Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.82

$$\int \frac{(d + ex^n)^2}{(a + cx^{2n})^2} dx \\ = \frac{x \left(ae^2 (1+n) \text{Hypergeometric2F1} \left(1, \frac{1}{2n}, \frac{1}{2} \left(2 + \frac{1}{n} \right), -\frac{cx^{2n}}{a} \right) + (cd^2 - ae^2) (1+n) \text{Hypergeometric2F1} \left(2, \frac{1}{2n}, \frac{1}{2} \left(2 + \frac{1}{n} \right), -\frac{cx^{2n}}{a} \right) \right)}{a^2 c (1+n)}$$

input `Integrate[(d + e*x^n)^2/(a + c*x^(2*n))^2, x]`

output
$$(x*(a*e^2*(1+n)*\text{Hypergeometric2F1}[1, 1/(2*n), (2+n^(-1))/2, -((c*x^(2*n))/a)] + (c*d^2 - a*e^2)*(1+n)*\text{Hypergeometric2F1}[2, 1/(2*n), (2+n^(-1))/2, -((c*x^(2*n))/a)] + 2*c*d*e*x^n*\text{Hypergeometric2F1}[2, (1+n)/(2*n), (3+n^(-1))/2, -((c*x^(2*n))/a)]))/((a^2*c*(1+n)))$$

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.22, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.095, Rules used = {1767, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^n)^2}{(a + cx^{2n})^2} dx \\ \downarrow 1767 \\ \int \left(\frac{-ae^2 + cd^2 + 2cdex^n}{c(a + cx^{2n})^2} + \frac{e^2}{c(a + cx^{2n})} \right) dx \\ \downarrow 2009$$

$$\begin{aligned}
 & -\frac{(1-2n)x(cd^2 - ae^2) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{2a^2 cn} \\
 & \frac{de(1-n)x^{n+1} \operatorname{Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a^2 n(n+1)} + \frac{x(-ae^2 + cd^2 + 2cdex^n)}{2acn(a + cx^{2n})} + \\
 & \frac{e^2 x \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{ac}
 \end{aligned}$$

input `Int[(d + e*x^n)^2/(a + c*x^(2*n))^2, x]`

output `(x*(c*d^2 - a*e^2 + 2*c*d*e*x^n))/(2*a*c*n*(a + c*x^(2*n))) + (e^2*x*Hypergeometric2F1[1, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)])/(a*c) - ((c*d^2 - a*e^2)*(1 - 2*n)*x*Hypergeometric2F1[1, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)])/(2*a^2*c*n) - (d*e*(1 - n)*x^(1 + n)*Hypergeometric2F1[1, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)])/(a^2*n*(1 + n))`

Definitions of rubi rules used

rule 1767 `Int[((d_) + (e_.)*(x_)^(n_))^(q_)*((a_) + (c_.)*(x_)^(n2_))^(p_), x_Symbol] :> Int[ExpandIntegrand[(d + e*x^n)^q*(a + c*x^(2*n))^p, x], x] /; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && NeQ[c*d^2 + a*e^2, 0] && ((IntegersQ[p, q] && !IntegerQ[n]) || IGtQ[p, 0] || (IGtQ[q, 0] && !IntegerQ[n]))`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [F]

$$\int \frac{(d + e x^n)^2}{(a + c x^{2n})^2} dx$$

input `int((d+e*x^n)^2/(a+c*x^(2*n))^2, x)`

output `int((d+e*x^n)^2/(a+c*x^(2*n))^2, x)`

Fricas [F]

$$\int \frac{(d + ex^n)^2}{(a + cx^{2n})^2} dx = \int \frac{(ex^n + d)^2}{(cx^{2n} + a)^2} dx$$

input `integrate((d+e*x^n)^2/(a+c*x^(2*n))^2,x, algorithm="fricas")`

output `integral((e^2*x^(2*n) + 2*d*e*x^n + d^2)/(c^2*x^(4*n) + 2*a*c*x^(2*n) + a^2), x)`

Sympy [F]

$$\int \frac{(d + ex^n)^2}{(a + cx^{2n})^2} dx = \int \frac{(d + ex^n)^2}{(a + cx^{2n})^2} dx$$

input `integrate((d+e*x**n)**2/(a+c*x**(2*n))**2,x)`

output `Integral((d + e*x**n)**2/(a + c*x**(2*n))**2, x)`

Maxima [F]

$$\int \frac{(d + ex^n)^2}{(a + cx^{2n})^2} dx = \int \frac{(ex^n + d)^2}{(cx^{2n} + a)^2} dx$$

input `integrate((d+e*x^n)^2/(a+c*x^(2*n))^2,x, algorithm="maxima")`

output `1/2*(2*c*d*e*x*x^n + (c*d^2 - a*e^2)*x)/(a*c^2*n*x^(2*n) + a^2*c*n) + integrate(1/2*(2*c*d*e*(n - 1)*x^n + c*d^2*(2*n - 1) + a*e^2)/(a*c^2*n*x^(2*n) + a^2*c*n), x)`

Giac [F]

$$\int \frac{(d + ex^n)^2}{(a + cx^{2n})^2} dx = \int \frac{(ex^n + d)^2}{(cx^{2n} + a)^2} dx$$

input `integrate((d+e*x^n)^2/(a+c*x^(2*n))^2,x, algorithm="giac")`

output `integrate((e*x^n + d)^2/(c*x^(2*n) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(d + ex^n)^2}{(a + cx^{2n})^2} dx = \int \frac{(d + e x^n)^2}{(a + c x^{2n})^2} dx$$

input `int((d + e*x^n)^2/(a + c*x^(2*n))^2,x)`

output `int((d + e*x^n)^2/(a + c*x^(2*n))^2, x)`

Reduce [F]

$$\begin{aligned} & \int \frac{(d + ex^n)^2}{(a + cx^{2n})^2} dx \\ &= \frac{x^{2n} \left(\int \frac{x^{2n}}{x^{4n}c^2 + 2x^{2n}ac + a^2} dx \right) ac e^2 + 2x^{2n} \left(\int \frac{x^{2n}}{x^{4n}c^2 + 2x^{2n}ac + a^2} dx \right) c^2 d^2 n - x^{2n} \left(\int \frac{x^{2n}}{x^{4n}c^2 + 2x^{2n}ac + a^2} dx \right) c^2 d^2 + 2x^{2n} \left(\int \frac{x^{2n}}{x^{4n}c^2 + 2x^{2n}ac + a^2} dx \right) a c e^2}{x^{4n}c^2 + 2x^{2n}ac + a^2} \end{aligned}$$

input `int((d+e*x^n)^2/(a+c*x^(2*n))^2,x)`

```
output
(x**(2*n)*int(x**2*n)/(x**4*n)*c**2 + 2*x**2*n*a*c + a**2),x)*a*c*e**2
+ 2*x**2*n*int(x**2*n)/(x**4*n)*c**2 + 2*x**2*n*a*c + a**2),x)*c**2
*d**2*n - x**2*n*int(x**2*n)/(x**4*n)*c**2 + 2*x**2*n*a*c + a**2),x)
*c**2*d**2 + 2*x**2*n*int(x**n/(x**4*n)*c**2 + 2*x**2*n*a*c + a**2),x
)*a*c*d*e + int(x**2*n)/(x**4*n)*c**2 + 2*x**2*n*a*c + a**2),x)*a**2*e
**2 + 2*int(x**2*n)/(x**4*n)*c**2 + 2*x**2*n*a*c + a**2),x)*a*c*d**2*n
- int(x**2*n)/(x**4*n)*c**2 + 2*x**2*n*a*c + a**2),x)*a*c*d**2 + 2*in
t(x**n/(x**4*n)*c**2 + 2*x**2*n*a*c + a**2),x)*a**2*d*e + d**2*x)/(a*(x
)**(2*n)*c + a))
```

3.34 $\int \frac{d+ex^n}{(a+cx^{2n})^2} dx$

Optimal result	263
Mathematica [A] (verified)	264
Rubi [A] (verified)	264
Maple [F]	266
Fricas [F]	266
Sympy [C] (verification not implemented)	266
Maxima [F]	267
Giac [F]	268
Mupad [F(-1)]	268
Reduce [F]	268

Optimal result

Integrand size = 19, antiderivative size = 134

$$\begin{aligned} \int \frac{d+ex^n}{(a+cx^{2n})^2} dx &= \frac{x(d+ex^n)}{2an(a+cx^{2n})} \\ &\quad - \frac{d(1-2n)x \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{2a^2n} \\ &\quad - \frac{e(1-n)x^{1+n} \operatorname{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{2a^2n(1+n)} \end{aligned}$$

output $1/2*x*(d+e*x^n)/a/n/(a+c*x^(2*n))-1/2*d*(1-2*n)*x*hypergeom([1, 1/2/n], [1+1/2/n], -c*x^(2*n)/a)/a^2/n-1/2*e*(1-n)*x^(1+n)*hypergeom([1, 1/2*(1+n)/n], [3/2+1/2/n], -c*x^(2*n)/a)/a^2/n/(1+n)$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.62

$$\int \frac{d + ex^n}{(a + cx^{2n})^2} dx = \frac{dx \text{Hypergeometric2F1}\left(2, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a^2} + \frac{ex^{1+n} \text{Hypergeometric2F1}\left(2, \frac{1+n}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a^2(1+n)}$$

input `Integrate[(d + e*x^n)/(a + c*x^(2*n))^2, x]`

output `(d*x*Hypergeometric2F1[2, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)])/a^2 + (e*x^(1 + n)*Hypergeometric2F1[2, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)])/(a^2*(1 + n))`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.99, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {1761, 1748, 778, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{d + ex^n}{(a + cx^{2n})^2} dx \\ & \quad \downarrow \textcolor{blue}{1761} \\ & \frac{x(d + ex^n)}{2an(a + cx^{2n})} - \frac{\int \frac{e(1-n)x^n + d(1-2n)}{cx^{2n}+a} dx}{2an} \\ & \quad \downarrow \textcolor{blue}{1748} \\ & \frac{x(d + ex^n)}{2an(a + cx^{2n})} - \frac{d(1-2n) \int \frac{1}{cx^{2n}+a} dx + e(1-n) \int \frac{x^n}{cx^{2n}+a} dx}{2an} \\ & \quad \downarrow \textcolor{blue}{778} \end{aligned}$$

$$\begin{aligned}
 & \frac{x(d + ex^n)}{2an(a + cx^{2n})} - \frac{e(1-n) \int \frac{x^n}{cx^{2n}+a} dx + \frac{d(1-2n)x \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a}}{2an} \\
 & \quad \downarrow \text{888} \\
 & \frac{x(d + ex^n)}{2an(a + cx^{2n})} - \\
 & \frac{\frac{d(1-2n)x \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a} + \frac{e(1-n)x^{n+1} \text{Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a(n+1)}}{2an}
 \end{aligned}$$

input `Int[(d + e*x^n)/(a + c*x^(2*n))^2, x]`

output `(x*(d + e*x^n))/(2*a*n*(a + c*x^(2*n))) - ((d*(1 - 2*n)*x*Hypergeometric2F1[1, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)])/a + (e*(1 - n)*x^(1 + n)*Hypergeometric2F1[1, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)])/(a*(1 + n)))/(2*a*n)`

Definitions of rubi rules used

rule 778 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F1[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] || GtQ[a, 0])`

rule 888 `Int[((c_)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 1748 `Int[((d_) + (e_.)*(x_)^(n_))/((a_) + (c_.)*(x_)^(n2_)), x_Symbol] := Simp[d Int[1/(a + c*x^(2*n)), x], x] + Simp[e Int[x^n/(a + c*x^(2*n)), x], x] /; FreeQ[{a, c, d, e, n}, x] && EqQ[n2, 2*n] && NeQ[c*d^2 + a*e^2, 0] && (PosQ[a*c] || !IntegerQ[n])`

rule 1761

```
Int[((d_) + (e_)*(x_)^(n_))*((a_) + (c_)*(x_)^(n2_))^(p_), x_Symbol] :> S
imp[(-x)*(d + e*x^n)*((a + c*x^(2*n))^(p + 1)/(2*a*n*(p + 1))), x] + Simp[1
/(2*a*n*(p + 1)) Int[(d*(2*n*p + 2*n + 1) + e*(2*n*p + 3*n + 1)*x^n)*(a +
c*x^(2*n))^(p + 1), x], x] /; FreeQ[{a, c, d, e, n}, x] && EqQ[n2, 2*n] &&
ILtQ[p, -1]
```

Maple [F]

$$\int \frac{d + e x^n}{(a + c x^{2n})^2} dx$$

input `int((d+e*x^n)/(a+c*x^(2*n))^2,x)`

output `int((d+e*x^n)/(a+c*x^(2*n))^2,x)`

Fricas [F]

$$\int \frac{d + e x^n}{(a + c x^{2n})^2} dx = \int \frac{e x^n + d}{(c x^{2n} + a)^2} dx$$

input `integrate((d+e*x^n)/(a+c*x^(2*n))^2,x, algorithm="fricas")`

output `integral((e*x^n + d)/(c^2*x^(4*n) + 2*a*c*x^(2*n) + a^2), x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 173.59 (sec) , antiderivative size = 994, normalized size of antiderivative = 7.42

$$\int \frac{d + e x^n}{(a + c x^{2n})^2} dx = \text{Too large to display}$$

input `integrate((d+e*x**n)/(a+c*x**2*n)**2,x)`

output

```
d*(2*a*a**1/(2*n))*a**(-2 - 1/(2*n))*n*x*lerchphi(c*x**2*n)*exp_polar(I*pi)/a, 1, 1/(2*n))*gamma(1/(2*n))/(8*a*n**3*gamma(1 + 1/(2*n)) + 8*c*n**3*x**2*n)*gamma(1 + 1/(2*n)) + 2*a*a**1/(2*n))*a**(-2 - 1/(2*n))*n*x*gamm a(1/(2*n))/(8*a*n**3*gamma(1 + 1/(2*n)) + 8*c*n**3*x**2*n)*gamma(1 + 1/(2*n)) - a*a**1/(2*n))*a**(-2 - 1/(2*n))*x*lerchphi(c*x**2*n)*exp_polar(I*pi)/a, 1, 1/(2*n))*gamma(1/(2*n))/(8*a*n**3*gamma(1 + 1/(2*n)) + 8*c*n**3*x**2*n)*gamma(1 + 1/(2*n)) + 2*a**1/(2*n))*a**(-2 - 1/(2*n))*c*n*x*x**2*n)*lerchphi(c*x**2*n)*exp_polar(I*pi)/a, 1, 1/(2*n))*gamma(1/(2*n))/(8*a*n**3*gamma(1 + 1/(2*n)) + 8*c*n**3*x**2*n)*gamma(1 + 1/(2*n)) - a**1/(2*n))*a**(-2 - 1/(2*n))*c*x*x**2*n)*lerchphi(c*x**2*n)*exp_polar(I*pi)/a, 1, 1/(2*n))*gamma(1/(2*n))/(8*a*n**3*gamma(1 + 1/(2*n)) + 8*c*n**3*x**2*n)*gamma(1 + 1/(2*n))) + e*(a*a**(-5/2 - 1/(2*n)))*a**1/2 + 1/(2*n))*n**2*x**2*(n + 1)*lerchphi(c*x**2*n)*exp_polar(I*pi)/a, 1, 1/2 + 1/(2*n))*g amma(1/2 + 1/(2*n))/(8*a*n**3*gamma(3/2 + 1/(2*n)) + 8*c*n**3*x**2*n)*gamma(3/2 + 1/(2*n)) + 2*a*a**(-5/2 - 1/(2*n)))*a**1/2 + 1/(2*n))*n**2*x**2*(n + 1)*gamma(1/2 + 1/(2*n))/(8*a*n**3*gamma(3/2 + 1/(2*n)) + 8*c*n**3*x**2*n)*gamma(3/2 + 1/(2*n)) + 2*a*a**(-5/2 - 1/(2*n)))*a**1/2 + 1/(2*n))*n*x*(n + 1)*gamma(1/2 + 1/(2*n))/(8*a*n**3*gamma(3/2 + 1/(2*n)) + 8*c*n**3*x**2*n)*gamma(3/2 + 1/(2*n)) - a*a**(-5/2 - 1/(2*n)))*a**1/2 + 1/(2*n))*x*(n + 1)*lerchphi(c*x**2*n)*exp_polar(I*pi)/a, 1, 1/2 + 1/(2*n))*gamma...
```

Maxima [F]

$$\int \frac{d + ex^n}{(a + cx^{2n})^2} dx = \int \frac{ex^n + d}{(cx^{2n} + a)^2} dx$$

input

```
integrate((d+e*x^n)/(a+c*x^(2*n))^2,x, algorithm="maxima")
```

output

```
1/2*(e*x*x^n + d*x)/(a*c*n*x^(2*n) + a^2*n) + integrate(1/2*(e*(n - 1)*x^n + d*(2*n - 1))/(a*c*n*x^(2*n) + a^2*n), x)
```

Giac [F]

$$\int \frac{d + ex^n}{(a + cx^{2n})^2} dx = \int \frac{ex^n + d}{(cx^{2n} + a)^2} dx$$

input `integrate((d+e*x^n)/(a+c*x^(2*n))^2,x, algorithm="giac")`

output `integrate((e*x^n + d)/(c*x^(2*n) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{d + ex^n}{(a + cx^{2n})^2} dx = \int \frac{d + e x^n}{(a + c x^{2n})^2} dx$$

input `int((d + e*x^n)/(a + c*x^(2*n))^2,x)`

output `int((d + e*x^n)/(a + c*x^(2*n))^2, x)`

Reduce [F]

$$\int \frac{d + ex^n}{(a + cx^{2n})^2} dx = \left(\int \frac{x^n}{x^{4n}c^2 + 2x^{2n}ac + a^2} dx \right) e + \left(\int \frac{1}{x^{4n}c^2 + 2x^{2n}ac + a^2} dx \right) d$$

input `int((d+e*x^n)/(a+c*x^(2*n))^2,x)`

output `int(x**n/(x**(4*n)*c**2 + 2*x**(2*n)*a*c + a**2),x)*e + int(1/(x**(4*n)*c**2 + 2*x**(2*n)*a*c + a**2),x)*d`

3.35 $\int \frac{1}{(d+ex^n)(a+cx^{2n})^2} dx$

Optimal result	269
Mathematica [A] (verified)	270
Rubi [A] (verified)	270
Maple [F]	272
Fricas [F]	272
Sympy [F(-1)]	272
Maxima [F]	273
Giac [F]	273
Mupad [F(-1)]	273
Reduce [F]	274

Optimal result

Integrand size = 21, antiderivative size = 249

$$\begin{aligned} & \int \frac{1}{(d+ex^n)(a+cx^{2n})^2} dx \\ &= \frac{cx(d-ex^n)}{2a(cd^2+ae^2)n(a+cx^{2n})} \\ & - \frac{cd(ae^2(1-4n)+cd^2(1-2n))x\text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{2a^2(cd^2+ae^2)^2n} \\ & + \frac{e^4x\text{Hypergeometric2F1}\left(1, \frac{1}{n}, 1+\frac{1}{n}, -\frac{ex^n}{d}\right)}{d(cd^2+ae^2)^2} \\ & + \frac{ce(ae^2(1-3n)+cd^2(1-n))x^{1+n}\text{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{2a^2(cd^2+ae^2)^2n(1+n)} \end{aligned}$$

output

```
1/2*c*x*(d-e*x^n)/a/(a*e^2+c*d^2)/n/(a+c*x^(2*n))-1/2*c*d*(a*e^2*(1-4*n)+c*d^2*(1-2*n))*x*hypergeom([1, 1/2/n], [1+1/2/n], -c*x^(2*n)/a)/a^2/(a*e^2+c*d^2)^2/n+e^4*x*hypergeom([1, 1/n], [1+1/n], -e*x^n/d)/d/(a*e^2+c*d^2)^2+1/2*c*e*(a*e^2*(1-3*n)+c*d^2*(1-n))*x^(1+n)*hypergeom([1, 1/2*(1+n)/n], [3/2+1/2/n], -c*x^(2*n)/a)/a^2/(a*e^2+c*d^2)^2/n/(1+n)
```

Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 227, normalized size of antiderivative = 0.91

$$\int \frac{1}{(d + ex^n)(a + cx^{2n})^2} dx \\ = \frac{x \left(acd^2 e^2 (1 + n) \text{Hypergeometric2F1} \left(1, \frac{1}{2n}, \frac{1}{2} \left(2 + \frac{1}{n} \right), -\frac{cx^{2n}}{a} \right) + a^2 e^4 (1 + n) \text{Hypergeometric2F1} \left(1, \frac{1}{n}, \frac{1}{2} \left(2 + \frac{1}{n} \right), -\frac{ex^n}{a} \right) \right)}{(a + cx^{2n})^2}$$

input `Integrate[1/((d + e*x^n)*(a + c*x^(2*n))^2),x]`

output
$$(x*(a*c*d^2*e^2*(1+n)*\text{Hypergeometric2F1}[1, 1/(2*n), (2+n^{-1})/2, -(c*x^(2*n))/a]) + a^2*e^4*(1+n)*\text{Hypergeometric2F1}[1, n^{-1}, 1+n^{-1}, -((e*x^n)/d)] + c*d*(-(a*e^3*x^n*\text{Hypergeometric2F1}[1, (1+n)/(2*n), (3+n^{-1})/2, -((c*x^(2*n))/a)]) + (c*d^2 + a*e^2)*(d*(1+n)*\text{Hypergeometric2F1}[2, 1/(2*n), (2+n^{-1})/2, -((c*x^(2*n))/a)] - e*x^n*\text{Hypergeometric2F1}[2, (1+n)/(2*n), (3+n^{-1})/2, -((c*x^(2*n))/a)])))/(a^2*d*(c*d^2 + a*e^2)^2*(1+n))$$

Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 333, normalized size of antiderivative = 1.34, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {1767, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + cx^{2n})^2 (d + ex^n)} dx \\ \downarrow 1767 \\ \int \left(-\frac{ce^2(ex^n - d)}{(ae^2 + cd^2)^2 (a + cx^{2n})} - \frac{c(ex^n - d)}{(ae^2 + cd^2)(a + cx^{2n})^2} + \frac{e^4}{(ae^2 + cd^2)^2 (d + ex^n)} \right) dx \\ \downarrow 2009$$

$$\begin{aligned}
& \frac{ce(1-n)x^{n+1} \text{Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{2a^2n(n+1)(ae^2+cd^2)} - \\
& \frac{cd(1-2n)x \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{2a^2n(ae^2+cd^2)} + \\
& \frac{cde^2x \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a(ae^2+cd^2)^2} + \frac{cx(d-ex^n)}{2an(ae^2+cd^2)(a+cx^{2n})} + \\
& \frac{e^4x \text{Hypergeometric2F1}\left(1, \frac{1}{n}, 1+\frac{1}{n}, -\frac{ex^n}{d}\right)}{d(ae^2+cd^2)^2} - \\
& \frac{ce^3x^{n+1} \text{Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a(n+1)(ae^2+cd^2)^2}
\end{aligned}$$

input `Int[1/((d + e*x^n)*(a + c*x^(2*n))^2), x]`

output
$$\begin{aligned}
& \frac{(c*x*(d - e*x^n))/(2*a*(c*d^2 + a*e^2)*n*(a + c*x^(2*n))) + (c*d*e^2*x*\text{Hypergeometric2F1}[1, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)])/(a*(c*d^2 + a*e^2)^2) - (c*d*(1 - 2*n)*x*\text{Hypergeometric2F1}[1, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)])/(2*a^2*(c*d^2 + a*e^2)*n) + (e^4*x*\text{Hypergeometric2F1}[1, n^(-1), 1 + n^(-1), -((e*x^n)/d)])/(d*(c*d^2 + a*e^2)^2) - (c*e^3*x^(1 + n)*\text{Hypergeometric2F1}[1, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)])/(a*(c*d^2 + a*e^2)^2*(1 + n)) + (c*e*(1 - n)*x^(1 + n)*\text{Hypergeometric2F1}[1, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)])/(2*a^2*(c*d^2 + a*e^2)*n*(1 + n))
\end{aligned}$$

Defintions of rubi rules used

rule 1767 `Int[((d_) + (e_.)*(x_)^(n_))^(q_)*((a_) + (c_.)*(x_)^(n2_))^(p_), x_Symbol] :> Int[ExpandIntegrand[(d + e*x^n)^q*(a + c*x^(2*n))^p, x], x]; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && NeQ[c*d^2 + a*e^2, 0] && ((IntegersQ[p, q] && !IntegerQ[n]) || IGtQ[p, 0] || (IGtQ[q, 0] && !IntegerQ[n]))`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [F]

$$\int \frac{1}{(d + e x^n) (a + c x^{2n})^2} dx$$

input `int(1/(d+e*x^n)/(a+c*x^(2*n))^2,x)`

output `int(1/(d+e*x^n)/(a+c*x^(2*n))^2,x)`

Fricas [F]

$$\int \frac{1}{(d + e x^n) (a + c x^{2n})^2} dx = \int \frac{1}{(c x^{2n} + a)^2 (e x^n + d)} dx$$

input `integrate(1/(d+e*x^n)/(a+c*x^(2*n))^2,x, algorithm="fricas")`

output `integral(1/(a^2*e*x^n + a^2*d + (c^2*e*x^n + c^2*d)*x^(4*n) + 2*(a*c*e*x^n + a*c*d)*x^(2*n)), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(d + e x^n) (a + c x^{2n})^2} dx = \text{Timed out}$$

input `integrate(1/(d+e*x**n)/(a+c*x**2*n)**2,x)`

output `Timed out`

Maxima [F]

$$\int \frac{1}{(d + ex^n)(a + cx^{2n})^2} dx = \int \frac{1}{(cx^{2n} + a)^2(ex^n + d)} dx$$

input `integrate(1/(d+e*x^n)/(a+c*x^(2*n))^2,x, algorithm="maxima")`

output `e^4*integrate(1/(c^2*d^5 + 2*a*c*d^3*e^2 + a^2*d*e^4 + (c^2*d^4*e + 2*a*c*d^2*e^3 + a^2*e^5)*x^n), x) - 1/2*(c*e*x*x^n - c*d*x)/(a^2*c*d^2*n + a^3*e^2*n + (a*c^2*d^2*n + a^2*c*e^2*n)*x^(2*n)) - integrate(-1/2*(a*c*d*e^2*(4*n - 1) + c^2*d^3*(2*n - 1) - (a*c*e^3*(3*n - 1) + c^2*d^2*e*(n - 1))*x^n)/(a^2*c^2*d^4*n + 2*a^3*c*d^2*e^2*n + a^4*e^4*n + (a*c^3*d^4*n + 2*a^2*c^2*d^2*e^2*n + a^3*c*e^4*n)*x^(2*n)), x)`

Giac [F]

$$\int \frac{1}{(d + ex^n)(a + cx^{2n})^2} dx = \int \frac{1}{(cx^{2n} + a)^2(ex^n + d)} dx$$

input `integrate(1/(d+e*x^n)/(a+c*x^(2*n))^2,x, algorithm="giac")`

output `integrate(1/((c*x^(2*n) + a)^2*(e*x^n + d)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex^n)(a + cx^{2n})^2} dx = \int \frac{1}{(a + cx^{2n})^2(d + ex^n)} dx$$

input `int(1/((a + c*x^(2*n))^2*(d + e*x^n)),x)`

output `int(1/((a + c*x^(2*n))^2*(d + e*x^n)), x)`

Reduce [F]

$$\int \frac{1}{(d + ex^n)(a + cx^{2n})^2} dx = \int \frac{1}{x^{5n}c^2e + x^{4n}c^2d + 2x^{3n}ace + 2x^{2n}acd + x^na^2e + a^2d} dx$$

input `int(1/(d+e*x^n)/(a+c*x^(2*n))^2,x)`

output `int(1/(x**(5*n)*c**2*e + x**(4*n)*c**2*d + 2*x**((3*n)*a*c*e + 2*x**((2*n)*a*c*d + x**n*a**2*e + a**2*d)),x)`

3.36 $\int \frac{1}{(d+ex^n)^2(a+cx^{2n})^2} dx$

Optimal result	275
Mathematica [A] (verified)	276
Rubi [A] (verified)	276
Maple [F]	278
Fricas [F]	278
Sympy [F(-1)]	279
Maxima [F]	279
Giac [F]	280
Mupad [F(-1)]	280
Reduce [F]	280

Optimal result

Integrand size = 21, antiderivative size = 347

$$\begin{aligned} & \int \frac{1}{(d+ex^n)^2(a+cx^{2n})^2} dx \\ &= -\frac{e^2(cd^2 - ae^2)x}{ad(cd^2 + ae^2)^2 n(d+ex^n)} + \frac{cx(d-ex^n)}{2a(cd^2 + ae^2)n(d+ex^n)(a+cx^{2n})} \\ &+ \frac{c(a^2e^4(1-4n) - c^2d^4(1-2n) + 6acd^2e^2n)x \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{2a^2(cd^2 + ae^2)^3 n} \\ &- \frac{e^4(cd^2(1-5n) + ae^2(1-n))x \text{Hypergeometric2F1}\left(1, \frac{1}{n}, 1+\frac{1}{n}, -\frac{ex^n}{d}\right)}{d^2(cd^2 + ae^2)^3 n} \\ &+ \frac{c^2de(ae^2(1-5n) + cd^2(1-n))x^{1+n} \text{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a^2(cd^2 + ae^2)^3 n(1+n)} \end{aligned}$$

output

```
-e^2*(-a*e^2+c*d^2)*x/a/d/(a*e^2+c*d^2)^2/n/(d+e*x^n)+1/2*c*x*(d-e*x^n)/a/
(a*e^2+c*d^2)/n/(d+e*x^n)/(a+c*x^(2*n))+1/2*c*(a^2*e^4*(1-4*n)-c^2*d^4*(1-
2*n)+6*a*c*d^2*e^2*n)*x*hypergeom([1, 1/2*n], [1+1/2/n], -c*x^(2*n)/a)/a^2/(
a*e^2+c*d^2)^3/n-e^4*(c*d^2*(1-5*n)+a*e^2*(1-n))*x*hypergeom([1, 1/n], [1+1
/n], -e*x^n/d)/d^2/(a*e^2+c*d^2)^3/n+c^2*d*e*(a*e^2*(1-5*n)+c*d^2*(1-n))*x^
(1+n)*hypergeom([1, 1/2*(1+n)/n], [3/2+1/2/n], -c*x^(2*n)/a)/a^2/(a*e^2+c*d^
2)^3/n/(1+n)
```

Mathematica [A] (verified)

Time = 0.65 (sec) , antiderivative size = 298, normalized size of antiderivative = 0.86

$$\int \frac{1}{(d + ex^n)^2 (a + cx^{2n})^2} dx \\ = \frac{x \left(\frac{ce^2 (3cd^2 - ae^2) \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a} + 4ce^4 \text{Hypergeometric2F1}\left(1, \frac{1}{n}, 1 + \frac{1}{n}, -\frac{ex^n}{d}\right) - \frac{4c^2 de^3 x}{a^2} \right)}{(d + ex^n)^2 (a + cx^{2n})^2}$$

input `Integrate[1/((d + e*x^n)^2*(a + c*x^(2*n))^2), x]`

output
$$(x*((c*e^2*(3*c*d^2 - a*e^2)*\text{Hypergeometric2F1}[1, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)])/a + 4*c*e^4*\text{Hypergeometric2F1}[1, n^(-1), 1 + n^(-1), -((e*x^n)/d)] - (4*c^2*d*e^3*x^n*\text{Hypergeometric2F1}[1, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)])/(a*(1 + n)) + (c*(c*d^2 - a*e^2)*(c*d^2 + a*e^2)*\text{Hypergeometric2F1}[2, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)])/a^2 + (e^4*(c*d^2 + a*e^2)*\text{Hypergeometric2F1}[2, n^(-1), 1 + n^(-1), -((e*x^n)/d)])/d^2 - (2*c^2*d*e*(c*d^2 + a*e^2)*x^n*\text{Hypergeometric2F1}[2, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)])/(a^2*(1 + n))))/(c*d^2 + a*e^2)^3$$

Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 410, normalized size of antiderivative = 1.18, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {1767, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + cx^{2n})^2 (d + ex^n)^2} dx \\ \downarrow \textcolor{blue}{1767}$$

$$\int \left(-\frac{ce^2(ae^2 - 3cd^2 + 4cdex^n)}{(ae^2 + cd^2)^3 (a + cx^{2n})} - \frac{c(ae^2 - cd^2 + 2cdex^n)}{(ae^2 + cd^2)^2 (a + cx^{2n})^2} + \frac{4cde^4}{(ae^2 + cd^2)^3 (d + ex^n)} + \frac{e^4}{(ae^2 + cd^2)^2 (d + ex^n)^2} \right) dx$$

$$\begin{aligned}
 & \downarrow \text{2009} \\
 & \frac{c^2 de(1-n)x^{n+1} \text{Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a^2 n(n+1)(ae^2 + cd^2)^2} - \\
 & \frac{c(1-2n)x(cd^2 - ae^2) \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{2a^2 n (ae^2 + cd^2)^2} - \\
 & \frac{4c^2 de^3 x^{n+1} \text{Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a(n+1)(ae^2 + cd^2)^3} + \\
 & \frac{ce^2 x(3cd^2 - ae^2) \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a(ae^2 + cd^2)^3} + \\
 & \frac{cx(-ae^2 + cd^2 - 2cdex^n)}{2an(ae^2 + cd^2)^2(a + cx^{2n})} + \frac{4ce^4 x \text{Hypergeometric2F1}\left(1, \frac{1}{n}, 1 + \frac{1}{n}, -\frac{ex^n}{d}\right)}{(ae^2 + cd^2)^3} + \\
 & \frac{e^4 x \text{Hypergeometric2F1}\left(2, \frac{1}{n}, 1 + \frac{1}{n}, -\frac{ex^n}{d}\right)}{d^2 (ae^2 + cd^2)^2}
 \end{aligned}$$

input `Int[1/((d + e*x^n)^2*(a + c*x^(2*n))^2),x]`

output

$$\begin{aligned}
 & \frac{(c*x*(c*d^2 - a*e^2 - 2*c*d*e*x^n))/(2*a*(c*d^2 + a*e^2)^2*n*(a + c*x^(2*n))) + (c*e^2*(3*c*d^2 - a*e^2)*x*\text{Hypergeometric2F1}[1, 1/(2*n), (2 + n^{-1})/2, -((c*x^(2*n))/a)])/(a*(c*d^2 + a*e^2)^3) - (c*(c*d^2 - a*e^2)*(1 - 2*n)*x*\text{Hypergeometric2F1}[1, 1/(2*n), (2 + n^{-1})/2, -((c*x^(2*n))/a)])/(2*a^2*(c*d^2 + a*e^2)^2*n) + (4*c*e^4*x*\text{Hypergeometric2F1}[1, n^{-1}, 1 + n^{-1}, -((e*x^n)/d)])/(c*d^2 + a*e^2)^3 - (4*c^2*d*e^3*x^(1 + n))*\text{Hypergeometric2F1}[1, (1 + n)/(2*n), (3 + n^{-1})/2, -((c*x^(2*n))/a)])/(a*(c*d^2 + a*e^2)^3*(1 + n)) + (c^2*d*e*(1 - n)*x^(1 + n))*\text{Hypergeometric2F1}[1, (1 + n)/(2*n), (3 + n^{-1})/2, -((c*x^(2*n))/a)])/(a^2*(c*d^2 + a*e^2)^2*n*(1 + n)) + (e^4*x*\text{Hypergeometric2F1}[2, n^{-1}, 1 + n^{-1}, -((e*x^n)/d)])/(d^2*(c*d^2 + a*e^2)^2)
 \end{aligned}$$

Defintions of rubi rules used

rule 1767 $\text{Int}[(d_+ + e_-) \cdot (x_-)^{(n_-)} \cdot (q_-) \cdot ((a_- + c_-) \cdot (x_-)^{(n_2_-)})^{(p_-)}, x_{\text{Symbol}}] \\ :> \text{Int}[\text{ExpandIntegrand}[(d + e \cdot x^n)^q \cdot (a + c \cdot x^{(2 \cdot n)})^p, x], x] /; \text{FreeQ}[\{a, c, d, e, n, p, q\}, x] \&& \text{EqQ}[n_2, 2 \cdot n] \&& \text{NeQ}[c \cdot d^2 + a \cdot e^2, 0] \&& ((\text{IntegersQ}[p, q] \&& !\text{IntegerQ}[n]) \mid\mid \text{IGtQ}[p, 0] \mid\mid (\text{IGtQ}[q, 0] \&& !\text{IntegerQ}[n]))$

rule 2009 $\text{Int}[u_-, x_{\text{Symbol}}] :> \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

Maple [F]

$$\int \frac{1}{(d + e x^n)^2 (a + c x^{2n})^2} dx$$

input $\text{int}(1/(d+e*x^n)^2/(a+c*x^(2*n))^2, x)$

output $\text{int}(1/(d+e*x^n)^2/(a+c*x^(2*n))^2, x)$

Fricas [F]

$$\int \frac{1}{(d + e x^n)^2 (a + c x^{2n})^2} dx = \int \frac{1}{(c x^{2n} + a)^2 (e x^n + d)^2} dx$$

input $\text{integrate}(1/(d+e*x^n)^2/(a+c*x^(2*n))^2, x, \text{algorithm}=\text{"fricas"})$

output $\text{integral}(1/(a^{2*}e^{2*}x^{(2*n)} + 2*a^{2*d}*e*x^n + a^{2*d^2} + (c^{2*}e^{2*}x^{(2*n)} + 2*c^{2*d}*e*x^n + c^{2*d^2})*x^{(4*n)} + 2*(a*c*e^{2*}x^{(2*n)} + 2*a*c*d*e*x^n + a*c*d^2)*x^{(2*n)}), x)$

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex^n)^2 (a + cx^{2n})^2} dx = \text{Timed out}$$

input `integrate(1/(d+e*x**n)**2/(a+c*x**(2*n))**2,x)`

output `Timed out`

Maxima [F]

$$\int \frac{1}{(d + ex^n)^2 (a + cx^{2n})^2} dx = \int \frac{1}{(cx^{2n} + a)^2 (ex^n + d)^2} dx$$

input `integrate(1/(d+e*x^n)^2/(a+c*x^(2*n))^2,x, algorithm="maxima")`

output `(c*d^2*e^4*(5*n - 1) + a*e^6*(n - 1))*integrate(1/(c^3*d^8*n + 3*a*c^2*d^6 *e^2*n + 3*a^2*c*d^4*e^4*n + a^3*d^2*e^6*n + (c^3*d^7*e*n + 3*a*c^2*d^5*e^3*n + 3*a^2*c*d^3*e^5*n + a^3*d^4*e^7*n)*x*n), x) - 1/2*(2*(c^2*d^2*e^2 - a*c^4)*x*x^(2*n) + (c^2*d^3*e + a*c*d^3*e^3)*x*x^(2*n) - (c^2*d^4 - a*c*d^2*e^2 + 2*a^2*e^4)*x)/(a^2*c^2*d^6*n + 2*a^3*c*d^4*e^2*n + a^4*d^2*e^4*n + (a*c^3*d^5*e*n + 2*a^2*c^2*d^3*e^3*n + a^3*c*d^5*n)*x^(3*n) + (a*c^3*d^6*n + 2*a^2*c^2*d^4*e^2*n + a^3*c*d^2*e^4*n)*x^(2*n) + (a^2*c^2*d^5*e*n + 2*a^3*c^3*e^3*n + a^4*d^3*e^5*n)*x^(2*n)) - integrate(1/2*(a^2*c*e^4*(4*n - 1) - c^3*d^4*(2*n - 1) - 6*a*c^2*d^2*e^2*n + 2*(a*c^2*d^3*(5*n - 1) + c^3*d^3*e^(n - 1))*x*n)/(a^2*c^3*d^6*n + 3*a^3*c^2*d^4*e^2*n + 3*a^4*c*d^2*e^4*n + a^5*e^6*n + (a*c^4*d^6*n + 3*a^2*c^3*d^4*e^2*n + 3*a^3*c^2*d^2*e^4*n + a^4*c^4*e^6*n)*x^(2*n)), x)`

Giac [F]

$$\int \frac{1}{(d + ex^n)^2 (a + cx^{2n})^2} dx = \int \frac{1}{(cx^{2n} + a)^2 (ex^n + d)^2} dx$$

input `integrate(1/(d+e*x^n)^2/(a+c*x^(2*n))^2,x, algorithm="giac")`

output `integrate(1/((c*x^(2*n) + a)^2*(e*x^n + d)^2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex^n)^2 (a + cx^{2n})^2} dx = \int \frac{1}{(a + cx^{2n})^2 (d + ex^n)^2} dx$$

input `int(1/((a + c*x^(2*n))^2*(d + e*x^n)^2),x)`

output `int(1/((a + c*x^(2*n))^2*(d + e*x^n)^2), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{1}{(d + ex^n)^2 (a + cx^{2n})^2} dx \\ &= \int \frac{1}{x^{6n}c^2e^2 + 2x^{5n}c^2de + 2x^{4n}ace^2 + x^{4n}c^2d^2 + 4x^{3n}acde + x^{2n}a^2e^2 + 2x^{2n}acd^2 + 2x^na^2de + a^2d^2} dx \end{aligned}$$

input `int(1/(d+e*x^n)^2/(a+c*x^(2*n))^2,x)`

output `int(1/(x**6*c**2*e**2 + 2*x**5*c**2*d*e + 2*x**4*a*c*e**2 + x**4*c**2*d**2 + 4*x**3*a*c*d*e + x**2*a**2*e**2 + 2*x**2*a*c*d**2 + 2*x**n*a**2*d*e + a**2*d**2),x)`

3.37 $\int \frac{(d+ex^n)^3}{(a+cx^{2n})^3} dx$

Optimal result	281
Mathematica [A] (verified)	282
Rubi [A] (verified)	282
Maple [F]	284
Fricas [F]	284
Sympy [F(-1)]	284
Maxima [F]	285
Giac [F]	285
Mupad [F(-1)]	285
Reduce [F]	286

Optimal result

Integrand size = 21, antiderivative size = 259

$$\begin{aligned} \int \frac{(d+ex^n)^3}{(a+cx^{2n})^3} dx &= \frac{x(d+ex^n)^3}{4an(a+cx^{2n})^2} \\ &- \frac{x(d(cd^2(1-4n)-3ae^2(1-2n))+e(3cd^2(1-3n)-ae^2(1-n))x^n)}{8a^2cn^2(a+cx^{2n})} \\ &- \frac{d(3ae^2-cd^2(1-4n))(1-2n)x \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{8a^3cn^2} \\ &+ \frac{e(1-n)(3cd^2(1-3n)-ae^2(1+n))x^{1+n} \text{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{8a^3cn^2(1+n)} \end{aligned}$$

output

```
1/4*x*(d+e*x^n)^3/a/n/(a+c*x^(2*n))^2-1/8*x*(d*(c*d^2*(1-4*n)-3*a*e^2*(1-2*n))+e*(3*c*d^2*(1-3*n)-a*e^2*(1-n))*x^n)/a^2/c/n^2/(a+c*x^(2*n))-1/8*d*(3*a*e^2-c*d^2*(1-4*n))*(1-2*n)*x*hypergeom([1, 1/2/n], [1+1/2/n], -c*x^(2*n)/a)/a^3/c/n^2+1/8*e*(1-n)*(3*c*d^2*(1-3*n)-a*e^2*(1+n))*x^(1+n)*hypergeom([1, 1/2*(1+n)/n], [3/2+1/2/n], -c*x^(2*n)/a)/a^3/c/n^2/(1+n)
```

Mathematica [A] (verified)

Time = 0.90 (sec) , antiderivative size = 188, normalized size of antiderivative = 0.73

$$\int \frac{(d + ex^n)^3}{(a + cx^{2n})^3} dx \\ = \frac{x \left(3ade^2 \text{Hypergeometric2F1} \left(2, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a} \right) + \frac{ae^3 x^n \text{Hypergeometric2F1} \left(2, \frac{1+n}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a} \right)}{1+n} \right) + d(c a e^2 x^{2n} \text{Hypergeometric2F1} \left(2, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a} \right) + \frac{ae^3 x^n \text{Hypergeometric2F1} \left(2, \frac{1+n}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a} \right)}{1+n})}{(a + cx^{2n})^3}$$

input `Integrate[(d + e*x^n)^3/(a + c*x^(2*n))^3, x]`

output
$$(x*(3*a*d*e^2*\text{Hypergeometric2F1}[2, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)] + (a*e^3*x^n*\text{Hypergeometric2F1}[2, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)]/(1 + n) + d*(c*d^2 - 3*a*e^2)*\text{Hypergeometric2F1}[3, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)] + (e*(3*c*d^2 - a*e^2)*x^n*\text{Hypergeometric2F1}[3, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)]/(1 + n)))/(a^3*c))$$

Rubi [A] (verified)

Time = 0.61 (sec) , antiderivative size = 424, normalized size of antiderivative = 1.64, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {1767, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^n)^3}{(a + cx^{2n})^3} dx \\ \downarrow 1767 \\ \int \left(\frac{x^n(3cd^2e - ae^3) - 3ade^2 + cd^3}{c(a + cx^{2n})^3} + \frac{e^2(3d + ex^n)}{c(a + cx^{2n})^2} \right) dx \\ \downarrow 2009$$

$$\begin{aligned}
& \frac{e(1-3n)(1-n)x^{n+1}(3cd^2 - ae^2) \text{Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{8a^3cn^2(n+1)} + \\
& \frac{d(1-4n)(1-2n)x(cd^2 - 3ae^2) \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{8a^3cn^2} - \\
& \frac{x(e(1-3n)x^n(3cd^2 - ae^2) + d(1-4n)(cd^2 - 3ae^2))}{8a^2cn^2(a + cx^{2n})} - \\
& \frac{3de^2(1-2n)x \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{2a^2cn} - \\
& \frac{e^3(1-n)x^{n+1} \text{Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{2a^2cn(n+1)} + \\
& \frac{x(ex^n(3cd^2 - ae^2) + d(cd^2 - 3ae^2))}{4acn(a + cx^{2n})^2} + \frac{e^2x(3d + ex^n)}{2acn(a + cx^{2n})}
\end{aligned}$$

input `Int[(d + e*x^n)^3/(a + c*x^(2*n))^3, x]`

output

$$\begin{aligned}
& \frac{(x*(d*(c*d^2 - 3*a*e^2) + e*(3*c*d^2 - a*e^2)*x^n))/(4*a*c*n*(a + c*x^(2*n))^2) + (e^2*x*(3*d + e*x^n))/(2*a*c*n*(a + c*x^(2*n))) - (x*(d*(c*d^2 - 3*a*e^2)*(1 - 4*n) + e*(3*c*d^2 - a*e^2)*(1 - 3*n)*x^n))/(8*a^2*c*n^2*(a + c*x^(2*n))) + (d*(c*d^2 - 3*a*e^2)*(1 - 4*n)*(1 - 2*n)*x*\text{Hypergeometric2F1}[1, 1/(2*n), (2 + n^{(-1)})/2, -((c*x^(2*n))/a)])/(8*a^3*c*n^2) - (3*d*e^2*(1 - 2*n)*x*\text{Hypergeometric2F1}[1, 1/(2*n), (2 + n^{(-1)})/2, -((c*x^(2*n))/a)])/(2*a^2*c*n) + (e*(3*c*d^2 - a*e^2)*(1 - 3*n)*(1 - n)*x^(1 + n)*\text{Hypergeometric2F1}[1, (1 + n)/(2*n), (3 + n^{(-1)})/2, -((c*x^(2*n))/a)])/(8*a^3*c*n^2*(1 + n)) - (e^3*(1 - n)*x^(1 + n)*\text{Hypergeometric2F1}[1, (1 + n)/(2*n), (3 + n^{(-1)})/2, -((c*x^(2*n))/a)])/(2*a^2*c*n*(1 + n))
\end{aligned}$$

Definitions of rubi rules used

rule 1767

```

Int[((d_) + (e_.)*(x_)^(n_))^(q_)*((a_) + (c_.)*(x_)^(n2_))^(p_), x_Symbol]
  :> Int[ExpandIntegrand[(d + e*x^n)^q*(a + c*x^(2*n))^p, x], x] /; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && NeQ[c*d^2 + a*e^2, 0] && ((IntegersQ[p, q] && !IntegerQ[n]) || IGtQ[p, 0] || (IGtQ[q, 0] && !IntegerQ[n]))
)

```

rule 2009

```

Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]

```

Maple [F]

$$\int \frac{(d + e x^n)^3}{(a + c x^{2n})^3} dx$$

input `int((d+e*x^n)^3/(a+c*x^(2*n))^3,x)`

output `int((d+e*x^n)^3/(a+c*x^(2*n))^3,x)`

Fricas [F]

$$\int \frac{(d + e x^n)^3}{(a + c x^{2n})^3} dx = \int \frac{(e x^n + d)^3}{(c x^{2n} + a)^3} dx$$

input `integrate((d+e*x^n)^3/(a+c*x^(2*n))^3,x, algorithm="fricas")`

output `integral((e^3*x^(3*n) + 3*d*e^2*x^(2*n) + 3*d^2*e*x^n + d^3)/(c^3*x^(6*n) + 3*a*c^2*x^(4*n) + 3*a^2*c*x^(2*n) + a^3), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{(d + e x^n)^3}{(a + c x^{2n})^3} dx = \text{Timed out}$$

input `integrate((d+e*x**n)**3/(a+c*x**2*n)**3,x)`

output `Timed out`

Maxima [F]

$$\int \frac{(d + ex^n)^3}{(a + cx^{2n})^3} dx = \int \frac{(ex^n + d)^3}{(cx^{2n} + a)^3} dx$$

input `integrate((d+e*x^n)^3/(a+c*x^(2*n))^3,x, algorithm="maxima")`

output
$$\begin{aligned} & 1/8*((3*c^2*d^2*e*(3*n - 1) + a*c*e^3*(n + 1))*x*x^(3*n) + (c^2*d^3*(4*n - 1) + 3*a*c*d*e^2)*x*x^(2*n) + (3*a*c*d^2*e*(5*n - 1) - a^2*e^3*(n - 1))*x*x^n + (a*c*d^3*(6*n - 1) - 3*a^2*d*e^2*(2*n - 1))*x)/(a^2*c^3*n^2*x^(4*n) + 2*a^3*c^2*n^2*x^(2*n) + a^4*c*n^2) + \text{integrate}(1/8*((8*n^2 - 6*n + 1)*c*d^3 + 3*a*d*e^2*(2*n - 1) + (3*(3*n^2 - 4*n + 1)*c*d^2*e + (n^2 - 1)*a*e^3)*x^n)/(a^2*c^2*n^2*x^(2*n) + a^3*c*n^2), x) \end{aligned}$$

Giac [F]

$$\int \frac{(d + ex^n)^3}{(a + cx^{2n})^3} dx = \int \frac{(ex^n + d)^3}{(cx^{2n} + a)^3} dx$$

input `integrate((d+e*x^n)^3/(a+c*x^(2*n))^3,x, algorithm="giac")`

output `integrate((e*x^n + d)^3/(c*x^(2*n) + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(d + ex^n)^3}{(a + cx^{2n})^3} dx = \int \frac{(d + e x^n)^3}{(a + c x^{2n})^3} dx$$

input `int((d + e*x^n)^3/(a + c*x^(2*n))^3,x)`

output `int((d + e*x^n)^3/(a + c*x^(2*n))^3, x)`

Reduce [F]

$$\int \frac{(d + ex^n)^3}{(a + cx^{2n})^3} dx = \text{too large to display}$$

input `int((d+e*x^n)^3/(a+c*x^(2*n))^3,x)`

output
$$\begin{aligned} & (x^{**}(4*n)*\int(x^{**}(3*n)/(x^{**}(6*n)*c^{**3*n} + x^{**}(6*n)*c^{**3} + 3*x^{**}(4*n)*a*c^{**} \\ & 2*n + 3*x^{**}(4*n)*a*c^{**2} + 3*x^{**}(2*n)*a^{**2}*c*n + 3*x^{**}(2*n)*a^{**2}*c + a^{**3}*n \\ & + a^{**3}), x)*a*c^{**2}*e^{**3*n**2} + 2*x^{**}(4*n)*\int(x^{**}(3*n)/(x^{**}(6*n)*c^{**3*n} + \\ & x^{**}(6*n)*c^{**3} + 3*x^{**}(4*n)*a*c^{**2}*n + 3*x^{**}(4*n)*a*c^{**2} + 3*x^{**}(2*n)*a^{**2}* \\ & c*n + 3*x^{**}(2*n)*a^{**2}*c + a^{**3}*n + a^{**3}), x)*a*c^{**2}*e^{**3*n} + x^{**}(4*n)*\int(x \\ & **(3*n)/(x^{**}(6*n)*c^{**3*n} + x^{**}(6*n)*c^{**3} + 3*x^{**}(4*n)*a*c^{**2}*n + 3*x^{**}(4*n) \\ &)*a*c^{**2} + 3*x^{**}(2*n)*a^{**2}*c*n + 3*x^{**}(2*n)*a^{**2}*c + a^{**3}*n + a^{**3}), x)*a*c \\ & **2*e^{**3} + 9*x^{**}(4*n)*\int(x^{**}(3*n)/(x^{**}(6*n)*c^{**3*n} + x^{**}(6*n)*c^{**3} + 3*x^{**} \\ & (4*n)*a*c^{**2}*n + 3*x^{**}(4*n)*a*c^{**2} + 3*x^{**}(2*n)*a^{**2}*c*n + 3*x^{**}(2*n)*a^{**} \\ & 2*c + a^{**3}*n + a^{**3}), x)*c^{**3*d^{**2}*e*n**2} + 6*x^{**}(4*n)*\int(x^{**}(3*n)/(x^{**}(6* \\ & n)*c^{**3*n} + x^{**}(6*n)*c^{**3} + 3*x^{**}(4*n)*a*c^{**2}*n + 3*x^{**}(4*n)*a*c^{**2} + 3*x^{**} \\ & (2*n)*a^{**2}*c*n + 3*x^{**}(2*n)*a^{**2}*c + a^{**3}*n + a^{**3}), x)*c^{**3*d^{**2}*e*n} - 3*x^{**} \\ & (4*n)*\int(x^{**}(3*n)/(x^{**}(6*n)*c^{**3*n} + x^{**}(6*n)*c^{**3} + 3*x^{**}(4*n)*a*c^{**} \\ & 2*n + 3*x^{**}(4*n)*a*c^{**2} + 3*x^{**}(2*n)*a^{**2}*c*n + 3*x^{**}(2*n)*a^{**2}*c + a^{**3}*n \\ & + a^{**3}), x)*c^{**3*d^{**2}*e} + 3*x^{**}(4*n)*\int(x^{**}(2*n)/(x^{**}(6*n)*c^{**3*n} + x^{**}(6* \\ & n)*c^{**3} + 3*x^{**}(4*n)*a*c^{**2}*n + 3*x^{**}(4*n)*a*c^{**2} + 3*x^{**}(2*n)*a^{**2}*c*n + \\ & 3*x^{**}(2*n)*a^{**2}*c + a^{**3}*n + a^{**3}), x)*a*c^{**2*d^{**2}*e^{**2}*n**2} + 6*x^{**}(4*n)*\int(x^{**} \\ & (2*n)/(x^{**}(6*n)*c^{**3*n} + x^{**}(6*n)*c^{**3} + 3*x^{**}(4*n)*a*c^{**2}*n + 3*x^{**}(4*n) \\ & *a*c^{**2} + 3*x^{**}(2*n)*a^{**2}*c*n + 3*x^{**}(2*n)*a^{**2}*c + a^{**3}*n + a^{**3}), x)*a*c^{**} \\ & 2*d^{**2}*e^{**2}*n + 3*x^{**}(4*n)*\int(x^{**}(2*n)/(x^{**}(6*n)*c^{**3*n} + x^{**}(6*n)*c^{**}... \end{aligned}$$

$$3.38 \quad \int \frac{(d+ex^n)^2}{(a+cx^{2n})^3} dx$$

Optimal result	287
Mathematica [A] (verified)	288
Rubi [A] (verified)	288
Maple [F]	290
Fricas [F]	290
Sympy [F(-1)]	290
Maxima [F]	291
Giac [F]	291
Mupad [F(-1)]	291
Reduce [F]	292

Optimal result

Integrand size = 21, antiderivative size = 221

$$\begin{aligned} & \int \frac{(d+ex^n)^2}{(a+cx^{2n})^3} dx \\ &= \frac{x(d+ex^n)^2}{4an(a+cx^{2n})^2} - \frac{x(cd^2(1-4n) - ae^2(1-2n) + 2cde(1-3n)x^n)}{8a^2cn^2(a+cx^{2n})} \\ &\quad - \frac{(ae^2 - cd^2(1-4n))(1-2n)x \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{8a^3cn^2} \\ &\quad + \frac{de(1-3n)(1-n)x^{1+n} \operatorname{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{4a^3n^2(1+n)} \end{aligned}$$

output

```
1/4*x*(d+e*x^n)^2/a/n/(a+c*x^(2*n))^2-1/8*x*(c*d^2*(1-4*n)-a*e^2*(1-2*n)+2*c*d*e*(1-3*n)*x^n)/a^2/c/n^2/(a+c*x^(2*n))-1/8*(a*e^2-c*d^2*(1-4*n))*(1-2*n)*x*hypergeom([1, 1/2/n], [1+1/2/n], -c*x^(2*n)/a)/a^3/c/n^2+1/4*d*e*(1-3*n)*(1-n)*x^(1+n)*hypergeom([1, 1/2*(1+n)/n], [3/2+1/2/n], -c*x^(2*n)/a)/a^3/n^2/(1+n)
```

Mathematica [A] (verified)

Time = 0.53 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.62

$$\int \frac{(d + ex^n)^2}{(a + cx^{2n})^3} dx \\ = \frac{x \left(ae^2 (1+n) \text{Hypergeometric2F1} \left(2, \frac{1}{2n}, \frac{1}{2} \left(2 + \frac{1}{n} \right), -\frac{cx^{2n}}{a} \right) + (cd^2 - ae^2) (1+n) \text{Hypergeometric2F1} \left(3, \frac{1}{2n}, \frac{1}{2} \left(3 + \frac{1}{n} \right), -\frac{ex^{2n}}{a} \right) \right)}{a^3 c (1+n)}$$

input `Integrate[(d + e*x^n)^2/(a + c*x^(2*n))^3, x]`

output
$$(x*(a*e^2*(1+n)*\text{Hypergeometric2F1}[2, 1/(2*n), (2+n^(-1))/2, -((c*x^(2*n))/a)] + (c*d^2 - a*e^2)*(1+n)*\text{Hypergeometric2F1}[3, 1/(2*n), (2+n^(-1))/2, -((c*x^(2*n))/a)] + 2*c*d*e*x^n*\text{Hypergeometric2F1}[3, (1+n)/(2*n), (3+n^(-1))/2, -((c*x^(2*n))/a)]))/((a^3*c*(1+n)))$$

Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 272, normalized size of antiderivative = 1.23, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.095, Rules used = {1767, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^n)^2}{(a + cx^{2n})^3} dx \\ \downarrow 1767 \\ \int \left(\frac{-ae^2 + cd^2 + 2cdex^n}{c(a + cx^{2n})^3} + \frac{e^2}{c(a + cx^{2n})^2} \right) dx \\ \downarrow 2009$$

$$\begin{aligned}
 & \frac{(1 - 4n)(1 - 2n)x(cd^2 - ae^2) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{8a^3cn^2} + \\
 & \frac{de(1 - 3n)(1 - n)x^{n+1} \operatorname{Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{8a^3n^2(n+1)} - \\
 & \frac{x((1 - 4n)(cd^2 - ae^2) + 2cde(1 - 3n)x^n)}{8a^2cn^2(a + cx^{2n})} + \\
 & \frac{e^2x \operatorname{Hypergeometric2F1}\left(2, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a^2c} + \frac{x(-ae^2 + cd^2 + 2cdex^n)}{4acn(a + cx^{2n})^2}
 \end{aligned}$$

input `Int[(d + e*x^n)^2/(a + c*x^(2*n))^3, x]`

output `(x*(c*d^2 - a*e^2 + 2*c*d*e*x^n))/(4*a*c*n*(a + c*x^(2*n))^2) - (x*((c*d^2 - a*e^2)*(1 - 4*n) + 2*c*d*e*(1 - 3*n)*x^n)/(8*a^2*c*n^2*(a + c*x^(2*n))) + ((c*d^2 - a*e^2)*(1 - 4*n)*(1 - 2*n)*x*Hypergeometric2F1[1, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)])/(8*a^3*c*n^2) + (d*e*(1 - 3*n)*(1 - n)*x^(1 + n)*Hypergeometric2F1[1, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)])/(4*a^3*n^2*(1 + n)) + (e^2*x*Hypergeometric2F1[2, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)])/(a^2*c)`

Definitions of rubi rules used

rule 1767 `Int[((d_) + (e_.)*(x_)^(n_))^(q_)*((a_) + (c_.)*(x_)^(n2_))^(p_), x_Symbol] :> Int[ExpandIntegrand[(d + e*x^n)^q*(a + c*x^(2*n))^p, x], x]; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && NeQ[c*d^2 + a*e^2, 0] && ((IntegersQ[p, q] && !IntegerQ[n]) || IGtQ[p, 0] || (IGtQ[q, 0] && !IntegerQ[n]))`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [F]

$$\int \frac{(d + e x^n)^2}{(a + c x^{2n})^3} dx$$

input `int((d+e*x^n)^2/(a+c*x^(2*n))^3,x)`

output `int((d+e*x^n)^2/(a+c*x^(2*n))^3,x)`

Fricas [F]

$$\int \frac{(d + e x^n)^2}{(a + c x^{2n})^3} dx = \int \frac{(e x^n + d)^2}{(c x^{2n} + a)^3} dx$$

input `integrate((d+e*x^n)^2/(a+c*x^(2*n))^3,x, algorithm="fricas")`

output `integral((e^2*x^(2*n) + 2*d*e*x^n + d^2)/(c^3*x^(6*n) + 3*a*c^2*x^(4*n) + 3*a^2*c*x^(2*n) + a^3), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{(d + e x^n)^2}{(a + c x^{2n})^3} dx = \text{Timed out}$$

input `integrate((d+e*x**n)**2/(a+c*x**2*n)**3,x)`

output `Timed out`

Maxima [F]

$$\int \frac{(d + ex^n)^2}{(a + cx^{2n})^3} dx = \int \frac{(ex^n + d)^2}{(cx^{2n} + a)^3} dx$$

input `integrate((d+e*x^n)^2/(a+c*x^(2*n))^3,x, algorithm="maxima")`

output $\frac{1}{8} \cdot (2c^2 \cdot d \cdot e \cdot (3n - 1) \cdot x \cdot x^{(3n)} + 2 \cdot a \cdot c \cdot d \cdot e \cdot (5n - 1) \cdot x \cdot x^n + (c^2 \cdot d^2 \cdot (4n - 1) + a \cdot c \cdot e^2) \cdot x \cdot x^{(2n)} + (a \cdot c \cdot d^2 \cdot (6n - 1) - a^2 \cdot e^2 \cdot (2n - 1)) \cdot x) / (a^2 \cdot c^3 \cdot n^2 \cdot x^{(4n)} + 2 \cdot a^3 \cdot c^2 \cdot n^2 \cdot x^{(2n)} + a^4 \cdot c \cdot n^2) + \text{integrate}(1/8 \cdot (2 \cdot (3n^2 - 4n + 1) \cdot c \cdot d \cdot e \cdot x^n + (8n^2 - 6n + 1) \cdot c \cdot d^2 + a \cdot e^2 \cdot (2n - 1)) / (a^2 \cdot c^2 \cdot n^2 \cdot x^{(2n)} + a^3 \cdot c \cdot n^2), x)$

Giac [F]

$$\int \frac{(d + ex^n)^2}{(a + cx^{2n})^3} dx = \int \frac{(ex^n + d)^2}{(cx^{2n} + a)^3} dx$$

input `integrate((d+e*x^n)^2/(a+c*x^(2*n))^3,x, algorithm="giac")`

output `integrate((e*x^n + d)^2/(c*x^(2*n) + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(d + ex^n)^2}{(a + cx^{2n})^3} dx = \int \frac{(d + e x^n)^2}{(a + c x^{2n})^3} dx$$

input `int((d + e*x^n)^2/(a + c*x^(2*n))^3,x)`

output `int((d + e*x^n)^2/(a + c*x^(2*n))^3, x)`

Reduce [F]

$$\int \frac{(d + ex^n)^2}{(a + cx^{2n})^3} dx \\ = \frac{x^{4n} \left(\int \frac{x^{2n}}{x^{6n}c^3 + 3x^{4n}a c^2 + 3x^{2n}a^2c + a^3} dx \right) a c^2 e^2 + 4x^{4n} \left(\int \frac{x^{2n}}{x^{6n}c^3 + 3x^{4n}a c^2 + 3x^{2n}a^2c + a^3} dx \right) c^3 d^2 n - x^{4n} \left(\int \frac{x^{2n}}{x^{6n}c^3 + 3x^{4n}a c^2 + 3x^{2n}a^2c + a^3} dx \right) c^3 d^2 n}{(a + cx^{2n})^3}$$

input `int((d+e*x^n)^2/(a+c*x^(2*n))^3,x)`

output `(x**4*n)*int(x**2*n)/(x**6*n)*c**3 + 3*x**4*n*a*c**2 + 3*x**2*n*a**2*c + a**3, x)*a*c**2*e**2 + 4*x**4*n*int(x**2*n)/(x**6*n)*c**3 + 3*x**4*n*a*c**2 + 3*x**2*n*a**2*c + a**3, x)*c**3*d**2*n - x**4*n*int(x**2*n)/(x**6*n)*c**3 + 3*x**4*n*a*c**2 + 3*x**2*n*a**2*c + a**3, x)*c**3*d**2 + 2*x**4*n*int(x**n)/(x**6*n)*c**3 + 3*x**4*n*a*c**2 + 3*x**2*n*a**2*c + a**3, x)*a*c**2*d*e + 2*x**2*n*int(x**2*n)/(x**6*n)*c**3 + 3*x**4*n*a*c**2 + 3*x**2*n*a**2*c + a**3, x)*a*c**2*d**2 + 2*x**2*n*int(x**2*n)/(x**6*n)*c**3 + 3*x**4*n*a*c**2 + 3*x**2*n*a**2*c + a**3, x)*a*c**2*d**2*n - 2*x**2*n*int(x**2*n)/(x**6*n)*c**3 + 3*x**4*n*a*c**2 + 3*x**2*n*a**2*c + a**3, x)*a*c**2*d**2 + 4*x**2*n*int(x**n)/(x**6*n)*c**3 + 3*x**4*n*a*c**2 + 3*x**2*n*a**2*c + a**3, x)*a*c**2*d**2*c + 4*x**2*n*int(x**n)/(x**6*n)*c**3 + 3*x**4*n*a*c**2 + 3*x**2*n*a**2*c + a**3, x)*a*c**2*d**2*c*d + e + int(x**2*n)/(x**6*n)*c**3 + 3*x**4*n*a*c**2 + 3*x**2*n*a**2*c + a**3, x)*a*c**3*e**2 + 4*int(x**2*n)/(x**6*n)*c**3 + 3*x**4*n*a*c**2 + 3*x**2*n*a**2*c + a**3, x)*a*c**2*d**2*n - int(x**2*n)/(x**6*n)*c**3 + 3*x**4*n*a*c**2 + 3*x**2*n*a**2*c + a**3, x)*a*c**2*d**2 + 2*int(x**n)/(x**6*n)*c**3 + 3*x**4*n*a*c**2 + 3*x**2*n*a**2*c + a**3, x)*a*c**3*d*e + d**2*x)/(a*(x**4*n)*c**2 + 2*x**2*n*a*c + a**2))`

3.39 $\int \frac{d+ex^n}{(a+cx^{2n})^3} dx$

Optimal result	293
Mathematica [A] (verified)	294
Rubi [A] (verified)	294
Maple [F]	296
Fricas [F]	297
Sympy [F(-1)]	297
Maxima [F]	297
Giac [F]	298
Mupad [F(-1)]	298
Reduce [F]	298

Optimal result

Integrand size = 19, antiderivative size = 184

$$\begin{aligned} & \int \frac{d + ex^n}{(a + cx^{2n})^3} dx \\ &= \frac{x(d + ex^n)}{4an(a + cx^{2n})^2} - \frac{x(d(1 - 4n) + e(1 - 3n)x^n)}{8a^2n^2(a + cx^{2n})} \\ &+ \frac{d(1 - 6n + 8n^2)x \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{8a^3n^2} \\ &+ \frac{e(1 - 3n)(1 - n)x^{1+n} \operatorname{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{8a^3n^2(1 + n)} \end{aligned}$$

output

```
1/4*x*(d+e*x^n)/a/n/(a+c*x^(2*n))^2-1/8*x*(d*(1-4*n)+e*(1-3*n)*x^n)/a^2/n^2/(a+c*x^(2*n))+1/8*d*(8*n^2-6*n+1)*x*hypergeom([1, 1/2/n],[1+1/2/n],-c*x^(2*n)/a)/a^3/n^2+1/8*e*(1-3*n)*(1-n)*x^(1+n)*hypergeom([1, 1/2*(1+n)/n],[3/2+1/2/n],-c*x^(2*n)/a)/a^3/n^2/(1+n)
```

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.45

$$\int \frac{d + ex^n}{(a + cx^{2n})^3} dx = \frac{dx \text{Hypergeometric2F1}\left(3, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a^3} + \frac{ex^{1+n} \text{Hypergeometric2F1}\left(3, \frac{1+n}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a^3(1+n)}$$

input `Integrate[(d + e*x^n)/(a + c*x^(2*n))^3, x]`

output `(d*x*Hypergeometric2F1[3, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)])/a^3 + (e*x^(1 + n)*Hypergeometric2F1[3, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)])/(a^3*(1 + n))`

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.05, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {1761, 1761, 1748, 778, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{d + ex^n}{(a + cx^{2n})^3} dx \\ & \downarrow \textcolor{blue}{1761} \\ & \frac{x(d + ex^n)}{4an(a + cx^{2n})^2} - \frac{\int \frac{e(1-3n)x^n + d(1-4n)}{(cx^{2n}+a)^2} dx}{4an} \\ & \downarrow \textcolor{blue}{1761} \\ & \frac{x(d + ex^n)}{4an(a + cx^{2n})^2} - \frac{\frac{x(d(1-4n)+e(1-3n)x^n)}{2an(a+cx^{2n})} - \frac{\int \frac{e(1-3n)(1-n)x^n + d(8n^2-6n+1)}{cx^{2n}+a} dx}{2an}}{4an} \end{aligned}$$

$$\begin{aligned}
 & \downarrow 1748 \\
 \frac{x(d+ex^n)}{4an(a+cx^{2n})^2} - \frac{\frac{x(d(1-4n)+e(1-3n)x^n)}{2an(a+cx^{2n})} - \frac{d(8n^2-6n+1) \int \frac{1}{cx^{2n}+a} dx + e(1-3n)(1-n) \int \frac{x^n}{cx^{2n}+a} dx}{2an}}{4an} \\
 & \downarrow 778 \\
 \frac{x(d+ex^n)}{4an(a+cx^{2n})^2} - \\
 \frac{x(d(1-4n)+e(1-3n)x^n)}{2an(a+cx^{2n})} - \frac{e(1-3n)(1-n) \int \frac{x^n}{cx^{2n}+a} dx + \frac{d(8n^2-6n+1)x \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a}}{2an} \\
 & \downarrow 888 \\
 \frac{x(d+ex^n)}{4an(a+cx^{2n})^2} - \\
 \frac{x(d(1-4n)+e(1-3n)x^n)}{2an(a+cx^{2n})} - \frac{\frac{d(8n^2-6n+1)x \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a} + \frac{e(1-3n)(1-n)x^{n+1} \text{Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3+\frac{1}{n}), -\frac{cx^{2n}}{a(n+1)}\right)}{2an}}{4an}
 \end{aligned}$$

input $\text{Int}[(d + e*x^n)/(a + c*x^(2*n))^3, x]$

output
$$\begin{aligned}
 & (x*(d + e*x^n))/(4*a*n*(a + c*x^(2*n))^2) - ((x*(d*(1 - 4*n) + e*(1 - 3*n) *x^n))/(2*a*n*(a + c*x^(2*n))) - ((d*(1 - 6*n + 8*n^2)*x*Hypergeometric2F1[1, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)])/a + (e*(1 - 3*n)*(1 - n)*x^(1 + n)*Hypergeometric2F1[1, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)])/(a*(1 + n)))/(2*a*n))/(4*a*n)
 \end{aligned}$$

Definitions of rubi rules used

rule 778 $\text{Int}[(a_.) + (b_.)*(x_.)^{(n_.)}^{(p_.)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[a^p * x^p * \text{Hypergeometric2F1}[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; \text{FreeQ}[\{a, b, n, p\}, x] \&& \text{!IGtQ}[p, 0] \&& \text{!IntegerQ}[1/n] \&& \text{!ILtQ}[\text{Simplify}[1/n + p], 0] \&& (\text{IntegerQ}[p] \mid \text{GtQ}[a, 0])$

rule 888 $\text{Int}[((c_.)*(x_.))^{(m_.)}*((a_.) + (b_.)*(x_.)^{(n_.)}^{(p_.)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[a^p * ((c*x)^(m+1)/(c*(m+1))) * \text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n + 1, (-b)*(x^n/a)], x] /; \text{FreeQ}[\{a, b, c, m, n, p\}, x] \&& \text{!IGtQ}[p, 0] \&& (\text{ILtQ}[p, 0] \mid \text{GtQ}[a, 0])$

rule 1748 $\text{Int}[((d_.) + (e_.)*(x_.)^{(n_.)})/((a_.) + (c_.)*(x_.)^{(n2_.)}), x_{\text{Symbol}}] \rightarrow \text{Simp}[d \text{Int}[1/(a + c*x^(2*n)), x], x] + \text{Simp}[e \text{Int}[x^n/(a + c*x^(2*n)), x], x] /; \text{FreeQ}[\{a, c, d, e, n\}, x] \&& \text{EqQ}[n2, 2*n] \&& \text{NeQ}[c*d^2 + a*e^2, 0] \&& (\text{PosQ}[a*c] \mid \text{!IntegerQ}[n])$

rule 1761 $\text{Int}[((d_.) + (e_.)*(x_.)^{(n_.)})*((a_.) + (c_.)*(x_.)^{(n2_.)}^{(p_.)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[-(d + e*x^n)*((a + c*x^(2*n))^(p+1)/(2*a*n*(p+1))), x] + \text{Simp}[1/(2*a*n*(p+1)) \text{Int}[(d*(2*n*p + 2*n + 1) + e*(2*n*p + 3*n + 1)*x^n)*(a + c*x^(2*n))^(p+1), x], x] /; \text{FreeQ}[\{a, c, d, e, n\}, x] \&& \text{EqQ}[n2, 2*n] \&& \text{ILtQ}[p, -1]$

Maple [F]

$$\int \frac{d + e x^n}{(a + c x^{2n})^3} dx$$

input $\text{int}((d+e*x^n)/(a+c*x^(2*n))^3, x)$

output $\text{int}((d+e*x^n)/(a+c*x^(2*n))^3, x)$

Fricas [F]

$$\int \frac{d + ex^n}{(a + cx^{2n})^3} dx = \int \frac{ex^n + d}{(cx^{2n} + a)^3} dx$$

input `integrate((d+e*x^n)/(a+c*x^(2*n))^3,x, algorithm="fricas")`

output `integral((e*x^n + d)/(c^3*x^(6*n) + 3*a*c^2*x^(4*n) + 3*a^2*c*x^(2*n) + a^3), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{d + ex^n}{(a + cx^{2n})^3} dx = \text{Timed out}$$

input `integrate((d+e*x**n)/(a+c*x**2*n)**3,x)`

output `Timed out`

Maxima [F]

$$\int \frac{d + ex^n}{(a + cx^{2n})^3} dx = \int \frac{ex^n + d}{(cx^{2n} + a)^3} dx$$

input `integrate((d+e*x^n)/(a+c*x^(2*n))^3,x, algorithm="maxima")`

output `1/8*(c*e*(3*n - 1)*x*x^(3*n) + c*d*(4*n - 1)*x*x^(2*n) + a*e*(5*n - 1)*x*x^n + a*d*(6*n - 1)*x)/(a^2*c^2*n^2*x^(4*n) + 2*a^3*c*n^2*x^(2*n) + a^4*n^2) + integrate(1/8*((3*n^2 - 4*n + 1)*e*x^n + (8*n^2 - 6*n + 1)*d)/(a^2*c*n^2*x^(2*n) + a^3*n^2), x)`

Giac [F]

$$\int \frac{d + ex^n}{(a + cx^{2n})^3} dx = \int \frac{ex^n + d}{(cx^{2n} + a)^3} dx$$

input `integrate((d+e*x^n)/(a+c*x^(2*n))^3,x, algorithm="giac")`

output `integrate((e*x^n + d)/(c*x^(2*n) + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{d + ex^n}{(a + cx^{2n})^3} dx = \int \frac{d + e x^n}{(a + c x^{2n})^3} dx$$

input `int((d + e*x^n)/(a + c*x^(2*n))^3,x)`

output `int((d + e*x^n)/(a + c*x^(2*n))^3, x)`

Reduce [F]

$$\begin{aligned} \int \frac{d + ex^n}{(a + cx^{2n})^3} dx &= \left(\int \frac{x^n}{x^{6n}c^3 + 3x^{4n}a c^2 + 3x^{2n}a^2c + a^3} dx \right) e \\ &\quad + \left(\int \frac{1}{x^{6n}c^3 + 3x^{4n}a c^2 + 3x^{2n}a^2c + a^3} dx \right) d \end{aligned}$$

input `int((d+e*x^n)/(a+c*x^(2*n))^3,x)`

output `int(x**n/(x**(6*n)*c**3 + 3*x**(4*n)*a*c**2 + 3*x**(2*n)*a**2*c + a**3),x)*e + int(1/(x**(6*n)*c**3 + 3*x**(4*n)*a*c**2 + 3*x**(2*n)*a**2*c + a**3),x)*d`

3.40 $\int \frac{1}{(d+ex^n)(a+cx^{2n})^3} dx$

Optimal result	299
Mathematica [A] (verified)	300
Rubi [A] (verified)	301
Maple [F]	302
Fricas [F]	303
Sympy [F(-1)]	303
Maxima [F]	303
Giac [F]	304
Mupad [F(-1)]	304
Reduce [F]	305

Optimal result

Integrand size = 21, antiderivative size = 462

$$\begin{aligned} \int \frac{1}{(d+ex^n)(a+cx^{2n})^3} dx &= \frac{cx(d-ex^n)}{4a(cd^2+ae^2)n(a+cx^{2n})^2} \\ &- \frac{cx(d(ae^2(1-8n)+cd^2(1-4n))-e(ae^2(1-7n)+cd^2(1-3n))x^n)}{8a^2(cd^2+ae^2)^2n^2(a+cx^{2n})} \\ &+ \frac{cde^4x \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a(cd^2+ae^2)^3} \\ &+ \frac{cd(ae^2(1-8n)+cd^2(1-4n))(1-2n)x \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{8a^3(cd^2+ae^2)^2n^2} \\ &+ \frac{e^6x \text{Hypergeometric2F1}\left(1, \frac{1}{n}, 1+\frac{1}{n}, -\frac{ex^n}{d}\right)}{d(cd^2+ae^2)^3} \\ &- \frac{ce^5x^{1+n} \text{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a(cd^2+ae^2)^3(1+n)} \\ &- \frac{ce(ae^2(1-7n)+cd^2(1-3n))(1-n)x^{1+n} \text{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{8a^3(cd^2+ae^2)^2n^2(1+n)} \end{aligned}$$

output

$$\begin{aligned} & \frac{1}{4} c x (d - e x^n) / a / (a e^2 + c d^2) / n / (a + c x^{(2n)})^2 - \frac{1}{8} c x (d (a e^{2(1-8n)} + c d^{2(1-4n)}) - e (a e^{2(1-7n)} + c d^{2(1-3n)}) x^n) / a^2 / (a e^2 + c d^2)^2 \\ & - \frac{2}{n^2} / (a + c x^{(2n)}) + c d e^4 x \operatorname{hypergeom}([1, 1/2/n], [1+1/2/n], -c x^{(2n)}/a) \\ & / a / (a e^2 + c d^2)^3 + \frac{1}{8} c d (a e^{2(1-8n)} + c d^{2(1-4n)}) (1-2n) x \operatorname{hypergeom}([1, 1/2/n], [1+1/2/n], -c x^{(2n)}/a) / a^3 / (a e^2 + c d^2)^2 \\ & - \frac{2}{n^2} e^6 / (a + c x^{(2n)}) + c d e^4 x \operatorname{hypergeom}([1, 1/n], [1+1/n], -e x^n/d) / d / (a e^2 + c d^2)^3 - c e^{5n} x^{(1+n)} \operatorname{hypergeom}([1, 1/2*(1+n)/n], [3/2+1/2/n], -c x^{(2n)}/a) / a / (a e^2 + c d^2)^3 / (1+n) - \frac{1}{8} c e^4 (a e^{2(1-7n)} + c d^{2(1-3n)}) (1-n) x^{(1+n)} \operatorname{hypergeom}([1, 1/2*(1+n)/n], [3/2+1/2/n], -c x^{(2n)}/a) / a^3 / (a e^2 + c d^2)^2 / (1+n) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.57 (sec), antiderivative size = 346, normalized size of antiderivative = 0.75

$$\begin{aligned} & \int \frac{1}{(d + e x^n) (a + c x^{2n})^3} dx \\ & = x \left(\frac{c d e^4 \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{c x^{2n}}{a}\right)}{a} + \frac{e^6 \operatorname{Hypergeometric2F1}\left(1, \frac{1}{n}, 1+\frac{1}{n}, -\frac{e x^n}{d}\right)}{d} - \frac{c e^5 x^n \operatorname{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}, -\frac{c x^{2n}}{a}\right)}{a(1+n)} \right) \end{aligned}$$

input

Integrate[1/((d + e*x^n)*(a + c*x^(2n))^3), x]

output

$$\begin{aligned} & (x ((c d e^4 \operatorname{Hypergeometric2F1}[1, 1/(2n), (2 + n^{(-1)})/2, -((c x^{(2n)})/a)])) / a + (e^6 \operatorname{Hypergeometric2F1}[1, n^{(-1)}, 1 + n^{(-1)}, -((e x^n)/d)]) / d - (c e^{5n} x \operatorname{Hypergeometric2F1}[1, (1 + n)/(2n), (3 + n^{(-1)})/2, -((c x^{(2n)})/a)]) / (a (1 + n)) + (c d e^2 (c d^2 + a e^2) \operatorname{Hypergeometric2F1}[2, 1/(2n), (2 + n^{(-1)})/2, -((c x^{(2n)})/a)]) / a^2 - (c e^3 (c d^2 + a e^2) x^n \operatorname{Hypergeometric2F1}[2, (1 + n)/(2n), (3 + n^{(-1)})/2, -((c x^{(2n)})/a)]) / (a^2 (1 + n)) + (c d (c d^2 + a e^2)^2 \operatorname{Hypergeometric2F1}[3, 1/(2n), (2 + n^{(-1)})/2, -((c x^{(2n)})/a)]) / a^3 - (c e^2 (c d^2 + a e^2)^2 x^n \operatorname{Hypergeometric2F1}[3, (1 + n)/(2n), (3 + n^{(-1)})/2, -((c x^{(2n)})/a)]) / (a^3 (1 + n))) / (c d^2 + a e^2)^3 \end{aligned}$$

Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 582, normalized size of antiderivative = 1.26, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {1767, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a + cx^{2n})^3 (d + ex^n)} dx \\
 & \quad \downarrow \textcolor{blue}{1767} \\
 & \int \left(-\frac{ce^2(ex^n - d)}{(ae^2 + cd^2)^2 (a + cx^{2n})^2} - \frac{c(ex^n - d)}{(ae^2 + cd^2)(a + cx^{2n})^3} + \frac{e^6}{(ae^2 + cd^2)^3 (d + ex^n)} - \frac{ce^4(ex^n - d)}{(ae^2 + cd^2)^3 (a + cx^{2n})} \right) \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & -\frac{ce(1 - 3n)(1 - n)x^{n+1} \text{Hypergeometric2F1} \left(1, \frac{n+1}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a} \right)}{8a^3n^2(n+1)(ae^2 + cd^2)} + \\
 & \quad \frac{cd(1 - 4n)(1 - 2n)x \text{Hypergeometric2F1} \left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a} \right)}{8a^3n^2(ae^2 + cd^2)} - \\
 & \quad \frac{cde^2(1 - 2n)x \text{Hypergeometric2F1} \left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a} \right)}{2a^2n(ae^2 + cd^2)^2} - \frac{cx(d(1 - 4n) - e(1 - 3n)x^n)}{8a^2n^2(ae^2 + cd^2)(a + cx^{2n})} + \\
 & \quad \frac{ce^3(1 - n)x^{n+1} \text{Hypergeometric2F1} \left(1, \frac{n+1}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a} \right)}{2a^2n(n+1)(ae^2 + cd^2)^2} + \frac{ce^2x(d - ex^n)}{2an(ae^2 + cd^2)^2(a + cx^{2n})} + \\
 & \quad \frac{cx(d - ex^n)}{4an(ae^2 + cd^2)(a + cx^{2n})^2} + \frac{e^6x \text{Hypergeometric2F1} \left(1, \frac{1}{n}, 1 + \frac{1}{n}, -\frac{ex^n}{d} \right)}{d(ae^2 + cd^2)^3} - \\
 & \quad \frac{ce^5x^{n+1} \text{Hypergeometric2F1} \left(1, \frac{n+1}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a} \right)}{a(n+1)(ae^2 + cd^2)^3} + \\
 & \quad \frac{cde^4x \text{Hypergeometric2F1} \left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a} \right)}{a(ae^2 + cd^2)^3}
 \end{aligned}$$

input `Int[1/((d + e*x^n)*(a + c*x^(2*n))^3),x]`

output

$$\begin{aligned} & \frac{(c*x*(d - e*x^n))/(4*a*(c*d^2 + a*e^2)*n*(a + c*x^(2*n))^2) + (c*e^2*x*(d - e*x^n))/(2*a*(c*d^2 + a*e^2)^2*n*(a + c*x^(2*n))) - (c*x*(d*(1 - 4*n) - e*(1 - 3*n)*x^n))/(8*a^2*(c*d^2 + a*e^2)*n^2*(a + c*x^(2*n))) + (c*d*e^4*x*Hypergeometric2F1[1, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)])/(a*(c*d^2 + a*e^2)^3) + (c*d*(1 - 4*n)*(1 - 2*n)*x*Hypergeometric2F1[1, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)])/(8*a^3*(c*d^2 + a*e^2)*n^2) - (c*d*e^2*(1 - 2*n)*x*Hypergeometric2F1[1, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)])/(2*a^2*(c*d^2 + a*e^2)^2*n) + (e^6*x*Hypergeometric2F1[1, n^(-1), 1 + n^(-1), -((e*x^n)/d)])/(d*(c*d^2 + a*e^2)^3) - (c*e^5*x^(1 + n)*Hypergeometric2F1[1, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)])/(a*(c*d^2 + a*e^2)^3*(1 + n)) - (c*e*(1 - 3*n)*(1 - n)*x^(1 + n)*Hypergeometric2F1[1, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)])/(8*a^3*(c*d^2 + a*e^2)*n^2*(1 + n)) + (c*e^3*(1 - n)*x^(1 + n)*Hypergeometric2F1[1, (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)])/(2*a^2*(c*d^2 + a*e^2)^2*n*(1 + n)) \end{aligned}$$

Defintions of rubi rules used

rule 1767

```
Int[((d_) + (e_)*(x_)^(n_))^(q_)*((a_) + (c_)*(x_)^(n2_))^(p_), x_Symbol]
  :> Int[ExpandIntegrand[(d + e*x^n)^q*(a + c*x^(2*n))^p, x], x] /; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && NeQ[c*d^2 + a*e^2, 0] && ((IntegersQ[p, q] && !IntegerQ[n]) || IGtQ[p, 0] || (IGtQ[q, 0] && !IntegerQ[n]))
```

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [F]

$$\int \frac{1}{(d + e x^n) (a + c x^{2n})^3} dx$$

input

```
int(1/(d+e*x^n)/(a+c*x^(2*n))^3,x)
```

output

```
int(1/(d+e*x^n)/(a+c*x^(2*n))^3,x)
```

Fricas [F]

$$\int \frac{1}{(d + ex^n)(a + cx^{2n})^3} dx = \int \frac{1}{(cx^{2n} + a)^3(ex^n + d)} dx$$

input `integrate(1/(d+e*x^n)/(a+c*x^(2*n))^3,x, algorithm="fricas")`

output `integral(1/(a^3*e*x^n + a^3*d + (c^3*e*x^n + c^3*d)*x^(6*n) + 3*(a*c^2*e*x^n + a*c^2*d)*x^(4*n) + 3*(a^2*c*e*x^n + a^2*c*d)*x^(2*n)), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex^n)(a + cx^{2n})^3} dx = \text{Timed out}$$

input `integrate(1/(d+e*x**n)/(a+c*x**2*n)**3,x)`

output `Timed out`

Maxima [F]

$$\int \frac{1}{(d + ex^n)(a + cx^{2n})^3} dx = \int \frac{1}{(cx^{2n} + a)^3(ex^n + d)} dx$$

input `integrate(1/(d+e*x^n)/(a+c*x^(2*n))^3,x, algorithm="maxima")`

output

```
e^6*integrate(1/(c^3*d^7 + 3*a*c^2*d^5*e^2 + 3*a^2*c*d^3*e^4 + a^3*d*e^6 +
(c^3*d^6*e + 3*a*c^2*d^4*e^3 + 3*a^2*c*d^2*e^5 + a^3*e^7)*x^n), x) - 1/8*
((a*c^2*e^3*(7*n - 1) + c^3*d^2*e*(3*n - 1))*x*x^(3*n) - (a*c^2*d*e^2*(8*n -
1) + c^3*d^3*(4*n - 1))*x*x^(2*n) + (a^2*c*e^3*(9*n - 1) + a*c^2*d^2*e^
(5*n - 1))*x*x^n - (a^2*c*d*e^2*(10*n - 1) + a*c^2*d^3*(6*n - 1))*x)/(a^4*c^
2*d^4*n^2 + 2*a^5*c*d^2*e^2*n^2 + a^6*e^4*n^2 + (a^2*c^4*d^4*n^2 + 2*a^3*c^
3*d^2*e^2*n^2 + a^4*c^2*e^4*n^2)*x^(4*n) + 2*(a^3*c^3*d^4*n^2 + 2*a^4*c^
2*d^2*e^2*n^2 + a^5*c*e^4*n^2)*x^(2*n)) - integrate(-1/8*((8*n^2 - 6*n +
1)*c^3*d^5 + 2*(12*n^2 - 8*n + 1)*a*c^2*d^3*e^2 + (24*n^2 - 10*n + 1)*a^2*c^
d*e^4 - ((3*n^2 - 4*n + 1)*c^3*d^4*e + 2*(5*n^2 - 6*n + 1)*a*c^2*d^2*e^3 +
(15*n^2 - 8*n + 1)*a^2*c*e^5)*x^n)/(a^3*c^3*d^6*n^2 + 3*a^4*c^2*d^4*e^2*n^
2 + 3*a^5*c*d^2*e^4*n^2 + a^6*e^6*n^2 + (a^2*c^4*d^6*n^2 + 3*a^3*c^3*d^4*e^
2*n^2 + 3*a^4*c^2*d^2*e^4*n^2 + a^5*c*e^6*n^2)*x^(2*n)), x)
```

Giac [F]

$$\int \frac{1}{(d + ex^n)(a + cx^{2n})^3} dx = \int \frac{1}{(cx^{2n} + a)^3(ex^n + d)} dx$$

input

```
integrate(1/(d+e*x^n)/(a+c*x^(2*n))^3,x, algorithm="giac")
```

output

```
integrate(1/((c*x^(2*n) + a)^3*(e*x^n + d)), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex^n)(a + cx^{2n})^3} dx = \int \frac{1}{(a + cx^{2n})^3(d + ex^n)} dx$$

input

```
int(1/((a + c*x^(2*n))^3*(d + e*x^n)),x)
```

output

```
int(1/((a + c*x^(2*n))^3*(d + e*x^n)), x)
```

Reduce [F]

$$\begin{aligned} & \int \frac{1}{(d + ex^n)(a + cx^{2n})^3} dx \\ &= \int \frac{1}{x^{7n}c^3e + x^{6n}c^3d + 3x^{5n}ac^2e + 3x^{4n}ac^2d + 3x^{3n}a^2ce + 3x^{2n}a^2cd + x^na^3e + a^3d} dx \end{aligned}$$

input `int(1/(d+e*x^n)/(a+c*x^(2*n))^3,x)`

output `int(1/(x**(7*n)*c**3*e + x**(6*n)*c**3*d + 3*x**(5*n)*a*c**2*e + 3*x**(4*n)*a*c**2*d + 3*x**(3*n)*a**2*c*e + 3*x**(2*n)*a**2*c*d + x**n*a**3*e + a**3*d),x)`

3.41 $\int \frac{1}{(d+ex^n)^2(a+cx^{2n})^3} dx$

Optimal result	306
Mathematica [A] (verified)	307
Rubi [A] (verified)	308
Maple [F]	310
Fricas [F]	310
Sympy [F(-1)]	310
Maxima [F]	311
Giac [F]	311
Mupad [F(-1)]	312
Reduce [F]	312

Optimal result

Integrand size = 21, antiderivative size = 660

$$\begin{aligned}
 \int \frac{1}{(d+ex^n)^2(a+cx^{2n})^3} dx &= \frac{cx(d-ex^n)}{4a(cd^2+ae^2)n(d+ex^n)(a+cx^{2n})^2} \\
 &+ \frac{cx(a^2e^4(1-8n)-c^2d^4(1-4n)+8acd^2e^2n+2cde(ae^2(1-9n)+cd^2(1-3n))x^n)}{8a^2(cd^2+ae^2)^3n^2(a+cx^{2n})} \\
 &- \frac{c(1-2n)(a^2e^4(1-8n)-c^2d^4(1-4n)+8acd^2e^2n)x\text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{8a^3(cd^2+ae^2)^3n^2} \\
 &- \frac{ce^2(acd^2e^2(1-12n)+c^2d^4(1-2n)+2a^2e^4n)x\text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{2a^2(cd^2+ae^2)^4n} \\
 &- \frac{ce^4(ae^2(1-13n)+cd^2(1-n))x\text{Hypergeometric2F1}\left(1, \frac{1}{n}, 1+\frac{1}{n}, -\frac{ex^n}{d}\right)}{2a(cd^2+ae^2)^4n} \\
 &- \frac{c^2de(ae^2(1-9n)+cd^2(1-3n))(1-n)x^{1+n}\text{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{4a^3(cd^2+ae^2)^3n^2(1+n)} \\
 &+ \frac{c^2de^3(ae^2(1-13n)+cd^2(1-n))x^{1+n}\text{Hypergeometric2F1}\left(1, \frac{1+n}{2n}, \frac{1}{2}(3+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{2a^2(cd^2+ae^2)^4n(1+n)} \\
 &- \frac{e^4(cd^2-2ae^2)x\text{Hypergeometric2F1}\left(2, \frac{1}{n}, 1+\frac{1}{n}, -\frac{ex^n}{d}\right)}{2ad^2(cd^2+ae^2)^3}
 \end{aligned}$$

output

$$\begin{aligned}
 & 1/4*c*x*(d-e*x^n)/a/(a*e^2+c*d^2)/n/(d+e*x^n)/(a+c*x^(2*n))^2 + 1/8*c*x*(a^2 \\
 & *e^4*(1-8*n)-c^2*d^4*(1-4*n)+8*a*c*d^2*e^2*n+2*c*d*e*(a*e^2*(1-9*n)+c*d^2* \\
 & (1-3*n))*x^n/a^2/(a*e^2+c*d^2)^3/n^2/(a+c*x^(2*n))-1/8*c*(1-2*n)*(a^2*e^4 \\
 & *(1-8*n)-c^2*d^4*(1-4*n)+8*a*c*d^2*e^2*n)*x*hypergeom([1, 1/2/n], [1+1/2/n] \\
 & , -c*x^(2*n)/a)/a^3/(a*e^2+c*d^2)^3/n^2-1/2*c*e^2*(a*c*d^2*e^2*(1-12*n)+c^2 \\
 & *d^4*(1-2*n)+2*a^2*e^4*n)*x*hypergeom([1, 1/2/n], [1+1/2/n], -c*x^(2*n)/a)/a \\
 & ^2/(a*e^2+c*d^2)^4/n-1/2*c*e^4*(a*e^2*(1-13*n)+c*d^2*(1-n))*x*hypergeom([1 \\
 & , 1/n], [1+1/n], -e*x^n/d)/a/(a*e^2+c*d^2)^4/n-1/4*c^2*d*e*(a*e^2*(1-9*n)+c* \\
 & d^2*(1-3*n))*(1-n)*x^(1+n)*hypergeom([1, 1/2*(1+n)/n], [3/2+1/2/n], -c*x^(2*n) \\
 & /a)/a^3/(a*e^2+c*d^2)^3/n^2/(1+n)+1/2*c^2*d*e^3*(a*e^2*(1-13*n)+c*d^2*(1 \\
 & -n))*x^(1+n)*hypergeom([1, 1/2*(1+n)/n], [3/2+1/2/n], -c*x^(2*n)/a)/a^2/(a*e \\
 & ^2+c*d^2)^4/n/(1+n)-1/2*e^4*(-2*a*e^2+c*d^2)*x*hypergeom([2, 1/n], [1+1/n], \\
 & -e*x^n/d)/a/d^2/(a*e^2+c*d^2)^3
 \end{aligned}$$
Mathematica [A] (verified)

Time = 1.02 (sec), antiderivative size = 426, normalized size of antiderivative = 0.65

$$\begin{aligned}
 & \int \frac{1}{(d+ex^n)^2(a+cx^{2n})^3} dx \\
 & = \frac{x \left(\frac{ce^4(5cd^2-ae^2) \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a} + 6ce^6 \text{Hypergeometric2F1}\left(1, \frac{1}{n}, 1+\frac{1}{n}, -\frac{ex^n}{d}\right) - \frac{6c^2de^5x}{a^2} \right)}{(d+ex^n)^2(a+cx^{2n})^3}
 \end{aligned}$$

input

```
Integrate[1/((d + e*x^n)^2*(a + c*x^(2*n))^3), x]
```

output

$$\begin{aligned}
 & (x*((c*e^4*(5*c*d^2 - a*e^2)*\text{Hypergeometric2F1}[1, 1/(2*n), (2 + n^(-1))/2, \\
 & -((c*x^(2*n))/a)])/a + 6*c*e^6*\text{Hypergeometric2F1}[1, n^(-1), 1 + n^(-1), \\
 & -(e*x^n)/d]) - (6*c^2*d*e^5*x^n*\text{Hypergeometric2F1}[1, (1 + n)/(2*n), (3 + n \\
 & ^(-1))/2, -((c*x^(2*n))/a)])/(a*(1 + n)) + (c*e^2*(3*c*d^2 - a*e^2)*(c*d^2 \\
 & + a*e^2)*\text{Hypergeometric2F1}[2, 1/(2*n), (2 + n^(-1))/2, -((c*x^(2*n))/a)]) \\
 & /a^2 + (e^6*(c*d^2 + a*e^2)*\text{Hypergeometric2F1}[2, n^(-1), 1 + n^(-1), -((e* \\
 & x^n)/d)])/d^2 - (4*c^2*d*e^3*(c*d^2 + a*e^2)*x^n*\text{Hypergeometric2F1}[2, (1 + \\
 & n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)])/(a^2*(1 + n)) + (c*(c*d^2 - \\
 & a*e^2)*(c*d^2 + a*e^2)^2*\text{Hypergeometric2F1}[3, 1/(2*n), (2 + n^(-1))/2, -((\\
 & c*x^(2*n))/a)])/a^3 - (2*c^2*d*e*(c*d^2 + a*e^2)^2*x^n*\text{Hypergeometric2F1}[3 \\
 & , (1 + n)/(2*n), (3 + n^(-1))/2, -((c*x^(2*n))/a)])/(a^3*(1 + n)))/(c*d^2 \\
 & + a*e^2)^4
 \end{aligned}$$

Rubi [A] (verified)

Time = 1.00 (sec) , antiderivative size = 701, normalized size of antiderivative = 1.06, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {1767, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a + cx^{2n})^3 (d + ex^n)^2} dx \\
 & \quad \downarrow \textcolor{blue}{1767} \\
 & \int \left(-\frac{ce^2(ae^2 - 3cd^2 + 4cdex^n)}{(ae^2 + cd^2)^3 (a + cx^{2n})^2} - \frac{c(ae^2 - cd^2 + 2cdex^n)}{(ae^2 + cd^2)^2 (a + cx^{2n})^3} + \frac{6cde^6}{(ae^2 + cd^2)^4 (d + ex^n)} + \frac{e^6}{(ae^2 + cd^2)^3 (d + ex^n)^2} \right. \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & - \frac{c^2 de(1 - 3n)(1 - n)x^{n+1} \text{Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{4a^3 n^2 (n + 1) (ae^2 + cd^2)^2} + \\
 & \frac{c(1 - 4n)(1 - 2n)x(cd^2 - ae^2) \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{8a^3 n^2 (ae^2 + cd^2)^2} + \\
 & \frac{2c^2 de^3(1 - n)x^{n+1} \text{Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a^2 n(n + 1) (ae^2 + cd^2)^3} - \\
 & \frac{ce^2(1 - 2n)x(3cd^2 - ae^2) \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{2a^2 n (ae^2 + cd^2)^3} - \\
 & \frac{cx((1 - 4n)(cd^2 - ae^2) - 2cde(1 - 3n)x^n)}{8a^2 n^2 (ae^2 + cd^2)^2 (a + cx^{2n})} - \\
 & \frac{6c^2 de^5 x^{n+1} \text{Hypergeometric2F1}\left(1, \frac{n+1}{2n}, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a(n + 1) (ae^2 + cd^2)^4} + \frac{ce^2 x(-ae^2 + 3cd^2 - 4cdex^n)}{2an (ae^2 + cd^2)^3 (a + cx^{2n})} + \\
 & \frac{cx(-ae^2 + cd^2 - 2cdex^n)}{4an (ae^2 + cd^2)^2 (a + cx^{2n})^2} + \frac{6ce^6 x \text{Hypergeometric2F1}\left(1, \frac{1}{n}, 1 + \frac{1}{n}, -\frac{ex^n}{d}\right)}{(ae^2 + cd^2)^4} + \\
 & \frac{e^6 x \text{Hypergeometric2F1}\left(2, \frac{1}{n}, 1 + \frac{1}{n}, -\frac{ex^n}{d}\right)}{d^2 (ae^2 + cd^2)^3} + \\
 & \frac{ce^4 x(5cd^2 - ae^2) \text{Hypergeometric2F1}\left(1, \frac{1}{2n}, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{a (ae^2 + cd^2)^4}
 \end{aligned}$$

input $\text{Int}[1/((d + e*x^n)^2*(a + c*x^{(2*n)})^3), x]$

output
$$\begin{aligned} & \frac{(c*x*(c*d^2 - a*e^2 - 2*c*d*e*x^n))/(4*a*(c*d^2 + a*e^2)^2*n*(a + c*x^{(2*n)}))}{(c*e^2*x*(3*c*d^2 - a*e^2 - 4*c*d*e*x^n))/(2*a*(c*d^2 + a*e^2)^3*n*(a + c*x^{(2*n)}))} \\ & - \frac{(c*x*((c*d^2 - a*e^2)*(1 - 4*n) - 2*c*d*e*(1 - 3*n)*x^n)/(8*a^2*(c*d^2 + a*e^2)^2*n^2*(a + c*x^{(2*n)})) + (c*e^4*(5*c*d^2 - a*e^2)*x*\text{Hypergeometric2F1}[1, 1/(2*n), (2 + n^{(-1)})/2, -((c*x^{(2*n)})/a)]/(a*(c*d^2 + a*e^2)^4) + (c*(c*d^2 - a*e^2)*(1 - 4*n)*(1 - 2*n)*x*\text{Hypergeometric2F1}[1, 1/(2*n), (2 + n^{(-1)})/2, -((c*x^{(2*n)})/a)]/(8*a^3*(c*d^2 + a*e^2)^2*n^2) - (c*e^2*(3*c*d^2 - a*e^2)*(1 - 2*n)*x*\text{Hypergeometric2F1}[1, 1/(2*n), (2 + n^{(-1)})/2, -((c*x^{(2*n)})/a)]/(2*a^2*(c*d^2 + a*e^2)^3*n) + (6*c*e^6*x*\text{Hypergeometric2F1}[1, n^{(-1)}, 1 + n^{(-1)}, -((e*x^n)/d)])/(c*d^2 + a*e^2)^4 - (6*c^2*d*e^5*x^{(1 + n)}*\text{Hypergeometric2F1}[1, (1 + n)/(2*n), (3 + n^{(-1)})/2, -((c*x^{(2*n)})/a)]/(a*(c*d^2 + a*e^2)^4*(1 + n)) - (c^2*d*e*(1 - 3*n)*(1 - n)*x^{(1 + n)}*\text{Hypergeometric2F1}[1, (1 + n)/(2*n), (3 + n^{(-1)})/2, -((c*x^{(2*n)})/a)]/(4*a^3*(c*d^2 + a*e^2)^2*n^2*(1 + n)) + (2*c^2*d*e^3*(1 - n)*x^{(1 + n)}*\text{Hypergeometric2F1}[1, (1 + n)/(2*n), (3 + n^{(-1)})/2, -((c*x^{(2*n)})/a)]/(a^2*(c*d^2 + a*e^2)^3*n*(1 + n)) + (e^6*x*\text{Hypergeometric2F1}[2, n^{(-1)}, 1 + n^{(-1)}, -((e*x^n)/d)])/(d^2*(c*d^2 + a*e^2)^3)} \end{aligned}$$

Definitions of rubi rules used

rule 1767
$$\begin{aligned} & \text{Int}[(d_ + e_)*(x_)^{(n_)}(q_)*(a_ + c_)*(x_)^{(n2_)}(p_), x_Symbol] \\ & :> \text{Int}[\text{ExpandIntegrand}[(d + e*x^n)^q*(a + c*x^{(2*n)})^p, x], x] /; \text{FreeQ}[\{a, c, d, e, n, p, q\}, x] \&& \text{EqQ}[n2, 2*n] \&& \text{NeQ}[c*d^2 + a*e^2, 0] \&& (\text{IntegersQ}[p, q] \&& \text{!IntegerQ}[n]) \text{ || } \text{IGtQ}[p, 0] \text{ || } (\text{IGtQ}[q, 0] \&& \text{!IntegerQ}[n]) \end{aligned}$$

rule 2009
$$\text{Int}[u_, x_Symbol] :> \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$$

Maple [F]

$$\int \frac{1}{(d + e x^n)^2 (a + c x^{2n})^3} dx$$

input `int(1/(d+e*x^n)^2/(a+c*x^(2*n))^3,x)`

output `int(1/(d+e*x^n)^2/(a+c*x^(2*n))^3,x)`

Fricas [F]

$$\int \frac{1}{(d + e x^n)^2 (a + c x^{2n})^3} dx = \int \frac{1}{(c x^{2n} + a)^3 (e x^n + d)^2} dx$$

input `integrate(1/(d+e*x^n)^2/(a+c*x^(2*n))^3,x, algorithm="fricas")`

output `integral(1/(a^3*e^2*x^(2*n) + 2*a^3*d*e*x^n + a^3*d^2 + (c^3*e^2*x^(2*n) + 2*c^3*d*e*x^n + c^3*d^2)*x^(6*n) + 3*(a*c^2*e^2*x^(2*n) + 2*a*c^2*d*e*x^n + a*c^2*d^2)*x^(4*n) + 3*(a^2*c*e^2*x^(2*n) + 2*a^2*c*d*e*x^n + a^2*c*d^2)*x^(2*n)), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(d + e x^n)^2 (a + c x^{2n})^3} dx = \text{Timed out}$$

input `integrate(1/(d+e*x**n)**2/(a+c*x**(2*n))**3,x)`

output `Timed out`

Maxima [F]

$$\int \frac{1}{(d + ex^n)^2 (a + cx^{2n})^3} dx = \int \frac{1}{(cx^{2n} + a)^3 (ex^n + d)^2} dx$$

input `integrate(1/(d+e*x^n)^2/(a+c*x^(2*n))^3,x, algorithm="maxima")`

output

```
(c*d^2*e^6*(7*n - 1) + a*e^8*(n - 1))*integrate(1/(c^4*d^10*n + 4*a*c^3*d^8*e^2*n + 6*a^2*c^2*d^6*e^4*n + 4*a^3*c*d^4*e^6*n + a^4*d^2*e^8*n + (c^4*d^9*e*n + 4*a*c^3*d^7*e^3*n + 6*a^2*c^2*d^5*e^5*n + 4*a^3*c*d^3*e^7*n + a^4*d*e^9*n)*x^n), x) - 1/8*(2*(a*c^3*d^2*e^4*(11*n - 1) + c^4*d^4*e^2*(3*n - 1) - 4*a^2*c^2*e^6*n)*x*x^(4*n) + (a^2*c^2*d*e^5*(8*n - 1) + 2*a*c^3*d^3*e^3*(5*n - 1) + c^4*d^5*e^2*(2*n - 1))*x*x^(3*n) + (a^2*c^2*d^2*e^4*(34*n - 3) - c^4*d^6*(4*n - 1) - 2*a*c^3*d^4*e^2*(n + 1) - 16*a^3*c*e^6*n)*x*x^(2*n) + (a^3*c*d*e^5*(10*n - 1) + 2*a^2*c^2*d^3*e^3*(7*n - 1) + a*c^3*d^5*e^4*(4*n - 1))*x*x^n + (a^3*c*d^2*e^4*(10*n - 1) - a*c^3*d^6*(6*n - 1) - 12*a^2*c^2*d^4*e^2*n - 8*a^4*e^6*n)*x)/(a^4*c^3*d^8*n^2 + 3*a^5*c^2*d^6*e^2*n^2 + 3*a^6*c*d^4*e^4*n^2 + a^7*d^2*e^6*n^2 + (a^2*c^5*d^7*n^2 + 3*a^3*c^4*d^6*e^2*n^2 + 3*a^4*c^3*d^3*e^5*n^2 + a^5*c^2*d^4*e^7*n^2)*x^(5*n) + (a^2*c^5*d^8*n^2 + 3*a^3*c^4*d^6*e^2*n^2 + 3*a^4*c^3*d^4*e^4*n^2 + a^5*c^2*d^2*e^6*n^2)*x^(4*n) + 2*(a^3*c^4*d^7*e*n^2 + 3*a^4*c^3*d^5*e^3*n^2 + 3*a^5*c^2*d^3*e^5*n^2 + a^6*c*d^4*e^7*n^2)*x^(3*n) + 2*(a^3*c^4*d^8*n^2 + 3*a^4*c^3*d^6*e^2*n^2 + 3*a^5*c^2*d^4*e^4*n^2 + a^6*c^2*d^2*e^6*n^2 + 3*a^5*c^2*d^7*e*n^2 + 3*a^5*c^2*d^5*e^3*n^2 + 3*a^6*c*d^3*e^5*n^2 + a^7*d^2*e^7*n^2)*x^(2*n) + (a^4*c^3*d^7*e*n^2 + 3*a^5*c^2*d^5*e^3*n^2 + 3*a^6*c*d^3*e^5*n^2 + a^7*d^2*e^7*n^2)*x^(1*n)) - integrate(-1/8*((8*n^2 - 6*n + 1)*c^4*d^6 + (32*n^2 - 18*n + 1)*a*c^3*d^4*e^2 + (48*n^2 - 2*n - 1)*a^2*c^2*d^2*e^4 - (24*n^2 - 10*n + 1)*a^3*c*e^6 - 2*((3*n^2 - 4*n + 1)*c^4*d^5*e + 2*(7*n^2 - 8*n + 1)*a*c^3*d^3*e^...)
```

Giac [F]

$$\int \frac{1}{(d + ex^n)^2 (a + cx^{2n})^3} dx = \int \frac{1}{(cx^{2n} + a)^3 (ex^n + d)^2} dx$$

input `integrate(1/(d+e*x^n)^2/(a+c*x^(2*n))^3,x, algorithm="giac")`

output `integrate(1/((c*x^(2*n) + a)^3*(e*x^n + d)^2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex^n)^2 (a + cx^{2n})^3} dx = \int \frac{1}{(a + cx^{2n})^3 (d + ex^n)^2} dx$$

input `int(1/((a + c*x^(2*n))^3*(d + e*x^n)^2),x)`

output `int(1/((a + c*x^(2*n))^3*(d + e*x^n)^2), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{1}{(d + ex^n)^2 (a + cx^{2n})^3} dx \\ &= \int \frac{1}{x^{8n}c^3e^2 + 2x^{7n}c^3de + 3x^{6n}a c^2e^2 + x^{6n}c^3d^2 + 6x^{5n}a c^2de + 3x^{4n}a^2ce^2 + 3x^{4n}a c^2d^2 + 6x^{3n}a^2cde + x^{2n}a^3c^2e^2} dx \end{aligned}$$

input `int(1/(d+e*x^n)^2/(a+c*x^(2*n))^3,x)`

output `int(1/(x**8*n)*c**3*e**2 + 2*x**7*n)*c**3*d*e + 3*x**6*n)*a*c**2*e**2 + x**6*n)*c**3*d**2 + 6*x**5*n)*a*c**2*d*e + 3*x**4*n)*a**2*c*e**2 + 3*x**4*n)*a*c**2*d**2 + 6*x**3*n)*a**2*c*d*e + x**2*n)*a**3*c**2*d**2 + 2*x**n*a**3*d*e + a**3*d**2),x)`

3.42 $\int \frac{(d+ex^n)^{3/2}}{a+cx^{2n}} dx$

Optimal result	313
Mathematica [F]	313
Rubi [A] (verified)	314
Maple [F]	315
Fricas [F]	316
Sympy [F]	316
Maxima [F]	316
Giac [F]	317
Mupad [F(-1)]	317
Reduce [F]	317

Optimal result

Integrand size = 23, antiderivative size = 146

$$\int \frac{(d+ex^n)^{3/2}}{a+cx^{2n}} dx = \frac{dx\sqrt{d+ex^n} \operatorname{AppellF1}\left(\frac{1}{n}, -\frac{3}{2}, 1, 1 + \frac{1}{n}, -\frac{ex^n}{d}, -\frac{\sqrt{cx^n}}{\sqrt{-a}}\right)}{2a\sqrt{1 + \frac{ex^n}{d}}} \\ + \frac{dx\sqrt{d+ex^n} \operatorname{AppellF1}\left(\frac{1}{n}, -\frac{3}{2}, 1, 1 + \frac{1}{n}, -\frac{ex^n}{d}, \frac{\sqrt{cx^n}}{\sqrt{-a}}\right)}{2a\sqrt{1 + \frac{ex^n}{d}}}$$

output
$$\frac{1/2*d*x*(d+e*x^n)^(1/2)*\operatorname{AppellF1}(1/n, 1, -3/2, 1+1/n, -c^(1/2)*x^n/(-a)^(1/2), -e*x^n/d)/a/(1+e*x^n/d)^(1/2)+1/2*d*x*(d+e*x^n)^(1/2)*\operatorname{AppellF1}(1/n, 1, -3/2, 1+1/n, c^(1/2)*x^n/(-a)^(1/2), -e*x^n/d)/a/(1+e*x^n/d)^(1/2)}$$

Mathematica [F]

$$\int \frac{(d+ex^n)^{3/2}}{a+cx^{2n}} dx = \int \frac{(d+ex^n)^{3/2}}{a+cx^{2n}} dx$$

input
$$\text{Integrate}[(d + e*x^n)^(3/2)/(a + c*x^(2*n)), x]$$

output Integrate[(d + e*x^n)^(3/2)/(a + c*x^(2*n)), x]

Rubi [A] (verified)

Time = 0.34 (sec), antiderivative size = 146, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.130, Rules used = {1759, 937, 936}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d + ex^n)^{3/2}}{a + cx^{2n}} dx \\
 & \quad \downarrow \textcolor{blue}{1759} \\
 & - \frac{\sqrt{c} \int \frac{(ex^n + d)^{3/2}}{\sqrt{-a}\sqrt{c}-cx^n} dx}{2\sqrt{-a}} - \frac{\sqrt{c} \int \frac{(ex^n + d)^{3/2}}{cx^n + \sqrt{-a}\sqrt{c}} dx}{2\sqrt{-a}} \\
 & \quad \downarrow \textcolor{blue}{937} \\
 & - \frac{\sqrt{cd}\sqrt{d+ex^n} \int \frac{\left(\frac{ex^n}{d}+1\right)^{3/2}}{\sqrt{-a}\sqrt{c}-cx^n} dx}{2\sqrt{-a}\sqrt{\frac{ex^n}{d}+1}} - \frac{\sqrt{cd}\sqrt{d+ex^n} \int \frac{\left(\frac{ex^n}{d}+1\right)^{3/2}}{cx^n + \sqrt{-a}\sqrt{c}} dx}{2\sqrt{-a}\sqrt{\frac{ex^n}{d}+1}} \\
 & \quad \downarrow \textcolor{blue}{936} \\
 & \frac{dx\sqrt{d+ex^n} \operatorname{AppellF1}\left(\frac{1}{n}, 1, -\frac{3}{2}, 1 + \frac{1}{n}, -\frac{\sqrt{c}x^n}{\sqrt{-a}}, -\frac{ex^n}{d}\right)}{2a\sqrt{\frac{ex^n}{d}+1}} + \\
 & \quad \frac{dx\sqrt{d+ex^n} \operatorname{AppellF1}\left(\frac{1}{n}, 1, -\frac{3}{2}, 1 + \frac{1}{n}, \frac{\sqrt{c}x^n}{\sqrt{-a}}, -\frac{ex^n}{d}\right)}{2a\sqrt{\frac{ex^n}{d}+1}}
 \end{aligned}$$

input Int[(d + e*x^n)^(3/2)/(a + c*x^(2*n)), x]

output

$$(d*x*sqrt[d + e*x^n]*AppellF1[n^(-1), 1, -3/2, 1 + n^(-1), -((sqrt[c]*x^n)/sqrt[-a]), -((e*x^n)/d)]/(2*a*sqrt[1 + (e*x^n)/d]) + (d*x*sqrt[d + e*x^n]*AppellF1[n^(-1), 1, -3/2, 1 + n^(-1), (sqrt[c]*x^n)/sqrt[-a], -((e*x^n)/d)]/(2*a*sqrt[1 + (e*x^n)/d]))$$

Definitions of rubi rules used

rule 936

$$\text{Int}[(a_.) + (b_.)*(x_)^{(n_.)}^{(p_.)}*((c_.) + (d_.)*(x_)^{(n_.)}^{(q_.)}), x_Symbol] \rightarrow \text{Simp}[a^p*c^q*x^{\text{AppellF1}[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)]}, x] /; \text{FreeQ}[\{a, b, c, d, n, p, q\}, x] \&& \text{NeQ}[b*c - a*d, 0] \&& \text{NeQ}[n, -1] \&& (\text{IntegerQ}[p] \&& \text{GtQ}[a, 0]) \&& (\text{IntegerQ}[q] \&& \text{GtQ}[c, 0])$$

rule 937

$$\text{Int}[(a_.) + (b_.)*(x_)^{(n_.)}^{(p_.)}*((c_.) + (d_.)*(x_)^{(n_.)}^{(q_.)}), x_Symbol] \rightarrow \text{Simp}[a^{\text{IntPart}[p]}*((a + b*x^n)^{\text{FracPart}[p]}/(1 + b*(x^n/a))^{\text{FracPart}[p]}) \text{Int}[(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /; \text{FreeQ}[\{a, b, c, d, n, p, q\}, x] \&& \text{NeQ}[b*c - a*d, 0] \&& \text{NeQ}[n, -1] \&& !(\text{IntegerQ}[p] \&& \text{GtQ}[a, 0])$$

rule 1759

$$\text{Int}[(d_.) + (e_.)*(x_)^{(n_.)}^{(q_.)}/((a_.) + (c_.)*(x_)^{(n2_.)}), x_Symbol] \rightarrow \text{With}[\{r = \text{Rt}[-a*c, 2]\}, \text{Simp}[-c/(2*r) \text{Int}[(d + e*x^n)^q/(r - c*x^n), x], x] - \text{Simp}[c/(2*r) \text{Int}[(d + e*x^n)^q/(r + c*x^n), x], x]] /; \text{FreeQ}[\{a, c, d, e, n, q\}, x] \&& \text{EqQ}[n2, 2*n] \&& \text{NeQ}[c*d^2 + a*e^2, 0] \&& !\text{IntegerQ}[q]]$$

Maple [F]

$$\int \frac{(d + e x^n)^{\frac{3}{2}}}{a + c x^{2n}} dx$$

input

$$\text{int}((d+e*x^n)^(3/2)/(a+c*x^(2*n)), x)$$

output

$$\text{int}((d+e*x^n)^(3/2)/(a+c*x^(2*n)), x)$$

Fricas [F]

$$\int \frac{(d + ex^n)^{3/2}}{a + cx^{2n}} dx = \int \frac{(ex^n + d)^{\frac{3}{2}}}{cx^{2n} + a} dx$$

input `integrate((d+e*x^n)^(3/2)/(a+c*x^(2*n)),x, algorithm="fricas")`

output `integral((e*x^n + d)^(3/2)/(c*x^(2*n) + a), x)`

Sympy [F]

$$\int \frac{(d + ex^n)^{3/2}}{a + cx^{2n}} dx = \int \frac{(d + ex^n)^{\frac{3}{2}}}{a + cx^{2n}} dx$$

input `integrate((d+e*x**n)**(3/2)/(a+c*x**(2*n)),x)`

output `Integral((d + e*x**n)**(3/2)/(a + c*x**(2*n)), x)`

Maxima [F]

$$\int \frac{(d + ex^n)^{3/2}}{a + cx^{2n}} dx = \int \frac{(ex^n + d)^{\frac{3}{2}}}{cx^{2n} + a} dx$$

input `integrate((d+e*x^n)^(3/2)/(a+c*x^(2*n)),x, algorithm="maxima")`

output `integrate((e*x^n + d)^(3/2)/(c*x^(2*n) + a), x)`

Giac [F]

$$\int \frac{(d + ex^n)^{3/2}}{a + cx^{2n}} dx = \int \frac{(ex^n + d)^{\frac{3}{2}}}{cx^{2n} + a} dx$$

input `integrate((d+e*x^n)^(3/2)/(a+c*x^(2*n)),x, algorithm="giac")`

output `integrate((e*x^n + d)^(3/2)/(c*x^(2*n) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(d + ex^n)^{3/2}}{a + cx^{2n}} dx = \int \frac{(d + e x^n)^{3/2}}{a + c x^{2n}} dx$$

input `int((d + e*x^n)^(3/2)/(a + c*x^(2*n)),x)`

output `int((d + e*x^n)^(3/2)/(a + c*x^(2*n)), x)`

Reduce [F]

$$\int \frac{(d + ex^n)^{3/2}}{a + cx^{2n}} dx = \left(\int \frac{\sqrt{x^n e + d}}{x^{2n} c + a} dx \right) d + \left(\int \frac{x^n \sqrt{x^n e + d}}{x^{2n} c + a} dx \right) e$$

input `int((d+e*x^n)^(3/2)/(a+c*x^(2*n)),x)`

output `int(sqrt(x**n*e + d)/(x**2*n*c + a),x)*d + int((x**n*sqrt(x**n*e + d))/((x**2*n*c + a),x)*e`

3.43 $\int \frac{\sqrt{d+ex^n}}{a+cx^{2n}} dx$

Optimal result	318
Mathematica [F]	318
Rubi [A] (verified)	319
Maple [F]	320
Fricas [F]	321
Sympy [F]	321
Maxima [F]	321
Giac [F]	322
Mupad [F(-1)]	322
Reduce [F]	322

Optimal result

Integrand size = 23, antiderivative size = 144

$$\begin{aligned} \int \frac{\sqrt{d+ex^n}}{a+cx^{2n}} dx &= \frac{x\sqrt{d+ex^n} \operatorname{AppellF1}\left(\frac{1}{n}, -\frac{1}{2}, 1, 1 + \frac{1}{n}, -\frac{ex^n}{d}, -\frac{\sqrt{cx^n}}{\sqrt{-a}}\right)}{2a\sqrt{1 + \frac{ex^n}{d}}} \\ &+ \frac{x\sqrt{d+ex^n} \operatorname{AppellF1}\left(\frac{1}{n}, -\frac{1}{2}, 1, 1 + \frac{1}{n}, -\frac{ex^n}{d}, \frac{\sqrt{cx^n}}{\sqrt{-a}}\right)}{2a\sqrt{1 + \frac{ex^n}{d}}} \end{aligned}$$

output
$$\begin{aligned} &1/2*x*(d+e*x^n)^(1/2)*\operatorname{AppellF1}(1/n, 1, -1/2, 1+1/n, -c^(1/2)*x^n/(-a)^(1/2), -e*x^n/d)/a/(1+e*x^n/d)^(1/2)+1/2*x*(d+e*x^n)^(1/2)*\operatorname{AppellF1}(1/n, 1, -1/2, 1+1/n, c^(1/2)*x^n/(-a)^(1/2), -e*x^n/d)/a/(1+e*x^n/d)^(1/2) \end{aligned}$$

Mathematica [F]

$$\int \frac{\sqrt{d+ex^n}}{a+cx^{2n}} dx = \int \frac{\sqrt{d+ex^n}}{a+cx^{2n}} dx$$

input $\text{Integrate}[\text{Sqrt}[d + e*x^n]/(a + c*x^{(2*n)}), x]$

output $\text{Integrate}[\text{Sqrt}[d + e*x^n]/(a + c*x^{(2*n)}), x]$

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.130, Rules used = {1759, 937, 936}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{d + ex^n}}{a + cx^{2n}} dx \\
 & \quad \downarrow 1759 \\
 & - \frac{\sqrt{c} \int \frac{\sqrt{ex^n + d}}{\sqrt{-a\sqrt{c} - cx^n}} dx}{2\sqrt{-a}} - \frac{\sqrt{c} \int \frac{\sqrt{ex^n + d}}{cx^n + \sqrt{-a}\sqrt{c}} dx}{2\sqrt{-a}} \\
 & \quad \downarrow 937 \\
 & - \frac{\sqrt{c}\sqrt{d + ex^n} \int \frac{\sqrt{\frac{ex^n}{d} + 1}}{\sqrt{-a\sqrt{c} - cx^n}} dx}{2\sqrt{-a}\sqrt{\frac{ex^n}{d} + 1}} - \frac{\sqrt{c}\sqrt{d + ex^n} \int \frac{\sqrt{\frac{ex^n}{d} + 1}}{cx^n + \sqrt{-a}\sqrt{c}} dx}{2\sqrt{-a}\sqrt{\frac{ex^n}{d} + 1}} \\
 & \quad \downarrow 936 \\
 & \frac{x\sqrt{d + ex^n} \text{AppellF1}\left(\frac{1}{n}, 1, -\frac{1}{2}, 1 + \frac{1}{n}, -\frac{\sqrt{c}x^n}{\sqrt{-a}}, -\frac{ex^n}{d}\right)}{2a\sqrt{\frac{ex^n}{d} + 1}} + \\
 & \frac{x\sqrt{d + ex^n} \text{AppellF1}\left(\frac{1}{n}, 1, -\frac{1}{2}, 1 + \frac{1}{n}, \frac{\sqrt{c}x^n}{\sqrt{-a}}, -\frac{ex^n}{d}\right)}{2a\sqrt{\frac{ex^n}{d} + 1}}
 \end{aligned}$$

input $\text{Int}[\text{Sqrt}[d + e*x^n]/(a + c*x^{(2*n)}), x]$

output

$$(x*\text{Sqrt}[d + e*x^n]*\text{AppellF1}[n^(-1), 1, -1/2, 1 + n^(-1), -((\text{Sqrt}[c]*x^n)/\text{Sqrt}[-a]), -((e*x^n)/d)])/(2*a*\text{Sqrt}[1 + (e*x^n)/d]) + (x*\text{Sqrt}[d + e*x^n]*\text{AppellF1}[n^(-1), 1, -1/2, 1 + n^(-1), (\text{Sqrt}[c]*x^n)/\text{Sqrt}[-a], -((e*x^n)/d)])/(2*a*\text{Sqrt}[1 + (e*x^n)/d])$$

Defintions of rubi rules used

rule 936

$$\begin{aligned} \text{Int}[(a_.) + (b_.)*(x_.)^{(n_.)}^{(p_.)}*((c_.) + (d_.)*(x_.)^{(n_.)}^{(q_.)}), x_Symbol] \\ :> \text{Simp}[a^p*c^q*x*\text{AppellF1}[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /; \text{FreeQ}[\{a, b, c, d, n, p, q\}, x] \&& \text{NeQ}[b*c - a*d, 0] \&& \text{NeQ}[n, -1] \\ \&\& (\text{IntegerQ}[p] \&& \text{GtQ}[a, 0]) \&& (\text{IntegerQ}[q] \&& \text{GtQ}[c, 0]) \end{aligned}$$

rule 937

$$\begin{aligned} \text{Int}[(a_.) + (b_.)*(x_.)^{(n_.)}^{(p_.)}*((c_.) + (d_.)*(x_.)^{(n_.)}^{(q_.)}), x_Symbol] \\ :> \text{Simp}[a^{\text{IntPart}[p]}*((a + b*x^n)^{\text{FracPart}[p]}/(1 + b*(x^n/a))^{\text{FracPart}[p]}) \\ \text{Int}[(1 + b*(x^n/a))^{p*}(c + d*x^n)^q, x], x] /; \text{FreeQ}[\{a, b, c, d, n, p, q\}, x] \&& \text{NeQ}[b*c - a*d, 0] \&& \text{NeQ}[n, -1] \&& !(\text{IntegerQ}[p] \&& \text{GtQ}[a, 0]) \end{aligned}$$

rule 1759

$$\begin{aligned} \text{Int}[(d_.) + (e_.)*(x_.)^{(n_.)}^{(q_.)}/((a_.) + (c_.)*(x_.)^{(n2_.)}), x_Symbol] :> \text{With}[\{r = \text{Rt}[(-a)*c, 2]\}, \text{Simp}[-c/(2*r) \text{Int}[(d + e*x^n)^q/(r - c*x^n), x], x] - \text{Simp}[c/(2*r) \text{Int}[(d + e*x^n)^q/(r + c*x^n), x], x]] /; \text{FreeQ}[\{a, c, d, e, n, q\}, x] \&& \text{EqQ}[n2, 2*n] \&& \text{NeQ}[c*d^2 + a*e^2, 0] \&& !\text{IntegerQ}[q]] \end{aligned}$$

Maple [F]

$$\int \frac{\sqrt{d + e x^n}}{a + c x^{2n}} dx$$

input

$$\text{int}((d+e*x^n)^(1/2)/(a+c*x^(2*n)), x)$$

output

$$\text{int}((d+e*x^n)^(1/2)/(a+c*x^(2*n)), x)$$

Fricas [F]

$$\int \frac{\sqrt{d + ex^n}}{a + cx^{2n}} dx = \int \frac{\sqrt{ex^n + d}}{cx^{2n} + a} dx$$

input `integrate((d+e*x^n)^(1/2)/(a+c*x^(2*n)),x, algorithm="fricas")`

output `integral(sqrt(e*x^n + d)/(c*x^(2*n) + a), x)`

Sympy [F]

$$\int \frac{\sqrt{d + ex^n}}{a + cx^{2n}} dx = \int \frac{\sqrt{d + ex^n}}{a + cx^{2n}} dx$$

input `integrate((d+e*x**n)**(1/2)/(a+c*x**(2*n)),x)`

output `Integral(sqrt(d + e*x**n)/(a + c*x**(2*n)), x)`

Maxima [F]

$$\int \frac{\sqrt{d + ex^n}}{a + cx^{2n}} dx = \int \frac{\sqrt{ex^n + d}}{cx^{2n} + a} dx$$

input `integrate((d+e*x^n)^(1/2)/(a+c*x^(2*n)),x, algorithm="maxima")`

output `integrate(sqrt(e*x^n + d)/(c*x^(2*n) + a), x)`

Giac [F]

$$\int \frac{\sqrt{d + ex^n}}{a + cx^{2n}} dx = \int \frac{\sqrt{ex^n + d}}{cx^{2n} + a} dx$$

input `integrate((d+e*x^n)^(1/2)/(a+c*x^(2*n)),x, algorithm="giac")`

output `integrate(sqrt(e*x^n + d)/(c*x^(2*n) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{d + ex^n}}{a + cx^{2n}} dx = \int \frac{\sqrt{d + e x^n}}{a + c x^{2n}} dx$$

input `int((d + e*x^n)^(1/2)/(a + c*x^(2*n)),x)`

output `int((d + e*x^n)^(1/2)/(a + c*x^(2*n)), x)`

Reduce [F]

$$\int \frac{\sqrt{d + ex^n}}{a + cx^{2n}} dx = \int \frac{\sqrt{x^n e + d}}{x^{2n} c + a} dx$$

input `int((d+e*x^n)^(1/2)/(a+c*x^(2*n)),x)`

output `int(sqrt(x**n*e + d)/(x**2*n*c + a),x)`

3.44 $\int \frac{1}{\sqrt{d+ex^n}(a+cx^{2n})} dx$

Optimal result	323
Mathematica [F]	323
Rubi [A] (verified)	324
Maple [F]	325
Fricas [F]	325
Sympy [F]	326
Maxima [F]	326
Giac [F]	326
Mupad [F(-1)]	327
Reduce [F]	327

Optimal result

Integrand size = 23, antiderivative size = 144

$$\int \frac{1}{\sqrt{d+ex^n}(a+cx^{2n})} dx = \frac{x \sqrt{1 + \frac{ex^n}{d}} \operatorname{AppellF1} \left(\frac{1}{n}, \frac{1}{2}, 1, 1 + \frac{1}{n}, -\frac{ex^n}{d}, -\frac{\sqrt{cx^n}}{\sqrt{-a}} \right)}{2a\sqrt{d+ex^n}} \\ + \frac{x \sqrt{1 + \frac{ex^n}{d}} \operatorname{AppellF1} \left(\frac{1}{n}, \frac{1}{2}, 1, 1 + \frac{1}{n}, -\frac{ex^n}{d}, \frac{\sqrt{cx^n}}{\sqrt{-a}} \right)}{2a\sqrt{d+ex^n}}$$

output $1/2*x*(1+e*x^n/d)^(1/2)*\operatorname{AppellF1}(1/n, 1, 1/2, 1+1/n, -c^(1/2)*x^n/(-a)^(1/2), -e*x^n/d)/a/(d+e*x^n)^(1/2)+1/2*x*(1+e*x^n/d)^(1/2)*\operatorname{AppellF1}(1/n, 1, 1/2, 1+1/n, c^(1/2)*x^n/(-a)^(1/2), -e*x^n/d)/a/(d+e*x^n)^(1/2)$

Mathematica [F]

$$\int \frac{1}{\sqrt{d+ex^n}(a+cx^{2n})} dx = \int \frac{1}{\sqrt{d+ex^n}(a+cx^{2n})} dx$$

input `Integrate[1/(Sqrt[d + e*x^n]*(a + c*x^(2*n))), x]`

output $\text{Integrate}[1/(\text{Sqrt}[d + e*x^n]*(a + c*x^{(2*n)})), x]$

Rubi [A] (verified)

Time = 0.32 (sec), antiderivative size = 144, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {1759, 937, 936}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a + cx^{2n}) \sqrt{d + ex^n}} dx \\
 & \quad \downarrow \textcolor{blue}{1759} \\
 & - \frac{\sqrt{c} \int \frac{1}{(\sqrt{-a}\sqrt{c}-cx^n)\sqrt{ex^n+d}} dx}{2\sqrt{-a}} - \frac{\sqrt{c} \int \frac{1}{(cx^n+\sqrt{-a}\sqrt{c})\sqrt{ex^n+d}} dx}{2\sqrt{-a}} \\
 & \quad \downarrow \textcolor{blue}{937} \\
 & - \frac{\sqrt{c}\sqrt{\frac{ex^n}{d}+1} \int \frac{1}{(\sqrt{-a}\sqrt{c}-cx^n)\sqrt{\frac{ex^n}{d}+1}} dx}{2\sqrt{-a}\sqrt{d+ex^n}} - \frac{\sqrt{c}\sqrt{\frac{ex^n}{d}+1} \int \frac{1}{(cx^n+\sqrt{-a}\sqrt{c})\sqrt{\frac{ex^n}{d}+1}} dx}{2\sqrt{-a}\sqrt{d+ex^n}} \\
 & \quad \downarrow \textcolor{blue}{936} \\
 & \frac{x\sqrt{\frac{ex^n}{d}+1} \text{AppellF1}\left(\frac{1}{n}, 1, \frac{1}{2}, 1 + \frac{1}{n}, -\frac{\sqrt{c}x^n}{\sqrt{-a}}, -\frac{ex^n}{d}\right)}{2a\sqrt{d+ex^n}} + \\
 & \quad \frac{x\sqrt{\frac{ex^n}{d}+1} \text{AppellF1}\left(\frac{1}{n}, 1, \frac{1}{2}, 1 + \frac{1}{n}, \frac{\sqrt{c}x^n}{\sqrt{-a}}, -\frac{ex^n}{d}\right)}{2a\sqrt{d+ex^n}}
 \end{aligned}$$

input $\text{Int}[1/(\text{Sqrt}[d + e*x^n]*(a + c*x^{(2*n)})), x]$

output $(x*\text{Sqrt}[1 + (e*x^n)/d]*\text{AppellF1}[n^{(-1)}, 1, 1/2, 1 + n^{(-1)}, -((\text{Sqrt}[c]*x^n)/\text{Sqrt}[-a]), -((e*x^n)/d)]/(2*a*\text{Sqrt}[d + e*x^n]) + (x*\text{Sqrt}[1 + (e*x^n)/d]*\text{AppellF1}[n^{(-1)}, 1, 1/2, 1 + n^{(-1)}, (\text{Sqrt}[c]*x^n)/\text{Sqrt}[-a], -((e*x^n)/d)]/(2*a*\text{Sqrt}[d + e*x^n]))$

Definitions of rubi rules used

rule 936 $\text{Int}[(a_+ + b_-) \cdot (x_-)^n \cdot (p_-) \cdot ((c_+ + d_-) \cdot (x_-)^n)^q, x_{\text{Symbol}}] \rightarrow \text{Simp}[a^p \cdot c^q \cdot x^* \text{AppellF1}[1/n, -p, -q, 1 + 1/n, (-b) \cdot (x^n/a), (-d) \cdot (x^n/c)], x] /; \text{FreeQ}[\{a, b, c, d, n, p, q\}, x] \& \text{NeQ}[b*c - a*d, 0] \& \text{NeQ}[n, -1] \& (\text{IntegerQ}[p] \text{ || } \text{GtQ}[a, 0]) \& (\text{IntegerQ}[q] \text{ || } \text{GtQ}[c, 0])$

rule 937 $\text{Int}[(a_+ + b_-) \cdot (x_-)^n \cdot (p_-) \cdot ((c_+ + d_-) \cdot (x_-)^n)^q, x_{\text{Symbol}}] \rightarrow \text{Simp}[a^{\text{IntPart}[p]} \cdot ((a + b*x^n)^{\text{FracPart}[p]} / (1 + b*(x^n/a))^{\text{FracPart}[p]}) \cdot \text{Int}[(1 + b*(x^n/a))^{p_*} \cdot (c + d*x^n)^q, x], x] /; \text{FreeQ}[\{a, b, c, d, n, p, q\}, x] \& \text{NeQ}[b*c - a*d, 0] \& \text{NeQ}[n, -1] \& !(\text{IntegerQ}[p] \text{ || } \text{GtQ}[a, 0])$

rule 1759 $\text{Int}[(d_+ + e_-) \cdot (x_-)^n \cdot (q_-) / ((a_+ + c_-) \cdot (x_-)^{n2}), x_{\text{Symbol}}] \rightarrow \text{With}[\{r = \text{Rt}[-a*c, 2]\}, \text{Simp}[-c/(2*r) \cdot \text{Int}[(d + e*x^n)^q / (r - c*x^n), x], x] - \text{Simp}[c/(2*r) \cdot \text{Int}[(d + e*x^n)^q / (r + c*x^n), x], x]] /; \text{FreeQ}[\{a, c, d, e, n, q\}, x] \& \text{EqQ}[n2, 2*n] \& \text{NeQ}[c*d^2 + a*e^2, 0] \& !\text{IntegerQ}[q]$

Maple [F]

$$\int \frac{1}{\sqrt{d + e x^n} (a + c x^{2n})} dx$$

input `int(1/(d+e*x^n)^(1/2)/(a+c*x^(2*n)),x)`

output `int(1/(d+e*x^n)^(1/2)/(a+c*x^(2*n)),x)`

Fricas [F]

$$\int \frac{1}{\sqrt{d + e x^n} (a + c x^{2n})} dx = \int \frac{1}{(c x^{2n} + a) \sqrt{e x^n + d}} dx$$

input `integrate(1/(d+e*x^n)^(1/2)/(a+c*x^(2*n)),x, algorithm="fricas")`

output `integral(sqrt(e*x^n + d)/(a*e*x^n + a*d + (c*e*x^n + c*d)*x^(2*n)), x)`

Sympy [F]

$$\int \frac{1}{\sqrt{d + ex^n} (a + cx^{2n})} dx = \int \frac{1}{(a + cx^{2n}) \sqrt{d + ex^n}} dx$$

input `integrate(1/(d+e*x**n)**(1/2)/(a+c*x**(2*n)), x)`

output `Integral(1/((a + c*x**(2*n))*sqrt(d + e*x**n)), x)`

Maxima [F]

$$\int \frac{1}{\sqrt{d + ex^n} (a + cx^{2n})} dx = \int \frac{1}{(cx^{2n} + a)\sqrt{ex^n + d}} dx$$

input `integrate(1/(d+e*x^n)^(1/2)/(a+c*x^(2*n)), x, algorithm="maxima")`

output `integrate(1/((c*x^(2*n) + a)*sqrt(e*x^n + d)), x)`

Giac [F]

$$\int \frac{1}{\sqrt{d + ex^n} (a + cx^{2n})} dx = \int \frac{1}{(cx^{2n} + a)\sqrt{ex^n + d}} dx$$

input `integrate(1/(d+e*x^n)^(1/2)/(a+c*x^(2*n)), x, algorithm="giac")`

output `integrate(1/((c*x^(2*n) + a)*sqrt(e*x^n + d)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{d+ex^n} (a+cx^{2n})} dx = \int \frac{1}{(a+cx^{2n}) \sqrt{d+ex^n}} dx$$

input `int(1/((a + c*x^(2*n))*(d + e*x^n)^(1/2)),x)`

output `int(1/((a + c*x^(2*n))*(d + e*x^n)^(1/2)), x)`

Reduce [F]

$$\int \frac{1}{\sqrt{d+ex^n} (a+cx^{2n})} dx = \int \frac{\sqrt{x^n e + d}}{x^{3n} ce + x^{2n} cd + x^n ae + ad} dx$$

input `int(1/(d+e*x^n)^(1/2)/(a+c*x^(2*n)),x)`

output `int(sqrt(x**n*e + d)/(x**(3*n)*c*e + x**(2*n)*c*d + x**n*a*e + a*d),x)`

3.45 $\int \frac{1}{(d+ex^n)^{3/2}(a+cx^{2n})} dx$

Optimal result	328
Mathematica [F]	328
Rubi [A] (verified)	329
Maple [F]	331
Fricas [F]	331
Sympy [F]	332
Maxima [F]	332
Giac [F]	332
Mupad [F(-1)]	333
Reduce [F]	333

Optimal result

Integrand size = 23, antiderivative size = 150

$$\int \frac{1}{(d+ex^n)^{3/2}(a+cx^{2n})} dx = \frac{x\sqrt{1+\frac{ex^n}{d}} \operatorname{AppellF1}\left(\frac{1}{n}, \frac{3}{2}, 1, 1 + \frac{1}{n}, -\frac{ex^n}{d}, -\frac{\sqrt{cx^n}}{\sqrt{-a}}\right)}{2ad\sqrt{d+ex^n}} \\ + \frac{x\sqrt{1+\frac{ex^n}{d}} \operatorname{AppellF1}\left(\frac{1}{n}, \frac{3}{2}, 1, 1 + \frac{1}{n}, -\frac{ex^n}{d}, \frac{\sqrt{cx^n}}{\sqrt{-a}}\right)}{2ad\sqrt{d+ex^n}}$$

output $1/2*x*(1+e*x^n/d)^(1/2)*\operatorname{AppellF1}(1/n, 1, 3/2, 1+1/n, -c^(1/2)*x^n/(-a)^(1/2), -e*x^n/d)/a/d/(d+e*x^n)^(1/2)+1/2*x*(1+e*x^n/d)^(1/2)*\operatorname{AppellF1}(1/n, 1, 3/2, 1+1/n, c^(1/2)*x^n/(-a)^(1/2), -e*x^n/d)/a/d/(d+e*x^n)^(1/2)$

Mathematica [F]

$$\int \frac{1}{(d+ex^n)^{3/2}(a+cx^{2n})} dx = \int \frac{1}{(d+ex^n)^{3/2}(a+cx^{2n})} dx$$

input `Integrate[1/((d + e*x^n)^(3/2)*(a + c*x^(2*n))), x]`

output Integrate[1/((d + e*x^n)^(3/2)*(a + c*x^(2*n))), x]

Rubi [A] (verified)

Time = 0.98 (sec), antiderivative size = 260, normalized size of antiderivative = 1.73, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.217, Rules used = {1757, 779, 778, 7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a + cx^{2n}) (d + ex^n)^{3/2}} dx \\
 & \quad \downarrow \textcolor{blue}{1757} \\
 & \frac{e^2 \int \frac{1}{(ex^n+d)^{3/2}} dx}{ae^2 + cd^2} + \frac{c \int \frac{d-ex^n}{\sqrt{ex^n+d(cx^{2n}+a)}} dx}{ae^2 + cd^2} \\
 & \quad \downarrow \textcolor{blue}{779} \\
 & \frac{e^2 \sqrt{\frac{ex^n}{d} + 1} \int \frac{1}{\left(\frac{ex^n}{d} + 1\right)^{3/2}} dx}{d(ae^2 + cd^2) \sqrt{d + ex^n}} + \frac{c \int \frac{d-ex^n}{\sqrt{ex^n+d(cx^{2n}+a)}} dx}{ae^2 + cd^2} \\
 & \quad \downarrow \textcolor{blue}{778} \\
 & \frac{c \int \frac{d-ex^n}{\sqrt{ex^n+d(cx^{2n}+a)}} dx}{ae^2 + cd^2} + \frac{e^2 x \sqrt{\frac{ex^n}{d} + 1} \text{Hypergeometric2F1}\left(\frac{3}{2}, \frac{1}{n}, 1 + \frac{1}{n}, -\frac{ex^n}{d}\right)}{d(ae^2 + cd^2) \sqrt{d + ex^n}} \\
 & \quad \downarrow \textcolor{blue}{7293} \\
 & \frac{c \int \left(\frac{-d - \frac{\sqrt{-a}e}{\sqrt{c}}}{2\sqrt{-a}\sqrt{c}(x^n + \frac{\sqrt{-a}}{\sqrt{c}})\sqrt{ex^n+d}} - \frac{d - \frac{\sqrt{-a}e}{\sqrt{c}}}{2\sqrt{-a}\sqrt{c}(\frac{\sqrt{-a}}{\sqrt{c}} - x^n)\sqrt{ex^n+d}} \right) dx}{ae^2 + cd^2} + \\
 & \quad \frac{e^2 x \sqrt{\frac{ex^n}{d} + 1} \text{Hypergeometric2F1}\left(\frac{3}{2}, \frac{1}{n}, 1 + \frac{1}{n}, -\frac{ex^n}{d}\right)}{d(ae^2 + cd^2) \sqrt{d + ex^n}} \\
 & \quad \downarrow \textcolor{blue}{2009}
 \end{aligned}$$

$$\begin{aligned}
& c \left(\frac{x \left(\frac{\sqrt{-a}e}{\sqrt{c}} + d \right) \sqrt{\frac{ex^n}{d} + 1} \operatorname{AppellF1} \left(\frac{1}{n}, 1, \frac{1}{2}, 1 + \frac{1}{n}, -\frac{\sqrt{c}x^n}{\sqrt{-a}}, -\frac{ex^n}{d} \right)}{2a\sqrt{d+ex^n}} + \frac{x \left(d - \frac{\sqrt{-a}e}{\sqrt{c}} \right) \sqrt{\frac{ex^n}{d} + 1} \operatorname{AppellF1} \left(\frac{1}{n}, 1, \frac{1}{2}, 1 + \frac{1}{n}, \frac{\sqrt{c}x^n}{\sqrt{-a}}, -\frac{ex^n}{d} \right)}{2a\sqrt{d+ex^n}} \right) + \\
& \frac{ae^2 + cd^2}{e^2 x \sqrt{\frac{ex^n}{d} + 1} \operatorname{Hypergeometric2F1} \left(\frac{3}{2}, \frac{1}{n}, 1 + \frac{1}{n}, -\frac{ex^n}{d} \right)} \\
& \frac{d (ae^2 + cd^2) \sqrt{d + ex^n}}{d (ae^2 + cd^2) \sqrt{d + ex^n}}
\end{aligned}$$

input `Int[1/((d + e*x^n)^(3/2)*(a + c*x^(2*n))), x]`

output

```
(c*((d + (Sqrt[-a]*e)/Sqrt[c])*x*Sqrt[1 + (e*x^n)/d]*AppellF1[n^(-1), 1, 1/2, 1 + n^(-1), -(Sqrt[c]*x^n)/Sqrt[-a]], -((e*x^n)/d))/(2*a*Sqrt[d + e*x^n]) + ((d - (Sqrt[-a]*e)/Sqrt[c])*x*Sqrt[1 + (e*x^n)/d]*AppellF1[n^(-1), 1, 1/2, 1 + n^(-1), (Sqrt[c]*x^n)/Sqrt[-a], -((e*x^n)/d)])/(2*a*Sqrt[d + e*x^n]))/(c*d^2 + a*e^2) + (e^2*x*Sqrt[1 + (e*x^n)/d]*Hypergeometric2F1[3/2, n^(-1), 1 + n^(-1), -((e*x^n)/d)])/(d*(c*d^2 + a*e^2)*Sqrt[d + e*x^n])
```

Definitions of rubi rules used

rule 778 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F1[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] || GtQ[a, 0])`

rule 779 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p]) Int[(1 + b*(x^n/a))^p, x], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && !(IntegerQ[p] || GtQ[a, 0])`

rule 1757 `Int[((d_) + (e_.)*(x_)^(n_))^(q_)/((a_) + (c_.)*(x_)^(n2_)), x_Symbol] := Simp[e^2/(c*d^2 + a*e^2) Int[(d + e*x^n)^q, x], x] + Simp[c/(c*d^2 + a*e^2) Int[(d + e*x^n)^(q + 1)*((d - e*x^n)/(a + c*x^(2*n))), x], x] /; FreeQ[{a, c, d, e, n}, x] && EqQ[n2, 2*n] && NeQ[c*d^2 + a*e^2, 0] && !IntegerQ[q] && LtQ[q, -1]`

rule 2009 $\text{Int}[u_, x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 7293 $\text{Int}[u_, x_{\text{Symbol}}] \rightarrow \text{With}[\{v = \text{ExpandIntegrand}[u, x]\}, \text{Int}[v, x] /; \text{SumQ}[v]]$

Maple [F]

$$\int \frac{1}{(d + e x^n)^{\frac{3}{2}} (a + c x^{2n})} dx$$

input $\text{int}(1/(d+e*x^n)^(3/2)/(a+c*x^(2*n)), x)$

output $\text{int}(1/(d+e*x^n)^(3/2)/(a+c*x^(2*n)), x)$

Fricas [F]

$$\int \frac{1}{(d + e x^n)^{3/2} (a + c x^{2n})} dx = \int \frac{1}{(c x^{2n} + a)(e x^n + d)^{\frac{3}{2}}} dx$$

input $\text{integrate}(1/(d+e*x^n)^(3/2)/(a+c*x^(2*n)), x, \text{algorithm}=\text{"fricas"})$

output $\text{integral}(\sqrt(e*x^n + d)/(a*e^2*x^(2*n) + 2*a*d*e*x^n + a*d^2 + (c*e^2*x^(2*n) + 2*c*d*e*x^n + c*d^2)*x^(2*n)), x)$

Sympy [F]

$$\int \frac{1}{(d + ex^n)^{3/2} (a + cx^{2n})} dx = \int \frac{1}{(a + cx^{2n}) (d + ex^n)^{\frac{3}{2}}} dx$$

input `integrate(1/(d+e*x**n)**(3/2)/(a+c*x**(2*n)),x)`

output `Integral(1/((a + c*x**(2*n))*(d + e*x**n)**(3/2)), x)`

Maxima [F]

$$\int \frac{1}{(d + ex^n)^{3/2} (a + cx^{2n})} dx = \int \frac{1}{(cx^{2n} + a)(ex^n + d)^{\frac{3}{2}}} dx$$

input `integrate(1/(d+e*x^n)^(3/2)/(a+c*x^(2*n)),x, algorithm="maxima")`

output `integrate(1/((c*x^(2*n) + a)*(e*x^n + d))^(3/2), x)`

Giac [F]

$$\int \frac{1}{(d + ex^n)^{3/2} (a + cx^{2n})} dx = \int \frac{1}{(cx^{2n} + a)(ex^n + d)^{\frac{3}{2}}} dx$$

input `integrate(1/(d+e*x^n)^(3/2)/(a+c*x^(2*n)),x, algorithm="giac")`

output `integrate(1/((c*x^(2*n) + a)*(e*x^n + d))^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex^n)^{3/2} (a + cx^{2n})} dx = \int \frac{1}{(a + cx^{2n}) (d + ex^n)^{3/2}} dx$$

input `int(1/((a + c*x^(2*n))*(d + e*x^n)^(3/2)),x)`

output `int(1/((a + c*x^(2*n))*(d + e*x^n)^(3/2)), x)`

Reduce [F]

$$\int \frac{1}{(d + ex^n)^{3/2} (a + cx^{2n})} dx = \int \frac{\sqrt{x^n e + d}}{x^{4n} c e^2 + 2x^{3n} c d e + x^{2n} a e^2 + x^{2n} c d^2 + 2x^n a d e + a d^2} dx$$

input `int(1/(d+e*x^n)^(3/2)/(a+c*x^(2*n)),x)`

output `int(sqrt(x**n*e + d)/(x**(4*n)*c*e**2 + 2*x**(3*n)*c*d*e + x**(2*n)*a*e**2 + x**(2*n)*c*d**2 + 2*x**n*a*d*e + a*d**2),x)`

3.46 $\int \frac{1}{(d+ex^n)\sqrt{a+cx^{2n}}} dx$

Optimal result	334
Mathematica [F]	335
Rubi [A] (verified)	335
Maple [F]	336
Fricas [F]	336
Sympy [F]	337
Maxima [F]	337
Giac [F]	337
Mupad [F(-1)]	338
Reduce [F]	338

Optimal result

Integrand size = 23, antiderivative size = 171

$$\begin{aligned} & \int \frac{1}{(d+ex^n)\sqrt{a+cx^{2n}}} dx \\ &= \frac{x\sqrt{1+\frac{cx^{2n}}{a}} \operatorname{AppellF1}\left(\frac{1}{2n}, \frac{1}{2}, 1, \frac{1}{2}(2+\frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2x^{2n}}{d^2}\right)}{d\sqrt{a+cx^{2n}}} \\ &\quad - \frac{ex^{1+n}\sqrt{1+\frac{cx^{2n}}{a}} \operatorname{AppellF1}\left(\frac{1+n}{2n}, \frac{1}{2}, 1, \frac{1}{2}(3+\frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2x^{2n}}{d^2}\right)}{d^2(1+n)\sqrt{a+cx^{2n}}} \end{aligned}$$

output $x*(1+c*x^(2*n)/a)^(1/2)*AppellF1(1/2/n, 1, 1/2, 1+1/2/n, e^2*x^(2*n)/d^2, -c*x^(2*n)/a)/d/(a+c*x^(2*n))^(1/2)-e*x^(1+n)*(1+c*x^(2*n)/a)^(1/2)*AppellF1(1/2*(1+n)/n, 1, 1/2, 3/2+1/2/n, e^2*x^(2*n)/d^2, -c*x^(2*n)/a)/d^2/(1+n)/(a+c*x^(2*n))^(1/2)$

Mathematica [F]

$$\int \frac{1}{(d + ex^n) \sqrt{a + cx^{2n}}} dx = \int \frac{1}{(d + ex^n) \sqrt{a + cx^{2n}}} dx$$

input `Integrate[1/((d + e*x^n)*Sqrt[a + c*x^(2*n)]), x]`

output `Integrate[1/((d + e*x^n)*Sqrt[a + c*x^(2*n)]), x]`

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {1768, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\sqrt{a + cx^{2n}} (d + ex^n)} dx \\ & \quad \downarrow 1768 \\ & \int \left(\frac{d}{\sqrt{a + cx^{2n}} (d^2 - e^2 x^{2n})} + \frac{ex^n}{\sqrt{a + cx^{2n}} (e^2 x^{2n} - d^2)} \right) dx \\ & \quad \downarrow 2009 \\ & \frac{x \sqrt{\frac{cx^{2n}}{a} + 1} \operatorname{AppellF1} \left(\frac{1}{2n}, \frac{1}{2}, 1, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2 x^{2n}}{d^2} \right)}{d \sqrt{a + cx^{2n}}} - \\ & \frac{ex^{n+1} \sqrt{\frac{cx^{2n}}{a} + 1} \operatorname{AppellF1} \left(\frac{n+1}{2n}, \frac{1}{2}, 1, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2 x^{2n}}{d^2} \right)}{d^2(n+1) \sqrt{a + cx^{2n}}} \end{aligned}$$

input `Int[1/((d + e*x^n)*Sqrt[a + c*x^(2*n)]), x]`

output
$$(x \operatorname{Sqrt}[1 + (c x^{2n})/a] \operatorname{AppellF1}[1/(2n), 1/2, 1, (2 + n^{-1})/2, -((c x^{2n})/a), (e^{2x^{2n}}/d^2)/(d \operatorname{Sqrt}[a + c x^{2n}]) - (e x^{1+n}) \operatorname{Sqr}t[1 + (c x^{2n})/a] \operatorname{AppellF1}[(1+n)/(2n), 1/2, 1, (3 + n^{-1})/2, -((c x^{2n})/a), (e^{2x^{2n}}/d^2)/(d^2(1+n) \operatorname{Sqr}t[a + c x^{2n}])]$$

Defintions of rubi rules used

rule 1768
$$\operatorname{Int}[(d_+ + e_-)(x_-)^{(n_-)} \cdot (a_+ + c_-)(x_-)^{(n_2_-)} \cdot (a_+ + c_-)(x_-)^{(p_-)}, x_{\text{Symbol}}] \rightarrow \operatorname{Int}[\operatorname{ExpandIntegrand}[(a + c x^{2n})^p, (d/(d^2 - e^{2x^{2n}}) - e(x^n/(d^2 - e^{2x^{2n}})))^{(-q)}, x], x] /; \operatorname{FreeQ}[\{a, c, d, e, n, p\}, x] \&& \operatorname{EqQ}[n, 2n] \&& \operatorname{NeQ}[c*d^2 + a*e^2, 0] \&& \operatorname{!IntegerQ}[p] \&& \operatorname{ILtQ}[q, 0]$$

rule 2009
$$\operatorname{Int}[u_, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] /; \operatorname{SumQ}[u]$$

Maple [F]

$$\int \frac{1}{(d + e x^n) \sqrt{a + c x^{2n}}} dx$$

input
$$\operatorname{int}(1/(d+e*x^n)/(a+c*x^(2n))^(1/2), x)$$

output
$$\operatorname{int}(1/(d+e*x^n)/(a+c*x^(2n))^(1/2), x)$$

Fricas [F]

$$\int \frac{1}{(d + e x^n) \sqrt{a + c x^{2n}}} dx = \int \frac{1}{\sqrt{c x^{2n} + a} (e x^n + d)} dx$$

input
$$\operatorname{integrate}(1/(d+e*x^n)/(a+c*x^(2n))^(1/2), x, \operatorname{algorithm}=\text{"fricas"})$$

output
$$\operatorname{integral}(\operatorname{sqrt}(c x^{2n} + a)/(a * e * x^n + a * d + (c * e * x^n + c * d) * x^(2n)), x)$$

Sympy [F]

$$\int \frac{1}{(d + ex^n) \sqrt{a + cx^{2n}}} dx = \int \frac{1}{\sqrt{a + cx^{2n}} (d + ex^n)} dx$$

input `integrate(1/(d+e*x**n)/(a+c*x**(2*n))**(1/2),x)`

output `Integral(1/(sqrt(a + c*x**(2*n))*(d + e*x**n)), x)`

Maxima [F]

$$\int \frac{1}{(d + ex^n) \sqrt{a + cx^{2n}}} dx = \int \frac{1}{\sqrt{cx^{2n} + a}(ex^n + d)} dx$$

input `integrate(1/(d+e*x^n)/(a+c*x^(2*n))^(1/2),x, algorithm="maxima")`

output `integrate(1/(sqrt(c*x^(2*n) + a)*(e*x^n + d)), x)`

Giac [F]

$$\int \frac{1}{(d + ex^n) \sqrt{a + cx^{2n}}} dx = \int \frac{1}{\sqrt{cx^{2n} + a}(ex^n + d)} dx$$

input `integrate(1/(d+e*x^n)/(a+c*x^(2*n))^(1/2),x, algorithm="giac")`

output `integrate(1/(sqrt(c*x^(2*n) + a)*(e*x^n + d)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex^n) \sqrt{a + cx^{2n}}} dx = \int \frac{1}{\sqrt{a + cx^{2n}} (d + e x^n)} dx$$

input `int(1/((a + c*x^(2*n))^(1/2)*(d + e*x^n)),x)`

output `int(1/((a + c*x^(2*n))^(1/2)*(d + e*x^n)), x)`

Reduce [F]

$$\int \frac{1}{(d + ex^n) \sqrt{a + cx^{2n}}} dx = \int \frac{\sqrt{x^{2n}c + a}}{x^{3n}ce + x^{2n}cd + x^nae + ad} dx$$

input `int(1/(d+e*x^n)/(a+c*x^(2*n))^(1/2),x)`

output `int(sqrt(x^(2*n)*c + a)/(x^(3*n)*c*e + x^(2*n)*c*d + x^n*a*e + a*d),x)`

3.47 $\int (d + ex^n)^3 (a + cx^{2n})^p dx$

Optimal result	339
Mathematica [A] (verified)	340
Rubi [A] (verified)	340
Maple [F]	342
Fricas [F]	342
Sympy [F(-1)]	342
Maxima [F]	343
Giac [F(-2)]	343
Mupad [F(-1)]	343
Reduce [F]	344

Optimal result

Integrand size = 21, antiderivative size = 263

$$\begin{aligned} \int (d + ex^n)^3 (a + cx^{2n})^p dx = & \frac{3de^2x(a + cx^{2n})^{1+p}}{c(1 + 2n(1 + p))} + \frac{e^3x^{1+n}(a + cx^{2n})^{1+p}}{c(1 + n(3 + 2p))} \\ & - \frac{d(3ae^2 - cd^2(1 + 2n(1 + p)))x(a + cx^{2n})^p \left(1 + \frac{cx^{2n}}{a}\right)^{-p} \text{Hypergeometric2F1}\left(\frac{1}{2n}, -p, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^2}{a}\right)}{c(1 + 2n(1 + p))} \\ & + e\left(\frac{3d^2}{1 + n} - \frac{ae^2}{c + cn(3 + 2p)}\right)x^{1+n}(a + cx^{2n})^p \left(1 + \frac{cx^{2n}}{a}\right)^{-p} \text{Hypergeometric2F1}\left(\frac{1 + n}{2n}, -p, \frac{1}{2}\left(3 + \frac{1}{n}\right), -\frac{cx^{2n}}{a}\right) \end{aligned}$$

```
output 3*d*e^2*x*(a+c*x^(2*n))^(p+1)/c/(1+2*n*(p+1))+e^3*x^(1+n)*(a+c*x^(2*n))^(p+1)/c/(1+n*(3+2*p))-d*(3*a*e^2-c*d^2*(1+2*n*(p+1)))*x*(a+c*x^(2*n))^p*hypergeom([-p, 1/2/n], [1+1/2/n], -c*x^(2*n)/a)/c/(1+2*n*(p+1))/((1+c*x^(2*n)/a)^p)+e*(3*d^2/(1+n)-a*e^2/(c+c*n*(3+2*p)))*x^(1+n)*(a+c*x^(2*n))^p*hypergeom([-p, 1/2*(1+n)/n], [3/2+1/2/n], -c*x^(2*n)/a)/((1+c*x^(2*n)/a)^p)
```

Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 213, normalized size of antiderivative = 0.81

$$\begin{aligned}
 & \int (d + ex^n)^3 (a + cx^{2n})^p dx \\
 &= x(a + cx^{2n})^p \left(1 \right. \\
 &\quad \left. + \frac{cx^{2n}}{a} \right)^{-p} \left(\frac{3de^2x^{2n} \text{Hypergeometric2F1} \left(\frac{1}{2}(2 + \frac{1}{n}), -p, \frac{1}{2}(4 + \frac{1}{n}), -\frac{cx^{2n}}{a} \right)}{1 + 2n} \right. \\
 &\quad \left. + \frac{e^3x^{3n} \text{Hypergeometric2F1} \left(\frac{1}{2}(3 + \frac{1}{n}), -p, \frac{1}{2}(5 + \frac{1}{n}), -\frac{cx^{2n}}{a} \right)}{1 + 3n} \right. \\
 &\quad \left. + d^2 \left(d \text{Hypergeometric2F1} \left(\frac{1}{2n}, -p, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a} \right) \right. \right. \\
 &\quad \left. \left. + \frac{3ex^n \text{Hypergeometric2F1} \left(\frac{1+n}{2n}, -p, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a} \right)}{1 + n} \right) \right)
 \end{aligned}$$

input `Integrate[(d + e*x^n)^3*(a + c*x^(2*n))^p, x]`

output `(x*(a + c*x^(2*n))^p*((3*d*e^2*x^(2*n)*Hypergeometric2F1[(2 + n^(-1))/2, -p, (4 + n^(-1))/2, -((c*x^(2*n))/a)]/(1 + 2*n) + (e^3*x^(3*n)*Hypergeometric2F1[(3 + n^(-1))/2, -p, (5 + n^(-1))/2, -((c*x^(2*n))/a)]/(1 + 3*n) + d^2*(d*Hypergeometric2F1[1/(2*n), -p, (2 + n^(-1))/2, -((c*x^(2*n))/a)] + (3*e*x^n*Hypergeometric2F1[(1 + n)/(2*n), -p, (3 + n^(-1))/2, -((c*x^(2*n))/a)]/(1 + (c*x^(2*n))/a))^p)/(1 + (c*x^(2*n))/a)))/(1 + (c*x^(2*n))/a)^p`

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 299, normalized size of antiderivative = 1.14, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {1767, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (d + ex^n)^3 (a + cx^{2n})^p dx \\
 & \quad \downarrow \textcolor{blue}{1767} \\
 & \int (d^3(a + cx^{2n})^p + 3d^2ex^n(a + cx^{2n})^p + 3de^2x^{2n}(a + cx^{2n})^p + e^3x^{3n}(a + cx^{2n})^p) dx \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & \frac{d^3x(a + cx^{2n})^p \left(\frac{cx^{2n}}{a} + 1\right)^{-p} \text{Hypergeometric2F1}\left(\frac{1}{2n}, -p, \frac{1}{2}\left(2 + \frac{1}{n}\right), -\frac{cx^{2n}}{a}\right) +}{n+1} \\
 & \quad \frac{3d^2ex^{n+1}(a + cx^{2n})^p \left(\frac{cx^{2n}}{a} + 1\right)^{-p} \text{Hypergeometric2F1}\left(\frac{n+1}{2n}, -p, \frac{1}{2}\left(3 + \frac{1}{n}\right), -\frac{cx^{2n}}{a}\right) +}{n+1} \\
 & \quad \frac{3de^2x^{2n+1}(a + cx^{2n})^p \left(\frac{cx^{2n}}{a} + 1\right)^{-p} \text{Hypergeometric2F1}\left(\frac{1}{2}\left(2 + \frac{1}{n}\right), -p, \frac{1}{2}\left(4 + \frac{1}{n}\right), -\frac{cx^{2n}}{a}\right) +}{2n+1} \\
 & \quad \frac{e^3x^{3n+1}(a + cx^{2n})^p \left(\frac{cx^{2n}}{a} + 1\right)^{-p} \text{Hypergeometric2F1}\left(\frac{1}{2}\left(3 + \frac{1}{n}\right), -p, \frac{1}{2}\left(5 + \frac{1}{n}\right), -\frac{cx^{2n}}{a}\right)}{3n+1}
 \end{aligned}$$

input `Int[(d + e*x^n)^3*(a + c*x^(2*n))^p, x]`

output

$$\begin{aligned}
 & (3*d*e^2*x^(1 + 2*n)*(a + c*x^(2*n))^p*\text{Hypergeometric2F1}[(2 + n^(-1))/2, -p, (4 + n^(-1))/2, -((c*x^(2*n))/a)])/((1 + 2*n)*(1 + (c*x^(2*n))/a)^p) + \\
 & (e^3*x^(1 + 3*n)*(a + c*x^(2*n))^p*\text{Hypergeometric2F1}[(3 + n^(-1))/2, -p, (5 + n^(-1))/2, -((c*x^(2*n))/a)])/((1 + 3*n)*(1 + (c*x^(2*n))/a)^p) + (d^3*x*(a + c*x^(2*n))^p*\text{Hypergeometric2F1}[1/(2*n), -p, (2 + n^(-1))/2, -((c*x^(2*n))/a)])/(1 + (c*x^(2*n))/a)^p + (3*d^2*x*(a + c*x^(2*n))^p*\text{Hypergeometric2F1}[(1 + n)/(2*n), -p, (3 + n^(-1))/2, -((c*x^(2*n))/a)])/((1 + n)*(1 + (c*x^(2*n))/a)^p)
 \end{aligned}$$

Definitions of rubi rules used

rule 1767

```

Int[((d_) + (e_.)*(x_)^(n_))^(q_)*((a_) + (c_.)*(x_)^(n2_))^(p_), x_Symbol]
  :> Int[ExpandIntegrand[(d + e*x^n)^q*(a + c*x^(2*n))^p, x], x] /; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && NeQ[c*d^2 + a*e^2, 0] && ((IntegersQ[p, q] && !IntegerQ[n]) || IGtQ[p, 0] || (IGtQ[q, 0] && !IntegerQ[n]))
)

```

rule 2009 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \ x], \ x] /; \ \text{SumQ}[u]$

Maple [F]

$$\int (d + e x^n)^3 (a + c x^{2n})^p dx$$

input $\text{int}((d+e*x^n)^3*(a+c*x^(2*n))^p, x)$

output $\text{int}((d+e*x^n)^3*(a+c*x^(2*n))^p, x)$

Fricas [F]

$$\int (d + e x^n)^3 (a + c x^{2n})^p dx = \int (e x^n + d)^3 (c x^{2n} + a)^p dx$$

input $\text{integrate}((d+e*x^n)^3*(a+c*x^(2*n))^p, x, \text{algorithm}=\text{"fricas"})$

output $\text{integral}(e^{3*x^(3*n)} + 3*d*e^{2*x^(2*n)} + 3*d^2*e*x^n + d^3)*(c*x^(2*n) + a)^p, x)$

Sympy [F(-1)]

Timed out.

$$\int (d + e x^n)^3 (a + c x^{2n})^p dx = \text{Timed out}$$

input $\text{integrate}((d+e*x**n)**3*(a+c*x**2*n)**p, x)$

output Timed out

Maxima [F]

$$\int (d + ex^n)^3 (a + cx^{2n})^p dx = \int (ex^n + d)^3 (cx^{2n} + a)^p dx$$

input `integrate((d+e*x^n)^3*(a+c*x^(2*n))^p,x, algorithm="maxima")`

output `integrate((e*x^n + d)^3*(c*x^(2*n) + a)^p, x)`

Giac [F(-2)]

Exception generated.

$$\int (d + ex^n)^3 (a + cx^{2n})^p dx = \text{Exception raised: TypeError}$$

input `integrate((d+e*x^n)^3*(a+c*x^(2*n))^p,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:Unable to divide, perhaps due to ro
unding error%%{96,[1,0,6,4,3,5,4,1,2]}+%%{480,[1,0,6,4,3,4,4,1,2]}
+%%{960,`

Mupad [F(-1)]

Timed out.

$$\int (d + ex^n)^3 (a + cx^{2n})^p dx = \int (a + cx^{2n})^p (d + ex^n)^3 dx$$

input `int((a + c*x^(2*n))^p*(d + e*x^n)^3,x)`

output `int((a + c*x^(2*n))^p*(d + e*x^n)^3, x)`

Reduce [F]

$$\int (d + ex^n)^3 (a + cx^{2n})^p dx = \text{too large to display}$$

input `int((d+e*x^n)^3*(a+c*x^(2*n))^p,x)`

output

```
(8*x**3*(3*n)*(x**2*c + a)**p*c*e**3*n**3*p**3*x + 12*x**3*(3*n)*(x**2*n)
*c + a)**p*c*e**3*n**3*p**2*x + 4*x**3*(3*n)*(x**2*c + a)**p*c*e**3*n**3
*p*x + 12*x**3*(3*n)*(x**2*c + a)**p*c*e**3*n**2*p**2*x + 12*x**3*(3*n)*(x
**2*c + a)**p*c*e**3*n**2*p*x + 2*x**3*(3*n)*(x**2*c + a)**p*c*e**3*
n**2*x + 6*x**3*(3*n)*(x**2*c + a)**p*c*e**3*n*p*x + 3*x**3*(3*n)*(x**2*n
)*c + a)**p*c*e**3*n*x + x**3*(3*n)*(x**2*c + a)**p*c*e**3*x + 24*x**2*(2*
n)*(x**2*c + a)**p*c*d*e**2*n**3*p**3*x + 48*x**2*(2*n)*(x**2*c + a)
**p*c*d*e**2*n**3*p**2*x + 18*x**2*(2*n)*(x**2*c + a)**p*c*d*e**2*n**3*p
*x + 36*x**2*(2*n)*(x**2*c + a)**p*c*d*e**2*n**2*p**2*x + 48*x**2*(2*n)*(x
**2*c + a)**p*c*d*e**2*n**2*p*x + 9*x**2*(2*n)*(x**2*c + a)**p*c*d*e
**2*n**2*x + 18*x**2*(2*n)*(x**2*c + a)**p*c*d*e**2*n*p*x + 12*x**2*(2*n)*
(x**2*c + a)**p*c*d*e**2*n*x + 3*x**2*(2*n)*(x**2*c + a)**p*c*d*e**2
*x + 8*x**n*(x**2*c + a)**p*a*e**3*n**3*p**3*x + 8*x**n*(x**2*c + a)
**p*a*e**3*n**3*p**2*x + 8*x**n*(x**2*c + a)**p*a*e**3*n**2*p**2*x +
4*x**n*(x**2*c + a)**p*a*e**3*n**2*p*x + 2*x**n*(x**2*c + a)**p*a
*e**3*n*p*x + 24*x**n*(x**2*c + a)**p*c*d**2*e*n**3*p**3*x + 60*x**n*(x
**2*c + a)**p*c*d**2*e*n**3*p**2*x + 36*x**n*(x**2*c + a)**p*c*d*
2*e*n**3*p*x + 36*x**n*(x**2*c + a)**p*c*d**2*e*n**2*p**2*x + 60*x**n
*(x**2*c + a)**p*c*d**2*e*n**2*p*x + 18*x**n*(x**2*c + a)**p*c*d**
2*e*n**2*x + 18*x**n*(x**2*c + a)**p*c*d**2*e*n*p*x + 15*x**n*(x**...
```

3.48 $\int (d + ex^n)^2 (a + cx^{2n})^p dx$

Optimal result	345
Mathematica [A] (verified)	346
Rubi [A] (verified)	346
Maple [F]	347
Fricas [F]	348
Sympy [F(-1)]	348
Maxima [F]	348
Giac [F(-2)]	349
Mupad [F(-1)]	349
Reduce [F]	349

Optimal result

Integrand size = 21, antiderivative size = 201

$$\begin{aligned} \int (d + ex^n)^2 (a + cx^{2n})^p dx &= \frac{e^2 x (a + cx^{2n})^{1+p}}{c(1 + 2n(1 + p))} \\ &- \frac{(ae^2 - cd^2(1 + 2n(1 + p))) x (a + cx^{2n})^p \left(1 + \frac{cx^{2n}}{a}\right)^{-p} \text{Hypergeometric2F1}\left(\frac{1}{2n}, -p, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{c(1 + 2n(1 + p))} \\ &+ \frac{2dex^{1+n}(a + cx^{2n})^p \left(1 + \frac{cx^{2n}}{a}\right)^{-p} \text{Hypergeometric2F1}\left(\frac{1+n}{2n}, -p, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{1 + n} \end{aligned}$$

output

```
e^2*x*(a+c*x^(2*n))^(p+1)/c/(1+2*n*(p+1))-(a*e^2-c*d^2*(1+2*n*(p+1)))*x*(a+c*x^(2*n))^p*hypergeom([-p, 1/2/n], [1+1/2/n], -c*x^(2*n)/a)/c/(1+2*n*(p+1))/((1+c*x^(2*n)/a)^p)+2*d*e*x^(1+n)*(a+c*x^(2*n))^p*hypergeom([-p, 1/2*(1+n)/n], [3/2+1/2/n], -c*x^(2*n)/a)/(1+n)/((1+c*x^(2*n)/a)^p)
```

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.85

$$\int (d + ex^n)^2 (a + cx^{2n})^p dx \\ = \frac{x(a + cx^{2n})^p \left(1 + \frac{cx^{2n}}{a}\right)^{-p} \left(e^2(1+n)x^{2n} \text{Hypergeometric2F1}\left(\frac{1}{2}(2+\frac{1}{n}), -p, \frac{1}{2}(4+\frac{1}{n}), -\frac{cx^{2n}}{a}\right) + d(1+n)x^{2n} \text{Hypergeometric2F1}\left(\frac{1}{2}(2+\frac{1}{n}), -p-1, \frac{1}{2}(4+\frac{1}{n}), -\frac{cx^{2n}}{a}\right)\right)}{x(a + cx^{2n})^p \left(1 + \frac{cx^{2n}}{a}\right)^{-p}}$$

input `Integrate[(d + e*x^n)^2*(a + c*x^(2*n))^p, x]`

output
$$(x*(a + c*x^{2n})^p*(e^{2*(1 + n)}*x^{(2*n)}*\text{Hypergeometric2F1}[(2 + n^{(-1)})/2, -p, (4 + n^{(-1)})/2, -((c*x^{(2*n)})/a)] + d*(1 + 2*n)*(d*(1 + n)*\text{Hypergeometric2F1}[1/(2*n), -p, (2 + n^{(-1)})/2, -((c*x^{(2*n)})/a)] + 2*e*x^{n}*Hyp\text{ypergeometric2F1}[(1 + n)/(2*n), -p, (3 + n^{(-1)})/2, -((c*x^{(2*n)})/a)]))/((1 + n)*(1 + 2*n)*(1 + (c*x^{(2*n)})/a)^p)}$$

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.08, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {1767, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex^n)^2 (a + cx^{2n})^p dx \\ \downarrow 1767 \\ \int (d^2(a + cx^{2n})^p + 2dex^n(a + cx^{2n})^p + e^2x^{2n}(a + cx^{2n})^p) dx \\ \downarrow 2009$$

$$\frac{d^2x(a+cx^{2n})^p \left(\frac{cx^{2n}}{a} + 1\right)^{-p} \text{Hypergeometric2F1}\left(\frac{1}{2n}, -p, \frac{1}{2}\left(2 + \frac{1}{n}\right), -\frac{cx^{2n}}{a}\right) + 2dex^{n+1}(a+cx^{2n})^p \left(\frac{cx^{2n}}{a} + 1\right)^{-p} \text{Hypergeometric2F1}\left(\frac{n+1}{2n}, -p, \frac{1}{2}\left(3 + \frac{1}{n}\right), -\frac{cx^{2n}}{a}\right)}{n+1} + \frac{e^2x^{2n+1}(a+cx^{2n})^p \left(\frac{cx^{2n}}{a} + 1\right)^{-p} \text{Hypergeometric2F1}\left(\frac{1}{2}\left(2 + \frac{1}{n}\right), -p, \frac{1}{2}\left(4 + \frac{1}{n}\right), -\frac{cx^{2n}}{a}\right)}{2n+1}$$

input `Int[(d + e*x^n)^2*(a + c*x^(2*n))^p, x]`

output $(e^{2*x^{(1 + 2*n)*(a + c*x^{(2*n)})^p}*\text{Hypergeometric2F1}[(2 + n^{(-1)})/2, -p, (4 + n^{(-1)})/2, -((c*x^{(2*n)})/a)]}/((1 + 2*n)*(1 + (c*x^{(2*n)})/a)^p) + (d^2*x*(a + c*x^{(2*n)})^p*\text{Hypergeometric2F1}[1/(2*n), -p, (2 + n^{(-1)})/2, -((c*x^{(2*n)})/a)]}/(1 + (c*x^{(2*n)})/a)^p + (2*d*e*x^{(1 + n)*(a + c*x^{(2*n)})^p}*\text{Hypergeometric2F1}[(1 + n)/(2*n), -p, (3 + n^{(-1)})/2, -((c*x^{(2*n)})/a)]}/((1 + n)*(1 + (c*x^{(2*n)})/a)^p))$

Definitions of rubi rules used

rule 1767 `Int[((d_) + (e_)*(x_)^(n_))^(q_)*((a_) + (c_)*(x_)^(n2_))^(p_), x_Symbol] :> Int[ExpandIntegrand[(d + e*x^n)^q*(a + c*x^(2*n))^p, x], x] /; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && NeQ[c*d^2 + a*e^2, 0] && ((IntegersQ[p, q] && !IntegerQ[n]) || IGtQ[p, 0] || (IGtQ[q, 0] && !IntegerQ[n]))`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [F]

$$\int (d + e x^n)^2 (a + c x^{2n})^p dx$$

input `int((d+e*x^n)^2*(a+c*x^(2*n))^p, x)`

output `int((d+e*x^n)^2*(a+c*x^(2*n))^p,x)`

Fricas [F]

$$\int (d + ex^n)^2 (a + cx^{2n})^p \, dx = \int (ex^n + d)^2 (cx^{2n} + a)^p \, dx$$

input `integrate((d+e*x^n)^2*(a+c*x^(2*n))^p,x, algorithm="fricas")`

output `integral((e^2*x^(2*n) + 2*d*e*x^n + d^2)*(c*x^(2*n) + a)^p, x)`

Sympy [F(-1)]

Timed out.

$$\int (d + ex^n)^2 (a + cx^{2n})^p \, dx = \text{Timed out}$$

input `integrate((d+e*x**n)**2*(a+c*x**2*n)**p,x)`

output `Timed out`

Maxima [F]

$$\int (d + ex^n)^2 (a + cx^{2n})^p \, dx = \int (ex^n + d)^2 (cx^{2n} + a)^p \, dx$$

input `integrate((d+e*x^n)^2*(a+c*x^(2*n))^p,x, algorithm="maxima")`

output `integrate((e*x^n + d)^2*(c*x^(2*n) + a)^p, x)`

Giac [F(-2)]

Exception generated.

$$\int (d + ex^n)^2 (a + cx^{2n})^p dx = \text{Exception raised: TypeError}$$

input `integrate((d+e*x^n)^2*(a+c*x^(2*n))^p,x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:Unable to divide, perhaps due to ro
unding error%{64,[1,0,4,3,1,4,3,1,1]}%+%{256,[1,0,4,3,1,3,3,1,1]}%
+%{384,

Mupad [F(-1)]

Timed out.

$$\int (d + ex^n)^2 (a + cx^{2n})^p dx = \int (a + cx^{2n})^p (d + ex^n)^2 dx$$

input `int((a + c*x^(2*n))^p*(d + e*x^n)^2,x)`

output `int((a + c*x^(2*n))^p*(d + e*x^n)^2, x)`

Reduce [F]

$$\int (d + ex^n)^2 (a + cx^{2n})^p dx = \text{too large to display}$$

input `int((d+e*x^n)^2*(a+c*x^(2*n))^p,x)`

output

```
(4*x**2*n)*(x**2*n*c + a)**p*c*e**2*n**2*p**2*x + 2*x**2*(2*n)*(x**2*n*c + a)**p*c*e**2*n**2*p*x + 4*x**2*(2*n)*(x**2*n*c + a)**p*c*e**2*n*p*x + x**2*(2*n)*(x**2*n*c + a)**p*c*e**2*n*x + x**2*(2*n)*(x**2*n*c + a)**p*c*e**2*x + 8*x**n*(x**2*n*c + a)**p*c*d*e*n**2*p**2*x + 8*x**n*(x**2*n*c + a)**p*c*d*e*n*p*x + 4*x**n*(x**2*n*c + a)**p*c*d*e*n*x + 2*x**n*(x**2*n*c + a)**p*c*d*e*x + 4*(x**2*n*c + a)**p*a*e**2*n**2*p**2*x + 2*(x**2*n*c + a)**p*a*e**2*n**2*p*x + 2*(x**2*n*c + a)**p*a*e**2*n*p*x + 4*(x**2*n*c + a)**p*c*d**2*n**2*p**2*x + 6*(x**2*n*c + a)**p*c*d**2*n**2*p*x + 2*(x**2*n*c + a)**p*c*d**2*n*p*x + 3*(x**2*n*c + a)**p*c*d**2*n*x + (x**2*n*c + a)**p*c*d**2*x - 32*int((x**2*n*c + a)**p/(8*x**2*n*c*n**3*p**3 + 12*x**2*(2*n)*c*n**3*p**2 + 4*x**2*(2*n)*c*n**3*p + 12*x**2*(2*n)*c*n**2*p**2 + 12*x**2*(2*n)*c*n**2*p + 2*x**2*(2*n)*c*n**2 + 6*x**2*(2*n)*c*n*p + 3*x**2*(2*n)*c*n + x**2*(2*n)*c + 8*a*n**3*p**3 + 12*a*n**3*p**2 + 4*a*n**3*p + 12*a*n**2*p**2 + 12*a*n**2*p + 2*a*n**2 + 6*a*n*p + 3*a*n + a), x)*a**2*e**2*n**5*p**5 - 64*int((x**2*n*c + a)**p/(8*x**2*(2*n)*c*n**3*p**3 + 12*x**2*(2*n)*c*n**3*p**2 + 4*x**2*(2*n)*c*n**2*p**2 + 12*x**2*(2*n)*c*n**2*p + 2*x**2*(2*n)*c*n**2 + 6*x**2*(2*n)*c*n*p + 3*x**2*(2*n)*c*n + x**2*(2*n)*c + 8*a*n**3*p**3 + 12*a*n**3*p**2 + 4*a*n**3*p + 12*a*n**2*p**2 + 12*a*n**2*p + 2*a*n**2 + 6*a*n*p + 3*a*n + a), x)*a**2*e**2*n...
```

3.49 $\int (d + ex^n) (a + cx^{2n})^p dx$

Optimal result	351
Mathematica [A] (verified)	351
Rubi [A] (verified)	352
Maple [F]	353
Fricas [F]	353
Sympy [C] (verification not implemented)	354
Maxima [F]	354
Giac [F]	355
Mupad [F(-1)]	355
Reduce [F]	355

Optimal result

Integrand size = 19, antiderivative size = 135

$$\begin{aligned} & \int (d + ex^n) (a + cx^{2n})^p dx \\ &= dx (a + cx^{2n})^p \left(1 + \frac{cx^{2n}}{a}\right)^{-p} \text{Hypergeometric2F1} \left(\frac{1}{2n}, -p, \frac{1}{2} \left(2 + \frac{1}{n}\right), -\frac{cx^{2n}}{a}\right) \\ &+ \frac{ex^{1+n} (a + cx^{2n})^p \left(1 + \frac{cx^{2n}}{a}\right)^{-p} \text{Hypergeometric2F1} \left(\frac{1+n}{2n}, -p, \frac{1}{2} \left(3 + \frac{1}{n}\right), -\frac{cx^{2n}}{a}\right)}{1+n} \end{aligned}$$

output $d*x*(a+c*x^(2*n))^p*hypergeom([-p, 1/2/n], [1+1/2/n], -c*x^(2*n)/a)/((1+c*x^(2*n)/a)^p)+e*x^(1+n)*(a+c*x^(2*n))^p*hypergeom([-p, 1/2*(1+n)/n], [3/2+1/2/n], -c*x^(2*n)/a)/(1+n)/((1+c*x^(2*n)/a)^p)$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.81

$$\begin{aligned} & \int (d + ex^n) (a + cx^{2n})^p dx \\ &= \frac{x(a + cx^{2n})^p \left(1 + \frac{cx^{2n}}{a}\right)^{-p} \left(d(1 + n) \text{Hypergeometric2F1} \left(\frac{1}{2n}, -p, \frac{1}{2} \left(2 + \frac{1}{n}\right), -\frac{cx^{2n}}{a}\right) + ex^n \text{Hypergeometric2F1} \left(\frac{1+n}{2n}, -p, \frac{1}{2} \left(3 + \frac{1}{n}\right), -\frac{cx^{2n}}{a}\right)\right)}{1+n} \end{aligned}$$

input `Integrate[(d + e*x^n)*(a + c*x^(2*n))^p, x]`

output
$$\frac{(x*(a + c*x^{(2*n)})^p*(d*(1 + n)*Hypergeometric2F1[1/(2*n), -p, (2 + n^{(-1)})/2, -((c*x^{(2*n)})/a)] + e*x^{n*Hypergeometric2F1[(1 + n)/(2*n), -p, (3 + n^{(-1)})/2, -((c*x^{(2*n)})/a)]}))/((1 + n)*(1 + (c*x^{(2*n)})/a)^p)}{(1 + n)*(1 + (c*x^{(2*n)})/a)^p}$$

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {1763, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (d + ex^n) (a + cx^{2n})^p dx \\ & \quad \downarrow 1763 \\ & \int (d(a + cx^{2n})^p + ex^n(a + cx^{2n})^p) dx \\ & \quad \downarrow 2009 \\ & \frac{dx(a + cx^{2n})^p \left(\frac{cx^{2n}}{a} + 1\right)^{-p} \text{Hypergeometric2F1}\left(\frac{1}{2n}, -p, \frac{1}{2}\left(2 + \frac{1}{n}\right), -\frac{cx^{2n}}{a}\right) + ex^{n+1}(a + cx^{2n})^p \left(\frac{cx^{2n}}{a} + 1\right)^{-p} \text{Hypergeometric2F1}\left(\frac{n+1}{2n}, -p, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}\right)}{n + 1} \end{aligned}$$

input `Int[(d + e*x^n)*(a + c*x^(2*n))^p, x]`

output
$$\frac{(d*x*(a + c*x^{(2*n)})^p*Hypergeometric2F1[1/(2*n), -p, (2 + n^{(-1)})/2, -((c*x^{(2*n)})/a)])/(1 + (c*x^{(2*n)})/a)^p + (e*x^{(1 + n)}*(a + c*x^{(2*n)})^p*Hypergeometric2F1[(1 + n)/(2*n), -p, (3 + n^{(-1)})/2, -((c*x^{(2*n)})/a)])/(1 + n)*(1 + (c*x^{(2*n)})/a)^p}{(1 + n)*(1 + (c*x^{(2*n)})/a)^p}$$

Definitions of rubi rules used

rule 1763 $\text{Int}[(d_+ + e_*)*(x_*)^{(n_*)}*((a_+ + c_*)*(x_*)^{(n2_*)})^p, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(d + e*x^n)*(a + c*x^{(2*n)})^p, x], x] /; \text{FreeQ}[\{a, c, d, e, n\}, x] \&& \text{EqQ}[n2, 2*n]$

rule 2009 $\text{Int}[u_*, x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

Maple [F]

$$\int (d + e x^n) (a + c x^{2n})^p dx$$

input $\text{int}((d+e*x^n)*(a+c*x^{(2*n)})^p, x)$

output $\text{int}((d+e*x^n)*(a+c*x^{(2*n)})^p, x)$

Fricas [F]

$$\int (d + e x^n) (a + c x^{2n})^p dx = \int (e x^n + d) (c x^{2n} + a)^p dx$$

input $\text{integrate}((d+e*x^n)*(a+c*x^{(2*n)})^p, x, \text{algorithm}=\text{"fricas"})$

output $\text{integral}((e*x^n + d)*(c*x^{(2*n)} + a)^p, x)$

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 138.22 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.07

$$\begin{aligned} & \int (d + ex^n) (a + cx^{2n})^p dx \\ &= \frac{a^{\frac{1}{2n}} a^{p - \frac{1}{2n}} dx \Gamma\left(\frac{1}{2n}\right) {}_2F_1\left(\begin{matrix} \frac{1}{2n}, -p \\ 1 + \frac{1}{2n} \end{matrix} \middle| \frac{cx^{2n} e^{i\pi}}{a}\right)}{2n \Gamma\left(1 + \frac{1}{2n}\right)} \\ &+ \frac{a^{\frac{1}{2} + \frac{1}{2n}} a^{p - \frac{1}{2} - \frac{1}{2n}} ex^{n+1} \Gamma\left(\frac{1}{2} + \frac{1}{2n}\right) {}_2F_1\left(\begin{matrix} -p, \frac{1}{2} + \frac{1}{2n} \\ \frac{3}{2} + \frac{1}{2n} \end{matrix} \middle| \frac{cx^{2n} e^{i\pi}}{a}\right)}{2n \Gamma\left(\frac{3}{2} + \frac{1}{2n}\right)} \end{aligned}$$

input `integrate((d+e*x**n)*(a+c*x**(2*n))**p,x)`

output `a**(1/(2*n))*a**(-p - 1/(2*n))*d*x*gamma(1/(2*n))*hyper((1/(2*n), -p), (1 + 1/(2*n),), c*x**(2*n)*exp_polar(I*pi)/a)/(2*n*gamma(1 + 1/(2*n))) + a**(-1/2 + 1/(2*n))*a**(-p - 1/2 - 1/(2*n))*e*x**(-n + 1)*gamma(1/2 + 1/(2*n))*hyper((-p, 1/2 + 1/(2*n)), (3/2 + 1/(2*n),), c*x**(2*n)*exp_polar(I*pi)/a)/(2*n*gamma(3/2 + 1/(2*n)))`

Maxima [F]

$$\int (d + ex^n) (a + cx^{2n})^p dx = \int (ex^n + d)(cx^{2n} + a)^p dx$$

input `integrate((d+e*x^n)*(a+c*x^(2*n))^p,x, algorithm="maxima")`

output `integrate((e*x^n + d)*(c*x^(2*n) + a)^p, x)`

Giac [F]

$$\int (d + ex^n) (a + cx^{2n})^p dx = \int (ex^n + d)(cx^{2n} + a)^p dx$$

input `integrate((d+e*x^n)*(a+c*x^(2*n))^p,x, algorithm="giac")`

output `integrate((e*x^n + d)*(c*x^(2*n) + a)^p, x)`

Mupad [F(-1)]

Timed out.

$$\int (d + ex^n) (a + cx^{2n})^p dx = \int (a + cx^{2n})^p (d + ex^n) dx$$

input `int((a + c*x^(2*n))^p*(d + e*x^n),x)`

output `int((a + c*x^(2*n))^p*(d + e*x^n), x)`

Reduce [F]

$$\int (d + ex^n) (a + cx^{2n})^p dx = \text{too large to display}$$

input `int((d+e*x^n)*(a+c*x^(2*n))^p,x)`

output

```
(2*x**n*(x**(2*n)*c + a)**p*e*n*p*x + x**n*(x**(2*n)*c + a)**p*e*x + 2*(x*(2*n)*c + a)**p*d*n*p*x + (x**(2*n)*c + a)**p*d*n*x + (x**(2*n)*c + a)**p*d*x + 16*int((x**(2*n)*c + a)**p/(4*x**(2*n)*c*n**2*p**2 + 2*x**(2*n)*c*n**2*p + 4*x**(2*n)*c*n*p + x**(2*n)*c*n + x**(2*n)*c + 4*a*n**2*p**2 + 2*a*n**2*p + 4*a*n*p + a*n + a),x)*a*d*n**4*p**4 + 16*int((x**(2*n)*c + a)**p/(4*x**(2*n)*c*n**2*p**2 + 2*x**(2*n)*c*n**2*p + 4*x**(2*n)*c*n*p + x**(2*n)*c*n + x**(2*n)*c + 4*a*n**2*p**2 + 2*a*n**2*p + 4*a*n*p + a*n + a),x)*a*d*n**4*p**3 + 4*int((x**(2*n)*c + a)**p/(4*x**(2*n)*c*n**2*p**2 + 2*x**(2*n)*c*n**2*p + 4*x**(2*n)*c*n*p + x**(2*n)*c*n + x**(2*n)*c + 4*a*n**2*p**2 + 2*a*n**2*p + 4*a*n*p + a*n + a),x)*a*d*n**4*p**2 + 24*int((x**(2*n)*c + a)**p/(4*x**(2*n)*c*n**2*p**2 + 2*x**(2*n)*c*n**2*p + 4*x**(2*n)*c*n*p + x**(2*n)*c*n + x**(2*n)*c + 4*a*n**2*p**2 + 2*a*n**2*p + 4*a*n*p + a*n + a),x)*a*d*n**3*p**3 + 16*int((x**(2*n)*c + a)**p/(4*x**(2*n)*c*n**2*p**2 + 2*x**(2*n)*c*n**2*p + 4*x**(2*n)*c*n*p + x**(2*n)*c*n + x**(2*n)*c + 4*a*n**2*p**2 + 2*a*n**2*p + 4*a*n*p + a*n + a),x)*a*d*n**3*p**2 + 2*int((x**(2*n)*c + a)**p/(4*x**(2*n)*c*n**2*p**2 + 2*x**(2*n)*c*n**2*p + 4*x**(2*n)*c*n*p + x**(2*n)*c*n + x**(2*n)*c + 4*a*n**2*p**2 + 2*a*n**2*p + 4*a*n*p + a*n + a),x)*a*d*n**3*p + 12*int((x**(2*n)*c + a)**p/(4*x**(2*n)*c*n**2*p**2 + 2*a*n**2*p + 4*a*n*p + a*n + a),x)*a*d*n**2*p**2 + 4*a*n**2*p**2 + 2*a*n**2*p + 4*a*n*p + a*n + a),x)*a*d*n**2*p**2 + 4*in...
```

$$3.50 \quad \int \frac{(a+cx^{2n})^p}{d+ex^n} dx$$

Optimal result	357
Mathematica [F]	358
Rubi [A] (verified)	358
Maple [F]	359
Fricas [F]	359
Sympy [F(-2)]	360
Maxima [F]	360
Giac [F]	360
Mupad [F(-1)]	361
Reduce [F]	361

Optimal result

Integrand size = 21, antiderivative size = 167

$$\begin{aligned} & \int \frac{(a+cx^{2n})^p}{d+ex^n} dx \\ &= \frac{x(a+cx^{2n})^p \left(1 + \frac{cx^{2n}}{a}\right)^{-p} \text{AppellF1} \left(\frac{1}{2n}, -p, 1, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2 x^{2n}}{d^2}\right)}{d} \\ &\quad - \frac{ex^{1+n}(a+cx^{2n})^p \left(1 + \frac{cx^{2n}}{a}\right)^{-p} \text{AppellF1} \left(\frac{1+n}{2n}, -p, 1, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2 x^{2n}}{d^2}\right)}{d^2(1+n)} \end{aligned}$$

output

```
x*(a+c*x^(2*n))^p*AppellF1(1/2/n,1,-p,1+1/2/n,e^2*x^(2*n)/d^2,-c*x^(2*n)/a)/d/((1+c*x^(2*n)/a)^p)-e*x^(1+n)*(a+c*x^(2*n))^p*AppellF1(1/2*(1+n)/n,1,-p,3/2+1/2/n,e^2*x^(2*n)/d^2,-c*x^(2*n)/a)/d^2/(1+n)/((1+c*x^(2*n)/a)^p)
```

Mathematica [F]

$$\int \frac{(a + cx^{2n})^p}{d + ex^n} dx = \int \frac{(a + cx^{2n})^p}{d + ex^n} dx$$

input `Integrate[(a + c*x^(2*n))^p/(d + e*x^n), x]`

output `Integrate[(a + c*x^(2*n))^p/(d + e*x^n), x]`

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {1768, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + cx^{2n})^p}{d + ex^n} dx \\ & \quad \downarrow 1768 \\ & \int \left(\frac{d(a + cx^{2n})^p}{d^2 - e^2 x^{2n}} + \frac{ex^n (a + cx^{2n})^p}{e^2 x^{2n} - d^2} \right) dx \\ & \quad \downarrow 2009 \\ & \frac{x (a + cx^{2n})^p \left(\frac{cx^{2n}}{a} + 1 \right)^{-p} \text{AppellF1} \left(\frac{1}{2n}, -p, 1, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2 x^{2n}}{d^2} \right)}{d} - \\ & \frac{ex^{n+1} (a + cx^{2n})^p \left(\frac{cx^{2n}}{a} + 1 \right)^{-p} \text{AppellF1} \left(\frac{n+1}{2n}, -p, 1, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2 x^{2n}}{d^2} \right)}{d^2(n+1)} \end{aligned}$$

input `Int[(a + c*x^(2*n))^p/(d + e*x^n), x]`

output
$$(x*(a + c*x^{(2*n)})^p * \text{AppellF1}[1/(2*n), -p, 1, (2 + n^{(-1)})/2, -((c*x^{(2*n)})/a), (e^{2*x^{(2*n)}}/d^2)/(d*(1 + (c*x^{(2*n)})/a)^p) - (e*x^{(1+n)}*(a + c*x^{(2*n)})^p * \text{AppellF1}[(1+n)/(2*n), -p, 1, (3 + n^{(-1)})/2, -((c*x^{(2*n)})/a), (e^{2*x^{(2*n)}}/d^2)]/(d^2*(1 + (c*x^{(2*n)})/a)^p)$$

Defintions of rubi rules used

rule 1768
$$\text{Int}[(d_0 + e_0*x^{n_0})^{q_0}*((a_0 + c_0*x^{(2*n)})^p, x_{\text{Symbol}}) :> \text{Int}[\text{ExpandIntegrand}[(a + c*x^{(2*n)})^p, (d/(d^2 - e^{2*x^{(2*n)}}) - e*(x^n/(d^2 - e^{2*x^{(2*n)}})))^{(-q)}, x], x] /; \text{FreeQ}[\{a, c, d, e, n, p\}, x] \&& \text{EqQ}[n_0, 2*n] \&& \text{NeQ}[c*d^2 + a*e^2, 0] \&& !\text{IntegerQ}[p] \&& \text{ILtQ}[q, 0]$$

rule 2009
$$\text{Int}[u_, x_{\text{Symbol}}] :> \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$$

Maple [F]

$$\int \frac{(a + cx^{2n})^p}{d + ex^n} dx$$

input $\text{int}((a+c*x^{(2*n)})^p/(d+e*x^{n}), x)$

output $\text{int}((a+c*x^{(2*n)})^p/(d+e*x^{n}), x)$

Fricas [F]

$$\int \frac{(a + cx^{2n})^p}{d + ex^n} dx = \int \frac{(cx^{2n} + a)^p}{ex^n + d} dx$$

input $\text{integrate}((a+c*x^{(2*n)})^p/(d+e*x^{n}), x, \text{algorithm}=\text{"fricas"})$

output $\text{integral}((c*x^{(2*n)} + a)^p/(e*x^{n} + d), x)$

Sympy [F(-2)]

Exception generated.

$$\int \frac{(a + cx^{2n})^p}{d + ex^n} dx = \text{Exception raised: HeuristicGCDFailed}$$

input `integrate((a+c*x**(2*n))**p/(d+e*x**n),x)`

output `Exception raised: HeuristicGCDFailed >> no luck`

Maxima [F]

$$\int \frac{(a + cx^{2n})^p}{d + ex^n} dx = \int \frac{(cx^{2n} + a)^p}{ex^n + d} dx$$

input `integrate((a+c*x^(2*n))^p/(d+e*x^n),x, algorithm="maxima")`

output `integrate((c*x^(2*n) + a)^p/(e*x^n + d), x)`

Giac [F]

$$\int \frac{(a + cx^{2n})^p}{d + ex^n} dx = \int \frac{(cx^{2n} + a)^p}{ex^n + d} dx$$

input `integrate((a+c*x^(2*n))^p/(d+e*x^n),x, algorithm="giac")`

output `integrate((c*x^(2*n) + a)^p/(e*x^n + d), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + cx^{2n})^p}{d + ex^n} dx = \int \frac{(a + cx^{2n})^p}{d + e x^n} dx$$

input `int((a + c*x^(2*n))^p/(d + e*x^n),x)`

output `int((a + c*x^(2*n))^p/(d + e*x^n), x)`

Reduce [F]

$$\int \frac{(a + cx^{2n})^p}{d + ex^n} dx = \int \frac{(x^{2n}c + a)^p}{x^n e + d} dx$$

input `int((a+c*x^(2*n))^p/(d+e*x^n),x)`

output `int((x^(2*n)*c + a)^p/(x^n*e + d),x)`

3.51 $\int \frac{(a+cx^{2n})^p}{(d+ex^n)^2} dx$

Optimal result	362
Mathematica [F]	363
Rubi [A] (verified)	363
Maple [F]	364
Fricas [F]	364
Sympy [F(-1)]	365
Maxima [F]	365
Giac [F]	365
Mupad [F(-1)]	366
Reduce [F]	366

Optimal result

Integrand size = 21, antiderivative size = 261

$$\begin{aligned} & \int \frac{(a+cx^{2n})^p}{(d+ex^n)^2} dx \\ &= \frac{e^2 x^{1+2n} (a+cx^{2n})^p \left(1 + \frac{cx^{2n}}{a}\right)^{-p} \text{AppellF1}\left(\frac{1}{2}(2 + \frac{1}{n}), -p, 2, \frac{1}{2}(4 + \frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2 x^{2n}}{d^2}\right)}{d^4(1+2n)} \\ &+ \frac{x(a+cx^{2n})^p \left(1 + \frac{cx^{2n}}{a}\right)^{-p} \text{AppellF1}\left(\frac{1}{2n}, -p, 2, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2 x^{2n}}{d^2}\right)}{d^2} \\ &- \frac{2ex^{1+n} (a+cx^{2n})^p \left(1 + \frac{cx^{2n}}{a}\right)^{-p} \text{AppellF1}\left(\frac{1+n}{2n}, -p, 2, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2 x^{2n}}{d^2}\right)}{d^3(1+n)} \end{aligned}$$

output $e^{2*x^(1+2*n)*(a+c*x^(2*n))^p}*\text{AppellF1}(1+1/2/n, 2, -p, 2+1/2/n, e^{2*x^(2*n)}/d^2, -c*x^(2*n)/a)/d^4/(1+2*n)/((1+c*x^(2*n)/a)^p)+x*(a+c*x^(2*n))^p*\text{AppellF1}(1/2/n, 2, -p, 1+1/2/n, e^{2*x^(2*n)}/d^2, -c*x^(2*n)/a)/d^2/((1+c*x^(2*n)/a)^p)-2*e*x^(1+n)*(a+c*x^(2*n))^p*\text{AppellF1}(1/2*(1+n)/n, 2, -p, 3/2+1/2/n, e^{2*x^(2*n)}/d^2, -c*x^(2*n)/a)/d^3/(1+n)/((1+c*x^(2*n)/a)^p)$

Mathematica [F]

$$\int \frac{(a + cx^{2n})^p}{(d + ex^n)^2} dx = \int \frac{(a + cx^{2n})^p}{(d + ex^n)^2} dx$$

input `Integrate[(a + c*x^(2*n))^p/(d + e*x^n)^2, x]`

output `Integrate[(a + c*x^(2*n))^p/(d + e*x^n)^2, x]`

Rubi [A] (verified)

Time = 0.47 (sec), antiderivative size = 261, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {1768, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + cx^{2n})^p}{(d + ex^n)^2} dx \\ & \quad \downarrow 1768 \\ & \int \left(\frac{e^2 x^{2n} (a + cx^{2n})^p}{(e^2 x^{2n} - d^2)^2} + \frac{d^2 (a + cx^{2n})^p}{(d^2 - e^2 x^{2n})^2} - \frac{2d e x^n (a + cx^{2n})^p}{(e^2 x^{2n} - d^2)^2} \right) dx \\ & \quad \downarrow 2009 \\ & \frac{x (a + cx^{2n})^p \left(\frac{cx^{2n}}{a} + 1 \right)^{-p} \text{AppellF1} \left(\frac{1}{2n}, -p, 2, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2 x^{2n}}{d^2} \right)}{d^2} + \\ & \frac{e^2 x^{2n+1} (a + cx^{2n})^p \left(\frac{cx^{2n}}{a} + 1 \right)^{-p} \text{AppellF1} \left(\frac{1}{2}(2 + \frac{1}{n}), -p, 2, \frac{1}{2}(4 + \frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2 x^{2n}}{d^2} \right)}{d^4(2n+1)} - \\ & \frac{2e x^{n+1} (a + cx^{2n})^p \left(\frac{cx^{2n}}{a} + 1 \right)^{-p} \text{AppellF1} \left(\frac{n+1}{2n}, -p, 2, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2 x^{2n}}{d^2} \right)}{d^3(n+1)} \end{aligned}$$

input `Int[(a + c*x^(2*n))^p/(d + e*x^n)^2, x]`

output

$$(e^{2x}(1 + 2n)(a + cx^{2n})^p) \operatorname{AppellF1}[(2 + n(-1))/2, -p, 2, (4 + n(-1))/2, -((c*x^{2n})/a), (e^{2x}(2n)/d^2)/(d^4(1 + 2n)(1 + (c*x^{2n})/a)^p) + (x(a + cx^{2n})^p) \operatorname{AppellF1}[1/(2n), -p, 2, (2 + n(-1))/2, -((c*x^{2n})/a), (e^{2x}(2n)/d^2)/(d^2(1 + (c*x^{2n})/a)^p) - (2e^x(1 + n)(a + cx^{2n})^p) \operatorname{AppellF1}[(1 + n)/(2n), -p, 2, (3 + n(-1))/2, -((c*x^{2n})/a), (e^{2x}(2n)/d^2)/(d^3(1 + n)(1 + (c*x^{2n})/a)^p)]$$

Definitions of rubi rules used

rule 1768

```
Int[((d_) + (e_)*(x_)^(n_))^(q_)*((a_) + (c_)*(x_)^(n2_))^(p_), x_Symbol]
  :> Int[ExpandIntegrand[(a + c*x^{2n})^p, (d/(d^2 - e^{2x}(2n)) - e*(x^n/(d^2 - e^{2x}(2n))))^{(-q)}, x], x] /; FreeQ[{a, c, d, e, n, p}, x] && EqQ[n, 2*n] && NeQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && ILtQ[q, 0]
```

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [F]

$$\int \frac{(a + cx^{2n})^p}{(d + ex^n)^2} dx$$

input

```
int((a+c*x^{2n})^p/(d+e*x^n)^2,x)
```

output

```
int((a+c*x^{2n})^p/(d+e*x^n)^2,x)
```

Fricas [F]

$$\int \frac{(a + cx^{2n})^p}{(d + ex^n)^2} dx = \int \frac{(cx^{2n} + a)^p}{(ex^n + d)^2} dx$$

input

```
integrate((a+c*x^{2n})^p/(d+e*x^n)^2,x, algorithm="fricas")
```

output `integral((c*x^(2*n) + a)^p/(e^2*x^(2*n) + 2*d*e*x^n + d^2), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + cx^{2n})^p}{(d + ex^n)^2} dx = \text{Timed out}$$

input `integrate((a+c*x**(2*n))**p/(d+e*x**n)**2,x)`

output `Timed out`

Maxima [F]

$$\int \frac{(a + cx^{2n})^p}{(d + ex^n)^2} dx = \int \frac{(cx^{2n} + a)^p}{(ex^n + d)^2} dx$$

input `integrate((a+c*x^(2*n))^p/(d+e*x^n)^2,x, algorithm="maxima")`

output `integrate((c*x^(2*n) + a)^p/(e*x^n + d)^2, x)`

Giac [F]

$$\int \frac{(a + cx^{2n})^p}{(d + ex^n)^2} dx = \int \frac{(cx^{2n} + a)^p}{(ex^n + d)^2} dx$$

input `integrate((a+c*x^(2*n))^p/(d+e*x^n)^2,x, algorithm="giac")`

output `integrate((c*x^(2*n) + a)^p/(e*x^n + d)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + cx^{2n})^p}{(d + ex^n)^2} dx = \int \frac{(a + cx^{2n})^p}{(d + ex^n)^2} dx$$

input `int((a + c*x^(2*n))^p/(d + e*x^n)^2,x)`

output `int((a + c*x^(2*n))^p/(d + e*x^n)^2, x)`

Reduce [F]

$$\int \frac{(a + cx^{2n})^p}{(d + ex^n)^2} dx = \int \frac{(x^{2n}c + a)^p}{x^{2n}e^2 + 2x^n de + d^2} dx$$

input `int((a+c*x^(2*n))^p/(d+e*x^n)^2,x)`

output `int((x**(2*n)*c + a)**p/(x**(2*n)*e**2 + 2*x**n*d*e + d**2),x)`

3.52 $\int \frac{(a+cx^{2n})^p}{(d+ex^n)^3} dx$

Optimal result	367
Mathematica [F]	368
Rubi [A] (verified)	368
Maple [F]	369
Fricas [F]	370
Sympy [F(-1)]	370
Maxima [F]	370
Giac [F]	371
Mupad [F(-1)]	371
Reduce [F]	371

Optimal result

Integrand size = 21, antiderivative size = 357

$$\begin{aligned} & \int \frac{(a+cx^{2n})^p}{(d+ex^n)^3} dx \\ &= \frac{3e^2x^{1+2n}(a+cx^{2n})^p \left(1 + \frac{cx^{2n}}{a}\right)^{-p} \text{AppellF1}\left(\frac{1}{2}(2 + \frac{1}{n}), -p, 3, \frac{1}{2}(4 + \frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2x^{2n}}{d^2}\right)}{d^5(1+2n)} \\ &\quad - \frac{e^3x^{1+3n}(a+cx^{2n})^p \left(1 + \frac{cx^{2n}}{a}\right)^{-p} \text{AppellF1}\left(\frac{1}{2}(3 + \frac{1}{n}), -p, 3, \frac{1}{2}(5 + \frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2x^{2n}}{d^2}\right)}{d^6(1+3n)} \\ &\quad + \frac{x(a+cx^{2n})^p \left(1 + \frac{cx^{2n}}{a}\right)^{-p} \text{AppellF1}\left(\frac{1}{2n}, -p, 3, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2x^{2n}}{d^2}\right)}{d^3} \\ &\quad - \frac{3ex^{1+n}(a+cx^{2n})^p \left(1 + \frac{cx^{2n}}{a}\right)^{-p} \text{AppellF1}\left(\frac{1+n}{2n}, -p, 3, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2x^{2n}}{d^2}\right)}{d^4(1+n)} \end{aligned}$$

output

```
3*e^2*x^(1+2*n)*(a+c*x^(2*n))^p*AppellF1(1+1/2/n,3,-p,2+1/2/n,e^2*x^(2*n)/
d^2,-c*x^(2*n)/a)/d^5/(1+2*n)/((1+c*x^(2*n)/a)^p)-e^3*x^(1+3*n)*(a+c*x^(2*
n))^p*AppellF1(3/2+1/2/n,3,-p,5/2+1/2/n,e^2*x^(2*n)/d^2,-c*x^(2*n)/a)/d^6/
(1+3*n)/((1+c*x^(2*n)/a)^p)+x*(a+c*x^(2*n))^p*AppellF1(1/2/n,3,-p,1+1/2/n,
e^2*x^(2*n)/d^2,-c*x^(2*n)/a)/d^3/((1+c*x^(2*n)/a)^p)-3*e*x^(1+n)*(a+c*x^(2*n))^p*AppellF1(1/2*(1+n)/n,3,-p,3/2+1/2/n,e^2*x^(2*n)/d^2,-c*x^(2*n)/a)/
d^4/(1+n)/((1+c*x^(2*n)/a)^p)
```

Mathematica [F]

$$\int \frac{(a + cx^{2n})^p}{(d + ex^n)^3} dx = \int \frac{(a + cx^{2n})^p}{(d + ex^n)^3} dx$$

input `Integrate[(a + c*x^(2*n))^p/(d + e*x^n)^3, x]`

output `Integrate[(a + c*x^(2*n))^p/(d + e*x^n)^3, x]`

Rubi [A] (verified)

Time = 0.61 (sec) , antiderivative size = 357, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {1768, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + cx^{2n})^p}{(d + ex^n)^3} dx \\
 & \quad \downarrow \textcolor{blue}{1768} \\
 & \int \left(-\frac{3de^2x^{2n}(a + cx^{2n})^p}{(e^2x^{2n} - d^2)^3} + \frac{3d^2ex^n(a + cx^{2n})^p}{(e^2x^{2n} - d^2)^3} + \frac{e^3x^{3n}(a + cx^{2n})^p}{(e^2x^{2n} - d^2)^3} + \frac{d^3(a + cx^{2n})^p}{(d^2 - e^2x^{2n})^3} \right) dx \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & -\frac{e^3x^{3n+1}(a + cx^{2n})^p \left(\frac{cx^{2n}}{a} + 1 \right)^{-p} \text{AppellF1} \left(\frac{1}{2}(3 + \frac{1}{n}), -p, 3, \frac{1}{2}(5 + \frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2x^{2n}}{d^2} \right)}{d^6(3n+1)} + \\
 & \quad -\frac{3e^2x^{2n+1}(a + cx^{2n})^p \left(\frac{cx^{2n}}{a} + 1 \right)^{-p} \text{AppellF1} \left(\frac{1}{2}(2 + \frac{1}{n}), -p, 3, \frac{1}{2}(4 + \frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2x^{2n}}{d^2} \right)}{d^5(2n+1)} - \\
 & \quad \frac{3ex^{n+1}(a + cx^{2n})^p \left(\frac{cx^{2n}}{a} + 1 \right)^{-p} \text{AppellF1} \left(\frac{n+1}{2n}, -p, 3, \frac{1}{2}(3 + \frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2x^{2n}}{d^2} \right)}{d^4(n+1)} + \\
 & \quad \frac{x(a + cx^{2n})^p \left(\frac{cx^{2n}}{a} + 1 \right)^{-p} \text{AppellF1} \left(\frac{1}{2n}, -p, 3, \frac{1}{2}(2 + \frac{1}{n}), -\frac{cx^{2n}}{a}, \frac{e^2x^{2n}}{d^2} \right)}{d^3}
 \end{aligned}$$

input $\text{Int}[(a + c*x^{(2*n)})^p/(d + e*x^n)^3, x]$

output
$$\begin{aligned} & \frac{(3e^{2x}(1+2n)(a+c*x^{(2n)})^p \text{AppellF1}[(2+n^{(-1)})/2, -p, 3, (4+n^{(-1)})/2, -((c*x^{(2n)})/a), (e^{2x}(2n)/d^2)/(d^{5*(1+2n)}(1+(c*x^{(2n)})/a)^p) - (e^{3x}(1+3n)(a+c*x^{(2n)})^p \text{AppellF1}[(3+n^{(-1)})/2, -p, 3, (5+n^{(-1)})/2, -((c*x^{(2n)})/a), (e^{2x}(2n)/d^2)/(d^{6*(1+3n)}(1+(c*x^{(2n)})/a)^p) + (x*(a+c*x^{(2n)})^p \text{AppellF1}[1/(2n), -p, 3, (2+n^{(-1)})/2, -((c*x^{(2n)})/a), (e^{2x}(2n)/d^2)/(d^{3*(1+(c*x^{(2n)})/a)^p}) - (3e*x^{(1+n)}(a+c*x^{(2n)})^p \text{AppellF1}[(1+n)/(2n), -p, 3, (3+n^{(-1)})/2, -((c*x^{(2n)})/a), (e^{2x}(2n)/d^2)/(d^{4*(1+(c*x^{(2n)})/a)^p})]}}{ } \end{aligned}$$

Definitions of rubi rules used

rule 1768 $\text{Int}[(d_+ + e_-)(x_-)^{n_-})^{q_-}((a_- + c_-)(x_-)^{n_2_-})^{p_-}, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + c*x^{(2n)})^p, (d/(d^2 - e^{2x}(2n)) - e*(x^n/(d^2 - e^{2x}(2n))))^{-q}, x], x] /; \text{FreeQ}[\{a, c, d, e, n, p\}, x] \&& \text{EqQ}[n, 2n] \&& \text{NeQ}[c*d^2 + a*e^2, 0] \&& \text{!IntegerQ}[p] \&& \text{ILtQ}[q, 0]$

rule 2009 $\text{Int}[u_, x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

Maple [F]

$$\int \frac{(a + cx^{2n})^p}{(d + ex^n)^3} dx$$

input $\text{int}((a+c*x^{(2n)})^p/(d+e*x^n)^3, x)$

output $\text{int}((a+c*x^{(2n)})^p/(d+e*x^n)^3, x)$

Fricas [F]

$$\int \frac{(a + cx^{2n})^p}{(d + ex^n)^3} dx = \int \frac{(cx^{2n} + a)^p}{(ex^n + d)^3} dx$$

input `integrate((a+c*x^(2*n))^p/(d+e*x^n)^3,x, algorithm="fricas")`

output `integral((c*x^(2*n) + a)^p/(e^3*x^(3*n) + 3*d*e^2*x^(2*n) + 3*d^2*e*x^n + d^3), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + cx^{2n})^p}{(d + ex^n)^3} dx = \text{Timed out}$$

input `integrate((a+c*x**2*n)**p/(d+e*x**n)**3,x)`

output `Timed out`

Maxima [F]

$$\int \frac{(a + cx^{2n})^p}{(d + ex^n)^3} dx = \int \frac{(cx^{2n} + a)^p}{(ex^n + d)^3} dx$$

input `integrate((a+c*x^(2*n))^p/(d+e*x^n)^3,x, algorithm="maxima")`

output `integrate((c*x^(2*n) + a)^p/(e*x^n + d)^3, x)`

Giac [F]

$$\int \frac{(a + cx^{2n})^p}{(d + ex^n)^3} dx = \int \frac{(cx^{2n} + a)^p}{(ex^n + d)^3} dx$$

input `integrate((a+c*x^(2*n))^p/(d+e*x^n)^3,x, algorithm="giac")`

output `integrate((c*x^(2*n) + a)^p/(e*x^n + d)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + cx^{2n})^p}{(d + ex^n)^3} dx = \int \frac{(a + cx^{2n})^p}{(d + ex^n)^3} dx$$

input `int((a + c*x^(2*n))^p/(d + e*x^n)^3,x)`

output `int((a + c*x^(2*n))^p/(d + e*x^n)^3, x)`

Reduce [F]

$$\int \frac{(a + cx^{2n})^p}{(d + ex^n)^3} dx = \int \frac{(x^{2n}c + a)^p}{x^{3n}e^3 + 3x^{2n}de^2 + 3x^nd^2e + d^3} dx$$

input `int((a+c*x^(2*n))^p/(d+e*x^n)^3,x)`

output `int((x^(2*n)*c + a)**p/(x^(3*n)*e**3 + 3*x^(2*n)*d*e**2 + 3*x**n*d**2*e + d**3),x)`

3.53 $\int (d + ex^n)^q (a + cx^{2n})^p dx$

Optimal result	372
Mathematica [N/A]	372
Rubi [N/A]	373
Maple [N/A]	373
Fricas [N/A]	374
Sympy [F(-1)]	374
Maxima [N/A]	374
Giac [N/A]	375
Mupad [N/A]	375
Reduce [N/A]	376

Optimal result

Integrand size = 21, antiderivative size = 21

$$\int (d + ex^n)^q (a + cx^{2n})^p dx = \text{Int}((d + ex^n)^q (a + cx^{2n})^p, x)$$

output `Defer(Int)((d+e*x^n)^q*(a+c*x^(2*n))^p,x)`

Mathematica [N/A]

Not integrable

Time = 0.38 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int (d + ex^n)^q (a + cx^{2n})^p dx = \int (d + ex^n)^q (a + cx^{2n})^p dx$$

input `Integrate[(d + e*x^n)^q*(a + c*x^(2*n))^p,x]`

output `Integrate[(d + e*x^n)^q*(a + c*x^(2*n))^p, x]`

Rubi [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {1770}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + cx^{2n})^p (d + ex^n)^q dx \xrightarrow{1770} \int (a + cx^{2n})^p (d + ex^n)^q dx$$

input `Int[(d + e*x^n)^q*(a + c*x^(2*n))^p, x]`

output `$Aborted`

Definitions of rubi rules used

rule 1770 `Int[((d_) + (e_.)*(x_)^(n_))^(q_)*((a_) + (c_.)*(x_)^(n2_))^(p_), x_Symbol] :> Unintegrable[(d + e*x^n)^q*(a + c*x^(2*n))^p, x] /; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n]`

Maple [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int (d + e x^n)^q (a + c x^{2n})^p dx$$

input `int((d+e*x^n)^q*(a+c*x^(2*n))^p, x)`

output `int((d+e*x^n)^q*(a+c*x^(2*n))^p,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int (d + ex^n)^q (a + cx^{2n})^p dx = \int (cx^{2n} + a)^p (ex^n + d)^q dx$$

input `integrate((d+e*x^n)^q*(a+c*x^(2*n))^p,x, algorithm="fricas")`

output `integral((c*x^(2*n) + a)^p*(e*x^n + d)^q, x)`

Sympy [F(-1)]

Timed out.

$$\int (d + ex^n)^q (a + cx^{2n})^p dx = \text{Timed out}$$

input `integrate((d+e*x**n)**q*(a+c*x**2*n)**p,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int (d + ex^n)^q (a + cx^{2n})^p dx = \int (cx^{2n} + a)^p (ex^n + d)^q dx$$

input `integrate((d+e*x^n)^q*(a+c*x^(2*n))^p,x, algorithm="maxima")`

output `integrate((c*x^(2*n) + a)^p*(e*x^n + d)^q, x)`

Giac [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int (d + ex^n)^q (a + cx^{2n})^p dx = \int (cx^{2n} + a)^p (ex^n + d)^q dx$$

input `integrate((d+e*x^n)^q*(a+c*x^(2*n))^p,x, algorithm="giac")`

output `integrate((c*x^(2*n) + a)^p*(e*x^n + d)^q, x)`

Mupad [N/A]

Not integrable

Time = 11.38 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int (d + ex^n)^q (a + cx^{2n})^p dx = \int (a + cx^{2n})^p (d + ex^n)^q dx$$

input `int((a + c*x^(2*n))^p*(d + e*x^n)^q,x)`

output `int((a + c*x^(2*n))^p*(d + e*x^n)^q, x)`

Reduce [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 820, normalized size of antiderivative = 39.05

$$\int (d + ex^n)^q (a + cx^{2n})^p dx \\ = \frac{(x^n e + d)^q (x^{2n} c + a)^p x + 4 \left(\int \frac{(x^n e + d)^q (x^{2n} c + a)^p}{2x^{3n} c e n p + x^{3n} c e + 2x^{2n} c d n p + x^{2n} c d + 2x^n a e n p + x^n a e + 2a d n p + a d} dx \right) a d n^2 p^2 + 2 \left(\int \frac{(x^n e + d)^q (x^{2n} c + a)^p}{2x^{3n}} dx \right) a d n^2 p^2}{2x^{3n}}$$

input `int((d+e*x^n)^q*(a+c*x^(2*n))^p,x)`

output

```
((x**n*e + d)**q*(x**((2*n)*c + a)**p*x + 4*int(((x**n*e + d)**q*(x**((2*n)*c + a)**p)/(2*x**((3*n)*c*e*n*p + x**((3*n)*c*e + 2*x**((2*n)*c*d*n*p + x**((2*n)*c*d + 2*x**((n*a*e*n*p + x**n*a*e + 2*a*d*n*p + a*d),x)*a*d*n**2*p**2 + 2*int(((x**n*e + d)**q*(x**((2*n)*c + a)**p)/(2*x**((3*n)*c*e*n*p + x**((3*n)*c*e + 2*x**((2*n)*c*d*n*p + x**((2*n)*c*d + 2*x**((n*a*e*n*p + x**n*a*e + 2*a*d*n*p + a*d),x)*a*d*n*p - 2*int((x**((3*n)*(x**n*e + d)**q*(x**((2*n)*c + a)**p)/(2*x**((3*n)*c*e*n*p + x**((3*n)*c*e + 2*x**((2*n)*c*d*n*p + x**((2*n)*c*d + 2*x**((n*a*e*n*p + x**n*a*e + 2*a*d*n*p + a*d),x)*c*e*n**2*p*q - int((x**((3*n)*(x**n*e + d)**q*(x**((2*n)*c + a)**p)/(2*x**((3*n)*c*e*n*p + x**((3*n)*c*e + 2*x**((2*n)*c*d*n*p + x**((2*n)*c*d + 2*x**((n*a*e*n*p + x**n*a*e + 2*a*d*n*p + a*d),x)*c*e*n*q + 4*int((x**n*(x**n*e + d)**q*(x**((2*n)*c + a)**p)/(2*x**((3*n)*c*e*n*p + x**((3*n)*c*e + 2*x**((2*n)*c*d*n*p + x**((2*n)*c*d + 2*x**((n*a*e*n*p + x**n*a*e + 2*a*d*n*p + a*d),x)*a*e*n**2*p**2 - 2*int((x**n*(x**n*e + d)**q*(x**((2*n)*c + a)**p)/(2*x**((3*n)*c*e*n*p + x**((3*n)*c*e + 2*x**((2*n)*c*d*n*p + x**((2*n)*c*d + 2*x**((n*a*e*n*p + x**n*a*e + 2*a*d*n*p + a*d),x)*a*e*n**2*p*q + 2*int((x**n*(x**n*e + d)**q*(x**((2*n)*c + a)**p)/(2*x**((3*n)*c*e*n*p + x**((3*n)*c*e + 2*x**((2*n)*c*d*n*p + x**((2*n)*c*d + 2*x**((n*a*e*n*p + x**n*a*e + 2*a*d*n*p + a*d),x)*a*e*n*p - int((x**n*(x**n*e + d)**q*(x**((2*n)*c + a)**p)/(2*x**((3*n)*c*e*n*p + x**((3*n)*c*e + 2*x**((2*n)*c*d*n*p + x**((2*n)*c*d + 2*x**((n*a*e*n*p + x**n*a*e + 2*a*d*n*p + a*d),x)*a*e*n*p + x**n*a*e + 2*a*d*n*p ...
```

3.54 $\int (d + ex^n)^q (cd^2 - ce^2x^{2n})^p dx$

Optimal result	377
Mathematica [F]	377
Rubi [A] (verified)	378
Maple [F]	379
Fricas [F]	379
Sympy [F(-1)]	380
Maxima [F]	380
Giac [F]	380
Mupad [F(-1)]	381
Reduce [F]	381

Optimal result

Integrand size = 29, antiderivative size = 99

$$\int (d + ex^n)^q (cd^2 - ce^2x^{2n})^p dx = x(d + ex^n)^q \left(1 - \frac{ex^n}{d}\right)^{-p} \left(1 + \frac{ex^n}{d}\right)^{-p-q} (cd^2 - ce^2x^{2n})^p \text{AppellF1} \left(\frac{1}{n}, -p, -p - q, 1 + \frac{1}{n}, \frac{ex^n}{d}, -\frac{ex^n}{d}\right)$$

output $x*(d+e*x^n)^q*(1+e*x^n/d)^{(-p-q)}*(c*d^2-c*e^2*x^(2*n))^p*\text{AppellF1}(1/n,-p,-p-q,1+1/n,e*x^n/d,-e*x^n/d)/((1-e*x^n/d)^p)$

Mathematica [F]

$$\int (d + ex^n)^q (cd^2 - ce^2x^{2n})^p dx = \int (d + ex^n)^q (cd^2 - ce^2x^{2n})^p dx$$

input `Integrate[(d + e*x^n)^q*(c*d^2 - c*e^2*x^(2*n))^p, x]`

output `Integrate[(d + e*x^n)^q*(c*d^2 - c*e^2*x^(2*n))^p, x]`

Rubi [A] (verified)

Time = 0.29 (sec), antiderivative size = 99, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.138$, Rules used = {1396, 937, 937, 936}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (d + ex^n)^q (cd^2 - ce^2 x^{2n})^p dx \\
 & \quad \downarrow \textcolor{blue}{1396} \\
 & (d + ex^n)^{-p} (cd - cex^n)^{-p} (cd^2 - ce^2 x^{2n})^p \int (ex^n + d)^{p+q} (cd - cex^n)^p dx \\
 & \quad \downarrow \textcolor{blue}{937} \\
 & (d + ex^n)^q (cd - cex^n)^{-p} (cd^2 - ce^2 x^{2n})^p \left(\frac{ex^n}{d} + 1 \right)^{-p-q} \int (cd - cex^n)^p \left(\frac{ex^n}{d} + 1 \right)^{p+q} dx \\
 & \quad \downarrow \textcolor{blue}{937} \\
 & \left(1 - \frac{ex^n}{d} \right)^{-p} (d + ex^n)^q (cd^2 - ce^2 x^{2n})^p \left(\frac{ex^n}{d} + 1 \right)^{-p-q} \int \left(1 - \frac{ex^n}{d} \right)^p \left(\frac{ex^n}{d} + 1 \right)^{p+q} dx \\
 & \quad \downarrow \textcolor{blue}{936} \\
 & x \left(1 - \frac{ex^n}{d} \right)^{-p} (d + ex^n)^q (cd^2 - ce^2 x^{2n})^p \left(\frac{ex^n}{d} + 1 \right)^{-p-q} \text{AppellF1} \left(\frac{1}{n}, -p, -p - q, 1 + \frac{1}{n}, \frac{ex^n}{d}, -\frac{ex^n}{d} \right)
 \end{aligned}$$

input `Int[(d + e*x^n)^q*(c*d^2 - c*e^2*x^(2*n))^p, x]`

output `(x*(d + e*x^n)^q*(1 + (e*x^n)/d)^(-p - q)*(c*d^2 - c*e^2*x^(2*n))^p*AppellF1[F1[n^(-1), -p, -p - q, 1 + n^(-1), (e*x^n)/d, -((e*x^n)/d)])/(1 - (e*x^n)/d)^p`

Definitions of rubi rules used

rule 936 $\text{Int}[(a_+ + b_-) \cdot (x_-)^{(n_-)} \cdot (p_-) \cdot ((c_+ + d_-) \cdot (x_-)^{(n_-)})^{(q_-)}, x_{\text{Symbol}}] \\ \rightarrow \text{Simp}[a^p \cdot c^q \cdot x \cdot \text{AppellF1}[1/n, -p, -q, 1 + 1/n, (-b) \cdot (x^n/a), (-d) \cdot (x^n/c)], x] /; \text{FreeQ}[\{a, b, c, d, n, p, q\}, x] \& \text{NeQ}[b*c - a*d, 0] \& \text{NeQ}[n, -1] \\ \& \& (\text{IntegerQ}[p] \mid\mid \text{GtQ}[a, 0]) \& (\text{IntegerQ}[q] \mid\mid \text{GtQ}[c, 0])$

rule 937 $\text{Int}[(a_+ + b_-) \cdot (x_-)^{(n_-)} \cdot (p_-) \cdot ((c_+ + d_-) \cdot (x_-)^{(n_-)})^{(q_-)}, x_{\text{Symbol}}] \\ \rightarrow \text{Simp}[a^{\text{IntPart}[p]} \cdot ((a + b \cdot x^n)^{\text{FracPart}[p]} / (1 + b \cdot (x^n/a))^{\text{FracPart}[p]}) \cdot \text{Int}[(1 + b \cdot (x^n/a))^{p_*} \cdot (c + d \cdot x^n)^q, x], x] /; \text{FreeQ}[\{a, b, c, d, n, p, q\}, x] \& \text{NeQ}[b*c - a*d, 0] \& \text{NeQ}[n, -1] \& !(\text{IntegerQ}[p] \mid\mid \text{GtQ}[a, 0])$

rule 1396 $\text{Int}[(u_-) \cdot ((a_+ + c_-) \cdot (x_-)^{(n2_-)})^{(p_-)} \cdot ((d_+ + e_-) \cdot (x_-)^{(n_-)})^{(q_-)}, x_{\text{Symbol}}] \\ \rightarrow \text{Simp}[(a + c \cdot x^{(2*n)})^{\text{FracPart}[p]} / ((d + e \cdot x^n)^{\text{FracPart}[p]} \cdot (a/d + c \cdot (x^n/e))^{\text{FracPart}[p]}) \cdot \text{Int}[u \cdot (d + e \cdot x^n)^{(p+q)} \cdot (a/d + (c/e) \cdot x^n)^p, x], x] /; \text{FreeQ}[\{a, c, d, e, n, p, q\}, x] \& \text{EqQ}[n2, 2*n] \& \text{EqQ}[c*d^2 + a*e^2, 0] \& !\text{IntegerQ}[p] \& !(\text{EqQ}[q, 1] \& \text{EqQ}[n, 2])$

Maple [F]

$$\int (d + e x^n)^q (c d^2 - c e^2 x^{2n})^p dx$$

input $\text{int}((d+e*x^n)^q * (c*d^2 - c*e^2*x^(2*n))^p, x)$

output $\text{int}((d+e*x^n)^q * (c*d^2 - c*e^2*x^(2*n))^p, x)$

Fricas [F]

$$\int (d + e x^n)^q (c d^2 - c e^2 x^{2n})^p dx = \int (-c e^2 x^{2n} + c d^2)^p (e x^n + d)^q dx$$

input $\text{integrate}((d+e*x^n)^q * (c*d^2 - c*e^2*x^(2*n))^p, x, \text{algorithm}=\text{"fricas"})$

output $\text{integral}((-c*e^{2*x^{(2*n)}} + c*d^2)^p * (e*x^n + d)^q, x)$

Sympy [F(-1)]

Timed out.

$$\int (d + ex^n)^q (cd^2 - ce^2 x^{2n})^p dx = \text{Timed out}$$

input $\text{integrate}((d+e*x^{(2*n)})^q * (c*d^2 - c*e^{2*x^{(2*n)}})^p, x)$

output Timed out

Maxima [F]

$$\int (d + ex^n)^q (cd^2 - ce^2 x^{2n})^p dx = \int (-ce^2 x^{2n} + cd^2)^p (ex^n + d)^q dx$$

input $\text{integrate}((d+e*x^n)^q * (c*d^2 - c*e^{2*x^{(2*n)}})^p, x, \text{algorithm}=\text{"maxima"})$

output $\text{integrate}((-c*e^{2*x^{(2*n)}} + c*d^2)^p * (e*x^n + d)^q, x)$

Giac [F]

$$\int (d + ex^n)^q (cd^2 - ce^2 x^{2n})^p dx = \int (-ce^2 x^{2n} + cd^2)^p (ex^n + d)^q dx$$

input $\text{integrate}((d+e*x^n)^q * (c*d^2 - c*e^{2*x^{(2*n)}})^p, x, \text{algorithm}=\text{"giac"})$

output $\text{integrate}((-c*e^{2*x^{(2*n)}} + c*d^2)^p * (e*x^n + d)^q, x)$

Mupad [F(-1)]

Timed out.

$$\int (d + ex^n)^q (cd^2 - ce^2 x^{2n})^p dx = \int (cd^2 - ce^2 x^{2n})^p (d + ex^n)^q dx$$

input `int((c*d^2 - c*e^2*x^(2*n))^p*(d + e*x^n)^q,x)`

output `int((c*d^2 - c*e^2*x^(2*n))^p*(d + e*x^n)^q, x)`

Reduce [F]

$$\int (d + ex^n)^q (cd^2 - ce^2 x^{2n})^p dx = \int (x^n e + d)^q (-x^{2n} c e^2 + c d^2)^p dx$$

input `int((d+e*x^n)^q*(c*d^2-c*e^2*x^(2*n))^p,x)`

output `int((x**n*e + d)**q*(-x**2*n*c*e**2 + c*d**2)**p,x)`

3.55 $\int (2 - ex^n)^p (2 + ex^n)^{p+q} dx$

Optimal result	382
Mathematica [B] (warning: unable to verify)	382
Rubi [A] (verified)	383
Maple [F]	384
Fricas [F]	384
Sympy [F]	384
Maxima [F]	385
Giac [F]	385
Mupad [F(-1)]	385
Reduce [F]	386

Optimal result

Integrand size = 22, antiderivative size = 44

$$\int (2 - ex^n)^p (2 + ex^n)^{p+q} dx = 2^{2p+q} x \text{AppellF1} \left(\frac{1}{n}, -p, -p - q, 1 + \frac{1}{n}, \frac{ex^n}{2}, -\frac{ex^n}{2} \right)$$

output $2^{(2*p+q)}*x*\text{AppellF1}(1/n, -p-q, -p, 1+1/n, -1/2*e*x^n, 1/2*e*x^n)$

Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 199 vs. $2(44) = 88$.

Time = 0.40 (sec), antiderivative size = 199, normalized size of antiderivative = 4.52

$$\begin{aligned} & \int (2 - ex^n)^p (2 + ex^n)^{p+q} dx \\ &= \frac{2(1+n)x(2 - ex^n)^p (2 + ex^n)^{p+q} \text{AppellF1} \left(\frac{1}{n}, -p, -p - q, 1 + \frac{1}{n}, \frac{ex^n}{2}, -\frac{ex^n}{2} \right)}{-enpx^n \text{AppellF1} \left(1 + \frac{1}{n}, 1 - p, -p - q, 2 + \frac{1}{n}, \frac{ex^n}{2}, -\frac{ex^n}{2} \right) + en(p + q)x^n \text{AppellF1} \left(1 + \frac{1}{n}, -p, 1 - p - q, 2 + \frac{1}{n}, \frac{ex^n}{2}, -\frac{ex^n}{2} \right)} \end{aligned}$$

input `Integrate[(2 - e*x^n)^p*(2 + e*x^n)^(p + q), x]`

output
$$(2*(1 + n)*x*(2 - e*x^n)^p*(2 + e*x^n)^(p + q)*AppellF1[n^(-1), -p, -p - q, 1 + n^(-1), (e*x^n)/2, -1/2*(e*x^n)])/(-(e*n*p*x^n*AppellF1[1 + n^(-1), 1 - p, -p - q, 2 + n^(-1), (e*x^n)/2, -1/2*(e*x^n)]) + e*n*(p + q)*x^n*AppellF1[1 + n^(-1), -p, 1 - p - q, 2 + n^(-1), (e*x^n)/2, -1/2*(e*x^n)] + 2*(1 + n)*AppellF1[n^(-1), -p, -p - q, 1 + n^(-1), (e*x^n)/2, -1/2*(e*x^n)])$$

Rubi [A] (verified)

Time = 0.18 (sec), antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.045$, Rules used = {936}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (2 - ex^n)^p (ex^n + 2)^{p+q} dx \\ & \quad \downarrow \textcolor{blue}{936} \\ & x^{2p+q} \operatorname{AppellF1}\left(\frac{1}{n}, -p, -p - q, 1 + \frac{1}{n}, \frac{ex^n}{2}, -\frac{ex^n}{2}\right) \end{aligned}$$

input $\operatorname{Int}[(2 - e*x^n)^p*(2 + e*x^n)^(p + q), x]$

output $2^{(2p + q)}*x*\operatorname{AppellF1}[n^(-1), -p, -p - q, 1 + n^(-1), (e*x^n)/2, -1/2*(e*x^n)]$

Definitions of rubi rules used

rule 936
$$\begin{aligned} & \operatorname{Int}[(a_0 + b_0)*(x_0)^{(n_0)})^{(p_0)}*((c_0) + (d_0)*(x_0)^{(n_0)})^{(q_0)}, x_{\text{Symbol}}] \\ & :> \operatorname{Simp}[a^{p_0}c^{q_0}x^{p_0+q_0}\operatorname{AppellF1}[1/n, -p_0, -q_0, 1 + 1/n, (-b_0)*(x^{n_0}/a_0), (-d_0)*(x^{n_0}/c_0)], x] /; \operatorname{FreeQ}[\{a_0, b_0, c_0, d_0, n_0, p_0, q_0\}, x] \&& \operatorname{NeQ}[b_0*c_0 - a_0*d_0, 0] \&& \operatorname{NeQ}[n_0, -1] \\ & \&& (\operatorname{IntegerQ}[p_0] \&& \operatorname{GtQ}[a_0, 0]) \&& (\operatorname{IntegerQ}[q_0] \&& \operatorname{GtQ}[c_0, 0]) \end{aligned}$$

Maple [F]

$$\int (2 - e x^n)^p (2 + e x^n)^{p+q} dx$$

input `int((2-e*x^n)^p*(2+e*x^n)^(p+q),x)`

output `int((2-e*x^n)^p*(2+e*x^n)^(p+q),x)`

Fricas [F]

$$\int (2 - ex^n)^p (2 + ex^n)^{p+q} dx = \int (ex^n + 2)^{p+q} (-ex^n + 2)^p dx$$

input `integrate((2-e*x^n)^p*(2+e*x^n)^(p+q),x, algorithm="fricas")`

output `integral((e*x^n + 2)^(p + q)*(-e*x^n + 2)^p, x)`

Sympy [F]

$$\int (2 - ex^n)^p (2 + ex^n)^{p+q} dx = \int (-ex^n + 2)^p (ex^n + 2)^{p+q} dx$$

input `integrate((2-e*x**n)**p*(2+e*x**n)**(p+q),x)`

output `Integral((-e*x**n + 2)**p*(e*x**n + 2)**(p + q), x)`

Maxima [F]

$$\int (2 - ex^n)^p (2 + ex^n)^{p+q} dx = \int (ex^n + 2)^{p+q} (-ex^n + 2)^p dx$$

input `integrate((2-e*x^n)^p*(2+e*x^n)^(p+q),x, algorithm="maxima")`

output `integrate((e*x^n + 2)^(p + q)*(-e*x^n + 2)^p, x)`

Giac [F]

$$\int (2 - ex^n)^p (2 + ex^n)^{p+q} dx = \int (ex^n + 2)^{p+q} (-ex^n + 2)^p dx$$

input `integrate((2-e*x^n)^p*(2+e*x^n)^(p+q),x, algorithm="giac")`

output `integrate((e*x^n + 2)^(p + q)*(-e*x^n + 2)^p, x)`

Mupad [F(-1)]

Timed out.

$$\int (2 - ex^n)^p (2 + ex^n)^{p+q} dx = \int (2 - e x^n)^p (e x^n + 2)^{p+q} dx$$

input `int((2 - e*x^n)^p*(e*x^n + 2)^(p + q),x)`

output `int((2 - e*x^n)^p*(e*x^n + 2)^(p + q), x)`

Reduce [F]

$$\int (2 - ex^n)^p (2 + ex^n)^{p+q} dx = \int (x^n e + 2)^{p+q} (-x^n e + 2)^p dx$$

input `int((2-e*x^n)^p*(2+e*x^n)^(p+q),x)`

output `int((x**n*e + 2)**(p + q)*(- x**n*e + 2)**p,x)`

3.56 $\int (2 + ex^n)^q (4 - e^2 x^{2n})^p dx$

Optimal result	387
Mathematica [F]	387
Rubi [A] (verified)	388
Maple [F]	389
Fricas [F]	389
Sympy [F(-1)]	389
Maxima [F]	390
Giac [F]	390
Mupad [F(-1)]	390
Reduce [F]	391

Optimal result

Integrand size = 24, antiderivative size = 44

$$\int (2 + ex^n)^q (4 - e^2 x^{2n})^p dx = 2^{2p+q} x \text{AppellF1} \left(\frac{1}{n}, -p, -p - q, 1 + \frac{1}{n}, \frac{ex^n}{2}, -\frac{ex^n}{2} \right)$$

output `2^(2*p+q)*x*AppellF1(1/n, -p-q, -p, 1+1/n, -1/2*e*x^n, 1/2*e*x^n)`

Mathematica [F]

$$\int (2 + ex^n)^q (4 - e^2 x^{2n})^p dx = \int (2 + ex^n)^q (4 - e^2 x^{2n})^p dx$$

input `Integrate[(2 + e*x^n)^q*(4 - e^2*x^(2*n))^p, x]`

output `Integrate[(2 + e*x^n)^q*(4 - e^2*x^(2*n))^p, x]`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {1388, 936}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (4 - e^2 x^{2n})^p (ex^n + 2)^q dx \\ & \quad \downarrow \textcolor{blue}{1388} \\ & \int (2 - ex^n)^p (ex^n + 2)^{p+q} dx \\ & \quad \downarrow \textcolor{blue}{936} \\ & x^{2p+q} \text{AppellF1} \left(\frac{1}{n}, -p, -p - q, 1 + \frac{1}{n}, \frac{ex^n}{2}, -\frac{ex^n}{2} \right) \end{aligned}$$

input `Int[(2 + e*x^n)^q*(4 - e^2*x^(2*n))^p, x]`

output `2^(2*p + q)*x*AppellF1[n^(-1), -p, -p - q, 1 + n^(-1), (e*x^n)/2, -1/2*(e*x^n)]`

Definitions of rubi rules used

rule 936 `Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] :> Simp[a^p*c^q*x*AppellF1[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`

rule 1388 `Int[(u_)*(a_ + (c_)*(x_)^(n2_))^(p_)*(d_ + (e_)*(x_)^(n_))^(q_), x_Symbol] :> Int[u*(d + e*x^n)^(p + q)*(a/d + (c/e)*x^n)^p, x] /; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[c*d^2 + a*e^2, 0] && (IntegerQ[p] || (GtQ[a, 0] && GtQ[d, 0]))`

Maple [F]

$$\int (2 + e^{x^n})^q (4 - e^2 x^{2n})^p dx$$

input `int((2+e*x^n)^q*(4-e^2*x^(2*n))^p,x)`

output `int((2+e*x^n)^q*(4-e^2*x^(2*n))^p,x)`

Fricas [F]

$$\int (2 + e^{x^n})^q (4 - e^2 x^{2n})^p dx = \int (-e^2 x^{2n} + 4)^p (e^{x^n} + 2)^q dx$$

input `integrate((2+e*x^n)^q*(4-e^2*x^(2*n))^p,x, algorithm="fricas")`

output `integral((-e^2*x^(2*n) + 4)^p*(e*x^n + 2)^q, x)`

Sympy [F(-1)]

Timed out.

$$\int (2 + e^{x^n})^q (4 - e^2 x^{2n})^p dx = \text{Timed out}$$

input `integrate((2+e*x**n)**q*(4-e**2*x**2*n)**p,x)`

output `Timed out`

Maxima [F]

$$\int (2 + ex^n)^q (4 - e^2 x^{2n})^p dx = \int (-e^2 x^{2n} + 4)^p (ex^n + 2)^q dx$$

input `integrate((2+e*x^n)^q*(4-e^2*x^(2*n))^p,x, algorithm="maxima")`

output `integrate((-e^2*x^(2*n) + 4)^p*(e*x^n + 2)^q, x)`

Giac [F]

$$\int (2 + ex^n)^q (4 - e^2 x^{2n})^p dx = \int (-e^2 x^{2n} + 4)^p (ex^n + 2)^q dx$$

input `integrate((2+e*x^n)^q*(4-e^2*x^(2*n))^p,x, algorithm="giac")`

output `integrate((-e^2*x^(2*n) + 4)^p*(e*x^n + 2)^q, x)`

Mupad [F(-1)]

Timed out.

$$\int (2 + ex^n)^q (4 - e^2 x^{2n})^p dx = \int (4 - e^2 x^{2n})^p (e x^n + 2)^q dx$$

input `int((4 - e^2*x^(2*n))^p*(e*x^n + 2)^q,x)`

output `int((4 - e^2*x^(2*n))^p*(e*x^n + 2)^q, x)`

Reduce [F]

$$\int (2 + ex^n)^q (4 - e^2 x^{2n})^p dx = \int (x^n e + 2)^q (-x^{2n} e^2 + 4)^p dx$$

input `int((2+e*x^n)^q*(4-e^2*x^(2*n))^p,x)`

output `int((x**n*e + 2)**q*(- x**(2*n)*e**2 + 4)**p,x)`

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions	392
4.2 Links to plain text integration problems used in this report for each CAS .	410

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```
(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leaf
expnResult = ExpnType[result];
expnOptimal = ExpnType[optimal];
leafCountResult = LeafCount[result];
leafCountOptimal = LeafCount[optimal];

(*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
If[expnResult<=expnOptimal,
  If[Not[FreeQ[result,Complex]], (*result contains complex*)
    If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A","");
        ,(*ELSE*)
        finalresult={"B","Both result and optimal contain complex but leaf count
      ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*) (*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A","");
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
    ,(*ELSE*) (*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "}
      ,
      finalresult={"C","Result contains higher order function than in optimal. Order "}
    ]
  ]
]
]
```

```
finalresult={"F","Contains unresolved integral."}
]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hypergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
If[AtomQ[expn],
  1,
  If[ListQ[expn],
    Max[Map[ExpnType, expn]],
    If[Head[expn] === Power,
      If[IntegerQ[expn[[2]]],
        ExpnType[expn[[1]]],
        If[Head[expn[[2]]] === Rational,
          If[IntegerQ[expn[[1]]] || Head[expn[[1]]] === Rational,
            1,
            Max[ExpnType[expn[[1]]], 2]],
          Max[ExpnType[expn[[1]]], ExpnType[expn[[2]]], 3]]],
      If[Head[expn] === Plus || Head[expn] === Times,
        Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
        If[ElementaryFunctionQ[Head[expn]],
          Max[3, ExpnType[expn[[1]]]],
          If[SpecialFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Max[ExpnType[expn[[1]]], ExpnType[expn[[2]]], 3]]]]]]]]]
```

```
Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],  
If[AppellFunctionQ[Head[expn]],  
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],  
If[Head[expn]==RootSum,  
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],  
If[Head[expn]==Integrate || Head[expn]==Int,  
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],  
9]]]]]]]]]]]
```

```
ElementaryFunctionQ[func_] :=  
MemberQ[{  
Exp, Log,  
Sin, Cos, Tan, Cot, Sec, Csc,  
ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,  
Sinh, Cosh, Tanh, Coth, Sech, Csch,  
ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch  
} , func]
```

```
SpecialFunctionQ[func_] :=  
MemberQ[{  
Erf, Erfc, Erfi,  
FresnelS, FresnelC,  
ExpIntegralE, ExpIntegralEi, LogIntegral,  

```

```
HypergeometricFunctionQ[func_] :=  
MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]
```

```
AppellFunctionQ[func_] :=  
MemberQ[{AppellF1}, func]
```

Maple grading function

```
# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#           if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#           see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

leaf_count_result:=leafcount(result);
#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issue
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
#   "F" if the result fails to integrate an expression that
#       is integrable
#   "C" if result involves higher level functions than necessary
#   "B" if result is more than twice the size of the optimal
```

```
#      antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A", " ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf
                end if
            else #result contains complex but optimal is not
                if debug then
                    print("result contains complex but optimal is not");
                fi;
                return "C","Result contains complex when optimal does not.";
            fi;
        else # result do not contain complex
            # this assumes optimal do not as well. No check is needed here.
            if debug then
                print("result do not contain complex, this assumes optimal do not as well");
            fi;
```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                           convert(leaf_count_result,string),\"$ vs. \$2(", 
                           convert(leaf_count_optimal,string),")=",convert(2*leaf_co
                           fi;
            fi;
        else #ExpnType(result) > ExpnType(optimal)
            if debug then
                print("ExpnType(result) > ExpnType(optimal)");
            fi;
            return "C",cat("Result contains higher order function than in optimal. Order ",
                           convert(ExpnType_result,string)," vs. order ",
                           convert(ExpnType_optimal,string),"."));
        fi;

    end proc:

    #

    # is_contains_complex(result)
    # takes expressions and returns true if it contains "I" else false
    #
    #Nasser 032417
    is_contains_complex:= proc(expression)
        return (has(expression,I));
    end proc:

    # The following summarizes the type number assigned an expression
    # based on the functions it involves
    # 1 = rational function
    # 2 = algebraic function
    # 3 = elementary function
    # 4 = special function
    # 5 = hypergeometric function

```

```
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
    if type(expn,'atomic') then
        1
    elif type(expn,'list') then
        apply(max,map(ExpnType,expn))
    elif type(expn,'sqrt') then
        if type(op(1,expn),'rational') then
            1
        else
            max(2,ExpnType(op(1,expn)))
        end if
    elif type(expn,'`^`') then
        if type(op(2,expn),'integer') then
            ExpnType(op(1,expn))
        elif type(op(2,expn),'rational') then
            if type(op(1,expn),'rational') then
                1
            else
                max(2,ExpnType(op(1,expn)))
            end if
        else
            max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
        end if
    elif type(expn,'`+`) or type(expn,'`*`) then
        max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif ElementaryFunctionQ(op(0,expn)) then
        max(3,ExpnType(op(1,expn)))
    elif SpecialFunctionQ(op(0,expn)) then
        max(4,apply(max,map(ExpnType,[op(expn)])))
    elif HypergeometricFunctionQ(op(0,expn)) then
        max(5,apply(max,map(ExpnType,[op(expn)])))
    elif AppellFunctionQ(op(0,expn)) then
        max(6,apply(max,map(ExpnType,[op(expn)])))
    elif op(0,expn)='int' then
        max(8,apply(max,map(ExpnType,[op(expn)]))) else
9
```

```
end if
end proc:

ElementaryFunctionQ := proc(func)
member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,
    arcsin,arccos,arctan,arccot,arcsec,arccsc,
    sinh,cosh,tanh,coth,sech,csch,
    arccsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
member(func,[
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
member(func,[Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
member(func,[AppellF1])
end proc:

# u is a sum or product.  rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
if nops(u)=2 then
    op(2,u)
else
    apply(op(0,u),op(2..nops(u),u))
end if
end proc:

#leafcount(u) returns the number of nodes in u.
```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma][LeafCount](u);
end proc:
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]
```

```
def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False
    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'`^`)
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'`+') or type(expn,'`*')
```

```

m1 = expnType(expn.args[0])
m2 = expnType(list(expn.args[1:]))
return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]]
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

#print ("Enter grade_antiderivative for sagemath")
#print("Enter grade_antiderivative, result=",result, " optimal=",optimal)

leaf_count_result = leaf_count(result)
leaf_count_optimal = leaf_count(optimal)

#print("leaf_count_result=",leaf_count_result)
#print("leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if str(result).find("Integral") != -1:
    grade = "F"
    grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. "+str(grade)
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(grade)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(grade)

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
# Albert Rich to use with Sagemath. This is used to
# grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
# 'arctan2','floor','abs','log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
# issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```
from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    """
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
                        'sin','cos','tan','cot','sec','csc',
                        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
                        'sinh','cosh','tanh','coth','sech','csch',
                        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
                        'arctan2','floor','abs'
                       ]
    if debug:
```

```

if m:
    print ("func ", func , " is elementary_function")
else:
    print ("func ", func , " is NOT elementary_function")

return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral'
'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
'elliptic_pi','exp_integral_e','log_integral',
'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
'weierstrassPPrime','weierstrassSigma']

if debug:
    print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):
    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__)
    return False


def expnType(expn):

    if debug:
        print (">>>>Enter expnType, expn=", expn)
        print (">>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0], Rational)):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1], Integer)

```

```

    return expnType(expn.operands()[0])  #expnType(expn.args[0])
elif type(expn.operands()[1]) == Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0]) == Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0]))  #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1]))  #max(3,expnType(expn.args[0]),expnType(expn.args[1]))
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinstance(expn,Mul)
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2)  #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(expn.args)))
    return max(4,m1)  #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(expn.args)))
    return max(5,m1)  #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(expn.args)))
    return max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(expn.args)))
    return max(8,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sageMath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation ="none"
            else:
                grade = "B"
                grade_annotation ="Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation ="Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation ="none"
        else:
            grade = "B"
            grade_annotation ="Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)+"/"+str(leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation ="Result contains higher order function than in optimal. Order "+str(expnType_result)+"/"+str(expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file