

Computer Algebra Independent Integration Tests

Summer 2024

1-Algebraic-functions/1.2-Trinomial/1.2.4-Nested-quadratic-
trinomial/135-1.2.4.1

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 10:32pm

Contents

1	Introduction	4
1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Time and leaf size Performance	10
1.4	Performance based on number of rules Rubi used	12
1.5	Performance based on number of steps Rubi used	13
1.6	Solved integrals histogram based on leaf size of result	14
1.7	Solved integrals histogram based on CPU time used	15
1.8	Leaf size vs. CPU time used	16
1.9	list of integrals with no known antiderivative	17
1.10	List of integrals solved by CAS but has no known antiderivative	17
1.11	list of integrals solved by CAS but failed verification	17
1.12	Timing	18
1.13	Verification	18
1.14	Important notes about some of the results	19
1.15	Current tree layout of integration tests	22
1.16	Design of the test system	23
2	detailed summary tables of results	24
2.1	List of integrals sorted by grade for each CAS	25
2.2	Detailed conclusion table per each integral for all CAS systems	29
2.3	Detailed conclusion table specific for Rubi results	42
3	Listing of integrals	44
3.1	$\int (x + \sqrt{-3 - 2x + x^2})^3 dx$	47
3.2	$\int (x + \sqrt{-3 - 2x + x^2})^2 dx$	53
3.3	$\int (x + \sqrt{-3 - 2x + x^2}) dx$	59
3.4	$\int \frac{1}{x + \sqrt{-3 - 2x + x^2}} dx$	64
3.5	$\int \frac{1}{(x + \sqrt{-3 - 2x + x^2})^2} dx$	70

3.6	$\int \frac{1}{(x+\sqrt{-3-2x+x^2})^3} dx$	76
3.7	$\int \left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^3 dx$	83
3.8	$\int \left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^2 dx$	92
3.9	$\int \left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right) dx$	101
3.10	$\int \frac{1}{d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}} dx$	107
3.11	$\int \frac{1}{\left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^2} dx$	114
3.12	$\int \frac{1}{\left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^3} dx$	122
3.13	$\int \left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^{5/2} dx$	130
3.14	$\int \left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^{3/2} dx$	139
3.15	$\int \sqrt{d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}} dx$	147
3.16	$\int \frac{1}{\sqrt{d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}}} dx$	155
3.17	$\int \frac{1}{\left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^{3/2}} dx$	163
3.18	$\int \frac{1}{\left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^{5/2}} dx$	171
3.19	$\int \left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^p dx$	180
3.20	$\int (x+\sqrt{3-2x-x^2})^2 dx$	186
3.21	$\int (x+\sqrt{3-2x-x^2}) dx$	192
3.22	$\int \frac{1}{x+\sqrt{3-2x-x^2}} dx$	197
3.23	$\int \frac{1}{(x+\sqrt{3-2x-x^2})^2} dx$	207
3.24	$\int \frac{1}{(x+\sqrt{3-2x-x^2})^3} dx$	215
3.25	$\int (d+ex+f\sqrt{a+bx+cx^2})^2 dx$	224
3.26	$\int (d+ex+f\sqrt{a+bx+cx^2}) dx$	232
3.27	$\int \frac{1}{d+ex+f\sqrt{a+bx+cx^2}} dx$	238
3.28	$\int \frac{1}{(d+ex+f\sqrt{a+bx+cx^2})^2} dx$	245
3.29	$\int \frac{1}{(d+ex+f\sqrt{a+bx+cx^2})^3} dx$	255
3.30	$\int (x+\sqrt{-3-2x+4x^2})^3 dx$	261
3.31	$\int (x+\sqrt{-3-2x+4x^2})^2 dx$	267

3.32	$\int (x + \sqrt{-3 - 2x + 4x^2}) dx$	273
3.33	$\int \frac{1}{x + \sqrt{-3 - 2x + 4x^2}} dx$	279
3.34	$\int \frac{1}{(x + \sqrt{-3 - 2x + 4x^2})^2} dx$	288
3.35	$\int \frac{1}{(x + \sqrt{-3 - 2x + 4x^2})^3} dx$	296
3.36	$\int (d + ex + f\sqrt{-a + bx + cx^2})^2 dx$	305
3.37	$\int (d + ex + f\sqrt{-a + bx + cx^2}) dx$	313
3.38	$\int \frac{1}{d + ex + f\sqrt{-a + bx + cx^2}} dx$	319
3.39	$\int \frac{1}{(d + ex + f\sqrt{-a + bx + cx^2})^2} dx$	326
3.40	$\int \frac{1}{(d + ex + f\sqrt{-a + bx + cx^2})^3} dx$	332
3.41	$\int (x + \sqrt{-3 - 4x - x^2})^2 dx$	338
3.42	$\int (x + \sqrt{-3 - 4x - x^2}) dx$	343
3.43	$\int \frac{1}{x + \sqrt{-3 - 4x - x^2}} dx$	348
3.44	$\int \frac{1}{(x + \sqrt{-3 - 4x - x^2})^2} dx$	358
3.45	$\int \frac{1}{(x + \sqrt{-3 - 4x - x^2})^3} dx$	365
3.46	$\int (d + ex + f\sqrt{-a + bx - cx^2})^2 dx$	373
3.47	$\int (d + ex + f\sqrt{-a + bx - cx^2}) dx$	381
3.48	$\int \frac{1}{d + ex + f\sqrt{-a + bx - cx^2}} dx$	387
3.49	$\int \frac{1}{(d + ex + f\sqrt{-a + bx - cx^2})^2} dx$	394
3.50	$\int \frac{1}{(d + ex + f\sqrt{-a + bx - cx^2})^3} dx$	401
4	Appendix	407
4.1	Listing of Grading functions	407
4.2	Links to plain text integration problems used in this report for each CAS425	

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Time and leaf size Performance	10
1.4	Performance based on number of rules Rubi used	12
1.5	Performance based on number of steps Rubi used	13
1.6	Solved integrals histogram based on leaf size of result	14
1.7	Solved integrals histogram based on CPU time used	15
1.8	Leaf size vs. CPU time used	16
1.9	list of integrals with no known antiderivative	17
1.10	List of integrals solved by CAS but has no known antiderivative	17
1.11	list of integrals solved by CAS but failed verification	17
1.12	Timing	18
1.13	Verification	18
1.14	Important notes about some of the results	19
1.15	Current tree layout of integration tests	22
1.16	Design of the test system	23

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [50]. This is test number [135].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Mathematica	100.00 (50)	0.00 (0)
Rubi	90.00 (45)	10.00 (5)
Maple	86.00 (43)	14.00 (7)
Fricas	86.00 (43)	14.00 (7)
Reduce	68.00 (34)	32.00 (16)
Giac	64.00 (32)	36.00 (18)
Sympy	38.00 (19)	62.00 (31)
Mupad	32.00 (16)	68.00 (34)
Maxima	24.00 (12)	76.00 (38)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	84.000	6.000	0.000	10.000
Mathematica	84.000	6.000	10.000	0.000
Fricas	66.000	20.000	0.000	14.000
Maple	42.000	32.000	12.000	14.000
Giac	42.000	22.000	0.000	36.000
Sympy	38.000	0.000	0.000	62.000
Maxima	24.000	0.000	0.000	76.000
Mupad	0.000	32.000	0.000	68.000
Reduce	0.000	68.000	0.000	32.000

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Mathematica	0	0.00	0.00	0.00
Rubi	5	100.00	0.00	0.00
Fricas	7	0.00	100.00	0.00
Maple	7	100.00	0.00	0.00
Reduce	16	100.00	0.00	0.00
Giac	18	38.89	11.11	50.00
Sympy	31	96.77	3.23	0.00
Mupad	34	0.00	100.00	0.00
Maxima	38	81.58	0.00	18.42

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.08
Reduce	0.26
Giac	0.29
Sympy	0.50
Rubi	0.94
Fricas	1.87
Mathematica	7.04
Mupad	9.62
Maple	14.97

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mupad	86.69	0.86	61.00	0.84
Maxima	86.75	1.03	65.00	1.03
Sympy	216.11	1.39	126.00	1.17
Giac	217.84	1.42	158.50	1.01
Rubi	233.82	1.16	165.00	1.02
Reduce	516.15	2.48	144.50	1.27
Mathematica	542.64	1.45	134.00	0.97
Fricas	841.05	2.94	222.00	1.41
Maple	425505.35	414.69	129.00	0.92

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {22, 23, 24}

Mathematica {28, 29, 39, 40, 49, 50}

Maple {27, 28, 29, 38, 39, 40, 48, 49, 50}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	25
2.2	Detailed conclusion table per each integral for all CAS systems	29
2.3	Detailed conclusion table specific for Rubi results	42

2.1 List of integrals sorted by grade for each CAS

Rubi	25
Mma	25
Maple	26
Fricas	26
Maxima	26
Giac	27
Mupad	27
Sympy	27
Reduce	28

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 30, 31, 32, 33, 34, 35, 36, 37, 41, 42, 43, 44, 45, 46, 47, 48 }

B grade { 27, 38, 49 }

C grade { }

F normal fail { 28, 29, 39, 40, 50 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 30, 31, 32, 33, 34, 35, 36, 37, 38, 41, 42, 43, 44, 45, 46, 47 }

B grade { 13, 28, 29 }

C grade { 39, 40, 48, 49, 50 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 5, 6, 8, 9, 20, 21, 25, 26, 30, 31, 32, 36, 37, 41, 42, 46, 47 }

B grade { 7, 10, 11, 12, 22, 27, 28, 29, 33, 38, 39, 40, 43, 48, 49, 50 }

C grade { 23, 24, 34, 35, 44, 45 }

F normal fail { 13, 14, 15, 16, 17, 18, 19 }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 13, 14, 15, 16, 20, 21, 23, 24, 25, 26, 30, 31, 32, 34, 35, 36, 37, 41, 42, 44, 45, 46, 47 }

B grade { 11, 12, 17, 18, 22, 28, 33, 39, 43, 49 }

C grade { }

F normal fail { }

F(-1) timedout fail { 19, 27, 29, 38, 40, 48, 50 }

F(-2) exception fail { }

Maxima

A grade { 1, 2, 3, 20, 21, 30, 31, 32, 36, 37, 41, 42 }

B grade { }

C grade { }

F normal fail { 4, 5, 6, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 22, 23, 24, 27, 28, 29, 33, 34, 35, 38, 39, 40, 43, 44, 45, 48, 49, 50 }

F(-1) timedout fail { }

F(-2) exception fail { 7, 8, 9, 25, 26, 46, 47 }

Giac

A grade { 1, 2, 3, 4, 7, 8, 9, 20, 21, 24, 25, 26, 30, 31, 32, 36, 37, 41, 42, 46, 47 }

B grade { 5, 6, 11, 22, 23, 33, 34, 35, 43, 44, 45 }

C grade { }

F normal fail { 13, 14, 15, 16, 17, 18, 19 }

F(-1) timeout fail { 10, 12 }

F(-2) exception fail { 27, 28, 29, 38, 39, 40, 48, 49, 50 }

Mupad

A grade { }

B grade { 1, 2, 3, 20, 21, 25, 26, 30, 31, 32, 36, 37, 41, 42, 46, 47 }

C grade { }

F normal fail { }

F(-1) timeout fail { 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 22, 23, 24, 27, 28, 29, 33, 34, 35, 38, 39, 40, 43, 44, 45, 48, 49, 50 }

F(-2) exception fail { }

Sympy

A grade { 1, 2, 3, 7, 8, 9, 20, 21, 25, 26, 30, 31, 32, 36, 37, 41, 42, 46, 47 }

B grade { }

C grade { }

F normal fail { 4, 5, 6, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 22, 23, 24, 27, 28, 29, 33, 34, 35, 38, 39, 40, 43, 44, 45, 48, 49 }

F(-1) timeout fail { 50 }

F(-2) exception fail { }

Reduce

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 20, 21, 22, 23, 24, 25, 26, 30, 31, 32, 33, 34, 35, 36, 37, 41, 42, 43, 44, 45, 46, 47 }

C grade { }

F normal fail { 13, 14, 15, 16, 17, 18, 19, 27, 28, 29, 38, 39, 40, 48, 49, 50 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	104	68	59	81	59	114	57	83	64
N.S.	1	1.04	0.68	0.59	0.81	0.59	1.14	0.57	0.83	0.64
time (sec)	N/A	0.215	0.232	0.363	0.031	0.092	0.523	0.117	0.163	0.510

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	86	63	54	70	54	60	54	70	52
N.S.	1	1.19	0.88	0.75	0.97	0.75	0.83	0.75	0.97	0.72
time (sec)	N/A	0.221	0.171	0.198	0.029	0.084	0.416	0.121	0.153	0.106

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	50	48	39	51	39	44	40	48	38
N.S.	1	1.00	0.96	0.78	1.02	0.78	0.88	0.80	0.96	0.76
time (sec)	N/A	0.177	0.094	0.141	0.028	0.077	0.271	0.110	0.152	20.969

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	67	78	71	0	77	0	81	49	0
N.S.	1	1.03	1.20	1.09	0.00	1.18	0.00	1.25	0.75	0.00
time (sec)	N/A	0.216	0.315	0.092	0.000	0.084	0.000	0.129	0.163	0.000

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	85	69	61	0	97	0	143	107	0
N.S.	1	1.02	0.83	0.73	0.00	1.17	0.00	1.72	1.29	0.00
time (sec)	N/A	0.221	0.346	0.068	0.000	0.079	0.000	0.123	0.154	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	105	83	69	0	129	0	184	172	0
N.S.	1	1.04	0.82	0.68	0.00	1.28	0.00	1.82	1.70	0.00
time (sec)	N/A	0.226	0.391	0.073	0.000	0.110	0.000	0.125	0.162	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	A	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	303	312	260	685	0	345	1363	397	870	0
N.S.	1	1.03	0.86	2.26	0.00	1.14	4.50	1.31	2.87	0.00
time (sec)	N/A	0.534	1.917	0.046	0.000	0.121	1.245	0.142	0.395	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	237	246	176	300	0	219	490	236	497	0
N.S.	1	1.04	0.74	1.27	0.00	0.92	2.07	1.00	2.10	0.00
time (sec)	N/A	0.442	1.328	1.214	0.000	0.103	0.741	0.131	0.179	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	118	118	180	123	0	123	197	119	207	0
N.S.	1	1.00	1.53	1.04	0.00	1.04	1.67	1.01	1.75	0.00
time (sec)	N/A	0.254	0.558	0.332	0.000	0.086	0.309	0.115	0.159	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	A	F	F(-1)	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	215	222	266	1263	0	371	0	0	1760	0
N.S.	1	1.03	1.24	5.87	0.00	1.73	0.00	0.00	8.19	0.00
time (sec)	N/A	0.450	1.045	0.051	0.000	2.244	0.000	0.000	0.176	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	266	275	427	6305	0	826	0	1618	2122	0
N.S.	1	1.03	1.61	23.70	0.00	3.11	0.00	6.08	7.98	0.00
time (sec)	N/A	0.467	2.170	0.067	0.000	1.362	0.000	4.934	0.226	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	330	339	300	29137	0	1954	0	0	6354	0
N.S.	1	1.03	0.91	88.29	0.00	5.92	0.00	0.00	19.25	0.00
time (sec)	N/A	0.523	10.852	0.172	0.000	10.873	0.000	0.000	0.401	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	370	377	985	0	0	923	0	0	28	0
N.S.	1	1.02	2.66	0.00	0.00	2.49	0.00	0.00	0.08	0.00
time (sec)	N/A	0.880	6.856	0.000	0.000	0.305	0.000	0.000	200.024	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	302	309	443	0	0	657	0	0	28	0
N.S.	1	1.02	1.47	0.00	0.00	2.18	0.00	0.00	0.09	0.00
time (sec)	N/A	0.685	3.615	0.000	0.000	0.220	0.000	0.000	200.026	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	233	243	327	0	0	692	0	0	28	0
N.S.	1	1.04	1.40	0.00	0.00	2.97	0.00	0.00	0.12	0.00
time (sec)	N/A	0.508	1.970	0.000	0.000	0.217	0.000	0.000	0.437	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	246	276	341	0	0	716	0	0	28	0
N.S.	1	1.12	1.39	0.00	0.00	2.91	0.00	0.00	0.11	0.00
time (sec)	N/A	0.456	2.083	0.000	0.000	0.237	0.000	0.000	200.026	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	269	281	395	0	0	1456	0	0	28	0
N.S.	1	1.04	1.47	0.00	0.00	5.41	0.00	0.00	0.10	0.00
time (sec)	N/A	0.502	3.341	0.000	0.000	0.622	0.000	0.000	200.033	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	335	354	557	0	0	2396	0	0	28	0
N.S.	1	1.06	1.66	0.00	0.00	7.15	0.00	0.00	0.08	0.00
time (sec)	N/A	0.689	4.776	0.000	0.000	1.709	0.000	0.000	200.372	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	166	168	134	0	0	0	0	0	29	0
N.S.	1	1.01	0.81	0.00	0.00	0.00	0.00	0.00	0.17	0.00
time (sec)	N/A	0.362	1.475	0.000	0.000	0.000	0.000	0.000	3.453	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	56	51	60	60	41	40	61	55
N.S.	1	1.00	0.97	0.88	1.03	1.03	0.71	0.69	1.05	0.95
time (sec)	N/A	0.255	0.166	0.237	0.121	0.078	0.453	0.130	0.164	0.140

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	52	34	43	52	32	31	41	32
N.S.	1	1.00	1.27	0.83	1.05	1.27	0.78	0.76	1.00	0.78
time (sec)	N/A	0.176	0.097	0.148	0.106	0.088	0.288	0.117	0.165	0.084

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F	B	B	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	180	197	111	359	0	372	0	287	67	0
N.S.	1	1.09	0.62	1.99	0.00	2.07	0.00	1.59	0.37	0.00
time (sec)	N/A	0.520	0.418	0.536	0.000	0.119	0.000	0.164	0.174	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	A	F	B	B	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	172	165	94	105	0	171	0	350	146	0
N.S.	1	0.96	0.55	0.61	0.00	0.99	0.00	2.03	0.85	0.00
time (sec)	N/A	0.372	0.442	0.151	0.000	0.112	0.000	0.174	0.163	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	A	F	A	B	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	311	305	115	129	0	223	0	452	389	0
N.S.	1	0.98	0.37	0.41	0.00	0.72	0.00	1.45	1.25	0.00
time (sec)	N/A	0.551	0.543	0.145	0.000	0.078	0.000	0.167	0.159	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	158	226	156	194	0	429	345	174	368	190
N.S.	1	1.43	0.99	1.23	0.00	2.72	2.18	1.10	2.33	1.20
time (sec)	N/A	0.477	1.933	1.681	0.000	0.107	0.701	0.125	0.174	21.252

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	85	77	0	216	131	78	143	74
N.S.	1	1.00	0.97	0.88	0.00	2.45	1.49	0.89	1.62	0.84
time (sec)	N/A	0.217	0.573	0.298	0.000	0.099	0.297	0.127	0.164	20.746

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	A	B	F	F(-1)	F	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	361	1087	323	4879	0	0	0	0	23	0
N.S.	1	3.01	0.89	13.52	0.00	0.00	0.00	0.00	0.06	0.00
time (sec)	N/A	10.420	3.165	0.148	0.000	0.000	0.000	0.000	200.026	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	B	B	F	B	F	F(-2)	F	F(-1)
verified	N/A	N/A	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	394	0	2162	616160	0	6899	0	0	23	0
N.S.	1	0.00	5.49	1563.86	0.00	17.51	0.00	0.00	0.06	0.00
time (sec)	N/A	0.000	19.100	1.725	0.000	19.398	0.000	0.000	200.650	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	B	B	F	F(-1)	F	F(-2)	F	F(-1)
verified	N/A	N/A	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	1477	0	3355	6017742	0	0	0	0	23	0
N.S.	1	0.00	2.27	4074.30	0.00	0.00	0.00	0.00	0.02	0.00
time (sec)	N/A	0.000	21.960	256.430	0.000	0.000	0.000	0.000	200.027	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	108	108	77	67	96	67	126	68	101	114
N.S.	1	1.00	0.71	0.62	0.89	0.62	1.17	0.63	0.94	1.06
time (sec)	N/A	0.325	0.232	0.394	0.113	0.062	0.555	0.140	0.161	0.512

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	70	60	79	60	65	61	83	58
N.S.	1	1.00	0.82	0.71	0.93	0.71	0.76	0.72	0.98	0.68
time (sec)	N/A	0.259	0.143	0.210	0.105	0.084	0.467	0.117	0.161	20.916

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	59	57	47	57	47	49	48	59	44
N.S.	1	1.00	0.97	0.80	0.97	0.80	0.83	0.81	1.00	0.75
time (sec)	N/A	0.184	0.088	0.151	0.109	0.081	0.313	0.135	0.157	0.122

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	112	98	427	0	281	0	275	239	0
N.S.	1	1.06	0.92	4.03	0.00	2.65	0.00	2.59	2.25	0.00
time (sec)	N/A	0.367	0.411	0.371	0.000	0.079	0.000	0.165	0.176	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	A	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	149	156	87	105	0	171	0	320	648	0
N.S.	1	1.05	0.58	0.70	0.00	1.15	0.00	2.15	4.35	0.00
time (sec)	N/A	0.329	0.494	0.164	0.000	0.083	0.000	0.175	0.297	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	A	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	218	230	110	128	0	223	0	419	1122	0
N.S.	1	1.06	0.50	0.59	0.00	1.02	0.00	1.92	5.15	0.00
time (sec)	N/A	0.373	0.602	0.189	0.000	0.101	0.000	0.171	2.612	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	237	237	162	205	286	438	352	179	380	203
N.S.	1	1.00	0.68	0.86	1.21	1.85	1.49	0.76	1.60	0.86
time (sec)	N/A	0.480	1.115	1.761	0.039	0.094	0.710	0.159	0.175	22.857

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	124	81	117	222	134	82	147	78
N.S.	1	1.00	1.35	0.88	1.27	2.41	1.46	0.89	1.60	0.85
time (sec)	N/A	0.217	0.299	0.298	0.030	0.089	0.290	0.113	0.164	22.957

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	A	B	F	F(-1)	F	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	300	1084	287	4829	0	0	0	0	25	0
N.S.	1	3.61	0.96	16.10	0.00	0.00	0.00	0.00	0.08	0.00
time (sec)	N/A	10.301	1.729	0.130	0.000	0.000	0.000	0.000	200.023	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	C	B	F	B	F	F(-2)	F	F(-1)
verified	N/A	N/A	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	385	0	2401	508167	0	6919	0	0	25	0
N.S.	1	0.00	6.24	1319.91	0.00	17.97	0.00	0.00	0.06	0.00
time (sec)	N/A	0.000	50.690	1.246	0.000	20.290	0.000	0.000	200.040	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	C	B	F	F(-1)	F	F(-2)	F	F(-1)
verified	N/A	N/A	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	896	0	3628	4527517	0	0	0	0	25	0
N.S.	1	0.00	4.05	5053.03	0.00	0.00	0.00	0.00	0.03	0.00
time (sec)	N/A	0.000	75.180	114.039	0.000	0.000	0.000	0.000	200.037	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	52	52	56	48	60	58	39	36	59	55
N.S.	1	1.00	1.08	0.92	1.15	1.12	0.75	0.69	1.13	1.06
time (sec)	N/A	0.251	0.167	0.234	0.113	0.089	0.485	0.144	0.153	0.143

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	52	32	41	52	29	29	37	30
N.S.	1	1.00	1.41	0.86	1.11	1.41	0.78	0.78	1.00	0.81
time (sec)	N/A	0.166	0.101	0.143	0.124	0.098	0.316	0.122	0.150	0.082

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	96	124	81	370	0	187	0	197	44	0
N.S.	1	1.29	0.84	3.85	0.00	1.95	0.00	2.05	0.46	0.00
time (sec)	N/A	0.338	0.349	0.518	0.000	0.087	0.000	0.124	0.157	0.000

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	A	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	100	84	110	0	121	0	263	92	0
N.S.	1	0.99	0.83	1.09	0.00	1.20	0.00	2.60	0.91	0.00
time (sec)	N/A	0.272	0.396	0.182	0.000	0.080	0.000	0.124	0.153	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	A	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	174	163	109	131	0	171	0	367	135	0
N.S.	1	0.94	0.63	0.75	0.00	0.98	0.00	2.11	0.78	0.00
time (sec)	N/A	0.333	0.490	0.305	0.000	0.096	0.000	0.141	0.154	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	243	243	165	210	0	450	357	191	617	215
N.S.	1	1.00	0.68	0.86	0.00	1.85	1.47	0.79	2.54	0.88
time (sec)	N/A	0.496	2.219	1.754	0.000	0.096	0.768	0.133	0.244	22.319

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	94	94	134	83	0	235	138	95	232	85
N.S.	1	1.00	1.43	0.88	0.00	2.50	1.47	1.01	2.47	0.90
time (sec)	N/A	0.213	0.444	0.309	0.000	0.106	0.322	0.121	0.167	0.199

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	F(-1)	F	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	621	397	1737	4815	0	0	0	0	26	0
N.S.	1	0.64	2.80	7.75	0.00	0.00	0.00	0.00	0.04	0.00
time (sec)	N/A	1.878	2.474	0.152	0.000	0.000	0.000	0.000	200.042	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	C	B	F	B	F	F(-2)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	463	965	2346	609986	0	6908	0	0	26	0
N.S.	1	2.08	5.07	1317.46	0.00	14.92	0.00	0.00	0.06	0.00
time (sec)	N/A	4.308	49.723	1.740	0.000	20.124	0.000	0.000	200.025	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	C	B	F	F(-1)	F(-1)	F(-2)	F	F(-1)
verified	N/A	N/A	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	1382	0	3558	5961416	0	0	0	0	26	0
N.S.	1	0.00	2.57	4313.62	0.00	0.00	0.00	0.00	0.02	0.00
time (sec)	N/A	0.000	72.553	255.152	0.000	0.000	0.000	0.000	200.029	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [43] had the largest ratio of [.666667000000000010]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	5	4	1.04	16	0.250
2	A	5	4	1.19	16	0.250
3	A	1	1	1.00	14	0.071
4	A	5	4	1.03	16	0.250
5	A	5	4	1.02	16	0.250
6	A	5	4	1.04	16	0.250
7	A	4	3	1.03	28	0.107
8	A	4	3	1.04	28	0.107
9	A	1	1	1.00	26	0.038
10	A	4	3	1.03	28	0.107
11	A	4	3	1.03	28	0.107
12	A	4	3	1.03	28	0.107
13	A	7	6	1.02	30	0.200
14	A	6	5	1.02	30	0.167
15	A	7	6	1.04	30	0.200
16	A	7	6	1.12	30	0.200
17	A	7	6	1.04	30	0.200
18	A	7	6	1.06	30	0.200
19	A	4	3	1.01	28	0.107
20	A	2	2	1.00	18	0.111
21	A	1	1	1.00	16	0.062

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	12	11	1.09	18	0.611
23	A	7	6	0.96	18	0.333
24	A	8	7	0.98	18	0.389
25	A	2	2	1.43	23	0.087
26	A	1	1	1.00	21	0.048
27	B	2	2	3.01	23	0.087
28	F	0	0	N/A	0.000	N/A
29	F	0	0	N/A	0.000	N/A
30	A	2	2	1.00	18	0.111
31	A	2	2	1.00	18	0.111
32	A	1	1	1.00	16	0.062
33	A	5	4	1.06	18	0.222
34	A	7	6	1.05	18	0.333
35	A	8	7	1.06	18	0.389
36	A	2	2	1.00	25	0.080
37	A	1	1	1.00	23	0.043
38	B	2	2	3.61	25	0.080
39	F	0	0	N/A	0.000	N/A
40	F	0	0	N/A	0.000	N/A
41	A	2	2	1.00	18	0.111
42	A	1	1	1.00	16	0.062
43	A	13	12	1.29	18	0.667
44	A	6	5	0.99	18	0.278
45	A	8	7	0.94	18	0.389
46	A	2	2	1.00	26	0.077
47	A	1	1	1.00	24	0.042
48	A	2	2	0.64	26	0.077
49	B	2	2	2.08	26	0.077
50	F	0	0	N/A	0.000	N/A

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int (x + \sqrt{-3 - 2x + x^2})^3 dx$	47
3.2	$\int (x + \sqrt{-3 - 2x + x^2})^2 dx$	53
3.3	$\int (x + \sqrt{-3 - 2x + x^2}) dx$	59
3.4	$\int \frac{1}{x + \sqrt{-3 - 2x + x^2}} dx$	64
3.5	$\int \frac{1}{(x + \sqrt{-3 - 2x + x^2})^2} dx$	70
3.6	$\int \frac{1}{(x + \sqrt{-3 - 2x + x^2})^3} dx$	76
3.7	$\int \left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^3 dx$	83
3.8	$\int \left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^2 dx$	92
3.9	$\int \left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right) dx$	101
3.10	$\int \frac{1}{d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}} dx$	107
3.11	$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^2} dx$	114
3.12	$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^3} dx$	122
3.13	$\int \left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{5/2} dx$	130
3.14	$\int \left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{3/2} dx$	139
3.15	$\int \sqrt{d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}} dx$	147
3.16	$\int \frac{1}{\sqrt{d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}}} dx$	155
3.17	$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{3/2}} dx$	163

3.18	$\int \frac{1}{\left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^{5/2}} dx$	171
3.19	$\int \left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^p dx$	180
3.20	$\int \left(x+\sqrt{3-2x-x^2}\right)^2 dx$	186
3.21	$\int \left(x+\sqrt{3-2x-x^2}\right) dx$	192
3.22	$\int \frac{1}{x+\sqrt{3-2x-x^2}} dx$	197
3.23	$\int \frac{1}{\left(x+\sqrt{3-2x-x^2}\right)^2} dx$	207
3.24	$\int \frac{1}{\left(x+\sqrt{3-2x-x^2}\right)^3} dx$	215
3.25	$\int \left(d+ex+f\sqrt{a+bx+cx^2}\right)^2 dx$	224
3.26	$\int \left(d+ex+f\sqrt{a+bx+cx^2}\right) dx$	232
3.27	$\int \frac{1}{d+ex+f\sqrt{a+bx+cx^2}} dx$	238
3.28	$\int \frac{1}{\left(d+ex+f\sqrt{a+bx+cx^2}\right)^2} dx$	245
3.29	$\int \frac{1}{\left(d+ex+f\sqrt{a+bx+cx^2}\right)^3} dx$	255
3.30	$\int \left(x+\sqrt{-3-2x+4x^2}\right)^3 dx$	261
3.31	$\int \left(x+\sqrt{-3-2x+4x^2}\right)^2 dx$	267
3.32	$\int \left(x+\sqrt{-3-2x+4x^2}\right) dx$	273
3.33	$\int \frac{1}{x+\sqrt{-3-2x+4x^2}} dx$	279
3.34	$\int \frac{1}{\left(x+\sqrt{-3-2x+4x^2}\right)^2} dx$	288
3.35	$\int \frac{1}{\left(x+\sqrt{-3-2x+4x^2}\right)^3} dx$	296
3.36	$\int \left(d+ex+f\sqrt{-a+bx+cx^2}\right)^2 dx$	305
3.37	$\int \left(d+ex+f\sqrt{-a+bx+cx^2}\right) dx$	313
3.38	$\int \frac{1}{d+ex+f\sqrt{-a+bx+cx^2}} dx$	319
3.39	$\int \frac{1}{\left(d+ex+f\sqrt{-a+bx+cx^2}\right)^2} dx$	326
3.40	$\int \frac{1}{\left(d+ex+f\sqrt{-a+bx+cx^2}\right)^3} dx$	332
3.41	$\int \left(x+\sqrt{-3-4x-x^2}\right)^2 dx$	338
3.42	$\int \left(x+\sqrt{-3-4x-x^2}\right) dx$	343
3.43	$\int \frac{1}{x+\sqrt{-3-4x-x^2}} dx$	348
3.44	$\int \frac{1}{\left(x+\sqrt{-3-4x-x^2}\right)^2} dx$	358
3.45	$\int \frac{1}{\left(x+\sqrt{-3-4x-x^2}\right)^3} dx$	365
3.46	$\int \left(d+ex+f\sqrt{-a+bx-cx^2}\right)^2 dx$	373
3.47	$\int \left(d+ex+f\sqrt{-a+bx-cx^2}\right) dx$	381

3.48	$\int \frac{1}{d+ex+f\sqrt{-a+bx-cx^2}} dx$	387
3.49	$\int \frac{1}{(d+ex+f\sqrt{-a+bx-cx^2})^2} dx$	394
3.50	$\int \frac{1}{(d+ex+f\sqrt{-a+bx-cx^2})^3} dx$	401

3.1 $\int (x + \sqrt{-3 - 2x + x^2})^3 dx$

Optimal result	47
Mathematica [A] (verified)	47
Rubi [A] (verified)	48
Maple [A] (verified)	49
Fricas [A] (verification not implemented)	50
Sympy [A] (verification not implemented)	50
Maxima [A] (verification not implemented)	51
Giac [A] (verification not implemented)	51
Mupad [B] (verification not implemented)	52
Reduce [B] (verification not implemented)	52

Optimal result

Integrand size = 16, antiderivative size = 100

$$\int (x + \sqrt{-3 - 2x + x^2})^3 dx = -\frac{2}{1 - x - \sqrt{-3 - 2x + x^2}} - 4(x + \sqrt{-3 - 2x + x^2}) - (x + \sqrt{-3 - 2x + x^2})^2 + \frac{1}{8}(x + \sqrt{-3 - 2x + x^2})^4 - 6 \log(1 - x - \sqrt{-3 - 2x + x^2})$$

output

```
-2/(1-x-(x^2-2*x-3)^(1/2))-4*x-4*(x^2-2*x-3)^(1/2)-(x+(x^2-2*x-3)^(1/2))^2
+1/8*(x+(x^2-2*x-3)^(1/2))^4-6*ln(1-x-(x^2-2*x-3)^(1/2))
```

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.68

$$\int (x + \sqrt{-3 - 2x + x^2})^3 dx = -\frac{9x^2}{2} - 2x^3 + x^4 + \frac{1}{2}\sqrt{-3 - 2x + x^2}(-9 - 7x - 2x^2 + 2x^3) - 12 \operatorname{arctanh}\left(\frac{\sqrt{-3 - 2x + x^2}}{-3 + x}\right)$$

input

```
Integrate[(x + Sqrt[-3 - 2*x + x^2])^3, x]
```


output

$$\frac{(-9x^2)/2 - 2x^3 + x^4 + (\text{Sqrt}[-3 - 2x + x^2]) * (-9 - 7x - 2x^2 + 2x^3)}{2} - 12 * \text{ArcTanh}[\text{Sqrt}[-3 - 2x + x^2]/(-3 + x)]$$
Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.04, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2541, 27, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\sqrt{x^2 - 2x - 3} + x)^3 dx$$

$$\downarrow 2541$$

$$2 \int -\frac{(x + \sqrt{x^2 - 2x - 3})^3 \left(-(x + \sqrt{x^2 - 2x - 3})^2 + 2(x + \sqrt{x^2 - 2x - 3}) + 3 \right)}{4(-x - \sqrt{x^2 - 2x - 3} + 1)^2} d(x + \sqrt{x^2 - 2x - 3})$$

$$\downarrow 27$$

$$-\frac{1}{2} \int \frac{(x + \sqrt{x^2 - 2x - 3})^3 \left(-(x + \sqrt{x^2 - 2x - 3})^2 + 2(x + \sqrt{x^2 - 2x - 3}) + 3 \right)}{(-x - \sqrt{x^2 - 2x - 3} + 1)^2} d(x + \sqrt{x^2 - 2x - 3})$$

$$\downarrow 1195$$

$$-\frac{1}{2} \int \left(-(x + \sqrt{x^2 - 2x - 3})^3 + 4(x + \sqrt{x^2 - 2x - 3}) + \frac{12}{x + \sqrt{x^2 - 2x - 3} - 1} + \frac{4}{(x + \sqrt{x^2 - 2x - 3} - 1)^2} \right) d(x + \sqrt{x^2 - 2x - 3})$$

$$\downarrow 2009$$

$$\frac{1}{2} \left(\frac{1}{4} (\sqrt{x^2 - 2x - 3} + x)^4 - 2(\sqrt{x^2 - 2x - 3} + x)^2 - 8(\sqrt{x^2 - 2x - 3} + x) - \frac{4}{-\sqrt{x^2 - 2x - 3} - x + 1} - 12 \right)$$

input `Int[(x + Sqrt[-3 - 2*x + x^2])^3,x]`

output `(-4/(1 - x - Sqrt[-3 - 2*x + x^2]) - 8*(x + Sqrt[-3 - 2*x + x^2]) - 2*(x + Sqrt[-3 - 2*x + x^2])^2 + (x + Sqrt[-3 - 2*x + x^2])^4/4 - 12*Log[1 - x - Sqrt[-3 - 2*x + x^2]])/2`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 1195 `Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))^(n_)*((a_.) + (b_.)*(x_.) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2541 `Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_.) + (f_.)*Sqrt[(a_.) + (b_.)*(x_.) + (c_.)*(x_)^2])^(n_))^(p_.), x_Symbol] := Simp[2 Subst[Int[(g + h*x^n)^p*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)/(-2*d*e + b*f^2 + 2*e*x)^2], x], x, d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]`

Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.59

method	result
trager	$\frac{(2x^2-4x-9)x^2}{2} + (x^3 - x^2 - \frac{7}{2}x - \frac{9}{2})\sqrt{x^2 - 2x - 3} + 6 \ln(1 - x + \sqrt{x^2 - 2x - 3})$
default	$\frac{(2x-2)(x^2-2x-3)^{\frac{3}{2}}}{8} + \frac{3(2x-2)\sqrt{x^2-2x-3}}{4} - 6 \ln(\sqrt{x^2 - 2x - 3} + x - 1) + x^4 - 2x^3 - \frac{9x^2}{2} + \frac{3x(x^2-2x-3)}{4}$

input `int((x+(x^2-2*x-3)^(1/2))^3,x,method=_RETURNVERBOSE)`

output `1/2*(2*x^2-4*x-9)*x^2+(x^3-x^2-7/2*x-9/2)*(x^2-2*x-3)^(1/2)+6*ln(1-x+(x^2-2*x-3)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.59

$$\int \left(x + \sqrt{-3 - 2x + x^2} \right)^3 dx = x^4 - 2x^3 - \frac{9}{2}x^2 + \frac{1}{2}(2x^3 - 2x^2 - 7x - 9)\sqrt{x^2 - 2x - 3} + 6 \log \left(-x + \sqrt{x^2 - 2x - 3} + 1 \right)$$

input `integrate((x+(x^2-2*x-3)^(1/2))^3,x, algorithm="fricas")`

output `x^4 - 2*x^3 - 9/2*x^2 + 1/2*(2*x^3 - 2*x^2 - 7*x - 9)*sqrt(x^2 - 2*x - 3) + 6*log(-x + sqrt(x^2 - 2*x - 3) + 1)`

Sympy [A] (verification not implemented)

Time = 0.52 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.14

$$\begin{aligned} \int \left(x + \sqrt{-3 - 2x + x^2} \right)^3 dx &= x^4 - 2x^3 - \frac{9x^2}{2} - 3 \left(\frac{x}{2} - \frac{1}{2} \right) \sqrt{x^2 - 2x - 3} \\ &\quad - 2 \left(\frac{x^2}{3} - \frac{x}{6} - \frac{3}{2} \right) \sqrt{x^2 - 2x - 3} \\ &\quad + 4\sqrt{x^2 - 2x - 3} \left(\frac{x^3}{4} - \frac{x^2}{12} - \frac{7x}{12} - \frac{9}{4} \right) \\ &\quad - 6 \log \left(2x + 2\sqrt{x^2 - 2x - 3} - 2 \right) \end{aligned}$$

input `integrate((x+(x**2-2*x-3)**(1/2))**3,x)`

output

```
x**4 - 2*x**3 - 9*x**2/2 - 3*(x/2 - 1/2)*sqrt(x**2 - 2*x - 3) - 2*(x**2/3
- x/6 - 3/2)*sqrt(x**2 - 2*x - 3) + 4*sqrt(x**2 - 2*x - 3)*(x**3/4 - x**2/
12 - 7*x/12 - 9/4) - 6*log(2*x + 2*sqrt(x**2 - 2*x - 3) - 2)
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.81

$$\int \left(x + \sqrt{-3 - 2x + x^2} \right)^3 dx = x^4 - 2x^3 + (x^2 - 2x - 3)^{\frac{3}{2}}x - \frac{9}{2}x^2 + (x^2 - 2x - 3)^{\frac{3}{2}} + \frac{3}{2}\sqrt{x^2 - 2x - 3}x - \frac{3}{2}\sqrt{x^2 - 2x - 3} - 6 \log \left(2x + 2\sqrt{x^2 - 2x - 3} - 2 \right)$$

input

```
integrate((x+(x^2-2*x-3)^(1/2))^3,x, algorithm="maxima")
```

output

```
x^4 - 2*x^3 + (x^2 - 2*x - 3)^(3/2)*x - 9/2*x^2 + (x^2 - 2*x - 3)^(3/2) +
3/2*sqrt(x^2 - 2*x - 3)*x - 3/2*sqrt(x^2 - 2*x - 3) - 6*log(2*x + 2*sqrt(x
^2 - 2*x - 3) - 2)
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.57

$$\int \left(x + \sqrt{-3 - 2x + x^2} \right)^3 dx = x^4 - 2x^3 - \frac{9}{2}x^2 + \frac{1}{2} \left((2(x-1)x - 7)x - 9 \right) \sqrt{x^2 - 2x - 3} + 6 \log \left(\left| -x + \sqrt{x^2 - 2x - 3} + 1 \right| \right)$$

input

```
integrate((x+(x^2-2*x-3)^(1/2))^3,x, algorithm="giac")
```

output

```
x^4 - 2*x^3 - 9/2*x^2 + 1/2*((2*(x - 1)*x - 7)*x - 9)*sqrt(x^2 - 2*x - 3)
+ 6*log(abs(-x + sqrt(x^2 - 2*x - 3) + 1))
```

Mupad [B] (verification not implemented)

Time = 0.51 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.64

$$\int \left(x + \sqrt{-3 - 2x + x^2}\right)^3 dx = x(x^2 - 2x - 3)^{3/2} - \frac{\sqrt{x^2 - 2x - 3}(-8x^2 + 4x + 36)}{8} - 6 \ln \left(x + \sqrt{x^2 - 2x - 3} - 1\right) - \frac{9x^2}{2} - 2x^3 + x^4$$

input `int((x + (x^2 - 2*x - 3)^(1/2))^3,x)`output `x*(x^2 - 2*x - 3)^(3/2) - ((x^2 - 2*x - 3)^(1/2)*(4*x - 8*x^2 + 36))/8 - 6*log(x + (x^2 - 2*x - 3)^(1/2) - 1) - (9*x^2)/2 - 2*x^3 + x^4`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.83

$$\int \left(x + \sqrt{-3 - 2x + x^2}\right)^3 dx = \sqrt{x^2 - 2x - 3}x^3 - \sqrt{x^2 - 2x - 3}x^2 - \frac{7\sqrt{x^2 - 2x - 3}x}{2} - \frac{9\sqrt{x^2 - 2x - 3}}{2} - 6 \log \left(\frac{\sqrt{x^2 - 2x - 3}}{2} + \frac{x}{2} - \frac{1}{2}\right) + x^4 - 2x^3 - \frac{9x^2}{2}$$

input `int((x+(x^2-2*x-3)^(1/2))^3,x)`output `(2*sqrt(x**2 - 2*x - 3)*x**3 - 2*sqrt(x**2 - 2*x - 3)*x**2 - 7*sqrt(x**2 - 2*x - 3)*x - 9*sqrt(x**2 - 2*x - 3) - 12*log((sqrt(x**2 - 2*x - 3) + x - 1)/2) + 2*x**4 - 4*x**3 - 9*x**2)/2`

3.2 $\int (x + \sqrt{-3 - 2x + x^2})^2 dx$

Optimal result	53
Mathematica [A] (verified)	53
Rubi [A] (verified)	54
Maple [A] (verified)	55
Fricas [A] (verification not implemented)	56
Sympy [A] (verification not implemented)	56
Maxima [A] (verification not implemented)	57
Giac [A] (verification not implemented)	57
Mupad [B] (verification not implemented)	57
Reduce [B] (verification not implemented)	58

Optimal result

Integrand size = 16, antiderivative size = 72

$$\int (x + \sqrt{-3 - 2x + x^2})^2 dx = -3x - x^2 + \frac{2x^3}{3} - (1 - x)\sqrt{-3 - 2x + x^2} + \frac{2}{3}(-3 - 2x + x^2)^{3/2} + 4\operatorname{arctanh}\left(\frac{1 - x}{\sqrt{-3 - 2x + x^2}}\right)$$

output

```
-3*x-x^2+2/3*x^3-(1-x)*(x^2-2*x-3)^(1/2)+2/3*(x^2-2*x-3)^(3/2)+4*arctanh((1-x)/(x^2-2*x-3)^(1/2))
```

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.88

$$\int (x + \sqrt{-3 - 2x + x^2})^2 dx = \frac{1}{3}x(-9 - 3x + 2x^2) + \frac{1}{3}\sqrt{-3 - 2x + x^2}(-9 - x + 2x^2) - 8\operatorname{arctanh}\left(\frac{\sqrt{-3 - 2x + x^2}}{-3 + x}\right)$$

input

```
Integrate[(x + Sqrt[-3 - 2*x + x^2])^2, x]
```

output

```
(x*(-9 - 3*x + 2*x^2))/3 + (Sqrt[-3 - 2*x + x^2]*(-9 - x + 2*x^2))/3 - 8*ArcTanh[Sqrt[-3 - 2*x + x^2]/(-3 + x)]
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.19, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2541, 27, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\sqrt{x^2 - 2x - 3} + x)^2 dx$$

$$\downarrow 2541$$

$$2 \int -\frac{(x + \sqrt{x^2 - 2x - 3})^2 \left(-(x + \sqrt{x^2 - 2x - 3})^2 + 2(x + \sqrt{x^2 - 2x - 3}) + 3 \right)}{4(-x - \sqrt{x^2 - 2x - 3} + 1)^2} d(x + \sqrt{x^2 - 2x - 3})$$

$$\downarrow 27$$

$$-\frac{1}{2} \int \frac{(x + \sqrt{x^2 - 2x - 3})^2 \left(-(x + \sqrt{x^2 - 2x - 3})^2 + 2(x + \sqrt{x^2 - 2x - 3}) + 3 \right)}{(-x - \sqrt{x^2 - 2x - 3} + 1)^2} d(x + \sqrt{x^2 - 2x - 3})$$

$$\downarrow 1195$$

$$-\frac{1}{2} \int \left(-(x + \sqrt{x^2 - 2x - 3})^2 + \frac{8}{x + \sqrt{x^2 - 2x - 3} - 1} + \frac{4}{(x + \sqrt{x^2 - 2x - 3} - 1)^2} + 4 \right) d(x + \sqrt{x^2 - 2x - 3})$$

$$\downarrow 2009$$

$$\frac{1}{2} \left(\frac{1}{3} (\sqrt{x^2 - 2x - 3} + x)^3 - 4(\sqrt{x^2 - 2x - 3} + x) - \frac{4}{-\sqrt{x^2 - 2x - 3} - x + 1} - 8 \log(-\sqrt{x^2 - 2x - 3} - x + 1) \right)$$

input `Int[(x + Sqrt[-3 - 2*x + x^2])^2,x]`

output `(-4/(1 - x - Sqrt[-3 - 2*x + x^2]) - 4*(x + Sqrt[-3 - 2*x + x^2]) + (x + Sqrt[-3 - 2*x + x^2])^3/3 - 8*Log[1 - x - Sqrt[-3 - 2*x + x^2]])/2`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 1195 `Int[((d_) + (e_)*(x_)^(m_))*((f_) + (g_)*(x_)^(n_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2541 `Int[((g_) + (h_))*((d_) + (e_)*(x_) + (f_)*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2])^(n_)^(p_), x_Symbol] := Simp[2 Subst[Int[(g + h*x^n)^p*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)/(-2*d*e + b*f^2 + 2*e*x)^2), x], x, d + e*x + f*Sqrt[a + b*x + c*x^2], x] /; FreeQ[{a, b, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]`

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.75

method	result	size
trager	$\frac{(2x^2-3x-9)x}{3} + \left(\frac{2}{3}x^2 - \frac{1}{3}x - 3\right) \sqrt{x^2 - 2x - 3} + 4 \ln(1 - x + \sqrt{x^2 - 2x - 3})$	54
default	$\frac{2x^3}{3} - x^2 - 3x + \frac{2(x^2-2x-3)^{\frac{3}{2}}}{3} + \frac{(2x-2)\sqrt{x^2-2x-3}}{2} - 4 \ln(\sqrt{x^2 - 2x - 3} + x - 1)$	60

input `int((x+(x^2-2*x-3)^(1/2))^2,x,method=_RETURNVERBOSE)`

output `1/3*(2*x^2-3*x-9)*x+(2/3*x^2-1/3*x-3)*(x^2-2*x-3)^(1/2)+4*ln(1-x+(x^2-2*x-3)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.75

$$\int \left(x + \sqrt{-3 - 2x + x^2} \right)^2 dx = \frac{2}{3}x^3 - x^2 + \frac{1}{3}(2x^2 - x - 9)\sqrt{x^2 - 2x - 3} - 3x + 4 \log \left(-x + \sqrt{x^2 - 2x - 3} + 1 \right)$$

input `integrate((x+(x^2-2*x-3)^(1/2))^2,x, algorithm="fricas")`

output `2/3*x^3 - x^2 + 1/3*(2*x^2 - x - 9)*sqrt(x^2 - 2*x - 3) - 3*x + 4*log(-x + sqrt(x^2 - 2*x - 3) + 1)`

Sympy [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.83

$$\int \left(x + \sqrt{-3 - 2x + x^2} \right)^2 dx = \frac{2x^3}{3} - x^2 - 3x + 2 \left(\frac{x^2}{3} - \frac{x}{6} - \frac{3}{2} \right) \sqrt{x^2 - 2x - 3} - 4 \log \left(2x + 2\sqrt{x^2 - 2x - 3} - 2 \right)$$

input `integrate((x+(x**2-2*x-3)**(1/2))**2,x)`

output `2*x**3/3 - x**2 - 3*x + 2*(x**2/3 - x/6 - 3/2)*sqrt(x**2 - 2*x - 3) - 4*log(2*x + 2*sqrt(x**2 - 2*x - 3) - 2)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.97

$$\int \left(x + \sqrt{-3 - 2x + x^2} \right)^2 dx = \frac{2}{3} x^3 - x^2 + \frac{2}{3} (x^2 - 2x - 3)^{\frac{3}{2}} + \sqrt{x^2 - 2x - 3} x - 3x - \sqrt{x^2 - 2x - 3} - 4 \log \left(2x + 2\sqrt{x^2 - 2x - 3} - 2 \right)$$

input `integrate((x+(x^2-2*x-3)^(1/2))^2,x, algorithm="maxima")`output `2/3*x^3 - x^2 + 2/3*(x^2 - 2*x - 3)^(3/2) + sqrt(x^2 - 2*x - 3)*x - 3*x - sqrt(x^2 - 2*x - 3) - 4*log(2*x + 2*sqrt(x^2 - 2*x - 3) - 2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.75

$$\int \left(x + \sqrt{-3 - 2x + x^2} \right)^2 dx = \frac{2}{3} x^3 - x^2 + \frac{1}{3} ((2x - 1)x - 9) \sqrt{x^2 - 2x - 3} - 3x + 4 \log \left(\left| -x + \sqrt{x^2 - 2x - 3} + 1 \right| \right)$$

input `integrate((x+(x^2-2*x-3)^(1/2))^2,x, algorithm="giac")`output `2/3*x^3 - x^2 + 1/3*((2*x - 1)*x - 9)*sqrt(x^2 - 2*x - 3) - 3*x + 4*log(abs(-x + sqrt(x^2 - 2*x - 3) + 1))`**Mupad [B] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.72

$$\int \left(x + \sqrt{-3 - 2x + x^2} \right)^2 dx = \frac{2x^3}{3} - 4 \ln \left(x + \sqrt{x^2 - 2x - 3} - 1 \right) - \frac{\sqrt{x^2 - 2x - 3} (-8x^2 + 4x + 36)}{12} - x^2 - 3x$$

input `int((x + (x^2 - 2*x - 3)^(1/2))^2,x)`

output `(2*x^3)/3 - 4*log(x + (x^2 - 2*x - 3)^(1/2) - 1) - ((x^2 - 2*x - 3)^(1/2)*
(4*x - 8*x^2 + 36))/12 - x^2 - 3*x`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.97

$$\int \left(x + \sqrt{-3 - 2x + x^2} \right)^2 dx = \frac{2\sqrt{x^2 - 2x - 3}x^2}{3} - \frac{\sqrt{x^2 - 2x - 3}x}{3} - 3\sqrt{x^2 - 2x - 3} \\ - 4 \log \left(\frac{\sqrt{x^2 - 2x - 3}}{2} + \frac{x}{2} - \frac{1}{2} \right) + \frac{2x^3}{3} - x^2 - 3x$$

input `int((x+(x^2-2*x-3)^(1/2))^2,x)`

output `(2*sqrt(x**2 - 2*x - 3)*x**2 - sqrt(x**2 - 2*x - 3)*x - 9*sqrt(x**2 - 2*x
- 3) - 12*log((sqrt(x**2 - 2*x - 3) + x - 1)/2) + 2*x**3 - 3*x**2 - 9*x)/3`

3.3 $\int (x + \sqrt{-3 - 2x + x^2}) dx$

Optimal result	59
Mathematica [A] (verified)	59
Rubi [A] (verified)	60
Maple [A] (verified)	61
Fricas [A] (verification not implemented)	61
Sympy [A] (verification not implemented)	62
Maxima [A] (verification not implemented)	62
Giac [A] (verification not implemented)	62
Mupad [B] (verification not implemented)	63
Reduce [B] (verification not implemented)	63

Optimal result

Integrand size = 14, antiderivative size = 50

$$\int (x + \sqrt{-3 - 2x + x^2}) dx = \frac{x^2}{2} - \frac{1}{2}(1 - x)\sqrt{-3 - 2x + x^2} + 2\operatorname{arctanh}\left(\frac{1 - x}{\sqrt{-3 - 2x + x^2}}\right)$$

output `1/2*x^2-1/2*(1-x)*(x^2-2*x-3)^(1/2)+2*arctanh((1-x)/(x^2-2*x-3)^(1/2))`

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.96

$$\int (x + \sqrt{-3 - 2x + x^2}) dx = \frac{x^2}{2} + \frac{1}{2}(-1 + x)\sqrt{-3 - 2x + x^2} - 4\operatorname{arctanh}\left(\frac{\sqrt{-3 - 2x + x^2}}{-3 + x}\right)$$

input `Integrate[x + Sqrt[-3 - 2*x + x^2],x]`

output

$$x^2/2 + ((-1 + x)*\text{Sqrt}[-3 - 2*x + x^2])/2 - 4*\text{ArcTanh}[\text{Sqrt}[-3 - 2*x + x^2]/(-3 + x)]$$
Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\sqrt{x^2 - 2x - 3} + x) dx$$

↓ 2009

$$2\text{arctanh}\left(\frac{1-x}{\sqrt{x^2-2x-3}}\right) + \frac{x^2}{2} - \frac{1}{2}(1-x)\sqrt{x^2-2x-3}$$

input

$$\text{Int}[x + \text{Sqrt}[-3 - 2*x + x^2], x]$$

output

$$x^2/2 - ((1 - x)*\text{Sqrt}[-3 - 2*x + x^2])/2 + 2*\text{ArcTanh}[(1 - x)/\text{Sqrt}[-3 - 2*x + x^2]]$$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.78

method	result	size
trager	$\frac{x^2}{2} + \left(\frac{x}{2} - \frac{1}{2}\right) \sqrt{x^2 - 2x - 3} - 2 \ln(\sqrt{x^2 - 2x - 3} + x - 1)$	39
default	$\frac{x^2}{2} + \frac{(2x-2)\sqrt{x^2-2x-3}}{4} - 2 \ln(\sqrt{x^2 - 2x - 3} + x - 1)$	40
parts	$\frac{x^2}{2} + \frac{(2x-2)\sqrt{x^2-2x-3}}{4} - 2 \ln(\sqrt{x^2 - 2x - 3} + x - 1)$	40

input `int(x+(x^2-2*x-3)^(1/2),x,method=_RETURNVERBOSE)`

output `1/2*x^2+(1/2*x-1/2)*(x^2-2*x-3)^(1/2)-2*ln((x^2-2*x-3)^(1/2)+x-1)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.78

$$\int \left(x + \sqrt{-3 - 2x + x^2} \right) dx = \frac{1}{2} x^2 + \frac{1}{2} \sqrt{x^2 - 2x - 3}(x - 1) + 2 \log \left(-x + \sqrt{x^2 - 2x - 3} + 1 \right)$$

input `integrate(x+(x^2-2*x-3)^(1/2),x, algorithm="fricas")`

output `1/2*x^2 + 1/2*sqrt(x^2 - 2*x - 3)*(x - 1) + 2*log(-x + sqrt(x^2 - 2*x - 3) + 1)`

Sympy [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.88

$$\int \left(x + \sqrt{-3 - 2x + x^2} \right) dx = \frac{x^2}{2} + \left(\frac{x}{2} - \frac{1}{2} \right) \sqrt{x^2 - 2x - 3} - 2 \log \left(2x + 2\sqrt{x^2 - 2x - 3} - 2 \right)$$

input `integrate(x+(x**2-2*x-3)**(1/2),x)`

output `x**2/2 + (x/2 - 1/2)*sqrt(x**2 - 2*x - 3) - 2*log(2*x + 2*sqrt(x**2 - 2*x - 3) - 2)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.02

$$\int \left(x + \sqrt{-3 - 2x + x^2} \right) dx = \frac{1}{2} x^2 + \frac{1}{2} \sqrt{x^2 - 2x - 3} x - \frac{1}{2} \sqrt{x^2 - 2x - 3} - 2 \log \left(2x + 2\sqrt{x^2 - 2x - 3} - 2 \right)$$

input `integrate(x+(x^2-2*x-3)^(1/2),x, algorithm="maxima")`

output `1/2*x^2 + 1/2*sqrt(x^2 - 2*x - 3)*x - 1/2*sqrt(x^2 - 2*x - 3) - 2*log(2*x + 2*sqrt(x^2 - 2*x - 3) - 2)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.80

$$\int \left(x + \sqrt{-3 - 2x + x^2} \right) dx = \frac{1}{2} x^2 + \frac{1}{2} \sqrt{x^2 - 2x - 3}(x - 1) + 2 \log \left(\left| -x + \sqrt{x^2 - 2x - 3} + 1 \right| \right)$$

input `integrate(x+(x^2-2*x-3)^(1/2),x, algorithm="giac")`

output $\frac{1}{2}x^2 + \frac{1}{2}\sqrt{x^2 - 2x - 3}(x - 1) + 2\log(\text{abs}(-x + \sqrt{x^2 - 2x - 3}) + 1)$

Mupad [B] (verification not implemented)

Time = 20.97 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.76

$$\int \left(x + \sqrt{-3 - 2x + x^2} \right) dx = \left(\frac{x}{2} - \frac{1}{2} \right) \sqrt{x^2 - 2x - 3} - 2 \ln \left(x + \sqrt{x^2 - 2x - 3} - 1 \right) + \frac{x^2}{2}$$

input `int(x + (x^2 - 2*x - 3)^(1/2),x)`

output $\left(\frac{x}{2} - \frac{1}{2}\right) \cdot (x^2 - 2x - 3)^{1/2} - 2 \cdot \log(x + (x^2 - 2x - 3)^{1/2} - 1) + x^2/2$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.96

$$\int \left(x + \sqrt{-3 - 2x + x^2} \right) dx = \frac{\sqrt{x^2 - 2x - 3} x}{2} - \frac{\sqrt{x^2 - 2x - 3}}{2} - 2 \log \left(\frac{\sqrt{x^2 - 2x - 3}}{2} + \frac{x}{2} - \frac{1}{2} \right) + \frac{x^2}{2}$$

input `int(x+(x^2-2*x-3)^(1/2),x)`

output $(\sqrt{x^2 - 2x - 3} \cdot x - \sqrt{x^2 - 2x - 3} - 4 \cdot \log((\sqrt{x^2 - 2x - 3} + x - 1)/2) + x^2)/2$

3.4 $\int \frac{1}{x + \sqrt{-3 - 2x + x^2}} dx$

Optimal result	64
Mathematica [A] (verified)	64
Rubi [A] (verified)	65
Maple [A] (verified)	66
Fricas [A] (verification not implemented)	67
Sympy [F]	67
Maxima [F]	68
Giac [A] (verification not implemented)	68
Mupad [F(-1)]	69
Reduce [B] (verification not implemented)	69

Optimal result

Integrand size = 16, antiderivative size = 65

$$\int \frac{1}{x + \sqrt{-3 - 2x + x^2}} dx = -\frac{2}{1 - x - \sqrt{-3 - 2x + x^2}} + 2 \log \left(1 - x - \sqrt{-3 - 2x + x^2} \right) - \frac{3}{2} \log \left(x + \sqrt{-3 - 2x + x^2} \right)$$

output
$$-2/(1-x-(x^2-2*x-3)^{(1/2)})+2*\ln(1-x-(x^2-2*x-3)^{(1/2)})-3/2*\ln(x+(x^2-2*x-3)^{(1/2)})$$

Mathematica [A] (verified)

Time = 0.32 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.20

$$\int \frac{1}{x + \sqrt{-3 - 2x + x^2}} dx = \frac{1}{2} \left(x - \sqrt{-3 - 2x + x^2} - \log \left(-1 - x + \sqrt{-3 - 2x + x^2} \right) + 4 \log \left(1 + x + \sqrt{-3 - 2x + x^2} \right) - 3 \log \left(3 + 3x + \sqrt{-3 - 2x + x^2} \right) \right)$$

input
$$\text{Integrate}[(x + \text{Sqrt}[-3 - 2*x + x^2])^{-1}, x]$$

output

$$(x - \text{Sqrt}[-3 - 2*x + x^2] - \text{Log}[-1 - x + \text{Sqrt}[-3 - 2*x + x^2]] + 4*\text{Log}[1 + x + \text{Sqrt}[-3 - 2*x + x^2]] - 3*\text{Log}[3 + 3*x + \text{Sqrt}[-3 - 2*x + x^2]])/2$$
Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.03, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2541, 27, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x^2 - 2x - 3} + x} dx$$

↓ 2541

$$2 \int -\frac{(x + \sqrt{x^2 - 2x - 3})^2 + 2(x + \sqrt{x^2 - 2x - 3}) + 3}{4(-x - \sqrt{x^2 - 2x - 3} + 1)^2(x + \sqrt{x^2 - 2x - 3})} d(x + \sqrt{x^2 - 2x - 3})$$

↓ 27

$$-\frac{1}{2} \int \frac{(x + \sqrt{x^2 - 2x - 3})^2 + 2(x + \sqrt{x^2 - 2x - 3}) + 3}{(-x - \sqrt{x^2 - 2x - 3} + 1)^2(x + \sqrt{x^2 - 2x - 3})} d(x + \sqrt{x^2 - 2x - 3})$$

↓ 1195

$$-\frac{1}{2} \int \left(\frac{3}{x + \sqrt{x^2 - 2x - 3}} - \frac{4}{x + \sqrt{x^2 - 2x - 3} - 1} + \frac{4}{(x + \sqrt{x^2 - 2x - 3} - 1)^2} \right) d(x + \sqrt{x^2 - 2x - 3})$$

↓ 2009

$$\frac{1}{2} \left(-\frac{4}{-\sqrt{x^2 - 2x - 3} - x + 1} + 4 \log(-\sqrt{x^2 - 2x - 3} - x + 1) - 3 \log(\sqrt{x^2 - 2x - 3} + x) \right)$$

input

$$\text{Int}[(x + \text{Sqrt}[-3 - 2*x + x^2])^{-1}, x]$$

```
output (-4/(1 - x - Sqrt[-3 - 2*x + x^2]) + 4*Log[1 - x - Sqrt[-3 - 2*x + x^2]] - 3*Log[x + Sqrt[-3 - 2*x + x^2]])/2
```

Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 1195 Int[((d_) + (e_)*(x_)^(m_))*((f_) + (g_)*(x_)^(n_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2541 Int[((g_) + (h_))*((d_) + (e_)*(x_) + (f_)*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2])^(n_))^(p_), x_Symbol] := Simp[2 Subst[Int[(g + h*x^n)^p*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)/(-2*d*e + b*f^2 + 2*e*x)^2), x], x, d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]
```

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.09

method	result	size
default	$-\frac{\sqrt{4(x+\frac{3}{2})^2-20x-21}}{4} + \frac{5 \ln\left(-1+x+\sqrt{(x+\frac{3}{2})^2-5x-\frac{21}{4}}\right)}{4} + \frac{3 \operatorname{arctanh}\left(\frac{-2-\frac{10x}{3}}{\sqrt{4(x+\frac{3}{2})^2-20x-21}}\right)}{4} + \frac{x}{2} - \frac{3 \ln(2x+3)}{4}$	71
trager	$\frac{x}{2} - \frac{\sqrt{x^2-2x-3}}{2} - \frac{\ln\left(\sqrt{x^2-2x-3}x^3-x^4+3x^2\sqrt{x^2-2x-3}-2x^3+\sqrt{x^2-2x-3}x+4x^2-3\sqrt{x^2-2x-3}+12x+9\right)}{2}$	93

```
input int(1/(x+(x^2-2*x-3)^(1/2)),x,method=_RETURNVERBOSE)
```

output

```
-1/4*(4*(x+3/2)^2-20*x-21)^(1/2)+5/4*ln(-1+x+((x+3/2)^2-5*x-21/4)^(1/2))+3/4*arctanh(2/3*(-3-5*x)/(4*(x+3/2)^2-20*x-21)^(1/2))+1/2*x-3/4*ln(2*x+3)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.18

$$\int \frac{1}{x + \sqrt{-3 - 2x + x^2}} dx = \frac{1}{2}x - \frac{1}{2}\sqrt{x^2 - 2x - 3} - \frac{3}{4}\log(2x + 3) - \frac{5}{4}\log(-x + \sqrt{x^2 - 2x - 3} + 1) + \frac{3}{4}\log(-x + \sqrt{x^2 - 2x - 3}) - \frac{3}{4}\log(-x + \sqrt{x^2 - 2x - 3} - 3)$$

input

```
integrate(1/(x+(x^2-2*x-3)^(1/2)),x, algorithm="fricas")
```

output

```
1/2*x - 1/2*sqrt(x^2 - 2*x - 3) - 3/4*log(2*x + 3) - 5/4*log(-x + sqrt(x^2 - 2*x - 3) + 1) + 3/4*log(-x + sqrt(x^2 - 2*x - 3)) - 3/4*log(-x + sqrt(x^2 - 2*x - 3) - 3)
```

Sympy [F]

$$\int \frac{1}{x + \sqrt{-3 - 2x + x^2}} dx = \int \frac{1}{x + \sqrt{x^2 - 2x - 3}} dx$$

input

```
integrate(1/(x+(x**2-2*x-3)**(1/2)),x)
```

output

```
Integral(1/(x + sqrt(x**2 - 2*x - 3)), x)
```

Maxima [F]

$$\int \frac{1}{x + \sqrt{-3 - 2x + x^2}} dx = \int \frac{1}{x + \sqrt{x^2 - 2x - 3}} dx$$

input `integrate(1/(x+(x^2-2*x-3)^(1/2)),x, algorithm="maxima")`

output `integrate(1/(x + sqrt(x^2 - 2*x - 3)), x)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.25

$$\begin{aligned} \int \frac{1}{x + \sqrt{-3 - 2x + x^2}} dx &= \frac{1}{2}x - \frac{1}{2}\sqrt{x^2 - 2x - 3} - \frac{3}{4}\log(|2x + 3|) \\ &\quad - \frac{5}{4}\log\left(\left|-x + \sqrt{x^2 - 2x - 3} + 1\right|\right) \\ &\quad + \frac{3}{4}\log\left(\left|-x + \sqrt{x^2 - 2x - 3}\right|\right) \\ &\quad - \frac{3}{4}\log\left(\left|-x + \sqrt{x^2 - 2x - 3} - 3\right|\right) \end{aligned}$$

input `integrate(1/(x+(x^2-2*x-3)^(1/2)),x, algorithm="giac")`

output `1/2*x - 1/2*sqrt(x^2 - 2*x - 3) - 3/4*log(abs(2*x + 3)) - 5/4*log(abs(-x + sqrt(x^2 - 2*x - 3) + 1)) + 3/4*log(abs(-x + sqrt(x^2 - 2*x - 3))) - 3/4*log(abs(-x + sqrt(x^2 - 2*x - 3) - 3))`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{x + \sqrt{-3 - 2x + x^2}} dx = \frac{x}{2} - \frac{3 \ln\left(x + \frac{3}{2}\right)}{4} - \int \frac{\sqrt{x^2 - 2x - 3}}{2x + 3} dx$$

input `int(1/(x + (x^2 - 2*x - 3)^(1/2)),x)`output `x/2 - (3*log(x + 3/2))/4 - int((x^2 - 2*x - 3)^(1/2)/(2*x + 3), x)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.75

$$\int \frac{1}{x + \sqrt{-3 - 2x + x^2}} dx = -\frac{\sqrt{x^2 - 2x - 3}}{2} - \frac{3 \log(\sqrt{x^2 - 2x - 3} + x)}{2} + 2 \log\left(\frac{\sqrt{x^2 - 2x - 3}}{2} + \frac{x}{2} - \frac{1}{2}\right) + \frac{x}{2} - \frac{1}{2}$$

input `int(1/(x+(x^2-2*x-3)^(1/2)),x)`output `(- sqrt(x**2 - 2*x - 3) - 3*log(sqrt(x**2 - 2*x - 3) + x) + 4*log((sqrt(x**2 - 2*x - 3) + x - 1)/2) + x - 1)/2`

3.5
$$\int \frac{1}{\left(x + \sqrt{-3 - 2x + x^2}\right)^2} dx$$

Optimal result	70
Mathematica [A] (verified)	70
Rubi [A] (verified)	71
Maple [A] (verified)	72
Fricas [A] (verification not implemented)	73
Sympy [F]	73
Maxima [F]	74
Giac [B] (verification not implemented)	74
Mupad [F(-1)]	75
Reduce [B] (verification not implemented)	75

Optimal result

Integrand size = 16, antiderivative size = 83

$$\int \frac{1}{\left(x + \sqrt{-3 - 2x + x^2}\right)^2} dx = -\frac{2}{1 - x - \sqrt{-3 - 2x + x^2}} + \frac{3}{2\left(x + \sqrt{-3 - 2x + x^2}\right)} + 4 \log\left(1 - x - \sqrt{-3 - 2x + x^2}\right) - 4 \log\left(x + \sqrt{-3 - 2x + x^2}\right)$$

output `-2/(1-x-(x^2-2*x-3)^(1/2))+3/(2*x+2*(x^2-2*x-3)^(1/2))+4*ln(1-x-(x^2-2*x-3)^(1/2))-4*ln(x+(x^2-2*x-3)^(1/2))`

Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.83

$$\int \frac{1}{\left(x + \sqrt{-3 - 2x + x^2}\right)^2} dx = \frac{-9 + 6x + 4x^2 - 4(3 + x)\sqrt{-3 - 2x + x^2} - 32(3 + 2x)\operatorname{arctanh}\left(\frac{1+x}{2+2x+\sqrt{-3-2x+x^2}}\right)}{4(3 + 2x)}$$

input `Integrate[(x + Sqrt[-3 - 2*x + x^2])^(-2), x]`

output `(-9 + 6*x + 4*x^2 - 4*(3 + x)*Sqrt[-3 - 2*x + x^2] - 32*(3 + 2*x)*ArcTanh[(1 + x)/(2 + 2*x + Sqrt[-3 - 2*x + x^2])])/(4*(3 + 2*x))`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2541, 27, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(\sqrt{x^2 - 2x - 3} + x)^2} dx$$

$$\downarrow 2541$$

$$2 \int -\frac{-(x + \sqrt{x^2 - 2x - 3})^2 + 2(x + \sqrt{x^2 - 2x - 3}) + 3}{4(-x - \sqrt{x^2 - 2x - 3} + 1)^2 (x + \sqrt{x^2 - 2x - 3})^2} d(x + \sqrt{x^2 - 2x - 3})$$

$$\downarrow 27$$

$$-\frac{1}{2} \int \frac{-(x + \sqrt{x^2 - 2x - 3})^2 + 2(x + \sqrt{x^2 - 2x - 3}) + 3}{(-x - \sqrt{x^2 - 2x - 3} + 1)^2 (x + \sqrt{x^2 - 2x - 3})^2} d(x + \sqrt{x^2 - 2x - 3})$$

$$\downarrow 1195$$

$$-\frac{1}{2} \int \left(\frac{8}{x + \sqrt{x^2 - 2x - 3}} + \frac{3}{(x + \sqrt{x^2 - 2x - 3})^2} - \frac{8}{x + \sqrt{x^2 - 2x - 3} - 1} + \frac{4}{(x + \sqrt{x^2 - 2x - 3} - 1)^2} \right) d(x + \sqrt{x^2 - 2x - 3})$$

$$\downarrow 2009$$

$$\frac{1}{2} \left(-\frac{4}{-\sqrt{x^2 - 2x - 3} - x + 1} + \frac{3}{\sqrt{x^2 - 2x - 3} + x} + 8 \log(-\sqrt{x^2 - 2x - 3} - x + 1) - 8 \log(\sqrt{x^2 - 2x - 3} + x) \right)$$

input `Int[(x + Sqrt[-3 - 2*x + x^2])^(-2), x]`

output `(-4/(1 - x - Sqrt[-3 - 2*x + x^2]) + 3/(x + Sqrt[-3 - 2*x + x^2]) + 8*Log[1 - x - Sqrt[-3 - 2*x + x^2]] - 8*Log[x + Sqrt[-3 - 2*x + x^2]])/2`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 1195 `Int[((d_) + (e_)*(x_)^(m_))*((f_) + (g_)*(x_)^(n_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2541 `Int[((g_) + (h_))*((d_) + (e_)*(x_) + (f_)*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2])^(n_)^(p_), x_Symbol] := Simp[2 Subst[Int[(g + h*x^n)^p*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)/(-2*d*e + b*f^2 + 2*e*x)^2), x], x, d + e*x + f*Sqrt[a + b*x + c*x^2], x] /; FreeQ[{a, b, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.73

method	result
trager	$\frac{(3+x)x}{2x+3} - \frac{(3+x)\sqrt{x^2-2x-3}}{2x+3} + 4 \ln \left(-\frac{\sqrt{x^2-2x-3}+3+x}{2x+3} \right)$
default	$-2 \ln(2x+3) + \frac{x}{2} - \frac{9}{4(2x+3)} - \frac{\left((x+\frac{3}{2})^2 - 5x - \frac{21}{4}\right)^{\frac{3}{2}}}{3(x+\frac{3}{2})} - \frac{2\sqrt{4(x+\frac{3}{2})^2 - 20x - 21}}{3} + 2 \ln \left(-1 + x + \sqrt{(x+\frac{3}{2})^2 - 5x - \frac{21}{4}} \right)$

input `int(1/(x+(x^2-2*x-3)^(1/2))^2,x,method=_RETURNVERBOSE)`

output `(3+x)*x/(2*x+3)-(3+x)/(2*x+3)*(x^2-2*x-3)^(1/2)+4*ln(-(x^2-2*x-3)^(1/2)+x)/(2*x+3)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.17

$$\int \frac{1}{(x + \sqrt{-3 - 2x + x^2})^2} dx$$

$$= \frac{4x^2 - 8(2x + 3) \log(x^2 - \sqrt{x^2 - 2x - 3}(x + 1) - 3) - 8(2x + 3) \log(2x + 3) + 8(2x + 3) \log(-x + \sqrt{x^2 - 2x - 3})}{4(2x + 3)}$$

input `integrate(1/(x+(x^2-2*x-3)^(1/2))^2,x, algorithm="fricas")`

output `1/4*(4*x^2 - 8*(2*x + 3)*log(x^2 - sqrt(x^2 - 2*x - 3)*(x + 1) - 3) - 8*(2*x + 3)*log(2*x + 3) + 8*(2*x + 3)*log(-x + sqrt(x^2 - 2*x - 3)) - 4*sqrt(x^2 - 2*x - 3)*(x + 3) + 2*x - 15)/(2*x + 3)`

Sympy [F]

$$\int \frac{1}{(x + \sqrt{-3 - 2x + x^2})^2} dx = \int \frac{1}{(x + \sqrt{x^2 - 2x - 3})^2} dx$$

input `integrate(1/(x+(x**2-2*x-3)**(1/2))**2,x)`

output `Integral((x + sqrt(x**2 - 2*x - 3))**(-2), x)`

Maxima [F]

$$\int \frac{1}{(x + \sqrt{-3 - 2x + x^2})^2} dx = \int \frac{1}{(x + \sqrt{x^2 - 2x - 3})^2} dx$$

input `integrate(1/(x+(x^2-2*x-3)^(1/2))^2,x, algorithm="maxima")`

output `integrate((x + sqrt(x^2 - 2*x - 3))^(-2), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 143 vs. 2(69) = 138.

Time = 0.12 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.72

$$\int \frac{1}{(x + \sqrt{-3 - 2x + x^2})^2} dx = \frac{1}{2}x - \frac{1}{2}\sqrt{x^2 - 2x - 3} - \frac{3(5x - 5\sqrt{x^2 - 2x - 3} + 3)}{4((x - \sqrt{x^2 - 2x - 3})^2 + 3x - 3\sqrt{x^2 - 2x - 3})} - \frac{9}{4(2x + 3)} - 2 \log(|2x + 3|) - 2 \log\left(\left|-x + \sqrt{x^2 - 2x - 3} + 1\right|\right) + 2 \log\left(\left|-x + \sqrt{x^2 - 2x - 3}\right|\right) - 2 \log\left(\left|-x + \sqrt{x^2 - 2x - 3} - 3\right|\right)$$

input `integrate(1/(x+(x^2-2*x-3)^(1/2))^2,x, algorithm="giac")`

output `1/2*x - 1/2*sqrt(x^2 - 2*x - 3) - 3/4*(5*x - 5*sqrt(x^2 - 2*x - 3) + 3)/((x - sqrt(x^2 - 2*x - 3))^2 + 3*x - 3*sqrt(x^2 - 2*x - 3)) - 9/4/(2*x + 3) - 2*log(abs(2*x + 3)) - 2*log(abs(-x + sqrt(x^2 - 2*x - 3) + 1)) + 2*log(abs(-x + sqrt(x^2 - 2*x - 3))) - 2*log(abs(-x + sqrt(x^2 - 2*x - 3) - 3))`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(x + \sqrt{-3 - 2x + x^2})^2} dx = \int \frac{1}{(x + \sqrt{x^2 - 2x - 3})^2} dx$$

input `int(1/(x + (x^2 - 2*x - 3)^(1/2))^2,x)`output `int(1/(x + (x^2 - 2*x - 3)^(1/2))^2, x)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.29

$$\int \frac{1}{(x + \sqrt{-3 - 2x + x^2})^2} dx$$

$$= \frac{-\sqrt{x^2 - 2x - 3} x - 3\sqrt{x^2 - 2x - 3} - 8 \log(\sqrt{x^2 - 2x - 3} + x) x - 12 \log(\sqrt{x^2 - 2x - 3} + x) + 8 \log(\sqrt{x^2 - 2x - 3} + x)}{2x + 3}$$

input `int(1/(x+(x^2-2*x-3)^(1/2))^2,x)`output `(- sqrt(x**2 - 2*x - 3)*x - 3*sqrt(x**2 - 2*x - 3) - 8*log(sqrt(x**2 - 2*x - 3) + x)*x - 12*log(sqrt(x**2 - 2*x - 3) + x) + 8*log((sqrt(x**2 - 2*x - 3) + x - 1)/2)*x + 12*log((sqrt(x**2 - 2*x - 3) + x - 1)/2) + x**2 - 5*x - 12)/(2*x + 3)`

3.6
$$\int \frac{1}{\left(x + \sqrt{-3 - 2x + x^2}\right)^3} dx$$

Optimal result	76
Mathematica [A] (verified)	76
Rubi [A] (verified)	77
Maple [A] (verified)	79
Fricas [A] (verification not implemented)	79
Sympy [F]	80
Maxima [F]	80
Giac [B] (verification not implemented)	80
Mupad [F(-1)]	81
Reduce [B] (verification not implemented)	81

Optimal result

Integrand size = 16, antiderivative size = 101

$$\int \frac{1}{\left(x + \sqrt{-3 - 2x + x^2}\right)^3} dx = -\frac{2}{1 - x - \sqrt{-3 - 2x + x^2}} + \frac{3}{4\left(x + \sqrt{-3 - 2x + x^2}\right)^2} + \frac{4}{x + \sqrt{-3 - 2x + x^2}} + 6 \log\left(1 - x - \sqrt{-3 - 2x + x^2}\right) - 6 \log\left(x + \sqrt{-3 - 2x + x^2}\right)$$

output

$$-2/(1-x-(x^2-2*x-3)^(1/2))+3/4/(x+(x^2-2*x-3)^(1/2))^2+4/(x+(x^2-2*x-3)^(1/2))+6*\ln(1-x-(x^2-2*x-3)^(1/2))-6*\ln(x+(x^2-2*x-3)^(1/2))$$

Mathematica [A] (verified)

Time = 0.39 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.82

$$\int \frac{1}{\left(x + \sqrt{-3 - 2x + x^2}\right)^3} dx = \frac{189 + 108x - 48x^2 - 16x^3 + 4\sqrt{-3 - 2x + x^2}(33 + 31x + 4x^2) + 96(3 + 2x)^2 \operatorname{arctanh}\left(\frac{1+x}{2+2x+\sqrt{-3-2x+x^2}}\right)}{8(3 + 2x)^2}$$

input `Integrate[(x + Sqrt[-3 - 2*x + x^2])^(-3), x]`

output
$$-1/8*(189 + 108*x - 48*x^2 - 16*x^3 + 4*\text{Sqrt}[-3 - 2*x + x^2]*(33 + 31*x + 4*x^2) + 96*(3 + 2*x)^2*\text{ArcTanh}[(1 + x)/(2 + 2*x + \text{Sqrt}[-3 - 2*x + x^2])]) / (3 + 2*x)^2$$

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.04, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2541, 27, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(\sqrt{x^2 - 2x - 3} + x)^3} dx$$

↓ 2541

$$2 \int -\frac{(x + \sqrt{x^2 - 2x - 3})^2 + 2(x + \sqrt{x^2 - 2x - 3}) + 3}{4(-x - \sqrt{x^2 - 2x - 3} + 1)^2 (x + \sqrt{x^2 - 2x - 3})^3} d(x + \sqrt{x^2 - 2x - 3})$$

↓ 27

$$-\frac{1}{2} \int \frac{(x + \sqrt{x^2 - 2x - 3})^2 + 2(x + \sqrt{x^2 - 2x - 3}) + 3}{(-x - \sqrt{x^2 - 2x - 3} + 1)^2 (x + \sqrt{x^2 - 2x - 3})^3} d(x + \sqrt{x^2 - 2x - 3})$$

↓ 1195

$$-\frac{1}{2} \int \left(\frac{12}{x + \sqrt{x^2 - 2x - 3}} + \frac{8}{(x + \sqrt{x^2 - 2x - 3})^2} + \frac{3}{(x + \sqrt{x^2 - 2x - 3})^3} - \frac{12}{x + \sqrt{x^2 - 2x - 3} - 1} + \frac{1}{x + \sqrt{x^2 - 2x - 3} - 1} \right) dx$$

↓ 2009

$$\frac{1}{2} \left(-\frac{4}{-\sqrt{x^2 - 2x - 3} - x + 1} + \frac{8}{\sqrt{x^2 - 2x - 3} + x} + \frac{3}{2(\sqrt{x^2 - 2x - 3} + x)^2} + 12 \log(-\sqrt{x^2 - 2x - 3} - x + 1) \right)$$

input `Int[(x + Sqrt[-3 - 2*x + x^2])^(-3), x]`

output `(-4/(1 - x - Sqrt[-3 - 2*x + x^2]) + 3/(2*(x + Sqrt[-3 - 2*x + x^2])^2) + 8/(x + Sqrt[-3 - 2*x + x^2]) + 12*Log[1 - x - Sqrt[-3 - 2*x + x^2]] - 12*Log[x + Sqrt[-3 - 2*x + x^2]])/2`

Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 1195 `Int[((d_) + (e_)*(x_)^(m_))*((f_) + (g_)*(x_)^(n_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2541 `Int[((g_) + (h_))*((d_) + (e_)*(x_) + (f_)*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2])^(n_)]^(p_), x_Symbol] := Simp[2 Subst[Int[(g + h*x^n)^p*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)/(-2*d*e + b*f^2 + 2*e*x^2), x], x, d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.68

method	result
trager	$\frac{(4x^2+33x+36)x}{2(2x+3)^2} - \frac{(4x^2+31x+33)\sqrt{x^2-2x-3}}{2(2x+3)^2} - 6 \ln(3+x-\sqrt{x^2-2x-3})$
default	$-\frac{9}{2x+3} - 3 \ln(2x+3) + \frac{x}{2} + \frac{27}{8(2x+3)^2} + \frac{\left((x+\frac{3}{2})^2-5x-\frac{21}{4}\right)^{\frac{3}{2}}}{4(x+\frac{3}{2})^2} - \frac{\left((x+\frac{3}{2})^2-5x-\frac{21}{4}\right)^{\frac{3}{2}}}{2(x+\frac{3}{2})} - \sqrt{4\left(x+\frac{3}{2}\right)^2-2}$

input `int(1/(x+(x^2-2*x-3)^(1/2))^3,x,method=_RETURNVERBOSE)`

output $\frac{1}{2}*(4*x^2+33*x+36)*x/(2*x+3)^2-1/2*(4*x^2+31*x+33)/(2*x+3)^2*(x^2-2*x-3)^(1/2)-6*\ln(3+x-(x^2-2*x-3)^(1/2))$

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.28

$$\int \frac{1}{(x + \sqrt{-3 - 2x + x^2})^3} dx$$

$$= \frac{8x^3 - 10x^2 - 12(4x^2 + 12x + 9) \log(x^2 - \sqrt{x^2 - 2x - 3}(x+1) - 3) - 12(4x^2 + 12x + 9) \log(2x + (x+1) - 3) - 12(4x^2 + 12x + 9) \log(-x + \sqrt{x^2 - 2x - 3}) - 2*(4*x^2 + 31*x + 33)*\sqrt{x^2 - 2*x - 3} - 156*x - 171}{4(4x^2 + 12x + 9)}$$

input `integrate(1/(x+(x^2-2*x-3)^(1/2))^3,x, algorithm="fricas")`

output $\frac{1}{4}*(8*x^3 - 10*x^2 - 12*(4*x^2 + 12*x + 9)*\log(x^2 - \sqrt{x^2 - 2*x - 3}*(x + 1) - 3) - 12*(4*x^2 + 12*x + 9)*\log(2*x + (x + 1) - 3) + 12*(4*x^2 + 12*x + 9)*\log(-x + \sqrt{x^2 - 2*x - 3}) - 2*(4*x^2 + 31*x + 33)*\sqrt{x^2 - 2*x - 3} - 156*x - 171)/(4*x^2 + 12*x + 9)$

Sympy [F]

$$\int \frac{1}{(x + \sqrt{-3 - 2x + x^2})^3} dx = \int \frac{1}{(x + \sqrt{x^2 - 2x - 3})^3} dx$$

input `integrate(1/(x+(x**2-2*x-3)**(1/2))**3,x)`

output `Integral((x + sqrt(x**2 - 2*x - 3))**(-3), x)`

Maxima [F]

$$\int \frac{1}{(x + \sqrt{-3 - 2x + x^2})^3} dx = \int \frac{1}{(x + \sqrt{x^2 - 2x - 3})^3} dx$$

input `integrate(1/(x+(x^2-2*x-3)^(1/2))^3,x, algorithm="maxima")`

output `integrate((x + sqrt(x^2 - 2*x - 3))^-3, x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 184 vs. 2(85) = 170.

Time = 0.12 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.82

$$\int \frac{1}{(x + \sqrt{-3 - 2x + x^2})^3} dx = \frac{1}{2}x - \frac{1}{2}\sqrt{x^2 - 2x - 3} - \frac{104(x - \sqrt{x^2 - 2x - 3})^3 + 315(x - \sqrt{x^2 - 2x - 3})^2 + 162x - 162\sqrt{x^2 - 2x - 3} + 27}{8((x - \sqrt{x^2 - 2x - 3})^2 + 3x - 3\sqrt{x^2 - 2x - 3})^2} - \frac{9(16x + 21)}{8(2x + 3)^2} - 3 \log(|2x + 3|) - 3 \log\left(\left|-x + \sqrt{x^2 - 2x - 3} + 1\right|\right) + 3 \log\left(\left|-x + \sqrt{x^2 - 2x - 3}\right|\right) - 3 \log\left(\left|-x + \sqrt{x^2 - 2x - 3} - 3\right|\right)$$

input `integrate(1/(x+(x^2-2*x-3)^(1/2))^3,x, algorithm="giac")`

output $\frac{1}{2}x - \frac{1}{2}\sqrt{x^2 - 2x - 3} - \frac{1}{8}(104(x - \sqrt{x^2 - 2x - 3})^3 + 315(x - \sqrt{x^2 - 2x - 3})^2 + 162x - 162\sqrt{x^2 - 2x - 3} + 27)/((x - \sqrt{x^2 - 2x - 3})^2 + 3x - 3\sqrt{x^2 - 2x - 3})^2 - \frac{9}{8}(16x + 21)/(2x + 3)^2 - 3\log(\text{abs}(2x + 3)) - 3\log(\text{abs}(-x + \sqrt{x^2 - 2x - 3} + 1)) + 3\log(\text{abs}(-x + \sqrt{x^2 - 2x - 3})) - 3\log(\text{abs}(-x + \sqrt{x^2 - 2x - 3} - 3))$

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(x + \sqrt{-3 - 2x + x^2})^3} dx = \int \frac{1}{(x + \sqrt{x^2 - 2x - 3})^3} dx$$

input `int(1/(x + (x^2 - 2*x - 3)^(1/2))^3,x)`

output `int(1/(x + (x^2 - 2*x - 3)^(1/2))^3, x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.70

$$\int \frac{1}{(x + \sqrt{-3 - 2x + x^2})^3} dx$$

$$= \frac{-8\sqrt{x^2 - 2x - 3}x^2 - 62\sqrt{x^2 - 2x - 3}x - 66\sqrt{x^2 - 2x - 3} - 96\log(\sqrt{x^2 - 2x - 3} + x)x^2 - 288\log(\sqrt{x^2 - 2x - 3} + x)}{\dots}$$

input `int(1/(x+(x^2-2*x-3)^(1/2))^3,x)`

output

```
( - 8*sqrt(x**2 - 2*x - 3)*x**2 - 62*sqrt(x**2 - 2*x - 3)*x - 66*sqrt(x**2
- 2*x - 3) - 96*log(sqrt(x**2 - 2*x - 3) + x)*x**2 - 288*log(sqrt(x**2 -
2*x - 3) + x)*x - 216*log(sqrt(x**2 - 2*x - 3) + x) + 96*log((sqrt(x**2 -
2*x - 3) + x - 1)/2)*x**2 + 288*log((sqrt(x**2 - 2*x - 3) + x - 1)/2)*x +
216*log((sqrt(x**2 - 2*x - 3) + x - 1)/2) + 8*x**3 + 6*x**2 - 108*x - 135)
/(4*(4*x**2 + 12*x + 9))
```

$$3.7 \quad \int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^3 dx$$

Optimal result	83
Mathematica [A] (verified)	84
Rubi [A] (verified)	85
Maple [B] (verified)	86
Fricas [A] (verification not implemented)	87
Sympy [A] (verification not implemented)	88
Maxima [F(-2)]	88
Giac [A] (verification not implemented)	89
Mupad [F(-1)]	90
Reduce [B] (verification not implemented)	90

Optimal result

Integrand size = 28, antiderivative size = 303

$$\begin{aligned} & \int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^3 dx \\ &= \frac{f^2(2de - bf^2)(4ae^2 - b^2f^2) \left(ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)}{8e^4} \\ &+ \frac{f^2(4ae^2 - b^2f^2) \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^2}{16e^3} \\ &+ \frac{\left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^4}{8e} - \frac{f^2(2de - bf^2)^3(4ae^2 - b^2f^2)}{32e^5 \left(bf^2 + 2e \left(ex + f \sqrt{a + \frac{x(bf^2 + e^2x)}{f^2}} \right) \right)} \\ &+ \frac{3f^2(2de - bf^2)^2(4ae^2 - b^2f^2) \log \left(bf^2 + 2e \left(ex + f \sqrt{a + \frac{x(bf^2 + e^2x)}{f^2}} \right) \right)}{32e^5} \end{aligned}$$

output

```
1/8*f^2*(-b*f^2+2*d*e)*(-b^2*f^2+4*a*e^2)*(e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2)
)/e^4+1/16*f^2*(-b^2*f^2+4*a*e^2)*(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^2/e^
3+1/8*(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^4/e-1/32*f^2*(-b*f^2+2*d*e)^3*(-
b^2*f^2+4*a*e^2)/e^5/(b*f^2+2*e*(e*x+f*(a+x*(b*f^2+e^2*x)/f^2)^(1/2)))+3/3
2*f^2*(-b*f^2+2*d*e)^2*(-b^2*f^2+4*a*e^2)*ln(b*f^2+2*e*(e*x+f*(a+x*(b*f^2+
e^2*x)/f^2)^(1/2)))/e^5
```

Mathematica [A] (verified)

Time = 1.92 (sec) , antiderivative size = 260, normalized size of antiderivative = 0.86

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^3 dx$$

$$= \frac{1}{16} \left(8x(2d^3 + 3d^2 ex + ex(3af^2 + 2x(bf^2 + e^2 x)) + d(6af^2 + x(3bf^2 + 4e^2 x))) \right. \\ \left. + \frac{\sqrt{a + x \left(b + \frac{e^2 x}{f^2} \right)} (3b^3 f^7 - 2b^2 e f^5 (6d + ex) + 4be^2 f^3 (3d^2 - 2af^2 + 2dex + 2e^2 x^2) + 8e^3 f (2af^2 (2d + \right. \\ \left. \frac{3(4ae^2 - b^2 f^2) (-2def + bf^3)^2 \operatorname{arctanh} \left(\frac{ex}{f(-\sqrt{a + \frac{e^2 x}{f^2}} + \sqrt{a + x(b + \frac{e^2 x}{f^2})})} \right))}{e^4}} \right)}{e^5} \right)$$

input

```
Integrate[(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^3,x]
```

output

```
(8*x*(2*d^3 + 3*d^2*e*x + e*x*(3*a*f^2 + 2*x*(b*f^2 + e^2*x)) + d*(6*a*f^2
+ x*(3*b*f^2 + 4*e^2*x))) + (Sqrt[a + x*(b + (e^2*x)/f^2)]*(3*b^3*f^7 - 2
*b^2*e*f^5*(6*d + e*x) + 4*b*e^2*f^3*(3*d^2 - 2*a*f^2 + 2*d*e*x + 2*e^2*x^
2) + 8*e^3*f*(2*a*f^2*(2*d + e*x) + e*x*(3*d^2 + 4*d*e*x + 2*e^2*x^2))))/e
^4 + (3*(4*a*e^2 - b^2*f^2)*(-2*d*e*f + b*f^3)^2*ArcTanh[(e*x)/(f*(-Sqrt[a
] + Sqrt[a + x*(b + (e^2*x)/f^2])]))]/e^5)/16
```

Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 312, normalized size of antiderivative = 1.03, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$, Rules used = {2541, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} + d + ex \right)^3 dx$$

↓ 2541

$$2 \int \frac{\left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)^3 \left(ed^2 - bf^2 d + aef^2 + e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)^2 - (2de - bf^2) \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right) \right)}{\left(-bf^2 + 2de - 2e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right) \right)^2} dx$$

↓ 1195

$$2 \int \left(\frac{f^2 (4ae^2 - b^2 f^2) (2de - bf^2)^3}{32e^4 \left(-bf^2 + 2de - 2e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right) \right)^2} + \frac{f^2 (4ae^2 - b^2 f^2) (2de - bf^2)}{16e^4} + \frac{\left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)}{4e} \right) dx$$

↓ 2009

$$2 \left(\frac{f^2 (4ae^2 - b^2 f^2) (2de - bf^2)^3}{64e^5 \left(-2e \left(f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} + d + ex \right) - bf^2 + 2de \right)} + \frac{3f^2 (4ae^2 - b^2 f^2) (2de - bf^2)^2 \log \left(-2e \left(f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} + d + ex \right) - bf^2 + 2de \right)}{64e^5} \right)$$

input

```
Int[(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^3,x]
```

output

```
2*((f^2*(2*d*e - b*f^2)*(4*a*e^2 - b^2*f^2)*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]))/(16*e^4) + (f^2*(4*a*e^2 - b^2*f^2)*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^2)/(32*e^3) + (d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^4/(16*e) + (f^2*(2*d*e - b*f^2)^3*(4*a*e^2 - b^2*f^2))/(64*e^5*(2*d*e - b*f^2 - 2*e*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]))) + (3*f^2*(2*d*e - b*f^2)^2*(4*a*e^2 - b^2*f^2)*Log[2*d*e - b*f^2 - 2*e*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])])/(64*e^5))
```

Defintions of rubi rules used

rule 1195

```
Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))^(n_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2541

```
Int[((g_) + (h_)*((d_) + (e_)*(x_) + (f_)*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2]))^(n_))^(p_), x_Symbol] := Simp[2 Subst[Int[(g + h*x^n)^p*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)/(-2*d*e + b*f^2 + 2*e*x)^2], x], x, d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 684 vs. $2(283) = 566$.

Time = 0.05 (sec) , antiderivative size = 685, normalized size of antiderivative = 2.26

$$\frac{3d^2 f^3 \sqrt{a + bx + \frac{e^2 x^2}{f^2}} b}{4e^2} + \frac{3f d^2 \ln \left(\frac{\frac{b}{2} + \frac{e^2 x}{f^2}}{\sqrt{\frac{e^2}{f^2}}} + \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right) a}{2\sqrt{\frac{e^2}{f^2}}} - \frac{3d b^2 f^5 \sqrt{a + bx + \frac{e^2 x^2}{f^2}}}{4e^3} + \frac{3f^5 b^2 \sqrt{a + bx}}{8e^2}$$

input

```
int((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^3,x)
```

output

```

3/4*d^2/e^2*f^3*(a+b*x+e^2*x^2/f^2)^(1/2)*b+3/2*f*d^2*ln((1/2*b+e^2*x/f^2)
/(e^2/f^2)^(1/2)+(a+b*x+e^2*x^2/f^2)^(1/2))/(e^2/f^2)^(1/2)*a-3/4*d/e^3*b^
2*f^5*(a+b*x+e^2*x^2/f^2)^(1/2)+3/8*f^5/e^2*b^2*(a+b*x+e^2*x^2/f^2)^(1/2)*
x-3/32*f^7/e^4*b^4*ln((1/2*b+e^2*x/f^2)/(e^2/f^2)^(1/2)+(a+b*x+e^2*x^2/f^2
)^(1/2))/(e^2/f^2)^(1/2)-3/2*d/e*b*f^3*ln((1/2*b+e^2*x/f^2)/(e^2/f^2)^(1/2
)+(a+b*x+e^2*x^2/f^2)^(1/2))/(e^2/f^2)^(1/2)*a+f^3*(a+b*x+e^2*x^2/f^2)^(3/
2)*x+e^3*x^4+1/4*d^4/e+3/8*f^5/e^2*a*ln((1/2*b+e^2*x/f^2)/(e^2/f^2)^(1/2)+
(a+b*x+e^2*x^2/f^2)^(1/2))/(e^2/f^2)^(1/2)*b^2-3/8*d^2/e^2*f^3*ln((1/2*b+e
^2*x/f^2)/(e^2/f^2)^(1/2)+(a+b*x+e^2*x^2/f^2)^(1/2))/(e^2/f^2)^(1/2)*b^2-3
/2*d/e*b*f^3*(a+b*x+e^2*x^2/f^2)^(1/2)*x+3/8*d/e^3*b^3*f^5*ln((1/2*b+e^2*x
/f^2)/(e^2/f^2)^(1/2)+(a+b*x+e^2*x^2/f^2)^(1/2))/(e^2/f^2)^(1/2)+2*d/e*f^3
*(a+b*x+e^2*x^2/f^2)^(3/2)+3/2*f*d^2*(a+b*x+e^2*x^2/f^2)^(1/2)*x-1/2*f^5/e
^2*(a+b*x+e^2*x^2/f^2)^(3/2)*b+3/16*f^7/e^4*b^3*(a+b*x+e^2*x^2/f^2)^(1/2)+
2*d*e^2*x^3+f^2*x^3*e*b+3/2*f^2*x^2*a*e+3/2*f^2*b*d*x^2+3*f^2*x*a*d+3/2*e*
x^2*d^2+x*d^3

```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 345, normalized size of antiderivative = 1.14

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^3 dx$$

$$= \frac{32 e^8 x^4 + 32 (b e^6 f^2 + 2 d e^7) x^3 + 48 (d^2 e^6 + (b d e^5 + a e^6) f^2) x^2 + 32 (3 a d e^5 f^2 + d^3 e^5) x + 3 (b^4 f^8 - 16 a d e^5 f^2 + 12 d^2 e^5 f - 4 (b^3 d e + a b^2 e^2) f^6 + 4 (b^2 d^2 e^2 + 4 a b d e^3) f^4) \log(-b f^2 - 2 e^2 x + 2 e f \sqrt{(b f^2 x + e^2 x^2 + a f^2) / f^2}) + 2 (3 b^3 e f^7 + 16 e^7 f x^3 - 4 (3 b^2 d e^2 + 2 a b e^3) f^5 + 4 (3 b d^2 e^3 + 8 a d e^4) f^3 + 8 (b e^5 f^3 + 4 d e^6 f) x^2 - 2 (b^2 e^3 f^5 - 12 d^2 e^5 f - 4 (b d e^4 + 2 a e^5) f^3) x) \sqrt{(b f^2 x + e^2 x^2 + a f^2) / f^2}}{e^5}$$

input

```
integrate((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^3,x, algorithm="fricas")
```

output

```

1/32*(32*e^8*x^4 + 32*(b*e^6*f^2 + 2*d*e^7)*x^3 + 48*(d^2*e^6 + (b*d*e^5 +
a*e^6)*f^2)*x^2 + 32*(3*a*d*e^5*f^2 + d^3*e^5)*x + 3*(b^4*f^8 - 16*a*d^2*
e^4*f^2 - 4*(b^3*d*e + a*b^2*e^2)*f^6 + 4*(b^2*d^2*e^2 + 4*a*b*d*e^3)*f^4)
*log(-b*f^2 - 2*e^2*x + 2*e*f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2)) + 2*(
3*b^3*e*f^7 + 16*e^7*f*x^3 - 4*(3*b^2*d*e^2 + 2*a*b*e^3)*f^5 + 4*(3*b*d^2*
e^3 + 8*a*d*e^4)*f^3 + 8*(b*e^5*f^3 + 4*d*e^6*f)*x^2 - 2*(b^2*e^3*f^5 - 12
*d^2*e^5*f - 4*(b*d*e^4 + 2*a*e^5)*f^3)*x)*sqrt((b*f^2*x + e^2*x^2 + a*f^2
)/f^2))/e^5

```


Sympy [A] (verification not implemented)

Time = 1.25 (sec) , antiderivative size = 1363, normalized size of antiderivative = 4.50

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^3 dx = \text{Too large to display}$$

input `integrate((d+e*x+f*(a+b*x+e**2*x**2/f**2)**(1/2))**3,x)`

output `3*a*d*f**2*x + 3*a*e*f**2*x**2/2 + a*f**3*Piecewise(((a/2 - b**2*f**2/(8*e**2))*Piecewise((log(b + 2*e**2*x/f**2 + 2*sqrt(e**2/f**2)*sqrt(a + b*x + e**2*x**2/f**2))/sqrt(e**2/f**2), Ne(a - b**2*f**2/(4*e**2), 0)), ((b*f**2/(2*e**2) + x)*log(b*f**2/(2*e**2) + x)/sqrt(e**2*(b*f**2/(2*e**2) + x)**2/f**2), True)) + (b*f**2/(4*e**2) + x/2)*sqrt(a + b*x + e**2*x**2/f**2), Ne(e**2/f**2, 0)), (2*(a + b*x)**(3/2)/(3*b), Ne(b, 0)), (sqrt(a)*x, True)) + 3*b*d*f**2*x**2/2 + b*e*f**2*x**3 + b*f**3*Piecewise(((a/2 - b**2*f**2/(8*e**2)) - b*f**2*(a/3 - b**2*f**2/(8*e**2))/(2*e**2))*Piecewise((log(b + 2*e**2*x/f**2 + 2*sqrt(e**2/f**2)*sqrt(a + b*x + e**2*x**2/f**2))/sqrt(e**2/f**2), Ne(a - b**2*f**2/(4*e**2), 0)), ((b*f**2/(2*e**2) + x)*log(b*f**2/(2*e**2) + x)/sqrt(e**2*(b*f**2/(2*e**2) + x)**2/f**2), True)) + sqrt(a + b*x + e**2*x**2/f**2)*(b*f**2*x/(12*e**2) + x**2/3 + f**2*(a/3 - b**2*f**2/(8*e**2))/e**2), Ne(e**2/f**2, 0)), (2*(-a*(a + b*x)**(3/2)/3 + (a + b*x)**(5/2)/5)/b**2, Ne(b, 0)), (sqrt(a)*x**2/2, True)) + d**3*x + 3*d**2*e*x**2/2 + 3*d**2*f*Piecewise(((a/2 - b**2*f**2/(8*e**2))*Piecewise((log(b + 2*e**2*x/f**2 + 2*sqrt(e**2/f**2)*sqrt(a + b*x + e**2*x**2/f**2))/sqrt(e**2/f**2), Ne(a - b**2*f**2/(4*e**2), 0)), ((b*f**2/(2*e**2) + x)*log(b*f**2/(2*e**2) + x)/sqrt(e**2*(b*f**2/(2*e**2) + x)**2/f**2), True)) + (b*f**2/(4*e**2) + x/2)*sqrt(a + b*x + e**2*x**2/f**2), Ne(e**2/f**2, 0)), (2*(a + b*x)**(3/2)/(3*b), Ne(b, 0)), (sqrt(a)*x, True)) + 2*d*e**2*x**3 + 6*d*e*f*...`

Maxima [F(-2)]

Exception generated.

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^3 dx = \text{Exception raised: ValueError}$$

input `integrate((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^3,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b^2*f^2-4*a*e^2>0)', see `assume ?` for mor

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 397, normalized size of antiderivative = 1.31

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^3 dx$$

$$= be^2 x^3 + e^3 x^4 + \frac{3}{2} bdf^2 x^2 + \frac{3}{2} aef^2 x^2 + 2de^2 x^3 + 3adf^2 x + \frac{3}{2} d^2 ex^2 + d^3 x$$

$$+ \frac{1}{16} \sqrt{bf^2 x + e^2 x^2 + af^2} \left(2 \left(4 \left(\frac{2e^2 x|f|}{f} + \frac{be^6 f^4 |f| + 4de^7 f^2 |f|}{e^6 f^3} \right) x - \frac{b^2 e^4 f^6 |f| - 4bde^5 f^4 |f| - 8ae^6 f^3}{e^6 f^3} \right. \right.$$

$$\left. \left. + \frac{3(b^4 f^7 |f| - 4b^3 de f^5 |f| - 4ab^2 e^2 f^5 |f| + 4b^2 d^2 e^2 f^3 |f| + 16abde^3 f^3 |f| - 16ad^2 e^4 f |f|) \log(|-bf^2 - 2e^2 x + af^2|)}{32e^4 |e|} \right) \right)$$

input `integrate((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^3,x, algorithm="giac")`

output `b*e*f^2*x^3 + e^3*x^4 + 3/2*b*d*f^2*x^2 + 3/2*a*e*f^2*x^2 + 2*d*e^2*x^3 + 3*a*d*f^2*x + 3/2*d^2*e*x^2 + d^3*x + 1/16*sqrt(b*f^2*x + e^2*x^2 + a*f^2) * (2*(4*(2*e^2*x*abs(f)/f + (b*e^6*f^4*abs(f) + 4*d*e^7*f^2*abs(f))/(e^6*f^3)))*x - (b^2*e^4*f^6*abs(f) - 4*b*d*e^5*f^4*abs(f) - 8*a*e^6*f^4*abs(f) - 12*d^2*e^6*f^2*abs(f))/(e^6*f^3))*x + (3*b^3*e^2*f^8*abs(f) - 12*b^2*d*e^3*f^6*abs(f) - 8*a*b*e^4*f^6*abs(f) + 12*b*d^2*e^4*f^4*abs(f) + 32*a*d*e^5*f^4*abs(f))/(e^6*f^3) + 3/32*(b^4*f^7*abs(f) - 4*b^3*d*e*f^5*abs(f) - 4*a*b^2*e^2*f^5*abs(f) + 4*b^2*d^2*e^2*f^3*abs(f) + 16*a*b*d*e^3*f^3*abs(f) - 16*a*d^2*e^4*f*abs(f))*log(abs(-b*f^2 - 2*(x*abs(e) - sqrt(b*f^2*x + e^2*x^2 + a*f^2))*abs(e)))/(e^4*abs(e))`

Mupad [F(-1)]

Timed out.

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^3 dx = \int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^3 dx$$

input `int((d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^3,x)`output `int((d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^3, x)`**Reduce [B] (verification not implemented)**

Time = 0.39 (sec) , antiderivative size = 870, normalized size of antiderivative = 2.87

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^3 dx$$

$$= \frac{6\sqrt{b f^2 x + e^2 x^2 + a f^2} b^3 e f^6 + 48\sqrt{b f^2 x + e^2 x^2 + a f^2} d^2 e^5 x + 64\sqrt{b f^2 x + e^2 x^2 + a f^2} d e^6 x^2 + 48 a e^6}{\dots}$$

input `int((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^3,x)`

output

```
( - 16*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*a*b*e**3*f**4 + 64*sqrt(a*f**2
+ b*f**2*x + e**2*x**2)*a*d*e**4*f**2 + 32*sqrt(a*f**2 + b*f**2*x + e**2*x
**2)*a*e**5*f**2*x + 6*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b**3*e*f**6 - 2
4*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b**2*d*e**2*f**4 - 4*sqrt(a*f**2 + b
*f**2*x + e**2*x**2)*b**2*e**3*f**4*x + 24*sqrt(a*f**2 + b*f**2*x + e**2*x
**2)*b*d**2*e**3*f**2 + 16*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b*d*e**4*f*
*2*x + 16*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b*e**5*f**2*x**2 + 48*sqrt(a
*f**2 + b*f**2*x + e**2*x**2)*d**2*e**5*x + 64*sqrt(a*f**2 + b*f**2*x + e
**2*x**2)*d*e**6*x**2 + 32*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*e**7*x**3 +
12*log((2*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*e + b*f**2 + 2*e**2*x)/(sqrt
(4*a*e**2 - b**2*f**2)*f))*a*b**2*e**2*f**6 - 48*log((2*sqrt(a*f**2 + b*f*
**2*x + e**2*x**2)*e + b*f**2 + 2*e**2*x)/(sqrt(4*a*e**2 - b**2*f**2)*f))*a
*b*d*e**3*f**4 + 48*log((2*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*e + b*f**2
+ 2*e**2*x)/(sqrt(4*a*e**2 - b**2*f**2)*f))*a*d**2*e**4*f**2 - 3*log((2*sq
rt(a*f**2 + b*f**2*x + e**2*x**2)*e + b*f**2 + 2*e**2*x)/(sqrt(4*a*e**2 -
b**2*f**2)*f))*b**4*f**8 + 12*log((2*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*e
+ b*f**2 + 2*e**2*x)/(sqrt(4*a*e**2 - b**2*f**2)*f))*b**3*d*e*f**6 - 12*l
og((2*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*e + b*f**2 + 2*e**2*x)/(sqrt(4*a
e**2 - b**2*f**2)*f))*b**2*d**2*e**2*f**4 + 96*a*d*e**5*f**2*x + 48*a*e**
6*f**2*x**2 + 48*b*d*e**5*f**2*x**2 + 32*b*e**6*f**2*x**3 + 32*d**3*e**...
```

3.8 $\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^2 dx$

Optimal result	92
Mathematica [A] (verified)	93
Rubi [A] (verified)	93
Maple [A] (verified)	95
Fricas [A] (verification not implemented)	96
Sympy [A] (verification not implemented)	97
Maxima [F(-2)]	98
Giac [A] (verification not implemented)	99
Mupad [F(-1)]	99
Reduce [B] (verification not implemented)	100

Optimal result

Integrand size = 28, antiderivative size = 237

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^2 dx$$

$$= \frac{f^2(4ae^2 - b^2 f^2) \left(ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)}{8e^3} + \frac{\left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^3}{6e}$$

$$- \frac{f^2(2de - bf^2)^2 (4ae^2 - b^2 f^2)}{16e^4 \left(bf^2 + 2e \left(ex + f \sqrt{a + \frac{x(bf^2 + e^2 x)}{f^2}} \right) \right)}$$

$$+ \frac{f^2(2de - bf^2) (4ae^2 - b^2 f^2) \log \left(bf^2 + 2e \left(ex + f \sqrt{a + \frac{x(bf^2 + e^2 x)}{f^2}} \right) \right)}{8e^4}$$

output

```
1/8*f^2*(-b^2*f^2+4*a*e^2)*(e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))/e^3+1/6*(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^3/e-1/16*f^2*(-b*f^2+2*d*e)^2*(-b^2*f^2+4*a*e^2)/e^4/(b*f^2+2*e*(e*x+f*(a+x*(b*f^2+e^2*x)/f^2)^(1/2)))+1/8*f^2*(-b*f^2+2*d*e)*(-b^2*f^2+4*a*e^2)*ln(b*f^2+2*e*(e*x+f*(a+x*(b*f^2+e^2*x)/f^2)^(1/2)))/e^4
```

Mathematica [A] (verified)

Time = 1.33 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.74

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^2 dx$$

$$= \frac{1}{12} \left(2x(6d^2 + 6af^2 + 6dex + x(3bf^2 + 4e^2x)) \right. \\ \left. + \frac{\sqrt{a + x \left(b + \frac{e^2 x}{f^2} \right)} (-3b^2 f^5 + 2be f^3 (3d + ex) + 4e^2 f (2af^2 + ex(3d + 2ex)))}{e^3} \right. \\ \left. + \frac{3f^2(-2de + bf^2)(-4ae^2 + b^2 f^2) \operatorname{arctanh} \left(\frac{ex}{f(-\sqrt{a} + \sqrt{a + x(b + \frac{e^2 x}{f^2})})} \right)}{e^4} \right)$$

input `Integrate[(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^2,x]`

output `(2*x*(6*d^2 + 6*a*f^2 + 6*d*e*x + x*(3*b*f^2 + 4*e^2*x)) + (Sqrt[a + x*(b + (e^2*x)/f^2)]*(-3*b^2*f^5 + 2*b*e*f^3*(3*d + e*x) + 4*e^2*f*(2*a*f^2 + e*x*(3*d + 2*e*x))))/e^3 + (3*f^2*(-2*d*e + b*f^2)*(-4*a*e^2 + b^2*f^2)*ArcTanh[(e*x)/(f*(-Sqrt[a] + Sqrt[a + x*(b + (e^2*x)/f^2])]))]/e^4)/12`

Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.04, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$, Rules used = {2541, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} + d + ex \right)^2 dx$$

↓ 2541

$$2 \int \frac{\left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)^2 \left(ed^2 - bf^2 d + aef^2 + e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)^2 - (2de - bf^2) \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right) \right)}{\left(-bf^2 + 2de - 2e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right) \right)^2} dx$$

↓ 1195

$$2 \int \left(-\frac{(2de - bf^2)(4ae^2 - b^2 f^2) f^2}{8e^3 \left(-bf^2 + 2de - 2e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right) \right)} + \frac{\left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)^2}{4e} + \frac{4ae^2 f^2 - b^2 f^2}{16e^3} \right) dx$$

↓ 2009

$$2 \left(\frac{f^2 (4ae^2 - b^2 f^2) (2de - bf^2)^2}{32e^4 \left(-2e \left(f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} + d + ex \right) - bf^2 + 2de \right)} + \frac{f^2 (4ae^2 - b^2 f^2) (2de - bf^2) \log \left(-2e \left(f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} + d + ex \right) - bf^2 + 2de \right)}{16e^4} \right)$$

input

```
Int[(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^2,x]
```

output

```
2*((f^2*(4*a*e^2 - b^2*f^2)*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]))/(16*e^3) + (d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^3/(12*e) + (f^2*(2*d*e - b*f^2)^2*(4*a*e^2 - b^2*f^2))/(32*e^4*(2*d*e - b*f^2 - 2*e*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]))) + (f^2*(2*d*e - b*f^2)*(4*a*e^2 - b^2*f^2)*Log[2*d*e - b*f^2 - 2*e*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]])/(16*e^4))
```

Definitions of rubi rules used

rule 1195

```
Int[((d_.) + (e_.)*(x_)^(m_.))*((f_.) + (g_.)*(x_)^(n_.))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2541

```
Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]))^(n_.))^((p_.), x_Symbol] := Simp[2 Subst[Int[(g + h*x^n)^p*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)/(-2*d*e + b*f^2 + 2*e*x)^2], x], x, d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]
```

Maple [A] (verified)

Time = 1.21 (sec) , antiderivative size = 300, normalized size of antiderivative = 1.27

method	result
default	$\frac{bx^2f^2}{2} + \frac{e^2x^3}{3} + af^2x + 2f \left(d \left(\frac{(b + \frac{2e^2x}{f^2})f^2\sqrt{a+bx+\frac{e^2x^2}{f^2}}}{4e^2} + \frac{(4e^2a - b^2)f^2 \ln\left(\frac{\frac{b}{2} + \frac{e^2x}{f^2}}{\sqrt{\frac{e^2}{f^2}}} + \sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)}{8e^2\sqrt{\frac{e^2}{f^2}}}\right) + e \right)$

input

```
int((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^2,x,method=_RETURNVERBOSE)
```


output

```
1/2*b*x^2*f^2+1/3*e^2*x^3+a*f^2*x+2*f*(d*(1/4*(b+2*e^2*x/f^2)/e^2*f^2*(a+b
*x+e^2*x^2/f^2)^(1/2)+1/8*(4*e^2/f^2*a-b^2)/e^2*f^2*ln((1/2*b+e^2*x/f^2)/(
e^2/f^2)^(1/2)+(a+b*x+e^2*x^2/f^2)^(1/2))/(e^2/f^2)^(1/2))+e*(1/3*(a+b*x+e
^2*x^2/f^2)^(3/2)/e^2*f^2-1/2*b/e^2*f^2*(1/4*(b+2*e^2*x/f^2)/e^2*f^2*(a+b*
x+e^2*x^2/f^2)^(1/2)+1/8*(4*e^2/f^2*a-b^2)/e^2*f^2*ln((1/2*b+e^2*x/f^2)/(e
^2/f^2)^(1/2)+(a+b*x+e^2*x^2/f^2)^(1/2))/(e^2/f^2)^(1/2))))+1/3*(e*x+d)^3/
e
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 219, normalized size of antiderivative = 0.92

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^2 dx$$

$$= \frac{16 e^6 x^3 + 12 (b e^4 f^2 + 2 d e^5) x^2 + 24 (a e^4 f^2 + d^2 e^4) x - 3 (b^3 f^6 + 8 a d e^3 f^2 - 2 (b^2 d e + 2 a b e^2) f^4) \log(-b$$

input

```
integrate((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^2,x, algorithm="fricas")
```

output

```
1/24*(16*e^6*x^3 + 12*(b*e^4*f^2 + 2*d*e^5)*x^2 + 24*(a*e^4*f^2 + d^2*e^4)
*x - 3*(b^3*f^6 + 8*a*d*e^3*f^2 - 2*(b^2*d*e + 2*a*b*e^2)*f^4)*log(-b*f^2
- 2*e^2*x + 2*e*f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2)) - 2*(3*b^2*e*f^5
- 8*e^5*f*x^2 - 2*(3*b*d*e^2 + 4*a*e^3)*f^3 - 2*(b*e^3*f^3 + 6*d*e^4*f)*x)
*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2))/e^4
```

Sympy [A] (verification not implemented)

Time = 0.74 (sec) , antiderivative size = 490, normalized size of antiderivative = 2.07

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^2 dx = af^2 x + \frac{bf^2 x^2}{2} + d^2 x + dex^2$$

$$+ 2df \left(\left(\frac{a}{2} - \frac{b^2 f^2}{8e^2} \right) \left(\begin{cases} \frac{\log \left(b + \frac{2e^2 x}{f^2} + 2\sqrt{\frac{e^2}{f^2}} \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)}{\sqrt{\frac{e^2}{f^2}}} & \text{for } a - \frac{b^2 f^2}{4e^2} \neq 0 \\ \frac{\left(\frac{bf^2}{2e^2} + x \right) \log \left(\frac{bf^2}{2e^2} + x \right)}{\sqrt{\frac{e^2 \left(\frac{bf^2}{2e^2} + x \right)^2}{f^2}}} & \text{otherwise} \end{cases} \right) + \left(\frac{bf^2}{4e^2} + \frac{x}{2} \right) \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right.$$

$$\left. \frac{2(a+bx)^{\frac{3}{2}}}{3b} \right)$$

$$+ \frac{2e^2 x^3}{3}$$

$$+ 2ef \left(\left(-\frac{abf^2}{12e^2} - \frac{bf^2 \left(\frac{a}{3} - \frac{b^2 f^2}{8e^2} \right)}{2e^2} \right) \left(\begin{cases} \frac{\log \left(b + \frac{2e^2 x}{f^2} + 2\sqrt{\frac{e^2}{f^2}} \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)}{\sqrt{\frac{e^2}{f^2}}} & \text{for } a - \frac{b^2 f^2}{4e^2} \neq 0 \\ \frac{\left(\frac{bf^2}{2e^2} + x \right) \log \left(\frac{bf^2}{2e^2} + x \right)}{\sqrt{\frac{e^2 \left(\frac{bf^2}{2e^2} + x \right)^2}{f^2}}} & \text{otherwise} \end{cases} \right) + \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)$$

$$\left(\frac{2 \left(-\frac{a(a+bx)^{\frac{3}{2}}}{3} + \frac{(a+bx)^{\frac{5}{2}}}{5} \right)}{b^2} \right)$$

$$\left(\frac{\sqrt{ax^2}}{2} \right)$$

input

```
integrate((d+e*x+f*(a+b*x+e**2*x**2/f**2)**(1/2))**2,x)
```

output

```
a*f**2*x + b*f**2*x**2/2 + d**2*x + d*e*x**2 + 2*d*f*Piecewise(((a/2 - b**
2*f**2/(8*e**2))*Piecewise((log(b + 2*e**2*x/f**2 + 2*sqrt(e**2/f**2)*sqrt
(a + b*x + e**2*x**2/f**2))/sqrt(e**2/f**2), Ne(a - b**2*f**2/(4*e**2), 0)
), ((b*f**2/(2*e**2) + x)*log(b*f**2/(2*e**2) + x)/sqrt(e**2*(b*f**2/(2*e
**2) + x)**2/f**2), True)) + (b*f**2/(4*e**2) + x/2)*sqrt(a + b*x + e**2*x*
*2/f**2), Ne(e**2/f**2, 0)), (2*(a + b*x)**(3/2)/(3*b), Ne(b, 0)), (sqrt(a
)*x, True)) + 2*e**2*x**3/3 + 2*e*f*Piecewise(((a*b*f**2/(12*e**2) - b*f*
*2*(a/3 - b**2*f**2/(8*e**2))/(2*e**2))*Piecewise((log(b + 2*e**2*x/f**2 +
2*sqrt(e**2/f**2)*sqrt(a + b*x + e**2*x**2/f**2))/sqrt(e**2/f**2), Ne(a -
b**2*f**2/(4*e**2), 0)), ((b*f**2/(2*e**2) + x)*log(b*f**2/(2*e**2) + x)/
sqrt(e**2*(b*f**2/(2*e**2) + x)**2/f**2), True)) + sqrt(a + b*x + e**2*x**
2/f**2)*(b*f**2*x/(12*e**2) + x**2/3 + f**2*(a/3 - b**2*f**2/(8*e**2))/e**
2), Ne(e**2/f**2, 0)), (2*(-a*(a + b*x)**(3/2)/3 + (a + b*x)**(5/2)/5)/b**
2, Ne(b, 0)), (sqrt(a)*x**2/2, True))
```

Maxima [F(-2)]

Exception generated.

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^2 dx = \text{Exception raised: ValueError}$$

input

```
integrate((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^2,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(b^2*f^2-4*a*e^2>0)', see `assume
?` for mor
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.00

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^2 dx = \frac{1}{2} b f^2 x^2 + \frac{2}{3} e^2 x^3 + a f^2 x + d e x^2 + d^2 x$$

$$+ \frac{1}{12} \sqrt{b f^2 x + e^2 x^2 + a f^2} \left(2 \left(\frac{4 e x |f|}{f} + \frac{b e^3 f^3 |f| + 6 d e^4 f |f|}{e^4 f^2} \right) x - \frac{3 b^2 e f^5 |f| - 6 b d e^2 f^3 |f| - 8 a e^3 f^3 |f|}{e^4 f^2} \right.$$

$$\left. - \frac{(b^3 f^5 |f| - 2 b^2 d e f^3 |f| - 4 a b e^2 f^3 |f| + 8 a d e^3 f |f|) \log(|-b f^2 - 2(x|e| - \sqrt{b f^2 x + e^2 x^2 + a f^2})|e|)}{8 e^3 |e|} \right)$$

input `integrate((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^2,x, algorithm="giac")`output `1/2*b*f^2*x^2 + 2/3*e^2*x^3 + a*f^2*x + d*e*x^2 + d^2*x + 1/12*sqrt(b*f^2*x + e^2*x^2 + a*f^2)*(2*(4*e*x*abs(f)/f + (b*e^3*f^3*abs(f) + 6*d*e^4*f*abs(f))/(e^4*f^2))*x - (3*b^2*e*f^5*abs(f) - 6*b*d*e^2*f^3*abs(f) - 8*a*e^3*f^3*abs(f))/(e^4*f^2) - 1/8*(b^3*f^5*abs(f) - 2*b^2*d*e*f^3*abs(f) - 4*a*b*e^2*f^3*abs(f) + 8*a*d*e^3*f*abs(f))*log(abs(-b*f^2 - 2*(x*abs(e) - sqrt(b*f^2*x + e^2*x^2 + a*f^2))*abs(e)))/(e^3*abs(e))`**Mupad [F(-1)]**

Timed out.

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^2 dx = \int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^2 dx$$

input `int((d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^2,x)`output `int((d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^2, x)`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 497, normalized size of antiderivative = 2.10

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^2 dx$$

$$= \frac{16\sqrt{b} f^2 x + e^2 x^2 + a f^2 a e^3 f^2 - 6\sqrt{b} f^2 x + e^2 x^2 + a f^2 b^2 e f^4 + 12\sqrt{b} f^2 x + e^2 x^2 + a f^2 b d e^2 f^2 + 4\sqrt{b} f^2 x + e^2 x^2 + a f^2 b^2 e f^4 + 12\sqrt{b} f^2 x + e^2 x^2 + a f^2 b d e^2 f^2 + 4\sqrt{b} f^2 x + e^2 x^2 + a f^2 b^2 e f^4}{24 e^4}$$

input

```
int((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^2,x)
```

output

```
(16*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*a*e**3*f**2 - 6*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b**2*e*f**4 + 12*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b*d*e**2*f**2 + 4*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b*e**3*f**2*x + 24*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*d*e**4*x + 16*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*e**5*x**2 - 12*log((2*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*e + b*f**2 + 2*e**2*x)/(sqrt(4*a*e**2 - b**2*f**2)*f))*a*b*e**2*f**4 + 24*log((2*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*e + b*f**2 + 2*e**2*x)/(sqrt(4*a*e**2 - b**2*f**2)*f))*a*d*e**3*f**2 + 3*log((2*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*e + b*f**2 + 2*e**2*x)/(sqrt(4*a*e**2 - b**2*f**2)*f))*b**3*f**6 - 6*log((2*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*e + b*f**2 + 2*e**2*x)/(sqrt(4*a*e**2 - b**2*f**2)*f))*b**2*d*e*f**4 + 24*a*e**4*f**2*x + 12*b*e**4*f**2*x**2 + 24*d**2*e**4*x + 24*d*e**5*x**2 + 16*e**6*x**3)/(24*e**4)
```

3.9 $\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right) dx$

Optimal result	101
Mathematica [A] (verified)	102
Rubi [A] (verified)	102
Maple [A] (verified)	103
Fricas [A] (verification not implemented)	104
Sympy [A] (verification not implemented)	104
Maxima [F(-2)]	105
Giac [A] (verification not implemented)	105
Mupad [F(-1)]	106
Reduce [B] (verification not implemented)	106

Optimal result

Integrand size = 26, antiderivative size = 118

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right) dx = dx + \frac{ex^2}{2} + \frac{f(bf^2 + 2e^2x) \sqrt{a + bx + \frac{e^2 x^2}{f^2}}}{4e^2} + \frac{f^2(4ae^2 - b^2f^2) \operatorname{arctanh} \left(\frac{bf^2 + 2e^2x}{2ef \sqrt{a + bx + \frac{e^2 x^2}{f^2}}} \right)}{8e^3}$$

output

```
d*x+1/2*e*x^2+1/4*f*(b*f^2+2*e^2*x)*(a+b*x+e^2*x^2/f^2)^(1/2)/e^2+1/8*f^2*(-b^2*f^2+4*a*e^2)*arctanh(1/2*(b*f^2+2*e^2*x)/e/f/(a+b*x+e^2*x^2/f^2)^(1/2))/e^3
```

Mathematica [A] (verified)

Time = 0.56 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.53

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right) dx$$

$$= \frac{8de^3 x + 4e^4 x^2 + 2bef^3 \sqrt{a + x \left(b + \frac{e^2 x}{f^2} \right)} + 4e^3 f x \sqrt{a + x \left(b + \frac{e^2 x}{f^2} \right)} + (-4ae^2 f^2 + b^2 f^4) \log \left(e^3 \left(\sqrt{a} f \right)}{8e^3}$$

input

```
Integrate[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2],x]
```

output

```
(8*d*e^3*x + 4*e^4*x^2 + 2*b*e*f^3*Sqrt[a + x*(b + (e^2*x)/f^2)] + 4*e^3*f*x*Sqrt[a + x*(b + (e^2*x)/f^2)] + (-4*a*e^2*f^2 + b^2*f^4)*Log[e^3*(Sqrt[a]*f + e*x - f*Sqrt[a + x*(b + (e^2*x)/f^2)])] + (4*a*e^2*f^2 - b^2*f^4)*Log[-(Sqrt[a]*f) + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])]/(8*e^3)
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.038$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} + d + ex \right) dx$$

↓ 2009

$$\frac{f^2(4ae^2 - b^2 f^2) \operatorname{arctanh} \left(\frac{bf^2 + 2e^2 x}{2ef \sqrt{a + bx + \frac{e^2 x^2}{f^2}}} \right)}{8e^3} + \frac{f(bf^2 + 2e^2 x) \sqrt{a + bx + \frac{e^2 x^2}{f^2}}}{4e^2} + dx + \frac{ex^2}{2}$$

input

```
Int[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2],x]
```

output

$$d*x + (e*x^2)/2 + (f*(b*f^2 + 2*e^2*x)*\text{Sqrt}[a + b*x + (e^2*x^2)/f^2])/(4*e^2) + (f^2*(4*a*e^2 - b^2*f^2)*\text{ArcTanh}[(b*f^2 + 2*e^2*x)/(2*e*f*\text{Sqrt}[a + b*x + (e^2*x^2)/f^2]])/(8*e^3)$$
Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x_Symbol] \text{ :> Simp}[\text{IntSum}[u, x], x] \text{ /; SumQ}[u]$$
Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.04

method	result	size
default	$dx + \frac{e x^2}{2} + f \left(\frac{\left(b + \frac{2e^2 x}{f^2}\right) f^2 \sqrt{a + b x + \frac{e^2 x^2}{f^2}}}{4e^2} + \frac{\left(\frac{4e^2 a}{f^2} - b^2\right) f^2 \ln\left(\frac{\frac{b}{2} + \frac{e^2 x}{f^2}}{\sqrt{\frac{e^2}{f^2}}} + \sqrt{a + b x + \frac{e^2 x^2}{f^2}}\right)}{8e^2 \sqrt{\frac{e^2}{f^2}}}\right)$	123
parts	$dx + \frac{e x^2}{2} + f \left(\frac{\left(b + \frac{2e^2 x}{f^2}\right) f^2 \sqrt{a + b x + \frac{e^2 x^2}{f^2}}}{4e^2} + \frac{\left(\frac{4e^2 a}{f^2} - b^2\right) f^2 \ln\left(\frac{\frac{b}{2} + \frac{e^2 x}{f^2}}{\sqrt{\frac{e^2}{f^2}}} + \sqrt{a + b x + \frac{e^2 x^2}{f^2}}\right)}{8e^2 \sqrt{\frac{e^2}{f^2}}}\right)$	123

input

$$\text{int}(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2), x, \text{method}=_RETURNVERBOSE)$$

output

$$d*x+1/2*e*x^2+f*(1/4*(b+2*e^2*x/f^2)/e^2*f^2*(a+b*x+e^2*x^2/f^2)^(1/2)+1/8*(4*e^2/f^2*a-b^2)/e^2*f^2*\ln((1/2*b+e^2*x/f^2)/(e^2/f^2)^(1/2)+(a+b*x+e^2*x^2/f^2)^(1/2)))/(e^2/f^2)^(1/2))$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.04

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right) dx$$

$$= \frac{4e^4 x^2 + 8de^3 x + (b^2 f^4 - 4ae^2 f^2) \log \left(-bf^2 - 2e^2 x + 2ef \sqrt{\frac{bf^2 x + e^2 x^2 + af^2}{f^2}} \right) + 2(bef^3 + 2e^3 fx) \sqrt{\frac{bf^2 x + e^2 x^2 + af^2}{f^2}}}{8e^3}$$

input `integrate(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2),x, algorithm="fricas")`

output `1/8*(4*e^4*x^2 + 8*d*e^3*x + (b^2*f^4 - 4*a*e^2*f^2)*log(-b*f^2 - 2*e^2*x + 2*e*f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2)) + 2*(b*e*f^3 + 2*e^3*f*x)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2))/e^3`

Sympy [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.67

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right) dx = dx + \frac{ex^2}{2}$$

$$+ f \left(\left(\frac{a}{2} - \frac{b^2 f^2}{8e^2} \right) \left(\begin{cases} \frac{\log \left(b + \frac{2e^2 x}{f^2} + 2\sqrt{\frac{e^2}{f^2}} \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)}{\sqrt{\frac{e^2}{f^2}}} & \text{for } a - \frac{b^2 f^2}{4e^2} \neq 0 \\ \frac{\left(\frac{bf^2}{2e^2} + x \right) \log \left(\frac{bf^2}{2e^2} + x \right)}{\sqrt{\frac{e^2 \left(\frac{bf^2}{2e^2} + x \right)^2}{f^2}}} & \text{otherwise} \end{cases} \right) + \left(\frac{bf^2}{4e^2} + \frac{x}{2} \right) \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right.$$

$$\left. \frac{2(a+bx)^{\frac{3}{2}}}{3b} \right)$$

$$\left. \sqrt{ax} \right)$$

input `integrate(d+e*x+f*(a+b*x+e**2*x**2/f**2)**(1/2),x)`

output

```
d*x + e*x**2/2 + f*Piecewise(((a/2 - b**2*f**2/(8*e**2))*Piecewise((log(b
+ 2*e**2*x/f**2 + 2*sqrt(e**2/f**2)*sqrt(a + b*x + e**2*x**2/f**2))/sqrt(e
**2/f**2), Ne(a - b**2*f**2/(4*e**2), 0)), ((b*f**2/(2*e**2) + x)*log(b*f*
**2/(2*e**2) + x)/sqrt(e**2*(b*f**2/(2*e**2) + x)**2/f**2), True)) + (b*f**
2/(4*e**2) + x/2)*sqrt(a + b*x + e**2*x**2/f**2), Ne(e**2/f**2, 0)), (2*(a
+ b*x)**(3/2)/(3*b), Ne(b, 0)), (sqrt(a)*x, True))
```

Maxima [F(-2)]

Exception generated.

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right) dx = \text{Exception raised: ValueError}$$

input

```
integrate(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(b^2*f^2-4*a*e^2>0)', see `assume
?` for mor
```

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.01

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right) dx = \frac{1}{2} ex^2 + dx + \frac{\left(2 \sqrt{bf^2x + e^2x^2 + af^2} \left(\frac{bf^2}{e^2} + 2x \right) + \frac{(b^2f^4 - 4ae^2f^2) \log\left(\frac{|-bf^2 - 2(x|e| - \sqrt{bf^2x + e^2x^2 + af^2})|e|}{e^2|e|} \right)}{8f} \right) |f|}{8f}$$

input

```
integrate(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2),x, algorithm="giac")
```

output

```
1/2*e*x^2 + d*x + 1/8*(2*sqrt(b*f^2*x + e^2*x^2 + a*f^2)*(b*f^2/e^2 + 2*x)
+ (b^2*f^4 - 4*a*e^2*f^2)*log(abs(-b*f^2 - 2*(x*abs(e) - sqrt(b*f^2*x + e
^2*x^2 + a*f^2))*abs(e)))/(e^2*abs(e))*abs(f)/f
```

Mupad [F(-1)]

Timed out.

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right) dx = \int d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} dx$$

input

```
int(d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2), x)
```

output

```
int(d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2), x)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.75

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right) dx$$

$$= \frac{2\sqrt{b f^2 x + e^2 x^2 + a f^2} b e f^2 + 4\sqrt{b f^2 x + e^2 x^2 + a f^2} e^3 x + 4 \log\left(\frac{2\sqrt{b f^2 x + e^2 x^2 + a f^2} e + b f^2 + 2e^2 x}{\sqrt{-b^2 f^2 + 4a e^2 f}}\right) a e^2 f^2 - 1}{8e^3}$$

input

```
int(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2), x)
```

output

```
(2*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b*e*f**2 + 4*sqrt(a*f**2 + b*f**2*x
+ e**2*x**2)*e**3*x + 4*log((2*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*e + b*
f**2 + 2*e**2*x)/(sqrt(4*a*e**2 - b**2*f**2)*f))*a*e**2*f**2 - log((2*sqrt
(a*f**2 + b*f**2*x + e**2*x**2)*e + b*f**2 + 2*e**2*x)/(sqrt(4*a*e**2 - b*
**2*f**2)*f))*b**2*f**4 + 8*d*e**3*x + 4*e**4*x**2)/(8*e**3)
```

$$3.10 \quad \int \frac{1}{d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}} dx$$

Optimal result	107
Mathematica [A] (verified)	108
Rubi [A] (verified)	108
Maple [B] (verified)	110
Fricas [A] (verification not implemented)	111
Sympy [F]	111
Maxima [F]	112
Giac [F(-1)]	112
Mupad [F(-1)]	112
Reduce [B] (verification not implemented)	113

Optimal result

Integrand size = 28, antiderivative size = 215

$$\begin{aligned} & \int \frac{1}{d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}} dx \\ &= -\frac{f^2(4ae^2-b^2f^2)}{2e(2de-bf^2)\left(bf^2+2e\left(ex+f\sqrt{a+\frac{x(bf^2+e^2x)}{f^2}}\right)\right)} \\ & \quad + \frac{2(d^2e-bdf^2+ae^2f^2)\log\left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)}{(2de-bf^2)^2} \\ & \quad - \frac{f^2(4ae^2-b^2f^2)\log\left(bf^2+2e\left(ex+f\sqrt{a+\frac{x(bf^2+e^2x)}{f^2}}\right)\right)}{2e(2de-bf^2)^2} \end{aligned}$$

output

```
-1/2*f^2*(-b^2*f^2+4*a*e^2)/e/(-b*f^2+2*d*e)/(b*f^2+2*e*(e*x+f*(a+x*(b*f^2+e^2*x)/f^2)^(1/2)))+2*(a*e*f^2-b*d*f^2+d^2*e)*ln(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))/(-b*f^2+2*d*e)^2-1/2*f^2*(-b^2*f^2+4*a*e^2)*ln(b*f^2+2*e*(e*x+f*(a+x*(b*f^2+e^2*x)/f^2)^(1/2)))/e/(-b*f^2+2*d*e)^2
```

Mathematica [A] (verified)

Time = 1.04 (sec) , antiderivative size = 266, normalized size of antiderivative = 1.24

$$\int \frac{1}{d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}} dx$$

$$= \frac{2e^2(2de - bf^2)x + 2ef(-2de + bf^2)\sqrt{a + x\left(b + \frac{e^2x}{f^2}\right)} - (-2de + bf^2)^2 \log\left(e\left(\sqrt{a}f + ex - f\sqrt{a + x\left(b + \frac{e^2x}{f^2}\right)}\right)\right)}{f^2}$$

input

```
Integrate[(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^(-1),x]
```

output

```
(2*e^2*(2*d*e - b*f^2)*x + 2*e*f*(-2*d*e + b*f^2)*Sqrt[a + x*(b + (e^2*x)/f^2)] - (-2*d*e + b*f^2)^2*Log[e*(Sqrt[a]*f + e*x - f*Sqrt[a + x*(b + (e^2*x)/f^2)])] - (4*a*e^2*f^2 - b^2*f^4)*Log[-(Sqrt[a]*f) + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]] + 4*e*(d^2*e - b*d*f^2 + a*e*f^2)*Log[-(a*f^2) + d*e*x - b*f^2*x - d*f*Sqrt[a + x*(b + (e^2*x)/f^2)] + Sqrt[a]*f*(d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])]/(2*e*(-2*d*e + b*f^2)^2)
```

Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.03, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$, Rules used = {2541, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex} dx$$

↓ 2541

$$2 \int \frac{ed^2 - bf^2d + aef^2 + e\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)^2 - (2de - bf^2)\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)}{\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)\left(-bf^2 + 2de - 2e\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)\right)^2} d\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)$$

↓ 1195

$$2 \int \left(\frac{ed^2 - bf^2d + aef^2}{(2de - bf^2)^2 \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a} \right)} + \frac{4ae^2f^2 - b^2f^4}{2(2de - bf^2)^2 \left(-bf^2 + 2de - 2e \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a} \right) \right)} \right) dx$$

↓ 2009

$$2 \left(\frac{f^2(4ae^2 - b^2f^2)}{4e(2de - bf^2) \left(-2e \left(f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex \right) - bf^2 + 2de \right)} - \frac{f^2(4ae^2 - b^2f^2) \log \left(-2e \left(f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex \right) - bf^2 + 2de \right)}{4e(2de - bf^2)^2} \right)$$

input `Int[(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^(-1),x]`

output `2*((f^2*(4*a*e^2 - b^2*f^2))/(4*e*(2*d*e - b*f^2)*(2*d*e - b*f^2 - 2*e*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]))) + ((d^2*e - b*d*f^2 + a*e*f^2)*Log[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]])/(2*d*e - b*f^2)^2 - (f^2*(4*a*e^2 - b^2*f^2)*Log[2*d*e - b*f^2 - 2*e*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]]))/(4*e*(2*d*e - b*f^2)^2)`

Defintions of rubi rules used

rule 1195 `Int[((d_.) + (e_.)*(x_))^(m_.)*((f_.) + (g_.)*(x_))^(n_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2541 `Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]))^(n_.)^(p_.), x_Symbol] := Simp[2 Subst[Int[(g + h*x^n)^p*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)/(-2*d*e + b*f^2 + 2*e*x)^2], x], x, d + e*x + f*Sqrt[a + b*x + c*x^2], x] /; FreeQ[{a, b, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1262 vs. $2(205) = 410$.

Time = 0.05 (sec) , antiderivative size = 1263, normalized size of antiderivative = 5.87

method	result	size
default	Expression too large to display	1263

input `int(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2)),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & f/(b*f^2-2*d*e)*((e^2*(x+(a*f^2-d^2)/(b*f^2-2*d*e))^2/f^2-(-b^2*f^4+2*a*e^2*f^2+2*b*d*e*f^2-2*d^2*e^2)/f^2/(b*f^2-2*d*e)*(x+(a*f^2-d^2)/(b*f^2-2*d*e))) \\ &)+(a^2*e^2*f^4-2*a*b*d*e*f^4+b^2*d^2*f^4+2*a*d^2*e^2*f^2-2*b*d^3*e*f^2+d^4*e^2)/f^2/(b*f^2-2*d*e)^2)^{(1/2)}-1/2*(-b^2*f^4+2*a*e^2*f^2+2*b*d*e*f^2-2*d^2*e^2)/f^2/(b*f^2-2*d*e)*\ln((-1/2*(-b^2*f^4+2*a*e^2*f^2+2*b*d*e*f^2-2*d^2*e^2)/f^2/(b*f^2-2*d*e)+e^2/f^2*(x+(a*f^2-d^2)/(b*f^2-2*d*e)))/(e^2/f^2)^{(1/2)}+(e^2*(x+(a*f^2-d^2)/(b*f^2-2*d*e))^2/f^2-(-b^2*f^4+2*a*e^2*f^2+2*b*d*e*f^2-2*d^2*e^2)/f^2/(b*f^2-2*d*e)*(x+(a*f^2-d^2)/(b*f^2-2*d*e)))+(a^2*e^2*f^4-2*a*b*d*e*f^4+b^2*d^2*f^4+2*a*d^2*e^2*f^2-2*b*d^3*e*f^2+d^4*e^2)/f^2/(b*f^2-2*d*e)^2)^{(1/2)})/(e^2/f^2)^{(1/2)}-(a^2*e^2*f^4-2*a*b*d*e*f^4+b^2*d^2*f^4+2*a*d^2*e^2*f^2-2*b*d^3*e*f^2+d^4*e^2)/f^2/(b*f^2-2*d*e)^2/((a^2*e^2*f^4-2*a*b*d*e*f^4+b^2*d^2*f^4+2*a*d^2*e^2*f^2-2*b*d^3*e*f^2+d^4*e^2)/f^2/(b*f^2-2*d*e)^2)^{(1/2)}*\ln((2*(a^2*e^2*f^4-2*a*b*d*e*f^4+b^2*d^2*f^4+2*a*d^2*e^2*f^2-2*b*d^3*e*f^2+d^4*e^2)/f^2/(b*f^2-2*d*e)^2-(-b^2*f^4+2*a*e^2*f^2+2*b*d*e*f^2-2*d^2*e^2)/f^2/(b*f^2-2*d*e)*(x+(a*f^2-d^2)/(b*f^2-2*d*e)))+2*((a^2*e^2*f^4-2*a*b*d*e*f^4+b^2*d^2*f^4+2*a*d^2*e^2*f^2-2*b*d^3*e*f^2+d^4*e^2)/f^2/(b*f^2-2*d*e)^2)^{(1/2)}*(e^2*(x+(a*f^2-d^2)/(b*f^2-2*d*e))^2/f^2-(-b^2*f^4+2*a*e^2*f^2+2*b*d*e*f^2-2*d^2*e^2)/f^2/(b*f^2-2*d*e)*(x+(a*f^2-d^2)/(b*f^2-2*d*e)))+(a^2*e^2*f^4-2*a*b*d*e*f^4+b^2*d^2*f^4+2*a*d^2*e^2*f^2-2*b*d^3*e*f^2+d^4*e^2)/f^2/(b*f^2-2*d*e)^2)^{(1/2)})/(x+(a*f^2-d^2)/(b*f^2-... \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 2.24 (sec) , antiderivative size = 371, normalized size of antiderivative = 1.73

$$\int \frac{1}{d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}} dx =$$

$$\frac{2 (be^2f^2 - 2de^3)x - 2(d^2e^2 - (bde - ae^2)f^2) \log\left((bd - 2ae)f^2 - (bef^2 - 2de^2)x + (bf^3 - 2def)\sqrt{\dots}\right)}{\dots}$$

input `integrate(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2)),x, algorithm="fricas")`

output `-1/2*(2*(b*e^2*f^2 - 2*d*e^3)*x - 2*(d^2*e^2 - (b*d*e - a*e^2)*f^2)*log((b*d - 2*a*e)*f^2 - (b*e*f^2 - 2*d*e^2)*x + (b*f^3 - 2*d*e*f)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2)) - 2*(d^2*e^2 - (b*d*e - a*e^2)*f^2)*log(a*f^2 - d^2 + (b*f^2 - 2*d*e)*x) + (b^2*f^4 + 2*d^2*e^2 - 2*(b*d*e + a*e^2)*f^2)*log(-b*f^2 - 2*e^2*x + 2*e*f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2)) + 2*(d^2*e^2 - (b*d*e - a*e^2)*f^2)*log(-e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) - d) - 2*(b*e*f^3 - 2*d*e^2*f)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) / (b^2*e*f^4 - 4*b*d*e^2*f^2 + 4*d^2*e^3)`

Sympy [F]

$$\int \frac{1}{d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}} dx = \int \frac{1}{d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}} dx$$

input `integrate(1/(d+e*x+f*(a+b*x+e**2*x**2/f**2)**(1/2)),x)`

output `Integral(1/(d + e*x + f*sqrt(a + b*x + e**2*x**2/f**2)), x)`

Maxima [F]

$$\int \frac{1}{d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}} dx = \int \frac{1}{ex + \sqrt{bx + \frac{e^2x^2}{f^2} + af + d}} dx$$

input `integrate(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2)),x, algorithm="maxima")`

output `integrate(1/(e*x + sqrt(b*x + e^2*x^2/f^2 + a)*f + d), x)`

Giac [F(-1)]

Timed out.

$$\int \frac{1}{d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}} dx = \text{Timed out}$$

input `integrate(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2)),x, algorithm="giac")`

output `Timed out`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}} dx = \int \frac{1}{d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}} dx$$

input `int(1/(d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2)),x)`

output `int(1/(d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2)), x)`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 1760, normalized size of antiderivative = 8.19

$$\int \frac{1}{d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}} dx = \text{Too large to display}$$

input `int(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2)),x)`

output

```
( - 4*sqrt(4*a*e**2 - b**2*f**2)*sqrt( - 4*a*e**2 + b**2*f**2)*atan((8*sqrt(a*f**2 + b*f**2*x + e**2*x**2))*a*e**3 - 2*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b**2*e*f**2 + 4*a*b*e**2*f**2 + 8*a*e**4*x - b**3*f**4 - 2*b**2*e**2*f**2*x)/(sqrt(4*a*e**2 - b**2*f**2)*sqrt( - 4*a*e**2 + b**2*f**2)*b*f**2 - 2*sqrt(4*a*e**2 - b**2*f**2)*sqrt( - 4*a*e**2 + b**2*f**2)*d*e))*a*e**2*f**2 + 4*sqrt(4*a*e**2 - b**2*f**2)*sqrt( - 4*a*e**2 + b**2*f**2)*atan((8*sqrt(a*f**2 + b*f**2*x + e**2*x**2))*a*e**3 - 2*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b**2*e*f**2 + 4*a*b*e**2*f**2 + 8*a*e**4*x - b**3*f**4 - 2*b**2*e**2*f**2*x)/(sqrt(4*a*e**2 - b**2*f**2)*sqrt( - 4*a*e**2 + b**2*f**2)*b*f**2 - 2*sqrt(4*a*e**2 - b**2*f**2)*sqrt( - 4*a*e**2 + b**2*f**2)*d*e))*b*d*e*f**2 - 4*sqrt(4*a*e**2 - b**2*f**2)*sqrt( - 4*a*e**2 + b**2*f**2)*atan((8*sqrt(a*f**2 + b*f**2*x + e**2*x**2))*a*e**3 - 2*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b**2*e*f**2 + 4*a*b*e**2*f**2 + 8*a*e**4*x - b**3*f**4 - 2*b**2*e**2*f**2*x)/(sqrt(4*a*e**2 - b**2*f**2)*sqrt( - 4*a*e**2 + b**2*f**2)*b*f**2 - 2*sqrt(4*a*e**2 - b**2*f**2)*sqrt( - 4*a*e**2 + b**2*f**2)*d*e))*d**2*e**2 + 8*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*a*b*e**3*f**2 - 16*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*a*d*e**4 - 2*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b**3*e*f**4 + 4*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b**2*d*e**2*f**2 + 8*log(4*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b*e*f**2 + 8*sqrt(a*f**2 + b*f**2*x + e**2*x**2))*e**3*x + 4*a*e**2*f**2 + 4*b*d*e*f**2 + 8*b*e**2*f...
```

$$3.11 \quad \int \frac{1}{\left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^2} dx$$

Optimal result	114
Mathematica [A] (verified)	115
Rubi [A] (verified)	116
Maple [B] (verified)	117
Fricas [B] (verification not implemented)	118
Sympy [F]	118
Maxima [F]	119
Giac [B] (verification not implemented)	119
Mupad [F(-1)]	120
Reduce [B] (verification not implemented)	121

Optimal result

Integrand size = 28, antiderivative size = 266

$$\begin{aligned} & \int \frac{1}{\left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^2} dx \\ &= -\frac{2(d^2e - bdf^2 + aef^2)}{(2de - bf^2)^2 \left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)} \\ & \quad - \frac{f^2(4ae^2 - b^2f^2)}{(2de - bf^2)^2 \left(bf^2 + 2e\left(ex+f\sqrt{a+\frac{x(bf^2+e^2x)}{f^2}}\right)\right)} \\ & \quad + \frac{2f^2(4ae^2 - b^2f^2) \log\left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)}{(2de - bf^2)^3} \\ & \quad - \frac{2f^2(4ae^2 - b^2f^2) \log\left(bf^2 + 2e\left(ex+f\sqrt{a+\frac{x(bf^2+e^2x)}{f^2}}\right)\right)}{(2de - bf^2)^3} \end{aligned}$$

output

```
(-2*a*e*f^2+2*b*d*f^2-2*d^2*e)/(-b*f^2+2*d*e)^2/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))-f^2*(-b^2*f^2+4*a*e^2)/(-b*f^2+2*d*e)^2/(b*f^2+2*e*(e*x+f*(a+x*(b*f^2+e^2*x)/f^2)^(1/2)))+2*f^2*(-b^2*f^2+4*a*e^2)*ln(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))/(-b*f^2+2*d*e)^3-2*f^2*(-b^2*f^2+4*a*e^2)*ln(b*f^2+2*e*(e*x+f*(a+x*(b*f^2+e^2*x)/f^2)^(1/2)))/(-b*f^2+2*d*e)^3
```

Mathematica [A] (verified)

Time = 2.17 (sec) , antiderivative size = 427, normalized size of antiderivative = 1.61

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^2} dx$$

$$= \frac{2(bf^3(-d + ex) + 2ef(af^2 - dex))\sqrt{a + x\left(b + \frac{e^2x}{f^2}\right)}}{(-2de + bf^2)^2(d^2 + 2dex - f^2(a + bx))}$$

$$- \frac{2(4a^2e^3f^2x + bdx(2d^2e^2 - 2bdef^2 + b^2f^4 + 2de^3x - be^2f^2x) + a(4d^3e^2 + 2be^3f^2x^2 + d^2(-4bef^2 + 4e^3x) + bd - 2ae)(-2de + bf^2)^2(-d^2 - 2dex + f^2(a + bx)))}{(bd - 2ae)(-2de + bf^2)^2(-d^2 - 2dex + f^2(a + bx))}$$

$$- \frac{2(4ae^2f^2 - b^2f^4)\log\left(-\sqrt{a}f + ex + f\sqrt{a + x\left(b + \frac{e^2x}{f^2}\right)}\right)}{(2de - bf^2)^3}$$

$$+ \frac{2(4ae^2f^2 - b^2f^4)\log\left(-af^2 + dex - bf^2x - df\sqrt{a + x\left(b + \frac{e^2x}{f^2}\right)} + \sqrt{a}f\left(d + ex + f\sqrt{a + x\left(b + \frac{e^2x}{f^2}\right)}\right)\right)}{(2de - bf^2)^3}$$

input

```
Integrate[(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^(-2), x]
```

output

```
(2*(b*f^3*(-d + e*x) + 2*e*f*(a*f^2 - d*e*x))*Sqrt[a + x*(b + (e^2*x)/f^2)
])/((-2*d*e + b*f^2)^2*(d^2 + 2*d*e*x - f^2*(a + b*x))) - (2*(4*a^2*e^3*f^
2*x + b*d*x*(2*d^2*e^2 - 2*b*d*e*f^2 + b^2*f^4 + 2*d*e^3*x - b*e^2*f^2*x)
+ a*(4*d^3*e^2 + 2*b*e^3*f^2*x^2 + d^2*(-4*b*e*f^2 + 4*e^3*x) + d*(b^2*f^4
- 6*b*e^2*f^2*x - 4*e^4*x^2)))/((b*d - 2*a*e)*(-2*d*e + b*f^2)^2*(-d^2 -
2*d*e*x + f^2*(a + b*x))) - (2*(4*a*e^2*f^2 - b^2*f^4)*Log[-(Sqrt[a]*f) +
e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2])])/(2*d*e - b*f^2)^3 + (2*(4*a*e^2*f
^2 - b^2*f^4)*Log[-(a*f^2) + d*e*x - b*f^2*x - d*f*Sqrt[a + x*(b + (e^2*x)
/f^2)] + Sqrt[a]*f*(d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])])/(2*d*e -
b*f^2)^3
```

Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.03, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$, Rules used = {2541, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\left(f\sqrt{a+bx+\frac{e^2x^2}{f^2}}+d+ex\right)^2} dx$$

↓ 2541

$$2 \int \frac{ed^2 - bf^2d + aef^2 + e\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)^2 - (2de - bf^2)\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)}{\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)^2 \left(-bf^2 + 2de - 2e\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)\right)^2} d\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)$$

↓ 1195

$$2 \int \left(\frac{ed^2 - bf^2d + aef^2}{(2de - bf^2)^2 \left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)^2} + \frac{4ae^2f^2 - b^2f^4}{(2de - bf^2)^3 \left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)} + \frac{1}{(2de - bf^2)^3} \right) d\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)$$

↓ 2009

$$2 \left(\frac{f^2(4ae^2 - b^2f^2)}{2(2de - bf^2)^2 \left(-2e\left(f\sqrt{a+bx+\frac{e^2x^2}{f^2}}+d+ex\right) - bf^2 + 2de\right)} + \frac{f^2(4ae^2 - b^2f^2) \log\left(f\sqrt{a+bx+\frac{e^2x^2}{f^2}}+d+ex\right)}{(2de - bf^2)^3} \right)$$

input `Int[(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^(-2),x]`

output `2*(-((d^2*e - b*d*f^2 + a*e*f^2)/((2*d*e - b*f^2)^2*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]))) + (f^2*(4*a*e^2 - b^2*f^2))/(2*(2*d*e - b*f^2)^2*(2*d*e - b*f^2 - 2*e*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]))) + (f^2*(4*a*e^2 - b^2*f^2)*Log[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]])/(2*d*e - b*f^2)^3 - (f^2*(4*a*e^2 - b^2*f^2)*Log[2*d*e - b*f^2 - 2*e*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]])/(2*d*e - b*f^2)^3)`

Defintions of rubi rules used

rule 1195

```
Int[((d_.) + (e_.)*(x_))^(m_.)*((f_.) + (g_.)*(x_))^(n_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2541

```
Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]))^(n_.))^p, x_Symbol] := Simp[2 Subst[Int[(g + h*x^n)^p*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)/(-2*d*e + b*f^2 + 2*e*x)^2), x], x, d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 6304 vs. $2(259) = 518$.

Time = 0.07 (sec) , antiderivative size = 6305, normalized size of antiderivative = 23.70

method	result	size
default	Expression too large to display	6305

input

```
int(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^2,x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 826 vs. $2(251) = 502$.

Time = 1.36 (sec) , antiderivative size = 826, normalized size of antiderivative = 3.11

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^2} dx = \text{Too large to display}$$

input `integrate(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^2,x, algorithm="fricas")`

output

```
-1/2*(a*b^2*f^6 + (3*b^2*d^2 - 14*a*b*d*e + 8*a^2*e^2)*f^4 - 2*(b*d^3*e -
4*a*d^2*e^2)*f^2 - 4*(b^2*e^2*f^4 - 4*b*d*e^3*f^2 + 4*d^2*e^4)*x^2 + (b^3*
f^6 - 8*b^2*d*e*f^4 + 20*b*d^2*e^2*f^2 - 16*d^3*e^3)*x - 2*(a*b^2*f^6 + 4*
a*d^2*e^2*f^2 - (b^2*d^2 + 4*a^2*e^2)*f^4 + (b^3*f^6 + 8*a*d*e^3*f^2 - 2*(
b^2*d*e + 2*a*b*e^2)*f^4)*x)*log(-4*a*d*e^2*f^2 - (b^2*d - 4*a*b*e)*f^4 +
4*(b*e^3*f^2 - 2*d*e^4)*x^2 + (3*b^2*e*f^4 - 4*(2*b*d*e^2 - a*e^3)*f^2)*x
- (b^2*f^5 - 4*(b*d*e - a*e^2)*f^3 + 4*(b*e^2*f^3 - 2*d*e^3*f)*x)*sqrt((b*
f^2*x + e^2*x^2 + a*f^2)/f^2)) - 2*(a*b^2*f^6 + 4*a*d^2*e^2*f^2 - (b^2*d^2
+ 4*a^2*e^2)*f^4 + (b^3*f^6 + 8*a*d*e^3*f^2 - 2*(b^2*d*e + 2*a*b*e^2)*f^4
)*x)*log(a*f^2 - d^2 + (b*f^2 - 2*d*e)*x) + 2*(a*b^2*f^6 + 4*a*d^2*e^2*f^2
- (b^2*d^2 + 4*a^2*e^2)*f^4 + (b^3*f^6 + 8*a*d*e^3*f^2 - 2*(b^2*d*e + 2*a
*b*e^2)*f^4)*x)*log(-e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) - d) -
4*((b^2*d - 2*a*b*e)*f^5 - 2*(b*d^2*e - 2*a*d*e^2)*f^3 - (b^2*e*f^5 - 4*b*
d*e^2*f^3 + 4*d^2*e^3*f)*x)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2))/(a*b^3*
f^8 + 8*d^5*e^3 - (b^3*d^2 + 6*a*b^2*d*e)*f^6 + 6*(b^2*d^3*e + 2*a*b*d^2*e
^2)*f^4 - 4*(3*b*d^4*e^2 + 2*a*d^3*e^3)*f^2 + (b^4*f^8 - 8*b^3*d*e*f^6 + 2
4*b^2*d^2*e^2*f^4 - 32*b*d^3*e^3*f^2 + 16*d^4*e^4)*x)
```

Sympy [F]

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^2} dx = \int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^2} dx$$

input `integrate(1/(d+e*x+f*(a+b*x+e**2*x**2/f**2)**(1/2))**2,x)`

output `Integral((d + e*x + f*sqrt(a + b*x + e**2*x**2/f**2))**(-2), x)`

Maxima [F]

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^2} dx = \int \frac{1}{\left(ex + \sqrt{bx + \frac{e^2x^2}{f^2} + af + d}\right)^2} dx$$

input `integrate(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^2,x, algorithm="maxima")`

output `integrate((e*x + sqrt(b*x + e^2*x^2/f^2 + a)*f + d)^(-2), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1618 vs. $2(251) = 502$.

Time = 4.93 (sec) , antiderivative size = 1618, normalized size of antiderivative = 6.08

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^2} dx = \text{Too large to display}$$

input `integrate(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^2,x, algorithm="giac")`

output

```

2*e^2*x/(b^2*f^4 - 4*b*d*e*f^2 + 4*d^2*e^2) + 1/5*(b^2*e*f^3*abs(f) - 4*a*
e^3*f*abs(f))*log(abs((x*abs(e) - sqrt(b*f^2*x + e^2*x^2 + a*f^2))^4*b^3*f
^6 - 2*(x*abs(e) - sqrt(b*f^2*x + e^2*x^2 + a*f^2))^2*b^3*d^2*f^6 + b^3*d^
4*f^6 + 4*(x*abs(e) - sqrt(b*f^2*x + e^2*x^2 + a*f^2))^2*a*b^2*d*e*f^6 - 4
*a*b^2*d^3*e*f^6 + 4*(x*abs(e) - sqrt(b*f^2*x + e^2*x^2 + a*f^2))^2*a^2*b*
e^2*f^6 + 4*a^2*b*d^2*e^2*f^6 + 4*(x*abs(e) - sqrt(b*f^2*x + e^2*x^2 + a*f
^2))^3*a*b^2*f^6*abs(e) - 4*(x*abs(e) - sqrt(b*f^2*x + e^2*x^2 + a*f^2))*a
*b^2*d^2*f^6*abs(e) + 8*(x*abs(e) - sqrt(b*f^2*x + e^2*x^2 + a*f^2))*a^2*b
*d*e*f^6*abs(e) - 4*(x*abs(e) - sqrt(b*f^2*x + e^2*x^2 + a*f^2))^4*b^2*d*e
*f^4 + 4*(x*abs(e) - sqrt(b*f^2*x + e^2*x^2 + a*f^2))^2*b^2*d^3*e*f^4 + 8*
(x*abs(e) - sqrt(b*f^2*x + e^2*x^2 + a*f^2))^4*a*b*e^2*f^4 - 24*(x*abs(e)
- sqrt(b*f^2*x + e^2*x^2 + a*f^2))^2*a*b*d^2*e^2*f^4 + 16*(x*abs(e) - sqrt
(b*f^2*x + e^2*x^2 + a*f^2))^2*a^2*d*e^3*f^4 + 2*(x*abs(e) - sqrt(b*f^2*x
+ e^2*x^2 + a*f^2))^5*b^2*f^4*abs(e) - 8*(x*abs(e) - sqrt(b*f^2*x + e^2*x^
2 + a*f^2))^3*b^2*d^2*f^4*abs(e) + 6*(x*abs(e) - sqrt(b*f^2*x + e^2*x^2 +
a*f^2))*b^2*d^4*f^4*abs(e) - 16*(x*abs(e) - sqrt(b*f^2*x + e^2*x^2 + a*f^2
))*a*b*d^3*e*f^4*abs(e) + 8*(x*abs(e) - sqrt(b*f^2*x + e^2*x^2 + a*f^2))^3
*a^2*e^2*f^4*abs(e) + 8*(x*abs(e) - sqrt(b*f^2*x + e^2*x^2 + a*f^2))*a^2*d
^2*e^2*f^4*abs(e) - 4*(x*abs(e) - sqrt(b*f^2*x + e^2*x^2 + a*f^2))^4*b*d^2
*e^2*f^2 + 12*(x*abs(e) - sqrt(b*f^2*x + e^2*x^2 + a*f^2))^2*b*d^4*e^2*...

```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^2} dx = \int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^2} dx$$

input

```
int(1/(d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^2,x)
```

output

```
int(1/(d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^2, x)
```

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 2122, normalized size of antiderivative = 7.98

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^2} dx = \text{Too large to display}$$

input `int(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^2,x)`

output

```
(2*(-2*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*a**2*b*e*f**6 + 4*sqrt(a*f**2
+ b*f**2*x + e**2*x**2)*a**2*d*e**2*f**4 + sqrt(a*f**2 + b*f**2*x + e**2*
x**2)*a*b**2*d*f**6 - sqrt(a*f**2 + b*f**2*x + e**2*x**2)*a*b**2*e*f**6*x
+ 4*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*a*b*d*e**2*f**4*x - 4*sqrt(a*f**2
+ b*f**2*x + e**2*x**2)*a*d**3*e**2*f**2 - 4*sqrt(a*f**2 + b*f**2*x + e**2
*x**2)*a*d**2*e**3*f**2*x - sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b**2*d**3*
f**4 + sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b**2*d**2*e*f**4*x + 2*sqrt(a*f
**2 + b*f**2*x + e**2*x**2)*b*d**4*e*f**2 - 4*sqrt(a*f**2 + b*f**2*x + e**
2*x**2)*b*d**3*e**2*f**2*x + 4*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*d**4*e*
*3*x - 4*log(-sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b*f**2 + 2*sqrt(a*f**2
+ b*f**2*x + e**2*x**2)*d*e + 2*a*e*f**2 - b*d*f**2 + b*e*f**2*x - 2*d*e*
*2*x)*a**3*e**2*f**6 + log(-sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b*f**2 +
2*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*d*e + 2*a*e*f**2 - b*d*f**2 + b*e*f
**2*x - 2*d*e**2*x)*a**2*b**2*f**8 - 4*log(-sqrt(a*f**2 + b*f**2*x + e**
2*x**2)*b*f**2 + 2*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*d*e + 2*a*e*f**2 -
b*d*f**2 + b*e*f**2*x - 2*d*e**2*x)*a**2*b*e**2*f**6*x + 8*log(-sqrt(a*f
**2 + b*f**2*x + e**2*x**2)*b*f**2 + 2*sqrt(a*f**2 + b*f**2*x + e**2*x**2)
*d*e + 2*a*e*f**2 - b*d*f**2 + b*e*f**2*x - 2*d*e**2*x)*a**2*d**2*e**2*f**
4 + 8*log(-sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b*f**2 + 2*sqrt(a*f**2 +
b*f**2*x + e**2*x**2)*d*e + 2*a*e*f**2 - b*d*f**2 + b*e*f**2*x - 2*d*e...
```

$$3.12 \quad \int \frac{1}{\left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^3} dx$$

Optimal result	122
Mathematica [A] (verified)	123
Rubi [A] (verified)	123
Maple [B] (verified)	125
Fricas [B] (verification not implemented)	126
Sympy [F]	127
Maxima [F]	127
Giac [F(-1)]	127
Mupad [F(-1)]	128
Reduce [B] (verification not implemented)	128

Optimal result

Integrand size = 28, antiderivative size = 330

$$\begin{aligned} & \int \frac{1}{\left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^3} dx \\ &= -\frac{d^2e - bdf^2 + aef^2}{(2de - bf^2)^2 \left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^2} \\ & \quad - \frac{2f^2(4ae^2 - b^2f^2)}{(2de - bf^2)^3 \left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)} \\ & \quad - \frac{2ef^2(4ae^2 - b^2f^2)}{(2de - bf^2)^3 \left(bf^2 + 2e\left(ex+f\sqrt{a+\frac{x(bf^2+e^2x)}{f^2}}\right)\right)} \\ & \quad + \frac{6ef^2(4ae^2 - b^2f^2) \log\left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)}{(2de - bf^2)^4} \\ & \quad - \frac{6ef^2(4ae^2 - b^2f^2) \log\left(bf^2 + 2e\left(ex+f\sqrt{a+\frac{x(bf^2+e^2x)}{f^2}}\right)\right)}{(2de - bf^2)^4} \end{aligned}$$

output

$$\begin{aligned}
 & -\frac{(a*ef^2-b*d*f^2+d^2*e)}{(-b*f^2+2*d*e)^2/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^{(1/2}))^2-2*f^2*(-b^2*f^2+4*a*e^2)/(-b*f^2+2*d*e)^3/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^{(1/2}))} \\
 & -2*e*f^2*(-b^2*f^2+4*a*e^2)/(-b*f^2+2*d*e)^3/(b*f^2+2*e*(e*x+f*(a+x*(b*f^2+e^2*x)/f^2)^{(1/2}))) \\
 & +6*e*f^2*(-b^2*f^2+4*a*e^2)*\ln(d+e*x+f*(a+b*x+e^2*x^2/f^2)^{(1/2}))/(-b*f^2+2*d*e)^4 \\
 & -6*e*f^2*(-b^2*f^2+4*a*e^2)*\ln(b*f^2+2*e*(e*x+f*(a+x*(b*f^2+e^2*x)/f^2)^{(1/2}))/(-b*f^2+2*d*e)^4
 \end{aligned}$$

Mathematica [A] (verified)

Time = 10.85 (sec) , antiderivative size = 300, normalized size of antiderivative = 0.91

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^3} dx = \frac{(-2de+bf^2)^2(d^2e-bdf^2+ae^2f^2)}{\left(d+ex+f\sqrt{a+x\left(b+\frac{e^2x}{f^2}\right)}\right)^2} + \frac{2f^2(-2de+bf^2)(-4ae^2+b^2f^2)}{d+ex+f\sqrt{a+x\left(b+\frac{e^2x}{f^2}\right)}} + \frac{2ef^2(2de-bf^2)(4ae^2-b^2f^2)}{bf^2+2e\left(ex+f\sqrt{a+x\left(b+\frac{e^2x}{f^2}\right)}\right)} - 6ef^2(4ae^2 - b^2f^2) \ln\left(\frac{d+ex+f\sqrt{a+x\left(b+\frac{e^2x}{f^2}\right)}}{d+ex+f\sqrt{a+x\left(b+\frac{e^2x}{f^2}\right)}}\right) - 6ef^2(4ae^2 - b^2f^2) \ln\left(\frac{d+ex+f\sqrt{a+x\left(b+\frac{e^2x}{f^2}\right)}}{d+ex+f\sqrt{a+x\left(b+\frac{e^2x}{f^2}\right)}}\right)$$

input

```
Integrate[(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^(-3), x]
```

output

$$\begin{aligned}
 & -\left(\frac{(-2*d*e + b*f^2)^2*(d^2*e - b*d*f^2 + a*e*f^2)}{(d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])^2 + (2*f^2*(-2*d*e + b*f^2)*(-4*a*e^2 + b^2*f^2))/(d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]) + (2*e*f^2*(2*d*e - b*f^2)*(4*a*e^2 - b^2*f^2))/(b*f^2 + 2*e*(e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]))} - 6*e*f^2*(4*a*e^2 - b^2*f^2)*Log[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]] + 6*e*f^2*(4*a*e^2 - b^2*f^2)*Log[-(b*f^2) - 2*e*(e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])]\right)/(-2*d*e + b*f^2)^4
 \end{aligned}$$

Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 339, normalized size of antiderivative = 1.03, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$, Rules used = {2541, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\left(f\sqrt{a+bx+\frac{e^2x^2}{f^2}}+d+ex\right)^3} dx$$

↓ 2541

$$2 \int \frac{ed^2 - bf^2d + aef^2 + e\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)^2 - (2de - bf^2)\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)}{\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)^3 \left(-bf^2 + 2de - 2e\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)\right)^2} d\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)$$

↓ 1195

$$2 \int \left(\frac{ed^2 - bf^2d + aef^2}{(2de - bf^2)^2 \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)^3} + \frac{3(4ae^3f^2 - b^2ef^4)}{(2de - bf^2)^4 \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)} + \frac{1}{(2de - bf^2)^4} \right) d\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)$$

↓ 2009

$$2 \left(-\frac{f^2(4ae^2 - b^2f^2)}{(2de - bf^2)^3 \left(f\sqrt{a+bx+\frac{e^2x^2}{f^2}}+d+ex\right)} + \frac{ef^2(4ae^2 - b^2f^2)}{(2de - bf^2)^3 \left(-2e\left(f\sqrt{a+bx+\frac{e^2x^2}{f^2}}+d+ex\right) - bf^2 + 2de\right)} \right)$$

input `Int[(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^(-3),x]`

output `2*(-1/2*(d^2*e - b*d*f^2 + a*e*f^2)/((2*d*e - b*f^2)^2*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^2) - (f^2*(4*a*e^2 - b^2*f^2))/((2*d*e - b*f^2)^3*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])) + (e*f^2*(4*a*e^2 - b^2*f^2))/((2*d*e - b*f^2)^3*(2*d*e - b*f^2 - 2*e*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]))) + (3*e*f^2*(4*a*e^2 - b^2*f^2)*Log[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]])/(2*d*e - b*f^2)^4 - (3*e*f^2*(4*a*e^2 - b^2*f^2)*Log[2*d*e - b*f^2 - 2*e*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]])/(2*d*e - b*f^2)^4)`

Defintions of rubi rules used

rule 1195

```
Int[((d_.) + (e_.)*(x_)^(m_.))*((f_.) + (g_.)*(x_)^(n_.))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2541

```
Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]))^(n_.))^p, x_Symbol] := Simp[2 Subst[Int[(g + h*x^n)^p*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)/(-2*d*e + b*f^2 + 2*e*x)^2), x], x, d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 29136 vs. $2(320) = 640$.

Time = 0.17 (sec) , antiderivative size = 29137, normalized size of antiderivative = 88.29

method	result	size
default	Expression too large to display	29137

input

```
int(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^3,x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1954 vs. $2(311) = 622$.

Time = 10.87 (sec) , antiderivative size = 1954, normalized size of antiderivative = 5.92

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^3} dx = \text{Too large to display}$$

input `integrate(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^3,x, algorithm="fricas")`

output

```
((3*a*b^3*d - 4*a^2*b^2*e)*f^8 - (b^3*d^3 + 4*a*b^2*d^2*e + 10*a^2*b*d*e^2
- 20*a^3*e^3)*f^6 - 4*(b^2*d^4*e - 8*a*b*d^3*e^2 + 6*a^2*d^2*e^3)*f^4 - 4
*(b^3*e^3*f^6 - 6*b^2*d*e^4*f^4 + 12*b*d^2*e^5*f^2 - 8*d^3*e^6)*x^3 + 2*(b
*d^5*e^2 - 6*a*d^4*e^3)*f^2 - (b^4*e*f^8 - 2*a*b^2*e^3*f^6 - 40*d^4*e^5 -
2*(11*b^2*d^2*e^3 - 4*a*b*d*e^4)*f^4 + 8*(7*b*d^3*e^4 - a*d^2*e^5)*f^2)*x^
2 + (16*d^5*e^4 + (3*b^4*d - 5*a*b^3*e)*f^8 - (7*b^3*d^2*e + 10*a*b^2*d*e^
2 - 28*a^2*b*e^3)*f^6 + 2*(5*b^2*d^3*e^2 + 22*a*b*d^2*e^3 - 28*a^2*d*e^4)*
f^4 - 8*(3*b*d^4*e^3 + a*d^3*e^4)*f^2)*x - 3*(a^2*b^2*e*f^8 - 4*a*d^4*e^3*
f^2 - 2*(a*b^2*d^2*e + 2*a^3*e^3)*f^6 + (b^2*d^4*e + 8*a^2*d^2*e^3)*f^4 +
(b^4*e*f^8 - 16*a*d^2*e^5*f^2 - 4*(b^3*d*e^2 + a*b^2*e^3)*f^6 + 4*(b^2*d^2
*e^3 + 4*a*b*d*e^4)*f^4)*x^2 + 2*(a*b^3*e*f^8 - 8*a*d^3*e^4*f^2 - (b^3*d^2
*e + 2*a*b^2*d*e^2 + 4*a^2*b*e^3)*f^6 + 2*(b^2*d^3*e^2 + 2*a*b*d^2*e^3 + 4
*a^2*d*e^4)*f^4)*x)*log(-4*a*d*e^2*f^2 - (b^2*d - 4*a*b*e)*f^4 + 4*(b*e^3*
f^2 - 2*d*e^4)*x^2 + (3*b^2*e*f^4 - 4*(2*b*d*e^2 - a*e^3)*f^2)*x - (b^2*f^
5 - 4*(b*d*e - a*e^2)*f^3 + 4*(b*e^2*f^3 - 2*d*e^3*f)*x)*sqrt((b*f^2*x + e
^2*x^2 + a*f^2)/f^2)) - 3*(a^2*b^2*e*f^8 - 4*a*d^4*e^3*f^2 - 2*(a*b^2*d^2*
e + 2*a^3*e^3)*f^6 + (b^2*d^4*e + 8*a^2*d^2*e^3)*f^4 + (b^4*e*f^8 - 16*a*d
^2*e^5*f^2 - 4*(b^3*d*e^2 + a*b^2*e^3)*f^6 + 4*(b^2*d^2*e^3 + 4*a*b*d*e^4)
*f^4)*x^2 + 2*(a*b^3*e*f^8 - 8*a*d^3*e^4*f^2 - (b^3*d^2*e + 2*a*b^2*d*e^2
+ 4*a^2*b*e^3)*f^6 + 2*(b^2*d^3*e^2 + 2*a*b*d^2*e^3 + 4*a^2*d*e^4)*f^4)...
```

Sympy [F]

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^3} dx = \int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^3} dx$$

input `integrate(1/(d+e*x+f*(a+b*x+e**2*x**2/f**2)**(1/2))**3,x)`

output `Integral((d + e*x + f*sqrt(a + b*x + e**2*x**2/f**2))**(-3), x)`

Maxima [F]

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^3} dx = \int \frac{1}{\left(ex + \sqrt{bx + \frac{e^2x^2}{f^2}} + af + d\right)^3} dx$$

input `integrate(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^3,x, algorithm="maxima")`

output `integrate((e*x + sqrt(b*x + e^2*x^2/f^2 + a)*f + d)^(-3), x)`

Giac [F(-1)]

Timed out.

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^3} dx = \text{Timed out}$$

input `integrate(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^3,x, algorithm="giac")`

output `Timed out`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^3} dx = \int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^3} dx$$

input `int(1/(d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^3,x)`

output `int(1/(d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^3, x)`

Reduce [B] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 6354, normalized size of antiderivative = 19.25

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^3} dx = \text{Too large to display}$$

input `int(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^3,x)`

output

```
(24*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*a**3*b*e**2*f**8 - 48*sqrt(a*f**2
+ b*f**2*x + e**2*x**2)*a**3*d*e**3*f**6 - 4*sqrt(a*f**2 + b*f**2*x + e**2
*x**2)*a**2*b**3*f**10 + 12*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*a**2*b**2*
d*e*f**8 + 36*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*a**2*b**2*e**2*f**8*x -
72*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*a**2*b*d**2*e**2*f**6 - 144*sqrt(a*
f**2 + b*f**2*x + e**2*x**2)*a**2*b*d*e**3*f**6*x + 128*sqrt(a*f**2 + b*f*
**2*x + e**2*x**2)*a**2*d**3*e**3*f**4 + 144*sqrt(a*f**2 + b*f**2*x + e**2*
x**2)*a**2*d**2*e**4*f**4*x - 4*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*a*b**4
*f**10*x + 4*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*a*b**3*d**2*f**8 + 12*sqr
t(a*f**2 + b*f**2*x + e**2*x**2)*a*b**3*d*e*f**8*x + 8*sqrt(a*f**2 + b*f**
2*x + e**2*x**2)*a*b**3*e**2*f**8*x**2 - 48*sqrt(a*f**2 + b*f**2*x + e**2*
x**2)*a*b**2*d**2*e**2*f**6*x - 48*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*a*b
**2*d*e**3*f**6*x**2 + 24*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*a*b*d**4*e**
2*f**4 + 176*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*a*b*d**3*e**3*f**4*x + 96
*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*a*b*d**2*e**4*f**4*x**2 - 80*sqrt(a*f
**2 + b*f**2*x + e**2*x**2)*a*d**5*e**3*f**2 - 192*sqrt(a*f**2 + b*f**2*x
+ e**2*x**2)*a*d**4*e**4*f**2*x - 64*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*a
*d**3*e**5*f**2*x**2 + 4*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b**4*d**2*f**
8*x - 12*sqrt(a*f**2 + b*f**2*x + e**2*x**2)*b**3*d**3*e*f**6*x - 8*sqrt(a
*f**2 + b*f**2*x + e**2*x**2)*b**3*d**2*e**2*f**6*x**2 - 12*sqrt(a*f**2...
```

$$3.13 \quad \int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx$$

Optimal result	130
Mathematica [B] (verified)	131
Rubi [A] (verified)	132
Maple [F]	135
Fricas [A] (verification not implemented)	135
Sympy [F]	136
Maxima [F]	137
Giac [F]	137
Mupad [F(-1)]	137
Reduce [F]	138

Optimal result

Integrand size = 30, antiderivative size = 370

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx = \frac{f^2(2de - bf^2)(4ae^2 - b^2 f^2) \sqrt{d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}}}}{4e^4} + \frac{f^2(4ae^2 - b^2 f^2) \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{3/2}}{12e^3} + \frac{\left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{7/2}}{7e} - \frac{f^2(2de - bf^2)^2(4ae^2 - b^2 f^2) \sqrt{d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}}}}{16e^4 \left(bf^2 + 2e \left(ex + f \sqrt{a + \frac{x(bf^2 + e^2 x)}{f^2}} \right) \right)} - \frac{5f^2(2de - bf^2)^{3/2} (4ae^2 - b^2 f^2) \operatorname{arctanh} \left(\frac{\sqrt{2}\sqrt{e} \sqrt{d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}}}}{\sqrt{2de - bf^2}} \right)}{16\sqrt{2}e^{9/2}}$$

output

```

1/4*f^2*(-b*f^2+2*d*e)*(-b^2*f^2+4*a*e^2)*(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2)/e^4+1/12*f^2*(-b^2*f^2+4*a*e^2)*(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(3/2)/e^3+1/7*(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(7/2)/e-1/16*f^2*(-b*f^2+2*d*e)^2*(-b^2*f^2+4*a*e^2)*(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2)/e^4/(b*f^2+2*e*(e*x+f*(a+x*(b*f^2+e^2*x)/f^2)^(1/2)))-5/32*f^2*(-b*f^2+2*d*e)^(3/2)*(-b^2*f^2+4*a*e^2)*arctanh(2^(1/2)*e^(1/2)*(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2)/(-b*f^2+2*d*e)^(1/2))*2^(1/2)/e^(9/2)

```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 985 vs. $2(370) = 740$.

Time = 6.86 (sec) , antiderivative size = 985, normalized size of antiderivative = 2.66

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx = \text{Too large to display}$$

input

```
Integrate[(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^(5/2),x]
```

output

```

((2*Sqrt[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2))]*(105*b^4*f^8 + 28*b^3*
e*f^6*(-10*d + 3*e*x + 5*f*Sqrt[a + x*(b + (e^2*x)/f^2)]) + 4*b^2*e^2*f^4*
(21*d^2 - 119*a*f^2 + 16*e*x*(2*e*x - f*Sqrt[a + x*(b + (e^2*x)/f^2)]) - 2
*d*(31*e*x + 49*f*Sqrt[a + x*(b + (e^2*x)/f^2)])) + 16*b*e^3*f^2*(3*d^3 +
79*a*d*f^2 + 36*d*e*x*(2*e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]) + 9*d^2*(3
*e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]) + 4*(15*e^3*x^3 - 8*a*f^3*Sqrt[a +
x*(b + (e^2*x)/f^2)] + 9*e^2*f*x^2*Sqrt[a + x*(b + (e^2*x)/f^2)])) + 16*e
^4*(20*a^2*f^4 + 6*(d + 2*e*x)^3*(e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]) +
a*f^2*(-3*d^2 + 4*e*x*(19*e*x + 13*f*Sqrt[a + x*(b + (e^2*x)/f^2)]) + 4*d
*(38*e*x + 29*f*Sqrt[a + x*(b + (e^2*x)/f^2)])))/((21*e^4*(b*f^2 + 2*e*(e
*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])) - (20*b^2*d^2*f^4*ArcTan[(Sqrt[2]*
Sqrt[e]*Sqrt[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])]/Sqrt[-2*d*e + b*f
^2]])/(e^(5/2)*Sqrt[-(d*e) + (b*f^2)/2]) - (80*a*b*d*f^4*ArcTan[(Sqrt[2]*S
qrt[e]*Sqrt[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])]/Sqrt[-2*d*e + b*f^
2]])/(e^(3/2)*Sqrt[-(d*e) + (b*f^2)/2]) + (20*b^3*d*f^6*ArcTan[(Sqrt[2]*Sq
rt[e]*Sqrt[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])]/Sqrt[-2*d*e + b*f^2
]])/(e^(7/2)*Sqrt[-(d*e) + (b*f^2)/2]) + (20*a*b^2*f^6*ArcTan[(Sqrt[2]*Sqr
t[e]*Sqrt[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])]/Sqrt[-2*d*e + b*f^2
]])/(e^(5/2)*Sqrt[-(d*e) + (b*f^2)/2]) - (5*b^4*f^8*ArcTan[(Sqrt[2]*Sqrt[e]
*Sqrt[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])]/Sqrt[-2*d*e + b*f^2]])...

```

Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 377, normalized size of antiderivative = 1.02, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2541, 1192, 1580, 25, 2341, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} + d + ex \right)^{5/2} dx$$

\downarrow 2541

$$\frac{2 \int \frac{\left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)^{5/2} \left(ed^2 - bf^2 d + aef^2 + e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)^2 - (2de - bf^2) \left(d + \right. \right.}{\left. \left. -bf^2 + 2de - 2e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right) \right)^2} dx}{1}$$

↓ 1192

$$4 \int \frac{\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)^3 \left(ed^2 - bf^2d + aef^2 + e\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)^2 - (2de - bf^2)\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)\right)}{\left(-bf^2 + 2de - 2e\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)\right)^2}$$

↓ 1580

$$4 \left(\int \frac{32\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)^4 e^5 - 16(2de-bf^2)\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)^3 e^4 + 8f^2(4ae^2-b^2f^2)\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)^2 e^3 + 4f^2(2de-bf^2)\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right) e^2}{-bf^2+2de-2e\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)} dx \right) \frac{1}{64e^5}$$

↓ 25

$$4 \left(\frac{f^2(4ae^2 - b^2f^2)(2de - bf^2)^2 \sqrt{f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex}}{64e^4 \left(-2e\left(f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex\right) - bf^2 + 2de\right)} - \int \frac{32\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)^4 e^5 - 16(2de-bf^2)\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)^3 e^4 - 4f^2(2de-bf^2)\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right) e^2}{-bf^2+2de-2e\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)} dx \right)$$

↓ 2341

$$4 \left(\frac{f^2(4ae^2 - b^2f^2)(2de - bf^2)^2 \sqrt{f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex}}{64e^4 \left(-2e\left(f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex\right) - bf^2 + 2de\right)} - \int \left(-16\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)^3 e^4 - 4f^2(2de - bf^2)\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right) e^2\right) dx \right)$$

↓ 2009

$$4 \left(\frac{f^2(4ae^2 - b^2f^2)(2de - bf^2)^2 \sqrt{f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex}}{64e^4 \left(-2e\left(f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex\right) - bf^2 + 2de\right)} - \frac{5\sqrt{e}f^2(4ae^2 - b^2f^2)(2de - bf^2)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{e}\sqrt{f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex}}{\sqrt{2}}\right)}{\sqrt{2}} \right)$$

input `Int[(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^(5/2),x]`

output `4*((f^2*(2*d*e - b*f^2)^2*(4*a*e^2 - b^2*f^2)*Sqrt[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]])/(64*e^4*(2*d*e - b*f^2 - 2*e*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])) - (-4*e*f^2*(2*d*e - b*f^2)*(4*a*e^2 - b^2*f^2)*Sqrt[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]] - (4*e^2*f^2*(4*a*e^2 - b^2*f^2)*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^(3/2))/3 - (16*e^4*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^(7/2))/7 + (5*Sqrt[e]*f^2*(2*d*e - b*f^2)^(3/2)*(4*a*e^2 - b^2*f^2)*ArcTanh[(Sqrt[2]*Sqrt[e]*Sqrt[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]])/Sqrt[2*d*e - b*f^2]]/Sqrt[2])/(64*e^5)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 1192 `Int[((d_.) + (e_.)*(x_)^(m_))*((f_.) + (g_.)*(x_)^(n_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Simp[2/e^(n + 2*p + 1) Subst[Int[x^(2*m + 1)*(e*f - d*g + g*x^2)^n*(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 + c*x^4)^p, x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && IGtQ[p, 0] && ILtQ[n, 0] && IntegerQ[m + 1/2]`

rule 1580 `Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := Simp[(-d)^(m/2 - 1)*(c*d^2 - b*d*e + a*e^2)^p*x*((d + e*x^2)^(q + 1)/(2*e^(2*p + m/2)*(q + 1))), x] + Simp[1/(2*e^(2*p + m/2)*(q + 1)) Int[(d + e*x^2)^(q + 1)*ExpandToSum[Together[(1/(d + e*x^2))*(2*e^(2*p + m/2)*(q + 1)*x^m*(a + b*x^2 + c*x^4)^p - (-d)^(m/2 - 1)*(c*d^2 - b*d*e + a*e^2)^p*(d + e*(2*q + 3)*x^2)]], x], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && ILtQ[q, -1] && IGtQ[m/2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2341 `Int[(Pq_)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[Pq*(a + b*x^2)^p, x], x] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

rule 2541

```
Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2])^(n_.))^(p_.), x_Symbol] :> Simp[2 Subst[Int[(g + h*x^n)^p*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)/(-2*d*e + b*f^2 + 2*e*x)^2), x], x, d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]
```

Maple [F]

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{\frac{5}{2}} dx$$

input

```
int((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(5/2),x)
```

output

```
int((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(5/2),x)
```

Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 923, normalized size of antiderivative = 2.49

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx = \text{Too large to display}$$

input

```
integrate((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(5/2),x, algorithm="fricas")
```


output

```
[1/672*(105*sqrt(1/2)*(b^3*f^6 + 8*a*d*e^3*f^2 - 2*(b^2*d*e + 2*a*b*e^2)*f^4)*sqrt(-(b*f^2 - 2*d*e)/e)*log(-b^2*f^4 + 4*(b*d*e - a*e^2)*f^2 - 4*(b*e^2*f^2 - 2*d*e^3)*x + 4*(2*sqrt(1/2)*e^2*f*sqrt(-(b*f^2 - 2*d*e)/e)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) - sqrt(1/2)*(b*e*f^2 + 2*e^3*x)*sqrt(-(b*f^2 - 2*d*e)/e))*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) + d) + 4*(b*e*f^3 - 2*d*e^2*f)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2)) + 2*(105*b^3*f^6 + 192*e^6*x^3 + 48*d^3*e^3 - 56*(5*b^2*d*e + 6*a*b*e^2)*f^4 + 4*(21*b*d^2*e^2 + 232*a*d*e^3)*f^2 + 144*(b*e^4*f^2 + 2*d*e^5)*x^2 + 2*(7*b^2*e^2*f^4 + 156*d^2*e^4 - 4*(3*b*d*e^3 - 32*a*e^4)*f^2)*x - 2*(35*b^2*e*f^5 - 96*e^5*f*x^2 + 12*d^2*e^3*f - 4*(21*b*d*e^2 + 20*a*e^3)*f^3 - 24*(b*e^3*f^3 + 6*d*e^4*f)*x)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) + d))/e^4, -1/336*(105*sqrt(1/2)*(b^3*f^6 + 8*a*d*e^3*f^2 - 2*(b^2*d*e + 2*a*b*e^2)*f^4)*sqrt((b*f^2 - 2*d*e)/e)*arctan(2*sqrt(1/2)*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) + d)*e*sqrt((b*f^2 - 2*d*e)/e)/(b*f^2 - 2*d*e) - (105*b^3*f^6 + 192*e^6*x^3 + 48*d^3*e^3 - 56*(5*b^2*d*e + 6*a*b*e^2)*f^4 + 4*(21*b*d^2*e^2 + 232*a*d*e^3)*f^2 + 144*(b*e^4*f^2 + 2*d*e^5)*x^2 + 2*(7*b^2*e^2*f^4 + 156*d^2*e^4 - 4*(3*b*d*e^3 - 32*a*e^4)*f^2)*x - 2*(35*b^2*e*f^5 - 96*e^5*f*x^2 + 12*d^2*e^3*f - 4*(21*b*d*e^2 + 20*a*e^3)*f^3 - 24*(b*e^3*f^3 + 6*d*e^4*f)*x)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2...
```

Sympy [F]

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx = \int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx$$

input

```
integrate((d+e*x+f*(a+b*x+e**2*x**2/f**2)**(1/2))**(5/2),x)
```

output

```
Integral((d + e*x + f*sqrt(a + b*x + e**2*x**2/f**2))**(5/2), x)
```

Maxima [F]

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx = \int \left(ex + \sqrt{bx + \frac{e^2 x^2}{f^2} + af + d} \right)^{5/2} dx$$

input `integrate((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(5/2),x, algorithm="maxima")`

output `integrate((e*x + sqrt(b*x + e^2*x^2/f^2 + a)*f + d)^(5/2), x)`

Giac [F]

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx = \int \left(ex + \sqrt{bx + \frac{e^2 x^2}{f^2} + af + d} \right)^{5/2} dx$$

input `integrate((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(5/2),x, algorithm="giac")`

output `integrate((e*x + sqrt(b*x + e^2*x^2/f^2 + a)*f + d)^(5/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx = \int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx$$

input `int((d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^(5/2),x)`

output `int((d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^(5/2), x)`

Reduce [F]

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx = \int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx$$

input `int((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(5/2),x)`

output `int((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(5/2),x)`

$$3.14 \quad \int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx$$

Optimal result	139
Mathematica [A] (verified)	140
Rubi [A] (verified)	141
Maple [F]	143
Fricas [A] (verification not implemented)	144
Sympy [F]	145
Maxima [F]	145
Giac [F]	145
Mupad [F(-1)]	146
Reduce [F]	146

Optimal result

Integrand size = 30, antiderivative size = 302

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx = \frac{f^2(4ae^2 - b^2 f^2) \sqrt{d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}}}}{4e^3} + \frac{\left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{5/2}}{5e} - \frac{f^2(2de - bf^2)(4ae^2 - b^2 f^2) \sqrt{d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}}}}{8e^3 \left(bf^2 + 2e \left(ex + f \sqrt{a + \frac{x(bf^2 + e^2 x)}{f^2}} \right) \right)} - \frac{3f^2 \sqrt{2de - bf^2} (4ae^2 - b^2 f^2) \operatorname{arctanh} \left(\frac{\sqrt{2} \sqrt{e} \sqrt{d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}}}}{\sqrt{2de - bf^2}} \right)}{8\sqrt{2}e^{7/2}}$$

output

$$\frac{1}{4}f^2(-b^2f^2+4ae^2)(d+ex+f(a+bx+e^2x^2/f^2)^{1/2})^{1/2}/e^{3+1/5}(d+ex+f(a+bx+e^2x^2/f^2)^{1/2})^{5/2}/e-1/8f^2(-bf^2+2de)(-b^2f^2+4ae^2)(d+ex+f(a+bx+e^2x^2/f^2)^{1/2})^{1/2}/e^3/(bf^2+2e(ex+f(a+x(bf^2+e^2x)/f^2)^{1/2}))-3/16f^2(-bf^2+2de)^{1/2}(-b^2f^2+4ae^2)\operatorname{arctanh}(2^{1/2}e^{1/2}(d+ex+f(a+bx+e^2x^2/f^2)^{1/2}))^{1/2}/(-bf^2+2de)^{1/2})^{1/2})^{1/2}/e^{7/2}$$
Mathematica [A] (verified)

Time = 3.62 (sec) , antiderivative size = 443, normalized size of antiderivative = 1.47

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx = \frac{\sqrt{d + ex + f \sqrt{a + x \left(b + \frac{e^2 x}{f^2} \right)}} \left(-15b^3 f^6 - 2b^2 e f^4 \left(-5d + 6ex + 10f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right) \right)}{2e^{3/2}} - \frac{3af^2 \sqrt{-de + \frac{bf^2}{2}} \operatorname{arctan} \left(\frac{\sqrt{2}\sqrt{e} \sqrt{d+ex+f \sqrt{a+x \left(b + \frac{e^2 x}{f^2} \right)}}}{\sqrt{-2de+bf^2}} \right)}{2e^{3/2}} + \frac{3b^2 f^4 \sqrt{-de + \frac{bf^2}{2}} \operatorname{arctan} \left(\frac{\sqrt{2}\sqrt{e} \sqrt{d+ex+f \sqrt{a+x \left(b + \frac{e^2 x}{f^2} \right)}}}{\sqrt{-2de+bf^2}} \right)}{8e^{7/2}}$$

input

`Integrate[(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^(3/2), x]`

output

```
(Sqrt[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2))]*(-15*b^3*f^6 - 2*b^2*e*f^4*(-5*d + 6*e*x + 10*f*Sqrt[a + x*(b + (e^2*x)/f^2)]) + 4*b*e^2*f^2*(2*d^2 + 17*a*f^2 + 8*e*x*(2*e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]) + 4*d*(3*e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)))) + 8*e^3*(2*(d + 2*e*x)^2*(e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]) + a*f^2*(-d + 16*e*x + 12*f*Sqrt[a + x*(b + (e^2*x)/f^2)])))/(40*e^3*(b*f^2 + 2*e*(e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]))) - (3*a*f^2*Sqrt[-(d*e) + (b*f^2)/2]*ArcTan[(Sqrt[2]*Sqrt[e]*Sqrt[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])]/Sqrt[-2*d*e + b*f^2])/(2*e^(3/2)) + (3*b^2*f^4*Sqrt[-(d*e) + (b*f^2)/2]*ArcTan[(Sqrt[2]*Sqrt[e]*Sqrt[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])]/Sqrt[-2*d*e + b*f^2])/(8*e^(7/2))
```

Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 309, normalized size of antiderivative = 1.02, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2541, 1192, 1580, 2341, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} + d + ex \right)^{3/2} dx$$

↓ 2541

$$2 \int \frac{\left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)^{3/2} \left(ed^2 - bf^2 d + aef^2 + e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)^2 - (2de - bf^2) \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right) \right)}{\left(-bf^2 + 2de - 2e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right) \right)^2} dx$$

↓ 1192

$$4 \int \frac{\left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)^2 \left(ed^2 - bf^2 d + aef^2 + e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)^2 - (2de - bf^2) \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right) \right)}{\left(-bf^2 + 2de - 2e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right) \right)^2} dx$$

↓ 1580

$$4 \left(\frac{f^2(4ae^2 - b^2f^2)(2de - bf^2) \sqrt{f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex}}{32e^3 \left(-2e \left(f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex \right) - bf^2 + 2de \right)} - \int \frac{16 \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a} \right)^3 e^{4-8(2de-bf^2)} \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a} \right)}{\dots} \right)$$

↓ 2341

$$4 \left(\frac{f^2(4ae^2 - b^2f^2)(2de - bf^2) \sqrt{f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex}}{32e^3 \left(-2e \left(f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex \right) - bf^2 + 2de \right)} - \int \left(-8 \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a} \right)^2 e^3 - 2f^2 \right) \dots \right)$$

↓ 2009

$$4 \left(\frac{f^2(4ae^2 - b^2f^2)(2de - bf^2) \sqrt{f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex}}{32e^3 \left(-2e \left(f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex \right) - bf^2 + 2de \right)} - \frac{3\sqrt{e}f^2(4ae^2 - b^2f^2)\sqrt{2de-bf^2}\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{e}\sqrt{f\sqrt{a+bx+\frac{e^2x^2}{f^2}}+d+ex}}{\sqrt{2de-bf^2}}\right)}{\sqrt{2}} \right)$$

input `Int[(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^(3/2),x]`

output `4*((f^2*(2*d*e - b*f^2)*(4*a*e^2 - b^2*f^2)*Sqrt[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]]/(32*e^3*(2*d*e - b*f^2 - 2*e*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]))) - (-2*e*f^2*(4*a*e^2 - b^2*f^2)*Sqrt[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]] - (8*e^3*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^(5/2))/5 + (3*Sqrt[e]*f^2*Sqrt[2*d*e - b*f^2]*(4*a*e^2 - b^2*f^2)*ArcTanh[(Sqrt[2]*Sqrt[e]*Sqrt[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]])/Sqrt[2*d*e - b*f^2])/Sqrt[2])/(32*e^4)`

Definitions of rubi rules used

rule 1192

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))^(n_)*((a_.) + (b_.)*(x_)
+ (c_.)*(x_)^2)^(p_.), x_Symbol] := Simp[2/e^(n + 2*p + 1) Subst[Int[x^(
2*m + 1)*(e*f - d*g + g*x^2)^n*(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 +
c*x^4)^p, x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, g}, x] &&
IGtQ[p, 0] && ILtQ[n, 0] && IntegerQ[m + 1/2]
```

rule 1580

```
Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)
^4)^(p_.), x_Symbol] := Simp[(-d)^(m/2 - 1)*(c*d^2 - b*d*e + a*e^2)^p*x*((d
+ e*x^2)^(q + 1)/(2*e^(2*p + m/2)*(q + 1))), x] + Simp[1/(2*e^(2*p + m/2)*
(q + 1)) Int[(d + e*x^2)^(q + 1)*ExpandToSum[Together[(1/(d + e*x^2))*(2*
e^(2*p + m/2)*(q + 1)*x^m*(a + b*x^2 + c*x^4)^p - (-d)^(m/2 - 1)*(c*d^2 - b
*d*e + a*e^2)^p*(d + e*(2*q + 3)*x^2)], x], x], x] /; FreeQ[{a, b, c, d, e
}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && ILtQ[q, -1] && IGtQ[m/2, 0]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2341

```
Int[(Pq_)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[Pq*
(a + b*x^2)^p, x], x] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && IGtQ[p, -2]
```

rule 2541

```
Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_.) + (b_.)*(x_) + (c
_.)*(x_)^2]))^(n_))^(p_.), x_Symbol] := Simp[2 Subst[Int[(g + h*x^n)^p*((d
^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)/(-2*d*e + b*f^2 + 2*e*x
)^2), x], x, d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e
, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]
```

Maple [F]

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{\frac{3}{2}} dx$$

input

```
int((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(3/2),x)
```


output `int((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(3/2),x)`

Fricas [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 657, normalized size of antiderivative = 2.18

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx = \frac{15 \sqrt{\frac{1}{2}} (b^2 f^4 - 4 a e^2 f^2) \sqrt{-\frac{b f^2 - 2 d e}{e}} \log \left(-b^2 f^4 + 4 (b d e - a e^2) f^2 - 4 (b e^2 f^2 - 2 d e^3) x - 4 (2 \sqrt{\frac{1}{2}} e^2 f \sqrt{-\frac{b f^2 - 2 d e}{e}} \sqrt{\frac{b f^2 x + e^2 x^2 + a f^2}{f^2}} - \sqrt{\frac{1}{2}} (b e f^2 + 2 e^3 x) \sqrt{-\frac{b f^2 - 2 d e}{e}} \sqrt{e x + f \sqrt{\frac{b f^2 x + e^2 x^2 + a f^2}{f^2}} + d} + 4 (b e f^3 - 2 d e^2 f) \sqrt{\frac{b f^2 x + e^2 x^2 + a f^2}{f^2}}) - 2 (15 b^2 f^4 - 16 e^4 x^2 - 8 d^2 e^2 - 2 (5 b d e + 24 a e^2) f^2 + 2 (b e^2 f^2 - 18 d e^3) x - 2 (5 b e f^3 + 8 e^3 f x - 2 d e^2 f) \sqrt{\frac{b f^2 x + e^2 x^2 + a f^2}{f^2}}) \sqrt{e x + f \sqrt{\frac{b f^2 x + e^2 x^2 + a f^2}{f^2}} + d} \right)}{e^3} + \frac{1}{40} (15 \sqrt{\frac{1}{2}} (b^2 f^4 - 4 a e^2 f^2) \sqrt{-\frac{b f^2 - 2 d e}{e}} \arctan(2 \sqrt{\frac{1}{2}} \sqrt{e x + f \sqrt{\frac{b f^2 x + e^2 x^2 + a f^2}{f^2}} + d}) \sqrt{-\frac{b f^2 - 2 d e}{e}} / (b f^2 - 2 d e)) - (15 b^2 f^4 - 16 e^4 x^2 - 8 d^2 e^2 - 2 (5 b d e + 24 a e^2) f^2 + 2 (b e^2 f^2 - 18 d e^3) x - 2 (5 b e f^3 + 8 e^3 f x - 2 d e^2 f) \sqrt{\frac{b f^2 x + e^2 x^2 + a f^2}{f^2}}) \sqrt{e x + f \sqrt{\frac{b f^2 x + e^2 x^2 + a f^2}{f^2}} + d} \right)}{e^3}$$

input `integrate((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(3/2),x, algorithm="fricas")`

output `[1/80*(15*sqrt(1/2)*(b^2*f^4 - 4*a*e^2*f^2)*sqrt(-(b*f^2 - 2*d*e)/e)*log(-b^2*f^4 + 4*(b*d*e - a*e^2)*f^2 - 4*(b*e^2*f^2 - 2*d*e^3)*x - 4*(2*sqrt(1/2)*e^2*f*sqrt(-(b*f^2 - 2*d*e)/e)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) - sqrt(1/2)*(b*e*f^2 + 2*e^3*x)*sqrt(-(b*f^2 - 2*d*e)/e))*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) + d) + 4*(b*e*f^3 - 2*d*e^2*f)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2)) - 2*(15*b^2*f^4 - 16*e^4*x^2 - 8*d^2*e^2 - 2*(5*b*d*e + 24*a*e^2)*f^2 + 2*(b*e^2*f^2 - 18*d*e^3)*x - 2*(5*b*e*f^3 + 8*e^3*f*x - 2*d*e^2*f)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) + d))/e^3, 1/40*(15*sqrt(1/2)*(b^2*f^4 - 4*a*e^2*f^2)*sqrt(-(b*f^2 - 2*d*e)/e)*arctan(2*sqrt(1/2)*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) + d))*sqrt(-(b*f^2 - 2*d*e)/e)/(b*f^2 - 2*d*e)) - (15*b^2*f^4 - 16*e^4*x^2 - 8*d^2*e^2 - 2*(5*b*d*e + 24*a*e^2)*f^2 + 2*(b*e^2*f^2 - 18*d*e^3)*x - 2*(5*b*e*f^3 + 8*e^3*f*x - 2*d*e^2*f)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) + d))/e^3]`

Sympy [F]

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx = \int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{\frac{3}{2}} dx$$

input `integrate((d+e*x+f*(a+b*x+e**2*x**2/f**2)**(1/2))**(3/2),x)`

output `Integral((d + e*x + f*sqrt(a + b*x + e**2*x**2/f**2))**(3/2), x)`

Maxima [F]

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx = \int \left(ex + \sqrt{bx + \frac{e^2 x^2}{f^2} + af + d} \right)^{\frac{3}{2}} dx$$

input `integrate((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(3/2),x, algorithm="maxima")`

output `integrate((e*x + sqrt(b*x + e^2*x^2/f^2 + a)*f + d)^(3/2), x)`

Giac [F]

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx = \int \left(ex + \sqrt{bx + \frac{e^2 x^2}{f^2} + af + d} \right)^{\frac{3}{2}} dx$$

input `integrate((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(3/2),x, algorithm="giac")`

output `integrate((e*x + sqrt(b*x + e^2*x^2/f^2 + a)*f + d)^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx = \int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx$$

input `int((d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^(3/2),x)`output `int((d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^(3/2), x)`**Reduce [F]**

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx = \int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{\frac{3}{2}} dx$$

input `int((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(3/2),x)`output `int((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(3/2),x)`

3.15
$$\int \sqrt{d + ex + f} \sqrt{a + bx + \frac{e^2 x^2}{f^2}} dx$$

Optimal result	147
Mathematica [A] (verified)	148
Rubi [A] (verified)	148
Maple [F]	151
Fricas [A] (verification not implemented)	152
Sympy [F]	153
Maxima [F]	153
Giac [F]	153
Mupad [F(-1)]	154
Reduce [F]	154

Optimal result

Integrand size = 30, antiderivative size = 233

$$\int \sqrt{d + ex + f} \sqrt{a + bx + \frac{e^2 x^2}{f^2}} dx$$

$$= \frac{\left(d + ex + f\sqrt{a + bx + \frac{e^2 x^2}{f^2}}\right)^{3/2}}{3e} - \frac{f^2 \left(4a - \frac{b^2 f^2}{e^2}\right) \sqrt{d + ex + f\sqrt{a + bx + \frac{e^2 x^2}{f^2}}}}{4 \left(bf^2 + 2e \left(ex + f\sqrt{a + \frac{x(bf^2 + e^2 x)}{f^2}}\right)\right)}$$

$$- \frac{f^2(4ae^2 - b^2 f^2) \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{e}\sqrt{d+ex+f\sqrt{a+bx+\frac{e^2 x^2}{f^2}}}}{\sqrt{2de-bf^2}}\right)}{4\sqrt{2}e^{5/2}\sqrt{2de-bf^2}}$$

output

```
1/3*(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(3/2)/e-f^2*(4*a-b^2*f^2/e^2)*(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2)/(4*b*f^2+8*e*(e*x+f*(a+x*(b*f^2+e^2*x)/f^2)^(1/2)))-1/8*f^2*(-b^2*f^2+4*a*e^2)*arctanh(2^(1/2)*e^(1/2)*(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2)/(-b*f^2+2*d*e)^(1/2))*2^(1/2)/e^(5/2)/(-b*f^2+2*d*e)^(1/2)
```

Mathematica [A] (verified)

Time = 1.97 (sec) , antiderivative size = 327, normalized size of antiderivative = 1.40

$$\int \sqrt{d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}}} dx$$

$$= \frac{\sqrt{d + ex + f \sqrt{a + x \left(b + \frac{e^2 x}{f^2} \right)}} \left(3b^2 f^4 + 4bef^2 \left(d + 3ex + f \sqrt{a + x \left(b + \frac{e^2 x}{f^2} \right)} \right) + 4e^2 \left(-af^2 + 2(d + 2ex) \right) \right)}{12e^2 \left(bf^2 + 2e \left(ex + f \sqrt{a + x \left(b + \frac{e^2 x}{f^2} \right)} \right) \right)}$$

$$+ \frac{af^2 \arctan \left(\frac{\sqrt{2}\sqrt{e}\sqrt{d+ex+f\sqrt{a+x\left(b+\frac{e^2x}{f^2}\right)}}}{\sqrt{-2de+bf^2}} \right)}{\sqrt{2}\sqrt{e}\sqrt{-2de+bf^2}} - \frac{b^2 f^4 \arctan \left(\frac{\sqrt{2}\sqrt{e}\sqrt{d+ex+f\sqrt{a+x\left(b+\frac{e^2x}{f^2}\right)}}}{\sqrt{-2de+bf^2}} \right)}{4e^{5/2}\sqrt{-4de+2bf^2}}$$

input

```
Integrate[Sqrt[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]], x]
```

output

```
(Sqrt[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])*(3*b^2*f^4 + 4*b*e*f^2*(d + 3*e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]) + 4*e^2*(-(a*f^2) + 2*(d + 2*e*x)*(e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])))/(12*e^2*(b*f^2 + 2*e*(e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]))) + (a*f^2*ArcTan[(Sqrt[2]*Sqrt[e]*Sqrt[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])]/Sqrt[-2*d*e + b*f^2]]/Sqrt[2]*Sqrt[e]*Sqrt[-2*d*e + b*f^2]) - (b^2*f^4*ArcTan[(Sqrt[2]*Sqrt[e]*Sqrt[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])]/Sqrt[-2*d*e + b*f^2]])/(4*e^(5/2)*Sqrt[-4*d*e + 2*b*f^2])
```

Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.04, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2541, 1192, 1580, 25, 1467, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} + d + ex} dx$$

↓ 2541

$$2 \int \frac{\sqrt{d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a}} \left(ed^2 - bf^2 d + aef^2 + e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)^2 - (2de - bf^2) \left(d + ex \right) \right)}{\left(-bf^2 + 2de - 2e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right) \right)^2}$$

↓ 1192

$$4 \int \frac{\left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right) \left(ed^2 - bf^2 d + aef^2 + e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)^2 - (2de - bf^2) \left(d + ex \right) \right)}{\left(-bf^2 + 2de - 2e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right) \right)^2}$$

↓ 1580

$$4 \left(\frac{\int \frac{8 \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)^2 e^3 - 4(2de - bf^2) \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right) e^2 + f^2 (4ae^2 - b^2 f^2) e}{-bf^2 + 2de - 2e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)} d \sqrt{d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a}}}{16e^3} + \dots \right)$$

↓ 25

$$4 \left(\frac{f^2 \left(4a - \frac{b^2 f^2}{e^2} \right) \sqrt{f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} + d + ex}}{16 \left(-2e \left(f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} + d + ex \right) - bf^2 + 2de \right)} - \frac{\int \frac{8 \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)^2 e^3 - 4(2de - bf^2) \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right) e^2 + f^2 (4ae^2 - b^2 f^2) e}{-bf^2 + 2de - 2e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)}}{16e^3} + \dots \right)$$

↓ 1467

$$4 \left(\frac{f^2 \left(4a - \frac{b^2 f^2}{e^2} \right) \sqrt{f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} + d + ex}}{16 \left(-2e \left(f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} + d + ex \right) - bf^2 + 2de \right)} - \frac{\int \left(\frac{ef^2 (4ae^2 - b^2 f^2)}{-bf^2 + 2de - 2e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)} - 4e^2 \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right) \right)}{16e^3} + \dots \right)$$

↓ 2009

$$4 \left(\frac{f^2 \left(4a - \frac{b^2 f^2}{e^2}\right) \sqrt{f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} + d + ex}}{16 \left(-2e \left(f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} + d + ex\right) - bf^2 + 2de\right)} - \frac{\sqrt{e} f^2 (4ae^2 - b^2 f^2) \operatorname{arctanh} \left(\frac{\sqrt{2} \sqrt{e} \sqrt{f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} + d + ex}}{\sqrt{2de - bf^2}} \right)}{\sqrt{2} \sqrt{2de - bf^2}} \right) - \frac{4}{3}$$

input `Int[Sqrt[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]],x]`

output `4*((f^2*(4*a - (b^2*f^2)/e^2)*Sqrt[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]])/(16*(2*d*e - b*f^2 - 2*e*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]))) - ((-4*e^2*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^(3/2))/3 + (Sqrt[e]*f^2*(4*a*e^2 - b^2*f^2)*ArcTanh[(Sqrt[2]*Sqrt[e]*Sqrt[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]])/Sqrt[2*d*e - b*f^2]])/(Sqrt[2]*Sqrt[2*d*e - b*f^2]))/(16*e^3)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 1192 `Int[((d_) + (e_)*(x_)^(m_))*((f_) + (g_)*(x_)^(n_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] :> Simp[2/e^(n + 2*p + 1) Subst[Int[x^(2*m + 1)*(e*f - d*g + g*x^2)^n*(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 + c*x^4)^p, x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && IGtQ[p, 0] && ILtQ[n, 0] && IntegerQ[m + 1/2]`

rule 1467 `Int[((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] :> Int[ExpandIntegrand[(d + e*x^2)^q*(a + b*x^2 + c*x^4)^p, x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && IGtQ[p, 0] && IGtQ[q, -2]`

rule 1580

```
Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := Simp[(-d)^(m/2 - 1)*(c*d^2 - b*d*e + a*e^2)^p*x*((d + e*x^2)^(q + 1)/(2*e^(2*p + m/2)*(q + 1))), x] + Simp[1/(2*e^(2*p + m/2)*(q + 1)) Int[(d + e*x^2)^(q + 1)*ExpandToSum[Together[(1/(d + e*x^2))*(2*e^(2*p + m/2)*(q + 1)*x^m*(a + b*x^2 + c*x^4)^p - (-d)^(m/2 - 1)*(c*d^2 - b*d*e + a*e^2)^p*(d + e*(2*q + 3)*x^2))], x], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && ILtQ[q, -1] && IGtQ[m/2, 0]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2541

```
Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2])^(n_.))^(p_.), x_Symbol] := Simp[2 Subst[Int[(g + h*x^n)^p*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)/(-2*d*e + b*f^2 + 2*e*x)^2), x], x, d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]
```

Maple [F]

$$\int \sqrt{d + ex + f} \sqrt{a + bx + \frac{e^2 x^2}{f^2}} dx$$

input

```
int((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2),x)
```

output

```
int((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2),x)
```


Fricas [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 692, normalized size of antiderivative = 2.97

$$\int \sqrt{d + ex + f} \sqrt{a + bx + \frac{e^2 x^2}{f^2}} dx$$

$$= \frac{3(b^2 f^4 - 4ae^2 f^2) \sqrt{-2bef^2 + 4de^2} \log\left(-b^2 f^4 + 4(bde - ae^2)f^2 - 4(be^2 f^2 - 2de^3)x - 2\left(2\sqrt{-2}\right)\right)}{\dots}$$

input `integrate((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2),x, algorithm="fricas")`

output `[-1/48*(3*(b^2*f^4 - 4*a*e^2*f^2)*sqrt(-2*b*e*f^2 + 4*d*e^2)*log(-b^2*f^4 + 4*(b*d*e - a*e^2)*f^2 - 4*(b*e^2*f^2 - 2*d*e^3)*x - 2*(2*sqrt(-2)*b*e*f^2 + 4*d*e^2)*e*f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) - sqrt(-2*b*e*f^2 + 4*d*e^2)*(b*f^2 + 2*e^2*x))*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) + d) + 4*(b*e*f^3 - 2*d*e^2*f)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2)) - 4*(3*b^2*e*f^4 - 2*b*d*e^2*f^2 - 8*d^2*e^3 + 10*(b*e^3*f^2 - 2*d*e^4)*x - 2*(b*e^2*f^3 - 2*d*e^3*f)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) + d))/(b*e^3*f^2 - 2*d*e^4), 1/24*(3*(b^2*f^4 - 4*a*e^2*f^2)*sqrt(2*b*e*f^2 - 4*d*e^2)*arctan(1/2*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) + d)*(sqrt(2*b*e*f^2 - 4*d*e^2)*f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) - sqrt(2*b*e*f^2 - 4*d*e^2)*(e*x + d))/(a*e*f^2 - d^2*e + (b*e*f^2 - 2*d*e^2)*x) + 2*(3*b^2*e*f^4 - 2*b*d*e^2*f^2 - 8*d^2*e^3 + 10*(b*e^3*f^2 - 2*d*e^4)*x - 2*(b*e^2*f^3 - 2*d*e^3*f)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) + d))/(b*e^3*f^2 - 2*d*e^4)]`

Sympy [F]

$$\int \sqrt{d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}} dx = \int \sqrt{d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}} dx$$

input `integrate((d+e*x+f*(a+b*x+e**2*x**2/f**2)**(1/2))**(1/2),x)`

output `Integral(sqrt(d + e*x + f*sqrt(a + b*x + e**2*x**2/f**2)), x)`

Maxima [F]

$$\int \sqrt{d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}} dx = \int \sqrt{ex + \sqrt{bx + \frac{e^2x^2}{f^2}} + af + d} dx$$

input `integrate((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(e*x + sqrt(b*x + e^2*x^2/f^2 + a)*f + d), x)`

Giac [F]

$$\int \sqrt{d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}} dx = \int \sqrt{ex + \sqrt{bx + \frac{e^2x^2}{f^2}} + af + d} dx$$

input `integrate((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(e*x + sqrt(b*x + e^2*x^2/f^2 + a)*f + d), x)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}}} dx = \int \sqrt{d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}}} dx$$

input `int((d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^(1/2),x)`

output `int((d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^(1/2), x)`

Reduce [F]

$$\int \sqrt{d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}}} dx = \int \sqrt{\sqrt{b f^2 x + e^2 x^2 + a f^2} + d + ex} dx$$

input `int((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2),x)`

output `int(sqrt(sqrt(a*f**2 + b*f**2*x + e**2*x**2) + d + e*x),x)`

3.16
$$\int \frac{1}{\sqrt{d+ex+f}\sqrt{a+bx+\frac{e^2x^2}{f^2}}} dx$$

Optimal result	155
Mathematica [A] (verified)	156
Rubi [A] (verified)	156
Maple [F]	159
Fricas [A] (verification not implemented)	160
Sympy [F]	161
Maxima [F]	161
Giac [F]	161
Mupad [F(-1)]	162
Reduce [F]	162

Optimal result

Integrand size = 30, antiderivative size = 246

$$\int \frac{1}{\sqrt{d+ex+f}\sqrt{a+bx+\frac{e^2x^2}{f^2}}} dx$$

$$= \frac{\sqrt{d+ex+f}\sqrt{a+bx+\frac{e^2x^2}{f^2}}}{e} - \frac{f^2(4ae^2 - b^2f^2)\sqrt{d+ex+f}\sqrt{a+bx+\frac{e^2x^2}{f^2}}}{2e(2de - bf^2)\left(bf^2 + 2e\left(ex + f\sqrt{a+\frac{x(bf^2+e^2x)}{f^2}}\right)\right)}$$

$$+ \frac{f^2(4ae^2 - b^2f^2)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{e}\sqrt{d+ex+f}\sqrt{a+bx+\frac{e^2x^2}{f^2}}}{\sqrt{2de-bf^2}}\right)}{2\sqrt{2}e^{3/2}(2de - bf^2)^{3/2}}$$

output

```
(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2)/e-1/2*f^2*(-b^2*f^2+4*a*e^2)*(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2)/e/(-b*f^2+2*d*e)/(b*f^2+2*e*(e*x+f*(a+x*(b*f^2+e^2*x)/f^2)^(1/2)))+1/4*f^2*(-b^2*f^2+4*a*e^2)*arctanh(2^(1/2)*e^(1/2)*(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2)/(-b*f^2+2*d*e)^(1/2))*2^(1/2)/e^(3/2)/(-b*f^2+2*d*e)^(3/2)
```

Mathematica [A] (verified)

Time = 2.08 (sec) , antiderivative size = 341, normalized size of antiderivative = 1.39

$$\int \frac{1}{\sqrt{d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}}} dx$$

$$= \frac{2\sqrt{e}\sqrt{d+ex+f\sqrt{a+x\left(b+\frac{e^2x}{f^2}\right)}}\left(-b^2f^4-4bef^2\left(-d+ex+f\sqrt{a+x\left(b+\frac{e^2x}{f^2}\right)}\right)+4e^2\left(-af^2+2dex+2df\sqrt{a+x\left(b+\frac{e^2x}{f^2}\right)}\right)\right)}{(2de-bf^2)\left(bf^2+2e\left(ex+f\sqrt{a+x\left(b+\frac{e^2x}{f^2}\right)}\right)\right)} + \frac{4\sqrt{2}ae^2f^2 \arctan\left(\frac{\sqrt{d+ex+f\sqrt{a+x\left(b+\frac{e^2x}{f^2}\right)}}}{\sqrt{a+x\left(b+\frac{e^2x}{f^2}\right)}}\right)}{4e^{3/2}}$$

input `Integrate[1/Sqrt[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]], x]`

output `((2*Sqrt[e]*Sqrt[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])*(-(b^2*f^4) - 4*b*e*f^2*(-d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]) + 4*e^2*(-a*f^2) + 2*d*e*x + 2*d*f*Sqrt[a + x*(b + (e^2*x)/f^2)])))/((2*d*e - b*f^2)*(b*f^2 + 2*e*(e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]))) + (4*Sqrt[2]*a*e^2*f^2*ArcTan[(Sqrt[2]*Sqrt[e]*Sqrt[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])]/Sqrt[-2*d*e + b*f^2]])/(-2*d*e + b*f^2)^(3/2) - (Sqrt[2]*b^2*f^4*ArcTan[(Sqrt[2]*Sqrt[e]*Sqrt[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])]/Sqrt[-2*d*e + b*f^2]])/(-2*d*e + b*f^2)^(3/2))/(4*e^(3/2))`

Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 276, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2541, 1192, 1471, 27, 299, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex}} dx$$

↓ 2541

$$2 \int \frac{ed^2 - bf^2d + aef^2 + e(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a})^2 - (2de - bf^2)(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a})}{\sqrt{d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}} \left(-bf^2 + 2de - 2e(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a})\right)^2} d(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a})$$

↓ 1192

$$4 \int \frac{ed^2 - bf^2d + aef^2 + e(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a})^2 - (2de - bf^2)(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a})}{\left(-bf^2 + 2de - 2e(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a})\right)^2} d\sqrt{d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}}$$

↓ 1471

$$4 \left(\frac{f^2(4ae - \frac{b^2f^2}{e}) \sqrt{f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex}}{8(2de - bf^2) \left(-2e\left(f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex\right) - bf^2 + 2de\right)} - \frac{\int -\frac{\frac{b^2f^4}{e} - 8bdf^2 + 4aef^2 + 8d^2e - 4(2de - bf^2)(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a})}{4(-bf^2 + 2de - 2e(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}))} d\sqrt{d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}}}{2(2de - bf^2)} \right)$$

↓ 27

$$4 \left(\frac{\int \frac{\frac{b^2f^4}{e} - 8bdf^2 + 4aef^2 + 8d^2e - 4(2de - bf^2)(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a})}{-bf^2 + 2de - 2e(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a})} d\sqrt{d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}}}{8(2de - bf^2)} + \frac{f^2(4ae - \frac{b^2f^2}{e})}{8(2de - bf^2) \left(-2e\left(f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex\right) - bf^2 + 2de\right)} \right)$$

↓ 299

$$4 \left(\frac{f^2(4ae^2 - b^2f^2) \int \frac{1}{-bf^2 + 2de - 2e(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a})} d\sqrt{d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}}}{e} + \frac{2(2de - bf^2) \sqrt{f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex}}{e} + \frac{f^2(4ae - \frac{b^2f^2}{e})}{8(2de - bf^2) \left(-2e\left(f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex\right) - bf^2 + 2de\right)} \right)$$

↓ 221

$$4 \left(\frac{f^2(4ae^2 - b^2f^2) \operatorname{arctanh} \left(\frac{\sqrt{2}\sqrt{e} \sqrt{f \sqrt{a+bx + \frac{e^2x^2}{f^2}} + d + ex}}{\sqrt{2de - bf^2}} \right)}{\sqrt{2}e^{3/2} \sqrt{2de - bf^2}} + \frac{2(2de - bf^2) \sqrt{f \sqrt{a+bx + \frac{e^2x^2}{f^2}} + d + ex}}{e} \right) + \frac{f^2 \left(4ae - \frac{b^2f^2}{e} \right) \sqrt{a+bx + \frac{e^2x^2}{f^2}}}{8(2de - bf^2) \left(-2e \left(f \sqrt{a+bx + \frac{e^2x^2}{f^2}} \right) \right)}$$

input `Int[1/Sqrt[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]],x]`

output `4*((f^2*(4*a*e - (b^2*f^2)/e)*Sqrt[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]])/(8*(2*d*e - b*f^2)*(2*d*e - b*f^2 - 2*e*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]))) + ((2*(2*d*e - b*f^2)*Sqrt[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]])/e + (f^2*(4*a*e^2 - b^2*f^2)*ArcTanh[(Sqrt[2]*Sqrt[e]*Sqrt[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]])/Sqrt[2*d*e - b*f^2]])/(Sqrt[2]*e^(3/2)*Sqrt[2*d*e - b*f^2]))/(8*(2*d*e - b*f^2))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 299 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[d*x*((a + b*x^2)^(p + 1)/(b*(2*p + 3))), x] - Simp[(a*d - b*c*(2*p + 3))/(b*(2*p + 3)) Int[(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && NeQ[2*p + 3, 0]`

rule 1192

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))^(n_)*((a_.) + (b_.)*(x_)
+ (c_.)*(x_)^2)^(p_.), x_Symbol] := Simp[2/e^(n + 2*p + 1) Subst[Int[x^(
2*m + 1)*(e*f - d*g + g*x^2)^n*(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 +
c*x^4)^p, x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, g}, x] &&
IGtQ[p, 0] && ILtQ[n, 0] && IntegerQ[m + 1/2]
```

rule 1471

```
Int[((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.),
x_Symbol] := With[{Qx = PolynomialQuotient[(a + b*x^2 + c*x^4)^p, d + e*x^2
, x], R = Coeff[PolynomialRemainder[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], x
, 0]}, Simp[(-R)*x*((d + e*x^2)^(q + 1)/(2*d*(q + 1))), x] + Simp[1/(2*d*(q
+ 1)) Int[(d + e*x^2)^(q + 1)*ExpandToSum[2*d*(q + 1)*Qx + R*(2*q + 3),
x], x], x]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^
2 - b*d*e + a*e^2, 0] && IGtQ[p, 0] && LtQ[q, -1]
```

rule 2541

```
Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_.) + (b_.)*(x_) + (c
_.)*(x_)^2])^(n_))^(p_.), x_Symbol] := Simp[2 Subst[Int[(g + h*x^n)^p*((d
^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)/(-2*d*e + b*f^2 + 2*e*x
)^2), x], x, d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e
, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]
```

Maple [F]

$$\int \frac{1}{\sqrt{d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}}} dx$$

input

```
int(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2),x)
```

output

```
int(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2),x)
```


Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 716, normalized size of antiderivative = 2.91

$$\int \frac{1}{\sqrt{d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}}} dx$$

$$= \frac{(b^2f^4 - 4ae^2f^2)\sqrt{-2bef^2 + 4de^2} \log\left(-b^2f^4 + 4(bde - ae^2)f^2 - 4(be^2f^2 - 2de^3)x + 2\left(2\sqrt{-2bef^2 + 4de^2}\right)\right)}{\dots}$$

input

```
integrate(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2),x, algorithm="fricas")
```

output

```
[1/8*((b^2*f^4 - 4*a*e^2*f^2)*sqrt(-2*b*e*f^2 + 4*d*e^2)*log(-b^2*f^4 + 4*(b*d*e - a*e^2)*f^2 - 4*(b*e^2*f^2 - 2*d*e^3)*x + 2*(2*sqrt(-2*b*e*f^2 + 4*d*e^2)*e*f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) - sqrt(-2*b*e*f^2 + 4*d*e^2)*(b*f^2 + 2*e^2*x))*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) + d) + 4*(b*e*f^3 - 2*d*e^2*f)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2)) + 4*(b^2*e*f^4 - 6*b*d*e^2*f^2 + 8*d^2*e^3 - 2*(b*e^3*f^2 - 2*d*e^4)*x + 2*(b*e^2*f^3 - 2*d*e^3*f)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) + d))/(b^2*e^2*f^4 - 4*b*d*e^3*f^2 + 4*d^2*e^4), 1/4*((b^2*f^4 - 4*a*e^2*f^2)*sqrt(2*b*e*f^2 - 4*d*e^2)*arctan(1/2*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) + d)*(sqrt(2*b*e*f^2 - 4*d*e^2)*f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) - sqrt(2*b*e*f^2 - 4*d*e^2)*(e*x + d))/(a*e*f^2 - d^2*e + (b*e*f^2 - 2*d*e^2)*x)) + 2*(b^2*e*f^4 - 6*b*d*e^2*f^2 + 8*d^2*e^3 - 2*(b*e^3*f^2 - 2*d*e^4)*x + 2*(b*e^2*f^3 - 2*d*e^3*f)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) + d))/(b^2*e^2*f^4 - 4*b*d*e^3*f^2 + 4*d^2*e^4)]
```

Sympy [F]

$$\int \frac{1}{\sqrt{d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}}} dx = \int \frac{1}{\sqrt{d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}}} dx$$

input `integrate(1/(d+e*x+f*(a+b*x+e**2*x**2/f**2)**(1/2))**(1/2),x)`

output `Integral(1/sqrt(d + e*x + f*sqrt(a + b*x + e**2*x**2/f**2)), x)`

Maxima [F]

$$\int \frac{1}{\sqrt{d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}}} dx = \int \frac{1}{\sqrt{ex+\sqrt{bx+\frac{e^2x^2}{f^2}}+af+d}} dx$$

input `integrate(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(e*x + sqrt(b*x + e^2*x^2/f^2 + a)*f + d), x)`

Giac [F]

$$\int \frac{1}{\sqrt{d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}}} dx = \int \frac{1}{\sqrt{ex+\sqrt{bx+\frac{e^2x^2}{f^2}}+af+d}} dx$$

input `integrate(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2),x, algorithm="giac")`

output `integrate(1/sqrt(e*x + sqrt(b*x + e^2*x^2/f^2 + a)*f + d), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}}} dx = \int \frac{1}{\sqrt{d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}}} dx$$

input `int(1/(d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^(1/2), x)`output `int(1/(d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^(1/2), x)`**Reduce [F]**

$$\int \frac{1}{\sqrt{d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}}} dx = \int \frac{1}{\sqrt{d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}}} dx$$

input `int(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2), x)`output `int(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(1/2), x)`

$$3.17 \quad \int \frac{1}{\left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^{3/2}} dx$$

Optimal result	163
Mathematica [A] (verified)	164
Rubi [A] (verified)	165
Maple [F]	167
Fricas [B] (verification not implemented)	168
Sympy [F]	169
Maxima [F]	169
Giac [F]	169
Mupad [F(-1)]	170
Reduce [F]	170

Optimal result

Integrand size = 30, antiderivative size = 269

$$\int \frac{1}{\left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^{3/2}} dx =$$

$$\frac{4(d^2e - bdf^2 + aef^2)}{(2de - bf^2)^2 \sqrt{d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}}}$$

$$\frac{f^2(4ae^2 - b^2f^2) \sqrt{d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}}}{(2de - bf^2)^2 \left(bf^2 + 2e \left(ex + f\sqrt{a+\frac{x(bf^2+e^2x)}{f^2}}\right)\right)}$$

$$+ \frac{3f^2(4ae^2 - b^2f^2) \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{e}\sqrt{d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}}}{\sqrt{2de-bf^2}}\right)}{\sqrt{2}\sqrt{e}(2de - bf^2)^{5/2}}$$

output

$$\begin{aligned} & \frac{(-4*a*e*f^2+4*b*d*f^2-4*d^2*e)/(-b*f^2+2*d*e)^2/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^{(1/2)})^{(1/2)}-f^2*(-b^2*f^2+4*a*e^2)*(d+e*x+f*(a+b*x+e^2*x^2/f^2)^{(1/2)})^{(1/2)}/(-b*f^2+2*d*e)^2/(b*f^2+2*e*(e*x+f*(a+x*(b*f^2+e^2*x)/f^2)^{(1/2)}))+3/2*f^2*(-b^2*f^2+4*a*e^2)*\operatorname{arctanh}(2^{(1/2)}*e^{(1/2)}*(d+e*x+f*(a+b*x+e^2*x^2/f^2)^{(1/2)})^{(1/2)})/(-b*f^2+2*d*e)^{(1/2)}*2^{(1/2)}/e^{(1/2)}/(-b*f^2+2*d*e)^{(5/2)}}{(-2de + bf^2)^2 \sqrt{d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}}}} \end{aligned}$$

Mathematica [A] (verified)

Time = 3.34 (sec) , antiderivative size = 395, normalized size of antiderivative = 1.47

$$\begin{aligned} \int \frac{1}{\left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}}\right)^{3/2}} dx &= \frac{b^2 f^4 \left(5d + ex + f \sqrt{a + x \left(b + \frac{e^2 x}{f^2}\right)}\right) - 4bef^2 \left(d^2 + af^2 - 2d \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}}\right)\right)}{(-2de + bf^2)^2 \sqrt{d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}}}} \\ & - \frac{6\sqrt{2}ae^{3/2}f^2 \arctan\left(\frac{\sqrt{2}\sqrt{e}\sqrt{d+ex+f\sqrt{a+x\left(b+\frac{e^2x}{f^2}\right)}}}{\sqrt{-2de+bf^2}}\right)}{(-2de + bf^2)^{5/2}} \\ & + \frac{3b^2 f^4 \arctan\left(\frac{\sqrt{2}\sqrt{e}\sqrt{d+ex+f\sqrt{a+x\left(b+\frac{e^2x}{f^2}\right)}}}{\sqrt{-2de+bf^2}}\right)}{\sqrt{2}\sqrt{e}(-2de + bf^2)^{5/2}} \end{aligned}$$

input

Integrate[(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^(-3/2), x]

output

$$\begin{aligned} & \frac{(b^2*f^4*(5*d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]) - 4*b*e*f^2*(d^2 + a*f^2 - 2*d*(e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2]))) - 4*e^2*(2*d^2*(e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]) + a*f^2*(d + 3*e*x + 3*f*Sqrt[a + x*(b + (e^2*x)/f^2)])))/((-2*d*e + b*f^2)^2*Sqrt[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])*(b*f^2 + 2*e*(e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])) - (6*Sqrt[2]*a*e^{(3/2)}*f^2*ArcTan[(Sqrt[2]*Sqrt[e]*Sqrt[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])]/Sqrt[-2*d*e + b*f^2]])/(-2*d*e + b*f^2)^{(5/2)} + (3*b^2*f^4*ArcTan[(Sqrt[2]*Sqrt[e]*Sqrt[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])]/Sqrt[-2*d*e + b*f^2]])/(Sqrt[2]*Sqrt[e]*(-2*d*e + b*f^2)^{(5/2))}}{(-2de + bf^2)^2 \sqrt{d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}}}} \end{aligned}$$

Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.04, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2541, 1192, 1582, 27, 359, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\left(f\sqrt{a+bx+\frac{e^2x^2}{f^2}}+d+ex\right)^{3/2}} dx$$

↓ 2541

$$2 \int \frac{ed^2 - bf^2d + aef^2 + e\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)^2 - (2de - bf^2)\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)}{\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)^{3/2} \left(-bf^2 + 2de - 2e\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)\right)^2} d\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)$$

↓ 1192

$$4 \int \frac{ed^2 - bf^2d + aef^2 + e\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)^2 - (2de - bf^2)\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)}{\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right) \left(-bf^2 + 2de - 2e\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)\right)^2} d\sqrt{d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}}$$

↓ 1582

$$4 \left(\frac{f^2(4ae^2 - b^2f^2) \sqrt{f\sqrt{a+bx+\frac{e^2x^2}{f^2}}+d+ex}}{4(2de - bf^2)^2 \left(-2e\left(f\sqrt{a+bx+\frac{e^2x^2}{f^2}}+d+ex\right) - bf^2 + 2de\right)} - \int \frac{2e^2 \left(4(2de - bf^2)(ed^2 - bf^2d + aef^2) - (3b^2f^4 - 4ae^2f^2 - 8bde f^2 + 8d^2e^2)\right)}{\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right) \left(-bf^2 + 2de - 2e\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)\right)} d\sqrt{d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}} \right)$$

↓ 27

$$4 \left(\frac{\int \frac{4(2de - bf^2)(ed^2 - bf^2d + aef^2) - (3b^2f^4 - 4ae^2f^2 - 8bde f^2 + 8d^2e^2) \left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)}{\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right) \left(-bf^2 + 2de - 2e\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)\right)} d\sqrt{d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}}}{4(2de - bf^2)^2} + \dots \right)$$

↓ 359

$$4 \left(\frac{3f^2(4ae^2 - b^2f^2) \int \frac{1}{-bf^2+2de-2e\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)} d\sqrt{d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}} - \frac{4(aef^2-bdf^2+d^2e)}{\sqrt{f\sqrt{a+bx+\frac{e^2x^2}{f^2}+d+ex}}} \right) +$$

↓ 221

$$4 \left(\frac{3f^2(4ae^2-b^2f^2) \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{e}\sqrt{f\sqrt{a+bx+\frac{e^2x^2}{f^2}+d+ex}}}{\sqrt{2de-bf^2}}\right)}{\sqrt{2}\sqrt{e}\sqrt{2de-bf^2}} - \frac{4(aef^2-bdf^2+d^2e)}{\sqrt{f\sqrt{a+bx+\frac{e^2x^2}{f^2}+d+ex}}} + \frac{f^2(4ae^2-b^2f^2)\sqrt{f\sqrt{a+bx+\frac{e^2x^2}{f^2}+d+ex}}}{4(2de-bf^2)^2(-2e\left(f\sqrt{a+bx+\frac{e^2x^2}{f^2}+d+ex}\right))} \right)$$

input `Int[(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^(-3/2),x]`

output `4*((f^2*(4*a*e^2 - b^2*f^2)*Sqrt[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]])/(4*(2*d*e - b*f^2)^2*(2*d*e - b*f^2 - 2*e*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]))) + ((-4*(d^2*e - b*d*f^2 + a*e*f^2))/Sqrt[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]]) + (3*f^2*(4*a*e^2 - b^2*f^2)*ArcTanh[(Sqrt[2]*Sqrt[e]*Sqrt[d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2]])/Sqrt[2*d*e - b*f^2]])/(Sqrt[2]*Sqrt[e]*Sqrt[2*d*e - b*f^2]))/(4*(2*d*e - b*f^2)^2)`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 359

```
Int[((e._)*(x._))^(m._)*((a._) + (b._)*(x._)^2)^(p._)*((c._) + (d._)*(x._)^2), x
_Symbol] := Simp[c*(e*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*e*(m + 1))), x] +
Simp[(a*d*(m + 1) - b*c*(m + 2*p + 3))/(a*e^2*(m + 1)) Int[(e*x)^(m + 2)*
(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0]
&& LtQ[m, -1] && !ILtQ[p, -1]
```

rule 1192

```
Int[((d._) + (e._)*(x._))^(m._)*((f._) + (g._)*(x._))^(n._)*((a._) + (b._)*(x._)
+ (c._)*(x._)^2)^(p._), x_Symbol] := Simp[2/e^(n + 2*p + 1) Subst[Int[x^(
2*m + 1)*(e*f - d*g + g*x^2)^n*(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 +
c*x^4)^p, x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, g}, x] &&
IGtQ[p, 0] && ILtQ[n, 0] && IntegerQ[m + 1/2]
```

rule 1582

```
Int[(x._)^(m._)*((d._) + (e._)*(x._)^2)^(q._)*((a._) + (b._)*(x._)^2 + (c._)*(x._)^
4)^(p._), x_Symbol] := Simp[(-d)^(m/2 - 1)*(c*d^2 - b*d*e + a*e^2)^p*x*((d
+ e*x^2)^(q + 1)/(2*e^(2*p + m/2)*(q + 1))), x] + Simp[(-d)^(m/2 - 1)/(2*e^
(2*p)*(q + 1)) Int[x^m*(d + e*x^2)^(q + 1)*ExpandToSum[Together[(1/(d + e
*x^2))*(2*(-d)^(-m/2 + 1)*e^(2*p)*(q + 1)*(a + b*x^2 + c*x^4)^p - ((c*d^2 -
b*d*e + a*e^2)^p/(e^(m/2)*x^m))*(d + e*(2*q + 3)*x^2)], x], x], x] /; Fre
eQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && ILtQ[q, -1]
&& ILtQ[m/2, 0]
```

rule 2541

```
Int[((g._) + (h._)*((d._) + (e._)*(x._) + (f._)*Sqrt[(a._) + (b._)*(x._) + (c
._)*(x._)^2]))^(n._))^(p._), x_Symbol] := Simp[2 Subst[Int[(g + h*x^n)^p*((d
^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)/(-2*d*e + b*f^2 + 2*e*x
)^2), x], x, d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e
, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]
```

Maple [F]

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{\frac{3}{2}}} dx$$

input

```
int(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(3/2),x)
```


output `int(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(3/2),x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 696 vs. 2(239) = 478.

Time = 0.62 (sec) , antiderivative size = 1456, normalized size of antiderivative = 5.41

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{3/2}} dx = \text{Too large to display}$$

input `integrate(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(3/2),x, algorithm="fricas")`

output `[-1/4*(3*(a*b^2*f^6 + 4*a*d^2*e^2*f^2 - (b^2*d^2 + 4*a^2*e^2)*f^4 + (b^3*f^6 + 8*a*d*e^3*f^2 - 2*(b^2*d*e + 2*a*b*e^2)*f^4)*x)*sqrt(-2*b*e*f^2 + 4*d*e^2)*log(-b^2*f^4 + 4*(b*d*e - a*e^2)*f^2 - 4*(b*e^2*f^2 - 2*d*e^3)*x + 2*(2*sqrt(-2*b*e*f^2 + 4*d*e^2)*e*f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) - sqrt(-2*b*e*f^2 + 4*d*e^2)*(b*f^2 + 2*e^2*x))*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) + d) + 4*(b*e*f^3 - 2*d*e^2*f)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2)) - 4*(a*b^2*e*f^6 - 8*d^4*e^3 - (5*b^2*d^2*e - 2*a*b*d*e^2)*f^4 + 2*(7*b*d^3*e^2 - 4*a*d^2*e^3)*f^2 + 2*(b^2*e^3*f^4 - 4*b*d*e^4*f^2 + 4*d^2*e^5)*x^2 + (b^3*e*f^6 - 4*d^3*e^4 - 2*(4*b^2*d*e^2 - 3*a*b*e^3)*f^4 + 2*(7*b*d^2*e^3 - 6*a*d*e^4)*f^2)*x + 2*(2*d^3*e^3*f + (2*b^2*d*e - 3*a*b*e^2)*f^5 - (5*b*d^2*e^2 - 6*a*d*e^3)*f^3 - (b^2*e^2*f^5 - 4*b*d*e^3*f^3 + 4*d^2*e^4*f)*x)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) + d))/(a*b^3*e*f^8 + 8*d^5*e^4 - (b^3*d^2*e + 6*a*b^2*d*e^2)*f^6 + 6*(b^2*d^3*e^2 + 2*a*b*d^2*e^3)*f^4 - 4*(3*b*d^4*e^3 + 2*a*d^3*e^4)*f^2 + (b^4*e*f^8 - 8*b^3*d*e^2*f^6 + 24*b^2*d^2*e^3*f^4 - 32*b*d^3*e^4*f^2 + 16*d^4*e^5)*x), -1/2*(3*(a*b^2*f^6 + 4*a*d^2*e^2*f^2 - (b^2*d^2 + 4*a^2*e^2)*f^4 + (b^3*f^6 + 8*a*d*e^3*f^2 - 2*(b^2*d*e + 2*a*b*e^2)*f^4)*x)*sqrt(2*b*e*f^2 - 4*d*e^2)*arctan(1/2*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) + d)*(sqrt(2*b*e*f^2 - 4*d*e^2)*f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) - sqrt(2*b*e*f^2 - 4*d*e^2)*(e*x + d...`

Sympy [F]

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{3/2}} dx = \int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{\frac{3}{2}}} dx$$

input `integrate(1/(d+e*x+f*(a+b*x+e**2*x**2/f**2)**(1/2))**(3/2),x)`

output `Integral((d + e*x + f*sqrt(a + b*x + e**2*x**2/f**2))**(-3/2), x)`

Maxima [F]

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{3/2}} dx = \int \frac{1}{\left(ex + \sqrt{bx + \frac{e^2x^2}{f^2}} + af + d\right)^{\frac{3}{2}}} dx$$

input `integrate(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(3/2),x, algorithm="maxima")`

output `integrate((e*x + sqrt(b*x + e^2*x^2/f^2 + a)*f + d)^(-3/2), x)`

Giac [F]

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{3/2}} dx = \int \frac{1}{\left(ex + \sqrt{bx + \frac{e^2x^2}{f^2}} + af + d\right)^{\frac{3}{2}}} dx$$

input `integrate(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(3/2),x, algorithm="giac")`

output `integrate((e*x + sqrt(b*x + e^2*x^2/f^2 + a)*f + d)^(-3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{3/2}} dx = \int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{3/2}} dx$$

input `int(1/(d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^(3/2), x)`output `int(1/(d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^(3/2), x)`**Reduce [F]**

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{3/2}} dx = \int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{\frac{3}{2}}} dx$$

input `int(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(3/2), x)`output `int(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(3/2), x)`

3.18
$$\int \frac{1}{\left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^{5/2}} dx$$

Optimal result	171
Mathematica [A] (verified)	172
Rubi [A] (verified)	173
Maple [F]	176
Fricas [B] (verification not implemented)	176
Sympy [F]	177
Maxima [F]	178
Giac [F]	178
Mupad [F(-1)]	178
Reduce [F]	179

Optimal result

Integrand size = 30, antiderivative size = 335

$$\int \frac{1}{\left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^{5/2}} dx =$$

$$\frac{4(d^2e - bdf^2 + aef^2)}{3(2de - bf^2)^2 \left(d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}\right)^{3/2}}$$

$$\frac{4f^2(4ae^2 - b^2f^2)}{(2de - bf^2)^3 \sqrt{d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}}}$$

$$\frac{2ef^2(4ae^2 - b^2f^2) \sqrt{d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}}}{(2de - bf^2)^3 \left(bf^2 + 2e\left(ex + f\sqrt{a+\frac{x(bf^2+e^2x)}{f^2}}\right)\right)}$$

$$+ \frac{5\sqrt{2}\sqrt{e}f^2(4ae^2 - b^2f^2) \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{e}\sqrt{d+ex+f\sqrt{a+bx+\frac{e^2x^2}{f^2}}}}{\sqrt{2de-bf^2}}\right)}{(2de - bf^2)^{7/2}}$$

output

$$\frac{1}{3} \frac{(-4ae^2f^2 + 4bd^2 - 4d^2e)}{(-bf^2 + 2de)^2} \frac{1}{(d+ex+f(a+bx+e^2x^2/f^2))^{3/2}} - 4f^2 \frac{(-b^2f^2 + 4ae^2)}{(-bf^2 + 2de)^3} \frac{1}{(d+ex+f(a+bx+e^2x^2/f^2))^{1/2}} - 2e^2 \frac{(-b^2f^2 + 4ae^2)}{(-bf^2 + 2de)^3} \frac{1}{(d+ex+f(a+bx+e^2x^2/f^2))^{1/2}} + 5e^{1/2} \frac{1}{(-bf^2 + 2de)^3} \frac{1}{(d+ex+f(a+bx+e^2x^2/f^2))^{1/2}} + 5e^{1/2} \frac{1}{(-bf^2 + 2de)^3} \frac{1}{(d+ex+f(a+bx+e^2x^2/f^2))^{1/2}} \operatorname{arctanh}\left(\frac{2}{(d+ex+f(a+bx+e^2x^2/f^2))^{1/2}}\right) - \frac{1}{(-bf^2 + 2de)^{7/2}}$$

Mathematica [A] (verified)

Time = 4.78 (sec) , antiderivative size = 557, normalized size of antiderivative = 1.66

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{5/2}} dx = \frac{2b^3f^6 \left(4d + 21ex + 6f\sqrt{a + x\left(b + \frac{e^2x}{f^2}\right)}\right) + 2b^2ef^4 \left(9d^2 + 17a\right)}{(-2de + bf^2)^{7/2}} + \frac{20\sqrt{2}ae^{5/2}f^2 \arctan\left(\frac{\sqrt{2}\sqrt{e}\sqrt{d+ex+f\sqrt{a+x\left(b+\frac{e^2x}{f^2}\right)}}}{\sqrt{-2de+bf^2}}\right)}{(-2de + bf^2)^{7/2}} - \frac{5\sqrt{2}b^2\sqrt{e}f^4 \arctan\left(\frac{\sqrt{2}\sqrt{e}\sqrt{d+ex+f\sqrt{a+x\left(b+\frac{e^2x}{f^2}\right)}}}{\sqrt{-2de+bf^2}}\right)}{(-2de + bf^2)^{7/2}}$$

input

Integrate[(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^(-5/2), x]

output

```
(2*b^3*f^6*(4*d + 21*e*x + 6*f*Sqrt[a + x*(b + (e^2*x)/f^2)]) + 2*b^2*e*f^4*(9*d^2 + 17*a*f^2 + 14*d*(e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]) + 30*e*x*(e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])) - 8*b*e^2*f^2*(d^3 + 7*a*d*f^2 - 3*d^2*(e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]) + 5*a*f^2*(4*e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])) - 8*e^3*(15*a^2*f^4 + 2*d^3*(e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]) + a*f^2*(3*d^2 + 20*d*(e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])) + 30*e*x*(e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])))/(3*(2*d*e - b*f^2)^3*(d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])^(3/2)*(b*f^2 + 2*e*(e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]))) + (20*Sqrt[2]*a*e^(5/2)*f^2*ArcTan[Sqrt[2]*Sqrt[e]*Sqrt[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]]]/Sqrt[-2*d*e + b*f^2])/(-2*d*e + b*f^2)^(7/2) - (5*Sqrt[2]*b^2*Sqrt[e]*f^4*ArcTan[Sqrt[2]*Sqrt[e]*Sqrt[d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)]]]/Sqrt[-2*d*e + b*f^2])/(-2*d*e + b*f^2)^(7/2)
```

Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 354, normalized size of antiderivative = 1.06, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2541, 1192, 1582, 27, 1584, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\left(f\sqrt{a+bx+\frac{e^2x^2}{f^2}}+d+ex\right)^{5/2}} dx$$

↓ 2541

$$2 \int \frac{ed^2 - bf^2d + aef^2 + e\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)^2 - (2de - bf^2)\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)}{\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)^{5/2}\left(-bf^2+2de-2e\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)\right)^2} d\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)$$

↓ 1192

$$4 \int \frac{ed^2 - bf^2d + aef^2 + e\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)^2 - (2de - bf^2)\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)}{\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)^2\left(-bf^2+2de-2e\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}\right)\right)^2} d\sqrt{d+ex+f\sqrt{\frac{e^2x^2}{f^2}+bx+a}}$$

↓ 1582

$$4 \left(\frac{\int \frac{f^2(4ae^2 - b^2f^2) \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)^2 e^3 + 2(2de - bf^2)^2 (ed^2 - bf^2d + aef^2)e^2 - 2(2de - bf^2) (b^2f^4 - 2ae^2f^2 - 2bdef^2 + 2d^2e^2) \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)}{\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)^2 \left(-bf^2 + 2de - 2e\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)\right)} dx}{8e^2(2de - bf^2)^3}$$

↓ 27

$$4 \left(\frac{\int \frac{f^2(4ae^2 - b^2f^2) \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)^2 e^3 + 2(2de - bf^2)^2 (ed^2 - bf^2d + aef^2)e^2 - 2(2de - bf^2) (b^2f^4 - 2ae^2f^2 - 2bdef^2 + 2d^2e^2) \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)}{\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)^2 \left(-bf^2 + 2de - 2e\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)\right)} dx}{2e^2(2de - bf^2)^3}$$

↓ 1584

$$4 \left(\frac{\int \left(\frac{2(2de - bf^2)(ed^2 - bf^2d + aef^2)e^2}{\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)^2} + \frac{2(4ae^4f^2 - b^2e^2f^4)}{d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}} + \frac{5(4ae^5f^2 - b^2e^3f^4)}{-bf^2 + 2de - 2e\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)} \right) d\sqrt{d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}}}{2e^2(2de - bf^2)^3}$$

↓ 2009

$$4 \left(\frac{5e^{5/2}f^2(4ae^2 - b^2f^2) \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{e}\sqrt{f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex}}{\sqrt{2de - bf^2}}\right)}{\sqrt{2}\sqrt{2de - bf^2}} - \frac{2e^2f^2(4ae^2 - b^2f^2)}{\sqrt{f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex}} - \frac{2e^2(2de - bf^2)(aef^2 - bdf^2 + d^2e)}{3\left(f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex\right)^{3/2}}}{2e^2(2de - bf^2)^3} + \frac{1}{2}$$

input

```
Int[(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^(-5/2), x]
```

output

$$4*((e*f^2*(4*a*e^2 - b^2*f^2)*\text{Sqrt}[d + e*x + f*\text{Sqrt}[a + b*x + (e^2*x^2)/f^2]])/(2*(2*d*e - b*f^2)^3*(2*d*e - b*f^2 - 2*e*(d + e*x + f*\text{Sqrt}[a + b*x + (e^2*x^2)/f^2]))) + ((-2*e^2*(2*d*e - b*f^2)*(d^2*e - b*d*f^2 + a*e*f^2))/(3*(d + e*x + f*\text{Sqrt}[a + b*x + (e^2*x^2)/f^2])^(3/2)) - (2*e^2*f^2*(4*a*e^2 - b^2*f^2))/\text{Sqrt}[d + e*x + f*\text{Sqrt}[a + b*x + (e^2*x^2)/f^2]]) + (5*e^(5/2)*f^2*(4*a*e^2 - b^2*f^2)*\text{ArcTanh}[(\text{Sqrt}[2]*\text{Sqrt}[e]*\text{Sqrt}[d + e*x + f*\text{Sqrt}[a + b*x + (e^2*x^2)/f^2]])/\text{Sqrt}[2*d*e - b*f^2]])/(\text{Sqrt}[2]*\text{Sqrt}[2*d*e - b*f^2]))/(2*e^2*(2*d*e - b*f^2)^3))$$

Defintions of rubi rules used

rule 27

$$\text{Int}[(a_*)(F_x), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \&\& \text{!MatchQ}[F_x, (b_*)(G_x) /; \text{FreeQ}[b, x]]$$

rule 1192

$$\text{Int}(((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))^(n_)*((a_.) + (b_.)*(x_)^2 + (c_.)*(x_)^2)^(p_.), x_Symbol] \rightarrow \text{Simp}[2/e^(n + 2*p + 1) \quad \text{Subst}[\text{Int}[x^(2*m + 1)*(e*f - d*g + g*x^2)^n*(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 + c*x^4)^p, x], x, \text{Sqrt}[d + e*x]], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g\}, x] \&\& \text{IGtQ}[p, 0] \&\& \text{ILtQ}[n, 0] \&\& \text{IntegerQ}[m + 1/2]$$

rule 1582

$$\text{Int}[(x_)^{(m_)*((d_.) + (e_.)*(x_)^2)^(q_)*((a_.) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] \rightarrow \text{Simp}[(-d)^(m/2 - 1)*(c*d^2 - b*d*e + a*e^2)^p*x*((d + e*x^2)^(q + 1)/(2*e^(2*p + m/2)*(q + 1))), x] + \text{Simp}[(-d)^(m/2 - 1)/(2*e^(2*p)*(q + 1)) \quad \text{Int}[x^m*(d + e*x^2)^(q + 1)*\text{ExpandToSum}[\text{Together}[(1/(d + e*x^2))*(2*(-d)^(-m/2 + 1)*e^(2*p)*(q + 1)*(a + b*x^2 + c*x^4)^p - ((c*d^2 - b*d*e + a*e^2)^p/(e^(m/2)*x^m))*(d + e*(2*q + 3)*x^2)], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{IGtQ}[p, 0] \&\& \text{ILtQ}[q, -1] \&\& \text{ILtQ}[m/2, 0]$$

rule 1584

$$\text{Int}(((f_.)*(x_))^(m_)*((d_.) + (e_.)*(x_)^2)^(q_)*((a_.) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(f*x)^m*(d + e*x^2)^q*(a + b*x^2 + c*x^4)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m, q\}, x] \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{IGtQ}[p, 0] \&\& \text{IGtQ}[q, -2]$$

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2541 `Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2])^(n_))^(p_.), x_Symbol] := Simp[2 Subst[Int[(g + h*x^n)^p*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)/(-2*d*e + b*f^2 + 2*e*x)^2), x], x, d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]`

Maple [F]

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{\frac{5}{2}}} dx$$

input `int(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(5/2),x)`

output `int(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(5/2),x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1105 vs. $2(298) = 596$.

Time = 1.71 (sec) , antiderivative size = 2396, normalized size of antiderivative = 7.15

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{5/2}} dx = \text{Too large to display}$$

input `integrate(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(5/2),x, algorithm="fricas")`

output

```

[-1/6*(15*sqrt(2)*(a^2*b^2*f^8 - 4*a*d^4*e^2*f^2 - 2*(a*b^2*d^2 + 2*a^3*e^
2)*f^6 + (b^2*d^4 + 8*a^2*d^2*e^2)*f^4 + (b^4*f^8 - 16*a*d^2*e^4*f^2 - 4*(
b^3*d*e + a*b^2*e^2)*f^6 + 4*(b^2*d^2*e^2 + 4*a*b*d*e^3)*f^4)*x^2 + 2*(a*b
^3*f^8 - 8*a*d^3*e^3*f^2 - (b^3*d^2 + 2*a*b^2*d*e + 4*a^2*b*e^2)*f^6 + 2*(
b^2*d^3*e + 2*a*b*d^2*e^2 + 4*a^2*d*e^3)*f^4)*x)*sqrt(-e/(b*f^2 - 2*d*e))*
log(-b^2*f^4 + 4*(b*d*e - a*e^2)*f^2 - 4*(b*e^2*f^2 - 2*d*e^3)*x - 2*(2*sq
rt(2)*(b*e*f^3 - 2*d*e^2*f)*sqrt(-e/(b*f^2 - 2*d*e))*sqrt((b*f^2*x + e^2*x
^2 + a*f^2)/f^2) - sqrt(2)*(b^2*f^4 - 2*b*d*e*f^2 + 2*(b*e^2*f^2 - 2*d*e^3
)*x)*sqrt(-e/(b*f^2 - 2*d*e)))*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f
^2)/f^2) + d) + 4*(b*e*f^3 - 2*d*e^2*f)*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f
^2)) - 4*(4*d^5*e^2 + (8*a*b^2*d - 5*a^2*b*e)*f^6 - 2*(2*b^2*d^3 + a*b*d^2*
e + 10*a^2*d*e^2)*f^4 - 6*(b^2*e^3*f^4 - 4*b*d*e^4*f^2 + 4*d^2*e^5)*x^3 -
(9*b*d^4*e - 32*a*d^3*e^2)*f^2 + (3*b^3*e*f^6 - 16*d^3*e^4 + 4*(b^2*d*e^2
- 10*a*b*e^3)*f^4 - 4*(3*b*d^2*e^3 - 20*a*d*e^4)*f^2)*x^2 + 2*(d^4*e^3 + (
4*b^3*d - a*b^2*e)*f^6 - (7*b^2*d^2*e + 6*a*b*d*e^2 + 15*a^2*e^3)*f^4 - 2*
(5*b*d^3*e^2 - 23*a*d^2*e^3)*f^2)*x - 2*(3*a*b^2*f^7 + d^4*e^2*f - (b^2*d
^2 + 2*a*b*d*e + 15*a^2*e^2)*f^5 - 2*(3*b*d^3*e - 11*a*d^2*e^2)*f^3 - 3*(b
^2*e^2*f^5 - 4*b*d*e^3*f^3 + 4*d^2*e^4*f)*x^2 + (3*b^3*f^7 + 40*a*d*e^3*f^3
- 8*d^3*e^3*f - 4*(b^2*d*e + 5*a*b*e^2)*f^5)*x)*sqrt((b*f^2*x + e^2*x^2 +
a*f^2)/f^2))*sqrt(e*x + f*sqrt((b*f^2*x + e^2*x^2 + a*f^2)/f^2) + d))/...

```

Sympy [F]

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{5/2}} dx = \int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{5/2}} dx$$

input

```
integrate(1/(d+e*x+f*(a+b*x+e**2*x**2/f**2)**(1/2))**(5/2), x)
```

output

```
Integral((d + e*x + f*sqrt(a + b*x + e**2*x**2/f**2))**(-5/2), x)
```

Maxima [F]

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{5/2}} dx = \int \frac{1}{\left(ex + \sqrt{bx + \frac{e^2x^2}{f^2}} + af + d\right)^{5/2}} dx$$

input `integrate(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(5/2),x, algorithm="maxima")`

output `integrate((e*x + sqrt(b*x + e^2*x^2/f^2 + a)*f + d)^(-5/2), x)`

Giac [F]

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{5/2}} dx = \int \frac{1}{\left(ex + \sqrt{bx + \frac{e^2x^2}{f^2}} + af + d\right)^{5/2}} dx$$

input `integrate(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(5/2),x, algorithm="giac")`

output `integrate((e*x + sqrt(b*x + e^2*x^2/f^2 + a)*f + d)^(-5/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{5/2}} dx = \int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{5/2}} dx$$

input `int(1/(d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^(5/2),x)`

output `int(1/(d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^(5/2), x)`

Reduce [F]

$$\int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{5/2}} dx = \int \frac{1}{\left(d + ex + f\sqrt{a + bx + \frac{e^2x^2}{f^2}}\right)^{\frac{5}{2}}} dx$$

input `int(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(5/2),x)`

output `int(1/(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(5/2),x)`

$$3.19 \quad \int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^p dx$$

Optimal result	180
Mathematica [A] (verified)	181
Rubi [A] (verified)	181
Maple [F]	183
Fricas [F(-1)]	183
Sympy [F]	184
Maxima [F]	184
Giac [F]	184
Mupad [F(-1)]	185
Reduce [F]	185

Optimal result

Integrand size = 28, antiderivative size = 166

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^p dx = \frac{\left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{1+p}}{2e(1+p)} + \frac{f^2(4ae^2 - b^2 f^2) \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^{1+p} \operatorname{Hypergeometric2F1} \left(2, 1+p, 2+p, \frac{2e \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)}{2de - bf^2} \right)}{2e(2de - bf^2)^2(1+p)}$$

output

```
1/2*(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(p+1)/e/(p+1)+1/2*f^2*(-b^2*f^2+4*
a*e^2)*(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^(p+1)*hypergeom([2, p+1], [2+p],
2*e*(d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))/(-b*f^2+2*d*e))/e/(-b*f^2+2*d*e)^2
/(p+1)
```

Mathematica [A] (verified)

Time = 1.47 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.81

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^p dx$$

$$= \frac{\left(d + ex + f \sqrt{a + x \left(b + \frac{e^2 x}{f^2} \right)} \right)^{1+p} \left((-2de + bf^2)^2 + (4ae^2 f^2 - b^2 f^4) \operatorname{Hypergeometric2F1} \left(2, 1 + p, \right. \right.}{2e(-2de + bf^2)^2(1+p)}$$

input `Integrate[(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^p,x]`

output `((d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2)])^(1 + p)*((-2*d*e + b*f^2)^2 + (4*a*e^2*f^2 - b^2*f^4)*Hypergeometric2F1[2, 1 + p, 2 + p, (2*e*(d + e*x + f*Sqrt[a + x*(b + (e^2*x)/f^2]))/(2*d*e - b*f^2)]))/(2*e*(-2*d*e + b*f^2)^2*(1 + p))`

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.01, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$, Rules used = {2541, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} + d + ex \right)^p dx$$

↓ 2541

$$2 \int \frac{\left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)^p \left(ed^2 - bf^2 d + aef^2 + e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right)^2 - (2de - bf^2) \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right) \right)}{\left(-bf^2 + 2de - 2e \left(d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + bx + a} \right) \right)^2} dx$$

↓ 1195

$$2 \int \left(\frac{\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)^p}{4e} + \frac{(4ae^2f^2 - b^2f^4) \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)^p}{4e \left(-bf^2 + 2de - 2e \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)\right)^2} \right) d \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)$$

↓ 2009

$$2 \frac{f^2(4ae^2 - b^2f^2) \left(f\sqrt{a + bx + \frac{e^2x^2}{f^2}} + d + ex\right)^{p+1} \text{Hypergeometric2F1} \left(2, p + 1, p + 2, \frac{2e \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + bx + a}\right)}{2de - bf^2}\right)}{4e(p + 1) (2de - bf^2)^2}$$

input `Int[(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^p,x]`

output `2*((d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^(1 + p)/(4*e*(1 + p)) + (f^2*(4*a*e^2 - b^2*f^2)*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])^(1 + p)*Hypergeometric2F1[2, 1 + p, 2 + p, (2*e*(d + e*x + f*Sqrt[a + b*x + (e^2*x^2)/f^2])/(2*d*e - b*f^2)])/(2*d*e - b*f^2))/(4*e*(2*d*e - b*f^2)^2*(1 + p))`

Defintions of rubi rules used

rule 1195 `Int[((d_.) + (e_.)*(x_))^(m_.)*((f_.) + (g_.)*(x_))^(n_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2541

```
Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2])^(n_.))^(p_.), x_Symbol] :> Simp[2 Subst[Int[(g + h*x^n)^p*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)/(-2*d*e + b*f^2 + 2*e*x)^2], x], x, d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]
```

Maple [F]

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^p dx$$

input

```
int((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^p,x)
```

output

```
int((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^p,x)
```

Fricas [F(-1)]

Timed out.

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^p dx = \text{Timed out}$$

input

```
integrate((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^p,x, algorithm="fricas")
```

output

```
Timed out
```


Sympy [F]

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^p dx = \int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^p dx$$

input `integrate((d+e*x+f*(a+b*x+e**2*x**2/f**2)**(1/2))**p,x)`

output `Integral((d + e*x + f*sqrt(a + b*x + e**2*x**2/f**2))**p, x)`

Maxima [F]

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^p dx = \int \left(ex + \sqrt{bx + \frac{e^2 x^2}{f^2} + af + d} \right)^p dx$$

input `integrate((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^p,x, algorithm="maxima")`

output `integrate((e*x + sqrt(b*x + e^2*x^2/f^2 + a)*f + d)^p, x)`

Giac [F]

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^p dx = \int \left(ex + \sqrt{bx + \frac{e^2 x^2}{f^2} + af + d} \right)^p dx$$

input `integrate((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^p,x, algorithm="giac")`

output `integrate((e*x + sqrt(b*x + e^2*x^2/f^2 + a)*f + d)^p, x)`

Mupad [F(-1)]

Timed out.

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^p dx = \int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^p dx$$

input `int((d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^p,x)`output `int((d + e*x + f*(a + b*x + (e^2*x^2)/f^2)^(1/2))^p, x)`**Reduce [F]**

$$\int \left(d + ex + f \sqrt{a + bx + \frac{e^2 x^2}{f^2}} \right)^p dx = \int \left(\sqrt{b f^2 x + e^2 x^2 + a f^2} + d + ex \right)^p dx$$

input `int((d+e*x+f*(a+b*x+e^2*x^2/f^2)^(1/2))^p,x)`output `int((sqrt(a*f**2 + b*f**2*x + e**2*x**2) + d + e*x)**p,x)`

3.20 $\int (x + \sqrt{3 - 2x - x^2})^2 dx$

Optimal result	186
Mathematica [A] (verified)	186
Rubi [A] (verified)	187
Maple [A] (verified)	188
Fricas [A] (verification not implemented)	188
Sympy [A] (verification not implemented)	189
Maxima [A] (verification not implemented)	189
Giac [A] (verification not implemented)	190
Mupad [B] (verification not implemented)	190
Reduce [B] (verification not implemented)	191

Optimal result

Integrand size = 18, antiderivative size = 58

$$\int (x + \sqrt{3 - 2x - x^2})^2 dx = 3x - x^2 - (1 + x)\sqrt{3 - 2x - x^2} - \frac{2}{3}(3 - 2x - x^2)^{3/2} + 4 \arcsin\left(\frac{1}{2}(-1 - x)\right)$$

output

```
3*x-x^2-(1+x)*(-x^2-2*x+3)^(1/2)-2/3*(-x^2-2*x+3)^(3/2)-4*arcsin(1/2+1/2*x)
```

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.97

$$\int (x + \sqrt{3 - 2x - x^2})^2 dx = -((-3 + x)x) + \frac{1}{3}\sqrt{3 - 2x - x^2}(-9 + x + 2x^2) + 8 \arctan\left(\frac{\sqrt{3 - 2x - x^2}}{3 + x}\right)$$

input

```
Integrate[(x + Sqrt[3 - 2*x - x^2])^2,x]
```

output $-\left((-3 + x)*x\right) + \left(\text{Sqrt}[3 - 2*x - x^2]*(-9 + x + 2*x^2)\right)/3 + 8*\text{ArcTan}[\text{Sqrt}[3 - 2*x - x^2]/(3 + x)]$

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\sqrt{-x^2 - 2x + 3} + x \right)^2 dx$$

↓ 7293

$$\int \left(2\sqrt{-x^2 - 2x + 3} + x - 2x + 3 \right) dx$$

↓ 2009

$$4 \arcsin\left(\frac{1}{2}(-x - 1)\right) - x^2 - \frac{2}{3}(-x^2 - 2x + 3)^{3/2} - (x + 1)\sqrt{-x^2 - 2x + 3} + 3x$$

input $\text{Int}[(x + \text{Sqrt}[3 - 2*x - x^2])^2, x]$

output $3*x - x^2 - (1 + x)*\text{Sqrt}[3 - 2*x - x^2] - (2*(3 - 2*x - x^2)^{(3/2)})/3 + 4*\text{ArcSin}[(-1 - x)/2]$

Defintions of rubi rules used

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ /; SumQ}[u]$

rule 7293 $\text{Int}[u_, x_Symbol] \rightarrow \text{With}[\{v = \text{ExpandIntegrand}[u, x]\}, \text{Int}[v, x] \text{ /; SumQ}[v]]$

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.88

method	result
default	$-x^2 + 3x - \frac{2(-x^2-2x+3)^{\frac{3}{2}}}{3} + \frac{(-2x-2)\sqrt{-x^2-2x+3}}{2} - 4 \arcsin\left(\frac{1}{2} + \frac{x}{2}\right)$
trager	$-x(x-3) + \left(\frac{2}{3}x^2 + \frac{1}{3}x - 3\right)\sqrt{-x^2-2x+3} + 4 \operatorname{RootOf}(_Z^2 + 1) \ln(\operatorname{RootOf}(_Z^2 + 1) x +$

input `int((x+(-x^2-2*x+3)^(1/2))^2,x,method=_RETURNVERBOSE)`output `-x^2+3*x-2/3*(-x^2-2*x+3)^(3/2)+1/2*(-2*x-2)*(-x^2-2*x+3)^(1/2)-4*arcsin(1/2+1/2*x)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.03

$$\int \left(x + \sqrt{3 - 2x - x^2}\right)^2 dx = -x^2 + \frac{1}{3}(2x^2 + x - 9)\sqrt{-x^2 - 2x + 3} + 3x + 4 \arctan\left(\frac{\sqrt{-x^2 - 2x + 3}(x + 1)}{x^2 + 2x - 3}\right)$$

input `integrate((x+(-x^2-2*x+3)^(1/2))^2,x, algorithm="fricas")`output `-x^2 + 1/3*(2*x^2 + x - 9)*sqrt(-x^2 - 2*x + 3) + 3*x + 4*arctan(sqrt(-x^2 - 2*x + 3)*(x + 1)/(x^2 + 2*x - 3))`

Sympy [A] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.71

$$\int \left(x + \sqrt{3 - 2x - x^2} \right)^2 dx = -x^2 + 3x + 2\sqrt{-x^2 - 2x + 3} \left(\frac{x^2}{3} + \frac{x}{6} - \frac{3}{2} \right) - 4 \operatorname{asin} \left(\frac{x}{2} + \frac{1}{2} \right)$$

input `integrate((x+(-x**2-2*x+3)**(1/2))**2,x)`

output `-x**2 + 3*x + 2*sqrt(-x**2 - 2*x + 3)*(x**2/3 + x/6 - 3/2) - 4*asin(x/2 + 1/2)`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.03

$$\int \left(x + \sqrt{3 - 2x - x^2} \right)^2 dx = -x^2 - \frac{2}{3} (-x^2 - 2x + 3)^{\frac{3}{2}} - \sqrt{-x^2 - 2x + 3}x + 3x - \sqrt{-x^2 - 2x + 3} + 4 \operatorname{arcsin} \left(-\frac{1}{2}x - \frac{1}{2} \right)$$

input `integrate((x+(-x^2-2*x+3)^(1/2))^2,x, algorithm="maxima")`

output `-x^2 - 2/3*(-x^2 - 2*x + 3)^(3/2) - sqrt(-x^2 - 2*x + 3)*x + 3*x - sqrt(-x^2 - 2*x + 3) + 4*arcsin(-1/2*x - 1/2)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.69

$$\int \left(x + \sqrt{3 - 2x - x^2} \right)^2 dx = -x^2 + \frac{1}{3} ((2x + 1)x - 9) \sqrt{-x^2 - 2x + 3} + 3x - 4 \arcsin \left(\frac{1}{2}x + \frac{1}{2} \right)$$

input `integrate((x+(-x^2-2*x+3)^(1/2))^2,x, algorithm="giac")`

output `-x^2 + 1/3*((2*x + 1)*x - 9)*sqrt(-x^2 - 2*x + 3) + 3*x - 4*arcsin(1/2*x + 1/2)`

Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.95

$$\int \left(x + \sqrt{3 - 2x - x^2} \right)^2 dx = 3x + \frac{\sqrt{-x^2 - 2x + 3} (8x^2 + 4x - 36)}{12} - x^2 + \ln \left(x + 1 - \sqrt{-x^2 - 2x + 3} \right) 4i$$

input `int((x + (3 - x^2 - 2*x)^(1/2))^2,x)`

output `3*x + log(x - (3 - x^2 - 2*x)^(1/2)*1i + 1)*4i + ((3 - x^2 - 2*x)^(1/2)*(4*x + 8*x^2 - 36))/12 - x^2`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.05

$$\int \left(x + \sqrt{3 - 2x - x^2}\right)^2 dx = -4\operatorname{asin}\left(\frac{x}{2} + \frac{1}{2}\right) + \frac{2\sqrt{-x^2 - 2x + 3}x^2}{3} + \frac{\sqrt{-x^2 - 2x + 3}x}{3} - 3\sqrt{-x^2 - 2x + 3} - x^2 + 3x + \frac{16}{3}$$

input `int((x+(-x^2-2*x+3)^(1/2))^2,x)`output `(- 12*asin((x + 1)/2) + 2*sqrt(- x**2 - 2*x + 3)*x**2 + sqrt(- x**2 - 2*x + 3)*x - 9*sqrt(- x**2 - 2*x + 3) - 3*x**2 + 9*x + 16)/3`

3.21 $\int (x + \sqrt{3 - 2x - x^2}) dx$

Optimal result	192
Mathematica [A] (verified)	192
Rubi [A] (verified)	193
Maple [A] (verified)	193
Fricas [A] (verification not implemented)	194
Sympy [A] (verification not implemented)	194
Maxima [A] (verification not implemented)	195
Giac [A] (verification not implemented)	195
Mupad [B] (verification not implemented)	195
Reduce [B] (verification not implemented)	196

Optimal result

Integrand size = 16, antiderivative size = 41

$$\int (x + \sqrt{3 - 2x - x^2}) dx = \frac{x^2}{2} + \frac{1}{2}(1 + x)\sqrt{3 - 2x - x^2} - 2 \arcsin\left(\frac{1}{2}(-1 - x)\right)$$

output

```
1/2*x^2+1/2*(1+x)*(-x^2-2*x+3)^(1/2)+2*arcsin(1/2+1/2*x)
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.27

$$\int (x + \sqrt{3 - 2x - x^2}) dx = \frac{x^2}{2} + \frac{1}{2}(1 + x)\sqrt{3 - 2x - x^2} - 4 \arctan\left(\frac{\sqrt{3 - 2x - x^2}}{3 + x}\right)$$

input

```
Integrate[x + Sqrt[3 - 2*x - x^2], x]
```

output

```
x^2/2 + ((1 + x)*Sqrt[3 - 2*x - x^2])/2 - 4*ArcTan[Sqrt[3 - 2*x - x^2]/(3 + x)]
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.062$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\sqrt{-x^2 - 2x + 3} + x \right) dx$$

↓ 2009

$$-2 \arcsin \left(\frac{1}{2}(-x - 1) \right) + \frac{x^2}{2} + \frac{1}{2}(x + 1)\sqrt{-x^2 - 2x + 3}$$

input `Int[x + Sqrt[3 - 2*x - x^2],x]`

output `x^2/2 + ((1 + x)*Sqrt[3 - 2*x - x^2])/2 - 2*ArcSin[(-1 - x)/2]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.83

method	result
default	$\frac{x^2}{2} - \frac{(-2x-2)\sqrt{-x^2-2x+3}}{4} + 2 \arcsin \left(\frac{1}{2} + \frac{x}{2} \right)$
parts	$\frac{x^2}{2} - \frac{(-2x-2)\sqrt{-x^2-2x+3}}{4} + 2 \arcsin \left(\frac{1}{2} + \frac{x}{2} \right)$
trager	$\frac{x^2}{2} + \left(\frac{1}{2} + \frac{x}{2} \right) \sqrt{-x^2 - 2x + 3} + 2 \text{RootOf} \left(_Z^2 + 1 \right) \ln \left(- \text{RootOf} \left(_Z^2 + 1 \right) x + \sqrt{-x^2 - 2x} \right)$

input `int(x+(-x^2-2*x+3)^(1/2),x,method=_RETURNVERBOSE)`

output `1/2*x^2-1/4*(-2*x-2)*(-x^2-2*x+3)^(1/2)+2*arcsin(1/2+1/2*x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.27

$$\int \left(x + \sqrt{3 - 2x - x^2} \right) dx = \frac{1}{2}x^2 + \frac{1}{2}\sqrt{-x^2 - 2x + 3}(x + 1) - 2 \arctan \left(\frac{\sqrt{-x^2 - 2x + 3}(x + 1)}{x^2 + 2x - 3} \right)$$

input `integrate(x+(-x^2-2*x+3)^(1/2),x, algorithm="fricas")`

output `1/2*x^2 + 1/2*sqrt(-x^2 - 2*x + 3)*(x + 1) - 2*arctan(sqrt(-x^2 - 2*x + 3) * (x + 1)/(x^2 + 2*x - 3))`

Sympy [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.78

$$\int \left(x + \sqrt{3 - 2x - x^2} \right) dx = \frac{x^2}{2} + \left(\frac{x}{2} + \frac{1}{2} \right) \sqrt{-x^2 - 2x + 3} + 2 \operatorname{asin} \left(\frac{x}{2} + \frac{1}{2} \right)$$

input `integrate(x+(-x**2-2*x+3)**(1/2),x)`

output `x**2/2 + (x/2 + 1/2)*sqrt(-x**2 - 2*x + 3) + 2*asin(x/2 + 1/2)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.05

$$\int \left(x + \sqrt{3 - 2x - x^2} \right) dx = \frac{1}{2} x^2 + \frac{1}{2} \sqrt{-x^2 - 2x + 3} x + \frac{1}{2} \sqrt{-x^2 - 2x + 3} - 2 \arcsin \left(-\frac{1}{2} x - \frac{1}{2} \right)$$

input `integrate(x+(-x^2-2*x+3)^(1/2),x, algorithm="maxima")`output `1/2*x^2 + 1/2*sqrt(-x^2 - 2*x + 3)*x + 1/2*sqrt(-x^2 - 2*x + 3) - 2*arcsin(-1/2*x - 1/2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.76

$$\int \left(x + \sqrt{3 - 2x - x^2} \right) dx = \frac{1}{2} x^2 + \frac{1}{2} \sqrt{-x^2 - 2x + 3}(x + 1) + 2 \arcsin \left(\frac{1}{2} x + \frac{1}{2} \right)$$

input `integrate(x+(-x^2-2*x+3)^(1/2),x, algorithm="giac")`output `1/2*x^2 + 1/2*sqrt(-x^2 - 2*x + 3)*(x + 1) + 2*arcsin(1/2*x + 1/2)`**Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.78

$$\int \left(x + \sqrt{3 - 2x - x^2} \right) dx = 2 \operatorname{asin} \left(\frac{x}{2} + \frac{1}{2} \right) + \left(\frac{x}{2} + \frac{1}{2} \right) \sqrt{-x^2 - 2x + 3} + \frac{x^2}{2}$$

input `int(x + (3 - x^2 - 2*x)^(1/2),x)`output `2*asin(x/2 + 1/2) + (x/2 + 1/2)*(3 - x^2 - 2*x)^(1/2) + x^2/2`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \left(x + \sqrt{3 - 2x - x^2} \right) dx = 2 \operatorname{asin} \left(\frac{x}{2} + \frac{1}{2} \right) + \frac{\sqrt{-x^2 - 2x + 3} x}{2} + \frac{\sqrt{-x^2 - 2x + 3}}{2} + \frac{x^2}{2}$$

input `int(x+(-x^2-2*x+3)^(1/2),x)`

output `(4*asin((x + 1)/2) + sqrt(- x**2 - 2*x + 3)*x + sqrt(- x**2 - 2*x + 3) + x**2)/2`

3.22 $\int \frac{1}{x + \sqrt{3 - 2x - x^2}} dx$

Optimal result	197
Mathematica [A] (verified)	198
Rubi [A] (warning: unable to verify)	198
Maple [B] (verified)	202
Fricas [B] (verification not implemented)	203
Sympy [F]	204
Maxima [F]	204
Giac [B] (verification not implemented)	205
Mupad [F(-1)]	206
Reduce [B] (verification not implemented)	206

Optimal result

Integrand size = 18, antiderivative size = 180

$$\int \frac{1}{x + \sqrt{3 - 2x - x^2}} dx = \arctan\left(\frac{\sqrt{3} - \sqrt{3 - 2x - x^2}}{x}\right) - \frac{1}{2} \log\left(-\frac{3 - x - \sqrt{3}\sqrt{3 - 2x - x^2}}{x^2}\right) + \frac{1}{14}(7 + \sqrt{7}) \log\left(1 + \sqrt{3} - \sqrt{7} - \frac{\sqrt{3}(\sqrt{3} - \sqrt{3 - 2x - x^2})}{x}\right) + \frac{1}{14}(7 - \sqrt{7}) \log\left(1 + \sqrt{3} + \sqrt{7} - \frac{\sqrt{3}(\sqrt{3} - \sqrt{3 - 2x - x^2})}{x}\right)$$

output

```
arctan((3^(1/2)-(-x^2-2*x+3)^(1/2))/x)-1/2*ln(-(3-x-3^(1/2)*(-x^2-2*x+3)^(1/2))/x^2)+1/14*(7+7^(1/2))*ln(1+3^(1/2)-7^(1/2)-3^(1/2)*(3^(1/2)-(-x^2-2*x+3)^(1/2))/x)+1/14*(7-7^(1/2))*ln(1+3^(1/2)+7^(1/2)-3^(1/2)*(3^(1/2)-(-x^2-2*x+3)^(1/2))/x)
```

Mathematica [A] (verified)

Time = 0.42 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.62

$$\int \frac{1}{x + \sqrt{3 - 2x - x^2}} dx = \frac{1}{14} \left(-14 \arctan \left(\frac{\sqrt{3 - 2x - x^2}}{3 + x} \right) - 7 \log(-1 + x) \right. \\ \left. - (-7 + \sqrt{7}) \log \left(-2 + \sqrt{7}(-1 + x) + 2x - \sqrt{3 - 2x - x^2} \right) \right. \\ \left. + (7 + \sqrt{7}) \log \left(2 + \sqrt{7}(-1 + x) - 2x + \sqrt{3 - 2x - x^2} \right) \right)$$

input

```
Integrate[(x + Sqrt[3 - 2*x - x^2])^(-1), x]
```

output

```
(-14*ArcTan[Sqrt[3 - 2*x - x^2]/(3 + x)] - 7*Log[-1 + x] - (-7 + Sqrt[7])*
Log[-2 + Sqrt[7]*(-1 + x) + 2*x - Sqrt[3 - 2*x - x^2]] + (7 + Sqrt[7])*Log
[2 + Sqrt[7]*(-1 + x) - 2*x + Sqrt[3 - 2*x - x^2]])/14
```

Rubi [A] (warning: unable to verify)Time = 0.52 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.09, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.611$, Rules used = {7285, 2142, 27, 452, 216, 240, 1142, 27, 1083, 219, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{-x^2 - 2x + 3} + x} dx$$

↓ 7285

$$2 \int \frac{-\frac{\sqrt{3}(\sqrt{3 - \sqrt{-x^2 - 2x + 3}})^2}{x^2} + \frac{2(\sqrt{3 - \sqrt{-x^2 - 2x + 3}})}{x} + \sqrt{3}}{\left(\frac{(\sqrt{3 - \sqrt{-x^2 - 2x + 3}})^2}{x^2} + 1 \right) \left(\frac{\sqrt{3}(\sqrt{3 - \sqrt{-x^2 - 2x + 3}})^2}{x^2} - \frac{2(1 + \sqrt{3})(\sqrt{3 - \sqrt{-x^2 - 2x + 3}})}{x} - \sqrt{3} + 2 \right)} dx \left(-\frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x} \right)$$

↓ 2142

$$2 \left(\frac{1}{32} \int -\frac{16 \left(1 - \frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x} \right)}{\frac{(\sqrt{3} - \sqrt{-x^2 - 2x + 3})^2}{x^2} + 1} d \left(-\frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x} \right) + \frac{1}{32} \int \frac{16 \left(-\frac{\sqrt{3}(\sqrt{3} - \sqrt{-x^2 - 2x + 3})}{x} + \right)}{\frac{\sqrt{3}(\sqrt{3} - \sqrt{-x^2 - 2x + 3})^2}{x^2} - \frac{2(1 + \sqrt{3})(\sqrt{3} - \sqrt{-x^2 - 2x + 3})}{x} + \sqrt{3} + 2} d \left(-\frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x} \right) - \frac{1}{2} \int \frac{1 - \sqrt{3}}{(\sqrt{3} - \sqrt{-x^2 - 2x + 3})} d \left(-\frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x} \right) \right)$$

↓ 27

$$2 \left(\frac{1}{2} \int \frac{-\frac{\sqrt{3}(\sqrt{3} - \sqrt{-x^2 - 2x + 3})}{x} + \sqrt{3} + 2}{\frac{\sqrt{3}(\sqrt{3} - \sqrt{-x^2 - 2x + 3})^2}{x^2} - \frac{2(1 + \sqrt{3})(\sqrt{3} - \sqrt{-x^2 - 2x + 3})}{x} - \sqrt{3} + 2} d \left(-\frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x} \right) - \frac{1}{2} \int \frac{1 - \sqrt{3}}{(\sqrt{3} - \sqrt{-x^2 - 2x + 3})} d \left(-\frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x} \right) \right)$$

↓ 452

$$2 \left(\frac{1}{2} \left(- \int \frac{1}{\frac{(\sqrt{3} - \sqrt{-x^2 - 2x + 3})^2}{x^2} + 1} d \left(-\frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x} \right) - \int -\frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x \left(\frac{(\sqrt{3} - \sqrt{-x^2 - 2x + 3})^2}{x^2} + 1 \right)} d \left(-\frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x} \right) \right) \right)$$

↓ 216

$$2 \left(\frac{1}{2} \left(\arctan \left(\frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x} \right) - \int -\frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x \left(\frac{(\sqrt{3} - \sqrt{-x^2 - 2x + 3})^2}{x^2} + 1 \right)} d \left(-\frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x} \right) \right) + \frac{1}{2} \int \frac{1 - \sqrt{3}}{(\sqrt{3} - \sqrt{-x^2 - 2x + 3})} d \left(-\frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x} \right) \right)$$

↓ 240

$$2 \left(\frac{1}{2} \int \frac{-\frac{\sqrt{3}(\sqrt{3} - \sqrt{-x^2 - 2x + 3})}{x} + \sqrt{3} + 2}{\frac{\sqrt{3}(\sqrt{3} - \sqrt{-x^2 - 2x + 3})^2}{x^2} - \frac{2(1 + \sqrt{3})(\sqrt{3} - \sqrt{-x^2 - 2x + 3})}{x} - \sqrt{3} + 2} d \left(-\frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x} \right) + \frac{1}{2} \left(\arctan \left(\frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x} \right) - \int -\frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x \left(\frac{(\sqrt{3} - \sqrt{-x^2 - 2x + 3})^2}{x^2} + 1 \right)} d \left(-\frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x} \right) \right) \right)$$

↓ 1142

$$2 \left(\frac{1}{2} \left(\int \frac{1}{\frac{\sqrt{3}(\sqrt{3} - \sqrt{-x^2 - 2x + 3})^2}{x^2} - \frac{2(1 + \sqrt{3})(\sqrt{3} - \sqrt{-x^2 - 2x + 3})}{x} - \sqrt{3} + 2} d \left(-\frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x} \right) + \frac{1}{2} \int \frac{1 - \sqrt{3}}{\sqrt{3}(\sqrt{3} - \sqrt{-x^2 - 2x + 3})} d \left(-\frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x} \right) \right) \right)$$

↓ 27

$$\begin{aligned}
 & 2 \left(\frac{1}{2} \left(\int \frac{1}{\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2} - \frac{2(1+\sqrt{3})(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - \sqrt{3}+2} d\left(-\frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{x}\right) + \int \frac{1}{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})} \right) \right. \\
 & \qquad \qquad \qquad \downarrow \text{1083} \\
 & 2 \left(\frac{1}{2} \left(\int \frac{-\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} + \sqrt{3}+1}{\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2} - \frac{2(1+\sqrt{3})(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - \sqrt{3}+2} d\left(-\frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{x}\right) - 2 \int \frac{1}{28 - (\sqrt{3}-\sqrt{-x^2-2x+3})} \right) \right. \\
 & \qquad \qquad \qquad \downarrow \text{219} \\
 & 2 \left(\frac{1}{2} \left(\int \frac{-\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} + \sqrt{3}+1}{\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2} - \frac{2(1+\sqrt{3})(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - \sqrt{3}+2} d\left(-\frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{x}\right) + \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{2\sqrt{7}x}\right)}{\sqrt{7}} \right) \right. \\
 & \qquad \qquad \qquad \downarrow \text{1103} \\
 & 2 \left(\frac{1}{2} \left(\arctan\left(\frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{x}\right) - \frac{1}{2} \log\left(\frac{(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2} + 1\right) \right) \right) + \frac{1}{2} \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{2\sqrt{7}x}\right)}{\sqrt{7}} \right)
 \end{aligned}$$

input

```
Int[(x + Sqrt[3 - 2*x - x^2])^(-1), x]
```

output

```
2*((ArcTan[(Sqrt[3] - Sqrt[3 - 2*x - x^2])/x] - Log[1 + (Sqrt[3] - Sqrt[3 - 2*x - x^2])^2/x^2])/2) + (ArcTanh[(Sqrt[3] - Sqrt[3 - 2*x - x^2])/(2*Sqrt[7]*x)]/Sqrt[7] + Log[2 - Sqrt[3] - (2*(1 + Sqrt[3])*(Sqrt[3] - Sqrt[3 - 2*x - x^2]))]/x + (Sqrt[3]*(Sqrt[3] - Sqrt[3 - 2*x - x^2])^2/x^2)/2)/2
```

Definitions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 216 $\text{Int}[((a_) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$
- rule 219 $\text{Int}[((a_) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 240 $\text{Int}[(x_)/((a_) + (b_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x^2, x]]/(2*b), x] /; \text{FreeQ}\{a, b\}, x]$
- rule 452 $\text{Int}[((c_) + (d_*)(x_))/((a_) + (b_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[c \text{ Int}[1/(a + b*x^2), x], x] + \text{Simp}[d \text{ Int}[x/(a + b*x^2), x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[b*c^2 + a*d^2, 0]$
- rule 1083 $\text{Int}[((a_) + (b_*)(x_) + (c_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[-2 \text{ Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}\{a, b, c\}, x]$
- rule 1103 $\text{Int}[((d_) + (e_*)(x_))/((a_) + (b_*)(x_) + (c_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$
- rule 1142 $\text{Int}[((d_) + (e_*)(x_))/((a_) + (b_*)(x_) + (c_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \text{ Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \text{ Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x]$

rule 2142

```
Int[(Px_)/(((a_) + (b_.)*(x_) + (c_.)*(x_)^2)*((d_) + (f_.)*(x_)^2)), x_Symbol]
:> With[{A = Coeff[Px, x, 0], B = Coeff[Px, x, 1], C = Coeff[Px, x, 2]},
q = c^2*d^2 + b^2*d*f - 2*a*c*d*f + a^2*f^2},
Simp[1/q Int[(A*c^2*d - a*c*C*d + A*b^2*f - a*b*B*f - a*A*c*f + a^2*C*f + c*(B*c*d - b*C*d + A*b*f -
a*B*f)*x)/(a + b*x + c*x^2), x], x] + Simp[1/q Int[(c*C*d^2 + b*B*d*f -
A*c*d*f - a*C*d*f + a*A*f^2 - f*(B*c*d - b*C*d + A*b*f - a*B*f)*x)/(d + f*x^2), x], x]
;/; NeQ[q, 0]] /; FreeQ[{a, b, c, d, f}, x] && PolyQ[Px, x, 2]
```

rule 7285

```
Int[u_, x_Symbol] :> With[{lst = FunctionOfSquareRootOfQuadratic[u, x]},
Simp[2 Subst[Int[lst[[1]], x], x, lst[[2]]], x] /; !FalseQ[lst] && EqQ[lst[[3]], 1]]
;/; EulerIntegrandQ[u, x]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 358 vs. 2(142) = 284.

Time = 0.54 (sec) , antiderivative size = 359, normalized size of antiderivative = 1.99

method	result
default	$\sqrt{7} \left(-\frac{\sqrt{-4\left(x+\frac{1}{2}-\frac{\sqrt{7}}{2}\right)^2+4(-1-\sqrt{7})\left(x+\frac{1}{2}-\frac{\sqrt{7}}{2}\right)+8-2\sqrt{7}}}{4} - \frac{(-1-\sqrt{7}) \arcsin\left(\frac{x+1}{\sqrt{2-\frac{\sqrt{7}}{2}+\frac{(-1-\sqrt{7})^2}{4}}}\right)}{4} + \frac{(2-\frac{\sqrt{7}}{2}) \operatorname{arctanh}\left(\frac{-\frac{1}{2}+\frac{\sqrt{7}}{2}}{-\frac{1}{2}+\frac{\sqrt{7}}{2}}\right)}{4} \right)$
trager	Expression too large to display

input

```
int(1/(x+(-x^2-2*x+3)^(1/2)),x,method=_RETURNVERBOSE)
```

output

```

1/7*7^(1/2)*(-1/4*(-4*(x+1/2-1/2*7^(1/2))^2+4*(-1-7^(1/2))*(x+1/2-1/2*7^(1/2))+8-2*7^(1/2))^(1/2)-1/4*(-1-7^(1/2))*arcsin(1/(2-1/2*7^(1/2)+1/4*(-1-7^(1/2))^2)^(1/2)*(x+1))+1/2*(2-1/2*7^(1/2))/(-1/2+1/2*7^(1/2))*arctanh((4-7^(1/2)+(-1-7^(1/2))*(x+1/2-1/2*7^(1/2)))/(-1/2+1/2*7^(1/2)))/(-4*(x+1/2-1/2*7^(1/2))^2+4*(-1-7^(1/2))*(x+1/2-1/2*7^(1/2))+8-2*7^(1/2))^(1/2))+1/7*7^(1/2)*(1/4*(-4*(x+1/2+1/2*7^(1/2))^2+4*(-1+7^(1/2))*(x+1/2+1/2*7^(1/2))+8+2*7^(1/2))^(1/2)+1/4*(-1+7^(1/2))*arcsin(1/(2+1/2*7^(1/2)+1/4*(-1+7^(1/2))^2)^(1/2)*(x+1))-1/2*(2+1/2*7^(1/2))/(1/2+1/2*7^(1/2))*arctanh((4+7^(1/2)+(-1+7^(1/2))*(x+1/2+1/2*7^(1/2)))/(1/2+1/2*7^(1/2)))/(-4*(x+1/2+1/2*7^(1/2))^2+4*(-1+7^(1/2))*(x+1/2+1/2*7^(1/2))+8+2*7^(1/2))^(1/2))+1/4*ln(2*x^2+2*x-3)+1/14*7^(1/2)*arctanh(1/14*(4*x+2)*7^(1/2))

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 372 vs. $2(136) = 272$.

Time = 0.12 (sec) , antiderivative size = 372, normalized size of antiderivative = 2.07

$$\begin{aligned}
& \int \frac{1}{x + \sqrt{3 - 2x - x^2}} dx \\
&= \frac{1}{56} \sqrt{7} \log \left(\frac{24x^4 + 62x^3 - 153x^2 + 2\sqrt{7}(3x^4 + x^3 - 45x^2 + 45x) - (14x^3 - 84x^2 + \sqrt{7}(8x^3 - 30x^2 - 12x + 9))}{4x^4 + 8x^3 - 8x^2 - 12x + 9} \right) \\
&+ \frac{1}{56} \sqrt{7} \log \left(\frac{24x^4 + 62x^3 - 153x^2 - 2\sqrt{7}(3x^4 + x^3 - 45x^2 + 45x) + (14x^3 - 84x^2 - \sqrt{7}(8x^3 - 30x^2 - 12x + 9))}{4x^4 + 8x^3 - 8x^2 - 12x + 9} \right) \\
&+ \frac{1}{28} \sqrt{7} \log \left(\frac{2x^2 + \sqrt{7}(2x + 1) + 2x + 4}{2x^2 + 2x - 3} \right) - \frac{1}{2} \arctan \left(\frac{\sqrt{-x^2 - 2x + 3}(x + 1)}{x^2 + 2x - 3} \right) \\
&+ \frac{1}{4} \log(2x^2 + 2x - 3) - \frac{1}{8} \log \left(\frac{2\sqrt{-x^2 - 2x + 3}x + 2x - 3}{x^2} \right) \\
&+ \frac{1}{8} \log \left(-\frac{2\sqrt{-x^2 - 2x + 3}x - 2x + 3}{x^2} \right)
\end{aligned}$$

input

```

integrate(1/(x+(-x^2-2*x+3)^(1/2)),x, algorithm="fricas")

```

output

```
1/56*sqrt(7)*log((24*x^4 + 62*x^3 - 153*x^2 + 2*sqrt(7)*(3*x^4 + x^3 - 45*
x^2 + 45*x) - (14*x^3 - 84*x^2 + sqrt(7)*(8*x^3 - 30*x^2 + 27*x - 27) + 12
6*x)*sqrt(-x^2 - 2*x + 3) + 180*x - 135)/(4*x^4 + 8*x^3 - 8*x^2 - 12*x + 9
)) + 1/56*sqrt(7)*log((24*x^4 + 62*x^3 - 153*x^2 - 2*sqrt(7)*(3*x^4 + x^3
- 45*x^2 + 45*x) + (14*x^3 - 84*x^2 - sqrt(7)*(8*x^3 - 30*x^2 + 27*x - 27)
+ 126*x)*sqrt(-x^2 - 2*x + 3) + 180*x - 135)/(4*x^4 + 8*x^3 - 8*x^2 - 12*
x + 9)) + 1/28*sqrt(7)*log((2*x^2 + sqrt(7)*(2*x + 1) + 2*x + 4)/(2*x^2 +
2*x - 3)) - 1/2*arctan(sqrt(-x^2 - 2*x + 3)*(x + 1)/(x^2 + 2*x - 3)) + 1/4
*log(2*x^2 + 2*x - 3) - 1/8*log((2*sqrt(-x^2 - 2*x + 3)*x + 2*x - 3)/x^2)
+ 1/8*log(-(2*sqrt(-x^2 - 2*x + 3)*x - 2*x + 3)/x^2)
```

Sympy [F]

$$\int \frac{1}{x + \sqrt{3 - 2x - x^2}} dx = \int \frac{1}{x + \sqrt{-x^2 - 2x + 3}} dx$$

input

```
integrate(1/(x+(-x**2-2*x+3)**(1/2)),x)
```

output

```
Integral(1/(x + sqrt(-x**2 - 2*x + 3)), x)
```

Maxima [F]

$$\int \frac{1}{x + \sqrt{3 - 2x - x^2}} dx = \int \frac{1}{x + \sqrt{-x^2 - 2x + 3}} dx$$

input

```
integrate(1/(x+(-x^2-2*x+3)^(1/2)),x, algorithm="maxima")
```

output

```
integrate(1/(x + sqrt(-x^2 - 2*x + 3)), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 287 vs. $2(136) = 272$.

Time = 0.16 (sec) , antiderivative size = 287, normalized size of antiderivative = 1.59

$$\int \frac{1}{x + \sqrt{3 - 2x - x^2}} dx$$

$$= -\frac{1}{28} \sqrt{7} \log \left(\frac{|4x - 2\sqrt{7} + 2|}{|4x + 2\sqrt{7} + 2|} \right) + \frac{1}{28} \sqrt{7} \log \left(\frac{\left| -2\sqrt{7} + \frac{6(\sqrt{-x^2 - 2x + 3} - 2)}{x+1} + 4 \right|}{\left| 2\sqrt{7} + \frac{6(\sqrt{-x^2 - 2x + 3} - 2)}{x+1} + 4 \right|} \right)$$

$$- \frac{1}{28} \sqrt{7} \log \left(\frac{\left| -2\sqrt{7} + \frac{2(\sqrt{-x^2 - 2x + 3} - 2)}{x+1} - 4 \right|}{\left| 2\sqrt{7} + \frac{2(\sqrt{-x^2 - 2x + 3} - 2)}{x+1} - 4 \right|} \right)$$

$$+ \frac{1}{2} \arcsin \left(\frac{1}{2}x + \frac{1}{2} \right) + \frac{1}{4} \log (|2x^2 + 2x - 3|)$$

$$+ \frac{1}{4} \log \left(\frac{4(\sqrt{-x^2 - 2x + 3} - 2)}{x+1} + \frac{3(\sqrt{-x^2 - 2x + 3} - 2)^2}{(x+1)^2} - 1 \right)$$

$$- \frac{1}{4} \log \left(-\frac{4(\sqrt{-x^2 - 2x + 3} - 2)}{x+1} + \frac{(\sqrt{-x^2 - 2x + 3} - 2)^2}{(x+1)^2} - 3 \right)$$

input `integrate(1/(x+(-x^2-2*x+3)^(1/2)),x, algorithm="giac")`

output `-1/28*sqrt(7)*log(abs(4*x - 2*sqrt(7) + 2)/abs(4*x + 2*sqrt(7) + 2)) + 1/28*sqrt(7)*log(abs(-2*sqrt(7) + 6*(sqrt(-x^2 - 2*x + 3) - 2)/(x + 1) + 4)/abs(2*sqrt(7) + 6*(sqrt(-x^2 - 2*x + 3) - 2)/(x + 1) + 4)) - 1/28*sqrt(7)*log(abs(-2*sqrt(7) + 2*(sqrt(-x^2 - 2*x + 3) - 2)/(x + 1) - 4)/abs(2*sqrt(7) + 2*(sqrt(-x^2 - 2*x + 3) - 2)/(x + 1) - 4)) + 1/2*arcsin(1/2*x + 1/2) + 1/4*log(abs(2*x^2 + 2*x - 3)) + 1/4*log(abs(4*(sqrt(-x^2 - 2*x + 3) - 2)/(x + 1) + 3*(sqrt(-x^2 - 2*x + 3) - 2)^2/(x + 1)^2 - 1)) - 1/4*log(abs(-4*(sqrt(-x^2 - 2*x + 3) - 2)/(x + 1) + (sqrt(-x^2 - 2*x + 3) - 2)^2/(x + 1)^2 - 3))`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{x + \sqrt{3 - 2x - x^2}} dx = \int \frac{1}{x + \sqrt{-x^2 - 2x + 3}} dx$$

input `int(1/(x + (3 - x^2 - 2*x)^(1/2)),x)`output `int(1/(x + (3 - x^2 - 2*x)^(1/2)), x)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.37

$$\int \frac{1}{x + \sqrt{3 - 2x - x^2}} dx = \frac{\operatorname{asin}\left(\frac{x}{2} + \frac{1}{2}\right)}{2} - \frac{\sqrt{7} \log\left(-\sqrt{7} + 3 \tan\left(\frac{\operatorname{asin}\left(\frac{x}{2} + \frac{1}{2}\right)}{2}\right) - 2\right)}{14}$$

$$+ \frac{\sqrt{7} \log\left(\sqrt{7} + 3 \tan\left(\frac{\operatorname{asin}\left(\frac{x}{2} + \frac{1}{2}\right)}{2}\right) - 2\right)}{14}$$

$$+ \frac{\log\left(\sqrt{-x^2 - 2x + 3} + x\right)}{2}$$

input `int(1/(x+(-x^2-2*x+3)^(1/2)),x)`output `(7*asin((x + 1)/2) - sqrt(7)*log(- sqrt(7) + 3*tan(asin((x + 1)/2)/2) - 2) + sqrt(7)*log(sqrt(7) + 3*tan(asin((x + 1)/2)/2) - 2) + 7*log(sqrt(- x* *2 - 2*x + 3) + x))/14`

3.23
$$\int \frac{1}{\left(x + \sqrt{3 - 2x - x^2}\right)^2} dx$$

Optimal result	207
Mathematica [A] (verified)	208
Rubi [A] (warning: unable to verify)	208
Maple [C] (verified)	211
Fricas [A] (verification not implemented)	211
Sympy [F]	212
Maxima [F]	212
Giac [B] (verification not implemented)	213
Mupad [F(-1)]	214
Reduce [B] (verification not implemented)	214

Optimal result

Integrand size = 18, antiderivative size = 172

$$\int \frac{1}{\left(x + \sqrt{3 - 2x - x^2}\right)^2} dx = \frac{2\left(4 - \sqrt{3} + \frac{3(\sqrt{3} - \sqrt{3 - 2x - x^2})}{x}\right)}{7\left(2 - \sqrt{3} - \frac{2(1 + \sqrt{3})(\sqrt{3} - \sqrt{3 - 2x - x^2})}{x} + \frac{\sqrt{3}(\sqrt{3} - \sqrt{3 - 2x - x^2})^2}{x^2}\right)} + \frac{8\operatorname{arctanh}\left(\frac{3 - x - \sqrt{3}x - \sqrt{3}\sqrt{3 - 2x - x^2}}{\sqrt{7}x}\right)}{7\sqrt{7}}$$

output

```
2*(4-3^(1/2))+3*(3^(1/2)-(-x^2-2*x+3)^(1/2))/x/(14-7*3^(1/2)-14*(1+3^(1/2))
)*(3^(1/2)-(-x^2-2*x+3)^(1/2))/x+7*3^(1/2)*(3^(1/2)-(-x^2-2*x+3)^(1/2))^2/
x^2)+8/49*arctanh(1/7*(3-x-x*3^(1/2)-3^(1/2)*(-x^2-2*x+3)^(1/2))*7^(1/2)/x
)*7^(1/2)
```


Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.55

$$\int \frac{1}{(x + \sqrt{3 - 2x - x^2})^2} dx = \frac{3 + 6\sqrt{3 - 2x - x^2} - 2x(4 + \sqrt{3 - 2x - x^2})}{14(-3 + 2x + 2x^2)} + \frac{8 \operatorname{arctanh}\left(\frac{2 - 2x + \sqrt{3 - 2x - x^2}}{\sqrt{7}(-1 + x)}\right)}{7\sqrt{7}}$$

input

```
Integrate[(x + Sqrt[3 - 2*x - x^2])^(-2), x]
```

output

```
(3 + 6*Sqrt[3 - 2*x - x^2] - 2*x*(4 + Sqrt[3 - 2*x - x^2]))/(14*(-3 + 2*x + 2*x^2)) + (8*ArcTanh[(2 - 2*x + Sqrt[3 - 2*x - x^2])/(Sqrt[7]*(-1 + x))])/(7*Sqrt[7])
```

Rubi [A] (warning: unable to verify)

Time = 0.37 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.96, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {7285, 25, 2191, 27, 1083, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(\sqrt{-x^2 - 2x + 3} + x)^2} dx$$

↓ 7285

$$2 \int -\frac{-\frac{\sqrt{3}(\sqrt{3} - \sqrt{-x^2 - 2x + 3})^2}{x^2} + \frac{2(\sqrt{3} - \sqrt{-x^2 - 2x + 3})}{x} + \sqrt{3}}{\left(\frac{\sqrt{3}(\sqrt{3} - \sqrt{-x^2 - 2x + 3})^2}{x^2} - \frac{2(1 + \sqrt{3})(\sqrt{3} - \sqrt{-x^2 - 2x + 3})}{x} - \sqrt{3} + 2\right)^2} d\left(-\frac{\sqrt{3} - \sqrt{-x^2 - 2x + 3}}{x}\right)$$

↓ 25

$$-2 \int \frac{-\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2} + \frac{2(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} + \sqrt{3}}{\left(\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2} - \frac{2(1+\sqrt{3})(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - \sqrt{3} + 2\right)^2} d\left(-\frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{x}\right)$$

↓ 2191

$$2 \left(\frac{1}{28} \int \frac{16}{\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2} - \frac{2(1+\sqrt{3})(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - \sqrt{3} + 2} d\left(-\frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{x}\right) + \frac{1}{7} \left(\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - \sqrt{3} + 2 \right) \right)$$

↓ 27

$$2 \left(\frac{4}{7} \int \frac{1}{\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2} - \frac{2(1+\sqrt{3})(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - \sqrt{3} + 2} d\left(-\frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{x}\right) + \frac{1}{7} \left(\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - \sqrt{3} + 2 \right) \right)$$

↓ 1083

$$2 \left(\frac{\frac{3(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - \sqrt{3} + 4}{7 \left(\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2} - \frac{2(1+\sqrt{3})(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - \sqrt{3} + 2 \right)} - \frac{8}{7} \int \frac{1}{28 - \frac{(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2}} d\left(2(1+\sqrt{3}) - \frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{x}\right) \right)$$

↓ 219

$$2 \left(\frac{4 \operatorname{arctanh}\left(\frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{2\sqrt{7}x}\right)}{7\sqrt{7}} + \frac{\frac{3(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - \sqrt{3} + 4}{7 \left(\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2} - \frac{2(1+\sqrt{3})(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - \sqrt{3} + 2 \right)} \right)$$

input `Int[(x + Sqrt[3 - 2*x - x^2])^(-2), x]`

output $2*((4 - \sqrt{3} + (3*(\sqrt{3} - \sqrt{3 - 2x - x^2}))/x)/(7*(2 - \sqrt{3} - (2*(1 + \sqrt{3})*(\sqrt{3} - \sqrt{3 - 2x - x^2}))/x + (\sqrt{3}*(\sqrt{3} - \sqrt{3 - 2x - x^2})^2)/x^2)) + (4*\text{ArcTanh}[(\sqrt{3} - \sqrt{3 - 2x - x^2})/(2*\sqrt{7}*x))]/(7*\sqrt{7}))$

Defintions of rubi rules used

rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$

rule 27 $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] /; \text{FreeQ}[\text{a}, \text{x}] \&\& \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_) /; \text{FreeQ}[\text{b}, \text{x}]$

rule 219 $\text{Int}[(\text{a}_) + (\text{b}_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2]*\text{Rt}[-\text{b}, 2]))*\text{ArcTanh}[\text{Rt}[-\text{b}, 2]*(x/\text{Rt}[\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{NegQ}[\text{a}/\text{b}] \&\& (\text{GtQ}[\text{a}, 0] \|\| \text{LtQ}[\text{b}, 0])$

rule 1083 $\text{Int}[(\text{a}_) + (\text{b}_.)*(x_) + (\text{c}_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[-2 \quad \text{Subst}[\text{Int}[1/\text{Simp}[\text{b}^2 - 4*\text{a}*c - x^2, \text{x}], \text{x}], \text{x}, \text{b} + 2*c*x], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}\}, \text{x}]$

rule 2191 $\text{Int}[(\text{Pq}_)*((\text{a}_.) + (\text{b}_.)*(x_) + (\text{c}_.)*(x_)^2)^{(\text{p}_)}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{Q} = \text{PolynomialQuotient}[\text{Pq}, \text{a} + \text{b}*x + \text{c}*x^2, \text{x}], \text{f} = \text{Coeff}[\text{PolynomialRemainder}[\text{Pq}, \text{a} + \text{b}*x + \text{c}*x^2, \text{x}], \text{x}, 0], \text{g} = \text{Coeff}[\text{PolynomialRemainder}[\text{Pq}, \text{a} + \text{b}*x + \text{c}*x^2, \text{x}], \text{x}, 1]\}, \text{Simp}[(\text{b}*f - 2*\text{a}*g + (2*c*f - \text{b}*g)*x)*((\text{a} + \text{b}*x + \text{c}*x^2)^{(p+1})/((p+1)*(b^2 - 4*a*c))), \text{x}] + \text{Simp}[1/((p+1)*(b^2 - 4*a*c)) \quad \text{Int}[(\text{a} + \text{b}*x + \text{c}*x^2)^{(p+1)}*\text{ExpandToSum}[(p+1)*(b^2 - 4*a*c)*Q - (2*p+3)*(2*c*f - \text{b}*g), \text{x}], \text{x}], \text{x}]] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}\}, \text{x}] \&\& \text{PolyQ}[\text{Pq}, \text{x}] \&\& \text{NeQ}[\text{b}^2 - 4*\text{a}*c, 0] \&\& \text{LtQ}[\text{p}, -1]$

rule 7285 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{With}[\{\text{lst} = \text{FunctionOfSquareRootOfQuadratic}[\text{u}, \text{x}]\}, \text{Simp}[2 \quad \text{Subst}[\text{Int}[\text{lst}[[1]], \text{x}], \text{x}, \text{lst}[[2]]], \text{x}] /; \text{!FalseQ}[\text{lst}] \&\& \text{EqQ}[\text{lst}[[3]], 1]] /; \text{EulerIntegrandQ}[\text{u}, \text{x}]$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.15 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.61

method	result
trager	$\frac{x(x-3)}{14x^2+14x-21} - \frac{(x-3)\sqrt{-x^2-2x+3}}{7(2x^2+2x-3)} - \frac{4 \operatorname{RootOf}(_Z^2-7) \ln\left(\frac{\operatorname{RootOf}(_Z^2-7)^{x-3} \operatorname{RootOf}(_Z^2-7)^{-7\sqrt{-x^2-2x+3}}}{\operatorname{RootOf}(_Z^2-7)^{x-x+3}}\right)}{49}$
default	Expression too large to display

input `int(1/(x+(-x^2-2*x+3)^(1/2))^2,x,method=_RETURNVERBOSE)`

output `1/7*(x-3)*x/(2*x^2+2*x-3)-1/7*(x-3)/(2*x^2+2*x-3)*(-x^2-2*x+3)^(1/2)-4/49*
RootOf(_Z^2-7)*ln((RootOf(_Z^2-7)*x-3*RootOf(_Z^2-7)-7*(-x^2-2*x+3)^(1/2))
/(RootOf(_Z^2-7)*x-x+3))`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.99

$$\int \frac{1}{(x + \sqrt{3 - 2x - x^2})^2} dx$$

$$= \frac{2\sqrt{7}(2x^2 + 2x - 3) \log\left(\frac{x^4 + 44x^3 - \sqrt{7}(3x^3 + x^2 - 45x + 45)\sqrt{-x^2 - 2x + 3} + 26x^2 - 276x + 207}{4x^4 + 8x^3 - 8x^2 - 12x + 9}\right) + 4\sqrt{7}(2x^2 + 2x - 3) \log\left(\frac{x^4 + 44x^3 - \sqrt{7}(3x^3 + x^2 - 45x + 45)\sqrt{-x^2 - 2x + 3} + 26x^2 - 276x + 207}{4x^4 + 8x^3 - 8x^2 - 12x + 9}\right)}{98(2x^2 + 2x - 3)}$$

input `integrate(1/(x+(-x^2-2*x+3)^(1/2))^2,x, algorithm="fricas")`

output `1/98*(2*sqrt(7)*(2*x^2 + 2*x - 3)*log((x^4 + 44*x^3 - sqrt(7)*(3*x^3 + x^2
- 45*x + 45)*sqrt(-x^2 - 2*x + 3) + 26*x^2 - 276*x + 207)/(4*x^4 + 8*x^3
- 8*x^2 - 12*x + 9)) + 4*sqrt(7)*(2*x^2 + 2*x - 3)*log((2*x^2 + sqrt(7)*(2
*x + 1) + 2*x + 4)/(2*x^2 + 2*x - 3)) - 14*sqrt(-x^2 - 2*x + 3)*(x - 3) -
56*x + 21)/(2*x^2 + 2*x - 3)`

Sympy [F]

$$\int \frac{1}{(x + \sqrt{3 - 2x - x^2})^2} dx = \int \frac{1}{(x + \sqrt{-x^2 - 2x + 3})^2} dx$$

input `integrate(1/(x+(-x**2-2*x+3)**(1/2))**2,x)`

output `Integral((x + sqrt(-x**2 - 2*x + 3))**(-2), x)`

Maxima [F]

$$\int \frac{1}{(x + \sqrt{3 - 2x - x^2})^2} dx = \int \frac{1}{(x + \sqrt{-x^2 - 2x + 3})^2} dx$$

input `integrate(1/(x+(-x^2-2*x+3)^(1/2))^2,x, algorithm="maxima")`

output `integrate((x + sqrt(-x^2 - 2*x + 3))^-2, x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 350 vs. $2(132) = 264$.

Time = 0.17 (sec) , antiderivative size = 350, normalized size of antiderivative = 2.03

$$\int \frac{1}{(x + \sqrt{3 - 2x - x^2})^2} dx$$

$$= -\frac{2}{49} \sqrt{7} \log \left(\frac{|4x - 2\sqrt{7} + 2|}{|4x + 2\sqrt{7} + 2|} \right) + \frac{2}{49} \sqrt{7} \log \left(\frac{\left| -2\sqrt{7} + \frac{6(\sqrt{-x^2 - 2x + 3} - 2)}{x+1} + 4 \right|}{\left| 2\sqrt{7} + \frac{6(\sqrt{-x^2 - 2x + 3} - 2)}{x+1} + 4 \right|} \right)$$

$$- \frac{2}{49} \sqrt{7} \log \left(\frac{\left| -2\sqrt{7} + \frac{2(\sqrt{-x^2 - 2x + 3} - 2)}{x+1} - 4 \right|}{\left| 2\sqrt{7} + \frac{2(\sqrt{-x^2 - 2x + 3} - 2)}{x+1} - 4 \right|} \right) - \frac{8x - 3}{14(2x^2 + 2x - 3)}$$

$$- \frac{8 \left(\frac{5(\sqrt{-x^2 - 2x + 3} - 2)}{x+1} + \frac{26(\sqrt{-x^2 - 2x + 3} - 2)^2}{(x+1)^2} + \frac{11(\sqrt{-x^2 - 2x + 3} - 2)^3}{(x+1)^3} - 6 \right)}{21 \left(\frac{8(\sqrt{-x^2 - 2x + 3} - 2)}{x+1} + \frac{26(\sqrt{-x^2 - 2x + 3} - 2)^2}{(x+1)^2} + \frac{8(\sqrt{-x^2 - 2x + 3} - 2)^3}{(x+1)^3} - \frac{3(\sqrt{-x^2 - 2x + 3} - 2)^4}{(x+1)^4} - 3 \right)}$$

input `integrate(1/(x+(-x^2-2*x+3)^(1/2))^2,x, algorithm="giac")`

output

```
-2/49*sqrt(7)*log(abs(4*x - 2*sqrt(7) + 2)/abs(4*x + 2*sqrt(7) + 2)) + 2/49*sqrt(7)*log(abs(-2*sqrt(7) + 6*(sqrt(-x^2 - 2*x + 3) - 2)/(x + 1) + 4)/abs(2*sqrt(7) + 6*(sqrt(-x^2 - 2*x + 3) - 2)/(x + 1) + 4)) - 2/49*sqrt(7)*log(abs(-2*sqrt(7) + 2*(sqrt(-x^2 - 2*x + 3) - 2)/(x + 1) - 4)/abs(2*sqrt(7) + 2*(sqrt(-x^2 - 2*x + 3) - 2)/(x + 1) - 4)) - 1/14*(8*x - 3)/(2*x^2 + 2*x - 3) - 8/21*(5*(sqrt(-x^2 - 2*x + 3) - 2)/(x + 1) + 26*(sqrt(-x^2 - 2*x + 3) - 2)^2/(x + 1)^2 + 11*(sqrt(-x^2 - 2*x + 3) - 2)^3/(x + 1)^3 - 6)/(8*(sqrt(-x^2 - 2*x + 3) - 2)/(x + 1) + 26*(sqrt(-x^2 - 2*x + 3) - 2)^2/(x + 1)^2 + 8*(sqrt(-x^2 - 2*x + 3) - 2)^3/(x + 1)^3 - 3*(sqrt(-x^2 - 2*x + 3) - 2)^4/(x + 1)^4 - 3)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(x + \sqrt{3 - 2x - x^2})^2} dx = \int \frac{1}{(x + \sqrt{-x^2 - 2x + 3})^2} dx$$

input `int(1/(x + (3 - x^2 - 2*x)^(1/2))^2,x)`output `int(1/(x + (3 - x^2 - 2*x)^(1/2))^2, x)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.85

$$\int \frac{1}{(x + \sqrt{3 - 2x - x^2})^2} dx$$

$$= \frac{-4\sqrt{-x^2 - 2x + 3}\sqrt{7}\log\left(-\sqrt{7} + 3\tan\left(\frac{\operatorname{asin}\left(\frac{x+1}{2}\right)}{2}\right) - 2\right) + 4\sqrt{-x^2 - 2x + 3}\sqrt{7}\log\left(\sqrt{7} + 3\tan\left(\frac{\operatorname{asin}\left(\frac{x+1}{2}\right)}{2}\right) - 2\right)}{49}$$

input `int(1/(x+(-x^2-2*x+3)^(1/2))^2,x)`output `(- 4*sqrt(- x**2 - 2*x + 3)*sqrt(7)*log(- sqrt(7) + 3*tan(asin((x + 1)/2)/2) - 2) + 4*sqrt(- x**2 - 2*x + 3)*sqrt(7)*log(sqrt(7) + 3*tan(asin((x + 1)/2)/2) - 2) - 56*sqrt(- x**2 - 2*x + 3) - 4*sqrt(7)*log(- sqrt(7) + 3*tan(asin((x + 1)/2)/2) - 2)*x + 4*sqrt(7)*log(sqrt(7) + 3*tan(asin((x + 1)/2)/2) - 2)*x - 49*x - 21)/(49*(sqrt(- x**2 - 2*x + 3) + x))`

3.24
$$\int \frac{1}{\left(x + \sqrt{3 - 2x - x^2}\right)^3} dx$$

Optimal result	215
Mathematica [A] (verified)	216
Rubi [A] (warning: unable to verify)	216
Maple [C] (verified)	219
Fricas [A] (verification not implemented)	220
Sympy [F]	220
Maxima [F]	221
Giac [A] (verification not implemented)	221
Mupad [F(-1)]	222
Reduce [B] (verification not implemented)	222

Optimal result

Integrand size = 18, antiderivative size = 311

$$\begin{aligned} & \int \frac{1}{\left(x + \sqrt{3 - 2x - x^2}\right)^3} dx \\ &= -\frac{4\left(9 - 5\sqrt{3} + \frac{(21+5\sqrt{3})(\sqrt{3}-\sqrt{3-2x-x^2})}{x}\right)}{21\left(2 - \sqrt{3} - \frac{2(1+\sqrt{3})(\sqrt{3}-\sqrt{3-2x-x^2})}{x} + \frac{\sqrt{3}(\sqrt{3}-\sqrt{3-2x-x^2})^2}{x^2}\right)^2} \\ & \quad -\frac{2\left(43 - 6\sqrt{3} + \frac{(49+6\sqrt{3})(\sqrt{3}-\sqrt{3-2x-x^2})}{x}\right)}{49\sqrt{3}\left(2 - \sqrt{3} - \frac{2(1+\sqrt{3})(\sqrt{3}-\sqrt{3-2x-x^2})}{x} + \frac{\sqrt{3}(\sqrt{3}-\sqrt{3-2x-x^2})^2}{x^2}\right)} \\ & \quad + \frac{12\operatorname{arctanh}\left(\frac{3-x-\sqrt{3}x-\sqrt{3}\sqrt{3-2x-x^2}}{\sqrt{7}x}\right)}{49\sqrt{7}} \end{aligned}$$

output

$$\frac{1}{21}(-36+20\sqrt{3}-4(21+5\sqrt{3})(\sqrt{3}-(-x^2-2x+3)^{1/2})/x)/(2-3^{1/2}-2(1+3^{1/2})(\sqrt{3}-(-x^2-2x+3)^{1/2})/x+3^{1/2})(\sqrt{3}-(-x^2-2x+3)^{1/2})^2/x^2-2/147(43-6\sqrt{3}+(49+6\sqrt{3})(\sqrt{3}-(-x^2-2x+3)^{1/2})/x)*3^{1/2}/(2-3^{1/2}-2(1+3^{1/2})(\sqrt{3}-(-x^2-2x+3)^{1/2})/x+3^{1/2})(\sqrt{3}-(-x^2-2x+3)^{1/2})^2/x^2+12/343\operatorname{arctanh}(1/7(3-x-x\sqrt{3}-3^{1/2})(\sqrt{3}-(-x^2-2x+3)^{1/2}))/x)*7^{1/2}$$

Mathematica [A] (verified)

Time = 0.54 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.37

$$\int \frac{1}{(x + \sqrt{3 - 2x - x^2})^3} dx$$

$$= \frac{7(-279+300x+26x^2-48x^3)}{(-3+2x+2x^2)^2} + \frac{14\sqrt{3-2x-x^2}(15+83x-58x^2-34x^3)}{(-3+2x+2x^2)^2} + 48\sqrt{7}\operatorname{arctanh}\left(\frac{2-2x+\sqrt{3-2x-x^2}}{\sqrt{7}(-1+x)}\right)$$

1372

input

Integrate[(x + Sqrt[3 - 2*x - x^2])^(-3), x]

output

$$\frac{((7*(-279 + 300*x + 26*x^2 - 48*x^3))/(-3 + 2*x + 2*x^2)^2 + (14*\operatorname{Sqrt}[3 - 2*x - x^2]*(15 + 83*x - 58*x^2 - 34*x^3))/(-3 + 2*x + 2*x^2)^2 + 48*\operatorname{Sqrt}[7]*\operatorname{ArcTanh}[(2 - 2*x + \operatorname{Sqrt}[3 - 2*x - x^2])/(\operatorname{Sqrt}[7]*(-1 + x))])}{1372}$$

Rubi [A] (warning: unable to verify)Time = 0.55 (sec) , antiderivative size = 305, normalized size of antiderivative = 0.98, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {7285, 2191, 27, 2191, 27, 1083, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(\sqrt{-x^2 - 2x + 3} + x)^3} dx$$

↓ 7285

$$2 \int \frac{-\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})^4}{x^4} + \frac{2(\sqrt{3}-\sqrt{-x^2-2x+3})^3}{x^3} + \frac{2(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} + \sqrt{3}}{\left(\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2} - \frac{2(1+\sqrt{3})(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - \sqrt{3} + 2\right)^3} d\left(-\frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{x}\right)$$

↓ 2191

$$2 \left(-\frac{1}{56} \int \frac{8 \left(-\frac{21(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2} - \frac{42(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} + 16\sqrt{3} + 21 \right)}{3 \left(\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2} - \frac{2(1+\sqrt{3})(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - \sqrt{3} + 2 \right)^2} d\left(-\frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{x}\right) - \frac{7}{21} \left(\frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{x} \right) \right)$$

↓ 27

$$2 \left(\frac{1}{21} \int \frac{-\frac{21(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2} - \frac{42(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} + 16\sqrt{3} + 21}{\left(\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2} - \frac{2(1+\sqrt{3})(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - \sqrt{3} + 2\right)^2} d\left(-\frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{x}\right) - \frac{7}{21} \left(\frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{x} \right) \right)$$

↓ 2191

$$2 \left(\frac{1}{21} \left(\frac{-\frac{(18+49\sqrt{3})(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - 43\sqrt{3} + 18}{7 \left(\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2} - \frac{2(1+\sqrt{3})(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - \sqrt{3} + 2 \right)} - \frac{1}{28} \int \frac{7}{\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2} - \frac{2(1+\sqrt{3})(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - \sqrt{3} + 2} d\left(-\frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{x}\right) + \frac{7}{21} \left(\frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{x} \right) \right) \right)$$

↓ 27

$$2 \left(\frac{1}{21} \left(\frac{18}{7} \int \frac{1}{\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2} - \frac{2(1+\sqrt{3})(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - \sqrt{3} + 2} d\left(-\frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{x}\right) + \frac{7}{21} \left(\frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{x} \right) \right) \right)$$

↓ 1083

$$2 \left(\frac{1}{21} \left(\frac{-\frac{(18+49\sqrt{3})(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - 43\sqrt{3} + 18}{7 \left(\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2} - \frac{2(1+\sqrt{3})(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - \sqrt{3} + 2 \right)} - \frac{36}{7} \int \frac{1}{28 - \frac{(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2}} dx \right) \right)$$

↓ 219

$$2 \left(\frac{1}{21} \left(\frac{18 \operatorname{arctanh} \left(\frac{\sqrt{3}-\sqrt{-x^2-2x+3}}{2\sqrt{7}x} \right)}{7\sqrt{7}} + \frac{-\frac{(18+49\sqrt{3})(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - 43\sqrt{3} + 18}{7 \left(\frac{\sqrt{3}(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2} - \frac{2(1+\sqrt{3})(\sqrt{3}-\sqrt{-x^2-2x+3})}{x} - \sqrt{3} + 2 \right)} \right) \right) - \frac{36}{7} \int \frac{1}{28 - \frac{(\sqrt{3}-\sqrt{-x^2-2x+3})^2}{x^2}} dx$$

input `Int[(x + Sqrt[3 - 2*x - x^2])^(-3), x]`

output `2*((-2*(9 - 5*Sqrt[3] + ((21 + 5*Sqrt[3])*(Sqrt[3] - Sqrt[3 - 2*x - x^2]))/x))/(21*(2 - Sqrt[3] - (2*(1 + Sqrt[3])*(Sqrt[3] - Sqrt[3 - 2*x - x^2]))/x + (Sqrt[3]*(Sqrt[3] - Sqrt[3 - 2*x - x^2])^2)/x^2) + ((18 - 43*Sqrt[3] - ((18 + 49*Sqrt[3])*(Sqrt[3] - Sqrt[3 - 2*x - x^2]))/x)/(7*(2 - Sqrt[3] - (2*(1 + Sqrt[3])*(Sqrt[3] - Sqrt[3 - 2*x - x^2]))/x + (Sqrt[3]*(Sqrt[3] - Sqrt[3 - 2*x - x^2])^2)/x^2)) + (18*ArcTanh[(Sqrt[3] - Sqrt[3 - 2*x - x^2])/(2*Sqrt[7]*x)]/(7*Sqrt[7])))/21)`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 1083 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`

rule 2191 `Int[(Pq_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := With[{Q = PolynomialQuotient[Pq, a + b*x + c*x^2, x], f = Coeff[PolynomialRemainder[Pq, a + b*x + c*x^2, x], x, 0], g = Coeff[PolynomialRemainder[Pq, a + b*x + c*x^2, x], x, 1]}, Simp[(b*f - 2*a*g + (2*c*f - b*g)*x)*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c))), x] + Simp[1/((p + 1)*(b^2 - 4*a*c)) Int[(a + b*x + c*x^2)^(p + 1)*ExpandToSum[(p + 1)*(b^2 - 4*a*c)*Q - (2*p + 3)*(2*c*f - b*g), x], x], x]] /; FreeQ[{a, b, c}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1]`

rule 7285 `Int[u_, x_Symbol] := With[{lst = FunctionOfSquareRootOfQuadratic[u, x]}, Simp[2 Subst[Int[lst[[1]], x], x, lst[[2]]], x] /; !FalseQ[lst] && EqQ[lst[[3]], 1] /; EulerIntegrandQ[u, x]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.14 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.41

method	result
trager	$\frac{(62x^3+100x^2-111x-36)x}{98(2x^2+2x-3)^2} - \frac{(34x^3+58x^2-83x-15)\sqrt{-x^2-2x+3}}{98(2x^2+2x-3)^2} - \frac{6\operatorname{RootOf}(_Z^2-7)\ln\left(\frac{\operatorname{RootOf}(_Z^2-7)^{x-3}\operatorname{RootOf}(_Z^2-7)}{\operatorname{RootOf}(_Z^2-7)}\right)}{343}$
default	Expression too large to display

input `int(1/(x+(-x^2-2*x+3)^(1/2))^3,x,method=_RETURNVERBOSE)`

output `1/98*(62*x^3+100*x^2-111*x-36)*x/(2*x^2+2*x-3)^2-1/98*(34*x^3+58*x^2-83*x-15)/(2*x^2+2*x-3)^2*(-x^2-2*x+3)^(1/2)-6/343*RootOf(_Z^2-7)*ln((RootOf(_Z^2-7)*x-3*RootOf(_Z^2-7)-7*(-x^2-2*x+3)^(1/2))/(RootOf(_Z^2-7)*x-3))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 223, normalized size of antiderivative = 0.72

$$\int \frac{1}{(x + \sqrt{3 - 2x - x^2})^3} dx = \frac{336x^3 - 6\sqrt{7}(4x^4 + 8x^3 - 8x^2 - 12x + 9) \log\left(\frac{x^4 + 44x^3 - \sqrt{7}(3x^3 + x^2 - 45x + 45)\sqrt{-x^2 - 2x + 3} + 26x^2 - 276x + 207}{4x^4 + 8x^3 - 8x^2 - 12x + 9}\right)}{1}$$

input `integrate(1/(x+(-x^2-2*x+3)^(1/2))^3,x, algorithm="fricas")`

output `-1/1372*(336*x^3 - 6*sqrt(7)*(4*x^4 + 8*x^3 - 8*x^2 - 12*x + 9)*log((x^4 + 44*x^3 - sqrt(7)*(3*x^3 + x^2 - 45*x + 45)*sqrt(-x^2 - 2*x + 3) + 26*x^2 - 276*x + 207)/(4*x^4 + 8*x^3 - 8*x^2 - 12*x + 9)) - 12*sqrt(7)*(4*x^4 + 8*x^3 - 8*x^2 - 12*x + 9)*log((2*x^2 + sqrt(7)*(2*x + 1) + 2*x + 4)/(2*x^2 + 2*x - 3)) - 182*x^2 + 14*(34*x^3 + 58*x^2 - 83*x - 15)*sqrt(-x^2 - 2*x + 3) - 2100*x + 1953)/(4*x^4 + 8*x^3 - 8*x^2 - 12*x + 9)`

Sympy [F]

$$\int \frac{1}{(x + \sqrt{3 - 2x - x^2})^3} dx = \int \frac{1}{(x + \sqrt{-x^2 - 2x + 3})^3} dx$$

input `integrate(1/(x+(-x**2-2*x+3)**(1/2))**3,x)`

output `Integral((x + sqrt(-x**2 - 2*x + 3))**(-3), x)`

Maxima [F]

$$\int \frac{1}{(x + \sqrt{3 - 2x - x^2})^3} dx = \int \frac{1}{(x + \sqrt{-x^2 - 2x + 3})^3} dx$$

input `integrate(1/(x+(-x^2-2*x+3)^(1/2))^3,x, algorithm="maxima")`

output `integrate((x + sqrt(-x^2 - 2*x + 3))^-3, x)`

Giac [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 452, normalized size of antiderivative = 1.45

$$\begin{aligned} & \int \frac{1}{(x + \sqrt{3 - 2x - x^2})^3} dx \\ &= -\frac{3}{343} \sqrt{7} \log \left(\frac{|4x - 2\sqrt{7} + 2|}{|4x + 2\sqrt{7} + 2|} \right) + \frac{3}{343} \sqrt{7} \log \left(\frac{\left| -2\sqrt{7} + \frac{6(\sqrt{-x^2-2x+3}-2)}{x+1} + 4 \right|}{\left| 2\sqrt{7} + \frac{6(\sqrt{-x^2-2x+3}-2)}{x+1} + 4 \right|} \right) \\ & - \frac{3}{343} \sqrt{7} \log \left(\frac{\left| -2\sqrt{7} + \frac{2(\sqrt{-x^2-2x+3}-2)}{x+1} - 4 \right|}{\left| 2\sqrt{7} + \frac{2(\sqrt{-x^2-2x+3}-2)}{x+1} - 4 \right|} \right) - \frac{48x^3 - 26x^2 - 300x + 279}{196(2x^2 + 2x - 3)^2} \\ & + \frac{4 \left(\frac{231(\sqrt{-x^2-2x+3}-2)}{x+1} + \frac{3286(\sqrt{-x^2-2x+3}-2)^2}{(x+1)^2} - \frac{4441(\sqrt{-x^2-2x+3}-2)^3}{(x+1)^3} - \frac{18906(\sqrt{-x^2-2x+3}-2)^4}{(x+1)^4} - \frac{12487(\sqrt{-x^2-2x+3}-2)^5}{(x+1)^5} \right)}{441 \left(\frac{8(\sqrt{-x^2-2x+3}-2)}{x+1} + \frac{26(\sqrt{-x^2-2x+3}-2)^2}{(x+1)^2} + \frac{8(\sqrt{-x^2-2x+3}-2)^3}{(x+1)^3} \right)} \end{aligned}$$

input `integrate(1/(x+(-x^2-2*x+3)^(1/2))^3,x, algorithm="giac")`

output

```
-3/343*sqrt(7)*log(abs(4*x - 2*sqrt(7) + 2)/abs(4*x + 2*sqrt(7) + 2)) + 3/
343*sqrt(7)*log(abs(-2*sqrt(7) + 6*(sqrt(-x^2 - 2*x + 3) - 2)/(x + 1) + 4)
/abs(2*sqrt(7) + 6*(sqrt(-x^2 - 2*x + 3) - 2)/(x + 1) + 4)) - 3/343*sqrt(7
)*log(abs(-2*sqrt(7) + 2*(sqrt(-x^2 - 2*x + 3) - 2)/(x + 1) - 4)/abs(2*sq
rt(7) + 2*(sqrt(-x^2 - 2*x + 3) - 2)/(x + 1) - 4)) - 1/196*(48*x^3 - 26*x^2
- 300*x + 279)/(2*x^2 + 2*x - 3)^2 + 4/441*(231*(sqrt(-x^2 - 2*x + 3) - 2
)/(x + 1) + 3286*(sqrt(-x^2 - 2*x + 3) - 2)^2/(x + 1)^2 - 4441*(sqrt(-x^2
- 2*x + 3) - 2)^3/(x + 1)^3 - 18906*(sqrt(-x^2 - 2*x + 3) - 2)^4/(x + 1)^4
- 12487*(sqrt(-x^2 - 2*x + 3) - 2)^5/(x + 1)^5 + 946*(sqrt(-x^2 - 2*x + 3
- 2)^6/(x + 1)^6 + 1977*(sqrt(-x^2 - 2*x + 3) - 2)^7/(x + 1)^7 - 414)/(8
*(sqrt(-x^2 - 2*x + 3) - 2)/(x + 1) + 26*(sqrt(-x^2 - 2*x + 3) - 2)^2/(x +
1)^2 + 8*(sqrt(-x^2 - 2*x + 3) - 2)^3/(x + 1)^3 - 3*(sqrt(-x^2 - 2*x + 3)
- 2)^4/(x + 1)^4 - 3)^2
```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(x + \sqrt{3 - 2x - x^2})^3} dx = \int \frac{1}{(x + \sqrt{-x^2 - 2x + 3})^3} dx$$

input

```
int(1/(x + (3 - x^2 - 2*x)^(1/2))^3,x)
```

output

```
int(1/(x + (3 - x^2 - 2*x)^(1/2))^3, x)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 389, normalized size of antiderivative = 1.25

$$\int \frac{1}{(x + \sqrt{3 - 2x - x^2})^3} dx = \text{Too large to display}$$

input

```
int(1/(x+(-x^2-2*x+3)^(1/2))^3,x)
```

output

```
(2*( - 81*sqrt(7)*log( - sqrt(7) + 3*tan(asin((x + 1)/2)/2) - 2)*tan(asin(
(x + 1)/2)/2)**4 + 216*sqrt(7)*log( - sqrt(7) + 3*tan(asin((x + 1)/2)/2) -
2)*tan(asin((x + 1)/2)/2)**3 - 90*sqrt(7)*log( - sqrt(7) + 3*tan(asin((x
+ 1)/2)/2) - 2)*tan(asin((x + 1)/2)/2)**2 - 72*sqrt(7)*log( - sqrt(7) + 3*
tan(asin((x + 1)/2)/2) - 2)*tan(asin((x + 1)/2)/2) - 9*sqrt(7)*log( - sqrt
(7) + 3*tan(asin((x + 1)/2)/2) - 2) + 81*sqrt(7)*log(sqrt(7) + 3*tan(asin(
(x + 1)/2)/2) - 2)*tan(asin((x + 1)/2)/2)**4 - 216*sqrt(7)*log(sqrt(7) + 3
*tan(asin((x + 1)/2)/2) - 2)*tan(asin((x + 1)/2)/2)**3 + 90*sqrt(7)*log(sq
rt(7) + 3*tan(asin((x + 1)/2)/2) - 2)*tan(asin((x + 1)/2)/2)**2 + 72*sqrt(
7)*log(sqrt(7) + 3*tan(asin((x + 1)/2)/2) - 2)*tan(asin((x + 1)/2)/2) + 9*
sqrt(7)*log(sqrt(7) + 3*tan(asin((x + 1)/2)/2) - 2) - 399*tan(asin((x + 1)
/2)/2)**4 + 770*tan(asin((x + 1)/2)/2)**2 - 140*tan(asin((x + 1)/2)/2) - 2
59))/(1029*(9*tan(asin((x + 1)/2)/2)**4 - 24*tan(asin((x + 1)/2)/2)**3 + 1
0*tan(asin((x + 1)/2)/2)**2 + 8*tan(asin((x + 1)/2)/2) + 1))
```


3.25 $\int \left(d + ex + f\sqrt{a + bx + cx^2} \right)^2 dx$

Optimal result	224
Mathematica [A] (verified)	225
Rubi [A] (verified)	225
Maple [A] (verified)	226
Fricas [A] (verification not implemented)	227
Sympy [A] (verification not implemented)	228
Maxima [F(-2)]	229
Giac [A] (verification not implemented)	229
Mupad [B] (verification not implemented)	230
Reduce [B] (verification not implemented)	230

Optimal result

Integrand size = 23, antiderivative size = 158

$$\begin{aligned} & \int \left(d + ex + f\sqrt{a + bx + cx^2} \right)^2 dx \\ &= (d^2 + af^2)x + \frac{1}{2}(2de + bf^2)x^2 + \frac{1}{3}(e^2 + cf^2)x^3 \\ &+ \frac{2ef(a + bx + cx^2)^{3/2}}{3c} - \frac{(-2cd + be)f(b + 2cx)\sqrt{a + x(b + cx)}}{4c^2} \\ &+ \frac{(b^2 - 4ac)(-2cd + be)f \operatorname{arctanh}\left(\frac{b + 2cx}{2\sqrt{c}\sqrt{a + x(b + cx)}}\right)}{8c^{5/2}} \end{aligned}$$

output

```
(a*f^2+d^2)*x+1/2*(b*f^2+2*d*e)*x^2+1/3*(c*f^2+e^2)*x^3+2/3*e*f*(c*x^2+b*x+a)^(3/2)/c-1/4*(b*e-2*c*d)*f*(2*c*x+b)*(a+x*(c*x+b))^(1/2)/c^2+1/8*(-4*a*c+b^2)*(b*e-2*c*d)*f*arctanh(1/2*(2*c*x+b)/c^(1/2)/(a+x*(c*x+b))^(1/2))/c^(5/2)
```

Mathematica [A] (verified)

Time = 1.93 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.99

$$\int (d + ex + f\sqrt{a + bx + cx^2})^2 dx$$

$$= \frac{1}{12} \left(2x(6d^2 + 6af^2 + 6dex + x(3bf^2 + 2(e^2 + cf^2)x)) \right.$$

$$+ \frac{f\sqrt{a + x(b + cx)}(-3b^2e + 2bc(3d + ex) + 4c(2ae + cx(3d + 2ex)))}{c^2}$$

$$\left. + \frac{3(b^2 - 4ac)(-2cd + be)f \operatorname{arctanh}\left(\frac{\sqrt{cx}}{-\sqrt{a + \sqrt{a + x(b + cx)}}}\right)}{c^{5/2}} \right)$$

input

```
Integrate[(d + e*x + f*Sqrt[a + b*x + c*x^2])^2,x]
```

output

```
(2*x*(6*d^2 + 6*a*f^2 + 6*d*e*x + x*(3*b*f^2 + 2*(e^2 + c*f^2)*x)) + (f*Sqrt[a + x*(b + c*x)]*(-3*b^2*e + 2*b*c*(3*d + e*x) + 4*c*(2*a*e + c*x*(3*d + 2*e*x))))/c^2 + (3*(b^2 - 4*a*c)*(-2*c*d + b*e)*f*ArcTanh[(Sqrt[c]*x)/(-Sqrt[a] + Sqrt[a + x*(b + c*x)])])/c^(5/2))/12
```

Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.43, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (f\sqrt{a + bx + cx^2} + d + ex)^2 dx$$

↓ 7293

$$\int \left(2df\sqrt{a + bx + cx^2} + 2efx\sqrt{a + bx + cx^2} + d^2 \left(\frac{af^2}{d^2} + 1 \right) + 2dex \left(\frac{bf^2}{2de} + 1 \right) + e^2x^2 \left(\frac{cf^2}{e^2} + 1 \right) \right) dx$$

$$\begin{aligned}
 & \downarrow \text{2009} \\
 & -\frac{df(b^2 - 4ac) \operatorname{arctanh}\left(\frac{b+2cx}{2\sqrt{c}\sqrt{a+bx+cx^2}}\right)}{4c^{3/2}} + \frac{bef(b^2 - 4ac) \operatorname{arctanh}\left(\frac{b+2cx}{2\sqrt{c}\sqrt{a+bx+cx^2}}\right)}{8c^{5/2}} - \\
 & \frac{bef(b+2cx)\sqrt{a+bx+cx^2}}{4c^2} + \frac{df(b+2cx)\sqrt{a+bx+cx^2}}{2c} + \frac{2ef(a+bx+cx^2)^{3/2}}{3c} + \\
 & x(af^2 + d^2) + \frac{1}{2}x^2(bf^2 + 2de) + \frac{1}{3}x^3(cf^2 + e^2)
 \end{aligned}$$

input `Int[(d + e*x + f*Sqrt[a + b*x + c*x^2])^2,x]`

output `(d^2 + a*f^2)*x + ((2*d*e + b*f^2)*x^2)/2 + ((e^2 + c*f^2)*x^3)/3 + (d*f*(b + 2*c*x)*Sqrt[a + b*x + c*x^2])/(2*c) - (b*e*f*(b + 2*c*x)*Sqrt[a + b*x + c*x^2])/(4*c^2) + (2*e*f*(a + b*x + c*x^2)^(3/2))/(3*c) - ((b^2 - 4*a*c)*d*f*ArcTanh[(b + 2*c*x)/(2*Sqrt[c]*Sqrt[a + b*x + c*x^2])])/(4*c^(3/2)) + (b*(b^2 - 4*a*c)*e*f*ArcTanh[(b + 2*c*x)/(2*Sqrt[c]*Sqrt[a + b*x + c*x^2])])/(8*c^(5/2))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

Maple [A] (verified)

Time = 1.68 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.23

method	result
default	$ f^2\left(\frac{1}{3}cx^3 + \frac{1}{2}bx^2 + xa\right) + 2f\left(d\left(\frac{(2cx+b)\sqrt{cx^2+bx+a}}{4c} + \frac{(4ac-b^2)\ln\left(\frac{\frac{b}{2}+cx}{\sqrt{c}}+\sqrt{cx^2+bx+a}\right)}{8c^{\frac{3}{2}}}\right)\right) + e\left(\frac{cx^2+bx+a}{3c}\right) $

input `int((d+e*x+f*(c*x^2+b*x+a)^(1/2))^2,x,method=_RETURNVERBOSE)`

output `f^2*(1/3*c*x^3+1/2*b*x^2+x*a)+2*f*(d*(1/4*(2*c*x+b)/c*(c*x^2+b*x+a)^(1/2)+1/8*(4*a*c-b^2)/c^(3/2)*ln((1/2*b+c*x)/c^(1/2)+(c*x^2+b*x+a)^(1/2)))+e*(1/3*(c*x^2+b*x+a)^(3/2)/c-1/2*b/c*(1/4*(2*c*x+b)/c*(c*x^2+b*x+a)^(1/2)+1/8*(4*a*c-b^2)/c^(3/2)*ln((1/2*b+c*x)/c^(1/2)+(c*x^2+b*x+a)^(1/2))))+1/3*(e*x+d)^3/e`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 429, normalized size of antiderivative = 2.72

$$\int \left(d + ex + f\sqrt{a + bx + cx^2} \right)^2 dx$$

$$= \left[\frac{16(c^4 f^2 + c^3 e^2)x^3 + 3(2(b^2 c - 4ac^2)d - (b^3 - 4abc)e)\sqrt{c}f \log(-8c^2 x^2 - 8bcx - b^2 + 4\sqrt{cx^2 + bx + a})}{1} \right]$$

input `integrate((d+e*x+f*(c*x^2+b*x+a)^(1/2))^2,x, algorithm="fricas")`

output `[1/48*(16*(c^4*f^2 + c^3*e^2)*x^3 + 3*(2*(b^2*c - 4*a*c^2)*d - (b^3 - 4*a*b*c)*e)*sqrt(c)*f*log(-8*c^2*x^2 - 8*b*c*x - b^2 + 4*sqrt(c*x^2 + b*x + a)*(2*c*x + b)*sqrt(c) - 4*a*c) + 24*(b*c^3*f^2 + 2*c^3*d*e)*x^2 + 48*(a*c^3*f^2 + c^3*d^2)*x + 4*(8*c^3*e*f*x^2 + 2*(6*c^3*d + b*c^2*e)*f*x + (6*b*c^2*d - (3*b^2*c - 8*a*c^2)*e)*f)*sqrt(c*x^2 + b*x + a))/c^3, 1/24*(8*(c^4*f^2 + c^3*e^2)*x^3 + 3*(2*(b^2*c - 4*a*c^2)*d - (b^3 - 4*a*b*c)*e)*sqrt(-c)*f*arctan(1/2*sqrt(c*x^2 + b*x + a)*(2*c*x + b)*sqrt(-c)/(c^2*x^2 + b*c*x + a*c)) + 12*(b*c^3*f^2 + 2*c^3*d*e)*x^2 + 24*(a*c^3*f^2 + c^3*d^2)*x + 2*(8*c^3*e*f*x^2 + 2*(6*c^3*d + b*c^2*e)*f*x + (6*b*c^2*d - (3*b^2*c - 8*a*c^2)*e)*f)*sqrt(c*x^2 + b*x + a))/c^3]`

Sympy [A] (verification not implemented)

Time = 0.70 (sec) , antiderivative size = 345, normalized size of antiderivative = 2.18

$$\int \left(d + ex + f\sqrt{a + bx + cx^2} \right)^2 dx = af^2x + \frac{bf^2x^2}{2} + \frac{cf^2x^3}{3} + d^2x + dex^2$$

$$+ 2df \left\{ \begin{array}{l} \left(\frac{a}{2} - \frac{b^2}{8c} \right) \left(\begin{array}{l} \frac{\log(b + 2\sqrt{c}\sqrt{a + bx + cx^2} + 2cx)}{\sqrt{c}} \\ \frac{(\frac{b}{2c} + x) \log(\frac{b}{2c} + x)}{\sqrt{c}(\frac{b}{2c} + x)^2} \end{array} \right) \text{ for } a - \frac{b^2}{4c} \neq 0 \\ \frac{2(a + bx)^{\frac{3}{2}}}{3b} \\ \sqrt{ax} \end{array} \right. + \left(\frac{b}{4c} + \frac{x}{2} \right) \sqrt{a + bx + cx^2} \quad \text{for } c \neq 0$$

$$+ \frac{e^2x^3}{3}$$

$$+ 2ef \left\{ \begin{array}{l} \left(-\frac{ab}{12c} - \frac{b(\frac{a}{3} - \frac{b^2}{8c})}{2c} \right) \left(\begin{array}{l} \frac{\log(b + 2\sqrt{c}\sqrt{a + bx + cx^2} + 2cx)}{\sqrt{c}} \\ \frac{(\frac{b}{2c} + x) \log(\frac{b}{2c} + x)}{\sqrt{c}(\frac{b}{2c} + x)^2} \end{array} \right) \text{ for } a - \frac{b^2}{4c} \neq 0 \\ \frac{2\left(-\frac{a(a + bx)^{\frac{3}{2}}}{3} + \frac{(a + bx)^{\frac{5}{2}}}{5}\right)}{b^2} \\ \frac{\sqrt{ax^2}}{2} \end{array} \right. + \sqrt{a + bx + cx^2} \left(\frac{bx}{12c} + \frac{x^2}{3} \right) \quad \text{for } b \neq 0$$

$$\text{otherwise}$$

```
input integrate((d+e*x+f*(c*x**2+b*x+a)**(1/2))**2,x)
```

```
output a*f**2*x + b*f**2*x**2/2 + c*f**2*x**3/3 + d**2*x + d*e*x**2 + 2*d*f*Piecewise(((a/2 - b**2/(8*c))*Piecewise((log(b + 2*sqrt(c)*sqrt(a + b*x + c*x**2) + 2*c*x)/sqrt(c), Ne(a - b**2/(4*c), 0)), ((b/(2*c) + x)*log(b/(2*c) + x)/sqrt(c*(b/(2*c) + x)**2), True)) + (b/(4*c) + x/2)*sqrt(a + b*x + c*x**2), Ne(c, 0)), (2*(a + b*x)**(3/2)/(3*b), Ne(b, 0)), (sqrt(a)*x, True)) + e**2*x**3/3 + 2*e*f*Piecewise(((a*b/(12*c) - b*(a/3 - b**2/(8*c)))/(2*c))*Piecewise((log(b + 2*sqrt(c)*sqrt(a + b*x + c*x**2) + 2*c*x)/sqrt(c), Ne(a - b**2/(4*c), 0)), ((b/(2*c) + x)*log(b/(2*c) + x)/sqrt(c*(b/(2*c) + x)**2), True)) + sqrt(a + b*x + c*x**2)*(b*x/(12*c) + x**2/3 + (a/3 - b**2/(8*c))/c), Ne(c, 0)), (2*(-a*(a + b*x)**(3/2)/3 + (a + b*x)**(5/2)/5)/b**2, Ne(b, 0)), (sqrt(a)*x**2/2, True))
```

Maxima [F(-2)]

Exception generated.

$$\int (d + ex + f\sqrt{a + bx + cx^2})^2 dx = \text{Exception raised: ValueError}$$

input `integrate((d+e*x+f*(c*x^2+b*x+a)^(1/2))^2,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a*c-b^2>0)', see `assume?` for more deta

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.10

$$\begin{aligned} & \int (d + ex + f\sqrt{a + bx + cx^2})^2 dx \\ &= \frac{1}{3}cf^2x^3 + \frac{1}{2}bf^2x^2 + \frac{1}{3}e^2x^3 + af^2x + dex^2 + d^2x \\ &+ \frac{1}{12}\sqrt{cx^2 + bx + a} \left(2 \left(4efx + \frac{6c^2df + bcef}{c^2} \right) x + \frac{6bcdf - 3b^2ef + 8acef}{c^2} \right) \\ &+ \frac{(2b^2cdf - 8ac^2df - b^3ef + 4abcef) \log(|2(\sqrt{cx} - \sqrt{cx^2 + bx + a})\sqrt{c} + b|)}{8c^{\frac{5}{2}}} \end{aligned}$$

input `integrate((d+e*x+f*(c*x^2+b*x+a)^(1/2))^2,x, algorithm="giac")`

output `1/3*c*f^2*x^3 + 1/2*b*f^2*x^2 + 1/3*e^2*x^3 + a*f^2*x + d*e*x^2 + d^2*x + 1/12*sqrt(c*x^2 + b*x + a)*(2*(4*e*f*x + (6*c^2*d*f + b*c*e*f)/c^2)*x + (6*b*c*d*f - 3*b^2*e*f + 8*a*c*e*f)/c^2) + 1/8*(2*b^2*c*d*f - 8*a*c^2*d*f - b^3*e*f + 4*a*b*c*e*f)*log(abs(2*(sqrt(c)*x - sqrt(c*x^2 + b*x + a))*sqrt(c) + b))/c^(5/2)`

Mupad [B] (verification not implemented)

Time = 21.25 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.20

$$\int (d + ex + f\sqrt{a + bx + cx^2})^2 dx$$

$$= x^3 \left(\frac{e^2}{3} + \frac{cf^2}{3} \right) + x^2 \left(\frac{bf^2}{2} + de \right) + x(d^2 + af^2) + 2df \left(\frac{x}{2} + \frac{b}{4c} \right) \sqrt{cx^2 + bx + a}$$

$$+ \frac{ef(-3b^2 + 2cxb + 8c(cx^2 + a)) \sqrt{cx^2 + bx + a}}{12c^2}$$

$$+ \frac{df \ln \left(\frac{\frac{b}{2} + cx}{\sqrt{c}} + \sqrt{cx^2 + bx + a} \right) \left(ac - \frac{b^2}{4} \right)}{c^{3/2}}$$

$$+ \frac{ef \ln \left(\frac{b+2cx}{\sqrt{c}} + 2\sqrt{cx^2 + bx + a} \right) (b^3 - 4abc)}{8c^{5/2}}$$

input `int((d + e*x + f*(a + b*x + c*x^2)^(1/2))^2,x)`output `x^3*((c*f^2)/3 + e^2/3) + x^2*(d*e + (b*f^2)/2) + x*(a*f^2 + d^2) + 2*d*f*(x/2 + b/(4*c))*(a + b*x + c*x^2)^(1/2) + (e*f*(8*c*(a + c*x^2) - 3*b^2 + 2*b*c*x)*(a + b*x + c*x^2)^(1/2))/(12*c^2) + (d*f*log((b/2 + c*x)/c^(1/2) + (a + b*x + c*x^2)^(1/2))*(a*c - b^2/4))/c^(3/2) + (e*f*log((b + 2*c*x)/c^(1/2) + 2*(a + b*x + c*x^2)^(1/2))*(b^3 - 4*a*b*c))/(8*c^(5/2))`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 368, normalized size of antiderivative = 2.33

$$\int (d + ex + f\sqrt{a + bx + cx^2})^2 dx$$

$$= \frac{16\sqrt{cx^2 + bx + a}ac^2ef - 6\sqrt{cx^2 + bx + a}b^2cef + 12\sqrt{cx^2 + bx + a}bc^2df + 4\sqrt{cx^2 + bx + a}bc^2efx - \dots}{\dots}$$

input `int((d+e*x+f*(c*x^2+b*x+a)^(1/2))^2,x)`

output

```
(16*sqrt(a + b*x + c*x**2)*a*c**2*e*f - 6*sqrt(a + b*x + c*x**2)*b**2*c*e*
f + 12*sqrt(a + b*x + c*x**2)*b*c**2*d*f + 4*sqrt(a + b*x + c*x**2)*b*c**2
*e*f*x + 24*sqrt(a + b*x + c*x**2)*c**3*d*f*x + 16*sqrt(a + b*x + c*x**2)*
c**3*e*f*x**2 - 12*sqrt(c)*log((2*sqrt(c)*sqrt(a + b*x + c*x**2) + b + 2*c
*x)/sqrt(4*a*c - b**2))*a*b*c*e*f + 24*sqrt(c)*log((2*sqrt(c)*sqrt(a + b*x
+ c*x**2) + b + 2*c*x)/sqrt(4*a*c - b**2))*a*c**2*d*f + 3*sqrt(c)*log((2*
sqrt(c)*sqrt(a + b*x + c*x**2) + b + 2*c*x)/sqrt(4*a*c - b**2))*b**3*e*f -
6*sqrt(c)*log((2*sqrt(c)*sqrt(a + b*x + c*x**2) + b + 2*c*x)/sqrt(4*a*c -
b**2))*b**2*c*d*f + 24*a*c**3*f**2*x + 12*b*c**3*f**2*x**2 + 8*c**4*f**2*
x**3 + 24*c**3*d**2*x + 24*c**3*d*e*x**2 + 8*c**3*e**2*x**3)/(24*c**3)
```


3.26 $\int \left(d + ex + f\sqrt{a + bx + cx^2} \right) dx$

Optimal result	232
Mathematica [A] (verified)	232
Rubi [A] (verified)	233
Maple [A] (verified)	234
Fricas [A] (verification not implemented)	234
Sympy [A] (verification not implemented)	235
Maxima [F(-2)]	235
Giac [A] (verification not implemented)	236
Mupad [B] (verification not implemented)	236
Reduce [B] (verification not implemented)	237

Optimal result

Integrand size = 21, antiderivative size = 88

$$\int \left(d + ex + f\sqrt{a + bx + cx^2} \right) dx = dx + \frac{ex^2}{2} + \frac{f(b + 2cx)\sqrt{a + bx + cx^2}}{4c} - \frac{(b^2 - 4ac) \operatorname{farctanh}\left(\frac{b+2cx}{2\sqrt{c}\sqrt{a+bx+cx^2}}\right)}{8c^{3/2}}$$

output

```
d*x+1/2*e*x^2+1/4*f*(2*c*x+b)*(c*x^2+b*x+a)^(1/2)/c-1/8*(-4*a*c+b^2)*f*arc
tanh(1/2*(2*c*x+b)/c^(1/2)/(c*x^2+b*x+a)^(1/2))/c^(3/2)
```

Mathematica [A] (verified)

Time = 0.57 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.97

$$\int \left(d + ex + f\sqrt{a + bx + cx^2} \right) dx = \frac{1}{4} \left(4dx + 2ex^2 + \frac{f(b + 2cx)\sqrt{a + x(b + cx)}}{c} + \frac{(b^2 - 4ac) \operatorname{farctanh}\left(\frac{\sqrt{cx}}{\sqrt{a - \sqrt{a + x(b + cx)}}}\right)}{c^{3/2}} \right)$$

input `Integrate[d + e*x + f*Sqrt[a + b*x + c*x^2], x]`

output `(4*d*x + 2*e*x^2 + (f*(b + 2*c*x)*Sqrt[a + x*(b + c*x)])/c + ((b^2 - 4*a*c)*f*ArcTanh[(Sqrt[c]*x)/(Sqrt[a] - Sqrt[a + x*(b + c*x)])])/c^(3/2))/4`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(f \sqrt{a + bx + cx^2} + d + ex \right) dx$$

↓ 2009

$$-\frac{f(b^2 - 4ac) \operatorname{arctanh}\left(\frac{b+2cx}{2\sqrt{c}\sqrt{a+bx+cx^2}}\right)}{8c^{3/2}} + \frac{f(b + 2cx)\sqrt{a + bx + cx^2}}{4c} + dx + \frac{ex^2}{2}$$

input `Int[d + e*x + f*Sqrt[a + b*x + c*x^2], x]`

output `d*x + (e*x^2)/2 + (f*(b + 2*c*x)*Sqrt[a + b*x + c*x^2])/(4*c) - ((b^2 - 4*a*c)*f*ArcTanh[(b + 2*c*x)/(2*Sqrt[c]*Sqrt[a + b*x + c*x^2])])/(8*c^(3/2))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.88

method	result	size
default	$dx + \frac{ex^2}{2} + f \left(\frac{(2cx+b)\sqrt{cx^2+bx+a}}{4c} + \frac{(4ac-b^2) \ln \left(\frac{\frac{b}{\sqrt{c}} + \sqrt{cx^2+bx+a}}{\sqrt{c}} \right)}{8c^{\frac{3}{2}}} \right)$	77
parts	$dx + \frac{ex^2}{2} + f \left(\frac{(2cx+b)\sqrt{cx^2+bx+a}}{4c} + \frac{(4ac-b^2) \ln \left(\frac{\frac{b}{\sqrt{c}} + \sqrt{cx^2+bx+a}}{\sqrt{c}} \right)}{8c^{\frac{3}{2}}} \right)$	77

input `int(d+e*x+f*(c*x^2+b*x+a)^(1/2),x,method=_RETURNVERBOSE)`

output `d*x+1/2*e*x^2+f*(1/4*(2*c*x+b)/c*(c*x^2+b*x+a)^(1/2)+1/8*(4*a*c-b^2)/c^(3/2)*ln((1/2*b+c*x)/c^(1/2)+(c*x^2+b*x+a)^(1/2)))`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 216, normalized size of antiderivative = 2.45

$$\int (d + ex + f\sqrt{a + bx + cx^2}) dx$$

$$= \frac{8c^2ex^2 + 16c^2dx - (b^2 - 4ac)\sqrt{c}f \log(-8c^2x^2 - 8bcx - b^2 - 4\sqrt{cx^2 + bx + a}(2cx + b)\sqrt{c} - 4ac) + 4c^2 \arctan\left(\frac{2cx + b}{\sqrt{cx^2 + bx + a}}\right)}{16c^2}$$

input `integrate(d+e*x+f*(c*x^2+b*x+a)^(1/2),x, algorithm="fricas")`

output `[1/16*(8*c^2*e*x^2 + 16*c^2*d*x - (b^2 - 4*a*c)*sqrt(c)*f*log(-8*c^2*x^2 - 8*b*c*x - b^2 - 4*sqrt(c*x^2 + b*x + a)*(2*c*x + b)*sqrt(c) - 4*a*c) + 4*(2*c^2*f*x + b*c*f)*sqrt(c*x^2 + b*x + a))/c^2, 1/8*(4*c^2*e*x^2 + 8*c^2*d*x + (b^2 - 4*a*c)*sqrt(-c)*f*arctan(1/2*sqrt(c*x^2 + b*x + a)*(2*c*x + b)*sqrt(-c)/(c^2*x^2 + b*c*x + a*c)) + 2*(2*c^2*f*x + b*c*f)*sqrt(c*x^2 + b*x + a))/c^2]`

Sympy [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.49

$$\int (d + ex + f\sqrt{a + bx + cx^2}) dx = dx + \frac{ex^2}{2} + f \left(\begin{cases} \left(\frac{a}{2} - \frac{b^2}{8c} \right) \left(\begin{cases} \frac{\log(b + 2\sqrt{c}\sqrt{a + bx + cx^2} + 2cx)}{\sqrt{c}} & \text{for } a - \frac{b^2}{4c} \neq 0 \\ \frac{(\frac{b}{2c} + x) \log(\frac{b}{2c} + x)}{\sqrt{c(\frac{b}{2c} + x)^2}} & \text{otherwise} \end{cases} \right) + \left(\frac{b}{4c} + \frac{x}{2} \right) \sqrt{a + bx + cx^2} & \text{for } c \neq 0 \\ \frac{2(a + bx)^{\frac{3}{2}}}{3b} & \text{for } b \neq 0 \\ \sqrt{ax} & \text{otherwise} \end{cases} \right)$$

input `integrate(d+e*x+f*(c*x**2+b*x+a)**(1/2),x)`output `d*x + e*x**2/2 + f*Piecewise(((a/2 - b**2/(8*c))*Piecewise((log(b + 2*sqrt(c)*sqrt(a + b*x + c*x**2) + 2*c*x)/sqrt(c), Ne(a - b**2/(4*c), 0)), ((b/(2*c) + x)*log(b/(2*c) + x)/sqrt(c*(b/(2*c) + x)**2), True)) + (b/(4*c) + x/2)*sqrt(a + b*x + c*x**2), Ne(c, 0)), (2*(a + b*x)**(3/2)/(3*b), Ne(b, 0)), (sqrt(a)*x, True))`**Maxima [F(-2)]**

Exception generated.

$$\int (d + ex + f\sqrt{a + bx + cx^2}) dx = \text{Exception raised: ValueError}$$

input `integrate(d+e*x+f*(c*x^2+b*x+a)^(1/2),x, algorithm="maxima")`output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a*c-b^2>0)', see `assume?` for more deta`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.89

$$\int (d + ex + f\sqrt{a + bx + cx^2}) dx = \frac{1}{2} ex^2 + \frac{1}{8} \left(2\sqrt{cx^2 + bx + a} \left(2x + \frac{b}{c} \right) + \frac{(b^2 - 4ac) \log(|2(\sqrt{cx} - \sqrt{cx^2 + bx + a})\sqrt{c} + b|)}{c^{\frac{3}{2}}} \right) f + dx$$

input `integrate(d+e*x+f*(c*x^2+b*x+a)^(1/2),x, algorithm="giac")`

output `1/2*e*x^2 + 1/8*(2*sqrt(c*x^2 + b*x + a)*(2*x + b/c) + (b^2 - 4*a*c)*log(abs(2*(sqrt(c)*x - sqrt(c*x^2 + b*x + a))*sqrt(c) + b))/c^(3/2))*f + d*x`

Mupad [B] (verification not implemented)

Time = 20.75 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.84

$$\int (d + ex + f\sqrt{a + bx + cx^2}) dx = dx + \frac{ex^2}{2} + f \left(\frac{x}{2} + \frac{b}{4c} \right) \sqrt{cx^2 + bx + a} + \frac{f \ln \left(\frac{\frac{b}{2} + cx}{\sqrt{c}} + \sqrt{cx^2 + bx + a} \right) \left(ac - \frac{b^2}{4} \right)}{2c^{3/2}}$$

input `int(d + e*x + f*(a + b*x + c*x^2)^(1/2),x)`

output `d*x + (e*x^2)/2 + f*(x/2 + b/(4*c))*(a + b*x + c*x^2)^(1/2) + (f*log((b/2 + c*x)/c^(1/2) + (a + b*x + c*x^2)^(1/2))*(a*c - b^2/4))/(2*c^(3/2))`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.62

$$\int (d + ex + f\sqrt{a + bx + cx^2}) dx$$

$$= \frac{2\sqrt{cx^2 + bx + a}bcf + 4\sqrt{cx^2 + bx + a}c^2fx + 4\sqrt{c}\log\left(\frac{2\sqrt{c}\sqrt{cx^2+bx+a}+b+2cx}{\sqrt{4ac-b^2}}\right)acf - \sqrt{c}\log\left(\frac{2\sqrt{c}\sqrt{cx^2+bx+a}}{\sqrt{4ac-b^2}}\right)}{8c^2}$$

input

```
int(d+e*x+f*(c*x^2+b*x+a)^(1/2),x)
```

output

```
(2*sqrt(a + b*x + c*x**2)*b*c*f + 4*sqrt(a + b*x + c*x**2)*c**2*f*x + 4*sqrt(c)*log((2*sqrt(c)*sqrt(a + b*x + c*x**2) + b + 2*c*x)/sqrt(4*a*c - b**2)))*a*c*f - sqrt(c)*log((2*sqrt(c)*sqrt(a + b*x + c*x**2) + b + 2*c*x)/sqrt(4*a*c - b**2))*b**2*f + 8*c**2*d*x + 4*c**2*e*x**2)/(8*c**2)
```

3.27 $\int \frac{1}{d+ex+f\sqrt{a+bx+cx^2}} dx$

Optimal result	238
Mathematica [A] (verified)	239
Rubi [B] (verified)	240
Maple [B] (warning: unable to verify)	241
Fricas [F(-1)]	242
Sympy [F]	243
Maxima [F]	243
Giac [F(-2)]	243
Mupad [F(-1)]	244
Reduce [F]	244

Optimal result

Integrand size = 23, antiderivative size = 361

$$\int \frac{1}{d+ex+f\sqrt{a+bx+cx^2}} dx$$

$$= \frac{2\sqrt{c}f \operatorname{arctanh}\left(\frac{\sqrt{a}-\sqrt{a+bx+cx^2}}{\sqrt{cx}}\right)}{e^2 - cf^2}$$

$$+ \frac{2(2cd - be) f \operatorname{arctanh}\left(\frac{2af+bf x+2d\sqrt{a+x(b+cx)}-2\sqrt{a}(d+ex+f\sqrt{a+x(b+cx)})}{\sqrt{-4bde+4ae^2+b^2f^2+4c(d^2-af^2)}x}\right)}{(e^2 - cf^2) \sqrt{-4bde + 4ae^2 + b^2f^2 + 4c(d^2 - af^2)}}$$

$$- \frac{e \log\left(c - \frac{(\sqrt{a}-\sqrt{a+bx+cx^2})^2}{x^2}\right)}{e^2 - cf^2}$$

$$+ \frac{e \log\left(be - c(d + \sqrt{a}f) + \frac{(2\sqrt{a}e-bf)(\sqrt{a}-\sqrt{a+bx+cx^2})}{x} + \frac{(d-\sqrt{a}f)(\sqrt{a}-\sqrt{a+bx+cx^2})^2}{x^2} \right)}{e^2 - cf^2}$$

output

$$2*c^{(1/2)}*f*\operatorname{arctanh}((a^{(1/2)}-(c*x^2+b*x+a)^{(1/2)})/c^{(1/2)}/x)/(-c*f^2+e^2)+2*(-b*e+2*c*d)*f*\operatorname{arctanh}((2*a*f+b*f*x+2*d*(a+x*(c*x+b))^{(1/2)}-2*a^{(1/2)}*(d+e*x+f*(a+x*(c*x+b))^{(1/2)}))/(-4*b*d*e+4*a*e^2+b^2*f^2+4*c*(-a*f^2+d^2))^{(1/2)}/x)/(-c*f^2+e^2)/(-4*b*d*e+4*a*e^2+b^2*f^2+4*c*(-a*f^2+d^2))^{(1/2)}-e*\ln(c-(a^{(1/2)}-(c*x^2+b*x+a)^{(1/2)})^2/x^2)/(-c*f^2+e^2)+e*\ln(b*e-c*(d+a^{(1/2)})*f)+(2*a^{(1/2)}*e-b*f)*(a^{(1/2)}-(c*x^2+b*x+a)^{(1/2)})/x+(d-a^{(1/2)}*f)*(a^{(1/2)}-(c*x^2+b*x+a)^{(1/2)})^2/x^2)/(-c*f^2+e^2)$$

Mathematica [A] (verified)

Time = 3.17 (sec) , antiderivative size = 323, normalized size of antiderivative = 0.89

$$\int \frac{1}{d+ex+f\sqrt{a+bx+cx^2}} dx$$

$$= \frac{2(2cd-be)f \arctan\left(\frac{\sqrt{-4cd^2+4bde-4ae^2-b^2f^2+4acf^2}x}{2af+bf x+2d\sqrt{a+bx+cx^2}-2\sqrt{a}(d+ex+f\sqrt{a+bx+cx^2})}\right)}{\sqrt{-4cd^2+4bde-4ae^2-b^2f^2+4acf^2}} + 2\sqrt{c}f \operatorname{arctanh}\left(\frac{\sqrt{cx}}{\sqrt{a}-\sqrt{a+bx+cx^2}}\right) + e\left(-\log\left(-2\right)\right)$$

input

Integrate[(d + e*x + f*Sqrt[a + b*x + c*x^2])^(-1),x]

output

$$((2*(2*c*d - b*e)*f*\operatorname{ArcTan}[(\operatorname{Sqrt}[-4*c*d^2 + 4*b*d*e - 4*a*e^2 - b^2*f^2 + 4*a*c*f^2]*x)/(2*a*f + b*f*x + 2*d*\operatorname{Sqrt}[a + x*(b + c*x)] - 2*\operatorname{Sqrt}[a]*(d + e*x + f*\operatorname{Sqrt}[a + x*(b + c*x)])))]/\operatorname{Sqrt}[-4*c*d^2 + 4*b*d*e - 4*a*e^2 - b^2*f^2 + 4*a*c*f^2] + 2*\operatorname{Sqrt}[c]*f*\operatorname{ArcTanh}[(\operatorname{Sqrt}[c]*x)/(\operatorname{Sqrt}[a] - \operatorname{Sqrt}[a + x*(b + c*x)])] + e*(-\operatorname{Log}[-2*a - b*x + 2*\operatorname{Sqrt}[a]*\operatorname{Sqrt}[a + x*(b + c*x)]] + \operatorname{Log}[-2*a^{(3/2)}*f + 2*a*(d + e*x + f*\operatorname{Sqrt}[a + x*(b + c*x)]) + b*x*(d + e*x + f*\operatorname{Sqrt}[a + x*(b + c*x)]) - 2*\operatorname{Sqrt}[a]*(b*f*x + c*f*x^2 + (d + e*x)*\operatorname{Sqrt}[a + x*(b + c*x)])]))/(e^2 - c*f^2)$$

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1087 vs. 2(361) = 722.

Time = 10.42 (sec) , antiderivative size = 1087, normalized size of antiderivative = 3.01, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$,

Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{f\sqrt{a+bx+cx^2}+d+ex} dx$$

↓ 7293

$$\int \left(\frac{f\sqrt{a+bx+cx^2}}{af^2-x(2de-bf^2)-x^2(e^2-cf^2)-d^2} + \frac{d+ex}{-af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2} \right) dx$$

↓ 2009

$$\frac{(2cd-be)\operatorname{farctanh}\left(\frac{-bf^2+2de+2(e^2-cf^2)x}{f\sqrt{4ae^2-4bde+b^2f^2+4c(d^2-af^2)}}\right)}{(e^2-cf^2)\sqrt{4ae^2-4bde+b^2f^2+4c(d^2-af^2)}} - \frac{\sqrt{c}\operatorname{farctanh}\left(\frac{b+2cx}{2\sqrt{c}\sqrt{cx^2+bx+a}}\right)}{e^2-cf^2} -$$

$$\frac{\sqrt{2ae^4-2bde^3+2cd^2e^2+b^2f^2e^2-2acf^2e^2-2bcd^2e-(2cd-be)f\sqrt{4ae^2-4bde+b^2f^2+4c(d^2-af^2)}}e + \sqrt{2}(e^2-cf^2)\sqrt{4ae^2-4bde+b^2f^2+4c(d^2-af^2)}}{\sqrt{2ae^4-2bde^3+2cd^2e^2+b^2f^2e^2-2acf^2e^2-2bcd^2e+(2cd-be)f\sqrt{4ae^2-4bde+b^2f^2+4c(d^2-af^2)}}e + \sqrt{2}(e^2-cf^2)\sqrt{4ae^2-4bde+b^2f^2+4c(d^2-af^2)}} -$$

$$\frac{e \log(d^2-af^2+(e^2-cf^2)x^2+(2de-bf^2)x)}{2(e^2-cf^2)}$$

input `Int[(d + e*x + f*Sqrt[a + b*x + c*x^2])^(-1),x]`

output

```

((2*c*d - b*e)*f*ArcTanh[(2*d*e - b*f^2 + 2*(e^2 - c*f^2)*x)/(f*Sqrt[-4*b*d
d*e + 4*a*e^2 + b^2*f^2 + 4*c*(d^2 - a*f^2)])]/((e^2 - c*f^2)*Sqrt[-4*b*d
*e + 4*a*e^2 + b^2*f^2 + 4*c*(d^2 - a*f^2)]) - (Sqrt[c]*f*ArcTanh[(b + 2*c
*x)/(2*Sqrt[c]*Sqrt[a + b*x + c*x^2)])]/(e^2 - c*f^2) - (Sqrt[2*c*d^2*e^2
- 2*b*d*e^3 + 2*a*e^4 + 2*c^2*d^2*f^2 - 2*b*c*d*e*f^2 + b^2*e^2*f^2 - 2*a*
c*e^2*f^2 - e*(2*c*d - b*e)*f*Sqrt[-4*b*d*e + 4*a*e^2 + b^2*f^2 + 4*c*(d^2
- a*f^2)])*ArcTanh[(2*b*d*e - 4*a*(e^2 - c*f^2) - b*f*(b*f + Sqrt[-4*b*d*
e + 4*a*e^2 + b^2*f^2 + 4*c*(d^2 - a*f^2)]) + 2*(2*c*d*e - b*e^2 - c*f*Sqr
t[-4*b*d*e + 4*a*e^2 + b^2*f^2 + 4*c*(d^2 - a*f^2)])*x)/(2*Sqrt[2]*Sqrt[2*
c*d^2*e^2 - 2*b*d*e^3 + 2*a*e^4 + 2*c^2*d^2*f^2 - 2*b*c*d*e*f^2 + b^2*e^2*
f^2 - 2*a*c*e^2*f^2 - e*(2*c*d - b*e)*f*Sqrt[-4*b*d*e + 4*a*e^2 + b^2*f^2
+ 4*c*(d^2 - a*f^2)])*Sqrt[a + b*x + c*x^2])]/(Sqrt[2]*(e^2 - c*f^2)*Sqrt
[-4*b*d*e + 4*a*e^2 + b^2*f^2 + 4*c*(d^2 - a*f^2)]) + (Sqrt[2*c*d^2*e^2 -
2*b*d*e^3 + 2*a*e^4 + 2*c^2*d^2*f^2 - 2*b*c*d*e*f^2 + b^2*e^2*f^2 - 2*a*c*
e^2*f^2 + e*(2*c*d - b*e)*f*Sqrt[-4*b*d*e + 4*a*e^2 + b^2*f^2 + 4*c*(d^2 -
a*f^2)])*ArcTanh[(2*b*d*e - 4*a*e^2 - b^2*f^2 + 4*a*c*f^2 + b*f*Sqrt[-4*b
*d*e + 4*a*e^2 + b^2*f^2 + 4*c*(d^2 - a*f^2)] + 2*(2*c*d*e - b*e^2 + c*f*S
qrt[-4*b*d*e + 4*a*e^2 + b^2*f^2 + 4*c*(d^2 - a*f^2)])*x)/(2*Sqrt[2]*Sqrt[
2*c*d^2*e^2 - 2*b*d*e^3 + 2*a*e^4 + 2*c^2*d^2*f^2 - 2*b*c*d*e*f^2 + b^2*e^
2*f^2 - 2*a*c*e^2*f^2 + e*(2*c*d - b*e)*f*Sqrt[-4*b*d*e + 4*a*e^2 + b^2...

```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 7293

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 4878 vs. $2(325) = 650$.

Time = 0.15 (sec) , antiderivative size = 4879, normalized size of antiderivative = 13.52

method	result	size
default	Expression too large to display	4879

input `int(1/(d+e*x+f*(c*x^2+b*x+a)^(1/2)),x,method=_RETURNVERBOSE)`

output

$$\frac{f \cdot (-2 \cdot (c \cdot f^2 - e^2)) / (-f^2 \cdot (4 \cdot a \cdot c \cdot f^2 - b^2 \cdot f^2 - 4 \cdot a \cdot e^2 + 4 \cdot b \cdot d \cdot e - 4 \cdot c \cdot d^2))^{1/2}}{(2 \cdot c \cdot f^2 - 2 \cdot e^2) \cdot (1/2 \cdot (4 \cdot (x + (b \cdot f^2 - 2 \cdot d \cdot e + (-f^2 \cdot (4 \cdot a \cdot c \cdot f^2 - b^2 \cdot f^2 - 4 \cdot a \cdot e^2 + 4 \cdot b \cdot d \cdot e - 4 \cdot c \cdot d^2))^{1/2})) / (2 \cdot c \cdot f^2 - 2 \cdot e^2))^{2 \cdot c - 4 \cdot (b \cdot e^2 - 2 \cdot c \cdot d \cdot e + (f^2 \cdot (-4 \cdot a \cdot c \cdot f^2 + b^2 \cdot f^2 + 4 \cdot a \cdot e^2 - 4 \cdot b \cdot d \cdot e + 4 \cdot c \cdot d^2))^{1/2}) \cdot c) / (c \cdot f^2 - e^2) \cdot (x + (b \cdot f^2 - 2 \cdot d \cdot e + (-f^2 \cdot (4 \cdot a \cdot c \cdot f^2 - b^2 \cdot f^2 - 4 \cdot a \cdot e^2 + 4 \cdot b \cdot d \cdot e - 4 \cdot c \cdot d^2))^{1/2})) / (2 \cdot c \cdot f^2 - 2 \cdot e^2)) - 2 \cdot (2 \cdot a \cdot c \cdot e^2 \cdot f^2 - b^2 \cdot e^2 \cdot f^2 + 2 \cdot b \cdot c \cdot d \cdot e \cdot f^2 - 2 \cdot c^2 \cdot d^2 \cdot f^2 - 2 \cdot e^4 \cdot a + 2 \cdot d \cdot e^3 \cdot b - 2 \cdot d^2 \cdot e^2 \cdot c - (f^2 \cdot (-4 \cdot a \cdot c \cdot f^2 + b^2 \cdot f^2 + 4 \cdot a \cdot e^2 - 4 \cdot b \cdot d \cdot e + 4 \cdot c \cdot d^2))^{1/2}) \cdot b \cdot e^2 + 2 \cdot (f^2 \cdot (-4 \cdot a \cdot c \cdot f^2 + b^2 \cdot f^2 + 4 \cdot a \cdot e^2 - 4 \cdot b \cdot d \cdot e + 4 \cdot c \cdot d^2))^{1/2} \cdot c \cdot d \cdot e) / (c \cdot f^2 - e^2)^2)^{1/2} - 1/2 \cdot (b \cdot e^2 - 2 \cdot c \cdot d \cdot e + (f^2 \cdot (-4 \cdot a \cdot c \cdot f^2 + b^2 \cdot f^2 + 4 \cdot a \cdot e^2 - 4 \cdot b \cdot d \cdot e + 4 \cdot c \cdot d^2))^{1/2}) \cdot c) / (c \cdot f^2 - e^2) \cdot \ln((-1/2 \cdot (b \cdot e^2 - 2 \cdot c \cdot d \cdot e + (f^2 \cdot (-4 \cdot a \cdot c \cdot f^2 + b^2 \cdot f^2 + 4 \cdot a \cdot e^2 - 4 \cdot b \cdot d \cdot e + 4 \cdot c \cdot d^2))^{1/2}) \cdot c) / (c \cdot f^2 - e^2) + c \cdot (x + (b \cdot f^2 - 2 \cdot d \cdot e + (-f^2 \cdot (4 \cdot a \cdot c \cdot f^2 - b^2 \cdot f^2 - 4 \cdot a \cdot e^2 + 4 \cdot b \cdot d \cdot e - 4 \cdot c \cdot d^2))^{1/2})) / (2 \cdot c \cdot f^2 - 2 \cdot e^2))) / c^{1/2} + ((x + (b \cdot f^2 - 2 \cdot d \cdot e + (-f^2 \cdot (4 \cdot a \cdot c \cdot f^2 - b^2 \cdot f^2 - 4 \cdot a \cdot e^2 + 4 \cdot b \cdot d \cdot e - 4 \cdot c \cdot d^2))^{1/2})) / (2 \cdot c \cdot f^2 - 2 \cdot e^2))^{2 \cdot c - (b \cdot e^2 - 2 \cdot c \cdot d \cdot e + (f^2 \cdot (-4 \cdot a \cdot c \cdot f^2 + b^2 \cdot f^2 + 4 \cdot a \cdot e^2 - 4 \cdot b \cdot d \cdot e + 4 \cdot c \cdot d^2))^{1/2}) \cdot c) / (c \cdot f^2 - e^2) \cdot (x + (b \cdot f^2 - 2 \cdot d \cdot e + (-f^2 \cdot (4 \cdot a \cdot c \cdot f^2 - b^2 \cdot f^2 - 4 \cdot a \cdot e^2 + 4 \cdot b \cdot d \cdot e - 4 \cdot c \cdot d^2))^{1/2})) / (2 \cdot c \cdot f^2 - 2 \cdot e^2)) - 1/2 \cdot (2 \cdot a \cdot c \cdot e^2 \cdot f^2 - b^2 \cdot e^2 \cdot f^2 + 2 \cdot b \cdot c \cdot d \cdot e \cdot f^2 - 2 \cdot c^2 \cdot d^2 \cdot f^2 - 2 \cdot e^4 \cdot a + 2 \cdot d \cdot e^3 \cdot b - 2 \cdot d^2 \cdot e^2 \cdot c - (f^2 \cdot (-4 \cdot a \cdot c \cdot f^2 + b^2 \cdot f^2 + 4 \cdot a \cdot e^2 - 4 \cdot b \cdot d \cdot e + 4 \cdot c \cdot d^2))^{1/2}) \cdot b \cdot e^2 + 2 \cdot (f^2 \cdot (-4 \cdot a \cdot c \cdot f^2 + b^2 \cdot f^2 + 4 \cdot a \cdot e^2 - 4 \cdot b \cdot d \cdot e + 4 \cdot c \cdot d^2))^{1/2} \cdot c \cdot d \cdot e) / (c \cdot f^2 - e^2)^2)^{1/2}}{c^{1/2} + (2 \cdot a \cdot c \cdot e^2 \cdot f^2 - b^2 \cdot e^2 \cdot f^2 + 2 \cdot b \cdot c \cdot d \cdot e \cdot f^2 - 2 \cdot c^2 \cdot d^2 \cdot f^2 - \dots$$

Fricas [F(-1)]

Timed out.

$$\int \frac{1}{d + ex + f\sqrt{a + bx + cx^2}} dx = \text{Timed out}$$

input `integrate(1/(d+e*x+f*(c*x^2+b*x+a)^(1/2)),x, algorithm="fricas")`

output Timed out

Sympy [F]

$$\int \frac{1}{d + ex + f\sqrt{a + bx + cx^2}} dx = \int \frac{1}{d + ex + f\sqrt{a + bx + cx^2}} dx$$

input `integrate(1/(d+e*x+f*(c*x**2+b*x+a)**(1/2)),x)`

output `Integral(1/(d + e*x + f*sqrt(a + b*x + c*x**2)), x)`

Maxima [F]

$$\int \frac{1}{d + ex + f\sqrt{a + bx + cx^2}} dx = \int \frac{1}{ex + \sqrt{cx^2 + bx + a}f + d} dx$$

input `integrate(1/(d+e*x+f*(c*x^2+b*x+a)^(1/2)),x, algorithm="maxima")`

output `integrate(1/(e*x + sqrt(c*x^2 + b*x + a)*f + d), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{1}{d + ex + f\sqrt{a + bx + cx^2}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(d+e*x+f*(c*x^2+b*x+a)^(1/2)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Error: Bad Argument Type`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{d + ex + f\sqrt{a + bx + cx^2}} dx = \int \frac{1}{d + ex + f\sqrt{cx^2 + bx + a}} dx$$

input `int(1/(d + e*x + f*(a + b*x + c*x^2)^(1/2)),x)`output `int(1/(d + e*x + f*(a + b*x + c*x^2)^(1/2)), x)`**Reduce [F]**

$$\int \frac{1}{d + ex + f\sqrt{a + bx + cx^2}} dx = \int \frac{1}{d + ex + f\sqrt{cx^2 + bx + a}} dx$$

input `int(1/(d+e*x+f*(c*x^2+b*x+a)^(1/2)),x)`output `int(1/(d+e*x+f*(c*x^2+b*x+a)^(1/2)),x)`

3.28
$$\int \frac{1}{\left(d+ex+f\sqrt{a+bx+cx^2}\right)^2} dx$$

Optimal result	245
Mathematica [B] (warning: unable to verify)	246
Rubi [F]	247
Maple [B] (warning: unable to verify)	251
Fricas [B] (verification not implemented)	252
Sympy [F]	252
Maxima [F]	253
Giac [F(-2)]	253
Mupad [F(-1)]	253
Reduce [F]	254

Optimal result

Integrand size = 23, antiderivative size = 394

$$\int \frac{1}{\left(d+ex+f\sqrt{a+bx+cx^2}\right)^2} dx =$$

$$\frac{2\left(2b(cd^2+ae^2)-4a^{3/2}cef-b^2(2de-\sqrt{a}ef)\right)+\frac{(4a^{3/2}e^2+4\sqrt{a}d(cd-be)+b^2df-4acdf)(\sqrt{a}-\sqrt{a+bx+cx^2})}{x}}{(d-\sqrt{a}f)(4bde-4ae^2-b^2f^2-4c(d^2-af^2))\left(be-c(d+\sqrt{a}f)+\frac{(2\sqrt{a}e-bf)(\sqrt{a}-\sqrt{a+bx+cx^2})}{x}+\frac{(d-\sqrt{a}f)(\sqrt{a}-\sqrt{a+bx+cx^2})}{x} \right)}$$

$$+\frac{4(b^2-4ac)f\operatorname{arctanh}\left(\frac{2af+bf x+2d\sqrt{a+bx+cx^2}-2\sqrt{a}(d+ex+f\sqrt{a+bx+cx^2})}{\sqrt{-4bde+4ae^2+b^2f^2+4c(d^2-af^2)}x}\right)}{(-4bde+4ae^2+b^2f^2+4c(d^2-af^2))^{3/2}}$$

output

```
(-4*b*(a*e^2+c*d^2)+8*a^(3/2)*c*e*f+2*b^2*(2*d*e-a^(1/2)*e*f)-2*(4*a^(3/2)*e^2+4*a^(1/2)*d*(-b*e+c*d)+b^2*d*f-4*a*c*d*f)*(a^(1/2)-(c*x^2+b*x+a)^(1/2))/x)/(d-a^(1/2)*f)/(4*b*d*e-4*a*e^2-b^2*f^2-4*c*(-a*f^2+d^2))/(b*e-c*(d+a^(1/2)*f)+(2*a^(1/2)*e-b*f)*(a^(1/2)-(c*x^2+b*x+a)^(1/2))/x+(d-a^(1/2)*f)*(a^(1/2)-(c*x^2+b*x+a)^(1/2))^2/x^2)+4*(-4*a*c+b^2)*f*arctanh((2*a*f+b*f*x+2*d*(a+x*(c*x+b))^(1/2)-2*a^(1/2)*(d+e*x+f*(a+x*(c*x+b))^(1/2)))/(-4*b*d*e+4*a*e^2+b^2*f^2+4*c*(-a*f^2+d^2))^(1/2)/x)/(-4*b*d*e+4*a*e^2+b^2*f^2+4*c*(-a*f^2+d^2))^(3/2)
```

Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 2162 vs. $2(394) = 788$.

Time = 19.10 (sec) , antiderivative size = 2162, normalized size of antiderivative = 5.49

$$\int \frac{1}{(d + ex + f\sqrt{a + bx + cx^2})^2} dx = \text{Result too large to show}$$

input `Integrate[(d + e*x + f*Sqrt[a + b*x + c*x^2])^(-2),x]`

output

```
(-2*(2*c*d^3*e - 2*b*d^2*e^2 + 2*a*d*e^3 + b*c*d^2*f^2 - 4*a*c*d*e*f^2 + a
*b*e^2*f^2 + 2*c*d^2*e^2*x - 2*b*d*e^3*x + 2*a*e^4*x + 2*c^2*d^2*f^2*x - 2
*b*c*d*e*f^2*x + b^2*e^2*f^2*x - 2*a*c*e^2*f^2*x))/((e^2 - c*f^2)*(4*c*d^2
- 4*b*d*e + 4*a*e^2 + b^2*f^2 - 4*a*c*f^2)*(d^2 - a*f^2 + 2*d*e*x - b*f^2
*x + e^2*x^2 - c*f^2*x^2)) + (2*(-(b*d*f) + 2*a*e*f - 2*c*d*f*x + b*e*f*x)
*Sqrt[a + b*x + c*x^2])/((4*c*d^2 - 4*b*d*e + 4*a*e^2 + b^2*f^2 - 4*a*c*f^
2)*(d^2 - a*f^2 + 2*d*e*x - b*f^2*x + e^2*x^2 - c*f^2*x^2)) - (2*(-b^2 + 4
*a*c)*f*ArcTanh[(2*d*e - b*f^2 + 2*e^2*x - 2*c*f^2*x)/(f*Sqrt[4*c*d^2 - 4*
b*d*e + 4*a*e^2 + b^2*f^2 - 4*a*c*f^2]])/(4*c*d^2 - 4*b*d*e + 4*a*e^2 + b
^2*f^2 - 4*a*c*f^2)^(3/2) - ((-b^2 + 4*a*c)*f*(-2*c*d*f + b*e*f + e*Sqrt[4
*c*d^2 - 4*b*d*e + 4*a*e^2 + b^2*f^2 - 4*a*c*f^2])*Log[2*d*e - b*f^2 - f*S
qrt[4*c*d^2 - 4*b*d*e + 4*a*e^2 + b^2*f^2 - 4*a*c*f^2] + 2*e^2*x - 2*c*f^2
*x])/(Sqrt[2]*(4*c*d^2 - 4*b*d*e + 4*a*e^2 + b^2*f^2 - 4*a*c*f^2)^(3/2)*Sqr
t[2*c*d^2*e^2 - 2*b*d*e^3 + 2*a*e^4 + 2*c^2*d^2*f^2 - 2*b*c*d*e*f^2 + b^2
*e^2*f^2 - 2*a*c*e^2*f^2 - 2*c*d*e*f*Sqrt[4*c*d^2 - 4*b*d*e + 4*a*e^2 + b^
2*f^2 - 4*a*c*f^2] + b*e^2*f*Sqrt[4*c*d^2 - 4*b*d*e + 4*a*e^2 + b^2*f^2 -
4*a*c*f^2]]) + ((-b^2 + 4*a*c)*f*(-2*c*d*f + b*e*f - e*Sqrt[4*c*d^2 - 4*b*
d*e + 4*a*e^2 + b^2*f^2 - 4*a*c*f^2])*Log[2*d*e - b*f^2 + f*Sqrt[4*c*d^2 -
4*b*d*e + 4*a*e^2 + b^2*f^2 - 4*a*c*f^2] + 2*e^2*x - 2*c*f^2*x])/(Sqrt[2]
*(4*c*d^2 - 4*b*d*e + 4*a*e^2 + b^2*f^2 - 4*a*c*f^2)^(3/2)*Sqrt[2*c*d^2...
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\left(f\sqrt{a+bx+cx^2}+d+ex\right)^2} dx$$

↓ 7293

$$\int \left(\frac{2f^2(a+bx+cx^2)}{\left(-af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2\right)^2} - \frac{2df\sqrt{a+bx+cx^2}}{\left(-af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2\right)^2} - \frac{1}{(-af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2)} \right) dx$$

↓ 7239

$$\int \frac{2d\left(ex-f\sqrt{a+x(b+cx)}\right)+x\left(-2ef\sqrt{a+x(b+cx)}+bf^2+cf^2x+e^2x\right)+d^2\left(\frac{af^2}{d^2}+1\right)}{\left(-af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2\right)^2} dx$$

↓ 7293

$$\int \left(\frac{af^2+d^2}{\left(-af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2\right)^2} + \frac{2d\left(ex-f\sqrt{a+bx+cx^2}\right)}{\left(-af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2\right)^2} + \frac{x\left(-2ef\sqrt{a+x(b+cx)}+bf^2+cf^2x+e^2x\right)}{\left(-af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2\right)^2} \right) dx$$

↓ 7239

$$\int \frac{2d\left(ex-f\sqrt{a+x(b+cx)}\right)+x\left(-2ef\sqrt{a+x(b+cx)}+bf^2+cf^2x+e^2x\right)+d^2\left(\frac{af^2}{d^2}+1\right)}{\left(-af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2\right)^2} dx$$

↓ 7293

$$\int \left(\frac{af^2+d^2}{\left(-af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2\right)^2} + \frac{2d\left(ex-f\sqrt{a+bx+cx^2}\right)}{\left(-af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2\right)^2} + \frac{x\left(-2ef\sqrt{a+x(b+cx)}+bf^2+cf^2x+e^2x\right)}{\left(-af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2\right)^2} \right) dx$$

↓ 7239

$$\int \frac{2d\left(ex-f\sqrt{a+x(b+cx)}\right)+x\left(-2ef\sqrt{a+x(b+cx)}+bf^2+cf^2x+e^2x\right)+d^2\left(\frac{af^2}{d^2}+1\right)}{\left(-af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2\right)^2} dx$$

↓ 7293

$$\int \left(\frac{af^2 + d^2}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{2d(ex - f\sqrt{a + bx + cx^2})}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{x(-2ef\sqrt{a + bx + cx^2})}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} \right) dx$$

↓ 7239

$$\int \frac{2d(ex - f\sqrt{a + x(b + cx)}) + x(-2ef\sqrt{a + x(b + cx)} + bf^2 + cf^2x + e^2x) + d^2\left(\frac{af^2}{d^2} + 1\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} dx$$

↓ 7293

$$\int \left(\frac{af^2 + d^2}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{2d(ex - f\sqrt{a + bx + cx^2})}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{x(-2ef\sqrt{a + bx + cx^2})}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} \right) dx$$

↓ 7239

$$\int \frac{2d(ex - f\sqrt{a + x(b + cx)}) + x(-2ef\sqrt{a + x(b + cx)} + bf^2 + cf^2x + e^2x) + d^2\left(\frac{af^2}{d^2} + 1\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} dx$$

↓ 7293

$$\int \left(\frac{af^2 + d^2}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{2d(ex - f\sqrt{a + bx + cx^2})}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{x(-2ef\sqrt{a + bx + cx^2})}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} \right) dx$$

↓ 7239

$$\int \frac{2d(ex - f\sqrt{a + x(b + cx)}) + x(-2ef\sqrt{a + x(b + cx)} + bf^2 + cf^2x + e^2x) + d^2\left(\frac{af^2}{d^2} + 1\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} dx$$

↓ 7293

$$\int \left(\frac{af^2 + d^2}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{2d(ex - f\sqrt{a + bx + cx^2})}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{x(-2ef\sqrt{a + bx + cx^2})}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} \right) dx$$

↓ 7239

$$\int \frac{2d\left(ex - f\sqrt{a+x(b+cx)}\right) + x\left(-2ef\sqrt{a+x(b+cx)} + bf^2 + cf^2x + e^2x\right) + d^2\left(\frac{af^2}{d^2} + 1\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} dx$$

↓ 7293

$$\int \left(\frac{af^2 + d^2}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{2d\left(ex - f\sqrt{a+bx+cx^2}\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{x\left(-2ef\sqrt{a+bx+cx^2}\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} \right) dx$$

↓ 7239

$$\int \frac{2d\left(ex - f\sqrt{a+x(b+cx)}\right) + x\left(-2ef\sqrt{a+x(b+cx)} + bf^2 + cf^2x + e^2x\right) + d^2\left(\frac{af^2}{d^2} + 1\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} dx$$

↓ 7293

$$\int \left(\frac{af^2 + d^2}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{2d\left(ex - f\sqrt{a+bx+cx^2}\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{x\left(-2ef\sqrt{a+bx+cx^2}\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} \right) dx$$

↓ 7239

$$\int \frac{2d\left(ex - f\sqrt{a+x(b+cx)}\right) + x\left(-2ef\sqrt{a+x(b+cx)} + bf^2 + cf^2x + e^2x\right) + d^2\left(\frac{af^2}{d^2} + 1\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} dx$$

↓ 7293

$$\int \left(\frac{af^2 + d^2}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{2d\left(ex - f\sqrt{a+bx+cx^2}\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{x\left(-2ef\sqrt{a+bx+cx^2}\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} \right) dx$$

↓ 7239

$$\int \frac{2d\left(ex - f\sqrt{a+x(b+cx)}\right) + x\left(-2ef\sqrt{a+x(b+cx)} + bf^2 + cf^2x + e^2x\right) + d^2\left(\frac{af^2}{d^2} + 1\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} dx$$

↓ 7293

$$\int \left(\frac{af^2 + d^2}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{2d\left(ex - f\sqrt{a+bx+cx^2}\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{x\left(-2ef\sqrt{a+bx+cx^2}\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} \right) dx$$

↓ 7239

$$\int \frac{2d\left(ex - f\sqrt{a + x(b + cx)}\right) + x\left(-2ef\sqrt{a + x(b + cx)} + bf^2 + cf^2x + e^2x\right) + d^2\left(\frac{af^2}{d^2} + 1\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} dx$$

↓ 7293

$$\int \left(\frac{af^2 + d^2}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{2d\left(ex - f\sqrt{a + bx + cx^2}\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{x\left(-2ef\sqrt{a + bx + cx^2}\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} \right) dx$$

↓ 7239

$$\int \frac{2d\left(ex - f\sqrt{a + x(b + cx)}\right) + x\left(-2ef\sqrt{a + x(b + cx)} + bf^2 + cf^2x + e^2x\right) + d^2\left(\frac{af^2}{d^2} + 1\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} dx$$

↓ 7293

$$\int \left(\frac{af^2 + d^2}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{2d\left(ex - f\sqrt{a + bx + cx^2}\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{x\left(-2ef\sqrt{a + bx + cx^2}\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} \right) dx$$

↓ 7239

$$\int \frac{2d\left(ex - f\sqrt{a + x(b + cx)}\right) + x\left(-2ef\sqrt{a + x(b + cx)} + bf^2 + cf^2x + e^2x\right) + d^2\left(\frac{af^2}{d^2} + 1\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} dx$$

↓ 7293

$$\int \left(\frac{af^2 + d^2}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{2d\left(ex - f\sqrt{a + bx + cx^2}\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{x\left(-2ef\sqrt{a + bx + cx^2}\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} \right) dx$$

↓ 7239

$$\int \frac{2d\left(ex - f\sqrt{a + x(b + cx)}\right) + x\left(-2ef\sqrt{a + x(b + cx)} + bf^2 + cf^2x + e^2x\right) + d^2\left(\frac{af^2}{d^2} + 1\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} dx$$

↓ 7293

$$\int \left(\frac{af^2 + d^2}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{2d(ex - f\sqrt{a + bx + cx^2})}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{x(-2ef\sqrt{a + bx + cx^2})}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} \right) dx$$

↓ 7239

$$\int \frac{2d(ex - f\sqrt{a + x(b + cx)}) + x(-2ef\sqrt{a + x(b + cx)} + bf^2 + cf^2x + e^2x) + d^2\left(\frac{af^2}{d^2} + 1\right)}{(-af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} dx$$

input `Int[(d + e*x + f*Sqrt[a + b*x + c*x^2])^(-2),x]`

output `$Aborted`

Defintions of rubi rules used

rule 7239 `Int[u_, x_Symbol] :=> With[{v = SimplifyIntegrand[u, x]}, Int[v, x] /; SimplifyIntegrandQ[v, u, x]]`

rule 7293 `Int[u_, x_Symbol] :=> With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.72 (sec) , antiderivative size = 616160, normalized size of antiderivative = 1563.86

method	result	size
default	Expression too large to display	616160

input `int(1/(d+e*x+f*(c*x^2+b*x+a)^(1/2))^2,x,method=_RETURNVERBOSE)`

output result too large to display

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2867 vs. 2(359) = 718.

Time = 19.40 (sec) , antiderivative size = 6899, normalized size of antiderivative = 17.51

$$\int \frac{1}{(d + ex + f\sqrt{a + bx + cx^2})^2} dx = \text{Too large to display}$$

input `integrate(1/(d+e*x+f*(c*x^2+b*x+a)^(1/2))^2,x, algorithm="fricas")`

output Too large to include

Sympy [F]

$$\int \frac{1}{(d + ex + f\sqrt{a + bx + cx^2})^2} dx = \int \frac{1}{(d + ex + f\sqrt{a + bx + cx^2})^2} dx$$

input `integrate(1/(d+e*x+f*(c*x**2+b*x+a)**(1/2))**2,x)`

output `Integral((d + e*x + f*sqrt(a + b*x + c*x**2))**(-2), x)`

Maxima [F]

$$\int \frac{1}{(d + ex + f\sqrt{a + bx + cx^2})^2} dx = \int \frac{1}{(ex + \sqrt{cx^2 + bx + a}f + d)^2} dx$$

input `integrate(1/(d+e*x+f*(c*x^2+b*x+a)^(1/2))^2,x, algorithm="maxima")`

output `integrate((e*x + sqrt(c*x^2 + b*x + a)*f + d)^(-2), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{1}{(d + ex + f\sqrt{a + bx + cx^2})^2} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(d+e*x+f*(c*x^2+b*x+a)^(1/2))^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);OUTPUT:Error: Bad Argument Type`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex + f\sqrt{a + bx + cx^2})^2} dx = \int \frac{1}{(d + ex + f\sqrt{cx^2 + bx + a})^2} dx$$

input `int(1/(d + e*x + f*(a + b*x + c*x^2)^(1/2))^2,x)`

output `int(1/(d + e*x + f*(a + b*x + c*x^2)^(1/2))^2, x)`

Reduce [F]

$$\int \frac{1}{(d + ex + f\sqrt{a + bx + cx^2})^2} dx = \int \frac{1}{(d + ex + f\sqrt{cx^2 + bx + a})^2} dx$$

input `int(1/(d+e*x+f*(c*x^2+b*x+a)^(1/2))^2,x)`

output `int(1/(d+e*x+f*(c*x^2+b*x+a)^(1/2))^2,x)`

$$3.29 \quad \int \frac{1}{(d+ex+f\sqrt{a+bx+cx^2})^3} dx$$

Optimal result	255
Mathematica [B] (warning: unable to verify)	256
Rubi [F]	257
Maple [B] (warning: unable to verify)	258
Fricas [F(-1)]	259
Sympy [F]	259
Maxima [F]	259
Giac [F(-2)]	260
Mupad [F(-1)]	260
Reduce [F]	260

Optimal result

Integrand size = 23, antiderivative size = 1477

$$\int \frac{1}{(d+ex+f\sqrt{a+bx+cx^2})^3} dx = \text{Too large to display}$$

output

```
(b^4*d*e*f^2-b^3*(d+a^(1/2)*f)*(2*d*e^2+a^(1/2)*e^2*f+c*d*f^2)-4*b*(2*a^(1/2)*c^2*d^3*f+a^(3/2)*c*d*f*(-c*f^2+3*e^2)+a*c*d^2*(-c*f^2+4*e^2)+a^2*(-c*e^2*f^2+2*e^4))+8*a*c*e*(a*e^2*(d+a^(1/2)*f)+c*(d^3-a^(3/2)*f^3))+2*b^2*(a*e^3*(5*d+a^(1/2)*f)+c*e*(d^3+6*a^(1/2)*d^2*f-2*a*d*f^2+a^(3/2)*f^3))+4*a^(1/2)*b*d*(-b*e+c*d)*(-b*f^2+2*d*e)-16*a^(5/2)*(-c*e^2*f^2+e^4)+a^2*(-4*b*c*e*f^3-8*c^2*d*f^3+8*b*e^3*f)-b^2*d*f*(2*c*d^2-b*(-b*f^2+3*d*e))+4*a^(3/2)*(4*c^2*d^2*f^2+b*e^2*(-b*f^2+6*d*e)-4*c*d*e*(b*f^2+d*e))-a*f*(8*c^2*d^3+b^2*e*(-b*f^2+12*d*e)-6*b*c*d*(b*f^2+2*d*e))*(a^(1/2)-(c*x^2+b*x+a)^(1/2))/x/(d-a^(1/2)*f)^3/(4*b*d*e-4*a*e^2-b^2*f^2-4*c*(-a*f^2+d^2))/(b*e-c*(d+a^(1/2)*f)+(2*a^(1/2)*e-b*f)*(a^(1/2)-(c*x^2+b*x+a)^(1/2))/x+(d-a^(1/2)*f)*(a^(1/2)-(c*x^2+b*x+a)^(1/2))^2/x^2)^2+((b^5*d*f^4-b^4*e*f^2*(5*d^2+6*a^(1/2)*d*f-a*f^2)-16*a*e*(2*a^2*e^4+4*a*c*e^2*(-a*f^2+d^2)+c^2*(d-a^(1/2)*f)^2*(2*d^2+7*a^(1/2)*d*f+2*a*f^2))-2*b^3*(3*a^(3/2)*e^2*f^3-a*d*f^2*(-7*c*f^2+18*e^2)+3*a^(1/2)*d^2*f*(-2*c*f^2+e^2)-d^3*(c*f^2+8*e^2))+8*b*(10*a^2*d*e^4+c^2*d*(d-a^(1/2)*f)^2*(2*d^2+4*a^(1/2)*d*f+5*a*f^2)+3*a*c*e^2*(4*d^3+a^(1/2)*d^2*f-6*a*d*f^2+a^(3/2)*f^3))-4*b^2*(4*a*e^3*(a*f^2+4*d^2)+c*e*(8*d^4-3*a^(1/2)*d^3*f+5*a*d^2*f^2-9*a^(3/2)*d*f^3-a^2*f^4))/(d-a^(1/2)*f)^2+2*(3*b^2*d^2*(-b*e+2*c*d)*f-12*a^2*c*(-b*e+2*c*d)*f^3-3*a*(-b*e+2*c*d)*f*(-b^2*f^2+4*c*d^2)+16*a^(5/2)*(-c*f^2+e^2)^2+8*a^(3/2)*(2*c^2*d^2*f^2-b*e^2*(-b*f^2+4*d*e)+c*(-b^2*f^4+b*d*e*f^2+4*d^2*e^2))+a^(1/2)*(16*c^2*d^4...
```

Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 3355 vs. $2(1477) = 2954$.

Time = 21.96 (sec) , antiderivative size = 3355, normalized size of antiderivative = 2.27

$$\int \frac{1}{(d + ex + f\sqrt{a + bx + cx^2})^3} dx = \text{Result too large to show}$$

input

```
Integrate[(d + e*x + f*Sqrt[a + b*x + c*x^2])^(-3),x]
```

output

```
(-2*(-4*c^2*d^4*e*f^2 + 5*b*c*d^3*e^2*f^2 - b^2*d^2*e^3*f^2 - 2*a*c*d^2*e^3*f^2 - a*b*d*e^4*f^2 + 2*a^2*e^5*f^2 - b*c^2*d^3*f^4 + 6*a*c^2*d^2*e*f^4 - 3*a*b*c*d*e^2*f^4 + a*b^2*e^3*f^4 - 2*a^2*c*e^3*f^4 - 6*c^2*d^3*e^2*f^2*x + 9*b*c*d^2*e^3*f^2*x - 3*b^2*d*e^4*f^2*x - 6*a*c*d*e^4*f^2*x + 3*a*b*e^5*f^2*x - 2*c^3*d^3*f^4*x + 3*b*c^2*d^2*e*f^4*x - 3*b^2*c*d*e^2*f^4*x + 6*a*c^2*d*e^2*f^4*x + b^3*e^3*f^4*x - 3*a*b*c*e^3*f^4*x))/((e^2 - c*f^2)^2*(4*c*d^2 - 4*b*d*e + 4*a*e^2 + b^2*f^2 - 4*a*c*f^2)*(d^2 - a*f^2 + 2*d*e*x - b*f^2*x + e^2*x^2 - c*f^2*x^2)^2) + (-8*c^2*d^4*e^3 + 16*b*c*d^3*e^4 - 8*b^2*d^2*e^5 - 16*a*c*d^2*e^5 + 16*a*b*d*e^6 - 8*a^2*e^7 - 24*c^3*d^4*e*f^2 + 48*b*c^2*d^3*e^2*f^2 - 34*b^2*c*d^2*e^3*f^2 - 8*a*c^2*d^2*e^3*f^2 + 7*b^3*d*e^4*f^2 + 20*a*b*c*d*e^4*f^2 - 4*a*b^2*e^5*f^2 - 8*a^2*c*e^5*f^2 - 6*b^2*c^2*d^2*e*f^4 + 24*a*c^3*d^2*e*f^4 + 12*b^3*c*d*e^2*f^4 - 48*a*b*c^2*d*e^2*f^4 - 2*b^4*e^3*f^4 - 2*a*b^2*c*e^3*f^4 + 40*a^2*c^2*e^3*f^4 - 3*b^3*c^2*d*f^6 + 12*a*b*c^3*d*f^6 + 6*a*b^2*c^2*e*f^6 - 24*a^2*c^3*e*f^6 - 6*b^2*c*d*e^4*f^2*x + 24*a*c^2*d*e^4*f^2*x + 3*b^3*e^5*f^2*x - 12*a*b*c*e^5*f^2*x + 12*b^2*c^2*d*e^2*f^4*x - 48*a*c^3*d*e^2*f^4*x - 6*b^3*c*e^3*f^4*x + 24*a*b*c^2*e^3*f^4*x - 6*b^2*c^3*d*f^6*x + 24*a*c^4*d*f^6*x + 3*b^3*c^2*e*f^6*x - 12*a*b*c^3*e*f^6*x))/((e^2 - c*f^2)^2*(4*c*d^2 - 4*b*d*e + 4*a*e^2 + b^2*f^2 - 4*a*c*f^2)^2*(d^2 - a*f^2 + 2*d*e*x - b*f^2*x + e^2*x^2 - c*f^2*x^2)) + Sqrt[a + b*x + c*x^2]*((2*(2*c*d^3*e*f - 2*b*d^2*e^2*f + 2*a...
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(f\sqrt{a+bx+cx^2} + d + ex)^3} dx$$

↓ 7293

$$\int \left(\frac{4f^2(d+ex)(a+bx+cx^2)}{(-af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2)^3} - \frac{3f\sqrt{a+bx+cx^2}}{(-af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2)^2} + \frac{1}{(-af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2)} \right) dx$$

↓ 7299

$$\int \left(\frac{4f^2(d+ex)(a+bx+cx^2)}{(-af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2)^3} - \frac{3f\sqrt{a+bx+cx^2}}{(-af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2)^2} + \frac{1}{(-af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2)} \right) dx$$

input `Int[(d + e*x + f*Sqrt[a + b*x + c*x^2])^(-3),x]`

output `$Aborted`

Defintions of rubi rules used

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]`
`]`

rule 7299 `Int[u_, x_] := CannotIntegrate[u, x]`

Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 256.43 (sec) , antiderivative size = 6017742, normalized size of antiderivative = 4074.30

method	result	size
default	Expression too large to display	6017742

input `int(1/(d+e*x+f*(c*x^2+b*x+a)^(1/2))^3,x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex + f\sqrt{a + bx + cx^2})^3} dx = \text{Timed out}$$

input `integrate(1/(d+e*x+f*(c*x^2+b*x+a)^(1/2))^3,x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{1}{(d + ex + f\sqrt{a + bx + cx^2})^3} dx = \int \frac{1}{(d + ex + f\sqrt{a + bx + cx^2})^3} dx$$

input `integrate(1/(d+e*x+f*(c*x**2+b*x+a)**(1/2))**3,x)`

output `Integral((d + e*x + f*sqrt(a + b*x + c*x**2))**(-3), x)`

Maxima [F]

$$\int \frac{1}{(d + ex + f\sqrt{a + bx + cx^2})^3} dx = \int \frac{1}{(ex + \sqrt{cx^2 + bx + af + d})^3} dx$$

input `integrate(1/(d+e*x+f*(c*x^2+b*x+a)^(1/2))^3,x, algorithm="maxima")`

output `integrate((e*x + sqrt(c*x^2 + b*x + a)*f + d)^(-3), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{1}{(d + ex + f\sqrt{a + bx + cx^2})^3} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(d+e*x+f*(c*x^2+b*x+a)^(1/2))^3,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);;OUTPUT:Error: Bad Argument Type`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex + f\sqrt{a + bx + cx^2})^3} dx = \int \frac{1}{(d + ex + f\sqrt{cx^2 + bx + a})^3} dx$$

input `int(1/(d + e*x + f*(a + b*x + c*x^2)^(1/2))^3,x)`

output `int(1/(d + e*x + f*(a + b*x + c*x^2)^(1/2))^3, x)`

Reduce [F]

$$\int \frac{1}{(d + ex + f\sqrt{a + bx + cx^2})^3} dx = \int \frac{1}{(d + ex + f\sqrt{cx^2 + bx + a})^3} dx$$

input `int(1/(d+e*x+f*(c*x^2+b*x+a)^(1/2))^3,x)`

output `int(1/(d+e*x+f*(c*x^2+b*x+a)^(1/2))^3,x)`

3.30 $\int (x + \sqrt{-3 - 2x + 4x^2})^3 dx$

Optimal result	261
Mathematica [A] (verified)	261
Rubi [A] (verified)	262
Maple [A] (verified)	263
Fricas [A] (verification not implemented)	263
Sympy [A] (verification not implemented)	264
Maxima [A] (verification not implemented)	264
Giac [A] (verification not implemented)	265
Mupad [B] (verification not implemented)	265
Reduce [B] (verification not implemented)	266

Optimal result

Integrand size = 18, antiderivative size = 108

$$\int (x + \sqrt{-3 - 2x + 4x^2})^3 dx = -\frac{9x^2}{2} - 2x^3 + \frac{13x^4}{4} + \frac{105}{512}(1 - 4x)\sqrt{-3 - 2x + 4x^2} + \frac{1}{64}(-3 - 2x + 4x^2)^{3/2} + \frac{7}{16}x(-3 - 2x + 4x^2)^{3/2} - \frac{1365 \operatorname{arctanh}\left(\frac{1-4x}{2\sqrt{-3-2x+4x^2}}\right)}{1024}$$

output

```
-9/2*x^2-2*x^3+13/4*x^4+105/512*(1-4*x)*(4*x^2-2*x-3)^(1/2)+1/64*(4*x^2-2*x-3)^(3/2)+7/16*x*(4*x^2-2*x-3)^(3/2)-1365/1024*arctanh(1/2*(1-4*x)/(4*x^2-2*x-3)^(1/2))
```

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.71

$$\int (x + \sqrt{-3 - 2x + 4x^2})^3 dx = \frac{1}{4}x^2(-18 - 8x + 13x^2) + \frac{1}{512}\sqrt{-3 - 2x + 4x^2}(81 - 1108x - 416x^2 + 896x^3) - \frac{1365 \log(1 - 4x + 2\sqrt{-3 - 2x + 4x^2})}{1024}$$

input `Integrate[(x + Sqrt[-3 - 2*x + 4*x^2])^3,x]`

output $(x^2*(-18 - 8*x + 13*x^2))/4 + (\text{Sqrt}[-3 - 2*x + 4*x^2]*(81 - 1108*x - 416*x^2 + 896*x^3))/512 - (1365*\text{Log}[1 - 4*x + 2*\text{Sqrt}[-3 - 2*x + 4*x^2]])/1024$

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\sqrt{4x^2 - 2x - 3} + x)^3 dx$$

$$\downarrow 7293$$

$$\int (13x^3 + 7\sqrt{4x^2 - 2x - 3}x^2 - 6x^2 - 2\sqrt{4x^2 - 2x - 3}x - 3\sqrt{4x^2 - 2x - 3} - 9x) dx$$

$$\downarrow 2009$$

$$-\frac{1365 \operatorname{arctanh}\left(\frac{1-4x}{2\sqrt{4x^2-2x-3}}\right)}{1024} + \frac{13x^4}{4} - 2x^3 - \frac{9x^2}{2} + \frac{7}{16}(4x^2 - 2x - 3)^{3/2}x + \frac{1}{64}(4x^2 - 2x - 3)^{3/2} + \frac{105}{512}(1 - 4x)\sqrt{4x^2 - 2x - 3}$$

input `Int[(x + Sqrt[-3 - 2*x + 4*x^2])^3,x]`

output $(-9*x^2)/2 - 2*x^3 + (13*x^4)/4 + (105*(1 - 4*x)*\text{Sqrt}[-3 - 2*x + 4*x^2])/512 + (-3 - 2*x + 4*x^2)^{(3/2)}/64 + (7*x*(-3 - 2*x + 4*x^2)^{(3/2)})/16 - (1365*\text{ArcTanh}[(1 - 4*x)/(2*\text{Sqrt}[-3 - 2*x + 4*x^2])])/1024$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]`
`]`

Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.62

method	result
trager	$\frac{(13x^2-8x-18)x^2}{4} + \left(\frac{7}{4}x^3 - \frac{13}{16}x^2 - \frac{277}{128}x + \frac{81}{512}\right) \sqrt{4x^2 - 2x - 3} - \frac{1365 \ln(1-4x+2\sqrt{4x^2-2x-3})}{1024}$
default	$\frac{(8x-2)(4x^2-2x-3)^{\frac{3}{2}}}{32} - \frac{105(8x-2)\sqrt{4x^2-2x-3}}{1024} + \frac{1365 \ln\left(\frac{(4x-1)\sqrt{4} + \sqrt{4x^2-2x-3}}{4}\right)\sqrt{4}}{2048} + \frac{13x^4}{4} - 2x^3 - \frac{9x^2}{2} + \frac{3x(4x^2-2x-3)}{4}$

input `int((x+(4*x^2-2*x-3)^(1/2))^3,x,method=_RETURNVERBOSE)`

output `1/4*(13*x^2-8*x-18)*x^2+(7/4*x^3-13/16*x^2-277/128*x+81/512)*(4*x^2-2*x-3)^(1/2)-1365/1024*ln(1-4*x+2*(4*x^2-2*x-3)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.62

$$\int \left(x + \sqrt{-3 - 2x + 4x^2}\right)^3 dx = \frac{13}{4}x^4 - 2x^3 - \frac{9}{2}x^2 + \frac{1}{512}(896x^3 - 416x^2 - 1108x + 81)\sqrt{4x^2 - 2x - 3} - \frac{1365}{1024} \log\left(-4x + 2\sqrt{4x^2 - 2x - 3} + 1\right)$$

input `integrate((x+(4*x^2-2*x-3)^(1/2))^3,x, algorithm="fricas")`

output

```
13/4*x^4 - 2*x^3 - 9/2*x^2 + 1/512*(896*x^3 - 416*x^2 - 1108*x + 81)*sqrt(
4*x^2 - 2*x - 3) - 1365/1024*log(-4*x + 2*sqrt(4*x^2 - 2*x - 3) + 1)
```

Sympy [A] (verification not implemented)

Time = 0.55 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.17

$$\int \left(x + \sqrt{-3 - 2x + 4x^2} \right)^3 dx = \frac{13x^4}{4} - 2x^3 - \frac{9x^2}{2} - 3\left(\frac{x}{2} - \frac{1}{8}\right) \sqrt{4x^2 - 2x - 3} - 2\left(\frac{x^2}{3} - \frac{x}{24} - \frac{9}{32}\right) \sqrt{4x^2 - 2x - 3} + 7\sqrt{4x^2 - 2x - 3} \left(\frac{x^3}{4} - \frac{x^2}{48} - \frac{41x}{384} - \frac{57}{512}\right) + \frac{1365 \log(8x + 4\sqrt{4x^2 - 2x - 3} - 2)}{1024}$$

input

```
integrate((x+(4*x**2-2*x-3)**(1/2))**3,x)
```

output

```
13*x**4/4 - 2*x**3 - 9*x**2/2 - 3*(x/2 - 1/8)*sqrt(4*x**2 - 2*x - 3) - 2*(
x**2/3 - x/24 - 9/32)*sqrt(4*x**2 - 2*x - 3) + 7*sqrt(4*x**2 - 2*x - 3)*(
**3/4 - x**2/48 - 41*x/384 - 57/512) + 1365*log(8*x + 4*sqrt(4*x**2 - 2*x
- 3) - 2)/1024
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.89

$$\int \left(x + \sqrt{-3 - 2x + 4x^2} \right)^3 dx = \frac{13}{4} x^4 - 2x^3 + \frac{7}{16} (4x^2 - 2x - 3)^{\frac{3}{2}} x - \frac{9}{2} x^2 + \frac{1}{64} (4x^2 - 2x - 3)^{\frac{3}{2}} - \frac{105}{128} \sqrt{4x^2 - 2x - 3} x + \frac{105}{512} \sqrt{4x^2 - 2x - 3} + \frac{1365}{1024} \log(8x + 4\sqrt{4x^2 - 2x - 3} - 2)$$

input

```
integrate((x+(4*x^2-2*x-3)^(1/2))^3,x, algorithm="maxima")
```

output

$$\begin{aligned} & 13/4*x^4 - 2*x^3 + 7/16*(4*x^2 - 2*x - 3)^{(3/2)}*x - 9/2*x^2 + 1/64*(4*x^2 \\ & - 2*x - 3)^{(3/2)} - 105/128*\text{sqrt}(4*x^2 - 2*x - 3)*x + 105/512*\text{sqrt}(4*x^2 - \\ & 2*x - 3) + 1365/1024*\log(8*x + 4*\text{sqrt}(4*x^2 - 2*x - 3) - 2) \end{aligned}$$
Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.63

$$\begin{aligned} \int \left(x + \sqrt{-3 - 2x + 4x^2} \right)^3 dx &= \frac{13}{4} x^4 - 2x^3 - \frac{9}{2} x^2 \\ &+ \frac{1}{512} (4(8(28x - 13)x - 277)x + 81)\sqrt{4x^2 - 2x - 3} \\ &- \frac{1365}{1024} \log \left(\left| -4x + 2\sqrt{4x^2 - 2x - 3} + 1 \right| \right) \end{aligned}$$

input

`integrate((x+(4*x^2-2*x-3)^(1/2))^3,x, algorithm="giac")`

output

$$\begin{aligned} & 13/4*x^4 - 2*x^3 - 9/2*x^2 + 1/512*(4*(8*(28*x - 13)*x - 277)*x + 81)*\text{sqrt} \\ & (4*x^2 - 2*x - 3) - 1365/1024*\log(\text{abs}(-4*x + 2*\text{sqrt}(4*x^2 - 2*x - 3) + 1)) \end{aligned}$$
Mupad [B] (verification not implemented)

Time = 0.51 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.06

$$\begin{aligned} \int \left(x + \sqrt{-3 - 2x + 4x^2} \right)^3 dx &= \frac{351 \ln \left(2x + \sqrt{4x^2 - 2x - 3} - \frac{1}{2} \right)}{256} \\ &- \frac{39 \ln \left(x + \frac{\sqrt{4x^2 - 2x - 3}}{2} - \frac{1}{4} \right)}{1024} \\ &- \frac{27 \left(\frac{x}{2} - \frac{1}{8} \right) \sqrt{4x^2 - 2x - 3}}{16} \\ &- \frac{\sqrt{4x^2 - 2x - 3} (-128x^2 + 16x + 108)}{2048} \\ &+ \frac{7x(4x^2 - 2x - 3)^{3/2}}{16} - \frac{9x^2}{2} - 2x^3 + \frac{13x^4}{4} \end{aligned}$$

input

`int((x + (4*x^2 - 2*x - 3)^(1/2))^3,x)`

output

```
(351*log(2*x + (4*x^2 - 2*x - 3)^(1/2) - 1/2))/256 - (39*log(x + (4*x^2 - 2*x - 3)^(1/2)/2 - 1/4))/1024 - (27*(x/2 - 1/8)*(4*x^2 - 2*x - 3)^(1/2))/16 - ((4*x^2 - 2*x - 3)^(1/2)*(16*x - 128*x^2 + 108))/2048 + (7*x*(4*x^2 - 2*x - 3)^(3/2))/16 - (9*x^2)/2 - 2*x^3 + (13*x^4)/4
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.94

$$\int \left(x + \sqrt{-3 - 2x + 4x^2} \right)^3 dx = \frac{7\sqrt{4x^2 - 2x - 3}x^3}{4} - \frac{13\sqrt{4x^2 - 2x - 3}x^2}{16} - \frac{277\sqrt{4x^2 - 2x - 3}x}{128} + \frac{81\sqrt{4x^2 - 2x - 3}}{512} + \frac{1365 \log\left(\frac{2\sqrt{4x^2 - 2x - 3} + 4x - 1}{\sqrt{13}}\right)}{1024} + \frac{13x^4}{4} - 2x^3 - \frac{9x^2}{2}$$

input

```
int((x+(4*x^2-2*x-3)^(1/2))^3,x)
```

output

```
(1792*sqrt(4*x**2 - 2*x - 3)*x**3 - 832*sqrt(4*x**2 - 2*x - 3)*x**2 - 2216*sqrt(4*x**2 - 2*x - 3)*x + 162*sqrt(4*x**2 - 2*x - 3) + 1365*log((2*sqrt(4*x**2 - 2*x - 3) + 4*x - 1)/sqrt(13)) + 3328*x**4 - 2048*x**3 - 4608*x**2)/1024
```

3.31 $\int (x + \sqrt{-3 - 2x + 4x^2})^2 dx$

Optimal result	267
Mathematica [A] (verified)	267
Rubi [A] (verified)	268
Maple [A] (verified)	269
Fricas [A] (verification not implemented)	269
Sympy [A] (verification not implemented)	270
Maxima [A] (verification not implemented)	270
Giac [A] (verification not implemented)	271
Mupad [B] (verification not implemented)	271
Reduce [B] (verification not implemented)	272

Optimal result

Integrand size = 18, antiderivative size = 85

$$\begin{aligned} \int (x + \sqrt{-3 - 2x + 4x^2})^2 dx &= -3x - x^2 + \frac{5x^3}{3} - \frac{1}{16}(1 - 4x)\sqrt{-3 - 2x + 4x^2} \\ &\quad + \frac{1}{6}(-3 - 2x + 4x^2)^{3/2} \\ &\quad + \frac{13}{32}\operatorname{arctanh}\left(\frac{1 - 4x}{2\sqrt{-3 - 2x + 4x^2}}\right) \end{aligned}$$

output

```
-3*x-x^2+5/3*x^3-1/16*(1-4*x)*(4*x^2-2*x-3)^(1/2)+1/6*(4*x^2-2*x-3)^(3/2)+
13/32*arctanh(1/2*(1-4*x)/(4*x^2-2*x-3)^(1/2))
```

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.82

$$\begin{aligned} \int (x + \sqrt{-3 - 2x + 4x^2})^2 dx &= \frac{1}{3}x(-9 - 3x + 5x^2) \\ &\quad + \frac{1}{48}\sqrt{-3 - 2x + 4x^2}(-27 - 4x + 32x^2) \\ &\quad + \frac{13}{32}\log\left(1 - 4x + 2\sqrt{-3 - 2x + 4x^2}\right) \end{aligned}$$

input `Integrate[(x + Sqrt[-3 - 2*x + 4*x^2])^2,x]`

output `(x*(-9 - 3*x + 5*x^2))/3 + (Sqrt[-3 - 2*x + 4*x^2]*(-27 - 4*x + 32*x^2))/4
8 + (13*Log[1 - 4*x + 2*Sqrt[-3 - 2*x + 4*x^2]])/32`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\sqrt{4x^2 - 2x - 3} + x \right)^2 dx$$

$$\downarrow \text{7293}$$

$$\int \left(5x^2 + 2\sqrt{4x^2 - 2x - 3}x - 2x - 3 \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{13}{32} \operatorname{arctanh} \left(\frac{1 - 4x}{2\sqrt{4x^2 - 2x - 3}} \right) + \frac{5x^3}{3} - x^2 + \frac{1}{6} (4x^2 - 2x - 3)^{3/2} - \frac{1}{16} (1 - 4x) \sqrt{4x^2 - 2x - 3} - 3x$$

input `Int[(x + Sqrt[-3 - 2*x + 4*x^2])^2,x]`

output `-3*x - x^2 + (5*x^3)/3 - ((1 - 4*x)*Sqrt[-3 - 2*x + 4*x^2])/16 + (-3 - 2*x
+ 4*x^2)^(3/2)/6 + (13*ArcTanh[(1 - 4*x)/(2*Sqrt[-3 - 2*x + 4*x^2]])/32`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]`
`]`

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.71

method	result	size
trager	$\frac{(5x^2-3x-9)x}{3} + \left(\frac{2}{3}x^2 - \frac{1}{12}x - \frac{9}{16}\right) \sqrt{4x^2 - 2x - 3} - \frac{13 \ln(2\sqrt{4x^2-2x-3}-1+4x)}{32}$	60
default	$\frac{5x^3}{3} - x^2 - 3x + \frac{(4x^2-2x-3)^{\frac{3}{2}}}{6} + \frac{(8x-2)\sqrt{4x^2-2x-3}}{32} - \frac{13 \ln\left(\frac{(4x-1)\sqrt{4} + \sqrt{4x^2-2x-3}}{4}\right)\sqrt{4}}{64}$	77

input `int((x+(4*x^2-2*x-3)^(1/2))^2,x,method=_RETURNVERBOSE)`

output `1/3*(5*x^2-3*x-9)*x+(2/3*x^2-1/12*x-9/16)*(4*x^2-2*x-3)^(1/2)-13/32*ln(2*(4*x^2-2*x-3)^(1/2)-1+4*x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.71

$$\int \left(x + \sqrt{-3 - 2x + 4x^2}\right)^2 dx = \frac{5}{3}x^3 - x^2 + \frac{1}{48}(32x^2 - 4x - 27)\sqrt{4x^2 - 2x - 3} - 3x + \frac{13}{32} \log\left(-4x + 2\sqrt{4x^2 - 2x - 3} + 1\right)$$

input `integrate((x+(4*x^2-2*x-3)^(1/2))^2,x, algorithm="fricas")`

output `5/3*x^3 - x^2 + 1/48*(32*x^2 - 4*x - 27)*sqrt(4*x^2 - 2*x - 3) - 3*x + 13/32*log(-4*x + 2*sqrt(4*x^2 - 2*x - 3) + 1)`

Sympy [A] (verification not implemented)

Time = 0.47 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.76

$$\int \left(x + \sqrt{-3 - 2x + 4x^2} \right)^2 dx = \frac{5x^3}{3} - x^2 - 3x + 2 \left(\frac{x^2}{3} - \frac{x}{24} - \frac{9}{32} \right) \sqrt{4x^2 - 2x - 3} - \frac{13 \log(8x + 4\sqrt{4x^2 - 2x - 3} - 2)}{32}$$

input `integrate((x+(4*x**2-2*x-3)**(1/2))**2,x)`output `5*x**3/3 - x**2 - 3*x + 2*(x**2/3 - x/24 - 9/32)*sqrt(4*x**2 - 2*x - 3) - 13*log(8*x + 4*sqrt(4*x**2 - 2*x - 3) - 2)/32`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.93

$$\int \left(x + \sqrt{-3 - 2x + 4x^2} \right)^2 dx = \frac{5}{3}x^3 - x^2 + \frac{1}{6}(4x^2 - 2x - 3)^{\frac{3}{2}} + \frac{1}{4}\sqrt{4x^2 - 2x - 3}x - 3x - \frac{1}{16}\sqrt{4x^2 - 2x - 3} - \frac{13}{32}\log(8x + 4\sqrt{4x^2 - 2x - 3} - 2)$$

input `integrate((x+(4*x^2-2*x-3)^(1/2))^2,x, algorithm="maxima")`output `5/3*x^3 - x^2 + 1/6*(4*x^2 - 2*x - 3)^(3/2) + 1/4*sqrt(4*x^2 - 2*x - 3)*x - 3*x - 1/16*sqrt(4*x^2 - 2*x - 3) - 13/32*log(8*x + 4*sqrt(4*x^2 - 2*x - 3) - 2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.72

$$\int \left(x + \sqrt{-3 - 2x + 4x^2} \right)^2 dx = \frac{5}{3}x^3 - x^2 + \frac{1}{48}(4(8x - 1)x - 27)\sqrt{4x^2 - 2x - 3} - 3x + \frac{13}{32} \log \left(\left| -4x + 2\sqrt{4x^2 - 2x - 3} + 1 \right| \right)$$

input `integrate((x+(4*x^2-2*x-3)^(1/2))^2,x, algorithm="giac")`output `5/3*x^3 - x^2 + 1/48*(4*(8*x - 1)*x - 27)*sqrt(4*x^2 - 2*x - 3) - 3*x + 13/32*log(abs(-4*x + 2*sqrt(4*x^2 - 2*x - 3) + 1))`**Mupad [B] (verification not implemented)**

Time = 20.92 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.68

$$\int \left(x + \sqrt{-3 - 2x + 4x^2} \right)^2 dx = \frac{5x^3}{3} - \frac{13 \ln \left(x + \frac{\sqrt{4x^2 - 2x - 3}}{2} - \frac{1}{4} \right)}{32} - \frac{\sqrt{4x^2 - 2x - 3}(-128x^2 + 16x + 108)}{192} - x^2 - 3x$$

input `int((x + (4*x^2 - 2*x - 3)^(1/2))^2,x)`output `(5*x^3)/3 - (13*log(x + (4*x^2 - 2*x - 3)^(1/2)/2 - 1/4))/32 - ((4*x^2 - 2*x - 3)^(1/2)*(16*x - 128*x^2 + 108))/192 - x^2 - 3*x`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.98

$$\int \left(x + \sqrt{-3 - 2x + 4x^2} \right)^2 dx = \frac{2\sqrt{4x^2 - 2x - 3} x^2}{3} - \frac{\sqrt{4x^2 - 2x - 3} x}{12} - \frac{9\sqrt{4x^2 - 2x - 3}}{16} - \frac{13 \log\left(\frac{2\sqrt{4x^2 - 2x - 3} + 4x - 1}{\sqrt{13}}\right)}{32} + \frac{5x^3}{3} - x^2 - 3x$$

input `int((x+(4*x^2-2*x-3)^(1/2))^2,x)`output `(64*sqrt(4*x**2 - 2*x - 3)*x**2 - 8*sqrt(4*x**2 - 2*x - 3)*x - 54*sqrt(4*x**2 - 2*x - 3) - 39*log((2*sqrt(4*x**2 - 2*x - 3) + 4*x - 1)/sqrt(13)) + 160*x**3 - 96*x**2 - 288*x)/96`

3.32 $\int (x + \sqrt{-3 - 2x + 4x^2}) dx$

Optimal result	273
Mathematica [A] (verified)	273
Rubi [A] (verified)	274
Maple [A] (verified)	275
Fricas [A] (verification not implemented)	275
Sympy [A] (verification not implemented)	276
Maxima [A] (verification not implemented)	276
Giac [A] (verification not implemented)	277
Mupad [B] (verification not implemented)	277
Reduce [B] (verification not implemented)	278

Optimal result

Integrand size = 16, antiderivative size = 59

$$\int (x + \sqrt{-3 - 2x + 4x^2}) dx = \frac{x^2}{2} - \frac{1}{8}(1 - 4x)\sqrt{-3 - 2x + 4x^2} + \frac{13}{16} \operatorname{arctanh}\left(\frac{1 - 4x}{2\sqrt{-3 - 2x + 4x^2}}\right)$$

output

```
1/2*x^2-1/8*(1-4*x)*(4*x^2-2*x-3)^(1/2)+13/16*arctanh(1/2*(1-4*x)/(4*x^2-2*x-3)^(1/2))
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.97

$$\int (x + \sqrt{-3 - 2x + 4x^2}) dx = \frac{x^2}{2} + \frac{1}{8}(-1 + 4x)\sqrt{-3 - 2x + 4x^2} + \frac{13}{16} \log\left(1 - 4x + 2\sqrt{-3 - 2x + 4x^2}\right)$$

input

```
Integrate[x + Sqrt[-3 - 2*x + 4*x^2], x]
```

output

$$x^2/2 + ((-1 + 4x)\sqrt{-3 - 2x + 4x^2})/8 + (13\text{Log}[1 - 4x + 2\sqrt{-3 - 2x + 4x^2}])/16$$
Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.062$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\sqrt{4x^2 - 2x - 3} + x) dx$$

↓ 2009

$$\frac{13}{16} \operatorname{arctanh}\left(\frac{1 - 4x}{2\sqrt{4x^2 - 2x - 3}}\right) + \frac{x^2}{2} - \frac{1}{8}(1 - 4x)\sqrt{4x^2 - 2x - 3}$$

input

$$\text{Int}[x + \text{Sqrt}[-3 - 2*x + 4*x^2], x]$$

output

$$x^2/2 - ((1 - 4x)\sqrt{-3 - 2x + 4x^2})/8 + (13\text{ArcTanh}[(1 - 4x)/(2\sqrt{-3 - 2x + 4x^2}))/16$$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.80

method	result	size
trager	$\frac{x^2}{2} + \left(\frac{x}{2} - \frac{1}{8}\right) \sqrt{4x^2 - 2x - 3} + \frac{13 \ln(1 - 4x + 2\sqrt{4x^2 - 2x - 3})}{16}$	47
default	$\frac{x^2}{2} + \frac{(8x-2)\sqrt{4x^2-2x-3}}{16} - \frac{13 \ln\left(\frac{(4x-1)\sqrt{4}}{4} + \sqrt{4x^2-2x-3}\right)\sqrt{4}}{32}$	55
parts	$\frac{x^2}{2} + \frac{(8x-2)\sqrt{4x^2-2x-3}}{16} - \frac{13 \ln\left(\frac{(4x-1)\sqrt{4}}{4} + \sqrt{4x^2-2x-3}\right)\sqrt{4}}{32}$	55

input `int(x+(4*x^2-2*x-3)^(1/2),x,method=_RETURNVERBOSE)`

output `1/2*x^2+(1/2*x-1/8)*(4*x^2-2*x-3)^(1/2)+13/16*ln(1-4*x+2*(4*x^2-2*x-3)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.80

$$\int \left(x + \sqrt{-3 - 2x + 4x^2} \right) dx = \frac{1}{2} x^2 + \frac{1}{8} \sqrt{4x^2 - 2x - 3} (4x - 1) + \frac{13}{16} \log \left(-4x + 2\sqrt{4x^2 - 2x - 3} + 1 \right)$$

input `integrate(x+(4*x^2-2*x-3)^(1/2),x, algorithm="fricas")`

output `1/2*x^2 + 1/8*sqrt(4*x^2 - 2*x - 3)*(4*x - 1) + 13/16*log(-4*x + 2*sqrt(4*x^2 - 2*x - 3) + 1)`

Sympy [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.83

$$\int \left(x + \sqrt{-3 - 2x + 4x^2} \right) dx = \frac{x^2}{2} + \left(\frac{x}{2} - \frac{1}{8} \right) \sqrt{4x^2 - 2x - 3} - \frac{13 \log(8x + 4\sqrt{4x^2 - 2x - 3} - 2)}{16}$$

input `integrate(x+(4*x**2-2*x-3)**(1/2),x)`output `x**2/2 + (x/2 - 1/8)*sqrt(4*x**2 - 2*x - 3) - 13*log(8*x + 4*sqrt(4*x**2 - 2*x - 3) - 2)/16`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.97

$$\int \left(x + \sqrt{-3 - 2x + 4x^2} \right) dx = \frac{1}{2} x^2 + \frac{1}{2} \sqrt{4x^2 - 2x - 3} x - \frac{1}{8} \sqrt{4x^2 - 2x - 3} - \frac{13}{16} \log(8x + 4\sqrt{4x^2 - 2x - 3} - 2)$$

input `integrate(x+(4*x^2-2*x-3)^(1/2),x, algorithm="maxima")`output `1/2*x^2 + 1/2*sqrt(4*x^2 - 2*x - 3)*x - 1/8*sqrt(4*x^2 - 2*x - 3) - 13/16*log(8*x + 4*sqrt(4*x^2 - 2*x - 3) - 2)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.81

$$\int \left(x + \sqrt{-3 - 2x + 4x^2} \right) dx = \frac{1}{2} x^2 + \frac{1}{8} \sqrt{4x^2 - 2x - 3}(4x - 1) + \frac{13}{16} \log \left(\left| -4x + 2\sqrt{4x^2 - 2x - 3} + 1 \right| \right)$$

input `integrate(x+(4*x^2-2*x-3)^(1/2),x, algorithm="giac")`

output `1/2*x^2 + 1/8*sqrt(4*x^2 - 2*x - 3)*(4*x - 1) + 13/16*log(abs(-4*x + 2*sqrt(4*x^2 - 2*x - 3) + 1))`

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.75

$$\int \left(x + \sqrt{-3 - 2x + 4x^2} \right) dx = \left(\frac{x}{2} - \frac{1}{8} \right) \sqrt{4x^2 - 2x - 3} - \frac{13 \ln \left(2x + \sqrt{4x^2 - 2x - 3} - \frac{1}{2} \right)}{16} + \frac{x^2}{2}$$

input `int(x + (4*x^2 - 2*x - 3)^(1/2),x)`

output `(x/2 - 1/8)*(4*x^2 - 2*x - 3)^(1/2) - (13*log(2*x + (4*x^2 - 2*x - 3)^(1/2) - 1/2))/16 + x^2/2`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00

$$\int \left(x + \sqrt{-3 - 2x + 4x^2} \right) dx = \frac{\sqrt{4x^2 - 2x - 3} x}{2} - \frac{\sqrt{4x^2 - 2x - 3}}{8} - \frac{13 \log\left(\frac{2\sqrt{4x^2 - 2x - 3} + 4x - 1}{\sqrt{13}}\right)}{16} + \frac{x^2}{2}$$

input `int(x+(4*x^2-2*x-3)^(1/2),x)`output `(8*sqrt(4*x**2 - 2*x - 3)*x - 2*sqrt(4*x**2 - 2*x - 3) - 13*log((2*sqrt(4*x**2 - 2*x - 3) + 4*x - 1)/sqrt(13)) + 8*x**2)/16`

3.33 $\int \frac{1}{x + \sqrt{-3 - 2x + 4x^2}} dx$

Optimal result	279
Mathematica [A] (verified)	280
Rubi [A] (verified)	280
Maple [B] (verified)	282
Fricas [B] (verification not implemented)	283
Sympy [F]	284
Maxima [F]	284
Giac [B] (verification not implemented)	285
Mupad [F(-1)]	286
Reduce [B] (verification not implemented)	286

Optimal result

Integrand size = 18, antiderivative size = 106

$$\int \frac{1}{x + \sqrt{-3 - 2x + 4x^2}} dx = -\frac{1}{30} (10 - \sqrt{10}) \log (1 - \sqrt{10} - 3(2x + \sqrt{-3 - 2x + 4x^2})) - \frac{1}{30} (10 + \sqrt{10}) \log (1 + \sqrt{10} - 3(2x + \sqrt{-3 - 2x + 4x^2})) + \log (1 - 2(2x + \sqrt{-3 - 2x + 4x^2}))$$

output

```
-1/30*(10-10^(1/2))*ln(1-10^(1/2)-6*x-3*(4*x^2-2*x-3)^(1/2))-1/30*(10+10^(1/2))*ln(1+10^(1/2)-6*x-3*(4*x^2-2*x-3)^(1/2))+ln(1-4*x-2*(4*x^2-2*x-3)^(1/2))
```


Mathematica [A] (verified)

Time = 0.41 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.92

$$\int \frac{1}{x + \sqrt{-3 - 2x + 4x^2}} dx = \frac{1}{30} \left((-10 + \sqrt{10}) \log(-1 + \sqrt{10} + 2x - \sqrt{-3 - 2x + 4x^2}) \right. \\ \left. - (10 + \sqrt{10}) \log(1 + \sqrt{10} - 2x + \sqrt{-3 - 2x + 4x^2}) \right. \\ \left. - 10 \log(1 - 4x + 2\sqrt{-3 - 2x + 4x^2}) \right)$$

input

```
Integrate[(x + Sqrt[-3 - 2*x + 4*x^2])^(-1), x]
```

output

```
((-10 + Sqrt[10])*Log[-1 + Sqrt[10] + 2*x - Sqrt[-3 - 2*x + 4*x^2]] - (10 + Sqrt[10])*Log[1 + Sqrt[10] - 2*x + Sqrt[-3 - 2*x + 4*x^2]] - 10*Log[1 - 4*x + 2*Sqrt[-3 - 2*x + 4*x^2]])/30
```

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.06, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {7286, 25, 2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{4x^2 - 2x - 3} + x} dx$$

↓ 7286

$$2 \int -\frac{-\left(2x + \sqrt{4x^2 - 2x - 3}\right)^2 + 2x + \sqrt{4x^2 - 2x - 3} + 3}{6\left(2x + \sqrt{4x^2 - 2x - 3}\right)^3 - 7\left(2x + \sqrt{4x^2 - 2x - 3}\right)^2 - 4\left(2x + \sqrt{4x^2 - 2x - 3}\right) + 3} d\left(2x + \sqrt{4x^2 - 2x - 3}\right)$$

↓ 25

$$-2 \int \frac{-(2x + \sqrt{4x^2 - 2x - 3})^2 + 2x + \sqrt{4x^2 - 2x - 3} + 3}{6(2x + \sqrt{4x^2 - 2x - 3})^3 - 7(2x + \sqrt{4x^2 - 2x - 3})^2 - 4(2x + \sqrt{4x^2 - 2x - 3}) + 3} d(2x + \sqrt{4x^2 - 2x - 3})$$

↓ 2462

$$-2 \int \left(\frac{2x + \sqrt{4x^2 - 2x - 3}}{3(2x + \sqrt{4x^2 - 2x - 3})^2 - 2(2x + \sqrt{4x^2 - 2x - 3}) - 3} + \frac{1}{1 - 2(2x + \sqrt{4x^2 - 2x - 3})} \right) d(2x + \sqrt{4x^2 - 2x - 3})$$

↓ 2009

$$2 \left(-\frac{1}{60} (10 - \sqrt{10}) \log \left(-3(\sqrt{4x^2 - 2x - 3} + 2x) - \sqrt{10} + 1 \right) - \frac{1}{60} (10 + \sqrt{10}) \log \left(-3(\sqrt{4x^2 - 2x - 3} + 2x) \right) \right)$$

input `Int[(x + Sqrt[-3 - 2*x + 4*x^2])^(-1), x]`

output `2*(-1/60*((10 - Sqrt[10])*Log[1 - Sqrt[10] - 3*(2*x + Sqrt[-3 - 2*x + 4*x^2])]) - ((10 + Sqrt[10])*Log[1 + Sqrt[10] - 3*(2*x + Sqrt[-3 - 2*x + 4*x^2])]))/60 + Log[1 - 2*(2*x + Sqrt[-3 - 2*x + 4*x^2])]/2)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2462 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegr and[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ [Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0] && RationalFunctionQ[u, x]`

rule 7286

```
Int[u_, x_Symbol] := With[{lst = FunctionOfSquareRootOfQuadratic[u, x]}, Si
mp[2 Subst[Int[lst[[1]], x], x, lst[[2]]], x] /; !FalseQ[lst] && EqQ[lst
[[3]], 2] /; EulerIntegrandQ[u, x]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 426 vs. 2(85) = 170.

Time = 0.37 (sec) , antiderivative size = 427, normalized size of antiderivative = 4.03

method	result
default	$3\sqrt{10} \left(-\frac{\sqrt{36\left(x-\frac{1}{3}-\frac{\sqrt{10}}{3}\right)^2+9\left(\frac{2}{3}+\frac{8\sqrt{10}}{3}\right)\left(x-\frac{1}{3}-\frac{\sqrt{10}}{3}\right)+11+2\sqrt{10}}}{9} - \frac{\left(\frac{2}{3}+\frac{8\sqrt{10}}{3}\right) \ln\left(\frac{(4x-1)\sqrt{4}+\sqrt{4\left(x-\frac{1}{3}-\frac{\sqrt{10}}{3}\right)^2+\left(\frac{2}{3}+\frac{8\sqrt{10}}{3}\right)}(x-\frac{1}{3}-\frac{\sqrt{10}}{3})\right)}{24} \right)$
trager	$-\frac{\ln\left(7009920 \operatorname{RootOf}\left(30_Z^2+20_Z+3\right)^2 \sqrt{4x^2-2x-3} x^2-1252800 \operatorname{RootOf}\left(30_Z^2+20_Z+3\right)^2 x^3+5257440 \operatorname{RootOf}\left(30_Z^2+20_Z+3\right)^2 x^2-1252800 \operatorname{RootOf}\left(30_Z^2+20_Z+3\right)^2 x+5257440 \operatorname{RootOf}\left(30_Z^2+20_Z+3\right)^2\right)}{20}$

input

```
int(1/(x+(4*x^2-2*x-3)^(1/2)),x,method=_RETURNVERBOSE)
```

output

```
-3/20*10^(1/2)*(-1/9*(36*(x-1/3-1/3*10^(1/2))^2+9*(2/3+8/3*10^(1/2))*(x-1/3-1/3*10^(1/2))+11+2*10^(1/2))^(1/2)-1/24*(2/3+8/3*10^(1/2))*ln(1/4*(4*x-1)*4^(1/2)+(4*(x-1/3-1/3*10^(1/2))^2+(2/3+8/3*10^(1/2))*(x-1/3-1/3*10^(1/2))+11/9+2/9*10^(1/2))^(1/2))*4^(1/2)+1/3*(11/9+2/9*10^(1/2))/(1/3+1/3*10^(1/2))*arctanh(3/2*(22/9+4/9*10^(1/2)+(2/3+8/3*10^(1/2))*(x-1/3-1/3*10^(1/2)))/(1/3+1/3*10^(1/2)))/(36*(x-1/3-1/3*10^(1/2))^2+9*(2/3+8/3*10^(1/2))*(x-1/3-1/3*10^(1/2))+11+2*10^(1/2))^(1/2))-3/20*10^(1/2)*(1/9*(36*(x-1/3+1/3*10^(1/2))^2+9*(2/3-8/3*10^(1/2))*(x-1/3+1/3*10^(1/2))+11-2*10^(1/2))^(1/2)+1/24*(2/3-8/3*10^(1/2))*ln(1/4*(4*x-1)*4^(1/2)+(4*(x-1/3+1/3*10^(1/2))^2+(2/3-8/3*10^(1/2))*(x-1/3+1/3*10^(1/2))+11/9-2/9*10^(1/2))^(1/2))*4^(1/2)-1/3*(11/9-2/9*10^(1/2))/(-1/3+1/3*10^(1/2))*arctanh(3/2*(22/9-4/9*10^(1/2)+(2/3-8/3*10^(1/2))*(x-1/3+1/3*10^(1/2)))/(-1/3+1/3*10^(1/2)))/(36*(x-1/3+1/3*10^(1/2))^2+9*(2/3-8/3*10^(1/2))*(x-1/3+1/3*10^(1/2))+11-2*10^(1/2))^(1/2))-1/6*ln(3*x^2-2*x-3)+1/30*10^(1/2)*arctanh(1/20*(6*x-2)*10^(1/2))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 281 vs. $2(83) = 166$.

Time = 0.08 (sec) , antiderivative size = 281, normalized size of antiderivative = 2.65

$$\int \frac{1}{x + \sqrt{-3 - 2x + 4x^2}} dx = \frac{1}{60} \sqrt{10} \log \left(\frac{9x^2 + 2\sqrt{10}(3x - 1) - 6x + 11}{3x^2 - 2x - 3} \right) \\ + \frac{1}{60} \sqrt{10} \log \left(-\frac{3x^2 - \sqrt{10}(6x^2 + 17x - 3) - \sqrt{4x^2 - 2x - 3}(\sqrt{10}(3x + 19) - 60x + 10) + 28x + 57}{3x^2 - 2x - 3} \right) \\ + \frac{1}{60} \sqrt{10} \log \left(-\frac{x^2 - \sqrt{10}(2x^2 + 5x - 3) - \sqrt{4x^2 - 2x - 3}(\sqrt{10}(x - 7) - 20x + 10) - 4x - 21}{3x^2 - 2x - 3} \right) \\ + \frac{1}{6} \log \left(12x^2 - \sqrt{4x^2 - 2x - 3}(6x - 1) - 5x - 6 \right) \\ - \frac{1}{6} \log \left(4x^2 - \sqrt{4x^2 - 2x - 3}(2x - 1) - 3x - 6 \right) \\ - \frac{1}{6} \log \left(3x^2 - 2x - 3 \right) - \frac{2}{3} \log \left(-4x + 2\sqrt{4x^2 - 2x - 3} + 1 \right)$$

input `integrate(1/(x+(4*x^2-2*x-3)^(1/2)),x, algorithm="fricas")`

output `1/60*sqrt(10)*log((9*x^2 + 2*sqrt(10)*(3*x - 1) - 6*x + 11)/(3*x^2 - 2*x - 3)) + 1/60*sqrt(10)*log(-(3*x^2 - sqrt(10)*(6*x^2 + 17*x - 3) - sqrt(4*x^2 - 2*x - 3)*(sqrt(10)*(3*x + 19) - 60*x + 10) + 28*x + 57)/(3*x^2 - 2*x - 3)) + 1/60*sqrt(10)*log(-(x^2 - sqrt(10)*(2*x^2 + 5*x - 3) - sqrt(4*x^2 - 2*x - 3)*(sqrt(10)*(x - 7) - 20*x + 10) - 4*x - 21)/(3*x^2 - 2*x - 3)) + 1/6*log(12*x^2 - sqrt(4*x^2 - 2*x - 3)*(6*x - 1) - 5*x - 6) - 1/6*log(4*x^2 - sqrt(4*x^2 - 2*x - 3)*(2*x - 1) - 3*x - 6) - 1/6*log(3*x^2 - 2*x - 3) - 2/3*log(-4*x + 2*sqrt(4*x^2 - 2*x - 3) + 1)`

Sympy [F]

$$\int \frac{1}{x + \sqrt{-3 - 2x + 4x^2}} dx = \int \frac{1}{x + \sqrt{4x^2 - 2x - 3}} dx$$

input `integrate(1/(x+(4*x**2-2*x-3)**(1/2)),x)`

output `Integral(1/(x + sqrt(4*x**2 - 2*x - 3)), x)`

Maxima [F]

$$\int \frac{1}{x + \sqrt{-3 - 2x + 4x^2}} dx = \int \frac{1}{x + \sqrt{4x^2 - 2x - 3}} dx$$

input `integrate(1/(x+(4*x^2-2*x-3)^(1/2)),x, algorithm="maxima")`

output `integrate(1/(x + sqrt(4*x^2 - 2*x - 3)), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 275 vs. $2(83) = 166$.

Time = 0.16 (sec) , antiderivative size = 275, normalized size of antiderivative = 2.59

$$\begin{aligned} & \int \frac{1}{x + \sqrt{-3 - 2x + 4x^2}} dx \\ &= -\frac{1}{60} \sqrt{10} \log \left(\frac{|6x - 2\sqrt{10} - 2|}{|6x + 2\sqrt{10} - 2|} \right) \\ &+ \frac{1}{60} \sqrt{10} \log \left(\frac{|-4x - 2\sqrt{10} + 2\sqrt{4x^2 - 2x - 3} + 2|}{|-4x + 2\sqrt{10} + 2\sqrt{4x^2 - 2x - 3} + 2|} \right) \\ &- \frac{1}{60} \sqrt{10} \log \left(\frac{|-12x - 2\sqrt{10} + 6\sqrt{4x^2 - 2x - 3} + 2|}{|-12x + 2\sqrt{10} + 6\sqrt{4x^2 - 2x - 3} + 2|} \right) \\ &+ \frac{1}{6} \log \left(\left| 3 \left(2x - \sqrt{4x^2 - 2x - 3} \right)^2 - 4x + 2\sqrt{4x^2 - 2x - 3} - 3 \right| \right) \\ &- \frac{1}{6} \log \left(\left| \left(2x - \sqrt{4x^2 - 2x - 3} \right)^2 - 4x + 2\sqrt{4x^2 - 2x - 3} - 9 \right| \right) \\ &- \frac{1}{6} \log (|3x^2 - 2x - 3|) - \frac{2}{3} \log \left(\left| -4x + 2\sqrt{4x^2 - 2x - 3} + 1 \right| \right) \end{aligned}$$

input `integrate(1/(x+(4*x^2-2*x-3)^(1/2)),x, algorithm="giac")`

output

```
-1/60*sqrt(10)*log(abs(6*x - 2*sqrt(10) - 2)/abs(6*x + 2*sqrt(10) - 2)) +
1/60*sqrt(10)*log(abs(-4*x - 2*sqrt(10) + 2*sqrt(4*x^2 - 2*x - 3) + 2)/abs
(-4*x + 2*sqrt(10) + 2*sqrt(4*x^2 - 2*x - 3) + 2)) - 1/60*sqrt(10)*log(abs
(-12*x - 2*sqrt(10) + 6*sqrt(4*x^2 - 2*x - 3) + 2)/abs(-12*x + 2*sqrt(10)
+ 6*sqrt(4*x^2 - 2*x - 3) + 2)) + 1/6*log(abs(3*(2*x - sqrt(4*x^2 - 2*x -
3))^2 - 4*x + 2*sqrt(4*x^2 - 2*x - 3) - 3)) - 1/6*log(abs((2*x - sqrt(4*x^
2 - 2*x - 3))^2 - 4*x + 2*sqrt(4*x^2 - 2*x - 3) - 9)) - 1/6*log(abs(3*x^2
- 2*x - 3)) - 2/3*log(abs(-4*x + 2*sqrt(4*x^2 - 2*x - 3) + 1))
```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{x + \sqrt{-3 - 2x + 4x^2}} dx = \int \frac{1}{x + \sqrt{4x^2 - 2x - 3}} dx$$

input `int(1/(x + (4*x^2 - 2*x - 3)^(1/2)), x)`output `int(1/(x + (4*x^2 - 2*x - 3)^(1/2)), x)`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 239, normalized size of antiderivative = 2.25

$$\begin{aligned} & \int \frac{1}{x + \sqrt{-3 - 2x + 4x^2}} dx \\ &= -\frac{\sqrt{10} \operatorname{atan}\left(\frac{6\sqrt{4x^2 - 2x - 3}i + 12ix - 3i}{2\sqrt{10} - 1}\right) i}{30} - \frac{\operatorname{atan}\left(\frac{6\sqrt{4x^2 - 2x - 3}i + 12ix - 3i}{2\sqrt{10} - 1}\right) i}{3} \\ & \quad - \frac{\sqrt{10} \log(144\sqrt{4x^2 - 2x - 3}x - 36\sqrt{4x^2 - 2x - 3} + 4\sqrt{10} + 288x^2 - 144x - 140)}{60} \\ & \quad + \frac{\sqrt{10} \log(6\sqrt{4x^2 - 2x - 3} + 2\sqrt{10} + 12x - 2)}{30} \\ & \quad - \frac{\log(144\sqrt{4x^2 - 2x - 3}x - 36\sqrt{4x^2 - 2x - 3} + 4\sqrt{10} + 288x^2 - 144x - 140)}{6} \\ & \quad - \frac{\log(6\sqrt{4x^2 - 2x - 3} + 2\sqrt{10} + 12x - 2)}{3} + \log\left(\frac{2\sqrt{4x^2 - 2x - 3} + 4x - 1}{\sqrt{13}}\right) \end{aligned}$$

input `int(1/(x+(4*x^2-2*x-3)^(1/2)), x)`

output

```
( - 2*sqrt(10)*atan((6*sqrt(4*x**2 - 2*x - 3)*i + 12*i*x - 3*i)/(2*sqrt(10)
) - 1))*i - 20*atan((6*sqrt(4*x**2 - 2*x - 3)*i + 12*i*x - 3*i)/(2*sqrt(10)
) - 1))*i - sqrt(10)*log(144*sqrt(4*x**2 - 2*x - 3)*x - 36*sqrt(4*x**2 - 2
*x - 3) + 4*sqrt(10) + 288*x**2 - 144*x - 140) + 2*sqrt(10)*log(6*sqrt(4*x
**2 - 2*x - 3) + 2*sqrt(10) + 12*x - 2) - 10*log(144*sqrt(4*x**2 - 2*x - 3
)*x - 36*sqrt(4*x**2 - 2*x - 3) + 4*sqrt(10) + 288*x**2 - 144*x - 140) - 2
0*log(6*sqrt(4*x**2 - 2*x - 3) + 2*sqrt(10) + 12*x - 2) + 60*log((2*sqrt(4
*x**2 - 2*x - 3) + 4*x - 1)/sqrt(13)))/60
```


3.34
$$\int \frac{1}{\left(x + \sqrt{-3 - 2x + 4x^2}\right)^2} dx$$

Optimal result	288
Mathematica [A] (verified)	289
Rubi [A] (verified)	289
Maple [C] (verified)	291
Fricas [A] (verification not implemented)	292
Sympy [F]	292
Maxima [F]	293
Giac [B] (verification not implemented)	293
Mupad [F(-1)]	294
Reduce [B] (verification not implemented)	294

Optimal result

Integrand size = 18, antiderivative size = 149

$$\begin{aligned} & \int \frac{1}{\left(x + \sqrt{-3 - 2x + 4x^2}\right)^2} dx \\ &= \frac{3 - 19(2x + \sqrt{-3 - 2x + 4x^2})}{15 \left(3 + 2(2x + \sqrt{-3 - 2x + 4x^2}) - 3(2x + \sqrt{-3 - 2x + 4x^2})^2\right)} \\ & \quad - \frac{13 \log(1 - \sqrt{10} - 3(2x + \sqrt{-3 - 2x + 4x^2}))}{10\sqrt{10}} \\ & \quad + \frac{13 \log(1 + \sqrt{10} - 3(2x + \sqrt{-3 - 2x + 4x^2}))}{10\sqrt{10}} \end{aligned}$$

output

```
(3-38*x-19*(4*x^2-2*x-3)^(1/2))/(45+60*x+30*(4*x^2-2*x-3)^(1/2)-45*(2*x+(4*x^2-2*x-3)^(1/2))^2)-13/100*ln(1-10^(1/2)-6*x-3*(4*x^2-2*x-3)^(1/2))*10^(1/2)+13/100*ln(1+10^(1/2)-6*x-3*(4*x^2-2*x-3)^(1/2))*10^(1/2)
```

Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.58

$$\int \frac{1}{(x + \sqrt{-3 - 2x + 4x^2})^2} dx = \frac{3 + 11x - 9\sqrt{-3 - 2x + 4x^2} - 3x\sqrt{-3 - 2x + 4x^2}}{90 + 60x - 90x^2} + \frac{13\operatorname{arctanh}\left(\frac{1-2x+\sqrt{-3-2x+4x^2}}{\sqrt{10}}\right)}{5\sqrt{10}}$$

input

```
Integrate[(x + Sqrt[-3 - 2*x + 4*x^2])^(-2), x]
```

output

```
(3 + 11*x - 9*Sqrt[-3 - 2*x + 4*x^2] - 3*x*Sqrt[-3 - 2*x + 4*x^2])/(90 + 60*x - 90*x^2) + (13*ArcTanh[(1 - 2*x + Sqrt[-3 - 2*x + 4*x^2])/Sqrt[10]])/(5*Sqrt[10])
```

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.05, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {7286, 27, 2191, 27, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(\sqrt{4x^2 - 2x - 3} + x)^2} dx$$

↓ 7286

$$2 \int -\frac{2\left(-\left(2x + \sqrt{4x^2 - 2x - 3}\right)^2 + 2x + \sqrt{4x^2 - 2x - 3} + 3\right)}{\left(-3\left(2x + \sqrt{4x^2 - 2x - 3}\right)^2 + 2\left(2x + \sqrt{4x^2 - 2x - 3}\right) + 3\right)^2} d\left(2x + \sqrt{4x^2 - 2x - 3}\right)$$

↓ 27

$$-4 \int \frac{-(2x + \sqrt{4x^2 - 2x - 3})^2 + 2x + \sqrt{4x^2 - 2x - 3} + 3}{(-3(2x + \sqrt{4x^2 - 2x - 3})^2 + 2(2x + \sqrt{4x^2 - 2x - 3}) + 3)^2} d(2x + \sqrt{4x^2 - 2x - 3})$$

↓ 2191

$$-4 \left(-\frac{1}{40} \int \frac{26}{-3(2x + \sqrt{4x^2 - 2x - 3})^2 + 2(2x + \sqrt{4x^2 - 2x - 3}) + 3} d(2x + \sqrt{4x^2 - 2x - 3}) - \frac{1}{60} \int \frac{1}{-3(2x + \sqrt{4x^2 - 2x - 3})^2 + 2(2x + \sqrt{4x^2 - 2x - 3}) + 3} d(2x + \sqrt{4x^2 - 2x - 3}) \right)$$

↓ 27

$$-4 \left(\frac{13}{20} \int \frac{1}{-3(2x + \sqrt{4x^2 - 2x - 3})^2 + 2(2x + \sqrt{4x^2 - 2x - 3}) + 3} d(2x + \sqrt{4x^2 - 2x - 3}) - \frac{1}{60} \int \frac{1}{-3(2x + \sqrt{4x^2 - 2x - 3})^2 + 2(2x + \sqrt{4x^2 - 2x - 3}) + 3} d(2x + \sqrt{4x^2 - 2x - 3}) \right)$$

↓ 1081

$$-4 \left(-\frac{39}{20} \int \left(\frac{1}{2\sqrt{10}(-3(2x + \sqrt{4x^2 - 2x - 3}) - \sqrt{10} + 1)} - \frac{1}{2\sqrt{10}(-3(2x + \sqrt{4x^2 - 2x - 3}) + \sqrt{10} + 1)} \right) d(2x + \sqrt{4x^2 - 2x - 3}) \right)$$

↓ 2009

$$-4 \left(-\frac{3 - 19(\sqrt{4x^2 - 2x - 3} + 2x)}{60(-3(\sqrt{4x^2 - 2x - 3} + 2x)^2 + 2(\sqrt{4x^2 - 2x - 3} + 2x) + 3)} - \frac{39}{20} \left(\frac{\log(-3(\sqrt{4x^2 - 2x - 3} + 2x))}{6\sqrt{10}} \right) \right)$$

input `Int[(x + Sqrt[-3 - 2*x + 4*x^2])^(-2), x]`

output `-4*(-1/60*(3 - 19*(2*x + Sqrt[-3 - 2*x + 4*x^2]))/(3 + 2*(2*x + Sqrt[-3 - 2*x + 4*x^2]) - 3*(2*x + Sqrt[-3 - 2*x + 4*x^2])^2) - (39*(-1/6*Log[1 - Sqrt[10] - 3*(2*x + Sqrt[-3 - 2*x + 4*x^2]])/Sqrt[10] + Log[1 + Sqrt[10] - 3*(2*x + Sqrt[-3 - 2*x + 4*x^2]])]/(6*Sqrt[10])))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 1081 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c Int[ExpandIntegrand[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x]] /; FreeQ[{a, b, c}, x] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2191 `Int[(Pq_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := With[{Q = PolynomialQuotient[Pq, a + b*x + c*x^2, x], f = Coeff[PolynomialRemainder[Pq, a + b*x + c*x^2, x], x, 0], g = Coeff[PolynomialRemainder[Pq, a + b*x + c*x^2, x], x, 1]}, Simp[(b*f - 2*a*g + (2*c*f - b*g)*x)*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c))), x] + Simp[1/((p + 1)*(b^2 - 4*a*c)) Int[(a + b*x + c*x^2)^(p + 1)*ExpandToSum[(p + 1)*(b^2 - 4*a*c)*Q - (2*p + 3)*(2*c*f - b*g), x], x], x]] /; FreeQ[{a, b, c}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1]`

rule 7286 `Int[u_, x_Symbol] := With[{lst = FunctionOfSquareRootOfQuadratic[u, x]}, Simp[2 Subst[Int[lst[[1]], x], x, lst[[2]]], x] /; !FalseQ[lst] && EqQ[lst[[3]], 2]] /; EulerIntegrandQ[u, x]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.16 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.70

method	result
trager	$-\frac{(3+x)x}{10(3x^2-2x-3)} + \frac{(3+x)\sqrt{4x^2-2x-3}}{30x^2-20x-30} - \frac{13\operatorname{RootOf}(_Z^2-10) \ln\left(\frac{\operatorname{RootOf}(_Z^2-10)^{x+3} \operatorname{RootOf}(_Z^2-10)^{-10}\sqrt{4x^2-2x-3}}{\operatorname{RootOf}(_Z^2-10)^{x-x-3}}\right)}{100}$
default	Expression too large to display

input `int(1/(x+(4*x^2-2*x-3)^(1/2))^2,x,method=_RETURNVERBOSE)`

output `-1/10*(3+x)*x/(3*x^2-2*x-3)+1/10*(3+x)/(3*x^2-2*x-3)*(4*x^2-2*x-3)^(1/2)-13/100*RootOf(_Z^2-10)*ln((RootOf(_Z^2-10)*x+3*RootOf(_Z^2-10)-10*(4*x^2-2*x-3)^(1/2))/(RootOf(_Z^2-10)*x-x-3))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.15

$$\int \frac{1}{(x + \sqrt{-3 - 2x + 4x^2})^2} dx$$

$$= \frac{13\sqrt{10}(3x^2 - 2x - 3) \log\left(\frac{1681x^2 + 4\sqrt{10}(41x^2 - 14x - 21) + 2\sqrt{4x^2 - 2x - 3}(41\sqrt{10}(x+3) + 40x + 120) - 574x - 861}{3x^2 - 2x - 3}\right) + 13\sqrt{10}}{200(3x^2 - 2x - 3)}$$

input `integrate(1/(x+(4*x^2-2*x-3)^(1/2))^2,x, algorithm="fricas")`

output `1/200*(13*sqrt(10)*(3*x^2 - 2*x - 3)*log((1681*x^2 + 4*sqrt(10)*(41*x^2 - 14*x - 21) + 2*sqrt(4*x^2 - 2*x - 3)*(41*sqrt(10)*(x + 3) + 40*x + 120) - 574*x - 861)/(3*x^2 - 2*x - 3)) + 13*sqrt(10)*(3*x^2 - 2*x - 3)*log((9*x^2 - 2*sqrt(10)*(3*x - 1) - 6*x + 11)/(3*x^2 - 2*x - 3)) + 40*x^2 + 20*sqrt(4*x^2 - 2*x - 3)*(x + 3) - 100*x - 60)/(3*x^2 - 2*x - 3)`

Sympy [F]

$$\int \frac{1}{(x + \sqrt{-3 - 2x + 4x^2})^2} dx = \int \frac{1}{(x + \sqrt{4x^2 - 2x - 3})^2} dx$$

input `integrate(1/(x+(4*x**2-2*x-3)**(1/2))**2,x)`

output `Integral((x + sqrt(4*x**2 - 2*x - 3))**(-2), x)`

Maxima [F]

$$\int \frac{1}{(x + \sqrt{-3 - 2x + 4x^2})^2} dx = \int \frac{1}{(x + \sqrt{4x^2 - 2x - 3})^2} dx$$

input `integrate(1/(x+(4*x^2-2*x-3)^(1/2))^2,x, algorithm="maxima")`

output `integrate((x + sqrt(4*x^2 - 2*x - 3))^(-2), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 320 vs. 2(121) = 242.

Time = 0.17 (sec) , antiderivative size = 320, normalized size of antiderivative = 2.15

$$\begin{aligned} \int \frac{1}{(x + \sqrt{-3 - 2x + 4x^2})^2} dx &= \frac{13}{200} \sqrt{10} \log \left(\frac{|6x - 2\sqrt{10} - 2|}{|6x + 2\sqrt{10} - 2|} \right) \\ &- \frac{13}{200} \sqrt{10} \log \left(\frac{|-4x - 2\sqrt{10} + 2\sqrt{4x^2 - 2x - 3} + 2|}{|-4x + 2\sqrt{10} + 2\sqrt{4x^2 - 2x - 3} + 2|} \right) \\ &+ \frac{13}{200} \sqrt{10} \log \left(\frac{|-12x - 2\sqrt{10} + 6\sqrt{4x^2 - 2x - 3} + 2|}{|-12x + 2\sqrt{10} + 6\sqrt{4x^2 - 2x - 3} + 2|} \right) \\ &- \frac{41(2x - \sqrt{4x^2 - 2x - 3})^3 - 28(2x - \sqrt{4x^2 - 2x - 3})^2 - 246x + 123\sqrt{4x^2 - 2x - 3}}{15(3(2x - \sqrt{4x^2 - 2x - 3})^4 - 8(2x - \sqrt{4x^2 - 2x - 3})^3 - 26(2x - \sqrt{4x^2 - 2x - 3})^2 + 48x - 2)} \\ &- \frac{11x + 3}{30(3x^2 - 2x - 3)} \end{aligned}$$

input `integrate(1/(x+(4*x^2-2*x-3)^(1/2))^2,x, algorithm="giac")`

output

```
13/200*sqrt(10)*log(abs(6*x - 2*sqrt(10) - 2)/abs(6*x + 2*sqrt(10) - 2)) -
13/200*sqrt(10)*log(abs(-4*x - 2*sqrt(10) + 2*sqrt(4*x^2 - 2*x - 3) + 2)/
abs(-4*x + 2*sqrt(10) + 2*sqrt(4*x^2 - 2*x - 3) + 2)) + 13/200*sqrt(10)*lo
g(abs(-12*x - 2*sqrt(10) + 6*sqrt(4*x^2 - 2*x - 3) + 2)/abs(-12*x + 2*sqrt
(10) + 6*sqrt(4*x^2 - 2*x - 3) + 2)) - 1/15*(41*(2*x - sqrt(4*x^2 - 2*x -
3))^3 - 28*(2*x - sqrt(4*x^2 - 2*x - 3))^2 - 246*x + 123*sqrt(4*x^2 - 2*x
- 3))/(3*(2*x - sqrt(4*x^2 - 2*x - 3))^4 - 8*(2*x - sqrt(4*x^2 - 2*x - 3))
^3 - 26*(2*x - sqrt(4*x^2 - 2*x - 3))^2 + 48*x - 24*sqrt(4*x^2 - 2*x - 3)
+ 27) - 1/30*(11*x + 3)/(3*x^2 - 2*x - 3)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(x + \sqrt{-3 - 2x + 4x^2})^2} dx = \int \frac{1}{(x + \sqrt{4x^2 - 2x - 3})^2} dx$$

input

```
int(1/(x + (4*x^2 - 2*x - 3)^(1/2))^2, x)
```

output

```
int(1/(x + (4*x^2 - 2*x - 3)^(1/2))^2, x)
```

Reduce [B] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 648, normalized size of antiderivative = 4.35

$$\int \frac{1}{(x + \sqrt{-3 - 2x + 4x^2})^2} dx = \text{Too large to display}$$

input

```
int(1/(x+(4*x^2-2*x-3)^(1/2))^2, x)
```

output

```
( - 642798*sqrt(10)*atan((2*sqrt(4*x**2 - 2*x - 3)*i + 4*i*x - i)/(2*sqrt(10) - 1))*i*x**2 + 428532*sqrt(10)*atan((2*sqrt(4*x**2 - 2*x - 3)*i + 4*i*x - i)/(2*sqrt(10) - 1))*i*x + 642798*sqrt(10)*atan((2*sqrt(4*x**2 - 2*x - 3)*i + 4*i*x - i)/(2*sqrt(10) - 1))*i + 329640*sqrt(4*x**2 - 2*x - 3)*x + 988920*sqrt(4*x**2 - 2*x - 3) + 642798*sqrt(10)*log(- sqrt(10) + 3*x - 1)*x**2 - 428532*sqrt(10)*log(- sqrt(10) + 3*x - 1)*x - 642798*sqrt(10)*log(- sqrt(10) + 3*x - 1) - 642798*sqrt(10)*log(sqrt(10) + 3*x - 1)*x**2 + 428532*sqrt(10)*log(sqrt(10) + 3*x - 1)*x + 642798*sqrt(10)*log(sqrt(10) + 3*x - 1) + 321399*sqrt(10)*log(16*sqrt(4*x**2 - 2*x - 3)*x - 4*sqrt(4*x**2 - 2*x - 3) + 4*sqrt(10) + 32*x**2 - 16*x - 52)*x**2 - 214266*sqrt(10)*log(16*sqrt(4*x**2 - 2*x - 3)*x - 4*sqrt(4*x**2 - 2*x - 3) + 4*sqrt(10) + 32*x**2 - 16*x - 52)*x - 321399*sqrt(10)*log(16*sqrt(4*x**2 - 2*x - 3)*x - 4*sqrt(4*x**2 - 2*x - 3) + 4*sqrt(10) + 32*x**2 - 16*x - 52) + 642798*sqrt(10)*log(6*sqrt(4*x**2 - 2*x - 3) - 2*sqrt(10) + 12*x - 2)*x**2 - 428532*sqrt(10)*log(6*sqrt(4*x**2 - 2*x - 3) - 2*sqrt(10) + 12*x - 2)*x - 642798*sqrt(10)*log(6*sqrt(4*x**2 - 2*x - 3) - 2*sqrt(10) + 12*x - 2) - 642798*sqrt(10)*log(6*sqrt(4*x**2 - 2*x - 3) + 2*sqrt(10) + 12*x - 2)*x**2 + 428532*sqrt(10)*log(6*sqrt(4*x**2 - 2*x - 3) + 2*sqrt(10) + 12*x - 2)*x + 642798*sqrt(10)*log(6*sqrt(4*x**2 - 2*x - 3) + 2*sqrt(10) + 12*x - 2) - 642798*sqrt(10)*log(2*sqrt(4*x**2 - 2*x - 3) - 2*sqrt(10) + 4*x - 2)*x**2 + 42853...
```


3.35 $\int \frac{1}{\left(x + \sqrt{-3 - 2x + 4x^2}\right)^3} dx$

Optimal result	296
Mathematica [A] (verified)	297
Rubi [A] (verified)	297
Maple [C] (verified)	300
Fricas [A] (verification not implemented)	301
Sympy [F]	301
Maxima [F]	302
Giac [B] (verification not implemented)	302
Mupad [F(-1)]	303
Reduce [B] (verification not implemented)	303

Optimal result

Integrand size = 18, antiderivative size = 218

$$\int \frac{1}{\left(x + \sqrt{-3 - 2x + 4x^2}\right)^3} dx$$

$$= \frac{123 + 2x + \sqrt{-3 - 2x + 4x^2}}{45 \left(3 + 2(2x + \sqrt{-3 - 2x + 4x^2}) - 3(2x + \sqrt{-3 - 2x + 4x^2})^2\right)^2}$$

$$+ \frac{917 - 351(2x + \sqrt{-3 - 2x + 4x^2})}{900 \left(3 + 2(2x + \sqrt{-3 - 2x + 4x^2}) - 3(2x + \sqrt{-3 - 2x + 4x^2})^2\right)}$$

$$- \frac{39 \log(1 - \sqrt{10} - 3(2x + \sqrt{-3 - 2x + 4x^2}))}{200\sqrt{10}}$$

$$+ \frac{39 \log(1 + \sqrt{10} - 3(2x + \sqrt{-3 - 2x + 4x^2}))}{200\sqrt{10}}$$

output

```
1/45*(123+2*x+(4*x^2-2*x-3)^(1/2))/(3+4*x+2*(4*x^2-2*x-3)^(1/2)-3*(2*x+(4*x^2-2*x-3)^(1/2))^2)+(917-702*x-351*(4*x^2-2*x-3)^(1/2))/(2700+3600*x+1800*(4*x^2-2*x-3)^(1/2))-2700*(2*x+(4*x^2-2*x-3)^(1/2))^2-39/2000*ln(1-10^(1/2)-6*x-3*(4*x^2-2*x-3)^(1/2))*10^(1/2)+39/2000*ln(1+10^(1/2)-6*x-3*(4*x^2-2*x-3)^(1/2))*10^(1/2)
```

Mathematica [A] (verified)

Time = 0.60 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.50

$$\int \frac{1}{(x + \sqrt{-3 - 2x + 4x^2})^3} dx$$

$$= \frac{15\sqrt{-3-2x+4x^2}(-87+113x+221x^2-231x^3)}{(3+2x-3x^2)^2} + \frac{5(-963-933x+949x^2+351x^3)}{(3+2x-3x^2)^2} + 117\sqrt{10}\operatorname{arctanh}\left(\frac{1-2x+\sqrt{-3-2x+4x^2}}{\sqrt{10}}\right)$$

$$\frac{\hspace{10em}}{3000}$$

input `Integrate[(x + Sqrt[-3 - 2*x + 4*x^2])^(-3), x]`

output `((15*Sqrt[-3 - 2*x + 4*x^2]*(-87 + 113*x + 221*x^2 - 231*x^3))/(3 + 2*x - 3*x^2)^2 + (5*(-963 - 933*x + 949*x^2 + 351*x^3))/(3 + 2*x - 3*x^2)^2 + 117*Sqrt[10]*ArcTanh[(1 - 2*x + Sqrt[-3 - 2*x + 4*x^2])/Sqrt[10]])/3000`

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.06, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {7286, 27, 2191, 27, 1159, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(\sqrt{4x^2 - 2x - 3} + x)^3} dx$$

$$\downarrow \text{7286}$$

$$2 \int -\frac{4(1 - 2(2x + \sqrt{4x^2 - 2x - 3})) \left(-(2x + \sqrt{4x^2 - 2x - 3})^2 + 2x + \sqrt{4x^2 - 2x - 3} + 3 \right)}{\left(-3(2x + \sqrt{4x^2 - 2x - 3})^2 + 2(2x + \sqrt{4x^2 - 2x - 3}) + 3 \right)^3} d(2x + \sqrt{4x^2 - 2x - 3})$$

$$\downarrow \text{27}$$

$$-8 \int \frac{\left(1 - 2(2x + \sqrt{4x^2 - 2x - 3})\right) \left(- (2x + \sqrt{4x^2 - 2x - 3})^2 + 2x + \sqrt{4x^2 - 2x - 3} + 3\right)}{\left(-3(2x + \sqrt{4x^2 - 2x - 3})^2 + 2(2x + \sqrt{4x^2 - 2x - 3}) + 3\right)^3} d(2x + \sqrt{4x^2 - 2x - 3})$$

↓ 2191

$$-8 \left(-\frac{1}{80} \int -\frac{2(197 - 240(2x + \sqrt{4x^2 - 2x - 3}))}{9 \left(-3(2x + \sqrt{4x^2 - 2x - 3})^2 + 2(2x + \sqrt{4x^2 - 2x - 3}) + 3\right)^2} d(2x + \sqrt{4x^2 - 2x - 3}) - \frac{1}{360} \int \frac{1}{\left(-3(2x + \sqrt{4x^2 - 2x - 3})^2 + 2(2x + \sqrt{4x^2 - 2x - 3}) + 3\right)^2} d(2x + \sqrt{4x^2 - 2x - 3}) \right)$$

↓ 27

$$-8 \left(\frac{1}{360} \int \frac{197 - 240(2x + \sqrt{4x^2 - 2x - 3})}{\left(-3(2x + \sqrt{4x^2 - 2x - 3})^2 + 2(2x + \sqrt{4x^2 - 2x - 3}) + 3\right)^2} d(2x + \sqrt{4x^2 - 2x - 3}) - \frac{1}{360} \int \frac{1}{\left(-3(2x + \sqrt{4x^2 - 2x - 3})^2 + 2(2x + \sqrt{4x^2 - 2x - 3}) + 3\right)^2} d(2x + \sqrt{4x^2 - 2x - 3}) \right)$$

↓ 1159

$$-8 \left(\frac{1}{360} \left(\frac{351}{20} \int \frac{1}{-3(2x + \sqrt{4x^2 - 2x - 3})^2 + 2(2x + \sqrt{4x^2 - 2x - 3}) + 3} d(2x + \sqrt{4x^2 - 2x - 3}) - \frac{1}{20} \int \frac{1}{\left(-3(2x + \sqrt{4x^2 - 2x - 3})^2 + 2(2x + \sqrt{4x^2 - 2x - 3}) + 3\right)} d(2x + \sqrt{4x^2 - 2x - 3}) \right) \right)$$

↓ 1081

$$-8 \left(\frac{1}{360} \left(-\frac{1053}{20} \int \left(\frac{1}{2\sqrt{10}(-3(2x + \sqrt{4x^2 - 2x - 3}) - \sqrt{10} + 1)} - \frac{1}{2\sqrt{10}(-3(2x + \sqrt{4x^2 - 2x - 3}) + \sqrt{10} + 1)} \right) d(2x + \sqrt{4x^2 - 2x - 3}) \right) \right)$$

↓ 2009

$$-8 \left(\frac{1}{360} \left(-\frac{917 - 351(\sqrt{4x^2 - 2x - 3} + 2x)}{20 \left(-3(\sqrt{4x^2 - 2x - 3} + 2x)^2 + 2(\sqrt{4x^2 - 2x - 3} + 2x) + 3\right)} - \frac{1053}{20} \left(\frac{\log(-3(\sqrt{4x^2 - 2x - 3} + 2x))}{6\sqrt{10}} \right) \right) \right)$$

input `Int[(x + Sqrt[-3 - 2*x + 4*x^2])^(-3), x]`

output `-8*(-1/360*(123 + 2*x + Sqrt[-3 - 2*x + 4*x^2])/(3 + 2*(2*x + Sqrt[-3 - 2*x + 4*x^2]) - 3*(2*x + Sqrt[-3 - 2*x + 4*x^2])^2)^2 + (-1/20*(917 - 351*(2*x + Sqrt[-3 - 2*x + 4*x^2]))/(3 + 2*(2*x + Sqrt[-3 - 2*x + 4*x^2]) - 3*(2*x + Sqrt[-3 - 2*x + 4*x^2])^2) - (1053*(-1/6*Log[1 - Sqrt[10] - 3*(2*x + Sqrt[-3 - 2*x + 4*x^2])]/Sqrt[10] + Log[1 + Sqrt[10] - 3*(2*x + Sqrt[-3 - 2*x + 4*x^2])]/(6*Sqrt[10])))/20)/360)`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 1081 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c Int[ExpandIntegrand[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x] /; FreeQ[{a, b, c}, x] && NiceSqrtQ[b^2 - 4*a*c]`

rule 1159 `Int[((d_) + (e_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[((b*d - 2*a*e + (2*c*d - b*e)*x)/((p + 1)*(b^2 - 4*a*c)))*(a + b*x + c*x^2)^(p + 1), x] - Simp[(2*p + 3)*((2*c*d - b*e)/((p + 1)*(b^2 - 4*a*c)))] Int[(a + b*x + c*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2191

```
Int[(Pq_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := With[{Q =
PolynomialQuotient[Pq, a + b*x + c*x^2, x], f = Coeff[PolynomialRemainder[P
q, a + b*x + c*x^2, x], x, 0], g = Coeff[PolynomialRemainder[Pq, a + b*x +
c*x^2, x], x, 1]}, Simp[(b*f - 2*a*g + (2*c*f - b*g)*x)*((a + b*x + c*x^2)^(
p + 1)/((p + 1)*(b^2 - 4*a*c))), x] + Simp[1/((p + 1)*(b^2 - 4*a*c)) Int
[(a + b*x + c*x^2)^(p + 1)*ExpandToSum[(p + 1)*(b^2 - 4*a*c)*Q - (2*p + 3)*
(2*c*f - b*g), x], x], x]] /; FreeQ[{a, b, c}, x] && PolyQ[Pq, x] && NeQ[b^
2 - 4*a*c, 0] && LtQ[p, -1]
```

rule 7286

```
Int[u_, x_Symbol] := With[{lst = FunctionOfSquareRootOfQuadratic[u, x]}, Si
mp[2 Subst[Int[lst[[1]], x], x, lst[[2]]], x] /; !FalseQ[lst] && EqQ[lst
[[3]], 2]] /; EulerIntegrandQ[u, x]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.19 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.59

method	result
trager	$\frac{(321x^3 - 311x^2 - 183x + 117)x}{200(3x^2 - 2x - 3)^2} - \frac{(231x^3 - 221x^2 - 113x + 87)\sqrt{4x^2 - 2x - 3}}{200(3x^2 - 2x - 3)^2} + \frac{39\text{RootOf}(_Z^2 - 10)\ln\left(-\frac{\text{RootOf}(_Z^2 - 10)x + \text{RootOf}(_Z^2 - 10)}{2000}\right)}{2000}$
default	Expression too large to display

input

```
int(1/(x+(4*x^2-2*x-3)^(1/2))^3,x,method=_RETURNVERBOSE)
```

output

```
1/200*(321*x^3-311*x^2-183*x+117)*x/(3*x^2-2*x-3)^2-1/200*(231*x^3-221*x^2
-113*x+87)/(3*x^2-2*x-3)^2*(4*x^2-2*x-3)^(1/2)+39/2000*RootOf(_Z^2-10)*ln(
-(RootOf(_Z^2-10)*x+10*(4*x^2-2*x-3)^(1/2)+3*RootOf(_Z^2-10))/(RootOf(_Z^2
-10)*x+x+3))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.02

$$\int \frac{1}{(x + \sqrt{-3 - 2x + 4x^2})^3} dx =$$

$$\frac{9240x^4 - 14660x^3 - 39\sqrt{10}(9x^4 - 12x^3 - 14x^2 + 12x + 9) \log\left(\frac{1681x^2 + 4\sqrt{10}(41x^2 - 14x - 21) + 2\sqrt{4x^2 - 2x - 3}}{3x^2 - 2x - 3}\right)}{}$$

input `integrate(1/(x+(4*x^2-2*x-3)^(1/2))^3,x, algorithm="fricas")`

output `-1/4000*(9240*x^4 - 14660*x^3 - 39*sqrt(10)*(9*x^4 - 12*x^3 - 14*x^2 + 12*x + 9)*log((1681*x^2 + 4*sqrt(10)*(41*x^2 - 14*x - 21) + 2*sqrt(4*x^2 - 2*x - 3)*(41*sqrt(10)*(x + 3) + 40*x + 120) - 574*x - 861)/(3*x^2 - 2*x - 3)) - 39*sqrt(10)*(9*x^4 - 12*x^3 - 14*x^2 + 12*x + 9)*log((9*x^2 - 2*sqrt(10)*(3*x - 1) - 6*x + 11)/(3*x^2 - 2*x - 3)) - 20700*x^2 + 20*(231*x^3 - 221*x^2 - 113*x + 87)*sqrt(4*x^2 - 2*x - 3) + 18540*x + 15660)/(9*x^4 - 12*x^3 - 14*x^2 + 12*x + 9)`

Sympy [F]

$$\int \frac{1}{(x + \sqrt{-3 - 2x + 4x^2})^3} dx = \int \frac{1}{(x + \sqrt{4x^2 - 2x - 3})^3} dx$$

input `integrate(1/(x+(4*x**2-2*x-3)**(1/2))**3,x)`

output `Integral((x + sqrt(4*x**2 - 2*x - 3))**(-3), x)`

Maxima [F]

$$\int \frac{1}{(x + \sqrt{-3 - 2x + 4x^2})^3} dx = \int \frac{1}{(x + \sqrt{4x^2 - 2x - 3})^3} dx$$

input `integrate(1/(x+(4*x^2-2*x-3)^(1/2))^3,x, algorithm="maxima")`

output `integrate((x + sqrt(4*x^2 - 2*x - 3))^(-3), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 419 vs. 2(181) = 362.

Time = 0.17 (sec) , antiderivative size = 419, normalized size of antiderivative = 1.92

$$\begin{aligned} \int \frac{1}{(x + \sqrt{-3 - 2x + 4x^2})^3} dx &= \frac{39}{4000} \sqrt{10} \log \left(\frac{|6x - 2\sqrt{10} - 2|}{|6x + 2\sqrt{10} - 2|} \right) \\ &- \frac{39}{4000} \sqrt{10} \log \left(\frac{|-4x - 2\sqrt{10} + 2\sqrt{4x^2 - 2x - 3} + 2|}{|-4x + 2\sqrt{10} + 2\sqrt{4x^2 - 2x - 3} + 2|} \right) \\ &+ \frac{39}{4000} \sqrt{10} \log \left(\frac{|-12x - 2\sqrt{10} + 6\sqrt{4x^2 - 2x - 3} + 2|}{|-12x + 2\sqrt{10} + 6\sqrt{4x^2 - 2x - 3} + 2|} \right) \\ &+ \frac{351(2x - \sqrt{4x^2 - 2x - 3})^7 + 9796(2x - \sqrt{4x^2 - 2x - 3})^6 - 31087(2x - \sqrt{4x^2 - 2x - 3})^5 - 63}{300(3(2x - \sqrt{4x^2 - 2x - 3})^4 - 8(2x} \\ &+ \frac{351x^3 + 949x^2 - 933x - 963}{600(3x^2 - 2x - 3)^2} \end{aligned}$$

input `integrate(1/(x+(4*x^2-2*x-3)^(1/2))^3,x, algorithm="giac")`

output

```
39/4000*sqrt(10)*log(abs(6*x - 2*sqrt(10) - 2)/abs(6*x + 2*sqrt(10) - 2))
- 39/4000*sqrt(10)*log(abs(-4*x - 2*sqrt(10) + 2*sqrt(4*x^2 - 2*x - 3) + 2)
)/abs(-4*x + 2*sqrt(10) + 2*sqrt(4*x^2 - 2*x - 3) + 2)) + 39/4000*sqrt(10)
*log(abs(-12*x - 2*sqrt(10) + 6*sqrt(4*x^2 - 2*x - 3) + 2)/abs(-12*x + 2*s
qrt(10) + 6*sqrt(4*x^2 - 2*x - 3) + 2)) + 1/300*(351*(2*x - sqrt(4*x^2 - 2
*x - 3))^7 + 9796*(2*x - sqrt(4*x^2 - 2*x - 3))^6 - 31087*(2*x - sqrt(4*x^
2 - 2*x - 3))^5 - 63608*(2*x - sqrt(4*x^2 - 2*x - 3))^4 + 131661*(2*x - sq
rt(4*x^2 - 2*x - 3))^3 + 131364*(2*x - sqrt(4*x^2 - 2*x - 3))^2 - 278154*x
+ 139077*sqrt(4*x^2 - 2*x - 3) - 129600)/(3*(2*x - sqrt(4*x^2 - 2*x - 3))
^4 - 8*(2*x - sqrt(4*x^2 - 2*x - 3))^3 - 26*(2*x - sqrt(4*x^2 - 2*x - 3))^
2 + 48*x - 24*sqrt(4*x^2 - 2*x - 3) + 27)^2 + 1/600*(351*x^3 + 949*x^2 - 9
33*x - 963)/(3*x^2 - 2*x - 3)^2
```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(x + \sqrt{-3 - 2x + 4x^2})^3} dx = \int \frac{1}{(x + \sqrt{4x^2 - 2x - 3})^3} dx$$

input

```
int(1/(x + (4*x^2 - 2*x - 3)^(1/2))^3,x)
```

output

```
int(1/(x + (4*x^2 - 2*x - 3)^(1/2))^3, x)
```

Reduce [B] (verification not implemented)

Time = 2.61 (sec) , antiderivative size = 1122, normalized size of antiderivative = 5.15

$$\int \frac{1}{(x + \sqrt{-3 - 2x + 4x^2})^3} dx = \text{Too large to display}$$

input

```
int(1/(x+(4*x^2-2*x-3)^(1/2))^3,x)
```


output

```
( - 2773519866*sqrt(10)*atan((2*sqrt(4*x**2 - 2*x - 3)*i + 4*i*x - i)/(2*sqrt(10) - 1))*i*x**4 + 3698026488*sqrt(10)*atan((2*sqrt(4*x**2 - 2*x - 3)*i + 4*i*x - i)/(2*sqrt(10) - 1))*i*x**3 + 4314364236*sqrt(10)*atan((2*sqrt(4*x**2 - 2*x - 3)*i + 4*i*x - i)/(2*sqrt(10) - 1))*i*x**2 - 3698026488*sqrt(10)*atan((2*sqrt(4*x**2 - 2*x - 3)*i + 4*i*x - i)/(2*sqrt(10) - 1))*i*x - 2773519866*sqrt(10)*atan((2*sqrt(4*x**2 - 2*x - 3)*i + 4*i*x - i)/(2*sqrt(10) - 1))*i - 36506158920*sqrt(4*x**2 - 2*x - 3)*x**3 + 34925805720*sqrt(4*x**2 - 2*x - 3)*x**2 + 17857991160*sqrt(4*x**2 - 2*x - 3)*x - 13749072840*sqrt(4*x**2 - 2*x - 3) + 2773519866*sqrt(10)*log(- sqrt(10) + 3*x - 1)*x**4 - 3698026488*sqrt(10)*log(- sqrt(10) + 3*x - 1)*x**3 - 4314364236*sqrt(10)*log(- sqrt(10) + 3*x - 1)*x**2 + 3698026488*sqrt(10)*log(- sqrt(10) + 3*x - 1)*x + 2773519866*sqrt(10)*log(- sqrt(10) + 3*x - 1) - 2773519866*sqrt(10)*log(sqrt(10) + 3*x - 1)*x**4 + 3698026488*sqrt(10)*log(sqrt(10) + 3*x - 1)*x**3 + 4314364236*sqrt(10)*log(sqrt(10) + 3*x - 1)*x**2 - 3698026488*sqrt(10)*log(sqrt(10) + 3*x - 1)*x - 2773519866*sqrt(10)*log(sqrt(10) + 3*x - 1) + 1386759933*sqrt(10)*log(16*sqrt(4*x**2 - 2*x - 3)*x - 4*sqrt(4*x**2 - 2*x - 3) + 4*sqrt(10) + 32*x**2 - 16*x - 52)*x**4 - 1849013244*sqrt(10)*log(16*sqrt(4*x**2 - 2*x - 3)*x - 4*sqrt(4*x**2 - 2*x - 3) + 4*sqrt(10) + 32*x**2 - 16*x - 52)*x**3 - 2157182118*sqrt(10)*log(16*sqrt(4*x**2 - 2*x - 3)*x - 4*sqrt(4*x**2 - 2*x - 3) + 4*sqrt(10) + 32*x**2 - ...
```

3.36 $\int (d + ex + f\sqrt{-a + bx + cx^2})^2 dx$

Optimal result	305
Mathematica [A] (verified)	306
Rubi [A] (verified)	306
Maple [A] (verified)	307
Fricas [A] (verification not implemented)	308
Sympy [A] (verification not implemented)	309
Maxima [A] (verification not implemented)	310
Giac [A] (verification not implemented)	310
Mupad [B] (verification not implemented)	311
Reduce [B] (verification not implemented)	312

Optimal result

Integrand size = 25, antiderivative size = 237

$$\int (d + ex + f\sqrt{-a + bx + cx^2})^2 dx = (d^2 - af^2)x + \frac{1}{2}(2de + bf^2)x^2 + \frac{1}{3}(e^2 + cf^2)x^3 + \frac{df(b + 2cx)\sqrt{-a + bx + cx^2}}{2c} - \frac{bef(b + 2cx)\sqrt{-a + bx + cx^2}}{4c^2} + \frac{2ef(-a + bx + cx^2)^{3/2}}{3c} - \frac{(b^2 + 4ac)df \operatorname{arctanh}\left(\frac{b+2cx}{2\sqrt{c}\sqrt{-a+bx+cx^2}}\right)}{4c^{3/2}} + \frac{b(b^2 + 4ac)ef \operatorname{arctanh}\left(\frac{b+2cx}{2\sqrt{c}\sqrt{-a+bx+cx^2}}\right)}{8c^{5/2}}$$

output

```
(-a*f^2+d^2)*x+1/2*(b*f^2+2*d*e)*x^2+1/3*(c*f^2+e^2)*x^3+1/2*d*f*(2*c*x+b)
*(c*x^2+b*x-a)^(1/2)/c-1/4*b*e*f*(2*c*x+b)*(c*x^2+b*x-a)^(1/2)/c^2+2/3*e*f
*(c*x^2+b*x-a)^(3/2)/c-1/4*(4*a*c+b^2)*d*f*arctanh(1/2*(2*c*x+b)/c^(1/2)/(
c*x^2+b*x-a)^(1/2))/c^(3/2)+1/8*b*(4*a*c+b^2)*e*f*arctanh(1/2*(2*c*x+b)/c^(
1/2)/(c*x^2+b*x-a)^(1/2))/c^(5/2)
```

Mathematica [A] (verified)

Time = 1.11 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.68

$$\int \left(d + ex + f\sqrt{-a + bx + cx^2} \right)^2 dx$$

$$= \frac{1}{24} \left(4x(6d^2 - 6af^2 + 6dex + x(3bf^2 + 2(e^2 + cf^2)x)) \right. \\ \left. + \frac{2f\sqrt{-a + x(b + cx)}(-3b^2e + 2bc(3d + ex) + 4c(-2ae + cx(3d + 2ex)))}{c^2} \right. \\ \left. + \frac{3(b^2 + 4ac)(2cd - be)f \log \left(c^2 \left(b + 2cx - 2\sqrt{c}\sqrt{-a + x(b + cx)} \right) \right)}{c^{5/2}} \right)$$

input

```
Integrate[(d + e*x + f*Sqrt[-a + b*x + c*x^2])^2,x]
```

output

```
(4*x*(6*d^2 - 6*a*f^2 + 6*d*e*x + x*(3*b*f^2 + 2*(e^2 + c*f^2)*x)) + (2*f*
Sqrt[-a + x*(b + c*x)]*(-3*b^2*e + 2*b*c*(3*d + e*x) + 4*c*(-2*a*e + c*x*(
3*d + 2*e*x))))/c^2 + (3*(b^2 + 4*a*c)*(2*c*d - b*e)*f*Log[c^2*(b + 2*c*x
- 2*Sqrt[c]*Sqrt[-a + x*(b + c*x)]])/c^(5/2))/24
```

Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(f\sqrt{-a + bx + cx^2} + d + ex \right)^2 dx$$

↓ 7293

$$\int \left(2df\sqrt{-a + bx + cx^2} + 2efx\sqrt{-a + bx + cx^2} + d^2 \left(1 - \frac{af^2}{d^2} \right) + 2dex \left(\frac{bf^2}{2de} + 1 \right) + e^2x^2 \left(\frac{cf^2}{e^2} + 1 \right) \right) dx$$

↓ 2009

$$\begin{aligned} & -\frac{df(4ac + b^2) \operatorname{arctanh}\left(\frac{b+2cx}{2\sqrt{c}\sqrt{-a+bx+cx^2}}\right)}{4c^{3/2}} + \frac{bef(4ac + b^2) \operatorname{arctanh}\left(\frac{b+2cx}{2\sqrt{c}\sqrt{-a+bx+cx^2}}\right)}{8c^{5/2}} - \\ & \frac{bef(b + 2cx)\sqrt{-a + bx + cx^2}}{4c^2} + \frac{df(b + 2cx)\sqrt{-a + bx + cx^2}}{2c} + \frac{2ef(-a + bx + cx^2)^{3/2}}{3c} + \\ & \quad x(d^2 - af^2) + \frac{1}{2}x^2(bf^2 + 2de) + \frac{1}{3}x^3(cf^2 + e^2) \end{aligned}$$

input `Int[(d + e*x + f*Sqrt[-a + b*x + c*x^2])^2,x]`

output `(d^2 - a*f^2)*x + ((2*d*e + b*f^2)*x^2)/2 + ((e^2 + c*f^2)*x^3)/3 + (d*f*(b + 2*c*x)*Sqrt[-a + b*x + c*x^2])/(2*c) - (b*e*f*(b + 2*c*x)*Sqrt[-a + b*x + c*x^2])/(4*c^2) + (2*e*f*(-a + b*x + c*x^2)^(3/2))/(3*c) - ((b^2 + 4*a*c)*d*f*ArcTanh[(b + 2*c*x)/(2*Sqrt[c]*Sqrt[-a + b*x + c*x^2])])/(4*c^(3/2)) + (b*(b^2 + 4*a*c)*e*f*ArcTanh[(b + 2*c*x)/(2*Sqrt[c]*Sqrt[-a + b*x + c*x^2])])/(8*c^(5/2))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

Maple [A] (verified)

Time = 1.76 (sec) , antiderivative size = 205, normalized size of antiderivative = 0.86

method	result
default	$f^2\left(\frac{1}{3}cx^3 + \frac{1}{2}bx^2 - xa\right) + 2f\left(d\left(\frac{(2cx+b)\sqrt{cx^2+bx-a}}{4c} + \frac{(-4ac-b^2)\ln\left(\frac{\frac{b}{2}+\frac{cx}{\sqrt{c}}+\sqrt{cx^2+bx-a}}{\sqrt{c}}\right)}{8c^{\frac{3}{2}}}\right)\right) + e\left(\frac{cx^2+}{\dots}\right)$

input `int((d+e*x+f*(c*x^2+b*x-a)^(1/2))^2,x,method=_RETURNVERBOSE)`

output `f^2*(1/3*c*x^3+1/2*b*x^2-x*a)+2*f*(d*(1/4*(2*c*x+b)/c*(c*x^2+b*x-a)^(1/2)+1/8*(-4*a*c-b^2)/c^(3/2)*ln((1/2*b+c*x)/c^(1/2)+(c*x^2+b*x-a)^(1/2)))+e*(1/3*(c*x^2+b*x-a)^(3/2)/c-1/2*b/c*(1/4*(2*c*x+b)/c*(c*x^2+b*x-a)^(1/2)+1/8*(-4*a*c-b^2)/c^(3/2)*ln((1/2*b+c*x)/c^(1/2)+(c*x^2+b*x-a)^(1/2))))+1/3*(e*x+d)^3/e`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 438, normalized size of antiderivative = 1.85

$$\int \left(d + ex + f\sqrt{-a + bx + cx^2} \right)^2 dx$$

$$= \left[\frac{16(c^4 f^2 + c^3 e^2)x^3 - 3(2(b^2 c + 4ac^2)d - (b^3 + 4abc)e)\sqrt{c}f \log(8c^2 x^2 + 8bcx + b^2 + 4\sqrt{cx^2 + bx - a})}{16(c^4 f^2 + c^3 e^2)x^3 - 3(2(b^2 c + 4ac^2)d - (b^3 + 4abc)e)\sqrt{c}f \log(8c^2 x^2 + 8bcx + b^2 + 4\sqrt{cx^2 + bx - a})} \right]$$

input `integrate((d+e*x+f*(c*x^2+b*x-a)^(1/2))^2,x, algorithm="fricas")`

output `[1/48*(16*(c^4*f^2 + c^3*e^2)*x^3 - 3*(2*(b^2*c + 4*a*c^2)*d - (b^3 + 4*a*b*c)*e)*sqrt(c)*f*log(8*c^2*x^2 + 8*b*c*x + b^2 + 4*sqrt(c*x^2 + b*x - a)*(2*c*x + b)*sqrt(c) - 4*a*c) + 24*(b*c^3*f^2 + 2*c^3*d*e)*x^2 - 48*(a*c^3*f^2 - c^3*d^2)*x + 4*(8*c^3*e*f*x^2 + 2*(6*c^3*d + b*c^2*e)*f*x + (6*b*c^2*d - (3*b^2*c + 8*a*c^2)*e)*f)*sqrt(c*x^2 + b*x - a))/c^3, 1/24*(8*(c^4*f^2 + c^3*e^2)*x^3 + 3*(2*(b^2*c + 4*a*c^2)*d - (b^3 + 4*a*b*c)*e)*sqrt(-c)*f*arctan(1/2*sqrt(c*x^2 + b*x - a)*(2*c*x + b)*sqrt(-c)/(c^2*x^2 + b*c*x - a*c)) + 12*(b*c^3*f^2 + 2*c^3*d*e)*x^2 - 24*(a*c^3*f^2 - c^3*d^2)*x + 2*(8*c^3*e*f*x^2 + 2*(6*c^3*d + b*c^2*e)*f*x + (6*b*c^2*d - (3*b^2*c + 8*a*c^2)*e)*f)*sqrt(c*x^2 + b*x - a))/c^3]`

Sympy [A] (verification not implemented)

Time = 0.71 (sec) , antiderivative size = 352, normalized size of antiderivative = 1.49

$$\int \left(d + ex + f\sqrt{-a + bx + cx^2} \right)^2 dx = -af^2x + \frac{bf^2x^2}{2} + \frac{cf^2x^3}{3} + d^2x + dex^2$$

$$+ 2df \left\{ \begin{array}{l} \left(-\frac{a}{2} - \frac{b^2}{8c} \right) \left(\begin{array}{l} \frac{\log(b + 2\sqrt{c}\sqrt{-a + bx + cx^2} + 2cx)}{\sqrt{c}} \text{ for } a + \frac{b^2}{4c} \neq 0 \\ \frac{(\frac{b}{2c} + x) \log(\frac{b}{2c} + x)}{\sqrt{c}(\frac{b}{2c} + x)^2} \text{ otherwise} \end{array} \right) + \left(\frac{b}{4c} + \frac{x}{2} \right) \sqrt{-a + bx + cx^2} \\ \frac{2(-a + bx)^{\frac{3}{2}}}{3b} \\ x\sqrt{-a} \end{array} \right. + \left(\frac{b}{4c} + \frac{x}{2} \right) \sqrt{-a + bx + cx^2} \left(\frac{bx}{12c} + \dots \right)$$

$$+ \frac{e^2x^3}{3}$$

$$+ 2ef \left\{ \begin{array}{l} \left(\frac{ab}{12c} - \frac{b(-\frac{a}{3} - \frac{b^2}{8c})}{2c} \right) \left(\begin{array}{l} \frac{\log(b + 2\sqrt{c}\sqrt{-a + bx + cx^2} + 2cx)}{\sqrt{c}} \text{ for } a + \frac{b^2}{4c} \neq 0 \\ \frac{(\frac{b}{2c} + x) \log(\frac{b}{2c} + x)}{\sqrt{c}(\frac{b}{2c} + x)^2} \text{ otherwise} \end{array} \right) + \sqrt{-a + bx + cx^2} \left(\frac{bx}{12c} + \dots \right) \\ \frac{2\left(\frac{a(-a + bx)^{\frac{3}{2}}}{3} + \frac{(-a + bx)^{\frac{5}{2}}}{5}\right)}{b^2} \\ \frac{x^2\sqrt{-a}}{2} \end{array} \right.$$

```
input integrate((d+e*x+f*(c*x**2+b*x-a)**(1/2))**2,x)
```

```
output -a*f**2*x + b*f**2*x**2/2 + c*f**2*x**3/3 + d**2*x + d*e*x**2 + 2*d*f*Piecewise((( -a/2 - b**2/(8*c))*Piecewise((log(b + 2*sqrt(c)*sqrt(-a + b*x + c*x**2) + 2*c*x)/sqrt(c), Ne(a + b**2/(4*c), 0)), ((b/(2*c) + x)*log(b/(2*c) + x)/sqrt(c*(b/(2*c) + x)**2), True)) + (b/(4*c) + x/2)*sqrt(-a + b*x + c*x**2), Ne(c, 0)), (2*(-a + b*x)**(3/2)/(3*b), Ne(b, 0)), (x*sqrt(-a), True)) + e**2*x**3/3 + 2*e*f*Piecewise(((a*b/(12*c) - b*(-a/3 - b**2/(8*c)))/(2*c))*Piecewise((log(b + 2*sqrt(c)*sqrt(-a + b*x + c*x**2) + 2*c*x)/sqrt(c), Ne(a + b**2/(4*c), 0)), ((b/(2*c) + x)*log(b/(2*c) + x)/sqrt(c*(b/(2*c) + x)**2), True)) + sqrt(-a + b*x + c*x**2)*(b*x/(12*c) + x**2/3 + (-a/3 - b**2/(8*c))/c), Ne(c, 0)), (2*(a*(-a + b*x)**(3/2)/3 + (-a + b*x)**(5/2)/5)/b**2, Ne(b, 0)), (x**2*sqrt(-a)/2, True))
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 286, normalized size of antiderivative = 1.21

$$\int \left(d + ex + f\sqrt{-a + bx + cx^2} \right)^2 dx = \frac{1}{3} e^2 x^3 - \frac{1}{24} \left(\frac{12\sqrt{cx^2 + bx - abx}}{c} - \frac{3b^3 \log(2cx + b + 2\sqrt{cx^2 + bx - a\sqrt{c}})}{c^{\frac{5}{2}}} - \frac{12ab \log(2cx + b + 2\sqrt{cx^2 + bx - a\sqrt{c}})}{c^{\frac{3}{2}}} \right) + \frac{1}{6} (2cx^3 + 3bx^2 - 6ax)f^2 + d^2x + \frac{1}{4} \left(4ex^2 + \left(4\sqrt{cx^2 + bx - ax} - \frac{b^2 \log(2cx + b + 2\sqrt{cx^2 + bx - a\sqrt{c}})}{c^{\frac{3}{2}}} - \frac{4a \log(2cx + b + 2\sqrt{cx^2 + bx - a\sqrt{c}})}{\sqrt{c}} \right) \right)$$

input `integrate((d+e*x+f*(c*x^2+b*x-a)^(1/2))^2,x, algorithm="maxima")`

output `1/3*e^2*x^3 - 1/24*(12*sqrt(c*x^2 + b*x - a)*b*x/c - 3*b^3*log(2*c*x + b + 2*sqrt(c*x^2 + b*x - a)*sqrt(c))/c^(5/2) - 12*a*b*log(2*c*x + b + 2*sqrt(c*x^2 + b*x - a)*sqrt(c))/c^(3/2) + 6*sqrt(c*x^2 + b*x - a)*b^2/c^2 - 16*(c*x^2 + b*x - a)^(3/2)/c)*e*f + 1/6*(2*c*x^3 + 3*b*x^2 - 6*a*x)*f^2 + d^2*x + 1/4*(4*e*x^2 + (4*sqrt(c*x^2 + b*x - a)*x - b^2*log(2*c*x + b + 2*sqrt(c*x^2 + b*x - a)*sqrt(c))/c^(3/2) - 4*a*log(2*c*x + b + 2*sqrt(c*x^2 + b*x - a)*sqrt(c))/sqrt(c) + 2*sqrt(c*x^2 + b*x - a)*b/c)*f)*d`

Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 179, normalized size of antiderivative = 0.76

$$\int \left(d + ex + f\sqrt{-a + bx + cx^2} \right)^2 dx = \frac{1}{3} cf^2x^3 + \frac{1}{2} bf^2x^2 + \frac{1}{3} e^2x^3 - af^2x + dex^2 + d^2x + \frac{1}{12} \sqrt{cx^2 + bx - a} \left(2 \left(4efx + \frac{6c^2df + bcef}{c^2} \right) x + \frac{6bcd f - 3b^2ef - 8acef}{c^2} \right) + \frac{(2b^2cdf + 8ac^2df - b^3ef - 4abcef) \log(|2(\sqrt{cx} - \sqrt{cx^2 + bx - a})\sqrt{c} + b|)}{8c^{\frac{5}{2}}}$$

input `integrate((d+e*x+f*(c*x^2+b*x-a)^(1/2))^2,x, algorithm="giac")`

output

```
1/3*c*f^2*x^3 + 1/2*b*f^2*x^2 + 1/3*e^2*x^3 - a*f^2*x + d*e*x^2 + d^2*x +
1/12*sqrt(c*x^2 + b*x - a)*(2*(4*e*f*x + (6*c^2*d*f + b*c*e*f)/c^2)*x + (6
*b*c*d*f - 3*b^2*e*f - 8*a*c*e*f)/c^2) + 1/8*(2*b^2*c*d*f + 8*a*c^2*d*f -
b^3*e*f - 4*a*b*c*e*f)*log(abs(2*(sqrt(c)*x - sqrt(c*x^2 + b*x - a))*sqrt(
c) + b))/c^(5/2)
```

Mupad [B] (verification not implemented)

Time = 22.86 (sec) , antiderivative size = 203, normalized size of antiderivative = 0.86

$$\int (d + ex + f\sqrt{-a + bx + cx^2})^2 dx$$

$$= x^3 \left(\frac{e^2}{3} + \frac{cf^2}{3} \right) + x^2 \left(\frac{bf^2}{2} + de \right) - x (af^2 - d^2)$$

$$+ 2df \left(\frac{x}{2} + \frac{b}{4c} \right) \sqrt{cx^2 + bx - a} - \frac{df \ln \left(\sqrt{cx^2 + bx - a} + \frac{\frac{b}{2} + cx}{\sqrt{c}} \right) \left(\frac{b^2}{4} + ac \right)}{c^{3/2}}$$

$$+ \frac{ef \ln \left(\frac{b+2cx}{\sqrt{c}} + 2\sqrt{cx^2 + bx - a} \right) (b^3 + 4acb)}{8c^{5/2}}$$

$$- \frac{ef(3b^2 - 2cxb + 8c(a - cx^2)) \sqrt{cx^2 + bx - a}}{12c^2}$$

input

```
int((d + f*(b*x - a + c*x^2)^(1/2) + e*x)^2,x)
```

output

```
x^3*((c*f^2)/3 + e^2/3) + x^2*(d*e + (b*f^2)/2) - x*(a*f^2 - d^2) + 2*d*f*
(x/2 + b/(4*c))*(b*x - a + c*x^2)^(1/2) - (d*f*log((b*x - a + c*x^2)^(1/2)
+ (b/2 + c*x)/c^(1/2))*(a*c + b^2/4))/c^(3/2) + (e*f*log((b + 2*c*x)/c^(1
/2) + 2*(b*x - a + c*x^2)^(1/2))*(b^3 + 4*a*b*c))/(8*c^(5/2)) - (e*f*(8*c*
(a - c*x^2) + 3*b^2 - 2*b*c*x)*(b*x - a + c*x^2)^(1/2))/(12*c^2)
```


Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 380, normalized size of antiderivative = 1.60

$$\int \left(d + ex + f\sqrt{-a + bx + cx^2} \right)^2 dx$$

$$= \frac{-16\sqrt{cx^2 + bx - a}ac^2ef - 6\sqrt{cx^2 + bx - a}b^2cef + 12\sqrt{cx^2 + bx - a}bc^2df + 4\sqrt{cx^2 + bx - a}b^2c^2ef}{}$$

input `int((d+e*x+f*(c*x^2+b*x-a)^(1/2))^2,x)`

output

```
( - 16*sqrt( - a + b*x + c*x**2)*a*c**2*e*f - 6*sqrt( - a + b*x + c*x**2)*
b**2*c*e*f + 12*sqrt( - a + b*x + c*x**2)*b*c**2*d*f + 4*sqrt( - a + b*x +
c*x**2)*b*c**2*e*f*x + 24*sqrt( - a + b*x + c*x**2)*c**3*d*f*x + 16*sqrt(
- a + b*x + c*x**2)*c**3*e*f*x**2 + 12*sqrt(c)*log((2*sqrt(c)*sqrt( - a +
b*x + c*x**2) + b + 2*c*x)/sqrt(4*a*c + b**2))*a*b*c*e*f - 24*sqrt(c)*log
((2*sqrt(c)*sqrt( - a + b*x + c*x**2) + b + 2*c*x)/sqrt(4*a*c + b**2))*a*c
**2*d*f + 3*sqrt(c)*log((2*sqrt(c)*sqrt( - a + b*x + c*x**2) + b + 2*c*x)/
sqrt(4*a*c + b**2))*b**3*e*f - 6*sqrt(c)*log((2*sqrt(c)*sqrt( - a + b*x +
c*x**2) + b + 2*c*x)/sqrt(4*a*c + b**2))*b**2*c*d*f - 24*a*c**3*f**2*x + 1
2*b*c**3*f**2*x**2 + 8*c**4*f**2*x**3 + 24*c**3*d**2*x + 24*c**3*d*e*x**2
+ 8*c**3*e**2*x**3)/(24*c**3)
```

3.37 $\int (d + ex + f\sqrt{-a + bx + cx^2}) dx$

Optimal result	313
Mathematica [A] (verified)	313
Rubi [A] (verified)	314
Maple [A] (verified)	315
Fricas [A] (verification not implemented)	315
Sympy [A] (verification not implemented)	316
Maxima [A] (verification not implemented)	317
Giac [A] (verification not implemented)	317
Mupad [B] (verification not implemented)	318
Reduce [B] (verification not implemented)	318

Optimal result

Integrand size = 23, antiderivative size = 92

$$\int (d + ex + f\sqrt{-a + bx + cx^2}) dx = dx + \frac{ex^2}{2} + \frac{f(b + 2cx)\sqrt{-a + bx + cx^2}}{4c} - \frac{(b^2 + 4ac) f \operatorname{arctanh}\left(\frac{b+2cx}{2\sqrt{c}\sqrt{-a+bx+cx^2}}\right)}{8c^{3/2}}$$

output

```
d*x+1/2*e*x^2+1/4*f*(2*c*x+b)*(c*x^2+b*x-a)^(1/2)/c-1/8*(4*a*c+b^2)*f*arctanh(1/2*(2*c*x+b)/c^(1/2)/(c*x^2+b*x-a)^(1/2))/c^(3/2)
```

Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.35

$$\begin{aligned} & \int (d + ex + f\sqrt{-a + bx + cx^2}) dx \\ &= dx + \frac{ex^2}{2} + \frac{f(b + 2cx)\sqrt{-a + x(b + cx)}}{4c} \\ & \quad + \frac{(-b^2 - 4ac) f \sqrt{-a + x(b + cx)} \operatorname{arctanh}\left(\frac{b+2cx}{2\sqrt{c}\sqrt{-a+bx+cx^2}}\right)}{8c^{3/2}\sqrt{-a + bx + cx^2}} \end{aligned}$$

input `Integrate[d + e*x + f*Sqrt[-a + b*x + c*x^2],x]`

output `d*x + (e*x^2)/2 + (f*(b + 2*c*x)*Sqrt[-a + x*(b + c*x)]/(4*c) + ((-b^2 - 4*a*c)*f*Sqrt[-a + x*(b + c*x)]*ArcTanh[(b + 2*c*x)/(2*Sqrt[c]*Sqrt[-a + b*x + c*x^2])])/(8*c^(3/2)*Sqrt[-a + b*x + c*x^2])`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(f \sqrt{-a + bx + cx^2} + d + ex \right) dx$$

↓ 2009

$$-\frac{f(4ac + b^2) \operatorname{arctanh}\left(\frac{b+2cx}{2\sqrt{c}\sqrt{-a+bx+cx^2}}\right)}{8c^{3/2}} + \frac{f(b+2cx)\sqrt{-a+bx+cx^2}}{4c} + dx + \frac{ex^2}{2}$$

input `Int[d + e*x + f*Sqrt[-a + b*x + c*x^2],x]`

output `d*x + (e*x^2)/2 + (f*(b + 2*c*x)*Sqrt[-a + b*x + c*x^2])/(4*c) - ((b^2 + 4*a*c)*f*ArcTanh[(b + 2*c*x)/(2*Sqrt[c]*Sqrt[-a + b*x + c*x^2])])/(8*c^(3/2))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.88

method	result	size
default	$dx + \frac{ex^2}{2} + f\left(\frac{(2cx+b)\sqrt{cx^2+bx-a}}{4c} + \frac{(-4ac-b^2)\ln\left(\frac{\frac{b}{2}+cx}{\sqrt{c}} + \sqrt{cx^2+bx-a}\right)}{8c^{\frac{3}{2}}}\right)$	81
parts	$dx + \frac{ex^2}{2} + f\left(\frac{(2cx+b)\sqrt{cx^2+bx-a}}{4c} + \frac{(-4ac-b^2)\ln\left(\frac{\frac{b}{2}+cx}{\sqrt{c}} + \sqrt{cx^2+bx-a}\right)}{8c^{\frac{3}{2}}}\right)$	81

input `int(d+e*x+f*(c*x^2+b*x-a)^(1/2),x,method=_RETURNVERBOSE)`

output `d*x+1/2*e*x^2+f*(1/4*(2*c*x+b)/c*(c*x^2+b*x-a)^(1/2)+1/8*(-4*a*c-b^2)/c^(3/2)*ln((1/2*b+c*x)/c^(1/2)+(c*x^2+b*x-a)^(1/2)))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 222, normalized size of antiderivative = 2.41

$$\int (d + ex + f\sqrt{-a + bx + cx^2}) dx$$

$$= \left[\frac{8c^2ex^2 + 16c^2dx + (b^2 + 4ac)\sqrt{c}f \log(8c^2x^2 + 8bcx + b^2 - 4\sqrt{cx^2 + bx - a}(2cx + b)\sqrt{c} - 4ac) + 4}{16c^2} \right]$$

input `integrate(d+e*x+f*(c*x^2+b*x-a)^(1/2),x, algorithm="fricas")`

output

```
[1/16*(8*c^2*e*x^2 + 16*c^2*d*x + (b^2 + 4*a*c)*sqrt(c)*f*log(8*c^2*x^2 + 8*b*c*x + b^2 - 4*sqrt(c*x^2 + b*x - a)*(2*c*x + b)*sqrt(c) - 4*a*c) + 4*(2*c^2*f*x + b*c*f)*sqrt(c*x^2 + b*x - a))/c^2, 1/8*(4*c^2*e*x^2 + 8*c^2*d*x + (b^2 + 4*a*c)*sqrt(-c)*f*arctan(1/2*sqrt(c*x^2 + b*x - a)*(2*c*x + b)*sqrt(-c)/(c^2*x^2 + b*c*x - a*c)) + 2*(2*c^2*f*x + b*c*f)*sqrt(c*x^2 + b*x - a))/c^2]
```

Sympy [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.46

$$\int (d + ex + f\sqrt{-a + bx + cx^2}) dx = dx + \frac{ex^2}{2} + f \begin{cases} \left(-\frac{a}{2} - \frac{b^2}{8c} \right) \begin{cases} \frac{\log(b + 2\sqrt{c}\sqrt{-a + bx + cx^2} + 2cx)}{\sqrt{c}} & \text{for } a + \frac{b^2}{4c} \neq 0 \\ \frac{(\frac{b}{2c} + x) \log(\frac{b}{2c} + x)}{\sqrt{c}(\frac{b}{2c} + x)^2} & \text{otherwise} \end{cases} + \left(\frac{b}{4c} + \frac{x}{2} \right) \sqrt{-a + bx + cx^2} & \text{for } c \neq 0 \\ \frac{2(-a + bx)^{\frac{3}{2}}}{3b} & \text{for } b \neq 0 \\ x\sqrt{-a} & \text{otherwise} \end{cases}$$

input

```
integrate(d+e*x+f*(c*x**2+b*x-a)**(1/2),x)
```

output

```
d*x + e*x**2/2 + f*Piecewise((( -a/2 - b**2/(8*c))*Piecewise((log(b + 2*sqrt(c)*sqrt(-a + b*x + c*x**2) + 2*c*x)/sqrt(c), Ne(a + b**2/(4*c), 0)), ((b/(2*c) + x)*log(b/(2*c) + x)/sqrt(c*(b/(2*c) + x)**2), True)) + (b/(4*c) + x/2)*sqrt(-a + b*x + c*x**2), Ne(c, 0)), (2*(-a + b*x)**(3/2)/(3*b), Ne(b, 0)), (x*sqrt(-a), True))
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.27

$$\int (d + ex + f\sqrt{-a + bx + cx^2}) dx = \frac{1}{2} ex^2 + \frac{1}{8} \left(4\sqrt{cx^2 + bx - ax} - \frac{b^2 \log(2cx + b + 2\sqrt{cx^2 + bx - a}\sqrt{c})}{c^{\frac{3}{2}}} - \frac{4a \log(2cx + b + 2\sqrt{cx^2 + bx - a}\sqrt{c})}{\sqrt{c}} + dx \right)$$

input `integrate(d+e*x+f*(c*x^2+b*x-a)^(1/2),x, algorithm="maxima")`output `1/2*e*x^2 + 1/8*(4*sqrt(c*x^2 + b*x - a)*x - b^2*log(2*c*x + b + 2*sqrt(c*x^2 + b*x - a)*sqrt(c))/c^(3/2) - 4*a*log(2*c*x + b + 2*sqrt(c*x^2 + b*x - a)*sqrt(c))/sqrt(c) + 2*sqrt(c*x^2 + b*x - a)*b/c)*f + d*x`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.89

$$\int (d + ex + f\sqrt{-a + bx + cx^2}) dx = \frac{1}{2} ex^2 + \frac{1}{8} \left(2\sqrt{cx^2 + bx - a} \left(2x + \frac{b}{c} \right) + \frac{(b^2 + 4ac) \log(|2(\sqrt{cx} - \sqrt{cx^2 + bx - a})\sqrt{c} + b|)}{c^{\frac{3}{2}}} \right) f + dx$$

input `integrate(d+e*x+f*(c*x^2+b*x-a)^(1/2),x, algorithm="giac")`output `1/2*e*x^2 + 1/8*(2*sqrt(c*x^2 + b*x - a)*(2*x + b/c) + (b^2 + 4*a*c)*log(abs(2*(sqrt(c)*x - sqrt(c*x^2 + b*x - a))*sqrt(c) + b))/c^(3/2))*f + d*x`

Mupad [B] (verification not implemented)

Time = 22.96 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.85

$$\int \left(d + ex + f\sqrt{-a + bx + cx^2} \right) dx = dx + \frac{ex^2}{2} + f \left(\frac{x}{2} + \frac{b}{4c} \right) \sqrt{cx^2 + bx - a} - \frac{f \ln \left(\sqrt{cx^2 + bx - a} + \frac{\frac{b}{2} + cx}{\sqrt{c}} \right) \left(\frac{b^2}{4} + ac \right)}{2c^{3/2}}$$

input `int(d + f*(b*x - a + c*x^2)^(1/2) + e*x,x)`output `d*x + (e*x^2)/2 + f*(x/2 + b/(4*c))*(b*x - a + c*x^2)^(1/2) - (f*log((b*x - a + c*x^2)^(1/2) + (b/2 + c*x)/c^(1/2))*(a*c + b^2/4))/(2*c^(3/2))`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.60

$$\int \left(d + ex + f\sqrt{-a + bx + cx^2} \right) dx = \frac{2\sqrt{cx^2 + bx - a}bcf + 4\sqrt{cx^2 + bx - a}c^2fx - 4\sqrt{c} \log \left(\frac{2\sqrt{c}\sqrt{cx^2 + bx - a} + b + 2cx}{\sqrt{4ac + b^2}} \right) acf - \sqrt{c} \log \left(\frac{2\sqrt{c}\sqrt{cx^2 + bx - a} + b + 2cx}{\sqrt{4ac + b^2}} \right) acf}{8c^2}$$

input `int(d+e*x+f*(c*x^2+b*x-a)^(1/2),x)`output `(2*sqrt(-a + b*x + c*x**2)*b*c*f + 4*sqrt(-a + b*x + c*x**2)*c**2*f*x - 4*sqrt(c)*log((2*sqrt(c)*sqrt(-a + b*x + c*x**2) + b + 2*c*x)/sqrt(4*a*c + b**2))*a*c*f - sqrt(c)*log((2*sqrt(c)*sqrt(-a + b*x + c*x**2) + b + 2*c*x)/sqrt(4*a*c + b**2))*b**2*f + 8*c**2*d*x + 4*c**2*e*x**2)/(8*c**2)`

3.38 $\int \frac{1}{d+ex+f\sqrt{-a+bx+cx^2}} dx$

Optimal result	319
Mathematica [A] (verified)	320
Rubi [B] (verified)	320
Maple [B] (warning: unable to verify)	322
Fricas [F(-1)]	323
Sympy [F]	324
Maxima [F]	324
Giac [F(-2)]	324
Mupad [F(-1)]	325
Reduce [F]	325

Optimal result

Integrand size = 25, antiderivative size = 300

$$\int \frac{1}{d+ex+f\sqrt{-a+bx+cx^2}} dx$$

$$= -\frac{2(2cd-be)f \operatorname{arctanh}\left(\frac{bf+2cfx+2e\sqrt{-a+x(b+cx)}+2\sqrt{c}(d+ex+f\sqrt{-a+x(b+cx)})}{\sqrt{-4bde-4ae^2+b^2f^2+4c(d^2+af^2)}}\right)}{(e^2-cf^2)\sqrt{-4bde-4ae^2+b^2f^2+4c(d^2+af^2)}} - \frac{\log\left(b+2cx+2\sqrt{c}\sqrt{-a+x(b+cx)}\right)}{e-\sqrt{cf}} + \frac{e \log\left(bd+a(e-\sqrt{cf})+(2\sqrt{cd}+bf)(\sqrt{cx}+\sqrt{-a+bx+cx^2})+(e+\sqrt{cf})(\sqrt{cx}+\sqrt{-a+bx+cx^2})\right)}{e^2-cf^2}$$

output

```
-2*(-b*e+2*c*d)*f*arctanh((b*f+2*c*f*x+2*e*(-a+x*(c*x+b))^(1/2)+2*c^(1/2)*(d+e*x+f*(-a+x*(c*x+b))^(1/2)))/(-4*b*d*e-4*a*e^2+b^2*f^2+4*c*(a*f^2+d^2))^(1/2))/(-c*f^2+e^2)/(-4*b*d*e-4*a*e^2+b^2*f^2+4*c*(a*f^2+d^2))^(1/2)-ln(b+2*c*x+2*c^(1/2)*(-a+x*(c*x+b))^(1/2))/(e-c^(1/2)*f)+e*ln(b*d+a*(e-c^(1/2)*f)+(2*c^(1/2)*d+b*f)*(c^(1/2)*x+(c*x^2+b*x-a)^(1/2)))+(e+c^(1/2)*f)*(c^(1/2)*x+(c*x^2+b*x-a)^(1/2))^2)/(-c*f^2+e^2)
```


Mathematica [A] (verified)

Time = 1.73 (sec) , antiderivative size = 287, normalized size of antiderivative = 0.96

$$\int \frac{1}{d + ex + f\sqrt{-a + bx + cx^2}} dx$$

$$= \frac{2(2cd - be)f \arctan\left(\frac{bf + 2cfx + 2e\sqrt{-a + x(b + cx)} - 2\sqrt{c}(d + ex + f\sqrt{-a + x(b + cx)})}{\sqrt{4bde + 4ae^2 - b^2f^2 - 4c(d^2 + af^2)}}\right)}{\sqrt{4bde + 4ae^2 - b^2f^2 - 4c(d^2 + af^2)}} + (e - \sqrt{cf}) \log\left(b + 2cx - 2\sqrt{c}\sqrt{-a + x(b + cx)}\right)$$

input

```
Integrate[(d + e*x + f*Sqrt[-a + b*x + c*x^2])^(-1), x]
```

output

```
((-2*(2*c*d - b*e)*f*ArcTan[(b*f + 2*c*f*x + 2*e*Sqrt[-a + x*(b + c*x)] - 2*Sqrt[c]*(d + e*x + f*Sqrt[-a + x*(b + c*x)])]/Sqrt[4*b*d*e + 4*a*e^2 - b^2*f^2 - 4*c*(d^2 + a*f^2)])]/Sqrt[4*b*d*e + 4*a*e^2 - b^2*f^2 - 4*c*(d^2 + a*f^2)] + (e - Sqrt[c]*f)*Log[b + 2*c*x - 2*Sqrt[c]*Sqrt[-a + x*(b + c*x)]] - e*Log[b*(d + e*x - 2*Sqrt[c]*f*x + f*Sqrt[-a + x*(b + c*x)]) + 2*Sqrt[c]*(a*f + (-d - e*x + Sqrt[c]*f*x)*(-(Sqrt[c]*x) + Sqrt[-a + x*(b + c*x)])))]/(-e^2 + c*f^2)
```

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1084 vs. 2(300) = 600.

Time = 10.30 (sec) , antiderivative size = 1084, normalized size of antiderivative = 3.61, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{f\sqrt{-a + bx + cx^2} + d + ex} dx$$

↓ 7293

$$\int \left(\frac{f\sqrt{-a+bx+cx^2}}{-af^2-x(2de-bf^2)-x^2(e^2-cf^2)-d^2} + \frac{d+ex}{af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2} \right) dx$$

↓ 2009

$$\frac{(2cd-be)f \operatorname{arctanh}\left(\frac{-bf^2+2de+2(e^2-cf^2)x}{f\sqrt{-4ae^2-4bde+b^2f^2+4c(d^2+af^2)}}\right)}{(e^2-cf^2)\sqrt{-4ae^2-4bde+b^2f^2+4c(d^2+af^2)}} - \frac{\sqrt{c}f \operatorname{arctanh}\left(\frac{b+2cx}{2\sqrt{c}\sqrt{cx^2+bx-a}}\right)}{e^2-cf^2} -$$

$$\frac{\sqrt{-2ae^4-2bde^3+2cd^2e^2+b^2f^2e^2+2acf^2e^2-2bcd f^2e-(2cd-be)f\sqrt{-4ae^2-4bde+b^2f^2+4c(d^2+af^2)}}}{\sqrt{2}(e^2-cf^2)\sqrt{-4a}}$$

$$\frac{\sqrt{-2ae^4-2bde^3+2cd^2e^2+b^2f^2e^2+2acf^2e^2-2bcd f^2e+(2cd-be)f\sqrt{-4ae^2-4bde+b^2f^2+4c(d^2+af^2)}}}{\sqrt{2}(e^2-cf^2)\sqrt{-4a}}$$

$$\frac{e \log(d^2+af^2+(e^2-cf^2)x^2+(2de-bf^2)x)}{2(e^2-cf^2)}$$

input

```
Int[(d + e*x + f*Sqrt[-a + b*x + c*x^2])^(-1),x]
```

output

```

((2*c*d - b*e)*f*ArcTanh[(2*d*e - b*f^2 + 2*(e^2 - c*f^2)*x)/(f*Sqrt[-4*b*d
e - 4*a*e^2 + b^2*f^2 + 4*c*(d^2 + a*f^2)])])/(e^2 - c*f^2)*Sqrt[-4*b*d
*e - 4*a*e^2 + b^2*f^2 + 4*c*(d^2 + a*f^2)] - (Sqrt[c]*f*ArcTanh[(b + 2*c
*x)/(2*Sqrt[c]*Sqrt[-a + b*x + c*x^2])])/(e^2 - c*f^2) - (Sqrt[2*c*d^2*e^2
- 2*b*d*e^3 - 2*a*e^4 + 2*c^2*d^2*f^2 - 2*b*c*d*e*f^2 + b^2*e^2*f^2 + 2*a
*c*e^2*f^2 - e*(2*c*d - b*e)*f*Sqrt[-4*b*d*e - 4*a*e^2 + b^2*f^2 + 4*c*(d^
2 + a*f^2)])*ArcTanh[(2*b*d*e + 4*a*e^2 - b^2*f^2 - 4*a*c*f^2 - b*f*Sqrt[-
4*b*d*e - 4*a*e^2 + b^2*f^2 + 4*c*(d^2 + a*f^2)] + 2*(2*c*d*e - b*e^2 - c*
f*Sqrt[-4*b*d*e - 4*a*e^2 + b^2*f^2 + 4*c*(d^2 + a*f^2)]*x)/(2*Sqrt[2]*Sq
rt[2*c*d^2*e^2 - 2*b*d*e^3 - 2*a*e^4 + 2*c^2*d^2*f^2 - 2*b*c*d*e*f^2 + b^2
*e^2*f^2 + 2*a*c*e^2*f^2 - e*(2*c*d - b*e)*f*Sqrt[-4*b*d*e - 4*a*e^2 + b^2
*f^2 + 4*c*(d^2 + a*f^2)])*Sqrt[-a + b*x + c*x^2])])/(Sqrt[2]*(e^2 - c*f^2
)*Sqrt[-4*b*d*e - 4*a*e^2 + b^2*f^2 + 4*c*(d^2 + a*f^2)]) + (Sqrt[2*c*d^2*
e^2 - 2*b*d*e^3 - 2*a*e^4 + 2*c^2*d^2*f^2 - 2*b*c*d*e*f^2 + b^2*e^2*f^2 +
2*a*c*e^2*f^2 + e*(2*c*d - b*e)*f*Sqrt[-4*b*d*e - 4*a*e^2 + b^2*f^2 + 4*c*
(d^2 + a*f^2)])*ArcTanh[(2*b*d*e + 4*a*e^2 - b^2*f^2 - 4*a*c*f^2 + b*f*Sqr
t[-4*b*d*e - 4*a*e^2 + b^2*f^2 + 4*c*(d^2 + a*f^2)] + 2*(2*c*d*e - b*e^2 +
c*f*Sqrt[-4*b*d*e - 4*a*e^2 + b^2*f^2 + 4*c*(d^2 + a*f^2)]*x)/(2*Sqrt[2]
*Sqrt[2*c*d^2*e^2 - 2*b*d*e^3 - 2*a*e^4 + 2*c^2*d^2*f^2 - 2*b*c*d*e*f^2 +
b^2*e^2*f^2 + 2*a*c*e^2*f^2 + e*(2*c*d - b*e)*f*Sqrt[-4*b*d*e - 4*a*e^2...

```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 7293

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 4828 vs. $2(270) = 540$.

Time = 0.13 (sec) , antiderivative size = 4829, normalized size of antiderivative = 16.10

method	result	size
default	Expression too large to display	4829

input `int(1/(d+e*x+f*(c*x^2+b*x-a)^(1/2)),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & f*(-2*(c*f^2-e^2)/(f^2*(4*a*c*f^2+b^2*f^2-4*a*e^2-4*b*d*e+4*c*d^2))^(1/2)/ \\ & (2*c*f^2-2*e^2)*(1/2*(4*(x+(b*f^2-2*d*e+(f^2*(4*a*c*f^2+b^2*f^2-4*a*e^2-4* \\ & b*d*e+4*c*d^2))^(1/2))/(2*c*f^2-2*e^2))^2*c-4*(b*e^2-2*c*d*e+(f^2*(4*a*c*f \\ & ^2+b^2*f^2-4*a*e^2-4*b*d*e+4*c*d^2))^(1/2)*c)/(c*f^2-e^2)*(x+(b*f^2-2*d*e+ \\ & (f^2*(4*a*c*f^2+b^2*f^2-4*a*e^2-4*b*d*e+4*c*d^2))^(1/2)))/(2*c*f^2-2*e^2))+ \\ & 2*(2*a*c*e^2*f^2+b^2*e^2*f^2-2*b*c*d*e*f^2+2*c^2*d^2*f^2-2*e^4*a-2*d*e^3*b \\ & +2*d^2*e^2*c+(f^2*(4*a*c*f^2+b^2*f^2-4*a*e^2-4*b*d*e+4*c*d^2))^(1/2)*b*e^2 \\ & -2*(f^2*(4*a*c*f^2+b^2*f^2-4*a*e^2-4*b*d*e+4*c*d^2))^(1/2)*c*d*e)/(c*f^2-e \\ & ^2)^2)^(1/2)-1/2*(b*e^2-2*c*d*e+(f^2*(4*a*c*f^2+b^2*f^2-4*a*e^2-4*b*d*e+4* \\ & c*d^2))^(1/2)*c)/(c*f^2-e^2)*ln((-1/2*(b*e^2-2*c*d*e+(f^2*(4*a*c*f^2+b^2*f \\ & ^2-4*a*e^2-4*b*d*e+4*c*d^2))^(1/2)*c)/(c*f^2-e^2)+c*(x+(b*f^2-2*d*e+(f^2*(\\ & 4*a*c*f^2+b^2*f^2-4*a*e^2-4*b*d*e+4*c*d^2))^(1/2)))/(2*c*f^2-2*e^2)))/c^(1/ \\ & 2))+((x+(b*f^2-2*d*e+(f^2*(4*a*c*f^2+b^2*f^2-4*a*e^2-4*b*d*e+4*c*d^2))^(1/2) \\ &))/(2*c*f^2-2*e^2))^2*c-(b*e^2-2*c*d*e+(f^2*(4*a*c*f^2+b^2*f^2-4*a*e^2-4*b \\ & *d*e+4*c*d^2))^(1/2)*c)/(c*f^2-e^2)*(x+(b*f^2-2*d*e+(f^2*(4*a*c*f^2+b^2*f^ \\ & 2-4*a*e^2-4*b*d*e+4*c*d^2))^(1/2)))/(2*c*f^2-2*e^2))+1/2*(2*a*c*e^2*f^2+b^2 \\ & *e^2*f^2-2*b*c*d*e*f^2+2*c^2*d^2*f^2-2*e^4*a-2*d*e^3*b+2*d^2*e^2*c+(f^2*(4 \\ & *a*c*f^2+b^2*f^2-4*a*e^2-4*b*d*e+4*c*d^2))^(1/2)*b*e^2-2*(f^2*(4*a*c*f^2+b \\ & ^2*f^2-4*a*e^2-4*b*d*e+4*c*d^2))^(1/2)*c*d*e)/(c*f^2-e^2)^2)^(1/2))/c^(1/2) \\ &)-1/2*(2*a*c*e^2*f^2+b^2*e^2*f^2-2*b*c*d*e*f^2+2*c^2*d^2*f^2-2*e^4*a-2*... \end{aligned}$$

Fricas [F(-1)]

Timed out.

$$\int \frac{1}{d+ex+f\sqrt{-a+bx+cx^2}} dx = \text{Timed out}$$

input `integrate(1/(d+e*x+f*(c*x^2+b*x-a)^(1/2)),x, algorithm="fricas")`

output Timed out

Sympy [F]

$$\int \frac{1}{d + ex + f\sqrt{-a + bx + cx^2}} dx = \int \frac{1}{d + ex + f\sqrt{-a + bx + cx^2}} dx$$

input `integrate(1/(d+e*x+f*(c*x**2+b*x-a)**(1/2)),x)`

output `Integral(1/(d + e*x + f*sqrt(-a + b*x + c*x**2)), x)`

Maxima [F]

$$\int \frac{1}{d + ex + f\sqrt{-a + bx + cx^2}} dx = \int \frac{1}{ex + \sqrt{cx^2 + bx - a}f + d} dx$$

input `integrate(1/(d+e*x+f*(c*x^2+b*x-a)^(1/2)),x, algorithm="maxima")`

output `integrate(1/(e*x + sqrt(c*x^2 + b*x - a)*f + d), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{1}{d + ex + f\sqrt{-a + bx + cx^2}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(d+e*x+f*(c*x^2+b*x-a)^(1/2)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Error: Bad Argument Type`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{d + ex + f\sqrt{-a + bx + cx^2}} dx = \int \frac{1}{d + f\sqrt{cx^2 + bx - a} + ex} dx$$

input `int(1/(d + f*(b*x - a + c*x^2)^(1/2) + e*x), x)`output `int(1/(d + f*(b*x - a + c*x^2)^(1/2) + e*x), x)`**Reduce [F]**

$$\int \frac{1}{d + ex + f\sqrt{-a + bx + cx^2}} dx = \int \frac{1}{d + ex + f\sqrt{cx^2 + bx - a}} dx$$

input `int(1/(d+e*x+f*(c*x^2+b*x-a)^(1/2)), x)`output `int(1/(d+e*x+f*(c*x^2+b*x-a)^(1/2)), x)`

3.39 $\int \frac{1}{\left(d+ex+f\sqrt{-a+bx+cx^2}\right)^2} dx$

Optimal result	326
Mathematica [C] (warning: unable to verify)	327
Rubi [F]	328
Maple [B] (warning: unable to verify)	329
Fricas [B] (verification not implemented)	329
Sympy [F]	330
Maxima [F]	330
Giac [F(-2)]	330
Mupad [F(-1)]	331
Reduce [F]	331

Optimal result

Integrand size = 25, antiderivative size = 385

$$\int \frac{1}{\left(d+ex+f\sqrt{-a+bx+cx^2}\right)^2} dx$$

$$= \frac{2(2b(cd^2 - ae^2) - 4ac^{3/2}df - b^2d(2e + \sqrt{cf}) + (4c^{3/2}d^2 - 4\sqrt{ce}(bd + ae) - b^2ef - 4ae^2))}{(e + \sqrt{cf})(4bde + 4ae^2 - b^2f^2 - 4c(d^2 + af^2))} \left(bd + a(e - \sqrt{cf}) + (2\sqrt{cd} + bf)(\sqrt{cx} + \sqrt{-a + bx + cx^2}) \right) - \frac{4(b^2 + 4ac) \operatorname{farctanh}\left(\frac{bf + 2cfx + 2e\sqrt{-a + x(b + cx)} + 2\sqrt{c}(d + ex + f\sqrt{-a + x(b + cx)})}{\sqrt{-4bde - 4ae^2 + b^2f^2 + 4c(d^2 + af^2)}}\right)}{(-4bde - 4ae^2 + b^2f^2 + 4c(d^2 + af^2))^{3/2}}$$

output

```
2*(2*b*(-a*e^2+c*d^2)-4*a*c^(3/2)*d*f-b^2*d*(2*e+c^(1/2)*f)+(4*c^(3/2)*d^2-4*c^(1/2)*e*(a*e+b*d)-b^2*e*f-4*a*c*e*f)*(c^(1/2)*x+(c*x^2+b*x-a)^(1/2)))/(e+c^(1/2)*f)/(4*b*d*e+4*a*e^2-b^2*f^2-4*c*(a*f^2+d^2))/(b*d+a*(e-c^(1/2)*f)+(2*c^(1/2)*d+b*f)*(c^(1/2)*x+(c*x^2+b*x-a)^(1/2))+(e+c^(1/2)*f)*(c^(1/2)*x+(c*x^2+b*x-a)^(1/2))^2-4*(4*a*c+b^2)*f*arctanh((b*f+2*c*f*x+2*e*(-a+x*(c*x+b))^(1/2)+2*c^(1/2)*(d+e*x+f*(-a+x*(c*x+b))^(1/2)))/(-4*b*d*e-4*a*e^2+b^2*f^2+4*c*(a*f^2+d^2))^(1/2))/(-4*b*d*e-4*a*e^2+b^2*f^2+4*c*(a*f^2+d^2))^(3/2)
```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 50.69 (sec) , antiderivative size = 2401, normalized size of antiderivative = 6.24

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx + cx^2})^2} dx = \text{Result too large to show}$$

input `Integrate[(d + e*x + f*Sqrt[-a + b*x + c*x^2])^(-2),x]`

output

```
(-2*(-2*c*d^3*e + 2*b*d^2*e^2 + 2*a*d*e^3 - b*c*d^2*f^2 - 4*a*c*d*e*f^2 +
a*b*e^2*f^2 - 2*c*d^2*e^2*x + 2*b*d*e^3*x + 2*a*e^4*x - 2*c^2*d^2*f^2*x +
2*b*c*d*e*f^2*x - b^2*e^2*f^2*x - 2*a*c*e^2*f^2*x))/((e^2 - c*f^2)*(-4*c*d
^2 + 4*b*d*e + 4*a*e^2 - b^2*f^2 - 4*a*c*f^2)*(d^2 + a*f^2 + 2*d*e*x - b*f
^2*x + e^2*x^2 - c*f^2*x^2)) + (2*(b*d*f + 2*a*e*f + 2*c*d*f*x - b*e*f*x)*
Sqrt[-a + b*x + c*x^2])/((-4*c*d^2 + 4*b*d*e + 4*a*e^2 - b^2*f^2 - 4*a*c*f
^2)*(d^2 + a*f^2 + 2*d*e*x - b*f^2*x + e^2*x^2 - c*f^2*x^2)) + (2*(b^2 + 4
*a*c)*f*ArcTan[(2*d*e - b*f^2 + 2*e^2*x - 2*c*f^2*x)/(f*Sqrt[-4*c*d^2 + 4*
b*d*e + 4*a*e^2 - b^2*f^2 - 4*a*c*f^2])])/(-4*c*d^2 + 4*b*d*e + 4*a*e^2 -
b^2*f^2 - 4*a*c*f^2)^(3/2) + ((b^2 + 4*a*c)*f*((-2*I)*c*d*f + I*b*e*f + e
Sqrt[-4*c*d^2 + 4*b*d*e + 4*a*e^2 - b^2*f^2 - 4*a*c*f^2])*Log[-((Sqrt[2]*(
e^2 - c*f^2)*(-4*c*d^2 + 4*b*d*e + 4*a*e^2 - b^2*f^2 - 4*a*c*f^2)^(3/2))*((-
2*I)*b*d*e - (4*I)*a*e^2 + I*b^2*f^2 + (4*I)*a*c*f^2 + b*f*Sqrt[-4*c*d^2
+ 4*b*d*e + 4*a*e^2 - b^2*f^2 - 4*a*c*f^2] - (4*I)*c*d*e*x + (2*I)*b*e^2*x
+ 2*c*f*Sqrt[-4*c*d^2 + 4*b*d*e + 4*a*e^2 - b^2*f^2 - 4*a*c*f^2]*x))/((b^
2 + 4*a*c)*f*(-2*c*d*f + b*e*f - I*e*Sqrt[-4*c*d^2 + 4*b*d*e + 4*a*e^2 - b
^2*f^2 - 4*a*c*f^2])*Sqrt[2*c*d^2*e^2 - 2*b*d*e^3 - 2*a*e^4 + 2*c^2*d^2*f^
2 - 2*b*c*d*e*f^2 + b^2*e^2*f^2 + 2*a*c*e^2*f^2 + (2*I)*c*d*e*f*Sqrt[-4*c*
d^2 + 4*b*d*e + 4*a*e^2 - b^2*f^2 - 4*a*c*f^2] - I*b*e^2*f*Sqrt[-4*c*d^2 +
4*b*d*e + 4*a*e^2 - b^2*f^2 - 4*a*c*f^2])*(2*d*e - b*f^2 + I*f*Sqrt[-4...
```


Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\left(f\sqrt{-a+bx+cx^2}+d+ex\right)^2} dx$$

↓ 7293

$$\int \left(\frac{2f^2(-a+bx+cx^2)}{(af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2)^2} - \frac{2df\sqrt{-a+bx+cx^2}}{(af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2)^2} - \frac{2e}{(af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2)^2} \right) dx$$

↓ 7239

$$\int \frac{2d\left(ex-f\sqrt{x(b+cx)-a}\right)+x\left(-2ef\sqrt{x(b+cx)-a}+bf^2+cf^2x+e^2x\right)+d^2\left(1-\frac{af^2}{d^2}\right)}{(af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2)^2} dx$$

↓ 7293

$$\int \left(\frac{d^2-af^2}{(af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2)^2} + \frac{2d\left(ex-f\sqrt{-a+bx+cx^2}\right)}{(af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2)^2} + \frac{x\left(-2ef\sqrt{-a+bx+cx^2}\right)}{(af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2)^2} \right) dx$$

↓ 7299

$$\int \left(\frac{d^2-af^2}{(af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2)^2} + \frac{2d\left(ex-f\sqrt{-a+bx+cx^2}\right)}{(af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2)^2} + \frac{x\left(-2ef\sqrt{-a+bx+cx^2}\right)}{(af^2+x(2de-bf^2)+x^2(e^2-cf^2)+d^2)^2} \right) dx$$

input

```
Int[(d + e*x + f*Sqrt[-a + b*x + c*x^2])^(-2),x]
```

output

```
$Aborted
```

Defintions of rubi rules used

rule 7239 `Int[u_, x_Symbol] :=> With[{v = SimplifyIntegrand[u, x]}, Int[v, x] /; SimplifyIntegrandQ[v, u, x]]`

rule 7293 `Int[u_, x_Symbol] :=> With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

rule 7299 `Int[u_, x_] :=> CannotIntegrate[u, x]`

Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.25 (sec) , antiderivative size = 508167, normalized size of antiderivative = 1319.91

method	result	size
default	Expression too large to display	508167

input `int(1/(d+e*x+f*(c*x^2+b*x-a)^(1/2))^2,x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2877 vs. $2(346) = 692$.

Time = 20.29 (sec) , antiderivative size = 6919, normalized size of antiderivative = 17.97

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx + cx^2})^2} dx = \text{Too large to display}$$

input `integrate(1/(d+e*x+f*(c*x^2+b*x-a)^(1/2))^2,x, algorithm="fricas")`

output Too large to include

Sympy [F]

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx + cx^2})^2} dx = \int \frac{1}{(d + ex + f\sqrt{-a + bx + cx^2})^2} dx$$

input `integrate(1/(d+e*x+f*(c*x**2+b*x-a)**(1/2))**2,x)`

output `Integral((d + e*x + f*sqrt(-a + b*x + c*x**2))**(-2), x)`

Maxima [F]

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx + cx^2})^2} dx = \int \frac{1}{(ex + \sqrt{cx^2 + bx - a}f + d)^2} dx$$

input `integrate(1/(d+e*x+f*(c*x^2+b*x-a)^(1/2))^2,x, algorithm="maxima")`

output `integrate((e*x + sqrt(c*x^2 + b*x - a)*f + d)^(-2), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx + cx^2})^2} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(d+e*x+f*(c*x^2+b*x-a)^(1/2))^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:Error: Bad Argument Type`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx + cx^2})^2} dx = \int \frac{1}{(d + f\sqrt{cx^2 + bx - a} + ex)^2} dx$$

input `int(1/(d + f*(b*x - a + c*x^2)^(1/2) + e*x)^2,x)`output `int(1/(d + f*(b*x - a + c*x^2)^(1/2) + e*x)^2, x)`**Reduce [F]**

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx + cx^2})^2} dx = \int \frac{1}{(d + ex + f\sqrt{cx^2 + bx - a})^2} dx$$

input `int(1/(d+e*x+f*(c*x^2+b*x-a)^(1/2))^2,x)`output `int(1/(d+e*x+f*(c*x^2+b*x-a)^(1/2))^2,x)`

$$3.40 \quad \int \frac{1}{\left(d+ex+f\sqrt{-a+bx+cx^2}\right)^3} dx$$

Optimal result	332
Mathematica [C] (warning: unable to verify)	333
Rubi [F]	334
Maple [B] (warning: unable to verify)	335
Fricas [F(-1)]	336
Sympy [F]	336
Maxima [F]	336
Giac [F(-2)]	337
Mupad [F(-1)]	337
Reduce [F]	337

Optimal result

Integrand size = 25, antiderivative size = 896

$$\int \frac{1}{\left(d+ex+f\sqrt{-a+bx+cx^2}\right)^3} dx =$$

$$\frac{b^3d(2e^2 + \sqrt{c}ef + cf^2) + 4bcd(2cd^2 - 4ae^2 + a\sqrt{c}ef + acf^2) - 2b^2(5cd^2e - ae^3 + c^{3/2}d^2f - acef^2) - (e + \sqrt{c}f)^2(4bde + 4ae^2 - b^2f^2 - 4c(d^2 + af^2))(bd + a}{8c(e + \sqrt{c}f)(bd + a(e - \sqrt{c}f))(4bde + 4ae^2 - b^2f^2 - 4c(d^2 + af^2)) + (2\sqrt{cd} + bf)(3b^3e^2f + 3b^3\sqrt{c}ef^2 - 2bcf(7bde - 2ae^2 - b^2f^2) + 8c^{5/2}(2d^3 - adf))} + \frac{(4bde + 4ae^2 - b^2f^2 - 4c(d^2 + af^2))^2 (bd + a(e - \sqrt{c}f) + (2\sqrt{cd} + bf))}{12(b^2 + 4ac)(2cd - be) \operatorname{farctanh}\left(\frac{bf + 2cfx + 2e\sqrt{-a+x(b+cx)} + 2\sqrt{c}(d+ex+f\sqrt{-a+x(b+cx)})}{\sqrt{-4bde - 4ae^2 + b^2f^2 + 4c(d^2 + af^2)}}\right)} + \frac{(-4bde - 4ae^2 + b^2f^2 + 4c(d^2 + af^2))^{5/2}}$$

output

```

-(b^3*d*(2*e^2+c^(1/2)*e*f+c*f^2)+4*b*c*d*(2*c*d^2-4*a*e^2+a*c^(1/2)*e*f+a
*c*f^2)-2*b^2*(5*c*d^2*e-a*e^3+c^(3/2)*d^2*f-a*c*e*f^2)-8*a*c*(a*e^3+c^(3/
2)*d^2*f-c*e*(a*f^2+d^2))-(-b*e+2*c*d)*(b^2*e*f+4*a*c*e*f-4*c^(3/2)*(a*f^2
+2*d^2)+c^(1/2)*(-b^2*f^2+8*a*e^2+8*b*d*e))*(c^(1/2)*x+(c*x^2+b*x-a)^(1/2)
))/((e+c^(1/2)*f)^2/(4*b*d*e+4*a*e^2-b^2*f^2-4*c*(a*f^2+d^2)))/(b*d+a*(e-c^(
1/2)*f)+(2*c^(1/2)*d+b*f)*(c^(1/2)*x+(c*x^2+b*x-a)^(1/2)))+(e+c^(1/2)*f)*(c
^(1/2)*x+(c*x^2+b*x-a)^(1/2))^2)-((8*c*(e+c^(1/2)*f)*(b*d+a*(e-c^(1/2)*f
))*(4*b*d*e+4*a*e^2-b^2*f^2-4*c*(a*f^2+d^2)))+(2*c^(1/2)*d+b*f)*(3*b^3*e^2*
f+3*b^3*c^(1/2)*e*f^2-2*b*c*f*(-b^2*f^2-2*a*e^2+7*b*d*e)+8*c^(5/2)*(-a*d*f
^2+2*d^3)-2*c^(3/2)*(-6*a*b*e*f^2+b^2*d*f^2+8*a*d*e^2+8*b*d^2*e)-8*c^2*f*(
3*a*d*e-b*(a*f^2+d^2))))/(e+c^(1/2)*f)^2-6*(4*a*c+b^2)*(-b*e+2*c*d)*f*(c^(
1/2)*x+(c*x^2+b*x-a)^(1/2)))/(4*b*d*e+4*a*e^2-b^2*f^2-4*c*(a*f^2+d^2))^2/(
b*d+a*(e-c^(1/2)*f)+(2*c^(1/2)*d+b*f)*(c^(1/2)*x+(c*x^2+b*x-a)^(1/2)))+(e+c
^(1/2)*f)*(c^(1/2)*x+(c*x^2+b*x-a)^(1/2))^2)-12*(4*a*c+b^2)*(-b*e+2*c*d)*f
*arctanh((b*f+2*c*f*x+2*e*(-a+x*(c*x+b))^(1/2)+2*c^(1/2)*(d+e*x+f*(-a+x*(c
*x+b))^(1/2)))/(-4*b*d*e-4*a*e^2+b^2*f^2+4*c*(a*f^2+d^2))^(1/2))/(-4*b*d*e
-4*a*e^2+b^2*f^2+4*c*(a*f^2+d^2))^(5/2)

```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 75.18 (sec) , antiderivative size = 3628, normalized size of antiderivative = 4.05

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx + cx^2})^3} dx = \text{Result too large to show}$$

input

```
Integrate[(d + e*x + f*Sqrt[-a + b*x + c*x^2])^(-3),x]
```

output

```
(2*(-4*c^2*d^4*e*f^2 + 5*b*c*d^3*e^2*f^2 - b^2*d^2*e^3*f^2 + 2*a*c*d^2*e^3*f^2 + a*b*d*e^4*f^2 + 2*a^2*e^5*f^2 - b*c^2*d^3*f^4 - 6*a*c^2*d^2*e*f^4 + 3*a*b*c*d*e^2*f^4 - a*b^2*e^3*f^4 - 2*a^2*c*e^3*f^4 - 6*c^2*d^3*e^2*f^2*x + 9*b*c*d^2*e^3*f^2*x - 3*b^2*d*e^4*f^2*x + 6*a*c*d*e^4*f^2*x - 3*a*b*e^5*f^2*x - 2*c^3*d^3*f^4*x + 3*b*c^2*d^2*e*f^4*x - 3*b^2*c*d*e^2*f^4*x - 6*a*c^2*d*e^2*f^4*x + b^3*e^3*f^4*x + 3*a*b*c*e^3*f^4*x))/((e^2 - c*f^2)^2*(-4*c*d^2 + 4*b*d*e + 4*a*e^2 - b^2*f^2 - 4*a*c*f^2)*(d^2 + a*f^2 + 2*d*e*x - b*f^2*x + e^2*x^2 - c*f^2*x^2)^2) + (-8*c^2*d^4*e^3 + 16*b*c*d^3*e^4 - 8*b^2*d^2*e^5 + 16*a*c*d^2*e^5 - 16*a*b*d*e^6 - 8*a^2*e^7 - 24*c^3*d^4*e*f^2 + 48*b*c^2*d^3*e^2*f^2 - 34*b^2*c*d^2*e^3*f^2 + 8*a*c^2*d^2*e^3*f^2 + 7*b^3*d*e^4*f^2 - 20*a*b*c*d*e^4*f^2 + 4*a*b^2*e^5*f^2 - 8*a^2*c*e^5*f^2 - 6*b^2*c^2*d^2*e*f^4 - 24*a*c^3*d^2*e*f^4 + 12*b^3*c*d*e^2*f^4 + 48*a*b*c^2*d*e^2*f^4 - 2*b^4*e^3*f^4 + 2*a*b^2*c*e^3*f^4 + 40*a^2*c^2*e^3*f^4 - 3*b^3*c^2*d*f^6 - 12*a*b*c^3*d*f^6 - 6*a*b^2*c^2*e*f^6 - 24*a^2*c^3*e*f^6 - 6*b^2*c*d*e^4*f^2*x - 24*a*c^2*d*e^4*f^2*x + 3*b^3*e^5*f^2*x + 12*a*b*c*e^5*f^2*x + 12*b^2*c^2*d*e^2*f^4*x + 48*a*c^3*d*e^2*f^4*x - 6*b^3*c*e^3*f^4*x - 24*a*b*c^2*e^3*f^4*x - 6*b^2*c^3*d*f^6*x - 24*a*c^4*d*f^6*x + 3*b^3*c^2*e*f^6*x + 12*a*b*c^3*e*f^6*x))/((e^2 - c*f^2)^2*(-4*c*d^2 + 4*b*d*e + 4*a*e^2 - b^2*f^2 - 4*a*c*f^2)^2*(d^2 + a*f^2 + 2*d*e*x - b*f^2*x + e^2*x^2 - c*f^2*x^2)) + Sqrt[-a + b*x + c*x^2]*((2*(-2*c*d^3*e*f + 2*b*d^2*e^2*f + ...
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(f\sqrt{-a + bx + cx^2} + d + ex)^3} dx$$

↓ 7293

$$\int \left(\frac{4f^2(d + ex)(-a + bx + cx^2)}{(af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^3} - \frac{3f\sqrt{-a + bx + cx^2}}{(af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{1}{af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2} \right) dx$$

↓ 7299

$$\int \left(\frac{4f^2(d + ex)(-a + bx + cx^2)}{(af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^3} - \frac{3f\sqrt{-a + bx + cx^2}}{(af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2)^2} + \frac{1}{af^2 + x(2de - bf^2) + x^2(e^2 - cf^2) + d^2} \right) dx$$

input `Int[(d + e*x + f*Sqrt[-a + b*x + c*x^2])^(-3),x]`

output `$Aborted`

Defintions of rubi rules used

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

rule 7299 `Int[u_, x_] := CannotIntegrate[u, x]`

Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 114.04 (sec) , antiderivative size = 4527517, normalized size of antiderivative = 5053.03

method	result	size
default	Expression too large to display	4527517

input `int(1/(d+e*x+f*(c*x^2+b*x-a)^(1/2))^3,x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx + cx^2})^3} dx = \text{Timed out}$$

input `integrate(1/(d+e*x+f*(c*x^2+b*x-a)^(1/2))^3,x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx + cx^2})^3} dx = \int \frac{1}{(d + ex + f\sqrt{-a + bx + cx^2})^3} dx$$

input `integrate(1/(d+e*x+f*(c*x**2+b*x-a)**(1/2))**3,x)`

output `Integral((d + e*x + f*sqrt(-a + b*x + c*x**2))**(-3), x)`

Maxima [F]

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx + cx^2})^3} dx = \int \frac{1}{(ex + \sqrt{cx^2 + bx - a}f + d)^3} dx$$

input `integrate(1/(d+e*x+f*(c*x^2+b*x-a)^(1/2))^3,x, algorithm="maxima")`

output `integrate((e*x + sqrt(c*x^2 + b*x - a)*f + d)^(-3), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx + cx^2})^3} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(d+e*x+f*(c*x^2+b*x-a)^(1/2))^3,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Error: Bad Argument Type`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx + cx^2})^3} dx = \int \frac{1}{(d + f\sqrt{cx^2 + bx - a} + ex)^3} dx$$

input `int(1/(d + f*(b*x - a + c*x^2)^(1/2) + e*x)^3,x)`

output `int(1/(d + f*(b*x - a + c*x^2)^(1/2) + e*x)^3, x)`

Reduce [F]

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx + cx^2})^3} dx = \int \frac{1}{(d + ex + f\sqrt{cx^2 + bx - a})^3} dx$$

input `int(1/(d+e*x+f*(c*x^2+b*x-a)^(1/2))^3,x)`

output `int(1/(d+e*x+f*(c*x^2+b*x-a)^(1/2))^3,x)`

3.41 $\int (x + \sqrt{-3 - 4x - x^2})^2 dx$

Optimal result	338
Mathematica [A] (verified)	338
Rubi [A] (verified)	339
Maple [A] (verified)	340
Fricas [A] (verification not implemented)	340
Sympy [A] (verification not implemented)	341
Maxima [A] (verification not implemented)	341
Giac [A] (verification not implemented)	341
Mupad [B] (verification not implemented)	342
Reduce [B] (verification not implemented)	342

Optimal result

Integrand size = 18, antiderivative size = 52

$$\int (x + \sqrt{-3 - 4x - x^2})^2 dx = -3x - 2x^2 - 2(2 + x)\sqrt{-3 - 4x - x^2} - \frac{2}{3}(-3 - 4x - x^2)^{3/2} - 2 \arcsin(2 + x)$$

output `-3*x-2*x^2-2*(2+x)*(-x^2-4*x-3)^(1/2)-2/3*(-x^2-4*x-3)^(3/2)-2*arcsin(2+x)`

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.08

$$\int (x + \sqrt{-3 - 4x - x^2})^2 dx = -x(3 + 2x) + \frac{2}{3}\sqrt{-3 - 4x - x^2}(-3 + x + x^2) + 4 \arctan\left(\frac{\sqrt{-3 - 4x - x^2}}{3 + x}\right)$$

input `Integrate[(x + Sqrt[-3 - 4*x - x^2])^2,x]`

output

$$-(x*(3 + 2*x)) + (2*\text{Sqrt}[-3 - 4*x - x^2]*(-3 + x + x^2))/3 + 4*\text{ArcTan}[\text{Sqrt}[-3 - 4*x - x^2]/(3 + x)]$$
Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\sqrt{-x^2 - 4x - 3} + x \right)^2 dx$$

↓ 7293

$$\int \left(2\sqrt{-x^2 - 4x - 3} - 3x - 4x - 3 \right) dx$$

↓ 2009

$$-2 \arcsin(x + 2) - 2x^2 - \frac{2}{3}(-x^2 - 4x - 3)^{3/2} - 2(x + 2)\sqrt{-x^2 - 4x - 3} - 3x$$

input

$$\text{Int}[(x + \text{Sqrt}[-3 - 4*x - x^2])^2, x]$$

output

$$-3*x - 2*x^2 - 2*(2 + x)*\text{Sqrt}[-3 - 4*x - x^2] - (2*(-3 - 4*x - x^2)^{(3/2)})/3 - 2*\text{ArcSin}[2 + x]$$
Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x_Symbol] \text{ :> } \text{Simp}[\text{IntSum}[u, x], x] \text{ /; } \text{SumQ}[u]$$

rule 7293

$$\text{Int}[u_, x_Symbol] \text{ :> } \text{With}[\{v = \text{ExpandIntegrand}[u, x]\}, \text{Int}[v, x] \text{ /; } \text{SumQ}[v]]$$

Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.92

method	result
default	$-2x^2 - 3x - \frac{2(-x^2-4x-3)^{\frac{3}{2}}}{3} + (-2x - 4)\sqrt{-x^2 - 4x - 3} - 2 \arcsin(2 + x)$
trager	$-(2x + 3)x + \left(\frac{2}{3}x^2 + \frac{2}{3}x - 2\right)\sqrt{-x^2 - 4x - 3} + 2 \operatorname{RootOf}(_Z^2 + 1) \ln(\operatorname{RootOf}(_Z^2 + 1) x)$

input `int((x+(-x^2-4*x-3)^(1/2))^2,x,method=_RETURNVERBOSE)`

output `-2*x^2-3*x-2/3*(-x^2-4*x-3)^(3/2)+(-2*x-4)*(-x^2-4*x-3)^(1/2)-2*arcsin(2+x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.12

$$\int \left(x + \sqrt{-3 - 4x - x^2}\right)^2 dx = -2x^2 + \frac{2}{3}(x^2 + x - 3)\sqrt{-x^2 - 4x - 3} - 3x + 2 \arctan\left(\frac{\sqrt{-x^2 - 4x - 3}(x + 2)}{x^2 + 4x + 3}\right)$$

input `integrate((x+(-x^2-4*x-3)^(1/2))^2,x, algorithm="fricas")`

output `-2*x^2 + 2/3*(x^2 + x - 3)*sqrt(-x^2 - 4*x - 3) - 3*x + 2*arctan(sqrt(-x^2 - 4*x - 3)*(x + 2)/(x^2 + 4*x + 3))`

Sympy [A] (verification not implemented)

Time = 0.48 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.75

$$\int \left(x + \sqrt{-3 - 4x - x^2} \right)^2 dx = -2x^2 - 3x + 2\sqrt{-x^2 - 4x - 3} \left(\frac{x^2}{3} + \frac{x}{3} - 1 \right) - 2 \operatorname{asin}(x + 2)$$

input `integrate((x+(-x**2-4*x-3)**(1/2))**2,x)`output `-2*x**2 - 3*x + 2*sqrt(-x**2 - 4*x - 3)*(x**2/3 + x/3 - 1) - 2*asin(x + 2)`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.15

$$\int \left(x + \sqrt{-3 - 4x - x^2} \right)^2 dx = -2x^2 - \frac{2}{3}(-x^2 - 4x - 3)^{\frac{3}{2}} - 2\sqrt{-x^2 - 4x - 3}x - 3x - 4\sqrt{-x^2 - 4x - 3} + 2 \operatorname{arcsin}(-x - 2)$$

input `integrate((x+(-x^2-4*x-3)^(1/2))^2,x, algorithm="maxima")`output `-2*x^2 - 2/3*(-x^2 - 4*x - 3)^(3/2) - 2*sqrt(-x^2 - 4*x - 3)*x - 3*x - 4*sqrt(-x^2 - 4*x - 3) + 2*arcsin(-x - 2)`**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.69

$$\int \left(x + \sqrt{-3 - 4x - x^2} \right)^2 dx = -2x^2 + \frac{2}{3}((x + 1)x - 3)\sqrt{-x^2 - 4x - 3} - 3x - 2 \operatorname{arcsin}(x + 2)$$

input `integrate((x+(-x^2-4*x-3)^(1/2))^2,x, algorithm="giac")`output `-2*x^2 + 2/3*((x + 1)*x - 3)*sqrt(-x^2 - 4*x - 3) - 3*x - 2*arcsin(x + 2)`

Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.06

$$\int \left(x + \sqrt{-3 - 4x - x^2}\right)^2 dx = \frac{\sqrt{-x^2 - 4x - 3}(8x^2 + 8x - 24)}{12} - 3x - 2x^2 + \ln\left(x + 2 - \sqrt{-x^2 - 4x - 3}\right) 2i$$

input `int((x + (- 4*x - x^2 - 3)^(1/2))^2,x)`output `log(x - (- 4*x - x^2 - 3)^(1/2)*1i + 2)*2i - 3*x + ((- 4*x - x^2 - 3)^(1/2))*(8*x + 8*x^2 - 24))/12 - 2*x^2`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.13

$$\int \left(x + \sqrt{-3 - 4x - x^2}\right)^2 dx = -2\operatorname{asin}(x + 2) + \frac{2\sqrt{-x^2 - 4x - 3}x^2}{3} + \frac{2\sqrt{-x^2 - 4x - 3}x}{3} - 2\sqrt{-x^2 - 4x - 3} - 2x^2 - 3x + \frac{2}{3}$$

input `int((x+(-x^2-4*x-3)^(1/2))^2,x)`output `(- 6*asin(x + 2) + 2*sqrt(- x**2 - 4*x - 3)*x**2 + 2*sqrt(- x**2 - 4*x - 3)*x - 6*sqrt(- x**2 - 4*x - 3) - 6*x**2 - 9*x + 2)/3`

3.42 $\int (x + \sqrt{-3 - 4x - x^2}) dx$

Optimal result	343
Mathematica [A] (verified)	343
Rubi [A] (verified)	344
Maple [A] (verified)	344
Fricas [A] (verification not implemented)	345
Sympy [A] (verification not implemented)	345
Maxima [A] (verification not implemented)	346
Giac [A] (verification not implemented)	346
Mupad [B] (verification not implemented)	346
Reduce [B] (verification not implemented)	347

Optimal result

Integrand size = 16, antiderivative size = 37

$$\int (x + \sqrt{-3 - 4x - x^2}) dx = \frac{x^2}{2} + \frac{1}{2}(2+x)\sqrt{-3 - 4x - x^2} + \frac{1}{2} \arcsin(2+x)$$

output

```
1/2*x^2+1/2*(2+x)*(-x^2-4*x-3)^(1/2)+1/2*arcsin(2+x)
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.41

$$\int (x + \sqrt{-3 - 4x - x^2}) dx = \frac{x^2}{2} + \frac{1}{2}(2+x)\sqrt{-3 - 4x - x^2} - \arctan\left(\frac{\sqrt{-3 - 4x - x^2}}{3+x}\right)$$

input

```
Integrate[x + Sqrt[-3 - 4*x - x^2], x]
```

output

```
x^2/2 + ((2 + x)*Sqrt[-3 - 4*x - x^2])/2 - ArcTan[Sqrt[-3 - 4*x - x^2]/(3 + x)]
```


Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.062$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\sqrt{-x^2 - 4x - 3} + x \right) dx$$

↓ 2009

$$\frac{1}{2} \arcsin(x+2) + \frac{x^2}{2} + \frac{1}{2}(x+2)\sqrt{-x^2 - 4x - 3}$$

input `Int[x + Sqrt[-3 - 4*x - x^2], x]`

output `x^2/2 + ((2 + x)*Sqrt[-3 - 4*x - x^2])/2 + ArcSin[2 + x]/2`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.86

method	result
default	$\frac{x^2}{2} - \frac{(-2x-4)\sqrt{-x^2-4x-3}}{4} + \frac{\arcsin(2+x)}{2}$
parts	$\frac{x^2}{2} - \frac{(-2x-4)\sqrt{-x^2-4x-3}}{4} + \frac{\arcsin(2+x)}{2}$
trager	$\frac{x^2}{2} + \left(1 + \frac{x}{2}\right) \sqrt{-x^2 - 4x - 3} + \frac{\text{RootOf}(_Z^2 + 1) \ln\left(-\text{RootOf}(_Z^2 + 1)x + \sqrt{-x^2 - 4x - 3} - 2\text{RootOf}(_Z^2 + 1)\right)}{2}$

input `int(x+(-x^2-4*x-3)^(1/2),x,method=_RETURNVERBOSE)`

output `1/2*x^2-1/4*(-2*x-4)*(-x^2-4*x-3)^(1/2)+1/2*arcsin(2+x)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.41

$$\int \left(x + \sqrt{-3 - 4x - x^2} \right) dx = \frac{1}{2} x^2 + \frac{1}{2} \sqrt{-x^2 - 4x - 3}(x + 2) - \frac{1}{2} \arctan \left(\frac{\sqrt{-x^2 - 4x - 3}(x + 2)}{x^2 + 4x + 3} \right)$$

input `integrate(x+(-x^2-4*x-3)^(1/2),x, algorithm="fricas")`

output `1/2*x^2 + 1/2*sqrt(-x^2 - 4*x - 3)*(x + 2) - 1/2*arctan(sqrt(-x^2 - 4*x - 3)*(x + 2)/(x^2 + 4*x + 3))`

Sympy [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.78

$$\int \left(x + \sqrt{-3 - 4x - x^2} \right) dx = \frac{x^2}{2} + \left(\frac{x}{2} + 1 \right) \sqrt{-x^2 - 4x - 3} + \frac{\operatorname{asin}(x + 2)}{2}$$

input `integrate(x+(-x**2-4*x-3)**(1/2),x)`

output `x**2/2 + (x/2 + 1)*sqrt(-x**2 - 4*x - 3) + asin(x + 2)/2`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.11

$$\int \left(x + \sqrt{-3 - 4x - x^2} \right) dx = \frac{1}{2} x^2 + \frac{1}{2} \sqrt{-x^2 - 4x - 3} x + \sqrt{-x^2 - 4x - 3} - \frac{1}{2} \arcsin(-x - 2)$$

input `integrate(x+(-x^2-4*x-3)^(1/2),x, algorithm="maxima")`output `1/2*x^2 + 1/2*sqrt(-x^2 - 4*x - 3)*x + sqrt(-x^2 - 4*x - 3) - 1/2*arcsin(-x - 2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.78

$$\int \left(x + \sqrt{-3 - 4x - x^2} \right) dx = \frac{1}{2} x^2 + \frac{1}{2} \sqrt{-x^2 - 4x - 3}(x + 2) + \frac{1}{2} \arcsin(x + 2)$$

input `integrate(x+(-x^2-4*x-3)^(1/2),x, algorithm="giac")`output `1/2*x^2 + 1/2*sqrt(-x^2 - 4*x - 3)*(x + 2) + 1/2*arcsin(x + 2)`**Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.81

$$\int \left(x + \sqrt{-3 - 4x - x^2} \right) dx = \frac{\arcsin(x + 2)}{2} + \left(\frac{x}{2} + 1 \right) \sqrt{-x^2 - 4x - 3} + \frac{x^2}{2}$$

input `int(x + (- 4*x - x^2 - 3)^(1/2),x)`output `asin(x + 2)/2 + (x/2 + 1)*(- 4*x - x^2 - 3)^(1/2) + x^2/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00

$$\int \left(x + \sqrt{-3 - 4x - x^2} \right) dx = \frac{\operatorname{asin}(x + 2)}{2} + \frac{\sqrt{-x^2 - 4x - 3} x}{2} + \sqrt{-x^2 - 4x - 3} + \frac{x^2}{2}$$

input `int(x+(-x^2-4*x-3)^(1/2),x)`

output `(asin(x + 2) + sqrt(- x**2 - 4*x - 3)*x + 2*sqrt(- x**2 - 4*x - 3) + x**2)/2`

3.43 $\int \frac{1}{x + \sqrt{-3 - 4x - x^2}} dx$

Optimal result	348
Mathematica [A] (verified)	349
Rubi [A] (verified)	349
Maple [B] (verified)	353
Fricas [B] (verification not implemented)	354
Sympy [F]	355
Maxima [F]	355
Giac [B] (verification not implemented)	356
Mupad [F(-1)]	357
Reduce [B] (verification not implemented)	357

Optimal result

Integrand size = 18, antiderivative size = 96

$$\int \frac{1}{x + \sqrt{-3 - 4x - x^2}} dx = -\arctan\left(\frac{\sqrt{-3 - 4x - x^2}}{3 + x}\right) - \sqrt{2} \arctan\left(\frac{1 - \frac{3\sqrt{-3 - 4x - x^2}}{3 + x}}{\sqrt{2}}\right) + \frac{1}{2} \log(3 + x) + \frac{1}{2} \log\left(\frac{x + \sqrt{-3 - 4x - x^2}}{3 + x}\right)$$

```
output -arctan((-x^2-4*x-3)^(1/2)/(3+x))-2^(1/2)*arctan(1/2*(1-3*(-x^2-4*x-3)^(1/2)/(3+x))*2^(1/2))+1/2*ln(3+x)+1/2*ln((x+(-x^2-4*x-3)^(1/2))/(3+x))
```

Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.84

$$\int \frac{1}{x + \sqrt{-3 - 4x - x^2}} dx = \frac{1}{2} \left(-2 \arctan \left(\frac{\sqrt{-3 - 4x - x^2}}{3 + x} \right) - 2\sqrt{2} \arctan \left(\frac{\sqrt{2}(1 + x)}{1 + x + \sqrt{-3 - 4x - x^2}} \right) + \log \left(x + \sqrt{-3 - 4x - x^2} \right) \right)$$

input

```
Integrate[(x + Sqrt[-3 - 4*x - x^2])^(-1), x]
```

output

```
(-2*ArcTan[Sqrt[-3 - 4*x - x^2]/(3 + x)] - 2*Sqrt[2]*ArcTan[(Sqrt[2]*(1 + x))/(1 + x + Sqrt[-3 - 4*x - x^2])]) + Log[x + Sqrt[-3 - 4*x - x^2]]/2
```

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.29, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {7287, 27, 1356, 27, 452, 216, 240, 1142, 27, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{-x^2 - 4x - 3} + x} dx$$

↓ 7287

$$2 \int \frac{2\sqrt{-x-1}}{\sqrt{x+3} \left(\frac{-x-1}{x+3} + 1 \right) \left(\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1 \right)} d \frac{\sqrt{-x-1}}{\sqrt{x+3}}$$

↓ 27

$$4 \int \frac{\sqrt{-x-1}}{\sqrt{x+3} \left(\frac{-x-1}{x+3} + 1 \right) \left(\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1 \right)} d \frac{\sqrt{-x-1}}{\sqrt{x+3}}$$

$$\begin{aligned}
& \downarrow 1356 \\
4 \left(\frac{1}{8} \int -\frac{2 \left(\frac{\sqrt{-x-1}}{\sqrt{x+3}} + 1 \right)}{\frac{-x-1}{x+3} + 1} d \frac{\sqrt{-x-1}}{\sqrt{x+3}} + \frac{1}{8} \int \frac{2 \left(\frac{3\sqrt{-x-1}}{\sqrt{x+3}} + 1 \right)}{\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1} d \frac{\sqrt{-x-1}}{\sqrt{x+3}} \right) \\
& \downarrow 27 \\
4 \left(\frac{1}{4} \int \frac{\frac{3\sqrt{-x-1}}{\sqrt{x+3}} + 1}{\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1} d \frac{\sqrt{-x-1}}{\sqrt{x+3}} - \frac{1}{4} \int \frac{\frac{\sqrt{-x-1}}{\sqrt{x+3}} + 1}{\frac{-x-1}{x+3} + 1} d \frac{\sqrt{-x-1}}{\sqrt{x+3}} \right) \\
& \downarrow 452 \\
4 \left(\frac{1}{4} \left(- \int \frac{1}{\frac{-x-1}{x+3} + 1} d \frac{\sqrt{-x-1}}{\sqrt{x+3}} - \int \frac{\sqrt{-x-1}}{\sqrt{x+3} \left(\frac{-x-1}{x+3} + 1 \right)} d \frac{\sqrt{-x-1}}{\sqrt{x+3}} \right) + \frac{1}{4} \int \frac{\frac{3\sqrt{-x-1}}{\sqrt{x+3}} + 1}{\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1} d \frac{\sqrt{-x-1}}{\sqrt{x+3}} \right) \\
& \downarrow 216 \\
4 \left(\frac{1}{4} \left(- \int \frac{\sqrt{-x-1}}{\sqrt{x+3} \left(\frac{-x-1}{x+3} + 1 \right)} d \frac{\sqrt{-x-1}}{\sqrt{x+3}} - \arctan \left(\frac{\sqrt{-x-1}}{\sqrt{x+3}} \right) \right) + \frac{1}{4} \int \frac{\frac{3\sqrt{-x-1}}{\sqrt{x+3}} + 1}{\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1} d \frac{\sqrt{-x-1}}{\sqrt{x+3}} \right) \\
& \downarrow 240 \\
4 \left(\frac{1}{4} \int \frac{\frac{3\sqrt{-x-1}}{\sqrt{x+3}} + 1}{\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1} d \frac{\sqrt{-x-1}}{\sqrt{x+3}} + \frac{1}{4} \left(- \arctan \left(\frac{\sqrt{-x-1}}{\sqrt{x+3}} \right) - \frac{1}{2} \log \left(\frac{-x-1}{x+3} + 1 \right) \right) \right) \\
& \downarrow 1142 \\
4 \left(\frac{1}{4} \left(2 \int \frac{1}{\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1} d \frac{\sqrt{-x-1}}{\sqrt{x+3}} + \frac{1}{2} \int -\frac{2 \left(1 - \frac{3\sqrt{-x-1}}{\sqrt{x+3}} \right)}{\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1} d \frac{\sqrt{-x-1}}{\sqrt{x+3}} \right) + \frac{1}{4} \left(- \arctan \left(\frac{\sqrt{-x-1}}{\sqrt{x+3}} \right) \right) \right) \\
& \downarrow 27 \\
4 \left(\frac{1}{4} \left(2 \int \frac{1}{\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1} d \frac{\sqrt{-x-1}}{\sqrt{x+3}} - \int \frac{1 - \frac{3\sqrt{-x-1}}{\sqrt{x+3}}}{\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1} d \frac{\sqrt{-x-1}}{\sqrt{x+3}} \right) + \frac{1}{4} \left(- \arctan \left(\frac{\sqrt{-x-1}}{\sqrt{x+3}} \right) \right) \right) \\
& \downarrow 1083
\end{aligned}$$

$$\begin{aligned}
& 4 \left(\frac{1}{4} \left(-4 \int \frac{1}{-\frac{x-1}{x+3} - 8} d \left(\frac{6\sqrt{-x-1}}{\sqrt{x+3}} - 2 \right) - \int \frac{1 - \frac{3\sqrt{-x-1}}{\sqrt{x+3}}}{\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1} d \frac{\sqrt{-x-1}}{\sqrt{x+3}} \right) + \frac{1}{4} \left(-\arctan \left(\frac{\sqrt{-x-1}}{\sqrt{x+3}} \right) \right. \right. \\
& \quad \downarrow \text{217} \\
& 4 \left(\frac{1}{4} \left(\sqrt{2} \arctan \left(\frac{6\sqrt{-x-1}}{2\sqrt{2}} - 2 \right) - \int \frac{1 - \frac{3\sqrt{-x-1}}{\sqrt{x+3}}}{\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1} d \frac{\sqrt{-x-1}}{\sqrt{x+3}} \right) + \frac{1}{4} \left(-\arctan \left(\frac{\sqrt{-x-1}}{\sqrt{x+3}} \right) - \frac{1}{2} \log \right. \right. \\
& \quad \downarrow \text{1103} \\
& 4 \left(\frac{1}{4} \left(-\arctan \left(\frac{\sqrt{-x-1}}{\sqrt{x+3}} \right) - \frac{1}{2} \log \left(\frac{-x-1}{x+3} + 1 \right) \right) + \frac{1}{4} \left(\sqrt{2} \arctan \left(\frac{6\sqrt{-x-1}}{2\sqrt{2}} - 2 \right) + \frac{1}{2} \log \left(\frac{3(-x-1)}{x+3} - \dots \right) \right. \right.
\end{aligned}$$

input `Int[(x + Sqrt[-3 - 4*x - x^2])^(-1), x]`

output `4*((-ArcTan[Sqrt[-1 - x]/Sqrt[3 + x]] - Log[1 + (-1 - x)/(3 + x)]/2)/4 + (Sqrt[2]*ArcTan[(-2 + (6*Sqrt[-1 - x])/Sqrt[3 + x])/(2*Sqrt[2])]) + Log[1 + (3*(-1 - x))/(3 + x) - (2*Sqrt[-1 - x])/Sqrt[3 + x]]/2)/4`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 240 $\text{Int}[(x_)/((a_) + (b_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x^2, x]]/(2*b), x] /; \text{FreeQ}[\{a, b\}, x]$

rule 452 $\text{Int}[((c_) + (d_)*(x_))/((a_) + (b_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[c \text{ Int}[1/(a + b*x^2), x], x] + \text{Simp}[d \text{ Int}[x/(a + b*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[b*c^2 + a*d^2, 0]$

rule 1083 $\text{Int}(((a_) + (b_)*(x_) + (c_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[-2 \text{ Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1103 $\text{Int}(((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$

rule 1142 $\text{Int}(((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \text{ Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \text{ Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

rule 1356 $\text{Int}(((g_) + (h_)*(x_))/(((a_) + (b_)*(x_) + (c_)*(x_)^2)*((d_) + (f_)*(x_)^2)), x_Symbol] \rightarrow \text{With}[\{q = \text{Simplify}[c^2*d^2 + b^2*d*f - 2*a*c*d*f + a^2*f^2]\}, \text{Simp}[1/q \text{ Int}[\text{Simp}[g*c^2*d + g*b^2*f - a*b*h*f - a*g*c*f + c*(h*c*d + g*b*f - a*h*f)*x, x]/(a + b*x + c*x^2), x], x] + \text{Simp}[1/q \text{ Int}[\text{Simp}[b*h*d*f - g*c*d*f + a*g*f^2 - f*(h*c*d + g*b*f - a*h*f)*x, x]/(d + f*x^2), x], x] /; \text{NeQ}[q, 0] /; \text{FreeQ}[\{a, b, c, d, f, g, h\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0]$

rule 7287 $\text{Int}[u_, x_Symbol] \rightarrow \text{With}[\{lst = \text{FunctionOfSquareRootOfQuadratic}[u, x]\}, \text{Simp}[2 \text{ Subst}[\text{Int}[lst[[1]], x], x, lst[[2]]], x] /; \text{!FalseQ}[lst] \ \&\& \ \text{EqQ}[lst[[3]], 3] /; \text{EulerIntegrandQ}[u, x]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 369 vs. 2(83) = 166.

Time = 0.52 (sec) , antiderivative size = 370, normalized size of antiderivative = 3.85

method	result
default	$\frac{\arcsin(2+x)}{2} - \frac{\sqrt{3}\sqrt{4}\sqrt{\frac{3x^2}{(-\frac{3}{2}-x)^2-12}}\left(\sqrt{2}\arctan\left(\frac{\sqrt{\frac{3x^2}{(-\frac{3}{2}-x)^2-12}}\sqrt{2}}{6}\right) - \operatorname{arctanh}\left(\frac{3x}{(-\frac{3}{2}-x)\sqrt{\frac{3x^2}{(-\frac{3}{2}-x)^2-12}}}\right)\right)}{12\sqrt{\frac{-x^2}{(-\frac{3}{2}-x)^2-4}}\left(\frac{-x}{-\frac{3}{2}-x}+1\right)} + \frac{\sqrt{3}\sqrt{4}}{2}$
trager	Expression too large to display

input `int(1/(x+(-x^2-4*x-3)^(1/2)),x,method=_RETURNVERBOSE)`

output

```

1/2*arcsin(2+x)-1/12*3^(1/2)*4^(1/2)*(3*x^2/(-3/2-x)^2-12)^(1/2)*(2^(1/2)*
arctan(1/6*(3*x^2/(-3/2-x)^2-12)^(1/2)*2^(1/2))-arctanh(3*x/(-3/2-x)/(3*x^
2/(-3/2-x)^2-12)^(1/2)))/((x^2/(-3/2-x)^2-4)/(x/(-3/2-x)+1)^2)^(1/2)/(x/(-
3/2-x)+1)+1/3*3^(1/2)*4^(1/2)/((x^2/(-3/2-x)^2-4)/(x/(-3/2-x)+1)^2)^(1/2)/
(x/(-3/2-x)+1)*(3*x^2/(-3/2-x)^2-12)^(1/2)*2^(1/2)*arctan(1/6*(3*x^2/(-3/2
-x)^2-12)^(1/2)*2^(1/2))-1/6*3^(1/2)*4^(1/2)*(3*x^2/(-3/2-x)^2-12)^(1/2)*(
2^(1/2)*arctan(1/6*(3*x^2/(-3/2-x)^2-12)^(1/2)*2^(1/2))+arctanh(3*x/(-3/2-
x)/(3*x^2/(-3/2-x)^2-12)^(1/2)))/((x^2/(-3/2-x)^2-4)/(x/(-3/2-x)+1)^2)^(1/
2)/(x/(-3/2-x)+1)+1/4*ln(2*x^2+4*x+3)-1/2*2^(1/2)*arctan(1/4*(4*x+4)*2^(1/
2))
    
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 187 vs. $2(83) = 166$.

Time = 0.09 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.95

$$\int \frac{1}{x + \sqrt{-3 - 4x - x^2}} dx = -\frac{1}{2} \sqrt{2} \arctan \left(\sqrt{2}(x + 1) \right) + \frac{1}{4} \sqrt{2} \arctan \left(\frac{\sqrt{2}x + 3 \sqrt{2} \sqrt{-x^2 - 4x - 3}}{2(2x + 3)} \right) + \frac{1}{4} \sqrt{2} \arctan \left(-\frac{\sqrt{2}x - 3 \sqrt{2} \sqrt{-x^2 - 4x - 3}}{2(2x + 3)} \right) - \frac{1}{2} \arctan \left(\frac{\sqrt{-x^2 - 4x - 3}(x + 2)}{x^2 + 4x + 3} \right) + \frac{1}{4} \log(2x^2 + 4x + 3) - \frac{1}{8} \log \left(-\frac{2\sqrt{-x^2 - 4x - 3}x + 4x + 3}{x^2} \right) + \frac{1}{8} \log \left(\frac{2\sqrt{-x^2 - 4x - 3}x - 4x - 3}{x^2} \right)$$

input `integrate(1/(x+(-x^2-4*x-3)^(1/2)),x, algorithm="fricas")`

output `-1/2*sqrt(2)*arctan(sqrt(2)*(x + 1)) + 1/4*sqrt(2)*arctan(1/2*(sqrt(2)*x + 3*sqrt(2)*sqrt(-x^2 - 4*x - 3))/(2*x + 3)) + 1/4*sqrt(2)*arctan(-1/2*(sqrt(2)*x - 3*sqrt(2)*sqrt(-x^2 - 4*x - 3))/(2*x + 3)) - 1/2*arctan(sqrt(-x^2 - 4*x - 3)*(x + 2)/(x^2 + 4*x + 3)) + 1/4*log(2*x^2 + 4*x + 3) - 1/8*log(-(2*sqrt(-x^2 - 4*x - 3)*x + 4*x + 3)/x^2) + 1/8*log((2*sqrt(-x^2 - 4*x - 3)*x - 4*x - 3)/x^2)`

Sympy [F]

$$\int \frac{1}{x + \sqrt{-3 - 4x - x^2}} dx = \int \frac{1}{x + \sqrt{-x^2 - 4x - 3}} dx$$

input `integrate(1/(x+(-x**2-4*x-3)**(1/2)),x)`

output `Integral(1/(x + sqrt(-x**2 - 4*x - 3)), x)`

Maxima [F]

$$\int \frac{1}{x + \sqrt{-3 - 4x - x^2}} dx = \int \frac{1}{x + \sqrt{-x^2 - 4x - 3}} dx$$

input `integrate(1/(x+(-x^2-4*x-3)^(1/2)),x, algorithm="maxima")`

output `integrate(1/(x + sqrt(-x^2 - 4*x - 3)), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 197 vs. $2(83) = 166$.

Time = 0.12 (sec) , antiderivative size = 197, normalized size of antiderivative = 2.05

$$\int \frac{1}{x + \sqrt{-3 - 4x - x^2}} dx = -\frac{1}{2} \sqrt{2} \arctan(\sqrt{2}(x+1))$$

$$+ \frac{1}{2} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2} \left(\frac{3(\sqrt{-x^2 - 4x - 3} - 1)}{x+2} + 1\right)\right)$$

$$+ \frac{1}{2} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2} \left(\frac{\sqrt{-x^2 - 4x - 3} - 1}{x+2} + 1\right)\right)$$

$$+ \frac{1}{2} \arcsin(x+2) + \frac{1}{4} \log(2x^2 + 4x + 3)$$

$$+ \frac{1}{4} \log\left(\frac{2(\sqrt{-x^2 - 4x - 3} - 1)}{x+2} + \frac{3(\sqrt{-x^2 - 4x - 3} - 1)^2}{(x+2)^2} + 1\right)$$

$$- \frac{1}{4} \log\left(\frac{2(\sqrt{-x^2 - 4x - 3} - 1)}{x+2} + \frac{(\sqrt{-x^2 - 4x - 3} - 1)^2}{(x+2)^2} + 3\right)$$

input `integrate(1/(x+(-x^2-4*x-3)^(1/2)),x, algorithm="giac")`

output `-1/2*sqrt(2)*arctan(sqrt(2)*(x + 1)) + 1/2*sqrt(2)*arctan(1/2*sqrt(2)*(3*(sqrt(-x^2 - 4*x - 3) - 1)/(x + 2) + 1)) + 1/2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(-x^2 - 4*x - 3) - 1)/(x + 2) + 1)) + 1/2*arcsin(x + 2) + 1/4*log(2*x^2 + 4*x + 3) + 1/4*log(2*(sqrt(-x^2 - 4*x - 3) - 1)/(x + 2) + 3*(sqrt(-x^2 - 4*x - 3) - 1)^2/(x + 2)^2 + 1) - 1/4*log(2*(sqrt(-x^2 - 4*x - 3) - 1)/(x + 2) + (sqrt(-x^2 - 4*x - 3) - 1)^2/(x + 2)^2 + 3)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{x + \sqrt{-3 - 4x - x^2}} dx = \int \frac{1}{x + \sqrt{-x^2 - 4x - 3}} dx$$

input `int(1/(x + (- 4*x - x^2 - 3)^(1/2)), x)`output `int(1/(x + (- 4*x - x^2 - 3)^(1/2)), x)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.46

$$\int \frac{1}{x + \sqrt{-3 - 4x - x^2}} dx = \frac{\operatorname{asin}(x + 2)}{2} - \sqrt{2} \operatorname{atan} \left(\frac{3 \tan \left(\frac{\operatorname{asin}(x+2)}{2} \right) - 1}{\sqrt{2}} \right) + \frac{\log(\sqrt{-x^2 - 4x - 3} + x)}{2}$$

input `int(1/(x+(-x^2-4*x-3)^(1/2)),x)`output `(asin(x + 2) - 2*sqrt(2)*atan((3*tan(asin(x + 2)/2) - 1)/sqrt(2)) + log(sqrt(- x**2 - 4*x - 3) + x))/2`

3.44 $\int \frac{1}{(x + \sqrt{-3 - 4x - x^2})^2} dx$

Optimal result	358
Mathematica [A] (verified)	358
Rubi [A] (verified)	359
Maple [C] (verified)	361
Fricas [A] (verification not implemented)	361
Sympy [F]	362
Maxima [F]	362
Giac [B] (verification not implemented)	362
Mupad [F(-1)]	363
Reduce [B] (verification not implemented)	363

Optimal result

Integrand size = 18, antiderivative size = 101

$$\int \frac{1}{(x + \sqrt{-3 - 4x - x^2})^2} dx = \frac{1 - \frac{\sqrt{-3 - 4x - x^2}}{3+x}}{1 - \frac{2\sqrt{-3 - 4x - x^2}}{3+x} - \frac{3(3+4x+x^2)}{(3+x)^2}} + \frac{\arctan\left(\frac{1 - \frac{3\sqrt{-3 - 4x - x^2}}{3+x}}{\sqrt{2}}\right)}{\sqrt{2}}$$

output

```
(1-(-x^2-4*x-3)^(1/2)/(3+x))/(1-2*(-x^2-4*x-3)^(1/2)/(3+x)-3*(x^2+4*x+3)/(3+x)^2)+1/2*2^(1/2)*arctan(1/2*(1-3*(-x^2-4*x-3)^(1/2)/(3+x))*2^(1/2))
```

Mathematica [A] (verified)

Time = 0.40 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.83

$$\int \frac{1}{(x + \sqrt{-3 - 4x - x^2})^2} dx = \frac{3 + x + (3 + 2x)\sqrt{-3 - 4x - x^2} + \sqrt{2}(3 + 4x + 2x^2) \arctan\left(\frac{\sqrt{2}(1+x)}{1+x+\sqrt{-3-4x-x^2}}\right)}{2(3 + 4x + 2x^2)}$$

input

```
Integrate[(x + Sqrt[-3 - 4*x - x^2])^(-2), x]
```

output

$$(3 + x + (3 + 2*x)*\text{Sqrt}[-3 - 4*x - x^2] + \text{Sqrt}[2]*(3 + 4*x + 2*x^2)*\text{ArcTan}[(\text{Sqrt}[2]*(1 + x))/((1 + x + \text{Sqrt}[-3 - 4*x - x^2]))])/(2*(3 + 4*x + 2*x^2))$$
Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.99, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {7287, 27, 1159, 1083, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(\sqrt{-x^2 - 4x - 3} + x)^2} dx$$

$$\downarrow 7287$$

$$2 \int -\frac{2\sqrt{-x-1}}{\sqrt{x+3} \left(\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1 \right)^2} d\frac{\sqrt{-x-1}}{\sqrt{x+3}}$$

$$\downarrow 27$$

$$-4 \int \frac{\sqrt{-x-1}}{\sqrt{x+3} \left(\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1 \right)^2} d\frac{\sqrt{-x-1}}{\sqrt{x+3}}$$

$$\downarrow 1159$$

$$-4 \left(\frac{1}{4} \int \frac{1}{\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1} d\frac{\sqrt{-x-1}}{\sqrt{x+3}} - \frac{1 - \frac{\sqrt{-x-1}}{\sqrt{x+3}}}{4 \left(\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1 \right)} \right)$$

$$\downarrow 1083$$

$$-4 \left(-\frac{1}{2} \int \frac{1}{-\frac{x-1}{x+3} - 8} d\left(\frac{6\sqrt{-x-1}}{\sqrt{x+3}} - 2 \right) - \frac{1 - \frac{\sqrt{-x-1}}{\sqrt{x+3}}}{4 \left(\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1 \right)} \right)$$

$$\downarrow 217$$

$$-4 \left(\frac{\arctan\left(\frac{6\sqrt{-x-1}-2}{2\sqrt{2}}\right)}{4\sqrt{2}} - \frac{1 - \frac{\sqrt{-x-1}}{\sqrt{x+3}}}{4 \left(\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1 \right)} \right)$$

input `Int[(x + Sqrt[-3 - 4*x - x^2])^(-2), x]`

output `-4*(-1/4*(1 - Sqrt[-1 - x]/Sqrt[3 + x])/(1 + (3*(-1 - x))/(3 + x) - (2*Sqrt[-1 - x])/Sqrt[3 + x]) + ArcTan[(-2 + (6*Sqrt[-1 - x])/Sqrt[3 + x])/(2*Sqrt[2])]/(4*Sqrt[2]))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1083 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`

rule 1159 `Int[((d_.) + (e_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[((b*d - 2*a*e + (2*c*d - b*e)*x)/((p + 1)*(b^2 - 4*a*c)))*(a + b*x + c*x^2)^(p + 1), x] - Simp[(2*p + 3)*((2*c*d - b*e)/((p + 1)*(b^2 - 4*a*c)))] Int[(a + b*x + c*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 7287 `Int[u_, x_Symbol] := With[{lst = FunctionOfSquareRootOfQuadratic[u, x]}, Simp[2 Subst[Int[lst[[1]], x], x, lst[[2]]], x] /; !FalseQ[lst] && EqQ[lst[[3]], 3] /; EulerIntegrandQ[u, x]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.18 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.09

method	result
trager	$-\frac{(2x+3)x}{2(2x^2+4x+3)} + \frac{(2x+3)\sqrt{-x^2-4x-3}}{4x^2+8x+6} - \frac{\text{RootOf}(_Z^2+2) \ln\left(\frac{2\text{RootOf}(_Z^2+2)x+3\text{RootOf}(_Z^2+2)+2\sqrt{-x^2-4x-3}}{\text{RootOf}(_Z^2+2)x-2x-3}\right)}{4}$
default	Expression too large to display

input `int(1/(x+(-x^2-4*x-3)^(1/2))^2,x,method=_RETURNVERBOSE)`

output `-1/2*(2*x+3)*x/(2*x^2+4*x+3)+1/2*(2*x+3)/(2*x^2+4*x+3)*(-x^2-4*x-3)^(1/2)-1/4*RootOf(_Z^2+2)*ln((2*RootOf(_Z^2+2)*x+3*RootOf(_Z^2+2)+2*(-x^2-4*x-3)^(1/2))/(RootOf(_Z^2+2)*x-2*x-3))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.20

$$\int \frac{1}{(x + \sqrt{-3 - 4x - x^2})^2} dx$$

$$= \frac{2\sqrt{2}(2x^2 + 4x + 3) \arctan(\sqrt{2}(x + 1)) - \sqrt{2}(2x^2 + 4x + 3) \arctan\left(\frac{\sqrt{2}(6x^2 + 20x + 15)\sqrt{-x^2 - 4x - 3}}{4(2x^3 + 11x^2 + 18x + 9)}\right) + 4\sqrt{2}}{8(2x^2 + 4x + 3)}$$

input `integrate(1/(x+(-x^2-4*x-3)^(1/2))^2,x, algorithm="fricas")`

output `1/8*(2*sqrt(2)*(2*x^2 + 4*x + 3)*arctan(sqrt(2)*(x + 1)) - sqrt(2)*(2*x^2 + 4*x + 3)*arctan(1/4*sqrt(2)*(6*x^2 + 20*x + 15)*sqrt(-x^2 - 4*x - 3)/(2*x^3 + 11*x^2 + 18*x + 9)) + 4*sqrt(-x^2 - 4*x - 3)*(2*x + 3) + 4*x + 12)/(8*(2*x^2 + 4*x + 3))`

Sympy [F]

$$\int \frac{1}{(x + \sqrt{-3 - 4x - x^2})^2} dx = \int \frac{1}{(x + \sqrt{-x^2 - 4x - 3})^2} dx$$

input `integrate(1/(x+(-x**2-4*x-3)**(1/2))**2,x)`

output `Integral((x + sqrt(-x**2 - 4*x - 3))**(-2), x)`

Maxima [F]

$$\int \frac{1}{(x + \sqrt{-3 - 4x - x^2})^2} dx = \int \frac{1}{(x + \sqrt{-x^2 - 4x - 3})^2} dx$$

input `integrate(1/(x+(-x^2-4*x-3)^(1/2))^2,x, algorithm="maxima")`

output `integrate((x + sqrt(-x^2 - 4*x - 3))^-2, x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 263 vs. $2(92) = 184$.

Time = 0.12 (sec) , antiderivative size = 263, normalized size of antiderivative = 2.60

$$\begin{aligned} & \int \frac{1}{(x + \sqrt{-3 - 4x - x^2})^2} dx \\ &= \frac{1}{4} \sqrt{2} \arctan(\sqrt{2}(x+1)) - \frac{1}{4} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2} \left(\frac{3(\sqrt{-x^2 - 4x - 3} - 1)}{x+2} + 1\right)\right) \\ & \quad - \frac{1}{4} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2} \left(\frac{\sqrt{-x^2 - 4x - 3} - 1}{x+2} + 1\right)\right) + \frac{x+3}{2(2x^2 + 4x + 3)} \\ & \quad - \frac{10(\sqrt{-x^2 - 4x - 3} - 1)}{x+2} + \frac{7(\sqrt{-x^2 - 4x - 3} - 1)^2}{(x+2)^2} - \frac{2(\sqrt{-x^2 - 4x - 3} - 1)^3}{(x+2)^3} + 3 \\ & \quad - \frac{3\left(\frac{8(\sqrt{-x^2 - 4x - 3} - 1)}{x+2} + \frac{14(\sqrt{-x^2 - 4x - 3} - 1)^2}{(x+2)^2} + \frac{8(\sqrt{-x^2 - 4x - 3} - 1)^3}{(x+2)^3} + \frac{3(\sqrt{-x^2 - 4x - 3} - 1)^4}{(x+2)^4} + 3\right)}{3} \end{aligned}$$

input `integrate(1/(x+(-x^2-4*x-3)^(1/2))^2,x, algorithm="giac")`

output `1/4*sqrt(2)*arctan(sqrt(2)*(x + 1)) - 1/4*sqrt(2)*arctan(1/2*sqrt(2)*(3*(sqrt(-x^2 - 4*x - 3) - 1)/(x + 2) + 1)) - 1/4*sqrt(2)*arctan(1/2*sqrt(2)*((sqrt(-x^2 - 4*x - 3) - 1)/(x + 2) + 1)) + 1/2*(x + 3)/(2*x^2 + 4*x + 3) - 1/3*(10*(sqrt(-x^2 - 4*x - 3) - 1)/(x + 2) + 7*(sqrt(-x^2 - 4*x - 3) - 1)^2/(x + 2)^2 - 2*(sqrt(-x^2 - 4*x - 3) - 1)^3/(x + 2)^3 + 3)/(8*(sqrt(-x^2 - 4*x - 3) - 1)/(x + 2) + 14*(sqrt(-x^2 - 4*x - 3) - 1)^2/(x + 2)^2 + 8*(sqrt(-x^2 - 4*x - 3) - 1)^3/(x + 2)^3 + 3*(sqrt(-x^2 - 4*x - 3) - 1)^4/(x + 2)^4 + 3)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(x + \sqrt{-3 - 4x - x^2})^2} dx = \int \frac{1}{(x + \sqrt{-x^2 - 4x - 3})^2} dx$$

input `int(1/(x + (- 4*x - x^2 - 3)^(1/2))^2,x)`

output `int(1/(x + (- 4*x - x^2 - 3)^(1/2))^2, x)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.91

$$\int \frac{1}{(x + \sqrt{-3 - 4x - x^2})^2} dx$$

$$= \frac{4\sqrt{-x^2 - 4x - 3} \sqrt{2} \operatorname{atan}\left(\frac{3 \tan\left(\frac{\operatorname{asin}(x+2)}{2}\right) - 1}{\sqrt{2}}\right) + 4\sqrt{2} \operatorname{atan}\left(\frac{3 \tan\left(\frac{\operatorname{asin}(x+2)}{2}\right) - 1}{\sqrt{2}}\right) x + 7\sqrt{-x^2 - 4x - 3} - x}{8\sqrt{-x^2 - 4x - 3} + 8x}$$

input `int(1/(x+(-x^2-4*x-3)^(1/2))^2,x)`

output

```
(4*sqrt(-x**2 - 4*x - 3)*sqrt(2)*atan((3*tan(asin(x + 2)/2) - 1)/sqrt(2)
) + 4*sqrt(2)*atan((3*tan(asin(x + 2)/2) - 1)/sqrt(2))*x + 7*sqrt(-x**2
- 4*x - 3) - x - 12)/(8*(sqrt(-x**2 - 4*x - 3) + x))
```

3.45
$$\int \frac{1}{\left(x + \sqrt{-3 - 4x - x^2}\right)^3} dx$$

Optimal result	365
Mathematica [A] (verified)	366
Rubi [A] (verified)	366
Maple [C] (verified)	369
Fricas [A] (verification not implemented)	369
Sympy [F]	370
Maxima [F]	370
Giac [B] (verification not implemented)	371
Mupad [F(-1)]	372
Reduce [B] (verification not implemented)	372

Optimal result

Integrand size = 18, antiderivative size = 174

$$\int \frac{1}{\left(x + \sqrt{-3 - 4x - x^2}\right)^3} dx = -\frac{2\left(2 - \frac{\sqrt{-3 - 4x - x^2}}{3+x}\right)}{9\left(1 - \frac{2\sqrt{-3 - 4x - x^2}}{3+x} - \frac{3(3+4x+x^2)}{(3+x)^2}\right)^2} - \frac{13 - \frac{27\sqrt{-3 - 4x - x^2}}{3+x}}{18\left(1 - \frac{2\sqrt{-3 - 4x - x^2}}{3+x} - \frac{3(3+4x+x^2)}{(3+x)^2}\right)} - \frac{3 \arctan\left(\frac{1 - \frac{3\sqrt{-3 - 4x - x^2}}{3+x}}{\sqrt{2}}\right)}{2\sqrt{2}}$$

output

```
1/9*(-4+2*(-x^2-4*x-3)^(1/2)/(3+x))/(1-2*(-x^2-4*x-3)^(1/2)/(3+x)-3*(x^2+4*x+3)/(3+x)^2)-
(13-27*(-x^2-4*x-3)^(1/2)/(3+x))/(18-36*(-x^2-4*x-3)^(1/2)/(3+x)-54*(x^2+4*x+3)/(3+x)^2)-
3/4*2^(1/2)*arctan(1/2*(1-3*(-x^2-4*x-3)^(1/2)/(3+x))*2^(1/2))
```

Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.63

$$\int \frac{1}{(x + \sqrt{-3 - 4x - x^2})^3} dx = \frac{9 + 15x + 16x^2 + 6x^3 + \sqrt{-3 - 4x - x^2}(15 + 26x + 22x^2 + 8x^3) + 3\sqrt{2}(3 + 4x + 2x^2)^2 \arctan\left(\frac{\sqrt{2}(1+x)}{1+x}\right)}{4(3 + 4x + 2x^2)^2}$$

input `Integrate[(x + Sqrt[-3 - 4*x - x^2])^(-3), x]`

output `-1/4*(9 + 15*x + 16*x^2 + 6*x^3 + Sqrt[-3 - 4*x - x^2]*(15 + 26*x + 22*x^2 + 8*x^3) + 3*Sqrt[2]*(3 + 4*x + 2*x^2)^2*ArcTan[(Sqrt[2]*(1 + x))/(1 + x + Sqrt[-3 - 4*x - x^2]])]/(3 + 4*x + 2*x^2)^2`

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.94, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {7287, 27, 2191, 27, 1159, 1083, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(\sqrt{-x^2 - 4x - 3} + x)^3} dx \\ & \quad \downarrow \text{7287} \\ & 2 \int \frac{2\sqrt{-x-1}\left(\frac{-x-1}{x+3} + 1\right)}{\sqrt{x+3}\left(\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1\right)^3} d\frac{\sqrt{-x-1}}{\sqrt{x+3}} \\ & \quad \downarrow \text{27} \\ & 4 \int \frac{\sqrt{-x-1}\left(\frac{-x-1}{x+3} + 1\right)}{\sqrt{x+3}\left(\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1\right)^3} d\frac{\sqrt{-x-1}}{\sqrt{x+3}} \end{aligned}$$

$$\begin{aligned}
& \downarrow 2191 \\
& 4 \left(\frac{1}{16} \int \frac{8 \left(\frac{6\sqrt{-x-1}}{\sqrt{x+3}} + 7 \right)}{9 \left(\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1 \right)^2} d \frac{\sqrt{-x-1}}{\sqrt{x+3}} - \frac{2 - \frac{\sqrt{-x-1}}{\sqrt{x+3}}}{18 \left(\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1 \right)^2} \right) \\
& \downarrow 27 \\
& 4 \left(\frac{1}{18} \int \frac{\frac{6\sqrt{-x-1}}{\sqrt{x+3}} + 7}{\left(\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1 \right)^2} d \frac{\sqrt{-x-1}}{\sqrt{x+3}} - \frac{2 - \frac{\sqrt{-x-1}}{\sqrt{x+3}}}{18 \left(\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1 \right)^2} \right) \\
& \downarrow 1159 \\
& 4 \left(\frac{1}{18} \left(\frac{27}{4} \int \frac{1}{\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1} d \frac{\sqrt{-x-1}}{\sqrt{x+3}} - \frac{13 - \frac{27\sqrt{-x-1}}{\sqrt{x+3}}}{4 \left(\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1 \right)} \right) - \frac{2 - \frac{\sqrt{-x-1}}{\sqrt{x+3}}}{18 \left(\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1 \right)^2} \right) \\
& \downarrow 1083 \\
& 4 \left(\frac{1}{18} \left(-\frac{27}{2} \int \frac{1}{-\frac{-x-1}{x+3} - 8} d \left(\frac{6\sqrt{-x-1}}{\sqrt{x+3}} - 2 \right) - \frac{13 - \frac{27\sqrt{-x-1}}{\sqrt{x+3}}}{4 \left(\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1 \right)} \right) - \frac{2 - \frac{\sqrt{-x-1}}{\sqrt{x+3}}}{18 \left(\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1 \right)^2} \right) \\
& \downarrow 217 \\
& 4 \left(\frac{1}{18} \left(\frac{27 \arctan \left(\frac{\frac{6\sqrt{-x-1}}{\sqrt{x+3}} - 2}{2\sqrt{2}} \right)}{4\sqrt{2}} - \frac{13 - \frac{27\sqrt{-x-1}}{\sqrt{x+3}}}{4 \left(\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1 \right)} \right) - \frac{2 - \frac{\sqrt{-x-1}}{\sqrt{x+3}}}{18 \left(\frac{3(-x-1)}{x+3} - \frac{2\sqrt{-x-1}}{\sqrt{x+3}} + 1 \right)^2} \right)
\end{aligned}$$

input

```
Int[(x + Sqrt[-3 - 4*x - x^2])^(-3), x]
```

output

```
4*(-1/18*(2 - Sqrt[-1 - x]/Sqrt[3 + x])/(1 + (3*(-1 - x))/(3 + x) - (2*Sqrt[-1 - x])/Sqrt[3 + x])^2 + (-1/4*(13 - (27*Sqrt[-1 - x])/Sqrt[3 + x])/(1 + (3*(-1 - x))/(3 + x) - (2*Sqrt[-1 - x])/Sqrt[3 + x]) + (27*ArcTan[(-2 + (6*Sqrt[-1 - x])/Sqrt[3 + x])/(2*Sqrt[2])])/(4*Sqrt[2]))/18)
```


Definitions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 217 $\text{Int}[((a_) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 1083 $\text{Int}[((a_) + (b_*)(x_) + (c_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[-2 \text{ Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1159 $\text{Int}[((d_) + (e_*)(x_))*((a_) + (b_*)(x_) + (c_*)(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[((b*d - 2*a*e + (2*c*d - b*e)*x)/((p + 1)*(b^2 - 4*a*c)))*(a + b*x + c*x^2)^{(p + 1)}, x] - \text{Simp}[(2*p + 3)*((2*c*d - b*e)/((p + 1)*(b^2 - 4*a*c)) \ \text{Int}[(a + b*x + c*x^2)^{(p + 1)}, x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \& \ \& \ \text{LtQ}[p, -1] \ \&\& \ \text{NeQ}[p, -3/2]$
- rule 2191 $\text{Int}[(Pq_)*((a_) + (b_*)(x_) + (c_*)(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{With}[\{Q = \text{PolynomialQuotient}[Pq, a + b*x + c*x^2, x], f = \text{Coeff}[\text{PolynomialRemainder}[Pq, a + b*x + c*x^2, x], x, 0], g = \text{Coeff}[\text{PolynomialRemainder}[Pq, a + b*x + c*x^2, x], x, 1]\}, \text{Simp}[(b*f - 2*a*g + (2*c*f - b*g)*x)*((a + b*x + c*x^2)^{(p + 1})/((p + 1)*(b^2 - 4*a*c))), x] + \text{Simp}[1/((p + 1)*(b^2 - 4*a*c)) \ \text{Int}[(a + b*x + c*x^2)^{(p + 1)}*\text{ExpandToSum}[(p + 1)*(b^2 - 4*a*c)*Q - (2*p + 3)*(2*c*f - b*g), x], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{PolyQ}[Pq, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{LtQ}[p, -1]$
- rule 7287 $\text{Int}[u_, x_Symbol] \rightarrow \text{With}[\{lst = \text{FunctionOfSquareRootOfQuadratic}[u, x]\}, \text{Simp}[2 \ \text{Subst}[\text{Int}[lst[[1]], x], x, lst[[2]]], x] /; \ !\text{FalseQ}[lst] \ \&\& \ \text{EqQ}[lst[[3]], 3] /; \text{EulerIntegrandQ}[u, x]$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.30 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.75

method	result
trager	$\frac{(4x^3+10x^2+12x+9)x}{4(2x^2+4x+3)^2} - \frac{(8x^3+22x^2+26x+15)\sqrt{-x^2-4x-3}}{4(2x^2+4x+3)^2} - \frac{3 \operatorname{RootOf}(_Z^2+2) \ln\left(-\frac{2 \operatorname{RootOf}(_Z^2+2)x+3 \operatorname{RootOf}(_Z^2+2)}{\operatorname{RootOf}(_Z^2+2)x}\right)}{8}$
default	Expression too large to display

input `int(1/(x+(-x^2-4*x-3)^(1/2))^3,x,method=_RETURNVERBOSE)`

output `1/4*(4*x^3+10*x^2+12*x+9)*x/(2*x^2+4*x+3)^2-1/4*(8*x^3+22*x^2+26*x+15)/(2*x^2+4*x+3)^2*(-x^2-4*x-3)^(1/2)-3/8*RootOf(_Z^2+2)*ln(-(2*RootOf(_Z^2+2)*x+3*RootOf(_Z^2+2)-2*(-x^2-4*x-3)^(1/2))/(RootOf(_Z^2+2)*x+2*x+3))`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.98

$$\int \frac{1}{(x + \sqrt{-3 - 4x - x^2})^3} dx = \frac{24x^3 + 6\sqrt{2}(4x^4 + 16x^3 + 28x^2 + 24x + 9) \arctan(\sqrt{2}(x + 1)) - 3\sqrt{2}(4x^4 + 16x^3 + 28x^2 + 24x + 9)}{16(4x^4 + 16x^3 + 28x^2 + 24x + 9)}$$

input `integrate(1/(x+(-x^2-4*x-3)^(1/2))^3,x, algorithm="fricas")`

output `-1/16*(24*x^3 + 6*sqrt(2)*(4*x^4 + 16*x^3 + 28*x^2 + 24*x + 9)*arctan(sqrt(2)*(x + 1)) - 3*sqrt(2)*(4*x^4 + 16*x^3 + 28*x^2 + 24*x + 9)*arctan(1/4*sqrt(2)*(6*x^2 + 20*x + 15)*sqrt(-x^2 - 4*x - 3)/(2*x^3 + 11*x^2 + 18*x + 9)) + 64*x^2 + 4*(8*x^3 + 22*x^2 + 26*x + 15)*sqrt(-x^2 - 4*x - 3) + 60*x + 36)/(4*x^4 + 16*x^3 + 28*x^2 + 24*x + 9)`

Sympy [F]

$$\int \frac{1}{(x + \sqrt{-3 - 4x - x^2})^3} dx = \int \frac{1}{(x + \sqrt{-x^2 - 4x - 3})^3} dx$$

input `integrate(1/(x+(-x**2-4*x-3)**(1/2))**3,x)`

output `Integral((x + sqrt(-x**2 - 4*x - 3))**(-3), x)`

Maxima [F]

$$\int \frac{1}{(x + \sqrt{-3 - 4x - x^2})^3} dx = \int \frac{1}{(x + \sqrt{-x^2 - 4x - 3})^3} dx$$

input `integrate(1/(x+(-x^2-4*x-3)^(1/2))^3,x, algorithm="maxima")`

output `integrate((x + sqrt(-x^2 - 4*x - 3))^-3, x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 367 vs. $2(154) = 308$.

Time = 0.14 (sec) , antiderivative size = 367, normalized size of antiderivative = 2.11

$$\int \frac{1}{(x + \sqrt{-3 - 4x - x^2})^3} dx$$

$$= -\frac{3}{8} \sqrt{2} \arctan(\sqrt{2}(x+1)) + \frac{3}{8} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2} \left(\frac{3(\sqrt{-x^2 - 4x - 3} - 1)}{x+2} + 1\right)\right)$$

$$+ \frac{3}{8} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2} \left(\frac{\sqrt{-x^2 - 4x - 3} - 1}{x+2} + 1\right)\right) - \frac{6x^3 + 16x^2 + 15x + 9}{4(2x^2 + 4x + 3)^2}$$

$$+ \frac{618(\sqrt{-x^2 - 4x - 3} - 1)}{x+2} + \frac{1547(\sqrt{-x^2 - 4x - 3} - 1)^2}{(x+2)^2} + \frac{2362(\sqrt{-x^2 - 4x - 3} - 1)^3}{(x+2)^3} + \frac{2223(\sqrt{-x^2 - 4x - 3} - 1)^4}{(x+2)^4} + \frac{1174(\sqrt{-x^2 - 4x - 3} - 1)^5}{(x+2)^5}$$

$$+ 18 \left(\frac{8(\sqrt{-x^2 - 4x - 3} - 1)}{x+2} + \frac{14(\sqrt{-x^2 - 4x - 3} - 1)^2}{(x+2)^2} + \frac{8(\sqrt{-x^2 - 4x - 3} - 1)^3}{(x+2)^3} + \frac{3(\sqrt{-x^2 - 4x - 3} - 1)^4}{(x+2)^4} + \frac{3(\sqrt{-x^2 - 4x - 3} - 1)^5}{(x+2)^5} \right)$$

input `integrate(1/(x+(-x^2-4*x-3)^(1/2))^3,x, algorithm="giac")`

output

```
-3/8*sqrt(2)*arctan(sqrt(2)*(x + 1)) + 3/8*sqrt(2)*arctan(1/2*sqrt(2)*(3*(sqrt(-x^2 - 4*x - 3) - 1)/(x + 2) + 1)) + 3/8*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(-x^2 - 4*x - 3) - 1)/(x + 2) + 1)) - 1/4*(6*x^3 + 16*x^2 + 15*x + 9)/(2*x^2 + 4*x + 3)^2 + 1/18*(618*(sqrt(-x^2 - 4*x - 3) - 1)/(x + 2) + 1547*(sqrt(-x^2 - 4*x - 3) - 1)^2/(x + 2)^2 + 2362*(sqrt(-x^2 - 4*x - 3) - 1)^3/(x + 2)^3 + 2223*(sqrt(-x^2 - 4*x - 3) - 1)^4/(x + 2)^4 + 1174*(sqrt(-x^2 - 4*x - 3) - 1)^5/(x + 2)^5 + 377*(sqrt(-x^2 - 4*x - 3) - 1)^6/(x + 2)^6 + 6*(sqrt(-x^2 - 4*x - 3) - 1)^7/(x + 2)^7 + 117)/(8*(sqrt(-x^2 - 4*x - 3) - 1)/(x + 2) + 14*(sqrt(-x^2 - 4*x - 3) - 1)^2/(x + 2)^2 + 8*(sqrt(-x^2 - 4*x - 3) - 1)^3/(x + 2)^3 + 3*(sqrt(-x^2 - 4*x - 3) - 1)^4/(x + 2)^4 + 3)^2
```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(x + \sqrt{-3 - 4x - x^2})^3} dx = \int \frac{1}{(x + \sqrt{-x^2 - 4x - 3})^3} dx$$

input `int(1/(x + (- 4*x - x^2 - 3)^(1/2))^3,x)`output `int(1/(x + (- 4*x - x^2 - 3)^(1/2))^3, x)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.78

$$\int \frac{1}{(x + \sqrt{-3 - 4x - x^2})^3} dx$$

$$= \frac{-36\sqrt{-x^2 - 4x - 3}\sqrt{2} \operatorname{atan}\left(\frac{3 \tan\left(\frac{\operatorname{asin}(x+2)}{2}\right) - 1}{\sqrt{2}}\right) x + 72\sqrt{2} \operatorname{atan}\left(\frac{3 \tan\left(\frac{\operatorname{asin}(x+2)}{2}\right) - 1}{\sqrt{2}}\right) x + 54\sqrt{2} \operatorname{atan}\left(\frac{3 \tan\left(\frac{\operatorname{asin}(x+2)}{2}\right) - 1}{\sqrt{2}}\right) x + 10\sqrt{-x^2 - 4x - 3} + 30\sqrt{-x^2 - 4x - 3} + 24x^2 + 86x + 33}{48\sqrt{-x^2 - 4x - 3} x - 96x - 72}$$

input `int(1/(x+(-x^2-4*x-3)^(1/2))^3,x)`output `(- 36*sqrt(- x**2 - 4*x - 3)*sqrt(2)*atan((3*tan(asin(x + 2)/2) - 1)/sqrt(2))*x + 72*sqrt(2)*atan((3*tan(asin(x + 2)/2) - 1)/sqrt(2))*x + 54*sqrt(2)*atan((3*tan(asin(x + 2)/2) - 1)/sqrt(2)) - 10*sqrt(- x**2 - 4*x - 3)*x + 30*sqrt(- x**2 - 4*x - 3) + 24*x**2 + 86*x + 33)/(24*(2*sqrt(- x**2 - 4*x - 3)*x - 4*x - 3))`

3.46 $\int (d + ex + f\sqrt{-a + bx - cx^2})^2 dx$

Optimal result	373
Mathematica [A] (verified)	374
Rubi [A] (verified)	374
Maple [A] (verified)	375
Fricas [A] (verification not implemented)	376
Sympy [A] (verification not implemented)	377
Maxima [F(-2)]	378
Giac [A] (verification not implemented)	378
Mupad [B] (verification not implemented)	379
Reduce [B] (verification not implemented)	379

Optimal result

Integrand size = 26, antiderivative size = 243

$$\int (d + ex + f\sqrt{-a + bx - cx^2})^2 dx = (d^2 - af^2)x + \frac{1}{2}(2de + bf^2)x^2 + \frac{1}{3}(e^2 - cf^2)x^3$$

$$- \frac{df(b - 2cx)\sqrt{-a + bx - cx^2}}{2c}$$

$$- \frac{bef(b - 2cx)\sqrt{-a + bx - cx^2}}{4c^2}$$

$$- \frac{2ef(-a + bx - cx^2)^{3/2}}{3c}$$

$$- \frac{(b^2 - 4ac)df \arctan\left(\frac{b - 2cx}{2\sqrt{c}\sqrt{-a + bx - cx^2}}\right)}{4c^{3/2}}$$

$$- \frac{b(b^2 - 4ac)ef \arctan\left(\frac{b - 2cx}{2\sqrt{c}\sqrt{-a + bx - cx^2}}\right)}{8c^{5/2}}$$

output

```
(-a*f^2+d^2)*x+1/2*(b*f^2+2*d*e)*x^2+1/3*(-c*f^2+e^2)*x^3-1/2*d*f*(-2*c*x+b)*(-c*x^2+b*x-a)^(1/2)/c-1/4*b*e*f*(-2*c*x+b)*(-c*x^2+b*x-a)^(1/2)/c^2-2/3*e*f*(-c*x^2+b*x-a)^(3/2)/c-1/4*(-4*a*c+b^2)*d*f*arctan(1/2*(-2*c*x+b)/c^(1/2)/(-c*x^2+b*x-a)^(1/2))/c^(3/2)-1/8*b*(-4*a*c+b^2)*e*f*arctan(1/2*(-2*c*x+b)/c^(1/2)/(-c*x^2+b*x-a)^(1/2))/c^(5/2)
```

Mathematica [A] (verified)

Time = 2.22 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.68

$$\int \left(d + ex + f\sqrt{-a + bx - cx^2} \right)^2 dx$$

$$= \frac{1}{12} \left(-2x(-6d^2 + 6af^2 - 6dex + x(-3bf^2 - 2e^2x + 2cf^2x)) \right. \\ \left. + \frac{f\sqrt{-a + x(b - cx)}(-3b^2e - 2bc(3d + ex) + 4c(2ae + cx(3d + 2ex)))}{c^2} \right. \\ \left. - \frac{3(b^2 - 4ac)(2cd + be)f \arctan\left(\frac{\sqrt{cx}}{\sqrt{-a} - \sqrt{-a + x(b - cx)}}\right)}{c^{5/2}} \right)$$

input

```
Integrate[(d + e*x + f*Sqrt[-a + b*x - c*x^2])^2,x]
```

output

```
(-2*x*(-6*d^2 + 6*a*f^2 - 6*d*e*x + x*(-3*b*f^2 - 2*e^2*x + 2*c*f^2*x)) +
(f*Sqrt[-a + x*(b - c*x)]*(-3*b^2*e - 2*b*c*(3*d + e*x) + 4*c*(2*a*e + c*x
*(3*d + 2*e*x))))/c^2 - (3*(b^2 - 4*a*c)*(2*c*d + b*e)*f*ArcTan[(Sqrt[c]*x
)/(Sqrt[-a] - Sqrt[-a + x*(b - c*x)])])/c^(5/2))/12
```

Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(f\sqrt{-a + bx - cx^2} + d + ex \right)^2 dx$$

↓ 7293

$$\int \left(2df\sqrt{-a + bx - cx^2} + 2efx\sqrt{-a + bx - cx^2} + d^2 \left(1 - \frac{af^2}{d^2} \right) + 2dex \left(\frac{bf^2}{2de} + 1 \right) + e^2x^2 \left(1 - \frac{cf^2}{e^2} \right) \right) dx$$

$$\begin{aligned}
 & \downarrow 2009 \\
 & -\frac{df(b^2 - 4ac) \arctan\left(\frac{b-2cx}{2\sqrt{c}\sqrt{-a+bx-cx^2}}\right)}{4c^{3/2}} - \frac{bef(b^2 - 4ac) \arctan\left(\frac{b-2cx}{2\sqrt{c}\sqrt{-a+bx-cx^2}}\right)}{8c^{5/2}} - \\
 & \frac{bef(b-2cx)\sqrt{-a+bx-cx^2}}{4c^2} - \frac{df(b-2cx)\sqrt{-a+bx-cx^2}}{2c} - \frac{2ef(-a+bx-cx^2)^{3/2}}{3c} + \\
 & x(d^2 - af^2) + \frac{1}{2}x^2(bf^2 + 2de) + \frac{1}{3}x^3(e^2 - cf^2)
 \end{aligned}$$

input `Int[(d + e*x + f*Sqrt[-a + b*x - c*x^2])^2,x]`

output `(d^2 - a*f^2)*x + ((2*d*e + b*f^2)*x^2)/2 + ((e^2 - c*f^2)*x^3)/3 - (d*f*(b - 2*c*x)*Sqrt[-a + b*x - c*x^2])/(2*c) - (b*e*f*(b - 2*c*x)*Sqrt[-a + b*x - c*x^2])/(4*c^2) - (2*e*f*(-a + b*x - c*x^2)^(3/2))/(3*c) - ((b^2 - 4*a*c)*d*f*ArcTan[(b - 2*c*x)/(2*Sqrt[c]*Sqrt[-a + b*x - c*x^2])])/(4*c^(3/2)) - (b*(b^2 - 4*a*c)*e*f*ArcTan[(b - 2*c*x)/(2*Sqrt[c]*Sqrt[-a + b*x - c*x^2])])/(8*c^(5/2))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

Maple [A] (verified)

Time = 1.75 (sec) , antiderivative size = 210, normalized size of antiderivative = 0.86

method	result
default	$ f^2\left(-\frac{1}{3}cx^3 + \frac{1}{2}bx^2 - xa\right) + 2f \left(d \left(-\frac{(-2cx+b)\sqrt{-cx^2+bx-a}}{4c} - \frac{(4ac-b^2) \arctan\left(\frac{\sqrt{c}\left(x-\frac{b}{2c}\right)}{\sqrt{-cx^2+bx-a}}\right)}{8c^{3/2}} \right) + e \left(-\right. \right. $

input `int((d+e*x+f*(-c*x^2+b*x-a)^(1/2))^2,x,method=_RETURNVERBOSE)`

output $f^2*(-1/3*c*x^3+1/2*b*x^2-x*a)+2*f*(d*(-1/4*(-2*c*x+b)/c*(-c*x^2+b*x-a)^(1/2)-1/8*(4*a*c-b^2)/c^(3/2)*arctan(c^(1/2)*(x-1/2*b/c)/(-c*x^2+b*x-a)^(1/2))))+e*(-1/3*(-c*x^2+b*x-a)^(3/2)/c+1/2*b/c*(-1/4*(-2*c*x+b)/c*(-c*x^2+b*x-a)^(1/2)-1/8*(4*a*c-b^2)/c^(3/2)*arctan(c^(1/2)*(x-1/2*b/c)/(-c*x^2+b*x-a)^(1/2))))+1/3*(e*x+d)^3/e$

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 450, normalized size of antiderivative = 1.85

$$\int (d + ex + f\sqrt{-a + bx - cx^2})^2 dx$$

$$= \left[-\frac{16(c^4 f^2 - c^3 e^2)x^3 - 3(2(b^2 c - 4ac^2)d + (b^3 - 4abc)e)\sqrt{-c}f \log(-8c^2 x^2 + 8bcx - b^2 - 4\sqrt{-cx^2 - a})}{8(c^4 f^2 - c^3 e^2)x^3 + 3(2(b^2 c - 4ac^2)d + (b^3 - 4abc)e)\sqrt{c}f \arctan\left(\frac{\sqrt{-cx^2 + bx - a}(2cx - b)\sqrt{c}}{2(c^2 x^2 - bcx + ac)}\right)} - 12(bc^3 f^2 - \dots) \right]$$

input `integrate((d+e*x+f*(-c*x^2+b*x-a)^(1/2))^2,x, algorithm="fricas")`

output $[-1/48*(16*(c^4*f^2 - c^3*e^2)*x^3 - 3*(2*(b^2*c - 4*a*c^2)*d + (b^3 - 4*a*b*c)*e)*sqrt(-c)*f*log(-8*c^2*x^2 + 8*b*c*x - b^2 - 4*sqrt(-c*x^2 + b*x - a)*(2*c*x - b)*sqrt(-c) - 4*a*c) - 24*(b*c^3*f^2 + 2*c^3*d*e)*x^2 + 48*(a*c^3*f^2 - c^3*d^2)*x - 4*(8*c^3*e*f*x^2 + 2*(6*c^3*d - b*c^2*e)*f*x - (6*b*c^2*d + (3*b^2*c - 8*a*c^2)*e)*f)*sqrt(-c*x^2 + b*x - a))/c^3, -1/24*(8*(c^4*f^2 - c^3*e^2)*x^3 + 3*(2*(b^2*c - 4*a*c^2)*d + (b^3 - 4*a*b*c)*e)*sqrt(c)*f*arctan(1/2*sqrt(-c*x^2 + b*x - a)*(2*c*x - b)*sqrt(c)/(c^2*x^2 - b*c*x + a*c)) - 12*(b*c^3*f^2 + 2*c^3*d*e)*x^2 + 24*(a*c^3*f^2 - c^3*d^2)*x - 2*(8*c^3*e*f*x^2 + 2*(6*c^3*d - b*c^2*e)*f*x - (6*b*c^2*d + (3*b^2*c - 8*a*c^2)*e)*f)*sqrt(-c*x^2 + b*x - a))/c^3]$

Sympy [A] (verification not implemented)

Time = 0.77 (sec) , antiderivative size = 357, normalized size of antiderivative = 1.47

$$\int \left(d + ex + f\sqrt{-a + bx - cx^2} \right)^2 dx = -af^2x + \frac{bf^2x^2}{2} - \frac{cf^2x^3}{3} + d^2x + dex^2$$

$$+ 2df \left(\begin{cases} \left(-\frac{a}{2} + \frac{b^2}{8c} \right) \left(\begin{cases} \frac{\log(b - 2cx + 2\sqrt{-c}\sqrt{-a + bx - cx^2})}{\sqrt{-c}} & \text{for } a - \frac{b^2}{4c} \neq 0 \\ \frac{(-\frac{b}{2c} + x) \log(-\frac{b}{2c} + x)}{\sqrt{-c}(-\frac{b}{2c} + x)^2} & \text{otherwise} \end{cases} \right) + \left(-\frac{b}{4c} + \frac{x}{2} \right) \sqrt{-a + bx - cx^2} \\ \frac{2(-a + bx)^{\frac{3}{2}}}{3b} \\ x\sqrt{-a} \end{cases} \right)$$

$$+ \frac{e^2x^3}{3}$$

$$+ 2ef \left(\begin{cases} \left(-\frac{ab}{12c} + \frac{b(-\frac{a}{3} + \frac{b^2}{8c})}{2c} \right) \left(\begin{cases} \frac{\log(b - 2cx + 2\sqrt{-c}\sqrt{-a + bx - cx^2})}{\sqrt{-c}} & \text{for } a - \frac{b^2}{4c} \neq 0 \\ \frac{(-\frac{b}{2c} + x) \log(-\frac{b}{2c} + x)}{\sqrt{-c}(-\frac{b}{2c} + x)^2} & \text{otherwise} \end{cases} \right) + \sqrt{-a + bx - cx^2} \left(-\frac{b}{12c} + \frac{x}{2} \right) \\ \frac{2 \left(\frac{a(-a + bx)^{\frac{3}{2}}}{3} + \frac{(-a + bx)^{\frac{5}{2}}}{5} \right)}{b^2} \\ \frac{x^2\sqrt{-a}}{2} \end{cases} \right)$$

input `integrate((d+e*x+f*(-c*x**2+b*x-a)**(1/2))**2,x)`output `-a*f**2*x + b*f**2*x**2/2 - c*f**2*x**3/3 + d**2*x + d*e*x**2 + 2*d*f*Piecewise(((-a/2 + b**2/(8*c))*Piecewise((log(b - 2*c*x + 2*sqrt(-c)*sqrt(-a + b*x - c*x**2))/sqrt(-c), Ne(a - b**2/(4*c), 0)), ((-b/(2*c) + x)*log(-b/(2*c) + x)/sqrt(-c*(-b/(2*c) + x)**2), True)) + (-b/(4*c) + x/2)*sqrt(-a + b*x - c*x**2), Ne(c, 0)), (2*(-a + b*x)**(3/2)/(3*b), Ne(b, 0)), (x*sqrt(-a), True)) + e**2*x**3/3 + 2*e*f*Piecewise(((-a*b/(12*c) + b*(-a/3 + b**2/(8*c))/(2*c))*Piecewise((log(b - 2*c*x + 2*sqrt(-c)*sqrt(-a + b*x - c*x**2))/sqrt(-c), Ne(a - b**2/(4*c), 0)), ((-b/(2*c) + x)*log(-b/(2*c) + x)/sqrt(-c*(-b/(2*c) + x)**2), True)) + sqrt(-a + b*x - c*x**2)*(-b*x/(12*c) + x**2/3 - (-a/3 + b**2/(8*c))/c), Ne(c, 0)), (2*(a*(-a + b*x)**(3/2)/3 + (-a + b*x)**(5/2)/5)/b**2, Ne(b, 0)), (x**2*sqrt(-a)/2, True))`

Maxima [F(-2)]

Exception generated.

$$\int (d + ex + f\sqrt{-a + bx - cx^2})^2 dx = \text{Exception raised: ValueError}$$

input `integrate((d+e*x+f*(-c*x^2+b*x-a)^(1/2))^2,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a*c-b^2>0)', see `assume?` for more deta

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.79

$$\begin{aligned} & \int (d + ex + f\sqrt{-a + bx - cx^2})^2 dx \\ &= -\frac{1}{3}cf^2x^3 + \frac{1}{2}bf^2x^2 + \frac{1}{3}e^2x^3 - af^2x + dex^2 + d^2x \\ &+ \frac{1}{12}\sqrt{-cx^2 + bx - a} \left(2 \left(4efx + \frac{6c^2df - bcef}{c^2} \right) x - \frac{6bcdf + 3b^2ef - 8acef}{c^2} \right) \\ &- \frac{(2b^2cdf - 8ac^2df + b^3ef - 4abcef) \log(|2(\sqrt{-cx} - \sqrt{-cx^2 + bx - a})\sqrt{-c + b}|)}{8\sqrt{-cc^2}} \end{aligned}$$

input `integrate((d+e*x+f*(-c*x^2+b*x-a)^(1/2))^2,x, algorithm="giac")`

output `-1/3*c*f^2*x^3 + 1/2*b*f^2*x^2 + 1/3*e^2*x^3 - a*f^2*x + d*e*x^2 + d^2*x + 1/12*sqrt(-c*x^2 + b*x - a)*(2*(4*e*f*x + (6*c^2*d*f - b*c*e*f)/c^2)*x - (6*b*c*d*f + 3*b^2*e*f - 8*a*c*e*f)/c^2) - 1/8*(2*b^2*c*d*f - 8*a*c^2*d*f + b^3*e*f - 4*a*b*c*e*f)*log(abs(2*(sqrt(-c)*x - sqrt(-c*x^2 + b*x - a))*sqrt(-c) + b))/(sqrt(-c)*c^2)`

Mupad [B] (verification not implemented)

Time = 22.32 (sec) , antiderivative size = 215, normalized size of antiderivative = 0.88

$$\begin{aligned}
& \int \left(d + ex + f\sqrt{-a + bx - cx^2} \right)^2 dx \\
&= x^2 \left(\frac{bf^2}{2} + de \right) - x^3 \left(\frac{cf^2}{3} - \frac{e^2}{3} \right) - x (af^2 - d^2) + 2df \left(\frac{x}{2} - \frac{b}{4c} \right) \sqrt{-cx^2 + bx - a} \\
&\quad - \frac{ef(3b^2 + 2cxb - 8c(cx^2 + a)) \sqrt{-cx^2 + bx - a}}{12c^2} \\
&\quad + \frac{df \ln \left(\sqrt{-cx^2 + bx - a} + \frac{\frac{b}{2} - cx}{\sqrt{-c}} \right) \left(ac - \frac{b^2}{4} \right)}{(-c)^{3/2}} \\
&\quad + \frac{ef \ln \left(2\sqrt{-cx^2 + bx - a} + \frac{b - 2cx}{\sqrt{-c}} \right) (b^3 - 4abc)}{8(-c)^{5/2}}
\end{aligned}$$

input `int((d + f*(b*x - a - c*x^2)^(1/2) + e*x)^2,x)`output `x^2*(d*e + (b*f^2)/2) - x^3*((c*f^2)/3 - e^2/3) - x*(a*f^2 - d^2) + 2*d*f*(x/2 - b/(4*c))*(b*x - a - c*x^2)^(1/2) - (e*f*(3*b^2 - 8*c*(a + c*x^2) + 2*b*c*x)*(b*x - a - c*x^2)^(1/2))/(12*c^2) + (d*f*log((b*x - a - c*x^2)^(1/2) + (b/2 - c*x)/(-c)^(1/2))*(a*c - b^2/4))/(-c)^(3/2) + (e*f*log(2*(b*x - a - c*x^2)^(1/2) + (b - 2*c*x)/(-c)^(1/2))*(b^3 - 4*a*b*c))/(8*(-c)^(5/2))`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 617, normalized size of antiderivative = 2.54

$$\begin{aligned}
& \int \left(d + ex + f\sqrt{-a + bx - cx^2} \right)^2 dx \\
&= \frac{-48\sqrt{c} \operatorname{asinh} \left(\frac{-2cx+bi}{\sqrt{-4ac+b^2}} \right) a^2 b c^2 e f i - 96\sqrt{c} \operatorname{asinh} \left(\frac{-2cx+bi}{\sqrt{-4ac+b^2}} \right) a^2 c^3 d f i + 24\sqrt{c} \operatorname{asinh} \left(\frac{-2cx+bi}{\sqrt{-4ac+b^2}} \right) a b^3 c e f i + \dots}{\dots}
\end{aligned}$$

input `int((d+e*x+f*(-c*x^2+b*x-a)^(1/2))^2,x)`

output

```
( - 48*sqrt(c)*asinh((b*i - 2*c*i*x)/sqrt( - 4*a*c + b**2))*a**2*b*c**2*e*
f*i - 96*sqrt(c)*asinh((b*i - 2*c*i*x)/sqrt( - 4*a*c + b**2))*a**2*c**3*d*
f*i + 24*sqrt(c)*asinh((b*i - 2*c*i*x)/sqrt( - 4*a*c + b**2))*a*b**3*c*e*f
*i + 48*sqrt(c)*asinh((b*i - 2*c*i*x)/sqrt( - 4*a*c + b**2))*a*b**2*c**2*d
*f*i - 3*sqrt(c)*asinh((b*i - 2*c*i*x)/sqrt( - 4*a*c + b**2))*b**5*e*f*i -
6*sqrt(c)*asinh((b*i - 2*c*i*x)/sqrt( - 4*a*c + b**2))*b**4*c*d*f*i + 16*
sqrt(a - b*x + c*x**2)*sqrt(4*a*c - b**2)*sqrt( - 4*a*c + b**2)*a*c**2*e*f
- 6*sqrt(a - b*x + c*x**2)*sqrt(4*a*c - b**2)*sqrt( - 4*a*c + b**2)*b**2*
c*e*f - 12*sqrt(a - b*x + c*x**2)*sqrt(4*a*c - b**2)*sqrt( - 4*a*c + b**2)
*b*c**2*d*f - 4*sqrt(a - b*x + c*x**2)*sqrt(4*a*c - b**2)*sqrt( - 4*a*c +
b**2)*b*c**2*e*f*x + 24*sqrt(a - b*x + c*x**2)*sqrt(4*a*c - b**2)*sqrt( -
4*a*c + b**2)*c**3*d*f*x + 16*sqrt(a - b*x + c*x**2)*sqrt(4*a*c - b**2)*sq
rt( - 4*a*c + b**2)*c**3*e*f*x**2 - 96*a**2*c**4*f**2*x + 24*a*b**2*c**3*f
**2*x + 48*a*b*c**4*f**2*x**2 - 32*a*c**5*f**2*x**3 + 96*a*c**4*d**2*x + 9
6*a*c**4*d*e*x**2 + 32*a*c**4*e**2*x**3 - 12*b**3*c**3*f**2*x**2 + 8*b**2*
c**4*f**2*x**3 - 24*b**2*c**3*d**2*x - 24*b**2*c**3*d*e*x**2 - 8*b**2*c**3
*e**2*x**3)/(24*c**3*(4*a*c - b**2))
```

3.47 $\int (d + ex + f\sqrt{-a + bx - cx^2}) dx$

Optimal result	381
Mathematica [A] (verified)	381
Rubi [A] (verified)	382
Maple [A] (verified)	383
Fricas [A] (verification not implemented)	383
Sympy [A] (verification not implemented)	384
Maxima [F(-2)]	385
Giac [A] (verification not implemented)	385
Mupad [B] (verification not implemented)	386
Reduce [B] (verification not implemented)	386

Optimal result

Integrand size = 24, antiderivative size = 94

$$\int (d + ex + f\sqrt{-a + bx - cx^2}) dx = dx + \frac{ex^2}{2} - \frac{f(b - 2cx)\sqrt{-a + bx - cx^2}}{4c} - \frac{(b^2 - 4ac) f \arctan\left(\frac{b - 2cx}{2\sqrt{c}\sqrt{-a + bx - cx^2}}\right)}{8c^{3/2}}$$

output

```
d*x+1/2*e*x^2-1/4*f*(-2*c*x+b)*(-c*x^2+b*x-a)^(1/2)/c-1/8*(-4*a*c+b^2)*f*a
rctan(1/2*(-2*c*x+b)/c^(1/2)/(-c*x^2+b*x-a)^(1/2))/c^(3/2)
```

Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.43

$$\begin{aligned} & \int (d + ex + f\sqrt{-a + bx - cx^2}) dx \\ &= dx + \frac{ex^2}{2} + \frac{f(-b + 2cx)\sqrt{-a + x(b - cx)}}{4c} \\ &+ \frac{(-b^2 + 4ac) f \sqrt{-a + x(b - cx)} \arctan\left(\frac{\sqrt{cx}}{\sqrt{-a} - \sqrt{-a + bx - cx^2}}\right)}{4c^{3/2}\sqrt{-a + bx - cx^2}} \end{aligned}$$

input `Integrate[d + e*x + f*Sqrt[-a + b*x - c*x^2],x]`

output `d*x + (e*x^2)/2 + (f*(-b + 2*c*x)*Sqrt[-a + x*(b - c*x)]/(4*c) + ((-b^2 + 4*a*c)*f*Sqrt[-a + x*(b - c*x)]*ArcTan[(Sqrt[c]*x)/(Sqrt[-a] - Sqrt[-a + b*x - c*x^2])])/(4*c^(3/2)*Sqrt[-a + b*x - c*x^2])`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(f \sqrt{-a + bx - cx^2} + d + ex \right) dx$$

$$\downarrow \text{2009}$$

$$-\frac{f(b^2 - 4ac) \arctan\left(\frac{b-2cx}{2\sqrt{c}\sqrt{-a+bx-cx^2}}\right)}{8c^{3/2}} - \frac{f(b-2cx)\sqrt{-a+bx-cx^2}}{4c} + dx + \frac{ex^2}{2}$$

input `Int[d + e*x + f*Sqrt[-a + b*x - c*x^2],x]`

output `d*x + (e*x^2)/2 - (f*(b - 2*c*x)*Sqrt[-a + b*x - c*x^2])/(4*c) - ((b^2 - 4*a*c)*f*ArcTan[(b - 2*c*x)/(2*Sqrt[c]*Sqrt[-a + b*x - c*x^2])])/(8*c^(3/2))`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.88

method	result	size
default	$dx + \frac{ex^2}{2} + f \left(-\frac{(-2cx+b)\sqrt{-cx^2+bx-a}}{4c} - \frac{(4ac-b^2) \arctan\left(\frac{\sqrt{c}\left(x-\frac{b}{2c}\right)}{\sqrt{-cx^2+bx-a}}\right)}{8c^{\frac{3}{2}}}\right)$	83
parts	$dx + \frac{ex^2}{2} + f \left(-\frac{(-2cx+b)\sqrt{-cx^2+bx-a}}{4c} - \frac{(4ac-b^2) \arctan\left(\frac{\sqrt{c}\left(x-\frac{b}{2c}\right)}{\sqrt{-cx^2+bx-a}}\right)}{8c^{\frac{3}{2}}}\right)$	83

input `int(d+e*x+f*(-c*x^2+b*x-a)^(1/2),x,method=_RETURNVERBOSE)`

output `d*x+1/2*e*x^2+f*(-1/4*(-2*c*x+b)/c*(-c*x^2+b*x-a)^(1/2)-1/8*(4*a*c-b^2)/c^(3/2)*arctan(c^(1/2)*(x-1/2*b/c)/(-c*x^2+b*x-a)^(1/2)))`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 235, normalized size of antiderivative = 2.50

$$\int (d + ex + f\sqrt{-a + bx - cx^2}) dx$$

$$= \left[\frac{8c^2ex^2 + 16c^2dx + (b^2 - 4ac)\sqrt{-c}f \log(-8c^2x^2 + 8bcx - b^2 - 4\sqrt{-cx^2 + bx - a}(2cx - b)\sqrt{-c} - 4}{16c^2} \right]$$

input `integrate(d+e*x+f*(-c*x^2+b*x-a)^(1/2),x, algorithm="fricas")`

output

```
[1/16*(8*c^2*e*x^2 + 16*c^2*d*x + (b^2 - 4*a*c)*sqrt(-c)*f*log(-8*c^2*x^2
+ 8*b*c*x - b^2 - 4*sqrt(-c*x^2 + b*x - a)*(2*c*x - b)*sqrt(-c) - 4*a*c) +
4*(2*c^2*f*x - b*c*f)*sqrt(-c*x^2 + b*x - a))/c^2, 1/8*(4*c^2*e*x^2 + 8*c
^2*d*x - (b^2 - 4*a*c)*sqrt(c)*f*arctan(1/2*sqrt(-c*x^2 + b*x - a)*(2*c*x
- b)*sqrt(c)/(c^2*x^2 - b*c*x + a*c)) + 2*(2*c^2*f*x - b*c*f)*sqrt(-c*x^2
+ b*x - a))/c^2]
```

Sympy [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.47

$$\int (d + ex + f\sqrt{-a + bx - cx^2}) dx = dx + \frac{ex^2}{2} + f \left(\begin{cases} \left(-\frac{a}{2} + \frac{b^2}{8c} \right) \left(\begin{cases} \frac{\log(b - 2cx + 2\sqrt{-c}\sqrt{-a + bx - cx^2})}{\sqrt{-c}} & \text{for } a - \frac{b^2}{4c} \neq 0 \\ \frac{(-\frac{b}{2c} + x) \log(-\frac{b}{2c} + x)}{\sqrt{-c}(-\frac{b}{2c} + x)^2} & \text{otherwise} \end{cases} \right) + \left(-\frac{b}{4c} + \frac{x}{2} \right) \sqrt{-a + bx - cx^2} & \text{for } a - \frac{b^2}{4c} \neq 0 \\ \frac{2(-a + bx)^{\frac{3}{2}}}{3b} & \text{for } a - \frac{b^2}{4c} = 0 \\ x\sqrt{-a} & \text{otherwise} \end{cases} \right)$$

input

```
integrate(d+e*x+f*(-c*x**2+b*x-a)**(1/2),x)
```

output

```
d*x + e*x**2/2 + f*Piecewise((( -a/2 + b**2/(8*c))*Piecewise((log(b - 2*c*x
+ 2*sqrt(-c)*sqrt(-a + b*x - c*x**2))/sqrt(-c), Ne(a - b**2/(4*c), 0)), (
(-b/(2*c) + x)*log(-b/(2*c) + x)/sqrt(-c*(-b/(2*c) + x)**2), True)) + (-b/
(4*c) + x/2)*sqrt(-a + b*x - c*x**2), Ne(c, 0)), (2*(-a + b*x)**(3/2)/(3*b
), Ne(b, 0)), (x*sqrt(-a), True))
```

Maxima [F(-2)]

Exception generated.

$$\int (d + ex + f\sqrt{-a + bx - cx^2}) dx = \text{Exception raised: ValueError}$$

input `integrate(d+e*x+f*(-c*x^2+b*x-a)^(1/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a*c-b^2>0)', see `assume?` for more deta

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.01

$$\int (d + ex + f\sqrt{-a + bx - cx^2}) dx = \frac{1}{2} ex^2 + \frac{1}{8} \left(2\sqrt{-cx^2 + bx - a} \left(2x - \frac{b}{c} \right) - \frac{(b^2 - 4ac) \log(|2(\sqrt{-cx} - \sqrt{-cx^2 + bx - a})\sqrt{-c} + b|)}{\sqrt{-cc}} \right) f + dx$$

input `integrate(d+e*x+f*(-c*x^2+b*x-a)^(1/2),x, algorithm="giac")`

output `1/2*e*x^2 + 1/8*(2*sqrt(-c*x^2 + b*x - a)*(2*x - b/c) - (b^2 - 4*a*c)*log(abs(2*(sqrt(-c)*x - sqrt(-c*x^2 + b*x - a))*sqrt(-c) + b))/(sqrt(-c)*c))*f + d*x`

Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.90

$$\int \left(d + ex + f\sqrt{-a + bx - cx^2} \right) dx = dx + \frac{ex^2}{2} + f \left(\frac{x}{2} - \frac{b}{4c} \right) \sqrt{-cx^2 + bx - a} + \frac{f \ln \left(\sqrt{-cx^2 + bx - a} + \frac{\frac{b}{2} - cx}{\sqrt{-c}} \right) \left(ac - \frac{b^2}{4} \right)}{2(-c)^{3/2}}$$

input `int(d + f*(b*x - a - c*x^2)^(1/2) + e*x,x)`output `d*x + (e*x^2)/2 + f*(x/2 - b/(4*c))*(b*x - a - c*x^2)^(1/2) + (f*log((b*x - a - c*x^2)^(1/2) + (b/2 - c*x)/(-c)^(1/2))*(a*c - b^2/4))/(2*(-c)^(3/2))`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 232, normalized size of antiderivative = 2.47

$$\int \left(d + ex + f\sqrt{-a + bx - cx^2} \right) dx = \frac{-16\sqrt{c} \operatorname{asinh}\left(\frac{-2cix+bi}{\sqrt{-4ac+b^2}}\right) a^2 c^2 fi + 8\sqrt{c} \operatorname{asinh}\left(\frac{-2cix+bi}{\sqrt{-4ac+b^2}}\right) a b^2 c fi - \sqrt{c} \operatorname{asinh}\left(\frac{-2cix+bi}{\sqrt{-4ac+b^2}}\right) b^4 fi - 2\sqrt{c} x^2 - \dots}{\dots}$$

input `int(d+e*x+f*(-c*x^2+b*x-a)^(1/2),x)`output `(- 16*sqrt(c)*asinh((b*i - 2*c*i*x)/sqrt(- 4*a*c + b**2))*a**2*c**2*f*i + 8*sqrt(c)*asinh((b*i - 2*c*i*x)/sqrt(- 4*a*c + b**2))*a*b**2*c*f*i - sqrt(c)*asinh((b*i - 2*c*i*x)/sqrt(- 4*a*c + b**2))*b**4*f*i - 2*sqrt(a - b*x + c*x**2)*sqrt(4*a*c - b**2)*sqrt(- 4*a*c + b**2)*b*c*f + 4*sqrt(a - b*x + c*x**2)*sqrt(4*a*c - b**2)*sqrt(- 4*a*c + b**2)*c**2*f*x + 32*a*c**3*d*x + 16*a*c**3*e*x**2 - 8*b**2*c**2*d*x - 4*b**2*c**2*e*x**2)/(8*c**2*(4*a*c - b**2))`

3.48 $\int \frac{1}{d+ex+f\sqrt{-a+bx-cx^2}} dx$

Optimal result	387
Mathematica [C] (verified)	388
Rubi [A] (verified)	389
Maple [B] (warning: unable to verify)	391
Fricas [F(-1)]	392
Sympy [F]	392
Maxima [F]	392
Giac [F(-2)]	393
Mupad [F(-1)]	393
Reduce [F]	393

Optimal result

Integrand size = 26, antiderivative size = 621

$$\int \frac{1}{d+ex+f\sqrt{-a+bx-cx^2}} dx = \frac{2\sqrt{c}f \arctan\left(\frac{2\sqrt{c}\sqrt{-a+bx-cx^2}}{b-\sqrt{b^2-4ac}-2cx}\right)}{e^2+cf^2}$$

$$+ \frac{2(2cd+be)f \arctan\left(\frac{\sqrt{b^2-4ac}f - \frac{4cd\sqrt{-a+bx-cx^2}}{b-\sqrt{b^2-4ac}-2cx} - \frac{2(b-\sqrt{b^2-4ac})e\sqrt{-a+bx-cx^2}}{b-\sqrt{b^2-4ac}-2cx}}{\sqrt{4bde+4ae^2-b^2f^2+4c(d^2+af^2)}}\right)}{(e^2+cf^2)\sqrt{4bde+4ae^2-b^2f^2+4c(d^2+af^2)}}$$

$$- \frac{e \log\left(\frac{c(b^2-4ac-b\sqrt{b^2-4ac}+2c\sqrt{b^2-4ac}x)}{(b-\sqrt{b^2-4ac}-2cx)^2}\right)}{e^2+cf^2}$$

$$+ \frac{e \log\left(\frac{c^2(b^2-4ac-b\sqrt{b^2-4ac})d+2c^3\sqrt{b^2-4ac}dx+b^2c^2ex-4ac^3ex-bc^2\sqrt{b^2-4ac}ex+2c^3\sqrt{b^2-4ac}ex^2+b^2c^2f\sqrt{-a+bx-cx^2}-4ac^3f\sqrt{-a+bx-cx^2}}{(b-\sqrt{b^2-4ac}-2cx)^2}\right)}{e^2+cf^2}$$

output

```

2*c^(1/2)*f*arctan(2*c^(1/2)*(-c*x^2+b*x-a)^(1/2)/(b-(-4*a*c+b^2)^(1/2)-2*
c*x))/(c*f^2+e^2)+2*(b*e+2*c*d)*f*arctan((f*(-4*a*c+b^2)^(1/2)-4*c*d*(-c*x
^2+b*x-a)^(1/2)/(b-(-4*a*c+b^2)^(1/2)-2*c*x))-2*(b-(-4*a*c+b^2)^(1/2))*e*(-
c*x^2+b*x-a)^(1/2)/(b-(-4*a*c+b^2)^(1/2)-2*c*x))/(4*b*d*e+4*a*e^2-b^2*f^2+
4*c*(a*f^2+d^2)^(1/2))/(c*f^2+e^2)/(4*b*d*e+4*a*e^2-b^2*f^2+4*c*(a*f^2+d^
2))^(1/2)-e*ln(c*(b^2-4*a*c-b*(-4*a*c+b^2)^(1/2)+2*(-4*a*c+b^2)^(1/2)*c*x)
/(b-(-4*a*c+b^2)^(1/2)-2*c*x)^2)/(c*f^2+e^2)+e*ln((c^2*(b^2-4*a*c-b*(-4*a*
c+b^2)^(1/2))*d+2*c^3*(-4*a*c+b^2)^(1/2)*d*x+b^2*c^2*e*x-4*a*c^3*e*x-b*c^2
*(-4*a*c+b^2)^(1/2)*e*x+2*c^3*(-4*a*c+b^2)^(1/2)*e*x^2+b^2*c^2*f*(-c*x^2+b
*x-a)^(1/2)-4*a*c^3*f*(-c*x^2+b*x-a)^(1/2)-b*c^2*(-4*a*c+b^2)^(1/2)*f*(-c*
x^2+b*x-a)^(1/2)+2*c^3*(-4*a*c+b^2)^(1/2)*f*x*(-c*x^2+b*x-a)^(1/2))/(b-(-4
*a*c+b^2)^(1/2)-2*c*x)^2)/(c*f^2+e^2)

```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 2.47 (sec) , antiderivative size = 1737, normalized size of antiderivative = 2.80

$$\int \frac{1}{d + ex + f\sqrt{-a + bx - cx^2}} dx = \text{Too large to display}$$

input

```
Integrate[(d + e*x + f*Sqrt[-a + b*x - c*x^2])^(-1),x]
```

output

```

(-2*Sqrt[c]*f*ArcTan[(Sqrt[c]*x)/(Sqrt[-a] - Sqrt[-a + x*(b - c*x)])] + 2*
e*Log[x] - e*Log[2*a - b*x + 2*Sqrt[-a]*Sqrt[-a + x*(b - c*x)]] - Sqrt[-a]
*(c*d + b*e)*(e^2 + c*f^2)*RootSum[c^2*d^2 + 2*b*c*d*e + b^2*e^2 + a*c^2*f
^2 + 2*b*c*d*f**1 + 2*b^2*e*f**1 - 4*a*c*e*f**1 + 2*c*d^2**1^2 + 2*b*d*e**
1^2 + 4*a*e^2**1^2 + b^2*f^2**1^2 - 2*a*c*f^2**1^2 + 2*b*d*f**1^3 + 4*a*e*
f**1^3 + d^2**1^4 + a*f^2**1^4 & , (-Log[x] + Log[Sqrt[-a] - Sqrt[-a + b*x
- c*x^2] + x**1])/(b*c*d*f + b^2*e*f - 2*a*c*e*f + 2*c*d^2**1 + 2*b*d*e**
1 + 4*a*e^2**1 + b^2*f^2**1 - 2*a*c*f^2**1 + 3*b*d*f**1^2 + 6*a*e*f**1^2 +
2*d^2**1^3 + 2*a*f^2**1^3) & ] + Sqrt[-a]*d*(e^2 + c*f^2)*RootSum[c^2*d^2
+ 2*b*c*d*e + b^2*e^2 + a*c^2*f^2 + 2*b*c*d*f**1 + 2*b^2*e*f**1 - 4*a*c*e
*f**1 + 2*c*d^2**1^2 + 2*b*d*e**1^2 + 4*a*e^2**1^2 + b^2*f^2**1^2 - 2*a*c*
f^2**1^2 + 2*b*d*f**1^3 + 4*a*e*f**1^3 + d^2**1^4 + a*f^2**1^4 & , (-Log[x]
**1^2) + Log[Sqrt[-a] - Sqrt[-a + b*x - c*x^2] + x**1]**1^2)/(b*c*d*f +
b^2*e*f - 2*a*c*e*f + 2*c*d^2**1 + 2*b*d*e**1 + 4*a*e^2**1 + b^2*f^2**1 -
2*a*c*f^2**1 + 3*b*d*f**1^2 + 6*a*e*f**1^2 + 2*d^2**1^3 + 2*a*f^2**1^3) &
] + RootSum[c^2*d^2 + 2*b*c*d*e + b^2*e^2 + a*c^2*f^2 + 2*b*c*d*f**1 + 2*b
^2*e*f**1 - 4*a*c*e*f**1 + 2*c*d^2**1^2 + 2*b*d*e**1^2 + 4*a*e^2**1^2 + b
^2*f^2**1^2 - 2*a*c*f^2**1^2 + 2*b*d*f**1^3 + 4*a*e*f**1^3 + d^2**1^4 + a*f
^2**1^4 & , -(c^2*d^2*f*Log[x]) - 2*b*c*d*e*f*Log[x] - b^2*e^2*f*Log[x] +
a*c*e^2*f*Log[x] + c^2*d^2*f*Log[Sqrt[-a] - Sqrt[-a + b*x - c*x^2] + x...

```

Rubi [A] (verified)

Time = 1.88 (sec) , antiderivative size = 397, normalized size of antiderivative = 0.64, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{f\sqrt{-a+bx-cx^2+d+ex}} dx$$

↓ 7293

$$\int \left(\frac{f\sqrt{-a+bx-cx^2}}{-af^2-x(2de-bf^2)-x^2(cf^2+e^2)-d^2} + \frac{d+ex}{af^2+x(2de-bf^2)+x^2(cf^2+e^2)+d^2} \right) dx$$

↓ 2009

$$\frac{f(be + 2cd) \arctan\left(\frac{2ae - x(be + 2cd) + bd}{\sqrt{-a + bx - cx^2} \sqrt{4c(af^2 + d^2) + 4ae^2 - b^2f^2 + 4bde}}\right)}{(cf^2 + e^2) \sqrt{4c(af^2 + d^2) + 4ae^2 - b^2f^2 + 4bde}} +$$

$$\frac{f(be + 2cd) \arctan\left(\frac{-bf^2 + 2x(cf^2 + e^2) + 2de}{f \sqrt{4c(af^2 + d^2) + 4ae^2 - b^2f^2 + 4bde}}\right)}{(cf^2 + e^2) \sqrt{4c(af^2 + d^2) + 4ae^2 - b^2f^2 + 4bde}} - \frac{\sqrt{c} f \arctan\left(\frac{b - 2cx}{2\sqrt{c}\sqrt{-a + bx - cx^2}}\right)}{cf^2 + e^2} +$$

$$\frac{\operatorname{earctanh}\left(\frac{d + ex}{f \sqrt{-a + bx - cx^2}}\right)}{cf^2 + e^2} + \frac{e \log(af^2 + x(2de - bf^2) + x^2(cf^2 + e^2) + d^2)}{2(cf^2 + e^2)}$$

input

```
Int[(d + e*x + f*Sqrt[-a + b*x - c*x^2])^(-1),x]
```

output

```
((2*c*d + b*e)*f*ArcTan[(2*d*e - b*f^2 + 2*(e^2 + c*f^2)*x)/(f*Sqrt[4*b*d*e + 4*a*e^2 - b^2*f^2 + 4*c*(d^2 + a*f^2)])]/((e^2 + c*f^2)*Sqrt[4*b*d*e + 4*a*e^2 - b^2*f^2 + 4*c*(d^2 + a*f^2)]) - (Sqrt[c]*f*ArcTan[(b - 2*c*x)/(2*Sqrt[c]*Sqrt[-a + b*x - c*x^2])]/(e^2 + c*f^2) + ((2*c*d + b*e)*f*ArcTan[(b*d + 2*a*e - (2*c*d + b*e)*x)/(Sqrt[4*b*d*e + 4*a*e^2 - b^2*f^2 + 4*c*(d^2 + a*f^2)]*Sqrt[-a + b*x - c*x^2])]/((e^2 + c*f^2)*Sqrt[4*b*d*e + 4*a*e^2 - b^2*f^2 + 4*c*(d^2 + a*f^2)]) + (e*ArcTanh[(d + e*x)/(f*Sqrt[-a + b*x - c*x^2])]/(e^2 + c*f^2) + (e*Log[d^2 + a*f^2 + (2*d*e - b*f^2)*x + (e^2 + c*f^2)*x^2])/(2*(e^2 + c*f^2)))
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 7293

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]
```

Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 4814 vs. $2(569) = 1138$.

Time = 0.15 (sec) , antiderivative size = 4815, normalized size of antiderivative = 7.75

method	result	size
default	Expression too large to display	4815

input `int(1/(d+e*x+f*(-c*x^2+b*x-a)^(1/2)),x,method=_RETURNVERBOSE)`

output

```
f*(2*(c*f^2+e^2)/(-f^2*(4*a*c*f^2-b^2*f^2+4*a*e^2+4*b*d*e+4*c*d^2))^(1/2)/
(2*c*f^2+2*e^2)*(1/2*(-4*(x+(-b*f^2+2*d*e+(-f^2*(4*a*c*f^2-b^2*f^2+4*a*e^2
+4*b*d*e+4*c*d^2))^(1/2))/(2*c*f^2+2*e^2))^2*c+4*(b*e^2+2*c*d*e+(f^2*(-4*a
*c*f^2+b^2*f^2-4*a*e^2-4*b*d*e-4*c*d^2))^(1/2)*c)/(c*f^2+e^2)*(x+(-b*f^2+2
*d*e+(-f^2*(4*a*c*f^2-b^2*f^2+4*a*e^2+4*b*d*e+4*c*d^2))^(1/2))/(2*c*f^2+2*
e^2))-2*(2*a*c*e^2*f^2-b^2*e^2*f^2-2*b*c*d*e*f^2-2*c^2*d^2*f^2+2*e^4*a+2*d
*e^3*b+2*d^2*e^2*c+(f^2*(-4*a*c*f^2+b^2*f^2-4*a*e^2-4*b*d*e-4*c*d^2))^(1/2
)*b*e^2+2*(f^2*(-4*a*c*f^2+b^2*f^2-4*a*e^2-4*b*d*e-4*c*d^2))^(1/2)*c*d*e)/
(c*f^2+e^2)^2)^(1/2)+1/2*(b*e^2+2*c*d*e+(f^2*(-4*a*c*f^2+b^2*f^2-4*a*e^2-4
*b*d*e-4*c*d^2))^(1/2)*c)/(c*f^2+e^2)/c^(1/2)*arctan(2*c^(1/2)*(x+(-b*f^2+
2*d*e+(-f^2*(4*a*c*f^2-b^2*f^2+4*a*e^2+4*b*d*e+4*c*d^2))^(1/2))/(2*c*f^2+2
*e^2)-1/2*(b*e^2+2*c*d*e+(f^2*(-4*a*c*f^2+b^2*f^2-4*a*e^2-4*b*d*e-4*c*d^2)
)^(1/2)*c)/(c*f^2+e^2)/c)/(-4*(x+(-b*f^2+2*d*e+(-f^2*(4*a*c*f^2-b^2*f^2+4*
a*e^2+4*b*d*e+4*c*d^2))^(1/2))/(2*c*f^2+2*e^2))^2*c+4*(b*e^2+2*c*d*e+(f^2*
(-4*a*c*f^2+b^2*f^2-4*a*e^2-4*b*d*e-4*c*d^2))^(1/2)*c)/(c*f^2+e^2)*(x+(-b*
f^2+2*d*e+(-f^2*(4*a*c*f^2-b^2*f^2+4*a*e^2+4*b*d*e+4*c*d^2))^(1/2))/(2*c*f
^2+2*e^2))-2*(2*a*c*e^2*f^2-b^2*e^2*f^2-2*b*c*d*e*f^2-2*c^2*d^2*f^2+2*e^4*
a+2*d*e^3*b+2*d^2*e^2*c+(f^2*(-4*a*c*f^2+b^2*f^2-4*a*e^2-4*b*d*e-4*c*d^2))
^(1/2)*b*e^2+2*(f^2*(-4*a*c*f^2+b^2*f^2-4*a*e^2-4*b*d*e-4*c*d^2))^(1/2)*c*
d*e)/(c*f^2+e^2)^2)^(1/2)+(2*a*c*e^2*f^2-b^2*e^2*f^2-2*b*c*d*e*f^2-2*c...
```


Fricas [F(-1)]

Timed out.

$$\int \frac{1}{d + ex + f\sqrt{-a + bx - cx^2}} dx = \text{Timed out}$$

input `integrate(1/(d+e*x+f*(-c*x^2+b*x-a)^(1/2)),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{1}{d + ex + f\sqrt{-a + bx - cx^2}} dx = \int \frac{1}{d + ex + f\sqrt{-a + bx - cx^2}} dx$$

input `integrate(1/(d+e*x+f*(-c*x**2+b*x-a)**(1/2)),x)`

output `Integral(1/(d + e*x + f*sqrt(-a + b*x - c*x**2)), x)`

Maxima [F]

$$\int \frac{1}{d + ex + f\sqrt{-a + bx - cx^2}} dx = \int \frac{1}{ex + \sqrt{-cx^2 + bx - a}f + d} dx$$

input `integrate(1/(d+e*x+f*(-c*x^2+b*x-a)^(1/2)),x, algorithm="maxima")`

output `integrate(1/(e*x + sqrt(-c*x^2 + b*x - a)*f + d), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{1}{d + ex + f\sqrt{-a + bx - cx^2}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(d+e*x+f*(-c*x^2+b*x-a)^(1/2)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:Error: Bad Argument Type`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{d + ex + f\sqrt{-a + bx - cx^2}} dx = \int \frac{1}{d + f\sqrt{-cx^2 + bx - a} + ex} dx$$

input `int(1/(d + f*(b*x - a - c*x^2)^(1/2) + e*x), x)`

output `int(1/(d + f*(b*x - a - c*x^2)^(1/2) + e*x), x)`

Reduce [F]

$$\int \frac{1}{d + ex + f\sqrt{-a + bx - cx^2}} dx = \int \frac{1}{d + ex + f\sqrt{-cx^2 + bx - a}} dx$$

input `int(1/(d+e*x+f*(-c*x^2+b*x-a)^(1/2)),x)`

output `int(1/(d+e*x+f*(-c*x^2+b*x-a)^(1/2)),x)`

$$3.49 \quad \int \frac{1}{\left(d+ex+f\sqrt{-a+bx-cx^2}\right)^2} dx$$

Optimal result	394
Mathematica [C] (warning: unable to verify)	395
Rubi [B] (verified)	396
Maple [B] (warning: unable to verify)	397
Fricas [B] (verification not implemented)	398
Sympy [F]	398
Maxima [F]	399
Giac [F(-2)]	399
Mupad [F(-1)]	399
Reduce [F]	400

Optimal result

Integrand size = 26, antiderivative size = 463

$$\int \frac{1}{\left(d+ex+f\sqrt{-a+bx-cx^2}\right)^2} dx$$

$$= \frac{4\sqrt{b^2-4ac}\left(2cd+(b+\sqrt{b^2-4ac})e-\frac{2c\sqrt{b^2-4ac}f\sqrt{-a+bx-cx^2}}{b-\sqrt{b^2-4ac}-2cx}\right)}{(4bde+4ae^2-b^2f^2+4c(d^2+af^2))\left(2cd+(b+\sqrt{b^2-4ac})e-\frac{4c\sqrt{b^2-4ac}f\sqrt{-a+bx-cx^2}}{b-\sqrt{b^2-4ac}-2cx}-\frac{4c(2cd+(b-\sqrt{b^2-4ac})e)}{(b-\sqrt{b^2-4ac}-2cx)}\right)}$$

$$+ \frac{4(b^2-4ac)f \arctan\left(\frac{\frac{\sqrt{b^2-4ac}f-\frac{4cd\sqrt{-a+bx-cx^2}}{b-\sqrt{b^2-4ac}-2cx}-\frac{2(b-\sqrt{b^2-4ac})e\sqrt{-a+bx-cx^2}}{b-\sqrt{b^2-4ac}-2cx}}{\sqrt{4bde+4ae^2-b^2f^2+4c(d^2+af^2)}}\right)}{(4bde+4ae^2-b^2f^2+4c(d^2+af^2))^{3/2}}$$

output

```
4*(-4*a*c+b^2)^(1/2)*(2*c*d+(b+(-4*a*c+b^2)^(1/2))*e-2*c*(-4*a*c+b^2)^(1/2)
)*f*(-c*x^2+b*x-a)^(1/2)/(b-(-4*a*c+b^2)^(1/2)-2*c*x)/(4*b*d*e+4*a*e^2-b^2
*f^2+4*c*(a*f^2+d^2))/(2*c*d+(b+(-4*a*c+b^2)^(1/2))*e-4*c*(-4*a*c+b^2)^(1
/2)*f*(-c*x^2+b*x-a)^(1/2)/(b-(-4*a*c+b^2)^(1/2)-2*c*x)-4*c*(2*c*d+(b-(-4*
a*c+b^2)^(1/2))*e)*(c*x^2-b*x+a)/(b-(-4*a*c+b^2)^(1/2)-2*c*x)+4*(-4*a*c
+b^2)*f*arctan((f*(-4*a*c+b^2)^(1/2)-4*c*d*(-c*x^2+b*x-a)^(1/2)/(b-(-4*a*c
+b^2)^(1/2)-2*c*x)-2*(b-(-4*a*c+b^2)^(1/2))*e*(-c*x^2+b*x-a)^(1/2)/(b-(-4*
a*c+b^2)^(1/2)-2*c*x))/(4*b*d*e+4*a*e^2-b^2*f^2+4*c*(a*f^2+d^2))^(1/2))/(4
*b*d*e+4*a*e^2-b^2*f^2+4*c*(a*f^2+d^2))^(3/2)
```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 49.72 (sec) , antiderivative size = 2346, normalized size of antiderivative = 5.07

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx - cx^2})^2} dx = \text{Result too large to show}$$

input `Integrate[(d + e*x + f*Sqrt[-a + b*x - c*x^2])^(-2),x]`

output

```
(-2*(2*c*d^3*e + 2*b*d^2*e^2 + 2*a*d*e^3 + b*c*d^2*f^2 + 4*a*c*d*e*f^2 + a
*b*e^2*f^2 + 2*c*d^2*e^2*x + 2*b*d*e^3*x + 2*a*e^4*x - 2*c^2*d^2*f^2*x - 2
*b*c*d*e*f^2*x - b^2*e^2*f^2*x + 2*a*c*e^2*f^2*x))/(e^2 + c*f^2)*(4*c*d^2
+ 4*b*d*e + 4*a*e^2 - b^2*f^2 + 4*a*c*f^2)*(d^2 + a*f^2 + 2*d*e*x - b*f^2
*x + e^2*x^2 + c*f^2*x^2)) + (2*(b*d*f + 2*a*e*f - 2*c*d*f*x - b*e*f*x)*Sqr
t[-a + b*x - c*x^2])/((4*c*d^2 + 4*b*d*e + 4*a*e^2 - b^2*f^2 + 4*a*c*f^2)
*(d^2 + a*f^2 + 2*d*e*x - b*f^2*x + e^2*x^2 + c*f^2*x^2)) - (2*(-b^2 + 4*a
*c)*f*ArcTan[(2*d*e - b*f^2 + 2*e^2*x + 2*c*f^2*x)/(f*Sqrt[4*c*d^2 + 4*b*d
*e + 4*a*e^2 - b^2*f^2 + 4*a*c*f^2])])/(4*c*d^2 + 4*b*d*e + 4*a*e^2 - b^2*
f^2 + 4*a*c*f^2)^(3/2) - ((-b^2 + 4*a*c)*f*((2*I)*c*d*f + I*b*e*f + e*Sqrt
[4*c*d^2 + 4*b*d*e + 4*a*e^2 - b^2*f^2 + 4*a*c*f^2])*Log[-((Sqrt[2]*(e^2 +
c*f^2)*(4*c*d^2 + 4*b*d*e + 4*a*e^2 - b^2*f^2 + 4*a*c*f^2)^(3/2)*((-2*I)*
b*d*e - (4*I)*a*e^2 + I*b^2*f^2 - (4*I)*a*c*f^2 + b*f*Sqrt[4*c*d^2 + 4*b*d
*e + 4*a*e^2 - b^2*f^2 + 4*a*c*f^2] + (4*I)*c*d*e*x + (2*I)*b*e^2*x - 2*c*
f*Sqrt[4*c*d^2 + 4*b*d*e + 4*a*e^2 - b^2*f^2 + 4*a*c*f^2]*x)))/((-b^2 + 4*a
*c)*f*(2*c*d*f + b*e*f - I*e*Sqrt[4*c*d^2 + 4*b*d*e + 4*a*e^2 - b^2*f^2 +
4*a*c*f^2])*Sqrt[-2*c*d^2*e^2 - 2*b*d*e^3 - 2*a*e^4 + 2*c^2*d^2*f^2 + 2*b*
c*d*e*f^2 + b^2*e^2*f^2 - 2*a*c*e^2*f^2 - (2*I)*c*d*e*f*Sqrt[4*c*d^2 + 4*b
*d*e + 4*a*e^2 - b^2*f^2 + 4*a*c*f^2] - I*b*e^2*f*Sqrt[4*c*d^2 + 4*b*d*e +
4*a*e^2 - b^2*f^2 + 4*a*c*f^2])*(2*d*e - b*f^2 + I*f*Sqrt[4*c*d^2 + 4*...
```

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 965 vs. $2(463) = 926$.

Time = 4.31 (sec) , antiderivative size = 965, normalized size of antiderivative = 2.08, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\left(f\sqrt{-a+bx-cx^2+d+ex}\right)^2} dx$$

↓ 7293

$$\int \left(\frac{2f^2(-a+bx-cx^2)}{(af^2+x(2de-bf^2)+x^2(cf^2+e^2)+d^2)^2} - \frac{2df\sqrt{-a+bx-cx^2}}{(af^2+x(2de-bf^2)+x^2(cf^2+e^2)+d^2)^2} - \frac{2e}{(af^2+x(2de-bf^2)+x^2(cf^2+e^2)+d^2)^2} \right) dx$$

↓ 2009

$$\frac{2e\sqrt{-cx^2+bx-a}(2(d^2+af^2)+(2de-bf^2)x)}{f(4ae^2+4bde-b^2f^2+4c(d^2+af^2))(d^2+af^2+(e^2+cf^2)x^2+(2de-bf^2)x)} - \frac{4(2ae^2+2bde-b^2f^2+2c(d^2+2af^2)) \arctan\left(\frac{-bf^2+2de+2(e^2+cf^2)x}{f\sqrt{4ae^2+4bde-b^2f^2+4c(d^2+af^2)}}\right)}{f(4ae^2+4bde-b^2f^2+4c(d^2+af^2))^{3/2}} + \frac{2 \arctan\left(\frac{-bf^2+2de+2(e^2+cf^2)x}{f\sqrt{4ae^2+4bde-b^2f^2+4c(d^2+af^2)}}\right)}{f\sqrt{4ae^2+4bde-b^2f^2+4c(d^2+af^2)}} + \frac{(b^2-4ac)e(bd+2ae)f \arctan\left(\frac{bd+2ae-(2cd+be)x}{\sqrt{4ae^2+4bde-b^2f^2+4c(d^2+af^2)}\sqrt{-cx^2+bx-a}}\right)}{(cd^2+e(bd+ae))(4ae^2+4bde-b^2f^2+4c(d^2+af^2))^{3/2}} + \frac{(b^2-4ac)d(2cd+be)f \arctan\left(\frac{bd+2ae-(2cd+be)x}{\sqrt{4ae^2+4bde-b^2f^2+4c(d^2+af^2)}\sqrt{-cx^2+bx-a}}\right)}{(cd^2+e(bd+ae))(4ae^2+4bde-b^2f^2+4c(d^2+af^2))^{3/2}} - \frac{2((2bd^2+2aed+abf^2)e^2+cd(2ed^2+bf^2d+4aef^2)-((2ae^2+2bde-b^2f^2)e^2)-2c(ed^2-bf^2d+aef^2))}{(e^2+cf^2)(4ae^2+4bde-b^2f^2+4c(d^2+af^2))(d^2+af^2+(e^2+cf^2)x^2+(2de-bf^2)x)} - \frac{2d(-bf^2+2de+2(e^2+cf^2)x)\sqrt{-cx^2+bx-a}}{f(4ae^2+4bde-b^2f^2+4c(d^2+af^2))(d^2+af^2+(e^2+cf^2)x^2+(2de-bf^2)x)}$$

input

```
Int[(d + e*x + f*Sqrt[-a + b*x - c*x^2])^(-2), x]
```

output

$$\begin{aligned} & (-2*(e^2*(2*b*d^2 + 2*a*d*e + a*b*f^2) + c*d*(2*d^2*e + b*d*f^2 + 4*a*e*f^2) \\ & - (2*c^2*d^2*f^2 - e^2*(2*b*d*e + 2*a*e^2 - b^2*f^2) - 2*c*e*(d^2*e - b \\ & *d*f^2 + a*e*f^2))*x)/((e^2 + c*f^2)*(4*b*d*e + 4*a*e^2 - b^2*f^2 + 4*c*(\\ & d^2 + a*f^2))*(d^2 + a*f^2 + (2*d*e - b*f^2)*x + (e^2 + c*f^2)*x^2)) + (2* \\ & e*(2*(d^2 + a*f^2) + (2*d*e - b*f^2)*x)*\text{Sqrt}[-a + b*x - c*x^2])/(f*(4*b*d* \\ & e + 4*a*e^2 - b^2*f^2 + 4*c*(d^2 + a*f^2))*(d^2 + a*f^2 + (2*d*e - b*f^2)* \\ & x + (e^2 + c*f^2)*x^2)) - (2*d*(2*d*e - b*f^2 + 2*(e^2 + c*f^2)*x)*\text{Sqrt}[-a \\ & + b*x - c*x^2])/(f*(4*b*d*e + 4*a*e^2 - b^2*f^2 + 4*c*(d^2 + a*f^2))*(d^2 \\ & + a*f^2 + (2*d*e - b*f^2)*x + (e^2 + c*f^2)*x^2)) + (2*\text{ArcTan}[(2*d*e - b* \\ & f^2 + 2*(e^2 + c*f^2)*x)/(f*\text{Sqrt}[4*b*d*e + 4*a*e^2 - b^2*f^2 + 4*c*(d^2 + \\ & a*f^2)])))/(f*\text{Sqrt}[4*b*d*e + 4*a*e^2 - b^2*f^2 + 4*c*(d^2 + a*f^2)]) - (4* \\ & (2*b*d*e + 2*a*e^2 - b^2*f^2 + 2*c*(d^2 + 2*a*f^2))*\text{ArcTan}[(2*d*e - b*f^2 \\ & + 2*(e^2 + c*f^2)*x)/(f*\text{Sqrt}[4*b*d*e + 4*a*e^2 - b^2*f^2 + 4*c*(d^2 + a*f^2 \\ & 2)])))/(f*(4*b*d*e + 4*a*e^2 - b^2*f^2 + 4*c*(d^2 + a*f^2))^(3/2)) + ((b^2 \\ & - 4*a*c)*e*(b*d + 2*a*e)*f*\text{ArcTan}[(b*d + 2*a*e - (2*c*d + b*e)*x)/(\text{Sqrt}[4 \\ & *b*d*e + 4*a*e^2 - b^2*f^2 + 4*c*(d^2 + a*f^2)]*\text{Sqrt}[-a + b*x - c*x^2])]) / \\ & ((c*d^2 + e*(b*d + a*e))*(4*b*d*e + 4*a*e^2 - b^2*f^2 + 4*c*(d^2 + a*f^2)) \\ & ^{(3/2)}) + ((b^2 - 4*a*c)*d*(2*c*d + b*e)*f*\text{ArcTan}[(b*d + 2*a*e - (2*c*d + \\ & b*e)*x)/(\text{Sqrt}[4*b*d*e + 4*a*e^2 - b^2*f^2 + 4*c*(d^2 + a*f^2)]*\text{Sqrt}[-a + b \\ & *x - c*x^2])]) / ((c*d^2 + e*(b*d + a*e))*(4*b*d*e + 4*a*e^2 - b^2*f^2 + \dots \end{aligned}$$

Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$$

rule 7293

$$\text{Int}[u_, x_Symbol] \rightarrow \text{With}[\{v = \text{ExpandIntegrand}[u, x]\}, \text{Int}[v, x] /; \text{SumQ}[v]]$$

Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.74 (sec) , antiderivative size = 609986, normalized size of antiderivative = 1317.46

method	result	size
default	Expression too large to display	609986

input `int(1/(d+e*x+f*(-c*x^2+b*x-a)^(1/2))^2,x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2873 vs. $2(424) = 848$.

Time = 20.12 (sec) , antiderivative size = 6908, normalized size of antiderivative = 14.92

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx - cx^2})^2} dx = \text{Too large to display}$$

input `integrate(1/(d+e*x+f*(-c*x^2+b*x-a)^(1/2))^2,x, algorithm="fricas")`

output `Too large to include`

Sympy [F]

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx - cx^2})^2} dx = \int \frac{1}{(d + ex + f\sqrt{-a + bx - cx^2})^2} dx$$

input `integrate(1/(d+e*x+f*(-c*x**2+b*x-a)**(1/2))**2,x)`

output `Integral((d + e*x + f*sqrt(-a + b*x - c*x**2))**(-2), x)`

Maxima [F]

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx - cx^2})^2} dx = \int \frac{1}{(ex + \sqrt{-cx^2 + bx - a} + d)^2} dx$$

input `integrate(1/(d+e*x+f*(-c*x^2+b*x-a)^(1/2))^2,x, algorithm="maxima")`

output `integrate((e*x + sqrt(-c*x^2 + b*x - a)*f + d)^(-2), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx - cx^2})^2} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(d+e*x+f*(-c*x^2+b*x-a)^(1/2))^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);OUTPUT:Error: Bad Argument Type`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx - cx^2})^2} dx = \int \frac{1}{(d + f\sqrt{-cx^2 + bx - a} + ex)^2} dx$$

input `int(1/(d + f*(b*x - a - c*x^2)^(1/2) + e*x)^2,x)`

output `int(1/(d + f*(b*x - a - c*x^2)^(1/2) + e*x)^2, x)`

Reduce [F]

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx - cx^2})^2} dx = \int \frac{1}{(d + ex + f\sqrt{-cx^2 + bx - a})^2} dx$$

input `int(1/(d+e*x+f*(-c*x^2+b*x-a)^(1/2))^2,x)`

output `int(1/(d+e*x+f*(-c*x^2+b*x-a)^(1/2))^2,x)`

$$3.50 \quad \int \frac{1}{\left(d+ex+f\sqrt{-a+bx-cx^2}\right)^3} dx$$

Optimal result	401
Mathematica [C] (warning: unable to verify)	402
Rubi [F]	403
Maple [B] (warning: unable to verify)	404
Fricas [F(-1)]	405
Sympy [F(-1)]	405
Maxima [F]	405
Giac [F(-2)]	406
Mupad [F(-1)]	406
Reduce [F]	406

Optimal result

Integrand size = 26, antiderivative size = 1382

$$\int \frac{1}{\left(d+ex+f\sqrt{-a+bx-cx^2}\right)^3} dx = \text{Too large to display}$$

output

```

-8*(-4*a*c+b^2)^(1/2)*(c^4*(8*a*c^2*d*f^2+e*(4*b*(-4*a*c+b^2)^(1/2)*d*e+4*
a*(-4*a*c+b^2)^(1/2)*e^2-b^3*f^2-b^2*(-4*a*c+b^2)^(1/2)*f^2)+c*(4*(-4*a*c+
b^2)^(1/2)*d^2*e-2*b^2*d*f^2+4*a*(b+(-4*a*c+b^2)^(1/2))*e*f^2))-2*c^4*f*(b
^2*(b-(-4*a*c+b^2)^(1/2))*e^2+4*c^2*((-4*a*c+b^2)^(1/2)*d^2-2*a*d*e+2*a*(-
4*a*c+b^2)^(1/2)*f^2)+c*(8*a*(-4*a*c+b^2)^(1/2)*e^2+4*b*e*((-4*a*c+b^2)^(1
/2)*d-a*e)+2*b^2*(d*e-(-4*a*c+b^2)^(1/2)*f^2)))*(-c*x^2+b*x-a)^(1/2)/(b-(-
4*a*c+b^2)^(1/2)-2*c*x)/(2*c*d+(b-(-4*a*c+b^2)^(1/2))*e)^2/(4*b*d*e+4*a*e
^2-b^2*f^2+4*c*(a*f^2+d^2))/(c*(2*c*d+(b+(-4*a*c+b^2)^(1/2))*e)-4*c^2*(-4*
a*c+b^2)^(1/2)*f*(-c*x^2+b*x-a)^(1/2)/(b-(-4*a*c+b^2)^(1/2)-2*c*x)-4*c^2*(
2*c*d+(b-(-4*a*c+b^2)^(1/2))*e)*(c*x^2-b*x+a)/(b-(-4*a*c+b^2)^(1/2)-2*c*x)
^2)^2+4*(-4*a*c+b^2)^(1/2)*(c^2*(3*b^3*(b-(-4*a*c+b^2)^(1/2))*e^2*f^2+16*c
^3*(2*a^2*f^4+a*d^2*f^2+2*d^4)+2*c*(16*a^2*e^4-2*b^3*d*e*f^2+b^4*f^4+2*a*b
*e^2*(16*d*e+3*(-4*a*c+b^2)^(1/2)*f^2)+b^2*e*(16*d^2*e-3*(-4*a*c+b^2)^(1/2
)*d*f^2-14*a*e*f^2))+c^2*(16*b*d*e*(a*f^2+4*d^2)-4*b^2*f^2*(4*a*f^2+d^2)+8
*a*e*(8*d^2*e+3*(-4*a*c+b^2)^(1/2)*d*f^2+8*a*e*f^2)))/(2*c*d+(b-(-4*a*c+b
^2)^(1/2))*e)^2+6*c^2*(b*e+2*c*d)*(b*(b-(-4*a*c+b^2)^(1/2))*e-2*c*((-4*a*c+
b^2)^(1/2)*d+2*a*e))*f*(-c*x^2+b*x-a)^(1/2)/(2*c*d+(b-(-4*a*c+b^2)^(1/2))*
e)/(b-(-4*a*c+b^2)^(1/2)-2*c*x)/(4*b*d*e+4*a*e^2-b^2*f^2+4*c*(a*f^2+d^2)
^2)/(c*(2*c*d+(b+(-4*a*c+b^2)^(1/2))*e)-4*c^2*(-4*a*c+b^2)^(1/2)*f*(-c*x^2+
b*x-a)^(1/2)/(b-(-4*a*c+b^2)^(1/2)-2*c*x)-4*c^2*(2*c*d+(b-(-4*a*c+b^2)^(1/2)

```

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 72.55 (sec) , antiderivative size = 3558, normalized size of antiderivative = 2.57

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx - cx^2})^3} dx = \text{Result too large to show}$$

input

```
Integrate[(d + e*x + f*Sqrt[-a + b*x - c*x^2])^(-3),x]
```

output

```
(2*(-4*c^2*d^4*e*f^2 - 5*b*c*d^3*e^2*f^2 - b^2*d^2*e^3*f^2 - 2*a*c*d^2*e^3*f^2 + a*b*d*e^4*f^2 + 2*a^2*e^5*f^2 - b*c^2*d^3*f^4 - 6*a*c^2*d^2*e*f^4 - 3*a*b*c*d*e^2*f^4 - a*b^2*e^3*f^4 + 2*a^2*c*e^3*f^4 - 6*c^2*d^3*e^2*f^2*x - 9*b*c*d^2*e^3*f^2*x - 3*b^2*d*e^4*f^2*x - 6*a*c*d*e^4*f^2*x - 3*a*b*e^5*f^2*x + 2*c^3*d^3*f^4*x + 3*b*c^2*d^2*e*f^4*x + 3*b^2*c*d*e^2*f^4*x - 6*a*c^2*d*e^2*f^4*x + b^3*e^3*f^4*x - 3*a*b*c*e^3*f^4*x))/((e^2 + c*f^2)^2*(4*c*d^2 + 4*b*d*e + 4*a*e^2 - b^2*f^2 + 4*a*c*f^2)*(d^2 + a*f^2 + 2*d*e*x - b*f^2*x + e^2*x^2 + c*f^2*x^2)^2) + (-8*c^2*d^4*e^3 - 16*b*c*d^3*e^4 - 8*b^2*d^2*e^5 - 16*a*c*d^2*e^5 - 16*a*b*d*e^6 - 8*a^2*e^7 + 24*c^3*d^4*e*f^2 + 48*b*c^2*d^3*e^2*f^2 + 34*b^2*c*d^2*e^3*f^2 + 8*a*c^2*d^2*e^3*f^2 + 7*b^3*d*e^4*f^2 + 20*a*b*c*d*e^4*f^2 + 4*a*b^2*e^5*f^2 + 8*a^2*c*e^5*f^2 - 6*b^2*c^2*d^2*e*f^4 + 24*a*c^3*d^2*e*f^4 - 12*b^3*c*d*e^2*f^4 + 48*a*b*c^2*d*e^2*f^4 - 2*b^4*e^3*f^4 - 2*a*b^2*c*e^3*f^4 + 40*a^2*c^2*e^3*f^4 - 3*b^3*c^2*d*f^6 + 12*a*b*c^3*d*f^6 - 6*a*b^2*c^2*e*f^6 + 24*a^2*c^3*e*f^6 + 6*b^2*c*d*e^4*f^2*x - 24*a*c^2*d*e^4*f^2*x + 3*b^3*e^5*f^2*x - 12*a*b*c*e^5*f^2*x + 12*b^2*c^2*d*e^2*f^4*x - 48*a*c^3*d*e^2*f^4*x + 6*b^3*c*e^3*f^4*x - 24*a*b*c^2*e^3*f^4*x + 6*b^2*c^3*d*f^6*x - 24*a*c^4*d*f^6*x + 3*b^3*c^2*e*f^6*x - 12*a*b*c^3*e*f^6*x))/((e^2 + c*f^2)^2*(4*c*d^2 + 4*b*d*e + 4*a*e^2 - b^2*f^2 + 4*a*c*f^2)^2*(d^2 + a*f^2 + 2*d*e*x - b*f^2*x + e^2*x^2 + c*f^2*x^2)) + Sqrt[-a + b*x - c*x^2]*((2*(2*c*d^3*e*f + 2*b*d^2*e^2*f + 2*a...
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(f\sqrt{-a+bx-cx^2}+d+ex)^3} dx$$

↓ 7293

$$\int \left(\frac{4f^2(d+ex)(-a+bx-cx^2)}{(af^2+x(2de-bf^2)+x^2(cf^2+e^2)+d^2)^3} - \frac{3f\sqrt{-a+bx-cx^2}}{(af^2+x(2de-bf^2)+x^2(cf^2+e^2)+d^2)^2} + \frac{1}{af^2+x(2de-bf^2)+x^2(cf^2+e^2)+d^2} \right) dx$$

↓ 7299

$$\int \left(\frac{4f^2(d+ex)(-a+bx-cx^2)}{(af^2+x(2de-bf^2)+x^2(cf^2+e^2)+d^2)^3} - \frac{3f\sqrt{-a+bx-cx^2}}{(af^2+x(2de-bf^2)+x^2(cf^2+e^2)+d^2)^2} + \frac{1}{af^2+x(2de-bf^2)+x^2(cf^2+e^2)+d^2} \right) dx$$

input `Int[(d + e*x + f*Sqrt[-a + b*x - c*x^2])^(-3),x]`

output `$Aborted`

Defintions of rubi rules used

rule 7293 `Int[u_, x_Symbol] :=> With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]`

rule 7299 `Int[u_, x_] :=> CannotIntegrate[u, x]`

Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 255.15 (sec) , antiderivative size = 5961416, normalized size of antiderivative = 4313.62

method	result	size
default	Expression too large to display	5961416

input `int(1/(d+e*x+f*(-c*x^2+b*x-a)^(1/2))^3,x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx - cx^2})^3} dx = \text{Timed out}$$

input `integrate(1/(d+e*x+f*(-c*x^2+b*x-a)^(1/2))^3,x, algorithm="fricas")`

output `Timed out`

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx - cx^2})^3} dx = \text{Timed out}$$

input `integrate(1/(d+e*x+f*(-c*x**2+b*x-a)**(1/2))**3,x)`

output `Timed out`

Maxima [F]

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx - cx^2})^3} dx = \int \frac{1}{(ex + \sqrt{-cx^2 + bx - af + d})^3} dx$$

input `integrate(1/(d+e*x+f*(-c*x^2+b*x-a)^(1/2))^3,x, algorithm="maxima")`

output `integrate((e*x + sqrt(-c*x^2 + b*x - a)*f + d)^(-3), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx - cx^2})^3} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(d+e*x+f*(-c*x^2+b*x-a)^(1/2))^3,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:Error: Bad Argument Type`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx - cx^2})^3} dx = \int \frac{1}{(d + f\sqrt{-cx^2 + bx - a} + ex)^3} dx$$

input `int(1/(d + f*(b*x - a - c*x^2)^(1/2) + e*x)^3,x)`

output `int(1/(d + f*(b*x - a - c*x^2)^(1/2) + e*x)^3, x)`

Reduce [F]

$$\int \frac{1}{(d + ex + f\sqrt{-a + bx - cx^2})^3} dx = \int \frac{1}{(d + ex + f\sqrt{-cx^2 + bx - a})^3} dx$$

input `int(1/(d+e*x+f*(-c*x^2+b*x-a)^(1/2))^3,x)`

output `int(1/(d+e*x+f*(-c*x^2+b*x-a)^(1/2))^3,x)`

CHAPTER 4

APPENDIX

4.1	Listing of Grading functions	407
4.2	Links to plain text integration problems used in this report for each CAS .	425

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```



```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
          finalresult={"C","Result contains complex when optimal does not."}
        ]
      ,(*ELSE*)(*result does not contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
          ]
        ]
      ,(*ELSE*)(*expnResult>expnOptimal*)
        If[FreeQ[result,Integrate] && FreeQ[result,Int],
          finalresult={"C","Result contains higher order function than in optimal. Order "
          ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType, expn]],
  If[Head[expn]===Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]===Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
        1,
        Max[ExpnType[expn[[1]], 2]],
      Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]],
    If[Head[expn]===Plus || Head[expn]===Times,
      Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3, ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
    If[HypergeometricFunctionQ[Head[expn]],

```


Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

#   antiderivative
#   "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

        end if
    else #result contains complex but optimal is not
        if debug then
            print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
    fi;
else # result do not contain complex
    # this assumes optimal do not as well. No check is needed here.
    if debug then
        print("result do not contain complex, this assumes optimal do not as well
    fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
            fi;
        fi;
    else #ExpnType(result) > ExpnType(optimal)
        if debug then
            print("ExpnType(result) > ExpnType(optimal)");
        fi;
        return "C",cat("Result contains higher order function than in optimal. Order ",
                        convert(ExpnType_result,string)," vs. order ",
                        convert(ExpnType_optimal,string),".");
    fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```



```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```



```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file