

# Computer Algebra Independent Integration Tests

Summer 2024

1-Algebraic-functions/1.2-Trinomial/1.2.4-Nested-quadratic-  
trinomial/137-1.2.4.3

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 10:42pm

# Contents

<b>1</b>	<b>Introduction</b>	<b>5</b>
1.1	Listing of CAS systems tested . . . . .	6
1.2	Results . . . . .	7
1.3	Time and leaf size Performance . . . . .	11
1.4	Performance based on number of rules Rubi used . . . . .	13
1.5	Performance based on number of steps Rubi used . . . . .	14
1.6	Solved integrals histogram based on leaf size of result . . . . .	15
1.7	Solved integrals histogram based on CPU time used . . . . .	16
1.8	Leaf size vs. CPU time used . . . . .	17
1.9	list of integrals with no known antiderivative . . . . .	18
1.10	List of integrals solved by CAS but has no known antiderivative . . . . .	18
1.11	list of integrals solved by CAS but failed verification . . . . .	18
1.12	Timing . . . . .	19
1.13	Verification . . . . .	19
1.14	Important notes about some of the results . . . . .	20
1.15	Current tree layout of integration tests . . . . .	23
1.16	Design of the test system . . . . .	24
<b>2</b>	<b>detailed summary tables of results</b>	<b>25</b>
2.1	List of integrals sorted by grade for each CAS . . . . .	26
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	30
2.3	Detailed conclusion table specific for Rubi results . . . . .	44
<b>3</b>	<b>Listing of integrals</b>	<b>46</b>
3.1	$\int \left( d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}} \right)^n dx$ . . . . .	49
3.2	$\int \left( d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}} \right)^3 dx$ . . . . .	55
3.3	$\int \left( d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}} \right)^2 dx$ . . . . .	63
3.4	$\int \left( d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}} \right) dx$ . . . . .	70

3.5	$\int \frac{1}{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}} dx$	76
3.6	$\int \frac{1}{\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)^2} dx$	83
3.7	$\int \frac{1}{\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)^3} dx$	90
3.8	$\int \left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)^{5/2} dx$	97
3.9	$\int \left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)^{3/2} dx$	104
3.10	$\int \sqrt{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}} dx$	111
3.11	$\int \frac{1}{\sqrt{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}}} dx$	118
3.12	$\int \frac{1}{\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)^{3/2}} dx$	125
3.13	$\int \frac{1}{\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)^{5/2}} dx$	132
3.14	$\int \sqrt{x-\sqrt{-4+x^2}} dx$	140
3.15	$\int \sqrt{ax+b\sqrt{c+\frac{a^2x^2}{b^2}}} dx$	145
3.16	$\int (a+x^2)^2 (x+\sqrt{a+x^2})^n dx$	150
3.17	$\int (a+x^2) (x+\sqrt{a+x^2})^n dx$	156
3.18	$\int (x+\sqrt{a+x^2})^n dx$	162
3.19	$\int \frac{(x+\sqrt{a+x^2})^n}{a+x^2} dx$	168
3.20	$\int \frac{(x+\sqrt{a+x^2})^n}{(a+x^2)^2} dx$	173
3.21	$\int (a+x^2)^2 (x-\sqrt{a+x^2})^n dx$	178
3.22	$\int (a+x^2) (x-\sqrt{a+x^2})^n dx$	184
3.23	$\int (x-\sqrt{a+x^2})^n dx$	189
3.24	$\int \frac{(x-\sqrt{a+x^2})^n}{a+x^2} dx$	194
3.25	$\int \frac{(x-\sqrt{a+x^2})^n}{(a+x^2)^2} dx$	199
3.26	$\int (a+x^2)^{5/2} (x+\sqrt{a+x^2})^n dx$	204
3.27	$\int (a+x^2)^{3/2} (x+\sqrt{a+x^2})^n dx$	210
3.28	$\int \sqrt{a+x^2} (x+\sqrt{a+x^2})^n dx$	216
3.29	$\int \frac{(x+\sqrt{a+x^2})^n}{\sqrt{a+x^2}} dx$	221
3.30	$\int \frac{(x+\sqrt{a+x^2})^n}{(a+x^2)^{3/2}} dx$	226

3.31	$\int \frac{(x+\sqrt{a+x^2})^n}{(a+x^2)^{5/2}} dx \dots\dots\dots$	231
3.32	$\int (a+x^2)^{5/2} (x-\sqrt{a+x^2})^n dx \dots\dots\dots$	236
3.33	$\int (a+x^2)^{3/2} (x-\sqrt{a+x^2})^n dx \dots\dots\dots$	242
3.34	$\int \sqrt{a+x^2} (x-\sqrt{a+x^2})^n dx \dots\dots\dots$	248
3.35	$\int \frac{(x-\sqrt{a+x^2})^n}{\sqrt{a+x^2}} dx \dots\dots\dots$	253
3.36	$\int \frac{(x-\sqrt{a+x^2})^n}{(a+x^2)^{3/2}} dx \dots\dots\dots$	258
3.37	$\int \frac{(x-\sqrt{a+x^2})^n}{(a+x^2)^{5/2}} dx \dots\dots\dots$	263
3.38	$\int \left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^2 \left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n dx \dots\dots\dots$	268
3.39	$\int \left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right) \left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n dx \dots\dots\dots$	276
3.40	$\int \left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n dx \dots\dots\dots$	283
3.41	$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}} dx \dots\dots\dots$	289
3.42	$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\left(a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}\right)^2} dx \dots\dots\dots$	295
3.43	$\int \left(d + ex + f\sqrt{\frac{af^2+ex(2d+ex)}{f^2}}\right)^n dx \dots\dots\dots$	301
3.44	$\int \frac{\left(d+ex+f\sqrt{\frac{af^2+ex(2d+ex)}{f^2}}\right)^n}{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}} dx \dots\dots\dots$	307
3.45	$\int \left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^{3/2} \left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n dx \dots\dots\dots$	314
3.46	$\int \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n dx \dots\dots\dots$	322
3.47	$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}} dx \dots\dots\dots$	329
3.48	$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\left(a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}\right)^{3/2}} dx \dots\dots\dots$	335
3.49	$\int \frac{\left(d+ex+f\sqrt{\frac{af^2+ex(2d+ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2+ex(2d+ex)}{f^2}}} dx \dots\dots\dots$	341
3.50	$\int \sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}} \left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n dx \dots\dots\dots$	347
3.51	$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\sqrt{ag+\frac{2degx}{f^2}+\frac{e^2gx^2}{f^2}}} dx \dots\dots\dots$	354

3.52  $\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\left(ag+\frac{2degx}{f^2}+\frac{e^2gx^2}{f^2}\right)^{3/2}} dx \dots\dots\dots 360$

3.53  $\int \frac{\left(d+ex+f\sqrt{\frac{af^2+ex(2d+ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2g+egx(2d+ex)}{f^2}}} dx \dots\dots\dots 367$

**4 Appendix 374**

4.1 Listing of Grading functions 374

4.2 Links to plain text integration problems used in this report for each CAS392

# CHAPTER 1

## INTRODUCTION

1.1	Listing of CAS systems tested . . . . .	6
1.2	Results . . . . .	7
1.3	Time and leaf size Performance . . . . .	11
1.4	Performance based on number of rules Rubi used . . . . .	13
1.5	Performance based on number of steps Rubi used . . . . .	14
1.6	Solved integrals histogram based on leaf size of result . . . . .	15
1.7	Solved integrals histogram based on CPU time used . . . . .	16
1.8	Leaf size vs. CPU time used . . . . .	17
1.9	list of integrals with no known antiderivative . . . . .	18
1.10	List of integrals solved by CAS but has no known antiderivative . . . . .	18
1.11	list of integrals solved by CAS but failed verification . . . . .	18
1.12	Timing . . . . .	19
1.13	Verification . . . . .	19
1.14	Important notes about some of the results . . . . .	20
1.15	Current tree layout of integration tests . . . . .	23
1.16	Design of the test system . . . . .	24

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [ 53 ]. This is test number [ 137 ].

## 1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

## 1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 ( 53 )	0.00 ( 0 )
Mathematica	100.00 ( 53 )	0.00 ( 0 )
Fricas	73.58 ( 39 )	26.42 ( 14 )
Reduce	62.26 ( 33 )	37.74 ( 20 )
Maple	20.75 ( 11 )	79.25 ( 42 )
Giac	16.98 ( 9 )	83.02 ( 44 )
Sympy	13.21 ( 7 )	86.79 ( 46 )
Mupad	11.32 ( 6 )	88.68 ( 47 )
Maxima	5.66 ( 3 )	94.34 ( 50 )

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.



grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> <li>1. antiderivative contains a hypergeometric function and the optimal antiderivative does not.</li> <li>2. antiderivative contains a special function and the optimal antiderivative does not.</li> <li>3. antiderivative contains the imaginary unit and the optimal antiderivative does not.</li> </ol>
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Mathematica	98.113	1.887	0.000	0.000
Fricas	67.925	5.660	0.000	26.415
Giac	15.094	1.887	0.000	83.019
Maple	9.434	7.547	3.774	79.245
Maxima	5.660	0.000	0.000	94.340
Sympy	5.660	7.547	0.000	86.792
Mupad	0.000	11.321	0.000	88.679
Reduce	0.000	62.264	0.000	37.736

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	14	100.00	0.00	0.00
Reduce	20	100.00	0.00	0.00
Maple	42	100.00	0.00	0.00
Giac	44	100.00	0.00	0.00
Sympy	46	69.57	19.57	10.87
Mupad	47	0.00	100.00	0.00
Maxima	50	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

### 1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.04
Fricas	0.12
Reduce	0.17
Maple	0.19
Giac	0.20
Rubi	0.34
Mathematica	1.08
Sympy	3.36
Mupad	22.11

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mupad	76.50	1.29	40.00	0.98
Giac	114.56	1.09	107.00	0.97
Mathematica	118.66	0.97	111.00	0.92
Rubi	119.75	0.98	116.00	0.99
Maxima	151.33	1.16	107.00	1.01
Reduce	179.85	1.13	103.00	0.95
Fricas	199.79	1.29	117.00	1.00
Maple	1463.36	8.71	167.00	1.32
Sympy	2655.43	30.03	311.00	2.62

Table 1.6: Leaf size performance for each CAS

# 1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the  $y$  axis is the percentage solved which Rubi itself needed the number of rules given the  $x$  axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

## 1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

## 1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution



## 1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

## 1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

## 1.9 list of integrals with no known antiderivative

{}

## 1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

## 1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {}

Maple {}

**Maxima** Verification phase not currently implemented.

**Fricas** Verification phase not currently implemented.

**Sympy** Verification phase not currently implemented.

**Giac** Verification phase not currently implemented.

**Reduce** Verification phase not currently implemented.

**Mupad** Verification phase not currently implemented.

## 1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

## 1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

## 1.14 Important notes about some of the results

### Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

## Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

## Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at [https://ask.sagemath.org/question/57123/could-we-have-a-leaf\\_count-function-in-base-sagemath/](https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/)

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

### Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand,the_variable)
```

Which gives  $\sin(x)^2/2$

# 1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree



# 1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



**High level overview of the CAS independent integration test build system**

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

*The following fields are present only in Rubi Table file*

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi  
January 13, 2024  
Design note

# CHAPTER 2

## DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS . . . . .	26
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	30
2.3	Detailed conclusion table specific for Rubi results . . . . .	44

## 2.1 List of integrals sorted by grade for each CAS

Rubi . . . . .	26
Mma . . . . .	26
Maple . . . . .	27
Fricas . . . . .	27
Maxima . . . . .	27
Giac . . . . .	28
Mupad . . . . .	28
Sympy . . . . .	28
Reduce . . . . .	29

### Rubi

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53 }

**B grade** { }

**C grade** { }

**F normal fail** { }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

### Mma

**A grade** { 1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53 }

**B grade** { 5 }

**C grade** { }

**F normal fail** { }

**F(-1) timedout fail** { }

**F(-2) exception fail { }**

## **Maple**

**A grade { 2, 3, 4, 29, 35 }**

**B grade { 5, 6, 7, 18 }**

**C grade { 16, 17 }**

**F normal fail { 1, 8, 9, 10, 11, 12, 13, 14, 15, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 30, 31, 32, 33, 34, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53 }**

**F(-1) timedout fail { }**

**F(-2) exception fail { }**

## **Fricas**

**A grade { 2, 3, 4, 5, 8, 9, 10, 11, 12, 14, 15, 16, 17, 18, 21, 22, 23, 26, 27, 28, 29, 32, 33, 34, 35, 38, 39, 40, 43, 45, 46, 47, 49, 50, 51, 53 }**

**B grade { 6, 7, 13 }**

**C grade { }**

**F normal fail { 1, 19, 20, 24, 25, 30, 31, 36, 37, 41, 42, 44, 48, 52 }**

**F(-1) timedout fail { }**

**F(-2) exception fail { }**

## **Maxima**

**A grade { 2, 3, 4 }**

**B grade { }**

**C grade { }**

**F normal fail { 1, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53 }**

**F(-1) timedout fail { }**

**F(-2) exception fail { }**

## Giac

**A grade** { 2, 3, 4, 6, 7, 14, 29, 35 }

**B grade** { 5 }

**C grade** { }

**F normal fail** { 1, 8, 9, 10, 11, 12, 13, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 30, 31, 32, 33, 34, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## Mupad

**A grade** { }

**B grade** { 3, 4, 29, 35, 47, 49 }

**C grade** { }

**F normal fail** { }

**F(-1) timedout fail** { 1, 2, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 30, 31, 32, 33, 34, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 48, 50, 51, 52, 53 }

**F(-2) exception fail** { }

## Sympy

**A grade** { 2, 3, 4 }

**B grade** { 17, 18, 29, 35 }

**C grade** { }

**F normal fail** { 1, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 19, 20, 21, 22, 23, 24, 25, 27, 28, 30, 31, 33, 34, 36, 37, 39, 40, 47, 50, 51 }

**F(-1) timedout fail** { 16, 38, 41, 42, 43, 44, 49, 52, 53 }

**F(-2) exception fail** { 26, 32, 45, 46, 48 }

## Reduce

**A grade** { }

**B grade** { 2, 3, 4, 5, 6, 7, 14, 15, 16, 17, 18, 21, 22, 23, 26, 27, 28, 29, 32, 33, 34, 35, 38, 39, 40, 43, 45, 46, 47, 49, 50, 51, 53 }

**C grade** { }

**F normal fail** { 1, 8, 9, 10, 11, 12, 13, 19, 20, 24, 25, 30, 31, 36, 37, 41, 42, 44, 48, 52 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## 2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as  $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$ . To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	121	116	86	0	0	0	0	0	23	0
N.S.	1	0.96	0.71	0.00	0.00	0.00	0.00	0.00	0.19	0.00
time (sec)	N/A	0.276	0.597	0.000	0.000	0.000	0.000	0.000	0.166	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	175	168	192	175	293	161	459	171	207	0
N.S.	1	0.96	1.10	1.00	1.67	0.92	2.62	0.98	1.18	0.00
time (sec)	N/A	0.303	1.044	0.059	0.038	0.097	0.652	0.151	0.155	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	131	113	124	107	114	185	107	133	210
N.S.	1	1.24	1.07	1.17	1.01	1.08	1.75	1.01	1.25	1.98
time (sec)	N/A	0.285	0.878	0.970	0.030	0.083	0.710	0.131	0.145	23.856

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	80	75	54	74	54	68	67	136
N.S.	1	1.00	1.18	1.10	0.79	1.09	0.79	1.00	0.99	2.00
time (sec)	N/A	0.205	0.213	0.245	0.042	0.074	1.057	0.126	0.148	22.835

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	<b>F</b>	A	<b>F</b>	B	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	111	337	651	0	187	0	299	132	0
N.S.	1	0.95	2.88	5.56	0.00	1.60	0.00	2.56	1.13	0.00
time (sec)	N/A	0.281	1.231	0.076	0.000	0.124	0.000	0.158	0.154	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F</b>	B	<b>F</b>	A	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	151	146	228	3182	0	284	0	146	365	0
N.S.	1	0.97	1.51	21.07	0.00	1.88	0.00	0.97	2.42	0.00
time (sec)	N/A	0.303	0.966	0.084	0.000	0.132	0.000	0.172	0.157	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F</b>	B	<b>F</b>	A	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	193	182	278	11352	0	536	0	182	767	0
N.S.	1	0.94	1.44	58.82	0.00	2.78	0.00	0.94	3.97	0.00
time (sec)	N/A	0.309	2.852	0.125	0.000	0.273	0.000	0.684	0.160	0.000



Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	225	219	212	0	0	467	0	0	25	0
N.S.	1	0.97	0.94	0.00	0.00	2.08	0.00	0.00	0.11	0.00
time (sec)	N/A	0.425	1.796	0.000	0.000	0.138	0.000	0.000	200.022	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	183	181	170	0	0	388	0	0	25	0
N.S.	1	0.99	0.93	0.00	0.00	2.12	0.00	0.00	0.14	0.00
time (sec)	N/A	0.389	1.559	0.000	0.000	0.127	0.000	0.000	200.030	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	147	147	142	0	0	352	0	0	22	0
N.S.	1	1.00	0.97	0.00	0.00	2.39	0.00	0.00	0.15	0.00
time (sec)	N/A	0.333	1.123	0.000	0.000	0.133	0.000	0.000	0.234	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	147	151	141	0	0	347	0	0	25	0
N.S.	1	1.03	0.96	0.00	0.00	2.36	0.00	0.00	0.17	0.00
time (sec)	N/A	0.292	1.083	0.000	0.000	0.130	0.000	0.000	200.022	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	158	160	169	0	0	538	0	0	25	0
N.S.	1	1.01	1.07	0.00	0.00	3.41	0.00	0.00	0.16	0.00
time (sec)	N/A	0.320	1.687	0.000	0.000	0.227	0.000	0.000	200.021	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	199	196	209	0	0	861	0	0	25	0
N.S.	1	0.98	1.05	0.00	0.00	4.33	0.00	0.00	0.13	0.00
time (sec)	N/A	0.384	1.823	0.000	0.000	0.343	0.000	0.000	200.020	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	A	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	41	45	41	0	0	26	0	25	39	0
N.S.	1	1.10	1.00	0.00	0.00	0.63	0.00	0.61	0.95	0.00
time (sec)	N/A	0.185	0.075	0.000	0.000	0.097	0.000	0.118	0.160	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	69	70	69	0	0	59	0	0	45	0
N.S.	1	1.01	1.00	0.00	0.00	0.86	0.00	0.00	0.65	0.00
time (sec)	N/A	0.225	0.732	0.000	0.000	0.079	0.000	0.000	0.158	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	<b>F</b>	A	<b>F(-1)</b>	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	164	155	138	216	0	158	0	0	219	0
N.S.	1	0.95	0.84	1.32	0.00	0.96	0.00	0.00	1.34	0.00
time (sec)	N/A	0.304	0.690	0.168	0.000	0.112	0.000	0.000	0.174	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	<b>F</b>	A	B	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	108	103	92	167	0	78	15302	0	94	0
N.S.	1	0.95	0.85	1.55	0.00	0.72	141.69	0.00	0.87	0.00
time (sec)	N/A	0.258	0.426	0.039	0.000	0.119	16.375	0.000	0.167	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F</b>	A	B	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	52	51	43	120	0	32	2236	0	30	0
N.S.	1	0.98	0.83	2.31	0.00	0.62	43.00	0.00	0.58	0.00
time (sec)	N/A	0.201	0.235	0.027	0.000	0.102	1.987	0.000	0.159	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	59	59	61	0	0	0	0	0	20	0
N.S.	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	0.34	0.00
time (sec)	N/A	0.229	0.291	0.000	0.000	0.000	0.000	0.000	0.155	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	59	59	61	0	0	0	0	0	28	0
N.S.	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	0.47	0.00
time (sec)	N/A	0.228	0.343	0.000	0.000	0.000	0.000	0.000	0.163	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	176	167	150	0	0	159	0	0	229	0
N.S.	1	0.95	0.85	0.00	0.00	0.90	0.00	0.00	1.30	0.00
time (sec)	N/A	0.297	0.717	0.000	0.000	0.096	0.000	0.000	0.172	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	116	111	100	0	0	79	0	0	103	0
N.S.	1	0.96	0.86	0.00	0.00	0.68	0.00	0.00	0.89	0.00
time (sec)	N/A	0.248	0.384	0.000	0.000	0.093	0.000	0.000	0.166	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	56	55	50	0	0	33	0	0	37	0
N.S.	1	0.98	0.89	0.00	0.00	0.59	0.00	0.00	0.66	0.00
time (sec)	N/A	0.200	0.235	0.000	0.000	0.088	0.000	0.000	0.157	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	65	0	0	0	0	0	22	0
N.S.	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	0.35	0.00
time (sec)	N/A	0.226	0.269	0.000	0.000	0.000	0.000	0.000	0.166	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	65	0	0	0	0	0	30	0
N.S.	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	0.48	0.00
time (sec)	N/A	0.221	0.281	0.000	0.000	0.000	0.000	0.000	0.166	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F(-2)</b>	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	187	176	157	0	0	201	0	0	269	0
N.S.	1	0.94	0.84	0.00	0.00	1.07	0.00	0.00	1.44	0.00
time (sec)	N/A	0.318	0.588	0.000	0.000	0.093	0.000	0.000	0.174	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	131	124	111	0	0	110	0	0	130	0
N.S.	1	0.95	0.85	0.00	0.00	0.84	0.00	0.00	0.99	0.00
time (sec)	N/A	0.285	0.382	0.000	0.000	0.092	0.000	0.000	0.174	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	75	72	65	0	0	48	0	0	47	0
N.S.	1	0.96	0.87	0.00	0.00	0.64	0.00	0.00	0.63	0.00
time (sec)	N/A	0.249	0.313	0.000	0.000	0.108	0.000	0.000	0.166	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	16	0	15	311	15	14	15
N.S.	1	1.00	1.00	0.94	0.00	0.88	18.29	0.88	0.82	0.88
time (sec)	N/A	0.204	0.039	0.128	0.000	0.076	1.753	0.137	0.157	21.016

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	59	59	61	0	0	0	0	0	34	0
N.S.	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	0.58	0.00
time (sec)	N/A	0.233	0.135	0.000	0.000	0.000	0.000	0.000	0.176	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	59	59	61	0	0	0	0	0	48	0
N.S.	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	0.81	0.00
time (sec)	N/A	0.237	0.148	0.000	0.000	0.000	0.000	0.000	0.167	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F(-2)</b>	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	201	190	173	0	0	204	0	0	279	0
N.S.	1	0.95	0.86	0.00	0.00	1.01	0.00	0.00	1.39	0.00
time (sec)	N/A	0.319	0.733	0.000	0.000	0.127	0.000	0.000	0.180	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	141	134	123	0	0	113	0	0	140	0
N.S.	1	0.95	0.87	0.00	0.00	0.80	0.00	0.00	0.99	0.00
time (sec)	N/A	0.291	0.377	0.000	0.000	0.157	0.000	0.000	0.172	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	81	78	73	0	0	51	0	0	57	0
N.S.	1	0.96	0.90	0.00	0.00	0.63	0.00	0.00	0.70	0.00
time (sec)	N/A	0.257	0.310	0.000	0.000	0.106	0.000	0.000	0.176	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	19	0	18	41	18	23	18
N.S.	1	1.00	1.00	0.95	0.00	0.90	2.05	0.90	1.15	0.90
time (sec)	N/A	0.212	0.043	0.171	0.000	0.084	0.952	0.113	0.167	20.931

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	65	0	0	0	0	0	36	0
N.S.	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	0.57	0.00
time (sec)	N/A	0.236	0.143	0.000	0.000	0.000	0.000	0.000	0.158	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	65	0	0	0	0	0	50	0
N.S.	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	0.79	0.00
time (sec)	N/A	0.232	0.160	0.000	0.000	0.000	0.000	0.000	0.167	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F(-1)</b>	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	365	327	280	0	0	654	0	0	1095	0
N.S.	1	0.90	0.77	0.00	0.00	1.79	0.00	0.00	3.00	0.00
time (sec)	N/A	0.539	6.286	0.000	0.000	0.168	0.000	0.000	0.284	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	239	217	186	0	0	239	0	0	324	0
N.S.	1	0.91	0.78	0.00	0.00	1.00	0.00	0.00	1.36	0.00
time (sec)	N/A	0.392	2.433	0.000	0.000	0.129	0.000	0.000	0.184	0.000



Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	107	104	89	0	0	80	0	0	66	0
N.S.	1	0.97	0.83	0.00	0.00	0.75	0.00	0.00	0.62	0.00
time (sec)	N/A	0.274	1.603	0.000	0.000	0.120	0.000	0.000	0.159	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	122	122	112	0	0	0	0	0	53	0
N.S.	1	1.00	0.92	0.00	0.00	0.00	0.00	0.00	0.43	0.00
time (sec)	N/A	0.407	1.964	0.000	0.000	0.000	0.000	0.000	0.166	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	122	122	112	0	0	0	0	0	91	0
N.S.	1	1.00	0.92	0.00	0.00	0.00	0.00	0.00	0.75	0.00
time (sec)	N/A	0.382	2.172	0.000	0.000	0.000	0.000	0.000	0.208	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F(-1)</b>	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	107	104	89	0	0	80	0	0	66	0
N.S.	1	0.97	0.83	0.00	0.00	0.75	0.00	0.00	0.62	0.00
time (sec)	N/A	0.332	0.015	0.000	0.000	0.139	0.000	0.000	0.164	0.000

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	122	122	112	0	0	0	0	0	53	0
N.S.	1	1.00	0.92	0.00	0.00	0.00	0.00	0.00	0.43	0.00
time (sec)	N/A	0.560	0.013	0.000	0.000	0.000	0.000	0.000	0.168	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F(-2)</b>	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	297	266	228	0	0	377	0	0	554	0
N.S.	1	0.90	0.77	0.00	0.00	1.27	0.00	0.00	1.87	0.00
time (sec)	N/A	0.503	8.814	0.000	0.000	0.134	0.000	0.000	0.200	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F(-2)</b>	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	171	156	135	0	0	122	0	0	128	0
N.S.	1	0.91	0.79	0.00	0.00	0.71	0.00	0.00	0.75	0.00
time (sec)	N/A	0.462	3.616	0.000	0.000	0.109	0.000	0.000	0.169	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	B	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	36	0	0	41	0	0	34	41
N.S.	1	1.00	0.88	0.00	0.00	1.00	0.00	0.00	0.83	1.00
time (sec)	N/A	0.369	0.492	0.000	0.000	0.097	0.000	0.000	0.158	22.056

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	122	122	112	0	0	0	0	0	110	0
N.S.	1	1.00	0.92	0.00	0.00	0.00	0.00	0.00	0.90	0.00
time (sec)	N/A	0.410	4.352	0.000	0.000	0.000	0.000	0.000	0.177	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F(-1)</b>	<b>F</b>	B	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	36	0	0	41	0	0	34	39
N.S.	1	1.00	0.88	0.00	0.00	1.00	0.00	0.00	0.83	0.95
time (sec)	N/A	0.543	0.013	0.000	0.000	0.099	0.000	0.000	0.172	21.989

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	327	208	175	0	0	231	0	0	130	0
N.S.	1	0.64	0.54	0.00	0.00	0.71	0.00	0.00	0.40	0.00
time (sec)	N/A	0.725	0.271	0.000	0.000	0.132	0.000	0.000	0.181	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	93	93	76	0	0	117	0	0	39	0
N.S.	1	1.00	0.82	0.00	0.00	1.26	0.00	0.00	0.42	0.00
time (sec)	N/A	0.646	0.056	0.000	0.000	0.105	0.000	0.000	0.164	0.000

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	177	177	152	0	0	0	0	0	117	0
N.S.	1	1.00	0.86	0.00	0.00	0.00	0.00	0.00	0.66	0.00
time (sec)	N/A	0.701	0.219	0.000	0.000	0.000	0.000	0.000	0.189	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F(-1)	F	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	93	93	76	0	0	117	0	0	39	0
N.S.	1	1.00	0.82	0.00	0.00	1.26	0.00	0.00	0.42	0.00
time (sec)	N/A	0.839	0.029	0.000	0.000	0.079	0.000	0.000	0.159	0.000

## 2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio  $\frac{\text{number of rules}}{\text{integrand size}}$  is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [14] had the largest ratio of [.235294000000000003]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	4	3	0.96	25	0.120
2	A	4	3	0.96	25	0.120
3	A	4	3	1.24	25	0.120
4	A	1	1	1.00	23	0.043
5	A	4	3	0.95	25	0.120
6	A	4	3	0.97	25	0.120
7	A	4	3	0.94	25	0.120
8	A	7	6	0.97	27	0.222
9	A	6	5	0.99	27	0.185
10	A	7	6	1.00	27	0.222
11	A	7	6	1.03	27	0.222
12	A	7	6	1.01	27	0.222
13	A	6	5	0.98	27	0.185
14	A	5	4	1.10	17	0.235
15	A	4	3	1.01	26	0.115
16	A	4	3	0.95	21	0.143
17	A	4	3	0.95	19	0.158
18	A	4	3	0.98	13	0.231
19	A	3	2	1.00	21	0.095
20	A	3	2	1.00	21	0.095
21	A	4	3	0.95	23	0.130

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	4	3	0.96	21	0.143
23	A	4	3	0.98	15	0.200
24	A	3	2	1.00	23	0.087
25	A	3	2	1.00	23	0.087
26	A	4	3	0.94	23	0.130
27	A	4	3	0.95	23	0.130
28	A	4	3	0.96	23	0.130
29	A	3	2	1.00	23	0.087
30	A	3	2	1.00	23	0.087
31	A	3	2	1.00	23	0.087
32	A	4	3	0.95	25	0.120
33	A	4	3	0.95	25	0.120
34	A	4	3	0.96	25	0.120
35	A	3	2	1.00	25	0.080
36	A	3	2	1.00	25	0.080
37	A	3	2	1.00	25	0.080
38	A	5	4	0.90	56	0.071
39	A	5	4	0.91	54	0.074
40	A	5	4	0.97	33	0.121
41	A	5	4	1.00	56	0.071
42	A	4	3	1.00	56	0.054
43	A	6	5	0.97	33	0.152
44	A	6	5	1.00	56	0.089
45	A	5	4	0.90	58	0.069
46	A	5	4	0.91	58	0.069
47	A	3	2	1.00	58	0.034
48	A	4	3	1.00	58	0.052
49	A	4	3	1.00	58	0.052
50	A	6	5	0.64	62	0.081
51	A	4	3	1.00	62	0.048
52	A	5	4	1.00	62	0.065
53	A	5	4	1.00	60	0.067

# CHAPTER 3

## LISTING OF INTEGRALS

3.1	$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^n dx \dots\dots\dots$	49
3.2	$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^3 dx \dots\dots\dots$	55
3.3	$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^2 dx \dots\dots\dots$	63
3.4	$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right) dx \dots\dots\dots$	70
3.5	$\int \frac{1}{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}} dx \dots\dots\dots$	76
3.6	$\int \frac{1}{\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)^2} dx \dots\dots\dots$	83
3.7	$\int \frac{1}{\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)^3} dx \dots\dots\dots$	90
3.8	$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx \dots\dots\dots$	97
3.9	$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx \dots\dots\dots$	104
3.10	$\int \sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}} dx \dots\dots\dots$	111
3.11	$\int \frac{1}{\sqrt{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}}} dx \dots\dots\dots$	118
3.12	$\int \frac{1}{\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)^{3/2}} dx \dots\dots\dots$	125
3.13	$\int \frac{1}{\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)^{5/2}} dx \dots\dots\dots$	132
3.14	$\int \sqrt{x - \sqrt{-4 + x^2}} dx \dots\dots\dots$	140
3.15	$\int \sqrt{ax + b} \sqrt{c + \frac{a^2 x^2}{b^2}} dx \dots\dots\dots$	145
3.16	$\int (a + x^2)^2 (x + \sqrt{a + x^2})^n dx \dots\dots\dots$	150

3.17	$\int (a+x^2)(x+\sqrt{a+x^2})^n dx$	156
3.18	$\int (x+\sqrt{a+x^2})^n dx$	162
3.19	$\int \frac{(x+\sqrt{a+x^2})^n}{a+x^2} dx$	168
3.20	$\int \frac{(x+\sqrt{a+x^2})^n}{(a+x^2)^2} dx$	173
3.21	$\int (a+x^2)^2(x-\sqrt{a+x^2})^n dx$	178
3.22	$\int (a+x^2)(x-\sqrt{a+x^2})^n dx$	184
3.23	$\int (x-\sqrt{a+x^2})^n dx$	189
3.24	$\int \frac{(x-\sqrt{a+x^2})^n}{a+x^2} dx$	194
3.25	$\int \frac{(x-\sqrt{a+x^2})^n}{(a+x^2)^2} dx$	199
3.26	$\int (a+x^2)^{5/2}(x+\sqrt{a+x^2})^n dx$	204
3.27	$\int (a+x^2)^{3/2}(x+\sqrt{a+x^2})^n dx$	210
3.28	$\int \sqrt{a+x^2}(x+\sqrt{a+x^2})^n dx$	216
3.29	$\int \frac{(x+\sqrt{a+x^2})^n}{\sqrt{a+x^2}} dx$	221
3.30	$\int \frac{(x+\sqrt{a+x^2})^n}{(a+x^2)^{3/2}} dx$	226
3.31	$\int \frac{(x+\sqrt{a+x^2})^n}{(a+x^2)^{5/2}} dx$	231
3.32	$\int (a+x^2)^{5/2}(x-\sqrt{a+x^2})^n dx$	236
3.33	$\int (a+x^2)^{3/2}(x-\sqrt{a+x^2})^n dx$	242
3.34	$\int \sqrt{a+x^2}(x-\sqrt{a+x^2})^n dx$	248
3.35	$\int \frac{(x-\sqrt{a+x^2})^n}{\sqrt{a+x^2}} dx$	253
3.36	$\int \frac{(x-\sqrt{a+x^2})^n}{(a+x^2)^{3/2}} dx$	258
3.37	$\int \frac{(x-\sqrt{a+x^2})^n}{(a+x^2)^{5/2}} dx$	263
3.38	$\int \left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^2 \left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n dx$	268
3.39	$\int \left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right) \left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n dx$	276
3.40	$\int \left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n dx$	283
3.41	$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}} dx$	289
3.42	$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\left(a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}\right)^2} dx$	295
3.43	$\int \left(d + ex + f\sqrt{\frac{af^2+ex(2d+ex)}{f^2}}\right)^n dx$	301



3.44	$\int \frac{\left(d+ex+f\sqrt{\frac{af^2+ex(2d+ex)}{f^2}}\right)^n}{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}} dx$	307
3.45	$\int \left(a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}\right)^{3/2} \left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n dx$	314
3.46	$\int \sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}} \left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n dx$	322
3.47	$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}} dx$	329
3.48	$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\left(a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}\right)^{3/2}} dx$	335
3.49	$\int \frac{\left(d+ex+f\sqrt{\frac{af^2+ex(2d+ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2+ex(2d+ex)}{f^2}}} dx$	341
3.50	$\int \sqrt{ag+\frac{2degx}{f^2}+\frac{e^2gx^2}{f^2}} \left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n dx$	347
3.51	$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\sqrt{ag+\frac{2degx}{f^2}+\frac{e^2gx^2}{f^2}}} dx$	354
3.52	$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\left(ag+\frac{2degx}{f^2}+\frac{e^2gx^2}{f^2}\right)^{3/2}} dx$	360
3.53	$\int \frac{\left(d+ex+f\sqrt{\frac{af^2+ex(2d+ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2g+egx(2d+ex)}{f^2}}} dx$	367

$$3.1 \quad \int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^n dx$$

Optimal result	49
Mathematica [A] (verified)	50
Rubi [A] (verified)	50
Maple [F]	52
Fricas [F]	52
Sympy [F]	52
Maxima [F]	53
Giac [F]	53
Mupad [F(-1)]	53
Reduce [F]	54

### Optimal result

Integrand size = 25, antiderivative size = 121

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^n dx = \frac{\left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{1+n}}{2e(1+n)} + \frac{af^2 \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{1+n} \operatorname{Hypergeometric2F1} \left( 2, 1+n, 2+n, \frac{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}}{d} \right)}{2d^2e(1+n)}$$

output

```
1/2*(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1+n)/e/(1+n)+1/2*a*f^2*(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1+n)*hypergeom([2, 1+n], [2+n], (d+e*x+f*(a+e^2*x^2/f^2)^(1/2))/d)/d^2/e/(1+n)
```

**Mathematica [A] (verified)**

Time = 0.60 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.71

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^n dx$$

$$= \frac{\left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{1+n} \left( d^2 + af^2 \operatorname{Hypergeometric2F1} \left( 2, 1 + n, 2 + n, \frac{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}}{d} \right) \right)}{2d^2 e(1 + n)}$$

input

```
Integrate[(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^n,x]
```

output

```
((d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(1 + n)*(d^2 + a*f^2*Hypergeometric
2F1[2, 1 + n, 2 + n, (d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])/d]))/(2*d^2*e*(
1 + n))
```

**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.96, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$ , Rules used = {2542, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( f \sqrt{a + \frac{e^2 x^2}{f^2}} + d + ex \right)^n dx$$

$$\downarrow \text{2542}$$

$$\int \frac{\left( d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + a} \right)^n \left( d^2 - 2 \left( d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + a} \right) d + af^2 + \left( d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + a} \right)^2 \right)}{\left( -\sqrt{\frac{e^2 x^2}{f^2} + a} f - ex \right)^2} dx$$

$$\downarrow \text{1195}$$

$$\frac{\int \left( \frac{af^2 \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right)^n}{\left( -\sqrt{\frac{e^2x^2}{f^2}+a}f-ex \right)^2} + \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right)^n \right) d \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right)}{2e} \xrightarrow{2009} \frac{af^2 \left( f\sqrt{a+\frac{e^2x^2}{f^2}}+d+ex \right)^{n+1} \operatorname{Hypergeometric2F1} \left( 2, n+1, n+2, \frac{d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}}{d} \right)}{d^2(n+1)} + \frac{\left( f\sqrt{a+\frac{e^2x^2}{f^2}}+d+ex \right)^{n+1}}{n+1}$$

input `Int[(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^n,x]`

output `((d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(1 + n)/(1 + n) + (a*f^2*(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(1 + n)*Hypergeometric2F1[2, 1 + n, 2 + n, (d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])/d])/(d^2*(1 + n)))/(2*e)`

### Defintions of rubi rules used

rule 1195 `Int[((d_.) + (e_.)*(x_.))^(m_.)*((f_.) + (g_.)*(x_.))^(n_.)*((a_.) + (b_.)*(x_.) + (c_.)*(x_.^2))^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2542 `Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2]))^(n_.))^(p_.), x_Symbol] := Simp[1/(2*e) Subst[Int[(g + h*x^n)^p*((d^2 + a*f^2 - 2*d*x + x^2)/(d - x)^2), x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]`

**Maple [F]**

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^n dx$$

input `int((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^n,x)`

output `int((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^n,x)`

**Fricas [F]**

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^n dx = \int \left( ex + \sqrt{\frac{e^2 x^2}{f^2} + af + d} \right)^n dx$$

input `integrate((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^n,x, algorithm="fricas")`

output `integral((e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d)^n, x)`

**Sympy [F]**

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^n dx = \int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^n dx$$

input `integrate((d+e*x+f*(a+e**2*x**2/f**2)**(1/2))**n,x)`

output `Integral((d + e*x + f*sqrt(a + e**2*x**2/f**2))**n, x)`

**Maxima [F]**

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^n dx = \int \left( ex + \sqrt{\frac{e^2 x^2}{f^2} + af + d} \right)^n dx$$

input `integrate((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^n,x, algorithm="maxima")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a)*f + d)^n, x)`

**Giac [F]**

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^n dx = \int \left( ex + \sqrt{\frac{e^2 x^2}{f^2} + af + d} \right)^n dx$$

input `integrate((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^n,x, algorithm="giac")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a)*f + d)^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^n dx = \int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^n dx$$

input `int((d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^n,x)`

output `int((d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^n, x)`

**Reduce [F]**

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^n dx = \int \left( \sqrt{e^2 x^2 + a f^2} + d + ex \right)^n dx$$

input `int((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^n,x)`

output `int((sqrt(a*f**2 + e**2*x**2) + d + e*x)**n,x)`

$$3.2 \quad \int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^3 dx$$

Optimal result . . . . .	55
Mathematica [A] (verified) . . . . .	56
Rubi [A] (verified) . . . . .	56
Maple [A] (verified) . . . . .	58
Fricas [A] (verification not implemented) . . . . .	58
Sympy [A] (verification not implemented) . . . . .	59
Maxima [A] (verification not implemented) . . . . .	60
Giac [A] (verification not implemented) . . . . .	61
Mupad [F(-1)] . . . . .	61
Reduce [B] (verification not implemented) . . . . .	62

**Optimal result**

Integrand size = 25, antiderivative size = 175

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^3 dx = -\frac{ad^3 f^2}{2e \left( ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)} + \frac{adf^2 \left( ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)}{e}$$

$$+ \frac{af^2 \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^2}{4e}$$

$$+ \frac{\left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^4}{8e}$$

$$+ \frac{3ad^2 f^2 \log \left( ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)}{2e}$$

output

```
-1/2*a*d^3*f^2/e/(e*x+f*(a+e^2*x^2/f^2)^(1/2))+a*d*f^2*(e*x+f*(a+e^2*x^2/f^2)^(1/2))/e+1/4*a*f^2*(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^2/e+1/8*(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^4/e+3/2*a*d^2*f^2*ln(e*x+f*(a+e^2*x^2/f^2)^(1/2))/e
```



**Mathematica [A] (verified)**

Time = 1.04 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.10

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^3 dx$$

$$= \frac{ex(2d^3 + 6adf^2 + 3d^2ex + 3aef^2x + 4de^2x^2 + 2e^3x^3) + \sqrt{a + \frac{e^2x^2}{f^2}}(2af^3(2d + ex) + efx(3d^2 + 4dex + 2e^2x^2))}{2e}$$

input

```
Integrate[(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^3,x]
```

output

```
(e*x*(2*d^3 + 6*a*d*f^2 + 3*d^2*e*x + 3*a*e*f^2*x + 4*d*e^2*x^2 + 2*e^3*x^3) + Sqrt[a + (e^2*x^2)/f^2]*(2*a*f^3*(2*d + e*x) + e*f*x*(3*d^2 + 4*d*e*x + 2*e^2*x^2)) - 3*a*d^2*f^2*Log[e*(Sqrt[a]*f + e*x - f*Sqrt[a + (e^2*x^2)/f^2]]) + 3*a*d^2*f^2*Log[-(Sqrt[a]*f) + e*x + f*Sqrt[a + (e^2*x^2)/f^2]])/(2*e)
```

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.96, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$ , Rules used = {2542, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( f \sqrt{a + \frac{e^2 x^2}{f^2}} + d + ex \right)^3 dx$$

↓ 2542

$$\int \frac{\left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right)^3 \left( d^2-2\left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right) d+af^2+\left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right)^2 \right)}{\left( -\sqrt{\frac{e^2x^2}{f^2}+a}f-ex \right)^2} d \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right)}{2e}$$

↓ 1195

$$\int \left( \frac{af^2d^3}{\left(-\sqrt{\frac{e^2x^2}{f^2}+af-ex}\right)^2} - \frac{3af^2d^2}{-\sqrt{\frac{e^2x^2}{f^2}+af-ex}} + 2af^2d + \left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^3 + af^2\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right) \right) dx$$

2e

↓ 2009

$$\frac{ad^3f^2}{f\left(-\sqrt{a+\frac{e^2x^2}{f^2}}\right)-ex} + 3ad^2f^2 \log\left(f\left(-\sqrt{a+\frac{e^2x^2}{f^2}}\right)-ex\right) + \frac{1}{4}\left(f\sqrt{a+\frac{e^2x^2}{f^2}}+d+ex\right)^4 + \frac{1}{2}af^2\left(f\sqrt{a+\frac{e^2x^2}{f^2}}+d+ex\right)$$

2e

input `Int[(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^3,x]`

output `((a*d^3*f^2)/(-(e*x) - f*Sqrt[a + (e^2*x^2)/f^2]) + 2*a*d*f^2*(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]) + (a*f^2*(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^2)/2 + (d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^4/4 + 3*a*d^2*f^2*Log[-(e*x) - f*Sqrt[a + (e^2*x^2)/f^2]])/(2*e)`

### Defintions of rubi rules used

rule 1195 `Int[((d_.) + (e_.)*(x_))^(m_.)*((f_.) + (g_.)*(x_))^(n_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2542 `Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2]))^(n_.)^(p_.), x_Symbol] := Simp[1/(2*e) Subst[Int[(g + h*x^n)^p*((d^2 + a*f^2 - 2*d*x + x^2)/(d - x)^2), x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]`

**Maple [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.00

$$f^3 x \left( a + \frac{e^2 x^2}{f^2} \right)^{\frac{3}{2}} + e^3 x^4 + 2d e^2 x^3 + \frac{3f^2 x^2 a e}{2} + 3f^2 x a d + \frac{3f d^2 x \sqrt{a + \frac{e^2 x^2}{f^2}}}{2} + \frac{3f d^2 a \ln \left( \frac{e^2 x}{f^2 \sqrt{\frac{e^2}{f^2}}} + \sqrt{a + \frac{e^2 x^2}{f^2}} \right)}{2 \sqrt{\frac{e^2}{f^2}}}$$

input `int((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^3,x)`output `f^3*x*(a+e^2*x^2/f^2)^(3/2)+e^3*x^4+2*d*e^2*x^3+3/2*f^2*x^2*a*e+3*f^2*x*a*d+3/2*f*d^2*x*(a+e^2*x^2/f^2)^(1/2)+3/2*f*d^2*a*ln(e^2*x/f^2/(e^2/f^2)^(1/2)+(a+e^2*x^2/f^2)^(1/2))/(e^2/f^2)^(1/2)+2*d/e*f^3*((e^2*x^2+a*f^2)/f^2)^(3/2)+3/2*e*x^2*d^2+x*d^3+1/4*d^4/e`**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 161, normalized size of antiderivative = 0.92

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^3 dx$$

$$= \frac{2e^4 x^4 + 4de^3 x^3 - 3ad^2 f^2 \log \left( -ex + f \sqrt{\frac{e^2 x^2 + af^2}{f^2}} \right) + 3(ae^2 f^2 + d^2 e^2) x^2 + 2(3adef^2 + d^3 e) x + (2e^3 f^2 + 3d^2 e^2) x^2}{2e}$$

input `integrate((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^3,x, algorithm="fricas")`output `1/2*(2*e^4*x^4 + 4*d*e^3*x^3 - 3*a*d^2*f^2*log(-e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2)) + 3*(a*e^2*f^2 + d^2*e^2)*x^2 + 2*(3*a*d*e*f^2 + d^3*e)*x + (2*e^3*f^2 + 3*d^2*e^2)*x^2 + 4*d*e^2*f*x^2 + 4*a*d*f^3 + (2*a*e*f^3 + 3*d^2*e*f)*x)*sqrt((e^2*x^2 + a*f^2)/f^2)/e`

**Sympy [A] (verification not implemented)**

Time = 0.65 (sec) , antiderivative size = 459, normalized size of antiderivative = 2.62

$$\begin{aligned}
& \int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^3 dx = 3adf^2x + \frac{3ae f^2 x^2}{2} \\
& + af^3 \left( \left( \frac{\log \left( \frac{2e^2 x + 2\sqrt{\frac{e^2}{f^2}} \sqrt{a + \frac{e^2 x^2}{f^2}}}{\sqrt{\frac{e^2}{f^2}}} \right)}{\sqrt{\frac{e^2 x^2}{f^2}}} \right) \text{ for } a \neq 0 \right. \\
& \left. \frac{x \log(x)}{\sqrt{\frac{e^2 x^2}{f^2}}} \text{ otherwise} \right) + \frac{x \sqrt{a + \frac{e^2 x^2}{f^2}}}{2} \text{ for } \frac{e^2}{f^2} \neq 0 \\
& \left. \frac{\sqrt{ax}}{2} \text{ otherwise} \right) \\
& + d^3x + \frac{3d^2ex^2}{2} \\
& + 3d^2f \left( \left( \frac{\log \left( \frac{2e^2 x + 2\sqrt{\frac{e^2}{f^2}} \sqrt{a + \frac{e^2 x^2}{f^2}}}{\sqrt{\frac{e^2}{f^2}}} \right)}{\sqrt{\frac{e^2 x^2}{f^2}}} \right) \text{ for } a \neq 0 \right. \\
& \left. \frac{x \log(x)}{\sqrt{\frac{e^2 x^2}{f^2}}} \text{ otherwise} \right) + \frac{x \sqrt{a + \frac{e^2 x^2}{f^2}}}{2} \text{ for } \frac{e^2}{f^2} \neq 0 \\
& \left. \frac{\sqrt{ax}}{2} \text{ otherwise} \right) \\
& + 2de^2x^3 + 6def \left( \left( \frac{\sqrt{a + \frac{e^2 x^2}{f^2}} \left( \frac{af^2}{3e^2} + \frac{x^2}{3} \right)}{\sqrt{\frac{e^2 x^2}{f^2}}} \right) \text{ for } \frac{e^2}{f^2} \neq 0 \right. \\
& \left. \frac{\sqrt{ax^2}}{2} \text{ otherwise} \right) + e^3x^4 \\
& + 4e^2f \left( \left( \frac{\log \left( \frac{2e^2 x + 2\sqrt{\frac{e^2}{f^2}} \sqrt{a + \frac{e^2 x^2}{f^2}}}{\sqrt{\frac{e^2}{f^2}}} \right)}{\sqrt{\frac{e^2 x^2}{f^2}}} \right) \text{ for } a \neq 0 \right. \\
& \left. \frac{x \log(x)}{\sqrt{\frac{e^2 x^2}{f^2}}} \text{ otherwise} \right) + \frac{\sqrt{a + \frac{e^2 x^2}{f^2}} \left( \frac{af^2 x}{8e^2} + \frac{x^3}{4} \right)}{8e^2} \text{ for } \frac{e^2}{f^2} \neq 0 \\
& \left. \frac{\sqrt{ax^3}}{3} \text{ otherwise} \right)
\end{aligned}$$

input

```
integrate((d+e*x+f*(a+e**2*x**2/f**2)**(1/2))**3,x)
```

output

```

3*a*d*f**2*x + 3*a*e*f**2*x**2/2 + a*f**3*Piecewise((a*Piecewise((log(2*e*
**2*x/f**2 + 2*sqrt(e**2/f**2)*sqrt(a + e**2*x**2/f**2))/sqrt(e**2/f**2), N
e(a, 0)), (x*log(x)/sqrt(e**2*x**2/f**2), True))/2 + x*sqrt(a + e**2*x**2/
f**2)/2, Ne(e**2/f**2, 0)), (sqrt(a)*x, True)) + d**3*x + 3*d**2*e*x**2/2
+ 3*d**2*f*Piecewise((a*Piecewise((log(2*e**2*x/f**2 + 2*sqrt(e**2/f**2)*s
qrt(a + e**2*x**2/f**2))/sqrt(e**2/f**2), Ne(a, 0)), (x*log(x)/sqrt(e**2*x
**2/f**2), True))/2 + x*sqrt(a + e**2*x**2/f**2)/2, Ne(e**2/f**2, 0)), (sq
rt(a)*x, True)) + 2*d*e**2*x**3 + 6*d*e*f*Piecewise((sqrt(a + e**2*x**2/f*
**2)*(a*f**2/(3*e**2) + x**2/3), Ne(e**2/f**2, 0)), (sqrt(a)*x**2/2, True))
+ e**3*x**4 + 4*e**2*f*Piecewise((-a**2*f**2*Piecewise((log(2*e**2*x/f**2
+ 2*sqrt(e**2/f**2)*sqrt(a + e**2*x**2/f**2))/sqrt(e**2/f**2), Ne(a, 0)),
(x*log(x)/sqrt(e**2*x**2/f**2), True))/(8*e**2) + sqrt(a + e**2*x**2/f**2
)*(a*f**2*x/(8*e**2) + x**3/4), Ne(e**2/f**2, 0)), (sqrt(a)*x**3/3, True))

```

### Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 293, normalized size of antiderivative = 1.67

$$\begin{aligned}
& \int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^3 dx \\
&= \frac{1}{4} e^3 x^4 + \frac{3 \left( \frac{e^2 x^2}{f^2} + a \right)^2 f^4}{4e} \\
&\quad - \frac{3}{8} \left( \frac{a^2 f^3 \operatorname{arsinh} \left( \frac{e^2 x}{\sqrt{ae^2} f} \right)}{\sqrt{e^2} e^2} - \frac{2 \left( \frac{e^2 x^2}{f^2} + a \right)^{\frac{3}{2}} f^2 x}{e^2} + \frac{\sqrt{\frac{e^2 x^2}{f^2} + a} a f^2 x}{e^2} \right) e^2 f \\
&\quad + \frac{1}{8} \left( \frac{3 a^2 f \operatorname{arsinh} \left( \frac{e^2 x}{\sqrt{ae^2} f} \right)}{\sqrt{e^2}} + 2 \left( \frac{e^2 x^2}{f^2} + a \right)^{\frac{3}{2}} x + 3 \sqrt{\frac{e^2 x^2}{f^2} + a} a x \right) f^3 \\
&\quad + d^3 x + \frac{3}{2} \left( ex^2 + \left( \frac{af \operatorname{arsinh} \left( \frac{e^2 x}{\sqrt{ae^2} f} \right)}{\sqrt{e^2}} + \sqrt{\frac{e^2 x^2}{f^2} + ax} \right) f \right) d^2 \\
&\quad + \left( e^2 x^3 + \frac{2 \left( \frac{e^2 x^2}{f^2} + a \right)^{\frac{3}{2}} f^3}{e} + \left( \frac{e^2 x^3}{f^2} + 3ax \right) f^2 \right) d
\end{aligned}$$

input

```

integrate((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^3,x, algorithm="maxima")

```

output

$$\begin{aligned} & 1/4*e^3*x^4 + 3/4*(e^2*x^2/f^2 + a)^2*f^4/e - 3/8*(a^2*f^3*arcsinh(e^2*x/( \\ & \text{sqrt}(a*e^2)*f)))/(sqrt(e^2)*e^2) - 2*(e^2*x^2/f^2 + a)^{(3/2)}*f^2*x/e^2 + \text{sq} \\ & \text{rt}(e^2*x^2/f^2 + a)*a*f^2*x/e^2)*e^2*f + 1/8*(3*a^2*f*arcsinh(e^2*x/(sqrt( \\ & a*e^2)*f)))/sqrt(e^2) + 2*(e^2*x^2/f^2 + a)^{(3/2)}*x + 3*sqrt(e^2*x^2/f^2 + \\ & a)*a*x)*f^3 + d^3*x + 3/2*(e*x^2 + (a*f*arcsinh(e^2*x/(sqrt(a*e^2)*f)))/\text{sq} \\ & \text{rt}(e^2) + sqrt(e^2*x^2/f^2 + a)*x)*f)*d^2 + (e^2*x^3 + 2*(e^2*x^2/f^2 + a)^ \\ & (3/2)*f^3/e + (e^2*x^3/f^2 + 3*a*x)*f^2)*d \end{aligned}$$
**Giac [A] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.98

$$\begin{aligned} & \int \left( d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}} \right)^3 dx = e^3x^4 + \frac{3}{2}aef^2x^2 + 2de^2x^3 \\ & + 3adf^2x + \frac{3}{2}d^2ex^2 - \frac{3ad^2f|f|\log(|-x|e| + \sqrt{e^2x^2 + af^2})}{2|e|} + d^3x \\ & + \frac{1}{2}\sqrt{e^2x^2 + af^2} \left( \frac{4adf|f|}{e} + \left( 2 \left( \frac{e^2x|f|}{f} + \frac{2de|f|}{f} \right) x + \frac{2ae^4f^4|f| + 3d^2e^4f^2|f|}{e^4f^3} \right) x \right) \end{aligned}$$

input

```
integrate((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^3,x, algorithm="giac")
```

output

$$\begin{aligned} & e^3*x^4 + 3/2*a*e*f^2*x^2 + 2*d*e^2*x^3 + 3*a*d*f^2*x + 3/2*d^2*e*x^2 - 3/ \\ & 2*a*d^2*f*abs(f)*log(abs(-x*abs(e) + sqrt(e^2*x^2 + a*f^2)))/abs(e) + d^3*x \\ & x + 1/2*sqrt(e^2*x^2 + a*f^2)*(4*a*d*f*abs(f)/e + (2*(e^2*x*abs(f)/f + 2*d \\ & *e*abs(f)/f)*x + (2*a*e^4*f^4*abs(f) + 3*d^2*e^4*f^2*abs(f))/(e^4*f^3))*x \end{aligned}$$
**Mupad [F(-1)]**

Timed out.

$$\int \left( d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}} \right)^3 dx = \int \left( d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}} \right)^3 dx$$

input

```
int((d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^3,x)
```

output `int((d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^3, x)`

### Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.18

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^3 dx$$

$$= \frac{4\sqrt{e^2 x^2 + a f^2} a d f^2 + 2\sqrt{e^2 x^2 + a f^2} a e f^2 x + 3\sqrt{e^2 x^2 + a f^2} d^2 e x + 4\sqrt{e^2 x^2 + a f^2} d e^2 x^2 + 2\sqrt{e^2 x^2 + a f^2} d^2 e x^2 + 2\sqrt{e^2 x^2 + a f^2} d e^2 x^3 + 2\sqrt{e^2 x^2 + a f^2} d^2 e x^3 + 2\sqrt{e^2 x^2 + a f^2} d e^2 x^4}{2e}$$

input `int((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^3,x)`

output `(4*sqrt(a*f**2 + e**2*x**2)*a*d*f**2 + 2*sqrt(a*f**2 + e**2*x**2)*a*e*f**2*x + 3*sqrt(a*f**2 + e**2*x**2)*d**2*e*x + 4*sqrt(a*f**2 + e**2*x**2)*d*e**2*x**2 + 2*sqrt(a*f**2 + e**2*x**2)*e**3*x**3 + 3*log((sqrt(a*f**2 + e**2*x**2) + e*x)/(sqrt(a)*f))*a*d**2*f**2 + 6*a*d*e*f**2*x + 3*a*e**2*f**2*x**2 + 2*d**3*e*x + 3*d**2*e**2*x**2 + 4*d*e**3*x**3 + 2*e**4*x**4)/(2*e)`

### 3.3 $\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^2 dx$

Optimal result . . . . .	63
Mathematica [A] (verified) . . . . .	64
Rubi [A] (verified) . . . . .	64
Maple [A] (verified) . . . . .	66
Fricas [A] (verification not implemented) . . . . .	66
Sympy [A] (verification not implemented) . . . . .	67
Maxima [A] (verification not implemented) . . . . .	68
Giac [A] (verification not implemented) . . . . .	68
Mupad [B] (verification not implemented) . . . . .	69
Reduce [B] (verification not implemented) . . . . .	69

#### Optimal result

Integrand size = 25, antiderivative size = 106

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^2 dx = (d^2 + af^2)x + dex^2 + \frac{2e^2 x^3}{3} + dfx \sqrt{a + \frac{e^2 x^2}{f^2}} + \frac{2f^3 \left( a + \frac{e^2 x^2}{f^2} \right)^{3/2}}{3e} + \frac{adf^2 \operatorname{arctanh} \left( \frac{ex}{f \sqrt{a + \frac{e^2 x^2}{f^2}}} \right)}{e}$$

output

```
(a*f^2+d^2)*x+d*e*x^2+2/3*e^2*x^3+d*f*x*(a+e^2*x^2/f^2)^(1/2)+2/3*f^3*(a+e^2*x^2/f^2)^(3/2)/e+a*d*f^2*arctanh(e*x/f/(a+e^2*x^2/f^2)^(1/2))/e
```



**Mathematica [A] (verified)**

Time = 0.88 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.07

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^2 dx = d^2 x + af^2 x + dex^2 + \frac{2e^2 x^3}{3} + \frac{\sqrt{a + \frac{e^2 x^2}{f^2}} (2af^3 + ef x (3d + 2ex))}{3e} + \frac{2adf^2 \operatorname{arctanh} \left( \frac{f \left( -\sqrt{a} + \sqrt{a + \frac{e^2 x^2}{f^2}} \right)}{ex} \right)}{e}$$

input

```
Integrate[(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^2,x]
```

output

```
d^2*x + a*f^2*x + d*e*x^2 + (2*e^2*x^3)/3 + (Sqrt[a + (e^2*x^2)/f^2]*(2*a*f^3 + e*f*x*(3*d + 2*e*x)))/(3*e) + (2*a*d*f^2*ArcTanh[(f*(-Sqrt[a] + Sqrt[a + (e^2*x^2)/f^2]))/(e*x))]/e
```

**Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.24, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$ , Rules used = {2542, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( f \sqrt{a + \frac{e^2 x^2}{f^2}} + d + ex \right)^2 dx$$

↓ 2542

$$\int \frac{\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^2 \left(d^2-2\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)d+af^2+\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^2\right)}{\left(-\sqrt{\frac{e^2x^2}{f^2}+af-ex}\right)^2} d\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)$$

$2e$   
↓ 1195

$$\int \left(af^2 - \frac{2adf^2}{-\sqrt{\frac{e^2x^2}{f^2}+af-ex}} + \frac{ad^2f^2}{\left(-\sqrt{\frac{e^2x^2}{f^2}+af-ex}\right)^2} + \left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^2\right) d\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)$$

$2e$   
↓ 2009

$$\frac{ad^2f^2}{f\left(-\sqrt{a+\frac{e^2x^2}{f^2}}\right)-ex} + \frac{1}{3}\left(f\sqrt{a+\frac{e^2x^2}{f^2}}+d+ex\right)^3 + af^2\left(f\sqrt{a+\frac{e^2x^2}{f^2}}+d+ex\right) + 2adf^2\log\left(f\left(-\sqrt{a+\frac{e^2x^2}{f^2}}\right)-ex\right)$$

$2e$

input `Int[(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^2,x]`

output `((a*d^2*f^2)/(-(e*x) - f*Sqrt[a + (e^2*x^2)/f^2]) + a*f^2*(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]) + (d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^3/3 + 2*a*d*f^2*Log[-(e*x) - f*Sqrt[a + (e^2*x^2)/f^2]])/(2*e)`

### Defintions of rubi rules used

rule 1195 `Int[((d_.) + (e_.)*(x_))^(m_.)*((f_.) + (g_.)*(x_))^(n_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2542

```
Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(
n_))^(p_.), x_Symbol] :> Simp[1/(2*e) Subst[Int[(g + h*x^n)^p*((d^2 + a*f
^2 - 2*d*x + x^2)/(d - x)^2), x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; Fr
eeQ[{a, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]
```

**Maple [A] (verified)**

Time = 0.97 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.17

method	result	size
default	$\frac{e^2 x^3}{3} + a f^2 x + 2f \left( d \left( \frac{x \sqrt{a + \frac{e^2 x^2}{f^2}}}{2} + \frac{a \ln \left( \frac{e^2 x}{f^2 \sqrt{\frac{e^2}{f^2}} + \sqrt{a + \frac{e^2 x^2}{f^2}}} \right)}{2 \sqrt{\frac{e^2}{f^2}}} \right) + \frac{f^2 \left( \frac{e^2 x^2 + f^2 a}{f^2} \right)^{\frac{3}{2}}}{3e} \right) + \frac{(ex+d)^3}{3e}$	124

input

```
int((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^2,x,method=_RETURNVERBOSE)
```

output

```
1/3*e^2*x^3+a*f^2*x+2*f*(d*(1/2*x*(a+e^2*x^2/f^2)^(1/2)+1/2*a*ln(e^2*x/f^2
/(e^2/f^2)^(1/2)+(a+e^2*x^2/f^2)^(1/2))/(e^2/f^2)^(1/2))+1/3/e*f^2*((e^2*x
^2+a*f^2)/f^2)^(3/2))+1/3*(e*x+d)^3/e
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.08

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^2 dx$$

$$= \frac{2e^3 x^3 + 3de^2 x^2 - 3adf^2 \log \left( -ex + f \sqrt{\frac{e^2 x^2 + af^2}{f^2}} \right) + 3(af^2 + d^2e)x + (2e^2 f x^2 + 2af^3 + 3defx) \sqrt{\frac{e^2 x^2 + af^2}{f^2}}}{3e}$$

input

```
integrate((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^2,x, algorithm="fricas")
```

output

```
1/3*(2*e^3*x^3 + 3*d*e^2*x^2 - 3*a*d*f^2*log(-e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2)) + 3*(a*e*f^2 + d^2*e)*x + (2*e^2*f*x^2 + 2*a*f^3 + 3*d*e*f*x)*sqrt((e^2*x^2 + a*f^2)/f^2))/e
```

**Sympy [A] (verification not implemented)**

Time = 0.71 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.75

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^2 dx$$

$$= af^2 x + d^2 x + dex^2$$

$$+ 2df \left( \frac{a \left( \begin{cases} \frac{\log\left(\frac{2e^2 x + 2\sqrt{\frac{e^2}{f^2}} \sqrt{a + \frac{e^2 x^2}{f^2}}}\right)}{\sqrt{\frac{e^2}{f^2}}} & \text{for } a \neq 0 \\ \frac{x \log(x)}{\sqrt{\frac{e^2 x^2}{f^2}}} & \text{otherwise} \end{cases} \right)}{2} + \frac{x \sqrt{a + \frac{e^2 x^2}{f^2}}}{2} \right) \begin{matrix} \text{for } \frac{e^2}{f^2} \neq 0 \\ \text{otherwise} \end{matrix}$$

$$+ \frac{2e^2 x^3}{3} + 2ef \left( \begin{matrix} \sqrt{a + \frac{e^2 x^2}{f^2}} \left( \frac{af^2}{3e^2} + \frac{x^2}{3} \right) & \text{for } \frac{e^2}{f^2} \neq 0 \\ \frac{\sqrt{ax^2}}{2} & \text{otherwise} \end{matrix} \right)$$

input

```
integrate((d+e*x+f*(a+e**2*x**2/f**2)**(1/2))**2,x)
```

output

```
a*f**2*x + d**2*x + d*e*x**2 + 2*d*f*Piecewise((a*Piecewise((log(2*e**2*x/f**2 + 2*sqrt(e**2/f**2)*sqrt(a + e**2*x**2/f**2))/sqrt(e**2/f**2), Ne(a, 0)), (x*log(x)/sqrt(e**2*x**2/f**2), True))/2 + x*sqrt(a + e**2*x**2/f**2)/2, Ne(e**2/f**2, 0)), (sqrt(a)*x, True)) + 2*e**2*x**3/3 + 2*e*f*Piecewise((sqrt(a + e**2*x**2/f**2)*(a*f**2/(3*e**2) + x**2/3), Ne(e**2/f**2, 0)), (sqrt(a)*x**2/2, True))
```

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.01

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^2 dx$$

$$= \frac{1}{3} e^2 x^3 + \frac{2 \left( \frac{e^2 x^2}{f^2} + a \right)^{\frac{3}{2}} f^3}{3e} + \frac{1}{3} \left( \frac{e^2 x^3}{f^2} + 3ax \right) f^2 + d^2 x$$

$$+ \left( ex^2 + \left( \frac{af \operatorname{arsinh} \left( \frac{e^2 x}{\sqrt{ae^2} f} \right)}{\sqrt{e^2}} + \sqrt{\frac{e^2 x^2}{f^2} + ax} \right) f \right) d$$

input `integrate((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^2,x, algorithm="maxima")`output `1/3*e^2*x^3 + 2/3*(e^2*x^2/f^2 + a)^(3/2)*f^3/e + 1/3*(e^2*x^3/f^2 + 3*a*x)*f^2 + d^2*x + (e*x^2 + (a*f*arcsinh(e^2*x/(sqrt(a*e^2)*f))/sqrt(e^2) + sqrt(e^2*x^2/f^2 + a)*x)*f)*d`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.01

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^2 dx = \frac{2}{3} e^2 x^3 + af^2 x + dex^2$$

$$- \frac{adf|f| \log(|-x|e| + \sqrt{e^2 x^2 + af^2})}{|e|} + d^2 x$$

$$+ \frac{1}{3} \sqrt{e^2 x^2 + af^2} \left( \left( \frac{2ex|f|}{f} + \frac{3d|f|}{f} \right) x + \frac{2af|f|}{e} \right)$$

input `integrate((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^2,x, algorithm="giac")`output `2/3*e^2*x^3 + a*f^2*x + d*e*x^2 - a*d*f*abs(f)*log(abs(-x*abs(e) + sqrt(e^2*x^2 + a*f^2)))/abs(e) + d^2*x + 1/3*sqrt(e^2*x^2 + a*f^2)*((2*e*x*abs(f)/f + 3*d*abs(f)/f)*x + 2*a*f*abs(f)/e)`

**Mupad [B] (verification not implemented)**

Time = 23.86 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.98

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^2 dx$$

$$= \left\{ \begin{array}{l} x(d + \sqrt{a}f)^2 \\ x(d^2 + af^2) + \frac{2e^2 x^3}{3} + dex^2 + \frac{2af^3 \sqrt{a + \frac{e^2 x^2}{f^2}}}{e} - \frac{2f \sqrt{a + \frac{e^2 x^2}{f^2}} (2af^2 - e^2 x^2)}{3e} + dfx \sqrt{a + \frac{e^2 x^2}{f^2}} + \frac{2adf \ln(x)}{3e} \end{array} \right.$$

input `int((d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^2,x)`output `piecewise(e == 0, x*(d + a^(1/2)*f)^2, e ~= 0, x*(a*f^2 + d^2) + (2*e^2*x^3)/3 + d*e*x^2 + (2*a*f^3*(a + (e^2*x^2)/f^2)^(1/2))/e - (2*f*(a + (e^2*x^2)/f^2)^(1/2)*(2*a*f^2 - e^2*x^2))/(3*e) + d*f*x*(a + (e^2*x^2)/f^2)^(1/2) + (2*a*d*f*log(x*(e^2/f^2)^(1/2) + (a + (e^2*x^2)/f^2)^(1/2)))/(e^2/f^2)^(1/2) - (a*d*e^2*log(2*x*(e^2/f^2)^(1/2) + 2*(a + (e^2*x^2)/f^2)^(1/2)))/(f*(e^2/f^2)^(3/2)))`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.25

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^2 dx$$

$$= \frac{2\sqrt{e^2 x^2 + a f^2} a f^2 + 3\sqrt{e^2 x^2 + a f^2} dex + 2\sqrt{e^2 x^2 + a f^2} e^2 x^2 + 3 \log\left(\frac{\sqrt{e^2 x^2 + a f^2} + ex}{\sqrt{a} f}\right) ad f^2 + 3ae f^2 x}{3e}$$

input `int((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^2,x)`output `(2*sqrt(a*f**2 + e**2*x**2)*a*f**2 + 3*sqrt(a*f**2 + e**2*x**2)*d*e*x + 2*sqrt(a*f**2 + e**2*x**2)*e**2*x**2 + 3*log((sqrt(a*f**2 + e**2*x**2) + e*x)/(sqrt(a)*f))*a*d*f**2 + 3*a*e*f**2*x + 3*d**2*e*x + 3*d*e**2*x**2 + 2*e**3*x**3)/(3*e)`

### 3.4 $\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right) dx$

Optimal result . . . . .	70
Mathematica [A] (verified) . . . . .	70
Rubi [A] (verified) . . . . .	71
Maple [A] (verified) . . . . .	72
Fricas [A] (verification not implemented) . . . . .	72
Sympy [A] (verification not implemented) . . . . .	73
Maxima [A] (verification not implemented) . . . . .	73
Giac [A] (verification not implemented) . . . . .	74
Mupad [B] (verification not implemented) . . . . .	74
Reduce [B] (verification not implemented) . . . . .	75

#### Optimal result

Integrand size = 23, antiderivative size = 68

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right) dx = dx + \frac{ex^2}{2} + \frac{1}{2}fx \sqrt{a + \frac{e^2 x^2}{f^2}} + \frac{af^2 \operatorname{arctanh} \left( \frac{ex}{f \sqrt{a + \frac{e^2 x^2}{f^2}}} \right)}{2e}$$

output

```
d*x+1/2*e*x^2+1/2*f*x*(a+e^2*x^2/f^2)^(1/2)+1/2*a*f^2*arctanh(e*x/f/(a+e^2*x^2/f^2)^(1/2))/e
```

#### Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.18

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right) dx = dx + \frac{ex^2}{2} + \frac{1}{2}fx \sqrt{a + \frac{e^2 x^2}{f^2}} + \frac{af^2 \operatorname{arctanh} \left( \frac{-\frac{\sqrt{a}f}{e} + \frac{f \sqrt{a + \frac{e^2 x^2}{f^2}}}{e}}{x} \right)}{e}$$

input `Integrate[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2],x]`

output `d*x + (e*x^2)/2 + (f*x*Sqrt[a + (e^2*x^2)/f^2])/2 + (a*f^2*ArcTanh[-((Sqrt[a]*f)/e) + (f*Sqrt[a + (e^2*x^2)/f^2])/e]/x))/e`

### Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( f \sqrt{a + \frac{e^2 x^2}{f^2}} + d + ex \right) dx$$

↓ 2009

$$\frac{a f^2 \operatorname{arctanh}\left(\frac{ex}{f \sqrt{a + \frac{e^2 x^2}{f^2}}}\right)}{2e} + \frac{1}{2} f x \sqrt{a + \frac{e^2 x^2}{f^2}} + dx + \frac{ex^2}{2}$$

input `Int[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2],x]`

output `d*x + (e*x^2)/2 + (f*x*Sqrt[a + (e^2*x^2)/f^2])/2 + (a*f^2*ArcTanh[(e*x)/(f*Sqrt[a + (e^2*x^2)/f^2]])]/(2*e)`



**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.10

method	result	size
default	$dx + \frac{ex^2}{2} + \frac{fx\sqrt{a + \frac{e^2x^2}{f^2}}}{2} + \frac{fa \ln\left(\frac{e^2x}{f^2\sqrt{\frac{e^2}{f^2}} + \sqrt{a + \frac{e^2x^2}{f^2}}}\right)}{2\sqrt{\frac{e^2}{f^2}}}$	75
parts	$dx + \frac{ex^2}{2} + \frac{fx\sqrt{a + \frac{e^2x^2}{f^2}}}{2} + \frac{fa \ln\left(\frac{e^2x}{f^2\sqrt{\frac{e^2}{f^2}} + \sqrt{a + \frac{e^2x^2}{f^2}}}\right)}{2\sqrt{\frac{e^2}{f^2}}}$	75

input `int(d+e*x+f*(a+e^2*x^2/f^2)^(1/2),x,method=_RETURNVERBOSE)`

output `d*x+1/2*e*x^2+1/2*f*x*(a+e^2*x^2/f^2)^(1/2)+1/2*f*a*ln(e^2*x/f^2/(e^2/f^2)^(1/2)+(a+e^2*x^2/f^2)^(1/2))/(e^2/f^2)^(1/2)`

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.09

$$\int \left( d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}} \right) dx$$

$$= \frac{e^2x^2 - af^2 \log\left(-ex + f\sqrt{\frac{e^2x^2 + af^2}{f^2}}\right) + efx\sqrt{\frac{e^2x^2 + af^2}{f^2}} + 2dex}{2e}$$

input `integrate(d+e*x+f*(a+e^2*x^2/f^2)^(1/2),x, algorithm="fricas")`

output

```
1/2*(e^2*x^2 - a*f^2*log(-e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2)) + e*f*x*sqrt((e^2*x^2 + a*f^2)/f^2) + 2*d*e*x)/e
```

**Sympy [A] (verification not implemented)**

Time = 1.06 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.79

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right) dx = dx + \frac{ex^2}{2} + f \left( \frac{\sqrt{ax} \sqrt{1 + \frac{e^2 x^2}{af^2}}}{2} + \frac{af \operatorname{asinh}\left(\frac{ex}{\sqrt{af}}\right)}{2e} \right)$$

input

```
integrate(d+e*x+f*(a+e**2*x**2/f**2)**(1/2),x)
```

output

```
d*x + e*x**2/2 + f*(sqrt(a)*x*sqrt(1 + e**2*x**2/(a*f**2))/2 + a*f*asinh(e*x/(sqrt(a)*f))/(2*e))
```

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.79

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right) dx = \frac{1}{2} ex^2 + \frac{1}{2} \left( \frac{af \operatorname{arsinh}\left(\frac{e^2 x}{\sqrt{ae^2} f}\right)}{\sqrt{e^2}} + \sqrt{\frac{e^2 x^2}{f^2} + ax} \right) f + dx$$

input

```
integrate(d+e*x+f*(a+e^2*x^2/f^2)^(1/2),x, algorithm="maxima")
```

output

```
1/2*e*x^2 + 1/2*(a*f*arcsinh(e^2*x/(sqrt(a*e^2)*f))/sqrt(e^2) + sqrt(e^2*x^2/f^2 + a)*x)*f + d*x
```

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.00

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right) dx = \frac{1}{2} ex^2 + dx - \frac{\left( \frac{af^2 \log\left(\frac{|-x|e| + \sqrt{e^2 x^2 + af^2}}{|e|}\right) - \sqrt{e^2 x^2 + af^2} x}{|f|} \right)}{2f}$$

input `integrate(d+e*x+f*(a+e^2*x^2/f^2)^(1/2),x, algorithm="giac")`output `1/2*e*x^2 + d*x - 1/2*(a*f^2*log(abs(-x*abs(e) + sqrt(e^2*x^2 + a*f^2)))/abs(e) - sqrt(e^2*x^2 + a*f^2)*x)*abs(f)/f`**Mupad [B] (verification not implemented)**

Time = 22.84 (sec) , antiderivative size = 136, normalized size of antiderivative = 2.00

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right) dx = \begin{cases} x(d + \sqrt{a}f) & \text{if } e = 0 \\ dx + \frac{ex^2}{2} + \frac{fx\sqrt{a + \frac{e^2 x^2}{f^2}}}{2} + \frac{ae^2 \ln\left(x\sqrt{\frac{e^2}{f^2} + \sqrt{a + \frac{e^2 x^2}{f^2}}}\right)}{f\left(\frac{e^2}{f^2}\right)^{3/2}} - \frac{ae^2 \ln\left(2x\sqrt{\frac{e^2}{f^2} + 2\sqrt{a + \frac{e^2 x^2}{f^2}}}\right)}{2f\left(\frac{e^2}{f^2}\right)^{3/2}} & \text{if } e \neq 0 \end{cases}$$

input `int(d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2),x)`output `piecewise(e == 0, x*(d + a^(1/2)*f), e ~= 0, d*x + (e*x^2)/2 + (f*x*(a + (e^2*x^2)/f^2)^(1/2))/2 + (a*e^2*log(x*(e^2/f^2)^(1/2) + (a + (e^2*x^2)/f^2)^(1/2)))/(f*(e^2/f^2)^(3/2)) - (a*e^2*log(2*x*(e^2/f^2)^(1/2) + 2*(a + (e^2*x^2)/f^2)^(1/2)))/(2*f*(e^2/f^2)^(3/2)))`

**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.99

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right) dx$$

$$= \frac{\sqrt{e^2 x^2 + a f^2} ex + \log\left(\frac{\sqrt{e^2 x^2 + a f^2} + ex}{\sqrt{a} f}\right) a f^2 + 2dex + e^2 x^2}{2e}$$

input

```
int(d+e*x+f*(a+e^2*x^2/f^2)^(1/2),x)
```

output

```
(sqrt(a*f**2 + e**2*x**2)*e*x + log((sqrt(a*f**2 + e**2*x**2) + e*x)/(sqrt
(a)*f))*a*f**2 + 2*d*e*x + e**2*x**2)/(2*e)
```

**3.5** 
$$\int \frac{1}{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}} dx$$

Optimal result	76
Mathematica [B] (verified)	76
Rubi [A] (verified)	77
Maple [B] (verified)	78
Fricas [A] (verification not implemented)	79
Sympy [F]	80
Maxima [F]	80
Giac [B] (verification not implemented)	81
Mupad [F(-1)]	81
Reduce [B] (verification not implemented)	82

**Optimal result**

Integrand size = 25, antiderivative size = 117

$$\int \frac{1}{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}} dx = -\frac{af^2}{2de\left(ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)} - \frac{af^2 \log\left(ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)}{2d^2e} + \frac{\left(1+\frac{af^2}{d^2}\right) \log\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)}{2e}$$

output `-1/2*a*f^2/d/e/(e*x+f*(a+e^2*x^2/f^2)^(1/2))-1/2*a*f^2*ln(e*x+f*(a+e^2*x^2/f^2)^(1/2))/d^2/e+1/2*(1+a*f^2/d^2)*ln(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))/e`

**Mathematica [B] (verified)**

Leaf count is larger than twice the leaf count of optimal. 337 vs. 2(117) = 234.

Time = 1.23 (sec) , antiderivative size = 337, normalized size of antiderivative = 2.88

$$\int \frac{1}{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}} dx = \frac{2de^2x - 2def\sqrt{a+\frac{e^2x^2}{f^2}} - \left(af^2\left(e - \sqrt{\frac{e^2}{f^2}}f\right) + d^2\left(e + \sqrt{\frac{e^2}{f^2}}f\right)\right) \log\left(-\sqrt{\frac{e^2}{f^2}}x + \sqrt{a+\frac{e^2x^2}{f^2}}\right) + \sqrt{\frac{e^2}{f^2}}f(a + \dots)}{\dots}$$

input `Integrate[(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(-1),x]`

output  $(2*d*e^2*x - 2*d*e*f*\text{Sqrt}[a + (e^2*x^2)/f^2] - (a*f^2*(e - \text{Sqrt}[e^2/f^2]*f) + d^2*(e + \text{Sqrt}[e^2/f^2]*f))*\text{Log}[-(\text{Sqrt}[e^2/f^2]*x) + \text{Sqrt}[a + (e^2*x^2)/f^2]] + \text{Sqrt}[e^2/f^2]*f*(d^2 + a*f^2)*\text{Log}[a*f + d*(-(\text{Sqrt}[e^2/f^2]*x) + \text{Sqrt}[a + (e^2*x^2)/f^2])] + e*(d^2 + a*f^2)*\text{Log}[d*e*(a*f + d*(-(\text{Sqrt}[e^2/f^2]*x) + \text{Sqrt}[a + (e^2*x^2)/f^2]))] - \text{Sqrt}[e^2/f^2]*f*(d^2 + a*f^2)*\text{Log}[d + f*(-(\text{Sqrt}[e^2/f^2]*x) + \text{Sqrt}[a + (e^2*x^2)/f^2])] + e*(d^2 + a*f^2)*\text{Log}[d^2*e*(d + f*(-(\text{Sqrt}[e^2/f^2]*x) + \text{Sqrt}[a + (e^2*x^2)/f^2]))]/(4*d^2*e^2)$

### Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.95, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$ , Rules used = {2542, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex} dx$$

↓ 2542

$$\int \frac{d^2 - 2\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)d + af^2 + \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)^2}{\left(-\sqrt{\frac{e^2x^2}{f^2} + a} - ex\right)^2 \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)} d\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)$$

2e

↓ 1195

$$\int \left( \frac{af^2}{d^2\left(-\sqrt{\frac{e^2x^2}{f^2} + a} - ex\right)} + \frac{af^2}{d\left(-\sqrt{\frac{e^2x^2}{f^2} + a} - ex\right)^2} + \frac{d^2 + af^2}{d^2\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)} \right) d\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)$$

2e

↓ 2009

$$\frac{-\frac{af^2 \log\left(f\left(-\sqrt{a+\frac{e^2x^2}{f^2}}\right)-ex\right)}{d^2} + \left(\frac{af^2}{d^2} + 1\right) \log\left(f\sqrt{a+\frac{e^2x^2}{f^2}} + d + ex\right) + \frac{af^2}{d\left(f\left(-\sqrt{a+\frac{e^2x^2}{f^2}}\right)-ex\right)}}{2e}$$

input `Int[(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(-1),x]`

output `((a*f^2)/(d*(-(e*x) - f*Sqrt[a + (e^2*x^2)/f^2])) - (a*f^2*Log[-(e*x) - f*Sqrt[a + (e^2*x^2)/f^2]])/d^2 + (1 + (a*f^2)/d^2)*Log[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]])/(2*e)`

### Defintions of rubi rules used

rule 1195 `Int[((d_.) + (e_.)*(x_))^(m_.)*((f_.) + (g_.)*(x_))^(n_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2542 `Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2]))^(n_)^(p_.), x_Symbol] := Simp[1/(2*e) Subst[Int[(g + h*x^n)^p*((d^2 + a*f^2 - 2*d*x + x^2)/(d - x)^2), x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]`

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 650 vs.  $2(105) = 210$ .

Time = 0.08 (sec) , antiderivative size = 651, normalized size of antiderivative = 5.56

method	result
default	$f \sqrt{\frac{4e^2 \left(x + \frac{-f^2 a + d^2}{2de}\right)^2}{f^2} + \frac{4e(f^2 a - d^2) \left(x + \frac{-f^2 a + d^2}{2de}\right)}{f^2 d} + \frac{a^2 f^4 + 2a d^2 f^2 + d^4}{d^2 f^2}} + \frac{e^{(f^2 a - d^2)} \ln\left(\frac{e^{(f^2 a - d^2)} + \frac{e^2 \left(x + \frac{-f^2 a + d^2}{2de}\right)}{2f^2 d} + \frac{e^2}{\sqrt{\frac{e^2}{f^2}}}}{f^2}\right) + \sqrt{\frac{e^2}{f^2}}}{4d^2 e}$

```
input int(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2)),x,method=_RETURNVERBOSE)
```

```
output -1/2*f/d/e*(1/2*(4*e^2*(x+1/2*(-a*f^2+d^2)/d/e)^2/f^2+4*e*(a*f^2-d^2)/f^2/d*(x+1/2*(-a*f^2+d^2)/d/e)+(a^2*f^4+2*a*d^2*f^2+d^4)/d^2/f^2)^(1/2)+1/2*e*(a*f^2-d^2)/f^2/d*ln((1/2*e*(a*f^2-d^2)/f^2/d+e^2/f^2*(x+1/2*(-a*f^2+d^2)/d/e))/(e^2/f^2)^(1/2)+(e^2*(x+1/2*(-a*f^2+d^2)/d/e)^2/f^2+e*(a*f^2-d^2)/f^2/d*(x+1/2*(-a*f^2+d^2)/d/e)+1/4*(a^2*f^4+2*a*d^2*f^2+d^4)/d^2/f^2)^(1/2))/(e^2/f^2)^(1/2)-1/2*(a^2*f^4+2*a*d^2*f^2+d^4)/d^2/f^2/((a^2*f^4+2*a*d^2*f^2+d^4)/d^2/f^2)^(1/2)*ln((1/2*(a^2*f^4+2*a*d^2*f^2+d^4)/d^2/f^2+e*(a*f^2-d^2)/f^2/d*(x+1/2*(-a*f^2+d^2)/d/e)+1/2*((a^2*f^4+2*a*d^2*f^2+d^4)/d^2/f^2)^(1/2))*(4*e^2*(x+1/2*(-a*f^2+d^2)/d/e)^2/f^2+4*e*(a*f^2-d^2)/f^2/d*(x+1/2*(-a*f^2+d^2)/d/e)+(a^2*f^4+2*a*d^2*f^2+d^4)/d^2/f^2)^(1/2))/(x+1/2*(-a*f^2+d^2)/d/e))+1/2*ln(a*f^2-2*d*e*x-d^2)/e-e*(-1/2*x/d/e+1/4*(-a*f^2+d^2)/d^2/e^2*ln(-a*f^2+2*d*e*x+d^2))
```

**Fricas [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.60

$$\int \frac{1}{d + ex + f\sqrt{a + \frac{e^2 x^2}{f^2}}} dx$$

$$= \frac{2 dex - 2 df \sqrt{\frac{e^2 x^2 + af^2}{f^2}} + (af^2 + d^2) \log\left(af^2 - dex + df \sqrt{\frac{e^2 x^2 + af^2}{f^2}}\right) + (af^2 + d^2) \log\left(-af^2 + 2 dex + \dots\right)}{4d^2 e}$$



input `integrate(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2)),x, algorithm="fricas")`

output `1/4*(2*d*e*x - 2*d*f*sqrt((e^2*x^2 + a*f^2)/f^2) + (a*f^2 + d^2)*log(a*f^2 - d*e*x + d*f*sqrt((e^2*x^2 + a*f^2)/f^2)) + (a*f^2 + d^2)*log(-a*f^2 + 2*d*e*x + d^2) - (a*f^2 + d^2)*log(-e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) - d) + (a*f^2 - d^2)*log(-e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2)))/(d^2*e)`

### Sympy [F]

$$\int \frac{1}{d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}} dx = \int \frac{1}{d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}} dx$$

input `integrate(1/(d+e*x+f*(a+e**2*x**2/f**2)**(1/2)),x)`

output `Integral(1/(d + e*x + f*sqrt(a + e**2*x**2/f**2)), x)`

### Maxima [F]

$$\int \frac{1}{d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}} dx = \int \frac{1}{ex + \sqrt{\frac{e^2x^2}{f^2} + af + d}} dx$$

input `integrate(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2)),x, algorithm="maxima")`

output `integrate(1/(e*x + sqrt(e^2*x^2/f^2 + a)*f + d), x)`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 299 vs.  $2(105) = 210$ .

Time = 0.16 (sec) , antiderivative size = 299, normalized size of antiderivative = 2.56

$$\int \frac{1}{d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}} dx$$

$$= \frac{x}{2d} + \frac{(af^2 + d^2) \log(|-af^2 + 2dex + d^2|)}{4d^2e}$$

$$- \frac{\sqrt{e^2x^2 + af^2}|f|}{2def} + \frac{(af^2|f| - d^2|f|) \log(|-x|e| + \sqrt{e^2x^2 + af^2}|)}{4d^2f|e|}$$

$$- \frac{(a^2e^2f^4|f| + 2ad^2e^2f^2|f| + d^4e^2|f|) \log\left(\frac{ae f^2 - d^2e - 2(x|e| - \sqrt{e^2x^2 + af^2})d|e| - |ae f^2 + d^2e|}{ae f^2 - d^2e - 2(x|e| - \sqrt{e^2x^2 + af^2})d|e| + |ae f^2 + d^2e|}\right)}{4d^2ef|ae f^2 + d^2e||e|}$$

input `integrate(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2)),x, algorithm="giac")`

output `1/2*x/d + 1/4*(a*f^2 + d^2)*log(abs(-a*f^2 + 2*d*e*x + d^2))/(d^2*e) - 1/2*sqrt(e^2*x^2 + a*f^2)*abs(f)/(d*e*f) + 1/4*(a*f^2*abs(f) - d^2*abs(f))*log(abs(-x*abs(e) + sqrt(e^2*x^2 + a*f^2)))/(d^2*f*abs(e)) - 1/4*(a^2*e^2*f^4*abs(f) + 2*a*d^2*e^2*f^2*abs(f) + d^4*e^2*abs(f))*log(abs(a*e*f^2 - d^2*e - 2*(x*abs(e) - sqrt(e^2*x^2 + a*f^2))*d*abs(e) - abs(a*e*f^2 + d^2*e))/abs(a*e*f^2 - d^2*e - 2*(x*abs(e) - sqrt(e^2*x^2 + a*f^2))*d*abs(e) + abs(a*e*f^2 + d^2*e)))/(d^2*e*f*abs(a*e*f^2 + d^2*e)*abs(e))`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}} dx = \int \frac{1}{d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}} dx$$

input `int(1/(d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2)),x)`

output `int(1/(d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2)), x)`

**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.13

$$\int \frac{1}{d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}} dx$$

$$= \frac{-\sqrt{e^2x^2 + af^2}d - \log\left(\frac{\sqrt{e^2x^2 + af^2} + ex}{\sqrt{af}}\right)af^2 + \log\left(\frac{\sqrt{e^2x^2 + af^2}a + ad + aex}{\sqrt{a}}\right)af^2 + \log\left(\frac{\sqrt{e^2x^2 + af^2}a + ad + aex}{\sqrt{a}}\right)d}{2d^2e}$$

input `int(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2)),x)`output `( - sqrt(a*f**2 + e**2*x**2)*d - log((sqrt(a*f**2 + e**2*x**2) + e*x)/(sqrt(a)*f))*a*f**2 + log((sqrt(a*f**2 + e**2*x**2)*a + a*d + a*e*x)/sqrt(a))*a*f**2 + log((sqrt(a*f**2 + e**2*x**2)*a + a*d + a*e*x)/sqrt(a))*d**2 + d*e*x)/(2*d**2*e)`

**3.6** 
$$\int \frac{1}{\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)^2} dx$$

Optimal result . . . . .	83
Mathematica [A] (verified) . . . . .	84
Rubi [A] (verified) . . . . .	84
Maple [B] (verified) . . . . .	86
Fricas [B] (verification not implemented) . . . . .	87
Sympy [F] . . . . .	87
Maxima [F] . . . . .	88
Giac [A] (verification not implemented) . . . . .	88
Mupad [F(-1)] . . . . .	89
Reduce [B] (verification not implemented) . . . . .	89

**Optimal result**

Integrand size = 25, antiderivative size = 151

$$\int \frac{1}{\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)^2} dx = -\frac{af^2}{2d^2e\left(ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)} - \frac{1+\frac{af^2}{d^2}}{2e\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)} - \frac{af^2\log\left(ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)}{d^3e} + \frac{af^2\log\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)}{d^3e}$$

output

```
-1/2*a*f^2/d^2/e/(e*x+f*(a+e^2*x^2/f^2)^(1/2))-1/2*(1+a*f^2/d^2)/e/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))-a*f^2*ln(e*x+f*(a+e^2*x^2/f^2)^(1/2))/d^3/e+a*f^2*ln(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))/d^3/e
```

### Mathematica [A] (verified)

Time = 0.97 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.51

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^2} dx$$

$$= \frac{-d\left(d^3 + d^2ex + af^2\left(ex - f\sqrt{a + \frac{e^2x^2}{f^2}}\right) + dex\left(-ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)\right) + af^2(-d^2 + af^2 - 2dex) \log\left(\frac{-d^3 + d^2ex + af^2\left(ex - f\sqrt{a + \frac{e^2x^2}{f^2}}\right) + dex\left(-ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right) + af^2(-d^2 + af^2 - 2dex)}{d^3 + d^2ex + af^2\left(ex - f\sqrt{a + \frac{e^2x^2}{f^2}}\right) + dex\left(-ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right) + af^2(-d^2 + af^2 - 2dex)}\right)}{d^3 + d^2ex + af^2\left(ex - f\sqrt{a + \frac{e^2x^2}{f^2}}\right) + dex\left(-ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right) + af^2(-d^2 + af^2 - 2dex)}$$

input `Integrate[(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(-2),x]`

output `(-(d*(d^3 + d^2*e*x + a*f^2*(e*x - f*Sqrt[a + (e^2*x^2)/f^2]) + d*e*x*(-(e*x) + f*Sqrt[a + (e^2*x^2)/f^2]))) + a*f^2*(-d^2 + a*f^2 - 2*d*e*x)*Log[-(Sqrt[a]*f) + e*x + f*Sqrt[a + (e^2*x^2)/f^2]] + a*f^2*(d^2 - a*f^2 + 2*d*e*x)*Log[-(a*f^2) + d*(e*x - f*Sqrt[a + (e^2*x^2)/f^2]) + Sqrt[a]*f*(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])])/(d^3*e*(d^2 - a*f^2 + 2*d*e*x))`

### Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.97, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$ , Rules used = {2542, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\left(f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex\right)^2} dx$$

↓ 2542

$$\int \frac{d^2 - 2\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)d + af^2 + \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)^2}{\left(-\sqrt{\frac{e^2x^2}{f^2} + a}f - ex\right)^2 \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)^2} d \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)$$

2e

↓ 1195

$$\frac{\int \left( \frac{2af^2}{d^3 \left( -\sqrt{\frac{e^2x^2}{f^2} + af - ex} \right)} + \frac{2af^2}{d^3 \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + a} \right)} + \frac{af^2}{d^2 \left( -\sqrt{\frac{e^2x^2}{f^2} + af - ex} \right)^2} + \frac{d^2+af^2}{d^2 \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + a} \right)^2} \right) d(d+ex+f\sqrt{\frac{e^2x^2}{f^2} + a})}{2e}$$

↓ 2009

$$\frac{-\frac{2af^2 \log \left( f \left( -\sqrt{a + \frac{e^2x^2}{f^2}} \right) - ex \right)}{d^3} + \frac{2af^2 \log \left( f \sqrt{a + \frac{e^2x^2}{f^2}} + d+ex \right)}{d^3} + \frac{af^2}{d^2 \left( f \left( -\sqrt{a + \frac{e^2x^2}{f^2}} \right) - ex \right)} - \frac{\frac{af^2}{d^2} + 1}{f \sqrt{a + \frac{e^2x^2}{f^2}} + d+ex}}{2e}$$

input

```
Int[(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(-2), x]
```

output

```
((a*f^2)/(d^2*(-(e*x) - f*Sqrt[a + (e^2*x^2)/f^2])) - (1 + (a*f^2)/d^2)/(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]) - (2*a*f^2*Log[-(e*x) - f*Sqrt[a + (e^2*x^2)/f^2]])/d^3 + (2*a*f^2*Log[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]])/d^3)/(2*e)
```

### Defintions of rubi rules used

rule 1195

```
Int[((d_.) + (e_.)*(x_.))^(m_.)*((f_.) + (g_.)*(x_.))^(n_.)*((a_.) + (b_.)*(x_.) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2542

```
Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.))^(p_.), x_Symbol] := Simp[1/(2*e) Subst[Int[(g + h*x^n)^p*((d^2 + a*f^2 - 2*d*x + x^2)/(d - x)^2), x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]
```

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 3181 vs.  $2(139) = 278$ .

Time = 0.08 (sec) , antiderivative size = 3182, normalized size of antiderivative = 21.07

method	result	size
default	Expression too large to display	3182

input `int(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^2,x,method=_RETURNVERBOSE)`

output

```
1/2*d/(a*f^2-2*d*e*x-d^2)/e+1/2*f^2*a/(a*f^2-2*d*e*x-d^2)/d/e+2*e^2*(1/4*x
/d^2/e^2-1/8*(a^2*f^4-2*a*d^2*f^2+d^4)/e^3/d^3/(-a*f^2+2*d*e*x+d^2)+1/4/d^
3/e^3*(a*f^2-d^2)*ln(-a*f^2+2*d*e*x+d^2))+2*d*e*(-1/4*(a*f^2-d^2)/(-a*f^2+
2*d*e*x+d^2)/e^2/d^2+1/4/d^2/e^2*ln(-a*f^2+2*d*e*x+d^2))-1/2/d*f/e^2*(-4/(
a^2*f^4+2*a*d^2*f^2+d^4)*d^2*f^2/(x+1/2*(-a*f^2+d^2)/d/e)*(e^2*(x+1/2*(-a*
f^2+d^2)/d/e)^2/f^2+e*(a*f^2-d^2)/f^2/d*(x+1/2*(-a*f^2+d^2)/d/e)+1/4*(a^2*
f^4+2*a*d^2*f^2+d^4)/d^2/f^2)^(3/2)+2*e*(a*f^2-d^2)*d/(a^2*f^4+2*a*d^2*f^2
+d^4)*(1/2*(4*e^2*(x+1/2*(-a*f^2+d^2)/d/e)^2/f^2+4*e*(a*f^2-d^2)/f^2/d*(x+
1/2*(-a*f^2+d^2)/d/e)+(a^2*f^4+2*a*d^2*f^2+d^4)/d^2/f^2)^(1/2)+1/2*e*(a*f^
2-d^2)/f^2/d*ln((1/2*e*(a*f^2-d^2)/f^2/d+e^2/f^2*(x+1/2*(-a*f^2+d^2)/d/e))
/(e^2/f^2)^(1/2)+(e^2*(x+1/2*(-a*f^2+d^2)/d/e)^2/f^2+e*(a*f^2-d^2)/f^2/d*(
x+1/2*(-a*f^2+d^2)/d/e)+1/4*(a^2*f^4+2*a*d^2*f^2+d^4)/d^2/f^2)^(1/2))/(e^2
/f^2)^(1/2)-1/2*(a^2*f^4+2*a*d^2*f^2+d^4)/d^2/f^2/((a^2*f^4+2*a*d^2*f^2+d^
4)/d^2/f^2)^(1/2)*ln((1/2*(a^2*f^4+2*a*d^2*f^2+d^4)/d^2/f^2+e*(a*f^2-d^2)/
f^2/d*(x+1/2*(-a*f^2+d^2)/d/e)+1/2*((a^2*f^4+2*a*d^2*f^2+d^4)/d^2/f^2)^(1/
2)*(4*e^2*(x+1/2*(-a*f^2+d^2)/d/e)^2/f^2+4*e*(a*f^2-d^2)/f^2/d*(x+1/2*(-a*
f^2+d^2)/d/e)+(a^2*f^4+2*a*d^2*f^2+d^4)/d^2/f^2)^(1/2))/(x+1/2*(-a*f^2+d^2
)/d/e))+8*e^2/(a^2*f^4+2*a*d^2*f^2+d^4)*d^2*(1/4*(2*e^2/f^2*(x+1/2*(-a*f^
2+d^2)/d/e)+e*(a*f^2-d^2)/f^2/d)/e^2*f^2*(e^2*(x+1/2*(-a*f^2+d^2)/d/e)^2/f
^2+e*(a*f^2-d^2)/f^2/d*(x+1/2*(-a*f^2+d^2)/d/e)+1/4*(a^2*f^4+2*a*d^2*f^...
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 284 vs.  $2(139) = 278$ .

Time = 0.13 (sec) , antiderivative size = 284, normalized size of antiderivative = 1.88

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^2} dx$$

$$= \frac{a^2f^4 - 2d^2e^2x^2 + ad^2f^2 - 2d^3ex + (a^2f^4 - 2adef^2x - ad^2f^2) \log\left(-aef^2x + 2de^2x^2 + adf^2 + (af^3 - \dots\right)}{\dots}$$

input `integrate(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^2,x, algorithm="fricas")`

output `1/2*(a^2*f^4 - 2*d^2*e^2*x^2 + a*d^2*f^2 - 2*d^3*e*x + (a^2*f^4 - 2*a*d*e*f^2*x - a*d^2*f^2)*log(-a*e*f^2*x + 2*d*e^2*x^2 + a*d*f^2 + (a*f^3 - 2*d*e*f*x)*sqrt((e^2*x^2 + a*f^2)/f^2)) + (a^2*f^4 - 2*a*d*e*f^2*x - a*d^2*f^2)*log(-a*f^2 + 2*d*e*x + d^2) - (a^2*f^4 - 2*a*d*e*f^2*x - a*d^2*f^2)*log(-e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) - d) - 2*(a*d*f^3 - d^2*e*f*x)*sqrt((e^2*x^2 + a*f^2)/f^2))/(a*d^3*e*f^2 - 2*d^4*e^2*x - d^5*e)`

**Sympy [F]**

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^2} dx = \int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^2} dx$$

input `integrate(1/(d+e*x+f*(a+e**2*x**2/f**2)**(1/2))**2,x)`

output `Integral((d + e*x + f*sqrt(a + e**2*x**2/f**2))**(-2), x)`



**Maxima [F]**

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^2} dx = \int \frac{1}{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + af + d}\right)^2} dx$$

input `integrate(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^2,x, algorithm="maxima")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a)*f + d)^(-2), x)`

**Giac [A] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.97

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^2} dx = \frac{af^2 \log(|-af^2 + 2dex + d^2|)}{2d^3e} + \frac{af|f| \log(|-x|e| + \sqrt{e^2x^2 + af^2}|)}{2d^3|e|} + \frac{x}{2d^2} - \frac{\sqrt{e^2x^2 + af^2}|f|}{2d^2ef} + \frac{a^2f^4 + 2ad^2f^2 + d^4}{4(af^2 - 2dex - d^2)d^3e}$$

input `integrate(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^2,x, algorithm="giac")`

output `1/2*a*f^2*log(abs(-a*f^2 + 2*d*e*x + d^2))/(d^3*e) + 1/2*a*f*abs(f)*log(abs(-x*abs(e) + sqrt(e^2*x^2 + a*f^2)))/(d^3*abs(e)) + 1/2*x/d^2 - 1/2*sqrt(e^2*x^2 + a*f^2)*abs(f)/(d^2*e*f) + 1/4*(a^2*f^4 + 2*a*d^2*f^2 + d^4)/((a*f^2 - 2*d*e*x - d^2)*d^3*e)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^2} dx = \int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^2} dx$$

input `int(1/(d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^2,x)`output `int(1/(d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^2, x)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 365, normalized size of antiderivative = 2.42

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^2} dx$$

$$= \frac{-\sqrt{e^2x^2 + af^2} a^2 d f^4 + \sqrt{e^2x^2 + af^2} a d^3 f^2 + \sqrt{e^2x^2 + af^2} a d^2 e f^2 x - \sqrt{e^2x^2 + af^2} d^4 e x + \log(\sqrt{e^2x^2 + af^2})}{\dots}$$

input `int(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^2,x)`output `( - sqrt(a*f**2 + e**2*x**2)*a**2*d*f**4 + sqrt(a*f**2 + e**2*x**2)*a*d**3*f**2 + sqrt(a*f**2 + e**2*x**2)*a*d**2*e*f**2*x - sqrt(a*f**2 + e**2*x**2)*d**4*e*x + log(sqrt(a*f**2 + e**2*x**2)*d + a*f**2 - d*e*x)*a**3*f**6 - 2*log(sqrt(a*f**2 + e**2*x**2)*d + a*f**2 - d*e*x)*a**2*d**2*f**4 - 2*log(sqrt(a*f**2 + e**2*x**2)*d + a*f**2 - d*e*x)*a**2*d*e*f**4*x + log(sqrt(a*f**2 + e**2*x**2)*d + a*f**2 - d*e*x)*a*d**4*f**2 + 2*log(sqrt(a*f**2 + e**2*x**2)*d + a*f**2 - d*e*x)*a*d**3*e*f**2*x + a**2*d*e*f**4*x - a*d**2*e**2*f**2*x**2 + d**5*e*x + d**4*e**2*x**2)/(d**3*e*(a**2*f**4 - 2*a*d**2*f**2 - 2*a*d*e*f**2*x + d**4 + 2*d**3*e*x))`

$$3.7 \quad \int \frac{1}{\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)^3} dx$$

Optimal result . . . . .	90
Mathematica [A] (verified) . . . . .	91
Rubi [A] (verified) . . . . .	91
Maple [B] (verified) . . . . .	93
Fricas [B] (verification not implemented) . . . . .	93
Sympy [F] . . . . .	94
Maxima [F] . . . . .	94
Giac [A] (verification not implemented) . . . . .	95
Mupad [F(-1)] . . . . .	95
Reduce [B] (verification not implemented) . . . . .	96

**Optimal result**

Integrand size = 25, antiderivative size = 193

$$\int \frac{1}{\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)^3} dx = -\frac{af^2}{2d^3e\left(ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)} - \frac{1+\frac{af^2}{d^2}}{4e\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)^2} - \frac{af^2}{d^3e\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)} - \frac{3af^2\log\left(ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)}{2d^4e} + \frac{3af^2\log\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)}{2d^4e}$$

output

```
-1/2*a*f^2/d^3/e/(e*x+f*(a+e^2*x^2/f^2)^(1/2))-1/4*(1+a*f^2/d^2)/e/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^2-a*f^2/d^3/e/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))-3/2*a*f^2*ln(e*x+f*(a+e^2*x^2/f^2)^(1/2))/d^4/e+3/2*a*f^2*ln(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))/d^4/e
```

**Mathematica [A] (verified)**

Time = 2.85 (sec) , antiderivative size = 278, normalized size of antiderivative = 1.44

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^3} dx = \frac{d\sqrt{a + \frac{e^2x^2}{f^2}}(3a^2f^5 + d^2efx(3d + 4ex) - adf^3(5d + 9ex))}{(d^2 - af^2 + 2dex)^2} + \frac{d(2d^5 + 6d^4ex - 3a^2ef^4x + 3d^3e^2x^2 + 9ade^2f^2x^2 + d^2(3aef^2x - 4e^3x^3))}{(d^2 - af^2 + 2dex)^2} + 3af^2$$

input `Integrate[(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(-3),x]`

output 
$$\frac{-1/2*((d*\text{Sqrt}[a + (e^2*x^2)/f^2]*(3*a^2*f^5 + d^2*e*f*x*(3*d + 4*e*x) - a*d*f^3*(5*d + 9*e*x)))/(d^2 - a*f^2 + 2*d*e*x)^2 + (d*(2*d^5 + 6*d^4*e*x - 3*a^2*e*f^4*x + 3*d^3*e^2*x^2 + 9*a*d*e^2*f^2*x^2 + d^2*(3*a*e*f^2*x - 4*e^3*x^3)))/(d^2 - a*f^2 + 2*d*e*x)^2 + 3*a*f^2*\text{Log}[-(\text{Sqrt}[a]*f) + e*x + f*\text{Sqrt}[a + (e^2*x^2)/f^2]] - 3*a*f^2*\text{Log}[-(a*f^2) + d*(e*x - f*\text{Sqrt}[a + (e^2*x^2)/f^2])] + \text{Sqrt}[a]*f*(d + e*x + f*\text{Sqrt}[a + (e^2*x^2)/f^2]))/(d^4*e)}$$

**Rubi [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.94, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$ , Rules used = {2542, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\left(f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex\right)^3} dx$$

↓ 2542

$$\int \frac{d^2 - 2\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)d + af^2 + \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)^2}{\left(-\sqrt{\frac{e^2x^2}{f^2} + a}f - ex\right)^2 \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)^3} d\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)$$

2e

↓ 1195

$$\int \left( \frac{3af^2}{d^4 \left( -\sqrt{\frac{e^2x^2}{f^2} + af - ex} \right)} + \frac{3af^2}{d^4 \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + a} \right)} + \frac{af^2}{d^3 \left( -\sqrt{\frac{e^2x^2}{f^2} + af - ex} \right)^2} + \frac{2af^2}{d^3 \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + a} \right)^2} + \frac{d^2+af^2}{d^2 \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + a} \right)} \right) dx$$

↓ 2009

$$-\frac{3af^2 \log \left( f \left( -\sqrt{a + \frac{e^2x^2}{f^2}} \right) - ex \right)}{d^4} + \frac{3af^2 \log \left( f \sqrt{a + \frac{e^2x^2}{f^2}} + d + ex \right)}{d^4} + \frac{af^2}{d^3 \left( f \left( -\sqrt{a + \frac{e^2x^2}{f^2}} \right) - ex \right)} - \frac{2af^2}{d^3 \left( f \sqrt{a + \frac{e^2x^2}{f^2}} + d + ex \right)} - \frac{2 \left( f \sqrt{a + \frac{e^2x^2}{f^2}} + d + ex \right)}{2e}$$

input `Int[(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(-3),x]`

output `((a*f^2)/(d^3*(-(e*x) - f*Sqrt[a + (e^2*x^2)/f^2])) - (1 + (a*f^2)/d^2)/(2*(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^2) - (2*a*f^2)/(d^3*(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])) - (3*a*f^2*Log[-(e*x) - f*Sqrt[a + (e^2*x^2)/f^2]])/d^4 + (3*a*f^2*Log[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]])/d^4)/(2*e)`

### Defintions of rubi rules used

rule 1195 `Int[((d_.) + (e_.)*(x_.))^(m_.)*((f_.) + (g_.)*(x_.))^(n_.)*((a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2542 `Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_.) + (f_.)*Sqrt[(a_.) + (c_.)*(x_.)^2]))^(n_.)^(p_.), x_Symbol] := Simp[1/(2*e) Subst[Int[(g + h*x^n)^p*((d^2 + a*f^2 - 2*d*x + x^2)/(d - x)^2), x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]`

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 11351 vs.  $2(175) = 350$ .

Time = 0.12 (sec) , antiderivative size = 11352, normalized size of antiderivative = 58.82

method	result	size
default	Expression too large to display	11352

input `int(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^3,x,method=_RETURNVERBOSE)`

output `result too large to display`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 536 vs.  $2(175) = 350$ .

Time = 0.27 (sec) , antiderivative size = 536, normalized size of antiderivative = 2.78

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^3} dx$$

$$= \frac{5a^3f^6 + 8d^3e^3x^3 - 6a^2d^2f^4 - 3ad^4f^2 + 2(ad^2e^2f^2 + 5d^4e^2)x^2 - 2(7a^2def^4 + ad^3ef^2 - 2d^5e)x + 3(\dots)}{\dots}$$

input `integrate(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^3,x, algorithm="fricas")`

output

```

1/4*(5*a^3*f^6 + 8*d^3*e^3*x^3 - 6*a^2*d^2*f^4 - 3*a*d^4*f^2 + 2*(a*d^2*e^
2*f^2 + 5*d^4*e^2)*x^2 - 2*(7*a^2*d*e*f^4 + a*d^3*e*f^2 - 2*d^5*e)*x + 3*(
a^3*f^6 + 4*a*d^2*e^2*f^2*x^2 - 2*a^2*d^2*f^4 + a*d^4*f^2 - 4*(a^2*d*e*f^4
- a*d^3*e*f^2)*x)*log(-a*e*f^2*x + 2*d*e^2*x^2 + a*d*f^2 + (a*f^3 - 2*d*e
*f*x)*sqrt((e^2*x^2 + a*f^2)/f^2)) + 3*(a^3*f^6 + 4*a*d^2*e^2*f^2*x^2 - 2*
a^2*d^2*f^4 + a*d^4*f^2 - 4*(a^2*d*e*f^4 - a*d^3*e*f^2)*x)*log(-a*f^2 + 2*
d*e*x + d^2) - 3*(a^3*f^6 + 4*a*d^2*e^2*f^2*x^2 - 2*a^2*d^2*f^4 + a*d^4*f^
2 - 4*(a^2*d*e*f^4 - a*d^3*e*f^2)*x)*log(-e*x + f*sqrt((e^2*x^2 + a*f^2)/f
^2) - d) - 2*(3*a^2*d*f^5 + 4*d^3*e^2*f*x^2 - 5*a*d^3*f^3 - 3*(3*a*d^2*e*f
^3 - d^4*e*f)*x)*sqrt((e^2*x^2 + a*f^2)/f^2))/(a^2*d^4*e*f^4 + 4*d^6*e^3*x
^2 - 2*a*d^6*e*f^2 + d^8*e - 4*(a*d^5*e^2*f^2 - d^7*e^2)*x)

```

**Sympy [F]**

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^3} dx = \int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^3} dx$$

input

```
integrate(1/(d+e*x+f*(a+e**2*x**2/f**2)**(1/2))**3,x)
```

output

```
Integral((d + e*x + f*sqrt(a + e**2*x**2/f**2))**(-3), x)
```

**Maxima [F]**

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^3} dx = \int \frac{1}{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + af + d}\right)^3} dx$$

input

```
integrate(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^3,x, algorithm="maxima")
```

output

```
integrate((e*x + sqrt(e^2*x^2/f^2 + a)*f + d)^(-3), x)
```

**Giac [A] (verification not implemented)**

Time = 0.68 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.94

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^3} dx$$

$$= \frac{3af^2 \log(|-af^2 + 2dex + d^2|)}{4d^4e} + \frac{3af|f| \log(|-x|e| + \sqrt{e^2x^2 + af^2|})}{4d^4|e|} + \frac{x}{2d^3}$$

$$- \frac{\sqrt{e^2x^2 + af^2}|f|}{2d^3ef} + \frac{5a^3f^6 - 3a^2d^2f^4 - 9ad^4f^2 - d^6 - 12(a^2def^4 + ad^3ef^2)x}{8(af^2 - 2dex - d^2)^2d^4e}$$

input `integrate(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^3,x, algorithm="giac")`

output `3/4*a*f^2*log(abs(-a*f^2 + 2*d*e*x + d^2))/(d^4*e) + 3/4*a*f*abs(f)*log(abs(-x*abs(e) + sqrt(e^2*x^2 + a*f^2)))/(d^4*abs(e)) + 1/2*x/d^3 - 1/2*sqrt(e^2*x^2 + a*f^2)*abs(f)/(d^3*e*f) + 1/8*(5*a^3*f^6 - 3*a^2*d^2*f^4 - 9*a*d^4*f^2 - d^6 - 12*(a^2*d*e*f^4 + a*d^3*e*f^2)*x)/((a*f^2 - 2*d*e*x - d^2)^2*d^4*e)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^3} dx = \int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^3} dx$$

input `int(1/(d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^3,x)`

output `int(1/(d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^3, x)`



**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 767, normalized size of antiderivative = 3.97

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^3} dx$$

$$= \frac{12 \log(\sqrt{e^2x^2 + a} f^2 d + a f^2 - dex) a^4 f^8 - 9a^3 d^2 f^6 + 3a^2 d^4 f^4 - 12d^6 e^2 x^2 - 16d^5 e^3 x^3 + 3a^4 f^8 - 12\sqrt{e^2x^2 + a}}{\dots}$$

input `int(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^3,x)`

output

```
( - 12*sqrt(a*f**2 + e**2*x**2)*a**3*d*f**6 + 32*sqrt(a*f**2 + e**2*x**2)*
a**2*d**3*f**4 + 36*sqrt(a*f**2 + e**2*x**2)*a**2*d**2*e*f**4*x - 20*sqrt(
a*f**2 + e**2*x**2)*a*d**5*f**2 - 48*sqrt(a*f**2 + e**2*x**2)*a*d**4*e*f**
2*x - 16*sqrt(a*f**2 + e**2*x**2)*a*d**3*e**2*f**2*x**2 + 12*sqrt(a*f**2 +
e**2*x**2)*d**6*e*x + 16*sqrt(a*f**2 + e**2*x**2)*d**5*e**2*x**2 + 12*log
(sqrt(a*f**2 + e**2*x**2)*d + a*f**2 - d*e*x)*a**4*f**8 - 36*log(sqrt(a*f*
*2 + e**2*x**2)*d + a*f**2 - d*e*x)*a**3*d**2*f**6 - 48*log(sqrt(a*f**2 +
e**2*x**2)*d + a*f**2 - d*e*x)*a**3*d*e*f**6*x + 36*log(sqrt(a*f**2 + e**2
*x**2)*d + a*f**2 - d*e*x)*a**2*d**4*f**4 + 96*log(sqrt(a*f**2 + e**2*x**2
)*d + a*f**2 - d*e*x)*a**2*d**3*e*f**4*x + 48*log(sqrt(a*f**2 + e**2*x**2)
*d + a*f**2 - d*e*x)*a**2*d**2*e**2*f**4*x**2 - 12*log(sqrt(a*f**2 + e**2*
x**2)*d + a*f**2 - d*e*x)*a*d**6*f**2 - 48*log(sqrt(a*f**2 + e**2*x**2)*d
+ a*f**2 - d*e*x)*a*d**5*e*f**2*x - 48*log(sqrt(a*f**2 + e**2*x**2)*d + a
f**2 - d*e*x)*a*d**4*e**2*f**2*x**2 + 3*a**4*f**8 - 9*a**3*d**2*f**6 + 3*a
**2*d**4*f**4 - 24*a**2*d**2*e**2*f**4*x**2 + a*d**6*f**2 + 12*a*d**4*e**2
*f**2*x**2 + 16*a*d**3*e**3*f**2*x**3 + 2*d**8 - 12*d**6*e**2*x**2 - 16*d*
*5*e**3*x**3)/(8*d**4*e*(a**3*f**6 - 3*a**2*d**2*f**4 - 4*a**2*d*e*f**4*x
+ 3*a*d**4*f**2 + 8*a*d**3*e*f**2*x + 4*a*d**2*e**2*f**2*x**2 - d**6 - 4*d
**5*e*x - 4*d**4*e**2*x**2))
```

**3.8**  $\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx$

Optimal result	97
Mathematica [A] (verified)	98
Rubi [A] (verified)	98
Maple [F]	101
Fricas [A] (verification not implemented)	101
Sympy [F]	102
Maxima [F]	102
Giac [F]	103
Mupad [F(-1)]	103
Reduce [F]	103

**Optimal result**

Integrand size = 27, antiderivative size = 225

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx = \frac{2adf^2 \sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}}}{e} - \frac{ad^2 f^2 \sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}}}{2e \left( ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)} + \frac{af^2 \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{3/2}}{3e} + \frac{\left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{7/2}}{7e} - \frac{5ad^{3/2} f^2 \operatorname{arctanh} \left( \frac{\sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}}}{\sqrt{d}} \right)}{2e}$$

output

```
2*a*d*f^2*(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2)/e-1/2*a*d^2*f^2*(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2)/e/(e*x+f*(a+e^2*x^2/f^2)^(1/2))+1/3*a*f^2*(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(3/2)/e+1/7*(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(7/2)/e-5/2*a*d^(3/2)*f^2*arctanh((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2)/d^(1/2))/e
```

**Mathematica [A] (verified)**

Time = 1.80 (sec) , antiderivative size = 212, normalized size of antiderivative = 0.94

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx = \frac{\sqrt{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}} \left( 20a^2f^4+6(d+2ex)^3 \left( ex+f\sqrt{a+\frac{e^2x^2}{f^2}} \right) + af^2 \left( -3d^2+4ex \left( 19ex+13f\sqrt{a+\frac{e^2x^2}{f^2}} \right) + 4d \left( ex+f\sqrt{a+\frac{e^2x^2}{f^2}} \right) \right) \right)}{42e}$$

input `Integrate[(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(5/2),x]`

output `((Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]]*(20*a^2*f^4 + 6*(d + 2*e*x)^3*(e*x + f*Sqrt[a + (e^2*x^2)/f^2]) + a*f^2*(-3*d^2 + 4*e*x*(19*e*x + 13*f*Sqrt[a + (e^2*x^2)/f^2]) + 4*d*(38*e*x + 29*f*Sqrt[a + (e^2*x^2)/f^2])))/(e*x + f*Sqrt[a + (e^2*x^2)/f^2]) - 105*a*d^(3/2)*f^2*ArcTanh[Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]]/Sqrt[d]])/(42*e)`

**Rubi [A] (verified)**

Time = 0.42 (sec) , antiderivative size = 219, normalized size of antiderivative = 0.97, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {2542, 1192, 1580, 25, 2341, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( f \sqrt{a + \frac{e^2 x^2}{f^2}} + d + ex \right)^{5/2} dx$$

↓ 2542

$$\int \frac{\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^{5/2} \left(d^2-2\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)d+af^2+\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^2\right)}{\left(-\sqrt{\frac{e^2x^2}{f^2}+af-ex}\right)^2} d\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)$$

---

2e

↓ 1192

$$\int \frac{\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^3 \left(d^2-2\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)d+af^2+\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^2\right)}{\left(-\sqrt{\frac{e^2x^2}{f^2}+af-ex}\right)^2} d\sqrt{d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}}$$

---

e

↓ 1580

$$\frac{1}{2} \int -\frac{2\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^4 - 2d\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^3 + 2af^2\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^2 + 2adf^2\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right) + ad^2f^2}{-\sqrt{\frac{e^2x^2}{f^2}+af-ex}} d\sqrt{d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}}$$

---

e

↓ 25

$$\frac{ad^2f^2\sqrt{f\sqrt{a+\frac{e^2x^2}{f^2}+d+ex}}}{2\left(f\left(-\sqrt{a+\frac{e^2x^2}{f^2}}\right)-ex\right)} - \frac{1}{2} \int \frac{2\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^4 - 2d\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^3 + 2af^2\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^2 + 2adf^2\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)}{-\sqrt{\frac{e^2x^2}{f^2}+af-ex}}$$

---

e

↓ 2341

$$\frac{ad^2f^2\sqrt{f\sqrt{a+\frac{e^2x^2}{f^2}+d+ex}}}{2\left(f\left(-\sqrt{a+\frac{e^2x^2}{f^2}}\right)-ex\right)} - \frac{1}{2} \int \left( -2\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^3 - 2af^2\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right) - 4adf^2 + \frac{5ad^2f}{-\sqrt{\frac{e^2x^2}{f^2}+a}} \right)$$

---

e

↓ 2009

$$\frac{1}{2} \left( -5ad^{3/2}f^2 \operatorname{arctanh} \left( \frac{\sqrt{f\sqrt{a+\frac{e^2x^2}{f^2}+d+ex}}}{\sqrt{d}} \right) + \frac{2}{7} \left( f\sqrt{a+\frac{e^2x^2}{f^2}+d+ex} \right)^{7/2} + \frac{2}{3} af^2 \left( f\sqrt{a+\frac{e^2x^2}{f^2}+d+ex} \right)^{3/2} + \dots \right)$$

---

e

input `Int[(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(5/2),x]`

output `((a*d^2*f^2*Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]])/(2*(-(e*x) - f*Sqrt[a + (e^2*x^2)/f^2])) + (4*a*d*f^2*Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]]) + (2*a*f^2*(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(3/2))/3 + (2*(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(7/2))/7 - 5*a*d^(3/2)*f^2*ArcTanh[Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]]/Sqrt[d]])/2)/e`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 1192 `Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))^(n_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Simp[2/e^(n + 2*p + 1) Subst[Int[x^(2*m + 1)*(e*f - d*g + g*x^2)^n*(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 + c*x^4)^p, x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && IGtQ[p, 0] && ILtQ[n, 0] && IntegerQ[m + 1/2]`

rule 1580 `Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := Simp[(-d)^(m/2 - 1)*(c*d^2 - b*d*e + a*e^2)^p*x*((d + e*x^2)^(q + 1)/(2*e^(2*p + m/2)*(q + 1))), x] + Simp[1/(2*e^(2*p + m/2)*(q + 1)) Int[(d + e*x^2)^(q + 1)*ExpandToSum[Together[(1/(d + e*x^2))*(2*e^(2*p + m/2)*(q + 1)*x^m*(a + b*x^2 + c*x^4)^p - (-d)^(m/2 - 1)*(c*d^2 - b*d*e + a*e^2)^p*(d + e*(2*q + 3)*x^2)]], x], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && ILtQ[q, -1] && IGtQ[m/2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2341 `Int[(Pq_)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[Pq*(a + b*x^2)^p, x], x] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

rule 2542

```
Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(
n_))^(p_.), x_Symbol] := Simp[1/(2*e) Subst[Int[(g + h*x^n)^p*((d^2 + a*f
^2 - 2*d*x + x^2)/(d - x)^2), x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; Fr
eeQ[{a, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]
```

**Maple [F]**

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{\frac{5}{2}} dx$$

input

```
int((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(5/2),x)
```

output

```
int((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(5/2),x)
```

**Fricas [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 467, normalized size of antiderivative = 2.08

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{\frac{5}{2}} dx = \frac{105 ad^{\frac{3}{2}} f^2 \log \left( af^2 - 2dex + 2df \sqrt{\frac{e^2 x^2 + af^2}{f^2 - d}} + 2 \left( \sqrt{d}ex - \sqrt{d}f \sqrt{\frac{e^2 x^2 + af^2}{f^2 - d}} \right) \sqrt{e} \right)}{105 a \sqrt{-d} df^2 \arctan \left( \frac{\sqrt{ex + f \sqrt{\frac{e^2 x^2 + af^2}{f^2 - d}} + d} \left( \sqrt{-d}f \sqrt{\frac{e^2 x^2 + af^2}{f^2 - d}} - (ex + d)\sqrt{-d} \right)}{af^2 - 2dex - d^2} \right)} - (24 e^3 x^3 + 36 de^2 x^2 + 116 adf$$

input `integrate((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(5/2),x, algorithm="fricas")`

output `[1/84*(105*a*d^(3/2)*f^2*log(a*f^2 - 2*d*e*x + 2*d*f*sqrt((e^2*x^2 + a*f^2)/f^2) + 2*(sqrt(d)*e*x - sqrt(d)*f*sqrt((e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d) + 2*(24*e^3*x^3 + 36*d*e^2*x^2 + 116*a*d*f^2 + 6*d^3 + (32*a*e*f^2 + 39*d^2*e)*x + (24*e^2*f*x^2 + 20*a*f^3 + 36*d*e*f*x - 3*d^2*f)*sqrt((e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d))/e, -1/42*(105*a*sqrt(-d)*d*f^2*arctan(-sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d)*(sqrt(-d)*f*sqrt((e^2*x^2 + a*f^2)/f^2) - (e*x + d)*sqrt(-d))/(a*f^2 - 2*d*e*x - d^2)) - (24*e^3*x^3 + 36*d*e^2*x^2 + 116*a*d*f^2 + 6*d^3 + (32*a*e*f^2 + 39*d^2*e)*x + (24*e^2*f*x^2 + 20*a*f^3 + 36*d*e*f*x - 3*d^2*f)*sqrt((e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d))/e]`

### Sympy [F]

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx = \int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{\frac{5}{2}} dx$$

input `integrate((d+e*x+f*(a+e**2*x**2/f**2)**(1/2))**(5/2),x)`

output `Integral((d + e*x + f*sqrt(a + e**2*x**2/f**2))**(5/2), x)`

### Maxima [F]

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx = \int \left( ex + \sqrt{\frac{e^2 x^2}{f^2} + af + d} \right)^{\frac{5}{2}} dx$$

input `integrate((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(5/2),x, algorithm="maxima")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a)*f + d)^(5/2), x)`

**Giac [F]**

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx = \int \left( ex + \sqrt{\frac{e^2 x^2}{f^2} + af + d} \right)^{5/2} dx$$

input `integrate((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(5/2),x, algorithm="giac")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a)*f + d)^(5/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx = \int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx$$

input `int((d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^(5/2),x)`

output `int((d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^(5/2), x)`

**Reduce [F]**

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx = \int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{5/2} dx$$

input `int((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(5/2),x)`

output `int((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(5/2),x)`



**3.9**  $\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx$

Optimal result	104
Mathematica [A] (verified)	105
Rubi [A] (verified)	105
Maple [F]	107
Fricas [A] (verification not implemented)	108
Sympy [F]	109
Maxima [F]	109
Giac [F]	109
Mupad [F(-1)]	110
Reduce [F]	110

**Optimal result**

Integrand size = 27, antiderivative size = 183

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx = \frac{af^2 \sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}}}{e} - \frac{adf^2 \sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}}}{2e \left( ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)} + \frac{\left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{5/2}}{5e} - \frac{3a\sqrt{d}f^2 \operatorname{arctanh} \left( \frac{\sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}}}{\sqrt{d}} \right)}{2e}$$

output

```
a*f^2*(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2)/e-1/2*a*d*f^2*(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2)/e/(e*x+f*(a+e^2*x^2/f^2)^(1/2))+1/5*(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(5/2)/e-3/2*a*d^(1/2)*f^2*arctanh((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2)/d^(1/2))/e
```

### Mathematica [A] (verified)

Time = 1.56 (sec) , antiderivative size = 170, normalized size of antiderivative = 0.93

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx = \frac{\sqrt{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}} \left( 2(d+2ex)^2 \left( ex+f\sqrt{a+\frac{e^2x^2}{f^2}} \right) + af^2 \left( -d+16ex+12f\sqrt{a+\frac{e^2x^2}{f^2}} \right) \right)}{ex+f\sqrt{a+\frac{e^2x^2}{f^2}}} - 15a\sqrt{d}f^2 \operatorname{arctanh} \left( \frac{\sqrt{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}}}{\sqrt{d}} \right)}{10e}$$

input

```
Integrate[(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(3/2), x]
```

output

```
((Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]])*(2*(d + 2*e*x)^2*(e*x + f*Sqrt[a + (e^2*x^2)/f^2]) + a*f^2*(-d + 16*e*x + 12*f*Sqrt[a + (e^2*x^2)/f^2]))/(e*x + f*Sqrt[a + (e^2*x^2)/f^2]) - 15*a*Sqrt[d]*f^2*ArcTanh[Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]]/Sqrt[d]]/(10*e)
```

### Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 181, normalized size of antiderivative = 0.99, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$ , Rules used = {2542, 1192, 1580, 2341, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( f \sqrt{a + \frac{e^2 x^2}{f^2}} + d + ex \right)^{3/2} dx$$

↓ 2542

$$\int \frac{\left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right)^{3/2} \left( d^2-2\left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right) d+af^2+\left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right)^2 \right)}{\left( -\sqrt{\frac{e^2x^2}{f^2}+a} f-ex \right)^2} d \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right)}{2e}$$

↓ 1192

$$\int \frac{\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^2 \left(d^2-2\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)d+af^2+\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^2\right)}{\left(-\sqrt{\frac{e^2x^2}{f^2}+a}f-ex\right)^2} d\sqrt{d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}}$$

e  
↓ 1580

$$\frac{adf^2\sqrt{f\sqrt{a+\frac{e^2x^2}{f^2}}+d+ex}}{2\left(f\left(-\sqrt{a+\frac{e^2x^2}{f^2}}\right)-ex\right)} - \frac{1}{2} \int \frac{2\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^3 - 2d\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^2 + 2af^2\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right) + adf^2}{-\sqrt{\frac{e^2x^2}{f^2}+a}f-ex} d\sqrt{d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}}$$

e  
↓ 2341

$$\frac{adf^2\sqrt{f\sqrt{a+\frac{e^2x^2}{f^2}}+d+ex}}{2\left(f\left(-\sqrt{a+\frac{e^2x^2}{f^2}}\right)-ex\right)} - \frac{1}{2} \int \left(-2af^2 + \frac{3adf^2}{-\sqrt{\frac{e^2x^2}{f^2}+a}f-ex} - 2\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^2\right) d\sqrt{d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}}$$

e  
↓ 2009

$$\frac{1}{2} \left( -3a\sqrt{d}f^2 \operatorname{arctanh}\left(\frac{\sqrt{f\sqrt{a+\frac{e^2x^2}{f^2}}+d+ex}}{\sqrt{d}}\right) + \frac{2}{5}\left(f\sqrt{a+\frac{e^2x^2}{f^2}}+d+ex\right)^{5/2} + 2af^2\sqrt{f\sqrt{a+\frac{e^2x^2}{f^2}}+d+ex} \right) + \frac{a}{2}$$

input `Int[(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(3/2),x]`

output `((a*d*f^2*Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]])/(2*(-(e*x) - f*Sqrt[a + (e^2*x^2)/f^2])) + (2*a*f^2*Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]] + (2*(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(5/2))/5 - 3*a*Sqrt[d]*f^2*ArcTanh[Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]]/Sqrt[d]])/2)/e`

## Definitions of rubi rules used

rule 1192

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))^(n_)*((a_.) + (b_.)*(x_)
+ (c_.)*(x_)^2)^(p_.), x_Symbol] := Simp[2/e^(n + 2*p + 1) Subst[Int[x^(
2*m + 1)*(e*f - d*g + g*x^2)^n*(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 +
c*x^4)^p, x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, g}, x] &&
IGtQ[p, 0] && ILtQ[n, 0] && IntegerQ[m + 1/2]
```

rule 1580

```
Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)
^4)^(p_.), x_Symbol] := Simp[(-d)^(m/2 - 1)*(c*d^2 - b*d*e + a*e^2)^p*x*((d
+ e*x^2)^(q + 1)/(2*e^(2*p + m/2)*(q + 1))), x] + Simp[1/(2*e^(2*p + m/2)*
(q + 1)) Int[(d + e*x^2)^(q + 1)*ExpandToSum[Together[(1/(d + e*x^2))*(2*
e^(2*p + m/2)*(q + 1)*x^m*(a + b*x^2 + c*x^4)^p - (-d)^(m/2 - 1)*(c*d^2 - b
*d*e + a*e^2)^p*(d + e*(2*q + 3)*x^2)], x], x], x] /; FreeQ[{a, b, c, d, e
}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && ILtQ[q, -1] && IGtQ[m/2, 0]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2341

```
Int[(Pq_)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[Pq*
(a + b*x^2)^p, x], x] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && IGtQ[p, -2]
```

rule 2542

```
Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(
n_))^(p_.), x_Symbol] := Simp[1/(2*e) Subst[Int[(g + h*x^n)^p*((d^2 + a*f
^2 - 2*d*x + x^2)/(d - x)^2), x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; Fr
eeQ[{a, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]
```

## Maple [F]

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{\frac{3}{2}} dx$$

input

```
int((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(3/2),x)
```

output `int((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(3/2),x)`

### Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 388, normalized size of antiderivative = 2.12

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx = \frac{15 a \sqrt{d} f^2 \log \left( a f^2 - 2 d e x + 2 d f \sqrt{\frac{e^2 x^2 + a f^2}{f^2}} + 2 \left( \sqrt{d} e x - \sqrt{d} f \sqrt{\frac{e^2 x^2 + a f^2}{f^2}} \right) \sqrt{e x + f \sqrt{a + \frac{e^2 x^2}{f^2}}} \right)}{10 e} - \frac{15 a \sqrt{-d} f^2 \arctan \left( -\frac{\sqrt{e x + f \sqrt{\frac{e^2 x^2 + a f^2}{f^2}} + d} \left( \sqrt{-d} f \sqrt{\frac{e^2 x^2 + a f^2}{f^2}} - (e x + d) \sqrt{-d} \right)}{a f^2 - 2 d e x - d^2} \right)}{10 e}$$

input `integrate((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(3/2),x, algorithm="fricas")`

output `[1/20*(15*a*sqrt(d)*f^2*log(a*f^2 - 2*d*e*x + 2*d*f*sqrt((e^2*x^2 + a*f^2)/f^2) + 2*(sqrt(d)*e*x - sqrt(d)*f*sqrt((e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d)) + 2*(4*e^2*x^2 + 12*a*f^2 + 9*d*e*x + 2*d^2 + (4*e*f*x - d*f)*sqrt((e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d))/e, -1/10*(15*a*sqrt(-d)*f^2*arctan(-sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d)*(sqrt(-d)*f*sqrt((e^2*x^2 + a*f^2)/f^2) - (e*x + d)*sqrt(-d))/(a*f^2 - 2*d*e*x - d^2)) - (4*e^2*x^2 + 12*a*f^2 + 9*d*e*x + 2*d^2 + (4*e*f*x - d*f)*sqrt((e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d))/e]`

**Sympy [F]**

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx = \int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{\frac{3}{2}} dx$$

input `integrate((d+e*x+f*(a+e**2*x**2/f**2)**(1/2))**(3/2),x)`

output `Integral((d + e*x + f*sqrt(a + e**2*x**2/f**2))**(3/2), x)`

**Maxima [F]**

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx = \int \left( ex + \sqrt{\frac{e^2 x^2}{f^2} + af + d} \right)^{\frac{3}{2}} dx$$

input `integrate((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(3/2),x, algorithm="maxima")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a)*f + d)^(3/2), x)`

**Giac [F]**

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx = \int \left( ex + \sqrt{\frac{e^2 x^2}{f^2} + af + d} \right)^{\frac{3}{2}} dx$$

input `integrate((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(3/2),x, algorithm="giac")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a)*f + d)^(3/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx = \int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx$$

input `int((d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^(3/2),x)`output `int((d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^(3/2), x)`**Reduce [F]**

$$\int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{3/2} dx = \int \left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{\frac{3}{2}} dx$$

input `int((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(3/2),x)`output `int((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(3/2),x)`

**3.10**  $\int \sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}} dx$

Optimal result	111
Mathematica [A] (verified)	112
Rubi [A] (verified)	112
Maple [F]	115
Fricas [A] (verification not implemented)	115
Sympy [F]	116
Maxima [F]	116
Giac [F]	117
Mupad [F(-1)]	117
Reduce [F]	117

**Optimal result**

Integrand size = 27, antiderivative size = 147

$$\int \sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}} dx = -\frac{af^2 \sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}}}{2e \left( ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)} + \frac{\left( d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}} \right)^{3/2}}{3e} - \frac{af^2 \operatorname{arctanh} \left( \frac{\sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}}}{\sqrt{d}} \right)}{2\sqrt{de}}$$

output

```
-1/2*a*f^2*(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2)/e/(e*x+f*(a+e^2*x^2/f^2)^(1/2))+1/3*(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(3/2)/e-1/2*a*f^2*arctanh((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2)/d^(1/2))/d^(1/2)/e
```



**Mathematica [A] (verified)**

Time = 1.12 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.97

$$\int \sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}} dx$$

$$= \frac{\sqrt{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}}\left(-af^2+2(d+2ex)\left(ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)\right)}{ex+f\sqrt{a+\frac{e^2x^2}{f^2}}} - \frac{3af^2\operatorname{arctanh}\left(\frac{\sqrt{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}}}{\sqrt{d}}\right)}{\sqrt{d}}$$

$$= \frac{\dots}{6e}$$

input `Integrate[Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]],x]`

output `((Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]]*(-(a*f^2) + 2*(d + 2*e*x)*(e*x + f*Sqrt[a + (e^2*x^2)/f^2])))/(e*x + f*Sqrt[a + (e^2*x^2)/f^2]) - (3*a*f^2*ArcTanh[Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]]/Sqrt[d]]/Sqrt[d])/(6*e)`

**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {2542, 1192, 1580, 25, 1467, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{f \sqrt{a + \frac{e^2 x^2}{f^2}} + d + ex} dx$$

↓ 2542

$$\int \frac{\sqrt{d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}} \left( d^2 - 2 \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right) d + af^2 + \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right)^2 \right)}{\left( -\sqrt{\frac{e^2x^2}{f^2}+af-ex} \right)^2} d \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right)$$

---

2e

↓ 1192

$$\int \frac{\left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right) \left( d^2 - 2 \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right) d + af^2 + \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right)^2 \right)}{\left( -\sqrt{\frac{e^2x^2}{f^2}+af-ex} \right)^2} d \sqrt{d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}}$$

---

e

↓ 1580

$$\frac{1}{2} \int - \frac{af^2 + 2 \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right)^2 - 2d \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right)}{-\sqrt{\frac{e^2x^2}{f^2}+af-ex}} d \sqrt{d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}} + \frac{af^2 \sqrt{f\sqrt{a+\frac{e^2x^2}{f^2}}+d+ex}}{2 \left( f \left( -\sqrt{a+\frac{e^2x^2}{f^2}} \right) - ex \right)}$$

---

e

↓ 25

$$\frac{af^2 \sqrt{f\sqrt{a+\frac{e^2x^2}{f^2}}+d+ex}}{2 \left( f \left( -\sqrt{a+\frac{e^2x^2}{f^2}} \right) - ex \right)} - \frac{1}{2} \int \frac{af^2 + 2 \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right)^2 - 2d \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right)}{-\sqrt{\frac{e^2x^2}{f^2}+af-ex}} d \sqrt{d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}}$$

---

e

↓ 1467

$$\frac{af^2 \sqrt{f\sqrt{a+\frac{e^2x^2}{f^2}}+d+ex}}{2 \left( f \left( -\sqrt{a+\frac{e^2x^2}{f^2}} \right) - ex \right)} - \frac{1}{2} \int \left( \frac{af^2}{-\sqrt{\frac{e^2x^2}{f^2}+af-ex}} - 2 \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a} \right) \right) d \sqrt{d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}}$$

---

e

↓ 2009

$$\frac{1}{2} \left( \frac{2}{3} \left( f \sqrt{a+\frac{e^2x^2}{f^2}} + d+ex \right)^{3/2} - \frac{af^2 \operatorname{arctanh} \left( \frac{\sqrt{f\sqrt{a+\frac{e^2x^2}{f^2}}+d+ex}}{\sqrt{d}} \right)}{\sqrt{d}} \right) + \frac{af^2 \sqrt{f\sqrt{a+\frac{e^2x^2}{f^2}}+d+ex}}{2 \left( f \left( -\sqrt{a+\frac{e^2x^2}{f^2}} \right) - ex \right)}$$

---

e

input `Int[Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]],x]`

output `((a*f^2*Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]])/(2*(-(e*x) - f*Sqrt[a + (e^2*x^2)/f^2])) + ((2*(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(3/2))/3 - (a*f^2*ArcTanh[Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]]/Sqrt[d]])/Sqrt[d])/2)/e`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 1192 `Int[((d_) + (e_)*(x_)^(m_))*((f_) + (g_)*(x_)^(n_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[2/e^(n + 2*p + 1) Subst[Int[x^(2*m + 1)*(e*f - d*g + g*x^2)^n*(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 + c*x^4)^p, x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && IGtQ[p, 0] && ILtQ[n, 0] && IntegerQ[m + 1/2]`

rule 1467 `Int[((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x^2)^q*(a + b*x^2 + c*x^4)^p, x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && IGtQ[p, 0] && IGtQ[q, -2]`

rule 1580 `Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[(-d)^(m/2 - 1)*(c*d^2 - b*d*e + a*e^2)^p*x*((d + e*x^2)^(q + 1)/(2*e^(2*p + m/2)*(q + 1))), x] + Simp[1/(2*e^(2*p + m/2)*(q + 1)) Int[(d + e*x^2)^(q + 1)*ExpandToSum[Together[(1/(d + e*x^2))*(2*e^(2*p + m/2)*(q + 1)*x^m*(a + b*x^2 + c*x^4)^p - (-d)^(m/2 - 1)*(c*d^2 - b*d*e + a*e^2)^p*(d + e*(2*q + 3)*x^2)], x], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && ILtQ[q, -1] && IGtQ[m/2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2542

```
Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(
n_))^(p_.), x_Symbol] := Simp[1/(2*e) Subst[Int[(g + h*x^n)^p*((d^2 + a*f
^2 - 2*d*x + x^2)/(d - x)^2), x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; Fr
eeQ[{a, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]
```

**Maple [F]**

$$\int \sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}} dx$$

input

```
int((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2),x)
```

output

```
int((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2),x)
```

**Fricas [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 352, normalized size of antiderivative = 2.39

$$\int \sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}} dx$$

$$= \frac{3a\sqrt{d}f^2 \log \left( af^2 - 2dex + 2df \sqrt{\frac{e^2 x^2 + af^2}{f^2}} + 2 \left( \sqrt{d}ex - \sqrt{d}f \sqrt{\frac{e^2 x^2 + af^2}{f^2}} \right) \sqrt{ex + f \sqrt{\frac{e^2 x^2 + af^2}{f^2}} + d} \right) + 3a\sqrt{-d}f^2 \arctan \left( -\frac{\sqrt{ex + f \sqrt{\frac{e^2 x^2 + af^2}{f^2}} + d} \left( \sqrt{-d}f \sqrt{\frac{e^2 x^2 + af^2}{f^2}} - (ex + d)\sqrt{-d} \right)}{af^2 - 2dex - d^2} \right) - \left( 5dex - df \sqrt{\frac{e^2 x^2 + af^2}{f^2}} + 2d^2 \right)}{12de}$$

6de

input `integrate((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2),x, algorithm="fricas")`

output `[1/12*(3*a*sqrt(d)*f^2*log(a*f^2 - 2*d*e*x + 2*d*f*sqrt((e^2*x^2 + a*f^2)/f^2) + 2*(sqrt(d)*e*x - sqrt(d)*f*sqrt((e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d)) + 2*(5*d*e*x - d*f*sqrt((e^2*x^2 + a*f^2)/f^2) + 2*d^2)*sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d)/(d*e), -1/6*(3*a*sqrt(-d)*f^2*arctan(-sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d)*(sqrt(-d)*f*sqrt((e^2*x^2 + a*f^2)/f^2) - (e*x + d)*sqrt(-d))/(a*f^2 - 2*d*e*x - d^2)) - (5*d*e*x - d*f*sqrt((e^2*x^2 + a*f^2)/f^2) + 2*d^2)*sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d)/(d*e)]`

### Sympy [F]

$$\int \sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}} dx = \int \sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}} dx$$

input `integrate((d+e*x+f*(a+e**2*x**2/f**2)**(1/2))**(1/2),x)`

output `Integral(sqrt(d + e*x + f*sqrt(a + e**2*x**2/f**2)), x)`

### Maxima [F]

$$\int \sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}} dx = \int \sqrt{ex + \sqrt{\frac{e^2 x^2}{f^2} + af} + d} dx$$

input `integrate((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(e*x + sqrt(e^2*x^2/f^2 + a)*f + d), x)`

**Giac [F]**

$$\int \sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}} dx = \int \sqrt{ex + \sqrt{\frac{e^2 x^2}{f^2} + af} + d} dx$$

input `integrate((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(e*x + sqrt(e^2*x^2/f^2 + a)*f + d), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}} dx = \int \sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}} dx$$

input `int((d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^(1/2),x)`

output `int((d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^(1/2), x)`

**Reduce [F]**

$$\int \sqrt{d + ex + f \sqrt{a + \frac{e^2 x^2}{f^2}}} dx = \int \sqrt{\sqrt{e^2 x^2 + a f^2} + d + ex} dx$$

input `int((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2),x)`

output `int(sqrt(sqrt(a*f**2 + e**2*x**2) + d + e*x),x)`

**3.11** 
$$\int \frac{1}{\sqrt{d+ex+f}\sqrt{a+\frac{e^2x^2}{f^2}}} dx$$

Optimal result	118
Mathematica [A] (verified)	119
Rubi [A] (verified)	119
Maple [F]	122
Fricas [A] (verification not implemented)	122
Sympy [F]	123
Maxima [F]	123
Giac [F]	123
Mupad [F(-1)]	124
Reduce [F]	124

**Optimal result**

Integrand size = 27, antiderivative size = 147

$$\int \frac{1}{\sqrt{d+ex+f}\sqrt{a+\frac{e^2x^2}{f^2}}} dx = \frac{\sqrt{d+ex+f}\sqrt{a+\frac{e^2x^2}{f^2}}}{e} - \frac{af^2\sqrt{d+ex+f}\sqrt{a+\frac{e^2x^2}{f^2}}}{2de\left(ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)} + \frac{af^2\operatorname{arctanh}\left(\frac{\sqrt{d+ex+f}\sqrt{a+\frac{e^2x^2}{f^2}}}{\sqrt{d}}\right)}{2d^{3/2}e}$$

output

```
(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2)/e-1/2*a*f^2*(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))/d/e/(e*x+f*(a+e^2*x^2/f^2)^(1/2))+1/2*a*f^2*arctanh((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2)/d^(1/2))/d^(3/2)/e
```

### Mathematica [A] (verified)

Time = 1.08 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.96

$$\int \frac{1}{\sqrt{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}}} dx$$

$$= \frac{\sqrt{d}\sqrt{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}}\left(-af^2+2d\left(ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)\right)}{ex+f\sqrt{a+\frac{e^2x^2}{f^2}}} + af^2 \operatorname{arctanh}\left(\frac{\sqrt{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}}}{\sqrt{d}}\right)}{2d^{3/2}e}$$

input `Integrate[1/Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]],x]`

output `((Sqrt[d]*Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]]*(-(a*f^2) + 2*d*(e*x + f*Sqrt[a + (e^2*x^2)/f^2])))/(e*x + f*Sqrt[a + (e^2*x^2)/f^2]) + a*f^2*ArcTanh[Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]]/Sqrt[d]])/(2*d^(3/2)*e)`

### Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.03, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {2542, 1192, 1471, 25, 299, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{f\sqrt{a+\frac{e^2x^2}{f^2}}+d+ex}} dx$$

↓ 2542

$$\int \frac{d^2-2\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)d+af^2+\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)^2}{\left(-\sqrt{\frac{e^2x^2}{f^2}+a}f-ex\right)^2\sqrt{d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}}} d\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+a}\right)$$

2e

↓ 1192



$$\int \frac{d^2 - 2\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)d + af^2 + \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)^2}{\left(-\sqrt{\frac{e^2x^2}{f^2} + a}f - ex\right)^2} d\sqrt{d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}}$$

e  
↓ 1471

$$\frac{af^2\sqrt{f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex}}{2d\left(f\left(-\sqrt{a + \frac{e^2x^2}{f^2}}\right) - ex\right)} - \frac{\int -\frac{2d^2 - 2\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)d + af^2}{-\sqrt{\frac{e^2x^2}{f^2} + a}f - ex} d\sqrt{d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}}}{2d}$$

e  
↓ 25

$$\frac{\int \frac{2d^2 - 2\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)d + af^2}{-\sqrt{\frac{e^2x^2}{f^2} + a}f - ex} d\sqrt{d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}}}{2d} + \frac{af^2\sqrt{f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex}}{2d\left(f\left(-\sqrt{a + \frac{e^2x^2}{f^2}}\right) - ex\right)}$$

e  
↓ 299

$$\frac{af^2 \int \frac{1}{-\sqrt{\frac{e^2x^2}{f^2} + a}f - ex} d\sqrt{d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}} + 2d\sqrt{f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex}}{2d} + \frac{af^2\sqrt{f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex}}{2d\left(f\left(-\sqrt{a + \frac{e^2x^2}{f^2}}\right) - ex\right)}$$

e  
↓ 219

$$\frac{af^2 \operatorname{arctanh}\left(\frac{\sqrt{f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex}}{\sqrt{d}}\right)}{\sqrt{d}} + \frac{2d\sqrt{f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex}}{2d} + \frac{af^2\sqrt{f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex}}{2d\left(f\left(-\sqrt{a + \frac{e^2x^2}{f^2}}\right) - ex\right)}$$

e

input `Int[1/Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]],x]`

output `((a*f^2*Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]])/(2*d*(-(e*x) - f*Sqrt[a + (e^2*x^2)/f^2])) + (2*d*Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]] + (a*f^2*ArcTanh[Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]]/Sqrt[d]])/Sqrt[d])/(2*d))/e`

## Definitions of rubi rules used

- rule 25  $\text{Int}[-(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 219  $\text{Int}[(\text{a}_) + (\text{b}_) * (\text{x}_)^2)^{-1}, \text{x\_Symbol}] \rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2] * \text{Rt}[-\text{b}, 2])) * \text{ArcTanh}[\text{Rt}[-\text{b}, 2] * (\text{x}/\text{Rt}[\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{NegQ}[\text{a}/\text{b}] \&\& (\text{GtQ}[\text{a}, 0] \mid\mid \text{LtQ}[\text{b}, 0])$
- rule 299  $\text{Int}[(\text{a}_) + (\text{b}_) * (\text{x}_)^2)^{\text{p}_} * ((\text{c}_) + (\text{d}_) * (\text{x}_)^2), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{d} * \text{x} * ((\text{a} + \text{b} * \text{x}^2)^{\text{p} + 1} / (\text{b} * (2 * \text{p} + 3))), \text{x}] - \text{Simp}[(\text{a} * \text{d} - \text{b} * \text{c} * (2 * \text{p} + 3)) / (\text{b} * (2 * \text{p} + 3)) \quad \text{Int}[(\text{a} + \text{b} * \text{x}^2)^{\text{p}}, \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \&\& \text{NeQ}[\text{b} * \text{c} - \text{a} * \text{d}, 0] \&\& \text{NeQ}[2 * \text{p} + 3, 0]$
- rule 1192  $\text{Int}[(\text{d}_) + (\text{e}_) * (\text{x}_)^2)^{\text{m}_} * ((\text{f}_) + (\text{g}_) * (\text{x}_)^2)^{\text{n}_} * ((\text{a}_) + (\text{b}_) * (\text{x}_) + (\text{c}_) * (\text{x}_)^2)^{\text{p}_}, \text{x\_Symbol}] \rightarrow \text{Simp}[2 / \text{e}^{(\text{n} + 2 * \text{p} + 1)} \quad \text{Subst}[\text{Int}[\text{x}^{(2 * \text{m} + 1) * (\text{e} * \text{f} - \text{d} * \text{g} + \text{g} * \text{x}^2)^{\text{n}} * (\text{c} * \text{d}^2 - \text{b} * \text{d} * \text{e} + \text{a} * \text{e}^2 - (2 * \text{c} * \text{d} - \text{b} * \text{e}) * \text{x}^2 + \text{c} * \text{x}^4)^{\text{p}}, \text{x}], \text{x}, \text{Sqrt}[\text{d} + \text{e} * \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{g}\}, \text{x}] \&\& \text{IGtQ}[\text{p}, 0] \&\& \text{ILtQ}[\text{n}, 0] \&\& \text{IntegerQ}[\text{m} + 1/2]$
- rule 1471  $\text{Int}[(\text{d}_) + (\text{e}_) * (\text{x}_)^2)^{\text{q}_} * ((\text{a}_) + (\text{b}_) * (\text{x}_)^2 + (\text{c}_) * (\text{x}_)^4)^{\text{p}_}, \text{x\_Symbol}] \rightarrow \text{With}[\{\text{Qx} = \text{PolynomialQuotient}[(\text{a} + \text{b} * \text{x}^2 + \text{c} * \text{x}^4)^{\text{p}}, \text{d} + \text{e} * \text{x}^2, \text{x}], \text{R} = \text{Coeff}[\text{PolynomialRemainder}[(\text{a} + \text{b} * \text{x}^2 + \text{c} * \text{x}^4)^{\text{p}}, \text{d} + \text{e} * \text{x}^2, \text{x}], \text{x}, 0]\}, \text{Simp}[(-\text{R}) * \text{x} * (\text{d} + \text{e} * \text{x}^2)^{\text{q} + 1} / (2 * \text{d} * (\text{q} + 1)), \text{x}] + \text{Simp}[1 / (2 * \text{d} * (\text{q} + 1)) \quad \text{Int}[(\text{d} + \text{e} * \text{x}^2)^{\text{q} + 1} * \text{ExpandToSum}[2 * \text{d} * (\text{q} + 1) * \text{Qx} + \text{R} * (2 * \text{q} + 3), \text{x}], \text{x}], \text{x}]] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}\}, \text{x}] \&\& \text{NeQ}[\text{b}^2 - 4 * \text{a} * \text{c}, 0] \&\& \text{NeQ}[\text{c} * \text{d}^2 - \text{b} * \text{d} * \text{e} + \text{a} * \text{e}^2, 0] \&\& \text{IGtQ}[\text{p}, 0] \&\& \text{LtQ}[\text{q}, -1]$
- rule 2542  $\text{Int}[(\text{g}_) + (\text{h}_) * ((\text{d}_) + (\text{e}_) * (\text{x}_) + (\text{f}_) * \text{Sqrt}[(\text{a}_) + (\text{c}_) * (\text{x}_)^2])^{\text{n}_})^{\text{p}_}, \text{x\_Symbol}] \rightarrow \text{Simp}[1 / (2 * \text{e}) \quad \text{Subst}[\text{Int}[(\text{g} + \text{h} * \text{x}^{\text{n}})^{\text{p}} * ((\text{d}^2 + \text{a} * \text{f}^2 - 2 * \text{d} * \text{x} + \text{x}^2) / (\text{d} - \text{x})^2), \text{x}], \text{x}, \text{d} + \text{e} * \text{x} + \text{f} * \text{Sqrt}[\text{a} + \text{c} * \text{x}^2], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{c}, \text{d}, \text{e}, \text{f}, \text{g}, \text{h}, \text{n}\}, \text{x}] \&\& \text{EqQ}[\text{e}^2 - \text{c} * \text{f}^2, 0] \&\& \text{IntegerQ}[\text{p}]$

**Maple [F]**

$$\int \frac{1}{\sqrt{d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}}} dx$$

input `int(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2),x)`

output `int(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2),x)`

**Fricas [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 347, normalized size of antiderivative = 2.36

$$\int \frac{1}{\sqrt{d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}}} dx$$

$$= \frac{a\sqrt{d}f^2 \log\left(a f^2 - 2dex + 2df\sqrt{\frac{e^2x^2+af^2}{f^2}} - 2\left(\sqrt{d}ex - \sqrt{d}f\sqrt{\frac{e^2x^2+af^2}{f^2}}\right)\sqrt{ex + f\sqrt{\frac{e^2x^2+af^2}{f^2}} + d}\right) + 2}{4d^2e}$$

input `integrate(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2),x, algorithm="fricas")`

output `[1/4*(a*sqrt(d)*f^2*log(a*f^2 - 2*d*e*x + 2*d*f*sqrt((e^2*x^2 + a*f^2)/f^2) - 2*(sqrt(d)*e*x - sqrt(d)*f*sqrt((e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d) + 2*(d*e*x - d*f*sqrt((e^2*x^2 + a*f^2)/f^2) + 2*d^2)*sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d)/(d^2*e), 1/2*(a*sqrt(-d)*f^2*arctan(-sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d)*(sqrt(-d)*f*sqrt((e^2*x^2 + a*f^2)/f^2) - (e*x + d)*sqrt(-d))/(a*f^2 - 2*d*e*x - d^2)) + (d*e*x - d*f*sqrt((e^2*x^2 + a*f^2)/f^2) + 2*d^2)*sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d)/(d^2*e)]`

**Sympy [F]**

$$\int \frac{1}{\sqrt{d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}}} dx = \int \frac{1}{\sqrt{d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}}} dx$$

input `integrate(1/(d+e*x+f*(a+e**2*x**2/f**2)**(1/2))**(1/2),x)`

output `Integral(1/sqrt(d + e*x + f*sqrt(a + e**2*x**2/f**2)), x)`

**Maxima [F]**

$$\int \frac{1}{\sqrt{d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}}} dx = \int \frac{1}{\sqrt{ex + \sqrt{\frac{e^2x^2}{f^2} + af} + d}} dx$$

input `integrate(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(e*x + sqrt(e^2*x^2/f^2 + a)*f + d), x)`

**Giac [F]**

$$\int \frac{1}{\sqrt{d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}}} dx = \int \frac{1}{\sqrt{ex + \sqrt{\frac{e^2x^2}{f^2} + af} + d}} dx$$

input `integrate(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2),x, algorithm="giac")`

output `integrate(1/sqrt(e*x + sqrt(e^2*x^2/f^2 + a)*f + d), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}}} dx = \int \frac{1}{\sqrt{d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}}} dx$$

input `int(1/(d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^(1/2),x)`output `int(1/(d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^(1/2), x)`**Reduce [F]**

$$\int \frac{1}{\sqrt{d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}}} dx = \int \frac{1}{\sqrt{d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}}} dx$$

input `int(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2),x)`output `int(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2),x)`

$$3.12 \quad \int \frac{1}{\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)^{3/2}} dx$$

Optimal result . . . . .	125
Mathematica [A] (verified) . . . . .	126
Rubi [A] (verified) . . . . .	126
Maple [F] . . . . .	129
Fricas [A] (verification not implemented) . . . . .	129
Sympy [F] . . . . .	130
Maxima [F] . . . . .	130
Giac [F] . . . . .	130
Mupad [F(-1)] . . . . .	131
Reduce [F] . . . . .	131

**Optimal result**

Integrand size = 27, antiderivative size = 158

$$\int \frac{1}{\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)^{3/2}} dx = -\frac{1+\frac{af^2}{d^2}}{e\sqrt{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}}} - \frac{af^2\sqrt{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}}}{2d^2e\left(ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)} + \frac{3af^2\operatorname{arctanh}\left(\frac{\sqrt{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}}}{\sqrt{d}}\right)}{2d^{5/2}e}$$

output

```
-(1+af^2/d^2)/e/(d+ex+f*(a+e^2*x^2/f^2)^(1/2))^(1/2)-1/2*af^2*(d+ex+f*(a+e^2*x^2/f^2)^(1/2))^(1/2)/d^2/e/(ex+f*(a+e^2*x^2/f^2)^(1/2))+3/2*af^2*arctanh((d+ex+f*(a+e^2*x^2/f^2)^(1/2))^(1/2)/d^(1/2))/d^(5/2)/e
```

### Mathematica [A] (verified)

Time = 1.69 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.07

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{3/2}} dx = \frac{\sqrt{d}\left(2d^2\left(ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)+af^2\left(d+3ex+3f\sqrt{a+\frac{e^2x^2}{f^2}}\right)\right)}{\left(ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)\sqrt{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}}} + 3af^2\operatorname{arctanh}\left(\sqrt{\frac{d+ex}{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}}}\right)$$

input

```
Integrate[(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(-3/2),x]
```

output

```
(-((Sqrt[d]*(2*d^2*(e*x + f*Sqrt[a + (e^2*x^2)/f^2]) + a*f^2*(d + 3*e*x + 3*f*Sqrt[a + (e^2*x^2)/f^2])))/((e*x + f*Sqrt[a + (e^2*x^2)/f^2])*Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]])) + 3*a*f^2*ArcTanh[Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]]/Sqrt[d]])/(2*d^(5/2)*e)
```

### Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.01, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {2542, 1192, 1582, 25, 359, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\left(f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex\right)^{3/2}} dx$$

↓ 2542

$$\int \frac{d^2 - 2\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)d + af^2 + \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)^2}{\left(-\sqrt{\frac{e^2x^2}{f^2} + a} - ex\right)^2 \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)^{3/2}} d\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)$$

2e

↓ 1192

$$\begin{aligned}
 & \frac{\int \frac{d^2 - 2\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right) d + af^2 + \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)^2}{\left(-\sqrt{\frac{e^2x^2}{f^2} + a}f - ex\right)^2 \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)} d\sqrt{d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}}}{e} \\
 & \quad \downarrow \text{1582} \\
 & \frac{af^2\sqrt{f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex} \int -\frac{2d(d^2 + af^2) - (2d^2 - af^2)\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)}{\left(-\sqrt{\frac{e^2x^2}{f^2} + a}f - ex\right)\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)} d\sqrt{d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}}}{2d^2\left(f\left(-\sqrt{a + \frac{e^2x^2}{f^2}}\right) - ex\right)} - \frac{2d^2}{2d^2} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \frac{2d(d^2 + af^2) - (2d^2 - af^2)\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)}{\left(-\sqrt{\frac{e^2x^2}{f^2} + a}f - ex\right)\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)} d\sqrt{d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}}}{2d^2} + \frac{af^2\sqrt{f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex}}{2d^2\left(f\left(-\sqrt{a + \frac{e^2x^2}{f^2}}\right) - ex\right)} \\
 & \quad \downarrow \text{359} \\
 & \frac{3af^2 \int \frac{1}{-\sqrt{\frac{e^2x^2}{f^2} + a}f - ex} d\sqrt{d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}} - \frac{2(af^2 + d^2)}{\sqrt{f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex}}}{2d^2} + \frac{af^2\sqrt{f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex}}{2d^2\left(f\left(-\sqrt{a + \frac{e^2x^2}{f^2}}\right) - ex\right)} \\
 & \quad \downarrow \text{219} \\
 & \frac{3af^2 \operatorname{arctanh}\left(\frac{\sqrt{f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex}}{\sqrt{d}}\right)}{\sqrt{d}} - \frac{2(af^2 + d^2)}{\sqrt{f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex}} + \frac{af^2\sqrt{f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex}}{2d^2\left(f\left(-\sqrt{a + \frac{e^2x^2}{f^2}}\right) - ex\right)} \\
 & \quad \downarrow e
 \end{aligned}$$

input `Int[(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(-3/2),x]`

output `((a*f^2*Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]])/(2*d^2*(-(e*x) - f*Sqrt[a + (e^2*x^2)/f^2])) + ((-2*(d^2 + a*f^2))/Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]]) + (3*a*f^2*ArcTanh[Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]]/Sqrt[d]])/Sqrt[d])/(2*d^2))/e`



## Defintions of rubi rules used

- rule 25  $\text{Int}[-(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 219  $\text{Int}[(\text{a}_) + (\text{b}_) * (\text{x}_)^2)^{-1}, \text{x\_Symbol}] \rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2] * \text{Rt}[-\text{b}, 2])) * \text{ArcTanh}[\text{Rt}[-\text{b}, 2] * (\text{x}/\text{Rt}[\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{NegQ}[\text{a}/\text{b}] \&\& (\text{GtQ}[\text{a}, 0] \parallel \text{LtQ}[\text{b}, 0])$
- rule 359  $\text{Int}[(\text{e}_) * (\text{x}_)]^{(\text{m}_)} * ((\text{a}_) + (\text{b}_) * (\text{x}_)^2)^{(\text{p}_)} * ((\text{c}_) + (\text{d}_) * (\text{x}_)^2), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{c} * (\text{e} * \text{x})^{(\text{m} + 1)} * ((\text{a} + \text{b} * \text{x}^2)^{(\text{p} + 1)} / (\text{a} * \text{e}^{(\text{m} + 1)})), \text{x}] + \text{Simp}[(\text{a} * \text{d} * (\text{m} + 1) - \text{b} * \text{c} * (\text{m} + 2 * \text{p} + 3)) / (\text{a} * \text{e}^{2 * (\text{m} + 1)}) \quad \text{Int}[(\text{e} * \text{x})^{(\text{m} + 2)} * (\text{a} + \text{b} * \text{x}^2)^{\text{p}}, \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{p}\}, \text{x}] \&\& \text{NeQ}[\text{b} * \text{c} - \text{a} * \text{d}, 0] \&\& \text{LtQ}[\text{m}, -1] \&\& !\text{ILtQ}[\text{p}, -1]$
- rule 1192  $\text{Int}[(\text{d}_) + (\text{e}_) * (\text{x}_)]^{(\text{m}_)} * ((\text{f}_) + (\text{g}_) * (\text{x}_)]^{(\text{n}_)} * ((\text{a}_) + (\text{b}_) * (\text{x}_) + (\text{c}_) * (\text{x}_)^2)^{(\text{p}_)}, \text{x\_Symbol}] \rightarrow \text{Simp}[2 / \text{e}^{(\text{n} + 2 * \text{p} + 1)} \quad \text{Subst}[\text{Int}[\text{x}^{(2 * \text{m} + 1)} * (\text{e} * \text{f} - \text{d} * \text{g} + \text{g} * \text{x}^2)^{\text{n}} * (\text{c} * \text{d}^2 - \text{b} * \text{d} * \text{e} + \text{a} * \text{e}^2 - (2 * \text{c} * \text{d} - \text{b} * \text{e}) * \text{x}^2 + \text{c} * \text{x}^4)^{\text{p}}, \text{x}], \text{x}, \text{Sqrt}[\text{d} + \text{e} * \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{g}\}, \text{x}] \&\& \text{IGtQ}[\text{p}, 0] \&\& \text{ILtQ}[\text{n}, 0] \&\& \text{IntegerQ}[\text{m} + 1/2]$
- rule 1582  $\text{Int}[(\text{x}_)]^{(\text{m}_)} * ((\text{d}_) + (\text{e}_) * (\text{x}_)^2)^{(\text{q}_)} * ((\text{a}_) + (\text{b}_) * (\text{x}_)^2 + (\text{c}_) * (\text{x}_)^4)^{(\text{p}_)}, \text{x\_Symbol}] \rightarrow \text{Simp}[(-\text{d})^{(\text{m}/2 - 1)} * (\text{c} * \text{d}^2 - \text{b} * \text{d} * \text{e} + \text{a} * \text{e}^2)^{\text{p}} * \text{x} * ((\text{d} + \text{e} * \text{x}^2)^{(\text{q} + 1)} / (2 * \text{e}^{(2 * \text{p} + \text{m}/2)} * (\text{q} + 1))), \text{x}] + \text{Simp}[(-\text{d})^{(\text{m}/2 - 1)} / (2 * \text{e}^{(2 * \text{p})} * (\text{q} + 1)) \quad \text{Int}[\text{x}^{\text{m}} * (\text{d} + \text{e} * \text{x}^2)^{(\text{q} + 1)} * \text{ExpandToSum}[\text{Together}[(1/(\text{d} + \text{e} * \text{x}^2)) * (2 * (-\text{d})^{(-\text{m}/2 + 1)} * \text{e}^{(2 * \text{p})} * (\text{q} + 1) * (\text{a} + \text{b} * \text{x}^2 + \text{c} * \text{x}^4)^{\text{p}} - ((\text{c} * \text{d}^2 - \text{b} * \text{d} * \text{e} + \text{a} * \text{e}^2)^{\text{p}} / (\text{e}^{(\text{m}/2)} * \text{x}^{\text{m}})) * (\text{d} + \text{e} * (2 * \text{q} + 3) * \text{x}^2)], \text{x}], \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}\}, \text{x}] \&\& \text{NeQ}[\text{b}^2 - 4 * \text{a} * \text{c}, 0] \&\& \text{IGtQ}[\text{p}, 0] \&\& \text{ILtQ}[\text{q}, -1] \&\& \text{ILtQ}[\text{m}/2, 0]$
- rule 2542  $\text{Int}[(\text{g}_) + (\text{h}_) * ((\text{d}_) + (\text{e}_) * (\text{x}_) + (\text{f}_) * \text{Sqrt}[(\text{a}_) + (\text{c}_) * (\text{x}_)^2])]^{(\text{n}_)} * ((\text{d}_) + (\text{e}_) * (\text{x}_) + (\text{f}_) * \text{Sqrt}[(\text{a}_) + (\text{c}_) * (\text{x}_)^2])^{(\text{p}_)}, \text{x\_Symbol}] \rightarrow \text{Simp}[1 / (2 * \text{e}) \quad \text{Subst}[\text{Int}[(\text{g} + \text{h} * \text{x}^{\text{n}})^{\text{p}} * ((\text{d}^2 + \text{a} * \text{f}^2 - 2 * \text{d} * \text{x} + \text{x}^2) / (\text{d} - \text{x})^2), \text{x}], \text{x}, \text{d} + \text{e} * \text{x} + \text{f} * \text{Sqrt}[\text{a} + \text{c} * \text{x}^2], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{c}, \text{d}, \text{e}, \text{f}, \text{g}, \text{h}, \text{n}\}, \text{x}] \&\& \text{EqQ}[\text{e}^2 - \text{c} * \text{f}^2, 0] \&\& \text{IntegerQ}[\text{p}]$

**Maple [F]**

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{\frac{3}{2}}} dx$$

input `int(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(3/2),x)`

output `int(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(3/2),x)`

**Fricas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 538, normalized size of antiderivative = 3.41

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{\frac{3}{2}}} dx = \left[ \frac{3(a^2f^4 - 2ade^2x - ad^2f^2)\sqrt{d} \log\left(af^2 - 2dex + 2df\sqrt{\frac{e^2x^2 + af^2}{f^2}}\right)}{\dots} \right]$$

input `integrate(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(3/2),x, algorithm="fricas")`

output `[1/4*(3*(a^2*f^4 - 2*a*d*e*f^2*x - a*d^2*f^2)*sqrt(d)*log(a*f^2 - 2*d*e*x + 2*d*f*sqrt((e^2*x^2 + a*f^2)/f^2) - 2*(sqrt(d)*e*x - sqrt(d)*f*sqrt((e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d) - 2*(2*d^2*e^2*x^2 - 2*a*d^2*f^2 - 2*d^4 - (3*a*d*e*f^2 + d^3*e)*x + (3*a*d*f^3 - 2*d^2*e*f*x + d^3*f)*sqrt((e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d)/(a*d^3*e*f^2 - 2*d^4*e^2*x - d^5*e), 1/2*(3*(a^2*f^4 - 2*a*d*e*f^2*x - a*d^2*f^2)*sqrt(-d)*arctan(-sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d)*(sqrt(-d)*f*sqrt((e^2*x^2 + a*f^2)/f^2) - (e*x + d)*sqrt(-d))/(a*f^2 - 2*d*e*x - d^2)) - (2*d^2*e^2*x^2 - 2*a*d^2*f^2 - 2*d^4 - (3*a*d*e*f^2 + d^3*e)*x + (3*a*d*f^3 - 2*d^2*e*f*x + d^3*f)*sqrt((e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d)/(a*d^3*e*f^2 - 2*d^4*e^2*x - d^5*e)]`

**Sympy [F]**

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{3/2}} dx = \int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{3/2}} dx$$

input `integrate(1/(d+e*x+f*(a+e**2*x**2/f**2)**(1/2))**(3/2),x)`

output `Integral((d + e*x + f*sqrt(a + e**2*x**2/f**2))**(-3/2), x)`

**Maxima [F]**

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{3/2}} dx = \int \frac{1}{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + af + d}\right)^{3/2}} dx$$

input `integrate(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(3/2),x, algorithm="maxima")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a)*f + d)^(-3/2), x)`

**Giac [F]**

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{3/2}} dx = \int \frac{1}{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + af + d}\right)^{3/2}} dx$$

input `integrate(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(3/2),x, algorithm="giac")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a)*f + d)^(-3/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{3/2}} dx = \int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{3/2}} dx$$

input `int(1/(d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^(3/2),x)`output `int(1/(d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^(3/2), x)`**Reduce [F]**

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{3/2}} dx = \int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{3/2}} dx$$

input `int(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(3/2),x)`output `int(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(3/2),x)`

**3.13** 
$$\int \frac{1}{\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)^{5/2}} dx$$

Optimal result . . . . .	132
Mathematica [A] (verified) . . . . .	133
Rubi [A] (verified) . . . . .	133
Maple [F] . . . . .	136
Fricas [B] (verification not implemented) . . . . .	136
Sympy [F] . . . . .	137
Maxima [F] . . . . .	138
Giac [F] . . . . .	138
Mupad [F(-1)] . . . . .	138
Reduce [F] . . . . .	139

**Optimal result**

Integrand size = 27, antiderivative size = 199

$$\int \frac{1}{\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)^{5/2}} dx = -\frac{1+\frac{af^2}{d^2}}{3e\left(d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)^{3/2}} - \frac{2af^2}{d^3e\sqrt{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}}} - \frac{af^2\sqrt{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}}}{2d^3e\left(ex+f\sqrt{a+\frac{e^2x^2}{f^2}}\right)} + \frac{5af^2\operatorname{arctanh}\left(\frac{\sqrt{d+ex+f\sqrt{a+\frac{e^2x^2}{f^2}}}}{\sqrt{d}}\right)}{2d^{7/2}e}$$

output

```
-1/3*(1+a*f^2/d^2)/e/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(3/2)-2*a*f^2/d^3/e/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2)-1/2*a*f^2*(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))/d^3/e/(e*x+f*(a+e^2*x^2/f^2)^(1/2))+5/2*a*f^2*arctanh((d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(1/2)/d^(1/2))/d^(7/2)/e
```

### Mathematica [A] (verified)

Time = 1.82 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.05

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{5/2}} dx = \frac{\sqrt{d}\left(15a^2f^4 + 2d^3\left(ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right) + af^2\left(3d^2 + 20d\left(ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right) + 30ex\left(ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)\right)\right)}{\left(ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{3/2}} \frac{1}{6d^{7/2}e}$$

input

```
Integrate[(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(-5/2), x]
```

output

```
(-((Sqrt[d]*(15*a^2*f^4 + 2*d^3*(e*x + f*Sqrt[a + (e^2*x^2)/f^2]) + a*f^2*(3*d^2 + 20*d*(e*x + f*Sqrt[a + (e^2*x^2)/f^2]) + 30*e*x*(e*x + f*Sqrt[a + (e^2*x^2)/f^2]))) / ((e*x + f*Sqrt[a + (e^2*x^2)/f^2])*(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(3/2))) + 15*a*f^2*ArcTanh[Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]]/Sqrt[d]]) / (6*d^(7/2)*e)
```

### Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.98, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$ , Rules used = {2542, 1192, 1582, 1584, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\left(f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex\right)^{5/2}} dx$$

↓ 2542

$$\int \frac{d^2 - 2\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)d + af^2 + \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)^2}{\left(-\sqrt{\frac{e^2x^2}{f^2} + a}f - ex\right)^2 \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)^{5/2}} d \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)$$

2e

↓ 1192

$$\int \frac{d^2 - 2\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)d + af^2 + \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)^2}{\left(-\sqrt{\frac{e^2x^2}{f^2} + a}f - ex\right)^2 \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)^2} d\sqrt{d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}}$$

e  
↓ 1582

$$\int \frac{2(d^2 + af^2)d^2 - 2(d^2 - af^2)\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)d + af^2\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)^2}{\left(-\sqrt{\frac{e^2x^2}{f^2} + a}f - ex\right)\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)^2} d\sqrt{d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}} + \frac{af^2\sqrt{f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex}}{2d^3\left(f\left(-\sqrt{a + \frac{e^2x^2}{f^2}}\right) - ex\right)}$$

e  
↓ 1584

$$\int \left( \frac{5af^2}{-\sqrt{\frac{e^2x^2}{f^2} + a}f - ex} + \frac{4af^2}{d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}} + \frac{2(d^3 + af^2d)}{\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}\right)^2} \right) d\sqrt{d + ex + f\sqrt{\frac{e^2x^2}{f^2} + a}} + \frac{af^2\sqrt{f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex}}{2d^3\left(f\left(-\sqrt{a + \frac{e^2x^2}{f^2}}\right) - ex\right)}$$

e  
↓ 2009

$$\frac{5af^2 \operatorname{arctanh}\left(\frac{\sqrt{f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex}}{\sqrt{d}}\right)}{\sqrt{d}} - \frac{2d(af^2 + d^2)}{3\left(f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex\right)^{3/2}} - \frac{4af^2}{\sqrt{f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex}} + \frac{af^2\sqrt{f\sqrt{a + \frac{e^2x^2}{f^2}} + d + ex}}{2d^3\left(f\left(-\sqrt{a + \frac{e^2x^2}{f^2}}\right) - ex\right)}$$

e

input `Int[(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(-5/2),x]`

output `((a*f^2*Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]])/(2*d^3*(-(e*x) - f*Sqrt[a + (e^2*x^2)/f^2])) + ((-2*d*(d^2 + a*f^2))/(3*(d + e*x + f*Sqrt[a + (e^2*x^2)/f^2])^(3/2)) - (4*a*f^2)/Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]] + (5*a*f^2*ArcTanh[Sqrt[d + e*x + f*Sqrt[a + (e^2*x^2)/f^2]]/Sqrt[d]])/Sqrt[d])/(2*d^3))/e`

## Definitions of rubi rules used

rule 1192

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))^(n_)*((a_.) + (b_.)*(x_)
+ (c_.)*(x_)^2)^(p_.), x_Symbol] := Simp[2/e^(n + 2*p + 1) Subst[Int[x^(
2*m + 1)*(e*f - d*g + g*x^2)^n*(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 +
c*x^4)^p, x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, g}, x] &&
IGtQ[p, 0] && ILtQ[n, 0] && IntegerQ[m + 1/2]
```

rule 1582

```
Int[(x_)^(m_)*((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^
4)^(p_.), x_Symbol] := Simp[(-d)^(m/2 - 1)*(c*d^2 - b*d*e + a*e^2)^p*x*((d
+ e*x^2)^(q + 1)/(2*e^(2*p + m/2)*(q + 1))), x] + Simp[(-d)^(m/2 - 1)/(2*e^
(2*p)*(q + 1)) Int[x^m*(d + e*x^2)^(q + 1)*ExpandToSum[Together[(1/(d + e
*x^2))*(2*(-d)^(-m/2 + 1)*e^(2*p)*(q + 1)*(a + b*x^2 + c*x^4)^p - ((c*d^2 -
b*d*e + a*e^2)^p/(e^(m/2)*x^m))*(d + e*(2*q + 3)*x^2)], x], x] /; Fre
eQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && ILtQ[q, -1]
&& ILtQ[m/2, 0]
```

rule 1584

```
Int[((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (
c_.)*(x_)^4)^(p_.), x_Symbol] := Int[ExpandIntegrand[(f*x)^m*(d + e*x^2)^q*
(a + b*x^2 + c*x^4)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, q}, x] && NeQ[
b^2 - 4*a*c, 0] && IGtQ[p, 0] && IGtQ[q, -2]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2542

```
Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(
n_))^(p_.), x_Symbol] := Simp[1/(2*e) Subst[Int[(g + h*x^n)^p*((d^2 + a*f
^2 - 2*d*x + x^2)/(d - x)^2), x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; Fr
eeQ[{a, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]
```



**Maple [F]**

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{5/2}} dx$$

input `int(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(5/2),x)`

output `int(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(5/2),x)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 426 vs. 2(171) = 342.

Time = 0.34 (sec) , antiderivative size = 861, normalized size of antiderivative = 4.33

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{5/2}} dx = \text{Too large to display}$$

input `integrate(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(5/2),x, algorithm="fricas")`

output

```
[1/12*(15*(a^3*f^6 + 4*a*d^2*e^2*f^2*x^2 - 2*a^2*d^2*f^4 + a*d^4*f^2 - 4*(a^2*d*e*f^4 - a*d^3*e*f^2)*x)*sqrt(d)*log(a*f^2 - 2*d*e*x + 2*d*f*sqrt((e^2*x^2 + a*f^2)/f^2)) - 2*(sqrt(d)*e*x - sqrt(d)*f*sqrt((e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d) + 2*(12*d^3*e^3*x^3 + 10*a^2*d^2*f^4 - 16*a*d^4*f^2 - 2*d^6 - 8*(5*a*d^2*e^2*f^2 - d^4*e^2)*x^2 + (15*a^2*d*e*f^4 - 46*a*d^3*e*f^2 - d^5*e)*x - (15*a^2*d*f^5 + 12*d^3*e^2*f*x^2 - 22*a*d^3*f^3 - d^5*f - 8*(5*a*d^2*e*f^3 - d^4*e*f)*x)*sqrt((e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d))/(a^2*d^4*e*f^4 + 4*d^6*e^3*x^2 - 2*a*d^6*e*f^2 + d^8*e - 4*(a*d^5*e^2*f^2 - d^7*e^2)*x), 1/6*(15*(a^3*f^6 + 4*a*d^2*e^2*f^2*x^2 - 2*a^2*d^2*f^4 + a*d^4*f^2 - 4*(a^2*d*e*f^4 - a*d^3*e*f^2)*x)*sqrt(-d)*arctan(-sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d)*(sqrt(-d)*f*sqrt((e^2*x^2 + a*f^2)/f^2) - (e*x + d)*sqrt(-d))/(a*f^2 - 2*d*e*x - d^2)) + (12*d^3*e^3*x^3 + 10*a^2*d^2*f^4 - 16*a*d^4*f^2 - 2*d^6 - 8*(5*a*d^2*e^2*f^2 - d^4*e^2)*x^2 + (15*a^2*d*e*f^4 - 46*a*d^3*e*f^2 - d^5*e)*x - (15*a^2*d*f^5 + 12*d^3*e^2*f*x^2 - 22*a*d^3*f^3 - d^5*f - 8*(5*a*d^2*e*f^3 - d^4*e*f)*x)*sqrt((e^2*x^2 + a*f^2)/f^2))*sqrt(e*x + f*sqrt((e^2*x^2 + a*f^2)/f^2) + d))/(a^2*d^4*e*f^4 + 4*d^6*e^3*x^2 - 2*a*d^6*e*f^2 + d^8*e - 4*(a*d^5*e^2*f^2 - d^7*e^2)*x)]
```

SymPy [F]

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{5/2}} dx = \int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{5/2}} dx$$

input

```
integrate(1/(d+e*x+f*(a+e**2*x**2/f**2)**(1/2))**(5/2),x)
```

output

```
Integral((d + e*x + f*sqrt(a + e**2*x**2/f**2))**(-5/2), x)
```

**Maxima [F]**

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{5/2}} dx = \int \frac{1}{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + af + d}\right)^{5/2}} dx$$

input `integrate(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(5/2),x, algorithm="maxima")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a)*f + d)^(-5/2), x)`

**Giac [F]**

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{5/2}} dx = \int \frac{1}{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + af + d}\right)^{5/2}} dx$$

input `integrate(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(5/2),x, algorithm="giac")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a)*f + d)^(-5/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{5/2}} dx = \int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{5/2}} dx$$

input `int(1/(d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^(5/2),x)`

output `int(1/(d + e*x + f*(a + (e^2*x^2)/f^2)^(1/2))^(5/2), x)`

Reduce [F]

$$\int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{5/2}} dx = \int \frac{1}{\left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{5/2}} dx$$

input `int(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(5/2),x)`

output `int(1/(d+e*x+f*(a+e^2*x^2/f^2)^(1/2))^(5/2),x)`

### 3.14 $\int \sqrt{x - \sqrt{-4 + x^2}} dx$

Optimal result	140
Mathematica [A] (verified)	140
Rubi [A] (verified)	141
Maple [F]	142
Fricas [A] (verification not implemented)	142
Sympy [F]	143
Maxima [F]	143
Giac [A] (verification not implemented)	143
Mupad [F(-1)]	144
Reduce [B] (verification not implemented)	144

#### Optimal result

Integrand size = 17, antiderivative size = 41

$$\int \sqrt{x - \sqrt{-4 + x^2}} dx = \frac{4}{\sqrt{x - \sqrt{-4 + x^2}}} + \frac{1}{3} \left( x - \sqrt{-4 + x^2} \right)^{3/2}$$

output  $4/(x-(x^2-4)^{(1/2)})^{(1/2)}+1/3*(x-(x^2-4)^{(1/2)})^{(3/2)}$

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \sqrt{x - \sqrt{-4 + x^2}} dx = \frac{4}{\sqrt{x - \sqrt{-4 + x^2}}} + \frac{1}{3} \left( x - \sqrt{-4 + x^2} \right)^{3/2}$$

input `Integrate[Sqrt[x - Sqrt[-4 + x^2]], x]`

output  $4/\text{Sqrt}[x - \text{Sqrt}[-4 + x^2]] + (x - \text{Sqrt}[-4 + x^2])^{(3/2)}/3$

**Rubi [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.10, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {2542, 25, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{x - \sqrt{x^2 - 4}} dx \\
 & \quad \downarrow \text{2542} \\
 & \frac{1}{2} \int -\frac{4 - (x - \sqrt{x^2 - 4})^2}{(x - \sqrt{x^2 - 4})^{3/2}} d(x - \sqrt{x^2 - 4}) \\
 & \quad \downarrow \text{25} \\
 & -\frac{1}{2} \int \frac{4 - (x - \sqrt{x^2 - 4})^2}{(x - \sqrt{x^2 - 4})^{3/2}} d(x - \sqrt{x^2 - 4}) \\
 & \quad \downarrow \text{244} \\
 & -\frac{1}{2} \int \left( \frac{4}{(x - \sqrt{x^2 - 4})^{3/2}} - \sqrt{x - \sqrt{x^2 - 4}} \right) d(x - \sqrt{x^2 - 4}) \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{2} \left( \frac{2}{3} (x - \sqrt{x^2 - 4})^{3/2} + \frac{8}{\sqrt{x - \sqrt{x^2 - 4}}} \right)
 \end{aligned}$$

input `Int[Sqrt[x - Sqrt[-4 + x^2]],x]`

output `(8/Sqrt[x - Sqrt[-4 + x^2]] + (2*(x - Sqrt[-4 + x^2])^(3/2))/3)/2`

**Defintions of rubi rules used**

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 244 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2542 `Int[((g_) + (h_)*((d_) + (e_)*(x_) + (f_)*Sqrt[(a_) + (c_)*(x_)^2])^(n_))^(p_), x_Symbol] := Simp[1/(2*e) Subst[Int[(g + h*x^n)^p*((d^2 + a*f^2 - 2*d*x + x^2)/(d - x)^2), x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]`

**Maple [F]**

$$\int \sqrt{x - \sqrt{x^2 - 4}} dx$$

input `int((x-(x^2-4)^(1/2))^(1/2),x)`

output `int((x-(x^2-4)^(1/2))^(1/2),x)`

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.63

$$\int \sqrt{x - \sqrt{-4 + x^2}} dx = \frac{2}{3} \left( 2x + \sqrt{x^2 - 4} \right) \sqrt{x - \sqrt{x^2 - 4}}$$

input `integrate((x-(x^2-4)^(1/2))^(1/2),x, algorithm="fricas")`

output `2/3*(2*x + sqrt(x^2 - 4))*sqrt(x - sqrt(x^2 - 4))`

### Sympy [F]

$$\int \sqrt{x - \sqrt{-4 + x^2}} dx = \int \sqrt{x - \sqrt{x^2 - 4}} dx$$

input `integrate((x-(x**2-4)**(1/2))**(1/2),x)`

output `Integral(sqrt(x - sqrt(x**2 - 4)), x)`

### Maxima [F]

$$\int \sqrt{x - \sqrt{-4 + x^2}} dx = \int \sqrt{x - \sqrt{x^2 - 4}} dx$$

input `integrate((x-(x^2-4)^(1/2))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(x - sqrt(x^2 - 4)), x)`

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.61

$$\int \sqrt{x - \sqrt{-4 + x^2}} dx = \frac{2}{3}(x + 2)\sqrt{\frac{1}{2}x + 1} - \frac{2}{3}(x - 2)\sqrt{\frac{1}{2}x - 1}$$

input `integrate((x-(x^2-4)^(1/2))^(1/2),x, algorithm="giac")`

output `2/3*(x + 2)*sqrt(1/2*x + 1) - 2/3*(x - 2)*sqrt(1/2*x - 1)`



**Mupad [F(-1)]**

Timed out.

$$\int \sqrt{x - \sqrt{-4 + x^2}} dx = \int \sqrt{x - \sqrt{x^2 - 4}} dx$$

input `int((x - (x^2 - 4)^(1/2))^(1/2), x)`output `int((x - (x^2 - 4)^(1/2))^(1/2), x)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.95

$$\int \sqrt{x - \sqrt{-4 + x^2}} dx = \frac{\sqrt{x+2}\sqrt{2}(-\sqrt{x^2-4}x + 2\sqrt{x^2-4} + x^2 + 4x + 4)}{3x+6}$$

input `int((x-(x^2-4)^(1/2))^(1/2), x)`output `(sqrt(x + 2)*sqrt(2)*(- sqrt(x**2 - 4)*x + 2*sqrt(x**2 - 4) + x**2 + 4*x + 4))/(3*(x + 2))`

$$3.15 \quad \int \sqrt{ax + b\sqrt{c + \frac{a^2x^2}{b^2}}} dx$$

Optimal result	145
Mathematica [A] (verified)	145
Rubi [A] (verified)	146
Maple [F]	147
Fricas [A] (verification not implemented)	147
Sympy [F]	148
Maxima [F]	148
Giac [F]	148
Mupad [F(-1)]	149
Reduce [B] (verification not implemented)	149

### Optimal result

Integrand size = 26, antiderivative size = 69

$$\int \sqrt{ax + b\sqrt{c + \frac{a^2x^2}{b^2}}} dx = -\frac{b^2c}{a\sqrt{ax + b\sqrt{c + \frac{a^2x^2}{b^2}}}} + \frac{\left(ax + b\sqrt{c + \frac{a^2x^2}{b^2}}\right)^{3/2}}{3a}$$

output

```
-b^2*c/a/(a*x+b*(c+a^2*x^2/b^2)^(1/2))^(1/2)+1/3*(a*x+b*(c+a^2*x^2/b^2)^(1/2))^(3/2)/a
```

### Mathematica [A] (verified)

Time = 0.73 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00

$$\int \sqrt{ax + b\sqrt{c + \frac{a^2x^2}{b^2}}} dx = -\frac{b^2c}{a\sqrt{ax + b\sqrt{c + \frac{a^2x^2}{b^2}}}} + \frac{\left(ax + b\sqrt{c + \frac{a^2x^2}{b^2}}\right)^{3/2}}{3a}$$

input

```
Integrate[Sqrt[a*x + b*Sqrt[c + (a^2*x^2)/b^2]],x]
```

output

$$-\left(\frac{b^2 c}{a \sqrt{a x + b \sqrt{c + \frac{a^2 x^2}{b^2}}}}\right) + \frac{a x + b \sqrt{c + \frac{a^2 x^2}{b^2}}}{3 a}$$

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.01, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$ , Rules used = {2542, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{b \sqrt{\frac{a^2 x^2}{b^2} + c} + a x} dx$$

$$\downarrow \text{2542}$$

$$\int \frac{cb^2 + \left(\sqrt{\frac{a^2 x^2}{b^2} + cb + ax}\right)^2}{\left(\sqrt{\frac{a^2 x^2}{b^2} + cb + ax}\right)^{3/2}} d\left(\sqrt{\frac{a^2 x^2}{b^2} + cb + ax}\right)$$

$$\frac{2a}{\downarrow \text{244}}$$

$$\int \left( \frac{cb^2}{\left(\sqrt{\frac{a^2 x^2}{b^2} + cb + ax}\right)^{3/2}} + \sqrt{\sqrt{\frac{a^2 x^2}{b^2} + cb + ax}} \right) d\left(\sqrt{\frac{a^2 x^2}{b^2} + cb + ax}\right)$$

$$\frac{2a}{\downarrow \text{2009}}$$

$$\frac{\frac{2}{3} \left( b \sqrt{\frac{a^2 x^2}{b^2} + c} + ax \right)^{3/2} - \frac{2b^2 c}{\sqrt{b \sqrt{\frac{a^2 x^2}{b^2} + c} + ax}}}{2a}$$

input

$$\text{Int}[\text{Sqrt}[a x + b \text{Sqrt}[c + (a^2 x^2)/b^2]], x]$$

output

$$\left(\frac{-2 b^2 c}{\text{Sqrt}[a x + b \text{Sqrt}[c + (a^2 x^2)/b^2]]} + \frac{2(a x + b \text{Sqrt}[c + (a^2 x^2)/b^2])^{3/2}}{3}\right)/(2 a)$$

### Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2542 `Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2]))^(n_)^(p_.), x_Symbol] := Simp[1/(2*e) Subst[Int[(g + h*x^n)^p*((d^2 + a*f^2 - 2*d*x + x^2)/(d - x)^2), x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]`

### Maple [F]

$$\int \sqrt{xa + b\sqrt{c + \frac{a^2x^2}{b^2}}} dx$$

input `int((x*a+b*(c+a^2*x^2/b^2)^(1/2))^(1/2),x)`

output `int((x*a+b*(c+a^2*x^2/b^2)^(1/2))^(1/2),x)`

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.86

$$\int \sqrt{ax + b\sqrt{c + \frac{a^2x^2}{b^2}}} dx = \frac{2 \left( 2ax - b\sqrt{\frac{a^2x^2 + b^2c}{b^2}} \right) \sqrt{ax + b\sqrt{\frac{a^2x^2 + b^2c}{b^2}}}{3a}$$

input `integrate((a*x+b*(c+a^2*x^2/b^2)^(1/2))^(1/2),x, algorithm="fricas")`

output  $\frac{2}{3}(2ax - b\sqrt{(a^2x^2 + b^2c)/b^2})\sqrt{ax + b\sqrt{(a^2x^2 + b^2c)/b^2}}/a$

### Sympy [F]

$$\int \sqrt{ax + b\sqrt{c + \frac{a^2x^2}{b^2}}} dx = \int \sqrt{ax + b\sqrt{\frac{a^2x^2}{b^2} + c}} dx$$

input `integrate((a*x+b*(c+a**2*x**2/b**2)**(1/2))**(1/2),x)`

output `Integral(sqrt(a*x + b*sqrt(a**2*x**2/b**2 + c)), x)`

### Maxima [F]

$$\int \sqrt{ax + b\sqrt{c + \frac{a^2x^2}{b^2}}} dx = \int \sqrt{ax + \sqrt{\frac{a^2x^2}{b^2} + cb}} dx$$

input `integrate((a*x+b*(c+a^2*x^2/b^2)^(1/2))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(a*x + sqrt(a^2*x^2/b^2 + c)*b), x)`

### Giac [F]

$$\int \sqrt{ax + b\sqrt{c + \frac{a^2x^2}{b^2}}} dx = \int \sqrt{ax + \sqrt{\frac{a^2x^2}{b^2} + cb}} dx$$

input `integrate((a*x+b*(c+a^2*x^2/b^2)^(1/2))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(a*x + sqrt(a^2*x^2/b^2 + c)*b), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \sqrt{ax + b\sqrt{c + \frac{a^2x^2}{b^2}}} dx = \int \sqrt{ax + b\sqrt{c + \frac{a^2x^2}{b^2}}} dx$$

input `int((a*x + b*(c + (a^2*x^2)/b^2)^(1/2))^(1/2), x)`output `int((a*x + b*(c + (a^2*x^2)/b^2)^(1/2))^(1/2), x)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.65

$$\int \sqrt{ax + b\sqrt{c + \frac{a^2x^2}{b^2}}} dx = \frac{2\sqrt{\sqrt{a^2x^2 + b^2c} + ax}(-\sqrt{a^2x^2 + b^2c} + 2ax)}{3a}$$

input `int((a*x+b*(c+a^2*x^2/b^2)^(1/2))^(1/2), x)`output `(2*sqrt(sqrt(a**2*x**2 + b**2*c) + a*x)*(- sqrt(a**2*x**2 + b**2*c) + 2*a*x))/(3*a)`

### 3.16 $\int (a + x^2)^2 \left(x + \sqrt{a + x^2}\right)^n dx$

Optimal result	150
Mathematica [A] (verified)	151
Rubi [A] (verified)	151
Maple [C] (verified)	153
Fricas [A] (verification not implemented)	153
Sympy [F(-1)]	154
Maxima [F]	154
Giac [F]	154
Mupad [F(-1)]	155
Reduce [B] (verification not implemented)	155

#### Optimal result

Integrand size = 21, antiderivative size = 164

$$\int (a + x^2)^2 \left(x + \sqrt{a + x^2}\right)^n dx = -\frac{a^5(x + \sqrt{a + x^2})^{-5+n}}{32(5 - n)} - \frac{5a^4(x + \sqrt{a + x^2})^{-3+n}}{32(3 - n)} - \frac{5a^3(x + \sqrt{a + x^2})^{-1+n}}{16(1 - n)} + \frac{5a^2(x + \sqrt{a + x^2})^{1+n}}{16(1 + n)} + \frac{5a(x + \sqrt{a + x^2})^{3+n}}{32(3 + n)} + \frac{(x + \sqrt{a + x^2})^{5+n}}{32(5 + n)}$$

output

```
-1/32*a^5*(x+(x^2+a)^(1/2))^(5-n)/(5-n)-5*a^4*(x+(x^2+a)^(1/2))^(3-n)/(96-32*n)-5*a^3*(x+(x^2+a)^(1/2))^(1+n)/(16-16*n)+5*a^2*(x+(x^2+a)^(1/2))^(1+n)/(16+16*n)+5*a*(x+(x^2+a)^(1/2))^(3+n)/(96+32*n)+(x+(x^2+a)^(1/2))^(5+n)/(160+32*n)
```

**Mathematica [A] (verified)**

Time = 0.69 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.84

$$\int (a+x^2)^2 (x+\sqrt{a+x^2})^n dx = \frac{1}{32} (x+\sqrt{a+x^2})^{-5+n} \left( \frac{a^5}{-5+n} + \frac{5a^4(x+\sqrt{a+x^2})^2}{-3+n} \right. \\ \left. + \frac{10a^3(x+\sqrt{a+x^2})^4}{-1+n} + \frac{10a^2(x+\sqrt{a+x^2})^6}{1+n} \right. \\ \left. + \frac{5a(x+\sqrt{a+x^2})^8}{3+n} + \frac{(x+\sqrt{a+x^2})^{10}}{5+n} \right)$$

input `Integrate[(a + x^2)^2*(x + Sqrt[a + x^2])^n,x]`

output `((x + Sqrt[a + x^2])^(-5 + n)*(a^5/(-5 + n) + (5*a^4*(x + Sqrt[a + x^2])^2)/(-3 + n) + (10*a^3*(x + Sqrt[a + x^2])^4)/(-1 + n) + (10*a^2*(x + Sqrt[a + x^2])^6)/(1 + n) + (5*a*(x + Sqrt[a + x^2])^8)/(3 + n) + (x + Sqrt[a + x^2])^10/(5 + n))/32`

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.95, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {2547, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a+x^2)^2 (\sqrt{a+x^2}+x)^n dx \\ \downarrow 2547 \\ \frac{1}{32} \int (x+\sqrt{x^2+a})^{n-6} \left( (x+\sqrt{x^2+a})^2 + a \right)^5 d(x+\sqrt{x^2+a}) \\ \downarrow 244$$



$$\frac{1}{32} \int \left( a^5 (x + \sqrt{x^2 + a})^{n-6} + 5a^4 (x + \sqrt{x^2 + a})^{n-4} + 10a^3 (x + \sqrt{x^2 + a})^{n-2} + 10a^2 (x + \sqrt{x^2 + a})^n + 5a \right) dx$$

↓ 2009

$$\frac{1}{32} \left( -\frac{a^5 (\sqrt{a+x^2} + x)^{n-5}}{5-n} - \frac{5a^4 (\sqrt{a+x^2} + x)^{n-3}}{3-n} - \frac{10a^3 (\sqrt{a+x^2} + x)^{n-1}}{1-n} + \frac{10a^2 (\sqrt{a+x^2} + x)^{n+1}}{n+1} + \dots \right)$$

input `Int[(a + x^2)^2*(x + Sqrt[a + x^2])^n,x]`

output `((-(a^5*(x + Sqrt[a + x^2])^(-5 + n))/(5 - n)) - (5*a^4*(x + Sqrt[a + x^2])^(-3 + n))/(3 - n) - (10*a^3*(x + Sqrt[a + x^2])^(-1 + n))/(1 - n) + (10*a^2*(x + Sqrt[a + x^2])^(1 + n))/(1 + n) + (5*a*(x + Sqrt[a + x^2])^(3 + n))/(3 + n) + (x + Sqrt[a + x^2])^(5 + n)/(5 + n))/32`

### Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2547 `Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

**Maple [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 3.

Time = 0.17 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.32

method	result
meijerg	$\frac{2^n x^{5+n} \operatorname{hypergeom}\left(\left[-\frac{n}{2}, -\frac{5}{2} - \frac{n}{2}, \frac{1}{2} - \frac{n}{2}\right], \left[1-n, -\frac{3}{2} - \frac{n}{2}\right], -\frac{a}{x^2}\right)}{5+n} + \frac{2^{1+n} a x^{3+n} \operatorname{hypergeom}\left(\left[-\frac{n}{2}, -\frac{3}{2} - \frac{n}{2}, \frac{1}{2} - \frac{n}{2}\right], \left[1-n, -\frac{1}{2} - \frac{n}{2}\right], -\frac{a}{x^2}\right)}{3+n}$

input `int((x^2+a)^2*(x+(x^2+a)^(1/2))^n,x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & 2^n/(5+n)*x^{(5+n)}*\operatorname{hypergeom}\left(\left[-1/2*n, -5/2-1/2*n, 1/2-1/2*n\right], \left[1-n, -3/2-1/2*n\right], \right. \\ & \left. -a/x^2\right)+2^{(1+n)}*a/(3+n)*x^{(3+n)}*\operatorname{hypergeom}\left(\left[-1/2*n, -3/2-1/2*n, 1/2-1/2*n\right], \left[1-n, -1/2-1/2*n\right], \right. \\ & \left. -a/x^2\right)+1/4*a^{(5/2+1/2*n)}/\operatorname{Pi}^{(1/2)}*n*(8*\operatorname{Pi}^{(1/2)}/(1+n)/n*x^{(1+n)}*a^{(-1/2-1/2*n)} \\ & *(a/x^{2*n+n-1})/(-2+2*n)*((1+a/x^2)^{(1/2)+1})^{(-1+n)}+4*\operatorname{Pi}^{(1/2)}/(1+n)/n*x^{(1+n)} \\ & *a^{(-1/2-1/2*n)}*(1+a/x^2)^{(1/2)}*((1+a/x^2)^{(1/2)+1})^{(-1+n)} \end{aligned}$$

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.96

$$\int (a+x^2)^2 \left(x + \sqrt{a+x^2}\right)^n dx = \frac{(5(n^4 - 10n^2 + 9)x^5 + 10(an^4 - 16an^2 + 15a)x^3 + 5(a^2n^4 - 22a^2n^2 + 45a^2)x - (a^2n^5 - 30a^2n^3 - n^6 + 35n^4 + 259n^2 - 225)) \sqrt{x^2 + a}}{n^6 - 35n^4 + 259n^2 - 225}$$

input `integrate((x^2+a)^2*(x+(x^2+a)^(1/2))^n,x, algorithm="fricas")`

output 
$$\begin{aligned} & -(5*(n^4 - 10*n^2 + 9)*x^5 + 10*(a*n^4 - 16*a*n^2 + 15*a)*x^3 + 5*(a^2*n^4 \\ & - 22*a^2*n^2 + 45*a^2)*x - (a^2*n^5 - 30*a^2*n^3 + (n^5 - 10*n^3 + 9*n)*x^4 \\ & + 149*a^2*n + 2*(a*n^5 - 20*a*n^3 + 19*a*n)*x^2)*\operatorname{sqrt}(x^2 + a))*(x + \operatorname{sqrt}(x^2 + a))^n \\ & / (n^6 - 35*n^4 + 259*n^2 - 225) \end{aligned}$$

**Sympy [F(-1)]**

Timed out.

$$\int (a + x^2)^2 (x + \sqrt{a + x^2})^n dx = \text{Timed out}$$

input `integrate((x**2+a)**2*(x+(x**2+a)**(1/2))**n,x)`

output `Timed out`

**Maxima [F]**

$$\int (a + x^2)^2 (x + \sqrt{a + x^2})^n dx = \int (x^2 + a)^2 (x + \sqrt{x^2 + a})^n dx$$

input `integrate((x^2+a)^2*(x+(x^2+a)^(1/2))^n,x, algorithm="maxima")`

output `integrate((x^2 + a)^2*(x + sqrt(x^2 + a))^n, x)`

**Giac [F]**

$$\int (a + x^2)^2 (x + \sqrt{a + x^2})^n dx = \int (x^2 + a)^2 (x + \sqrt{x^2 + a})^n dx$$

input `integrate((x^2+a)^2*(x+(x^2+a)^(1/2))^n,x, algorithm="giac")`

output `integrate((x^2 + a)^2*(x + sqrt(x^2 + a))^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (a + x^2)^2 (x + \sqrt{a + x^2})^n dx = \int (x^2 + a)^2 (x + \sqrt{x^2 + a})^n dx$$

input `int((a + x^2)^2*(x + (a + x^2)^(1/2))^n,x)`output `int((a + x^2)^2*(x + (a + x^2)^(1/2))^n, x)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.34

$$\int (a + x^2)^2 (x + \sqrt{a + x^2})^n dx$$

$$= \frac{(\sqrt{x^2 + a} + x)^n (\sqrt{x^2 + a} a^2 n^5 - 30\sqrt{x^2 + a} a^2 n^3 + 149\sqrt{x^2 + a} a^2 n + 2\sqrt{x^2 + a} a n^5 x^2 - 40\sqrt{x^2 + a} a n^3 x - 10\sqrt{x^2 + a} a n^3 x^3 + 9\sqrt{x^2 + a} a n^3 x^4 - 5a^2 n^4 x + 110a^2 n^4 x^2 - 225a^2 n^4 x^3 - 10a^2 n^4 x^4 + 160a^2 n^4 x^5 - 150a^2 n^4 x^6 - 5n^4 x^5 + 50n^4 x^5 - 45x^5)}{(n^6 - 35n^4 + 259n^2 - 225)}$$

input `int((x^2+a)^2*(x+(x^2+a)^(1/2))^n,x)`output `((sqrt(a + x**2) + x)**n*(sqrt(a + x**2)*a**2*n**5 - 30*sqrt(a + x**2)*a**2*n**3 + 149*sqrt(a + x**2)*a**2*n + 2*sqrt(a + x**2)*a*n**5*x**2 - 40*sqrt(a + x**2)*a*n**3*x**3 + 38*sqrt(a + x**2)*a*n*x**2 + sqrt(a + x**2)*n**5*x**4 - 10*sqrt(a + x**2)*n**3*x**4 + 9*sqrt(a + x**2)*n*x**4 - 5*a**2*n**4*x + 110*a**2*n**2*x - 225*a**2*x - 10*a*n**4*x**3 + 160*a*n**2*x**3 - 150*a*x**3 - 5*n**4*x**5 + 50*n**2*x**5 - 45*x**5))/(n**6 - 35*n**4 + 259*n**2 - 225)`

### 3.17 $\int (a + x^2) \left(x + \sqrt{a + x^2}\right)^n dx$

Optimal result	156
Mathematica [A] (verified)	156
Rubi [A] (verified)	157
Maple [C] (verified)	158
Fricas [A] (verification not implemented)	159
Sympy [B] (verification not implemented)	159
Maxima [F]	160
Giac [F]	161
Mupad [F(-1)]	161
Reduce [B] (verification not implemented)	161

#### Optimal result

Integrand size = 19, antiderivative size = 108

$$\int (a + x^2) \left(x + \sqrt{a + x^2}\right)^n dx = -\frac{a^3(x + \sqrt{a + x^2})^{-3+n}}{8(3 - n)} - \frac{3a^2(x + \sqrt{a + x^2})^{-1+n}}{8(1 - n)} + \frac{3a(x + \sqrt{a + x^2})^{1+n}}{8(1 + n)} + \frac{(x + \sqrt{a + x^2})^{3+n}}{8(3 + n)}$$

output

$$-1/8*a^3*(x+(x^2+a)^(1/2))^{(-3+n)/(3-n)}-3*a^2*(x+(x^2+a)^(1/2))^{(-1+n)/(8-8*n)}+3*a*(x+(x^2+a)^(1/2))^{(1+n)/(8+8*n)}+(x+(x^2+a)^(1/2))^{(3+n)/(24+8*n)}$$

#### Mathematica [A] (verified)

Time = 0.43 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.85

$$\int (a + x^2) \left(x + \sqrt{a + x^2}\right)^n dx = \frac{1}{8} \left(x + \sqrt{a + x^2}\right)^{-3+n} \left( \frac{a^3}{-3 + n} + \frac{3a^2(x + \sqrt{a + x^2})^2}{-1 + n} + \frac{3a(x + \sqrt{a + x^2})^4}{1 + n} + \frac{(x + \sqrt{a + x^2})^6}{3 + n} \right)$$

input

```
Integrate[(a + x^2)*(x + Sqrt[a + x^2])^n,x]
```

output

$$\left( (x + \sqrt{a + x^2})^{-3 + n} (a^3 / (-3 + n) + (3a^2(x + \sqrt{a + x^2}))^2 / (-1 + n) + (3a(x + \sqrt{a + x^2}))^4 / (1 + n) + (x + \sqrt{a + x^2})^6 / (3 + n)) \right) / 8$$

**Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.95, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$ , Rules used = {2547, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + x^2) (\sqrt{a + x^2} + x)^n dx$$

$$\downarrow 2547$$

$$\frac{1}{8} \int (x + \sqrt{x^2 + a})^{n-4} \left( (x + \sqrt{x^2 + a})^2 + a \right)^3 d(x + \sqrt{x^2 + a})$$

$$\downarrow 244$$

$$\frac{1}{8} \int \left( a^3 (x + \sqrt{x^2 + a})^{n-4} + 3a^2 (x + \sqrt{x^2 + a})^{n-2} + 3a (x + \sqrt{x^2 + a})^n + (x + \sqrt{x^2 + a})^{n+2} \right) d(x + \sqrt{x^2 + a})$$

$$\downarrow 2009$$

$$\frac{1}{8} \left( -\frac{a^3 (\sqrt{a + x^2} + x)^{n-3}}{3 - n} - \frac{3a^2 (\sqrt{a + x^2} + x)^{n-1}}{1 - n} + \frac{3a (\sqrt{a + x^2} + x)^{n+1}}{n + 1} + \frac{(\sqrt{a + x^2} + x)^{n+3}}{n + 3} \right)$$

input

$$\text{Int}[(a + x^2)*(x + \text{Sqrt}[a + x^2])^n, x]$$

output

$$\left( -((a^3(x + \text{Sqrt}[a + x^2])^{-3 + n}) / (3 - n)) - (3a^2(x + \text{Sqrt}[a + x^2])^{-1 + n}) / (1 - n) + (3a(x + \text{Sqrt}[a + x^2])^{1 + n}) / (1 + n) + (x + \text{Sqrt}[a + x^2])^{3 + n} / (3 + n) \right) / 8$$

**Defintions of rubi rules used**

```
rule 244 Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[Expand
Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p
, 0]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2547 Int[((g_) + (i_)*(x_)^2)^(m_)*((d_) + (e_)*(x_) + (f_)*Sqrt[(a_) + (c_
.)*(x_)^2])^(n_), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m
Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))],
x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n},
x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (Intege
rQ[m] || GtQ[i/c, 0])
```

**Maple [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 3.

Time = 0.04 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.55

method	result
meijerg	$\frac{2^n x^{3+n} \operatorname{hypergeom}\left(\left[-\frac{n}{2}, -\frac{3}{2} - \frac{n}{2}, \frac{1}{2} - \frac{n}{2}\right], \left[1-n, -\frac{1}{2} - \frac{n}{2}\right], -\frac{a}{x^2}\right)}{3+n} + \frac{a^{\frac{3}{2} + \frac{n}{2}} n \left( \frac{8\sqrt{\pi} x^{1+n} a^{-\frac{1}{2} - \frac{n}{2}} \left(\frac{an}{x^2} + n - 1\right) \left(\sqrt{1 + \frac{a}{x^2}} + 1\right)^{-1+n}}{(1+n)n(-2+2n)} + \dots \right)}{4\sqrt{\pi}}$

```
input int((x^2+a)*(x+(x^2+a)^(1/2))^n,x,method=_RETURNVERBOSE)
```

```
output 2^n/(3+n)*x^(3+n)*hypergeom([-1/2*n, -3/2-1/2*n, 1/2-1/2*n], [1-n, -1/2-1/2*n]
, -a/x^2)+1/4*a^(3/2+1/2*n)/Pi^(1/2)*n*(8*Pi^(1/2)/(1+n)/n*x^(1+n)*a^(-1/2-
1/2*n)*(a/x^2+n-1)/(-2+2*n)*((1+a/x^2)^(1/2)+1)^(-1+n)+4*Pi^(1/2)/(1+n)/
n*x^(1+n)*a^(-1/2-1/2*n)*(1+a/x^2)^(1/2)*((1+a/x^2)^(1/2)+1)^(-1+n))
```

**Fricas [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.72

$$\int (a + x^2) \left( x + \sqrt{a + x^2} \right)^n dx = \frac{(3(n^2 - 1)x^3 + 3(an^2 - 3a)x - (an^3 + (n^3 - n)x^2 - 7an)\sqrt{x^2 + a})(x + \sqrt{x^2 + a})^n}{n^4 - 10n^2 + 9}$$

input `integrate((x^2+a)*(x+(x^2+a)^(1/2))^n,x, algorithm="fricas")`

output `-(3*(n^2 - 1)*x^3 + 3*(a*n^2 - 3*a)*x - (a*n^3 + (n^3 - n)*x^2 - 7*a*n)*sqrt(x^2 + a))*(x + sqrt(x^2 + a))^n/(n^4 - 10*n^2 + 9)`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 2237 vs. 2(85) = 170.

Time = 16.38 (sec) , antiderivative size = 15302, normalized size of antiderivative = 141.69

$$\int (a + x^2) \left( x + \sqrt{a + x^2} \right)^n dx = \text{Too large to display}$$

input `integrate((x**2+a)*(x+(x**2+a)**(1/2))**n,x)`



output

```

a*Piecewise((2*a**(9/2)*a**(n/2 + 1/2)*n*cosh(n*asinh(x/sqrt(a)) + asinh(x
/sqrt(a)))*gamma(1 - n/2)/(2*a**(9/2)*n**2*gamma(1 - n/2) - 2*a**(9/2)*gam
ma(1 - n/2) + 2*a**(7/2)*n**2*x**2*gamma(1 - n/2) - 2*a**(7/2)*x**2*gamma(
1 - n/2)) - 2*a**(9/2)*a**(n/2 + 1/2)*n*gamma(1 - n/2)/(2*a**(9/2)*n**2*ga
mma(1 - n/2) - 2*a**(9/2)*gamma(1 - n/2) + 2*a**(7/2)*n**2*x**2*gamma(1 -
n/2) - 2*a**(7/2)*x**2*gamma(1 - n/2)) - 2*a**(7/2)*a**(n/2 + 1/2)*n*x**2*
sqrt(a/x**2 + 1)*sinh(n*asinh(x/sqrt(a)) + asinh(x/sqrt(a)))*gamma(1 - n/2
)/(2*a**(9/2)*n**2*gamma(1 - n/2) - 2*a**(9/2)*gamma(1 - n/2) + 2*a**(7/2)
*n**2*x**2*gamma(1 - n/2) - 2*a**(7/2)*x**2*gamma(1 - n/2)) + 4*a**(7/2)*a
**(n/2 + 1/2)*n*x**2*cosh(n*asinh(x/sqrt(a)) + asinh(x/sqrt(a)))*gamma(1 -
n/2)/(2*a**(9/2)*n**2*gamma(1 - n/2) - 2*a**(9/2)*gamma(1 - n/2) + 2*a**(
7/2)*n**2*x**2*gamma(1 - n/2) - 2*a**(7/2)*x**2*gamma(1 - n/2)) - 2*a**(7/
2)*a**(n/2 + 1/2)*n*x**2*gamma(1 - n/2)/(2*a**(9/2)*n**2*gamma(1 - n/2) -
2*a**(9/2)*gamma(1 - n/2) + 2*a**(7/2)*n**2*x**2*gamma(1 - n/2) - 2*a**(7/
2)*x**2*gamma(1 - n/2)) - 2*a**(7/2)*a**(n/2 + 1/2)*x**2*sqrt(a/x**2 + 1)*
sinh(n*asinh(x/sqrt(a)) + asinh(x/sqrt(a)))*gamma(1 - n/2)/(2*a**(9/2)*n**
2*gamma(1 - n/2) - 2*a**(9/2)*gamma(1 - n/2) + 2*a**(7/2)*n**2*x**2*gamma(
1 - n/2) - 2*a**(7/2)*x**2*gamma(1 - n/2)) + 2*a**(7/2)*a**(n/2 + 1/2)*x**
2*cosh(n*asinh(x/sqrt(a)) + asinh(x/sqrt(a)))*gamma(1 - n/2)/(2*a**(9/2)*n
**2*gamma(1 - n/2) - 2*a**(9/2)*gamma(1 - n/2) + 2*a**(7/2)*n**2*x**2*g...

```

## Maxima [F]

$$\int (a + x^2) (x + \sqrt{a + x^2})^n dx = \int (x^2 + a) (x + \sqrt{x^2 + a})^n dx$$

input

```
integrate((x^2+a)*(x+(x^2+a)^(1/2))^n,x, algorithm="maxima")
```

output

```
integrate((x^2 + a)*(x + sqrt(x^2 + a))^n, x)
```

**Giac [F]**

$$\int (a + x^2) (x + \sqrt{a + x^2})^n dx = \int (x^2 + a) (x + \sqrt{x^2 + a})^n dx$$

input `integrate((x^2+a)*(x+(x^2+a)^(1/2))^n,x, algorithm="giac")`

output `integrate((x^2 + a)*(x + sqrt(x^2 + a))^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (a + x^2) (x + \sqrt{a + x^2})^n dx = \int (x^2 + a) (x + \sqrt{x^2 + a})^n dx$$

input `int((a + x^2)*(x + (a + x^2)^(1/2))^n,x)`

output `int((a + x^2)*(x + (a + x^2)^(1/2))^n, x)`

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.87

$$\int (a + x^2) (x + \sqrt{a + x^2})^n dx = \frac{(\sqrt{x^2 + a} + x)^n (\sqrt{x^2 + a} a n^3 - 7\sqrt{x^2 + a} a n + \sqrt{x^2 + a} n^3 x^2 - \sqrt{x^2 + a} n x^2 - 3a n^2 x + 9a x - 3n^2 x^3)}{n^4 - 10n^2 + 9}$$

input `int((x^2+a)*(x+(x^2+a)^(1/2))^n,x)`

output `((sqrt(a + x**2) + x)**n*(sqrt(a + x**2)*a*n**3 - 7*sqrt(a + x**2)*a*n + sqrt(a + x**2)*n**3*x**2 - sqrt(a + x**2)*n*x**2 - 3*a*n**2*x + 9*a*x - 3*n**2*x**3 + 3*x**3))/(n**4 - 10*n**2 + 9)`

### 3.18 $\int \left(x + \sqrt{a + x^2}\right)^n dx$

Optimal result	162
Mathematica [A] (verified)	162
Rubi [A] (verified)	163
Maple [B] (verified)	164
Fricas [A] (verification not implemented)	165
Sympy [B] (verification not implemented)	165
Maxima [F]	166
Giac [F]	167
Mupad [F(-1)]	167
Reduce [B] (verification not implemented)	167

#### Optimal result

Integrand size = 13, antiderivative size = 52

$$\int \left(x + \sqrt{a + x^2}\right)^n dx = -\frac{a(x + \sqrt{a + x^2})^{-1+n}}{2(1-n)} + \frac{(x + \sqrt{a + x^2})^{1+n}}{2(1+n)}$$

output `-1/2*a*(x+(x^2+a)^(1/2))^(1+n)/(1+n)+(x+(x^2+a)^(1/2))^(1+n)/(2+2*n)`

#### Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.83

$$\int \left(x + \sqrt{a + x^2}\right)^n dx = \frac{(x + \sqrt{a + x^2})^{-1+n} (an + (-1 + n)x(x + \sqrt{a + x^2}))}{-1 + n^2}$$

input `Integrate[(x + Sqrt[a + x^2])^n,x]`

output `((x + Sqrt[a + x^2])^(-1 + n)*(a*n + (-1 + n)*x*(x + Sqrt[a + x^2])))/(-1 + n^2)`

**Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.98, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {2542, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\sqrt{a+x^2} + x)^n dx$$

$$\downarrow \text{2542}$$

$$\frac{1}{2} \int (x + \sqrt{x^2 + a})^{n-2} \left( (x + \sqrt{x^2 + a})^2 + a \right) d(x + \sqrt{x^2 + a})$$

$$\downarrow \text{244}$$

$$\frac{1}{2} \int \left( a(x + \sqrt{x^2 + a})^{n-2} + (x + \sqrt{x^2 + a})^n \right) d(x + \sqrt{x^2 + a})$$

$$\downarrow \text{2009}$$

$$\frac{1}{2} \left( \frac{(\sqrt{a+x^2} + x)^{n+1}}{n+1} - \frac{a(\sqrt{a+x^2} + x)^{n-1}}{1-n} \right)$$

input `Int[(x + Sqrt[a + x^2])^n,x]`

output `((-(a*(x + Sqrt[a + x^2])^(-1 + n))/(1 - n)) + (x + Sqrt[a + x^2])^(1 + n))/(1 + n)/2`

## Definitions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2542 `Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2]))^(n_)^(p_.), x_Symbol] := Simp[1/(2*e) Subst[Int[(g + h*x^n)^p*((d^2 + a*f^2 - 2*d*x + x^2)/(d - x)^2), x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]`

## Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 119 vs.  $2(45) = 90$ .

Time = 0.03 (sec) , antiderivative size = 120, normalized size of antiderivative = 2.31

method	result	size
meijerg	$\frac{a^{\frac{1}{2} + \frac{n}{2}} n \left( \frac{8\sqrt{\pi} x^{1+n} a^{-\frac{1}{2} - \frac{n}{2}} \left( \frac{a^{\frac{n}{2}} + n - 1}{x^2} \right) \left( \sqrt{1 + \frac{a}{x^2}} + 1 \right)^{-1+n}}{(1+n)n(-2+2n)} + \frac{4\sqrt{\pi} x^{1+n} a^{-\frac{1}{2} - \frac{n}{2}} \sqrt{1 + \frac{a}{x^2}} \left( \sqrt{1 + \frac{a}{x^2}} + 1 \right)^{-1+n}}{(1+n)n} \right)}{4\sqrt{\pi}}$	120

input `int((x+(x^2+a)^(1/2))^n,x,method=_RETURNVERBOSE)`

output `1/4*a^(1/2+1/2*n)/Pi^(1/2)*n*(8*Pi^(1/2)/(1+n)/n*x^(1+n)*a^(-1/2-1/2*n)*(a/x^2*n+n-1)/(-2+2*n)*((1+a/x^2)^(1/2)+1)^(-1+n)+4*Pi^(1/2)/(1+n)/n*x^(1+n)*a^(-1/2-1/2*n)*(1+a/x^2)^(1/2)*((1+a/x^2)^(1/2)+1)^(-1+n)`

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.62

$$\int \left(x + \sqrt{a + x^2}\right)^n dx = \frac{(\sqrt{x^2 + a} - x)(x + \sqrt{x^2 + a})^n}{n^2 - 1}$$

input `integrate((x+(x^2+a)^(1/2))^n,x, algorithm="fricas")`

output `(sqrt(x^2 + a)*n - x)*(x + sqrt(x^2 + a))^n/(n^2 - 1)`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 2236 vs. 2(37) = 74.

Time = 1.99 (sec) , antiderivative size = 2236, normalized size of antiderivative = 43.00

$$\int \left(x + \sqrt{a + x^2}\right)^n dx = \text{Too large to display}$$

input `integrate((x+(x**2+a)**(1/2))**n,x)`

output

```
Piecewise((2*a**(9/2)*a**(n/2 + 1/2)*n*cosh(n*asinh(x/sqrt(a)) + asinh(x/sqrt(a)))*gamma(1 - n/2)/(2*a**(9/2)*n**2*gamma(1 - n/2) - 2*a**(9/2)*gamma(1 - n/2) + 2*a**(7/2)*n**2*x**2*gamma(1 - n/2) - 2*a**(7/2)*x**2*gamma(1 - n/2)) - 2*a**(9/2)*a**(n/2 + 1/2)*n*gamma(1 - n/2)/(2*a**(9/2)*n**2*gamma(1 - n/2) - 2*a**(9/2)*gamma(1 - n/2) + 2*a**(7/2)*n**2*x**2*gamma(1 - n/2) - 2*a**(7/2)*x**2*gamma(1 - n/2)) - 2*a**(7/2)*a**(n/2 + 1/2)*n*x**2*sqrt(a/x**2 + 1)*sinh(n*asinh(x/sqrt(a)) + asinh(x/sqrt(a)))*gamma(1 - n/2)/(2*a**(9/2)*n**2*gamma(1 - n/2) - 2*a**(9/2)*gamma(1 - n/2) + 2*a**(7/2)*n**2*x**2*gamma(1 - n/2) - 2*a**(7/2)*x**2*gamma(1 - n/2)) + 4*a**(7/2)*a**(n/2 + 1/2)*n*x**2*cosh(n*asinh(x/sqrt(a)) + asinh(x/sqrt(a)))*gamma(1 - n/2)/(2*a**(9/2)*n**2*gamma(1 - n/2) - 2*a**(9/2)*gamma(1 - n/2) + 2*a**(7/2)*n**2*x**2*gamma(1 - n/2) - 2*a**(7/2)*x**2*gamma(1 - n/2)) - 2*a**(7/2)*a**(n/2 + 1/2)*n*x**2*gamma(1 - n/2)/(2*a**(9/2)*n**2*gamma(1 - n/2) - 2*a**(9/2)*gamma(1 - n/2) + 2*a**(7/2)*n**2*x**2*gamma(1 - n/2) - 2*a**(7/2)*x**2*gamma(1 - n/2)) - 2*a**(7/2)*a**(n/2 + 1/2)*x**2*sqrt(a/x**2 + 1)*sinh(n*asinh(x/sqrt(a)) + asinh(x/sqrt(a)))*gamma(1 - n/2)/(2*a**(9/2)*n**2*gamma(1 - n/2) - 2*a**(9/2)*gamma(1 - n/2) + 2*a**(7/2)*n**2*x**2*gamma(1 - n/2) - 2*a**(7/2)*x**2*gamma(1 - n/2)) + 2*a**(7/2)*a**(n/2 + 1/2)*x**2*cosh(n*asinh(x/sqrt(a)) + asinh(x/sqrt(a)))*gamma(1 - n/2)/(2*a**(9/2)*n**2*gamma(1 - n/2) - 2*a**(9/2)*gamma(1 - n/2) + 2*a**(7/2)*n**2*x**2*gam...
```

**Maxima [F]**

$$\int \left(x + \sqrt{a + x^2}\right)^n dx = \int \left(x + \sqrt{x^2 + a}\right)^n dx$$

input

```
integrate((x+(x^2+a)^(1/2))^n,x, algorithm="maxima")
```

output

```
integrate((x + sqrt(x^2 + a))^n, x)
```

**Giac [F]**

$$\int (x + \sqrt{a + x^2})^n dx = \int (x + \sqrt{x^2 + a})^n dx$$

input `integrate((x+(x^2+a)^(1/2))^n,x, algorithm="giac")`

output `integrate((x + sqrt(x^2 + a))^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (x + \sqrt{a + x^2})^n dx = \int (x + \sqrt{x^2 + a})^n dx$$

input `int((x + (a + x^2)^(1/2))^n,x)`

output `int((x + (a + x^2)^(1/2))^n, x)`

**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.58

$$\int (x + \sqrt{a + x^2})^n dx = \frac{(\sqrt{x^2 + a} + x)^n (\sqrt{x^2 + a} n - x)}{n^2 - 1}$$

input `int((x+(x^2+a)^(1/2))^n,x)`

output `((sqrt(a + x**2) + x)**n*(sqrt(a + x**2)*n - x))/(n**2 - 1)`



**3.19**  $\int \frac{(x + \sqrt{a + x^2})^n}{a + x^2} dx$

Optimal result	168
Mathematica [A] (verified)	168
Rubi [A] (verified)	169
Maple [F]	170
Fricas [F]	170
Sympy [F]	171
Maxima [F]	171
Giac [F]	171
Mupad [F(-1)]	172
Reduce [F]	172

**Optimal result**

Integrand size = 21, antiderivative size = 59

$$\int \frac{(x + \sqrt{a + x^2})^n}{a + x^2} dx = \frac{2(x + \sqrt{a + x^2})^{1+n} \operatorname{Hypergeometric2F1}\left(1, \frac{1+n}{2}, \frac{3+n}{2}, -\frac{(x + \sqrt{a + x^2})^2}{a}\right)}{a(1 + n)}$$

output

```
2*(x+(x^2+a)^(1/2))^(1+n)*hypergeom([1, 1/2+1/2*n],[3/2+1/2*n],-(x+(x^2+a)^(1/2))^2/a)/a/(1+n)
```

**Mathematica [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.03

$$\int \frac{(x + \sqrt{a + x^2})^n}{a + x^2} dx = \frac{2(x + \sqrt{a + x^2})^{1+n} \operatorname{Hypergeometric2F1}\left(1, \frac{1+n}{2}, 1 + \frac{1+n}{2}, -\frac{(x + \sqrt{a + x^2})^2}{a}\right)}{a(1 + n)}$$

input `Integrate[(x + Sqrt[a + x^2])^n/(a + x^2), x]`

output `(2*(x + Sqrt[a + x^2])^(1 + n)*Hypergeometric2F1[1, (1 + n)/2, 1 + (1 + n)/2, -((x + Sqrt[a + x^2])^2/a)])/(a*(1 + n))`

### Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {2547, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(\sqrt{a+x^2}+x)^n}{a+x^2} dx$$

$$\downarrow 2547$$

$$2 \int \frac{(x+\sqrt{x^2+a})^n}{(x+\sqrt{x^2+a})^2+a} d(x+\sqrt{x^2+a})$$

$$\downarrow 278$$

$$\frac{2(\sqrt{a+x^2}+x)^{n+1} \text{Hypergeometric2F1}\left(1, \frac{n+1}{2}, \frac{n+3}{2}, -\frac{(x+\sqrt{x^2+a})^2}{a}\right)}{a(n+1)}$$

input `Int[(x + Sqrt[a + x^2])^n/(a + x^2), x]`

output `(2*(x + Sqrt[a + x^2])^(1 + n)*Hypergeometric2F1[1, (1 + n)/2, (3 + n)/2, -((x + Sqrt[a + x^2])^2/a)])/(a*(1 + n))`

## Definitions of rubi rules used

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 2547 `Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

## Maple [F]

$$\int \frac{(x + \sqrt{x^2 + a})^n}{x^2 + a} dx$$

input `int((x+(x^2+a)^(1/2))^n/(x^2+a),x)`

output `int((x+(x^2+a)^(1/2))^n/(x^2+a),x)`

## Fricas [F]

$$\int \frac{(x + \sqrt{a + x^2})^n}{a + x^2} dx = \int \frac{(x + \sqrt{x^2 + a})^n}{x^2 + a} dx$$

input `integrate((x+(x^2+a)^(1/2))^n/(x^2+a),x, algorithm="fricas")`

output `integral((x + sqrt(x^2 + a))^n/(x^2 + a), x)`

**Sympy [F]**

$$\int \frac{(x + \sqrt{a + x^2})^n}{a + x^2} dx = \int \frac{(x + \sqrt{a + x^2})^n}{a + x^2} dx$$

input `integrate((x+(x**2+a)**(1/2))**n/(x**2+a), x)`

output `Integral((x + sqrt(a + x**2))**n/(a + x**2), x)`

**Maxima [F]**

$$\int \frac{(x + \sqrt{a + x^2})^n}{a + x^2} dx = \int \frac{(x + \sqrt{x^2 + a})^n}{x^2 + a} dx$$

input `integrate((x+(x^2+a)^(1/2))^n/(x^2+a), x, algorithm="maxima")`

output `integrate((x + sqrt(x^2 + a))^n/(x^2 + a), x)`

**Giac [F]**

$$\int \frac{(x + \sqrt{a + x^2})^n}{a + x^2} dx = \int \frac{(x + \sqrt{x^2 + a})^n}{x^2 + a} dx$$

input `integrate((x+(x^2+a)^(1/2))^n/(x^2+a), x, algorithm="giac")`

output `integrate((x + sqrt(x^2 + a))^n/(x^2 + a), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(x + \sqrt{a + x^2})^n}{a + x^2} dx = \int \frac{(x + \sqrt{x^2 + a})^n}{x^2 + a} dx$$

input `int((x + (a + x^2)^(1/2))^n/(a + x^2), x)`output `int((x + (a + x^2)^(1/2))^n/(a + x^2), x)`**Reduce [F]**

$$\int \frac{(x + \sqrt{a + x^2})^n}{a + x^2} dx = \int \frac{(\sqrt{x^2 + a} + x)^n}{x^2 + a} dx$$

input `int((x+(x^2+a)^(1/2))^n/(x^2+a), x)`output `int((sqrt(a + x**2) + x)**n/(a + x**2), x)`

**3.20**  $\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^2} dx$

Optimal result	173
Mathematica [A] (verified)	173
Rubi [A] (verified)	174
Maple [F]	175
Fricas [F]	175
Sympy [F]	176
Maxima [F]	176
Giac [F]	176
Mupad [F(-1)]	177
Reduce [F]	177

**Optimal result**

Integrand size = 21, antiderivative size = 59

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^2} dx = \frac{8(x + \sqrt{a + x^2})^{3+n} \operatorname{Hypergeometric2F1}\left(3, \frac{3+n}{2}, \frac{5+n}{2}, -\frac{(x + \sqrt{a + x^2})^2}{a}\right)}{a^3(3 + n)}$$

output `8*(x+(x^2+a)^(1/2))^(3+n)*hypergeom([3, 3/2+1/2*n],[5/2+1/2*n],-(x+(x^2+a)^(1/2))^2/a)/a^3/(3+n)`

**Mathematica [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.03

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^2} dx = \frac{8(x + \sqrt{a + x^2})^{3+n} \operatorname{Hypergeometric2F1}\left(3, \frac{3+n}{2}, 1 + \frac{3+n}{2}, -\frac{(x + \sqrt{a + x^2})^2}{a}\right)}{a^3(3 + n)}$$

input `Integrate[(x + Sqrt[a + x^2])^n/(a + x^2)^2,x]`

output `(8*(x + Sqrt[a + x^2])^(3 + n)*Hypergeometric2F1[3, (3 + n)/2, 1 + (3 + n)/2, -((x + Sqrt[a + x^2])^2/a)])/(a^3*(3 + n))`

### Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {2547, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(\sqrt{a+x^2}+x)^n}{(a+x^2)^2} dx$$

↓ 2547

$$8 \int \frac{(x + \sqrt{x^2 + a})^{n+2}}{\left( (x + \sqrt{x^2 + a})^2 + a \right)^3} d(x + \sqrt{x^2 + a})$$

↓ 278

$$\frac{8(\sqrt{a+x^2}+x)^{n+3} \text{Hypergeometric2F1}\left(3, \frac{n+3}{2}, \frac{n+5}{2}, -\frac{(x+\sqrt{x^2+a})^2}{a}\right)}{a^3(n+3)}$$

input `Int[(x + Sqrt[a + x^2])^n/(a + x^2)^2,x]`

output `(8*(x + Sqrt[a + x^2])^(3 + n)*Hypergeometric2F1[3, (3 + n)/2, (5 + n)/2, -((x + Sqrt[a + x^2])^2/a)])/(a^3*(3 + n))`

## Definitions of rubi rules used

rule 278

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])
```

rule 2547

```
Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])
```

## Maple [F]

$$\int \frac{(x + \sqrt{x^2 + a})^n}{(x^2 + a)^2} dx$$

input

```
int((x+(x^2+a)^(1/2))^n/(x^2+a)^2,x)
```

output

```
int((x+(x^2+a)^(1/2))^n/(x^2+a)^2,x)
```

## Fricas [F]

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^2} dx = \int \frac{(x + \sqrt{x^2 + a})^n}{(x^2 + a)^2} dx$$

input

```
integrate((x+(x^2+a)^(1/2))^n/(x^2+a)^2,x, algorithm="fricas")
```

output

```
integral((x + sqrt(x^2 + a))^n/(x^4 + 2*a*x^2 + a^2), x)
```



**Sympy [F]**

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^2} dx = \int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^2} dx$$

input `integrate((x+(x**2+a)**(1/2))**n/(x**2+a)**2,x)`

output `Integral((x + sqrt(a + x**2))**n/(a + x**2)**2, x)`

**Maxima [F]**

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^2} dx = \int \frac{(x + \sqrt{x^2 + a})^n}{(x^2 + a)^2} dx$$

input `integrate((x+(x^2+a)^(1/2))^n/(x^2+a)^2,x, algorithm="maxima")`

output `integrate((x + sqrt(x^2 + a))^n/(x^2 + a)^2, x)`

**Giac [F]**

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^2} dx = \int \frac{(x + \sqrt{x^2 + a})^n}{(x^2 + a)^2} dx$$

input `integrate((x+(x^2+a)^(1/2))^n/(x^2+a)^2,x, algorithm="giac")`

output `integrate((x + sqrt(x^2 + a))^n/(x^2 + a)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^2} dx = \int \frac{(x + \sqrt{x^2 + a})^n}{(x^2 + a)^2} dx$$

input `int((x + (a + x^2)^(1/2))^n/(a + x^2)^2,x)`output `int((x + (a + x^2)^(1/2))^n/(a + x^2)^2, x)`**Reduce [F]**

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^2} dx = \int \frac{(\sqrt{x^2 + a} + x)^n}{x^4 + 2ax^2 + a^2} dx$$

input `int((x+(x^2+a)^(1/2))^n/(x^2+a)^2,x)`output `int((sqrt(a + x**2) + x)**n/(a**2 + 2*a*x**2 + x**4),x)`

### 3.21 $\int (a + x^2)^2 \left(x - \sqrt{a + x^2}\right)^n dx$

Optimal result	178
Mathematica [A] (verified)	179
Rubi [A] (verified)	179
Maple [F]	181
Fricas [A] (verification not implemented)	181
Sympy [F]	181
Maxima [F]	182
Giac [F]	182
Mupad [F(-1)]	182
Reduce [B] (verification not implemented)	183

#### Optimal result

Integrand size = 23, antiderivative size = 176

$$\int (a + x^2)^2 \left(x - \sqrt{a + x^2}\right)^n dx = -\frac{a^5(x - \sqrt{a + x^2})^{-5+n}}{32(5 - n)} - \frac{5a^4(x - \sqrt{a + x^2})^{-3+n}}{32(3 - n)} - \frac{5a^3(x - \sqrt{a + x^2})^{-1+n}}{16(1 - n)} + \frac{5a^2(x - \sqrt{a + x^2})^{1+n}}{16(1 + n)} + \frac{5a(x - \sqrt{a + x^2})^{3+n}}{32(3 + n)} + \frac{(x - \sqrt{a + x^2})^{5+n}}{32(5 + n)}$$

output

```
-1/32*a^5*(x-(x^2+a)^(1/2))^(5-n)/(5-n)-5*a^4*(x-(x^2+a)^(1/2))^(3-n)/(96-32*n)-5*a^3*(x-(x^2+a)^(1/2))^(1+n)/(16-16*n)+5*a^2*(x-(x^2+a)^(1/2))^(1+n)/(16+16*n)+5*a*(x-(x^2+a)^(1/2))^(3+n)/(96+32*n)+(x-(x^2+a)^(1/2))^(5+n)/(160+32*n)
```

**Mathematica [A] (verified)**

Time = 0.72 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.85

$$\int (a+x^2)^2 (x-\sqrt{a+x^2})^n dx = \frac{1}{32} (x-\sqrt{a+x^2})^{-5+n} \left( \frac{a^5}{-5+n} + \frac{5a^4(x-\sqrt{a+x^2})^2}{-3+n} \right. \\ \left. + \frac{10a^3(x-\sqrt{a+x^2})^4}{-1+n} + \frac{10a^2(x-\sqrt{a+x^2})^6}{1+n} \right. \\ \left. + \frac{5a(x-\sqrt{a+x^2})^8}{3+n} + \frac{(x-\sqrt{a+x^2})^{10}}{5+n} \right)$$

input `Integrate[(a + x^2)^2*(x - Sqrt[a + x^2])^n,x]`

output `((x - Sqrt[a + x^2])^(-5 + n)*(a^5/(-5 + n) + (5*a^4*(x - Sqrt[a + x^2])^2)/(-3 + n) + (10*a^3*(x - Sqrt[a + x^2])^4)/(-1 + n) + (10*a^2*(x - Sqrt[a + x^2])^6)/(1 + n) + (5*a*(x - Sqrt[a + x^2])^8)/(3 + n) + (x - Sqrt[a + x^2])^10/(5 + n))/32`

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.95, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {2547, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a+x^2)^2 (x-\sqrt{a+x^2})^n dx \\ \downarrow 2547 \\ \frac{1}{32} \int (x-\sqrt{x^2+a})^{n-6} \left( (x-\sqrt{x^2+a})^2 + a \right)^5 d(x-\sqrt{x^2+a}) \\ \downarrow 244$$

$$\frac{1}{32} \int \left( a^5 (x - \sqrt{x^2 + a})^{n-6} + 5a^4 (x - \sqrt{x^2 + a})^{n-4} + 10a^3 (x - \sqrt{x^2 + a})^{n-2} + 10a^2 (x - \sqrt{x^2 + a})^n + 5a \right) dx$$

↓ 2009

$$\frac{1}{32} \left( -\frac{a^5 (x - \sqrt{a + x^2})^{n-5}}{5-n} - \frac{5a^4 (x - \sqrt{a + x^2})^{n-3}}{3-n} - \frac{10a^3 (x - \sqrt{a + x^2})^{n-1}}{1-n} + \frac{10a^2 (x - \sqrt{a + x^2})^{n+1}}{n+1} + \dots \right)$$

input `Int[(a + x^2)^2*(x - Sqrt[a + x^2])^n,x]`

output `((-(a^5*(x - Sqrt[a + x^2])^(-5 + n))/(5 - n)) - (5*a^4*(x - Sqrt[a + x^2])^(-3 + n))/(3 - n) - (10*a^3*(x - Sqrt[a + x^2])^(-1 + n))/(1 - n) + (10*a^2*(x - Sqrt[a + x^2])^(1 + n))/(1 + n) + (5*a*(x - Sqrt[a + x^2])^(3 + n))/(3 + n) + (x - Sqrt[a + x^2])^(5 + n)/(5 + n))/32`

### Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2547 `Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

**Maple [F]**

$$\int (x^2 + a)^2 (x - \sqrt{x^2 + a})^n dx$$

input `int((x^2+a)^2*(x-(x^2+a)^(1/2))^n,x)`

output `int((x^2+a)^2*(x-(x^2+a)^(1/2))^n,x)`

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 159, normalized size of antiderivative = 0.90

$$\int (a + x^2)^2 (x - \sqrt{a + x^2})^n dx = \frac{(5(n^4 - 10n^2 + 9)x^5 + 10(an^4 - 16an^2 + 15a)x^3 + 5(a^2n^4 - 22a^2n^2 + 45a^2)x + (a^2n^5 - 30a^2n^3 - n^6 - 35n^4 + 259n^2 - 225))}{n^6 - 35n^4 + 259n^2 - 225}$$

input `integrate((x^2+a)^2*(x-(x^2+a)^(1/2))^n,x, algorithm="fricas")`

output `-(5*(n^4 - 10*n^2 + 9)*x^5 + 10*(a*n^4 - 16*a*n^2 + 15*a)*x^3 + 5*(a^2*n^4 - 22*a^2*n^2 + 45*a^2)*x + (a^2*n^5 - 30*a^2*n^3 + (n^5 - 10*n^3 + 9*n)*x^4 + 149*a^2*n + 2*(a*n^5 - 20*a*n^3 + 19*a*n)*x^2)*sqrt(x^2 + a))*(x - sqrt(x^2 + a))^n/(n^6 - 35*n^4 + 259*n^2 - 225)`

**Sympy [F]**

$$\int (a + x^2)^2 (x - \sqrt{a + x^2})^n dx = \int (a + x^2)^2 (x - \sqrt{a + x^2})^n dx$$

input `integrate((x**2+a)**2*(x-(x**2+a)**(1/2))**n,x)`

output `Integral((a + x**2)**2*(x - sqrt(a + x**2))**n, x)`

**Maxima [F]**

$$\int (a + x^2)^2 (x - \sqrt{a + x^2})^n dx = \int (x^2 + a)^2 (x - \sqrt{x^2 + a})^n dx$$

input `integrate((x^2+a)^2*(x-(x^2+a)^(1/2))^n,x, algorithm="maxima")`

output `integrate((x^2 + a)^2*(x - sqrt(x^2 + a))^n, x)`

**Giac [F]**

$$\int (a + x^2)^2 (x - \sqrt{a + x^2})^n dx = \int (x^2 + a)^2 (x - \sqrt{x^2 + a})^n dx$$

input `integrate((x^2+a)^2*(x-(x^2+a)^(1/2))^n,x, algorithm="giac")`

output `integrate((x^2 + a)^2*(x - sqrt(x^2 + a))^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (a + x^2)^2 (x - \sqrt{a + x^2})^n dx = \int (x - \sqrt{x^2 + a})^n (x^2 + a)^2 dx$$

input `int((x - (a + x^2)^(1/2))^n*(a + x^2)^2,x)`

output `int((x - (a + x^2)^(1/2))^n*(a + x^2)^2, x)`

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.30

$$\int (a + x^2)^2 (x - \sqrt{a + x^2})^n dx$$

$$= \frac{a^n (-1)^n (-\sqrt{x^2 + a} a^2 n^5 + 30\sqrt{x^2 + a} a^2 n^3 - 149\sqrt{x^2 + a} a^2 n - 2\sqrt{x^2 + a} a n^5 x^2 + 40\sqrt{x^2 + a} a n^3 x^2 - 38\sqrt{x^2 + a} a n x^2 - \sqrt{x^2 + a} n^5 x^4 + 10\sqrt{x^2 + a} n^3 x^4 - 9\sqrt{x^2 + a} n x^4 - 5a^2 n^4 x + 110a^2 n^2 x - 225a^2 x - 10a n^4 x^3 + 160a n^2 x^3 - 150a x^3 - 5n^4 x^5 + 50n^2 x^5 - 45x^5)}{(\sqrt{x^2 + a} + x)^n (n^6 - 35n^4 + 259n^2 - 225)}$$

input

```
int((x^2+a)^2*(x-(x^2+a)^(1/2))^n,x)
```

output

```
(a**n*( - 1)**n*(- sqrt(a + x**2)*a**2*n**5 + 30*sqrt(a + x**2)*a**2*n**3
- 149*sqrt(a + x**2)*a**2*n - 2*sqrt(a + x**2)*a*n**5*x**2 + 40*sqrt(a +
x**2)*a*n**3*x**2 - 38*sqrt(a + x**2)*a*n*x**2 - sqrt(a + x**2)*n**5*x**4
+ 10*sqrt(a + x**2)*n**3*x**4 - 9*sqrt(a + x**2)*n*x**4 - 5*a**2*n**4*x +
110*a**2*n**2*x - 225*a**2*x - 10*a*n**4*x**3 + 160*a*n**2*x**3 - 150*a*x*
*3 - 5*n**4*x**5 + 50*n**2*x**5 - 45*x**5))/((sqrt(a + x**2) + x)**n*(n**6
- 35*n**4 + 259*n**2 - 225))
```



### 3.22 $\int (a + x^2) \left(x - \sqrt{a + x^2}\right)^n dx$

Optimal result	184
Mathematica [A] (verified)	184
Rubi [A] (verified)	185
Maple [F]	186
Fricas [A] (verification not implemented)	186
Sympy [F]	187
Maxima [F]	187
Giac [F]	187
Mupad [F(-1)]	188
Reduce [B] (verification not implemented)	188

#### Optimal result

Integrand size = 21, antiderivative size = 116

$$\int (a + x^2) \left(x - \sqrt{a + x^2}\right)^n dx = -\frac{a^3(x - \sqrt{a + x^2})^{-3+n}}{8(3 - n)} - \frac{3a^2(x - \sqrt{a + x^2})^{-1+n}}{8(1 - n)} + \frac{3a(x - \sqrt{a + x^2})^{1+n}}{8(1 + n)} + \frac{(x - \sqrt{a + x^2})^{3+n}}{8(3 + n)}$$

output

$$-1/8*a^3*(x-(x^2+a)^(1/2))^{(-3+n)/(3-n)}-3*a^2*(x-(x^2+a)^(1/2))^{(-1+n)/(8-8*n)}+3*a*(x-(x^2+a)^(1/2))^{(1+n)/(8+8*n)}+(x-(x^2+a)^(1/2))^{(3+n)/(24+8*n)}$$

#### Mathematica [A] (verified)

Time = 0.38 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.86

$$\int (a + x^2) \left(x - \sqrt{a + x^2}\right)^n dx = \frac{1}{8} \left(x - \sqrt{a + x^2}\right)^{-3+n} \left( \frac{a^3}{-3 + n} + \frac{3a^2(x - \sqrt{a + x^2})^2}{-1 + n} + \frac{3a(x - \sqrt{a + x^2})^4}{1 + n} + \frac{(x - \sqrt{a + x^2})^6}{3 + n} \right)$$

input

```
Integrate[(a + x^2)*(x - Sqrt[a + x^2])^n,x]
```

output

$$\frac{((x - \sqrt{a + x^2})^{-3 + n})(a^3/(-3 + n) + (3*a^2*(x - \sqrt{a + x^2}))^2)/(-1 + n) + (3*a*(x - \sqrt{a + x^2})^4)/(1 + n) + (x - \sqrt{a + x^2})^6/(3 + n))}{8}$$

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.96, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {2547, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + x^2) (x - \sqrt{a + x^2})^n dx$$

$$\downarrow 2547$$

$$\frac{1}{8} \int (x - \sqrt{x^2 + a})^{n-4} \left( (x - \sqrt{x^2 + a})^2 + a \right)^3 d(x - \sqrt{x^2 + a})$$

$$\downarrow 244$$

$$\frac{1}{8} \int \left( a^3 (x - \sqrt{x^2 + a})^{n-4} + 3a^2 (x - \sqrt{x^2 + a})^{n-2} + 3a (x - \sqrt{x^2 + a})^n + (x - \sqrt{x^2 + a})^{n+2} \right) d(x - \sqrt{x^2 + a})$$

$$\downarrow 2009$$

$$\frac{1}{8} \left( -\frac{a^3 (x - \sqrt{a + x^2})^{n-3}}{3 - n} - \frac{3a^2 (x - \sqrt{a + x^2})^{n-1}}{1 - n} + \frac{3a (x - \sqrt{a + x^2})^{n+1}}{n + 1} + \frac{(x - \sqrt{a + x^2})^{n+3}}{n + 3} \right)$$

input

$$\text{Int}[(a + x^2)*(x - \text{Sqrt}[a + x^2])^n, x]$$

output

$$\frac{(-(a^3*(x - \text{Sqrt}[a + x^2])^{-3 + n})/(3 - n)) - (3*a^2*(x - \text{Sqrt}[a + x^2])^{-1 + n})/(1 - n) + (3*a*(x - \text{Sqrt}[a + x^2])^{1 + n})/(1 + n) + (x - \text{Sqrt}[a + x^2])^{3 + n}/(3 + n))/8}$$

**Defintions of rubi rules used**

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2547 `Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

**Maple [F]**

$$\int (x^2 + a) (x - \sqrt{x^2 + a})^n dx$$

input `int((x^2+a)*(x-(x^2+a)^(1/2))^n,x)`

output `int((x^2+a)*(x-(x^2+a)^(1/2))^n,x)`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.68

$$\int (a + x^2) (x - \sqrt{a + x^2})^n dx = \frac{(3(n^2 - 1)x^3 + 3(an^2 - 3a)x + (an^3 + (n^3 - n)x^2 - 7an)\sqrt{x^2 + a})(x - \sqrt{x^2 + a})^n}{n^4 - 10n^2 + 9}$$

input `integrate((x^2+a)*(x-(x^2+a)^(1/2))^n,x, algorithm="fricas")`

output  $-(3*(n^2 - 1)*x^3 + 3*(a*n^2 - 3*a)*x + (a*n^3 + (n^3 - n)*x^2 - 7*a*n)*\text{sqrt}(x^2 + a))*(x - \text{sqrt}(x^2 + a))^n/(n^4 - 10*n^2 + 9)$

### Sympy [F]

$$\int (a + x^2) (x - \sqrt{a + x^2})^n dx = \int (a + x^2) (x - \sqrt{a + x^2})^n dx$$

input `integrate((x**2+a)*(x-(x**2+a)**(1/2))**n,x)`

output `Integral((a + x**2)*(x - sqrt(a + x**2))**n, x)`

### Maxima [F]

$$\int (a + x^2) (x - \sqrt{a + x^2})^n dx = \int (x^2 + a) (x - \sqrt{x^2 + a})^n dx$$

input `integrate((x^2+a)*(x-(x^2+a)^(1/2))^n,x, algorithm="maxima")`

output `integrate((x^2 + a)*(x - sqrt(x^2 + a))^n, x)`

### Giac [F]

$$\int (a + x^2) (x - \sqrt{a + x^2})^n dx = \int (x^2 + a) (x - \sqrt{x^2 + a})^n dx$$

input `integrate((x^2+a)*(x-(x^2+a)^(1/2))^n,x, algorithm="giac")`

output `integrate((x^2 + a)*(x - sqrt(x^2 + a))^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (a + x^2) (x - \sqrt{a + x^2})^n dx = \int (x - \sqrt{x^2 + a})^n (x^2 + a) dx$$

input `int((x - (a + x^2)^(1/2))^n*(a + x^2), x)`output `int((x - (a + x^2)^(1/2))^n*(a + x^2), x)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.89

$$\int (a + x^2) (x - \sqrt{a + x^2})^n dx$$

$$= \frac{a^n (-1)^n (-\sqrt{x^2 + a} a n^3 + 7\sqrt{x^2 + a} a n - \sqrt{x^2 + a} n^3 x^2 + \sqrt{x^2 + a} n x^2 - 3a n^2 x + 9a x - 3n^2 x^3 + 3x^3)}{(\sqrt{x^2 + a} + x)^n (n^4 - 10n^2 + 9)}$$

input `int((x^2+a)*(x-(x^2+a)^(1/2))^n,x)`output `(a**n*(-1)**n*(-sqrt(a+x**2)*a*n**3+7*sqrt(a+x**2)*a*n-sqrt(a+x**2)*n**3*x**2+sqrt(a+x**2)*n*x**2-3*a*n**2*x+9*a*x-3*n**2*x**3+3*x**3))/((sqrt(a+x**2)+x)**n*(n**4-10*n**2+9))`

### 3.23 $\int \left(x - \sqrt{a + x^2}\right)^n dx$

Optimal result	189
Mathematica [A] (verified)	189
Rubi [A] (verified)	190
Maple [F]	191
Fricas [A] (verification not implemented)	191
Sympy [F]	192
Maxima [F]	192
Giac [F]	192
Mupad [F(-1)]	193
Reduce [B] (verification not implemented)	193

#### Optimal result

Integrand size = 15, antiderivative size = 56

$$\int \left(x - \sqrt{a + x^2}\right)^n dx = -\frac{a(x - \sqrt{a + x^2})^{-1+n}}{2(1 - n)} + \frac{(x - \sqrt{a + x^2})^{1+n}}{2(1 + n)}$$

output

```
-1/2*a*(x-(x^2+a)^(1/2))^(1+n)/(1+n)+(x-(x^2+a)^(1/2))^(1+n)/(2+2*n)
```

#### Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.89

$$\int \left(x - \sqrt{a + x^2}\right)^n dx = \frac{1}{2} \left(x - \sqrt{a + x^2}\right)^{-1+n} \left(\frac{a}{-1 + n} + \frac{(x - \sqrt{a + x^2})^2}{1 + n}\right)$$

input

```
Integrate[(x - Sqrt[a + x^2])^n,x]
```

output

```
((x - Sqrt[a + x^2])^(-1 + n)*(a/(-1 + n) + (x - Sqrt[a + x^2])^2/(1 + n)))/2
```

**Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.98, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {2542, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (x - \sqrt{a + x^2})^n dx$$

$$\downarrow 2542$$

$$\frac{1}{2} \int (x - \sqrt{x^2 + a})^{n-2} \left( (x - \sqrt{x^2 + a})^2 + a \right) d(x - \sqrt{x^2 + a})$$

$$\downarrow 244$$

$$\frac{1}{2} \int \left( a(x - \sqrt{x^2 + a})^{n-2} + (x - \sqrt{x^2 + a})^n \right) d(x - \sqrt{x^2 + a})$$

$$\downarrow 2009$$

$$\frac{1}{2} \left( \frac{(x - \sqrt{a + x^2})^{n+1}}{n+1} - \frac{a(x - \sqrt{a + x^2})^{n-1}}{1-n} \right)$$

input `Int[(x - Sqrt[a + x^2])^n,x]`

output `((-(a*(x - Sqrt[a + x^2])^(-1 + n))/(1 - n)) + (x - Sqrt[a + x^2])^(1 + n))/(1 + n)/2`

**Defintions of rubi rules used**

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2542 `Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.))^(p_.), x_Symbol] := Simp[1/(2*e) Subst[Int[(g + h*x^n)^p*((d^2 + a*f^2 - 2*d*x + x^2)/(d - x)^2), x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]`

**Maple [F]**

$$\int (x - \sqrt{x^2 + a})^n dx$$

input `int((x-(x^2+a)^(1/2))^n,x)`

output `int((x-(x^2+a)^(1/2))^n,x)`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.59

$$\int (x - \sqrt{a + x^2})^n dx = -\frac{(\sqrt{x^2 + a}n + x)(x - \sqrt{x^2 + a})^n}{n^2 - 1}$$

input `integrate((x-(x^2+a)^(1/2))^n,x, algorithm="fricas")`

output `-(sqrt(x^2 + a)*n + x)*(x - sqrt(x^2 + a))^n/(n^2 - 1)`



**Sympy [F]**

$$\int (x - \sqrt{a + x^2})^n dx = \int (x - \sqrt{a + x^2})^n dx$$

input `integrate((x-(x**2+a)**(1/2))**n,x)`

output `Integral((x - sqrt(a + x**2))**n, x)`

**Maxima [F]**

$$\int (x - \sqrt{a + x^2})^n dx = \int (x - \sqrt{x^2 + a})^n dx$$

input `integrate((x-(x^2+a)^(1/2))^n,x, algorithm="maxima")`

output `integrate((x - sqrt(x^2 + a))^n, x)`

**Giac [F]**

$$\int (x - \sqrt{a + x^2})^n dx = \int (x - \sqrt{x^2 + a})^n dx$$

input `integrate((x-(x^2+a)^(1/2))^n,x, algorithm="giac")`

output `integrate((x - sqrt(x^2 + a))^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (x - \sqrt{a + x^2})^n dx = \int (x - \sqrt{x^2 + a})^n dx$$

input `int((x - (a + x^2)^(1/2))^n,x)`output `int((x - (a + x^2)^(1/2))^n, x)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.66

$$\int (x - \sqrt{a + x^2})^n dx = -\frac{a^n(-1)^n(\sqrt{x^2 + a}n + x)}{(\sqrt{x^2 + a} + x)^n(n^2 - 1)}$$

input `int((x-(x^2+a)^(1/2))^n,x)`output `( - a**n*( - 1)**n*(sqrt(a + x**2)*n + x))/((sqrt(a + x**2) + x)**n*(n**2 - 1))`

**3.24** 
$$\int \frac{(x - \sqrt{a+x^2})^n}{a+x^2} dx$$

Optimal result	194
Mathematica [A] (verified)	194
Rubi [A] (verified)	195
Maple [F]	196
Fricas [F]	196
Sympy [F]	197
Maxima [F]	197
Giac [F]	197
Mupad [F(-1)]	198
Reduce [F]	198

**Optimal result**

Integrand size = 23, antiderivative size = 63

$$\int \frac{(x - \sqrt{a+x^2})^n}{a+x^2} dx = \frac{2(x - \sqrt{a+x^2})^{1+n} \text{Hypergeometric2F1}\left(1, \frac{1+n}{2}, \frac{3+n}{2}, -\frac{(x - \sqrt{a+x^2})^2}{a}\right)}{a(1+n)}$$

output

```
2*(x-(x^2+a)^(1/2))^(1+n)*hypergeom([1, 1/2+1/2*n],[3/2+1/2*n],-(x-(x^2+a)^(1/2))^2/a)/a/(1+n)
```

**Mathematica [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.03

$$\int \frac{(x - \sqrt{a+x^2})^n}{a+x^2} dx = \frac{2(x - \sqrt{a+x^2})^{1+n} \text{Hypergeometric2F1}\left(1, \frac{1+n}{2}, 1 + \frac{1+n}{2}, -\frac{(x - \sqrt{a+x^2})^2}{a}\right)}{a(1+n)}$$

input `Integrate[(x - Sqrt[a + x^2])^n/(a + x^2), x]`

output `(2*(x - Sqrt[a + x^2])^(1 + n)*Hypergeometric2F1[1, (1 + n)/2, 1 + (1 + n)/2, -((x - Sqrt[a + x^2])^2/a)])/(a*(1 + n))`

### Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$ , Rules used = {2547, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(x - \sqrt{a + x^2})^n}{a + x^2} dx$$

$$\downarrow 2547$$

$$2 \int \frac{(x - \sqrt{x^2 + a})^n}{(x - \sqrt{x^2 + a})^2 + a} d(x - \sqrt{x^2 + a})$$

$$\downarrow 278$$

$$\frac{2(x - \sqrt{a + x^2})^{n+1} \text{Hypergeometric2F1}\left(1, \frac{n+1}{2}, \frac{n+3}{2}, -\frac{(x - \sqrt{x^2 + a})^2}{a}\right)}{a(n+1)}$$

input `Int[(x - Sqrt[a + x^2])^n/(a + x^2), x]`

output `(2*(x - Sqrt[a + x^2])^(1 + n)*Hypergeometric2F1[1, (1 + n)/2, (3 + n)/2, -((x - Sqrt[a + x^2])^2/a)])/(a*(1 + n))`

## Definitions of rubi rules used

rule 278

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])
```

rule 2547

```
Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])
```

## Maple [F]

$$\int \frac{(x - \sqrt{x^2 + a})^n}{x^2 + a} dx$$

input

```
int((x-(x^2+a)^(1/2))^n/(x^2+a),x)
```

output

```
int((x-(x^2+a)^(1/2))^n/(x^2+a),x)
```

## Fricas [F]

$$\int \frac{(x - \sqrt{a + x^2})^n}{a + x^2} dx = \int \frac{(x - \sqrt{x^2 + a})^n}{x^2 + a} dx$$

input

```
integrate((x-(x^2+a)^(1/2))^n/(x^2+a),x, algorithm="fricas")
```

output

```
integral((x - sqrt(x^2 + a))^n/(x^2 + a), x)
```

**Sympy [F]**

$$\int \frac{(x - \sqrt{a + x^2})^n}{a + x^2} dx = \int \frac{(x - \sqrt{a + x^2})^n}{a + x^2} dx$$

input `integrate((x-(x**2+a)**(1/2))**n/(x**2+a), x)`

output `Integral((x - sqrt(a + x**2))**n/(a + x**2), x)`

**Maxima [F]**

$$\int \frac{(x - \sqrt{a + x^2})^n}{a + x^2} dx = \int \frac{(x - \sqrt{x^2 + a})^n}{x^2 + a} dx$$

input `integrate((x-(x^2+a)^(1/2))^n/(x^2+a), x, algorithm="maxima")`

output `integrate((x - sqrt(x^2 + a))^n/(x^2 + a), x)`

**Giac [F]**

$$\int \frac{(x - \sqrt{a + x^2})^n}{a + x^2} dx = \int \frac{(x - \sqrt{x^2 + a})^n}{x^2 + a} dx$$

input `integrate((x-(x^2+a)^(1/2))^n/(x^2+a), x, algorithm="giac")`

output `integrate((x - sqrt(x^2 + a))^n/(x^2 + a), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(x - \sqrt{a + x^2})^n}{a + x^2} dx = \int \frac{(x - \sqrt{x^2 + a})^n}{x^2 + a} dx$$

input `int((x - (a + x^2)^(1/2))^n/(a + x^2), x)`output `int((x - (a + x^2)^(1/2))^n/(a + x^2), x)`**Reduce [F]**

$$\int \frac{(x - \sqrt{a + x^2})^n}{a + x^2} dx = \int \frac{(-\sqrt{x^2 + a} + x)^n}{x^2 + a} dx$$

input `int((x-(x^2+a)^(1/2))^n/(x^2+a), x)`output `int((-sqrt(a + x**2) + x)**n/(a + x**2), x)`

**3.25**  $\int \frac{(x - \sqrt{a+x^2})^n}{(a+x^2)^2} dx$

Optimal result	199
Mathematica [A] (verified)	199
Rubi [A] (verified)	200
Maple [F]	201
Fricas [F]	201
Sympy [F]	202
Maxima [F]	202
Giac [F]	202
Mupad [F(-1)]	203
Reduce [F]	203

**Optimal result**

Integrand size = 23, antiderivative size = 63

$$\int \frac{(x - \sqrt{a+x^2})^n}{(a+x^2)^2} dx = \frac{8(x - \sqrt{a+x^2})^{3+n} \operatorname{Hypergeometric2F1}\left(3, \frac{3+n}{2}, \frac{5+n}{2}, -\frac{(x - \sqrt{a+x^2})^2}{a}\right)}{a^3(3+n)}$$

output `8*(x-(x^2+a)^(1/2))^(3+n)*hypergeom([3, 3/2+1/2*n],[5/2+1/2*n],-(x-(x^2+a)^(1/2))^2/a)/a^3/(3+n)`

**Mathematica [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.03

$$\int \frac{(x - \sqrt{a+x^2})^n}{(a+x^2)^2} dx = \frac{8(x - \sqrt{a+x^2})^{3+n} \operatorname{Hypergeometric2F1}\left(3, \frac{3+n}{2}, 1 + \frac{3+n}{2}, -\frac{(x - \sqrt{a+x^2})^2}{a}\right)}{a^3(3+n)}$$



input `Integrate[(x - Sqrt[a + x^2])^n/(a + x^2)^2,x]`

output `(8*(x - Sqrt[a + x^2])^(3 + n)*Hypergeometric2F1[3, (3 + n)/2, 1 + (3 + n)/2, -((x - Sqrt[a + x^2])^2/a)])/(a^3*(3 + n))`

### Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$ , Rules used = {2547, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^2} dx$$

↓ 2547

$$8 \int \frac{(x - \sqrt{x^2 + a})^{n+2}}{\left( (x - \sqrt{x^2 + a})^2 + a \right)^3} d(x - \sqrt{x^2 + a})$$

↓ 278

$$\frac{8(x - \sqrt{a + x^2})^{n+3} \text{Hypergeometric2F1}\left(3, \frac{n+3}{2}, \frac{n+5}{2}, -\frac{(x - \sqrt{x^2 + a})^2}{a}\right)}{a^3(n + 3)}$$

input `Int[(x - Sqrt[a + x^2])^n/(a + x^2)^2,x]`

output `(8*(x - Sqrt[a + x^2])^(3 + n)*Hypergeometric2F1[3, (3 + n)/2, (5 + n)/2, -((x - Sqrt[a + x^2])^2/a)])/(a^3*(3 + n))`

## Definitions of rubi rules used

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 2547 `Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

## Maple [F]

$$\int \frac{(x - \sqrt{x^2 + a})^n}{(x^2 + a)^2} dx$$

input `int((x-(x^2+a)^(1/2))^n/(x^2+a)^2,x)`

output `int((x-(x^2+a)^(1/2))^n/(x^2+a)^2,x)`

## Fricas [F]

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^2} dx = \int \frac{(x - \sqrt{x^2 + a})^n}{(x^2 + a)^2} dx$$

input `integrate((x-(x^2+a)^(1/2))^n/(x^2+a)^2,x, algorithm="fricas")`

output `integral((x - sqrt(x^2 + a))^n/(x^4 + 2*a*x^2 + a^2), x)`

**Sympy [F]**

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^2} dx = \int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^2} dx$$

input `integrate((x-(x**2+a)**(1/2))**n/(x**2+a)**2,x)`

output `Integral((x - sqrt(a + x**2))**n/(a + x**2)**2, x)`

**Maxima [F]**

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^2} dx = \int \frac{(x - \sqrt{x^2 + a})^n}{(x^2 + a)^2} dx$$

input `integrate((x-(x^2+a)^(1/2))^n/(x^2+a)^2,x, algorithm="maxima")`

output `integrate((x - sqrt(x^2 + a))^n/(x^2 + a)^2, x)`

**Giac [F]**

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^2} dx = \int \frac{(x - \sqrt{x^2 + a})^n}{(x^2 + a)^2} dx$$

input `integrate((x-(x^2+a)^(1/2))^n/(x^2+a)^2,x, algorithm="giac")`

output `integrate((x - sqrt(x^2 + a))^n/(x^2 + a)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^2} dx = \int \frac{(x - \sqrt{x^2 + a})^n}{(x^2 + a)^2} dx$$

input `int((x - (a + x^2)^(1/2))^n/(a + x^2)^2,x)`output `int((x - (a + x^2)^(1/2))^n/(a + x^2)^2, x)`**Reduce [F]**

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^2} dx = \int \frac{(-\sqrt{x^2 + a} + x)^n}{x^4 + 2ax^2 + a^2} dx$$

input `int((x-(x^2+a)^(1/2))^n/(x^2+a)^2,x)`output `int((-sqrt(a + x**2) + x)**n/(a**2 + 2*a*x**2 + x**4),x)`

### 3.26 $\int (a + x^2)^{5/2} \left(x + \sqrt{a + x^2}\right)^n dx$

Optimal result	204
Mathematica [A] (verified)	205
Rubi [A] (verified)	205
Maple [F]	207
Fricas [A] (verification not implemented)	207
Sympy [F(-2)]	207
Maxima [F]	208
Giac [F]	208
Mupad [F(-1)]	208
Reduce [B] (verification not implemented)	209

#### Optimal result

Integrand size = 23, antiderivative size = 187

$$\int (a + x^2)^{5/2} \left(x + \sqrt{a + x^2}\right)^n dx = -\frac{a^6(x + \sqrt{a + x^2})^{-6+n}}{64(6 - n)} - \frac{3a^5(x + \sqrt{a + x^2})^{-4+n}}{32(4 - n)} - \frac{15a^4(x + \sqrt{a + x^2})^{-2+n}}{64(2 - n)} + \frac{5a^3(x + \sqrt{a + x^2})^n}{16n} + \frac{15a^2(x + \sqrt{a + x^2})^{2+n}}{64(2 + n)} + \frac{3a(x + \sqrt{a + x^2})^{4+n}}{32(4 + n)} + \frac{(x + \sqrt{a + x^2})^{6+n}}{64(6 + n)}$$

output

```
-1/64*a^6*(x+(x^2+a)^(1/2))^(6-n)/(6-n)-3*a^5*(x+(x^2+a)^(1/2))^(4-n)/(128-32*n)-15*a^4*(x+(x^2+a)^(1/2))^(2-n)/(128-64*n)+5/16*a^3*(x+(x^2+a)^(1/2))^n/n+15*a^2*(x+(x^2+a)^(1/2))^(2+n)/(128+64*n)+3*a*(x+(x^2+a)^(1/2))^(4+n)/(128+32*n)+(x+(x^2+a)^(1/2))^(6+n)/(384+64*n)
```

**Mathematica [A] (verified)**

Time = 0.59 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.84

$$\int (a+x^2)^{5/2} (x+\sqrt{a+x^2})^n dx = \frac{1}{64} (x+\sqrt{a+x^2})^n \left( \frac{20a^3}{n} \right. \\ \left. + \frac{a^6}{(-6+n)(x+\sqrt{a+x^2})^6} + \frac{6a^5}{(-4+n)(x+\sqrt{a+x^2})^4} \right. \\ \left. + \frac{15a^4}{(-2+n)(x+\sqrt{a+x^2})^2} + \frac{15a^2(x+\sqrt{a+x^2})^2}{2+n} \right. \\ \left. + \frac{6a(x+\sqrt{a+x^2})^4}{4+n} + \frac{(x+\sqrt{a+x^2})^6}{6+n} \right)$$

input

```
Integrate[(a + x^2)^(5/2)*(x + Sqrt[a + x^2])^n,x]
```

output

```
((x + Sqrt[a + x^2])^n*((20*a^3)/n + a^6/((-6 + n)*(x + Sqrt[a + x^2])^6)
+ (6*a^5)/((-4 + n)*(x + Sqrt[a + x^2])^4) + (15*a^4)/((-2 + n)*(x + Sqrt[
a + x^2])^2) + (15*a^2*(x + Sqrt[a + x^2])^2)/(2 + n) + (6*a*(x + Sqrt[a +
x^2])^4)/(4 + n) + (x + Sqrt[a + x^2])^6/(6 + n)))/64
```

**Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.94, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {2547, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a+x^2)^{5/2} (\sqrt{a+x^2}+x)^n dx \\ \downarrow 2547 \\ \frac{1}{64} \int (x+\sqrt{x^2+a})^{n-7} \left( (x+\sqrt{x^2+a})^2+a \right)^6 d(x+\sqrt{x^2+a})$$

↓ 244

$$\frac{1}{64} \int \left( a^6 (x + \sqrt{x^2 + a})^{n-7} + 6a^5 (x + \sqrt{x^2 + a})^{n-5} + 15a^4 (x + \sqrt{x^2 + a})^{n-3} + 20a^3 (x + \sqrt{x^2 + a})^{n-1} + \dots \right)$$

↓ 2009

$$\frac{1}{64} \left( -\frac{a^6 (\sqrt{a+x^2}+x)^{n-6}}{6-n} - \frac{6a^5 (\sqrt{a+x^2}+x)^{n-4}}{4-n} - \frac{15a^4 (\sqrt{a+x^2}+x)^{n-2}}{2-n} + \frac{20a^3 (\sqrt{a+x^2}+x)^n}{n} + \dots \right)$$

input `Int[(a + x^2)^(5/2)*(x + Sqrt[a + x^2])^n,x]`

output `(-((a^6*(x + Sqrt[a + x^2])^(-6 + n))/(6 - n)) - (6*a^5*(x + Sqrt[a + x^2])^(-4 + n))/(4 - n) - (15*a^4*(x + Sqrt[a + x^2])^(-2 + n))/(2 - n) + (20*a^3*(x + Sqrt[a + x^2])^n)/n + (15*a^2*(x + Sqrt[a + x^2])^(2 + n))/(2 + n) + (6*a*(x + Sqrt[a + x^2])^(4 + n))/(4 + n) + (x + Sqrt[a + x^2])^(6 + n)/(6 + n))/64`

### Defintions of rubi rules used

rule 244 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2547 `Int[((g_) + (i_)*(x_)^2)^(m_)*((d_) + (e_)*(x_) + (f_)*Sqrt[(a_) + (c_)*(x_)^2])^(n_), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

**Maple [F]**

$$\int (x^2 + a)^{\frac{5}{2}} (x + \sqrt{x^2 + a})^n dx$$

input `int((x^2+a)^(5/2)*(x+(x^2+a)^(1/2))^n,x)`

output `int((x^2+a)^(5/2)*(x+(x^2+a)^(1/2))^n,x)`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.07

$$\int (a + x^2)^{5/2} (x + \sqrt{a + x^2})^n dx = \frac{(a^3 n^6 - 50 a^3 n^4 + (n^6 - 20 n^4 + 64 n^2) x^6 + 544 a^3 n^2 + 3 (a n^6 - 30 a n^4 + 104 a n^2) x^4 - 720 a^3 + 3 (a^2 n^6 - 40 a^2 n^4 + 264 a^2 n^2) x^2 - 6 ((n^5 - 20 n^3 + 64 n) x^5 + 2 (a n^5 - 30 a n^3 + 104 a n) x^3 + (a^2 n^5 - 40 a^2 n^3 + 264 a^2 n) x) \sqrt{x^2 + a}) (x + \sqrt{x^2 + a})^n}{(n^7 - 56 n^5 + 784 n^3 - 2304 n)}$$

input `integrate((x^2+a)^(5/2)*(x+(x^2+a)^(1/2))^n,x, algorithm="fricas")`

output `(a^3*n^6 - 50*a^3*n^4 + (n^6 - 20*n^4 + 64*n^2)*x^6 + 544*a^3*n^2 + 3*(a*n^6 - 30*a*n^4 + 104*a*n^2)*x^4 - 720*a^3 + 3*(a^2*n^6 - 40*a^2*n^4 + 264*a^2*n^2)*x^2 - 6*((n^5 - 20*n^3 + 64*n)*x^5 + 2*(a*n^5 - 30*a*n^3 + 104*a*n)*x^3 + (a^2*n^5 - 40*a^2*n^3 + 264*a^2*n)*x)*sqrt(x^2 + a)*(x + sqrt(x^2 + a))^n/(n^7 - 56*n^5 + 784*n^3 - 2304*n)`

**Sympy [F(-2)]**

Exception generated.

$$\int (a + x^2)^{5/2} (x + \sqrt{a + x^2})^n dx = \text{Exception raised: HeuristicGCDFailed}$$

input `integrate((x**2+a)**(5/2)*(x+(x**2+a)**(1/2))**n,x)`

output `Exception raised: HeuristicGCDFailed >> no luck`



**Maxima [F]**

$$\int (a + x^2)^{5/2} (x + \sqrt{a + x^2})^n dx = \int (x^2 + a)^{5/2} (x + \sqrt{x^2 + a})^n dx$$

input `integrate((x^2+a)^(5/2)*(x+(x^2+a)^(1/2))^n,x, algorithm="maxima")`

output `integrate((x^2 + a)^(5/2)*(x + sqrt(x^2 + a))^n, x)`

**Giac [F]**

$$\int (a + x^2)^{5/2} (x + \sqrt{a + x^2})^n dx = \int (x^2 + a)^{5/2} (x + \sqrt{x^2 + a})^n dx$$

input `integrate((x^2+a)^(5/2)*(x+(x^2+a)^(1/2))^n,x, algorithm="giac")`

output `integrate((x^2 + a)^(5/2)*(x + sqrt(x^2 + a))^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (a + x^2)^{5/2} (x + \sqrt{a + x^2})^n dx = \int (x^2 + a)^{5/2} (x + \sqrt{x^2 + a})^n dx$$

input `int((a + x^2)^(5/2)*(x + (a + x^2)^(1/2))^n,x)`

output `int((a + x^2)^(5/2)*(x + (a + x^2)^(1/2))^n, x)`

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.44

$$\int (a + x^2)^{5/2} \left( x + \sqrt{a + x^2} \right)^n dx = \frac{(\sqrt{x^2 + a} + x)^n \left( -6\sqrt{x^2 + a} a^2 n^5 x + 240\sqrt{x^2 + a} a^2 n^3 x - 1584\sqrt{x^2 + a} a^2 n x - 12\sqrt{x^2 + a} a^2 \right)}{n^6 - 56n^4 + 784n^2 - 2304}$$

input `int((x^2+a)^(5/2)*(x+(x^2+a)^(1/2))^n,x)`output `((sqrt(a + x**2) + x)**n*( - 6*sqrt(a + x**2)*a**2*n**5*x + 240*sqrt(a + x**2)*a**2*n**3*x - 1584*sqrt(a + x**2)*a**2*n*x - 12*sqrt(a + x**2)*a*n**5*x**3 + 360*sqrt(a + x**2)*a*n**3*x**3 - 1248*sqrt(a + x**2)*a*n*x**3 - 6*sqrt(a + x**2)*n**5*x**5 + 120*sqrt(a + x**2)*n**3*x**5 - 384*sqrt(a + x**2)*n*x**5 + a**3*n**6 - 50*a**3*n**4 + 544*a**3*n**2 - 720*a**3 + 3*a**2*n**6*x**2 - 120*a**2*n**4*x**2 + 792*a**2*n**2*x**2 + 3*a*n**6*x**4 - 90*a*n**4*x**4 + 312*a*n**2*x**4 + n**6*x**6 - 20*n**4*x**6 + 64*n**2*x**6))/(n*(n**6 - 56*n**4 + 784*n**2 - 2304))`

### 3.27 $\int (a + x^2)^{3/2} \left(x + \sqrt{a + x^2}\right)^n dx$

Optimal result	210
Mathematica [A] (verified)	211
Rubi [A] (verified)	211
Maple [F]	213
Fricas [A] (verification not implemented)	213
Sympy [F]	213
Maxima [F]	214
Giac [F]	214
Mupad [F(-1)]	214
Reduce [B] (verification not implemented)	215

#### Optimal result

Integrand size = 23, antiderivative size = 131

$$\int (a + x^2)^{3/2} \left(x + \sqrt{a + x^2}\right)^n dx = -\frac{a^4(x + \sqrt{a + x^2})^{-4+n}}{16(4 - n)} - \frac{a^3(x + \sqrt{a + x^2})^{-2+n}}{4(2 - n)} + \frac{3a^2(x + \sqrt{a + x^2})^n}{8n} + \frac{a(x + \sqrt{a + x^2})^{2+n}}{4(2 + n)} + \frac{(x + \sqrt{a + x^2})^{4+n}}{16(4 + n)}$$

output

```
-1/16*a^4*(x+(x^2+a)^(1/2))^(4-n)/(4-n)-a^3*(x+(x^2+a)^(1/2))^(2-n)/(8-4*n)+3/8*a^2*(x+(x^2+a)^(1/2))^n/n+a*(x+(x^2+a)^(1/2))^(2+n)/(8+4*n)+(x+(x^2+a)^(1/2))^(4+n)/(64+16*n)
```

**Mathematica [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.85

$$\int (a + x^2)^{3/2} (x + \sqrt{a + x^2})^n dx = \frac{1}{16} (x + \sqrt{a + x^2})^n \left( \frac{6a^2}{n} + \frac{a^4}{(-4 + n)(x + \sqrt{a + x^2})^4} + \frac{4a^3}{(-2 + n)(x + \sqrt{a + x^2})^2} + \frac{4a(x + \sqrt{a + x^2})^2}{2 + n} + \frac{(x + \sqrt{a + x^2})^4}{4 + n} \right)$$

input `Integrate[(a + x^2)^(3/2)*(x + Sqrt[a + x^2])^n,x]`

output `((x + Sqrt[a + x^2])^n*((6*a^2)/n + a^4/((-4 + n)*(x + Sqrt[a + x^2])^4) + (4*a^3)/((-2 + n)*(x + Sqrt[a + x^2])^2) + (4*a*(x + Sqrt[a + x^2])^2)/(2 + n) + (x + Sqrt[a + x^2])^4/(4 + n))/16`

**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.95, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {2547, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + x^2)^{3/2} (\sqrt{a + x^2} + x)^n dx$$

$$\downarrow 2547$$

$$\frac{1}{16} \int (x + \sqrt{x^2 + a})^{n-5} \left( (x + \sqrt{x^2 + a})^2 + a \right)^4 d(x + \sqrt{x^2 + a})$$

$$\downarrow 244$$

$$\frac{1}{16} \int \left( a^4 (x + \sqrt{x^2 + a})^{n-5} + 4a^3 (x + \sqrt{x^2 + a})^{n-3} + 6a^2 (x + \sqrt{x^2 + a})^{n-1} + 4a (x + \sqrt{x^2 + a})^{n+1} + (x + \sqrt{x^2 + a})^{n+3} \right) dx$$

↓ 2009

$$\frac{1}{16} \left( -\frac{a^4 (\sqrt{a+x^2}+x)^{n-4}}{4-n} - \frac{4a^3 (\sqrt{a+x^2}+x)^{n-2}}{2-n} + \frac{6a^2 (\sqrt{a+x^2}+x)^n}{n} + \frac{4a (\sqrt{a+x^2}+x)^{n+2}}{n+2} + \frac{(\sqrt{a+x^2}+x)^{n+4}}{n+4} \right)$$

input `Int[(a + x^2)^(3/2)*(x + Sqrt[a + x^2])^n,x]`

output `(-((a^4*(x + Sqrt[a + x^2])^(-4 + n))/(4 - n)) - (4*a^3*(x + Sqrt[a + x^2])^(-2 + n))/(2 - n) + (6*a^2*(x + Sqrt[a + x^2])^n)/n + (4*a*(x + Sqrt[a + x^2])^(2 + n))/(2 + n) + (x + Sqrt[a + x^2])^(4 + n)/(4 + n))/16`

### Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2547 `Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

**Maple [F]**

$$\int (x^2 + a)^{\frac{3}{2}} (x + \sqrt{x^2 + a})^n dx$$

input `int((x^2+a)^(3/2)*(x+(x^2+a)^(1/2))^n,x)`

output `int((x^2+a)^(3/2)*(x+(x^2+a)^(1/2))^n,x)`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.84

$$\int (a + x^2)^{3/2} (x + \sqrt{a + x^2})^n dx = \frac{(a^2 n^4 + (n^4 - 4n^2)x^4 - 16a^2 n^2 + 2(an^4 - 10an^2)x^2 + 24a^2 - 4((n^3 - 4n)x^3 + (an^3 + \dots))}{n^5 - 20n^3 + 64n}$$

input `integrate((x^2+a)^(3/2)*(x+(x^2+a)^(1/2))^n,x, algorithm="fricas")`

output `(a^2*n^4 + (n^4 - 4*n^2)*x^4 - 16*a^2*n^2 + 2*(a*n^4 - 10*a*n^2)*x^2 + 24*a^2 - 4*((n^3 - 4*n)*x^3 + (a*n^3 - 10*a*n)*x)*sqrt(x^2 + a))*(x + sqrt(x^2 + a))^n/(n^5 - 20*n^3 + 64*n)`

**Sympy [F]**

$$\int (a + x^2)^{3/2} (x + \sqrt{a + x^2})^n dx = \int (a + x^2)^{\frac{3}{2}} (x + \sqrt{a + x^2})^n dx$$

input `integrate((x**2+a)**(3/2)*(x+(x**2+a)**(1/2))**n,x)`

output `Integral((a + x**2)**(3/2)*(x + sqrt(a + x**2))**n, x)`

**Maxima [F]**

$$\int (a + x^2)^{3/2} (x + \sqrt{a + x^2})^n dx = \int (x^2 + a)^{\frac{3}{2}} (x + \sqrt{x^2 + a})^n dx$$

input `integrate((x^2+a)^(3/2)*(x+(x^2+a)^(1/2))^n,x, algorithm="maxima")`

output `integrate((x^2 + a)^(3/2)*(x + sqrt(x^2 + a))^n, x)`

**Giac [F]**

$$\int (a + x^2)^{3/2} (x + \sqrt{a + x^2})^n dx = \int (x^2 + a)^{\frac{3}{2}} (x + \sqrt{x^2 + a})^n dx$$

input `integrate((x^2+a)^(3/2)*(x+(x^2+a)^(1/2))^n,x, algorithm="giac")`

output `integrate((x^2 + a)^(3/2)*(x + sqrt(x^2 + a))^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (a + x^2)^{3/2} (x + \sqrt{a + x^2})^n dx = \int (x^2 + a)^{3/2} (x + \sqrt{x^2 + a})^n dx$$

input `int((a + x^2)^(3/2)*(x + (a + x^2)^(1/2))^n,x)`

output `int((a + x^2)^(3/2)*(x + (a + x^2)^(1/2))^n, x)`

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.99

$$\int (a + x^2)^{3/2} (x + \sqrt{a + x^2})^n dx = \frac{(\sqrt{x^2 + a} + x)^n (-4\sqrt{x^2 + a} a n^3 x + 40\sqrt{x^2 + a} a n x - 4\sqrt{x^2 + a} n^3 x^3 + 16\sqrt{x^2 + a} n a x^2 - 16a^2 n^2 + 24a^2 + 2a n^4 x^2 - 20a n^2 x^2 + n^4 x^4 - 4n^2 x^4)}{n(n^4 - 20n^2 + 64)}$$

input `int((x^2+a)^(3/2)*(x+(x^2+a)^(1/2))^n,x)`output `((sqrt(a + x**2) + x)**n*(- 4*sqrt(a + x**2)*a*n**3*x + 40*sqrt(a + x**2)*a*n*x - 4*sqrt(a + x**2)*n**3*x**3 + 16*sqrt(a + x**2)*n*x**3 + a**2*n**4 - 16*a**2*n**2 + 24*a**2 + 2*a*n**4*x**2 - 20*a*n**2*x**2 + n**4*x**4 - 4*n**2*x**4))/(n*(n**4 - 20*n**2 + 64))`



### 3.28 $\int \sqrt{a+x^2} \left(x + \sqrt{a+x^2}\right)^n dx$

Optimal result	216
Mathematica [A] (verified)	216
Rubi [A] (verified)	217
Maple [F]	218
Fricas [A] (verification not implemented)	218
Sympy [F]	219
Maxima [F]	219
Giac [F]	219
Mupad [F(-1)]	220
Reduce [B] (verification not implemented)	220

#### Optimal result

Integrand size = 23, antiderivative size = 75

$$\int \sqrt{a+x^2} \left(x + \sqrt{a+x^2}\right)^n dx = -\frac{a^2(x + \sqrt{a+x^2})^{-2+n}}{4(2-n)} + \frac{a(x + \sqrt{a+x^2})^n}{2n} + \frac{(x + \sqrt{a+x^2})^{2+n}}{4(2+n)}$$

output

```
-1/4*a^2*(x+(x^2+a)^(1/2))^(2-n)/(2-n)+1/2*a*(x+(x^2+a)^(1/2))^n/n+(x+(x^2+a)^(1/2))^(2+n)/(8+4*n)
```

#### Mathematica [A] (verified)

Time = 0.31 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.87

$$\int \sqrt{a+x^2} \left(x + \sqrt{a+x^2}\right)^n dx = \frac{1}{4} \left(x + \sqrt{a+x^2}\right)^n \left( \frac{2a}{n} + \frac{a^2}{(-2+n)(x + \sqrt{a+x^2})^2} + \frac{(x + \sqrt{a+x^2})^2}{2+n} \right)$$

input

```
Integrate[Sqrt[a + x^2]*(x + Sqrt[a + x^2])^n,x]
```

output  $\frac{((x + \text{Sqrt}[a + x^2])^n * ((2*a)/n + a^2/((-2 + n)*(x + \text{Sqrt}[a + x^2])^2) + (x + \text{Sqrt}[a + x^2])^2/(2 + n)))/4}$

### Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.96, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {2547, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{a+x^2} (\sqrt{a+x^2}+x)^n dx$$

$$\downarrow 2547$$

$$\frac{1}{4} \int (x + \sqrt{x^2+a})^{n-3} \left( (x + \sqrt{x^2+a})^2 + a \right)^2 d(x + \sqrt{x^2+a})$$

$$\downarrow 244$$

$$\frac{1}{4} \int \left( a^2 (x + \sqrt{x^2+a})^{n-3} + 2a (x + \sqrt{x^2+a})^{n-1} + (x + \sqrt{x^2+a})^{n+1} \right) d(x + \sqrt{x^2+a})$$

$$\downarrow 2009$$

$$\frac{1}{4} \left( -\frac{a^2 (\sqrt{a+x^2}+x)^{n-2}}{2-n} + \frac{2a (\sqrt{a+x^2}+x)^n}{n} + \frac{(\sqrt{a+x^2}+x)^{n+2}}{n+2} \right)$$

input  $\text{Int}[\text{Sqrt}[a + x^2]*(x + \text{Sqrt}[a + x^2])^n, x]$

output  $\frac{-((a^2*(x + \text{Sqrt}[a + x^2])^{(-2 + n)})/(2 - n)) + (2*a*(x + \text{Sqrt}[a + x^2])^n)/n + (x + \text{Sqrt}[a + x^2])^{(2 + n)}/(2 + n))/4}$

**Defintions of rubi rules used**

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2547 `Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

**Maple [F]**

$$\int \sqrt{x^2 + a} \left( x + \sqrt{x^2 + a} \right)^n dx$$

input `int((x^2+a)^(1/2)*(x+(x^2+a)^(1/2))^n,x)`

output `int((x^2+a)^(1/2)*(x+(x^2+a)^(1/2))^n,x)`

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.64

$$\int \sqrt{a + x^2} \left( x + \sqrt{a + x^2} \right)^n dx = \frac{(n^2 x^2 + a n^2 - 2 \sqrt{x^2 + a} n x - 2 a) (x + \sqrt{x^2 + a})^n}{n^3 - 4 n}$$

input `integrate((x^2+a)^(1/2)*(x+(x^2+a)^(1/2))^n,x, algorithm="fricas")`

output  $(n^2x^2 + an^2 - 2\sqrt{x^2 + a})n^2x - 2a)(x + \sqrt{x^2 + a})^n / (n^3 - 4n)$

### Sympy [F]

$$\int \sqrt{a+x^2} (x + \sqrt{a+x^2})^n dx = \int \sqrt{a+x^2} (x + \sqrt{a+x^2})^n dx$$

input `integrate((x**2+a)**(1/2)*(x+(x**2+a)**(1/2))**n,x)`

output `Integral(sqrt(a + x**2)*(x + sqrt(a + x**2))**n, x)`

### Maxima [F]

$$\int \sqrt{a+x^2} (x + \sqrt{a+x^2})^n dx = \int \sqrt{x^2+a} (x + \sqrt{x^2+a})^n dx$$

input `integrate((x^2+a)^(1/2)*(x+(x^2+a)^(1/2))^n,x, algorithm="maxima")`

output `integrate(sqrt(x^2 + a)*(x + sqrt(x^2 + a))^n, x)`

### Giac [F]

$$\int \sqrt{a+x^2} (x + \sqrt{a+x^2})^n dx = \int \sqrt{x^2+a} (x + \sqrt{x^2+a})^n dx$$

input `integrate((x^2+a)^(1/2)*(x+(x^2+a)^(1/2))^n,x, algorithm="giac")`

output `integrate(sqrt(x^2 + a)*(x + sqrt(x^2 + a))^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \sqrt{a+x^2} (x + \sqrt{a+x^2})^n dx = \int \sqrt{x^2+a} (x + \sqrt{x^2+a})^n dx$$

input `int((a + x^2)^(1/2)*(x + (a + x^2)^(1/2))^n,x)`

output `int((a + x^2)^(1/2)*(x + (a + x^2)^(1/2))^n, x)`

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.63

$$\int \sqrt{a+x^2} (x + \sqrt{a+x^2})^n dx = \frac{(\sqrt{x^2+a} + x)^n (-2\sqrt{x^2+a}nx + an^2 - 2a + n^2x^2)}{n(n^2 - 4)}$$

input `int((x^2+a)^(1/2)*(x+(x^2+a)^(1/2))^n,x)`

output `((sqrt(a + x**2) + x)**n*(- 2*sqrt(a + x**2)*n*x + a*n**2 - 2*a + n**2*x**2))/(n*(n**2 - 4))`

$$3.29 \quad \int \frac{(x + \sqrt{a + x^2})^n}{\sqrt{a + x^2}} dx$$

Optimal result	221
Mathematica [A] (verified)	221
Rubi [A] (verified)	222
Maple [A] (verified)	223
Fricas [A] (verification not implemented)	223
Sympy [B] (verification not implemented)	223
Maxima [F]	224
Giac [A] (verification not implemented)	224
Mupad [B] (verification not implemented)	225
Reduce [B] (verification not implemented)	225

### Optimal result

Integrand size = 23, antiderivative size = 17

$$\int \frac{(x + \sqrt{a + x^2})^n}{\sqrt{a + x^2}} dx = \frac{(x + \sqrt{a + x^2})^n}{n}$$

output  $(x + \sqrt{a + x^2})^n / n$

### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{(x + \sqrt{a + x^2})^n}{\sqrt{a + x^2}} dx = \frac{(x + \sqrt{a + x^2})^n}{n}$$

input `Integrate[(x + Sqrt[a + x^2])^n/Sqrt[a + x^2], x]`

output  $(x + \sqrt{a + x^2})^n / n$

**Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$ , Rules used = {2547, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(\sqrt{a+x^2}+x)^n}{\sqrt{a+x^2}} dx$$

↓ 2547

$$\int (\sqrt{a+x^2}+x)^{n-1} d(\sqrt{a+x^2}+x)$$

↓ 15

$$\frac{(\sqrt{a+x^2}+x)^n}{n}$$

input `Int[(x + Sqrt[a + x^2])^n/Sqrt[a + x^2],x]`

output `(x + Sqrt[a + x^2])^n/n`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2547 `Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

**Maple [A] (verified)**

Time = 0.13 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

method	result	size
derivativedivides	$\frac{(x+\sqrt{x^2+a})^n}{n}$	16
default	$\frac{(x+\sqrt{x^2+a})^n}{n}$	16

input `int((x+(x^2+a)^(1/2))^n/(x^2+a)^(1/2),x,method=_RETURNVERBOSE)`

output `(x+(x^2+a)^(1/2))^n/n`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{(x + \sqrt{a + x^2})^n}{\sqrt{a + x^2}} dx = \frac{(x + \sqrt{x^2 + a})^n}{n}$$

input `integrate((x+(x^2+a)^(1/2))^n/(x^2+a)^(1/2),x, algorithm="fricas")`

output `(x + sqrt(x^2 + a))^n/n`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 311 vs.  $2(12) = 24$ .

Time = 1.75 (sec) , antiderivative size = 311, normalized size of antiderivative = 18.29

$$\int \frac{(x + \sqrt{a + x^2})^n}{\sqrt{a + x^2}} dx$$

$$= \begin{cases} \frac{\sqrt{a} a^{\frac{n}{2}} \sinh\left(n \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right) - \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right)\right)}{n x \sqrt{\frac{a}{x^2} + 1}} - \frac{2 a^{\frac{n}{2}} \cosh\left(n \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right)\right) \Gamma\left(1 - \frac{n}{2}\right)}{n^2 \Gamma\left(-\frac{n}{2}\right)} + \frac{a^{\frac{n}{2}} x \cosh\left(n \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right) - \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right)\right)}{\sqrt{a n}} + \frac{a^{\frac{n}{2}} x \sinh\left(n \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right) - \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right)\right)}{\sqrt{a n}} \\ \frac{a^{\frac{n}{2}} \sinh\left(n \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right) - \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right)\right)}{n x \sqrt{1 + \frac{x^2}{a}}} - \frac{2 a^{\frac{n}{2}} \cosh\left(n \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right)\right) \Gamma\left(1 - \frac{n}{2}\right)}{n^2 \Gamma\left(-\frac{n}{2}\right)} + \frac{a^{\frac{n}{2}} x^2 \sinh\left(n \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right) - \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right)\right)}{a n \sqrt{1 + \frac{x^2}{a}}} + \frac{a^{\frac{n}{2}} x \cosh\left(n \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right) - \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right)\right)}{a n \sqrt{1 + \frac{x^2}{a}}} \end{cases}$$



input `integrate((x+(x**2+a)**(1/2))**n/(x**2+a)**(1/2),x)`

output `Piecewise((sqrt(a)*a**(n/2)*sinh(n*asinh(x/sqrt(a))) - asinh(x/sqrt(a)))/(n*x*sqrt(a/x**2 + 1)) - 2*a**(n/2)*cosh(n*asinh(x/sqrt(a)))*gamma(1 - n/2)/(n**2*gamma(-n/2)) + a**(n/2)*x*cosh(n*asinh(x/sqrt(a)) - asinh(x/sqrt(a)))/(sqrt(a)*n) + a**(n/2)*x*sinh(n*asinh(x/sqrt(a)) - asinh(x/sqrt(a)))/(sqrt(a)*n*sqrt(a/x**2 + 1)), Abs(x**2/a) > 1), (a**(n/2)*sinh(n*asinh(x/sqrt(a)) - asinh(x/sqrt(a)))/(n*sqrt(1 + x**2/a)) - 2*a**(n/2)*cosh(n*asinh(x/sqrt(a)))*gamma(1 - n/2)/(n**2*gamma(-n/2)) + a**(n/2)*x**2*sinh(n*asinh(x/sqrt(a)) - asinh(x/sqrt(a)))/(a*n*sqrt(1 + x**2/a)) + a**(n/2)*x*cosh(n*asinh(x/sqrt(a)) - asinh(x/sqrt(a)))/(sqrt(a)*n), True))`

## Maxima [F]

$$\int \frac{(x + \sqrt{a + x^2})^n}{\sqrt{a + x^2}} dx = \int \frac{(x + \sqrt{x^2 + a})^n}{\sqrt{x^2 + a}} dx$$

input `integrate((x+(x^2+a)^(1/2))^n/(x^2+a)^(1/2),x, algorithm="maxima")`

output `integrate((x + sqrt(x^2 + a))^n/sqrt(x^2 + a), x)`

## Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{(x + \sqrt{a + x^2})^n}{\sqrt{a + x^2}} dx = \frac{(x + \sqrt{x^2 + a})^n}{n}$$

input `integrate((x+(x^2+a)^(1/2))^n/(x^2+a)^(1/2),x, algorithm="giac")`

output `(x + sqrt(x^2 + a))^n/n`

**Mupad [B] (verification not implemented)**

Time = 21.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{(x + \sqrt{a + x^2})^n}{\sqrt{a + x^2}} dx = \frac{(x + \sqrt{x^2 + a})^n}{n}$$

input `int((x + (a + x^2)^(1/2))^n/(a + x^2)^(1/2), x)`output `(x + (a + x^2)^(1/2))^n/n`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \frac{(x + \sqrt{a + x^2})^n}{\sqrt{a + x^2}} dx = \frac{(\sqrt{x^2 + a} + x)^n}{n}$$

input `int((x+(x^2+a)^(1/2))^n/(x^2+a)^(1/2), x)`output `(sqrt(a + x**2) + x)**n/n`

**3.30**  $\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^{3/2}} dx$

Optimal result	226
Mathematica [A] (verified)	226
Rubi [A] (verified)	227
Maple [F]	228
Fricas [F]	228
Sympy [F]	229
Maxima [F]	229
Giac [F]	229
Mupad [F(-1)]	230
Reduce [F]	230

**Optimal result**

Integrand size = 23, antiderivative size = 59

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^{3/2}} dx = \frac{4(x + \sqrt{a + x^2})^{2+n} \text{Hypergeometric2F1}\left(2, \frac{2+n}{2}, \frac{4+n}{2}, -\frac{(x + \sqrt{a + x^2})^2}{a}\right)}{a^2(2 + n)}$$

output

```
4*(x+(x^2+a)^(1/2))^(2+n)*hypergeom([2, 1+1/2*n], [2+1/2*n], -(x+(x^2+a)^(1/2))^2/a)/a^2/(2+n)
```

**Mathematica [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.03

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^{3/2}} dx = \frac{4(x + \sqrt{a + x^2})^{2+n} \text{Hypergeometric2F1}\left(2, \frac{2+n}{2}, 1 + \frac{2+n}{2}, -\frac{(x + \sqrt{a + x^2})^2}{a}\right)}{a^2(2 + n)}$$

input

```
Integrate[(x + Sqrt[a + x^2])^n/(a + x^2)^(3/2), x]
```

output

$$(4*(x + \text{Sqrt}[a + x^2])^{(2 + n)}*\text{Hypergeometric2F1}[2, (2 + n)/2, 1 + (2 + n)/2, -((x + \text{Sqrt}[a + x^2])^2/a)])/(a^2*(2 + n))$$
**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$ , Rules used = {2547, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(\sqrt{a+x^2}+x)^n}{(a+x^2)^{3/2}} dx$$

$$\downarrow 2547$$

$$4 \int \frac{(x + \sqrt{x^2 + a})^{n+1}}{\left( (x + \sqrt{x^2 + a})^2 + a \right)^2} d(x + \sqrt{x^2 + a})$$

$$\downarrow 278$$

$$\frac{4(\sqrt{a+x^2}+x)^{n+2} \text{Hypergeometric2F1}\left(2, \frac{n+2}{2}, \frac{n+4}{2}, -\frac{(x+\sqrt{x^2+a})^2}{a}\right)}{a^2(n+2)}$$

input

$$\text{Int}[(x + \text{Sqrt}[a + x^2])^n/(a + x^2)^{(3/2)}, x]$$

output

$$(4*(x + \text{Sqrt}[a + x^2])^{(2 + n)}*\text{Hypergeometric2F1}[2, (2 + n)/2, (4 + n)/2, -((x + \text{Sqrt}[a + x^2])^2/a)])/(a^2*(2 + n))$$

## Definitions of rubi rules used

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 2547 `Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

## Maple [F]

$$\int \frac{(x + \sqrt{x^2 + a})^n}{(x^2 + a)^{\frac{3}{2}}} dx$$

input `int((x+(x^2+a)^(1/2))^n/(x^2+a)^(3/2),x)`

output `int((x+(x^2+a)^(1/2))^n/(x^2+a)^(3/2),x)`

## Fricas [F]

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^{3/2}} dx = \int \frac{(x + \sqrt{x^2 + a})^n}{(x^2 + a)^{\frac{3}{2}}} dx$$

input `integrate((x+(x^2+a)^(1/2))^n/(x^2+a)^(3/2),x, algorithm="fricas")`

output `integral(sqrt(x^2 + a)*(x + sqrt(x^2 + a))^n/(x^4 + 2*a*x^2 + a^2), x)`

**Sympy [F]**

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^{3/2}} dx = \int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^{\frac{3}{2}}} dx$$

input `integrate((x+(x**2+a)**(1/2))**n/(x**2+a)**(3/2),x)`

output `Integral((x + sqrt(a + x**2))**n/(a + x**2)**(3/2), x)`

**Maxima [F]**

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^{3/2}} dx = \int \frac{(x + \sqrt{x^2 + a})^n}{(x^2 + a)^{\frac{3}{2}}} dx$$

input `integrate((x+(x^2+a)^(1/2))^n/(x^2+a)^(3/2),x, algorithm="maxima")`

output `integrate((x + sqrt(x^2 + a))^n/(x^2 + a)^(3/2), x)`

**Giac [F]**

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^{3/2}} dx = \int \frac{(x + \sqrt{x^2 + a})^n}{(x^2 + a)^{\frac{3}{2}}} dx$$

input `integrate((x+(x^2+a)^(1/2))^n/(x^2+a)^(3/2),x, algorithm="giac")`

output `integrate((x + sqrt(x^2 + a))^n/(x^2 + a)^(3/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^{3/2}} dx = \int \frac{(x + \sqrt{x^2 + a})^n}{(x^2 + a)^{3/2}} dx$$

input `int((x + (a + x^2)^(1/2))^n/(a + x^2)^(3/2),x)`output `int((x + (a + x^2)^(1/2))^n/(a + x^2)^(3/2), x)`**Reduce [F]**

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^{3/2}} dx = \int \frac{(\sqrt{x^2 + a} + x)^n}{\sqrt{x^2 + a} a + \sqrt{x^2 + a} x^2} dx$$

input `int((x+(x^2+a)^(1/2))^n/(x^2+a)^(3/2),x)`output `int((sqrt(a + x**2) + x)**n/(sqrt(a + x**2)*a + sqrt(a + x**2)*x**2),x)`

**3.31** 
$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^{5/2}} dx$$

Optimal result	231
Mathematica [A] (verified)	231
Rubi [A] (verified)	232
Maple [F]	233
Fricas [F]	233
Sympy [F]	234
Maxima [F]	234
Giac [F]	234
Mupad [F(-1)]	235
Reduce [F]	235

**Optimal result**

Integrand size = 23, antiderivative size = 59

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^{5/2}} dx = \frac{16(x + \sqrt{a + x^2})^{4+n} \text{Hypergeometric2F1}\left(4, \frac{4+n}{2}, \frac{6+n}{2}, -\frac{(x + \sqrt{a + x^2})^2}{a}\right)}{a^4(4 + n)}$$

output

```
16*(x+(x^2+a)^(1/2))^(4+n)*hypergeom([4, 2+1/2*n], [3+1/2*n], -(x+(x^2+a)^(1/2))^2/a)/a^4/(4+n)
```

**Mathematica [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.03

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^{5/2}} dx = \frac{16(x + \sqrt{a + x^2})^{4+n} \text{Hypergeometric2F1}\left(4, \frac{4+n}{2}, 1 + \frac{4+n}{2}, -\frac{(x + \sqrt{a + x^2})^2}{a}\right)}{a^4(4 + n)}$$

input

```
Integrate[(x + Sqrt[a + x^2])^n/(a + x^2)^(5/2), x]
```



output

```
(16*(x + Sqrt[a + x^2])^(4 + n)*Hypergeometric2F1[4, (4 + n)/2, 1 + (4 + n)
]/2, -((x + Sqrt[a + x^2])^2/a)))/(a^4*(4 + n))
```

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$ , Rules used = {2547, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(\sqrt{a+x^2}+x)^n}{(a+x^2)^{5/2}} dx$$

$$\downarrow 2547$$

$$16 \int \frac{(x + \sqrt{x^2 + a})^{n+3}}{\left( (x + \sqrt{x^2 + a})^2 + a \right)^4} d(x + \sqrt{x^2 + a})$$

$$\downarrow 278$$

$$\frac{16(\sqrt{a+x^2}+x)^{n+4} \text{Hypergeometric2F1}\left(4, \frac{n+4}{2}, \frac{n+6}{2}, -\frac{(x+\sqrt{x^2+a})^2}{a}\right)}{a^4(n+4)}$$

input

```
Int[(x + Sqrt[a + x^2])^n/(a + x^2)^(5/2), x]
```

output

```
(16*(x + Sqrt[a + x^2])^(4 + n)*Hypergeometric2F1[4, (4 + n)/2, (6 + n)/2,
-((x + Sqrt[a + x^2])^2/a)))/(a^4*(4 + n))
```

## Definitions of rubi rules used

rule 278

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])
```

rule 2547

```
Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])
```

## Maple [F]

$$\int \frac{(x + \sqrt{x^2 + a})^n}{(x^2 + a)^{\frac{5}{2}}} dx$$

input

```
int((x+(x^2+a)^(1/2))^n/(x^2+a)^(5/2),x)
```

output

```
int((x+(x^2+a)^(1/2))^n/(x^2+a)^(5/2),x)
```

## Fricas [F]

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^{5/2}} dx = \int \frac{(x + \sqrt{x^2 + a})^n}{(x^2 + a)^{\frac{5}{2}}} dx$$

input

```
integrate((x+(x^2+a)^(1/2))^n/(x^2+a)^(5/2),x, algorithm="fricas")
```

output

```
integral(sqrt(x^2 + a)*(x + sqrt(x^2 + a))^n/(x^6 + 3*a*x^4 + 3*a^2*x^2 + a^3), x)
```

**Sympy [F]**

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^{5/2}} dx = \int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^{\frac{5}{2}}} dx$$

input `integrate((x+(x**2+a)**(1/2))**n/(x**2+a)**(5/2),x)`

output `Integral((x + sqrt(a + x**2))**n/(a + x**2)**(5/2), x)`

**Maxima [F]**

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^{5/2}} dx = \int \frac{(x + \sqrt{x^2 + a})^n}{(x^2 + a)^{\frac{5}{2}}} dx$$

input `integrate((x+(x^2+a)^(1/2))^n/(x^2+a)^(5/2),x, algorithm="maxima")`

output `integrate((x + sqrt(x^2 + a))^n/(x^2 + a)^(5/2), x)`

**Giac [F]**

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^{5/2}} dx = \int \frac{(x + \sqrt{x^2 + a})^n}{(x^2 + a)^{\frac{5}{2}}} dx$$

input `integrate((x+(x^2+a)^(1/2))^n/(x^2+a)^(5/2),x, algorithm="giac")`

output `integrate((x + sqrt(x^2 + a))^n/(x^2 + a)^(5/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^{5/2}} dx = \int \frac{(x + \sqrt{x^2 + a})^n}{(x^2 + a)^{5/2}} dx$$

input `int((x + (a + x^2)^(1/2))^n/(a + x^2)^(5/2), x)`output `int((x + (a + x^2)^(1/2))^n/(a + x^2)^(5/2), x)`**Reduce [F]**

$$\int \frac{(x + \sqrt{a + x^2})^n}{(a + x^2)^{5/2}} dx = \int \frac{(\sqrt{x^2 + a} + x)^n}{\sqrt{x^2 + a} a^2 + 2\sqrt{x^2 + a} a x^2 + \sqrt{x^2 + a} x^4} dx$$

input `int((x+(x^2+a)^(1/2))^n/(x^2+a)^(5/2), x)`output `int((sqrt(a + x**2) + x)**n/(sqrt(a + x**2)*a**2 + 2*sqrt(a + x**2)*a*x**2 + sqrt(a + x**2)*x**4), x)`

### 3.32 $\int (a + x^2)^{5/2} \left(x - \sqrt{a + x^2}\right)^n dx$

Optimal result	236
Mathematica [A] (verified)	237
Rubi [A] (verified)	237
Maple [F]	239
Fricas [A] (verification not implemented)	239
Sympy [F(-2)]	239
Maxima [F]	240
Giac [F]	240
Mupad [F(-1)]	240
Reduce [B] (verification not implemented)	241

#### Optimal result

Integrand size = 25, antiderivative size = 201

$$\int (a + x^2)^{5/2} \left(x - \sqrt{a + x^2}\right)^n dx = \frac{a^6(x - \sqrt{a + x^2})^{-6+n}}{64(6 - n)} + \frac{3a^5(x - \sqrt{a + x^2})^{-4+n}}{32(4 - n)} + \frac{15a^4(x - \sqrt{a + x^2})^{-2+n}}{64(2 - n)} - \frac{5a^3(x - \sqrt{a + x^2})^n}{16n} - \frac{15a^2(x - \sqrt{a + x^2})^{2+n}}{64(2 + n)} - \frac{3a(x - \sqrt{a + x^2})^{4+n}}{32(4 + n)} - \frac{(x - \sqrt{a + x^2})^{6+n}}{64(6 + n)}$$

output

```
a^6*(x-(x^2+a)^(1/2))^(6-n)/(384-64*n)+3*a^5*(x-(x^2+a)^(1/2))^(4-n)/(128-32*n)+15*a^4*(x-(x^2+a)^(1/2))^(2-n)/(128-64*n)-5/16*a^3*(x-(x^2+a)^(1/2))^n/n-15*a^2*(x-(x^2+a)^(1/2))^(2+n)/(128+64*n)-3*a*(x-(x^2+a)^(1/2))^(4+n)/(128+32*n)-(x-(x^2+a)^(1/2))^(6+n)/(384+64*n)
```

**Mathematica [A] (verified)**

Time = 0.73 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.86

$$\int (a+x^2)^{5/2} (x-\sqrt{a+x^2})^n dx = \frac{1}{64} (x-\sqrt{a+x^2})^n \left( -\frac{20a^3}{n} - \frac{a^6}{(-6+n)(x-\sqrt{a+x^2})^6} - \frac{6a^5}{(-4+n)(x-\sqrt{a+x^2})^4} - \frac{15a^4}{(-2+n)(x-\sqrt{a+x^2})^2} - \frac{15a^2(x-\sqrt{a+x^2})^2}{2+n} - \frac{6a(x-\sqrt{a+x^2})^4}{4+n} - \frac{(x-\sqrt{a+x^2})^6}{6+n} \right)$$

input `Integrate[(a + x^2)^(5/2)*(x - Sqrt[a + x^2])^n,x]`

output `((x - Sqrt[a + x^2])^n*((-20*a^3)/n - a^6/((-6 + n)*(x - Sqrt[a + x^2])^6) - (6*a^5)/((-4 + n)*(x - Sqrt[a + x^2])^4) - (15*a^4)/((-2 + n)*(x - Sqrt[a + x^2])^2) - (15*a^2*(x - Sqrt[a + x^2])^2)/(2 + n) - (6*a*(x - Sqrt[a + x^2])^4)/(4 + n) - (x - Sqrt[a + x^2])^6/(6 + n))/64`

**Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 190, normalized size of antiderivative = 0.95, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$ , Rules used = {2547, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a+x^2)^{5/2} (x-\sqrt{a+x^2})^n dx$$

↓ 2547

$$-\frac{1}{64} \int (x-\sqrt{x^2+a})^{n-7} \left( (x-\sqrt{x^2+a})^2 + a \right)^6 d(x-\sqrt{x^2+a})$$

↓ 244

$$-\frac{1}{64} \int \left( a^6 (x - \sqrt{x^2 + a})^{n-7} + 6a^5 (x - \sqrt{x^2 + a})^{n-5} + 15a^4 (x - \sqrt{x^2 + a})^{n-3} + 20a^3 (x - \sqrt{x^2 + a})^{n-1} - \dots \right)$$

↓ 2009

$$\frac{1}{64} \left( \frac{a^6 (x - \sqrt{a + x^2})^{n-6}}{6-n} + \frac{6a^5 (x - \sqrt{a + x^2})^{n-4}}{4-n} + \frac{15a^4 (x - \sqrt{a + x^2})^{n-2}}{2-n} - \frac{20a^3 (x - \sqrt{a + x^2})^n}{n} - \frac{15a^2 (x - \sqrt{a + x^2})^{n+2}}{n+2} + \dots \right)$$

input `Int[(a + x^2)^(5/2)*(x - Sqrt[a + x^2])^n,x]`

output `((a^6*(x - Sqrt[a + x^2])^(-6 + n))/(6 - n) + (6*a^5*(x - Sqrt[a + x^2])^(-4 + n))/(4 - n) + (15*a^4*(x - Sqrt[a + x^2])^(-2 + n))/(2 - n) - (20*a^3*(x - Sqrt[a + x^2])^n)/n - (15*a^2*(x - Sqrt[a + x^2])^(2 + n))/(2 + n) - (6*a*(x - Sqrt[a + x^2])^(4 + n))/(4 + n) - (x - Sqrt[a + x^2])^(6 + n)/(6 + n))/64`

### Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2547 `Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

**Maple [F]**

$$\int (x^2 + a)^{\frac{5}{2}} (x - \sqrt{x^2 + a})^n dx$$

input `int((x^2+a)^(5/2)*(x-(x^2+a)^(1/2))^n,x)`

output `int((x^2+a)^(5/2)*(x-(x^2+a)^(1/2))^n,x)`

**Fricas [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.01

$$\int (a + x^2)^{5/2} (x - \sqrt{a + x^2})^n dx =$$

$$\frac{(a^3 n^6 - 50 a^3 n^4 + (n^6 - 20 n^4 + 64 n^2)x^6 + 544 a^3 n^2 + 3(an^6 - 30 an^4 + 104 an^2)x^4 - 720 a^3 + 3(a^2 n^6$$

input `integrate((x^2+a)^(5/2)*(x-(x^2+a)^(1/2))^n,x, algorithm="fricas")`

output `-(a^3*n^6 - 50*a^3*n^4 + (n^6 - 20*n^4 + 64*n^2)*x^6 + 544*a^3*n^2 + 3*(a*n^6 - 30*a*n^4 + 104*a*n^2)*x^4 - 720*a^3 + 3*(a^2*n^6 - 40*a^2*n^4 + 264*a^2*n^2)*x^2 + 6*((n^5 - 20*n^3 + 64*n)*x^5 + 2*(a*n^5 - 30*a*n^3 + 104*a*n)*x^3 + (a^2*n^5 - 40*a^2*n^3 + 264*a^2*n)*x)*sqrt(x^2 + a))*(x - sqrt(x^2 + a))^n/(n^7 - 56*n^5 + 784*n^3 - 2304*n)`

**Sympy [F(-2)]**

Exception generated.

$$\int (a + x^2)^{5/2} (x - \sqrt{a + x^2})^n dx = \text{Exception raised: HeuristicGCDFailed}$$

input `integrate((x**2+a)**(5/2)*(x-(x**2+a)**(1/2))**n,x)`

output `Exception raised: HeuristicGCDFailed >> no luck`



**Maxima [F]**

$$\int (a + x^2)^{5/2} (x - \sqrt{a + x^2})^n dx = \int (x^2 + a)^{5/2} (x - \sqrt{x^2 + a})^n dx$$

input `integrate((x^2+a)^(5/2)*(x-(x^2+a)^(1/2))^n,x, algorithm="maxima")`

output `integrate((x^2 + a)^(5/2)*(x - sqrt(x^2 + a))^n, x)`

**Giac [F]**

$$\int (a + x^2)^{5/2} (x - \sqrt{a + x^2})^n dx = \int (x^2 + a)^{5/2} (x - \sqrt{x^2 + a})^n dx$$

input `integrate((x^2+a)^(5/2)*(x-(x^2+a)^(1/2))^n,x, algorithm="giac")`

output `integrate((x^2 + a)^(5/2)*(x - sqrt(x^2 + a))^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (a + x^2)^{5/2} (x - \sqrt{a + x^2})^n dx = \int (x - \sqrt{x^2 + a})^n (x^2 + a)^{5/2} dx$$

input `int((x - (a + x^2)^(1/2))^n*(a + x^2)^(5/2), x)`

output `int((x - (a + x^2)^(1/2))^n*(a + x^2)^(5/2), x)`

**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 279, normalized size of antiderivative = 1.39

$$\int (a + x^2)^{5/2} (x - \sqrt{a + x^2})^n dx = \frac{a^n (-1)^n (-6\sqrt{x^2 + a} a^2 n^5 x + 240\sqrt{x^2 + a} a^2 n^3 x - 1584\sqrt{x^2 + a} a^2 n x - 12\sqrt{x^2 + a} a^2 n^5 x^3 + 360\sqrt{x^2 + a} a^2 n^3 x^3 - 1248\sqrt{x^2 + a} a^2 n x^3 - 6\sqrt{x^2 + a} a^2 n^5 x^5 + 120\sqrt{x^2 + a} a^2 n^3 x^5 - 384\sqrt{x^2 + a} a^2 n x^5 - a^3 n^6 x^2 + 50 a^3 n^4 x^2 - 544 a^3 n^2 x^2 + 720 a^3 x^2 - 3 a^2 n^6 x^2 + 120 a^2 n^4 x^2 - 792 a^2 n^2 x^2 - 3 a n^6 x^4 + 90 a n^4 x^4 - 312 a n^2 x^4 - n^6 x^6 + 20 n^4 x^6 - 64 n^2 x^6)}{(a + x^2)^{n+1}}$$

input

```
int((x^2+a)^(5/2)*(x-(x^2+a)^(1/2))^n,x)
```

output

```
(a**n*(-1)**n*(-6*sqrt(a+x**2)*a**2*n**5*x+240*sqrt(a+x**2)*a**2
*n**3*x-1584*sqrt(a+x**2)*a**2*n*x-12*sqrt(a+x**2)*a*n**5*x**3+3
60*sqrt(a+x**2)*a*n**3*x**3-1248*sqrt(a+x**2)*a*n*x**3-6*sqrt(a+
x**2)*n**5*x**5+120*sqrt(a+x**2)*n**3*x**5-384*sqrt(a+x**2)*n*x**5
-a**3*n**6+50*a**3*n**4-544*a**3*n**2+720*a**3-3*a**2*n**6*x**2
+120*a**2*n**4*x**2-792*a**2*n**2*x**2-3*a*n**6*x**4+90*a*n**4*x**4
-312*a*n**2*x**4-n**6*x**6+20*n**4*x**6-64*n**2*x**6)/((sqrt(a+
x**2)+x)**n*n*(n**6-56*n**4+784*n**2-2304))
```

### 3.33 $\int (a + x^2)^{3/2} \left(x - \sqrt{a + x^2}\right)^n dx$

Optimal result	242
Mathematica [A] (verified)	242
Rubi [A] (verified)	243
Maple [F]	244
Fricas [A] (verification not implemented)	245
Sympy [F]	245
Maxima [F]	245
Giac [F]	246
Mupad [F(-1)]	246
Reduce [B] (verification not implemented)	246

#### Optimal result

Integrand size = 25, antiderivative size = 141

$$\int (a + x^2)^{3/2} \left(x - \sqrt{a + x^2}\right)^n dx = \frac{a^4(x - \sqrt{a + x^2})^{-4+n}}{16(4 - n)} + \frac{a^3(x - \sqrt{a + x^2})^{-2+n}}{4(2 - n)} - \frac{3a^2(x - \sqrt{a + x^2})^n}{8n} - \frac{a(x - \sqrt{a + x^2})^{2+n}}{4(2 + n)} - \frac{(x - \sqrt{a + x^2})^{4+n}}{16(4 + n)}$$

output

```
a^4*(x-(x^2+a)^(1/2))^(4-n)/(64-16*n)+a^3*(x-(x^2+a)^(1/2))^(2-n)/(8-4*n)-3/8*a^2*(x-(x^2+a)^(1/2))^n/n-a*(x-(x^2+a)^(1/2))^(2+n)/(8+4*n)-(x-(x^2+a)^(1/2))^(4+n)/(64+16*n)
```

#### Mathematica [A] (verified)

Time = 0.38 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.87

$$\int (a + x^2)^{3/2} \left(x - \sqrt{a + x^2}\right)^n dx = \frac{1}{16} \left(x - \sqrt{a + x^2}\right)^n \left( -\frac{6a^2}{n} - \frac{a^4}{(-4 + n)(x - \sqrt{a + x^2})^4} - \frac{4a^3}{(-2 + n)(x - \sqrt{a + x^2})^2} - \frac{4a(x - \sqrt{a + x^2})^2}{2 + n} - \frac{(x - \sqrt{a + x^2})^4}{4 + n} \right)$$

input `Integrate[(a + x^2)^(3/2)*(x - Sqrt[a + x^2])^n,x]`

output `((x - Sqrt[a + x^2])^n*((-6*a^2)/n - a^4/((-4 + n)*(x - Sqrt[a + x^2])^4) - (4*a^3)/((-2 + n)*(x - Sqrt[a + x^2])^2) - (4*a*(x - Sqrt[a + x^2])^2)/(2 + n) - (x - Sqrt[a + x^2])^4/(4 + n))/16`

### Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.95, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$ , Rules used = {2547, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + x^2)^{3/2} (x - \sqrt{a + x^2})^n dx$$

$$\downarrow 2547$$

$$-\frac{1}{16} \int (x - \sqrt{x^2 + a})^{n-5} \left( (x - \sqrt{x^2 + a})^2 + a \right)^4 d(x - \sqrt{x^2 + a})$$

$$\downarrow 244$$

$$-\frac{1}{16} \int \left( a^4 (x - \sqrt{x^2 + a})^{n-5} + 4a^3 (x - \sqrt{x^2 + a})^{n-3} + 6a^2 (x - \sqrt{x^2 + a})^{n-1} + 4a (x - \sqrt{x^2 + a})^{n+1} + (x - \sqrt{x^2 + a})^{n+3} \right) dx$$

$$\downarrow 2009$$

$$\frac{1}{16} \left( \frac{a^4 (x - \sqrt{a + x^2})^{n-4}}{4 - n} + \frac{4a^3 (x - \sqrt{a + x^2})^{n-2}}{2 - n} - \frac{6a^2 (x - \sqrt{a + x^2})^n}{n} - \frac{4a (x - \sqrt{a + x^2})^{n+2}}{n + 2} - \frac{(x - \sqrt{a + x^2})^{n+4}}{n + 4} \right)$$

input `Int[(a + x^2)^(3/2)*(x - Sqrt[a + x^2])^n,x]`

output 
$$\frac{(a^4(x - \sqrt{a + x^2})^{-4 + n})}{(4 - n)} + \frac{(4a^3(x - \sqrt{a + x^2})^{-2 + n})}{(2 - n)} - \frac{(6a^2(x - \sqrt{a + x^2})^n)}{n} - \frac{(4a(x - \sqrt{a + x^2})^{2 + n})}{(2 + n)} - \frac{(x - \sqrt{a + x^2})^{4 + n}}{(4 + n)} / 16$$

### Defintions of rubi rules used

rule 244 
$$\text{Int}[\{(c\_)*(x\_)\}^{(m\_)}*\{(a\_)+(b\_)*(x\_)^2\}^{(p\_)}, x\_Symbol] \text{ :> Int[Expand Integrand}[\{(c*x)^m*(a + b*x^2)^p, x], x] \text{ /; FreeQ}\{a, b, c, m\}, x\} \&\& \text{IGtQ}\{p, 0\}$$

rule 2009 
$$\text{Int}[u_, x\_Symbol] \text{ :> Simp[IntSum}[u, x], x] \text{ /; SumQ}[u]$$

rule 2547 
$$\text{Int}[\{(g\_)+(i\_)*(x\_)^2\}^{(m\_)}*\{(d\_)+(e\_)*(x\_)+(f\_)*\text{Sqrt}[a\_+(c\_)*(x\_)^2]\}^{(n\_)}, x\_Symbol] \text{ :> Simp}[\{1/(2^{2*m+1}*e*f^{2*m})\}*(i/c)^m \text{ Subst[Int}[x^n*\{(d^2 + a*f^2 - 2*d*x + x^2)\}^{(2*m+1)}/\{-d + x\}^{(2*(m+1))}, x], x, d + e*x + f*\text{Sqrt}[a + c*x^2]], x] \text{ /; FreeQ}\{a, c, d, e, f, g, i, n\}, x\} \&\& \text{EqQ}\{e^2 - c*f^2, 0\} \&\& \text{EqQ}\{c*g - a*i, 0\} \&\& \text{IntegerQ}[2*m] \&\& (\text{IntegerQ}[m] \text{ || GtQ}\{i/c, 0\})$$

### Maple [F]

$$\int (x^2 + a)^{\frac{3}{2}} (x - \sqrt{x^2 + a})^n dx$$

input 
$$\text{int}((x^2+a)^{(3/2)}*(x-(x^2+a)^{(1/2)})^n,x)$$

output 
$$\text{int}((x^2+a)^{(3/2)}*(x-(x^2+a)^{(1/2)})^n,x)$$

**Fricas [A] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.80

$$\int (a + x^2)^{3/2} (x - \sqrt{a + x^2})^n dx = \frac{(a^2 n^4 + (n^4 - 4n^2)x^4 - 16a^2 n^2 + 2(an^4 - 10an^2)x^2 + 24a^2 + 4((n^3 - 4n)x^3 + (an^3 - 10an)x)\sqrt{x^2 + a})}{n^5 - 20n^3 + 64n}$$

input `integrate((x^2+a)^(3/2)*(x-(x^2+a)^(1/2))^n,x, algorithm="fricas")`

output `-(a^2*n^4 + (n^4 - 4*n^2)*x^4 - 16*a^2*n^2 + 2*(a*n^4 - 10*a*n^2)*x^2 + 24*a^2 + 4*((n^3 - 4*n)*x^3 + (a*n^3 - 10*a*n)*x)*sqrt(x^2 + a))*(x - sqrt(x^2 + a))^n/(n^5 - 20*n^3 + 64*n)`

**Sympy [F]**

$$\int (a + x^2)^{3/2} (x - \sqrt{a + x^2})^n dx = \int (a + x^2)^{\frac{3}{2}} (x - \sqrt{a + x^2})^n dx$$

input `integrate((x**2+a)**(3/2)*(x-(x**2+a)**(1/2))**n,x)`

output `Integral((a + x**2)**(3/2)*(x - sqrt(a + x**2))**n, x)`

**Maxima [F]**

$$\int (a + x^2)^{3/2} (x - \sqrt{a + x^2})^n dx = \int (x^2 + a)^{\frac{3}{2}} (x - \sqrt{x^2 + a})^n dx$$

input `integrate((x^2+a)^(3/2)*(x-(x^2+a)^(1/2))^n,x, algorithm="maxima")`

output `integrate((x^2 + a)^(3/2)*(x - sqrt(x^2 + a))^n, x)`

**Giac [F]**

$$\int (a + x^2)^{3/2} (x - \sqrt{a + x^2})^n dx = \int (x^2 + a)^{\frac{3}{2}} (x - \sqrt{x^2 + a})^n dx$$

input `integrate((x^2+a)^(3/2)*(x-(x^2+a)^(1/2))^n,x, algorithm="giac")`

output `integrate((x^2 + a)^(3/2)*(x - sqrt(x^2 + a))^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (a + x^2)^{3/2} (x - \sqrt{a + x^2})^n dx = \int (x - \sqrt{x^2 + a})^n (x^2 + a)^{3/2} dx$$

input `int((x - (a + x^2)^(1/2))^n*(a + x^2)^(3/2), x)`

output `int((x - (a + x^2)^(1/2))^n*(a + x^2)^(3/2), x)`

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.99

$$\int (a + x^2)^{3/2} (x - \sqrt{a + x^2})^n dx = \frac{a^n (-1)^n (-4\sqrt{x^2 + a} a n^3 x + 40\sqrt{x^2 + a} a n x - 4\sqrt{x^2 + a} n^3 x^3 + 16\sqrt{x^2 + a} n x^3 - a^2 n^3 x^5 + 4a^2 n^3 x^3 - 4a^2 n^3 x) (\sqrt{x^2 + a} + x)^n (n^4 - 20n^2 + 16)}{(\sqrt{x^2 + a} + x)^n (n^4 - 20n^2 + 16)}$$

input `int((x^2+a)^(3/2)*(x-(x^2+a)^(1/2))^n,x)`

output

```
(a**n*( - 1)**n*( - 4*sqrt(a + x**2)*a*n**3*x + 40*sqrt(a + x**2)*a*n*x -  
4*sqrt(a + x**2)*n**3*x**3 + 16*sqrt(a + x**2)*n*x**3 - a**2*n**4 + 16*a**  
2*n**2 - 24*a**2 - 2*a*n**4*x**2 + 20*a*n**2*x**2 - n**4*x**4 + 4*n**2*x**  
4))/((sqrt(a + x**2) + x)**n*n*(n**4 - 20*n**2 + 64))
```



### 3.34 $\int \sqrt{a+x^2} \left(x - \sqrt{a+x^2}\right)^n dx$

Optimal result	248
Mathematica [A] (verified)	248
Rubi [A] (verified)	249
Maple [F]	250
Fricas [A] (verification not implemented)	250
Sympy [F]	251
Maxima [F]	251
Giac [F]	251
Mupad [F(-1)]	252
Reduce [B] (verification not implemented)	252

#### Optimal result

Integrand size = 25, antiderivative size = 81

$$\int \sqrt{a+x^2} \left(x - \sqrt{a+x^2}\right)^n dx = \frac{a^2(x - \sqrt{a+x^2})^{-2+n}}{4(2-n)} - \frac{a(x - \sqrt{a+x^2})^n}{2n} - \frac{(x - \sqrt{a+x^2})^{2+n}}{4(2+n)}$$

output

$a^2*(x-(x^2+a)^{(1/2)})^{(-2+n)}/(8-4*n)-1/2*a*(x-(x^2+a)^{(1/2)})^n/n-(x-(x^2+a)^{(1/2)})^{(2+n)}/(8+4*n)$

#### Mathematica [A] (verified)

Time = 0.31 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.90

$$\int \sqrt{a+x^2} \left(x - \sqrt{a+x^2}\right)^n dx = \frac{1}{4} \left(x - \sqrt{a+x^2}\right)^n \left( -\frac{2a}{n} - \frac{a^2}{(-2+n)(x - \sqrt{a+x^2})^2} - \frac{(x - \sqrt{a+x^2})^2}{2+n} \right)$$

input

`Integrate[Sqrt[a + x^2]*(x - Sqrt[a + x^2])^n,x]`

output  $\frac{((x - \sqrt{a + x^2})^n)^{(-2*a)/n - a^2/((-2 + n)*(x - \sqrt{a + x^2})^2) - (x - \sqrt{a + x^2})^{2/(2 + n))}}{4}$

### Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.96, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$ , Rules used = {2547, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{a + x^2} (x - \sqrt{a + x^2})^n dx$$

$$\downarrow 2547$$

$$-\frac{1}{4} \int (x - \sqrt{x^2 + a})^{n-3} \left( (x - \sqrt{x^2 + a})^2 + a \right)^2 d(x - \sqrt{x^2 + a})$$

$$\downarrow 244$$

$$-\frac{1}{4} \int \left( a^2 (x - \sqrt{x^2 + a})^{n-3} + 2a (x - \sqrt{x^2 + a})^{n-1} + (x - \sqrt{x^2 + a})^{n+1} \right) d(x - \sqrt{x^2 + a})$$

$$\downarrow 2009$$

$$\frac{1}{4} \left( \frac{a^2 (x - \sqrt{a + x^2})^{n-2}}{2 - n} - \frac{2a (x - \sqrt{a + x^2})^n}{n} - \frac{(x - \sqrt{a + x^2})^{n+2}}{n + 2} \right)$$

input `Int[Sqrt[a + x^2]*(x - Sqrt[a + x^2])^n,x]`

output  $\frac{((a^2*(x - \sqrt{a + x^2})^{(-2 + n))}/(2 - n) - (2*a*(x - \sqrt{a + x^2})^n)/n - (x - \sqrt{a + x^2})^{(2 + n)/(2 + n))}}{4}$

**Defintions of rubi rules used**

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2547 `Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

**Maple [F]**

$$\int \sqrt{x^2 + a} \left( x - \sqrt{x^2 + a} \right)^n dx$$

input `int((x^2+a)^(1/2)*(x-(x^2+a)^(1/2))^n,x)`

output `int((x^2+a)^(1/2)*(x-(x^2+a)^(1/2))^n,x)`

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.63

$$\int \sqrt{a + x^2} \left( x - \sqrt{a + x^2} \right)^n dx = -\frac{(n^2 x^2 + a n^2 + 2 \sqrt{x^2 + a} n x - 2 a) (x - \sqrt{x^2 + a})^n}{n^3 - 4 n}$$

input `integrate((x^2+a)^(1/2)*(x-(x^2+a)^(1/2))^n,x, algorithm="fricas")`

output  $-(n^2x^2 + a n^2 + 2\sqrt{x^2 + a}nx - 2a)(x - \sqrt{x^2 + a})^n / (n^3 - 4n)$

### Sympy [F]

$$\int \sqrt{a+x^2} (x - \sqrt{a+x^2})^n dx = \int \sqrt{a+x^2} (x - \sqrt{a+x^2})^n dx$$

input `integrate((x**2+a)**(1/2)*(x-(x**2+a)**(1/2))**n,x)`

output `Integral(sqrt(a + x**2)*(x - sqrt(a + x**2))**n, x)`

### Maxima [F]

$$\int \sqrt{a+x^2} (x - \sqrt{a+x^2})^n dx = \int \sqrt{x^2+a} (x - \sqrt{x^2+a})^n dx$$

input `integrate((x^2+a)^(1/2)*(x-(x^2+a)^(1/2))^n,x, algorithm="maxima")`

output `integrate(sqrt(x^2 + a)*(x - sqrt(x^2 + a))^n, x)`

### Giac [F]

$$\int \sqrt{a+x^2} (x - \sqrt{a+x^2})^n dx = \int \sqrt{x^2+a} (x - \sqrt{x^2+a})^n dx$$

input `integrate((x^2+a)^(1/2)*(x-(x^2+a)^(1/2))^n,x, algorithm="giac")`

output `integrate(sqrt(x^2 + a)*(x - sqrt(x^2 + a))^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \sqrt{a+x^2} (x - \sqrt{a+x^2})^n dx = \int (x - \sqrt{x^2+a})^n \sqrt{x^2+a} dx$$

input `int((x - (a + x^2)^(1/2))^n*(a + x^2)^(1/2), x)`

output `int((x - (a + x^2)^(1/2))^n*(a + x^2)^(1/2), x)`

**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.70

$$\int \sqrt{a+x^2} (x - \sqrt{a+x^2})^n dx = \frac{a^n (-1)^n (-2\sqrt{x^2+a} nx - a n^2 + 2a - n^2 x^2)}{(\sqrt{x^2+a} + x)^n n (n^2 - 4)}$$

input `int((x^2+a)^(1/2)*(x-(x^2+a)^(1/2))^n,x)`

output `(a**n*(-1)**n*(-2*sqrt(a+x**2)*n*x - a*n**2 + 2*a - n**2*x**2))/((sqrt(a+x**2) + x)**n*n*(n**2 - 4))`

**3.35** 
$$\int \frac{(x - \sqrt{a+x^2})^n}{\sqrt{a+x^2}} dx$$

Optimal result . . . . .	253
Mathematica [A] (verified) . . . . .	253
Rubi [A] (verified) . . . . .	254
Maple [A] (verified) . . . . .	255
Fricas [A] (verification not implemented) . . . . .	255
Sympy [B] (verification not implemented) . . . . .	255
Maxima [F] . . . . .	256
Giac [A] (verification not implemented) . . . . .	256
Mupad [B] (verification not implemented) . . . . .	257
Reduce [B] (verification not implemented) . . . . .	257

**Optimal result**

Integrand size = 25, antiderivative size = 20

$$\int \frac{(x - \sqrt{a+x^2})^n}{\sqrt{a+x^2}} dx = -\frac{(x - \sqrt{a+x^2})^n}{n}$$

output

`-(x-(x^2+a)^(1/2))^n/n`

**Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(x - \sqrt{a+x^2})^n}{\sqrt{a+x^2}} dx = -\frac{(x - \sqrt{a+x^2})^n}{n}$$

input

`Integrate[(x - Sqrt[a + x^2])^n/Sqrt[a + x^2],x]`

output

`-((x - Sqrt[a + x^2])^n/n)`

**Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$ , Rules used = {2547, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(x - \sqrt{a + x^2})^n}{\sqrt{a + x^2}} dx$$

↓ 2547

$$- \int (x - \sqrt{x^2 + a})^{n-1} d(x - \sqrt{x^2 + a})$$

↓ 15

$$-\frac{(x - \sqrt{a + x^2})^n}{n}$$

input `Int[(x - Sqrt[a + x^2])^n/Sqrt[a + x^2],x]`

output `-((x - Sqrt[a + x^2])^n/n)`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2547 `Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

**Maple [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

method	result	size
derivativedivides	$-\frac{(x-\sqrt{x^2+a})^n}{n}$	19
default	$-\frac{(x-\sqrt{x^2+a})^n}{n}$	19

input `int((x-(x^2+a)^(1/2))^n/(x^2+a)^(1/2),x,method=_RETURNVERBOSE)`

output `-(x-(x^2+a)^(1/2))^n/n`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{(x - \sqrt{a + x^2})^n}{\sqrt{a + x^2}} dx = -\frac{(x - \sqrt{x^2 + a})^n}{n}$$

input `integrate((x-(x^2+a)^(1/2))^n/(x^2+a)^(1/2),x, algorithm="fricas")`

output `-(x - sqrt(x^2 + a))^n/n`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 41 vs. 2(14) = 28.

Time = 0.95 (sec) , antiderivative size = 41, normalized size of antiderivative = 2.05

$$\int \frac{(x - \sqrt{a + x^2})^n}{\sqrt{a + x^2}} dx = \begin{cases} -\frac{(x - \sqrt{a + x^2})^n}{n} & \text{for } n \neq 0 \\ \begin{cases} \log(2x + 2\sqrt{a + x^2}) & \text{for } a \neq 0 \\ \frac{x \log(x)}{\sqrt{x^2}} & \text{otherwise} \end{cases} & \text{otherwise} \end{cases}$$



input `integrate((x-(x**2+a)**(1/2))**n/(x**2+a)**(1/2),x)`

output `Piecewise((-x - sqrt(a + x**2))**n/n, Ne(n, 0)), (Piecewise((log(2*x + 2*sqrt(a + x**2)), Ne(a, 0)), (x*log(x)/sqrt(x**2), True)), True))`

### Maxima [F]

$$\int \frac{(x - \sqrt{a + x^2})^n}{\sqrt{a + x^2}} dx = \int \frac{(x - \sqrt{x^2 + a})^n}{\sqrt{x^2 + a}} dx$$

input `integrate((x-(x^2+a)^(1/2))^n/(x^2+a)^(1/2),x, algorithm="maxima")`

output `integrate((x - sqrt(x^2 + a))^n/sqrt(x^2 + a), x)`

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{(x - \sqrt{a + x^2})^n}{\sqrt{a + x^2}} dx = -\frac{(x - \sqrt{x^2 + a})^n}{n}$$

input `integrate((x-(x^2+a)^(1/2))^n/(x^2+a)^(1/2),x, algorithm="giac")`

output `-(x - sqrt(x^2 + a))^n/n`

**Mupad [B] (verification not implemented)**

Time = 20.93 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{(x - \sqrt{a + x^2})^n}{\sqrt{a + x^2}} dx = -\frac{(x - \sqrt{x^2 + a})^n}{n}$$

input `int((x - (a + x^2)^(1/2))^n/(a + x^2)^(1/2), x)`output `-(x - (a + x^2)^(1/2))^n/n`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.15

$$\int \frac{(x - \sqrt{a + x^2})^n}{\sqrt{a + x^2}} dx = -\frac{a^n(-1)^n}{(\sqrt{x^2 + a} + x)^n n}$$

input `int((x-(x^2+a)^(1/2))^n/(x^2+a)^(1/2), x)`output `( - a**n*( - 1)**n)/((sqrt(a + x**2) + x)**n*n)`

**3.36** 
$$\int \frac{(x - \sqrt{a+x^2})^n}{(a+x^2)^{3/2}} dx$$

Optimal result	258
Mathematica [A] (verified)	258
Rubi [A] (verified)	259
Maple [F]	260
Fricas [F]	260
Sympy [F]	261
Maxima [F]	261
Giac [F]	261
Mupad [F(-1)]	262
Reduce [F]	262

**Optimal result**

Integrand size = 25, antiderivative size = 63

$$\int \frac{(x - \sqrt{a+x^2})^n}{(a+x^2)^{3/2}} dx = \frac{4(x - \sqrt{a+x^2})^{2+n} \text{Hypergeometric2F1}\left(2, \frac{2+n}{2}, \frac{4+n}{2}, -\frac{(x - \sqrt{a+x^2})^2}{a}\right)}{a^2(2+n)}$$

output `-4*(x-(x^2+a)^(1/2))^(2+n)*hypergeom([2, 1+1/2*n], [2+1/2*n], -(x-(x^2+a)^(1/2))^2/a)/a^2/(2+n)`

**Mathematica [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.03

$$\int \frac{(x - \sqrt{a+x^2})^n}{(a+x^2)^{3/2}} dx = \frac{4(x - \sqrt{a+x^2})^{2+n} \text{Hypergeometric2F1}\left(2, \frac{2+n}{2}, 1 + \frac{2+n}{2}, -\frac{(x - \sqrt{a+x^2})^2}{a}\right)}{a^2(2+n)}$$

input `Integrate[(x - Sqrt[a + x^2])^n/(a + x^2)^(3/2),x]`

output `(-4*(x - Sqrt[a + x^2])^(2 + n)*Hypergeometric2F1[2, (2 + n)/2, 1 + (2 + n)/2, -((x - Sqrt[a + x^2])^2/a)]/(a^2*(2 + n))`

### Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$ , Rules used = {2547, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^{3/2}} dx$$

$$\downarrow 2547$$

$$-4 \int \frac{(x - \sqrt{x^2 + a})^{n+1}}{\left( (x - \sqrt{x^2 + a})^2 + a \right)^2} d(x - \sqrt{x^2 + a})$$

$$\downarrow 278$$

$$\frac{4(x - \sqrt{a + x^2})^{n+2} \text{Hypergeometric2F1}\left(2, \frac{n+2}{2}, \frac{n+4}{2}, -\frac{(x - \sqrt{x^2 + a})^2}{a}\right)}{a^2(n+2)}$$

input `Int[(x - Sqrt[a + x^2])^n/(a + x^2)^(3/2),x]`

output `(-4*(x - Sqrt[a + x^2])^(2 + n)*Hypergeometric2F1[2, (2 + n)/2, (4 + n)/2, -((x - Sqrt[a + x^2])^2/a)]/(a^2*(2 + n))`

## Definitions of rubi rules used

rule 278

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])
```

rule 2547

```
Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])
```

## Maple [F]

$$\int \frac{(x - \sqrt{x^2 + a})^n}{(x^2 + a)^{\frac{3}{2}}} dx$$

input

```
int((x-(x^2+a)^(1/2))^n/(x^2+a)^(3/2),x)
```

output

```
int((x-(x^2+a)^(1/2))^n/(x^2+a)^(3/2),x)
```

## Fricas [F]

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^{3/2}} dx = \int \frac{(x - \sqrt{x^2 + a})^n}{(x^2 + a)^{\frac{3}{2}}} dx$$

input

```
integrate((x-(x^2+a)^(1/2))^n/(x^2+a)^(3/2),x, algorithm="fricas")
```

output

```
integral(sqrt(x^2 + a)*(x - sqrt(x^2 + a))^n/(x^4 + 2*a*x^2 + a^2), x)
```

**Sympy [F]**

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^{3/2}} dx = \int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^{\frac{3}{2}}} dx$$

input `integrate((x-(x**2+a)**(1/2))**n/(x**2+a)**(3/2),x)`

output `Integral((x - sqrt(a + x**2))**n/(a + x**2)**(3/2), x)`

**Maxima [F]**

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^{3/2}} dx = \int \frac{(x - \sqrt{x^2 + a})^n}{(x^2 + a)^{\frac{3}{2}}} dx$$

input `integrate((x-(x^2+a)^(1/2))^n/(x^2+a)^(3/2),x, algorithm="maxima")`

output `integrate((x - sqrt(x^2 + a))^n/(x^2 + a)^(3/2), x)`

**Giac [F]**

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^{3/2}} dx = \int \frac{(x - \sqrt{x^2 + a})^n}{(x^2 + a)^{\frac{3}{2}}} dx$$

input `integrate((x-(x^2+a)^(1/2))^n/(x^2+a)^(3/2),x, algorithm="giac")`

output `integrate((x - sqrt(x^2 + a))^n/(x^2 + a)^(3/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^{3/2}} dx = \int \frac{(x - \sqrt{x^2 + a})^n}{(x^2 + a)^{3/2}} dx$$

input `int((x - (a + x^2)^(1/2))^n/(a + x^2)^(3/2),x)`output `int((x - (a + x^2)^(1/2))^n/(a + x^2)^(3/2), x)`**Reduce [F]**

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^{3/2}} dx = \int \frac{(-\sqrt{x^2 + a} + x)^n}{\sqrt{x^2 + a} a + \sqrt{x^2 + a} x^2} dx$$

input `int((x-(x^2+a)^(1/2))^n/(x^2+a)^(3/2),x)`output `int((-sqrt(a + x**2) + x)**n/(sqrt(a + x**2)*a + sqrt(a + x**2)*x**2),x)`

**3.37** 
$$\int \frac{(x - \sqrt{a+x^2})^n}{(a+x^2)^{5/2}} dx$$

Optimal result	263
Mathematica [A] (verified)	263
Rubi [A] (verified)	264
Maple [F]	265
Fricas [F]	265
Sympy [F]	266
Maxima [F]	266
Giac [F]	266
Mupad [F(-1)]	267
Reduce [F]	267

**Optimal result**

Integrand size = 25, antiderivative size = 63

$$\int \frac{(x - \sqrt{a+x^2})^n}{(a+x^2)^{5/2}} dx = \frac{16(x - \sqrt{a+x^2})^{4+n} \operatorname{Hypergeometric2F1}\left(4, \frac{4+n}{2}, \frac{6+n}{2}, -\frac{(x - \sqrt{a+x^2})^2}{a}\right)}{a^4(4+n)}$$

```
output -16*(x-(x^2+a)^(1/2))^(4+n)*hypergeom([4, 2+1/2*n],[3+1/2*n],-(x-(x^2+a)^(1/2))^2/a)/a^4/(4+n)
```

**Mathematica [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.03

$$\int \frac{(x - \sqrt{a+x^2})^n}{(a+x^2)^{5/2}} dx = \frac{16(x - \sqrt{a+x^2})^{4+n} \operatorname{Hypergeometric2F1}\left(4, \frac{4+n}{2}, 1 + \frac{4+n}{2}, -\frac{(x - \sqrt{a+x^2})^2}{a}\right)}{a^4(4+n)}$$



input `Integrate[(x - Sqrt[a + x^2])^n/(a + x^2)^(5/2),x]`

output `(-16*(x - Sqrt[a + x^2])^(4 + n)*Hypergeometric2F1[4, (4 + n)/2, 1 + (4 + n)/2, -((x - Sqrt[a + x^2])^2/a)])/(a^4*(4 + n))`

### Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$ , Rules used = {2547, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^{5/2}} dx$$

$$\downarrow 2547$$

$$-16 \int \frac{(x - \sqrt{x^2 + a})^{n+3}}{\left( (x - \sqrt{x^2 + a})^2 + a \right)^4} d(x - \sqrt{x^2 + a})$$

$$\downarrow 278$$

$$\frac{16(x - \sqrt{a + x^2})^{n+4} \operatorname{Hypergeometric2F1}\left(4, \frac{n+4}{2}, \frac{n+6}{2}, -\frac{(x - \sqrt{x^2 + a})^2}{a}\right)}{a^4(n+4)}$$

input `Int[(x - Sqrt[a + x^2])^n/(a + x^2)^(5/2),x]`

output `(-16*(x - Sqrt[a + x^2])^(4 + n)*Hypergeometric2F1[4, (4 + n)/2, (6 + n)/2, -((x - Sqrt[a + x^2])^2/a)])/(a^4*(4 + n))`

## Definitions of rubi rules used

rule 278

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])
```

rule 2547

```
Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])
```

## Maple [F]

$$\int \frac{(x - \sqrt{x^2 + a})^n}{(x^2 + a)^{\frac{5}{2}}} dx$$

input

```
int((x-(x^2+a)^(1/2))^n/(x^2+a)^(5/2),x)
```

output

```
int((x-(x^2+a)^(1/2))^n/(x^2+a)^(5/2),x)
```

## Fricas [F]

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^{5/2}} dx = \int \frac{(x - \sqrt{x^2 + a})^n}{(x^2 + a)^{\frac{5}{2}}} dx$$

input

```
integrate((x-(x^2+a)^(1/2))^n/(x^2+a)^(5/2),x, algorithm="fricas")
```

output

```
integral(sqrt(x^2 + a)*(x - sqrt(x^2 + a))^n/(x^6 + 3*a*x^4 + 3*a^2*x^2 + a^3), x)
```

**Sympy [F]**

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^{5/2}} dx = \int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^{\frac{5}{2}}} dx$$

input `integrate((x-(x**2+a)**(1/2))**n/(x**2+a)**(5/2),x)`

output `Integral((x - sqrt(a + x**2))**n/(a + x**2)**(5/2), x)`

**Maxima [F]**

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^{5/2}} dx = \int \frac{(x - \sqrt{x^2 + a})^n}{(x^2 + a)^{\frac{5}{2}}} dx$$

input `integrate((x-(x^2+a)^(1/2))^n/(x^2+a)^(5/2),x, algorithm="maxima")`

output `integrate((x - sqrt(x^2 + a))^n/(x^2 + a)^(5/2), x)`

**Giac [F]**

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^{5/2}} dx = \int \frac{(x - \sqrt{x^2 + a})^n}{(x^2 + a)^{\frac{5}{2}}} dx$$

input `integrate((x-(x^2+a)^(1/2))^n/(x^2+a)^(5/2),x, algorithm="giac")`

output `integrate((x - sqrt(x^2 + a))^n/(x^2 + a)^(5/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^{5/2}} dx = \int \frac{(x - \sqrt{x^2 + a})^n}{(x^2 + a)^{5/2}} dx$$

input `int((x - (a + x^2)^(1/2))^n/(a + x^2)^(5/2), x)`output `int((x - (a + x^2)^(1/2))^n/(a + x^2)^(5/2), x)`**Reduce [F]**

$$\int \frac{(x - \sqrt{a + x^2})^n}{(a + x^2)^{5/2}} dx = \int \frac{(-\sqrt{x^2 + a} + x)^n}{\sqrt{x^2 + a} a^2 + 2\sqrt{x^2 + a} a x^2 + \sqrt{x^2 + a} x^4} dx$$

input `int((x-(x^2+a)^(1/2))^n/(x^2+a)^(5/2), x)`output `int((-sqrt(a + x**2) + x)**n/(sqrt(a + x**2)*a**2 + 2*sqrt(a + x**2)*a*x**2 + sqrt(a + x**2)*x**4), x)`

**3.38**  $\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^2 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$

Optimal result	268
Mathematica [A] (verified)	269
Rubi [A] (verified)	270
Maple [F]	272
Fricas [A] (verification not implemented)	272
Sympy [F(-1)]	273
Maxima [F]	273
Giac [F]	274
Mupad [F(-1)]	274
Reduce [B] (verification not implemented)	275

**Optimal result**

Integrand size = 56, antiderivative size = 365

$$\begin{aligned} & \int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^2 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx \\ &= \frac{(d^2 - af^2)^5 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{-5+n}}{32ef^4(5 - n)} \\ & \quad - \frac{5(d^2 - af^2)^4 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{-3+n}}{32ef^4(3 - n)} \\ & \quad + \frac{5(d^2 - af^2)^3 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{-1+n}}{16ef^4(1 - n)} \\ & \quad + \frac{5(d^2 - af^2)^2 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{1+n}}{16ef^4(1 + n)} \\ & \quad - \frac{5(d^2 - af^2) \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{3+n}}{32ef^4(3 + n)} \\ & \quad + \frac{\left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{5+n}}{32ef^4(5 + n)} \end{aligned}$$

output

$$\begin{aligned} & \frac{1}{32}(-af^2+d^2)^5(d+ex+f(a+2d*ex/f^2+e^2*x^2/f^2))^{(1/2)}^{(-5+n)}/e/f \\ & ^4/(5-n)-5/32(-af^2+d^2)^4(d+ex+f(a+2d*ex/f^2+e^2*x^2/f^2))^{(1/2)}^{(-3+n)}/e/f^4/(3-n)+5/16(-af^2+d^2)^3(d+ex+f(a+2d*ex/f^2+e^2*x^2/f^2))^{(1/2)}^{(-1+n)}/e/f^4/(1-n)+5/16(-af^2+d^2)^2(d+ex+f(a+2d*ex/f^2+e^2*x^2/f^2))^{(1/2)}^{(1+n)}/e/f^4/(1+n)-5/32(-af^2+d^2)(d+ex+f(a+2d*ex/f^2+e^2*x^2/f^2))^{(1/2)}^{(3+n)}/e/f^4/(3+n)+1/32(d+ex+f(a+2d*ex/f^2+e^2*x^2/f^2))^{(1/2)}^{(5+n)}/e/f^4/(5+n) \end{aligned}$$

**Mathematica [A] (verified)**

Time = 6.29 (sec) , antiderivative size = 280, normalized size of antiderivative = 0.77

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^2 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$\left( d + ex + f \sqrt{a + \frac{ex(2d+ex)}{f^2}} \right)^{-5+n} \left( -\frac{(d^2-af^2)^5}{-5+n} + \frac{5(d^2-af^2)^4 \left( d+ex+f \sqrt{a + \frac{ex(2d+ex)}{f^2}} \right)^2}{-3+n} - \frac{10(d^2-af^2)^3 \left( d+ex+f \sqrt{a + \frac{ex(2d+ex)}{f^2}} \right)}{-1+n} \right)$$

32

input

```
Integrate[(a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2)^2*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n,x]
```

output

```
((d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^(-5 + n))*(-(d^2 - a*f^2)^5/(-5 + n)) + (5*(d^2 - a*f^2)^4*(d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^2)/(-3 + n) - (10*(d^2 - a*f^2)^3*(d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^4)/(-1 + n) + (10*(d^2 - a*f^2)^2*(d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^6)/(1 + n) - (5*(d^2 - a*f^2)*(d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^8)/(3 + n) + (d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^10/(5 + n))/(32*e*f^4)
```

### Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 327, normalized size of antiderivative = 0.90, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$ , Rules used = {2546, 27, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^2 \left( f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex \right)^n dx$$

↓ 2546

$$2 \int \frac{\left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-6} \left( d^2-af^2 - \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^2 \right)^5}{64e} d \left( d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right) dx$$

↓ 27

$$\int \frac{\left( d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-6} \left( d^2 - af^2 - \left( d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^2 \right)^5}{32ef^4} d \left( d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right) dx$$

↓ 244

$$\int \left( (d^2 - af^2)^5 \left( d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-6} - 5(d^2 - af^2)^4 \left( d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-4} + 10(d^2 - af^2)^3 \left( d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-2} - 5(d^2 - af^2)^2 \left( d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-2} + 10(d^2 - af^2) \left( d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-2} - 5 \left( d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-2} \right) dx$$

↓ 2009

$$\frac{(d^2-af^2)^5 \left( f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex \right)^{n-5}}{5-n} + \frac{5(d^2-af^2)^4 \left( f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex \right)^{n-3}}{3-n} - \frac{10(d^2-af^2)^3 \left( f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex \right)^{n-1}}{1-n} + \frac{5(d^2-af^2)^2 \left( f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex \right)^{n-1}}{1-n} - \frac{10(d^2-af^2) \left( f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex \right)^{n-1}}{1-n} + \frac{5 \left( f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex \right)^{n-1}}{1-n}$$

input

```
Int[(a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2)^2*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n,x]
```

output

$$\begin{aligned}
& -1/32 * ( - ( (d^2 - a*f^2)^5 * (d + e*x + f*\text{Sqrt}[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^{(-5 + n)} / (5 - n) ) \\
& + (5*(d^2 - a*f^2)^4 * (d + e*x + f*\text{Sqrt}[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^{(-3 + n)} / (3 - n) - (10*(d^2 - a*f^2)^3 * (d + e*x \\
& + f*\text{Sqrt}[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^{(-1 + n)} / (1 - n) - (10*(d^2 - a*f^2)^2 * (d + e*x + f*\text{Sqrt}[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^{(1 + n)} \\
& / (1 + n) + (5*(d^2 - a*f^2) * (d + e*x + f*\text{Sqrt}[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^{(3 + n)} / (3 + n) - (d + e*x + f*\text{Sqrt}[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^{(5 + n)} / (5 + n) ) / (e*f^4)
\end{aligned}$$

### Defintions of rubi rules used

rule 27

$$\text{Int}[(a_*)(F_x), x\_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_*)(G_x)] /; \text{FreeQ}[b, x]$$

rule 244

$$\text{Int}[(c_*)(x_)^{(m_*)} * ((a_) + (b_*)(x_)^2)^{(p_)}, x\_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m * (a + b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, m\}, x] \ \&\& \ \text{IGtQ}[p, 0]$$

rule 2009

$$\text{Int}[u_, x\_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$$

rule 2546

$$\begin{aligned}
& \text{Int}[(g_.) + (h_*)(x_) + (i_*)(x_)^2)^{(m_*)} * ((d_.) + (e_*)(x_) + (f_*)*\text{Sqrt}[(a_.) + (b_*)(x_) + (c_*)(x_)^2])^{(n_*)}, x\_Symbol] \rightarrow \text{Simp}[(2/f^{(2*m)}) * (i/c)^m \quad \text{Subst}[\text{Int}[x^n * ((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)^{(2*m + 1)} / (-2*d*e + b*f^2 + 2*e*x)^{(2*(m + 1))}], x], x, d + e*x + f*\text{Sqrt}[a + b*x + c*x^2]], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, h, i, n\}, x] \ \&\& \ \text{EqQ}[e^2 - c*f^2, 0] \ \&\& \ \text{EqQ}[c*g - a*i, 0] \ \&\& \ \text{EqQ}[c*h - b*i, 0] \ \&\& \ \text{IntegerQ}[2*m] \ \&\& \ (\text{IntegerQ}[m] \ || \ \text{GtQ}[i/c, 0])
\end{aligned}$$



**Maple [F]**

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^2 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

input

```
int((a+2*d*e*x/f^2+e^2*x^2/f^2)^2*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x)
```

output

```
int((a+2*d*e*x/f^2+e^2*x^2/f^2)^2*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x)
```

**Fricas [A] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 654, normalized size of antiderivative = 1.79

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^2 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx =$$

$$\frac{\left( 5a^2df^4n^4 + 225a^2df^4 - 300ad^3f^2 + 5(e^5n^4 - 10e^5n^2 + 9e^5)x^5 + 120d^5 + 25(de^4n^4 - 10de^4n^2 + \dots \right)}{\dots}$$

input

```
integrate((a+2*d*e*x/f^2+e^2*x^2/f^2)^2*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x, algorithm="fricas")
```

output

```

-(5*a^2*d*f^4*n^4 + 225*a^2*d*f^4 - 300*a*d^3*f^2 + 5*(e^5*n^4 - 10*e^5*n^
2 + 9*e^5)*x^5 + 120*d^5 + 25*(d*e^4*n^4 - 10*d*e^4*n^2 + 9*d*e^4)*x^4 + 1
0*(15*a*e^3*f^2 + 30*d^2*e^3 + (a*e^3*f^2 + 4*d^2*e^3)*n^4 - 2*(8*a*e^3*f^
2 + 17*d^2*e^3)*n^2)*x^3 - 10*(11*a^2*d*f^4 - 6*a*d^3*f^2)*n^2 + 10*(45*a*
d*e^2*f^2 + (3*a*d*e^2*f^2 + 2*d^3*e^2)*n^4 - 2*(24*a*d*e^2*f^2 + d^3*e^2)
*n^2)*x^2 + 5*(45*a^2*e*f^4 + (a^2*e*f^4 + 4*a*d^2*e*f^2)*n^4 - 2*(11*a^2*
e*f^4 + 26*a*d^2*e*f^2 - 12*d^4*e)*n^2)*x - (a^2*f^5*n^5 + (e^4*f*n^5 - 10
*e^4*f*n^3 + 9*e^4*f*n)*x^4 - 10*(3*a^2*f^5 - 2*a*d^2*f^3)*n^3 + 4*(d*e^3*
f*n^5 - 10*d*e^3*f*n^3 + 9*d*e^3*f*n)*x^3 + 2*((a*e^2*f^3 + 2*d^2*e^2*f)*n
^5 - 10*(2*a*e^2*f^3 + d^2*e^2*f)*n^3 + (19*a*e^2*f^3 + 8*d^2*e^2*f)*n)*x^
2 + (149*a^2*f^5 - 260*a*d^2*f^3 + 120*d^4*f)*n + 4*(a*d*e*f^3*n^5 - 10*(2
*a*d*e*f^3 - d^3*e*f)*n^3 + (19*a*d*e*f^3 - 10*d^3*e*f)*n)*x)*sqrt((e^2*x^
2 + a*f^2 + 2*d*e*x)/f^2))*(e*x + f*sqrt((e^2*x^2 + a*f^2 + 2*d*e*x)/f^2)
+ d)^n/(e*f^4*n^6 - 35*e*f^4*n^4 + 259*e*f^4*n^2 - 225*e*f^4)

```

**Sympy [F(-1)]**

Timed out.

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^2 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx = \text{Timed out}$$

input

```

integrate((a+2*d*e*x/f**2+e**2*x**2/f**2)**2*(d+e*x+f*(a+2*d*e*x/f**2+e**2
*x**2/f**2)**(1/2))**n,x)

```

output

Timed out

**Maxima [F]**

$$\begin{aligned} & \int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^2 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx \\ &= \int \left( \frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2} \right)^2 \left( ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}} f + d \right)^n dx \end{aligned}$$

input `integrate((a+2*d*e*x/f^2+e^2*x^2/f^2)^2*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x, algorithm="maxima")`

output `integrate((e^2*x^2/f^2 + a + 2*d*e*x/f^2)^2*(e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2)*f + d)^n, x)`

### Giac [F]

$$\begin{aligned} & \int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^2 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx \\ &= \int \left( \frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2} \right)^2 \left( ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}} f + d \right)^n dx \end{aligned}$$

input `integrate((a+2*d*e*x/f^2+e^2*x^2/f^2)^2*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x, algorithm="giac")`

output `integrate((e^2*x^2/f^2 + a + 2*d*e*x/f^2)^2*(e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2)*f + d)^n, x)`

### Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^2 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx \\ &= \int \left( d + f \sqrt{a + \frac{e^2x^2}{f^2} + \frac{2dex}{f^2}} + ex \right)^n \left( a + \frac{e^2x^2}{f^2} + \frac{2dex}{f^2} \right)^2 dx \end{aligned}$$

input `int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^2,x)`

output

```
int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^2, x)
```

**Reduce [B] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 1095, normalized size of antiderivative = 3.00

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^2 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx = \text{Too large to display}$$

input

```
int((a+2*d*e*x/f^2+e^2*x^2/f^2)^2*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x)
```

output

```
((sqrt(a*f**2 + 2*d*e*x + e**2*x**2) + d + e*x)**n*(sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*a**2*f**4*n**5 - 30*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*a**2*f**4*n**3 + 149*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*a**2*f**4*n + 20*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*a*d**2*f**2*n**3 - 260*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*a*d**2*f**2*n + 4*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*a*d*e*f**2*n**5*x - 80*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*a*d*e*f**2*n**3*x + 76*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*a*d*e*f**2*n*x + 2*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*a*e**2*f**2*n**5*x**2 - 40*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*a*e**2*f**2*n**3*x**2 + 38*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*a*e**2*f**2*n*x**2 + 120*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*d**4*n + 40*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*d**3*e*n**3*x - 40*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*d**3*e*n*x + 4*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*d**2*e**2*n**5*x**2 - 20*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*d**2*e**2*n**3*x**2 + 16*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*d**2*e**2*n*x**2 + 4*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*d*e**3*n**5*x**3 - 40*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*d*e**3*n**3*x**3 + 36*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*d*e**3*n*x**3 + sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*e**4*n**5*x**4 - 10*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*e**4*n**3*x**4 + 9*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*e**4*n*x**4 - 5*a**2*d*f**4*n**4 + 110*a**2*d*f**4*n**2 - 225*a**2*d*f**4 - 5*a**2*e*f**4*n**4*x + 110*a**2*e*f**4*n**2*x - 225*a**2*e*f**4*x - 60*a*d**3*f**2*n...
```

**3.39** 
$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right) \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

Optimal result	276
Mathematica [A] (verified)	277
Rubi [A] (verified)	277
Maple [F]	279
Fricas [A] (verification not implemented)	280
Sympy [F]	280
Maxima [F]	281
Giac [F]	281
Mupad [F(-1)]	282
Reduce [B] (verification not implemented)	282

**Optimal result**

Integrand size = 54, antiderivative size = 239

$$\begin{aligned} & \int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right) \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx \\ &= \frac{(d^2 - af^2)^3 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{-3+n}}{8ef^2(3 - n)} \\ & \quad - \frac{3(d^2 - af^2)^2 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{-1+n}}{8ef^2(1 - n)} \\ & \quad - \frac{3(d^2 - af^2) \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{1+n}}{8ef^2(1 + n)} \\ & \quad + \frac{\left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{3+n}}{8ef^2(3 + n)} \end{aligned}$$

output

```
1/8*(-a*f^2+d^2)^3*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^(3-n)/e/f^2/(3-n)-3/8*(-a*f^2+d^2)^2*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^(1+n)/e/f^2/(1-n)-3/8*(-a*f^2+d^2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^(1+n)/e/f^2/(1+n)+1/8*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^(3+n)/e/f^2/(3+n)
```

### Mathematica [A] (verified)

Time = 2.43 (sec) , antiderivative size = 186, normalized size of antiderivative = 0.78

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right) \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \frac{\left( d + ex + f \sqrt{a + \frac{ex(2d+ex)}{f^2}} \right)^{-3+n} \left( -\frac{(d^2-af^2)^3}{-3+n} + \frac{3(d^2-af^2)^2 \left( d+ex+f \sqrt{a + \frac{ex(2d+ex)}{f^2}} \right)^2}{-1+n} - \frac{3(d^2-af^2) \left( d+ex+f \sqrt{a + \frac{ex(2d+ex)}{f^2}} \right)}{1+n} \right)}{8ef^2}$$

input

```
Integrate[(a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2)*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n,x]
```

output

```
((d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^(-3 + n)*(-(d^2 - a*f^2)^3/(-3 + n)) + (3*(d^2 - a*f^2)^2*(d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^2)/(-1 + n) - (3*(d^2 - a*f^2)*(d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^4)/(1 + n) + (d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^6/(3 + n))/(8*e*f^2)
```

### Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 217, normalized size of antiderivative = 0.91, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.074$ , Rules used = {2546, 27, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right) \left( f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex \right)^n dx$$

↓ 2546

$$2 \int \frac{\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+\frac{2dex}{f^2}+a}\right)^{n-4} \left(d^2-af^2-\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+\frac{2dex}{f^2}+a}\right)^2\right)^3}{16ef^2} d\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+\frac{2dex}{f^2}+a}\right)$$

↓ 27

$$\int \frac{\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+\frac{2dex}{f^2}+a}\right)^{n-4} \left(d^2-af^2-\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+\frac{2dex}{f^2}+a}\right)^2\right)^3}{8ef^2} d\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+\frac{2dex}{f^2}+a}\right)$$

↓ 244

$$\int \frac{\left(d^2-af^2\right)^3 \left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+\frac{2dex}{f^2}+a}\right)^{n-4} - 3\left(d^2-af^2\right)^2 \left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+\frac{2dex}{f^2}+a}\right)^{n-2} + 3\left(d^2-af^2\right) \left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+\frac{2dex}{f^2}+a}\right)^{n-2}}{8ef^2} dx$$

↓ 2009

$$\frac{\left(d^2-af^2\right)^3 \left(f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}+d+ex\right)^{n-3}}{3-n} + \frac{3\left(d^2-af^2\right)^2 \left(f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}+d+ex\right)^{n-1}}{1-n} + \frac{3\left(d^2-af^2\right) \left(f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}+d+ex\right)^{n-2}}{n+1}$$

input

```
Int[(a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2)*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n,x]
```

output

```
-1/8*(-(((d^2 - a*f^2)^3*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^(-3 + n))/(3 - n) + (3*(d^2 - a*f^2)^2*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^(-1 + n))/(1 - n) + (3*(d^2 - a*f^2)*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^(1 + n))/(1 + n) - (d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^(3 + n)/(3 + n))/(e*f^2)
```

## Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 244 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2546 `Int[((g_) + (h_)*(x_) + (i_)*(x_)^2)^(m_)*((d_) + (e_)*(x_) + (f_)Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2])^(n_), x_Symbol] := Simp[(2/f^(2*m))*(i/c)^m Subst[Int[x^n*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)^(2*m + 1)/(-2*d*e + b*f^2 + 2*e*x)^(2*(m + 1))), x], x, d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && EqQ[c*h - b*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

## Maple [F]

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right) \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

input `int((a+2*d*e*x/f^2+e^2*x^2/f^2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x)`

output `int((a+2*d*e*x/f^2+e^2*x^2/f^2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x)`



**Fricas [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 239, normalized size of antiderivative = 1.00

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right) \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx =$$

$$\frac{(3adf^2n^2 - 9adf^2 + 3(e^3n^2 - e^3)x^3 + 6d^3 + 9(de^2n^2 - de^2)x^2 - 3(3aef^2 - (aef^2 + 2d^2e)n^2)x - (ef^2n^2 - 3a^2ef^2))}{f^2}$$

input `integrate((a+2*d*e*x/f^2+e^2*x^2/f^2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x, algorithm="fricas")`

output `-(3*a*d*f^2*n^2 - 9*a*d*f^2 + 3*(e^3*n^2 - e^3)*x^3 + 6*d^3 + 9*(d*e^2*n^2 - d*e^2)*x^2 - 3*(3*a*e*f^2 - (a*e*f^2 + 2*d^2*e)*n^2)*x - (a*f^3*n^3 + (e^2*f*n^3 - e^2*f*n)*x^2 - (7*a*f^3 - 6*d^2*f)*n + 2*(d*e*f*n^3 - d*e*f*n)*x)*sqrt((e^2*x^2 + a*f^2 + 2*d*e*x)/f^2)*(e*x + f*sqrt((e^2*x^2 + a*f^2 + 2*d*e*x)/f^2) + d)^n/(e*f^2*n^4 - 10*e*f^2*n^2 + 9*e*f^2)`

**Sympy [F]**

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right) \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \frac{\int a f^2 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx + \int e^2 x^2 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx + \int 2dex \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx}{f^2}$$

input `integrate((a+2*d*e*x/f**2+e**2*x**2/f**2)*(d+e*x+f*(a+2*d*e*x/f**2+e**2*x**2/f**2)**(1/2))**n,x)`

output `(Integral(a*f**2*(d + e*x + f*sqrt(a + 2*d*e*x/f**2 + e**2*x**2/f**2))**n, x) + Integral(e**2*x**2*(d + e*x + f*sqrt(a + 2*d*e*x/f**2 + e**2*x**2/f**2))**n, x) + Integral(2*d*e*x*(d + e*x + f*sqrt(a + 2*d*e*x/f**2 + e**2*x**2/f**2))**n, x))/f**2`

**Maxima [F]**

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right) \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \int \left( \frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2} \right) \left( ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}} f + d \right)^n dx$$

input `integrate((a+2*d*e*x/f^2+e^2*x^2/f^2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x, algorithm="maxima")`

output `integrate((e^2*x^2/f^2 + a + 2*d*e*x/f^2)*(e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2)*f + d)^n, x)`

**Giac [F]**

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right) \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \int \left( \frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2} \right) \left( ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}} f + d \right)^n dx$$

input `integrate((a+2*d*e*x/f^2+e^2*x^2/f^2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x, algorithm="giac")`

output `integrate((e^2*x^2/f^2 + a + 2*d*e*x/f^2)*(e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2)*f + d)^n, x)`

### Mupad [F(-1)]

Timed out.

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right) \left( d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \int \left( d + f\sqrt{a + \frac{e^2x^2}{f^2} + \frac{2dex}{f^2}} + ex \right)^n \left( a + \frac{e^2x^2}{f^2} + \frac{2dex}{f^2} \right) dx$$

input `int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2), x)`

output `int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2), x)`

### Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 324, normalized size of antiderivative = 1.36

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right) \left( d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \frac{(\sqrt{e^2x^2 + af^2 + 2dex} + d + ex)^n (\sqrt{e^2x^2 + af^2 + 2dex} a f^2 n^3 - 7\sqrt{e^2x^2 + af^2 + 2dex} a f^2 n + 6\sqrt{e^2x^2 + af^2 + 2dex} a f^2)}{...}$$

input `int((a+2*d*e*x/f^2+e^2*x^2/f^2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x)`

output `((sqrt(a*f**2 + 2*d*e*x + e**2*x**2) + d + e*x)**n*(sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*a*f**2*n**3 - 7*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*a*f**2*n + 6*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*d**2*n + 2*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*d*e*n**3*x - 2*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*d*e*n*x + sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*e**2*n**3*x**2 - sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*e**2*n*x**2 - 3*a*d*f**2*n**2 + 9*a*d*f**2 - 3*a*e*f**2*n**2*x + 9*a*e*f**2*x - 6*d**3 - 6*d**2*e*n**2*x - 9*d*e**2*n**2*x**2 + 9*d*e**2*x**2 - 3*e**3*n**2*x**3 + 3*e**3*x**3))/(e*f**2*(n**4 - 10*n**2 + 9))`

$$3.40 \quad \int \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

Optimal result	283
Mathematica [A] (verified)	283
Rubi [A] (verified)	284
Maple [F]	285
Fricas [A] (verification not implemented)	286
Sympy [F]	286
Maxima [F]	286
Giac [F]	287
Mupad [F(-1)]	287
Reduce [B] (verification not implemented)	287

### Optimal result

Integrand size = 33, antiderivative size = 107

$$\int \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \frac{(d^2 - af^2) \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{-1+n}}{2e(1-n)} + \frac{\left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{1+n}}{2e(1+n)}$$

output

```
1/2*(-a*f^2+d^2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^(1+n)/e/(1-n)
)+1/2*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^(1+n)/e/(1+n)
```

### Mathematica [A] (verified)

Time = 1.60 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.83

$$\int \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \frac{\left( d + ex + f \sqrt{a + \frac{ex(2d+ex)}{f^2}} \right)^{-1+n} \left( \frac{-d^2+af^2}{-1+n} + \frac{\left( d+ex+f \sqrt{a + \frac{ex(2d+ex)}{f^2}} \right)^2}{1+n} \right)}{2e}$$

input `Integrate[(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n,x]`

output `((d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^(-1 + n)*((-d^2 + a*f^2)/(-1 + n) + (d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^2/(1 + n))/(2*e)`

### Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.97, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.121$ , Rules used = {2541, 27, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex \right)^n dx$$

↓ 2541

$$2 \int -\frac{\left( d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-2} \left( d^2 - af^2 - \left( d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^2 \right)}{4e} d \left( d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)$$

↓ 27

$$\frac{\int \left( d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-2} \left( d^2 - af^2 - \left( d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^2 \right) d \left( d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)}{2e}$$

↓ 244

$$\frac{\int \left( (d^2 - af^2) \left( d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-2} - \left( d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^n \right) d \left( d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)}{2e}$$

↓ 2009

$$\frac{\frac{(d^2 - af^2) \left( f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex \right)^{n-1}}{1-n} - \frac{\left( f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex \right)^{n+1}}{n+1}}{2e}$$

input `Int[(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n,x]`

output `-1/2*(-(((d^2 - a*f^2)*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^(-1 + n))/(1 - n)) - (d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^(1 + n)/(1 + n))/e`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 244 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2541 `Int[((g_) + (h_)*((d_) + (e_)*(x_) + (f_)*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2])^(n_))^(p_), x_Symbol] := Simp[2 Subst[Int[(g + h*x^n)^p*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)/(-2*d*e + b*f^2 + 2*e*x)^2), x], x, d + e*x + f*Sqrt[a + b*x + c*x^2], x] /; FreeQ[{a, b, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]`

### Maple [F]

$$\int \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

input `int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x)`

output `int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x)`

**Fricas [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.75

$$\int \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \frac{\left( fn \sqrt{\frac{e^2x^2 + af^2 + 2dex}{f^2}} - ex - d \right) \left( ex + f \sqrt{\frac{e^2x^2 + af^2 + 2dex}{f^2}} + d \right)^n}{en^2 - e}$$

input `integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x, algorithm="fricas")`

output `(f*n*sqrt((e^2*x^2 + a*f^2 + 2*d*e*x)/f^2) - e*x - d)*(e*x + f*sqrt((e^2*x^2 + a*f^2 + 2*d*e*x)/f^2) + d)^n/(e*n^2 - e)`

**Sympy [F]**

$$\int \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx = \int \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

input `integrate((d+e*x+f*(a+2*d*e*x/f**2+e**2*x**2/f**2)**(1/2))**n,x)`

output `Integral((d + e*x + f*sqrt(a + 2*d*e*x/f**2 + e**2*x**2/f**2))**n, x)`

**Maxima [F]**

$$\int \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx = \int \left( ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}} f + d \right)^n dx$$

input `integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x, algorithm="maxima")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2)*f + d)^n, x)`

### Giac [F]

$$\int \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx = \int \left( ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}} f + d \right)^n dx$$

input `integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x, algorithm="giac")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2)*f + d)^n, x)`

### Mupad [F(-1)]

Timed out.

$$\int \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx = \int \left( d + f \sqrt{a + \frac{e^2x^2}{f^2} + \frac{2dex}{f^2}} + ex \right)^n dx$$

input `int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n,x)`

output `int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n, x)`

### Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.62

$$\begin{aligned} & \int \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx \\ &= \frac{(\sqrt{e^2x^2 + af^2 + 2dex} + d + ex)^n (\sqrt{e^2x^2 + af^2 + 2dex} n - d - ex)}{e(n^2 - 1)} \end{aligned}$$



input `int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x)`

output `((sqrt(a*f**2 + 2*d*e*x + e**2*x**2) + d + e*x)**n*(sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*n - d - e*x))/(e*(n**2 - 1))`

$$3.41 \quad \int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}} dx$$

Optimal result	289
Mathematica [A] (verified)	290
Rubi [A] (verified)	290
Maple [F]	292
Fricas [F]	292
Sympy [F(-1)]	293
Maxima [F]	293
Giac [F]	293
Mupad [F(-1)]	294
Reduce [F]	294

### Optimal result

Integrand size = 56, antiderivative size = 122

$$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}} dx = \frac{2f^2\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^{1+n} \operatorname{Hypergeometric2F1}\left(1, \frac{1+n}{2}, \frac{3+n}{2}, \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^2}{d^2-af^2}\right)}{e(d^2-af^2)(1+n)}$$

output

```
-2*f^2*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^(1+n)*hypergeom([1, 1/2+1/2*n], [3/2+1/2*n], (d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^2/(-a*f^2+d^2))/e/(-a*f^2+d^2)/(1+n)
```

### Mathematica [A] (verified)

Time = 1.96 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.92

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} dx = \frac{2f^2 \left(d + ex + f\sqrt{a + \frac{ex(2d+ex)}{f^2}}\right)^{1+n} \operatorname{Hypergeometric2F1}\left(1, \frac{1+n}{2}, \frac{3+n}{2}, \frac{\left(d+ex+f\sqrt{a+\frac{ex(2d+ex)}{f^2}}\right)^2}{d^2-af^2}\right)}{e(d^2 - af^2)(1+n)}$$

input

```
Integrate[(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n/(a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2),x]
```

output

```
(-2*f^2*(d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^(1 + n)*Hypergeometric2F1[1, (1 + n)/2, (3 + n)/2, (d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^2/(d^2 - a*f^2)]/(e*(d^2 - a*f^2)*(1 + n))
```

### Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$ , Rules used = {2546, 25, 27, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\left(f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex\right)^n}{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} dx$$

↓ 2546

$$2f^2 \int -\frac{\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^n}{e\left(d^2 - af^2 - \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^2\right)} d\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)$$

↓ 25

$$\begin{aligned}
 & -2f^2 \int \frac{\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^n}{e\left(d^2 - af^2 - \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^2\right)} d\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right) \\
 & \quad \downarrow 27 \\
 & \frac{2f^2 \int \frac{\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^n}{d^2 - af^2 - \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^2} d\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)}{e} \\
 & \quad \downarrow 278 \\
 & \frac{2f^2 \left(f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex\right)^{n+1} \operatorname{Hypergeometric2F1}\left(1, \frac{n+1}{2}, \frac{n+3}{2}, \frac{\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^2}{d^2 - af^2}\right)}{e(n+1)(d^2 - af^2)}
 \end{aligned}$$

input `Int[(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n/(a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2), x]`

output `(-2*f^2*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^(1 + n)*Hypergeometric2F1[1, (1 + n)/2, (3 + n)/2, (d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^2/(d^2 - a*f^2)]/(e*(d^2 - a*f^2)*(1 + n))`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 2546

```
Int[((g_.) + (h_.)*(x_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*S
qrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2])^(n_.), x_Symbol] :> Simp[(2/f^(2*m
))*(i/c)^m Subst[Int[x^n*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e
*x^2)^(2*m + 1)/(-2*d*e + b*f^2 + 2*e*x)^(2*(m + 1))), x], x, d + e*x + f*S
qrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, n}, x] && Eq
Q[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && EqQ[c*h - b*i, 0] && IntegerQ[2*m
] && (IntegerQ[m] || GtQ[i/c, 0])
```

**Maple [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} dx$$

input

```
int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f
^2),x)
```

output

```
int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f
^2),x)
```

**Fricas [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} dx = \int \frac{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}}f + d\right)^n}{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}} dx$$

input

```
integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2
*x^2/f^2),x, algorithm="fricas")
```

output

```
integral((e*x + f*sqrt((e^2*x^2 + a*f^2 + 2*d*e*x)/f^2) + d)^n*f^2/(e^2*x^
2 + a*f^2 + 2*d*e*x), x)
```

**Sympy [F(-1)]**

Timed out.

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} dx = \text{Timed out}$$

input `integrate((d+e*x+f*(a+2*d*e*x/f**2+e**2*x**2/f**2)**(1/2))**n/(a+2*d*e*x/f**2+e**2*x**2/f**2),x)`

output `Timed out`

**Maxima [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} dx = \int \frac{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}}f + d\right)^n}{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}} dx$$

input `integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f^2),x, algorithm="maxima")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2)*f + d)^n/(e^2*x^2/f^2 + a + 2*d*e*x/f^2), x)`

**Giac [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} dx = \int \frac{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}}f + d\right)^n}{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}} dx$$

input `integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f^2),x, algorithm="giac")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2))*f + d)^n/(e^2*x^2/f^2 + a + 2*d*e*x/f^2), x)`

### Mupad [F(-1)]

Timed out.

$$\int \frac{\left(d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} dx = \int \frac{\left(d + f \sqrt{a + \frac{e^2x^2}{f^2} + \frac{2dex}{f^2}} + ex\right)^n}{a + \frac{e^2x^2}{f^2} + \frac{2dex}{f^2}} dx$$

input `int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n/(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2), x)`

output `int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n/(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2), x)`

### Reduce [F]

$$\int \frac{\left(d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} dx = \left( \int \frac{\left(\sqrt{e^2x^2 + af^2 + 2dex} + d + ex\right)^n}{e^2x^2 + af^2 + 2dex} dx \right) f^2$$

input `int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f^2), x)`

output `int((sqrt(a*f**2 + 2*d*e*x + e**2*x**2) + d + e*x)**n/(a*f**2 + 2*d*e*x + e**2*x**2), x)*f**2`

$$3.42 \quad \int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\left(a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}\right)^2} dx$$

Optimal result	295
Mathematica [A] (verified)	296
Rubi [A] (verified)	296
Maple [F]	298
Fricas [F]	298
Sympy [F(-1)]	299
Maxima [F]	299
Giac [F]	299
Mupad [F(-1)]	300
Reduce [F]	300

### Optimal result

Integrand size = 56, antiderivative size = 122

$$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\left(a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}\right)^2} dx =$$

$$\frac{8f^4\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^{3+n} \operatorname{Hypergeometric2F1}\left(3, \frac{3+n}{2}, \frac{5+n}{2}, \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^2}{d^2-af^2}\right)}{e(d^2-af^2)^3(3+n)}$$

output

```
-8*f^4*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^(3+n)*hypergeom([3, 3/2+1/2*n], [5/2+1/2*n], (d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^2/(-a*f^2+d^2))/e/(-a*f^2+d^2)^3/(3+n)
```



**Mathematica [A] (verified)**

Time = 2.17 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.92

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^2} dx =$$

$$\frac{8f^4 \left(d + ex + f\sqrt{a + \frac{ex(2d+ex)}{f^2}}\right)^{3+n} \operatorname{Hypergeometric2F1}\left(3, \frac{3+n}{2}, \frac{5+n}{2}, \frac{\left(d+ex+f\sqrt{a+\frac{ex(2d+ex)}{f^2}}\right)^2}{d^2-af^2}\right)}{e(d^2 - af^2)^3(3+n)}$$

input

```
Integrate[(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n/(a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2)^2,x]
```

output

```
(-8*f^4*(d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^(3 + n)*Hypergeometric2F1[3, (3 + n)/2, (5 + n)/2, (d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^2/(d^2 - a*f^2)])/(e*(d^2 - a*f^2)^3*(3 + n))
```

**Rubi [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.054$ , Rules used = {2546, 27, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\left(f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex\right)^n}{\left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^2} dx$$

↓ 2546

$$2f^4 \int -\frac{4\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^{n+2}}{e\left(d^2 - af^2 - \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^2\right)^3} d\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)$$

$$8f^4 \int \frac{\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+\frac{2dex}{f^2}+a}\right)^{n+2}}{\left(d^2-af^2-\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+\frac{2dex}{f^2}+a}\right)^2\right)^3} d\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+\frac{2dex}{f^2}+a}\right)$$

↓ 27

↓ 278

$$\frac{8f^4 \left(f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}+d+ex\right)^{n+3} \operatorname{Hypergeometric2F1}\left(3, \frac{n+3}{2}, \frac{n+5}{2}, \frac{\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2}+\frac{2dex}{f^2}+a}\right)^2}{d^2-af^2}\right)}{e(n+3)(d^2-af^2)^3}$$

input `Int[(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n/(a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2)^2,x]`

output `(-8*f^4*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^(3 + n)*Hypergeometric2F1[3, (3 + n)/2, (5 + n)/2, (d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^2/(d^2 - a*f^2)]/(e*(d^2 - a*f^2)^3*(3 + n))`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 2546

```
Int[((g_.) + (h_.)*(x_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*S
qrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2])^(n_.), x_Symbol] :> Simp[(2/f^(2*m)
)*(i/c)^m Subst[Int[x^n*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e
*x^2)^(2*m + 1)/(-2*d*e + b*f^2 + 2*e*x)^(2*(m + 1))), x], x, d + e*x + f*S
qrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, n}, x] && Eq
Q[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && EqQ[c*h - b*i, 0] && IntegerQ[2*m
] && (IntegerQ[m] || GtQ[i/c, 0])
```

**Maple [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^2} dx$$

input

```
int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f
^2)^2,x)
```

output

```
int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f
^2)^2,x)
```

**Fricas [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^2} dx = \int \frac{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}}f + d\right)^n}{\left(\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}\right)^2} dx$$

input

```
integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2
*x^2/f^2)^2,x, algorithm="fricas")
```

output

```
integral((e*x + f*sqrt((e^2*x^2 + a*f^2 + 2*d*e*x)/f^2) + d)^n*f^4/(e^4*x^
4 + 4*d*e^3*x^3 + a^2*f^4 + 4*a*d*e*f^2*x + 2*(a*e^2*f^2 + 2*d^2*e^2)*x^2)
, x)
```

**Sympy [F(-1)]**

Timed out.

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^2} dx = \text{Timed out}$$

input `integrate((d+e*x+f*(a+2*d*e*x/f**2+e**2*x**2/f**2)**(1/2))**n/(a+2*d*e*x/f**2+e**2*x**2/f**2)**2,x)`

output Timed out

**Maxima [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^2} dx = \int \frac{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}}f + d\right)^n}{\left(\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}\right)^2} dx$$

input `integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f^2)^2,x, algorithm="maxima")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2)*f + d)^n/(e^2*x^2/f^2 + a + 2*d*e*x/f^2)^2, x)`

**Giac [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^2} dx = \int \frac{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}}f + d\right)^n}{\left(\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}\right)^2} dx$$

input `integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f^2)^2,x, algorithm="giac")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2))*f + d)^n/(e^2*x^2/f^2 + a + 2*d*e*x/f^2)^2, x)`

### Mupad [F(-1)]

Timed out.

$$\int \frac{\left(d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^2} dx = \int \frac{\left(d + f \sqrt{a + \frac{e^2x^2}{f^2} + \frac{2dex}{f^2}} + ex\right)^n}{\left(a + \frac{e^2x^2}{f^2} + \frac{2dex}{f^2}\right)^2} dx$$

input `int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n/(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^2,x)`

output `int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n/(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^2, x)`

### Reduce [F]

$$\int \frac{\left(d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^2} dx$$

$$= \left( \int \frac{\left(\sqrt{e^2x^2 + af^2 + 2dex} + d + ex\right)^n}{e^4x^4 + 2ae^2f^2x^2 + 4de^3x^3 + a^2f^4 + 4ade f^2x + 4d^2e^2x^2} dx \right) f^4$$

input `int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f^2)^2,x)`

output `int((sqrt(a*f**2 + 2*d*e*x + e**2*x**2) + d + e*x)**n/(a**2*f**4 + 4*a*d*e*f**2*x + 2*a*e**2*f**2*x**2 + 4*d**2*e**2*x**2 + 4*d*e**3*x**3 + e**4*x**4),x)*f**4`

$$3.43 \quad \int \left( d + ex + f \sqrt{\frac{af^2 + ex(2d + ex)}{f^2}} \right)^n dx$$

Optimal result	301
Mathematica [A] (verified)	301
Rubi [A] (verified)	302
Maple [F]	304
Fricas [A] (verification not implemented)	304
Sympy [F(-1)]	305
Maxima [F]	305
Giac [F]	305
Mupad [F(-1)]	306
Reduce [B] (verification not implemented)	306

**Optimal result**

Integrand size = 33, antiderivative size = 107

$$\int \left( d + ex + f \sqrt{\frac{af^2 + ex(2d + ex)}{f^2}} \right)^n dx$$

$$= \frac{(d^2 - af^2) \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{-1+n}}{2e(1 - n)} + \frac{\left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{1+n}}{2e(1 + n)}$$

```
output 1/2*(-a*f^2+d^2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^(1+n)/e/(1-n)
+1/2*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^(1+n)/e/(1+n)
```

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.83

$$\int \left( d + ex + f \sqrt{\frac{af^2 + ex(2d + ex)}{f^2}} \right)^n dx$$

$$= \frac{\left( d + ex + f \sqrt{a + \frac{ex(2d+ex)}{f^2}} \right)^{-1+n} \left( \frac{-d^2+af^2}{-1+n} + \frac{\left( d+ex+f \sqrt{a + \frac{ex(2d+ex)}{f^2}} \right)^2}{1+n} \right)}{2e}$$

input `Integrate[(d + e*x + f*Sqrt[(a*f^2 + e*x*(2*d + e*x))/f^2])^n,x]`

output `((d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^(-1 + n)*((-d^2 + a*f^2)/(-1 + n) + (d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^2/(1 + n))/(2*e)`

### Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.97, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.152$ , Rules used = {2543, 2541, 27, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( f \sqrt{\frac{af^2 + ex(2d + ex)}{f^2}} + d + ex \right)^n dx$$

↓ 2543

$$\int \left( f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex \right)^n dx$$

↓ 2541

$$2 \int -\frac{\left( d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-2} \left( d^2 - af^2 - \left( d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^2 \right)}{4e} d \left( d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)$$

↓ 27

$$\frac{\int \left( d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-2} \left( d^2 - af^2 - \left( d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^2 \right) d \left( d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)}{2e}$$

↓ 244

$$\frac{\int \left( (d^2 - af^2) \left( d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-2} - \left( d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^n \right) d \left( d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)}{2e}$$

$$\frac{(d^2 - af^2) \left( f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex \right)^{n-1}}{1-n} - \frac{\left( f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex \right)^{n+1}}{n+1}$$

↓ 2009

$$2e$$

input `Int[(d + e*x + f*Sqrt[(a*f^2 + e*x*(2*d + e*x))/f^2])^n,x]`

output `-1/2*(-(((d^2 - a*f^2)*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^(-1 + n))/(1 - n)) - (d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^(1 + n)/(1 + n))/e`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 244 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2541 `Int[((g_) + (h_)*((d_) + (e_)*(x_) + (f_)*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2])^(n_))^(p_), x_Symbol] := Simp[2 Subst[Int[(g + h*x^n)^p*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)/(-2*d*e + b*f^2 + 2*e*x)^2], x], x, d + e*x + f*Sqrt[a + b*x + c*x^2], x] /; FreeQ[{a, b, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]`



rule 2543

```
Int[((g_.) + (h_.)*((u_) + (f_.)*Sqrt[v_])^(n_))^(p_.), x_Symbol] := Int[(g
+ h*(ExpandToSum[u, x] + f*Sqrt[ExpandToSum[v, x]])^n)^p, x] /; FreeQ[{f,
g, h, n}, x] && LinearQ[u, x] && QuadraticQ[v, x] && !(LinearMatchQ[u, x]
&& QuadraticMatchQ[v, x]) && EqQ[Coefficient[u, x, 1]^2 - Coefficient[v, x,
2]*f^2, 0] && IntegerQ[p]
```

**Maple [F]**

$$\int \left( d + ex + f \sqrt{\frac{f^2 a + ex(ex + 2d)}{f^2}} \right)^n dx$$

input

```
int((d+e*x+f*((f^2*a+e*x*(e*x+2*d))/f^2)^(1/2))^n,x)
```

output

```
int((d+e*x+f*((f^2*a+e*x*(e*x+2*d))/f^2)^(1/2))^n,x)
```

**Fricas [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.75

$$\int \left( d + ex + f \sqrt{\frac{af^2 + ex(2d + ex)}{f^2}} \right)^n dx$$

$$= \frac{\left( fn \sqrt{\frac{e^2 x^2 + af^2 + 2dex}{f^2}} - ex - d \right) \left( ex + f \sqrt{\frac{e^2 x^2 + af^2 + 2dex}{f^2}} + d \right)^n}{en^2 - e}$$

input

```
integrate((d+e*x+f*((a*f^2+e*x*(e*x+2*d))/f^2)^(1/2))^n,x, algorithm="fric
as")
```

output

```
(f*n*sqrt((e^2*x^2 + a*f^2 + 2*d*e*x)/f^2) - e*x - d)*(e*x + f*sqrt((e^2*x
^2 + a*f^2 + 2*d*e*x)/f^2) + d)^n/(e*n^2 - e)
```

**Sympy [F(-1)]**

Timed out.

$$\int \left( d + ex + f \sqrt{\frac{af^2 + ex(2d + ex)}{f^2}} \right)^n dx = \text{Timed out}$$

input `integrate((d+e*x+f*((a*f**2+e*x*(e*x+2*d))/f**2)**(1/2))**n,x)`

output `Timed out`

**Maxima [F]**

$$\begin{aligned} & \int \left( d + ex + f \sqrt{\frac{af^2 + ex(2d + ex)}{f^2}} \right)^n dx \\ &= \int \left( ex + f \left( \frac{\sqrt{af^2 + (ex + 2d)ex}}{f} \right) + d \right)^n dx \end{aligned}$$

input `integrate((d+e*x+f*((a*f^2+e*x*(e*x+2*d))/f^2)^(1/2))^n,x, algorithm="maxima")`

output `integrate((e*x + d + sqrt(a*f^2 + (e*x + 2*d)*e*x))^n, x)`

**Giac [F]**

$$\int \left( d + ex + f \sqrt{\frac{af^2 + ex(2d + ex)}{f^2}} \right)^n dx = \int \left( ex + f \sqrt{\frac{af^2 + (ex + 2d)ex}{f^2}} + d \right)^n dx$$

input `integrate((d+e*x+f*((a*f^2+e*x*(e*x+2*d))/f^2)^(1/2))^n,x, algorithm="giac")`

output `integrate((e*x + f*sqrt((a*f^2 + (e*x + 2*d)*e*x)/f^2) + d)^n, x)`

### Mupad [F(-1)]

Timed out.

$$\int \left( d + ex + f \sqrt{\frac{af^2 + ex(2d + ex)}{f^2}} \right)^n dx$$

$$= \int \left( d + ex + f \sqrt{\frac{af^2 + ex(2d + ex)}{f^2}} \right)^n dx$$

input `int((d + e*x + f*((a*f^2 + e*x*(2*d + e*x))/f^2)^(1/2))^n,x)`

output `int((d + e*x + f*((a*f^2 + e*x*(2*d + e*x))/f^2)^(1/2))^n, x)`

### Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.62

$$\int \left( d + ex + f \sqrt{\frac{af^2 + ex(2d + ex)}{f^2}} \right)^n dx$$

$$= \frac{(\sqrt{e^2x^2 + af^2 + 2dex} + d + ex)^n (\sqrt{e^2x^2 + af^2 + 2dex} n - d - ex)}{e(n^2 - 1)}$$

input `int((d+e*x+f*((a*f^2+e*x*(e*x+2*d))/f^2)^(1/2))^n,x)`

output `((sqrt(a*f**2 + 2*d*e*x + e**2*x**2) + d + e*x)**n*(sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*n - d - e*x))/(e*(n**2 - 1))`

$$3.44 \quad \int \frac{\left(d+ex+f\sqrt{\frac{af^2+ex(2d+ex)}{f^2}}\right)^n}{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}} dx$$

Optimal result	307
Mathematica [A] (verified)	308
Rubi [A] (verified)	308
Maple [F]	310
Fricas [F]	311
Sympy [F(-1)]	311
Maxima [F]	311
Giac [F]	312
Mupad [F(-1)]	312
Reduce [F]	313

### Optimal result

Integrand size = 56, antiderivative size = 122

$$\int \frac{\left(d+ex+f\sqrt{\frac{af^2+ex(2d+ex)}{f^2}}\right)^n}{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}} dx = \frac{2f^2\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^{1+n} \operatorname{Hypergeometric2F1}\left(1, \frac{1+n}{2}, \frac{3+n}{2}, \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^2}{d^2-af^2}\right)}{e(d^2-af^2)(1+n)}$$

output

```
-2*f^2*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^(1+n)*hypergeom([1, 1/2+1/2*n], [3/2+1/2*n], (d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^2/(-a*f^2+d^2))/e/(-a*f^2+d^2)/(1+n)
```

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.92

$$\int \frac{\left(d + ex + f\sqrt{\frac{af^2 + ex(2d+ex)}{f^2}}\right)^n}{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} dx =$$

$$\frac{2f^2 \left(d + ex + f\sqrt{a + \frac{ex(2d+ex)}{f^2}}\right)^{1+n} \operatorname{Hypergeometric2F1}\left(1, \frac{1+n}{2}, \frac{3+n}{2}, \frac{\left(d+ex+f\sqrt{a+\frac{ex(2d+ex)}{f^2}}\right)^2}{d^2-af^2}\right)}{e(d^2 - af^2)(1+n)}$$

input

```
Integrate[(d + e*x + f*Sqrt[(a*f^2 + e*x*(2*d + e*x))/f^2])^n/(a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2),x]
```

output

```
(-2*f^2*(d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^(1 + n)*Hypergeometric2F1[1, (1 + n)/2, (3 + n)/2, (d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^2/(d^2 - a*f^2)]/(e*(d^2 - a*f^2)*(1 + n))
```

**Rubi [A] (verified)**

Time = 0.56 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.089$ , Rules used = {2552, 2546, 25, 27, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\left(f\sqrt{\frac{af^2 + ex(2d+ex)}{f^2}} + d + ex\right)^n}{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} dx$$

↓ 2552

$$\int \frac{\left(f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex\right)^n}{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} dx$$

↓ 2546

$$2f^2 \int -\frac{\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^n}{e\left(d^2 - af^2 - \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^2\right)} d\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)$$

↓ 25

$$-2f^2 \int \frac{\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^n}{e\left(d^2 - af^2 - \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^2\right)} d\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)$$

↓ 27

$$2f^2 \int \frac{\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^n}{d^2 - af^2 - \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^2} d\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)$$

e

↓ 278

$$\frac{2f^2 \left(f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex\right)^{n+1} \text{Hypergeometric2F1}\left(1, \frac{n+1}{2}, \frac{n+3}{2}, \frac{\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^2}{d^2 - af^2}\right)}{e(n+1)(d^2 - af^2)}$$

input

```
Int[(d + e*x + f*Sqrt[(a*f^2 + e*x*(2*d + e*x))/f^2])^n/(a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2), x]
```

output

```
(-2*f^2*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^(1 + n)*Hypergeometric2F1[1, (1 + n)/2, (3 + n)/2, (d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^2/(d^2 - a*f^2)]/(e*(d^2 - a*f^2)*(1 + n))
```

### Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

rule 278

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((
c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (
-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0
] || GtQ[a, 0])
```

rule 2546

```
Int[((g_.) + (h_.)*(x_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*S
qrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(2/f^(2*m)
)*(i/c)^m Subst[Int[x^n*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e
*x^2)^(2*m + 1)/(-2*d*e + b*f^2 + 2*e*x)^(2*(m + 1))), x], x, d + e*x + f*S
qrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, n}, x] && Eq
Q[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && EqQ[c*h - b*i, 0] && IntegerQ[2*m
] && (IntegerQ[m] || GtQ[i/c, 0])
```

rule 2552

```
Int[((u_) + (f_.)*((j_.) + (k_.)*Sqrt[v_]))^(n_.)*(w_)^(m_.), x_Symbol] :=
Int[ExpandToSum[w, x]^m*(ExpandToSum[u + f*j, x] + f*k*Sqrt[ExpandToSum[v,
x]])^n, x] /; FreeQ[{f, j, k, m, n}, x] && LinearQ[u, x] && QuadraticQ[{v,
w}, x] && !(LinearMatchQ[u, x] && QuadraticMatchQ[{v, w}, x] && (EqQ[j, 0]
|| EqQ[f, 1])) && EqQ[Coefficient[u, x, 1]^2 - Coefficient[v, x, 2]*f^2*k^
2, 0]
```

## Maple [F]

$$\int \frac{\left(d + ex + f \sqrt{\frac{f^2 a + ex(ex + 2d)}{f^2}}\right)^n}{a + \frac{2dex}{f^2} + \frac{e^2 x^2}{f^2}} dx$$

input

```
int((d+e*x+f*((f^2*a+e*x*(e*x+2*d))/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f
^2),x)
```

output

```
int((d+e*x+f*((f^2*a+e*x*(e*x+2*d))/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f
^2),x)
```

**Fricas [F]**

$$\int \frac{\left(d + ex + f \sqrt{\frac{af^2 + ex(2d+ex)}{f^2}}\right)^n}{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} dx = \int \frac{\left(ex + f \sqrt{\frac{af^2 + (ex+2d)ex}{f^2}} + d\right)^n}{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}} dx$$

input `integrate((d+e*x+f*((a*f^2+e*x*(e*x+2*d))/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f^2),x, algorithm="fricas")`

output `integral((e*x + f*sqrt((e^2*x^2 + a*f^2 + 2*d*e*x)/f^2) + d)^n*f^2/(e^2*x^2 + a*f^2 + 2*d*e*x), x)`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{\left(d + ex + f \sqrt{\frac{af^2 + ex(2d+ex)}{f^2}}\right)^n}{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} dx = \text{Timed out}$$

input `integrate((d+e*x+f*((a*f**2+e*x*(e*x+2*d))/f**2)**(1/2))**n/(a+2*d*e*x/f**2+e**2*x**2/f**2),x)`

output `Timed out`

**Maxima [F]**

$$\int \frac{\left(d + ex + f \sqrt{\frac{af^2 + ex(2d+ex)}{f^2}}\right)^n}{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} dx = \int \frac{\left(ex + d + \sqrt{af^2 + (ex + 2d)ex}\right)^n}{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}} dx$$

input `integrate((d+e*x+f*((a*f^2+e*x*(e*x+2*d))/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f^2),x, algorithm="maxima")`



output

```
integrate((e*x + d + sqrt(a*f^2 + (e*x + 2*d)*e*x))^n/(e^2*x^2/f^2 + a + 2*d*e*x/f^2), x)
```

**Giac [F]**

$$\int \frac{\left(d + ex + f \sqrt{\frac{af^2 + ex(2d + ex)}{f^2}}\right)^n}{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} dx = \int \frac{\left(ex + f \sqrt{\frac{af^2 + (ex + 2d)ex}{f^2}} + d\right)^n}{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}} dx$$

input

```
integrate((d+e*x+f*((a*f^2+e*x*(e*x+2*d))/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f^2),x, algorithm="giac")
```

output

```
integrate((e*x + f*sqrt((a*f^2 + (e*x + 2*d)*e*x)/f^2) + d)^n/(e^2*x^2/f^2 + a + 2*d*e*x/f^2), x)
```

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\left(d + ex + f \sqrt{\frac{af^2 + ex(2d + ex)}{f^2}}\right)^n}{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} dx = \int \frac{\left(d + ex + f \sqrt{\frac{af^2 + ex(2d + ex)}{f^2}}\right)^n}{a + \frac{e^2x^2}{f^2} + \frac{2dex}{f^2}} dx$$

input

```
int((d + e*x + f*((a*f^2 + e*x*(2*d + e*x))/f^2)^(1/2))^n/(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2), x)
```

output

```
int((d + e*x + f*((a*f^2 + e*x*(2*d + e*x))/f^2)^(1/2))^n/(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2), x)
```

**Reduce [F]**

$$\int \frac{\left(d + ex + f \sqrt{\frac{af^2 + ex(2d + ex)}{f^2}}\right)^n}{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} dx = \left( \int \frac{(\sqrt{e^2x^2 + af^2 + 2dex} + d + ex)^n}{e^2x^2 + af^2 + 2dex} dx \right) f^2$$

input

```
int((d+e*x+f*((a*f^2+e*x*(e*x+2*d))/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f^2),x)
```

output

```
int((sqrt(a*f**2 + 2*d*e*x + e**2*x**2) + d + e*x)**n/(a*f**2 + 2*d*e*x + e**2*x**2),x)*f**2
```

**3.45**  $\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^{3/2} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)$

Optimal result	314
Mathematica [A] (verified)	315
Rubi [A] (verified)	316
Maple [F]	317
Fricas [A] (verification not implemented)	318
Sympy [F(-2)]	318
Maxima [F]	319
Giac [F]	319
Mupad [F(-1)]	320
Reduce [B] (verification not implemented)	320

**Optimal result**

Integrand size = 58, antiderivative size = 297

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^{3/2} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx =$$

$$\frac{(d^2 - af^2)^4 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{-4+n}}{16ef^3(4 - n)}$$

$$+ \frac{(d^2 - af^2)^3 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{-2+n}}{4ef^3(2 - n)}$$

$$+ \frac{3(d^2 - af^2)^2 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n}{8ef^3n}$$

$$- \frac{(d^2 - af^2) \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{2+n}}{4ef^3(2 + n)}$$

$$+ \frac{\left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{4+n}}{16ef^3(4 + n)}$$

output

$$\begin{aligned}
& -1/16*(-a*f^2+d^2)^4*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^{(-4+n)}/e/ \\
& f^3/(4-n)+1/4*(-a*f^2+d^2)^3*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^{(- \\
& -2+n)}/e/f^3/(2-n)+3/8*(-a*f^2+d^2)^2*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^( \\
& (1/2))^{n}/e/f^3/n-1/4*(-a*f^2+d^2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/ \\
& 2))^{(2+n)}/e/f^3/(2+n)+1/16*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^{(4+ \\
& n)}/e/f^3/(4+n)
\end{aligned}$$

**Mathematica [A] (verified)**

Time = 8.81 (sec) , antiderivative size = 228, normalized size of antiderivative = 0.77

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^{3/2} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx = \frac{\left( d + ex + f \sqrt{a + \frac{ex(2d+ex)}{f^2}} \right)^n \left( \frac{6(d^2-af^2)^2}{n} + \frac{(d^2-af^2)^4}{(-4+n) \left( d+ex+f \sqrt{a + \frac{ex(2d+ex)}{f^2}} \right)^4} \right)}{16e^3}$$

input

```
Integrate[(a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2)^(3/2)*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n,x]
```

output

```
((d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^n*((6*(d^2 - a*f^2)^2)/n + (d^2 - a*f^2)^4/((-4 + n)*(d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^4) - (4*(d^2 - a*f^2)^3)/((-2 + n)*(d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^2) - (4*(d^2 - a*f^2)*(d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^2)/(2 + n) + (d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^4/(4 + n))/(16*e*f^3)
```

**Rubi [A] (verified)**

Time = 0.50 (sec) , antiderivative size = 266, normalized size of antiderivative = 0.90, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.069$ , Rules used = {2546, 27, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^{3/2} \left( f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex \right)^n dx$$

↓ 2546

$$2 \int \frac{\left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-5} \left( d^2-af^2 - \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^2 \right)^4}{32e} d \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)}{f^3}$$

↓ 27

$$\int \frac{\left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-5} \left( d^2-af^2 - \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^2 \right)^4}{16ef^3} d \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)}{16ef^3}$$

↓ 244

$$\int \frac{\left( d^2-af^2 \right)^4 \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-5} - 4 \left( d^2-af^2 \right)^3 \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-3} + 6 \left( d^2-af^2 \right)^2 \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-1}}{16ef^3}$$

↓ 2009

$$\frac{\left( d^2-af^2 \right)^4 \left( f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d+ex \right)^{n-4}}{4-n} + \frac{4 \left( d^2-af^2 \right)^3 \left( f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d+ex \right)^{n-2}}{2-n} + \frac{6 \left( d^2-af^2 \right)^2 \left( f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d+ex \right)^n}{16ef^3}$$

input

```
Int[(a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2)^(3/2)*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n,x]
```

output

```
(-(((d^2 - a*f^2)^4*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^
(-4 + n))/(4 - n)) + (4*(d^2 - a*f^2)^3*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^
2 + (e^2*x^2)/f^2])^(-2 + n))/(2 - n) + (6*(d^2 - a*f^2)^2*(d + e*x + f*Sq
rt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n)/n - (4*(d^2 - a*f^2)*(d + e*x +
f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^(2 + n))/(2 + n) + (d + e*x + f
*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^(4 + n)/(4 + n))/(16*e*f^3)
```

### Defintions of rubi rules used

rule 27

```
Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]
```

rule 244

```
Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[Expand
Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p
, 0]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2546

```
Int[((g_) + (h_)*(x_) + (i_)*(x_)^2)^(m_)*((d_) + (e_)*(x_) + (f_) *S
qrt[(a_) + (b_)*(x_) + (c_)*(x_)^2])^(n_), x_Symbol] := Simp[(2/f^(2*m)
)*(i/c)^m Subst[Int[x^n*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e
*x^2)^(2*m + 1)/(-2*d*e + b*f^2 + 2*e*x)^(2*(m + 1))), x], x, d + e*x + f*S
qrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, n}, x] && Eq
Q[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && EqQ[c*h - b*i, 0] && IntegerQ[2*m
] && (IntegerQ[m] || GtQ[i/c, 0])
```

### Maple [F]

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^{\frac{3}{2}} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

input

```
int((a+2*d*e*x/f^2+e^2*x^2/f^2)^(3/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)
^(1/2))^n,x)
```

output

```
int((a+2*d*e*x/f^2+e^2*x^2/f^2)^(3/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x)
```

**Fricas [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 377, normalized size of antiderivative = 1.27

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^{3/2} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx = \frac{\left( a^2 f^4 n^4 + 24 a^2 f^4 - 48 a d^2 f^2 + (e^4 n^4 - 4 e^4 n^2) x^4 + 24 d^4 + 4 (d e^3 n^4 - 4 d e^3 n^2) x^2 + 4 d^2 e^3 n^2 x \right) \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}}{(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2})^{3/2}}$$

input

```
integrate((a+2*d*e*x/f^2+e^2*x^2/f^2)^(3/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x, algorithm="fricas")
```

output

```
(a^2*f^4*n^4 + 24*a^2*f^4 - 48*a*d^2*f^2 + (e^4*n^4 - 4*e^4*n^2)*x^4 + 24*d^4 + 4*(d*e^3*n^4 - 4*d*e^3*n^2)*x^3 - 4*(4*a^2*f^4 - 3*a*d^2*f^2)*n^2 + 2*((a*e^2*f^2 + 2*d^2*e^2)*n^4 - 2*(5*a*e^2*f^2 + d^2*e^2)*n^2)*x^2 + 4*(a*d*e*f^2*n^4 - 2*(5*a*d*e*f^2 - 3*d^3*e)*n^2)*x - 4*(a*d*f^3*n^3 + (e^3*f*n^3 - 4*e^3*f*n)*x^3 + 3*(d*e^2*f*n^3 - 4*d*e^2*f*n)*x^2 - 2*(5*a*d*f^3 - 3*d^3*f)*n + ((a*e*f^3 + 2*d^2*e*f)*n^3 - 2*(5*a*e*f^3 + d^2*e*f)*n)*x)*sqrt((e^2*x^2 + a*f^2 + 2*d*e*x)/f^2)*(e*x + f*sqrt((e^2*x^2 + a*f^2 + 2*d*e*x)/f^2) + d)^n/(e*f^3*n^5 - 20*e*f^3*n^3 + 64*e*f^3*n)
```

**Sympy [F(-2)]**

Exception generated.

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^{3/2} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx = \text{Exception raised: HeuristicGCDFailed}$$

input `integrate((a+2*d*e*x/f**2+e**2*x**2/f**2)**(3/2)*(d+e*x+f*(a+2*d*e*x/f**2+e**2*x**2/f**2)**(1/2))**n,x)`

output Exception raised: HeuristicGCDFailed >> no luck

### Maxima [F]

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^{3/2} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx = \int \left( \frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2} \right)^{\frac{3}{2}} \left( ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}} f + d \right)^n dx$$

input `integrate((a+2*d*e*x/f^2+e^2*x^2/f^2)^(3/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x, algorithm="maxima")`

output `integrate((e^2*x^2/f^2 + a + 2*d*e*x/f^2)^(3/2)*(e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2)*f + d)^n, x)`

### Giac [F]

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^{3/2} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx = \int \left( \frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2} \right)^{\frac{3}{2}} \left( ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}} f + d \right)^n dx$$

input `integrate((a+2*d*e*x/f^2+e^2*x^2/f^2)^(3/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x, algorithm="giac")`

output `integrate((e^2*x^2/f^2 + a + 2*d*e*x/f^2)^(3/2)*(e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2)*f + d)^n, x)`



**Mupad [F(-1)]**

Timed out.

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^{3/2} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx = \int \left( d + f \sqrt{a + \frac{e^2x^2}{f^2} + \frac{2dex}{f^2}} + ex \right)^n \left( a + \frac{e^2x^2}{f^2} + \frac{2dex}{f^2} \right)^{3/2} dx$$

input

```
int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(3/2), x)
```

output

```
int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(3/2), x)
```

**Reduce [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 554, normalized size of antiderivative = 1.87

$$\int \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)^{3/2} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx = \frac{(\sqrt{e^2x^2 + af^2 + 2dex} + d + ex)^n (a^2f^4n^4 + e^4n^4x^4 - 4\sqrt{e^2x^2 + af^2 + 2dex})}{2(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2})^{3/2}}$$

input

```
int((a+2*d*e*x/f^2+e^2*x^2/f^2)^(3/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x)
```

output

```

((sqrt(a*f**2 + 2*d*e*x + e**2*x**2) + d + e*x)**n*( - 4*sqrt(a*f**2 + 2*d
*e*x + e**2*x**2)*a*d*f**2*n**3 + 40*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*a
*d*f**2*n - 4*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*a*e*f**2*n**3*x + 40*sqrt(
a*f**2 + 2*d*e*x + e**2*x**2)*a*e*f**2*n*x - 24*sqrt(a*f**2 + 2*d*e*x + e
**2*x**2)*d**3*n - 8*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*d**2*e*n**3*x + 8*s
qrt(a*f**2 + 2*d*e*x + e**2*x**2)*d**2*e*n*x - 12*sqrt(a*f**2 + 2*d*e*x +
e**2*x**2)*d*e**2*n**3*x**2 + 48*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*d*e**2
*n*x**2 - 4*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*e**3*n**3*x**3 + 16*sqrt(a*
f**2 + 2*d*e*x + e**2*x**2)*e**3*n*x**3 + a**2*f**4*n**4 - 16*a**2*f**4*n*
*2 + 24*a**2*f**4 + 12*a*d**2*f**2*n**2 - 48*a*d**2*f**2 + 4*a*d*e*f**2*n*
*4*x - 40*a*d*e*f**2*n**2*x + 2*a*e**2*f**2*n**4*x**2 - 20*a*e**2*f**2*n**
2*x**2 + 24*d**4 + 24*d**3*e*n**2*x + 4*d**2*e**2*n**4*x**2 - 4*d**2*e**2*
n**2*x**2 + 4*d*e**3*n**4*x**3 - 16*d*e**3*n**2*x**3 + e**4*n**4*x**4 - 4*
e**4*n**2*x**4))/(e*f**3*n*(n**4 - 20*n**2 + 64))

```

**3.46**  $\int \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$

Optimal result	322
Mathematica [A] (verified)	323
Rubi [A] (verified)	323
Maple [F]	325
Fricas [A] (verification not implemented)	325
Sympy [F(-2)]	326
Maxima [F]	326
Giac [F]	327
Mupad [F(-1)]	327
Reduce [B] (verification not implemented)	328

**Optimal result**

Integrand size = 58, antiderivative size = 171

$$\int \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= -\frac{(d^2 - af^2)^2 \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{-2+n}}{4ef(2 - n)}$$

$$- \frac{(d^2 - af^2) \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n}{2efn} + \frac{\left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{2+n}}{4ef(2 + n)}$$

output

```
-1/4*(-a*f^2+d^2)^2*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^(2-n)/e/f
/(2-n)-1/2*(-a*f^2+d^2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/e/f/
n+1/4*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^(2+n)/e/f/(2+n)
```

### Mathematica [A] (verified)

Time = 3.62 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.79

$$\int \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \frac{\left( d + ex + f \sqrt{a + \frac{ex(2d+ex)}{f^2}} \right)^n \left( \frac{2(-d^2+af^2)}{n} + \frac{(d^2-af^2)^2}{(-2+n) \left( d+ex+f \sqrt{a + \frac{ex(2d+ex)}{f^2}} \right)^2} + \frac{\left( d+ex+f \sqrt{a + \frac{ex(2d+ex)}{f^2}} \right)^2}{2+n} \right)}{4ef}$$

input

```
Integrate[Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2]*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n,x]
```

output

```
((d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^n*((2*(-d^2 + a*f^2))/n + (d^2 - a*f^2)^2/((-2 + n)*(d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^2) + (d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^2/(2 + n)))/(4*e*f)
```

### Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.91, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.069$ , Rules used = {2546, 27, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \left( f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex \right)^n dx$$

↓ 2546

$$2 \int \frac{\left( d+ex+f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-3} \left( d^2-af^2 - \left( d+ex+f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^2 \right)^2}{8e} d \left( d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)$$

f

↓ 27

$$\frac{\int \left( d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-3} \left( d^2 - af^2 - \left( d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^2 \right)^2 d \left( d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + \frac{2dex}{f^2} + a} \right)}{4ef}$$

↓ 244

$$\frac{\int \left( (d^2 - af^2)^2 \left( d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-3} - 2(d^2 - af^2) \left( d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n-1} + \left( d + ex + f \sqrt{\frac{e^2 x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n+1} \right)}{4ef}$$

↓ 2009

$$\frac{-\frac{(d^2 - af^2)^2 \left( f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2 x^2}{f^2}} + d + ex \right)^{n-2}}{2-n} - \frac{2(d^2 - af^2) \left( f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2 x^2}{f^2}} + d + ex \right)^n}{n} + \frac{\left( f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2 x^2}{f^2}} + d + ex \right)^{n+2}}{n+2}}{4ef}$$

input

```
Int[Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2]*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n,x]
```

output

```
(-(((d^2 - a*f^2)^2*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^(-2 + n))/(2 - n) - (2*(d^2 - a*f^2)*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n)/n + (d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^(2 + n)/(2 + n))/(4*e*f)
```

### Defintions of rubi rules used

rule 27

```
Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]
```

rule 244

```
Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2546

```
Int[((g_.) + (h_.)*(x_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*S
qrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(2/f^(2*m)
)*(i/c)^m Subst[Int[x^n*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e
*x^2)^(2*m + 1)/(-2*d*e + b*f^2 + 2*e*x)^(2*(m + 1))), x], x, d + e*x + f*S
qrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, n}, x] && Eq
Q[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && EqQ[c*h - b*i, 0] && IntegerQ[2*m
] && (IntegerQ[m] || GtQ[i/c, 0])
```

**Maple [F]**

$$\int \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

input

```
int((a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)
^(1/2))^n,x)
```

output

```
int((a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)
^(1/2))^n,x)
```

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.71

$$\int \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \frac{\left( e^2n^2x^2 + af^2n^2 + 2den^2x - 2af^2 + 2d^2 - 2(efnx + dfn) \sqrt{\frac{e^2x^2 + af^2 + 2dex}{f^2}} \right) \left( ex + f \sqrt{\frac{e^2x^2 + af^2 + 2dex}{f^2}} + \right)}{efn^3 - 4efn}$$

input

```
integrate((a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)
^(1/2))^n,x, algorithm="fricas")
```

output

```
(e^2*n^2*x^2 + a*f^2*n^2 + 2*d*e*n^2*x - 2*a*f^2 + 2*d^2 - 2*(e*f*n*x + d
*f*n)*sqrt((e^2*x^2 + a*f^2 + 2*d*e*x)/f^2))*(e*x + f*sqrt((e^2*x^2 + a*f^2
+ 2*d*e*x)/f^2) + d)^n/(e*f*n^3 - 4*e*f*n)
```

**Sympy [F(-2)]**

Exception generated.

$$\int \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

= Exception raised: HeuristicGCDFailed

input

```
integrate((a+2*d*e*x/f**2+e**2*x**2/f**2)**(1/2)*(d+e*x+f*(a+2*d*e*x/f**2+
e**2*x**2/f**2)**(1/2))**n,x)
```

output

```
Exception raised: HeuristicGCDFailed >> no luck
```

**Maxima [F]**

$$\int \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \int \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}} \left( ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}} f + d \right)^n dx$$

input

```
integrate((a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^
2/f^2)^(1/2))^n,x, algorithm="maxima")
```

output

```
integrate(sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2)*(e*x + sqrt(e^2*x^2/f^2 + a
+ 2*d*e*x/f^2)*f + d)^n, x)
```

**Giac [F]**

$$\int \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \int \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}} \left( ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}} f + d \right)^n dx$$

input

```
integrate((a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x, algorithm="giac")
```

output

```
integrate(sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2)*(e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2)*f + d)^n, x)
```

**Mupad [F(-1)]**

Timed out.

$$\int \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \int \left( d + f \sqrt{a + \frac{e^2x^2}{f^2} + \frac{2dex}{f^2}} + ex \right)^n \sqrt{a + \frac{e^2x^2}{f^2} + \frac{2dex}{f^2}} dx$$

input

```
int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2),x)
```

output

```
int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2), x)
```



**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.75

$$\int \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \frac{(\sqrt{e^2x^2 + af^2 + 2dex} + d + ex)^n (-2\sqrt{e^2x^2 + af^2 + 2dex} dn - 2\sqrt{e^2x^2 + af^2 + 2dex} enx + af^2n^2 - efn(n^2 - 4))}{efn(n^2 - 4)}$$

input

```
int((a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x)
```

output

```
((sqrt(a*f**2 + 2*d*e*x + e**2*x**2) + d + e*x)**n*(- 2*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*d*n - 2*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*e*n*x + a*f**2*n**2 - 2*a*f**2 + 2*d**2 + 2*d*e*n**2*x + e**2*n**2*x**2))/(e*f*n*(n**2 - 4))
```

$$3.47 \quad \int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}} dx$$

Optimal result	329
Mathematica [A] (verified)	329
Rubi [A] (verified)	330
Maple [F]	331
Fricas [A] (verification not implemented)	331
Sympy [F]	332
Maxima [F]	332
Giac [F]	333
Mupad [B] (verification not implemented)	333
Reduce [B] (verification not implemented)	333

### Optimal result

Integrand size = 58, antiderivative size = 41

$$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}} dx = \frac{f\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{en}$$

output `f*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/e/n`

### Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.88

$$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}} dx = \frac{f\left(d+ex+f\sqrt{a+\frac{ex(2d+ex)}{f^2}}\right)^n}{en}$$

input `Integrate[(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n/Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2], x]`

output  $(f*(d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^n)/(e*n)$

### Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.034$ , Rules used = {2546, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\left(f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex\right)^n}{\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}} dx$$

↓ 2546

$$2f \int \frac{\left(d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^{n-1}}{2e} d\left(d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)$$

↓ 15

$$\frac{f \left(f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex\right)^n}{en}$$

input  $\text{Int}[(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n/Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2],x]$

output  $(f*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n)/(e*n)$

## Definitions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2546 `Int[((g_.) + (h_.)*(x_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(2/f^(2*m))* (i/c)^m Subst[Int[x^n*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)^(2*m + 1)/(-2*d*e + b*f^2 + 2*e*x)^(2*(m + 1))), x], x, d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && EqQ[c*h - b*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

## Maple [F]

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}} dx$$

input `int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2),x)`

output `int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2),x)`

## Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}} dx = \frac{\left(ex + f\sqrt{\frac{e^2x^2 + af^2 + 2dex}{f^2}} + d\right)^n f}{en}$$

input `integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2),x, algorithm="fricas")`

output  $(e*x + f*\text{sqrt}((e^2*x^2 + a*f^2 + 2*d*e*x)/f^2) + d)^n*f/(e*n)$

### Sympy [F]

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}} dx = \int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}} dx$$

input `integrate((d+e*x+f*(a+2*d*e*x/f**2+e**2*x**2/f**2)**(1/2))**n/(a+2*d*e*x/f**2+e**2*x**2/f**2)**(1/2),x)`

output `Integral((d + e*x + f*sqrt(a + 2*d*e*x/f**2 + e**2*x**2/f**2))**n/sqrt(a + 2*d*e*x/f**2 + e**2*x**2/f**2), x)`

### Maxima [F]

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}} dx = \int \frac{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}}f + d\right)^n}{\sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}}} dx$$

input `integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2),x, algorithm="maxima")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2)*f + d)^n/sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2), x)`

**Giac [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}} dx = \int \frac{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}}f + d\right)^n}{\sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}}} dx$$

input `integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2),x, algorithm="giac")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2)*f + d)^n/sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2), x)`

**Mupad [B] (verification not implemented)**

Time = 22.06 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}} dx = \frac{f \left(d + ex + f\sqrt{\frac{e^2x^2 + 2dex + af^2}{f^2}}\right)^n}{en}$$

input `int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n/(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2),x)`

output `(f*(d + e*x + f*((a*f^2 + e^2*x^2 + 2*d*e*x)/f^2)^(1/2))^n)/(e*n)`

**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.83

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}} dx = \frac{(\sqrt{e^2x^2 + af^2 + 2dex} + d + ex)^n f}{en}$$

input `int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2),x)`

output `((sqrt(a*f**2 + 2*d*e*x + e**2*x**2) + d + e*x)**n*f)/(e*n)`

**3.48** 
$$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\left(a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}\right)^{3/2}} dx$$

Optimal result	335
Mathematica [A] (verified)	336
Rubi [A] (verified)	336
Maple [F]	338
Fricas [F]	338
Sympy [F(-2)]	339
Maxima [F]	339
Giac [F]	339
Mupad [F(-1)]	340
Reduce [F]	340

**Optimal result**

Integrand size = 58, antiderivative size = 122

$$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\left(a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}\right)^{3/2}} dx = \frac{4f^3\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^{2+n} \operatorname{Hypergeometric2F1}\left(2, \frac{2+n}{2}, \frac{2+n}{2}+1, \frac{e\left(d^2-af^2\right)^2}{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^2}\right)}{e\left(d^2-af^2\right)^2(2+n)}$$

output

```
4*f^3*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^(2+n)*hypergeom([2, 1+1/2*n], [2+1/2*n], (d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^2/(-a*f^2+d^2))/e/(-a*f^2+d^2)^(2+n)
```



**Mathematica [A] (verified)**

Time = 4.35 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.92

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^{3/2}} dx = \frac{4f^3 \left(d + ex + f\sqrt{a + \frac{e^2x^2}{f^2}}\right)^{2+n} \text{Hypergeometric2F1}\left(2, \frac{2+n}{2}\right)}{e(d^2 - af^2)^2(2+n)}$$

input

```
Integrate[(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n/(a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2)^(3/2), x]
```

output

```
(4*f^3*(d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^(2 + n)*Hypergeometric2F1[2, (2 + n)/2, (4 + n)/2, (d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^2/(d^2 - a*f^2)])/(e*(d^2 - a*f^2)^2*(2 + n))
```

**Rubi [A] (verified)**Time = 0.41 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.052$ , Rules used = {2546, 27, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\left(f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex\right)^n}{\left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^{3/2}} dx$$

↓ 2546

$$2f^3 \int \frac{2\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2}} + a\right)^{n+1}}{e\left(d^2 - af^2 - \left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2}} + a\right)^2\right)^2 d\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2}} + a\right)} dx$$

↓ 27

$$4f^3 \int \frac{\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^{n+1}}{\left(d^2-af^2-\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^2\right)^2} d\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)$$

e  
↓  
278

$$\frac{4f^3 \left(f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex\right)^{n+2} \operatorname{Hypergeometric2F1}\left(2, \frac{n+2}{2}, \frac{n+4}{2}, \frac{\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^2}{d^2-af^2}\right)}{e(n+2)(d^2-af^2)^2}$$

input

```
Int[(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n/(a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2)^(3/2), x]
```

output

```
(4*f^3*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^(2 + n)*Hypergeometric2F1[2, (2 + n)/2, (4 + n)/2, (d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^2/(d^2 - a*f^2)]/(e*(d^2 - a*f^2)^2*(2 + n))
```

### Defintions of rubi rules used

rule 27

```
Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

rule 278

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])
```

rule 2546

```
Int[((g_.) + (h_.)*(x_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*S
qrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2])^(n_.), x_Symbol] :> Simp[(2/f^(2*m)
)*(i/c)^m Subst[Int[x^n*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e
*x^2)^(2*m + 1)/(-2*d*e + b*f^2 + 2*e*x)^(2*(m + 1))), x], x, d + e*x + f*S
qrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, n}, x] && Eq
Q[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && EqQ[c*h - b*i, 0] && IntegerQ[2*m
] && (IntegerQ[m] || GtQ[i/c, 0])
```

**Maple [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^{\frac{3}{2}}} dx$$

input

```
int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f
^2)^(3/2),x)
```

output

```
int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f
^2)^(3/2),x)
```

**Fricas [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^{3/2}} dx = \int \frac{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}}f + d\right)^n}{\left(\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}\right)^{\frac{3}{2}}} dx$$

input

```
integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2
*x^2/f^2)^(3/2),x, algorithm="fricas")
```

output

```
integral((e*x + f*sqrt((e^2*x^2 + a*f^2 + 2*d*e*x)/f^2) + d)^n*f^4*sqrt((e
^2*x^2 + a*f^2 + 2*d*e*x)/f^2)/(e^4*x^4 + 4*d*e^3*x^3 + a^2*f^4 + 4*a*d*e*
f^2*x + 2*(a*e^2*f^2 + 2*d^2*e^2)*x^2), x)
```

**Sympy [F(-2)]**

Exception generated.

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^{3/2}} dx = \text{Exception raised: HeuristicGCDFailed}$$

input `integrate((d+e*x+f*(a+2*d*e*x/f**2+e**2*x**2/f**2)**(1/2))**n/(a+2*d*e*x/f**2+e**2*x**2/f**2)**(3/2),x)`

output `Exception raised: HeuristicGCDFailed >> no luck`

**Maxima [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^{3/2}} dx = \int \frac{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}}f + d\right)^n}{\left(\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}\right)^{3/2}} dx$$

input `integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f^2)^(3/2),x, algorithm="maxima")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2)*f + d)^n/(e^2*x^2/f^2 + a + 2*d*e*x/f^2)^(3/2), x)`

**Giac [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^{3/2}} dx = \int \frac{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}}f + d\right)^n}{\left(\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}\right)^{3/2}} dx$$

input `integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f^2)^(3/2),x, algorithm="giac")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2))*f + d)^n/(e^2*x^2/f^2 + a + 2*d*e*x/f^2)^(3/2), x)`

### Mupad [F(-1)]

Timed out.

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^{3/2}} dx = \int \frac{\left(d + f\sqrt{a + \frac{e^2x^2}{f^2} + \frac{2dex}{f^2}} + ex\right)^n}{\left(a + \frac{e^2x^2}{f^2} + \frac{2dex}{f^2}\right)^{3/2}} dx$$

input `int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n/(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(3/2),x)`

output `int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n/(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(3/2), x)`

### Reduce [F]

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)^{3/2}} dx = \left( \int \frac{(\sqrt{e^2x^2 + af^2 + 2dex} + d + ex)^n}{\sqrt{e^2x^2 + af^2 + 2dex} a f^2 + 2\sqrt{e^2x^2 + af^2 + 2dex} dex + \sqrt{e^2x^2 + af^2 + 2dex}}$$

input `int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a+2*d*e*x/f^2+e^2*x^2/f^2)^(3/2),x)`

output `int((sqrt(a*f**2 + 2*d*e*x + e**2*x**2) + d + e*x)**n/(sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*a*f**2 + 2*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*d*e*x + sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*e**2*x**2),x)*f**3`

**3.49** 
$$\int \frac{\left(d+ex+f\sqrt{\frac{af^2+ex(2d+ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2+ex(2d+ex)}{f^2}}} dx$$

Optimal result	341
Mathematica [A] (verified)	341
Rubi [A] (verified)	342
Maple [F]	343
Fricas [A] (verification not implemented)	344
Sympy [F(-1)]	344
Maxima [F]	344
Giac [F]	345
Mupad [B] (verification not implemented)	345
Reduce [B] (verification not implemented)	346

**Optimal result**

Integrand size = 58, antiderivative size = 41

$$\int \frac{\left(d+ex+f\sqrt{\frac{af^2+ex(2d+ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2+ex(2d+ex)}{f^2}}} dx = \frac{f\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{en}$$

output `f*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/e/n`

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.88

$$\int \frac{\left(d+ex+f\sqrt{\frac{af^2+ex(2d+ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2+ex(2d+ex)}{f^2}}} dx = \frac{f\left(d+ex+f\sqrt{a+\frac{ex(2d+ex)}{f^2}}\right)^n}{en}$$

input `Integrate[(d + e*x + f*Sqrt[(a*f^2 + e*x*(2*d + e*x))/f^2])^n/Sqrt[(a*f^2 + e*x*(2*d + e*x))/f^2],x]`

output  $(f*(d + e*x + f*\text{Sqrt}[a + (e*x*(2*d + e*x))/f^2])^n)/(e*n)$

### Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.052$ , Rules used = {2552, 2546, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\left(f \sqrt{\frac{af^2 + ex(2d+ex)}{f^2}} + d + ex\right)^n}{\sqrt{\frac{af^2 + ex(2d+ex)}{f^2}}} dx$$

↓ 2552

$$\int \frac{\left(f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex\right)^n}{\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}} dx$$

↓ 2546

$$2f \int \frac{\left(d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^{n-1}}{2e} d \left(d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)$$

↓ 15

$$\frac{f \left(f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex\right)^n}{en}$$

input  $\text{Int}[(d + e*x + f*\text{Sqrt}[(a*f^2 + e*x*(2*d + e*x))/f^2])^n/\text{Sqrt}[(a*f^2 + e*x*(2*d + e*x))/f^2], x]$

output  $(f*(d + e*x + f*\text{Sqrt}[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n)/(e*n)$

## Definitions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2546 `Int[((g_.) + (h_.)*(x_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(2/f^(2*m))* (i/c)^m Subst[Int[x^n*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)^(2*m + 1)/(-2*d*e + b*f^2 + 2*e*x)^(2*(m + 1))), x], x, d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && EqQ[c*h - b*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

rule 2552 `Int[((u_) + (f_.)*((j_.) + (k_.)*Sqrt[v_]))^(n_.)*(w_)^(m_.), x_Symbol] := Int[ExpandToSum[w, x]^m*(ExpandToSum[u + f*j, x] + f*k*Sqrt[ExpandToSum[v, x]])^n, x] /; FreeQ[{f, j, k, m, n}, x] && LinearQ[u, x] && QuadraticQ[{v, w}, x] && !(LinearMatchQ[u, x] && QuadraticMatchQ[{v, w}, x] && (EqQ[j, 0] || EqQ[f, 1])) && EqQ[Coefficient[u, x, 1]^2 - Coefficient[v, x, 2]*f^2*k^2, 0]`

## Maple [F]

$$\int \frac{\left(d + ex + f\sqrt{\frac{f^2a + ex(ex+2d)}{f^2}}\right)^n}{\sqrt{\frac{f^2a + ex(ex+2d)}{f^2}}} dx$$

input `int((d+e*x+f*((f^2*a+e*x*(e*x+2*d))/f^2)^(1/2))^n/((f^2*a+e*x*(e*x+2*d))/f^2)^(1/2),x)`

output `int((d+e*x+f*((f^2*a+e*x*(e*x+2*d))/f^2)^(1/2))^n/((f^2*a+e*x*(e*x+2*d))/f^2)^(1/2),x)`



**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \frac{\left(d + ex + f \sqrt{\frac{af^2 + ex(2d+ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2 + ex(2d+ex)}{f^2}}} dx = \frac{\left(ex + f \sqrt{\frac{e^2x^2 + af^2 + 2dex}{f^2}} + d\right)^n f}{en}$$

input `integrate((d+e*x+f*((a*f^2+e*x*(e*x+2*d))/f^2)^(1/2))^n/((a*f^2+e*x*(e*x+2*d))/f^2)^(1/2),x, algorithm="fricas")`

output `(e*x + f*sqrt((e^2*x^2 + a*f^2 + 2*d*e*x)/f^2) + d)^n*f/(e*n)`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{\left(d + ex + f \sqrt{\frac{af^2 + ex(2d+ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2 + ex(2d+ex)}{f^2}}} dx = \text{Timed out}$$

input `integrate((d+e*x+f*((a*f**2+e*x*(e*x+2*d))/f**2)**(1/2))**n/((a*f**2+e*x*(e*x+2*d))/f**2)**(1/2),x)`

output `Timed out`

**Maxima [F]**

$$\int \frac{\left(d + ex + f \sqrt{\frac{af^2 + ex(2d+ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2 + ex(2d+ex)}{f^2}}} dx = \int \frac{\left(ex + d + \sqrt{af^2 + (ex + 2d)ex}\right)^n}{\frac{\sqrt{af^2 + (ex + 2d)ex}}{f}} dx$$

input `integrate((d+e*x+f*((a*f^2+e*x*(e*x+2*d))/f^2)^(1/2))^n/((a*f^2+e*x*(e*x+2*d))/f^2)^(1/2),x, algorithm="maxima")`

output `f*integrate((e*x + d + sqrt(a*f^2 + (e*x + 2*d)*e*x))^n/sqrt(a*f^2 + (e*x + 2*d)*e*x), x)`

### Giac [F]

$$\int \frac{\left(d + ex + f \sqrt{\frac{af^2 + ex(2d + ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2 + ex(2d + ex)}{f^2}}} dx = \int \frac{\left(ex + f \sqrt{\frac{af^2 + (ex + 2d)ex}{f^2}} + d\right)^n}{\sqrt{\frac{af^2 + (ex + 2d)ex}{f^2}}} dx$$

input `integrate((d+e*x+f*((a*f^2+e*x*(e*x+2*d))/f^2)^(1/2))^n/((a*f^2+e*x*(e*x+2*d))/f^2)^(1/2),x, algorithm="giac")`

output `integrate((e*x + f*sqrt((a*f^2 + (e*x + 2*d)*e*x)/f^2) + d)^n/sqrt((a*f^2 + (e*x + 2*d)*e*x)/f^2), x)`

### Mupad [B] (verification not implemented)

Time = 21.99 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.95

$$\int \frac{\left(d + ex + f \sqrt{\frac{af^2 + ex(2d + ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2 + ex(2d + ex)}{f^2}}} dx = \frac{f \left(d + ex + f \sqrt{\frac{af^2 + ex(2d + ex)}{f^2}}\right)^n}{en}$$

input `int((d + e*x + f*((a*f^2 + e*x*(2*d + e*x))/f^2)^(1/2))^n/((a*f^2 + e*x*(2*d + e*x))/f^2)^(1/2),x)`

output `(f*(d + e*x + f*((a*f^2 + e*x*(2*d + e*x))/f^2)^(1/2))^n)/(e*n)`

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.83

$$\int \frac{\left(d + ex + f \sqrt{\frac{af^2 + ex(2d + ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2 + ex(2d + ex)}{f^2}}} dx = \frac{(\sqrt{e^2x^2 + af^2 + 2dex} + d + ex)^n f}{en}$$

input

```
int((d+e*x+f*((a*f^2+e*x*(e*x+2*d))/f^2)^(1/2))^n/((a*f^2+e*x*(e*x+2*d))/f^2)^(1/2),x)
```

output

```
((sqrt(a*f**2 + 2*d*e*x + e**2*x**2) + d + e*x)**n*f)/(e*n)
```

**3.50** 
$$\int \sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}} \left( d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)$$

Optimal result	347
Mathematica [A] (verified)	348
Rubi [A] (verified)	348
Maple [F]	350
Fricas [A] (verification not implemented)	351
Sympy [F]	351
Maxima [F]	352
Giac [F]	352
Mupad [F(-1)]	353
Reduce [B] (verification not implemented)	353

**Optimal result**

Integrand size = 62, antiderivative size = 327

$$\int \sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}} \left( d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= -\frac{(d^2 - af^2)^2 \sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}} \left( d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{-2+n}}{4ef(2 - n)\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}}$$

$$- \frac{(d^2 - af^2) \sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}} \left( d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n}{2efn\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}}$$

$$+ \frac{\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}} \left( d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^{2+n}}{4ef(2 + n)\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}}$$

output

```
-1/4*(-a*f^2+d^2)^2*(a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(1/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^(-2+n)/e/f/(2-n)/(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2)-1/2*(-a*f^2+d^2)*(a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(1/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/e/f/n/(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2)+1/4*(a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(1/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^(2+n)/e/f/(2+n)/(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2)
```

### Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.54

$$\int \sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \frac{\sqrt{g \left( a + \frac{ex(2d+ex)}{f^2} \right)} \left( d + ex + f \sqrt{a + \frac{ex(2d+ex)}{f^2}} \right)^n \left( \frac{2(-d^2+af^2)}{n} + \frac{(d^2-af^2)^2}{(-2+n) \left( d+ex+f \sqrt{a + \frac{ex(2d+ex)}{f^2}} \right)^2} + \frac{(d+ex+f \sqrt{a + \frac{ex(2d+ex)}{f^2}})^2}{4ef \sqrt{a + \frac{ex(2d+ex)}{f^2}}} \right)}{4ef \sqrt{a + \frac{ex(2d+ex)}{f^2}}}$$

input `Integrate[Sqrt[a*g + (2*d*e*g*x)/f^2 + (e^2*g*x^2)/f^2]*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n,x]`

output `(Sqrt[g*(a + (e*x*(2*d + e*x))/f^2)]*(d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2]))^n*((2*(-d^2 + a*f^2))/n + (d^2 - a*f^2)^2/((-2 + n)*(d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^2) + (d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^2/(2 + n))/(4*e*f*Sqrt[a + (e*x*(2*d + e*x))/f^2])`

### Rubi [A] (verified)

Time = 0.73 (sec) , antiderivative size = 208, normalized size of antiderivative = 0.64, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.081$ , Rules used = {2548, 2546, 27, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}} \left( f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex \right)^n dx$$

↓ 2548

$$\frac{\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}} \int \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \left( d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^n dx}{\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}}$$

↓ 2546

$$\frac{2\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}} \int \frac{\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^{n-3} \left(d^2-af^2 - \left(d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^2\right)^2}{8e} d\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)}{f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}}$$

↓ 27

$$\frac{\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}} \int \left(d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^{n-3} \left(d^2-af^2 - \left(d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^2\right)^2 d\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)}{4ef\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}}$$

↓ 244

$$\frac{\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}} \int \left((d^2-af^2)^2 \left(d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^{n-3} - 2(d^2-af^2) \left(d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^2\right) d\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)}{4ef\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}}$$

↓ 2009

$$\frac{\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}} \left( -\frac{(d^2-af^2)^2 \left(f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex\right)^{n-2}}{2-n} - \frac{2(d^2-af^2) \left(f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex\right)^n}{n} + \frac{\left(f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex\right)^{n+1}}{n+1} \right)}{4ef\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}}$$

input

```
Int[Sqrt[a*g + (2*d*e*g*x)/f^2 + (e^2*g*x^2)/f^2]*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n,x]
```

output

```
(Sqrt[a*g + (2*d*e*g*x)/f^2 + (e^2*g*x^2)/f^2]*(-(((d^2 - a*f^2)^2*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^(-2 + n))/(2 - n) - (2*(d^2 - a*f^2)*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n)/n + (d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^(2 + n)/(2 + n)))/(4*e*f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])
```

## Definitions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 244 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2546 `Int[((g_) + (h_)*(x_) + (i_)*(x_)^2)^(m_)*((d_) + (e_)*(x_) + (f_)*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2])^(n_), x_Symbol] := Simp[(2/f^(2*m))*(i/c)^m Subst[Int[x^n*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)^(2*m + 1)/(-2*d*e + b*f^2 + 2*e*x)^(2*(m + 1))), x], x, d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && EqQ[c*h - b*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

rule 2548 `Int[((g_) + (h_)*(x_) + (i_)*(x_)^2)^(m_)*((d_) + (e_)*(x_) + (f_)*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2])^(n_), x_Symbol] := Simp[(i/c)^(m - 1/2)*(Sqrt[g + h*x + i*x^2]/Sqrt[a + b*x + c*x^2]) Int[(a + b*x + c*x^2)^m*(d + e*x + f*Sqrt[a + b*x + c*x^2])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && EqQ[c*h - b*i, 0] && IGtQ[m + 1/2, 0] && !GtQ[i/c, 0]`

Maple **[F]**

$$\int \sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

input `int((a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(1/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x)`

output

```
int((a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(1/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x)
```

**Fricas [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 231, normalized size of antiderivative = 0.71

$$\int \sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx =$$

$$\frac{(2e^3nx^3 + 6de^2nx^2 + 2adf^2n + 2(aef^2 + 2d^2e)nx - (e^2fn^2x^2 + af^3n^2 + 2defn^2x - 2af^3 + 2d^2f^2n))}{aef^2n^3 - 4aef^2n + (e^3n^3 - 4e^3n)x^2 + 2(de^2n^2x^2 + 2defn^2x - 2af^3 + 2d^2f^2n)}$$

input

```
integrate((a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(1/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n,x, algorithm="fricas")
```

output

```
-(2*e^3*n*x^3 + 6*d*e^2*n*x^2 + 2*a*d*f^2*n + 2*(a*e*f^2 + 2*d^2*e)*n*x - (e^2*f*n^2*x^2 + a*f^3*n^2 + 2*d*e*f*n^2*x - 2*a*f^3 + 2*d^2*f)*sqrt((e^2*x^2 + a*f^2 + 2*d*e*x)/f^2))*(e*x + f*sqrt((e^2*x^2 + a*f^2 + 2*d*e*x)/f^2) + d)^n*sqrt((e^2*g*x^2 + a*f^2*g + 2*d*e*g*x)/f^2)/(a*e*f^2*n^3 - 4*a*e*f^2*n + (e^3*n^3 - 4*e^3*n)*x^2 + 2*(d*e^2*n^3 - 4*d*e^2*n)*x)
```

**Sympy [F]**

$$\int \sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \int \sqrt{g \left( a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2} \right)} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

input

```
integrate((a*g+2*d*e*g*x/f**2+e**2*g*x**2/f**2)**(1/2)*(d+e*x+f*(a+2*d*e*x/f**2+e**2*x**2/f**2)**(1/2))**n,x)
```



output

```
Integral(sqrt(g*(a + 2*d*e*x/f**2 + e**2*x**2/f**2))*(d + e*x + f*sqrt(a +
2*d*e*x/f**2 + e**2*x**2/f**2))**n, x)
```

**Maxima [F]**

$$\int \sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}} \left( d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \int \sqrt{\frac{e^2gx^2}{f^2} + ag + \frac{2degx}{f^2}} \left( ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}}f + d \right)^n dx$$

input

```
integrate((a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(1/2)*(d+e*x+f*(a+2*d*e*x/f^2+
e^2*x^2/f^2)^(1/2))^n,x, algorithm="maxima")
```

output

```
integrate(sqrt(e^2*g*x^2/f^2 + a*g + 2*d*e*g*x/f^2)*(e*x + sqrt(e^2*x^2/f^
2 + a + 2*d*e*x/f^2)*f + d)^n, x)
```

**Giac [F]**

$$\int \sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}} \left( d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \int \sqrt{\frac{e^2gx^2}{f^2} + ag + \frac{2degx}{f^2}} \left( ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}}f + d \right)^n dx$$

input

```
integrate((a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(1/2)*(d+e*x+f*(a+2*d*e*x/f^2+
e^2*x^2/f^2)^(1/2))^n,x, algorithm="giac")
```

output

```
integrate(sqrt(e^2*g*x^2/f^2 + a*g + 2*d*e*g*x/f^2)*(e*x + sqrt(e^2*x^2/f^
2 + a + 2*d*e*x/f^2)*f + d)^n, x)
```

**Mupad [F(-1)]**

Timed out.

$$\int \sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \int \left( d + f \sqrt{a + \frac{e^2x^2}{f^2} + \frac{2dex}{f^2}} + ex \right)^n \sqrt{ag + \frac{e^2gx^2}{f^2} + \frac{2degx}{f^2}} dx$$

input

```
int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n*(a*g + (e^2*
g*x^2)/f^2 + (2*d*e*g*x)/f^2)^(1/2), x)
```

output

```
int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n*(a*g + (e^2*
g*x^2)/f^2 + (2*d*e*g*x)/f^2)^(1/2), x)
```

**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.40

$$\int \sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}} \left( d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \right)^n dx$$

$$= \frac{\sqrt{g} (\sqrt{e^2x^2 + af^2 + 2dex} + d + ex)^n (-2\sqrt{e^2x^2 + af^2 + 2dex} dn - 2\sqrt{e^2x^2 + af^2 + 2dex} enx + af^2n)}{efn(n^2 - 4)}$$

input

```
int((a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(1/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^
2/f^2)^(1/2))^n,x)
```

output

```
(sqrt(g)*(sqrt(a*f**2 + 2*d*e*x + e**2*x**2) + d + e*x)**n*( - 2*sqrt(a*f*
*2 + 2*d*e*x + e**2*x**2)*d*n - 2*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*e*n*x
+ a*f**2*n**2 - 2*a*f**2 + 2*d**2 + 2*d*e*n**2*x + e**2*n**2*x**2))/(e*f*
n*(n**2 - 4))
```

$$3.51 \quad \int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\sqrt{ag+\frac{2degx}{f^2}+\frac{e^2gx^2}{f^2}}} dx$$

Optimal result	354
Mathematica [A] (verified)	354
Rubi [A] (verified)	355
Maple [F]	357
Fricas [A] (verification not implemented)	357
Sympy [F]	358
Maxima [F]	358
Giac [F]	358
Mupad [F(-1)]	359
Reduce [B] (verification not implemented)	359

### Optimal result

Integrand size = 62, antiderivative size = 93

$$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\sqrt{ag+\frac{2degx}{f^2}+\frac{e^2gx^2}{f^2}}} dx = \frac{f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{en\sqrt{ag+\frac{2degx}{f^2}+\frac{e^2gx^2}{f^2}}}$$

output

```
f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/e/n/(a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(1/2)
```

### Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.82

$$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\sqrt{ag+\frac{2degx}{f^2}+\frac{e^2gx^2}{f^2}}} dx = \frac{f\sqrt{a+\frac{ex(2d+ex)}{f^2}}\left(d+ex+f\sqrt{a+\frac{ex(2d+ex)}{f^2}}\right)^n}{en\sqrt{g\left(a+\frac{ex(2d+ex)}{f^2}\right)}}$$

input

```
Integrate[(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n/Sqrt[a*g
+ (2*d*e*g*x)/f^2 + (e^2*g*x^2)/f^2], x]
```

output

```
(f*Sqrt[a + (e*x*(2*d + e*x))/f^2]*(d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))
/f^2])^n)/(e*n*Sqrt[g*(a + (e*x*(2*d + e*x))/f^2)])
```

**Rubi [A] (verified)**

Time = 0.65 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$ , Rules used = {2550, 2546, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\left(f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex\right)^n}{\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}}} dx$$

$$\downarrow \text{2550}$$

$$\frac{\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \int \frac{\left(d+ex+f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^n}{\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}} dx}{\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}}}$$

$$\downarrow \text{2546}$$

$$\frac{2f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \int \frac{\left(d+ex+f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^{n-1}}{2e} d\left(d + ex + f \sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)}{\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}}}$$

$$\downarrow \text{15}$$

$$\frac{f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \left(f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex\right)^n}{en \sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}}}$$

input

```
Int[(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n/Sqrt[a*g + (2*d*e*g*x)/f^2 + (e^2*g*x^2)/f^2],x]
```

output

```
(f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2]*(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n/(e*n*Sqrt[a*g + (2*d*e*g*x)/f^2 + (e^2*g*x^2)/f^2])
```

### Defintions of rubi rules used

rule 15

```
Int[(a_.)*(x_)^(m_.), x_Symbol] :> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]
```

rule 2546

```
Int[((g_.) + (h_.)*(x_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2])^(n_.), x_Symbol] :> Simp[(2/f^(2*m))*(i/c)^m Subst[Int[x^n*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)^(2*m + 1)/(-2*d*e + b*f^2 + 2*e*x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && EqQ[c*h - b*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])
```

rule 2550

```
Int[((g_.) + (h_.)*(x_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2])^(n_.), x_Symbol] :> Simp[(i/c)^(m + 1/2)*(Sqrt[a + b*x + c*x^2]/Sqrt[g + h*x + i*x^2]) Int[(a + b*x + c*x^2)^m*(d + e*x + f*Sqrt[a + b*x + c*x^2])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && EqQ[c*h - b*i, 0] && ILtQ[m - 1/2, 0] && !GtQ[i/c, 0]
```

**Maple [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}}} dx$$

input `int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(1/2),x)`

output `int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(1/2),x)`

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.26

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}}} dx$$

$$= \frac{\left(ex + f\sqrt{\frac{e^2x^2 + af^2 + 2dex}{f^2}} + d\right)^n f^3 \sqrt{\frac{e^2gx^2 + af^2g + 2degx}{f^2}} \sqrt{\frac{e^2x^2 + af^2 + 2dex}{f^2}}}{e^3gnx^2 + aef^2gn + 2de^2gnx}$$

input `integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(1/2),x, algorithm="fricas")`

output `(e*x + f*sqrt((e^2*x^2 + a*f^2 + 2*d*e*x)/f^2) + d)^n*f^3*sqrt((e^2*g*x^2 + a*f^2*g + 2*d*e*g*x)/f^2)*sqrt((e^2*x^2 + a*f^2 + 2*d*e*x)/f^2)/(e^3*g*n*x^2 + a*e*f^2*g*n + 2*d*e^2*g*n*x)`

**Sympy [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}}} dx = \int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\sqrt{g\left(a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}\right)}} dx$$

input `integrate((d+e*x+f*(a+2*d*e*x/f**2+e**2*x**2/f**2)**(1/2))**n/(a*g+2*d*e*g*x/f**2+e**2*g*x**2/f**2)**(1/2), x)`

output `Integral((d + e*x + f*sqrt(a + 2*d*e*x/f**2 + e**2*x**2/f**2))**n/sqrt(g*(a + 2*d*e*x/f**2 + e**2*x**2/f**2)), x)`

**Maxima [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}}} dx = \int \frac{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}}f + d\right)^n}{\sqrt{\frac{e^2gx^2}{f^2} + ag + \frac{2degx}{f^2}}} dx$$

input `integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(1/2), x, algorithm="maxima")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2)*f + d)^n/sqrt(e^2*g*x^2/f^2 + a*g + 2*d*e*g*x/f^2), x)`

**Giac [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}}} dx = \int \frac{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}}f + d\right)^n}{\sqrt{\frac{e^2gx^2}{f^2} + ag + \frac{2degx}{f^2}}} dx$$

input `integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(1/2),x, algorithm="giac")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2))*f + d)^n/sqrt(e^2*g*x^2/f^2 + a*g + 2*d*e*g*x/f^2), x)`

### Mupad [F(-1)]

Timed out.

$$\int \frac{\left(d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}}} dx = \int \frac{\left(d + f \sqrt{a + \frac{e^2x^2}{f^2} + \frac{2dex}{f^2}} + ex\right)^n}{\sqrt{ag + \frac{e^2gx^2}{f^2} + \frac{2degx}{f^2}}} dx$$

input `int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n/(a*g + (e^2*g*x^2)/f^2 + (2*d*e*g*x)/f^2)^(1/2),x)`

output `int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n/(a*g + (e^2*g*x^2)/f^2 + (2*d*e*g*x)/f^2)^(1/2), x)`

### Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.42

$$\int \frac{\left(d + ex + f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}}} dx = \frac{\sqrt{g} (\sqrt{e^2x^2 + af^2 + 2dex} + d + ex)^n f}{egn}$$

input `int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(1/2),x)`

output `(sqrt(g)*(sqrt(a*f**2 + 2*d*e*x + e**2*x**2) + d + e*x)**n*f)/(e*g*n)`



**3.52** 
$$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\left(ag+\frac{2degx}{f^2}+\frac{e^2gx^2}{f^2}\right)^{3/2}} dx$$

Optimal result	360
Mathematica [A] (verified)	361
Rubi [A] (verified)	361
Maple [F]	363
Fricas [F]	364
Sympy [F(-1)]	364
Maxima [F]	365
Giac [F]	365
Mupad [F(-1)]	365
Reduce [F]	366

**Optimal result**

Integrand size = 62, antiderivative size = 177

$$\int \frac{\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{\left(ag+\frac{2degx}{f^2}+\frac{e^2gx^2}{f^2}\right)^{3/2}} dx = \frac{4f^3\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^{2+n}}{e(d^2-af^2)^2g(2+n)\sqrt{ag+\frac{2degx}{f^2}+\frac{e^2gx^2}{f^2}}} \text{Hypergeom}$$

output

```
4*f^3*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^(2+n)*hypergeom([2, 1+1/2*n], [2+1/2*n], (d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^2/(-a*f^2+d^2))/e/(-a*f^2+d^2)^2/g/(2+n)/(a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.86

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}\right)^{3/2}} dx = \frac{4f^3 \left(a + \frac{ex(2d+ex)}{f^2}\right)^{3/2} \left(d + ex + f\sqrt{a + \frac{ex(2d+ex)}{f^2}}\right)^{2+n} \text{Hypergeometric}}{e(d^2 - af^2)^2(2+n) \left(g \left(a + \frac{ex(2d+ex)}{f^2}\right)\right)^{2+n}}$$

input

```
Integrate[(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n/(a*g + (2*d*e*g*x)/f^2 + (e^2*g*x^2)/f^2)^(3/2),x]
```

output

```
(4*f^3*(a + (e*x*(2*d + e*x))/f^2)^(3/2)*(d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^(2 + n)*Hypergeometric2F1[2, (2 + n)/2, (4 + n)/2, (d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^2/(d^2 - a*f^2)]/(e*(d^2 - a*f^2)^2*(2 + n)*(g*(a + (e*x*(2*d + e*x))/f^2))^(3/2))
```

**Rubi [A] (verified)**Time = 0.70 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.065$ , Rules used = {2550, 2546, 27, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\left(f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex\right)^n}{\left(ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}\right)^{3/2}} dx$$

↓ 2550

$$\frac{\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \int \frac{\left(d + ex + f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^n}{\left(\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a\right)^{3/2}} dx}{g\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}}}$$

↓ 2546

$$\frac{2f^3 \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \int \frac{2 \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n+1}}{e \left( d^2 - af^2 - \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^2 \right)^2} d \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)}{g \sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}}}$$

↓ 27

$$\frac{4f^3 \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \int \frac{\left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^{n+1}}{\left( d^2 - af^2 - \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^2 \right)^2} d \left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)}{eg \sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}}}$$

↓ 278

$$\frac{4f^3 \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \left( f \sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex \right)^{n+2} \operatorname{Hypergeometric2F1} \left( 2, \frac{n+2}{2}, \frac{n+4}{2}, \frac{\left( d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a} \right)^2}{d^2 - af^2} \right)}{eg(n+2)(d^2 - af^2)^2 \sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}}}$$

input

```
Int[(d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^n/(a*g + (2*d*e*
g*x)/f^2 + (e^2*g*x^2)/f^2)^(3/2),x]
```

output

```
(4*f^3*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2]*(d + e*x + f*Sqrt[a + (2*d*
e*x)/f^2 + (e^2*x^2)/f^2])^(2 + n)*Hypergeometric2F1[2, (2 + n)/2, (4 + n)
/2, (d + e*x + f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2])^2/(d^2 - a*f^2)]
)/(e*(d^2 - a*f^2)^2*g*(2 + n)*Sqrt[a*g + (2*d*e*g*x)/f^2 + (e^2*g*x^2)/f
^2])
```

## Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x)] /; FreeQ[b, x]`

rule 278 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 2546 `Int[((g_) + (h_)*(x_) + (i_)*(x_)^2)^(m_)*((d_) + (e_)*(x_) + (f_)*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2])^(n_), x_Symbol] := Simp[(2/f^(2*m))* (i/c)^m Subst[Int[x^n*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e*x^2)^(2*m + 1)/(-2*d*e + b*f^2 + 2*e*x)^(2*(m + 1))), x], x, d + e*x + f*Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && EqQ[c*h - b*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

rule 2550 `Int[((g_) + (h_)*(x_) + (i_)*(x_)^2)^(m_)*((d_) + (e_)*(x_) + (f_)*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2])^(n_), x_Symbol] := Simp[(i/c)^(m + 1/2)*(Sqrt[a + b*x + c*x^2]/Sqrt[g + h*x + i*x^2]) Int[(a + b*x + c*x^2)^m*(d + e*x + f*Sqrt[a + b*x + c*x^2])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && EqQ[c*h - b*i, 0] && ILtQ[m - 1/2, 0] && !GtQ[i/c, 0]`

## Maple [F]

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}\right)^{\frac{3}{2}}} dx$$

input `int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(3/2),x)`

output

```
int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(3/2),x)
```

**Fricas [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}\right)^{3/2}} dx = \int \frac{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}}f + d\right)^n}{\left(\frac{e^2gx^2}{f^2} + ag + \frac{2degx}{f^2}\right)^{3/2}} dx$$

input

```
integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(3/2),x, algorithm="fricas")
```

output

```
integral((e*x + f*sqrt((e^2*x^2 + a*f^2 + 2*d*e*x)/f^2) + d)^n*f^4*sqrt((e^2*g*x^2 + a*f^2*g + 2*d*e*g*x)/f^2)/(e^4*g^2*x^4 + 4*d*e^3*g^2*x^3 + a^2*f^4*g^2 + 4*a*d*e*f^2*g^2*x + 2*(a*e^2*f^2 + 2*d^2*e^2)*g^2*x^2), x)
```

**Sympy [F(-1)]**

Timed out.

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}\right)^{3/2}} dx = \text{Timed out}$$

input

```
integrate((d+e*x+f*(a+2*d*e*x/f**2+e**2*x**2/f**2)**(1/2))**n/(a*g+2*d*e*g*x/f**2+e**2*g*x**2/f**2)**(3/2),x)
```

output

Timed out

**Maxima [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}\right)^{3/2}} dx = \int \frac{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}}f + d\right)^n}{\left(\frac{e^2gx^2}{f^2} + ag + \frac{2degx}{f^2}\right)^{3/2}} dx$$

input `integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(3/2),x, algorithm="maxima")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2)*f + d)^n/(e^2*g*x^2/f^2 + a*g + 2*d*e*g*x/f^2)^(3/2), x)`

**Giac [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}\right)^{3/2}} dx = \int \frac{\left(ex + \sqrt{\frac{e^2x^2}{f^2} + a + \frac{2dex}{f^2}}f + d\right)^n}{\left(\frac{e^2gx^2}{f^2} + ag + \frac{2degx}{f^2}\right)^{3/2}} dx$$

input `integrate((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(3/2),x, algorithm="giac")`

output `integrate((e*x + sqrt(e^2*x^2/f^2 + a + 2*d*e*x/f^2)*f + d)^n/(e^2*g*x^2/f^2 + a*g + 2*d*e*g*x/f^2)^(3/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}\right)^{3/2}} dx = \int \frac{\left(d + f\sqrt{a + \frac{e^2x^2}{f^2} + \frac{2dex}{f^2}} + ex\right)^n}{\left(ag + \frac{e^2gx^2}{f^2} + \frac{2degx}{f^2}\right)^{3/2}} dx$$

input

```
int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n/(a*g + (e^2*
g*x^2)/f^2 + (2*d*e*g*x)/f^2)^(3/2), x)
```

output

```
int((d + f*(a + (e^2*x^2)/f^2 + (2*d*e*x)/f^2)^(1/2) + e*x)^n/(a*g + (e^2*
g*x^2)/f^2 + (2*d*e*g*x)/f^2)^(3/2), x)
```

**Reduce [F]**

$$\int \frac{\left(d + ex + f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}}\right)^n}{\left(ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}\right)^{3/2}} dx = \frac{\left(\int \frac{\left(\sqrt{e^2x^2 + af^2 + 2dex} + d + ex\right)^n}{\sqrt{e^2x^2 + af^2 + 2dex} a f^2 + 2\sqrt{e^2x^2 + af^2 + 2dex} dex + \sqrt{e^2x^2 + af^2 + 2dex} e^2x^2} dx\right)}{\sqrt{g} g}$$

input

```
int((d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/(a*g+2*d*e*g*x/f^2+e^2*
g*x^2/f^2)^(3/2), x)
```

output

```
(int((sqrt(a*f**2 + 2*d*e*x + e**2*x**2) + d + e*x)**n/(sqrt(a*f**2 + 2*d*
e*x + e**2*x**2)*a*f**2 + 2*sqrt(a*f**2 + 2*d*e*x + e**2*x**2)*d*e*x + sqr
t(a*f**2 + 2*d*e*x + e**2*x**2)*e**2*x**2), x)*f**3)/(sqrt(g)*g)
```

**3.53** 
$$\int \frac{\left(d+ex+f\sqrt{\frac{af^2+ex(2d+ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2g+egx(2d+ex)}{f^2}}} dx$$

Optimal result	367
Mathematica [A] (verified)	368
Rubi [A] (verified)	368
Maple [F]	370
Fricas [A] (verification not implemented)	370
Sympy [F(-1)]	371
Maxima [F]	371
Giac [F]	372
Mupad [F(-1)]	372
Reduce [B] (verification not implemented)	372

**Optimal result**

Integrand size = 60, antiderivative size = 93

$$\int \frac{\left(d+ex+f\sqrt{\frac{af^2+ex(2d+ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2g+egx(2d+ex)}{f^2}}} dx$$

$$= \frac{f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\left(d+ex+f\sqrt{a+\frac{2dex}{f^2}+\frac{e^2x^2}{f^2}}\right)^n}{en\sqrt{ag+\frac{2degx}{f^2}+\frac{e^2gx^2}{f^2}}}$$

output

```
f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2)*(d+e*x+f*(a+2*d*e*x/f^2+e^2*x^2/f^2)^(1/2))^n/e/n/(a*g+2*d*e*g*x/f^2+e^2*g*x^2/f^2)^(1/2)
```



**Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.82

$$\int \frac{\left(d + ex + f\sqrt{\frac{af^2 + ex(2d+ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2g + egx(2d+ex)}{f^2}}} dx = \frac{f\sqrt{a + \frac{ex(2d+ex)}{f^2}} \left(d + ex + f\sqrt{a + \frac{ex(2d+ex)}{f^2}}\right)^n}{en\sqrt{g\left(a + \frac{ex(2d+ex)}{f^2}\right)}}$$

input

```
Integrate[(d + e*x + f*Sqrt[(a*f^2 + e*x*(2*d + e*x))/f^2])^n/Sqrt[(a*f^2*g + e*g*x*(2*d + e*x))/f^2],x]
```

output

```
(f*Sqrt[a + (e*x*(2*d + e*x))/f^2]*(d + e*x + f*Sqrt[a + (e*x*(2*d + e*x))/f^2])^n)/(e*n*Sqrt[g*(a + (e*x*(2*d + e*x))/f^2)])
```

**Rubi [A] (verified)**

Time = 0.84 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$ , Rules used = {2552, 2550, 2546, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\left(f\sqrt{\frac{af^2 + ex(2d+ex)}{f^2}} + d + ex\right)^n}{\sqrt{\frac{af^2g + egx(2d+ex)}{f^2}}} dx$$

↓ 2552

$$\int \frac{\left(f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex\right)^n}{\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}}} dx$$

↓ 2550

$$\begin{aligned}
& \frac{\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \int \frac{\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^n}{\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}} dx}{\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}}} \\
& \quad \downarrow \text{2546} \\
& \frac{2f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \int \frac{\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)^{n-1}}{2e} d\left(d+ex+f\sqrt{\frac{e^2x^2}{f^2} + \frac{2dex}{f^2} + a}\right)}{\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}}} \\
& \quad \downarrow \text{15} \\
& \frac{f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} \left(f\sqrt{a + \frac{2dex}{f^2} + \frac{e^2x^2}{f^2}} + d + ex\right)^n}{en\sqrt{ag + \frac{2degx}{f^2} + \frac{e^2gx^2}{f^2}}}
\end{aligned}$$

input

```
Int[(d + e*x + f*Sqrt[(a*f^2 + e*x*(2*d + e*x))/f^2])^n/Sqrt[(a*f^2*g + e*
g*x*(2*d + e*x))/f^2],x]
```

output

```
(f*Sqrt[a + (2*d*e*x)/f^2 + (e^2*x^2)/f^2]*(d + e*x + f*Sqrt[a + (2*d*e*x)
/f^2 + (e^2*x^2)/f^2])^n)/(e*n*Sqrt[a*g + (2*d*e*g*x)/f^2 + (e^2*g*x^2)/f
^2])
```

### Defintions of rubi rules used

rule 15

```
Int[(a_.)*(x_)^(m_.), x_Symbol] :> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[
{a, m}, x] && NeQ[m, -1]
```

rule 2546

```
Int[((g_.) + (h_.)*(x_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*S
qrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2])^(n_.), x_Symbol] :> Simp[(2/f^(2*m)
)*(i/c)^m Subst[Int[x^n*((d^2*e - (b*d - a*e)*f^2 - (2*d*e - b*f^2)*x + e
*x^2)^(2*m + 1)/(-2*d*e + b*f^2 + 2*e*x)^(2*(m + 1))], x], x, d + e*x + f*S
qrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, n}, x] && Eq
Q[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && EqQ[c*h - b*i, 0] && IntegerQ[2*m
] && (IntegerQ[m] || GtQ[i/c, 0])
```

rule 2550

```
Int[((g_.) + (h_.)*(x_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(i/c)^(m + 1/2)*(Sqrt[a + b*x + c*x^2]/Sqrt[g + h*x + i*x^2]) Int[(a + b*x + c*x^2)^m*(d + e*x + f*Sqrt[a + b*x + c*x^2])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && EqQ[c*h - b*i, 0] && ILtQ[m - 1/2, 0] && !GtQ[i/c, 0]
```

rule 2552

```
Int[((u_) + (f_.)*((j_.) + (k_.)*Sqrt[v_]))^(n_.)*(w_)^(m_.), x_Symbol] := Int[ExpandToSum[w, x]^m*(ExpandToSum[u + f*j, x] + f*k*Sqrt[ExpandToSum[v, x]])^n, x] /; FreeQ[{f, j, k, m, n}, x] && LinearQ[u, x] && QuadraticQ[{v, w}, x] && !(LinearMatchQ[u, x] && QuadraticMatchQ[{v, w}, x] && (EqQ[j, 0] || EqQ[f, 1])) && EqQ[Coefficient[u, x, 1]^2 - Coefficient[v, x, 2]*f^2*k^2, 0]
```

## Maple [F]

$$\int \frac{\left(d + ex + f \sqrt{\frac{f^2 a + ex(ex+2d)}{f^2}}\right)^n}{\sqrt{\frac{af^2g + egx(ex+2d)}{f^2}}} dx$$

input

```
int((d+e*x+f*((f^2*a+e*x*(e*x+2*d))/f^2)^(1/2))^n/((a*f^2*g+e*g*x*(e*x+2*d))/f^2)^(1/2),x)
```

output

```
int((d+e*x+f*((f^2*a+e*x*(e*x+2*d))/f^2)^(1/2))^n/((a*f^2*g+e*g*x*(e*x+2*d))/f^2)^(1/2),x)
```

## Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.26

$$\begin{aligned} & \int \frac{\left(d + ex + f \sqrt{\frac{af^2 + ex(2d+ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2g + egx(2d+ex)}{f^2}}} dx \\ &= \frac{\left(ex + f \sqrt{\frac{e^2x^2 + af^2 + 2dex}{f^2}} + d\right)^n f^3 \sqrt{\frac{e^2gx^2 + af^2g + 2degx}{f^2}} \sqrt{\frac{e^2x^2 + af^2 + 2dex}{f^2}}}{e^3gnx^2 + ae f^2gn + 2de^2gnx} \end{aligned}$$

input `integrate((d+e*x+f*((a*f^2+e*x*(e*x+2*d))/f^2)^(1/2))^n/((a*f^2*g+e*g*x*(e*x+2*d))/f^2)^(1/2),x, algorithm="fricas")`

output  $(e*x + f*\sqrt{(e^2*x^2 + a*f^2 + 2*d*e*x)/f^2} + d)^n*f^3*\sqrt{(e^2*g*x^2 + a*f^2*g + 2*d*e*g*x)/f^2}*\sqrt{(e^2*x^2 + a*f^2 + 2*d*e*x)/f^2}/(e^3*g*n*x^2 + a*e*f^2*g*n + 2*d*e^2*g*n*x)$

### Sympy [F(-1)]

Timed out.

$$\int \frac{\left(d + ex + f\sqrt{\frac{af^2 + ex(2d + ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2g + egx(2d + ex)}{f^2}}} dx = \text{Timed out}$$

input `integrate((d+e*x+f*((a*f**2+e*x*(e*x+2*d))/f**2)**(1/2))**n/((a*f**2*g+e*g*x*(e*x+2*d))/f**2)**(1/2),x)`

output Timed out

### Maxima [F]

$$\int \frac{\left(d + ex + f\sqrt{\frac{af^2 + ex(2d + ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2g + egx(2d + ex)}{f^2}}} dx = \int \frac{\left(ex + d + \sqrt{af^2 + (ex + 2d)ex}\right)^n}{\frac{\sqrt{af^2g + (ex + 2d)egx}}{f}} dx$$

input `integrate((d+e*x+f*((a*f^2+e*x*(e*x+2*d))/f^2)^(1/2))^n/((a*f^2*g+e*g*x*(e*x+2*d))/f^2)^(1/2),x, algorithm="maxima")`

output `f*integrate((e*x + d + sqrt(a*f^2 + (e*x + 2*d)*e*x))^n/sqrt(a*f^2*g + (e*x + 2*d)*e*g*x), x)`

**Giac [F]**

$$\int \frac{\left(d + ex + f \sqrt{\frac{af^2 + ex(2d+ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2g + egx(2d+ex)}{f^2}}} dx = \int \frac{\left(ex + f \sqrt{\frac{af^2 + (ex+2d)ex}{f^2}} + d\right)^n}{\sqrt{\frac{af^2g + (ex+2d)egx}{f^2}}} dx$$

input `integrate((d+e*x+f*((a*f^2+e*x*(e*x+2*d))/f^2)^(1/2))^n/((a*f^2*g+e*g*x*(e*x+2*d))/f^2)^(1/2),x, algorithm="giac")`

output `integrate((e*x + f*sqrt((a*f^2 + (e*x + 2*d)*e*x)/f^2) + d)^n/sqrt((a*f^2*g + (e*x + 2*d)*e*g*x)/f^2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\left(d + ex + f \sqrt{\frac{af^2 + ex(2d+ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2g + egx(2d+ex)}{f^2}}} dx = \int \frac{\left(d + ex + f \sqrt{\frac{af^2 + ex(2d+ex)}{f^2}}\right)^n}{\sqrt{\frac{agf^2 + egx(2d+ex)}{f^2}}} dx$$

input `int((d + e*x + f*((a*f^2 + e*x*(2*d + e*x))/f^2)^(1/2))^n/((a*f^2*g + e*g*x*(2*d + e*x))/f^2)^(1/2),x)`

output `int((d + e*x + f*((a*f^2 + e*x*(2*d + e*x))/f^2)^(1/2))^n/((a*f^2*g + e*g*x*(2*d + e*x))/f^2)^(1/2), x)`

**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.42

$$\int \frac{\left(d + ex + f \sqrt{\frac{af^2 + ex(2d+ex)}{f^2}}\right)^n}{\sqrt{\frac{af^2g + egx(2d+ex)}{f^2}}} dx = \frac{\sqrt{g} (\sqrt{e^2x^2 + af^2 + 2dex} + d + ex)^n f}{egn}$$

input `int((d+e*x+f*((a*f^2+e*x*(e*x+2*d))/f^2)^(1/2))^n/((a*f^2*g+e*g*x*(e*x+2*d))/f^2)^(1/2),x)`

output `(sqrt(g)*(sqrt(a*f**2 + 2*d*e*x + e**2*x**2) + d + e*x)**n*f)/(e*g*n)`

# CHAPTER 4

## APPENDIX

4.1	Listing of Grading functions . . . . .	374
4.2	Links to plain text integration problems used in this report for each CAS .	392

### 4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

#### Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
  ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "
  ,

```



```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]===Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]===Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]===Plus || Head[expn]===Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```

```

  Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
  If[AppellFunctionQ[Head[expn]],
    Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
  If[Head[expn]===RootSum,
    Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
  If[Head[expn]===Integrate || Head[expn]===Int,
    Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
  9]]]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=

```

```

  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

```

```

SpecialFunctionQ[func_] :=

```

```

  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

```

```

HypergeometricFunctionQ[func_] :=

```

```

  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

```

```

AppellFunctionQ[func_] :=

```

```

  MemberQ[{AppellF1}, func]

```

## Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

#   antiderivative
#   "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
            fi;
        fi;
    else #ExpnType(result) > ExpnType(optimal)
        if debug then
            print("ExpnType(result) > ExpnType(optimal)");
        fi;
        return "C",cat("Result contains higher order function than in optimal. Order ",
                        convert(ExpnType_result,string)," vs. order ",
                        convert(ExpnType_optimal,string),".");
    fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

## Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```



```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

## SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```



## 4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file