

# Computer Algebra Independent Integration Tests

Summer 2024

1-Algebraic-functions/1.5-Polynomial/146-1.7.1

Nasser M. Abbasi

May 17, 2024

Compiled on May 17, 2024 at 9:05pm

# Contents

<b>1</b>	<b>Introduction</b>	<b>4</b>
1.1	Listing of CAS systems tested . . . . .	5
1.2	Results . . . . .	6
1.3	Time and leaf size Performance . . . . .	10
1.4	Performance based on number of rules Rubi used . . . . .	12
1.5	Performance based on number of steps Rubi used . . . . .	13
1.6	Solved integrals histogram based on leaf size of result . . . . .	14
1.7	Solved integrals histogram based on CPU time used . . . . .	15
1.8	Leaf size vs. CPU time used . . . . .	16
1.9	list of integrals with no known antiderivative . . . . .	17
1.10	List of integrals solved by CAS but has no known antiderivative . . . . .	17
1.11	list of integrals solved by CAS but failed verification . . . . .	17
1.12	Timing . . . . .	18
1.13	Verification . . . . .	18
1.14	Important notes about some of the results . . . . .	19
1.15	Current tree layout of integration tests . . . . .	22
1.16	Design of the test system . . . . .	23
<b>2</b>	<b>detailed summary tables of results</b>	<b>24</b>
2.1	List of integrals sorted by grade for each CAS . . . . .	25
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	29
2.3	Detailed conclusion table specific for Rubi results . . . . .	43
<b>3</b>	<b>Listing of integrals</b>	<b>46</b>
3.1	$\int (a + bx + cx^2 + dx^3) dx$ . . . . .	48
3.2	$\int (-x^3 + x^4) dx$ . . . . .	53
3.3	$\int (-1 + x^5) dx$ . . . . .	58
3.4	$\int (7 + 4x) dx$ . . . . .	63
3.5	$\int (4x + \pi x^3) dx$ . . . . .	68
3.6	$\int (2x + 5x^2) dx$ . . . . .	73

3.7	$\int \left( \frac{x^2}{2} + \frac{x^3}{3} \right) dx$ . . . . .	78
3.8	$\int (3 - 5x + 2x^2) dx$ . . . . .	83
3.9	$\int (-2x + x^2 + x^3) dx$ . . . . .	88
3.10	$\int (1 - x^2 - 3x^5) dx$ . . . . .	93
3.11	$\int (5 + 2x + 3x^2 + 4x^3) dx$ . . . . .	98
3.12	$\int (3 - 19x^2 + 32x^4 - 16x^6)^4 dx$ . . . . .	103
3.13	$\int (3 - 19x^2 + 32x^4 - 16x^6)^3 dx$ . . . . .	110
3.14	$\int (3 - 19x^2 + 32x^4 - 16x^6)^2 dx$ . . . . .	117
3.15	$\int (3 - 19x^2 + 32x^4 - 16x^6) dx$ . . . . .	123
3.16	$\int \frac{1}{3-19x^2+32x^4-16x^6} dx$ . . . . .	128
3.17	$\int \frac{1}{(3-19x^2+32x^4-16x^6)^2} dx$ . . . . .	134
3.18	$\int \frac{1}{(3-19x^2+32x^4-16x^6)^3} dx$ . . . . .	140
3.19	$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3 dx$ . . . . .	147
3.20	$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2 dx$ . . . . .	154
3.21	$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5) dx$ . . . . .	160
3.22	$\int \frac{1}{a^5+5a^4bx+10a^3b^2x^2+10a^2b^3x^3+5ab^4x^4+b^5x^5} dx$ . . . . .	166
3.23	$\int \frac{1}{(a^5+5a^4bx+10a^3b^2x^2+10a^2b^3x^3+5ab^4x^4+b^5x^5)^2} dx$ . . . . .	172
3.24	$\int \frac{1}{(a^5+5a^4bx+10a^3b^2x^2+10a^2b^3x^3+5ab^4x^4+b^5x^5)^3} dx$ . . . . .	178
3.25	$\int \frac{1}{1+x^2+x^3+x^5} dx$ . . . . .	184
3.26	$\int \frac{1}{2+3(1+x)^5} dx$ . . . . .	189
3.27	$\int \frac{1}{5+15x+30x^2+30x^3+15x^4+3x^5} dx$ . . . . .	200
3.28	$\int \frac{1}{2+2(1+x)^3-3(1+x)^6} dx$ . . . . .	210
3.29	$\int \frac{1}{1-12x-39x^2-58x^3-45x^4-18x^5-3x^6} dx$ . . . . .	225
3.30	$\int \frac{1}{(a+bx^2)^4} dx$ . . . . .	241
3.31	$\int \frac{1}{(a^2+2abx^2+b^2x^4)^2} dx$ . . . . .	247
3.32	$\int \frac{1}{a^4+4a^3bx^2+6a^2b^2x^4+4ab^3x^6+b^4x^8} dx$ . . . . .	254
3.33	$\int \frac{1}{(a+b(s+x)^2)^4} dx$ . . . . .	261
3.34	$\int \frac{1}{(a^2+2ab(s+x)^2+b^2(s+x)^4)^2} dx$ . . . . .	269
3.35	$\int \frac{1}{a^4+4a^3b(s+x)^2+6a^2b^2(s+x)^4+4ab^3(s+x)^6+b^4(s+x)^8} dx$ . . . . .	277
3.36	$\int \frac{1}{(a+bs^2+2bsx+bx^2)^4} dx$ . . . . .	285
3.37	$\int \frac{1}{((a+bs^2)^2+4bs(a+bs^2)x+2b(a+3bs^2)x^2+4b^2sx^3+b^2x^4)^2} dx$ . . . . .	293
3.38	$\int \frac{1}{(a+bs^2)^4+8bs(a+bs^2)^3x+4b(a+bs^2)^2(a+7bs^2)x^2+8b^2s(3a^2+10abs^2+7b^2s^4)x^3+2b^2(3a^2+30abs^2+35b^2s^4)x^4+8b^3s(3a+7bs^2)x^5+b^4s^4x^6} dx$ . . . . .	311
3.39	$\int \left( a + \frac{d}{x^3} + \frac{c}{x^2} + \frac{b}{x} \right) dx$ . . . . .	311
3.40	$\int \left( \frac{1}{x^5} + x + x^5 \right) dx$ . . . . .	316
3.41	$\int \left( \frac{1}{x^3} + \frac{1}{x^2} + \frac{1}{x} \right) dx$ . . . . .	321
3.42	$\int \left( -\frac{2}{x^2} + \frac{3}{x} \right) dx$ . . . . .	326

---

3.43	$\int \left(-\frac{1}{7x^6} + x^6\right) dx$	331
3.44	$\int \left(1 + \frac{1}{x} + x\right) dx$	336
3.45	$\int \left(-\frac{3}{x^3} + \frac{4}{x^2}\right) dx$	341
3.46	$\int \left(\frac{1}{x} + 2x + x^2\right) dx$	346
3.47	$\int \left(x^{5/6} - x^3\right) dx$	351
3.48	$\int \left(33 + \sqrt[3]{x}\right) dx$	356
3.49	$\int \left(\frac{1}{2\sqrt{x}} + 2\sqrt{x}\right) dx$	361
3.50	$\int \left(-\frac{1}{x^2} + \frac{10}{x} + 6\sqrt{x}\right) dx$	366
3.51	$\int \left(\frac{1}{x^{3/2}} + x^{3/2}\right) dx$	371
3.52	$\int \left(-5x^{3/2} + 7x^{5/2}\right) dx$	376
3.53	$\int \left(\frac{2}{\sqrt{x}} + \sqrt{x} - \frac{x}{2}\right) dx$	381
3.54	$\int \left(-\frac{2}{x} + \frac{\sqrt{x}}{5} + x^{3/2}\right) dx$	386
<b>4</b>	<b>Appendix</b>	<b>391</b>
4.1	Listing of Grading functions	391
4.2	Links to plain text integration problems used in this report for each CAS409	

# CHAPTER 1

## INTRODUCTION

1.1	Listing of CAS systems tested . . . . .	5
1.2	Results . . . . .	6
1.3	Time and leaf size Performance . . . . .	10
1.4	Performance based on number of rules Rubi used . . . . .	12
1.5	Performance based on number of steps Rubi used . . . . .	13
1.6	Solved integrals histogram based on leaf size of result . . . . .	14
1.7	Solved integrals histogram based on CPU time used . . . . .	15
1.8	Leaf size vs. CPU time used . . . . .	16
1.9	list of integrals with no known antiderivative . . . . .	17
1.10	List of integrals solved by CAS but has no known antiderivative . . . . .	17
1.11	list of integrals solved by CAS but failed verification . . . . .	17
1.12	Timing . . . . .	18
1.13	Verification . . . . .	18
1.14	Important notes about some of the results . . . . .	19
1.15	Current tree layout of integration tests . . . . .	22
1.16	Design of the test system . . . . .	23

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [ 54 ]. This is test number [ 146 ].

## 1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

## 1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 ( 54 )	0.00 ( 0 )
Mathematica	100.00 ( 54 )	0.00 ( 0 )
Maple	100.00 ( 54 )	0.00 ( 0 )
Fricas	100.00 ( 54 )	0.00 ( 0 )
Mupad	100.00 ( 54 )	0.00 ( 0 )
Sympy	100.00 ( 54 )	0.00 ( 0 )
Giac	92.59 ( 50 )	7.41 ( 4 )
Maxima	92.59 ( 50 )	7.41 ( 4 )
Reduce	92.59 ( 50 )	7.41 ( 4 )

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> <li>1. antiderivative contains a hypergeometric function and the optimal antiderivative does not.</li> <li>2. antiderivative contains a special function and the optimal antiderivative does not.</li> <li>3. antiderivative contains the imaginary unit and the optimal antiderivative does not.</li> </ol>
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.

Table 1.2: Description of grading applied to integration result


Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.

System	% A grade	% B grade	% C grade	% F grade
Rubi	96.296	3.704	0.000	0.000
Mathematica	92.593	1.852	5.556	0.000
Maple	88.889	3.704	7.407	0.000
Giac	85.185	7.407	0.000	7.407
Sympy	75.926	24.074	0.000	0.000
Fricas	72.222	27.778	0.000	0.000
Maxima	68.519	24.074	0.000	7.407
Mupad	0.000	100.000	0.000	0.000
Reduce	0.000	92.593	0.000	7.407


Table 1.3: Antiderivative Grade distribution of each CAS



The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	0	0.00	0.00	0.00
Maple	0	0.00	0.00	0.00
Mupad	0	0.00	0.00	0.00
Sympy	0	0.00	0.00	0.00
Giac	4	100.00	0.00	0.00
Maxima	4	100.00	0.00	0.00
Reduce	4	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

### 1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Mathematica	0.02
Maxima	0.05
Maple	0.06
Fricas	0.08
Giac	0.13
Reduce	0.19
Sympy	0.19
Rubi	0.40
Mupad	8.74

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mathematica	40.87	1.00	21.00	1.00
Giac	44.50	1.34	16.00	0.82
Maple	46.13	1.21	16.50	0.84
Rubi	68.85	1.13	19.00	1.00
Sympy	76.65	1.95	20.00	0.85
Mupad	77.19	1.69	21.00	0.83
Maxima	80.04	2.69	17.50	0.85
Reduce	147.88	2.75	19.00	0.85
Fricas	172.35	2.80	24.50	0.84

Table 1.6: Leaf size performance for each CAS

## 1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the  $y$  axis is the percentage solved which Rubi itself needed the number of rules given the  $x$  axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

## 1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram shows that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

## 1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

## 1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used



## 1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

## 1.9 list of integrals with no known antiderivative

{}

## 1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

## 1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {}

Maple {}

**Maxima** Verification phase not currently implemented.

**Fricas** Verification phase not currently implemented.

**Sympy** Verification phase not currently implemented.

**Giac** Verification phase not currently implemented.

**Reduce** Verification phase not currently implemented.

**Mupad** Verification phase not currently implemented.

## 1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

## 1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

## 1.14 Important notes about some of the results

### Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

## Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

## Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at [https://ask.sagemath.org/question/57123/could-we-have-a-leaf\\_count-function-in-base-sagemath/](https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/)

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

### Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand,the_variable)
```

Which gives  $\sin(x)^2/2$

# 1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

# 1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



**High level overview of the CAS independent integration test build system**

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

*The following fields are present only in Rubi Table file*

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi  
January 13, 2024  
Design note



# CHAPTER 2

## DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS . . . . .	25
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	29
2.3	Detailed conclusion table specific for Rubi results . . . . .	43

## 2.1 List of integrals sorted by grade for each CAS

Rubi . . . . .	25
Mma . . . . .	25
Maple . . . . .	26
Fricas . . . . .	26
Maxima . . . . .	26
Giac . . . . .	27
Mupad . . . . .	27
Sympy . . . . .	27
Reduce . . . . .	28

### Rubi

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 36, 37, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54 }

**B grade** { 35, 38 }

**C grade** { }

**F normal fail** { }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

### Mma

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 23, 24, 25, 26, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54 }

**B grade** { 21 }

**C grade** { 27, 28, 29 }

**F normal fail** { }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## Maple

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 21, 22, 23, 24, 25, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54 }

**B grade** { 19, 20 }

**C grade** { 26, 27, 28, 29 }

**F normal fail** { }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## Fricas

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 25, 26, 27, 28, 29, 30, 31, 32, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54 }

**B grade** { 16, 17, 18, 19, 20, 21, 22, 23, 24, 33, 34, 35, 36, 37, 38 }

**C grade** { }

**F normal fail** { }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## Maxima

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 17, 18, 25, 30, 31, 32, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54 }

**B grade** { 16, 19, 20, 21, 22, 23, 24, 33, 34, 35, 36, 37, 38 }

**C grade** { }

**F normal fail** { 26, 27, 28, 29 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## Giac

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 17, 18, 22, 23, 24, 25, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54 }

**B grade** { 16, 19, 20, 21 }

**C grade** { }

**F normal fail** { 26, 27, 28, 29 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## Mupad

**A grade** { }

**B grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54 }

**C grade** { }

**F normal fail** { }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## Sympy

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 17, 18, 25, 26, 27, 28, 29, 30, 31, 32, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54 }

**B grade** { 16, 19, 20, 21, 22, 23, 24, 33, 34, 35, 36, 37, 38 }

**C grade** { }

**F normal fail** { }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## Reduce

**A grade** { }

**B grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54 }

**C grade** { }

**F normal fail** { 26, 27, 28, 29 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## 2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as  $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$ . To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	28	23	22	22	22	22	23	22
N.S.	1	1.00	1.00	0.82	0.79	0.79	0.79	0.79	0.82	0.79
time (sec)	N/A	0.253	0.000	0.021	0.031	0.064	0.018	0.114	0.189	0.037

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	11	11	11	8	11	10	10
N.S.	1	1.00	1.00	0.73	0.73	0.73	0.53	0.73	0.67	0.67
time (sec)	N/A	0.235	0.000	0.020	0.030	0.066	0.017	0.112	0.142	0.022

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	11	9	9	9	5	9	8	8
N.S.	1	1.00	1.00	0.82	0.82	0.82	0.45	0.82	0.73	0.73
time (sec)	N/A	0.231	0.000	0.020	0.033	0.078	0.017	0.118	0.157	0.016

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	9	8	9	9	7	9	7	7
N.S.	1	1.00	0.82	0.73	0.82	0.82	0.64	0.82	0.64	0.64
time (sec)	N/A	0.212	0.000	0.015	0.025	0.057	0.018	0.105	0.156	0.030

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	13	12	12	10	12	12	12
N.S.	1	1.00	1.00	0.93	0.86	0.86	0.71	0.86	0.86	0.86
time (sec)	N/A	0.238	0.000	0.046	0.034	0.076	0.019	0.110	0.139	0.020

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	11	10	9	9	8	9	10	10
N.S.	1	1.00	1.00	0.91	0.82	0.82	0.73	0.82	0.91	0.91
time (sec)	N/A	0.233	0.000	0.021	0.025	0.063	0.018	0.106	0.150	0.019

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	9	11	11	8	11	8	8
N.S.	1	1.00	1.00	0.60	0.73	0.73	0.53	0.73	0.53	0.53
time (sec)	N/A	0.241	0.000	0.022	0.026	0.083	0.019	0.914	0.153	0.021

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	18	14	14	14	15	14	13	13
N.S.	1	1.00	1.00	0.78	0.78	0.78	0.83	0.78	0.72	0.72
time (sec)	N/A	0.234	0.000	0.022	0.032	0.060	0.018	0.097	0.145	0.024

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	16	16	16	12	16	15	15
N.S.	1	1.00	1.00	0.80	0.80	0.80	0.60	0.80	0.75	0.75
time (sec)	N/A	0.237	0.000	0.022	0.031	0.077	0.017	0.110	0.178	0.026

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	13	12	12	10	12	15	12
N.S.	1	1.00	1.00	0.81	0.75	0.75	0.62	0.75	0.94	0.75
time (sec)	N/A	0.247	0.000	0.023	0.024	0.062	0.018	0.113	0.260	0.023

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	13	12	13	13	12	13	11	13
N.S.	1	1.00	1.00	0.92	1.00	1.00	0.92	1.00	0.85	1.00
time (sec)	N/A	0.248	0.000	0.022	0.030	0.059	0.019	0.112	0.227	0.027



Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	84	84	65	64	64	80	64	65	64
N.S.	1	1.00	1.00	0.77	0.76	0.76	0.95	0.76	0.77	0.76
time (sec)	N/A	1.447	0.005	0.041	0.029	0.061	0.031	0.109	0.211	0.113

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	63	50	49	49	60	49	50	49
N.S.	1	1.00	1.00	0.79	0.78	0.78	0.95	0.78	0.79	0.78
time (sec)	N/A	1.494	0.005	0.037	0.026	0.076	0.030	0.106	0.228	0.079

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	44	44	35	34	34	41	34	35	34
N.S.	1	1.00	1.00	0.80	0.77	0.77	0.93	0.77	0.80	0.77
time (sec)	N/A	1.053	0.002	0.035	0.031	0.067	0.027	0.109	0.216	0.025

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	25	20	19	19	22	19	20	19
N.S.	1	1.00	1.00	0.80	0.76	0.76	0.88	0.76	0.80	0.76
time (sec)	N/A	0.254	0.000	0.042	0.028	0.060	0.019	0.104	0.242	0.034

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	62	42	54	56	63	62	53	27
N.S.	1	1.00	2.00	1.35	1.74	1.81	2.03	2.00	1.71	0.87
time (sec)	N/A	0.315	0.025	0.051	0.109	0.081	0.077	0.109	0.215	0.072

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	89	103	84	89	177	104	97	296	64
N.S.	1	1.19	1.37	1.12	1.19	2.36	1.39	1.29	3.95	0.85
time (sec)	N/A	0.419	0.082	0.077	0.107	0.094	0.850	0.114	0.232	23.145

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	161	137	108	119	282	134	112	536	93
N.S.	1	1.38	1.17	0.92	1.02	2.41	1.15	0.96	4.58	0.79
time (sec)	N/A	0.563	0.108	0.092	0.107	0.110	0.918	0.110	0.214	23.173

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	164	592	163	185	163	164	163
N.S.	1	1.00	1.00	11.71	42.29	11.64	13.21	11.64	11.71	11.64
time (sec)	N/A	0.271	0.004	0.044	0.039	0.073	0.040	0.114	0.242	0.194

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	109	228	108	114	108	109	108
N.S.	1	1.00	1.00	7.79	16.29	7.71	8.14	7.71	7.79	7.71
time (sec)	N/A	0.271	0.003	0.033	0.037	0.081	0.030	0.114	0.162	0.092

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	61	13	53	53	60	53	54	53
N.S.	1	1.00	4.36	0.93	3.79	3.79	4.29	3.79	3.86	3.79
time (sec)	N/A	0.275	0.000	0.029	0.034	0.064	0.022	0.114	0.152	0.025

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	13	46	46	49	12	45	48
N.S.	1	1.00	1.00	0.93	3.29	3.29	3.50	0.86	3.21	3.43
time (sec)	N/A	0.280	0.005	0.045	0.035	0.073	0.138	0.104	0.157	0.046

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	13	101	101	109	12	100	103
N.S.	1	1.00	1.00	0.93	7.21	7.21	7.79	0.86	7.14	7.36
time (sec)	N/A	0.265	0.004	0.086	0.029	0.089	0.295	0.116	0.145	22.510

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	13	156	156	168	12	155	158
N.S.	1	1.00	1.00	0.93	11.14	11.14	12.00	0.86	11.07	11.29
time (sec)	N/A	0.266	0.005	0.148	0.037	0.074	0.452	0.111	0.147	23.377

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	38	31	30	30	29	31	30	36
N.S.	1	1.00	1.00	0.82	0.79	0.79	0.76	0.82	0.79	0.95
time (sec)	N/A	0.321	0.011	0.070	0.110	0.080	0.073	0.109	0.159	0.053

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	<b>F</b>	A	A	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	305	307	248	59	0	255	15	0	29	251
N.S.	1	1.01	0.81	0.19	0.00	0.84	0.05	0.00	0.10	0.82
time (sec)	N/A	0.977	0.341	0.033	0.000	0.104	0.090	0.000	0.145	23.007

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	<b>F</b>	A	A	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	305	307	69	59	0	255	15	0	29	251
N.S.	1	1.01	0.23	0.19	0.00	0.84	0.05	0.00	0.10	0.82
time (sec)	N/A	0.871	0.020	0.029	0.000	0.124	0.084	0.000	0.154	22.543

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	337	310	45	71	0	260	29	0	36	463
N.S.	1	0.92	0.13	0.21	0.00	0.77	0.09	0.00	0.11	1.37
time (sec)	N/A	0.947	0.020	0.045	0.000	0.088	0.094	0.000	0.156	24.949

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	337	310	83	71	0	260	29	0	36	463
N.S.	1	0.92	0.25	0.21	0.00	0.77	0.09	0.00	0.11	1.37
time (sec)	N/A	0.885	0.022	0.035	0.000	0.102	0.092	0.000	0.144	23.075

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	95	66	78	80	254	129	56	162	77
N.S.	1	1.20	0.84	0.99	1.01	3.22	1.63	0.71	2.05	0.97
time (sec)	N/A	0.304	0.049	0.141	0.108	0.078	0.213	0.113	0.145	22.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	114	66	78	80	254	129	56	162	77
N.S.	1	1.44	0.84	0.99	1.01	3.22	1.63	0.71	2.05	0.97
time (sec)	N/A	0.344	0.008	0.052	0.107	0.107	0.221	0.113	0.158	21.885

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	95	66	78	80	254	129	56	162	77
N.S.	1	1.20	0.84	0.99	1.01	3.22	1.63	0.71	2.05	0.97
time (sec)	N/A	0.350	0.007	0.053	0.111	0.119	0.209	0.121	0.141	21.966

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	109	76	149	299	970	354	148	808	91
N.S.	1	1.17	0.82	1.60	3.22	10.43	3.81	1.59	8.69	0.98
time (sec)	N/A	0.316	0.078	0.375	0.112	0.115	1.017	0.109	0.153	22.195

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	128	76	149	299	970	354	148	808	91
N.S.	1	1.38	0.82	1.60	3.22	10.43	3.81	1.59	8.69	0.98
time (sec)	N/A	0.376	0.009	0.132	0.117	0.098	0.975	0.114	0.168	22.201

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	311	76	149	299	970	354	148	808	91
N.S.	1	3.34	0.82	1.60	3.22	10.43	3.81	1.59	8.69	0.98
time (sec)	N/A	0.798	0.009	0.141	0.124	0.098	0.927	0.117	0.145	21.792

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	142	76	149	299	970	354	148	808	297
N.S.	1	1.21	0.65	1.27	2.56	8.29	3.03	1.26	6.91	2.54
time (sec)	N/A	0.455	0.007	0.273	0.117	0.106	0.956	0.113	0.152	22.064

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	128	76	149	299	970	354	148	808	297
N.S.	1	1.38	0.82	1.60	3.22	10.43	3.81	1.59	8.69	3.19
time (sec)	N/A	0.426	0.009	0.128	0.114	0.094	0.950	0.111	0.161	21.618

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	311	76	149	299	970	354	148	808	297
N.S.	1	3.34	0.82	1.60	3.22	10.43	3.81	1.59	8.69	3.19
time (sec)	N/A	0.846	0.010	0.151	0.114	0.096	0.956	0.117	0.165	21.873

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	22	21	20	27	20	21	27	20
N.S.	1	1.00	1.00	0.95	0.91	1.23	0.91	0.95	1.23	0.91
time (sec)	N/A	0.250	0.010	0.026	0.026	0.064	0.092	0.111	0.184	21.646

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	22	17	16	17	15	16	17	17
N.S.	1	1.00	1.00	0.77	0.73	0.77	0.68	0.73	0.77	0.77
time (sec)	N/A	0.238	0.002	0.024	0.035	0.067	0.043	0.116	0.209	0.030

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	13	13	17	14	14	17	11
N.S.	1	1.00	1.00	0.87	0.87	1.13	0.93	0.93	1.13	0.73
time (sec)	N/A	0.243	0.004	0.023	0.025	0.062	0.041	0.108	0.158	0.030

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	10	11	10	11	7	11	11	10
N.S.	1	1.00	1.00	1.10	1.00	1.10	0.70	1.10	1.10	1.00
time (sec)	N/A	0.239	0.003	0.022	0.030	0.065	0.043	0.114	0.366	21.890

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	12	11	12	10	11	12	12
N.S.	1	1.00	1.00	0.80	0.73	0.80	0.67	0.73	0.80	0.80
time (sec)	N/A	0.242	0.002	0.027	0.030	0.095	0.038	0.109	0.234	0.026



Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	11	10	9	9	8	10	9	9
N.S.	1	1.00	1.00	0.91	0.82	0.82	0.73	0.91	0.82	0.82
time (sec)	N/A	0.233	0.001	0.019	0.030	0.062	0.028	0.114	0.215	0.022

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	13	10	11	10	8	11	10	10
N.S.	1	1.00	1.00	0.77	0.85	0.77	0.62	0.85	0.77	0.77
time (sec)	N/A	0.238	0.004	0.023	0.025	0.095	0.035	0.114	0.237	0.025

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	13	12	11	11	10	12	11	11
N.S.	1	1.00	1.00	0.92	0.85	0.85	0.77	0.92	0.85	0.85
time (sec)	N/A	0.239	0.001	0.019	0.025	0.065	0.038	0.114	0.218	0.025

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	12	11	11	12	11	14	11
N.S.	1	1.00	1.00	0.71	0.65	0.65	0.71	0.65	0.82	0.65
time (sec)	N/A	0.230	0.003	0.063	0.024	0.066	0.020	0.117	0.232	21.816

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	13	10	9	9	10	9	8	8
N.S.	1	1.00	1.00	0.77	0.69	0.69	0.77	0.69	0.62	0.62
time (sec)	N/A	0.230	0.002	0.188	0.026	0.099	0.025	0.127	0.217	0.024

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	14	10	9	10	12	9	9	10
N.S.	1	1.00	0.93	0.67	0.60	0.67	0.80	0.60	0.60	0.67
time (sec)	N/A	0.230	0.017	0.037	0.034	0.073	0.022	0.118	0.238	0.023

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	18	14	14	18	15
N.S.	1	1.00	1.00	0.93	0.87	1.20	0.93	0.93	1.20	1.00
time (sec)	N/A	0.242	0.027	0.089	0.024	0.078	0.022	0.109	0.224	0.035

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	14	11	11	10	14	11	11	12
N.S.	1	1.00	0.82	0.65	0.65	0.59	0.82	0.65	0.65	0.71
time (sec)	N/A	0.232	0.023	0.042	0.036	0.077	0.019	0.116	0.243	21.778

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	10	9	11	14	12	11	10	8
N.S.	1	1.00	0.67	0.60	0.73	0.93	0.80	0.73	0.67	0.53
time (sec)	N/A	0.235	0.017	0.049	0.025	0.072	0.020	0.107	0.143	0.027

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	24	17	16	14	19	16	15	15
N.S.	1	1.00	1.00	0.71	0.67	0.58	0.79	0.67	0.62	0.62
time (sec)	N/A	0.239	0.006	0.067	0.026	0.066	0.021	0.114	0.154	0.031

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	23	16	15	19	20	16	17	17
N.S.	1	1.00	1.00	0.70	0.65	0.83	0.87	0.70	0.74	0.74
time (sec)	N/A	0.235	0.009	0.075	0.025	0.086	0.021	0.110	0.155	0.034

## 2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio  $\frac{\text{number of rules}}{\text{integrand size}}$  is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [26] had the largest ratio of [.818181999999999965]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	1	1	1.00	15	0.067
2	A	1	1	1.00	9	0.111
3	A	1	1	1.00	5	0.200
4	A	1	1	1.00	5	0.200
5	A	1	1	1.00	9	0.111
6	A	1	1	1.00	9	0.111
7	A	1	1	1.00	15	0.067
8	A	1	1	1.00	10	0.100
9	A	1	1	1.00	10	0.100
10	A	1	1	1.00	12	0.083
11	A	1	1	1.00	15	0.067
12	A	5	5	1.00	19	0.263
13	A	7	7	1.00	19	0.368
14	A	5	5	1.00	19	0.263
15	A	1	1	1.00	17	0.059
16	A	2	2	1.00	19	0.105
17	A	2	2	1.19	19	0.105
18	A	2	2	1.38	19	0.105
19	A	2	2	1.00	51	0.039
20	A	2	2	1.00	51	0.039
21	A	2	2	1.00	49	0.041

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	2	2	1.00	51	0.039
23	A	2	2	1.00	51	0.039
24	A	2	2	1.00	51	0.039
25	A	2	2	1.00	13	0.154
26	A	10	9	1.01	11	0.818
27	A	10	9	1.01	27	0.333
28	A	11	10	0.92	18	0.556
29	A	11	10	0.92	32	0.312
30	A	4	4	1.20	9	0.444
31	A	5	5	1.44	20	0.250
32	A	5	5	1.20	42	0.119
33	A	6	5	1.17	11	0.455
34	A	7	6	1.38	24	0.250
35	B	2	2	3.34	50	0.040
36	A	6	5	1.21	19	0.263
37	A	7	6	1.38	54	0.111
38	B	2	2	3.34	159	0.013
39	A	1	1	1.00	17	0.059
40	A	1	1	1.00	8	0.125
41	A	1	1	1.00	10	0.100
42	A	1	1	1.00	11	0.091
43	A	1	1	1.00	11	0.091
44	A	1	1	1.00	6	0.167
45	A	1	1	1.00	11	0.091
46	A	1	1	1.00	10	0.100
47	A	1	1	1.00	11	0.091
48	A	1	1	1.00	7	0.143
49	A	1	1	1.00	17	0.059
50	A	1	1	1.00	18	0.056
51	A	1	1	1.00	11	0.091
52	A	1	1	1.00	15	0.067
53	A	1	1	1.00	18	0.056

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
54	A	1	1	1.00	20	0.050

# CHAPTER 3

## LISTING OF INTEGRALS

3.1	$\int (a + bx + cx^2 + dx^3) dx$ . . . . .	48
3.2	$\int (-x^3 + x^4) dx$ . . . . .	53
3.3	$\int (-1 + x^5) dx$ . . . . .	58
3.4	$\int (7 + 4x) dx$ . . . . .	63
3.5	$\int (4x + \pi x^3) dx$ . . . . .	68
3.6	$\int (2x + 5x^2) dx$ . . . . .	73
3.7	$\int \left(\frac{x^2}{2} + \frac{x^3}{3}\right) dx$ . . . . .	78
3.8	$\int (3 - 5x + 2x^2) dx$ . . . . .	83
3.9	$\int (-2x + x^2 + x^3) dx$ . . . . .	88
3.10	$\int (1 - x^2 - 3x^5) dx$ . . . . .	93
3.11	$\int (5 + 2x + 3x^2 + 4x^3) dx$ . . . . .	98
3.12	$\int (3 - 19x^2 + 32x^4 - 16x^6)^4 dx$ . . . . .	103
3.13	$\int (3 - 19x^2 + 32x^4 - 16x^6)^3 dx$ . . . . .	110
3.14	$\int (3 - 19x^2 + 32x^4 - 16x^6)^2 dx$ . . . . .	117
3.15	$\int (3 - 19x^2 + 32x^4 - 16x^6) dx$ . . . . .	123
3.16	$\int \frac{1}{3-19x^2+32x^4-16x^6} dx$ . . . . .	128
3.17	$\int \frac{1}{(3-19x^2+32x^4-16x^6)^2} dx$ . . . . .	134
3.18	$\int \frac{1}{(3-19x^2+32x^4-16x^6)^3} dx$ . . . . .	140
3.19	$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3 dx$ . . . . .	147
3.20	$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2 dx$ . . . . .	154
3.21	$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5) dx$ . . . . .	160
3.22	$\int \frac{1}{a^5+5a^4bx+10a^3b^2x^2+10a^2b^3x^3+5ab^4x^4+b^5x^5} dx$ . . . . .	166
3.23	$\int \frac{1}{(a^5+5a^4bx+10a^3b^2x^2+10a^2b^3x^3+5ab^4x^4+b^5x^5)^2} dx$ . . . . .	172
3.24	$\int \frac{1}{(a^5+5a^4bx+10a^3b^2x^2+10a^2b^3x^3+5ab^4x^4+b^5x^5)^3} dx$ . . . . .	178
3.25	$\int \frac{1}{1+x^2+x^3+x^5} dx$ . . . . .	184
3.26	$\int \frac{1}{2+3(1+x)^5} dx$ . . . . .	189

3.27	$\int \frac{1}{5+15x+30x^2+30x^3+15x^4+3x^5} dx$	200
3.28	$\int \frac{1}{2+2(1+x)^3-3(1+x)^6} dx$	210
3.29	$\int \frac{1}{1-12x-39x^2-58x^3-45x^4-18x^5-3x^6} dx$	225
3.30	$\int \frac{1}{(a+bx^2)^4} dx$	241
3.31	$\int \frac{1}{(a^2+2abx^2+b^2x^4)^2} dx$	247
3.32	$\int \frac{1}{a^4+4a^3bx^2+6a^2b^2x^4+4ab^3x^6+b^4x^8} dx$	254
3.33	$\int \frac{1}{(a+b(s+x)^2)^4} dx$	261
3.34	$\int \frac{1}{(a^2+2ab(s+x)^2+b^2(s+x)^4)^2} dx$	269
3.35	$\int \frac{1}{a^4+4a^3b(s+x)^2+6a^2b^2(s+x)^4+4ab^3(s+x)^6+b^4(s+x)^8} dx$	277
3.36	$\int \frac{1}{(a+bs^2+2bsx+bx^2)^4} dx$	285
3.37	$\int \frac{1}{((a+bs^2)^2+4bs(a+bs^2)x+2b(a+3bs^2)x^2+4b^2sx^3+b^2x^4)^2} dx$	293
3.38	$\int \frac{1}{(a+bs^2)^4+8bs(a+bs^2)^3x+4b(a+bs^2)^2(a+7bs^2)x^2+8b^2s(3a^2+10abs^2+7b^2s^4)x^3+2b^2(3a^2+30abs^2+35b^2s^4)x^4+8b^3s(3a+7bs^2)x^5+}$	
3.39	$\int \left(a + \frac{d}{x^3} + \frac{c}{x^2} + \frac{b}{x}\right) dx$	311
3.40	$\int \left(\frac{1}{x^5} + x + x^5\right) dx$	316
3.41	$\int \left(\frac{1}{x^3} + \frac{1}{x^2} + \frac{1}{x}\right) dx$	321
3.42	$\int \left(-\frac{2}{x^2} + \frac{3}{x}\right) dx$	326
3.43	$\int \left(-\frac{1}{7x^6} + x^6\right) dx$	331
3.44	$\int \left(1 + \frac{1}{x} + x\right) dx$	336
3.45	$\int \left(-\frac{3}{x^3} + \frac{4}{x^2}\right) dx$	341
3.46	$\int \left(\frac{1}{x} + 2x + x^2\right) dx$	346
3.47	$\int \left(x^{5/6} - x^3\right) dx$	351
3.48	$\int \left(33 + \sqrt[3]{x}\right) dx$	356
3.49	$\int \left(\frac{1}{2\sqrt{x}} + 2\sqrt{x}\right) dx$	361
3.50	$\int \left(-\frac{1}{x^2} + \frac{10}{x} + 6\sqrt{x}\right) dx$	366
3.51	$\int \left(\frac{1}{x^{3/2}} + x^{3/2}\right) dx$	371
3.52	$\int \left(-5x^{3/2} + 7x^{5/2}\right) dx$	376
3.53	$\int \left(\frac{2}{\sqrt{x}} + \sqrt{x} - \frac{x}{2}\right) dx$	381
3.54	$\int \left(-\frac{2}{x} + \frac{\sqrt{x}}{5} + x^{3/2}\right) dx$	386



### 3.1 $\int (a + bx + cx^2 + dx^3) dx$

Optimal result . . . . .	48
Mathematica [A] (verified) . . . . .	48
Rubi [A] (verified) . . . . .	49
Maple [A] (verified) . . . . .	50
Fricas [A] (verification not implemented) . . . . .	50
Sympy [A] (verification not implemented) . . . . .	51
Maxima [A] (verification not implemented) . . . . .	51
Giac [A] (verification not implemented) . . . . .	51
Mupad [B] (verification not implemented) . . . . .	52
Reduce [B] (verification not implemented) . . . . .	52

#### Optimal result

Integrand size = 15, antiderivative size = 28

$$\int (a + bx + cx^2 + dx^3) dx = ax + \frac{bx^2}{2} + \frac{cx^3}{3} + \frac{dx^4}{4}$$

output

```
a*x+1/2*b*x^2+1/3*c*x^3+1/4*d*x^4
```

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int (a + bx + cx^2 + dx^3) dx = ax + \frac{bx^2}{2} + \frac{cx^3}{3} + \frac{dx^4}{4}$$

input

```
Integrate[a + b*x + c*x^2 + d*x^3,x]
```

output

```
a*x + (b*x^2)/2 + (c*x^3)/3 + (d*x^4)/4
```

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx + cx^2 + dx^3) dx$$

↓ 2009

$$ax + \frac{bx^2}{2} + \frac{cx^3}{3} + \frac{dx^4}{4}$$

input `Int[a + b*x + c*x^2 + d*x^3,x]`

output `a*x + (b*x^2)/2 + (c*x^3)/3 + (d*x^4)/4`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.82

method	result	size
gospers	$xa + \frac{1}{2}bx^2 + \frac{1}{3}cx^3 + \frac{1}{4}x^4d$	23
default	$xa + \frac{1}{2}bx^2 + \frac{1}{3}cx^3 + \frac{1}{4}x^4d$	23
norman	$xa + \frac{1}{2}bx^2 + \frac{1}{3}cx^3 + \frac{1}{4}x^4d$	23
risch	$xa + \frac{1}{2}bx^2 + \frac{1}{3}cx^3 + \frac{1}{4}x^4d$	23
parallelrisc	$xa + \frac{1}{2}bx^2 + \frac{1}{3}cx^3 + \frac{1}{4}x^4d$	23
parts	$xa + \frac{1}{2}bx^2 + \frac{1}{3}cx^3 + \frac{1}{4}x^4d$	23
orering	$\frac{x(3dx^3+4cx^2+6bx+12a)}{12}$	24

input `int(d*x^3+c*x^2+b*x+a,x,method=_RETURNVERBOSE)`output `x*a+1/2*b*x^2+1/3*c*x^3+1/4*x^4*d`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int (a + bx + cx^2 + dx^3) dx = \frac{1}{4}dx^4 + \frac{1}{3}cx^3 + \frac{1}{2}bx^2 + ax$$

input `integrate(d*x^3+c*x^2+b*x+a,x, algorithm="fricas")`output `1/4*d*x^4 + 1/3*c*x^3 + 1/2*b*x^2 + a*x`

**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int (a + bx + cx^2 + dx^3) dx = ax + \frac{bx^2}{2} + \frac{cx^3}{3} + \frac{dx^4}{4}$$

input `integrate(d*x**3+c*x**2+b*x+a,x)`output `a*x + b*x**2/2 + c*x**3/3 + d*x**4/4`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int (a + bx + cx^2 + dx^3) dx = \frac{1}{4} dx^4 + \frac{1}{3} cx^3 + \frac{1}{2} bx^2 + ax$$

input `integrate(d*x^3+c*x^2+b*x+a,x, algorithm="maxima")`output `1/4*d*x^4 + 1/3*c*x^3 + 1/2*b*x^2 + a*x`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int (a + bx + cx^2 + dx^3) dx = \frac{1}{4} dx^4 + \frac{1}{3} cx^3 + \frac{1}{2} bx^2 + ax$$

input `integrate(d*x^3+c*x^2+b*x+a,x, algorithm="giac")`output `1/4*d*x^4 + 1/3*c*x^3 + 1/2*b*x^2 + a*x`

**Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int (a + bx + cx^2 + dx^3) dx = \frac{dx^4}{4} + \frac{cx^3}{3} + \frac{bx^2}{2} + ax$$

input `int(a + b*x + c*x^2 + d*x^3,x)`

output `a*x + (b*x^2)/2 + (c*x^3)/3 + (d*x^4)/4`

**Reduce [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.82

$$\int (a + bx + cx^2 + dx^3) dx = \frac{x(3dx^3 + 4cx^2 + 6bx + 12a)}{12}$$

input `int(d*x^3+c*x^2+b*x+a,x)`

output `(x*(12*a + 6*b*x + 4*c*x**2 + 3*d*x**3))/12`

## 3.2 $\int (-x^3 + x^4) dx$

Optimal result	53
Mathematica [A] (verified)	53
Rubi [A] (verified)	54
Maple [A] (verified)	55
Fricas [A] (verification not implemented)	55
Sympy [A] (verification not implemented)	56
Maxima [A] (verification not implemented)	56
Giac [A] (verification not implemented)	56
Mupad [B] (verification not implemented)	57
Reduce [B] (verification not implemented)	57

### Optimal result

Integrand size = 9, antiderivative size = 15

$$\int (-x^3 + x^4) dx = -\frac{x^4}{4} + \frac{x^5}{5}$$

output

```
-1/4*x^4+1/5*x^5
```

### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int (-x^3 + x^4) dx = -\frac{x^4}{4} + \frac{x^5}{5}$$

input

```
Integrate[-x^3 + x^4,x]
```

output

```
-1/4*x^4 + x^5/5
```

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (x^4 - x^3) dx$$

$$\downarrow \text{2009}$$

$$\frac{x^5}{5} - \frac{x^4}{4}$$

input `Int[-x^3 + x^4,x]`

output `-1/4*x^4 + x^5/5`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

method	result	size
gospers	$\frac{x^4(4x-5)}{20}$	11
default	$-\frac{1}{4}x^4 + \frac{1}{5}x^5$	12
norman	$-\frac{1}{4}x^4 + \frac{1}{5}x^5$	12
risch	$-\frac{1}{4}x^4 + \frac{1}{5}x^5$	12
parallelrisch	$-\frac{1}{4}x^4 + \frac{1}{5}x^5$	12
parts	$-\frac{1}{4}x^4 + \frac{1}{5}x^5$	12
orering	$\frac{x(4x-5)(x^4-x^3)}{20x-20}$	23

input `int(x^4-x^3,x,method=_RETURNVERBOSE)`output `1/20*x^4*(4*x-5)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int (-x^3 + x^4) dx = \frac{1}{5}x^5 - \frac{1}{4}x^4$$

input `integrate(x^4-x^3,x, algorithm="fricas")`output `1/5*x^5 - 1/4*x^4`



**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.53

$$\int (-x^3 + x^4) dx = \frac{x^5}{5} - \frac{x^4}{4}$$

input `integrate(x**4-x**3,x)`

output `x**5/5 - x**4/4`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int (-x^3 + x^4) dx = \frac{1}{5}x^5 - \frac{1}{4}x^4$$

input `integrate(x^4-x^3,x, algorithm="maxima")`

output `1/5*x^5 - 1/4*x^4`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int (-x^3 + x^4) dx = \frac{1}{5}x^5 - \frac{1}{4}x^4$$

input `integrate(x^4-x^3,x, algorithm="giac")`

output `1/5*x^5 - 1/4*x^4`

**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.67

$$\int (-x^3 + x^4) dx = \frac{x^4(4x - 5)}{20}$$

input `int(x^4 - x^3,x)`

output `(x^4*(4*x - 5))/20`

**Reduce [B] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.67

$$\int (-x^3 + x^4) dx = \frac{x^4(4x - 5)}{20}$$

input `int(x^4-x^3,x)`

output `(x**4*(4*x - 5))/20`

### 3.3 $\int (-1 + x^5) dx$

Optimal result . . . . .	58
Mathematica [A] (verified) . . . . .	58
Rubi [A] (verified) . . . . .	59
Maple [A] (verified) . . . . .	60
Fricas [A] (verification not implemented) . . . . .	60
Sympy [A] (verification not implemented) . . . . .	61
Maxima [A] (verification not implemented) . . . . .	61
Giac [A] (verification not implemented) . . . . .	61
Mupad [B] (verification not implemented) . . . . .	62
Reduce [B] (verification not implemented) . . . . .	62

#### Optimal result

Integrand size = 5, antiderivative size = 11

$$\int (-1 + x^5) dx = -x + \frac{x^6}{6}$$

output `-x+1/6*x^6`

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int (-1 + x^5) dx = -x + \frac{x^6}{6}$$

input `Integrate[-1 + x^5,x]`

output `-x + x^6/6`

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (x^5 - 1) dx$$

↓ 2009

$$\frac{x^6}{6} - x$$

input `Int[-1 + x^5,x]`

output `-x + x^6/6`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

method	result	size
gospers	$\frac{x(x^5-6)}{6}$	9
default	$-x + \frac{1}{6}x^6$	10
norman	$-x + \frac{1}{6}x^6$	10
risch	$-x + \frac{1}{6}x^6$	10
parallelrisch	$-x + \frac{1}{6}x^6$	10
parts	$-x + \frac{1}{6}x^6$	10
orering	$\frac{x(x^5-6)(x^5-1)}{6(x-1)(x^4+x^3+x^2+x+1)}$	33

input `int(x^5-1,x,method=_RETURNVERBOSE)`

output `1/6*x*(x^5-6)`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int (-1 + x^5) dx = \frac{1}{6}x^6 - x$$

input `integrate(x^5-1,x, algorithm="fricas")`

output `1/6*x^6 - x`

**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.45

$$\int (-1 + x^5) dx = \frac{x^6}{6} - x$$

input `integrate(x**5-1,x)`

output `x**6/6 - x`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int (-1 + x^5) dx = \frac{1}{6} x^6 - x$$

input `integrate(x^5-1,x, algorithm="maxima")`

output `1/6*x^6 - x`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int (-1 + x^5) dx = \frac{1}{6} x^6 - x$$

input `integrate(x^5-1,x, algorithm="giac")`

output `1/6*x^6 - x`

**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int (-1 + x^5) dx = \frac{x(x^5 - 6)}{6}$$

input `int(x^5 - 1,x)`

output `(x*(x^5 - 6))/6`

**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int (-1 + x^5) dx = \frac{x(x^5 - 6)}{6}$$

input `int(x^5-1,x)`

output `(x*(x**5 - 6))/6`

### 3.4 $\int (7 + 4x) dx$

Optimal result . . . . .	63
Mathematica [A] (verified) . . . . .	63
Rubi [A] (verified) . . . . .	64
Maple [A] (verified) . . . . .	65
Fricas [A] (verification not implemented) . . . . .	65
Sympy [A] (verification not implemented) . . . . .	66
Maxima [A] (verification not implemented) . . . . .	66
Giac [A] (verification not implemented) . . . . .	66
Mupad [B] (verification not implemented) . . . . .	67
Reduce [B] (verification not implemented) . . . . .	67

#### Optimal result

Integrand size = 5, antiderivative size = 11

$$\int (7 + 4x) dx = \frac{1}{8}(7 + 4x)^2$$

output `1/8*(7+4*x)^2`

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int (7 + 4x) dx = 7x + 2x^2$$

input `Integrate[7 + 4*x,x]`

output `7*x + 2*x^2`



**Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (4x + 7) dx$$

$$\downarrow 17$$

$$\frac{1}{8}(4x + 7)^2$$

input `Int[7 + 4*x,x]`

output `(7 + 4*x)^2/8`

**Defintions of rubi rules used**

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] :> Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

**Maple [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

method	result	size
orering	$x(2x + 7)$	8
gosper	$2x^2 + 7x$	10
default	$2x^2 + 7x$	10
norman	$2x^2 + 7x$	10
risch	$2x^2 + 7x$	10
parallelrisch	$2x^2 + 7x$	10
parts	$2x^2 + 7x$	10

input `int(7+4*x,x,method=_RETURNVERBOSE)`

output `x*(2*x+7)`

**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int (7 + 4x) dx = 2x^2 + 7x$$

input `integrate(7+4*x,x, algorithm="fricas")`

output `2*x^2 + 7*x`

**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.64

$$\int (7 + 4x) dx = 2x^2 + 7x$$

input `integrate(7+4*x,x)`

output `2*x**2 + 7*x`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int (7 + 4x) dx = 2x^2 + 7x$$

input `integrate(7+4*x,x, algorithm="maxima")`

output `2*x^2 + 7*x`

**Giac [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int (7 + 4x) dx = 2x^2 + 7x$$

input `integrate(7+4*x,x, algorithm="giac")`

output `2*x^2 + 7*x`

**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.64

$$\int (7 + 4x) dx = x(2x + 7)$$

input `int(4*x + 7,x)`

output `x*(2*x + 7)`

**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.64

$$\int (7 + 4x) dx = x(2x + 7)$$

input `int(7+4*x,x)`

output `x*(2*x + 7)`

### 3.5 $\int (4x + \pi x^3) dx$

Optimal result	68
Mathematica [A] (verified)	68
Rubi [A] (verified)	69
Maple [A] (verified)	70
Fricas [A] (verification not implemented)	70
Sympy [A] (verification not implemented)	71
Maxima [A] (verification not implemented)	71
Giac [A] (verification not implemented)	71
Mupad [B] (verification not implemented)	72
Reduce [B] (verification not implemented)	72

#### Optimal result

Integrand size = 9, antiderivative size = 14

$$\int (4x + \pi x^3) dx = 2x^2 + \frac{\pi x^4}{4}$$

output

```
2*x^2+1/4*Pi*x^4
```

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int (4x + \pi x^3) dx = 2x^2 + \frac{\pi x^4}{4}$$

input

```
Integrate[4*x + Pi*x^3,x]
```

output

```
2*x^2 + (Pi*x^4)/4
```

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\pi x^3 + 4x) dx$$

$$\downarrow \text{2009}$$

$$\frac{\pi x^4}{4} + 2x^2$$

input `Int[4*x + Pi*x^3,x]`

output `2*x^2 + (Pi*x^4)/4`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

method	result	size
gosper	$\frac{x^2(\pi x^2+8)}{4}$	13
norman	$2x^2 + \frac{1}{4}\pi x^4$	13
risch	$2x^2 + \frac{1}{4}\pi x^4$	13
parallelrisch	$2x^2 + \frac{1}{4}\pi x^4$	13
parts	$2x^2 + \frac{1}{4}\pi x^4$	13
default	$\frac{(\pi x^2+4)^2}{4\pi}$	15
orering	$\frac{x(\pi x^2+8)(\pi x^3+4x)}{4\pi x^2+16}$	29

input `int(Pi*x^3+4*x,x,method=_RETURNVERBOSE)`output `1/4*x^2*(Pi*x^2+8)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int (4x + \pi x^3) dx = \frac{1}{4} \pi x^4 + 2x^2$$

input `integrate(pi*x^3+4*x,x, algorithm="fricas")`output `1/4*pi*x^4 + 2*x^2`

**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int (4x + \pi x^3) dx = \frac{\pi x^4}{4} + 2x^2$$

input `integrate(pi*x**3+4*x,x)`

output `pi*x**4/4 + 2*x**2`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int (4x + \pi x^3) dx = \frac{1}{4} \pi x^4 + 2x^2$$

input `integrate(pi*x^3+4*x,x, algorithm="maxima")`

output `1/4*pi*x^4 + 2*x^2`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int (4x + \pi x^3) dx = \frac{1}{4} \pi x^4 + 2x^2$$

input `integrate(pi*x^3+4*x,x, algorithm="giac")`

output `1/4*pi*x^4 + 2*x^2`



**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int (4x + \pi x^3) dx = \frac{\pi x^4}{4} + 2x^2$$

input `int(4*x + Pi*x^3,x)`

output `(Pi*x^4)/4 + 2*x^2`

**Reduce [B] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int (4x + \pi x^3) dx = \frac{x^2(\pi x^2 + 8)}{4}$$

input `int(Pi*x^3+4*x,x)`

output `(x**2*(pi*x**2 + 8))/4`

### 3.6 $\int (2x + 5x^2) dx$

Optimal result	73
Mathematica [A] (verified)	73
Rubi [A] (verified)	74
Maple [A] (verified)	75
Fricas [A] (verification not implemented)	75
Sympy [A] (verification not implemented)	76
Maxima [A] (verification not implemented)	76
Giac [A] (verification not implemented)	76
Mupad [B] (verification not implemented)	77
Reduce [B] (verification not implemented)	77

#### Optimal result

Integrand size = 9, antiderivative size = 11

$$\int (2x + 5x^2) dx = x^2 + \frac{5x^3}{3}$$

output `x^2+5/3*x^3`

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int (2x + 5x^2) dx = x^2 + \frac{5x^3}{3}$$

input `Integrate[2*x + 5*x^2,x]`

output `x^2 + (5*x^3)/3`

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (5x^2 + 2x) dx$$

↓ 2009

$$\frac{5x^3}{3} + x^2$$

input `Int [2*x + 5*x^2,x]`

output `x^2 + (5*x^3)/3`

**Defintions of rubi rules used**

rule 2009 `Int [u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

method	result	size
default	$x^2 + \frac{5}{3}x^3$	10
norman	$x^2 + \frac{5}{3}x^3$	10
risch	$x^2 + \frac{5}{3}x^3$	10
parallelrisch	$x^2 + \frac{5}{3}x^3$	10
parts	$x^2 + \frac{5}{3}x^3$	10
gosper	$\frac{x^2(5x+3)}{3}$	11
orering	$\frac{x(5x+3)(5x^2+2x)}{15x+6}$	25

input `int(5*x^2+2*x,x,method=_RETURNVERBOSE)`output `x^2+5/3*x^3`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int (2x + 5x^2) dx = \frac{5}{3}x^3 + x^2$$

input `integrate(5*x^2+2*x,x, algorithm="fricas")`output `5/3*x^3 + x^2`

**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int (2x + 5x^2) dx = \frac{5x^3}{3} + x^2$$

input `integrate(5*x**2+2*x,x)`

output `5*x**3/3 + x**2`

**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int (2x + 5x^2) dx = \frac{5}{3}x^3 + x^2$$

input `integrate(5*x^2+2*x,x, algorithm="maxima")`

output `5/3*x^3 + x^2`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int (2x + 5x^2) dx = \frac{5}{3}x^3 + x^2$$

input `integrate(5*x^2+2*x,x, algorithm="giac")`

output `5/3*x^3 + x^2`

**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

$$\int (2x + 5x^2) dx = \frac{x^2(5x + 3)}{3}$$

input `int(2*x + 5*x^2,x)`

output `(x^2*(5*x + 3))/3`

**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

$$\int (2x + 5x^2) dx = \frac{x^2(5x + 3)}{3}$$

input `int(5*x^2+2*x,x)`

output `(x**2*(5*x + 3))/3`

### 3.7 $\int \left( \frac{x^2}{2} + \frac{x^3}{3} \right) dx$

Optimal result	78
Mathematica [A] (verified)	78
Rubi [A] (verified)	79
Maple [A] (verified)	80
Fricas [A] (verification not implemented)	80
Sympy [A] (verification not implemented)	81
Maxima [A] (verification not implemented)	81
Giac [A] (verification not implemented)	81
Mupad [B] (verification not implemented)	82
Reduce [B] (verification not implemented)	82

#### Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \left( \frac{x^2}{2} + \frac{x^3}{3} \right) dx = \frac{x^3}{6} + \frac{x^4}{12}$$

output `1/6*x^3+1/12*x^4`

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \left( \frac{x^2}{2} + \frac{x^3}{3} \right) dx = \frac{x^3}{6} + \frac{x^4}{12}$$

input `Integrate[x^2/2 + x^3/3,x]`

output `x^3/6 + x^4/12`

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( \frac{x^3}{3} + \frac{x^2}{2} \right) dx$$

↓ 2009

$$\frac{x^4}{12} + \frac{x^3}{6}$$

input `Int[x^2/2 + x^3/3,x]`

output `x^3/6 + x^4/12`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`



**Maple [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.60

method	result	size
gospers	$\frac{x^3(2+x)}{12}$	9
default	$\frac{1}{6}x^3 + \frac{1}{12}x^4$	12
norman	$\frac{1}{6}x^3 + \frac{1}{12}x^4$	12
risch	$\frac{1}{6}x^3 + \frac{1}{12}x^4$	12
parallelrisch	$\frac{1}{6}x^3 + \frac{1}{12}x^4$	12
parts	$\frac{1}{6}x^3 + \frac{1}{12}x^4$	12
orering	$\frac{x(2+x)(\frac{1}{2}x^2 + \frac{1}{3}x^3)}{4x+6}$	25

input `int(1/2*x^2+1/3*x^3,x,method=_RETURNVERBOSE)`output `1/12*x^3*(2+x)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \left( \frac{x^2}{2} + \frac{x^3}{3} \right) dx = \frac{1}{12}x^4 + \frac{1}{6}x^3$$

input `integrate(1/2*x^2+1/3*x^3,x, algorithm="fricas")`output `1/12*x^4 + 1/6*x^3`

**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.53

$$\int \left( \frac{x^2}{2} + \frac{x^3}{3} \right) dx = \frac{x^4}{12} + \frac{x^3}{6}$$

input `integrate(1/2*x**2+1/3*x**3,x)`

output `x**4/12 + x**3/6`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \left( \frac{x^2}{2} + \frac{x^3}{3} \right) dx = \frac{1}{12} x^4 + \frac{1}{6} x^3$$

input `integrate(1/2*x^2+1/3*x^3,x, algorithm="maxima")`

output `1/12*x^4 + 1/6*x^3`

**Giac [A] (verification not implemented)**

Time = 0.91 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \left( \frac{x^2}{2} + \frac{x^3}{3} \right) dx = \frac{1}{12} x^4 + \frac{1}{6} x^3$$

input `integrate(1/2*x^2+1/3*x^3,x, algorithm="giac")`

output `1/12*x^4 + 1/6*x^3`

**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.53

$$\int \left( \frac{x^2}{2} + \frac{x^3}{3} \right) dx = \frac{x^3(x+2)}{12}$$

input `int(x^2/2 + x^3/3,x)`

output `(x^3*(x + 2))/12`

**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.53

$$\int \left( \frac{x^2}{2} + \frac{x^3}{3} \right) dx = \frac{x^3(x+2)}{12}$$

input `int(1/2*x^2+1/3*x^3,x)`

output `(x**3*(x + 2))/12`

### 3.8 $\int (3 - 5x + 2x^2) dx$

Optimal result . . . . .	83
Mathematica [A] (verified) . . . . .	83
Rubi [A] (verified) . . . . .	84
Maple [A] (verified) . . . . .	85
Fricas [A] (verification not implemented) . . . . .	85
Sympy [A] (verification not implemented) . . . . .	86
Maxima [A] (verification not implemented) . . . . .	86
Giac [A] (verification not implemented) . . . . .	86
Mupad [B] (verification not implemented) . . . . .	87
Reduce [B] (verification not implemented) . . . . .	87

#### Optimal result

Integrand size = 10, antiderivative size = 18

$$\int (3 - 5x + 2x^2) dx = 3x - \frac{5x^2}{2} + \frac{2x^3}{3}$$

output `3*x-5/2*x^2+2/3*x^3`

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (3 - 5x + 2x^2) dx = 3x - \frac{5x^2}{2} + \frac{2x^3}{3}$$

input `Integrate[3 - 5*x + 2*x^2,x]`

output `3*x - (5*x^2)/2 + (2*x^3)/3`

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (2x^2 - 5x + 3) dx$$

↓ 2009

$$\frac{2x^3}{3} - \frac{5x^2}{2} + 3x$$

input `Int[3 - 5*x + 2*x^2,x]`

output `3*x - (5*x^2)/2 + (2*x^3)/3`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

method	result	size
gospers	$\frac{x(4x^2-15x+18)}{6}$	14
default	$3x - \frac{5}{2}x^2 + \frac{2}{3}x^3$	15
norman	$3x - \frac{5}{2}x^2 + \frac{2}{3}x^3$	15
risch	$3x - \frac{5}{2}x^2 + \frac{2}{3}x^3$	15
parallelrisch	$3x - \frac{5}{2}x^2 + \frac{2}{3}x^3$	15
parts	$3x - \frac{5}{2}x^2 + \frac{2}{3}x^3$	15
orering	$\frac{x(4x^2-15x+18)(2x^2-5x+3)}{6(x-1)(2x-3)}$	36

input `int(2*x^2-5*x+3,x,method=_RETURNVERBOSE)`output `1/6*x*(4*x^2-15*x+18)`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int (3 - 5x + 2x^2) dx = \frac{2}{3}x^3 - \frac{5}{2}x^2 + 3x$$

input `integrate(2*x^2-5*x+3,x, algorithm="fricas")`output `2/3*x^3 - 5/2*x^2 + 3*x`

**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int (3 - 5x + 2x^2) dx = \frac{2x^3}{3} - \frac{5x^2}{2} + 3x$$

input `integrate(2*x**2-5*x+3,x)`

output `2*x**3/3 - 5*x**2/2 + 3*x`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int (3 - 5x + 2x^2) dx = \frac{2}{3}x^3 - \frac{5}{2}x^2 + 3x$$

input `integrate(2*x^2-5*x+3,x, algorithm="maxima")`

output `2/3*x^3 - 5/2*x^2 + 3*x`

**Giac [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int (3 - 5x + 2x^2) dx = \frac{2}{3}x^3 - \frac{5}{2}x^2 + 3x$$

input `integrate(2*x^2-5*x+3,x, algorithm="giac")`

output `2/3*x^3 - 5/2*x^2 + 3*x`

**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.72

$$\int (3 - 5x + 2x^2) dx = \frac{x(4x^2 - 15x + 18)}{6}$$

input `int(2*x^2 - 5*x + 3,x)`

output `(x*(4*x^2 - 15*x + 18))/6`

**Reduce [B] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.72

$$\int (3 - 5x + 2x^2) dx = \frac{x(4x^2 - 15x + 18)}{6}$$

input `int(2*x^2-5*x+3,x)`

output `(x*(4*x**2 - 15*x + 18))/6`



### 3.9 $\int (-2x + x^2 + x^3) dx$

Optimal result	88
Mathematica [A] (verified)	88
Rubi [A] (verified)	89
Maple [A] (verified)	90
Fricas [A] (verification not implemented)	90
Sympy [A] (verification not implemented)	91
Maxima [A] (verification not implemented)	91
Giac [A] (verification not implemented)	91
Mupad [B] (verification not implemented)	92
Reduce [B] (verification not implemented)	92

#### Optimal result

Integrand size = 10, antiderivative size = 20

$$\int (-2x + x^2 + x^3) dx = -x^2 + \frac{x^3}{3} + \frac{x^4}{4}$$

output

```
-x^2+1/3*x^3+1/4*x^4
```

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int (-2x + x^2 + x^3) dx = -x^2 + \frac{x^3}{3} + \frac{x^4}{4}$$

input

```
Integrate[-2*x + x^2 + x^3,x]
```

output

```
-x^2 + x^3/3 + x^4/4
```

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (x^3 + x^2 - 2x) dx$$

↓ 2009

$$\frac{x^4}{4} + \frac{x^3}{3} - x^2$$

input `Int[-2*x + x^2 + x^3,x]`

output `-x^2 + x^3/3 + x^4/4`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

method	result	size
gospers	$\frac{x^2(3x^2+4x-12)}{12}$	16
default	$-x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4$	17
norman	$-x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4$	17
risch	$-x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4$	17
parallelrisch	$-x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4$	17
parts	$-x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4$	17
orering	$\frac{x(3x^2+4x-12)(x^3+x^2-2x)}{12(2+x)(x-1)}$	34

input `int(x^3+x^2-2*x,x,method=_RETURNVERBOSE)`output `1/12*x^2*(3*x^2+4*x-12)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int (-2x + x^2 + x^3) dx = \frac{1}{4}x^4 + \frac{1}{3}x^3 - x^2$$

input `integrate(x^3+x^2-2*x,x, algorithm="fricas")`output `1/4*x^4 + 1/3*x^3 - x^2`

**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.60

$$\int (-2x + x^2 + x^3) dx = \frac{x^4}{4} + \frac{x^3}{3} - x^2$$

input `integrate(x**3+x**2-2*x,x)`

output `x**4/4 + x**3/3 - x**2`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int (-2x + x^2 + x^3) dx = \frac{1}{4}x^4 + \frac{1}{3}x^3 - x^2$$

input `integrate(x^3+x^2-2*x,x, algorithm="maxima")`

output `1/4*x^4 + 1/3*x^3 - x^2`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int (-2x + x^2 + x^3) dx = \frac{1}{4}x^4 + \frac{1}{3}x^3 - x^2$$

input `integrate(x^3+x^2-2*x,x, algorithm="giac")`

output `1/4*x^4 + 1/3*x^3 - x^2`

**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

$$\int (-2x + x^2 + x^3) dx = \frac{x^2(3x^2 + 4x - 12)}{12}$$

input `int(x^2 - 2*x + x^3,x)`

output `(x^2*(4*x + 3*x^2 - 12))/12`

**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

$$\int (-2x + x^2 + x^3) dx = \frac{x^2(3x^2 + 4x - 12)}{12}$$

input `int(x^3+x^2-2*x,x)`

output `(x**2*(3*x**2 + 4*x - 12))/12`

### 3.10 $\int (1 - x^2 - 3x^5) dx$

Optimal result	93
Mathematica [A] (verified)	93
Rubi [A] (verified)	94
Maple [A] (verified)	95
Fricas [A] (verification not implemented)	95
Sympy [A] (verification not implemented)	96
Maxima [A] (verification not implemented)	96
Giac [A] (verification not implemented)	96
Mupad [B] (verification not implemented)	97
Reduce [B] (verification not implemented)	97

#### Optimal result

Integrand size = 12, antiderivative size = 16

$$\int (1 - x^2 - 3x^5) dx = x - \frac{x^3}{3} - \frac{x^6}{2}$$

output

```
x-1/3*x^3-1/2*x^6
```

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int (1 - x^2 - 3x^5) dx = x - \frac{x^3}{3} - \frac{x^6}{2}$$

input

```
Integrate[1 - x^2 - 3*x^5,x]
```

output

```
x - x^3/3 - x^6/2
```

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (-3x^5 - x^2 + 1) dx$$

↓ 2009

$$-\frac{x^6}{2} - \frac{x^3}{3} + x$$

input `Int[1 - x^2 - 3*x^5,x]`

output `x - x^3/3 - x^6/2`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.81

method	result	size
default	$x - \frac{1}{3}x^3 - \frac{1}{2}x^6$	13
norman	$x - \frac{1}{3}x^3 - \frac{1}{2}x^6$	13
risch	$x - \frac{1}{3}x^3 - \frac{1}{2}x^6$	13
parallelrisc	$x - \frac{1}{3}x^3 - \frac{1}{2}x^6$	13
parts	$x - \frac{1}{3}x^3 - \frac{1}{2}x^6$	13
gospers	$-\frac{x(3x^5+2x^2-6)}{6}$	16
orering	$\frac{x(3x^5+2x^2-6)(-3x^5-x^2+1)}{18x^5+6x^2-6}$	40

input `int(-3*x^5-x^2+1,x,method=_RETURNVERBOSE)`output `x-1/3*x^3-1/2*x^6`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int (1 - x^2 - 3x^5) dx = -\frac{1}{2}x^6 - \frac{1}{3}x^3 + x$$

input `integrate(-3*x^5-x^2+1,x, algorithm="fricas")`output `-1/2*x^6 - 1/3*x^3 + x`



**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.62

$$\int (1 - x^2 - 3x^5) dx = -\frac{x^6}{2} - \frac{x^3}{3} + x$$

input `integrate(-3*x**5-x**2+1,x)`

output `-x**6/2 - x**3/3 + x`

**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int (1 - x^2 - 3x^5) dx = -\frac{1}{2}x^6 - \frac{1}{3}x^3 + x$$

input `integrate(-3*x^5-x^2+1,x, algorithm="maxima")`

output `-1/2*x^6 - 1/3*x^3 + x`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int (1 - x^2 - 3x^5) dx = -\frac{1}{2}x^6 - \frac{1}{3}x^3 + x$$

input `integrate(-3*x^5-x^2+1,x, algorithm="giac")`

output `-1/2*x^6 - 1/3*x^3 + x`

**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int (1 - x^2 - 3x^5) dx = -\frac{x^6}{2} - \frac{x^3}{3} + x$$

input `int(1 - 3*x^5 - x^2,x)`

output `x - x^3/3 - x^6/2`

**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int (1 - x^2 - 3x^5) dx = \frac{x(-3x^5 - 2x^2 + 6)}{6}$$

input `int(-3*x^5-x^2+1,x)`

output `(x*( - 3*x**5 - 2*x**2 + 6))/6`

### 3.11 $\int (5 + 2x + 3x^2 + 4x^3) dx$

Optimal result	98
Mathematica [A] (verified)	98
Rubi [A] (verified)	99
Maple [A] (verified)	99
Fricas [A] (verification not implemented)	100
Sympy [A] (verification not implemented)	100
Maxima [A] (verification not implemented)	101
Giac [A] (verification not implemented)	101
Mupad [B] (verification not implemented)	101
Reduce [B] (verification not implemented)	102

#### Optimal result

Integrand size = 15, antiderivative size = 13

$$\int (5 + 2x + 3x^2 + 4x^3) dx = 5x + x^2 + x^3 + x^4$$

output `x^4+x^3+x^2+5*x`

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int (5 + 2x + 3x^2 + 4x^3) dx = 5x + x^2 + x^3 + x^4$$

input `Integrate[5 + 2*x + 3*x^2 + 4*x^3,x]`

output `5*x + x^2 + x^3 + x^4`

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (4x^3 + 3x^2 + 2x + 5) dx$$

$$\downarrow \text{2009}$$

$$x^4 + x^3 + x^2 + 5x$$

input `Int[5 + 2*x + 3*x^2 + 4*x^3,x]`

output `5*x + x^2 + x^3 + x^4`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
orering	$x(x^3 + x^2 + x + 5)$	12
gosper	$x^4 + x^3 + x^2 + 5x$	14
default	$x^4 + x^3 + x^2 + 5x$	14
norman	$x^4 + x^3 + x^2 + 5x$	14
risch	$x^4 + x^3 + x^2 + 5x$	14
parallelrisch	$x^4 + x^3 + x^2 + 5x$	14
parts	$x^4 + x^3 + x^2 + 5x$	14

input `int(4*x^3+3*x^2+2*x+5,x,method=_RETURNVERBOSE)`

output `x*(x^3+x^2+x+5)`

### **Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int (5 + 2x + 3x^2 + 4x^3) dx = x^4 + x^3 + x^2 + 5x$$

input `integrate(4*x^3+3*x^2+2*x+5,x, algorithm="fricas")`

output `x^4 + x^3 + x^2 + 5*x`

### **Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

$$\int (5 + 2x + 3x^2 + 4x^3) dx = x^4 + x^3 + x^2 + 5x$$

input `integrate(4*x**3+3*x**2+2*x+5,x)`

output `x**4 + x**3 + x**2 + 5*x`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int (5 + 2x + 3x^2 + 4x^3) dx = x^4 + x^3 + x^2 + 5x$$

input `integrate(4*x^3+3*x^2+2*x+5,x, algorithm="maxima")`

output `x^4 + x^3 + x^2 + 5*x`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int (5 + 2x + 3x^2 + 4x^3) dx = x^4 + x^3 + x^2 + 5x$$

input `integrate(4*x^3+3*x^2+2*x+5,x, algorithm="giac")`

output `x^4 + x^3 + x^2 + 5*x`

**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int (5 + 2x + 3x^2 + 4x^3) dx = x^4 + x^3 + x^2 + 5x$$

input `int(2*x + 3*x^2 + 4*x^3 + 5,x)`

output `5*x + x^2 + x^3 + x^4`

**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int (5 + 2x + 3x^2 + 4x^3) dx = x(x^3 + x^2 + x + 5)$$

input `int(4*x^3+3*x^2+2*x+5,x)`

output `x*(x**3 + x**2 + x + 5)`

### 3.12 $\int (3 - 19x^2 + 32x^4 - 16x^6)^4 dx$

Optimal result . . . . .	103
Mathematica [A] (verified) . . . . .	104
Rubi [A] (verified) . . . . .	104
Maple [A] (verified) . . . . .	106
Fricas [A] (verification not implemented) . . . . .	106
Sympy [A] (verification not implemented) . . . . .	107
Maxima [A] (verification not implemented) . . . . .	107
Giac [A] (verification not implemented) . . . . .	108
Mupad [B] (verification not implemented) . . . . .	108
Reduce [B] (verification not implemented) . . . . .	109

#### Optimal result

Integrand size = 19, antiderivative size = 84

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^4 dx = 81x - 684x^3 + 4590x^5 - \frac{149700x^7}{7} + \frac{634321x^9}{9} - \frac{1841600x^{11}}{11} + \frac{3764416x^{13}}{13} - \frac{1094656x^{15}}{3} + \frac{5633536x^{17}}{17} - \frac{4014080x^{19}}{19} + \frac{1884160x^{21}}{21} - \frac{524288x^{23}}{23} + \frac{65536x^{25}}{25}$$

output

```
81*x-684*x^3+4590*x^5-149700/7*x^7+634321/9*x^9-1841600/11*x^11+3764416/13
*x^13-1094656/3*x^15+5633536/17*x^17-4014080/19*x^19+1884160/21*x^21-52428
8/23*x^23+65536/25*x^25
```



**Mathematica [A] (verified)**

Time = 0.00 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.00

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^4 dx = 81x - 684x^3 + 4590x^5 - \frac{149700x^7}{7} + \frac{634321x^9}{9} - \frac{1841600x^{11}}{11} + \frac{3764416x^{13}}{13} - \frac{1094656x^{15}}{3} + \frac{5633536x^{17}}{17} - \frac{4014080x^{19}}{19} + \frac{1884160x^{21}}{21} - \frac{524288x^{23}}{23} + \frac{65536x^{25}}{25}$$

input

```
Integrate[(3 - 19*x^2 + 32*x^4 - 16*x^6)^4, x]
```

output

```
81*x - 684*x^3 + 4590*x^5 - (149700*x^7)/7 + (634321*x^9)/9 - (1841600*x^11)/11 + (3764416*x^13)/13 - (1094656*x^15)/3 + (5633536*x^17)/17 - (4014080*x^19)/19 + (1884160*x^21)/21 - (524288*x^23)/23 + (65536*x^25)/25
```

**Rubi [A] (verified)**

Time = 1.45 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {2464, 2036, 2036, 396, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (-16x^6 + 32x^4 - 19x^2 + 3)^4 dx$$

$$\downarrow 2464$$

$$\int (x-1)^4(x+1)^4(2x-1)^4(2x+1)^4(4x^2-3)^4 dx$$

$$\downarrow 2036$$

$$\int (2x-1)^4(2x+1)^4(x^2-1)^4(4x^2-3)^4 dx$$

$$\begin{array}{c}
 \downarrow 2036 \\
 \int (x^2 - 1)^4 (4x^2 - 3)^4 (4x^2 - 1)^4 dx \\
 \downarrow 396 \\
 \int (65536x^{24} - 524288x^{22} + 1884160x^{20} - 4014080x^{18} + 5633536x^{16} - 5473280x^{14} + 3764416x^{12} - 1841600x^{10} \\
 \downarrow 2009 \\
 \frac{65536x^{25}}{25} - \frac{524288x^{23}}{23} + \frac{1884160x^{21}}{21} - \frac{4014080x^{19}}{19} + \frac{5633536x^{17}}{17} - \frac{1094656x^{15}}{15} + \\
 \frac{3764416x^{13}}{13} - \frac{1841600x^{11}}{11} + \frac{634321x^9}{9} - \frac{149700x^7}{7} + 4590x^5 - 684x^3 + 81x
 \end{array}$$

input `Int[(3 - 19*x^2 + 32*x^4 - 16*x^6)^4,x]`

output `81*x - 684*x^3 + 4590*x^5 - (149700*x^7)/7 + (634321*x^9)/9 - (1841600*x^11)/11 + (3764416*x^13)/13 - (1094656*x^15)/15 + (5633536*x^17)/17 - (4014080*x^19)/19 + (1884160*x^21)/21 - (524288*x^23)/23 + (65536*x^25)/25`

### Defintions of rubi rules used

rule 396 `Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_)*((e_) + (f_)*(x_)^2)^(r_), x_Symbol] := Int[ExpandIntegrand[(a + b*x^2)^p*(c + d*x^2)^q*(e + f*x^2)^r, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IGtQ[q, 0] && IGtQ[r, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2036 `Int[(u_)*((c_) + (d_)*(x_)^(n_))^(q_)*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*(x_)^(non2_))^(p_), x_Symbol] := Int[u*(a1*a2 + b1*b2*x^n)^p*(c + d*x^n)^q, x] /; FreeQ[{a1, b1, a2, b2, c, d, n, p, q}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))`

rule 2464

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[u*Qx^p, x] /
; !SumQ[NonfreeFactors[Qx, x]]] /; PolyQ[Px, x] && GtQ[Expon[Px, x], 2] &&
!BinomialQ[Px, x] && !TrinomialQ[Px, x] && IGtQ[p, 1]
```

**Maple [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.77

method	result
default	$81x - 684x^3 + 4590x^5 - \frac{149700}{7}x^7 + \frac{634321}{9}x^9 - \frac{1841600}{11}x^{11} + \frac{3764416}{13}x^{13} - \frac{1094656}{3}x^{15} + \frac{563353}{17}$
norman	$81x - 684x^3 + 4590x^5 - \frac{149700}{7}x^7 + \frac{634321}{9}x^9 - \frac{1841600}{11}x^{11} + \frac{3764416}{13}x^{13} - \frac{1094656}{3}x^{15} + \frac{563353}{17}$
risch	$81x - 684x^3 + 4590x^5 - \frac{149700}{7}x^7 + \frac{634321}{9}x^9 - \frac{1841600}{11}x^{11} + \frac{3764416}{13}x^{13} - \frac{1094656}{3}x^{15} + \frac{563353}{17}$
paralelrisch	$81x - 684x^3 + 4590x^5 - \frac{149700}{7}x^7 + \frac{634321}{9}x^9 - \frac{1841600}{11}x^{11} + \frac{3764416}{13}x^{13} - \frac{1094656}{3}x^{15} + \frac{563353}{17}$
gospers	$\frac{x(4386184298496x^{24} - 38140733030400x^{22} + 150122379264000x^{20} - 353491826688000x^{18} + 554471344627200x^{16} - 61052487}{17}$
orering	$\frac{x(4386184298496x^{24} - 38140733030400x^{22} + 150122379264000x^{20} - 353491826688000x^{18} + 554471344627200x^{16} - 61052487}{17}$

input

```
int((-16*x^6+32*x^4-19*x^2+3)^4,x,method=_RETURNVERBOSE)
```

output

```
81*x-684*x^3+4590*x^5-149700/7*x^7+634321/9*x^9-1841600/11*x^11+3764416/13
*x^13-1094656/3*x^15+5633536/17*x^17-4014080/19*x^19+1884160/21*x^21-52428
8/23*x^23+65536/25*x^25
```

**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.76

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^4 dx = \frac{65536}{25}x^{25} - \frac{524288}{23}x^{23} + \frac{1884160}{21}x^{21} - \frac{4014080}{19}x^{19} + \frac{5633536}{17}x^{17} - \frac{1094656}{3}x^{15} + \frac{3764416}{13}x^{13} - \frac{1841600}{11}x^{11} + \frac{634321}{9}x^9 - \frac{149700}{7}x^7 + 4590x^5 - 684x^3 + 81x$$

input `integrate((-16*x^6+32*x^4-19*x^2+3)^4,x, algorithm="fricas")`

output  $65536/25*x^{25} - 524288/23*x^{23} + 1884160/21*x^{21} - 4014080/19*x^{19} + 5633536/17*x^{17} - 1094656/3*x^{15} + 3764416/13*x^{13} - 1841600/11*x^{11} + 634321/9*x^9 - 149700/7*x^7 + 4590*x^5 - 684*x^3 + 81*x$

### Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.95

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^4 dx = \frac{65536x^{25}}{25} - \frac{524288x^{23}}{23} + \frac{1884160x^{21}}{21} - \frac{4014080x^{19}}{19} + \frac{5633536x^{17}}{17} - \frac{1094656x^{15}}{3} + \frac{3764416x^{13}}{13} - \frac{1841600x^{11}}{11} + \frac{634321x^9}{9} - \frac{149700x^7}{7} + 4590x^5 - 684x^3 + 81x$$

input `integrate((-16*x**6+32*x**4-19*x**2+3)**4,x)`

output  $65536*x^{25}/25 - 524288*x^{23}/23 + 1884160*x^{21}/21 - 4014080*x^{19}/19 + 5633536*x^{17}/17 - 1094656*x^{15}/3 + 3764416*x^{13}/13 - 1841600*x^{11}/11 + 634321*x^9/9 - 149700*x^7/7 + 4590*x^5 - 684*x^3 + 81*x$

### Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.76

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^4 dx = \frac{65536}{25} x^{25} - \frac{524288}{23} x^{23} + \frac{1884160}{21} x^{21} - \frac{4014080}{19} x^{19} + \frac{5633536}{17} x^{17} - \frac{1094656}{3} x^{15} + \frac{3764416}{13} x^{13} - \frac{1841600}{11} x^{11} + \frac{634321}{9} x^9 - \frac{149700}{7} x^7 + 4590 x^5 - 684 x^3 + 81 x$$

input `integrate((-16*x^6+32*x^4-19*x^2+3)^4,x, algorithm="maxima")`

output `65536/25*x^25 - 524288/23*x^23 + 1884160/21*x^21 - 4014080/19*x^19 + 5633536/17*x^17 - 1094656/3*x^15 + 3764416/13*x^13 - 1841600/11*x^11 + 634321/9*x^9 - 149700/7*x^7 + 4590*x^5 - 684*x^3 + 81*x`

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.76

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^4 dx = \frac{65536}{25} x^{25} - \frac{524288}{23} x^{23} + \frac{1884160}{21} x^{21} - \frac{4014080}{19} x^{19} + \frac{5633536}{17} x^{17} - \frac{1094656}{3} x^{15} + \frac{3764416}{13} x^{13} - \frac{1841600}{11} x^{11} + \frac{634321}{9} x^9 - \frac{149700}{7} x^7 + 4590 x^5 - 684 x^3 + 81 x$$

input `integrate((-16*x^6+32*x^4-19*x^2+3)^4,x, algorithm="giac")`

output `65536/25*x^25 - 524288/23*x^23 + 1884160/21*x^21 - 4014080/19*x^19 + 5633536/17*x^17 - 1094656/3*x^15 + 3764416/13*x^13 - 1841600/11*x^11 + 634321/9*x^9 - 149700/7*x^7 + 4590*x^5 - 684*x^3 + 81*x`

### Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.76

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^4 dx = \frac{65536 x^{25}}{25} - \frac{524288 x^{23}}{23} + \frac{1884160 x^{21}}{21} - \frac{4014080 x^{19}}{19} + \frac{5633536 x^{17}}{17} - \frac{1094656 x^{15}}{3} + \frac{3764416 x^{13}}{13} - \frac{1841600 x^{11}}{11} + \frac{634321 x^9}{9} - \frac{149700 x^7}{7} + 4590 x^5 - 684 x^3 + 81 x$$

input `int((19*x^2 - 32*x^4 + 16*x^6 - 3)^4,x)`

output `81*x - 684*x^3 + 4590*x^5 - (149700*x^7)/7 + (634321*x^9)/9 - (1841600*x^11)/11 + (3764416*x^13)/13 - (1094656*x^15)/3 + (5633536*x^17)/17 - (4014080*x^19)/19 + (1884160*x^21)/21 - (524288*x^23)/23 + (65536*x^25)/25`

### Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.77

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^4 dx$$

$$= \frac{x(4386184298496x^{24} - 38140733030400x^{22} + 150122379264000x^{20} - 353491826688000x^{18} + 554471344627200x^{16} - 610524871756800x^{14} + 484508289988800x^{12} - 280123520040000x^{10} + 117927076992725x^8 - 35782502827500x^6 + 7679972049750x^4 - 1144466423100x^2 + 135528918525)}{673196525}$$

input `int((-16*x^6+32*x^4-19*x^2+3)^4,x)`

output `(x*(4386184298496*x**24 - 38140733030400*x**22 + 150122379264000*x**20 - 353491826688000*x**18 + 554471344627200*x**16 - 610524871756800*x**14 + 484508289988800*x**12 - 280123520040000*x**10 + 117927076992725*x**8 - 35782502827500*x**6 + 7679972049750*x**4 - 1144466423100*x**2 + 135528918525))/673196525`

### 3.13 $\int (3 - 19x^2 + 32x^4 - 16x^6)^3 dx$

Optimal result . . . . .	110
Mathematica [A] (verified) . . . . .	110
Rubi [A] (verified) . . . . .	111
Maple [A] (verified) . . . . .	113
Fricas [A] (verification not implemented) . . . . .	113
Sympy [A] (verification not implemented) . . . . .	114
Maxima [A] (verification not implemented) . . . . .	114
Giac [A] (verification not implemented) . . . . .	115
Mupad [B] (verification not implemented) . . . . .	115
Reduce [B] (verification not implemented) . . . . .	116

#### Optimal result

Integrand size = 19, antiderivative size = 63

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^3 dx = 27x - 171x^3 + \frac{4113x^5}{5} - 2605x^7 + \frac{16448x^9}{3} - \frac{84912x^{11}}{11} + \frac{93440x^{13}}{13} - \frac{21248x^{15}}{5} + \frac{24576x^{17}}{17} - \frac{4096x^{19}}{19}$$

output

```
27*x-171*x^3+4113/5*x^5-2605*x^7+16448/3*x^9-84912/11*x^11+93440/13*x^13-21248/5*x^15+24576/17*x^17-4096/19*x^19
```

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^3 dx = 27x - 171x^3 + \frac{4113x^5}{5} - 2605x^7 + \frac{16448x^9}{3} - \frac{84912x^{11}}{11} + \frac{93440x^{13}}{13} - \frac{21248x^{15}}{5} + \frac{24576x^{17}}{17} - \frac{4096x^{19}}{19}$$

input

```
Integrate[(3 - 19*x^2 + 32*x^4 - 16*x^6)^3,x]
```

output

$$27*x - 171*x^3 + (4113*x^5)/5 - 2605*x^7 + (16448*x^9)/3 - (84912*x^{11})/11 + (93440*x^{13})/13 - (21248*x^{15})/5 + (24576*x^{17})/17 - (4096*x^{19})/19$$

**Rubi [A] (verified)**

Time = 1.49 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$ , Rules used = {2464, 25, 25, 2036, 2036, 396, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (-16x^6 + 32x^4 - 19x^2 + 3)^3 dx \\ & \quad \downarrow 2464 \\ & \int -(x-1)^3(x+1)^3(2x-1)^3(2x+1)^3(4x^2-3)^3 dx \\ & \quad \downarrow 25 \\ & - \int -(1-2x)^3(1-x)^3(x+1)^3(2x+1)^3(3-4x^2)^3 dx \\ & \quad \downarrow 25 \\ & \int (1-2x)^3(1-x)^3(x+1)^3(2x+1)^3(3-4x^2)^3 dx \\ & \quad \downarrow 2036 \\ & \int (1-x)^3(x+1)^3(1-4x^2)^3(3-4x^2)^3 dx \\ & \quad \downarrow 2036 \\ & \int (1-4x^2)^3(3-4x^2)^3(1-x^2)^3 dx \\ & \quad \downarrow 396 \\ & \int (-4096x^{18} + 24576x^{16} - 63744x^{14} + 93440x^{12} - 84912x^{10} + 49344x^8 - 18235x^6 + 4113x^4 - 513x^2 + 27) dx \\ & \quad \downarrow 2009 \end{aligned}$$



$$-\frac{4096x^{19}}{19} + \frac{24576x^{17}}{17} - \frac{21248x^{15}}{5} + \frac{93440x^{13}}{171x^3 + 27x} - \frac{84912x^{11}}{11} + \frac{16448x^9}{3} - 2605x^7 + \frac{4113x^5}{5} -$$

input `Int[(3 - 19*x^2 + 32*x^4 - 16*x^6)^3,x]`

output `27*x - 171*x^3 + (4113*x^5)/5 - 2605*x^7 + (16448*x^9)/3 - (84912*x^11)/11 + (93440*x^13)/13 - (21248*x^15)/5 + (24576*x^17)/17 - (4096*x^19)/19`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 396 `Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_)*((e_) + (f_)*(x_)^2)^(r_), x_Symbol] := Int[ExpandIntegrand[(a + b*x^2)^p*(c + d*x^2)^q*(e + f*x^2)^r, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IGtQ[q, 0] && IGtQ[r, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2036 `Int[(u_)*((c_) + (d_)*(x_)^(n_))^(q_)*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*(x_)^(non2_))^(p_), x_Symbol] := Int[u*(a1*a2 + b1*b2*x^n)^p*(c + d*x^n)^q, x] /; FreeQ[{a1, b1, a2, b2, c, d, n, p, q}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))`

rule 2464 `Int[(u_)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[u*Qx^p, x] /; !SumQ[NonfreeFactors[Qx, x]]] /; PolyQ[Px, x] && GtQ[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && IGtQ[p, 1]`

**Maple [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.79

method	result
default	$27x - 171x^3 + \frac{4113}{5}x^5 - 2605x^7 + \frac{16448}{3}x^9 - \frac{84912}{11}x^{11} + \frac{93440}{13}x^{13} - \frac{21248}{5}x^{15} + \frac{24576}{17}x^{17} - \frac{4096}{19}x^{19}$
norman	$27x - 171x^3 + \frac{4113}{5}x^5 - 2605x^7 + \frac{16448}{3}x^9 - \frac{84912}{11}x^{11} + \frac{93440}{13}x^{13} - \frac{21248}{5}x^{15} + \frac{24576}{17}x^{17} - \frac{4096}{19}x^{19}$
risch	$27x - 171x^3 + \frac{4113}{5}x^5 - 2605x^7 + \frac{16448}{3}x^9 - \frac{84912}{11}x^{11} + \frac{93440}{13}x^{13} - \frac{21248}{5}x^{15} + \frac{24576}{17}x^{17} - \frac{4096}{19}x^{19}$
paralelrisch	$27x - 171x^3 + \frac{4113}{5}x^5 - 2605x^7 + \frac{16448}{3}x^9 - \frac{84912}{11}x^{11} + \frac{93440}{13}x^{13} - \frac{21248}{5}x^{15} + \frac{24576}{17}x^{17} - \frac{4096}{19}x^{19}$
gosper	$\frac{x(149360640x^{18} - 1001594880x^{16} + 2944271616x^{14} - 4979884800x^{12} + 5348182320x^{10} - 3798583360x^8 + 1804835175x^6 - 569920x^4 + 692835x^2 - 4096)}{692835}$
orering	$\frac{x(149360640x^{18} - 1001594880x^{16} + 2944271616x^{14} - 4979884800x^{12} + 5348182320x^{10} - 3798583360x^8 + 1804835175x^6 - 569920x^4 + 692835x^2 - 4096)}{692835(2x-1)^3(2x+1)^3(4x^2-3)^3(x+1)^3(x-1)^3}$

input `int((-16*x^6+32*x^4-19*x^2+3)^3,x,method=_RETURNVERBOSE)`output `27*x-171*x^3+4113/5*x^5-2605*x^7+16448/3*x^9-84912/11*x^11+93440/13*x^13-21248/5*x^15+24576/17*x^17-4096/19*x^19`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.78

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^3 dx = -\frac{4096}{19}x^{19} + \frac{24576}{17}x^{17} - \frac{21248}{5}x^{15} + \frac{93440}{13}x^{13} - \frac{84912}{11}x^{11} + \frac{16448}{3}x^9 - 2605x^7 + \frac{4113}{5}x^5 - 171x^3 + 27x$$

input `integrate((-16*x^6+32*x^4-19*x^2+3)^3,x, algorithm="fricas")`output `-4096/19*x^19 + 24576/17*x^17 - 21248/5*x^15 + 93440/13*x^13 - 84912/11*x^11 + 16448/3*x^9 - 2605*x^7 + 4113/5*x^5 - 171*x^3 + 27*x`

**Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.95

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^3 dx = -\frac{4096x^{19}}{19} + \frac{24576x^{17}}{17} - \frac{21248x^{15}}{5} + \frac{93440x^{13}}{13} - \frac{84912x^{11}}{11} + \frac{16448x^9}{3} - 2605x^7 + \frac{4113x^5}{5} - 171x^3 + 27x$$

input `integrate((-16*x**6+32*x**4-19*x**2+3)**3,x)`output `-4096*x**19/19 + 24576*x**17/17 - 21248*x**15/5 + 93440*x**13/13 - 84912*x**11/11 + 16448*x**9/3 - 2605*x**7 + 4113*x**5/5 - 171*x**3 + 27*x`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.78

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^3 dx = -\frac{4096}{19}x^{19} + \frac{24576}{17}x^{17} - \frac{21248}{5}x^{15} + \frac{93440}{13}x^{13} - \frac{84912}{11}x^{11} + \frac{16448}{3}x^9 - 2605x^7 + \frac{4113}{5}x^5 - 171x^3 + 27x$$

input `integrate((-16*x^6+32*x^4-19*x^2+3)^3,x, algorithm="maxima")`output `-4096/19*x^19 + 24576/17*x^17 - 21248/5*x^15 + 93440/13*x^13 - 84912/11*x^11 + 16448/3*x^9 - 2605*x^7 + 4113/5*x^5 - 171*x^3 + 27*x`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.78

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^3 dx = -\frac{4096}{19}x^{19} + \frac{24576}{17}x^{17} - \frac{21248}{5}x^{15} + \frac{93440}{13}x^{13} - \frac{84912}{11}x^{11} + \frac{16448}{3}x^9 - 2605x^7 + \frac{4113}{5}x^5 - 171x^3 + 27x$$

input `integrate((-16*x^6+32*x^4-19*x^2+3)^3,x, algorithm="giac")`output `-4096/19*x^19 + 24576/17*x^17 - 21248/5*x^15 + 93440/13*x^13 - 84912/11*x^11 + 16448/3*x^9 - 2605*x^7 + 4113/5*x^5 - 171*x^3 + 27*x`**Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.78

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^3 dx = -\frac{4096x^{19}}{19} + \frac{24576x^{17}}{17} - \frac{21248x^{15}}{5} + \frac{93440x^{13}}{13} - \frac{84912x^{11}}{11} + \frac{16448x^9}{3} - 2605x^7 + \frac{4113x^5}{5} - 171x^3 + 27x$$

input `int(-(19*x^2 - 32*x^4 + 16*x^6 - 3)^3,x)`output `27*x - 171*x^3 + (4113*x^5)/5 - 2605*x^7 + (16448*x^9)/3 - (84912*x^11)/11 + (93440*x^13)/13 - (21248*x^15)/5 + (24576*x^17)/17 - (4096*x^19)/19`

**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.79

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^3 dx$$
$$= \frac{x(-149360640x^{18} + 1001594880x^{16} - 2944271616x^{14} + 4979884800x^{12} - 5348182320x^{10} + 3798583360x^8 - 1804835175x^6 + 569926071x^4 - 118474785x^2 + 18706545)}{692835}$$

input `int((-16*x^6+32*x^4-19*x^2+3)^3,x)`

output `(x*( - 149360640*x**18 + 1001594880*x**16 - 2944271616*x**14 + 4979884800*x**12 - 5348182320*x**10 + 3798583360*x**8 - 1804835175*x**6 + 569926071*x**4 - 118474785*x**2 + 18706545))/692835`

### 3.14 $\int (3 - 19x^2 + 32x^4 - 16x^6)^2 dx$

Optimal result . . . . .	117
Mathematica [A] (verified) . . . . .	117
Rubi [A] (verified) . . . . .	118
Maple [A] (verified) . . . . .	119
Fricas [A] (verification not implemented) . . . . .	120
Sympy [A] (verification not implemented) . . . . .	120
Maxima [A] (verification not implemented) . . . . .	121
Giac [A] (verification not implemented) . . . . .	121
Mupad [B] (verification not implemented) . . . . .	121
Reduce [B] (verification not implemented) . . . . .	122

#### Optimal result

Integrand size = 19, antiderivative size = 44

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^2 dx = 9x - 38x^3 + \frac{553x^5}{5} - \frac{1312x^7}{7} + \frac{544x^9}{3} - \frac{1024x^{11}}{11} + \frac{256x^{13}}{13}$$

output

```
9*x-38*x^3+553/5*x^5-1312/7*x^7+544/3*x^9-1024/11*x^11+256/13*x^13
```

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^2 dx = 9x - 38x^3 + \frac{553x^5}{5} - \frac{1312x^7}{7} + \frac{544x^9}{3} - \frac{1024x^{11}}{11} + \frac{256x^{13}}{13}$$

input

```
Integrate[(3 - 19*x^2 + 32*x^4 - 16*x^6)^2,x]
```

output

```
9*x - 38*x^3 + (553*x^5)/5 - (1312*x^7)/7 + (544*x^9)/3 - (1024*x^11)/11 +
(256*x^13)/13
```

**Rubi [A] (verified)**

Time = 1.05 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {2464, 2036, 2036, 396, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (-16x^6 + 32x^4 - 19x^2 + 3)^2 dx \\
 & \quad \downarrow \text{2464} \\
 & \int (x-1)^2(x+1)^2(2x-1)^2(2x+1)^2(4x^2-3)^2 dx \\
 & \quad \downarrow \text{2036} \\
 & \int (2x-1)^2(2x+1)^2(x^2-1)^2(4x^2-3)^2 dx \\
 & \quad \downarrow \text{2036} \\
 & \int (x^2-1)^2(4x^2-3)^2(4x^2-1)^2 dx \\
 & \quad \downarrow \text{396} \\
 & \int (256x^{12} - 1024x^{10} + 1632x^8 - 1312x^6 + 553x^4 - 114x^2 + 9) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{256x^{13}}{13} - \frac{1024x^{11}}{11} + \frac{544x^9}{3} - \frac{1312x^7}{7} + \frac{553x^5}{5} - 38x^3 + 9x
 \end{aligned}$$

input

```
Int[(3 - 19*x^2 + 32*x^4 - 16*x^6)^2,x]
```

output

```
9*x - 38*x^3 + (553*x^5)/5 - (1312*x^7)/7 + (544*x^9)/3 - (1024*x^11)/11 + (256*x^13)/13
```

## Defintions of rubi rules used

rule 396

```
Int[((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.)*((e_) + (f_.)*(
x_)^2)^(r_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x^2)^p*(c + d*x^2)^q*
(e + f*x^2)^r, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IGtQ
[q, 0] && IGtQ[r, 0]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2036

```
Int[(u_.)*((c_) + (d_.)*(x_)^(n_.))^(q_.)*((a1_) + (b1_.)*(x_)^(non2_.))^(p
_.)*((a2_) + (b2_.)*(x_)^(non2_.))^(p_.), x_Symbol] := Int[u*(a1*a2 + b1*b2
*x^n)^p*(c + d*x^n)^q, x] /; FreeQ[{a1, b1, a2, b2, c, d, n, p, q}, x] && E
qQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && Gt
Q[a2, 0]))
```

rule 2464

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[u*Qx^p, x] /
; !SumQ[NonfreeFactors[Qx, x]]] /; PolyQ[Px, x] && GtQ[Expon[Px, x], 2] &&
!BinomialQ[Px, x] && !TrinomialQ[Px, x] && IGtQ[p, 1]
```

## Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.80

method	result	size
default	$9x - 38x^3 + \frac{553}{5}x^5 - \frac{1312}{7}x^7 + \frac{544}{3}x^9 - \frac{1024}{11}x^{11} + \frac{256}{13}x^{13}$	35
norman	$9x - 38x^3 + \frac{553}{5}x^5 - \frac{1312}{7}x^7 + \frac{544}{3}x^9 - \frac{1024}{11}x^{11} + \frac{256}{13}x^{13}$	35
risch	$9x - 38x^3 + \frac{553}{5}x^5 - \frac{1312}{7}x^7 + \frac{544}{3}x^9 - \frac{1024}{11}x^{11} + \frac{256}{13}x^{13}$	35
parallelrisch	$9x - 38x^3 + \frac{553}{5}x^5 - \frac{1312}{7}x^7 + \frac{544}{3}x^9 - \frac{1024}{11}x^{11} + \frac{256}{13}x^{13}$	35
gospers	$\frac{x(295680x^{12} - 1397760x^{10} + 2722720x^8 - 2814240x^6 + 1660659x^4 - 570570x^2 + 135135)}{15015}$	36
orering	$\frac{x(295680x^{12} - 1397760x^{10} + 2722720x^8 - 2814240x^6 + 1660659x^4 - 570570x^2 + 135135)(-16x^6 + 32x^4 - 19x^2 + 3)^2}{15015(x-1)^2(x+1)^2(4x^2-3)^2(2x+1)^2(2x-1)^2}$	88

input

```
int((-16*x^6+32*x^4-19*x^2+3)^2,x,method=_RETURNVERBOSE)
```



output  $9*x-38*x^3+553/5*x^5-1312/7*x^7+544/3*x^9-1024/11*x^11+256/13*x^13$

### Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.77

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^2 dx = \frac{256}{13} x^{13} - \frac{1024}{11} x^{11} + \frac{544}{3} x^9 - \frac{1312}{7} x^7 + \frac{553}{5} x^5 - 38x^3 + 9x$$

input `integrate((-16*x^6+32*x^4-19*x^2+3)^2,x, algorithm="fricas")`

output  $256/13*x^13 - 1024/11*x^11 + 544/3*x^9 - 1312/7*x^7 + 553/5*x^5 - 38*x^3 + 9*x$

### Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.93

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^2 dx = \frac{256x^{13}}{13} - \frac{1024x^{11}}{11} + \frac{544x^9}{3} - \frac{1312x^7}{7} + \frac{553x^5}{5} - 38x^3 + 9x$$

input `integrate((-16*x**6+32*x**4-19*x**2+3)**2,x)`

output  $256*x**13/13 - 1024*x**11/11 + 544*x**9/3 - 1312*x**7/7 + 553*x**5/5 - 38*x**3 + 9*x$

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.77

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^2 dx = \frac{256}{13} x^{13} - \frac{1024}{11} x^{11} + \frac{544}{3} x^9 - \frac{1312}{7} x^7 + \frac{553}{5} x^5 - 38x^3 + 9x$$

input `integrate((-16*x^6+32*x^4-19*x^2+3)^2,x, algorithm="maxima")`output `256/13*x^13 - 1024/11*x^11 + 544/3*x^9 - 1312/7*x^7 + 553/5*x^5 - 38*x^3 + 9*x`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.77

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^2 dx = \frac{256}{13} x^{13} - \frac{1024}{11} x^{11} + \frac{544}{3} x^9 - \frac{1312}{7} x^7 + \frac{553}{5} x^5 - 38x^3 + 9x$$

input `integrate((-16*x^6+32*x^4-19*x^2+3)^2,x, algorithm="giac")`output `256/13*x^13 - 1024/11*x^11 + 544/3*x^9 - 1312/7*x^7 + 553/5*x^5 - 38*x^3 + 9*x`**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.77

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^2 dx = \frac{256 x^{13}}{13} - \frac{1024 x^{11}}{11} + \frac{544 x^9}{3} - \frac{1312 x^7}{7} + \frac{553 x^5}{5} - 38 x^3 + 9 x$$

input `int((19*x^2 - 32*x^4 + 16*x^6 - 3)^2,x)`

output  $9x - 38x^3 + \frac{553x^5}{5} - \frac{1312x^7}{7} + \frac{544x^9}{3} - \frac{1024x^{11}}{11} + \frac{256x^{13}}{13}$

### Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.80

$$\int (3 - 19x^2 + 32x^4 - 16x^6)^2 dx$$

$$= \frac{x(295680x^{12} - 1397760x^{10} + 2722720x^8 - 2814240x^6 + 1660659x^4 - 570570x^2 + 135135)}{15015}$$

input `int((-16*x^6+32*x^4-19*x^2+3)^2,x)`

output  $(x(295680x^{12} - 1397760x^{10} + 2722720x^8 - 2814240x^6 + 1660659x^4 - 570570x^2 + 135135))/15015$

### 3.15 $\int (3 - 19x^2 + 32x^4 - 16x^6) dx$

Optimal result	123
Mathematica [A] (verified)	123
Rubi [A] (verified)	124
Maple [A] (verified)	125
Fricas [A] (verification not implemented)	125
Sympy [A] (verification not implemented)	126
Maxima [A] (verification not implemented)	126
Giac [A] (verification not implemented)	126
Mupad [B] (verification not implemented)	127
Reduce [B] (verification not implemented)	127

#### Optimal result

Integrand size = 17, antiderivative size = 25

$$\int (3 - 19x^2 + 32x^4 - 16x^6) dx = 3x - \frac{19x^3}{3} + \frac{32x^5}{5} - \frac{16x^7}{7}$$

output

```
3*x-19/3*x^3+32/5*x^5-16/7*x^7
```

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int (3 - 19x^2 + 32x^4 - 16x^6) dx = 3x - \frac{19x^3}{3} + \frac{32x^5}{5} - \frac{16x^7}{7}$$

input

```
Integrate[3 - 19*x^2 + 32*x^4 - 16*x^6, x]
```

output

```
3*x - (19*x^3)/3 + (32*x^5)/5 - (16*x^7)/7
```

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (-16x^6 + 32x^4 - 19x^2 + 3) dx$$

$$\downarrow \text{2009}$$

$$-\frac{16x^7}{7} + \frac{32x^5}{5} - \frac{19x^3}{3} + 3x$$

input `Int[3 - 19*x^2 + 32*x^4 - 16*x^6,x]`

output `3*x - (19*x^3)/3 + (32*x^5)/5 - (16*x^7)/7`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

method	result	size
default	$3x - \frac{19}{3}x^3 + \frac{32}{5}x^5 - \frac{16}{7}x^7$	20
norman	$3x - \frac{19}{3}x^3 + \frac{32}{5}x^5 - \frac{16}{7}x^7$	20
risch	$3x - \frac{19}{3}x^3 + \frac{32}{5}x^5 - \frac{16}{7}x^7$	20
parallelrisch	$3x - \frac{19}{3}x^3 + \frac{32}{5}x^5 - \frac{16}{7}x^7$	20
parts	$3x - \frac{19}{3}x^3 + \frac{32}{5}x^5 - \frac{16}{7}x^7$	20
gospers	$-\frac{x(240x^6 - 672x^4 + 665x^2 - 315)}{105}$	21
orering	$\frac{x(240x^6 - 672x^4 + 665x^2 - 315)(-16x^6 + 32x^4 - 19x^2 + 3)}{105(x-1)(x+1)(4x^2-3)(2x+1)(2x-1)}$	71

input `int(-16*x^6+32*x^4-19*x^2+3,x,method=_RETURNVERBOSE)`output `3*x-19/3*x^3+32/5*x^5-16/7*x^7`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int (3 - 19x^2 + 32x^4 - 16x^6) dx = -\frac{16}{7}x^7 + \frac{32}{5}x^5 - \frac{19}{3}x^3 + 3x$$

input `integrate(-16*x^6+32*x^4-19*x^2+3,x, algorithm="fricas")`output `-16/7*x^7 + 32/5*x^5 - 19/3*x^3 + 3*x`

**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int (3 - 19x^2 + 32x^4 - 16x^6) dx = -\frac{16x^7}{7} + \frac{32x^5}{5} - \frac{19x^3}{3} + 3x$$

input `integrate(-16*x**6+32*x**4-19*x**2+3,x)`

output `-16*x**7/7 + 32*x**5/5 - 19*x**3/3 + 3*x`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int (3 - 19x^2 + 32x^4 - 16x^6) dx = -\frac{16}{7}x^7 + \frac{32}{5}x^5 - \frac{19}{3}x^3 + 3x$$

input `integrate(-16*x^6+32*x^4-19*x^2+3,x, algorithm="maxima")`

output `-16/7*x^7 + 32/5*x^5 - 19/3*x^3 + 3*x`

**Giac [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int (3 - 19x^2 + 32x^4 - 16x^6) dx = -\frac{16}{7}x^7 + \frac{32}{5}x^5 - \frac{19}{3}x^3 + 3x$$

input `integrate(-16*x^6+32*x^4-19*x^2+3,x, algorithm="giac")`

output `-16/7*x^7 + 32/5*x^5 - 19/3*x^3 + 3*x`

**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int (3 - 19x^2 + 32x^4 - 16x^6) dx = -\frac{16x^7}{7} + \frac{32x^5}{5} - \frac{19x^3}{3} + 3x$$

input `int(32*x^4 - 19*x^2 - 16*x^6 + 3,x)`

output `3*x - (19*x^3)/3 + (32*x^5)/5 - (16*x^7)/7`

**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int (3 - 19x^2 + 32x^4 - 16x^6) dx = \frac{x(-240x^6 + 672x^4 - 665x^2 + 315)}{105}$$

input `int(-16*x^6+32*x^4-19*x^2+3,x)`

output `(x*( - 240*x**6 + 672*x**4 - 665*x**2 + 315))/105`



### 3.16 $\int \frac{1}{3-19x^2+32x^4-16x^6} dx$

Optimal result	128
Mathematica [A] (verified)	128
Rubi [A] (verified)	129
Maple [A] (verified)	130
Fricas [B] (verification not implemented)	130
Sympy [B] (verification not implemented)	131
Maxima [B] (verification not implemented)	131
Giac [B] (verification not implemented)	132
Mupad [B] (verification not implemented)	132
Reduce [B] (verification not implemented)	133

#### Optimal result

Integrand size = 19, antiderivative size = 31

$$\int \frac{1}{3-19x^2+32x^4-16x^6} dx = \frac{\operatorname{arctanh}(x)}{3} + \frac{1}{3}\operatorname{arctanh}(2x) - \frac{\operatorname{arctanh}\left(\frac{2x}{\sqrt{3}}\right)}{\sqrt{3}}$$

output

```
1/3*arctanh(x)+1/3*arctanh(2*x)-1/3*arctanh(2/3*x*3^(1/2))*3^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 62, normalized size of antiderivative = 2.00

$$\int \frac{1}{3-19x^2+32x^4-16x^6} dx = \frac{1}{6} \left( \sqrt{3} \log(\sqrt{3}-2x) - \sqrt{3} \log(\sqrt{3}+2x) - \log(1-3x+2x^2) + \log(1+3x+2x^2) \right)$$

input

```
Integrate[(3 - 19*x^2 + 32*x^4 - 16*x^6)^(-1), x]
```

output

```
(Sqrt[3]*Log[Sqrt[3] - 2*x] - Sqrt[3]*Log[Sqrt[3] + 2*x] - Log[1 - 3*x + 2*x^2] + Log[1 + 3*x + 2*x^2])/6
```

**Rubi [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$ , Rules used = {2460, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{-16x^6 + 32x^4 - 19x^2 + 3} dx$$

$$\downarrow 2460$$

$$\int \left( \frac{2}{4x^2 - 3} - \frac{2}{3(4x^2 - 1)} - \frac{1}{3(x^2 - 1)} \right) dx$$

$$\downarrow 2009$$

$$\frac{\operatorname{arctanh}(x)}{3} + \frac{1}{3} \operatorname{arctanh}(2x) - \frac{\operatorname{arctanh}\left(\frac{2x}{\sqrt{3}}\right)}{\sqrt{3}}$$

input `Int[(3 - 19*x^2 + 32*x^4 - 16*x^6)^(-1),x]`

output `ArcTanh[x]/3 + ArcTanh[2*x]/3 - ArcTanh[(2*x)/Sqrt[3]]/Sqrt[3]`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2460 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px /. x -> Sqrt[x]]}, Int[ExpandIntegrand[u*(Qx /. x -> x^2)^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x^2] && GtQ[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0] && RationalFunctionQ[u, x]`

**Maple [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.35

method	result	size
default	$-\frac{\ln(x-1)}{6} - \frac{\operatorname{arctanh}\left(\frac{2\sqrt{3}x}{3}\right)\sqrt{3}}{3} + \frac{\ln(2x+1)}{6} + \frac{\ln(x+1)}{6} - \frac{\ln(2x-1)}{6}$	42
risch	$\frac{\sqrt{3} \ln(-\sqrt{3}+2x)}{6} - \frac{\sqrt{3} \ln(\sqrt{3}+2x)}{6} + \frac{\ln(2x^2+3x+1)}{6} - \frac{\ln(2x^2-3x+1)}{6}$	56

input `int(1/(-16*x^6+32*x^4-19*x^2+3),x,method=_RETURNVERBOSE)`

output `-1/6*ln(x-1)-1/3*arctanh(2/3*3^(1/2)*x)*3^(1/2)+1/6*ln(2*x+1)+1/6*ln(x+1)-1/6*ln(2*x-1)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 56 vs. 2(23) = 46.

Time = 0.08 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.81

$$\int \frac{1}{3 - 19x^2 + 32x^4 - 16x^6} dx = \frac{1}{6} \sqrt{3} \log \left( \frac{4x^2 - 4\sqrt{3}x + 3}{4x^2 - 3} \right) + \frac{1}{6} \log(2x^2 + 3x + 1) - \frac{1}{6} \log(2x^2 - 3x + 1)$$

input `integrate(1/(-16*x^6+32*x^4-19*x^2+3),x, algorithm="fricas")`

output `1/6*sqrt(3)*log((4*x^2 - 4*sqrt(3)*x + 3)/(4*x^2 - 3)) + 1/6*log(2*x^2 + 3*x + 1) - 1/6*log(2*x^2 - 3*x + 1)`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 63 vs.  $2(29) = 58$ .

Time = 0.08 (sec) , antiderivative size = 63, normalized size of antiderivative = 2.03

$$\int \frac{1}{3 - 19x^2 + 32x^4 - 16x^6} dx = \frac{\sqrt{3} \log\left(x - \frac{\sqrt{3}}{2}\right)}{6} - \frac{\sqrt{3} \log\left(x + \frac{\sqrt{3}}{2}\right)}{6} - \frac{\log\left(x^2 - \frac{3x}{2} + \frac{1}{2}\right)}{6} + \frac{\log\left(x^2 + \frac{3x}{2} + \frac{1}{2}\right)}{6}$$

input `integrate(1/(-16*x**6+32*x**4-19*x**2+3),x)`

output `sqrt(3)*log(x - sqrt(3)/2)/6 - sqrt(3)*log(x + sqrt(3)/2)/6 - log(x**2 - 3*x/2 + 1/2)/6 + log(x**2 + 3*x/2 + 1/2)/6`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 54 vs.  $2(23) = 46$ .

Time = 0.11 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.74

$$\int \frac{1}{3 - 19x^2 + 32x^4 - 16x^6} dx = \frac{1}{6} \sqrt{3} \log\left(\frac{2x - \sqrt{3}}{2x + \sqrt{3}}\right) + \frac{1}{6} \log(2x + 1) - \frac{1}{6} \log(2x - 1) + \frac{1}{6} \log(x + 1) - \frac{1}{6} \log(x - 1)$$

input `integrate(1/(-16*x^6+32*x^4-19*x^2+3),x, algorithm="maxima")`

output `1/6*sqrt(3)*log((2*x - sqrt(3))/(2*x + sqrt(3))) + 1/6*log(2*x + 1) - 1/6*log(2*x - 1) + 1/6*log(x + 1) - 1/6*log(x - 1)`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 62 vs.  $2(23) = 46$ .

Time = 0.11 (sec) , antiderivative size = 62, normalized size of antiderivative = 2.00

$$\int \frac{1}{3 - 19x^2 + 32x^4 - 16x^6} dx = \frac{1}{6} \sqrt{3} \log \left( \frac{|8x - 4\sqrt{3}|}{|8x + 4\sqrt{3}|} \right) + \frac{1}{6} \log(|2x + 1|) - \frac{1}{6} \log(|2x - 1|) + \frac{1}{6} \log(|x + 1|) - \frac{1}{6} \log(|x - 1|)$$

input `integrate(1/(-16*x^6+32*x^4-19*x^2+3),x, algorithm="giac")`

output `1/6*sqrt(3)*log(abs(8*x - 4*sqrt(3))/abs(8*x + 4*sqrt(3))) + 1/6*log(abs(2*x + 1)) - 1/6*log(abs(2*x - 1)) + 1/6*log(abs(x + 1)) - 1/6*log(abs(x - 1))`

**Mupad [B] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.87

$$\int \frac{1}{3 - 19x^2 + 32x^4 - 16x^6} dx = \frac{\operatorname{atanh}\left(\frac{x}{4608\left(\frac{x^2}{6912} + \frac{1}{13824}\right)}\right)}{3} - \frac{\sqrt{3} \operatorname{atanh}\left(\frac{2\sqrt{3}x}{3}\right)}{3}$$

input `int(-1/(19*x^2 - 32*x^4 + 16*x^6 - 3),x)`

output `atanh(x/(4608*(x^2/6912 + 1/13824)))/3 - (3^(1/2)*atanh((2*3^(1/2)*x)/3))/3`

**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.71

$$\int \frac{1}{3 - 19x^2 + 32x^4 - 16x^6} dx = \frac{\sqrt{3} \log(-\sqrt{3} + 2x)}{6} - \frac{\sqrt{3} \log(\sqrt{3} + 2x)}{6} - \frac{\log(2x - 1)}{6} \\ + \frac{\log(2x + 1)}{6} - \frac{\log(x - 1)}{6} + \frac{\log(x + 1)}{6}$$

input `int(1/(-16*x^6+32*x^4-19*x^2+3),x)`

output `(sqrt(3)*log(-sqrt(3)+2*x)-sqrt(3)*log(sqrt(3)+2*x)-log(2*x-1)+log(2*x+1)-log(x-1)+log(x+1))/6`

**3.17**  $\int \frac{1}{(3-19x^2+32x^4-16x^6)^2} dx$

Optimal result	134
Mathematica [A] (verified)	134
Rubi [A] (verified)	135
Maple [A] (verified)	136
Fricas [B] (verification not implemented)	136
Sympy [A] (verification not implemented)	137
Maxima [A] (verification not implemented)	137
Giac [A] (verification not implemented)	138
Mupad [B] (verification not implemented)	138
Reduce [B] (verification not implemented)	139

**Optimal result**

Integrand size = 19, antiderivative size = 75

$$\int \frac{1}{(3-19x^2+32x^4-16x^6)^2} dx = \frac{2x}{9(1-4x^2)} + \frac{2x}{3(3-4x^2)} + \frac{x}{18(1-x^2)} + \frac{67\operatorname{arctanh}(x)}{54} - \frac{7}{27}\operatorname{arctanh}(2x) - \frac{5\operatorname{arctanh}\left(\frac{2x}{\sqrt{3}}\right)}{3\sqrt{3}}$$

output

```
2*x/(-36*x^2+9)+2*x/(-12*x^2+9)+x/(-18*x^2+18)+67/54*arctanh(x)-7/27*arctanh(2*x)-5/9*arctanh(2/3*x*3^(1/2))*3^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.37

$$\int \frac{1}{(3-19x^2+32x^4-16x^6)^2} dx = \frac{1}{108} \left( -\frac{6x(27-104x^2+80x^4)}{-3+19x^2-32x^4+16x^6} + 14\log(1-2x) + 30\sqrt{3}\log(\sqrt{3}-2x) - 67\log(1-x) + 67\log(1+x) - 14\log(1+2x) - 30\sqrt{3}\log(\sqrt{3}+2x) \right)$$

input `Integrate[(3 - 19*x^2 + 32*x^4 - 16*x^6)^(-2), x]`

output `((-6*x*(27 - 104*x^2 + 80*x^4))/(-3 + 19*x^2 - 32*x^4 + 16*x^6) + 14*Log[1 - 2*x] + 30*Sqrt[3]*Log[Sqrt[3] - 2*x] - 67*Log[1 - x] + 67*Log[1 + x] - 14*Log[1 + 2*x] - 30*Sqrt[3]*Log[Sqrt[3] + 2*x])/108`

### Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.19, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$ , Rules used = {2460, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(-16x^6 + 32x^4 - 19x^2 + 3)^2} dx$$

↓ 2460

$$\int \left( \frac{4}{4x^2 - 3} + \frac{14}{27(4x^2 - 1)} + \frac{4}{(4x^2 - 3)^2} - \frac{67}{54(x^2 - 1)} + \frac{1}{36(x - 1)^2} + \frac{1}{36(x + 1)^2} + \frac{1}{9(2x - 1)^2} + \frac{1}{9(2x + 1)^2} \right) dx$$

↓ 2009

$$\frac{67 \operatorname{arctanh}(x)}{54} - \frac{7}{27} \operatorname{arctanh}(2x) - \frac{5 \operatorname{arctanh}\left(\frac{2x}{\sqrt{3}}\right)}{3\sqrt{3}} + \frac{2x}{3(3 - 4x^2)} + \frac{1}{18(1 - 2x)} + \frac{1}{36(1 - x)} - \frac{1}{36(x + 1)} - \frac{1}{18(2x + 1)}$$

input `Int[(3 - 19*x^2 + 32*x^4 - 16*x^6)^(-2), x]`

output `1/(18*(1 - 2*x)) + 1/(36*(1 - x)) - 1/(36*(1 + x)) - 1/(18*(1 + 2*x)) + (2*x)/(3*(3 - 4*x^2)) + (67*ArcTanh[x])/54 - (7*ArcTanh[2*x])/27 - (5*ArcTanh[(2*x)/Sqrt[3]])/(3*Sqrt[3])`



### Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2460 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px /. x -> Sqrt[x]]}, Int[ExpandIntegrand[u*(Qx /. x -> x^2)^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]]] /; PolyQ[Px, x^2] && GtQ[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0] && RationalFunctionQ[u, x]`

### Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.12

method	result
default	$-\frac{1}{36(x-1)} - \frac{67\ln(x-1)}{108} - \frac{x}{6(x^2-\frac{3}{4})} - \frac{5 \operatorname{arctanh}\left(\frac{2\sqrt{3}x}{3}\right)\sqrt{3}}{9} - \frac{1}{18(2x+1)} - \frac{7\ln(2x+1)}{54} - \frac{1}{36(x+1)} + \frac{67\ln(x+1)}{108}$
risch	$\frac{-\frac{5}{18}x^5 + \frac{13}{36}x^3 - \frac{3}{32}x}{x^6 - 2x^4 + \frac{19}{16}x^2 - \frac{3}{16}} + \frac{7\ln(2x-1)}{54} + \frac{5\sqrt{3}\ln(-\sqrt{3}+2x)}{18} - \frac{5\sqrt{3}\ln(\sqrt{3}+2x)}{18} + \frac{67\ln(x+1)}{108} - \frac{7\ln(2x+1)}{54} - \frac{67\ln(x-1)}{108}$

input `int(1/(-16*x^6+32*x^4-19*x^2+3)^2,x,method=_RETURNVERBOSE)`

output  $-1/36/(x-1)-67/108*\ln(x-1)-1/6*x/(x^2-3/4)-5/9*\operatorname{arctanh}(2/3*3^{(1/2)*x})*3^{(1/2)}-1/18/(2*x+1)-7/54*\ln(2*x+1)-1/36/(x+1)+67/108*\ln(x+1)-1/18/(2*x-1)+7/54*\ln(2*x-1)$

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 177 vs. 2(57) = 114.

Time = 0.09 (sec) , antiderivative size = 177, normalized size of antiderivative = 2.36

$$\int \frac{1}{(3 - 19x^2 + 32x^4 - 16x^6)^2} dx =$$

$$\frac{480x^5 - 624x^3 - 30\sqrt{3}(16x^6 - 32x^4 + 19x^2 - 3) \log\left(\frac{4x^2 - 4\sqrt{3}x + 3}{4x^2 - 3}\right) + 14(16x^6 - 32x^4 + 19x^2 - 3)}{\dots}$$

input `integrate(1/(-16*x^6+32*x^4-19*x^2+3)^2,x, algorithm="fricas")`

output 
$$\begin{aligned} & -1/108*(480*x^5 - 624*x^3 - 30*\sqrt{3}*(16*x^6 - 32*x^4 + 19*x^2 - 3)*\log( \\ & (4*x^2 - 4*\sqrt{3}*x + 3)/(4*x^2 - 3)) + 14*(16*x^6 - 32*x^4 + 19*x^2 - 3) \\ & *\log(2*x + 1) - 14*(16*x^6 - 32*x^4 + 19*x^2 - 3)*\log(2*x - 1) - 67*(16*x^6 \\ & - 32*x^4 + 19*x^2 - 3)*\log(x + 1) + 67*(16*x^6 - 32*x^4 + 19*x^2 - 3)*\log(x - 1) \\ & + 162*x)/(16*x^6 - 32*x^4 + 19*x^2 - 3) \end{aligned}$$

### Sympy [A] (verification not implemented)

Time = 0.85 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.39

$$\begin{aligned} \int \frac{1}{(3 - 19x^2 + 32x^4 - 16x^6)^2} dx &= \frac{-80x^5 + 104x^3 - 27x}{288x^6 - 576x^4 + 342x^2 - 54} - \frac{67 \log(x - 1)}{108} \\ &+ \frac{7 \log(x - \frac{1}{2})}{54} - \frac{7 \log(x + \frac{1}{2})}{54} + \frac{67 \log(x + 1)}{108} \\ &+ \frac{5\sqrt{3} \log(x - \frac{\sqrt{3}}{2})}{18} - \frac{5\sqrt{3} \log(x + \frac{\sqrt{3}}{2})}{18} \end{aligned}$$

input `integrate(1/(-16*x**6+32*x**4-19*x**2+3)**2,x)`

output 
$$\begin{aligned} & (-80*x**5 + 104*x**3 - 27*x)/(288*x**6 - 576*x**4 + 342*x**2 - 54) - 67*\log(x - 1)/108 \\ & + 7*\log(x - 1/2)/54 - 7*\log(x + 1/2)/54 + 67*\log(x + 1)/108 + \\ & 5*\sqrt{3}*\log(x - \sqrt{3}/2)/18 - 5*\sqrt{3}*\log(x + \sqrt{3}/2)/18 \end{aligned}$$

### Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.19

$$\begin{aligned} \int \frac{1}{(3 - 19x^2 + 32x^4 - 16x^6)^2} dx &= \frac{5}{18} \sqrt{3} \log\left(\frac{2x - \sqrt{3}}{2x + \sqrt{3}}\right) \\ & - \frac{80x^5 - 104x^3 + 27x}{18(16x^6 - 32x^4 + 19x^2 - 3)} - \frac{7}{54} \log(2x + 1) \\ & + \frac{7}{54} \log(2x - 1) + \frac{67}{108} \log(x + 1) - \frac{67}{108} \log(x - 1) \end{aligned}$$

input `integrate(1/(-16*x^6+32*x^4-19*x^2+3)^2,x, algorithm="maxima")`

output 
$$\frac{5}{18}\sqrt{3}\log\left(\frac{2x - \sqrt{3}}{2x + \sqrt{3}}\right) - \frac{1}{18}(80x^5 - 104x^3 + 27x)/(16x^6 - 32x^4 + 19x^2 - 3) - \frac{7}{54}\log(2x + 1) + \frac{7}{54}\log(2x - 1) + \frac{67}{108}\log(x + 1) - \frac{67}{108}\log(x - 1)$$

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.29

$$\int \frac{1}{(3 - 19x^2 + 32x^4 - 16x^6)^2} dx = \frac{5}{18} \sqrt{3} \log\left(\frac{|8x - 4\sqrt{3}|}{|8x + 4\sqrt{3}|}\right) - \frac{80x^5 - 104x^3 + 27x}{18(16x^6 - 32x^4 + 19x^2 - 3)} - \frac{7}{54} \log(|2x + 1|) + \frac{7}{54} \log(|2x - 1|) + \frac{67}{108} \log(|x + 1|) - \frac{67}{108} \log(|x - 1|)$$

input `integrate(1/(-16*x^6+32*x^4-19*x^2+3)^2,x, algorithm="giac")`

output 
$$\frac{5}{18}\sqrt{3}\log(\text{abs}(8x - 4\sqrt{3})/\text{abs}(8x + 4\sqrt{3})) - \frac{1}{18}(80x^5 - 104x^3 + 27x)/(16x^6 - 32x^4 + 19x^2 - 3) - \frac{7}{54}\log(\text{abs}(2x + 1)) + \frac{7}{54}\log(\text{abs}(2x - 1)) + \frac{67}{108}\log(\text{abs}(x + 1)) - \frac{67}{108}\log(\text{abs}(x - 1))$$

### Mupad [B] (verification not implemented)

Time = 23.14 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.85

$$\int \frac{1}{(3 - 19x^2 + 32x^4 - 16x^6)^2} dx = -\frac{\text{atan}(x \text{ li})}{54} \frac{67i}{27} + \frac{\text{atan}(x \text{ 2i})}{27} \frac{7i}{9} - \frac{\frac{5x^5}{18} - \frac{13x^3}{36} + \frac{3x}{32}}{x^6 - 2x^4 + \frac{19x^2}{16} - \frac{3}{16}} + \frac{\sqrt{3} \text{atan}\left(\frac{\sqrt{3}x \text{ 2i}}{3}\right)}{9} \frac{5i}{9}$$

input `int(1/(19*x^2 - 32*x^4 + 16*x^6 - 3)^2,x)`

output `(atan(x*2i)*7i)/27 - (atan(x*1i)*67i)/54 - ((3*x)/32 - (13*x^3)/36 + (5*x^5)/18)/((19*x^2)/16 - 2*x^4 + x^6 - 3/16) + (3^(1/2)*atan((3^(1/2)*x*2i)/3)*5i)/9`

### Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 296, normalized size of antiderivative = 3.95

$$\int \frac{1}{(3 - 19x^2 + 32x^4 - 16x^6)^2} dx$$

$$= \frac{-162x + 480\sqrt{3} \log(-\sqrt{3} + 2x) x^6 - 960\sqrt{3} \log(-\sqrt{3} + 2x) x^4 + 570\sqrt{3} \log(-\sqrt{3} + 2x) x^2 - 480\sqrt{3} \log(-\sqrt{3} + 2x)}{(3 - 19x^2 + 32x^4 - 16x^6)^2}$$

input `int(1/(-16*x^6+32*x^4-19*x^2+3)^2,x)`

output `(480*sqrt(3)*log(-sqrt(3)+2*x)*x**6 - 960*sqrt(3)*log(-sqrt(3)+2*x)*x**4 + 570*sqrt(3)*log(-sqrt(3)+2*x)*x**2 - 90*sqrt(3)*log(-sqrt(3)+2*x) - 480*sqrt(3)*log(sqrt(3)+2*x)*x**6 + 960*sqrt(3)*log(sqrt(3)+2*x)*x**4 - 570*sqrt(3)*log(sqrt(3)+2*x)*x**2 + 90*sqrt(3)*log(sqrt(3)+2*x) + 224*log(2*x-1)*x**6 - 448*log(2*x-1)*x**4 + 266*log(2*x-1)*x**2 - 42*log(2*x-1) - 224*log(2*x+1)*x**6 + 448*log(2*x+1)*x**4 - 266*log(2*x+1)*x**2 + 42*log(2*x+1) - 1072*log(x-1)*x**6 + 2144*log(x-1)*x**4 - 1273*log(x-1)*x**2 + 201*log(x-1) + 1072*log(x+1)*x**6 - 2144*log(x+1)*x**4 + 1273*log(x+1)*x**2 - 201*log(x+1) - 480*x**5 + 624*x**3 - 162*x)/(108*(16*x**6 - 32*x**4 + 19*x**2 - 3))`

**3.18**  $\int \frac{1}{(3-19x^2+32x^4-16x^6)^3} dx$

Optimal result . . . . . 140  
 Mathematica [A] (verified) . . . . . 141  
 Rubi [A] (verified) . . . . . 141  
 Maple [A] (verified) . . . . . 142  
 Fricas [B] (verification not implemented) . . . . . 143  
 Sympy [A] (verification not implemented) . . . . . 144  
 Maxima [A] (verification not implemented) . . . . . 144  
 Giac [A] (verification not implemented) . . . . . 145  
 Mupad [B] (verification not implemented) . . . . . 145  
 Reduce [B] (verification not implemented) . . . . . 146

**Optimal result**

Integrand size = 19, antiderivative size = 117

$$\int \frac{1}{(3-19x^2+32x^4-16x^6)^3} dx = \frac{2x}{27(1-4x^2)^2} - \frac{7x}{27(1-4x^2)} - \frac{2x}{3(3-4x^2)^2} + \frac{5x}{3(3-4x^2)} + \frac{x}{108(1-x^2)^2} + \frac{67x}{216(1-x^2)} + \frac{3913\operatorname{arctanh}(x)}{648} + \frac{67}{162}\operatorname{arctanh}(2x) - \frac{67\operatorname{arctanh}\left(\frac{2x}{\sqrt{3}}\right)}{6\sqrt{3}}$$

output

```
2/27*x/(-4*x^2+1)^2-7*x/(-108*x^2+27)-2/3*x/(-4*x^2+3)^2+5*x/(-12*x^2+9)+1/108*x/(-x^2+1)^2+67*x/(-216*x^2+216)+3913/648*arctanh(x)+67/162*arctanh(2*x)-67/18*arctanh(2/3*x*3^(1/2))*3^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.11 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.17

$$\int \frac{1}{(3 - 19x^2 + 32x^4 - 16x^6)^3} dx$$

$$= \frac{\frac{36x(27-104x^2+80x^4)}{(3-19x^2+32x^4-16x^6)^2} - \frac{6x(345-2384x^2+2288x^4)}{-3+19x^2-32x^4+16x^6} - 268 \log(1-2x) + 2412\sqrt{3} \log(\sqrt{3}-2x) - 3913 \log(1-x)}{1296}$$

input

```
Integrate[(3 - 19*x^2 + 32*x^4 - 16*x^6)^(-3), x]
```

output

```
((36*x*(27 - 104*x^2 + 80*x^4))/(3 - 19*x^2 + 32*x^4 - 16*x^6)^2 - (6*x*(345 - 2384*x^2 + 2288*x^4))/(-3 + 19*x^2 - 32*x^4 + 16*x^6) - 268*Log[1 - 2*x] + 2412*Sqrt[3]*Log[Sqrt[3] - 2*x] - 3913*Log[1 - x] + 3913*Log[1 + x] + 268*Log[1 + 2*x] - 2412*Sqrt[3]*Log[Sqrt[3] + 2*x])/1296
```

**Rubi [A] (verified)**

Time = 0.56 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.38, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$ , Rules used = {2460, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(-16x^6 + 32x^4 - 19x^2 + 3)^3} dx$$

$$\downarrow \text{2460}$$

$$\int \left( \frac{24}{4x^2 - 3} - \frac{67}{81(4x^2 - 1)} + \frac{12}{(4x^2 - 3)^2} + \frac{8}{(4x^2 - 3)^3} - \frac{3913}{648(x^2 - 1)} + \frac{67}{432(x - 1)^2} + \frac{67}{432(x + 1)^2} - \frac{7}{54(2x - 1)} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{3913 \operatorname{arctanh}(x)}{648} + \frac{67}{162} \operatorname{arctanh}(2x) - 4\sqrt{3} \operatorname{arctanh}\left(\frac{2x}{\sqrt{3}}\right) + \frac{5 \operatorname{arctanh}\left(\frac{2x}{\sqrt{3}}\right)}{6\sqrt{3}} + \frac{5x}{3(3-4x^2)} - \frac{1}{3(3-4x^2)^2} - \frac{7}{108(1-2x)} + \frac{67}{432(1-x)} - \frac{67}{432(x+1)} + \frac{1}{108(2x+1)} + \frac{1}{108(1-2x)^2} + \frac{1}{432(1-x)^2} - \frac{1}{432(x+1)^2} - \frac{1}{108(2x+1)^2}$$

input `Int[(3 - 19*x^2 + 32*x^4 - 16*x^6)^(-3), x]`

output `1/(108*(1 - 2*x)^2) - 7/(108*(1 - 2*x)) + 1/(432*(1 - x)^2) + 67/(432*(1 - x)) - 1/(432*(1 + x)^2) - 67/(432*(1 + x)) - 1/(108*(1 + 2*x)^2) + 7/(108*(1 + 2*x)) - (2*x)/(3*(3 - 4*x^2)^2) + (5*x)/(3*(3 - 4*x^2)) + (3913*ArcTanh[x])/648 + (67*ArcTanh[2*x])/162 + (5*ArcTanh[(2*x)/Sqrt[3]])/(6*Sqrt[3]) - 4*Sqrt[3]*ArcTanh[(2*x)/Sqrt[3]]`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2460 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px /. x -> Sqrt[x]]}, Int[ExpandIntegrand[u*(Qx /. x -> x^2)^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x^2] && GtQ[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0] && RationalFunctionQ[u, x]`

**Maple [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.92

method	result
risch	$\frac{-\frac{4576}{27}x^{11} + \frac{4640}{9}x^9 - 580x^7 + \frac{7960}{27}x^5 - \frac{4777}{72}x^3 + \frac{133}{24}x}{(16x^6 - 32x^4 + 19x^2 - 3)^2} - \frac{3913 \ln(x-1)}{1296} + \frac{67\sqrt{3} \ln(-\sqrt{3}+2x)}{36} - \frac{67\sqrt{3} \ln(\sqrt{3}+2x)}{36} - \frac{67 \ln(2)}{32}$
default	$\frac{1}{432(x-1)^2} - \frac{67}{432(x-1)} - \frac{3913 \ln(x-1)}{1296} + \frac{-\frac{20}{3}x^3 + \frac{13}{3}x}{(4x^2-3)^2} - \frac{67 \operatorname{arctanh}\left(\frac{2\sqrt{3}x}{3}\right)\sqrt{3}}{18} - \frac{1}{108(2x+1)^2} + \frac{7}{108(2x+1)} + \frac{67 \ln(2)}{32}$

input `int(1/(-16*x^6+32*x^4-19*x^2+3)^3,x,method=_RETURNVERBOSE)`

output 
$$\frac{256*(-143/216*x^{11}+145/72*x^9-145/64*x^7+995/864*x^5-4777/18432*x^3+133/6144*x)}{(16*x^6-32*x^4+19*x^2-3)^2-3913/1296*\ln(x-1)+67/36*3^{(1/2)}*\ln(-3^{(1/2)}+2*x)-67/36*3^{(1/2)}*\ln(3^{(1/2)}+2*x)-67/324*\ln(2*x-1)+67/324*\ln(2*x+1)+3913/1296*\ln(x+1)}$$

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 282 vs.  $2(91) = 182$ .

Time = 0.11 (sec) , antiderivative size = 282, normalized size of antiderivative = 2.41

$$\int \frac{1}{(3 - 19x^2 + 32x^4 - 16x^6)^3} dx =$$


---


$$\frac{219648 x^{11} - 668160 x^9 + 751680 x^7 - 382080 x^5 + 85986 x^3 - 2412 \sqrt{3}(256 x^{12} - 1024 x^{10} + 1632 x^8 - 1312 x^6 + 553 x^4 - 114 x^2 + 9) \log((4 x^2 - 4 \sqrt{3} x + 3)/(4 x^2 - 3)) - 268(256 x^{12} - 1024 x^{10} + 1632 x^8 - 1312 x^6 + 553 x^4 - 114 x^2 + 9) \log(2 x + 1) + 268(256 x^{12} - 1024 x^{10} + 1632 x^8 - 1312 x^6 + 553 x^4 - 114 x^2 + 9) \log(2 x - 1) - 3913(256 x^{12} - 1024 x^{10} + 1632 x^8 - 1312 x^6 + 553 x^4 - 114 x^2 + 9) \log(x + 1) + 3913(256 x^{12} - 1024 x^{10} + 1632 x^8 - 1312 x^6 + 553 x^4 - 114 x^2 + 9) \log(x - 1) - 7182 x}{(256 x^{12} - 1024 x^{10} + 1632 x^8 - 1312 x^6 + 553 x^4 - 114 x^2 + 9)^3}$$

input `integrate(1/(-16*x^6+32*x^4-19*x^2+3)^3,x, algorithm="fricas")`

output 
$$\frac{-1/1296*(219648*x^{11} - 668160*x^9 + 751680*x^7 - 382080*x^5 + 85986*x^3 - 2412*\sqrt{3}*(256*x^{12} - 1024*x^{10} + 1632*x^8 - 1312*x^6 + 553*x^4 - 114*x^2 + 9)*\log((4*x^2 - 4*\sqrt{3}*x + 3)/(4*x^2 - 3)) - 268*(256*x^{12} - 1024*x^{10} + 1632*x^8 - 1312*x^6 + 553*x^4 - 114*x^2 + 9)*\log(2*x + 1) + 268*(256*x^{12} - 1024*x^{10} + 1632*x^8 - 1312*x^6 + 553*x^4 - 114*x^2 + 9)*\log(2*x - 1) - 3913*(256*x^{12} - 1024*x^{10} + 1632*x^8 - 1312*x^6 + 553*x^4 - 114*x^2 + 9)*\log(x + 1) + 3913*(256*x^{12} - 1024*x^{10} + 1632*x^8 - 1312*x^6 + 553*x^4 - 114*x^2 + 9)*\log(x - 1) - 7182*x}{(256*x^{12} - 1024*x^{10} + 1632*x^8 - 1312*x^6 + 553*x^4 - 114*x^2 + 9)^3}$$



**Sympy [A] (verification not implemented)**

Time = 0.92 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.15

$$\int \frac{1}{(3 - 19x^2 + 32x^4 - 16x^6)^3} dx$$

$$= -\frac{36608x^{11} - 111360x^9 + 125280x^7 - 63680x^5 + 14331x^3 - 1197x}{55296x^{12} - 221184x^{10} + 352512x^8 - 283392x^6 + 119448x^4 - 24624x^2 + 1944}$$

$$- \frac{3913 \log(x-1)}{1296} - \frac{67 \log(x-\frac{1}{2})}{324} + \frac{67 \log(x+\frac{1}{2})}{324}$$

$$+ \frac{3913 \log(x+1)}{1296} + \frac{67\sqrt{3} \log(x-\frac{\sqrt{3}}{2})}{36} - \frac{67\sqrt{3} \log(x+\frac{\sqrt{3}}{2})}{36}$$

input `integrate(1/(-16*x**6+32*x**4-19*x**2+3)**3,x)`output `-(36608*x**11 - 111360*x**9 + 125280*x**7 - 63680*x**5 + 14331*x**3 - 1197*x)/(55296*x**12 - 221184*x**10 + 352512*x**8 - 283392*x**6 + 119448*x**4 - 24624*x**2 + 1944) - 3913*log(x - 1)/1296 - 67*log(x - 1/2)/324 + 67*log(x + 1/2)/324 + 3913*log(x + 1)/1296 + 67*sqrt(3)*log(x - sqrt(3)/2)/36 - 67*sqrt(3)*log(x + sqrt(3)/2)/36`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.02

$$\int \frac{1}{(3 - 19x^2 + 32x^4 - 16x^6)^3} dx$$

$$= \frac{67}{36} \sqrt{3} \log\left(\frac{2x - \sqrt{3}}{2x + \sqrt{3}}\right)$$

$$- \frac{36608x^{11} - 111360x^9 + 125280x^7 - 63680x^5 + 14331x^3 - 1197x}{216(256x^{12} - 1024x^{10} + 1632x^8 - 1312x^6 + 553x^4 - 114x^2 + 9)}$$

$$+ \frac{67}{324} \log(2x+1) - \frac{67}{324} \log(2x-1) + \frac{3913}{1296} \log(x+1) - \frac{3913}{1296} \log(x-1)$$

input `integrate(1/(-16*x^6+32*x^4-19*x^2+3)^3,x, algorithm="maxima")`

output

```
67/36*sqrt(3)*log((2*x - sqrt(3))/(2*x + sqrt(3))) - 1/216*(36608*x^11 - 1
11360*x^9 + 125280*x^7 - 63680*x^5 + 14331*x^3 - 1197*x)/(256*x^12 - 1024*
x^10 + 1632*x^8 - 1312*x^6 + 553*x^4 - 114*x^2 + 9) + 67/324*log(2*x + 1)
- 67/324*log(2*x - 1) + 3913/1296*log(x + 1) - 3913/1296*log(x - 1)
```

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.96

$$\int \frac{1}{(3 - 19x^2 + 32x^4 - 16x^6)^3} dx$$

$$= \frac{67}{36} \sqrt{3} \log \left( \frac{|8x - 4\sqrt{3}|}{|8x + 4\sqrt{3}|} \right)$$

$$- \frac{36608x^{11} - 111360x^9 + 125280x^7 - 63680x^5 + 14331x^3 - 1197x}{216(16x^6 - 32x^4 + 19x^2 - 3)^2}$$

$$+ \frac{67}{324} \log(|2x + 1|) - \frac{67}{324} \log(|2x - 1|) + \frac{3913}{1296} \log(|x + 1|) - \frac{3913}{1296} \log(|x - 1|)$$

input

```
integrate(1/(-16*x^6+32*x^4-19*x^2+3)^3,x, algorithm="giac")
```

output

```
67/36*sqrt(3)*log(abs(8*x - 4*sqrt(3))/abs(8*x + 4*sqrt(3))) - 1/216*(3660
8*x^11 - 111360*x^9 + 125280*x^7 - 63680*x^5 + 14331*x^3 - 1197*x)/(16*x^6
- 32*x^4 + 19*x^2 - 3)^2 + 67/324*log(abs(2*x + 1)) - 67/324*log(abs(2*x
- 1)) + 3913/1296*log(abs(x + 1)) - 3913/1296*log(abs(x - 1))
```

**Mupad [B] (verification not implemented)**

Time = 23.17 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.79

$$\int \frac{1}{(3 - 19x^2 + 32x^4 - 16x^6)^3} dx = \frac{-\frac{143x^{11}}{216} + \frac{145x^9}{72} - \frac{145x^7}{64} + \frac{995x^5}{864} - \frac{4777x^3}{18432} + \frac{133x}{6144}}{x^{12} - 4x^{10} + \frac{51x^8}{8} - \frac{41x^6}{8} + \frac{553x^4}{256} - \frac{57x^2}{128} + \frac{9}{256}}$$

$$- \frac{\operatorname{atan}(x 2i) 67i}{162} - \frac{\operatorname{atan}(x 1i) 3913i}{648}$$

$$+ \frac{\sqrt{3} \operatorname{atan}\left(\frac{\sqrt{3}x 2i}{3}\right) 67i}{18}$$

input `int(-1/(19*x^2 - 32*x^4 + 16*x^6 - 3)^3,x)`

output `((133*x)/6144 - (4777*x^3)/18432 + (995*x^5)/864 - (145*x^7)/64 + (145*x^9)/72 - (143*x^11)/216)/((553*x^4)/256 - (57*x^2)/128 - (41*x^6)/8 + (51*x^8)/8 - 4*x^10 + x^12 + 9/256) - (atan(x*2i)*67i)/162 - (atan(x*1i)*3913i)/648 + (3^(1/2)*atan((3^(1/2)*x*2i)/3)*67i)/18`

### Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 536, normalized size of antiderivative = 4.58

$$\int \frac{1}{(3 - 19x^2 + 32x^4 - 16x^6)^3} dx = \text{Too large to display}$$

input `int(1/(-16*x^6+32*x^4-19*x^2+3)^3,x)`

output `(617472*sqrt(3)*log(-sqrt(3)+2*x)*x**12 - 2469888*sqrt(3)*log(-sqrt(3)+2*x)*x**10 + 3936384*sqrt(3)*log(-sqrt(3)+2*x)*x**8 - 3164544*sqrt(3)*log(-sqrt(3)+2*x)*x**6 + 1333836*sqrt(3)*log(-sqrt(3)+2*x)*x**4 - 274968*sqrt(3)*log(-sqrt(3)+2*x)*x**2 + 21708*sqrt(3)*log(-sqrt(3)+2*x) - 617472*sqrt(3)*log(sqrt(3)+2*x)*x**12 + 2469888*sqrt(3)*log(sqrt(3)+2*x)*x**10 - 3936384*sqrt(3)*log(sqrt(3)+2*x)*x**8 + 3164544*sqrt(3)*log(sqrt(3)+2*x)*x**6 - 1333836*sqrt(3)*log(sqrt(3)+2*x)*x**4 + 274968*sqrt(3)*log(sqrt(3)+2*x)*x**2 - 21708*sqrt(3)*log(sqrt(3)+2*x) - 68608*log(2*x-1)*x**12 + 274432*log(2*x-1)*x**10 - 437376*log(2*x-1)*x**8 + 351616*log(2*x-1)*x**6 - 148204*log(2*x-1)*x**4 + 30552*log(2*x-1)*x**2 - 2412*log(2*x-1) + 68608*log(2*x+1)*x**12 - 274432*log(2*x+1)*x**10 + 437376*log(2*x+1)*x**8 - 351616*log(2*x+1)*x**6 + 148204*log(2*x+1)*x**4 - 30552*log(2*x+1)*x**2 + 2412*log(2*x+1) - 1001728*log(x-1)*x**12 + 4006912*log(x-1)*x**10 - 6386016*log(x-1)*x**8 + 5133856*log(x-1)*x**6 - 2163889*log(x-1)*x**4 + 446082*log(x-1)*x**2 - 35217*log(x-1) + 1001728*log(x+1)*x**12 - 4006912*log(x+1)*x**10 + 6386016*log(x+1)*x**8 - 5133856*log(x+1)*x**6 + 2163889*log(x+1)*x**4 - 446082*log(x+1)*x**2 + 35217*log(x+1) - 219648*x**11 + 668160*x**9 - 751680*x**7 + 382080*x**5 - 85986*x**3 + 7182*x)/(1296*(256*x**12 - 1024*x**10 + 1632*x**8 - 1312*x**6 + 553*x**4 - 114*x**2 + 9))`

### 3.19 $\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)$

Optimal result	147
Mathematica [A] (verified)	147
Rubi [A] (verified)	148
Maple [B] (verified)	149
Fricas [B] (verification not implemented)	149
Sympy [B] (verification not implemented)	150
Maxima [B] (verification not implemented)	151
Giac [B] (verification not implemented)	152
Mupad [B] (verification not implemented)	153
Reduce [B] (verification not implemented)	153

#### Optimal result

Integrand size = 51, antiderivative size = 14

$$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3 dx = \frac{(a + bx)^{16}}{16b}$$

output `1/16*(b*x+a)^16/b`

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3 dx = \frac{(a + bx)^{16}}{16b}$$

input `Integrate[(a^5 + 5*a^4*b*x + 10*a^3*b^2*x^2 + 10*a^2*b^3*x^3 + 5*a*b^4*x^4 + b^5*x^5)^3, x]`

output `(a + b*x)^16/(16*b)`

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.039$ , Rules used = {2006, 17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3 dx$$

↓ 2006

$$\int (a + bx)^{15} dx$$

↓ 17

$$\frac{(a + bx)^{16}}{16b}$$

input

```
Int[(a^5 + 5*a^4*b*x + 10*a^3*b^2*x^2 + 10*a^2*b^3*x^3 + 5*a*b^4*x^4 + b^5*x^5)^3,x]
```

output

```
(a + b*x)^16/(16*b)
```

**Defintions of rubi rules used**

rule 17

```
Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] :> Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]
```

rule 2006

```
Int[(u_.)*(Px_), x_Symbol] :> With[{a = Rt[Coeff[Px, x, 0], Expon[Px, x]], b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Int[u*(a + b*x)^Expon[Px, x], x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; PolyQ[Px, x] && GtQ[Expon[Px, x], 1] && NeQ[Coeff[Px, x, 0], 0] && !MatchQ[Px, (a_.)*(v_)^Expon[Px, x]] /; FreeQ[a, x] && LinearQ[v, x]]
```

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 163 vs.  $2(12) = 24$ .

Time = 0.04 (sec) , antiderivative size = 164, normalized size of antiderivative = 11.71

method	result
default	$a^{15}x + \frac{15}{2}a^{14}bx^2 + 35a^{13}b^2x^3 + \frac{455}{4}a^{12}b^3x^4 + 273a^{11}b^4x^5 + \frac{1001}{2}a^{10}b^5x^6 + 715a^9b^6x^7 + \frac{6435}{8}$
norman	$a^{15}x + \frac{15}{2}a^{14}bx^2 + 35a^{13}b^2x^3 + \frac{455}{4}a^{12}b^3x^4 + 273a^{11}b^4x^5 + \frac{1001}{2}a^{10}b^5x^6 + 715a^9b^6x^7 + \frac{6435}{8}$
risch	$a^{15}x + \frac{15}{2}a^{14}bx^2 + 35a^{13}b^2x^3 + \frac{455}{4}a^{12}b^3x^4 + 273a^{11}b^4x^5 + \frac{1001}{2}a^{10}b^5x^6 + 715a^9b^6x^7 + \frac{6435}{8}$
parallelrisch	$a^{15}x + \frac{15}{2}a^{14}bx^2 + 35a^{13}b^2x^3 + \frac{455}{4}a^{12}b^3x^4 + 273a^{11}b^4x^5 + \frac{1001}{2}a^{10}b^5x^6 + 715a^9b^6x^7 + \frac{6435}{8}$
gosper	$\frac{x(b^{15}x^{15} + 16ab^{14}x^{14} + 120a^2b^{13}x^{13} + 560a^3b^{12}x^{12} + 1820a^4b^{11}x^{11} + 4368a^5b^{10}x^{10} + 8008a^6b^9x^9 + 11440a^7b^8x^8 + 12870a^8b^7x^7 + 6435a^9b^6x^6 + 273a^{10}b^5x^5 + 715a^{11}b^4x^4 + 1001a^{12}b^3x^3 + 455a^{13}b^2x^2 + 15a^{14}bx + a^{15})^3}{16}$
orering	$\frac{x(b^{15}x^{15} + 16ab^{14}x^{14} + 120a^2b^{13}x^{13} + 560a^3b^{12}x^{12} + 1820a^4b^{11}x^{11} + 4368a^5b^{10}x^{10} + 8008a^6b^9x^9 + 11440a^7b^8x^8 + 12870a^8b^7x^7 + 6435a^9b^6x^6 + 273a^{10}b^5x^5 + 715a^{11}b^4x^4 + 1001a^{12}b^3x^3 + 455a^{13}b^2x^2 + 15a^{14}bx + a^{15})^3}{16}$

input

```
int((b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5)^3,x,
method=_RETURNVERBOSE)
```

output

```
a^15*x+15/2*a^14*b*x^2+35*a^13*b^2*x^3+455/4*a^12*b^3*x^4+273*a^11*b^4*x^5
+1001/2*a^10*b^5*x^6+715*a^9*b^6*x^7+6435/8*a^8*b^7*x^8+715*a^7*b^8*x^9+10
01/2*a^6*b^9*x^10+273*a^5*b^10*x^11+455/4*a^4*b^11*x^12+35*a^3*b^12*x^13+1
5/2*a^2*b^13*x^14+a*b^14*x^15+1/16*b^15*x^16
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 163 vs.  $2(12) = 24$ .

Time = 0.07 (sec) , antiderivative size = 163, normalized size of antiderivative = 11.64

$$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3 dx$$

$$= \frac{1}{16}b^{15}x^{16} + ab^{14}x^{15} + \frac{15}{2}a^2b^{13}x^{14} + 35a^3b^{12}x^{13} + \frac{455}{4}a^4b^{11}x^{12}$$

$$+ 273a^5b^{10}x^{11} + \frac{1001}{2}a^6b^9x^{10} + 715a^7b^8x^9 + \frac{6435}{8}a^8b^7x^8 + 715a^9b^6x^7$$

$$+ \frac{1001}{2}a^{10}b^5x^6 + 273a^{11}b^4x^5 + \frac{455}{4}a^{12}b^3x^4 + 35a^{13}b^2x^3 + \frac{15}{2}a^{14}bx^2 + a^{15}x$$

input

```
integrate((b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5)^3,x, algorithm="fricas")
```

output

```
1/16*b^15*x^16 + a*b^14*x^15 + 15/2*a^2*b^13*x^14 + 35*a^3*b^12*x^13 + 455/4*a^4*b^11*x^12 + 273*a^5*b^10*x^11 + 1001/2*a^6*b^9*x^10 + 715*a^7*b^8*x^9 + 6435/8*a^8*b^7*x^8 + 715*a^9*b^6*x^7 + 1001/2*a^10*b^5*x^6 + 273*a^11*b^4*x^5 + 455/4*a^12*b^3*x^4 + 35*a^13*b^2*x^3 + 15/2*a^14*b*x^2 + a^15*x
```

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 185 vs.  $2(8) = 16$ .

Time = 0.04 (sec) , antiderivative size = 185, normalized size of antiderivative = 13.21

$$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3 dx$$

$$= a^{15}x + \frac{15a^{14}bx^2}{2} + 35a^{13}b^2x^3 + \frac{455a^{12}b^3x^4}{4} + 273a^{11}b^4x^5 + \frac{1001a^{10}b^5x^6}{2}$$

$$+ 715a^9b^6x^7 + \frac{6435a^8b^7x^8}{8} + 715a^7b^8x^9 + \frac{1001a^6b^9x^{10}}{2} + 273a^5b^{10}x^{11}$$

$$+ \frac{455a^4b^{11}x^{12}}{4} + 35a^3b^{12}x^{13} + \frac{15a^2b^{13}x^{14}}{2} + ab^{14}x^{15} + \frac{b^{15}x^{16}}{16}$$

input

```
integrate((b**5*x**5+5*a*b**4*x**4+10*a**2*b**3*x**3+10*a**3*b**2*x**2+5*a**4*b*x+a**5)**3,x)
```

output

```
a**15*x + 15*a**14*b*x**2/2 + 35*a**13*b**2*x**3 + 455*a**12*b**3*x**4/4 + 273*a**11*b**4*x**5 + 1001*a**10*b**5*x**6/2 + 715*a**9*b**6*x**7 + 6435*a**8*b**7*x**8/8 + 715*a**7*b**8*x**9 + 1001*a**6*b**9*x**10/2 + 273*a**5*b**10*x**11 + 455*a**4*b**11*x**12/4 + 35*a**3*b**12*x**13 + 15*a**2*b**13*x**14/2 + a*b**14*x**15 + b**15*x**16/16
```

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 592 vs.  $2(12) = 24$ .

Time = 0.04 (sec) , antiderivative size = 592, normalized size of antiderivative = 42.29

$$\begin{aligned}
 & \int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3 dx \\
 &= \frac{1}{16} b^{15} x^{16} + ab^{14} x^{15} + \frac{75}{14} a^2 b^{13} x^{14} + \frac{125}{13} a^3 b^{12} x^{13} + 100 a^6 b^9 x^{10} + \frac{1000}{7} a^9 b^6 x^7 \\
 &+ \frac{125}{4} a^{12} b^3 x^4 + a^{15} x + \frac{1}{2} (b^5 x^6 + 6 ab^4 x^5 + 15 a^2 b^3 x^4 + 20 a^3 b^2 x^3 + 15 a^4 b x^2) a^{10} \\
 &+ \frac{25}{56} (21 b^5 x^8 + 120 ab^4 x^7 + 280 a^2 b^3 x^6 + 336 a^3 b^2 x^5) a^8 b^2 \\
 &+ \frac{5}{3} (18 b^5 x^{10} + 100 ab^4 x^9 + 225 a^2 b^3 x^8) a^6 b^4 + \frac{25}{11} (11 b^5 x^{12} + 60 ab^4 x^{11}) a^4 b^6 \\
 &+ \frac{1}{462} (126 b^{10} x^{11} + 1386 ab^9 x^{10} + 3850 a^2 b^8 x^9 + 19800 a^4 b^6 x^7 + 27720 a^6 b^4 x^5 + 11550 a^8 b^2 x^3 + 330 (6 b^5 x^6 + 6 ab^4 x^5 + 15 a^2 b^3 x^4 + 20 a^3 b^2 x^3 + 15 a^4 b x^2) a^{10} \\
 &+ \frac{5}{308} (77 b^{10} x^{12} + 840 ab^9 x^{11} + 4158 a^2 b^8 x^{10} + 12320 a^3 b^7 x^9 + 23100 a^4 b^6 x^8 + 26400 a^5 b^5 x^7 + 15400 a^6 b^4 x^6 + 5600 a^7 b^3 x^5 + 560 a^8 b^2 x^4 + 56 a^9 b x^3 + 56 a^{10} x^2) a^2 b^2 \\
 &+ \frac{5}{429} (198 b^{10} x^{13} + 2145 ab^9 x^{12} + 10530 a^2 b^8 x^{11} + 25740 a^3 b^7 x^{10} + 28600 a^4 b^6 x^9) a^3 b^2 \\
 &+ \frac{5}{182} (78 b^{10} x^{14} + 840 ab^9 x^{13} + 2275 a^2 b^8 x^{12}) a^2 b^3
 \end{aligned}$$

input `integrate((b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5)^3,x, algorithm="maxima")`



output

```

1/16*b^15*x^16 + a*b^14*x^15 + 75/14*a^2*b^13*x^14 + 125/13*a^3*b^12*x^13
+ 100*a^6*b^9*x^10 + 1000/7*a^9*b^6*x^7 + 125/4*a^12*b^3*x^4 + a^15*x + 1/
2*(b^5*x^6 + 6*a*b^4*x^5 + 15*a^2*b^3*x^4 + 20*a^3*b^2*x^3 + 15*a^4*b*x^2)
*a^10 + 25/56*(21*b^5*x^8 + 120*a*b^4*x^7 + 280*a^2*b^3*x^6 + 336*a^3*b^2*
x^5)*a^8*b^2 + 5/3*(18*b^5*x^10 + 100*a*b^4*x^9 + 225*a^2*b^3*x^8)*a^6*b^4
+ 25/11*(11*b^5*x^12 + 60*a*b^4*x^11)*a^4*b^6 + 1/462*(126*b^10*x^11 + 13
86*a*b^9*x^10 + 3850*a^2*b^8*x^9 + 19800*a^4*b^6*x^7 + 27720*a^6*b^4*x^5 +
11550*a^8*b^2*x^3 + 330*(6*b^5*x^7 + 35*a*b^4*x^6 + 84*a^2*b^3*x^5 + 105*
a^3*b^2*x^4)*a^4*b + 165*(21*b^5*x^8 + 120*a*b^4*x^7 + 280*a^2*b^3*x^6)*a^
3*b^2 + 385*(8*b^5*x^9 + 45*a*b^4*x^8)*a^2*b^3)*a^5 + 5/308*(77*b^10*x^12
+ 840*a*b^9*x^11 + 4158*a^2*b^8*x^10 + 12320*a^3*b^7*x^9 + 23100*a^4*b^6*x
^8 + 26400*a^5*b^5*x^7 + 15400*a^6*b^4*x^6)*a^4*b + 5/429*(198*b^10*x^13 +
2145*a*b^9*x^12 + 10530*a^2*b^8*x^11 + 25740*a^3*b^7*x^10 + 28600*a^4*b^6
*x^9)*a^3*b^2 + 5/182*(78*b^10*x^14 + 840*a*b^9*x^13 + 2275*a^2*b^8*x^12)*
a^2*b^3

```

### Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 163 vs.  $2(12) = 24$ .

Time = 0.11 (sec) , antiderivative size = 163, normalized size of antiderivative = 11.64

$$\begin{aligned}
 & \int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3 dx \\
 &= \frac{1}{16} b^{15} x^{16} + ab^{14} x^{15} + \frac{15}{2} a^2 b^{13} x^{14} + 35 a^3 b^{12} x^{13} + \frac{455}{4} a^4 b^{11} x^{12} \\
 &+ 273 a^5 b^{10} x^{11} + \frac{1001}{2} a^6 b^9 x^{10} + 715 a^7 b^8 x^9 + \frac{6435}{8} a^8 b^7 x^8 + 715 a^9 b^6 x^7 \\
 &+ \frac{1001}{2} a^{10} b^5 x^6 + 273 a^{11} b^4 x^5 + \frac{455}{4} a^{12} b^3 x^4 + 35 a^{13} b^2 x^3 + \frac{15}{2} a^{14} b x^2 + a^{15} x
 \end{aligned}$$

input

```

integrate((b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5
)^3,x, algorithm="giac")

```

output

```

1/16*b^15*x^16 + a*b^14*x^15 + 15/2*a^2*b^13*x^14 + 35*a^3*b^12*x^13 + 455
/4*a^4*b^11*x^12 + 273*a^5*b^10*x^11 + 1001/2*a^6*b^9*x^10 + 715*a^7*b^8*x
^9 + 6435/8*a^8*b^7*x^8 + 715*a^9*b^6*x^7 + 1001/2*a^10*b^5*x^6 + 273*a^11
*b^4*x^5 + 455/4*a^12*b^3*x^4 + 35*a^13*b^2*x^3 + 15/2*a^14*b*x^2 + a^15*x

```

**Mupad [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 163, normalized size of antiderivative = 11.64

$$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3 dx$$

$$= a^{15}x + \frac{15a^{14}bx^2}{2} + 35a^{13}b^2x^3 + \frac{455a^{12}b^3x^4}{4} + 273a^{11}b^4x^5 + \frac{1001a^{10}b^5x^6}{2}$$

$$+ 715a^9b^6x^7 + \frac{6435a^8b^7x^8}{8} + 715a^7b^8x^9 + \frac{1001a^6b^9x^{10}}{2} + 273a^5b^{10}x^{11}$$

$$+ \frac{455a^4b^{11}x^{12}}{4} + 35a^3b^{12}x^{13} + \frac{15a^2b^{13}x^{14}}{2} + ab^{14}x^{15} + \frac{b^{15}x^{16}}{16}$$

input

```
int((a^5 + b^5*x^5 + 5*a*b^4*x^4 + 10*a^3*b^2*x^2 + 10*a^2*b^3*x^3 + 5*a^4*b*x)^3,x)
```

output

```
a^15*x + (b^15*x^16)/16 + (15*a^14*b*x^2)/2 + a*b^14*x^15 + 35*a^13*b^2*x^3 + (455*a^12*b^3*x^4)/4 + 273*a^11*b^4*x^5 + (1001*a^10*b^5*x^6)/2 + 715*a^9*b^6*x^7 + (6435*a^8*b^7*x^8)/8 + 715*a^7*b^8*x^9 + (1001*a^6*b^9*x^10)/2 + 273*a^5*b^10*x^11 + (455*a^4*b^11*x^12)/4 + 35*a^3*b^12*x^13 + (15*a^2*b^13*x^14)/2
```

**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 164, normalized size of antiderivative = 11.71

$$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3 dx$$

$$= \frac{x(b^{15}x^{15} + 16ab^{14}x^{14} + 120a^2b^{13}x^{13} + 560a^3b^{12}x^{12} + 1820a^4b^{11}x^{11} + 4368a^5b^{10}x^{10} + 8008a^6b^9x^9 + 11440a^7b^8x^8 + 8008a^8b^7x^7 + 4368a^9b^6x^6 + 1820a^{10}b^5x^5 + 120a^{11}b^4x^4 + 16a^{12}b^3x^3 + 15a^{13}b^2x^2 + 15a^{14}bx + a^{15})}{16}$$

input

```
int((b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5)^3,x)
```

output

```
(x*(16*a**15 + 120*a**14*b*x + 560*a**13*b**2*x**2 + 1820*a**12*b**3*x**3 + 4368*a**11*b**4*x**4 + 8008*a**10*b**5*x**5 + 11440*a**9*b**6*x**6 + 12870*a**8*b**7*x**7 + 11440*a**7*b**8*x**8 + 8008*a**6*b**9*x**9 + 4368*a**5*b**10*x**10 + 1820*a**4*b**11*x**11 + 560*a**3*b**12*x**12 + 120*a**2*b**13*x**13 + 16*a*b**14*x**14 + b**15*x**15))/16
```

### 3.20 $\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2 dx$

Optimal result	154
Mathematica [A] (verified)	154
Rubi [A] (verified)	155
Maple [B] (verified)	156
Fricas [B] (verification not implemented)	156
Sympy [B] (verification not implemented)	157
Maxima [B] (verification not implemented)	157
Giac [B] (verification not implemented)	158
Mupad [B] (verification not implemented)	159
Reduce [B] (verification not implemented)	159

#### Optimal result

Integrand size = 51, antiderivative size = 14

$$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2 dx = \frac{(a + bx)^{11}}{11b}$$

output `1/11*(b*x+a)^11/b`

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2 dx = \frac{(a + bx)^{11}}{11b}$$

input `Integrate[(a^5 + 5*a^4*b*x + 10*a^3*b^2*x^2 + 10*a^2*b^3*x^3 + 5*a*b^4*x^4 + b^5*x^5)^2, x]`

output `(a + b*x)^11/(11*b)`

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.039$ , Rules used = {2006, 17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2 dx$$

$$\downarrow \text{2006}$$

$$\int (a + bx)^{10} dx$$

$$\downarrow \text{17}$$

$$\frac{(a + bx)^{11}}{11b}$$

input

```
Int[(a^5 + 5*a^4*b*x + 10*a^3*b^2*x^2 + 10*a^2*b^3*x^3 + 5*a*b^4*x^4 + b^5*x^5)^2,x]
```

output

```
(a + b*x)^11/(11*b)
```

**Defintions of rubi rules used**

rule 17

```
Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]
```

rule 2006

```
Int[(u_.)*(Px_), x_Symbol] := With[{a = Rt[Coeff[Px, x, 0], Expon[Px, x]], b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Int[u*(a + b*x)^Expon[Px, x], x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; PolyQ[Px, x] && GtQ[Expon[Px, x], 1] && NeQ[Coeff[Px, x, 0], 0] && !MatchQ[Px, (a_.)*(v_)^Expon[Px, x]] /; FreeQ[a, x] && LinearQ[v, x]]
```

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 108 vs.  $2(12) = 24$ .

Time = 0.03 (sec) , antiderivative size = 109, normalized size of antiderivative = 7.79

method	result
default	$\frac{1}{11}b^{10}x^{11} + ab^9x^{10} + 5a^2b^8x^9 + 15a^3b^7x^8 + 30a^4b^6x^7 + 42a^5b^5x^6 + 42a^6b^4x^5 + 30a^7b^3x^4 +$
norman	$\frac{1}{11}b^{10}x^{11} + ab^9x^{10} + 5a^2b^8x^9 + 15a^3b^7x^8 + 30a^4b^6x^7 + 42a^5b^5x^6 + 42a^6b^4x^5 + 30a^7b^3x^4 +$
risch	$\frac{1}{11}b^{10}x^{11} + ab^9x^{10} + 5a^2b^8x^9 + 15a^3b^7x^8 + 30a^4b^6x^7 + 42a^5b^5x^6 + 42a^6b^4x^5 + 30a^7b^3x^4 +$
parallelrisch	$\frac{1}{11}b^{10}x^{11} + ab^9x^{10} + 5a^2b^8x^9 + 15a^3b^7x^8 + 30a^4b^6x^7 + 42a^5b^5x^6 + 42a^6b^4x^5 + 30a^7b^3x^4 +$
gosper	$\frac{x(b^{10}x^{10} + 11ab^9x^9 + 55a^2b^8x^8 + 165a^3b^7x^7 + 330a^4b^6x^6 + 462a^5b^5x^5 + 462a^6b^4x^4 + 330a^7b^3x^3 + 165a^8b^2x^2 + 55a^9bx + 11a^{10})}{11}$
orering	$\frac{x(b^{10}x^{10} + 11ab^9x^9 + 55a^2b^8x^8 + 165a^3b^7x^7 + 330a^4b^6x^6 + 462a^5b^5x^5 + 462a^6b^4x^4 + 330a^7b^3x^3 + 165a^8b^2x^2 + 55a^9bx + 11a^{10})}{11(bx+a)^{10}}$

input

```
int((b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5)^2,x,
method=_RETURNVERBOSE)
```

output

```
1/11*b^10*x^11+a*b^9*x^10+5*a^2*b^8*x^9+15*a^3*b^7*x^8+30*a^4*b^6*x^7+42*a^5*b^5*x^6+42*a^6*b^4*x^5+30*a^7*b^3*x^4+15*a^8*b^2*x^3+5*a^9*b*x^2+a^10*x
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 108 vs.  $2(12) = 24$ .

Time = 0.08 (sec) , antiderivative size = 108, normalized size of antiderivative = 7.71

$$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2 dx$$

$$= \frac{1}{11}b^{10}x^{11} + ab^9x^{10} + 5a^2b^8x^9 + 15a^3b^7x^8 + 30a^4b^6x^7$$

$$+ 42a^5b^5x^6 + 42a^6b^4x^5 + 30a^7b^3x^4 + 15a^8b^2x^3 + 5a^9bx^2 + a^{10}x$$

input

```
integrate((b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5)^2,x, algorithm="fricas")
```

output

```
1/11*b^10*x^11 + a*b^9*x^10 + 5*a^2*b^8*x^9 + 15*a^3*b^7*x^8 + 30*a^4*b^6*
x^7 + 42*a^5*b^5*x^6 + 42*a^6*b^4*x^5 + 30*a^7*b^3*x^4 + 15*a^8*b^2*x^3 +
5*a^9*b*x^2 + a^10*x
```

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 114 vs.  $2(8) = 16$ .

Time = 0.03 (sec) , antiderivative size = 114, normalized size of antiderivative = 8.14

$$\begin{aligned} & \int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2 dx \\ &= a^{10}x + 5a^9bx^2 + 15a^8b^2x^3 + 30a^7b^3x^4 + 42a^6b^4x^5 + 42a^5b^5x^6 \\ & \quad + 30a^4b^6x^7 + 15a^3b^7x^8 + 5a^2b^8x^9 + ab^9x^{10} + \frac{b^{10}x^{11}}{11} \end{aligned}$$

input

```
integrate((b**5*x**5+5*a*b**4*x**4+10*a**2*b**3*x**3+10*a**3*b**2*x**2+5*a
**4*b*x+a**5)**2,x)
```

output

```
a**10*x + 5*a**9*b*x**2 + 15*a**8*b**2*x**3 + 30*a**7*b**3*x**4 + 42*a**6*
b**4*x**5 + 42*a**5*b**5*x**6 + 30*a**4*b**6*x**7 + 15*a**3*b**7*x**8 + 5*
a**2*b**8*x**9 + a*b**9*x**10 + b**10*x**11/11
```

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 228 vs.  $2(12) = 24$ .

Time = 0.04 (sec) , antiderivative size = 228, normalized size of antiderivative = 16.29

$$\begin{aligned} & \int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2 dx \\ &= \frac{1}{11} b^{10}x^{11} + ab^9x^{10} + \frac{25}{9} a^2b^8x^9 + \frac{100}{7} a^4b^6x^7 + 20 a^6b^4x^5 + \frac{25}{3} a^8b^2x^3 \\ & \quad + a^{10}x + \frac{1}{3} (b^5x^6 + 6ab^4x^5 + 15a^2b^3x^4 + 20a^3b^2x^3 + 15a^4bx^2) a^5 \\ & \quad + \frac{5}{21} (6b^5x^7 + 35ab^4x^6 + 84a^2b^3x^5 + 105a^3b^2x^4) a^4b \\ & \quad + \frac{5}{42} (21b^5x^8 + 120ab^4x^7 + 280a^2b^3x^6) a^3b^2 + \frac{5}{18} (8b^5x^9 + 45ab^4x^8) a^2b^3 \end{aligned}$$

input `integrate((b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5)^2,x, algorithm="maxima")`

output  $1/11*b^{10}*x^{11} + a*b^9*x^{10} + 25/9*a^2*b^8*x^9 + 100/7*a^4*b^6*x^7 + 20*a^6*b^4*x^5 + 25/3*a^8*b^2*x^3 + a^{10}*x + 1/3*(b^5*x^6 + 6*a*b^4*x^5 + 15*a^2*b^3*x^4 + 20*a^3*b^2*x^3 + 15*a^4*b*x^2)*a^5 + 5/21*(6*b^5*x^7 + 35*a*b^4*x^6 + 84*a^2*b^3*x^5 + 105*a^3*b^2*x^4)*a^4*b + 5/42*(21*b^5*x^8 + 120*a*b^4*x^7 + 280*a^2*b^3*x^6)*a^3*b^2 + 5/18*(8*b^5*x^9 + 45*a*b^4*x^8)*a^2*b^3$

### Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 108 vs.  $2(12) = 24$ .

Time = 0.11 (sec) , antiderivative size = 108, normalized size of antiderivative = 7.71

$$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2 dx$$

$$= \frac{1}{11} b^{10} x^{11} + ab^9 x^{10} + 5 a^2 b^8 x^9 + 15 a^3 b^7 x^8 + 30 a^4 b^6 x^7 + 42 a^5 b^5 x^6 + 42 a^6 b^4 x^5 + 30 a^7 b^3 x^4 + 15 a^8 b^2 x^3 + 5 a^9 b x^2 + a^{10} x$$

input `integrate((b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5)^2,x, algorithm="giac")`

output  $1/11*b^{10}*x^{11} + a*b^9*x^{10} + 5*a^2*b^8*x^9 + 15*a^3*b^7*x^8 + 30*a^4*b^6*x^7 + 42*a^5*b^5*x^6 + 42*a^6*b^4*x^5 + 30*a^7*b^3*x^4 + 15*a^8*b^2*x^3 + 5*a^9*b*x^2 + a^{10}*x$

**Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 108, normalized size of antiderivative = 7.71

$$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2 dx$$

$$= a^{10}x + 5a^9bx^2 + 15a^8b^2x^3 + 30a^7b^3x^4 + 42a^6b^4x^5 + 42a^5b^5x^6$$

$$+ 30a^4b^6x^7 + 15a^3b^7x^8 + 5a^2b^8x^9 + ab^9x^{10} + \frac{b^{10}x^{11}}{11}$$

input

```
int((a^5 + b^5*x^5 + 5*a*b^4*x^4 + 10*a^3*b^2*x^2 + 10*a^2*b^3*x^3 + 5*a^4
*b*x)^2,x)
```

output

```
a^10*x + (b^10*x^11)/11 + 5*a^9*b*x^2 + a*b^9*x^10 + 15*a^8*b^2*x^3 + 30*a
^7*b^3*x^4 + 42*a^6*b^4*x^5 + 42*a^5*b^5*x^6 + 30*a^4*b^6*x^7 + 15*a^3*b^7
*x^8 + 5*a^2*b^8*x^9
```

**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 109, normalized size of antiderivative = 7.79

$$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2 dx$$

$$= \frac{x(b^{10}x^{10} + 11ab^9x^9 + 55a^2b^8x^8 + 165a^3b^7x^7 + 330a^4b^6x^6 + 462a^5b^5x^5 + 462a^6b^4x^4 + 330a^7b^3x^3 + 165a^8b^2x^2 + 5a^9bx + a^{10})}{11}$$

input

```
int((b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5)^2,x)
```

output

```
(x*(11*a**10 + 55*a**9*b*x + 165*a**8*b**2*x**2 + 330*a**7*b**3*x**3 + 462
*a**6*b**4*x**4 + 462*a**5*b**5*x**5 + 330*a**4*b**6*x**6 + 165*a**3*b**7*
x**7 + 55*a**2*b**8*x**8 + 11*a*b**9*x**9 + b**10*x**10))/11
```



### 3.21 $\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)$

Optimal result	160
Mathematica [B] (verified)	160
Rubi [A] (verified)	161
Maple [A] (verified)	162
Fricas [B] (verification not implemented)	162
Sympy [B] (verification not implemented)	163
Maxima [B] (verification not implemented)	163
Giac [B] (verification not implemented)	164
Mupad [B] (verification not implemented)	164
Reduce [B] (verification not implemented)	165

#### Optimal result

Integrand size = 49, antiderivative size = 14

$$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5) dx = \frac{(a + bx)^6}{6b}$$

output `1/6*(b*x+a)^6/b`

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 61 vs.  $2(14) = 28$ .

Time = 0.00 (sec) , antiderivative size = 61, normalized size of antiderivative = 4.36

$$\begin{aligned} & \int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5) dx \\ &= a^5x + \frac{5}{2}a^4bx^2 + \frac{10}{3}a^3b^2x^3 + \frac{5}{2}a^2b^3x^4 + ab^4x^5 + \frac{b^5x^6}{6} \end{aligned}$$

input `Integrate[a^5 + 5*a^4*b*x + 10*a^3*b^2*x^2 + 10*a^2*b^3*x^3 + 5*a*b^4*x^4 + b^5*x^5, x]`

output

$$a^5x + (5a^4bx^2)/2 + (10a^3b^2x^3)/3 + (5a^2b^3x^4)/2 + ab^4x^5 + (b^5x^6)/6$$

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.041$ , Rules used = {2006, 17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5) dx$$

$$\downarrow 2006$$

$$\int (a + bx)^5 dx$$

$$\downarrow 17$$

$$\frac{(a + bx)^6}{6b}$$

input

```
Int[a^5 + 5*a^4*b*x + 10*a^3*b^2*x^2 + 10*a^2*b^3*x^3 + 5*a*b^4*x^4 + b^5*x^5, x]
```

output

```
(a + b*x)^6/(6*b)
```

**Defintions of rubi rules used**

rule 17

```
Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]
```

rule 2006

```
Int[(u_.)*(Px_), x_Symbol] := With[{a = Rt[Coeff[Px, x, 0], Expon[Px, x]],
b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Int[u*(a + b*x)^Expon[Px,
x], x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; PolyQ[Px, x] && GtQ[Expon[P
x, x], 1] && NeQ[Coeff[Px, x, 0], 0] && !MatchQ[Px, (a_.)*(v_)^Expon[Px, x
] /; FreeQ[a, x] && LinearQ[v, x]]
```

**Maple [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

method	result	size
default	$\frac{(bx+a)^6}{6b}$	13
norman	$\frac{1}{6}b^5x^6 + ab^4x^5 + \frac{5}{2}a^2b^3x^4 + \frac{10}{3}a^3b^2x^3 + \frac{5}{2}a^4bx^2 + xa^5$	54
risch	$\frac{1}{6}b^5x^6 + ab^4x^5 + \frac{5}{2}a^2b^3x^4 + \frac{10}{3}a^3b^2x^3 + \frac{5}{2}a^4bx^2 + xa^5$	54
parallelrisch	$\frac{1}{6}b^5x^6 + ab^4x^5 + \frac{5}{2}a^2b^3x^4 + \frac{10}{3}a^3b^2x^3 + \frac{5}{2}a^4bx^2 + xa^5$	54
parts	$\frac{1}{6}b^5x^6 + ab^4x^5 + \frac{5}{2}a^2b^3x^4 + \frac{10}{3}a^3b^2x^3 + \frac{5}{2}a^4bx^2 + xa^5$	54
gosper	$\frac{x(b^5x^5+6b^4ax^4+15a^2b^3x^3+20b^2x^2a^3+15a^4bx+6a^5)}{6}$	55
orering	$\frac{x(b^5x^5+6b^4ax^4+15a^2b^3x^3+20b^2x^2a^3+15a^4bx+6a^5)(b^5x^5+5b^4ax^4+10a^2b^3x^3+10b^2x^2a^3+5a^4bx+a^5)}{6(bx+a)^5}$	111

input

```
int(b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5,x,method=
_RETURNVERBOSE)
```

output

```
1/6*(b*x+a)^6/b
```

**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 53 vs.  $2(12) = 24$ .

Time = 0.06 (sec) , antiderivative size = 53, normalized size of antiderivative = 3.79

$$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5) dx$$

$$= \frac{1}{6}b^5x^6 + ab^4x^5 + \frac{5}{2}a^2b^3x^4 + \frac{10}{3}a^3b^2x^3 + \frac{5}{2}a^4bx^2 + a^5x$$

input `integrate(b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5, x, algorithm="fricas")`

output `1/6*b^5*x^6 + a*b^4*x^5 + 5/2*a^2*b^3*x^4 + 10/3*a^3*b^2*x^3 + 5/2*a^4*b*x^2 + a^5*x`

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 60 vs.  $2(8) = 16$ .

Time = 0.02 (sec) , antiderivative size = 60, normalized size of antiderivative = 4.29

$$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5) dx$$

$$= a^5x + \frac{5a^4bx^2}{2} + \frac{10a^3b^2x^3}{3} + \frac{5a^2b^3x^4}{2} + ab^4x^5 + \frac{b^5x^6}{6}$$

input `integrate(b**5*x**5+5*a*b**4*x**4+10*a**2*b**3*x**3+10*a**3*b**2*x**2+5*a**4*b*x+a**5,x)`

output `a**5*x + 5*a**4*b*x**2/2 + 10*a**3*b**2*x**3/3 + 5*a**2*b**3*x**4/2 + a*b**4*x**5 + b**5*x**6/6`

### Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 53 vs.  $2(12) = 24$ .

Time = 0.03 (sec) , antiderivative size = 53, normalized size of antiderivative = 3.79

$$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5) dx$$

$$= \frac{1}{6}b^5x^6 + ab^4x^5 + \frac{5}{2}a^2b^3x^4 + \frac{10}{3}a^3b^2x^3 + \frac{5}{2}a^4bx^2 + a^5x$$

input `integrate(b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5, x, algorithm="maxima")`

output

$$\frac{1}{6}b^5x^6 + ab^4x^5 + \frac{5}{2}a^2b^3x^4 + \frac{10}{3}a^3b^2x^3 + \frac{5}{2}a^4bx^2 + a^5x$$

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 53 vs.  $2(12) = 24$ .

Time = 0.11 (sec) , antiderivative size = 53, normalized size of antiderivative = 3.79

$$\begin{aligned} & \int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5) dx \\ &= \frac{1}{6}b^5x^6 + ab^4x^5 + \frac{5}{2}a^2b^3x^4 + \frac{10}{3}a^3b^2x^3 + \frac{5}{2}a^4bx^2 + a^5x \end{aligned}$$

input

```
integrate(b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5,
x, algorithm="giac")
```

output

$$\frac{1}{6}b^5x^6 + ab^4x^5 + \frac{5}{2}a^2b^3x^4 + \frac{10}{3}a^3b^2x^3 + \frac{5}{2}a^4bx^2 + a^5x$$

**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 53, normalized size of antiderivative = 3.79

$$\begin{aligned} & \int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5) dx \\ &= a^5x + \frac{5a^4bx^2}{2} + \frac{10a^3b^2x^3}{3} + \frac{5a^2b^3x^4}{2} + ab^4x^5 + \frac{b^5x^6}{6} \end{aligned}$$

input

```
int(a^5 + b^5*x^5 + 5*a*b^4*x^4 + 10*a^3*b^2*x^2 + 10*a^2*b^3*x^3 + 5*a^4*
b*x,x)
```

output

$$a^5x + (b^5x^6)/6 + (5a^4bx^2)/2 + ab^4x^5 + (10a^3b^2x^3)/3 + (5a^2b^3x^4)/2$$

**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 54, normalized size of antiderivative = 3.86

$$\int (a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5) dx$$
$$= \frac{x(b^5x^5 + 6ab^4x^4 + 15a^2b^3x^3 + 20a^3b^2x^2 + 15a^4bx + 6a^5)}{6}$$

input `int(b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5,x)`

output `(x*(6*a**5 + 15*a**4*b*x + 20*a**3*b**2*x**2 + 15*a**2*b**3*x**3 + 6*a*b**4*x**4 + b**5*x**5))/6`

$$3.22 \quad \int \frac{1}{a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5} dx$$

Optimal result . . . . .	166
Mathematica [A] (verified) . . . . .	166
Rubi [A] (verified) . . . . .	167
Maple [A] (verified) . . . . .	168
Fricas [B] (verification not implemented) . . . . .	168
Sympy [B] (verification not implemented) . . . . .	169
Maxima [B] (verification not implemented) . . . . .	169
Giac [A] (verification not implemented) . . . . .	170
Mupad [B] (verification not implemented) . . . . .	170
Reduce [B] (verification not implemented) . . . . .	170

### Optimal result

Integrand size = 51, antiderivative size = 14

$$\int \frac{1}{a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5} dx = -\frac{1}{4b(a + bx)^4}$$

output

```
-1/4/b/(b*x+a)^4
```

### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5} dx = -\frac{1}{4b(a + bx)^4}$$

input

```
Integrate[(a^5 + 5*a^4*b*x + 10*a^3*b^2*x^2 + 10*a^2*b^3*x^3 + 5*a*b^4*x^4 + b^5*x^5)^(-1), x]
```

output

```
-1/4*1/(b*(a + b*x)^4)
```

**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.039$ , Rules used = {2007, 17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5} dx$$

↓ 2007

$$\int \frac{1}{(a + bx)^5} dx$$

↓ 17

$$-\frac{1}{4b(a + bx)^4}$$

input

```
Int[(a^5 + 5*a^4*b*x + 10*a^3*b^2*x^2 + 10*a^2*b^3*x^3 + 5*a*b^4*x^4 + b^5*x^5)^(-1),x]
```

output

```
-1/4*1/(b*(a + b*x)^4)
```

**Defintions of rubi rules used**

rule 17

```
Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]
```

rule 2007

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{a = Rt[Coeff[Px, x, 0], Expon[Px, x]], b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Int[u*(a + b*x)^(Expon[Px, x]*p), x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; IntegerQ[p] && PolyQ[Px, x] && GtQ[Expon[Px, x], 1] && NeQ[Coeff[Px, x, 0], 0]
```



**Maple [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

method	result	size
default	$-\frac{1}{4b(bx+a)^4}$	13
norman	$-\frac{1}{4b(bx+a)^4}$	13
gosper	$-\frac{1}{4b(b^4x^4+4ab^3x^3+6a^2b^2x^2+4ba^3x+a^4)}$	46
risch	$-\frac{1}{4b(b^4x^4+4ab^3x^3+6a^2b^2x^2+4ba^3x+a^4)}$	46
parallelrisc	$-\frac{1}{4b(b^4x^4+4ab^3x^3+6a^2b^2x^2+4ba^3x+a^4)}$	46
orering	$-\frac{bx+a}{4b(b^5x^5+5b^4ax^4+10a^2b^3x^3+10b^2x^2a^3+5a^4bx+a^5)}$	62

input `int(1/(b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5),x,method=_RETURNVERBOSE)`

output `-1/4/b/(b*x+a)^4`

**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 46 vs.  $2(12) = 24$ .

Time = 0.07 (sec) , antiderivative size = 46, normalized size of antiderivative = 3.29

$$\int \frac{1}{a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5} dx$$

$$= -\frac{1}{4(b^5x^4 + 4ab^4x^3 + 6a^2b^3x^2 + 4a^3b^2x + a^4b)}$$

input `integrate(1/(b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5),x,algorithm="fricas")`

output `-1/4/(b^5*x^4 + 4*a*b^4*x^3 + 6*a^2*b^3*x^2 + 4*a^3*b^2*x + a^4*b)`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 49 vs.  $2(12) = 24$ .

Time = 0.14 (sec) , antiderivative size = 49, normalized size of antiderivative = 3.50

$$\int \frac{1}{a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5} dx$$

$$= -\frac{1}{4a^4b + 16a^3b^2x + 24a^2b^3x^2 + 16ab^4x^3 + 4b^5x^4}$$

input

```
integrate(1/(b**5*x**5+5*a*b**4*x**4+10*a**2*b**3*x**3+10*a**3*b**2*x**2+5*a**4*b*x+a**5),x)
```

output

```
-1/(4*a**4*b + 16*a**3*b**2*x + 24*a**2*b**3*x**2 + 16*a*b**4*x**3 + 4*b**5*x**4)
```

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 46 vs.  $2(12) = 24$ .

Time = 0.03 (sec) , antiderivative size = 46, normalized size of antiderivative = 3.29

$$\int \frac{1}{a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5} dx$$

$$= -\frac{1}{4(b^5x^4 + 4ab^4x^3 + 6a^2b^3x^2 + 4a^3b^2x + a^4b)}$$

input

```
integrate(1/(b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5),x, algorithm="maxima")
```

output

```
-1/4/(b^5*x^4 + 4*a*b^4*x^3 + 6*a^2*b^3*x^2 + 4*a^3*b^2*x + a^4*b)
```

**Giac [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5} dx = -\frac{1}{4(bx + a)^4b}$$

input `integrate(1/(b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5),x, algorithm="giac")`

output `-1/4/((b*x + a)^4*b)`

**Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 48, normalized size of antiderivative = 3.43

$$\int \frac{1}{a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5} dx$$

$$= -\frac{1}{4a^4b + 16a^3b^2x + 24a^2b^3x^2 + 16ab^4x^3 + 4b^5x^4}$$

input `int(1/(a^5 + b^5*x^5 + 5*a*b^4*x^4 + 10*a^3*b^2*x^2 + 10*a^2*b^3*x^3 + 5*a^4*b*x), x)`

output `-1/(4*a^4*b + 4*b^5*x^4 + 16*a^3*b^2*x + 16*a*b^4*x^3 + 24*a^2*b^3*x^2)`

**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 45, normalized size of antiderivative = 3.21

$$\int \frac{1}{a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5} dx$$

$$= -\frac{1}{4b(b^4x^4 + 4ab^3x^3 + 6a^2b^2x^2 + 4a^3bx + a^4)}$$

input `int(1/(b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5), x)`

output  $(-1)/(4*b*(a**4 + 4*a**3*b*x + 6*a**2*b**2*x**2 + 4*a*b**3*x**3 + b**4*x**4))$

**3.23**  $\int \frac{1}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2} dx$

Optimal result . . . . .	172
Mathematica [A] (verified) . . . . .	172
Rubi [A] (verified) . . . . .	173
Maple [A] (verified) . . . . .	174
Fricas [B] (verification not implemented) . . . . .	174
Sympy [B] (verification not implemented) . . . . .	175
Maxima [B] (verification not implemented) . . . . .	175
Giac [A] (verification not implemented) . . . . .	176
Mupad [B] (verification not implemented) . . . . .	176
Reduce [B] (verification not implemented) . . . . .	177

**Optimal result**

Integrand size = 51, antiderivative size = 14

$$\int \frac{1}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2} dx = -\frac{1}{9b(a + bx)^9}$$

output -1/9/b/(b\*x+a)^9

**Mathematica [A] (verified)**

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2} dx = -\frac{1}{9b(a + bx)^9}$$

input Integrate[(a^5 + 5\*a^4\*b\*x + 10\*a^3\*b^2\*x^2 + 10\*a^2\*b^3\*x^3 + 5\*a\*b^4\*x^4 + b^5\*x^5)^(-2), x]

output -1/9\*1/(b\*(a + b\*x)^9)

**Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.039$ , Rules used = {2007, 17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2} dx$$

↓ 2007

$$\int \frac{1}{(a + bx)^{10}} dx$$

↓ 17

$$-\frac{1}{9b(a + bx)^9}$$

input

```
Int[(a^5 + 5*a^4*b*x + 10*a^3*b^2*x^2 + 10*a^2*b^3*x^3 + 5*a*b^4*x^4 + b^5*x^5)^(-2), x]
```

output

```
-1/9*1/(b*(a + b*x)^9)
```

**Defintions of rubi rules used**

rule 17

```
Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]
```

rule 2007

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{a = Rt[Coeff[Px, x, 0], Expon[Px, x]], b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Int[u*(a + b*x)^(Expon[Px, x]*p), x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; IntegerQ[p] && PolyQ[Px, x] && GtQ[Expon[Px, x], 1] && NeQ[Coeff[Px, x, 0], 0]
```

**Maple [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

method	result	size
default	$-\frac{1}{9b(bx+a)^9}$	13
norman	$-\frac{1}{9b(bx+a)^9}$	13
risch	$-\frac{1}{9b(b^4x^4+4ab^3x^3+6a^2b^2x^2+4ba^3x+a^4)^2(bx+a)}$	53
orering	$-\frac{bx+a}{9b(b^5x^5+5b^4ax^4+10a^2b^3x^3+10b^2x^2a^3+5a^4bx+a^5)^2}$	62
gospers	$-\frac{1}{9(b^4x^4+4ab^3x^3+6a^2b^2x^2+4ba^3x+a^4)(b^5x^5+5b^4ax^4+10a^2b^3x^3+10b^2x^2a^3+5a^4bx+a^5)b}$	97
parallelrisch	$-\frac{1}{9(b^4x^4+4ab^3x^3+6a^2b^2x^2+4ba^3x+a^4)(b^5x^5+5b^4ax^4+10a^2b^3x^3+10b^2x^2a^3+5a^4bx+a^5)b}$	97

input `int(1/(b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5)^2, x, method=_RETURNVERBOSE)`

output `-1/9/b/(b*x+a)^9`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 101 vs. 2(12) = 24.

Time = 0.09 (sec) , antiderivative size = 101, normalized size of antiderivative = 7.21

$$\int \frac{1}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2} dx =$$

$$-\frac{1}{9(b^{10}x^9 + 9ab^9x^8 + 36a^2b^8x^7 + 84a^3b^7x^6 + 126a^4b^6x^5 + 126a^5b^5x^4 + 84a^6b^4x^3 + 36a^7b^3x^2 + 9a^8b^2x + a^9b)}$$

input `integrate(1/(b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5)^2,x, algorithm="fricas")`

output `-1/9/(b^10*x^9 + 9*a*b^9*x^8 + 36*a^2*b^8*x^7 + 84*a^3*b^7*x^6 + 126*a^4*b^6*x^5 + 126*a^5*b^5*x^4 + 84*a^6*b^4*x^3 + 36*a^7*b^3*x^2 + 9*a^8*b^2*x + a^9*b)`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 109 vs.  $2(12) = 24$ .

Time = 0.30 (sec) , antiderivative size = 109, normalized size of antiderivative = 7.79

$$\int \frac{1}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2} dx =$$

$$\frac{1}{9a^9b + 81a^8b^2x + 324a^7b^3x^2 + 756a^6b^4x^3 + 1134a^5b^5x^4 + 1134a^4b^6x^5 + 756a^3b^7x^6 + 324a^2b^8x^7 + 81ab^9x^8 + 9b^{10}x^9}$$

input

```
integrate(1/(b**5*x**5+5*a*b**4*x**4+10*a**2*b**3*x**3+10*a**3*b**2*x**2+5*a**4*b*x+a**5)**2,x)
```

output

```
-1/(9*a**9*b + 81*a**8*b**2*x + 324*a**7*b**3*x**2 + 756*a**6*b**4*x**3 + 1134*a**5*b**5*x**4 + 1134*a**4*b**6*x**5 + 756*a**3*b**7*x**6 + 324*a**2*b**8*x**7 + 81*a*b**9*x**8 + 9*b**10*x**9)
```

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 101 vs.  $2(12) = 24$ .

Time = 0.03 (sec) , antiderivative size = 101, normalized size of antiderivative = 7.21

$$\int \frac{1}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2} dx =$$

$$\frac{1}{9(b^{10}x^9 + 9ab^9x^8 + 36a^2b^8x^7 + 84a^3b^7x^6 + 126a^4b^6x^5 + 126a^5b^5x^4 + 84a^6b^4x^3 + 36a^7b^3x^2 + 9a^8b^2x + a^9b)}$$

input

```
integrate(1/(b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5)^2,x, algorithm="maxima")
```

output

```
-1/9/(b^10*x^9 + 9*a*b^9*x^8 + 36*a^2*b^8*x^7 + 84*a^3*b^7*x^6 + 126*a^4*b^6*x^5 + 126*a^5*b^5*x^4 + 84*a^6*b^4*x^3 + 36*a^7*b^3*x^2 + 9*a^8*b^2*x + a^9*b)
```



**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2} dx = -\frac{1}{9(bx + a)^9b}$$

input

```
integrate(1/(b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5)^2,x, algorithm="giac")
```

output

```
-1/9/((b*x + a)^9*b)
```

**Mupad [B] (verification not implemented)**

Time = 22.51 (sec) , antiderivative size = 103, normalized size of antiderivative = 7.36

$$\int \frac{1}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2} dx = \frac{1}{9a^9b + 81a^8b^2x + 324a^7b^3x^2 + 756a^6b^4x^3 + 1134a^5b^5x^4 + 1134a^4b^6x^5 + 756a^3b^7x^6 + 324a^2b^8x^7}$$

input

```
int(1/(a^5 + b^5*x^5 + 5*a*b^4*x^4 + 10*a^3*b^2*x^2 + 10*a^2*b^3*x^3 + 5*a^4*b*x)^2,x)
```

output

```
-1/(9*a^9*b + 9*b^10*x^9 + 81*a^8*b^2*x + 81*a*b^9*x^8 + 324*a^7*b^3*x^2 + 756*a^6*b^4*x^3 + 1134*a^5*b^5*x^4 + 1134*a^4*b^6*x^5 + 756*a^3*b^7*x^6 + 324*a^2*b^8*x^7)
```

**Reduce [B] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 100, normalized size of antiderivative = 7.14

$$\int \frac{1}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^2} dx =$$

$$\frac{1}{9b(b^9x^9 + 9ab^8x^8 + 36a^2b^7x^7 + 84a^3b^6x^6 + 126a^4b^5x^5 + 126a^5b^4x^4 + 84a^6b^3x^3 + 36a^7b^2x^2 + 9a^8bx + a^9)}$$

input

```
int(1/(b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5)^2,
x)
```

output

```
( - 1)/(9*b*(a**9 + 9*a**8*b*x + 36*a**7*b**2*x**2 + 84*a**6*b**3*x**3 + 1
26*a**5*b**4*x**4 + 126*a**4*b**5*x**5 + 84*a**3*b**6*x**6 + 36*a**2*b**7*
x**7 + 9*a*b**8*x**8 + b**9*x**9))
```

$$3.24 \quad \int \frac{1}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3} dx$$

Optimal result	178
Mathematica [A] (verified)	178
Rubi [A] (verified)	179
Maple [A] (verified)	180
Fricas [B] (verification not implemented)	180
Sympy [B] (verification not implemented)	181
Maxima [B] (verification not implemented)	181
Giac [A] (verification not implemented)	182
Mupad [B] (verification not implemented)	182
Reduce [B] (verification not implemented)	183

### Optimal result

Integrand size = 51, antiderivative size = 14

$$\int \frac{1}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3} dx = -\frac{1}{14b(a + bx)^{14}}$$

output `-1/14/b/(b*x+a)^14`

### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3} dx = -\frac{1}{14b(a + bx)^{14}}$$

input `Integrate[(a^5 + 5*a^4*b*x + 10*a^3*b^2*x^2 + 10*a^2*b^3*x^3 + 5*a*b^4*x^4 + b^5*x^5)^(-3), x]`

output `-1/14*1/(b*(a + b*x)^14)`

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.039$ , Rules used = {2007, 17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3} dx$$

↓ 2007

$$\int \frac{1}{(a + bx)^{15}} dx$$

↓ 17

$$-\frac{1}{14b(a + bx)^{14}}$$

input

```
Int[(a^5 + 5*a^4*b*x + 10*a^3*b^2*x^2 + 10*a^2*b^3*x^3 + 5*a*b^4*x^4 + b^5*x^5)^(-3), x]
```

output

```
-1/14*1/(b*(a + b*x)^14)
```

**Defintions of rubi rules used**

rule 17

```
Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]
```

rule 2007

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{a = Rt[Coeff[Px, x, 0], Expon[Px, x]], b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Int[u*(a + b*x)^(Expon[Px, x]*p), x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; IntegerQ[p] && PolyQ[Px, x] && GtQ[Expon[Px, x], 1] && NeQ[Coeff[Px, x, 0], 0]
```

**Maple [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

method	result	size
default	$-\frac{1}{14b(bx+a)^{14}}$	13
norman	$-\frac{1}{14b(bx+a)^{14}}$	13
risch	$-\frac{1}{14b(b^4x^4+4ab^3x^3+6a^2b^2x^2+4ba^3x+a^4)^3(bx+a)^2}$	53
orering	$-\frac{bx+a}{14b(b^5x^5+5b^4ax^4+10a^2b^3x^3+10b^2x^2a^3+5a^4bx+a^5)^3}$	62
gospers	$-\frac{1}{14(b^4x^4+4ab^3x^3+6a^2b^2x^2+4ba^3x+a^4)(b^5x^5+5b^4ax^4+10a^2b^3x^3+10b^2x^2a^3+5a^4bx+a^5)^2b}$	97
parallelrisch	$-\frac{1}{14(b^4x^4+4ab^3x^3+6a^2b^2x^2+4ba^3x+a^4)(b^5x^5+5b^4ax^4+10a^2b^3x^3+10b^2x^2a^3+5a^4bx+a^5)^2b}$	97

input `int(1/(b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5)^3, x, method=_RETURNVERBOSE)`

output `-1/14/b/(b*x+a)^14`

**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 156 vs.  $2(12) = 24$ .

Time = 0.07 (sec) , antiderivative size = 156, normalized size of antiderivative = 11.14

$$\int \frac{1}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3} dx =$$

$$-\frac{1}{14(b^{15}x^{14} + 14ab^{14}x^{13} + 91a^2b^{13}x^{12} + 364a^3b^{12}x^{11} + 1001a^4b^{11}x^{10} + 2002a^5b^{10}x^9 + 3003a^6b^9x^8 + 3003a^7b^8x^7 + 1401a^8b^7x^6 + 2802a^9b^6x^5 + 1401a^{10}b^5x^4 + 2802a^{11}b^4x^3 + 1401a^{12}b^3x^2 + 2802a^{13}b^2x + 1401a^{14}b)x^{13} + \dots}$$

input `integrate(1/(b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5)^3,x, algorithm="fricas")`

output

$$-1/14/(b^{15}x^{14} + 14a*b^{14}x^{13} + 91a^2*b^{13}x^{12} + 364a^3*b^{12}x^{11} + 1001a^4*b^{11}x^{10} + 2002a^5*b^{10}x^9 + 3003a^6*b^9x^8 + 3432a^7*b^8x^7 + 3003a^8*b^7x^6 + 2002a^9*b^6x^5 + 1001a^{10}b^5x^4 + 364a^{11}b^4x^3 + 91a^{12}b^3x^2 + 14a^{13}b^2x + a^{14}b)$$

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 168 vs.  $2(12) = 24$ .

Time = 0.45 (sec) , antiderivative size = 168, normalized size of antiderivative = 12.00

$$\int \frac{1}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3} dx =$$

$$\frac{-14a^{14}b + 196a^{13}b^2x + 1274a^{12}b^3x^2 + 5096a^{11}b^4x^3 + 14014a^{10}b^5x^4 + 28028a^9b^6x^5 + 42042a^8b^7x^6 + 48048a^7b^8x^7 + 42042a^6b^9x^8 + 28028a^5b^{10}x^9 + 14014a^4b^{11}x^{10} + 5096a^3b^{12}x^{11} + 1274a^2b^{13}x^{12} + 196ab^{14}x^{13} + 14b^{15}x^{14}}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3}$$

input

```
integrate(1/(b**5*x**5+5*a*b**4*x**4+10*a**2*b**3*x**3+10*a**3*b**2*x**2+
*a**4*b*x+a**5)**3,x)
```

output

$$-1/(14a^{14}b + 196a^{13}b^2x + 1274a^{12}b^3x^2 + 5096a^{11}b^4x^3 + 14014a^{10}b^5x^4 + 28028a^9b^6x^5 + 42042a^8b^7x^6 + 48048a^7b^8x^7 + 42042a^6b^9x^8 + 28028a^5b^{10}x^9 + 14014a^4b^{11}x^{10} + 5096a^3b^{12}x^{11} + 1274a^2b^{13}x^{12} + 196a^{14}b + 14b^{15}x^{14})$$

### Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 156 vs.  $2(12) = 24$ .

Time = 0.04 (sec) , antiderivative size = 156, normalized size of antiderivative = 11.14

$$\int \frac{1}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3} dx =$$

$$\frac{-14(b^{15}x^{14} + 14ab^{14}x^{13} + 91a^2b^{13}x^{12} + 364a^3b^{12}x^{11} + 1001a^4b^{11}x^{10} + 2002a^5b^{10}x^9 + 3003a^6b^9x^8 + 3432a^7b^8x^7 + 3003a^8b^7x^6 + 2002a^9b^6x^5 + 1001a^{10}b^5x^4 + 364a^{11}b^4x^3 + 91a^{12}b^3x^2 + 14a^{13}b^2x + a^{14}b)}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3}$$

input

```
integrate(1/(b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a
^5)^3,x, algorithm="maxima")
```

output

```
-1/14/(b^15*x^14 + 14*a*b^14*x^13 + 91*a^2*b^13*x^12 + 364*a^3*b^12*x^11 +
1001*a^4*b^11*x^10 + 2002*a^5*b^10*x^9 + 3003*a^6*b^9*x^8 + 3432*a^7*b^8*
x^7 + 3003*a^8*b^7*x^6 + 2002*a^9*b^6*x^5 + 1001*a^10*b^5*x^4 + 364*a^11*b
^4*x^3 + 91*a^12*b^3*x^2 + 14*a^13*b^2*x + a^14*b)
```

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3} dx = -\frac{1}{14(bx + a)^{14}b}$$

input

```
integrate(1/(b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a
^5)^3,x, algorithm="giac")
```

output

```
-1/14/((b*x + a)^14*b)
```

**Mupad [B] (verification not implemented)**

Time = 23.38 (sec) , antiderivative size = 158, normalized size of antiderivative = 11.29

$$\int \frac{1}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3} dx =$$

$$-\frac{1}{14a^{14}b + 196a^{13}b^2x + 1274a^{12}b^3x^2 + 5096a^{11}b^4x^3 + 14014a^{10}b^5x^4 + 28028a^9b^6x^5 + 42042a^8b^7x^6 + 3432a^7b^8x^7 + 2002a^6b^9x^8 + 91a^5b^{10}x^9 + 14a^4b^{11}x^{10} + a^3b^{12}x^{11} + 14a^2b^{13}x^{12} + a^{14}b^{14}}$$

input

```
int(1/(a^5 + b^5*x^5 + 5*a*b^4*x^4 + 10*a^3*b^2*x^2 + 10*a^2*b^3*x^3 + 5*a
^4*b*x)^3,x)
```

output

```
-1/(14*a^14*b + 14*b^15*x^14 + 196*a^13*b^2*x + 196*a*b^14*x^13 + 1274*a^1
2*b^3*x^2 + 5096*a^11*b^4*x^3 + 14014*a^10*b^5*x^4 + 28028*a^9*b^6*x^5 + 4
2042*a^8*b^7*x^6 + 48048*a^7*b^8*x^7 + 42042*a^6*b^9*x^8 + 28028*a^5*b^10*
x^9 + 14014*a^4*b^11*x^10 + 5096*a^3*b^12*x^11 + 1274*a^2*b^13*x^12)
```

**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 155, normalized size of antiderivative = 11.07

$$\int \frac{1}{(a^5 + 5a^4bx + 10a^3b^2x^2 + 10a^2b^3x^3 + 5ab^4x^4 + b^5x^5)^3} dx =$$

$$\frac{1}{14b(b^{14}x^{14} + 14ab^{13}x^{13} + 91a^2b^{12}x^{12} + 364a^3b^{11}x^{11} + 1001a^4b^{10}x^{10} + 2002a^5b^9x^9 + 3003a^6b^8x^8 + 3432a^7b^7x^7 + 3003a^8b^6x^6 + 1001a^9b^5x^5 + 14a^{10}b^4x^4 + 14a^{11}b^3x^3 + 14a^{12}b^2x^2 + 14a^{13}bx + 14a^{14})}$$

input

```
int(1/(b^5*x^5+5*a*b^4*x^4+10*a^2*b^3*x^3+10*a^3*b^2*x^2+5*a^4*b*x+a^5)^3,
x)
```

output

```
( - 1)/(14*b*(a**14 + 14*a**13*b*x + 91*a**12*b**2*x**2 + 364*a**11*b**3*x**3 + 1001*a**10*b**4*x**4 + 2002*a**9*b**5*x**5 + 3003*a**8*b**6*x**6 + 3432*a**7*b**7*x**7 + 3003*a**6*b**8*x**8 + 2002*a**5*b**9*x**9 + 1001*a**4*b**10*x**10 + 364*a**3*b**11*x**11 + 91*a**2*b**12*x**12 + 14*a*b**13*x**13 + b**14*x**14))
```



### 3.25 $\int \frac{1}{1+x^2+x^3+x^5} dx$

Optimal result . . . . .	184
Mathematica [A] (verified) . . . . .	184
Rubi [A] (verified) . . . . .	185
Maple [A] (verified) . . . . .	186
Fricas [A] (verification not implemented) . . . . .	186
Sympy [A] (verification not implemented) . . . . .	186
Maxima [A] (verification not implemented) . . . . .	187
Giac [A] (verification not implemented) . . . . .	187
Mupad [B] (verification not implemented) . . . . .	187
Reduce [B] (verification not implemented) . . . . .	188

#### Optimal result

Integrand size = 13, antiderivative size = 38

$$\int \frac{1}{1+x^2+x^3+x^5} dx = \frac{\arctan(x)}{2} + \frac{1}{6} \log(1+x) + \frac{1}{4} \log(1+x^2) - \frac{1}{3} \log(1-x+x^2)$$

output `1/2*arctan(x)+1/6*ln(1+x)+1/4*ln(x^2+1)-1/3*ln(x^2-x+1)`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x^2+x^3+x^5} dx = \frac{\arctan(x)}{2} + \frac{1}{6} \log(1+x) + \frac{1}{4} \log(1+x^2) - \frac{1}{3} \log(1-x+x^2)$$

input `Integrate[(1 + x^2 + x^3 + x^5)^(-1),x]`

output `ArcTan[x]/2 + Log[1 + x]/6 + Log[1 + x^2]/4 - Log[1 - x + x^2]/3`

**Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^5 + x^3 + x^2 + 1} dx$$

$$\downarrow \text{2462}$$

$$\int \left( \frac{1-2x}{3(x^2-x+1)} + \frac{x+1}{2(x^2+1)} + \frac{1}{6(x+1)} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{\arctan(x)}{2} + \frac{1}{4} \log(x^2+1) - \frac{1}{3} \log(x^2-x+1) + \frac{1}{6} \log(x+1)$$

input

```
Int[(1 + x^2 + x^3 + x^5)^(-1),x]
```

output

```
ArcTan[x]/2 + Log[1 + x]/6 + Log[1 + x^2]/4 - Log[1 - x + x^2]/3
```

**Defintions of rubi rules used**

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2462

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegr
and[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ
[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0
] && RationalFunctionQ[u, x]
```

**Maple [A] (verified)**

Time = 0.07 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.82

method	result	size
default	$\frac{\arctan(x)}{2} + \frac{\ln(x+1)}{6} + \frac{\ln(x^2+1)}{4} - \frac{\ln(x^2-x+1)}{3}$	31
risch	$\frac{\arctan(x)}{2} + \frac{\ln(x+1)}{6} + \frac{\ln(x^2+1)}{4} - \frac{\ln(x^2-x+1)}{3}$	31
parallelrisc	$\frac{\ln(x+1)}{6} + \frac{\ln(x-i)}{4} - \frac{i \ln(x-i)}{4} + \frac{\ln(x+i)}{4} + \frac{i \ln(x+i)}{4} - \frac{\ln(x^2-x+1)}{3}$	49

input `int(1/(x^5+x^3+x^2+1),x,method=_RETURNVERBOSE)`output `1/2*arctan(x)+1/6*ln(x+1)+1/4*ln(x^2+1)-1/3*ln(x^2-x+1)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.79

$$\int \frac{1}{1+x^2+x^3+x^5} dx = \frac{1}{2} \arctan(x) - \frac{1}{3} \log(x^2-x+1) + \frac{1}{4} \log(x^2+1) + \frac{1}{6} \log(x+1)$$

input `integrate(1/(x^5+x^3+x^2+1),x, algorithm="fricas")`output `1/2*arctan(x) - 1/3*log(x^2 - x + 1) + 1/4*log(x^2 + 1) + 1/6*log(x + 1)`**Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.76

$$\int \frac{1}{1+x^2+x^3+x^5} dx = \frac{\log(x+1)}{6} + \frac{\log(x^2+1)}{4} - \frac{\log(x^2-x+1)}{3} + \frac{\operatorname{atan}(x)}{2}$$

input `integrate(1/(x**5+x**3+x**2+1),x)`output `log(x + 1)/6 + log(x**2 + 1)/4 - log(x**2 - x + 1)/3 + atan(x)/2`

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.79

$$\int \frac{1}{1+x^2+x^3+x^5} dx = \frac{1}{2} \arctan(x) - \frac{1}{3} \log(x^2-x+1) + \frac{1}{4} \log(x^2+1) + \frac{1}{6} \log(x+1)$$

input `integrate(1/(x^5+x^3+x^2+1),x, algorithm="maxima")`output `1/2*arctan(x) - 1/3*log(x^2 - x + 1) + 1/4*log(x^2 + 1) + 1/6*log(x + 1)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.82

$$\int \frac{1}{1+x^2+x^3+x^5} dx = \frac{1}{2} \arctan(x) - \frac{1}{3} \log(x^2-x+1) + \frac{1}{4} \log(x^2+1) + \frac{1}{6} \log(|x+1|)$$

input `integrate(1/(x^5+x^3+x^2+1),x, algorithm="giac")`output `1/2*arctan(x) - 1/3*log(x^2 - x + 1) + 1/4*log(x^2 + 1) + 1/6*log(abs(x + 1))`**Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.95

$$\int \frac{1}{1+x^2+x^3+x^5} dx = \frac{\ln(x+1)}{6} - \frac{\ln(x^2-x+1)}{3} + \ln(x-i) \left( \frac{1}{4} - \frac{1}{4}i \right) + \ln(x+1i) \left( \frac{1}{4} + \frac{1}{4}i \right)$$

input `int(1/(x^2 + x^3 + x^5 + 1),x)`

output  $\log(x + 1)/6 + \log(x - 1i)*(1/4 - 1i/4) + \log(x + 1i)*(1/4 + 1i/4) - \log(x^2 - x + 1)/3$

### Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.79

$$\int \frac{1}{1 + x^2 + x^3 + x^5} dx = \frac{\operatorname{atan}(x)}{2} - \frac{\log(x^2 - x + 1)}{3} + \frac{\log(x^2 + 1)}{4} + \frac{\log(x + 1)}{6}$$

input `int(1/(x^5+x^3+x^2+1),x)`

output  $(6*\operatorname{atan}(x) - 4*\log(x^2 - x + 1) + 3*\log(x^2 + 1) + 2*\log(x + 1))/12$

### 3.26 $\int \frac{1}{2+3(1+x)^5} dx$

Optimal result	189
Mathematica [A] (verified)	190
Rubi [A] (verified)	191
Maple [C] (verified)	195
Fricas [A] (verification not implemented)	196
Sympy [A] (verification not implemented)	197
Maxima [F]	197
Giac [F]	197
Mupad [B] (verification not implemented)	198
Reduce [F]	199

#### Optimal result

Integrand size = 11, antiderivative size = 305

$$\int \frac{1}{2+3(1+x)^5} dx = -\frac{\sqrt{5-\sqrt{5}} \arctan\left(\sqrt{\frac{1}{5}}(5+2\sqrt{5}) - \frac{2 \cdot 2^{3/10} \sqrt[5]{3}(1+x)}{\sqrt{5-\sqrt{5}}}\right)}{10 \cdot 2^{3/10} \sqrt[5]{3}} + \frac{\sqrt{5+\sqrt{5}} \arctan\left(\sqrt{\frac{1}{5}}(5-2\sqrt{5}) + \frac{2 \cdot 2^{3/10} \sqrt[5]{3}(1+x)}{\sqrt{5+\sqrt{5}}}\right)}{10 \cdot 2^{3/10} \sqrt[5]{3}} + \frac{\log\left(\sqrt[5]{2} + \sqrt[5]{3}(1+x)\right)}{5 \cdot 2^{4/5} \sqrt[5]{3}} - \frac{(1-\sqrt{5}) \log\left(2\sqrt[5]{2} - \sqrt[5]{3}(1-\sqrt{5})(1+x) + 2^{4/5} 3^{2/5}(1+x)^2\right)}{20 \cdot 2^{4/5} \sqrt[5]{3}} - \frac{(1+\sqrt{5}) \log\left(2\sqrt[5]{2} - \sqrt[5]{3}(1+\sqrt{5})(1+x) + 2^{4/5} 3^{2/5}(1+x)^2\right)}{20 \cdot 2^{4/5} \sqrt[5]{3}}$$

output

```
1/60*(5-5^(1/2))^(1/2)*arctan(-1/5*(25+10*5^(1/2))^(1/2)+2*2^(3/10)*3^(1/5)
)*(1+x)/(5-5^(1/2))^(1/2)*2^(7/10)*3^(4/5)+1/60*(5+5^(1/2))^(1/2)*arctan(
1/5*(25-10*5^(1/2))^(1/2)+2*2^(3/10)*3^(1/5)*(1+x)/(5+5^(1/2))^(1/2))*2^(7
/10)*3^(4/5)+1/30*ln(2^(1/5)+3^(1/5)*(1+x))*2^(1/5)*3^(4/5)-1/120*(-5^(1/2
)+1)*ln(2*2^(1/5)-3^(1/5)*(-5^(1/2)+1)*(1+x)+2^(4/5)*3^(2/5)*(1+x)^2*2^(1
/5)*3^(4/5)-1/120*(5^(1/2)+1)*ln(2*2^(1/5)-3^(1/5)*(5^(1/2)+1)*(1+x)+2^(4/
5)*3^(2/5)*(1+x)^2)*2^(1/5)*3^(4/5)
```

**Mathematica [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 248, normalized size of antiderivative = 0.81

$$\int \frac{1}{2+3(1+x)^5} dx$$

$$= \frac{2\sqrt{2(5+\sqrt{5})} \arctan\left(\frac{-1+2^{4/5}\sqrt[5]{3}+\sqrt{5}+2^{4/5}\sqrt[5]{3}x}{\sqrt{2(5+\sqrt{5})}}\right) + 2\sqrt{10-2\sqrt{5}} \arctan\left(\sqrt{\frac{1}{10}(5+\sqrt{5})}\left(\frac{1}{2}(-1-\sqrt{5})\right)\right)}{20 \cdot 2^{4/5} \cdot 3^{1/5}}$$

input

```
Integrate[(2 + 3*(1 + x)^5)^(-1), x]
```

output

```
(2*Sqrt[2*(5 + Sqrt[5])]*ArcTan[(-1 + 2*2^(4/5)*3^(1/5) + Sqrt[5] + 2*2^(4
/5)*3^(1/5)*x)/Sqrt[2*(5 + Sqrt[5])]] + 2*Sqrt[10 - 2*Sqrt[5]]*ArcTan[Sqrt
[(5 + Sqrt[5])/10]*((-1 - Sqrt[5])/2 + 2^(4/5)*3^(1/5)*(1 + x))] + 4*Log[2
+ 2^(4/5)*3^(1/5)*(1 + x)] + (-1 + Sqrt[5])*Log[2 + (3/2)^(1/5)*(-1 + Sqr
t[5])*(1 + x) + 2^(3/5)*3^(2/5)*(1 + x)^2] - (1 + Sqrt[5])*Log[2 - (3/2)^(
1/5)*(1 + Sqrt[5])*(1 + x) + 2^(3/5)*3^(2/5)*(1 + x)^2])/(20*2^(4/5)*3^(1/
5))
```

**Rubi [A] (verified)**

Time = 0.98 (sec) , antiderivative size = 307, normalized size of antiderivative = 1.01, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.818$ , Rules used = {239, 751, 16, 27, 1142, 25, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{3(x+1)^5 + 2} dx \\
 & \quad \downarrow \text{239} \\
 & \int \frac{1}{3(x+1)^5 + 2} d(x+1) \\
 & \quad \downarrow \text{751} \\
 & \frac{\int \frac{1}{\sqrt[5]{3}(x+1) + \sqrt[5]{2}} d(x+1)}{5 \cdot 2^{4/5}} + \frac{1}{5} \sqrt[5]{2} \int \frac{4\sqrt[5]{2} - \sqrt[5]{3}(1 - \sqrt{5})(x+1)}{2 \left( 2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1 - \sqrt{5})(x+1) + 2 \cdot 2^{2/5} \right)} d(x+1) \\
 & \quad + \frac{1}{5} \sqrt[5]{2} \int \frac{4\sqrt[5]{2} - \sqrt[5]{3}(1 + \sqrt{5})(x+1)}{2 \left( 2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1 + \sqrt{5})(x+1) + 2 \cdot 2^{2/5} \right)} d(x+1) \\
 & \quad \downarrow \text{16} \\
 & \frac{1}{5} \sqrt[5]{2} \int \frac{4\sqrt[5]{2} - \sqrt[5]{3}(1 - \sqrt{5})(x+1)}{2 \left( 2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1 - \sqrt{5})(x+1) + 2 \cdot 2^{2/5} \right)} d(x+1) + \\
 & \frac{1}{5} \sqrt[5]{2} \int \frac{4\sqrt[5]{2} - \sqrt[5]{3}(1 + \sqrt{5})(x+1)}{2 \left( 2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1 + \sqrt{5})(x+1) + 2 \cdot 2^{2/5} \right)} d(x+1) + \frac{\log \left( \sqrt[5]{3}(x+1) + \sqrt[5]{2} \right)}{5 \cdot 2^{4/5} \sqrt[5]{3}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{4\sqrt[5]{2} - \sqrt[5]{3}(1 - \sqrt{5})(x+1)}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1 - \sqrt{5})(x+1) + 2 \cdot 2^{2/5}} d(x+1)}{5 \cdot 2^{4/5}} + \\
 & \frac{\int \frac{4\sqrt[5]{2} - \sqrt[5]{3}(1 + \sqrt{5})(x+1)}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1 + \sqrt{5})(x+1) + 2 \cdot 2^{2/5}} d(x+1)}{5 \cdot 2^{4/5}} + \frac{\log \left( \sqrt[5]{3}(x+1) + \sqrt[5]{2} \right)}{5 \cdot 2^{4/5} \sqrt[5]{3}} \\
 & \quad \downarrow \text{1142}
 \end{aligned}$$



$$\frac{(5+\sqrt{5}) \int \frac{1}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1-\sqrt{5})(x+1)+2} \cdot \frac{1}{2^{2/5}} d(x+1)}{2^{4/5}} - \frac{(1-\sqrt{5}) \int \frac{\sqrt[5]{6}(1-\sqrt{5})^{-4} \cdot 3^{2/5}(x+1)}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1-\sqrt{5})(x+1)+2} \cdot \frac{1}{2^{2/5}} d(x+1)}{4 \sqrt[5]{3}}$$


---


$$\frac{5 \cdot 2^{4/5}}{4 \sqrt[5]{3}} +$$

$$\frac{(5-\sqrt{5}) \int \frac{1}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1+\sqrt{5})(x+1)+2} \cdot \frac{1}{2^{2/5}} d(x+1)}{2^{4/5}} - \frac{(1+\sqrt{5}) \int \frac{\sqrt[5]{6}(1+\sqrt{5})^{-4} \cdot 3^{2/5}(x+1)}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1+\sqrt{5})(x+1)+2} \cdot \frac{1}{2^{2/5}} d(x+1)}{4 \sqrt[5]{3}}$$


---


$$\frac{5 \cdot 2^{4/5}}{4 \sqrt[5]{3}} +$$

$$\frac{\log\left(\sqrt[5]{3}(x+1) + \sqrt[5]{2}\right)}{5 \cdot 2^{4/5} \sqrt[5]{3}}$$

↓ 25

$$\frac{(5+\sqrt{5}) \int \frac{1}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1-\sqrt{5})(x+1)+2} \cdot \frac{1}{2^{2/5}} d(x+1)}{2^{4/5}} + \frac{(1-\sqrt{5}) \int \frac{\sqrt[5]{6}(1-\sqrt{5})^{-4} \cdot 3^{2/5}(x+1)}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1-\sqrt{5})(x+1)+2} \cdot \frac{1}{2^{2/5}} d(x+1)}{4 \sqrt[5]{3}}$$


---


$$\frac{5 \cdot 2^{4/5}}{4 \sqrt[5]{3}} +$$

$$\frac{(5-\sqrt{5}) \int \frac{1}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1+\sqrt{5})(x+1)+2} \cdot \frac{1}{2^{2/5}} d(x+1)}{2^{4/5}} + \frac{(1+\sqrt{5}) \int \frac{\sqrt[5]{6}(1+\sqrt{5})^{-4} \cdot 3^{2/5}(x+1)}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1+\sqrt{5})(x+1)+2} \cdot \frac{1}{2^{2/5}} d(x+1)}{4 \sqrt[5]{3}}$$


---


$$\frac{5 \cdot 2^{4/5}}{4 \sqrt[5]{3}} +$$

$$\frac{\log\left(\sqrt[5]{3}(x+1) + \sqrt[5]{2}\right)}{5 \cdot 2^{4/5} \sqrt[5]{3}}$$

↓ 1083

$$\frac{(1-\sqrt{5}) \int \frac{\sqrt[5]{6}(1-\sqrt{5})^{-4} \cdot 3^{2/5}(x+1)}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1-\sqrt{5})(x+1)+2} \cdot \frac{1}{2^{2/5}} d(x+1)}{4 \sqrt[5]{3}} - \frac{\sqrt[5]{2}(5+\sqrt{5}) \int \frac{1}{-\left(4 \cdot 3^{2/5}(x+1) - \sqrt[5]{6}(1-\sqrt{5})\right)^2 - 2 \cdot 6^{2/5}(5+\sqrt{5})} d\left(4 \cdot 3^{2/5}(x+1) - \sqrt[5]{6}(1-\sqrt{5})\right)}{5 \cdot 2^{4/5}}$$


---


$$\frac{(1+\sqrt{5}) \int \frac{\sqrt[5]{6}(1+\sqrt{5})^{-4} \cdot 3^{2/5}(x+1)}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1+\sqrt{5})(x+1)+2} \cdot \frac{1}{2^{2/5}} d(x+1)}{4 \sqrt[5]{3}} - \frac{\sqrt[5]{2}(5-\sqrt{5}) \int \frac{1}{-\left(4 \cdot 3^{2/5}(x+1) - \sqrt[5]{6}(1+\sqrt{5})\right)^2 - 2 \cdot 6^{2/5}(5-\sqrt{5})} d\left(4 \cdot 3^{2/5}(x+1) - \sqrt[5]{6}(1+\sqrt{5})\right)}{5 \cdot 2^{4/5}}$$


---


$$\frac{\log\left(\sqrt[5]{3}(x+1) + \sqrt[5]{2}\right)}{5 \cdot 2^{4/5} \sqrt[5]{3}}$$

↓ 217

$$\begin{aligned}
 & \frac{(1-\sqrt{5}) \int \frac{\sqrt[5]{6}(1-\sqrt{5})-4 \cdot 3^{2/5}(x+1)}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1-\sqrt{5})(x+1)+2 \cdot 2^{2/5}} d(x+1)}{4 \sqrt[5]{3}} + \frac{\sqrt{\frac{1}{2}(5+\sqrt{5})} \arctan\left(\frac{4 \cdot 3^{2/5}(x+1) - \sqrt[5]{6}(1-\sqrt{5})}{2^{7/10} \sqrt[5]{3} \sqrt{5+\sqrt{5}}}\right)}{\sqrt[5]{3}} \\
 & \frac{(1+\sqrt{5}) \int \frac{\sqrt[5]{6}(1+\sqrt{5})-4 \cdot 3^{2/5}(x+1)}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1+\sqrt{5})(x+1)+2 \cdot 2^{2/5}} d(x+1)}{4 \sqrt[5]{3}} + \frac{\sqrt{\frac{1}{2}(5-\sqrt{5})} \arctan\left(\frac{4 \cdot 3^{2/5}(x+1) - \sqrt[5]{6}(1+\sqrt{5})}{2^{7/10} \sqrt[5]{3} \sqrt{5-\sqrt{5}}}\right)}{\sqrt[5]{3}} \\
 & \frac{5 \cdot 2^{4/5} \log\left(\sqrt[5]{3}(x+1) + \sqrt[5]{2}\right)}{5 \cdot 2^{4/5} \sqrt[5]{3}} \\
 & \quad \downarrow \text{1103} \\
 & \frac{\sqrt{\frac{1}{2}(5+\sqrt{5})} \arctan\left(\frac{4 \cdot 3^{2/5}(x+1) - \sqrt[5]{6}(1-\sqrt{5})}{2^{7/10} \sqrt[5]{3} \sqrt{5+\sqrt{5}}}\right)}{\sqrt[5]{3}} - \frac{(1-\sqrt{5}) \log\left(2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1-\sqrt{5})(x+1)+2 \cdot 2^{2/5}\right)}{4 \sqrt[5]{3}} \\
 & \frac{\sqrt{\frac{1}{2}(5-\sqrt{5})} \arctan\left(\frac{4 \cdot 3^{2/5}(x+1) - \sqrt[5]{6}(1+\sqrt{5})}{2^{7/10} \sqrt[5]{3} \sqrt{5-\sqrt{5}}}\right)}{\sqrt[5]{3}} - \frac{(1+\sqrt{5}) \log\left(2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1+\sqrt{5})(x+1)+2 \cdot 2^{2/5}\right)}{4 \sqrt[5]{3}} \\
 & \frac{5 \cdot 2^{4/5} \log\left(\sqrt[5]{3}(x+1) + \sqrt[5]{2}\right)}{5 \cdot 2^{4/5} \sqrt[5]{3}}
 \end{aligned}$$

input `Int[(2 + 3*(1 + x)^5)^(-1),x]`

output `Log[2^(1/5) + 3^(1/5)*(1 + x)]/(5*2^(4/5)*3^(1/5)) + ((Sqrt[(5 + Sqrt[5])/2]*ArcTan[(-(6^(1/5)*(1 - Sqrt[5])) + 4*3^(2/5)*(1 + x))/(2^(7/10)*3^(1/5)*Sqrt[5 + Sqrt[5]])])/3^(1/5) - ((1 - Sqrt[5])*Log[2*2^(2/5) - 6^(1/5)*(1 - Sqrt[5])*(1 + x) + 2*3^(2/5)*(1 + x)^2])/(4*3^(1/5)))/(5*2^(4/5)) + ((Sqrt[(5 - Sqrt[5])/2]*ArcTan[(-(6^(1/5)*(1 + Sqrt[5])) + 4*3^(2/5)*(1 + x))/(2^(7/10)*3^(1/5)*Sqrt[5 - Sqrt[5]])])/3^(1/5) - ((1 + Sqrt[5])*Log[2*2^(2/5) - 6^(1/5)*(1 + Sqrt[5])*(1 + x) + 2*3^(2/5)*(1 + x)^2])/(4*3^(1/5)))/(5*2^(4/5))`

## Definitions of rubi rules used

- rule 16  $\text{Int}[(c\_)/((a\_)+(b\_)*(x\_)), x\_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 25  $\text{Int}[-(Fx\_), x\_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[Fx, x], x]$
- rule 27  $\text{Int}[(a\_)*(Fx\_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b\_)*(Gx\_)] /; \text{FreeQ}[b, x]$
- rule 217  $\text{Int}[(a\_)+(b\_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 239  $\text{Int}[(a\_)+(b\_)*(v_)^{(n_)})^{(p_)}, x\_Symbol] \rightarrow \text{Simp}[1/\text{Coefficient}[v, x, 1] \text{ Subst}[\text{Int}[(a + b*x^n)^p, x], x, v], x] /; \text{FreeQ}[\{a, b, n, p\}, x] \ \&\& \ \text{LinearQ}[v, x] \ \&\& \ \text{NeQ}[v, x]$
- rule 751  $\text{Int}[(a\_)+(b\_)*(x_)^{(n_)})^{-1}, x\_Symbol] \rightarrow \text{Module}[\{r = \text{Numerator}[\text{Rt}[a/b, n]], s = \text{Denominator}[\text{Rt}[a/b, n]], k, u\}, \text{Simp}[u = \text{Int}[(r - s*\text{Cos}[(2*k - 1)*(Pi/n)]*x)/(r^2 - 2*r*s*\text{Cos}[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x]; r/(a*n) \text{ Int}[1/(r + s*x), x] + 2*(r/(a*n)) \text{ Sum}[u, \{k, 1, (n - 1)/2\}], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{IGtQ}[(n - 3)/2, 0] \ \&\& \ \text{PosQ}[a/b]$
- rule 1083  $\text{Int}[(a\_)+(b\_)*(x_)+(c\_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[-2 \text{ Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103  $\text{Int}[(d\_)+(e\_)*(x_))/((a\_)+(b\_)*(x_)+(c\_)*(x_)^2), x\_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$

rule 1142

```
Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c)
Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]
```

**Maple [C] (verified)**

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.03 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.19

method	result	size
default	$\frac{\left( \sum_{R=\text{RootOf}(3Z^5+15Z^4+30Z^3+30Z^2+15Z+5)} \frac{\ln(x-R)}{R^4+4R^3+6R^2+4R+1} \right)}{15}$	59
risch	$\frac{\left( \sum_{R=\text{RootOf}(3Z^5+15Z^4+30Z^3+30Z^2+15Z+5)} \frac{\ln(x-R)}{R^4+4R^3+6R^2+4R+1} \right)}{15}$	59

input

```
int(1/(2+3*(x+1)^5),x,method=_RETURNVERBOSE)
```

output

```
1/15*sum(1/(_R^4+4*_R^3+6*_R^2+4*_R+1)*ln(x-_R),_R=RootOf(3*_Z^5+15*_Z^4+30*_Z^3+30*_Z^2+15*_Z+5))
```

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 255, normalized size of antiderivative = 0.84

$$\begin{aligned}
& \int \frac{1}{2+3(1+x)^5} dx \\
&= -\frac{1}{960} \left( 48^{\frac{4}{5}} - 60 \cdot 48^{\frac{3}{10}} \sqrt{\frac{1}{15}} \right) \log \left( 48x^2 - 48^{\frac{4}{5}}(x+1) + 60 \cdot 48^{\frac{3}{10}} \sqrt{\frac{1}{15}}(x+1) \right. \\
&\quad \left. + 96x + 4 \cdot 48^{\frac{3}{5}} + 48 \right) - \frac{1}{960} \left( 48^{\frac{4}{5}} + 60 \cdot 48^{\frac{3}{10}} \sqrt{\frac{1}{15}} \right) \log \left( 48x^2 - 48^{\frac{4}{5}}(x+1) - 60 \right. \\
&\quad \left. \cdot 48^{\frac{3}{10}} \sqrt{\frac{1}{15}}(x+1) + 96x + 4 \cdot 48^{\frac{3}{5}} + 48 \right) + \frac{1}{240} \cdot 48^{\frac{4}{5}} \log \left( 24x + 48^{\frac{4}{5}} + 24 \right) \\
&\quad - \frac{1}{8} \sqrt{\frac{2}{15} \cdot 48^{\frac{3}{5}} + \frac{8}{5} \cdot 48^{\frac{1}{10}} \sqrt{\frac{1}{15}}} \arctan \left( \frac{5}{64} \left( 48^{\frac{3}{10}} \sqrt{\frac{1}{15}} \left( 48^{\frac{3}{5}}(x+1) - 3 \cdot 48^{\frac{2}{5}} \right) - 4 \cdot 48^{\frac{2}{5}}(x+1) + 4 \cdot 48^{\frac{1}{5}} \right) \right. \\
&\quad \left. + \frac{1}{8} \sqrt{\frac{2}{15} \cdot 48^{\frac{3}{5}} - \frac{8}{5} \cdot 48^{\frac{1}{10}} \sqrt{\frac{1}{15}}} \arctan \left( \frac{5}{64} \left( 48^{\frac{3}{10}} \sqrt{\frac{1}{15}} \left( 48^{\frac{3}{5}}(x+1) - 3 \cdot 48^{\frac{2}{5}} \right) + 4 \cdot 48^{\frac{2}{5}}(x+1) - 4 \cdot 48^{\frac{1}{5}} \right) \right)
\end{aligned}$$

```
input integrate(1/(2+3*(1+x)^5),x, algorithm="fricas")
```

```
output -1/960*(48^(4/5) - 60*48^(3/10)*sqrt(1/15))*log(48*x^2 - 48^(4/5)*(x + 1)
+ 60*48^(3/10)*sqrt(1/15)*(x + 1) + 96*x + 4*48^(3/5) + 48) - 1/960*(48^(4
/5) + 60*48^(3/10)*sqrt(1/15))*log(48*x^2 - 48^(4/5)*(x + 1) - 60*48^(3/10
)*sqrt(1/15)*(x + 1) + 96*x + 4*48^(3/5) + 48) + 1/240*48^(4/5)*log(24*x +
48^(4/5) + 24) - 1/8*sqrt(2/15*48^(3/5) + 8/5*48^(1/10)*sqrt(1/15))*arcta
n(5/64*(48^(3/10)*sqrt(1/15)*(48^(3/5)*(x + 1) - 3*48^(2/5)) - 4*48^(2/5)*
(x + 1) + 4*48^(1/5))*sqrt(2/15*48^(3/5) + 8/5*48^(1/10)*sqrt(1/15))) + 1/
8*sqrt(2/15*48^(3/5) - 8/5*48^(1/10)*sqrt(1/15))*arctan(5/64*(48^(3/10)*sq
rt(1/15)*(48^(3/5)*(x + 1) - 3*48^(2/5)) + 4*48^(2/5)*(x + 1) - 4*48^(1/5)
)*sqrt(2/15*48^(3/5) - 8/5*48^(1/10)*sqrt(1/15)))
```

**Sympy [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.05

$$\int \frac{1}{2 + 3(1 + x)^5} dx = \text{RootSum}(150000t^5 - 1, (t \mapsto t \log(10t + x + 1)))$$

input `integrate(1/(2+3*(1+x)**5),x)`

output `RootSum(150000*_t**5 - 1, Lambda(_t, _t*log(10*_t + x + 1)))`

**Maxima [F]**

$$\int \frac{1}{2 + 3(1 + x)^5} dx = \int \frac{1}{3(x + 1)^5 + 2} dx$$

input `integrate(1/(2+3*(1+x)^5),x, algorithm="maxima")`

output `integrate(1/(3*(x + 1)^5 + 2), x)`

**Giac [F]**

$$\int \frac{1}{2 + 3(1 + x)^5} dx = \int \frac{1}{3(x + 1)^5 + 2} dx$$

input `integrate(1/(2+3*(1+x)^5),x, algorithm="giac")`

output `integrate(1/(3*(x + 1)^5 + 2), x)`

**Mupad [B] (verification not implemented)**

Time = 23.01 (sec) , antiderivative size = 251, normalized size of antiderivative = 0.82

$$\int \frac{1}{2 + 3(1+x)^5} dx = \frac{162^{1/5} \ln\left(x + \frac{162^{1/5}}{3} + 1\right)}{30} - \frac{3^{4/5} \ln\left(x - \frac{3^{4/5} \left(2^{1/5} (\sqrt{5}+1) + 2^{7/10} \sqrt{\sqrt{5}-5}\right)}{12} + 1\right) \left(2^{1/5} (\sqrt{5}+1) + 2^{7/10} \sqrt{\sqrt{5}-5}\right)}{120} - \frac{3^{4/5} \ln\left(x - \frac{3^{4/5} \left(2^{1/5} (\sqrt{5}+1) - 2^{7/10} \sqrt{\sqrt{5}-5}\right)}{12} + 1\right) \left(2^{1/5} (\sqrt{5}+1) - 2^{7/10} \sqrt{\sqrt{5}-5}\right)}{120} + \frac{3^{4/5} \ln\left(x + \frac{3^{4/5} \left(2^{7/10} \sqrt{-\sqrt{5}-5} + 2^{1/5} (\sqrt{5}-1)\right)}{12} + 1\right) \left(2^{7/10} \sqrt{-\sqrt{5}-5} + 2^{1/5} (\sqrt{5}-1)\right)}{120} - \frac{3^{4/5} \ln\left(x - \frac{3^{4/5} \left(2^{7/10} \sqrt{-\sqrt{5}-5} - 2^{1/5} (\sqrt{5}-1)\right)}{12} + 1\right) \left(2^{7/10} \sqrt{-\sqrt{5}-5} - 2^{1/5} (\sqrt{5}-1)\right)}{120}$$

input `int(1/(3*(x + 1)^5 + 2),x)`

output

```
(162^(1/5)*log(x + 162^(1/5)/3 + 1))/30 - (3^(4/5)*log(x - (3^(4/5)*(2^(1/5)*(5^(1/2) + 1) + 2^(7/10)*(5^(1/2) - 5)^(1/2)))/12 + 1)*(2^(1/5)*(5^(1/2) + 1) + 2^(7/10)*(5^(1/2) - 5)^(1/2)))/120 - (3^(4/5)*log(x - (3^(4/5)*(2^(1/5)*(5^(1/2) + 1) - 2^(7/10)*(5^(1/2) - 5)^(1/2)))/12 + 1)*(2^(1/5)*(5^(1/2) + 1) - 2^(7/10)*(5^(1/2) - 5)^(1/2)))/120 + (3^(4/5)*log(x + (3^(4/5)*(2^(7/10)*(-5^(1/2) - 5)^(1/2) + 2^(1/5)*(5^(1/2) - 1)))/12 + 1)*(2^(7/10)*(-5^(1/2) - 5)^(1/2) + 2^(1/5)*(5^(1/2) - 1)))/120 - (3^(4/5)*log(x - (3^(4/5)*(2^(7/10)*(-5^(1/2) - 5)^(1/2) - 2^(1/5)*(5^(1/2) - 1)))/12 + 1)*(2^(7/10)*(-5^(1/2) - 5)^(1/2) - 2^(1/5)*(5^(1/2) - 1)))/120
```

Reduce [F]

$$\int \frac{1}{2 + 3(1 + x)^5} dx = \int \frac{1}{3x^5 + 15x^4 + 30x^3 + 30x^2 + 15x + 5} dx$$

input `int(1/(2+3*(1+x)^5),x)`

output `int(1/(3*x**5 + 15*x**4 + 30*x**3 + 30*x**2 + 15*x + 5),x)`



$$3.27 \quad \int \frac{1}{5+15x+30x^2+30x^3+15x^4+3x^5} dx$$

Optimal result	200
Mathematica [C] (verified)	201
Rubi [A] (verified)	201
Maple [C] (verified)	206
Fricas [A] (verification not implemented)	206
Sympy [A] (verification not implemented)	207
Maxima [F]	207
Giac [F]	208
Mupad [B] (verification not implemented)	208
Reduce [F]	209

### Optimal result

Integrand size = 27, antiderivative size = 305

$$\int \frac{1}{5 + 15x + 30x^2 + 30x^3 + 15x^4 + 3x^5} dx$$

$$= -\frac{\sqrt{5 - \sqrt{5}} \arctan\left(\sqrt{\frac{1}{5}}(5 + 2\sqrt{5}) - \frac{2 \cdot 2^{3/10} \sqrt[5]{3}(1+x)}{\sqrt{5 - \sqrt{5}}}\right)}{10 \cdot 2^{3/10} \sqrt[5]{3}} + \frac{\sqrt{5 + \sqrt{5}} \arctan\left(\sqrt{\frac{1}{5}}(5 - 2\sqrt{5}) + \frac{2 \cdot 2^{3/10} \sqrt[5]{3}(1+x)}{\sqrt{5 + \sqrt{5}}}\right)}{10 \cdot 2^{3/10} \sqrt[5]{3}} + \frac{\log\left(\sqrt[5]{2} + \sqrt[5]{3}(1+x)\right)}{5 \cdot 2^{4/5} \sqrt[5]{3}}$$

$$- \frac{(1 - \sqrt{5}) \log\left(2\sqrt[5]{2} - \sqrt[5]{3}(1 - \sqrt{5})(1+x) + 2^{4/5} 3^{2/5}(1+x)^2\right)}{20 \cdot 2^{4/5} \sqrt[5]{3}}$$

$$- \frac{(1 + \sqrt{5}) \log\left(2\sqrt[5]{2} - \sqrt[5]{3}(1 + \sqrt{5})(1+x) + 2^{4/5} 3^{2/5}(1+x)^2\right)}{20 \cdot 2^{4/5} \sqrt[5]{3}}$$

output

```
1/60*(5-5^(1/2))^(1/2)*arctan(-1/5*(25+10*5^(1/2))^(1/2)+2*2^(3/10)*3^(1/5)
)*(1+x)/(5-5^(1/2))^(1/2)*2^(7/10)*3^(4/5)+1/60*(5+5^(1/2))^(1/2)*arctan(
1/5*(25-10*5^(1/2))^(1/2)+2*2^(3/10)*3^(1/5)*(1+x)/(5+5^(1/2))^(1/2))*2^(7
/10)*3^(4/5)+1/30*ln(2^(1/5)+3^(1/5)*(1+x))*2^(1/5)*3^(4/5)-1/120*(-5^(1/2
)+1)*ln(2*2^(1/5)-3^(1/5)*(-5^(1/2)+1)*(1+x)+2^(4/5)*3^(2/5)*(1+x)^2)*2^(1
/5)*3^(4/5)-1/120*(5^(1/2)+1)*ln(2*2^(1/5)-3^(1/5)*(5^(1/2)+1)*(1+x)+2^(4/
5)*3^(2/5)*(1+x)^2)*2^(1/5)*3^(4/5)
```

**Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.02 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.23

$$\int \frac{1}{5 + 15x + 30x^2 + 30x^3 + 15x^4 + 3x^5} dx$$

$$= \frac{1}{15} \text{RootSum} \left[ 5 + 15\#1 + 30\#1^2 + 30\#1^3 + 15\#1^4 + 3\#1^5 \&, \frac{\log(x - \#1)}{1 + 4\#1 + 6\#1^2 + 4\#1^3 + \#1^4} \& \right]$$

input

```
Integrate[(5 + 15*x + 30*x^2 + 30*x^3 + 15*x^4 + 3*x^5)^(-1), x]
```

output

```
RootSum[5 + 15*#1 + 30*#1^2 + 30*#1^3 + 15*#1^4 + 3*#1^5 & , Log[x - #1]/(1 + 4*#1 + 6*#1^2 + 4*#1^3 + #1^4) & ]/15
```

**Rubi [A] (verified)**

Time = 0.87 (sec) , antiderivative size = 307, normalized size of antiderivative = 1.01, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2458, 751, 16, 27, 1142, 25, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{3x^5 + 15x^4 + 30x^3 + 30x^2 + 15x + 5} dx$$

$$\downarrow \text{2458}$$

$$\int \frac{1}{3(x+1)^5 + 2} d(x+1)$$

$$\downarrow \text{751}$$

$$\frac{\int \frac{1}{\sqrt[5]{3(x+1)+\sqrt[5]{2}}} d(x+1)}{5 \cdot 2^{4/5}} + \frac{1}{5} \sqrt[5]{2} \int \frac{4\sqrt[5]{2} - \sqrt[5]{3}(1-\sqrt{5})(x+1)}{2 \left( 2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1-\sqrt{5})(x+1) + 2 \cdot 2^{2/5} \right)} d(x+1) +$$

$$1) + \frac{1}{5} \sqrt[5]{2} \int \frac{4\sqrt[5]{2} - \sqrt[5]{3}(1+\sqrt{5})(x+1)}{2 \left( 2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1+\sqrt{5})(x+1) + 2 \cdot 2^{2/5} \right)} d(x+1)$$

↓ 16

$$\frac{1}{5} \sqrt[5]{2} \int \frac{4\sqrt[5]{2} - \sqrt[5]{3}(1-\sqrt{5})(x+1)}{2 \left( 2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1-\sqrt{5})(x+1) + 2 \cdot 2^{2/5} \right)} d(x+1) +$$

$$\frac{1}{5} \sqrt[5]{2} \int \frac{4\sqrt[5]{2} - \sqrt[5]{3}(1+\sqrt{5})(x+1)}{2 \left( 2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1+\sqrt{5})(x+1) + 2 \cdot 2^{2/5} \right)} d(x+1) + \frac{\log \left( \sqrt[5]{3}(x+1) + \sqrt[5]{2} \right)}{5 \cdot 2^{4/5} \sqrt[5]{3}}$$

↓ 27

$$\frac{\int \frac{4\sqrt[5]{2} - \sqrt[5]{3}(1-\sqrt{5})(x+1)}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1-\sqrt{5})(x+1) + 2 \cdot 2^{2/5}} d(x+1)}{5 \cdot 2^{4/5}} +$$

$$\frac{\int \frac{4\sqrt[5]{2} - \sqrt[5]{3}(1+\sqrt{5})(x+1)}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1+\sqrt{5})(x+1) + 2 \cdot 2^{2/5}} d(x+1)}{5 \cdot 2^{4/5}} + \frac{\log \left( \sqrt[5]{3}(x+1) + \sqrt[5]{2} \right)}{5 \cdot 2^{4/5} \sqrt[5]{3}}$$

↓ 1142

$$\frac{(5+\sqrt{5}) \int \frac{1}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1-\sqrt{5})(x+1) + 2 \cdot 2^{2/5}} d(x+1)}{2^{4/5}} - \frac{(1-\sqrt{5}) \int \frac{\sqrt[5]{6}(1-\sqrt{5}) - 4 \cdot 3^{2/5}(x+1)}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1-\sqrt{5})(x+1) + 2 \cdot 2^{2/5}} d(x+1)}{4 \sqrt[5]{3}}$$

$$\frac{5 \cdot 2^{4/5}}{5 \cdot 2^{4/5}} +$$

$$\frac{(5-\sqrt{5}) \int \frac{1}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1+\sqrt{5})(x+1) + 2 \cdot 2^{2/5}} d(x+1)}{2^{4/5}} - \frac{(1+\sqrt{5}) \int \frac{\sqrt[5]{6}(1+\sqrt{5}) - 4 \cdot 3^{2/5}(x+1)}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1+\sqrt{5})(x+1) + 2 \cdot 2^{2/5}} d(x+1)}{4 \sqrt[5]{3}}$$

$$\frac{5 \cdot 2^{4/5}}{5 \cdot 2^{4/5}} +$$

$$\frac{\log \left( \sqrt[5]{3}(x+1) + \sqrt[5]{2} \right)}{5 \cdot 2^{4/5} \sqrt[5]{3}}$$

↓ 25

$$\frac{(5+\sqrt{5}) \int \frac{1}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1-\sqrt{5})(x+1)+2} \cdot \frac{1}{2^{2/5}} d(x+1)}{2^{4/5}} + \frac{(1-\sqrt{5}) \int \frac{\sqrt[5]{6}(1-\sqrt{5})^{-4} \cdot 3^{2/5}(x+1)}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1-\sqrt{5})(x+1)+2} \cdot \frac{1}{2^{2/5}} d(x+1)}{4 \sqrt[5]{3}}$$


---


$$\frac{5 \cdot 2^{4/5}}{4 \sqrt[5]{3}} +$$

$$\frac{(5-\sqrt{5}) \int \frac{1}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1+\sqrt{5})(x+1)+2} \cdot \frac{1}{2^{2/5}} d(x+1)}{2^{4/5}} + \frac{(1+\sqrt{5}) \int \frac{\sqrt[5]{6}(1+\sqrt{5})^{-4} \cdot 3^{2/5}(x+1)}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1+\sqrt{5})(x+1)+2} \cdot \frac{1}{2^{2/5}} d(x+1)}{4 \sqrt[5]{3}}$$


---


$$\frac{5 \cdot 2^{4/5}}{4 \sqrt[5]{3}} +$$

$$\frac{\log\left(\sqrt[5]{3}(x+1) + \sqrt[5]{2}\right)}{5 \cdot 2^{4/5} \sqrt[5]{3}}$$

↓ 1083

$$\frac{(1-\sqrt{5}) \int \frac{\sqrt[5]{6}(1-\sqrt{5})^{-4} \cdot 3^{2/5}(x+1)}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1-\sqrt{5})(x+1)+2} \cdot \frac{1}{2^{2/5}} d(x+1)}{4 \sqrt[5]{3}} - \frac{\sqrt[5]{2}(5+\sqrt{5}) \int \frac{1}{-\left(4 \cdot 3^{2/5}(x+1) - \sqrt[5]{6}(1-\sqrt{5})\right)^2 - 2 \cdot 6^{2/5}(5+\sqrt{5})} d\left(4 \cdot 3^{2/5}(x+1) - \sqrt[5]{6}(1-\sqrt{5})\right)}{5 \cdot 2^{4/5}}$$


---


$$\frac{(1+\sqrt{5}) \int \frac{\sqrt[5]{6}(1+\sqrt{5})^{-4} \cdot 3^{2/5}(x+1)}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1+\sqrt{5})(x+1)+2} \cdot \frac{1}{2^{2/5}} d(x+1)}{4 \sqrt[5]{3}} - \frac{\sqrt[5]{2}(5-\sqrt{5}) \int \frac{1}{-\left(4 \cdot 3^{2/5}(x+1) - \sqrt[5]{6}(1+\sqrt{5})\right)^2 - 2 \cdot 6^{2/5}(5-\sqrt{5})} d\left(4 \cdot 3^{2/5}(x+1) - \sqrt[5]{6}(1+\sqrt{5})\right)}{5 \cdot 2^{4/5}}$$


---


$$\frac{\log\left(\sqrt[5]{3}(x+1) + \sqrt[5]{2}\right)}{5 \cdot 2^{4/5} \sqrt[5]{3}}$$

↓ 217

$$\frac{(1-\sqrt{5}) \int \frac{\sqrt[5]{6}(1-\sqrt{5})^{-4} \cdot 3^{2/5}(x+1)}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1-\sqrt{5})(x+1)+2} \cdot \frac{1}{2^{2/5}} d(x+1)}{4 \sqrt[5]{3}} + \frac{\sqrt{\frac{1}{2}(5+\sqrt{5})} \arctan\left(\frac{4 \cdot 3^{2/5}(x+1) - \sqrt[5]{6}(1-\sqrt{5})}{2^{7/10} \sqrt[5]{3} \sqrt{5+\sqrt{5}}}\right)}{5 \sqrt[5]{3}}$$


---


$$\frac{5 \cdot 2^{4/5}}{4 \sqrt[5]{3}} +$$

$$\frac{(1+\sqrt{5}) \int \frac{\sqrt[5]{6}(1+\sqrt{5})^{-4} \cdot 3^{2/5}(x+1)}{2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1+\sqrt{5})(x+1)+2} \cdot \frac{1}{2^{2/5}} d(x+1)}{4 \sqrt[5]{3}} + \frac{\sqrt{\frac{1}{2}(5-\sqrt{5})} \arctan\left(\frac{4 \cdot 3^{2/5}(x+1) - \sqrt[5]{6}(1+\sqrt{5})}{2^{7/10} \sqrt[5]{3} \sqrt{5-\sqrt{5}}}\right)}{5 \sqrt[5]{3}}$$


---


$$\frac{5 \cdot 2^{4/5}}{4 \sqrt[5]{3}} +$$

$$\frac{\log\left(\sqrt[5]{3}(x+1) + \sqrt[5]{2}\right)}{5 \cdot 2^{4/5} \sqrt[5]{3}}$$

↓ 1103

$$\frac{\sqrt{\frac{1}{2}(5+\sqrt{5})} \arctan\left(\frac{4 \cdot 3^{2/5}(x+1) - \sqrt[5]{6}(1-\sqrt{5})}{2^{7/10} \sqrt[5]{3} \sqrt{5+\sqrt{5}}}\right)}{\sqrt[5]{3}} - \frac{(1-\sqrt{5}) \log\left(2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1-\sqrt{5})(x+1) + 2 \cdot 2^{2/5}\right)}{4 \sqrt[5]{3}}$$

$$+ \frac{\sqrt{\frac{1}{2}(5-\sqrt{5})} \arctan\left(\frac{4 \cdot 3^{2/5}(x+1) - \sqrt[5]{6}(1+\sqrt{5})}{2^{7/10} \sqrt[5]{3} \sqrt{5-\sqrt{5}}}\right)}{\sqrt[5]{3}} - \frac{(1+\sqrt{5}) \log\left(2 \cdot 3^{2/5}(x+1)^2 - \sqrt[5]{6}(1+\sqrt{5})(x+1) + 2 \cdot 2^{2/5}\right)}{4 \sqrt[5]{3}}$$

$$+ \frac{5 \cdot 2^{4/5} \log\left(\sqrt[5]{3}(x+1) + \sqrt[5]{2}\right)}{5 \cdot 2^{4/5} \sqrt[5]{3}}$$

input

```
Int[(5 + 15*x + 30*x^2 + 30*x^3 + 15*x^4 + 3*x^5)^(-1),x]
```

output

```
Log[2^(1/5) + 3^(1/5)*(1 + x)]/(5*2^(4/5)*3^(1/5)) + ((Sqrt[(5 + Sqrt[5])/2]*ArcTan[(-6^(1/5)*(1 - Sqrt[5])) + 4*3^(2/5)*(1 + x)]/(2^(7/10)*3^(1/5)*Sqrt[5 + Sqrt[5]])]/3^(1/5) - ((1 - Sqrt[5])*Log[2*2^(2/5) - 6^(1/5)*(1 - Sqrt[5])*(1 + x) + 2*3^(2/5)*(1 + x)^2]/(4*3^(1/5)))/(5*2^(4/5)) + ((Sqrt[(5 - Sqrt[5])/2]*ArcTan[(-6^(1/5)*(1 + Sqrt[5])) + 4*3^(2/5)*(1 + x)]/(2^(7/10)*3^(1/5)*Sqrt[5 - Sqrt[5]])]/3^(1/5) - ((1 + Sqrt[5])*Log[2*2^(2/5) - 6^(1/5)*(1 + Sqrt[5])*(1 + x) + 2*3^(2/5)*(1 + x)^2]/(4*3^(1/5)))/(5*2^(4/5))
```

### Defintions of rubi rules used

rule 16

```
Int[(c_)/((a_) + (b_)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]
```

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 751 `Int[((a_) + (b_.)*(x_)^(n_))^(k_), x_Symbol] := Module[{r = Numerator[Rt[a/b, n]], s = Denominator[Rt[a/b, n]], k, u}, Simp[u = Int[(r - s*Cos[(2*k - 1)*(Pi/n)]*x)/(r^2 - 2*r*s*Cos[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x]; r/(a*n) Int[1/(r + s*x), x] + 2*(r/(a*n)) Sum[u, {k, 1, (n - 1)/2}], x] /; FreeQ[{a, b}, x] && IGtQ[(n - 3)/2, 0] && PosQ[a/b]`

rule 1083 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1142 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 2458 `Int[(Pn_)^(p_), x_Symbol] := With[{S = Coeff[Pn, x, Expon[Pn, x] - 1]/(Expon[Pn, x]*Coeff[Pn, x, Expon[Pn, x]])}, Subst[Int[ExpandToSum[Pn /. x -> x - S, x]^p, x], x, x + S] /; BinomialQ[Pn /. x -> x - S, x] || (IntegerQ[Expon[Pn, x]/2] && TrinomialQ[Pn /. x -> x - S, x])] /; FreeQ[p, x] && PolyQ[Pn, x] && GtQ[Expon[Pn, x], 2] && NeQ[Coeff[Pn, x, Expon[Pn, x] - 1], 0]`

**Maple [C] (verified)**

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.03 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.19

method	result	size
default	$\frac{\sum_{R=\text{RootOf}(3Z^5+15Z^4+30Z^3+30Z^2+15Z+5)} \frac{\ln(x-R)}{R^4+4R^3+6R^2+4R+1}}{15}$	59
risch	$\frac{\sum_{R=\text{RootOf}(3Z^5+15Z^4+30Z^3+30Z^2+15Z+5)} \frac{\ln(x-R)}{R^4+4R^3+6R^2+4R+1}}{15}$	59

input `int(1/(3*x^5+15*x^4+30*x^3+30*x^2+15*x+5),x,method=_RETURNVERBOSE)`

output `1/15*sum(1/(_R^4+4*_R^3+6*_R^2+4*_R+1)*ln(x-_R),_R=RootOf(3*_Z^5+15*_Z^4+30*_Z^3+30*_Z^2+15*_Z+5))`

**Fricas [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 255, normalized size of antiderivative = 0.84

$$\int \frac{1}{5 + 15x + 30x^2 + 30x^3 + 15x^4 + 3x^5} dx$$

$$= -\frac{1}{960} \left( 48^{\frac{4}{5}} - 60 \cdot 48^{\frac{3}{10}} \sqrt{\frac{1}{15}} \right) \log \left( 48x^2 - 48^{\frac{4}{5}}(x+1) + 60 \cdot 48^{\frac{3}{10}} \sqrt{\frac{1}{15}}(x+1) \right. \\ \left. + 96x + 4 \cdot 48^{\frac{3}{5}} + 48 \right) - \frac{1}{960} \left( 48^{\frac{4}{5}} + 60 \cdot 48^{\frac{3}{10}} \sqrt{\frac{1}{15}} \right) \log \left( 48x^2 - 48^{\frac{4}{5}}(x+1) - 60 \right. \\ \left. \cdot 48^{\frac{3}{10}} \sqrt{\frac{1}{15}}(x+1) + 96x + 4 \cdot 48^{\frac{3}{5}} + 48 \right) + \frac{1}{240} \cdot 48^{\frac{4}{5}} \log \left( 24x + 48^{\frac{4}{5}} + 24 \right) \\ - \frac{1}{8} \sqrt{\frac{2}{15} \cdot 48^{\frac{3}{5}} + \frac{8}{5} \cdot 48^{\frac{1}{10}} \sqrt{\frac{1}{15}}} \arctan \left( \frac{5}{64} \left( 48^{\frac{3}{10}} \sqrt{\frac{1}{15}} \left( 48^{\frac{3}{5}}(x+1) - 3 \cdot 48^{\frac{2}{5}} \right) - 4 \cdot 48^{\frac{2}{5}}(x+1) + 4 \cdot 48 \right) \right. \\ \left. + \frac{1}{8} \sqrt{\frac{2}{15} \cdot 48^{\frac{3}{5}} - \frac{8}{5} \cdot 48^{\frac{1}{10}} \sqrt{\frac{1}{15}}} \arctan \left( \frac{5}{64} \left( 48^{\frac{3}{10}} \sqrt{\frac{1}{15}} \left( 48^{\frac{3}{5}}(x+1) - 3 \cdot 48^{\frac{2}{5}} \right) + 4 \cdot 48^{\frac{2}{5}}(x+1) - 4 \cdot 48 \right) \right)$$

input `integrate(1/(3*x^5+15*x^4+30*x^3+30*x^2+15*x+5),x, algorithm="fricas")`

output 
$$\begin{aligned} & -1/960*(48^{(4/5)} - 60*48^{(3/10)}*\sqrt{1/15})*\log(48*x^2 - 48^{(4/5)}*(x + 1) \\ & + 60*48^{(3/10)}*\sqrt{1/15}*(x + 1) + 96*x + 4*48^{(3/5)} + 48) - 1/960*(48^{(4/5)} \\ & + 60*48^{(3/10)}*\sqrt{1/15})*\log(48*x^2 - 48^{(4/5)}*(x + 1) - 60*48^{(3/10)} \\ & )*\sqrt{1/15}*(x + 1) + 96*x + 4*48^{(3/5)} + 48) + 1/240*48^{(4/5)}*\log(24*x + \\ & 48^{(4/5)} + 24) - 1/8*\sqrt{2/15*48^{(3/5)} + 8/5*48^{(1/10)}*\sqrt{1/15}}*\arctan \\ & n(5/64*(48^{(3/10)}*\sqrt{1/15}*(48^{(3/5)}*(x + 1) - 3*48^{(2/5)}) - 4*48^{(2/5)}* \\ & (x + 1) + 4*48^{(1/5)})*\sqrt{2/15*48^{(3/5)} + 8/5*48^{(1/10)}*\sqrt{1/15}})) + 1/ \\ & 8*\sqrt{2/15*48^{(3/5)} - 8/5*48^{(1/10)}*\sqrt{1/15}}*\arctan(5/64*(48^{(3/10)}*\sqrt{1/15} \\ & *(48^{(3/5)}*(x + 1) - 3*48^{(2/5)}) + 4*48^{(2/5)}*(x + 1) - 4*48^{(1/5)} \\ & )*\sqrt{2/15*48^{(3/5)} - 8/5*48^{(1/10)}*\sqrt{1/15}})) \end{aligned}$$

### Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.05

$$\begin{aligned} & \int \frac{1}{5 + 15x + 30x^2 + 30x^3 + 15x^4 + 3x^5} dx \\ & = \text{RootSum}(150000t^5 - 1, (t \mapsto t \log(10t + x + 1))) \end{aligned}$$

input `integrate(1/(3*x**5+15*x**4+30*x**3+30*x**2+15*x+5),x)`

output `RootSum(150000*_t**5 - 1, Lambda(_t, _t*log(10*_t + x + 1)))`

### Maxima [F]

$$\begin{aligned} & \int \frac{1}{5 + 15x + 30x^2 + 30x^3 + 15x^4 + 3x^5} dx \\ & = \int \frac{1}{3x^5 + 15x^4 + 30x^3 + 30x^2 + 15x + 5} dx \end{aligned}$$

input `integrate(1/(3*x^5+15*x^4+30*x^3+30*x^2+15*x+5),x, algorithm="maxima")`



output `integrate(1/(3*x^5 + 15*x^4 + 30*x^3 + 30*x^2 + 15*x + 5), x)`

### Giac [F]

$$\int \frac{1}{5 + 15x + 30x^2 + 30x^3 + 15x^4 + 3x^5} dx$$

$$= \int \frac{1}{3x^5 + 15x^4 + 30x^3 + 30x^2 + 15x + 5} dx$$

input `integrate(1/(3*x^5+15*x^4+30*x^3+30*x^2+15*x+5),x, algorithm="giac")`

output `integrate(1/(3*x^5 + 15*x^4 + 30*x^3 + 30*x^2 + 15*x + 5), x)`

### Mupad [B] (verification not implemented)

Time = 22.54 (sec) , antiderivative size = 251, normalized size of antiderivative = 0.82

$$\int \frac{1}{5 + 15x + 30x^2 + 30x^3 + 15x^4 + 3x^5} dx = \frac{162^{1/5} \ln \left( x + \frac{162^{1/5}}{3} + 1 \right)}{30}$$

$$- \frac{3^{4/5} \ln \left( x - \frac{3^{4/5} \left( 2^{1/5} (\sqrt{5}+1) + 2^{7/10} \sqrt{\sqrt{5}-5} \right)}{12} + 1 \right) \left( 2^{1/5} (\sqrt{5}+1) + 2^{7/10} \sqrt{\sqrt{5}-5} \right)}{120}$$

$$- \frac{3^{4/5} \ln \left( x - \frac{3^{4/5} \left( 2^{1/5} (\sqrt{5}+1) - 2^{7/10} \sqrt{\sqrt{5}-5} \right)}{12} + 1 \right) \left( 2^{1/5} (\sqrt{5}+1) - 2^{7/10} \sqrt{\sqrt{5}-5} \right)}{120}$$

$$+ \frac{3^{4/5} \ln \left( x + \frac{3^{4/5} \left( 2^{7/10} \sqrt{-\sqrt{5}-5} + 2^{1/5} (\sqrt{5}-1) \right)}{12} + 1 \right) \left( 2^{7/10} \sqrt{-\sqrt{5}-5} + 2^{1/5} (\sqrt{5}-1) \right)}{120}$$

$$- \frac{3^{4/5} \ln \left( x - \frac{3^{4/5} \left( 2^{7/10} \sqrt{-\sqrt{5}-5} - 2^{1/5} (\sqrt{5}-1) \right)}{12} + 1 \right) \left( 2^{7/10} \sqrt{-\sqrt{5}-5} - 2^{1/5} (\sqrt{5}-1) \right)}{120}$$

input `int(1/(15*x + 30*x^2 + 30*x^3 + 15*x^4 + 3*x^5 + 5),x)`

output

```
(162^(1/5)*log(x + 162^(1/5)/3 + 1))/30 - (3^(4/5)*log(x - (3^(4/5)*(2^(1/5)*5^(1/2) + 1) + 2^(7/10)*(5^(1/2) - 5)^(1/2)))/12 + 1)*(2^(1/5)*5^(1/2) + 1) + 2^(7/10)*(5^(1/2) - 5)^(1/2))/120 - (3^(4/5)*log(x - (3^(4/5)*(2^(1/5)*5^(1/2) + 1) - 2^(7/10)*(5^(1/2) - 5)^(1/2)))/12 + 1)*(2^(1/5)*5^(1/2) + 1) - 2^(7/10)*(5^(1/2) - 5)^(1/2))/120 + (3^(4/5)*log(x + (3^(4/5)*(2^(7/10)*(- 5^(1/2) - 5)^(1/2) + 2^(1/5)*5^(1/2) - 1)))/12 + 1)*(2^(7/10)*(- 5^(1/2) - 5)^(1/2) + 2^(1/5)*5^(1/2) - 1))/120 - (3^(4/5)*log(x - (3^(4/5)*(2^(7/10)*(- 5^(1/2) - 5)^(1/2) - 2^(1/5)*5^(1/2) - 1)))/12 + 1)*(2^(7/10)*(- 5^(1/2) - 5)^(1/2) - 2^(1/5)*5^(1/2) - 1))/120
```

**Reduce [F]**

$$\int \frac{1}{5 + 15x + 30x^2 + 30x^3 + 15x^4 + 3x^5} dx = \int \frac{1}{3x^5 + 15x^4 + 30x^3 + 30x^2 + 15x + 5} dx$$

input

```
int(1/(3*x^5+15*x^4+30*x^3+30*x^2+15*x+5),x)
```

output

```
int(1/(3*x**5 + 15*x**4 + 30*x**3 + 30*x**2 + 15*x + 5),x)
```

### 3.28 $\int \frac{1}{2+2(1+x)^3-3(1+x)^6} dx$

Optimal result	210
Mathematica [C] (verified)	211
Rubi [A] (verified)	211
Maple [C] (verified)	220
Fricas [A] (verification not implemented)	221
Sympy [A] (verification not implemented)	222
Maxima [F]	222
Giac [F]	223
Mupad [B] (verification not implemented)	223
Reduce [F]	224

#### Optimal result

Integrand size = 18, antiderivative size = 337

$$\begin{aligned}
 & \int \frac{1}{2+2(1+x)^3-3(1+x)^6} dx \\
 &= \frac{\sqrt[6]{3} \arctan\left(\frac{1-2\sqrt[3]{\frac{3}{-1+\sqrt{7}}(1+x)}}{\sqrt{3}}\right)}{2\sqrt{7}(-1+\sqrt{7})^{2/3}} + \frac{\sqrt[6]{3} \arctan\left(\frac{1+2\sqrt[3]{\frac{3}{1+\sqrt{7}}(1+x)}}{\sqrt{3}}\right)}{2\sqrt{7}(1+\sqrt{7})^{2/3}} \\
 & - \frac{\log\left(\sqrt[3]{1+\sqrt{7}} - \sqrt[3]{3}(1+x)\right)}{2\sqrt[3]{3}\sqrt{7}(1+\sqrt{7})^{2/3}} + \frac{\log\left(\sqrt[3]{-1+\sqrt{7}} + \sqrt[3]{3}(1+x)\right)}{2\sqrt[3]{3}\sqrt{7}(-1+\sqrt{7})^{2/3}} \\
 & - \frac{\log\left((-1+\sqrt{7})^{2/3} - \sqrt[3]{3}(-1+\sqrt{7})(1+x) + 3^{2/3}(1+x)^2\right)}{4\sqrt[3]{3}\sqrt{7}(-1+\sqrt{7})^{2/3}} \\
 & + \frac{\log\left((1+\sqrt{7})^{2/3} + \sqrt[3]{3}(1+\sqrt{7})(1+x) + 3^{2/3}(1+x)^2\right)}{4\sqrt[3]{3}\sqrt{7}(1+\sqrt{7})^{2/3}}
 \end{aligned}$$

output

$$\begin{aligned}
& -1/14*3^{(1/6)}*\arctan(1/3*(1-2*3^{(1/3)}*(1/(-1+7^{(1/2))})^{(1/3)}*(1+x))*3^{(1/2)} \\
& )*7^{(1/2)/(-1+7^{(1/2)})^{(2/3)}+1/14*3^{(1/6)}*\arctan(1/3*(1+2*3^{(1/3)}*(1/(1+7 \\
& ^{(1/2))})^{(1/3)}*(1+x))*3^{(1/2)})*7^{(1/2)/(1+7^{(1/2)})^{(2/3)}-1/42*\ln((1+7^{(1/2)} \\
& )^{(1/3)}-3^{(1/3)}*(1+x))*3^{(2/3)}*7^{(1/2)/(1+7^{(1/2)})^{(2/3)}+1/42*\ln((-1+7^{(1/2)} \\
& )^{(1/3)}+3^{(1/3)}*(1+x))*3^{(2/3)}*7^{(1/2)/(-1+7^{(1/2)})^{(2/3)}-1/84*\ln((-1+7 \\
& ^{(1/2)})^{(2/3)}-(-3+3*7^{(1/2)})^{(1/3)}*(1+x)+3^{(2/3)}*(1+x)^2)*3^{(2/3)}*7^{(1/2)/ \\
& (-1+7^{(1/2)})^{(2/3)}+1/84*\ln((1+7^{(1/2)})^{(2/3)}+(3+3*7^{(1/2)})^{(1/3)}*(1+x)+3^{(2/3)}*(1+x)^2)*3^{(2/3)}*7^{(1/2)/(1+7^{(1/2)})^{(2/3)}
\end{aligned}$$
**Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.02 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.13

$$\begin{aligned}
& \int \frac{1}{2 + 2(1+x)^3 - 3(1+x)^6} dx \\
& = -\frac{1}{6} \text{RootSum} \left[ -2 - 2\#1^3 + 3\#1^6 \&, \frac{\log(1+x-\#1)}{-\#1^2 + 3\#1^5} \& \right]
\end{aligned}$$

input

$$\text{Integrate}[(2 + 2*(1 + x)^3 - 3*(1 + x)^6)^{-1}, x]$$

output

$$-1/6*\text{RootSum}[-2 - 2*\#1^3 + 3*\#1^6 \& , \text{Log}[1 + x - \#1]/(-\#1^2 + 3*\#1^5) \& ]$$
**Rubi [A] (verified)**

Time = 0.95 (sec) , antiderivative size = 310, normalized size of antiderivative = 0.92, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$ , Rules used = {1687, 1685, 750, 16, 25, 1142, 25, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{-3(x+1)^6 + 2(x+1)^3 + 2} dx$$

↓ 1687

$$\begin{aligned}
 & \int \frac{1}{-3(x+1)^6 + 2(x+1)^3 + 2} d(x+1) \\
 & \quad \downarrow 1685 \\
 & \frac{3 \int \frac{1}{-3(x+1)^3 + \sqrt{7} + 1} d(x+1)}{2\sqrt{7}} - \frac{3 \int \frac{1}{-3(x+1)^3 - \sqrt{7} + 1} d(x+1)}{2\sqrt{7}} \\
 & \quad \downarrow 750 \\
 & \frac{3 \left( \frac{\int \frac{1}{\sqrt[3]{1+\sqrt{7}} - \sqrt[3]{3(x+1)}} d(x+1)}{3(1+\sqrt{7})^{2/3}} + \frac{\int \frac{\sqrt[3]{3(x+1)+2}\sqrt[3]{1+\sqrt{7}}}{3^{2/3(x+1)+2} + \sqrt[3]{3(1+\sqrt{7})^{(x+1)+(1+\sqrt{7})^{2/3}}}} d(x+1)}{3(1+\sqrt{7})^{2/3}} \right)}{2\sqrt{7}} - \\
 & \frac{3 \left( \frac{\int \frac{1}{-\sqrt[3]{3(x+1)} - \sqrt[3]{-1+\sqrt{7}}} d(x+1)}{3(\sqrt{7}-1)^{2/3}} + \frac{\int -\frac{2\sqrt[3]{-1+\sqrt{7}} - \sqrt[3]{3(x+1)}}{3^{2/3(x+1)+2} - \sqrt[3]{3(-1+\sqrt{7})^{(x+1)+(-1+\sqrt{7})^{2/3}}}} d(x+1)}{3(\sqrt{7}-1)^{2/3}} \right)}{2\sqrt{7}} \\
 & \quad \downarrow 16 \\
 & \frac{3 \left( \frac{\int \frac{\sqrt[3]{3(x+1)+2}\sqrt[3]{1+\sqrt{7}}}{3^{2/3(x+1)+2} + \sqrt[3]{3(1+\sqrt{7})^{(x+1)+(1+\sqrt{7})^{2/3}}}} d(x+1)}{3(1+\sqrt{7})^{2/3}} - \frac{\log\left(\sqrt[3]{1+\sqrt{7}} - \sqrt[3]{3(x+1)}\right)}{3\sqrt[3]{3}(1+\sqrt{7})^{2/3}} \right)}{2\sqrt{7}} - \\
 & \frac{3 \left( \frac{\int -\frac{2\sqrt[3]{-1+\sqrt{7}} - \sqrt[3]{3(x+1)}}{3^{2/3(x+1)+2} - \sqrt[3]{3(-1+\sqrt{7})^{(x+1)+(-1+\sqrt{7})^{2/3}}}} d(x+1)}{3(\sqrt{7}-1)^{2/3}} - \frac{\log\left(\sqrt[3]{3(x+1)+\sqrt[3]{\sqrt{7}-1}}\right)}{3\sqrt[3]{3}(\sqrt{7}-1)^{2/3}} \right)}{2\sqrt{7}}
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow 25 \\
 3 \left( \frac{\int \frac{\sqrt[3]{3(x+1)+2} \sqrt[3]{1+\sqrt{7}}}{3^{2/3}(x+1)^2 + \sqrt[3]{3(1+\sqrt{7})(x+1)+(1+\sqrt{7})^{2/3}}} d(x+1)}{3(1+\sqrt{7})^{2/3}} - \frac{\log\left(\sqrt[3]{1+\sqrt{7}} - \sqrt[3]{3(x+1)}\right)}{3\sqrt[3]{3(1+\sqrt{7})^{2/3}}} \right) \\
 \hline
 2\sqrt{7} \\
 3 \left( -\frac{\int \frac{\sqrt[3]{-1+\sqrt{7}} - \sqrt[3]{3(x+1)}}{3^{2/3}(x+1)^2 - \sqrt[3]{3(-1+\sqrt{7})(x+1)+(-1+\sqrt{7})^{2/3}}} d(x+1)}{3(\sqrt{7}-1)^{2/3}} - \frac{\log\left(\sqrt[3]{3(x+1)} + \sqrt[3]{\sqrt{7}-1}\right)}{3\sqrt[3]{3(\sqrt{7}-1)^{2/3}}} \right) \\
 \hline
 2\sqrt{7} \\
 \downarrow 1142
 \end{array}$$

$$3 \left( \frac{\frac{3}{2} \sqrt[3]{1 + \sqrt{7}} \int \frac{1}{3^{2/3(x+1)^2 + \sqrt[3]{3(1 + \sqrt{7})(x+1) + (1 + \sqrt{7})^{2/3}}} dx + \frac{2 \cdot 3^{2/3(x+1)} \sqrt[3]{3(1 + \sqrt{7})} \int \frac{1}{3^{2/3(x+1)^2 + \sqrt[3]{3(1 + \sqrt{7})(x+1) + (1 + \sqrt{7})^{2/3}}} dx}{2 \sqrt[3]{3}}}{3(1 + \sqrt{7})^{2/3}} \right) \log$$

$$3 \left( \frac{\frac{3}{2} \sqrt[3]{\sqrt{7} - 1} \int \frac{1}{3^{2/3(x+1)^2 - \sqrt[3]{3(-1 + \sqrt{7})(x+1) + (-1 + \sqrt{7})^{2/3}}} dx + \frac{2\sqrt{7} \sqrt[3]{3(-1 + \sqrt{7})} \int \frac{1}{3^{2/3(x+1)^2 - \sqrt[3]{3(-1 + \sqrt{7})(x+1) + (-1 + \sqrt{7})^{2/3}}} dx}{2 \sqrt[3]{3}}}{3(\sqrt{7} - 1)^{2/3}} \right)$$

$2\sqrt{7}$

↓ 25

$$\left( \begin{array}{l}
 \frac{3}{2} \sqrt[3]{1 + \sqrt{7}} f \frac{1}{3^{2/3}(x+1)^2 + \sqrt[3]{3(1 + \sqrt{7})}^{(x+1)+(1+\sqrt{7})^{2/3}}} d(x+1) + \frac{2 \cdot 3^{2/3}(x+1) + \sqrt[3]{3(1 + \sqrt{7})}}{3^{2/3}(x+1)^2 + \sqrt[3]{3(1 + \sqrt{7})}^{(x+1)+(1+\sqrt{7})^{2/3}}} d(x+1) \\
 \frac{2 \sqrt[3]{3}}{3(1+\sqrt{7})^{2/3}} \log
 \end{array} \right)$$

$$\left( \begin{array}{l}
 \frac{3}{2} \sqrt[3]{\sqrt{7} - 1} f \frac{1}{3^{2/3}(x+1)^2 - \sqrt[3]{3(-1 + \sqrt{7})}^{(x+1)+(-1+\sqrt{7})^{2/3}}} d(x+1) + \frac{2\sqrt{7} \sqrt[3]{3(-1 + \sqrt{7})}^{-2} 3^{2/3}(x+1)}{3^{2/3}(x+1)^2 - \sqrt[3]{3(-1 + \sqrt{7})}^{(x+1)+(-1+\sqrt{7})^{2/3}}} d(x+1) \\
 \frac{2\sqrt{7}}{3(\sqrt{7}-1)^{2/3}}
 \end{array} \right)$$

$2\sqrt{7}$

↓ 1082



$$\left( \frac{\int \frac{2 \cdot 3^{2/3(x+1)} \sqrt[3]{3(1+\sqrt{7})}}{3^{2/3(x+1)^2 + \sqrt[3]{3(1+\sqrt{7})(x+1) + (1+\sqrt{7})^{2/3}}} dx}{2 \sqrt[3]{3}} - 3^{2/3} \int \frac{1}{\left(2 \sqrt[3]{\frac{3}{1+\sqrt{7}}(x+1)+1}\right)^2} d\left(2 \sqrt[3]{\frac{3}{1+\sqrt{7}}(x+1)+1}\right) - \log\left(\sqrt[3]{\frac{3}{1+\sqrt{7}}(x+1)+1}\right)}{3(1+\sqrt{7})^{2/3}} \right)$$

$2\sqrt{7}$

$$\left( \frac{\int \frac{\sqrt[3]{3(-1+\sqrt{7})} \cdot 3^{2/3(x+1)}}{3^{2/3(x+1)^2 - \sqrt[3]{3(-1+\sqrt{7})(x+1) + (-1+\sqrt{7})^{2/3}}} dx}{2 \sqrt[3]{3}} + 3^{2/3} \int \frac{1}{\left(1-2 \sqrt[3]{\frac{3}{-1+\sqrt{7}}(x+1)}\right)^2} d\left(1-2 \sqrt[3]{\frac{3}{-1+\sqrt{7}}(x+1)}\right) - \log\left(\sqrt[3]{\frac{3}{-1+\sqrt{7}}(x+1)}\right)}{3(\sqrt{7}-1)^{2/3}} \right)$$

$2\sqrt{7}$

$$3 \left( \frac{\int \frac{2 \cdot 3^{2/3(x+1)} + \sqrt[3]{3(1+\sqrt{7})}}{3^{2/3(x+1)^2 + \sqrt[3]{3(1+\sqrt{7})(x+1) + (1+\sqrt{7})^{2/3}}} dx}{2 \sqrt[3]{3}} + \sqrt[6]{3} \arctan \left( \frac{\sqrt[2]{3} \sqrt[3]{\frac{3}{1+\sqrt{7}}(x+1)+1}}{\sqrt{3}} \right) - \frac{\log \left( \sqrt[3]{1+\sqrt{7}} - \sqrt[3]{3(x+1)} \right)}{3 \sqrt[3]{3}(1+\sqrt{7})^{2/3}} \right)$$

$2\sqrt{7}$

$$3 \left( \frac{\int \frac{\sqrt[3]{3(-1+\sqrt{7})} - 2 \cdot 3^{2/3(x+1)}}{3^{2/3(x+1)^2 - \sqrt[3]{3(-1+\sqrt{7})(x+1) + (-1+\sqrt{7})^{2/3}}} dx}{2 \sqrt[3]{3}} - \sqrt[6]{3} \arctan \left( \frac{1-2 \sqrt[3]{\frac{3}{\sqrt{7}-1}}(x+1)}}{\sqrt{3}} \right) - \frac{\log \left( \sqrt[3]{3(x+1)} + \sqrt[3]{\sqrt{7}-1} \right)}{3 \sqrt[3]{3}(\sqrt{7}-1)^{2/3}} \right)$$

$2\sqrt{7}$

↓ 1103

$$\frac{3 \left( \frac{\sqrt[6]{3} \arctan \left( \frac{\sqrt[2]{3} \sqrt[3]{\frac{3}{1+\sqrt{7}}(x+1)+1}}{\sqrt{3}} \right) + \frac{\log \left( 3^{2/3}(x+1)^2 + \sqrt[3]{3(1+\sqrt{7})}^{(x+1)+(1+\sqrt{7})^{2/3}} \right)}{2\sqrt[3]{3}} \right)}{3(1+\sqrt{7})^{2/3}} - \frac{\log \left( \sqrt[3]{1+\sqrt{7}} - \sqrt[3]{3(x+1)} \right)}{3\sqrt[3]{3}(1+\sqrt{7})^{2/3}} \right)}{2\sqrt{7}}$$

$$\frac{3 \left( -\sqrt[6]{3} \arctan \left( \frac{1-2\sqrt[3]{\frac{3}{\sqrt{7}-1}}(x+1)}}{\sqrt{3}} \right) - \frac{\log \left( 3^{2/3}(x+1)^2 - \sqrt[3]{3(\sqrt{7}-1)}^{(x+1)+(\sqrt{7}-1)^{2/3}} \right)}{2\sqrt[3]{3}} \right)}{3(\sqrt{7}-1)^{2/3}} - \frac{\log \left( \sqrt[3]{3(x+1)} + \sqrt[3]{\sqrt{7}-1} \right)}{3\sqrt[3]{3}(\sqrt{7}-1)^{2/3}} \right)}{2\sqrt{7}}$$

input `Int[(2 + 2*(1 + x)^3 - 3*(1 + x)^6)^(-1),x]`

output `(-3*(-1/3*Log[(-1 + Sqrt[7])^(1/3) + 3^(1/3)*(1 + x)]/(3^(1/3)*(-1 + Sqrt[7])^(2/3)) - (-3^(1/6)*ArcTan[(1 - 2*(3/(-1 + Sqrt[7]))^(1/3)*(1 + x))/Sqrt[3]]) - Log[(-1 + Sqrt[7])^(2/3) - (3*(-1 + Sqrt[7]))^(1/3)*(1 + x) + 3^(2/3)*(1 + x)^2]/(2*3^(1/3)))/(3*(-1 + Sqrt[7])^(2/3)))/(2*Sqrt[7]) + (3*(-1/3*Log[(1 + Sqrt[7])^(1/3) - 3^(1/3)*(1 + x)]/(3^(1/3)*(1 + Sqrt[7])^(2/3)) + (3^(1/6)*ArcTan[(1 + 2*(3/(1 + Sqrt[7]))^(1/3)*(1 + x))/Sqrt[3]]) + Log[(1 + Sqrt[7])^(2/3) + (3*(1 + Sqrt[7]))^(1/3)*(1 + x) + 3^(2/3)*(1 + x)^2]/(2*3^(1/3)))/(3*(1 + Sqrt[7])^(2/3)))/(2*Sqrt[7])`

## Defintions of rubi rules used

- rule 16  $\text{Int}[(c\_)/((a\_)+(b\_)*(x\_)), x\_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 25  $\text{Int}[-(Fx\_), x\_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[Fx, x], x]$
- rule 217  $\text{Int}[(a\_)+(b\_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(-(\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b] \& \& (\text{LtQ}[a, 0] \parallel \text{LtQ}[b, 0])]$
- rule 750  $\text{Int}[(a\_)+(b\_)*(x_)^3)^{-1}, x\_Symbol] \rightarrow \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[(2*\text{Rt}[a, 3] - \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$
- rule 1082  $\text{Int}[(a\_)+(b\_)*(x_)+(c\_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S \text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \&\& (\text{EqQ}[q^2, 1] \parallel \text{!RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103  $\text{Int}[(d\_)+(e\_)*(x_)/((a\_)+(b\_)*(x_)+(c\_)*(x_)^2), x\_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{EqQ}[2*c*d - b*e, 0]$
- rule 1142  $\text{Int}[(d\_)+(e\_)*(x_)/((a\_)+(b\_)*(x_)+(c\_)*(x_)^2), x\_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \text{ Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \text{ Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$
- rule 1685  $\text{Int}[(a\_)+(b\_)*(x_)^{(n\_)}+(c\_)*(x_)^{(n2\_)}^{-1}, x\_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b^2 - 4*a*c, 2]\}, \text{Simp}[c/q \text{ Int}[1/(b/2 - q/2 + c*x^n), x], x] - \text{Simp}[c/q \text{ Int}[1/(b/2 + q/2 + c*x^n), x], x]] /; \text{FreeQ}[\{a, b, c\}, x] \&\& \text{EqQ}[n2, 2*n] \&\& \text{NeQ}[b^2 - 4*a*c, 0]$

rule 1687

```
Int[((a_) + (c_.)*(u_)^(n2_.) + (b_.)*(u_)^(n_))^(p_), x_Symbol] := Simp[1/
Coefficient[u, x, 1] Subst[Int[(a + b*x^n + c*x^(2*n))^p, x], x, u], x] /
; FreeQ[{a, b, c, n, p}, x] && EqQ[n2, 2*n] && LinearQ[u, x] && NeQ[u, x]
```

**Maple [C] (verified)**

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.04 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.21

method	result	size
default	$-\frac{\left( \sum_{R=\text{RootOf}(3Z^6+18Z^5+45Z^4+58Z^3+39Z^2+12Z-1)} \frac{\ln(x-R)}{3R^5+15R^4+30R^3+29R^2+13R+2} \right)}{6}$	71
risch	$-\frac{\left( \sum_{R=\text{RootOf}(3Z^6+18Z^5+45Z^4+58Z^3+39Z^2+12Z-1)} \frac{\ln(x-R)}{3R^5+15R^4+30R^3+29R^2+13R+2} \right)}{6}$	71

input

```
int(1/(2+2*(x+1)^3-3*(x+1)^6),x,method=_RETURNVERBOSE)
```

output

```
-1/6*sum(1/(3*_R^5+15*_R^4+30*_R^3+29*_R^2+13*_R+2)*ln(x-_R),_R=RootOf(3*_
Z^6+18*_Z^5+45*_Z^4+58*_Z^3+39*_Z^2+12*_Z-1))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 260, normalized size of antiderivative = 0.77

$$\begin{aligned}
& \int \frac{1}{2+2(1+x)^3-3(1+x)^6} dx = \\
& -\frac{1}{12} \left( \frac{2}{49} \sqrt{7} + \frac{1}{14} \right)^{\frac{1}{3}} (\sqrt{-3} + 1) \log \left( \left( \sqrt{7}(\sqrt{-3} + 1) - 7\sqrt{-3} - 7 \right) \left( \frac{2}{49} \sqrt{7} + \frac{1}{14} \right)^{\frac{1}{3}} \right. \\
& \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \left. + 6x + 6 \right) \\
& + \frac{1}{12} \left( \frac{2}{49} \sqrt{7} + \frac{1}{14} \right)^{\frac{1}{3}} (\sqrt{-3} - 1) \log \left( - \left( \sqrt{7}(\sqrt{-3} - 1) - 7\sqrt{-3} + 7 \right) \left( \frac{2}{49} \sqrt{7} + \frac{1}{14} \right)^{\frac{1}{3}} \right. \\
& \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \left. + 6x + 6 \right) \\
& - \frac{1}{12} \left( -\frac{2}{49} \sqrt{7} + \frac{1}{14} \right)^{\frac{1}{3}} (\sqrt{-3} + 1) \log \left( - \left( \sqrt{7}(\sqrt{-3} + 1) + 7\sqrt{-3} + 7 \right) \left( -\frac{2}{49} \sqrt{7} + \frac{1}{14} \right)^{\frac{1}{3}} \right. \\
& \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \left. + 6x + 6 \right) \\
& + \frac{1}{12} \left( -\frac{2}{49} \sqrt{7} + \frac{1}{14} \right)^{\frac{1}{3}} (\sqrt{-3} - 1) \log \left( \left( \sqrt{7}(\sqrt{-3} - 1) + 7\sqrt{-3} - 7 \right) \left( -\frac{2}{49} \sqrt{7} + \frac{1}{14} \right)^{\frac{1}{3}} \right. \\
& \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \left. + 6x + 6 \right) + \frac{1}{6} \left( \frac{2}{49} \sqrt{7} + \frac{1}{14} \right)^{\frac{1}{3}} \log \left( - \left( \sqrt{7} - 7 \right) \left( \frac{2}{49} \sqrt{7} + \frac{1}{14} \right)^{\frac{1}{3}} + 3x + 3 \right) \\
& + \frac{1}{6} \left( -\frac{2}{49} \sqrt{7} + \frac{1}{14} \right)^{\frac{1}{3}} \log \left( \left( \sqrt{7} + 7 \right) \left( -\frac{2}{49} \sqrt{7} + \frac{1}{14} \right)^{\frac{1}{3}} + 3x + 3 \right)
\end{aligned}$$

input `integrate(1/(2+2*(1+x)^3-3*(1+x)^6),x, algorithm="fricas")`

output

```
-1/12*(2/49*sqrt(7) + 1/14)^(1/3)*(sqrt(-3) + 1)*log((sqrt(7)*(sqrt(-3) + 1) - 7*sqrt(-3) - 7)*(2/49*sqrt(7) + 1/14)^(1/3) + 6*x + 6) + 1/12*(2/49*sqrt(7) + 1/14)^(1/3)*(sqrt(-3) - 1)*log(-(sqrt(7)*(sqrt(-3) - 1) - 7*sqrt(-3) + 7)*(2/49*sqrt(7) + 1/14)^(1/3) + 6*x + 6) - 1/12*(-2/49*sqrt(7) + 1/14)^(1/3)*(sqrt(-3) + 1)*log(-(sqrt(7)*(sqrt(-3) + 1) + 7*sqrt(-3) + 7)*(-2/49*sqrt(7) + 1/14)^(1/3) + 6*x + 6) + 1/12*(-2/49*sqrt(7) + 1/14)^(1/3)*(sqrt(-3) - 1)*log((sqrt(7)*(sqrt(-3) - 1) + 7*sqrt(-3) - 7)*(-2/49*sqrt(7) + 1/14)^(1/3) + 6*x + 6) + 1/6*(2/49*sqrt(7) + 1/14)^(1/3)*log(-(sqrt(7) - 7)*(2/49*sqrt(7) + 1/14)^(1/3) + 3*x + 3) + 1/6*(-2/49*sqrt(7) + 1/14)^(1/3)*log((sqrt(7) + 7)*(-2/49*sqrt(7) + 1/14)^(1/3) + 3*x + 3)
```

**Sympy [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.09

$$\int \frac{1}{2 + 2(1+x)^3 - 3(1+x)^6} dx$$

$$= -\text{RootSum}\left(7112448t^6 + 4704t^3 - 1, \left(t \mapsto t \log\left(-10584t^4 - \frac{35t}{2} + x + 1\right)\right)\right)$$

input

```
integrate(1/(2+2*(1+x)**3-3*(1+x)**6),x)
```

output

```
-RootSum(7112448*_t**6 + 4704*_t**3 - 1, Lambda(_t, _t*log(-10584*_t**4 - 35*_t/2 + x + 1)))
```

**Maxima [F]**

$$\int \frac{1}{2 + 2(1+x)^3 - 3(1+x)^6} dx = \int -\frac{1}{3(x+1)^6 - 2(x+1)^3 - 2} dx$$

input

```
integrate(1/(2+2*(1+x)^3-3*(1+x)^6),x, algorithm="maxima")
```

output

```
-integrate(1/(3*(x + 1)^6 - 2*(x + 1)^3 - 2), x)
```

**Giac [F]**

$$\int \frac{1}{2 + 2(1+x)^3 - 3(1+x)^6} dx = \int -\frac{1}{3(x+1)^6 - 2(x+1)^3 - 2} dx$$

input `integrate(1/(2+2*(1+x)^3-3*(1+x)^6),x, algorithm="giac")`

output `integrate(-1/(3*(x + 1)^6 - 2*(x + 1)^3 - 2), x)`

**Mupad [B] (verification not implemented)**

Time = 24.95 (sec) , antiderivative size = 463, normalized size of antiderivative = 1.37

$$\begin{aligned} & \int \frac{1}{2 + 2(1+x)^3 - 3(1+x)^6} dx \\ &= \frac{\ln \left( x + \frac{2^{2/3} 7^{1/3} (7-4\sqrt{7})^{1/3}}{6} + \frac{2^{2/3} 7^{5/6} (7-4\sqrt{7})^{1/3}}{42} + 1 \right) (196 - 112\sqrt{7})^{1/3}}{84} \\ & - \frac{\ln \left( x - \frac{2^{2/3} 7^{1/3} (4\sqrt{7}-7)^{1/3}}{6} - \frac{2^{2/3} 7^{5/6} (4\sqrt{7}-7)^{1/3}}{42} + 1 \right) (112\sqrt{7} - 196)^{1/3}}{84} \\ & + \frac{\ln \left( x + \frac{2^{2/3} 7^{1/3} (4\sqrt{7}+7)^{1/3}}{6} - \frac{2^{2/3} 7^{5/6} (4\sqrt{7}+7)^{1/3}}{42} + 1 \right) (112\sqrt{7} + 196)^{1/3}}{84} \\ & - \frac{2^{2/3} 7^{1/3} \ln \left( x - \frac{2^{2/3} 7^{1/3} (7-4\sqrt{7})^{1/3}}{12} - \frac{2^{2/3} 7^{5/6} (7-4\sqrt{7})^{1/3}}{84} + 1 - \frac{2^{2/3} \sqrt{3} 7^{1/3} (7-4\sqrt{7})^{1/3}}{12} \operatorname{li} - \frac{2^{2/3} \sqrt{3} 7^{5/6} (7-4\sqrt{7})^{1/3}}{84} \right)}{168} \\ & - \frac{2^{2/3} 7^{1/3} \ln \left( x - \frac{2^{2/3} 7^{1/3} (4\sqrt{7}+7)^{1/3}}{12} + \frac{2^{2/3} 7^{5/6} (4\sqrt{7}+7)^{1/3}}{84} + 1 - \frac{2^{2/3} \sqrt{3} 7^{1/3} (4\sqrt{7}+7)^{1/3}}{12} \operatorname{li} + \frac{2^{2/3} \sqrt{3} 7^{5/6} (4\sqrt{7}+7)^{1/3}}{84} \right)}{168} \\ & + \frac{2^{2/3} 7^{1/3} \ln \left( x - \frac{2^{2/3} 7^{1/3} (4\sqrt{7}+7)^{1/3}}{12} + \frac{2^{2/3} 7^{5/6} (4\sqrt{7}+7)^{1/3}}{84} + 1 + \frac{2^{2/3} \sqrt{3} 7^{1/3} (4\sqrt{7}+7)^{1/3}}{12} \operatorname{li} - \frac{2^{2/3} \sqrt{3} 7^{5/6} (4\sqrt{7}+7)^{1/3}}{84} \right)}{168} \end{aligned}$$

168



input `int(1/(2*(x + 1)^3 - 3*(x + 1)^6 + 2),x)`

output  $(\log(x + (2^{2/3} \cdot 7^{1/3}) \cdot (7 - 4 \cdot 7^{1/2}))^{1/3})/6 + (2^{2/3} \cdot 7^{5/6}) \cdot (7 - 4 \cdot 7^{1/2})^{1/3}/42 + 1) \cdot (196 - 112 \cdot 7^{1/2})^{1/3}/84 - (\log(x - (2^{2/3} \cdot 7^{1/3}) \cdot (4 \cdot 7^{1/2} - 7)^{1/3})/6 - (2^{2/3} \cdot 7^{5/6}) \cdot (4 \cdot 7^{1/2} - 7)^{1/3})/42 + 1) \cdot (112 \cdot 7^{1/2} - 196)^{1/3}/84 + (\log(x + (2^{2/3} \cdot 7^{1/3}) \cdot (4 \cdot 7^{1/2} + 7)^{1/3})/6 - (2^{2/3} \cdot 7^{5/6}) \cdot (4 \cdot 7^{1/2} + 7)^{1/3})/42 + 1) \cdot (112 \cdot 7^{1/2} + 196)^{1/3}/84 - (2^{2/3} \cdot 7^{1/3}) \cdot \log(x - (2^{2/3} \cdot 7^{1/3}) \cdot (7 - 4 \cdot 7^{1/2}))^{1/3})/12 - (2^{2/3} \cdot 7^{5/6}) \cdot (7 - 4 \cdot 7^{1/2})^{1/3}/84 - (2^{2/3} \cdot 3^{1/2} \cdot 7^{1/3}) \cdot (7 - 4 \cdot 7^{1/2})^{1/3} \cdot i/12 - (2^{2/3} \cdot 3^{1/2} \cdot 7^{5/6}) \cdot (7 - 4 \cdot 7^{1/2})^{1/3} \cdot i/84 + 1) \cdot (3^{1/2} \cdot i + 1) \cdot (7 - 4 \cdot 7^{1/2})^{1/3})/168 - (2^{2/3} \cdot 7^{1/3}) \cdot \log(x - (2^{2/3} \cdot 7^{1/3}) \cdot (4 \cdot 7^{1/2} + 7)^{1/3})/12 + (2^{2/3} \cdot 7^{5/6}) \cdot (4 \cdot 7^{1/2} + 7)^{1/3})/84 - (2^{2/3} \cdot 3^{1/2} \cdot 7^{1/3}) \cdot (4 \cdot 7^{1/2} + 7)^{1/3} \cdot i/12 + (2^{2/3} \cdot 3^{1/2} \cdot 7^{5/6}) \cdot (4 \cdot 7^{1/2} + 7)^{1/3} \cdot i/84 + 1) \cdot (3^{1/2} \cdot i + 1) \cdot (4 \cdot 7^{1/2} + 7)^{1/3})/168 + (2^{2/3} \cdot 7^{1/3}) \cdot \log(x - (2^{2/3} \cdot 7^{1/3}) \cdot (4 \cdot 7^{1/2} + 7)^{1/3})/12 + (2^{2/3} \cdot 7^{5/6}) \cdot (4 \cdot 7^{1/2} + 7)^{1/3})/84 + (2^{2/3} \cdot 3^{1/2} \cdot 7^{1/3}) \cdot (4 \cdot 7^{1/2} + 7)^{1/3} \cdot i/12 - (2^{2/3} \cdot 3^{1/2} \cdot 7^{5/6}) \cdot (4 \cdot 7^{1/2} + 7)^{1/3} \cdot i/84 + 1) \cdot (3^{1/2} \cdot i - 1) \cdot (4 \cdot 7^{1/2} + 7)^{1/3})/168$

## Reduce [F]

$$\int \frac{1}{2 + 2(1+x)^3 - 3(1+x)^6} dx$$

$$= - \left( \int \frac{1}{3x^6 + 18x^5 + 45x^4 + 58x^3 + 39x^2 + 12x - 1} dx \right)$$

input `int(1/(2+2*(1+x)^3-3*(1+x)^6),x)`

output `- int(1/(3*x**6 + 18*x**5 + 45*x**4 + 58*x**3 + 39*x**2 + 12*x - 1),x)`

**3.29**  $\int \frac{1}{1-12x-39x^2-58x^3-45x^4-18x^5-3x^6} dx$

Optimal result	225
Mathematica [C] (verified)	226
Rubi [A] (verified)	226
Maple [C] (verified)	235
Fricas [A] (verification not implemented)	236
Sympy [A] (verification not implemented)	237
Maxima [F]	237
Giac [F]	238
Mupad [B] (verification not implemented)	239
Reduce [F]	240

**Optimal result**

Integrand size = 32, antiderivative size = 337

$$\int \frac{1}{1-12x-39x^2-58x^3-45x^4-18x^5-3x^6} dx$$

$$= -\frac{\sqrt[6]{3} \arctan\left(\frac{1-2\sqrt[3]{\frac{3}{-1+\sqrt{7}}(1+x)}}{\sqrt{3}}\right)}{2\sqrt{7}(-1+\sqrt{7})^{2/3}} + \frac{\sqrt[6]{3} \arctan\left(\frac{1+2\sqrt[3]{\frac{3}{1+\sqrt{7}}(1+x)}}{\sqrt{3}}\right)}{2\sqrt{7}(1+\sqrt{7})^{2/3}}$$

$$- \frac{\log\left(\sqrt[3]{1+\sqrt{7}} - \sqrt[3]{3}(1+x)\right)}{2\sqrt[3]{3}\sqrt{7}(1+\sqrt{7})^{2/3}} + \frac{\log\left(\sqrt[3]{-1+\sqrt{7}} + \sqrt[3]{3}(1+x)\right)}{2\sqrt[3]{3}\sqrt{7}(-1+\sqrt{7})^{2/3}}$$

$$- \frac{\log\left((-1+\sqrt{7})^{2/3} - \sqrt[3]{3}(-1+\sqrt{7})(1+x) + 3^{2/3}(1+x)^2\right)}{4\sqrt[3]{3}\sqrt{7}(-1+\sqrt{7})^{2/3}}$$

$$+ \frac{\log\left((1+\sqrt{7})^{2/3} + \sqrt[3]{3}(1+\sqrt{7})(1+x) + 3^{2/3}(1+x)^2\right)}{4\sqrt[3]{3}\sqrt{7}(1+\sqrt{7})^{2/3}}$$

output

$$\begin{aligned}
& -1/14*3^{(1/6)}*\arctan(1/3*(1-2*3^{(1/3)}*(1/(-1+7^{(1/2))})^{(1/3)}*(1+x))*3^{(1/2)} \\
& )*7^{(1/2)/(-1+7^{(1/2)})^{(2/3)}+1/14*3^{(1/6)}*\arctan(1/3*(1+2*3^{(1/3)}*(1/(1+7^{(1/2)}))^{(1/3)}*(1+x))*3^{(1/2)})*7^{(1/2)/(1+7^{(1/2)})^{(2/3)}-1/42*\ln((1+7^{(1/2)})^{(1/3)}-3^{(1/3)}*(1+x))*3^{(2/3)}*7^{(1/2)/(-1+7^{(1/2)})^{(2/3)}+1/42*\ln((-1+7^{(1/2)})^{(1/3)}+3^{(1/3)}*(1+x))*3^{(2/3)}*7^{(1/2)/(-1+7^{(1/2)})^{(2/3)}-1/84*\ln((-1+7^{(1/2)})^{(2/3)}-(-3+3*7^{(1/2)})^{(1/3)}*(1+x)+3^{(2/3)}*(1+x)^2)*3^{(2/3)}*7^{(1/2)/(-1+7^{(1/2)})^{(2/3)}+1/84*\ln((1+7^{(1/2)})^{(2/3)}+(3+3*7^{(1/2)})^{(1/3)}*(1+x)+3^{(2/3)}*(1+x)^2)*3^{(2/3)}*7^{(1/2)/(1+7^{(1/2)})^{(2/3)}}
\end{aligned}$$
**Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.02 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.25

$$\begin{aligned}
& \int \frac{1}{1 - 12x - 39x^2 - 58x^3 - 45x^4 - 18x^5 - 3x^6} dx \\
& = -\frac{1}{6}\text{RootSum}\left[-1 + 12\#1 + 39\#1^2 + 58\#1^3 + 45\#1^4 + 18\#1^5 \right. \\
& \quad \left. + 3\#1^6 \&, \frac{\log(x - \#1)}{2 + 13\#1 + 29\#1^2 + 30\#1^3 + 15\#1^4 + 3\#1^5} \& \right]
\end{aligned}$$

input

$$\text{Integrate}[(1 - 12*x - 39*x^2 - 58*x^3 - 45*x^4 - 18*x^5 - 3*x^6)^{-1}, x]$$

output

$$\begin{aligned}
& -1/6*\text{RootSum}[-1 + 12*\#1 + 39*\#1^2 + 58*\#1^3 + 45*\#1^4 + 18*\#1^5 + 3*\#1^6 \& \\
& , \text{Log}[x - \#1]/(2 + 13*\#1 + 29*\#1^2 + 30*\#1^3 + 15*\#1^4 + 3*\#1^5) \& ]
\end{aligned}$$
**Rubi [A] (verified)**

Time = 0.88 (sec) , antiderivative size = 310, normalized size of antiderivative = 0.92, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$ , Rules used = {2458, 1685, 750, 16, 25, 1142, 25, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{-3x^6 - 18x^5 - 45x^4 - 58x^3 - 39x^2 - 12x + 1} dx \\
 & \quad \downarrow 2458 \\
 & \int \frac{1}{-3(x+1)^6 + 2(x+1)^3 + 2} d(x+1) \\
 & \quad \downarrow 1685 \\
 & \frac{3 \int \frac{1}{-3(x+1)^3 + \sqrt{7} + 1} d(x+1)}{2\sqrt{7}} - \frac{3 \int \frac{1}{-3(x+1)^3 - \sqrt{7} + 1} d(x+1)}{2\sqrt{7}} \\
 & \quad \downarrow 750 \\
 & \frac{3 \left( \int \frac{1}{\sqrt[3]{1 + \sqrt{7} - \sqrt[3]{3}(x+1)}} d(x+1)}{3(1 + \sqrt{7})^{2/3}} + \int \frac{\sqrt[3]{3}(x+1)^2 \sqrt[3]{1 + \sqrt{7}}}{3^{2/3}(x+1)^2 + \sqrt[3]{3(1 + \sqrt{7})(x+1) + (1 + \sqrt{7})^{2/3}}} d(x+1)}{3(1 + \sqrt{7})^{2/3}} \right)}{2\sqrt{7}} - \\
 & \frac{3 \left( \int \frac{1}{-\sqrt[3]{3}(x+1) - \sqrt[3]{-1 + \sqrt{7}}} d(x+1)}{3(\sqrt{7} - 1)^{2/3}} + \int \frac{\sqrt[3]{-1 + \sqrt{7} - \sqrt[3]{3}(x+1)}}{3^{2/3}(x+1)^2 - \sqrt[3]{3(-1 + \sqrt{7})(x+1) + (-1 + \sqrt{7})^{2/3}}} d(x+1)}{3(\sqrt{7} - 1)^{2/3}} \right)}{2\sqrt{7}} \\
 & \quad \downarrow 16
 \end{aligned}$$

$$\begin{array}{c}
 \left( \int \frac{\sqrt[3]{3^{(x+1)+2}} \sqrt[3]{1+\sqrt{7}}}{3^{2/3(x+1)^2+} \sqrt[3]{3(1+\sqrt{7})^{(x+1)+(1+\sqrt{7})^{2/3}}}} d(x+1) - \frac{\log\left(\sqrt[3]{1+\sqrt{7}} - \sqrt[3]{3^{(x+1)}}\right)}{3 \sqrt[3]{3(1+\sqrt{7})^{2/3}}} \right) \\
 \hline
 2\sqrt{7} \\
 \left( \int -\frac{2 \sqrt[3]{-1+\sqrt{7}} - \sqrt[3]{3^{(x+1)}}}{3^{2/3(x+1)^2-} \sqrt[3]{3(-1+\sqrt{7})^{(x+1)+(-1+\sqrt{7})^{2/3}}}} d(x+1) - \frac{\log\left(\sqrt[3]{3^{(x+1)}} + \sqrt[3]{\sqrt{7}-1}\right)}{3 \sqrt[3]{3(\sqrt{7}-1)^{2/3}}} \right) \\
 \hline
 2\sqrt{7} \\
 \downarrow 25 \\
 \left( \int \frac{\sqrt[3]{3^{(x+1)+2}} \sqrt[3]{1+\sqrt{7}}}{3^{2/3(x+1)^2+} \sqrt[3]{3(1+\sqrt{7})^{(x+1)+(1+\sqrt{7})^{2/3}}}} d(x+1) - \frac{\log\left(\sqrt[3]{1+\sqrt{7}} - \sqrt[3]{3^{(x+1)}}\right)}{3 \sqrt[3]{3(1+\sqrt{7})^{2/3}}} \right) \\
 \hline
 2\sqrt{7} \\
 \left( \int \frac{2 \sqrt[3]{-1+\sqrt{7}} - \sqrt[3]{3^{(x+1)}}}{3^{2/3(x+1)^2-} \sqrt[3]{3(-1+\sqrt{7})^{(x+1)+(-1+\sqrt{7})^{2/3}}}} d(x+1) - \frac{\log\left(\sqrt[3]{3^{(x+1)}} + \sqrt[3]{\sqrt{7}-1}\right)}{3 \sqrt[3]{3(\sqrt{7}-1)^{2/3}}} \right) \\
 \hline
 2\sqrt{7} \\
 \downarrow 1142
 \end{array}$$

$$3 \left( \frac{\frac{3}{2} \sqrt[3]{1 + \sqrt{7}} \int \frac{1}{3^{2/3}(x+1)^2 + \sqrt[3]{3(1 + \sqrt{7})(x+1) + (1 + \sqrt{7})^{2/3}}} dx + \frac{2 \cdot 3^{2/3}(x+1) + \sqrt[3]{3(1 + \sqrt{7})}}{2 \sqrt[3]{3}} \int \frac{d(x+1)}{3^{2/3}(x+1)^2 + \sqrt[3]{3(1 + \sqrt{7})(x+1) + (1 + \sqrt{7})^{2/3}}} \right) \log$$

$$3 \left( \frac{\frac{3}{2} \sqrt[3]{\sqrt{7} - 1} \int \frac{1}{3^{2/3}(x+1)^2 - \sqrt[3]{3(-1 + \sqrt{7})(x+1) + (-1 + \sqrt{7})^{2/3}}} dx + \frac{2\sqrt{7}}{2 \sqrt[3]{3}} \int \frac{\sqrt[3]{3(-1 + \sqrt{7})} - 2 \cdot 3^{2/3}(x+1)}{3^{2/3}(x+1)^2 - \sqrt[3]{3(-1 + \sqrt{7})(x+1) + (-1 + \sqrt{7})^{2/3}}} dx \right)$$

$2\sqrt{7}$

↓ 25

$$3 \left( \frac{\frac{3}{2} \sqrt[3]{1 + \sqrt{7}} \int \frac{1}{3^{2/3(x+1)2 + \sqrt[3]{3(1 + \sqrt{7})(x+1) + (1 + \sqrt{7})^{2/3}}} dx + \frac{2 \cdot 3^{2/3(x+1)} \sqrt[3]{3(1 + \sqrt{7})}}{2 \sqrt[3]{3}} \int \frac{3^{2/3(x+1)} \sqrt[3]{3(1 + \sqrt{7})}}{3^{2/3(x+1)2 + \sqrt[3]{3(1 + \sqrt{7})(x+1) + (1 + \sqrt{7})^{2/3}}} dx}{3(1 + \sqrt{7})^{2/3}} \right) \log$$

$$3 \left( \frac{\frac{3}{2} \sqrt[3]{\sqrt{7} - 1} \int \frac{1}{3^{2/3(x+1)2 - \sqrt[3]{3(-1 + \sqrt{7})(x+1) + (-1 + \sqrt{7})^{2/3}}} dx + \frac{2\sqrt{7}}{3^{2/3(x+1)}} \int \frac{\sqrt[3]{3(-1 + \sqrt{7})} \cdot 3^{2/3(x+1)}}{3^{2/3(x+1)2 - \sqrt[3]{3(-1 + \sqrt{7})(x+1) + (-1 + \sqrt{7})^{2/3}}} dx}{3(\sqrt{7} - 1)^{2/3}} \right)$$

$2\sqrt{7}$

↓ 1082

$$\left( \frac{\int \frac{2 \cdot 3^{2/3(x+1)} \sqrt[3]{3(1+\sqrt{7})}}{3^{2/3(x+1)^2} \sqrt[3]{3(1+\sqrt{7})^{(x+1)}(1+\sqrt{7})^{2/3}}} dx}{2 \sqrt[3]{3}} - 3^{2/3} \int \frac{1}{\left(2 \sqrt[3]{\frac{3}{1+\sqrt{7}}^{(x+1)+1}}\right)^2} d\left(2 \sqrt[3]{\frac{3}{1+\sqrt{7}}^{(x+1)+1}}\right) - \log\left(\sqrt[3]{\frac{3}{1+\sqrt{7}}^{(x+1)+1}}\right)}{3(1+\sqrt{7})^{2/3}} \right)$$

$$\left( \frac{\int \frac{\sqrt[3]{3(-1+\sqrt{7})} \cdot 3^{2/3(x+1)}}{3^{2/3(x+1)^2} \sqrt[3]{3(-1+\sqrt{7})^{(x+1)}(-1+\sqrt{7})^{2/3}}} dx}{2 \sqrt[3]{3}} + 3^{2/3} \int \frac{1}{\left(1-2 \sqrt[3]{\frac{3}{-1+\sqrt{7}}^{(x+1)}}\right)^2} d\left(1-2 \sqrt[3]{\frac{3}{-1+\sqrt{7}}^{(x+1)}}\right)}{3(\sqrt{7}-1)^{2/3}} \right)$$

$2\sqrt{7}$

↓ 217



$$3 \left( \frac{\int \frac{2 \cdot 3^{2/3(x+1)} + \sqrt[3]{3(1+\sqrt{7})}}{3^{2/3(x+1)^2 + \sqrt[3]{3(1+\sqrt{7})(x+1) + (1+\sqrt{7})^{2/3}}} dx}{2 \sqrt[3]{3}} + \sqrt[6]{3} \arctan \left( \frac{\sqrt[2]{3} \sqrt[3]{\frac{3}{1+\sqrt{7}}(x+1)+1}}{\sqrt{3}} \right) - \frac{\log \left( \sqrt[3]{1+\sqrt{7}} - \sqrt[3]{3(x+1)} \right)}{3 \sqrt[3]{3}(1+\sqrt{7})^{2/3}} \right)$$

$2\sqrt{7}$

$$3 \left( \frac{\int \frac{\sqrt[3]{3(-1+\sqrt{7})} - 2 \cdot 3^{2/3(x+1)}}{3^{2/3(x+1)^2 - \sqrt[3]{3(-1+\sqrt{7})(x+1) + (-1+\sqrt{7})^{2/3}}} dx}{2 \sqrt[3]{3}} - \sqrt[6]{3} \arctan \left( \frac{1-2 \sqrt[3]{\frac{3}{\sqrt{7}-1}}(x+1)}}{\sqrt{3}} \right) - \frac{\log \left( \sqrt[3]{3(x+1)} + \sqrt[3]{\sqrt{7}-1} \right)}{3 \sqrt[3]{3}(\sqrt{7}-1)^{2/3}} \right)$$

$2\sqrt{7}$

↓ 1103

$$\frac{3 \left( \frac{\sqrt[6]{3} \arctan \left( \frac{\sqrt[2]{3} \sqrt[3]{\frac{3}{1+\sqrt{7}}(x+1)+1}}{\sqrt{3}} \right) + \frac{\log \left( 3^{2/3}(x+1)^2 + \sqrt[3]{3(1+\sqrt{7})}^{(x+1)+(1+\sqrt{7})^{2/3}} \right)}{2\sqrt[3]{3}}}{3(1+\sqrt{7})^{2/3}} - \frac{\log \left( \sqrt[3]{1+\sqrt{7}} - \sqrt[3]{3(x+1)} \right)}{3\sqrt[3]{3}(1+\sqrt{7})^{2/3}} \right)}{2\sqrt{7}} - \frac{3 \left( -\sqrt[6]{3} \arctan \left( \frac{1-2\sqrt[3]{\frac{3}{\sqrt{7}-1}}(x+1)}}{\sqrt{3}} \right) - \frac{\log \left( 3^{2/3}(x+1)^2 - \sqrt[3]{3(\sqrt{7}-1)}^{(x+1)+(\sqrt{7}-1)^{2/3}} \right)}{2\sqrt[3]{3}}}{3(\sqrt{7}-1)^{2/3}} - \frac{\log \left( \sqrt[3]{3(x+1)} + \sqrt[3]{\sqrt{7}-1} \right)}{3\sqrt[3]{3}(\sqrt{7}-1)^{2/3}} \right)}{2\sqrt{7}}$$

input `Int[(1 - 12*x - 39*x^2 - 58*x^3 - 45*x^4 - 18*x^5 - 3*x^6)^(-1),x]`

output `(-3*(-1/3*Log[(-1 + Sqrt[7])^(1/3) + 3^(1/3)*(1 + x)]/(3^(1/3)*(-1 + Sqrt[7])^(2/3)) - (-3^(1/6)*ArcTan[(1 - 2*(3/(-1 + Sqrt[7]))^(1/3)*(1 + x))/Sqrt[3]]) - Log[(-1 + Sqrt[7])^(2/3) - (3*(-1 + Sqrt[7]))^(1/3)*(1 + x) + 3^(2/3)*(1 + x)^2]/(2*3^(1/3)))/(3*(-1 + Sqrt[7])^(2/3)))/(2*Sqrt[7]) + (3*(-1/3*Log[(1 + Sqrt[7])^(1/3) - 3^(1/3)*(1 + x)]/(3^(1/3)*(1 + Sqrt[7])^(2/3)) + (3^(1/6)*ArcTan[(1 + 2*(3/(1 + Sqrt[7]))^(1/3)*(1 + x))/Sqrt[3]]) + Log[(1 + Sqrt[7])^(2/3) + (3*(1 + Sqrt[7]))^(1/3)*(1 + x) + 3^(2/3)*(1 + x)^2]/(2*3^(1/3)))/(3*(1 + Sqrt[7])^(2/3)))/(2*Sqrt[7])`

## Defintions of rubi rules used

- rule 16  $\text{Int}[(c\_)/((a\_)+(b\_)*(x\_)), x\_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 25  $\text{Int}[-(Fx\_), x\_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[Fx, x], x]$
- rule 217  $\text{Int}[(a\_)+(b\_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(-(\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b] \& \& (\text{LtQ}[a, 0] \parallel \text{LtQ}[b, 0])]$
- rule 750  $\text{Int}[(a\_)+(b\_)*(x_)^3)^{-1}, x\_Symbol] \rightarrow \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[(2*\text{Rt}[a, 3] - \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$
- rule 1082  $\text{Int}[(a\_)+(b\_)*(x_)+(c\_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S \text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \&\& (\text{EqQ}[q^2, 1] \parallel \text{!RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103  $\text{Int}[(d\_)+(e\_)*(x_)/((a\_)+(b\_)*(x_)+(c\_)*(x_)^2), x\_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{EqQ}[2*c*d - b*e, 0]$
- rule 1142  $\text{Int}[(d\_)+(e\_)*(x_)/((a\_)+(b\_)*(x_)+(c\_)*(x_)^2), x\_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \text{ Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \text{ Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$
- rule 1685  $\text{Int}[(a\_)+(b\_)*(x_)^{(n\_)}+(c\_)*(x_)^{(n2\_)}^{-1}, x\_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b^2 - 4*a*c, 2]\}, \text{Simp}[c/q \text{ Int}[1/(b/2 - q/2 + c*x^n), x], x] - \text{Simp}[c/q \text{ Int}[1/(b/2 + q/2 + c*x^n), x], x]] /; \text{FreeQ}[\{a, b, c\}, x] \&\& \text{EqQ}[n2, 2*n] \&\& \text{NeQ}[b^2 - 4*a*c, 0]$

rule 2458

```
Int[(Pn_)^(p_), x_Symbol] := With[{S = Coeff[Pn, x, Expon[Pn, x] - 1]/(Exp
on[Pn, x]*Coeff[Pn, x, Expon[Pn, x]])}, Subst[Int[ExpandToSum[Pn /. x -> x
- S, x]^p, x], x, x + S] /; BinomialQ[Pn /. x -> x - S, x] || (IntegerQ[Exp
on[Pn, x]/2] && TrinomialQ[Pn /. x -> x - S, x])] /; FreeQ[p, x] && PolyQ[P
n, x] && GtQ[Expon[Pn, x], 2] && NeQ[Coeff[Pn, x, Expon[Pn, x] - 1], 0]
```

### Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.04 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.21

method	result	size
default	$-\frac{\left( \sum_{R=\text{RootOf}(3Z^6+18Z^5+45Z^4+58Z^3+39Z^2+12Z-1)} \frac{\ln(x-R)}{3R^5+15R^4+30R^3+29R^2+13R+2} \right)}{6}$	71
risch	$-\frac{\left( \sum_{R=\text{RootOf}(3Z^6+18Z^5+45Z^4+58Z^3+39Z^2+12Z-1)} \frac{\ln(x-R)}{3R^5+15R^4+30R^3+29R^2+13R+2} \right)}{6}$	71

input

```
int(1/(-3*x^6-18*x^5-45*x^4-58*x^3-39*x^2-12*x+1),x,method=_RETURNVERBOSE)
```

output

```
-1/6*sum(1/(3*_R^5+15*_R^4+30*_R^3+29*_R^2+13*_R+2)*ln(x-_R),_R=RootOf(3*_
Z^6+18*_Z^5+45*_Z^4+58*_Z^3+39*_Z^2+12*_Z-1))
```



output

```
-1/12*(2/49*sqrt(7) + 1/14)^(1/3)*(sqrt(-3) + 1)*log((sqrt(7)*(sqrt(-3) + 1) - 7*sqrt(-3) - 7)*(2/49*sqrt(7) + 1/14)^(1/3) + 6*x + 6) + 1/12*(2/49*sqrt(7) + 1/14)^(1/3)*(sqrt(-3) - 1)*log(-(sqrt(7)*(sqrt(-3) - 1) - 7*sqrt(-3) + 7)*(2/49*sqrt(7) + 1/14)^(1/3) + 6*x + 6) - 1/12*(-2/49*sqrt(7) + 1/14)^(1/3)*(sqrt(-3) + 1)*log(-(sqrt(7)*(sqrt(-3) + 1) + 7*sqrt(-3) + 7)*(-2/49*sqrt(7) + 1/14)^(1/3) + 6*x + 6) + 1/12*(-2/49*sqrt(7) + 1/14)^(1/3)*(sqrt(-3) - 1)*log((sqrt(7)*(sqrt(-3) - 1) + 7*sqrt(-3) - 7)*(-2/49*sqrt(7) + 1/14)^(1/3) + 6*x + 6) + 1/6*(2/49*sqrt(7) + 1/14)^(1/3)*log(-(sqrt(7) - 7)*(2/49*sqrt(7) + 1/14)^(1/3) + 3*x + 3) + 1/6*(-2/49*sqrt(7) + 1/14)^(1/3)*log((sqrt(7) + 7)*(-2/49*sqrt(7) + 1/14)^(1/3) + 3*x + 3)
```

**Sympy [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.09

$$\int \frac{1}{1 - 12x - 39x^2 - 58x^3 - 45x^4 - 18x^5 - 3x^6} dx$$

$$= -\text{RootSum}\left(7112448t^6 + 4704t^3 - 1, \left(t \mapsto t \log\left(-10584t^4 - \frac{35t}{2} + x + 1\right)\right)\right)$$

input

```
integrate(1/(-3*x**6-18*x**5-45*x**4-58*x**3-39*x**2-12*x+1),x)
```

output

```
-RootSum(7112448*_t**6 + 4704*_t**3 - 1, Lambda(_t, _t*log(-10584*_t**4 - 35*_t/2 + x + 1)))
```

**Maxima [F]**

$$\int \frac{1}{1 - 12x - 39x^2 - 58x^3 - 45x^4 - 18x^5 - 3x^6} dx$$

$$= \int -\frac{1}{3x^6 + 18x^5 + 45x^4 + 58x^3 + 39x^2 + 12x - 1} dx$$

input

```
integrate(1/(-3*x^6-18*x^5-45*x^4-58*x^3-39*x^2-12*x+1),x, algorithm="maxima")
```

output `-integrate(1/(3*x^6 + 18*x^5 + 45*x^4 + 58*x^3 + 39*x^2 + 12*x - 1), x)`

**Giac** [F]

$$\int \frac{1}{1 - 12x - 39x^2 - 58x^3 - 45x^4 - 18x^5 - 3x^6} dx$$
$$= \int -\frac{1}{3x^6 + 18x^5 + 45x^4 + 58x^3 + 39x^2 + 12x - 1} dx$$

input `integrate(1/(-3*x^6-18*x^5-45*x^4-58*x^3-39*x^2-12*x+1),x, algorithm="giac")`

output `integrate(-1/(3*x^6 + 18*x^5 + 45*x^4 + 58*x^3 + 39*x^2 + 12*x - 1), x)`

**Mupad [B] (verification not implemented)**

Time = 23.07 (sec) , antiderivative size = 463, normalized size of antiderivative = 1.37

$$\begin{aligned}
& \int \frac{1}{1 - 12x - 39x^2 - 58x^3 - 45x^4 - 18x^5 - 3x^6} dx \\
& \ln \left( x + \frac{2^{2/3} 7^{1/3} (7-4\sqrt{7})^{1/3}}{6} + \frac{2^{2/3} 7^{5/6} (7-4\sqrt{7})^{1/3}}{42} + 1 \right) (196 - 112\sqrt{7})^{1/3} \\
= & \frac{\ln \left( x - \frac{2^{2/3} 7^{1/3} (4\sqrt{7}-7)^{1/3}}{6} - \frac{2^{2/3} 7^{5/6} (4\sqrt{7}-7)^{1/3}}{42} + 1 \right) (112\sqrt{7} - 196)^{1/3}}{84} \\
& - \frac{\ln \left( x + \frac{2^{2/3} 7^{1/3} (4\sqrt{7}+7)^{1/3}}{6} - \frac{2^{2/3} 7^{5/6} (4\sqrt{7}+7)^{1/3}}{42} + 1 \right) (112\sqrt{7} + 196)^{1/3}}{84} \\
& + \frac{2^{2/3} 7^{1/3} \ln \left( x - \frac{2^{2/3} 7^{1/3} (7-4\sqrt{7})^{1/3}}{12} - \frac{2^{2/3} 7^{5/6} (7-4\sqrt{7})^{1/3}}{84} + 1 - \frac{2^{2/3} \sqrt{3} 7^{1/3} (7-4\sqrt{7})^{1/3} \operatorname{li}}{12} - \frac{2^{2/3} \sqrt{3} 7^{5/6} (7-4\sqrt{7})^{1/3}}{84} \right)}{84} \\
& - \frac{2^{2/3} 7^{1/3} \ln \left( x - \frac{2^{2/3} 7^{1/3} (4\sqrt{7}+7)^{1/3}}{12} + \frac{2^{2/3} 7^{5/6} (4\sqrt{7}+7)^{1/3}}{84} + 1 - \frac{2^{2/3} \sqrt{3} 7^{1/3} (4\sqrt{7}+7)^{1/3} \operatorname{li}}{12} + \frac{2^{2/3} \sqrt{3} 7^{5/6} (4\sqrt{7}+7)^{1/3}}{84} \right)}{168} \\
& - \frac{2^{2/3} 7^{1/3} \ln \left( x - \frac{2^{2/3} 7^{1/3} (4\sqrt{7}+7)^{1/3}}{12} + \frac{2^{2/3} 7^{5/6} (4\sqrt{7}+7)^{1/3}}{84} + 1 + \frac{2^{2/3} \sqrt{3} 7^{1/3} (4\sqrt{7}+7)^{1/3} \operatorname{li}}{12} - \frac{2^{2/3} \sqrt{3} 7^{5/6} (4\sqrt{7}+7)^{1/3}}{84} \right)}{168} \\
& + \frac{2^{2/3} 7^{1/3} \ln \left( x - \frac{2^{2/3} 7^{1/3} (7-4\sqrt{7})^{1/3}}{12} - \frac{2^{2/3} 7^{5/6} (7-4\sqrt{7})^{1/3}}{84} + 1 - \frac{2^{2/3} \sqrt{3} 7^{1/3} (7-4\sqrt{7})^{1/3} \operatorname{li}}{12} - \frac{2^{2/3} \sqrt{3} 7^{5/6} (7-4\sqrt{7})^{1/3}}{84} \right)}{168}
\end{aligned}$$

input `int(-1/(12*x + 39*x^2 + 58*x^3 + 45*x^4 + 18*x^5 + 3*x^6 - 1),x)`



output

```
(log(x + (2^(2/3)*7^(1/3)*(7 - 4*7^(1/2))^(1/3))/6 + (2^(2/3)*7^(5/6)*(7 -
4*7^(1/2))^(1/3))/42 + 1)*(196 - 112*7^(1/2))^(1/3))/84 - (log(x - (2^(2/
3)*7^(1/3)*(4*7^(1/2) - 7)^(1/3))/6 - (2^(2/3)*7^(5/6)*(4*7^(1/2) - 7)^(1/
3))/42 + 1)*(112*7^(1/2) - 196)^(1/3))/84 + (log(x + (2^(2/3)*7^(1/3)*(4*7
^(1/2) + 7)^(1/3))/6 - (2^(2/3)*7^(5/6)*(4*7^(1/2) + 7)^(1/3))/42 + 1)*(11
2*7^(1/2) + 196)^(1/3))/84 - (2^(2/3)*7^(1/3)*log(x - (2^(2/3)*7^(1/3)*(7
- 4*7^(1/2))^(1/3))/12 - (2^(2/3)*7^(5/6)*(7 - 4*7^(1/2))^(1/3))/84 - (2^(
2/3)*3^(1/2)*7^(1/3)*(7 - 4*7^(1/2))^(1/3)*1i)/12 - (2^(2/3)*3^(1/2)*7^(5/
6)*(7 - 4*7^(1/2))^(1/3)*1i)/84 + 1)*(3^(1/2)*1i + 1)*(7 - 4*7^(1/2))^(1/3
))/168 - (2^(2/3)*7^(1/3)*log(x - (2^(2/3)*7^(1/3)*(4*7^(1/2) + 7)^(1/3))/
12 + (2^(2/3)*7^(5/6)*(4*7^(1/2) + 7)^(1/3))/84 - (2^(2/3)*3^(1/2)*7^(1/3)
*(4*7^(1/2) + 7)^(1/3)*1i)/12 + (2^(2/3)*3^(1/2)*7^(5/6)*(4*7^(1/2) + 7)^(
1/3)*1i)/84 + 1)*(3^(1/2)*1i + 1)*(4*7^(1/2) + 7)^(1/3))/168 + (2^(2/3)*7
^(1/3)*log(x - (2^(2/3)*7^(1/3)*(4*7^(1/2) + 7)^(1/3))/12 + (2^(2/3)*7^(5/6
)*(4*7^(1/2) + 7)^(1/3))/84 + (2^(2/3)*3^(1/2)*7^(1/3)*(4*7^(1/2) + 7)^(1/
3)*1i)/12 - (2^(2/3)*3^(1/2)*7^(5/6)*(4*7^(1/2) + 7)^(1/3)*1i)/84 + 1)*(3^
(1/2)*1i - 1)*(4*7^(1/2) + 7)^(1/3))/168
```

**Reduce [F]**

$$\int \frac{1}{1 - 12x - 39x^2 - 58x^3 - 45x^4 - 18x^5 - 3x^6} dx$$

$$= - \left( \int \frac{1}{3x^6 + 18x^5 + 45x^4 + 58x^3 + 39x^2 + 12x - 1} dx \right)$$

input

```
int(1/(-3*x^6-18*x^5-45*x^4-58*x^3-39*x^2-12*x+1),x)
```

output

```
- int(1/(3*x**6 + 18*x**5 + 45*x**4 + 58*x**3 + 39*x**2 + 12*x - 1),x)
```

### 3.30 $\int \frac{1}{(a+bx^2)^4} dx$

Optimal result	241
Mathematica [A] (verified)	241
Rubi [A] (verified)	242
Maple [A] (verified)	243
Fricas [A] (verification not implemented)	244
Sympy [A] (verification not implemented)	244
Maxima [A] (verification not implemented)	245
Giac [A] (verification not implemented)	245
Mupad [B] (verification not implemented)	245
Reduce [B] (verification not implemented)	246

#### Optimal result

Integrand size = 9, antiderivative size = 79

$$\int \frac{1}{(a+bx^2)^4} dx = \frac{x}{6a(a+bx^2)^3} + \frac{5x}{24a^2(a+bx^2)^2} + \frac{5x}{16a^3(a+bx^2)} + \frac{5 \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{16a^{7/2}\sqrt{b}}$$

output

```
1/6*x/a/(b*x^2+a)^3+5/24*x/a^2/(b*x^2+a)^2+5/16*x/a^3/(b*x^2+a)+5/16*arctan(b^(1/2)*x/a^(1/2))/a^(7/2)/b^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.84

$$\int \frac{1}{(a+bx^2)^4} dx = \frac{33a^2x + 40abx^3 + 15b^2x^5}{48a^3(a+bx^2)^3} + \frac{5 \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{16a^{7/2}\sqrt{b}}$$

input

```
Integrate[(a + b*x^2)^(-4), x]
```

output

```
(33*a^2*x + 40*a*b*x^3 + 15*b^2*x^5)/(48*a^3*(a + b*x^2)^3) + (5*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(16*a^(7/2)*Sqrt[b])
```

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.20, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$ , Rules used = {215, 215, 215, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a + bx^2)^4} dx \\
 & \quad \downarrow \text{215} \\
 & \frac{5 \int \frac{1}{(bx^2+a)^3} dx}{6a} + \frac{x}{6a(a + bx^2)^3} \\
 & \quad \downarrow \text{215} \\
 & \frac{5 \left( \frac{3 \int \frac{1}{(bx^2+a)^2} dx}{4a} + \frac{x}{4a(a+bx^2)^2} \right)}{6a} + \frac{x}{6a(a + bx^2)^3} \\
 & \quad \downarrow \text{215} \\
 & \frac{5 \left( \frac{3 \left( \frac{\int \frac{1}{bx^2+a} dx}{2a} + \frac{x}{2a(a+bx^2)} \right)}{4a} + \frac{x}{4a(a+bx^2)^2} \right)}{6a} + \frac{x}{6a(a + bx^2)^3} \\
 & \quad \downarrow \text{218} \\
 & \frac{5 \left( \frac{3 \left( \frac{\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{2a^{3/2}\sqrt{b}} + \frac{x}{2a(a+bx^2)} \right)}{4a} + \frac{x}{4a(a+bx^2)^2} \right)}{6a} + \frac{x}{6a(a + bx^2)^3}
 \end{aligned}$$

input `Int[(a + b*x^2)^(-4), x]`

output 
$$\frac{x}{6a(a + bx^2)^3} + \frac{5x}{4a(a + bx^2)^2} + \frac{3x}{2a(a + bx^2)} + \frac{\text{ArcTan}\left[\frac{\sqrt{b}x}{\sqrt{a}}\right]}{(2a^{3/2}\sqrt{b})} \Big/ (4a) \Big/ (6a)$$

### Defintions of rubi rules used

rule 215 
$$\text{Int}[(a_ + (b_)(x_)^2)^{(p_)}, x\_Symbol] \rightarrow \text{Simp}[(-x)((a + bx^2)^{(p + 1)} / (2a^{p + 1}))], x] + \text{Simp}[(2p + 3)/(2a^{p + 1}) \text{Int}[(a + bx^2)^{(p + 1)}, x], x] /; \text{FreeQ}\{a, b\}, x\} \ \&\& \ \text{LtQ}\{p, -1\} \ \&\& \ (\text{IntegerQ}\{4p\} \ || \ \text{IntegerQ}\{6p\})$$

rule 218 
$$\text{Int}[(a_ + (b_)(x_)^2)^{(-1)}, x\_Symbol] \rightarrow \text{Simp}[(\text{Rt}\{a/b, 2\}/a) \text{ArcTan}\{x/\text{Rt}\{a/b, 2\}\}], x] /; \text{FreeQ}\{a, b\}, x\} \ \&\& \ \text{PosQ}\{a/b\}$$

### Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.99

method	result	size
default	$\frac{x}{6a(bx^2+a)^3} + \frac{\frac{5x}{24a(bx^2+a)^2} + \frac{5\left(\frac{3x}{8a(bx^2+a)} + \frac{3\arctan\left(\frac{bx}{\sqrt{ab}}\right)}{8a\sqrt{ab}}\right)}{6a}}{a}$	78
risch	$\frac{\frac{5b^2x^5}{16a^3} + \frac{5bx^3}{6a^2} + \frac{11x}{16a}}{(bx^2+a)^3} - \frac{5\ln(bx+\sqrt{-ab})}{32\sqrt{-ab}a^3} + \frac{5\ln(-bx+\sqrt{-ab})}{32\sqrt{-ab}a^3}$	84

input `int(1/(b*x^2+a)^4,x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{6}x/a/(bx^2+a)^3 + \frac{5}{6}x/a/(bx^2+a)^2 + \frac{3}{4}x/a/(bx^2+a) + \frac{1}{2}a/(ab)^{(1/2)} \text{arctan}(bx/(ab)^{(1/2)})$$

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 254, normalized size of antiderivative = 3.22

$$\int \frac{1}{(a + bx^2)^4} dx$$

$$= \left[ \frac{30 ab^3 x^5 + 80 a^2 b^2 x^3 + 66 a^3 b x - 15 (b^3 x^6 + 3 ab^2 x^4 + 3 a^2 b x^2 + a^3) \sqrt{-ab} \log\left(\frac{bx^2 - 2\sqrt{-ab}x - a}{bx^2 + a}\right) + 15 ab^3 x}{96 (a^4 b^4 x^6 + 3 a^5 b^3 x^4 + 3 a^6 b^2 x^2 + a^7 b)} \right],$$

input `integrate(1/(b*x^2+a)^4,x, algorithm="fricas")`output `[1/96*(30*a*b^3*x^5 + 80*a^2*b^2*x^3 + 66*a^3*b*x - 15*(b^3*x^6 + 3*a*b^2*x^4 + 3*a^2*b*x^2 + a^3)*sqrt(-a*b)*log((b*x^2 - 2*sqrt(-a*b)*x - a)/(b*x^2 + a)))/(a^4*b^4*x^6 + 3*a^5*b^3*x^4 + 3*a^6*b^2*x^2 + a^7*b), 1/48*(15*a*b^3*x^5 + 40*a^2*b^2*x^3 + 33*a^3*b*x + 15*(b^3*x^6 + 3*a*b^2*x^4 + 3*a^2*b*x^2 + a^3)*sqrt(a*b)*arctan(sqrt(a*b)*x/a))/(a^4*b^4*x^6 + 3*a^5*b^3*x^4 + 3*a^6*b^2*x^2 + a^7*b)]`**Sympy [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.63

$$\int \frac{1}{(a + bx^2)^4} dx = -\frac{5\sqrt{-\frac{1}{a^7b}} \log\left(-a^4\sqrt{-\frac{1}{a^7b}} + x\right)}{32} + \frac{5\sqrt{-\frac{1}{a^7b}} \log\left(a^4\sqrt{-\frac{1}{a^7b}} + x\right)}{32}$$

$$+ \frac{33a^2x + 40abx^3 + 15b^2x^5}{48a^6 + 144a^5bx^2 + 144a^4b^2x^4 + 48a^3b^3x^6}$$

input `integrate(1/(b*x**2+a)**4,x)`output `-5*sqrt(-1/(a**7*b))*log(-a**4*sqrt(-1/(a**7*b)) + x)/32 + 5*sqrt(-1/(a**7*b))*log(a**4*sqrt(-1/(a**7*b)) + x)/32 + (33*a**2*x + 40*a*b*x**3 + 15*b**2*x**5)/(48*a**6 + 144*a**5*b*x**2 + 144*a**4*b**2*x**4 + 48*a**3*b**3*x**6)`

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.01

$$\int \frac{1}{(a + bx^2)^4} dx = \frac{15b^2x^5 + 40abx^3 + 33a^2x}{48(a^3b^3x^6 + 3a^4b^2x^4 + 3a^5bx^2 + a^6)} + \frac{5 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{16\sqrt{aba^3}}$$

input `integrate(1/(b*x^2+a)^4,x, algorithm="maxima")`output `1/48*(15*b^2*x^5 + 40*a*b*x^3 + 33*a^2*x)/(a^3*b^3*x^6 + 3*a^4*b^2*x^4 + 3*a^5*b*x^2 + a^6) + 5/16*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*a^3)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.71

$$\int \frac{1}{(a + bx^2)^4} dx = \frac{5 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{16\sqrt{aba^3}} + \frac{15b^2x^5 + 40abx^3 + 33a^2x}{48(bx^2 + a)^3a^3}$$

input `integrate(1/(b*x^2+a)^4,x, algorithm="giac")`output `5/16*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*a^3) + 1/48*(15*b^2*x^5 + 40*a*b*x^3 + 33*a^2*x)/((b*x^2 + a)^3*a^3)`**Mupad [B] (verification not implemented)**

Time = 22.00 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.97

$$\int \frac{1}{(a + bx^2)^4} dx = \frac{\frac{11x}{16a} + \frac{5bx^3}{6a^2} + \frac{5b^2x^5}{16a^3}}{a^3 + 3a^2bx^2 + 3ab^2x^4 + b^3x^6} + \frac{5 \operatorname{atan}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{16a^{7/2}\sqrt{b}}$$

input `int(1/(a + b*x^2)^4,x)`

output

```
((11*x)/(16*a) + (5*b*x^3)/(6*a^2) + (5*b^2*x^5)/(16*a^3))/(a^3 + b^3*x^6
+ 3*a^2*b*x^2 + 3*a*b^2*x^4) + (5*atan((b^(1/2)*x)/a^(1/2)))/(16*a^(7/2)*b
^(1/2))
```

**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 162, normalized size of antiderivative = 2.05

$$\int \frac{1}{(a + bx^2)^4} dx$$

$$= \frac{15\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right) a^3 + 45\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right) a^2 b x^2 + 45\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right) a b^2 x^4 + 15\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right) a^3}{48a^4 b (b^3 x^6 + 3a b^2 x^4 + 3a^2 b x^2 + a^3)}$$

input

```
int(1/(b*x^2+a)^4,x)
```

output

```
(15*sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*sqrt(a)))*a**3 + 45*sqrt(b)*sqrt(a)
)*atan((b*x)/(sqrt(b)*sqrt(a)))*a**2*b*x**2 + 45*sqrt(b)*sqrt(a)*atan((b*x
)/(sqrt(b)*sqrt(a)))*a*b**2*x**4 + 15*sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*
sqrt(a)))*b**3*x**6 + 33*a**3*b*x + 40*a**2*b**2*x**3 + 15*a*b**3*x**5)/(4
8*a**4*b*(a**3 + 3*a**2*b*x**2 + 3*a*b**2*x**4 + b**3*x**6))
```

### 3.31 $\int \frac{1}{(a^2 + 2abx^2 + b^2x^4)^2} dx$

Optimal result	247
Mathematica [A] (verified)	247
Rubi [A] (verified)	248
Maple [A] (verified)	250
Fricas [A] (verification not implemented)	250
Sympy [A] (verification not implemented)	251
Maxima [A] (verification not implemented)	251
Giac [A] (verification not implemented)	252
Mupad [B] (verification not implemented)	252
Reduce [B] (verification not implemented)	252

#### Optimal result

Integrand size = 20, antiderivative size = 79

$$\int \frac{1}{(a^2 + 2abx^2 + b^2x^4)^2} dx = \frac{x}{6a(a + bx^2)^3} + \frac{5x}{24a^2(a + bx^2)^2} + \frac{5x}{16a^3(a + bx^2)} + \frac{5 \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{16a^{7/2}\sqrt{b}}$$

output

```
1/6*x/a/(b*x^2+a)^3+5/24*x/a^2/(b*x^2+a)^2+5/16*x/a^3/(b*x^2+a)+5/16*arctan(b^(1/2)*x/a^(1/2))/a^(7/2)/b^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.84

$$\int \frac{1}{(a^2 + 2abx^2 + b^2x^4)^2} dx = \frac{33a^2x + 40abx^3 + 15b^2x^5}{48a^3(a + bx^2)^3} + \frac{5 \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{16a^{7/2}\sqrt{b}}$$

input

```
Integrate[(a^2 + 2*a*b*x^2 + b^2*x^4)^(-2), x]
```



output

$$(33a^2x + 40abx^3 + 15b^2x^5)/(48a^3(a + bx^2)^3) + (5\text{ArcTan}[(\text{Sqrt}[b]x)/\text{Sqrt}[a]])/(16a^{7/2}\text{Sqrt}[b])$$
**Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.44, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {1379, 215, 215, 215, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a^2 + 2abx^2 + b^2x^4)^2} dx$$

$$\downarrow 1379$$

$$b^4 \int \frac{1}{(b^2x^2 + ab)^4} dx$$

$$\downarrow 215$$

$$b^4 \left( \frac{5 \int \frac{1}{(b^2x^2 + ab)^3} dx}{6ab} + \frac{x}{6ab^4(a + bx^2)^3} \right)$$

$$\downarrow 215$$

$$b^4 \left( \frac{5 \left( \frac{3 \int \frac{1}{(b^2x^2 + ab)^2} dx}{4ab} + \frac{x}{4ab^3(a + bx^2)^2} \right)}{6ab} + \frac{x}{6ab^4(a + bx^2)^3} \right)$$

$$\downarrow 215$$

$$b^4 \left( \frac{5 \left( \frac{3 \left( \frac{\int \frac{1}{b^2x^2 + ab} dx}{2ab} + \frac{x}{2ab^2(a + bx^2)} \right)}{4ab} + \frac{x}{4ab^3(a + bx^2)^2} \right)}{6ab} + \frac{x}{6ab^4(a + bx^2)^3} \right)$$

$$\begin{array}{c}
 \downarrow 218 \\
 b^4 \left( \frac{5 \left( \frac{3 \left( \frac{\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{2a^{3/2}b^{5/2}} + \frac{x}{2ab^2(a+bx^2)} \right)}{4ab} + \frac{x}{4ab^3(a+bx^2)^2} \right)}{6ab} + \frac{x}{6ab^4(a+bx^2)^3} \right)
 \end{array}$$

input `Int[(a^2 + 2*a*b*x^2 + b^2*x^4)^(-2), x]`

output `b^4*(x/(6*a*b^4*(a + b*x^2)^3) + (5*(x/(4*a*b^3*(a + b*x^2)^2) + (3*(x/(2*a*b^2*(a + b*x^2)) + ArcTan[(Sqrt[b]*x)/Sqrt[a]]/(2*a^(3/2)*b^(5/2)))))/(4*a*b))/(6*a*b))`

### Defintions of rubi rules used

rule 215 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && LtQ[p, -1] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 1379 `Int[((a_) + (c_.)*(x_)^(n2_.) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/c^p Int[(b/2 + c*x^n)^(2*p), x], x] /; FreeQ[{a, b, c, n, p}, x] && EqQ[n2, 2*n] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p] && NeQ[p, 1]`

**Maple [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.99

method	result	size
default	$\frac{x}{6a(bx^2+a)^3} + \frac{\frac{5x}{24a(bx^2+a)^2} + \frac{5\left(\frac{3x}{8a(bx^2+a)} + \frac{3\arctan\left(\frac{bx}{\sqrt{ab}}\right)}{8a\sqrt{ab}}\right)}{6a}}{a}$	78
risch	$\frac{\frac{5b^2x^5}{16a^3} + \frac{5bx^3}{6a^2} + \frac{11x}{16a}}{(b^2x^4+2abx^2+a^2)(bx^2+a)} - \frac{5\ln(bx+\sqrt{-ab})}{32\sqrt{-ab}a^3} + \frac{5\ln(-bx+\sqrt{-ab})}{32\sqrt{-ab}a^3}$	104

input `int(1/(b^2*x^4+2*a*b*x^2+a^2)^2,x,method=_RETURNVERBOSE)`

output  $\frac{1}{6}x/a/(bx^2+a)^3 + 5/6/a*(1/4*x/a/(bx^2+a)^2 + 3/4/a*(1/2*x/a/(bx^2+a) + 1/2/a/(a*b)^{(1/2)}*\arctan(b*x/(a*b)^{(1/2)}))$

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 254, normalized size of antiderivative = 3.22

$$\int \frac{1}{(a^2 + 2abx^2 + b^2x^4)^2} dx$$

$$= \left[ \frac{30ab^3x^5 + 80a^2b^2x^3 + 66a^3bx - 15(b^3x^6 + 3ab^2x^4 + 3a^2bx^2 + a^3)\sqrt{-ab} \log\left(\frac{bx^2 - 2\sqrt{-ab}x - a}{bx^2 + a}\right)}{96(a^4b^4x^6 + 3a^5b^3x^4 + 3a^6b^2x^2 + a^7b)}, \frac{15ab^3x^5}{96(a^4b^4x^6 + 3a^5b^3x^4 + 3a^6b^2x^2 + a^7b)} \right]$$

input `integrate(1/(b^2*x^4+2*a*b*x^2+a^2)^2,x, algorithm="fricas")`

output  $[1/96*(30*a*b^3*x^5 + 80*a^2*b^2*x^3 + 66*a^3*b*x - 15*(b^3*x^6 + 3*a*b^2*x^4 + 3*a^2*b*x^2 + a^3)*\sqrt{-a*b}*\log((b*x^2 - 2*\sqrt{-a*b}*x - a)/(b*x^2 + a)))/(a^4*b^4*x^6 + 3*a^5*b^3*x^4 + 3*a^6*b^2*x^2 + a^7*b), 1/48*(15*a*b^3*x^5 + 40*a^2*b^2*x^3 + 33*a^3*b*x + 15*(b^3*x^6 + 3*a*b^2*x^4 + 3*a^2*b*x^2 + a^3)*\sqrt{a*b}*\arctan(\sqrt{a*b}*x/a))/(a^4*b^4*x^6 + 3*a^5*b^3*x^4 + 3*a^6*b^2*x^2 + a^7*b)]$

**Sympy [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.63

$$\int \frac{1}{(a^2 + 2abx^2 + b^2x^4)^2} dx = -\frac{5\sqrt{-\frac{1}{a^7b}} \log\left(-a^4\sqrt{-\frac{1}{a^7b}} + x\right)}{32} + \frac{5\sqrt{-\frac{1}{a^7b}} \log\left(a^4\sqrt{-\frac{1}{a^7b}} + x\right)}{32} + \frac{33a^2x + 40abx^3 + 15b^2x^5}{48a^6 + 144a^5bx^2 + 144a^4b^2x^4 + 48a^3b^3x^6}$$

input `integrate(1/(b**2*x**4+2*a*b*x**2+a**2)**2,x)`output `-5*sqrt(-1/(a**7*b))*log(-a**4*sqrt(-1/(a**7*b)) + x)/32 + 5*sqrt(-1/(a**7*b))*log(a**4*sqrt(-1/(a**7*b)) + x)/32 + (33*a**2*x + 40*a*b*x**3 + 15*b**2*x**5)/(48*a**6 + 144*a**5*b*x**2 + 144*a**4*b**2*x**4 + 48*a**3*b**3*x**6)`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.01

$$\int \frac{1}{(a^2 + 2abx^2 + b^2x^4)^2} dx = \frac{15b^2x^5 + 40abx^3 + 33a^2x}{48(a^3b^3x^6 + 3a^4b^2x^4 + 3a^5bx^2 + a^6)} + \frac{5 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{16\sqrt{aba^3}}$$

input `integrate(1/(b^2*x^4+2*a*b*x^2+a^2)^2,x, algorithm="maxima")`output `1/48*(15*b^2*x^5 + 40*a*b*x^3 + 33*a^2*x)/(a^3*b^3*x^6 + 3*a^4*b^2*x^4 + 3*a^5*b*x^2 + a^6) + 5/16*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*a^3)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.71

$$\int \frac{1}{(a^2 + 2abx^2 + b^2x^4)^2} dx = \frac{5 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{16 \sqrt{aba^3}} + \frac{15 b^2 x^5 + 40 abx^3 + 33 a^2 x}{48 (bx^2 + a)^3 a^3}$$

input `integrate(1/(b^2*x^4+2*a*b*x^2+a^2)^2,x, algorithm="giac")`output `5/16*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*a^3) + 1/48*(15*b^2*x^5 + 40*a*b*x^3 + 33*a^2*x)/((b*x^2 + a)^3*a^3)`**Mupad [B] (verification not implemented)**

Time = 21.88 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.97

$$\int \frac{1}{(a^2 + 2abx^2 + b^2x^4)^2} dx = \frac{\frac{11x}{16a} + \frac{5bx^3}{6a^2} + \frac{5b^2x^5}{16a^3}}{a^3 + 3a^2bx^2 + 3ab^2x^4 + b^3x^6} + \frac{5 \operatorname{atan}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{16 a^{7/2} \sqrt{b}}$$

input `int(1/(a^2 + b^2*x^4 + 2*a*b*x^2)^2,x)`output `((11*x)/(16*a) + (5*b*x^3)/(6*a^2) + (5*b^2*x^5)/(16*a^3))/(a^3 + b^3*x^6 + 3*a^2*b*x^2 + 3*a*b^2*x^4) + (5*atan((b^(1/2)*x)/a^(1/2)))/(16*a^(7/2)*b^(1/2))`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 162, normalized size of antiderivative = 2.05

$$\int \frac{1}{(a^2 + 2abx^2 + b^2x^4)^2} dx = \frac{15\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right) a^3 + 45\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right) a^2 b x^2 + 45\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right) a b^2 x^4 + 15\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right) a^3}{48a^4b(b^3x^6 + 3ab^2x^4 + 3a^2bx^2 + a^3)}$$

input `int(1/(b^2*x^4+2*a*b*x^2+a^2)^2,x)`

output `(15*sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*sqrt(a)))*a**3 + 45*sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*sqrt(a)))*a**2*b*x**2 + 45*sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*sqrt(a)))*a*b**2*x**4 + 15*sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*sqrt(a)))*b**3*x**6 + 33*a**3*b*x + 40*a**2*b**2*x**3 + 15*a*b**3*x**5)/(48*a**4*b*(a**3 + 3*a**2*b*x**2 + 3*a*b**2*x**4 + b**3*x**6))`

**3.32**  $\int \frac{1}{a^4 + 4a^3bx^2 + 6a^2b^2x^4 + 4ab^3x^6 + b^4x^8} dx$

Optimal result . . . . .	254
Mathematica [A] (verified) . . . . .	254
Rubi [A] (verified) . . . . .	255
Maple [A] (verified) . . . . .	257
Fricas [A] (verification not implemented) . . . . .	257
Sympy [A] (verification not implemented) . . . . .	258
Maxima [A] (verification not implemented) . . . . .	258
Giac [A] (verification not implemented) . . . . .	259
Mupad [B] (verification not implemented) . . . . .	259
Reduce [B] (verification not implemented) . . . . .	260

**Optimal result**

Integrand size = 42, antiderivative size = 79

$$\int \frac{1}{a^4 + 4a^3bx^2 + 6a^2b^2x^4 + 4ab^3x^6 + b^4x^8} dx$$

$$= \frac{x}{6a(a + bx^2)^3} + \frac{5x}{24a^2(a + bx^2)^2} + \frac{5x}{16a^3(a + bx^2)} + \frac{5 \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{16a^{7/2}\sqrt{b}}$$

output

```
1/6*x/a/(b*x^2+a)^3+5/24*x/a^2/(b*x^2+a)^2+5/16*x/a^3/(b*x^2+a)+5/16*arctan(b^(1/2)*x/a^(1/2))/a^(7/2)/b^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.84

$$\int \frac{1}{a^4 + 4a^3bx^2 + 6a^2b^2x^4 + 4ab^3x^6 + b^4x^8} dx$$

$$= \frac{33a^2x + 40abx^3 + 15b^2x^5}{48a^3(a + bx^2)^3} + \frac{5 \arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{16a^{7/2}\sqrt{b}}$$

input

```
Integrate[(a^4 + 4*a^3*b*x^2 + 6*a^2*b^2*x^4 + 4*a*b^3*x^6 + b^4*x^8)^(-1),x]
```

output

$$(33*a^2*x + 40*a*b*x^3 + 15*b^2*x^5)/(48*a^3*(a + b*x^2)^3) + (5*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(16*a^{(7/2)}*Sqrt[b])$$
**Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.20, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.119$ , Rules used = {2070, 215, 215, 215, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{a^4 + 4a^3bx^2 + 6a^2b^2x^4 + 4ab^3x^6 + b^4x^8} dx$$

↓ 2070

$$\int \frac{1}{(a + bx^2)^4} dx$$

↓ 215

$$\frac{5 \int \frac{1}{(bx^2+a)^3} dx}{6a} + \frac{x}{6a(a + bx^2)^3}$$

↓ 215

$$\frac{5 \left( \frac{3 \int \frac{1}{(bx^2+a)^2} dx}{4a} + \frac{x}{4a(a+bx^2)^2} \right)}{6a} + \frac{x}{6a(a + bx^2)^3}$$

↓ 215

$$\frac{5 \left( \frac{3 \left( \frac{\int \frac{1}{bx^2+a} dx}{2a} + \frac{x}{2a(a+bx^2)} \right)}{4a} + \frac{x}{4a(a+bx^2)^2} \right)}{6a} + \frac{x}{6a(a + bx^2)^3}$$

↓ 218



$$\frac{5 \left( \frac{3 \left( \frac{\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{2a^{3/2}\sqrt{b}} + \frac{x}{2a(a+bx^2)} \right)}{4a} + \frac{x}{4a(a+bx^2)^2} \right)}{6a} + \frac{x}{6a(a+bx^2)^3}$$

input `Int[(a^4 + 4*a^3*b*x^2 + 6*a^2*b^2*x^4 + 4*a*b^3*x^6 + b^4*x^8)^(-1),x]`

output `x/(6*a*(a + b*x^2)^3) + (5*(x/(4*a*(a + b*x^2)^2) + (3*(x/(2*a*(a + b*x^2)) + ArcTan[(Sqrt[b]*x)/Sqrt[a]]/(2*a^(3/2)*Sqrt[b])))/(4*a)))/(6*a)`

### Definitions of rubi rules used

rule 215 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && LtQ[p, -1] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 2070 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{a = Rt[Coeff[Px, x^2, 0], Expon[Px, x^2]], b = Rt[Coeff[Px, x^2, Expon[Px, x^2]], Expon[Px, x^2]]}, Int[u*(a + b*x^2)^(Expon[Px, x^2]*p), x] /; EqQ[Px, (a + b*x^2)^Expon[Px, x^2]]] /; IntegerQ[p] && PolyQ[Px, x^2] && GtQ[Expon[Px, x^2], 1] && NeQ[Coeff[Px, x^2, 0], 0]`

**Maple [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.99

method	result	size
default	$\frac{x}{6a(bx^2+a)^3} + \frac{\frac{5x}{24a(bx^2+a)^2} + \frac{5\left(\frac{3x}{8a(bx^2+a)} + \frac{3\arctan\left(\frac{bx}{\sqrt{ab}}\right)}{8a\sqrt{ab}}\right)}{6a}}{a}$	78
risch	$\frac{\frac{5b^2x^5}{16a^3} + \frac{5bx^3}{6a^2} + \frac{11x}{16a}}{b^3x^6 + 3ab^2x^4 + 3ba^2x^2 + a^3} - \frac{5\ln(bx + \sqrt{-ab})}{32\sqrt{-ab}a^3} + \frac{5\ln(-bx + \sqrt{-ab})}{32\sqrt{-ab}a^3}$	106

input `int(1/(b^4*x^8+4*a*b^3*x^6+6*a^2*b^2*x^4+4*a^3*b*x^2+a^4),x,method=_RETURN  
VERBOSE)`

output `1/6*x/a/(b*x^2+a)^3+5/6/a*(1/4*x/a/(b*x^2+a)^2+3/4/a*(1/2*x/a/(b*x^2+a)+1/  
2/a/(a*b)^(1/2)*arctan(b*x/(a*b)^(1/2))))`

**Fricas [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 254, normalized size of antiderivative = 3.22

$$\int \frac{1}{a^4 + 4a^3bx^2 + 6a^2b^2x^4 + 4ab^3x^6 + b^4x^8} dx$$

$$= \left[ \frac{30ab^3x^5 + 80a^2b^2x^3 + 66a^3bx - 15(b^3x^6 + 3ab^2x^4 + 3a^2bx^2 + a^3)\sqrt{-ab} \log\left(\frac{bx^2 - 2\sqrt{-ab}x - a}{bx^2 + a}\right)}{96(a^4b^4x^6 + 3a^5b^3x^4 + 3a^6b^2x^2 + a^7b)}, \frac{15ab^3x}{96(a^4b^4x^6 + 3a^5b^3x^4 + 3a^6b^2x^2 + a^7b)} \right]$$

input `integrate(1/(b^4*x^8+4*a*b^3*x^6+6*a^2*b^2*x^4+4*a^3*b*x^2+a^4),x, algorit  
hm="fricas")`

output `[1/96*(30*a*b^3*x^5 + 80*a^2*b^2*x^3 + 66*a^3*b*x - 15*(b^3*x^6 + 3*a*b^2*x  
x^4 + 3*a^2*b*x^2 + a^3)*sqrt(-a*b)*log((b*x^2 - 2*sqrt(-a*b)*x - a)/(b*x^2  
2 + a)))/(a^4*b^4*x^6 + 3*a^5*b^3*x^4 + 3*a^6*b^2*x^2 + a^7*b), 1/48*(15*a  
*b^3*x^5 + 40*a^2*b^2*x^3 + 33*a^3*b*x + 15*(b^3*x^6 + 3*a*b^2*x^4 + 3*a^2  
*b*x^2 + a^3)*sqrt(a*b)*arctan(sqrt(a*b)*x/a))/(a^4*b^4*x^6 + 3*a^5*b^3*x^4  
4 + 3*a^6*b^2*x^2 + a^7*b)]`

**Sympy [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.63

$$\int \frac{1}{a^4 + 4a^3bx^2 + 6a^2b^2x^4 + 4ab^3x^6 + b^4x^8} dx$$

$$= -\frac{5\sqrt{-\frac{1}{a^7b}} \log\left(-a^4\sqrt{-\frac{1}{a^7b}} + x\right)}{32} + \frac{5\sqrt{-\frac{1}{a^7b}} \log\left(a^4\sqrt{-\frac{1}{a^7b}} + x\right)}{32}$$

$$+ \frac{33a^2x + 40abx^3 + 15b^2x^5}{48a^6 + 144a^5bx^2 + 144a^4b^2x^4 + 48a^3b^3x^6}$$

input `integrate(1/(b**4*x**8+4*a*b**3*x**6+6*a**2*b**2*x**4+4*a**3*b*x**2+a**4), x)`

output `-5*sqrt(-1/(a**7*b))*log(-a**4*sqrt(-1/(a**7*b)) + x)/32 + 5*sqrt(-1/(a**7*b))*log(a**4*sqrt(-1/(a**7*b)) + x)/32 + (33*a**2*x + 40*a*b*x**3 + 15*b**2*x**5)/(48*a**6 + 144*a**5*b*x**2 + 144*a**4*b**2*x**4 + 48*a**3*b**3*x**6)`

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.01

$$\int \frac{1}{a^4 + 4a^3bx^2 + 6a^2b^2x^4 + 4ab^3x^6 + b^4x^8} dx$$

$$= \frac{15b^2x^5 + 40abx^3 + 33a^2x}{48(a^3b^3x^6 + 3a^4b^2x^4 + 3a^5bx^2 + a^6)} + \frac{5 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{16\sqrt{aba^3}}$$

input `integrate(1/(b^4*x^8+4*a*b^3*x^6+6*a^2*b^2*x^4+4*a^3*b*x^2+a^4), x, algorithm="maxima")`

output `1/48*(15*b^2*x^5 + 40*a*b*x^3 + 33*a^2*x)/(a^3*b^3*x^6 + 3*a^4*b^2*x^4 + 3*a^5*b*x^2 + a^6) + 5/16*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*a^3)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.71

$$\int \frac{1}{a^4 + 4a^3bx^2 + 6a^2b^2x^4 + 4ab^3x^6 + b^4x^8} dx$$

$$= \frac{5 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{16 \sqrt{aba^3}} + \frac{15b^2x^5 + 40abx^3 + 33a^2x}{48(bx^2 + a)^3a^3}$$

input

```
integrate(1/(b^4*x^8+4*a*b^3*x^6+6*a^2*b^2*x^4+4*a^3*b*x^2+a^4),x, algorithm="giac")
```

output

```
5/16*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*a^3) + 1/48*(15*b^2*x^5 + 40*a*b*x^3 + 33*a^2*x)/((b*x^2 + a)^3*a^3)
```

**Mupad [B] (verification not implemented)**

Time = 21.97 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.97

$$\int \frac{1}{a^4 + 4a^3bx^2 + 6a^2b^2x^4 + 4ab^3x^6 + b^4x^8} dx$$

$$= \frac{\frac{11x}{16a} + \frac{5bx^3}{6a^2} + \frac{5b^2x^5}{16a^3}}{a^3 + 3a^2bx^2 + 3ab^2x^4 + b^3x^6} + \frac{5 \operatorname{atan}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{16a^{7/2}\sqrt{b}}$$

input

```
int(1/(a^4 + b^4*x^8 + 4*a^3*b*x^2 + 4*a*b^3*x^6 + 6*a^2*b^2*x^4),x)
```

output

```
((11*x)/(16*a) + (5*b*x^3)/(6*a^2) + (5*b^2*x^5)/(16*a^3))/(a^3 + b^3*x^6 + 3*a^2*b*x^2 + 3*a*b^2*x^4) + (5*atan((b^(1/2)*x)/a^(1/2)))/(16*a^(7/2)*b^(1/2))
```

**Reduce [B] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 162, normalized size of antiderivative = 2.05

$$\int \frac{1}{a^4 + 4a^3bx^2 + 6a^2b^2x^4 + 4ab^3x^6 + b^4x^8} dx$$

$$= \frac{15\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right) a^3 + 45\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right) a^2bx^2 + 45\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right) ab^2x^4 + 15\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right) a^3}{48a^4b(b^3x^6 + 3ab^2x^4 + 3a^2bx^2 + a^3)}$$

input `int(1/(b^4*x^8+4*a*b^3*x^6+6*a^2*b^2*x^4+4*a^3*b*x^2+a^4),x)`

output

```
(15*sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*sqrt(a)))*a**3 + 45*sqrt(b)*sqrt(a)
)*atan((b*x)/(sqrt(b)*sqrt(a)))*a**2*b*x**2 + 45*sqrt(b)*sqrt(a)*atan((b*x
)/(sqrt(b)*sqrt(a)))*a*b**2*x**4 + 15*sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*
sqrt(a)))*b**3*x**6 + 33*a**3*b*x + 40*a**2*b**2*x**3 + 15*a*b**3*x**5)/(4
8*a**4*b*(a**3 + 3*a**2*b*x**2 + 3*a*b**2*x**4 + b**3*x**6))
```

### 3.33 $\int \frac{1}{(a+b(s+x)^2)^4} dx$

Optimal result . . . . .	261
Mathematica [A] (verified) . . . . .	261
Rubi [A] (verified) . . . . .	262
Maple [A] (verified) . . . . .	264
Fricas [B] (verification not implemented) . . . . .	264
Sympy [B] (verification not implemented) . . . . .	265
Maxima [B] (verification not implemented) . . . . .	266
Giac [A] (verification not implemented) . . . . .	267
Mupad [B] (verification not implemented) . . . . .	267
Reduce [B] (verification not implemented) . . . . .	268

#### Optimal result

Integrand size = 11, antiderivative size = 93

$$\int \frac{1}{(a+b(s+x)^2)^4} dx = \frac{s+x}{6a(a+b(s+x)^2)^3} + \frac{5(s+x)}{24a^2(a+b(s+x)^2)^2} + \frac{5(s+x)}{16a^3(a+b(s+x)^2)} + \frac{5 \arctan\left(\frac{\sqrt{b}(s+x)}{\sqrt{a}}\right)}{16a^{7/2}\sqrt{b}}$$

output

```
1/6*(s+x)/a/(a+b*(s+x)^2)^3+5/24*(s+x)/a^2/(a+b*(s+x)^2)^2+5/16*(s+x)/a^3/(a+b*(s+x)^2)+5/16*arctan(b^(1/2)*(s+x)/a^(1/2))/a^(7/2)/b^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.82

$$\int \frac{1}{(a+b(s+x)^2)^4} dx = \frac{(s+x)(33a^2+40ab(s+x)^2+15b^2(s+x)^4)}{48a^3(a+b(s+x)^2)^3} + \frac{5 \arctan\left(\frac{\sqrt{b}(s+x)}{\sqrt{a}}\right)}{16a^{7/2}\sqrt{b}}$$

input

```
Integrate[(a + b*(s + x)^2)^(-4), x]
```

output

$$\frac{((s+x)(33a^2 + 40ab(s+x)^2 + 15b^2(s+x)^4))/(48a^3(a+b(s+x)^2)^3) + (5\text{ArcTan}[\sqrt{b}(s+x)/\sqrt{a}])/(16a^{7/2}\sqrt{b})}{1}$$
**Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.17, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$ , Rules used = {239, 215, 215, 215, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a+b(s+x)^2)^4} dx$$

$$\downarrow 239$$

$$\int \frac{1}{(a+b(s+x)^2)^4} d(s+x)$$

$$\downarrow 215$$

$$\frac{5 \int \frac{1}{(b(s+x)^2+a)^3} d(s+x)}{6a} + \frac{s+x}{6a(a+b(s+x)^2)^3}$$

$$\downarrow 215$$

$$\frac{5 \left( \frac{3 \int \frac{1}{(b(s+x)^2+a)^2} d(s+x)}{4a} + \frac{s+x}{4a(a+b(s+x)^2)^2} \right)}{6a} + \frac{s+x}{6a(a+b(s+x)^2)^3}$$

$$\downarrow 215$$

$$\frac{5 \left( \frac{3 \left( \frac{\int \frac{1}{b(s+x)^2+a} d(s+x)}{2a} + \frac{s+x}{2a(a+b(s+x)^2)} \right)}{4a} + \frac{s+x}{4a(a+b(s+x)^2)^2} \right)}{6a} + \frac{s+x}{6a(a+b(s+x)^2)^3}$$

$$\downarrow 218$$

$$\frac{5 \left( \frac{3 \left( \frac{\arctan\left(\frac{\sqrt{b}(s+x)}{\sqrt{a}}\right)}{2a^{3/2}\sqrt{b}} + \frac{s+x}{2a(a+b(s+x)^2)} \right)}{4a} + \frac{s+x}{4a(a+b(s+x)^2)^2} \right)}{6a} + \frac{s+x}{6a(a+b(s+x)^2)^3}$$

input `Int[(a + b*(s + x)^2)^(-4), x]`

output `(s + x)/(6*a*(a + b*(s + x)^2)^3) + (5*((s + x)/(4*a*(a + b*(s + x)^2)^2) + (3*((s + x)/(2*a*(a + b*(s + x)^2)) + ArcTan[(Sqrt[b]*(s + x))/Sqrt[a]]/(2*a^(3/2)*Sqrt[b])))/(4*a)))/(6*a)`

### Defintions of rubi rules used

rule 215 `Int[((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && LtQ[p, -1] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 239 `Int[((a_.) + (b_.)*(v_)^(n_))^(p_), x_Symbol] := Simp[1/Coefficient[v, x, 1] Subst[Int[(a + b*x^n)^p, x], x, v], x] /; FreeQ[{a, b, n, p}, x] && LinearQ[v, x] && NeQ[v, x]`



**Maple [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.60

method	result
default	$\frac{2bs+2bx}{12ab(b^2s^2+2bsx+bx^2+a)^3} + \frac{5(2bs+2bx)}{48ab(b^2s^2+2bsx+bx^2+a)^2} + \frac{5 \left( \frac{3(2bs+2bx)}{16ab(b^2s^2+2bsx+bx^2+a)} + \frac{3 \arctan\left(\frac{2bs+2bx}{2\sqrt{ab}}\right)}{8a\sqrt{ab}} \right)}{6a}$
risch	$\frac{\frac{5b^2x^5}{16a^3} + \frac{25b^2sx^4}{16a^3} + \frac{5b(15bs^2+4a)x^3}{24a^3} + \frac{5bs(5bs^2+4a)x^2}{8a^3} + \frac{(25b^2s^4+40abs^2+11a^2)x}{16a^3} + \frac{s(15b^2s^4+40abs^2+33a^2)}{48a^3}}{(b^2s^2+2bsx+bx^2+a)^3} - \frac{5 \ln(bs+bx+\sqrt{-ab})}{32\sqrt{-ab}a^3}$

input `int(1/(a+b*(s+x)^2)^4,x,method=_RETURNVERBOSE)`

output `1/12*(2*b*s+2*b*x)/a/b/(b*s^2+2*b*s*x+b*x^2+a)^3+5/6/a*(1/8*(2*b*s+2*b*x)/a/b/(b*s^2+2*b*s*x+b*x^2+a)^2+3/4/a*(1/4*(2*b*s+2*b*x)/a/b/(b*s^2+2*b*s*x+b*x^2+a)+1/2/a/(a*b)^(1/2)*arctan(1/2*(2*b*s+2*b*x)/(a*b)^(1/2)))`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 465 vs. 2(77) = 154.

Time = 0.11 (sec) , antiderivative size = 970, normalized size of antiderivative = 10.43

$$\int \frac{1}{(a+b(s+x)^2)^4} dx = \text{Too large to display}$$

input `integrate(1/(a+b*(s+x)^2)^4,x, algorithm="fricas")`

output

```
[1/96*(30*a*b^3*s^5 + 150*a*b^3*s*x^4 + 30*a*b^3*x^5 + 80*a^2*b^2*s^3 + 66
*a^3*b*s + 20*(15*a*b^3*s^2 + 4*a^2*b^2)*x^3 + 60*(5*a*b^3*s^3 + 4*a^2*b^2
*s)*x^2 - 15*(b^3*s^6 + 6*b^3*s*x^5 + b^3*x^6 + 3*a*b^2*s^4 + 3*a^2*b*s^2
+ 3*(5*b^3*s^2 + a*b^2)*x^4 + 4*(5*b^3*s^3 + 3*a*b^2*s)*x^3 + a^3 + 3*(5*b
^3*s^4 + 6*a*b^2*s^2 + a^2*b)*x^2 + 6*(b^3*s^5 + 2*a*b^2*s^3 + a^2*b*s)*x)
*sqrt(-a*b)*log((b*s^2 + 2*b*s*x + b*x^2 - 2*sqrt(-a*b)*(s + x) - a)/(b*s^
2 + 2*b*s*x + b*x^2 + a)) + 6*(25*a*b^3*s^4 + 40*a^2*b^2*s^2 + 11*a^3*b)*x
)/(a^4*b^4*s^6 + 6*a^4*b^4*s*x^5 + a^4*b^4*x^6 + 3*a^5*b^3*s^4 + 3*a^6*b^2
*s^2 + a^7*b + 3*(5*a^4*b^4*s^2 + a^5*b^3)*x^4 + 4*(5*a^4*b^4*s^3 + 3*a^5*
b^3*s)*x^3 + 3*(5*a^4*b^4*s^4 + 6*a^5*b^3*s^2 + a^6*b^2)*x^2 + 6*(a^4*b^4*
s^5 + 2*a^5*b^3*s^3 + a^6*b^2*s)*x), 1/48*(15*a*b^3*s^5 + 75*a*b^3*s*x^4 +
15*a*b^3*x^5 + 40*a^2*b^2*s^3 + 33*a^3*b*s + 10*(15*a*b^3*s^2 + 4*a^2*b^2
)*x^3 + 30*(5*a*b^3*s^3 + 4*a^2*b^2*s)*x^2 + 15*(b^3*s^6 + 6*b^3*s*x^5 + b
^3*x^6 + 3*a*b^2*s^4 + 3*a^2*b*s^2 + 3*(5*b^3*s^2 + a*b^2)*x^4 + 4*(5*b^3*
s^3 + 3*a*b^2*s)*x^3 + a^3 + 3*(5*b^3*s^4 + 6*a*b^2*s^2 + a^2*b)*x^2 + 6*(
b^3*s^5 + 2*a*b^2*s^3 + a^2*b*s)*x)*sqrt(a*b)*arctan(sqrt(a*b)*(s + x)/a)
+ 3*(25*a*b^3*s^4 + 40*a^2*b^2*s^2 + 11*a^3*b)*x)/(a^4*b^4*s^6 + 6*a^4*b^4
*s*x^5 + a^4*b^4*x^6 + 3*a^5*b^3*s^4 + 3*a^6*b^2*s^2 + a^7*b + 3*(5*a^4*b^
4*s^2 + a^5*b^3)*x^4 + 4*(5*a^4*b^4*s^3 + 3*a^5*b^3*s)*x^3 + 3*(5*a^4*b^4*
s^4 + 6*a^5*b^3*s^2 + a^6*b^2)*x^2 + 6*(a^4*b^4*s^5 + 2*a^5*b^3*s^3 + a...
```

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 354 vs.  $2(92) = 184$ .

Time = 1.02 (sec) , antiderivative size = 354, normalized size of antiderivative = 3.81

$$\int \frac{1}{(a + b(s + x))^4} dx$$

$$= -\frac{5\sqrt{-\frac{1}{a^7b}} \log\left(-a^4\sqrt{-\frac{1}{a^7b}} + s + x\right)}{32} + \frac{5\sqrt{-\frac{1}{a^7b}} \log\left(a^4\sqrt{-\frac{1}{a^7b}} + s + x\right)}{32}$$

$$+ \frac{33a^2s + 40abs^3 + 15b^2s^5 + 75b^2sx^4 + 15b^2x^5 + x^3 \cdot (40ab + 1)}{48a^6 + 144a^5bs^2 + 144a^4b^2s^4 + 48a^3b^3s^6 + 288a^3b^3sx^5 + 48a^3b^3x^6 + x^4 \cdot (144a^4b^2 + 720a^3b^3s^2) + x^3}$$

input

```
integrate(1/(a+b*(s+x)**2)**4,x)
```

output

```
-5*sqrt(-1/(a**7*b))*log(-a**4*sqrt(-1/(a**7*b)) + s + x)/32 + 5*sqrt(-1/(
a**7*b))*log(a**4*sqrt(-1/(a**7*b)) + s + x)/32 + (33*a**2*s + 40*a*b*s**3
+ 15*b**2*s**5 + 75*b**2*s*x**4 + 15*b**2*x**5 + x**3*(40*a*b + 150*b**2*
s**2) + x**2*(120*a*b*s + 150*b**2*s**3) + x*(33*a**2 + 120*a*b*s**2 + 75*
b**2*s**4))/(48*a**6 + 144*a**5*b*s**2 + 144*a**4*b**2*s**4 + 48*a**3*b**3
*s**6 + 288*a**3*b**3*s*x**5 + 48*a**3*b**3*x**6 + x**4*(144*a**4*b**2 + 7
20*a**3*b**3*s**2) + x**3*(576*a**4*b**2*s + 960*a**3*b**3*s**3) + x**2*(1
44*a**5*b + 864*a**4*b**2*s**2 + 720*a**3*b**3*s**4) + x*(288*a**5*b*s + 5
76*a**4*b**2*s**3 + 288*a**3*b**3*s**5))
```

### Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 299 vs.  $2(77) = 154$ .

Time = 0.11 (sec) , antiderivative size = 299, normalized size of antiderivative = 3.22

$$\int \frac{1}{(a + b(s + x)^2)^4} dx$$

$$= \frac{15b^2s^5 + 75b^2sx^4 + 15b^2x^5 + 40abs^3 + 10(15b^2s^2 + 4ab)x^3 + 33a^2s + 30(5b^2s^2 + 4ab^2)x + 5a^3b^3s^3 + 3a^4b^2s^2 + 3a^5bs^2 + a^6 + 3(5a^3b^3s^2 + a^4b^2)x^4 + 4(5a^3b^3s^3 + 3a^4b^2s^2 + 3a^5bs^2 + a^6)}{48(a^3b^3s^6 + 6a^3b^3sx^5 + a^3b^3x^6 + 3a^4b^2s^4 + 3a^5bs^2 + a^6 + 3(5a^3b^3s^2 + a^4b^2)x^4 + 4(5a^3b^3s^3 + 3a^4b^2s^2 + 3a^5bs^2 + a^6))} + \frac{5 \arctan\left(\frac{bs+bx}{\sqrt{ab}}\right)}{16\sqrt{aba^3}}$$

input

```
integrate(1/(a+b*(s+x)^2)^4,x, algorithm="maxima")
```

output

```
1/48*(15*b^2*s^5 + 75*b^2*s*x^4 + 15*b^2*x^5 + 40*a*b*s^3 + 10*(15*b^2*s^2
+ 4*a*b)*x^3 + 33*a^2*s + 30*(5*b^2*s^3 + 4*a*b*s)*x^2 + 3*(25*b^2*s^4 +
40*a*b*s^2 + 11*a^2)*x)/(a^3*b^3*s^6 + 6*a^3*b^3*s*x^5 + a^3*b^3*x^6 + 3*a
^4*b^2*s^4 + 3*a^5*b*s^2 + a^6 + 3*(5*a^3*b^3*s^2 + a^4*b^2)*x^4 + 4*(5*a
^3*b^3*s^3 + 3*a^4*b^2*s)*x^3 + 3*(5*a^3*b^3*s^4 + 6*a^4*b^2*s^2 + a^5*b)*x
^2 + 6*(a^3*b^3*s^5 + 2*a^4*b^2*s^3 + a^5*b*s)*x) + 5/16*arctan((b*s + b*x
)/sqrt(a*b))/(sqrt(a*b)*a^3)
```

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.59

$$\int \frac{1}{(a + b(s + x)^2)^4} dx = \frac{5 \arctan\left(\frac{bs+bx}{\sqrt{ab}}\right)}{16 \sqrt{ab} a^3} + \frac{15 b^2 s^5 + 75 b^2 s^4 x + 150 b^2 s^3 x^2 + 150 b^2 s^2 x^3 + 75 b^2 s x^4 + 15 b^2 x^5 + 40 a b s^3 + 120 a b s^2 x + 120 a b s x^2 + 40 a^2 b x^3 + 33 a^2 s + 33 a^2 x}{48 (b s^2 + 2 b s x + b x^2 + a)^3 a^3}$$

input `integrate(1/(a+b*(s+x)^2)^4,x, algorithm="giac")`output `5/16*arctan((b*s + b*x)/sqrt(a*b))/(sqrt(a*b)*a^3) + 1/48*(15*b^2*s^5 + 75*b^2*s^4*x + 150*b^2*s^3*x^2 + 150*b^2*s^2*x^3 + 75*b^2*s*x^4 + 15*b^2*x^5 + 40*a*b*s^3 + 120*a*b*s^2*x + 120*a*b*s*x^2 + 40*a*b*x^3 + 33*a^2*s + 33*a^2*x)/((b*s^2 + 2*b*s*x + b*x^2 + a)^3*a^3)`**Mupad [B] (verification not implemented)**

Time = 22.19 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.98

$$\int \frac{1}{(a + b(s + x)^2)^4} dx = \frac{\frac{11(s+x)}{16a} + \frac{5b(s+x)^3}{6a^2} + \frac{5b^2(s+x)^5}{16a^3}}{a^3 + b^3(s+x)^6 + 3a^2b(s+x)^2 + 3ab^2(s+x)^4} + \frac{5 \operatorname{atan}\left(\frac{\sqrt{b}(s+x)}{\sqrt{a}}\right)}{16 a^{7/2} \sqrt{b}}$$

input `int(1/(a + b*(s + x)^2)^4,x)`output `((11*(s + x))/(16*a) + (5*b*(s + x)^3)/(6*a^2) + (5*b^2*(s + x)^5)/(16*a^3))/((a^3 + b^3*(s + x)^6 + 3*a^2*b*(s + x)^2 + 3*a*b^2*(s + x)^4) + (5*atan((b^(1/2)*(s + x))/a^(1/2)))/(16*a^(7/2)*b^(1/2))`

**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 808, normalized size of antiderivative = 8.69

$$\int \frac{1}{(a + b(s + x)^2)^4} dx = \text{Too large to display}$$

input `int(1/(a+b*(s+x)^2)^4,x)`

output

```
(30*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**3*s + 90*sqrt(b)
)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**2*b*s**3 + 180*sqrt(b)*sq
rt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**2*b*s**2*x + 90*sqrt(b)*sqrt(
a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**2*b*s*x**2 + 90*sqrt(b)*sqrt(a)*
atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**5 + 360*sqrt(b)*sqrt(a)*atan
((b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**4*x + 540*sqrt(b)*sqrt(a)*atan((
b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**3*x**2 + 360*sqrt(b)*sqrt(a)*atan(
(b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**2*x**3 + 90*sqrt(b)*sqrt(a)*atan(
(b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s*x**4 + 30*sqrt(b)*sqrt(a)*atan((b*
s + b*x)/(sqrt(b)*sqrt(a)))*b**3*s**7 + 180*sqrt(b)*sqrt(a)*atan((b*s + b*
x)/(sqrt(b)*sqrt(a)))*b**3*s**6*x + 450*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(
sqrt(b)*sqrt(a)))*b**3*s**5*x**2 + 600*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(s
qrt(b)*sqrt(a)))*b**3*s**4*x**3 + 450*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sq
rt(b)*sqrt(a)))*b**3*s**3*x**4 + 180*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqr
t(b)*sqrt(a)))*b**3*s**2*x**5 + 30*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(
b)*sqrt(a)))*b**3*s*x**6 - 5*a**4 + 51*a**3*b*s**2 + 36*a**3*b*s*x - 15*a*
*3*b*x**2 + 65*a**2*b**2*s**4 + 180*a**2*b**2*s**3*x + 150*a**2*b**2*s**2*
x**2 + 20*a**2*b**2*s*x**3 - 15*a**2*b**2*x**4 + 25*a*b**3*s**6 + 120*a*b*
*3*s**5*x + 225*a*b**3*s**4*x**2 + 200*a*b**3*s**3*x**3 + 75*a*b**3*s**2*x
**4 - 5*a*b**3*x**6)/(96*a**4*b*s*(a**3 + 3*a**2*b*s**2 + 6*a**2*b*s*x ...
```

**3.34**  $\int \frac{1}{(a^2+2ab(s+x)^2+b^2(s+x)^4)^2} dx$

Optimal result	269
Mathematica [A] (verified)	269
Rubi [A] (verified)	270
Maple [A] (verified)	272
Fricas [B] (verification not implemented)	272
Sympy [B] (verification not implemented)	273
Maxima [B] (verification not implemented)	274
Giac [A] (verification not implemented)	275
Mupad [B] (verification not implemented)	275
Reduce [B] (verification not implemented)	276

**Optimal result**

Integrand size = 24, antiderivative size = 93

$$\int \frac{1}{(a^2 + 2ab(s+x)^2 + b^2(s+x)^4)^2} dx = \frac{s+x}{6a(a+b(s+x)^2)^3} + \frac{5(s+x)}{24a^2(a+b(s+x)^2)^2} + \frac{5(s+x)}{16a^3(a+b(s+x)^2)} + \frac{5 \arctan\left(\frac{\sqrt{b}(s+x)}{\sqrt{a}}\right)}{16a^{7/2}\sqrt{b}}$$

output

```
1/6*(s+x)/a/(a+b*(s+x)^2)^3+5/24*(s+x)/a^2/(a+b*(s+x)^2)^2+5/16*(s+x)/a^3/(a+b*(s+x)^2)+5/16*arctan(b^(1/2)*(s+x)/a^(1/2))/a^(7/2)/b^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.82

$$\int \frac{1}{(a^2 + 2ab(s+x)^2 + b^2(s+x)^4)^2} dx = \frac{(s+x)(33a^2 + 40ab(s+x)^2 + 15b^2(s+x)^4)}{48a^3(a+b(s+x)^2)^3} + \frac{5 \arctan\left(\frac{\sqrt{b}(s+x)}{\sqrt{a}}\right)}{16a^{7/2}\sqrt{b}}$$

input

```
Integrate[(a^2 + 2*a*b*(s + x)^2 + b^2*(s + x)^4)^(-2),x]
```

output

$$\frac{((s+x)(33a^2 + 40ab(s+x)^2 + 15b^2(s+x)^4))/(48a^3(a+b(s+x)^2)^3) + (5\text{ArcTan}[\sqrt{b}(s+x)]/\sqrt{a})/(16a^{7/2}\sqrt{b})}{1}$$
**Rubi [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.38, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {1687, 1379, 215, 215, 215, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a^2 + 2ab(s+x)^2 + b^2(s+x)^4)^2} dx$$

↓ 1687

$$\int \frac{1}{(a^2 + 2ab(s+x)^2 + b^2(s+x)^4)^2} d(s+x)$$

↓ 1379

$$b^4 \int \frac{1}{(b^2(s+x)^2 + ab)^4} d(s+x)$$

↓ 215

$$b^4 \left( \frac{5 \int \frac{1}{(b^2(s+x)^2 + ab)^3} d(s+x)}{6ab} + \frac{s+x}{6ab^4 (a + b(s+x)^2)^3} \right)$$

↓ 215

$$b^4 \left( \frac{5 \left( \frac{3 \int \frac{1}{(b^2(s+x)^2 + ab)^2} d(s+x)}{4ab} + \frac{s+x}{4ab^3 (a + b(s+x)^2)^2} \right)}{6ab} + \frac{s+x}{6ab^4 (a + b(s+x)^2)^3} \right)$$

↓ 215

$$b^4 \left( \frac{5 \left( \frac{3 \left( \frac{\int \frac{1}{b^2(s+x)^2+ab} d(s+x)}{2ab} + \frac{s+x}{2ab^2(a+b(s+x)^2)} \right)}{4ab} + \frac{s+x}{4ab^3(a+b(s+x)^2)^2} \right)}{6ab} + \frac{s+x}{6ab^4(a+b(s+x)^2)^3} \right)$$

↓ 218

$$b^4 \left( \frac{5 \left( \frac{3 \left( \frac{\arctan\left(\frac{\sqrt{b}(s+x)}{\sqrt{a}}\right)}{2a^{3/2}b^{5/2}} + \frac{s+x}{2ab^2(a+b(s+x)^2)} \right)}{4ab} + \frac{s+x}{4ab^3(a+b(s+x)^2)^2} \right)}{6ab} + \frac{s+x}{6ab^4(a+b(s+x)^2)^3} \right)$$

input `Int[(a^2 + 2*a*b*(s + x)^2 + b^2*(s + x)^4)^(-2),x]`

output `b^4*((s + x)/(6*a*b^4*(a + b*(s + x)^2)^3) + (5*((s + x)/(4*a*b^3*(a + b*(s + x)^2)^2) + (3*((s + x)/(2*a*b^2*(a + b*(s + x)^2))) + ArcTan[(Sqrt[b]*(s + x))/Sqrt[a]]/(2*a^(3/2)*b^(5/2))))/(4*a*b)))/(6*a*b)`

### Defintions of rubi rules used

rule 215 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && LtQ[p, -1] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`



rule 1379

```
Int[((a_) + (c_)*(x_)^(n2_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/
c^p Int[(b/2 + c*x^n)^(2*p), x], x] /; FreeQ[{a, b, c, n, p}, x] && EqQ[n
2, 2*n] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p] && NeQ[p, 1]
```

rule 1687

```
Int[((a_) + (c_)*(u_)^(n2_) + (b_)*(u_)^(n_))^(p_), x_Symbol] := Simp[1/
Coefficient[u, x, 1] Subst[Int[(a + b*x^n + c*x^(2*n))^p, x], x, u], x] /
; FreeQ[{a, b, c, n, p}, x] && EqQ[n2, 2*n] && LinearQ[u, x] && NeQ[u, x]
```

### Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.60

method	result
default	$\frac{2bs+2bx}{12ab(b^2s^2+2bsx+bx^2+a)^3} + \frac{5(2bs+2bx)}{48ab(b^2s^2+2bsx+bx^2+a)^2} + \frac{5 \left( \frac{3(2bs+2bx)}{16ab(b^2s^2+2bsx+bx^2+a)} + \frac{3 \arctan\left(\frac{2bs+2bx}{2\sqrt{ab}}\right)}{8a\sqrt{ab}} \right)}{6a}$
risch	$\frac{5b^2x^5}{16a^3} + \frac{25b^2sx^4}{16a^3} + \frac{5b(15bs^2+4a)x^3}{24a^3} + \frac{5bs(5bs^2+4a)x^2}{8a^3} + \frac{(25b^2s^4+40abs^2+11a^2)x}{16a^3} + \frac{s(15b^2s^4+40abs^2+33a^2)}{48a^3} - \frac{5 \ln(bs+bx+\sqrt{-ab})}{32\sqrt{-ab}a^3}$

input

```
int(1/(a^2+2*a*b*(s+x)^2+b^2*(s+x)^4)^2,x,method=_RETURNVERBOSE)
```

output

```
1/12*(2*b*s+2*b*x)/a/b/(b*s^2+2*b*s*x+b*x^2+a)^3+5/6/a*(1/8*(2*b*s+2*b*x)/
a/b/(b*s^2+2*b*s*x+b*x^2+a)^2+3/4/a*(1/4*(2*b*s+2*b*x)/a/b/(b*s^2+2*b*s*x+
b*x^2+a)+1/2/a/(a*b)^(1/2)*arctan(1/2*(2*b*s+2*b*x)/(a*b)^(1/2)))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 465 vs. 2(77) = 154.

Time = 0.10 (sec) , antiderivative size = 970, normalized size of antiderivative = 10.43

$$\int \frac{1}{(a^2 + 2ab(s+x)^2 + b^2(s+x)^4)^2} dx = \text{Too large to display}$$

input

```
integrate(1/(a^2+2*a*b*(s+x)^2+b^2*(s+x)^4)^2,x, algorithm="fricas")
```

output

```
[1/96*(30*a*b^3*s^5 + 150*a*b^3*s*x^4 + 30*a*b^3*x^5 + 80*a^2*b^2*s^3 + 66
*a^3*b*s + 20*(15*a*b^3*s^2 + 4*a^2*b^2)*x^3 + 60*(5*a*b^3*s^3 + 4*a^2*b^2
*s)*x^2 - 15*(b^3*s^6 + 6*b^3*s*x^5 + b^3*x^6 + 3*a*b^2*s^4 + 3*a^2*b*s^2
+ 3*(5*b^3*s^2 + a*b^2)*x^4 + 4*(5*b^3*s^3 + 3*a*b^2*s)*x^3 + a^3 + 3*(5*b
^3*s^4 + 6*a*b^2*s^2 + a^2*b)*x^2 + 6*(b^3*s^5 + 2*a*b^2*s^3 + a^2*b*s)*x)
*sqrt(-a*b)*log((b*s^2 + 2*b*s*x + b*x^2 - 2*sqrt(-a*b)*(s + x) - a)/(b*s^
2 + 2*b*s*x + b*x^2 + a)) + 6*(25*a*b^3*s^4 + 40*a^2*b^2*s^2 + 11*a^3*b)*x
)/(a^4*b^4*s^6 + 6*a^4*b^4*s*x^5 + a^4*b^4*x^6 + 3*a^5*b^3*s^4 + 3*a^6*b^2
*s^2 + a^7*b + 3*(5*a^4*b^4*s^2 + a^5*b^3)*x^4 + 4*(5*a^4*b^4*s^3 + 3*a^5*
b^3*s)*x^3 + 3*(5*a^4*b^4*s^4 + 6*a^5*b^3*s^2 + a^6*b^2)*x^2 + 6*(a^4*b^4*
s^5 + 2*a^5*b^3*s^3 + a^6*b^2*s)*x), 1/48*(15*a*b^3*s^5 + 75*a*b^3*s*x^4 +
15*a*b^3*x^5 + 40*a^2*b^2*s^3 + 33*a^3*b*s + 10*(15*a*b^3*s^2 + 4*a^2*b^2
)*x^3 + 30*(5*a*b^3*s^3 + 4*a^2*b^2*s)*x^2 + 15*(b^3*s^6 + 6*b^3*s*x^5 + b
^3*x^6 + 3*a*b^2*s^4 + 3*a^2*b*s^2 + 3*(5*b^3*s^2 + a*b^2)*x^4 + 4*(5*b^3*
s^3 + 3*a*b^2*s)*x^3 + a^3 + 3*(5*b^3*s^4 + 6*a*b^2*s^2 + a^2*b)*x^2 + 6*(
b^3*s^5 + 2*a*b^2*s^3 + a^2*b*s)*x)*sqrt(a*b)*arctan(sqrt(a*b)*(s + x)/a)
+ 3*(25*a*b^3*s^4 + 40*a^2*b^2*s^2 + 11*a^3*b)*x)/(a^4*b^4*s^6 + 6*a^4*b^4
*s*x^5 + a^4*b^4*x^6 + 3*a^5*b^3*s^4 + 3*a^6*b^2*s^2 + a^7*b + 3*(5*a^4*b^
4*s^2 + a^5*b^3)*x^4 + 4*(5*a^4*b^4*s^3 + 3*a^5*b^3*s)*x^3 + 3*(5*a^4*b^4*
s^4 + 6*a^5*b^3*s^2 + a^6*b^2)*x^2 + 6*(a^4*b^4*s^5 + 2*a^5*b^3*s^3 + a...
```

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 354 vs.  $2(92) = 184$ .

Time = 0.97 (sec) , antiderivative size = 354, normalized size of antiderivative = 3.81

$$\int \frac{1}{(a^2 + 2ab(s+x)^2 + b^2(s+x)^4)^2} dx$$

$$= -\frac{5\sqrt{-\frac{1}{a^7b}} \log\left(-a^4\sqrt{-\frac{1}{a^7b}} + s + x\right)}{32} + \frac{5\sqrt{-\frac{1}{a^7b}} \log\left(a^4\sqrt{-\frac{1}{a^7b}} + s + x\right)}{32}$$

$$+ \frac{33a^2s + 40abs^3 + 15b^2s^5 + 75b^2sx^4 + 15b^2x^5 + x^3 \cdot (40ab + 1)}{48a^6 + 144a^5bs^2 + 144a^4b^2s^4 + 48a^3b^3s^6 + 288a^3b^3sx^5 + 48a^3b^3x^6 + x^4 \cdot (144a^4b^2 + 720a^3b^3s^2) + x^3}$$

input

```
integrate(1/(a**2+2*a*b*(s+x)**2+b**2*(s+x)**4)**2,x)
```

output

```
-5*sqrt(-1/(a**7*b))*log(-a**4*sqrt(-1/(a**7*b)) + s + x)/32 + 5*sqrt(-1/(
a**7*b))*log(a**4*sqrt(-1/(a**7*b)) + s + x)/32 + (33*a**2*s + 40*a*b*s**3
+ 15*b**2*s**5 + 75*b**2*s*x**4 + 15*b**2*x**5 + x**3*(40*a*b + 150*b**2*
s**2) + x**2*(120*a*b*s + 150*b**2*s**3) + x*(33*a**2 + 120*a*b*s**2 + 75*
b**2*s**4))/(48*a**6 + 144*a**5*b*s**2 + 144*a**4*b**2*s**4 + 48*a**3*b**3
*s**6 + 288*a**3*b**3*s*x**5 + 48*a**3*b**3*x**6 + x**4*(144*a**4*b**2 + 7
20*a**3*b**3*s**2) + x**3*(576*a**4*b**2*s + 960*a**3*b**3*s**3) + x**2*(1
44*a**5*b + 864*a**4*b**2*s**2 + 720*a**3*b**3*s**4) + x*(288*a**5*b*s + 5
76*a**4*b**2*s**3 + 288*a**3*b**3*s**5))
```

### Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 299 vs.  $2(77) = 154$ .

Time = 0.12 (sec) , antiderivative size = 299, normalized size of antiderivative = 3.22

$$\int \frac{1}{(a^2 + 2ab(s+x)^2 + b^2(s+x)^4)^2} dx$$

$$= \frac{15b^2s^5 + 75b^2sx^4 + 15b^2x^5 + 40abs^3 + 10(15b^2s^2 + 4ab)x^3 + 33a^2s + 30(5b^2s^2 + 4a^2b^2)x^2 + 33a^2s + 30(5b^2s^2 + 4a^2b^2)x^2 + 33a^2s + 30(5b^2s^2 + 4a^2b^2)x^2}{48(a^3b^3s^6 + 6a^3b^3sx^5 + a^3b^3x^6 + 3a^4b^2s^4 + 3a^5bs^2 + a^6 + 3(5a^3b^3s^2 + a^4b^2)x^4 + 4(5a^3b^3s^3 + 3a^4b^2s^2)x^3 + 3(5a^3b^3s^4 + 6a^4b^2s^2 + a^5b)x^2 + 6(a^3b^3s^5 + 2a^4b^2s^3 + a^5b)s + a^6) + \frac{5 \arctan\left(\frac{bs+bx}{\sqrt{ab}}\right)}{16\sqrt{aba^3}}$$

input

```
integrate(1/(a^2+2*a*b*(s+x)^2+b^2*(s+x)^4)^2,x, algorithm="maxima")
```

output

```
1/48*(15*b^2*s^5 + 75*b^2*s*x^4 + 15*b^2*x^5 + 40*a*b*s^3 + 10*(15*b^2*s^2
+ 4*a*b)*x^3 + 33*a^2*s + 30*(5*b^2*s^3 + 4*a*b*s)*x^2 + 3*(25*b^2*s^4 +
40*a*b*s^2 + 11*a^2)*x)/(a^3*b^3*s^6 + 6*a^3*b^3*s*x^5 + a^3*b^3*x^6 + 3*a
^4*b^2*s^4 + 3*a^5*b*s^2 + a^6 + 3*(5*a^3*b^3*s^2 + a^4*b^2)*x^4 + 4*(5*a
^3*b^3*s^3 + 3*a^4*b^2*s)*x^3 + 3*(5*a^3*b^3*s^4 + 6*a^4*b^2*s^2 + a^5*b)*x
^2 + 6*(a^3*b^3*s^5 + 2*a^4*b^2*s^3 + a^5*b*s)*x) + 5/16*arctan((b*s + b*x
)/sqrt(a*b))/(sqrt(a*b)*a^3)
```

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.59

$$\int \frac{1}{(a^2 + 2ab(s+x)^2 + b^2(s+x)^4)^2} dx = \frac{5 \arctan\left(\frac{bs+bx}{\sqrt{ab}}\right)}{16 \sqrt{ab} a^3} + \frac{15 b^2 s^5 + 75 b^2 s^4 x + 150 b^2 s^3 x^2 + 150 b^2 s^2 x^3 + 75 b^2 s x^4 + 15 b^2 x^5 + 40 a b s^3 + 120 a b s^2 x + 120 a b s x^2 + 40 a^2 b x^3 + 33 a^2 s + 33 a^2 x}{48 (b s^2 + 2 b s x + b x^2 + a)^3 a^3}$$

input `integrate(1/(a^2+2*a*b*(s+x)^2+b^2*(s+x)^4)^2,x, algorithm="giac")`output `5/16*arctan((b*s + b*x)/sqrt(a*b))/(sqrt(a*b)*a^3) + 1/48*(15*b^2*s^5 + 75*b^2*s^4*x + 150*b^2*s^3*x^2 + 150*b^2*s^2*x^3 + 75*b^2*s*x^4 + 15*b^2*x^5 + 40*a*b*s^3 + 120*a*b*s^2*x + 120*a*b*s*x^2 + 40*a*b*x^3 + 33*a^2*s + 33*a^2*x)/((b*s^2 + 2*b*s*x + b*x^2 + a)^3*a^3)`**Mupad [B] (verification not implemented)**

Time = 22.20 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.98

$$\int \frac{1}{(a^2 + 2ab(s+x)^2 + b^2(s+x)^4)^2} dx = \frac{\frac{11(s+x)}{16a} + \frac{5b(s+x)^3}{6a^2} + \frac{5b^2(s+x)^5}{16a^3}}{a^3 + b^3(s+x)^6 + 3a^2b(s+x)^2 + 3ab^2(s+x)^4} + \frac{5 \operatorname{atan}\left(\frac{\sqrt{b}(s+x)}{\sqrt{a}}\right)}{16 a^{7/2} \sqrt{b}}$$

input `int(1/(a^2 + b^2*(s + x)^4 + 2*a*b*(s + x)^2)^2,x)`output `((11*(s + x))/(16*a) + (5*b*(s + x)^3)/(6*a^2) + (5*b^2*(s + x)^5)/(16*a^3))/((a^3 + b^3*(s + x)^6 + 3*a^2*b*(s + x)^2 + 3*a*b^2*(s + x)^4) + (5*atan((b^(1/2)*(s + x))/a^(1/2)))/(16*a^(7/2)*b^(1/2)))`

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 808, normalized size of antiderivative = 8.69

$$\int \frac{1}{(a^2 + 2ab(s+x)^2 + b^2(s+x)^4)^2} dx = \text{Too large to display}$$

input `int(1/(a^2+2*a*b*(s+x)^2+b^2*(s+x)^4)^2,x)`

output

```
(30*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**3*s + 90*sqrt(b)
)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**2*b*s**3 + 180*sqrt(b)*sq
rt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**2*b*s**2*x + 90*sqrt(b)*sqrt(
a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**2*b*s*x**2 + 90*sqrt(b)*sqrt(a)*
atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**5 + 360*sqrt(b)*sqrt(a)*atan
((b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**4*x + 540*sqrt(b)*sqrt(a)*atan((
b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**3*x**2 + 360*sqrt(b)*sqrt(a)*atan(
(b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**2*x**3 + 90*sqrt(b)*sqrt(a)*atan(
(b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s*x**4 + 30*sqrt(b)*sqrt(a)*atan((b*
s + b*x)/(sqrt(b)*sqrt(a)))*b**3*s**7 + 180*sqrt(b)*sqrt(a)*atan((b*s + b*
x)/(sqrt(b)*sqrt(a)))*b**3*s**6*x + 450*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(
sqrt(b)*sqrt(a)))*b**3*s**5*x**2 + 600*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(s
qrt(b)*sqrt(a)))*b**3*s**4*x**3 + 450*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sq
rt(b)*sqrt(a)))*b**3*s**3*x**4 + 180*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqr
t(b)*sqrt(a)))*b**3*s**2*x**5 + 30*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(
b)*sqrt(a)))*b**3*s*x**6 - 5*a**4 + 51*a**3*b*s**2 + 36*a**3*b*s*x - 15*a*
*3*b*x**2 + 65*a**2*b**2*s**4 + 180*a**2*b**2*s**3*x + 150*a**2*b**2*s**2*
x**2 + 20*a**2*b**2*s*x**3 - 15*a**2*b**2*x**4 + 25*a*b**3*s**6 + 120*a*b*
*3*s**5*x + 225*a*b**3*s**4*x**2 + 200*a*b**3*s**3*x**3 + 75*a*b**3*s**2*x
**4 - 5*a*b**3*x**6)/(96*a**4*b*s*(a**3 + 3*a**2*b*s**2 + 6*a**2*b*s*x ...
```

### 3.35 $\int \frac{1}{a^4+4a^3b(s+x)^2+6a^2b^2(s+x)^4+4ab^3(s+x)^6+b^4(s+x)^8} dx$

Optimal result . . . . .	277
Mathematica [A] (verified) . . . . .	277
Rubi [B] (verified) . . . . .	278
Maple [A] (verified) . . . . .	279
Fricas [B] (verification not implemented) . . . . .	280
Sympy [B] (verification not implemented) . . . . .	281
Maxima [B] (verification not implemented) . . . . .	281
Giac [A] (verification not implemented) . . . . .	282
Mupad [B] (verification not implemented) . . . . .	283
Reduce [B] (verification not implemented) . . . . .	283

#### Optimal result

Integrand size = 50, antiderivative size = 93

$$\int \frac{1}{a^4 + 4a^3b(s+x)^2 + 6a^2b^2(s+x)^4 + 4ab^3(s+x)^6 + b^4(s+x)^8} dx$$

$$= \frac{s+x}{6a(a+b(s+x)^2)^3} + \frac{5(s+x)}{24a^2(a+b(s+x)^2)^2}$$

$$+ \frac{5(s+x)}{16a^3(a+b(s+x)^2)} + \frac{5 \arctan\left(\frac{\sqrt{b}(s+x)}{\sqrt{a}}\right)}{16a^{7/2}\sqrt{b}}$$

output

```
1/6*(s+x)/a/(a+b*(s+x)^2)^3+5/24*(s+x)/a^2/(a+b*(s+x)^2)^2+5/16*(s+x)/a^3/
(a+b*(s+x)^2)+5/16*arctan(b^(1/2)*(s+x)/a^(1/2))/a^(7/2)/b^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.82

$$\int \frac{1}{a^4 + 4a^3b(s+x)^2 + 6a^2b^2(s+x)^4 + 4ab^3(s+x)^6 + b^4(s+x)^8} dx$$

$$= \frac{(s+x)(33a^2 + 40ab(s+x)^2 + 15b^2(s+x)^4)}{48a^3(a+b(s+x)^2)^3} + \frac{5 \arctan\left(\frac{\sqrt{b}(s+x)}{\sqrt{a}}\right)}{16a^{7/2}\sqrt{b}}$$

input

```
Integrate[(a^4 + 4*a^3*b*(s + x)^2 + 6*a^2*b^2*(s + x)^4 + 4*a*b^3*(s + x)^6 + b^4*(s + x)^8)^(-1),x]
```

output

```
((s + x)*(33*a^2 + 40*a*b*(s + x)^2 + 15*b^2*(s + x)^4))/(48*a^3*(a + b*(s + x)^2)^3) + (5*ArcTan[(Sqrt[b]*(s + x))/Sqrt[a]])/(16*a^(7/2)*Sqrt[b])
```

## Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 311 vs. 2(93) = 186.

Time = 0.80 (sec) , antiderivative size = 311, normalized size of antiderivative = 3.34, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$ , Rules used = {2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{a^4 + 4a^3b(s+x)^2 + 6a^2b^2(s+x)^4 + 4ab^3(s+x)^6 + b^4(s+x)^8} dx$$

↓ 2462

$$\int \left( -\frac{5b}{8a^3(2\sqrt{-a}\sqrt{b} - 2bs - 2bx)^2} - \frac{5b}{8a^3(2\sqrt{-a}\sqrt{b} + 2bs + 2bx)^2} + \frac{b^2}{a^2(2\sqrt{-a}\sqrt{b} - 2bs - 2bx)^4} + \frac{b^2}{a^2(2\sqrt{-a}\sqrt{b} + 2bs + 2bx)^4} \right) dx$$

↓ 2009

$$\begin{aligned} & -\frac{5}{32a^3\sqrt{b}(\sqrt{-a} - \sqrt{bs} - \sqrt{bx})} + \frac{5}{32a^3\sqrt{b}(\sqrt{-a} + \sqrt{bs} + \sqrt{bx})} + \\ & \frac{1}{48a^2\sqrt{b}(\sqrt{-a} - \sqrt{bs} - \sqrt{bx})^3} - \frac{1}{48a^2\sqrt{b}(\sqrt{-a} + \sqrt{bs} + \sqrt{bx})^3} + \\ & \frac{1}{16(-a)^{5/2}\sqrt{b}(\sqrt{-a} - \sqrt{bs} - \sqrt{bx})^2} - \frac{1}{16(-a)^{5/2}\sqrt{b}(\sqrt{-a} + \sqrt{bs} + \sqrt{bx})^2} - \\ & \frac{5 \log(\sqrt{-a} - \sqrt{bs} - \sqrt{bx})}{32(-a)^{7/2}\sqrt{b}} + \frac{5 \log(\sqrt{-a} + \sqrt{bs} + \sqrt{bx})}{32(-a)^{7/2}\sqrt{b}} \end{aligned}$$

input `Int[(a^4 + 4*a^3*b*(s + x)^2 + 6*a^2*b^2*(s + x)^4 + 4*a*b^3*(s + x)^6 + b^4*(s + x)^8)^(-1),x]`

output `1/(48*a^2*Sqrt[b]*(Sqrt[-a] - Sqrt[b]*s - Sqrt[b]*x)^3) + 1/(16*(-a)^(5/2)*Sqrt[b]*(Sqrt[-a] - Sqrt[b]*s - Sqrt[b]*x)^2) - 5/(32*a^3*Sqrt[b]*(Sqrt[-a] - Sqrt[b]*s - Sqrt[b]*x)) - 1/(48*a^2*Sqrt[b]*(Sqrt[-a] + Sqrt[b]*s + Sqrt[b]*x)^3) - 1/(16*(-a)^(5/2)*Sqrt[b]*(Sqrt[-a] + Sqrt[b]*s + Sqrt[b]*x)^2) + 5/(32*a^3*Sqrt[b]*(Sqrt[-a] + Sqrt[b]*s + Sqrt[b]*x)) - (5*Log[Sqrt[-a] - Sqrt[b]*s - Sqrt[b]*x])/(32*(-a)^(7/2)*Sqrt[b]) + (5*Log[Sqrt[-a] + Sqrt[b]*s + Sqrt[b]*x])/(32*(-a)^(7/2)*Sqrt[b])`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2462 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegr and[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]]] /; PolyQ[Px, x] && GtQ [Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0 ] && RationalFunctionQ[u, x]`

**Maple [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.60

method	result
default	$\frac{2bs+2bx}{12ab(bs^2+2bsx+bx^2+a)^3} + \frac{5(2bs+2bx)}{48ab(bs^2+2bsx+bx^2+a)^2} + \frac{5 \left( \frac{3(2bs+2bx)}{16ab(bs^2+2bsx+bx^2+a)} + \frac{3 \arctan\left(\frac{2bs+2bx}{2\sqrt{ab}}\right)}{8a\sqrt{ab}} \right)}{6a}$
risch	$\frac{5b^2x^5}{16a^3} + \frac{25b^2sx^4}{16a^3} + \frac{5b(15bs^2+4a)x^3}{24a^3} + \frac{5bs(5bs^2+4a)x^2}{8a^3} + \frac{(25b^2s^4+40abs^2+11a^2)x}{16a^3} + \frac{s(15b^2s^4+40abs^2+33a^2)}{48a^3}$ $b^3s^6+6b^3s^5x+15b^3s^4x^2+20b^3s^3x^3+15b^3s^2x^4+6b^3sx^5+b^3x^6+3ab^2s^4+12ab^2s^3x+18ab^2s^2x^2+12ab^2sx^3+3ab^2x^4+3a^2bs^2+6a^2bx+3a^2$

input `int(1/(a^4+4*a^3*b*(s+x)^2+6*a^2*b^2*(s+x)^4+4*a*b^3*(s+x)^6+b^4*(s+x)^8), x,method=_RETURNVERBOSE)`



output

```
1/12*(2*b*s+2*b*x)/a/b/(b*s^2+2*b*s*x+b*x^2+a)^3+5/6/a*(1/8*(2*b*s+2*b*x)/
a/b/(b*s^2+2*b*s*x+b*x^2+a)^2+3/4/a*(1/4*(2*b*s+2*b*x)/a/b/(b*s^2+2*b*s*x+
b*x^2+a)+1/2/a/(a*b)^(1/2)*arctan(1/2*(2*b*s+2*b*x)/(a*b)^(1/2)))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 465 vs.  $2(77) = 154$ .

Time = 0.10 (sec) , antiderivative size = 970, normalized size of antiderivative = 10.43

$$\int \frac{1}{a^4 + 4a^3b(s+x)^2 + 6a^2b^2(s+x)^4 + 4ab^3(s+x)^6 + b^4(s+x)^8} dx$$

= Too large to display

input

```
integrate(1/(a^4+4*a^3*b*(s+x)^2+6*a^2*b^2*(s+x)^4+4*a*b^3*(s+x)^6+b^4*(s+
x)^8),x, algorithm="fricas")
```

output

```
[1/96*(30*a*b^3*s^5 + 150*a*b^3*s*x^4 + 30*a*b^3*x^5 + 80*a^2*b^2*s^3 + 66
*a^3*b*s + 20*(15*a*b^3*s^2 + 4*a^2*b^2)*x^3 + 60*(5*a*b^3*s^3 + 4*a^2*b^2
*s)*x^2 - 15*(b^3*s^6 + 6*b^3*s*x^5 + b^3*x^6 + 3*a*b^2*s^4 + 3*a^2*b*s^2
+ 3*(5*b^3*s^2 + a*b^2)*x^4 + 4*(5*b^3*s^3 + 3*a*b^2*s)*x^3 + a^3 + 3*(5*b
^3*s^4 + 6*a*b^2*s^2 + a^2*b)*x^2 + 6*(b^3*s^5 + 2*a*b^2*s^3 + a^2*b*s)*x)
*sqrt(-a*b)*log((b*s^2 + 2*b*s*x + b*x^2 - 2*sqrt(-a*b)*(s + x) - a)/(b*s^
2 + 2*b*s*x + b*x^2 + a)) + 6*(25*a*b^3*s^4 + 40*a^2*b^2*s^2 + 11*a^3*b)*x
)/(a^4*b^4*s^6 + 6*a^4*b^4*s*x^5 + a^4*b^4*x^6 + 3*a^5*b^3*s^4 + 3*a^6*b^2
*s^2 + a^7*b + 3*(5*a^4*b^4*s^2 + a^5*b^3)*x^4 + 4*(5*a^4*b^4*s^3 + 3*a^5*
b^3*s)*x^3 + 3*(5*a^4*b^4*s^4 + 6*a^5*b^3*s^2 + a^6*b^2)*x^2 + 6*(a^4*b^4*
s^5 + 2*a^5*b^3*s^3 + a^6*b^2*s)*x), 1/48*(15*a*b^3*s^5 + 75*a*b^3*s*x^4 +
15*a*b^3*x^5 + 40*a^2*b^2*s^3 + 33*a^3*b*s + 10*(15*a*b^3*s^2 + 4*a^2*b^2
)*x^3 + 30*(5*a*b^3*s^3 + 4*a^2*b^2*s)*x^2 + 15*(b^3*s^6 + 6*b^3*s*x^5 + b
^3*x^6 + 3*a*b^2*s^4 + 3*a^2*b*s^2 + 3*(5*b^3*s^2 + a*b^2)*x^4 + 4*(5*b^3*
s^3 + 3*a*b^2*s)*x^3 + a^3 + 3*(5*b^3*s^4 + 6*a*b^2*s^2 + a^2*b)*x^2 + 6*(
b^3*s^5 + 2*a*b^2*s^3 + a^2*b*s)*x)*sqrt(a*b)*arctan(sqrt(a*b)*(s + x)/a)
+ 3*(25*a*b^3*s^4 + 40*a^2*b^2*s^2 + 11*a^3*b)*x)/(a^4*b^4*s^6 + 6*a^4*b^4
*s*x^5 + a^4*b^4*x^6 + 3*a^5*b^3*s^4 + 3*a^6*b^2*s^2 + a^7*b + 3*(5*a^4*b^
4*s^2 + a^5*b^3)*x^4 + 4*(5*a^4*b^4*s^3 + 3*a^5*b^3*s)*x^3 + 3*(5*a^4*b^4*
s^4 + 6*a^5*b^3*s^2 + a^6*b^2)*x^2 + 6*(a^4*b^4*s^5 + 2*a^5*b^3*s^3 + a...
```

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 354 vs.  $2(92) = 184$ .

Time = 0.93 (sec) , antiderivative size = 354, normalized size of antiderivative = 3.81

$$\int \frac{1}{a^4 + 4a^3b(s+x)^2 + 6a^2b^2(s+x)^4 + 4ab^3(s+x)^6 + b^4(s+x)^8} dx$$

$$= -\frac{5\sqrt{-\frac{1}{a^7b}} \log\left(-a^4\sqrt{-\frac{1}{a^7b}} + s+x\right)}{32} + \frac{5\sqrt{-\frac{1}{a^7b}} \log\left(a^4\sqrt{-\frac{1}{a^7b}} + s+x\right)}{32}$$

$$+ \frac{33a^2s + 40abs^3 + 15b^2s^5 + 75b^2sx^4 + 15b^2x^5 + x^3 \cdot (40ab + 15b^2)}{48a^6 + 144a^5bs^2 + 144a^4b^2s^4 + 48a^3b^3s^6 + 288a^3b^3sx^5 + 48a^3b^3x^6 + x^4 \cdot (144a^4b^2 + 720a^3b^3s^2) + x^3}$$

input `integrate(1/(a**4+4*a**3*b*(s+x)**2+6*a**2*b**2*(s+x)**4+4*a*b**3*(s+x)**6+b**4*(s+x)**8), x)`

output `-5*sqrt(-1/(a**7*b))*log(-a**4*sqrt(-1/(a**7*b)) + s + x)/32 + 5*sqrt(-1/(a**7*b))*log(a**4*sqrt(-1/(a**7*b)) + s + x)/32 + (33*a**2*s + 40*a*b*s**3 + 15*b**2*s**5 + 75*b**2*s*x**4 + 15*b**2*x**5 + x**3*(40*a*b + 15*b**2*s**2) + x**2*(120*a*b*s + 150*b**2*s**3) + x*(33*a**2 + 120*a*b*s**2 + 75*b**2*s**4))/(48*a**6 + 144*a**5*b*s**2 + 144*a**4*b**2*s**4 + 48*a**3*b**3*s**6 + 288*a**3*b**3*s*x**5 + 48*a**3*b**3*x**6 + x**4*(144*a**4*b**2 + 720*a**3*b**3*s**2) + x**3*(576*a**4*b**2*s + 960*a**3*b**3*s**3) + x**2*(144*a**5*b + 864*a**4*b**2*s**2 + 720*a**3*b**3*s**4) + x*(288*a**5*b*s + 76*a**4*b**2*s**3 + 288*a**3*b**3*s**5))`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 299 vs.  $2(77) = 154$ .

Time = 0.12 (sec) , antiderivative size = 299, normalized size of antiderivative = 3.22

$$\int \frac{1}{a^4 + 4a^3b(s+x)^2 + 6a^2b^2(s+x)^4 + 4ab^3(s+x)^6 + b^4(s+x)^8} dx$$

$$= \frac{15b^2s^5 + 75b^2sx^4 + 15b^2x^5 + 40abs^3 + 10(15b^2s^2 + 4ab)x^3 + 33a^2s + 30(5b^2s^2 + 4ab^2)}{48(a^3b^3s^6 + 6a^3b^3sx^5 + a^3b^3x^6 + 3a^4b^2s^4 + 3a^5bs^2 + a^6 + 3(5a^3b^3s^2 + a^4b^2)x^4 + 4(5a^3b^3s^3 + 3a^4b^2s^2 + 3a^5bs^2 + a^6))} + \frac{5 \arctan\left(\frac{bs+bx}{\sqrt{ab}}\right)}{16\sqrt{aba^3}}$$

input `integrate(1/(a^4+4*a^3*b*(s+x)^2+6*a^2*b^2*(s+x)^4+4*a*b^3*(s+x)^6+b^4*(s+x)^8),x, algorithm="maxima")`

output 
$$\frac{1}{48} \cdot (15b^2s^5 + 75b^2s^4x + 15b^2x^5 + 40ab^3s^3 + 10(15b^2s^2 + 4ab^2)sx^3 + 33a^2s + 30(5b^2s^3 + 4ab^2s)x^2 + 3(25b^2s^4 + 40ab^2s^2 + 11a^2)x) / (a^3b^3s^6 + 6a^3b^3sx^5 + a^3b^3x^6 + 3a^4b^2s^4 + 3a^4b^2sx^2 + a^4 + 3(5a^3b^3s^2 + a^4b^2)x^4 + 4(5a^3b^3s^3 + 3a^4b^2s)x^3 + 3(5a^3b^3s^4 + 6a^4b^2s^2 + a^5b)x^2 + 6(a^3b^3s^5 + 2a^4b^2s^3 + a^5b)s)x + 5/16 \arctan((bs + bx)/\sqrt{ab}) / (\sqrt{ab})a^3$$

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.59

$$\int \frac{1}{a^4 + 4a^3b(s+x)^2 + 6a^2b^2(s+x)^4 + 4ab^3(s+x)^6 + b^4(s+x)^8} dx = \frac{5 \arctan\left(\frac{bs+bx}{\sqrt{ab}}\right)}{16 \sqrt{ab} a^3} + \frac{15b^2s^5 + 75b^2s^4x + 150b^2s^3x^2 + 150b^2s^2x^3 + 75b^2sx^4 + 15b^2x^5 + 40abs^3 + 120abs^2x + 120absx^2 - 48(bs^2 + 2bsx + bx^2 + a)^3 a^3}{48 (bs^2 + 2bsx + bx^2 + a)^3 a^3}$$

input `integrate(1/(a^4+4*a^3*b*(s+x)^2+6*a^2*b^2*(s+x)^4+4*a*b^3*(s+x)^6+b^4*(s+x)^8),x, algorithm="giac")`

output 
$$5/16 \arctan((bs + bx)/\sqrt{ab}) / (\sqrt{ab})a^3 + 1/48 \cdot (15b^2s^5 + 75b^2s^4x + 150b^2s^3x^2 + 150b^2s^2x^3 + 75b^2s^2x^4 + 15b^2x^5 + 40ab^3s^3 + 120ab^2s^2x + 120ab^2sx^2 + 40ab^2x^3 + 33a^2s + 33a^2sx) / ((b^2s^2 + 2b^2sx + b^2x^2 + a)^3 a^3)$$

**Mupad [B] (verification not implemented)**

Time = 21.79 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.98

$$\int \frac{1}{a^4 + 4a^3b(s+x)^2 + 6a^2b^2(s+x)^4 + 4ab^3(s+x)^6 + b^4(s+x)^8} dx$$

$$= \frac{\frac{11(s+x)}{16a} + \frac{5b(s+x)^3}{6a^2} + \frac{5b^2(s+x)^5}{16a^3}}{a^3 + b^3(s+x)^6 + 3a^2b(s+x)^2 + 3ab^2(s+x)^4} + \frac{5 \operatorname{atan}\left(\frac{\sqrt{b}(s+x)}{\sqrt{a}}\right)}{16a^{7/2}\sqrt{b}}$$

input

```
int(1/(a^4 + b^4*(s + x)^8 + 4*a^3*b*(s + x)^2 + 4*a*b^3*(s + x)^6 + 6*a^2
*b^2*(s + x)^4), x)
```

output

```
((11*(s + x))/(16*a) + (5*b*(s + x)^3)/(6*a^2) + (5*b^2*(s + x)^5)/(16*a^3
))/ (a^3 + b^3*(s + x)^6 + 3*a^2*b*(s + x)^2 + 3*a*b^2*(s + x)^4) + (5*atan
((b^(1/2)*(s + x))/a^(1/2)))/(16*a^(7/2)*b^(1/2))
```

**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 808, normalized size of antiderivative = 8.69

$$\int \frac{1}{a^4 + 4a^3b(s+x)^2 + 6a^2b^2(s+x)^4 + 4ab^3(s+x)^6 + b^4(s+x)^8} dx$$

= Too large to display

input

```
int(1/(a^4+4*a^3*b*(s+x)^2+6*a^2*b^2*(s+x)^4+4*a*b^3*(s+x)^6+b^4*(s+x)^8),
x)
```

output

```
(30*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**3*s + 90*sqrt(b)
)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**2*b*s**3 + 180*sqrt(b)*sq
rt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**2*b*s**2*x + 90*sqrt(b)*sqrt(
a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**2*b*s*x**2 + 90*sqrt(b)*sqrt(a)*
atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**5 + 360*sqrt(b)*sqrt(a)*atan
((b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**4*x + 540*sqrt(b)*sqrt(a)*atan((
b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**3*x**2 + 360*sqrt(b)*sqrt(a)*atan(
(b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**2*x**3 + 90*sqrt(b)*sqrt(a)*atan(
(b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s*x**4 + 30*sqrt(b)*sqrt(a)*atan((b*
s + b*x)/(sqrt(b)*sqrt(a)))*b**3*s**7 + 180*sqrt(b)*sqrt(a)*atan((b*s + b*
x)/(sqrt(b)*sqrt(a)))*b**3*s**6*x + 450*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(
sqrt(b)*sqrt(a)))*b**3*s**5*x**2 + 600*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(s
qrt(b)*sqrt(a)))*b**3*s**4*x**3 + 450*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sq
rt(b)*sqrt(a)))*b**3*s**3*x**4 + 180*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqr
t(b)*sqrt(a)))*b**3*s**2*x**5 + 30*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(
b)*sqrt(a)))*b**3*s*x**6 - 5*a**4 + 51*a**3*b*s**2 + 36*a**3*b*s*x - 15*a*
*3*b*x**2 + 65*a**2*b**2*s**4 + 180*a**2*b**2*s**3*x + 150*a**2*b**2*s**2*
x**2 + 20*a**2*b**2*s*x**3 - 15*a**2*b**2*x**4 + 25*a*b**3*s**6 + 120*a*b*
*3*s**5*x + 225*a*b**3*s**4*x**2 + 200*a*b**3*s**3*x**3 + 75*a*b**3*s**2*x
**4 - 5*a*b**3*x**6)/(96*a**4*b*s*(a**3 + 3*a**2*b*s**2 + 6*a**2*b*s*x ...
```

### 3.36 $\int \frac{1}{(a+bs^2+2bsx+bx^2)^4} dx$

Optimal result	285
Mathematica [A] (verified)	286
Rubi [A] (verified)	286
Maple [A] (verified)	288
Fricas [B] (verification not implemented)	288
Sympy [B] (verification not implemented)	289
Maxima [B] (verification not implemented)	290
Giac [A] (verification not implemented)	291
Mupad [B] (verification not implemented)	291
Reduce [B] (verification not implemented)	292

#### Optimal result

Integrand size = 19, antiderivative size = 117

$$\int \frac{1}{(a + bs^2 + 2bsx + bx^2)^4} dx = \frac{s + x}{6a(a + bs^2 + 2bsx + bx^2)^3} + \frac{5(s + x)}{24a^2(a + bs^2 + 2bsx + bx^2)^2} + \frac{5(s + x)}{16a^3(a + bs^2 + 2bsx + bx^2)} + \frac{5 \arctan\left(\frac{\sqrt{b}(s+x)}{\sqrt{a}}\right)}{16a^{7/2}\sqrt{b}}$$

output

```
1/6*(s+x)/a/(b*s^2+2*b*s*x+b*x^2+a)^3+5/24*(s+x)/a^2/(b*s^2+2*b*s*x+b*x^2+a)^2+5/16*(s+x)/a^3/(b*s^2+2*b*s*x+b*x^2+a)+5/16*arctan(b^(1/2)*(s+x)/a^(1/2))/a^(7/2)/b^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.65

$$\int \frac{1}{(a + bs^2 + 2bsx + bx^2)^4} dx = \frac{(s + x)(33a^2 + 40ab(s + x)^2 + 15b^2(s + x)^4)}{48a^3(a + b(s + x)^2)^3} + \frac{5 \arctan\left(\frac{\sqrt{b}(s+x)}{\sqrt{a}}\right)}{16a^{7/2}\sqrt{b}}$$

input `Integrate[(a + b*s^2 + 2*b*s*x + b*x^2)^(-4), x]`

output `((s + x)*(33*a^2 + 40*a*b*(s + x)^2 + 15*b^2*(s + x)^4)/(48*a^3*(a + b*(s + x)^2)^3) + (5*ArcTan[(Sqrt[b]*(s + x))/Sqrt[a]])/(16*a^(7/2)*Sqrt[b])`

**Rubi [A] (verified)**

Time = 0.45 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.21, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {1086, 1086, 1086, 1083, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(a + bs^2 + 2bsx + bx^2)^4} dx \\ & \quad \downarrow 1086 \\ & \frac{5 \int \frac{1}{(bs^2 + 2bsx + bx^2 + a)^3} dx}{6a} + \frac{s + x}{6a(a + bs^2 + 2bsx + bx^2)^3} \\ & \quad \downarrow 1086 \\ & \frac{5 \left( \frac{3 \int \frac{1}{(bs^2 + 2bsx + bx^2 + a)^2} dx}{4a} + \frac{s + x}{4a(a + bs^2 + 2bsx + bx^2)^2} \right)}{6a} + \frac{s + x}{6a(a + bs^2 + 2bsx + bx^2)^3} \\ & \quad \downarrow 1086 \end{aligned}$$

$$\begin{aligned}
 & \frac{5 \left( \frac{3 \left( \frac{\int \frac{1}{bs^2+2bsx+bx^2+a} dx}{2a} + \frac{s+x}{2a(a+bs^2+2bsx+bx^2)} \right)}{4a} + \frac{s+x}{4a(a+bs^2+2bsx+bx^2)^2} \right)}{\frac{6a}{s+x}} + \\
 & \frac{6a}{6a(a+bs^2+2bsx+bx^2)^3} \\
 & \quad \downarrow \text{1083} \\
 & \frac{5 \left( \frac{3 \left( \frac{s+x}{2a(a+bs^2+2bsx+bx^2)} - \frac{\int \frac{1}{-(2bs+2bx)^2-4ab} d(2bs+2bx)}{a} \right)}{4a} + \frac{s+x}{4a(a+bs^2+2bsx+bx^2)^2} \right)}{\frac{6a}{s+x}} + \\
 & \frac{6a}{6a(a+bs^2+2bsx+bx^2)^3} \\
 & \quad \downarrow \text{217} \\
 & \frac{5 \left( \frac{3 \left( \frac{\arctan\left(\frac{2bs+2bx}{2\sqrt{a}\sqrt{b}}\right)}{2a^{3/2}\sqrt{b}} + \frac{s+x}{2a(a+bs^2+2bsx+bx^2)} \right)}{4a} + \frac{s+x}{4a(a+bs^2+2bsx+bx^2)^2} \right)}{6a} + \frac{s+x}{6a(a+bs^2+2bsx+bx^2)^3}
 \end{aligned}$$

input `Int[(a + b*s^2 + 2*b*s*x + b*x^2)^(-4), x]`

output `(s + x)/(6*a*(a + b*s^2 + 2*b*s*x + b*x^2)^3) + (5*((s + x)/(4*a*(a + b*s^2 + 2*b*s*x + b*x^2)^2) + (3*((s + x)/(2*a*(a + b*s^2 + 2*b*s*x + b*x^2)) + ArcTan[(2*b*s + 2*b*x)/(2*Sqrt[a]*Sqrt[b])]/(2*a^(3/2)*Sqrt[b])))/(4*a)))/(6*a)`

**Defintions of rubi rules used**

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])`



```
rule 1083 Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]
```

```
rule 1086 Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c))), x] - Simp[2*c*((2*p + 3)/((p + 1)*(b^2 - 4*a*c))) Int[(a + b*x + c*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && ILtQ[p, -1]
```

### Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.27

method	result
default	$\frac{2bs+2bx}{12ab(b^2s^2+2bsx+bx^2+a)^3} + \frac{5(2bs+2bx)}{48ab(b^2s^2+2bsx+bx^2+a)^2} + \frac{5\left(\frac{3(2bs+2bx)}{16ab(b^2s^2+2bsx+bx^2+a)} + \frac{3\arctan\left(\frac{2bs+2bx}{2\sqrt{ab}}\right)}{8a\sqrt{ab}}\right)}{6a}$
risch	$\frac{5b^2x^5}{16a^3} + \frac{25b^2sx^4}{16a^3} + \frac{5b(15bs^2+4a)x^3}{24a^3} + \frac{5bs(5bs^2+4a)x^2}{8a^3} + \frac{(25b^2s^4+40abs^2+11a^2)x}{16a^3} + \frac{s(15b^2s^4+40abs^2+33a^2)}{48a^3} - \frac{5\ln(bs+bx+\sqrt{-ab})}{32\sqrt{-ab}a^3}$

```
input int(1/(b*s^2+2*b*s*x+b*x^2+a)^4,x,method=_RETURNVERBOSE)
```

```
output 1/12*(2*b*s+2*b*x)/a/b/(b*s^2+2*b*s*x+b*x^2+a)^3+5/6/a*(1/8*(2*b*s+2*b*x)/a/b/(b*s^2+2*b*s*x+b*x^2+a)^2+3/4/a*(1/4*(2*b*s+2*b*x)/a/b/(b*s^2+2*b*s*x+b*x^2+a)+1/2/a/(a*b)^(1/2)*arctan(1/2*(2*b*s+2*b*x)/(a*b)^(1/2)))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 465 vs. 2(101) = 202.

Time = 0.11 (sec) , antiderivative size = 970, normalized size of antiderivative = 8.29

$$\int \frac{1}{(a + bs^2 + 2bsx + bx^2)^4} dx = \text{Too large to display}$$

input `integrate(1/(b*s^2+2*b*s*x+b*x^2+a)^4,x, algorithm="fricas")`

output

```
[1/96*(30*a*b^3*s^5 + 150*a*b^3*s*x^4 + 30*a*b^3*x^5 + 80*a^2*b^2*s^3 + 66
*a^3*b*s + 20*(15*a*b^3*s^2 + 4*a^2*b^2)*x^3 + 60*(5*a*b^3*s^3 + 4*a^2*b^2
*s)*x^2 - 15*(b^3*s^6 + 6*b^3*s*x^5 + b^3*x^6 + 3*a*b^2*s^4 + 3*a^2*b*s^2
+ 3*(5*b^3*s^2 + a*b^2)*x^4 + 4*(5*b^3*s^3 + 3*a*b^2*s)*x^3 + a^3 + 3*(5*b
^3*s^4 + 6*a*b^2*s^2 + a^2*b)*x^2 + 6*(b^3*s^5 + 2*a*b^2*s^3 + a^2*b*s)*x)
*sqrt(-a*b)*log((b*s^2 + 2*b*s*x + b*x^2 - 2*sqrt(-a*b)*(s + x) - a)/(b*s^
2 + 2*b*s*x + b*x^2 + a)) + 6*(25*a*b^3*s^4 + 40*a^2*b^2*s^2 + 11*a^3*b)*x
)/(a^4*b^4*s^6 + 6*a^4*b^4*s*x^5 + a^4*b^4*x^6 + 3*a^5*b^3*s^4 + 3*a^6*b^2
*s^2 + a^7*b + 3*(5*a^4*b^4*s^2 + a^5*b^3)*x^4 + 4*(5*a^4*b^4*s^3 + 3*a^5*
b^3*s)*x^3 + 3*(5*a^4*b^4*s^4 + 6*a^5*b^3*s^2 + a^6*b^2)*x^2 + 6*(a^4*b^4*
s^5 + 2*a^5*b^3*s^3 + a^6*b^2*s)*x), 1/48*(15*a*b^3*s^5 + 75*a*b^3*s*x^4 +
15*a*b^3*x^5 + 40*a^2*b^2*s^3 + 33*a^3*b*s + 10*(15*a*b^3*s^2 + 4*a^2*b^2
)*x^3 + 30*(5*a*b^3*s^3 + 4*a^2*b^2*s)*x^2 + 15*(b^3*s^6 + 6*b^3*s*x^5 + b
^3*x^6 + 3*a*b^2*s^4 + 3*a^2*b*s^2 + 3*(5*b^3*s^2 + a*b^2)*x^4 + 4*(5*b^3*
s^3 + 3*a*b^2*s)*x^3 + a^3 + 3*(5*b^3*s^4 + 6*a*b^2*s^2 + a^2*b)*x^2 + 6*(
b^3*s^5 + 2*a*b^2*s^3 + a^2*b*s)*x)*sqrt(a*b)*arctan(sqrt(a*b)*(s + x)/a)
+ 3*(25*a*b^3*s^4 + 40*a^2*b^2*s^2 + 11*a^3*b)*x)/(a^4*b^4*s^6 + 6*a^4*b^4
*s*x^5 + a^4*b^4*x^6 + 3*a^5*b^3*s^4 + 3*a^6*b^2*s^2 + a^7*b + 3*(5*a^4*b^
4*s^2 + a^5*b^3)*x^4 + 4*(5*a^4*b^4*s^3 + 3*a^5*b^3*s)*x^3 + 3*(5*a^4*b^4*
s^4 + 6*a^5*b^3*s^2 + a^6*b^2)*x^2 + 6*(a^4*b^4*s^5 + 2*a^5*b^3*s^3 + a...
```

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 354 vs.  $2(122) = 244$ .

Time = 0.96 (sec) , antiderivative size = 354, normalized size of antiderivative = 3.03

$$\int \frac{1}{(a + bs^2 + 2bsx + bx^2)^4} dx$$

$$= -\frac{5\sqrt{-\frac{1}{a^7b}} \log\left(-a^4\sqrt{-\frac{1}{a^7b}} + s + x\right)}{32} + \frac{5\sqrt{-\frac{1}{a^7b}} \log\left(a^4\sqrt{-\frac{1}{a^7b}} + s + x\right)}{32}$$

$$+ \frac{33a^2s + 40abs^3 + 15b^2s^5 + 75b^2sx^4 + 15b^2x^5 + x^3 \cdot (40ab + 1}{48a^6 + 144a^5bs^2 + 144a^4b^2s^4 + 48a^3b^3s^6 + 288a^3b^3sx^5 + 48a^3b^3x^6 + x^4 \cdot (144a^4b^2 + 720a^3b^3s^2) + x^3}$$

input `integrate(1/(b*s**2+2*b*s*x+b*x**2+a)**4,x)`

output

```
-5*sqrt(-1/(a**7*b))*log(-a**4*sqrt(-1/(a**7*b)) + s + x)/32 + 5*sqrt(-1/(
a**7*b))*log(a**4*sqrt(-1/(a**7*b)) + s + x)/32 + (33*a**2*s + 40*a*b*s**3
+ 15*b**2*s**5 + 75*b**2*s*x**4 + 15*b**2*x**5 + x**3*(40*a*b + 150*b**2*
s**2) + x**2*(120*a*b*s + 150*b**2*s**3) + x*(33*a**2 + 120*a*b*s**2 + 75*
b**2*s**4))/(48*a**6 + 144*a**5*b*s**2 + 144*a**4*b**2*s**4 + 48*a**3*b**3
*s**6 + 288*a**3*b**3*s*x**5 + 48*a**3*b**3*x**6 + x**4*(144*a**4*b**2 + 7
20*a**3*b**3*s**2) + x**3*(576*a**4*b**2*s + 960*a**3*b**3*s**3) + x**2*(1
44*a**5*b + 864*a**4*b**2*s**2 + 720*a**3*b**3*s**4) + x*(288*a**5*b*s + 5
76*a**4*b**2*s**3 + 288*a**3*b**3*s**5))
```

### Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 299 vs.  $2(101) = 202$ .

Time = 0.12 (sec) , antiderivative size = 299, normalized size of antiderivative = 2.56

$$\int \frac{1}{(a + bs^2 + 2bsx + bx^2)^4} dx$$

$$= \frac{15b^2s^5 + 75b^2sx^4 + 15b^2x^5 + 40abs^3 + 10(15b^2s^2 + 4ab)x^3 + 33a^2s + 30(5b^2s^2 + 4a^2b^2)x + 33a^2s + 30(5b^2s^2 + 4a^2b^2)}{48(a^3b^3s^6 + 6a^3b^3sx^5 + a^3b^3x^6 + 3a^4b^2s^4 + 3a^5bs^2 + a^6 + 3(5a^3b^3s^2 + a^4b^2)x^4 + 4(5a^3b^3s^3 + 3a^4b^2s^2 + 3a^5bs^2 + a^6))} + \frac{5 \arctan\left(\frac{bs+bx}{\sqrt{ab}}\right)}{16\sqrt{aba^3}}$$

input

```
integrate(1/(b*s^2+2*b*s*x+b*x^2+a)^4,x, algorithm="maxima")
```

output

```
1/48*(15*b^2*s^5 + 75*b^2*s*x^4 + 15*b^2*x^5 + 40*a*b*s^3 + 10*(15*b^2*s^2
+ 4*a*b)*x^3 + 33*a^2*s + 30*(5*b^2*s^3 + 4*a*b*s)*x^2 + 3*(25*b^2*s^4 +
40*a*b*s^2 + 11*a^2)*x)/(a^3*b^3*s^6 + 6*a^3*b^3*s*x^5 + a^3*b^3*x^6 + 3*a
^4*b^2*s^4 + 3*a^5*b*s^2 + a^6 + 3*(5*a^3*b^3*s^2 + a^4*b^2)*x^4 + 4*(5*a
^3*b^3*s^3 + 3*a^4*b^2*s)*x^3 + 3*(5*a^3*b^3*s^4 + 6*a^4*b^2*s^2 + a^5*b)*x
^2 + 6*(a^3*b^3*s^5 + 2*a^4*b^2*s^3 + a^5*b*s)*x) + 5/16*arctan((b*s + b*x
)/sqrt(a*b))/(sqrt(a*b)*a^3)
```

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.26

$$\int \frac{1}{(a + bs^2 + 2bsx + bx^2)^4} dx = \frac{5 \arctan\left(\frac{bs+bx}{\sqrt{ab}}\right)}{16 \sqrt{aba^3}} + \frac{15b^2s^5 + 75b^2s^4x + 150b^2s^3x^2 + 150b^2s^2x^3 + 75b^2sx^4 + 15b^2x^5 + 40abs^3 + 120abs^2x + 120absx^2 + 40a^2s^3 + 120a^2s^2x + 120a^2sx^2 + 40a^2x^3 + 33a^2s + 33a^2x}{48(bs^2 + 2bsx + bx^2 + a)^3 a^3}$$

input `integrate(1/(b*s^2+2*b*s*x+b*x^2+a)^4,x, algorithm="giac")`output `5/16*arctan((b*s + b*x)/sqrt(a*b))/(sqrt(a*b)*a^3) + 1/48*(15*b^2*s^5 + 75*b^2*s^4*x + 150*b^2*s^3*x^2 + 150*b^2*s^2*x^3 + 75*b^2*s*x^4 + 15*b^2*x^5 + 40*a*b*s^3 + 120*a*b*s^2*x + 120*a*b*s*x^2 + 40*a*b*x^3 + 33*a^2*s + 33*a^2*x)/((b*s^2 + 2*b*s*x + b*x^2 + a)^3*a^3)`**Mupad [B] (verification not implemented)**

Time = 22.06 (sec) , antiderivative size = 297, normalized size of antiderivative = 2.54

$$\int \frac{1}{(a + bs^2 + 2bsx + bx^2)^4} dx = \frac{33a^2s + 40abs^3 + 15b^2s^5}{48a^3} + \frac{5x^3(15b^2s^2 + 4ab)}{24a^3} + \frac{5b^2x^5}{16a^3} + \frac{5x^2(5b^2s^3 + 4abs)}{8a^3} + \frac{x(20b^3s^3 + 15b^2s^2 + 12abs + 3a^2b)}{16a^3} + \frac{5 \operatorname{atan}\left(\frac{16a^3\left(\frac{5\sqrt{b}s}{16a^{7/2}} + \frac{5\sqrt{b}x}{16a^{7/2}}\right)}{5}\right)}{16a^{7/2}\sqrt{b}}$$

input `int(1/(a + b*s^2 + b*x^2 + 2*b*s*x)^4,x)`

output

```
((33*a^2*s + 15*b^2*s^5 + 40*a*b*s^3)/(48*a^3) + (5*x^3*(4*a*b + 15*b^2*s^2))/(24*a^3) + (5*b^2*x^5)/(16*a^3) + (5*x^2*(5*b^2*s^3 + 4*a*b*s))/(8*a^3) + (x*(11*a^2 + 25*b^2*s^4 + 40*a*b*s^2))/(16*a^3) + (25*b^2*s*x^4)/(16*a^3))/(x^4*(3*a*b^2 + 15*b^3*s^2) + x^2*(3*a^2*b + 15*b^3*s^4 + 18*a*b^2*s^2) + x*(6*b^3*s^5 + 12*a*b^2*s^3 + 6*a^2*b*s) + x^3*(20*b^3*s^3 + 12*a*b^2*s) + a^3 + b^3*s^6 + b^3*x^6 + 3*a^2*b*s^2 + 3*a*b^2*s^4 + 6*b^3*s*x^5) + (5*atan((16*a^3*((5*b^(1/2)*s)/(16*a^(7/2)) + (5*b^(1/2)*x)/(16*a^(7/2))))/5))/(16*a^(7/2)*b^(1/2))
```

**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 808, normalized size of antiderivative = 6.91

$$\int \frac{1}{(a + bs^2 + 2bsx + bx^2)^4} dx = \text{Too large to display}$$

input

```
int(1/(b*s^2+2*b*s*x+b*x^2+a)^4,x)
```

output

```
(30*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**3*s + 90*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**2*b*s**3 + 180*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**2*b*s**2*x + 90*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**2*b*s*x**2 + 90*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**5 + 360*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**4*x + 540*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**3*x**2 + 360*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**2*x**3 + 90*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s*x**4 + 30*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*b**3*s**7 + 180*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*b**3*s**6*x + 450*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*b**3*s**5*x**2 + 600*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*b**3*s**4*x**3 + 450*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*b**3*s**3*x**4 + 180*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*b**3*s**2*x**5 + 30*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*b**3*s*x**6 - 5*a**4 + 51*a**3*b*s**2 + 36*a**3*b*s*x - 15*a**3*b*x**2 + 65*a**2*b**2*s**4 + 180*a**2*b**2*s**3*x + 150*a**2*b**2*s**2*x**2 + 20*a**2*b**2*s*x**3 - 15*a**2*b**2*x**4 + 25*a*b**3*s**6 + 120*a*b**3*s**5*x + 225*a*b**3*s**4*x**2 + 200*a*b**3*s**3*x**3 + 75*a*b**3*s**2*x**4 - 5*a*b**3*x**6)/(96*a**4*b*s*(a**3 + 3*a**2*b*s**2 + 6*a**2*b*s*x ...
```

**3.37** 
$$\int \frac{1}{\left( (a+bs^2)^2 + 4bs(a+bs^2)x + 2b(a+3bs^2)x^2 + 4b^2sx^3 + b^2x^4 \right)^2} dx$$

Optimal result . . . . .	293
Mathematica [A] (verified) . . . . .	293
Rubi [A] (verified) . . . . .	294
Maple [A] (verified) . . . . .	296
Fricas [B] (verification not implemented) . . . . .	297
Sympy [B] (verification not implemented) . . . . .	298
Maxima [B] (verification not implemented) . . . . .	298
Giac [A] (verification not implemented) . . . . .	299
Mupad [B] (verification not implemented) . . . . .	300
Reduce [B] (verification not implemented) . . . . .	300

**Optimal result**

Integrand size = 54, antiderivative size = 93

$$\int \frac{1}{\left( (a+bs^2)^2 + 4bs(a+bs^2)x + 2b(a+3bs^2)x^2 + 4b^2sx^3 + b^2x^4 \right)^2} dx$$

$$= \frac{s+x}{6a(a+b(s+x)^2)^3} + \frac{5(s+x)}{24a^2(a+b(s+x)^2)^2}$$

$$+ \frac{5(s+x)}{16a^3(a+b(s+x)^2)} + \frac{5 \arctan\left(\frac{\sqrt{b}(s+x)}{\sqrt{a}}\right)}{16a^{7/2}\sqrt{b}}$$

output 1/6\*(s+x)/a/(a+b\*(s+x)^2)^3+5/24\*(s+x)/a^2/(a+b\*(s+x)^2)^2+5/16\*(s+x)/a^3/(a+b\*(s+x)^2)+5/16\*arctan(b^(1/2)\*(s+x)/a^(1/2))/a^(7/2)/b^(1/2)

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.82

$$\int \frac{1}{\left( (a+bs^2)^2 + 4bs(a+bs^2)x + 2b(a+3bs^2)x^2 + 4b^2sx^3 + b^2x^4 \right)^2} dx$$

$$= \frac{(s+x)(33a^2 + 40ab(s+x)^2 + 15b^2(s+x)^4)}{48a^3(a+b(s+x)^2)^3} + \frac{5 \arctan\left(\frac{\sqrt{b}(s+x)}{\sqrt{a}}\right)}{16a^{7/2}\sqrt{b}}$$

input

```
Integrate[((a + b*s^2)^2 + 4*b*s*(a + b*s^2)*x + 2*b*(a + 3*b*s^2)*x^2 + 4
*b^2*s*x^3 + b^2*x^4)^(-2),x]
```

output

```
((s + x)*(33*a^2 + 40*a*b*(s + x)^2 + 15*b^2*(s + x)^4))/(48*a^3*(a + b*(s
+ x)^2)^3) + (5*ArcTan[(Sqrt[b]*(s + x))/Sqrt[a]])/(16*a^(7/2)*Sqrt[b])
```

### Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.38, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {2458, 1379, 215, 215, 215, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\left(2bx^2(a + 3bs^2) + 4bsx(a + bs^2) + (a + bs^2)^2 + 4b^2sx^3 + b^2x^4\right)^2} dx \\
 & \quad \downarrow 2458 \\
 & \int \frac{1}{(a^2 + 2ab(s+x)^2 + b^2(s+x)^4)^2} d(s+x) \\
 & \quad \downarrow 1379 \\
 & b^4 \int \frac{1}{(b^2(s+x)^2 + ab)^4} d(s+x) \\
 & \quad \downarrow 215 \\
 & b^4 \left( \frac{5 \int \frac{1}{(b^2(s+x)^2 + ab)^3} d(s+x)}{6ab} + \frac{s+x}{6ab^4(a + b(s+x)^2)^3} \right) \\
 & \quad \downarrow 215 \\
 & b^4 \left( \frac{5 \left( \frac{3 \int \frac{1}{(b^2(s+x)^2 + ab)^2} d(s+x)}{4ab} + \frac{s+x}{4ab^3(a + b(s+x)^2)^2} \right)}{6ab} + \frac{s+x}{6ab^4(a + b(s+x)^2)^3} \right)
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow \text{215} \\
 b^4 \left( \frac{5 \left( \frac{3 \left( \frac{\int \frac{1}{b^2(s+x)^2+ab} d(s+x)}{2ab} + \frac{s+x}{2ab^2(a+b(s+x)^2)} \right)}{4ab} + \frac{s+x}{4ab^3(a+b(s+x)^2)^2} \right)}{6ab} + \frac{s+x}{6ab^4(a+b(s+x)^2)^3} \right)
 \end{array}$$

$$\begin{array}{c}
 \downarrow \text{218} \\
 b^4 \left( \frac{5 \left( \frac{3 \left( \frac{\arctan\left(\frac{\sqrt{b}(s+x)}{\sqrt{a}}\right)}{2a^{3/2}b^{5/2}} + \frac{s+x}{2ab^2(a+b(s+x)^2)} \right)}{4ab} + \frac{s+x}{4ab^3(a+b(s+x)^2)^2} \right)}{6ab} + \frac{s+x}{6ab^4(a+b(s+x)^2)^3} \right)
 \end{array}$$

input

```
Int[((a + b*s^2)^2 + 4*b*s*(a + b*s^2)*x + 2*b*(a + 3*b*s^2)*x^2 + 4*b^2*s*x^3 + b^2*x^4)^(-2),x]
```

output

```
b^4*((s + x)/(6*a*b^4*(a + b*(s + x)^2)^3) + (5*((s + x)/(4*a*b^3*(a + b*(s + x)^2)^2) + (3*((s + x)/(2*a*b^2*(a + b*(s + x)^2)) + ArcTan[(Sqrt[b]*(s + x))/Sqrt[a]]/(2*a^(3/2)*b^(5/2))))/(4*a*b)))/(6*a*b))
```

### Defintions of rubi rules used

rule 215

```
Int[((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && LtQ[p, -1] && (IntegerQ[4*p] || IntegerQ[6*p])
```

rule 218

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```



rule 1379

```
Int[((a_) + (c_.)*(x_)^(n2_.) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/
c^p Int[(b/2 + c*x^n)^(2*p), x], x] /; FreeQ[{a, b, c, n, p}, x] && EqQ[n
2, 2*n] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p] && NeQ[p, 1]
```

rule 2458

```
Int[(Pn_)^(p_.), x_Symbol] := With[{S = Coeff[Pn, x, Expon[Pn, x] - 1]/(Exp
on[Pn, x]*Coeff[Pn, x, Expon[Pn, x]])}, Subst[Int[ExpandToSum[Pn /. x -> x
- S, x]^p, x], x, x + S] /; BinomialQ[Pn /. x -> x - S, x] || (IntegerQ[Exp
on[Pn, x]/2] && TrinomialQ[Pn /. x -> x - S, x]) /; FreeQ[p, x] && PolyQ[P
n, x] && GtQ[Expon[Pn, x], 2] && NeQ[Coeff[Pn, x, Expon[Pn, x] - 1], 0]
```

### Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.60

method	result
default	$\frac{2bs+2bx}{12ab(bs^2+2bsx+bx^2+a)^3} + \frac{5(2bs+2bx)}{48ab(bs^2+2bsx+bx^2+a)^2} + \frac{5\left(\frac{3(2bs+2bx)}{16ab(bs^2+2bsx+bx^2+a)} + \frac{3\arctan\left(\frac{2bs+2bx}{2\sqrt{ab}}\right)}{8a\sqrt{ab}}\right)}{6a}$
risch	$\frac{5b^2x^5}{16a^3} + \frac{25b^2sx^4}{16a^3} + \frac{5b(15bs^2+4a)x^3}{24a^3} + \frac{5bs(5bs^2+4a)x^2}{8a^3} + \frac{(25b^2s^4+40abs^2+11a^2)x}{16a^3} + \frac{s(15b^2s^4+40abs^2+33a^2)}{48a^3} - \frac{5\ln(bs+bx+\sqrt{-ab})}{32\sqrt{-ab}a^3}$

input

```
int(1/((b*s^2+a)^2+4*b*s*(b*s^2+a)*x+2*b*(3*b*s^2+a)*x^2+4*b^2*s*x^3+b^2*x
^4)^2,x,method=_RETURNVERBOSE)
```

output

```
1/12*(2*b*s+2*b*x)/a/b/(b*s^2+2*b*s*x+b*x^2+a)^3+5/6/a*(1/8*(2*b*s+2*b*x)/
a/b/(b*s^2+2*b*s*x+b*x^2+a)^2+3/4/a*(1/4*(2*b*s+2*b*x)/a/b/(b*s^2+2*b*s*x+
b*x^2+a)+1/2/a/(a*b)^(1/2)*arctan(1/2*(2*b*s+2*b*x)/(a*b)^(1/2)))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 465 vs.  $2(77) = 154$ .

Time = 0.09 (sec) , antiderivative size = 970, normalized size of antiderivative = 10.43

$$\int \frac{1}{((a + bs^2)^2 + 4bs(a + bs^2)x + 2b(a + 3bs^2)x^2 + 4b^2sx^3 + b^2x^4)^2} dx$$

= Too large to display

input

```
integrate(1/((b*s^2+a)^2+4*b*s*(b*s^2+a)*x+2*b*(3*b*s^2+a)*x^2+4*b^2*s*x^3
+b^2*x^4)^2,x, algorithm="fricas")
```

output

```
[1/96*(30*a*b^3*s^5 + 150*a*b^3*s*x^4 + 30*a*b^3*x^5 + 80*a^2*b^2*s^3 + 66
*a^3*b*s + 20*(15*a*b^3*s^2 + 4*a^2*b^2)*x^3 + 60*(5*a*b^3*s^3 + 4*a^2*b^2
*s)*x^2 - 15*(b^3*s^6 + 6*b^3*s*x^5 + b^3*x^6 + 3*a*b^2*s^4 + 3*a^2*b*s^2
+ 3*(5*b^3*s^2 + a*b^2)*x^4 + 4*(5*b^3*s^3 + 3*a*b^2*s)*x^3 + a^3 + 3*(5*b
^3*s^4 + 6*a*b^2*s^2 + a^2*b)*x^2 + 6*(b^3*s^5 + 2*a*b^2*s^3 + a^2*b*s)*x)
*sqrt(-a*b)*log((b*s^2 + 2*b*s*x + b*x^2 - 2*sqrt(-a*b)*(s + x) - a)/(b*s^
2 + 2*b*s*x + b*x^2 + a)) + 6*(25*a*b^3*s^4 + 40*a^2*b^2*s^2 + 11*a^3*b)*x
)/(a^4*b^4*s^6 + 6*a^4*b^4*s*x^5 + a^4*b^4*x^6 + 3*a^5*b^3*s^4 + 3*a^6*b^2
*s^2 + a^7*b + 3*(5*a^4*b^4*s^2 + a^5*b^3)*x^4 + 4*(5*a^4*b^4*s^3 + 3*a^5*
b^3*s)*x^3 + 3*(5*a^4*b^4*s^4 + 6*a^5*b^3*s^2 + a^6*b^2)*x^2 + 6*(a^4*b^4*
s^5 + 2*a^5*b^3*s^3 + a^6*b^2*s)*x), 1/48*(15*a*b^3*s^5 + 75*a*b^3*s*x^4 +
15*a*b^3*x^5 + 40*a^2*b^2*s^3 + 33*a^3*b*s + 10*(15*a*b^3*s^2 + 4*a^2*b^2
)*x^3 + 30*(5*a*b^3*s^3 + 4*a^2*b^2*s)*x^2 + 15*(b^3*s^6 + 6*b^3*s*x^5 + b
^3*x^6 + 3*a*b^2*s^4 + 3*a^2*b*s^2 + 3*(5*b^3*s^2 + a*b^2)*x^4 + 4*(5*b^3*
s^3 + 3*a*b^2*s)*x^3 + a^3 + 3*(5*b^3*s^4 + 6*a*b^2*s^2 + a^2*b)*x^2 + 6*(
b^3*s^5 + 2*a*b^2*s^3 + a^2*b*s)*x)*sqrt(a*b)*arctan(sqrt(a*b)*(s + x)/a)
+ 3*(25*a*b^3*s^4 + 40*a^2*b^2*s^2 + 11*a^3*b)*x)/(a^4*b^4*s^6 + 6*a^4*b^4
*s*x^5 + a^4*b^4*x^6 + 3*a^5*b^3*s^4 + 3*a^6*b^2*s^2 + a^7*b + 3*(5*a^4*b^
4*s^2 + a^5*b^3)*x^4 + 4*(5*a^4*b^4*s^3 + 3*a^5*b^3*s)*x^3 + 3*(5*a^4*b^4*
s^4 + 6*a^5*b^3*s^2 + a^6*b^2)*x^2 + 6*(a^4*b^4*s^5 + 2*a^5*b^3*s^3 + a...
```

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 354 vs.  $2(92) = 184$ .

Time = 0.95 (sec) , antiderivative size = 354, normalized size of antiderivative = 3.81

$$\int \frac{1}{((a + bs^2)^2 + 4bs(a + bs^2)x + 2b(a + 3bs^2)x^2 + 4b^2sx^3 + b^2x^4)^2} dx$$

$$= -\frac{5\sqrt{-\frac{1}{ab}} \log\left(-a^4\sqrt{-\frac{1}{ab}} + s + x\right)}{32} + \frac{5\sqrt{-\frac{1}{ab}} \log\left(a^4\sqrt{-\frac{1}{ab}} + s + x\right)}{32}$$

$$+ \frac{33a^2s + 40abs^3 + 15b^2s^5 + 75b^2sx^4 + 15b^2x^5 + x^3 \cdot (40ab + 15b^2)}{48a^6 + 144a^5bs^2 + 144a^4b^2s^4 + 48a^3b^3s^6 + 288a^3b^3sx^5 + 48a^3b^3x^6 + x^4 \cdot (144a^4b^2 + 720a^3b^3s^2) + x^3 \cdot (40ab + 15b^2)}$$

input `integrate(1/((b*s**2+a)**2+4*b*s*(b*s**2+a)*x+2*b*(3*b*s**2+a)*x**2+4*b**2*s*x**3+b**2*x**4)**2,x)`

output `-5*sqrt(-1/(a**7*b))*log(-a**4*sqrt(-1/(a**7*b)) + s + x)/32 + 5*sqrt(-1/(a**7*b))*log(a**4*sqrt(-1/(a**7*b)) + s + x)/32 + (33*a**2*s + 40*a*b*s**3 + 15*b**2*s**5 + 75*b**2*s*x**4 + 15*b**2*x**5 + x**3*(40*a*b + 150*b**2*s**2) + x**2*(120*a*b*s + 150*b**2*s**3) + x*(33*a**2 + 120*a*b*s**2 + 75*b**2*s**4))/(48*a**6 + 144*a**5*b*s**2 + 144*a**4*b**2*s**4 + 48*a**3*b**3*s**6 + 288*a**3*b**3*s*x**5 + 48*a**3*b**3*x**6 + x**4*(144*a**4*b**2 + 720*a**3*b**3*s**2) + x**3*(576*a**4*b**2*s + 960*a**3*b**3*s**3) + x**2*(144*a**5*b + 864*a**4*b**2*s**2 + 720*a**3*b**3*s**4) + x*(288*a**5*b*s + 576*a**4*b**2*s**3 + 288*a**3*b**3*s**5))`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 299 vs.  $2(77) = 154$ .

Time = 0.11 (sec) , antiderivative size = 299, normalized size of antiderivative = 3.22

$$\int \frac{1}{((a + bs^2)^2 + 4bs(a + bs^2)x + 2b(a + 3bs^2)x^2 + 4b^2sx^3 + b^2x^4)^2} dx$$

$$= \frac{15b^2s^5 + 75b^2sx^4 + 15b^2x^5 + 40abs^3 + 10(15b^2s^2 + 4ab)x^3 + 33a^2s + 30(5b^2s^2 + 4ab)x^2 + 4(5a^3b^3s^3 + 3a^4b^2s^2)}{48(a^3b^3s^6 + 6a^3b^3sx^5 + a^3b^3x^6 + 3a^4b^2s^4 + 3a^5bs^2 + a^6 + 3(5a^3b^3s^2 + a^4b^2)x^4 + 4(5a^3b^3s^3 + 3a^4b^2s^2) + 33a^2s + 30(5b^2s^2 + 4ab)x^2 + 4(5a^3b^3s^3 + 3a^4b^2s^2))} + \frac{5 \arctan\left(\frac{bs+bx}{\sqrt{ab}}\right)}{16\sqrt{aba^3}}$$

input `integrate(1/((b*s^2+a)^2+4*b*s*(b*s^2+a)*x+2*b*(3*b*s^2+a)*x^2+4*b^2*s*x^3+b^2*x^4)^2,x, algorithm="maxima")`

output 
$$\frac{1}{48} \frac{(15b^2s^5 + 75b^2s^4x + 15b^2x^5 + 40ab^2s^3 + 10(15b^2s^2 + 4ab)sx^3 + 33a^2s + 30(5b^2s^3 + 4ab^2s)x^2 + 3(25b^2s^4 + 40ab^2s^2 + 11a^2)sx)/(a^3b^3s^6 + 6a^3b^3s^5x + a^3b^3s^4x^2 + 3a^4b^2s^4 + 3a^5b^2s^3 + a^6 + 3(5a^3b^3s^2 + a^4b^2)x^4 + 4(5a^3b^3s^3 + 3a^4b^2s)x^3 + 3(5a^3b^3s^4 + 6a^4b^2s^2 + a^5b)x^2 + 6(a^3b^3s^5 + 2a^4b^2s^3 + a^5b)sx) + 5/16 \arctan((bs + bx)/\sqrt{ab})/(\sqrt{ab})a^3}{(a + bs^2)^2 + 4bs(a + bs^2)x + 2b(a + 3bs^2)x^2 + 4b^2sx^3 + b^2x^4} dx$$

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.59

$$\int \frac{1}{((a + bs^2)^2 + 4bs(a + bs^2)x + 2b(a + 3bs^2)x^2 + 4b^2sx^3 + b^2x^4)^2} dx$$

$$= \frac{5 \arctan\left(\frac{bs+bx}{\sqrt{ab}}\right)}{16\sqrt{ab}a^3} + \frac{15b^2s^5 + 75b^2s^4x + 150b^2s^3x^2 + 150b^2s^2x^3 + 75b^2sx^4 + 15b^2x^5 + 40abs^3 + 120abs^2x + 120absx^2 - 48(bs^2 + 2bsx + bx^2 + a)^3a^3}{48(bs^2 + 2bsx + bx^2 + a)^3a^3}$$

input `integrate(1/((b*s^2+a)^2+4*b*s*(b*s^2+a)*x+2*b*(3*b*s^2+a)*x^2+4*b^2*s*x^3+b^2*x^4)^2,x, algorithm="giac")`

output 
$$\frac{5}{16} \frac{\arctan((bs + bx)/\sqrt{ab})/(\sqrt{ab})}{a^3} + \frac{1}{48} \frac{(15b^2s^5 + 75b^2s^4x + 150b^2s^3x^2 + 150b^2s^2x^3 + 75b^2sx^4 + 15b^2x^5 + 40ab^2s^3 + 120ab^2s^2x + 120ab^2sx^2 + 40ab^2x^3 + 33a^2s + 33a^2sx)/(b^3s^6 + 6a^3b^3s^5x + a^3b^3s^4x^2 + 3a^4b^2s^4 + 3a^5b^2s^3 + a^6 + 3(5a^3b^3s^2 + a^4b^2)x^4 + 4(5a^3b^3s^3 + 3a^4b^2s)x^3 + 3(5a^3b^3s^4 + 6a^4b^2s^2 + a^5b)x^2 + 6(a^3b^3s^5 + 2a^4b^2s^3 + a^5b)sx) + 5/16 \arctan((bs + bx)/\sqrt{ab})/(\sqrt{ab})a^3}{(a + bs^2)^2 + 4bs(a + bs^2)x + 2b(a + 3bs^2)x^2 + 4b^2sx^3 + b^2x^4} dx$$

**Mupad [B] (verification not implemented)**

Time = 21.62 (sec) , antiderivative size = 297, normalized size of antiderivative = 3.19

$$\int \frac{1}{((a + bs^2)^2 + 4bs(a + bs^2)x + 2b(a + 3bs^2)x^2 + 4b^2sx^3 + b^2x^4)^2} dx$$

$$= \frac{33a^2s + 40abs^3 + 15b^2s^5}{48a^3} + \frac{5x^3(15b^2s^2 + 4ab)}{24a^3} + \frac{5b^2x^5}{16a^3} + \frac{5x^2(5b^2s^3 + 4abs)}{8a^3} + \frac{x(15b^3s^2 + 3ab^2)}{x^4(15b^3s^2 + 3ab^2)} + \frac{x^2(3a^2b + 18ab^2s^2 + 15b^3s^4)}{x^2(3a^2b + 18ab^2s^2 + 15b^3s^4)} + \frac{x(6a^2bs + 12ab^2s^3 + 6b^3s^5)}{x(6a^2bs + 12ab^2s^3 + 6b^3s^5)} + \frac{x^3(20b^3s^3)}{x^3(20b^3s^3)}$$

$$+ \frac{5 \operatorname{atan}\left(\frac{16a^3\left(\frac{5\sqrt{b}s}{16a^{7/2}} + \frac{5\sqrt{b}x}{16a^{7/2}}\right)}{5}\right)}{16a^{7/2}\sqrt{b}}$$

input `int(1/((a + b*s^2)^2 + b^2*x^4 + 4*b^2*s*x^3 + 2*b*x^2*(a + 3*b*s^2) + 4*b*s*x*(a + b*s^2))^2,x)`

output `((33*a^2*s + 15*b^2*s^5 + 40*a*b*s^3)/(48*a^3) + (5*x^3*(4*a*b + 15*b^2*s^2))/(24*a^3) + (5*b^2*x^5)/(16*a^3) + (5*x^2*(5*b^2*s^3 + 4*a*b*s))/(8*a^3) + (x*(11*a^2 + 25*b^2*s^4 + 40*a*b*s^2))/(16*a^3) + (25*b^2*s*x^4)/(16*a^3))/((x^4*(3*a*b^2 + 15*b^3*s^2) + x^2*(3*a^2*b + 15*b^3*s^4 + 18*a*b^2*s^2) + x*(6*b^3*s^5 + 12*a*b^2*s^3 + 6*a^2*b*s) + x^3*(20*b^3*s^3 + 12*a*b^2*s*s) + a^3 + b^3*s^6 + b^3*x^6 + 3*a^2*b*s^2 + 3*a*b^2*s^4 + 6*b^3*s*x^5) + (5*atan((16*a^3*((5*b^(1/2)*s)/(16*a^(7/2)) + (5*b^(1/2)*x)/(16*a^(7/2))))/5))/(16*a^(7/2)*b^(1/2))`

**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 808, normalized size of antiderivative = 8.69

$$\int \frac{1}{((a + bs^2)^2 + 4bs(a + bs^2)x + 2b(a + 3bs^2)x^2 + 4b^2sx^3 + b^2x^4)^2} dx$$

= Too large to display

input `int(1/((b*s^2+a)^2+4*b*s*(b*s^2+a)*x+2*b*(3*b*s^2+a)*x^2+4*b^2*s*x^3+b^2*x^4)^2,x)`

output

```
(30*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**3*s + 90*sqrt(b)
)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**2*b*s**3 + 180*sqrt(b)*sq
rt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**2*b*s**2*x + 90*sqrt(b)*sqrt(
a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**2*b*s*x**2 + 90*sqrt(b)*sqrt(a)*
atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**5 + 360*sqrt(b)*sqrt(a)*atan
((b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**4*x + 540*sqrt(b)*sqrt(a)*atan((
b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**3*x**2 + 360*sqrt(b)*sqrt(a)*atan(
(b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**2*x**3 + 90*sqrt(b)*sqrt(a)*atan(
(b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s*x**4 + 30*sqrt(b)*sqrt(a)*atan((b*
s + b*x)/(sqrt(b)*sqrt(a)))*b**3*s**7 + 180*sqrt(b)*sqrt(a)*atan((b*s + b*
x)/(sqrt(b)*sqrt(a)))*b**3*s**6*x + 450*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(
sqrt(b)*sqrt(a)))*b**3*s**5*x**2 + 600*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(s
qrt(b)*sqrt(a)))*b**3*s**4*x**3 + 450*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sq
rt(b)*sqrt(a)))*b**3*s**3*x**4 + 180*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqr
t(b)*sqrt(a)))*b**3*s**2*x**5 + 30*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(
b)*sqrt(a)))*b**3*s*x**6 - 5*a**4 + 51*a**3*b*s**2 + 36*a**3*b*s*x - 15*a*
*3*b*x**2 + 65*a**2*b**2*s**4 + 180*a**2*b**2*s**3*x + 150*a**2*b**2*s**2*
x**2 + 20*a**2*b**2*s*x**3 - 15*a**2*b**2*x**4 + 25*a*b**3*s**6 + 120*a*b*
*3*s**5*x + 225*a*b**3*s**4*x**2 + 200*a*b**3*s**3*x**3 + 75*a*b**3*s**2*x
**4 - 5*a*b**3*x**6)/(96*a**4*b*s*(a**3 + 3*a**2*b*s**2 + 6*a**2*b*s*x ...
```



input

```
Integrate[((a + b*s^2)^4 + 8*b*s*(a + b*s^2)^3*x + 4*b*(a + b*s^2)^2*(a +
7*b*s^2)*x^2 + 8*b^2*s*(3*a^2 + 10*a*b*s^2 + 7*b^2*s^4)*x^3 + 2*b^2*(3*a^2
+ 30*a*b*s^2 + 35*b^2*s^4)*x^4 + 8*b^3*s*(3*a + 7*b*s^2)*x^5 + 4*b^3*(a +
7*b*s^2)*x^6 + 8*b^4*s*x^7 + b^4*x^8)^(-1),x]
```

output

```
((s + x)*(33*a^2 + 40*a*b*(s + x)^2 + 15*b^2*(s + x)^4))/(48*a^3*(a + b*(s
+ x)^2)^3) + (5*ArcTan[(Sqrt[b]*(s + x))/Sqrt[a]])/(16*a^(7/2)*Sqrt[b])
```

### Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 311 vs.  $2(93) = 186$ .

Time = 0.85 (sec) , antiderivative size = 311, normalized size of antiderivative = 3.34,  
 number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.013$ , Rules  
 used = {2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{2b^2x^4(3a^2 + 30abs^2 + 35b^2s^4) + 8b^2sx^3(3a^2 + 10abs^2 + 7b^2s^4) + 4b^3x^6(a + 7bs^2) + 8b^3sx^5(3a + 7bs^2) + 4b^4x^8} dx$$

↓ 2462

$$\int \left( -\frac{5b}{8a^3(2\sqrt{-a}\sqrt{b} - 2bs - 2bx)^2} - \frac{5b}{8a^3(2\sqrt{-a}\sqrt{b} + 2bs + 2bx)^2} + \frac{b^2}{a^2(2\sqrt{-a}\sqrt{b} - 2bs - 2bx)^4} + \frac{b^2}{a^2(2\sqrt{-a}\sqrt{b} + 2bs + 2bx)^4} \right) dx$$

↓ 2009



$$\begin{aligned} & -\frac{5}{32a^3\sqrt{b}\left(\sqrt{-a}-\sqrt{bs}-\sqrt{bx}\right)} + \frac{5}{32a^3\sqrt{b}\left(\sqrt{-a}+\sqrt{bs}+\sqrt{bx}\right)} + \\ & \frac{1}{48a^2\sqrt{b}\left(\sqrt{-a}-\sqrt{bs}-\sqrt{bx}\right)^3} - \frac{1}{48a^2\sqrt{b}\left(\sqrt{-a}+\sqrt{bs}+\sqrt{bx}\right)^3} + \\ & \frac{16(-a)^{5/2}\sqrt{b}\left(\sqrt{-a}-\sqrt{bs}-\sqrt{bx}\right)^2}{5\log\left(\sqrt{-a}-\sqrt{bs}-\sqrt{bx}\right)} - \frac{16(-a)^{5/2}\sqrt{b}\left(\sqrt{-a}+\sqrt{bs}+\sqrt{bx}\right)^2}{5\log\left(\sqrt{-a}+\sqrt{bs}+\sqrt{bx}\right)} - \\ & \frac{5\log\left(\sqrt{-a}-\sqrt{bs}-\sqrt{bx}\right)}{32(-a)^{7/2}\sqrt{b}} + \frac{5\log\left(\sqrt{-a}+\sqrt{bs}+\sqrt{bx}\right)}{32(-a)^{7/2}\sqrt{b}} \end{aligned}$$

input

```
Int[((a + b*s^2)^4 + 8*b*s*(a + b*s^2)^3*x + 4*b*(a + b*s^2)^2*(a + 7*b*s^2)*x^2 + 8*b^2*s*(3*a^2 + 10*a*b*s^2 + 7*b^2*s^4)*x^3 + 2*b^2*(3*a^2 + 30*a*b*s^2 + 35*b^2*s^4)*x^4 + 8*b^3*s*(3*a + 7*b*s^2)*x^5 + 4*b^3*(a + 7*b*s^2)*x^6 + 8*b^4*s*x^7 + b^4*x^8)^(-1), x]
```

output

```
1/(48*a^2*Sqrt[b]*(Sqrt[-a] - Sqrt[b]*s - Sqrt[b]*x)^3) + 1/(16*(-a)^(5/2)*Sqrt[b]*(Sqrt[-a] - Sqrt[b]*s - Sqrt[b]*x)^2) - 5/(32*a^3*Sqrt[b]*(Sqrt[-a] - Sqrt[b]*s - Sqrt[b]*x)) - 1/(48*a^2*Sqrt[b]*(Sqrt[-a] + Sqrt[b]*s + Sqrt[b]*x)^3) - 1/(16*(-a)^(5/2)*Sqrt[b]*(Sqrt[-a] + Sqrt[b]*s + Sqrt[b]*x)^2) + 5/(32*a^3*Sqrt[b]*(Sqrt[-a] + Sqrt[b]*s + Sqrt[b]*x)) - (5*Log[Sqrt[-a] - Sqrt[b]*s - Sqrt[b]*x])/(32*(-a)^(7/2)*Sqrt[b]) + (5*Log[Sqrt[-a] + Sqrt[b]*s + Sqrt[b]*x])/(32*(-a)^(7/2)*Sqrt[b])
```

### Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2462

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegrand[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0] && RationalFunctionQ[u, x]
```

### Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.60

method	result
default	$\frac{2bs+2bx}{12ab(b^2s^2+2bsx+bx^2+a)^3} + \frac{5(2bs+2bx)}{48ab(b^2s^2+2bsx+bx^2+a)^2} + \frac{5 \left( \frac{3(2bs+2bx)}{16ab(b^2s^2+2bsx+bx^2+a)} + \frac{3 \arctan\left(\frac{2bs+2bx}{2\sqrt{ab}}\right)}{8a\sqrt{ab}} \right)}{6a}$
risch	$\frac{\frac{5b^2x^5}{16a^3} + \frac{25b^2sx^4}{16a^3} + \frac{5b(15bs^2+4a)x^3}{24a^3} + \frac{5bs(5bs^2+4a)x^2}{8a^3} + \frac{(25b^2s^4+40abs^2+11a^2)x}{16a^3} + \frac{s(15b^2s^4+40abs^2+33a^2)}{48a^3}}{b^3s^6+6b^3s^5x+15b^3s^4x^2+20b^3s^3x^3+15b^3s^2x^4+6b^3sx^5+b^3x^6+3ab^2s^4+12ab^2s^3x+18ab^2s^2x^2+12ab^2sx^3+3ab^2x^4+3a^2bs^2+6a^2bx+3a^2}$

```
input int(1/((b*s^2+a)^4+8*b*s*(b*s^2+a)^3*x+4*b*(b*s^2+a)^2*(7*b*s^2+a)*x^2+8*b^2*s*(7*b^2*s^4+10*a*b*s^2+3*a^2)*x^3+2*b^2*(35*b^2*s^4+30*a*b*s^2+3*a^2)*x^4+8*b^3*s*(7*b*s^2+3*a)*x^5+4*b^3*(7*b*s^2+a)*x^6+8*b^4*s*x^7+b^4*x^8),x,method=_RETURNVERBOSE)
```

```
output 1/12*(2*b*s+2*b*x)/a/b/(b*s^2+2*b*s*x+b*x^2+a)^3+5/6/a*(1/8*(2*b*s+2*b*x)/a/b/(b*s^2+2*b*s*x+b*x^2+a)^2+3/4/a*(1/4*(2*b*s+2*b*x)/a/b/(b*s^2+2*b*s*x+b*x^2+a)+1/2/a/(a*b)^(1/2)*arctan(1/2*(2*b*s+2*b*x)/(a*b)^(1/2))))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 465 vs. 2(77) = 154.

Time = 0.10 (sec) , antiderivative size = 970, normalized size of antiderivative = 10.43

$$\int \frac{1}{(a + bs^2)^4 + 8bs(a + bs^2)^3x + 4b(a + bs^2)^2(a + 7bs^2)x^2 + 8b^2s(3a^2 + 10abs^2 + 7b^2s^4)x^3 + 2b^2(3a^2 + 10abs^2 + 7b^2s^4)x^4 + 8b^3s^2(3a^2 + 10abs^2 + 7b^2s^4)x^5 + 4b^3s^2(3a^2 + 10abs^2 + 7b^2s^4)x^6 + 8b^4s^3(3a^2 + 10abs^2 + 7b^2s^4)x^7 + b^4s^3(3a^2 + 10abs^2 + 7b^2s^4)x^8} dx$$

= Too large to display

```
input integrate(1/((b*s^2+a)^4+8*b*s*(b*s^2+a)^3*x+4*b*(b*s^2+a)^2*(7*b*s^2+a)*x^2+8*b^2*s*(7*b^2*s^4+10*a*b*s^2+3*a^2)*x^3+2*b^2*(35*b^2*s^4+30*a*b*s^2+3*a^2)*x^4+8*b^3*s*(7*b*s^2+3*a)*x^5+4*b^3*(7*b*s^2+a)*x^6+8*b^4*s*x^7+b^4*x^8),x,algorithm="fricas")
```

output

```
[1/96*(30*a*b^3*s^5 + 150*a*b^3*s*x^4 + 30*a*b^3*x^5 + 80*a^2*b^2*s^3 + 66
*a^3*b*s + 20*(15*a*b^3*s^2 + 4*a^2*b^2)*x^3 + 60*(5*a*b^3*s^3 + 4*a^2*b^2
*s)*x^2 - 15*(b^3*s^6 + 6*b^3*s*x^5 + b^3*x^6 + 3*a*b^2*s^4 + 3*a^2*b*s^2
+ 3*(5*b^3*s^2 + a*b^2)*x^4 + 4*(5*b^3*s^3 + 3*a*b^2*s)*x^3 + a^3 + 3*(5*b
^3*s^4 + 6*a*b^2*s^2 + a^2*b)*x^2 + 6*(b^3*s^5 + 2*a*b^2*s^3 + a^2*b*s)*x)
*sqrt(-a*b)*log((b*s^2 + 2*b*s*x + b*x^2 - 2*sqrt(-a*b)*(s + x) - a)/(b*s^
2 + 2*b*s*x + b*x^2 + a)) + 6*(25*a*b^3*s^4 + 40*a^2*b^2*s^2 + 11*a^3*b)*x
)/(a^4*b^4*s^6 + 6*a^4*b^4*s*x^5 + a^4*b^4*x^6 + 3*a^5*b^3*s^4 + 3*a^6*b^2
*s^2 + a^7*b + 3*(5*a^4*b^4*s^2 + a^5*b^3)*x^4 + 4*(5*a^4*b^4*s^3 + 3*a^5*
b^3*s)*x^3 + 3*(5*a^4*b^4*s^4 + 6*a^5*b^3*s^2 + a^6*b^2)*x^2 + 6*(a^4*b^4*
s^5 + 2*a^5*b^3*s^3 + a^6*b^2*s)*x), 1/48*(15*a*b^3*s^5 + 75*a*b^3*s*x^4 +
15*a*b^3*x^5 + 40*a^2*b^2*s^3 + 33*a^3*b*s + 10*(15*a*b^3*s^2 + 4*a^2*b^2
)*x^3 + 30*(5*a*b^3*s^3 + 4*a^2*b^2*s)*x^2 + 15*(b^3*s^6 + 6*b^3*s*x^5 + b
^3*x^6 + 3*a*b^2*s^4 + 3*a^2*b*s^2 + 3*(5*b^3*s^2 + a*b^2)*x^4 + 4*(5*b^3*
s^3 + 3*a*b^2*s)*x^3 + a^3 + 3*(5*b^3*s^4 + 6*a*b^2*s^2 + a^2*b)*x^2 + 6*(
b^3*s^5 + 2*a*b^2*s^3 + a^2*b*s)*x)*sqrt(a*b)*arctan(sqrt(a*b)*(s + x)/a)
+ 3*(25*a*b^3*s^4 + 40*a^2*b^2*s^2 + 11*a^3*b)*x)/(a^4*b^4*s^6 + 6*a^4*b^4
*s*x^5 + a^4*b^4*x^6 + 3*a^5*b^3*s^4 + 3*a^6*b^2*s^2 + a^7*b + 3*(5*a^4*b^
4*s^2 + a^5*b^3)*x^4 + 4*(5*a^4*b^4*s^3 + 3*a^5*b^3*s)*x^3 + 3*(5*a^4*b^4*
s^4 + 6*a^5*b^3*s^2 + a^6*b^2)*x^2 + 6*(a^4*b^4*s^5 + 2*a^5*b^3*s^3 + a...
```

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 354 vs.  $2(92) = 184$ .

Time = 0.96 (sec) , antiderivative size = 354, normalized size of antiderivative = 3.81

$$\int \frac{1}{(a + bs^2)^4 + 8bs(a + bs^2)^3 x + 4b(a + bs^2)^2(a + 7bs^2)x^2 + 8b^2s(3a^2 + 10abs^2 + 7b^2s^4)x^3 + 2b^2(3a^2 + 5\sqrt{-\frac{1}{a^7b}} \log\left(-a^4\sqrt{-\frac{1}{a^7b}} + s + x\right) + 5\sqrt{-\frac{1}{a^7b}} \log\left(a^4\sqrt{-\frac{1}{a^7b}} + s + x\right))} dx$$

$$= -\frac{5\sqrt{-\frac{1}{a^7b}} \log\left(-a^4\sqrt{-\frac{1}{a^7b}} + s + x\right)}{32} + \frac{5\sqrt{-\frac{1}{a^7b}} \log\left(a^4\sqrt{-\frac{1}{a^7b}} + s + x\right)}{32}$$

$$+ \frac{33a^2s + 40abs^3 + 15b^2s^5 + 75b^2sx^4 + 15b^2x^5 + x^3 \cdot (40ab + 1)}{48a^6 + 144a^5bs^2 + 144a^4b^2s^4 + 48a^3b^3s^6 + 288a^3b^3sx^5 + 48a^3b^3x^6 + x^4 \cdot (144a^4b^2 + 720a^3b^3s^2) + x^3}$$

input

```
integrate(1/((b*s**2+a)**4+8*b*s*(b*s**2+a)**3*x+4*b*(b*s**2+a)**2*(7*b*s*
*2+a)*x**2+8*b**2*s*(7*b**2*s**4+10*a*b*s**2+3*a**2)*x**3+2*b**2*(35*b**2*
s**4+30*a*b*s**2+3*a**2)*x**4+8*b**3*s*(7*b*s**2+3*a)*x**5+4*b**3*(7*b*s**
2+a)*x**6+8*b**4*s*x**7+b**4*x**8), x)
```

output

```
-5*sqrt(-1/(a**7*b))*log(-a**4*sqrt(-1/(a**7*b)) + s + x)/32 + 5*sqrt(-1/(
a**7*b))*log(a**4*sqrt(-1/(a**7*b)) + s + x)/32 + (33*a**2*s + 40*a*b*s**3
+ 15*b**2*s**5 + 75*b**2*s*x**4 + 15*b**2*x**5 + x**3*(40*a*b + 150*b**2*
s**2) + x**2*(120*a*b*s + 150*b**2*s**3) + x*(33*a**2 + 120*a*b*s**2 + 75*
b**2*s**4))/(48*a**6 + 144*a**5*b*s**2 + 144*a**4*b**2*s**4 + 48*a**3*b**3
*s**6 + 288*a**3*b**3*s*x**5 + 48*a**3*b**3*x**6 + x**4*(144*a**4*b**2 + 7
20*a**3*b**3*s**2) + x**3*(576*a**4*b**2*s + 960*a**3*b**3*s**3) + x**2*(1
44*a**5*b + 864*a**4*b**2*s**2 + 720*a**3*b**3*s**4) + x*(288*a**5*b*s + 5
76*a**4*b**2*s**3 + 288*a**3*b**3*s**5))
```

### Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 299 vs.  $2(77) = 154$ .

Time = 0.11 (sec) , antiderivative size = 299, normalized size of antiderivative = 3.22

$$\int \frac{1}{(a + bs^2)^4 + 8bs(a + bs^2)^3 x + 4b(a + bs^2)^2(a + 7bs^2)x^2 + 8b^2s(3a^2 + 10abs^2 + 7b^2s^4)x^3 + 2b^2(3a^2 + 15b^2s^5 + 75b^2sx^4 + 15b^2x^5 + 40abs^3 + 10(15b^2s^2 + 4ab)x^3 + 33a^2s + 30(5b^2s^2 + 4ab^2s^2))} dx$$

$$= \frac{5 \arctan\left(\frac{bs+bx}{\sqrt{ab}}\right)}{16\sqrt{aba^3}}$$

input

```
integrate(1/((b*s^2+a)^4+8*b*s*(b*s^2+a)^3*x+4*b*(b*s^2+a)^2*(7*b*s^2+a)*x
^2+8*b^2*s*(7*b^2*s^4+10*a*b*s^2+3*a^2)*x^3+2*b^2*(35*b^2*s^4+30*a*b*s^2+3
*a^2)*x^4+8*b^3*s*(7*b*s^2+3*a)*x^5+4*b^3*(7*b*s^2+a)*x^6+8*b^4*s*x^7+b^4*
x^8),x, algorithm="maxima")
```

output

```
1/48*(15*b^2*s^5 + 75*b^2*s*x^4 + 15*b^2*x^5 + 40*a*b*s^3 + 10*(15*b^2*s^2
+ 4*a*b)*x^3 + 33*a^2*s + 30*(5*b^2*s^3 + 4*a*b*s)*x^2 + 3*(25*b^2*s^4 +
40*a*b*s^2 + 11*a^2)*x)/(a^3*b^3*s^6 + 6*a^3*b^3*s*x^5 + a^3*b^3*x^6 + 3*a
^4*b^2*s^4 + 3*a^5*b*s^2 + a^6 + 3*(5*a^3*b^3*s^2 + a^4*b^2)*x^4 + 4*(5*a^
3*b^3*s^3 + 3*a^4*b^2*s)*x^3 + 3*(5*a^3*b^3*s^4 + 6*a^4*b^2*s^2 + a^5*b)*x
^2 + 6*(a^3*b^3*s^5 + 2*a^4*b^2*s^3 + a^5*b*s)*x) + 5/16*arctan((b*s + b*x
)/sqrt(a*b))/(sqrt(a*b)*a^3)
```

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.59

$$\int \frac{1}{(a + bs^2)^4 + 8bs(a + bs^2)^3 x + 4b(a + bs^2)^2(a + 7bs^2)x^2 + 8b^2s(3a^2 + 10abs^2 + 7b^2s^4)x^3 + 2b^2(3a^2 + 5 \arctan\left(\frac{bs+bx}{\sqrt{ab}}\right))} dx$$

$$= \frac{16\sqrt{aba^3} + 15b^2s^5 + 75b^2s^4x + 150b^2s^3x^2 + 150b^2s^2x^3 + 75b^2sx^4 + 15b^2x^5 + 40abs^3 + 120abs^2x + 120absx^2 - 48(bs^2 + 2bsx + bx^2 + a)^3a^3}{48(bs^2 + 2bsx + bx^2 + a)^3a^3}$$

input

```
integrate(1/((b*s^2+a)^4+8*b*s*(b*s^2+a)^3*x+4*b*(b*s^2+a)^2*(7*b*s^2+a)*x^2+8*b^2*s*(7*b^2*s^4+10*a*b*s^2+3*a^2)*x^3+2*b^2*(35*b^2*s^4+30*a*b*s^2+3*a^2)*x^4+8*b^3*s*(7*b*s^2+3*a)*x^5+4*b^3*(7*b*s^2+a)*x^6+8*b^4*s*x^7+b^4*x^8),x, algorithm="giac")
```

output

```
5/16*arctan((b*s + b*x)/sqrt(a*b))/(sqrt(a*b)*a^3) + 1/48*(15*b^2*s^5 + 75*b^2*s^4*x + 150*b^2*s^3*x^2 + 150*b^2*s^2*x^3 + 75*b^2*s*x^4 + 15*b^2*x^5 + 40*a*b*s^3 + 120*a*b*s^2*x + 120*a*b*s*x^2 + 40*a*b*x^3 + 33*a^2*s + 33*a^2*x)/((b*s^2 + 2*b*s*x + b*x^2 + a)^3*a^3)
```

**Mupad [B] (verification not implemented)**

Time = 21.87 (sec) , antiderivative size = 297, normalized size of antiderivative = 3.19

$$\int \frac{1}{(a + bs^2)^4 + 8bs(a + bs^2)^3 x + 4b(a + bs^2)^2(a + 7bs^2)x^2 + 8b^2s(3a^2 + 10abs^2 + 7b^2s^4)x^3 + 2b^2(3a^2 + 33a^2s + 33a^2x) \arctan\left(\frac{5\sqrt{b}s}{16a^{7/2}} + \frac{5\sqrt{b}x}{16a^{7/2}}\right)} dx$$

$$= \frac{\frac{33a^2s+40ab^3s^3+15b^2s^5}{48a^3} + \frac{5x^3(15b^2s^2+4ab)}{24a^3} + \frac{5b^2x^5}{16a^3} + \frac{5x^2(5b^2s^3+4abs)}{8a^3} + \frac{x(20b^3s^3)}{8a^3}}{x^4(15b^3s^2+3ab^2) + x^2(3a^2b+18ab^2s^2+15b^3s^4) + x(6a^2bs+12ab^2s^3+6b^3s^5) + x^3(20b^3s^3) + \frac{5 \operatorname{atan}\left(\frac{16a^3\left(\frac{5\sqrt{b}s}{16a^{7/2}} + \frac{5\sqrt{b}x}{16a^{7/2}}\right)}{5}\right)}{16a^{7/2}\sqrt{b}}}$$

input

```
int(1/((a + b*s^2)^4 + b^4*x^8 + 4*b^3*x^6*(a + 7*b*s^2) + 2*b^2*x^4*(3*a^2 + 35*b^2*s^4 + 30*a*b*s^2) + 8*b^4*s*x^7 + 8*b^3*s*x^5*(3*a + 7*b*s^2) + 8*b*s*x*(a + b*s^2)^3 + 4*b*x^2*(a + b*s^2)^2*(a + 7*b*s^2) + 8*b^2*s*x^3*(3*a^2 + 7*b^2*s^4 + 10*a*b*s^2)),x)
```

output

```
((33*a^2*s + 15*b^2*s^5 + 40*a*b*s^3)/(48*a^3) + (5*x^3*(4*a*b + 15*b^2*s^2))/(24*a^3) + (5*b^2*x^5)/(16*a^3) + (5*x^2*(5*b^2*s^3 + 4*a*b*s))/(8*a^3) + (x*(11*a^2 + 25*b^2*s^4 + 40*a*b*s^2))/(16*a^3) + (25*b^2*s*x^4)/(16*a^3))/(x^4*(3*a*b^2 + 15*b^3*s^2) + x^2*(3*a^2*b + 15*b^3*s^4 + 18*a*b^2*s^2) + x*(6*b^3*s^5 + 12*a*b^2*s^3 + 6*a^2*b*s) + x^3*(20*b^3*s^3 + 12*a*b^2*s) + a^3 + b^3*s^6 + b^3*x^6 + 3*a^2*b*s^2 + 3*a*b^2*s^4 + 6*b^3*s*x^5) + (5*atan((16*a^3*((5*b^(1/2)*s)/(16*a^(7/2)) + (5*b^(1/2)*x)/(16*a^(7/2))))/5))/(16*a^(7/2)*b^(1/2))
```

**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 808, normalized size of antiderivative = 8.69

$$\int \frac{1}{(a + bs^2)^4 + 8bs(a + bs^2)^3 x + 4b(a + bs^2)^2(a + 7bs^2)x^2 + 8b^2s(3a^2 + 10abs^2 + 7b^2s^4)x^3 + 2b^2(3a^2 + 10abs^2 + 7b^2s^4)x^4 + 8b^3s(7b^2s^2 + 3a)x^5 + 4b^3(7b^2s^2 + a)x^6 + 8b^4s^2x^7 + b^4x^8} dx$$

= Too large to display

input

```
int(1/((b*s^2+a)^4+8*b*s*(b*s^2+a)^3*x+4*b*(b*s^2+a)^2*(7*b*s^2+a)*x^2+8*b^2*s*(7*b^2*s^4+10*a*b*s^2+3*a^2)*x^3+2*b^2*(35*b^2*s^4+30*a*b*s^2+3*a^2)*x^4+8*b^3*s*(7*b*s^2+3*a)*x^5+4*b^3*(7*b*s^2+a)*x^6+8*b^4*s*x^7+b^4*x^8),x)
```

output

```
(30*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**3*s + 90*sqrt(b)
)*sqrt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**2*b*s**3 + 180*sqrt(b)*sq
rt(a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**2*b*s**2*x + 90*sqrt(b)*sqrt(
a)*atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a**2*b*s*x**2 + 90*sqrt(b)*sqrt(a)*
atan((b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**5 + 360*sqrt(b)*sqrt(a)*atan
((b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**4*x + 540*sqrt(b)*sqrt(a)*atan((
b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**3*x**2 + 360*sqrt(b)*sqrt(a)*atan(
(b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s**2*x**3 + 90*sqrt(b)*sqrt(a)*atan(
(b*s + b*x)/(sqrt(b)*sqrt(a)))*a*b**2*s*x**4 + 30*sqrt(b)*sqrt(a)*atan((b*
s + b*x)/(sqrt(b)*sqrt(a)))*b**3*s**7 + 180*sqrt(b)*sqrt(a)*atan((b*s + b*
x)/(sqrt(b)*sqrt(a)))*b**3*s**6*x + 450*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(
sqrt(b)*sqrt(a)))*b**3*s**5*x**2 + 600*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(s
qrt(b)*sqrt(a)))*b**3*s**4*x**3 + 450*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sq
rt(b)*sqrt(a)))*b**3*s**3*x**4 + 180*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqr
t(b)*sqrt(a)))*b**3*s**2*x**5 + 30*sqrt(b)*sqrt(a)*atan((b*s + b*x)/(sqrt(
b)*sqrt(a)))*b**3*s*x**6 - 5*a**4 + 51*a**3*b*s**2 + 36*a**3*b*s*x - 15*a*
*3*b*x**2 + 65*a**2*b**2*s**4 + 180*a**2*b**2*s**3*x + 150*a**2*b**2*s**2*
x**2 + 20*a**2*b**2*s*x**3 - 15*a**2*b**2*x**4 + 25*a*b**3*s**6 + 120*a*b*
*3*s**5*x + 225*a*b**3*s**4*x**2 + 200*a*b**3*s**3*x**3 + 75*a*b**3*s**2*x
**4 - 5*a*b**3*x**6)/(96*a**4*b*s*(a**3 + 3*a**2*b*s**2 + 6*a**2*b*s*x ...
```

### 3.39 $\int \left( a + \frac{d}{x^3} + \frac{c}{x^2} + \frac{b}{x} \right) dx$

Optimal result	311
Mathematica [A] (verified)	311
Rubi [A] (verified)	312
Maple [A] (verified)	312
Fricas [A] (verification not implemented)	313
Sympy [A] (verification not implemented)	313
Maxima [A] (verification not implemented)	314
Giac [A] (verification not implemented)	314
Mupad [B] (verification not implemented)	314
Reduce [B] (verification not implemented)	315

#### Optimal result

Integrand size = 17, antiderivative size = 22

$$\int \left( a + \frac{d}{x^3} + \frac{c}{x^2} + \frac{b}{x} \right) dx = -\frac{d}{2x^2} - \frac{c}{x} + ax + b \log(x)$$

output

```
-1/2*d/x^2-c/x+a*x+b*ln(x)
```

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \left( a + \frac{d}{x^3} + \frac{c}{x^2} + \frac{b}{x} \right) dx = -\frac{d}{2x^2} - \frac{c}{x} + ax + b \log(x)$$

input

```
Integrate[a + d/x^3 + c/x^2 + b/x,x]
```

output

```
-1/2*d/x^2 - c/x + a*x + b*Log[x]
```



**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( a + \frac{b}{x} + \frac{c}{x^2} + \frac{d}{x^3} \right) dx$$

$$\downarrow \text{2009}$$

$$ax + b \log(x) - \frac{c}{x} - \frac{d}{2x^2}$$

input `Int[a + d/x^3 + c/x^2 + b/x,x]`

output `-1/2*d/x^2 - c/x + a*x + b*Log[x]`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.95

method	result	size
default	$-\frac{d}{2x^2} - \frac{c}{x} + xa + b \ln(x)$	21
risch	$-\frac{d}{2x^2} - \frac{c}{x} + xa + b \ln(x)$	21
norman	$\frac{ax^3 - \frac{1}{2}d - cx}{x^2} + b \ln(x)$	23
parallelrisch	$\frac{2b \ln(x)x^2 - 2cx - d}{2x^2} + xa$	26

input `int(a+d/x^3+c/x^2+b/x,x,method=_RETURNVERBOSE)`

output `-1/2*d/x^2-c/x+x*a+b*ln(x)`

### **Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.23

$$\int \left( a + \frac{d}{x^3} + \frac{c}{x^2} + \frac{b}{x} \right) dx = \frac{2ax^3 + 2bx^2 \log(x) - 2cx - d}{2x^2}$$

input `integrate(a+d/x^3+c/x^2+b/x,x, algorithm="fricas")`

output `1/2*(2*a*x^3 + 2*b*x^2*log(x) - 2*c*x - d)/x^2`

### **Sympy [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \left( a + \frac{d}{x^3} + \frac{c}{x^2} + \frac{b}{x} \right) dx = ax + b \log(x) + \frac{-2cx - d}{2x^2}$$

input `integrate(a+d/x**3+c/x**2+b/x,x)`

output `a*x + b*log(x) + (-2*c*x - d)/(2*x**2)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \left( a + \frac{d}{x^3} + \frac{c}{x^2} + \frac{b}{x} \right) dx = ax + b \log(x) - \frac{c}{x} - \frac{d}{2x^2}$$

input `integrate(a+d/x^3+c/x^2+b/x,x, algorithm="maxima")`output `a*x + b*log(x) - c/x - 1/2*d/x^2`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.95

$$\int \left( a + \frac{d}{x^3} + \frac{c}{x^2} + \frac{b}{x} \right) dx = ax + b \log(|x|) - \frac{c}{x} - \frac{d}{2x^2}$$

input `integrate(a+d/x^3+c/x^2+b/x,x, algorithm="giac")`output `a*x + b*log(abs(x)) - c/x - 1/2*d/x^2`**Mupad [B] (verification not implemented)**

Time = 21.65 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \left( a + \frac{d}{x^3} + \frac{c}{x^2} + \frac{b}{x} \right) dx = ax - \frac{\frac{d}{2} + cx}{x^2} + b \ln(x)$$

input `int(a + b/x + c/x^2 + d/x^3,x)`output `a*x - (d/2 + c*x)/x^2 + b*log(x)`

**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.23

$$\int \left( a + \frac{d}{x^3} + \frac{c}{x^2} + \frac{b}{x} \right) dx = \frac{2 \log(x) b x^2 + 2a x^3 - 2cx - d}{2x^2}$$

input `int(a+d/x^3+c/x^2+b/x,x)`

output `(2*log(x)*b*x**2 + 2*a*x**3 - 2*c*x - d)/(2*x**2)`

### 3.40 $\int \left( \frac{1}{x^5} + x + x^5 \right) dx$

Optimal result	316
Mathematica [A] (verified)	316
Rubi [A] (verified)	317
Maple [A] (verified)	318
Fricas [A] (verification not implemented)	318
Sympy [A] (verification not implemented)	319
Maxima [A] (verification not implemented)	319
Giac [A] (verification not implemented)	319
Mupad [B] (verification not implemented)	320
Reduce [B] (verification not implemented)	320

#### Optimal result

Integrand size = 8, antiderivative size = 22

$$\int \left( \frac{1}{x^5} + x + x^5 \right) dx = -\frac{1}{4x^4} + \frac{x^2}{2} + \frac{x^6}{6}$$

output

```
-1/4/x^4+1/2*x^2+1/6*x^6
```

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \left( \frac{1}{x^5} + x + x^5 \right) dx = -\frac{1}{4x^4} + \frac{x^2}{2} + \frac{x^6}{6}$$

input

```
Integrate[x^(-5) + x + x^5,x]
```

output

```
-1/4*1/x^4 + x^2/2 + x^6/6
```

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( x^5 + \frac{1}{x^5} + x \right) dx$$

↓ 2009

$$\frac{x^6}{6} - \frac{1}{4x^4} + \frac{x^2}{2}$$

input

`Int[x^(-5) + x + x^5,x]`

output

`-1/4*1/x^4 + x^2/2 + x^6/6`**Defintions of rubi rules used**

rule 2009

`Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

method	result	size
default	$-\frac{1}{4x^4} + \frac{x^2}{2} + \frac{x^6}{6}$	17
norman	$\frac{\frac{1}{6}x^{10} + \frac{1}{2}x^6 - \frac{1}{4}}{x^4}$	17
risch	$-\frac{1}{4x^4} + \frac{x^2}{2} + \frac{x^6}{6}$	17
gospers	$\frac{2x^{10} + 6x^6 - 3}{12x^4}$	18
parallelrisch	$\frac{2x^{10} + 6x^6 - 3}{12x^4}$	18
orering	$\frac{(2x^{10} + 6x^6 - 3)x \left( \frac{1}{x^5} + x + x^5 \right)}{12x^{10} + 12x^6 + 12}$	34

input `int(1/x^5+x+x^5,x,method=_RETURNVERBOSE)`output `-1/4/x^4+1/2*x^2+1/6*x^6`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

$$\int \left( \frac{1}{x^5} + x + x^5 \right) dx = \frac{2x^{10} + 6x^6 - 3}{12x^4}$$

input `integrate(1/x^5+x+x^5,x, algorithm="fricas")`output `1/12*(2*x^10 + 6*x^6 - 3)/x^4`

**Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.68

$$\int \left( \frac{1}{x^5} + x + x^5 \right) dx = \frac{x^6}{6} + \frac{x^2}{2} - \frac{1}{4x^4}$$

input `integrate(1/x**5+x+x**5,x)`

output `x**6/6 + x**2/2 - 1/(4*x**4)`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \left( \frac{1}{x^5} + x + x^5 \right) dx = \frac{1}{6} x^6 + \frac{1}{2} x^2 - \frac{1}{4 x^4}$$

input `integrate(1/x^5+x+x^5,x, algorithm="maxima")`

output `1/6*x^6 + 1/2*x^2 - 1/4/x^4`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \left( \frac{1}{x^5} + x + x^5 \right) dx = \frac{1}{6} x^6 + \frac{1}{2} x^2 - \frac{1}{4 x^4}$$

input `integrate(1/x^5+x+x^5,x, algorithm="giac")`

output `1/6*x^6 + 1/2*x^2 - 1/4/x^4`



**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

$$\int \left( \frac{1}{x^5} + x + x^5 \right) dx = \frac{2x^{10} + 6x^6 - 3}{12x^4}$$

input `int(x + 1/x^5 + x^5,x)`

output `(6*x^6 + 2*x^10 - 3)/(12*x^4)`

**Reduce [B] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

$$\int \left( \frac{1}{x^5} + x + x^5 \right) dx = \frac{2x^{10} + 6x^6 - 3}{12x^4}$$

input `int(1/x^5+x+x^5,x)`

output `(2*x**10 + 6*x**6 - 3)/(12*x**4)`

### 3.41 $\int \left( \frac{1}{x^3} + \frac{1}{x^2} + \frac{1}{x} \right) dx$

Optimal result	321
Mathematica [A] (verified)	321
Rubi [A] (verified)	322
Maple [A] (verified)	322
Fricas [A] (verification not implemented)	323
Sympy [A] (verification not implemented)	323
Maxima [A] (verification not implemented)	324
Giac [A] (verification not implemented)	324
Mupad [B] (verification not implemented)	324
Reduce [B] (verification not implemented)	325

#### Optimal result

Integrand size = 10, antiderivative size = 15

$$\int \left( \frac{1}{x^3} + \frac{1}{x^2} + \frac{1}{x} \right) dx = -\frac{1}{2x^2} - \frac{1}{x} + \log(x)$$

output

```
-1/2/x^2-1/x+ln(x)
```

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \left( \frac{1}{x^3} + \frac{1}{x^2} + \frac{1}{x} \right) dx = -\frac{1}{2x^2} - \frac{1}{x} + \log(x)$$

input

```
Integrate[x^(-3) + x^(-2) + x^(-1),x]
```

output

```
-1/2*1/x^2 - x^(-1) + Log[x]
```

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( \frac{1}{x^3} + \frac{1}{x^2} + \frac{1}{x} \right) dx$$

$$\downarrow \text{2009}$$

$$-\frac{1}{2x^2} - \frac{1}{x} + \log(x)$$

input `Int[x^(-3) + x^(-2) + x^(-1),x]`

output `-1/2*1/x^2 - x^(-1) + Log[x]`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

method	result	size
norman	$-\frac{\frac{1}{2}-x}{x^2} + \ln(x)$	13
default	$-\frac{1}{2x^2} - \frac{1}{x} + \ln(x)$	14
risch	$-\frac{1}{2x^2} - \frac{1}{x} + \ln(x)$	14
parallelrisc	$\frac{2\ln(x)x^2-2x-1}{2x^2}$	18

input `int(1/x^3+1/x^2+1/x,x,method=_RETURNVERBOSE)`

output `(-1/2-x)/x^2+ln(x)`

### **Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \left( \frac{1}{x^3} + \frac{1}{x^2} + \frac{1}{x} \right) dx = \frac{2x^2 \log(x) - 2x - 1}{2x^2}$$

input `integrate(1/x^3+1/x^2+1/x,x, algorithm="fricas")`

output `1/2*(2*x^2*log(x) - 2*x - 1)/x^2`

### **Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \left( \frac{1}{x^3} + \frac{1}{x^2} + \frac{1}{x} \right) dx = \log(x) + \frac{-2x - 1}{2x^2}$$

input `integrate(1/x**3+1/x**2+1/x,x)`

output `log(x) + (-2*x - 1)/(2*x**2)`

**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \left( \frac{1}{x^3} + \frac{1}{x^2} + \frac{1}{x} \right) dx = -\frac{1}{x} - \frac{1}{2x^2} + \log(x)$$

input `integrate(1/x^3+1/x^2+1/x,x, algorithm="maxima")`output `-1/x - 1/2/x^2 + log(x)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \left( \frac{1}{x^3} + \frac{1}{x^2} + \frac{1}{x} \right) dx = -\frac{1}{x} - \frac{1}{2x^2} + \log(|x|)$$

input `integrate(1/x^3+1/x^2+1/x,x, algorithm="giac")`output `-1/x - 1/2/x^2 + log(abs(x))`**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \left( \frac{1}{x^3} + \frac{1}{x^2} + \frac{1}{x} \right) dx = \ln(x) - \frac{x + \frac{1}{2}}{x^2}$$

input `int(1/x + 1/x^2 + 1/x^3,x)`output `log(x) - (x + 1/2)/x^2`

**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \left( \frac{1}{x^3} + \frac{1}{x^2} + \frac{1}{x} \right) dx = \frac{2 \log(x) x^2 - 2x - 1}{2x^2}$$

input `int(1/x^3+1/x^2+1/x,x)`

output `(2*log(x)*x**2 - 2*x - 1)/(2*x**2)`

### 3.42 $\int \left(-\frac{2}{x^2} + \frac{3}{x}\right) dx$

Optimal result	326
Mathematica [A] (verified)	326
Rubi [A] (verified)	327
Maple [A] (verified)	327
Fricas [A] (verification not implemented)	328
Sympy [A] (verification not implemented)	328
Maxima [A] (verification not implemented)	329
Giac [A] (verification not implemented)	329
Mupad [B] (verification not implemented)	329
Reduce [B] (verification not implemented)	330

#### Optimal result

Integrand size = 11, antiderivative size = 10

$$\int \left(-\frac{2}{x^2} + \frac{3}{x}\right) dx = \frac{2}{x} + 3 \log(x)$$

output `2/x+3*ln(x)`

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \left(-\frac{2}{x^2} + \frac{3}{x}\right) dx = \frac{2}{x} + 3 \log(x)$$

input `Integrate[-2/x^2 + 3/x,x]`

output `2/x + 3*Log[x]`

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( \frac{3}{x} - \frac{2}{x^2} \right) dx$$

↓ 2009

$$\frac{2}{x} + 3 \log(x)$$

input `Int[-2/x^2 + 3/x,x]`

output `2/x + 3*Log[x]`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

method	result	size
default	$\frac{2}{x} + 3 \ln(x)$	11
norman	$\frac{2}{x} + 3 \ln(x)$	11
risch	$\frac{2}{x} + 3 \ln(x)$	11
parallelrisc	$\frac{3 \ln(x)x+2}{x}$	12



input `int(-2/x^2+3/x,x,method=_RETURNVERBOSE)`

output `2/x+3*ln(x)`

### **Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

$$\int \left( -\frac{2}{x^2} + \frac{3}{x} \right) dx = \frac{3x \log(x) + 2}{x}$$

input `integrate(-2/x^2+3/x,x, algorithm="fricas")`

output `(3*x*log(x) + 2)/x`

### **Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int \left( -\frac{2}{x^2} + \frac{3}{x} \right) dx = 3 \log(x) + \frac{2}{x}$$

input `integrate(-2/x**2+3/x,x)`

output `3*log(x) + 2/x`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \left( -\frac{2}{x^2} + \frac{3}{x} \right) dx = \frac{2}{x} + 3 \log(x)$$

input `integrate(-2/x^2+3/x,x, algorithm="maxima")`

output `2/x + 3*log(x)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

$$\int \left( -\frac{2}{x^2} + \frac{3}{x} \right) dx = \frac{2}{x} + 3 \log(|x|)$$

input `integrate(-2/x^2+3/x,x, algorithm="giac")`

output `2/x + 3*log(abs(x))`

**Mupad [B] (verification not implemented)**

Time = 21.89 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \left( -\frac{2}{x^2} + \frac{3}{x} \right) dx = 3 \ln(x) + \frac{2}{x}$$

input `int(3/x - 2/x^2,x)`

output `3*log(x) + 2/x`

**Reduce [B] (verification not implemented)**

Time = 0.37 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

$$\int \left( -\frac{2}{x^2} + \frac{3}{x} \right) dx = \frac{3 \log(x) x + 2}{x}$$

input `int(-2/x^2+3/x,x)`

output `(3*log(x)*x + 2)/x`

### 3.43 $\int \left(-\frac{1}{7x^6} + x^6\right) dx$

Optimal result	331
Mathematica [A] (verified)	331
Rubi [A] (verified)	332
Maple [A] (verified)	333
Fricas [A] (verification not implemented)	333
Sympy [A] (verification not implemented)	334
Maxima [A] (verification not implemented)	334
Giac [A] (verification not implemented)	334
Mupad [B] (verification not implemented)	335
Reduce [B] (verification not implemented)	335

#### Optimal result

Integrand size = 11, antiderivative size = 15

$$\int \left(-\frac{1}{7x^6} + x^6\right) dx = \frac{1}{35x^5} + \frac{x^7}{7}$$

output

```
1/35/x^5+1/7*x^7
```

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \left(-\frac{1}{7x^6} + x^6\right) dx = \frac{1}{35x^5} + \frac{x^7}{7}$$

input

```
Integrate[-1/7*1/x^6 + x^6,x]
```

output

```
1/(35*x^5) + x^7/7
```

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( x^6 - \frac{1}{7x^6} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{x^7}{7} + \frac{1}{35x^5}$$

input `Int[-1/7*1/x^6 + x^6,x]`

output `1/(35*x^5) + x^7/7`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

method	result	size
default	$\frac{1}{35x^5} + \frac{x^7}{7}$	12
norman	$\frac{\frac{1}{35} + \frac{x^{12}}{7}}{x^5}$	12
risch	$\frac{1}{35x^5} + \frac{x^7}{7}$	12
gosper	$\frac{5x^{12}+1}{35x^5}$	13
parallelrisc	$\frac{5x^{12}+1}{35x^5}$	13
orering	$\frac{(5x^{12}+1)x\left(-\frac{1}{7x^6}+x^6\right)}{35x^{12}-5}$	29

input `int(-1/7/x^6+x^6,x,method=_RETURNVERBOSE)`output `1/35/x^5+1/7*x^7`**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

$$\int \left( -\frac{1}{7x^6} + x^6 \right) dx = \frac{5x^{12} + 1}{35x^5}$$

input `integrate(-1/7/x^6+x^6,x, algorithm="fricas")`output `1/35*(5*x^12 + 1)/x^5`

**Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.67

$$\int \left( -\frac{1}{7x^6} + x^6 \right) dx = \frac{x^7}{7} + \frac{1}{35x^5}$$

input `integrate(-1/7/x**6+x**6,x)`output `x**7/7 + 1/(35*x**5)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \left( -\frac{1}{7x^6} + x^6 \right) dx = \frac{1}{7} x^7 + \frac{1}{35 x^5}$$

input `integrate(-1/7/x^6+x^6,x, algorithm="maxima")`output `1/7*x^7 + 1/35/x^5`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \left( -\frac{1}{7x^6} + x^6 \right) dx = \frac{1}{7} x^7 + \frac{1}{35 x^5}$$

input `integrate(-1/7/x^6+x^6,x, algorithm="giac")`output `1/7*x^7 + 1/35/x^5`

**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

$$\int \left( -\frac{1}{7x^6} + x^6 \right) dx = \frac{5x^{12} + 1}{35x^5}$$

input `int(x^6 - 1/(7*x^6),x)`output `(5*x^12 + 1)/(35*x^5)`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

$$\int \left( -\frac{1}{7x^6} + x^6 \right) dx = \frac{5x^{12} + 1}{35x^5}$$

input `int(-1/7/x^6+x^6,x)`output `(5*x**12 + 1)/(35*x**5)`



### 3.44 $\int \left(1 + \frac{1}{x} + x\right) dx$

Optimal result	336
Mathematica [A] (verified)	336
Rubi [A] (verified)	337
Maple [A] (verified)	337
Fricas [A] (verification not implemented)	338
Sympy [A] (verification not implemented)	338
Maxima [A] (verification not implemented)	339
Giac [A] (verification not implemented)	339
Mupad [B] (verification not implemented)	339
Reduce [B] (verification not implemented)	340

#### Optimal result

Integrand size = 6, antiderivative size = 11

$$\int \left(1 + \frac{1}{x} + x\right) dx = x + \frac{x^2}{2} + \log(x)$$

output

```
x+1/2*x^2+ln(x)
```

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \left(1 + \frac{1}{x} + x\right) dx = x + \frac{x^2}{2} + \log(x)$$

input

```
Integrate[1 + x^(-1) + x,x]
```

output

```
x + x^2/2 + Log[x]
```

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( x + \frac{1}{x} + 1 \right) dx$$

↓ 2009

$$\frac{x^2}{2} + x + \log(x)$$

input `Int[1 + x^(-1) + x,x]`

output `x + x^2/2 + Log[x]`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

method	result	size
default	$x + \frac{x^2}{2} + \ln(x)$	10
norman	$x + \frac{x^2}{2} + \ln(x)$	10
risch	$x + \frac{x^2}{2} + \ln(x)$	10
parallelrisch	$x + \frac{x^2}{2} + \ln(x)$	10

input `int(1+1/x+x,x,method=_RETURNVERBOSE)`

output `x+1/2*x^2+ln(x)`

### **Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \left( 1 + \frac{1}{x} + x \right) dx = \frac{1}{2} x^2 + x + \log(x)$$

input `integrate(1+1/x+x, x, algorithm="fricas")`

output `1/2*x^2 + x + log(x)`

### **Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int \left( 1 + \frac{1}{x} + x \right) dx = \frac{x^2}{2} + x + \log(x)$$

input `integrate(1+1/x+x, x)`

output `x**2/2 + x + log(x)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \left(1 + \frac{1}{x} + x\right) dx = \frac{1}{2}x^2 + x + \log(x)$$

input `integrate(1+1/x+x,x, algorithm="maxima")`output `1/2*x^2 + x + log(x)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

$$\int \left(1 + \frac{1}{x} + x\right) dx = \frac{1}{2}x^2 + x + \log(|x|)$$

input `integrate(1+1/x+x,x, algorithm="giac")`output `1/2*x^2 + x + log(abs(x))`**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \left(1 + \frac{1}{x} + x\right) dx = x + \ln(x) + \frac{x^2}{2}$$

input `int(x + 1/x + 1,x)`output `x + log(x) + x^2/2`

**Reduce [B] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \left( 1 + \frac{1}{x} + x \right) dx = \log(x) + \frac{x^2}{2} + x$$

input `int(1+1/x+x,x)`

output `(2*log(x) + x**2 + 2*x)/2`

### 3.45 $\int \left(-\frac{3}{x^3} + \frac{4}{x^2}\right) dx$

Optimal result	341
Mathematica [A] (verified)	341
Rubi [A] (verified)	342
Maple [A] (verified)	343
Fricas [A] (verification not implemented)	343
Sympy [A] (verification not implemented)	344
Maxima [A] (verification not implemented)	344
Giac [A] (verification not implemented)	344
Mupad [B] (verification not implemented)	345
Reduce [B] (verification not implemented)	345

#### Optimal result

Integrand size = 11, antiderivative size = 13

$$\int \left(-\frac{3}{x^3} + \frac{4}{x^2}\right) dx = \frac{3}{2x^2} - \frac{4}{x}$$

output  $3/2/x^2-4/x$

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \left(-\frac{3}{x^3} + \frac{4}{x^2}\right) dx = \frac{3}{2x^2} - \frac{4}{x}$$

input `Integrate[-3/x^3 + 4/x^2,x]`

output  $3/(2*x^2) - 4/x$

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( \frac{4}{x^2} - \frac{3}{x^3} \right) dx$$

↓ 2009

$$\frac{3}{2x^2} - \frac{4}{x}$$

input `Int[-3/x^3 + 4/x^2,x]`

output `3/(2*x^2) - 4/x`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

method	result	size
norman	$-\frac{4x+\frac{3}{2}}{x^2}$	10
gospers	$-\frac{8x-3}{2x^2}$	11
parallelrisc	$-\frac{8x+3}{2x^2}$	11
default	$\frac{3}{2x^2} - \frac{4}{x}$	12
risc	$\frac{3}{2x^2} - \frac{4}{x}$	12
orering	$-\frac{(8x-3)x\left(-\frac{3}{x^3} + \frac{4}{x^2}\right)}{2(4x-3)}$	27

input `int(-3/x^3+4/x^2,x,method=_RETURNVERBOSE)`output `(-4*x+3/2)/x^2`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \left( -\frac{3}{x^3} + \frac{4}{x^2} \right) dx = -\frac{8x-3}{2x^2}$$

input `integrate(-3/x^3+4/x^2,x, algorithm="fricas")`output `-1/2*(8*x - 3)/x^2`



**Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.62

$$\int \left( -\frac{3}{x^3} + \frac{4}{x^2} \right) dx = \frac{3 - 8x}{2x^2}$$

input `integrate(-3/x**3+4/x**2,x)`

output `(3 - 8*x)/(2*x**2)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \left( -\frac{3}{x^3} + \frac{4}{x^2} \right) dx = -\frac{4}{x} + \frac{3}{2x^2}$$

input `integrate(-3/x^3+4/x^2,x, algorithm="maxima")`

output `-4/x + 3/2/x^2`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \left( -\frac{3}{x^3} + \frac{4}{x^2} \right) dx = -\frac{4}{x} + \frac{3}{2x^2}$$

input `integrate(-3/x^3+4/x^2,x, algorithm="giac")`

output `-4/x + 3/2/x^2`

**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \left( -\frac{3}{x^3} + \frac{4}{x^2} \right) dx = -\frac{8x - 3}{2x^2}$$

input `int(4/x^2 - 3/x^3,x)`output `-(8*x - 3)/(2*x^2)`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \left( -\frac{3}{x^3} + \frac{4}{x^2} \right) dx = \frac{-8x + 3}{2x^2}$$

input `int(-3/x^3+4/x^2,x)`output `( - 8*x + 3)/(2*x**2)`

### 3.46 $\int \left(\frac{1}{x} + 2x + x^2\right) dx$

Optimal result . . . . .	346
Mathematica [A] (verified) . . . . .	346
Rubi [A] (verified) . . . . .	347
Maple [A] (verified) . . . . .	347
Fricas [A] (verification not implemented) . . . . .	348
Sympy [A] (verification not implemented) . . . . .	348
Maxima [A] (verification not implemented) . . . . .	349
Giac [A] (verification not implemented) . . . . .	349
Mupad [B] (verification not implemented) . . . . .	349
Reduce [B] (verification not implemented) . . . . .	350

#### Optimal result

Integrand size = 10, antiderivative size = 13

$$\int \left(\frac{1}{x} + 2x + x^2\right) dx = x^2 + \frac{x^3}{3} + \log(x)$$

output

```
x^2+1/3*x^3+ln(x)
```

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \left(\frac{1}{x} + 2x + x^2\right) dx = x^2 + \frac{x^3}{3} + \log(x)$$

input

```
Integrate[x^(-1) + 2*x + x^2,x]
```

output

```
x^2 + x^3/3 + Log[x]
```

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( x^2 + 2x + \frac{1}{x} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{x^3}{3} + x^2 + \log(x)$$

input `Int[x^(-1) + 2*x + x^2,x]`

output `x^2 + x^3/3 + Log[x]`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
default	$x^2 + \frac{x^3}{3} + \ln(x)$	12
norman	$x^2 + \frac{x^3}{3} + \ln(x)$	12
risch	$x^2 + \frac{x^3}{3} + \ln(x)$	12
parallelrisc	$x^2 + \frac{x^3}{3} + \ln(x)$	12

input `int(1/x+2*x+x^2,x,method=_RETURNVERBOSE)`

output `x^2+1/3*x^3+ln(x)`

### **Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \left( \frac{1}{x} + 2x + x^2 \right) dx = \frac{1}{3}x^3 + x^2 + \log(x)$$

input `integrate(1/x+2*x+x^2,x, algorithm="fricas")`

output `1/3*x^3 + x^2 + log(x)`

### **Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \left( \frac{1}{x} + 2x + x^2 \right) dx = \frac{x^3}{3} + x^2 + \log(x)$$

input `integrate(1/x+2*x+x**2,x)`

output `x**3/3 + x**2 + log(x)`

**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \left( \frac{1}{x} + 2x + x^2 \right) dx = \frac{1}{3}x^3 + x^2 + \log(x)$$

input `integrate(1/x+2*x+x^2,x, algorithm="maxima")`output `1/3*x^3 + x^2 + log(x)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

$$\int \left( \frac{1}{x} + 2x + x^2 \right) dx = \frac{1}{3}x^3 + x^2 + \log(|x|)$$

input `integrate(1/x+2*x+x^2,x, algorithm="giac")`output `1/3*x^3 + x^2 + log(abs(x))`**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \left( \frac{1}{x} + 2x + x^2 \right) dx = \ln(x) + x^2 + \frac{x^3}{3}$$

input `int(2*x + 1/x + x^2,x)`output `log(x) + x^2 + x^3/3`

**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \left( \frac{1}{x} + 2x + x^2 \right) dx = \log(x) + \frac{x^3}{3} + x^2$$

input `int(1/x+2*x+x^2,x)`

output `(3*log(x) + x**3 + 3*x**2)/3`

### 3.47 $\int (x^{5/6} - x^3) dx$

Optimal result	351
Mathematica [A] (verified)	351
Rubi [A] (verified)	352
Maple [A] (verified)	353
Fricas [A] (verification not implemented)	353
Sympy [A] (verification not implemented)	354
Maxima [A] (verification not implemented)	354
Giac [A] (verification not implemented)	354
Mupad [B] (verification not implemented)	355
Reduce [B] (verification not implemented)	355

#### Optimal result

Integrand size = 11, antiderivative size = 17

$$\int (x^{5/6} - x^3) dx = \frac{6x^{11/6}}{11} - \frac{x^4}{4}$$

output `6/11*x^(11/6)-1/4*x^4`

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int (x^{5/6} - x^3) dx = \frac{6x^{11/6}}{11} - \frac{x^4}{4}$$

input `Integrate[x^(5/6) - x^3,x]`

output `(6*x^(11/6))/11 - x^4/4`



**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (x^{5/6} - x^3) dx$$

$$\downarrow \text{2009}$$

$$\frac{6x^{11/6}}{11} - \frac{x^4}{4}$$

input `Int[x^(5/6) - x^3,x]`

output `(6*x^(11/6))/11 - x^4/4`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

method	result	size
derivativedivides	$\frac{6x^{\frac{11}{6}}}{11} - \frac{x^4}{4}$	12
default	$\frac{6x^{\frac{11}{6}}}{11} - \frac{x^4}{4}$	12
risch	$\frac{6x^{\frac{11}{6}}}{11} - \frac{x^4}{4}$	12
parts	$\frac{6x^{\frac{11}{6}}}{11} - \frac{x^4}{4}$	12
trager	$-\frac{(x^3+x^2+x+1)(x-1)}{4} + \frac{6x^{\frac{11}{6}}}{11}$	21
orering	$\frac{29x(x^{\frac{5}{6}}-x^3)}{44} - \frac{3x^2\left(\frac{5}{6x^{\frac{6}}}-3x^2\right)}{22}$	30

input `int(x^(5/6)-x^3,x,method=_RETURNVERBOSE)`output `6/11*x^(11/6)-1/4*x^4`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int (x^{5/6} - x^3) dx = -\frac{1}{4}x^4 + \frac{6}{11}x^{\frac{11}{6}}$$

input `integrate(x^(5/6)-x^3,x, algorithm="fricas")`output `-1/4*x^4 + 6/11*x^(11/6)`

**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

$$\int (x^{5/6} - x^3) dx = \frac{6x^{11/6}}{11} - \frac{x^4}{4}$$

input `integrate(x**(5/6)-x**3,x)`

output `6*x**(11/6)/11 - x**4/4`

**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int (x^{5/6} - x^3) dx = -\frac{1}{4}x^4 + \frac{6}{11}x^{11/6}$$

input `integrate(x^(5/6)-x^3,x, algorithm="maxima")`

output `-1/4*x^4 + 6/11*x^(11/6)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int (x^{5/6} - x^3) dx = -\frac{1}{4}x^4 + \frac{6}{11}x^{11/6}$$

input `integrate(x^(5/6)-x^3,x, algorithm="giac")`

output `-1/4*x^4 + 6/11*x^(11/6)`

**Mupad [B] (verification not implemented)**

Time = 21.82 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int (x^{5/6} - x^3) dx = \frac{6x^{11/6}}{11} - \frac{x^4}{4}$$

input `int(x^(5/6) - x^3,x)`

output `(6*x^(11/6))/11 - x^4/4`

**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int (x^{5/6} - x^3) dx = \frac{x(24x^{5/6} - 11x^3)}{44}$$

input `int(x^(5/6)-x^3,x)`

output `(x*(24*x**(5/6) - 11*x**3))/44`

### 3.48 $\int (33 + \sqrt[33]{x}) dx$

Optimal result	356
Mathematica [A] (verified)	356
Rubi [A] (verified)	357
Maple [A] (verified)	358
Fricas [A] (verification not implemented)	358
Sympy [A] (verification not implemented)	359
Maxima [A] (verification not implemented)	359
Giac [A] (verification not implemented)	359
Mupad [B] (verification not implemented)	360
Reduce [B] (verification not implemented)	360

#### Optimal result

Integrand size = 7, antiderivative size = 13

$$\int (33 + \sqrt[33]{x}) dx = 33x + \frac{33x^{34/33}}{34}$$

output `33*x+33/34*x^(34/33)`

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int (33 + \sqrt[33]{x}) dx = 33x + \frac{33x^{34/33}}{34}$$

input `Integrate[33 + x^(1/33),x]`

output `33*x + (33*x^(34/33))/34`

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\sqrt[33]{x} + 33) dx$$

$$\downarrow \text{2009}$$

$$\frac{33x^{34/33}}{34} + 33x$$

input `Int[33 + x^(1/33), x]`

output `33*x + (33*x^(34/33))/34`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

method	result	size
derivativedivides	$33x + \frac{33x^{34}}{34}$	10
default	$33x + \frac{33x^{34}}{34}$	10
risch	$33x + \frac{33x^{34}}{34}$	10
parts	$33x + \frac{33x^{34}}{34}$	10
trager	$33x - 33 + \frac{33x^{34}}{34}$	11
orering	$x\left(33 + x^{\frac{1}{33}}\right) - \frac{x^{34}}{34}$	14

input `int(33+x^(1/33),x,method=_RETURNVERBOSE)`output `33*x+33/34*x^(34/33)`**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int (33 + \sqrt[33]{x}) dx = \frac{33}{34} x^{\frac{34}{33}} + 33x$$

input `integrate(33+x^(1/33),x, algorithm="fricas")`output `33/34*x^(34/33) + 33*x`

**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int (33 + \sqrt[33]{x}) dx = \frac{33x^{34}}{34} + 33x$$

input `integrate(33+x**(1/33),x)`

output `33*x**(34/33)/34 + 33*x`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int (33 + \sqrt[33]{x}) dx = \frac{33}{34} x^{34} + 33x$$

input `integrate(33+x^(1/33),x, algorithm="maxima")`

output `33/34*x^(34/33) + 33*x`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int (33 + \sqrt[33]{x}) dx = \frac{33}{34} x^{34} + 33x$$

input `integrate(33+x^(1/33),x, algorithm="giac")`

output `33/34*x^(34/33) + 33*x`



**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.62

$$\int (33 + \sqrt[33]{x}) dx = \frac{33x(x^{1/33} + 34)}{34}$$

input `int(x^(1/33) + 33,x)`

output `(33*x*(x^(1/33) + 34))/34`

**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.62

$$\int (33 + \sqrt[33]{x}) dx = \frac{33x(x^{\frac{1}{33}} + 34)}{34}$$

input `int(33+x^(1/33),x)`

output `(33*x*(x**(1/33) + 34))/34`

$$3.49 \quad \int \left( \frac{1}{2\sqrt{x}} + 2\sqrt{x} \right) dx$$

Optimal result	361
Mathematica [A] (verified)	361
Rubi [A] (verified)	362
Maple [A] (verified)	363
Fricas [A] (verification not implemented)	363
Sympy [A] (verification not implemented)	364
Maxima [A] (verification not implemented)	364
Giac [A] (verification not implemented)	364
Mupad [B] (verification not implemented)	365
Reduce [B] (verification not implemented)	365

### Optimal result

Integrand size = 17, antiderivative size = 15

$$\int \left( \frac{1}{2\sqrt{x}} + 2\sqrt{x} \right) dx = \sqrt{x} + \frac{4x^{3/2}}{3}$$

output `x^(1/2)+4/3*x^(3/2)`

### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \left( \frac{1}{2\sqrt{x}} + 2\sqrt{x} \right) dx = \frac{1}{3}\sqrt{x}(3 + 4x)$$

input `Integrate[1/(2*Sqrt[x]) + 2*Sqrt[x],x]`

output `(Sqrt[x]*(3 + 4*x))/3`

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( 2\sqrt{x} + \frac{1}{2\sqrt{x}} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{4x^{3/2}}{3} + \sqrt{x}$$

input `Int[1/(2*Sqrt[x]) + 2*Sqrt[x],x]`

output `Sqrt[x] + (4*x^(3/2))/3`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.67

method	result	size
derivativedivides	$\sqrt{x} + \frac{4x^{\frac{3}{2}}}{3}$	10
default	$\sqrt{x} + \frac{4x^{\frac{3}{2}}}{3}$	10
risch	$\sqrt{x} + \frac{4x^{\frac{3}{2}}}{3}$	10
gosper	$\frac{\sqrt{x}(3+4x)}{3}$	11
trager	$\frac{(2+\frac{8x}{3})\sqrt{x}}{2}$	11
orering	$\frac{2(3+4x)x(\frac{1}{2\sqrt{x}}+2\sqrt{x})}{3(4x+1)}$	27

input `int(1/2/x^(1/2)+2*x^(1/2),x,method=_RETURNVERBOSE)`output `x^(1/2)+4/3*x^(3/2)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.67

$$\int \left( \frac{1}{2\sqrt{x}} + 2\sqrt{x} \right) dx = \frac{1}{3} (4x + 3)\sqrt{x}$$

input `integrate(1/2/x^(1/2)+2*x^(1/2),x, algorithm="fricas")`output `1/3*(4*x + 3)*sqrt(x)`

**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

$$\int \left( \frac{1}{2\sqrt{x}} + 2\sqrt{x} \right) dx = \frac{4x^{\frac{3}{2}}}{3} + \sqrt{x}$$

input `integrate(1/2/x**(1/2)+2*x**(1/2),x)`

output `4*x**(3/2)/3 + sqrt(x)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.60

$$\int \left( \frac{1}{2\sqrt{x}} + 2\sqrt{x} \right) dx = \frac{4}{3} x^{\frac{3}{2}} + \sqrt{x}$$

input `integrate(1/2/x^(1/2)+2*x^(1/2),x, algorithm="maxima")`

output `4/3*x^(3/2) + sqrt(x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.60

$$\int \left( \frac{1}{2\sqrt{x}} + 2\sqrt{x} \right) dx = \frac{4}{3} x^{\frac{3}{2}} + \sqrt{x}$$

input `integrate(1/2/x^(1/2)+2*x^(1/2),x, algorithm="giac")`

output `4/3*x^(3/2) + sqrt(x)`

**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.67

$$\int \left( \frac{1}{2\sqrt{x}} + 2\sqrt{x} \right) dx = \frac{\sqrt{x}(4x + 3)}{3}$$

input `int(1/(2*x^(1/2)) + 2*x^(1/2),x)`

output `(x^(1/2)*(4*x + 3))/3`

**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.60

$$\int \left( \frac{1}{2\sqrt{x}} + 2\sqrt{x} \right) dx = \frac{\sqrt{x}(4x + 3)}{3}$$

input `int(1/2/x^(1/2)+2*x^(1/2),x)`

output `(sqrt(x)*(4*x + 3))/3`

$$3.50 \quad \int \left( -\frac{1}{x^2} + \frac{10}{x} + 6\sqrt{x} \right) dx$$

Optimal result	366
Mathematica [A] (verified)	366
Rubi [A] (verified)	367
Maple [A] (verified)	367
Fricas [A] (verification not implemented)	368
Sympy [A] (verification not implemented)	368
Maxima [A] (verification not implemented)	369
Giac [A] (verification not implemented)	369
Mupad [B] (verification not implemented)	369
Reduce [B] (verification not implemented)	370

### Optimal result

Integrand size = 18, antiderivative size = 15

$$\int \left( -\frac{1}{x^2} + \frac{10}{x} + 6\sqrt{x} \right) dx = \frac{1}{x} + 4x^{3/2} + 10 \log(x)$$

output `1/x+4*x^(3/2)+10*ln(x)`

### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \left( -\frac{1}{x^2} + \frac{10}{x} + 6\sqrt{x} \right) dx = \frac{1}{x} + 4x^{3/2} + 10 \log(x)$$

input `Integrate[-x^(-2) + 10/x + 6*Sqrt[x],x]`

output `x^(-1) + 4*x^(3/2) + 10*Log[x]`

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( -\frac{1}{x^2} + 6\sqrt{x} + \frac{10}{x} \right) dx$$

$$\downarrow \text{2009}$$

$$4x^{3/2} + \frac{1}{x} + 10 \log(x)$$

input `Int[-x^(-2) + 10/x + 6*Sqrt[x],x]`

output `x^(-1) + 4*x^(3/2) + 10*Log[x]`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

method	result	size
derivativedivides	$\frac{1}{x} + 4x^{\frac{3}{2}} + 10 \ln(x)$	14
default	$\frac{1}{x} + 4x^{\frac{3}{2}} + 10 \ln(x)$	14
risch	$\frac{1}{x} + 4x^{\frac{3}{2}} + 10 \ln(x)$	14
trager	$-\frac{x-1}{x} + 4x^{\frac{3}{2}} + 10 \ln(x)$	19



input `int(-1/x^2+10/x+6*x^(1/2),x,method=_RETURNVERBOSE)`

output `1/x+4*x^(3/2)+10*ln(x)`

### **Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.20

$$\int \left( -\frac{1}{x^2} + \frac{10}{x} + 6\sqrt{x} \right) dx = \frac{4x^{\frac{5}{2}} + 20x \log(\sqrt{x}) + 1}{x}$$

input `integrate(-1/x^2+10/x+6*x^(1/2),x, algorithm="fricas")`

output `(4*x^(5/2) + 20*x*log(sqrt(x)) + 1)/x`

### **Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \left( -\frac{1}{x^2} + \frac{10}{x} + 6\sqrt{x} \right) dx = 4x^{\frac{3}{2}} + 10 \log(x) + \frac{1}{x}$$

input `integrate(-1/x**2+10/x+6*x**(1/2),x)`

output `4*x**(3/2) + 10*log(x) + 1/x`

**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \left( -\frac{1}{x^2} + \frac{10}{x} + 6\sqrt{x} \right) dx = 4x^{\frac{3}{2}} + \frac{1}{x} + 10 \log(x)$$

input `integrate(-1/x^2+10/x+6*x^(1/2),x, algorithm="maxima")`output `4*x^(3/2) + 1/x + 10*log(x)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \left( -\frac{1}{x^2} + \frac{10}{x} + 6\sqrt{x} \right) dx = 4x^{\frac{3}{2}} + \frac{1}{x} + 10 \log(|x|)$$

input `integrate(-1/x^2+10/x+6*x^(1/2),x, algorithm="giac")`output `4*x^(3/2) + 1/x + 10*log(abs(x))`**Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \left( -\frac{1}{x^2} + \frac{10}{x} + 6\sqrt{x} \right) dx = 20 \ln(\sqrt{x}) + \frac{1}{x} + 4x^{3/2}$$

input `int(10/x - 1/x^2 + 6*x^(1/2),x)`output `20*log(x^(1/2)) + 1/x + 4*x^(3/2)`

**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.20

$$\int \left( -\frac{1}{x^2} + \frac{10}{x} + 6\sqrt{x} \right) dx = \frac{4\sqrt{x} x^2 + 10 \log(x) x + 1}{x}$$

input `int(-1/x^2+10/x+6*x^(1/2),x)`

output `(4*sqrt(x)*x**2 + 10*log(x)*x + 1)/x`

### 3.51 $\int \left( \frac{1}{x^{3/2}} + x^{3/2} \right) dx$

Optimal result	371
Mathematica [A] (verified)	371
Rubi [A] (verified)	372
Maple [A] (verified)	373
Fricas [A] (verification not implemented)	373
Sympy [A] (verification not implemented)	374
Maxima [A] (verification not implemented)	374
Giac [A] (verification not implemented)	374
Mupad [B] (verification not implemented)	375
Reduce [B] (verification not implemented)	375

#### Optimal result

Integrand size = 11, antiderivative size = 17

$$\int \left( \frac{1}{x^{3/2}} + x^{3/2} \right) dx = -\frac{2}{\sqrt{x}} + \frac{2x^{5/2}}{5}$$

output `-2/x^(1/2)+2/5*x^(5/2)`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \left( \frac{1}{x^{3/2}} + x^{3/2} \right) dx = \frac{2(-5 + x^3)}{5\sqrt{x}}$$

input `Integrate[x^(-3/2) + x^(3/2), x]`

output `(2*(-5 + x^3))/(5*Sqrt[x])`

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( x^{3/2} + \frac{1}{x^{3/2}} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{2x^{5/2}}{5} - \frac{2}{\sqrt{x}}$$

input `Int[x^(-3/2) + x^(3/2),x]`

output `-2/Sqrt[x] + (2*x^(5/2))/5`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

method	result	size
gospers	$\frac{2x^3-2}{5\sqrt{x}}$	11
trager	$\frac{2x^3-2}{5\sqrt{x}}$	11
derivativdivides	$-\frac{2}{\sqrt{x}} + \frac{2x^{\frac{5}{2}}}{5}$	12
default	$-\frac{2}{\sqrt{x}} + \frac{2x^{\frac{5}{2}}}{5}$	12
risch	$-\frac{2}{\sqrt{x}} + \frac{2x^{\frac{5}{2}}}{5}$	12
orering	$\frac{2(x^3-5)x\left(\frac{1}{3}+x^{\frac{3}{2}}\right)}{5(x+1)(x^2-x+1)}$	31

input `int(1/x^(3/2)+x^(3/2),x,method=_RETURNVERBOSE)`output `2/5*(x^3-5)/x^(1/2)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.59

$$\int \left( \frac{1}{x^{3/2}} + x^{3/2} \right) dx = \frac{2(x^3 - 5)}{5\sqrt{x}}$$

input `integrate(1/x^(3/2)+x^(3/2),x, algorithm="fricas")`output `2/5*(x^3 - 5)/sqrt(x)`

**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \left( \frac{1}{x^{3/2}} + x^{3/2} \right) dx = \frac{2x^{5/2}}{5} - \frac{2}{\sqrt{x}}$$

input `integrate(1/x**(3/2)+x**(3/2),x)`

output `2*x**(5/2)/5 - 2/sqrt(x)`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int \left( \frac{1}{x^{3/2}} + x^{3/2} \right) dx = \frac{2}{5} x^{5/2} - \frac{2}{\sqrt{x}}$$

input `integrate(1/x^(3/2)+x^(3/2),x, algorithm="maxima")`

output `2/5*x^(5/2) - 2/sqrt(x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int \left( \frac{1}{x^{3/2}} + x^{3/2} \right) dx = \frac{2}{5} x^{5/2} - \frac{2}{\sqrt{x}}$$

input `integrate(1/x^(3/2)+x^(3/2),x, algorithm="giac")`

output `2/5*x^(5/2) - 2/sqrt(x)`

**Mupad [B] (verification not implemented)**

Time = 21.78 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

$$\int \left( \frac{1}{x^{3/2}} + x^{3/2} \right) dx = \frac{2x^3 - 10}{5\sqrt{x}}$$

input `int(1/x^(3/2) + x^(3/2),x)`output `(2*x^3 - 10)/(5*x^(1/2))`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int \left( \frac{1}{x^{3/2}} + x^{3/2} \right) dx = \frac{2x^3 - 2}{\sqrt{x}}$$

input `int(1/x^(3/2)+x^(3/2),x)`output `(2*(x**3 - 5))/(5*sqrt(x))`



### 3.52 $\int (-5x^{3/2} + 7x^{5/2}) dx$

Optimal result	376
Mathematica [A] (verified)	376
Rubi [A] (verified)	377
Maple [A] (verified)	378
Fricas [A] (verification not implemented)	378
Sympy [A] (verification not implemented)	379
Maxima [A] (verification not implemented)	379
Giac [A] (verification not implemented)	379
Mupad [B] (verification not implemented)	380
Reduce [B] (verification not implemented)	380

#### Optimal result

Integrand size = 15, antiderivative size = 15

$$\int (-5x^{3/2} + 7x^{5/2}) dx = -2x^{5/2} + 2x^{7/2}$$

output `-2*x^(5/2)+2*x^(7/2)`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.67

$$\int (-5x^{3/2} + 7x^{5/2}) dx = 2(-1 + x)x^{5/2}$$

input `Integrate[-5*x^(3/2) + 7*x^(5/2), x]`

output `2*(-1 + x)*x^(5/2)`

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (7x^{5/2} - 5x^{3/2}) dx$$

$$\downarrow \text{2009}$$

$$2x^{7/2} - 2x^{5/2}$$

input `Int[-5*x^(3/2) + 7*x^(5/2),x]`

output `-2*x^(5/2) + 2*x^(7/2)`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.60

method	result	size
gospers	$2x^{\frac{5}{2}}(x-1)$	9
trager	$2x^{\frac{5}{2}}(x-1)$	9
derivativdivides	$-2x^{\frac{5}{2}} + 2x^{\frac{7}{2}}$	12
default	$-2x^{\frac{5}{2}} + 2x^{\frac{7}{2}}$	12
risch	$-2x^{\frac{5}{2}} + 2x^{\frac{7}{2}}$	12
parts	$-2x^{\frac{5}{2}} + 2x^{\frac{7}{2}}$	12
orering	$\frac{2x(x-1)(-5x^{\frac{3}{2}}+7x^{\frac{5}{2}})}{7x-5}$	25

input `int(-5*x^(3/2)+7*x^(5/2),x,method=_RETURNVERBOSE)`output `2*x^(5/2)*(x-1)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int (-5x^{3/2} + 7x^{5/2}) dx = 2(x^3 - x^2)\sqrt{x}$$

input `integrate(-5*x^(3/2)+7*x^(5/2),x, algorithm="fricas")`output `2*(x^3 - x^2)*sqrt(x)`

**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

$$\int (-5x^{3/2} + 7x^{5/2}) dx = 2x^{7/2} - 2x^{5/2}$$

input `integrate(-5*x**(3/2)+7*x**(5/2),x)`

output `2*x**(7/2) - 2*x**(5/2)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int (-5x^{3/2} + 7x^{5/2}) dx = 2x^{7/2} - 2x^{5/2}$$

input `integrate(-5*x^(3/2)+7*x^(5/2),x, algorithm="maxima")`

output `2*x^(7/2) - 2*x^(5/2)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int (-5x^{3/2} + 7x^{5/2}) dx = 2x^{7/2} - 2x^{5/2}$$

input `integrate(-5*x^(3/2)+7*x^(5/2),x, algorithm="giac")`

output `2*x^(7/2) - 2*x^(5/2)`

**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.53

$$\int (-5x^{3/2} + 7x^{5/2}) dx = 2x^{5/2}(x - 1)$$

input `int(7*x^(5/2) - 5*x^(3/2),x)`

output `2*x^(5/2)*(x - 1)`

**Reduce [B] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.67

$$\int (-5x^{3/2} + 7x^{5/2}) dx = 2\sqrt{x}x^2(x - 1)$$

input `int(-5*x^(3/2)+7*x^(5/2),x)`

output `2*sqrt(x)*x**2*(x - 1)`

$$\mathbf{3.53} \quad \int \left( \frac{2}{\sqrt{x}} + \sqrt{x} - \frac{x}{2} \right) dx$$

Optimal result	381
Mathematica [A] (verified)	381
Rubi [A] (verified)	382
Maple [A] (verified)	383
Fricas [A] (verification not implemented)	383
Sympy [A] (verification not implemented)	384
Maxima [A] (verification not implemented)	384
Giac [A] (verification not implemented)	384
Mupad [B] (verification not implemented)	385
Reduce [B] (verification not implemented)	385

### Optimal result

Integrand size = 18, antiderivative size = 24

$$\int \left( \frac{2}{\sqrt{x}} + \sqrt{x} - \frac{x}{2} \right) dx = 4\sqrt{x} + \frac{2x^{3/2}}{3} - \frac{x^2}{4}$$

output `4*x^(1/2)+2/3*x^(3/2)-1/4*x^2`

### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \left( \frac{2}{\sqrt{x}} + \sqrt{x} - \frac{x}{2} \right) dx = 4\sqrt{x} + \frac{2x^{3/2}}{3} - \frac{x^2}{4}$$

input `Integrate[2/Sqrt[x] + Sqrt[x] - x/2,x]`

output `4*Sqrt[x] + (2*x^(3/2))/3 - x^2/4`

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( -\frac{x}{2} + \sqrt{x} + \frac{2}{\sqrt{x}} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{2x^{3/2}}{3} - \frac{x^2}{4} + 4\sqrt{x}$$

input `Int[2/Sqrt[x] + Sqrt[x] - x/2,x]`

output `4*Sqrt[x] + (2*x^(3/2))/3 - x^2/4`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.71

method	result	size
derivativedivides	$4\sqrt{x} + \frac{2x^{\frac{3}{2}}}{3} - \frac{x^2}{4}$	17
default	$4\sqrt{x} + \frac{2x^{\frac{3}{2}}}{3} - \frac{x^2}{4}$	17
risch	$4\sqrt{x} + \frac{2x^{\frac{3}{2}}}{3} - \frac{x^2}{4}$	17
trager	$-\frac{(x-1)(x+1)}{4} + \frac{(\frac{4x}{3}+8)\sqrt{x}}{2}$	20
orering	$\frac{x(5x+54)\left(\frac{2}{\sqrt{x}}+\sqrt{x}-\frac{x}{2}\right)}{6x+36} - \frac{x^2(x+18)\left(-\frac{1}{3}+\frac{1}{2\sqrt{x}}-\frac{1}{2}\right)}{3(x+6)}$	52

input `int(2/x^(1/2)+x^(1/2)-1/2*x,x,method=_RETURNVERBOSE)`output `4*x^(1/2)+2/3*x^(3/2)-1/4*x^2`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.58

$$\int \left( \frac{2}{\sqrt{x}} + \sqrt{x} - \frac{x}{2} \right) dx = -\frac{1}{4}x^2 + \frac{2}{3}(x+6)\sqrt{x}$$

input `integrate(2/x^(1/2)+x^(1/2)-1/2*x,x, algorithm="fricas")`output `-1/4*x^2 + 2/3*(x + 6)*sqrt(x)`



**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.79

$$\int \left( \frac{2}{\sqrt{x}} + \sqrt{x} - \frac{x}{2} \right) dx = \frac{2x^{\frac{3}{2}}}{3} + 4\sqrt{x} - \frac{x^2}{4}$$

input `integrate(2/x**(1/2)+x**(1/2)-1/2*x,x)`output `2*x**(3/2)/3 + 4*sqrt(x) - x**2/4`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int \left( \frac{2}{\sqrt{x}} + \sqrt{x} - \frac{x}{2} \right) dx = -\frac{1}{4}x^2 + \frac{2}{3}x^{\frac{3}{2}} + 4\sqrt{x}$$

input `integrate(2/x^(1/2)+x^(1/2)-1/2*x,x, algorithm="maxima")`output `-1/4*x^2 + 2/3*x^(3/2) + 4*sqrt(x)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int \left( \frac{2}{\sqrt{x}} + \sqrt{x} - \frac{x}{2} \right) dx = -\frac{1}{4}x^2 + \frac{2}{3}x^{\frac{3}{2}} + 4\sqrt{x}$$

input `integrate(2/x^(1/2)+x^(1/2)-1/2*x,x, algorithm="giac")`output `-1/4*x^2 + 2/3*x^(3/2) + 4*sqrt(x)`

**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.62

$$\int \left( \frac{2}{\sqrt{x}} + \sqrt{x} - \frac{x}{2} \right) dx = \frac{\sqrt{x} (8x - 3x^{3/2} + 48)}{12}$$

input `int(2/x^(1/2) - x/2 + x^(1/2),x)`output `(x^(1/2)*(8*x - 3*x^(3/2) + 48))/12`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.62

$$\int \left( \frac{2}{\sqrt{x}} + \sqrt{x} - \frac{x}{2} \right) dx = \frac{2\sqrt{x}x}{3} + 4\sqrt{x} - \frac{x^2}{4}$$

input `int(2/x^(1/2)+x^(1/2)-1/2*x,x)`output `(8*sqrt(x)*x + 48*sqrt(x) - 3*x**2)/12`

$$3.54 \quad \int \left( -\frac{2}{x} + \frac{\sqrt{x}}{5} + x^{3/2} \right) dx$$

Optimal result . . . . .	386
Mathematica [A] (verified) . . . . .	386
Rubi [A] (verified) . . . . .	387
Maple [A] (verified) . . . . .	387
Fricas [A] (verification not implemented) . . . . .	388
Sympy [A] (verification not implemented) . . . . .	388
Maxima [A] (verification not implemented) . . . . .	389
Giac [A] (verification not implemented) . . . . .	389
Mupad [B] (verification not implemented) . . . . .	389
Reduce [B] (verification not implemented) . . . . .	390

### Optimal result

Integrand size = 20, antiderivative size = 23

$$\int \left( -\frac{2}{x} + \frac{\sqrt{x}}{5} + x^{3/2} \right) dx = \frac{2x^{3/2}}{15} + \frac{2x^{5/2}}{5} - 2\log(x)$$

output `2/15*x^(3/2)+2/5*x^(5/2)-2*ln(x)`

### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \left( -\frac{2}{x} + \frac{\sqrt{x}}{5} + x^{3/2} \right) dx = \frac{2x^{3/2}}{15} + \frac{2x^{5/2}}{5} - 2\log(x)$$

input `Integrate[-2/x + Sqrt[x]/5 + x^(3/2), x]`

output `(2*x^(3/2))/15 + (2*x^(5/2))/5 - 2*Log[x]`

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( x^{3/2} + \frac{\sqrt{x}}{5} - \frac{2}{x} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{2x^{5/2}}{5} + \frac{2x^{3/2}}{15} - 2\log(x)$$

input `Int[-2/x + Sqrt[x]/5 + x^(3/2),x]`

output `(2*x^(3/2))/15 + (2*x^(5/2))/5 - 2*Log[x]`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

**Maple [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.70

method	result	size
derivativedivides	$\frac{2x^{3/2}}{15} + \frac{2x^{5/2}}{5} - 2\ln(x)$	16
default	$\frac{2x^{3/2}}{15} + \frac{2x^{5/2}}{5} - 2\ln(x)$	16
trager	$\frac{2x^{3/2}(1+3x)}{15} - 2\ln(x)$	16
risch	$\frac{2x^{3/2}}{15} + \frac{2x^{5/2}}{5} - 2\ln(x)$	16

input `int(-2/x+1/5*x^(1/2)+x^(3/2),x,method=_RETURNVERBOSE)`

output `2/15*x^(3/2)+2/5*x^(5/2)-2*ln(x)`

### **Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \left( -\frac{2}{x} + \frac{\sqrt{x}}{5} + x^{3/2} \right) dx = \frac{2}{15} (3x^2 + x)\sqrt{x} - 4 \log(\sqrt{x})$$

input `integrate(-2/x+1/5*x^(1/2)+x^(3/2),x, algorithm="fricas")`

output `2/15*(3*x^2 + x)*sqrt(x) - 4*log(sqrt(x))`

### **Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \left( -\frac{2}{x} + \frac{\sqrt{x}}{5} + x^{3/2} \right) dx = \frac{2x^{5/2}}{5} + \frac{2x^{3/2}}{15} - 2 \log(x)$$

input `integrate(-2/x+1/5*x**(1/2)+x**(3/2),x)`

output `2*x**(5/2)/5 + 2*x**(3/2)/15 - 2*log(x)`

**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int \left( -\frac{2}{x} + \frac{\sqrt{x}}{5} + x^{3/2} \right) dx = \frac{2}{5} x^{5/2} + \frac{2}{15} x^{3/2} - 2 \log(x)$$

input `integrate(-2/x+1/5*x^(1/2)+x^(3/2),x, algorithm="maxima")`output `2/5*x^(5/2) + 2/15*x^(3/2) - 2*log(x)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.70

$$\int \left( -\frac{2}{x} + \frac{\sqrt{x}}{5} + x^{3/2} \right) dx = \frac{2}{5} x^{5/2} + \frac{2}{15} x^{3/2} - 2 \log(|x|)$$

input `integrate(-2/x+1/5*x^(1/2)+x^(3/2),x, algorithm="giac")`output `2/5*x^(5/2) + 2/15*x^(3/2) - 2*log(abs(x))`**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \left( -\frac{2}{x} + \frac{\sqrt{x}}{5} + x^{3/2} \right) dx = \frac{2x^{3/2}}{15} - 4 \ln(\sqrt{x}) + \frac{2x^{5/2}}{5}$$

input `int(x^(1/2)/5 - 2/x + x^(3/2),x)`output `(2*x^(3/2))/15 - 4*log(x^(1/2)) + (2*x^(5/2))/5`

**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \left( -\frac{2}{x} + \frac{\sqrt{x}}{5} + x^{3/2} \right) dx = \frac{2\sqrt{x}x^2}{5} + \frac{2\sqrt{x}x}{15} - 2\log(x)$$

input `int(-2/x+1/5*x^(1/2)+x^(3/2),x)`

output `(2*(3*sqrt(x)*x**2 + sqrt(x)*x - 15*log(x)))/15`

# CHAPTER 4

## APPENDIX

4.1	Listing of Grading functions . . . . .	391
4.2	Links to plain text integration problems used in this report for each CAS .	409

### 4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

#### Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```



```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
  ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "
  ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]===Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]===Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]===Plus || Head[expn]===Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```

```

    Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
    If[AppellFunctionQ[Head[expn]],
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
      If[Head[expn]===RootSum,
        Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
        If[Head[expn]===Integrate || Head[expn]===Int,
          Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
          9]]]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

```

```

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

```

```

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

```

```

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

## Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022  add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result    := ExpnType(result);
      ExpnType_optimal   := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
#     is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

#   antiderivative
#   "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
            fi;
        fi;
    else #ExpnType(result) > ExpnType(optimal)
        if debug then
            print("ExpnType(result) > ExpnType(optimal)");
        fi;
        return "C",cat("Result contains higher order function than in optimal. Order ",
                        convert(ExpnType_result,string)," vs. order ",
                        convert(ExpnType_optimal,string),".");
    fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```



```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

## Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

## SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#      Albert Rich to use with Sagemath. This is used to
#      grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#      'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#      issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```



```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

## 4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file